

Random Process Background

Young W Lim

Nov 07, 2023

Copyright (c) 2023 - 2018 Young W. Lim. Permission is granted to copy, distribute and/or modify this document under the terms of the GNU Free Documentation License, Version 1.2 or any later version published by the Free Software Foundation; with no Invariant Sections, no Front-Cover Texts, and no Back-Cover Texts. A copy of the license is included in the section entitled "GNU Free Documentation License".

This work is licensed under a Creative Commons "Attribution-NonCommercial-ShareAlike 3.0 Unported" license.



Based on
Probability, Random Variables and Random Signal Principles,
P.Z. Peebles,Jr. and B. Shi

Outline

- 1 Open Sets and Classes
 - Open Set
 - Filter
 - Class
- 2 Borel Sets
 - Measurable Space
 - Topological Space
 - Borel Sets
- 3 Stochastic Process

Outline

- 1 Open Sets and Classes
 - Open Set
 - Filter
 - Class
- 2 Borel Sets
 - Measurable Space
 - Topological Space
 - Borel Sets
- 3 Stochastic Process

Open set examples

- The *circle* represents the set of points (x, y) satisfying $x^2 + y^2 = r^2$.
the *circle* set is its **boundary set**
- The *disk* represents the set of points (x, y) satisfying $x^2 + y^2 < r^2$.
The *disk* set is an **open set**
- the **union** of the *circle* and *disk* sets is a **closed set**.

https://en.wikipedia.org/wiki/Open_set

Open set (1)

- an **open set** is a generalization of an **open interval** in the real line.
- a **metric space** is a **set** along with a **distance** defined between any two **points**
- in a **metric space**, an **open set** is a **set** that, along with every **point** P , contains all **points** that are *sufficiently near* to P
 - all **points** whose **distance** to P is less than some value depending on P

https://en.wikipedia.org/wiki/Open_set

Open set (2)

- More generally, an **open set** is a **member** of a given collection of **subsets** of a given set a **collection** that has the property of **containing**

- every union of its **members**
- every finite intersection of its members
- the **empty set**
- the **whole set** itself

https://en.wikipedia.org/wiki/Open_set

Open set (3)

- These conditions are very loose, and allow enormous flexibility in the choice of **open sets**.
- For example,
 - every **subset** can be **open** (the **discrete topology**)
 - no **subset** can be **open** (the **indiscrete topology**) except
 - the space itself and
 - the empty set

https://en.wikipedia.org/wiki/Open_set

Open set (4)

- A **set** in which such a **collection** is given is called a **topological space**, and the **collection** is called a **topology**.
 - A **set** is a **collection** of distinct **objects**.
 - Given a **set** A , we say that a is an **element** of A if a is one of the distinct **objects** in A , and we write $a \in A$ to denote this
 - Given two **sets** A and B , we say that A is a **subset** of B if every element of A is also an element of B write $A \subseteq B$ to denote this.

<https://ximera.osu.edu/mooculus/calculusE/continuityOfFunctionsOfSeveralVariables/digInOpenAnd>

Open set (5) Open Balls


- An **open ball** $B_r(\mathbf{a})$ in \mathbb{R}^n
centered at $\mathbf{a} = (a_1, \dots, a_n) \in \mathbb{R}^n$ with radius r
is the set of all points $\mathbf{x} = (x_1, \dots, x_n) \in \mathbb{R}^n$
such that the distance between \mathbf{x} and \mathbf{a} is less than r
- In \mathbb{R}^2 an **open ball** is often called an **open disk**

We give these definitions in general, for when one is working in \mathbb{R}^n
since they are really not all that different to define in \mathbb{R}^n than in \mathbb{R}^2


<https://ximera.osu.edu/mooculus/calculusE/continuityOfFunctionsOfSeveralVariables/digInOpenAnd>

Open set (6) Interior points

- Suppose that $S \subseteq \mathbb{R}^n$
- A point $\mathbf{p} \in S$ is an **interior point** of S if there exists an **open ball** $B_r(\mathbf{p}) \subseteq S$
- Intuitively, \mathbf{p} is an **interior point** of S if we can squeeze an entire **open ball** centered at \mathbf{p} within S




an interior point




a boundary point

Open set (7) Boundary points

- A point $\mathbf{p} \in \mathbb{R}^n$ is a **boundary point** of S if all **open balls** centered at \mathbf{p} contain both **points** in S and **points** not in S
- The **boundary** of S is the **set** ∂S that consists of all of the **boundary points** of S .



an interior point



a boundary point

Open set (8) Open and Closed Sets

- A set $O \subseteq \mathbb{R}^n$ is **open** if every point in O is an **interior point**.
- A set $C \subseteq \mathbb{R}^n$ is **closed** if it contains all of its **boundary points**.

<https://ximera.osu.edu/mooculus/calculusE/continuityOfFunctionsOfSeveralVariables/digInOpenAnd>

Open set (9) Bounded and Unbounded

- A set S is **bounded** if there is an **open ball** $B_M(0)$ such that

$$S \subseteq B.$$

intuitively, this means that we can enclose all of the **set** S within a large enough **ball** centered at the origin, $B_M(0)$

- A **set** that is not **bounded** is called **unbounded**

<https://ximera.osu.edu/mooculus/calculusE/continuityOfFunctionsOfSeveralVariables/digInOpenAnd>

Family of sets (1)

- a **collection** F of **subsets** of a given **set** S is called a **family** of **subsets** of S , or a **family** of **sets** over S .
- More generally, a **collection** of any **sets** whatsoever is called a **family** of **sets**, **set family**, or a **set system**

https://en.wikipedia.org/wiki/Family_of_sets

Family of sets (2)

- The term "**collection**" is used here because,
 - in some contexts,
a **family** of **sets** may be allowed
to contain repeated copies of any given **member**, and
 - in other contexts
it may form a **proper class** rather than a **set**.

https://en.wikipedia.org/wiki/Family_of_sets

Examples of family of sets (1)

- The **set** of all **subsets** of a given **set** S is called the **power set** of S and is denoted by $\wp(S)$.

The **power set** $\wp(S)$ of a given **set** S is a **family** of **sets** over S .

- A **subset** of S having k elements is called a **k -subset** of S .

The **k -subset** $S^{(k)}$ of a set S form a **family** of **sets**.

https://en.wikipedia.org/wiki/Family_of_sets

Examples of family of sets (2)

- Let $S = \{a, b, c, 1, 2\}$.

An example of a **family** of **sets** over S

(in the multiset sense) is given by $F = \{A_1, A_2, A_3, A_4\}$, where

$A_1 = \{a, b, c\}$, $A_2 = \{1, 2\}$, $A_3 = \{1, 2\}$, and $A_4 = \{a, b, 1\}$.

https://en.wikipedia.org/wiki/Family_of_sets

Neighbourhood basis (1)

- A **neighbourhood basis** or **local basis** (or **neighbourhood base** or **local base**) for a **point** x is a **filter base** of the **neighbourhood filter**;
- this means that it is a **subset** $\mathcal{B} \subseteq \mathcal{N}(x)$ such that for all $V \in \mathcal{N}(x)$, there exists some $B \in \mathcal{B}$ such that $B \subseteq V$.
That is, for any **neighbourhood** V we can find a **neighbourhood** B in the **neighbourhood basis** that is contained in V .

https://en.wikipedia.org/wiki/Neighbourhood_system#Neighbourhood_basis

Neighbourhood basis (2)

- Equivalently, \mathcal{B} is a local basis at x if and only if the neighbourhood filter \mathcal{N} can be recovered from \mathcal{B} in the sense that the following equality holds:

$$\mathcal{N}(x) = \{V \subseteq X : B \subseteq V \text{ for some } B \in \mathcal{B}\}$$

- A family $\mathcal{B} \subseteq \mathcal{N}(x)$ is a neighbourhood basis for x if and only if \mathcal{B} is a cofinal subset of $(\mathcal{N}(x), \supseteq)$ with respect to the partial order \supseteq (importantly, this partial order is the superset relation and not the subset relation).

https://en.wikipedia.org/wiki/Neighbourhood_system#Neighbourhood_basis

A collection of sets around x

- In general, one refers to the family of **sets** containing 0, used to **approximate** 0, as a **neighborhood basis**;
- a **member** of this **neighborhood basis** is referred to as an **open set**.
- In fact, one may generalize these notions to an arbitrary set (X); rather than just the **real numbers**.
- In this case, given a **point** (x) of that **set** (X), one may define a **collection** of **sets** "**around**" (that is, containing) x , used to **approximate** x .

https://en.wikipedia.org/wiki/Open_set

Smaller sets containing x

- Of course, this **collection** would have to *satisfy* certain properties (known as **axioms**) for otherwise we may not have a *well-defined method* to measure **distance**.
- For example, every **point** in X should **approximate** x to some **degree** of **accuracy**.
- Thus X should be in this **family**.
- Once we begin to define "smaller" **sets** containing x , we tend to **approximate** x to a greater **degree** of **accuracy**.
- Bearing this in mind, one may define the remaining **axioms** that the **family** of **sets** about x is required to satisfy.

https://en.wikipedia.org/wiki/Open_set

Open ball (1)

- a **ball** is the solid figure bounded by a **sphere**;
it is also called a **solid sphere**.
 - a **closed ball**
includes the *boundary points* that constitute the sphere
 - an **open ball**
excludes them

[https://en.wikipedia.org/wiki/Ball_\(mathematics\)](https://en.wikipedia.org/wiki/Ball_(mathematics))

Open ball (2)

- A **ball** in n dimensions is called a **hyperball** or **n-ball** and is bounded by a **hypersphere** or $(n - 1)$ -sphere
- One may talk about **balls** in any **topological space** X , not necessarily induced by a **metric**.
- An n -dimensional **topological ball** of X is any **subset** of X which is **homeomorphic** to an **Euclidean n-ball**.

[https://en.wikipedia.org/wiki/Ball_\(mathematics\)](https://en.wikipedia.org/wiki/Ball_(mathematics))

Outline

- 1 Open Sets and Classes
 - Open Set
 - **Filter**
 - Class
- 2 Borel Sets
 - Measurable Space
 - Topological Space
 - Borel Sets
- 3 Stochastic Process

Homogeneous Relation

- a homogeneous relation (also called endorelation) on a set X is a binary relation between X and itself, i.e. it is a subset of the Cartesian product $X \times X$.
- This is commonly phrased as "a relation on X " or "a (binary) relation over X ".
- An example of a homogeneous relation is the relation of kinship, where the relation is between people.

https://en.wikipedia.org/wiki/Homogeneous_relation

Binary Relation (1)

- a binary relation associates elements of one set, called the domain, with elements of another set, called the codomain.
- A binary relation over sets X and Y is a new set of ordered pairs (x,y) consisting of elements x from X and y from Y .

https://en.wikipedia.org/wiki/Binary_relation

Binary Relation (2)

- It is a generalization of a unary function.
- It encodes the common concept of relation:
- an element x is related to an element y ,
if and only if the pair (x, y) belongs
to the set of ordered pairs that defines the binary relation.
- A binary relation is the most studied special case $n = 2$
of an n -ary relation over sets X_1, \dots, X_n ,
which is a subset of the Cartesian product $X_1 \times \dots \times X_n$.

https://en.wikipedia.org/wiki/Binary_relation

Partially Ordered Set (1-1)

- a **partial order** on a **set** is an arrangement such that, for certain **pairs** of elements, one precedes the other.
- The word **partial** is used to indicate that not every **pair** of elements needs to be comparable; that is, there may be **pairs** for which neither element precedes the other.
- **Partial orders** thus generalize **total orders**, in which every pair is comparable.

https://en.wikipedia.org/wiki/Partially_ordered_set

Partially Ordered Set (1-2)

- Formally, a **partial order** is a **homogeneous binary relation** that is **reflexive**, **transitive** and **antisymmetric**.
- A **partially ordered set** (**poset** for short) is a set on which a **partial order** is defined.
- A **reflexive**, **weak**, or **non-strict partial order**, commonly referred to simply as a **partial order**, is a **homogeneous relation** \leq on a **set** P that is **reflexive**, **antisymmetric**, and **transitive**.

https://en.wikipedia.org/wiki/Partially_ordered_set

Partially Ordered Set (2)

- a **homogeneous relation** \leq on a **set** P that is **reflexive**, **antisymmetric**, and **transitive**.
- That is, for all $a, b, c \in P$, it must satisfy:
 - **Reflexivity**:
 $a \leq a$, i.e. every element is related to itself.
 - **Antisymmetry**:
if $a \leq b$ and $b \leq a$ then $a = b$,
i.e. no two distinct elements precede each other.
 - **Transitivity**:
if $a \leq b$ and $b \leq c$ then $a \leq c$.
- A **non-strict partial order** is also known as an **antisymmetric preorder**.

https://en.wikipedia.org/wiki/Partially_ordered_set

Filter in Set Theory (1-1)

- A **filter** on a **set** may be thought of as representing a "**collection** of *large subsets*", one intuitive example being the **neighborhood filter**.
- keep *large* grains excluding *small* impurities

[https://en.wikipedia.org/wiki/Filter_\(set_theory\)#filter_base](https://en.wikipedia.org/wiki/Filter_(set_theory)#filter_base)

Filter in Set Theory (1-2)

- When you put a filter in your sink, the idea is that you filter out the *big* chunks of food, and let the water and the *smaller* chunks go through (which can, in principle, be washed through the pipes) .
- You filter out the *larger parts*.
- A filter filters out the *larger sets*.
- It is a way to say "these *sets* are '*large*'"

[https://en.wikipedia.org/wiki/Filter_\(set_theory\)#filter_base](https://en.wikipedia.org/wiki/Filter_(set_theory)#filter_base)

Filter in Set Theory (1-3)

- a **filter** on a **set** X is a **family** \mathcal{B} of **subsets** such that:

- 1 $X \in \mathcal{B}$ and $\emptyset \notin \mathcal{B}$
- 2 if $A \in \mathcal{B}$ and $B \in \mathcal{B}$,
then $A \cap B \in \mathcal{B}$
- 3 If $A, B \subset X, A \in \mathcal{B}$, and $A \subset B$,
then $B \in \mathcal{B}$

[https://en.wikipedia.org/wiki/Filter_\(set_theory\)#filter_base](https://en.wikipedia.org/wiki/Filter_(set_theory)#filter_base)

Filter in Set Theory (1-4)

- The set of "everything" is definitely *large*

$$X \in \mathcal{B}$$

- and "nothing" is definitely not;

$$\emptyset \notin \mathcal{B}$$

- if something is *larger* than a *large set*, then it is also *large*;

$$\text{If } A, B \subset X, A \in \mathcal{B}, \text{ and } A \subset B, \text{ then } B \in \mathcal{B}$$

- and two *large sets intersect* on a *large set*.

$$\text{If } A \in \mathcal{B} \text{ and } B \in \mathcal{B}, \text{ then } A \cap B \in \mathcal{B}$$

[https://en.wikipedia.org/wiki/Filter_\(set_theory\)#filter_base](https://en.wikipedia.org/wiki/Filter_(set_theory)#filter_base)



Filter in Set Theory (1-5)

- you can think about this as
 - being **co-finite**,
 - or being of **measure 1** on the **unit interval**,
 - or having a **dense open subset** (again on the unit interval).
- These are examples of ways where a [set](#) can be thought of as "almost everything". and that is the idea behind a filter.

[https://en.wikipedia.org/wiki/Filter_\(set_theory\)#filter_base](https://en.wikipedia.org/wiki/Filter_(set_theory)#filter_base)

Co-finite

- a **cofinite subset** of a set X is a subset A whose complement in X is a finite set.
- a subset A contains all but *finitely many* elements of X
- If the complement is not finite, but is countable, then one says the set is **cocountable**.
- These arise naturally when generalizing structures on finite sets to infinite sets, particularly on infinite products, as in the **product topology** or direct sum.
- This use of the prefix "co" to describe a property possessed by a set's complement is *consistent* with its use in other terms such as "comeagre set".

<https://en.wikipedia.org/wiki/Cofiniteness>

Unit interval

- the **unit interval** is the **closed interval** $[0,1]$, that is, the set of all real numbers that are greater than or equal to 0 and less than or equal to 1.
- It is often denoted I (capital letter I).
- In addition to its role in **real analysis**, the **unit interval** is used to study **homotopy theory** in the field of **topology**.
- the term "**unit interval**" is sometimes applied to the other shapes that an interval from 0 to 1 could take: $(0,1]$, $[0,1)$, and $(0,1)$.
- However, the notation I is most commonly reserved for the **closed interval** $[0,1]$.

Dense set

- In **topology**, a **subset** A of a topological space X is said to be **dense** in X if every **point** of X either belongs to A or else is arbitrarily "close" to a **member** of A
 - for instance, the **rational numbers** are a **dense** subset of the **real numbers** because every **real number** either is a **rational number** or has a rational number arbitrarily close to it (see **Diophantine approximation**).
- Formally, A is **dense** in X if the *smallest* **closed subset** of X containing A is X itself.
- The **density** of a **topological space** X is the **least cardinality** of a **dense subset** of X .

https://en.wikipedia.org/wiki/Dense_set

Ultrafilter (1)

- an ultrafilter on a given partially ordered set (or "poset") P is a certain subset of P , namely a maximal filter on P ; that is, a proper filter on P that cannot be enlarged to a bigger proper filter on P .
- If X is an arbitrary set, its power set $\mathcal{P}(X)$, ordered by set inclusion, is always a Boolean algebra and hence a poset, and ultrafilters on $\mathcal{P}(X)$ are usually called ultrafilter on the set X .
- An ultrafilter on a set X may be considered as a finitely additive measure on X .
- In this view, every subset of X is either considered "almost everything" (has measure 1) or "almost nothing" (has measure 0), depending on whether it belongs to the given ultrafilter or not

<https://en.wikipedia.org/wiki/Ultrafilter>

Ultrafilter on partial orders (1)

- In order theory, an ultrafilter is a subset of a partially ordered set that is maximal among all proper filters. This implies that any filter that properly contains an ultrafilter has to be equal to the whole poset.
- Formally, if P is a set, partially ordered by \leq then

<https://en.wikipedia.org/wiki/Ultrafilter>

Ultrafilter on partial orders (2)

- a subset $F \subseteq P$ is called a filter on P if F is nonempty, for every $x, y \in F$, there exists some element $z \in F$ such that $z \leq x$ and $z \leq y$, and for every $x \in F$ and $y \in P$, $x \leq y$ implies that y is in F too; a proper subset U of P is called an ultrafilter on P if U is a filter on P , and there is no proper filter F on P that properly extends U (that is, such that U is a proper subset of F).

<https://en.wikipedia.org/wiki/Ultrafilter>

Filter in Set Theory (2-1)

- Let $X = 1, 2, 3$
Choose some element from X say $F = 1, 1, 2, 1, 3, 1, 2, 3$
- Then every **intersection** of an element of F with another element in F is in F again.
Examples: $1 \cap 1, 2, 3 = 1$ $1, 2 \cap 1, 2, 3 = 1, 2$
 $1, 3 \cap 1, 2, 3 = 1, 3$ $1, 2, 3 \cap 1, 2, 3 = 1, 2, 3$
- Also the original $X = 1, 2, 3$ is also in F .
Here $F = 1, 1, 2, 1, 3, 1, 2, 3$ is called the **filter** on $X = 1, 2, 3$

<https://math.stackexchange.com/questions/2816362/meaning-behind-filter-in-set-theory>

Filter in Set Theory (2-2)

- Suppose we have the **collection** $G = 1, 1, 2, 1, 3, 2, 3, 1, 2, 3$
- Then we have $1, 3 \cap 2, 3 = 3$ but 3 isn't in G .
So this G is not called a **filter**.
- Now with $F = 1, 1, 2, 1, 3, 1, 2, 3$
can we put as any other **element** in it
so that after placing the **extra element** it is still a **filter**?
Probably not in this case.
So on $X = 1, 2, 3$, $F = 1, 1, 2, 1, 3, 1, 2, 3$ is an **Ultrafilter**.

<https://math.stackexchange.com/questions/2816362/meaning-behind-filter-in-set-theory>

Filter in Set Theory (3-1)

- If we have started say with $H = 1, 1, 2, 1, 2, 3$
this is still a **filter** on $X = 1, 2, 3$
but we can still add $1, 3$
and it will still be classified as **filter**.
- So on $X = 1, 2, 3$
 $F = 1, 1, 2, 1, 3, 1, 2, 3$ is an **Ultrafilter**
but $H = 1, 1, 2, 1, 2, 3$ is a filter but not an **Ultrafilter**.

<https://math.stackexchange.com/questions/2816362/meaning-behind-filter-in-set-theory>

Filter in Set Theory (3-2)

- Now suppose we have $X = 1, 2, 3, 4$
Let $F = 1, 4, 1, 2, 4, 1, 3, 4, 1, 2, 3, 4$
- Every in intersection of element of F is in F again.
We have as examples $1, 4 \cap 1, 4 = 1, 4$ $1, 4 \cap 1, 2, 4 = 1, 4$
 $1, 4 \cap 1, 3, 4 = 1, 4$ $1, 2, 4 \cap 1, 2, 4 = 1, 2, 4$ $1, 2, 4 \cap 1, 3, 4 = 1, 4$
 $1, 3, 4 \cap 1, 3, 4 = 1, 3, 4$ $1, 2, 3, 4 \cap 1, 2, 3, 4 = 1, 2, 3, 4$
- Also $X = 1, 2, 3, 4$ is also in $F = 1, 4, 1, 2, 4, 1, 3, 4, 1, 2, 3, 4$
and the null element $\emptyset =$ is not in F .

<https://math.stackexchange.com/questions/2816362/meaning-behind-filter-in-set-theory>

Filter in Set Theory (3-3)

- We call F a filter but not an Ultrafilter on $X = 1, 2, 3, 4$
- We can still add element in it and it will still be a filter for instance by adding the element 1 from $X = 1, 2, 3, 4$ we can have the filter $F = 1, 1, 4, 1, 2, 4, 1, 3, 4, 1, 2, 3, 4$
- This is an Ultrafilter on $X = 1, 2, 3, 4$ as we cannot add any further element from $X = 1, 2, 3, 4$ that satisfies closures on intersection.

<https://math.stackexchange.com/questions/2816362/meaning-behind-filter-in-set-theory>

Filter in Set Theory (4)

- There is another collection of sets taken from $X=1,2,3,4$ Which is the powerset $P=,1,2,3,4,1,2,1,3,1,4,2,3,2,4,3,4,1,2,3,1,2,4,1,3,4,2,3,4,1,2,3,4$ This contain the null element $\emptyset=$ so we cannot call this as Ultrafilter. This is not a proper filter according to the article in Wikipedia. In the powerset every intersection of element is again in the powerset again but it contains the null element $\emptyset=$ and isn't classified as proper filter.

<https://math.stackexchange.com/questions/2816362/meaning-behind-filter-in-set-theory>

Outline

- 1 Open Sets and Classes
 - Open Set
 - Filter
 - Class
- 2 Borel Sets
 - Measurable Space
 - Topological Space
 - Borel Sets
- 3 Stochastic Process

Class (1)

- a **class** is a **collection** of **sets**
(or sometimes other **mathematical objects**)
that can be unambiguously defined
by a **property** that all its members share.
- **Classes** act as a way to have **set-like collections**
while **differing** from **sets** so as to **avoid Russell's paradox**

[https://en.wikipedia.org/wiki/Class_\(set_theory\)](https://en.wikipedia.org/wiki/Class_(set_theory))

Class (2)

- A class *that is not a set* is called a **proper class**, and
- a class *that is a set* is sometimes called a **small class**.
- the followings are **proper classes** in many formal systems
 - the **class** of all sets
 - the **class** of all ordinal numbers
 - the **class** of all cardinal numbers

[https://en.wikipedia.org/wiki/Class_\(set_theory\)](https://en.wikipedia.org/wiki/Class_(set_theory))

Class (3)

- consider "the **set** of all **sets** with **property** X ."
- especially when dealing with **categories**, since the **objects** of a **concrete category** are all **sets** with certain additional **structure**.
- However, **if** we're not *careful* about this we can get into serious *trouble* –

<https://www.quora.com/In-set-theory-what-is-the-difference-between-a-set-of-objects-and-a-class-of-objects>

Class (4)

- let X be the **set** of all **sets** which do not contain *themselves*
- Since X is a **set**, we can ask whether X is an element of *itself*.
- But then we run into a **paradox** – **if** X contains *itself* as an element, **then** it does not, and vice versa.

<https://www.quora.com/In-set-theory-what-is-the-difference-between-a-set-of-objects-and-a-class-of-objects>

Class (5)

- In order to avoid this **paradox**, we cannot consider the **collection** of all **sets** to be itself a **set**.
- This means we have to *throw out* the whole "the **set** of all **sets** with **property** X" construction. But we wanted that.
- So the way we get around it is to say that there's something called a **class**, which is like a **set** but not a **set**.

<https://www.quora.com/In-set-theory-what-is-the-difference-between-a-set-of-objects-and-a-class-of-objects>

Class (6)

- Then we can talk about "the class X of all sets with property Y ."
- Since X is not a set, it can't be an element of itself, and we're fine.
- Of course, if we need to talk about the collection of all classes, we need to create another name that goes another step back, and so forth.

<https://www.quora.com/In-set-theory-what-is-the-difference-between-a-set-of-objects-and-a-class-of-objects>

Class Examples (1)

- The **collection** of all **algebraic structures** of a given type will usually be a **proper class**.
(a **class** *that is not a set* is called a **proper class**)
 - the **class** of all **groups**
 - the **class** of all **vector spaces**
 - and many others.
- Within set theory, many **collections** of **sets** turn out to be **proper classes**.

[https://en.wikipedia.org/wiki/Class_\(set_theory\)](https://en.wikipedia.org/wiki/Class_(set_theory))

Class Examples (2)

- One way to *prove* that a **class** is **proper** is to place it in **bijection** with the **class** of all ordinal numbers.
 - **Cardinal numbers** indicate an amount how many of something we have: one, two, three, four, five.
 - **Ordinal numbers** indicate position in a series: first, second, third, fourth, fifth.

[https://en.wikipedia.org/wiki/Class_\(set_theory\)](https://en.wikipedia.org/wiki/Class_(set_theory))
<https://editarians.com/cardinals-ordinals/>

Class Paradoxes (1)

- The **paradoxes** of naive set theory can be explained in terms of the *inconsistent tacit assumption* that "all **classes** are **sets**".
- These **paradoxes** do not arise with **classes** because there is no notion of **classes** containing **classes**.
- Otherwise, one could, for example, define a **class** of all **classes** that do not contain themselves, which would lead to a **Russell paradox** for classes.

[https://en.wikipedia.org/wiki/Class_\(set_theory\)](https://en.wikipedia.org/wiki/Class_(set_theory))

Class Paradoxes (2)

- With a rigorous foundation, these **paradoxes** instead *suggest proofs* that certain **classes** are **proper** (i.e., that they are not **sets**).
 - **Russell's paradox** *suggests a proof* that the **class** of all **sets** which do not contain themselves is **proper**
 - the **Burali-Forti paradox** *suggests* that the **class** of all ordinal numbers is **proper**.

[https://en.wikipedia.org/wiki/Class_\(set_theory\)](https://en.wikipedia.org/wiki/Class_(set_theory))

Russell's Paradox (1)

- According to the unrestricted comprehension principle, for any sufficiently well-defined **property**, there is the **set** of **all** and only the **objects** that have that **property**.

https://en.wikipedia.org/wiki/Russell%27s_paradox

Russell's Paradox (2)

- Let R be the **set of all sets** ($R = \{x \mid x \notin x\}$) that are not members of themselves ($R \notin R$).
 - *if* R is not a **member** of itself ($R \notin R$), *then* its definition (the **set of all sets**) entails that it is a **member** of itself ($R \in R$);
 - yet, *if* it is a **member** of itself ($R \in R$), *then* it is not a **member** of itself ($R \notin R$), since it is the **set of all sets** that are not members of themselves ($R \notin R$)
- the resulting **contradiction** is **Russell's paradox**.
- Let $R = \{x \mid x \notin x\}$, then $R \in R \iff R \notin R$

https://en.wikipedia.org/wiki/Russell%27s_paradox

Russell's Paradox (3)

- Most *sets* commonly encountered are not *members* of themselves.
- For example, consider the *set* of all squares in a plane.
- This *set* is not itself a square in the plane, thus it is not a *member* of itself.
- Let us call a *set* "**normal**" if it is not a *member* of itself, and "**abnormal**" if it is a *member* of itself.

https://en.wikipedia.org/wiki/Russell%27s_paradox

Russell's Paradox (4)

- Clearly every **set** must be either **normal** or **abnormal**.
- The **set** of squares in the plane is **normal**.
- In contrast, the **complementary set** that contains everything which is not a square in the plane is itself not a square in the plane, and so it is one of its own **members** and is therefore **abnormal**.

https://en.wikipedia.org/wiki/Russell%27s_paradox

Russell's Paradox (5)

- Now we consider the **set** of all **normal sets**, R , and try to determine whether R is **normal** or **abnormal**.
 - *If* R were **normal**, it would be contained in the **set** of all **normal sets** (itself), and therefore be **abnormal**;
 - on the other hand *if* R were **abnormal**, it would not be contained in the **set** of all **normal sets** (itself), and therefore be **normal**.
- This leads to the conclusion that R is neither **normal** nor **abnormal**: **Russell's paradox**.

https://en.wikipedia.org/wiki/Russell%27s_paradox

Outline

- 1 Open Sets and Classes
 - Open Set
 - Filter
 - Class
- 2 Borel Sets
 - Measurable Space
 - Topological Space
 - Borel Sets
- 3 Stochastic Process

Mathematical objects (1)

- a **mathematical object** is an **abstract concept** arising in mathematics.
- an **mathematical object** is anything that has been (or could be) **formally defined**, and with which one may do
 - **deductive reasoning**
 - **mathematical proofs**

https://en.wikipedia.org/wiki/Mathematical_object

Mathematical objects (2)

- typically, a **mathematical object**
 - can be a value that can be assigned to a variable
 - therefore can be involved in formulas

https://en.wikipedia.org/wiki/Mathematical_object

Mathematical objects (3)

- commonly encountered **mathematical objects** include
 - numbers
 - sets
 - functions
 - expressions
 - geometric objects
 - transformations of other mathematical objects
 - spaces

https://en.wikipedia.org/wiki/Mathematical_object

Mathematical objects (4)

- **Mathematical objects** can be very *complex*;
 - for example, the followings are considered as **mathematical objects** in **proof theory**.
 - theorems
 - proofs
 - theories

https://en.wikipedia.org/wiki/Mathematical_object

Structure (1)

- a **structure** is a **set** endowed with some *additional features* on the **set**
 - an *operation*
 - *relation*
 - *metric*
 - *topology*
- often, the *additional features* are attached or related to the **set**, so as to provide it with some *additional meaning* or *significance*.

<https://www.localmaxradio.com/questions/what-is-a-mathematical-space>

Structure (2)

- A partial list of possible **structures** are
 - measures
 - algebraic structures (groups, fields, etc.)
 - topologies
 - metric structures (geometries)
 - orders
 - events
 - equivalence relations
 - differential structures
 - categories.

<https://www.localmaxradio.com/questions/what-is-a-mathematical-space>

Space (1)

- A **space** consists of selected **mathematical objects** that are treated as **points**, and selected **relationships** between these **points**.
 - the *nature* of the **points** can vary widely:
for example, the **points** can be
 - elements of a set
 - functions on another space
 - subspaces of another space
 - It is the **relationships** between **points** that define the *nature* of the **space**.

[https://en.wikipedia.org/wiki/Space_\(mathematics\)](https://en.wikipedia.org/wiki/Space_(mathematics))

Space (2)

- *modern mathematics* uses many types of **spaces**, such as
 - Euclidean spaces
 - linear spaces
 - topological spaces
 - Hilbert spaces
 - probability spaces
- *modern mathematics* does not define the notion of **space** itself.

[https://en.wikipedia.org/wiki/Space_\(mathematics\)](https://en.wikipedia.org/wiki/Space_(mathematics))

Space (3)

- a **space** is
a **set** (or a **universe**) with some added **features**
- it is not always clear
whether a given **mathematical object** should be considered
as a **geometric space**, or an **algebraic structure**
- a general definition of **structure** embraces
all common types of **space**

[https://en.wikipedia.org/wiki/Space_\(mathematics\)](https://en.wikipedia.org/wiki/Space_(mathematics))

Mathematical space (1)

- A **mathematical space** is, informally, a **collection** of **mathematical objects** under consideration.
- The **universe** of **mathematical objects** within a **space** are *precisely defined entities* whose **rules** of *interaction* come baked into the **rules** of the **space**.

<https://www.localmaxradio.com/questions/what-is-a-mathematical-space>

Mathematical space (2)

- A **space** differs from a **mathematical set** in several important ways:
 - A **mathematical set** is also a **collection** of **objects**
 - but these **objects** are being pulled from a **space** (or **universe**) of **objects** where the **rules** and **definitions** have already been agreed upon

<https://www.localmaxradio.com/questions/what-is-a-mathematical-space>

Mathematical space (3)

- A **space** differs from a **mathematical set** in several important ways:
 - a **mathematical set** has no **internal structure**,
 - a **space** usually has some **internal structure**.

<https://www.localmaxradio.com/questions/what-is-a-mathematical-space>

Mathematical space (4)

- having some **internal structure** could mean a variety of things, but typically it involves
 - *interactions* and *relationships* between **elements** of the **space**
 - *rules* on how to *create* and *define* **new elements** of the **space**

<https://www.localmaxradio.com/questions/what-is-a-mathematical-space>

Measurable space (1)

- A measurable space is any space with a sigma-algebra which can then be equipped with a measure
 - collection of subsets of the space following certain rules with a way to assign sizes to those sets.

<https://www.quora.com/What-is-a-measurable-space-and-probability-space-intuitively-What-differences-do-they-have>

Measurable space (2)

- Intuitively, certain **sets** belonging to a **measurable space** can be given a **size** in a *consistent way*.

consistent way means that certain **axioms** are met:

- the **empty set** is given a **size** of **zero**
- if a measurable set is **contained** inside another one, then its **size** is **less than** or **equal to** the size of the **containing set**
- the size of a **disjoint union** of sets is the **sum** of the individual sets' **sizes**

<https://www.quora.com/What-is-a-measurable-space-and-probability-space-intuitively-What-differences-do-they-have>

The set of all real numbers

- In the [set](#) of all [real numbers](#), one has the natural [Euclidean metric](#); that is, a function which *measures* the [distance](#) between two [real numbers](#): $d(x, y) = |x - y|$.

https://en.wikipedia.org/wiki/Open_set

All points close to a real number x

- Therefore, given a **real number** x , one can speak of the **set** of all **points close** to that **real number** x ; that is, **within** ε of x .
- In essence, **points within** ε of x **approximate** x to an **accuracy** of **degree** ε .
- Note that $\varepsilon > 0$ always, but as ε becomes *smaller* and *smaller*, one obtains **points** that **approximate** x to a *higher* and *higher* **degree** of **accuracy**.

https://en.wikipedia.org/wiki/Open_set

The points within ε of x

- For example, if $x = 0$ and $\varepsilon = 1$, the **points** within ε of x are precisely the **points** of the interval $(-1, 1)$;
- However, with $\varepsilon = 0.5$, the **points** within ε of x are precisely the **points** of $(-0.5, 0.5)$.
- Clearly, these **points** approximate x to a *greater degree* of **accuracy** than when $\varepsilon = 1$.

https://en.wikipedia.org/wiki/Open_set

without a concrete Euclidean metric

- The previous examples shows, for the case $x = 0$, that one may approximate x to *higher and higher* degrees of accuracy by defining ε to be *smaller and smaller*.
- In particular, sets of the form $(-\varepsilon, \varepsilon)$ give us a lot of information about **points close** to $x = 0$.
- Thus, rather than speaking of a concrete Euclidean metric, one may use **sets** to describe **points close** to x .

https://en.wikipedia.org/wiki/Open_set

Different collections of sets containing 0

- This innovative idea has far-reaching consequences; in particular, by defining

different collections of sets containing 0
(distinct from the sets $(-\varepsilon, \varepsilon)$),
one may find different results
regarding the distance
between 0 and other real numbers.

https://en.wikipedia.org/wiki/Open_set

A set for measuring distance

- For example, if we were to define R as the *only* such set for "*measuring distance*", all points are close to 0
- since there is only one possible degree of accuracy one may achieve in approximating 0: being a member of R .

https://en.wikipedia.org/wiki/Open_set

The measure as a binary condition

- Thus, we find that in some sense, every real number is **distance** 0 away from 0.
- It may help in this case to think of the **measure** as being a **binary condition**:
 - all things in \mathbf{R} are equally close to 0,
 - while any item that is not in \mathbf{R} is not close to 0.

https://en.wikipedia.org/wiki/Open_set

Probability space

- A **probability space** is simply a **measurable space** equipped with a **probability measure**.
- A **probability measure** has the special property of giving the entire space a size of **1**.
 - this then implies that the **size** of any disjoint union of sets (the sum of the **sizes** of the sets) in the **probability space** is less than or equal to **1**

<https://www.quora.com/What-is-a-measurable-space-and-probability-space-intuitively-What-differences-do-they-have>

Euclidean space definition (1)

- A **subset** U of the **Euclidean n-space** \mathbb{R}^n is open
if, for every **point** x in U ,
there exists a positive **real number** ε
(depending on x)
such that any **point** in \mathbb{R}^n
whose **Euclidean distance** from x is smaller than ε
belongs to U

https://en.wikipedia.org/wiki/Open_set

Euclidean space definition (2)

- Equivalently, a subset U of \mathbb{R}^n is open if every point in U is the center of an open ball contained in U
- An example of a subset of \mathbb{R} that is not open is the closed interval $[0, 1]$, since neither $0 - \varepsilon$ nor $1 + \varepsilon$ belongs to $[0, 1]$ for any $\varepsilon > 0$, no matter how small.

https://en.wikipedia.org/wiki/Open_set

Metric space definition (1)

- A **subset** U of a **metric space** (M, d) is called **open**
if, for any **point** x in U , there exists a **real number** $\varepsilon > 0$
such that any **point** $y \in M$ satisfying $d(x, y) < \varepsilon$ belongs to U .
- Equivalently, U is **open**
if every **point** in U
has a **neighborhood** contained in U .
- This generalizes the **Euclidean space** example,
since **Euclidean space** with the **Euclidean distance**
is a **metric space**.

https://en.wikipedia.org/wiki/Open_set

Metric space definition (2)

- formally, a **metric space** is an **ordered pair** (M, d) where M is a **set** and d is a **metric** on M , i.e., a **function**

$$d : M \times M \rightarrow \mathbb{R}$$

satisfying the following **axioms** for all points $x, y, z \in M$:

- $d(x, x) = 0$.
- if $x \neq y$, then $d(x, y) > 0$.
- $d(x, y) = d(y, x)$.
- $d(x, z) \leq d(x, y) + d(y, z)$.

https://en.wikipedia.org/wiki/Open_set

Metric space definition (3)

- satisfying the following **axioms** for all points $x, y, z \in M$:
 - the distance from a point *to itself* is zero:
 - (**Positivity**) the **distance** between two distinct points is always **positive**:
 - (**Symmetry**) the **distance** from x to y is always the same as the **distance** from y to x :
 - (**Triangle inequality**) you can arrive at z from x by taking a detour through y , but this will not make your journey any faster than the shortest path.
- If the **metric** d is unambiguous, one often refers by abuse of notation to "the **metric space** M ".

https://en.wikipedia.org/wiki/Open_set

Outline

- 1 Open Sets and Classes
 - Open Set
 - Filter
 - Class
- 2 Borel Sets
 - Measurable Space
 - Topological Space
 - Borel Sets
- 3 Stochastic Process

Topology (1)

- **topology**
from the Greek words
τόπος, 'place, location',
and λόγος, 'study'

<https://en.wikipedia.org/wiki/Topology>

Topology (2)

- **topology** is concerned with the properties of a **geometric object** that are preserved
 - under **continuous deformations** such as
 - stretching
 - twisting
 - crumpling
 - bending
 - that is, without
 - closing holes
 - opening holes
 - tearing
 - gluing
 - passing through itself

<https://en.wikipedia.org/wiki/Topology>

Topological space (1)

- a **topological space** is, roughly speaking,
a **geometrical space**
in which **closeness** is defined
but cannot necessarily be **measured**
by a **numeric distance**.

https://en.wikipedia.org/wiki/Borel_set

Topological space (2)

- More specifically, a **topological space** is
 - a set whose elements are called points,
 - along with an additional structure called a topology,
- which can be defined as
 - a set of neighbourhoods for each point
 - that satisfy some axioms
formalizing the concept of closeness.

https://en.wikipedia.org/wiki/Borel_set

Topological space (3)

- There are several *equivalent definitions* of a **topology**, the most commonly used of which is the *definition* through *open sets*, which is easier than the others to manipulate.

https://en.wikipedia.org/wiki/Borel_set

Topological space (4)

- A **topological space** is the most **general type** of a **mathematical space** that allows for the definition of
 - **limits**
 - **continuity**
 - **connectedness**
- Although very **general**, the concept of **topological spaces** is **fundamental**, and used in virtually every branch of modern mathematics.
- The study of **topological spaces** in their own right is called **point-set topology** or **general topology**.

https://en.wikipedia.org/wiki/Topological_space

Topological space (5)

- Common types of **topological spaces** include
 - **Euclidean spaces** : a **set** of **points** satisfying certain **relationships**, expressible in terms of **distance** and **angles**.
 - **metric spaces** : a **set** together with a notion of **distance** between **points**. The **distance** is measured by a function called a **metric** or **distance function**.
 - **manifolds** : a topological space that *locally* resembles **Euclidean space** near each point. More precisely, an **n-manifold** is a topological space with the property that each **point** has a **neighborhood** that is **homeomorphic** to an **open subset** of **n-dimensional Euclidean space**.

https://en.wikipedia.org/wiki/Topological_space

Discrete Topology

- a **discrete space** is a **topological space**,
in which the **points** form a **discontinuous sequence**,
meaning they are isolated from each other in a certain sense.
- The **discrete topology** is
the finest **topology** that can be given on a **set**.
 - every **subset** is **open**
 - every **singleton subset** is an **open set**

https://en.wikipedia.org/wiki/Discrete_space

Singleton

- a **singleton**, also known as a **unit set** or **one-point set**, is a **set** with exactly one element.
- for example, the **set** $\{0\}$ is a **singleton** whose single element is 0

https://en.wikipedia.org/wiki/Discrete_space

Indiscrete Space (1)

- a **topological space** with the **trivial topology** is one where the only **open sets** are the **empty set** and the **entire space**.
- Such spaces are commonly called **indiscrete**, **anti-discrete**, **concrete** or **codiscrete**.
 - every **subset** can be **open** (the **discrete topology**), or
 - no **subset** can be **open** (the **indiscrete topology**) except the space itself and the empty set .

https://en.wikipedia.org/wiki/Discrete_space

Indiscrete Space (2)

- Intuitively, this has the consequence that all **points** of the **space** are "**lumped together**" and cannot be **distinguished** by topological means (not topologically **distinguishable** points)
- Every **indiscrete space** is a **pseudometric space** in which the **distance** between any two **points** is zero.

https://en.wikipedia.org/wiki/Discrete_space

T_0 Space

- a **topological space** X is a T_0 **space** or **if** for every **pair** of distinct points of X , at least one of them has a neighborhood not containing the other.
- In a T_0 **space**, all **points** are topologically distinguishable.
- This condition, called the T_0 **condition**, is the weakest of the **separation axioms**.
- Nearly all topological spaces *normally* studied in mathematics are T_0 **space**.

https://en.wikipedia.org/wiki/Kolmogorov_space

Topologically distinguishable points

- Intuitively, an **open set** provides a *method* to *distinguish* two **points**.
- two points in a **topological space**, there exists an **open set**
 - containing one point but
 - not containing the other (distinct) point
 - the two points are **topologically distinguishable**.

https://en.wikipedia.org/wiki/Open_set

Topologically distinguishable points

- Intuitively, an **open set** provides a *method* to *distinguish* two **points**.
- two points in a **topological space**, there exists an **open set**
 - containing one point but
 - not containing the other (distinct) point
 - the two points are **topologically distinguishable**.

https://en.wikipedia.org/wiki/Open_set

Metric spaces

- In this manner, one may speak of whether two **points**, or more generally two **subsets**, of a **topological space** are "**near**" without concretely defining a **distance**.
- Therefore, **topological spaces** may be seen as a generalization of **spaces** equipped with a notion of **distance**, which are called **metric spaces**.







https://en.wikipedia.org/wiki/Open_set

Examples of topology (1)

- Let τ be denoted with the circles, here are four examples **(a)**, **(b)**, **(c)**, **(d)**, and two non-examples **(e)**, **(f)** of topologies on the three-point set $\{1, 2, 3\}$.
- **(e)** is not a topology because the union of $\{2\}$ and $\{3\}$ [i.e. $\{2, 3\}$] is missing;
- **(f)** is not a topology because the intersection of $\{1, 2\}$ and $\{2, 3\}$ [i.e. $\{2\}$], is missing.

https://en.wikipedia.org/wiki/Topological_space

Examples of topology (2)



Every union of (c)

(c) is a topology $\{\{\}, \{1\}, \{2\}, \{1,2\}, \{1,2,3\}\}$
every union of (c)

\cup	$\{\}$	$\{1\}$	$\{2\}$	$\{1,2\}$	$\{1,2,3\}$
$\{\}$	$\{\}$	$\{1\}$	$\{2\}$	$\{1,2\}$	$\{1,2,3\}$
$\{1\}$	$\{1\}$	$\{1\}$	$\{1,2\}$	$\{1,2\}$	$\{1,2,3\}$
$\{2\}$	$\{2\}$	$\{1,2\}$	$\{2\}$	$\{1,2\}$	$\{1,2,3\}$
$\{1,2\}$	$\{1,2\}$	$\{1,2\}$	$\{1,2\}$	$\{1,2\}$	$\{1,2,3\}$
$\{1,2,3\}$	$\{1,2,3\}$	$\{1,2,3\}$	$\{1,2,3\}$	$\{1,2,3\}$	$\{1,2,3\}$

https://en.wikipedia.org/wiki/Topological_space

Every intersection of (c)

(c) is a topology $\{\{\}, \{1\}, \{2\}, \{1,2\}, \{1,2,3\}\}$
every intersection of (c)

\cap	$\{\}$	$\{1\}$	$\{2\}$	$\{1,2\}$	$\{1,2,3\}$
$\{\}$	$\{\}$	$\{\}$	$\{\}$	$\{\}$	$\{\}$
$\{1\}$	$\{\}$	$\{1\}$	$\{\}$	$\{1\}$	$\{1\}$
$\{2\}$	$\{\}$	$\{\}$	$\{2\}$	$\{2\}$	$\{2\}$
$\{1,2\}$	$\{\}$	$\{1\}$	$\{2\}$	$\{1,2\}$	$\{1,2\}$
$\{1,2,3\}$	$\{\}$	$\{1\}$	$\{2\}$	$\{1,2\}$	$\{1,2,3\}$

https://en.wikipedia.org/wiki/Topological_space

Every union of (f)

(f) is not a topology $\{\{\}, \{1,2\}, \{2,3\}, \{1,2,3\}\}$
every union of (f)

\cup	$\{\}$	$\{1,2\}$	$\{2,3\}$	$\{1,2,3\}$
$\{\}$	$\{\}$	$\{1,2\}$	$\{2,3\}$	$\{1,2,3\}$
$\{1,2\}$	$\{1,2\}$	$\{1,2\}$	$\{1,2,3\}$	$\{1,2,3\}$
$\{2,3\}$	$\{2,3\}$	$\{1,2,3\}$	$\{2,3\}$	$\{1,2,3\}$
$\{1,2,3\}$	$\{1,2,3\}$	$\{1,2,3\}$	$\{1,2,3\}$	$\{1,2,3\}$

https://en.wikipedia.org/wiki/Topological_space

Every intersection of (f)

(f) is not a topology $\{\{\}, \{1,2\}, \{2,3\}, \{1,2,3\}\}$
every intersection of (f)

\cap	$\{\}$	$\{1,2\}$	$\{2,3\}$	$\{1,2,3\}$
$\{\}$	$\{\}$	$\{\}$	$\{\}$	$\{\}$
$\{1,2\}$	$\{\}$	$\{1,2\}$	$\{2\}$	$\{1,2\}$
$\{2,3\}$	$\{\}$	$\{2\}$	$\{2,3\}$	$\{2,3\}$
$\{1,2,3\}$	$\{\}$	$\{1,2\}$	$\{2,3\}$	$\{1,2,3\}$

https://en.wikipedia.org/wiki/Topological_space

Examples of topology (3)

- Given $X = \{1, 2, 3, 4\}$,
the *trivial* or *indiscrete topology* on X is
the family $\tau = \{\{\}, \{1, 2, 3, 4\}\} = \{\emptyset, X\}$
consisting of only the two subsets of X
required by the axioms
forms a topology of X .

https://en.wikipedia.org/wiki/Topological_space

Examples of topology (4)

- Given $X = \{1, 2, 3, 4\}$,
the family $\tau = \{\{\}, \{2\}, \{1, 2\}, \{2, 3\}, \{1, 2, 3\}, \{1, 2, 3, 4\}\}$
 $= \{\emptyset, \{2\}, \{1, 2\}, \{2, 3\}, \{1, 2, 3\}, X\}$
of six **subsets** of X forms another **topology** of X .

https://en.wikipedia.org/wiki/Topological_space

Examples of topology (5)

- Given $X = \{1, 2, 3, 4\}$,
the *discrete topology* on X is
the *power set* of X , which is the family $\tau = \wp(X)$
consisting of *all possible subsets* of X .
the family

$$\begin{aligned}\tau = & \{\{\}, \{1\}, \{2\}, \{3\}, \{4\} \\ & \{1, 2\}, \{1, 3\}, \{1, 4\}, \{2, 3\}, \{2, 4\}, \{3, 4\}, \\ & \{1, 2, 3\}, \{1, 2, 4\}, \{1, 3, 4\}, \{2, 3, 4\}, \{1, 2, 3, 4\}\}\end{aligned}$$

- In this case the topological space (X, τ)
is called a *discrete space*.

https://en.wikipedia.org/wiki/Topological_space

Examples of topology (6)

- Given $X = \mathbb{Z}$, the set of integers, the family τ of all finite subsets of the integers plus \mathbb{Z} itself is not a topology, because (for example) the union of all finite sets not containing zero is not finite but is also not all of \mathbb{Z} , and so it cannot be in τ .

https://en.wikipedia.org/wiki/Topological_space

Definition via Open Sets (1)

- A **topology** τ on a **set** X is a **set** of **subsets** of X with the *properties* below.
 - a **topology** τ on a **set** X : a **set** of **subsets** of X
 - **members** of τ : **subsets** of X
- each **member** of τ is called an **open set**.
- X together with τ is called a **topological space**

https://en.wikipedia.org/wiki/Open_set

Definition via Open Sets (2)

- topology τ : a set of subsets of X has the *properties* below
 - $X \in \tau$ and $\emptyset \in \tau$
 - any union of sets in τ belong to τ :
any union of subsets of X belong to τ :
if $\{U_i : i \in I\} \subseteq \tau$ then

$$\bigcup_{i \in I} U_i \in \tau$$

- any finite intersection of sets in τ belong to τ
any finite intersection of subsets of X belong to τ :
if $U_1, \dots, U_n \in \tau$ then

$$U_1 \cap \dots \cap U_n \in \tau$$

https://en.wikipedia.org/wiki/Open_set

Definition via Open Sets (3)

- Infinite intersections of open sets need not be open.
- For example, the intersection of all intervals of the form $(-1/n, 1/n)$, where n is a positive integer, is the set $\{0\}$ which is not open in the real line.
- A metric space is a topological space, whose topology consists of the collection of all subsets that are unions of open balls.
- There are, however, topological spaces that are not metric spaces.

https://en.wikipedia.org/wiki/Open_set

Definition via Open Sets (4)

- A **topology** on a set X may be defined as a **collection** τ of **subsets** of X , called **open sets** and satisfying the following **axioms**:
 - The **empty set** and X itself belong to τ .
 - any arbitrary (**finite** or **infinite**) **union** of members of τ belongs to τ .
 - the **intersection** of any **finite** number of members of τ belongs to τ .

https://en.wikipedia.org/wiki/Topological_space

Definition via Open Sets (5)

- As this definition of a topology is the most commonly used, the set τ of the **open sets** is commonly called a **topology** on X .
- A **subset** $C \subseteq X$ is said to be **closed** in (X, τ) if its complement $X \setminus C$ is an **open set**.

https://en.wikipedia.org/wiki/Topological_space

Definition via Neighborhoods (1)

- This axiomatization is due to Felix Hausdorff.
- Let X be a **set**;
- the **elements** of X are usually called **points**, though they can be any mathematical object.
- We allow X to be **empty**.

https://en.wikipedia.org/wiki/Topological_space

Definition via Neighborhoods (2)

- Let \mathcal{N} be a **function** assigning to each x (**point**) in X a non-empty **collection** $\mathcal{N}(x)$ of **subsets** of X .
- The **elements** of $\mathcal{N}(x)$ will be called **neighbourhoods** of x with respect to \mathcal{N} (or, simply, **neighbourhoods** of x).
- The **function** \mathcal{N} is called a neighbourhood topology if *the axioms* below are satisfied; and
- then X with \mathcal{N} is called a **topological space**.

https://en.wikipedia.org/wiki/Topological_space

Definition via Neighborhoods (3)

- If N is a neighbourhood of x (i.e., $N \in \mathcal{N}(x)$), then $x \in N$.
In other words, each point belongs to every one of its neighbourhoods.
- If N is a subset of X and includes a neighbourhood of x , then N is a neighbourhood of x . I.e., every superset of a neighbourhood of a point $x \in X$ is again a neighbourhood of x .
- The intersection of two neighbourhoods of x is a neighbourhood of x .
- Any neighbourhood \mathcal{N} of x includes a neighbourhood \mathcal{M} of x such that \mathcal{M} is a neighbourhood of each point of M .

https://en.wikipedia.org/wiki/Topological_space

Definition via Neighborhoods (4)

- The first three axioms for neighbourhoods have a clear meaning.
- The fourth axiom has a very important use in the structure of the theory, that of linking together the neighbourhoods of different points of X .
- A standard example of such a system of neighbourhoods is for the real line \mathbb{R} , where a subset N of \mathbb{R} is defined to be a neighbourhood of a real number x if it includes an open interval containing x .

https://en.wikipedia.org/wiki/Topological_space

Definition via Neighborhoods (5)

- Given such a **structure**, a **subset** U of X is defined to be **open** if U is a **neighbourhood** of all **points** in U .
- The **open sets** then satisfy the **axioms** given below.
- Conversely, when given the **open sets** of a **topological space**, the **neighbourhoods** satisfying the above **axioms** can be recovered by defining N to be a **neighbourhood** of x if N includes an open set U such that $x \in U$.

https://en.wikipedia.org/wiki/Topological_space

Definitions via Closed Sets (1)

- Using **de Morgan's laws**, the above axioms defining **open sets** become axioms defining **closed sets**:
- The **empty set** and X are **closed**.
 - The **intersection** of any **collection** of **closed sets** is also **closed**.
 - The **union** of any finite number of **closed sets** is also **closed**.
- Using these **axioms**, another way to define a **topological space** is as a set X together with a **collection** τ of **closed subsets** of X . Thus the **sets** in the **topology** τ are the **closed sets**, and their complements in X are the **open sets**.

https://en.wikipedia.org/wiki/Open_set

Homeomorphism (1)

- a **homeomorphism**

(from Greek ὁμοιος (homoios) 'similar, same',

and μορφή (morphē) 'shape, form',

named by Henri Poincaré), **topological isomorphism**,

or **bicontinuous function** is

a **bijjective** and **continuous** function

between topological spaces

that has a **continuous inverse** function.

<https://en.wikipedia.org/wiki/Homeomorphism>

Homeomorphism (2)

- **Homeomorphisms** are the **isomorphisms** in the category of **topological spaces** – the **mappings** that **preserve** all the **topological properties** of a given space.
- Two **spaces** with a **homeomorphism** between them are called **homeomorphic**, and from a topological viewpoint they are the same.

<https://en.wikipedia.org/wiki/Homeomorphism>

Homeomorphism (3)

- Very roughly speaking,
a **topological space** is a **geometric object**,
and the **homeomorphism** is
a *continuous* **stretching** and **bending**
of the object into a *new* **shape**.

<https://en.wikipedia.org/wiki/Homeomorphism>

Homeomorphism (4)

- Thus, a *square* and a *circle* are **homeomorphic** to each other, but a *sphere* and a *torus* are not.
- However, this description can be misleading.
- Some *continuous deformations* are not **homeomorphisms**, such as the *deformation* of a *line* into a *point*.
- Some **homeomorphisms** are not *continuous deformations*, such as the homeomorphism between a *trefoil knot* and a *circle*.

<https://en.wikipedia.org/wiki/Homeomorphism>

Outline

- 1 Open Sets and Classes
 - Open Set
 - Filter
 - Class
- 2 Borel Sets
 - Measurable Space
 - Topological Space
 - Borel Sets
- 3 Stochastic Process

Sigma algebra (1)

- We term the **structures** which allow us to use **measure** to be **sigma algebras**
- the only requirements for **sigma algebras** (on a set X) are:
 - the $\{\}$ and X are in the **set**.
 - if A is in the **set**, *complement*(A) is in the **set**.
 - for any **sets** E_i in the set,
 $\bigcup_i E_i$ is in the **set** (for countable i).

<https://medium.com/intuition/measure-theory-for-beginners-an-intuitive-approach-f5cea0cc2e7>

Sigma algebra (2)

- The most intuitive way to think about a **sigma algebra** is that it is the kind of **structure** we can do **probability** on.
 - for example, we can assign ratios of areas and length, so the **measure** on such a **set** X tells something about the **probability** of its **subsets**.
 - we can find the **probability** of **subsets** A and B because we know their ratios with respect to a **set** X ;
 - we also know that
 - (the measure of) their **complements** are defined, and
 - their **unions** and **intersections** are defined,
 - so we know how to find the **probability** of things in this set X .

<https://medium.com/intuition/measure-theory-for-beginners-an-intuitive-approach-f5cea0cc2e7>

Sigma algebra (3)

- The **sigma algebra** which contains the **standard topology** on \mathbb{R} (that is, *all open sets* on \mathbb{R}) is called the **Borel Sigma Algebra**, and the elements of this **set** are called **Borel sets**.
- What this gives us, is the set of **sets** on which outer measure gives our list of dreams. That is, if we take a **Borel set** and we check that length follows translation, additivity, and interval length, it will always hold.

<https://medium.com/intuition/measure-theory-for-beginners-an-intuitive-approach-f5cea0cc2e7>

Sigma algebra (4)

- The **set** of Lebesgue measurable sets is the **set** of **Borel sets**, along with (union) all the sets which differ from a Borel set by a **set of measure 0**.
- More intuitively, it is all the sets we can normally measure, plus a bunch of stuff that doesn't affect our ideas of area or volume (think about the **border** of the circle above).

<https://medium.com/intuition/measure-theory-for-beginners-an-intuitive-approach-f5cea0cc2e7>

Borel Sets (1-1)

- a **Borel set** is any **set** in a **topological space** that can be formed from **open sets** (or, equivalently, from **closed sets**) through the operations of
 - countable union,
 - countable intersection, and
 - relative complement.

https://en.wikipedia.org/wiki/Borel_set

Borel Sets (1-2)

- For a **topological space X** , the collection of all Borel sets on X forms a σ -algebra, known as the **Borel algebra** or **Borel σ -algebra**.
- The **Borel algebra on X** is the smallest **σ -algebra** containing all open sets (or, equivalently, all closed sets).

https://en.wikipedia.org/wiki/Borel_set

Borel Sets (1-3)

- **Borel sets** are important in measure theory, since any measure defined on the open sets of a space, or on the closed sets of a space, must also be defined on all Borel sets of that space.
- Any measure defined on the Borel sets is called a **Borel measure**.
- **Borel sets** and the associated **Borel hierarchy** also play a fundamental role in descriptive set theory.

https://en.wikipedia.org/wiki/Borel_set

Borel Sets (2)

- **Borel sets** are those obtained from intervals by means of the operations allowed in a **σ -algebra**. So we may construct them in a (transfinite) "sequence" of steps:
- ... And again and again.

<https://math.stackexchange.com/questions/220248/understanding-borel-sets>

Borel Sets (3-1)

1. Start with **finite unions** of **closed-open intervals**.
These sets are completely **elementary**, and they form an **algebra**.
2. **Adjoin countable unions** and **intersections** of elementary sets.
What you get already includes **open sets** and **closed sets**, **intersections** of an open set and a closed set, and so on.
Thus you obtain an **algebra**, that is still not a **σ -algebra**.

<https://math.stackexchange.com/questions/220248/understanding-borel-sets>

Borel Sets (3)

3. Again, **adjoin countable unions** and **intersections** to 2.
Observe that you get a strictly larger class, since a **countable intersection** of **countable unions** of intervals is not necessarily included in 2.
Explicit examples of sets in 3 but not in 2 include F_σ sets, like, say, the set of *rational numbers*.
4. And do the same again.

<https://math.stackexchange.com/questions/220248/understanding-borel-sets>

Borel Sets (4-1)

- And even after a sequence of steps we are not yet finished. Take, say, a countable union of a set constructed at step 1, a set constructed at step 2, and so on. This union may very well not have been constructed at any step yet. By axioms of σ -algebra, you should include it as well - if you want, as step ∞

<https://math.stackexchange.com/questions/220248/understanding-borel-sets>

Borel Sets (4-2)

- (or, technically, the first infinite ordinal, if you know what that means).
- And then continue in the same way until you reach the first uncountable ordinal. And only then will you finally obtain the generated σ -algebra.

<https://math.stackexchange.com/questions/220248/understanding-borel-sets>

Stochastic Process (1)

In probability theory and related fields, a **stochastic** (/stou'kæstɪk/) or **random** process is a mathematical object usually defined as a family of **random variables**.

The word stochastic in English was originally used as an adjective with the definition "pertaining to **conjecturing**", and stemming from a Greek word meaning "to aim at a mark, guess", and the Oxford English Dictionary gives the year 1662 as its earliest occurrence.

From Ancient Greek στοχαστικός (stokhastikós), from στοχάζομαι (stokhá-zomai, "aim at a target, guess"), from στόχος (stókhos, "an aim, a guess").

<https://en.wikipedia.org/wiki/Stochastic>
<https://en.wiktionary.org/wiki/stochastic>

Stochastic Process (2)

The definition of a **stochastic process** varies, but a **stochastic process** is *traditionally* defined as a collection of **random variables** indexed by some set.

The terms **random process** and **stochastic process** are considered synonyms and are used interchangeably, without the **index set** being precisely specified.

Both "**collection**", or "**family**" are used while instead of "**index set**", sometimes the terms "**parameter set**" or "**parameter space**" are used.

https://en.wikipedia.org/wiki/Stochastic_process

Stochastic Process (3)

The term **random function** is also used to refer to a **stochastic** or **random process**, though sometimes it is only used when the stochastic process takes real values.

This term is also used when the **index sets** are **mathematical spaces** other than the **real line**,

while the terms **stochastic process** and **random process** are usually used when the **index set** is interpreted as time,

and other terms are used such as **random field** when the **index set** is n -dimensional **Euclidean space** \mathbb{R}^n or a manifold

https://en.wikipedia.org/wiki/Stochastic_process

Stochastic Process (4)

A **stochastic process** can be denoted, by $\{X(t)\}_{t \in T}$, $\{X_t\}_{t \in T}$, $\{X(t)\}$, $\{X_t\}$ or simply as X or $X(t)$, although $X(t)$ is regarded as an abuse of function notation.

For example, $X(t)$ or X_t are used to refer to the **random variable** with the **index** t , and not the entire **stochastic process**.

If the **index set** is $T = [0, \infty)$, then one can write, for example, $(X_t, t \geq 0)$ to denote the **stochastic process**.

https://en.wikipedia.org/wiki/Stochastic_process

Stochastic Process Definition (1)

A **stochastic process** is defined as a collection of **random variables** defined on a common **probability space** (Ω, \mathcal{F}, P) ,

- Ω is a **sample space**,
- \mathcal{F} is a σ -**algebra**,
- P is a **probability measure**;
- the **random variables**, indexed by some set T ,
- all take values in the same **mathematical space** S , which must be **measurable** with respect to some σ -algebra Σ

https://en.wikipedia.org/wiki/Stochastic_process

Stochastic Process Definition (2)

In other words, for a given **probability space** (Ω, \mathcal{F}, P) and a **measurable space** (S, Σ) , a **stochastic process** is a **collection** of S -valued **random variables**, which can be written as:

$$\{X(t) : t \in T\}.$$

https://en.wikipedia.org/wiki/Stochastic_process

Stochastic Process Definition (3)

Historically, in many problems from the natural sciences a point $t \in T$ had the meaning of time, so $X(t)$ is a **random variable** representing a value observed at time t .

A **stochastic process** can also be written as $\{X(t, \omega) : t \in T\}$ to reflect that it is actually a function of two variables, $t \in T$ and $\omega \in \Omega$.

https://en.wikipedia.org/wiki/Stochastic_process

Stochastic Process Definition (4)

There are other ways to consider a stochastic process, with the above definition being considered the traditional one.

For example, a stochastic process can be interpreted or defined as a S^T -valued **random variable**, where S^T is the space of all the possible functions from the set T into the space S .

However this alternative definition as a "**function-valued random variable**" in general requires additional regularity assumptions to be **well-defined**.

https://en.wikipedia.org/wiki/Stochastic_process

Index set (1)

The set T is called the **index set** or **parameter set** of the **stochastic process**.

Often this set is some subset of the real line, such as the natural numbers or an interval, giving the set T the interpretation of time.

https://en.wikipedia.org/wiki/Stochastic_process

Index set (2)

In addition to these sets, the index set T can be another set with a **total order** or a more general set, such as the Cartesian plane R^2 or n -dimensional **Euclidean space**, where an element $t \in T$ can represent a point in space.

That said, many results and theorems are only possible for **stochastic processes** with a **totally ordered index set**.

https://en.wikipedia.org/wiki/Stochastic_process

State space

The **mathematical space** S of a **stochastic process** is called its **state space**.

This mathematical space can be defined using integers, real lines, n -dimensional Euclidean spaces, complex planes, or more abstract mathematical spaces.

The **state space** is defined using elements that reflect the different values that the **stochastic process** can take.

https://en.wikipedia.org/wiki/Stochastic_process

Sample function (1)

A **sample function** is a single outcome of a **stochastic process**, so it is formed by taking a single possible value of each **random variable** of the **stochastic process**.

More precisely, if $\{X(t, \omega) : t \in T\}$ is a **stochastic process**, then for any point $\omega \in \Omega$, the mapping $X(\cdot, \omega) : T \rightarrow S$, is called a **sample function**, a **realization**, or, particularly when T is interpreted as time, a **sample path** of the **stochastic process** $\{X(t, \omega) : t \in T\}$.

https://en.wikipedia.org/wiki/Stochastic_process

Sample function (2)

This means that for a fixed $\omega \in \Omega$,
there exists a **sample function**
that maps the **index set** T to the **state space** S .

Other names for a **sample function** of a **stochastic process**
include **trajectory**, **path function** or **path**

https://en.wikipedia.org/wiki/Stochastic_process

