

Formelsammlung Statistik

Uli Schell

de.wikibooks.org

23. Mai 2014

On the 28th of April 2012 the contents of the English as well as German Wikibooks and Wikipedia projects were licensed under Creative Commons Attribution-ShareAlike 3.0 Unported license. A URI to this license is given in the list of figures on page 57. If this document is a derived work from the contents of one of these projects and the content was still licensed by the project under this license at the time of derivation this document has to be licensed under the same, a similar or a compatible license, as stated in section 4b of the license. The list of contributors is included in chapter Contributors on page 55. The licenses GPL, LGPL and GFDL are included in chapter Licenses on page 63, since this book and/or parts of it may or may not be licensed under one or more of these licenses, and thus require inclusion of these licenses. The licenses of the figures are given in the list of figures on page 57. This PDF was generated by the \LaTeX typesetting software. The \LaTeX source code is included as an attachment (`source.7z.txt`) in this PDF file. To extract the source from the PDF file, you can use the `pdfdetach` tool including in the `poppler` suite, or the <http://www.pdfplabs.com/tools/pdftk-the-pdf-toolkit/> utility. Some PDF viewers may also let you save the attachment to a file. After extracting it from the PDF file you have to rename it to `source.7z`. To uncompress the resulting archive we recommend the use of <http://www.7-zip.org/>. The \LaTeX source itself was generated by a program written by Dirk Hünninger, which is freely available under an open source license from http://de.wikibooks.org/wiki/Benutzer:Dirk_Huenniger/wb2pdf.

Inhaltsverzeichnis

1	Grundlagen	3
1.1	Mengenlehre und DE MORGANSche Regeln	3
1.2	Kombinatorik	3
1.3	Definition der Wahrscheinlichkeit	4
1.4	Additionssatz	4
1.5	BAYES Theorem	5
2	Zufallsvariablen und Verteilungsmodelle	7
2.1	diskrete Zufallsvariablen	7
2.2	diskrete Verteilungsmodelle	9
2.3	stetige Zufallsvariablen	10
2.4	stetige Verteilungsmodelle	11
2.5	Approximation von Verteilungen	15
2.6	Grenzwertsatz	16
3	Deskriptive Statistik	17
3.1	Skalenniveaus	17
3.2	Lageparameter	18
4	Parameterschätzung	23
4.1	Konfidenzintervall für den Erwartungswert μ	23
4.2	Konfidenzintervalle für den Anteilswert einer dichotomen Grundgesamtheit	24
5	Hypothesentests	25
5.1	Vorgehen beim Hypothesentest	25
5.2	Tests auf Lageparameter (Erwartungswert, Median, Anteilswert)	25
5.3	Tests auf Streuung	31
5.4	Tests auf Zusammenhangs- und Assoziationsparameter	32
5.5	Anpassungs- oder Verteilungstests	32
6	Varianzanalyse	35
7	Regressionsrechnung	39
8	Zeitreihenanalyse	41
8.1	Komponentenunterteilung bei Zeitreihen	41
9	Symbolverzeichnis	45
9.1	allgemein	45
9.2	Verteilungsmodelle	45

10 Tabellen	47
11 Autoren	55
Abbildungsverzeichnis	57
12 Licenses	63
12.1 GNU GENERAL PUBLIC LICENSE	63
12.2 GNU Free Documentation License	64
12.3 GNU Lesser General Public License	65

1 Grundlagen

1.1 Mengenlehre und DE MORGANSche Regeln

$$P(\bar{A} \cup \bar{B}) = P(\overline{A \cap B})$$

und

$$P(\bar{A} \cap \bar{B}) = P(\overline{A \cup B})$$

1.2 Kombinatorik

1.2.1 Fakultät

$$n! = 1 \cdot 2 \cdot \dots \cdot (n-1) \cdot n$$

$$0! = 1$$

1.2.2 Binomialkoeffizient

$$\binom{n}{k} = \frac{n!}{k!(n-k)!}, \quad k, n \in \mathbb{N}, \quad k, n \geq 0.$$

$$\binom{n}{0} = 1.$$

1.2.3 Zufallsstichproben

Anzahl der möglichen Stichproben vom Umfang n aus einer Grundgesamtheit vom Umfang N :

	Ohne Zurücklegen	Mit Zurücklegen
Mit	$\frac{N!}{(N-n)!}$	N^n
Ohne Berücksichtigung der Reihenfolge	$\binom{N}{n}$	$\binom{N+n-1}{n}$

1.3 Definition der Wahrscheinlichkeit

(**Symmetrieprinzip** oder **Prinzip nach LAPLACE**) Jedes Ergebnis A aus der Ergebnismenge Ω sei gleich häufig. $|A|$ ist die Zahl der Ergebnisse,

die durch A belegt werden (Anzahl der günstigen Ergebnisse), $|\Omega|$ ist die Zahl aller möglichen Ergebnisse. Es ist

$$P(A) = \frac{|A|}{|\Omega|}.$$

Axiome der Wahrscheinlichkeiten (Kolmogoroff): Gegeben sind zwei Ereignisse $A, B \subset \Omega$.

1. $P(A) \geq 0$. **Nichtnegativität**
2. $P(\Omega) = 1$. **Normiertheit**
3. $P(A \cup B) = P(A) + P(B)$, falls A und B disjunkt sind.

1.4 Additionssatz

Für zwei Ereignisse A, B aus Ω gilt :

$$P(A \cup B) = P(A) + P(B) - P(A \cap B).$$

Für drei Ereignisse A, B, C aus Ω gilt analog :

$$P(A \cup B \cup C) = P(A) + P(B) + P(C) - P(A \cap B) - P(A \cap C) - P(B \cap C) + P(A \cap B \cap C).$$

Falls die Ereignisse **disjunkt** sind, gilt

$$P(A \cup B) = P(A) + P(B).$$

$$P(A \cup B \cup C) = P(A) + P(B) + P(C).$$

1.4.1 Bedingte Wahrscheinlichkeit

$$P(A|B) = \frac{P(A \cap B)}{P(B)}$$

1.4.2 Unabhängigkeit von Ereignissen

Ein Ereignis A ist unabhängig von B , wenn

$$P(A|B) = P(A|\bar{B}) = P(A)$$

1.4.3 Totale Wahrscheinlichkeit

Sei $A_1 \dots A_k$ eine disjunkte Zerlegung von Ω . Dann gilt für $B \subset \Omega$: $P(B) = \sum_{i=1}^k P(B|A_i) \cdot P(A_i)$.

1.5 BAYES Theorem

Für zwei Ereignisse A und B mit $P(B) > 0$ lässt sich die Wahrscheinlichkeit von A unter der Bedingung, dass B eingetreten ist, angeben durch die Wahrscheinlichkeit von B unter der Bedingung, dass A eingetreten ist:

$$P(A | B) = \frac{P(B | A) \cdot P(A)}{P(B)}$$

Hierbei ist

$P(A | B)$ die (bedingte) Wahrscheinlichkeit des Ereignisses A unter der Bedingung, dass B eingetreten ist,

$P(B | A)$ die (bedingte) Wahrscheinlichkeit des Ereignisses B unter der Bedingung, dass A eingetreten ist,

$P(A)$ die A-priori-Wahrscheinlichkeit des Ereignisses A und

$P(B)$ die A-priori-Wahrscheinlichkeit des Ereignisses B .

Endlich viele Ereignisse:

Wenn $A_i, i = 1, \dots, N$ eine Zerlegung der Ergebnismenge in disjunkte Ereignisse ist, gilt für die A-posteriori-Wahrscheinlichkeit $P(A_i | B)$

$$P(A_i | B) = \frac{P(B | A_i) \cdot P(A_i)}{P(B)} = \frac{P(B | A_i) \cdot P(A_i)}{\sum_{j=1}^N P(B | A_j) \cdot P(A_j)}$$

Den letzten Umformungsschritt bezeichnet man auch als Marginalisierung.

Da ein Ereignis A und sein Komplement \bar{A} stets eine Zerlegung der Ergebnismenge darstellen, gilt insbesondere

$$P(A | B) = \frac{P(B | A) \cdot P(A)}{P(B | A) \cdot P(A) + P(B | \bar{A}) \cdot P(\bar{A})}$$

2 Zufallsvariablen und Verteilungsmodelle

2.1 diskrete Zufallsvariablen

Ein Merkmal X , das aufgrund zufälliger Ereignisse eine (endliche) Menge von Ausprägungen $x_1, x_2 \dots$ annehmen kann, nennt man diskrete Zufallsvariable X .

2.1.1 Eindimensionale Zufallsvariablen

Wahrscheinlichkeitsfunktion:

$$f(x) = \begin{cases} P(X = x_i) = p_i, & x = x_i \in \{x_1, x_2, \dots, x_k\} \\ 0 & \text{sonst} \end{cases}$$

Verteilungsfunktion:

$$F(x) = P(X \leq x) = \sum_{i: x_i \leq x} f(x_i).$$

Normiertheit:

$$\sum_{i=1}^k p_i = 1.$$

Erwartungswert

$$E(X) = \mu = \sum_{i=1}^k x_i \cdot p_i = \sum_{i=1}^k x_i \cdot f(x_i),$$

Varianz

$$Var(X) = \sum_{i=1}^k (x_i - E(X))^2 \cdot f(x_i).$$

bzw. mit dem Verschiebungssatz

$$\text{Var}(X) = \left(\sum_{i=1}^k x_i^2 \cdot f(x_i) \right) - E(X^2) =$$

Standardabweichung

$$\sigma = +\sqrt{\text{Var}(X)}.$$

Varianz der Summe unabhängiger Zufallsvariablen

$$\text{Var}(X + Y) = \text{Var}(X) + \text{Var}(Y).$$

2.1.2 Mehrdimensionale Zufallsvariablen

Einzelwahrscheinlichkeit

$$P(X = x_1) = f_X(x_1) = \sum_{j=1}^m f_{X,Y}(x_1; y_j)$$

Kovarianz

$$\text{cov}XY = \sum_{i=1}^n \sum_{j=1}^m (x_i - E(X))(y_j - E(Y)) f_{X,Y}(x_i; y_j)$$

bzw. mit dem Verschiebungssatz

$$\text{cov}XY = \sum_{i=1}^n \sum_{j=1}^m x_i \cdot y_j \cdot f_{X,Y}(x_i; y_j) - E(X) \cdot E(Y)$$

Korrelationskoeffizient r_{xy} nach Bravais-Pearson

für metrisch skalierte Merkmale zweier statistischer Variablen x und y

$$r = \frac{\sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y})}{\sqrt{\sum_{i=1}^n (x_i - \bar{x})^2} \cdot \sqrt{\sum_{i=1}^n (y_i - \bar{y})^2}},$$

mit $\bar{x} = \frac{1}{n} \cdot \sum_{i=1}^n x_i$ als dem arithmetischen Mittel des Merkmals x. Mit Hilfe des Verschiebungssatzes:

$$r = \frac{\sum_{i=1}^n x_i \cdot y_i - n \cdot \bar{x} \cdot \bar{y}}{\sqrt{(\sum_{i=1}^n x_i^2 - n \cdot (\bar{x})^2) \cdot (\sum_{i=1}^n y_i^2 - n \cdot (\bar{y})^2)}}$$

Rangkorrelationskoeffizient nach Spearman

- für Variablen, die stark von der Normalverteilung abweichen
- sowie ordinalskalierte Variablen

Nach Ordnung der einzelnen Beobachtungen von x bzw. y der Größe nach wird jedem Wert wird seine Rangzahl $rg(x_i)$ und $rg(y_i)$ zugewiesen. Damit:

$$r_{SP} = \frac{\sum_i (rg(x_i) - \overline{rg(x)})(rg(y_i) - \overline{rg(y)})}{\sqrt{\sum_i (rg(x_i) - \overline{rg(x)})^2} \sqrt{\sum_i (rg(y_i) - \overline{rg(y)})^2}}$$

2.2 diskrete Verteilungsmodelle

2.2.1 Binomialverteilung

Für eine binomialverteilte Zufallsvariable X mit den Parametern n und θ ($0 \leq \theta \leq 1$) lautet die Wahrscheinlichkeitsfunktion

$$P(X = x) = b(x|n; \theta) = \begin{cases} \binom{n}{x} \theta^x (1 - \theta)^{n-x} & \text{falls } x = 0, 1, \dots, n \\ 0 & \text{sonst.} \end{cases}$$

Erwartungswert

$$E(X) = n \cdot \theta$$

Varianz

$$Var(X) = n \cdot \theta \cdot (1 - \theta)$$

2.2.2 Hypergeometrische Verteilung

Eine Zufallsvariable X ist **hypergeometrisch** verteilt mit den Parametern N (Grundgesamtheit), M ("Kugeln der ersten Sorte") und n (Stichprobenumfang), wenn ihre Wahrscheinlichkeitsfunktion lautet

$$P(X = x) = h(x|N; M; n) = \begin{cases} \frac{\binom{M}{x} \cdot \binom{N-M}{n-x}}{\binom{N}{n}} & \text{für } x = 0, 1, \dots, n \\ 0 & \text{sonst} \end{cases}$$

Erwartungswert

$$E(X) = n \cdot \frac{M}{N} = n \cdot \Theta$$

Varianz

$$Var(X) = n \cdot \frac{M}{N} \cdot \left(1 - \frac{M}{N}\right) \frac{N-n}{N-1}.$$

Der Bruch $\frac{N-n}{N-1}$ wird *Korrekturfaktor* genannt.

2.2.3 Poissonverteilung**Wahrscheinlichkeitsfunktion ($\lambda > 0$)**

$$P(X = x) = p(x|\lambda) = \begin{cases} \frac{e^{-\lambda} \cdot \lambda^x}{x!} & \text{für } x = 0, 1, \dots \\ 0 & \text{sonst} \end{cases}$$

Erwartungswert und Varianz

$$E(X) = Var(X) = \lambda$$

2.3 stetige Zufallsvariablen

Eine stetige Zufallsvariable kann in jedem beschränkten Intervall unendlich viele Ausprägungen annehmen.

Ihre Verteilung lässt sich durch eine Dichtefunktion $f(x)$ beschreiben.

($f(x)$ ist hier keine Wahrscheinlichkeit, sondern eine Dichte !)

Verteilungsfunktion

$$P(X \leq a) = F(a) = \int_{-\infty}^a f(x) dx$$

- Es gilt: $P(X = a) = 0$.
- Wegen $P(X = a) = 0$ ist $P(X \leq a) = P(X < a)$ und $P(X > a) = P(X \geq a)$

Die **Dichtefunktion $f(x)$** ist die erste Ableitung der Verteilungsfunktion, falls diese an der Stelle x differenzierbar ist.

- Die Dichtefunktion $f(a)$ kann auch größer als 1 werden.
- Ausgehend von $P(X \leq x) = p$ ist das p -Quantil $x(p)$ der Wert x , der zu einer gegebenen Wahrscheinlichkeit p gehört. Speziell $x(0,5)$ ist der Median.

Erwartungswert

$$E(X) = \int_{-\infty}^{\infty} x \cdot f(x) dx,$$

falls $E(X)$ existiert, d.h. nicht unendlich wird.

Varianz

$$Var(X) = \int_{-\infty}^{\infty} (x - E(X))^2 \cdot f(x) dx$$

wobei auch hier der Verschiebungssatz angewendet werden kann:

$$Var(X) = \left(\int_{-\infty}^{\infty} x^2 f(x) dx \right) - (E(X))^2$$

2.4 stetige Verteilungsmodelle

2.4.1 Stetige Gleichverteilung (Rechteckverteilung)

Dichtefunktion der Gleichverteilung im Intervall $[a, b]$

$$f(x) = \begin{cases} \frac{1}{b-a} & \text{für } a \leq x \leq b \\ 0 & \text{sonst.} \end{cases}$$

Erwartungswert

$$E(X) = \frac{a+b}{2}$$

Varianz

$$Var(X) = \lambda \int_a^b x^2 \cdot \frac{1}{b-a} dx = \frac{(b-a)^2}{12}$$

2.4.2 Exponentialverteilung

Dichtefunktion der Exponentialverteilung

$$f(x) = \begin{cases} \lambda \cdot e^{-\lambda x} & \text{für } x \geq 0 \\ 0 & \text{für } x < 0 \end{cases}$$

Erwartungswert

$$E(X) = \lambda \int_0^{\infty} x \cdot e^{-\lambda x} dx = \frac{1}{\lambda}$$

Varianz

$$Var(X) = \frac{1}{\lambda^2}$$

2.4.3 Normalverteilung

Für eine Zufallsvariable $X \propto N(\mu, \sigma^2)$ lautet die **Dichtefunktion der NV**

$$f(x) = \frac{1}{\sqrt{2\pi} \cdot \sigma} \cdot e^{-\frac{(x-\mu)^2}{2\sigma^2}} \text{ für } x \in \mathbb{R}$$

Normierung mit $z = \frac{x-\mu}{\sigma}$ ergibt die **Standardnormalverteilung** mit der Dichtefunktion $\phi_x(z) \propto N(0, 1)$:

$$\phi_x(z) = \frac{1}{\sqrt{2 \cdot \pi}} \cdot e^{-\frac{1}{2}z^2}$$

Anm.: Es wird auch die Schreibweise $\phi_x(z|\mu, \sigma^2)$ anstelle $N(\mu, \sigma^2)$ verwendet

Erwartungswert

$$E(X) = \mu$$

Varianz

$$Var(X) = \sigma^2$$

p-Quantil

Der zu einer gegebenen Wahrscheinlichkeit p zugehörige z -Wert $z(p)$

$$P(Z \leq z(p)) = p$$




Abb. 1 97,5%-Quantil der Standardnormalverteilung

Beispielsweise ist $z(0,975) = 1,96$.

Linearkombinationen normalverteilter Zufallsvariablen

Für n normalverteilte Zufallsvariablen $X_i (i = 1, \dots, n)$, mit $X_i \propto N(\mu_i; \sigma_i^2)$ ist die Linearkombination

$$Y = a_0 + a_1 X_1 + a_2 X_2 + \dots + a_n X_n = a_0 + \sum_{i=1}^n a_i X_i$$

ebenfalls normalverteilt mit dem Erwartungswert

$$E(Y) = a_0 + \sum_{i=1}^n a_i E(X_i) = a_0 + \sum_{i=1}^n a_i \mu_i$$

Falls die $X_i (i = 1, \dots, n)$ stochastisch unabhängig sind, gilt für die Varianz

$$Var(Y) = \sum_{i=1}^n a_i^2 \cdot (X_i) = \sum_{i=1}^n a_i^2 \sigma_i^2$$

Die Varianz muss größer Null sein, deshalb muss zudem $a_j \neq 0$ für mindestens ein $j \in \{1, \dots, n\}$ gelten.

Verteilung des Stichprobendurchschnitts

Sind die n Zufallsvariablen $X_i (i = 1, \dots, n)$ sämtlich normalverteilt mit gleichem μ und gleichem σ^2 , ist die Linearkombination

$\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i$ mit $a_0 = 0, a_1 = a_2 = \dots = a_n = 1/n$, also $\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i$

normalverteilt dem **Erwartungswert**

$$E(\bar{X}) = \frac{1}{n} \sum_{i=1}^n \mu = \mu$$

und, falls die X_i ($i = 1, \dots, n$) stochastisch unabhängig sind, mit der **Varianz**

$$Var(\bar{X}) = \frac{1}{n^2} \sum_{i=1}^n \sigma^2 = \frac{\sigma^2}{n}$$

2.4.4 CHI-Quadrat-verteilung

Die X_1, X_2, \dots, X_n seien unabhängige standardnormalverteilte Zufallsvariablen.

Dann ist die Verteilung der Zufallsvariablen $Z = X_1^2 + X_2^2 + \dots + X_n^2$

chi-quadrat verteilt mit n Freiheitsgraden $Z \propto \chi^2(n)$

Erwartungswert:

$$E(Z) = n$$

Varianz

$$Var(Z) = 2n$$

Anm.: Die Gruppe der Hypothesentests mit χ^2 -Verteilung bezeichnet man als χ^2 -**Test**.

Hierunter sind mehrere Tests zu verstehen:

Verteilungstest oder Anpassungstest: Hier wird geprüft, ob vorliegende Daten auf eine bestimmte Weise verteilt sind.

Unabhängigkeitstest: Hier wird geprüft, ob zwei Merkmale stochastisch unabhängig sind.

Homogenitätstest: Hier wird geprüft, ob zwei oder mehr Stichproben derselben Verteilung bzw. einer homogenen Grundgesamtheit entstammen.

2.4.5 t- (Student-) Verteilung

Für die unabhängigen Variablen X (standardnormalverteilt) und Z ($Z \propto \chi^2(n)$) ist die Variable

$$T = \frac{X}{\sqrt{Z/n}}$$

t-verteilt ($T \propto t(n)$) mit n Freiheitsgraden.

Erwartungswert

$$E(T) = 0 \text{ für } (m \geq 2)$$

Varianz

$$Var(T) = \frac{n}{n-2} \text{ für } (n \geq 3)$$

2.4.6 Fisher- Verteilung

Für die unabhängigen Variablen $X \propto \chi^2(m)$ und $Y \propto \chi^2(n)$ ist die Verteilung der Variablen

$$Z = \frac{X/m}{Y/n}$$

Fisher- oder F-verteilt ($Z \propto F(m,n)$) mit den Freiheitsgraden m und n.

Erwartungswert

$$E(T) = \frac{n}{n-2} \text{ für } (n \geq 3)$$

Varianz

$$Var(Z) = \frac{2n^2(n+m-2)}{m(n-4)(n-2)^2} \text{ für } (n \geq 3)$$

2.5 Approximation von Verteilungen

Gesuchte Verteilung	Approximation durch		
	Binomial	Poisson	Normal
$P(X \leq x)$ Binomial $B(x n\theta) \approx$	---	$P(x n\theta)$ falls $n \geq 50$ und $\theta \leq 0,05$	$\Phi(x + 0,5 n \cdot \theta; n \cdot \theta \cdot (1 - \theta))$ falls $n > \frac{9}{\theta(1-\theta)}$
Hypergeometrische $H(x N; M; n) \approx$	$B(x n \frac{M}{N})$ falls $\frac{n}{N} < 0,05$	über Binomialverteilung	$\Phi(x + 0,5 n \cdot \frac{M}{N}; n \cdot \frac{M}{N} \cdot (1 - \frac{M}{N}) \cdot \frac{N-n}{N-1})$ falls $n > \frac{9}{\frac{M}{N} \cdot (1 - \frac{M}{N})}$ und $\frac{n}{N} < 0,05$
Poisson $P(x \lambda) \approx$	---	---	$\Phi(x + 0,5 \lambda; \lambda)$ falls $\lambda > 9$
χ^2 -Verteilung $\chi^2(x n) \rightarrow$ $P(\sqrt{2X} \leq \sqrt{2x}) \approx$	---	---	$\Phi(\sqrt{2x} \sqrt{2n-1}; 1)$ falls $n > 30$
t-Verteilung $t(x n) \approx$	---	---	$\Phi(x 0; 1)$ falls $n > 30$
F-Verteilung $F(x m; n) \approx$	---	---	$\Phi(x 0; 1)$ falls $m > 30$ und $n > 30$

2.6 Grenzwertsatz

2.6.1 Gesetz der großen Zahlen

Für ein beliebig kleines $c > 0$ gilt

$$P(|\bar{X}_n - \mu| \leq c) \rightarrow 1 \text{ für } (n \rightarrow \infty)$$

2.6.2 Theorem von Bernoulli

Die relative Häufigkeit, mit der ein Ereignis A bei n unabhängigen Wiederholungen eines Zufallsereignisses eintritt, konvergiert nach Wahrscheinlichkeit gegen $P(A)$

2.6.3 Hauptsatz der Statistik

Für eine Zufallsvariable X mit der Verteilungsfunktion $F(x)$ gilt für die Verteilungsfunktion $F_n(x)$

für die unabhängigen wie identisch wie X verteilten $X_1 \dots X_n$ ($x \in \mathbb{R}$)

$$P(\sup |F_n(x) - F(x)| \leq c) \rightarrow 1 \text{ für } (n \rightarrow \infty)$$

(sup: Maximale Abweichung zwischen $\hat{F}_n(x)$ und $\hat{F}(x)$).

2.6.4 Zentraler Grenzwertsatz

Für unabhängig identisch verteilte Zufallsvariablen $X_1 \dots X_n$ mit $E(X_i) = \mu$

und $\text{Var}(X_i) = \sigma^2 > 0$ konvergiert die Verteilungsfunktion $F_n(z) = P(Z_n \leq z)$

der standardisierten Summe

$$Z_n = \frac{X_1 + \dots + X_n - n \cdot \mu}{\sqrt{n} \sigma} = \frac{1}{\sqrt{n}} \sum_{i=1}^n \frac{X_i - \mu}{\sigma}$$

für $n \rightarrow \infty$ an jeder Stelle $z \in \mathbb{R}$ gegen die Verteilungsfunktion $\phi_x(z)$ der Standardnormalverteilung

$$F_n(z) \Rightarrow \phi_x(z)$$

2.6.5 Grenzwertsatz von De Moivre

Die Verteilung der standardisierten absoluten Häufigkeit $\frac{H_n - n \cdot \pi}{\sqrt{n \pi (1 - \pi)}}$ der Standardnormalverteilung

konvergiert für $n \rightarrow \infty$ gegen eine Standardnormalverteilung.

3 Deskriptive Statistik

3.1 Skalenniveaus

3.1.1 Nominalskala

Die Ausprägungen des nominalskalierten Merkmals können nicht **geordnet** werden, man kann sie nur **vergleichen** und **abzählen**.

Es handelt sich um **qualitative** Merkmale. Erhalten die Ausprägungen Ziffern zugeordnet, handelt es sich nur um eine **Verschlüsselung** (Codierung): 1 = männlich, 2 = weiblich.

3.1.2 Ordinalskala

Zwischen den Ausprägungen des ordinalskalierten (rangskalierten) Merkmals existiert eine Beziehung

der Form mehr oder weniger, < oder >, besser oder schlechter.

Eine Quotientenbildung macht wenig Sinn (Beispiel Noten: 1, 2, 3, 4, 5).

3.1.3 Intervallskala

Die Abstände zwischen den Ausprägungen des (quantitativen) Merkmals der Intervallskala können gemessen werden. Es handelt sich bei den Ausprägungen um (reelle) Zahlen.

Beispiel: Kinderzahl, Temperatur.

3.1.4 Verhältnisskala

Sowohl die Abstände als auch Verhältnisse zwischen den Ausprägungen des (quantitativen) Merkmals

können gemessen werden. Es handelt sich bei den Ausprägungen um (reelle) Zahlen. Beispiel: Einkommen.

3.1.5 Zweig-Blätter-(stem-leaf-) Diagramm

Die linke Spalte enthält als „Stämme“ die Äquivalenzklassen, in die die auf der rechten Seite als „Blätter“

dargestellten Merkmale eingeteilt werden. Beispiel: Gegeben sind die Werte 0,3 0,4 2,5 2,5 2,6 2,7 2,8 3,5 3,7.

Wählt man die natürlichen Zahlen als Klasseneinteilung, ergibt sich folgendes Stamm-Blatt-Diagramm:

3	5	7			
2	5	5	6	7	8
1					
0	3	4			

3.2 Lageparameter

3.2.1 Arithmetisches Mittel

$$\bar{X} = \frac{1}{n} \sum_{i=1}^n x_i$$

3.2.2 Median (Zentralwert)

Sind die Beobachtungswerte der Größe nach geordnet, ist der Median z die Stelle, die die Teilgesamtheit in zwei gleiche Hälften teilt.

$$z = \begin{cases} x_{[\frac{n+1}{2}]} & \text{n ungerade} \\ \frac{1}{2}(x_{[\frac{n}{2}]} + x_{[\frac{n}{2}+1]}) & \text{n gerade} \end{cases}$$

3.2.3 geometrisches Mittel

$$\bar{X}_{geom} = \sqrt[n]{\prod_{i=1}^n x_i}$$

3.2.4 harmonisches Mittel

$$\bar{X}_{harm} = \frac{n}{\sum_{i=1}^n \frac{1}{x_i}}$$

3.2.5 Modalwert

Der am häufigsten auftretende Wert

3.2.6 Varianz

Grundgesamtheit:

$$\sigma^2 = \frac{1}{n} \sum_{i=1}^n (x_i - \mu)^2$$

Stichprobe:

$$\bar{s}^2 = \frac{1}{n-1} \sum_{i=1}^n (x_i - \bar{X})^2$$

3.2.7 Verschiebungssatz

Für jedes $c \in \mathbb{R}$ gilt

$$\sum_{i=1}^n (x_i - c)^2 = \sum_{i=1}^n (x_i - \bar{X})^2 + n(\bar{X} - c)^2$$

Damit erhält man als Varianz

$$s^2 = \frac{1}{n} \left(\sum_{i=1}^n x_i^2 - n \cdot \bar{X}^2 \right)$$

3.2.8 Variationskoeffizient

$$v = \frac{\bar{s}}{\bar{X}}, \bar{X} > 0$$

3.2.9 Konzentrationsmasse

Konzentrationsrate

Die Konzentrationsrate¹ CR_n ist die Summe der Marktanteile der n größten Unternehmen eines relevanten Marktes. Im GWB² werden die Raten CR₁, CR₃ und CR₅ herangezogen.

¹ <http://de.wikipedia.org/wiki/Konzentrationsrate>

² http://de.wikipedia.org/wiki/Gesetz_gegen_Wettbewerbsbeschr%25C3%25A4nkungen

Lorenzkurve

Für eine geordnete Urliste $x_1 \leq x_2 \dots \leq x_n$ trägt man die kumulierte relative Merkmalssumme

$$q_i = \frac{\sum_{k=1}^j x_k}{\sum_{k=1}^n x_k}$$

über den Anteil der Merkmalsträger $p_j = \frac{j}{n}$ auf.

Liegen die Merkmale in gruppierter Form vor, trägt man die kumulierte relative Merkmalssumme

$$q_i = \frac{\sum_{k=1}^j x_k}{\sum_{k=1}^n x_k}$$

über der Häufigkeit $p_i = \frac{1}{n} \cdot \sum_{j=1}^i h_j$ auf.

Zwischen (0;0) und (1;1) wird die Winkelhalbierende des Koordinatensystems eingetragen.

Gini-Koeffizient

Als Ginikoeffizient G bezeichnet man das Verhältnis der Fläche zwischen Winkelhalbierender und der Lorenzkurve

zur Gesamtfläche unter der Winkelhalbierenden (= 1/2).

Die Fläche unterhalb der Lorenzkurve kann man einfach aus Teil-Trapezflächen zusammensetzen:

$$G = \left(\frac{1}{2} - \frac{1}{2} \sum_{i=1}^n \Delta p_i \cdot (q_{i-1} + q_i) \right) : \frac{1}{2} = 1 - \sum_{i=1}^n (p_i - p_{i-1}) \cdot (q_{i-1} + q_i)$$

($p_0 = 0$; $q_0 = 0$):




Abb. 2 Ginikoeffizient: Ermittlung einer Trapezfläche für $i=3$




Abb. 3 Ginikoeffizient

Herfindahl-Index

$$H = \sum_{i=1}^n p_i^2, \text{ wobei } p_i = \frac{x_i}{\sum_{j=1}^n x_j}$$

4 Parameterschätzung

4.1 Konfidenzintervall für den Erwartungswert μ

4.1.1 Normalverteiltes Merkmal mit bekannter Varianz

Das Zufallsintervall enthält mit einer Wahrscheinlichkeit $1-\alpha$ den Parameter:

$$P\left(\bar{x} - z_{1-\alpha/2} \cdot \frac{\sigma}{\sqrt{n}} \leq \mu \leq \bar{x} + z_{1-\alpha/2} \cdot \frac{\sigma}{\sqrt{n}}\right) = 1 - \alpha$$

Konfidenzintervall

$$\left[\bar{x} - z\left(1 - \frac{\alpha}{2}\right) \frac{\sigma}{\sqrt{n}}; \bar{x} + z\left(1 - \frac{\alpha}{2}\right) \frac{\sigma}{\sqrt{n}}\right].$$

(Quantil z aus Normalverteilungstabelle)

4.1.2 Normalverteiltes Merkmal mit unbekannter Varianz

Für normalverteilte Merkmale und unbekannter Varianz muss die Varianz durch s^2 geschätzt werden.

$$P\left(\bar{x} - t\left(1 - \frac{\alpha}{2}; n - 1\right) \frac{s}{\sqrt{n}} \leq \mu \leq \bar{x} + t\left(1 - \frac{\alpha}{2}; n - 1\right) \frac{s}{\sqrt{n}}\right) = 1 - \alpha.$$

Konfidenzintervall

$$\left[\bar{x} - t\left(1 - \frac{\alpha}{2}; n - 1\right) \frac{s}{\sqrt{n}}; \bar{x} + t\left(1 - \frac{\alpha}{2}; n - 1\right) \frac{s}{\sqrt{n}}\right].$$

(Quantil $t(1 - \frac{\alpha}{2}; n - 1)$ aus der t-Verteilungstabelle bei Freiheitsgrad $n-1$).

4.1.3 Merkmal mit unbekannter Verteilung und bekannter Varianz

Konfidenzintervall

$$\left[\bar{x} - z\left(1 - \frac{\alpha}{2}\right) \frac{\sigma}{\sqrt{n}}; \bar{x} + z\left(1 - \frac{\alpha}{2}\right) \frac{\sigma}{\sqrt{n}}\right]. \text{ für } n > 30.$$

4.1.4 Merkmal mit unbekannter Verteilung und unbekannter Varianz

Konfidenzintervall

$$\left[\bar{x} - z\left(1 - \frac{\alpha}{2}\right) \frac{s}{\sqrt{n}} ; \bar{x} + z\left(1 - \frac{\alpha}{2}\right) \frac{s}{\sqrt{n}} \right]. \text{ für } n > 50$$

4.2 Konfidenzintervalle für den Anteilswert einer dichotomen Grundgesamtheit

4.2.1 Modell mit Zurücklegen

Beschreibung durch den geschätztem Anteilswert $\hat{p} = \frac{x}{n}$. Für $n > 100$ und $n\hat{p}(1 - \hat{p}) \geq 9$ erhält man das $1 - \alpha$ -Konfidenzintervall für p durch eine Approximation der Binomialverteilung mit Hilfe der Normalverteilung:

$$\left[\hat{p} - z\left(1 - \frac{\alpha}{2}\right) \sqrt{\frac{\hat{p}(1 - \hat{p})}{n}} ; \hat{p} + z\left(1 - \frac{\alpha}{2}\right) \sqrt{\frac{\hat{p}(1 - \hat{p})}{n}} \right].$$

4.2.2 Modell ohne Zurücklegen

Für $n > \frac{9}{p(1-p)}$, $n > 100n/N \leq 0,05$ kann die hypergeometrische Verteilung durch die Normalverteilung approximiert werden:

$(1 - \alpha)$ -Konfidenzintervall für θ :

$$\left[p - z\left(1 - \frac{\alpha}{2}\right) \sqrt{\frac{p(1-p)}{n}} \sqrt{\frac{N-n}{N-1}} ; p + z\left(1 - \frac{\alpha}{2}\right) \sqrt{\frac{p(1-p)}{n}} \sqrt{\frac{N-n}{N-1}} \right].$$

5 Hypothesentests

5.1 Vorgehen beim Hypothesentest

- I. Feststellung der Verteilung des Merkmals in der Grundgesamtheit
- II. Aufstellen der Nullhypothese
- III. Festlegen der Testfunktion T
- IV. Festlegen des Annahmebereichs ("Nichtablehnungsbereichs") (für ein zu bestimmendes Signifikanzniveau)

Fällt die Prüfgröße \bar{x} in den Bereich $[\bar{x}_u; \bar{x}_o]$,

wird H_0 nicht abgelehnt. Es soll sein

$$P(\bar{x}_u \leq \bar{X} \leq \bar{x}_o) = 1 - \alpha$$

(beachte: ein- oder zweiseitig)

α : Signifikanzniveau oder α -Fehler

V. Stichprobe erheben

VI. Entscheidung treffen

	H₀ ist wirklich wahr	H₁ ist wirklich wahr
H ₀ wird beibehalten	richtige Entscheidung (1- α)	Fehler 2. Art (β -Fehler)
H ₁ wird angenommen	Fehler 1. Art (α -Fehler)	richtige Entscheidung (1- β)

5.2 Tests auf Lageparameter (Erwartungswert, Median, Anteilswert)

5.2.1 Test auf Erwartungswert

Test	H_0	H_1
zweiseitig	$\mu = \mu_0$	$\mu \neq \mu_0$
rechtsseitig	$\mu \leq \mu_0$	$\mu > \mu_0$
linksseitig	$\mu \geq \mu_0$	$\mu < \mu_0$




Abb. 4 Zweiseitiger Test für \bar{x}




Abb. 5 linksseitiger Test für \bar{x}




Abb. 6 Rechtsseitiger Test für \bar{x}

1. **X ist normalverteilt, σ ist bekannt** bei beliebigem n bzw. näherungsweise normalverteilt bei $n > 30$

Testfunktion

$$T = \frac{\bar{X}_n - \mu_0}{\sigma} \cdot \sqrt{n} \sim N(0;1) \text{ (Gauß-Test):}$$

	Ablehnungsbereich
zweiseitig	$ T > z_{1-\alpha/2}$
rechtsseitig	$T > z_{1-\alpha}$
linksseitig	$T < -z_{1-\alpha}$

2. X ist normalverteilt, σ ist unbekannt bei beliebigem n

Testfunktion

$$T = \frac{\bar{X}_n - \mu_0}{S} \cdot \sqrt{n} \sim t(n-1) \text{ (t-Test).}$$

	Ablehnungsbereich
zweiseitig	$ T > t_{1-n, 1-\alpha/2}$
rechtsseitig	$ T > t_{n-1, 1-\alpha}$
linksseitig	$ T < -t_{n-1, 1-\alpha}$

3. X ist näherungsweise normalverteilt, σ ist unbekannt bei n > 30

Testfunktion

$$T = \frac{\bar{X}_n - \mu_0}{S} \cdot \sqrt{n} \approx N(0; 1) \text{ (Gauß-Test) .}$$

	Ablehnungsbereich
zweiseitig	$ T > t_{1-n, 1-\alpha/2}$
rechtsseitig	$ T > t_{n-1, 1-\alpha}$
linksseitig	$ T < -t_{n-1, 1-\alpha}$

5.2.2 Vorzeichentest

Einstichprobenproblem

	Einseitig	Zweiseitig
H_0	$P(X \geq \theta_0) \geq 1/2$	$P(X \geq \theta_0) \leq 1/2$
H_1	$P(X \geq \theta_0) < 1/2$	$P(X \geq \theta_0) > 1/2$
H_0	$\theta \geq \theta_0$	$\theta \leq \theta_0$
H_1	$\theta < \theta_0$	$\theta > \theta_0$

Die Stichprobenwerte, die größer als der hypothetische Median θ_0 sind, bekommen ein "+" zugeordnet;

Werte, die kleiner sind, ein "-". Die Anzahl der positiven Vorzeichen wird gezählt und dient als Teststatistik.

Zweistichprobenproblem

Die n Beobachtungspaare dürfen nicht voneinander abhängen, d.h. das Wertepaar (x_{1i}, x_{2i}) muss unabhängig

vom Wertepaar $(x_{1j}, x_{2j}), \forall i \neq j$ sein.

Besitzen beide Grundgesamtheiten den gleichen Median, gilt $P(X_{11} > X_{12}) = P(X_{11} < X_{12})$.

Folgende Hypothesen können mit dem Vorzeichentest geprüft werden:

	Einseitig		Zweiseitig
H_0	$P(X_1 \geq X_2) \geq 1/2$	$P(X_1 \geq X_2) \leq 1/2$	$P(X_1 \geq X_2) = 1/2$
H_1 :	$P(X_1 \geq X_2) < 1/2$	$P(X_1 \geq X_2) > 1/2$	$P(X_1 \geq X_2) \neq 1/2$

Die Wertepaare der Stichproben, bei denen $x_{i1} > x_{i2}$ gilt, bekommen ein "+" zugeordnet; Wertepaare, für die $x_{i1} < x_{i2}$ gilt, ein "-". Die Anzahl der positiven Vorzeichen wird gezählt und dient als Teststatistik. Die Teststatistik entspricht der Anzahl der positiven Vergleiche (Differenzen der Werte bzw. Ränge):

$$V = \sum_{i=1}^{n'} I(x_{i1} > x_{i2}) \sim B(\pi = 0,5, n')$$

mit

$$I(x_{i1} > x_{i2}) = \begin{cases} 1, & \text{wenn } x_{i1} > x_{i2} \\ 0, & \text{sonst} \end{cases}$$

Für das Einstichprobenproblem sind die Werte der zweiten Stichprobe durch den hypothetischen Median zu ersetzen.

Bei Gültigkeit der Nullhypothese H_0 ist die Summe der positiven Differenzen binomialverteilt mit $\pi = 0,5$,

da der Median dem 50 %-Quantil entspricht. n' bezeichnet den nach Behandlung von Ties (Null Differenzen, Rangbindungen, s.u.)

verbleibenden Stichprobenumfang. Bei Gültigkeit der Nullhypothese ist die Verteilung der Prüfgröße symmetrisch.

Approximation durch die Normalverteilung

Mit $n \rightarrow \infty$ nähert sich die Binomialverteilung einer Normalverteilung mit $N(np, np(1-p))$, als Faustregel $np(1-p) \geq 9$ ($H_0 : p = 1/2$).

Mit $\frac{1}{4}n \geq 9$ bzw. $n \geq 36$ ist die z-standardisierte Größe

$$z_V = \frac{\sum_{i=1}^{n'} - \frac{1}{2} \cdot n'}{\frac{1}{2} \sqrt{n'}} \approx N(0, 1)$$

näherungsweise standardnormalverteilt.

Bindungen (Null Differenzen) Sind im Zweistichprobenproblem die Werte von Beobachtungen von der ersten zur zweiten Stichprobe unverändert

oder im Einstichprobenproblem einige Werte gleich dem Median, ergeben sich Null Differenzen bzw. Bindungen (Ties),

die man so behandeln kann:

- Beobachtungen mit Rangbindungen werden eliminiert, d.h. der Stichprobenumfang wird reduziert.
- Die Beobachtungen werden zu gleichen Teilen den Gruppen zugeordnet. Bei ungerader Anzahl von Bindungen wird ein Beobachtungspaar eliminiert.
- Die Beobachtungen werden jeweils mit einer Wahrscheinlichkeit von 0,5 einer der beiden Gruppen (+ oder -) zugeordnet.

5.2.3 Test auf Anteilswert (Binomialtest)

Der Anteilswert θ wird geschätzt durch

$$\hat{\theta} = p = \frac{x}{n}$$

Mit dem Binomialtest können folgende Hypothesenpaare für θ getestet werden:

Test	H_0	H_1
zweiseitig	$\theta = \theta_0$	$\theta \neq \theta_0$
rechtsseitig	$\theta \leq \theta_0$	$\theta > \theta_0$
linksseitig	$\theta \geq \theta_0$	$\theta < \theta_0$

für $n > 30$, $n\theta_0 \geq 10$ $n(1-\theta_0) \geq 10$

kann man durch die Gauß-Verteilung approximieren:

Testfunktion

$$T = \frac{\hat{\theta} - \theta_0}{\sqrt{\theta_0(1-\theta_0)}} \cdot \sqrt{n} \approx N(0;1) \text{ (Gauß-Test) .}$$

	Ablehnungsbereich
zweiseitig	$ T > z_{1-\alpha/2}$
rechtsseitig	$T > z - 1 - \alpha$
linksseitig	$T < -z - 1 - \alpha$

für $n < 30$ oder $n\theta_0 < 10$ oder $n(1-\theta_0) < 10$

ist der exakte Binomialtest anzuwenden:

Testfunktion

Die Teststatistik X gibt an, wie oft das Merkmal in einer zufälligen Stichprobe vom Umfang n aufgetreten ist.

Unter der Nullhypothese $H_0: \theta = \theta_0$ ist die Teststatistik $B(\theta_0, n)$ -verteilt, das heißt

$$P(X = i) = B(i|\theta_0, n) = \binom{n}{i} \theta_0^i (1 - \theta_0)^{n-i}$$

Ablehnungsbereich




Abb. 7 Teststatistik für den Binomialtest, die roten Balken gehören zum kritischen Bereich.

Da die Teststatistik diskret verteilt ist, kann das vorgegebene Signifikanzniveau α in der Regel nicht eingehalten werden.

Daher wird gefordert, die kritischen Werte so zu wählen, dass für ein möglichst großes *exaktes* Signifikanzniveau α_{ex} gilt $\alpha_{\text{ex}} \leq \alpha$.

Für den zweiseitigen Test werden daher als kritische Werte das größte c_1 und das kleinste c_2 bestimmt, für die gilt

- $\sum_{i=0}^{c_1} B(i|\theta_0, n) \leq \alpha/2$ und
- $\sum_{i=c_2}^n B(i|\theta_0, n) \leq \alpha/2$.

Das exakte Signifikanzniveau ergibt sich als

$$\alpha_{\text{ex}} = \sum_{i=0}^{c_1} B(i|\theta_0, n) + \sum_{i=c_2}^n B(i|\theta_0, n).$$

Für die beiden einseitigen Tests wird analog verfahren.

Test	Kritische Werte	Kritischer Bereich	Grenze(n)
zweiseitig	$c_1 + 1$ und $c_2 - 1$	$\{0, \dots, c_1\} \cup \{c_2, \dots, n\}$	
rechtsseitig	$c - 1$	$\{c, \dots, n\}$	$c =$ kleinster Wert, für den $\sum_{i=c}^n B(i \theta_0, n) = \alpha_{\text{ex}} \leq \alpha$

Test	Kritische Werte	Kritischer Bereich	Grenze(n)
linksseitig	$c + 1$	$\{0, \dots, c\}$	$c = \text{größter Wert, für den } \sum_{i=0}^c B(i \theta_0, n) = \alpha_{\text{ex}} \leq \alpha$

5.3 Tests auf Streuung

5.3.1 Test auf Varianz

Test	H_0	H_1
zweiseitig	$\sigma^2 = \sigma_0^2$	$\sigma^2 \neq \sigma_0^2$
rechtsseitig	$\sigma^2 \leq \sigma_0^2$	$\sigma^2 > \sigma_0^2$
linksseitig	$\sigma^2 \geq \sigma_0^2$	$\sigma^2 < \sigma_0^2$

1. X ist normalverteilt, μ ist unbekannt, n beliebig

Testfunktion

$$T = \frac{(n-1)S^2}{\sigma_0^2} = \frac{1}{\sigma_0^2} \sum_{i=1}^n (X_i - \bar{X})^2 \sim \chi^2(n-1)$$

	Ablehnungsbereich
zweiseitig	$T < \chi_{n-1, \alpha/2}^2$ oder $T > \chi_{n-1, 1-\alpha/2}^2$
rechtsseitig	$T > \chi_{n-1, 1-\alpha}^2$
linksseitig	$T < \chi_{n-1, \alpha}^2$

2. X ist normalverteilt, μ ist bekannt, n beliebig

Testfunktion

$$T = \frac{(n-1)\tilde{S}^2}{\sigma_0^2} = \frac{1}{\sigma_0^2} \sum_{i=1}^n (X_i - \mu)^2 \sim \chi^2(n)$$

	Ablehnungsbereich
zweiseitig	$T < \chi_{n, \alpha/2}^2$ oder $T > \chi_{n, 1-\alpha/2}^2$
rechtsseitig	$T > \chi_{n, 1-\alpha}^2$
linksseitig	$T < \chi_{n, \alpha}^2$

5.4 Tests auf Zusammenhangs- und Assoziationsparameter

5.4.1 Chi-Quadrat-Unabhängigkeitstest

Nullhypothese

H_0 : Die Merkmale X und Y sind stochastisch unabhängig.

Die Beobachtungen der Merkmale X und Y liegen paarweise in m bzw. r Klassen vor.

Es gibt insgesamt n paarweise Beobachtungen von X und Y , die sich auf $m \cdot r$ Kategorien verteilen. Aufstellung z. B. in einer Häufigkeitstabelle:

	Merkmal Y						Summe Σ
Merkmal X	1	2	...	k	...	r	$n_{j\cdot}$
1	n_{11}	n_{12}	...	n_{1k}	...	n_{1r}	$n_{1\cdot}$
2	n_{21}	n_{22}	...	n_{2k}	...	n_{2r}	$n_{2\cdot}$
...
j	n_{jk}	$n_{j\cdot}$
...
m	n_{m1}	n_{m2}	...	n_{mk}	...	n_{mr}	$n_{m\cdot}$
Summe Σ	$n_{\cdot 1}$	$n_{\cdot 2}$...	$n_{\cdot k}$...	$n_{\cdot r}$	n

Absolute Randhäufigkeiten $n_{j\cdot}$ bzw. $n_{\cdot k}$

$$n_{j\cdot} = \sum_{k=1}^r n_{jk} \text{ und } n_{\cdot k} = \sum_{j=1}^m n_{jk}$$

Prüfgröße für den Unabhängigkeitstest:

$$\chi^2 = \sum_{j=1}^m \sum_{k=1}^r \frac{(n_{jk} - n_{jk}^*)^2}{n_{jk}^*}.$$

Mit $n_{jk}^* = \frac{n_{j\cdot} \cdot n_{\cdot k}}{n}$,

H_0 wird abgelehnt, wenn $\chi^2 > \chi^2(1 - \alpha; (m - 1)(r - 1))$ ist.

5.5 Anpassungs- oder Verteilungstests

5.5.1 Chi-Quadrat-Anpassungs- oder Verteilungstest

Die Wahrscheinlichkeiten eines Merkmals X seien in der Grundgesamtheit unbekannt.

Nullhypothese: H_0 : Das Merkmal X besitzt die Wahrscheinlichkeitsverteilung $F_0(x)$

Für n unabhängige Beobachtungen x_1, \dots, x_n des Merkmals X wird die Zahl der Beobachtungen in der j -ten Klasse ist die beobachtete Häufigkeit N_j .

Im Vergleich dazu wird die hypothetische Verteilung bestimmt aufgrund der Wahrscheinlichkeit p_{0j} ,

dass eine Ausprägung von X in die Kategorie j fällt. Die unter H_0 zu erwartende Häufigkeit ist:

$$n_{0j} = p_{0j} \cdot n$$

Die Prüfgröße (Größe der Abweichung)

$$\chi^2 = \sum_{j=1}^m \frac{(N_j - n_{0j})^2}{n_{0j}}$$

ist bei ausreichend großen N_j annähernd chi-Quadrat-verteilt mit $m - 1$ Freiheitsgraden. H_0 wird abgelehnt, wenn $\chi^2 > \chi^2_{(1-\alpha; m-1)}$ gilt.

5.5.2 Kolmogorow-Smirnow-Anpassungstest

Test auf Übereinstimmung zweier Wahrscheinlichkeitsverteilungen.

Man betrachtet ein statistisches Merkmal X , dessen Verteilung in der Grundgesamtheit unbekannt ist.

$H_0 : F_X(x) = F_0(x)$ (Die Zufallsvariable X besitzt die Wahrscheinlichkeitsverteilung F_0 .)

$H_1 : F_X(x) \neq F_0(x)$ (Die Zufallsvariable X besitzt eine andere Wahrscheinlichkeitsverteilung als F_0 .)

Der Kolmogorow-Smirnow-Test vergleicht die empirische Verteilungsfunktion F_n mit F_0 mittels der Teststatistik

$$d_n = \|F_n - F_0\| = \sup_x |F_n(x) - F_0(x)|, \text{ (sup: Supremum)}$$

Die Teststatistik ist unabhängig von der hypothetischen Verteilung F_0 .

Ist der Wert der Teststatistik größer als der entsprechende tabellierte kritische Wert, so wird die Nullhypothese verworfen.

Einstichprobenproblem

Von einer reellen Zufallsvariablen X liegen n aufsteigend sortierte Beobachtungswerte x_i ($i = 1, \dots, n$) vor.

Von diesen Beobachtungen wird die relative Summenhäufigkeit $S(x_i)$ mit der entsprechenden hypothetischen

Verteilung der Grundgesamtheit $F_0(x_i)$ verglichen. Voraussetzung: F_0 ist stetig.

Für jedes $i = 1, \dots, n$ werden die absoluten Differenzen

$$d_{oi} = |S(x_i) - F_0(x_i)| \quad \text{und} \quad d_{ui} = |S(x_{i-1}) - F_0(x_i)|$$

berechnet, wobei $S(x_0) := 0$ gesetzt wird. Wenn die größte Differenz d_{\max} aus allen Differenzen d_{oi}, d_{ui}

einen kritischen Wert d_α übersteigt, wird die Hypothese abgelehnt.

Bis $n=40$ greift man auf Tabellen zurück (s. Anhang). Für größere n werden sie über $d_\alpha = \frac{\sqrt{\ln(\frac{2}{\alpha})}}{\sqrt{2n}}$ angenähert.

Zweistichprobenproblem =

Liegt nun zusätzlich zur Zufallsvariablen X eine entsprechende Zufallsvariable Y vor (mit m geordneten Werten y_i),

so kann durch den Zweistichprobentest überprüft werden, ob X und Y derselben Verteilungsfunktion folgen.

Von beiden Beobachtungen werden die die Differenzen der relativen Summenfunktionen $S_X(x_i)$ bzw. $S_Y(y_i)$ ermittelt:

$$d(z) = |S_X(z) - S_Y(z)| \quad \text{und} \quad d_{\max} = \sup_z d(z) \quad .$$

Die Nullhypothese wird abgelehnt, falls d_{\max} den kritischen Wert $d_{\text{krit}}(\alpha, n, m)$ überschreitet.

Für kleine Werte von n und m greift man auf Tabellen zurück.

Für große Werte von n und m wird die Nullhypothese abgelehnt, falls

$$\sqrt{\frac{nm}{n+m}} d_{\max} > K_\alpha$$

wobei K_α für große n und m näherungsweise als $K_\alpha = \sqrt{\frac{\ln(\frac{2}{\alpha})}{2}}$ berechnet werden kann.

6 Varianzanalyse

univariate Varianzanalyse (ANOVA)

Man untersucht man den Einfluss einer unabhängigen Variable (Faktor) mit k verschiedenen Stufen (Gruppen) auf die Ausprägungen einer Zufallsvariablen.

Dazu werden die k Mittelwerte der Ausprägungen für die Gruppen miteinander verglichen, und zwar vergleicht man die Varianz zwischen den Gruppen mit der Varianz innerhalb der Gruppen.

Weil sich die totale Varianz aus den zwei genannten Komponenten zusammensetzt, spricht man von Varianzanalyse.

Die einfaktorielle ANOVA ist die Verallgemeinerung des t-Tests bei mehr als zwei Gruppen. Für $k=2$ ist sie äquivalent mit dem t-Test.

Es sei μ_i der Erwartungswert der abhängigen Variable in der i . Gruppe.

$H_0 : \mu_1 = \mu_2 = \dots = \mu_k$ (Es besteht kein Unterschied zwischen den Erwartungswerten der Gruppen.)

$H_1 : \exists i, j : \mu_i \neq \mu_j$ (Es besteht zwischen *mindestens* zwei Erwartungswerten ein Unterschied.)

→ Wir wissen dann nur mit einer bestimmten Wahrscheinlichkeit, dass mindestens zwei Ausprägungen einen bedeutsamen Unterschied aufweisen.

Effektdarstellung :

$$X_{ij} = \mu + \alpha_i + \varepsilon_{ij}, \quad i = 1, \dots, k, \quad j = 1, \dots, n_i.$$

Darin sind:

X_{ij} : Zielvariable; annahmegemäß in den Gruppen normalverteilt

k : Anzahl der Faktorstufen des betrachteten Faktors

n_i : Stichprobenumfänge für die einzelnen Faktorstufen

μ : arithmetisches Mittel der Erwartungswerte in den Gruppen

α_i : Effekt der i -ten Faktorstufe

ε_{ij} : Störvariablen, unabhängig und normalverteilt mit Erwartungswert 0 und gleicher (unbekannter) Varianz σ^2 .

Erwartungswert in der i . Gruppe: $\mu_i = \mu + \alpha_i$

$$\sum_{i=1}^k n_i \alpha_i = 0.$$

Betrachtung der **Quadratsummen** (Variabilität)

Die gesamte Variabilität QST (gesamte quadratische Abweichung vom Mittelwert) lässt sich in zwei Teile zerlegen:

$$QST = \sum (X_{ij} - \bar{X})^2 = QSA + QSE$$

Der erste Teil QSA (Gruppenzugehörigkeit) entspricht der ('Inter-')Variabilität zwischen den Gruppen.

$$QSA = \sum_i n_i (\bar{X}_i - \bar{X})^2,$$

Der Rest QSE entspricht der Variabilität innerhalb der Gruppen (gesamte 'Intra'-Abweichung von den Mittelwerten in den Gruppen, der 'Zufall'):

$$QSE = \sum_{i,j} (X_{ij} - \bar{X}_i)^2.$$

Die zwei Quadratsummen QSA und QSE sind stochastisch unabhängig.

Im Fall von k Gruppen mit gleichem Umfang $b=n/k$ gilt unter der Nullhypothese außerdem:

QSA/σ^2 folgt einer Chi-Quadrat-Verteilung mit $k-1$ Freiheitsgraden,

und

QSE/σ^2 folgt einer Chi-Quadrat-Verteilung mit $n-k$ Freiheitsgraden.

mittlere Quadratsummen:

$$MQSA = \frac{1}{k-1} QSA, \text{ und } MQSE = \frac{1}{n-k} QSE.$$

Prüfgröße:

$$F = \frac{MQSA}{MQSE}.$$

Gruppen gleicher Größe

Im Falle Gruppen gleicher Größe ist F unter der Nullhypothese F-verteilt mit $k - 1$ Freiheitsgraden im Zähler und $k \cdot (b - 1)$ Freiheitsgraden im Nenner.

Wenn die Prüfgröße

$$F = \frac{MQSA}{MQSE} = \frac{\frac{1}{k-1} \cdot b \cdot \sum_i (\bar{X}_i - \bar{\bar{X}})^2}{\frac{1}{k \cdot (b-1)} \cdot \sum_{i,j} (X_{ij} - \bar{X}_i)^2}.$$

signifikant (d.h. $F > F_{krit}(\alpha, k-1, k \cdot (b-1))$) wird, unterscheiden sich mindestens zwei Faktoren ('Gruppen') voneinander.

In Post-Hoc-Tests kann dann berechnet werden, zwischen welchen einzelnen Gruppen der Unterschied liegt.

7 Regressionsrechnung

7.0.3 Lineare Regression

Methode der kleinsten Quadrate

$$RSS = \sum_{i=1}^n d_i^2 = \sum_{i=1}^n (y_i - (a + bx_i))^2 \rightarrow \min!$$

bezüglich a und b.

Nach Ausmultiplikation, Ableiten und Nullsetzen

$$\frac{\partial S}{\partial a} = -2 \sum_{i=1}^n y_i + 2na + 2b \sum_{i=1}^n x_i = 0,$$

$$\frac{\partial S}{\partial b} = -2 \sum_{i=1}^n x_i y_i + 2a \sum_{i=1}^n x_i + 2b \sum_{i=1}^n x_i^2 = 0,$$

erhält man die gesuchten Regressionskoeffizienten als die Lösungen

$$b = \frac{\sum_{i=1}^n x_i y_i - n\bar{x}\bar{y}}{\sum_{i=1}^n x_i^2 - n\bar{x}^2}$$

und

$$a = \bar{y} - b\bar{x},, \text{ wobei } \bar{x} = \frac{1}{n} \sum_{i=1}^n x_i.$$

Mit dem Verschiebungssatz:

$$b = \frac{\sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y})}{\sum_{i=1}^n (x_i - \bar{x})^2}$$

Schätzungen \hat{y}

$$\hat{y}_i = a + bx_i$$

Residuen r_i :

$$\begin{aligned} y_i &= a + bx_i + d_i = \hat{y}_i + d_i \\ \Rightarrow d_i &= y_i - \hat{y}_i \end{aligned}$$

Stichprobenvarianz der Residuen:

$$s^2 = \frac{1}{n-2} \sum_i d_i^2$$

Bestimmtheitsmaß

$$r^2 = \frac{\frac{1}{n} \sum_{i=1}^n (\hat{y}_i - \bar{y})^2}{\frac{1}{n} \sum_{i=1}^n (y_i - \bar{y})^2} = \frac{(\sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y}))^2}{\sum_{i=1}^n (x_i - \bar{x})^2 \sum_{i=1}^n (y_i - \bar{y})^2},$$

mit dem Verschiebungssatz :

$$r^2 = \frac{(\sum_{i=1}^n x_i y_i - n \cdot \bar{x} \cdot \bar{y})^2}{(\sum_{i=1}^n x_i^2 - n \cdot \bar{x}^2)(\sum_{i=1}^n y_i^2 - n \cdot \bar{y}^2)}.$$

$$0 \leq r^2 \leq 1$$

Varianz der Residuen

$$s^2 = \frac{1}{n-2} (1 - r^2) \cdot \sum_{i=1}^n (y_i - \bar{y})^2$$

7.0.4 Variablentransformation

Funktion	u	v
$y = a + b \cdot x^n$	x^n	$v = a + b \cdot u$
$y = \frac{a}{b+x}$	x	$v = \frac{1}{y} = \frac{b}{a} + \frac{1}{b} \cdot u$
$y = a \cdot x^b$	$\ln(x)$	$v = \ln(y) = \ln(a) + b \cdot \ln(x)$
$y = a \cdot b^x$	x	$v = \ln(y) = \ln(a) + x \cdot \ln(b)$
$y = a \cdot e^{b \cdot x}$	x	$v = \ln(y) = \ln(a) + b \cdot x$

Die Methode kann auf weitere Parameter erweitert werden.

8 Zeitreihenanalyse

8.1 Komponentenunterteilung bei Zeitreihen

Mögliche Aufteilung einer Zeitreihe in Komponenten:

- Trend Q
- Konjunkturelle Schwankung K
- Saisonale Schwankung S
- Restschwankung r

Bei Unabhängigkeit dieser Komponenten kann man ein additives Modell annehmen:

$$y = Q + K + S + r$$

Nehmen beispielsweise zyklische Schwankungen mit steigendem Trend zu, könnte ein multiplikatives Modell

$$y_t = Q_t \cdot K_t \cdot r_t$$

angebracht sein. Variablentransformation durch Logarithmieren

$$\log y_t = \log Q_t + \log S_t + \log r_t$$

8.1.1 Schätzung des Trends durch Regression

“Regressionsmodell“

$$\hat{y}_t = a + bt \text{ bzw. } y_t = a + bt + d_t \quad (t = 1, 2, \dots, T; y_t = y_1, y_2, \dots, y_T)$$

mit den Lösungen

$$\begin{aligned} b &= \frac{\sum_{t=1}^T (t - \bar{t})(y_t - \bar{y})}{\sum_{t=1}^T (t - \bar{t})^2} \\ &= \frac{\sum_{t=1}^T t \cdot y_t - T \cdot \bar{t} \cdot \bar{y}}{\sum_{t=1}^T t^2 - T \cdot \bar{t}^2} \\ &= \frac{\sum_{t=1}^T t y_t - \frac{T(T+1)}{2} \bar{y}}{\frac{1}{12}(T^3 - T)} \end{aligned}$$

und

$$a = \bar{y} - b \cdot \bar{t}$$

$$= \bar{y} - b \cdot \frac{T+1}{2}$$

Die Trendwerte Q_t sind dann

$$Q_t = \hat{y}_t = a + bt$$

Nichtlinearer Trendverlauf: Lösung über Variablentransformation oder Anwendung eines nichtlinearen Regressionsansatzes

8.1.2 Schätzung der Saisonkomponente

Additives Modell

$$y_t = Q_t + S_t + r_t$$

Nach Schätzung der Trendkomponente Q_t bleibt noch die Abweichung

$$d_t = y_t - Q_t$$

und

$$d_t = S_t + r_t$$

d_t : trendbereinigter Zeitreihenwert

Bestimmung der saisonalen Komponente S_t über Fourieranalyse oder (einfacher)

Bildung des arithmetischen Durchschnitts aller Werte d_t , die die gleiche Saison betreffen, als Schätzung für die saisonale Komponente. Dann bleibt die nichterklärte Restschwankung

$$r_t = y_t - Q_t - S_t$$

Prognose für den Zeitpunkt $T+k$ (mit S_t als Wert in der Saison $T+k$)

$$\hat{y}_{T+k} = Q_{T+k} + S_{T+k},$$

8.1.3 Schätzung der glatten Komponente mit gleitenden Mittelwerten

Lässt sich die Trendkomponente des Zeitreihenmodells offensichtlich durch keine funktionale lineare oder

nichtlineare Beziehung darstellen, kann man eine **glatte Komponente** mit Hilfe gleitender Mittelwerte bestimmen.

einfacher gleitender Mittelwert

Beispiel: Mittelwert dritter Ordnung:

$$Y_k = \frac{1}{3} \cdot (Y_{k-1} + Y_k + Y_{k+1}) = \frac{1}{3} \sum_{i=k-1}^{k+1} Y_i$$

Die Ordnung des Mittelwerts sollte so gewählt werden, daß möglichst genau eine Periode umfasst wird.

Zur Prognose über den Beobachtungszeitraum hinaus sind gleitende Mittelwerte bedingt geeignet,

da die Randwerte der Zeitreihe nicht geschätzt werden.

gewichteter gleitender Mittelwert

Beispiel: Mittelwert dritter Ordnung mit z.B. $w_1 = \frac{1}{4}, w_2 = \frac{1}{2}, w_3 = \frac{1}{4}, \sum_i w_i = 1$

$$Y_k = w_1 \cdot Y_{k-1} + w_2 \cdot Y_k + w_3 \cdot Y_{k+1}$$

8.1.4 Exponentielle Glättung

Gewichtung durch den Glättungsfaktor α mit $0 \leq \alpha \leq 1$:

Geglätteter Schätzwert y_t^* als gewichteter Durchschnitt aus dem aktuellen Zeitreihenwert y_t

und dem Schätzwert der Vorperiode y_{t-1}^* (y_0^* geeignet wählen):

$$y_t^* = \alpha \cdot y_t + (1 - \alpha) \cdot y_{t-1}^*$$

Auflösung der Rekursivität:

$$y_t^* = \alpha y_t + \alpha(1 - \alpha)y_{t-1} + \alpha(1 - \alpha)^2 y_{t-2} + \dots + \alpha(1 - \alpha)^{t-1} y_1 + (1 - \alpha)^t y_0.$$

Für die Wahl des Glättungsfaktors wird häufig 0,2 bis 0,3 empfohlen. Man kann aber auch mit Hilfe der Regressionsanalyse den Glättungsfaktor schätzen.

Exponentielle Glättung bei trendbehafteten Werten

Bei Trend werden die Zeitreihenwerte systematisch unter- bzw. überschätzt. Abhilfe bieten ggf. gleitende Durchschnitte zweiter Ordnung.

Die bereits einmal geglätteten Werte erneut einer Glättung unterzogen. Man erhält den Schätzwert y^{**} , der sich analog zu oben berechnet aus

$$y_t^{**} = \alpha \cdot y_t^* + (1 - \alpha) \cdot y_{t-1}^{**}$$

Für einen brauchbaren Prognosewert für Periode $t+1$ muss man dann bestimmen

$$\hat{y}_{t+1} = 2 \cdot y_t^* - y_{t-1}^{**}$$

9 Symbolverzeichnis

9.1 allgemein

Symbol	Verwendung
A,B	Ereignisse
$\Omega = \{A,B,C\dots\}$	Ereignisraum
$ A $	Anzahl der Ereignisse A
$P(A)$	Wahrscheinlichkeit für das Eintreten von A
$\cup \cap$	Und-Verknüpfung (Konjunktion) / Oder-Verknüpfung (Disjunktion)
$P(A B)$	Bedingte Wahrscheinlichkeit (A wenn B)
$E(X)$	Erwartungswert von X
$f(X)$	Wahrscheinlichkeitsfunktion
$F(X)$	Verteilungsfunktion
N	Grundgesamtheit
n	Stichprobe
X	Zufallsvariable
$Var(X)$	Varianz von X
Θ	Anteilswert einer Grundgesamtheit

9.2 Verteilungsmodelle

Symbol	Verwendung
$b(x n;\theta)$	Binomialverteilung
$F(m,n)$	Fisher-Verteilung
$h(x N; M; n)$	Hypergeometrische Verteilung
$p(x \lambda)$	Poissonverteilung
$N(\mu, \sigma^2)$	Normalverteilung
$t(n)$	t- (Student-) Verteilung
$\phi_x(z)$	Standardnormalverteilung

10 Tabellen

Binomialverteilung (Wahrscheinlichkeitsfunktion)								
n	x	p= 0,01	... 0,05	0,1	0,2	0,25	0,3	0,5
2	0	0,9801	0,9025	0,81	0,64	0,5625	0,49	0,25
	1	0,0198	0,095	0,18	0,32	0,375	0,42	0,5
	2	0,0001	0,0025	0,01	0,04	0,0625	0,09	0,25
3	0	0,9703	0,8574	0,729	0,512	0,4219	0,343	0,125
	1	0,0294	0,1354	0,243	0,384	0,4219	0,441	0,375
	2	0,0003	0,0071	0,027	0,0960	0,1406	0,189	0,375
	3	0	0,0001	0,001	0,008	0,0156	0,027	0,125
4	0	0,9606	0,8145	0,6561	0,4096	0,3164	0,2401	0,0625
	1	0,0388	0,1715	0,2916	0,4096	0,4219	0,4116	0,25
	2	0,0006	0,0135	0,0486	0,1536	0,2109	0,2646	0,375
	3	0	0,0005	0,0036	0,0256	0,0469	0,0756	0,25
	4	0	0	0,0001	0,0016	0,0039	0,0081	0,0625
5	0	0,951	0,7738	0,5905	0,3277	0,2373	0,1681	0,0313
	1	0,048	0,2036	0,3281	0,4096	0,3955	0,3602	0,1563
	2	0,001	0,0214	0,0729	0,2048	0,2637	0,3087	0,3125
	3	0	0,0011	0,0081	0,0512	0,0879	0,1323	0,3125
	4	0	0	0,0005	0,0064	0,0146	0,0284	0,1563
	5	0	0	0	0,0003	0,001	0,0024	0,0313
6	0	0,9415	0,7351	0,5314	0,2621	0,178	0,1176	0,0156
	1	0,0571	0,2321	0,3543	0,3932	0,356	0,3025	0,0938
	2	0,0014	0,0305	0,0984	0,2458	0,2966	0,3241	0,2344
	3	0	0,0021	0,0146	0,0819	0,1318	0,1852	0,3125
	4	0	0,0001	0,0012	0,0154	0,033	0,0595	0,2344
	5	0	0	0,0001	0,0015	0,0044	0,0102	0,0938
	6	0	0	0	0,0001	0,0002	0,0007	0,0156
7	0	0,9321	0,6983	0,4783	0,2097	0,1335	0,0824	0,0078
	1	0,0659	0,2573	0,372	0,367	0,3115	0,2471	0,0547
	2	0,002	0,0406	0,124	0,2753	0,3115	0,3177	0,1641
	3	0	0,0036	0,023	0,1147	0,173	0,2269	0,2734
	4	0	0,0002	0,0026	0,0287	0,0577	0,0972	0,2734
	5	0	0	0,0002	0,0043	0,0115	0,025	0,1641
	6	0	0	0	0,0004	0,0013	0,0036	0,0547
	7	0	0	0	0	0,0001	0,0002	0,0078

Binomialverteilung (Wahrscheinlichkeitsfunktion)								
n	x	p= 0,01	... 0,05	0,1	0,2	0,25	0,3	0,5
8	0	0,9227	0,6634	0,4305	0,1678	0,1001	0,0576	0,0039
	1	0,0746	0,2793	0,3826	0,3355	0,267	0,1977	0,0313
	2	0,0026	0,0515	0,1488	0,2936	0,3115	0,2965	0,1094
	3	0,0001	0,0054	0,0331	0,1468	0,2076	0,2541	0,2188
	4	0	0,0004	0,0046	0,0459	0,0865	0,1361	0,2734
	5	0	0	0,0004	0,0092	0,0231	0,0467	0,2188
	6	0	0	0	0,0011	0,0038	0,01	0,1094
	7	0	0	0	0,0001	0,0004	0,0012	0,0313
	8	0	0	0	0	0	0,0001	0,0039
9	0	0,9135	0,6302	0,3874	0,1342	0,0751	0,0404	0,002
	1	0,083	0,2985	0,3874	0,302	0,2253	0,1556	0,0176
	2	0,0034	0,0629	0,1722	0,302	0,3003	0,2668	0,0703
	3	0,0001	0,0077	0,0446	0,1762	0,2336	0,2668	0,1641
	4	0	0,0006	0,0074	0,0661	0,1168	0,1715	0,2461
	5	0	0	0,0008	0,0165	0,0389	0,0735	0,2461
	6	0	0	0,0001	0,0028	0,0087	0,021	0,1641
	7	0	0	0	0,0003	0,0012	0,0039	0,0703
	8	0	0	0	0	0,0001	0,0004	0,0176
	9	0	0	0	0	0	0	0,002
10	0	0,9044	0,5987	0,3487	0,1074	0,0563	0,0282	0,001
	1	0,0914	0,3151	0,3874	0,2684	0,1877	0,1211	0,0098
	2	0,0042	0,0746	0,1937	0,302	0,2816	0,2335	0,0439
	3	0,0001	0,0105	0,0574	0,2013	0,2503	0,2668	0,1172
	4	0	0,001	0,0112	0,0881	0,146	0,2001	0,2051
	5	0	0,0001	0,0015	0,0264	0,0584	0,1029	0,2461
	6	0	0	0,0001	0,0055	0,0162	0,0368	0,2051
	7	0	0	0	0,0008	0,0031	0,009	0,1172
	8	0	0	0	0,0001	0,0004	0,0014	0,0439
	9	0	0	0	0	0	0,0001	0,0098
	10	0	0	0	0	0	0	0,001

Weitere Werte können mittels =BINOMVERT(x;n;p;0) bei Tabellenkalkulationsprogrammen

oder der R-Funktion¹ dbinom(x, n , p) bestimmt² werden

¹ http://de.wikibooks.org/wiki/GNU_R%20

² <http://netmath.vcrp.de/downloads/Systeme/R.html>

Standard-Normalverteilung (Verteilungsfunktion) $\phi_x(z)$										
z	0,00	0,01	0,02	0,03	0,04	0,05	0,06	0,07	0,08	0,09
0,0	0,5	0,504	0,508	0,512	0,516	0,5199	0,5239	0,5279	0,5319	0,5359
0,1	0,5398	0,5438	0,5478	0,5517	0,5557	0,5596	0,5636	0,5675	0,5714	0,5753
0,2	0,5793	0,5832	0,5871	0,591	0,5948	0,5987	0,6026	0,6064	0,6103	0,6141
0,3	0,6179	0,6217	0,6255	0,6293	0,6331	0,6368	0,6406	0,6443	0,648	0,6517
0,4	0,6554	0,6591	0,6628	0,6664	0,67	0,6736	0,6772	0,6808	0,6844	0,6879
0,5	0,6915	0,695	0,6985	0,7019	0,7054	0,7088	0,7123	0,7157	0,719	0,7224
0,6	0,7257	0,7291	0,7324	0,7357	0,7389	0,7422	0,7454	0,7486	0,7517	0,7549
0,7	0,758	0,7611	0,7642	0,7673	0,7704	0,7734	0,7764	0,7794	0,7823	0,7852
0,8	0,7881	0,791	0,7939	0,7967	0,7995	0,8023	0,8051	0,8078	0,8106	0,8133
0,9	0,8159	0,8186	0,8212	0,8238	0,8264	0,8289	0,8315	0,834	0,8365	0,8389
1,0	0,8413	0,8438	0,8461	0,8485	0,8508	0,8531	0,8554	0,8577	0,8599	0,8621
1,1	0,8643	0,8665	0,8686	0,8708	0,8729	0,8749	0,877	0,879	0,881	0,883
1,2	0,8849	0,8869	0,8888	0,8907	0,8925	0,8944	0,8962	0,898	0,8997	0,9015
1,3	0,9032	0,9049	0,9066	0,9082	0,9099	0,9115	0,9131	0,9147	0,9162	0,9177
1,4	0,9192	0,9207	0,9222	0,9236	0,9251	0,9265	0,9279	0,9292	0,9306	0,9319
1,5	0,9332	0,9345	0,9357	0,937	0,9382	0,9394	0,9406	0,9418	0,9429	0,9441
1,6	0,9452	0,9463	0,9474	0,9484	0,9495	0,9505	0,9515	0,9525	0,9535	0,9545
1,7	0,9554	0,9564	0,9573	0,9582	0,9591	0,9599	0,9608	0,9616	0,9625	0,9633
1,8	0,9641	0,9649	0,9656	0,9664	0,9671	0,9678	0,9686	0,9693	0,9699	0,9706
1,9	0,9713	0,9719	0,9726	0,9732	0,9738	0,9744	0,975	0,9756	0,9761	0,9767
2,0	0,9772	0,9778	0,9783	0,9788	0,9793	0,9798	0,9803	0,9808	0,9812	0,9817
2,1	0,9821	0,9826	0,983	0,9834	0,9838	0,9842	0,9846	0,985	0,9854	0,9857
2,2	0,9861	0,9864	0,9868	0,9871	0,9875	0,9878	0,9881	0,9884	0,9887	0,989
2,3	0,9893	0,9896	0,9898	0,9901	0,9904	0,9906	0,9909	0,9911	0,9913	0,9916
2,4	0,9918	0,992	0,9922	0,9925	0,9927	0,9929	0,9931	0,9932	0,9934	0,9936
2,5	0,9938	0,994	0,9941	0,9943	0,9945	0,9946	0,9948	0,9949	0,9951	0,9952
2,6	0,9953	0,9955	0,9956	0,9957	0,9959	0,996	0,9961	0,9962	0,9963	0,9964
2,7	0,9965	0,9966	0,9967	0,9968	0,9969	0,997	0,9971	0,9972	0,9973	0,9974
2,8	0,9974	0,9975	0,9976	0,9977	0,9977	0,9978	0,9979	0,9979	0,998	0,9981
2,9	0,9981	0,9982	0,9982	0,9983	0,9984	0,9984	0,9985	0,9985	0,9986	0,9986
3,0	0,9987	0,9987	0,9987	0,9988	0,9988	0,9989	0,9989	0,9989	0,999	0,999
3,1	0,999	0,9991	0,9991	0,9991	0,9992	0,9992	0,9992	0,9992	0,9993	0,9993
3,2	0,9993	0,9993	0,9994	0,9994	0,9994	0,9994	0,9994	0,9995	0,9995	0,9995
3,3	0,9995	0,9995	0,9995	0,9996	0,9996	0,9996	0,9996	0,9996	0,9996	0,9997
3,4	0,9997	0,9997	0,9997	0,9997	0,9997	0,9997	0,9997	0,9997	0,9997	0,9998
3,5	0,9998	0,9998	0,9998	0,9998	0,9998	0,9998	0,9998	0,9998	0,9998	0,9998
3,6	0,9998	0,9998	0,9999	0,9999	0,9999	0,9999	0,9999	0,9999	0,9999	0,9999

Zur Bildung von z ist der Wert von linker Spalte und oberer Zeile zu addieren.

Ablesebeispiel: $\phi_x(1,75) = 0,9599$

weitere Werte: =NORM.S.VERT(z ;WAHR) bzw. (R-Aufruf³): pnorm(z)

3 <http://netmath.vcrp.de/downloads/Systeme/R.html>

χ^2 -Verteilung (Wahrscheinlichkeitsfunktion)											
n	$\alpha=$ 0,005	0,05	0,1	0,5	0,9	0,95	0,975	0,99	0,995
1	0,000	0,000	0,001	0,004	0,016	0,455	2,706	3,841	5,024	6,635	7,879
2	0,010	0,020	0,051	0,103	0,211	1,386	4,605	5,991	7,378	9,21	10,597
3	0,072	0,115	0,216	0,352	0,584	2,366	6,251	7,815	9,348	11,345	12,838
4	0,207	0,297	0,484	0,711	1,064	3,357	7,779	9,488	11,143	13,277	14,86
5	0,412	0,554	0,831	1,145	1,61	4,351	9,236	11,07	12,833	15,086	16,75
6	0,676	0,872	1,237	1,635	2,204	5,348	10,645	12,592	14,449	16,812	18,548
7	0,989	1,239	1,69	2,167	2,833	6,346	12,017	14,067	16,013	18,475	20,278
8	1,344	1,646	2,18	2,733	3,49	7,344	13,362	15,507	17,535	20,09	21,955
9	1,735	2,088	2,7	3,325	4,168	8,343	14,684	16,919	19,023	21,666	23,589
10	2,156	2,558	3,247	3,94	4,865	9,342	15,987	18,307	20,483	23,209	25,188
11	2,603	3,053	3,816	4,575	5,578	10,341	17,275	19,675	21,92	24,725	26,757
12	3,074	3,571	4,404	5,226	6,304	11,34	18,549	21,026	23,337	26,217	28,3
13	3,565	4,107	5,009	5,892	7,042	12,34	19,812	22,362	24,736	27,688	29,819
14	4,075	4,66	5,629	6,571	7,79	13,339	21,064	23,685	26,119	29,141	31,319
15	4,601	5,229	6,262	7,261	8,547	14,339	22,307	24,996	27,488	30,578	32,801
16	5,142	5,812	6,908	7,962	9,312	15,338	23,542	26,296	28,845	32	34,267
17	5,697	6,408	7,564	8,672	10,085	16,338	24,769	27,587	30,191	33,409	35,718
18	6,265	7,015	8,231	9,39	10,865	17,338	25,989	28,869	31,526	34,805	37,156
19	6,844	7,633	8,907	10,117	11,651	18,338	27,204	30,144	32,852	36,191	38,582
20	7,434	8,26	9,591	10,851	12,443	19,337	28,412	31,41	34,17	37,566	39,997
21	8,034	8,897	10,283	11,591	13,24	20,337	29,615	32,671	35,479	38,932	41,401
22	8,643	9,542	10,982	12,338	14,041	21,337	30,813	33,924	36,781	40,289	42,796
23	9,26	10,196	11,689	13,091	14,848	22,337	32,007	35,172	38,076	41,638	44,181
24	9,886	10,856	12,401	13,848	15,659	23,337	33,196	36,415	39,364	42,98	45,559
25	10,52	11,524	13,12	14,611	16,473	24,337	34,382	37,652	40,646	44,314	46,928
26	11,16	12,198	13,844	15,379	17,292	25,336	35,563	38,885	41,923	45,642	48,29
27	11,808	12,879	14,573	16,151	18,114	26,336	36,741	40,113	43,195	46,963	49,645
28	12,461	13,565	15,308	16,928	18,939	27,336	37,916	41,337	44,461	48,278	50,993
29	13,121	14,256	16,047	17,708	19,768	28,336	39,087	42,557	45,722	49,588	52,336
30	13,787	14,953	16,791	18,493	20,599	29,336	40,256	43,773	46,979	50,892	53,672
31	14,458	15,655	17,539	19,281	21,434	30,336	41,422	44,985	48,232	52,191	55,003
32	15,134	16,362	18,291	20,072	22,271	31,336	42,585	46,194	49,48	53,486	56,328
33	15,815	17,074	19,047	20,867	23,11	32,336	43,745	47,4	50,725	54,776	57,648
34	16,501	17,789	19,806	21,664	23,952	33,336	44,903	48,602	51,966	56,061	58,964
35	17,192	18,509	20,569	22,465	24,797	34,336	46,059	49,802	53,203	57,342	60,275
37	18,586	19,96	22,106	24,075	26,492	36,336	48,363	52,192	55,668	59,893	62,883
38	19,289	20,691	22,878	24,884	27,343	37,335	49,513	53,384	56,896	61,162	64,181
39	19,996	21,426	23,654	25,695	28,196	38,335	50,66	54,572	58,12	62,428	65,476
40	20,707	22,164	24,433	26,509	29,051	39,335	51,805	55,758	59,342	63,691	66,766

weitere Werte: =CHIQU.INV(n;p) bzw. (R-Aufruf⁴): qchisq(p,n)

4 <http://netmath.vcrp.de/downloads/Systeme/R.html>

Kolmogorov-Smirnov- (KS-) Verteilung						
n	$\alpha=$ 0,2	...	0,05	0,02	0,01	0,005
1	0,9	0,95	0,975	0,99	0,995	0,9975
2	0,68377	0,77638	0,84187	0,89998	0,92925	0,94995
3	0,56481	0,63604	0,70758	0,78452	0,82895	0,86419
4	0,49265	0,56521	0,62392	0,68884	0,73417	0,77628
5	0,44697	0,50944	0,56326	0,62715	0,66848	0,70533
6	0,41035	0,46799	0,51925	0,57738	0,61655	0,65277
7	0,38145	0,43606	0,48341	0,53841	0,57576	0,60966
8	0,35828	0,40962	0,45426	0,50652	0,54174	0,5742
9	0,33907	0,38746	0,43	0,47957	0,51327	0,54435
10	0,32257	0,36866	0,40924	0,4566	0,48889	0,51864
11	0,30825	0,35241	0,39121	0,43668	0,46766	0,49631
12	0,29573	0,33814	0,37542	0,41916	0,449	0,47664
13	0,28466	0,32548	0,36142	0,4036	0,43243	0,45914
14	0,27477	0,31416	0,34889	0,38968	0,41758	0,44345
15	0,26585	0,30397	0,33759	0,37711	0,40416	0,42927
16	0,25774	0,29471	0,32733	0,36569	0,39197	0,41637
17	0,25035	0,28626	0,31796	0,35526	0,38083	0,40458
18	0,24356	0,2785	0,30935	0,34568	0,37059	0,39374
19	0,23731	0,27135	0,30142	0,33684	0,36114	0,38373
20	0,23152	0,26473	0,29407	0,32864	0,35238	0,37445
21	0,22614	0,25857	0,28724	0,32103	0,34423	0,36582
22	0,22111	0,25283	0,28086	0,31392	0,33663	0,35776
23	0,21642	0,24746	0,2749	0,30727	0,32951	0,35021
24	0,21201	0,24241	0,2693	0,30102	0,32283	0,34313
25	0,20786	0,23767	0,26404	0,29515	0,31654	0,33646
26	0,20396	0,2332	0,25907	0,28961	0,3106	0,33016
27	0,20026	0,22897	0,25437	0,28437	0,30499	0,32421
28	0,19676	0,22497	0,24993	0,2794	0,29968	0,31857
29	0,19344	0,22117	0,24571	0,27469	0,29463	0,31322
30	0,19029	0,21756	0,2417	0,27021	0,28984	0,30813
31	0,18728	0,21412	0,23788	0,26595	0,28527	0,30328
32	0,18442	0,21084	0,23424	0,26188	0,28091	0,29865
33	0,18168	0,20771	0,23076	0,25799	0,27675	0,29423
34	0,17906	0,20471	0,22743	0,25428	0,27276	0,29
35	0,17655	0,20184	0,22424	0,25072	0,26895	0,28595

Für $n > 35$ kann näherungsweise die Formel $d_\alpha = \frac{\sqrt{\ln(\frac{2}{\alpha})}}{\sqrt{2n}}$ verwendet werden.

Werte berechnet nach George Marsaglia et al . Evaluating Kolmogorov's DistributionEvaluating Kolmogorov's Distribution⁵. *Journal of Statistical Software* , **8** (18):1-4Nov 2003

⁵ www.jstatsoft.org/v08/i18/

F-Verteilung (Wahrscheinlichkeitsfunktion) Teil 1										
n_1	α	$n_2=1$	$n_2=2$...	4	5	6	7	8	9
1	0,950	161,448	18,513	10,128	7,709	6,608	5,987	5,591	5,318	5,117
	0,975	647,789	38,506	17,443	12,218	10,007	8,813	8,073	7,571	7,209
	0,990	4052,1	98,503	34,116	21,198	16,258	13,745	12,246	11,259	10,561
	0,995	16210,7	198,501	55,552	31,333	22,785	18,635	16,236	14,688	13,614
2	0,950	199,5	19,0	9,552	6,944	5,786	5,143	4,737	4,459	4,256
	0,975	799,5	39,0	16,044	10,649	8,434	7,26	6,542	6,059	5,715
	0,990	4999,5	99,0	30,817	18,0	13,274	10,925	9,547	8,649	8,022
	0,995	19999,5	199,0	49,799	26,284	18,314	14,544	12,404	11,042	10,107
3	0,950	215,707	19,164	9,277	6,591	5,409	4,757	4,347	4,066	3,863
	0,975	864,163	39,165	15,439	9,979	7,764	6,599	5,89	5,416	5,078
	0,990	5403,3	99,166	29,457	16,694	12,06	9,78	8,451	7,591	6,992
	0,995	21614,7	199,166	47,467	24,259	16,53	12,917	10,882	9,596	8,717
4	0,950	224,583	19,296	8,941	6,094	4,818	4,099	3,637	3,313	3,073
	0,975	899,583	39,298	14,735	9,074	6,757	5,523	4,761	4,243	3,868
	0,990	5624,58	99,299	27,911	14,976	10,289	7,976	6,62	5,734	5,111
	0,995	22499,6	199,3	44,838	21,622	13,961	10,391	8,38	7,104	6,227
5	0,950	230,162	19,296	9,013	6,256	5,05	4,387	3,972	3,687	3,482
	0,975	921,848	39,298	14,885	9,364	7,146	5,988	5,285	4,817	4,484
	0,990	5763,6	99,299	28,237	15,522	10,967	8,746	7,46	6,632	6,057
	0,995	23055,8	199,3	45,392	22,456	14,94	11,464	9,522	8,302	7,471
6	0,950	233,986	19,33	8,941	6,163	4,95	4,284	3,866	3,581	3,374
	0,975	937,111	39,331	14,735	9,197	6,978	5,82	5,119	4,652	4,32
	0,990	5859	99,333	27,911	15,207	10,672	8,466	7,191	6,371	5,802
	0,995	23437,1	199,333	44,838	21,975	14,513	11,073	9,155	7,952	7,134
7	0,950	236,768	19,353	8,887	6,094	4,876	4,207	3,787	3,5	3,293
	0,975	948,217	39,355	14,624	9,074	6,853	5,695	4,995	4,529	4,197
	0,990	5928,4	99,356	27,672	14,976	10,456	8,26	6,993	6,178	5,613
	0,995	23714,6	199,357	44,434	21,622	14,2	10,786	8,885	7,694	6,885
8	0,950	238,883	19,371	8,845	6,041	4,818	4,147	3,726	3,438	3,23
	0,975	956,656	39,373	14,54	8,98	6,757	5,6	4,899	4,433	4,102
	0,990	5981,1	99,374	27,489	14,799	10,289	8,102	6,84	6,029	5,467
	0,995	23925,4	199,375	44,126	21,352	13,961	10,566	8,678	7,496	6,693
9	0,950	240,543	19,385	8,812	5,999	4,772	4,099	3,677	3,388	3,179
	0,975	963,285	39,387	14,473	8,905	6,681	5,523	4,823	4,357	4,026
	0,990	6022,5	99,388	27,345	14,659	10,158	7,976	6,719	5,911	5,351
	0,995	24091	199,388	43,882	21,139	13,772	10,391	8,514	7,339	6,541
10	0,950	241,882	19,396	8,786	5,964	4,735	4,06	3,637	3,347	3,137
	0,975	968,627	39,398	14,419	8,844	6,619	5,461	4,761	4,295	3,964
	0,990	6055,8	99,399	27,229	14,546	10,051	7,874	6,62	5,814	5,257
	0,995	24224,5	199,4	43,686	20,967	13,618	10,25	8,38	7,211	6,417
11	0,950	242,983	19,405	8,763	5,936	4,704	4,027	3,603	3,313	3,102
	0,975	973,025	39,407	14,374	8,794	6,568	5,41	4,709	4,243	3,912
	0,990	6083,3	99,408	27,133	14,452	9,963	7,79	6,538	5,734	5,178
	0,995	24334,4	199,409	43,524	20,824	13,491	10,133	8,27	7,104	6,314

F-Verteilung (Wahrscheinlichkeitsfunktion) Teil 2										
n_1	α	$n_2=1$	$n_2=2$...	4	5	6	7	8	9
12	0,950	243,906	19,413	8,745	5,912	4,678	4,0	3,575	3,284	3,073
	0,975	976,708	39,415	14,337	8,751	6,525	5,366	4,666	4,2	3,868
	0,990	6106,3	99,416	27,052	14,374	9,888	7,718	6,469	5,667	5,111
	0,995	24426,4	199,416	43,387	20,705	13,384	10,034	8,176	7,015	6,227
13	0,950	244,69	19,419	8,729	5,891	4,655	3,976	3,55	3,259	3,048
	0,975	979,837	39,421	14,304	8,715	6,488	5,329	4,628	4,162	3,831
	0,990	6125,9	99,422	26,983	14,307	9,825	7,657	6,41	5,609	5,055
	0,995	24504,5	199,423	43,271	20,603	13,293	9,95	8,097	6,938	6,153
14	0,950	245,364	19,424	8,715	5,873	4,636	3,956	3,529	3,237	3,025
	0,975	982,528	39,427	14,277	8,684	6,456	5,297	4,596	4,13	3,798
	0,990	6142,7	99,428	26,924	14,249	9,77	7,605	6,359	5,559	5,005
	0,995	24571,8	199,428	43,172	20,515	13,215	9,877	8,028	6,872	6,089
15	0,950	245,95	19,429	8,703	5,858	4,619	3,938	3,511	3,218	3,006
	0,975	984,867	39,431	14,253	8,657	6,428	5,269	4,568	4,101	3,769
	0,990	6157,3	99,433	26,872	14,198	9,722	7,559	6,314	5,515	4,962
	0,995	24630,2	199,433	43,085	20,438	13,146	9,814	7,968	6,814	6,032
20	0,950	248,013	19,446	8,66	5,803	4,558	3,874	3,445	3,15	2,936
	0,975	993,103	39,448	14,167	8,56	6,329	5,168	4,467	3,999	3,667
	0,990	6208,7	99,449	26,69	14,02	9,553	7,396	6,155	5,359	4,808
	0,995	24836	199,45	42,778	20,167	12,903	9,589	7,754	6,608	5,832
40	0,950	251,143	19,471	8,594	5,717	4,464	3,774	3,34	3,043	2,826
	0,975	1005,59	39,473	14,037	8,411	6,175	5,012	4,309	3,84	3,505
	0,990	6286,8	99,474	26,411	13,745	9,291	7,143	5,908	5,116	4,567
	0,995	25148,2	199,475	42,308	19,752	12,53	9,241	7,422	6,288	5,519
50	0,950	251,774	19,476	8,581	5,699	4,444	3,754	3,319	3,02	2,803
	0,975	1008,11	39,478	14,01	8,381	6,144	4,98	4,276	3,807	3,472
	0,990	6302,5	99,479	26,354	13,69	9,238	7,091	5,858	5,065	4,517
	0,995	25211,1	199,48	42,213	19,667	12,454	9,17	7,354	6,222	5,454
75	0,950	252,618	19,482	8,563	5,676	4,418	3,726	3,29	2,99	2,771
	0,975	1011,48	39,485	13,974	8,34	6,101	4,937	4,232	3,762	3,426
	0,990	6323,6	99,486	26,278	13,615	9,166	7,022	5,789	4,998	4,449
	0,995	25295,3	199,486	42,086	19,554	12,351	9,074	7,263	6,133	5,367
100	0,950	253,041	19,486	8,554	5,664	4,405	3,712	3,275	2,975	2,756
	0,975	1013,17	39,488	13,956	8,319	6,08	4,915	4,21	3,739	3,403
	0,990	6334,1	99,489	26,24	13,577	9,13	6,987	5,755	4,963	4,415
	0,995	25337,5	199,49	42,022	19,497	12,3	9,026	7,217	6,088	5,322
150	0,950	253,465	19,489	8,545	5,652	4,392	3,698	3,26	2,959	2,739
	0,975	1014,86	39,491	13,938	8,299	6,059	4,893	4,188	3,716	3,38
	0,990	6344,7	99,492	26,202	13,539	9,094	6,951	5,72	4,929	4,38
	0,995	25379,7	199,493	41,957	19,44	12,248	8,977	7,17	6,042	5,278

weitere Werte: =F.INV(p;f1;f2) bzw. (R-Aufruf⁶): qf(p, f1 , f2)

⁶ <http://netmath.vcrp.de/downloads/Systeme/R.html>

t-(Student-)Verteilung (Wahrscheinlichkeitsfunktion)									
n	p= 0,75	... 0,90	0,95	0,975	0,99	0,995	0,999	0,9995	0,9999
1	1,00	3,078	6,314	12,706	31,821	63,657	318,31	636,62	3183,1
2	0,816	1,886	2,92	4,303	6,965	9,925	22,327	31,599	70,700
3	0,765	1,638	2,353	3,182	4,541	5,841	10,215	12,924	22,204
4	0,741	1,533	2,132	2,776	3,747	4,604	7,173	8,61	13,034
5	0,727	1,476	2,015	2,571	3,365	4,032	5,893	6,869	9,678
6	0,718	1,44	1,943	2,447	3,143	3,707	5,208	5,959	8,025
7	0,711	1,415	1,895	2,365	2,998	3,499	4,785	5,408	7,063
8	0,706	1,397	1,86	2,306	2,896	3,355	4,501	5,041	6,442
9	0,703	1,383	1,833	2,262	2,821	3,25	4,297	4,781	6,01
10	0,7	1,372	1,812	2,228	2,764	3,169	4,144	4,587	5,694
11	0,697	1,363	1,796	2,201	2,718	3,106	4,025	4,437	5,453
12	0,695	1,356	1,782	2,179	2,681	3,055	3,93	4,318	5,263
13	0,694	1,35	1,771	2,16	2,65	3,012	3,852	4,221	5,111
14	0,692	1,345	1,761	2,145	2,624	2,977	3,787	4,14	4,985
15	0,691	1,341	1,753	2,131	2,602	2,947	3,733	4,073	4,88
16	0,69	1,337	1,746	2,12	2,583	2,921	3,686	4,015	4,791
17	0,689	1,333	1,74	2,11	2,567	2,898	3,646	3,965	4,714
18	0,688	1,33	1,734	2,101	2,552	2,878	3,61	3,922	4,648
19	0,688	1,328	1,729	2,093	2,539	2,861	3,579	3,883	4,59
20	0,687	1,325	1,725	2,086	2,528	2,845	3,552	3,85	4,539
21	0,686	1,323	1,721	2,08	2,518	2,831	3,527	3,819	4,493
22	0,686	1,321	1,717	2,074	2,508	2,819	3,505	3,792	4,452
23	0,685	1,319	1,714	2,069	2,5	2,807	3,485	3,768	4,415
24	0,685	1,318	1,711	2,064	2,492	2,797	3,467	3,745	4,382
25	0,684	1,316	1,708	2,06	2,485	2,787	3,45	3,725	4,352
26	0,684	1,315	1,706	2,056	2,479	2,779	3,435	3,707	4,324
27	0,684	1,314	1,703	2,052	2,473	2,771	3,421	3,69	4,299
28	0,683	1,313	1,701	2,048	2,467	2,763	3,408	3,674	4,275
29	0,683	1,311	1,699	2,045	2,462	2,756	3,396	3,659	4,254
30	0,683	1,31	1,697	2,042	2,457	2,75	3,385	3,646	4,234
31	0,682	1,309	1,696	2,04	2,453	2,744	3,375	3,633	4,216
32	0,682	1,309	1,694	2,037	2,449	2,738	3,365	3,622	4,198
33	0,682	1,308	1,692	2,035	2,445	2,733	3,356	3,611	4,182
34	0,682	1,307	1,691	2,032	2,441	2,728	3,348	3,601	4,167
35	0,682	1,306	1,69	2,03	2,438	2,724	3,34	3,591	4,153
37	0,681	1,305	1,687	2,026	2,431	2,715	3,326	3,574	4,127
38	0,681	1,304	1,686	2,024	2,429	2,712	3,319	3,566	4,116
39	0,681	1,304	1,685	2,023	2,426	2,708	3,313	3,558	4,105
40	0,681	1,303	1,684	2,021	2,423	2,704	3,307	3,551	4,094

weitere Werte: =T.INV(p;n) bzw. (R-Aufruf⁷): qt(p,n)

⁷ <http://netmath.vcrp.de/downloads/Systeme/R.html>

11 Autoren

Edits	User
10	Dirk Hünninger ¹
2	Juetho ²
166	Uli.schell ³

¹ http://de.wikibooks.org/wiki/Benutzer:Dirk_H%25C3%25BCnninger

² <http://de.wikibooks.org/wiki/Benutzer:Juetho>

³ <http://de.wikibooks.org/wiki/Benutzer:Uli.schell>

Abbildungsverzeichnis

- GFDL: Gnu Free Documentation License. <http://www.gnu.org/licenses/fdl.html>
- cc-by-sa-3.0: Creative Commons Attribution ShareAlike 3.0 License. <http://creativecommons.org/licenses/by-sa/3.0/>
- cc-by-sa-2.5: Creative Commons Attribution ShareAlike 2.5 License. <http://creativecommons.org/licenses/by-sa/2.5/>
- cc-by-sa-2.0: Creative Commons Attribution ShareAlike 2.0 License. <http://creativecommons.org/licenses/by-sa/2.0/>
- cc-by-sa-1.0: Creative Commons Attribution ShareAlike 1.0 License. <http://creativecommons.org/licenses/by-sa/1.0/>
- cc-by-2.0: Creative Commons Attribution 2.0 License. <http://creativecommons.org/licenses/by/2.0/>
- cc-by-2.0: Creative Commons Attribution 2.0 License. <http://creativecommons.org/licenses/by/2.0/deed.en>
- cc-by-2.5: Creative Commons Attribution 2.5 License. <http://creativecommons.org/licenses/by/2.5/deed.en>
- cc-by-3.0: Creative Commons Attribution 3.0 License. <http://creativecommons.org/licenses/by/3.0/deed.en>
- GPL: GNU General Public License. <http://www.gnu.org/licenses/gpl-2.0.txt>
- LGPL: GNU Lesser General Public License. <http://www.gnu.org/licenses/lgpl.html>
- PD: This image is in the public domain.
- ATTR: The copyright holder of this file allows anyone to use it for any purpose, provided that the copyright holder is properly attributed. Redistribution, derivative work, commercial use, and all other use is permitted.
- EURO: This is the common (reverse) face of a euro coin. The copyright on the design of the common face of the euro coins belongs to the European Commission. Authorised is reproduction in a format without relief (drawings, paintings, films) provided they are not detrimental to the image of the euro.
- LFK: Lizenz Freie Kunst. <http://artlibre.org/licence/lal/de>
- CFR: Copyright free use.

- EPL: Eclipse Public License. <http://www.eclipse.org/org/documents/epl-v10.php>

Copies of the GPL, the LGPL as well as a GFDL are included in chapter Licenses⁴. Please note that images in the public domain do not require attribution. You may click on the image numbers in the following table to open the webpage of the images in your webbrowser.

⁴ Kapitel 12 auf Seite 63

1	Dirk Huenniger, MichaelFrey, Philipendula	
2	Philipendula ⁵ , Philipendula ⁶	CC-BY-SA-3.0
3	Philipendula ⁷ , Philipendula ⁸	CC-BY-SA-3.0
4	Szczepan ⁹ dyskusja ¹⁰ Mail ¹¹ 16:25, 9 January 2007 (UTC), Szczepan ¹² dyskusja ¹³ Mail ¹⁴ 16:25, 9 January 2007 (UTC)	PD
5	Szczepan ¹⁵ dyskusja ¹⁶ Mail ¹⁷ 16:36, 9 January 2007 (UTC), Szczepan ¹⁸ dyskusja ¹⁹ Mail ²⁰ 16:36, 9 January 2007 (UTC)	PD
6	Szczepan ²¹ dyskusja ²² Mail ²³ 16:35, 9 January 2007 (UTC), Szczepan ²⁴ dyskusja ²⁵ Mail ²⁶ 16:35, 9 January 2007 (UTC)	PD
7	Sigbert ²⁷ , Sigbert ²⁸	PD

- 5 <http://commons.wikimedia.org/wiki/User:Philipendula>
6 <http://wiki/User:Philipendula>
7 <http://commons.wikimedia.org/wiki/User:Philipendula>
8 <http://wiki/User:Philipendula>
9 <http://commons.wikimedia.org/wiki/User:Szczepan1990>
10 http://commons.wikimedia.org/wiki/User_talk:Szczepan1990
11 <http://commons.wikimedia.org/wiki/Special:EmailUser/Szczepan1990>
12 <http://wiki/User:Szczepan1990>
13 http://wiki/User_talk:Szczepan1990
14 <http://wiki/Special:EmailUser/Szczepan1990>
15 <http://commons.wikimedia.org/wiki/User:Szczepan1990>
16 http://commons.wikimedia.org/wiki/User_talk:Szczepan1990
17 <http://commons.wikimedia.org/wiki/Special:EmailUser/Szczepan1990>
18 <http://wiki/User:Szczepan1990>
19 http://wiki/User_talk:Szczepan1990
20 <http://wiki/Special:EmailUser/Szczepan1990>
21 <http://commons.wikimedia.org/wiki/User:Szczepan1990>
22 http://commons.wikimedia.org/wiki/User_talk:Szczepan1990
23 <http://commons.wikimedia.org/wiki/Special:EmailUser/Szczepan1990>
24 <http://wiki/User:Szczepan1990>
25 http://wiki/User_talk:Szczepan1990
26 <http://wiki/Special:EmailUser/Szczepan1990>
27 <http://commons.wikimedia.org/w/index.php?title=User:Sigbert&action=edit&redlink=1>
28 <http://w/index.php?title=User:Sigbert&action=edit&redlink=1>

Index

- Additionssatz, 4
- ANOVA, 35
- Arithmetisches Mittel, 18
- Axiome der Wahrscheinlichkeiten (Kolmogoroff), 4

- BAYES Theorem, 5
- Berücksichtigung der Reihenfolge, 3
- Binomialkoeffizient , 3
- Binomialtest, 29
- Binomialverteilung, Tabelle der, 47
- Binomialverteilung, 9

- Chi-Quadrat-Anpassungs- oder Verteilungstest, 32
- Chi-Quadrat-Unabhängigkeitstest, 32
- CHI-Quadrat-verteilung, 14
- CHI-Quadrat-Verteilung, Tabelle der, 49

- DE MORGANSche Regeln , 3
- Dichtefunktion, 10
- diskrete Zufallsvariable, 7

- einfacher gleitender Mittelwert, 43
- Erwartungswert, 7
- Erwartungswert einer stetigen Zufallsvariablen, 10
- Exponentialverteilung, 11
- Exponentielle Glättung, 43
- Exponentielle Glättung bei trendbehafteten Werten, 44

- F-Verteilung, Tabelle der, 51
- Fakultät , 3
- Fisher- Verteilung, 15

- geometrisches Mittel, 18
- gewichteter gleitender Mittelwert, 43
- Gini-Koeffizient, 20
- Grenzwertsatz, 16
- Grenzwertsatz von De Moivre, 16

- harmonisches Mittel, 18
- Hauptsatz der Statistik, 16
- Herfindahl-Index, 21
- Hypergeometrische Verteilung, 9
- Hypothesentest, 25

- Intervallskala, 17

- Kolmogorov-Smirnov-Verteilung, Tabelle der, 50
- Kolmogorow-Smirnow-Anpassungstest, 33
- Konfidenzintervall; Merkmal mit unbekannter Verteilung und bekannter Varianz, 23
- Konfidenzintervall; Merkmal mit unbekannter Verteilung und unbekannter Varianz, 24
- Konfidenzintervall; Normalverteiltes Merkmal mit bekannter Varianz, 23
- Konfidenzintervall; Normalverteiltes Merkmal mit unbekannter Varianz, 23
- Konfidenzintervalle für den Anteilswert einer dichotomen Grundgesamtheit, 24
- Konzentrationsmasse, 19
- Konzentrationsrate, 19
- Korrelationskoeffizient nach Bravais-Pearson, 8
- Kovarianz, 8

- Lineare Regression, 39
- Lorenzkurve, 20

- Median, 18
- Methode der kleinsten Quadrate, 39
- Modalwert, 19

- Nominalskala, 17
- Normalverteilung, 12
- Normalverteilung, Tabelle der, 48

- Ordinalskala, 17

- Poissonverteilung, 10
- Prinzip nach LAPLACE, 4

- Rangkorrelationskoeffizient nach Spearman, 9
- Rechteckverteilung, 11
- Regressionsrechnung, 39

- Schätzung der glatten Komponente mit gleitenden Mittelwerten, 43
- Schätzung der Saisonkomponente, 42
- Schätzung des Trends durch Regression, 41
- Skalenniveaus, 17
- stem-leaf-Diagramm, 18
- Stetige Gleichverteilung, 11

Student-Verteilung, 14
Student-Verteilung, Tabelle der, 53
Symmetrieprinzip, 4

t-Verteilung, 14
t-Verteilung, Tabelle der, 53
Test auf Anteilswert , 29
Test auf Erwartungswert, 25
Test auf Varianz, 31
Tests auf Zusammenhangs- und Assoziationsparameter, 32
Theorem von Bernoulli, 16
Totale Wahrscheinlichkeit, 5

Variabilität, 36
Variablentransformation, 40
Varianz, 7, 19
Varianz der Summe unabhängiger Zufallsvariablen, 8
Varianzanalyse, 35
Variationskoeffizient, 19
Verhältnisskala, 17
Verschiebungssatz, 7, 19
Verteilungsfunktion, 10
Vorzeichentest; Einstichprobenproblem, 27
Vorzeichentest; Zweistichprobenproblem , 27

Zeitreihenanalyse, 41
Zentraler Grenzwertsatz, 16
Zentralwert, 18
Zweig-Blätter-Diagramm, 18

12 Licenses

12.1 GNU GENERAL PUBLIC LICENSE

Version 3, 29 June 2007

Copyright © 2007 Free Software Foundation, Inc.
<<http://fsf.org/>>

Everyone is permitted to copy and distribute verbatim copies of this license document, but changing it is not allowed. Preamble

The GNU General Public License is a free, copyleft license for software and other kinds of works.

The licenses for most software and other practical works are designed to take away your freedom to share and change the works. By contrast, the GNU General Public License is intended to guarantee your freedom to share and change all versions of a program—to make sure it remains free software for all its users. We, the Free Software Foundation, use the GNU General Public License for most of our software; it applies also to any other work released this way by its authors. You can apply it to your programs, too.

When we speak of free software, we are referring to freedom, not price. Our General Public Licenses are designed to make sure that you have the freedom to distribute copies of free software (and charge for them if you wish), that you receive source code or can get it if you want it, that you can change the software or use pieces of it in new free programs, and that you know you can do these things.

To protect your rights, we need to prevent others from denying you these rights or asking you to surrender the rights. Therefore, you have certain responsibilities if you distribute copies of the software, or if you modify it: responsibilities to respect the freedom of others.

For example, if you distribute copies of such a program, whether gratis or for a fee, you must pass on to the recipients the same freedoms that you received. You must make sure that they, too, receive or can get the source code. And you must show them these terms so they know their rights.

Developers that use the GNU GPL protect your rights with two steps: (1) assert copyright on the software, and (2) offer you this License giving you legal permission to copy, distribute and/or modify it.

For the developers' and authors' protection, the GPL clearly explains that there is no warranty for this free software. For both users' and authors' sake, the GPL requires that modified versions be marked as changed, so that their problems will not be attributed erroneously to authors of previous versions.

Some devices are designed to deny users access to install or run modified versions of the software inside them, although the manufacturer can do so. This is fundamentally incompatible with the aim of protecting users' freedom to change the software. The systematic pattern of such abuse occurs in the area of products for individuals to use, which is precisely where it is most unacceptable. Therefore, we have designed this version of the GPL to prohibit the practice for those products. If such problems arise substantially in other domains, we stand ready to extend this provision to those domains in future versions of the GPL, as needed to protect the freedom of users.

Finally, every program is threatened constantly by software patents. States should not allow patents to restrict development and use of software on general-purpose computers, but in those that do, we wish to avoid the special danger that patents applied to a free program could make it effectively proprietary. To prevent this, the GPL assures that patents cannot be used to render the program non-free.

The precise terms and conditions for copying, distribution and modification follow. TERMS AND CONDITIONS 0. Definitions.

"This License" refers to version 3 of the GNU General Public License.

"Copyright" also means copyright-like laws that apply to other kinds of works, such as semiconductor masks.

"The Program" refers to any copyrightable work licensed under this License. Each licensee is addressed as "you". "Licensees" and "recipients" may be individuals or organizations.

To "modify" a work means to copy from or adapt all or part of the work in a fashion requiring copyright permission, other than the making of an exact copy. The resulting work is called a "modified version" of the earlier work or a work "based on" the earlier work.

A "covered work" means either the unmodified Program or a work based on the Program.

To "propagate" a work means to do anything with it that, without permission, would make you directly or secondarily liable for infringement under applicable copyright law, except executing it on a computer or modifying a private copy. Propagation includes copying, distribution (with or without modification), making available to the public, and in some countries other activities as well.

To "convey" a work means any kind of propagation that enables other parties to make or receive copies. Mere interaction with a user through a computer network, with no transfer of a copy, is not conveying.

An interactive user interface displays "Appropriate Legal Notices" to the extent that it includes a convenient and prominently visible feature that (1) displays an appropriate copyright notice, and (2) tells the user that there is no warranty for the work (except to the extent that warranties are provided), that licensees may convey the work under this License, and how to view a copy of this License. If the interface presents a list of user commands or options, such as a menu, a prominent item in the list meets this criterion. 1. Source Code.

The "source code" for a work means the preferred form of the work for making modifications to it. "Object code" means any non-source form of a work.

A "Standard Interface" means an interface that either is an official standard defined by a recognized standards body, or, in the case of interfaces specified for a particular programming language, one that is widely used among developers working in that language.

The "System Libraries" of an executable work include anything, other than the work as a whole, that (a) is included in the normal form of packaging a Major Component, but which is not part of that Major Component, and (b) serves only to enable use of the work with that Major Component, or to implement a Standard Interface for which an implementation is available to the public in source code form. A "Major Component", in this context, means a major essential component (kernel, window system, and so on) of the specific operating system (if any) on which the executable work runs, or a compiler used to produce the work, or an object code interpreter used to run it.

The "Corresponding Source" for a work in object code form means all the source code needed to generate, install, and (for an executable work) run the object code and to modify the work, including scripts to control those activities. However, it does not include the work's System Libraries, or general-purpose tools or generally available free programs which are used unmodified in performing those activities but which are not part of the work. For example, Corresponding Source includes interface definition files associated with source files for the work, and the source code for shared libraries and dynamically linked subprograms that the work is specifically designed to require, such as by intimate data communication or control flow between those subprograms and other parts of the work.

The Corresponding Source need not include anything that users can regenerate automatically from other parts of the Corresponding Source.

The Corresponding Source for a work in source code form is that same work. 2. Basic Permissions.

All rights granted under this License are granted for the term of copyright on the Program, and are irrevocable provided the stated conditions are met. This License explicitly affirms your unlimited permission to run the unmodified Program. The output from running a covered work is covered by this License only if the output, given its content, constitutes a covered work. This License acknowledges your rights of fair use or other equivalent, as provided by copyright law.

You may make, run and propagate covered works that you do not convey, without conditions so long as your license otherwise remains in force. You may convey covered works to others for the sole purpose of having them make modifications exclusively for you, or provide you with facilities for running those works, provided that you comply with the terms of this License in conveying all material for which you do not control copyright. Those thus making or running the covered works for you must do so exclusively on your behalf, under your direction and control, on terms that prohibit them from making any copies of your copyrighted material outside their relationship with you.

Conveying under any other circumstances is permitted solely under the conditions stated below. Sublicensing is not allowed; section 10 makes it unnecessary. 3. Protecting Users' Legal Rights From Anti-Circumvention Law.

No covered work shall be deemed part of an effective technological measure under any applicable law fulfilling obligations under article 11 of the WIPO copyright treaty adopted on 20 December 1996, or similar laws prohibiting or restricting circumvention of such measures.

When you convey a covered work, you waive any legal power to forbid circumvention of technological measures to the extent such circumvention is effected by exercising rights under this License with respect to the covered work, and you disclaim any intention to limit operation or modification of the work as a means of enforcing, against the work's users, your or third parties' legal rights to forbid circumvention of technological measures. 4. Conveying Verbatim Copies.

You may convey verbatim copies of the Program's source code as you receive it, in any medium, provided that you conspicuously and appropriately publish on each copy an appropriate copyright notice; keep intact all notices stating that this License and any non-permissive terms added in accord with section 7 apply to the code; keep intact all notices of

the absence of any warranty; and give all recipients a copy of this License along with the Program.

You may charge any price or no price for each copy that you convey, and you may offer support or warranty protection for a fee. 5. Conveying Modified Source Versions.

You may convey a work based on the Program, or the modifications to produce it from the Program, in the form of source code under the terms of section 4, provided that you also meet all of these conditions:

* a) The work must carry prominent notices stating that you modified it, and giving a relevant date. * b) The work must carry prominent notices stating that it is released under this License and any conditions added under section 7. This requirement modifies the requirement in section 4 to "keep intact all notices". * c) You must license the entire work, as a whole, under this License to anyone who comes into possession of a copy. This License will therefore apply, along with any applicable section 7 additional terms, to the whole of the work, and all its parts, regardless of how they are packaged. This License gives no permission to license the work in any other way, but it does not invalidate such permission if you have separately received it. * d) If the work has interactive user interfaces, each must display Appropriate Legal Notices; however, if the Program has interactive interfaces that do not display Appropriate Legal Notices, your work need not make them do so.

A compilation of a covered work with other separate and independent works, which are not by their nature extensions of the covered work, and which are not combined with it such as to form a larger program, in or on a volume of a storage or distribution medium, is called an "aggregate" if the compilation and its resulting copyright are not used to limit the access or legal rights of the compilation's users beyond what the individual works permit. Inclusion of a covered work in an aggregate does not cause this License to apply to the other parts of the aggregate. 6. Conveying Non-Source Forms.

You may convey a covered work in object code form under the terms of sections 4 and 5, provided that you also convey the machine-readable Corresponding Source under the terms of this License, in one of these ways:

* a) Convey the object code in, or embodied in, a physical product (including a physical distribution medium), accompanied by the Corresponding Source fixed on a durable physical medium customarily used for software interchange. * b) Convey the object code in, or embodied in, a physical product (including a physical distribution medium), accompanied by a written offer, valid for at least three years and valid for as long as you offer spare parts or customer support for that product model, to give anyone who possesses the object code either (1) a copy of the Corresponding Source for all the software in the product that is covered by this License, on a durable physical medium customarily used for software interchange, for a price no more than your reasonable cost of physically performing this conveying of source, or (2) access to copy the Corresponding Source from a network server at no charge. * c) Convey individual copies of the object code with a copy of the written offer to provide the Corresponding Source. This alternative is allowed only occasionally and noncommercially, and only if you received the object code with such an offer, in accord with subsection 6b. * d) Convey the object code by offering access from a designated place (gratis or for a charge), and offer equivalent access to the Corresponding Source in the same way through the same place at no further charge. You need not require recipients to copy the Corresponding Source along with the object code. If the place to copy the object code is a network server, the Corresponding Source may be on a different server (operated by you or a third party) that supports equivalent copying facilities, provided you maintain clear directions next to the object code saying where to find the Corresponding Source. Regardless of what server hosts the Corresponding Source, you remain obligated to ensure that it is available for as long as needed to satisfy these requirements. * e) Convey the object code using peer-to-peer transmission, provided you inform other peers where the object code and Corresponding Source of the work are being offered to the general public at no charge under subsection 6d.

A separable portion of the object code, whose source code is excluded from the Corresponding Source as a System Library, need not be included in conveying the object code work.

A "User Product" is either (1) a "consumer product", which means any tangible personal property which is normally used for personal, family, or household purposes, or (2) anything designed or sold for incorporation into a dwelling. In determining whether a product is a consumer product, doubtful cases shall be resolved in favor of coverage. For a particular product received by a particular user, "normally used" refers to a typical or common use of that class of product, regardless of the status of the particular user or of the way in which the particular user actually uses, or expects or is expected to use, the product. A product is a consumer product regardless of whether the product has substantial commercial, industrial or non-consumer uses, unless such uses represent the only significant mode of use of the product.

"Installation Information" for a User Product means any methods, procedures, authorization keys, or other information required to install and execute modified versions of a covered work in that User

Product from a modified version of its Corresponding Source. The information must suffice to ensure that the continued functioning of the modified object code is in no case prevented or interfered with solely because modification has been made.

If you convey an object code work under this section in, or with, or specifically for use in, a User Product, and the conveying occurs as part of a transaction in which the right of possession and use of the User Product is transferred to the recipient in perpetuity or for a fixed term (regardless of how the transaction is characterized), the Corresponding Source conveyed under this section must be accompanied by the Installation Information. But this requirement does not apply if neither you nor any third party retains the ability to install modified object code on the User Product (for example, the work has been installed in ROM).

The requirement to provide Installation Information does not include a requirement to continue to provide support service, warranty, or updates for a work that has been modified or installed by the recipient, or for the User Product in which it has been modified or installed. Access to a network may be denied when the modification itself materially and adversely affects the operation of the network or violates the rules and protocols for communication across the network.

Corresponding Source conveyed, and Installation Information provided, in accord with this section must be in a format that is publicly documented (and with an implementation available to the public in source code form), and must require no special password or key for unpacking, reading or copying. 7. Additional Terms.

"Additional permissions" are terms that supplement the terms of this License by making exceptions from one or more of its conditions. Additional permissions that are applicable to the entire Program shall be treated as though they were included in this License, to the extent that they are valid under applicable law. If additional permissions apply only to part of the Program, that part may be used separately under those permissions, but the entire Program remains governed by this License without regard to the additional permissions.

When you convey a copy of a covered work, you may at your option remove any additional permissions from that copy, or from any part of it. (Additional permissions may be written to require their own removal in certain cases when you modify the work.) You may place additional permissions on material, added by you to cover work for which you have or can give appropriate copyright permission.

Notwithstanding any other provision of this License, for material you add to a covered work, you may (if authorized by the copyright holders of that material) supplement the terms of this License with terms:

* a) Disclaiming warranty or limiting liability differently from the terms of sections 15 and 16 of this License; or * b) Requiring preservation of specified reasonable legal notices or author attributions in that material or in the Appropriate Legal Notices displayed by works containing it; or * c) Prohibiting misrepresentation of the origin of that material, or requiring that modified versions of such material be marked in reasonable ways as different from the original version; or * d) Limiting the use for publicity purposes of names of licensors or authors of the material; or * e) Declining to grant rights under trademark law for use of some trade names, trademarks, or service marks; or * f) Requiring indemnification of licensors and authors of that material by anyone who conveys the material (or modified versions of it) with contractual assumptions of liability to the recipient, for any liability that these contractual assumptions directly impose on those licensors and authors.

All other non-permissive additional terms are considered "further restrictions" within the meaning of section 10. If the Program as you received it, or any part of it, contains a notice stating that it is governed by this License along with a term that is a further restriction, you may remove that term. If a license document contains a further restriction but permits relicensing or conveying under this License, you may add to a covered work material governed by the terms of that license document, provided that the further restriction does not survive such relicensing or conveying.

If you add terms to a covered work in accord with this section, you must place, in the relevant source files, a statement of the additional terms that apply to those files, or a notice indicating where to find the applicable terms.

Additional terms, permissive or non-permissive, may be stated in the form of a separately written license, or stated as exceptions; the above requirements apply either way. 8. Termination.

You may not propagate or modify a covered work except as expressly provided under this License. Any attempt otherwise to propagate or modify it is void, and will automatically terminate your rights under this License (including any patent licenses granted under the third paragraph of section 11).

However, if you cease all violation of this License, then your license from a particular copyright holder is reinstated (a) provisionally, unless and until the copyright holder explicitly and finally terminates your license, and (b) permanently, if the copyright holder fails to notify you of the violation by some reasonable means prior to 60 days after the cessation.

Moreover, your license from a particular copyright holder is reinstated permanently if the copyright holder notifies you of the violation by some reasonable means, this is the first time you have received notice of violation of this License (for any work) from that copyright holder, and you cure the violation prior to 30 days after your receipt of the notice.

Termination of your rights under this section does not terminate the licenses of parties who have received copies or rights from you under this License. If your rights have been terminated and not permanently reinstated, you do not qualify to receive new licenses for the same material under section 10. 9. Acceptance Not Required for Having Copies.

You are not required to accept this License in order to receive or run a copy of the Program. Ancillary propagation of a covered work occurring solely as a consequence of using peer-to-peer transmission to receive a copy likewise does not require acceptance. However, nothing other than this License grants you permission to propagate or modify any covered work. These actions bring copyright infringement if you do not accept this License. Therefore, by modifying or propagating a covered work, you indicate your acceptance of this License to do so. 10. Automatic Licensing of Downstream Recipients.

Each time you convey a covered work, the recipient automatically receives a license from the original licensors, to run, modify and propagate that work, subject to this License. You are not responsible for enforcing compliance by third parties with this License.

An "entity transaction" is a transaction transferring control of an organization, or substantially all assets of one, or subdividing an organization, or merging organizations. If propagation of a covered work results from an entity transaction, each party to that transaction who receives a copy of the work also receives whatever licenses to the work the party's predecessor in interest had or could give under the previous paragraph, plus a right to possession of the Corresponding Source of the work from the predecessor in interest, if the predecessor has it or can get it with reasonable efforts.

You may not impose any further restrictions on the exercise of the rights granted or affirmed under this License. For example, you may not impose a license fee, royalty, or other charge for exercise of rights granted under this License, and you may not initiate litigation (including a cross-claim or counterclaim in a lawsuit) alleging that any patent claim is infringed by making, using, selling, offering for sale, or importing the Program or any portion of it. 11. Patents.

A "contributor" is a copyright holder who authorizes use under this License of the Program or a work on which the Program is based. The work thus licensed is called the contributor's "contributor version".

A contributor's "essential patent claims" are all patent claims owned or controlled by the contributor, whether already acquired or hereafter acquired, that would be infringed by some manner, permitted by this License, of making, using, or selling its contributor version, but do not include claims that would be infringed only as a consequence of further modification of the contributor version. For purposes of this definition, "control" includes the right to grant patent sublicenses in a manner consistent with the requirements of this License.

Each contributor grants you a non-exclusive, worldwide, royalty-free patent license under the contri-

12.2 GNU Free Documentation License

Version 1.3, 3 November 2008

Copyright © 2000, 2001, 2002, 2007, 2008 Free Software Foundation, Inc. <<http://fsf.org/>>

Everyone is permitted to copy and distribute verbatim copies of this license document, but changing it is not allowed. 0. PREAMBLE

The purpose of this License is to make a manual, textbook, or other functional and useful document "free" in the sense of freedom: to assure everyone the effective freedom to copy and redistribute it, with or without modifying it, either commercially or noncommercially. Secondly, this License preserves for the author and publisher a way to get credit for their work, while not being considered responsible for modifications made by others.

This License is a kind of "copyleft", which means that derivative works of the document must themselves be free in the same sense. It complements the GNU General Public License, which is a copyleft license designed for free software.

We have designed this License in order to use it for manuals for free software, because free software needs free documentation; a free program should come with manuals providing the same freedoms that the software does. But this License is not limited to software manuals; it can be used for any textual work, regardless of subject matter or whether it is published as a printed book. We recommend this License principally for works whose purpose is instruction or reference. 1. APPLICABILITY AND DEFINITIONS

This License applies to any manual or other work, in any medium, that contains a notice placed by the copyright holder saying it can be distributed under the terms of this License. Such a notice grants a world-wide, royalty-free license, unlimited in duration, to use that work under the conditions stated herein. The "Document", below, refers to any such manual or work. Any member of the public is a licensee, and is addressed as "you". You accept the license if you copy, modify or distribute the work in a way requiring permission under copyright law.

A "Modified Version" of the Document means any work containing the Document or a portion of it, either copied verbatim, or with modifications and/or translated into another language.

A "Secondary Section" is a named appendix or a front-matter section of the Document that deals ex-

clusively with the relationship of the publishers or authors of the Document to the Document's overall subject (or to related matters) and contains nothing that could fall directly within that overall subject. (Thus, if the Document is in part a textbook of mathematics, a Secondary Section may not explain any mathematics.) The relationship could be a matter of historical connection with the subject or with related matters, or of legal, commercial, philosophical, ethical or political position regarding them.

The Invariant Sections are certain Secondary Sections whose titles are designated, as being those of Invariant Sections, in the notice that says that the Document is released under this License. If a section does not fit the above definition of Secondary then it is not allowed to be designated as Invariant. The Document may contain zero Invariant Sections. If the Document does not identify any Invariant Sections then there are none.

The "Cover Texts" are certain short passages of text that are listed, as Front-Cover Texts or Back-Cover Texts, in the notice that says that the Document is released under this License. A Front-Cover Text may be at most 5 words, and a Back-Cover Text may be at most 25 words.

A "Transparent" copy of the Document means a machine-readable copy, represented in a format whose specification is available to the general public, that is suitable for revising the document straightforwardsly with generic text editors or (for images composed of pixels) with paint programs or (for drawings) some widely available drawing editor, and that is suitable for input to text formatters or for automatic translation to a variety of formats suitable for input to text formatters. A copy made in an otherwise Transparent file format whose markup, or absence of markup, has been arranged to thwart or discourage subsequent modification by readers is not Transparent. An image format is not Transparent if used for any substantial amount of text. A copy that is not "Transparent" is called "Opaque".

Examples of suitable formats for Transparent copies include plain ASCII without markup, Texinfo input format, LaTeX input format, SGML or XML using a publicly available DTD, and standard-conforming simple HTML, PostScript or PDF designed for human modification. Examples of transparent image formats include PNG, XCF and JPG. Opaque formats include proprietary formats that can be read and edited only by proprietary word processors, SGML or XML for which the DTD and/or processing tools are not generally available,

and the machine-generated HTML, PostScript or PDF produced by some word processors for output purposes only.

The "Title Page" means, for a printed book, the title page itself, plus such following pages as are needed to hold together the material this License requires to appear in the title page. For works in formats which do not have any title page as such, "Title Page" means the text near the most prominent appearance of the work's title, preceding the beginning of the body of the text.

The "publisher" means any person or entity that distributes copies of the Document to the public.

A section Entitled "XYZ" means a named subunit of the Document whose title either is precisely XYZ or contains XYZ in parentheses following text that translates XYZ in another language. (Here XYZ stands for a specific section name mentioned below, such as Acknowledgements, "Dedications", Endorsements", or "History".) To "Preserve the Title" of such a section when you modify the Document means that it remains a section Entitled "XYZ" according to this definition.

The Document may include Warranty Disclaimers next to the notice which states that this License applies to the Document. These Warranty Disclaimers are considered to be included by reference in this License, but only as regards disclaiming warranties; any other implication that these Warranty Disclaimers may have is void and has no effect on the meaning of this License. 2. VERBATIM COPYING

You may copy and distribute the Document in any medium, either commercially or noncommercially, provided that the material this License requires to appear in the title page, and the license notice saying this License applies to the Document are reproduced in all copies, and that you add no other conditions whatsoever to those of this License. You may not use technical measures to obstruct or control the reading or further copying of the copies you make or distribute. However, you may accept compensation in exchange for copies. If you distribute a large enough number of copies you must also follow the conditions in section 3.

You may also lend copies, under the same conditions stated above, and you may publicly display copies. 3. COPYING IN QUANTITY

If you publish printed copies (or copies in media that commonly have printed covers) of the Docu-

ment, numbering more than 100, and the Document's license notice requires Cover Texts, you must enclose the copies in covers that carry, clearly and legibly, all these Cover Texts: Front-Cover Texts on the front cover, and Back-Cover Texts on the back cover. Both covers must also clearly and legibly identify you as the publisher of these copies. The front cover must present the full title with all words of the title equally prominent and visible. You may add other material on the covers in addition. Copying with changes limited to the covers, as long as they preserve the title of the Document and satisfy these conditions, can be treated as verbatim copying in other respects.

If the required texts for either cover are too voluminous to fit legibly, you should put the first ones listed (as many as fit reasonably) on the actual cover, and continue the rest onto adjacent pages.

If you publish or distribute Opaque copies of the Document numbering more than 100, you must either include a machine-readable Transparent copy along with each Opaque copy, or state in or with each Opaque copy a computer-network location from which the general network-using public has access to download using public-standard network protocols a complete Transparent copy of the Document, free of added material. If you use the latter option, you must take reasonably prudent steps, when you begin distribution of Opaque copies in quantity, to ensure that this Transparent copy will remain thus accessible at the stated location until at least one year after the last time you distribute an Opaque copy (directly or through your agents or retailers) of that edition to the public.

If it is requested, but not required, that you contact the authors of the Document well before redistributing any large number of copies, to give them a chance to provide you with an updated version of the Document. 4. MODIFICATIONS

You may copy and distribute a Modified Version of the Document under the conditions of sections 2 and 3 above, provided that you release the Modified Version under precisely this License, with the Modified Version filling the role of the Document, thus licensing distribution and modification of the Modified Version to whoever possesses a copy of it. In addition, you must do these things in the Modified Version:

* A. Use in the Title Page (and on the covers, if any) a title distinct from that of the Document, and from those of previous versions (which should, if there were any, be listed in the History section

Notwithstanding any other provision of this License, you have permission to link or combine any covered work with a work licensed under version 3 of the GNU Affero General Public License into a single combined work, and to convey the resulting work. The terms of this License will continue to apply to the part which is the covered work, but the special requirements of the GNU Affero General Public License, section 13, concerning interaction through a network will apply to the combination as such. 14. Revised Versions of this License.

The Free Software Foundation may publish revised and/or new versions of the GNU General Public License from time to time. Such new versions will be similar in spirit to the present version, but may differ in detail to address new problems or concerns.

Each version is given a distinguishing version number. If the Program specifies that a certain numbered version of the GNU General Public License "or any later version" applies to it, you have the option of following the terms and conditions either of that numbered version or of any later version published by the Free Software Foundation. If the Program does not specify a version number of the GNU General Public License, you may choose any version ever published by the Free Software Foundation.

If the Program specifies that a proxy can decide which future versions of the GNU General Public License can be used, that proxy's public statement of acceptance of a version permanently authorizes you to choose that version for the Program.

Later license versions may give you additional or different permissions. However, no additional obligations are imposed on any author or copyright holder as a result of your choosing to follow a later version. 15. Disclaimer of Warranty.

There is NO WARRANTY for the program, to the extent permitted by applicable law, except when otherwise stated in writing the copyright holders and/or other parties provide the program "as is" without warranty of any kind, either expressed or implied, including, but not limited to, the implied warranties of merchantability and fitness for a particular purpose. The entire risk as to the quality and performance of the program is with you. Should the program prove defective, you assume the cost of all necessary servicing, repair or correction. 16. Limitation of Liability.

IN NO EVENT UNLESS REQUIRED BY APPLICABLE LAW OR AGREED TO IN WRITING WILL ANY COPYRIGHT HOLDER, OR ANY OTHER PARTY WHO MODIFIES AND/OR CONVEYS THE PROGRAM AS PERMITTED ABOVE, BE LIABLE TO YOU FOR DAMAGES, INCLUDING ANY GENERAL, SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES ARISING OUT OF THE USE OR INABILITY TO USE THE PROGRAM (INCLUDING BUT NOT LIMITED TO LOSS OF DATA OR DATA BEING RENDERED INACCURATE OR LOSSES SUSTAINED BY YOU OR THIRD PARTIES OR A FAILURE OF THE PROGRAM TO OPERATE WITH ANY OTHER PROGRAMS), EVEN IF SUCH HOLDER OR OTHER PARTY HAS BEEN ADVISED OF THE POSSIBILITY OF SUCH DAMAGES. 17. Interpretation of Sections 15 and 16.

If the disclaimer of warranty and limitation of liability provided above cannot be given local legal effect according to their terms, reviewing courts shall

apply local law that most closely approximates an absolute waiver of all civil liability in connection with the Program, unless a warranty or assumption of liability accompanies a copy of the Program in return for a fee.

END OF TERMS AND CONDITIONS How to Apply These Terms to Your New Programs

If you develop a new program, and you want it to be of the greatest possible use to the public, the best way to achieve this is to make it free software which everyone can redistribute and change under these terms.

To do so, attach the following notices to the program. It is safest to attach them to the start of each source file to most effectively state the exclusion of warranty; and each file should have at least the "copyright" line and a pointer to where the full notice is found.

<one line to give the program's name and a brief idea of what it does.> Copyright (C) <year> <name of author>

This program is free software; you can redistribute it and/or modify it under the terms of the GNU General Public License as published by the Free Software Foundation, either version 3 of the License, or (at your option) any later version.

This program is distributed in the hope that it will be useful, but WITHOUT ANY WARRANTY; without even the implied warranty of MERCHANTABILITY or FITNESS FOR A PARTICULAR PURPOSE. See the GNU General Public License for more details.

You should have received a copy of the GNU General Public License along with this program. If not, see <<http://www.gnu.org/licenses/>>.

Also add information on how to contact you by electronic and paper mail.

If the program does terminal interaction, make it output a short notice like this when it starts in an interactive mode:

```
<program> Copyright (C) <year> <name of author>
This program comes with ABSOLUTELY NO WARRANTY; for details type `show w'. This is free software, and you are welcome to redistribute it under certain conditions; type `show c' for details.
```

The hypothetical commands 'show w' and 'show c' should show the appropriate parts and the General Public License. Of course, your program's commands might be different; for a GUI interface, you would use an "about box".

You should also get your employer (if you work as a programmer) or school, if any, to sign a "copyright disclaimer" for the program, if necessary. For more information on this, and how to apply and follow the GNU GPL, see <<http://www.gnu.org/licenses/>>.

The GNU General Public License does not prevent incorporating your program into proprietary programs. If your program is a subroutine library, you may consider it more useful to permit linking proprietary applications with the library. If this is what you want to do, use the GNU Lesser General Public License instead of this License. But first, please read <<http://www.gnu.org/philosophy/why-not-lgpl.html>>.

If you publish printed copies (or copies in media that commonly have printed covers) of the Docu-

of the Document). You may use the same title as a previous version if the original publisher of that version gives permission. * B. List on the Title Page, as authors, one or more persons or entities responsible for authorship of the modifications in the Modified Version, together with at least five of the principal authors of the Document (all of its principal authors, if it has fewer than five), unless they release you from this requirement. * C. State on the Title page the name of the publisher of the Modified Version, as the publisher. * D. Preserve all the copyright notices of the Document. * E. Add an appropriate copyright notice for your modifications adjacent to the other copyright notices. * F. Include, immediately after the copyright notices, a license notice giving the public permission to use the Modified Version under the terms of this License, in the form shown in the Addendum below. * G. Preserve in that license notice the full lists of Invariant Sections and required Cover Texts given in the Document's license notice. * H. Include an unaltered copy of this License. * I. Preserve the section entitled "History", Preserve its Title, and add to it an item stating at least the title, year, new authors, and publisher of the Modified Version as given on the Title Page. If there is no section entitled "History" in the Document, create one stating the title, year, authors, and publisher of the Document as given on its Title Page, then add an item describing the Modified Version as stated in the previous sentence. * J. Preserve the network location, if any, given in the Document for public access to a Transparent copy of the Document, and likewise the network locations given in the Document for previous versions it was based on. These may be placed in the "History" section. You may omit a network location for a work that was published at least four years before the Document itself, or if the original publisher of the version it refers to gives permission. * K. For any section entitled "Acknowledgements" or "Dedications", Preserve the Title of the section, and preserve in the section all the substance and tone of each of the contributor acknowledgements and/or dedications given therein. * L. Preserve all the Invariant Sections of the Document, unaltered in their text and in their titles. Section numbers or the equivalent are not considered part of the section titles. * M. Delete any section entitled "Endorsements". Such a section may not be included in the Modified Version. * N. Do not retitle any existing section to be entitled "Endorsement" or to conflict in title with any Invariant Section. * O. Preserve any Warranty Disclaimers.

If the Modified Version includes new front-matter sections or appendices that qualify as Secondary Sections and contain no material copied from the Document, you may at your option designate some or all of these sections as invariant. To do this, add their titles to the list of Invariant Sections in the Modified Version's license notice. These titles must be distinct from any other section titles.

You may add a section entitled "Endorsements", provided it contains nothing but endorsements of your Modified Version by various parties—for example, statements of peer review or that the text has been approved by an organization as the authoritative definition of a standard.

You may add a passage of up to five words as a Front-Cover Text, and a passage of up to 25 words as a Back-Cover Text, to the end of the list of Cover Texts in the Modified Version. Only one passage of Front-Cover Text and one of Back-Cover Text may be added by (or through arrangements made by) any one entity, if the Document already includes a cover text for the same cover, previously added

by you or by arrangement made by the same entity you are acting on behalf of, you may not add another; but you may replace the old one, on explicit permission from the previous publisher that added the old one.

The author(s) and publisher(s) of the Document do not by this License give permission to use their names for publicity for or to assert or imply endorsement of any Modified Version. 5. COMBINING DOCUMENTS

You may combine the Document with other documents released under this License, under the terms defined in section 4 above for modified versions, provided that you include in the combination all of the Invariant Sections of all of the original documents, unmodified, and list them all as Invariant Sections of your combined work in its license notice, and that you preserve all their Warranty Disclaimers.

The combined work need only contain one copy of this License, and multiple identical Invariant Sections may be replaced with a single copy. If there are multiple Invariant Sections with the same name but different contents, make the title of each such section unique by adding at the end of it, in parentheses, the name of the original author or publisher of that section if known, or else a unique number. Make the same adjustment to the section titles in the list of Invariant Sections in the license notice of the combined work.

In the combination, you must combine any sections entitled "History" in the various original documents, forming one section entitled "History"; likewise combine any sections entitled "Acknowledgements", and any sections entitled "Dedications". You must delete all sections entitled "Endorsements". 6. COLLECTIONS OF DOCUMENTS

You may make a collection consisting of the Document and other documents released under this License, and replace the individual copies of this License in the various documents with a single copy that is included in the collection, provided that you follow the rules of this License for verbatim copying of each of the documents in all other respects.

You may extract a single document from such a collection, and distribute it individually under this License, provided you insert a copy of this License into the extracted document, and follow this License in all other respects regarding verbatim copying of that document. 7. AGGREGATION WITH INDEPENDENT WORKS

A compilation of the Document or its derivatives with other separate and independent documents or works, in or on a volume of a storage or distribution medium, is called an aggregate if the copyright resulting from the compilation is not used to limit the legal rights of the compilation's users beyond what the individual works permit. When the Document is included in an aggregate, this License does not apply to the other works in the aggregate which are not themselves derivative works of the Document.

If the Cover Text requirement of section 3 is applicable to these copies of the Document, then if the Document is less than one half of the entire aggregate, the Document's Cover Texts may be placed on covers that bracket the Document within the aggregate, or the electronic equivalent of covers if the Document is in electronic form. Otherwise

The "Corresponding Application Code" for a Combined Work means the object code and/or source code for the Application, including any data and utility programs needed for reproducing the Combined Work from the Application, but excluding the System Libraries of the Combined Work. 1. Exception to Section 3 of the GNU GPL.

You may convey a covered work under sections 3 and 4 of this License without being bound by section 3 of the GNU GPL. 2. Conveying Modified Versions.

If you modify a copy of the Library, and, in your modifications, a facility refers to a function or data to be supplied by an Application that uses the facility (other than as an argument passed when the facility is invoked), then you may convey a copy of the modified version:

* a) under this License, provided that you make a good faith effort to ensure that, in the event an Application does not supply the function or data, the facility still operates, and performs whatever part of its purpose remains meaningful, or * b) under the GNU GPL, with none of the additional permissions of this License applicable to that copy.

3. Object Code Incorporating Material from Library Header Files.

The object code form of an Application may incorporate material from a header file that is part of the Library. You may convey such object code under terms of your choice, provided that, if the incorporated material is not limited to numerical parameters, data structure layouts and accessors, or small macros, inline functions and templates (ten or fewer lines in length), you do both of the following:

* a) Give prominent notice with each copy of the object code that the Library is used in it and that the Library and its use are covered by this License. * b) Accompany the object code with a copy of the GNU GPL and this license document.

they must appear on printed covers that bracket the whole aggregate. 8. TRANSLATION

Translation is considered a kind of modification, so you may distribute translations of the Document under the terms of section 4. Replacing Invariant Sections with translations requires special permission from their copyright holders, but you may include translations of some or all Invariant Sections in addition to the original versions of these Invariant Sections. You may include a translation of this License, and all the license notices in the Document, and any Warranty Disclaimers, provided that you also include the original English version of this License and the original versions of those notices and disclaimers. In case of a disagreement between the translation and the original version of this License or a notice or disclaimer, the original version will prevail.

If a section in the Document is entitled "Acknowledgements", "Dedications", or "History", the requirement (section 4) to Preserve its Title (section 1) will typically require changing the actual title. 9. TERMINATION

You may not copy, modify, sublicense, or distribute the Document except as expressly provided under this License. Any attempt otherwise to copy, modify, sublicense, or distribute it is void, and will automatically terminate your rights under this License.

However, if you cease all violation of this License, then your license from a particular copyright holder is reinstated (a) provisionally, unless and until the copyright holder explicitly and finally terminates your license, and (b) permanently, if the copyright holder fails to notify you of the violation by some reasonable means prior to 60 days after the cessation.

Moreover, your license from a particular copyright holder is reinstated permanently if the copyright holder notifies you of the violation by some reasonable means, this is the first time you have received notice of violation of this License (for any work) from that copyright holder, and you cure the violation prior to 30 days after your receipt of the notice.

Termination of your rights under this section does not terminate the licenses of parties who have received copies or rights from you under this License. If your rights have been terminated and not permanently reinstated, receipt of a copy of some or all of the same material does not give you any rights to use it. 10. FUTURE REVISIONS OF THIS LICENSE

The Free Software Foundation may publish new, revised versions of the GNU Free Documentation License from time to time. Such new versions will be similar in spirit to the present version, but may differ in detail to address new problems or concerns. See <http://www.gnu.org/copyleft/>.

Each version of the License is given a distinguishing version number. If the Document specifies that a particular numbered version of this License or any later version applies to it, you have the option of following the terms and conditions either of that specified version or of any later version that has been published (not as a draft) by the Free Software Foundation. If the Document does not specify a version number of this License, you may choose

4. Combined Works.

You may convey a Combined Work under terms of your choice that, taken together, effectively do not restrict modification of the portions of the Library contained in the Combined Work and reverse engineering for debugging such modifications, if you also do each of the following:

* a) Give prominent notice with each copy of the Combined Work that the Library is used in it and that the Library and its use are covered by this License. * b) Accompany the Combined Work with a copy of the GNU GPL and this license document. * c) For a Combined Work that displays copyright notices during execution, include the copyright notice for the Library among these notices, as well as a reference directing the user to the copies of the GNU GPL and this license document. * d) Do one of the following: o 0) Convey the Minimal Corresponding Source under the terms of this License, and the Corresponding Application Code in a form suitable for, and under terms that permit, the user to recombine or relink the Application with a modified version of the Linked Version to produce a modified Combined Work, in the manner specified by section 6 of the GNU GPL for conveying Corresponding Source. o 1) Use a suitable shared library mechanism for linking with the Library. A suitable mechanism is one that (a) uses at run time a copy of the Library already present on the user's computer system, and (b) will operate properly with a modified version of the Library that is interface-compatible with the Linked Version. * e) Provide Installation Information, but only if you would otherwise be required to provide such information under section 6 of the GNU GPL, and only to the extent that such information is necessary to install and execute a modified version of the Combined Work produced by recombining or relinking the Application with a modified version of the Linked Version. (If you use option 4d, the Installation Information must accompany the Minimal Corresponding Source and Corresponding Application Code. If you use option 4e, you must provide the Installation Information in the manner specified by section 6 of the GNU GPL for conveying Corresponding Source.)

any version ever published (not as a draft) by the Free Software Foundation. If the Document specifies that a proxy can decide which future versions of this License can be used, that proxy's public statement of acceptance of a version permanently authorizes you to choose that version for the Document. 11. RELICENSING

"Massive Multiauthor Collaboration Site" (or "MMC Site") means any World Wide Web server that publishes copyrightable works and also provides prominent facilities for anybody to edit those works. A public wiki that anybody can edit is an example of such a server. A "Massive Multiauthor Collaboration" (or "MMC") contained in the site means any set of copyrightable works thus published on the MMC site.

"CC-BY-SA" means the Creative Commons Attribution-Share Alike 3.0 license published by Creative Commons Corporation, a not-for-profit corporation with a principal place of business in San Francisco, California, as well as future copyleft versions of that license published by that same organization.

Incorporate" means to publish or republish a Document, in whole or in part, as part of another Document.

An MMC is eligible for relicensing if it is licensed under this License, and if all works that were first published under this License somewhere other than this MMC, and subsequently incorporated in whole or in part into the MMC, (1) had no cover texts or invariant sections, and (2) were thus incorporated prior to November 1, 2008.

The operator of an MMC Site may republish an MMC contained in the site under CC-BY-SA on the same site at any time before August 1, 2009, provided the MMC is eligible for relicensing. ADDENDUM: How to use this License for your documents

To use this License in a document you have written, include a copy of the License in the document and put the following copyright and license notices just after the title page:

Copyright (C) YEAR YOUR NAME. Permission is granted to copy, distribute and/or modify this document under the terms of the GNU Free Documentation License, Version 1.3 or any later version published by the Free Software Foundation; with no Invariant Sections, no Front-Cover Texts, and no Back-Cover Texts. A copy of the license is included in the section entitled "GNU Free Documentation License".

If you have Invariant Sections, Front-Cover Texts and Back-Cover Texts, replace the "with ... Texts." line with this:

with the Invariant Sections being LIST THEIR TITLES, with the Front-Cover Texts being LIST, and with the Back-Cover Texts being LIST.

If you have Invariant Sections without Cover Texts, or some other combination of the three, merge those two alternatives to suit the situation.

If your document contains nontrivial examples of program code, we recommend releasing these examples in parallel under your choice of free software license, such as the GNU General Public License, to permit their use in free software.

12.3 GNU Lesser General Public License

GNU LESSER GENERAL PUBLIC LICENSE

Version 3, 29 June 2007

Copyright © 2007 Free Software Foundation, Inc. <<http://fsf.org/>>

Everyone is permitted to copy and distribute verbatim copies of this license document, but changing it is not allowed.

This version of the GNU Lesser General Public License incorporates the terms and conditions of version 3 of the GNU General Public License, supplemented by the additional permissions listed below. 0. Additional Definitions.

As used herein, "this License" refers to version 3 of the GNU Lesser General Public License, and the "GNU GPL" refers to version 3 of the GNU General Public License.

"The Library" refers to a covered work governed by this License, other than an Application or a Combined Work as defined below.

An "Application" is any work that makes use of an interface provided by the Library, but which is not otherwise based on the Library. Defining a subclass of a class defined by the Library is deemed a mode of using an interface provided by the Library.

A "Combined Work" is a work produced by combining or linking an Application with the Library. The particular version of the Library with which the Combined Work was made is also called the "Linked Version".

The "Minimal Corresponding Source" for a Combined Work means the Corresponding Source for the Combined Work, excluding any source code for portions of the Combined Work that, considered in isolation, are based on the Application, and not on the Linked Version.

If the Library as you received it specifies that a proxy can decide whether future versions of the GNU Lesser General Public License shall apply, that proxy's public statement of acceptance of any version is permanent authorization for you to choose that version for the Library.