

Embrenagra rufivirgata, *Lawr.* Nov.
 Calamospiza bicolor, *Bonap.* Nov.
 Euspiza americana, *Bonap.* Aug.
 Guiraca cærulea, *Sw.* May, June, Oct.
 Cyanospiza ciris, *Baird.* April to Aug. Abundant.
 Pyrrhuloxia sinuata, *Bonap.* April to Sept., Dec. to Feb. Abundant.
 Cardinalis virginianus, *Bonap.* May to Aug., Nov., Dec. Abundant.
 Pipilo chlorura, *Baird.* Dec., Jan.
 Molothrus pecoris, *Sw.* April to July, Jan. Abundant.
 Xanthocephalus icterocephalus. April, May.
 Sturnella neglecta, *Aud.* Oct. and Nov. Abundant.
 Sturnella magna, *Sw.* Dec.
 Icterus cucullatus, *Sw.* June.
 Icterus spurius, *Bonap.* June, Aug.
 Icterus bullockii, *Bonap.* May to Aug. Abundant.
 Scolecophagus cyanocephalus, *Sw.* Nov., Feb.
 Quiscalus macroura, *Sw.* June.
 Corvus cryptoleucus, *Couch.* Feb., March.
 Zenaidura carolinensis, *Bonap.* April to July.
 Scardafella inca, *Bonap.* May, Sept. to Nov., Dec., Feb., March. Abundant.
 Ortyx texanus, *Lawr.* Oct., Nov., Dec.
 Callipepla squamata, *Gray.* April to Nov., Feb. Abundant.
 Grus canadensis, *Temm.* Jan.
 Ardea herodias, *Linn.* Jan.
 Nyctherodius violaceus, *Reich.* Sept.
 Aegialitis vociferus, *Cass.* April, May, Aug., Sept.
 Tringa wilsonii, *Nuttall.* Oct., Dec.
 Ereunetes petrificatus, *Ill.*
 Actiturus bartramius, *Bonap.* Aug.
 Tringites rufescens, *Cab.*
 Pelecanus fuscus, *Linn.* May.

—

Descriptions of four new species of Exotic UNIONIDÆ.

BY ISAAC LEA.

ANODONTA STREBELII.—Testa lævi, elliptica, inæquilaterali, postice obtuse angulata, antice rotundata; valvulis pertenuibus; natibus prominulis; epidermide tenebroso-viridi, radiis capillaris indutis; margarita cærulea et valde iridescente.

Hab.—Vera Cruz, Mexico; Dr. G. Strebel, per Smithsonian Institution.

UNIO VERACRUZENSIS.—Testa lævi, elliptica, subcompressa, inæquilaterali, postice obtuse angulata, antice rotundata; valvulis tenuibus; natibus prominulis; epidermide tenebroso-fusca, radiata politaque; dentibus cardinalibus parvis, compressis, crenulatis, in utroque valvulo duplicibus; lateralibus longis, rectis lamellatisque; margarita cærulea et valde iridescente.

Hab.—Vera Cruz, Mexico; Dr. G. Strebel, per Smithsonian Institution.

UNIO PRUNOIDES.—Testa lævi, elliptica, valde ventricosa, inæquilaterali, antice et postice rotundata; valvulis subcrassis, antice aliquanto crassioribus; natibus prominulis; epidermide tenebroso-fusca, eradiata; dentibus cardinalibus compressis, obliquis et valde crenulatis; lateralibus longis, lamellatis corrugatisque; margarita argentea.

UNIO CHINENSIS.—Testa nodulosa, subelliptica, inflata, fere æquilaterali, postice obtuse angulata, antice rotundata; valvulis crassiusculis, antice crassioribus; natibus subprominentibus, ad apices corrugatis; epidermide virido-lutea, radiis viridis undique indutis; dentibus cardinalibus erectis, compressis, striatis, crenulatis et in utroque valvulo subduplicibus; lateralibus sublongis subrectisque; margarita argentea et valde iridescente.

[May,