## II. Principles of Public Health (17 Hrs.)

| | | | |
|---|---|---|---|
| 1. | Outline of Public Health ..... | 2 | Saito |
| 2. | Health Center Activities ..... | 1.5 | Kimura |
| 3. | Food Control Administration ..... | 3 | Asomura |
| 4. | Communicable Disease Control ..... | 1.5 | Banba |
| 5. | T. B. Control ..... | 1.5 | Tsuda |
| 6. | Animal Infection Disease Control ..... | 3 | Tanaka |
| 7. | Health Education ..... | 1.5 | Shiga |
| 8. | Problems of the Budget ..... | 3 | Hinoue |

## III. Public Health Statistics (12 Hrs.)

| | | Lecture | Practice | |
|---|---|---|---|---|
| 1. | Complication of raw data ..... | 1 | 2 | Kawakami Shindo |
| 2. | Average, ratio and rates ..... | 1 | 2 | " |
| 3. | Sampling, Variation, measures of Unreliability ..... | 1 | 2 | " |
| 4. | Other health data ..... | 1 | 2 | Kawakami Shindo |

## IV. Epidemiology and Bacteriology (24 Hrs.)

| | | | | |
|---|---|---|---|---|
| 1. | Epidemiology introduction ..... | 3 | - | Kubo |
| 2. | Bacteriology and immunology introduction ..... | 3 | - | Someya |
| 3. | Bacteriology and immunology ..... | 12 | - | Ochi |
| 4. | Food Poisoning ..... | 6 | - | Hatta Kinbara |



DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 773013

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 773 013

V. Nutrition (6 Hrs.)

| | Lecture | Practice | |
|---|---|---|---|
| 1. Administration of nutrition ..... | 3 | - | Arimoto |
| 2. Nutrition General ..... | 3 | - | Nakagawa |

VI. Meat Hygiene (33 Hrs.)

| | | | |
|---|---|---|---|
| 1. Administrative aspects of slaughtering and of law relating processing plant of dead animal and others ..... | 3 | - | Asomura |
| 2. Sanitation of Slaughter house ..... | 3 | - | Ikeda |
| 3. Abnormal condition and disease of slaughter animals ..... | 15 | - | Yamamoto |
| 4. Examination of slaughter animals ..... | 3 | - | Dr. Pegg |
| 5. Production of food meat products and treatment of by-products from slaughtering ..... | 3 | - | Sasaki |
| 6. Examination of meat and meat products ..... | - | 6 | Matsui<br>Nakano |

VII. Milk Hygiene (24 Hrs.)

| | | | |
|---|---|---|---|
| 1. Feeding and management of dairy animals ..... | 3 | - | Okoshi |
| 2. Bacteriology and Chemistry of milk ..... | 6 | 9 | Kawasaki<br>Matsui<br>Nakano<br>Miyao |
| 3. Manufacturing of milk products and sanitation of those factory ..... | 3 | | Tsugo |
| 4. Inspection of dairy farm and milk plant ..... | 3 | | Matsui |

VIII. Sea food Hygiene (39 Hrs.)

| | | | |
|---|---|---|---|
| 1. Administration of sea food ..... | 3 | - | Asomura |

| | | | |
|---|---|---|---|
| 5. Production of food meat products and treatment of by-products from slaughtering ..... | 3 | - | Sasaki |
| 6. Examination of meat and meat products ..... | - | 6 | Matsui<br>Nakano |

VII. Milk Hygiene (24 Hrs.)

| | | | |
|---|---|---|---|
| 1. Feeding and management of dairy animals ..... | 3 | - | Okoshi |
| 2. Bacteriology and Chemistry of milk ..... | 6 | 9 | Kawasaki<br>Matsui<br>Nakano<br>Miyao |
| 3. Manufacturing of milk products and sanitation of those factory ..... | 3 | | Tsugo |
| 4. Inspection of dairy farm and milk plant ..... | 3 | | Matsui |

VIII. Sea food Hygiene (39 Hrs.)

| | | | |
|---|---|---|---|
| 1. Administration of sea food ..... | 3 | - | Asomura |
| 2. Sea food General ..... | 3 | - | Hiyama |
| 3. Bacteriology and Chemistry of sea food ..... | 15 | 12 | Amano<br>Toyama<br>Matsui<br>Nakano<br>Miyao |
| 4. Sanitation of Fresh fish and sea food products . | 3 | - | Dr. Beehwood |
| 5. Construction of sea food handling establishment & equipment ..... | 1.5 | - | |



DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 77 013

| | | Lecture | Practice | |
|---|---|---|---|---|
| 6. | Sanitation of sea food mfg. Plants ..... | 1.5 | - | |

IX. Food Inspection (6 Hrs.)

| | | | | |
|---|---|---|---|---|
| 1. | Food inspection system ..... | 3 | - | Kitaura |
| 2. | General food Inspection Practice ..... | 3 | - | |

X. Rabies Control (6 Hrs.)

| | | | | |
|---|---|---|---|---|
| 1. | Diagnosis and Vaccination of Rabies ..... | 3 | 3 | Kondo |

XI. ~~Sanitary Engineering~~ SANITATION (24 Hrs.)

| | | | | |
|---|---|---|---|---|
| 1. | Water ..... | 3 | - | Sago |
| 2. | Sewage ..... | 3 | 3 | Yorasawa |
| 3. | Theory and application of refrigeration ..... | 3 | 3 | Wakuri |
| 4. | Insect and rodent control ..... | 3 | 3 | Yamaguchi |
| 5. | Principles of detergents ..... | 3 | - | Motoyama |

XII. Field Practice (42 Hrs.)

| | | Hours |
|---|---|---|
| 1. | Examination of slaughter animal at slaughter house .... | 12 |
| 2. | Inspection of sea food ..... | 12 |
| 3. | Practical aspects of inspection (Health Center) ..... | 9 |
| 4. | Dairy farm & milk plant Inspection ..... | 9 |

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775013

The Institute of Public Health
6th 3-month Course for Sanitary Engineers
(April 10 - June 30, 1950)

Subjects and Hours

| | |
|---|---|
| A - Introductory Lectures | 6 |
| B - Principles of Public Health and its Administration | 7 |
| C - Vital Statistics | 12 |
| D - Administration and Management of Sanitary Engineering | 12 |
| E - Outline of Epidemiology and Infectious Diseases | 12 |
| F - Water and Sewage Works Engineering | 60 |
| G - Water Treatment | 33 |
| H - Sewage Treatment and its Utilization | 24 |
| I - Water and Sewage Examination | 87 |
| J - Night Soil: Collection, Treatment and Utilization | 9 |
| K - Refuse: Collection, Treatment and Utilization | 6 |
| L - Insect and Rodent Control | 21 |
| M - Industrial Sanitation | 5 |
| N - Architectural Hygiene | 15 |
| O - Food Sanitation | 3 |
| P - Field Practice | 30 |
| Q - Discussion | 24 |
| Total | 366 |

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775013

| | Lect. | Pract. | Total | Instructor |
|---|---|---|---|---|
| A - Introductory Lectures. | 6 | 0 | 6 | |
| 1) Public health and sanitary engineering. | (2) | (0) | (2) | McCoy |
| 2) General remarks to sanitary engineering | (2) | (0) | (2) | Maclaren |
| 3) Duties of sanitary engineers. | (1) | (0) | (1) | Tanaka |
| 4) On the course of sanitary engineering. | (1) | (0) | (1) | Horasawa |
| B - Principles of Public Health and its Administration. | 7 | 0 | 7 | |
| 1) Outline of public health. | | | | |
| a - Introduction of public health. | (1) | (0) | (1) | Saito |
| b - Outline of environmental hygiene. | (3) | (0) | (3) | Matsuoka |
| 2) Outline of public health administration | (3) | (0) | (3) | Shiga |
| C - Vital Statistics. | 4 | 8 | 12 | Kawakami |
| D - Administration and Management of Sanitary Engineering. | 12 | 0 | 12 | |
| 1) Administration and management of water and sewerage. | | | | |
| a - Administration of water works and sewerage. | (3) | (0) | (3) | Tanaka |
| b - Management of water and sewage works | (6) | (0) | (6) | Kawaguchi |
| 2) Administration of environmental sanitation. | (3) | (0) | (3) | Matsuda |
| E - Outline of Epidemiology and Infectious Diseases. | 13 | 0 | 13 | |
| 1) Generals of epidemiology | (2) | (0) | (2) | Kubo |
| 2) Disinfection. | (3) | (0) | (3) | Someya |
| 3) Epidemiology and control of intestinal infectious diseases. | (3) | (0) | (3) | Shigematsu |

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775013

| | | | | |
|---|---|---|---|---|
| 2) Outline of public health administration | (3) | (0) | (3) | Shiga |
| C - Vital Statistics. | 4 | 8 | 12 | Kawakami |
| D - Administration and Management of Sanitary Engineering. | 12 | 0 | 12 | |
| 1) Administration and management of water and sewerage. | | | | |
| a - Administration of water works and sewerage. | (3) | (0) | (3) | Tanaka |
| b - Management of water and sewage works | (6) | (0) | (6) | Kawaguchi |
| 2) Administration of environmental sanitation. | (3) | (0) | (3) | Matsuda |
| E - Outline of Epidemiology and Infectious Diseases. | 13 | 0 | 13 | |
| 1) Generals of epidemiology | (2) | (0) | (2) | Kubo |
| 2) Disinfection. | (3) | (0) | (3) | Someya |
| 3) Epidemiology and control of intestinal infectious diseases. | (3) | (0) | (3) | Shigematsu |
| a - Water-born diseases. | | | | |
| b - Food-born diseases. | | | | |
| 4) Epidemiology and control of insect-born diseases. | (2) | (0) | (2) | Hirayama |
| 5) Administration of infectious diseases | (3) | (0) | (3) | Banba |



DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 773 013

| | Lect. | Pract. | Total | Instructor |
|---|---|---|---|---|
| F - Water and Sewage Works Engineering | 34 | 26 | 60 | |
| 1) a - Outline of Water Works engineering. | (6) | (0) | (6) | Iwai |
| b - Outline of sewage works engineering. | (3) | (0) | (3) | Sago |
| c - Control of leakage | (3) | (3) | (6) | Iwai |
| d - Pump | (3) | (3) | (6) | Ishida |
| e - Piping | (3) | (0) | (3) | Ikeda |
| f - Concrete pipe | (3) | (3) | (6) | Itokura |
| g - Iron pipe | (6) | (3) | (9) | Ikeda |
| h - Meter | (1) | (2) | (3) | do. |
| 2) Hydraulics. | (6) | (3) | (9) | Sago. |
| 3) Plumbing. | (3) | (6) | (9) | Sato |
| G - Water Treatment | 21 | 12 | 33 | |
| 1) Sedimentation | (6) | (0) | (6) | Sago |
| 2) Sand filtration | (9) | (0) | (9) | Hirose |
| 3) Disinfection and control of algae | (3) | (3) | (6) | Horasawa |
| 4) Inspection tour to chlorinator manufacturing company | (0) | (3) | (3) | Iwato |
| 5) Operation | (0) | (6) | (6) | Matsumoto |
| (app.) Swimming pool, bath and hot spring | (3) | (0) | (3) | Iwato |
| H - Sewage Treatment and its Utilization. | 18 | 6 | 24 | |
| 1) Sewage and public health. | (15) | (0) | (15) | Horasawa |
| 2) Composition of sewage and its characteristics. | | | | |
| a - Noticeable substance and organisms. | | | | |
| b - Decomposition and its significance on water. | | | | |

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 773013

| | | | | |
|---|---|---|---|---|
| 3) Plumbing. | (3) | (6) | (9) | Sato |
| G - Water Treatment | 21 | 12 | 33 | |
| 1) Sedimentation | (6) | (0) | (6) | Sago |
| 2) Sand filtration | (9) | (0) | (9) | Hirose |
| 3) Disinfection and control of algae | (3) | (3) | (6) | Horasawa |
| 4) Inspection tour to chlorinator manufacturing company | (0) | (3) | (3) | Iwato |
| 5) Operation | (0) | (6) | (6) | Matsumoto |
| (app.) Swimming pool, bath and hot spring | (3) | (0) | (3) | Iwato |
| H - Sewage Treatment and its Utilization. | 18 | 6 | 24 | |
| 1) Sewage and public health. | (15) | (0) | (15) | Horasawa |

2) Composition of sewage and its characteristics.

a - Noticeable substance and organisms.

b - Decomposition and its significance on water.

3) Significance of sewage composition upon the treatment.

4) Sewage treatment.

a - Dilution..

b - Irrigation.

c - Preliminary treatment.

d - Biological treatment.



DECLASSIFIED E.O. 12065 SECTION 3-402/NNDC NO. 773 013

| | | | | |
|---|---|---|---|---|
| 5) Sewage sludge. | | | | |
| a - Composition of sludge. | | | | |
| b - Treatment of sewage sludge. | | | | |
| 6) Utilization of sewage. | | | | |
| 7) Maintenance and inspection. | | | | |
| 8) Induatrial waste. | (3) | (0) | (3) | Sibata |
| 9) Operation. | (0) | (6) | (6) | Nakajima |
| I - Water and Sewage Examination. | 42 | 45 | 87 | |
| 1) General principle of assaying water examination results. | (3) | (0) | (3) | Horasawa |
| 2) Chemical eremination of water and sewage. | | | | |
| a - Outline of water and sewage chemistry. | (6) | (0) | (6) | Shibata |
| b - Examination of drinking water. | | | | |
| Theory ..... | (9) | (0) | (9) | Aizawa |
| Laboratory ..... | (0) | (15) | (15) | |
| c - Examination of sewage. | (9) | (12) | (21) | Shibata Hagiwara |
| 3) Bacteriological examination of water and sewage. | (9) | (12) | (21) | Iwato |
| a - Outline of bacteriology of water and sewage. | | | | |
| b - Examination of water and sewage. | | | | |
| 4) Biological examination of water and sewage. | (6) | (6) | (12) | Horasawa Nakajima |
| a - Outline of biology of water and sewage. | | | | |
| b - Microscopical examination of water and sewage. | | | | |

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 743 013

| | | | | |
|---|---|---|---|---|
| Laboratory ..... | (0) | (15) | (15) | |
| c - Examination of sewage. | (9) | (12) | (21) | Shibata<br>Hagiwara |
| 3) Bacteriological examination of water and sewage. | (9) | (12) | (21) | Iwato |
| a - Outline of bacteriology of water and sewage. | | | | |
| b - Examination of water and sewage. | | | | |
| 4) Biological examination of water and sewage. | (6) | (6) | (12) | Horasawa<br>Nakajima |
| a - Outline of biology of water and sewage. | | | | |
| b - Microscopical examination of water and sewage. | | | | |
| J - Night soil : Collection, Treatment and Utilization | 6 | 3 | 9 | |
| 1) Night soil and public health. | (6) | (0) | (6) | Gonda |
| 2) Privy. | | | | |
| a - Sanitary privy. | | | | |
| b - Cess-pool of Japanese pattern. | | | | |
| c - Water closet. | | | | |
| d - Public privy. | | | | |



DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775013

| | | | | |
|---|---|---|---|---|
| 3) Collection and transportation. | | | | |
| 4) Treatment. | | | | |
| 5) Utilization. | | | | |
| 6) Operation. | (0) | (3) | (3) | Miyao<br>Matsumoto |
| K - Refuse: Collection, Treatment and Utilization | 3 | 3 | 6 | |
| 1) Refuse and public health. | (3) | (0) | (3) | Gonda |
| 2) Refuse bin. | | | | |
| 3) Collection and transportation. | | | | |
| 4) Treatment. | | | | |
| 5) Utilization. | | | | |
| 6) Operation. | (0) | (3) | (3) | Miyao<br>Matsumoto |
| L - Insect and Rodent Control. | 9 | 12 | 21 | |
| 1) Arthropods and redents of medical importance. | (3) | (3) | (6) | Horasawa |
| a - Taxomomy and habits. | | | | |
| b - Significance on public health. | | | | |
| 2) Insecticide and rodenticide. | (3) | (0) | (3) | Suenaga<br>Motoyama |
| 3) Rodent control: Practice. | (0) | (3) | (3) | Gonda |
| 4) Insect control. | | | | |
| a - Outline. | (3) | (0) | (3) | Wheeler |
| b - Practice; | (0) | (6) | (6) | Yamaguchi<br>Matsumoto |
| M - Industrial Sanitation. | 5 | 0 | 5 | Suzuki |
| 1) The scope of industrial sanitation. | | | | |

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 773 013

| | | | | |
|---|---|---|---|---|
| 1) Arthropods and rodents | | | | |
| a - Taxomomy and habits. | | | | |
| b - Significance on public health. | | | | |
| 2) Insecticide and rodenticide. | (3) | (0) | (3) | Suenaga<br>Motoyama |
| 3) Rodent control: Practice. | (0) | (3) | (3) | Gonda |
| 4) Insect control. | | | | |
| a - Outline. | (3) | (0) | (3) | Wheeler |
| b - Practice) | (0) | (6) | (6) | Yamaguchi<br>Matsumoto |
| | 5 | 0 | 5 | Suzuki |

M - Industrial Sanitation.

1) The scope of industrial sanitation.

2) General standard.

3) Control of dust, fumes, and toxic substances.

4) Special industrial sanitation problems.

N - Architectural Hygiene.

1) General description.

2) Housing problem and town planning.

3) Housing standards.

4) Standards of residental districts.

5) Clearance and re-development areas.



DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775013

| | | | | |
|---|---|---|---|---|
| O - Food Sanitation. | 3 | 0 | 3 | Matsui |
| P - Field Practice. | 0 | 30 | 30 | |
| 1) Metropolitan water works reservoir. | | | | Nakajima |
| 2) Yodobashi filter plant. | | | | Tanaka, Sato |
| 3) Kanamachi filter plant. | | | | Takemae, Uchida |
| 4) Tamagawa filter plant. | | | | Okumura, Takeda |
| 5) Sakai filter plant. | | | | Yamada, Funatsu |
| 6) Shibaura sewage disposal plant. | | | | Ishizaka, Nakajima |
| 7) Mikawashima sewage disposal plant. | | | | Shibachi, Sawada |
| 8) Metropolitan water works bureau. | | | | Sato, Iwasaki |
| Q - Discussion. | 32 | 0 | 32 | |

Total ..... 366

Lecture ..... 215

Practice ..... 151

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 773013

Contents of Medical Officers' Course
(July 10 - Sept. 27, 1950)

| | Lecture | | Laboratory Practice | | Field Trips | | Others | | Total | |
|---|---|---|---|---|---|---|---|---|---|---|
| | Units | Hrs. | Units | Hrs. | Units | Hrs. | Units | Hrs. | Units | Hrs. |
| Introduction & General lectures | 6 | 9.0 | | | | | 5 | 8.0 | 11 | 17.0 |
| P.H. Statistics | 11 | 13.0 | 7 | 17.5 | | | | | 18 | 30.5 |
| Epidemiology | 2 | 6.0 | 1 | 3.0 | | | | | 3 | 9.0 |
| Comm. & Parasitic diseases | 24 | 45.0 | 10 | 30.0 | 2 | 9.5 | | | 36 | 84.5 |
| Environmental Sanitation and housing | 10 | 20.5 | 2 | 6.0 | 2 | 6.0 | | | 14 | 32.5 |
| Physiological & Industrial Hygiene | 6 | 12.0 | 5 | 15.0 | 1 | 6.5 | | | 12 | 33.5 |
| Biochemistry & Pharmacology | 3 | 4.0 | 1 | 2.0 | | | | | 4 | 6.0 |
| Nutrition & Food Sanitation | 4 | 9.0 | 1 | 3.0 | | | | | 5 | 12.0 |
| P. H. Administration | 34 | 54.0 | | | 1 | 3.0 | | | 35 | 57.0 |
| Maternal Child & School Hygiene | 8 | 14.5 | 1 | 3.0 | 3 | 19.5 | | | 12 | 37.0 |
| P. H. Nursing | 2 | 3.5 | | | | | | | 2 | 3.5 |
| Health Education | 1 | 2.0 | 1 | 3.0 | | | | | 2 | 5.0 |
| Psychology & Mental Hygiene | 4 | 9.0 | | | 1 | 6.5 | | | 5 | 15.5 |
| P.H. Demography | 6 | 11.5 | | | | | | | 6 | 11.5 |
| Field Work in Health Center | | | | | 5 | 32.5 | | | 5 | 32.5 |
| | | | | | | | 4 | 6.0 | 4 | 6.0 |

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 773 013

| | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|
| Industrial Hygiene | 6 | 12.0 | [illegible] | [illegible] | | | | | 4 | 6.0 |
| Biochemistry & Pharmacology | 3 | 4.0 | 1 | 2.0 | | | | | 5 | 12.0 |
| Nutrition & Food Sanitation | 4 | 9.0 | 1 | 3.0 | | | | | 35 | 57.0 |
| P. H. Administration | 34 | 54.0 | | | 1 | 3.0 | | | 12 | 37.0 |
| Maternal Child & School Hygiene | 8 | 14.5 | 1 | 3.0 | 3 | 19.5 | | | 2 | 3.5 |
| P. H. Nursing | 2 | 3.5 | | | | | | | 2 | 5.0 |
| Health Education | 1 | 2.0 | 1 | 3.0 | | | | | 5 | 15.5 |
| Psychology & Mental Hygiene | 4 | 9.0 | | | 1 | 6.5 | | | 6 | 11.5 |
| P.H. Demography | 6 | 11.5 | | | | | | | 5 | 32.5 |
| Field Work in Health Center | | | | | 5 | 32.5 | | | 4 | 6.0 |
| Seminars | | | | | | | 4 | 6.0 | 8 | 24.0 |
| Elected Study | | | | | | | 8 | 24.0 | | |
| Total | 121 | 213.0 | 29 | 82.5 | 15 | 83.5 | 17 | 38.0 | 182 | 417.0 |

* Nine lectures totaling 15.5 hours are giving by G.H.Q. personnel

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 773 013

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775013

Institute of Public Health
Schedule of Courses
April 1950 to March 1951

1) Medical Health Officers --- Regular Course
10 April 1950 --- 28 March 1951

2) Medical Health Officers --- 3 months course

| | | |
|---|---|---|
| 1st | 10 April - 30 June | 1950 |
| 2nd | 10 July - 27 September | " |
| 3rd | 2 October - 20 December | " |
| 4th | 8 January - 28 March | 1951 |

3) Sanitarians --- 3 months

| | | |
|---|---|---|
| 1st | 10 April - 30 June | 1950 |
| 2nd | 10 July - 27 September | " |
| 3rd | 2 October - 20 December | " |
| 4th | 8 January - 28 March | 1951 |

4) Public Health Veterinarians --- 2 months

| | | |
|---|---|---|
| 1st | 27 February - 22 April | 1950 |
| 2nd | 3 July - 26 August | " |
| 3rd | 4 September - 28 October | " |
| 4th | 4 December - 27 January | 1951 |

5) Public Health Nutritionists --- 2 months

| | | |
|---|---|---|
| 1st | 27 February - 22 April | 1950 |
| 2nd | 29 May - 22 July | " |
| 3rd | 4 September - 28 October | " |
| 4th | 4 December - 27 January | 1951 |

6) Public Health Nurses --- 4 months

| | | |
|---|---|---|
| 1st | 3 April - 29 July | 1950 |
| 2nd | 5 August - 25 November | " |
| 3rd | 2 December - 24 March | 1951 |

7) Sanitary Engineers --- 3 months

| | | |
|---|---|---|
| 1st | 10 April - 30 June | 1950 |
| 2nd | 8 January - 28 March | 1951 |

8) Public Health Statistics Officers --- 2 months

| | | |
|---|---|---|
| 1st | 3 July - 26 August | 1950 |
| 2nd | 4 September - 28 October | " |
| 3rd | 8 January - 3 March | 1951 |

9) Short Course for the directors of Prefectural Health Department Laboratories --- 2 weeks
1 May - 13 May 1950

10) Bacteriology, Serology and Parasitology --- 3 months

4th 8 January - 28 March 1951

3) Sanitarians --- 3 months
1st 10 April - 30 June 1950
2nd 10 July - 27 September "
3rd 2 October - 20 December "
4th 8 January - 28 March 1951

4) Public Health Veterinarians --- 2 months
1st 27 February - 22 April 1950
2nd 3 July - 26 August "
3rd 4 September - 28 October "
4th 4 December - 27 January 1951

5) Public Health Nutritionists --- 2 months
1st 27 February - 22 April 1950
2nd 29 May - 22 July "
3rd 4 September - 28 October "
4th 4 December - 27 January 1951

6) Public Health Nurses --- 4 months
1st 3 April - 29 July 1950
2nd 5 August - 25 November "
3rd 2 December - 24 March 1951

7) Sanitary Engineers --- 3 months
1st 10 April - 30 June 1950
2nd 8 January - 28 March 1951

8) Public Health Statistics Officers --- 2 months
1st 3 July - 26 August 1950
2nd 4 September - 28 October "
3rd 8 January - 3 March 1951

9) Short Course for the directors of Prefectural Health Department Laboratories --- 2 weeks
1 May - 13 May 1950

10) Bacteriology, Serology and Parasitology --- 3 months
15 May - 5 August

11) Clinical Microscopy and Pathology --- 2 months
4 September - 28 October

12) Public Health Pharmacists --- 2 months
1st 3 April - 27 May
2nd 30 October - 23 December

Note : Courses 10, 11 and 12 are for personnel employed in prefectural health department laboratories.

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775013

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775 013

SECURITY DIVISION
TOKYO QUARTERMASTER DEPOT
APO 1051

29 May 1950

The Tokyo Public Health Bureau is authorized to carry supplies in and out of ALL CHECK POINTS, Tokyo QM Depot. They are engaged insect rodent control work and will carry some American insecticides and equipment. THIS IS NOT A DEPOT PASS. IT WILL REMAIN IN POSSESSION OF DRIVER AT ALL TIMES.

Ennis E Boyd

ENNIS E BOYD
1st Lt. CMP
Security Division

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775 013

SECURITY DIVISION
TOKYO QUARTERMASTER DEPOT
APO 1051

29 May 1950

The Tokyo Public Health Bureau is authorized to carry supplies in and out of ALL CHECK POINTS, Tokyo QM Depot. They are engaged insect rodent control work and will carry some American insecticides and equipment. THIS IS NOT A DEPOT PASS. IT WILL REMAIN IN POSSESSION OF DRIVER AT ALL TIMES.

Ennis E Boyd

ENNIS E BOYD
1st Lt. CMP
Security Division

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775 013

SECURITY DIVISION
TOKYO QUARTERMASTER DEPOT
APO 1051

29 May 1950

The Tokyo Public Health Bureau is authorized to carry supplies in and out of ALL CHECK POINTS, Tokyo QM Depot. They are engaged, insect rodent control work and will carry some American insecticides and equipment. THIS IS NOT A DEPOT PASS. IT WILL REMAIN IN POSSESSION OF DRIVER AT ALL TIMES.

Ennis E Boyd
ENNIS E BOYD
1st Lt. CMP
Security Division

昭和25年5月29日

東京補給係倉庫保安部(APO 1051)

東京都衛生局には東京補給係倉庫の全ての検問所を通って資材を持込むこと、持ち去ることの許可が與えられてゐる。
彼等は昆虫鼠族駆除作業に従事しており、アメリカ製の殺虫剤や器具を運ぶであらう。これは倉庫証ではない。
常に運転手が所持し居るものである。

CMP 保安部
一等中尉
エニスE ボイド

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775 013

GENERAL HEADQUARTERS
SUPREME COMMANDER FOR THE ALLIED POWERS
Public Health and Welfare Section

# B U L L E T I N

No. 174

For Period

15th - 28th February

1951


| SECTION | | PAGE |
|---|---|---|
| I | GENERAL | 1 |
| II | PREVENTIVE MEDICINE | 2 |
| III | NURSING AFFAIRS | 2 |
| IV | VETERINARY AFFAIRS | 2 |
| V | SUPPLY | 5 |
| VI | NARCOTIC CONTROL | 6 |
| VII | WELFARE | 7 |

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775 013

## SECTION I GENERAL

Effective 1 March, publication of the Public Health and Welfare Section Bulletin will be on a monthly basis. The next issue will contin a summary of activities and reports for the month of March 1951.

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775 013

## SECTION II PREVENTIVE MEDICINE

### Smallpox

Following a year in which only five scattered smallpox cases were reported in Japan, an outbreak involving approximately 10 persons occurred, starting 7 February in a Military Laundry in Kobe. The first crop of cases had onsets varying from the 7th to the 13th. Four other suspect cases with onsets up to the 24th of February are under observation. The route of infection has not as yet been clearly determined. All efforts to find a possible preceding case have failed. An intensive immunization campaign is being conducted in Kobe and vicinity and has been in a large part completed.

In addition to the Kobe incident one case of smallpox has been reported in a child from Nagoya. In this case the diagnosis is extremely questionable. The onset was 11 February. Preventive measures have been taken and there have been no secondary cases.

On 22 February word was received that an epidemic of suspected smallpox involving 12 people, of which seven are infants, had been going on since 28 January in Kamesaki-mura, Murakata-gun, Fukuoka-ken. There was no doctor in this village and the cases came to the attention of the health authorities by accident. Further investigation is being conducted.

## SECTION III NURSING AFFAIRS

### National Nurses Examination

Attached as Incls. 1 is a copy of Ministry of Welfare Notification No. 24, dated 23 Feb 1951, concerning the 1951 schedule for application for the national examination for class A nurses.

## SECTION IV VETERINARY AFFAIRS

### Rabies Control Program

A nation-wide survey of enforcement measures taken by prefectural officials since the inception of the Rabies Control Law No. 247, 26 August 1950 up to 31 December 1950 revealed the fact that 855,830 dogs had been registered, and 847,503 dogs immunized. In addition 786 dog catchers had been employed and 339 dog pounds put into operation. While the results so far are considered commendable, the program is aimed at having the estimated total of two million dogs registered and immunized immediately. A total of 800 dog pounds, one for each health center district, with the permanent employment of at least 1,600 dog catchers is proposed. An effort is also being made to obtain transportation for the dog catchers in the form of foot driven or motor driven tricycles with cage carriers attached.

Although the Kanto region has been the seat of rabies infection, three cases were recently reported in Hyogo prefecture. These are the first cases reported in this prefecture and only the second time that rabies has been confirmed, outside of the Kanto region, since the beginning of the occupation.

### Rinderpest Chick Embryo Vaccine Research

Work on the adaptation of bovine rinderpest virus to the chick embryo is progressing favorably under the direction of Dr. Kanematsu Fukusho, Chief of the Kyushu Branch of the National Experimental Station for Animal Hyginne. The virus is now in the 130th C.A. membrance passage, and separately in the 53rd yolk sac passage. The C.A. membrane-passaged virus seems to be gradually attenuated, in consequence of longer period of time between inoculation and death, and lessened clinical symptoms in test calves. The test calf inoculated with the 119th passage virus recovered after showing typical sumptoms.

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775 013

The present yolk sac strain was passaged into a bovine at the 88 C.A. membrane passage, then the bovine's lymphnode virus passaged back once more into the C.A. membrance and then into the yolk sac. Up to the present 53rd yolk sac generation, the virus has faded out in yolk sac passage three times, but has been brought back each time by one alternate passage in the bovine. Japanese calves inoculated with the virus at the 30th and 35th passages died of pneumonia complications, but those inoculated at the 40th, 45th passages all survived. Holstein calves inoculated with every fifth passage between 30th and 50th passage have all recovered. The virus is still considered too virulent for use as a field vaccine, but the virulence seems already to be lower than that of lapinized virus. The work is being continued, and it is believed that it should result in the production of a practical vaccine after 25 - 50 more yolk sac passages.

Weekly Rabies Report

The Veterinary Affairs Section, Ministry of Welfare submitted the following report on rabies cases occurring during the period 28 January - 17 February 1951.

| Prefecture | No. of Cases (28 Jan-3 Feb) | No. of Cases (4 - 10 Feb) | No. of Cases (11 - 17 Feb) |
|---|---|---|---|
| Tokyo | 1 | 4 | 0 |
| Kanagawa | 3 | 3 | 1 |
| Chiba | 0 | 0 | 0 |
| Saitama | 0 | 0 | 0 |
| Ibaragi | 5 | 3 | 1 |
| Gumma | 1 | 2 | 2 |
| Tochigi | 0 | 2 | 0 |
| Shizuoka | 1 | 0 | 0 |
| Hyogo | 0 | 1 | 1 |
| Total | 11 | 15 | 5 |

Japanese statistics for period of 1 January - 17 February 1951 disclose a total of 48 cases of rabies in dogs, 137 humans have reported being bitten by dogs with 5 deaths.

Weekly Animal Disease Report

The Animal Hygiene Section, Ministry of Agriculture and Forestry, submitted the following outbreaks of animal diseases for the period 10 - 23 February 1951.

| Prefecture | Diseases | No. of Cases (10 - 16 Feb) | No. of Cases (17 - 23 Feb) |
|---|---|---|---|
| Chiba | Swine erysipelas | 1 | |
| Hiroshima | Swine cholera | | 31 |
| Kanagawa | Swine cholera | 2 | 1 |
| Mie | Anthrax | 1 | |

Cumulative Total - 1 January to date of report.

| | |
|---|---|
| Anthrax | 3 |
| Swine cholera | 103 |
| Swine erysipelas | 9 |
| Swine plague | 4 |

Monthly Meat Inspection Report

Prefectural veterinary meat inspectors conducted ante and post-mortem inspections on the following number of livestock during the month of December 1950.

| | Number Slaughtered |
|---|---|
| Cattle | 49,294 |
| Calves | 3,053 |
| Horses | 13,589 |

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775 013

Numbers Slaughtered

| | |
|---|---|
| Swine | 113,194 |
| Sheep | 638 |
| Goats | 415 |

Sanitary inspections were conducted on the following number of meat processing establishments by Prefectural veterinary meat inspectors during the month of December 1950.

Meat Processing Plants

| | | |
|---|---|---|
| Numbers of establishments | | 355 |
| " " " inspected | | 181 |
| Sanitary Grades | A | 19 |
| | B | 44 |
| | C | 68 |
| | D | 47 |
| | Sub-standard | 3 |
| Total number of inspections | | 308 |
| Amount of meat & meat-food products examined | | 65,800 kgs |
| " " " " condemned | | 49 " |

Meat Retail Shops

| | | |
|---|---|---|
| Number of establishments | | 18,657 |
| " " " inspected | | 9,288 |
| Sanitary Grades | A | 912 |
| | B | 2,519 |
| | C | 3,790 |
| | D | 1,805 |
| | Sub-standard | 262 |
| Total number of inspections | | 11,399 |
| Amount of meat products inspected | | 851,224 kgs |
| " " " " condemned | | 470 " |

SOURCE: Ministry of Welfare

Monthly Seafood Inspection Report

Prefectural veterinary seafood inspectors conducted sanitary inspectins on the following seafood establishments during the month of December 1950.

Wholesale Seafood Market

| | | |
|---|---|---|
| Number of establishments | | 1,483 |
| " " " inspected | | 828 |
| Sanitary Grades | A | 59 |
| | B | 123 |
| | C | 354 |
| | D | 281 |
| | Sub-standard | 71 |

Seafood Processing Plants

| | | |
|---|---|---|
| Number of establishments | | 16,250 |
| " " " inspected | | 4,432 |
| Sanitary Grades | A | 206 |
| | B | 747 |
| | C | 1,554 |
| | D | 1,609 |
| | Sub-standard | 316 |

Seafood Retail Shops

| | | |
|---|---|---|
| Number of establishments | | 53,348 |
| " " " inspected | | 20,388 |

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775 013

| | |
|---|---|
| Sanitary Grades A | 1,192 |
| B | 3,357 |
| C | 8,348 |
| D | 5,822 |
| Sub-standard | 1,669 |

SOURCE: Ministry of Welfare

Monthly Milk Inspection Report for December 1950

Milk Production

| Type of Milk | No. of Inspection | No. of Farms | No. of Animals | Amount Produced |
|---|---|---|---|---|
| Special | 31 | 6 | 178 | 44,769 Lit. |
| Ordinary | 8,227 | 74,737 | 137,166 | 31,646,026 " |
| Goat | 76 | 478 | 1,245 | 51,105 " |

Bottled Milk

| Type of Milk | No. of Inspection | No. of Plants | Raw Milk | Pasteurized | Sterilized |
|---|---|---|---|---|---|
| Special | 22 | 6 | 21,058 Lit. | 19,328 Lit. | - |
| Ordinary | 3,239 | 2,991 | - | 8,667,857 " | 6,442,160 Lit. |

Note: All milk bottled in 1 go bottles having the equivalent of 180 cc.

Dairy Products

| | |
|---|---|
| Whole powdered milk | 103.191 kgs |
| Modified powdered milk (15% added sugar) | 1,100,234 " |
| " " " (35% added sugar) | 191,194 " |
| Sweetened condensed milk | 1,774,980 " |
| Evaporated milk | 35,448 " |
| Butter | 604,441 " |
| Cheese | 81,927 " |

SOURCE: Ministry of Welfare

Monthly Food Sanitation Report

Prefectural veterinary and food sanitation inspectors conducted sanitary inspections on those establishments where food and beverages of other than animal origin were processed, served, or sold during the month of December 1950.

| | |
|---|---|
| Number of food establishments | 1,256,782 |
| " " " inspected | 218,130 |
| Total number of inspections | 315,422 |
| Number of cases of food condemnations | 7,629 |
| " " suspensions | 347 |
| " " permits revoked | 70 |
| " " violation cases in civil court | 41 |

A total of 29 outbreaks of food or beverage poisoning was reported involving 233 people with 14 deaths resulting.

SECTION V SUPPLY

Benzedrine

The Japanese pharmaceutical industry produces limited quantities of propamine and methyl-propamine types of benzedrine. This drug is considered a

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775 013

habit forming central nervous system stimulent, and has been the basis of much concern and considerable study on the part of both SCAP and the Ministry of Welfare to restrict its use exclusively for medical purposes. Information was recently received in Japan concerning the discovery of a benzedrine-type drug in the United States which has all the medical properties of benzedrine and none of its ill effects. The Ministry of Welfare is investigating the possibility of manufacturing such a drug in Japan, which would help solve the problem of benzedrine addiction.

Production of Medical Supplies

Production of medical supplies during January totalled 3,822,570,100 yen as follows.

| | (Yen value) |
|---|---|
| Controlled Medicines | 169,065,520 |
| Non-Controlled Medicines | 2,401,086,398 |
| Patent Medicines | 647,265,410 |
| Biologics | 140,540,137 |
| Dental Instruments | 24,803,927 |
| Dental Materials | 15,418,664 |
| Textile Sanitary Materials | 375,621,188 |
| Rubber Sanitary Goods | 48,768,860 |
| Total | 3,822,570,104 |

Streptomycin Production

Production of indigenous streptomycin during January totalled 65,189 grams, an increase of 24 percent over reported production for December. The entire quantity passed assay and 5,836 grams were still under assay 1 February. Production by manufacturer follows:

(Unit = gram)

| Producer | Type Sale | Production |
|---|---|---|
| Meiji Seika | H Cl | 33,000 |
| Nihon Seibutsu | H Cl | 13,560 |
| Kyowa Hakko | $Ca Cl_2$ | 9,760 |
| Kagaku Kenkyusho | H Cl | 6,144 |
| Shionogi Kagaku | H Cl | 2,725 |
| | | 65,189 grams. |

## SECTION VI NARCOTIC CONTROL

Enforcement

Ministry of Welfare Narcotic Agents, stationed in Hyogo prefecture, seized 3,878 grams of heroin ranging in purity from 84 to 94%. The seizure represents approximately one-third of the total amount of heroin seized in Japan during 1950.

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775 013

The heroin was seized in the room of the owner of a large Chinese restaurant-hotel in Kobe and was allegedly smuggled aboard a ship bound from Hongkong to Kure.

## SECTION VII WELFARE

Prohibition Against Japanese Civil Courts Exercising Jurisdiction Over Occupation Persons.

Attention of Civil Affairs Welfare Officers is called to the provisions of SCAPIN 2127, 18 October 1951, paragraph 8, which precludes the Japanese courts from exercising civil jurisdiction over any Occupation person. The SCAPIN is quoted as follows:

> "Paragraph 8. Civil Jurisdiction. Japanese Courts will exercise no civil jurisdiction of any sort with respect to any case in which any one of the parties is a person within the category designated as Occupation personnel--."

The prohibition provided in the SCAPIN has been correctly interpreted by the Japanese as precluding the Family Courts from accepting petitions from O... Occupation persons for adoption in Japan of children of Japanese nationality.

However, the prohibition in the SCAPIN with respect to jurisdiction of the civil courts has been incorrectly interpreted by some of the Japanese Child Welfare Centers as also precluding the acceptance of requests by Occupation persons for approval for placement in their homes of Japanese children who are under the jurisdiction of the Centers. Nothing in the SCAPIN precludes a Child Welfare Center from approving the home of an Occupation person as a foster home in which a child of Japanese nationality may be placed in accordance with Article 27, paragraph 3, of the Child Welfare Law. Approval of the home of an Occupation person as a foster home under the provisions of Article 27 is an administrative and not a judicial procedure requiring approval of the Family Court.

Occupation persons inquiring at Civil Affairs concerning the procedures necessary for taking Japanese children from Japan for purposes of adoption in the United States may be referred to the United States Consular authorities. Consular authorities and the Law Division, Legal Section, SCAP, have been of assistance to Occupation persons in several such cases having come to the attention of PH&W during recent months, particulary concerning the preparation and presentation of necessary bills to Congress authorizing admission of the children into the United States.

Social Affairs Bureau Directives

| Date | File Number | From | To | Subject |
|---|---|---|---|---|
| 17 Jan 51 | Sha-otsu-hatsu No. 6 | Chief, Social Affairs Bureau | All Prefectural Governors | Issuance of this directive was prompted by an inquiry made by Wakayama Prefecture concerning the eligibility of an applicant for medical aid under the Daily Life Security Law. Aid originally was denied because applicant had applied in a locality other than the residence of her parents. The Ministry advised the original application was proper as subject had obtained permanent employment away from home and was maintaining |

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775 013

| Date | Fiel Number | From | To | Subject |
|---|---|---|---|---|
| | | | | a separate residence at the time of her illness. |

Ministry Directive Concerning Temporary Protection of Children by Police.

As Incl. No. 2, for information of Civil Affairs Welfare Officers, is a translation of Children's Bureau Directive, Ji-hatsu No. 12, dated 17 January 1951, subject: Temporary Protection of Children by Police Officers. The Directive transmits a copy of Kei-hatsu-bo No. 15, date 13 November 1950, from the Criminal Affairs Department, National Rural Police, detailing the procedures under which local police may accept temporary custody of children.

Child Welfare Institutions

For information in the field there is reproduced the following table of Children's institutions as compiled by the Children's Bureau. Figures on capacity and number of inmates may be considered as average for the year 1950 and are approximate.

| Kind of Institutions | Type Institution Public | Private | Total Institutions | Total Inmates | Total Capacity |
|---|---|---|---|---|---|
| Mother's Homes | 150 | 137 | 287 | 17,222 | 15,378 |
| Day Nurseries | 765 | 2,106 | 2,871 | 249,166 | 215,377 |
| Lying-In Agencies | 29 | 184 | 213 | 608 | 1,953 |
| Infant Homes | 35 | 75 | 110 | 1,659 | 2,843 |
| Homes for Physically Weak Children | 4 | 6 | 10 | 368 | 455 |
| Homes for Crippled Children | 1 | 1 | 2 | 62 | 110 |
| Homes for Dependent Children | 57 | 325 | 382 | 19,815 | 19,496 |
| Homes for Feeble Minded Children | 10 | 22 | 32 | 1,453 | 1,267 |
| Homes for Blind, Deaf and Dumb Children | 32 | 7 | 39 | 2,138 | 2,138 |
| Homes for Juvenile Training and Education | 48 | 11 | 59 | 4,564 | 4,517 |
| Totals | 1,131 | 2,874 | 4,005 | 297,065 | 263,534 |

UNICEF

After some two years as Chief, UNICEF Mission in Japan, attached to PH&W, Miss Margherita Strahler has resigned to return to the United States. Miss Strahler is being replaced by Mr. Bernard Fraser who will assume joint responsibilities as Chief, UNICEF Missions in Japan and in Korea. Mr. Fraser has had extensive experience in the UNICEF program recently having served as UNICEF Liaison Officer with the International Tuberculosis Campaign, Headquarters, Copenhagen.

GS Mollohan Col MC

for: CRAWFORD F. SAMS
Brigadier General, Medical Corps
Chief

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775 013

11 Incls:

1. Ministry of Welfare, Notification #24, 23 Feb. '51
2. Letter, Ji-Hatsu #12, 17 Jan. '51
3. Results of Nutrition Surveys - Japan, Nov. '50
4. Weekly C. D. Digest, 17 Feb. '51
5. Weekly C. D. Report, 17 Feb. '51
6. Weekly C. D. Digest, 24 Feb. '51
7. Weekly C. D. Report, 24 Feb. '51
8. Monthly C. D. Digest, 24 Feb. '51
9. Monthly C. D. Report, 24 Feb. '51
10. Japanese Hospital Digest, Jan. '51
11. Japanese Hospital Report, Jan. '51

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775 013

MINISTRY OF WELFARE
Notification No. 24

23 February 1951

The examination places, date and the time for application for the second National Examination for class A nurses in accordance with the provisions of Article 19 of the Public Health Nurse, Midwife and Nurse Law Enforcement Regulations have been decided as follows.

1. Place ; Sapporo City, Obihiro City, Sendai City, Fukushima City, Urawa City, Tokyo, Yokohama City, Niigata City, Kanazawa City, Shizuoka City, Nagoya-City, Kyoto City, Ohsaka City, Kobe City, Matsue City, Okayama City, Bofu City, Kochi City, Fukuoka City, Kumamoto City, Kagoshima City.

2. Date ; May 12 (Saturday) and 13 (Sunday), 1951.

3. Time for Application ; March 1, 1951 through March 20,1951. Applications submitted after the dead line shall not be accepted regardless of reasosns. However mailed applications with the post mark March 20 on them will be accepted.

4. Address to which applications will be submitted ;
   National Examination Division, Nursing Section
   Medical Affairs Bureau, Welfare Ministry
   1-2 Kasumigaseki, Chiyoka-ku Tokyo-to.

5. When an application is mailed, the name of the examination place shall be written on the envelope in red ink, like "Examination place such and such City".

6. Each applicant shall recieve an examination card after her application has been accepted.

7. An applicant can take her examination only at the examination place indicated on her examination card.

8. A revenue stamp of 350 yen shall be affixed to the application as examination fee, and it shall not bear any cancellation stamp.

9. As to the form of application and accompanying papers, those who come under the provisions of Article 21 of the Public Health Nurse, Midwife and Nurse Law shall refer to the provisions of Article 23 of the Enforcement Regulations of the Law and those who come under the provisions of Item 1, Article 53 of the same Law (those who recieved licensure from the prefectural governor accoroing to the old nurses' regulations) shall refer to the provisions of Item 3 of the Supplementary Regulations of the Enforcement Regulations of the Law, so that there will be no mistake about them.

10. The size of the applicants photograph wich will be affixed to the application shall be "tefudagata" (1½" by 2") and the photograph shall not be mounted.

11. The name of the examination place written on the application cannot be changed regardless of reasons after the application has been submitted.

12. The applicant's name written on the application shall be accompanied with "kana".

Incls. No. 1

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775 013

... ... place of protection.

When a police office is to give a child temporary protection, the protective room shall be used and, if the room should be locked, enough space must be allowed for the child to move around. The protective room to be used for the temporary protection of a child must be something like a regular room and any detention room shall never be used for a substitute.

III. Length of time to give protection.

The length of time the temporary protection is to be given must not exceed, as a rule 24 hours after the time the protection was begun: Provided that it may be extended in case there is a transportation or other inevitable reason. The temporary protection of the child after 24 hours passed is still under the consignment of the superintendent of the child welfare center according to the law and not subject to Article 3 par. 3 of the Law Concerning Execution of Duties of Police Officials.

IV. Expense.

As the expense required for the temporary protection of a child is to be paid by the prefecture as provided under Article 50 par. 1 item(9) of the law, the preliminary liaison with the prefectural section concerned as to the method of payment and other details must be made.

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775 013

Ji-Hatsu No. 12 17 January 1951

TO : Each of To, Do, Fu and Prefectures.

FROM : Director, Children's Bureau, Ministry of Welfare.

SUBJECT: Temporary Protection of children by Police Office.

Regarding the compulsory temporary protection to be taken for a child in accordance with Articles 33 of the Child Welfare Law, we have sent you before the notification by Ji-Hatsu, No. 505, "Re: The case of Compulsory Measure to be Taken for a child under the Child Welfare Law" dated 31 July 1950. The attached copy of the instruction given by the Criminal Affairs Dept. National Rural Police Headquarter - Kei-Hatsu-Bo, No. 15, dated on 13 November 1950 to each of the directors of Regional Headquarter of Police, chieves of Prefectural District Police and chief police of To and five largest cities regarding the case of temporary protection of a child consigned with a police office by the supperintendent of the child welfare center in accordance with the above notification should serve for your good team work with them. Your cooperation with the police organs concerned for the temporary protection of children is highly appreciated.

It must be noted that the notification of 31 July 1950-JiHatsu, No. 505 has been cleared with the National Rural Police Headquarters.

Kei-Hatsu*bo, No. 15 13 November 1950

TO: : Director, Regional Headquarter of Police Chef, Each Prefectural District Police (Chief Police of To and 5 largest cities).

FROM : Chief, Criminal Affairs Dep't. National Rural Police Headquarters.

SUBJECT: The Temporary Protection of Children to be Given by the Police Office.

The temporary protection of children by the police office with the authority delegated by the superintendent of the child welfare center under the provision of Article 33 par. 1 of the Child Welfare Law has been given in accordance with the instructions - 1948, June, Kei-Bo-Hatsu, No. 12 (Re: The Protection after the Repeal of the Administrative Execution Law) and 1948 December, Kei-Bo-Hatsu No. 109 (Re: The Disposition of Juvenile Cases) issued by the name of chief of Criminal Affairs Dep't. Regarding the nature of this temporary protection the attached notification "Re: The compulosry Measure to be taken for a child under the Chicl Welfare Law" dated on 31 July, 1950, issued by the director of Children's Bureau, Welfare Ministry is to serve for your understanding and your full cooperation is highly appreciated.

Your are requested to notify this to the municipal police offices and cooperations of all of you are much appreciated.

N O T E

I. When a police office is to give temporary protection of a child. The police office may give a temporary protection when a national or local police officer has discovered a child in need of temporary protection as provided under Article 33 of the Child Welfare Law or when such child is referred to the police office by any one at large, the child welfare center is located in distance, at night or when the child welfare center is not able to get such child immediately.

The consignment of the temporary protection by the superintendent of the child welfare center at such instances shall be understood as has received beforhand all inclusively. (This is cleared out with the Welfare Ministry).

Incl. 2

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 773 013

RESULTS OF NUTRITION SURVEYS- JAPAN, NOVEMBER 1950

NUTRIENTS IN GRAMS AND CALORIES PER CAPITA PER DAY

| | All Japan | 1/ Urban | TOKYO Total | TOKYO Farmer | TOKYO Non-farmer | 2/ ELEVEN CITIES Total | ELEVEN CITIES Farmer | ELEVEN CITIES Non-farmer | 3/ OTHER CITIES Total | OTHER CITIES Farmer | OTHER CITIES Non-farmer |
|---|---|---|---|---|---|---|---|---|---|---|---|
| Number of Families | 7,876 | 4,047 | 944 | 25 | 919 | 1,147 | 151 | 996 | 1,956 | 228 | 1,728 |
| Number of persons | 41,571 | 20,142 | 4,739 | 161 | 4,578 | 5,774 | 975 | 4,799 | 9,629 | 1,406 | 8,223 * |
| Adult Units: | | | | | | | | | | | |
| Protein | 0.854 | 0.860 | 0.867 | 0.876 | 0.867 | 0.860 | 0.872 | 0.857 | 0.859 | 0.853 | 0.860 |
| Calories | 0.847 | 0.817 | 0.802 | 0.899 | 0.799 | 0.815 | 0.868 | 0.805 | 0.822 | 0.884 | 0.812 |
| Protein (Gram) | | | | | | | | | | | |
| Total | 69.8 | 72.5 | 70.3 | 66.3 | 70.4 | 71.9 | 74.3 | 71.4 | 73.4 | 70.5 | 73.9 |
| Animal | 18.6 | 23.0 | 22.1 | 15.1 | 22.4 | 22.8 | 18.2 | 23.8 | 23.4 | 16.0 | 24.7 |
| Vegetable | 51.2 | 49.5 | 48.2 | 51.2 | 48.0 | 49.1 | 56.1 | 47.6 | 50.0 | 54.5 | 49.2 |
| Fat (Gram) | 18.6 | 21.7 | 22.8 | 19.1 | 23.0 | 20.6 | 17.8 | 21.1 | 21.8 | 17.7 | 22.5 |
| Carbohydrate (Gram) | 435.6 | 405.2 | 377.9 | 458.5 | 375.0 | 408.8 | 503.4 | 389.6 | 411.2 | 467.1 | 402.0 |
| Calories | | | | | | | | | | | |
| Total | 2,188 | 2,107 | 1,997 | 2,277 | 1,987 | 2,111 | 2,474 | 2,037 | 2,136 | 2,299 | 2,103 |
| Ration | 672 | 1,202 | 1,342 | 427 | 1,375 | 1,118 | 118 | 1,321 | 1,198 | 179 | 1,376 |
| Free market | 343 | 568 | 567 | 301 | 576 | 560 | 199 | 634 | 572 | 246 | 628 |
| Home production | 1,149 | 308 | 60 | 1,528 | 8 | 419 | 2,153 | 66 | 330 | 1,858 | 64 |
| Others | 24 | 29 | 28 | 21 | 28 | 14 | 4 | 16 | 36 | 16 | 40 |

x Correction August 1950 Survey - Change from 21,499 to 8,267

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775 013

RESULTS OF NUTRITION SURVEYS - JAPAN, NOVEMBER 1950

NUTRIENTS IN GRAMS AND CALORIES PER CAPITA PER DAY

| | RURAL AREAS 4/ | | | TOKYO RAILWAYS | AKITA COPPER MINE | THREE COAL MINES |
|---|---|---|---|---|---|---|
| | Total | Farmer | Non-farmer | | | |
| Number of Families | 3,829 | 2,648 | 1,181 | 88 | 100 | 280 |
| Number of Persons | x 21,429 | 15,752 | 5,677 | 427 | 472 | 1,285 |
| Adult Units: | | | | | | |
| Protein | 0.851 | 0.852 | 0.846 | 0.826 | 0.819 | 0.839 |
| Calories | 0.862 | 0.883 | 0.803 | 0.760 | 0.803 | 0.842 |
| Protein (Gram) | | | | | | |
| Total | 68.5 | 68.2 | 69.1 | 69.1 | 82.0 | 76.7 |
| Animal | 16.5 | 15.3 | 19.7 | 21.9 | 26.7 | 25.1 |
| Vegetable | 52.0 | 52.9 | 49.4 | 47.2 | 55.3 | 51.6 |
| Fat (Gram) | 17.1 | 16.8 | 17.9 | 21.7 | 18.9 | 22.8 |
| Carbohydrate (Gram) | 450.7 | 460.8 | 421.7 | 377.0 | 449.7 | 426.3 |
| Calories | | | | | | |
| Total | 2,228 | 2,265 | 2,124 | 1,981 | 2,296 | 2,220 |
| Ration | 409 | 127 | 1,213 | 1,327 | 1,632 | 1,517 |
| Free market | 232 | 164 | 427 | 610 | 633 | 599 |
| Home production | 1,565 | 1,959 | 443 | 22 | 6 | 73 |
| Others | 22 | 15 | 40 | 22 | 25 | 31 |

1/ All cities 30,000 population and over
2/ Nagoya, Osaka, Kure, Fukuoka, Sapporo, Sendai, Kanazawa, Matsuyama, Yokohama, Kyoto and Kobe
3/ All cities 30,000 population and over excluding Tokyo and 11 cities
4/ Rural areas in all prefectures

x Correction August 1950 survey - Change from 15,672 to 21,499

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775 013

GRAMS OF VARIOUS CLASSES OF FOOD CONSUMED PER CAPITA PER DAY FROM NUTRITION SURVEYS - JAPAN, NOV. 1950

| | All Japan | Urban | TOKYO Total | TOKYO Farmer | TOKYO Nonfarmer |
|---|---|---|---|---|---|
| Grains - Total | 475.9 | 467.9 | 496.6 | 513.9 | 496.0 |
| Rice | 343.1 | 305.8 | 283.0 | 409.3 | 278.5 |
| Wheat | 64.5 | 106.6 | 187.2 | 58.4 | 191.8 |
| Barley | 65.1 | 53.5 | 25.2 | 46.2 | 24.5 |
| Corn meal | 0.1 | 0.2 | 0.0 | - | 0.0 |
| Others | 3.1 | 1.8 | 1.2 | - | 1.2 |
| Nuts, Etc. | 1.6 | 1.2 | 0.9 | 0.5 | 0.9 |
| Potatoes - Total | 191.5 | 137.9 | 97.7 | 232.1 | 92.9 |
| Sweet | 136.9 | 97.1 | 55.6 | 118.0 | 53.4 |
| White | 19.0 | 18.5 | 24.8 | 61.5 | 23.5 |
| Others | 35.6 | 22.3 | 17.3 | 52.6 | 16.0 |
| Sugars | 7.1 | 10.1 | 11.2 | 5.0 | 11.5 |
| Oils | 2.6 | 3.5 | 4.5 | 3.1 | 4.6 |
| Legumes - Total | 56.0 | 59.9 | 58.2 | 46.6 | 58.7 |
| Soya bean | 1.9 | 1.3 | 0.3 | - | 0.4 |
| Miso | 32.2 | 25.3 | 25.5 | 33.0 | 25.3 |
| Soya products | 16.4 | 27.7 | 27.3 | 12.7 | 27.8 |
| Other beans | 5.5 | 5.6 | 5.1 | 0.9 | 5.2 |
| Animal Foods - Total | 89.8 | 122.0 | 120.5 | 79.1 | 121.9 |
| Fish - fresh | 61.2 | 81.3 | 76.9 | 65.0 | 77.3 |
| Fish - dried | 9.5 | 10.1 | 5.9 | 6.5 | 5.8 |
| Meat, Poultry | 9.1 | 13.8 | 16.8 | 5.6 | 17.2 |
| Eggs | 3.8 | 6.3 | 7.5 | 1.7 | 7.7 |
| Milk | 6.2 | 10.5 | 13.4 | 0.3 | 13.9 |
| Leafy, Green & Yellow Vegetables | 75.8 | 77.7 | 58.7 | 40.9 | 59.3 |
| Other Fruits & Vegetables - Total | 157.3 | 147.8 | 119.0 | 80.4 | 120.4 |
| Citrus, Tomatoes | 8.5 | 14.2 | 11.8 | 3.4 | 12.1 |
| Other Fruits | 34.4 | 43.8 | 35.0 | 3.4 | 36.2 |
| Other Vegetables | 114.4 | 89.8 | 72.2 | 73.6 | 72.1 |
| Seaweeds | 2.5 | 3.5 | 3.4 | 1.9 | 3.4 |
| Processed Vege. - Total | 41.7 | 41.0 | 34.4 | 69.6 | 33.1 |
| Dried | 0.6 | 0.5 | 0.5 | 0.4 | 0.5 |
| Pickled | 41.1 | 40.5 | 33.9 | 69.2 | 32.6 |
| Flavours | 31.7 | 33.7 | 26.2 | 24.6 | 26.2 |
| Total | 1,133.5 | 1,106.2 | 1,031.3 | 1,097.7 | 1,028.9 |

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775 013

GRAMS OF VARIOUS CLASSES OF FOOD CONSUMED PER CAPITA PER DAY FROM NUTRITION SURVEYS - JAPAN, NOV. 1950

| | ELEVEN CITIES | | | OTHER CITIES | | |
|---|---|---|---|---|---|---|
| | Total | Farmer | Non-farmer | Total | Farmer | Non-farmer |
| Grains - Total | 465.8 | 524.0 | 453.9 | 461.0 | 496.8 | 454.9 |
| Rice | 321.7 | 435.2 | 298.6 | 305.6 | 334.5 | 300.6 |
| Wheat | 90.8 | 33.8 | 102.4 | 91.1 | 68.5 | 95.1 |
| Barley | 52.1 | 54.6 | 51.6 | 61.8 | 92.0 | 56.6 |
| Corn meal | 0.2 | 0.4 | 0.1 | 0.2 | 1.4 | 0.0 |
| Others | 1.0 | - | 1.2 | 2.3 | 0.4 | 2.6 |
| Nuts, Etc. | 1.1 | 2.6 | 0.8 | 1.3 | 1.5 | 1.3 |
| Potatoes - Total | 154.9 | 289.6 | 127.4 | 141.9 | 174.5 | 136.1 |
| Sweet | 110.3 | 211.7 | 89.6 | 103.0 | 110.4 | 101.7 |
| White | 19.3 | 26.7 | 17.8 | 16.5 | 29.9 | 14.1 |
| Others | 25.3 | 51.2 | 20.0 | 22.4 | 34.2 | 20.3 |
| Sugars | 9.4 | 5.5 | 10.1 | 10.0 | 5.3 | 10.8 |
| Oils | 3.2 | 1.3 | 3.6 | 3.3 | 2.6 | 3.4 |
| Legumes - Total | 60.9 | 61.0 | 60.9 | 60.0 | 50.0 | 61.7 |
| Soya bean | 0.8 | 1.3 | 0.7 | 1.7 | 2.3 | 1.6 |
| Miso | 23.6 | 34.8 | 21.3 | 26.0 | 31.9 | 25.0 |
| Soya products | 31.2 | 17.2 | 34.1 | 26.4 | 8.6 | 29.4 |
| Other beans | 5.3 | 7.7 | 4.8 | 5.9 | 7.2 | 5.7 |
| Animal Foods - Total | 125.2 | 95.1 | 131.4 | 121.1 | 85.5 | 127.3 |
| Fish - fresh | 82.5 | 74.4 | 84.1 | 82.1 | 54.4 | 86.9 |
| Fish - dried | 9.3 | 8.1 | 9.6 | 11.6 | 9.7 | 11.9 |
| Meat, Poultry | 17.0 | 6.8 | 19.1 | 11.6 | 4.2 | 12.9 |
| Eggs | 6.7 | 2.6 | 7.5 | 5.8 | 3.2 | 6.3 |
| Milk | 9.7 | 3.2 | 11.1 | 10.0 | 14.0 | 9.3 |
| Leafy, Green & Yellow vegetables | 68.1 | 50.1 | 71.7 | 86.7 | 163.2 | 73.4 |
| Other Fruits & Vegetables - Total | 163.3 | 149.6 | 166.1 | 149.4 | 187.8 | 142.3 |
| Citrus, Tomatoes | 16.7 | 11.4 | 17.8 | 13.9 | 5.5 | 15.4 |
| Other Fruits | 45.1 | 26.5 | 48.9 | 45.7 | 59.3 | 43.3 |
| Other Vegetables | 101.5 | 111.7 | 99.4 | 89.8 | 123.0 | 84.1 |
| Seaweeds | 4.2 | 3.1 | 4.5 | 3.2 | 2.1 | 3.4 |
| Processed Vege. - Total | 37.6 | 49.6 | 35.2 | 44.2 | 82.1 | 37.6 |
| Dried | 0.4 | 0.4 | 0.5 | 0.5 | 0.1 | 0.6 |
| Pickled | 37.2 | 49.2 | 34.7 | 43.7 | 82.0 | 37.0 |
| Flavours | 34.3 | 39.8 | 33.3 | 35.5 | 32.2 | 36.0 |
| Total | 1,128.0 | 1,271.3 | 1,098.9 | 1,117.6 | 1,283.6 | 1,088.7 |

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775 013

GRAMS OF VARIOUS CLASSES OF FOOD CONSUMED PER CAPITA PER DAY FROM NUTRITION SURVEYS - JAPAN, NOV. 1950

| | RURAL AREAS | | | | | |
|---|---|---|---|---|---|---|
| | Total | Farmer | Non-Farmer | Tokyo Railways | Akita Copper Mine | Three Coal Mines |
| Grains - Total | 479.9 | 491.8 | 445.8 | 475.5 | 487.2 | 451.3 |
| Rice | 361.6 | 377.2 | 317.0 | 267.5 | 410.5 | 334.7 |
| Wheat | 43.6 | 39.1 | 56.4 | 175.0 | 39.7 | 69.6 |
| Barley | 70.8 | 71.1 | 70.1 | 32.7 | 36.3 | 46.2 |
| Corn meal | 0.1 | 0.1 | 0.0 | - | - | - |
| Others | 3.8 | 4.3 | 2.3 | 0.3 | 0.7 | 0.8 |
| Nuts, Etc. | 1.8 | 1.4 | 2.8 | 0.5 | 0.1 | 0.1 |
| Potatoes - Total | 218.0 | 220.1 | 211.7 | 120.8 | 116.7 | 151.9 |
| Sweet | 156.6 | 158.9 | 150.0 | 91.4 | 90.9 | 107.1 |
| White | 19.3 | 18.6 | 21.1 | 15.3 | 9.5 | 20.7 |
| Others | 42.1 | 42.6 | 40.6 | 14.1 | 16.3 | 24.1 |
| Sugars | 5.6 | 5.1 | 7.1 | 10.3 | 6.2 | 13.3 |
| Oils | 2.2 | 2.2 | 2.3 | 3.4 | 2.9 | 4.4 |
| Legumes - Total | 54.1 | 53.5 | 55.7 | 51.4 | 87.2 | 63.2 |
| Soya bean | 2.2 | 2.3 | 2.1 | 0.4 | 1.1 | 2.2 |
| Miso | 35.6 | 37.4 | 30.3 | 26.4 | 50.7 | 31.5 |
| Soya products | 10.8 | 8.0 | 18.7 | 21.0 | 30.0 | 23.9 |
| Other beans | 5.5 | 5.8 | 4.6 | 3.6 | 5.4 | 5.6 |
| Animal Foods - Total | 73.9 | 67.5 | 92.4 | 113.6 | 144.8 | 129.8 |
| Fish - fresh | 51.3 | 46.9 | 63.8 | 82.9 | 127.2 | 107.7 |
| Fish - dried | 9.2 | 8.7 | 10.4 | 6.9 | 5.2 | 6.7 |
| Meat, Poultry | 6.8 | 6.0 | 9.2 | 11.0 | 10.0 | 7.9 |
| Eggs | 2.5 | 2.2 | 3.7 | 5.8 | 1.8 | 3.1 |
| Milk | 4.1 | 3.7 | 5.3 | 7.0 | 0.6 | 4.4 |
| Leafy, Green & Yellow Vegetables | 74.9 | 76.0 | 71.8 | 53.3 | 108.2 | 91.0 |
| Other Fruits & Vegetables - Total | 162.0 | 163.7 | 157.2 | 138.4 | 232.0 | 209.2 |
| Citrus, Tomatoes | 5.7 | 4.3 | 9.7 | 16.3 | 1.6 | 6.2 |
| Other Fruits | 29.7 | 28.6 | 33.0 | 33.2 | 132.8 | 36.5 |
| Other Vegetables | 126.6 | 130.8 | 114.5 | 88.9 | 97.6 | 166.5 |
| Seaweeds | 2.0 | 1.7 | 2.6 | 1.9 | 1.1 | 1.6 |
| Processed Vegetables - Total | 41.9 | 44.6 | 34.5 | 26.4 | 82.2 | 56.4 |
| Dried | 0.6 | 0.7 | 0.4 | 0.0 | 0.1 | 0.3 |
| Pickled | 41.3 | 43.9 | 34.1 | 26.4 | 82.1 | 56.1 |
| Flavours | 30.7 | 30.5 | 31.3 | 32.3 | 40.2 | 44.0 |
| Total | 1,147.0 | 1,158.1 | 1,115.2 | 1,027.8 | 1,308.8 | 1,216.2 |

DISTRIBUTION OF FAMILIES ACORDING TO CONSUMED CALORIES PER CAPITA PER DAY, NOV. 1950

| Class | TOKYO | | | ELEVEN CITIES | | | OTHER CITIES | | | RURAL AREAS | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | Total | Farmer | Non-farmer | Total | Farmer | Non-farmer | Total | Farmer | Non-farmer | Total | Farmer | Non-farmer |
| Total | 944 | 25 | 919 | 1,147 | 151 | 996 | 1,956 | 228 | 1,728 | 3,829 | 2,648 | 1,181 |
| Under 1000 | - | - | - | - | - | - | - | - | - | 2 | 1 | 1 |
| 1000 - 1099 | 1 | - | 1 | 3 | - | 3 | 5 | - | 5 | 1 | - | 1 |
| 1100 - 1199 | 4 | - | 4 | 6 | - | 6 | 13 | - | 13 | 11 | 6 | 5 |
| 1200 - 1299 | 11 | - | 11 | 7 | - | 7 | 12 | 1 | 11 | 24 | 14 | 10 |
| 1300 - 1399 | 24 | - | 24 | 12 | - | 12 | 15 | 1 | 14 | 48 | 30 | 18 |
| 1400 - 1499 | 37 | 1 | 36 | 30 | 2 | 28 | 53 | 3 | 50 | 88 | 49 | 39 |
| 1500 - 1599 | 53 | - | 53 | 52 | 1 | 51 | 86 | 7 | 79 | 120 | 73 | 47 |
| 1600 - 1699 | 72 | 1 | 71 | 73 | 1 | 72 | 119 | 9 | 110 | 170 | 99 | 71 |
| 1700 - 1799 | 91 | 1 | 90 | 101 | 4 | 97 | 137 | 13 | 124 | 214 | 134 | 80 |
| 1800 - 1899 | 92 | 1 | 91 | 129 | 6 | 123 | 157 | 14 | 143 | 280 | 171 | 109 |
| 1900 - 1999 | 90 | 1 | 89 | 101 | 8 | 93 | 183 | 14 | 169 | 329 | 210 | 119 |
| 2000 - 2099 | 98 | 4 | 94 | 107 | 10 | 97 | 175 | 21 | 154 | 378 | 260 | 118 |
| 2100 - 2199 | 68 | 1 | 67 | 103 | 13 | 90 | 182 | 14 | 168 | 332 | 225 | 107 |
| 2200 - 2299 | 58 | 2 | 56 | 92 | 16 | 76 | 175 | 18 | 157 | 312 | 218 | 94 |
| 2300 - 2399 | 61 | 4 | 57 | 71 | 13 | 58 | 136 | 20 | 116 | 269 | 197 | 72 |
| 2400 - 2499 | 49 | - | 49 | 63 | 9 | 54 | 117 | 22 | 95 | 242 | 189 | 53 |
| 2500 - 2599 | 37 | 4 | 33 | 51 | 14 | 37 | 107 | 17 | 90 | 223 | 163 | 60 |
| 2600 - 2699 | 28 | - | 28 | 34 | 8 | 26 | 78 | 8 | 70 | 182 | 141 | 41 |
| 2700 - 2799 | 24 | 1 | 23 | 21 | 4 | 17 | 53 | 9 | 44 | 125 | 95 | 30 |
| 2800 - 2899 | 18 | - | 18 | 29 | 9 | 20 | 34 | 8 | 26 | 114 | 84 | 30 |
| 2900 - 2999 | 8 | - | 8 | 23 | 12 | 11 | 36 | 6 | 30 | 91 | 68 | 23 |
| 3000 - over | 20 | 4 | 16 | 39 | 21 | 18 | 83 | 23 | 60 | 274 | 221 | 53 |



DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775013

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 73 013

## NUTRITIVE VALUE OF FOOD CONSUMED - NOVEMBER 1950

| Area | Calories | Protein Total gram | Protein Animal gram | Fat gram | Carbohydrate gram | Calcium gram | Phosphorus gram | Iron mgm | VITAMINS A I.U. | VITAMINS B1 mgm | VITAMINS B2 mgm | VITAMINS Niacin mgm | VITAMINS C mgm |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| All Japan | 2,188 | 69.8 | 18.6 | 18.6 | 436 | .28 | 1.98 | 48.5 | 2,413 | 1.56 | .75 | 9.11 | 124 |
| Urban Areas | 2,107 | 72.5 | 23.0 | 21.7 | 405 | .29 | 2.01 | 52.1 | 2,563 | 1.58 | .79 | 9.26 | 110 |
| Tokyo | 1,997 | 70.3 | 22.1 | 22.8 | 378 | .28 | 2.02 | 52.1 | 2,002 | 1.59 | .79 | 8.61 | 82 |
| 11 Cities | 2,111 | 71.9 | 22.8 | 20.6 | 409 | .29 | 2.01 | 52.7 | 2,326 | 1.54 | .78 | 9.67 | 113 |
| Other Cities | 2,136 | 73.4 | 23.4 | 21.8 | 411 | .29 | 2.01 | 52.0 | 2,808 | 1.56 | .79 | 9.27 | 117 |
| Rural Areas | 2,288 | 68.5 | 16.5 | 17.1 | 451 | .27 | 1.96 | 46.6 | 2,339 | 1.53 | .72 | 9.01 | 130 |
| Tokyo Railways | 1,981 | 69.1 | 21.9 | 21.7 | 377 | .27 | 1.99 | 49.6 | 1,820 | 1.59 | .73 | 8.55 | 92 |
| Akita Copper Mine | 2,296 | 82.0 | 26.7 | 18.9 | 450 | .34 | 2.10 | 64.7 | 3,360 | 1.48 | .92 | 11.31 | 143 |
| 3 Coal Mines | 2,220 | 76.7 | 25.1 | 22.8 | 426 | .30 | 2.04 | 52.9 | 2,832 | 1.50 | .82 | 10.06 | 133 |

* Before Cooking

Approximate Loss Percent of Vitamins and Minerals by Cooking

| | | |
|---|---|---|
| Vitamins | A | 10 - 20 |
| | $B_1$ | 40 - 50 |
| | $B_2$ | 20 - 30 |
| | C | 50 - 60 |
| Minerals | | 20 - 30 |

Japanese Standard Requirement (per Capita)

| | | | | | |
|---|---|---|---|---|---|
| Calories | - | 2150 Cal | Vitamins | | |
| Protein | - | 75 Grams | A | - | 3,000 I.U. |
| Fat | - | 25 Grams | $B_1$ | - | 1.0 mgm |
| Calcium | - | 1 Gram | $B_2$ | - | 1.0 mgm |
| Iron | - | 10 Grams | Niacin | | 10 mgm |
| | | | C | | 40 mgm |

PERCENT OF PERSONS SHOWING NUTRITIONAL DEFICIENCY SYMPTOMS ACCORDING TO NUTRITION SURVEYS, NOV. 1950

| | No. Persons Examined | 1/ Anemia | 2/ Hyper-keratosis | 3/ Xeroph-thalmia | 4/ Chei-losis | 5/ Glos-sitis | 6/ Loss of Knee Jerk | Edema | 8/ Chronic Diarhea | 9/ Brady Cardia | 10/ Delayed Menstruation | 11/ Deficient Lactation | No Symptoms | One or more Symptoms |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| All Japan | - | 3.0 | 1.9 | 0.2 | 8.3 | 1.5 | 7.5 | 1.0 | 0.4 | 2.9 | 8.5 | *29.4 | 77.2 | 22.8 |
| Urban | - | 2.7 | 2.0 | 0.2 | 4.6 | 1.4 | 7.5 | 1.2 | 0.4 | 2.7 | 8.3 | 24.9 | 80.8 | 19.2 |
| Tokyo | 4,008 | 2.2 | 1.0 | 0.1 | 3.2 | 0.3 | 3.8 | 1.0 | 0.4 | 1.3 | 7.6 | 20.0 | 87.5 | 12.5 |
| 11 Cities | 4,848 | 2.3 | 3.1 | 0.1 | 4.1 | 2.0 | 9.6 | 1.2 | 0.5 | 2.3 | 7.8 | 23.1 | 78.1 | 21.9 |
| Other Cities | 8,211 | 3.1 | 1.8 | 0.2 | 5.3 | 1.4 | 7.6 | 1.2 | 0.4 | 3.2 | 8.5 | 27.0 | 80.1 | 19.9 |
| Rural Areas | 18,394 | 3.2 | 1.9 | 0.2 | 10.1 | 1.5 | 7.5 | 0.9 | 0.4 | 3.0 | 8.7 | **31.6 | 75.4 | 24.6 |
| Tokyo Railways | 385 | 0.8 | - | - | 7.0 | 0.3 | 5.2 | 0.5 | - | 3.3 | 6.0 | - | 83.1 | 16.9 |
| Akita Copper Mine | 450 | - | - | - | 3.8 | 0.4 | 1.8 | - | 0.9 | 8.2 | 9.4 | 14.3 | 87.6 | 12.4 |
| 3 Coal Mines | 1,140 | 1.7 | 2.9 | - | 6.6 | 0.1 | 2.3 | - | 0.4 | 0.4 | 18.6 | 27.3 | 82.7 | 17.3 |

1/ Deficiency of quality or quantity of blood due to wasting disease, loss of blood and imperfect nutrition, marked by paleness of the skin mucous membranes, loss of energy, palpitation of the heart, etc.
2/ Hypertrophy corneous layer in the skin.
3/ Conjunctivitis, eye disease, due to deficiency of Vitamin A.
4/ A condition marked by leisons on the lips and corners of the mouth.
5/ Inflamation of the tounge.
6/ Absence of reflex contraction of the quardriceps muscle.
7/ Excessive fluid in body tissues.
8/ Abnormally frequent and and loose fecal discharges for a long period.
9/ Abnormal slowness of the heart beat, as evidences by slowing of the pulse rate. This examination included only persons over 13 years of age.
10/ Infrequency of the sanguinous discharge of women 17 - 45 years of age.
11/ Inability of mother of furnish sufficient milk to breast feed her baby under six months.

* Correction February 1950 Survey -Change from 7.8 to 17.0
** Correction February 1950 Survey - Change from 7.0 to 14.6



DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 77 5 013

TOKYO

NUMBER OF PERSONS IN NUTRITION SURVEYS, PERCENT OF DISTRIBUTION ACCORDING TO OVERWEIGHT OR UNDERWEIGHT AND AVERAGE WEIGHT DEVIATION IN KILOGRAMS ABOVE AND BELOW THE TEN PERCENT LEVEL MEASURED FROM THE STANDARD WEIGHT BY AGE GROUPS, NOVEMBER 1950

| Classification | All Ages | Under 2 years | 2 - 5 years | 6 - 10 years | 11 - 15 years | 16 - 20 years | 21 - 30 years | 31 - 40 years | 41 - 50 years | 51 years and over |
|---|---|---|---|---|---|---|---|---|---|---|
| No. persons examined in each age group | 3,986 | 165 | 467 | 608 | 411 | 376 | 550 | 531 | 434 | 444 |
| Percent more than 10% below standard weight | 13.9 | 18.2 | 13.1 | 5.4 | 5.6 | 8.8 | 17.5 | 17.7 | 19.4 | 23.0 |
| Percent between 10% above and below standard weight | 63.4 | 48.5 | 64.6 | 56.9 | 63.3 | 76.3 | 68.0 | 62.9 | 61.5 | 61.9 |
| Percent more than 10% above standard weight | 22.7 | 33.3 | 22.3 | 37.7 | 31.1 | 14.9 | 14.5 | 19.4 | 19.1 | 15.1 |
| Average number of kilograms below standard weight | 4.7 | 1.0 | 1.4 | 2.1 | 3.4 | 5.2 | 5.7 | 5.7 | 5.7 | 5.7 |
| Average number of kilograms above standard weight | 3.9 | 1.2 | 1.5 | 2.7 | 3.9 | 5.2 | 5.6 | 5.7 | 5.7 | 5.4 |



DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 773013

ELEVEN CITIES

NUMBER OF PERSONS IN NUTRITION SURVEYS, PERCENT OF DISTRIBUTION ACCORDING TO OVERWEIGHT OR UNDERWEIGHT AND AVERAGE WEIGHT DEVIATION IN KILOGRAMS ABOVE AND BELOW THE TEN PERCENT LEVEL MEASURED FROM THE STANDARD WEIGHT BY AGE GROUPS, NOVEMBER 1950

| Classification | All Ages | Under 2 years | 2 - 5 years | 6 - 10 years | 11 - 15 years | 16 - 20 years | 21 - 30 years | 31 - 40 years | 41 - 50 years | 51 years and over |
|---|---|---|---|---|---|---|---|---|---|---|
| No. persons examined in each age group | 4,808 | 248 | 530 | 651 | 562 | 451 | 621 | 594 | 560 | 591 |
| Percent more than 10% below standard weight | 13.5 | 14.9 | 11.7 | 6.6 | 5.3 | 7.3 | 16.7 | 16.8 | 18.9 | 22.5 |
| Percent between 10% above and below standard weight | 62.8 | 51.6 | 63.8 | 58.8 | 61.4 | 69.9 | 66.4 | 63.3 | 62.5 | 62.9 |
| Percent more than 10% above standard weight | 23.7 | 33.5 | 24.5 | 34.6 | 33.3 | 22.8 | 16.9 | 19.9 | 18.6 | 14.6 |
| Average number of kilograms below standard weight | 4.6 | 0.9 | 1.4 | 2.0 | 3.6 | 5.1 | 5.6 | 5.6 | 5.7 | 5.6 |
| Average number of kilograms above standard weight | 4.0 | 1.1 | 1.5 | 2.6 | 4.2 | 5.5 | 5.6 | 5.7 | 5.6 | 5.3 |



DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 773013

OTHER CITIES

NUMBER OF PERSONS IN NUTRITION SURVEYS, PERCENT OF DISTRIBUTION ACCORDING TO OVERWEIGHT OR UNDERWEIGHT AND AVERAGE WEIGHT DEVIATION IN KILOGRAMS ABOVE AND BELOW THE TEN PERCENT LEVEL MEASURED FROM THE STANDARD WEIGHT BY AGE GROUPS, NOVEMBER 1950

| Classification | All Ages | Under 2 years | 2 - 5 years | 6 - 10 years | 11 - 15 years | 16 - 20 years | 21 - 30 years | 31 - 40 years | 41 - 50 years | 51 years and over |
|---|---|---|---|---|---|---|---|---|---|---|
| No. persons examined in each age group | 8,152 | 376 | 893 | 1,218 | 1,010 | 674 | 968 | 1,078 | 943 | 992 |
| Percent more than 10% below standard weight | 14.6 | 19.1 | 13.0 | 8.5 | 6.8 | 10.1 | 15.8 | 17.9 | 21.9 | 21.5 |
| Percent between 10% above and below standard weight | 63.6 | 44.7 | 64.6 | 55.1 | 64.1 | 73.6 | 69.3 | 66.3 | 63.8 | 64.2 |
| Percent more than 10% above standard weight | 21.8 | 36.2 | 22.4 | 36.4 | 29.1 | 16.3 | 14.9 | 15.8 | 14.3 | 14.3 |
| Average number of kilograms below standard weight | 4.5 | 1.0 | 1.4 | 2.2 | 3.6 | 5.2 | 5.5 | 5.6 | 5.8 | 5.6 |
| Average number of kilograms above standard weight | 3.8 | 1.1 | 1.5 | 2.8 | 4.0 | 5.3 | 5.5 | 5.6 | 5.6 | 5.3 |



DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 715013

RURAL AREAS

NUMBER OF PERSONS IN NUTRITION SURVEYS, PERCENT OF DISTRIBUTION ACCORDING TO OVERWEIGHT OR UNDERWEIGHT AND AVERAGE WEIGHT DEVIATION IN KILOGRAMS ABOVE AND BELOW THE TEN PERCENT LEVEL MEASURED FROM THE STANDARD WEIGHT BY AGE GROUPS, NOVEMBER 1950

| Classification | All Ages | Under 2 years | 2 - 5 years | 6 - 10 years | 11 - 15 years | 16 - 20 years | 21 - 30 years | 31 - 40 years | 41 - 50 years | 51 years and over |
|---|---|---|---|---|---|---|---|---|---|---|
| No. persons examined in each age group | 18,243 | 946 | 1,961 | 2,482 | 2,238 | 1,643 | 2,272 | 1,997 | 1,912 | 2,792 |
| Percent more than 10% below standard weight | 11.8 | 20.5 | 11.6 | 6.1 | 4.7 | 5.4 | 8.9 | 12.7 | 14.5 | 21.9 |
| Percent between 10% above and below standard weight | 62.6 | 46.7 | 68.3 | 53.2 | 55.1 | 64.7 | 69.1 | 67.3 | 68.2 | 66.0 |
| Percent more than 10% above standard weight | 25.6 | 32.8 | 20.1 | 40.7 | 40.2 | 29.9 | 22.0 | 20.0 | 17.3 | 12.1 |
| Average number of kilograms below standard weight | 4.3 | 1.0 | 1.4 | 2.1 | 3.6 | 5.1 | 5.3 | 5.4 | 5.4 | 5.5 |
| Average number of kilograms above standard weight | 4.1 | 1.1 | 1.5 | 2.9 | 4.2 | 5.7 | 5.7 | 5.7 | 5.5 | 5.3 |



DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775013

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775 013

# DIGEST OF WEEKLY REPORT OF COMMUNICABLE DISEASES IN JAPAN FOR THE WEEK ENDED 17 FEBRUARY 1951

During the seventh week, ended 17 February 1951, there were 21,064 cases of the 31 communicable diseases (exclusive of the four venereal diseases) now being reported compared with 22,822 cases reported for the same diseases last week. Totals for nine diseases (diphtheria, typhoid fever, paratyphoid fever, malaria, scarlet fever, whooping cough, influenza, puerperal infection, and leprosy) were lower this week than in either last week or the seventh week of last year. Five diseases (dysentery, smallpox, epidemic meningitis, measles, and pneumonia) were higher currently than last week or in the corresponding week of 1949. No cases of Japanese "B" encephalitis, cholera, plague, yellow fever, anthrax, glanders, dengue fever, or tsutsugamsuhi were reported during any of the three periods. Present totals for the remaining nine diseases fell between those reported last week and in the seventh week of last year.

Diphtheria cases this week (277) were 6 percent lower than in the preceding week (296). Deaths were about the same this week (28) as last week (27). The current case total was about 5 percent below that (290) for the same week in 1950 and almost 30 percent lower than in the corresponding week of 1949 (381). There were increases over last week in 22 prefectures, decreases in 16, and no change in 5. Kochi, the one remaining prefecture, reported no cases either week. Hokkaido reported the greatest number of cases (25). In the 44 remainig prefectures case totals ranged from 1 to 16. The current and cumulative case rates were 17.2 and 17.4 respectively. Corresponding death rates were 1.7 and 2.2.

The number of dysentery cases this week (220) was 22 percent higher than last week (180), and deaths increased from 27 to 37. The present case figure was almost three times that (75) recorded for the same week of last year and almost seven times the total (31) for the corresponding week of 1949. Case totals increased over last week in 19 prefectures, decreased in 12, and remained the same in 7. There have been no cases for two or more weeks in the 8 remaining prefectures. Tokyo-to continued to report the highest number of cases (61), more than one-fourth of the national total. Among the 31 other prefectures reporting current cases totals ranged from 1 to 17. Of the total dysentery reported this week, 211 cases and all 37 deaths were listed as bacillary and the remaining 9 cases as amebic. The current and cumulative case rates for all dysentery were 13.7 and 11.9 respectively. Corresponding death rates were 2.3 and 1.7.

There were 35 cases of typhoid fever this week, 20 percent less than last week (44), and deaths decreased from 4 to 2. This week's case total was about 40 percent lower than that (59) recorded for the same period last year and 60 percent below the number (88) in the corresponding week of 1949. No cases have been reported for two or more weeks in 21 prefectures. There were decreases from last week in 12 prefectures, increases in 6, and no change in 5. Seventeen prefectures reported cases this week, from one to five each. Current and cumulative case rates were both 2.2. Corresponding death rates were 0.1 and 0.2.

Paratyphoid fever cases decreased from 14 to 12 and deaths from 2 to zero. There were 16 cases in the seventh week of last year and 31 in the corresponding period of 1949. There have been no cases reported for two or more weeks in 29 prefectures, while cases decreased in 9, increased in 6, and remained the same in 2 prefectures. This week's cases occurred in 8 prefectures with from one to four cases each. The current and cumulative case rates were both 0.7. The cumulative death rate was less than 0.1.

There were seven cases of smallpox reported this week, the first to be reported this year. No deaths have been reported so far this year. No cases were recorded for the seventh weeks of 1950 or 1949. All current cases occurred in Hyogo Prefecture. The current and cumulative case rates were 0.4 and 0.1 respectively.

One case of typhus fever was reported from Osaka Prefecture both this week and last week. There have been no deaths reported so far this year. During the seventh week of last year there were 241 cases, and during the same period of

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775 013

1949 there were 4. The current and cumulative case rates were 0.1 and less than 0.1 respectively.

Malaria cases decreased from five to four while deaths increased from zero to three. During the seventh weeks of 1950 and 1949 there were 17 and 30 cases respectively. There have been no cases for two or more weeks in 39 prefectures while there were decreases from last week in 3, increases in 2, and no change in 2. Four prefectures accounted for this week's total. The current and cumulative case rates were 0.2 and 0.4 respectively. Corresponding death rates were 0.2 and 0.1.

There have been no cases of or deaths from Japanese "B" encephalitis for two weeks. There were no cases during the seventh weeks of 1950 or 1949. The cumulative case and death rates as of 17 February 1951 were both less than 0.1.

The number of scarlet fever cases decreased 11 percent, from 79 to 70. Deaths increased from zero to one. Cases were currently 8 percent lower than in the same week of last year (76) and about the same as in the corresponding 1949 period (69). There were decreases from last week's totals in 18 prefectures, increases in 11, and no change in 2, while 15 prefectures have reported no cases for two or more weeks. This week's cases were reported by 20 prefectures with from 1 to 14 each. The current and cumulative case rates were 4.4 and 4.9 respectvily. Corresponding death rates were 0.1 and less than 0.1.

There were 24 cases of epidemic meningitis this week compared with 21 last week. Deaths increased from 4 to 9. During the seventh weeks of 1950 and 1949 cases numbered 22 and 20 respectively. No cases have been reported in about half (24) of the prefectures for two weeks or longer. There were increases over last week, in ten prefectures, decreases in nine, and no change in three. Six of this week's cases were reported from Tokyo-to, the remainder from 14 prefectures with from one to three each. The current and cumulative case rates were 1.5 and 1.3 respectively. Corresponding death rates were 0.6 and 0.3.

The number of measles cases this week (2,121) was 5 percent higher than last week (2,011). It was well over twice the total for the seventh week of last year (935) and slightly higher than in the corresponding period of 1949 (2,066). There were increases over last week in one-half (23) of the prefectures and decreases in 22. The one remaining prefecture, Nara, has reported no cases since last October. The number of cases in Tokyo-to increased from 267 to 356, and in Hyogo from 61 to 131. These two prefectures together with Hokkaido (313) and Kanagawa (287) accounted for slightly more than half of all cases in Japan. In the 40 remaining prefectures reporting current cases totals ranged from 1 to 110. The current and cumulative case rates were 132.0 and 91.0 respectively.

Whooping cough cases (1,264) were 10 percent lower than last week (1,400). They were more than 50 percent below the total (2,725) for the seventh week of 1950 and about 20 percent lower than in the same period of 1949 (1,596). There were decreases from last week in 25 prefectures and increases in 17 while the four remaining prefectures reported the same number both weeks. The largest decreases occurred in Fukuoka (from 100 to 37) and Saitama (112 to 55). Hokkaido (118) accounted for nearly 10 percent of the total cases. Case totals ranged from 3 to 66 in the other 45 prefectures. The current and cumulative case rates were 78.6 and 72.9 respectively.

There were 8,426 cases of tuberculosis reported this week compared with 9,198 last week, a decrease of 8 percent. Cases in the present week were 17 percent higher than those (7,219) in the same period of 1950 and 3 percent higher than in the corresponding week of 1949 (8,184). There were decreases from last week in two-thirds (30) of the prefectures and increases in 15, while one reported the same number both weeks. The most notable decreases occurred in Kanagawa (from 463 to 286)* Yamaguchi (202 to 95), and Fukui (146 to 45). The only comparable increase was in Fukushima Prefecture (from 92 to 230). More than one-fourth of all cases were reported from Tokyo-to (887), Hokkaido (685), and Osaka (673). In the 43 remaining prefectures totals ranged from 25 to 304. Cases of respiratory tuberculosis (7,632) accounted for 91 percent of the total tuberculosis. The current and cumulative case rates for all forms of tuberculosis were 524.3 and 505.6 respectively.

* Hokkaido (794 to 685),

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775 013

The number of pneumonia cases was slightly higher this week (6,124) than last week (6,005). It was nearly 30 percent above the total (4,750) in the seventh week of last year and well over 50 percent above the corresponding figure (3,930) for 1949. There were increases over last week in 27 prefectures and decreases in 19. The most notable increase was in Toyama Prefecture, from 260 to 426, while a comparable decrease occurred in Tokyo-to (from 391 to 290). More than one-fifth of all current cases were reported from Saitama (505), Toyama (426), and Hokkaido (410). In the remaining 43 prefectures totals ranged from 12 to 297. The current and cumulative case rates were 381.1 and 346.5 respectively.

There was a decrease of about 80 percent in the number of influenza cases from 1,374 last week to 265 currently. The present total was more than 90 percent below that (2,997) for the seventh week of last year but nearly seven times the corresponding total (39) for 1949. There have been no cases reported for two or more weeks by 20 prefectures. Totals decreased from last week in 14 prefectures, increased in 10, and remained the same in 2. The most notable decreases were in Gifu (from 944 to 1) and Niigata (350 to 84). The only marked increase (from zero to 104) occurred in Ibaraki Prefecture. Ibaraki and Niigata together accounted for approximately 70 percent of the current total. The remainder was accounted for by 19 prefectures with from 1 to 23 cases each. The current and cumulative case rates were 16.5 and 47.4 respectively.

Poliomyelitis cases this week (30) were slightly lower than last week (32). There were 17 and 21 cases respectively during the corresponding weeks of 1950 and 1949. About one-half (22) of the prefectures have reported no cases for two or more weeks. Increases were reported by 10, decreases by 9 and no change by 5 prefectures. The current cases were reported by 19 prefectures with from one to five each. Current and cumulative case rates were both 1.9.

There were 20 cases of tetanus this week compared with 14 last week. In the seventh weeks of 1950 and 1949 there were 21 and 30 cases respectively. Slightly over half (24) of the prefectures have reported no cases for two weeks or longer, 11 reported increases, 7 decreases, and 4 no change. There were 16 prefectures currently reporting one to three cases each. The current and cumulative case rates were 1.2 and 1.6 respectively.

The number of cases of puerperal infection decreased from 20 last week to 16 currently. During the comparable periods of 1950 and 1949 there were 17 and 25 cases respectively. No cases have been reported for two or more weeks by 28 prefectures, increases over last week by 7, decreases by 6, and no change by 5. Twelve prefectures, all on the island of Honshu, reported the current cases, from one to four each. The current and cumulative case rates were both 1.0.

Rabies cases decreased from one to zero. No cases were recorded for the seventh weeks of 1950 or 1949. The cumulative case rate was 0.1.

There have been no cases of anthrax for four weeks. There were no cases during the seventh weeks of 1950 or 1949. The cumulative case rate as of 17 February 1951 was less than 0.1.

Seven cases of leprosy were reported currently compared with nine last week. There were thirteen and two cases respectively during the corresponding periods of last year and 1949. About three-fourths (35) of the prefectures have reported no cases for two or more weeks. Five prefectures reported increases over last week, four decreases, and two no change. This week's cases were reported by seven prefectures. The current and cumulative case rates were 0.4 and 0.5 respectively.

The number of trachoma cases was slightly higher this week (2,149) than last week (2,115). The current total was 13 percent lower than the corresponding totals for both 1950 (2,463) and 1949 (2,465). One-half (23) of the prefectures reported increases over last week, 22 decreases, and the one remaining prefecture the same number both weeks. Current totals ranged from 2 to 208. The current and cumulative case rates were 133.7 and 111.9 respectively.

There have been no cases of infectious diarrhea for five weeks. During the seventh week of 1950 there were nine cases and during the same week of 1949

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775 013

there was one. The cumulative case rate as of 17 February 1951 was less than 0.1.

Schistosomiasis cases increased from 3 to 8. There were nine cases during the seventh week of 1950. All current cases occurred in Yamanashi Prefecture. The current and cumulative case rates were 0.5 and 0.4 respectively.

There were four cases of filariasis this week compared with none last week. During the seventh week of 1950 there were 5. Three of the cases occurred in Kagoshima and one in Kumamoto Prefecture. The current and cumulative case rates were 0.2 and 0.1 respectively.

The four venereal diseases accounted for 5,096 cases currently compared with 6,049 cases last week. Current and cumulative numbers of syphilis cases were 1,592 and 11,067 respectively; gonorrhea, 3,247 and 22,518; chancroid, 251 and 1,954; and lymphogranuloma venereum, 6 and 4. All totals were lower this week than last week except lymphogranuloma venereum (syphilis, 1,965; gonorrhea, 3,753; chancroid, 326; and lymphogranuloma venereum, 4). Totals for all diseases except gonorrhea were lower this week than in the seventh week of 1950 (syphilis, 2,416; gonorrhea, 2,945; chancroid, 310; and lymphogranuloma venereum, 9). The current and cumulative case rates for each of these diseases as of 17 February 1951 were: syphilis, 99.1 and 98.4 respectively; gonorrhea, 202.0 and 200.2; chancroid, 15.6 and 17.4; and lymphogranuloma venereum, 0.4 and 0.3.

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775 013

# SUMMARY REPORT OF CASES AND DEATHS FROM COMMUNICABLE DISEASES IN JAPAN
## WEEK ENDED 17 FEBRUARY 1951

| PREFECTURE | DIPHTHERIA | | | | DYSENTERY | | | |
|---|---|---|---|---|---|---|---|---|
| | Current | | Cumulative | | Current | | Cumulative | |
| | Cases | Deaths | Cases | Deaths | Cases | Deaths | Cases | Deaths |
| HOKKAIDO | 25 | 1 | 123 | 13 | 7 | - | 37 | 1 |
| AOMORI | 0 | 1 | 41 | 6 | - | - | 7 | - |
| IWATE | 5 | - | 28 | 4 | 3 | - | 13 | 2 |
| MIYAGI | 6 | - | 38 | 4 | 17 | 1 | 46 | 6 |
| AKITA | 7 | - | 50 | 3 | 3 | 1 | 18 | 8 |
| YAMAGATA | 2 | - | 28 | 5 | 2 | - | 25 | - |
| FUKUSHIMA | 15 | 1 | 71 | 12 | 8 | 2 | 29 | 5 |
| IBARAKI | 3 | - | 25 | 2 | 4 | - | 20 | 8 |
| TOCHIGI | 5 | - | 31 | - | - | - | 9 | 2 |
| GUNMA | 5 | - | 15 | 1 | 3 | - | 10 | 1 |
| SAITAMA | 4 | - | 25 | 2 | 16 | 1 | 85 | 12 |
| CHIBA | 2 | 1 | 23 | 6 | 10 | 3 | 61 | 14 |
| TOKYO | 16 | 3 | 136 | 23 | 61 | 9 | 439 | 38 |
| KANAGAWA | 5 | 1 | 36 | 3 | 9 | 2 | 77 | 17 |
| NIIGATA | 6 | - | 102 | 16 | 13 | 5 | 114 | 7 |
| TOYAMA | 4 | - | 31 | 5 | 1 | - | 6 | - |
| ISHIKAWA | 7 | 2 | 30 | 5 | 2 | - | 11 | - |
| FUKUI | 5 | - | 19 | 3 | - | - | - | - |
| YAMANASHI | 2 | - | 5 | 1 | - | - | 2 | - |
| NAGANO | 2 | - | 14 | 1 | 1 | 1 | 8 | 2 |
| GIFU | 7 | - | 25 | 5 | 2 | 1 | 9 | 2 |
| SHIZUOKA | 4 | 1 | 24 | 2 | 1 | - | 19 | 6 |
| AICHI | 3 | - | 44 | 5 | 10 | 1 | 52 | 8 |
| MIE | 4 | 1 | 28 | 8 | - | - | 19 | 3 |
| SHIGA | 1 | - | 12 | 2 | - | - | 1 | - |
| KYOTO | 6 | - | 55 | 3 | 4 | - | 22 | 2 |
| OSAKA | 9 | 1 | 85 | 12 | 7 | 2 | 49 | 11 |
| HYOGO | 15 | 4 | 66 | 9 | 10 | 3 | 40 | 4 |
| NARA | 2 | - | 18 | 4 | - | - | - | - |
| WAKAYAMA | 4 | - | 14 | - | - | - | 3 | 1 |
| TOTTORI | 2 | 1 | 6 | 2 | 1 | - | 4 | - |
| SHIMANE | 3 | - | 20 | 1 | - | - | - | - |
| OKAYAMA | 1 | - | 15 | - | 2 | 1 | 11 | 1 |
| HIROSHIMA | 10 | 3 | 51 | 8 | 1 | - | 11 | 4 |
| YAMAGUCHI | 14 | 2 | 42 | 5 | - | - | 3 | - |
| TOKUSHIMA | 6 | - | 22 | 2 | - | - | 1 | - |
| KAGAWA | 4 | - | 20 | - | 1 | - | 9 | 1 |
| EHIME | 5 | - | 27 | 3 | 6 | 1 | 8 | 2 |
| KOCHI | - | - | 9 | 2 | - | - | 2 | - |
| FUKUOKA | 15 | 1 | 124 | 12 | 4 | - | 20 | 1 |
| SAGA | 2 | - | *46 | 4 | 3 | 2 | 8 | 7 |
| NAGASAKI | 2 | - | 99 | 2 | - | - | 4 | - |
| KUMAMOTO | 8 | 3 | 45 | 11 | 3 | - | 17 | 2 |
| OITA | 5 | - | 48 | 6 | 4 | - | 4 | - |
| MIYAZAKI | 9 | 1 | 69 | 9 | 1 | 1 | 2 | 2 |
| KAGOSHIMA | 2 | - | 70 | 12 | - | - | - | 6 |
| TOTAL | 277 | 28 | *1,957 | 244 | 220 | 37 | 1,335 | 186 |
| RATE | | | | | | | | |
| Current | 17.2 | 1.7 | 17.4 | 2.2 | 13.7 | 2.3 | 11.9 | 1.7 |
| Previous | 18.4 | 1.7 | | | 11.2 | 1.7 | | |

See footnotes at end of table.

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775 013



| PREFECTURE | TYPHOID FEVER | | | | PARATYPHOID FEVER | | | |
|---|---|---|---|---|---|---|---|---|
| | Current | | Cumulative | | Current | | Cumulative | |
| | Cases | Deaths | Cases | Deaths | Cases | Deaths | Cases | Deaths |
| HOKKAIDO | - | - | 2 | - | - | - | 1 | - |
| AOMORI | - | - | 2 | 1 | - | - | 1 | - |
| IWATE | - | - | 4 | 1 | - | - | - | - |
| MIYAGI | 2 | - | 7 | 2 | - | - | 4 | - |
| AKITA | - | - | - | - | - | - | - | - |
| YAMAGATA | - | - | - | - | - | - | 1 | - |
| FUKUSHIMA | - | - | 3 | - | - | - | - | - |
| IBARAKI | - | - | 2 | 1 | 1 | - | 1 | - |
| TOCHIGI | 1 | - | 3 | 1 | - | - | - | - |
| GUNMA | - | - | 3 | - | - | - | - | - |
| SAITAMA | - | - | 7 | 3 | - | - | 3 | - |
| CHIBA | 2 | - | 4 | - | - | - | - | - |
| TOKYO | 5 | - | 31 | 3 | 4 | - | 11 | - |
| KANAGAWA | 1 | - | 6 | - | - | - | 3 | - |
| NIIGATA | 1 | - | 7 | - | - | - | 6 | 1 |
| TOYAMA | - | - | 2 | - | - | - | 1 | - |
| ISHIKAWA | - | - | - | - | 1 | - | 1 | - |
| FUKUI | 3 | - | 5 | - | - | - | 2 | - |
| YAMANASHI | - | - | 3 | - | - | - | - | - |
| NAGANO | - | - | 1 | - | - | - | - | - |
| GIFU | 1 | - | 13 | - | - | - | 1 | - |
| SHIZUOKA | - | - | 8 | - | - | - | 3 | - |
| AICHI | - | 1 | 6 | 1 | - | - | 2 | - |
| MIE | - | - | 8 | 1 | 1 | - | 4 | - |
| SHIGA | - | - | 2 | - | - | - | - | - |
| KYOTO | 5 | - | 16 | - | 2 | - | 3 | 1 |
| OSAKA | 4 | - | 10 | 1 | - | - | 7 | - |
| HYOGO | 2 | - | 10 | 2 | - | - | 3 | - |
| NARA | - | - | 3 | - | - | - | - | - |
| WAKAYAMA | - | - | 1 | 2 | - | - | 2 | - |
| TOTTORI | - | - | 1 | - | - | - | - | - |
| SHIMANE | - | - | 5 | - | 1 | - | 3 | - |
| OKAYAMA | - | - | 5 | 1 | - | - | 1 | - |
| HIROSHIMA | 2 | 1 | 25 | 1 | - | - | 3 | 1 |
| YAMAGUCHI | 1 | - | 3 | - | 1 | - | 1 | - |
| TOKUSHIMA | 2 | - | 5 | - | - | - | 5 | - |
| KAGAWA | - | - | 1 | 1 | 1 | - | 1 | - |
| EHIME | - | - | 2 | - | - | - | - | - |
| KOCHI | 1 | - | 6 | - | - | - | - | - |
| FUKUOKA | 1 | - | 9 | - | - | - | - | - |
| SAGA | - | - | 1 | - | - | - | - | - |
| NAGASAKI | - | - | 7 | - | - | - | - | - |
| KUMAMOTO | - | - | 2 | - | - | - | 2 | - |
| OITA | 1 | - | 3 | - | - | - | - | - |
| MIYAZAKI | - | - | - | - | - | - | 1 | - |
| KAGOSHIMA | - | - | - | - | - | - | 1 | - |
| TOTAL | 35 | 2 | 244 | 22 | 12 | - | 78 | 3 |
| RATE | | | | | | | | |
| Current | 2.2 | 0.1 | 2.2 | 0.2 | 0.7 | - | 0.7 | 0.0 |
| Previous | 2.7 | 0.2 | | | 0.9 | 0.1 | | |

See footnotes at end of table.

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775 013



| PREFECTURE | SMALLPOX Current Cases | Current Deaths | Cumulative Cases | Cumulative Deaths |
|---|---|---|---|---|
| HOKKAIDO | - | - | - | - |
| AOMORI | - | - | - | - |
| IWATE | - | - | - | - |
| MIYAGI | - | - | - | - |
| AKITA | - | - | - | - |
| YAMAGATA | - | - | - | - |
| FUKUSHIMA | - | - | - | - |
| IBARAKI | - | - | - | - |
| TOCHIGI | - | - | - | - |
| GUNMA | - | - | - | - |
| SAITAMA | - | - | - | - |
| CHIBA | - | - | - | - |
| TOKYO | - | - | - | - |
| KANAGAWA | - | - | - | - |
| NIIGATA | - | - | - | - |
| TOYAMA | - | - | - | - |
| ISHIKAWA | - | - | - | - |
| FUKUI | - | - | - | - |
| YAMANASHI | - | - | - | - |
| NAGANO | - | - | - | - |
| GIFU | - | - | - | - |
| SHIZUOKA | - | - | - | - |
| AICHI | - | - | - | - |
| MIE | - | - | - | - |
| SHIGA | - | - | - | - |
| KYOTO | - | - | - | - |
| OSAKA | - | - | - | - |
| HYOGO | 7 | - | 7 | - |
| NARA | - | - | - | - |
| WAKAYAMA | - | - | - | - |
| TOTTORI | - | - | - | - |
| SHIMANE | - | - | - | - |
| OKAYAMA | - | - | - | - |
| HIROSHIMA | - | - | - | - |
| YAMAGUCHI | - | - | - | - |
| TOKUSHIMA | - | - | - | - |
| KAGAWA | - | - | - | - |
| EHIME | - | - | - | - |
| KOCHI | - | - | - | - |
| FUKUOKA | - | - | - | - |
| SAGA | - | - | - | - |
| NAGASAKI | - | - | - | - |
| KUMAMOTO | - | - | - | - |
| OITA | - | - | - | - |
| MIYAZAKI | - | - | - | - |
| KAGOSHIMA | - | - | - | - |
| TOTAL | 7 | - | 7 | - |
| RATE | | | | |
| Current | 0.4 | - | 0.1 | - |
| Previous | - | - | | |

See footnotes at end of table.

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775 013



| PREFECTURE | TYPHUS FEVER | | | | MALARIA | | | |
|---|---|---|---|---|---|---|---|---|
| | Current | | Cumulative | | Current | | Cumulative | |
| | Cases | Deaths | Cases | Deaths | Cases | Deaths | Cases | Deaths |
| HOKKAIDO | - | - | - | - | 1 | - | 4 | - |
| AOMORI | - | - | - | - | - | - | 2 | - |
| IWATE | - | - | - | - | - | - | - | - |
| MIYAGI | - | - | - | - | - | - | - | - |
| AKITA | - | - | - | - | - | - | - | - |
| YAMAGATA | - | - | - | - | - | - | 1 | - |
| FUKUSHIMA | - | - | - | - | - | - | 1 | - |
| IBARAKI | - | - | - | - | - | 1 | - | 1 |
| TOCHIGI | - | - | - | - | - | - | - | - |
| GUNMA | - | - | - | - | 1 | - | 3 | - |
| SAITAMA | - | - | - | - | - | - | - | - |
| CHIBA | - | - | - | - | - | - | 1 | 1 |
| TOKYO | - | - | - | - | 1 | 1 | 1 | 1 |
| KANAGAWA | - | - | - | - | - | - | 4 | - |
| NIIGATA | - | - | - | - | - | - | 1 | - |
| TOYAMA | - | - | - | - | - | - | - | - |
| ISHIKAWA | - | - | - | - | - | - | 5 | 1 |
| FUKUI | - | - | - | - | - | - | 1 | - |
| YAMANASHI | - | - | - | - | - | - | 1 | - |
| NAGANO | - | - | - | - | - | - | 2 | - |
| GIFU | - | - | - | - | - | - | - | - |
| SHIZUOKA | - | - | - | - | - | - | - | - |
| AICHI | - | - | - | - | - | - | 2 | 1 |
| MIE | - | - | - | - | - | - | 3 | - |
| SHIGA | - | - | - | - | - | - | 1 | - |
| KYOTO | - | - | - | - | - | - | - | - |
| OSAKA | 1 | - | 4 | - | - | - | 1 | - |
| HYOGO | - | - | - | - | - | - | 2 | - |
| NARA | - | - | - | - | - | - | - | - |
| WAKAYAMA | - | - | - | - | - | - | - | - |
| TOTTORI | - | - | - | - | - | - | - | - |
| SHIMANE | - | - | - | - | - | - | 1 | - |
| OKAYAMA | - | - | - | - | - | - | 1 | - |
| HIROSHIMA | - | - | - | - | - | - | 2 | 1 |
| YAMAGUCHI | - | - | - | - | - | - | - | - |
| TOKUSHIMA | - | - | - | - | - | - | - | - |
| KAGAWA | - | - | - | - | - | - | - | - |
| EHIME | - | - | - | - | - | - | - | - |
| KOCHI | - | - | - | - | - | - | - | - |
| FUKUOKA | - | - | - | - | - | - | 1 | 1 |
| SAGA | - | - | - | - | - | - | 1 | - |
| NAGASAKI | - | - | - | - | - | - | - | - |
| KUMAMOTO | - | - | - | - | 1 | 1 | 4 | 1 |
| OITA | - | - | - | - | - | - | - | - |
| MIYAZAKI | - | - | - | - | - | - | - | - |
| KAGOSHIMA | - | - | - | - | - | - | - | - |
| TOTAL | 1 | - | 4 | - | 4 | 3 | 46 | 8 |
| RATE | | | | | | | | |
| Current | 0.1 | - | 0.0 | - | 0.2 | 0.2 | 0.4 | 0.1 |
| Previous | 0.1 | - | | | 0.3 | - | | |

See footnotes at end of table.

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775 013

Weekly report - 17 February 1951
Continued

| PREFECTURE | JAPANESE "B" ENCEPHALITIS | | | | SCARLET FEVER | | | |
|---|---|---|---|---|---|---|---|---|
| | Current | | Cumulative | | Current | | Cumulative | |
| | Cases | Deaths | Cases | Deaths | Cases | Deaths | Cases | Deaths |
| HOKKAIDO | - | - | - | - | 7 | - | 28 | - |
| AOMORI | - | - | - | - | 1 | - | 10 | - |
| IWATE | - | - | - | - | - | - | 2 | - |
| MIYAGI | - | - | - | - | 3 | - | 8 | - |
| AKITA | - | - | - | - | 1 | - | 5 | - |
| YAMAGATA | - | - | - | - | 1 | - | 2 | - |
| FUKUSHIMA | - | - | - | - | 1 | - | 5 | - |
| IBARAKI | - | - | - | - | - | - | 3 | - |
| TOCHIGI | - | - | - | - | - | - | 3 | - |
| GUNMA | - | - | - | - | - | - | 2 | - |
| SAITAMA | - | - | - | - | 1 | - | 6 | - |
| CHIBA | - | - | - | - | 1 | - | 4 | - |
| TOKYO | - | - | - | - | 14 | - | 110 | - |
| KANAGAWA | - | - | - | - | 3 | - | 26 | - |
| NIIGATA | - | - | - | - | - | - | 5 | - |
| TOYAMA | - | - | - | - | - | - | 3 | - |
| ISHIKAWA | - | - | - | - | - | - | - | - |
| FUKUI | - | - | - | - | - | - | 2 | - |
| YAMANASHI | - | - | - | - | - | - | 3 | - |
| NAGANO | - | - | - | - | - | - | 13 | - |
| GIFU | - | - | - | - | - | - | 5 | - |
| SHIZUOKA | - | - | - | - | - | - | 10 | 1 |
| AICHI | - | - | - | - | 2 | - | 29 | - |
| MIE | - | - | - | - | - | - | 10 | - |
| SHIGA | - | - | - | - | 3 | - | 11 | - |
| KYOTO | - | - | - | - | 7 | - | 38 | - |
| OSAKA | - | - | - | - | 11 | - | 77 | 1 |
| HYOGO | - | - | - | - | 6 | - | 36 | - |
| NARA | - | - | - | - | - | - | 13 | - |
| WAKAYAMA | - | - | - | - | - | - | 3 | - |
| TOTTORI | - | - | 1 | - | - | - | - | - |
| SHIMANE | - | - | - | - | 1 | - | 4 | - |
| OKAYAMA | - | - | - | - | - | - | 3 | - |
| HIROSHIMA | - | - | - | - | 1 | - | 6 | - |
| YAMAGUCHI | - | - | - | - | - | - | 1 | - |
| TOKUSHIMA | - | - | - | - | - | - | 1 | - |
| KAGAWA | - | - | - | - | 1 | - | 7 | - |
| EHIME | - | - | - | - | 1 | - | 2 | - |
| KOCHI | - | - | - | 1 | - | - | 2 | - |
| FUKUOKA | - | - | 1 | - | 4 | 1 | 27 | 1 |
| SAGA | - | - | - | - | - | - | 1 | - |
| NAGASAKI | - | - | - | - | - | - | 12 | - |
| KUMAMOTO | - | - | - | - | - | - | - | - |
| OITA | - | - | - | - | - | - | - | - |
| MIYAZAKI | - | - | - | - | - | - | 8 | - |
| KAGOSHIMA | - | - | - | - | - | - | 2 | - |
| TOTAL | - | - | 2 | 1 | 70 | 1 | 548 | 3 |
| RATE | | | | | | | | |
| Current | - | - | 0.0 | 0.0 | 4.4 | 0.1 | 4.9 | 0.0 |
| Previous | - | - | | | 4.9 | - | | |

See footnotes at end of table.

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775 013



| PREFECTURE | EPIDEMIC MENINGITIS Current Cases | Current Deaths | Cumulative Cases | Cumulative Deaths | MEASLES Current Cases | Cumulative Cases |
|---|---|---|---|---|---|---|
| HOKKAIDO | 1 | - | 13 | 3 | 313 | 1,711 |
| AOMORI | 1 | - | 4 | 2 | 8 | 31 |
| IWATE | - | - | 1 | - | 25 | 256 |
| MIYAGI | 1 | - | 5 | 1 | 39 | 262 |
| AKITA | - | - | 3 | - | 15 | 102 |
| YAMAGATA | 3 | 1 | 8 | 1 | 10 | 109 |
| FUKUSHIMA | 1 | - | 6 | 2 | 15 | 55 |
| IBARAKI | 1 | 1 | 5 | 2 | 41 | 92 |
| TOCHIGI | - | - | - | - | 16 | 37 |
| GUNMA | - | - | 1 | 1 | 19 | 69 |
| SAITAMA | - | 1 | 2 | 2 | 35 | 121 |
| CHIBA | - | - | - | - | 38 | 77 |
| TOKYO | 6 | 2 | 24 | 7 | 356 | 1,331 |
| KANAGAWA | - | - | 8 | 3 | 207 | 1,350 |
| NIIGATA | - | - | 3 | - | 19 | 113 |
| TOYAMA | 1 | - | 2 | - | 3 | 22 |
| ISHIKAWA | - | - | 1 | - | 36 | 167 |
| FUKUI | - | - | 1 | 1 | 110 | 637 |
| YAMANASHI | - | - | - | - | 6 | 124 |
| NAGANO | 1 | - | 2 | - | 3 | 18 |
| GIFU | - | - | 1 | - | 48 | 308 |
| SHIZUOKA | - | - | 6 | - | 71 | 313 |
| AICHI | - | - | 2 | - | 52 | 203 |
| MIE | 1 | - | 1 | - | 5 | 17 |
| SHIGA | - | - | - | - | 7 | 27 |
| KYOTO | 2 | 2 | 9 | 2 | 1 | 11 |
| OSAKA | 1 | 2 | 16 | 4 | 60 | 185 |
| HYOGO | - | - | 2 | - | 131 | 423 |
| NARA | - | - | - | - | - | - |
| WAKAYAMA | - | - | - | - | 2 | 21 |
| TOTTORI | - | - | - | - | 2 | 8 |
| SHIMANE | 1 | - | 2 | 1 | - | 1 |
| OKAYAMA | 1 | - | 3 | - | 36 | 216 |
| HIROSHIMA | - | - | 1 | 1 | 50 | 153 |
| YAMAGUCHI | - | - | 4 | 2 | 20 | 144 |
| TOKUSHIMA | - | - | - | - | 1 | 7 |
| KAGAWA | - | - | - | - | 22 | 66 |
| EHIME | - | - | - | - | 46 | 278 |
| KOCHI | - | - | 1 | - | 17 | 111 |
| FUKUOKA | 2 | - | 4 | - | 74 | 500 |
| SAGA | - | - | 1 | - | 44 | 207 |
| NAGASAKI | - | - | 2 | - | 14 | 186 |
| KUMAMOTO | - | - | 2 | - | 3 | 10 |
| OITA | - | - | - | - | 1 | 27 |
| MIYAZAKI | - | - | 1 | - | 1 | 13 |
| KAGOSHIMA | - | - | 1 | 1 | 19 | 100 |
| TOTAL | 24 | 9 | 148 | 36 | 2,121 | 10,241 |
| RATE | | | | | | |
| Current | 1.5 | 0.6 | 1.3 | 0.3 | 132.0 | 91.0 |
| Previous | 1.3 | 0.2 | | | 125.1 | |

See footnotes at end of table.

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775 013

Weekly Report - 17 February 1951
Continued

| PREFECTURE | WHOOPING COUGH Current Cases | WHOOPING COUGH Cumulative Cases | TUBERCULOSIS Current Cases | TUBERCULOSIS Cumulative Cases |
|---|---|---|---|---|
| HOKKAIDO | 118 | 499 | 685 | 4,219 |
| AOMORI | 40 | 184 | 129 | 709 |
| IWATE | 30 | 214 | 159 | 983 |
| MIYAGI | 63 | 339 | 182 | 1,102 |
| AKITA | 25 | 178 | 149 | 894 |
| YAMAGATA | 13 | 87 | 139 | 834 |
| FUKUSHIMA | 37 | 237 | 230 | 890 |
| IBARAKI | 15 | 94 | 81 | 546 |
| TOCHIGI | 8 | 143 | 92 | 516 |
| GUNMA | 26 | 174 | 143 | 747 |
| SAITAMA | 55 | 456 | 295 | 1,398 |
| CHIBA | 30 | 115 | 120 | 804 |
| TOKYO | 39 | 315 | 887 | 5,853 |
| KANAGAWA | 30 | 214 | 286 | 2,071 |
| NIIGATA | 41 | 363 | 130 | 1,186 |
| TOYAMA | 30 | 153 | 146 | 1,079 |
| ISHIKAWA | 5 | 60 | 69 | 484 |
| FUKUI | 14 | 170 | 45 | 547 |
| YAMANASHI | 3 | 28 | 32 | 262 |
| NAGANO | 66 | 374 | 253 | 1,293 |
| GIFU | 32 | 177 | 146 | 998 |
| SHIZUOKA | 21 | 196 | 294 | 1,775 |
| AICHI | 55 | 337 | 384 | 2,898 |
| MIE | 17 | 103 | 121 | 843 |
| SHIGA | 35 | 163 | 80 | 550 |
| KYOTO | 17 | 115 | 200 | 1,397 |
| OSAKA | 31 | 178 | 673 | 4,293 |
| HYOGO | 32 | 160 | 319 | 2,379 |
| NARA | 6 | 12 | 38 | 315 |
| WAKAYAMA | 8 | 97 | 76 | 626 |
| TOTTORI | 8 | 50 | 25 | 299 |
| SHIMANE | 39 | 80 | 102 | 514 |
| OKAYAMA | 33 | 171 | 117 | 1,075 |
| HIROSHIMA | 30 | 137 | 174 | 1,494 |
| YAMAGUCHI | 4 | 25 | 95 | 783 |
| TOKUSHIMA | 3 | 22 | 36 | 460 |
| KAGAWA | 10 | 99 | 63 | 398 |
| EHIME | 19 | 131 | 92 | 737 |
| KOCHI | 7 | 93 | 78 | 504 |
| FUKUOKA | 37 | 408 | 363 | 2,900 |
| SAGA | 20 | 149 | 134 | 807 |
| NAGASAKI | 6 | 116 | 134 | 1,240 |
| KUMAMOTO | 47 | 244 | 159 | 942 |
| OITA | 3 | 46 | 90 | 615 |
| MIYAZAKI | 31 | 313 | 91 | 834 |
| KAGOSHIMA | 25 | 180 | 90 | 787 |
| TOTAL | 1,264 | 6,199 | 8,426 | 56,080 |
| RATE | | | | |
| Current | 78.6 | 72.9 | 524.3 | 505.6 |
| Previous | 87.1 | | 572.3 | |

See footnotes at end of table.

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775 013



| PREFECTURE | PNEUMONIA Current Cases | PNEUMONIA Cumulative Cases | INFLUENZA Current Cases | INFLUENZA Cumulative Cases |
|---|---|---|---|---|
| HOKKAIDO | 410 | 2,549 | 23 | 618 |
| AOMORI | 61 | 374 | - | - |
| IWATE | 103 | 653 | - | - |
| MIYAGI | 118 | 844 | - | 10 |
| AKITA | 50 | 442 | - | - |
| YAMAGATA | 126 | 855 | 1 | 17 |
| FUKUSHIMA | 168 | 895 | - | 1 |
| IBARAKI | 139 | 920 | 104 | 105 |
| TOCHIGI | 130 | 907 | - | - |
| GUNMA | 283 | 1,714 | 1 | 100 |
| SAITAMA | 505 | 3,110 | 4 | 128 |
| CHIBA | 136 | 740 | 3 | 125 |
| TOKYO | 290 | 2,510 | 2 | 192 |
| KANAGAWA | 275 | 2,074 | - | - |
| NIIGATA | 180 | 1,280 | 84 | 853 |
| TOYAMA | 426 | 1,936 | 11 | 344 |
| ISHIKAWA | 32 | 326 | - | 7 |
| FUKUI | 33 | 418 | - | 58 |
| YAMANASHI | 47 | 394 | - | 4 |
| NAGANO | 297 | 1,445 | - | 1 |
| GIFU | 160 | 808 | 1 | 1,133 |
| SHIZUOKA | 149 | 901 | 2 | 343 |
| AICHI | 183 | 1,334 | 8 | 192 |
| MIE | 71 | 466 | 2 | 116 |
| SHIGA | 124 | 546 | 2 | 8 |
| KYOTO | 129 | 500 | - | 8 |
| OSAKA | 190 | 889 | - | 10 |
| HYOGO | 159 | 803 | 1 | 77 |
| NARA | 12 | 89 | - | 3 |
| WAKAYAMA | 52 | 252 | - | 43 |
| TOTTORI | 21 | 147 | - | 1 |
| SHIMANE | 15 | 181 | - | 24 |
| OKAYAMA | 100 | 732 | - | 37 |
| HIROSHIMA | 122 | 639 | - | - |
| YAMAGUCHI | 47 | 227 | 6 | 227 |
| TOKUSHIMA | 37 | 197 | - | - |
| KAGAWA | 43 | 305 | 2 | 25 |
| EHIME | 120 | 790 | 1 | 383 |
| KOCHI | 27 | 278 | - | 1 |
| FUKUOKA | 125 | 1,253 | - | 36 |
| SAGA | 86 | 710 | - | - |
| NAGASAKI | 45 | 517 | 3 | 73 |
| KUMAMOTO | 128 | 907 | - | 13 |
| OITA | 37 | 237 | 1 | 4 |
| MIYAZAKI | 93 | 445 | 3 | 12 |
| KAGOSHIMA | 40 | 416 | - | - |
| TOTAL | 6,124 | 38,985 | 265 | 5,332 |
| RATE | | | | |
| Current | 301.1 | 346.5 | 16.5 | 47.4 |
| Previous | 373.6 | | 65.5 | |

See footnotes at end of table.

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775 013



| PREFECTURE | POLIOMYELITIS Current Cases | POLIOMYELITIS Cumulative Cases | TETANUS Current Cases | TETANUS Cumulative Cases |
|---|---|---|---|---|
| HOKKAIDO | 2 | 11 | - | 1 |
| AOMORI | - | - | - | 1 |
| IWATE | - | - | - | 2 |
| MIYAGI | 2 | 13 | - | 2 |
| AKITA | - | 1 | 1 | 4 |
| YAMAGATA | - | 7 | - | 4 |
| FUKUSHIMA | - | 3 | - | - |
| IBARAKI | - | 2 | - | 6 |
| TOCHIGI | - | - | - | 3 |
| GUNMA | - | 3 | 1 | 6 |
| SAITAMA | 3 | 20 | 3 | 9 |
| CHIBA | 1 | 3 | - | 12 |
| TOKYO | - | 16 | 1 | 9 |
| KANAGAWA | - | 3 | - | 2 |
| NIIGATA | 1 | 7 | 3 | 4 |
| TOYAMA | - | 1 | - | - |
| ISHIKAWA | - | - | - | 1 |
| FUKUI | - | 3 | - | - |
| YAMANASHI | - | 3 | 1 | 4 |
| NAGANO | 1 | 2 | - | 4 |
| GIFU | 1 | 3 | 1 | 3 |
| SHIZUOKA | 3 | 6 | 1 | 14 |
| AICHI | - | 2 | 1 | 6 |
| MIE | 1 | 5 | - | 4 |
| SHIGA | 1 | 2 | - | - |
| KYOTO | - | 2 | 1 | 5 |
| OSAKA | 5 | 43 | 1 | 5 |
| HYOGO | 1 | 11 | - | 4 |
| NARA | 1 | 7 | - | 3 |
| WAKAYAMA | 1 | 1 | - | - |
| TOTTORI | - | - | - | - |
| SHIMANE | - | - | - | 2 |
| OKAYAMA | 1 | 4 | - | 3 |
| HIROSHIMA | - | 1 | 1 | 6 |
| YAMAGUCHI | - | - | - | - |
| TOKUSHIMA | - | 5 | - | 2 |
| KAGAWA | - | - | 1 | 2 |
| EHIME | 2 | 7 | 1 | 8 |
| KOCHI | - | - | - | 2 |
| FUKUOKA | 1 | 7 | 1 | 7 |
| SAGA | - | 1 | 1 | 3 |
| NAGASAKI | - | 1 | - | - |
| KUMAMOTO | 1 | 5 | - | 8 |
| OITA | - | - | - | 3 |
| MIYAZAKI | - | 2 | - | 7 |
| KAGOSHIMA | 1 | 3 | - | 7 |
| TOTAL | 30 | 216 | 20 | 180 |
| RATE | | | | |
| Current | 1.9 | 1.9 | 1.2 | 1.6 |
| Previous | 2.0 | | 0.9 | |

See footnotes at end of table.

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775 013



| PREFECTURE | PUERPERAL INFECTION Current Cases | PUERPERAL INFECTION Cumulative Cases | RABIES Current Cases | RABIES Cumulative Cases |
|---|---|---|---|---|
| HOKKAIDO | - | 13 | - | - |
| AOMORI | 1 | 3 | - | - |
| [illegible] | - | - | - | - |
| MIYAGI | - | - | - | - |
| AKITA | - | - | - | - |
| YAMAGATA | - | 2 | - | - |
| FUKUSHIMA | 1 | 2 | - | - |
| IBARAKI | - | 2 | - | 2 |
| TOCHIGI | - | 2 | - | - |
| GUNMA | 1 | 4 | - | 1 |
| SAITAMA | - | 8 | - | 1 |
| CHIBA | - | 3 | - | - |
| TOKYO | 1 | 4 | - | 1 |
| KANAGAWA | - | - | - | 1 |
| NIIGATA | - | 3 | - | - |
| TOYAMA | 1 | 2 | - | - |
| ISHIKAWA | 1 | 1 | - | - |
| FUKUI | - | 1 | - | - |
| YAMANASHI | - | 4 | - | - |
| NAGANO | 4 | 7 | - | - |
| GIFU | 1 | 5 | - | - |
| SHIZUOKA | 1 | 2 | - | 1 |
| AICHI | - | 3 | - | - |
| MIE | - | - | - | - |
| SHIGA | 1 | 6 | - | - |
| KYOTO | - | - | - | - |
| OSAKA | 1 | 3 | - | - |
| HYOGO | - | - | - | - |
| NARA | - | - | - | - |
| WAKAYAMA | - | 1 | - | - |
| TOTTORI | - | - | - | - |
| SHIMANE | - | 2 | - | - |
| OKAYAMA | - | 3 | - | - |
| HIROSHIMA | 2 | 10 | - | - |
| [illegible] | - | - | - | - |
| [illegible] | - | - | - | - |
| [illegible] | - | 4 | - | - |
| EHIME | - | - | - | - |
| KOCHI | - | 3 | - | - |
| [illegible] | - | - | - | - |
| SAGA | - | 1 | - | - |
| KUMAMOTO | - | - | - | - |
| KAGOSHIMA | - | 1 | - | - |
| MIYAZAKI | - | 1 | - | - |
| OITA | - | 1 | - | - |
| NAGASAKI | - | 2 | - | - |
| TOTAL | 16 | 109 | - | 7 |
| RATE | | | | |
| Current | 1.0 | 1.0 | - | 0.1 |
| Previous | 1.2 | | 0.1 | |

See footnotes at end of table.

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775 013



| PREFECTURE | ANTHRAX Current Cases | ANTHRAX Cumulative Cases | LEPROSY Current Cases | LEPROSY Cumulative Cases |
|---|---|---|---|---|
| HOKKAIDO | - | - | - | - |
| AOMORI | - | - | - | - |
| IWATE | - | - | - | - |
| MIYAGI | - | - | - | 3 |
| AKITA | - | - | 1 | 3 |
| YAMAGATA | - | - | - | - |
| FUKUSHIMA | - | - | - | 7 |
| IBARAKI | - | - | - | 2 |
| TOCHIGI | - | - | - | 2 |
| GUNMA | - | - | - | 3 |
| SAITAMA | - | - | - | 1 |
| CHIBA | - | - | - | - |
| TOKYO | - | - | 1 | 3 |
| KANAGAWA | - | - | - | 1 |
| NIIGATA | - | - | - | - |
| TOYAMA | - | - | - | - |
| ISHIKAWA | - | - | - | 1 |
| FUKUI | - | - | - | - |
| YAMANASHI | - | - | - | 2 |
| NAGANO | - | - | 1 | 1 |
| GIFU | - | - | - | - |
| SHIZUOKA | - | - | - | 4 |
| AICHI | - | 1 | - | 4 |
| MIE | - | - | - | 2 |
| SHIGA | - | - | - | - |
| KYOTO | - | - | 1 | 4 |
| OSAKA | - | - | - | 1 |
| HYOGO | - | - | 1 | 4 |
| NARA | - | - | - | - |
| WAKAYAMA | - | - | - | 1 |
| TOTTORI | - | - | - | - |
| SHIMANE | - | - | - | - |
| OKAYAMA | - | - | - | - |
| HIROSHIMA | - | - | - | 1 |
| YAMAGUCHI | - | - | - | - |
| TOKUSHIMA | - | - | 1 | 1 |
| KAGAWA | - | - | - | 3 |
| EHIME | - | - | - | 1 |
| KOCHI | - | - | - | - |
| FUKUOKA | - | - | - | 2 |
| SAGA | - | - | - | - |
| NAGASAKI | - | - | - | 1 |
| KUMAMOTO | - | - | - | - |
| OITA | - | - | - | - |
| MIYAZAKI | - | - | - | - |
| KAGOSHIMA | - | - | 1 | 1 |
| TOTAL | - | 1 | 7 | 59 |
| RATE | | | | |
| Current | - | 0.0 | 0.4 | 0.5 |
| Previous | - | | 0.6 | |

See footnotes at end of table.

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775 013



| PREFECTURE | TRACHOMA Current Cases | TRACHOMA Cumulative Cases | INFECTIOUS DIARRHEA Current Cases | INFECTIOUS DIARRHEA Cumulative Cases |
|---|---|---|---|---|
| HOKKAIDO | 160 | 780 | - | 5 |
| AOMORI | 64 | 422 | - | - |
| IWATE | 76 | 591 | - | - |
| MIYAGI | 74 | 438 | - | - |
| AKITA | 208 | 598 | - | - |
| YAMAGATA | 20 | 160 | - | - |
| FUKUSHIMA | 162 | 296 | - | - |
| IBARAKI | 25 | 258 | - | - |
| TOCHIGI | 14 | 109 | - | - |
| GUNMA | 30 | 231 | - | - |
| SAITAMA | 52 | 322 | - | - |
| CHIBA | 40 | 261 | - | - |
| TOKYO | 46 | 316 | - | - |
| KANAGAWA | 13 | 241 | - | - |
| NIIGATA | 74 | 201 | - | - |
| TOYAMA | 44 | 171 | - | - |
| ISHIKAWA | 2 | 50 | - | - |
| FUKUI | 22 | 120 | - | - |
| YAMANASHI | 33 | 188 | - | - |
| NAGANO | 52 | 209 | - | - |
| GIFU | 45 | 153 | - | - |
| SHIZUOKA | 22 | 165 | - | - |
| AICHI | 203 | 712 | - | - |
| MIE | 14 | 92 | - | - |
| SHIGA | 16 | 61 | - | - |
| KYOTO | 29 | 248 | - | - |
| OSAKA | 88 | 730 | - | - |
| HYOGO | 122 | 786 | - | - |
| NARA | 7 | 70 | - | - |
| WAKAYAMA | 37 | 248 | - | - |
| TOTTORI | 8 | 415 | - | - |
| SHIMANE | 8 | 63 | - | - |
| OKAYAMA | 32 | 395 | - | - |
| HIROSHIMA | 54 | 315 | - | - |
| YAMAGUCHI | 9 | 37 | - | - |
| TOKUSHIMA | 9 | 63 | - | - |
| KAGAWA | 10 | 182 | - | - |
| EHIME | 41 | 177 | - | - |
| KOCHI | 5 | 31 | - | - |
| FUKUOKA | 89 | 745 | - | - |
| SAGA | 17 | 172 | - | - |
| NAGASAKI | 10 | 118 | - | - |
| KUMAMOTO | 22 | 149 | - | - |
| OITA | 5 | 256 | - | - |
| MIYAZAKI | 29 | 179 | - | - |
| KAGOSHIMA | 7 | 63 | - | - |
| TOTAL | 2,149 | 12,587 | - | 5 |
| RATE | | | | |
| Current | 133.7 | 111.9 | - | 0.0 |
| Previous | 131.6 | | - | |

See footnotes at end of table.

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775 013

Weekly Report - 17 February 1951
Continued

| PREFECTURE | SCHISTOSOMIASIS Current Cases | SCHISTOSOMIASIS Cumulative Cases | FILARIASIS Current Cases | FILARIASIS Cumulative Cases |
|---|---|---|---|---|
| HOKKAIDO | - | - | - | - |
| AOMORI | - | - | - | - |
| IWATE | - | - | - | - |
| MIYAGI | - | - | - | - |
| AKITA | - | - | - | - |
| YAMAGATA | - | - | - | - |
| FUKUSHIMA | - | - | - | - |
| IBARAKI | - | - | - | - |
| TOCHIGI | - | - | - | - |
| GUNMA | - | - | - | - |
| SAITAMA | - | - | - | - |
| CHIBA | - | - | - | 1 |
| TOKYO | - | - | - | - |
| KANAGAWA | - | - | - | - |
| NIIGATA | - | - | - | - |
| TOYAMA | - | - | - | - |
| ISHIKAWA | - | - | - | - |
| FUKUI | - | - | - | 1 |
| YAMANASHI | 8 | 37 | - | 1 |
| NAGANO | - | - | - | - |
| GIFU | - | - | - | - |
| SHIZUOKA | - | - | - | - |
| AICHI | - | - | - | - |
| MIE | - | - | - | - |
| SHIGA | - | - | - | - |
| KYOTO | - | - | - | - |
| OSAKA | - | - | - | - |
| HYOGO | - | - | - | - |
| NARA | - | - | - | - |
| WAKAYAMA | - | - | - | - |
| TOTTORI | - | - | - | - |
| SHIMANE | - | - | - | - |
| OKAYAMA | - | - | - | - |
| HIROSHIMA | - | 2 | - | - |
| YAMAGUCHI | - | - | - | - |
| TOKUSHIMA | - | - | - | - |
| KAGAWA | - | - | - | - |
| EHIME | - | - | - | - |
| KOCHI | - | - | - | - |
| FUKUOKA | - | 3 | - | 2 |
| SAGA | - | 2 | - | 1 |
| NAGASAKI | - | - | - | - |
| KUMAMOTO | - | - | 1 | 3 |
| OITA | - | - | - | - |
| MIYAZAKI | - | - | - | - |
| KAGOSHIMA | - | - | 3 | 5 |
| TOTAL | 8 | 44 | 4 | 14 |
| RATE | | | | |
| Current | 0.5 | 0.4 | 0.2 | 0.1 |
| Previous | 0.2 | | | |

See footnotes at end of table.

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775 013

NUMBERS OF CASES AND DEATHS OF COMMUNICABLE DISEASES DURING COMPARABLE PERIODS IN 1949, 1950 AND 1951

| Diseases | Week Ended 17 Feb. 1951 | Week Ended 18 Feb. 1950 | Week Ended 12 Feb. 1949 | Cumulative Number for First 7 Weeks 1951 | Cumulative Number for First 7 Weeks 1950 | Cumulative Number for First 7 Weeks 1949 |
|---|---|---|---|---|---|---|
| **Cases** | | | | | | |
| Diphtheria | 277 | 290 | 381 | 1,957 | 2,187 | 2,624 |
| Dysentery | 220 | 75 | 31 | 1,335 | 511 | 224 |
| Typhoid Fever | 35 | 59 | 88 | 244 | 444 | 719 |
| Paratyphoid Fever | 12 | 16 | 31 | 78 | 130 | 291 |
| Smallpox | 7 | - | - | 7 | 1 | 4 |
| Typhus Fever | 1 | 241 | 4 | 4 | 382 | 40 |
| Malaria | 4 | 17 | 30 | 46 | 76 | 166 |
| Japanese "B" Encephalitis | - | - | - | 2 | - | 1 |
| Scarlet Fever | 70 | 76 | 69 | 548 | 615 | 670 |
| Epidemic Meningitis | 24 | 22 | 20 | 148 | 155 | 159 |
| Cholera | - | - | - | - | - | - |
| Plague | - | - | - | - | - | - |
| Measles | 2,121 | 935 | 2,066 | 10,241 | 6,278 | 9,727 |
| Whooping Cough | 1,264 | 2,725 | 1,596 | 8,199 | 18,890 | 8,638 |
| Tuberculosis | 8,426 | 7,219 | 8,184 | 56,880 | 48,739 | 46,010 |
| Pneumonia | 6,124 | 4,750 | 3,930 | 38,985 | 34,161 | 21,857 |
| Influenza | 265 | 2,997 | 39 | 5,332 | 12,222 | 253 |
| Poliomyelitis | 30 | 17 | 21 | 216 | 209 | 175 |
| Yellow Fever | - | - | - | - | - | - |
| Tetanus | 20 | 21 | 30 | 180 | 180 | 227 |
| Puerperal Infection | 16 | 17 | 25 | 109 | 127 | 154 |
| Rabies | - | - | - | 7 | 8 | 4 |
| Anthrax | - | - | - | 1 | - | 1 |
| Glanders | - | - | - | - | - | - |
| Leprosy | 7 | 13 | 2 | 59 | 54 | 71 |
| Trachoma | 2,149 | 2,463 | 2,465 | 12,587 | 13,680 | 12,866 |
| Infectious Diarrhea | - | 9 | 1 | 5 | 10 | 16 |
| Dengue Fever | - | - | - | - | 1 | 1 |
| Tsutsugamushi Dis. | - | - | NA | - | - | NA |
| Schistosomiasis | 8 | 9 | NA | 44 | 32 | NA |
| Filariasis | 4 | 5 | NA | 14 | 16 | NA |
| **Deaths** | | | | | | |
| Diphtheria | 28 | 29 | 34 | 244 | 231 | 309 |
| Dysentery | 37 | 19 | 5 | 186 | 135 | 70 |
| Typhoid Fever | 2 | 14 | 13 | 22 | 76 | 70 |
| Paratyphoid Fever | - | - | 1 | 3 | 4 | 6 |
| Smallpox | - | - | - | - | - | - |
| Typhus Fever | - | 12 | - | - | 25 | - |
| Malaria | 3 | - | - | 8 | 9 | 5 |
| Japanese "B" Encephalitis | - | - | - | 1 | - | - |
| Scarlet Fever | 1 | - | - | 3 | 4 | 12 |
| Epidemic Meningitis | 9 | 8 | 5 | 36 | 33 | 37 |
| Cholera | - | - | - | - | - | - |
| Plague | - | - | - | - | - | - |

See footnotes at end of table.

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775 013

CASE AND DEATH RATES FOR COMMUNICABLE DISEASES
DURING COMPARABLE PERIODS IN 1949, 1950 AND 1951

| Diseases | Week Ended 17 Feb. 1951 | Week Ended 18 Feb. 1950 | Week Ended 12 Feb. 1949 | Cumulative Rates for First 7 Weeks 1951 | Cumulative Rates for First 7 Weeks 1950 | Cumulative Rates for First 7 Weeks 1949 |
|---|---|---|---|---|---|---|
| **Case Rates** | | | | | | |
| Diphtheria | 17.2 | 18.0 | 24.6 | 17.4 | 19.4 | 24.2 |
| Dysentery | 13.7 | 4.7 | 2.0 | 11.9 | 4.5 | 2.1 |
| Typhoid fever | 2.2 | 3.7 | 5.7 | 2.2 | 3.9 | 6.6 |
| Paratyphoid fever | 0.7 | 1.0 | 2.0 | 0.7 | 1.2 | 2.7 |
| Smallpox | 0.4 | - | - | 0.1 | 0.0 | 0.0 |
| Typhus fever | 0.1 | 15.0 | 0.3 | 0.0 | 3.4 | 0.4 |
| Malaria | 0.2 | 1.1 | 1.9 | 0.4 | 0.7 | 1.5 |
| Japanese "B" encephalitis | - | - | - | 0.0 | - | 0.0 |
| Scarlet fever | 4.4 | 4.7 | 4.4 | 4.9 | 5.5 | 6.2 |
| Epidemic meningitis | 1.5 | 1.4 | 1.3 | 1.3 | 1.4 | 1.5 |
| Cholera | - | - | - | - | - | - |
| Plague | - | - | - | - | - | - |
| Measles | 132.0 | 58.2 | 133.2 | 91.0 | 55.8 | 89.6 |
| Whooping cough | 78.6 | 169.6 | 102.9 | 72.9 | 167.9 | 79.6 |
| Tuberculosis | 524.3 | 449.2 | 527.7 | 505.6 | 433.2 | 423.8 |
| Pneumonia | 381.1 | 295.6 | 253.4 | 346.5 | 303.7 | 201.3 |
| Influenza | 16.5 | 186.5 | 2.5 | 47.4 | 108.6 | 2.3 |
| Poliomyelitis | 1.9 | 1.1 | 1.4 | 1.9 | 1.9 | 1.6 |
| Yellow fever | - | - | - | - | - | - |
| Tetanus | 1.2 | 1.3 | 1.[illegible] | 1.6 | 1.6 | 2.1 |
| Puerperal infection | 1.0 | 1.1 | 1.6 | 1.0 | 1.1 | 1.4 |
| Rabies | - | - | - | 0.1 | 0.1 | 0.0 |
| Anthrax | - | - | - | 0.0 | - | 0.0 |
| Glanders | - | - | - | - | - | - |
| Leprosy | 0.4 | 0.8 | 0.8 | 0.5 | 0.5 | 0.7 |
| Trachoma | 133.7 | 153.3 | 158.9 | 111.9 | 121.6 | 116.5 |
| Infectious diarrhea | - | 0.6 | 0.1[illegible] | 0.0 | 0.1 | 0.1 |
| Dengue fever | - | - | - | - | 0.0 | 0.0 |
| Tsutsugamushi Dis. | - | - | NA | - | - | NA |
| Schistosomiasis | 0.5 | 0.6 | NA | 0.4 | 0.3 | NA |
| Filariasis | 0.2 | 0.3 | NA | 0.1 | 0.1 | NA |
| **Deaths Rates** | | | | | | |
| Diphtheria | 1.7 | 1.8 | 2.2 | 2.2 | 2.1 | 2.8 |
| Dysentery | 3.3 | 1.2 | 0.3 | 1.7 | 1.2 | 0.6 |
| Typhoid fever | 0.1 | 0.9 | 0.8 | 0.2 | 0.7 | 0.6 |
| Paratyphoid fever | - | - | 0.1 | 0.0 | 0.0 | 0.1 |
| Smallpox | - | - | - | - | - | - |
| Typhus fever | - | 0.7 | - | - | 0.2 | - |
| Malaria | 0.2 | - | - | 0.1 | 0.1 | 0.05 |
| Japanese "B" encephalitis | - | - | - | 0.0 | - | - |
| Scarlet fever | 0.1 | - | - | 0.0 | 0.0 | 0.1 |
| Epidemic meningitis | 0.6 | 0.5 | 0.3 | 0.3 | 0.3 | 0.3 |
| Cholera | - | - | - | - | - | - |
| Plague | - | - | - | - | - | - |

See footnotes at end of table.

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775 013



| PREFECTURE | SYPHILIS Current Cases | SYPHILIS Cumulative Cases | GONORRHEA Current Cases | GONORRHEA Cumulative Cases |
|---|---|---|---|---|
| HOKKAIDO | 93 | 538 | 318 | 1,755 |
| AOMORI | 10 | 106 | 34 | 179 |
| IWATE | 12 | 89 | 9 | 80 |
| MIYAGI | 18 | 119 | 40 | 184 |
| AKITA | 17 | 124 | 7 | 71 |
| YAMAGATA | 35 | 181 | 25 | 123 |
| FUKUSHIMA | 25 | 312 | 76 | 344 |
| IBARAKI | 14 | 124 | 7 | 100 |
| TOCHIGI | 12 | 126 | 48 | 204 |
| GUNMA | 28 | 126 | 27 | 207 |
| SAITAMA | 27 | 130 | 40 | 246 |
| CHIBA | 29 | 166 | 11 | 165 |
| TOKYO | 82 | 612 | 229 | 1,685 |
| KANAGAWA | 109 | 772 | 259 | 2,667 |
| NIIGATA | 31 | 329 | 21 | 389 |
| TOYAMA | 11 | *163 | 56 | *338 |
| ISHIKAWA | 26 | 118 | 28 | 208 |
| FUKUI | 16 | 86 | 26 | 214 |
| YAMANASHI | 9 | 52 | 7 | 36 |
| NAGANO | 34 | 155 | 48 | 241 |
| GIFU | 25 | 222 | 64 | 353 |
| SHIZUOKA | 70 | 305 | 87 | 436 |
| AICHI | 41 | 375 | 83 | 724 |
| MIE | 7 | 155 | 15 | 215 |
| SHIGA | 13 | 67 | 29 | 156 |
| KYOTO | 42 | 235 | 62 | 494 |
| OSAKA | 121 | 742 | 112 | 686 |
| HYOGO | 44 | 409 | 107 | 730 |
| NARA | 5 | 33 | 31 | 164 |
| WAKAYAMA | 24 | 117 | 36 | 393 |
| TOTTORI | 24 | 122 | 36 | 179 |
| SHIMANE | 6 | 41 | 11 | 39 |
| OKAYAMA | 20 | 223 | 32 | 314 |
| HIROSHIMA | 37 | 293 | 180 | 1,083 |
| YAMAGUCHI | 50 | 298 | 170 | 756 |
| TOKUSHIMA | 4 | 70 | 1 | 81 |
| KAGAWA | 17 | 77 | 50 | 163 |
| EHIME | 31 | 144 | 16 | 135 |
| KOCHI | 9 | 79 | 9 | 153 |
| FUKUOKA | 140 | 980 | 406 | 3,523 |
| SAGA | 43 | 272 | 60 | 363 |
| NAGASAKI | 90 | 753 | 171 | 843 |
| KUMAMOTO | 23 | 174 | 94 | 410 |
| OITA | 12 | 160 | 30 | 259 |
| MIYAZAKI | 38 | 147 | 15 | 204 |
| KAGOSHIMA | 18 | 146 | 24 | 226 |
| TOTAL | 1,592 | *11,067 | 3,247 | *22,518 |
| RATE | | | | |
| Current | 99.1 | 98.4 | 202.0 | 200.2 |
| Previous | 122.8 | | 233.5 | |

See footnotes at end of table.

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775 013

Weekly Report - 17 February 1951
Continued

| PREFECTURE | CHANCROID Current Cases | CHANCROID Cumulative Cases | LYMPHOGRANULOMA VENEREUM Current Cases | LYMPHOGRANULOMA VENEREUM Cumulative Cases |
|---|---|---|---|---|
| HOKKAIDO | 8 | 67 | - | 1 |
| AOMORI | - | 11 | - | - |
| IWATE | - | 3 | - | - |
| MIYAGI | - | 1 | - | - |
| AKITA | - | 3 | - | - |
| YAMAGATA | 1 | 2 | - | - |
| FUKUSHIMA | 1 | 15 | - | - |
| IBARAKI | 1 | 17 | - | - |
| TOCHIGI | 1 | 14 | - | - |
| GUNMA | 5 | 16 | - | - |
| SAITAMA | 1 | 17 | - | - |
| CHIBA | 2 | 20 | - | 1 |
| TOKYO | 27 | 181 | 2 | 3 |
| KANAGAWA | 38 | 336 | - | 3 |
| NIIGATA | 2 | 25 | - | - |
| TOYAMA | 4 | 19 | 1 | 2 |
| ISHIKAWA | 2 | 12 | - | 2 |
| FUKUI | 3 | 19 | - | 2 |
| YAMANASHI | - | 2 | - | - |
| NAGANO | 2 | 3 | - | - |
| GIFU | 8 | 35 | - | - |
| SHIZUOKA | 4 | 30 | - | 1 |
| AICHI | 7 | 55 | - | - |
| MIE | 1 | 19 | 1 | 1 |
| SHIGA | 6 | 43 | - | - |
| KYOTO | 17 | 118 | - | 4 |
| OSAKA | 18 | 143 | 1 | 3 |
| HYOGO | 5 | 73 | - | 2 |
| NARA | 9 | 35 | - | 2 |
| WAKAYAMA | 1 | 31 | - | 1 |
| TOTTORI | 1 | 17 | - | - |
| SHIMANE | - | 3 | - | - |
| OKAYAMA | 5 | 27 | - | 1 |
| HIROSHIMA | 12 | 100 | - | - |
| YAMAGUCHI | 3 | 18 | - | - |
| TOKUSHIMA | - | 4 | - | - |
| KAGAWA | 2 | 22 | - | 1 |
| EHIME | 1 | 12 | - | - |
| KOCHI | 1 | 17 | - | - |
| FUKUOKA | 26 | 225 | - | 1 |
| SAGA | 3 | 22 | 1 | 1 |
| NAGASAKI | 19 | 86 | - | 1 |
| KUMAMOTO | 3 | 15 | - | 1 |
| OITA | 1 | 14 | - | - |
| MIYAZAKI | - | 3 | - | - |
| KAGOSHIMA | - | 4 | - | - |
| TOTAL | 251 | 1,954 | 6 | 34 |
| RATE | | | | |
| Current | 15.6 | 17.4 | 0.4 | 0.3 |
| Previous | 20.3 | | 0.2 | |

See footnotes at end of table.

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775 013

NUMBERS OF CASES AND CASE RATES FOR
VENEREAL DISEASES IN JAPAN DURING
COMPARABLE PERIODS IN 1949, 1950 AND 1951

| DISEASES | WEEK ENDED 17 Feb. 1951 | WEEK ENDED 18 Feb. 1950 | WEEK ENDED 12 Feb. 1949 | CUMULATIVE NUMBERS FOR FIRST 7 WEEKS 1951 | 1950 | 1949 |
|---|---|---|---|---|---|---|
| NUMBERS | | | | | | |
| SYPHILIS | 1,592 | 2,416 | 4,437 | 11,067 | 16,526 | 24,164 |
| GONORRHEA | 3,247 | 2,945 | 3,596 | 22,518 | 21,094 | 22,274 |
| CHANCROID | 251 | 310 | 563 | 1,954 | 2,245 | 3,673 |
| LYMPHOGRANULOMA VENEREUM | 6 | 9 | 12 | 34 | 81 | 95 |
| RATES | | | | | | |
| SYPHILIS | 99.1 | 150.3 | 286.1 | 98.4 | 146.9 | 222.6 |
| GONORRHEA | 202.0 | 183.2 | 231.9 | 200.2 | 187.5 | 205.2 |
| CHANCROID | 15.6 | 19.3 | 36.3 | 17.4 | 20.0 | 33.8 |
| LYMPHOGRANULOMA VENEREUM | 0.4 | 0.6 | 0.8 | 0.3 | 0.7 | 0.9 |

FOOTNOTES HERE:

1. There were no cases or deaths reported for cholera, plague, yellow fever, glanders, dengue fever, or tsutsugamushi.

2. Rates are the numbers of cases or deaths per 100,000 population, estimated as of 1 July 1950, and are computed on an annual basis.

3. A dash (-) indicates that no cases or deaths were reported and that the case or death rate was zero.

4. A rate of 0.0 indicates that there were some cases or deaths but that the rate was less than 0.1.

5. "NA" indicates that data are not available.

6. * Cumulative figures adjusted for delayed and corrected reports.

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775 013

# DIGEST OF WEEKLY REPORT OF COMMUNICABLE DISEASES IN JAPAN FOR THE WEEK ENDED 24 FEBRUARY 1951

During the eighth week, ended 24 February 1951, there were 21,999 cases of the 31 communicable diseases (exclusive of the four venereal diseases) now being reported compared with 21,082 cases reported for the same diseases last week. Some corrections were received this week for preceding weeks in the current year. Totals for eight diseases (typhoid fever, paratyphoid fever, scarlet fever, whooping cough, influenza, puerperal infection, trachoma and filariasis) were lower this week than in either last week or in the eighth week of last year. Seven diseases (diphtheria, dysentery, epidemic meningitis, measles, tuberculosis, pneumonia and tetanus) were higher currently than last week or in the corresponding week of 1950. Numbers for leprosy were the same in all three periods. No cases of Japanese "B" encephalitis, cholera, plague, yellow fever, anthrax, glanders, infectious diarrhea, dengue fever or tsutsugamushi were reported during any of the three periods. Present totals for the remaining six diseases fell between those reported last week and in the eighth week of last year.

Diphtheria cases this week (314) were 14 percent higher than in the preceding week (276). Deaths increased from 28 to 30 currently. The present case total was 13 percent above that (278) for the same week in 1950 but 21 percent below the corresponding week of 1949 (398). There were increases over last week in 24 prefectures and decreases in 22. Prefectural case totals ranged from zero to 22. The current and cumulative case rates were 19.5 and 17.7 respectively. The corresponding death rates were 1.9 and 2.2.

The number of dysentery cases this week (271) was 23 percent higher than last week (220), while deaths remained the same (37). The present case figure was over two and a half times that (105) recorded for the same week of last year and over eight times the total (32) for the corresponding week of 1949. Case totals increased over last week in 18 prefectures, decreased in 15 and remained the same in 4. There have been no cases for two or more weeks in the nine remaining prefectures. Tokyo-to (62 cases) and Niigata (59) accounted for more than 40 percent of the national total. Among the 30 other prefectures reporting current cases totals ranged from 1 to 20. Of the total dysentery reported this week, 262 cases and 36 deaths were listed as bacillary and the 9 remaining cases and 1 death as amebic. The current and cumulative case rates for all dysentery were 16.9 and 12.5 respectively. Corresponding death rates were 2.3 and 1.7.

There were 34 cases of typhoid fever this week compared with 35 last week. Deaths increased from 2 to 3. This week's case total was 33 percent lower than that (51) recorded for the same period last year and 59 percent below the number (83) in the corresponding week of 1949. No cases have been reported for two or more weeks in almost one-half (22) of the prefectures, while cases decreased in 10, increased in 8 and remained the same in 6. This week's cases occurred in 18 prefectures with from 1 to 6 cases each. The current and cumulative case rates were 2.1 and 2.2 respectively. Corresponding death rates were both 0.2.

Paratyphoid fever cases decreased from 11 to 9. There were no deaths either week. There were 10 cases in the eighth week of last year and 23 in the corresponding period of 1949. No cases have been reported for two or more weeks in 34 prefectures. There were decreases from last week in 6, increases in 5 and no change in 1. Six prefectures, in central Honshu, reported cases this week, one or two each. The current and cumulative case rates were 0.6 and 0.7 respectively, and the cumulative death rate was less than 0.1.

There were two cases of smallpox reported this week compared to seven last week. There was one death currently, the first this year. There were no cases in the corresponding weeks of 1950 and 1949. All current cases and the one death occurred in Hyogo Prefecture. The current and cumulative case rates were both 0.1. The corresponding death rates were 0.1 and less than 0.1 respectively.

One case of typhus fever was reported from Osaka Prefecture both this week and last week. There have been no deaths reported so for this year. During the eighth week of last year there were 113 cases, and during the same period of 1949 there were 6. The current and cumulative case rates were 0.1 and less than 0.1 respectively.

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775 013

Malaria cases increased from four to eight this week while deaths decreased from three to zero. During the eighth weeks of 1950 and 1949 there were 11 and 17 cases respectively. There have been no cases for two or more weeks in 37 prefectures while there were increases from last week in 6, decreases in 2 and no change in 1. Seven prefectures with one or two cases each accounted for this week's total. The current and cumulative case rates were 0.5 and 0.4 respectively. The cumulative death rate was 0.1.

There have been no cases of Japanese "B" encephalitis for three weeks but one death was reported currently compared with none last week. There were no cases during the eighth weeks of 1950 and 1949. The cumulative case rate as of 24 February 1951 was less than 0.1. The current and cumulative death rates were 0.1 and less than 0.1 respectively.

The number of scarlet fever cases decreased 10 percent, from 70 to 63. Deaths remained the same (1). Cases were currently 17 percent lower than in the same week of last year (76) and 29 percent below the corresponding 1949 period (89). Nineteen prefectures have reported no cases for two or more weeks, increases were shown by 10 and decreases by the same number. The remaining 7 prefectures did not change. Twelve cases in Tokyo-to, and 11 in Osaka accounted for more than one-third the present total. The remaining 19 prefectures currently reporting had from 1 to 7 cases each. The current and cumulative case rates were 3.9 and 4.8 respectively. Corresponding death rates were 0.1 and less than 0.1.

There were 29 cases of epidemic meningitis this week compared with 24 last week. Deaths decreased from 9 to 2. During the eighth weeks of 1950 and 1949 cases numbered 20 and 30 respectively. No cases have been reported in 20 prefectures for two weeks or longer. There were increases over last week in 13 prefectures, decreases in 11 and no change in 2. This week's cases were reported by 17 prefectures with from one to seven each. The current and cumulative case rates were 1.8 and 1.4 respectively. Corresponding death rates were 0.1 and 0.3.

The number of measles cases this week (2,911) was 37 percent higher than last week (2,121). It was more than two and a half times the total for the eighth week of last year (1,088) and six percent higher than in the corresponding period of 1949 (2,742). There were increases over last week in 28 prefectures, decreases in 14 and no change in 3. The remaining prefecture, Nara, has reported no cases since last October. All notable changes were increases, and occurred in Tokyo-to (from 356 to 546), Fukuoka (from 74 to 201), Fukui (110 to 209) and Kanagawa (287 to 385). These four prefectures together with Hokkaido (275) accounted for more than half of all cases in Japan. In the 40 remaining prefectures reporting current cases totals ranged from 1 to 141. The current and cumulative case rates were 181.1 and 102.3 respectively.

Whooping cough cases (1,245) were slightly lower than last week (1,264). They were 54 percent below the total (2,679) for the eighth week of 1950 and 17 percent below the same period of 1949 (1,498). There were decreases from last week in 24 prefectures and increases in 22. The only notable change was a decrease from 118 to 63 in Hokkaido. Case totals ranged from 2 to 70 this week. The current and cumulative case rates were 77.5 and 73.5 respectively.

There were 8,778 cases of tuberculosis reported this week compared with 8,426 last week, an increase of 4 percent. Cases in the present week were 10 percent higher than those (7,984) in the same period of 1950 and 6 percent higher than in the corresponding week of 1949 (8,276). There were increases over last week in 24 prefectures, decreases in 21 and no change in 1. The most notable increases were in Tokyo-to (from 887 to 1,183), Fukui Prefecture (from 45 to 217) and Nagasaki (from 134 to 291). The only comparable decrease was in Hokkaido (from 685 to 514). About one-fourth of all cases were reported from Tokyo-to (1,183), Osaka (569) and Hokkaido (514). In the remaining 43 prefectures totals ranged from 41 to 444. Cases of respiratory tuberculosis (7,889) accounted for 90 percent of the total tuberculosis. The current and cumulative case rates for all forms of tuberculosis were 546.2 and 511.2 respectively.

The number of pneumonia cases was slightly higher this week (6,328) than last week (6,124). It was more than 25 percent above the total (4,921) in the eighth week of last year and more than 50 percent above the corresponding figure

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775 013

(4,065) for 1949. There were increases over last week in 29 prefectures and decreases in 17. The most notable changes were decreases in Toyama Prefecture (from 426 to 215), Nagano (297 to 164) and Hokkaido (410 to 319). The most nearly comparable increases occurred in Fukui (33 to 124), Akita (50 to 133), and Mie (71 to 147). Saitama Prefecture (524), Kanagawa (321) and Hokkaido (319) accounted for nearly 20 percent of the national total. In the remaining 43 prefectures totals ranged from 21 to 294. The current and cumulative case rates were 387.5 and 352.4 respectively.

There was a decrease of 65 percent in the number of influenza cases, from 265 last week to 92 currently. The present total was about 95 percent below that (2,096) for the eighth week of last year but approximately 65 percent more than the corresponding total (55) for 1949. There have been no cases reported for two or more weeks by 20 prefectures. Totals decreases in 12, increased in 11 and remained the same in 3. The most notable decrease was in Ibaraki Prefecture (from 104 to zero). Niigata Prefecture reported the highest number of cases (22). Totals in the remaining 20 prefectures reporting cases this week ranged from 1 to 11. The current and cumulative case rates were 5.7 and 42.2 respectively.

Poliomyelitis cases this week increased from 30 to 37. There were 38 and 28 cases respectively in the eighth weeks of 1950 and 1949. Eighteen prefectures have reported no cases for two or more weeks, 14 increases, 11 decreases and 3 no change. The current cases were reported by 17 prefectures with from one to seven each. Current and cumulative case rates were 2.3 and 2.0 respectively.

There were 26 cases of tetanus this week compared with 20 last week. In the eighth weeks of 1950 and 1949 there were 23 and 41 cases respectively. Almost half (22) of the prefectures have reported no cases for at least two weeks. Twelve reported increases, nine decreases and three no change. There were 16 prefectures currently reporting one to three cases each. The current and cumulative case rates were both 1.6.

The number of cases of puerperal infection decreased from 16 last week to 9 currently. During the comparable periods of 1950 and 1949 there were 16 and 13 cases respectively. Two-thirds (30) of the prefectures have reported no cases for two or more weeks, 10 decreases, 5 increases and 1 no change. Seven prefectures, all on the island of Honshu, reported the current cases, with one or two each. The current and cumulative case rates were 0.6 and 0.9 respectively.

There were no rabies cases this week or last week, but one each in the corresponding weeks of 1950 and 1949. The cumulative case rate as of 24 February 1951 was 0.1.

There have been no cases of anthrax for five weeks and there were no cases in the eighth weeks of 1950 and 1949. The cumulative case rate was less than 0.1.

Seven cases of leprosy were reported this week, last week, and in the eighth week of 1950. In the corresponding period of 1949 there were 10 cases. About three-fourths (35) of the prefectures have reported no cases for at least two weeks. Six prefectures reported decreases and five increases. This week's cases were reported by five prefectures, one to three cases each. The current and cumulative case rates were 0.4 and 0.5 respectively.

The number of trachoma cases was 10 percent lower this week (1,927) than last week (2,149). The current total was about 25 percent lower than the corresponding total for 1950 (2,488) and approximately 15 percent below that (2,213) of 1949 for the same period. There were increases in one-half (23) of the prefectures and decreases in 22. Numbers did not change in the one remaining prefecture. Current totals ranged from 4 to 198. The current and cumulative case rates were 119.9 and 112.8 respectively.

There have been no cases of infectious diarrhea for six weeks. During the eighth week of 1950 there were also no cases, but 8 cases were reported during the same period of 1949. The cumulative case rate as of 24 February 1951 was less than 0.1

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775 013

Schistosomiasis cases decreased from 8 to 6. There were 5 cases during the eighth week of 1950. Yamanashi and Fukuoka with three cases each accounted for the current total. The current and cumulative case rates were both 0.4.

There were two cases of filariasis this week compared with 4 last week and 5 in the eighth week of 1950. Saga and Nagasaki Prefectures reported one case each. The current and cumulative case rates were both 0.1.

The four venereal diseases accounted for 6,056 cases currently compared with 5,096 cases last week. Current and cumulative numbers of syphilis cases were 1,874 and 12,939 respectively; gonorrhea, 3,869 and 26,406; chancroid, 301 and 2,257; and lymphogranuloma venereum, 12 and 46. All totals were higher this week than last week (syphilis, 1,592; gonorrhea, 3,247; chancroid, 251; and lymphogranuloma venereum, 6). Totals for syphilis and chancroid were lower this week than in the eighth week of 1950 (2,568 and 305 respectively), gonorrhea and lymphogranuloma were higher (3,024 and 8). The current and cumulative case rates for each of these diseases as of 24 February 1951 were: syphilis, 116.6 and 100.6 respectively; gonorrhea, 240.7 and 205.4; chancroid, 18.7 and 17.6; and lymphogranuloma venereum, 0.7 and 0.4.

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775 013

SUMMARY REPORT OF CASES AND DEATHS FROM
COMMUNICABLE DISEASES IN JAPAN
WEEK ENDED 24 FEBRUARY 1951

| PREFECTURE | DIPHTHERIA | | | | DYSENTERY | | | |
|---|---|---|---|---|---|---|---|---|
| | Current | | Cumulative | | Current | | Cumulative | |
| | Cases | Deaths | Cases | Deaths | Cases | Deaths | Cases | Deaths |
| HOKKAIDO | 20 | 2 | 143 | 15 | 5 | - | 42 | 1 |
| AOMORI | 4 | 1 | 45 | 7 | 9 | - | *13 | - |
| IWATE | 3 | 1 | 31 | 5 | - | 1 | 13 | 3 |
| MIYAGI | 10 | - | 48 | 4 | 2 | 1 | 48 | 7 |
| AKITA | 5 | 1 | 55 | 4 | 3 | - | 21 | 8 |
| YAMAGATA | 1 | - | 29 | 5 | 6 | 1 | 31 | 1 |
| FUKUSHIMA | 9 | 1 | 80 | 13 | 7 | - | 36 | 5 |
| IBARAKI | 1 | - | 26 | 2 | 2 | 3 | 22 | 11 |
| TOCHIGI | 3 | - | 34 | - | 8 | 4 | 17 | 6 |
| GUMMA | 2 | 1 | 17 | 2 | 6 | - | 16 | 1 |
| SAITAMA | 9 | 1 | *35 | 3 | 20 | 5 | *110 | 17 |
| CHIBA | 5 | 1 | 28 | 7 | 5 | 1 | 66 | 15 |
| TOKYO | 21 | - | 159 | 23 | 62 | 4 | 501 | 42 |
| KANAGAWA | 6 | 1 | 42 | 4 | 10 | 3 | 87 | 20 |
| NIIGATA | 12 | - | 114 | 16 | 59 | 4 | 173 | 11 |
| TOYAMA | 6 | 1 | 37 | 6 | 2 | - | 8 | - |
| ISHIKAWA | 2 | 1 | 32 | 6 | 4 | - | 15 | - |
| FUKUI | - | 1 | 19 | 4 | 1 | - | 1 | - |
| YAMANASHI | 1 | - | 6 | 1 | - | - | 2 | - |
| NAGANO | 4 | 1 | 18 | 2 | 2 | - | 10 | 2 |
| GIFU | 1 | 1 | 26 | 6 | 2 | - | 11 | 2 |
| SHIZUOKA | 1 | 2 | 25 | 4 | 5 | 1 | 24 | 7 |
| AICHI | 10 | 1 | 54 | 6 | 19 | 4 | 71 | 12 |
| MIE | 10 | - | 38 | 8 | 1 | - | 20 | 3 |
| SHIGA | - | - | 12 | 2 | - | - | 1 | - |
| KYOTO | 9 | - | *63 | 3 | 4 | - | 26 | 2 |
| OSAKA | 11 | - | 96 | 12 | 2 | - | 51 | 11 |
| HYOGO | 12 | 1 | 78 | 10 | 3 | 1 | 43 | 5 |
| NARA | 6 | 1 | *23 | 5 | 1 | - | 1 | - |
| WAKAYAMA | 5 | 1 | 19 | 1 | - | - | 3 | 1 |
| TOTTORI | - | - | 6 | 2 | - | - | 4 | - |
| SHIMANE | 5 | - | *24 | 1 | - | - | - | - |
| OKAYAMA | - | - | 15 | - | 1 | 1 | 12 | 2 |
| HIROSHIMA | 13 | 3 | 64 | 11 | 4 | 1 | 15 | 5 |
| YAMAGUCHI | 7 | - | 49 | 5 | - | - | 3 | - |
| TOKUSHIMA | 3 | - | 25 | 2 | - | - | 1 | - |
| KAGAWA | 2 | - | 22 | - | - | - | 9 | 1 |
| EHIME | 9 | - | 36 | 3 | 1 | - | 9 | 2 |
| KOCHI | 4 | - | 13 | 2 | - | - | 2 | - |
| FUKUOKA | 22 | - | *153 | 12 | 13 | - | 33 | 1 |
| SAGA | 7 | - | 53 | 4 | - | - | 8 | 7 |
| NAGASAKI | 12 | 1 | 111 | 3 | - | - | 4 | - |
| KUMAMOTO | 11 | - | *53 | *14 | 1 | 1 | 18 | 3 |
| OITA | 17 | - | 65 | 6 | - | - | 4 | - |
| MIYAZAKI | 6 | 5 | *74 | 14 | 1 | 1 | 3 | 3 |
| KAGOSHIMA | 7 | - | 77 | 12 | - | - | - | 6 |
| TOTAL | 314 | 30 | *2,272 | *277 | 271 | 37 | *1,608 | 223 |
| RATE | | | | | | | | |
| Current | 19.5 | 1.9 | 17.7 | 2.2 | 16.9 | 2.3 | 12.5 | 1.7 |
| Previous | 17.2 | 1.7 | | | 13.7 | 2.3 | | |

See footnotes at end of table.

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775 013

Weekly Reports - 24 February 1951
Continued

| PREFECTURE | TYPHOID FEVER Current Cases | Current Deaths | Cumulative Cases | Cumulative Deaths | PARATYPHOID FEVER Current Cases | Current Deaths | Cumulative Cases | Cumulative Deaths |
|---|---|---|---|---|---|---|---|---|
| HOKKAIDO | 1 | - | 3 | - | - | - | 1 | - |
| AOMORI | - | - | 2 | 1 | - | - | 1 | - |
| IWATE | - | - | 4 | 1 | - | - | - | - |
| MIYAGI | 2 | 2 | 9 | 4 | - | - | 4 | - |
| AKITA | - | - | - | - | - | - | - | - |
| YAMAGATA | - | - | - | - | - | - | 1 | - |
| FUKUSHIMA | - | - | 3 | - | - | - | - | - |
| IBARAKI | - | - | 2 | 1 | - | - | 1 | - |
| TOCHIGI | 1 | - | 4 | 1 | - | - | - | - |
| GUMMA | - | - | 3 | - | - | - | - | - |
| SAITAMA | 1 | - | 8 | 3 | - | - | 3 | - |
| CHIBA | 1 | - | 5 | - | - | - | - | - |
| TOKYO | 4 | - | 35 | 3 | - | - | 11 | - |
| KANAGAWA | 1 | - | 7 | - | 2 | - | 5 | - |
| NIIGATA | - | - | *8 | - | 2 | - | *7 | 1 |
| TOYAMA | - | - | 2 | - | - | - | 1 | - |
| ISHIKAWA | - | - | - | - | - | - | 1 | - |
| FUKUI | - | - | 5 | - | - | - | 2 | - |
| YAMANASHI | - | - | 3 | - | - | - | - | - |
| NAGANO | - | - | 1 | - | - | - | - | - |
| GIFU | - | - | 13 | - | 1 | - | 2 | - |
| SHIZUOKA | - | - | 8 | - | 2 | - | 5 | - |
| AICHI | 3 | - | 9 | 1 | - | - | 2 | - |
| MIE | - | - | 8 | 1 | 1 | - | 5 | - |
| SHIGA | - | - | 2 | - | - | - | - | - |
| KYOTO | 1 | - | 17 | - | - | - | 3 | 1 |
| OSAKA | 4 | - | 14 | 1 | 1 | - | 8 | - |
| HYOGO | 1 | - | 11 | 2 | - | - | 3 | - |
| NARA | - | - | 3 | - | - | - | - | - |
| WAKAYAMA | 2 | - | 3 | 2 | - | - | 2 | - |
| TOTTORI | - | - | 1 | - | - | - | - | - |
| SHIMANE | 1 | - | 6 | - | - | - | *2 | - |
| OKAYAMA | 1 | - | 6 | 1 | - | - | 1 | - |
| HIROSHIMA | 6 | 1 | 31 | 2 | - | - | 3 | 1 |
| YAMAGUCHI | 1 | - | 4 | *1 | - | - | 1 | - |
| TOKUSHIMA | 2 | - | 7 | - | - | - | *4 | - |
| KAGAWA | - | - | 1 | 1 | - | - | 1 | - |
| EHIME | 1 | - | 3 | - | - | - | - | - |
| KOCHI | - | - | 6 | - | - | - | - | - |
| FUKUOKA | - | - | 9 | - | - | - | - | - |
| SAGA | - | - | 1 | - | - | - | - | - |
| NAGASAKI | - | - | 7 | - | - | - | - | - |
| KUMAMOTO | - | - | 2 | - | - | - | 2 | - |
| OITA | - | - | 3 | - | - | - | - | - |
| MIYAZAKI | - | - | - | - | - | - | 1 | - |
| KAGOSHIMA | - | - | - | - | - | - | 1 | - |
| TOTAL | 34 | 3 | *279 | *26 | 9 | - | *84 | 3 |
| RATE | | | | | | | | |
| Current | 2.1 | 0.2 | 2.2 | 0.2 | 0.6 | - | 0.7 | 0.0 |
| Previous | 2.2 | 0.1 | | | 0.7 | - | | |

See footnotes at end of teble.

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775 013

Weekly Report - 24 February 1951
Continued

| PREFECTURE | SMALLPOX | | | | TYPHUS FEVER | | | |
|---|---|---|---|---|---|---|---|---|
| | Current | | Cumulative | | Current | | Cumulative | |
| | Cases | Deaths | Cases | Deaths | Cases | Deaths | Cases | Deaths |
| HOKKAIDO | - | - | - | - | - | - | - | - |
| AOMORI | - | - | - | - | - | - | - | - |
| IWATE | - | - | - | - | - | - | - | - |
| MIYAGI | - | - | - | - | - | - | - | - |
| AKITA | - | - | - | - | - | - | - | - |
| YAMAGATA | - | - | - | - | - | - | - | - |
| FUKUSHIMA | - | - | - | - | - | - | - | - |
| IBARAKI | - | - | - | - | - | - | - | - |
| TOCHIGI | - | - | - | - | - | - | - | - |
| GUMMA | - | - | - | - | - | - | - | - |
| SAITAMA | - | - | - | - | - | - | - | - |
| CHIBA | - | - | - | - | - | - | - | - |
| TOKYO | - | - | - | - | - | - | - | - |
| KANAGAWA | - | - | - | - | - | - | - | - |
| NIIGATA | - | - | - | - | - | - | - | - |
| TOYAMA | - | - | - | - | - | - | - | - |
| ISHIKAWA | - | - | - | - | - | - | - | - |
| FUKUI | - | - | - | - | - | - | - | - |
| YAMANASHI | - | - | - | - | - | - | - | - |
| NAGANO | - | - | - | - | - | - | - | - |
| GIFU | - | - | - | - | - | - | - | - |
| SHIZUOKA | - | - | - | - | - | - | - | - |
| AICHI | - | - | - | - | - | - | - | - |
| MIE | - | - | - | - | - | - | - | - |
| SHIGA | - | - | - | - | - | - | - | - |
| KYOTO | - | - | - | - | - | - | - | - |
| OSAKA | - | - | - | - | 1 | - | *3 | - |
| HYOGO | 2 | 1 | 9 | 1 | - | - | - | - |
| NARA | - | - | - | - | - | - | - | - |
| WAKAYAMA | - | - | - | - | - | - | - | - |
| TOTTORI | - | - | - | - | - | - | - | - |
| SHIMANE | - | - | - | - | - | - | - | - |
| OKAYAMA | - | - | - | - | - | - | - | - |
| HIROSHIMA | - | - | - | - | - | - | - | - |
| YAMAGUCHI | - | - | - | - | - | - | - | - |
| TOKUSHIMA | - | - | - | - | - | - | - | - |
| KAGAWA | - | - | - | - | - | - | - | - |
| EHIME | - | - | - | - | - | - | - | - |
| KOCHI | - | - | - | - | - | - | - | - |
| FUKUOKA | - | - | - | - | - | - | - | - |
| SAGA | - | - | - | - | - | - | - | - |
| NAGASAKI | - | - | - | - | - | - | - | - |
| KUMAMOTO | - | - | - | - | - | - | - | - |
| OITA | - | - | - | - | - | - | - | - |
| MIYAZAKI | - | - | - | - | - | - | - | - |
| KAGOSHIMA | - | - | - | - | - | - | - | - |
| TOTAL | 2 | 1 | 9 | 1 | 1 | - | *3 | - |
| RATE | | | | | | | | |
| Current | 0.1 | 0.1 | 0.1 | 0.0 | 0.1 | - | 0.0 | - |
| Previous | 0.4 | - | | | 0.1 | - | | |

See footnotes at end of table.

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775 013



| PREFECTURE | MALARIA | | | | JAPANESE "B" ENCEPHALITIS | | | |
|---|---|---|---|---|---|---|---|---|
| | Current | | Cumulative | | Current | | Cumulative | |
| | Cases | Deaths | Cases | Deaths | Cases | Deaths | Cases | Deaths |
| HOKKAIDO | - | - | 4 | - | - | - | - | - |
| AOMORI | - | - | 2 | - | - | - | - | - |
| IWATE | - | - | - | - | - | - | - | - |
| MIYAGI | - | - | - | - | - | - | - | - |
| AKITA | - | - | - | - | - | - | - | - |
| YAMAGATA | - | - | 1 | - | - | - | - | - |
| FUKUSHIMA | - | - | 1 | - | - | - | - | - |
| IBARAKI | 1 | - | 1 | 1 | - | - | - | - |
| TOCHIGI | - | - | - | - | - | - | - | - |
| GUMMA | 1 | - | 4 | - | - | - | - | - |
| SAITAMA | 1 | - | 1 | - | - | - | - | - |
| CHIBA | - | - | 1 | 1 | - | - | - | - |
| TOKYO | 2 | - | 3 | 1 | - | - | - | - |
| KANAGAWA | - | - | 4 | - | - | - | - | - |
| NIIGATA | - | - | 1 | - | - | - | - | - |
| TOYAMA | - | - | - | - | - | - | - | - |
| ISHIKAWA | - | - | 5 | 1 | - | - | - | - |
| FUKUI | - | - | 1 | - | - | - | - | - |
| YAMANASHI | - | - | 1 | - | - | - | - | - |
| NAGANO | - | - | 2 | - | - | - | - | - |
| GIFU | - | - | - | - | - | - | - | - |
| SHIZUOKA | - | - | - | - | - | 1 | - | 1 |
| AICHI | - | - | 2 | 1 | - | - | - | - |
| MIE | - | - | 3 | - | - | - | - | - |
| SHIGA | - | - | 1 | - | - | - | - | - |
| KYOTO | 1 | - | 1 | - | - | - | - | - |
| OSAKA | 1 | - | 2 | - | - | - | - | - |
| HYOGO | - | - | 2 | - | - | - | - | - |
| NARA | - | - | - | - | - | - | - | - |
| WAKAYAMA | - | - | - | - | - | - | - | - |
| TOTTORI | - | - | - | - | - | - | * - | - |
| SHIMANE | - | - | 1 | - | - | - | - | - |
| OKAYAMA | - | - | 1 | - | - | - | - | - |
| HIROSHIMA | 1 | - | 3 | 1 | - | - | - | - |
| YAMAGUCHI | - | - | - | - | - | - | - | - |
| TOKUSHIMA | - | - | - | - | - | - | - | - |
| KAGAWA | - | - | - | - | - | - | - | - |
| EHIME | - | - | - | - | - | - | - | - |
| KOCHI | - | - | - | - | - | - | - | 1 |
| FUKUOKA | - | - | 1 | 1 | - | - | 1 | - |
| SAGA | - | - | 1 | - | - | - | - | - |
| NAGASAKI | - | - | - | - | - | - | - | - |
| KUMAMOTO | - | - | 4 | 1 | - | - | - | - |
| OITA | - | - | - | - | - | - | - | - |
| MIYAZAKI | - | - | - | - | - | - | - | - |
| KAGOSHIMA | - | - | - | - | - | - | - | - |
| TOTAL | 8 | - | 54 | 8 | - | 1 | *1 | 2 |
| RATE | | | | | | | | |
| Current | 0.5 | - | 0.4 | 0.1 | - | 0.1 | 0.0 | 0.0 |
| Previous | 0.2 | 0.2 | | | - | - | | |

See footnotes at end of table.

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775 013

Weekly Report - 24 February 1951
Continued

| PREFECTURE | SCARLET FEVER | | | | EPIDEMIC MENINGITIS | | | |
|---|---|---|---|---|---|---|---|---|
| | Current | | Cumulative | | Current | | Cumulative | |
| | Cases | Deaths | Cases | Deaths | Cases | Deaths | Cases | Deaths |
| HOKKAIDO | 1 | - | *28 | - | - | - | 13 | 3 |
| AOMORI | - | - | 10 | - | - | - | 4 | 2 |
| IWATE | 1 | - | 3 | - | - | - | 1 | - |
| MIYAGI | - | - | 8 | - | 3 | 1 | 8 | 2 |
| AKITA | 1 | - | 6 | - | 2 | - | 5 | - |
| YAMAGATA | - | - | 2 | - | 1 | - | 9 | 1 |
| FUKUSHIMA | 1 | - | 6 | - | 1 | - | 7 | 2 |
| IBARAKI | - | - | 3 | - | - | - | 5 | 2 |
| TOCHIGI | - | - | 3 | - | - | - | - | - |
| GUMMA | - | - | 2 | - | 1 | - | 2 | 1 |
| SAITAMA | - | - | 6 | - | - | - | 2 | 2 |
| CHIBA | 1 | - | 5 | - | 1 | - | 1 | - |
| TOKYO | 12 | - | 122 | - | 7 | - | 31 | 7 |
| KANAGAWA | 3 | - | 29 | - | 2 | - | 10 | 3 |
| NIIGATA | - | - | 5 | - | 1 | - | 4 | - |
| TOYAMA | - | - | 3 | - | - | - | 2 | - |
| ISHIKAWA | - | - | - | - | - | - | 1 | - |
| FUKUI | 1 | - | 3 | - | - | - | 1 | 1 |
| YAMANASHI | - | - | *5 | - | - | - | - | - |
| NAGANO | 1 | - | 14 | - | - | - | 2 | - |
| GIFU | - | - | 5 | - | - | - | 1 | - |
| SHIZUOKA | 1 | - | 11 | 1 | - | - | 6 | - |
| AICHI | 5 | - | 34 | - | 2 | - | *5 | - |
| MIE | 1 | - | 11 | - | - | - | 1 | - |
| SHIGA | 3 | - | 14 | - | - | - | - | - |
| KYOTO | 6 | - | 44 | - | 1 | - | 10 | 2 |
| OSAKA | 11 | - | 88 | 1 | 1 | - | 17 | 4 |
| HYOGO | 7 | - | 43 | - | 1 | - | 3 | - |
| NARA | - | - | 13 | - | - | - | - | - |
| WAKAYAMA | - | - | 3 | - | - | - | - | - |
| TOTTORI | - | - | - | - | 1 | - | *2 | - |
| SHIMANE | - | - | 4 | - | - | - | 2 | 1 |
| OKAYAMA | - | - | 3 | - | - | - | 3 | - |
| HIROSHIMA | 2 | - | 8 | - | 2 | - | 3 | 1 |
| YAMAGUCHI | - | - | 1 | - | - | - | 4 | 2 |
| TOKUSHIMA | 1 | - | 2 | - | - | - | - | - |
| KAGAWA | 1 | - | 8 | - | - | - | - | - |
| EHIME | - | - | 2 | - | - | - | - | - |
| KOCHI | - | - | 2 | - | 1 | - | 2 | - |
| FUKUOKA | 2 | - | *30 | 1 | - | 1 | *6 | 1 |
| SAGA | - | - | 1 | - | - | - | 1 | - |
| NAGASAKI | - | - | 12 | - | 1 | - | 3 | - |
| KUMAMOTO | - | - | - | - | - | - | 2 | - |
| OITA | - | - | - | - | - | - | - | - |
| MIYAZAKI | 1 | 1 | 9 | 1 | - | - | 1 | - |
| KAGOSHIMA | - | - | 2 | - | - | - | 1 | 1 |
| TOTAL | 63 | 1 | *613 | 4 | 29 | 2 | *181 | 38 |
| RATE | | | | | | | | |
| Current | 3.9 | 0.1 | 4.8 | 0.0 | 1.8 | 0.1 | 1.4 | 0.3 |
| Previous | 4.4 | 0.1 | | | 1.5 | 0.6 | | |

See footnotes at end of table.

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775 013



| PREFECTURE | MEASLES | | WHOOPING COUGH | |
|---|---|---|---|---|
| | Current Cases | Cumulative Cases | Current Cases | Cumulative Cases |
| HOKKAIDO | 275 | 1,986 | 63 | 562 |
| AOMORI | 6 | 37 | 12 | 196 |
| IWATE | 38 | 294 | 13 | 227 |
| MIYAGI | 59 | 321 | 52 | 391 |
| AKITA | 11 | 113 | 19 | 197 |
| YAMAGATA | 13 | 122 | 15 | 102 |
| FUKUSHIMA | 24 | 79 | 28 | 265 |
| IBARAKI | 21 | 113 | 4 | 98 |
| TOCHIGI | 12 | 49 | 13 | 156 |
| GUMMA | 47 | 116 | 30 | 204 |
| SAITAMA | 43 | 164 | 48 | 504 |
| CHIBA | 27 | 104 | 9 | 124 |
| TOKYO | 546 | 1,877 | 49 | 364 |
| KANAGAWA | 385 | 1,735 | 36 | 250 |
| NIIGATA | 15 | 128 | 69 | 432 |
| TOYAMA | 20 | 42 | 13 | 166 |
| ISHIKAWA | 62 | 229 | 7 | 67 |
| FUKUI | 209 | 846 | 33 | 203 |
| YAMANASHI | 9 | 133 | 2 | 30 |
| NAGANO | 1 | 19 | 65 | 439 |
| GIFU | 83 | 391 | 17 | 194 |
| SHIZUOKA | 91 | 404 | 16 | 212 |
| AICHI | 45 | 248 | 47 | 384 |
| MIE | 11 | 28 | 39 | 142 |
| SHIGA | 7 | 34 | 25 | 188 |
| KYOTO | 6 | 17 | 39 | 154 |
| OSAKA | 78 | 263 | 70 | 248 |
| HYOGO | 141 | 564 | 26 | 186 |
| NARA | - | - | 7 | 19 |
| WAKAYAMA | 1 | 22 | 5 | 102 |
| TOTTORI | 4 | 12 | 19 | 69 |
| SHIMANE | 2 | 3 | 25 | *108 |
| OKAYAMA | 72 | 288 | 17 | 188 |
| HIROSHIMA | 68 | 221 | 40 | 177 |
| YAMAGUCHI | 8 | 152 | 3 | 28 |
| TOKUSHIMA | 1 | 8 | 4 | 26 |
| KAGAWA | 19 | 107 | 14 | 113 |
| EHIME | 51 | 329 | 24 | 155 |
| KOCHI | 13 | 124 | 18 | *110 |
| FUKUOKA | 201 | *699 | 69 | *479 |
| SAGA | 81 | 288 | 24 | 173 |
| NAGASAKI | 76 | 262 | 16 | 132 |
| KUMAMOTO | 5 | 15 | 36 | 280 |
| OITA | 1 | 28 | 12 | 58 |
| MIYAZAKI | 8 | 21 | 30 | 343 |
| KAGOSHIMA | 15 | 115 | 23 | 203 |
| TOTAL | 2,911 | *13,150 | 1,245 | *9,448 |
| RATE | | | | |
| Current | 181.1 | 102.3 | 77.5 | 73.5 |
| Previous | 132.0 | | 78.6 | |

See footnotes at end of teble.

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775 013

Weekly Report - 24 February 1951
Continued

| PREFECTURE | TUBERCULOSIS Current Cases | TUBERCULOSIS Cumulative Cases | PNEUMONIA Current Cases | PNEUMONIA Cumulative Cases |
|---|---|---|---|---|
| HOKKAIDO | 514 | 4,733 | 319 | 2,868 |
| AOMORI | 158 | 867 | 66 | 440 |
| IWATE | 89 | 1,072 | 80 | 763 |
| MIYAGI | 123 | 1,225 | 97 | 941 |
| AKITA | 185 | 1,079 | 133 | 575 |
| YAMAGATA | 148 | 982 | 135 | 990 |
| FUKUOKA | 138 | 1,028 | 148 | 1,043 |
| IBARAKI | 43 | 589 | 86 | 1,006 |
| TOCHIGI | 63 | 579 | 134 | 1,041 |
| GUMMA | 133 | 880 | 275 | 1,989 |
| SAITAMA | 180 | 1,578 | 524 | 3,634 |
| CHIBA | 119 | 923 | 152 | 892 |
| TOKYO | 1,183 | 7,036 | 294 | 2,804 |
| KANAGAWA | 288 | 2,359 | 321 | 2,395 |
| NIIGATA | 159 | 1,345 | 183 | 1,463 |
| TOYAMA | 135 | 1,214 | 211 | 2,147 |
| ISHIKAWA | 93 | 577 | 67 | 393 |
| FUKUI | 217 | 764 | 124 | 542 |
| YAMANASHI | 41 | 303 | 61 | 455 |
| NAGANO | 179 | 1,472 | 164 | 1,609 |
| GIFU | 121 | 1,119 | 133 | 941 |
| SHIZUOKA | 223 | 1,998 | 177 | 1,078 |
| AICHI | 383 | *3,296 | 193 | 1,527 |
| MIE | 147 | 990 | 147 | 613 |
| SHIGA | 62 | 612 | 141 | 687 |
| KYOTO | 265 | 1,662 | 132 | 632 |
| OSAKA | 569 | 4,862 | 208 | 1,097 |
| HYOGO | 216 | 2,595 | 124 | 927 |
| NARA | 84 | *361 | 36 | *114 |
| WAKAYAMA | 72 | 698 | 36 | 288 |
| TOTTORI | 60 | 359 | 54 | 201 |
| SHIMANE | 103 | *606 | 21 | *203 |
| OKAYAMA | 143 | *1,318 | 140 | 872 |
| HIROSHIMA | 198 | 1,692 | 96 | 735 |
| YAMAGUCHI | 179 | 962 | 26 | 253 |
| TOKUSHIMA | 61 | 521 | 34 | 231 |
| KAGAWA | 150 | 548 | 87 | 392 |
| EHIME | 116 | 853 | 118 | 908 |
| KOCHI | 63 | 567 | 39 | 317 |
| FUKUOKA | 444 | 3,344 | 193 | *1,546 |
| SAGA | 134 | 941 | 120 | 830 |
| NAGASAKI | 291 | 1,531 | 96 | 613 |
| KUMAMOTO | 157 | 1,099 | 118 | 1,025 |
| OITA | 68 | 683 | 70 | 307 |
| MIYAZAKI | 102 | 936 | 61 | 506 |
| KAGOSHIMA | 179 | 966 | 54 | 470 |
| TOTAL | 8,778 | *65,724 | 6,228 | *45,303 |
| RATE | | | | |
| Current | 546.2 | 511.2 | 387.5 | 352.4 |
| Previous | 524.3 | | 381.1 | |

See footnotes at end of table.

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775 013

Weekly Report - 24 February 1951
Continued

| PREFECTURE | INFLUENZA Current Cases | INFLUENZA Cumulative Cases | POLIOMYELITIS Current Cases | POLIOMYELITIS Cumulative Cases |
|---|---|---|---|---|
| HOKKAIDO | 7 | 625 | 2 | 13 |
| AOMORI | - | - | - | - |
| IWATE | - | - | - | - |
| MIYAGI | - | 10 | - | 13 |
| AKITA | - | - | 1 | 2 |
| YAMAGATA | - | 17 | - | 7 |
| FUKUSHIMA | - | 1 | - | 3 |
| IBARAKI | - | 105 | 1 | 3 |
| TOCHIGI | - | - | - | - |
| GUMMA | 3 | 103 | 3 | 6 |
| SAITAMA | 8 | 136 | 1 | 21 |
| CHIBA | 1 | 126 | - | 3 |
| TOKYO | 1 | 193 | 6 | 22 |
| KANAGAWA | - | - | - | 3 |
| NIIGATA | 22 | 875 | 2 | 9 |
| TOYAMA | 1 | 345 | - | 1 |
| ISHIKAWA | 1 | 8 | - | - |
| FUKUI | 3 | 61 | 3 | 6 |
| YAMANASHI | - | 4 | - | 3 |
| NAGANO | - | *- | - | 2 |
| GIFU | 3 | 1,136 | 2 | 5 |
| SHIZUOKA | 5 | 348 | - | 6 |
| AICHI | 6 | 198 | - | 2 |
| MIE | - | 116 | - | 5 |
| SHIGA | - | 8 | - | 2 |
| KYOTO | 1 | 9 | 2 | 4 |
| OSAKA | - | 10 | 7 | 50 |
| HYOGO | 1 | 78 | - | 11 |
| NARA | 2 | *4 | 2 | *8 |
| WAKAYAMA | - | 43 | - | 1 |
| TOTTORI | - | 1 | - | - |
| SHIMANE | - | 24 | 1 | 1 |
| OKAYAMA | - | 37 | 1 | 5 |
| HIROSHIMA | - | - | - | 1 |
| YAMAGUCHI | 3 | 230 | 1 | 1 |
| TOKUSHIMA | - | - | - | 5 |
| KAGAWA | 2 | 27 | - | - |
| EHIME | 1 | 384 | - | 7 |
| KOCHI | - | 1 | - | - |
| FUKUOKA | - | 36 | - | 7 |
| SAGA | - | - | - | 1 |
| NAGASAKI | 11 | 84 | - | 1 |
| KUMAMOTO | 1 | 14 | 1 | 6 |
| OITA | 9 | 13 | - | - |
| MIYAZAKI | - | 12 | 1 | 3 |
| KAGOSHIMA | - | - | - | 3 |
| TOTAL | 92 | *5,422 | 37 | *252 |
| RATE | | | | |
| Current | 5.7 | 42.2 | 2.3 | 2.0 |
| Previous | 16.5 | | 1.9 | |

See footnotes at end of table.

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775 013



| PREFECTURE | TETANUS | | PUERPERAL INFECTION | |
|---|---|---|---|---|
| | Current Cases | Cumulative Cases | Current Cases | Cumulative Cases |
| HOKKAIDO | - | 1 | - | 13 |
| AOMORI | - | 1 | - | 3 |
| IWATE | - | 2 | - | - |
| MIYAGI | - | 2 | - | - |
| AKITA | - | 4 | - | - |
| YAMAGATA | - | 4 | 1 | 3 |
| FUKUSHIMA | - | - | - | 2 |
| IBARAKI | 2 | 8 | - | 2 |
| TOCHIGI | 2 | 5 | 1 | 3 |
| GUMMA | 1 | 7 | - | 4 |
| SAITAMA | 1 | 10 | - | 8 |
| CHIBA | - | 12 | - | 3 |
| TOKYO | 2 | 11 | - | 4 |
| KANAGAWA | - | 2 | - | - |
| NIIGATA | - | 4 | 2 | 5 |
| TOYAMA | - | - | 1 | 3 |
| ISHIKAWA | - | 1 | - | 1 |
| FUKUI | - | - | - | 1 |
| YAMANASHI | - | 4 | - | 4 |
| NAGANO | - | 4 | - | 7 |
| GIFU | - | 3 | - | 5 |
| SHIZUOKA | 1 | 15 | - | 2 |
| AICHI | 2 | 10 | - | 3 |
| MIE | - | 4 | - | - |
| SHIGA | - | - | 2 | 8 |
| KYOTO | - | 5 | - | - |
| OSAKA | - | 5 | - | 3 |
| HYOGO | 2 | 6 | - | - |
| NARA | - | *2 | - | - |
| WAKAYAMA | - | - | - | 1 |
| TOTTORI | - | - | - | - |
| SHIMANE | - | 2 | - | *3 |
| OKAYAMA | - | 3 | - | 3 |
| HIROSHIMA | - | 6 | 1 | 11 |
| YAMAGUCHI | 2 | 2 | 1 | 1 |
| TOKUSHIMA | 2 | 4 | - | - |
| KAGAWA | - | 2 | - | 4 |
| EHIME | 2 | 10 | - | - |
| KOCHI | 1 | 3 | - | 3 |
| FUKUOKA | 3 | 10 | - | - |
| SAGA | 1 | 4 | - | 1 |
| NAGASAKI | - | - | - | - |
| KUMAMOTO | - | 8 | - | 1 |
| OITA | - | 3 | - | 1 |
| MIYAZAKI | 1 | 8 | - | 1 |
| KAGOSHIMA | 1 | 8 | - | 2 |
| TOTAL | 26 | *205 | 9 | *119 |
| RATE | | | | |
| Current | 1.6 | 1.6 | 0.6 | 0.9 |
| Previous | 1.2 | | 1.0 | |

See footnotes at end of table.

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775 013



| PREFECTURE | RABIES Current Cases | RABIES Cumulative Cases | ANTHRAX Current Cases | ANTHRAX Cumulative Cases |
|---|---|---|---|---|
| HOKKAIDO | - | - | - | - |
| AOMORI | - | - | - | - |
| IWATE | - | - | - | - |
| MIYAGI | - | - | - | - |
| AKITA | - | - | - | - |
| YAMAGATA | - | - | - | - |
| FUKUSHIMA | - | - | - | - |
| IBARAKI | - | 2 | - | - |
| TOCHIGI | - | - | - | - |
| GUMMA | - | 1 | - | - |
| SAITAMA | - | 1 | - | - |
| CHIBA | - | - | - | - |
| TOKYO | - | 1 | - | - |
| KANAGAWA | - | 1 | - | - |
| NIIGATA | - | - | - | - |
| TOYAMA | - | - | - | - |
| ISHIKAWA | - | - | - | - |
| FUKUI | - | - | - | - |
| YAMANASHI | - | - | - | - |
| NAGANO | - | - | - | - |
| GIFU | - | - | - | - |
| SHIZUOKA | - | 1 | - | - |
| AICHI | - | - | - | 1 |
| MIE | - | - | - | - |
| SHIGA | - | - | - | - |
| KYOTO | - | - | - | - |
| OSAKA | - | - | - | - |
| HYOGO | - | - | - | - |
| NARA | - | - | - | - |
| WAKAYAMA | - | - | - | - |
| TOTTORI | - | - | - | - |
| SHIMANE | - | - | - | - |
| OKAYAMA | - | - | - | - |
| HIROSHIMA | - | - | - | - |
| YAMAGUCHI | - | - | - | - |
| TOKUSHIMA | - | - | - | - |
| KAGAWA | - | - | - | - |
| EHIME | - | - | - | - |
| KOCHI | - | - | - | - |
| FUKUOKA | - | - | - | - |
| SAGA | - | - | - | - |
| NAGANO | - | - | - | - |
| KUMAMOTO | - | - | - | - |
| OITA | - | - | - | - |
| MIYAZAKI | - | - | - | - |
| KAGOSHIMA | - | - | - | - |
| TOTAL | - | 7 | - | 1 |
| RATE | | | | |
| Current | - | 0.1 | - | 0.0 |
| Previous | - | | - | |

See footnotes at end of table.

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775 013



| PREFECTURE | LEPROSY Current Cases | LEPROSY Cumulative Cases | TRACHOMA Current Cases | TRACHOMA Cumulative Cases |
|---|---|---|---|---|
| HOKKAIDO | - | - | 148 | 928 |
| AOMORI | - | - | 83 | 505 |
| IWATE | - | - | 62 | 653 |
| MIYAGI | - | 3 | 69 | 507 |
| AKITA | - | 3 | 27 | 625 |
| YAMAGATA | - | - | 42 | 202 |
| FUKUSHIMA | - | 7 | 30 | 326 |
| IBARAKI | - | 2 | 11 | 269 |
| TOCHIGI | - | 2 | 8 | 117 |
| GUMMA | 1 | 4 | 38 | 269 |
| SAITAMA | - | 1 | 58 | 380 |
| CHIBA | - | - | 20 | 281 |
| TOKYO | - | 3 | 88 | 404 |
| KANAGAWA | - | 1 | 44 | 285 |
| NIIGATA | - | - | 37 | 238 |
| TOYAMA | - | - | 15 | 186 |
| ISHIKAWA | - | 1 | 5 | 55 |
| FUKUI | - | - | 30 | 150 |
| YAMANASHI | - | 2 | 21 | 209 |
| NAGANO | - | 1 | 73 | 282 |
| GIFU | - | - | 17 | 170 |
| SHIZUOKA | - | 4 | 37 | 202 |
| AICHI | - | *5 | 78 | 790 |
| MIE | 1 | 3 | 15 | 107 |
| SHIGA | - | - | 12 | 73 |
| KYOTO | 3 | 7 | 40 | 288 |
| OSAKA | - | 1 | 161 | 891 |
| HYOGO | - | 4 | 198 | 984 |
| NARA | - | - | 13 | 66 |
| WAKAYAMA | - | 1 | 34 | 282 |
| TOTTORI | - | - | 4 | 419 |
| SHIMANE | - | - | 7 | *72 |
| OKAYAMA | - | - | 26 | 421 |
| HIROSHIMA | - | 1 | 42 | 357 |
| YAMAGUCHI | 1 | 1 | 4 | 41 |
| TOKUSHIMA | - | 1 | 7 | 70 |
| KAGAWA | - | 3 | 18 | 200 |
| EHIME | - | 1 | 35 | 212 |
| KOCHI | - | - | 26 | 57 |
| FUKUOKA | - | 2 | 218 | 873 |
| SAGA | - | - | 25 | 197 |
| NAGASAKI | - | 1 | 30 | 148 |
| KUMAMOTO | - | - | 28 | 172 |
| OITA | 1 | 1 | 10 | 266 |
| MIYAZAKI | - | - | 16 | 195 |
| KAGOSHIMA | - | 1 | 7 | 70 |
| TOTAL | 7 | *67 | 1,927 | *14,499 |
| RATE | | | | |
| Current | 0.4 | 0.5 | 119.9 | 112.8 |
| Previous | 0.4 | | 133.7 | |

See footnotes at end of table.

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775 013



| PREFECTURE | INFECTIOUS DIARRHEA Current Cases | INFECTIOUS DIARRHEA Cumulative Cases | SCHISTOSOMIASIS Current Cases | SCHISTOSOMIASIS Cumulative Cases |
|---|---|---|---|---|
| HOKKAIDO | - | 5 | - | - |
| AOMORI | - | - | - | - |
| IWATE | - | - | - | - |
| MIYAGI | - | - | - | - |
| AKITA | - | - | - | - |
| YAMAGATA | - | - | - | - |
| FUKUSHIMA | - | - | - | - |
| IBARAKI | - | - | - | - |
| TOCHIGI | - | - | - | - |
| GUMMA | - | - | - | - |
| SAITAMA | - | - | - | - |
| CHIBA | - | - | - | - |
| TOKYO | - | - | - | - |
| KANAGAWA | - | - | - | - |
| NIIGATA | - | - | - | - |
| TOYAMA | - | - | - | - |
| ISHIKAWA | - | - | - | - |
| FUKUI | - | - | - | - |
| YAMANASHI | - | - | 3 | 40 |
| NAGANO | - | - | - | - |
| GIFU | - | - | - | - |
| SHIZUOKA | - | - | - | - |
| AICHI | - | - | - | - |
| MIE | - | - | - | - |
| SHIGA | - | - | - | - |
| KYOTO | - | - | - | - |
| OSAKA | - | - | - | - |
| HYOGO | - | - | - | - |
| NARA | - | - | - | - |
| WAKAYAMA | - | - | - | - |
| TOTTORI | - | - | - | - |
| SHIMANE | - | - | - | - |
| OKAYAMA | - | - | - | - |
| HIROSHIMA | - | - | - | 2 |
| YAMAGUCHI | - | - | - | - |
| TOKUSHIMA | - | - | - | - |
| KAGAWA | - | - | - | - |
| EHIME | - | - | - | - |
| KOCHI | - | - | - | - |
| FUKUOKA | - | - | 3 | 6 |
| SAGA | - | - | - | 2 |
| NAGASAKI | - | - | - | - |
| KUMAMOTO | - | - | - | - |
| OITA | - | - | - | - |
| MIYAZAKI | - | - | - | - |
| KAGOSHIMA | - | - | - | - |
| TOTAL | - | 5 | 6 | 50 |
| RATE | | | | |
| Current | - | 0.0 | 0.4 | 0.4 |
| Previous | - | | 0.5 | |

See footnotes at end of table.

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775 013



| PREFECTURE | FILARIASIS Current Cases | FILARIASIS Cumulative Cases | SYPHILIS Current Cases | SYPHILIS Cumulative Cases |
|---|---|---|---|---|
| HOKKAIDO | - | - | 75 | 613 |
| AOMORI | - | - | 14 | 180 |
| IWATE | - | - | 4 | 93 |
| MIYAGI | - | - | 25 | 144 |
| AKITA | - | - | 12 | 136 |
| YAMAGATA | - | - | 38 | 219 |
| FUKUSHIMA | - | - | 37 | 349 |
| IBARAKI | - | - | 18 | 142 |
| TOCHIGI | - | - | 17 | 143 |
| GUMMA | - | - | 27 | 153 |
| SAITAMA | - | - | 25 | 155 |
| CHIBA | - | 1 | 10 | 176 |
| TOKYO | - | - | 71 | 683 |
| KANAGAWA | - | - | 126 | 898 |
| NIIGATA | - | - | 47 | 376 |
| TOYAMA | - | - | 46 | 209 |
| ISHIKAWA | - | - | 28 | 146 |
| FUKUI | - | 1 | 22 | 108 |
| YAMANASHI | - | 1 | 8 | 60 |
| NAGANO | - | - | 21 | 176 |
| GIFU | - | - | 11 | 233 |
| SHIZUOKA | - | - | 98 | 403 |
| AICHI | - | - | 99 | 474 |
| MIE | - | - | 21 | 176 |
| SHIGA | - | - | 13 | 80 |
| KYOTO | - | - | 46 | 281 |
| OSAKA | - | - | 124 | 868 |
| HYOGO | - | - | 70 | 479 |
| NARA | - | - | 17 | 50 |
| WAKAYAMA | - | - | 19 | 136 |
| TOTTORI | - | - | 24 | 146 |
| SHIMANE | - | - | 15 | 56 |
| OKAYAMA | - | - | 27 | 250 |
| HIROSHIMA | - | - | 72 | *366 |
| YAMAGUCHI | - | - | 47 | 345 |
| TOKUSHIMA | - | - | 3 | 73 |
| KAGAWA | - | - | 1 | 78 |
| EHIME | - | - | 14 | 158 |
| KOCHI | - | - | 16 | 95 |
| FUKUOKA | - | 2 | 219 | 1,199 |
| SAGA | 1 | 2 | 25 | 297 |
| NAGASAKI | 1 | 1 | 134 | *885 |
| KUMAMOTO | - | 3 | 35 | *208 |
| OITA | - | - | 18 | 178 |
| MIYAZAKI | - | - | 13 | 160 |
| KAGOSHIMA | - | 5 | 22 | 168 |
| TOTAL | 2 | 16 | 1,874 | *12,939 |
| RATE | | | | |
| Current | 0.1 | 0.1 | 116.6 | 100.6 |
| Previous | 0.2 | | 99.1 | |

See footnotes at end of table.

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775 013



| PREFECTURE | GONORRHEA Current Cases | GONORRHEA Cumulative Cases | CHANCROID Current Cases | CHANCROID Cumulative Cases |
|---|---|---|---|---|
| HOKKAIDO | 161 | 1,916 | 6 | 73 |
| AOMORI | 17 | 196 | 3 | 14 |
| IWATE | 14 | 94 | - | 3 |
| MIYAGI | 36 | 220 | - | 1 |
| AKITA | 12 | 83 | 1 | 4 |
| YAMAGATA | 20 | 143 | - | 2 |
| FUKUSHIMA | 35 | 379 | 3 | 18 |
| IBARAKI | 16 | 116 | 1 | 18 |
| TOCHIGI | 32 | 236 | 3 | 17 |
| GUMMA | 19 | 226 | 2 | 18 |
| SAITAMA | 39 | 285 | 2 | 19 |
| CHIBA | 29 | 194 | - | 20 |
| TOKYO | 211 | 1,896 | 12 | 193 |
| KANAGAWA | 503 | 3,170 | 52 | 388 |
| NIIGATA | 66 | 455 | 2 | 27 |
| TOYAMA | 41 | 379 | 2 | 21 |
| ISHIKAWA | 48 | 256 | 3 | 15 |
| FUKUI | 66 | 280 | 2 | 21 |
| YAMANASHI | 3 | 39 | 1 | 3 |
| NAGANO | 40 | 281 | 2 | 5 |
| GIFU | 18 | 371 | 2 | 37 |
| SHIZUOKA | 72 | 508 | 5 | 35 |
| AICHI | 120 | 844 | 5 | 60 |
| MIE | 46 | 261 | 4 | 23 |
| SHIGA | 21 | 177 | 2 | 45 |
| KYOTO | 113 | 607 | 17 | 135 |
| OSAKA | 127 | 813 | 19 | 162 |
| HYOGO | 135 | 865 | 14 | 87 |
| NARA | 47 | 211 | 8 | 43 |
| WAKAYAMA | 35 | 428 | 4 | 35 |
| TOTTORI | 36 | 215 | 1 | 18 |
| SHIMANE | 10 | 49 | 1 | 4 |
| OKAYAMA | 56 | 370 | 12 | 39 |
| HIROSHIMA | 255 | *1,334 | 17 | 117 |
| YAMAGUCHI | 113 | 869 | 4 | 22 |
| TOKUSHIMA | 21 | *101 | - | *5 |
| KAGAWA | 4 | 167 | - | 22 |
| EHIME | 24 | 159 | 3 | 15 |
| KOCHI | 53 | 206 | 5 | 22 |
| FUKUOKA | 745 | 4,268 | 53 | 278 |
| SAGA | 62 | 425 | 3 | 25 |
| NAGASAKI | 134 | *981 | 17 | 103 |
| KUMAMOTO | 93 | *523 | 2 | *18 |
| OITA | 30 | 289 | 2 | 16 |
| MIYAZAKI | 43 | 247 | - | 3 |
| KAGOSHIMA | 48 | 274 | 4 | 8 |
| TOTAL | 3,869 | *26,406 | 301 | *2,257 |
| RATE | | | | |
| Current | 240.7 | 205.4 | 18.7 | 17.6 |
| Previous | 202.0 | | 15.6 | |

See footnotes at end of table.

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775 013

NUMBERS OF CASES AND DEATHS OF COMMUNICABLE DISEASES DURING COMPARABLE PERIODS IN 1949, 1950 AND 1951

| Diseases | Week Ended 24 Feb 1951 | Week Ended 25 Feb 1950 | Week Ended 19 Feb 1949 | Cumulative Number for First 8 Weeks 1951 | Cumulative Number for First 8 Weeks 1950 | Cumulative Number for First 8 Weeks 1949 |
|---|---|---|---|---|---|---|
| **Cases** | | | | | | |
| Diphtheria | 314 | 278 | 398 | 2,272 | 2,465 | 3,022 |
| Dysentery | 271 | 105 | 32 | 1,608 | 616 | 256 |
| Typhoid Fever | 34 | 51 | 83 | 279 | 495 | 802 |
| Paratyphoid Fever | 9 | 10 | 23 | 84 | 140 | 314 |
| Smallpox | 2 | - | - | 9 | 1 | 4 |
| Typhus Fever | 1 | 113 | 6 | 3 | 495 | 46 |
| Malaria | 8 | 11 | 17 | 54 | 87 | 183 |
| Japanese "B" Encephalitis | - | - | - | 1 | - | 1 |
| Scarlet Fever | 63 | 76 | 89 | 613 | 691 | 759 |
| Epidemic Meningitis | 29 | 20 | 30 | 181 | 175 | 189 |
| Cholera | - | - | - | - | - | - |
| Plague | - | - | - | - | - | - |
| Measles | 2,911 | 1,088 | 2,742 | 13,150 | 7,366 | 12,469 |
| Whooping Cough | 1,245 | 2,679 | 1,498 | 9,448 | 21,569 | 10,136 |
| Tuberculosis | 8,778 | 7,984 | 8,276 | 65,724 | 56,723 | 54,286 |
| Pneumonia | 6,228 | 4,921 | 4,065 | 45,303 | 39,082 | 25,922 |
| Influenza | 92 | 2,096 | 55 | 5,422 | 14,318 | 308 |
| Poliomyelitis | 37 | 38 | 28 | 252 | 247 | 203 |
| Yellow Fever | - | - | - | - | - | - |
| Tetanus | 26 | 23 | 41 | 205 | 203 | 268 |
| Puerperal Infection | 9 | 16 | 13 | 119 | 143 | 167 |
| Rabies | - | 1 | 1 | 7 | 9 | 5 |
| Anthrax | - | - | - | 1 | - | 1 |
| Glanders | - | - | - | - | - | - |
| Leprosy | 7 | 7 | 10 | 67 | 61 | 81 |
| Trachoma | 1,927 | 2,488 | 2,213 | 14,499 | 16,168 | 15,079 |
| Infectious Diarrhea | - | - | 8 | 5 | 10 | 24 |
| Dengue Fever | - | - | - | - | 1 | 1 |
| Tustusgamushi Diseases | - | - | NA | - | - | NA |
| Schistosomiasis | 6 | 5 | NA | 50 | 37 | NA |
| Filariasis | 2 | 5 | NA | 16 | 21 | NA |
| **Deaths** | | | | | | |
| Diphtheria | 30 | 31 | 44 | 277 | 262 | 353 |
| Dysentery | 37 | 25 | 10 | 223 | 160 | 80 |
| Typhoid Fever | 3 | 6 | 8 | 26 | 82 | 78 |
| Paratyphoid Fever | - | 1 | 1 | 3 | 5 | 7 |
| Smallpox | 1 | - | - | 1 | - | - |
| Typhus Fever | - | 4 | - | - | 29 | - |
| Malaria | - | 1 | - | 8 | 10 | 5 |
| Japanese "B" Encephalitis | 1 | - | - | 2 | - | - |
| Scarlet Fever | 1 | - | 4 | 4 | 4 | 16 |
| Epidemic Meningitis | 2 | 9 | 10 | 38 | 42 | 47 |
| Cholera | - | - | - | - | - | - |
| Plague | - | - | - | - | - | - |

See footnotes at end of table.

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775 013

CASE AND DEATH RATES FOR COMMUNICABLE DISEASES
DURING COMPARABLE PERIODS IN 1949, 1950 AND 1951

| Diseases | Week Ended | | | Cumulative Rates for First 8 Weeks | | |
|---|---|---|---|---|---|---|
| | 24 Feb 1951 | 25 Feb 1950 | 19 Feb 1949 | 1951 | 1950 | 1949 |
| **Case Rates** | | | | | | |
| Diphtheria | 19.5 | 17.3 | 25.7 | 17.7 | 19.2 | 24.4 |
| Dysentery | 16.9 | 6.5 | 2.1 | 12.5 | 4.8 | 2.1 |
| Typhoid Fever | 2.1 | 3.2 | 5.4 | 2.2 | 3.9 | 6.5 |
| Paratyphoid Fever | 0.6 | 0.6 | 1.5 | 0.7 | 1.1 | 2.5 |
| Smallpox | 0.1 | - | - | 0.1 | 0.0 | 0.0 |
| Typhus Fever | 0.1 | 7.0 | 0.4 | 0.0 | 3.9 | 0.4 |
| Malaria | 0.5 | 0.7 | 1.1 | 0.4 | 0.7 | 1.5 |
| Japanese "B" Encephalitis | - | - | - | 0.0 | - | 0.0 |
| Scarlet Fever | 3.9 | 4.7 | 5.7 | 4.8 | 5.4 | 6.1 |
| Epidemic meningitis | 1.8 | 1.2 | 1.9 | 1.4 | 1.4 | 1.5 |
| Cholera | - | - | - | - | - | - |
| Plague | - | - | - | - | - | - |
| Measles | 181.1 | 67.7 | 176.8 | 102.3 | 57.3 | 100.5 |
| Whooping cough | 77.5 | 166.7 | 96.6 | 73.5 | 167.8 | 81.7 |
| Tuberculosis | 546.2 | 496.8 | 533.6 | 511.2 | 441.2 | 437.5 |
| Pneumonia | 387.5 | 306.2 | 262.1 | 352.4 | 304.0 | 208.9 |
| Influenza | 5.7 | 130.4 | 3.5 | 42.2 | 111.4 | 2.5 |
| Poliomyelitis | 2.3 | 2.4 | 1.8 | 2.0 | 1.9 | 1.6 |
| Yellow Fever | - | - | - | - | - | - |
| Tetanus | 1.6 | 1.4 | 2.6 | 1.6 | 1.6 | 2.2 |
| Puerperal Infection | 0.6 | 1.0 | 0.8 | 0.9 | 1.1 | 1.3 |
| Rabies | - | 0.1 | 0.1 | 0.1 | 0.1 | 0.0 |
| Anthrax | - | - | - | 0.0 | - | 0.0 |
| Glanders | - | - | - | - | - | - |
| Leprosy | 0.4 | 0.4 | 0.6 | 0.5 | 0.5 | 0.7 |
| Trachoma | 119.9 | 154.8 | 142.7 | 112.8 | 125.8 | 121.5 |
| Infectious diarrhea | - | - | 0.5 | 0.0 | 0.1 | 0.2 |
| Dengue Fever | - | - | - | - | 0.0 | 0.0 |
| Tustsugamushi Diseases | - | - | NA | - | - | NA |
| Schistosomiasis | 0.4 | 0.3 | NA | 0.4 | 0.3 | NA |
| Filariasis | 0.1 | 0.3 | NA | 0.1 | 0.2 | NA |
| **Death Rates** | | | | | | |
| Diphtheria | 1.9 | 1.9 | 2.8 | 2.2 | 2.0 | 2.8 |
| Dysentery | 2.3 | 1.6 | 0.6 | 1.7 | 1.2 | 0.6 |
| Typhoid Fever | 0.2 | 0.4 | 0.5 | 0.2 | 0.6 | 0.6 |
| Paratyphoid Fever | - | 0.1 | 0.1 | 0.0 | 0.0 | 0.1 |
| Smallpox | 0.1 | - | - | 0.0 | - | - |
| Typhus Fever | - | 0.2 | - | - | 0.2 | - |
| Malaria | - | 0.1 | - | 0.1 | 0.1 | 0.0 |
| Japanese "B" Encephalitis | 0.1 | - | - | 0.0 | - | - |
| Scarlet Fever | 0.1 | - | 0.3 | 0.0 | 0.0 | 0.1 |
| Epidemic Meningitis | 0.1 | 0.6 | 0.6 | 0.3 | 0.3 | 0.4 |
| Cholera | - | - | - | - | - | - |
| Plague | - | - | - | - | - | - |

See footnotes at end of table.

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775 013

Weekly Report - 24 February 1951
Continued

| PREFECTURE | LYMPHOGRANULOMA VENEREUM Current Cases | Cumulative Cases |
|---|---|---|
| HOKKAIDO | - | 1 |
| AOMORI | - | - |
| IWATE | - | - |
| MIYAGI | - | - |
| AKITA | - | - |
| YAMAGATA | - | - |
| FUKUSHIMA | 1 | 1 |
| IBARAKI | - | - |
| TOCHIGI | - | - |
| GUMMA | - | - |
| SAITAMA | - | - |
| CHIBA | - | 1 |
| TOKYO | 2 | 5 |
| KANAGAWA | 2 | 5 |
| NIIGATA | - | - |
| TOYAMA | - | 2 |
| ISHIKAWA | - | 2 |
| FUKUI | - | 2 |
| YAMANASHI | - | - |
| NAGANO | - | - |
| GIFU | - | - |
| SHIZUOKA | - | 1 |
| AICHI | - | - |
| MIE | - | 1 |
| SHIGA | - | - |
| KYOTO | 3 | 7 |
| OSAKA | 1 | 4 |
| HYOGO | - | 2 |
| NARA | - | 2 |
| WAKAYAMA | - | 1 |
| TOTTORI | - | - |
| SHIMANE | - | - |
| OKAYAMA | - | 1 |
| HIROSHIMA | - | - |
| YAMAGUCHI | 2 | 2 |
| TOKUSHIMA | - | - |
| KAGAWA | - | 1 |
| EHIME | 1 | 1 |
| KOCHI | - | - |
| FUKUOKA | - | 1 |
| SAGA | - | 1 |
| NAGASAKI | - | 1 |
| KUMAMOTO | - | 1 |
| OITA | - | - |
| MIYAZAKI | - | - |
| KAGOSHIMA | - | - |
| TOTAL | 12 | 46 |
| RATE | | |
| Current | 0.7 | 0.4 |
| Previous | 0.4 | |

See footnotes at end of table.

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775 013

NUMBERS OF CASES AND CASE RATES FOR
VENEREAL DISEASES IN JAPAN DURING
COMPARABLE PERIODS IN 1949, 1950 AND 1951

| DISEASES | WEEK ENDED 24 Feb 1951 | WEEK ENDED 25 Feb 1950 | WEEK ENDED 19 Feb 1949 | CUMULATIVE NUMBERS FOR FIRST 8 WEEKS 1951 | 1950 | 1949 |
|---|---|---|---|---|---|---|
| NUMBERS | | | | | | |
| SYPHILIS | 1,874 | 2,568 | 3,890 | 12,939 | 19,096 | 28,054 |
| GONORRHEA | 3,869 | 3,024 | 3,397 | 26,406 | 24,118 | 25,671 |
| CHANCROID | 301 | 305 | 549 | 2,257 | 2,550 | 4,222 |
| LYMPHOGRANULOMA VENEREUM | 12 | 8 | 11 | 46 | 89 | 106 |
| RATES | | | | | | |
| SYPHILIS | 116.6 | 159.8 | 250.8 | 100.6 | 148.5 | 226.1 |
| GONORRHEA | 240.7 | 188.2 | 219.0 | 205.4 | 187.6 | 206.9 |
| CHANCROID | 18.7 | 19.0 | 35.4 | 17.6 | 19.8 | 34.0 |
| LYMPHOGRANULOMA VENEREUM | 0.7 | 0.5 | 0.7 | 0.4 | 0.7 | 0.9 |

FOOTNOTES:

1. There were no cases or deaths reported for cholra, plague, yellow fever, glanders, dengue fever or tsutsugamushi disease.

2. Rates are the numbers of cases or deaths per 100,000 population, estimated as of 1 July 1950, and are computed on an annual basis.

3. A dash (-) indicates that no cases or deaths were reported and that the case or death rate was zero.

4. A rate of 0.0 indicates that there were some cases or deaths but that the rate was less than 0.1.

5. "NA" indicates that data are not available.

6. * Cumulative figures adjusted for delayed and corrected reports.

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775 013

# DIGEST OF MONTHLY REPORT OF COMMUNICABLE DISEASES IN JAPAN FOR THE FOUR WEEK PERIOD ENDED 24 FEBRUARY 1951

During the four weeks ended 24 February 1951 the 12 1/ communicable diseases included in this digest for which reports on deaths as well as cases are available accounted for 2,701 cases and 284 deaths. Cases were higher and deaths lower 2/ than in January when they numbered 2,397 and 294 respectively. The 19 additional diseases included herein (not counting the four venereal diseases) accounted for 87,030 cases compared with 67,309 last month. Rates for seven diseases (dysentery, smallpox, epidemic meningitis, measles, tuberculosis, pneumonia, and leprosy) were higher this month than in both last month and February 1950, while rates for three (scarlet fever, rabies, and infectious diarrhea) were lower. There were no cases of Cholera, plague, yellow fever, glanders, or tsutsugamushi during any of the three periods. February rates for the remaining 16 diseases fell between those reported last month and in February 1950.

The diphtheria case rate was 10 percent higher this month (18.5) than last month (16.8), but the death rate decreased from 2.- to 1.9. This month's case rate was the lowest 3/ ever recorded for February. It was 7 percent less than that recorded for February 1950 (19.8) and 28 percent below the comparable rate (25.7) in 1949. Rates increased over January in 27 prefectures and decreased in 15, while the 4 remaining prefectures reported the same rates both months. As in January, the highest rates were reported from the island of Kyushu. Rates in Nagasaki (48.0), Miyazaki (40.3), and Oita (37.2) were more than twice the national average. Rates in the remaining 43 prefectures ranged from 3.1 to 28.6.

The case rate for dysentery (13.5) was 17 percent higher than in the preceding month (11.5), and the death rate increased from 1.6 to 1.9. The current case rate was the highest February rate since 1942. It was two and a half times as great as the February 1950 rate (5.4) and over six times the corresponding 1949 rate (2.2). About half (24) of the prefectures reported higher rates than in January, 17 reported lower, and 3 the same rates both months. The two remaining prefectures reported no cases either month. The most notable change was an increase in rates in Niigata Prefecture from 23.7 to 67.3, the current rate being about five times the national average. The current rate in Tokyo-to (49.3) was nearly four times the national average. In the 39 additional prefectures reporting current cases, rates ranged from 0.8 to 31.4. Of the total dysentery this month, 826 cases and 117 deaths were listed as bacillary, the remaining 39 cases and 3 deaths as amebic.

The typhoid fever case rate increased from 1.8 in January to 2.5 in February, while the death rate was the same (0.2) both months. The current rate was the lowest 3/ on record for February. It was more than 25 percent below the rate (3.4) in the same month last year, and nearly 60 percent below the rate (6.0) in February 1949. There were 21 prefectures reporting increases over last month, 17 decreases, and 3 the same rate both months. The five remaining prefectures have reported no cases for two months or longer. The current rate in Hiroshima Prefecture (13.1) was over five times, and that in Kyoto Prefecture (9.9) about four times the national average. Among the 35 other prefectures reporting current cases, rates ranged from 0.3 to 8.8.

The case rate for paratyphoid fever increased from 0.6 to 0.7, and the death rate from zero to less than 0.1. The case rate was the lowest 3/ on record for February. Rates in February 1950 and 1949 were 0.9 and 1.9 respectively. There have been no cases reported in two or more months by 17 prefectures. Increases were reported by 14, decreases by 11, and no change by 4 prefectures. Among the 22 prefectures reporting cases in February rates ranged from 0.3 to 3.5.

Smallpox had a case rate of 0.1 and a death rate of less than 0.1, compared with no cases or deaths in January. There were no cases during February 1950, but a rate of less than 0.1 during the comparable period of 1949. All current cases were reported by Hyogo Prefecture, where the case rate was 3.5.

The case rate for typhus fever was less than 0.1 in both January and February 1941, and there were no deaths either month. The current case rate was the

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775 013

lowest recorded for February since 1943. Comparable rates during 1950 and 1949 were 7.4 and 0.4 respectively. All current cases were reported by Hyogo Prefecture, where the case rate was 1.0.

The malaria case rate was the same (0.4) in February as in January, while the death rate decreased from 0.1 to less than 0.1. Case rates in February 1950 and 1949 were 0.8 and 1.4 respectively. There have been no cases for two or more months in 19 prefectures, while 13 reported higher and 10 lower rates in February than in January. The four remaining prefectures reported the same rates both months. Among the 19 prefectures reporting current cases rates ranged from 0.4 to 2.4.

There were no cases of Japanese "B" encephalitis reported in February. The January case rate was less than 0.1. The current death rate was less than 0.1 compared with no deaths in January. No cases were reported during the comparable periods of 1950 or 1949.

The case rate for scarlet fever was approximately the same in February (4.7) as in January (4.8). The death rate was less than 0.1 both months. The currnet case rate was slightly lower than comparable rates in 1950 (4.9) and 1949 (5.1). Nineteen prefectures reported lower rates than in January, 18 higher, and 5 no change. The four remaining prefectures have reported no cases for two or more months. The highest prefectural rates, from three to four times the national average, were reported by three neighboring prefectures, Kyoto (18.4), Shiga (18.0), and Osaka (14.1). At the other extreme, no cases were reported in eight prefectures.

The case rate for epidemic meningitis increased from 1.2 in January to 1.5 in February, and the death rate from 0.2 to 0.3. Case rates in February 1950 and 1949 were 1.4 and 1.8 respectively. Rates were higher than in January for 17 prefectures and lower for 11. Among the remaining 18 there were 9 who have reported no cases for two or more months, and an equal number with the same rate in February as in January. Current rates ranged from zero in 14 prefecturres to a high of 5.7.

The current measles case rate (136.1) was about twice as high as the January rate (68.5). It was a little more than twice the rate (63.8) in February 1950, but 8 percent below the corresponding rate (148.4) in 1949. Rates increased in 39 prefectures, decreased in 5 and remained the same in 1. The one remaining prefecture (Nara) has reported no cases since last October. The most outstanding changes were increases in Fukui (from 538.4 to 916.9) and Kanagawa (286.6 to 615.9). Current rates in these two prefectures were approximately seven and five times respectively as high as the national. Rates in the 43 other prefectures ranged from 4.3 to 352.5.

The case rate for whooping cough increased 31 percent from January (63.7) to February (83.2). The present rate was 55 percent below that (183.4) recorded for February 1950, and 16 percent below the rate (99.2) for the same month of 1949. Rates increased over last months in tow-thirds (30) of the prefectures and decreased in 15. The one remaining prefecture reported the same rate both months. The most notable changes were all increases, and occurred in Saitama (from 104.3 to 199.6), Tottori (36.7 to 112.1), and Shimane (36.9 to 112.0). The highest rate, as in January, was reported by Miyazaki Prefecture (207.5), two and a half times the national average. Rates in Saitama (199.6) and Fukui (194.4) were nearly as high. In the remaining 43 prefectures rates ranged from 10.9 to 163.3.

The tuberculosis case rate in February (535.7) was about a fourth higher than in January (457.3). It was almost a sixth above the rate (486.4) in February 1950 and approximately a tenth above the corresponding 1949 rate (519.2). Rates increased in all but six prefectures over last month. The greatest increases occurred in Kagawa (from 162.8 to 586.9) and Kanagawa (429.7 to 797.5). The most nearly comparable decrease was in Hiroshima (580.6 to 471.2). The highest rates, from 60 to 40 percent above the national average, were recorded by Osaka (913.0), Tokyo-to (843.4), Toyama (814.7), and Fukui (806.8). Rates in the 42 remaining prefectures ranged from 247.3 to 797.5. Of the total tuberculosis cases reported this month (36,365), 90 percent (32,794) were designated as respiratory tuberculosis.

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775 013

The pneumonia case rate in February (388.1) was 22 percent higher than in January (316.9). It was 25 percent higher than the February 1950 rate (310.5) and 52 percent above the corresponding 1949 rate (255.2). Three-fourths (34) of the prefectures reported increases in rates over January, the remaining 12 reported decreases. The most notable increases occurred in Toyama (from 1,177.7 to 1,576.7), Gumma (610.9 to 996.4), Kagawa (104.0 to 432.3), and Saitama (932.2 to 1,259.0). The current rate in Toyama Prefecture was four times, and that in Saitama three times as high as the national average. At the other extreme, rates in Shimane (112.0) and Yamaguchi (125.1) were about 70 percent below the national average.

The February case rate for influenza (46.0) was 20 percent above the January rate (38.3). It was about 75 percent below the rate (178.0) in February 1950. The rate in February 1949 was 3.4. There were decreases in rates from last month in 27 prefectures, increases in 8, and no change in 1. There have been no cases for two or more months in the 10 remaining prefectures. The increase in the national rate is primarily attributable to increases in Gifu (from 140.8 to 811.1), Niigata (64.7 to 395.5), Shizuoka (1.6 to 180.7), and Hokkaido (37.4 to 151.0). Comparable decreases occurred in Toyama (from 408.0 to 34.6), Ehime (318.9 to 7.7), and Yamaguchi (from 183.1 to 10.1). Excluding the four prefectures mentioned above with large increases, 26 additional prefectures reported current cases with rates ranging from 0.6 to 66.0.

The poliomyelitis case rate this month (1.9) was about the same as last month (2.0), February of last year (1.8), and February 1949 (1.8). Rates increased over last month in 19 prefectures, decreased in 14, and did not change in 5. The eight remaining prefectures have reported no cases for two months or longer. Fifteen prefectures reported no current cases. Rates in the 31 prefectures reporting ranged from 0.4 to 10.2.

The current case rate for tetanus was 1.t compared with 1.7 last month, 1.4 in February of last year, and 2.2 in the same period of 1949. Rates decreased in 19 prefectures and increased in 13. The 14 remaining prefectures were equally divided between those reporting the same rate both months and those reporting no cases either month. For the 31 prefectures reporting current cases rates ranged from 0.5 to 5.9.

The case rate for puerperal infection increased from 0.8 in January to 1.0 in February. Rates in February 1950 and 1949 were 1.1 and 1.2 respectively. Rates decreased from January in 15 prefectures, increased in 12, and remained the same in 6. There have been no cases for two or more months in 13 prefectures. For the second consecutive month the highest prefectural rate was in Shiga (6.0). Rates in the 25 additional prefectures reporting current cases ranged from 0.4 to 4.4.

The rabies case rate was less than 0.1 in February 1951 and in February 1949. In January 1951 and February 1950 it was 0.1. All current cases were reported from Saitama Prefecture, where the rate was 0.6.

There were no cases of anthrax reported in February compared with one in January. There were also no cases in February 1950 or 1949.

The current case rate for leprosy was 0.6, compared with 0.5 both last month and in February of last year, and 0.9 in February 1949. There have been no cases reported for two or more months in 18 prefectures. Rates increased over last month in 15 prefectures, decreased in 11, and remained the same in 2. For the 22 prefectures reporting current cases rates ranged from 0.2 to 4.1.

The trachoma case rate was a third higher in February (129.1) than in January (96.7), but about one-eighth below the rates for February 1950 (146.0) and 1949 (147.9). Three-fourths (35) of the prefectures reported increases in rates over January and the remaining 11 decreases. Current prefectural rates ranged from 16.8 to 843.0.

There were no current cases of infectious diarrhea reported. During the comparable periods rates were 0.1 in January, 0.2 in February 1950, and 0.3 in February 1949.

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775 013

The case rate for schistosomiasis increased from 0.2 to 0.5. The rate was also 0.5 in February 1950. Data are not available for 1949. Current cases were reported by Yamanashi (41.5), Fukuoka (2.2), and Hiroshima (1.2).

The case rate for filariasis was the same (0.1) in February as in January. In February 1950 the rate was 0.2. All current cases were reported from the island of Kyushu, where prefectural rates were 2.9 in Kagoshima, 2.7 in Saga, 1.4 in Kumamoto, 0.8 in Nagasaki, and zero in the other three prefectures.

There were 22,949 cases of the four venereal diseases compared with 18,680 last month. The current case rate for gonorrhea was higher than in either last month or February of last year. Rates for the other three diseases were higher than in January but lower than in the previous February.

The case rate for syphilis in February (114.7) was about a third above the January rate (86.6), but about a third below the rate (168.9) in February 1950, and well under half of the corresponding 1949 rate (267.6). Prefectural rates ranged from 33.9 to 347.6.

The gonorrhea case rate was nearly 20 percent higher in February (223.0) than in January (187.5) and about 10 percent higher than in February of last year (201.8), but slightly lower than in February 1949 (228.7). Prefectural rates ranged from 38.3 to 886.2.

The current case rate for chancroid (18.9) was higher than in January (16.2) but lower than in February 1950 and 1949 (20.3 and 37.9 respectively.) Prefectural rates ranged from zero to 114.4

The rate for lymphogranuloma venereum was 0.4 currently compared with 0.3 last month. Rates for February 1950 and 1940 were 0.7 and 0.8 respectively. Two-thirds (30) of the prefectures reported no current cases. Rates ranged from 0.3 to 4.2 in the remaining 16.

Footnotes:

1/ These diseases are diphtheria, dysentery, typhoid fever, paratyphoid fever, smallpox, typhus fever, malaria, Japanese "B" encephalitis, scarlet fever, epidemic meningitis, cholera, and plague.

2/ February 1951, 1950, and 1949 and January 1951 were all four week periods. Rates for 1951 and 1950 are based upon the estimated population as of 1 July 1950, and for 1949 on the estimated population as of 1 July 1949. Comparisons of data should be based upon rates rather than numbers where there are differences in the base population.

3/ Monthly data are not available for the period 1928-1934 inclusive.

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775 013

SUMMARY REPORT OF CASES AND DEATHS FROM
COMMUNICABLE DISEASES IN JAPAN [illegible]
FOUR WEEK PERIOD ENDED 24 FEBRUARY 1951

| PREFECTURE | DIPHTHERIA Cases Number | DIPHTHERIA Cases Rate | DIPHTHERIA Deaths Number | DIPHTHERIA Deaths Rate | DYSENTERY Cases Number | DYSENTERY Cases Rate | DYSENTERY Deaths Number | DYSENTERY Deaths Rate |
|---|---|---|---|---|---|---|---|---|
| HOKKAIDO | 95 | 28.6 | 7 | 2.1 | 23 | 6.9 | - | - |
| AOMORI | 20 | 20.2 | 2 | 2.0 | 10 | 10.1 | - | - |
| IWATE | 14 | 13.5 | 2 | 1.9 | 10 | 9.6 | 3 | 2.9 |
| MIYAGI | 30 | 23.3 | 2 | 1.6 | 26 | 20.2 | 2 | 1.6 |
| AKITA | 28 | 27.7 | 2 | 2.0 | 9 | 8.9 | 1 | 1.0 |
| YAMAGATA | 8 | 7.6 | - | - | 14 | 13.3 | 1 | 1.0 |
| FUKUSHIMA | 42 | 26.4 | 6 | 3.8 | 28 | 17.6 | 3 | 1.9 |
| IBARAKI | 13 | 8.2 | 1 | 0.6 | 12 | 7.6 | 5 | 3.2 |
| TOCHIGI | 18 | 15.0 | - | - | 11 | 9.2 | 4 | 3.3 |
| GUMMA | 12 | 9.7 | 2 | 1.6 | 12 | 9.7 | - | - |
| SAITAMA | 21 | 12.7 | 1 | 0.6 | 52 | 31.4 | 7 | 4.2 |
| CHIBA | 18 | 10.9 | 3 | 1.8 | 35 | 21.2 | 7 | 4.2 |
| TOKYO | 87 | 17.9 | 11 | 2.3 | 239 | 49.3 | 22 | 4.5 |
| KANAGAWA | 29 | 15.1 | 3 | 1.6 | 43 | 22.4 | 12 | 6.2 |
| NIIGATA | 41 | 21.6 | 7 | 3.7 | 128 | 67.3 | 11 | 5.8 |
| TOYAMA | 19 | 24.4 | 2 | 2.6 | 4 | 5.1 | - | - |
| ISHIKAWA | 13 | 17.6 | 3 | 4.1 | 9 | 12.2 | - | - |
| FUKUI | 6 | 10.3 | 1 | 1.7 | 1 | 1.7 | - | - |
| YAMANASHI | 5 | 8.0 | - | - | - | - | - | - |
| NAGANO | 10 | 6.3 | 2 | 1.3 | 6 | 3.8 | 1 | 0.6 |
| GIFU | 13 | 10.9 | 2 | 1.7 | 5 | 4.2 | 1 | 0.8 |
| SHIZUOKA | 18 | 9.4 | 4 | 2.1 | 9 | 4.7 | 3 | 1.6 |
| AICHI | 28 | 10.7 | 2 | 0.8 | 40 | 15.3 | 8 | 3.1 |
| MIE | 26 | 23.0 | 1 | 0.9 | 5 | 4.4 | 1 | 0.9 |
| SHIGA | 7 | 10.5 | 1 | 1.5 | - | - | - | - |
| KYOTO | 33 | 23.3 | 1 | 0.7 | 14 | 9.9 | 1 | 0.7 |
| OSAKA | 46 | 15.4 | 5 | 1.7 | 24 | 8.1 | 8 | 2.7 |
| HYOGO | 38 | 14.9 | 7 | 2.7 | 22 | 8.6 | 4 | 1.6 |
| NARA | 15 | 25.4 | 3 | 5.1 | 1 | 1.7 | - | - |
| WAKAYAMA | 15 | 19.8 | 1 | 1.3 | - | - | - | - |
| TOTTORI | 3 | 6.5 | 1 | 2.2 | 3 | 6.5 | - | - |
| SHIMANE | 11 | 15.6 | - | - | - | - | - | - |
| OKAYAMA | 4 | 3.1 | - | - | 5 | 3.9 | 2 | 1.6 |
| HIROSHIMA | 32 | 19.9 | 7 | 4.4 | 8 | 5.0 | 5 | 3.1 |
| YAMAGUCHI | 32 | 26.9 | 3 | 2.5 | 1 | 0.8 | - | - |
| TOKUSHIMA | 17 | 25.0 | 1 | 1.5 | 1 | 1.5 | - | - |
| KAGAWA | 15 | 20.5 | - | - | 3 | 4.1 | 1 | 1.4 |
| EHIME | 25 | 21.3 | 1 | 0.9 | 9 | 7.7 | 2 | 1.7 |
| KOCHI | 6 | 8.9 | - | - | 1 | 1.5 | - | - |
| FUKUOKA | 66 | 24.2 | 3 | 1.1 | 25 | 9.2 | - | - |
| SAGA | 20 | 27.4 | 1 | 1.4 | 3 | 4.1 | 2 | 2.7 |
| NAGASAKI | 61 | 48.0 | 1 | 0.8 | 2 | 1.6 | - | - |
| KUMAMOTO | 29 | 20.5 | 6 | 4.2 | 6 | 4.2 | 1 | 0.7 |
| OITA | 36 | 37.2 | 1 | 1.0 | 4 | 4.1 | - | - |
| MIYAZAKI | 34 | 40.3 | 7 | 8.3 | 2 | 2.4 | 2 | 2.4 |
| KAGOSHIMA | 28 | 20.1 | 3 | 2.2 | - | - | - | - |
| *Feb 1951 | 1,187 | 18.5 | 119 | 1.9 | 865 | 13.5 | 120 | 1.9 |
| *Jan 1951 | 1,083 | 16.8 | 155 | 2.4 | 741 | 11.5 | 103 | 1.6 |
| *Feb 1950 | 1,276 | 19.8 | 167 | 2.6 | 344 | 5.4 | 115 | 1.8 |

See footnotes at end of table.

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775 013



| PREFECTURE | TYPHOID FEVER Cases Number | Cases Rate | Deaths Number | Deaths Rate | PARATYPHOID FEVER Cases Number | Cases Rate | Deaths Number | Deaths Rate |
|---|---|---|---|---|---|---|---|---|
| HOKKAIDO | 1 | 0.3 | - | - | 1 | 0.3 | - | - |
| AOMORI | - | - | - | - | - | - | - | - |
| IWATE | 1 | 1.0 | - | - | - | - | - | - |
| MIYAGI | 5 | 8.9 | 3 | 2.3 | 2 | 1.6 | - | - |
| AKITA | - | - | - | - | - | - | - | - |
| YAMAGATA | - | - | - | - | - | - | - | - |
| FUKUSHIMA | - | - | - | - | - | - | - | - |
| IBARAKI | 2 | 1.3 | - | - | 1 | 0.6 | - | - |
| TOCHIGI | 3 | 2.5 | - | - | - | - | - | - |
| GUMMA | 1 | 0.8 | - | - | - | - | - | - |
| SAITAMA | 5 | 3.0 | 2 | 1.2 | 1 | 0.6 | - | - |
| CHIBA | 4 | 2.4 | - | - | - | - | - | - |
| TOKYO | 24 | 4.9 | 1 | 0.2 | 6 | 1.2 | - | - |
| KANAGAWA | 3 | 1.6 | - | - | 5 | 2.6 | - | - |
| NIIGATA | 5 | 2.6 | - | - | 2 | 1.1 | 1 | 0.5 |
| TOYAMA | 1 | 1.3 | - | - | - | - | - | - |
| ISHIKAWA | - | - | - | - | 1 | 1.4 | - | - |
| FUKUI | 4 | 6.9 | - | - | 1 | 1.7 | - | - |
| YAMANASHI | 1 | 1.6 | - | - | - | - | - | - |
| NAGANO | 1 | 0.6 | - | - | - | - | - | - |
| GIFU | 5 | 4.2 | - | - | 1 | 0.8 | - | - |
| SHIZUOKA | 6 | 3.1 | - | - | 4 | 2.1 | - | - |
| AICHI | 6 | 2.3 | 1 | 0.4 | - | - | - | - |
| MIE | 3 | 2.7 | - | - | 4 | 3.5 | - | - |
| SHIGA | - | - | - | - | - | - | - | - |
| KYOTO | 14 | 9.9 | - | - | 3 | 2.1 | - | - |
| OSAKA | 8 | 2.7 | - | - | 5 | 1.7 | 1 | 0.3 |
| HYOGO | 5 | 2.0 | - | - | 1 | 0.4 | - | - |
| NARA | 1 | 1.7 | - | - | - | - | - | - |
| WAKAYAMA | 2 | 2.6 | 1 | 1.3 | - | - | - | - |
| TOTTORI | 1 | 2.2 | - | - | - | - | - | - |
| SHIMANE | 1 | 1.4 | - | - | 1 | 1.4 | - | - |
| OKAYAMA | 1 | 0.8 | - | - | 1 | 0.8 | - | - |
| HIROSHIMA | 21 | 13.1 | 2 | 1.2 | 2 | 1.2 | 1 | 0.6 |
| YAMAGUCHI | 2 | 1.7 | - | - | 1 | 0.8 | - | - |
| TOKUSHIMA | 6 | 8.8 | - | - | 1 | 1.5 | - | - |
| KAGAWA | 1 | 1.4 | 1 | 1.4 | 1 | 1.4 | - | - |
| EHIME | 1 | 0.9 | - | - | - | - | - | - |
| KOCHI | 5 | 7.4 | - | - | - | - | - | - |
| FUKUOKA | 5 | 1.8 | - | - | - | - | - | - |
| SAGA | - | - | - | - | - | - | - | - |
| NAGASAKI | 5 | 3.9 | - | - | - | - | - | - |
| KUMAMOTO | 1 | 0.7 | - | - | - | - | - | - |
| OITA | 1 | 1.0 | - | - | - | - | - | - |
| MIYAZAKI | - | - | - | - | 1 | 1.2 | - | - |
| KAGOSHIMA | - | - | - | - | - | - | - | - |
| *Feb 1951 | 162 | 2.5 | 11 | 0.2 | 46 | 0.7 | 3 | 0.0 |
| *Jan 1951 | 117 | 1.8 | 14 | 0.2 | 39 | 0.6 | - | - |
| *Feb 1950 | 218 | 3.4 | 41 | 0.6 | 60 | 0.9 | 2 | 0.0 |

See footnotes at end of table.

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775 013



| PREFECTURE | SMALLPOX | | | | TYPHUS FEVER | | | |
|---|---|---|---|---|---|---|---|---|
| | Cases | | Deaths | | Cases | | Deaths | |
| | Number | Rate | Number | Rate | Number | Rate | Number | Rate |
| HOKKAIDO | - | - | - | - | - | - | - | - |
| AOMORI | - | - | - | - | - | - | - | - |
| IWATE | - | - | - | - | - | - | - | - |
| MIYAGI | - | - | - | - | - | - | - | - |
| AKITA | - | - | - | - | - | - | - | - |
| YAMAGATA | - | - | - | - | - | - | - | - |
| FUKUSHIMA | - | - | - | - | - | - | - | - |
| IBARAKI | - | - | - | - | - | - | - | - |
| TOCHIGI | - | - | - | - | - | - | - | - |
| GUMMA | - | - | - | - | - | - | - | - |
| SAITAMA | - | - | - | - | - | - | - | - |
| CHIBA | - | - | - | - | - | - | - | - |
| TOKYO | - | - | - | - | - | - | - | - |
| KANAGAWA | - | - | - | - | - | - | - | - |
| NIIGATA | - | - | - | - | - | - | - | - |
| TOYAMA | - | - | - | - | - | - | - | - |
| ISHIKAWA | - | - | - | - | - | - | - | - |
| FUKUI | - | - | - | - | - | - | - | - |
| YAMANASHI | - | - | - | - | - | - | - | - |
| NAGANO | - | - | - | - | - | - | - | - |
| GIFU | - | - | - | - | - | - | - | - |
| SHIZUOKA | - | - | - | - | - | - | - | - |
| AICHI | - | - | - | - | - | - | - | - |
| MIE | - | - | - | - | - | - | - | - |
| SHIGA | - | - | - | - | - | - | - | - |
| KYOTO | - | - | - | - | - | - | - | - |
| OSAKA | - | - | - | - | 3 | 1.0 | - | - |
| HYOGO | 9 | 3.5 | 1 | 0.4 | - | - | - | - |
| NARA | - | - | - | - | - | - | - | - |
| WAKAYAMA | - | - | - | - | - | - | - | - |
| TOTTORI | - | - | - | - | - | - | - | - |
| SHIMANE | - | - | - | - | - | - | - | - |
| OKAYAMA | - | - | - | - | - | - | - | - |
| HIROSHIMA | - | - | - | - | - | - | - | - |
| YAMAGUCHI | - | - | - | - | - | - | - | - |
| TOKUSHIMA | - | - | - | - | - | - | - | - |
| KAGAWA | - | - | - | - | - | - | - | - |
| EHIME | - | - | - | - | - | - | - | - |
| KOCHI | - | - | - | - | - | - | - | - |
| FUKUOKA | - | - | - | - | - | - | - | - |
| SAGA | - | - | - | - | - | - | - | - |
| NAGASAKI | - | - | - | - | - | - | - | - |
| KUMAMOTO | - | - | - | - | - | - | - | - |
| OITA | - | - | - | - | - | - | - | - |
| MIYAZAKI | - | - | - | - | - | - | - | - |
| KAGOSHIMA | - | - | - | - | - | - | - | - |
| *Feb 1951 | 9 | 0.1 | 1 | 0.0 | 3 | 0.5 | - | - |
| *Jan 1951 | - | - | - | - | 2 | 0.0 | - | - |
| *Feb 1950 | - | - | 1 | 0.0 | 476 | 7.4 | 24 | 0.4 |

See footnotes at end of table.

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775 013



| PREFECTURE | MALARIA Cases Number | MALARIA Cases Rate | MALARIA Deaths Number | MALARIA Deaths Rate | JAPANESE "B" ENCEPHALITIS Cases Number | JAPANESE "B" ENCEPHALITIS Cases Rate | JAPANESE "B" ENCEPHALITIS Deaths Number | JAPANESE "B" ENCEPHALITIS Deaths Rate |
|---|---|---|---|---|---|---|---|---|
| HOKKAIDO | 2 | 0.6 | - | - | - | - | - | - |
| AOMORI | - | - | - | - | - | - | - | - |
| IWATE | - | - | - | - | - | - | - | - |
| MIYAGI | - | - | - | - | - | - | - | - |
| AKITA | - | - | - | - | - | - | - | - |
| YAMAGATA | - | - | - | - | - | - | - | - |
| FUKUSHIMA | - | - | - | - | - | - | - | - |
| IBARAKI | 1 | 0.6 | 1 | 0.6 | - | - | - | - |
| TOCHIGI | - | - | - | - | - | - | - | - |
| GUMMA | 3 | 2.4 | - | - | - | - | - | - |
| SAITAMA | 1 | 0.6 | - | - | - | - | - | - |
| CHIBA | - | - | - | - | - | - | - | - |
| TOKYO | 3 | 0.6 | 1 | 0.2 | - | - | - | - |
| KANAGAWA | 1 | 0.5 | - | - | - | - | - | - |
| NIIGATA | 1 | 0.5 | - | - | - | - | - | - |
| TOYAMA | - | - | - | - | - | - | - | - |
| ISHIKAWA | - | - | - | - | - | - | - | - |
| FUKUI | - | - | - | - | - | - | - | - |
| YAMANASHI | 1 | 1.6 | - | - | - | - | - | - |
| NAGANO | 1 | 0.6 | - | - | - | - | - | - |
| GIFU | - | - | - | - | - | - | 1 | 0.5 |
| SHIZUOKA | - | - | - | - | - | - | - | - |
| AICHI | 1 | 0.4 | - | - | - | - | - | - |
| MIE | 1 | 0.9 | - | - | - | - | - | - |
| SHIGA | - | - | - | - | - | - | - | - |
| KYOTO | 1 | 0.7 | - | - | - | - | - | - |
| OSAKA | 2 | 0.7 | - | - | - | - | - | - |
| HYOGO | 1 | 0.4 | - | - | - | - | - | - |
| NARA | - | - | - | - | - | - | - | - |
| WAKAYAMA | - | - | - | - | - | - | - | - |
| TOTTORI | - | - | - | - | - | - | - | - |
| SHIMANE | 1 | 1.4 | - | - | - | - | - | - |
| OKAYAMA | 1 | 0.8 | - | - | - | - | - | - |
| HIROSHIMA | 2 | 1.2 | - | - | - | - | - | - |
| YAMAGUCHI | - | - | - | - | - | - | - | - |
| TOKUSHIMA | - | - | - | - | - | - | - | - |
| KAGAWA | - | - | - | - | - | - | - | - |
| EHIME | - | - | - | - | - | - | 1 | 1.5 |
| KOCHI | - | - | - | - | - | - | - | - |
| FUKUOKA | - | - | - | - | - | - | - | - |
| SAGA | 1 | 1.4 | - | - | - | - | - | - |
| NAGASAKI | - | - | - | - | - | - | - | - |
| KUMAMOTO | 3 | 2.1 | 1 | 0.7 | - | - | - | - |
| OITA | - | - | - | - | - | - | - | - |
| MIYAZAKI | - | - | - | - | - | - | - | - |
| KAGOSHIMA | - | - | - | - | - | - | - | - |
| *Feb 1951 | 28 | 0.4 | 3 | 0.0 | - | - | 2 | 0.0 |
| *Jan 1951 | 26 | 0.4 | 5 | 0.1 | 1 | 0.0 | - | - |
| *Feb 1950 | 51 | 0.8 | 4 | 0.1 | - | - | 3 | 0.0 |

See footnotes at end of table.

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775 013



| PREFECTURE | SCARLET FEVER Cases Number | SCARLET FEVER Cases Rate | SCARLET FEVER Deaths Number | SCARLET FEVER Deaths Rate | EPIDEMIC MENINGITIS Cases Number | EPIDEMIC MENINGITIS Cases Rate | EPIDEMIC MENINGITIS Deaths Number | EPIDEMIC MENINGITIS Deaths Rate |
|---|---|---|---|---|---|---|---|---|
| HOKKAIDO | 14 | 4.2 | - | - | 3 | 0.9 | - | - |
| AOMORI | 3 | 3.0 | - | - | 1 | 1.0 | - | - |
| IWATE | 2 | 1.9 | - | - | - | - | - | - |
| MIYAGI | 5 | 3.9 | - | - | 4 | 3.1 | 2 | 1.6 |
| AKITA | 4 | 4.0 | - | - | 3 | 3.0 | - | - |
| YAMAGATA | 2 | 1.9 | - | - | 6 | 5.7 | 1 | 1.0 |
| FUKUSHIMA | 3 | 1.9 | - | - | 3 | 1.9 | 1 | 0.6 |
| IBARAKI | - | - | - | - | 4 | 2.5 | 1 | 0.6 |
| TOCHIGI | 2 | 1.7 | - | - | - | - | - | - |
| GUMMA | 1 | 0.8 | - | - | 1 | 0.8 | 1 | 0.8 |
| SAITAMA | 6 | 3.6 | - | - | 1 | 0.6 | 2 | 1.2 |
| CHIBA | 5 | 3.0 | - | - | 1 | 0.6 | - | - |
| TOKYO | 59 | 12.2 | - | - | 21 | 4.3 | 4 | 0.8 |
| KANAGAWA | 14 | 7.3 | - | - | 6 | 3.1 | 2 | 1.0 |
| NIIGATA | 4 | 2.1 | - | - | 2 | 1.1 | - | - |
| TOYAMA | 1 | 1.3 | - | - | 1 | 1.3 | - | - |
| ISHIKAWA | - | - | - | - | - | - | - | - |
| FUKUI | 2 | 3.4 | - | - | - | - | - | - |
| YAMANASHI | - | - | - | - | - | - | - | - |
| NAGANO | 6 | 3.8 | - | - | 2 | 1.3 | - | - |
| GIFU | 2 | 1.7 | - | - | 1 | 0.8 | - | - |
| SHIZUOKA | 5 | 2.6 | - | - | 2 | 1.0 | - | - |
| AICHI | 18 | 6.9 | - | - | 3 | 1.1 | - | - |
| MIE | 7 | 6.2 | - | - | 1 | 0.9 | - | - |
| SHIGA | 12 | 18.0 | - | - | - | - | - | - |
| KYOTO | 26 | 18.4 | - | - | 8 | 5.6 | 2 | 1.4 |
| OSAKA | 42 | 14.1 | 1 | 0.3 | 7 | 2.3 | 3 | 1.0 |
| HYOGO | 19 | 7.4 | - | - | 2 | 0.8 | - | - |
| NARA | 2 | 3.4 | - | - | - | - | - | - |
| WAKAYAMA | 2 | 2.6 | - | - | - | - | - | - |
| TOTTORI | - | - | - | - | 2 | 4.3 | - | - |
| SHIMANE | 1 | 1.4 | - | - | 1 | 1.4 | 1 | 1.4 |
| OKAYAMA | - | - | - | - | 1 | 0.8 | - | - |
| HIROSHIMA | 3 | 1.9 | - | - | 2 | 1.2 | - | - |
| YAMAGUCHI | 1 | 0.8 | - | - | 2 | 1.7 | 1 | 0.8 |
| TOKUSHIMA | 2 | 2.9 | - | - | - | - | - | - |
| KAGAWA | 4 | 5.5 | - | - | - | - | - | - |
| EHIME | 1 | 0.9 | - | - | - | - | - | - |
| KOCHI | 1 | 1.5 | - | - | 1 | 1.5 | - | - |
| FUKUOKA | 15 | 5.5 | 1 | 0.4 | 2 | 0.7 | 1 | 0.4 |
| SAGA | 1 | 1.4 | - | - | - | 1.4 | - | - |
| NAGASAKI | 3 | 2.4 | - | - | 2 | 1.6 | - | - |
| KUMAMOTO | - | - | - | - | - | - | - | - |
| OITA | - | - | - | - | - | - | - | - |
| MIYAZAKI | 3 | 3.6 | 1 | 1.2 | 1 | 1.2 | - | - |
| KAGOSHIMA | - | - | - | - | - | - | - | - |
| *Feb 1951 | 303 | 4.7 | 3 | 0.0 | 98 | 1.5 | 22 | 0.3 |
| *Jan 1951 | 308 | 4.8 | 1 | 0.0 | 80 | 1.2 | 16 | 0.2 |
| *Feb 1950 | 316 | 4.9 | - | - | 90 | 1.4 | 21 | 0.3 |

See footnotes at end of table.

DECLASSIFIED E.O. 12065 SECTION 3-402/NNDG NO. 775 013



| PREFECTURE | MEASLES Cases | | WHOOPING COUGH Cases | |
|---|---|---|---|---|
| | Number | Rate | Number | Rate |
| HOKKAIDO | 1,170 | 352.5 | 361 | 108.8 |
| AOMORI | 19 | 19.2 | 92 | 92.8 |
| IWATE | 139 | 133.6 | 133 | 127.8 |
| MIYAGI | 180 | 140.1 | 242 | 188.3 |
| AKITA | 64 | 63.3 | 89 | 88.0 |
| YAMAGATA | 44 | 42.0 | 51 | 48.6 |
| FUKUSHIMA | 57 | 35.8 | 148 | 92.9 |
| IBARAKI | 93 | 59.0 | 46 | 29.2 |
| TOCHIGI | 35 | 29.2 | 83 | 69.3 |
| GUMMA | 96 | 77.6 | 121 | 97.8 |
| SAITAMA | 138 | 83.2 | 331 | 199.6 |
| CHIBA | 86 | 52.0 | 89 | 53.8 |
| TOKYO | 1,407 | 290.2 | 189 | 39.0 |
| KANAGAWA | 1,184 | 615,9 | 148 | 77.0 |
| NIIGATA | 54 | 28.4 | 215 | 113.1 |
| TOYAMA | 36 | 46.2 | 82 | 105.2 |
| ISHIKAWA | 145 | 196.0 | 26 | 35.2 |
| FUKUI | 533 | 916.9 | 113 | 194.4 |
| YAMANASHI | 35 | 55.8 | 12 | 19.1 |
| NAGANO | 17 | 10.7 | 260 | 163.3 |
| GIFU | 259 | 217.0 | 99 | 83.0 |
| SHIZUOKA | 315 | 164.9 | 92 | 48.2 |
| AICHI | 167 | 63.7 | 198 | 75.6 |
| MIE | 22 | 19.5 | 101 | 89.5 |
| SHIGA | 22 | 33.1 | 105 | 157.8 |
| KYOTO | 13 | 9.2 | 85 | 60.0 |
| OSAKA | 207 | 69.5 | 156 | 52.3 |
| HYOGO | 412 | 161.1 | 112 | 43.8 |
| NARA | - | - | 18 | 30.5 |
| WAKAYAMA | 22 | 29.0 | 52 | 68.5 |
| TOTTORI | 10 | 21.6 | 52 | 112.1 |
| SHIMANE | 3 | 4.3 | 79 | 112.0 |
| OKAYAMA | 185 | 144.1 | 1110 | 85.7 |
| HIROSHIMA | 170 | 105.7 | 97 | 60.3 |
| YAMAGUCHI | 79 | 66.4 | 13 | 10.9 |
| TOKUSHIMA | 4 | 5.9 | 11 | 16.2 |
| KAGAWA | 52 | 71.1 | 69 | 94.4 |
| EHIME | 197 | 167.5 | 76 | 64.6 |
| KOCHI | 87 | 128.8 | 51 | 75.5 |
| FUKUOKA | 484 | 177.5 | 278 | 101.9 |
| SAGA | 211 | 288.9 | 85 | 116.4 |
| NAGASAKI | 162 | 127.4 | 60 | 47.2 |
| KUMAMOTO | 10 | 7.1 | 183 | 129.6 |
| OITA | 28 | 28.9 | 27 | 27.9 |
| MIYAZAKI | 18 | 21.3 | 175 | 207.5 |
| KAGOSHIMA | 78 | 56.0 | 131 | 94.0 |
| *Feb 1951 | 8,749 | 136.1 | 5,346 | 83.2 |
| *Jan 1951 | 4,403 | 68.5 | 4,098 | 63.7 |
| *Feb 1950 | 4,109 | 63.8 | 11,792 | 183.4 |

See footnotes at end of table.