





C 6.10

R41278

Digitized by the Internet Archive
in 2015

https://archive.org/details/b21901272_0003

Ex libris coll. Reg. Med. L.

FLORA BRITANNICA,

AUCTORE

J. E. S.

JACOBO EDVARDO SMITH, M. D.

SOCIETATIS LINNEANÆ PRÆSIDE,

IMP. NAT. CUR. REGIÆ LONDINENSIS, HOLM. UPSAL.

TAURIN. OLYSSIP. PHILADELPH.—PHYSIOGR.

LUND. BEROLIN. PARIS.

ALIARUMQUE SOCIETATUM SOCIO.

“ Nullius addictus jurare in verba magistri.”

HOR.

VOL. III.

LONDINI:

TYPIS J. TAYLOR, BLACK-HORSE-COURT.

VENEUNT APUD J. WHITE, FLEET-STREET.

MDCCCLIV.



Classis XX.

GYNANDRIA.

Stamina pistillo, supra germen, inserta.

DI'ANDRIA.

Stamina 2.

- 379. ORCHIS. *Nectarium corniculatum.*
- 380. SATYRIUM. *Nectarium inflato-didymum.*
- 381. OPHRYS. *Nectarium subcarinatum, deflexum.*
- 382. MALAXIS. *Nectarium cordatum, erectum; fructificalia amplectens.*
- 384. SERAPIAS. *Nectarium ovatum, gibbum, labio ovato.*
- 383. CYPRIPEDIUM. *Nectarium bilabiatum; labium inferius ventricosum, inflatum.*

HEXANDRIA.

Stamina 6.

- 385. ARISTOLOCHIA. *Stigmata 6. Cal. nulus. Cor. monopetala. Caps. sexocularis.*

XX. GYNANDRIA.

I. DIANDRIA.

379. ORCHIS. *G. Pl. 461. Juss. 65.*

Nectarium corniforme, pone florem.

* *Bulbis indivisis.*

- bifolia.* 1. ORCHIS bulbis indivisis, nectarii labio lanceo-lato integerrimo: cornu longissimo, petalis lateralibus patentibus.
Orchis bifolia. *Sp. Pl. 1331. Huds. 382. With. 21. Hull. 194. Relb. 331. Sibth. 9. Abbot. 191. Curt. Lond. fasc. 6. t. 65. Dicks. H. Sicc. fasc. 3. 16. Engl. Bot. t. 22.*
O. n. 1285. Hall. Hist. v. 2. 146. t. 35. f. 2.
O. hermaphroditica bifolia. Raii Syn. 380. Baub. Hist. v. 2. 772.
O. hermaphroditica. Ger. em. 211.
Satyrion trifolium. Fuchs. Hist. 710.
Testiculi species tertia. Matth. Valgr. v. 2. 233. f. 2.
T. species quinta. Camer. Epit. 625.
β. Orchis bifolia. *Fl. Dan. t. 235.*
O. alba bifolia minor, calcari oblongo. Raii Syn. 380.
O. alba, calcari longo. Baub. Hist. v. 2. 771.
Testiculus psycodes. Ger. em. 211.
Angl. Butterfly Orchis.
In dumetis et nemorosis, solo lutoso humidiusculo.
β in pascais elatioribus.

Perennis. Fl. Junio.

Bulbi ovati, singulus biennis, ut in aliis speciebus. Herba læte

lætè virens, glaberrima, immaculata. Caulis crectus, pedalis, angulatus, basi diphyllos, interdùm triphyllus, supernè foliolosus. Folia petiolata, vaginantia, elliptica, nervosa. Spica laxa. Bracteæ lanceolatæ, longitudo germinum. Flores circiter 12, alterni, sessiles, albi, mane et vesperi imprimis fragrantes. Pctala tria superiorea conniventia; duo lateralia patentissima. Nectarii cornu germine duplo longius, subclavatum; labium oblongum, indivisum, virescens, dependens. Stamina lateralia, distantia; filamenta basi articulata; antheræ clavatae. Germen triquetrum, contortum, suctatum; stigma concaviusculum, virescens, supernè carinatum. Capsula lateraliter dehiscentia. Semina numerosissima, scobiformia, ut in toto ordine naturali. β magnitudine minori tantum differt.

2. ORCHIS bulbis indivisis, nectarii labio trifido *pyramidalis.*
æquali integerrimo supernè bicorni: calcari elongato filiformi.

Orchis pyramidalis. *Sp. Pl.* 1332. *Huds.* 382.
Wittb. 23. *Hull.* 194. *Rehb.* 332. *Sibth.* 9.
Abbot. 191. *Engl. Bot.* t. 110. *Jacq. Austr.*
t. 266.

O. n. 1286. *Hall. Hist.* v. 2. 146. t. 35. f. 1.
O. purpurea, spicâ congestâ pyramidalis. *Raii Syn.* 377. t. 18.
O. parvo flore rubro sive phœnicio. *Baub. Hist.*
v. 2. 764.

Angl. Pyramidal Orchis.

In pascuis siccis, præsertim cretaccis apricis.

Perennis. Fl. Julio.

Bulbi subrotundi. Herba lætè virens, immaculata, glaberrima, sericeo-nitida. Folia lanceolata, acuta, vaginantia, superiora sensim minora. Spica densa, multiflora, pyramidalis, demùm elongata et obtusior, ut in ieone Hallerianâ depingitur. Flores vividè punicei, unicolores, subindè lâctei. Petala tria exteriora patentissima, ovata; interiora lanceolata, conniventia. Nectarii

tarii cornu germine longius, filiforme, acutum; labium trifidum, lacinias æqualibus, integerrimis, basi superne gibbis duobus notabile. Stamina approxinata, propriis foveolis linearibus recondita. Germen gracillimum.

Morio. 3. ORCHIS bulbis indivisis, nectarii labio quadrifido crenulato: cornu obtuso adscendente, petalis conniventibus multinervibus.

Orechis Morio. Sp. Pl. 1333. Huds. 383. Willd. 23. Hull. 194. Relb. 332. Sibth. 10. Abbot. 192. Curt. Lond. fasc. 3. t. 59. Fl. Dan. t. 253, haud bene.

O. n. 1281. Hall. Hist. v. 2. 143. t. 30. f. 1.

O. morio sœmina. Raii Syn. 377. Faill. Par. t. 31. f. 13, 14.

O. minor purpurea et aliorum colorum, cum aliis virentibus. Baub. Hist. v. 2. 761.

Cynosorchis morio sœmina. Ger. em. 208.

Triorchis serapias mas. Fuchs. Hist. 559.

Angl. Meadow Orchis.

In pratis et pascuis humidiusculis.

Perennis. Fl. Maio, Junio.

Bulbi subrotundi, altero remotiusculo. Folia lanceolata, saturatè viridia, immaculata; subtùs pallidiora, nitida. Spica laxiuscula. Bractæ lanceolatæ, membranaceæ, coloratæ, longitudine germinum. Flores purpurei, violacei, vel incarnati, petalis obtusis, conniventibus, lateralibus multinervibus, viridi lineatis. Nectarii cornu fursum curvatum, apice incrassatum, germine parum brevius; labium dilatatum, maculatum, quadrifidum, lacinias lateralibus majoribus, crenatis. Antheræ bifidae.

mascula. 4. ORCHIS bulbis indivisis, nectarii labio quadrifido crenulato: cornu obtuso, petalis exterioribus reflexis trinervibus.

Orchis

- Orchis mascula.* *Sp. Pl.* 1333. *Huds.* 383. *With.* 24. *Hull.* 194. *Relb.* 333. *Sibth.* 10. *Abbot.* 192. *Curt. Lond. fusc.* 2. t. 62. *Dicks. H. Sicc. fusc.* 11. 14. *Woodv. Med. Bot.* t. 90. *Engl. Bot.* t. 631. *Fl. Dan.* t. 457. *Jacq. Ic. rar.* v. 1. t. 180.
O. n. 1283. *Hall. Hist.* v. 2. 144. t. 33. f. 1.
O. morio mas, foliis inaculatis. *Ruii Syn.* 376.
Vaill. Par. t. 31. f. 11, 12.
O. major tota purpurea, maculofo folio. *Bauh.* *Hist.* v. 2. 763.
O. mas angustifolia. *Fuchs. Hist.* 555.
Cynosorchis morio mas. *Ger. em.* 208.
Angl. Early Purple Orchis.

In pascuis et nemorosis copiosè.

Perennis. Fl. Aprili, Maio.

Bulbi ovati, magni. Folia latè-lanceolata, nitida, magis vel minus colore atro-purpureo maculata. Caulis pedalis, teres. Spica laxiuscula, multiflora. Bractæ lanceolatæ, coloratæ, vix longitudine germinis. Flores purpurei, petalis ovatis, trinervibus, unicoloribus, lateralibus reflexis. Nectarii cornu obtusum, horizontale, vel parùm adscendens, germini æquale; labium dependens, basi album, purpureo punctatum, margine quadratum, subcrenatum, laciniis lateralibus majoribus, interinediis denticulo centrali interstineto. Antheræ simplices, flavæ, thecis violaceis inclusæ.

5. ORCHIS bulbis indivisis, nectarii labio quadri-*ustulata*, fido punctis scabro, cornu obtuso brevissimo, petalis distinctis.

Orchis ustulata. *Sp. Pl.* 1333. *Huds.* 384. *With.* 25. *Hull.* 194. *Relb.* 333. *Sibth.* 10. *Abbot.* 192. *Dicks. H. Sicc. fusc.* 3. 17. *Engl. Bot.* t. 18. *Fl. Dan.* t. 103.

O. n. 1273. *Hall. Hist.* v. 2. 138. t. 28. f. 2.

- O. pannonica* quarta. *Chuf. Hist.* v. 1. 268.
Raii Syn. 377.
O. parvis floribus, multis punctis notatis. *Bauh.*
Hist. v. 2. 765.
O. militaris praten sis humilior. *Vaill. Par.* t. 31.
f. 35, 36.
Cynosorchis minor pannonica. *Ger. em.* 207.
Angl. Dwarf Orchis.

In pascuis et collibus siccioribus, crctaccis.

Percnnis. Fl. Junio.

Bulbi ovato-subrotundi. Caulis palmaris, aut vix spithameus. Folia lanceolata, immaculata, subglaucia. Spica brevis, densa, obtusa, demum laxa. Bracteæ ovatae, purpureæ, germine duplo breviores. Flores parvi, petalis distinctis, conniventibus, extus fusco-purpureis. Nectarii cornu germine duplo brevius, incurvum, obtusum, compressissimum; labium dependens, quadrilobum, lobis æqualibus, integerrimis, elevato-punctatis, duobus interioribus subinde cum denticulo interjecto.

militaris. 6. ORCHIS bulbis indivisis, nectarii labio quinqüefido punctis scabro, cornu obtuso brevissimo, petalis confluentibus.

Orchis militaris. *Sp. Pl.* 1333. *Huds.* 384.
With. 26. var. 3. *Hull.* 194. var. 3. *Sibth.*
10. *Sm. Tour on the Continent,* v. 2. 295.
Jacq. Ic. rar. v. 3. t. 598.

O. n. 1277. *Hall. Hist.* v. 2. 140. t. 28. f. 1.
O. galea et alis ferè cinereis. *Bauh. Hist.* v. 2.
757. *Raii Syn.* 378.
O. altera oreades cercopithecophora. *Column.*
Eccphr. 320. f. 2. *Rudb. Elys.* v. 2. 194. f. 8.
O. cercopithecum exprimens. *Rudb. Elys.* v. 2.
194. f. 9.
O. mas latifolia. *Fuchs. Hist.* 554.
O. major. *Cord. Hist.* 128. 2.

O. Vaill. Par. t. 31. f. 21—26.

Cynosorchis major altera. Ger. em. 205.

Satyrion mas. Brumf. Hist. v. 1. 104. Trag. Hist. 778.

β . *Huds. 384.*

Orchis militaris. Engl. Bot. t. 16. With. 25.
var. 2. Hull. 194. var. 2.

O. purpurea. Huds. ed. 1. 334.

O. fusca. Curt. Lond. fasc. 6. t. 64. Jacq. Austr.
t. 307.

O. moravica. Jacq. Ic. rar. v. 1. t. 182.

O. n. 1276. Hall. Hist. v. 2. 140. t. 31.

O. magna, latis foliis, galeâ fuscâ vel nigricante.
Dill. in Raii Syn. 378. t. 19. f. 2.

O. militaris major. Vaill. Par. t. 31. f. 27, 28.

Angl. Military Orchis.

In pratis et pascuis, præcipue solo cretaceo.

α , on Cawsham hills by the Thames' side not far from Reading. *Ray.* Caversham warren, Oxfordshire. *Sibth.* β , at Northfleet near Gravesend; I. Sherard. *Dill.* On chalky banks abounding with Milkwort and Juniper, near woods, and in the woods themselves, in many parts of Kent, especially about Rochester, in great abundance. *Curtis.*

Perennis. Fl. Maio.

Bulbi ovati. Caulis pedalis, vel sesquipedalis. Folia latè lanceolata, lætè viridia, nec glauca, neque maculata, subtùs pallidiora nitida. Spica multiflora, laxiuscula, obtusa. Bracteæ parvæ, breves, membranaceæ, interdùm setaceo-acuminatæ. Flores purpurascentes, vel carnei, petalis connatis, conniventibus; in α acutioribus, pallidis; in β obtusioribus, latioribus, et viridi-fuscis. Nectarii cornu germine duplò brevius, incurvum, obtusum, subinflatum; labium dependens, punctato-scabrum, quinqulobum; lobo centrali minimo; lateralibus latitudine et figurâ valdè variantibus; in α

linearibus, retusis, quandoque angustissimis et elongatis; in β intermediis præcipue dilatatis et crenulatis. Herba dum exsiccatur Anthoxanthi vel Asperulae odoreni spargit. Variedades α et β , cum pluribus intermediis, in solo natali attentè scrutatus sum, atque earum differentias omnino evanescentes, quantumvis primo intuitu notabiles, inveni.

** *Bulbis palmatis.*

latifolia. 7. ORCHIS bulbis subpalmatis, nectarii cornu conico: labio trifido, bracteis flore duplo longioribus.

Orchis latifolia. *Sp. Pl.* 1334. *Huds.* 385. *With.* 26. *Hull.* 194. *Relb.* 334. *Sibth.* 11. *Abbot.* 192. *Curt. Lond. fasc.* 5. t. 65. *Fl. Dan.* t. 266.

O. n. 1279. *Hall. Hist.* v. 2. 142. t. 32. f. 2.

O. palmata pratensis latifolia, longis calcaribus. *Raii Syn.* 380. *Vaill. Par.* t. 31. f. 1—5.

O. palmata palustris latifolia. *Rudb. Elys.* v. 2. 214. f. 12.

Palma Christi mas. *Ger. em.* 220.

Palmata non maculata. *Baub. Hist.* v. 2. 774.

P. sive Serapias palustris latifolia, flore albo subpurpurascente. *Baub. Hist.* v. 2. 775.

Serapias palustris latifolia. *Ger. em.* 222. *Lob.* *Ic.* v. 1. 190.

Cynosorchis palustris platyphylla. *Dalech. Hist.* 1562.

β . Orchis palmata palustris, tota rubra. *Dill. in Raii Syn.* 382.

Palmata floribus impensè rubris. *Baub. Hist.* v. 2. 777.

Angl. Marsh Orchis.

In pratis humidis et palustribus.

Perennis.

Perennis. Fl. Maio, Junio.

Bulbi palmati, lobis inæqualibus, subulatis, rectiusculis. Caulis fistulosus, foliosus. Folia vaginantia, latè lanceolata, superiora parùm minora, omnia apud nos immaculata. Spica multiflora, densa, acutiuscula. Bractæ floribus duplò longiores, erectæ, lanceolatæ, acutæ, foliaceæ, subeoloratæ. Flores incarnati, rosei, vel sanguinci, petalis lateralibus patentibus, maculatis. Nectarii cornu conicum, deflexum, germine aliquantulum brevius; labium rotundatum, apice trifidum, lobis magnitudine et formâ variis, antice pallidum, flavescens, punctatum, lateribus demùm reflexis.

8. ORCHIS bulbis palmatis divaricatis, nectarii *maculata*.
cornu germinibus breviore; labio trilobo plano, petalis patentibus.

Orchis maculata. *Sp. Pl.* 1335. *Huds.* 385.

With. 27. *Hull.* 195. *Rehb.* 334. *Sibth.* 11.

Abbot. 193. *Engl. Bot.* t. 632. *Fl. Dan.*

t. 933.

O. n. 1278. *Hall. Hist.* v. 2. 141. t. 32. f. 1.

O. fœmina altera. *Trag. Hist.* 781.

Palmata speciosiore thyrso, folio maculato. *Raii*
Syn. 381. *Bauh. Hist.* v. 2. 774.

Palina Christi. *Cord. Hist.* 130. 2.

P. Christi fœmina. *Ger. em.* 220.

Satyrium basilicum fœmina. *Fuchs. Hist.* 713.

Angl. Spotted Palmate Orchis.

In pratis, pascuis, et nemoribus, quandoque in fiscis sterilioribus, vulgaris.

Perennis. Fl. Junio, Julio.

Bulbi palmati, lobis elongatis, divaricatis. Caulis solidus, gracilis, foliosus. Folia lanceolata, omnia nigro-maculata. Spica breviuscula, in solo sterili brevissima, subpyramidata, densa. Bractæ vix longitudine germinuni, parùm coloratæ. Flores incarnato-albi, maculis purpureis polymorphis varii, petalis tribus exterioribus patentibus. Nectarii cornu subulatum, ger-

mine

mine brevius, deflexum; labium planum, obcordatum, trilobum, plerumque crenatum.

conopsea. 9. ORCHIS bulbis palmatis, nectarii cornu setaceo germinibus duplò longiore: labio trifido integrimo.

Orchis conopsea. Sp. Pl. 1335. Huds. 385.

With. 28. Hull. 195. Relb. 335. Sibth. 11.

Abbot. 193. Dicks. H. Sicc. fasc. 4. 13. Engl.

Bot. t. 10. Fl. Dan. t. 224.

O. n. 1287. Hell. Hist. v. 2. 147. t. 29. f. 2.

O. palmata minor calcaribus oblongis. Vaill. Par. t. 30. f. 8. Rudb. Elys. v. 2. 212. f. 5.

O. palmata angustifolia minor. Baub. Pin. 85. Rudb. Elys. v. 2. 212. f. 6.

O. palmata montana maxima. Baub. Prod. 31.

Palmata rubella, cum longis calcaribus rubellis. Raii Syn. 381. Baub. Hist. v. 2. 778.

Palma Christi major. Matth. Valgr. v. 2. 237. Camer. Epit. 626.

Satyrium basilicum mas. Fuchs. Hist. 712.

S. foemina. Brunf. Herb. v. 1. 106.

Serapias minor nitente flore. Ger. em. 222.

Angl. Aromatic Orchis.

In pratis et pascuis humidioribus, submontosis.

Perennis. Fl. Junio.

Bulbi palmati, laciniis gracilibus. Caulis ferè sesquipedalis. Folia lanceolata vel linear-lanceolata, immaculata. Spica laxa, elongata, multiflora. Bracteæ germinibus parùm longiores, acuminatæ. Flores purpurei, unicolores, parvi, odore suavissimo caryophyllosum, quandoque albi. Nectarii cornu filiforme, pendens, germe duplò circuitè longius; labium æqualiter trifidum, integrimum.

Ab O. odoratissimâ Linnæi longitudine calcaris dignoseitur.

Obs. O. coriophoram et O. abortivam de industriâ omisi.

O. barbata fœtida minor, flore albo, Raii, quam ad priorem

priorem retulit Hudsonus, varietas Satyrii hircini, quantum ex imperfecto charactere suspicari licet, videatur, neque cum O. coriophorâ convenit. Limodorum Austriacum, sic dictum, a D. Goodyer in Southamptonæ comitatu inventum, (*Raii Syn.* 383), Orobanche cærulea, *Engl. Bot.* t. 423, esse, mihi luce clarius patet. Confer *Ger. em.* 228. et *Tr. of Linn. Soc.* v. 4. 164—169.

380. SATYRIUM. *G. Pl.* 461. *Juss.* 65.

Nectarium inflato-didymum, pone florem.

1. SATYRIUM bulbis indivisis, foliis lanceolatis, *hircinum*.
nectarii labio trifido; laciniâ intermediâ linearis elongatâ tortuosâ.

Satyrium hircinum. *Sp. Pl.* 1337. *Huds.* 386.
With. 29. *Hull.* 195. *Engl. Bot.* t. 34. *Jacq.*
Austr. t. 367.

Orchis n. 1268. *Hall. Hist.* v. 2. 135. t. 25.

O. barbata fœtida. *Raii Syn.* 376. *Bauh. Hist.*
v. 2. 756. *Vaill. Par.* t. 30. f. 6.

O. barbata odore hirci, breviore latiore que folio,
et *O. odore hirci* longiore angustiore que folio.

Rudb. Elys. v. 2. 195.

Tragorchis maximus, et *T. mas.* *Ger. em.* 210.

β. *Orchis barbata fœtida minor*, flore albo. *Raii*
Syn. 376. *ed.* 2. 236.

Angl. Lizard Satyrion.

In pascuis et dumetis, solo cretaceo, rarissimè.

In several places about Dartford; Kent, observed by
Mr. Rouse. *Ray.* About the bushy borders of
some hilly pastures at Darent 2 miles from Dart-
ford, pointed out to me in 1791 by the late Mr.
Lewin.

Perennis. Fl. Julio.

Bulbi subrotundi, magni. Caulis sesquipedalis, quando-
que tripedalis, foliosus, teres, fistulosus. Folia lato-
lanceolata,

lanceolata, glauco-viridia, immaculata. Spica cylindracea, densiuscula, 20—60-flora. Bracteæ linear-lanceolatæ, germine duplò longiores. Flores pedicellati, tristè purpurascentes, odore gravi et viroso. Petala conniventia, viridia, purpureo marginata et lineata. Nectarii sacculus germine triplò brevior, compressus, incurvus; labium trifidum, antice album, sanguineo guttatum, pubescentis, laciniis linearibus, contortis, purpureo-plumbeis, centrali longissimâ, apice saepius prænitorsâ. Germinis ellipticum.

Orchidearum nostratum maxima et rarissima.

- viride.* 2. SATYRIUM bulbis palmatis, foliis ovatis obtusiusculis, nectarii labio lineari trifido: lacinia intermedia minimâ.

Satyrium viride. *Sp. Pl.* 1337. *Huds.* 386. *With.* 30. *Hull.* 195. *Rehb.* 335. *Sibth.* 11. *Abbot.* 193. *Dickf. H. Sicc. fasc.* 4. 14. *Engl. Bot.* t. 94. *Fl. Dan.* t. 77.

Orchis p. 1269. *Hall. Hist.* v. 2. 136. t. 26. f. 2. O. palmata minor, flore luteo-viridi. *Raii Syn.* 381.

O. palmata, flore galericulato dilutè viridi. *Loes.* *Pruß.* 182. t. 59.

O. palmata batrachites. *Rudb. Elysf.* v. 2. 213. f. 10. *Baub. Hist.* v. 2. 772.

O. palmata flore viridi. *Rudb. Elysf.* v. 2. 216. f. 17.

Serapias batrachites. *Ger. em.* 224? *Angl.* Frog Satyriion.

In pratis et pascuis, haud infrequens,

Perennis. Fl. Junio, Julio.

Bulbi palmati. Caulis vix spithameus, foliosus. Folia immaculata; infima latè ovata, obtusa; superiora lanceolata, acutiora. Spica multiflora. Bracteæ floribus duplò longiores, foliaceæ. Flores tristes. Petala conniventia, viridia, apice fusco-purpurascens. Nectarii sacculus

ſaeculus brevissimus; labium longitudine germinis, dependens, linearι-oblongum, luteo-virens vel fuscum, apice trifidum, lacinia intermedia magnitudine variā, ſæpius minutā.

3. SATYRIUM bulbis fasciculatis, foliis lanceo-*albidum*.
latis, nectarii labio trifido acuto: lacinia intermedia longiore.

Satyrium albidum. *Sp. Pl.* 1338. *Huds.* 387.
Witb. 31. *Hull.* 195. *Lightf.* 519. *Dicks.*
H. Sicc. fasc. 1. 15. *Engl. Bot.* t. 505. *Fl. Dan.* t. 115.

Orchis n. 1270. *Hall. Hist.* v. 2. 137. t. 26. f. 1.
O. pusilla alba odorata, radice palmata. *Raii Cat. Pl. Engl. ed.* 2. 217. *Syn. ed.* 1. 173.
ed. 2. 239. *ed.* 3. 381.

Pseudo-*orchis alpina*, flore herbaceo. *Mich. Gen.* 30. t. 26. f. A. B. C.

β. *Orchis palmata*, thyrso specioso, longo, densè stipato, ex viridi albente. *Dill. in Raii Syn.* 382.

Angl. White Satyrion.

In apricis herbidis alpinis. β in pratis alpinis humidis.

ε, on the back of Snowdon, in the way from Llanberis to Caernarvon. *Ray.* At Hafod, Cardiganshire. *Mr. Todd.* In dry hilly pastures of Argyllshire, and in several of the Hebrides. *Lightf.*

β, in moist meadows, not only in Wales, but also about Malham, Yorkshire. *Dr. Richardson.*

Perennis. Fl. Junio.

Bulbi per paria fasciculati, cylindracei, acuti, tertio anno florentes. Caulis spithameus, foliosus. Folia lætè viridia, lanceolata; inferiora obtusa. Spica densa, multiflora. Bracteæ longitudine germinum. Flores

parvi, parùm odorati. Petala albida, subpatentia. Nectarii sacculus elongatus, haud didymus, incurvus; labium virens, trilobum, lobis omnibus plerumque acutis, intermedio longiori, rariùs retuso.

β non nisi magnitudine, ut videtur, differt.

repens. 4. SATYRIUM radicibus fibrosis repentibus, foliis ovatis radicalibus, floribus secundis, nectarii labio indiviso.

Satyrium repens. *Sp. Pl.* 1339. *Huds.* 387.

With. 31. *Hull.* 195. *Lightf.* 520. t. 22.

Dicks. Dr. Pl. 85. *Engl. Bot.* t. 289. *Jacq.*

Austr. t. 369. *Fl. Dan.* t. 812.

Epipactis n. 1295. *Hall. Hist.* v. 2. 153. t. 22. f. 4.

Orchis radice repente. *Camer. Ic.* 35. *Baub.* *Hist.* v. 2. 770. fig. 2, nec 1.

Pseudo-orchis. *Rudb. Elys.* v. 2. 209. f. 8.

Palma Christi radice repente. *Ger. em.* 227.

Pyrola acutifolia polyanthos, radice geniculatâ. *Loës. Pruss.* 210. t. 68.

Angl. Creeping Satyrion.

In sylvis alpinis muscosis Scotiæ, rariùs.

In an old birch wood called Ca bue, or Yellow hill, about 2 miles from the head of Little Loch Broom, Ross-shire. *Lightf.* In a wood opposite to Moy ha' l near Inverness. *Dr. Hope.* About Brodie house in Moray. *J. Brodie, Eyz.*

Perennis. Fl. Julio.

Radix omnino repens, sarmientosa, pubescens. Caules erecti, spithamei, teretes, pubescentes, ferè aphylli, bracteati. Folia radicalia, petiolata, ovata, nervosa, glabra, venis reticulata, subindè purpureo tessellata. Petioli dilatati. Spica multiflora, acuta, densa, pubescens. Bracteæ lanceolatæ, acutæ, floribus longiores. Flores secundi, spiraliter digesti, carnei, fragrantes, extus pubescentes. Petala tria superiora conglutinata; lateralia

teralia duo deflexa, remota. Nectarii sacculus ventricosus, lutescens, petalis subæqualis; labium lanceolatum, roseum, indivisum, sacculo brevius.

381. OPHRYS. G. Pl. 462. Juss. 65.

Nectarium subtus subcarinatum, deflexum.

* *Bulbis ramosis.*

1. OPHRYS bulbis fibroso-fasciculatis, caule vaginato aphylo, nectarii labio bifido. *Nidus avis.*

Ophrys *Nidus avis.* Sp. Pl. 1339. Huds. 388.
With. 32. Hull. 196. Relb. 336. Sibth. 12.
Abbot. 194. Dicks. H. Sicc. fasc. 9. 10. Engl.
Bot. t. 48. Fl. Dan. t. 181.

Epipactis n. 1290. Hall. Hist. v. 2. 149. t. 37.
Nidus avis. Raii Syn. 382. Dalech. Hist. 1073.

Lob. Ic. v. 1. 195.

Pseudoleimodoron. Cluf. Hist. v. 1. 270.

Satyrium abortivum, fide *Nidus avis.* Ger. em.
228.

Angl. Bird's-nest Ophrys.

In sylvis opacis, solo calcareo, sed rariū.

In Kent and Sussex. Ray. Huds. In several parts
of the North of England, also in Norfolk and
Suffolk occasionally.

Perennis. Fl. Maio, Junio.

Planta denso in nemore latens, tota fusca. Caulis fistulosus, angulato-teres, pedalis aut sesquipedalis, squamis alternis vaginatus. Folia nulla. Spica infernè laxa, supernè densa, multiflora. Flores pedicellati. Bracteæ pedicellis duplo circiter longiores. Petala patentiuscula, concava. Nectarium concaviusculum, subtus subcarinatum, anticè elongatum, dependens, bilobum, lobis divaricatis, basi unidentatis. Stamina operculo mobili calyprata.

2. OPIIRYS

- corallo-*
rhiza.
2. OPHRYS bulbis ramosis flexuosis divaricatis,
caule vaginato aphylo, nectarii labio indiviso.
Ophrys corallorrhiza. Sp. Pl. 1339. Hudſ. 388.
With. 32. Hull. 196. Lightf. 523. t. 23.
Fl. Dan. t. 451.
Corallorrhiza n. 1301. Hall. Hist. v. 2. 159.
t. 44.
Orobanche spuria, seu corallorrhiza. Rupp. Jen.
ed. 1. 284. t. 2. f. 3.
O. Sueonum, radice coralloide flore albo. Rudb.
Elys. v. 2. 234. f. 16.
Angl. Coral-rooted Ophrys.

In sylvis alpinis paludosis rarissimè.

Observed only in one place, and there sparingly,
in a moist hanging wood, on the south side,
near the head of Little Loch Broom, Ross-shire.
Lightf.

Perennis. Fl. Junio, Julio.

Radix carnosa, albida, ramosissima, lobis divaricatis. Herba tenella, spithamea, tota luteo-virens. Caulis squamis alternis vaginatus. Folia nulla. Spica paniciflora, laxa. Bracteæ exiguae. Flores pedicellati, parvi. Petala lanceolata, acuta; tria superiora conniventia; duo inferiora deflexa, divaricata, angustiora, longitudine nectarii. Labium ovatum, acutum, certè indivisum. Capsulæ latè ellipticæ, pendulæ, petalis persistentibus coronatæ.

- ovata.*
3. OPHRYS bulbis fibrosis, caule bifolio, foliis ellipticis, nectarii labio linearis bifido.
Ophrys ovata. Sp. Pl. 1340. Hudſ. 388. With.
34. Hull. 196. Relb. 356. Sibth. 12. Abbot.
195. Curt. Lond. fusc. 3. t. 60. Fl. Dan.
t. 127.
Ophris. Fuchs. Hist. 566. Matib. Falgr. v. 2.
565. Camer. Epit. 943.

O. bifolia.

O. bifolia. Ger. em. 402. Rudb. Elys. v. 2.
226. f. 1. etiam f. 2, *trifoliata*.
Epipactis n. 1291. Hall. Hist. v. 2. 150. t. 37.
Bifolium majus, seu *Ophrys major* quibusdāin:
Raii Syn. 385. Baub. Hist. v. 3. p. 2. 533.
Perfoliata mascula et fœmina. Brunf. Herb. v. 1:
182, 183.
Angl. Common Twayblade. Ovate Ophrys.

In nemorosis, dumetis; pratis et pascuis.

Perennis. Fl. Junio.

Radix e fasciculis duobus fibrarum simplicium numerosa-
rum. Herba tota virens. Caulis pedalis vel ultra,
pubescens. Folia medium versus caulis, bina, rariūs
terna, opposita, elliptico-ovata, nervosa, patentia. Spica
triuncialis, gracilis, multiflora. Flores pedicellati, vi-
rides. Petala patentia, inflexa, omnia sursum versus
disposita. Nectarii labium petalis duplo vel triplō lon-
gius, dependens, lineare, apice bilobum, lobis rectis.
Capsula elliptico-globosa, petalis diu immutatis coro-
nata.

4. OPHRYS bulbis fibrosis; caule bifolio; foliis *cordata*,
cordatis, nectarii labio quadrilobo.

Ophrys cordata. Sp. Pl. 1340. Hudf. 388.

With. 35. Hull. 196. Dicks. H. Sicc. fasc. 3.

19. Engl. Bot. t. 358.

Ophrys minima. Baub. Prod. 31. Rudb. Elys.
v. 2. 227. f. 4.

Epipactis n. 1292. Hall. Hist. v. 2. 151. t. 22.
f. 3.

Bifolium minimum. Raii Syn. 385. Baub. Hist.
v. 3. p. 2. 534.

Angl. Least Twayblade. Heart-leaved Moun-
tain Ophrys.

In ericetis spongiosis alpinis:

VOL. III.

C

I

In many parts of Derbyshire, Yorkshire, Northumberland and Lancashire. *Ray.* In the Highlands in many places. *Lightf.* Near Hawick in the south of Scotland. *Mr. Sibbald.* On Ingleborough hill, and on moors between Chatsworth and Sheffield.

Perennis. Fl. Julio.

Habitus omnino præcedentis, sed triplò vel quadruplò minor. Caulis palmaris, aut vix spithameus. Folia bipinnata, cordata. Spica uncialis. Petala undique patentia, virentia, interdum fuscescentia. Nectarii labium quadrilobum, lobis duobus ad basin brevioribus, divaricato-horizontalibus; duobus terminalibus porrectis, elongatis. Capsula globosa.

** *Bulbis rotundatis.*

spiralis. 5. OPHRYS bulbis aggregatis oblongis, caule subfolioso, floribus spiraliter secundis, nectarii labio indiviso crenulato.

Ophrys spiralis. Sp. Pl. 1340. Huds. 389. With. 33. Hull. 196. Relh. 337. Sibth. 12. Abbot. 194. Curt. Lond. fasc. 4. t. 59. Dicks. H. Sicc. fasc. 10. 17. Engl. Bot. t. 541. Fl. Dan. t. 387.

Orchis spiralis alba odorata. Raii Syn. 378.

Triorchis. Ger. em. 218.

T. alba odorata minor. Rudb. Elys. v. 2. 209.
f. 7.

Satyrion odoriferum. Brunf. Herb. v. 1. 105.
Angl. Spiral Ophrys. Ladics' Traces.

In pascuis apricis cretaceis vel glareofisis, quandoque in pratis.

Perennis. Fl. Augusto, Septembri.

Bulbi plures, ovato-oblongi, villosi. Caulis vix spithameus, bracteatus, glaucus, superne pubescens, viscidus.

dus. Folia omnia ferè radicalia, petiolata, ovata, nervosa, glaucescentia. Spica linearis, spiralis; multiflorus. Bracteæ lanceolatæ, acuminatæ, inflexæ, concavæ, extùs pubescentes. Flores secundi, sessiles, parvi, albi, crystallino-micantes, suaveolentes. Petala tria superiora conniventia, lanceolata, obtusa; duo lateralia deflexa, latiora. Nectarii labium oblongum, petalis æquale, obsoletè crenatum, medio virens. Capsula obovata, pilosa.

6. OPHRYS bulbo ovato basi radicato, scapo tri-*Loefelii*, gono, foliis lanceolatis, petalis linearibus, germine obovato.

Ophrys Loefelii. *Sp. Pl.* 1341. *With.* 35.
Hull. 196. *Dicks. H. Sicc. fasc.* 9, 11. *Engl. Bot.* t. 47.

O. lilifolia. *Huds.* 389. *Relb.* 337.

O. paludosa. *Fl. Dan.* t. 877.

O. diphyllos bulbosa. *Loef. Pruss.* 180. t. 58.

O. bifolia bulbosa. *Ger. em.* 403.

Orchis lilifolius minor fabuletorum Zelandiæ et Bataviæ. *Baub. Hist.* v. 2. 770. f. 1, nec 2.
Raii Cant. 105.

Pseudo-orchis bifolia palustris. *Raii Syn.* 382.

Bifolium bulbosum. *Dod. Pempt.* 242.

Angl. Dwarf Ophrys.

In palustribus arcnosis inter Juncos.

On Hinton and Teversham moors near Cambridge.

Ray. Relb. On St. Faith's Newton bogs near Norwich. *Mr. Pitchford.*

Perennis. Fl. Julio.

Bulbus ovatus, squamosus, virens, basi fibrosus et radicatus. Folia radicalia, plerumque bina, petiolata, elliptico-lanceolata, nervosa, lætè viridia, glabra. Petoli dilatati, carinati, equitantes. Scapus palmaris, nudus, trigonus. Racemus laxus, pauciflorus. Bracteæ breves. Flores pedicellati, citrini. Petala linearia,

patentia; tria superiora latiora et parum breviora; duo lateralia angustissima, longiora, labio æqualia. Nectarii labium ovato-rotundatum, carinatum, obsoletè crenatum, luteum. Germen obovatum, retusum.

O. lilifolia Linnæi, huic maximè affinis, discrepat foliis latè ovatis, floribus duplò majoribus, labio fusco, germine elongato, linearie.

monorchis. 7. *OPHRY'S* bulbis globosis altero remoto, cæule nudiusculo, neclarii labio petalisque interioribus trifidis.

Ophrys monorchis. *Sp. Pl.* 1342. *Huds.* 390. *With.* 36. *Hull.* 196. *Rehb.* 338. *Dicks.* *H. Sicc. fasc.* 10. 18. *Engl. Bot. t.* 71. *Fl. Dan. t.* 102.

Orchis n. 1262. *Hall. Hist. v.* 2. 132. *t.* 22. *f.* 2.

O. odorata moschata, sive *Monorchis*. *Raii Syn.* 378.

O. parva autumnalis lutea. *Baub. Hist. v.* 2. 768.

O. colco unico, seu *Monorchis flosculis pallidè viridibus*. *Loef. Pruss.* 184. *t.* 61.

O. trifolia, floribus spicatis herbaceis. *Segu. Veron.* *v.* 2. 131. *t.* 16. *f.* 15.

O. odorata moschata. *Rudb. Elys.* *v.* 2. 207. *f.* 1.

Monorchis. *Mich. Gen.* 30. *t.* 26. *f.* 3.
Angl. Musk Ophrys.

In apricis cretaceis, at minus frequens.

Perennis. Fl. Junio, Julio.

Bulbi subrotundi, altero remoto, pedunculato, minore, Herba latè virens, palmaris. Caulis nudiusculus, teres. Folia lanceolata, plerumque bina radicalia, cum altero minore cauli insidente. Spica densa, multiflora. Bracteæ lanceolatæ, longitudine germinis. Flores parvi, luteo-virides, odore melleo, moschato. Petala tria exteriora latè ovata, integerrima, patentia; duo interiora trifida.

irisida. Nectarii labium petalis interioribus conforme, sed parùm elongatum, et basi gibbum. Capsula elliptica.

8. OPHRYS bulbis subrotundis, caule subfolioso, *anthro-*
nectarii labio linearis tripartito : laciniâ mediâ pophora-
elongatâ bifidâ.

Ophrys anthropophora. *Sp. Pl.* 1343. *Huds.*
 390. *With.* 37. *Hull.* 196. *Relb.* 338. *Curt.*
Lond. fasc. 6. *t.* 66. *Dicks. H. Sicc. fasc.* 15.
 16. *Engl. Bot.* *t.* 29.

Orchis n. 1264. *Hall. Hist.* *v.* 2. 133. *t.* 23.

O. anthropophora oreades. *Raii Syn.* 379. *Co-*
lumn. *Ecphr.* *t.* 320. *f.* 1.

O. Vaill. Par. *t.* 31. *f.* 19, 20.

O. flore nudi hominis effigiem repræsentans.
Rudb. Elys. *v.* 2. 193. *f.* 6.

Angl. Green Man Ophrys.

In pratis et pascuis cretaceis.

Frequent in Kent. *Huds.* At Ashwelthorpe near
 Norwich. *Mr. Crowe.* At Fornacet, Norfolk.
Mr. Fox.

Perennis. Fl. Junio.

Bulbi ovato-subrotundi, villosi. Herba Iætè virens, vel
 subglaucia, glabra. Caulis spithameus, basi foliosus.
 Folia radicalia ovato-lanceolata; caulinum solitarium,
 vaginans. Spica multiflora, elongata. Bracteæ lan-
 ceolatae, flavescentes, vix germini æquales. Petala con-
 niventia, fornicata, purpureo-viridia. Nectarii labium
 dependens, germine duplò ferè longius, flavescens,
 subindè ferrugincum, planum, quadrilobum, lobis linea-
 ribus, obtusiusculis, intermediis brevioribus, basi con-
 natis.

9. OPHRYS bulbis subrotundis, caule folioso, *muscifera.*
nectarii labio quadrilobo elongato convexius-
cule subpubescente, columnâ obtusâ.

- Ophrys muscifera. *Huds.* 391. *With.* 37. *Hull.* 196. *Relb.* 339. *Abbot.* 195. *Dicks. H. Sicc.* fasc. 15. 15. *Engl. Bot.* t. 64.
O. insectifera α , myodes. *Sp. Pl.* 1343.
Orchis n. 1265. *Hall. Hist.* 133. t. 24. f. 2.
O. myodes galeâ et alis herbidis. *Raii Syn.* 379.
Bauh. Hist. v. 2. 767.
O. muscæ corpus referens minor. *Rudb. Elys.* v. 2. 201. f. 11. *Vaill. Par.* t. 31. f. 17, 18.
O. myodes. *Ger. em.* 213.
Vulpinus Testiculus. *Lob. Obs.* 91. f. 1, nec 2.
 β . *O. myodes* major. *Raii Syn.* 379. *Herb. Buddl.*
Angl. Fly Ophrys, or Orchis.

In pratis et pascuis calcareis, rariùs.

In Cambridgeshire, Essex and Suffolk. *Ray.* Plentiful in Kent. *Huds.* On the bath hills at Bungay, and in Earsham wood, Norfolk. *Mr. Woodward.* St. Vincent's rocks, Bristol. *With.* Near Tacolnestone church, Norfolk. *Mr. Humphrey.* About Roche Abbey, Yorkshire. *Mr. Salt.*

Perennis. Fl. Junio.

Bulbi subrotundi. Caulis pedalis, teres, basi foliosus. Folia lanceolata, subglauca. Spica laxa, sœpius sexflora. Bracteæ lanceolatæ, foliaceæ, germinibus longiores. Flores sessiles, muscam pulcherrimè simulantes. Petala tria exteriora viridia, ovata; duo interiora purpurea, linearia, angustissima, subpubescentia; omnia patentia. Nectarii labium dependens, oblongum, convexum, atro-sanguineum, subsericeum, quadrilobum, maculâ cœrulecente in disco, tuberculis duobus nitidis ad basin. Columna staminum obtusa, mutica, fusca.

β in *Herb. Buddleiano* vix differt ab α .

apifera. 10. OPHRYS bulbis subrotundis, caule folioso, nectarii labio convexo villoso quinquefido: laciñiâ terminali recurvata sibulatâ.

Ophrys

Ophrys apifera. *Huds.* 391. *With.* 38. *Hull.* 196. *Relb.* 339. *Sibth.* 13. *Abbot.* 195. *Curt.* *Lond. fasc.* 1. *t.* 66. *Engl. Bot.* *t.* 383.

Orchis fuciflora, galeâ et alis purpurascensibus. *Raii Syn.* 391. *Baub. Hist.* *v.* 2. 766, quoad descriptionem; et *Orchis* sive *Testiculus* sphegodes hirsuto flore. *Ibid.* 767, quoad iconem; optimè monente *Vaillantio*.

O. fucum referens major, foliolis superioribus candidis et purpurascensibus. *Baub. Pin.* 83. *Rudb. Elysf.* *v.* 2. 199. *f.* 7. *Vaill. Par.* 146. *t.* 30. *f.* 9.

Testiculus vulpinus secundus sphegodes. *Ger.* *em.* 212. *Lob. Ic.* *v.* 1. 179.
Angl. Bee *Ophrys*, or *Orchis*.

In pascuis et pratis, solo cretaceo, haud infrequens.

Perennis. Fl. Julio.

Habitus praecedentis, sed major et robustior. Folia latè lanceolata. Bracteæ maximæ. Flores apiformes. Petala tria exteriora reflexa, latè ovata, dilutè purpurea, rariùs alba, carinâ margineque viridi; duo interiora duplò breviora, lanceolata, obtusa, virescentia, intùs villosa. Nectarii labium rotundatum, convexum, ventricosum, villosum, purpureo-ferrugineum, flavo areolatum, punctatum et marginatum; margine quinquefidum, laciniis reflexis, terminali elongatâ, subulatâ, acutâ, iterùm anticè recurvata. Columna staminum porrecta, fornicata, mucronata, virens.

II. OPHRYS bulbis subrotundis, caule folioso, *aranifera*.
nectarii labio convexo villoso quadrifido mucico emarginato.

Ophrys aranifera. *Huds.* 392. *With.* 39. *Hull.* 196. *Relb.* 340. *Sibth.* 13. *Engl. Bot.* *t.* 65.

O. fucifera. *Curt. Lond. fasc.* 6. *t.* 67.

Orchis sive *Testiculus* sphegodes hirsuto flore.

Raii Syn. 380. *Bauk. Hist.* v. 2. 767, quoad descriptionem optimam, nec iconem.

O. fucum referens, colore rubiginoso. *Vaill. Par.* 146. t. 34. f. 15, 16.

O. fucum referens. *Rudb. Elys.* v. 2. 205. f. 25. *Testiculus vulpinus* secundus. *Lob. Obs.* 88. f. 4. *Angl. Spider Ophrys, or Orchis,*

In pascuis cretaccis, vel glareofisis.

In Cambridgeshire and Kent. *Ray. Relb.* Near Bury. *Sir T. G. Cullum*, and *W. Matthew, Esq.* Stone quarries in Oxfordshire. *Sibth.*

Perennis. Fl. Aprilij.

Præcedente humilior, floribus plerumque paucioribus, Petala omnia perpetuò viridia, interioribus glabriusculis, parùmque brevioribus. Nectarii labium rotundatum, convexum, villosum, fusco-ferrugineum, disco lineis duâbus glabratis, lividis, parallelis, cum transversali connexis, notatum, extùs flavescens, margine quadrifidum, reflexum, apice emarginatum. Columna staminum acutiuscula, porrecta.

Ophrys adrachnites, *Tour on the Contin.* v. 2. 295, *Hall. Hist.* n. 1266, alia species est, in Angliâ nondum inventa; aliæ plures etiam inter varietates *O. insestiferæ* Linnæi latenter confusæ.

382. MALAXIS. Swartz.

Nectarium cordatum, concavum, erectum, fructificalia amplectens. Cor. resupinata.

paludosa. 1. **MALAXIS** caule pentagono, foliis pluribus spatulatis apice scabris, racemo multiflora.

Malaxis paludosa. *Swartz. Aet. Holm. ann.* 1789. 127. t. 6. f. 2. *Engl. Bot.* t. 72. *With.* 39. *Hull.* 197.

Ophrys paludosa. *Sp. Pl.* 1341. *Huds.* 389. *Relb.* 337. *Abbot.* 196. *Rose Elem. append.*

450. t. 2. f. 3. *Dicks. H. Sicc. fasc.* 7. 16.

Ophrys

Ophrys bifolia palustris nostras. Pluk. Phyt. t. 247.
f. 2.

Orchis minima bulbosa. Dill. in Raii Syn. 378.
Raii Hist. v. 3. 587.

Bifolium palustre. Raii Syn. ed. 2. 243. ed. 3.
385.

Angl. Marsh Tender-tway-blade.

In paludibus spongiosis sterilibus.

On boggy ground near Gamlingay, Cambridgeshire, also between Hatfield and St. Albans, and at Tunbridge wells. Ray. On Felthorpe bogs near Norwich. Mr. C. Bryant. Near Potton, Bedfordshire. Rev. Mr. Abbot.

Perennis. Fl. Julio.

Bulbi ovati, parvi, basi radicati, squamis vaginati. Caulis vix palmaris, nudiusculus, pentagonus. Folia ovato-spatulata, pauca, alterna; basi dilatato-vaginantia; apice obtusa, scabriuscula. Racemus cylindraceus, multiflorus, densiusculus, virens. Bracteæ ovatæ, parvæ. Flores pedicellati, luteo-virides. Corolla resupinata, petalis tribus exterioribus duplo majoribus, quorum duo superiora erecta æqualia, impar deflexum; duabus interioribus lateralibus, reflexis. Nectarium petalis duplo brevius, erectum, ovatum, concavum, obtusiusculum, columnam fructificationis amplectens.

Orchidearum nostratum minimæ.

383. CYPRIPEDIUM. G. Pl. 464. Juss. 65.

Nectarium bilabiatum; labium inferius ventricosum, inflatum, cavum.

i. CYPRIPEDIUM radicibus fibrosis, caule fo- *Calceo-*
lioso, petalis quaternis lanceolatis acuminatis, *lus.*
labio superiore elliptico canaliculato.

Cypripedium Calceolus. Sp. Pl. 1346. Huds.
392. Willb. 43. Hull. 197. Engl. Bot. t. 1.
Salisb.

Salisb. Tr. of Linn. Soc. v. 1. 76. t. 2. f. 1,
Mill. Ic. t. 242.

Calceolus n. 1300. *Hall. Hist. v. 2. 157. t. 43.*

C. Cypripedium. *Gmel. Sib. v. 1. 2. t. 1.*

C. Mariæ. *Raii Syn. 385.* *Ger. em. 443.*

Damasonii species quibusdam, sive *Calceolus D.*

Mariæ. *Baub. Hist. v. 3. 518.*

Elleborine ferruginea. *Dalech. Hist. 1146.*

Angl. Ladies' Slipper.

In nemorosis et dumetis montosis rarissimè.

In soine woods in Lancashire, and in Helk's wood by Ingleborough, Yorkshire. *Ray.* In the county of Durham. *Rev. Mr. Rudston.*

Perennis. Fl. Junio.

Radix repens, tuberosa, fibris simplicibus, carnosus. Caulis erectus, simplex, pedalis, foliosus, subpubescens, uniflorus, subinde biflorus. Folia alterna, amplexicaulia, sessilia, elliptica, acuminata, multinervia, scabriuscula, tenuia, minimè carnosa. Flos terminalis, subsessilis, nutans, magnus, formosus. Petala quatuor, patentia, lanceolata, acuminata, nervosa, fusco-purpurea, quorum supremum latius, insimum subinde apice bifidum est. Nectarii labium superius petalis triplo brevius, pedicellatum, carnosum, flavum, rubro punctatum, ellipticum, saepius retusum, supra canaliculatum, subiùs carinatum, basi subinde angulatum; inferius longitudine ferè petalorum, dependens, ovatum, ventricosum, flavum, venosum, supernè hians. Columna utrinque flaminisera, apice in stylum ovatum umbilicatum desinens, sub tegmine labii superioris. Germen oblongum, attenuatum, prismaticum, scabrum.

384. SERAPIAS. G. Pl. 462. Juss. 65.

Gærtn. t. 14.

Nectarium ovatum, gibbum, labio ovato.

latifolia. 1. SERAPIAS foliis ovatis amplexicaulibus, floribus cernuis, labio integrum acuminato petalis breviore.

Scrapias

Scrapias latifolia. *Syst. Veg. ed. 14.* 814. *Huds.* 393. *With.* 40. *Hull.* 197. *Lightf.* 526. *Reb.* 340. *Sibth.* 13. *Abbot.* 196. *Engl. Bot. t. 269.* *Fl. Dan. t. 811.* *Dicks. H. Sicc. fasc. 15.* 17. *Ehrb. Phyt. n.* 37. *Freeman. Ic. t. 3.*

S. Helleborine α . *Sp. Pl.* 1344.

Epipactis n. 1297. *Hall. Hist. v. 2.* 154. *t. 40.* *f. 2.*

Helleborine. *Ger. em.* 442. *Dalech. Hist.* 1312.

H. latifolia montana. *Raii Syn.* 383.

β . *H. altera atro rubente flore.* *Raii Syn.* 383.
Herb. Buddle.

Angl. Broad-leaved Helleborine.

In sylvis et nemorosis opacis montosis.

β . On the sides of mountains about Malham and Settle, and other places in the north. *Ray. Mr. Woodward, and Mr. D. Turner.*

Perennis. Fl. Julio, Augusto.

Radix repens, fibris simplicibus, carnosis, pubescentibus. Caulis erectus, ferè bipedalis, simplex, foliosus, pubescens. Folia alterna, amplexicaulia, latè elliptica, acuta, multinervia, tenuia, rigidiuscula, nequaquam carnosa; superiora angustiora et lanceolata. Spica solitaria, terminalis, erecta, multiflora. Bractæ lanceolatæ; inferiores floribus longiores. Flores alterni, pedicellati, cernui, viridi vel fusco purpurascentes. Petala ovata, acuta, concava, conniventia, herbacea. Nectarii labium deflexum, petalis brevius, cordatum, acuminatum, integrimum, purpureum, disco flavescens, lave. Germen obovatum, breve, pubescens.

Herba in β magis pubescens, floribus saturatè rubentibus, ferotinis.

2. SERAPIAS foliis lanceolatis amplexicaulibus, *palustris.* floribus cernuis, labio crenato obtuso petalis æquali.

Serapias

Serapias palustris. *Scop. Carn.* v. 2. 204. *Lightf.* 527. *Curt. Cat.* n. 108. *Engl. Bot.* t. 270. *S. latifolia* γ, *palustris.* *Huds.* 393. *S. longifolia.* *Syst. Veg.* ed. 14. 815. *Sp. Pl.* 1345, γ. *With.* 41. *Hull.* 197. *Relb.* 341. *Sibth.* 14. *Abbot.* 196. *Ehrb. Phyt.* n. 47. *Epipactis* n. 1296. *Hall. Hist.* v. 2. 154. t. 39. *Helleborine.* *Fl. Dan.* t. 267. *H. palustris nostras.* *Raii Syn.* 384. *Angl.* Marsh Helleborine.

In palustribus, solo calcareo vel glaroso.

Perennis. Fl. Julio, Augusto.

Præcedente humilior. Folia ovato-lanceolata. Flores pauciores, cernui, demum penduli, majores quam in priore, magnisque speciosi. Bracteæ floribus breviores. Petala patentia; tria exteriora virentia; duo interiora alba; omnia purpureo-variata. Nectarii labium patens, petalis subæquale, cordatum, obtusum, margine dilatatum, undique crenatum, album, purpureo-lineatum; supernè callo dentato ad basin.

Huic nomen *palustris* aptissimum est, *longifolia* reverè inceptum,

grandi-flora. 3. SERAPIAS foliis elliptico-lanceolatis bracteis germine longioribus, floribus erectis labio obtuso petalis vix æquali.

Serapias grandiflora. *With.* 42. *Lightf.* 528. *Relb.* 341. *Sibth.* 14. *Dicks. H. Sicc. fasc.* 6. 18. *Engl. Bot.* t. 271.

S. Lonchophyllum. *Suppl.* 405. *Ehrb. Phyt.* 57.

S. longifolia. *Huds.* 393.

S. lancifolia. *Hull.* 197.

Epipactis n. 1298. *Hall. Hist.* v. 2. 155. t. 41.

Helleborine flore albo. *Raii Syn.* 383. *Moris. sect.* 12. t. 11. f. 12.

H. latifolia, flore albo clauso. *Raii Syn.* 384.

Alijsq[ue] quorundam. *Cord. Hist.* 150. 2.

Angl.

Angl. White Helleborine.

In sylvis et dumetis, imprimis solo calcareo.

In Stoken-church woods, Oxfordshire. *Ray. Sibth.*

In the midland counties frequent. *Huds. Gloucestershire. Rev. Mr. Baker.* Isle of Arran.

Lightf.

Perennis. Fl. Junio.

Radix repens. Caulis pedalis, glaber. Folia amplexicaulia, elliptico-lanceolata. Flores a 3 ad 8, erecti, magni, albi. Bracteæ infimæ maximæ, foliis conformes; reliquæ lanceolatæ, angustiores, longitudine ferè florum. Petala subpatentia, demum clausa, nivea, immaculata. Nectarii labium porrectum, petalis parum brevius, cordatum, obtusum, integerrimum, lineis tribus, elevatis, flavis, longitudinalibus, supernè notatum. Germen gracile, sulcatum, subpubescens.

4. SERAPIAS foliis ensiformibus, bracteis minus *ensifolia.* tis germine brevioribus, floribus erectiusculis, labio petalis duplo breviore.

Serapias ensifolia. *Syst. Veg. ed. 14. 815,* synonymis confusis. *With. 42. Hull. 197.* *Engl. Bot. t. 494.*

S. longifolia γ. *Huds. 394.*

S. grandiflora. *Fl. Dan. t. 506.*

S. Xiphophyllum. *Suppl. 404. Ebrh. Phyt. 67.*

Epipactis n. 1298 β. *Hall. Hist. v. 2. 155.*

Helleborine foliis prælongis angustis acutis.

Raii Syn. 384. Herb. Buddle.

Angl. Narrow-leaved Helleborine.

In sylvis montosis, sed rariùs.

In a wood near Ingleton, Yorkshire. *Dr. Richardson.* On the top of Aberley hill, and in Wire forest, Worcestershire. *W.A. Moseley, Esq.*

Perennis. Fl. Junio.

Radix e fibris carnosis, vix repens. Caulis pedalis. Folia amplexi-

amplexicaulia, subdisticha, lineari-lanceolata, elongata. Flores erecti usculi, praecedenti similes, albi. Bractea infima longitudine floris; reliquæ minimæ, acutæ, germinibus quadruplo breviores. Petala acutiuscula. Nectarium basi subcalcaratum; labio petalis duplo breviore, obtusiusculo, sulcato, anticè maculâ flavâ. Germen contortum, serè lineare.

rubra. 5. SERAPIAS foliis lanceolatis, bracteis germine longioribus, floribus erectis, labio acuto lineis elevatis undulatis.

Serapias rubra. Syst. Veg. ed. 14. 816. With. 42. Hull. 197. n. 5. Engl. Bot. t. 437..

S. longifolia δ. Huds. 394.

Epipactis n. 1299. Hall. Hist. v. 2. 156. t. 42.

Helleborine. Fl. Dán. t. 345.

H. montana angustifolia purpurascens. Dill. in Razii Syn. 384.

H. angustifolia 6 Clusii. Ger. em. 442.

Damasonium purpureum dilutum, sive Elleborine 4 (6) Clusii. Bauh. Hist. v. 3. 516.

Angl. Purple Helleborine.

In nemorosis petrosis, montofisis, rariūs.

In Ireland. Phukenet. Gloucestershire. Rev. Mr. Baker. On a steep stony bank, sloping to the south, on Hampton Common in that county. Mrs. Smith.

Perennis. Fl. Junio.

Radix repens, carnosa. Caulis pedalis, pubescens. Folia amplexicaulia, lanceolata, acuta. Flores erecti, purpurei, elegantes. Bracteæ lanceolatae, angustæ, longitudine florum, vel parùm breviores. Petala sursum curva, acuminata, purpurea. Nectarium subcalcaratum; labio longitudine petalorum, dependente, lanceolato; basi auriculato, lobato, albido; apice acuto, purpureo; antieè lineis elevatis, undulatis, flavecentibus notato. Germen gracile.

II. HEX-

II. HEXANDRIA.

385. ARISTOLOCHIA. *G. Pl.* 467. *Juss.* 73.
Gærtn. t. 14.

Stigmata 6. *Cal.* nullus. *Cor.* monopetala, lin-gulata, integra. *Caps.* infera, sexlocularis.

1. ARISTOLOCHIA foliis cordatis, caule erecto, *Clematis*-
floribus axillaribus confertis. *titis.*

Aristolochia Clematitis. *Sp. Pl.* 1364. *Huds.*

394. *With.* 360. *Hull.* 198. *Relb.* 342.

Sibth. 112. *Dicks. Dr. Pl.* 14. *Woodv. Suppl.*

t. 238. *Engl. Bot. t.* 398. *Mill. Illustr. t.* 75.

A. Saracenica. *Ger. em.* 847.

A. rotunda. *Fuchs. Hist.* 90.

A. longa. *Trag. Hist.* 178.


Angl. Common Birthwort.

In sylvis, dumetis, et ruderatis.

In a wood 2 miles from Thorndon, Essex; also in Cambridgeshire. *Blackstone.* Near Maidstone, and in other parts of Kent. *Huds.* At Godstow nunnery. *Sibth.* At Sturston, Suffolk, near Diss. *Mr. Woodward.* Among the ruins of Carrow abbey, Norwich. *Rev. Mr. Sutton.* Near Kencot, Oxfordshire. *Rev. Dr. Goodenough.*

Perennis. Fl. Julio, Augusto.

Radix repens, gracilis. Herba glabra, lætè virens. Caules erecti, simplices, bipedales, subflexuosi, foliosi, teretes, striati, multiflori. Folia alterna, petiolata, exstipulata, cordata, integrerima, glabra, venosa, venis pedato-ramosis. Flores axillares, conferti, pedunculati, pedunculis simplicibus, ebraeatis. Corolla citrina, viridi lineata, basi globosa, medio constricta, apice hians. Gerumen inferum. Stylus brevissimus. Stigmata sex, obtusa, radiantia. Stamina brevissima, styllo inserita, crassa, extùs antherifera. Capsulam nondum vidi.



Classis XXI.

MONOECIA.

Flores masculi et fœminei in eâdem
plantâ.

MONANDRIA.

Stamen 1.

386. ZANNICHELLIA. Masc. *Cal.* nullus. *Cor.*
nulla.

Fœm. *Cal.* monophyllus. *Cor.* nulla. *Pist.*
quatuor. *Sem.* quatuor.

Callitrichæ. *Chara?*

DIANDRIA.

Stamina 2.

387. LEMNA. Masc. *Cal.* monophyllus. *Cor.*
nulla.

Fœm. *Cal.* monophyllus. *Cor.* nulla. *Stylus* 1.
Caps. polysperma.

TRIANDRIA.

Stamina 3.

391. ERIOCAULON. Masc. *Cal. communis polyphyllus :*
proprius di- vel tri- phyllus. *Corollulæ* bi-
 vel tri-partitæ.
Fl. Masc. in medio : *fæm.* in radio. *Stigmata*
 2 vel 3. *Caps.* bi- vel 3-cocca. *Sem.* fo-
 litaria.
389. SPARGANIUM. Masc. *Cal. triphyllus.*
Cor. nulla.
Fœm. *Cal.* triphyllus. *Cor.* nulla. *Drupa ex-*
sueca, monosperma.
390. CAREX. Masc. *Amentum imbricatum. Cal.*
gluma univalvis. *Cor.* nulla.
Fœm. *Ament.* imbricatum. *Cal.* gluma unival-
 vis. *Cor.* nulla. *Stigmata* 2 vel 3. *Semen*
 arillo ventricoso tectum.
388. TYPHA. Masc. *Ament.* cylindraceum, pi-
 losum. *Anth.* subternæ in filamento com-
 muni.
Fœm. *Ament.* cylindraceum. *Sem.* 1, pedi-
 cello papposo.

TETRANDRIA.

Stamina 4.

392. LITTORELLA. Masc. *Cal. tetraphyllus.*
Cor. quadrifida. *Stam.* longissima.
Fœm. *Cal.* nullus. *Cor.* inæqualiter trifida.
Stylus longissimus. *Nux* unilocularis.

395. URTICA. Masc. *Cal.* tetrphyllus. *Cor.* nulla. *Germinis* rudimentum cyathiforme.
 Fœm. *Cal.* diphyllo. *Cor.* nulla. *Sem.* 1, superum, nitidum.
394. BUXUS. Masc. *Cal.* triphyllus. *Pet.* 2.
Germinis rudimentum:
 Fœm. *Cal.* tetrphyllus. *Pet.* 3. *Styli* 3.
Caps. trirostris, trilocularis. *Sem.* 2.
393. BETULA. Masc. *Cal.* amenti squama monophylla, trifida, triflora. *Cor.* quadripartita.
 Fœm. *Cal.* amenti squama monophylla, subtrifida, biflora. *Styli* 2. *Semina* compressa.
Eriocaulon. *Myrica Gale.*

PENTANDRIA.

Stamina 5.

396. XANTHIUM. Masc. *Cal.* communis imbricatus. *Cor.* monopetalæ, infundibuliformes, quinquesfidæ. *Recept.* paleaceum.
 Fœm. *Cal.* diphyllo, biflorus. *Cor.* nulla.
Drupa sicca, muricata, bifida. *Nucleus* bilocularis.
397. AMARANTHUS. Masc. *Cal.* triphyllus.
Cor. nulla. *Stam.* 3 five 5.
 Fœm. *Cal.* triphyllus. *Cor.* nulla. *Styli* 3.
Caps. unilocularis, circumscissa. *Sem.* 1.

398. BRYONIA. Masc. *Cal.* quinquedentatus.
Cor. quinquefida. *Filam.* 3. *Anth.* 5.
 Fœm. *Cal.* quinquedentatus. *Cor.* quinquefida. *Styl.* trifidus. *Bacca* infera, polysperma.

Fagus sylvatica. *Quercus.* *Atriplex portulacoides.*

HEXANDRIA.

Stamina 6.

Rumices. *Quercus.*

POLYANDRIA.

Stamina 8 vel plura.

400. MYRIOPHYLLUM. Masc. *Cal.* tetraphyllus. *Pet.* 4. *Stam.* 8.
 Fœm. *Cal.* tetraphyllus. *Pet.* 4. *Stigmata* 4, sessilia. *Sem.* 4, corticata.
403. POTERIUM. Masc. *Cal.* tetraphyllus. *Cor.* quadripartita. *Stam.* 30—40.
 Fœm. *Cal.* tetraphyllus. *Cor.* quadripartita. *Pist.* 2. *Nux* bilocularis, corticata.
401. SAGITTARIA. Masc. *Cal.* triphyllus. *Pet.* 3. *Stam.* ferè 24.
 Fœm. *Cal.* triphyllus. *Pet.* 3. *Pist.* plurima. *Caps.* plurimæ, ventricosæ, monospermæ.

399. CERATOPHYLLUM. Masc. *Cal.* multipartitus. *Cor.* nulla. *Stam.* 16—20.
 Fœm. *Cal.* multipartitus. *Cor.* nulla. *Stigma* subsessile. *Sem.* 1, corticatum.
405. FAGUS. Masc. *Cal.* campanulatus, quinquefidus. *Cor.* nulla. *Stam.* 5—12.
 Fœm. *Cal.* quadrifidus. *Cor.* nulla. *Styli* 2 vel 3, trifidi. *Sem.* 2 vel 3, calyce coriaceo, muricato, tecta.
404. QUERCUS. Masc. *Cal.* campanulatus, subquinquefidus. *Cor.* nulla. *Stam.* 5—10.
 Fœm. *Cal.* campanulatus, integerrimus, scaber. *Cor.* nulla. *Stylus* 1. *Stigmata* 3. *Nux* supera, coriacea, monosperma.
407. CORYLUS. Masc. *Cal.* amenti squama trifida. *Cor.* nulla. *Stam.* 8.
 Fœm. *Cal.* bifidus, lacer. *Cor.* nulla. *Styli* 2. *Nux* ovata, lævis, unilocularis, calyce coriaceo tecta.
406. CARPINUS. Masc. *Cal.* amenti squama subrotunda. *Cor.* nulla. *Stam.* 8—20.
 Fœm. *Cal.* amenti squama oblonga. *Cor.* nulla. *Germina* 2. *Styli* singulis 2. *Nux* angulosa, unilocularis.
402. ARUM. *Spatha* monophylla. *Cor.* nulla. *Spadix* androgynus, medio staminiferus, basi germiniferus. *Baccæ* uniloculares.

MONADELPHIA.

Filamenta infernè in unum corpus
coalita.

408. PINUS. Masc. *Cal.* amenti squama peltata. *Cor.* nulla. *Antheræ* squamis adnatae, sessiles.

Fœm. *Cal.* amenti squama biflora. *Cor.* nulla.
Pist. 1. *Nux* alata.

Typba?

XXI. MONOECIA.

I. MONANDRIA.

386. ZANNICHELLIA. G. Pl. 476. Juss. 19.
Gærtn. t. 19.

Masc. *Cal.* nullus. *Cor.* nulla. *Stam.* Filamentum unicum, longum, erectum. *Anthera* ovata, erecta.

Fœm. *Cal.* monophyllus. *Cor.* nulla. *Germina* 4, vel plura. *Sem.* totidem, pedicellata. *Stigmata* peltata.

1. ZANNICHELLIA antherâ quadriloculari, stig-palustris. matibus integerrimis.

Zannichellia palustris. Sp. Pl. 1375. Huds. 397. With. 6. Hull. 202. Relb. 344. Sibth. 3. Mill. Illustr. t. 77. Fl. Dan. t. 67.

Z. palustris major, foliis gramineis acutis, flore cum apice quadricapsulari, embryonis clypeolis integris, et vasculo non barbato, capsulis seminum ad costam dentatis. Mich. Gen. 71. t. 34. f. 1.

Aponogeton aquaticum graminifolium, staminibus singularibus. Raii Syn. 135.

Potamogeto similis, graminifolia, ramosa, ad genicula polyceratos. Pluk. Phyt. t. 102. f. 7.

Angl. Horned Pondweed. Pet. H. Brit. t. 6. f. 2.

In fossis et stagnis.

Annua. Fl. Julio.

Radix fibrosa, tenuissima. Caulis filiformis, ramosissimus, foliosus, glaber. Folia subverticillata, linearia, integerrima, acuta, angustissima, graminea. Bractea axillaris, tubulosa, membranacea. Flores gemini, masculi

lus cum fœmineo, intrà bracteam. Stamen ad basin pedicelli floris fœminæ, capillare, longissimum. Anthera flava, quadrilocularis. Flos fœmineus pedicellatus, calyce campanulato, ore inæqualiter dentato. Germina 4 vel 5, pedicellata, oblonga, compresa, extùs plerumque dentata vel corrugata. Styli breves. Stigmata magna, ovata, peltata, margine integerrima aut vix repanda, minimè dentata. Semina germinibus conformia.

Alia species, ut videtur, *Mich. Gen. t. 34. f. 2*, nunquam, me sciente, in Angliâ detecta est. Vulgarem certè descripsit Cel. Woodwardius apud Witheringium.

II. DIANDRIA.

387. LEMNA. *G. Pl. 478. Wiggers. Holsat. 66.*
Lenticula. *Juss. 19.*

Masc. *Cal. monophyllus. Cor. nulla.*

Fœm. *Cal. inonophyllus. Cor. nulla. Stylus 1.*
Caps. polysperma.

trisulca. 1. LEMNA foliis petiolatis lanceolatis proliferis.
Lemna trisulca. Sp. Pl. 1376. Huds. 399. With.
43. Hull. 202. Relb. 346. Sibth. 14. Ab-
bot. 198. Engl. Bot. t. 926.

Lenticula aquatica trisulca. Raii Syn. 129.
Baub. Hist. v. 3. 777. 2.

Lenticularia ramosa monorrhiza, foliis oblongis,
pediculis longioribus donatis. Mich. Gen. 16.
t. 11. f. 5.

Hederula aquatica. Ger. em. 830.
Angl. Ivy-leaved Duck-weed.

In fossis et stagnis.

Annua. Fl. Junio. D. Turner.

Herba natans, pellucida, glabra, pallidè virens. Radices
filiformes,

filiformes, solitariæ, simplices, ad extremitates (ut in omnibus speciebus, monente Michelio) vaginatæ. Folia petiolata, lanceolata, integerrima, subrepanda, subtus prolifera. Flores ex rimâ marginali foliorum, solitarii, hermaphroditici et masculini. Calyx integer. Stamina adscendentia, alba. Antheræ didymæ, flavæ. Germen ovatum. Stigma bifidum.

2. LEMNA foliis sessilibus utrinque planiusculis, *minor.*
radicibus solitariis.

Lemna minor. Sp. Pl. 1376. Huds. 399. With. 44. Hull. 202. Relb. 346. Sibth. 14. Abbot. 198. Engl. Bot. t. 1095.

Lenticula palustris vulgaris. Vaill. Par. t. 20.
f. 3.

Lenticularia minor, monorrhiza, foliis subrotundis utrinque viridibus. Mich. Gen. 16. t. 11. f. 3.

Lens palustris. Raii Syn. 129. t. 4. f. 1. Camer. Epit. 852. Valisn. Oper. v. 2. 88. t. 14, 15.
Angl. Lesser Duck-weed.

In fossis et stagnis vulgatissima.

Annua. Fl. Junio, Julio. D. Borrer.

Folia natantia, 2 vel 3 basi connexa, obovata, parva, vix duarum linearum longitudine; suprà plana, lètè viridia; subtus convexiuscula, pallida. Radices longissimæ, solitariæ, centrales. Flores ex foliorum margine, hermaphroditici. Calyx bilobus. Stamina et pistillum ut in priore.

3. LEMNA foliis sessilibus convexiusculis subtus *gibba.*
hemisphæricis, radicibus solitariis.

Lemna gibba. Sp. Pl. 1377. With. 44. Hull. 202. Relb. 346. Sibth. 15. Abbot. 199. Engl. Bot. t. 1233.

L. minor β. Huds. 399.

Lenticula

Lenticula palustris major, infernè magis convexa, fructu polyspermo. *Mich. Gen.* 15. *t. 11. f. 1.*

Angl. Gibbous Duck-weed.

In fossis et stagnis minus frequens.

Annua. Fl. Junio, Julio. D. Borrer.

Præcedente parùm major, foliis suprà purpurascens, subtus valdè gibbis et spongiosis, sueco plenis. Flores hermaphroditici. Calyx nullus, nisi margo, citò evanescens, rimæ. Stamina ut in præcedentibus. Stigma obtusum. Huic semina a 4 ad 6 tribuit Michelius, cuius icon stirpem nostram sat bene refert. Wiggers, qui forsitan secundam speciem Michelii invenit, fructum dispermum descripsit.

polyrrhiza.

4. *LEMNA* foliis sessilibus, radicibus confertis.

Lemna polyrrhiza. Sp. Pl. 1377. Huds. 399. Witb. 44. Hull. 202. Relb. 347. Sibth. 15. Abbot. 199.

Lenticularia major polyrrhiza, infernè atro-purpurea. *Mich. Gen.* 16. *t. 11. f. 1.*

Lenticula palustris major. *Dill. in Raii Syn.* 129. *t. 4. f. 2.* *Vaill. Par. t. 20. f. 2.*

Angl. Greater Duck-weed.

In fossis.

Annua. Fl. Julio, Augusto.

Præcedentibus duplò vel triplò major. Folia rotundiora, obtusiora; subtus convexa, atro-purpurea. Radices plurimæ, centrales, confertæ.

“ Flores inter duplicaturam laminæ utriusque, eo latere ubi unum folium altero insinuat. Fructus convenit cum priori, sed major est, et magis compressus.” *Graverus apud Wiggers.*

Genus forsitan ad Diandriam Monogyniam removendum.

III. TRIANDRIA.

388. TYPHA. *G. Pl. 479. Juss. 25.
Gærtn. t. 2.*

Masc. *Amentum cylindraceum, pilosum. Anthereæ subternæ, in filamento communi.*

Fœm. *Ament. cylindraceum. Sem. unicum, pedicello papposo.*

1. TYPHA foliis subensiformibus, spicâ masculâ *latifolia*. fœmineâque approximatis.

Typha latifolia. Sp. Pl. 1377. Huds. 400. With. 111. Hull. 203. Relb. 347. Sibth. 25. Abbot. 199. Leers. 207.

T. major. Curt. Lond. fasc. 3. t. 61.

Typha. Raii Syn. 436. Fuchs. Hist. 823. Ger. em. 46. Matth. Valgr. v. 2. 216.

*T. palustris major. Bauh. Hist. v. 2. 539.
Angl. Great Cat's-tail or Reed-mace.*

In stagnis, fossis, vel rivulis lentè fluentibus.

Perennis. Fl. Julio.

Radix repens. Herba glabra. Culmus ferè sexpedalis, erectus, simplicissimus, enodis, teres, basi foliosus. Folia erecta, culmo ferè æqualia, linearia, acutiuscula, subtùs convexiuscula. Amentum terminale, erectum, multiflorum, continuum, supernè masculinum, subfoliolosum, flavum; infernè fœmineum, fuscum. Antherae 1 ad 4 in filamento communi.

2. TYPHA foliis semicylindraceo-planiusculis cul- *angusti-*
mum æquantibus, spicâ masculâ fœmineâque folia,
remotis.

*Typha angustifolia. Sp. Pl. 1377. Huds. 400.
With.*

Witb. 111. *Hull.* 203. *Relb.* 348. *Abbot.*
199. *Fl. Dan.* t. 815.

T. minor. *Curt. Lond. fasc.* 3. t. 62.
T. palustris media. *Raii Syn.* 436. *Baub. Hist.*
v. 2. 540. *Moris. sect.* 8. t. 13. f. 2.
Angl. Lesser Cat's-tail or Reed-mace.

In fossis et stagnis rariūs.

In the middle of Woolwich common. *Curt.* In
clay-pits near Bungay. *Mr. Woodward.* Be-
tween Norwich and Hingham.

Perennis. Fl. Junio, Julio.

Priore duplō minor et grāciliōr. Folia infernē semicy-
lindracea. Spica interrupta, amento masculino a fœ-
mineo parūm distante.

minor. 3. **TYPHA** foliis cnsiformibus culmo duplō bre-
vioribus, spicā interruptā abbreviatā incrassatā,
culmo gracillimo.

Typha minor. *Baub. Hist.* v. 2. 540. *Lob. Ic.*
81. *Tabern. Kreuterb.* 559.

T. angustifolia B. *Sp. Pl.* 1378. *Huds.* 400.

T. palustris minor. *Dill. in Raii Syn.* 436.

T. palustris minima, dupli clavā. *Moris. sect.* 8.
t. 13. f. 3.

Angl. Dwarf Cat's-tail or Reed-macc.

In paluſtribus.

Found by Mr. Dandridge on Hounslow-heath.
Dill.

Perennis. Fl. Julio.

Culmus ſequipedalis, triplō quam in T. angustifoliā gra-
ciliōr. Folia plana, latiuscula, vix longitudine dimidii
culmi. Spica triplō erassior quām in T. angustifoliā,
abbreviata, foliolosa, interrupta, amento ſuperiore vel
omniō masculino, ut in praecedentibus, vel infernē
femineo.

fœmineo. Hanc speciem, certè distinctissimam, ex Dillenii fide hie admisi. Ex agro Genevensi habui.

389. SPARGANIUM. *G. Pl.* 480. *Juss.* 26.
Gærtm. t. 19.

Masc. *Cal.* triphyllus. *Cor.* nulla.

Fœm. *Cal.* triphyllus. *Cor.* nulla. *Drupa* ex-succa, monosperma.

1. SPARGANIUM foliis basi triquetris lateribus ramo-concavis, pedunculo communi ramoso, stig-summate linearari.

Sparganium ramosum. *Huds.* 401. *With.* 112.

Hull. 203. *Sibth.* 25. *Curt. Lond. fasc.* 5.

t. 66. *Raii Syn.* 437. *Bauh. Theatr.* 228.

Ger. em. 45. *Engl. Bot. t.* 744.

S. erectum. *Sp. Pl.* 1378. *Reh.* 348. *Abbot.* 200.

S. quibusdam. *Bauh. Hist.* v. 2. 541.

Angl. Branched Bur-reed.

In fossis, et ad ripas fluviorum, vulgaris.

Perennis. Fl. Julio, Augusto.

Radix repens. Caulis eretus, subtripedalis, teres, foliosus, glaber, apice paniculato-ramosus. Folia radicalia basi triquetra, lateribus concavis, superne ensiformia, ereta, glabra; eaulina amplexicaulia, erecto-patentia, abbreviata, plana. Pedunculi partiales alterni, racemosi, multiflori, basi flexuosi. Florum capitula alterna, sessilia, multiflora; inferiora fœminea; superiora masculina, approximata, magisque numeroosa. Calycis foliola spatulata, fusca. Stamina calyce longiora. Antheræ ochroleueæ, oblongæ, bifidæ. Germin biloculare, obovatum. Stylus brevis. Stigma obliquum, elongatum, lineare, hinc pubescens, plerumque solitarium, quandoque geminum. Drupa saepius monospernia, interdum disperma.

2. SPAR-

simplex. 2. SPARGANIUM foliis basi triquetris lateribus planis, pedunculo communi simplici, stigmate linearie.

Sparganium simplex. *Huds.* 401. *With.* 112.
Hull. 203. *Sibth.* 25. *Abbot.* 200. *Curt. Lond.*
fasc. 5. *t.* 67. *Dicks. H. Sicc. fasc.* 6. 7. *Engl.*
Bot. t. 745.

S. erectum B. *Sp. Pl.* 1378.

S. n. 345*. *Fl. Lapp.* *ed.* 1. 271. *ed.* 2. 280.
Herb. Linn.

S. non ramosum. *Raii Syn.* 437. *Baub. Theatr.*
231.

S. latifolium. *Ger. em.* 45.

S. alterum. *Baub. Hist.* v. 2. 541. *Lob. Ic.* 80.
Angl. Unbranched Upright Bur-reed.

In stagnis, imprimis solo glareoso.

Perennis. Fl. Julio, Augusto.

Præcedente minor. Folia radicalia basi triquetra, lateribus, optimè monente Curtisio, planis. Caulis simplissimus, nec paniculatus, pedunculo communi simplici. Capitulum infimum saepius pedicellatum, reliqua sessilia. Calyx virens, nec fuscescens. Stigma elongatum, lineare, plerumque simplex.

natans. 3. SPARGANIUM foliis decumbentibus planis, pedunculo communi simplici, stigmate ovato brevissimo, capitulo masculino subsolitario.

Sparganium natans. *Sp. Pl.* 1378. *With.* 112.
Hull. 203. *Lightf.* 540. *Engl. Bot. t.* 273.
Dicks. H. Sicc. fasc. 6. 8. *Fl. Dan. t.* 260.

S. simplex B. *Huds.* 401.

S. minimum. *Raii Syn.* 437.

Angl. Floating Bur-reed.

In rivulis et stagnis, solo lutoso.

Burwell fens, Cambridgeshire. *Rev. Mr. Hemsted.*

In the Norwich river. *Mr. Rose.* Scotland.

Dickson. Near Manchester. *Dr. Hull.*

Perenniz.

Perennis. Fl. Julio.

Herba natans, foliis planis, duplò minoribus quam in S. simplici. Racemus solitarius, simplex, pauciflorus. Capitula foeminea duo vel tria, infimum pedicellatum; masculinum plerumque solitarium, terminale. Calyx albido-virens. Stigma brevissimum, ovatum, peltatum, simplex. Drupa monosperma.

390. CAREX. G. Pl. 482. Juss. 26.
Gærtn. t. 2.

Masc. *Amentum imbricatum.* Cal. gluma univalvis. Cor. nulla.

Fœm. *Ament. imbricatum.* Cal. gluma univalvis. Cor. nulla. Stigmata 2 vel 3. Semen arillo ventricoso tectum.

* *Spicâ unicâ, simplici.*

1. CAREX spicâ simplici dioicâ, fructibus ovatis dioica. nervosis adscendentibus margine serrulatis.

Carex dioica. Sp. Pl. 1379. Gooden. Tr. of Linn. Soc. v. 2. 139. Hudf. 401. With. 86. Hull. 204. Relh. 349. Sibth. 26. Abbot. 201. Lightf. 541. Engl. Bot. t. 543. Fl. Dan. t. 369. Ehrb. Phyt. n. 6.

Masc. Gramen cyperoides, spicâ simplici cassâ. Raii Syn. ed. 2. 267.

G. cyper. minimum, spicâ cassâ simplici. Moris. sect. 8. t. 12. f. 22.

Fœm. G. cyper. minimum, Ranunculi capitulo rotundo. Raii Syn. 425. Moris. sect. 8. t. 12. f. 36.

Cyperoides parvum, caulis et foliis tenuissimis, triangularibus, spicâ subrotundâ, capsulis oblongis in angustum collum, vix bifidum, attenuatis. Mich. Gen. 56. t. 32. f. 2.

β. Gramen

B. Gramen cyperoides minus, Ranunculi capitulo longiore. *Raii Syn.* 425.

Angl. Common Separate-headed Carex.

In palustribus spongiosis.

Perennis. Fl. Maio, Junio.

Radix repens. Herba glabra, vix spithamæa. Culmi simplicissimi, triquetri, læviusculi, monostachyi, basi foliosi. Folia setacea, triquetra, culmo breviora, basi vaginantia. Spicæ dioicæ, lineares, erectæ, multifloræ, vix unciales, glumis dorso viridibus, margine scarriosis, spicæ fœmineæ latioribus. Stamina capillaria, exserta, antheris flavis, linearibus. Fructus ovati, extus gibbi atque nervosi, margine apicem versùs serrulati, ore scarioso, indiviso, demùm erecto-patentes. Stigmatea duo, linearia, pubescentia. Semina lenticularia, glabra, luteola, mucronulata.

Variat rariū spicâ androgynâ.

C. capitata, *Fl. Dan.* t. 372, discrepat spicis semper androgynis, floribus utriusque sexûs paucioribus, glumis vix dorso virentibus, arillo enervi, margine integerrimo. Hæc è Florâ Britannicâ exulat.

Daval-liana. 2. CAREX spicâ simplici dioicâ, fructibus lanceolato-triquetris nervosis patenti-deflexis : angulis apice scabris.

Carex Davalliana. *Tr. of Linn. Soc.* v. 5. 266.

C. dioica. *Wilden. Car. Berol.* 16, ex descr.

C. n. 1350. *Hall. Hist.* v. 2. 182. synonymis confusis.

Gramen cyperoides, spicâ simplici cassâ. *Scheuchz.*

Agr. 497. t. 11. f. 9, 10; synonymis omnino erroreis. Confer *Gooden. Tr. of Linn. Soc.* v. 2, ad calcem.

Angl. Prickly Separate-headed Carex.

In uliginosis.

Discovered in marshy ground in Mearns-shire,

North Britain, by Professor James Beattie jun. of Aberdeen.

Perennis. Fl. Maio, Junio.

Radix fibrosa, cæspitosa. Culmi simplicissimi, triquetri, scabriuscui, quandoque scaberrimi, monostachyi, basâ foliosi, parùm altiores quam in præcedente. Folia setacea, triqueta, culmo triplò breviora. Spicæ dioicæ, lineares, erectæ, multifloræ, magnitudine prioris, glumis fuscis, carinatis, acutis, margine scariosis. Stamina ut in C. dioicâ. Fructus lanceolato-triquetra, nec gibbi, rostrati, undique nervosi, angulis apice scabris, ore scariso, integro, demùm deflexo-patentes, acuminati, unde spicæ seminiferæ squarrosæ fiunt.

3. CAREX spicâ simplici androgynâ : supernè pulicaris : masculâ, fructibus divaricatis retroflexis utrinque acuminatis, stigmatibus binis.

Carex pulicaris. Sp. Pl. 1380. Gooden. Tr. of Linn. Soc. v. 2. 142. Huds. 402. With. 86. Hull. 204. Relb. 349. Sibth. 26. Abbot. 203. Lightf. 543. Leers. 194. t. 14. f. 1. Fl. Dan. t. 166.

C. psyllophora. Suppl. 413. Ehrb. Phyt. n. 7.

C. minima, caulis et foliis capillaccis, capitulo singulari, tenuiori, capsulis oblongis utrinque acuminatis et deorsum reflexis. Mich. Gen. 66. t. 33. f. 1.

Gramen cypcroides minimum, seminibus deorsum reflexis puliciformibus. Raii Syn. 424.

G. cyper. pulicare. Moris. sect. 8. t. 12. f. 21. Angl. Flea Carex.

In paludosis spongiosis et limosis.

Perennis. Fl. Junio.

Radix fibrosa, fusca. Culmi plerumque ferè pedales, filiformes, glabri. Folia setacea, supernè canaliculata, lævia, culmo subæqualia. Spica simplex, linearis, erecta.

Flores masculi plurimi, superiores, glumis fusca demum adpressis; foemini circiter totidem, laxè imbricati, demum refraeti, glumis decidui. Fructus elliptico-lanceolati, basi apiceque attenuati, utrinque carinati, fusi, nitidi, enerves. Stigmata duo. Semina elliptica, fusca.

Icon in *Fl. Dan.* ad plantam floribus immaturis delineata est.

pauciflora. 4. CAREX spicâ simplici androgynâ pauciflorâ : supernè masculâ, fructibus patentibus subulatis, stigmatibus ternis.

Carex pauciflora. Lightf. 543. t. 6. f. 2. Gooden. Tr. of Linn. Soc. v. 2. 143. With. 87. Hull. 204. Dicks. H. Sicc. fasc. 1. 16.

C. patula. Hudsf. 402 et 657.

C. Leucoglochin. Suppl. 413. Ehrb. Phyt. n. 8. Angl. Few-flowered Carex.

In paludibus alpinis rariū.

About half way up the mountain of Goat-field in the Isle of Arran, in a peat bog. Lightf. Near Glasgow. Hudsf. Wet moors in the Highlands. Dickson.

Perennis. Fl. Junio.

Radix repens. Culmi 3—5-unciales, adscendentis, glabri, subtriquetri. Folia plerumque bina, canaliculata, laevia, culmo duplo breviora. Spica laxa, floribus 4 ad 6, quorum terminalis masculus, erectus, inferiores foemini, patuli, demum refraeti. Fructus subulati, lutei, subtriquetri, sulcati. Stigmata tria.

Variat rariū stigmatibus binis. Lightf.

** *Spicâ compositâ, androgynâ*.

stellulata. 5. CAREX spiculis subternis remotis, fructibus divergentibus acuminatis ore integris.

Carex stellulata. Gooden. Tr. of Linn. Soc. v. 2. 144. With. 87. Hull. 204.

C. muri-

C. muricata. *Huds.* 406. *Lightf.* 549. *Relb.* 350.
Abbot. 202. *Fl. Dan.* t. 284. *Leers.* 196.
t. 14. f. 8.

C. echinata. *Sibth.* 28.

Gramañ cyperoides spicatum minimum, spicâ di-
vulsâ, aculeatâ. *Raii Syn.* 424. *Schenckz.*
Agr. 485. *t. 11. f. 3.*

G. cyper. nemorosum, spicis parvis asperis. *Mo-*
rif. sect. 8. *t. 12. f. 26.*

Angl. Little Prickly Carex.

In palustribus.

Perennis. Fl. Maio, Junio.

Radix fibrosa. Culmus 6—12-uncialis, triqueter, angulis
asperiusculis, basi foliosus. Folia plana, carinata, culmo
breviora, erectiuscula, margine carinâque apiceni versus
scabriuscula. Spica composita. Spiculæ tres vel qua-
tuor, androgynæ, alternæ, subrotundæ, omnes æquali-
tè remotiusculæ; infima sæpiùs bracteata. Flores in-
feriores masculi, pauciores; superiores foeminei, plures.
Glumæ ovatæ, fuscaæ; margine scariosæ, albæ; carinâ
latâ, virente. Fructus undique divaricati, ovati, ner-
vosi, mucronati, apice acuto, vix emarginati, nec bi-
fidi, utrinque scabriuseuli. Stigmata duo.

6. CAREX spiculis subfenis ellipticis remotiusculis *curta*.
nudis, glumis ovatis acutiusculis membra-
naceis, arillo integro.

Carex curta. *Gooden. Tr. of Linn. Soc.* v. 2.
145. *Wittb.* 88. *Hull.* 204. *Wilden. Car.*
Berol. 19. *t. 2. f. 3.*

C. brizoides. *Huds.* 406.

C. canescens. *Lightf.* 550. *Fl. Dan.* t. 285.

C. elongata. *Leers.* 197. *t. 14. f. 7.*

Gramañ cyperoides palustre elegans, spicâ com-
positâ asperiore. *Raii Syn.* 423.

G. cyper. elegans, spicâ compositâ molli. *Dill.*

in Raii Syn. 423. *Plak. Phyt.* t. 34. f. 4.
Rel. Rudb. 2. f. 34.

G. cyper. spicis curtis divulsis. *Loef. Pruss.* 117.
t. 32.

Angl. White Carex.

In aquosis rariūs.

In a pool at Middleton, Warwickshire, towards Cole's hill; also near Wrexham, Denbighshire, and in other places. *Ray.* In bogs in Isla, about 2 miles from the Sound; also in ditches by Loch Leven. *Herb. Jightf.* At Virginia water, Windsor forest. *Rev. Dr. Goodenough.*

Perennis. Pl. Junio.

Radix subrepens. Culmus pedalis, triqueter, angulis glabriusculis. Folia linearia, plana, angusta, glaucescentia, margine carinâque aspera, culmo parum breviora. Spiculae 4 ad 6, vel plures, alternæ, rariūs oppositæ, ellipticæ, obtusæ, multifloræ, argenteo-albidæ; infima subindè bracteata. Flores inferiores masculi, pauci; feminei plurimi. Glumæ ovatæ, acutiusculæ, nitidæ, membranaceæ, albæ, nervo tenuissimo, viridi. Fructus ovati, compressi, acuti, glabri, apice indiviso. Stigmata duo. Semen exactè ellipticum, stylo mucronatum.

ovalis. 7. CAREX spiculis subfusis ovalibus approximatis alternis, glumis lanceolatis arillum æquantibus.

Carex ovalis. *Gooden. Tr. of Linn. Soc.* v. 2.
148. *Huth.* 88. *Hull.* 205. *Engl. Bot.*
t. 306.

C. leporina. *Huds.* 404. *Relb. Suppl.* 1. 15.
Sibth. 27. *Leers.* 195. t. 14. f. 6. *Ehrb.*
Phyt. n. 38.

Gramen cyperoides spicâ e pluribus spicis brevi-
bus

bus mollibus composita. *Raii Syn.* 422.

Scheuchz. Agr. 456. t. 10. f. 15.

G. cyper. palustre majus, spicâ divisâ. *Moris.*

sect. 8. t. 12. f. 29.

Angl. Oval-spiked Carex.

In paluatribus et pratis humidis.

Percennis. Fl. Junio.

Radix repens. Culmus erexitus, pedalis vel altior, acutè triquetus, angulis scabris, basi foliosus. Folia saturatè viridia, plana, carinata, culmo subæqualia, basi vaginantia, alterna, margine carinâque scabriuscula. Spiculæ 4 ad 6, ellipticæ, androgynæ, e viridi rubiginosæ. Bractea foliacea ad basin spiculæ insimilæ. Flores inferiores masculi, pauci; fœminei numerosi. Glumæ ovato-lanceolatæ, rubiginosæ, margine scarioso, carinâ viridi. Fructus lanceolati, acuminati, hinc convexiusculi, ore subintegri, margine scabri. Stigmata duo. Semen elliptico-rotundatum.

Distinctissima, optimè monente D. Goodenough, a C. leporinâ Linnæi, quæ summorum Alpium incola est, et glumas habet obtusiores et breviores, corollamque margine lœvem, et spiculas ferè globosas.

8. CAREX spiculis solitariis remotis subsessilibus, *remota.*

· bracteis longissimis culmam superantibus,
arillo subintegro.

Carex remota. *Sp. Pl.* 1383. *Gooden. Tr. of Linn. Soc.* v. 3. 150. *Huds.* 407. *With.* 88. *Hull.* 205. *Relb.* 351. *Sibth.* 28. *Abbot.* 203. *Fl. Dan.* t. 370. *Leers.* 197. t. 15. f. 1. *Rel. Rudb.* 1. f. 27.

C. axillaris. *Sp. Pl.* 1382. *Linn. Mss. Sp. Pl.*

C. angustifolia, caule triquetro, capitulis pulchellis. *Mich. Gen.* 70. n. 2, 3. t. 33. f. 15, 16.

Gramen cyperoides angustifolium, spicis parvis sessilibus in foliorum aliis. *Raii Syn.* 424.

Pluk. Phyt. t. 34. f. 3.

G. cyper. paniculis pluribus sessilibus in foliorum alis. *Morif. se&t. 8. t. 12. f. 17.*

Angl. Remote Carex.

In nemoribus humidis, et aggeribus fossarum uidis umbrofis.

Perennis. Fl. Maio, Junio.

Radix fibrosa. Culmus pedalis, gracilis, debilis, triqueter, superne asper, basi glaber et terciusculus. Folia angusta, pallide viridia, margine scabra. Spiculae alternæ, solitariæ, circiter 6 vel 8, ovatæ, albidæ, androgynæ, quarum tres vel quatuor inferiores remotissimæ, bracteis foliaceis, culnum æquantibus vel superantibus. Flores masculi inferiores. Glumæ latè ovatæ, scariosæ, albæ, carinâ viridi. Fructus ovati, acuminati, glumâ longiores, marginibus lævibus, vel supernè tantum scarificulæ, apice plerumque integro. Stigmata duo. Semen lenticulare.

axillaris. 9. CAREX spiculis subternatis remotis sessilibus, bracteis elongatis, arillo apice bifido.

Carex axillaris. *Gooden. Tr. of Linn. Soc. v. 2. 151. t. 19. f. 1.* *With. 89.* *Hull. 205.* nec Linn.

Angl. Axillary Clustered Carex.

In palustribus rariùs.

Found near Putney by Mr. Curtis. *Rev. Dr. Goodenough.*

Perennis. Fl. Maio, Junio.

Radix fibrosa. Culmus erectus, firmus, striatus, 1 ad 3-pedalis, triqueter, angulis undique asperimis. Folia culmo breviora, marginibus, carinæque apice, aspera. Spiculae inferiores aggregatæ, subternæ, quandoque quaternæ vel quinæ, superiores remotæ vel remotissimæ, solitariæ. Bractea insima culnum superans; cæteræ breviores. Flores inferiores masculi. Glumæ ovatæ, acutæ, scariosæ, fuscescentes, carinâ virenti. Fructus ovati, acuminati, glumâ parùm longiores, marginibus

marginibus apicem versùs scabris, ore bifido. Stigmata duo. *Goodenough.*

10. CAREX spiculis congestis capitatis infernè *incurva*, fœmincis, bracteis scariosis, culmo teretiusculo glabro, foliis canaliculatis.

Carex incurva. Lightf. 544. t. 24. f. 1. Gooden. *Tr. of Linn. Soc. v. 2. 152.* Huds. 650. With. 89. Hull. 205.

C. *Fl. Dan.* t. 432.

C. *juncifolia.* Allion. *Ped. v. 2. 264.* t. 92. f. 4. Sm. *Tour on the Contin.* v. 3. 136.

Angl. Curved Carex.

In æstuariis, vel fluminum alpinorum ostiis, solo arenoso, rariùs.

In deep loose sea sand at the mouth of the water of Naver, and near Skelherry in Dunrossness in Shetland. Dr. Hope. Lightf.

Perennis. Fl. Julio, Augusto.

Radix repens. Culmi adscendentes, palmares, nudi, teretiusculi, læves, in maritimis arcuati, in irriguis alpinis fæpiùs recti. Folia radicalia, culmo breviora, linearia, canaliculata, lævia, nervosa. Spica composita, capitata, subconica, brevis. Spiculæ congestæ, subrotundæ, paucifloræ, fuscescentes. Bracteæ spiculis breviores, concavæ, scariosæ, ellipticæ, obtusæ, obsoletè carinatæ. Flores inferiores fœminei, superiores (parùm numerosiores) masculi. Glumæ ovatæ, scariosæ, fuscæ, margine albidae, carinâ tenuissimâ. Fructus ovati, acuti, virentes, apice fuso, indiviso, marginibus lævibus. Stamina capillaria, longissima, alba. Stigmata duo, fuscæ. Semen lenticulare.

Ne cum hâc C. fœtidam Allionii confundas, speciem distinctissimam.

11. CAREX spiculis congestis spicatis submonoi- *arenaria* cis,

cis, bracteis scariosis : inferioribus foliaceis,
culmo triquetro, foliis planis.

Carex arenaria. *Sp. Pl.* 1381. *Fl. Suec.* ed. 2.
325. *Gooden. Tr. of Linn. Soc.* v. 2. 153.
Huds. 405. *With.* 90. t. 20. *Hull.* 205.
Lightf. 545. *Dicks. H. Sicc. fasc.* 5. 14.
Ehrb. Phyt. n. 17.

C. maritima humilis, radice repente, caule trilatero, spicâ spadiceâ nonnihil foliosâ, capitulis crassioribus, capsulis marginatis. *Mich. Gen.* 67. t. 33. f. 4.

Gramini cyperoidi ex monte Ballon simile humilius, in maritimis et arenosis naescens. *Raii Syn.* 423. *Pluk. Phyt.* t. 34. f. 8.

Gramen cyperoides minus, repens, spicâ divisâ. *Loef. Pruss.* t. 31.

Angl. Sea Carex.

In arenosis maritimis copiosè.

Pereennis. Fl. Junio.

Radix valde repens, longè latèque extensa, squamosa, fibris tomentosis. Culni pedales, crebro-incurvi, nudí, triquetri, angulis acutis, apicè præcipue scabris. Folia basin culmi vaginantia, erecta, acuminata, plana, margine scabra. Spica ovata, composita, compressa. Spiculæ alternæ, approximatæ, ovato-subrotundæ, fuscae, multifloræ. Bracteæ lanceolatæ, carinatæ, margine scariosæ, inferiores in acumen foliaceum productæ. Flores inferiorcs fœmininei, in spiculis inferioribus numerosissimi ; superiores masculi, in spiculis superioribus longè magis numerosi. Glumæ ovato-lanceolatæ, fuscae, vix scariosæ, longius acuminatae. Fructus ovati, acuti, nervosi, apicem versus utrinque alati, margine que scabri, ore bifido. Stigmata duo. Semen lenticulare, vel obsoletè triquetrum.

interme- 12. CAREX spiculis congestis spicatis : infimis dia. termi-

terminalique fœmineis : intermediis masculis,
culmo triquetro erecto.

Carcx intermedia. *Gooden. Tr. of Linn. Soc. v. 2.*
154. *With. 91. Hull. 205. Dicks. H. Sicc.*
fasc. 7. 15.

C. disticha. *Huds. 403. Relb. 350. Sibb. 26.*
Abbot. 202.

C. arenaria. *Leers. 195. t. 14. f. 2.*

Gramini cyperoidi ex monte Ballon simile, spicâ
totali e pluribus spicis compositâ. *Raii Syn.*
423. *Phuk. Phyt. t. 34. f. 7. malè.*

G. cyperoides, spicâ subfuscâ molli, minor. *Mo-*
rif. seq. 8. t. 12. f. 32.

Angl. Soft brown Carex.

In palustribus et aquosis.

Perennis. Fl. Maio, Junio.

Radix repens, profundè descendens. Culmi erecti, stricti,
pedales et ultra, nudi, acutè triquetri, angulis scabri.
Folia basin culmi vaginantia, erecta, vix culmo æqua-
lia, plana, margine carinâque asperiuscula. Spica ob-
longa, obtusa, composita, fusco-ferruginea, mollis,
parùm compressa, nequaquam disticha. Spiculæ ap-
proximatæ, numerosæ, sursùm imbricatæ, ovatæ, qua-
rum duæ vel tres infimæ, cum terminali, omnino
ferè fœmineæ, cæteræ intermediæ submasculæ, “in
fœmineis scilicet sæpius flos unus aut alter masculus, et
in masculis fœmineus conspiciendus est.” *D. Good-*
enough. Flores superiores semper masculi, inferiores
fœminei. Bracteæ ovato-lanceolatæ, scariosæ, carina-
tæ, fuscæ, margine albo ; inferiores in mucronem foliaceum setaceum definentes. Glumæ ovatæ, fuscæ,
nervo pallido, margine scariofo, albo. Fruktus ovati,
acuti, utrinque alati, margine scabri, ore bisido. Stig-
mata duo ; interdùm tria inveni.

13. CAREX spicâ subdecompositâ spiculis omni- *divisa*
bus androgynis, bracteâ foliaceaâ erectâ, fruc-
tibus adpressis, radice repente.

Carex

Carex divisa. *Huds.* 405. *Gooden. Tr. of Linn. Soc.* v. 2. 157. t. 19. f. 2. *With.* 92. *Hull.* 205.

Gramen cyperoides ex monte Ballon, spicâ divulsa. *Raii Syn.* 423.

Angl. Bracteated Marsh Carex.

In palustribus, imprimis maritimis.

Near the Hithe at Colchester, and elsewhere; Mr. Newton. *Ray.* By Hithe in Kent; Mr. J. Sheppard. *Dill.* Near Acle bridge, Norfolk. Mr. C. Bryant. At Cley. Mr. Woodward.

Perennis. Fl. Maio, Junio.

Radix valde repens, crassa, tortuosa. Culmi erecti, debiles, pedales et ultra, nudi, triquetri, angulis supernè scabriusculis. Folia angusta, erecta, lètè viridia, margine carinâque scabra, longitudine varia. Spica uncialis, ovata, composita, sèpè basi decomposita, nigroferruginea, densa, compressiuscula. Spiculæ plures, irregulariter congestæ, ovatæ, acutæ, omnes androgynæ, floribus fœmineis inferioribus, numerosioribus, atque preeocioribus. Bractea ad spicæ basin foliaeea, erecta, acutè triquetra, plerunque longissima, spicam longè superans, interdùm spicâ brevior. Glumæ ovato-ellipticæ, mucronatæ, fuscæ, margine parùm scariosæ, carinâ pallidâ. Fructus glumis breviores, ovati, subalati, margine scabri, ore acuminato, bifido. Stigmata duo, longissima.

muricata. 14. CAREX spicâ oblongâ subdecompositâ squarrosâ, fructibus divergentibus acuminatis bifidis, radice fibrosâ.

Carex muricata. *Sp. Pl.* 1382. *Gooden. Tr. of Linn. Soc.* v. 2. 159. *With.* 92. *Hull.* 205.

C. spicata. *Huds.* 405. *Rehb.* 352. *Sibth.* 27. *Abbot.* 201. *Lightf.* 548.

C. palustris media, radice fibrosâ, caule exqui-

fitè triangulari, spicâ brevi, compaetiori. *Mich.*
Gen. 69. *t.* 33. *f.* 14.

Gramen cyperoides spicatum minus. *Raii Syn.*
424.

G. sylvaticum tenuifolium rigidiusculum. *Morif.*
seet. 8. *t.* 12. *f.* 27.

β . *G. cyperoides* spicis minoribus, minùsque compactis. *Scheuchz. Agr.* 488. *t.* 11. *f.* 5.

Angl. Greater Prickly Carex.

In sylvis et pratis humidis.

β . Near Eaton, Shropshire, on the banks of the driest ditches. *Rev. Mr. Williams. Gooden. Tr. of Linn. Soc. v. 3.* 79.

Perennis. Fl. Maio, Junio.

Radix fibrosa. Herba lœtè virens. Culmus crectus, pedalis vel sesquipedalis, nudus, triqueter, angulis superne asperis. Folia angusta, culmum plerumq[ue] superantia, margine carinâque scabra. Spica oblonga, uncialis vel sesquiuncialis, cylindracea, obtusa; basi interrupta, et subinde ramosa. Spiculæ circiter octo vel decem, subrotundæ, androgynæ, floribus masculis superioribus. Bracteæ ovatæ, scariosæ, carinatæ, in foliolum setaceum, asperum, desinentes, spiculas (saltè inferiorés) superantes, patentes. Glumæ ovatæ, scariosæ, ferrugineæ, mucronatae, carinâ viridi. Fructus divaricati, ovati, hinc convexi, virentes, demùm fusi, apice bifidi, lobis subdivaricatis, margine apicem versus scabri. Stigmata duo. Semen orbiculare, lenticulare.

An varicias β species sit distincta, spicâ compositâ, nec decompositâ, culmo glabriuscule, videant autoptæ.

15. CAREX spicâ elongatâ subdecompositâ basi *divulsa*. subramosâ; spiculis inferioribus remotis, fructibus crectiusculis margine lœviusculis.

Carex divulsa. Gooden. Tr. of Linn. Soc. v. 2.

160. *With.* 93. *Hull.* 205. *Sibth.* 28.

Abbot. 203. *Engl. Bot. t.* 629?

C. canef-

- C. canescens. *Huds.* 405. *Relb.* 351.
 C. nemorosa, fibrosâ radice, caule exquisitè triangulare, spicâ longâ, divulsâ seu interruptâ, capitulis solitariis præterquam ultimo. *Mich. Gen.* 69. *t.* 33. *f.* 10.
 Gramen cyperoides spicatum minus, spicâ longâ divulsâ seu interruptâ. *Raii Syn.* 424.
Angl. Grey Carex.

In nemorosis humidis.

Perennis. Fl. Maio.

Radix fibrosa. Culmus debilis, subreclinatus, pedalis aut sesquipedalis, nudus, acutè triquetus, angulis superne præcipue asperis. Folia angusta, culmo longiora, margine carinâque aspera. Spica elongata, infernè interrupta, albido-virens, basi subramosa. Spiculæ androgynæ, inferiores remotæ, subindè geminæ, infima sæpè composita. Bractæ parvæ, latè ovatæ, acutè carinatæ, in foliolum setaceum, asperum, spiculis plerumque longius, desinentes. Flores superiores masculi. Glumæ ovatæ, acutæ, albæ; inferiores præcipue carinâ viridi. Fructus erecto-patuli, ovati, hinc convexi, enerves, glabri, albidi, apice virentes, bifidi, margine carinato, omnino ferè lœvi, nec, nisi ad ipsum mucronem, ferrulato-scabro. Stigmata duo. Semen orbiculare, coquissimum.

vulpina. 16. CAREX spicâ supradecompositâ coarctatâ obtusâ, fructibus divergentibus, glumis acuminate, culmi angulis compressis acutissimis.

Carex vulpina. Sp. Pl. 1382. Gooden. Tr. of Linn. Soc. v. 2. 161. *Huds.* 404. *With.* 93. *Hull.* 206. *Relb.* 349. *Sibth.* 27. *Abbot.* 201. *Engl. Bot.* t. 307. *Leers.* 196. *t.* 14. *f.* 5. *Fl. Dan.* t. 308.

C. palustris major, radice fibrosâ, caule exquisitè triangulare, spicâ brevi, habitiori, compacâ, et spicâ

spicâ longâ, divulsâ, seu interruptâ. *Micb.*
Gen. 69. *t.* 33. *f.* 13.

Gramen cyperoides palustre majus, spicâ com-
pactâ. *Raii Syn.* 423. *Baub. Theatr.* 87.
Morif. sect. 8. *t.* 12. *f.* 24.

G. cyper. palustre triquetrum, spicâ integrâ.
Baub. Hist. v. 2. 497.

G. palustre cyperoides. *Ger. em.* 21.
Angl. Great Spiked Carex.

In paluatribus, et ad ripas fluviorum.

Perennis. Fl. Maio.

Radix fibrosa, cæspitosa. Culmus bipedalis, erectus,
strictus, firmus, rachi spicæ auctior, (quod in hâc sin-
gulare admodum), acutè triquierter, lateribus excavatis,
angulis compressis, asperrimis. Folia saturatè viridia,
lata, culmum superantia, margine carinâque seabra.
Spica decomposita vel supradecomposita, oblonga, ob-
tusa, coarctata, subindè parùm interrupta. Spiculæ
ovatæ, androgynæ, supernè masculæ. Bracteæ basi
ovatæ, apice setaceæ, asperæ, patentæ, exsertæ. Glu-
mæ ovatæ, acuminatæ, scariosæ, fuscæ, carinâ virente.
Fructus divergentes, ovati, acuminati, hinc convexi,
nervosi, marginati, apice bisidi, margine apieem versùs
scabri. Stigmata duo. Scimen elliptico-orbiculare.

17. CAREX spicâ subsupradecompositâ coarctatâ *teretius-*
acutiusculâ, spiculis glomeratis, fructibus pa-
tentibus gibbis, culmo *teretiusculo*.

Carex teretiuscula. *Gooden. Tr. of Linn. Soc.*
v. 2. 163. *t.* 19. *f.* 3. *Wiltb.* 95. *Hull.* 206.
Angl. Lesser Panicle Carex.

In paluatribus.

On St. Faith's bogs near Norwich, and at Barton
Mills. *Mr. Crowe.* In a bog to the north of
Queen's ferry. *Mr. J. Mackay.* Near Aberdeen.
Prof. Beattie.

Perennis.

Perennis. Fl. Maio.

Radix fibrosa, subrepens, sparsa, nec cæspitosa. Culmus pedalis vel sesquipedalis, erctus, acutè triquetus; lateribus prominulis, nec excavatis, unde terciusculus evadit. Folia ercta, rigidiuscula, longitudine circiter culmi, marginibus carinâque aspera. Spica ovato-oblonga, coarctata, decomposita, aliquando supradecomposita, acutiuscula. Spiculæ ovatæ, supernè masculæ. Bractæ ovatæ, scariosæ, insima mucronc foliacco, aspero, plerumque brevissimo. Glumæ ovatæ, fuscæ, margine scarioso, albo, carinâ lœvissimâ. Fructus fusi, ovati, acuminati, hinc planiusculi, extus valde convexi, basi utrinque gibbi, acumine utrinque ferrato-hispido, virente, apice bisido. Stigmata duo. Semen orbiculatum, extus gibbum.

Hanc post sequentis descriptionem, ut videtur, notavit Raius in *Syn. ed. 2.* 268. *ed. 3.* 422.

paniculata. 18. CAREX spicâ supradecompositâ paniculataramosâ acutâ interruptâ, fructibus patentibus acuminatis, culmo acutè triquetro.

Carex paniculata. *Sp. Pl.* 1383. *Gooden. Tr. of Linn. Soc.* v. 2. 164. *Huds.* 403. *With.* 94. *Hull.* 206. *Relb.* 351. *Sibth.* 28. *Abbot.* 203. *Leers.* 198. t. 14. f. 4.

C. radice repente, caule exquisitè triangulari, spicâ multiplici ferrugineâ; et spicâ multiplici fuscâ. *Mich. Gen.* 68. t. 33. f. 7.

Gramen cyperoides palustre elatius, spicâ longiore laxâ. *Raii Syn.* 422. *Moris. sect.* 8. t. 12. f. 23.

Cyperus alpinus longus inodorus, paniculâ ferrugineâ minus sparsâ. *Scheuchz. Prod.* 27. t. 8. f. 2.

Angl. Great Paniced Carex.

In palustribus spongiosis et aquosis.

Perennis. Fl. Junio.

Radix

Radix fibrosa, cæspites maximos constituens. Culmus 2- vel 3-pedalis, erectus, acutè triquierter; angulis afferrimis; lateribus planis, multincrvosif. Folia acuminata, ercta, marginibus carinâque aspera. Spica supradecomposita, paniculato-ramosa, ramulis alternis, subpatentibus, remotiusculis. Spiculæ ovatæ, floribus inferioribus fœmineis, paucioribus. Bracteæ ovatæ, acutæ, fuscæ, carinâ scabré, margine scarioso, albo; insima mucrone setaceo, scabro. Glumæ bracteis conformes, at carinâ lœvissimâ. Fructus virentes, ovato-lanceolati, hinc convexi, longius acuminati, bifidi, apicem versùs utrinque ciliato-scabri. Stigmata duo. Semen ovatum, extus convexum.

Variat spicâ magis vel minùs ramosa.

** *Spicis sexu distinctis: masculâ unicâ: bracteis membranaceis.*

19. CAREX bracteis membranaceis subaphyllis *digitata*, vaginantibus, spicis linearibus laxis erectis: masculâ breviori, foliis planis.

Carex digitata. Sp. Pl. 1384. Gooden. Tr. of Linn. Soc. v. 2. 166. Huds. 409. With. 95. Hull. 206. Engl. Bot. t. 615. Leers. 119. t. 16. f. 4.

Grainen caryophyllum polycarpon, fructu triangulo. Loef. Pruss. 112. t. 27.

G. caryophyllum montanum, spicâ variâ. Bauh. Prod. 9. Theatr. 48. Scheuchz. Agr. 448. t. 10. f. 14.

Cyperoides montanum nemorosum, caule triquetro-compresso, spicis ferrugineis tenuioribus, inter se distantibus, capsulis rariis dispositis, oblongis, turbinatis, trilateris. Mich. Gen. 65. t. 32. f. 9.

Angl. Fingered Carex.

In sylvis et umbrosis, solo calcareo, rariis.

Near Bath. Mr. Sole. Under St. Vincent's rocks, Brisiel,

Bristol, on the south side of the river. *Mr. W. Clayfield.* In the woods at Thorp-Arch, Yorkshire. *Sir T. Frankland.* Abundantly in Mackershaw wood, near Ripon. *Mr. W. Brunton jun.*

Perennis. Fl. Maio.

Radix fibrofa, nigra. Culmi adscendentes, patuli, spithamæi, teretiuseculi, vel obsoletè triquetri, læves; basi squamis cylindraceis, fusco-rubentibus, aphyllis, vaginati. Folia lætè-viridia, plana, marginibus ad basin retrorsum asperis, ad medium lævibus, apicem versùs antrorsum sebris, carinâ lævissimâ. Spicæ erectæ, lineares, laxæ, suboctofloræ; fœmineæ duæ vel tres, pedunculatæ, inferiores, sed masculam apice superantes; mascula sessilis, brevior. Bractæ ad basin pedunculorum, iisque duplò breviores, vaginantes, ore obliquæ, scariosæ, mucronulatæ, rariùs in foliolum brevissimum desinentes. Glumæ obovatæ, retusæ, fusco-rutilantes; carinâ viridi; margine scarioso, argenteo, imprimis in floribus masculinis. Fructus clavato-triquetri, virides, pubescentes, ore integro. Stigmata tria. Semen acutè triquetrum, fuscum.

clandestina. 20. CAREX bracteis membranaceis subaphyllis tina.

vaginantibus, spicis fœmineis remotis paucifloris vaginâ inclusis, foliis canaliculatis.

Carex clandestina. Gooden. *Tr. of Linn. Soc.* v. 2. 167. *With.* 96. *Hull.* 206.

C. n. 1370. *Hall. Hist.* v. 2. 189.

C. humilis. Leyser. *Hal.* 175.

C. prostrata. Allion. *Ped.* v. 2. 267.

Cyperoides montanum humile angustifolium, culmo veluti folioso spicis obsesto. Scheuchz.

Agr. 407. *t.* 10. *f.* 1.

Angl. Dwarf Silvery Carex.

In apricis petrosis rariùs.

On exposed sunny spots about St. Vincent's rocks, Bristol, where it was found by Mr. Sole.

Perennis.

Percennis. Fl. Maio.

Radix fibrosa. Culmi cæspitosi, unciales vel biunciales, erectiuseculi, obsoletè triquetri, glabri. Folia culmis triplo longiora, erecta, canaliculata, saturatè viridia, dorso sulcata, margine asperrima. Spica mascula terminalis, solitaria, erecta, lanceolata, multiflora, glumis imbricatis, lanceolatis, fuscis, margine scarioso, albo: femineæ duæ vel tres, inferiores, remotæ, alternæ, parvæ, paucifloræ, intrà bracteas proprias reconditæ. Bracteæ vaginantes, scariosæ, obtusæ. Fructus obovati, tricorni, ore indiviso. Stigmata tria. Semen triquetrum.

*** *Spicis sexu distinctis: masculâ unicâ, rariùs geminâ: bracteis foliaceis, et plerumque vaginantibus.*

21. CAREX vaginis elongatis pedunculo sub-pendula, æqualibus, spicis cylindricis longissimis nutantibus, fructibus confertissimis ovatis acutis.

Carex pendula. Huds. 411. Gooden. Tr. of Linn. Soc. v. 2. 168. With. 96. Hull. 206. Lightf. 564. Relh. 355. Abbot. 206. Sibth. 413. Curt. Lond. fasc. 3. t. 63.

C. Agastachys. Suppl. 414. Ehrh. Phyt. n. 19.

C. maxima. Scop. Carn. v. 2. 229.

C. n. 1396. Hall. Hist. v. 2. 196.

Gramen cyperoides, spicâ pendulâ longiore.

Raii Syn. 420. Moris. sect. 8. t. 12. f. 4.

G. cyper. latifolium, typhâ pendulâ longiore. Barell. Ic. t. 45.

Angl. Great Pendulous Carex.

In sylvis et sepibus humidiusculis.

About Braintree, Essex, and in other places. Ray.

About Hampstead and Highgate. Huds. Curt.

Near Woodbridge, Suffolk. Mr. Woodward.

In Worcestershire and Shropshire. Dr. Stokes.

By the river Tees near Croft; Mr. Robson.

With. In Breadalbane. *Rev. Mr. Stuart.* Near Hastings, Sussex. *Rev. Dr. Goodenough.*

Perennis. Fl. Maio, Junio.

Radix fibrosa. Culmus 3—6-pedalis, erectus, firmus, foliosus, triquierter, glaber, angulis apicem versus scabriusculis. Folia lata, saturatè viridia, rigida, dura, margine carinâque aspera, subtùs glaucescentia. Bracteæ foliaceæ, inferiores longissimæ, arcuato-dependentes, basi vaginantes, vaginis plerumque longitudine peduncularum. Spicæ 5 ad 7, pedunculatæ, arcuato-pendulæ, cylindraceæ, multifloræ, densissimæ; summa mascula, apice interdùm fœminea; cæteræ omnino ferè fœminæ. Glumæ lanceolatæ, acuminatæ, fuscæ, carinâ viridi. Fructus ovati, triquetri, subinflati, glabri, mucronulati, demùm emarginati. Semen triquetrum, fuscum.

Strigosa. 22. CAREX vaginis elongatis pedunculo subæqualibus, spicis filiformibus laxis sub nutantibus, fructibus lanceolatis triquetris nervosis.

Carex strigosa. *Huds.* 411. *Gooden. Tr. of Linn. Soc. v. 2. 169.* t. 20. f. 4. *With.* 96. *Hull.* 206. *Sibth.* 30. *Abbot.* 205. *Dicks. Dr. Pl.* 86.

C. Leptostachys. *Suppl.* 414. *Ehrb. Phyt.* n. 48.

Gramen cyperoides polysachyon majusculum latifolium, spicis multis longis strigosis. *Raii Syn.* 419.

Angl. Loose Pendulous Carex.

In nemorosis et sepibus.

In a lane at Black Notley; Essex; Mr. Dalc. *Ray.* Witham wood near Oxford. *Rev. Dr. Sheffield.* Noke wood. *Sibth.* Bedfordshire. *Abbot.* Hedenham wood, Norfolk. *Mr. Stone.*

Perennis.

Perennis. Fl. Aprili, Maio.

Radix fibrosa. Culmus erectus, bipedalis, gracilis, foliosus, triqueter, glaber. Folia latiuscula, tenuiora, lètè viridia, in margine carinâque scabra. Bracteæ foliis similes, basi vaginantes, vaginis longitudine ferè peduncularum, apice membranâ fuscâ, amplexicauli, evidenter marginatis. Spicæ 6 vel 8, pedunculatae, mox arcuato-pendulæ, filiformes, multifloræ, laxæ; summa mascula; reliquæ fœmineæ. Glumæ lanceolatae, viscentes. Fructus lanceolati, triquetri, acuti, nervosi, virentes, glabri. Semen elliptico-triquetrum.

23. CAREX vaginis pedunculo duplo brevioribus, *sylvatica*: spicis filiformibus laxiusculis nutantibus; fructibus ovatis triquetris rostratis.

Carex sylvatica. *Huds.* 411. *Gooden. Tr. of Linn. Soc.* v. 2. 183. *With.* 102. *Hull.* 207. *Lightf.* 562. *Rehb.* 356. *Sibth.* 30. *Abbot.* 205. *Dicks. H. Sicc. fasc.* 9. 13. *Fl. Dan.* t. 404. *Fl. Lapp. ed.* 2. 263.

C. vesicaria β. *Sp. Pl.* 1389.

C. Drymeia. *Suppl.* 414. *Ehrb. Phyt.* n. 58. *Wilden. Car. Berol.* 26. t. 3. f. 3.

C. patula. *Pollich. v.* 2. 597.

C. capillaris. *Leers.* 202. t. 15. f. 2.

C. n. 1395. *Hall. Hist.* v. 2. 196.

Gramen cyperoides sylvarum tenuius spicatum.

Raii Syn. 419. *Moris. sect.* 8. t. 12. f. 9.

Angl. Pendulous Wood Carex.

In sylvis frequens.

Perennis. Fl. Maio, Junio.

Radix fibrosa. Culmus 2—3-pedalis, erectus, foliosus, glaber, triqueter, lateribus convexis. Folia lètè viridia, suprà scabriuscula, marginibus carinâque aspera. Bracteæ foliis similes, longius vaginatæ. Spicæ 6 vel 8, post breve tempus arcuato-pendulæ, graciles, laxiusculæ, quarum una, rarissimè duæ, masculæ, omnes pendunculatae,

dunculatæ, pedunculis gracillimis, scabris, vaginâ propriâ plusquam duplò longioribus. Glumæ ovatæ, acuminatæ, scariosæ, albidæ, carinâ viridi, asperâ. Fructus ovato-triquetri, longius rostrati, glabri, apice emarginati. Semen triquetrum, fuscum.

depauperata 24. CAREX vaginis pedunculo triplò breviori-rata. spicis fœmineis remotis erectis paucifloris, fructibus inflatis rostratis.

Carex depauperata. Gooden. *Tr. of Linn. Soc.* v. 2. 181. *With.* 102. *Hull.* 207.

C. ventricosa. Curt. *Lond. fasc.* 6. t. 68.

C. triflora. Wilden. *Phytogr. fasc.* 1. 2. t. 1. f. 2.

Cyperoides sylvarum angustifolium, caule trilatero, spicis parvis, capsulis rariùs dispositis obefis, in angustum collum, vix ab unâ parte, ab alterâ verò magis, bifidum, coarctatis. Mich. *Gen.* 56. t. 32. f. 5.

C. vesicarium humile, loculis rarioribus. Tourn. *Inst.* 530. *Herb. Tourn.*

Angl. Starved Wood Carex.

In sylvis rariùs.

In Charlton wood. Rev. Dr. Goodenough. In dry woods near Godalmin, Surrey. Mr. Dickson.

Perennis. Fl. Maio, Junio.

Radix fibrosa. Culmus sesquipedalis, erectus, foliosus, obsoletè triquetè, striatus, glaber. Folia longius vaginantia, pallidè viridia, marginibus carinâque asperis. Bractæ foliis simillimæ, sed breviùs vaginatæ, erectæ. Spica mascula terminalis, erecta, multiflora, linearis, arctè imbricata, glumis ellipticis, obtusis, ferrugineis, margine scarioso albo, carinâ viridi glabré. Spicæ fœmineæ 3 ad 5, remotæ, laxæ, erectæ; pedunculis triquetris, scabris, vaginâ triplò longioribus; floribus 2 ad 4 vel 5, erectis, glumis obovatis, mucronatis, carinâ glabris. Fructus ovato-triquetri, nervosi, glabri, longius

giùs rostrati, ore scarioso, subintegro, demùm ventricosi.
Semen obovato-triquetrum.

25. CAREX vaginâ communi pedunculis duplò *capillaris*-
breviore, spicis fœmineis ovatis laxiusculis nu-
tantibus, fructibus acuminatis.

Carex capillaris. *Sp. Pl.* 1386. *Gooden. Tr. of Linn. Soc.* v. 2. 180. *Lightf.* 557. *Huds.* 651. *With.* 101. *Hull.* 207. *Fl. Dan.* t. 168. *Scop. Carn.* v. 2: 220. t. 59. *Dicks. H. Sicc. fasc.* 9. 14.

C. filiformis, ex errore. *Sm. Tour on the Contin.*
v. 3. 133.

Cyperoides alpinum, spicis seminiferis pendulis,
binis in summo caule. *Seguier. Veron.* v. 3.
83. t. 3. f. 1.

Angl. Capillary Carex.

In montibus scotieis.

On Ben-teskerney and other mountains in Breadal-
bane. *Rev. Mr. Stuart.*

Perennis. Fl. Julio, Augusto.

Radix fibrosa. Culmus erectus, palmaris, teretusculus,
striatus, glaber, nudus. Folia radicalia, culmo ferè
triplò breviora, angusta, acuta, planiuscula, margine
apicem versus scabra. Bractea communis figurâ folio-
rum sed minor, erecta, culmum terminans, basi vagi-
nans; interiores paucæ, minores. Pedunculi vaginâ
duplò, vix triplò, longiores, capillares, triquetri, sca-
bri, terni vel quaterni, ferè umbellati: quandoque,
præ rachi communi elongatâ, demùm distantes. Spica
mascula erecta, linearis, argenteo-fusca, subdecemflora:
fœmineæ 2 vel 3, nutantes sive pendulæ, ovatæ, breves,
laxiusculæ, ferè duodecimfloræ. Glumæ omnes ovatæ,
fuscæ; apice scariosæ, albæ. Fructus ovati, triquetri,
fuscæ, glabri, acuminati, exstriati, parùm ventricosi.
Semen ellipticum, acutè triquetrum.

Pseudo-cyperus. 26. CAREX vaginis subnullis, spicis cylindricis pedunculatis pendulis multifloris, fructibus patentibus sulcatis rostratis bifurcis.

Carex Pseudo-cyperus. *Sp. Pl.* 1387. *Gooden. Tr. of Linn. Soc.* v. 2. 188. *Huds.* 410. *With.* 104. *Hull.* 208. *Rehb.* 354. *Dicks. H. Sicc. fasc.* 7. 14. *Engl. Bot.* t. 242.

Gramen cyperoides spicâ pendulâ breviore. *Raii Syn.* 419. *Moris. sect.* 8. t. 12. f. 5. *Baub. Theatr.* 85.

Graminis cyperoidis genus, *pseudo-cyperus* *Lobelio*, spicis vel panniculis pendentibus ex longis pediculis. *Baub. Hist.* v. 2. 496.

Pseudo-cyperus. *Lob. Ic.* 76. *Ger. em.* 29. *Angl.* *Bastard-cyperus* *Carex.*

In nemorosis humidis et ad ripas fluyiorum vel foscarum.

On Hounslow heath. *Rev. Dr. Goodenough.* On St. Faith's bogs, near Norwich.

Perennis. Fl. Junio.

Radix fibrosa. Culmus pedalis et ultra, erectus, foliosus, acutè triquierter, angulis scabris. Folia pallidè viridia, vaginantia, suprà scabra, marginibus asperrima. Bracteæ foliis minores et haud vaginatæ. Pedunculi longissimi, triquetri, graciles, scabri; intermedii gemini. Spicæ cylindricæ, multifloræ, densæ; mascula erecta, rufa; fœmineæ pendulæ, virescentes. Glumæ ovatae, breves, longius aristatae, aristis scaberrimis. Fructus patentes, lanceolato-triquetri, nervosi, glabri, rostrati, apice bifurci. Semen elliptico-triquetrum, albidum.

limosa. 27. CAREX vaginis abbreviatissimis subnullis, spicis fœmineis oyatis pendulis, fructibus ellipticis compressis, radice repenti.

Carex limosa. *Sp. Pl.* 1386. *Gooden. Tr. of Linn. Soc.* v. 2. 187. *Huds.* 409. *With.* 103. *Hull.*

Hull. 208. *Lightf.* 556. *Fl. Dan.* t. 646.

Wilden. Car. Berolin. 26.

Cyperoides spicâ pendulâ, breviore, squamis e
spadiceo vel fusco rutilante viridibus. *Schenckz.*

Agr. 443. t. 10. f. 13.

Angl. Green and gold Carex.

In paludibus putridis spongiosis rariūs.

On turfey bogs in Yorkshire, Lancashire and Westmoreland. *Huds.* At Tullybanchar half a mile west of Comrie, near Crief, and in other places.

Rev. Mr. Stuart. Near Heydon, Norfolk, in a very deep rotten soil. *Rev. Mr. Bryant.* On St. Faith's Newton bogs. *Mr. Woodward.*

Perennis. Fl. Junio.

Radix repens, stolonifera, fibris densissimè tomentosis, fuscis. Herba subglauca. Culmi adscendentes, vix pedales, triquetri, striati, scabri, basi foliosi. Folia linearia, angusta, plana, culmo breviora, margine aspera, basi squamis fuscis vaginata. Bractea erecta, foliis simillima sed minor, basi brevissimè vaginata, vaginâ fuscâ, retusâ, margine scariosâ, albâ. Spica mascula culmum terminans, erecta, lanceolata, e fusco rutilans, multiflora, densa, glumis ovatis, acutis; fœmineæ duæ, sèpè solitariæ, pedunculatæ, cernuæ, mox pendulæ, ovatæ, multifloræ, densæ, formosæ, glumis latè ellipticis, mucronulatis, dorso viridibus, margine aureo-ferrugineis. Pedunculi longissimi, triquetri, læves. Fructus imbricati, glumis parùm longiores, elliptici, compressi, cæsi, nervosi, glabri, ore integerrimi, haud rostrati. Semen elliptico-triquetrum, vix compressum, fuscum.

Stolonibus latè propagatur, at rariūs flores profert.

28. CAREX diandra, vaginis subnullis, spicis ova-*atrata*.
tis pedunculatis pendulis : terminali infernè
masculâ, fructibus compressis.

Carex atrata. Sp. Pl. 1386. Fl. Suec. ed. 2.

330. Fl. Lapp. n. 324. Gooden. Tr. of Linn.

Soc. v. 2. 189. Huds. 409. With. 105. Hull. 208. Lightf. 555. Dicks. Dr. Pl. 87. Fl. Dan. t. 158.

Cyperoides alpinum pulchrum, foliis caryophyl-læis, spicis atris et tumentibus. Scheuchz. Agr. 481. t. 11. f. 1, 2.
Angl. Black Carex.

In pratis alpinis.

On the Welch mountains; plentiful about Llan-berris, *Huds.* Frequent on the highland moun-tains. *Lightf.*

Perennis. Fl. Junio, Julio.

Radix fibrosa. Culmus erectus, pedalis et ultrà, trique-ter, striatus, angulis læviusculis, basi præcipue foliosus. Folia lætè viridia, lata, plana, culmo breviora, mar-gine scabra. Bracteæ foliaceæ, amplexicaules, vix vagi-natæ, patentes; insima culmum plerumque superans. Spicæ sæpius quatuor, pedunculatæ, erectæ, mox pen-dulæ, ovatæ, densæ, multifloræ; terminalis infernè mascula, floribus diandris, supernè sœminea; cæteræ sœ-mineæ, floribus aliquot hermaphroditicis diandris inter-sinætis. Pedunculi triquetri, læves. Glumæ omnes ova-tæ, nigræ. Fructus imbricati, glumis parùm breviores, latè elliptici, conpresso, flavi, euerves, glabri, vix rostrati, ore emarginati. Semen triquetrum, breve, fuscum. Stigmata tria.

Archetypum certè fidendum in herbario Linnæano desi-deratur. Ad sunt profectò plura exemplaria, e Lappo-niâ nuper allata, hujuscce nomine perperam insignita, quibus omnibus spicæ sexu distinctæ sunt, et arillus longè a verâ plantâ Linnæi atque Scheuchzeri di-verlus.

pulla. 29. CAREX digyna, vaginis nullis, spicis ovatis: inferiori pedunculatâ, fructibus ellipticis infla-tis subrostratis emarginatis.

Carex pulla. Gooden. Tr. of Linn. Soc. v. 3. 78. t. 14. Hull. 208.

Angl.

Angl. Russel Carex.

In pascuis alpinis.

On the mountains of Scotland. *Mr. Dickson.*

Found on Ben Lawers in 1793, by Mr. J. Mackay.

Perennis. Fl. Julio.

Radix repens. Culmus erectiusculus, plerumque spithameus, triqueter, striatus, angulis superne scabris. Folia subhereta, plana, margine scabra. Bractea foliacea, aniplexicaulis, vix vaginata, margine aspera. Spica mascula terminalis, erecta, lanceolata, multiflora, triandra; foemineae saepius duæ, remotæ, ovatae, erectæ, quarum inferior longius pedunculata, superior subsessilis atque abbreviata, saepe nulla. Pedunculus scaber. Glumæ omnes elliptico-lanceolatae, obtusæ, atro-fuscæ, margine tenuissimo albido, nervo obsoleto. Fructus patuli, gluniis longiorcs, elliptici, inflati, glabri, enerves, brevè rostrati, emarginati, basi pallidi, apice atro-fusci, nitidi. Semen triquetrum. Stigmata duo.

30. CAREX vaginis brevissimis, spicis cylindricis *palleascens*, pedunculatis: fructiferis pendulis, fructibus ellipticis inflatis obtusis.

Carex palleascens. *Sp. Pl.* 1386. *Gooden. Tr. of Linn. Soc.* v. 2. 186. *Huds.* 410. *With.* 103. *Hull.* 208. *Lightf.* 558. *Relb. Suppl.* 3. 7. *Sibth.* 29. *Abbot.* 204. *Dicks. H. Sicc. fasc.* 4. 16. *Fl. Dan.* t. 1050. *Leers.* 203. t. 15. f. 4. *Rel. Rudb.* 2.

Gramen cyperoides polystachyon flavicans, spicis brevibus prope summitatem caulis. *Raii Syn.* 419. *Phuk. Phyt.* t. 34. f. 5. *Mich. Gen.* 61. t. 32. f. 13.

Angl. Pale Carex.

In pratis, pascuis et nemorosis humidis frequens.

Perennis. Fl. Maio, Junio.

Radix.

Radix fibrosa. Culmus erectus, pedalis aut sesquipedalis, triquierter, striatus, angulis superne acutioribus, asperis, basi foliosus. Folia plana, pallida, subpilosa, margine scabriuscula. Bracteæ foliaceæ, erectæ, culmum longè superantes, brevissimè vaginatæ. Spica mascula terminalis, lanceolata, erecta, densa, pallidè ferruginea; fœminæ tres, pedunculatæ, ovatæ, erectæ, mox pendulæ, cylindraceæ, obtusæ. Pedunculi triquetri, læves. Glumæ ovatæ, acuminatæ, lutescentes, carinâ viridi. Fructus longitudine glumarum, elliptici, inflati, obtusissimi, mutici, enerves, glabri, pallidè virides. Semen obovato-triquetrum.

flava. 31. *CAREX vaginis abbreviatis pedunculo subæqualibus, spicis fœminis subrotundis, fructibus rostratis arcuato-deflexis, culmo læviusculo.*

Carex flava. Sp. Pl. 1384. Gooden. Tr. of Linn. Soc. v. 2. 173. Huds. 407. With. 98. Hull. 207. Lightf. 551. Relb. 352. Sibth. 29. Abbot. 204. Dicks. H. Sicc. fasc. 3. 14. Fl. Dan. t. 1047. Leers. 198. t. 15. f. 6.

Gramen palustre echinatum. Raii Syn. 421. Ger. em. 17. Baub. Hist. v. 2. 497.

G. cyperoides palustre aculeatum germanicum. Moris. sedl. 8. t. 12. f. 19. Baub. Theatr. 109.

Angl. Yellow Carex.

In paludosis frequens.

Perennis. Fl. Maio, Junio.

Radix repens. Culmus erectus, spithameus, vel sæpius pedalis, acutè triquierter, angulis plerumque lævibus, interdum scabriusculis, infernè foliosus. Folia amœnè viridia, erecta, latiuscula, plana, culmo longiora, margine carinâque aspera. Bracteæ foliaceæ, culmum longè superantes; superiores divaricato-deflexæ, brevissimè vaginatæ; insinua longius vaginata, niinùsque patens. Spica mascula lanceolata, erecta, densa, plerumque

rumque solitaria, rarissimè gemina; fœmineæ circiter tres, ovato-subrotundæ, patentes; superiores plerumque approximatæ, ferè sessiles; infima remotiuscula, pendunculata, pedunculo vaginam parùm superante. Glumæ ovatæ, fulvæ vel xerampelinæ, nervo virenti, margine scarioso, albo. Fructus glumis longiores, luteo-virides, undique patentes, arcuato-deflexi, ovati, rostrati, inflati, subtriquetri, glabri, nervosi, emarginati, angulis omnino lœvibus. Semen parvum, obovato-triquetrum, nigrum.

32. CAREX vaginis elongatis pedunculo breviori-*fulva*, bus, spicis fœmineis ovatis, fructibus rostratis rectis, culmo scabro,

Carex fulva. *Gooden. Tr. of Linn. Soc.*, v. 2, 177,
t. 20, f. 6. v. 3. 77.

Angl. Tawny Carex.

In paludosis.

Near Eaton, not far from Shrewsbury; Rev. Mr. Williams. *Gooden*. Very common in Mearns-shire, North Britain. *Prof. James Beattie, jun.*

Perennis. Fl. Junio, Julio.

Radix repens. Culmus erectus, gracilis, ferè pedalis, triqueter, angulis supernè præcipue acutis, scaberrimus. Folia erecta, angusta, culmo breviora, marginibus catinâque aspera. Bracteæ foliaceæ, erectæ, culmo breviores, vaginis longis, apice fuscescentes. Spica mascula linearis, erecta, densa, multiflora, glumis fuscis, obtusis, margine scarioso, albo, nervo obloseto, concolori; fœmineæ duæ, rariùs tres, ovatæ, erectæ, remotissimæ, pedunculatæ, pedunculis vaginâ circiter sesqui-longioribus, glumis ovatis, fuscis, margine subscariosis, omnino muticis, nervo pallido, obsoleto. Fructus glumis longiores, fusco-virides, erecto-patentes, ovati, rostrati, subtriquetri, glabri, obsoletè nervosi, emarginati, angulis apicem versus scabriusculis. Semen triquetro-subrotundum, luteo-fuscum, duplò majus quam in præcedente.

Videtur

Videtur species a C. flavâ distinctissima. Glumis omnino mutieis a C. distante differt.

extensa. 33. CAREX vaginis pedunculisque brevissimis, bracteis longissimis, spicis foemineis subrotundis, fructibus ovatis, culmo laevi.

Carex extensa. Gooden. Tr. of Linn. Soc. v. 2.
175. t. 21. f. 7. With. 99. Hull. 207.

Cyperoides echinatum majus. Dill. in Raff. Syn.
421.

Angl. Long-bracted Carex.

In maritimis.

Found on Cley beach, Norfolk, June 18th 1776, by the Rev. H. Bryant. *Herb. Rose.* Near Harwich, and on the marshy part of Braunton Burrows, Devonshire. *Rev. Dr. Goodenough.* On the coast to the east of St. Andrews. *Mr. J. Mackay.*

Perennis. Fl. Junio.

Radix fibrosa. Culmus erectus, vel ascendens, subindè ineurvus, spithameus vel pedalis, obtuse triqueter, undique laevissimus. Folia recurva, angusta, eanalieulata, margine apicem versus seabra. Bracteæ foliaceæ, longissimæ, patentissimæ, arcuatae, vaginis dilatatis, brevissimis. Spica maseula plerumque solitaria, laneo-lata, obtusiseula; foemineæ duæ vel tres, approximatæ, subfessiles, erectæ, ovatae, rariùs cylindraceaæ, demum subrotundæ. Glumæ omnes latè ovatae, trinerves, mucronulatae, ferrugineaæ, dorso virides. Fructus glumis longiores, erecto-patentes, ovati, subtriquetri, fusco-virides, nervosi, glabri, haud rostrati, emarginati, angulis undique laevissimiis. Semen triquetrum, gibbum, fuscum.

Varietatem spicis elongato-cylindraceis invenit D. Dickson.

diflans. 34. CAREX vaginis elongatis pedunculo subæqualibus,

æqualibus, spicis oblongis remotissimis, glumis mucronulatis, culmo lœvi.

Carex distans. *Sp. Pl.* 1387. *Gooden. Tr. of Linn. Soc.* v. 2. 178. *Huds.* 412. *With.* 100. *Hull.* 207. *Relb.* 355. *Sibth.* 31. *Abbot.* 262. *Fl. Dan.* t. 1049.

C. n. 1382. *Hall. Hist.* v. 2. 192. Davall.

Cyperoides spicis parvis, longe distantibus. *Seguier. Veron.* v. 3. 77. *Herb. Linn. Scheuchz. Agr.* 431.

Gramen cyperoides spicis parvis longissimè distantibus. *Raii Syn.* 420.

Angl. Loose Carex.

In palustribus, præcipue maritimis.

Perennis. Fl. Junio.

Radix fibrosa. Culmus pedalis aut sesquipedalis, erectiusculus, debilis, triquierter, lœvis, vel angulis apicem versùs quandoque scabriusculis. Folia latiuscula, brevia, plana, margine aspera. Bracteæ foliis similcs, longius vaginatæ, remotissimæ, vel per caulem sparsæ. Spica mascula lanceolata, obtusa, multiflora, glumis obovatis, obtusis, acutè carinatis, obsoletè mucronulatis; fœmineæ duæ vel tres, rariùs quatuor, sparsæ, remotissimæ, erectæ, ovato-oblongæ, pedunculatæ, pedunculis inferioribus vaginâ sesquialongioribus, superioribus longitudine circiter vaginæ. Glumæ fœminearum latè ovatæ, trinerves, aristatæ, seu mucronulatæ, ferrugineæ, carinâ virenti, interdùm scabriæ. Fructus glumis longiores, ovati, fusi, multinervosi, glabri, brevè rostrati, bifidi, angulis lœvissimis. Semen triquetrum, fuscum, oculo armato granulatum.

[Floræ Danicæ synonymon huc retuli ob glumam aristatam.

35. CAREX *vaginis elongatis* pedunculo breviori- *binervis.* *bus,* spicis cylindraceis remotis subcompositis, *glumis mucronulatis,* fructibus binervibus.

Carex

Carex binervis. *Tr. of Linn. Soc. v. 5.* 268.

C. distans. *Lightf. 561,* ex descr.

Angl. Green-ribbed Carex.

In ericetis siccioribus.

Very common on the driest moors about Aberdeen.

Prof. Beattie. Near Edinburgh. *Mr. J. Mackay.*

Perennis. Fl. Junio.

Præcedente major. Culmus erectus, firmus, sesquipedalis, bipedalis, vel tripedalis, obtusè triquetter, lævis, apicem versus hinc scaber. Folia erecta, latiuscula, acuminata, glaucescentia, marginibus carinâque aspera. Bractæ foliis simillimæ, elongatæ, erectæ, longius vaginatæ, remotæ. Spica mascula utrinque attenuata, sœpè sesquiuncialis, multiflora, glumis densissimè imbricatis, ellipticis, obtusis, submucronulatis, nigranticibus, carinâ acutâ, virenti; fœmineæ tres aut quatuor, rarissimè quinque, sparsæ, cylindraceæ, erectæ, inferiores remotissimæ, longius pedunculatæ, atque sœpius basi compositæ seu ramosæ, quandoque omnes superne masculæ. Glumæ fœminearum ovatæ, nigræ, mucronulatæ, carinâ virenti, glabrâ, mucronulo scabro. Fructus glumis longiores, ovati, vix rostrati, glabri, nitidi, intus apiceque sanguineo-fusci, extus pallidi, nervis duobus lateralibus viridibus præcipue conspicuis. Semen acutè triquetrum, albidum.

præcox. 36. CAREX vaginis abbreviatis pedunculo subæqualibus, spicis ovatis approximatis, glamis mucronulatis, fructibus subrotundis pubescens tibus.

Carex præcox. *Jacq. Austr. t. 446.* *Gooden.*
Tr. of Linn. Soc. v. 2. 170. *With. 97.* *Hull.*
206. *Sibth. 29.* *Abbot. 205.* *Widen. Car.*
Berol. 23.

C. saxatilis. *Huds. 408.*

C. montana. *Rehb. 353.* *Lightf. 551.*

Gramen cyperoides vernum minimum. *Raii*
Syn. 421.

G. cyperoides spicatum. *Ger. em.* 22. *Morif.*
sect. 8. *t.* 12. *f.* 11.
Angl. Vernal Carex.

In apricis, ericetis, et pascuis siccioribus, vulgaris.

Perennis. Fl. Aprili.

Radix stolonibus reptantibus. Culmus adscendens, palmaris aut spithameus, nudus, obtusè triqueter, lævis. Folia brevia, recurva, subrevoluta, margine aspera, superè quandoque scabra. Bracteæ breves, vaginis brevibus, superne dilatatis, obtusis, longitudine ferè peduncularum. Spica mascula obovata vel subclavata, multiflora, glumis ellipticis, ferrugineis, unicoloribus, submuticis, antheris sulphureis, valdè conspicuis; fœmineæ plerumque duæ, approximatæ, brevitè pedunculatæ, ovatae, erectiæ, glumis latè ovatis, mucronulatis, ferrugineis, carinâ virenti. Fructus vix longitudine glumarum, subrotundi, vel triquetro-pyriformes, pubescentes, virentes, apice fusi. Semen albidum.

37. CAREX vaginis nullis, spicis fœmineis sessili-pilulifera. bus confertis subrotundis, glumis mucronulatis, fructibus subrotundis villosis.

Carex pilulifera. *Sp. Pl.* 1385. *Gooden. Tr. of Linn. Soc.* v. 2. 190. *Huds.* 408. *With.* 105. *Hull.* 208. *Relb.* 353. *Sibth.* 29. *Dicks. H. Sicc. fasc.* 9. 12. *Lightf.* 554. *Wilden. Car. Berol.* 24. *t.* 2. *f.* 2.

C. montana. *Sp. Pl.* 1385. *Huds.* 407?

C. filiformis. *Fl. Dan.* *t.* 1048.

Gramen cyperoides, spicis brevibus congestis, folio molli. *Raii Syn.* 421.

G. cyperoides tenuifolium, spicis ad summum caulem sessilibus, globulorum ænulatis. *Pluk. Phyt.* *t.* 91. *f.* 8. *Raii Syn.* 422.

Angl. Round-headed Carex.

In pascuis et ericetis frequens.

Perennis,

Perennis. Fl. Aprili, Maio.

Radix fibrosa. Culmus sæpius recumbens, incurvus, do-
dratalis, debilis, nudus, triquetter, apice scaber. Folia
lætè viridia, suprà marginibusque carinâque scabra. Braëteæ setaceæ, breves, triquetrae, vaginis omnino
ferè nullis. Spica mascula lanceolata, acuta, glumis
lanceolatis, acutis, ferrugineis, carinâ viridi, margine
scarioso, albo; fœmineæ duæ vel tres, sessiles, congo-
meratæ, subrotundæ, patentæ, glumis ovatis, acutis,
mucronulatis, colore masculinarum. Fruëtus glumis
breviores, subrotundi, obsoletè triquetri, virides, undi-
que villosi. Semen ferè globosum, obsoletè triquetrum,
flavescens.

Variat glumis muticis, minùs conspicuè carinatis.

tomentosa. 38. CAREX vaginis brevissimis, spicis fœmineis
subsessilibus cylindraceis obtusis, glumis ellip-
ticis acutis, fructibus tomentosis.

Carex tomentosa. Mant. 123. Leers. 200. t. 15.

f. 7. Wilden. Car. Berol. 24. Dicks. Dr. Pl.

43. Tr. of Linn. Soc. v. 5. 269.

Angl. Downy-fruited Carex.

In pratis rariùs.

In meadows near Merston Meafey, Wiltshire. Mr.
Teesdale.

Perennis. Fl. Junio.

Radix repens. Culmus pedalis, crectus, nudus, acutè
triqueter, angulis supernè scabris. Folia culmo breviora,
erecta, plana, lætè viridia, utrinque marginique scabra. Braëteæ foliaceæ, erecto-patentes, culmum vix super-
antes, vaginâ brevissimâ, aut ferè nullâ. Spica mascula
lanceolata, obtusifcula, glumis lanceolatis, ferrugineo-
fuscis, carinâ virenti, superioribus quandoque mucronu-
latis; fœmineæ plerumque duæ, parùm remotæ, bre-
vissimè pedunculatæ, cylindraceæ, obtusæ, longitudine
variæ, glumis elliptico-ovatis, vix mucronulatis, ferru-
ginco-fuscis, carinâ latè virenti. Fruëtus longitudine
circiter glumarum, dense imbricati, subrotundi, parùm
compressi, vix triquetri, virides, tomento densissimo,
brevi,

brevi, albido, demùm aureo, undique vestiti. Semen albidum, obsoletè triquetrum.
Diversa a C. tomentosâ Lightf. atque Hudſ. quæ C. filiformis est.

39. CAREX digyna, vaginis nullis, spicis ovatis: *rigida*.
superioribus sessilibus, foliis subrecurvis rigidis, fructibus compressiusculis.

Carex rigida. Gooden. *Tr. of Linn. Soc.* v. 2.
193. t. 22. f. 10. *With.* 106. *Hall.* 208.

C. saxatilis. Fl. Dan. t. 159.

C. n. 1378. *Hall. Hist.* v. 2. 191, ex Herb.
Jacq.

Cyperoides Germanicum, foliis brevibus rigidis
acutis, caule rotundo-triquetro, spicis parvis,
quamis obtusè mucronatis, capsulis oblongis
turbanatis, in angustum et longiusculum api-
cem attenuatis. Mich. Gen. 61. t. 32. f. 4.

Angl. Rigid Carex.

In summis alpibus.

On the top of Snowdon, Wales. Mr. Hudson.

Found on the top of Ben Lomond in 1782.—It
has since been observed by Mr. Dickson and Mr.
Mackay in abundance on the summits of several
of the highland mountains.

Perennis. Fl. Junio, Julio.

Radix crassa, repens. Culmus sæpiùs incurvus, palmaris
vel spithameus, nudus, rigidus, acutè triqueter, angu-
lis asperis. Folia rigida, recurva, glaucescentia, mar-
gine carinâque aspera. Bracteæ foliaceæ, culmo bre-
viore, sæpiùs recurvæ, basi auriculatæ. Spica mascula
brevis, ovata, obtusiuscula, crassa, rariùs gemina, glu-
mis obtusissiniis, muticis, nigris, carinâ tenui, viridulâ;
fœmineæ plerumque tres, ovatae, quarum duæ supe-
riores sessiles terminali approximatæ, inferior remotius-
cula, breviùs pedunculata, erecta, glumis ellipticis, ob-
tusiusculis, muticis, nigris, margine albido, carinâ te-

nni, viridi. Fructus glumis longiores, densè imbricati, elliptici, compressiusculi, hinc planiusculi, glabri, vi- rentes, extùs fuscæ; subindè (optimè monente Goode-novio) apice producti et recurvi, semper ore indivisi. Stigmata duo, crassiuscula.

panicea. 40. CAREX vaginis elongatis pedunculo subdiu-
nio brevioribus, spicis fœmincis laxiusculis re-
motis, fructibus inflatis, culmo lævi.

Carex panicea. *Sp. Pl.* 1387. *Gooden. Tr. of
Linn. Soc.* v. 2. 179. *Huds.* 410. *With.* 100.
Hull. 207. *Rehb.* 354. *Sibth.* 30. *Abbot.* 205.
Leers. 203. t. 15. f. 5. *Fl. Dan.* t. 261, haud
benè. *Rel. Rudb.* 2.

Cyperoides foliis caryophylleis, caule rotundo-
triquetro, spicis e rarioribus et tumidioribus
vesicis compositis. *Mich. Gen.* 61. t. 32. f. 11.

Gramen cyperoides foliis caryophylleis, spicis e
rarioribus et tumidioribus granis compositis.

Raii Syn. 418. *Pluk. Phyt.* t. 91. f. 7.

Angl. Pink-leaved Carex.

In pratis et pascuis humidis.

Perennis. Fl. Maio, Junio.

Radix repens. Culmus erectus, spithameus aut pedalis,
nudus, triqueter, striatus, angulis lævibus. Folia glauca,
brevia, erectiuscula, marginibus carinâque apicem ver-
sûs scabriusculis. Bracteæ foliaceæ, acuminatæ, cul-
mo breviores, vaginis longiusculis, sulcatis, subindè
flavescensibus. Spica mascula sæpè gemina, lanceo-
lata, acuta, densa, multiflora, glumis ferrugineo-fuscis,
obtusis, latitudine variis, carinâ virentibus; fœmineæ
plerumque duæ, subindè apice masculæ, remotæ, erec-
tæ, pedunculatæ, cylindricæ, glumis laxè imbricatis,
ovatis, ferrugineo-fuscis, acutis, muticis, margine sca-
riofo, carinâ viridi. Fructus ovati, inflati, obtusifus-
culi, glabri, subnervosi, e viridi fuscescentes, ore indivisi.
Semini breve, triquetruim, fuscum, usilagini frequentè
obnoxium.

C. *panicea*

C. panicea et recurva sub adolescentiâ valdè similes—at in hâc vagina valdè abbreviata, in illâ pedunculo inferiori dimidio brevior. *D. Goodenough.*

41. CAREX vaginis abbreviatis, spicis fœmineis *recurva*: cylindricis pendulis, fructibus elliptico-triquetris scabriusculis, radice repente.

Carex recurva. *Huds.* 413. *Gooden. Tr. of Linn.*
Soc. v. 2. 184. *Wittb.* 102. *Hull.* 208. *Relh.*
356. *Sibth.* 31. *Abbot.* 206. *Fl. Dan.*
t. 1051.

C. glauca. *Scop. Carn.* v. 2. 223. *Pollich.* v. 2.
594.

C. limosa β . *Leers.* 201. t. 15. f. 3.

Gramen cyperoides, foliis caryophylleis, spicis oblongis e pediculis longioribus pendulis. *Raii*
Syn. 418.

G. cyperoides nemorosum, spicâ subnigrâ re-curvâ. *Morif. seet.* 8. t. 12. f. 14.

Angl. Glaucous Heath Carex.

In pratis, pascuis, nemoribus et ericetis humidi-oribus.

Perennis. Fl. Maio, Junio.

Radix repens, squamis fuscis vaginata. Herba glauca. Culmus erectus, pedalis, triqueter, lateribus sœpè convexis, supernè scaber. Folia erecta, culmo breviora, latiuscula, plana, marginie carinâque scabra. Bracteæ foliaceæ, erectæ, culmo subæquales, vaginis brevibus, auriculatis. Spica mascula linear-lanceolata, erecta, subinde gemina, quandoque multiplicata, glumis ob-ovatis, obtusis, fuscis, carinâ luteolâ; fœmineæ sœpiùs duæ, longius pedunculatæ, erectæ, mox pendulæ, cylindraceæ, densæ, multifloræ, parùm remotæ, glumis ovatis, acutis, submucronatis, nigricantibus, margine albidis, nervo lutescentibus. Fructus imbricati, elliptici, obtuse triquetri, subgibbosæ, ferè enerves, plûs minus scabriusculi, vel subpubescentes, ferrugineo-virentes, demum nigri, ore indiviso. Semen triquetrum,

brevè, retusum, fuscum aut nigrum, angulis carinatis, pallidis.

Varietatem spicis masculinis numerosis, septenis, misit D. Beattie.

cæspitosa. 42. CAREX digyna, vaginis nullis, spicis sessilibus cylindricis obtusis, bracteis auriculatis, fructibus persistentibus.

Carex cæspitosa. Sp. Pl. 1388. Gooden. Tr. of Lim. Soc. v. 2. 195. t. 21. f. 8. With. 106. Hull. 208. Sibth. 31. Lightf. 560.

Gramen caryophylleum, angustissimis foliis, spicis sessilibus brevioribus erectis non compactis. Dill. in Raii Syn. 418. Moris. v. 3. 243. Angl. Tufted Bog Carex.

In palustribus, et nemorosis humidis.

Percennis. Fl. Maio.

Radix repens, implexa, cæspitosa. Culmus erectus, spithameus vel pedalis, acutè triqueter, angulis superne asperis. Folia erecta, culmum subaquantia, latè viridia, parùm glauca, molliuscula, acuminata, margine carinâque seabra. Bracteæ foliis similes, sessiles, basi utrinque auriculâ rotundatâ magnitudine variâ, vaginâ nullâ. Spica mascula solitaria, rarissimè gemina, lanceolata, acutiuscula, erecta, glumis fuscis, obtusis, nervo abbreviato, luteo; fœmineæ duæ vel tres, sessiles, erectæ, cylindricæ, obtuseæ, densissimæ, multifloræ, approxiniatæ, glumis ellipticis, obtusis, adpressis, nigris, nervo viridi. Fructus imbricati, latè elliptici, compressi, nervosi, virentes, glabri, persistentes. Semen lenticulare, compressum, fuscum.

striata. 43. CAREX digyna, vaginis nullis, spicis subsessilibus cylindricis elongatis acutis, bracteis subauriculatis, fructibus deciduis.

Carex striata. Gooden. Tr. of Lim. Soc. v. 2. 196. t. 21. f. 9. With. 107. Hull. 208.

C. cæspitosa. Huds. 412. Lightf. 561, β.

C. acuta

C. acuta α . *Leers.* 204. *t.* 16. *f.* 1.

C. n. 1400. *Hall. Hist.* *v.* 2. 198. *Davall.*

Gramen cyperoides foliis caryophylleis, spicis erectis sessilibus, e seminibus confertis compositis. *Raii Syn.* 418.

G. cyperoides palustre spicâ pendulâ. *Locf.* *Pruss.* 116. *t.* 30.

Angl. Glaucous Straight-leaved Carex.

In palustribus.

Near Norwich. *Mr. Pitchford.* In various other places not uncommon.

Perennis. Fl. Aprili.

Radix repens. Herba glaucescens, præcedente duplo major. Culmus erectus, sesquipedalis aut bipedalis, acutè triqueter, angulis supernè asperis. Folia erecta, stricta, aeuminata, culmo breviora, margine carinâque aspera, inferiora vaginis demùm laceris, fibrosis. Bracteæ foliis simillimæ, sessiles, erectæ, strictæ, basi utrinque subauriculatæ, auriculâ oblongâ, mox oblitteratâ, vaginâ nullâ. Spicæ brevissimè pedunculatæ, erectæ, sesquiunciales aut biunciales, lineari lanceolatæ, acutæ, subapproximatæ, densissimæ, multifloræ, glumis omnibus elliptico lanceolatis, obtusis, adpressis, nigris, nervo viridi; masculæ sæpius duæ; fœmineæ plerumque tres, apice frequentiùs masculæ. Stigmata duo. Fructus imbricati, ovati, acuti, compressi, nervosi, virentes, glabri, decidui. Semen ferè præcedentis, orbiculato-lenticulare.

**** Spicis sexu distinctis : masculis pluribus.

44. CAREX digyna, spicis filiformibus : floren- *acuta*. tibus cernuis : fructiferis erectis, fructibus ellipticis obtusis apice indivisis.

Carex acuta. *Sp. Pl.* 1388, β . *Fl. Suec.* 334, α .

Gooden. Tr. of Linn. Soc. *v.* 2. 203. *With.* 109.

Hull. 209. *Huds.* 413, β . *Engl. Bot.* *t.* 580.

G 3

C. gracilis.

C. gracilis. *Curt. Lond. fasc. 4. t. 62.* *Rell. 357.*
Sibth. 32.

Gramen cyperoides majus angustifolium. *Raii*
Syn. 417.

Angl. Slender-spiked Carex.

In aquosis, pratisve graminosis humidis, frequens.

Perennis. Fl. Maio.

Radix latè repens. Culmus altitudine varius, sèpiùs bipedalis et ultra, erectus, apice nutans, demùm strictus, acutè triquierter, angulis asperis. Folia latè viridia, erecta, apice recurvato-pendula, margine carináque aspera, culmo breviora. Bracteæ foliaceæ, elongatae, sessiles, basi subauriculatae, vaginis nullis. Spicæ subsessiles (fœmineæ inferiores demùm breviùs pedunculatæ), filiformes, graciles, elongatae, densæ, multifloræ, florentes cernuæ, fructiferæ erectæ, glumis oblongis, acutis, muticis, nigris, nervo pallido; masculæ duæ vel tres; fœmineæ tres aut quatuor, apice sèpè masculæ. Stigmata duo. Fructus imbricati, elliptici, obtusi, compressi, nervosi, virentes, glabri, decidui. Semen parvum, obovatum, bicarinatum, nec triquetrum.

Culmus mirè variat pro loco natali et cœli temperie, triangularis, tripedalis. *D. Goodenough.*

paludosa. 45. CAREX spicis cylindricis obtusiusculis erectis, glumis fœminearum aristatis; muscularum obtusis, fructibus ellipticis emarginatis.

Carex paludosa. *Gooden.* *Tr. of Linn. Soc.*
v. 2. 202. *Wth. 108.* *Hull. 209.*

C. acuta. *Curt. Lond. fasc. 4. t. 61.* *Huds.*
413, γ. *Lightf. 566, β.*

Gramen cyperoides minus angustifolium. *Dill.*
in Raii Syn. 418.

Angl. Lesser Common Carex.

In aquosis, et ad ripas fluviorum vel fossarum
frequens.

Perennis.

Perennis. Fl. Maio.

Radix valde repens. Herba glaucescens. Culmus sesquipedalis vel bipedalis, erectus, acutè et inæqualiter triqueter, angulis asperimis. Folia latiuscula, erecta, margine carinaque aspera. Bracteæ foliaceæ, haud vaginatæ, rariùs auriculatæ, culmum plerumque superantes. Spicæ cylindricæ, obtusifusculæ, erectæ; masculæ circiter tres; fœmineæ totidem, breviùs pedunculatæ, apice sèpè masculæ. Glumæ mascularum oblongæ, obtusæ, fuscæ, nitidæ, carinâ viridi; fœminarum lanceolatæ, acutæ, sèpiùs mucronulatæ, mucronulo scabro, fuscæ, carinâ virenti. Stigmata tria. Fructus imbricati, elliptico-lanceolati, acutiusculi, nervosi, apice demùm emarginati. Semen triquetrum.

46. CAREX spicis erectis: masculis triquetris: *riparia*.
fœmineis cylindricis, glumis omnibus acuminateis, fructibus rostratis bifurcatis.

Carex riparia. Curt. Lond. fasc. 4. t. 60.
Gooden. Tr. of Linn. Soc. v. 2. 200. With. 108.
Hull. 209. Relb. 356. Sibth. 31. Abbot. 206.
Engl. Bot. t. 579.

C. acuta. Hulst. 413. Lightf. 565.

Cyperoides aquaticum maximum, &c. Mich.
Gen. 57. n. 10, 11. t. 32. f. 6, 7.

Gramen cyperoides cum paniculis nigris. Raii
Syn. 417. Bauh. Hist. v. 2. 49+.

G. cyperoides majus latifolium, spicâ rufâ, caule
triangulo. Moris. sect. 8. t. 12. f. 1.

Angl. Great Common Carex.

In aquosis, et ad ripas fossarum fluviorumque
vulgaris.

Perennis. Fl. Aprili, Maio.

Radix valde repens, crassa. Herba atro-virens, parùm
glauca. Culmus tripedalis, erectus, firmus, acutè tri-
queter, angulis asperimis. Folia lata, erecta, margine
carinaque aspera. Bracteæ foliaceæ, culmum superan-
tes, brevissimè vaginatæ, plerumque auriculatæ, auricu-

lis rotundatis, pallidis. Spicæ masculæ tres aut quinque, erectæ, triquetræ, acutæ, nigro-fuscæ, glumis lan-
ceolatis, acuminatis, vel mucronatis, fuscis, carinâ con-
colori; fœmineæ tres aut quatuor, erectæ, plus minus
pedunculatæ, cylindricæ, apice saepe masculæ, glumis
ovatis, longius mucronatis, mucrone scabro, fuscæ, ca-
rinâ pallidiusculâ. Stigmata tria. Fructus imbricati,
patentiusculi, ovati, ventricosi, multinerves, brevius
rostrati, apice bifurei. Semen triquetrum, parvum,
albidum.

Micheli- 47. CAREX spicis erectis cylindricis: fœmineis
ana. pedunculatis, glumis omnibus obtusis muticis,
fructibus obovatis obtusissimis.

Carex Micheliana. *Tr. of Linn. Soc. v. 5. 270.*
Cyperoides foliis caryophylleis, caule exquisitè
triangulari, spicis habitioribus, squamis curtis
obtuse mucronatis, capsulis turbinatis brevibus
confertis. *Mich. Gen. 62. t. 32. f. 12.*

Angl. Blunt-fruited Black Carex.

In aquosis.

Near Aberdeen. *Prof. James Beattie, jun.*

Perennis. Fl. Maio?

Radix . . . Culmus crectus, pedalis aut sesquipedalis, tri-
queter, striatus, vix scaber. Folia erecta, acuminata,
marginibus aspera, carinâ leviusculâ. Bractæ folia-
ceæ, erectæ, culnum superantes, haud vaginatae, basi
auriculatæ, auriculis magnis, rotundatis, connatis, fuscis,
apice pallidis. Spicæ cylindricæ, obtusiusculæ.
erectæ; masculæ tres aut quatuor, subseffiles, graciles,
glumis obtusissimis, muticis, glauco-fuscis, margine
tenui, scarioso, niveo; fœmineæ duæ, longius pedun-
culatæ, crassiores, superior apice mascula, inferior basi
interdùm composita, glumis ellipticis, obtusissimis, mu-
ticis, fuscis, margine scarioso albo, carinâ flavescente
obtusâ. Stigmata tria. Fructus glumis multò brevi-
ores, obovato-triquetri, dilatati, serè encerves, glabri,
virescentes, apice obtusissimi, indivisi. Semen trique-
trum, breve, fuscum, angulis pallidis, ferè Caricis re-
curvæ,

curvæ, cui affinis est hæc species, sed differt spicis masculis numerosis, fructu glaberrimo, glumis breviore, compressiusculo, minimè gibbo.

48. CAREX spicis cylindricis: fœmineis pedunculatis, vaginis longissimis, glumis acuminate, fructibus triquetris rostratis bifurcatis.

Carex lœvigata. *Tr. of Linn. Soc. v. 5. 272.*
Angl. Smooth-stalked Beaked Carex.

In paludibus.

In a marsh near Glasgow, 1793. *Mr. J. Mackay.*
Marshes near Aberdeen. *Prof. J. Beattie, jun.*

Perennis. Fl. Maio.

Radix fibrosa. Culmus erectus, 2—4-pedalis, triqueter, lœvis, infernè foliosus. Folia ercta, latiuscula, lœte viridia, lœvia, longissimè vaginata; superiora margine aspera. Bracteæ foliaceæ, erectæ, culmum superantes, acuminatæ, acumine aspero, vaginis longissimis, pedunculo tamen brevioribus, supremis quandoque brevissimis et ferè nullis. Spicæ cylindraceæ, elongatæ, acutæ, graciles, ercta; masculæ spæniùs duæ, glumis lanceolatis, acutis, frequentiùs mueronulatis; fœmineæ duæ vel tres, longissimè pedunculatæ, pedunculis capillaribus, lœviusculis, glumis lanceolatis, mucronulatis, ferrugineis, dorso viridibus, mucrone scabro. Stigmata tria. Fructus glumis longiores, lanceolato-triquetri, ercto-patentes, nervosi, haud inflati, virides, glabri, in rostrum attenuati compressum, apice profundè bifurcum. Semen fructum implens, triquetrum, breve, fuscum, pedicellatum.

49. CAREX spicis fœmineis cylindricis abbrevia vesicatis subsessilibus, vaginis nullis, fructibus inflatis rostratis bifurcatis.

Carex vesicaria. *Sp. Pl. 1388, α et γ. Gooden.*
Tr. of Linn. Soc. v. 2. 205. With. 110.
Hull. 209. Sibth. 32. Fl. Dan. t. 647. Dicks.
H. Sicc.

*H. Sicc. fasc. 14. 18. Leers. 205. var. 3. t. 16.
f. 2. III.*

C. inflata. Huds. 412. Lightf. 567.

*Gramen cyperoides majus præcox, spicis turgidis teretibus flavescentibus. Dill. in Raii
Syn. 420.*

*G. cyperoides majus præcox, spicis turgidis teretibus flavescentibus. Moris. v. 3. 242. sect. 8.
t. 12. f. 6.*

Angl. Short-spiked Bladder Carex.

In palustribus.

In Wales and the North of England. Huds. In Breadalbane and other places. Lightf. About Oxford, but rarely. Bobart. By the water-works at Pimlico, and in other wet places about London.

Perennis. Fl. Maio.

Radix repens. Culmus erectus, bipedalis, triqueter, angulis acutissimis, asperis. Folia erēcta, pallidē viridia, angusta, acuminata, margine aspera. Bracteæ foliaceæ, culmum superantes, sessiles, vaginis nullis. Spicæ masculæ 1—3, filiformes, graciles, acutæ, glumiis linearī-lanceolatis, acutis, vel acutiusculis; fœmineæ tres aut quatuor, subsessiles, patentiusculæ, cylindricæ, breves (ferè unciales) obtusæ, glumis lanceolatis, vel linearī-lanceolatis, acutis, fuscis, dorso viridibus. Stigmata tria. Fruictus laxiusculè patentes, glumis longiores, ovati, acuminati, nervosi, inflati, glaberrimii, lutescentes, ore bifido. Semen parvum, triquetrum, breve, sessile, stylo longissimo mucronatum.

*ampulla-50. CAREX spicis fœmineis cylindricis elongatis
cea. subsessilibus, vaginis nullis, fructibus inflatis
rostro linearī bifurco.*

*Carex ampullacea. Gooden. Tr. of Linn. Soc.
v. 2. 207. With. 110. Hull. 209.*

C. ves-

- C. vesicaria.* *Huds.* 413. *Lightf.* 566.
C. rostrata. *With.* ed. 2. 1059. *Sibth.* 32.. *Abbot.* 206.
Gramen cyperoides polystachyon majus, spicis teretibus erectis. *Raii Syn.* 419.
G. cyperoides angustifolium, spicis erectis luteolis teretibus. *Moris. sect.* 8. *t.* 12. *f.* 8.
G. cyperoides angustifolium, spicis longis erectis. *Bauh. Theatr.* 84.
Angl. Slender-beaked Bladder Carex.

In paluſtribus et aquofis, haud vulgaris.

About Middleton in Warwickshire. *Ray.* Oxfordshire. *Dill.* *Sibth.* Rare in Bedfordshire. *Abbot.* Near Bungay. *Mr. Woodward.* Gloucestershire. *With.* At Virginia Water. *Rev. Dr. Goodenough.* Common in the north. *Curt.* Scotland. *Lightf.* and *Prof. James Beattie, jun.*

Perennis. Fl. Maio.

Radix repens. Culmus erectus, pedalis vel bipedalis, triquetus, angulis inferne glabris, supra spicam infimam scabris. Folia erecta, angusta, glauca, acuta, margine carinaque apicem versus scabra. Bracteae foliaceae, erectae, angustae, vaginis nullis, infima plerumque culmum superans. Spicæ masculæ duæ vel tres, remotiusculæ, filiformes, acutæ, glumis elliptico-lanceolatis, obtusis, fuscis, carinâ luteâ, margine scarioſo; fœmineæ duæ, rarius tres, erectæ, cylindricæ, duplo ferè longiores quam in præcedente multoque tenuiores, obtusæ, inferiores brevius pedunculatae, glumis muscularum, at ſæpe acutioribus. Stigmata tria. Fructus imbricato-patentes, glumis longiores, ferè globosi, inflati, nervosi, glabri, lutescentes, rostro linearis, apice bifido. Semen parvum, elliptico-triquetrum, ſeffile, stylo longissimo mucronatum.

51. CAREX pilosa, spicis abbreviatis cylindricis *birtu.*
remotis,

remotis, vaginis pedunculo subæqualibus, glumis aristatis, fructibus hirtis.

Carex hirta. *Sp. Pl.* 1389. *Gooden. Tr. of Linn. Soc. v. 2.* 208. *Huds.* 414. *With.* 111. *Hull.* 209. *Lightf.* 568. *Rehb.* 357. *Sibth.* 33. *Abbot.* 207. *Dicks. H. Sicc. fasc.* 11. 16. *Engl. Bot. t.* 685. *Leers.* 206. *t. 16. f. 3.* *Fl. Dan. t.* 379, quoad spicam e caule separatam.

Gramen cyperoides polystachyon lanuginosum.
Raii Syn. 418. *Morif. sect. 8. t. 12. f. 10.*
Angl. Hairy Carex.

In pratis humidis, aquosis et palustribus.

Perennis. Fl. Maio, Junio.

Radix latè repens, fibris villosissimis. Culmus erectus, bipedalis, foliosus, triquetus, angulis acutis, asperis. Folia culnum vix æquantia, erecta, plana, acuminata, subtus præcipue pilosa, margine aspera, vaginis apicem versus hirsutissimis. Bractæ spicarum fœminearum foliis omnino similes, culmo à quales, vaginis villosis pedunculum ferè totum continentibus. Spicæ masculæ duæ, remotiusculæ, magnitudine inæquales, lanceolatæ, glumis ovatis, mucronulatis, fuscis, villosis, carinâ luteâ; fœmineæ tres, remotæ, pedunculatæ, crectæ, cylindricæ, breves, laxiusculæ, glumis glabris, ovatis, longissimè aristatis, margine lato, scarioso, albo, dorso viridi. Fructus laxè imbricati, ovati, acuti, nervosi, undique hirti, apice produeto, bifurco. Semen obovato-triquetrum, stylò mucronatum.

In aquosis planta variat glabra, at vaginæ hinc ad apicem semper pilos quosdam rariores gerunt. D. Goodenough.

filiformis.

52. CAREX spicis ovatis, vaginis pedunculo subæqualibus, glumis mucronulatis, fructibus villosis, foliis canaliculatis glabris.

Carex filiformis. *Sp. Pl.* 1385. *Gooden. Tr. of Linn. Soc. v. 2.* 172. *t. 20. f. 5.* *With.* 97. *Hull.* 207. *Fl. Lapp. ed. 2.* 267.

C. angust-

- C. angustifolia. *Linn. MSS. in Sp. Pl. ed. 1.* 975.
 C. tomentosa. *Lightf.* 553. *Huds.* 650. *Herb.*
Lightf.
 C. hirta. *Fl. Dan.* t. 379, exclusâ spicâ seorsim
 delineatâ.
Angl. Slender-leaved Carex.

In palustribus rariū.

Plentifully at the south end of Air Links. *Dr. Hope.* In the Moss of Restenet, and other places in Scotland. *Mr. J. Mackay.* Near Eaton, Shropshire. *Rev. Mr. Williams.* Near Stoke, Norfolk. *Rev. Mr. Forby.*

Perennis. Fl. Junio.

Radix repens. Herba glabra. Culmus erectus, nudus, bipedalis, strictus, junceus, teretiusculus, lævis, apicem versùs sæpè triquierter, angulis scabris. Folia erecta, stricta, angusta, canaliculata, acuminata, ecarinata, lævia, margine scabra. Bracteæ foliis simillimæ, culmum superantes, breviūs vaginatæ, vaginis apice subauriculatis. Spicæ masculæ duæ (rariissimè solitariæ), remotiusculæ, magnitudine inæquales, lanceolatæ, glumis elliptico-oblongis, acutiusculis, nigricantibus, nervo lateo; fœmineæ duæ (rariūs solitariæ) remotiusculæ, ovatæ, erectæ, pedunculis vix vaginam excedentibus, glumis muscularum at sæpiùs brevissimè mucronulatis, glabris. Fruktus imbricati, ovati, nervosi, nigricantes, tomento brevi helvolo undique vestiti, apice latiusculo, bifurco. Semen elliptico-triquetrum, stylo brevi terminatum.

Icon apud Plukenetium, t. 34. f. 6, hanc potius quam præcedentem refert.

391. ERIOCaulon. G. Pl. 40. Juss. 44.
Gærtn. t. 83.

Cal. communis polyphyllus, multiflorus: *proprius* 2- vel 3-phyllos. *Corollulæ* 2- vel 3-partitæ. *Fl. Masc.* in medio disci. *Fœm.* in radio. *Stigmata*

mata 2 vel 3. Caps. 2- vel 3-cocca. Sem. solitaria.

Septan- 1. ERIOCAULON caule septangulari, foliis acuminatis cellulo-so-reticulatis, flosculis masculinis monopetalis tetrandris.

Eriocaulon septangulare. *With.* 184. *Sym.*
Syn. 41. *Engl. Bot.* t. 733.

E. decangulare. *Hull.* 29. *Lightf.* 569. *Penn.*
Voy. to the Hebrides, v. 1. t. 39. *Hope in Phil.*
Transf. v. 59. 243. t. 12.

Nasmythia articulata. *Huds.* 415.
Angl. Jointed Pipewort.

In lacubus insulæ Skyæ.

In a small lake in the isle of Skye, discovered there by Mr. Robertson. *Dr. Hope.* In two or three small fresh-water lakes about a mile west of Loch Sligachan, in the isle of Skye. *Lightf.* *Rev. Mr. Stuart and Mr. J. Mackay.* Plentifully in Loch-na-Caiplich in the same island. *Lightf.*

Perennis. Fl. Septembri.

Radix e fibris simplicibus, albidis, nitidis, artieulato-eellulosis. Herba tota congeries videtur tubularum, isthmis numerosis interceptarum. Caulis spithameus, erectus, implex, solitarius, aphyllus, tortus, septangularis, glaber, vaginâ tubulosâ ad basim. Folia radicalia, plurima, ensiformia, glabra, apice setacea. Capitulum terminale, solitarium, hemisphaericum, multiflorum, albo nigroque varium. Calycis communis foliola obovata, obtusa, scariosa, glabra. Flosculi nivei, apice extus ramentaceo-plumosi, intus nigrobiguttati; onines calyce 2—3-phyllo, obovato: masculini in disco plurimi, tetrandri, corolla monopetalâ, limbo bipartito, staminibus tubo insertis, antheris nigris: fœminei pauciores in radio, dipetali. Germina supera, didyma, stylo bipartito, stigmatibus acutis. Capsulæ dicoccæ. Semina solitaria.

IV. TETRANDRIA.

392. LITTORELLA. *Mant.* 160. *Juss.* 90.

Masc. *Cal.* tetraphyllus. *Cor.* monopetala, quadrifida. *Stam.* longissima.

Fœm. *Cal.* nullus. *Cor.* monopetala, inæquilaterè trifida. *Stylus* filiformis, longissimus. *Nux* unilocularis.

I. LITTORELLA.

lacustris.

Littorella lacustris. *Mant.* 295. *Huds.* 415.

With. 194. *Hull.* 210. *Relb.* 358. *Engl.*
Bot. t. 468. *Lightf.* 571. *Dicks.* H. *Sicc.*
fasc. 14. 14.

Plantago uniflora. *Sp. Pl.* 167. *Fl. Dan.* t. 170.

P. palustris gramineo folio monanthos, Parisiensis. *Raii Syn.* 316.

Gramen junceum minimum, capitulis longissimis filamentis donatis. *Moris. sect.* 8. t. 9. f. 30.

Angl. Plantain Shore-weed.

In inundatis arenofisis.

Perennis. Fl. Junio.

Radix fusiformis. Herba glabra, acaulis. Folia linearia, integerrima, subtùs convexa. Flores masculini pedunculati, solitarii, erecti, albidi, plantaginei; fœminei radicales, sessiles, stylo erecto, exerto, filiformi.

393. BETULA. *G. Pl.* 485. *Juss.* 409. *Gærtn.* t. 90.

Alnus. *Gærtn.* t. 90.

Masc. *Cal.* amenti squama monophylla, trifida, triflora. *Cor.* quadripartita.

Fœm. *Cal.* amenti squama monophylla, subtrifida, biflora. *Styli* duo. *Semina* compressa.

I. BE-

alba. 1. **BETULA** foliis ovatis acutis serratis glabriusculis.

Betula alba. *Sp. Pl.* 1393. *Huds.* 416. *With.* 206. *Hull.* 210. *Rehb.* 358. *Sibth.* 64. *Abbot.* 207. *Ait. H. Kew.* v. 3. 336.

Betula. *Raii Syn.* 443. *Baub. Hist.* v. 1. p. 2. 149. *Ger. em.* 1478. *Matth. Valgr.* v. 1. 128. *Camer. Epit.* 69. *Trag. Hist.* 1113.

Birch Tree. *Hunt. Evel. Sylv. tab.* ad p. 225.

♂. *Betula pendula.* *Gmel. Syst.* 282.

B. penduliflora virguliflora. *Loef. Pruss.* 26.

Angl. Common Birch. ♂ Weeping Birch.

In sylvis et sepibus umbrosis.

Arbor. Fl. Aprili, Maio.

Cortex niveus, epidermide chartaceâ. Rami viminei, tortuosi, leviusculi, apice pubescentes. Folia alterna, petiolata, ovato-deltoidea, inæqualiter ferrata, subtus subpubescentia, sero autumno aurea. Amenta cylindracea, pedunculata, cernua, squamis deciduis. Semina alata.

Ramuli in ♂ penduli, elongati, tuberculato-scabri, foliis omnino glabris.

nana. 2. **BETULA** foliis orbiculatis crenatis.

Betula nana. *Sp. Pl.* 1394. *Fl. Lapp.* ed. 2. 274. t. 6. f. 4. *Huds.* 416. *With.* 207. *Hull.* 210. *Lightf.* 575. t. 25. *Pallas. Ross.* v. 1. p. 1. 63. t. 40. f. D—G. *Fl. Dan.* t. 91? *Dicks. H. Sicc. fasc.* 8. 16.

Angl. Dwarf Birch.

In ericetis spongiosis montosis Scotiæ.

In Tweedale, found by Sir James Nasmyth, Bart. *Huds.* In Breadalbane, and Ross shire; also in the lowlands. *Lightf.*

Frutex. Fl. Maio.

Arbuscula tripedalis, erecta, ramosissima, rigida, ramulis sub-

subpubescentibus. Folia brevius petiolata, orbiculata, sæpè retusa, et ferè reniformia, inæqualiter crenata, utrinque glabra. Amenta pedunculata, lateralia, solitaria, erecta, ovato-cylindracea, obtusa, glabra.

3. BETULA pedunculis ramosis, foliis subrotundatis, cuneiformibus repandis serratis glutinosis : axillis venarum subtus villosis.
- Betula *Alnus*. *Sp. Pl.* 1394. *α. Huds.* 416.
With. 206. *Hull.* 210. *Relb.* 359. *Sibth.* 64.
Abbot. 207. *Ait. H. Kew.* v. 3. 338.
Alnus. Raii Syn. 442. *Ger. em.* 1477. *Matth.*
Valgr. v. 1. 127. *Camer. Epit.* 68. *Trag.*
Hist. 1084.
Angl. Common Alder. *Hunt. Evel. Sylv. tab.*
ad p. 240.

In aquosis.

Arbor. Fl. Martio.

Rami tortuosæ, glabri, patentes. Folia rotundata, obtusa, sæpè retusa, repanda, ferrata, viscidæ, saturatæ viridia, nitida, subtus pallidiora, venis rectis, parallelis, basi anticæ villosæ. Pedunculi terminales, raniosi. Amenta masculina pendula, elongata, cylindracea ; fœminea erectiuscula, ovata, obtusa, squamis persistentibus.

394. EUXUS. *G. Pl.* 486. *Juss.* 388. *Gærtn. t.* 108.

Masc. *Cal. triphyllus.* *Petala 2.* *Germinis rudimentum.*

Fœm. *Cal. tetraphyllus.* *Petala 3.* *Styli 3.* *Caps. trirostris, trilocularis.* *Sem. 2.*

1. BUXUS.

Buxus sempervirens. *Sp. Pl.* 1394. *Huds.* 417. *semper-*
With. 210. *Hull.* 210. *Wilden. Baumz.* 49. *virens.*

Buxus. Raii Syn. 445. *Ger. em.* 1410. *Hall.*

VOL. III.

H

Hist.

Hist. n. 1610. v. 2. 283. *Matth. Valgr.* v. 1.

172. *Camer. Epit.* 101. *Dalech. Hist.* 165.

B. B. angustifolia. *Dill. in Raii Syn.* 445.

Angl. Box-trec.

In collibus calcareis.

On Box hill, near Dorking, Surrey; Boxwell, Gloucestershire; and Boxley, Kent. *Ray.* In plenty on the chalk hills near Dunstable. *Mr. Woodward.*

Arbuscula. Fl. Aprili.

Arbor humilis, glabra, sempervirens, ligno duro, tenaci, ponderoso, flavescente. Folia opposita, subsessilia, elliptica, emarginata, integerrima, latitudine varia, lucida, hyeme sèpè miniata. Flores axillares, congesti, luteoli. Petala concava, staminibus breviora. Capsula globosa, virens, stylis crassis, persistentibus, divaricatis, coronata.

395. URTICA. *G. Pl.* 486. *Juss.* 403.
Gertn. t. 119.

Masc. *Cal.* tetraphyllus. *Cor.* nulla. *Germinis rudiumentum cyathiforme.*

Fœm. *Cal.* diphyllo. *Cor.* nulla. *Semen* 1, superum, nitidum.

*piluli-
fera.*

1. URTICA foliis oppositis ovatis serratis, floribus fœmineis capitatis.

Urtica pilulifera. *Sp. Pl.* 1395. *Huds.* 417.
With. 202. *Hull.* 210. *Engl. Bot.* t. 148.
Mill. Illustr. t. 79.

U. pilulifera, folio profundiùs Urticæ majoris in modum ferrato, semine magno lini. *Raii Syn.* 140.

U. romana. *Ger. em.* 706. *Fuchs. Hist.* 106.
Trag. Hist. 3.

U. prima.

U. prima. *Matth. Valgr. v. 2.* 469.

Angl. Roman Nettle. *Pet. H. Brit. t. 1. f. 11.*

In ruderatis maritimis.

At Yarmouth, Norfolk, and Aldborough, Suffolk.

Ray. About Lowestoft and Bungay, Suffolk.

Annua. Fl. Junio, Julio.

Herba ramosa, bipedalis, undique setis acerrimè venenatis hispida. Caulis cærulescens. Folia petiolata, ovata, vel subcordata, acuta, latè ferrata. Pedunculi axillares, plerumque gemini; quorum alter paniculatus flores masculinos sparsos, alter capitatus flores fœmineos in globum digestos, gerit. Calyx masculinorum patens, tetraphyllus. Stamina recurvato-patentia, rudimentum germinis circumcingentia. Cal. fœminorum diphyllos, concavus, clausus, extùs hispidus. Germen ovatum, superum. Stigma pubescens, sessile. Semen solitarium, nudum, fuscum, nitidum.

2. URTICA foliis oppositis ellipticis subquinque-urens: nervibus, racemis subsimplicibus.

Urtica urens. *Sp. Pl.* 1396. *Huds.* 417.

With. 202. *Hull.* 210. *Relb.* 359. *Sibth.* 62.

Abbot. 208. *Curt. Lond. fasc.* 6. *t. 70.* *Fl. Dan.* *t. 739.*

U. minor. *Raii Syn.* 140. *Ger. em.* 707.

Fuchs. Hist. 108. *Brunf. Herb.* *v. 1.* 154.

U. tertia. *Matth. Valgr. v. 2.* 471.

Angl. Small Nettle. *Pet. H. Brit. t. 1. f. 10.*

In cultis, vitium summum.

Annua. Fl. Junio—Octobri.

Radix fibrosa. Herba lètè virens, setis urentibus undique tecta. Folia elliptica, tri- vel quinque-nervia, scabra. Stipulæ parvæ, reflexæ. Racemi axillares, androgyni, subsimplices, petiolis plerumque breviores. Calyx fœmineus ovatus, compressiusculus.

dioica. 3. URTICA foliis oppositis cordatis, racemis ramosissimis geminis subdioicis.

Urtica dioica. *Sp. Pl.* 1396. *Huds.* 417.
With. 201. *Hull.* 210. *Rehb.* 359. *Sibth.* 62.
Abbat. 208. *Curt. Lond. fasc.* 6. *t. 69.* *Fl. Dan.* *t. 746.*

U. racemifera major perennis. *Raii Syn.* 139.

U. urens. *Ger. em.* 706.

U. major. *Fuchs. Hist.* 107. *Brunf. Herb.*
v. 1. 151.

U. secunda. *Matth. Valgr.* *v. 2. 470.*

Urtica, herba. *Pharmac. Lond.*

Angl. Great Nettle. *Pet. H. Brit.* *t. 1. f. 9.*

In ruderatis et ad scpes vulgaris.

Perennis. Fl. Julio, Augusto.

Radix repens. Caules erecti, tripedales. Herba pubescens, setisque urentibus vestita. Folia cordato-ovata, acuminata. Stipulae ovatae, acutae, erectae, membranaceae. Racemi gemini, ramosissimi, patentissimi, multiflori. Flores plerumque dioici, saepè monoici.

V. PENTANDRIA.

396. XANTHIUM. G. Pl. 487. Juss. 191.
Gærtn. t. 164.

Masc. *Cal.* communis imbricatus. *Cor.* monopetalæ, infundibuliformes, quinquefidæ. *Receptaculum* paleaceum.

Fœm. *Cal.* diphyllus, biflorus. *Cor.* nulla. *Drupa* sicca, muricata, bifida. *Nucleus* bilocularis.

I. XANTHIUM caule inermi, foliis cordatis basi *strumari-*
trinervibus.

Xanthium strumarium. Sp. Pl. 1400. Huds. 418. Wth. 283. Hull. 211. Fl. Dan. t. 970.

X. seu Lappa minor. Raii Syn. 140. Bauh. Hist. v. 3. 572. Matth. Valgr. v. 2. 545.

Xanthium. Fuchs. Hist. 579.

Bardana minor. Ger. em. 809.

Angl. Small Burdock. Pet. H. Brit. t. 1. f. 12.

In fimetis et pinguibus rariis.

Three miles from Portsmouth towards London,
and about Dulwich. Ray.

Annua. Fl. Augusto, Septembri.

Radix fibrosa. Caulis ramosus. Folia alterna, longius petiolata, cordata, lobata, ferrata, pubescentia, scabra. Racemi axillares, foliosi, pauciflori, superne masculi. Flores virides; masculini compositi, antheris discretis, nec syngenesiis. Fructus elliptici, pubescentes, undique muricati setis rigidis, uncinatis.

397. AMARANTHUS. *G. Pl.* 490. *Juss.* 88.
Gærtn. t. 128.

Masc. *Cal.* triphyllus. *Cor.* nulla. *Stam.* 3
five 5.

Fœm. *Cal.* triphyllus. *Cor.* nulla. *Styli* 3.
Caps. unilocularis, circumscissa. *Sem.* 1.

Blitum. 1. AMARANTHUS glomerulis lateralibus, floribus trifidis triandris, foliis ovatis, caule diffuso.

Amaranthus Blitum. *Sp. Pl.* 1405. *Huds.* 418.
With. 174. *Hull.* 211. *Relb.* 360.

Blitum rubrum minus. *Dill. in Raii Syn.* 157.
Camer. Epit. 235. *Bauß. Hist.* v. 2. 967.
Ger. em. 321.

Small Garden Blite. *Pet. H. Brit.* t. 7. f. 9.
Angl. Wild Amaranth.

In cultis et simeatis, haud vulgaris.

Battersea fields. *Mr. Dickson.* Ripton, Huntingdonshire. *Mr. Woodward.* Cambridge-shire. *Relb.*

Annua. Fl. Augusto.

Habitus Chenopodii seu Atriplicis. Caulis procumbens, ramosus, foliosus, latè diffusus. Folia alterna, longius petiolata, ovata, acuta, interdùm retusa, integerrima, margine scabriuscula, utrinque glabra. Racemi axillares, foliosi, floribus conglomeratis, viridibus, trifidis. Capsula elliptica, membranacea, stylis coronata. Semen nitidum, nigrum, lenticulare.

398. BRYONIA. *G. Pl.* 508. *Juss.* 394.
Gærtn. t. 88.

Masc. *Cal.* 5-dentatus. *Cor.* 5-sida. *Filam.* 3.
Anth. 5.

Fœm. *Cal.* 5-dentatus. *Cor.* 5-sida. *Stylus* 3-sidus.

3-fidus. *Bacca* infera, subglobosa, polyperma.

1. BRYONIA foliis palmatis utrinque calloso-*dioica*. scabris, floribus dioicis.
Bryonia dioica. *Jacq. Austr.* t. 199. *With.* 67.
Hull. 214. *Sibth.* 82. *Abbot.* 217. *Engl. Bot.*
t. 439.
B. alba. *Huds.* 437. *Reh.* 375. *Raii Syn.* 261.
Woodv. Med. Bot. t. 189. *Ger. em.* 869.
B. ruderalis. *Salisb. Prodri.* 158.
B. aspera sive *alba*, baccis rubris. *Bauh. Pin.*
297. *Mill. Ic.* t. 71.
Vitis alba sive *Bryonia*. *Matth. Valgr.* v. 2.
620. *Camer. Epit.* 987. *Fuchs. Hist.* 94.
Angl. Red-Berried Bryony.

In sepibus et dumetis vulgaris.

Perennis. Fl. Maio—Septembri.

Herba *dioica*; “adultior sèpè monoica.” *Miller*.

Radix magna, carnosa, alba, acris et cathartica. Caules annui, cirrhis scandentes, angulati, scabri, foliosi, subramosi, longissimi. Folia alterna, petiolata, palmata, quinqueloba, angulata, utrinque callis minutis scabra. Stipulae nullæ. Cirrhi solitarii, ad latera petiolorum, simplices. Pedunculi axillares, subumbellati, pauciflori. Corolla albida, venis virentibus elegantè picta. Antheræ per paria connatæ cum impari, omnes basi subconnexæ. Baccæ globosæ, coccineæ, glabræ. Semina sex, elliptica, maculata.

Folia autumno moschum redolent.

VI. POLYANDRIA.

399. CERATOPHYLLUM. *G. Pl.* 493.
Juss. 18. *Gærtn.* t. 44.

Masc. *Cal.* multipartitus. *Cor.* nulla. *Stam.*
 16—20.

Fœm. *Cal.* multipartitus. *Cor.* nulla. *Stigma*
 subsessile. *Sem.* 1, corticatum.

- demer-* 1. CERATOPHYLLUM fructibus trispinosis.
sum. Ceratophyllum demersum. *Sp. Pl.* 1409. *Huds.*
 419, α . *With.* 440. *Hut.* 211. *Gærtn.* v. 1.
 212. t. 44. f. ult. *Engl. Bot.* t. 947.
 C. n. 1599. *Hall. Hist.* v. 2. 275.
 Hydroceratophyllum folio aspero, quatuor cor-
 nibus armato. *Vaill. Aet. Paris. Ann.* 1719.
 t. 2. f. 1. *Raii Syn.* 135.
Angl. Common Hornwort.

In fossis et piscinis frequens.

Perennis. Fl. Augusto, Septembri.

Herba natans, aquâ immersa. Caulis ramosus, filiformis,
 foliosus. Folia verticillata, octona, dichotoma, laciinis
 saepius quatuor, linearibus, canaliculatis, dorso den-
 tatis, spinulosis. Flores axillares, solitarii, sessiles.
 Fructus ellipticus, teres, spinâ rectâ, longissimâ, termi-
 nali, duabusque basin versus brevioribus, divaricatis,
 longitudine maximè variantibus.

- submer-* 2. CERATOPHYLLUM fructibus muticis.
sum. Ceratophyllum submersum. *Sp. Pl.* 1409.
With. 441. *Hut.* 211. *Fl. Dan.* t. 510.
Engl. Bot. t. 679.
 C. demersum β . *Huds.* 419.
 Hydroceratophyllum folio laxi, octo cornibus ar-
 mato.

mato. *Vaill. Aet. Paris. Ann. 1719. t. 2. f. 2.*

Dill. in Raii Syn. 135.

Angl. Unarmed Hornwort.

In fossis rariūs.

In ditches by the road from Chichester to Selsey island. *Dill.* Between Yarmouth and Gorleston. *Mr. D. Turner.*

Perennis. Fl. Septembri.

Habitus prioris, sed foliis plerumque magis conpositis, tenuioribusque, et saepius laevibus. Præcipue differt fructu ovato, minori, omnino mutico et inermi, laciniosque calycinibus apice dentatis.

400. MYRIOPHYLLUM. *G. Pl. 493. Juss. 18.*
Gærtn. t. 68.

Masc. *Cal. tetraphyllum.* *Pet. 4. Stam. 8.*

Fœm. *Cal. tetraphyllum.* *Pet. 4. Stigmata 4,*
sessilia. *Sem. 4, corticata.*

1. MYRIOPHYLLUM floribus interruptè spi-*spicatum*, catis aphyllis.

Myriophyllum spicatum. *Sp. Pl. 1409. Huds. 419. With. 389. Hull. 212. Relb. 361. Sibth. 132. Abbot. 208. Fl. Dan. t. 681. Engl. Bot. t. 83.*

Potamogiton foliis pennatis. *Raii Syn. 150.*

Millefolium aquaticum pennatum spicatum.
Baub. Prodr. 73.

Feather Pondweed. *Pet. H. Brit. t. 6. f. 5.*

2. M. aquaticum pennatum minus, foliolis singularibus latiusculis flosculis subiectis donatum. *Dill. in Raii Syn. 151. Moris. Sect. 15. t. 4. f. 7.*

Fine Feather Pondweed. *Pet. H. Brit. t. 6. f. 6.*

Angl. Spiked Water-Millsoil.

In fossis et stagnis.

Perennis. Fl. Julio, Augusto.

Herba natans, immersa, spicis tantummodo exsertis. Caules flagelliformes, ramosi, supernè foliosi. Folia verticillata, quaterna, patentia, pectinata, glabra. Spicæ terminales, solitariæ, verticillatæ, bracteis ovatis, integris, quaternis, singulo verticillo subjectis. Petala in masculinis floribus calyce duplò longiora, concava, erecta, rubentia. Antheræ exsertæ, luteæ. Flores fœminei inferiores, petalis minoribus, stigmatibus villosis. Semina corticata, ovata.

verticil- 2. **MYRIOPHYLLUM** floribus omnibus axillatum.

Myriophyllum verticillatum. Sp. Pl. 1410.
Huds. 420. With. 390. Hull. 212. Relb. 361. Sibth. 132. Abbot. 209. Fl. Dan. t. 1046. Engl. Bot. t. 218.

M. aquaticum minus. Clus. Hist. v. 2. 252.

Pentapterophyllum aquaticum, flosculis ad foliorum nodos. Dill. Giss. 112. Append. 125. t. 7. Ruij Syn. 316.

Knotted Pondweed. Pet. H. Brit. t. 6. f. 8.

Angl. Verticillate Water-Millfoil.

In stagnis et piscinis.

Near Cambridge. Ray. At Bungay, Suffolk, and Hedenham, Norfolk. Mr. Woodward. At Lakenham, near Norwich. Mr. Crowe.

Perennis. Fl. Julio.

Herba natans, caules floriferi suprà aquam erecti, ad apicem usque foliosi. Folia quina, superiora minora. Flores axillares, verticillati; superiores masculini, numerosiores; inferiores fœminei; intermedii subindè hermaphroditici. Petala albida, parva. Stigmata crassa, apice plumosa.

401. SAGITTARIA. *G. Pl.* 494. *Juss.* 46.
Gærtn. t. 84.

Masc. *Cal.* triphyllus. *Pet.* 3. *Stamina* ferè 24.

Fœm. *Cal.* triphyllus. *Pet.* 3. *Pistilla* plurima.

Caps. plurimæ, ventricosæ, monospermæ.

1. SAGITTARIA foliis sagittatis acutis. *sagitti-*
Sagittaria sagittifolia. *Sp. Pl.* 1410. *Huds.* 420. *folia.*
With. 512. *Hull.* 212. *Relb.* 362. *Sibth.*
178. *Abbot.* 209. *Engl. Bot.* t. 84. *Dicks.*
H. Sicc. fasc. 15. 14. *Fl. Dan.* t. 172.
Sagitta. *Raii Syn.* 258. *Baub. Hist.* v. 3. p. 2.
779. *Camer. Epit.* 874.
S. minor. *Matth. Valgr.* v. 2. 483.
S. major et minor. *Ger. em.* 416.
Narrow Arrow-head. *Pet. H. Brit.* t. 43. f. 11.
β. Sagitta aquatica omnium minima. *Raii Syn.*
258.
Small Arrow-head. *Pet. H. Brit.* t. 43. f. 12.
Angl. Common Arrow-head.

In fluviis, fossis, et stagnis.

Perennis. Fl. Julio, Augusto.

Radix tuberosa. Herba glabra, laetescens. Scapus triquetus, erectus, supernè verticillatus, multiflorus. Folia radicalia, longè petiolata, sagittata, nervosa, integerima, lobis acutis, latitudine variis. Pedicelli terni, uniflori, bracteis ad basin solitariis, quandoque connatis. Flores superiores masculi, numerosiores; inferiores fœminei. Peta alba, basi violacea, decidua. Capsulæ in globum digestæ.

402. ARUM. *G. Pl.* 470. *Juss.* 24.
Gærtn. t. 84.

Spatha monophylla, basi convoluta. *Spadix* cylindricus, androgynus, suprà nudus, medio staminiferus,

staminiferus, basi germiniferus. *Baccæ uniloculares.*

macula- 1. ARUM acaule, soliis hastatis integerrimis, spatulum. ARUM acaule, foliis hastatis integerrimis, spatulo clavato obtuso.

Arum maculatum. *Sp. Pl.* 1370. *Huds.* 395. *With.* 497. *Hull.* 198. *Rehb.* 342. *Sibth.* 177. *Abbot.* 197. *Curt. Lond. fasc.* 2. t. 63. *Woodv. Med. Bot.* t. 25. *Mill. Illustr.* t. 76. *Fl. Dan.* t. 505.

Arum. *Raii Syn.* 266. *Riv. Monop. Irr.* t. 124. *Fuchs. Hist.* 69. *Matth. Valgr.* v. 1. 545. *Camer. Epit.* 365, 366.

A. vulgare. *Ger. em.* 834.

Arum, radix recens. *Pharmac. Lond.*
Angl. Cuckow-pint. Wake Robin.

In dumctis et ad sepes passim.

Perennis. Fl. Maio.

Radix tuberosa, acris, mucilaginosa. Herba acaulis, glabra, lucida. Folia petiolata, hastata, saepius nigromaculata. Scapus simplex, erectus. Spatha erecta, ventricosa, cucullata, acuta, pallide virens, subinde maculata. Spadix supernè clavatus, obtusus, frequenter purpureus; medium versus filamentis plurimis sterilibus, et paullò infrà antheris numerosis, sessilibus, armillatus; basi germinibus etiam numerosis, congestis, testus. Antheræ biporosæ. Stigmata simplicia, sessilia. *Baccæ miniatæ, succosæ.*

Flores nequaquam gynandri sunt. Locus in hâc classe sibi vindicat organis masculinis, respectu fœminorum, supernè aut internè positis, ut in plantis monoicis floribusque compositis universis ferè conspicendum est, nec externe aut infernè, ut in omni flore simplici.

403. PTERIUM. *G. Pl.* 495. *Juss.* 336.
Pimpinella. *Gærtn.* t. 32.

Masc. *Cal.* triphyllus. *Cor.* quadripartita. *Stam.* 30—40. *Fœm.*

Fœm. *Cal. triphyllus. Cor. quadripartita. Pist. 1*
five 2. Nux bilocularis, tubo corollæ indurato
corticata.

i. PTERIUM inerme, caulis subangulosis. *Sanguis-*
Poterium Sanguisorba. Sp. Pl. 1411. Hud. forba.

421. With. 493. Hull. 212. Relb. 362.
Sibth. 168. Abbot. 209. Curt. Lond. fasc. 2.
t. 64. Mart. Fl. Rust. t. 69. Dicks. H. Sicc.
fasc. 9. 15. Engl. Bot. t. 860.

Sanguisorba minor. Raii Syn. 203. Baub. Hist.
v. 3. p. 2. 115. Fuchs. Hist. 789.

Pimpinella hortensis. Ger. em. 1045.

P. five Sanguisorba minor. Matth. Valgr. v. 2.
381.

Angl. Common Burnet.

In montosis calcareis copiosè.

Perennis. Fl. Julio.

Radix lignosa. Caules ramosi, foliosi, angulati, multi-
flori, glabri. Folia impari-pinnata, foliolis elliptico-
subrotundis, ferratis, glabris, glaucescentibus. Pedun-
culi terminales, elongati, nudi. Capitula solitaria,
multiflora. Flores inferiores masculi, quod singulare,
staminibus longissimis, capillaribus, purpureis, depen-
dentibus, rarius pistillo abortivo muniti; superiores
fœminei, stigmate radiato, purpureo, exerto. Fructus
rugosus.

404. QUERCUS. *G. Pl. 495. Juss. 410.*
Gærtn. t. 37.

Masc. *Cal. campanulatus, lobatus. Cor. nulla.*
Stam. 5—10.

Fœm. *Cal. campanulatus, integerrimus, scaber.*
Cor. nulla. Stylis 1. Stigmata 3. Nux supera,
coriacea, monosperma.

i. QUER-

- Robur.* 1. **QUERCUS** foliis deciduis oblongis supernè latioribus : sinubus acutioribus : angulis obtusis, pedunculis fructiferis elongatis.
- Quercus Robur.** *Sp. Pl.* 1414. *Huds.* 421, α .
Relb. 363. *Sibth.* 133. *Woodv. Med. Bot.* t. 126.
Mart. Fl. Rust. t. 10. var. *pedunculata*.
- Q. longæva.** *Salisb. Prodr.* 392.
- Q. fœmina.** *With.* 387. *Fl. Dan.* t. 1180.
- Q. pedunculata.** *Hull.* 212. *Abbot.* 210. *Wilden. Baumz.* 278.
- Q. latifolia.** *Raii Syn.* 440.
- Q. hemeris.** *Dalech. Hist.* 4.
- Q. cum longo pediculo.** *Bauh. Pin.* 420.
- Quercus.** *Matth. Valgr.* v. 1. 184. *Duham. Arb.* v. 2. t. 47. *Tabern. Kreuterb.* 1374.
- Oak Tree.** *Hunt. Evel. Sylv. tab.* ad p. 69.
- Quercus, cortex.** *Pharmac. Lond.*
- Angl. Common British Oak.**

In sylvis et sepibus ubique.

Arbor. Fl. Aprili.

Arbor formosa, ligno utilissimo, duro, tenaci. Folia alterna, subsessilia, obovato-oblonga, sinuata, lobis obtusis ; suprà lævia, nitida ; subtùs pallidiora, subglauca. Racemi axillares ; masculini laxi, penduli, multiflori, flavi ; fœminei longius pedunculati, subtriflori. Calyx in flore masculino membranaceus, campanulatus, saepius quinquefidus ; in fœmineo coriaceus, squamosus, pubescens, demum hemisphæricus, integerrimus, ligneus. Stamina circiter decem, calyce longiora. Germen ovatum. Stylus brevis, cylindraceus. Stigmata tria. Nux elliptica, coriacea, lævis, calyci insidens, demum decidua.

*sessili-
flora.*

2. **QUERCUS** foliis petiolatis deciduis oblongis : sinubus acutioribus oppositis, fructibus sessilibus.

Quercus

- Quercus sessiliflora. *Salisb. Prodr.* 392.
 Q. Robur. *With.* 387. *Hull.* 212. *Abbot.*
 210. *Wilden. Baumz.* 277. *Huds.* 421, β.
Mart. Fl. Rust. t. 11, var. *sessilis*.
 Q. latifolia mas, quæ brevi pediculo est. *Raii*
Syn. 440.
 Q. platyphyllos mas et fœmina. *Dalech.*
Hist. 2, 3.
 β. Durmäst Oak. *Mart. Fl. Rust.* t. 12.
Angl. Seffile-fruited Oak.

In sylvis.

Bagley wood, and about Newberry. *Ray.* About Norwood in Surrey. *Martyn.* Common, but less so than the preceding. *Mr. Woodward.* In many parts of Norfolk, also about London.
 β. In the New Forest. *Martyn.*

Arbor. Fl. Aprili, Maio.

A priore differt ligno minoris pretii, foliis petiolatis, magis æqualiter ac regulariter pinnatifidis; in β subtus pubescens; floribus fœmineis fructibusque omnino ferè sessilibus.

405. FAGUS. *G. Pl.* 496. *Juss.* 409.

Gærtn. t. 37.

Castanea. *Gærtn.* t. 37.

Masc. *Cal.* campanulatus, quinquefidus. *Cor.* nulla. *Stam.* 5—12.

Fœm. *Cal.* quadrifidus. *Cor.* nulla. *Styli* 2—3, trifidi. *Sem.* 2 vel 3, calyce coriaceo, muri- cato, tecta.

1. FAGUS foliis lanceolatis acuminato-ferratis *Casta-* subtilis nudis, aculeis fructus compositis im- *nea.* plexis.

Fagus Castanea. *Sp. Pl.* 1416. *Huds.* 422.
With.

IWith. 443. *Hull.* 213. *Abbot.* 211. *Engl.*
Bot. t. 886. *Wilden. Baumz.* 111.
Castanca. *Raii Syn.* 440. *Baub. Hist.* v. 1.
p. 2. 121. *Matth. Valgr.* v. 1. 191. *Camer.*
Epit. 118. *Dubam. Arb.* v. 1. t. 50. *Ger.*
em. 1442. *Mill. Ic.* t. 84.
Angl. Chesnut tree. *Hunt. Evel. Sylv.* tab. ad
p. 159.

In sylvis.

Arbor. Fl. Maio.

Arbor vasta, longæva, eortice latè rimoso. Folia alterna,
 breviùs petiolata, lanceolato-oblonga, acuta, ferrata,
 glabra, deeidua, venis numerosis, parallelis, ferraturis
 acuminatis. Raeemi moniliformes, multiflori, eernui;
 superiores terminales, fœminei; inferiores numero-
 fiore, axillares, solitarii, masculini. Flores sessiles, in
 glomerulis alternis digesti. Calyees pilosi. Stamina
 capillaria, exserta, circiter duodecim. Stigmata su-
 bulata, rigida, glabra. Calyx fructus globosus, mag-
 nus, undique murieatus, aculeis stellato-ramosis, im-
 plexis. Semina plerunque duo, badia, glabra, hinc
 complanata, extus gibba, esculenta.

sylvatica. 2. *FAGUS* foliis ovatis obsoletè serratis, aculeis
 fructus simplicibus.

Fagus sylvatica. *Sp. Pl.* 1416. *Huds.* 422.

IWith. 444. *Hull.* 213. *Abbot.* 210. *Wil-*
den. Baumz. 113.

Fagus. *Raii Syn.* 439. *Matth. Valgr.* v. 1.
185. *Camer. Epit.* 112. *Baub. Hist.* v. 1.
p. 2. 117. *Ger. em.* 1444.

Angl. Beech tree. *Hunt. Evel. Sylv.* tab. ad
p. 136.

In sylvis et sepibus solo calcareo.

Arbor. Fl. Aprili, Maio.

Arbor

Arbor patula, formosa, cortice lœvi, æquali. Folia alterna, petiolata, latè ovata, vel subrhombéa, acuta, undulata, vix ferrata, nitida, ciliata, marcescentia, venis paucis, rectis, in axillis præcipue pilosis. Capitula axillaria et terminalia, pedunculata, villosa; superiora masculina, multiflora, bracteolata. Stamina 5 vel 6. Calyx fructus ovatus, sericeus, undique mureatus, aculeis simplicibus, mollibus, recurvis. Semina fusca, acutè triquetra, acuminata.

406. CARPINUS. G. Pl. 497. Juss. 409.
Gærtn. t. 89.

Masc. Cal. amenti squama subrotunda. Cor. nulla. Stam. 8—20.

Fœm. Cal. amenti squama oblonga. Cor. nulla. Germinalia 2. Styli singulis 2. Nux angulosa, unilocularis.

1. CARPINUS squamis strobilorum planis oblongis dentatis. *Betulus.*

Carpinus Betulus. Sp. Pl. 1416. Huds. 422. With. 442. Hull. 213. Relb. 363. Wilden. Baumz. 52.

Carpinus. Matth. Valgr. v. 1. 131. Camer. Epit. 71.

Osiyra Ulmo similis, fructu in umbilicis foliaceis. Raii Syn. 451.

Betulus sive Carpinius. Ger. em. 1479. Angl. Horn-beam.

In sylvis et sepibus, solo tenaci, sterili.

Arbor. Fl. Maio.

Arbor facie Ulmi, cortice lœvi, albido. Folia alterna, petiolata, ovata, acuminata, duplicato-ferrata, glabra, venis rectissimis, parallelis, ad axillas villosis. Amenta masculina lateralia, squamis ferrugineis, concavis, ciliatis, staminibus 8—16 (Linn.); fœminea terminalia, solitaria, laxa, squamis linearis-elongatis, planis, recurvis,

curvis, ciliatis, germinibus geminis, setosis, digynis.
Nuces solitariæ, ovatæ, compressæ, angulatæ, in axillis
squamarum persistentium, amplissimarum, ferratarum.

407. CORYLUS. *G. Pl.* 498. *Juss.* 410.
Gærtn. t. 89.

Masc. *Cal.* amenti squama trifida. *Cor.* nulla.
Stam. 8.

Fœm. *Cal.* bifidus, lacer. *Cor.* nulla. *Styli* 2.
Nux ovata, lævis, unilocularis, calyce ventri-
coso, coriaceo, tecta.

Avella-na. 1. **CORYLUS** stipulis ovatis obtusis, foliis subro-
tundis cordatis acuminatis, ramulis pilosis.

Corylus Avellana. *Sp. Pl.* 1417. *Ait. H. Kew.*
v. 3. 363. *Huds.* 423. *Witb.* 378. *Hull.* 213.
Relb. 364. *Sibtb.* 127. *Abbot.* 211. *Engl.*
Bot. t. 723. *Wilden. Baumz.* 79.

C. sylvestris. *Raii Syn.* 439. *Ger. em.* 1438.

Nux Avellana. *Matth. Valgr.* v. 1. 255. *Ca-*
mer. *Epit.* 177. *Trag. Hist.* 1096. *Fuchs.*
Hist. 398.

Angl. Hasel-nut tree. *Hunt. Evel. Sylv. tab. ad*
p. 220.

In sylvis et sepibus vulgaris.

Arbor. Fl. Martio, Aprili.

Arbor humilis, ramosa, ramulis piloso-scabris. Folia
alterna, breviūs petiolata, rotundata, acuminata, du-
plicato-ferrata, basi cordata et angustata, subtūs præ-
cipue pubescentia, omnia omnino post flores expansa.
Amenta mascula terminalia, subpaniculata, cylindra-
cea, pendula, tremula, multiiflora, flavescentia, squamis
extūs pubescentibus, receptaculo communi infixis, nec
deciduis. Flores fœminei pauci intra singulas gemmas
alternas, laterales, imbricatas, ovatas. Styli sanguinei,
exserti. Calyx exiguus, demùm ampliatus, coriaceus,
bilobus, apice lacer, nucem ovatam, lævem, duram,
esculentam, basi amplectens.

VII. MONADELPHIA.

408. PINUS. *G. Pl.* 499. *Juss.* 414.
Gærtn. t. 91.

Masc. *Cal.* amenti squama peltata. *Cor.* nulla.

Antheræ squamis adnatæ, sessiles.

Fœm. *Cal.* amenti squama biflora. *Cor.* nulla.

Nux unilocularis, alata.

1. PINUS foliis geminis rigidis, strobilis juniori-*sylvestris*.
bus pedunculatis recurvis, antherarum cristâ
exiguâ.

Pinus sylvestris. *Sp. Pl.* 1418. *Ait. H. Kew.*
v. 3. 366. *Huds.* 423. *With.* 615. *Hull.*
213. *Lightf.* 587. *Mill. Illustr.* t. 82. *Pall.*
Ross. v. 1. 5. t. 2. f. II. *Ger. em.* 1356. *Dalech.* *Hift.* 44. *Woodv. Med. Bot.* t. 207. *Wil-*
den. Baumz. 205. *Lambert. Pin.* 1. t. 1.

P. sylvestris, foliis brevibus glaucis, conis parvis
albentibus. *Raii Syn.* 442. *Dubam. Arb.*
v. 2. 125. t. 30.

P. sylvestris montana. *Matth. Valgr.* v. 1. 89.
Angl. Scotch Fir. *Hunt. Evel. Sylv.* tab. ad
p. 266.

In alpibus Scoticis copiosè. *Lightf.*

Arbor. Fl. Maio.

Arbor excelsa, recta, determinatè ramosa, semper, at
tristè, virens. Folia ex singulis geminis membranaceis,
ordine quincunciali digestis, bina, recta, linearia, acuta,
rigida, glabra, supra canaliculata, subtùs convexa.

Amenta masculina densè spicata, bracteata, pedicellata, elliptica, obtusa, fusco-sulphurea, squamis numerosis, suprà cristatis, subtùs antheram sessilem, bivalvem, gerentibus; fœminea inferiora, sæpiùs terna, pedunculata; mox cernua, ovata, viridia, glabra, demùm mutata in strobilos ligneos, tessellatos, tuberculatos, semina duo, alata, intùs ad singulam squamam gerentes.

Hæc sola hujus generis species patriæ nostræ indigena est.

Classis XXII.

DIOECIA.

Flores masculi et fœminei in diversis
plantis.

MONANDRIA.

Stamen 1.

Salix purpurea et Helix.

DIANDRIA.

Stamina 2.

409. SÁLIX. Masc. *Cal.* amenti squama. *Cor.*
nulla. *Glandula* baseos nectarifera. *Stam.*
1—5.

Fœm. *Cal.* amenti squama. *Cor.* nulla. *Stig-*
mata 2. *Caps.* supera, unilocularis, bivalvis.
Sem. papposa.

TRIANDRIA.

Stamina 3.

- 410. EMPETRUM.** Masc. *Cal.* tripartitus:
Pet. 3. *Stam.* capillaria, 3—9.
 Fœm. *Cal.* tripartitus. *Pet.* 3. *Stigmata* 9.
Bacca supera, enneasperma.
- 411. RUSCUS.** Masc. *Cal.* hexaphyllus. *Cor.*
 nulla. *Nectarium* ovatum, tubulosum, intus
 flaminiferum.
 Fœm. *Cal.* *Cor.* et *Nect.* maris. *Stam.* nulla.
Stylus 1. *Bacca* supera, trilocularis. *Sem.* ge-
 mina.

- *Valeriana dioica.* *Salix triandra et amygdalina.*

TETRANDRIA.

Stamina 4.

- 413. HIPPOPHAE.** Masc. *Cal.* bipartitus. *Cor.*
 nulla.
 Fœm. *Cal.* tubulosus, bifidus. *Stylus* 1. *Bacca*
 supera, monosperma. *Semen* arillo dupli-
 tectum.
- 414. MYRICA.** Masc. *Cal.* amenti squama con-
 cava. *Cor.* nulla.
 Fœm. *Cal.* amenti squama concava. *Cor.* nulla.
Styli 2. *Bacca* monosperma.

412. VIS-

412. VISCUM. Masc. *Cal.* nullus. *Pet.* 4, basi dilatata, connata, calycina. *Antheræ* sessiles, petalis adnatæ.

Fœm. *Cal.* submarginatus. *Pet.* 4, basi dilatata. *Stylus* nullus. *Bacca* infera, mono-sperma.

Rhamnus catharticus. *Urtica dioica.*

PENTANDRIA.

Stamina 5.

415. HUMULUS. Masc. *Cal.* pentaphyllus.
Cor. nulla. *Antb.* apice biporosæ.

Fœm. *Cal.* amenti squama obliqua, *integra*.
Cor. nulla. *Styli* 2. *Semen* 1, tunicatum, *calyce* alatum.

Ribes alpinum. *Pimpinella dioica.*
Bryonia dioica. *Salix pentandra.*

HEXANDRIA.

Stamina 6.

416. TAMUS. Masc. *Cal.* sexpartitus. *Cor.* nulla.

Fœm. *Cal.* sexpartitus. *Cor.* nulla. *Stylus* trifidus. *Bacca* infera, trilocularis. *Sem.* bina.

Rumex Acetosa et Acetosella.

OCTANDRIA.

Stamina 8.

418. RHODIOLA. Masc. *Cal.* quadripartitus.
Pet. 4. *Nectaria* 4, emarginata.
 Fœm. *Cal.* quadripartitus. *Pet.* 4. *Nect.* 4,
 emarginata. *Pistilla* 4. *Caps.* 4, poly-
 spermæ.
417. POPULUS. Masc. *Cal.* amenti squama
 lacera. *Cor.* turbinata, obliqua, integra.
 Fœm. *Cal.* amenti squama lacera. *Cor.* turbi-
 nata, integra. *Stigmata* 4. *Caps.* supera, bi-
 locularis, bivalvis. *Sem.* pappoia.

ENNEANDRIA.

Stamina 9.

419. MERCURIALIS. Masc. *Cal.* tripartitus.
Cor. nulla. *Stam.* 9—12. *Anth.* globosæ, di-
 dymæ.
 Fœm. *Cal.* tripartitus. *Cor.* nulla. *Styli* 2.
Caps. dicocca, bilocularis. *Sem.* solitaria.

420. HY-

420. HYDROCHARIS. Masc. *Cal.* trifidus.
Pet. 3. *Filamenta* tria interiora appendicu-
lata.

Fœm. *Cal.* trifidus. *Pet.* 3. *Styli* 6. *Caps.*
sexlocularis, polysperma, infera.

Empetrum nigrum.

DECANDRIA.

Stamina 10.

Silene Otites. *Lychnis dioica.*

ICOSANDRIA.

Stamina plura, calyci inserta.

Rubus Chamæmorus.

POLYANDRIA.

Stamina plura, receptaculo inserta.

Stratiotes aloides. *Populus nigra.*

MONA-

MONADELPHIA.

Filamenta infernè in unum corpus
coalita.

421. JUNIPERUS. Masc. *Cal.* amenti squamæ.
Cor. nulla. *Stam.* 3.

Fœm. *Cal.* amenti squamæ pauciores, demùm
carnosæ, coalitæ in *baccam* trispermam.

422. TAXUS. Masc. *Cal.* nullus. *Cor.* nulla.
Stam. plurima. *Anth.* peltatæ, octofidæ.

Fœm. *Cal.* urceolatus, integerrimus. *Stylus*
nullus. *Sem.* 1, calyci baccato impositum.

Salix fissa, rubra?, et *Croweana*.

XXII. DIOECIA.

I. DIANDRIA.

409. SALIX. G. Pl. 514. Juss. 408.
Gærtn. t. 90.

Masc. *Cal.* amenti squama. *Cor.* nulla. *Glandula* baseos nectarifera. *Stam.* 1—5.

Fœm. *Cal.* amenti squama. *Cor.* nulla. *Stigmata* duo. *Caps.* supera, unilocularis, bivalvis. *Sem.* papposa.

* *Foliis serratis glabriusculis.*

1. SALIX monandra decumbens, foliis obovato-*purpurea*, lanceolatis serratis glabris, stigmatibus brevissimis ovatis subsessilibus.

Salix purpurea. Sp. Pl. 1444. Huds. 427.
Relb. 366. Tr. of Linn. Soc. v. 6. 113.

S. monandra. Arduin. Mem. 1. 67. t. 11. Ehrb.
Arb. 58.

Angl. Bitter purple Willow.

In palustribus et ad fluvios, rariūs.

In King-street meadows, Norwich. Mr. Crowe.

Frutex. Floret Martio.

Frutex tri- aut quadri-pedalis, decumbens. Rami viminei, adscendentes, tenacissimi, teretes, glaberrimi, lucidi, sœpè purpurascentes. Folia opposita vel alterna, breviūs petiolata, obovato-lanceolata, vel sublinearia, apicem

apicem versùs præcipue ferrata, breviùs acuminata, glabra, subtùs glauca; gustui valde amara. Stipulae nullæ. Amenta mascula alterna vel opposita, solitaria, subsessilia, basi bracteata, cylindracea, vix uncialia, gracillima, obtusa, multiflora, supernè primò florentia. Squamæ apice nigricantes, pilosæ. Glandulæ nectariæ solitariæ, squamis oppositæ. Stamen unicum. Anthera gemina, seu quadriloba, fulva. Amenta fœminea in distincta planta, masculis similia. Germen sessile, parvum, ovato-ellipticum, sericeum. Stylus brevissimus, ut ferè nullus. Stigmata parva, sessilia, subovata, indivisa, sulco supernè exarata. Capsula ovata, parva, sericea.

Helix. 2. *SALIX monandra* erecta, foliis lanceolatis acuminatis ferrulatis glabris, stylo elongato filiformi, stigmatibus linearibus.

Salix Helix. *Sp. Pl.* 1444. *Huds.* 427. *Relh.* 366. *Dalech. Hist.* 277. *f. 2.* *Tr. of Linn. Soc. v. 6.* 114.

S. monandra. *With.* 45. *Hull.* 218. *Sibth.* 16. *Abbot.* 212. *Curt. Lond. fasc.* 6. *t. 71.* *Hoffm.* *Sal.* *v. 1.* 18. *t. 1.* *f. 1,* 2. *t. 5.* *f. 1.* *t. 23.* *f. 1.* *Villars.* *Dauph.* *v. 3.* 767.

S. humilior, foliis angustis subcæruleis, ex adverso binis. *Raii Syn.* 448. *Cant.* 144. *n.* 5.

S. n. 1640. *Hall. Hist.* *v. 2.* 306.

Salicis racemi seu nucamenta, rosæ et capitula squamata. *Baub. Hist.* *v. 1.* *p. 2.* 213. *Angl.* Rose Willow.

In salicetis et palustribus.

Arbor. Fl. Martio, Aprili.

Arbor gracilis, erecta, humilis, vix decempedalis. Ramæ viminei, teretes, tenaces, glaberrimi, cinerei aut purpureascentes, nitidi. Folia opposita vel alterna, petiolata, linear-lanceolata, acuminata, utrinque attenuata, apicem versùs ferrulata, glabra, subtùs glaucescentia, gustui minùs

minus amara. Stipulae nullae. Amenta mascula nondum vidi: staminum characterem ex auctorum descriptione, confirmante D. Crowe, mutuatus sum. Amenta foeminea longiora et duplo crassiora quam in praecedente, longiusque pedunculata, squamis oblongis. Germen sessile, ovatum, sericeum. Stylus elongatus, filiformis, glabratus. Stigmata ovato-linearia, elongata, patentia, sulcata, emarginata. Capsula ovata, lanuginosa, albida.

Amentorum crassitudine, caule erecto arboreo, at praecipue styli atque stigmatum formâ, a priore satis discrepat, monente D. Crowe.

3. **SALIX monandra erecta, foliis obovato-lanceo-** *Lambertiana*.
latis acutis serratis glabris, stigmatibus brevisimis ovatis emarginatis.

Angl. Boyton Willow.

In salicetis et palustribus.

On the banks of the river Willy at Boyton, Wiltshire, for 16 miles; also in osier-holts near Stains. *Mr. Lambert.* About Lackford bridge near Icklingham, Suffolk. *Mr. Crowe.*

Arbor. Fl. Martio, Aprili.

Magnitudo et forma praecedentis. Folia subopposita vel alterna, breviora quam in praecedente, (vix sesquicentaria,) breviusque petiolata, obovato-lanceolata, acuminate, apicem versus ferrata, glabra, subtus glaucescentia. Stipulae nullae. Amenta minora quam in S. Helice, squamis orbiculatis, nigris. Germen sessile, ovato-ellipticum, crassum, sericeum. Stylus brevissimus. Stigmata parva, ovata, patentia, sulcata, emarginata.

4. **SALIX monandra erecta, foliis substipulatis** *Forbyana*.
lanceolatis acutis denticulatis glabris subtus *ana.*
glaucis, stigmatibus linearibus.

Salix fissa. Tr. of Linn. Soc. v. 6. 115, excluso
synonymo.

Angl. Basket Osier.

In

In salicetis.

At Fincham, Norfolk. *Rev. Joseph Forby.* In several osier-grounds near Lynn, Norfolk. *Mr. Crowe.* At Prickwillow, near Ely. *Rev. Mr. Hemsted.*

Frutex. Fl. Aprili.

Rami viminei, erecti, flexiles et valde tenaces, glabri, cinereo-lutei, vel purpurascentes. Folia alterna, rarius subopposita, petiolata, lanceolata, acuminaia, bi- vel triuncialia, apicem versus præcipue denticulata vel subserrata, utrinque glaberrima; suprà saturatè viridia; subtùs glauca; juniora parùm pubescentia. Stipulæ parvæ, lanceolatæ, acutæ, rarius obviae. Amenta breviùs pedunculata, magnitudine et formâ S. *Helicis.* Squamæ obovatæ, nigræ, quandoque emarginatæ. Germen sessile, ovatum, sericeum. Stylus elongatus. Stigmata ovato-linearia, emarginata, patentia.

Flosculum unum vel alterum, masculinum, monandrum, ad basin amentorum fœmineorum, quandoque inveni.

Ad corbes concinniores texendas præstantissima est hæc species. *D. Crowe.*

zulra. 5. *SALIX monadelpha?* foliis linear-lanceolatis elongatis acutis denticulatis glabris: subtùs concoloribus.

Salix rubra. *Huds.* 428. *With.* 49. *Hull.* 218.

Relb. Suppl. 3. 8. *Tr. of Linn. Soc.* v. 6. 116.

Engl. Bot. t. 1145.

S. virescens. *Villars.* *Dauph.* v. 3. 785. t. 51. f. 30.

S. fistla. *Ehrb. Arb.* 29.

S. minimè fragilis, foliis longissimis utrinque viridibus non serratis. *Raii Syn.* 449.

S. nerii folio utrinque vidente. *Vaill. Paris.* 175. *Herb. Sherard.*

Angl. Green Osier.

In

In salicetis rariūs.

Between Maidenhead and Windsor, and near Salisbury. *J. Sherard.* In an osier-holt near Ely. *Rev. Dr. Goodenough.* At Prickwillow, near Ely. *Rev. Mr. Hemsted.* In Cambridgeshire. *Relhan.* Near Bedford. *Abbot.*

Frutex. Fl. Aprili, Maio.

Rami viminei, longissimi, tenaces, glabri, cinerei, vel purpurascens. Folia alterna, petiolata, linear-lanceolata, acuta, quadriuncialia, denticulata vel ferrulata, nequamquam integerrima, utrinque læte viridia, glabra, vel subtus rariūs obsoletè pilosa. Stipulæ linear-lanceolatæ, denticulatæ; sæpiūs verò nullæ. Amenta masculina nondam yidi: fœminea ferè ut in præcedente, sed stigmatibus aliquantulam crassioribus et ferè ovatis.

Differ a *S. Lambertianâ* foliis longioribus, duplòque angustioribus, figurâ ferè *S. viminalis*, at utrinque læte viridibus, neque glaucis neque incanis. Folia juniora reverè pubescunt ut in aliis speciebus.

6. *SALIX monadelpha*, foliis ellipticis subserratis *Croweana*.
glaberrimis: subtus glaucis.

Salix Croweana. *Tr. of Linn. Soc. v. 6. 117.*
Engl. Bot. t. 1146.

Angl. Broad-leaved Monadelphous Willow.

In palustribus.

At Cranberry Fen, in the parish of East Winch,
and in other parts of Norfolk. *Mr. Crowe.*

Arbuscula. Fl. Aprili, Maio.

Rami brevès, patuli, fragiliores, luteoli vel purpurascens, nitidi. Folia alterna, petiolata, patentia, vix sesquiuncialia, elliptica, vel obovato-elliptica, acutiuscula, obsoletè ferrata, vel subcrenata, utrinque glaberrima; suprà læte viridia, nitida; subtus glauca, venosa. Amenta subsessilia, ovato-cylindracea, brevia. Squamæ obovatæ,

obovatæ, nigræ, pilosissimæ. Stamina citrina, elongata.
filamentis plus minus connatis, antheris rubicundis.
Flores fœmineis adhuc desiderantur.

triandra. 7. **SALIX triandra**, foliis linear-i-oblongis serratis
glabris, germinibus pedicellatis.

Salix triandra. *Sp. Pl.* 1442. *Huds.* 425. *With.* 45.

Hull. 217. *Relb. Suppl.* 3. 7. *Sibth.* 15.

Abbot. 212. *Curt. Lond. fasc.* 6. t. 72.

Hoffm. Sal. v. 1. 45. t. 9, 10. t. 23. f. 2.

Villars. Dauph. v. 3. 762. *Tr. of Linn. Soc.*
v. 6. 118.

S. n. 9. Gmel. Sib. v. 1. 155. t. 34. f. 3.
Herb. Linn.

S. n. 1637. Hall. Hist. v. 2. 305.

*S. folio amygdalino utrinque aurito, corticem
abjiciens. Raii Syn.* 448.

Angl. Long-leaved Triandrous Willow.

In salicetis et ad ripas fluviorum frequens.

Arbor. Fl. Maio, etiam Augusto.

Arbor 30-pedalis. Cortex caudicis et ramorum deciduus,
ferè ut in Platano. Ramuli erecti, longi, viminei, tenaces,
fuscescentes, glabri, ad genicula subfragiles. Folia petio-
lata, triuncialia, linear-i-oblonga, basi angustata, late-
ribus æqualibus, acuminata, crebrè ferrata, ferraturis
inflexo-rotundatis, subglandulosis, utrinque glaberrima ;
suprà lætè viridia ; subtùs glaucescentia. Stipulæ ova-
to-obliquæ, crenatæ, venosæ, glabræ, sèpè nullæ.
Amenta ramulos novellos terminantia, erecta, cylin-
dracea, gracilia, flavicantia, squamis villosis, obtusis.
Nectarium obtusum. Stamina plerumque tria, rarissi-
mè in codem amento bina. Antheræ flavæ. Germen
pedicellatum, ovatum, acuminatum, tuberculatum.
Stigmata brevia, patentia, emarginata. Capsula glaber-
raria, virens.

Inter viminalia præstantiora numeranda est.

8: *SALIX triandra*, foliis ovatis obliquis ferratis *amygdalis*-
glabris, germinibus pedicellatis, stipulis maxi-lina-
mis.

Salix amygdalina. *Sp. Pl.* 1443. *Huds.* 426.
With. 47. *Hull.* 218. *Relh.* 365. *Lightf.*
596. *Tr. of Linn. Soc.* v. 6. 119.

S. folio auriculato splendente flexilis. *Raii Syn.*
448. *Cant.* 144:

Angl. Broad-leaved Triandrous Willow. Al-
mond-leaved Willow.

In salicetis et palustribus.

On Badley Moor, by Dereham, Norfolk. *Mr.*
Crowe. In osier-grounds, but not common,
being a bad osier.

Frutex. Fl. Aprili, Maio.

Nunquam arborescit. Cortex deciduus. Rami patentius-
culi; glabri, lutescentes, fragiles. Folia petiolata, ferè
biuncialia, latè ovata, basi rotundata, obliqua, sive la-
teribus inæqualibus, acuminata, undique ferrata, fer-
raturis inflexis, utrinque glaberrima; suprà latè viri-
dia; subtùs glauca. Stipulae maximæ, figuræ variæ, ro-
tundatæ, sive semicordatæ, crenatæ, deciduæ. Flores
(ex D. Crowe) triandri. Amenta fœminea præce-
dente crassiora, squamis glabris, capsulis pedicellatis,
glaberrimis.

Figuræ foliorum a præcedente satis differt, ex observa-
tionibus Dom. Crowe.

9. *SALIX triandra?*, foliis lanceolatis acumi- *Russel-*
natis ferratis glabris; germinibus pedicellatis *liana*.
subulatis lœvibus.

Angl. Bedford Willow. Leicestershire or Dish-
ley Willow.

In palustribus et salicetis.

Communicated by His Grace John Duke of Bed-
ford, VOL. III. K

ford, who, with his late illustrious brother, has paid much attention to the cultivation and uses of this hitherto nondescript species.

Arbor. Fl. Aprili, Maio.

Arbor excelsa, vegetissima, sub *S. fragili* adhuc, ni fallor, confusa. Rami nitidi, tenacissimi, et valde flexiles. Folia lanceolata, elongata, acuminata, glabra, nitida, undique ferrata, subtus glaucescentia. Stipulae exiguæ, semicordatae, dentatae. Amenta mascula nondum vidi; fœminea laxa, rachi tomentosa, squamis linearibus, pilosis, deciduis. Nectarium obtusissimum, apice croceum. Germen brevius pedicellatum, ovato-subulatum, laxe. Stylus parum elongatus, crassus. Stigmata bipartita.

Cortex ad rem coriariam omnium præstantissimus.

D. Biggin.

pentan- 10. *SALIX pentandra*, foliis elliptico-lanceolatis crenulatis glabris, germinibus glabris subsessilibus.

Salix pentandra. Sp. Pl. 1442. Huds. 426. With. 46. Hull. 218. Lightf. 595. Dicks. H. Sicc. fasc. 3. 15. Wilden. Baumz. 333. Tr. of Linn. Soc. v. 6. 120. Ehrb. Arb. 48. Pl. Off. n. 309.

S. n. 7. Gmel. Sib. v. 1. 153. t. 34. f. 1.
S. folio laureo, seu lato glabro odorato. Raii.

Syn. 449.

Angl. Sweet Willow. Bay-leaved Willow.

Ad rivos Angliae septentrionalis et Scotiae australis. Common in Westmoreland and the mountainous parts of Yorkshire, as remarked by Ray. About Moffat and other parts of the lowlands of Scotland.

Arbuscula. Fl. Maio, Junio.

Rami

Rami lœves, nitidi. Folia breviūs petiolata, elliptico-lanceolata, seu latè ovata, acuminata, crenulata, crenulis glandulosis, resinâ flavâ, odoratâ, scatentibus, utrinque lœtè viridia, nitida, glaberrima. Stipulæ vix ullæ. Amenta raimulos terminantia, solitaria, subnuntantia, cylindrica, crassiuscula, obtusa, flava, odorata, rachi villosâ. Squamæ rotundatæ, basi villosæ. Stamina circiter quinque, basi hirsuta. Germen subfessile, lanceolatum, glabrum, nitidum. Stigmata patentia, subrecurva, emarginata.

11. *SALIX* foliis elliptico-lanceolatis crenatis gla-nigribris subtùs glaucis, germinibus pedicellatis cans. lanceolatis acuminatis sericeis.

Salix nigricans. *Tr. of Linn. Soc.* v. 6. 120.
Engl. Bot. t. 1213.

S. phylicifolia B. *Sp. Pl.* 1442. *Fl. Lapp.* ed. 2.
291. t. 8. f. c. n. 350.

Angl. Dark broad-leaved Willow.

In salicetis.

At Wrongay Fen, Norfolk, and in osier-grounds in other places not uncommon. *Mr. Crowe*. Near Shobden Court, Herefordshire. *Mr. Dickson*.

Frutex. Fl. Aprili.

Vix arborescit. Rami rectiusculi, teretes, fragiliores, cortice nigricanti, glabro. Folia bi- vel tri-uncialia, elliptico-lanceolata, acuta, basi parùm rotundata, undique ferè crenata; suprà saturatè viridia, glaberrima, exsiccatione nigricantia; subtùs glauca, venosa, interdum pilosiuscula. Petioli basi dilatati. Stipulæ, si adhuc, majusculæ, obliquè cordatæ, serratæ, glabræ. Amenta uncialia vel sesqu'uncialia, crassiuscula, squamis lanceolatis, vel obovatis, obtusis, fuscis, pilosis. Stamina bina, capillaria, flava, basi plûs minus villosa. Antheræ oblongæ, aureæ. Capsulæ pedicellatæ, subulatæ, elongatæ, sericeo-albæ, uti pedicelli et rachis. Species a *S. phylicifolia* verâ proculdubio distincta.

bicolor. 12. **SALIX** foliis ellipticis acutis denticulato-serratis glabriusculis subtus glaucis, germinibus pedicellatis lanceolatis sericeis.

Salix bicolor. *Ehrb. Arb.* 118. *Wilden. Berl. Baumz.* 339.

S. laurina. *Tr. of Linn. Soc.* v. 6. 122.
Angl. Shining dark-green Willow.

In sylvis.

Frutex. Fl. Aprili, Maio.

Præcedenti affinis, prorsus verò distincta. Rami teretes, fragiles, cortice fusco, glabro. Folia bi- vel tri-uncialia, erecta, elliptica, subacuminata, subrevoluta, undique denticulata sive subserrata; suprà saturatè viridia, glabra, nitida, exsiccatione minus nigricantia; subtus valde glauca, venosa, pilosiuscula. Petioli basi maximè dilatati, triquetri. Stipulæ parvæ, lunato-semicordatae, dentatae, glabræ, subtus glaucæ. Amenta vix uncialia, squamis obovatis, obtusissimis, fuscis, pilosis. Stamina bina, flava, capillaria. Antheræ subrotundatae, luteolæ. Germina pedicellata, lanceolata, minus quam in priore attenuata vel acuminata, sericeo-alba, uti pedicelli et rachis. Stylus breviusculus, glaber. Stigmata crassa, ovata, connata, indivisa vel parùm emarginata.

Amentis masculinis duplo minoribus, staminibus brevioribus, a *S. nigricanti* certè dignoscitur; ut et colore foliorum lætiori, margineque subrevoluto denticulato, nec crenato.

petiolaris. 13. **SALIX** foliis lanceolatis serratis glabris subtus glaucis, germinibus pedicellatis ovatis scriccis, stigmatibus sessilibus bilobis.

Salix petiolaris. *Tr. of Linn. Soc.* v. 6. 122.
Engl. Bot. t. 1147.

Angl. Dark long-leaved Willow.

In salicetis et palustribus. *D. Dickson.*

Frutex.

Frutex. Fl. Aprili.

Magnitudo præcedentium. Rami viminei, graciles, teretes, glabri, atro-purpurei sive fusci. Folia 4- aut 5-uncialia, unciam ferè lata, lanceolata, acuminata, undique ferrata, subrevoluta, basi plerumque inæqualia, suprà lætè viridia, glaberrima, nitidiuscula; subtùs glauca, interdùm pilosiuscula; exsiccatione purpureo-nigrificantia. Petioli elongati, graciles, lineares, supernè sericei. Stipulæ parvæ, lunatæ, dentatæ, glabrae. Amenta mascula nondum vidi; fœminea vix uncialia, squamis obovatis, obtusis, sèrpè emarginatis, nigris, pilosis. Germina longius pedicellata, ovata, parva, sericea, stigmatibus omnino sessilibus, ovatis, obtusis, bilobis.

14. *SALIX* foliis lanceolatis undulato-crenatis *phylici-* glabris subtùs glaucis, stipulis sublunatis. *folia.*

Salix phylicifolia. *Tr. of Linn. Soc. v. 6. 123.*

S. phylicifolia α. *Sp. Pl. 1442. Fl. Lapp. ed. 2.*
291. t. 8. f. d. n. 351.

Angl. Tea-leaved Willow.

In alpibus Scoticis.

At Finlarig, Breadalbane. *Rev. Mr. Stuart.*

Herb. Lightf.

Arbuscula. Fl. Maio.

Rami læves, tenues, patentes. Folia alterna, petiolata, omnino lanceolata, acutiuscula, medium versùs præcipue inæqualiter et latius crenata, inter crenas sub-sinuato-undulata, margine aliquantulum incrassata; basi apiceque integerrima, eglandulosa; utrinque vel prorsus glaberrima, vel pilis spartis, vix conspicuis, subpubescentia; suprà saturatè viridia; subtùs glauca, reticulato-venosa. Petioli attenuati, læves. Stipulæ lunatæ, sèpius ferratæ, magnitudine variæ. Amenta non vidi.

Diversa a *S. phylicifoliâ* Jacquinii.

Arbuscula. 15. **SALIX** foliis lanceolatis obsoletè denticulatis
glabris exstipulatis, ramulis pubescentibus.

Salix Arbuscula. *Fl. Lapp.* ed. 2. 297. t. 8.

f. m. *Sp. Pl.* 1445, γ.

Angl. Little Tree Willow.

In alpibus Scoticis. *D. Dickson.*

Arbuscula. *Fl. . . .*

Caulis erectus, pedalis, gracilis. Rami erecto-patentes,
filiformes, juniores pubescentes. Folia alterna, petiolata,
lanceolata, utrinque acuta, denticulata, denticulis
minutis, sparsis, venosa, venis rectiusculis, omnia te-
nuia et laxiuscula; juniora subtilis pilosa, pilis adpressis,
sparsis. Petioli breviusculi, crassiusculi, demum glabri.
Stipulæ nullæ. Amenta

vitellina. 16. **SALIX** foliis lanceolatis acutis exstipulatis
ferratis suprà glabris: serraturis cartilagineis,
stigmatibus emarginatis.

Salix vitellina. *Sp. Pl.* 1442. *Huds.* 426.

With. 47. *Hull.* 218. *Sibth.* 16. *Hoffm.*

Sal. v. 1. 57. t. 11, 12. t. 24. f. 1. *Ehrb.*

Arb. 78.

S. folio longo subluteo, non auriculata, vimi-
nibus luteis: eademque viminibus rubris.

Raii Syn. 450.

S. folio utrinque glauco, viminibus rubris. *Raii*
Cant. 142.

Angl. Yellow Willow.

In salicetis et palustribus.

At Ovington, near Watton, Norfolk, abundantly,
and truly wild. *Mr. Crowe.*

Arbor, *Fl. Maio.*

Arbor mediocris. Rami erectiusculi, lutei, nitidi. Folia
subsessilia,

subsessilia, lanceolata, acuminata, ferrata, serraturis ercbris, parvis, cartilagineis; suprà glabra, nitida; subtùs cæstia, subsericea; juniora utrinque sericea; primordialia obtusa et integerima. Stipulæ nullæ. Amenta cylindraeea, acuta, squamis ovato-lanceolatis, extùs pubescentibus. Stamina duo, distincta. Germen sessile, lanceolatum. Stigmata emarginata.

Hanc speciem ex auctoriis conscripsi, nec fructificationem ejus adhuc satis examinavi. Hujus affinitatem cum *S. albâ* jam dudum notavit Hallerus, quam posteriores plerumque nec confirmant neque impugnant.

Liber in hâc specie luteus. *Hoffm.*

17. *SALIX* foliis lanceolatis acuminatis undique *fragilis*.
serratis glaberrimis, petiolis dentato-glandulosis, nectario masculinorum gemino.

Salix fragilis. *Sp. Pl.* 1443. *Huds.* 426.

With. 47. *Hull.* 218. *Rehb.* 365. *Sibth.* 16.

Abbot. 212. *Woodv. Med. Bot.* t. 198.

S. decipiens. *Hoffm. Sal.* v. 2. 9. t. 31.

S. folio longo latoque splendente, fragilis. *Raij*
Syn. 448. *Cant.* 143. n. 3.

Willow. *Hunt. Evel. Sylv. tab.* ad p. 245.

Angl. Crack Willow.

In salicetis et ad ripas fluviorum.

Arbor. Fl. Aprili, Maio.

Rami erecti, lœves, luteolo-fusci, basi fragillimi. Folia petiolata, 3—5-uncialia, lanceolata, acuminata, a basi ad apicem ferrata, serraturis obtusis, inflexis, glandulosis; utrinque glaberrima; suprà nitida: juniora apice ciliata: primordialia obovata, reflexa. Stipulæ rotundatæ, obsoletè dentatæ, majusculæ, at fæpiùs nullæ. Amenta masculina pallida, cylindracea, laxiuscula, squamis obovatis, villosis. Nectarium e glandulis (optimè monente Hoffmanno) duabus, luteis, quarum major inter stamina et rachin, minor e contrario inter stamina et squamam collocatur. Stanina duo,

duo, filiformia, glabra. Amenta fœminea nondum vidi.

Sub hoc nomine plures species apud nos forsitan lateant, (confer Hoffm. loc. cit.) at nondum rem probandi aut improbandi occasio data fuit.

tenuifolia. 18. *SALIX* foliis ellipticis acutis serratis glabriusculis subtus glaucis, stipulis obsoletis, capsulis glaberrimis.

Salix tenuifolia. *Fl. Lapp.* ed. 2. 292. t. 8. f. c. *Herb. Linn.*

S. Pontederæ. *Villars. Dauph.* v. 3. 766. t. 50. f. 8?

S. Fl. Dan. t. 1052?

Angl. Thin-leaved Willow.

In petrofisi ad ripas fluviorum.

Above the bridge at Kirkby Lonsdale, Westmoreland. 1783.

Frutex. Fl. Maio, Junio.

Frutex pedalis aut bipedalis, ramosissimus, patens. Rami rubicundi; juniores pubescentes, tomento brevissimo, incurvo. Folia haud sesquiuncialia, alterna, petiolata, elliptica, basi apice ue acutiuscula, medio præcipue serrata, quandoque subundulata ut in *S. phylicifoliæ*, tenuia, utrinque vel omniùm glabra, vel pilis tenuissimiis, albis, adpressis, sparsis, pubescentia; suprà pulchræ ac lè viridia; subtus glauco-albida, venis prominulis anastomosantibus reticulata, nervo subinde piloso; primordialia aut inferiora minora, angustiora, sèpius integririma. Stipulae exiguae, subfalcatae, serratae, glaberrimæ, sèpè nullæ. Amenta masculina flavelcentia, vix uncialia, squamis linearibus, angustis, canaliculatis, extùs pilosissimis, diandris; fœminea demùm sesquiuncialia, capsulis sessilibus, ovato-lanceolatis, glabris. Stylus elongatus. Stigmata crassissimula, emarginata.

Varietatem vel distinctam speciem eodem in loco legimus, foliis utrinque pilosioribus, etiam subtus villosis, minùsque

minùsque plerumque glaucis, squamis masculinis obovatis, planis. Hanc ulteriori in locis spontaneis investigationi, modo Deus nobis hæc otia fecerit, relinquimus.

19. *SALIX* foliis elliptico-oblongis dentatis re-malifor-pandis scariosis glaberrimis, stipulis cordatis *lia*. maximis.

Angl. Apple-leaved Willow.

In montosis?

Gathered by Mr. Crowe, in what particular spot is unknown.

Frutex. Fl.

Species distinctissima, habitu Pyri cujusdam, aut Mali, notabilis. Caulis tripedalis, ramosus, erectus. Ramī nigricantes; juniores hirti. Folia biuncialia, alterna, breviūs petiolata, elliptico-oblonga, acutiuscula, undique et crebrè dentata, plūs minūs undulata vel repanda, scariosa, rigidula, utrinque glaberrima; suprà lātē viridia; subtūs glaucescentia, venosa, venis divaricatis, anastomosantibus, pallidis, nervo albido aut basi rubicundo. Petoli crassi, basi latissimi. Stipulæ magnæ, cordatae vel semicordatae, recurvæ, dentatæ, venosæ, colore foliorum.

Variat foliis basi productis.

20. *SALIX* foliis elliptico-lanceolatis acutis inæ-radicans. qualitèr crenatis glaberrimis, ramis elongatis decumbentibus radicantibus.

Angl. Rooting-branched Willow.

In palustribus. *D. Dickson.*

Arbuscula. Fl.

Caulis tripedalis, ramis undique patentibus, decumbentibus, elongatis, radicantibus, glabris, foliosis; juniores fuscescentes. Folia sparsa, breviūs petiolata, stricta, biuncialia, elliptico-lanceolata, utrinque acuta, variè et inæqualitèr crenata vel subserrata, parùm repanda; suprà

suprà lätè viridia; subtùs glauca, venis flexuosis, anastomosantibus; utrinque glaberrima; juniora quandoque subpubescentia. Stipulæ exiguæ, ovatæ, arcuatæ, mox fuscescentes.

- myrsi
nites.* 21. **SALIX** foliis ellipticis ferratis glabris venosis utrinque nitidis, ramulis hirtis, capsulis subulatis pubescentibus.

*Salix myrsinoides. Sp. Pl. 1445. Fl. Lapp. ed. 2.
293. t. 8. f. f. t. 7. f. 6. Villars. Dauph.
v. 3. 769. t. 50. f. 12. Fl. Dan. t. 1054.
Hoffm. Sal. v. 1. 71. t. 18, et forsitan 17.*

*S. retusa. Dicks. Tr. of Linn. Soc. v. 2. 288.
ex ipso auctore. With. 48. Hull. 218.*

Angl. Green Whortle-leaved Willow.

In alpibus Scoticis.

Mountains of Glen-co. Rev. Mr. Stuart. Herb.
Ligbtf.

Frutex. Fl. Maio, Junio.

Caulis pedalis vel bipedalis, ramosissimus. Ramuli divaricati, densè foliosi, hirti, pilis porrectis, apice floriferi. Folia alterna, breviùs petiolata, vix uncialia, elliptica, undique ferrulata, rigidula, marcescentia, reticulato-venosa, utrinque lätè viridia, nitida, nequaquam glauca; inferiora obtusa; juniora subsericea. Stipulæ ferratæ, glabræ, formâ et magnitudine maximè variæ. Amenta terminalia, erecta, solitaria, pedunculata, crassa, obtusa, densa, squamis obovatis, obtusis, pilosis, rariùs denudatis. Antheræ plumbeæ. Capsulæ subulatæ, pube molli, brevi, densâ, vestitæ, demum ferè glabratæ. Stylus elongatus, glaber. Stigmata magna, bipartita.

- prunifo-
lia.* 22. **SALIX** foliis ovatis ferratis glabris: suprà lævibus; subtùs glaucis, ramulis subpubescentibus, capsulis ovatis sericeis.

Salix

Salix myrsinites. *Lightf.* 599. *Herb. Lightf.*
Huds. 427? *Hoffm. Sal.* v. 1. t. 19. et t. 24.
f. 2.

Angl. Plum-leaved Willow.

In alpibus Scoticis frequens.

Found by the Rev. Mr. Stuart and Rev. Dr. Walker in several parts of the Highlands.

Frutex. Fl. Maio, Junio.

Caulis tripedalis, ramosissimus. Rami patentes, sæpè rubicundi, foliosi, apice pubescentes, pilis brevissimis, curvis. Folia uncialia, alterna, petiolata, ovata, acuta, undique ferrulata; suprà saturatè viridia, glabra, lævia, nitida; subtùs glauca, venosa, opaca, et subsericea, imprimis juniora atque inferiora. Stipule exiguæ, semiovatæ, dentatæ, sæpè nullæ. Amenta e ramulis lateralibus brevioribus, solitaria, erecta, triplò quam in præcedente tenuiora, et longitudine vix uncialia, squamis subrotundis, extùs hirsutis. Capsulæ duplò minores quam in præcedente, ovatæ, undique sericeo-pilosæ, demùm glabriuscculæ, badiæ. Stylus brevis. Stigmata parva, bipartita.

Varietatem, ut videtur, foliis omnibus subtùs sericeis ex amicissimo D. Stuartio accepi.

23. *SALIX* foliis ovatis serratis glabris : suprà re-*venulosa*.
ticulato-venosis; subtùs glaucescentibus, cap-
sulis

Angl. Veiny-leaved Willow.

In alpibus Scoticis. *D. Dickson et D. Winch.*

Frutex. Fl.

Forma præcedentis; diserepat verò foliis suprà concinnè *venulosis*, venis prominentibus, parallelis, marginem versùs reticulato-anastomosantibus; subtùs pallidè *viridibus*, minusque glaucis.

24. *SALIX* foliis ovatis denticulatis glabris com-*carinata*.
presso-carinatis tenuissimè *venulosis*.

Angl.

Angl. Folded-leaved Willow.

In alpibus Scoticis. *D. Dickson.*

Frutex. Fl.

Præcedentibus duabus paulò major et erectior. Rami juniores vix pubescentes, stricti. Folia sesquiuncialia, breviis petiolata, ovato-elliptica, acuta, denticulata, nec ferrata, arcuato-recurva, compresso-carinata; suprà viridia, nitida, tenuissimè venulosa; subtùs pallidiora, opaca, vix glauca; utrinque glaberrima; juniora subtùs obsoletè sericea.

myrtill-
loides.

25. **SALIX** foliis ellipticis acutis obsoletè denticulatis glabris subtùs glaucis, ramulis glaberrimis.

Salix myrtilloides. *Sp. Pl.* 1446. *Fl. Lapp.*
ed. 2. 295. *t.* 8. *f.* i. *k.*

Angl. Glauous Bilberry-leaved Willow.

In alpibus Scoticis. *D. Dickson et D. Winch.*

Frutex. Fl.

Frutex pedalis, erectus, ramosus. Ramuli fusco-virides, glaberrimi. Folia vix uncialia, patentia, plana, elliptica, acuta, obsoletè denticulata vel subferrata, denticulis apice glandulosis, haud integerrima, aliquantulum revoluta; suprà lævissima, lætè viridia; subtùs glauca, tenuissimè venulosa; utrinque glaberrima. Petioli breviusculi, basi incrassati.

herbacea. 26. **SALIX** foliis orbiculatis serratis glaberrimis reticulato-venosis utrinque nitidis, capsulis glabris.

Salix herbacea. *Sp. Pl.* 1445. *Fl. Lapp.* *ed.* 2.
294. *t.* 8. *f.* h. *t.* 7. *f.* 3, 4. *Huds.* 427.
With. 48. *Hull.* 218. *Lightf.* 600. *Fl.*
Dan. *t.* 117. *Hoffm. Sal.* v. 1. 74. *t.* 20.

S. n. 1649. *Hall. Hist.* v. 2. 309.

S. alpina,

S. alpina, *alni rotundo folio repens*. *Dall. in Raii Syn.* 448. *Bocc. Mus.* 19. t. 1.
Angl. Least Willow.

In summis alpibus, solo micaceo.

On Snowdon. *Sherard.* On the tops of all the Highland Alps. *Lightf.* On Skiddaw, Cumberland. *Mr. Crowe and Mr. Woodward.*

Fruticulus. Fl. Junio.

Radix longè latèque repens, lignosa, ramosissima. Caules vix unciales, lignosi, subramosi, foliosi, glabri, fusi. Folia brevius petiolata, semuncialia, orbiculata, obtusa, quandoque emarginata, undique ferrata, rigidula, reticulato-venosa, utrinque latè viridia, nitida, glaberrima. Stipulæ nullæ. Amenta terminalia, solitaria, pedunculata, pauciflora, squamis obovatis, obtusis, subciliatis, extus glabratis. Stamina flava. Capsulæ ovato-lanceolatae, glabræ, rubicundæ. Stigmata parvæ, bifida.

** *Foliis integerrimis glabriusculis.*

27. *SALIX* foliis elliptico-orbiculatis obtusis in-reticulato-venosis subtus lata. glaucis, capsulis villosis.

Salix reticulata. Sp. Pl. 1446. Fl. Lapp. ed. 2. 296. t. 8. f. 1. t. 7. f. 1, 2. Huds. 428. With. 49. Hull. 218. Lightf. 601. Dicks. Dr. Pl. 44. Fl. Dan. t. 212. Hoffm. Sal. v. 2. 3. t. 25, 26, 27.

S. n. 1650. Hall. Hist. v. 2. 309.

S. pumila, folio rotundo. Raii Syn. 449. Bauh. Hist. v. 1. p. 2. 217.
Angl. Wrinkled Willow.

In summis alpibus.

On Ingleborough and Whern-side, Yorkshire; T. Willisel. Ray. On the highest mountains of

of Wales. *Ray.* On many of the Highland mountains. *Lightf.*

Fruticulus. Fl. Junio.

Radix lignosa. Caules ramosissimi, lignosi, validi, diffusi, cinerei, glabri, ramulis rubicundis, foliosis. Folia longius petiolata, patentia, præcedente duplo majora, elliptico-subrotunda, vel obovata, obtusa, integerrima, subrevoluta, rigida, reticulato-venosa, glabra; suprà viridia; subtus albido-glaуca, venis rubris; juniora pilosa. Petioli graciles, rubri. Stipulæ nullæ. Amenta terminalia, solitaria, longius pedunculata, brevia, subclavata, densa, squamis parvis, obovatis, obtusis, subpubescentibus. Stamina rubra. Capsulæ ovatæ, villosæ. Stigmata subsessilia, bipartita.

*** *Foliis villosis.*

arenaria. 28. *SALIX* foliis subintegerrimis ovatis acutis: suprà subvillosis; subtus densissimè lanatis.

Salix arenaria. Sp. Pl. 1447. Fl. Lapp. ed. 2. 298. t. 8. f. o. q.

S. lapponum. Lightf. 604. Huds. 651. With. 51. Hull. 218.

S. helvetica. Villars. Dauph. v. 3. 783.

S. n. 20. Gmel. Sib. v. 1. 164. t. 36. f. 1. Herb. Linn.

Angl. Downy Mountain Willow.

In alpibus Scoticis.

On the highland mountains of Breadalbane. Rev. Mr. Stuart. Herb. Lightf.

Frutex. Fl. Junio.

Caulis 4—5-pedalis, ramosus. Rami crassiusculi, rubicundi, sublanati. Folia petiolata, erectiuscula, subsessilia, ovata, acuta, parùm revoluta, plerumque integerrima, subiudè obsoletè ferrata; suprà atro-viridia, tomento raro, laxo, tecta, et demùm glabrata; subtus dense lanata, nivea, venis prominentibus; in plantâ

plantā masculinā angustiora, magisque subtūs sericea, ad S. lapponum veram aliquantulum accendentia. Stipulae ut videtur omnino nullae. Amenta brevia, villosissima, obtusa. Stamina lutea. Capsulae lanatae.

Gmelinus ut videtur Linnæum in errorem duxit de synymo Dillenii apud Raii Syn. quod ad sequentem spectat.

29. SALIX foliis integerrimis ellipticis subrevolute mucronulatis aduncis: suprà subvillosis; ^{tea.} subtūs ramulisque sericeo-argenteis.

Salix repens γ. *Huds.* 429.

S. fusca. *With.* 51. *Hull.* 219.

S. arenaria. *Lightf.* 604.

S. pumila, foliis utrinque candidantibus et lanuginosis. *Dill. in Raii Syn.* 447. t. 19. f. 3.

Angl. Silky Sand Willow.

In arenosis maritimis.

By Sandon Castle, Isle of Wight. *J. Sherard.*

On the sea shores of Scotland, among loose blowing sand. *Lightf.* On the sand burrows at Laugharn, South Wales. *Mr. Hurlock.* About Walmer Castle and Deal, Kent. *Rev. Dr. Goodenough.*

Frutex. Fl. Maio.

Caules diffusi, vel prostrati, ramis elongatis, patentibus, densè foliosis, subangulatis, sericeis. Folia sparsa, petiolata, ferè uncialia, elliptica, integerrima, subrevoluta, mucronulo reflexo terminata; suprà tomento raro, adpresso, seu pilis subsericeis, demùm deciduis, tecta; subtūs undique sericeo-nitidis. Stipulae exiguae, villosæ, glandulo mucronatae, sèpè nullæ. Amenta ramorum apicem versùs, cylindracea, obtusa, squamis ellipticis, obtusis, sericeis. Capsulae pedicellatae, ovato-lanceolatae, sericeæ, demùm glabratæ. Stigmata bipartita.

Hanc speciem aut planè ignoravit Linnæus, aut cum *S. fusca*,

S fuscâ, ut ego in Fl. Lapp. ed. 2, confudit. Differt à fuscâ foliis rotundioribus, nunquam non integerrimis, apice aduncis; ramulis crassioribus, densius foliosis; amentis subterminalibus; habituque longè magis sericeo ac nitido.

prostra-ta. 30. **SALIX** foliis subdentatis ellipticis acutis: subtûs glaucis sericeis, caule prostrato.

Salix polymorpha. Ehrb. Arb. 49?

Angl. Prostrate Dwarf Willow.

In montosis humidis. *D. Dickson.*

Fruticulus. Fl.

Caules prostrati, ramis elongatis, attenuatis; junioribus subsericeis. Folia sparsa, brevius petiolata, vix uncialia, elliptica sive elliptico-lanceolata, hinc inde dentata, vix revoluta, acuta; suprà saturatè viridia, obsoletè venosa, parùm pubescentia; subtûs glauca, subsericea. Stipulæ nullæ.

A præcedente differt caule ramiisque omnino prostratis, minùsque pubescentibus; foliis angustioribus, hinc inde denticulatis vel subfetratis, longè minùs nitidis, vix revolutis, vel apice aduncis. A *S. fuscâ*, præter caulem prostratum, petiolis brevibus incrassatis distinguatur.

fusca. 31. **SALIX** foliis subdenticulatis elliptico-oblongis acutis: suprà glabris; subtûs glaucis sericeis, petiolis attenuatis.

Salix fusca. Sp. Pl. 1447. *Fl. Lapp. ed. 2.* 299.

t. 8. f. r. Fl. Suec. ed. 2. 351.

Angl. Brownish Dwarf Willow.

In sylvis et ericetis humidiusculis.

In Breadalbane. *Rev. Mr. Stuart.*

Fruticulus. Fl. Maio.

Caules erecto-patentes, ramulis rectinsculis, pubescentibus, subangulatis. Folia sparsa, petiolata, vix uncialia, elliptico-

elliptico-oblonga, acutiuscula, apice recta; margine haud revoluta, nequaquam integerrima, sed denticulis raris, sparsis, glandulosis, in adultioribus et superioribus magis conspicuis et numerosis, munita; supra fusco-viridia, vel omnino glaberrima, vel pilis raris, exiguis, adpresso, albis, conspersa; subtus glauca, subsericea, vernalis tenuissimis, anastomosantibus. Petioli demum elongati, pilosi, multo magis quam in praecedente attenuati. Stipulae vix illae. Ramuli floriferi laterales, breves, ut in S. repente. Amenta cylindracea, obtusa, vix uncialia, squamis pilosis; masculinis obovatis, obtusis, pallidis, apice fuscoc-sanguineis; foemineis lanceolatis, angustioribus, unicoloribus. Stamina flava. Capsulae pedicellatae, subulatae, sericeo-villose. Stigmata bipartita.

Exemplaria a D. Stuartio missa cum archetypo originali, in Fl. Lapponicâ descripto, prorsus convenientia. In herbario Linnæi aliud exemplar adest, nomine S. incubaceæ insignitum, quod. S. fusca foliis adultioribus mihi omnino videtur; at hoc nullius est auctoritatis ad Salicem incubaceam (quæ mihi valde obscura) determinandam, cum neque signo necessario antiquitatis, ad priorem Spec. Plant. editionem spectante, gaudeat, neque hujusce characteri specifico benè respondeat.

S. fusca Hoffmanni parvitate suâ, amentis terminalibus, solitariis, foliisque utrinque glabris, saturatè viridibus, ab hac et praecedenti facilè dignoscenda videtur. Faciem insuper habet S. repentis.

32. SALIX foliis integrimis elliptico-lanceolatis *repens*.
submucronulatis: supra nudiusculis; subtus
glaucis sericeisve, caule depresso.

Salix repens. Sp. Pl. 1447. Huds. 428. With.
51. Hull. 218. Kelb. 366. Abbot. 213.
Engl. Bot. t. 183.

S. depressa. Hoffm. Sal. v. i. 63. t. 15, 16.
Sibth. 16.

S. pumila angustifolia, pronâ parte cinerea.
Ruij Syn. 447. Bauh. Hist. v. 1. p. 2. 213.
VOL. III. L Chamæitæ,

Chamæitea, sive *Salix pumila*. *Ger. em.*

^{1391.}

β. *Salix pumila angustifolia*, infernè lanuginosa.
Raii Syn. 447.

S. humilis repens. *Ger. em. 1391.*

γ. S. alpina pumila, rotundifolia repens, infernè
subcinerea. *Dill. in Raii Syn. 448.*

S. humilis. *Ger. em. 1391.*

S. humilis repens. *Lob. Ic. v. 2. 138.*

Angl. Creeping Dwarf Willow.

In ericetis et palustribus arenofis.

Fruticulus. Fl. Maio.

Caules procumbentes, repentes et radicantes, ramosissimi,
lignosi et rigidi, ramis divaricatis, implexis, glabris, vel
apice tantum sericeo-pubescentibus. Folia brevius petiolata,
elliptico-lanceolata, longitudine varia, integerima,
subrevoluta, mucronulo brevi, recto, terminata;
suprà omnino glabra vel rarius pilosa; subtus glauca,
atque saepius sericea. Stipulae nullæ. Amenta numero-
sa, lateralia, alterna, brevia, obtusa, squamis obovatis,
obtusis, pilosis, rubicundis. Capsule pedicellatae, lan-
ceolatae, pubescentes, demum glabratæ, fuscæ. Stigmata
bipartita.

Varietatum limites vix inveniuntur. Differt a præceden-
tibus magnitudine minori, habitu vix sericeo, amentis
 numerosioribus, ramulis implexis, depresso-
ribus; at præcipue a S. fuscâ discrepat foliis omnibus
semper integerrimis, nec marginे scilicet glandulosis,
neque denticulatis.

rosmarini- 33. *SALIX* foliis integerrimis linear-lanceolatis
nifolia. strictis: subtus sericeis, caule erecto, stipulis
erectis planis.

Salix rosmarinifolia. *Sp. Pl. 1448. Fl. Suec.*
ed. 2. 352. Witb. 52. Hull. 219. Ehrb.
Arb. 119.

S. repens s. *Huds. 429.*

S. pumila,

S. pumila, rhamni secundi Clusii folio. *Dill. in Raii Syn.* 447.

S. humilis repens angustifolia. *Lob. Ic. v. 2.*
137. *Baub. Hist.* v. 1. p. 2. 214.

Angl. Rosemary-leaved Willow.

In arenosis humidis.

Found by J. Sherard. *Dill.* A specimen in Mr. Rose's herbarium agrees exactly with that of Linnæus from Abo.

Frutex. Fl. Aprili, Maio.

Caulis tripedalis, erectus, vel erectiusculus, ramosus. Rami viminei, recti, dense foliosi; juniores sericei. Folia brevius petiolata, erecta, stricta, subsesquiuincialia, linear-lanceolata, acuta, integerima, margine subindè glandulosa; supra pubescentia, ætate plus minus glabra; subtus sericeo-villosa. Stipule erectæ, lanceolatæ, planæ, subintegræ. Amenta lateralia, brevia, cernua.

34. *SALIX*, foliis subintegerrimis obovato-lanceo-*cinerea*. latis: subtus reticulato-venosis glaucis subvillosis, stipulis semicordatis serratis.

Salix cinerea. *Sp. Pl.* 1449. *Fl. Suec.* ed. 2. 353. *Fl. Lapp.* ed. 2. 296, excluso syn. *Villarsii*.

S. daphnoides. *Villars. Dauph.* v. 3. 765. t. 50. f. 7. ex ipso auctore.

Angl. Grey Willow.

In sylvis.

Specimens from Cumberland and from Fream Wood are in Mr. Lightfoot's herbarium.

Arbor. Fl. Maio.

Arbor 20- aut 30-pedalis. Rami læves, nitidi, rubicundi; juniores pubescentes. Folia petiolata, sesquiuincialia, obovato-lanceolata, acuta; supra glabra, vi-

ridia, nitida, lævia; subtùs glauco-cinerea, brevissimè villosa, subrubiginosa, venis prominentibus, parallelis, venulis reticulatis: inferiora omnino integerrima; superiora plùs minùs ferrata. Stipulæ parvæ, semi-cor datæ, ferratæ, margine reflexæ, utrinque villosæ uti petioli, deciduæ. "Amenta brevissima, densissima."

Villars.

S. cinerea Witheringii ab hâc planè diversa est.

aurita. 35. SALIX foliis subserratis obovatîs obtusis mucronulo adunco: utrinque villosis reticulato-venosis, stipulis polymorphis.

Salix aurita. Sp. Pl. 1446. Fl. Lapp. ed. 2. 303. t. 8. f. y. Lightf. 602. With. 50. Hull. 218. Sibth. 17. Abbot. 212. Ehrb. Arb. 39. Hoffm. Sal. t. 22. f. 1.

S. caprea δ. Hudſ. 430.

S. folio rotundo minore. Dill. in Raii Syn. 450. Giff. 38. append. 36. Herb. Sherard.

S. caprea rotundifolia. Ger. em. 1390.

ꝝ *S. caprea* ε. Hudſ. 430.

S. caprea pumila, folio subrotundo, subtùs incano. Dill. in Raii Syn. 450. Herb. Sherard.

Angl. Round-eared Willow. Trailing Sallow.

In sylvis et sepibus, imprimis submontosis humidis.

Frutex. Fl. Aprili, Maio.

Arbuscula ramosissima, altitudine varia, plerumque erecta, ramis patentibus, elongatis, fusco-cinereis, incanis. Folia alterna, breviùs petiolata, obovata, obtusissima, mucronulata, mucronulo reflexo, reticulato-venosa; suprà cinereo-virentia, mollè pubescentia; subtùs villosa, incana, villo venis inferto, undique divaricato, subdepresso; juniora subintegerrima; adūltiora medium versùs ferrata, crispa, et undulata. Stipulæ formâ et magnitudine variæ, sæpiùs rotundatæ, dentatæ, villosæ, venosæ, persistentes. Amenta e ramulis laterilibus

libus brevissimis, solitaria, cylindracea, multiflora, squamis parvis, fuscis, plūs minūs pilosis, sēpiās obtusis. Stamina flava, interdūm pilosa. Capsulæ longius pedicellatæ, subulatæ, villosæ. Stigmata subsessilia, crassa, indivisa.

Huc potius spectat *S. ulmifolia* *Villars.* *Dauph.* v. 3. 776.

t. 51. f. 20. quam ad *S. cinereum* in *Fl. Lapp.* ed. 2.
ex errore retuli.

36. *SALIX* foliis subserratis obovato-ellipticis *aquatica*: pubescentibus planis: subtūs glaucescentibus, stipulis rotundatis dentatis.

Salix caprea α . *Huds.* 429.

S. cinerea. *With.* 54. *Hull.* 219. *Relb.* 387.

S. aurita. *Hoffm. Sal.* t. 5. f. 3.

S. folio ex rotunditate acuminato. *Raii Syn.* 449.

Cant. 145. n. 1.

Angl. Common Willow. Water Sallow.

In sepibus et dumetis humidioribus ubique.

Arbuscula, Fl. Aprili,

Frutex vel arbor humilis, ramosissima, erecta, ramis junioribus incanis. Folia breviūs petiolata, obovata, vel elliptica, plūs minūs acuta, plana, vix rugosa, medium versùs ferrata; inferiora sēpè integerrima; omnia mollia, cinereo-virentia, suprà demūm glabrata, subtūs glaucescentia, venulosa, pilosuscula. Stipulæ rotundatæ, fornicatæ, dentatæ, præcedente minores. Gemmæ incanæ. Amenta cylindracea, squamis parvis, fuscis, pilosis. Stamina flava, hirta. Capsulæ pedicellatæ, subulatæ, villosæ. Stigmata ferè sessilia, indivisa.

S. auritæ propinqua, at distincta species,

37. *SALIX* foliis obovato-lanceolatis planis denticulatis acutis: subtūs glaucis hirtis, stipulis exiguis.

Angl. Olive-leaved Willow.

In dumetis. *D. Dickson.*

Frutex. Fl.

Frutex quadripedalis, ramis sparsis, laxè patentibus et divaricatis, tenuissimè pubescentibus. Folia patentia, rigidula, obovato-lanceolata, acuta, margine denticulato-scabra; suprà lätè viridia, tenuissimè pubescentia, plana; subtùs glauca, reticulato-venosa, pilis brevibus, ferrugineis, hirta. Petioli mediocres, crassiusculi, suprà pubescentes. Stipulæ exiguæ, rotundatæ, subfalcatae, planæ, adpressæ, vix dentatae. Gemmæ incanæ.

cotinifo- 38. *SALIX* foliis elliptico-orbiculatis obsoletè denticulatis; subtùs villosis; venis rectangulis.

Angl. Quince-leaved Willow.

In dumetis montosis. *D. Dickson.*

Frutex. Fl. Aprili.

Bipedalis, ramis patentibus, elongatis, pubescentibus, undique foliosis. Folia patentia, rigida, latè elliptica, siccè orbiculata, plana, mucronulata, obsolete denticulata; suprà tristè viridia, pubescentia, pilis raris, adpressis; subtùs pallida vel glauca, hirta, reticulato-venosa, venis rectangulis, divaricatis, prominentibus. Petioli crassi, pilosi. Stipulæ exiguæ, membranaceæ, marcescentes. Gemmæ villosæ. Amenta fœminea semiuncialia, ovata, squamis obtusis, brevibus, pilosis, apice nigris. Germen breviùs pedicellatum, lanceolatum, villosum. Stylus semibifidus. Stigmata obtusa, hinc fissa.

sphacelata. 39. *SALIX* foliis integerrimis ellipticis planis utrinque pubescentibus apice subsphacelatis, stipulis obsoletis, capsulis subulatis.

Salix lanata. *Lightf.* 602. *Herb.* *Lightf.* *With.* 50. *Hull.* 218.

S. capreæ varietas. *Hoffm.* *Sal.* v. 1. 28. t. 5. f. 4. t. 21. f. d.

Angl.

Angl. Withered-pointed Willow.

In convallibus alpinis Scottiæ.

At Finlarig at the head of Loch Tay. Rev.
Mr. Stuart.

Arbuscula. Fl. Aprili, Maio.

Rami ramulique subpubescentes, patentes, rubicundi. Folia petiolata, uncialia, vel sesquiuncialia, elliptica, superficie plana sive æquivalia, subtenuia, integerrima, vel rarissimæ et obsoletæ crenata; apice plerumque sphacelata, fusca; utrinque pubescentia, at subtus densius tomentosa; suprà demum ferè glabrata, in primis apicem versus. Stipulæ vel nullæ, vel exiguae, sublunatæ, glandulosæ, demum glabratæ. Petioli graciles, tomentosi. Amenta lateralia, cylindrica, uncialia, multiflora, squamis fuscis, ellipticis vel lanceolatis, pilosis. Stainina flava. Capsulæ longius pedicellatae, subulatae, attenuatae, villosæ, demum penè denudatae. Stigmata subseffilia, indivisa.

Hujus varietas videtur S. n. 19. *Gmel. Sib.* v. i. 163, S. caprea γ, *Sp. Pl.* 1448, *Herb. Linn.* foliis angustioribus, brevius aliquantulum petiolatis, subtusque magis sericeis.

S. lanata Linnæi certo certius discrepat capsulis subseffilibus, ovatis, glabrioribus, stylo elongato, stigmatibus bipartitis, quæ omnia sectionem diversam generis indicant. Huc retulit Linnæus Salicem suam depresso, *Fl. Suec.* 352, cuius folia S. nostram sphacelatam valde referunt; capsulæ vero, quæ hanc litem facile dirimarent, in exemplario Linnæano desunt.

40. SALIX foliis ovatis acuminatis serratis undu- *caprea.*
latis: subtus tomentosis, stipulis sublunatis,
capsulis ventricosis.

Salix caprea. *Sp. Pl.* 1448, α. *Huds.* 429, γ.
Witb. 53. *Hull.* 219. *Relb.* 367. *Sibtb.* 17.
Abbot. 212. *Lightf.* 607. *Hoffm. Sal.* v. i.
25. t. 3. f. 1, 2. t. 21. f. a, b, c. *Erb.*
Arb. 98.

S. latifolia rotunda. *Raii Syn.* 449.

S. folio subrotundo auriculata, *Raii Cant.* 145.
n. 2.

S. latifolia infernè hirsuta. *Bauh. Hist.* v. 1.
p. 2. 215.

Angl. Great Round-leaved Sallow.

In sylvis et sepibus minus humidis, vulgaris.

Arbor. Fl. Aprili.

Rami teretes, laeves, nitidi, subfuscæ, ramiulis pubescens-tibus. Folia alterna, petiolata, figurâ varia, at plerumque elliptica vel subrotunda, acuminata, magna, undulata, repanda, vel ferrata; suprà laeviuscula, glabra; subtus densè tomentosa, venosa. Stipulæ lunatæ vel subrotundæ, recurvæ, repandæ, minus tomentosæ. Petioli lineares, suprà quandoque densius villosi. Amenta ante folia expansa, ovata, crassa, multiflora, squamis obovatis, pilosissimis. Stamina flava. Capsulæ pedicellatæ, ovatae, basi ventricosæ, villosæ. Stigmata subsessilia, indivisa, demum subinde fissa.

acumi- 41. *SALIX* foliis lanceolato-oblongis acuminatis
nata. undulatis denticulatis: subtus tomentosis,
stipulis reniformibus, capsulis ovato-subulatis.

Salix acuminata. *Mill. Diet.* ed. 8. n. 14. *Hoffm.*
Sal. v. 1. 39. t. 6. f. 1, 2. t. 22. f. 2.
With. 53. *Hult.* 219. *Sibth.* 17. *Wilt.*
Baumz. 346.

S. caprea varietas. *Lichtf.* 608. excluso syn.
Sp. Pl.

S. caprea, acuto longoque folia. *Raii Syn.* 450.

S. caprea latifolia. *Ger. em.* 1390.

Angl. Long-leaved Sallow.

In sylvis et sepibus humidiusculis frequens.

Arbor. Fl. Aprili.

Habitus præcedentis, sed plerumque humilior. Rami teretes,

teretes, subvaginei, glabriusculi; juniores subsericeo-pubescentes. Folia alterna, petiolata, triuncialia, lanceolato- five ovato-oblonga, acuminata, magis minusve repanda et undulata, apicem versus præcipue denticulata: suprà lærè viridia, lævia, glabrata; subtùs tomentosa, nervo rubicundo, venis arcuatis, parallelis. Stipulæ reniformes, sessiles, dentatæ, venosæ, ad folia præcipue inferiora. Petioli ut in priore. Amenta ovato-cylindracea, obtusa, ante folia enata, squamis ovatis, acutiusculis, pilosis. Stamina flava. Nectarium conicum, constrictum. Capsulæ longius pedicellatæ, ovatæ, attenuatæ, pubescentes. Stylus brevissimus. Stigmata obtusa, indivisa, subindè bifida.

42. SALIX foliis lanceolatis acuminatis obsoletè *stipularis*-
crenatis: subtùs tomentosis, stipulis semicor-*ris*.
datis maximis, nectario cylindraceo.

Salix stipularis. *Engl. Bot. t. 1214.*

Angl. Auricled Osier.

In salicetis.

Near Bury. *Mr. Crowe.*

Frutex. Fl. Martio.

Rami erecti, elongati, teretes, tomentosi. Folia alterna, petiolata, erectiuscula, ferè spithamea, lanceolata, acuminata, inæqualiter et obsoletè crenata; suprà viridia, glabra; subtùs tomentosa, nervo parùm rubicundo, venis arcuatis, parallelis. Stipulæ ultra semunciales, subpetiolatæ, semicordatæ, longius acuminatæ, basi præcipue dentato-lobatæ; suprà glabræ; subtùs tonientosæ. Petioli crassiusculi, basi ampliati. Amenta ferè præcedentis, at stigmatibus linearibus, indivisis. Nectarium cylindraceum, elongatum.

Diffr. a præcedente stipularum magnitudine et formâ, foliis angustioribus duplòque longioribus, nectario, stigmatibusque linearibus. Habitu ad S. viminalem et mollissimam accedit, sed in usus œconomicos longè minus præstat,

43. SALIX

mollissi- 43. SALIX foliis lanceolatis acuminatis subcrenatis : subtus sericco-incanis, stipulis lunatis minimis.

Salix mollissima. Ehrb. Arb. 79. Hoffm. Fl.

Germ. 343. Widen. Baumz. 349.

Angl. Velvet Osier.

In salicetis.

In osier-grounds about Bury. *Mr. Crowe.*

Frutex. Fl. . . .

Rami erecti, viminei, tenacissimi, teretes, glabri. Folia alterna, petiolata, ovato-lanceolata, acuminata, regulariter at levius crenata, tenuia; suprad viridia, laevia et glaberrima; subtus sericea, tactu molliissima, nervo rubro, prominulo, venis arcuato-divaricatis. Stipulae exiguae, lunatae, obtusae, dentatae, pilosae. Petioli attenuati, pubescentes, basi paululum dilatati. Amenta nondum apud nos protulit.

Vimina optima et copiosissima praebet. *D. Crowe.*

rimina- 44. SALIX foliis lanceolato-linearibus longissimis acuminatis integerrimis subtus sericeis, ramis virgatis, stylo elongato.

Salix viiminalis. Sp. Pl. 1448. Huds. 430.

With. 54. Hull. 219. Relb. 367. Sibth. 17.

Abbot. 213. Hoffm. Sal. v. 1. 22. t. 2. f. 1,

2. t. 5. f. 2. t. 21. f. e, f, g: Ehrb. Arb. 69.

S. folio longissimo. Raii Syn. 450. Cant. 146.

S. angustis et longissimis foliis crispis, subtus albicantibus. Bauh. Hist. v. 1. p. 2. 212.

Elæagnus. Dalech. Hist. 278.

Angl. Common Osier.

In salicetis et palusribus.

Fruter. Fl. Aprili, Maio.

Rami longissimi, erecti, viminei, teretes, sericei. Folia sparsa,

sparsa, brevius petiolata, erectiuscula, ferè spithamæa, linear-lanceolata, acuminata, integerrima, subundulata; suprà viridia, glabra; subtus sericeo-argentea, nervo prominente, venis numerosissimis, parallelis. Stipulæ linear-lanceolatæ, sæpius nullæ. Amenta cylindracea, antè folia erumpentia, squamis subrotundis, obtusis, pilosissimis. Stamina lutea, antheris subindè fulvis. Capsulæ sessiles, ovatæ, subsericeæ. Stylus elongatus, filiformis. Stigmata acuta, tenuia, indivisa.

45. SALIX foliis lanceolatis acuminatis serratis *alba*, utrinque sericeis; serraturis infimis glandulosis, stigmatibus bipartitis.

Salix alba. *Sp. Pl.* 1449. *Huds.* 430. *Wittb.* 55. *Hull.* 219. *Relb.* 367. *Sibth.* 18. *Abbot.* 213. *Hoffm.* *Sal.* v. 1. 41. t. 7, 8. *Wilden.* *Baumz.* 350.

Salix. *Raii Syn.* 447. *Ger. em.* 1389. *Matth.* *Valgr.* v. 1. 180. *Camer. Epit.* 107. *S.* folio utrinque glauco, viminibus albidioribus. *Raii Cant.* 142. *Angl.* Common White Willow.

In sylvis et pratis humidis.

Arbor. Fl. Aprili, Maio.

Arbor procera, cortice rimoso. Rami ramosissimi, teretes, patentes; juniores sericei. Folia alterna, brevius petiolata, lanceolata, acuminata, acutè et regulariter ferrulata, serraturis inferioribus remotioribus, glandulosis; utrinque pubescentia; subtus præcipue sericea; juniora omnino argentea, convoluta. Stipulæ nullæ. Amenta terminalia, cylindracea, elongata, gracilia, multiflora, squamis elliptico-lanceolatis, fuscis, pubescibus. Stamina flava, squamis parùm longiora. Capsulæ subsessiles, ovatæ, fuscæ, glabratæ. Stylus breviusculus. Stigmata bipartita, crassiuscula.

Liber, monente Hoffmanno, in hac specie viridis, in *S.* vitellinâ luteus est.

Varietatem

Varietatem habemus foliis majoribus, cærulecentibus, citissimè crescentem, notatu dignissimam, at vix charactere specifico distinguendam.

Obs. Hoc genus ad germina sessilia vel pedicellata, stylum elongatum, brevissimum, aut nullum, stigmata integra vel bipartita, in sectiones magis naturales quam ad pubescentiam vel marginem foliorum, ni fallor, dividì potuerit.

II. TRIANDRIA.

410. EMPETRUM. *G. Pl.* 515. *Juss.* 162.
Gærtn. t. 106.

Masc. *Cal.* tripartitus. *Pet.* 3. *Stam.* capillaria, 3—9.

Fœm. *Cal.* tripartitus. *Pet.* 3. *Stigmata* 9.
Bacca supera, enneasperma.

nigrum. 1. EMPETRUM caulibus procumbentibus.

Empetrum nigrum. *Sp. Pl.* 1450. *Huds.* 431.
With. 176. *Hull.* 220. *Dicks.* *H. Sicc.*
fasc. 2. 10. *Engl. Bot. t.* 526. *Mill. Illustr.*
t. 86. *Fl. Dan. t.* 975.

E. montanum, fructu nigro. *Raii Syn.* 444.

Erica. *Matth. Valgr.* v. 1. 139.

E. baccifera. *Camer. Epit.* 77.

E. baccifera procumbens. *Ger. em.* 1383.

Angl. Black Crow- or Crake-Berry.

In ericetis montofisis elatioribus borealibus copiose.

Frutex.

Frutex. Fl. Maio.

Fruticulus depressus, ericoides, spithamæus, ramosus, nitidus, ramulis erectis, foliosis, rubicundis. Folia sparsim imbricata, sœpè verticillata, patentiuscula, brevissimè petiolata, oblonga, obtusa, revoluta, marginibus ciliata, cæterum glaberrima. Flores apicem versùs raniorum præcedentis anni, axillares, solitarii, subfessiles, dioici, rariùs hermaphroditici vel monoici, carnei. Petala tria, vel subindè plura, elliptica. Stamina tenuissima, plerumque tria, quandoque novem. Antheræ rubræ. Germen orbiculato-depressum. Stylus brevis, simplex. Stigmata 9, radiata. Bacca globosa, nigra. Semina 9, in orbem digesta.

411. RUSCUS. G. Pl. 534. Juss. 42.

Gærtn. t. 16.

Masc. *Cal.* hexaphyllus. *Cor.* nulla. *Nectarium* ovatum, tubulosum, intùs staminiferum.

Fœm. *Cal.* *Cor.* et *Nect.* maris. *Stam.* nulla.

Stylus 1. *Bacca* supera, trilocularis. *Sem.* gemina.

1. RUSCUS foliis mucronato-pungentibus suprà aculeatus.
floriferis nudis.

Ruscus aculeatus. Sp. Pl. 1474. Huds. 437.

With. 68. Hull. 223. Relb. 374. Sibth. 22.

Engl. Bot. t. 560. Woodv. Suppl. t. 237.

Mill. Illustr. t. 96.

Ruscus. Raii Syn. 262. Ger. em. 907. Matth.

Valgr. v. 2. 555. Camer. Epit. 935. Trag.

Hist. 919.

β. R. laxus. Tr. of Lim. Soc. v. 3. 334.

Angl. Butcher's Broom.

In ericetis dumosis, et nemoribus, præcipue solo
glareoso.

β. at Stoke, near Gosport, plentifully. Mr.
G. Caley.

Perennis.

Perennis. Fl. Martio, Aprili.

Radix ramosa, multiceps, perennis. Herba valde rigida, biennis, sempervirens, erecta, ramosissima, glabra. Folia alterna, sessilia, elliptico-ovata, integerrima, nervosa, dura, mucronato-pungentia, atro-viridia, supra, medium versus, florifera. Flores sessiles, parvi, albivirentes, nectario pallide violaceo, ore luteolo. Antheræ sessiles. Bacca globosa, miniata, glabra. Semina sæpius duo, magna, cornea.

Var. β gaudet raniis laxioribus, foliis ellipticis, utrinque attenuatis. Hujusc specimina varia, e solo natali, communicavit D. Caley, in quibus varietatis tantum, nec distinctæ speciei, characteres lubenter agnoscō.

III. TETRANDRIA.

412. VISCUM. G. Pl. 517. Juss. 212.
Gærtn. t. 27.

Masc. *Cal.* nullus. *Pet.* 4, basi dilatata, con-nata, calycina. *Antheræ* sessiles, petalis adnatæ.

Fœm. *Cal.* submarginatus. *Pet.* 4, basi dilatata. *Stylus* nullus. *Bacca* infera, monosperma.

album. 1. VISCUM foliis lanceolatis obtusis, caule di-chotomo, capitulis axillaribus.

Viscum album. Sp. Pl. 1451. Huds. 431.
With. 203. Hull. 220. Relb. 368. Sibth. 63.
Abbot. 214. Mill. Illustr. t. 87. Woodv.
Suppl. t. 270.

Viscum.

Viscum. *Raii Syn.* 464. *Ger. em.* 1350. *Matth.*
Valgr. v. 2. 161. *Camer. Epit.* 555.
Angl. Mistletoe.

In arboribus parasitica.

Frutex. Fl. Maio.

Caulis ramosissimus, dichotomus, articulatus, teres, lœvis, pallidè virens. Folia colore caulis, opposita, lingulata, integerrima, lœvia, coriacea. Capitula axillaria, sessilia, pauciflora. Flores luteo-virentes, receptaculo communi carnoſo inserti. Baccæ magnitudine pisi, globosæ, margaritaceæ, succo viscosissimo subdulci repletæ.

413. HIPPOPHAE. *G. Pl.* 517. *Juss.* 75.
Gærtn. t. 42.

Masc. *Cal.* bipartitus. *Cor.* nulla.

Fœm. *Cal.* tubulosus, bifidus. *Stylus* 1. *Bacca* supera, monosperma. *Semen* arillo dupli- teclum.

1. HIPPOPHAE foliis lanceolatis.

rhamnoi-

Hippophae rhamnoides. *Sp. Pl.* 1452. *Huds. aes.*

431. *With.* 204. *Hull.* 220. *Engl. Bot.*

t. 425. *Pall. Roff.* v. 1. t. 68. *Fl. Dan.* t. 265.

Wilden. Baumz. 142.

Rhamnoides fructifera, foliis salicis, baccis leviter flavescentibus. *Raii Syn.* 445.

Rhamnus secundus Clufii. *Ger. em.* 1334.

Rhamni species. *Camer. Epit.* 81.

Angl. Sallow Thorn. Sea Buckthorn.

In aggeribus arenosis maritimis.

On various parts of the sandy eastern coast, from Kent to Yorkshire. Plentiful on the cliffs between Yarmouth and Cromer, Norfolk.

Arluscula. Fl. Maio.

Rami

Rami divaricati, duri, rigidi, apice spinoscentes; juniores rubiginosi. Folia sparsa, brevius petiolata, lanceolata, integerrima; supra saturate viridia; subtus squamosa, argentea, subrubiginosa. Flores axillares, ad folia primordialia sive inferiora, solitarii, virides. Antheræ flavæ, subsessiles. Stigma obliquum, virens, exsertum. Baccæ globosæ, vel subellipticæ, mucronulatæ; fulvæ, squamis stellatis conspersæ, acidæ, edules, sed ingratæ. Semen oblongum, nigrum, nitidum, arillo duplice, coriaceo, fusco, vestitum.

Frutex ob foliorum pulchritudinem in hortis sœpè colitur, at baccas nunquam nisi in solo natali persicit.

414. MYRICA. *G. Pl.* 518. *Juss.* 409.
Gærtn. t. 39.

Masc. *Cal.* amenti squama concava. *Cor.* nulla.
Fœm. *Cal.* amenti squama concava. *Cor.* nulla.
Styli 2. *Bacca* monosperma.

Gale. 1. MYRICA foliis lanceolatis subserratis, caule fruticoso.

Myrica Gale. *Sp. Pl.* 1453. *Huds.* 432.
With. 208. *Hull.* 220. *Kell.* 368. *Engl.*
Bot. t. 562. *Fl. Dan.* t. 327. *Wilden.*
Baumz. 198.

Gale frutex odoratus septentrionalium, Eleagnus Cordo. *Raii Syn.* 443. *Baub. Hist.* v. 1.
p. 2. 224.

Eleagnus. *Cord. Hist.* 212. 2.

Myrtus brabantica, sive Eleagnus Cordi. *Ger.*
em. 1414.

Angl. Sweet Gale. Dutch Myrtle.

In palustribus, imprimis solo glareoso.

Frutex. Fl. Maio.

Frutex ramosissimus, erectus, tripedalis, odoratus, ramulis subflexuosis, teretibus, glabris. Folia alterna, subsessilia, lanceolata, acuta, glabriuscula, apicem versus ferrata,

ferrata, basi attenuata. Amenta ex axillis foliorum præcedentis anni, sessilia, ovata, rubicunda, scariosa. Baccæ parvæ, extus resinoso-punctatae, valde fragrantes.

Variat floribus monoicis. *D. Templeton.*

IV. PENTANDRIA.

415. HUMULUS. *G. Pl.* 522. *Juss.* 404.

Masc. *Cal.* pentaphyllus. *Cor.* nulla. *Anth.*
apice biporosæ.

Fœm. *Cal.* amenti squama obliqua, integra.
Cor. nulla. *Styli* 2. *Semen* unicum, tunicatum.

I. HUMULUS.

Lupulus.

Humulus Lupulus. *Sp. Pl.* 1457. *Huds.* 433.
Witb. 277. *Hull.* 220. *Relb.* 369. *Sibth.* 90.
Abbot. 214. *Engl. Bot.* t. 427. *Mill. Illustr.*
t. 88.

Lupulus mas et fœmina. *Raii Syn.* 137. *Camer.*
Epit. 933, 934.

Lupulus. *Matth.* *Valgr.* v. 2. 553.

Lupus salictarius. *Gér. em.* 885. *Fuchs. Hist.*
164.

Angl. Hop.

In dumetis et sepibus, præcipue solo tenaci humidi sculo.

Perennis. Fl. Julio.

Radix ramosa, multiceps. Caules annui, scandentes,
VOL. III. M longissimi,

longissimi, ramosi, angulati, spinuloso-asperi. Folia opposita, petiolata, cordata, ferrata, integra vel lobata, scabra. Stipulæ intrafoliaceæ, integerrimæ, reflexæ, glabré. Flores axillares, bracteati; masculi paniculati, luteo-albidi, cernui; amenta fœminea solitaria, pendula, ovata, pallidè viridia, aromaticæ, amara, demùm scariosa. Semen tunicâ membranaceâ, hinc fissâ, vestitum.

V. HEXANDRIA.

416. TAMUS. *G. Pl.* 524.
Tamnus. Juss. 43.

Masc. *Cal.* sexpartitus. *Cor.* nulla.

Fœm. *Cal.* sexpartitus. *Cor.* nulla. *Stylus* trifidus. *Bacca* infera, trilocularis. *Sem.* bina.

commu- 1. TAMUS foliis cordatis indivisis.

nis. Tamus communis. *Sp. Pl.* 1458. *Huds.* 433.
With. 344. *Hull.* 221. *Relb.* 370. *Sibth.* 117.
Abbot. 214. *Engl. Bot.* t. 91. *Mill. Illustr.*
t. 89.

Tamnus racemosa, flore minore luteo-pallecente.

Raii Syn. 262.

Bryonia nigra. *Ger. em.* 871.

Vitis nigra. *Matth. Valgr.* v. 2. 622. *Camer.*
Epit. 988.

Angl. Black Bryony.

In sylvis et sepibus umbrosis.

Perennis. Fl. Junio.

Radix

Radix carnosa, crassa, extus nigricans. Caules annui, scandentes, volubiles, ramosi, teretiusculi, laeves, longissimi. Folia alterna, longius petiolata, cordata, acuta, integerrima, nervosa, glaberrima. Stipulae parvae, acutae, reflexae. Racemi axillares; maleuli longiores. Flores parvi, virescentes. Baccæ fulvæ, insipidæ. Semina ovata, fusca, laevia.

VI. OCTANDRIA.

417. POPULUS. G. Pl. 526. Juss. 409.
Gærtn. t. 90.

Masc. Cal. amenti squama lacera. Cor. turbinata, obliqua, integra.

Fœm. Cal. amenti squamia lacera. Cor. turbinata, integra. Stigmata 4. Caps. supera, bilocularis, bivalvis. Sem. papposa.

1. POPULUS soliis cordato-subrotundis lobatis *alba*.
dentatis subtus tomentoso-niveis, amentis
ovatis.

Populus alba. Sp. Pl. 1463. Hudf. 433. With.
375. Hull. 221. Relb. 370. Sibth. 126.
Abbæt. 215. Lightf. 616. Raii Syn. 446.
Ger. em. 1486. Baub. Hist. v. 1. p. 2. 160.
Matth. Valgr. v. 1. 123. Camer. Epit. 65.

P. *alba* β nivea. Ait. Hort. Kew. v. 3. 405.

P. *major*. Mill. Diet. ed. 8. n. 4.

P. *nivea*. Wilden. Baumz. 227.

Angl. Great White Poplar. Abele-tree.

In sylvis humidis ; quandoque in montosis sicci-
oribus.

Arbor. Fl. Martio.

Cortex lœviusculus. Rami horizontales. Ramuli tomentoso-nivei. Folia alterna, petiolata, cordato-subrotunda, obtusè acuminata, plūs minùs lobata et angulata, obsoletè dentata, venosa ; suprà atro-viridia, glabrata, nitida ; subtùs densissimè tomentoso-nivea : in ramulis junioribus et valdè luxuriantibus ferè palmata. Petioli foliis triplò breviores, lanati. Stipulæ in ramulis junioribus tantùm inveniuntur, linearis-lanceolatæ, subfalcatae, obsoletè dentatæ, subtùs lanatæ. Amenta uncialia vel paulò longiora, ovata, obtusa, densa, squamis nitidis, lutescentibus, subpilosus. Stigmata linearia, glabra, luteo-viridia.

canescens. 2. **POPULUS** foliis subrotundis angulato-repandis
dentatis subtùs tomentoso-incanis, amentis
cylindraceis laxis.

P. alba. Mill. Diet. ed. 8. n. 1. Wilden.
Baumz. 227.

P. alba α , canescens. Ait. Hort. Kew. v. 3.
405.

P. nigra. Mill. Illustr. t. 90.

P. alba foliis minoribus. Raii Syn. 446. Ger.
em. 1487. Bauh. Hist. v. 1. p. 2. 160. Lob.
Ic. v. 2. 193.

Angl. Common White Poplar. Grey Poplar.

Ad fluvios et in aquofis.

Arbor. Fl. Martio.

Radix latissimè repens. Cortex lœvis, cinerascens, nitidus. Rami adscendentes. Ramuli tomentosi, cinerei. Folia ovato-orbiculata, repanda nec lobata, inæqualiter ac sinuato-dentata, venosa ; suprà lœtè viridia, glabra, nitidiuscula ; subtùs tomentoso-incana, cinerascentia, nec nivea, quandoque glabrata. Petioli longitudine ferè prioris. Stipulæ linearis-lanceolatæ, falcatae, dentatae,

tatæ, canescentes. Amenta ferè biuncialia, cylindracea, laxa, squamis fuscis, ciliato-pilosissimis.

3. *POPULUS* foliis suborbiculatis dentatis utrin- *tremula*.
que glabris, petiolis compressis, ramulis hirtis.
Populus tremula. *Sp. Pl.* 1464. *Huds.* 434.
With. 375. *Hull.* 221. *Relb.* 371. *Sibth.* 126.
Abbot. 215. *Wilden. Baumz.* 228.
P. Lybica. *Raii Syn.* 446. *Ger. em.* 1487. *Baub.*
Hift. v. 1. p. 2. 163. *Matth. Valgr.* v. 1. 125.
Camer. Epit. 67.
Angl. Asp. Aspen. Trembling Poplar.

In sylvis uliginosis.

Arbor. Fl. Martio, Aprili.

Arbor excelsa, lævis. Ramuli glabri; juniores luxuriantes hirti, nec incani. Folia, e petiolis elongatis compressis rigidulis, tremula, ferè orbiculata, mucronulata, dentato-repanda, basi rotundata et integerrima, utrinque glabra, venosa. Stipulæ setaceæ, hirtæ, deciduæ. Amenta biuncialia, ovato-cylindracea, hirsuta, squamis fuscis, pilis canis sive lutescentibus. Stigmata sanguinea.

4. *POPULUS* foliis deltoidibus acuminatis fer- *nigra*
ratis utrinque glabris.

Populus nigra. *Sp. Pl.* 1464. *Huds.* 434.
With. 376. *Hull.* 221. *Relb.* 371. *Sibth.* 126.
Abbot. 215. *Raii Syn.* 446. *Ger. em.* 1486.
Matth. Valgr. v. 1. 124. *Camer. Epit.* 66.
Wilden. Baumz. 229.
Angl. Black Poplar.

Ad fluvios et in aquofis.

Arbor. Fl. Martio.

Arbor excelsa, glabra. Petioli foliis duplò breviores, subindè hirsuti. Folia saturatè viridia, glaberrima, deltoidea, plūs minùs acuminata, ferrata; basi integrina

gerrima et subindè producta. Amenta pedunculata,
squamis glabris.

Stamina 16. *Linn. Leers.*

418. RHODIOLA. *G. Pl.* 526. *Juss.* 307.

Masc. *Cal.* quadripartitus. *Pet.* 4. *Nectaria* 4,
emarginata.

Fœm. *Cal.* quadripartitus. *Pet.* 4. *Nect.* 4.
emarginata. *Pistilla* 4. *Caps.* 4, polyspermæ.

rosea.

I. RHODIOLA.

Rhodiola rosea. *Sp. Pl.* 1465. *Huds.* 434.
With. 389. *Hull.* 221. *Lightf.* 619. *Engl.*
Bat. t. 508. *Fl. Dan.* t. 183.

Anacampseros radice rosam spirante major. *Raii*
Syn. 269.

Rhodia radix. *Ger. èm.* 532. *Matth. Valgr.*
v. 2. 372. *Camer. Epit.* 769.

Angl. Rose-root. *Pet. H. Brit.* t. 42. f. 2.

In fissuris rupium alpinarum, quandoque in præ-
ruptis maritimis.

Plentiful on the highest mountains of Wales and
the north of Yorkshire. *Ray.* On almost all
the Highland Alps, and very frequently by the
sea-side upon bold rocky shores of the Hebrides.
Lightf. On the northern coast of Ireland.
Mr. Templeton. On Maze Beck Scar, West-
moreland. *Rev. Mr. Harriman and Mr. Oliver.*

Perennis. Fl. Maio, Junio.

Radix carnosa, exsiccata odore aquæ destillatæ rosarum
spargens. *Caules* simplicissimi, erecti, spithamæi,
dense foliosi. *Folia imbricata*, sessilia, obcvata, acuta,
carnosa, glauca, lævia, apicem versùs dentata. *Cyma*
terminalis, sessilis, multiflora. *Flores* flavi; masculi
pistillorum, fœminei staminum, rudimentis sœpè gau-
dentes. *Calyces* et petala fœniineorum minores,
rubroque punctati. *Variat* floribus quinquefidis.

VII. EN-

VII. ENNEANDRIA.

419. MERCURIALIS. *G. Pl.* 527. *Juss.* 385.
Gærtn. t. 107.

Masc. *Cal.* tripartitus. *Cor.* nulla. *Stam.* 9—12.
Antæræ globosæ, didymæ.

Fœm. *Cal.* tripartitus. *Cor.* nulla. *Styli* 2.
Caps. dicocca, bilocularis. *Sem.* solitaria.

1. MERCURIALIS caule simplicissimo, foliis *perennis.*
scabris, radice repente.

Mercurialis perennis. *Sp. Pl.* 1465. *Huds.* 435.
With. 392. *Hull.* 222. *Relb.* 372. *Sibth.* 133.
Abbot. 216. *Curt. Lond. fasc.* 2. *t.* 65. *Mill.*
Illuſtr. *t.* 91. *Fl. Dan.* *t.* 400.

M. perennis repens, Cynocrambe dicta. *Raii*
Syn. 138.

Cynocrambe. *Matth. Valgr.* *v.* 2. 635. *Camer.*
Epit. 998, 999. *Ger. em.* 333. *Fuchs.*
Hift. 444.

Dog's Mercury. *Pet. H. Brit.* *t.* 1. *f.* 5, 6.
Angl. Perennial Mercury.

In dumetis vulgaris.

Perennis. Fl. Aprili, Maio.

Radix repens, albida, ramosissima. Caules erecti, sim-
plicissimi, pedales aut sesquipedales, teretes, apicem
versus densius foliosi. Folia opposita, petiolata, ovata,
acuta, crenata, piloso-scabra. Stipulae intrafoliaceæ,
membranaceæ, acutæ, integerrimæ, reflexæ. Racemi
axillares, pedunculati. Flores virides, semper dioici.
Squama lanceolata utrinque ad latus germinis. Cap-
fulæ piloso-scabræ.

- annua.* 2. MERCURIALIS caule brachiato, foliis glabris,
floribus racemosis, radice fibrosa.
Mercurialis annua. *Sp. Pl.* 1465. *Huds.* 435.
With. 392. *Hull.* 222. *Relb.* 372. *Sibth.* 134.
Curt. Lond. fasc. 5. t. 68. *Engl. Bot.* t. 559.
M. annua glabra vulgaris. *Raii Syn.* 139.
M. mas et fœmina. *Ger. em.* 332. *Matth.*
Valgr. v. 2. 633, 634. *Camer. Epit.* 996,
997. *Fuchs. Hist.* 475, 476.
French Mercury. *Pet. H. Brit.* t. 1. f. 7, 8.
Angl. Annual Mercury.

In ruderatis et cultis suburbanis.

Very frequent about London, Norwich, and other towns.

Annua. Fl. Julio, Septembri.

Radix fibrosa. Herba lætè virens, lævis, nitida, ramo-sissima, erecta, fœtida. Folia ovata, acuta, ferrata. Racemi ut in priore. Squamæ ad latera germinis nullæ. Capsula muricata.

Flores aliquot fœmineos inter masculinos aliquando in-venerunt D. D. Robson, Sowerby, aliique.

420. HYDROCHARIS. G. Pl. 527. Juss. 67.

Masc. *Cal.* trifidus. *Pet.* 3. *Filamenta tria*
interiora appendiculata.

Fœm! *Cal.* trifidus. *Pet.* 3. *Styli 6.* *Caps.*
sexlocularis, polysperma, infera.

Morsus 1. HYDROCHARIS.

- ranæ.* Hydrocharis *Morsus ranæ.* *Sp. Pl.* 1466.
Huds. 436. *With.* 394. *Hull.* 222. *Relb.* 373.
Sibth. 135. *Abbot.* 216. *Curt. Lond.* fasc. 3.
t. 64. *Fl. Dan.* t. 878. *Engl. Bot.* t. 808.
Stratiotes foliis Afari, semine rotundo. *Raii*
Syn. 290.

Morsus

Morsus Ranæ. *Ger. em.* 818.

Nymphæa parva. *Mattb. Valgr.* v. 2. 247.

Camer. Epit. 636.

Angl. Common Frog-bit.

In fossis et aquis pigroribus.

Perennis. Fl. Julio.

Herba natans, stolonifera, glabra, acaulis. Folia petiolata, reniformia, integerrima, lævia, subtus purpurascenscentia. Pedunculi fœminei simplicissimi, uniflori, basi vaginati; masculini umbellati, 3- aut 4-flori, bracteis binis ad basin pedicellorum. Flores albi, petalis majusculis, rotundatis, corrugatis, basi luteis.

Variat flore pleno, odoratissimo. *Raius.*

Genus Stratotæ, p. 579, proximum.

VIII. MONADELPHIA.

421. JUNIPERUS. *G. Pl.* 531. *Juss.* 413.
Gærtn. t. 91.

Masc. *Cal.* amenti squamæ. *Cor.* nulla. *Stam.* 3.

Fœm. *Cal* amenti squamæ pauciores, demum carnosæ, coalitæ in *baccam* trispermam.

1. JUNIPERUS foliis ternis patentibus mucro-communato-spinosis baccâ longioribus.

mis.

Juniperus communis. *Sp. Pl.* 1470. *Huds.* 436.

With. 599. *Hull.* 222. *Lightf.* 623. *Rehb.* 373.

Sibth. 210. *Woodv. Med. Bot.* t. 95. *Mill.*

Illustr. t. 95. *Pall. Roff.* v. 1. p. 2. 12. t. 54.

Engl. Bot. t. 1100.

J. vulgaris,

- J. vulgaris*, baccis parvis purpureis. *Raii Syn.* 444. *Baub. Hist.* v. 1. p. 2. 293.
Juniperus. *Ger. em.* 1372. *Matth. Valgr.* v. 1. 109. *Camer. Epit.* 53.
J. minor. *Fuchs. Hist.* 78.
 β . *J. alpina*. *Raii Syn.* 444. *Baub. Hist.* v. 1. p. 2. 301.
J. alpina minor. *Ger. em.* 1372.
Juniperus, baccæ, cacumen. *Pharmac. Lond.* *Angl.* Common Juniper.

In ericetis et montofis calcareis.

β in alpibus.

Frutex. Fl. Maio.

Frutex humilis, ramosissimus, rigidus, glaber, semper-virens. Succus proprius totius plantæ terebinthinus. Folia acerosa, opposita, sessilia, canaliculata, integrifolia, mucronato-pungentia, supra glauca. Amenta axillaria, sessilia, solitaria, ovata, parva, bracteata; masculina flava, polline copiosissimo, deinde fusca; fœminea luteo-viridia, minora. Baccæ globosa, atropurpurea cum rore glauco, subdulcis, aromaticæ, e squamis amenti omnino coalitis, pulposis, composita. Semina tria, angulata.

Varietas *alpina* β humilior est, atque magis depressa, foliis latioribus, baccis subellipticis.

422. TAXUS. *G. Pl.* 532. *Juss.* 412.
Gærtn. t. 91.

Masc. *Cal.* nullus. *Cor.* nulla. *Stam.* plurima.
Anth. peltatæ, octofidæ.

Fœm. *Cal.* urceolatus, integerrimus. *Stylus* nullus. *Sem.* 1, calyci baccato impositum.

baccata. 1. TAXUS foliis approximatis.

Taxus baccata. *Sp. Pl.* 1472. *Huds.* 437.
 4 *With.*

With. 614. *Hull.* 222. *Rehb.* 374. *Sibth.* 216.
Engl. Bot. t. 746. *Wilden. Baumz.* 382.
Taxus. *Raii Syn.* 445. *Ger. em.* 1370. *Matth.*
Valgr. v. 2. 444. *Camer. Epit.* 840.
Angl. Yew Tree. *Hunt. Evel. Sylv. tab.*

In sylvis montosis et clivis calcareis.

Arbor. Fl. Martio, Aprili.

Caudex rectus, cortice deciduo, lævi. Rami horizontales, distichi. Folia sparsa, subfessilia, linearia, mucronulata, integerrima, subrevoluta, atro-viridia, glabra, nitida, sempervirentia. Flores axillares, sparsi, fessiles, solitarii, bracteis imbricatis, membranaceis, interioribus majoribus; masculini calyce planè destituti, staminibus receptaculo communi, columnari, affixis; fœminei calyce monophyllo, integerimo, lævi, demum ampliato, succoso, pulcherrimè rubro, semen amplectente.

Classis

Classis XXIII.

POLYGAMIA.

Flores hermaphroditici et masculi
aut fœminei, structurâ partium
inter se diversi, in eadem specie.

MONOECIA.

Fl. hermaphr. et masc. aut fœm.
in eadem planta.

423. ATRIPLEX. Hermaphr. *Cal.* quinque-partitus, inferus. *Cor.* nulla. *Stam.* 5. *Stylus* bipartitus. *Sem.* 1, depresso.

Fœm. *Cal.* diphyllus. *Cor.* nulla. *Stylus* bipartitus. *Sem.* 1, compressum.

XXIII. POLYGAMIA.

I. MONOECIA.

423. ATRIPLEX. G. Pl. 545. Juss. 85.
Gærtn. t. 75.

Hermaphr. Cal. quinquepartitus, inferus. Cor. nulla. Stam. 5. Stylus bipartitus. Sem. 1, depresso.

Fœm. Cal. diphyllus. Cor. nulla. Stylus bipartitus. Sem. 1, compressum.

portula- 1. ATRIPLEX caule fruticoso, foliis oppositis obovato-lanceolatis, floribus submonoicis.
caoides.

Atriplex portulacoides. Sp. Pl. 1493. Huds. 442. With. 274. Hull. 226. Relb. 378. Engl. Bot. t. 261.

A. maritima fruticosa, Halimus et Portulaca marina dicta, angustifolia. Raii Syn. 153.

Halimus vulgaris. Ger. em. 523. Matth. Valgr. v. 1. 145.

Sea Purslane. Pet. H. Brit. t. 7. f. 7.
Angl. Shrubby Orache.

In maritimis cœniosis.

Fruticulus. Fl. Julio, Augusto.

Radix lignosa, multiceps, subrepens. Caules fruticosi, ramosi, adscendentis, sesquipedales, foliosi, tetragoni, pulvрulento-incani, uti tota planta. Folia opposita, petiolata, obovata vel ferè lanceolata, integerrima, basi attenuata. Flores glomerati, spicati, terminales, flavescentes, omnes ferè masculi aut fœminei, vix aut ne vix hermaphroditici. Calyx floris fœminei demum auctus, deltoideus, basi rotundatus, margine trilobus, clausus.

lacinia- 2. ATRIPLEX caule herbaceo patente, foliis deltoidibus sinuato-dentatis subtus argenteis.
ta. Atriplex

Atriplex laciniata. *Sp. Pl.* 1494. *Huds.* 442.
With. 274. *Hull.* 226. *Lightf.* 636. *Dicks.*
H. Sicc. fasc. 4. 15. *Engl. Bot.* t. 165.

A. maritima. *Raii Syn.* 152. *Baub. Hist.*
v. 2. 974.

A. marina. *Matth. Valgr.* v. 1. 421. *Camer.*
Epit. 244. *Ger. em.* 325.

β . *A. maritima* nostras procerior, foliis angulosis
 incanis, admodum sinuatibus. *Pluk. Almag.* 60.
Dill. in Raii Syn. 152.

Angl. Frosted Sea Orache. *Pet. H. Brit.* t. 7.
 f. 3.

In maritimis arenosis haud vulgaris.

In Essex. *Ray.* At Harwich. *Mr. Francis Smith.*
 At New Haven near Edinburgh.

Annua. Fl. Julio.

Radix fibrosa. Caules patuli, ramosissimi, sæpius pro-
 cumbentes, teretiusculi, foliosi. Folia alterna, petio-
 lata, deltoideo-rhombea; basi angustata, integerima;
 anticè inæqualiter dentato-sinuata; utrinque, at im-
 primis subtus, argenteo-pulverulenta. Flores her-
 maphroditici in spicas densas terminales conglomerati,
 germinibus sæpius abortientibus; fœminei axillares,
 pedicellati, conferti, calyce demum magno, rhombeo,
 acutè denticulato, pulverulento, basi venoso, semine
 compreso, subcochleato.

Varietas β , nota mihi nomine tantum, distinctissima ab
A. tatarica Linnæi videtur.

3. ATRIPLEX caule herbaceo patulo, foliis del- *patula*,
 toideo-lanceolatis subhastatis, calycibus semi-
 num disco submucicatis.

Atriplex patula. *Sp. Pl.* 1494. *Engl. Bot.* t. 936.
A. hastata. *Huds.* 443. *With.* 274. *Hull.* 226.
Relb. 379. *Sibth.* 90. *Abbot.* 219. *Curt.*
Lond. fasc. 2. t. 66. *Lightf.* 636.

A. syl-

- A. *sylvestris*, folio hastato seu deltoide. *Raii*
Syn. 151.
- A. folio deltoide sinuato et mucronato, hastæ
 cuspidi simili. *Moris. sect. 5. t. 32. f. 14.*
- A. *sylvestris vulgaris*. *Ger. em. 326.*
 Delt Orrach. *Pet. H. Brit. t. 7. f. 1.*
- β. *Atriplex maritima perennis*, folio deltoide seu
 triangulari, minùs incano. *Raii Syn. 152.*
- γ. *Atriplex maritima*, ad foliorum basin auriculata,
 procumbens et ne vix sinuata. *Dill. in Raii*
Syn. 152.
- A. *marinæ species Valerando*. *Baub. Hist. v. 2.*
 974. *Chabr. Hist. 306. f. 4.*
 Delt Sea Orrach. *Pet. H. Brit. t. 7. f. 2.*
 Angl. Spreading Halberd-leaved Orache.
- In ruderatis et cultis vulgaris. β et γ in maritimis
 arenosis.

Annua. Fl. Junio—Augusto.

- Radix fibrosa, omnino annua, nec perennis. Caulis di-
 varicato-patulus, ramosissimus, erectiusculus; in β et
 γ procumbens, sèpè rubicundus. Folia alterna, lon-
 giùs petiolata, subtùs plerumque pulverulenta; inferiora
 deltoidea, hastata, sinuato-dentata; superiora angus-
 tiora, lanceolata et ferè integerrima; omnia in γ car-
 nosa, rubicunda, minùsque dentata. Racemi terminales
 et axillares, foliosi. Valvulae calycum fœmineorum
 deltoideo-rhombeæ, acutæ, margine angulum versus
 denticulatae, disco hinc vel utrinque tuberculato-muri-
 catæ. Semen magnum, subcochleatum, punctulatum.
- A. *hastata Linnæi* abundè discrepat, valvulis fructùs
 maximis, reticulato-venosis, submembranaceis, mar-
 gine sinuato-dentatis, dentibus apice setaceis.

angusti- 4. ATRIPLEX caule herbaceo divaricato, foliis
folia. lanceolatis integerimis; infimis subhastatis,
calycibus seminum hastatis læviusculis.

Atriplex

- Atriplex patula.* *Huds.* 443. *With.* 275.
Hull. 226. *Rehb.* 379. *Sibth.* 90. *Abbot.* 219.
Lightf. 637.
- A. sylvestris angustifolia.* *Raii Syn.* 151. *Ger.*
em. 326.
- A. sylvestris Polygoni aut Helxines soliis.* *Lob.*
Ic. 257.
- A. sylvestris prima.* *Matth. Valgr.* v. 1. 418.
- A. vulgaris angustifolia.* *Bauh. Hist.* v. 2. 973.
Spear Orrach. *Pet. H. Brit.* t. 7. f. 5.
Angl. Spreading Narrow-leaved Orache.

In ruderatis et cultis vulgaris.

Annua. Fl. Junio—Augusto.

Habitus præcedentis, at magis divaricata. Folia brevijs petiolata, omnia ferè lanceolata, integerrima, horizontalia; infima latiora, et subindè hastata, nec dentata. Valvulæ seminiferæ hastatæ, deltoideæ; apice elongato-lanceolatæ, acutæ; margine integerrimæ; disco vix unquam muricatæ. Semen duplò minus quam in præcedente, vix punctulatum.

5. **ATRIPLEX** caule herbaceo erecto, foliis ova-*erecta*, to-lanceolatis; inferioribus sinuatis, calycibus seminum undique muricatis.

Atriplex erecta. *Huds. ed.* 1. 376.

A. patula ♂. *Huds. ed.* 2. 444.

A. angustifolia laciniata. *Dill. in Raii Syn.* 152
Angl. Upright Spear-leaved Orache.

In ruderatis.

On the entrance into Battersea field from Nine Elms; Prof. Martyn senior. *Dill.* In Mr. Rose's herbarium.

Annua. Fl. Augusto.

Caulis ramosus, erectus, firmus. Folia petiolata, sub-pulverulenta, ovato-lanceolata, acuta; superiora ferè integerrima;

integerima; inferiora sinuato-dentata. Racemi terminales, compositi, multiflori, subaphylli. Valvulae seminiferæ triplè minores quam in præcedente, deltoideæ, acutæ, gibbosæ, disco utrinque muricatæ dentibus numerosis, porrectis, validis. Semen ferè præcedentis, sed inius.

Distinctissima species, fructu parvo, maximè muricato, copiosissimo, facile recognoscenda.

littoralis. 6. ATRIPLEX caule herbaceo erecto, folijs omnibus linearibus integris dentatisve, calycibus seminum muricatis sinuatis.

Atriplex littoralis. Sp. Pl. 1494. Huds. 444.

With. 275. Hull. 226. Engl. Bot. t. 708.

A. maritima, Scopariæ folio. Dill. in Raii Syn. 153.

A. angustifolia maritima minima. Bocc. Sic. 30. t. 15. f. 1.

A. angustissimo et longissimo folio. Dill. in Raii Syn. 153.

A. maritima angustifolia, obtusiore folio. Dill. in Raii Syn. 153.

Grass Sea Orrach. Pet. H. Brit. t. 7. f. 6.

6. *Atriplex serrata,* Huds. 444.

A. angustifolia maritima dentata. Raii Syn. 152.

A. angustifolia laciniata minor. Baub. Hist. v. 2. 972.

Jagged Long Orrach. Pet. H. Brit. t. 7. f. 4.

Angl. Gras-leaved Sea Orache.

In palustribus maritimis cœnosis.

Annua. Fl. Augusto, Septembri.

Caulis erectus, ramosus, angulatus, foliosus, ramis erectis. Folia pectiolata, linearia, magis vel minus denticulata, subinde sinuato-dentata, longitudine et latitudine varia, subtus præcipue pulverulenta. Spicæ ovato-cylindraceæ, obtuse, dense, multifloræ, subaphyllæ.

aphyllæ. Valvulæ feminiferæ ovatæ, acutæ, margine inæqualiter sinuatæ, disco muricatæ.

7. ATRIPLEX caule herbaceo flexuoso divari-pedunculato, foliis obovatis integerrimis, floribus *lata*. fœmineis pedunculatis cuneiformibus.

Atriplex pedunculata. *Sp. Pl.* 1675. *Huds.* 444.
With. 276. *Hull.* 227. *Engl. Bot.* t. 232.
Fl. Dan. t. 304.

A. marina semine lato. *Raii Syn.* 153.

Halimus humiliš ercta, semine folliculis membranaceis bivalvibus, in latitudinem porrectis et utrinque recurvis, longo pedunculo insidentibus, clauso. *Pluk. Phyt.* t. 36. f. 1.

Heart Sea Purslain. *Pet. H. Brit.* t. 7. f. 8.

♂. *Atriplex maritima nostras*, *Ocymi minoris* folio.
Raii Syn. 153.

Angl. Pedunculated Sea Orache.

In maritimis cœniosis Angliæ orientalibus, rariū.

Near Boston; Plukenet. *Ray*. In the isle of Thanet, near the ferry, in great plenty. *Sherard*. Just below Lynn, and at Yarmouth.

♂. Near Lynn, Plukenet. *Ray*.

Annua. Fl. Augusto, Septembri.

Caulis spithameus, subindè pedalis, erectiusculus, ramosus, angulatus, flexuosus, foliosus. Folia subsessilia, patentia, obovata, integerrimâ, utrinque, ut et tota planta, pulverulento-incana. Flores terminales, racemoſi, congesti; fœminei pedunculati, deflexi, cuneiformes, compressi, trilobi, lobis lateralibus recurvatis, disco plani, æquales, demùm maturescente semine aucti, ut in reliquis speciebus.

Var. ♂ non nisi caule humiliori, magisque diffuso, differt.

Classis XXIV.

CRYPTOGAMIA.

Fructificatio anomala vel obscura.

FILICES.

Herba frondosa. Fructificatio vel dorsalis, vel terminalis, vel sub-radicalis.

* *Annulatæ*; capsulis annulo elastico, valvulis contrario, cinctis.

428. POLYPODIUM. *Fructificationes* in punctis subrotundis, sparsis. *Involucrum* nullum.
429. ASPIDIUM. *Fructif.* in punctis subrotundis, sparsis. *Invol.* umbilicatum.
430. ASPLENIUM. *Fructif.* in lineolis sparsis. *Invol.* costam versus dehiscens.
431. SCOLOPENDRIUM. *Fructif.* in lineolis geminis, interveniis. *Involucra* sibi invicem incumbentia.

432. BLECHNUM. *Fructif.* in lineis continuis, costæ adjacentibus. *Invol.* costam versus dehiscens.
433. PTERIS. *Fructif.* in linea marginali. *Invol.* e margine frondis inflexo, continuo.
434. ADIANTUM. *Fructif.* in punctis marginalibus. *Involucra* e margine frondis inflexo, interrupto.
435. CYATHEA. *Fructif.* sparsæ, subrotundæ, calyci apice dehiscenti insidentes.
436. HYMENOPHYLLUM. *Fructif.* margini frondis insertæ. *Involucra* bivalvia, exterius hiantia.

** *Exannulatæ.*

424. EQUISETUM. *Frons* amentacea. *Semina* filamentis 4, polliniferis, circumPLICATA.
425. OPHIOGLOSSUM. *Spica* disticha. *Capsulae* immersæ.
426. OSMUNDA. *Spica* ramosa. *Caps.* nudæ, globosæ.
427. LYCOPodium. *Capsulae* axillares, solitariæ, compressæ.
437. PILULARIA. *Receptaculum commune* quadriloculare, flores occultans. *Sem.* corticata.
438. ISOETES. *Recept.* commune intra basin frondis, uniloculare. *Sem.* scabra.

MUSCI.

Herba foliosa. Corolla calyptrifor-
mis, styligera, capsulam, demùm
pedicellatam, operculatam, te-
gens.

* *Peristomio nullo.*

441. PHASCUM. *Capsula clausa operculo non dehiscente.*
 439. SPHAGNUM. *Caps. ore nudo. Calyptra circumscissa, capsulam basi cingens.*
 442. GYMNSTOMUM. *Caps. ore nudo. Calyptra integra e basi secedens.*

** *Peristomio simplici.*

445. TETRAPHIS. *Caps. oblonga. Perist. dentibus 4, erectis, liberis.*
 444. ANDRÆA. *Caps. brevissima. Perist. dentibus 4, incurvatis, apice cohærentibus.*
 443. SPLACHNUM. *Caps. cylindrica, apophysī carnosæ insidens. Perist. dentibus 16, basi dilatatis.*
 446. ENCALYPTA. *Perist. dentibus 16, linearibus, erectis. Calyptra campanulata, laxa.*
 452. PTEROGONIUM. *Perist. dentibus 16, linearibus, erectis. Fl. axillares.*
 447. GRIMVIA. *Perist. dentibus 16, basi latioribus. Fl. terminales.*
 448. DICRANUM. *Perist. dentibus 16, inflexis, semibifidis.*

449. TRICHOSTOMUM. *Perist.* dentibus 32, filiformibus, rectiusculis, per paria vel approximatis, vel basi connexis.

450. TORTULA. *Perist.* dentibus filiformibus, spiraliter ac multipliciter convolutis.

Orthotrichum. Buxbaumia.

*** *Peristomio dupli.*

451. ORTHOTRICHUM. *Caps.* terminalis. *Perist.* exterius dentibus 16: interius dent. 8—16, filiformibus, quandoque nullum. *Calyptra* sulcata.

453. NECKERA. *Caps.* e perichætio lateral. *Perist.* exter. dentibus 16: inter. dentibus 16, filiformibus. *Calyptra* lævis.

456. FUNARIA. *Caps.* obovata. *Perist.* exter. dentibus 16, obliquis, apice cohærentibus: inter. dentibus 16, planis. *Calyptra* quadrangula.

440. BUXBAUMIA. *Caps.* obliqua, hinc gibba. *Perist.* exter. dentibus 16, brevissimis: inter. membranaceum, plicatum.

457. BARTRAMIA. *Caps.* sphærica, sulcata. *Perist.* exter. dentibus 16, subulatis: inter. membranaceum, laciniatum.

458. MNIUM. *Caps.* cylindracea, sulcata. *Perist.* exter. dentibus 16, fulcatis: inter. membranaceum, laciniatum.

459. BRYUM. *Caps.* ovato-oblonga, lævis. *Perist.* exter. dentibus 16, basi dilatatis: inter. membranaceum, dentatum. *Flores* terminales.

454. HYPNUM. *Caps.* ovato-oblonga. *Perist.* exter. dentibus 16, basi dilatatis: inter. mem-

membranaceum, dentatum. *Pedicellus* e
perichætio squamoso, laterali.

455. FONTINALIS. *Caps.* perichætio squamoso
obvoluta. *Perist. exter.* dentibus 16, basi
dilatatis: *inter.* cancellatum.
460. POLYTRICHUM. *Perist. exter.* dentibus
32—64: *inter.* membrana plana, indivisa.

HEPATICÆ.

Herba frondosa. Capsulæ operculo
destitutæ.

ALGAE.

Herba frondosa. Semina nidulantia.

FUNGI.

Herba nulla. Fructificatio in cor-
pore carnoso.

With. 754. *Hull.* 233. *Relb.* 403. *Sibth.* 264.

Abbot. 222. *Bolt. Fil.* 64. *t.* 35. *Raii*

Syn. 131. *Ger. em.* 1114. *Fl. Dan.* *t.* 1183.

Hippuris minor. *Trag. Hist.* 694.

$\beta.$ *Equisetum palustre*, tenuissimis et longissimis setis. *Dill. in Raii Syn.* 131.

$\gamma.$ *E. palustre minus*, *polystachion*. *Baub. Prodr.* 24. *Dill. in Raii Syn.* 131. *t.* 5. *f.* 3.

Angl. Marsh Horsetail.

In palustribus spongiosis.

Perennis. Fl. Junio, Julio.

Caulis profundè fulcatus, et indè angulatus, undique ramosus, apice florifer, ramis erectis, elongatis, simplicibus, articulatis, minute scabratiss, vaginis omnibus vircentibus, ore scarioso-albis. Amentum terminale, solitarium, nigricans, gracile, e squainis majoribus, paucioribus, demum laxè distantibus.

fluvia-
tile.

4. *EQUISETUM* caulis sterilibus ramosis : ramis scabriusculis numerosissimis ; fructificantibus simplicibus, vaginis approximatis setaceo-multifidis.

Equisetum fluviatile. *Sp. Pl.* 1517. *Huds.* 448.

With. 755. *Hull.* 233. *Relb.* 404. *Sibth.* 264.

Abbot. 222. *Bolt. Fil.* 66. *t.* 36, 37.

E. majus. *Raii Syn.* 130. *Ger. em.* 1113.

E. primum. *Dalech. Hist.* 1069.

E. palustre longioribus setis. *Baub. Theatr.* 241.

Angl. Great Water Horsetail.

In palustribus, aquosis, et ad fluvios.

Perennis. Fl. Aprili.

Caules floriferi præcociores, simplicissimi, ut in *E. arvensi*, sed vaginis approximatis, nec remotis, undique fuscis, margine setaceo-multifidis, amento maximo, terminali ; steriles ferè sexpedales, erecti, formosi, undique ramosi

ramosi seu verticillati, ramis patentissimis, elongatis, numerosissimis, oculo armato scabris, plerumque simplicibus, dentibus vaginalium nigris.

5. EQUISETUM caule nudiusculo ramisque lœ-limosum.
vibus, amento terminali.

Equisetum limosum. *Sp. Pl.* 1517. *Huds.* 448.
With. 755. *Hull.* 234. *Relb.* 404. *Sibth.* 264.
Abbot. 222. *Bolt. Fil.* 68. *t.* 38. *Engl. Bot.* *t.* 929.

E. fluviatile. *Fl. Dan.* *t.* 1184.

E. nudum lœvius nostras. *Raii Syn.* 131. *t. 5.*
f. 2.

β. E. foliis nudum, ramosum. *Raii Syn.* 132. *Bolt. Fil. præf.* 19.

Angl. Smooth Naked Horsetail.

In aquosis et palustribus frequens.

Perennis. Fl. Junio, Julio.

Caulis bipedales, glaberrimi, ramis paucis, sparsis, foliariis, erectis, glabriusculis, rarissime verticillatis, amento terminali, magno, fusco, ovato; steriles plurimque simplicissimi, apice attenuati.

Videtur species a præcedente distinctissima.

6. EQUISETUM caule nudo scaberrimo basi *hyemale*.
subramoso, vaginalis albidi basi apiceque nigra, amento terminali.

Equisetum hyemale. *Sp. Pl.* 1517. *Huds.* 448.
With. 756. *Hull.* 234. *Lightf.* 650. *Relb.* 404.
Abbot. 222. *Bolt. Fil.* 70. *t.* 39. *Engl. Bot.* *t.* 915.

E. n. 1679. *Hall. Hist.* *v. 3.* 3.

E. nudum. *Raii Syn.* 131. *Ger. em.* 1113.

Angl. Rough Horsetail. Shave-grass.

In palustribus et sylvis humidis.

In

In Warwickshire, Wiltshire and Lancashire. *Ray.*
Merret. Arminghall wood near Norwich. *Mr.*
Crowe. Hedenham near Bungay. *Mr. Stone.*
 Near Lanerk, Scotland. *Dr. Stokes.* Gamlingay
 bogs. *Relb.* Bedfordshire. *Abbot.* Near Ingleton and Halifax. *Bolt.* On St. Faith's bogs, Norwich, and between Lafwade and Meavis bank, Edinburgh.

Perennis. Fl. Julio, Augusto.

Caules sesquipedales, subglauci, sempervirentes, scaberrimi, fistulosi, omnino nudi, basi subramosi, vaginis brevibus, crenatis, albidis, basi margineque nigris, summa fimbriata. Amentum terminale, ovatum, nigrum.

Epidermis in hâc specie terrâ siliceâ abundat. *D. Davy.*

425. OPHIOGLOSSUM. G. Pl. 559. Juss. 14.

Spica disticha. *Capulæ* bivalves, immersæ, transversè dehiscentes.

vulga-
tum. 1. OPHIOGLOSSUM fronde ovatâ aeniâ.

Ophioglossum vulgatum. *Sp. Pl.* 1518. *Huds.*
 449. *Witb.* 761. *Hull.* 234. *Relb.* 405.
Sibth. 266. *Abbot.* 223. *Engl. Bot.* t. 108.
Bolt. Fil. 2. t. 3. *Dicks. H. Sicc. fasc.* 9. 18.
Fl. Dan. t. 147. *Hedw. Theor.* 44. t. 4.
 f. 20—23. *Roth. Germ.* v. 3. 36.

Ophioglossum. *Raii Syn.* 128. *Fuchs. Hist.* 577.
Matth. Valgr. 543. *Camer. Epit.* 364. *Ger.*
em. 404.

Angl. Common Adder's-tongue.

In pratis et pascuis humidis.

Perennis. Fl. Maio.

Radix e fibris simplicibus. Herba vix spithamæa, lœvis atque glaberrima, undique virens. Frons simplicissima, solitaria,

solitaria, petiolata, ovata, integerrima, avenia, basi spicifera. Spica pedunculata, erecta, lanceolata, acuta, compressa, subarticulata, multicapsularis, demum, ex utroque latere, per foramina capsularum dehiscentium, semina pulverulenta spargens.

Variat rarius spicâ bifidâ, aut geminâ.

426. OSMUNDA. *G. Pl.* 559. *Juss.* 15.

Spica ramosa. *Capsulæ* bivalves, nudæ, globosæ.

1. OSMUNDA fronde pinnatâ basi spiciferâ : *Lunaria*.
pinnulis lunulatis crenatis.

Osmunda Lunaria. *Sp. Pl.* 1519. *Huds.* 449.

With. 762. *Hull.* 234. *Relh.* 496. *Sibth.* 266.

Abbot. 223. *Engl. Bot.* t. 348. *Bolt. Fil.* 4.

t. 4. *Dicks. H. dicc. fasc.* 5. 17. *Fl. Dan.*

t. 18. *Roth. Germ.* v. 3. 32.

O. foliis lunatis. *Garid.* 345. t. 78.

Lunaria minor. *Raii Syn.* 128. *Ger. em.* 405.

Matth. Valgr. 254. *Camer. Epit.* 643. *Fuchs.*

Hist. 482.

Epimedium Dioscoridis. *Column. Phytob.* 80.

2. *Lunaria minor ramosa.* *Raii Syn.* 129. *Camer.*

Epit. 644.

γ. *L. minor foliis dissectis.* *Raii Syn.* 129.

L. racemosa minor, Matricariæ folio. *Breyn.*

Cent. 94.

Fl. Dan. t. 18.

Angl. Common Moonwort.

In pratis et pascuis, imprimis submontosis.

Perennis. *Fl. Junio.*

Radix e fibris simplicibus. Herba vix spithamæa, lœvis,

virens. Frons solitaria, pinnata, pinnis lunato-cuneatis,

inciso-crenatis, basi spicifera. Spica pedunculata,

erecta,

erecta, ramosa, ramulis distichis, utroque margine capsuliferis. Capsulæ globosæ, læves, fuscæ.
Variat frondibus spicisque multiplicatis (β), vel fronde decompositâ, spicâ solitariâ (γ).

regalis. 2. OSMUNDA fronde bipinnatâ apice spiciferâ : pinnulis cordato-lanceolatis glabris.

Osmunda regalis. *Sp. Pl.* 1521. *Huds.* 449.
With. 763. *Hull.* 235. *Relb.* 406. *Abbot.* 224.
Lightf. 653. *Engl. Bot.* t. 209. *Fl. Dan.*
t. 217. *Bolt. Fil.* 6. t. 5. *Roth. Germ.*
v. 3. 34.

Filix ramosa non dentata florida. *Raii Syn.* 125.

F. florida sive Osmunda regalis. *Ger. em.* 1131.

F. majoris alterum genus. *Trag. Hist.* 543.

F. latifolia. *Cord. Hist.* 134.

Angl. Royal Moonwort. Osmund Royal.

In palustribus nemorosis, haud vulgaris.

Perennis. Fl. Junio, Julio.

Radix carnosa. Frondes plures, erectæ, tripedales, glabrae, lætè virentes, formosæ, fraxinæ, bipinnatæ, pinnis lanceolatis, subferrulatis, basi cordatis. Spicæ terminales, modo pinnarum dispositæ, at decompositæ. Capsulæ conglomeratæ, minores quam in priori, glabrae.

427. LYCOPODIUM. G. *Pl.* 561. *Juss.* 12.

Capsule axillares, solitariæ, bivalves, nudæ, subreniformes, compressæ.

clava-
tum. 1. LYCOPODIUM foliis sparsis filamentosis ; floralibus dilatato-membranaceis, ramis floriferis infernè strigosis : apice bifidis trifidisve.

Lycopodium clavatum. *Sp. Pl.* 1564. *Huds.* 462.
With. 756. *Hull.* 235. *Kelb.* 405. *Sibth.* 265.
Abbot.

Abbot. 228. *Engl. Bot.* t. 224. *Dicks. H.*
Sicc. fasc. 7. 18. *Fl. Dan.* t. 126. *Roth.*
Germ. v. 3. 12.

Lycopodium. *Raii Syn.* 107.

L. vulgare pilosum, amfragosum et repens. *Dill.*
Musc. 441. t. 58. f. 1.

Muscus terrestris. *Matth.* *Valgr.* v. 1. 57.
Camer. Epit. 32. *Trag. Hist.* 555.

M. clavatus, sive Lycopodium. *Ger. em.* 1562.

Chamæpeuce. *Cord. Hist.* 111. 2.

Angl. Common Club-moss.

In ericetis montosis.

Perennis. *Fl. Julio, Augusto.*

Caules flagelliformes, prostrati, ramosi, radicantes, un-
dique foliosi. Folia densa, patentia, linear-lanceolata,
subferrata, plana, apice pilifera, quandoque secunda.
Rami floriferi erecti, palmares, laxè foliosi, apice in
spicas duas vel tres, unciales, cylindraceas, foliosas,
divisi. Folia floralia basi dilatata, membranacea,
flavescentia. Capulæ axillares, sessiles, solitariæ,
flavescentes, membranaceæ. Semina minutissima, ful-
phurea, copiosa.

2. LYCOPODIUM foliis sparsis ciliatis lanceola- *Selagi-*
tis; floralibus majoribus patentibus. *noides.*

Lycopodium Selaginoides. *Sp. Pl.* 1565. *Huds.*
 463. *With.* 757. *Hull.* 236. *Lightf.* 686.
Dicks. H. *Sicc. fasc.* 7. 19. *Dr. Pl.* 90. *Fl.*
Dan. t. 70. *Engl. Bot.* t. 1148. *Roth. Germ.*
 v. 3. 13.

L. n. 1717. *Hall. Hist.* v. 3. t. 46. f. 1.

Selaginoides foliis spinosis. *Raii Syn.* 106. *Dill.*
Musc. 460. t. 68.

Angl. Prickly Club-moss.

VOL. III.

O

In

In ericetis alpinis irriguis.

In Westmoreland, Wales and Scotland.

Perennis. Fl. Augusto.

Caules basi graciles, ramosi, decumbentes; superne erecti, biunciales; undique foliosi, apicem versus fructiferi. Folia lætè virentia, splendentia, lanceolata, dentato-ciliata, enervia, concaviuscula, erectiuscula; floralia majora, atque horizontaliter patentia. Capsulæ axillares; superiores formâ præcedentis; inferiores tetracoccæ, corpuscula quatuor, globoſa, nivea, contingen-tes, cuius natura nobis nondum patuit.

inunda-tum. 3. **LYCOPODIUM** foliis sparsis integrerrimis linear-i-lanceolatis muticis; floralibus basi dilatatis patentiusculis.

Lycopodium inundatum. Sp. Pl. 1565. Huds. 463. Wth. 758. Hull. 236. Relb. 405. Abbot. 228. Lightf. 687. Engl. Bot. t. 239. Dicks. H. Sicc. fasc. 4. 18. Roth. Germ. v. 3. 14.

L. palustre repens, clavâ singulari. Dill. Musc. 452. t. 62. f. 7.

Muscus terrestris repens, clavis singularibus foliosis erectis. Raii Syn. 108.

Angl. Marsh Club-moss.

In ericetis inundatis, præcipue arenosis, rariūs.

On Hampstead and Bagshot heaths, and in the north. Ray. On St. Faith's heath, Norwich.

Perennis. Fl. Junio, Julio.

Caules omnino prostrati, repentes, breves, foliosi, ramis floriferis erectis, biuncialibus, simplicissimis. Folia densa, linear-i-lanceolata, integrerrima, mutica; floralia parùm majora, subpatentia, b̄si valde dilatata et s̄pē utrinque unidentata. Capsulæ reniformes, parvæ.

4. LYCOPODIUM foliis sparsis integerrimis lan- *Selago.*
ceolatis muticis uniformibus octofariis, caule
dichotomo erecto fastigiato.

Lycopodium Selago. *Sp. Pl.* 1565. *Huds.* 463.
With. 758. *Hull.* 236. *Sibth.* 265. *Engl.*
Bot. t. 233. *Fl. Dan.* t. 104. *Roth. Germ.*
v. 3. 14.

Selago foliis et facie Abietis. *Raii Syn.* 106.
S. vulgaris, Abietis rubræ facie. *Dill. Musc.* 435.
t. 56. f. 1.

Muscus terrestris Abietiformis. *Moris. v. 3.* 624.
sect. 15. t. 5. f. 9.
Angl. Fir Club-moss.

In ericetis montosis humidiusculis, præcipue al-
pinis.

On the high mountains of Wales, Yorkshire,
Derbyshire, &c. *Ray.* In Sussex. *Dill.* On
Felthorpe bogs, Norfolk. *Mr. J. Fox.*

Perennis. *Fl. Junio, Augusto.*

Radix fibrosa. Caules subspithamæi, erecti, dichotomi,
fastigiati, undique foliosi. Folia densa, uniformia,
octofaria, lanceolata, integerrima, acuta, mutica, sa-
turate viridia, nitida. Capsulæ axillares, sparsæ, parvæ,
reniformes, fusco-luteæ.

5. LYCOPODIUM foliis sparsis subserratis mu- *annoti-*
ticis quinquefariis; floralibus dilatatis abbre- *num.*
viatis, ramis floriferis annotino-articulatis.

Lycopodium annofinum. *Sp. Pl.* 1566. *Huds.*
464. *With.* 759. *Hull.* 236. *Lightf.* 689.
Fl. Dan. t. 127. *Roth. Germ.* v. 3. 16.

L. elatius juniperinum, clavis singularibus, sine
pediculis. *Dill. Musc.* 455. t. 63. f. 9. *Raii*
Syn. 107. *Moris. v. 3.* 624. sect. 15. t. 5. f. 3.

Angl. Interrupted Club-moss.

O 2

In

In Alpibus.

In Carnarvonshire. *Mr. Lhwyd.* On the sides of the Highland mountains, but not common. *Lightf.*

Perennis. Fl. Junio—Augusto.

Caules repentes, foliosi, rami floriferis erectis, dichotomis, spithamæis, foliosis, annotino-interruptis. Folia patentia, lanceolata, ferrulata, mutica, lætè viridia; floralia arctè imbricata, minora, dilatata, submembranacea, flavescentia. Capsulæ reniformes.

alpinum. 6. LYCOPodium foliis quadrifariàm imbricatis acutis carinatis; floralibus ovato-lanceolatis planis, ramulis fastigiatis erectis.

Lycopodium alpinum. Sp. Pl. 1567. Fl. Lapp. ed. 2. 338. t. 11. f. 6. Huds. 464. With. 759. Hull. 235. Lightf. 690. Engl. Bot. t. 234. Fl. Dan. t. 79. Dicks. Dr. Pl. 46. Roth. Germ. v. 3. 17.

L. Sabinæ facie. Raii Syn. 108. Dill. Musc. 445. t. 58. f. 2.

Angl. Savin-leaved Club-moss. *

In alpinis petrosis.

On the high mountains of Wales and the north of Yorkshire. *Ray.* Near the summits of almost all the high Scottish mountains. *Lightf.* On the Pentland hills, Edinburgh.

Perennis. Fl. Augusto.

Caules prostrati, repentes, flagelliformes, teretes, laxè foliosi, longissimi; rami floriferi alterni, erecti, ramosissimi, fastigiati, densè foliosi. Folia imbricata, quadrifaria, parva, acuta, carinata, glauca, integerrima; floralia membranacea, plana, basin versùs dilatata, flavescentia, acuminata, subserrata. Capsulæ reniformes, flavæ.

428. POLYPODIUM. G. Pl. 560. Juss. 15.
Act. Taurin. v. 5. 408. Swartz. in Schrad.
Journ. ann. 1800. v. 2. 18.

Fructificationes in punctis subrotundis, sparsis,
 non marginalibus. *Involucrum* nullum.

* *Fronde pinnatifidâ.*

1. POLYPODIUM fronde pinnatifidâ : lobis ob-vulgare.
 longis subserratis obtusis, radice squamatâ.

Polypodium vulgare. Sp. Pl. 1544. Hudf. 455.
With. 773. *Hull.* 236. *Relb.* 409. *Sibth.* 269.
Abbot. 226. *Curt. Lond. fasc.* 1. t. 68.
Woodv. Suppl. t. 271. *Bolt. Fil.* 32. t. 18.
Fl. Dan. t. 1060. *Sw. in Schr. Journ. ann.*
 1800. v. 2. 24. *Engl. Bot.* t. 1149. *Ehrb.*
Crypt. 121. *Roth. Germ.* v. 3. 41.

Polypodium. *Raii Syn.* 117. *Ger. em.* 1132.
f. 1, 2. *Matth. Valgr.* v. 2. 628, 629. *Fuchs.*
Hist. 588. *Cord. Hist.* 171. 2. *Tillands.*
Ic. 79.

β. P. murale, pinnulis serratis. *Dill. in Raii*
Syn. 117.

P. majus, ferrato folio. *Barrel. Ic.* t. 38.

γ: P. vulgare, lobis proliferis. *Bolt. Fil.* 33. t. 2.
f. 5. b.

δ. P. cambricum. Sp. Pl. 1546. *Bolt. Fil.* t. 2.
f. 5. a.

P. cambrobritannicum, pinnulis ad margines
 laciniatis. *Raii Syn.* 117. *Moris. sect.* 14.
t. 2. f. 8.

Filix amplissima, lobis foliorum laciniatis, cam-
 brica. *Pluk. Phyt.* t. 30. f. 1.

Angl. Common Polypody.

- In muris, tectis, et arborum truncis.
 β. On the walls of Windsor Castle. *Rev. Mr. Manningham.* In Worcestershire. *Dr. Stokes.*
 γ. Wood near Bingley. *Dr. Alexander.*
 δ. On a rock in a wood near Dennys Powys Castle, not far from Caerdiff, Glamorganshire. *Ray.* Rocks in North Wales. *Mr. Griffith.* Near Kidderminster. *With.*

Perennis. Fl: Maio—Octobri.

Radix repens, tortuosa, squamata, fibris villosis, aurato-fuscis, ut in omnibus ferè hujusce ordinis naturalis. Frondes spithamææ aut pedales, erectæ, lanceolatæ, acutæ, pinnatifidæ, glabræ; pinnis parùm connatis, linearis-oblongis, obtusiusculis, integerrimis, crenatis, serratisve; in δ variè pinnatisido-laciniatis, majoribus, atque sterilibus. Maculæ fructiferæ simplici serie utrinque, costam versùs, digestæ, subrotundæ, convexæ, fulvæ, omnino nudæ. Capsulæ paryæ, albidæ, annulo fulvo.

** *Fronde subbipinnata.*

*sonta-
num.*

2. POLYPODIUM? fronde linearis-lanceolata acutâ pinnatâ: pinnis cordatis, obtusiusculis pinnatifidis crenatis suprà glabris.

Polypodium fontanum. Sp. Pl. 1550. (exclusâ descriptione.) *Huds.* 456. *With.* 777. *Hull.* 238. *Bolt. Fil.* 38. t. 21.

Filix saxatilis non ramosa, minima. *Pluk. Phyt.* t. 89. f. 2.

Angl. Smooth Rock Polypody.

In muris antiquis et rupibus. *Huds.*

In stony places above Wybourn, Westmoreland, and on Agmondesham church, Buckinghamshire. *Huds.*

Perennis. Fl. Junio.

Radix cæspitosa, fibris longissimis, nigris, glabris. Frondes 2—3-unciales, rectæ, linear-lanceolatæ, acutæ, stipitibus basi squamis magnis, scariosis, interstinctis. Pinnæ numeroſæ, densæ, ſepiùs alternæ, breves, cordatæ, obtusiusculæ, pinnatifidæ, basi dilatatæ, margine plus minus lobatæ vel crenatæ, muticæ; ſuprà glaberrimæ, lætè virentes; ſubtùs maculis fructificantibus rotundis, demùm coacervatis, undique ferè tectæ. Capsulæ pilis interſertæ.

Ab Aſpido fontano Swartzii distinctum.

3. POLYPODIUM fronde lanceolatâ obtusius-*arvoniculâ* pinnatâ: pinnis cordatis pinnatifidis *cum utrinque pilosis*: lobis rotundatis repandis!

Polypodium arvonicum. *With.* 774. *Hull.* 238.

Sym. Syn. 191.

P. ilvense. *With.* 774. *Hull.* 237.

P. hyperboreum. *Sw. in Schr. Journ. ann.* 1800.

v. 2. 27.

Acrostichum ilvense. *Huds.* 451. *Dicks. H. Sicc. fasc.* 8. 17. nequaquam Linnæi.

A. alpinum. *Bolt. Fil.* 76. *t. 42.* opt.

Filix alpina, *Pedicularis rubræ* foliis ſubtùs villosis. *Raii Syn.* 118.

Filicula pumila, *Lonchitidis Maranthæ species,* *Cambrobritannica.* *Pluk. Phyt.* *t. 89. f. 5.*

Angl. Hairy Alpine Polypody.

In alpibus Cambrobritanicis et Scoticis rariùs.

On Snowdon, upon a moist black rock almost at the top of Clogwyn y Garnedh, facing the North West, directly above the lower lake. *Lbwyd, Richardson, Huds. and Rev. Hugh. Davies.* On Ben Lawers. *Dicks.*

Perennis. Fl. Junio.

Radix cæſpitosa. Frondes patulæ, 2—4 unciales, lanceolatæ,

ceolatæ, obtusiusculæ, utrinque, rachique, pilosæ. Pinnæ minùs approximatæ, sæpiùs suboppositæ, cordatæ, pinnatifidæ, basi dilatatæ, lobis rotundatis, convexiusculis, margine repandis, vix crenatis, inuticis. Maculæ fructificantes subrotundæ, pilosæ, numerosæ, demùm approximatæ, fuscæ.

- Phego-* 4. **POLYPÓDIUM** fronde pinnatâ : pinnis lanceolatis acuminatis pinnatifidis basi coadunatis ; infimis reflexis.

Polypodium Phegopteris. *Sp. Pl.* 1550. *Huds.* 456. *With.* 775. *Hull.* 238. *Lightf.* 669. *Balt. Fil.* 36. *t.* 20. *Sw. in Schr. Journ. ann.* 1800. *v.* 2. 28. *Erbh. Crypt.* 131.

Filix minor Britannica pediculo pallidiore, alis inferioribus deorsum spectantibus. *Dill. in Raii Syn.* 122.

Angl. Pale Mountain Polypody.

In rimis petrarum, et montosis humidis, Angliæ borealis et Scotiæ australis.

In Westmoreland, but not very common.

Perennis. Fl. Junio, Julio.

Radix repens. Frondes erctæ, pedales, pallidè virentes, stipite longissimo, glabro, pallido. Pinnæ oppositæ, connatæ, lanceolatæ, acuminatæ, regulariter pinnatifidæ, molles, pilosiusculæ, margine subcrenatae ; infimæ reflexæ. Maculæ fructiferæ laciniarum marginem versùs, distinctæ, sulphureæ. Rachis communis ut et partiales hirsutæ.

*** *Fronde supradecompositâ.*

- Dryo-*
pteris.

5. **POLYPODIUM** fronde ternatâ bipinnatâ patulo-deflexâ ; laciñiis obtusis subcrenatis, radice filiformi,

Polypodium Dryopteris. *Sp. Pl.* 1555. *Huds.* 460.

460. *With.* 780. *Hull.* 240. *Bolt. Fil.* 52.
t. 28. *Engl. Bot.* *t.* 616. *Hoffm. Germ.*
v. 2. 10. *Sw. in Schr. Journ. ann.* 1800.
v. 2. 29. *Ehrb. Crypt.* 102.

Filix ramosa minor. *Raii Syn.* 125. *Baub. Hist.*

v. 3. 733.

F. arborea. *Trag: Hist.* 538.

Filicula petræa fœmina quarta. *Tabern. Kreuterb.* 1182.

Angl. Tender Three-branched Polypody.

In umbrosis ad latera montium.

Near Tintern abbey, Monmouthshire. *Ray.* In
 the north of Yorkshire. *Richardson.* *Mr. T. Forster* and *Mr. W. Brunton.* Staffordshire. *Rev. T. Gisborne.*

Perennis. Fl. Julio.

Radix repens, filiformis, tenuis. Stipes gracillimus,
 erectus, subpedalis, lœvis. Frondes ternatæ, patulæ,
 plus minus deflexæ, tenerrimæ, lætè virentes, pal-
 mares, subbipinnatæ; impari paululum majore, et
 quandoque spithamææ. Pinnæ sessiles; inferiores ma-
 jores, magisque compositæ. Pinnulæ oppositæ vel
 alternæ, oblongæ, obtusiusculæ, crenato-dentatæ, mu-
 ticæ, sæpius subrevolutæ. Rachis ferè capillaris, ca-
 naliculata, pubescentula. Maculæ fructiferæ submar-
 ginales, numerosæ, parvæ, pilis omnino destitutæ,
 nunquam confluentes.

6. POLYPODIUM fronde ternatâ bipinnatâ erec- *calcare-*
 tâ, rigidulâ; laciniis obtusis subcrenatis, ma- *um.*
 culis fructiferis confluentibus.

Polyodium Dryopteris. *Dicks. Dr. Pl.* 16.
Bolt. Fil. 53. *t.* 1.

Dryopteris Tragi. *Ger. em.* 1135.

Filix pumila saxatilis prima. *Clus. Hist.* *v.* 2. 212.

Angl. Rigid Three-branched Polypody.

In

In ericetis montosis, solo calcareo.
About Matlock bath, Derbyshire.

Perennis. Fl. Julio.

Radix repens, sed brevior et crassior quam in præcedente; differt etiam fronde magis rigidâ, stipite basi squamofo, foliis duplò minoribus, firmis, erectiusculis, nec laxè deflexis, capsularum fasciculis demum confluentibus, totam frondis paginam occupantibus.

429. ASPIDIUM. Swartz. in Schrad. Journ. ann. 1800. v. 2. 29.

Polypodium. Act. Taurin. v. 5. 408.

Fructificationes in punetis subrotundis, sparsis, non marginalibus. *Involucrum umbilicatum*, undique ferè dehiscens.

* *Fronde pinnatâ.*

Lonchitis. 1. ASPIDIUM fronde pinnatâ: pinnis lunulatis ciliato-serratis declinatis, stipitibus squamatis.

Aspidium Lonchitis. Sw. in Schr. Journ. ann. 1800. v. 2. 30.

Polypodium Lonchitis. Sp. Pl. 1548. Huds. 455. Willb. 773. Hull. 237. Lightf. 668. Fl. Dan. t. 497. Bolt. Fil. 34. t. 19.

Polystichum Lonchitis. Roth. Germ. v. 3. 71.

Lonchitis aspera major. Raii Syn. 118. Ger. em. 1140. Matth. Valgr. v. 2. 273. Camer. Epit. 664.

Angl. Rough Alpine Shield-fern.

In fissuris rupium alpinarum.

In Carnarvonshire. Mr. Lhwyd: Breadalbane. Rev. Mr. Stuart. Near Stirling. Dr. Buckannan.

Perennis. Fl. Maio, Junio.

Frondes spithameæ, erectæ, lanceolatae, angustæ. Pinnæ numerosæ,

numerofæ, subfessiles, alternæ, approximatæ, declinatæ, acutæ, margine spinulosæ; basi supernè productæ, infernè excisæ; subtus ramentacco-subvillofæ: superiores tantum fructiferæ. Maculæ capsularum sœpè approximato-confluentes, involucris membranaceis, umbilicatis, circinatis.

Ab A. aculeato certè distinctum.

** *Fronde subbipinnatâ.*

2. ASPIDIUM fronde pinnatâ : pinnis pinnati-*Thely-*
fidis subcrenatis basi distinctis decussantibus, pteris:
punctis fructiferis confluentibus.

Aspidium Thelypteris. Sw. in Schr. Journ. ann.

1800. v. 2. 40.

Polypodium Thelypteris. Mant. 505. Syst.

Veg. ed. 14. 937. With. 776. Hull. 239.

Relb. 410. Abbot. 227. Fl. Dan. t. 760.

Dicks. H. Sicc. fasc. 6. 15. Engl. Bot. t. 1018.

Polystichum Thelypteris. Roth. Germ. v. 3. 77.

Acrostichum Thelypteris. Sp. Pl. 1528. Bolt.

Fil. 78. t. 43, 44.

Thelipteris non ramosa. Schmidel Ic. t. 11.

Filix minor palustris repens. Raii Syn. 122.

F. tenuissimè et profundè denticulata Montbel-
gardica. Baub. Hist. v. 3. 731, benè.

F. minor non ramosa. Ibid. 732, ex descr.

Angl. Marsh Shield-fern.

In paludosis rariùs.

On St. Faith's Newton bogs, Norwich. Mr. Pitch-
ford. Herb. D. Rose. At Lound near Yarmouth.

Mr. D. Turner. In Potton marshes, Bedford-
shire. Abbot. Cambridgeshire. Relhan. Scot-
land. Dickson. Terrington car, Yorkshire.

Mr. Teesdale.

Perennis. Fl. Julio.

Radix repens. Frondes erectæ, pedales, lanceolatæ, lœtè
virentes,

virentes, stipite glabro, subelongato. Pinnæ oppositæ, patentes, sessiles, nec connatæ, lanceolatæ, acuminatæ, pinnatifidæ, sæpius glabræ, quandoque pilis raris hirsutæ; lobis apice subcrenatis, infimis per paria decussantibus, quod icon Bauhiniana optimè repræsentat. Maculæ fructiferæ numerosæ, demum confluentes, et discum ferè totum occupantes. Rachis communis glaberrima; partiales rariùs pilosiusculæ.

Oreopteris. 3. **ASPIDIUM** fronde pinnatâ: pinnis pinnatifidis integerrinis subtûs resinoso-glandulosis, punctis fructiferis marginalibus confluentibus. *Aspidium Oreopteris.* Sw. in Scbr. Journ. ann. 1800. v. 2. 35.

Polypodium *Oreopteris.* Ehrb. Crypt. n. 22. Dicks. Tr. of L. Soc. v. 1. 181. H. Sicc. fasc. 1. 18. Wub. 775. Hull. 238. Sibth. 270. Teesdale Tr. of L. Soc. v. 2. 113. Fl. Dan. t. 1121. Hoffm. Germ. v. 2. 5. Engl. Bot. t. 1019.

P. *Thelypteris.* Huds. 457. Bolt. Fil. 40. t. 22. f. 1, 2. Hedw. Theor. 44. t. 6.

P. *fragrans.* Huds. 457? P. *limbospermum.* Bellardi in Act. Taurin. v. 5. 253. ex ipso auctore.

Polystichum *montanum.* Roth. Germ. v. 3. 74. *Filix* non *ramosa nostras*, pinnulis brevibus, acutioribus, integris, nonnihil falcatis, punctis ferrugineis ad oras pulverulentis. Pluk. Amalb. 91.

F. *maris vulgaris* varietas. Dood. in Raii Syn. ed. 2. 341. ed. 3. 122.

3. F. *pumila saxatilis* altera Clusii. Raii Syn. 122. Angl. Heath Shield-fern.

In montosis, ericetis, et sylvis siccioribus.

Most plentiful in Scotland. Dicks. North of England.

land. *With.* Near Southampton. *Mr. Lambert.* At Hafod, Cardiganshire.

Perennis. Fl. Julio.

Præcedente triplò plerumque majus. Radix magna, fibrosa, squamosa. Frondes lanceolatæ, fere ad basin usque pinnatæ, rachi lævi, fulcatâ. Pinnæ suboppositæ, sessiles, haud decusatae, lanceolatæ, acuminatæ, pinnatifidæ, glabriuscultæ, subtus punctulis resinosis, aureis, undique conspersæ, lobis plerumque integerimis, rariùsve apice subcrenatis. Maculæ fructiferæ numerosæ, simplici serie propè marginem digestæ, approximatæ, demùm confluentes, et lineari monili-formem, fuscam, constituentes. Involucrum tenuissimum, umbilicatum, lacerum, citò evanidum.

β. magnitudine minori tantùm differre videtur.

Synonymon *Hedwigii* hûc amandavi, ob puncta resinosa quibus hæc species a præcedente, etiamsi non fructifera, dignoscitur. Ex hisce forsitan odorem suavem, a Teesdalio memoratum, spargit.

4. **ASPIDIUM** fronde bipinnatâ : pinnulis ob-*Filix* tusis ferratis, stipite paleaceo, involucris cir-mas-cinatis.

Aspidium Filix mas. *Sw.* in *Schr. Journ. ann.* 1800. v. 2. 38.

Polypodium Filix mas. *Sp. Pl.* 1551. *Huds.* 458. *With.* 775. *Hull.* 238. *Rehb.* 411. *Sibth.* 270. *Abbot.* 226. *Bolt. Fil.* 44. t. 24. *Woodv.*

Med. Bot. t. 49. *Dicks. H. Sicc.* fasc. 3. 19.

Polystichum Filix mas. *Roth. Germ.* v. 3. 82.

Filix mas vulgaris. *Raii Syn.* 120. *Ger. em.* 1128. *Fuchs. Hist.* 595. *Matth. Vulgr.* v. 2. 626. *Camer. Epit.* 991.

Filix, radix. *Pharmac. Lond.*

Angl. Male Fern. *Male Shield fern.*

In sylvis, aggeribusve umbrosis, vulgaris.

Perennis.

Perennis. Fl. Junio, Julio.

Radix cæspitosa, magna, squamosa. Frondes lanceolatae, tripedales, stipite rachique paleaceis. Pinnæ lanceolatae, elongatae, numerosæ, pinnulatæ, pinnulis subalternis, numerosis, oblongis, obtusis, undique crenato-ferratis, utrinque glabris, superioribus confluentibus. Maculæ fructiferæ propè nervos, imprimis basin versùs pinnularum, simplici serie utrinque digestæ, magnæ, prominentes, copiosæ. Involucra umbilicata, undique circinata, hinc fissa, margine plicata.

aculeatum. 5. **ASPIDIUM** fronde bipinnatâ: pinnulis ovato-lunulatis ciliato-spinosis subtus pilosis, rachibus paleaceis.

Aspidium aculeatum. Sw. in Schr. Journ. ann. 1800. v. 2. 37.

Polypodium aculeatum. Sp. Pl. 1552. Huds. 459. Wth. 777. Hull. 239. Relb. 412. Sibth. 271. Abbot. 227. Bolt. Fil. 48. t. 26. Mill. Illustr. t. 101.

Polystichum aculeatum. Roth. Germ. v. 3. 79. *Filix* mas non rainosa, pinnulis latis auriculatis spinosis. Raii Syn. 121. Moris. sect. 14. t. 3: f. 15.

β. *F. mas aculeata nostras, alis expansis, muscosâ lanugine aspersâ.* Pluk. Phyt. t. 180. f. 1. Dill. in Raii Syn. 121.

γ. *F. lonchitidi affinis.* Raii Syn. 121.
F. aculeata, lonchitidis ænula. Pluk. Phyt. t. 180. f. 3.
Angl. Common Prickly Shield-fern.

In sylvis et umbrosis sœpium aggeribus, præsertim humidis et petrosis.

γ. On the Welch mountains. Mr. Lbwyd.

Perennis. Fl. Junio.

Radix

Radix cæspitosa. Frondes bipedales, elegantè bipinnatæ, saturatè virentes, rachi generali ut et partialibus maximè palmaceis. Pinnæ alternæ, parùm distantes, lanceolatæ, pinnulatæ, pinnulis alternis, subdistantibus, ovato-lunulatis, dentato-ferratis, spinulosis, anticè ad basin plus minus lobatis vel auriculatis; inferioribus interdùm pinnatifidis; omnibus subtùs pallidioribus et pilosis. Maculæ fruetiferae per discum sparsæ. Involucra omnino orbiculata, umbilicata, nec fissa.

Figuræ apud Plukenetium, t. 179. f. 6. et Bolt. Fil. t. 26. f. 2. plantam juniores exhibent.

Stirps solitæ magnitudinis ac formæ, e loco sylvoso in hortum apricum translata, secundo anno in varietatem mutatur, ut clare demonstrant exemplaria a D. Gulielmo Brunton fil. communicata.

6. ASPIDIUM fronde bipinnatâ: pinnulis ellip-lobatum, tico-lunulatis ciliato-spinosis subtùs pilosis; infimis anticis maximis, rachibus paleaceis.

Aspidium lobatum. Sw. in Schr. Journ. ann. 1800. v. 2. 37.

Polypodium lobatum. Huds. 459.

Filix aculeata major, pinnulis auriculatis crebrioribus, foliis integris angustioribus. Raii Syn. 121, descriptione optimâ. D. E. Forster.

F. aculeata, Lonchitidis æmula nostras. Pluk. Phyt. t. 180. f. 3.

Angl. Close-leaved Prickly Shield-fern.

In aggeribus sœpium umbrosis.

Frequent in Essex. Ray. Mr. E. Forster junior.

Perennis. Fl. Junio.

Similis præcedenti, at rigidior, pallidior, magisque nitida. Forma totius frondis angustior est, pinnis valde approximatis, pinnulis magis ellipticis, minusque incisis, quarum anticæ infimæ (seu rachi proximæ) reliquis duplo majores sunt, hinc angulato-lobatæ, ultrà pinnæ proxime superioris basin plerumque extensæ. Confer Raii Synopsin.

Filix 7. **ASPIDIUM** fronde bipinnatâ: pinnulis pinnatifidis ferratis inuticis, rachi lævi, involucris reniformibus.

Aspidium Filix fœmina. *Sw.* in *Schr. Journ. ann. 1800. v. 2. 41.*

Polypodium Filix fœmina. *Sp. Pl. 1551.*
Huds. 458. *With. 778.* *Hull. 239.* *Relb. 411.* *Sibth. 271.* *Abbot. 226.* *Bolt. Fil. 46. t. 25.* *Dicks. H. Sicc. fasc. 5. 18.*

Filix mas non ramosa, pinnulis angustis rarís, profundè dentatis. *Raii Syn. 121.* *Pluk. Phyt. t. 180. f. 4.* *Moris. sect. 14. t. 3. f. 8.*
Angl. Female Shield-fern.

In palustribus umbrosis, frequens.

Perennis. Fl. Junio, Julio.

Radix magna, cæspitosa. Frondes sesquipedales, latè lanceolatæ, saturatè virides, glabrae, rachi lævissimâ. Pinnæ alternæ, lanceolatæ, acuminatæ, pinnulatæ, pinnulis elegantè ac tenuè pinnatifidis, incisis, inuticis; superioribus præcipue decurrentibus. Maculæ fructiferæ eodem modo ac in A. Filice mari dispositæ, sed multò minores, et oblongæ. Involucra reniformia, marginè lacera.

spinulo- 8. **ASPIDIUM** fronde subbipinnatâ: pinnulis decurrentibus ellipticis pinnatifido-ferratis spinulosis, rachi lævi, nervis flexuosis.

Polypodium spinulosum. *Retz. Prod. 250. Fl. Dan. t. 707.* *With. 778.* *Hull. 239.* *Sym. 191.*

P. n. 841. *Mull. Fl. Fridr. 193. t. 2. f. 2.*

Filix pumila saxatilis altera. *Pluk. Phyt. t. 179. f. 5:* benè.

Angl. Lesser crested Shield-fern.

In paludosis.

Bogs on Birmingham heath, and in Devonshire.
With. In Mr. Rose's herbarium.

Perenn is.

Perennis. Fl. Junio, Julio.

Radix repens. Frondes pedales, longius stipitatæ, subovatae, læte virentes, glabrate, tenues et subpellucidæ; stipite, ut et rachi, glaberrimo. Pinnæ subalternæ, late lanceolatae, subpinnulatae, vel profundè pinnatifidæ, inferiores majores. Pinnulæ oppositæ, decurrentes, uniformes, ellipticæ, obtusiusculæ, pinnatifido-serratae seu lobatae, lobis incurvis, spinulosis, nervo maximè flexuoso. Maculæ fructiferæ distantes, parvæ, lobi axillas versus collocatæ, rotundatæ. Involucra circinata, hinc fissa.

Distincta species, a Cl. Witheringio benè elucidata.

Diversa videtur a *Polysticho spinoso* Roth. *Fl. Germ.* v. 3. 91, e Germaniâ ad amicis. D. Turnerum misso.

9. *ASPIDIUM* fronde bipinnatâ: pinnulis pinna-dilata-tifidis inciso-serratis spinulosis, rachi squamosâ: tum. involucro circinato.

Polypodium dilatatum. Hoffm. *Germ.* v. 2. 7.

P. aristatum. Villars. *Dauph.* v. 3. 844. Bel-lardi in *Act. Taurin.* v. 5. 255.

P. cristatum. Huds. 457. With. 778. Hull. 239. Relb. 410. Sibth. 270. Abbot. 226. Bolt. *Fil.* 42. t. 23.

P. n. 1705. Hall. *Hist.* v. 3. 14.

P. n. 845. Mull. *Fl. Frider.* 193. t. 2. f. 4.

Polystichum multiflorum. Roth. *Germ.* v. 3. 87.

Filix mas ramosa, pinnulis dentatis. Raii *Syn.* 124. *Pluk. Phyt.* t. 181. f. 2.

F. mas, pinnulis cristatis. Moris. *sect.* 14. t. 3. f. 11.

F. ramosa dentata, ramulis et pinnis longius ab invicem distantibus. Mapp. *Alsat.* 106. f. 8.

F. tenuissimè secta ex monte Ballon. Baub. *Hist.* v. 3. 731.

Angl. Great Crested Shield-fern.

In umbrosis irriguis, præsertim solo sabuloſo, vel
in sylvis petroſis humidioribus.

Perennis. Fl. Junio, Julio.

Radix tuberosa, vix repens. Frondes ſæpius bipedales,
erecto-patentes, latæ, acutæ, lætè virentes, bipinnatæ,
vel subtripinnatæ, stipite et rachibus squamosis. Pinnæ
suboppositæ, lanceolatæ, acutæ. Pinnulæ subalternæ,
basi pinnatæ, supernæ pinnatifidæ, laciñiis oblongis,
obtusis, apice præcipue ferratæ, spinulosæ, nervo ferè
recto. Maculæ fructiferæ numerosæ, distinctæ, re-
niformi-subrotundæ. Involucra hinc fissa.

Polypodium rhæticum, *Huds.* 458, hujus varietas minor
videtur.

P. cristatum *Linnæi* distinctissima species est, Sibiriæ
incola.

430. **ASPLENIUM.** *G. Pl.* 560. *Juss.* 15.
Act. Taurin. v. 5. 409.

Fructif. in lineolis sparsis. *Involucr.* e venâ
lateraliter ortum ducens, interius (sive costam
versus) dehiscens.

Trichomanes. 1. **ASPLENIUM** fronde pinnatâ: pinnis ovato-
subrotundis crenatis, rachi nitidâ subtûs ca-
rinatâ.

Asplenium Trichomanes. *Sp. Pl.* 1540. *Huds.*
452. *Wittb.* 768. *Hull.* 241. *Rehb.* 408.
Sibth. 268. *Abbot.* 225. *Bolt. Fil.* 22. t. 13.
Woodv: Med. Bot. t. 204. *Fl. Dan.* t. 119.
Dickf. H. Sicc. fasc. 4. 17. *Engl. Bot.* t. 576.
Roth. Germ. v. 3. 55.

Trichomanes. *Raii Syn.* 119. *Fuchs. Hist.* 796.
Matth. Valgr. v. 2. 543. *Camer. Epit.* 925.
Cord. Hist. 170. *Tillands. Ic.* 77.

T. mas. *Ger. em.* 1146.

Adiantum album. *Cord. Hist.* 172. f. 1.

β. Tricho-

3. *Trichomanes* foliis elegantè incisis. *Dill.* in
Raii Syn. 120. *Tourn. Inst.* t. 315. f. C.

Adiantum maritimum, segmentis rotundioribus.
Pluk. Phyt. t. 73. f. 6.

Angl. Common Maidenhair Spleenwort.

In rupibus umbrosis, et muris antiquis, frequens.

Perennis. Fl. toto ferè anno.

Radix cæspitosa, villosa, nigra. Frondes numerosæ, palmares, linear-lanceolatæ; breviùs stipitatæ, pinnatae, rachi atro-purpurea, glabra, nitida, suprà canaliculata, subtùs carinata. Pinnæ oppositæ, subsessiles, ovato-subrotundæ, crenatæ; basi obliquè truncatæ; suprà atro-virides, glabræ, aeniæ; subtùs pallidiores. Lineæ fructiferæ 6 vel 8, obliquæ, alternæ. Involucra linearia, albida, subintegerrima. Capsulæ fuscæ.

Pinnæ in β pinnat ifidæ, quandoque elongatæ, sunt.

Varietatem fronde apice ramosissimâ invenit D. Teesdale in comitatu Hantoniensi.

2. *ASPLENIUM* fronde pinnatâ: pinnis alternis viride. elliptico-subrotundis crenatis, rachi subtùs complanatâ.

Asplenium viride. *Huds.* 453. *With.* 768.
Hull. 241. *Bolt. Fil.* 24. t. 14. *Dickf. H.*
Sicc. fasc. 3. 18. *Lightf.* 663. *Roth. Germ.*
v. 3. 56.

3. A. *Trichomanes ramosum*. *Sp. Pl.* 1541. *Bolt.*
Fil. 25. t. 2. f. 3.

Trichomanes ramosum. *Bauh. Hist.* v. 3. 747.
Raii Syn. ed. 2. 46. *ed. 3.* 119.

T. minus et tenerius. *Moris. sect.* 14. t. 3.
f. 11.

Angl. Green Maidenhair Spleenwort.

In rupibus alpinis.

In Wales, Yorkshire, and Westmoreland. *Ray.*
Huds. Bolt. Scotland. *Lightf. Dicks.*

Perennis. Fl. Junio, Julio.

Magnitudine et formâ præcedenti similis, sed differt rachi viridi, tetraquetrâ, sive subtûs complanatâ; pinnis pallidioribus, alternis, basi magis cuneatis; involucris capsulisque flavescentibus. Rachis in β divisa.

Hanc speciem optimè designavit D. Lhwyd nomine *T. costâ viridi*, quandoque bisidâ, foliis lenticularibus crenatis. Conf. *Raii Syn.*

mari-
num.

3. **ASPLENIUM** fronde pinnatâ: pinnis ovatis obliquis ferratis obtusis basi inæqualibus cuneatis.

Asplenium marinum. *Sp. Pl.* 1540. *Huds.* 453. *With.* 769. *Hull.* 242. *Lightf.* 664. *Bolt. Fil.* 26. t. 15. *Engl. Bot.* t. 392. *Dicks.* *H. Sicc. fasc.* 7. 17.

Adiantum nigrum, foliis Lunariæ minoris. *Dill.* in *Raii Syn.* 120?

Chamæfilix marina anglica. *Raii Syn.* 119. *Moris. sect.* 14. t. 3. f. 25. *Pluk. Phyt.* t. 253. f. 5. *Ger. em.* 1143.

β . *Adiantum trapeziforme.* *Huds.* 460. nequaquam Linn.

A. majus Coriandri folio, et *Adianto vero affinis minor Scotica.* *Raii Syn.* 124. *Sibb. Scot.* 7, 8. t. 3. f. 1, 2.

Angl. Sea Spleenwort.

In rupibus marinis.

In Anglesea, Wales, Sussex and other parts of the southern coast. *Ray.* On many parts of the Scottish coast. *Lightf.* Near Hastings. *Rev. Dr. Goodenough.*

β . In the coves at Weems. *Sibb.* *Lightf.*

Perennis. Fl. Junio—Octobri.

Radix cæspitosa, squamosa, nigra. Frondes magnitudine variæ, lanceolatæ, stipite nigro, lævi. Pinnæ alternæ, ovatæ, obtusæ, saturatè virides, venosæ, crenatæ, serratæ, interdùm laciniatæ; basi cuneatæ, obliquæ, superne angulatæ; superiores plerumque confluentes. Rachis alata, subcomplanata. Lineæ fructiferæ suboppositæ. Involucra fusca, integerima.

In fissuris rupium profundioribus aliquando mirè luxuriat.

D. Goodenough.

β pinnis lobatis seu laciniatis discrepat.

4. *ASPLENIUM* fronde pinnato-trifidâ: pinnis *septentrionalis* linearibus apice laceris. *trionale.*

Asplenium septentrionale. *Hull.* 241. *Engl. Bot.* t. 1017.

Acrostichum septentrionale. *Sp. Pl.* 1524. *Huds.* 450. *Willb.* 764. *Lightf.* 656. *Bolt. Fil.* 12. t. 8. *Fl. Dan.* t. 60. *Dicks. Dr. Pl.* 45. *Ehrb. Crypt.* 101.

Scolopendrium septentrionale. *Roth. Germ.* v. 3. 49.

Felix saxatilis Tragi. *Raii Syn.* 120. *Baub. Hist.* v. 3. 747. *Dalech. Hist.* 1226.

F. nuda seu *saxatilis*. *Trag. Hist.* 537. n. 2. f. 1.

Muscus corniculatus. *Ger. em.* 1561. *Baub. Hist.* v. 3. 759.

Angl. Forked Spleenwort.

In fissuris rupium borealium.

In Carnarvonshire. *Ray.* On Ingleborough, Yorkshire; and above Ambleside, Westmoreland. *Huds.* At the Hermitage near Edinburgh, and about Arthur's Seat, where it was found by T. Willisel in the time of Ray.

Perennis. Fl. Junio—Octobri.

Radix cæspitosa, nigra. Frondes plurimæ, palmares, erectæ, confertæ, atro-virentes, glabræ; basi attenuatæ; supernè tripartitæ, sive in pinnas tres, alternas, petiolatas, lincares, apice laceras, suprà lincatas, divisæ. Lincæ fructiferæ approximatæ, parallelæ, fuscæ, demùm totum discum angustum occupantes. Involuta integerrima, superficiaria nec marginalia, interius dehiscentia, ut in omnibus veris Asplenii.

alterni-folium. 5. **ASPLENIUM** fronde pinnatâ: pinnis alternis cuneiformibus erectis apice erosis.

Asplenium alternifolium. *Jacq. Misc. v. 2. 51.*
t. 5. f. 2. Murray Syst. Veg. ed. 14. 933.
Dicks. Tr. of Linn. Soc. v. 2. 290. With. 768,
Hull. 241.

Scolopendrium alternifolium. *Roth. Germ. v. 3.*
53.

Adiantum novum germanicum, Rutæ murariæ facie. *Breyn. Cent. 1. 97.*

Angl. Alternate-leaved Spleenwort.

In rupibus subalpinis.

On rocks in the south of Scotland. *Dicks.*

Perennis. Fl. Junio—Octobri.

Radix præcedentis. Frondes numerosæ, palmares, erectæ, confertæ, lætè virentes, glabræ. Pinnæ alternæ, petiolatae, erectæ, cuneiformes, nervosæ, apice erosæ vel laceræ; inferiores rariū lobatæ, vel rarissimè compositæ. Lineæ fructificantes paucæ, breves, pallidæ.

Ruta-muraria. 6. **ASPLENIUM** fronde alternatîm decompositâ: pinnis rhomboideo-cunciformibus patentibus apice erosis.

Asplenium Ruta-muraria. *Sp. Pl. 1541. Huds.*
453. With. 769. Hull. 242. Relb. 408.
Sibth. 268. Abbot. 225. Bolt. Fil. 28. t. 16.
Engl. Bot. t. 150. Fl. Dan. t. 190.

Scolo-

Scolopendrium Ruta muraria. Roth. Germ.

v. 3. 52.

Ruta muraria. Raii Syn. 122. Ger. em. 1144.

Baub. Hist. v. 3. 737.

Saxifraga. Fuchs. Hist. 730.

Capillus Veneris. Brunf. Herb. v. 1. 219.

Paronychia. Camer. Epit. 785.

Trichomanes. Cord. Hist. 170. f. 2.

β. *Adiantum, an album tenuifolium, Rutæ mura-*
riæ accedens, J. B.? Raii Syn. 123.

A. foliis minutè in oblongum scissis, pediculo
viridi. Pluk. Phyt. t. 3. f. 3.

Angl. Wall-rue Spleenwort. White Maiden-
hair.

In muris antiquis, et rimis petrarum humidis.

Perennis. Fl. Junio—Octobri.

Radix atro-fusca. Frondes plurimæ, palmares, patentes,
confertæ, atro-virentes, subglaucescentes, glabræ, ter-
nato-decompositæ, pinnis subalternis, petiolatis, cu-
neiformibus, vel subindè rhombeis, horizontaliter
patentibus vel deflexis, apice erosionis atque crenatis.
Lineæ fructificantes plures, demùm confluentes. In-
volucra albida, margine crenulata.

7. *ASPLENIUM* fronde subdeltoideâ alternatîm *Adian-*
tripinnatâ : pinnulis lanceolatis acutiusculis *tum ni-*
grum. inciso-serratis.

Asplenium Adiantum-nigrum. Sp. Pl. 1541.

Huds. 454. With. 770. Hull. 242. Relb. 409.

Sibth. 269. Abbot. 225. Bolt. Fil. 30. t. 17.

Fl. Dan. t. 250. Roth. Germ. v. 3. 57.

Adiantum nigrum officinarum. Raii Syn. 126.

n. 10; etiam 127. n. 12, 13.

Onopteris mas. Ger. em. 1137.

Dryopteris nigra. Dod. Pempt. 466. Dalech.
Hist. 1228.

$\beta.$ *Filix minor longisolia*, tarsis raris, pinnulis longis, tenuissimis et oblongis laciniis fimbriatis.

Raii Syn. 127. *Herb. Sherard. Pluk. Phyt.*
t. 282. f. 3.

Angl. Black Maidenhair Spleenwort.

Ad radices et truncos arborum, inque muris antiquis vel aggeribus umbrosis, frequens.

$\beta.$ In a dark cave on the mountains of Mourne, Ireland. *Sherard.*

Perennis. Fl. Junio—Octobri,

Radix nigra. Frondes erectiusculæ, pedales, deltoideo-oblongæ, acutæ, atro-virides, glabræ, alternatim subtripinnatæ, pinnulis elliptico-laneeolatis, aequitiusculis; basi cuneatis, integerrimis; superne ineiso-ferratis. Stipes elongatus, atro-purpureus, nitidus. Lineæ fructificantes plures, demum confluentes. Involucra tenuia integerrima.

Pinnulæ in β steriles, membranaeæ, elegantè palmato-laciniatæ.

lanceola- 8. **ASPLENIUM** fronde lanceolatâ bipinnatâ :
tum. pinnulis obovatis dentato-incisis: dentibus acuminatis.

Asplenium lancoolatum. *Huds.* 454. *With.* 770.

Hull. 242. *Engl. Bot.* t. 240.

Filix elegans, *Adianto nigro accedens*, segmentis rotundioribus. *Raii Syn.* 127.

Dryopteris candida. *Dodon. Pempt.* 465. *Ger.*
em. 1135. quoad iconem.

Angl. Lanceolate Spleenwort.

In muris antiquis et rupibus Angliæ australioris.
Plentiful about St. Ives, Cornwall. *Huds.* On the great rocks at Tunbridge wells. *Mr. T. F. Forster.*

Perennis. Fl. Junio—Octobri.

Radix nigra, squamosa. Frondes erectæ, vix spithameæ,
lanceo-

lanceolatæ, acutæ, lætè virentes, glabræ, bipinnatæ, pinnis suboppositis. Pinnulæ subalternæ, obovato-subrotundæ, basi cuneatæ, supernè acutè dentatæ, dentibus quandoque subaristatis. Stipes brevis, glaber. Lineæ fructificantes 3 vel 5, breves, demùm confluentes. Involucra albida, margine lacera.

431. SCOLOPENDRIUM. *Act. Taurin.* v. 5.

410. t. 9. f. 2.

Asplenium. G. Pl. 560. Juss. 15.

Fructif. in lineolis sparsis, geminis, interveniis.
Involucr. superficiaria, sibi invicem longitudinaliter incumbentia, futurâ longitudinali dehiscentia.

1. SCOLOPENDRIUM fronde simplici cordato-*vulgare*.
lingulatâ subtûs lævi.

Scolopendrium vulgare. Sym. Syn. 193. Engl.
Bot. t. 1150.

Scolopendrium. Brunf. Herb. v. 2. 40.

S. Phyllitis. Roth. Germ. v. 3. 47.

Scolopendria vulgaris. Trag. Hist. 549.

Asplenium Scolopendrium. Sp. Pl. 1537. Huds.

452. *With. 766. Hull. 241. Relb. 408.*

Sibth. 268. Abbot. 225. Bolt. Fil. 18. t. 11.

Curt. Lond. fasc. 1. t. 67. Woodv. Suppl.

t. 272. Ehrb. Crypt. 111.

Phyllitis. Raii Syn. 116. Ger. em. 1138. Matth.

Valgr. v. 2. 186. Camer. Epit. 579. Dalech.

Hist. 1219.

Hemionitis. Fuchs. Hist. 294.

Angl. Common Hart's-tongue.

In saxosis humidis, muris antiquis, puteis, aggeribusque umbrosis, frequens.

Perennis. Fl. Julio.

Radix

Radix fibrosa, fusca. Frondes pedales aut sesquipedales, erectæ, simplices, lanceolatæ, acutæ, basi cordato-excisiæ, margine paululum repandæ, utrinque lœtè virides, glabræ; stipitibus fuscis, sæpius squamosis. Lineæ fructiferæ numerosæ, obliquæ, parallelæ, luteofuscæ, distinctæ. Involucra linea longitudinali dehiscentia, integerrima, demum reflexa, capsulis prominentibus occultata.

Variat vel frondibus margine crassis, vel apice lobatis, rariusve stipite ramoso. Conf. Moris. sect. 14. t. 1.

Ceterach.

2. SCOLOPENDRIUM fronde pinnatifidâ lanceolatâ subtus squamosâ.

Scolopendrium Ceterach. *Syn. Syn.* 193. *Roth.*
Germ. v. 3. 48.

Scolopendron. *Cord. Hist.* 171. 2.

Scolopendria vera. *Trag. Hist.* 551. *Tabern. Kreuterb.* 1191.

Asplenium Ceterach. *Sp. Pl.* 1538. *Huds.* 452.
With. 766. *Hull.* 241. *Lightf.* 661. *Bolt. Fil.* 20. t. 12.

A. sive Ceterach. *Raii Syn.* 118. *Ger. em.* 1140.

A. vulgare. *Barel. Ic.* t. 1043.

Asplenium. *Matth. Valgr.* v. 2. 251. *Camer. Epit.* 640. *Dalech. Hist.* 1215.

Angl. Common Scaly Spleenwort.

In rupibus calcareis murisve antiquis.

Plentiful about Bristol. *Ray.* Malham, Yorkshire, and in Wales. *Richardson.* On Heydon church, Norfolk. *Rev. Mr. Bryant.* On the hill of Kinoul, Perthshire. *Lightf.* At Nunney, Somersetshire. *Rev. Dr. Goodenough.* Plentiful in the north of England, and about Bristol hot-wells.

Perennis. Fl. toto ferè anno.

Radix fibrosa, nigra. Frondes numerosæ, palmares, patentæ, lanceolatæ, pinnatifidæ, lobis alternis, obtusis, integerrimis;

integerrimis; suprà lèves; subtùs squamis membranaceis, lanceolatis, acutis, ferrulatis, reticulatis, fuscis, nitidis, undique tectæ; stipitibus etiam squamosis. Lineæ fructificantes numerosæ, obliquæ, squamis immersæ. Involucra margine integræ. Capsulæ aureo-fuscæ.

432. BLECHNUM. G. Pl. 560. Juss. 15.
Act. Taurin. v. 5. 411.

Fructif. in lineis longitudinalibus, continuis, costæ adjacentibus. *Involucr.* superficiarium, continuum, costam versus dehiscens.

1. BLECHNUM fronde pinnatâ glabrâ: pinnis boreale. linearibus obtusiusculis integerrimis basi subæqualibus.

Blechnum boreale. Swartz. in Schrad. Journ. ann. 1800. v. 2. 75. Engl. Bot. t. 1159.

B. Spicant. With. 765. Hull. 242. Sym. Syn. 193. Relb. 407. Roth. Germ. v. 3. 44.

Osmunda borealis. Salisb. Prodr. 402.

O. Spicant. Sp. Pl. 1522. Huds. 450. Abbot. 224. Bölt. Fil. 8. t. 6. Curt. Lond. fasc. 2. t. 67. Fl. Dan. t. 99. Hedw. Theor. 43. t. 5.

Acrostichum Spicant. Sibth. 267.

Lonchitis aspera. Raii Syn. 118. Ger. em. 1140.

Asplenon sylvestre. Trag. Hist. 550. Dalech. Hist. 1216.

Struthiopteris. Cord. Hist. 170. 2.

Angl. Northern Blechnum. Rough Spleenwort.

In humidioribus nemorosis, asperis, et ericetis.

Perennis. Fl. Julio.

Radix squamosa, nigra. Frondes pedales vel sesquipedales, erectæ, lanceolatæ, acutæ, saturatè virides, glabré, pectinato-pinnatæ. Pinnæ numerosissimæ, subalternæ,

alternæ, parallelæ, lineares, obtusiusculæ cum mucronulo, integerrimæ, basi parùm dilatatae, nequaquam auriculatae vel rotundatae; fructiferæ in distincta fronde, angustiores. Lineæ fructificantes a marginibus pinnum omnino remotæ, costis approximatæ et parallelæ, utrinque solitariae. Involucra integerrima, continua. Capsulæ fuscæ.

Habitu et fructificationis dispositione omnino Blechnum est, ut ex charactere et descriptione lucc clarius patet. Nomen absonum *Spicant* inventoribus relinquo.

433. PTERIS. *G. Pl.* 559. *Juss.* 15. *Act. Taurin.*
v. 5. 412.

Fructif. in lineâ marginali, continuâ. *Involucrum* e margine ipsius frondis inflexo, continuum, interiùs dehiscens.

aquilina. I. PTERIS fronde supradecompositâ: pinnulis lanceolatis acutiusculis: infimis pinnatifidis; superioribus minoribus.

Pteris aquilina. *Sp. Pl.* 1533. *Huds.* 451. *With.* 765. *Hull.* 242. *Reh.* 407. *Sibth.* 267. *Abbot.* 224. *Bolt. Fil.* 16. *t.* 10, malè. *Roth.* *Germ.* *v.* 3. 42.

Felix foemina. *Raii Syn.* 124. *Ger. em.* 1128. *Mattb. Valgr.* *v.* 2. 627. *Fuchs. Hist.* 596. *Tabern. Kreuterb.* 1181.

Felix. *Tillands. Ic.* 78.

F. majoris primum genus. *Trag. Hist.* 542.

β. F. faxatilis ramosa maritima nostras. *Raii Syn.* 125. *Pluk. Phyt.* *t.* 182. *f.* 1.

Angl. Common Brakes.

In pascuis sterilioribus, ericetis, et dumctis, ubique. β ad muros humidos.

Perennis. Fl. Julio.

Radix fusiformis, longa. Frondes tripedales, patulæ, pinnato-

pinnato-supradecompositæ, glabræ. Pinnæ oppositæ. Pinnulæ ultimæ alternæ, sessiles, latè lanceolatæ, acutiusculæ, integerrimæ, striatæ, lineâ fructificante undique marginatæ. Involucrum continuum, compressum, crenulatum, fuscum.

Varietas β primâ facie omnino alienam sese exhibit, habitu tenerimo Trichomanis, pinnulis rotundatis, confluentibus, sterilibus; at in solum felicius translatæ solitam induit formam.

2. PTERIS fronde supradecompositâ : pinnulis *crispa*.
sterilibus cuneatis incisis; fertilibus ellipticis
obtusis convexis.

Pteris crispa. *Linn. MSS. in Sp. Pl. 1522. With.*
764. Hull. 243. Engl. Bot. t. 1160.

Osmunda crispa. *Sp. Pl. 1522. Huds. 450.*
Lightf. 655. Bolt. Fil. 10. t. 7. Fl. Dan.
t. 496.

Onoclea crispa. *Roth. Germ. v. 3. 39.*

Adiantum album crispum alpinum. *Raë Syn.*
126.

A. album floridum. *Pluk. Phyt. t. 3. f. 2.*

Filicula petræa florida Anglicæ, foliis plurifariam
divisis. *Moris. sect. 14. t. 4. f. 4.*

Angl. Curled Stone Fern. Rock Brakes.

In petrosis elatis apricis Angliæ borealis.

Common in Wales. *Richardson*. Plentiful in Westmoreland and Scotland.

Perennis. Fl. Julio.

Radix fibrosa. Frondes longius stipitatæ, spithamæ aut pedales, glabræ, pallide virides, subdeltoideæ, duplikato- vel triplicato-pinnatæ. Pinnæ petiolatæ; steriles cuneatæ, nervosæ, planæ, apice incisæ; fertiles in distinctâ fronde, ellipticæ, obtusæ, integræ, convexæ. Involucrum latum, turgidum, crenatum.

434. ADIANTUM. *G. Pl. 560. Juss. 15.*
Act. Taurin. v. 5. 414.

Fructif. in punctis subrotundis, marginalibus, distinctis. Involucr. squamiformia, e margine ipsius frondis inflexo, distincta, interius dehiscentia.

Capillus Veneris. 1. ADIANTUM fronde alternatim decompositâ :

pinnulis petiolatis cuneiformibus lobatis.
Adiantum Capillus Veneris. Sp. Pl. 1558. Huds.
460. Wth. 781. Hull. 243. Lightf. 679.
Bolt. Fil. 24. t. 29. Dicks. H. Sicc. fasc. 6. 16.
Jacq. Misc. v. 2. 77. t. 7.

Adiantum. Matth. Valgr. v. 2. 542. Camer.
Epit. 924. Fuchs. Hist. 82.

Capillus Veneris verus. Dill. in Raii Syn. 123.
Ger. em. 1143.

Angl. True Maidenhair.

In rupibus et muris humidis, præcipue maritimis,
 rarijùs.

Found by Mr. Lhwyd at Barry Island and Port
 Kirig, Glamorganshire. *Dill.* Isle of Arran near
 Galloway; Mr. Stonestreet. *Herb. Sherard.*

Perennis. Fl. Maio—Septembri.

Radix fibrosa. Frondes palmates, spithameæ, quandoque
pedales, erecto-patulæ, formosæ, glabræ, alternatim
decompositæ, stipitibus rachibusque glaberrimis, nitidis,
atro-purpureis. Pinnulæ majusculæ, flabelliformes,
striatæ; basi cuneatæ; extùs lobatæ, incisæ, rotundatæ,
fructiferæ. Involucra reniformia, adpressa, pallida, demùm nigra.

435. CYATHEA. *Act. Taurin. v. 5. 416.*
Polypodium. G. Pl. 560. Juss. 15.

Fructif. sparsæ, subrotundæ, calyci hemisphærico,

rico, apice dehiscenti absque operculo, insidentes.

1. CYATHEA fronde bipinnatâ : pinnulis pin- *fragilis*. natifidis inciso-serratis, floribus sparsis, calyce lacero, rachi alatâ.

Cyathea fragilis. *Act. Taurin.* v. 5. 417. *Sym.* *Syn.* 194. *Roth. Germ.* v. 3. 94.

Polypodium fragile. *Sp. Pl.* 1553. *Huds.* 459. *With.* 779. *Hull.* 240. *Lightf.* 677. *Dicks.* *Dr. Pl.* 15. *Bolt. Fil.* 50. t. 27, & 46. *Ebrh. Crypt.* 151.

Filix saxatilis, caule tenui fragili. *Raii Syn.* 125. *Pluk. Phyt.* t. 180. f. 5.

2. *Polypodium rhæticum.* *With.* 780. *Hull.* 240. *Dicks. H. Sicc. fasc.* 1. 17. *Bolt. Fil.* 80. t. 45. nec Linn.

P. polymorphum A. *Villars. Dauph.* v. 3. 846. t. 53. f. A.

Filix montana ramosa minor, argutè denticulata. *Raii Syn.* 124.

F. alpina Myrrhidis facie Cambrobritannica. *Pluk. Phyt.* t. 89. f. 4.

3. *Polypodium Ilvense.* *Raii Syn.* 117. *Angl.* Brittle Cup-fern.

In rimis petrarum, vel muris antiquis, præcipue locis subalpinis, cōpiosè.

Perennis. Fl. Junio, Julio.

Radix squamosa, fusca. Frondes spithamææ vel pedales ; stipite fusco, nitido, lœvi ; lanceolatæ, acutæ, glabræ, bipinnatæ, pinnis subalternis. Pinnulæ incisæ, vel sæpius pinnatifidæ, obovatæ, vel obovato-lanceolatæ, acutè serratæ, acuminatæ, nervis flexuosis, rachibus generalibus et partialibus tenuè alatis. Maculæ fructiferæ numerosæ, pinnularum costas versùs, alternæ, globosæ, paryæ, albidae. Involucrum cyathiforme, capsulas

capsulas subtūs amplectens, membranaceum, lateraliter dehiscens, demūm lacerum, reflexum.

Frons in β major, et tenuius incisa. Pinnulæ inferiores huic quandoque magnitudine inæquales sunt, ut etiam in α , et plurimis aliis. γ monente D. H. Davies huc retuli.

regia. 2. CYATHEA fronde bipinnatâ : pinnulis lobatis obtusis muticis margine integerrimis, calyce lacero, rachi alatâ.

Cyathea regia. *Syn. Syn.* 194.

C. incisa. *Engl. Bot.* t. 163.

Polypodium regium. *Sp. Pl.* 1553. *Herb. Cliff.* *Hull.* 240.

P. trifidum. *With.* 779.

P. polymorphum C. *Villars. Dauph.* 847. t. 53. f. C.

Adiantum nigrum, pinnulis Cicutariæ divisurâ. *Raii Syn.* 126; ex exemplario a D. H. Davies benevolè communicato.

Angl. Laciniated Cup-fern.

In muris ; etiam in rupibus alpinis.

On a wall at Walthamstow. *Mr. T. F. Forster.* On Snowdon, but very rare. *Rev. H. Davies*, and *J. W. Griffith, Esq.*

Perennis. Fl. Junio.

Radix squamosa. Frondes palmatae, lanceolatae, glabrae, laetè virides, bipinnatae, pinnis suboppositis. Pinnulae variè lobatae, laciinis elliptico-oblongis, obtusis, muticis, margine integerrimis, nec ferratis neque dentatis, nervis rectiusculis. Raches generales et partiales tenuitèr alatae. Maculae fructiferæ ferè ut in præcedenti, sed minores. Involucrum demùm lateraliter reflexum, concavum, margine lacerum.

Archetypo in *Herb. Clifftonian* accuratiis inspecto, synonymon *Linnæanum* extrà dubium confirmatum manet.

Annotata

Annotata Richardsoniana in *Raii Syn.* 126, n. 8, huic speciei nequaquam conveniunt; monente D. Davies.

3. CYATHEA fronde subbipinnatâ: pinnulis ova-*dentata*, tis obtusis inciso-dentatis muticis.

Polypodium dentatum. *Dickf. Crypt. fasc.* 3. 1.
t. 7. f. 1. *H. Sicc. fasc.* 5. 16. *With.* 776.
Hull. 238.

Angl. Toothed Cup-fern.

In fissuris rupium alpinarum.

Rocks in the Highlands of Scotland. *Dickson.* At the foot of the walls of Castle Dinas Bran, Flintshire; and near Llangollen, Denbighshire. *Rev. H. Davies.*

Perennis. Fl. Julio.

Magnitudo ferè præcedentis. Frondes lanceolatæ, acutæ, glabræ, bipinnatæ vel subbipinnatæ. Pinnulæ ovato-subrotundæ, obtusæ, undique inciso-dentatæ, muticæ, nervis flexuosis. Rachis apicem versùs subinde alata. Maculæ fructiferæ numerosæ, approximatæ, demùm confluentes. Involucrum dimidiatum, margine fimbriatum.

436. HYMENOPHYLLUM. *Act. Taurin.* v. 5.

418. t. 9. f. 8.

Trichomanes. *G. Pl.* 560. *Juss.* 16.

Fructif. márgini frondis insertæ, distinctæ. *Involucr.* bivalvia, planiuscula, recta, exterius hiantia. *Columellæ inclusæ.*

ii. HYMENOPHYLLUM fronde alternatim bi-*Tunbrid-* pinnatifidâ laciniis involucroque ferratis, fruc-*gense,* tificationibus solitariis margini axillarum pin- narum insidentibus.

WOL. III.

Q

Hymen-

Hymenophyllum Tunbridgense. *Engl. Bot.*
t. 162. Sym. Syn. 195.

Trichomanes Tunbridgense. *Sp. Pl. 1561.*
Huds. 461. With. 781. Hull. 243. Bolt.
Fil. 58. t. 31. Fl. Dan. t. 954.

Adiantum petræum perpusillum Anglicum, fo-
liis bifidis vel trifidis. *Raii Syn. 123.*

A. radieosum humisparsum. *Pluk. Phyt. t. 3.*
f. 5; et erectius, f. 6.

β . *Trichomanes pyxidiferum.* *Huds. 461. Bolt.*
Fil. 56. t. 30. With. 781. Hull. 243.

T. Tunbridgense var. 3. *With. 782. Hull. 244.*
Filix humilis repens, foliis pellueidis et splen-
dentibus, caule alato. *Dill. in Raii Syn. 127.*
t. 3. f. 3, 4.

Angl. Filmy-leaved Fern.

In saxosis humidis vel umbrosis.

Near Tunbridge. *Mr. Dare, and Mr. T. F. For-*
ster. Amongst the pebbles at Coekbush on the
coast of Sussex. *Rand. Sherard.* In moist,
shady, and sometimes exposed, places on the
mountains of Scotland, Wales, Westmoreland,
Yorkshire, Cumberland, Devonshire, &c.

β at Belbank, scarcely half a mile from Bingley,
Yorkshire, at the head of a remarkable spring.
Richardson, Bolton, and Mr. Teesdale. On rocks
by Powerscourt cascade near Dublin. *Dr. Whitley*
Stokes.

Perennis. Fl. Maio, Junio.

Radix repens, capillaris, implexa. Frondes cæspitosæ,
decumbentes vel erectæ, magnitudine variæ, fusco-
virentes, membranaceæ, pellucidæ, alternatim bipin-
natifidæ, laciniis linearibus, obtusis, plerumque acutè
ferratis, (in β reverâ integerrimis,) nervo prominulo,
valido,

valido, fusco, medium percurrente. Fructificationes axillares ad pinnarum marginem superiorem, solitariæ, sessiles vel brevius pedunculatae, lenticulares, compressæ. Involucrum bivalve, margine erosum vel serratum. Capsulae plurimæ, fuscæ, globofæ, bivalves, annulo elastico cinctæ, columellæ brevi insidentes.

Varietas β a Tr. pyxidifero vero Linnæi et Plumierii omnino diversa est, nec specie a vulgari differre videtur, optimè monentibus D. D. Bolton et Griffith apud With.

437. PILULARIA. G. Pl. 561. Juss. 16.

Receptaculum commune globosum, quadriloculare, quadrivalve, intus antheris numerosis, superioribus, et germinibus plurimis, inferioribus, repletum. Sem. corticata.

I. PILULARIA.

Pilularia globulifera. Sp. Pl. 1563. Huds. 462. ^{globuli-} fera. With. 760. Hull. 244. Bolt. Fil. 72. t. 40. Engl. Bot. t. 521. Fl. Dan. t. 223. Roth. Germ. v. 3. 24.

P. palustris juncifolia. Vaill. Par. 159. t. 15. f. 6. Dill. Musc. 538. t. 79.

Graminifolia palustris repens, vasculis granorum piperis æmulis. Raii Syn. 136. Pluk. Phyt. t. 48. f. 1.

Pepper Grass. Pet. H. Brit. t. 9. f. 8.
Angl. Pillwort.

In inundatis glareosis.

Near Streatham wells, and on Hounslow heath.

Ray. At Hainford and St. Faith's Newton.

Mr. Crowe and Mr. Pitchford. About 2 miles from Mold on the north side of the Chester road; Mr. Griffith. With. Near Yarmouth.

Mr. D. Turner. On Hillingdon Common, Middlesex.

Perennis. Fl. Junio, Julio.

Frons universalis depreesa, filiformis, subramosa, repens, radicibus fibrosis; partiales alternæ vel fasciculatæ, erectæ, subulatæ, glabræ, evolutione circinales ut in veris filicibus. Fructificationes solitariæ, subsessiles ad basin frondium partialium, globosæ, villosæ, fuscæ, clausæ, intùs quadriloculares, parietibus tectis superne antheris clavatis, flavis, infernè germinibus subrotundis, albidis. Semina tunica fuscâ vestita.

Anne ad Monoeciam Polyandriam in Systemate artificiali transferenda?

438. ISOETES. G. Pl. 561. Juss. 17.

Masc. intra basin frondis interioris. *Anthera* solitaria, receptaculo obcordato inserta.

Fœm. intra basin frondis exterioris. *Germen* receptaculo obcordato insertum. *Capsula* membranacea, polysperma. *Sem.* angulata, scabra.

lacustris. I. ISOETES frondibus subulatis semicylindraccis, capsulis subrotundis bilocularibus, seminibus undique granulatis.

Isoetes lacustris. Sp. Pl. 1563. Huds. 462. With. 760. Hull. 245. Lightf. 683. Bolt. Fil. 74. t. 41. Fl. Dan. t. 191. Engl. Bot. t. 1084. Roth. Germ. v. 3. 25.

Calamaria folio breviore et crassiore. Dill. Musc. 540. t. 80. f. 1.

Subularia vulgaris erecta, folio rigidissimo. Raii Syn. 306.

S. lacustris. Raii Syn. ed. 1. 210. t. 2.

β. Calamaria folio longiore et graciliore. Dill. Musc. 541. t. 80. f. 2.

Subularia fragilis, folio longiore et tenuiore. Dill. in Raii Syn. 307.

Angl. Quillwort.

In

In fundo lacuum alpinorum.

In lakes of North Wales. *Lbwyd*, Richardson.

Loch Tay, Scotland. Mr. A. Menzies. Der-
wentwater, Cumberland. Mr. Crowe and Mr.
Woodward.

Perennis. Fl. Maio, Junio.

Radix tuberosa, fibris numerosis, simplicibus. Frondes
numeroſæ, simplices, subulatæ, acutæ, semieyindra-
ceæ, quadriloculares, articulatæ, glabræ, palmates
vel spithamææ, erectiusculæ; basi dilatatæ, membranaceæ,
intùs floriferæ; interiores masculæ, exteriores
femineæ. Capsula frondis basi immersa, magnitudine
pisi, subrotunda, compressiuscula, membranacea, bi-
locularis, polysperma. Semina parva, nivea, extùs
rotundato-gibba, intùs pyramidato-triquetra, undique
minutè granulata, nequaquam annulata.

Meliùs forsitan in Monoeciâ Monandriâ locum sibi vin-
dicaret.

II. MUSCI.

439. SPHAGNUM. G. Pl. 562. Juss. 12.

Schreb. Gen. 758. Hedw. Fund. v. 2. 85.

Capsula ore nudo. *Calyptra* circumscissa, cap-
sulam basi cingens. *Antheræ* annulo cinctæ.

i. SPHAGNUM ramis deflexis turgidis, foliis *latifo-*
ovatis obtusis ventricosis. *lium*.

Sphagnum latifolium. Hedw. Sp. Musc. 27.

S. cymbifolium. Swartz. Musc. Suec. 19.

S. obtusifolium. Hoffm. Germ. v. 2. 21. Ehrb.
Crypt. 241. Roth. Germ. v. 3. 118.

Q 3

S. palustre.

- S. palustre* α . *Sp. Pl.* 1569. *Huds.* 465. *With.* 783. *Hull.* 245. *Relb.* 412. *Sibth.* 273. *Abbot.* 228. *Fl. Dan.* t. 474. *Hedw. Fund.* v. 1. t. 1. f. 1. v. 2. t. 3. f. 9. *Theor.* t. 12. f. 43—45. t. 13. f. 46, 47. *Fl. Dan.* t. 474. *S. palustre* molle deflexum, squamis cymbiformibus. *Dill. Musc.* 240. t. 32. f. 1. *S. cauliferum* et ramosum palustre molle candidans, reflexis ramulis, foliis latioribus. *Dill. in Raii Syn.* 104. *Muscus palustris* in ericetis nascens floridus. *Raii Syn. ed.* 1. 241. *Pluk. Phyt.* t. 101. f. 1. *Vaill. Par.* t. 23. f. 3. *Angl.* Broad-leaved Bog-moss.

In palustribus turfosis copiosè.

Perennis. Fl. Junio.

Herba erecta, cæspitosa, mollis, scariosa, albida, interdum rubicunda. Rami subfasciculati, deflexi, foliosi, turgidi, apice attenuati; superiores congesti. Folia imbricata, latè ovata, obtusa, integerrima, ecarinata, enervia, ventricosa, inflexa. Pedicelli fructiferi terminales, erecti, breves, foliosi, apice strumosi. Capsula subcylindracea, brevis, fusca, operculo planiusculo.

- capilli-* 2. **SPHAGNUM** ramis deflexis filiformibus, foliis *farium,* ovato-lanceolatis planiusculis acuminatis adpresso.
- Sphagnum capillifolium. *Hedw. Sp. Musc.* 28. *Swärtz. Act. Holm. ann.* 1795. 231. *S. capillaceum.* *Swärtz. Musc. Suec.* 18. *S. acutifolium.* *Schrad. Spicil.* 58. *Ehrb. Crypt.* 72. *S. intermedium.* *Hoffm. Germ.* v. 2. 22. *Roth. Germ.* v. 3. 119. *S. palustre*

S. palustre β. *Sp. Pl.* 1569. *Huds.* 465. *With.* 783. *Hull.* 245. *Hedw. Fund.* v. 1. t. 3. f. 13—15.

S. palustre molle deflexum, squamis capillaceis. *Dill. Musc.* 243. t. 32. f. 2. A.

S. caulinferum et *ramosum* *palustre* molle candicans, réflexis ramulis, foliis angustioribus. *Dill. in Raii Syn.* 104.

Angl. Slender Bog-moss.

In ericetis udis montosis.

Perennis. Fl. Junio.

Color præcedentis, at sæpius rubicundus. Rami graciles, filiformes, penduli. Folia arctè imbricata, parva, lanceolata, sive ovato-lanceolata, plùs minùs acuminata, planiuscula, marginibus subinflexis, vix concava, nequaquam undulata. Capsula ovata, longius pedicellata, pedicello apice strumoso.

3. *SPHAGNUM* ramis deflexo-patentibus, foliis *cuspida-* lanceolatis acuminatis undulatis laxis. *tum.*

Sphagnum cuspidatum. *Hoffm. Germ.* v. 2. 22.
Roth. Germ. v. 3. 120. *Ehrh. Crypt.* 251.

S. palustris mollis deflexi, squamis capillaceis, varietas fluitans. *Dill. Musc.* 244. t. 32. f. 2. B.

Angl. Long-leaved floating Bog-moss.

In aquosis montosis.

Perennis. Fl. Junio.

Albido virescens. Caulès elongati, natantes, ramis laxè deflexis, apice attenuatis. Folia lanceolata, elongata, planiuscula, margine undulata, apice subulato-acuminata, patentiuscula. Capsula ovata, longius pedicellata, pedicello apice strumoso, ut in reliquis.

Species a præcedentibus distinctissima.

440. BUXBAUMIA. *G. Pl.* 562. *Juss.* 12.
Schreb. Gen. 762. *Hedw. Fund.* v. 2. 95.

Caps. ovata, obliqua, hinc gibba. *Peristomium* duplex; *exterius brevissimum*; *interius membranaceum*, plicatum.

foliosa. I. BUXBAUMIA acaulis, capsulâ subsessili, foliis linearî lanceolatis.

Buxbaumia foliosa. *Swartz. in Linn. Fil. Meth. Musc.* 33. t. 1. f. 4. *Musc. Suec.* 74. *Hedw. Sp. Musc.* 166. *With.* 790. *Hull.* 276. *Engl. Bot.* t. 329. *Dicks. H. Sicc. fasc.* 2. 20. *Roth. Germ.* v. 3. 344.

B. sessilis. *Schmidel. Buxb.* 26. t. 1. f. 1—xiv. *Hedw. Fund.* v. 2. 40. t. 9. f. 51.

Phascum. *Fl. Dan.* t. 249. f. *superior.*

P. montanum. *Huds.* 466.

P. maximum. *Lightf.* 693.

Sphagnum acaulon maximum, foliis in centro ciliaribus. *Dill. Musc.* 253. t. 32. f. 13.

S. n. 1725. *Hall. Hist.* v. 3. 24. t. 46. f. 3. *Angl.* Leafy Buxbaumia.

In rupibus umbrosis, subalpinis.

Near Llanberris. *Dill.* At Beddgelart near Snowdon plentifully. *Mr. D. Turner.* On Hart-fell near Moffat; *Mr. Sheffield. Huds.* On rocks about Tunbridge wells. *Mr. T. F. Forster.* At Hafod, Cardiganshire, on moist rocks near the river, below the house.

Annua. Fl. Julio.

Caulis brevissimus, simplicissimus, frè nullus. Folia plurima, radicalia, lanceolata, plana, carinata, acuminate; interiora apice in versus utrinque dente fetaceo. Capsula solitaria, subsessilis, magnitudine seminis Tritici hybneri, ovata, obliqua, glabra, virens. Calyptra subulata,

subulata, rubra. Operculum conicum, striatum. Peristomium exterius obsoletum; interius membranaceum, album, plicatum, clausum.

Ex errore apud Engl. Bot. calyptram pro operculo descripti.

441. PHASCUM. *G. Pl.* 562. *Juss.* 11. *Schreb.*
Gen. 758. *Hedw. Fund.* v. 2. 85.

Caps. ovata, clausa, decidua, operculo adnato,
non dehiscente.

i. PHASCUM caulescens, capsulâ subsessili, foliis *subulatis* patulis basi dilatatis apice capillaceis. *tum.*

Phascum subulatum. *Sp. Pl.* 1570. *Hedw. Sp. Musc.* 19. *Crypt.* v. 1. 93. t. 35. *Huds.* 466. *With.* 785. *Hull.* 251. *Relb.* 413. *Sibth.* 272. *Abbot.* 229. *Curt. Lond. fasc.* 4. t. 67. *Fl. Dan.* t. 249. f. 2. *Roth. Germ.* v. 3. 109. *Ehrb. Crypt.* 182.

Sphagnum acaulon trichodes. *Dill. Musc.* 251.
t. 32. f. 10.

Muscus trichoides minor acaulos, capillaceis foliis. *Vaill. Par.* 128. t. 29. f. 4.

Angl. Awl-leaved Earth-moss.

In ericetis et aggeribus arenosis.

Annua. Fl. Novembri; capsulas Martio perficit.

Radix fibrosa. Caulis brevis, simplex, foliosus, basi radicans. Folia imbricata, subulata, integerrima, basi dilatata, apice tenuissima, patula, lutescentia. Capsula ovata, subpedicellata, nitide fusca, apice lutescens.

22. PHASCUM caulescens, capsulâ ellipticâ mox *axillare*,
axillari pedicellatâ, foliis subulatis carinatis
patulis.

Phascum axillare. *Dicks. Crypt. fasc. 1. 2. t. 1.*
f. 3. Engl. Bot. t. 1036.

P. nitidum. *Hedw. Sp. Musc. 19. Crypt. v. 1.*
91. t. 34. Willd. 787. Hull. 252. Hoffm.
Germ. v. 2. 20. Roth. Germ. v. 2. 113.
Angl. Lateral-fruited Earth-moss.

In uliginosis ericetorum. *Dickson.*

In Kensington gardens. *Mr. Sowerby.*

Annua. Fructum maturat Julio—Septembri.

Radix fibrosa. Caulis erectus, subsimplex, foliosus. Folia alterna, patentia, subulata, carinata, nitica, integerim, subpellucida, lætè viridia. Pedicellus primo ortu terminalis, at mox, producto caulis apice, lateralis, axillaris, foliis brevior, subindè geminus. Capsula elliptica, fusca, mucronulo obliquo, pallido, demùm cernua.

Nomen Dicksonianum, certè aptissimum, et quia antiquius, Hedwigiano prætuli.

patens. 3. *PHASCUM* subcaulescens, capsulâ globosâ, foliis lanceolatis patentibus apice ferratis.

Phascum patens. *Hedw. Sp. Musc. 20. Crypt. v. 1. 28. t. 10. Roth. Germ. v. 3. 112. Dicks. Crypt. fasc. 4. 2.*

β . *P. recurvisolium.* *Dicks. Crypt. fasc. 4. 1. t. 10.*
f. 2.

Angl. Spreading Earth-moss.

In limosis.

Annua. Fructum maturat Martio, Aprili.

Prioribus minus. Caulis erectus, brevissimus, apice foliosus. Folia lancolata, patent-recurva, nitida, apicem versus plerumque ferrata, uninervia. Pedicellus terminalis, brevis, rectus, longius vaginatus. Capsula erecta, fusca, mucronulo pallido.

Folia in β angustiora, recurvata. *D. Turner.*

4. PHASCUM subbeaulescens, capsulâ ovatâ, foliis *strictum*.
subulatis erectis strictis obsoletè serratis:

Phaseum strictum. *Dicks. Crypt. fasc. 4. t. 1. f. 1.*

Angl. Upright Earth-moss.

In paludosis alpinis Scotiæ. *Dickson.*

Annua. Fl.

Præcedente minus, colore atro-virente, lurido, a congeneribus distinctum. Caulis vix ullus. Folia erecta, stricta, vel paululùm incurvata, subulata, uninervia, oculo armato subferrata, capsulam longè superantia. Pedicellus erectus, brevis. Capsula ovata, atro-fusca.

5. PHASCUM caulescens, foliis ovato-oblongis *piliferum*.
erectis setacco-aristatis, capsulâ globosâ erectâ.

Phascum piliferum. *Schreb. Phasc. 8. t. 1.*

f. 6—10. Hedw. Sp. Musc. 20. Dicks. Crypt. fasc. 2. t. 1. Roth. Germ. v. 3. 110. Hull. 252.

Angl. Bearded Earth-moss.

In grainiosis sterilibus.

In barren grassy places near London. *Dicks.*

Annua. Fructum profert Aprili.

Priore duplò majus. Radix fibrosa, cæspitosa. Caules breves, simplices vel ramosi. Folia ovata, seu ovato-oblonga, integerima, carinata, erecta, dilutè viridia, sæpè rubicunda vel ferruginea, apice pilo terminata, exsiccatione conniventia et paululùm contorta. Capsula pedicellata, globosa, erecta.

P. piliferum Witheringii est *Gymnostomum ciliatum*, fide exemplarii apud D. Turnerum.

6. PHASCUM caulescens ramosum, foliis caulinis *crispum*.
minimis alternis; floralibus subulatis arefac-

tione contortis, capitulo pluribus.

Phaseum crispum. *Hedw. Sp. Musc. 21. Crypt. v. 1.*

*v. i. 25. t. 9. Dicks. Crypt. fasc. 4. 2. Hoffm.
Germ. v. 2. 20. Roth. Germ. v. 3. 111.
Angl. Curled-leaved Earth-moss.*

In graminosis humidis pinguioribus.

By the sides of ditches in a marshy pasture at Beverley, Yorkshire. Mr. Teesdale. Near Kilcullen bridge, Ireland. Mr. R. Brown. Near Bedford. Rev. Dr. Abbot.

Annua. Martio capsulas ferens.

Caules cæspitosi, plerumque ramosi, patentes, foliosi. Folia luteo-viridia, integerima; inferiora exigua, alterna, lanceolata, concava; superiora majora, conferta; floralia maxima, basi dilatata, albida et subpellucida, apice elongata, subulata, arefactione incurvato-tortilia. Capsulæ terminales, subsessiles, erectæ, fuscæ, ellipticæ, mucronulo obliquo. Calyptra pellucida, longius acuminata.

*multi-
capsu-
lare.*

7. *PHASCUM caulescens ramosum, foliis caulinis alternis; floralibus linear-lanceolatis rectis, capsulis numerosis.*

Phascum sphærocarpum. Abbot. 230.

*P. crispum. Swartz. Musc. Suec. 17. Engl.
Bot. 618.*

Angl. Many-fruited Earth-moss.

In nemorosis.

In cart-ruts in Clapham Park wood, near Clapham Ford end, and other places near Bedford. Rev. Dr. Abbot.

Annua. Capsulas toto hyeme proferens.

Præcedenti proximum, at distinctum videtur colore tristiori minùsque lutescente, capsulis numerosioribus, denique foliis floralibus latioribus, lanceolatis, rectis, nunquam exsiccatione crispis, neque e basi dilatata statim in acumen subulatum contractis.

8. *PHAS-*

8. PHASCUM acaule, foliis lanceolatis aristatis *curvicol-*
patentibus, capsulâ ellipticâ, pedicello incur- *lum-*
vato.

Phascum curvicollum. *Hedw. Sp. Musc.* 21.

Crypt. v. 1. 31. t. 11. Dickf. Crypt. fasc. 2. 1.

H. Sicc. fasc. 4. 20. With. 786. Hull. 252.

Relb. 413. Abbot. 230? Schrad. Spicil. 58.

Roth. Germ. v. 3. 114. Engl. Bot. t. 905

nec 330.

Angl. Crooked-stalked Earth-moss.

In graminosis sterilibus.

Near Croydon, Surrey. Mr. Dickson.

Annua. Fructum fert Maio.

Planta exigua, cæspitosa, acaulis, simplex, radice ramo-
 sissimâ, densâ. Folia lanceolata, integerrima, carinata,
 aristata, apice subrecurva, carinâ rubrâ; sæpè tota ru-
 bicunda. Pedicellus longitudine foliorum, terminalis,
 solitarius, fuscescens, mox recurvus. Capsula exquisitè
 elliptica, fusca, reticulato-punctata, mucronulo subob-
 liquo.

9. PHASCUM acaule, foliis ovatis muticis *rectum-*
tibus, capsulâ ellipticâ utrinque subattenuatâ,
pedicello erecto.

Phascum rectum. *With. 787. t. 18. f. 1. Hull.*

252. Engl. Bot. t. 330, p. 905. Relb. 414.

Angl. Straight-stalked Earth-moss.

In aggeribus et campis aridis.

Near Garn house, Denbighshire. Mr. Griffith. On
 a dry bank near Newmarket. Rev. Mr. Hemsted.
 Near Norwich in the road to Lakenham. Mr.
 D. Turner. At Cullen's wood bridge near
 Dublin. Dr. Whitley Stokes.

Annua. Fructum fert Aprili vel Maio.

Statura præcedentis, at folia latiora, ovata, mutica. Pe-
 dicellus

dicellus foliis duplò longior, erectus, spadiceus. Capsula erecta, latè elliptica, rufa, basi apiceque subattenuata, mucronulo obliquo.

bryoides. 10. PHASCUM caulescens simplex, foliis ovatis aristatis erectis, capsulâ ellipticâ rostratâ, pedicello erecto stricto.

Phascum bryoides. *Dicks. Crypt. fasc.* 4. 3.
t. 10. *f.* 3.

Angl. Tall Earth-moss.

In ericetis et sylvis. *Dicks.*

At Clapham springs, Bedfordshire. *Rev. Dr. Abbot.*

Annua. Fructum fert vere.

Præcedente duplò majus. Caulis elongatus, erectus, simplex, foliosus. Folia imbricata, erecta, ovata, integerrima, aristato-acuminata, carinata. Pedicellus terminalis, erectus, strictus, crassiusculus, fusco-ferrugineus. Capsula erecta, fusca, latè elliptica, longius mucronulata.

Facies Gymnostomi truncatuli, vel Encalyptæ lanceolatæ, sed capsulâ latiori, ellipticâ, non dehiscente, discrepat.

curvise-tum. 11. PHASCUM caule brevissimo subsimplici, foliis oblongis aristatis, capsulâ globosâ, pedicello brevissimo geniculato.

Phascum curvisetum. *Dicks. Crypt. fasc.* 4. 2.
t. 10. *f.* 4.

Angl. Short bent-stalked Earth-moss.

In cultis prope Croydon. *Dickson.*

Near Bedford. *Rev. Dr. Abbot.*

Annua. Fructum fert Martio.

Caules cæspitosi, brevissimi, subinde bifidi. Folia congesta, ovato-oblonga, concava, integerrima, erecto-patentia, aristata, carinata. Pedicelli terminales, solitarii, brevissimi, mox geniculato-refracti. Capsula globosa, rufa, foliis obvoluta.

12. PHAS-

12. PHASCUM subcaulescens, foliis ovatis aris-*cuspida-*
tato-acuminatis: superioribus conniventibus, *tum.*
capsulâ latè ellipticâ subsessili.

Phascum cuspidatum. *Hedw. Sp. Musc.* 22.

Relh. 413. *Sibth.* 273. *Abbot.* 229. *Ehrb.*

Crypt. 162.

P. acaulon. *Sp. Pl.* 1570, α . *Huds.* 466, α .

With. 784. *Hull.* 251. *Curt. Lond. fasc.* 4.

t. 66. *Fl. Dan.* *t.* 249. *f.* 3.

Sphagnum acaulon bulbiforme majus. *Dill.*

Musc. 251. *t.* 32. *f.* 11.

S. acaulon, foliis in bulbi formam congestis,
majus. *Raii Syn.* 105.

Angl. Sharp-leaved Dwarf Earth-moss.

In ericetis, aggeribus et ambulacris, præsertim solo
arenoso, copiosè.

Annua? Fl. Novembri; capsulas Martio perficit.

Radix, monente Dillenio, sobolifera; anne igitur peren-
nis? Caulis brevissimus, simplicissimus. Folia con-
gesta, ovata, concava, integerrima, mucronata, con-
niventia; inferiora patula, minora; omnia pallidè vi-
ridia, basi albidiaria. Capsula subsessilis, terminalis,
fusca, latè elliptica vel subrotunda, foliis omnino tecta.

13. PHASCUM caulescens ramosum, foliis ob-*Schrebe-*
longis aristatis erectis, capsulâ globosâ sub-*rianum.*
sessili.

Phascum Schreberianum. *Dicks. Crypt. fasc.* 4. 2.

P. cuspidatum. *Schreb. Phasc.* 8. *t.* 1. *f.* 1—5,
excluso synymo Dillenii.

Angl. Schreberian Earth-moss.

In ericetis et cultis.

Annua. Capsulas vere perficit.

Priore május, colore saturatiori. Caules breves, at sæpius
ramosi, densè foliosi. Folia oblonga, plana, aristata,
erecta,

erecta, nec conniventia, atro-virentia, basi albida. Capsula ferè sessilis, globosa, fusca, foliis immersa. Folia valdè quam in reliquis longiora. *D. Turner.*

*muti-
cum.*

14. PHASCUM acaule, foliis ovatis muticis concavis conniventibus, capsulâ globosâ sessili.

Phascum muticum. Schreb. *Phasc.* 8. t. 1. f. 11—14. *Linn. MSS. in Sp. Pl.* 1570. *Hedw. Sp. Musc.* 23. *With.* 784. *Hull.* 251. *Roth. Germ.* v. 3. 114. *Swartz. Musc. Suec.* 18. *Ehrb. Crypt.* 172.

P. acaulon β. *Sp. Pl.* 1570. *Huds.* 466.

Sphagnum acaulon bulbiforme minus. *Dill. Musc.* 252. t. 32. f. 12.

S. acaulon, foliis in bulbi formam congestis, minus. *Dill. in Raii Syn.* 105.

Muscus trichoides acaulos minor latifolius. *Vaill. Par.* 128. t. 27. f. 2.

Angl. Common Dwarf Earth-moss.

Ad scps et in aggeribus frequens.

Annua. Capsulas profert hyeme.

Penultimo duplò minus. Caulis omnino nullus. Herba tota, e foliis conniventibus, ovata, luteo-virens, nitida. Folia ovata, concava, acutiuscula, mutica. Capsula sessilis, subrotunda, rubicunda, nitida, foliis obvoluta.

*serra-
tum.*

15. PHASCUM acaule, surculis patentibus decompositis capillaribus articulatis sterilibus, foliis lanceolatis acutis serratis.

Phascum serratum. Schreb. *Phasc.* 9. t. 2. *Hedw. Sp. Musc.* 23. *Dicks. Crypt. fasc.* 1. t. 1. f. 1. *With.* 785. *Hull.* 251. *Rehb.* 412. *Engl. Bot.* t. 460. *Roth. Germ.* v. 3. 115.

P. velutinum. Hoffm. *Germ.* v. 2. 20.

Angl. Capillary branched Earth-moss.

In collibus glareosis roridis.

On the north side of Muswell hill near Highgate.

Dickson. Cambridgeshire. *Rev. Mr. Hemsted.*

Acle' wood, Norfolk. Mr. D. Turner.

Annua. Fructum fert Martio vel Aprili.

Radix fibrosa, nequaquam repens. Herba omnium ferè minutissima, monocarpa. Surculi patentes, non radicantes, ramosi, alternatim decompositi, fusco-virides, capillares, tenuissimi, Confervæ instar articulati. Folia radicalia, conferta, erecto-patula, lanceolata, acuminata, pellucida, acutè ferrata. Capsula subsessilis, erecta, ovata, acuminata, fusca. Calyptra parva, dilutè fusca.

16. *PHASCUM acaule, surculis repentibus ra-stolonimosis articulatis, foliis ovato-lanceolatis acu-ferum. minatis dentatis.*

Phascum stoloniferum. Dicks. Crypt. fasc. 3. 1.

t. 7. f. 2. Hedw. Sp. Musc. 24. With. 786.

Hull. 252.

Angl. Branching-rooted Earth-moss.

In argillâ nudâ.

Near Walthamstow. *Mr. E. Forster junior.*

Perennis? Capsulas Aprili profert.

Radix e surculis ramosis, lævibus, subfuscis, articulatis, pellucidis. Caulis nullus. Folia conferta, erecta, ovato-lanceolata, concava, acuminata, dentata. Capsula sessilis inter folia, solitaria, subrotunda, acuminata, fusca.

Præcedentis varietatem censet D. Turner.

17. *PHASCUM caulescens ramosum, ramis sim-alterniplicibus: sterilibus altioribus erectis, foliis foliūm. subulatis alternis.*

Phascum alternifolium. Dicks. Crypt. fasc. 1. 2.

t. 1. f. 2. Hedw. Sp. Musc. 24. With. 786.

Hull. 252. Relh. 413. Abbot. 229.

Angl. Alternate-leaved Earth-moss.

In uliginosis.

On Gamlingay bogs, Cambridgeshire. *Relb.* On Stevington bogs, Bedfordshire. *Abbot.*

Annua. Capsulas profert Aprili.

Caules cæspitosi, adscendentes, foliosi, plerumque bifidi. Rami simplices; steriles elongati; fructiferi brevissimi, monocarpi. Folia alterna, subulata, brevia, uniformia, lætè viridia. Capsula sessilis, elliptica, foliis erectis, duplò longioribus, cincta.

442. GYMNOSTOMUM. *Hedw.* *Fund.*

v. 2. 87. *Schreb.* *Gen.* 758.

Hedwigia. *Hedw.* *Fund.* *v.* 2. 86.

Anictangium. *Hedw.* *Sp. Musc.* 40.

Caps. ore nuda, operculo deciduo. *Calyptra* integra e basi secedens.

truncatulum. 1. **GYMNOSTOMUM** foliis ovatis acuminatis planis integerrimis, capsulâ turbinatâ truncatâ.

Gymnostomum truncatulum. *Hoffm.* *Germ.* *v.* 2. 27. *Sibth.* 274.

G. truncatum. *Hedw.* *Sp. Musc.* 30. *Crypt.* *v.* 1. 13. *t. 5.* *Swartz.* *Musc. Suec.* 20. *Roth.* *Germ.* *v.* 3. 124.

Bryum truncatulum. *Sp. Pl.* 1584. *Huds.* 477. *With.* 818. *Hull.* 258. *Relb.* 424. *Abbot.*

242. *Curt.* *Lond.* *fasc.* 2. *t.* 71.

B. exiguum, creberrimis capsulis rufis. *Dill.* *Musc.* 347. *t.* 45. *f.* 7. *F—K.* *Herb. Dill.*

B. parvum, erectis subrotundis majusculis capitulis subfuscis, foliis serpilli pellucidis. *Dill.* *in Raii Syn.* 93.

Angl. Little Blunt-fruited Beardless-moss.

In

In pascuis, aggeribus, et ad fossas.

Annua. Fl. hyeme; fructum fert primo vere.

Caules cæspitosi, breves, plerumque simplices. Folia ovata, plana, patentia, integerrima, acuta, mucronulata, haud aristata, saturatè viridia. Pedicellus terminalis, erectus, caule longior, plerumque solitarius. Capsula fusca, basi ovata, apice truncata, ore dilatata, patula. Operculum magnum, rubrum, supra basin subcomplanatum, mucrone obliquo, teretiusculo, obtuso, longitudine ferè capsulæ.

Folia siccitate subtortilia et neutiquam conniventia.

D. Turner.

2. **GYMNOSTOMUM** foliis ovato-lanceolatis *intermixtis* acuminatis planis integerrimis, capsulâ ellip-dium, tico-oblongâ truncatâ.

Bryum truncatulum. *Ehrb. Crypt.* 14.

B. exiguum, creberrimis capsulis rufis. *Dill. Musc.* 347. t. 45. f. 7. A—E.

B. exiguum, erectis parvis subrotundis creberimis capitulis rufis, foliolis serpilli angustis pellucidis. *Dill. in Raii Syn.* 94.

Angl. Longer Blunt-fruited Beardless-moss.

In graminosis et aggeribus.

Annua. Fruct. fert ineunte vere.

Præcedenti valdè affine, at diversum est foliis angustioribus, pallidioribus, siccitate magis incurvis et conniventibus, pedicello saepius altiore; at præcipue capsulâ ferè ellipticâ, nec turbinatâ, operculo longitudine vix dimidii capsulæ. Magnitudine variat.

3. **GYMNOSTOMUM** foliis ovato-oblongis acutis planis subintegerrimis, capsulâ obovatâ truncatâ, operculo subulato.

Gymnostomum obtusum. *Hedw. Sp. Musc.* 34. t. 2. f. 1—3.

R 2

Bryum.

Bryum obtusum. *Dicks. Crypt. fasc. 2. 5. t. 4.*
f. 7. With. 811. Hull. 256.
Angl. Dark Beardless-moss.

In saxis et rupibus Scotiæ, etiam in arenosis Angliæ.

On Ben Crechan. *Dicks.* Extremely common about Yarmouth. *Mr. D. Turner.*

Annua. Fruct. fert Junio, Julio.

Prioribus paulò majus. Caules breves, cæspitosi, parùm ramosi. Folia ovato-oblonga, acuta, plana, nervo valido, pleruinque integerrima, aut vix denticulata, atro-viridia; inferiora decolora, persistentia. Pedicellus firmus, erectus, ruber, caule longior. Capsula rufo-fusca, obovata, brevis, truncata, ore patulo. Operculum convexisculum, acumine subulato, arcuato, capsulâ paululùm breviore.

ovatum. 4. **GYMNOSTOMUM** foliis ovatis obtusis integrimis concavis aristatis, capsulâ ovatâ.

Gymnostomum ovatum. *Hedw. Sp. Musc. 31.*
Crypt. v. 1. 16. t. 6. Hoffm. Germ. v. 2. 27.
Swartz. Musc. Suec. 20. Roth. Germ. v. 3.
123. Sibth. 274.

Bryum ovatum. *Dicks. Crypt. fasc. 2. 4. H. Sicc. fasc. 3. 20. With. 811. Hull. 256.*
Relb. 424. Abbot. 240.

Muscus coronatus minimus, foliolis et capitulis oblongis in pediculis brevissimis, *Moris. v. 3.*
631. sect. 15. t. 7. f. 18. Herb. Bobart.

M. capillaceus, omnium minimus. *Vaill. Par.*
t. 26. f. 2.

Angl. Hairy-leaved Beardless-moss.

In aggeribus, murisve e limo constructis.

Annua. Fl. Februario; fruct. fert Aprili.

Prioribus minus. Caulis brevissimus, simplicissimus. *Folia*

lia ovata, obtusa, coneava, integerrima, pilo albo satis conspicuo aristata. Pedicellus solitarius. Capsula fusca, ovata, basi subproducta, apice truncata, nec tamen patula. Operculum complanatum, mucrone rectiusculo, subulato, pallido, longitudine totius ferè capsulæ. Calyptra fusca.

5. **GYMNOSTOMUM** caule simplicissimo elongato basi nudo, foliis acutis distichis, capsulâ daceum, subrotundâ, operculo mutico.

Gymnostomum osmundaceum. Hoffm. Germ.
v. 2. 28.

G. *pennatum.* Hedw. Sp. Musc. 31. Crypt.
v. 1. 77. t. 29. Roth. Germ. v. 3. 126.
Bridel. Muscol. v. 1. 43.

Mnium osmundaceum. Dicks. Crypt. fasc. 1. 3.
t. 1. f. 4. H. Sicc. fasc. 12. 21.

Bryum pennatum. With. 821. Hull. 239.

Dicksonia pusilla. Ehrh. Crypt. 65.

Angl. Fern-leaved Beardless-moss.

In cavitatibus aggerum, sub sepibus vetustioribus,
solo pingui, Angliae australis.

In the road from Zele to South Tawton church, 4
miles from Okehampton, Devonshire; Rev.
Mr. Newberry. Dicks.

Annua. Fl. vere.

(Caules semiunciales, erecti, simplicissimi, filiformes, basi aphylli. Folia lanceolata, acuta, integerrima, mutica, plana, enervia, disticha, verticalia, decurrenti-connata, filicina; superiora angustiora, sparsa. Flores dioici, terminales, solitarii; masculus foliis rosaceis, acutis, cinctus. Pedicellus erectus, capillaris, apice paululum incrassatus. Capsula ferè globosa, ereta, ore fulvo. Operculum convexiusculum, muticum.

Nomen triviale ab inventore datum optimè restituit Cel.
Hoffmannus.

Griffithianum. 6. **GYMNOSTOMUM** foliis spatulatis obtusis enervibus carnosis, capsulâ obovatâ, operculo mutico.

Bryum Griffithianum. *Dicks. Crypt. fasc. 4. 8.*
t. 10. f. 10.

Splachnum froelichianum. *With. 794.* *Hull. 246;* nequaquam *Hedwigii.*

Angl. Succulent Beardless-moss.

In alpibus Cambrobritannicis.

On the east side of Snowdon near the summit.

Mr. Griffith.

Annua? Fl. æstate,

Caules basi ramosi, breves, foliosi. Folia congesta, oblongata, obtusa, integerrima, mutica, enervia, carnosæ, atro-virentia. Pedicellus solitarius, erectus, pallidus, crassus et carnosus. Capsula erecta, obovata, fusca. Operculum hemisphæricum, muticum.

Habitus Splachni.

Heimii. 7. **GYMNOSTOMUM** foliis elliptico-lanceolatis acutis apice serrulatis, capsulâ cylindraceâ, operculo obliquè rostrato.

Gymnostomum Heimii. *Hedw. Sp. Musc. 32.*
Crypt. v. 1. 80. t. 30. *Hoffm. Germ. v. 2.*
28. *Roth. Germ. v. 3. 123.* *Swartz. Musc. Suec. 21.*

Bryum Heimii. *Dicks. Crypt. fasc. 2. 4.* *With.*
815. *Hull. 257.*

Angl. Long-stalked Beardless-moss.

In arenosis. *Dicks.*

In Anglesea. *Rev. H. Davies.* Near Forfar.
Mr. G. Donn.

Annua, Fl. ineunte vere.

Caulis brevis, simplex. Folia latè lanceolata, acuta, plana, reticulata, latè viridia, apice tantùm ferrulata. Pedi-

Pedicellus uncialis, basi ruber. Capsula erecta, fusca, cylindracea, retusa. Operculum basi planum, muerone tereti, arcuato, gracili, capsulâ parùm breviore. Calyptra fusco-rufescens.

8. **GYMNOSTOMUM** caule ramoso cæspitoso, æruginis foliis subulatis carinatis integerrimis, capsulâ nosum. campanulatâ, operculo
Angl. Verdigrise Beardless-moss.

In palustribus alpinis.

Gathered in North Wales by J. W. Griffith, Esq.

Perennis. Fl. vere; fruct. fert æstate.

Caules dense cæspitosi; ramosi, erecti, foliis persistentibus undique vestiti. Folia imbricata, patentiuscula, exsiccatione incurva, subulata, canaliculata, integririma, lætè viridia et quasi æruginosa, nervo prominente; nequaquam enervia. Pedicelli erecti, capillares, parùm flexuosi. Capsula erecta, ovata, retusa, sive campanulata, fusca, nitida, orâ nudâ. Operculum non vidi.

9. **GYMNOSTOMUM** caule ramoso cæspitoso, luteo-foliis lanceolatis concavis carinatis, capsulâ lutea oblongâ, operculo hemisphærico mutico.

Gymnostomum æstivum. *Hedw. Sp. Musc.* 32.
t. 2. f. 4--7.

Angl. Yellowish Beardless-moss.

In alpibus Cambrobritannicis.

Sent by the Rev. H. Davies.

Perennis. Fl. . . .

Habitus præcedentis, at pluribus notis differt. Folia lutescentia, nec æruginosa, lanceolata, nec subulata, exsiccatione multò magis quam in priore incurva et contorta. Capsula elongato-cylindracea, pallidè fusca, minimè campanulata. Opereulum hemisphæricum cum exiguo umbone.

curvirostrum. 10. **GYMNOSTOMUM** caule ramoso cæspitoso, foliis subulatis recurvatis interruptè fasciculatis, operculo subulato curvato.

Gymnostomum curvirostrum. *Hedw. Sp. Musc.* 33. *Crypt. v. 2. 68. t. 24.* *Hoffm. Germ. v. 2. 28. Roth. Germ. v. 3. 125.*

Bryum æstivum. *Sp. Pl. 1585. Wiltb. 826. Hull. 254.*

B. *palustre*. *Huds. ed. 1. 411.*

B. *palustre æstivum*, conservæ facie. *Dill. Musc. 375. t. 47. f. 36.* *Herb. Dill.*

B. *angustissimis* foliis crebrioribus, capitulis erectis brevibus, pediculis e surculis novis et longis enascentibus. *Dill. in Raii Syn. 99.*

B. *verticillatum*. *Dicks. H. Sicc. fasc. 5. 19.*

Pottia curvirostra. *Ehrh. Crypt. 93.*

Angl. Curve-beaked Beardless-moss.

In rupibus gypsaceis, et palustribus alpinis.

On rocks in Cumberland. *Dickson.* Communicated to Dillenius by Dr. Richardson.

Perennis. Fl. vere.

Caules densè cæspitosi, elongati, terrâ calcareâ obducti. Folia alterna, densa, hinc indè fasciculata, subulata, recurva, canaliculata, carinata, lætè viridia, demùm pallidè lutescentia, persistentia. Pedicelli e fasciculis lateralibus foliorum erecti, rectiusculi. Capsula elliptica, retusa, basi margineque rubra. Operculum longitudine capsulæ, subulatum, obliquè curvatum. Peristomium omnino nullum.

stelligerum. 11. **GYMNOSTOMUM** caule ramoso, foliis linear-lanceolatis patentibus : summis stellatis, capsulâ hemisphæricâ, operculo obliquo.

Gymnostomum stelligerum. *Schrad. Journ. v. 2. 55. Bridel. Muscol. v. 1. 46.*

Bryum

Bryum stelligerum. *Dicks. Crypt. fasc.* 2. 3.
t. 4. f. 4. With. 823. *Hull.* 253.
Angl. Star-topped Beardless-moss.

In sylvis alpinis Scotiæ. *Dicks.*

Perennis. Fl. . . .

Caules erecti, patuli, ramosi. Folia alterna, linearilanceolata, integerrima, carinata, patentirecurva, lætè viridia; summa verticillato-stellata. Pedicelli terminales, solitarii, erecti, rubri. Capsula hemisphærica, ore patulo. Operculum complanatum, rostro subulato, subcurvato, longitudine capsulæ.

12. **GYMNOSTOMUM** caule simplicissimo, foliis microstomatis subulatis arefactione incurvis, operculo stomum subulato, capsulâ ellipticâ ore coarctato.

Gymnostomum microstomum. *Hedw. Sp. Musc.*

33. *Crypt.* v. 3. 71. t. 30, B. *Swaritz.*

Musc. Suec. 21. *Hoffm. Germ.* v. 2. 29. *Roth.*

Germ. v. 3. 128.

Bryum microstomum. *Dicks. fasc.* 4. 9.

Angl. Small-mouthed Beardless-moss.

In paucis. *Dickson.*

Perennis. Fl. vere.

Caules breves, cæspitosi, simplices. Folia densa, pallidè viridia, subulata, carinata, integerrima, basi dilatata, exsiccatione involuto-crispa. Pedicellus errectus, capillaris, caule longior. Capsula elliptico-oblonga, fusco-virens, ore rubro, valde coarctato. Operculum subulatum, arcuatum, breve.

Gymnostomorum minimum. *D. Turner.*

13. **GYMNOSTOMUM** caule simplici brevissimofasciculato, foliis ovatis acutis ferrulatis, capsulâ pyriformi, operculo convexo mutico.

Gymnostomum fasciculare. *Engl. Bot.* t. 1245.

Bryum fasciculare. *Dicks.* *Crypt. fasc.* 3. 3.
t. 7. f. 5. *With.* 815. *Hull.* 257.
B. Ægypti. *Hasselquist.* *It.* 502. *ed. Engl.* 263.
Hyssopus Salomonis. *Ibid.* 614. *Linn. Sp. Pl.*
1584. *sub Bryo truncatulo.* *Herb. Linn.*
Angl. Blunt Pear-shaped Beardleß-moss.

In pascuis sterilibus, montibusque.

On barren sandy ground both in England and Scotland. *Dicks.* In Angus-shire. *Mr. G. Donn.* North Wales. *Rev. H. Davies.* Ireland. *Mr. Templeton.*

Annua. Fl. hyeme et vere.

Caules breves, simplices, cæspitosi, radicibus copiosis, fuscis. Folia congesta, patēntia, ovata seu ovato-lanceolata, acutissima, ferrulata, arefactione torta, lœtè at saturatè viridia, oculo armato pulcherrimè reticulata. Pedicellus erectus, tenuis, nitidus, ruber. Capsula omnino pyriformis, basi rubicunda, apice fusco-virens, ore coarctato. Operculum pallidum, convexiusculum, muticum. Calyptra subulata, longissima.

pyrifor- 14. *GYMNOSTOMUM* caule simplicissimo brevissimo, foliis ovatis acutis denticulatis capsulâ obovatâ, operculo obtusè mucronato.

Gymnostomum pyriforme. *Hedw.* *Sp. Musc.* 38.

Hoffm. *Germ.* v. 2. 27. *Roth.* *Germ.* v. 3.

121. *Swartz.* *Musc. Suec.* 20. *Sibth.* 274.

G. fasciculare. *Hedw.* *Sp. Musc.* 38. t. 4. f. 5—8.

Bryum pyriforme. *Sp. Pl.* 1580. *Huds.* 475.

With. 812. *Hull.* 257. *Rehb.* 424. *Abbot.*

241. *Dicks.* *H. Sicc. fasc.* 4. 22. *Engl. Bot.*

t. 413. *Fl. Dan.* t. 537. f. 1.

B. parvum, erectis piriformibus majusculis capitulis, foliis serpilli pellucidis. *Raii Syn.* 93.

B. serpillifolium pellucidum, capsulis pyriformibus. *Dill. Musc.* 345. t. 44. f. 6.

Muscus capillaceus minimus, capitulis pyriformibus turgidis. *Vaill. Par.* t. 29. f. 3.

Angl. Pointed Pear-shaped Beardless-moss.

In aggeribus, campis et ericetis humidiusculis.

Annua. Fl. imprimis hyeme; fructum perficit Aprili, Maio.

Planta pallidè at lätè virens. Caulis ferè nullus. Folia latè ovata, concaviuscula, acuta, denticulata, pellucida et reticulata. Pedicellus firmus, nitidus, fulvus, basi incarnatus. Capsula viridis, magna, obovata, ore coarctato. Operculum pallidum, convexum, mucrone brevi, recto, obtuso. Calyptre subulata.

15. GYMNSTOMUM caule ramoso, foliis linear-lanceolatis canaliculatis a refactione cris-pis, capsulis truncatis sulcatis.

Gymnostomum lapponicum. *Hedw. Crypt.*

v. 3. 13. t. 5. A. *Swartz. Musc. Suec.* 20.

Schrad. Spicil. 59. *Ehrb. Beitr. fasc.* 7. 182.

Hoffm. Germ. v. 2. 29. *Roth. Germ.* v. 3. 128.

Anictangium lapponicum. *Hedw. Sp. Musc.* 40.

Bryum lapponicum. *Dicks. Crypt. fasc.* 4. 10.

Angl. Lapland Beardless-moss.

In faxibus grandioribus in alpibus Scoticis. *Dicks.*

On Ben Lawers. *Mr. R. Brown.* In Cwm Id-well, North Wales. *Mr. Griffith.*

Perennis. Fl. . . .

Caules unciales, erecti, cæspitosi, sæpiùs ramosissimi, densè foliosi. Folia saturatè viridia, imbricata, linear-lanceolata, canaliculata, carinata, integerrima, mutica, exsiccatione involuto-crispa. Pedicelli terminales, folitarii,

litarii, erecti, breves. Capsula erecta, cylindracea, brevis, truncata, longitudinaliter sulcata, et medio contracta. Operculum nondum vidi.

ciliatum. 16. **GYMNOSTOMUM** caule ramosissimo diffuso, foliis ovatis enervibus apice diaphanis, capsulis campanulatis subsessilibus.

Gymnostomum ciliatum. Swartz. *Musc. Suec.* 19. Roth. *Germ.* v. 3. 129. *Engl. Bot.* t. 1179.

G. Hedwigia. Hoffm. *Germ.* v. 2. 28.

Hedwigia ciliata. Hedw. *Crypt.* v. 1. 107. t. 40.

H. Anodon. Ehrb. *Crypt.* 192.

Anictangium ciliatum. Hedw. *Sp. Musc.* 40.

Bryum apocarpum β. *Sp. Pl.* 1579. *Huds.* 474. *With.* 809.

B. ciliatum. Dicks. *Crypt. fasc.* 4. 6.

Fontinalis albicans. Weber. *Gott.* 38. *Hull.* 275.

Phascum piliferum. *With.* 786 !

Hypnum n. 1792. *Hall. Hist.* v. 3. 40.

Sphagnum nodosum, hirsutum et incanum. Dill. *Musc.* 246. t. 32. f. 5.

S. cauliferum et ramosum saxatile hirsutum incanum, capitulis virentibus. Raii *Syn.* 105.

Muscus squamosus saxatilis, tortuosus ac nodosus. Vaill. *Par.* t. 27. f. 18. *Herb. Vaill.*

Angl. Branched Hoary Beardleſs-moss.

In saxis elatioribus.

Perennis. Fl: autumno.

Caules longitudine varii, ramosissimi, proliferi, diffusi et implexi, dense foliosi, demum basi denudati, nigri. Folia pallidè viridia, arctè imbricata, ovata, integerima, plana, encrvia, mucronata, mucrone subulato, plano, laxo, albo ac diaphano, ferrulato. Capsulæ terminales, solitariæ, ferè sessiles, ovatæ, retusæ, breves,

ves, superficie æquales, castaneæ, foliis majoribus, erectis, apicibus laceris, obvolutæ. Operculum convexum, vix mucronulatum.

443. SPLACHNUM. *G. Pl. 563. Juss. 11.
Hedw. Fund. v. 2. 88. t. 7. f. 33.*

*Caps. cylindrica, apophysi carnosæ insidens.
Peristomium simplex, dentibus sedecim, per paria approximatis.*

1. SPLACHNUM apophysi obconicâ viridi, foliis *mnioides*, elliptico-lanceolatis aristatis integerrimis.

Splachnum mnioides. Linn. Fil. Meth. Musc. 26.

Hedw. Sp. Musc. 51. Crypt. v. 2. 35. t. 11.

*Dicks. Crypt. fasc. 1. 2. With. 793. Hull.
246.*

Angl. Green Tapering Gland-moss.

In locis humentibus alpinis.

On the mountains of Scotland. *Dicks.* About Snowdon, on the dung of foxes and badgers. *Mr. Griffith.*

Annua. Fl. Maio, Junio.

Caules vix unciales, ramosi. Folia lætè viridia, erectiuscula, elliptico-lanceolata, acuta, integerrima, haud revoluta, in aristam flexuosam, lutescentem, desinentia. Pedicellus strictus, luteus, longitudine circitè caulis. Apophysis parva, obconica, virens, longitudine capsulae. Capsula vix apophysi angustior, lutco-fusca, ore dilatato, patente, nec, ut in iconе Hedwigianâ, coaretato. Operculum flavum, convexum, obtusum. Peristomium fuscum, inflexum.

2. SPLACHNUM apophysi obovatâ fuscescente, *angusta-* foliis lanceolatis acuminatis aristatis: *super-tum.* *rioribus ferratis.*

Splachnum

Splachnum angustatum. *Linn. Fil. Meth. Musc.*

33. *Hedw. Sp. Musc.* 51. *Crypt. v. 2.* 37.

t. 12. *Dicks. Crypt. fasc. 2.* 3. *With.* 792.

Hull. 246. *Engl. Bot. t.* 1132.

Angl. Narrow-leaved Gland-moss.

In humidis alpinis Scotiæ. *Dicks.*

By the side of Lochawen, on cow-dung. *Mr. G. Donn.*

Annua? Fl.

Caules simplices, dense cæspitosi, altitudine varii, longissimè radicati. Folia lætè viridia, lanceolata vel linear-lanceolata, acuminata ac longius aristata; superiora ferrata. Pedicellus præ cæteris hujus generis brevissimus, vix semiuncialis, fusco-luteus, rectus. Apophysis obovata, fusca, capsulâ duplò brevior, paucumque crassior. Capsula fusca, ore patulo. Peristomium pallidè flavebens.

urceola- 3. **SPLACHNUM apophysi obovatâ viridi-fuscescens.** foliis ellipticis concavis integerrimis aristatis.

Splachnum urceolatum. *Hedw. Sp. Musc.* 52.

Crypt. v. 2. 39. t. 13. *Dicks. Crypt. fasc. 2.* 2.

With. 793. *Hull.* 246. *Roth. Germ. v. 3.*

134.

S. ampullaceum. *Fl. Dan. t.* 192.

S. bryoides. *Zoeg. Fl. Island.* 12. *Von Troil.*

Isl. 119. *Herb. Linn.*

Angl. Elliptical Gland-moss.

In alpibus Scoticis.

On Ben High. *Dicks.*

Perennis. Fl. . . .

Caules densissimè cæspitosi, simplices et divisi. Folia elliptica, obtusa, ventricosa, integerrima, longius aristata;

tata, fusco-viridia. Pedicellus longitudine varius, quandoque uncialis, fulvus, rectus, crassiusculus. Apophysis obovata, viridis, demum fusca. Capsula fusca, demum medio constricta, apophysi maturâ angustior et duplò brevior. Operculum hemisphæricum. Peristomium flavescens.

4. **SPLACHNUM** apophysi obconicâ attenuatâ *tenue*.
fuscâ, foliis ovatis acutis serratis muticis.
Splachnum tenue. *Dicks. Crypt. fasc. 2. 2.*
With. 793. Hull. 246.
S. purpureum. *With. 794. t. 18: f. 9. Hull.*
246.
Angl. Slender Gland-moss.

In stercore vaccino in alpibus Cambroritannicis
et Scoticis.

At Leney near Stirling. *Dr. F. Buchannan.* 1782.
On Ben Lawers. *Mr. Dickson.* On the Corum
Gorum mountains of Inverness-shire. *Mr.*
G. Donn. On cow-dung about Llyn Idwell,
North Wales. *Mr. Griffith.*

Annua. Fl. Junio, Julio.

Caules semiunciales, subsimplices, cæspitosi. Folia ovata,
seu obovata, acuta, mutica, ferrulata, plana, lætè vi-
ridia, pulcherrimè ac manifestè reticulata, pellucida,
nervo validiore. Pedicellus solitarius, ferè biuncialis,
erectus, pulcherrimè rubens, apice flavus, totus de-
mum atro-sanguineus. Apophysis parva, obconica,
fusca vel atro-sanguinea. Capsula ovato-cylindracea,
apophysi crassior, atro-fusca. Operculum conicum,
breve, fuscescens. Columella exserta. Peristomii
dentes breves, fusi. Calyptra conica, gracilis, pal-
lida, margine fimbriata, apice fusca.

5. **SPLACHNUM** apophysi obovato - oblongâ *fastigia-*
atro-tum.

atro-sanguineâ, foliis lanceolatis acutis aristatis integerrimis.

Splachnum fastigiatum. *Dicks. Crypt. fasc. 3. 2.*
Swartz. Musc. Suec. 24. *Hull. 246.*

S. Brewerianum. *Hedw. Sp. Musc. 53.* *Crypt. v. 2. 105. t. 38.*

S. Breweri. *With. 792.*

S. mnioides. *Engl. Bot. t. 786,* apophysi haud satîs rubente.

Bryum ampullaceum, foliis et ampullis angustioribus. *Dill. Musc. 345.* *t. 44. f. 5.* *Herb. Dill.*

Angl. Dark Tapering Gland-moss.

In ericetis humentibus alpinis, e fimo vaccino putrido.

Discovered in North Wales by Mr. Brewer. *Dill.*

In the Highlands of Scotland. *Mr. Dickson* and *Mr. J. Mackay.*

Annua. Fl. Maio, Junio.

Caules densè cæspitosi, ramosi. Folia lanceolata, acuta, integerrima, subrevoluta, aristata, aristâ flexuosa, lutescente, longitudine variâ. Pedicellus sesquiuncialis, rectiusculus, fulvus. Apophysis obovata, gracilis, atro-sanguinea, exsiccatione nigricans et opaca, ut in aliis. Capsula diametro ferè apophysis, sed duplò brevior, rufo-fusca, ore plûs minus patulo. Operculum convexum, obtusè mucronatum. Peristomium inflexum, fuscum, haud rubrum.

Ex hujus exemplari in Herb. Linn. manu quâdam alienâ ad *S. mnioides* relato, error in synonymiam apud *Engl. Bot.* irrepsit. Specimina propria cum Dillenianis contuli.

ovatum. 6. **SPLACHNUM** apophysi ovatâ fuscescente, foliis ovatis acutis integerrimis.

Splach-

Splachnum ovatum. *Hedw. Sp. Musc.* 54. t. 8.
f. 4—6. *Dicks. Crypt. fasc.* 2. 2. *With.* 793.
Hull. 246.

Phascum pedunculatum. *Huds. ed.* 1. 397.
Bryum ampullaceum, foliis serpylli pellucidis,
 collo crassiore. *Dill. Musc.* 344. t. 44. f. 4.
B. erectis gigartinis capitulis, foliis serpilli pel-
 lucidis, obtusis. *Dill. in Raii Syn.* 93. t. 3.
 . 2.

Angl. Ovate Gland-moss.

In palustribus spongiosis alpinis.

In Lancashire and Montgomeryshire. *Dill.* In
 the north of England. *Richardson.* At Leney
 near Stirling. *Dr. F. Buchanan.* On Ben Nevis.
Mr. Dickson. Near Forfar. *Mr. G. Donn.*

Annua. Fl. Aprili.

Caulis brevissimus, simplicissimus. Folia latè ovata,
 mucronulata, integerrima, reticulata, nervo validiore.
 Pedicellus uncialis, strictus, firmus, fulvus, apice lu-
 teo-virens. Apophysis ovata, viridis, maturitate fus-
 cescens. Capsula medio constricta, apophysi brevior
 atque angustior, peristomio reflexo.

7. *SPLACHNUM* apophysi globosâ demùm cor- *rugosum.*
rugatâ fuscâ, foliis latè ovatis acutis integer-
 rimis.

Splachnum rugosum. *Dicks. Crypt. fasc.* 4. 3.
t. 10. f. 7.

Angl. Rugged Gland-moss.

In alpibus Scoticis. *Dicks.*

Annua. Fl. . . .

Caulis breves, subsimplices, basi denudati. Folia pa-
 tula, latè ovata, integerrima, carinata, evidentiùs re-
 ticulata, acuta, vix acuminata. Pedicellus erectus,
 semiuncialis, crassiusculus, rigidus, purpureo-ferru-
 gineus.

gineus. Apophysis rufo-nigricans, maturitate rugosa, capsulâ duplò crassior. Capsula cylindracea, fusca.

sphæri- 8. **SPLACHNUM** apophysi globosâ viridi, foliis cum ovato-lanceolatis acuminatis integerrimis, pedicello capillari.

Splachnum sphæricum. *Linn. Fil. Meth. Musc.*

33. t. 1. f. 1. Hedw. Sp. Musc. 55. Crypt. v. 2. 46. t. 16. Dicks. Crypt. fasc. 1. 3. With. 791. Hull. 245. Engl. Bot. t. 785. Roth. Germ. v. 3. 136.

Angl. Green Globular Gland-moss.

In fimo vaccino, locis alpinis Scotiæ atque Cambrobritanniæ.

On a mountain near Stirling. *Dr. F. Buchanan.*

On cow-dung near the old castle at Llanberris; also about Llyn Aled, Denbighshire. *Mr. Griffith.*

Annua. Fl. Junio.

Caules cæspitosi, erecti, simplices, semiunciales. Folia alterna, ovato-lancolata, sive ovato-spatulata, latitudine varia, longius acuminata, integerrima, carinata, dilutè viridia. Flores masculi solitarii, terminales, disciformes, foliis longioribus et angustioribus obvallati. Pedicellus in distincta planta, terminalis, erctus, longissimus (sæpè triuncialis) capillaris, rectiusculus, luteo-purpurascens; apice dilatatus et rubicundus. Apophysis globosa, viridis, lævis. Capsula longitudine apophysis at paulò angustior, dilutè fusca, medio aliquantulum constricta. Peristomium fusco-flavescens, dentibus erecto-inflexis. Calyptra conica, obtusa, integerrima, flava. Operculum convexum, muticum.

gracile. 9. **SPLACHNUM** apophysi globosâ ferrugineaâ, foliis ovatis acuminatis apice ferrulatis, pedicello flexuoſo.

Splachnum

Splachnum gracile. *Dicks. Crypt. fasc.* 4. 3.
t. 10. f. 5.

S. vasculosum. *Huds.* 469. *Lightf.* 697. *With.*
791. (descriptionibus falsis). *Hull.* 245.
Angl. Zigzag Gland-moss.

In ericetis montosis humidis, e' fimo vaccino.

In the North of England and in Wales. *Huds.* On Ben Lomond and in the Isle of Skye. *Lightf.* On cow-dung upon several wet moors about Kirkmichael, Dumfriesshire. *Rev. Dr. Burgess,* 1782. On mountains near Lough Bray, Ireland. *Mr. D. Turner.*

Annua. Fl. æstate.

Præcedente duplò minus. Caules cæspitosi, brevissimi. Folia latè ovata, breviùs acuminata, apicem versùs obsoletè ferrata. Pedicellus sesquiuncialis, laxus et flexuofus, fulvus. Apophysis obovato-globofa, rufo-ferruginea. Capsula cylindracea, apophysi paullò brevior, fusca. Operculum hemisphæricum, obtusissimè mucronulatum.

10. **SPLACHNUM** apophysi obconicâ obtusâ vi-ampul-ridi-purpureâ capsulâ triplò crassiore, foliis laceum. lanceolatis acutis ferratis.

Splachnum ampullaceum. *Sp. Pl.* 1572. *Hedw.*
Sp. Musc. 55. *Crypt.* v. 2. 41. *t. 14.* *Huds.*
468. *With.* 792. *Hull.* 246. *Lightf.* 696.
Relb. 414. *Sibth.* 275. *Abbot.* 231. *Engl.*
Bot. *t. 144.* *Fl. Dan.* *t. 822.*

S. n. 3. *Amœn. Acad.* v. 2. 280.

Bryum ampullaceum, foliis thymi pellucidis, collo strictiore. *Dill. Musc.* 343. *t. 44.* *f. 3.*

B. erectis gigartinis capitulis, foliis serpylli pellucidis acutis. *Raii Syn.* 93.

Muscus aureus capillaris minor, capitulis gemi-

natis erectis, mutuo incubitu adnatis. *Moris.*

Sect. 15. 629. t. 6. f. 10.

M. capillaceus minor, capitulis geminatis. Vaill.

Par. 130. t. 26. f. 4.

Angl. Purple Gland-moss.

In paludibus turfosis, et ericetis humidis, fimo vaccino putrido præcipue innascens.

Annua. Fl. Martio—Julio.

Caules densè cæspitosi, ramosi. Folia lanceolata, latitudine varia, interdùm ferè linearia, carinata, acuta, ferrata. Pedicellus biuncialis, rectiusculus, ruber. Apophysis magna, obovata, seu pyriformis, obtusa, pellucida, viridis, mox apud nos purpurascens. Capsula parva, gracilis, lutea. Calyptra conica, fusca. Operculum convexum. Peristomium patens, fuscescens.

*Turne-
rianum.* 11. **SPLACHNUM** apophysi obovatâ viridi, capsulâ ovato-cylindracea pedicellatâ, foliis lanceolatis acutis subserratis.

*Splachnum Turnerianum. Dicks. Crypt. fasc. 4.
3. t. 10. f. 11. Engl. Bot. t. 1116.*

*S. sagittifolium. Wiltb. 791. t. 18. f. 2, malè,
foliis alienis.*

Angl. Turnerian Gland-moss.

In paludibus montosis.

On a hill 3 miles from Exeter towards Sydmouth.

*Mr. D. Turner and Mr. Sowerby. In North
Wales. Mr. Griffith.*

Annua. Fl. Maio, Junio.

Caules brevissimi, simplicissimi. Folia lanceolata, acuta, medio dilatata, reticulata; superiora apicem versus ferrata. Pedicellus haud uncialis, rectus, ruber, apice virens. Apophysis dilatè viridis, obovata, gracilis. Capsula fusca, ovato-cylindracea, crassitudine ferè apo-

apophysis, pedicello brevi fusco insidens. Operculum rufum, planiusculum. Peristomium luteum, demum reflexum.

Præcedentis varietatem censet Cel. Swartz.

12. **SPLACHNUM** apophysi obconicâ capsulâ *lingula-*
angustiore fuscâ, foliis lingulatis obtusis in-*tum*.
tegerrimis muticis.

Splachnum lingulatum. *Dicks. Crypt. fasc. 4. 4.*
t. 10. f. 6.

Angl. Tongue-leaved Gland-moss.

In alpibus Scoticis. *Dicks.*

On Ben Lawers. *Mr. G. Donn.*

Annua? Fl.

Caules simplices vel divisi, vix unciales. Folia saturatè viridia, reticulata, lingulata, concaviuscula, obtusa, mutica, integerrima, uninervia, haud carinata; inferiora acutiora. Pedicellus uncialis, rectiusculus, castaneus. Apophysis parva, obconica, fusca sive castanea. Capsula cylindracea, apophysi paulò crassior, apice dilatata. Peristomium inflexum, breve, fuscum.

13. **SPLACHNUM** apophysi obconicâ capsulâ *reticula-*
angustiore fuscâ, foliis ovatis acutis apice *ser-tum*.
ratis muticis.

Splachnum reticulatum. *Swartz. Musc. Suec. 24?*

Bryum reticulatum. *Dicks. Crypt. fasc. 2. 4.*
t. 4. f. 6. With. 821. Hull. 255.

Angl. Reticulated Gland-moss.

In alpibus Scoticis.

On Ben High. *Dicks.*

Perennis. Fl.

Caules divisi, semiunciales. Folia ovata, acutiuscula, mutica, concava, pellucida, insigniter reticulata, apice versus ferrata. Pedicellus e basi caulis. Apophysis parva, obconica, fusca. Capsula obovata, apophysi crassior et longior. Peristomium inflexum. *Dicks.*

A Splachno Frœlichiano Hedwigii, quocum Cel. Swartzius conjungendum censet, distinctum videtur.

444. ANDRÆA. *Ehrb. Beitr. v. I. 15 et 180.*
Hedw. Sp. Musc. 47.

Capſ. breviflora. Peristomium dentibus quatuor, incurvatis, apice cohærentibus, operculum et calyptram gerentibus.

rupeſtris. 1. ANDRÆA foliis lanceolatis carinatis falcato-secundis.

Andreæa rupeſtris. *Hedw. Sp. Musc. 47. t. 7. f. 2.*

Jungermannia rupeſtris. Sp. Pl. 1601. Huds. 516. With. 882. Hull. 281. Lightf. 787. Dicks. H. Sicc. fasc. 7. 23.

Lichenastrum alpinum nigricans, foliis capillaceis reflexis. Dill. Musc. 507. t. 73. f. 40.

Angl. Dusky Rock Andræa.

In rupibus alpinis, udis.

In Wales and the North of England. *Huds.* Frequent on the Scottish mountains upon moist rocks. *Lightf.*

Perennis. Fl. vere et æstate.

Caulis cæſpitosi, erectiusculi, ramosi, foliosi, vix unciales. Folia tristè viridia, vel atro-fusciflora, imbricata, falcato-secunda, ovato-lanceolata, integerrima, carinata, enervia, apice ferè subulata, nec obtusa. Pedicelli terminales, solitarii, erecti, luteoli, exsiccatione demùm fuscescentes, foliis majoribus, arctè imbricatis, ad summitatem usque inclusi. Capsula cotyloformis, brevis, erecta, albida. Peristomium e dentibus quatuor, fuscis, lanceolatis, acutis, capsulâ duplò longioribus, medio distantibus, apice incurvatis et cohærentibus. Operculum in apicibus peristomii persistens, calyptrâ conicâ, albida, mucronatâ, tectum.

Distinct-

Distinctissimum Muscorum genus, optimè a Jungermaniis ab acutissimo Ehrharto sejunctum.

2. ANDRÆA foliis ovatis concavis undique im- *alpina*.
bricatis.

Andræa alpina. *Hedw. Sp. Musc.* 49.

A. petrophila. *Ebrh. Crypt.* 67. *Beitr. v. 1.*
192. *Roth. Germ.* v. 3. 359.

Jungermannia alpina. *Sp. Pl.* 1601. *Huds.* 517.
With. 882. *Hull.* 281. *Lightf.* 787. *Dicks.*
H. Sicc. fasc. 3. 23.

Lichenastrum alpinum atro-rubens teres, calyci-
bus squamosis. *Dill. Musc.* 506. t. 73. f. 39.

Angl. Chocolate Alpine Andræa.

In rupibus alpinis haud infrequens.

Perennis. Fl. Julio.

Differt a præcedente colore atro-purpureo. Folia undique
imbricata (nec secunda), brevia, lata, ovata, acuta,
concava. Pedicelli imprimis apice atro-purpurei. Cap-
sula albida. Peristomium saturatè fuscum.

445. TETRAPHIS. *Hedw. Fund.* v. 2. 87. t. 7.
f. 32. *Schreb. Gen.* 758.

Capf. oblonga. *Perist.* dentibus quatuor, py-
ramidalis, erectis, liberis.

1. TETRAPHIS.

Tetraphis pellucida. *Hedw. Sp. Musc.* 45. t. 7. f. 1. da.
a—f. *Swartz. Musc. Suec.* 21. *Roth. Germ.*

v. 3. 132. *Sibth.* 275. *Engl. Bot.* t. 1020.

Mnium pellucidum. *Sp. Pl.* 1574. *Huds.* 472.
With. 799. *Hull.* 249. *Rehb.* 417.

M. serpilli foliis tenuibus pellucidis. *Dill. Musc.*
232. t. 31. f. 2.

S 4

M. minus

M. minus non ramosum, angustioribus et pelucidis foliis. *Dill. in Raii Syn.* 78.

M. n. 1853. *Hall. Hist. v. 3. 56. t. 45. f. 8.*

Bryum pellucidum. *Abbot.* 237.

Georgia Mnemosynum. *Ehrb. Crypt.* 202.

Angl. Transparent Four-toothed-moss.

In umbrosis udis ad radices arborum.

Annua. Fl. vere; fruct. fert Maio.

Radices cæspitosæ, densissimè fibrosæ. Caules erécti, subsimplices, flexuosi, foliosi. Folia alterna, ovato-lanceolata, integerrima, mutica, tenera, pellueida, lætè virentia; inferiora et superiora minora. Flores terminales, solitarii, dioici; quandoque, monente Hedwigio, hermaphroditiæ: masculus in caule elatiori, teneriore, disciformis, bracteis latè ovatis cinctus. Pedicellus fœmineus rectus, uncialis, capillaris. Capsula cylindracea, fusca. Peristomium quadridentatum, rectum, magnum, fusco-nitidum, acutum. Operculum conicum, rufum. Calyptra conica, plicata, basi lacera.

446. ENCALYPTA. *Schreb. Gen.* 759. *Hedw.*
Sp. Musc. 60.

Leersia. *Hedw. Fnd.* v. 2. 88.

Capf. cylindracea. *Perist.* dentibus sedecim linearibus, erectis. *Calyptra* campanulata, inflata, laxa.

vulgaris. 1. ENCALYPTA calyprâ margine integerrimâ lævi, caule subsimplici, foliis lanceolatis.

Encalypta vulgaris. *Hedw. Sp. Musc.* 60. *Roth.*
Germ. v. 3. 151. *Sibth.* 277.

E. extintoria. *Swartz. Musc. Suec.* 24.

Leersia vulgaris. *Hedw. Crypt.* v. 1. 46. t. 18.
Ehrb. Crypt. 113.

Bryum

Bryum extlectorium. *Sp. Pl.* 1581. *Huds.* 475.
With. 816. *Hull.* 258. *Relb.* 422. *Abbot.*
239. *Dicks. H. Sicc. fasc.* 8. 18. *Engl. Bot.*
t. 558.

B. calyprâ extlectorii figurâ minus. *Dill.*
Musc. 349. *t. 45. f. 8.*
B. erectis capitulis, calyprâ laxâ conicâ, foliis
serpylli pellucidis angustioribus. *Raii Syn.* 92.
Angl. Common Extinguisher-moss.

In fissuris rupium, muris aggeribusque umbrosis.
Under Salisbury craigs, Edinburgh. *Dr. Younge.*
At Bungay castle. *Mr. Stone.* Burgh castle.
Mr. D. Turner.

Annua. Fructum fert Aprili, Maio.

Caules breves, plerumque simplicissimi, cæspitosi. Fo-
lia lanceolata, seu elliptico-lanceolata, acuta, concava,
integerrima, carinata, exsiccatione incurva. Pedicel-
lus rectus, vix ultrâ semiuncialis, rubicundus. Cap-
sula erecta, cylindracea, lævis. Operculum subulatum,
album. Peristomium exiguum, deciduum. Calyptra
maxima, capsulam maturam ultra basin operiens, cam-
panulata, indivisa, citrina, pellucida, apice subulata,
margine integerrima, lævis.

2. ENCALYPTA calyprâ margine dentatâ, caule *ciliata.*
ramoso, foliis lanceolatis.

Encalypta ciliata. *Hedw. Sp. Musc.* 61. *Roth.*
Germ. v. 3. 153. *Swartz. Musc. Suec.* 25.

Leersia ciliata. *Hedw. Crypt.* v. 1. 49. *t. 19.*
Ehrb. Crypt. 123. *Fl. Dan.* *t. 1001. f. 2.*

Bryum extlectorium β. *Sp. Pl.* 1581. *Huds.*
476. *With.* 817. *Hull.* 258. *Lightf.* 719.

B. calyprâ extlectorii figurâ, majus et ramosum.
Dill. Musc. 350. *t. 45. f. 9.* *Herb. Dill.*

Angl. Greater Extinguisher-moss.

In rupibus montosis.

On rocks about Ludlow Castle. *L. Brown.*

Perennis. Fructum fert æstate.

Priore major. Caules ramosi, ferè unciales. Folia numerosa. Pedicellus sæpiùs uncialis. Calyptra longius ac tenuius acuminata, margine dentato-fimbriata.

strepto- 3. **ENCALYPTA** calyprâ margine constrictâ lacerâ, caule ramoso, foliis oblongis, capsulâ spiralitè sulcatâ.

carpa. *Encalypta streptocarpa.* *Hedw.* *Sp. Musc.* 62. *t. 10. f. 10—15.*

E. grandis. *Swartz.* in *Schrad. Journ.* v. 2. 172.

Bryum contortum. *Wulfen.* in *Jacq. Coll.* v. 2. 226, excluso Dillenii synonymo.

B. ciliare. *Dicks. Crypt. fasc.* 4. 15. *D. Turner.*

B. n. 1828. *Hall. Hist.* v. 3. 48. *t. 45. f. 3.*

Hypnum saxatile erectum, ramulis teretibus, foliis subrotundis saturatè viridibus. *Dill. Musc.* 335. *t. 43. f. 71.* *Herb. Dill.*

Angl. Spiral-fruited Extinguisher-moss.

In montosis. *Dicks.*

Perennis. Fruct. fert autumno.

Caules unciales, biunciales, et ultrà, cæspitosi, simplices vel ramosi, undique foliosi. Folia saturatè viridia, undique imbricata, lanceolato-oblonga, obtusiuscula, integerrima, nervo valido pallenscente carinata, siccitate incurva et tortuosa. Pedicelli terminales, mox laterales, solitarii erecti, unciales, rigidi, purpureo-fusci, basi nudi. Capsula erecta, cylindracea, gracilis, egregiè fulcata, quasi in spiram contorta, fusca. Operculum subulatum, rectum, rufum, longitudine capsulæ, basi coccineum. Calyptra fructu duplò major, cylindracea, flavescentia, lœvis, basi paululum constricta, et margine lacera; apice acuminata, fusca. Peristomium rectum, rubrum, deciduum.

4. ENCALYPTA calyprâ sulcatâ margine la- *Daviesii*.
cerâ, caule ramoso, foliis subulatis.

Bryum Daviesii. *Dicks. Crypt. fasc. 3. 3. t. 7.*
f. 6. With. 835. Hull. 264.

Angl. Least Extinguisher-moss.

In rupibus.

In Anglesea. *Rev. H. Davies*. On the fallen co-
lumns of the Giant's causeway, Ireland, abun-
dantly. *Mr. R. Brown*.

Perennis. Fl. . . .

Caules vix semiunciales, cæspitosi, ramosi, densè foliosi.
Folia subulata, carinata, integerrima, fusco-viridia,
exsiccatione incurvato-crispa. Pedicelli terminales,
solitarii, erecti, breves, fulvi, basi vaginati. Capsula
ovato-cylindracea, truncata. Operculum convexum,
mucronatum, rectum. Peristomium breve, inflexum.
Calyptra flava, nitida, longitudinaliter sulcata, apice
fusca, basi demù lacera.

447. GRIMMIA. *Hedw. Fund. v. 2. 89.*
Schreb. Gen. 759.

Weisia. *Hedw. Fund. v. 2. 90.*

Caps. ovata. *Perist. dentibus sedecim, simpli-*
cibus, basi latioribus. Flores terminales. Ca-
lyptra cylindracea.

1. GRIMMIA foliis setaceis, capsulâ pyriformi *recurva-*
lævi ore dilatatâ, pedicello arcuato-recuivo. ta.

Grimmia recurvata. *Hedw. Sp. Musc. 75. Crypt.*
v. 1. 102. t. 38. Roth. Germ. v. 3. 141.
Host. Syn. 566.

Bryum recurvatum. *Dicks. Crypt. fasc. 2. 7.*
With. 838.

B. setaceum. *Wulf. in Jacq. Misc. v. 2. 96.*
t. 12. f. 1. Hull. 265.

Angl.

Angl. Recurved Grimmia.In graminosis sterilibus. *Dicks.*Near Forfar. *Mr. G. Donn.**Annua.* Fl. vere.

Planta exigua, eaule brevissimo, simplici, folioso. Folia numerosa, latè viridia, nee (ut in icona Hedwigianâ) glauca; tenuissima, setacea, integerrima, patentia, nunquam ineurva. Pedicellus solitarius, terminalis, capillaris, tenuissimus, primò erectus, mature scente fructu, recurvatus. Capsula olivacea, pyriformis, demum campanulata, lèvis. Operculum subulatum, incurvum, longitudine dimidii capsulae. Peristomii dentes breves, saturatè rubri, patuli, basi dilatati.

pusilla. 2. GRIMMIA subacaulis, foliis setaceis, capsulâ ovatâ erectâ lèvi ore dilatatâ.

Grimmia pusilla. *Roth. Germ.* v. 3. 147. *Schrad.* *Journ.* v. 2. 56.

Wcisia pusilla. *Hedw. Sp. Musc.* 64. *Crypt.* v. 2. 78. t. 29. A.

Afzelia pusilla. *Ehrh. Crypt.* 183.

Bryum paludosum. *Sp. Pl.* 1584. *Huds.* 478. *With.* 813. *Hull.* 257.

B. trichodes acaulon palustre minimum, setis et capsulis brevissimis. *Dill. Musc.* 387. t. 49. f. 53.

Angl. Dwarf Grimmia.

In palustribus montosis.

In the North of Yorkshire. *Dr. Richardson.* North Wales. *Mr. Griffith.* Near Ambleside, Westmoreland.*Annua.* Fructum fert Aprili—Junio.

Priore minor. Caulis vix ullus. Folia setacea, patentia, fusco-viridia. Pedicellus erectus, flavus, subflexuosus. Capsula ovata, erecta, lèvis, pallidè fusca, ore patulo, rubro.

rubro. Peristomii dentes rubri, patuli. Operculum subulatum, rectiusculum.

3. GRIMMIA foliis linearibus acutis carinatis *striata*.
arefactione crispis, capsulâ campanulatâ sulcatâ, operculo setaceo obliquo.
Grimmia striata. *Schrad. Journ.* v. 2. 57.
G. crispata. *Roth. Germ.* v. 3. 145.
Weisia fugax. *Hedw. Sp. Musc.* 64. t. 13.
f. 5—10.
Bryum crispatum. *Dicks. Crypt. fasc.* 3. 3. t. 7.
f. 4. *With.* 833. *Hull.* 264.
Angl. Striated Grimmia.

In paludosis Scotiæ. *Dicks.*

Annua. Fruct. fert Maio, Junio.

Præcedentibüs vix major. Caules dense cæspitosi, ramosi, undique foliosi. Folia pallidè viridia, linearia, integerrima, carinata, siccitate crispa et incurva. Pedicellus erectus, strictus, filiformis, luteus. Capsula matura campanulata, longitudinaliter sulcata, ore patulo. Operculum setaceum, tenuissimum, obliquum, longitudine capsulæ.

4. GRIMMIA foliis subulatis carinatis arefactione *Schisti*.
incurvis, capsulâ campanulatâ sulcatâ basi strumosâ, operculo conico.

Bryum Schisti. *Retz. Prod.* 261.

B. Fl. Dan. t. 538. f. 2.

Angl. Slate Grimmia.

In humidiusculis alpinis.

On the mountains of Scotland; Mr. G. Donn.

Mr. D. Turner.

Annua? Fruct. fert æstate.

Prioribus duplò major. Caules laxè cæspitosi, graciles, semiunciales, foliosi, subdivisi. Folia læte viridia, subu-

subulata, integerima, carinata, siccitate incurva. Pedicellus erectus, luteolus, nitidus, subflexuosus. Capsula campanulata, fulcata, ore patentiusculo, basi strumosa, vel apophysi obsoletâ instructa. Operculum conicum, nec fetaceum, obliquum, lutescens, capsulâ duplò brevius. Calyptra e Floræ Danicæ auctoritate nigra.

Starke- 5. **GRIMMIA** foliis ovatis submuticis, capsulâ *ana.* ellipticâ erectâ operculo conico obtuso.

Grimmia Starkeana. *Roth. Germ.* v. 3. 146.
Weisia Starkeana. *Hedw. Sp. Musc.* 65. *Crypt.*
v. 3. 83. t. 34. B.

Bryum Starkianum. *Hoffm. Germ.* v. 2. 32.
B. minutum. *Dicks. Crypt. fasc.* 4. 7. t. 10.
f. 17.

Angl. Minute Oval-leaved Grimmia.

In pasuis. *Dicks.*

At Rathmines bridge near Dublin; Dr. W. Stokes.
Mr. D. Turner.

Perennis? Fruet. fert autumno.

Planta exigua. Caulis brevissimus. Folia imbricato-patentia, ovata, integerima, carinata, apice submu-cronulata. Pedicellus erectus, strictus. Capsula el-liptica, fusca, lœvis. Operculum, ex auctoritate Hed-wigii, conicum, obtusum, breve.

lanceola- 6. **GRIMMIA** foliis lanceolatis concavis aristatis,
ta. capsulâ ovatâ, operculo subulato obliquo.

Lcersia lanceolata. *Hedw. Sp. Musc.* 63. *Crypt.*
v. 2. 65. t. 23. *Ehrh. Crypt.* 133.

Encalypta lanceolata. *Roth. Germ.* v. 3. 150.
Sibth. 278.

Bryum lanceolatum. *Dicks. Crypt. fasc.* 3. 4.
H. Sicc. fasc. 11. 18. *With.* 824. *Hull.* 260.
Relb. 422. *Abbot.* 240.

Angl.

Angl. Lanceolate Bearded Grimmia.

In ericetis et aggeribus aridis.

Perennis. Fruct. fert Maio.

Caules conferti, simplices, foliosi. Folia lato-lanceolata, obtusiuscula, integerrima, pilo aristata. Pedicellus erectus, ruber, caule parum longior, siccitate tortus. Capsula erecta, ovata, fusca, laevis, ore subcoarctata. Operculum subulatum, arcuato-obliquum, capsulâ duplò brevius.

7. GRIMMIA foliis cylindraceis muticis basi *calcarea*, latatis, capsulâ turbinatâ, operculo subulato arcuato.

Weisia calcarea. *Hedw.* *Sp. Musc.* 66. t. 11. f. 1—6.

Bryum calcareum. *Dicks. Crypt. fasc.* 2. 3. t. 4. f. 3. *H. Sicc. fasc.* 2. 21. *With.* 812. *Hull.* 257. *Reh.* 427. *Abbot.* 243. *Engl. Bot.* t. 191.

Angl. Chalk Grimmia.

In collibus et clivis cretaceis.

On Gogmagog hills, and near Newmarket. *Mr. Dickson.* In the chalk-pits at Dartford in great plenty. *Mr. Sowerby.* On Barton hill, Bedfordshire. *Rev. Dr. Abbot.*

Perennis. Fruct. fert vere et aestate.

Planta exigua, simplicissima. Folia lætè viridia, integerrima; primordialia ovata, concava; adultiora in acumen cylindraceum, rectum, obtusum, muticium, producta. Pedicellus rectus, foliis duplò altior. Capsula parva, brevis, turbinata. Peristomii dentes plani, basi dilatati. Operculum curvatum, longitudine capsulae.

8. GRIMMIA foliis subulatis siccitate crispis basi *contro-dilatatis*, capsulâ ellipticâ, operculo subulato *versa*, arcuato.

Grimmia controversa. *Sibth.* 277.

G. crispa. *Roth. Germ.* v. 3. 142.

Weisia controversa. *Hedw. Sp. Musc.* 67. *Crypt.* v. 3. 13. t. 5. B. *Swartz. Musc. Suec.* 26.

Afzelia crispa. *Erbh. Crypt.* 222, proculdubio.

Bryum virens. *Dicks. Crypt. fasc.* 1. 4. *H. Sicc.* *fasc.* 8. 20. *Reh.* 425. *Abbot.* 243.

B. viridulum. *Huds.* 478. *With.* 817. *Hull.* 258. *Curt. Lond. fasc.* 2. t. 70. f. 1. *Lightf.* 731. *Weis.* 193.

B. controversum. *Hoffm. Germ.* v. 2. 34.

B. capillaceum breve, pallidè et lætè virens,
capsulis ovatis. *Dill. Musc.* 380. t. 48. f. 43.

B. trichoides exile, erectis capitulis in pediculis
brevissimis. *Dill. in Raii Syn.* 97; excluso
Richardsonii synon.

Muscus coronatus minimus capillaceis foliis, ca-
pitulis oblongis. *Morif.* v. 3. 631. *sect.* 15.
t. 7. f. 19.

Angl. Little Green Grimmia.

Ad sepes et aggeres fabulosos.

Perennis. Capsulas fert Februario.

Caules breves, simplices, erecti, cæspitosi, foliosi, de-
mum basi denudati. Folia pulchrè luteolo-viridia,
subulata, acuta, integerrima, basi dilatata et concava,
exsiccatione involuto-crispa; inferiora breviora. Pe-
dicellus rectus, capillaris, flavescens, pellucidus. Cap-
sula exquisitè at latè elliptica, virens, mox fusca,
lævissima, nitida, ore rubro, haud patulo. Opercu-
lum setaceum, arcuatum, capsulâ brevius. Peristo-
rium incurvum, rubicundum, deciduum. Calyptra
subulata, pallida, pellucida.

Dickso- ni. 9. **GRIMMIA** foliis linear-lanceolatis carinatis
siccitate crispis, capsulâ elliptico-cylindracea
coarctatâ, operculo filiformi.

Weisia

Weisia Dicksoni. *Wilden. Berol.* 303, e Dicksoni auctoritate.

W. cirrata. *Hedw. Sp. Musc.* 69. t. 12. f. 7—
12.

Encalypta cirrata. *Swartz. Musc. Suec.* 25.

Afzelia cirrhata. *Ehrb. Crypt.* 232.

Mnium cirratum. *Sp. Pl.* 1576.

Bryum Dicksoni. *Dicks. Crypt. fasc.* 3. 4. *With.*
826. *Hull.* 261.

B. cirratum et stellatum, tenuioribus foliis. *Dill.*
Musc. 379 t. 48. f. 42.

B. trichoides exile, erectis capitulis in pediculis
longioribus rubris. *Dill. in Raii Syn.* 97.

Hypnum n. 1786. *Hall. Hist.* v. 3. 38.

Angl. Dicksonian Grimmia.

In aggeribus sepium, in arborum truncis et ad
palos, situ humidiori.

Perennis. Fructum fert Aprili.

Caules dense cæspitosi, fastigiati, ramosi, semiunciales
vel unciales, undique foliosi. Folia imbricata, linear-
lanceolata, acuta, integerrima, carinata, lætè viridia,
arefactione contorta et crispa; inferiora persistentia
atque nigricantia. Pedicelli numerosi, semiunciales;
recti, capillares, flavescentes. Capsula elliptico-cy-
lindracea, erecta, lævis, fusca, nitidiuscula, ore valde
coarctato, rubro. Peristomium exiguum, rubrum.
Operculum setaceum, rectiusculum, flavum, basi ru-
brum, longitudine dimidii capsulæ. Calyptora cylin-
dracea, fuica.

Præcedente triplò major; quacum a Celeberrimo Wil-
denovio confunditur; hinc synonymorum apud auctores
confusio.

10. GRIMMIA foliis setaceis carinatis siccitate *cirrata*,
crispis, capsulâ ovatâ ore dilatato patulo.

VOL. III.

T

Bryum

- Bryum cirratum. *Huds.* 485? *With.* 824.
Hull. 260. *Abbot.* 238. *Lightf.* 728?
 Muscus muralis minimus roseus, sive stellaris,
 capitulis longiusculis acutis erectis. *Vaill.*
Par. t. 24. f. 8; ex iconē, nec charactere.
Angl. Frizzled Grimmia.

In truncis arborum et ericetis.

Perennis. Fructum fert vere.

Præcedenti maximè affinis; differt verò foliis magis lu-
 tescentibus, duplòque angustioribus, omnino setaceis,
 basi aliquantulum dilatatis; at præcipue capsulis bre-
 vioribus, ovatis, pallidis, ore dilatato, patentissimo;
 nequaquam coarctato. Operculum nondum mihi vi-
 deri contigit.

- recurvi-* 11. GRIMMIA foliis linear-lanceolatis carinatis,
rostra. capsulâ cylindraceâ, operculo conico-cylin-
 draceo obtuso recurvo.
 Grimmia recurvirostra. *Roth. Germ.* v. 3. 144.
Sibth. 277.
 Weisia recurvirostra. *Hedw. Sp. Musc.* 71.
Crypt. v. 1. 19. t. 7.
 Bryum curvirostrum. *Dicks. Crypt. fasc.* 2. 7.
Dr. Pl. 95. *Hull.* 262. *Hoffm. Germ.* v. 2.
 32.
 B. recurvirostrum. *With.* 829.
 B. setaceum β. *Huds.* 482.
 B. stellare lacustre, foliis rubris capillaceis. *Dill.*
Musc. 382. t. 48. f. 45. *Herb. Dill.*
Angl. Blunt-curved-beaked Grimmia.

In sylvis et montosis.

On the summit of Glyder mountain, North Wales,
 a little above the lake Llyn y cwn. *Dill.* In
 woods

woods in dry places. *Dicks.* Near Forfar, but rare. *Mr. G. Donn.* Ireland. *Dr. Scott.*

Perennis. Fructum fert Augusto.

Caules erecti, cæspitosi, simplices vel ramosi, altitudine varii, undique foliosi. Folia patentia, laxa, linearilanceolata, acuta, carinata, integerrima, saturatè viridia, basi vaginantia, a refactione plūs minus contorta; inferiora demum rufo-ferruginea. Pedicelli rubri, capillares, rectiusculi, vix unciales. Capsula ovato-cylindracea, gracilis, erecta, rufa, nitida, ore nec patulo nec coarctato. Peristomium rubrum. Operculum breve, conicum, apice subcylindraceum, obtusum, plerumque recurvum.

12. **GRIMMIA** foliis linear-lanceolatis acutis *verticillata*.
verticillato-fasciculatis, capsulâ ovatâ truncata,
 catâ, operculo subulato arcuato.

Bryum verticillatum. *Sp. Pl.* 1585. *Huds.* 485.
With. 821. *Hull.* 259.

B. fasciculatum. *Dicks. Crypt. fasc.* 3. 3, ex-
 cluso synomo. *With.* 828. *Hull.* 261.
Swartz. in Schrad. Journ. v. 2. 172.

B. pilosum verticillatum. *Dill. Musc.* 374. t. 47.
 f. 35.

B. trichoides brevifolium, angustis cauliculis,
 capitulis erectis parvis et minus aduncis. *Dill.*
in Raii Syn. 98.

Angl. Whorled Grimmia.

In irriguis subalpinis, solo calcareo.

In Yorkshire. *Richardson.* On rocks by the sea
 near Bangor. *S. Brewer and Rev. H. Davies.*

Perennis. Fruct. fert Augusto.

Caules dense cæspitosi, graciles, ramosi, foliosi, unciales
 vel biunciales, inferne foliis persistentibus, decoloratis,
 terrâ calcareâ granulatâ saepius obductis, vestiti. Folia
 imbricata, erecto-patula, hinc et hinc fasciculata, quasi

T 2
verti-

verticillata, linear-lanceolata, angusta, integerima, acuta, mutica, uninervia, planiuscula, lätè viridia, subglaucouscentia, siccitate vix incurva. Pedicelli solitarii, terminales, mox laterales, erecti, capillares, luteo-fuscescentes, a refactione contorti. Capsula eretta, parva, ovata, brevis, truncata, parùm obliqua, rufo-fusca, nitida, demùm subcorrugata, ore patulo, rubro. Peristomium rubrum, siccitate obliquè decumbens. Operculum rubrum, conicum, acumine subulato, arcuato, capsulâ brevius.

Muscus Hallero, Hedwigio, Lightfootio, aliisque plurimis botanicis planè ignotus, nec nisi Dillenii fide a Linnæo in *Sp. Pl.* recensus.

crispula. 13. GRIMMIA foliis lanceolatis carinatis siccitate incurvato-crispis, capsulâ ellipticâ.

Weisia crispula. *Hedw. Sp. Musc.* 68. t. 12, f. 1—6.

Angl. Lesser Curled Griminia.

In turfosis.

In Anglesea. *Rev. H. Davies.*

Perennis. Fruct. fert æstate.

Caules densè cæspitosi, ramosi, fastigiati, foliosi, unciales. Folia imbricata, fusco-viridia, lanceolata, concaviuscula, carinata, integerima, acutiuscula, a refactione incurvato-crispa. Pedicelli breves, recti, flavescentes, demùm sæpè denigrati. Capsula parva, eretta, exactè elliptica, dilutè fusca, lævis, tenuis, ore nigricante, haud patulo. Operculum

acuta. 14. GRIMMIA foliis subulatis apice capillaribus, capsulâ ovatâ ore dilatatâ, operculo setaceo obliquo.

Weisia acuta. *Hedw. Sp. Musc.* 71. *Crypt. v. 3.* 85. *t. 35.* *Swartz. Musc. Suec.* 26.

Bryum acutum. *Huds.* 484. *With.* 823. *Hull. 260.* *Dicks. H. Sicc. fasc.* 17. 20.

B. verti-

B. verticillatum. *Ligbtf.* 733. *Huds. ed.* 1.
411, β.

B. pilosum, Sphagni subulati facie. *Dill. Musc.*
374. t. 47. f. 34.

Angl. Sharp-pointed Grimmia.

In alpibus Cambrobritannicis et Scoticis.

In Ireland. *Dr. Scott.*

Perennis. Fructum fert Augusto.

Caules ferè triunciales, cæspitosi, graciles, apice subdivisi ac densius foliosi. Folia subsecunda, luteo-viridia, fuscescentia, subulata, integerrima; basi dilatata; apice acuminato-capillaria, arcuata, nunquam arefactione crispa. Pedicelli mox laterales, foliis basi vaginati, erecti, graciles, siccitate spiraliè torti. Capsula ova-ta, brevis, fusca, laevis, ore patulo, basi subproducta. Operculum longitudine capsulæ, subulatum, tenue, arcuatum. Peristomium rubrum. Calyptra fusca.

Bryum splachnoides amicissimi Dicksoni ab hâc specie mihi non videtur differre.

15. GRIMMIA foliis setaceis basi dilatatis vagi-nantibus, capsulâ ovatâ obliquâ cernuâ, oper-ta-culo conico.

Swartzia inclinata. *Hedw. Crypt.* v. 2. 74.
t. 27.

Afzelia inclinata. *Erbh. Crypt.* 193.

Didymodon inclinatum. *Swartz. Musc. Suec.* 28.

Cynontodium inclinatum. *Hedw. Sp. Musc.* 58.

Bryum inclinatum. *Dicks. Crypt. fasc.* 3. 9.
With. 835. *Hull.* 264.

Angl. Curve-fruited Grimmia.

In paludibus turfosis Scotiæ. *Dicks.*

Perennis. Fruct. fert æstate.

(Caules breves, cæspitosi, basi ramosi, supernè simplices, foliosi. Folia lutescentia, mox fusca, setacea, patula,
T 3 basi

basi dilatata, caulem vaginantia. Pedicellus terminalis, erectus, ferè uncialis, rectiusculus, purpureus. Capsula ovata, obliquè arcuata, subcernua, lævis, fusca, ore truncato, parùm dilatato. Peristomii dentes sedecim, lanceolati, rubri, perforati, æquidistantes, nec per paria approximati. Operculum conicum, breye, pallidum.

heteromalla. 16. GRIMMIA foliis setaceis secundis basi dilatatis, capsulâ ovato-cylindraceâ erectâ, operculo conico.

Grimmia heteromalla. Roth. Germ. v. 3, 145.
Weisia heteromalla. Hedw. Sp. Musc. 71. Crypt. v. 1. 22. t. 8.

Afzelia heteromalla. Ehrb. Crypt. 173.
Bryum Weisia. Dicks. Crypt. fasc. 2. 5. Dr. Pl. 94. With. 827. Hull. 261. Hoffm. Germ. v. 2. 33.

Angl. Curve-leaved Grimmia.

In sylvis et arenosis. Dicks.

On heaths and commons about Forfar, but not frequent. Mr. G. Donn. In Ireland. Dr. Scott.

Perennis? Fruct. fert Julio.

Caules cæspitosi, breves, subsimplices, rigidi, atro-fusci, foliosi. Folia luteo-viridia, arcuato-secunda, setacea, uninervia, basi latissimâ. Pedicellus terminalis, rectiusculus, ferè uncialis, purpureus; apice pallidus. Capsula erecta, ovato-cylindracea, lævis, fusca, ore subcoarctato. Operculum conicum, acutum, rectiusculum, pallidum, basi rubrum; in iconc Hedwigianâ nimis obtusum. Peristomii dentes sedecim, æquidistantes, quâ notâ a sequente differt.

homomalla. 17. GRIMMIA foliis setaceis secundis basi dilatatis, capsulâ ovatâ erectâ, operculo conico, peristomii dentibus geminatis.

Didy-

Didymodon homomallum. *Hedw. Sp. Musc.*

105. t. 23. f. 1—7.

Bryum pallidum. *With. 818*, exclusis synonymis. *Hull. 258.*

Angl. Pale Grimmia.

In turfosis et nemorosis subalpinis.

In Garn Dingle, and about peat holes near Lyn Aled, North Wales. *Mr. Griffith.* At Bedd-gelart, at the foot of Snowdon. *Mr. D. Turner.*

Perennis? Fruct. fert Junio, Julio.

Præcedenti tam arctè affinis ut vix nisi peristomii dentibus per paria approximatis, sive geminatis, distinguenda. Caulis fere nullus. Folia pallidè viridia, forma prioris, sed multò magis arcuato-secunda. Pedicellus purpureus, apice pallidè virens. Capsula omnino ovata; erecta. Operculum breve, conicum; acutum; paululùm incurvatum, pallidum, basi rubrum. Peristomii dentes sedecim, per paria approximati, at basi distincti.

18. GRIMMIA. foliis lanceolatis basi dilatatis, *nigrita*. capsulâ obovatâ cernuâ, operculo hemisphærico umbonato.

Grimmia nigrita. *Roth. Germ. v. 3. 146.*

Weisia nigrita. *Hedw. Sp. Musc. 72. Crypt. v. 3. 97. t. 39.*

Bryum nigritum. *Dicks. Crypt. fasc. 3. 9. With. 840. Hull. 266.*

Angl. Black-fruited Grimmia.

In ericetis turfosis Scotiæ: *Dicks.*

Perennis. Fl. . . .

Caulès cæspitosi, subdivisi, foliosi, ferrugineo-tomentosi.

Folia lætè viridia, lanceolata, acuminata, integrifima, subrevoluta, basi dilatata, siccitatem incurva. Pedicellus capillaris, ruber, paululùm flexuosus, vix uncialis. Capsula parva, obovata, arcuato-cernua, demùm atrata,

fulcata. Operculum convexum, mucrone brevissimo, obtusissimo.

Forsteri. 19. GRIMMIA ? foliis ovatis concavis muticis, capsulâ ovato-oblongâ erectâ, operculo subulato arcuato.

Bryum Forsteri. *Dicks. Crypt. fasc. 3. 4. t. 7. f. 8.*

Angl. Forsterian Grimmia.

In arborum truncis.

On the trunks of felled trees at Walthamstow.

Mr. T. F. Forster junior.

Perennis? Fl.

Præcedente vix major. Caules brevissimi, cæspitosi. Folia saturatè viridia, ovata, concava, integerrima, acutiuscula, mutica. Pedicellus flavescent, subtortus. Capsula ovato-oblonga, ereta. Operculum subulatum, arcuatum, longitudine ferè capsulæ. Peristomium non vidi.

conosto- 20. GRIMMIA foliis lanceolatis ferrulatis aristatis ma. quinquesariâm imbricatis, capsulâ subrotundâ, peristomio connivente.

Grimmia conostoma. *Engl. Bot. t. 1135.*

- Bryum tetragonum. *Dicks. Crypt. fasc. 2. 8. t. 4. f. 9. With. 835. Hull. 264.*

Conostomum arcticum. *Swartz. in epistolis.*

Angl. Cone-mouthed Grimmia.

In alpibus Scoticis.

On Ben Lomond. *Dicks.* On the summits of the Corum-gorum mountains of Inverness-shire; also on Ben Lawers and Loch-ne-gare. *Mr. G. Donn.*

Perennis. Fruct. fert Julio et Augusto.

Caules cæspitosi, unciales, vel biunciales, ramosi, fastigiati,

giati, e foliis undique quinquefariam imbricatis, concinnè pentagoni. Folia adpressa, lætè viridia, lanceolata, carinata, ferrulata, brevius aristata seu mucronulata. Pedicellus uncialis, subflexuosus, ruber. Capsula erectiuscula, subrotunda, parùm obliqua, demùm longitudinaliter corrugata, luteo-ferruginea. Peristomium rubrum, dentibus apice subulatis, conniventibus. Operculum conicum, incurvum, capsulâ duplò brevius.

Folia, optimè monente Swartzio, quinquefariam nec quadrifariam, imbricata sunt.

21. **GRIMMIA** foliis lanceolatis integerrimis acu-nuda.
tis muticis, capsulâ subrotundâ inclinatâ lævi,
caule nullo.

Bryum nudum. *Dicks. Crypt. fasc. 4. 7. t. 10.*
f. 15.

Angl. Naked Grimmia.

In humidis argillosis.

On the banks of ditches in a clay soil near Manchester; Mr. G. Caley. *Dicks.* On clay banks by the side of the Tay near Perth. *Mr. G. Donn.*

Annua? Fl.

Caulis omnino nullus. Folia pauca, radicalia, erecta, dilutè ac tristè viridia, pellucida, reticulata, lanceolata, acuta, integerima. Pedicellus solitarius, erectus, filiformis, æqualis, ruber, ultra unciam longus. Capsula cernua, ferè globosa, truncata, lævis, pallide fusca, ore rubro. Peristomii dentes rubri, erectiusculi, acuti. Operculum subulatum, arcuatum, breve.

22. **GRIMMIA** foliis spatulatis ferrulatis, cap-splach-
sulâ globosâ lævi, pedicello apice incrassato. *noides.*
Splachnum longicollum. *Dicks. fasc. 4. 4. t. 10.*

f. 6.

Weisia splachnoides. *Swartz. in epist.*
Angl. Splachnoid Grimmia.

In

In alpibus Scoticis. *Dicks.*

Perennis? Fl. . . .

Caules erecti, subsimplices, sesquiunciales, foliosi. Folia lætè ac dilutè viridia, reticulata, imbricato-patentia, oblongo-spatulata, acuta, serrulata. Pedicellus uncialis vel sesquiuncialis, flexuosus, purpureus, apice pallidior, et valdè, apophysis instar, sub capsula incrassatus. Capsula erectiuscula, globosa, pallidè ferruginea, lævissima, ore patentiusculo, rubro. Peristomii dentes sedecim, æquidistantes, breves, inflexi, rufi. Operculum convexum, breve, muticum. Calyptre cylindraceo-conica.

Muscus Weisiæ radianti *Hedw. Sp. Musc.* 73, ut videtur, propinquus, habitu *Splachnum* charactere omnino Grimmia.

Donni- 23. GRIMMIA foliis lanceolatis acutis apice pi-
ana. liferis, capsula ellipticâ; peristomio imperforato, operculo recto.

Angl. Bearded Alpine Grimmia.

In alpibus Scoticis rarissimè.

Upon large stones near a water-fall on a mountain in Angusshire, 18 miles north of Forfar. Mr. G. Donn.

Perennis. Fl. . . .

Caules breves, cæspitosi, parùm ramosi. Folia imbricata, lanceolata, acuminata, carinata, integerrima, colore ferè atra, apice pilo cano, erecto, elongato, terminata. Pedicellus pallidus, brevis, erectus, filiformis. Capsula erecta, elliptica, lævis, fusco-virens, demùm nigra, ore haud patulo. Peristomium rubrum, dentibus brevibus, imperforatis. Operculum convexum, mucrone brevi, recto, cylindraceo, obtuso. Grimmia cribrosa Hedwigii discrepat colore minus atro, capsula brevi ferè sessili, ore patentissimo, denique peristomii dentibus elegantè pertusis.

24. GRIMMIA foliis ovato-lanceolatis obtusis *alpicola*.
muticis, capsulâ urceolatâ subsessili, peristomio imperforato, operculo obliquo.

Grimmia alpicola. Swartz. Musc. Suec. 27 et 81.
t. I. f. I. Hedw. Sp. Musc. 77. t. 15. f. 1—5.
Angl. Beardless Alpine *Grimmia*.

In saxis ad rivulos alpinos.

On the mountains of Ireland. Dr. Scott.

Perennis. Fl. . . .

Caules unciales, cæspitosi, subdivisi. Folia atro-viridia, imbricata, ovato-lanceolata, carinata, integerima, subrevoluta, obtusiuscula, mutica, nunquam incurva. Capsulæ subsessiles, solitariæ, erectæ, urceolatæ, læves, ferrugineæ, ore patentissimo, rufo. Peristomii dentes saturatè rubri, demùm patentes. Operculum rubrum, mucrone brevi, arcuato.

Peristomii dentes rariùs foramine uno alterove pertusos invenit Cel. Swartz, qui primus hanc speciem a G. apocarpâ, cui valdè affinis, disjunxit.

25. GRIMMIA foliis lanceolatis acutis muticis *mariti-*
siccitate incurvis, capsulâ urceolatâ, peristomio imperforato, operculo recto brevissimo.

Angl. Sea *Grimmia*.

In rupibus maritimis.

On rocks exposed to spring tides about the ferry at Dundee, also on the coast of Fife. Mr. G. Donn.
In Ireland. Dr. Scott.

Perennis. Capsulas fert Septembri, Octobri.

Habitus præcedentis. Caules densè cæspitosi, ramosi. Folia lanceolata, integerima, apice attenuata, acuta, siccitate incurva, saturatè viridia, minusque quam in præcedente atra; juniora lætè virentia. Capsulæ terminales, vix pedicellatae, erectiusculæ, urceolatæ, læves, fuscae, foliis tectæ, ore dilatato. Peristomii dentes

dentes breves, imperforati, fusi. Operculum capsula latius, eonvexum, muerone brevissimo, obtuso, reeto. A G. plagiopodiâ Hedwigii differt eaule breviore, foliis angustioribus, nunquam piliferis, monente D. Turnero.

rivularis. 26. GRIMMIA foliis ovatis acutiusculis muticis, capsula urceolatâ sessili, operculo recto, caule ramosissimo.

Grimmia rivularis. Bridel?

Angl. River Grimmia.

In rivulis alpinis.

In Ireland. Dr. Scott. In Scotland. Mr. G. Donn.

Perennis. Fl. . . .

Caules biunciales et ultrà, ramosissimi, laxè patentes, rigiduli, nigri, foliosi; infernè demùm denudati, vel omnino glabri, vel e foliorum nervis persistentibus fetosi. Folia erecto-patula, atro-viridia, ovata, integrifima, carinata, acutiuscula, minimè acuminata, omnia mutica. Capsulae solitariæ, terminales, sessiles, foliis circumvallantibus superatæ, ovato-ureolatæ, lœves, fuscæ, ore patulo. Operculum hemisphærium, mucrone brevi, reeto. Peristomii dentes imperforati, incurvi, fusi.

apocarpa. 27. GRIMMIA foliis ovatis acutis; superioribus apice piliferis, capsula ovatâ sessili, caule ramosissimo.

Grimmia apocarpa. Hedw. Sp. Musc. 76. Crypt. v. 1. 104. t. 39. Sibth. 276. Roth. Germ.

v. 3. 138. Engl. Bot. t. 1134. . . .

G. polyodon. Ehrh. Crypt. 212.

Bryum apocarpum. Sp. Pl. 1579, α. Hud. 474, β. With. 809, var. 1. Relh. 424. Abbot. 236. Fl. Dan. t. 480.

Fontinalis

Fontinalis apocarpa. *Hull.* 274.

Hypnum n. 1793. *Hall. Hist.* v. 3. 40.

Sphagnum caulerum et ramosum saxatile hirsutum virescens, capitulis obscurè rubris. *Raii Syn.* 104.

S. subhirsutum obscurè virens, capsulis rubellis.
Dill. Musc. 245. t. 32. f. 4.

Muscus apocarpos hirsutus, saxis adnascens, capitulis obscurè rubris. *Vaill. Par.* 129.
t. 27. f. 15. *Herb. Vaill.*

Angl. Sessile Bearded Grimmia.

In rupibus et arborum truncis.

Pérennis. Fructum fert vere.

Caules altitudine varii, unciales vel biunciales, laxè cæspitosi, ramosissimi, undique foliosi. Folia fusco-viridia, ovata, integerrima, carinata, acuta; superiora, quandoque omnia, apice pilifera. Capsulæ terminales, solitariæ, sessiles, erectæ, ovatæ, breves, dilutè fuscæ, lèves, ore haud patulo. Peristomii dentes erecti, pulcherrimè rubri, acuminati, imperforati. Operculum hemisphæricum, rubrum, mucrone brevi, acuto, saepius incurvato. Calyptra magnitudine tantùm operculi.

448. DICRANUM. *Hedw. Fund.* v. 2. 91. t. 8.
f. 41, 42. *Schreb. Gen.* 759.

Fissidens. *Hedw. Fund.* v. 2. 91.

Caps. oblonga. *Perist.* dentibus sedecim, complanatis, subinflexis, semibifidis.

* *Foliis secundis vel patentibus.*

1. DICRANUM caule ramoso, foliis secundis scopo-ovato-lanceolatis acuminatis carinatis falcatis, rium. pedicellis vaginatis solitariis.

Dicranum

- Dicranum scoparium.* *Hedw.* *Sp. Musc.* 126.
Swartz. Musc. Suec. 34. *Sibth.* 281. *Roth.*
Germ. v. 3. 158. *Ehrb. Crypt.* 104.
Bryum scoparium. *Sp. Pl.* 1582. *Huds.* 483.
Hull. 261. *Abbot.* 241. *Curt. Lond. fasc.* 1.
t. 69. *Engl. Bot. t.* 354. *Dicks. H. Sicc. fasc.* 11. 17.
B. reclinatum, foliis falcatis, scoparum effigie.
Dill. Musc. 357. *t. 46.* *f. 16.* A. B. C.
B. erectis capitulis angustifolium, caule reclinato.
Raii Syn. 95.
Hypnum n. 1777. *Hall. Helvet.* *v. 3.* 36.
Mnium scoparium. *With.* 799. *Relb.* 418.
Angl. Broom Fork-moss.

In ericetis et sylvis vulgaris.

Perennis. Fruct. fert hyemē et verē, etiam aestate,
 Caules 2—4-unciales, cæspitosi, undique foliosi, dense
 villosi. Folia luteo-viridia, nitida, secunda, falcata,
 lanceolata seu ovato-lanceolata, coiceaviuscula, cari-
 nata, integerima, uninervia, superficie æqualia; nec
 undulata, apice acuminato-setacea. Pedicelli solitarii,
 erecti, sesquiunciales, basi foliis pluribus vaginantibus,
 brevissimè mucronulatis, vestiti. Capsula erecta, ru-
 fescens, cylindraceo-clavata, gracilis, paululum ar-
 cuata, basi attenuata, sub ore constricta, levigata nec
 sulcata. Peristomium fuscum, parvum, deciduum.
 Operculum setaceum, rectum, longitudine capsulae.
 Calyptra basi coarctata.

majus. 2. **DICRANUM** caule ramoso, foliis secundis
 ovatis acuminatis falcatis levibus, pedicellis
 vaginatis aggregatis.

Bryum reclinatum, foliis falcatis scoparum effi-
 gie, setis pluribus. *Dill. Musc.* 358. *t. 46.*
f. 16. D.

Angl. Greater Clustered Fork-moss.

In rupibus et montosis sylvaticis.

Sent by Dr. Richardson. *Dilleniuss.* On marine rocks near Bangor. *Mr. D. Turner.*

Perennis. Fruct. fert Julio.

Priore majus. Folia omnia prorsus secunda, falcata, basi ovata, nervo tenuiore, apice longius acuminato-setacea. Pedicelli bini vel terni ex eadem vaginâ. Capsula brevior, crassior, magisque quam in præcedente arcuata, saturatè fusca. Operculum subulatum, longitudine capsulæ.

3. **DICRANUM** caule ramoso, foliis subsecundis *undulato-*
ovato-lanceolatis acuminatis undulatis apice tuni-
serrulatis, pedicellis vaginatis aggregatis.

Dicranum undulatum. Schrad. *Spicil.* 59. Ehrb.
Crypt. 271. *Rotb. Germ.* v. 3. 167.

D. polysetum. Swartz. *Musc. Suec.* 34. et 87.
t. 3. f. 5.

Bryum rugosum. Hoffm. *Germ.* v. 2. 39.

Museus capillaceus, foliis pallidis, pediculis pluribus e communi thecâ egredientibus. Buxb.
Cent. 2. 8. t. 4. f. 1.

Angl. Waved-leaved Clustered Fork-moss.

In sylvis et uliginosis alpinis.

In Scotland. *Mr. Dickson.* In fir woods near Forfar. *Mr. G. Donn.*

Perennis. Fruct. fert æstate.

Magnitudine cum penultimâ specie convenit, discrepat verò foliis undique frè patentibus, (vix secundis,) ovatis, plus minus transversè corrugatis, brevius acuminatis, apice serrulatis. Pedicelli duo vel plures, quandoque octo, e communi vaginâ. Capsulæ cylindraceæ, arcuato-cernuæ, læves, minores quam in penultimâ. Operculum setaceum, capsulâ ferè longius.

Hanc

Hanc speciem cum D. scopario confuderunt Linnæus et Hedwigius, etiam, ut e speciminibus ex Helvetiâ missis patet, Hallerus. Synonymon Buxbaumianum male ad D. majus retulit Dillenius, et inde foliorum descriptionem minus aptè carpit.

- fusces-* 4. **DICRANUM** caule ramoſo, foliis secundis
cens. subulatis carinatis, pedicellis subvaginatis fo-
litariis, capsulâ obovatâ arcuatâ fulcatâ.
Angl. Brown Fork-moss.

In nemoribus et ericetis.

Near Edinburgh.

Pennis. Fruct. fert æstate.

Priori duplò minus. Folia fuscescentia, subulata, basi tantum dilatata, nervo validiore carinata. Pedicelli solitarii, erecti, foliis vaginantibus paucioribus, stric- tioribus, longiusque aristatis. Capsula brevis, obovata, arcuato-cernua, inæqualiter sulcata, fusca, ore patulo, peristomio brevi, concolore. Operculum longitudine capsulae, fetaceum, pallidum. Calyptra fusca.

Hanc speciem sub nomine D. falcati ex amicissimo Schradero olim accepi, at frustrâ in Spicilegio suo quæsivi.

- hetero-* 5. **DICRANUM** caule subsimplici, foliis capilla-
mallum. ribus secundis uninervibus basi dilatatis, cap-
sulâ obovatâ gibbosâ.

Dicranum heteromallum. *Hedw. Sp. Musc.* 128.
Crypt. v. 1. 68. t. 26. *Swartz. Musc. Suec.*

37. *Roth. Germ.* v. 3. 160. *Sibth.* 282.

Bryum heteromallum. *Sp. Pl.* 1583. *Huds.* 484:
Hull. 263. *Abbot.* 240. *Dicks. H. Sicc. fasc.*

12. 23. *Dill. Musc.* 375. t. 47. f. 37.

B. trichoides, reclinatis caulinulis, capitulis erec-
tis acutis. *Raii Syn.* 96.

Mnium heteromallum. *With.* 801. *Relb.* 418.

Muscus

Muscus capillaceus, foliis unam partem spectantibus. *Vaill. Paris.* 132. t. 27. f. 7...
Angl. Silky Heath Fork-moss.

In ericetis glareosis vel umbrosis fabulosis.

Annua. Fl. autumno et hyeme, fruct. fert aestate.
Magnitudine variat. Caules plerumque simplices, semiunciales. Folia secunda, falcata, elongata, laxa et mollia, pulchrè viridia, basi sensim dilatata, nervo validiusculo carinata. Pedicelli capillares, torti, flavi. Capsula nitida, spadicea, gibbosa, indè subcernua. Operculum setaceum, longitudine et colore capsulæ, incurvum. Peristomium rubrum, formosum, magnum, laciniis capillaribus, albidis.

6. **DICRANUM** caule subramoso, foliis capilla-*interrup-*
ribus uninervibus interruptè secundis basi di-tum.
latatis siccitate crispis, capsulâ ovatâ.

Dicranum interruptum. *Hedw. Sp. Musc.* 129.
t. 29. f. 8—12.

Bryum heteromallum β. *Huds.* 484. *Hull.* 263.
B. pilosum, interruptè falcatum. *Dill. Musc.*
376. t. 47. f. 38.

Mnium heteromallum, var. 2. *With.* 801.

Angl. Interrupt Fork-moss.

In montibus.

On Snowdon. *Dill. and Mr. Griffith*. Near New-
 castle. *Mr. Winch.*

Perennis? Fl. vere.

Differt a præcedente foliis per intervalla majoribus et minoribus, indè fasciculatim interruptis, inferioribus distichis, omnibus mox suprà basin ampliatam contractis, arefactione tortilibus; capsulis ovato-cylindraceis, erctis; peristomio minore.

Exemplaria e Cel. Hedwigio filio ad amicissimum D.

Dawsonum Turner missa a nostris paululum discrepant,

VOL. III.

U

foliis

foliis haud tortilibus et vix interruptis ; fructu reverâ conveniunt.

flagel- 7. **DICRANUM** caule ramoso, foliis setaceis se-
lare. **DICRANUM** caule ramoso, foliis setaceis se-
cundis subenervibus basi dilatatis, capsulâ
cylindraceâ sulcatâ erectâ.

Dicranum flagellare. *Hedw.* *Sp. Musc.* 130.
Crypt. v. 3. t. 1. *Schrad. Spicil.* 59. *Roth.*
Germ. v. 3. 165, excluso syn. *Dillenii.*
Bryum flagellare. *Dicks. Crypt. fasc.* 3. 6. *With.*
831. *Hull.* 262. *Hoffm. Germ.* v. 2. 37.
Angl. Slender-branched Fork-moss.

In rupibus et ericetis montosis.

On Ben Nevis. *Dicks.* On Cromford moor near
Matlock.

Perennis. Fl. . . .

Caules cæspitosi, 1—3-unciales, ramosi, foliosi, dense
tomentosi, tomento ferrugineo vel sanguineo, ramulis
subinde squamosis aphyllis. Folia plus minus secunda,
subulata, lutescentia, nitidiuscula, basi dilatata, con-
cava, nervo obsoleto, vel prorsus nullo, apice tenuis-
sima, subflexuosa. Pedicelli unciales, stricti, basi
foliolis setaceo-acuminatis vaginati. Capsula erecta,
stricta, cylindracea, sulcata, rufo-fusca.

subula- 8. **DICRANUM** caule simplici, foliis subulato-
capillaribus secundis basi dilatatis, capsulâ
ovatâ sulcatâ cernuâ.

Dicranum subulatum. *Hedw.* *Sp. Musc.* 123.
t. 34. f. 1—5.

Angl. Awl-leaved Fork-moss.

In subhumidis.

At Rathmines near Dublin. *Dr. W. Stokes.*

Annua. Fruct. fert autumno.

Caules

Caules erecti, simplices, stricti, foliosi, semiunciales, nigricantes. Folia luteo-viridia, areuato-secunda, tenuissima, acuta; basi dilatata, uninervia. Pedicellus capillaris, ruber, tortilis, haud uncialis. Capsula ovato-cylindracea, arcuato-cernua, demum obsoletè sulcata. Peristomium rubrum. Operculum exemplaribus deest.

97. DICRANUM caule simplici, foliis capillaribus *crispum*.
laxis enervibus basi dilatatis, capsulâ ovatâ
sulcatâ erectâ.

Dicranum crispum. Hedw. Sp. Musc. 132.

Crypt. v. 2. 91. t. 33. Swartz. Musc. Suec.

37. Engl. Bot. t. 1151.

Bryum vaginale. Dicks. Crypt. fasc. 3. 8. With.

827. Hull. 261.

Angl. Curled Fork-moss.

In paludosis Scotiæ. *Dicks.*

In Ireland on the moist banks of rivers. *Mr. Templeton* and *Dr. Scott.*

Perennis. Fruct. fert Julio.

Caules cæspitosi, breves, simplices, foliosi. Folia dilutæ ac flavo-viridia, undique ferrè patentia, laxa, capillaria, undulato-crispa, basi dilatata, amplexicaulia, vaginantia, enervia. Pedicellus capillaris, ruber, apice tortilis, vix uncialis. Capsula erecta, ovata, brevis, rufofuscâ, demum sulcata. Operculum setaceum, obliquum, rubrum, capsulâ plerumque longius. Peristomium rubrum.

98. DICRANUM caule subsimplici, foliis setaccis *uncinata*-
carinatis enervibus secundis, capsulâ cernuâ, *tum.*
apophysi hinc gibbosâ.

Bryum uncinatum. Dicks. Crypt. fasc. 4. 11.
t. 11. f. 8.

Angl. Hooked Fork-moss.

In terrâ nudâ turfosâ alpium Scoticarum. *Dicks.*

Perennis? Fl.

Magnitudine D. crispi. “ Caules quandoque subpollicares.” *Dicks.* Folia lætè viridia, secunda, latiora quam in præcedente, apice tamen gracillima; inferiora nigricantia; omnia carinata, nervo destituta, basi sensim dilatata. Pedicellus terminalis, flavescens, demum flexuosus. Capsula obovato-cylindracea, incurva, lævis, fusca, apophysi parvæ, hinc gibbosæ, insidens. Operculum subulatum, inflexum, capsulâ longius. Peristomium, quod non ipse vidi, dentibus bifidis, inflexis, descripsit amicissimus Dicksonus.

falcatum.

II. DICRANUM caule subsimplici, foliis setaceis secundis uninervibus, capsulâ obovatâ cernuâ basi strumulosâ.

Dicranum falcatum. *Hedw. Sp. Musc.* 150. t. 32. f. 1—7.

Bryum longifolium. *Dicks. Crypt. fasc.* 3. 7. *With.* 834. *Hull.* 264.

Angl. Sickle-leaved Fork-moss.

Ad rivulorum ripas in alpibus Scoticis.

On Ben Nevis. *Dicks.*

Perennis. Fruct. fert æstate.

Caules unciales vel sesquiunciales, cæspitosi, erectiusculi. Folia secunda, plus minus falcata, luteo-viridia, uninervia; basi lanceolata, concava; apice capillaria. Pedicellus semiuncialis, lutescens, apice tortus. Capsula subcernua, parva, obovata, lævis, castanea, basi hinc gibbosa vel strumosa. Peristomium majuscum, rubrum. Operculum subulatum, obliquum, capsulâ brevius.

Specimina Dicksoniana cum Hedwigianis contuli. D. longifolium verum differt magnitudine majori, foliis longioribus, laxis, nervo orbatis, capsulis cylindraceis, gracilioribus, sulcatis, fusco-nigricantibus, basi æquilibus,

12. DICRANUM caule ramoso, foliis setaccis *fulvelum*.
secundis uninervibus, capsulâ turbinatâ sulcatâ erectâ.

Bryum fulvellum. *Dicks. Crypt. fasc. 4.* 10.
t. 11. f. 1.

β. *B. immersum.* *Dicks. Crypt. fasc. 4.* 15, ex-
clusis synonymis.

Angl. Tawny Fork-moss.

In rupibus alpinis Scotiæ.

On Ben More. *Dicks.*

Perennis. Fl. . . .

Facies omnino Grimmiae acutæ, sed foliis longius acuminatis, capsulâ demum sulcatâ, denique peristomio Dicrani discrepat. Caulés sesquiunciales, ramosi, crecti, subflexuosi. Folia falcato-secunda, fulvo-viridia, basi dilatata, apice capillaria. Pedicellus terminalis, brevis, rectus. Capsula erecta, parva, castanea, turbinata, ore dilatata, haud ventricosa, demum profundè sulcata. Peristomii dentes, monente amicissimo D. Turnero, bifidi.

13. DICRANUM caule simplici sculo, foliis ca- *varium.*
pillaribus subsecundis apice flexuosis, capsulâ
urceolatâ subcernuâ basi attenuatâ.

Dicranum varium. *Hedw. Sp. Musc.* 133. *Crypt.*
v. 2. 93. t. 34. *Roth. Germ.* v. 3. 170. *Engl.*
Bot. t. 1215.

D. simplex. *Sibth.* 282.

Bryum simplex. *Sp. Pl.* 1587. *Huds.* 486.
With. 800. *Hull.* 254. *Abbot.* 239. *Dicks.*
H. Sicc. *fasc. 14.* 20. et 17. 21.

B. rubrum. *Huds.* *ed. 1.* 413.

B. trichodes, capsulis rubris cernuis. *Dill. Musc.*
390. *t. 50. f. 59.*

Mnium simplex. *Rehb.* 418.

Angl. Variable Fork-moss.

In gramineis arenosis, minus frequens.

Annua. Fl. hyeme; fruct. fert vere et aestate.

Prioribus longè minus. Caulcs plerumque simplices, quandoque varie subramosi. Folia capillaria, magis minusve secunda, apice laxiuscula nec rigidula, basi dilatata, integerrima, late viridia, oculo armato obsoletè reticulata. Pedicelli terminales, fulvi, nitidi. Capsula clavata, ore patulo, incurvata et demum cernua. Operculum incurvatum, subulatum, basi incastratum, capsulâ vix brevius. Peristomium magnum, rubrum.

rufescens. 14. DICRANUM caule simplici, foliis setaceis laxis secundis reticulatis subserratis, capsulâ urceolatâ erectâ.

Dicranum rufescens. Engl. Bot. t. 1216.

Bryum rufescens. Dicks. Crypt. fasc. 3. 6. t. 8, f. 1. With. 818. Hull. 258.

Angl. Red Fork-moss.

In lutosis vel arenosis humidis Scotiæ.

Discovered in 1782 on a moist bank by the side of the rivulet at the Hermitage a mile south of Edinburgh.

It has since been observed by Mr. G. Donn in wet stubble fields, and in fields sown with grass, near Forfar.

Annua. Fl. autumno.

Species a praecedente distinctissima, capsulis semper erectis, foliis, microscopio subjectis, longè evidenter reticulatis, et (è reticulis marginalibus concatenatis) quasi ferratis. Caulis simplicissimus. Folia omnia secunda, laxè patentia; superiora majora. Pedicelli terminales, lutescentes, minus nitidi. Capsula subcylindracea, erecta, ore patulo. Operculum longitudine capsulæ, rufum, arcuatum.

15. DICRANUM caule simplici, foliis lanceo-*callisto-*
lato-subulatis erectis, capsulâ ovato-urceolatâ *mum,*
incurvâ.

Bryum callistomum. *Dicks. Crypt. fasc. 3. 5.*

t. 7. f. 10. With. 818. Hull. 258.

Angl. Urn-shaped Fork-moss.

In sterilibus vel argillosis alpinis.

On barren rocks in the Highlands of Scotland.

Mr. Dickson. In Anglesea. *Rev. H. Davies.*

Perennis. Fl. vere.

Caules cæspitosi, erecti, foliosi, simplices, sæpiùs vix
lineam longi, at nonnunquam semiunciales evadunt.
Folia lutescentia, numerosa, erecto-patula, acutè ca-
rinata, angusta, integerrima; basi dilatata; apice bre-
viùs mucronata. Pedicelli semiunciales, erecti, rubri,
nitidi. Capsula erectiuscula, ovata, brevis, turgidula,
fusca, ex ore dilatato urceolata; demùm incurvato-
cernua. Calyptra subulata, longitudine capsulæ cum
operculo, decidua. Operculum longitudine capsulæ,
rufum, basi conicum, apice subulatum, curvatum.
Peristomium magnum, formosum, longitudine dimidii
capsulæ, erecto-patulum, rubrum, transversè striatum,
apice albidum.

16. DICRANUM caule erecto demùm ramoso, *rigidu-*
foliis subulatis aeuminatis: rameis secundis, *lam.*
capsulâ urceolatâ incurvâ.

Dicranum rigidulum. *Swartz. Musc. Suec. 38*
et 89. t. 3. f. 7. Hedw. Sp. Musc. 134. t. 32.
f. 8—12.

Angl. Rigid-leaved Fork-moss.

In locis argillosis.

In North Wales. *Rev. H. Davies.* Near the cir-
cular branch of the grand canal, Dublin, *Dr.*
W. Stokes.

Perennis. Fl. . . .

Caules cæspitosi, erecti, foliosi, demum ramosi, ramis erectis, pedicellum superantibus. Folia luteo-viridia; caulina subulata, basi dilatata, carinata, integerrinia, subsecunda, rigidula; ramea falcato-secunda, laxa; omnia apice acuminato-setacea. Pedicelli semiunciales, erecti, ex apice caulinum primordialium, solitarii, rubri. Capsula subcernua, ovato-urecolata, incurva, fusca. Operculum ut in præcedente. Peristomium rubrum, dentibus apice pallidis.

Dicrano vario perquam affine.

acicu-
lare.

17. **DICRANUM** caule erectiusculo ramoso, foliis lanceolatis obtusis patulis subsecundis, capsula cylindraceâ, operculo subulato.

Dicranum aciculare. Hedw. Sp. Musc. 135.
Crypt. v. 3. 79. t. 33. Swartz. Musc. Suec. 35. Roth. Germ. v. 3. 164.

Bryum aciculare. Sp. Pl. 1583. Huds. 479. Hull. 261. Lightf. 725. Dicks. H. Sicc. fasc. 6. 20.

B. montanum hemiheterophyllum, operculis acutis. Dill. Musc. 366. t. 46. f. 25.

B. hypnoides erectum montanum, erectis capitulis acutis. Dill. in Raii Syn. 94.

Mnium aciculare. With. 798.

B. *Bryum hypnoides aquaticum*, calyptris nigris acutis. Dill. Musc. 367. t. 46. f. 26. Fl. Dan. t. 1001. f. 1.

B. hypnoides repens aquaticum, erectis capitulis acutis. Raii Syn. 94.

Angl. Dark Mountain Fork-moss.

In rupibus humidis et ad rivulos alpinos.

Perennis. Fl. autumno, capsulas vere perficiens.

Caules laxè cæspitosi, bi- triunciales, ramosi, cylindracei, undique foliosi, folia saturatè viridia, opaca, imbricata,

cata, arcuato-subsecunda, ovato-lanceolata, obtusa, mutica, subrevoluta, concaviuscula, carinata. Pedicelli apicem versus ramorum, pauci, laterales, nigricantes, torti. Capsula erecta, elliptico-cylindracea. Calyptra subulata, basi dentato-lacera. Operculum rectum, subulatum, basi rubrum.

β vix differt nisi caulibus fluitantibus, coloreque atro.

18. DICRANUM caule erectiusculo, ramulis patens, tentibus, foliis linear-lanceolatis acutis strictis, capsula subovata.

Bryum patens. *Dicks. Crypt. fasc. 2. 6. t. 4.
f. 8.*

Angl. Spreading-branched Fork-moss.

Ad rupes inundatas alpinas.

On stones in rivulets on Ben Nevis. *Dicks.* On stones by a lake near the top of Snowdon. *Mr. D. Turner.* Plentiful in Inverness-shire and Angus-shire. *Mr. G. Donn.*

Peregrinis. Fruct. fert Junio, Julio.

Habitus praecedentis, sed laxè patens, caulibus basi denudatis, gracillimis, ramulis divaricato-patentibus, brevibus. Folia erecto-adpressa, ferè linearia, acuta, mutica, subrevoluta, carinata, atro-viridia; ramulorum dilutè ac lètè virentia. Pedicelli breves, demùm flexuosi vel tortuosi. Capsula erecta, ovata, seu ob-ovata, fusca, peristomio rubente. Calyptra basi dentata. Operculum conicum, rectum.

19. DICRANUM caulibus subsimplicibus cæspitelliptosis, foliis subulatis muticis basi ovatis, capsula ellipticâ coarctatâ erectâ.

Bryi n. 29 varietas atra. *Dill. Musc. 371.*

Angl. Elliptical Fork-moss.

In rupibus alpinis.

On the summit of Snowdon. *Mr. D. Turner.* In

Cwm

Cwm Idwell. *Mr. Griffith.* In Ireland. *Mr. Templeton.*

Perennis. Fruct. fert Junio, Julio.

Caules densè cæspitosi, haud unciales, vix ramosi. Folia undique imbricata, fusco-viridia; basi ovata, concava; apice subulata, elongata, mutica; juniora lætè virentia, subarcuato-secunda, conniventia. Pedicellus brevis, erectus, tortus. Capsula erecta, fusca, lævis, exactè elliptica, brevis, ore coarctato. Peristomii dentes arctè inflexi, fusci.

ovale. 20. **DICRANUM** caulis ramosis fastigiatis, foliis lanceolatis piliferis, capsulâ ovatâ erectâ ore patulo.

Dicranum ovalc. *Hedw. Sp. Musc.* 140. *Crypt.* v. 3. 81. t. 34, A. *Roth. Germ.* v. 3. 172.

D. ovatum. *Swartz. Musc. Suec.* 35.

Bryum ovale. *Dicks. Crypt. fasc.* 4. 14.

Angl. Oval Fork-moss.

In rupibus. *Dicks.*

In Ireland. *Dr. Scott* and *Mr. Templeton.*

Perennis. Fruct. fert vere et æstate.

Facies prioris. Caulcs unciales vel biunciales, ramosi, densè cæspitosi, fastigiati. Folia imbricata, fusco-viridia, opaca, lanceolata, carinata, subrevoluta; superiora pilo cano terminata. Pedicelli breves, torti, pallidi, erecti. Capsula erecta, ovata, subrugosa, demum fusca, ore patulo. Peristomium rubrum.

pulvina- 21. **DICRANUM** caulis ramosis fastigiatis, foliis lanceolatis piliferis, capsulâ ellipticâ nutante, operculo subulato.

Dicranum pulvinatum. *Swartz. Musc. Suec.* 32. *Roth. Germ.* v. 3. 173.

Fissidens pulvinatus α. *Hedw. Sp. Musc.* 158. t. 40. f. 1—3.

Encalypta

Encalypta pulvinata. *Sibth.* 278.

Afzelia pulvinata. *Ebrh. Crypt.* 163.

Bryum pulvinatum. *Sp. Pl.* 1586. *Huds.* 487.
With. 840. *Hull.* 267. *Relb.* 422. *Abbot.*
243.

B. orbiculare pulvinatum, hirsutie canescens,
capsulis immersis. *Dill. Musc.* 395. t. 50:
f. 65.

B. trichoides hirsutie canescens, capitulis subro-
tundis reflexis, in perbrevibus pediculis. *Raii*
Syn. 100.

Angl. Grey Cushion Fork-moss.

In tectis, muris, lapidibusque, vulgaris.

Perennis. Fruct. fert vere.

Caules breves, ramosi, in pulvilos magnitudine varios
congesti. Folia atro-viridia, latè lanceolata, concava,
carinata, subrevoluta, pilo cano, flexuoso, proprii lon-
gitudinis, magis quam in præcedente conspicuo, ter-
minata. Pedicelli terminales, arcuati, torti. Capsula
nutans, elliptica, brevis, fusca, angulosa. Operculum
cylindraceo-subulatum, rectum, capsulâ duplo brevius.

22. *DICRANUM* caule ramoso repente, foliis *sciuroi-*
imbricatis lanceolatis sulcatis muticis, capsulâ des.
cylindraceâ, operculo conico.

Dicranum sciuroides. *Swartz. Musc. Suec.* 32.
Sibth. 280. *Roth. Germ.* v. 3. 185.

Fissidens sciuroides. *Hedw. Sp. Musc.* 161.

Hypnum sciuroides. *Sp. Pl.* 1596. *Huds.* 507.
With. 846. *Hull.* 274. *Relb.* 429. *Abbot.*
251.

H. arboreum sciuroides. *Dill. Musc.* 319. t. 41.
f. 54.

H. trichoides erectum, ramulis recurvis, obscuri
coloris. *Raii Syn.* 83.

Angl.

Angl. Squirrel-tail Fork-moss.

In truncis arborum.

Perennis. Fl. autumno, sed rariūs. Fructum vere perficit.

Caules repentes, ramosissimi, laxè cæspitosi. Rami erecti, exsiccatione incurvi, simplices vel ramosi, cylindracei; undique foliosi. Folia arctè imbricata, saturatè viridia, vel sæpiùs luteo-fuscescentia, ovato-lanceolata, acuta, integerrima, longitudinaliter trifida, at enervia. Pedicelli e perichætiis lateralibus, gracilibus, fulvi, mox fusi, perichætio haud multùm longiores. Capsulæ crecta, cylindraceæ, ore coarctato. Operculum conicum, breve, rectum. Peristomium exile, dentibus ultrà medium fissis.

Foliorum axillæ rariūs, gemmiparæ sunt. *Dill. et D. Crowe.*

Peristomii formâ, dentibus ad basin ferè bipartitis, a Dierano aliquantulùm recedit. Habitu gaudet Pterogonii, neutiquam verò charactere.

glaucum. 23. DICRANUM caule ramoso fastigiato, foliis imbricatis ovato-lanceolatis enervibus albidis, capsulâ ovatâ sulcatâ.

Dieranum glaucum. *Hedw. Sp. Musc.* 135.
Swartz. Musc. Suec. 35. *Roth. Germ. v.* 3.
161. *Sibth. 281.*

Bryum glaucum. *Sp. Pl.* 1582. *Huds.* 479.
Hull. 263. *Abbot.* 239. *Lightf.* 723. *Ehrb.*

Crypt. 4.

B. albidum et *glaucum* fragile majus, foliis erectis, setis brevibus. *Dill. Musc.* 362. t. 46. f. 20.

B. trichoides, erectis capitulis, albidum fragile. *Raii Syn.* 97.

Muscus trichodes montanus albidus fragilis. *Vaill. Paris.* t. 26. f. 13.

Mnium

Mnium glaucum. *With.* 801.
Angl. White Fork-moss.

In ericetis et pascuis montosis.

Perennis. Fl. æstate, sed rariūs.

Caules fastigiati, 3, 4-unciales, cæspites latos formantes.

Folia ferè alba, mollia, spongiosa, opaca, arctè imbricata, ovato-lanceolata, integerrima, planiuscula, obtusiuscula, mucronulata. Pedicelli semiunciales, rectiusculi, fusco-rufi. Capsula ovata, brevis, subarcta, sulcata, nigricans, ore patulo. Peristomium rubrum.

24. *DICRANUM* caule dichotomo, foliis lanceo-purpureis, latis siccitate tortuosis, capsulâ ellipticâ sulcata, operculo conico.

Dicranum purpureum. *Hedw.* *Sp. Musc.* 136.

t. 36. *Swartz. Musc. Suec.* 36. *Roth. Germ.*

v. 3. 156. *Sibth.* 281.

Bryum purpureum. *Huds.* 482. *Hull.* 254.

B. tenue stellatum, setis purpureis. *Dill. Musc.* 386. *t.* 49. *f.* 51.

B. perangustis foliis et cauliculis, foliis crebrioribus et circa summitates magis congestis, capitulis erectis e surculis annotinis egreditibus, pediculis purpureis. *Raii. Syn.* 99.

Mnium purpureum. *Sp. Pl.* 1575. *With.* 802.

Relb. 417. *Abbot.* 233.

3. *Bryum tenue.* *Dicks. Crypt. fast.* 3. 8. *With.* 828. *Hull.* 262, nec *Weberi.*

B. purpureum β . *Huds.* 482, exclusis synonymis.

Angl. Purple-stalked Fork-moss. Redshank moss.

In muris, ericetis et glareosis copiosè.

Annua. Fruct. fert vère.

Caules cæspitosi, semiunciales, dichotomi, foliosi. Folia lætè viridia, imbricato-patula, lanceolata sive ovato-lanceolata, acuminata, carinata, uninervia, subrevoluta, arefactione incurva et tortuosa. Pedicelli plerumque e dichotomiâ caulis, solitarii, numerosi, unciales, amœnè purpurei, nitidissimi, basi foliis pluribus vaginati, quorum intimum, optimè notante Hedwigio, enerve est. Capsula erecta, elliptica, rutilans, demùm profundè fulcata, subarcuata, basi obsoletè quadrangula et constricta. Operculum conicum, obtusum, oblique, purpureum, capsulâ quadruplò brevius. Peristomium rubrum, dentibus ultrà medium bifidis.

biparti- 25. **DICRANUM** caule subdichotomo, foliis ovatis revolutis, pedicellis subgeminis, capsulâ ellipticâ striatâ, operculo conico.

Dicranum bipartitum. Roth. Germ. v. 3. 158.

Bryum bipartitum. Dicks. Crypt. fasc. 2. 7.
With. 835. Hull. 265. Hoffm. Germ. v. 2. 36.

B. parvum, surculis et setis geminatis. Dill.
Musc. 385. t. 49. f. 50.

B. parvum trichoides ramosum, erectis capitulis subfuscis, in pediculis obscurè rubris. Dill.
in Raii Syn. 96.

Angl. Twin-stalked Fork-moss.

Ad sepes, in hortis murisq[ue]. Dill. Dicks.

Annua. Fruct. fert primo vere.

Præcedente minus et humilius. Folia latiora, ovata, revoluta, breviùs acuminata, arefactione minùs incurva. Pedicelli pallidiores, sæpè geminati. Capsula striata, nec profundè fulcata, neque basi quadrangula. Operculum ut in priore, vel paulò angustius. Peristomium pallidum.

strictum. 26. **DICRANUM** caule dichotomo, foliis ovatis apice subulatis, capsulâ cylindraccâ sulcatâ inclinatâ, operculo conico.

Dicranum

- Dicranum purpureum β . *Roth. Germ.* v. 3. 158.
 D. intermedium. *Hedw. Sp. Musc.* 138. t. 31.
 f. 1—6?
 Bryum strictum. *Hoffm. Germ.* v. 2. 36. *Dicks.*
Crypt. fasc. 4. 13.
 B. purpureum γ . *Huds.* 483.
 B. n. 1818. *Hall. Hist.* v. 3. 46.
 B. polytrichodes palustre, setis longioribus rubris
 sericeis. *Dill. Musc.* 387. t. 49. f. 52.
 B. peranguſtis crebrioribus foliis, capitulis erec-
 tis, longiusculis pediculis e ſurculis annotinis
 innascentibus. *Dill. in Raii Syn.* 99.
 Mnium purpureum, var. 2. *With.* 802.
Angl. Tall Shining-stalked Fork-moss.

In paludibus montosis.

On Emmott moor, Lancashire. *Richardson.* Shob-
 don marsh, Herefordshire. *L. Brown.* Near
 Llyn Aled lake, North Wales. *Mr. Griffith.*
 Near Newcastle. *Mr. Winch.*

Annua? Fruct. fert Junio.

A proximis præcedentibus distinguitur caulis altioribus;
 foliis basi ovatis, apice longius acuminatis, ſubulatis,
 ſiccitate minus tortis; pedicellis biuncialibus, tenuiſ-
 simis, nitidiflumis, ſæpè geminis aut ternis.

Cum D. intermedii apud Hedwigium descriptione et
 icone, etiam loco, fat bene convenit, at horum ex-
 emplaria nondum contuli.

27. DICRANUM caule simplici, foliis capillari-*pusil-*
 bus enervibus erectiusculis, capsula ovata cer-*lum.*
 nuâ, apophysi hinc gibbosâ.

Dicranum pusillum. *Hedw. Sp. Musc.* 139.
Crypt. v. 2. 80. t. 29, B. *Swartz. Musc.*
Suec. 38. *Roth. Germ.* v. 3. 168. *Sibth.* 282.

Bryum

Bryum parvulum. *Dicks.* *Crypt. fasc.* 3. 7.
With. 832. *Hull.* 263. *Hoffm. Germ.* v. 2.

34.

Angl. Dwarf Fork-moss.

In pastures.

In pastures in Scotland. *Dicks.* On Shotover hill, Oxfordshire. *Sibth.* At Bradwell, Suffolk. *Mr. D. Turner.*

Perennis. Fruct. fert Februario, Martio.

Caules simplices, breves, cæspitosi. Folia undique erecto-patentia, pallidè viridia, demùm fuscescentia, angustissima, concava, vix carinata, enervia, basi dilatata. Pedicelli luteoli, nitidi, gracillimi, basi vaginati, demùm atrati. Capsula ovata, retusa, arcuato-cernua, dilutè fusca, lævis, apophysi rubrâ, exiguâ, hinc gibbosâ. Operculi rostrum setaceum, inflexum, longitudine capsulæ. Peristomium rubrum.

cervicu- 28. **DICRANUM** caule simplici, foliis capillari-latum. bus enervibus patentibus laxis, capsulâ ventricosâ erectiusculâ, apophysi hinc gibbosâ.

Dicranum cerviculatum. *Hedw. Sp. Musc.* 149.
Crypt. v. 3. 89. t. 37 (nec 36), A. *Swartz.*
Musc. Suec. 36.

Bryum cerviculatum. *Dicks.* *Crypt. fasc.* 3. 7.
With. 813. *Hull.* 257. *Abbot.* 237.

Angl. Red-necked Fork-moss.

In ericetis turfosis.

On the dry turfey parts of Hampstead heath near London. *Dicks.* At Ampthill, Bedfordshire. *Abbot.* In Anglesea. *Rev. H. Davies.*

Perennis. Fruct. fert Julio.

Præcedente duplò majus, sed arctè affinc, et differentia specifica difficillima est. Malè omnino in *Hedwigii* opere

opere recentissimo longè separantur. Pedicellus utriusque basi vaginatus. Folia in D. cerviculato laxiora, magisque patula; capsula ventricosior, nitida, apophysi sæpè ruberrimâ.

29. DICRANUM caule simplicissimo brevissimo, foliis lanceolatis acuminatis uninervibus, capsulâ sulcatâ basi strumosâ. *Celsii.*

Dicranum Celsii. *Hedw. Sp. Musc.* 149. t. 33.

f. 1—4. *Swartz. Musc. Suec.* 36.

Bryum Celsii. *Sp. Pl.* 1585. *Dicks. Crypt. fasc.* 3. 7. *With.* 832. *Hull.* 263.

B. trichodes acaulon minimum, setis et capsulis oblongis. *Dill. Musc.* 388. t. 49. f. 54.

Angl. Celsian Fork-moss.

In rupibus alpium Scoticarum. *Dicks.*

Annua? Fl. . . .

Caules cæspitosi, brevissimi. Folia arctè congesta, patula, ovato-lanceolata, revoluta, carinata, uninervia, setaceo-acuminata, luteo-viridia; inferiora nigricantia. Pedicellus terminalis, haud uncialis, rubicundus, basi vaginatus; vaginâ strictâ, muticâ. Capsula paululum inclinata, ovato-cylindracea, fusca, nitidiuscula, sulcata. Operculum conicum, breve. Peristomium rufo-fuscum, dentibus linearibus, ultrà medium bipartitis, ferè ut in *Trichostomo*.

30. DICRANUM caule simplicissimo brevissimo, foliis capillaribus basi dilatatis vaginatum, cylindracea lævi. *cylindricum.*

Trichostomum cylindricum. *Hedw. Sp. Musc.*

107. t. 24. f. 7—13.

Bryum paucifolium. *Dicks. Crypt. fasc.* 4. 7. t. 11. f. 3, ex herbario D. Turneri.

Angl. Cylindrical Fork-moss.

In fragmentis laterum in ruderatis.
Near Wetherby, Yorkshire. *Dickson.*

Annua? Fl. . . .

Caules brevissimi, cæspitosi. Folia undique patentia, capillaria, acutiuscula, enervia; basi dilatata, vaginantia. Pedicellus terminalis, erectus, haud uncialis. Capsula erecta, cylindracea, interdùm sub ore contracta, fusca, lœvis. Operculum subulatum, breviusculum. Peristomii dentes ultrà medium bipartiti.

spurium. 31. **DICRANUM** caule ramoso fastigiato, foliis ovatis acuminatis concavis carinatis, capsulâ cylindracea arcuatâ sulcatâ.

Dicranum spurium. *Hedw. Sp. Musc.* 141.
Crypt. v. 2. 82. t. 30. *Swartz. Musc. Suec.* 33. *Roth. Germ. v. 3. 178.*

Bryum spurium. *Hoffm. Germ. v. 2. 38.* *Dicks.*
Crypt. fasc. 4. 13.

Angl. Broad-leaved Fork-moss.

In paludibus turfosis rarissimè.

On Houghton moor and Barnby moor, Yorkshire,
among *Mnium palustre*. *Mr. Teesdale.*

Perennis. Fruct. fert Maio, Junio, at rariūs.

Caules 4- 5-unciales, cæspitosi, ramosi, fastigiati, undique foliosi, radiculo-villosi. Folia undique imbricato-patula, latè ovata, concava, ventricosa, integrifima, acuminata, carinata, uninervia, nitidiuscula, arefactione repanda et tortuosa; superiora dilutè ac pulchre viridia; inferiora fusca vel nigricantia. Pedicelli pauci, solitarii, sesquiunciales, recti, lutei, nitidi, graciles, basi vaginati. Capsula cylindracea, gracilis, arcuato-cernua, dilutè fusca, inæqualiter fulcata. Operculum subulatum. Peristomium rubrum, breviusculum.

Magnitudine cum primis speciebus certat, habituque quodammodo convenit, sed folia non habet secunda.

32. DICRANUM caule ramoso fastigiato, foliis *pellucidi*-
ovato-lanceolatis acuminatis siccitate incur-dum-
vis, capsulâ ovatâ lævi.
Dicranum pellucidum. *Hedw. Sp. Musc.* 142.
Swartz. Musc. Suec. 35. *Roth. Germ.* v. 3.
 176.
D. aquaticum. *Ehrb. Crypt.* 213. *Roth. Germ.*
 v. 3. 177.
Bryum pellucidum. *Sp. Pl.* 1583. *Huds.* 481.
Hull. 264. *Abbot.* 237. *Dicks. H. Sicc. fasc.*
 12. 22.
B. aquaticum. *Hoffm. Germ.* v. 2. 35.
B. palustre pellucidum, capsulis et foliis bre-
vibus recurvis. *Dill. Musc.* 364. t. 46. f. 23.
B. erectis capitulis subrotundis fuscis, foliis mi-
noribus pellucidis rugosis. *Raii Syn.* 96.
Mnium pellucens. *With.* 802.
Muscus polytrichoides elatior, foliis angustis
pellucidis et ferè membranaceis. *Pluk. Phyt.*
 t. 44. f. 7.
β. Muscus polytrichoides angustifolius pellucidus
ramosus. *Pluk. Phyt.* t. 49. f. 1.
Angl. Pellucid Fork-moss.

In umbrosis humidis, aquosis, et palustribus al-
pinis.

Perennis. Fl. vere.

Caules erecti, ferrugineo-tomentosi, graciles, foliosi, unciales vel biunciales, simplices, deniū ramosi. Folia remotiuscula, alterna, ovata, seu ovato-lanceolata, pellucida, carinata, uninervia, acuminata, integerrima; arefactione variè flexa. Pedicelli unciales, subflexuosi, flavescentes. Capsula ovata, brevis, subarcuata, lævis nec fulcata, ore patulo. Operculum longitudine capsulæ, conico-subulatum, incurvum. Peristomium purpurascens, apice flavum.

flaves-
cens. 33. **DICRANUM** caule erecto subramoso, foliis
lineari-lanceolatis siccitate flexuosis, capsula
turbanatâ erectâ lævi.

Bryum flavescentia. *Dicks. Crypt. fasc.* 2. 4. t. 4.
f. 5. *With.* 830. *Hull.* 254.
Angl. Yellowish Fork-moss.

In paludosis alpinis Scotiæ. *Dicks.*

Perennis? Fl. . . .

Caules erectiuseuli, bi- sive tri-unciales, simplices vel ramosi, undique foliosi, læves. Folia linearis lanceolata, acuta, carinata, integerrima, patula, arefactione undulata nec tortilia; basi dilatata; juniora flavescentia. Pedicelli semiunciales, recti. Capsula erecta, turbinata, seu ovato-cylindracea ore dilatata, lævis, haud arcuata, demum atro-fusca.

Diversum, ni fallor, a *Bryo flavescente* Scopolii. Capsula in icona Dicksonianâ nimis aliquantulum elongata videtur.

viridissi- 34. **DICRANUM** caule ramoso fastigiato; foliis
mum. confertissimi lanceolatis reflexo patulis, cap-
sula ovatâ sulcatâ erectâ.

Bryum viridissimum. *Dicks. Crypt. fasc.* 4. 9.
t. 10. f. 18.

Angl. Bright-green Fork-moss.

In pastuis, arborum truncis, et muris Scotiæ, Hi-
berniæ, et Cambrobritanniæ.

In pastures in Scotland. *Mr. Dickson.* Near Kilkullen bridge, Ireland. *Mr. R. Brown.* North Wales. *Rev. H. Davies.*

Perennis. Fl. . . .

Caules vix unciales, cæspitosi, ramosi, fastigiati, den-
sissimè foliosi, inferè ferrugineo-tomentosi. Folia
imbricato-patula, lanceolata, acuta, integerrima, uni-
nervia, subcanaliculata, apice reflexa; arefactione ad-
pressa

pressa et subtortuosa; juniora lætissimè virentia, opaca, nec nitida; omnia demum ferruginco-fusca. Pedicelli semiunciales, contorti, pallide fusci. Capsula erecta, ovato-cylindracea, longitudinaliter sulcata, pallidè fusca, oris coarctata. Operculum conicum, capsulâ brevius. Peristomium exiguum, deciduum.

35. DICRANUM caule ramoso fastigiato, foliis squarro-lanceolatis recurvato-squarrosis, capsulâ ova-sum. tâ inclinatâ lævi.

Dicranum squarrosum. Schrad. *Journ. ann.*

1802. fasc. 1. 68.

Bryum palustre. Dicks. *Crypt. fasc.* 4. 11.

B. pellucidum G. Sp. Pl. 1583.

B. erectis capitulis brevibus, foliis reflexis. Dill.

Musc. 365, t. 46. f. 24. Raii Syn. 95.

Angl. Drooping-leaved Fork-moss.

In paludibus montosis.

At North Bierley near Bradford, Yorkshire. Richardson. On the borders of lakes upon Glyder and Snowdon mountains. Dill. In Scotland. Dr. F. Buchanan, Mr. Dickson and Mr. G. Donn.

Perennis. Fl. vcre.

Caulis cæspitosi, unciales vel biunciales, simplices vel ramosi, undique foliosi, inferne radiculosi, ferruginei. Folia laxè imbricata, arcuato-recurva, lanceolata, acuta, integrinaria, pellucida, luteo-viridia, mox ferrugineo-fuscescentia; exsiccatione vix undulata aut tortuosa. Pedicelli unciales, crassiusculi, purpurei, subtortuosi. Capsula arcuato-cernua, ovata, lævis, rufo-fusca. Operculum longitudine capsulae, subulatum, arcuatum, rufum. Peristomium atro-purpureum, dentium laciniis capillaribus.

36. DICRANUM caule ramoso fastigiato, foliis poly-

X 3

lanceo-phyllosum.

lanceolatis apice ferrulatis siccitate tortilibus,
capsulâ ellipticâ erectâ.

Dicranum polyphyllum. *Engl. Bot.* t. 1217.

Bryum polyphyllum. *Dicks. Crypt. fasc.* 3. 7.
With. 825. *Hull.* 260.

B. cirratum β. *Huds.* 486.

B. cirratum, setis et capsulis brevioribus et plu-
ribus. *Dill. Musc.* 378. t. 48. f. 41.

B. trichoides, erectis sublongis capitulis, extre-
mitatibus per siccitatem stellatis. *Raii Syn.*
98.

Angl. Many-leaved Fork-moss.

In sylvis montosis.

On Snowdon. *Dill.* At Aber near Bangor. *Rev.*
H. Davies. Ireland. *Mr. D. Turner.* At
Hafod, Cardiganshire.

Perennis. Fruct. fert autumno.

Caules unciales, ramosi, fastigiati, undique densissimè
foliosi. Folia imbricata, linear-lanceolata, acuta, ca-
rinata, uninervia, apicem versus ferrulata; juniora
dilutè ac luteo viridia; reliqua atrata, persistentia;
omnia siccitate crispa et intorta. Pedicelli mox late-
rales, semiunciales, erecti, tenues, luteoli. Capsula
erecta, elliptica, fusca, lævis. Operculum subulatum,
rectum, longitudine capsulæ. Peristomii dentes pur-
purei, apice capillares, pallidi. Calyptra striata, basi
dentato-lacera.

Scotia- 37. *DICRANUM* caule ramoso fastigiato, foliis
num. subulatis integerrimis siccitate subincurvis,
capsulâ ellipticâ, pedicello arcuato.

Angl. Curve-stalked Fork-moss.

In montosis.

In Ireland. *Dr. Scott.*

Perennis.

Perennis. Fl. . . .

Cum præcedentc convenit habitu, magnitudinc ut et foliorum colorc denigrato, at differt foliis angustioribus, integerriinis, siccitate paululùm incurvis, nec tortuosis. Pedicelli fulvi, arcuati. Capsula elliptico-cylindracea, recta, subfulcata. Operculum subulatum, gracillimum, obliquum, longitudine capsulæ, basi purpurascens.

38. DICRANUM? caule simplici, foliis capilla-*hyperboreis* confertis strictis, capsulâ ovatâ erctâ, reum. pedicello brevi.

Bryum hyperboreum. *Dicks. Crypt. fasc. 3. 9.*

With. 828. Hull. 261. Fl. Dan. t. 538. f. 1.

Angl. Northern Fork-moss.

In rupibus alpinis Scotiæ.

On Ben Nevis. *Dicks.*

Perennis? Fl. . . .

Hanc speciem nondum vidi. Ex facie proximis præcedentibus affinis videtur.

39. DICRANUM caule ramoso, foliis linearis-fusculatis siccitate tortilibus, capsulâ obovatâ *carpum.* rectiusculâ sulcatâ.

Dicranum polycarpum. *Ehrh. Crypt. 84.*

Fissidens polycarpos. *Hedw. Sp. Musc. 159.*

Crypt. v. 2. 85. t. 31.

Angl. Many-headed Fork-moss.

In alpibus Scoticis. *D. G. Donn.*

Perennis. Fl. vere; fructum fert æstate.

Habitus sequentis. Caules cæspitosi, ramosi, undique foliosi, ferrugineo-tomentosi. Folia lætè viridia, linearia, integerrima, carinata, apice subulata, siccitate incurvato-tortuosa. Pedicelli unciales, graciles, fulvi, numerosi. Capsula fusca, obovata, obtusa, erecta,

rectiuscula, vel paululùm incurva, sulcata, basi attenuata, simplex nec gibbosa neque apophysi suffulta. Operculum basi conicum, apice subulatum, gracile, incurvum. Peristomium rubrum.

strumi- 40. **DICRANUM** caule ramoso, foliis linearifoliatis siccitate tortilibus, capsulâ arcuatâ fulcatâ basi strumosâ.

Dicranum strumiferum. *Erb.* *Crypt.* 74.

Schrad. *Spicil.* 59. *Swartz.* *Musc.* *Suec.* 33.

Roth. *Germ.* v. 3. 186.

Fissidens strumifer. *Hedw.* *Sp. Musc.* 160.
Crypt. v. 2. 88. t. 32.

Bryum strumiferum. *Dicks.* *Crypt. fasc.* 3. 8.
With. 833. *Hull.* 264.
Angl. Swelling Fork-moss.

In pratis uliginosis Scotiæ. *Dicks.*

Perennis. Fl. autumno et hyeme, fructum æstate perficiens.

Caules unciales vel biunciales, erecti, cæspitosi, ramosi, undique foliosi, infernè ferruginco-tomentosi. Folia læte viridia, linearia, integerrima, carinata; apicè subulata, clongata, patula; basi dilatata; arefactione incurvato-tortuosa. Pedicelli unciales, graciles, fulvi. Capsula pallidè fusca, obovato-oblonga, arcuato-incurva, sulcata, basi attenuata, apophysi hinc gibbosæ insidens. Calyptra flava, apicè nigra. Operculum exactè subulatum. Peristomium dilutè fuscum.

Fructus formâ, et apophysi, a præcedente præcipue differt.

monta- 41. **DICRANUM** caule ramoso, foliis linearilanceolatis apice subulatis siccitate crispis, capsulâ cylindraceâ rectâ.

Dicranum montanum. *Hedw.* *Sp. Musc.* 143.
t. 35. f. 8—13.

Angl. Mountain Curled Fork-moss.

In paludosis alpinis.

On bogs in Scotland, but rare. *Mr. G. Donn.*

Perennis. Fl. . . .

Caules plerumque haud semiunciales, cæspitosi, ramosi, dense foliosi. Folia linear-lanceolata, integerrima, uninervia, pallidè viridia, mox fuscescentia, siccitate crispa; basi dilatata, ferè ovata; apice elongata, setacea. Pedicelli haud unciales, pallidè fuscæ, recti. Capsula ovato-cylindracea, recta, dilutè fusca, lævis, vel ætate provectioni subfulcata, basi vix strumosa. Operculum subulatum, breve. Peristomium fuscum.

42. DICRANUM caule subramoso, foliis setaceis *flexuo-*
rectiusculis, capsulâ ovato-cylindracea fulcatâ, *sum.*
pedicello flexuoso.

Dicranum flexuosum. *Hedw.* *Sp. Musc.* 145.

t. 38. f. 1—6. *Swartz.* *Musc. Suec.* 34.

Roth. *Germ. v.* 3. 162.

Bryum flexuosum. *Sp. Pl.* 1583. *Huds.* 484.
With. 834. *Hull.* 264. *Lightf.* 725. *Dicks.*
H. Sicc. fasc. 9. 19.

B. pilosum molle, setis intortis. *Dill. Musc.* 373.
t. 47. f. 33. A—D.

B. trichoides, capitulis erectis, pediculis intortis
 tenuibus virentibus. *Raii Syn.* 97.

β. *B. fragile.* *Dicks. Crypt. fasc.* 3. 5.

B. pilosi mollis, setis intortis, varietas minor.
Dill. loc. cit. f. F—G.

γ. *Sphagnum alpinum.* *Sp. Pl.* 1570. *Huds.* 465.
With. 783. *Hull.* 245.

S. subulatum viridiissimum, capsulis ovatis. *Dill.*
Musc. 245. *t. 32. f. 3.*

Angl. Zigzag Fork-moss.

In rupibus alpinis: γ in paludibus alpinis.

In

In Ireland. *Sherard.* North Wales. *Dill. Mr. Griffith, and Rev. H. Davies.* Scotland. *Lightf. Dicks.*

Perennis. Fl. Aprili.

Caules simplices vel divisi, erecti, cæspitosi, foliosi, altitudine varii; in β vix semiunciales. Folia undique imbricata, erecta, stricta, rariùs subsecunda, lutescentia, nitida, subulata, concava, enervia; basi dilatata; apice tenuissima, recta: inferiora nigricantia. Pedicelli variè flexuosi ac tortuosi, rigiduli, semiunciales. Capsula fusco-virens, elliptico cylindracea, concinna sulcata, parùm arcuata, basi obsoletè gibbosa. Operculum tubulatum, tenue, rufum, capsulâ vix brevius, basi nigricans. Peristomium rubrum.

Icon *Hedwigiana* varietatem β potius refert. Folia enervia, ut apud Cel. Rothium describuntur, inveni, nec ut apud *Hedwigium*, “ ductorum fasciculò instructa.”

Folia in γ mucrone diaphano terminata. Hæc varietas semper sterilis invenitur. Capsulæ sic dictæ, in exemplari Dilleniano, corpora extranea sunt.

* * *Foliis compressis, distichis.*

viridu- 43. *DICRANUM* caule adscendente, foliis sub-
lum. distichis imbricatis elliptico-lanceolatis, pedicello terminali, operculo basi ventricoso.

Dicranum viridulum. *Swartz. Musc. Suec.* 31.
et 84. t. 2. f. 3.

D. bryoides. *Swartz. Act. Holm. ann.* 1795. 240.

Fissidens exilis. *Hedw. Sp. Musc.* 152 t. 38.
f. 7—10.

Bryum viridulum. *Sp. Pl.* 1584. *Dicks. Crypt.*
fasc. 1. 3. t. 1. f. 5. et *fasc.* 4. 8.

B. paucisolium. *Wuth.* 814. *Hull.* 257.

Grimmia viridula. *Roth. Germ.* v. 3. 141?
Angl. Smallest Fern Fork-moss.

Inuliginosis et umbrosis.

On

On Enfield Chace. *Dickson.* At Burgh near Yarmouth. *Mr. D. Turner.* On shady clay banks near Forfar. *Mr. G. Donn.*

Annua. Fruct. fert hyeme et vere.

Radix fibrosa, villosa, fusca. Herba exigua. Caulis simplicissimus, brevissimus, obliquè adscendens. Folia pauca, minus absolutè disticha, imbricata, lètè viridia, elliptico-lanceolata, acuta, integerima, plana, uninervia, exsiccatione subrepanda; summa majora, sèpiùs arcuato-scunda. Pedicellus terminalis, caule duplò vel triplò altior, rectiusculus, fulvus. Capsula erecta, ovata, sub apice constricta, (nec, ut in icono Hedwigianâ videri est, elliptica,) ore dilatata. Operculum fulvum, longitudine vix capsulæ, basi ventricosum et capsulâ latius, apice subulatum, acutissimum, paulò incurvum. Peristomium rubrum, dentibus apice attenuatis, elongatis, subpatentibus.

Witheringii et Hedwigii dubitationes de synonymo Linnaeano minimè probatae sunt.

44. DICRANUM caule adscendente, foliis distichis imbricatis ovatis, pedicello terminali, difolium. capsulâ cernuâ, operculo conico.

Hypnum tamarindifolium. D. G. Donn. in epist.
Angl. Tamarind-leaved Fern Fork-moss.

In uliginosis.

On the banks of the lake of Forfar, but very sparingly. *Mr. G. Donn.*

Annua? Fruct. fert Januario.

Priore paulò majus. Caulis simplex, adscendens, foliosus, bilinearisp. Folia plura, distichè imbricata, verticalia, pallidè viridia, ovata, integerima, mucronulata, subrepanda, uninervia; superiora falcato-cernua. Pedicellus e summitate caulis brevioris, semiuncialis, flexuosus, ruber. Capsula cernua, vel omnino pendula, ovata, subarcuata, ore patulo. Operculum exactè conicum, rubrum, apiee haud subulato.

Fissidenti

Fissidenti palmato Hedwigii proxima at distincta species videtur. Discrepat enim foliis latioribus magisque numerosis, operculo breviore, conico, nec subulato.

bryoides. 45. DICRANUM caule adscendente, foliis distichis remotiusculis lancolatis siccitate repandis, pedicello terminali, capsulâ erectâ.

Dicranum bryoides. *Sibth.* 279. *Roth. Germ.* v. 3. 181..

Fissidens bryoides. *Hedw. Sp. Musc.* 153. *Crypt.* v. 3. 67. t. 29.

Hypnum bryoides. *Sp. Pl.* 1588. *Huds.* 493. *With.* 842. *Hull.* 267. *Relb.* 428. *Abbot.* 244. *Engl. Bot.* t. 625. *Fl. Dan.* t. 473. f. 1.

H. taxiforme exiguum, versùs summitatem capsuliferum. *Dill. Musc.* 262. t. 34. f. 1.

H. repens filicifolium non ramosum, pediculis brevioribus, versùs foliorum summitatem egredientibus. *Raii Syn.* 88.

Muscus polytrichoides, exiguis capitulis in summis surculis, seu foliis subrotundis erexit. *Moris.* v. 3. 629. sect. 15. t. 6. f. 11. *Vaill.* *Paris.* t. 24. f. 13.

Angl. Distant-leaved Fern Fork-moss.

In umbrosis humidis.

Annua. Fruct. fert hyeme et vere.

Præcedentibus duobus duplò vel triplò altius. Caulis simplicissimus, quandoque uncialis. Folia lætè viridia, numcrofa, verticalia, remotiuscula, nec imbricata, lanceolata, acuta, siccitate undulata, perfectè disticha, basi supernè sisca at vix vaginantia, subinde secunda. Pedicellus frè uncialis, terminalis, subflexuofus, fulvus. Capsula erecta, ovata, ore paululum coarctata. Operculum capfulâ brevius et angustius, conicum, subincurvum,

curvum, rubrum. Calyptra parva, albida, apice nigra. Peristomium rubrum, dentibus elongatis, attenuatis.

46. DICRANUM caule erecto, foliis distichis *osmundi-*
imbricatis vaginantibus elliptico-lanceolatis *oides*.
compresso-planis, pedicello terminali.

Dicranum *osmundioides*. Swartz. *Act. Holm.*
ann. 1795. 240.

D. *bryoides*. Swartz. *Musc. Suec.* 32. et 86.
t. 2. f. 4.

Fissidens osmundioides. Hedw. *Sp. Musc.* 153.
t. 40. f. 7—11.

Hypnum *asplenoides*. Dicks. *Crypt. fasc.* 2. 10.
t. 5. f. 5, exculo Swartzii synonymo. H.
Sicc. fasc. 1. 21. With. 843. Hull. 268.

H. *bryoides*. Ehrb. *Crypt.* 124.

Angl. Close-leaved Fern Fork-moss.

In palustribus et nemorosis alpinis.

In Scotland. Mr. Dickson. At Nant Francon,
Carnarvonshire. Mr. D. Turner.

Perennis. Fruct. fert vere et aestate.

(Caulis erectus, strictus, atro-fuscus, uncialis vel biuncialis, plerumque simplex, rarius ramum e medio exserens, vel basin versus furcatus. Folia numerosa, fusco-viridia, disticha, verticalia, arcte imbricata, basi superne fissâ vaginaria, elliptico-lanceolata, compresso-plana, obtusiuscula, subinde mucronulata et adunca; inferiora sensim minora, obtusiora et remota. Pedicellus terminalis, semiuncialis, atro-rubens. Capsula erecta, ovata, fusco-nigricans, ore dilatata.

47. DICRANUM caule erectiusculo basi fructi- *taxifoli-*
fero, foliis distichis imbricatis vaginantibus *um.*
ovato-lanceolatis compresso-planis.

Dicranum *taxifolium*. Swartz. *Musc. Suec.* 31.
Roth. Germ. v. 3. 183. *Sibth. 279.*

Fissidens

Fissidens taxifolius. *Hedw. Sp. Musc.* 155. t. 39.
f. 1—5.

Hypnum taxifolium. *Sp. Pl.* 1587. *Huds.* 493.
With. 845. *Hull.* 267. *Relb.* 428. *Abbot.*
244. *Engl. Bot.* t. 426. *Fl. Dan.* t. 473. f. 2.
H. taxiforme minus, basi capsulifera. *Dill.*
Musc. 263. t. 34. f. 2.

*H. repens filicifolium non ramosum, pediculis
brevioribus, ad radicem egredientibus.* *Raii*
Syn. 88.

Muscus pennatus, capitulis adianthi. *Vaill. Par.*
t. 24. f. 11.

Angl. Yew-leaved Fern Fork-moss.

In umbrosis humidis.

Perennis. Fruct. fert vere.

Magnitudine variat. Caulis saepius semiuncialis, siun-
plicissimus, undique densè foliosus. Folia lætè at
saturatè viridia, disticha, arctè imbricata, superne
fissa, vaginantia, ovata, seu ovato-lanceolata, obtu-
siuseula, compresso-plana. Pedicellus ex perichætio
squamofo, radicali, solitarius, caule altior, flexuosus,
ruber. Capsula ovato-cylindracea, inclinata, ore pa-
tulo. Operculum subulatum, tenuè, basi rubrum.

adiantoi- 48. *DICRANUM* caule erecto subramoso medium
des. versùs fructifero, foliis distichis vaginantibus
lanceolatis compresso-planis.

Dicranum adiantoides. *Swartz. Musc. Suec.* 31.
Roth. Germ. v. 3. 184. *Sibth.* 280.

Fissidens adiantoides. *Hedw. Sp. Musc.* 157.
Crypt. v. 3. 62. t. 26.

Hypnum adiantoides. *Sp. Pl.* 1588. *Huds.*
494. *With.* 844. *Hull.* 268. *Relb.* 429.
Abbot. 244. *Engl. Bot.* t. 264. *Dicks. II.*
Sicc. fasc. 10. 22.

H. taxi-

H. taxiforme palustre ramosum, majus et erectum. *Dill. Musc. 264. t. 34. f. 3.*

H. erectum filicifolium ramosum pinnulis acutis. *Raii Syn. 87.*

Muscus taxiformis ramosus. *Vaill. Par. t. 28. f. 5.*

Angl. Greater Fern Fork-moss.

Ad rivulos et in umbrosis paludosis.

Perennis. Fruct. fert vere.

Habitus duorum præcedentium. Caulis biuncialis, sæpius ramosus, undique foliosus. Folia oblongo-lanceolata, acutiuscula, approximata, vix imbricata, saturatè viridia; juniora lutescentia. Pedicelli solitarii, e perichaetio laterali medium versus ramorum, flexuosi, fulvi. Capsula ovata, subcernua, demùm curvata, nigricans. Operculum subulatum, gracile.

449. TRICHOSTOMUM. *Hedw. Fund. v. 2. 90. t. 8. f. 43, 44,* male. *Schreb. Gen. 760. Engl. Bot. t. 1152.*

Didymodon. *Hedw. Crypt. v. 3. 8. Schreb. Gen. 760.*

Cynontodium. *Hedw. Sp. Musc. 57.*

Caps. oblonga. Perist. dentibus triginta duo, filiformibus, rectius ulis, per paria vel approximatis vel basi connexis.

* *Dentibus peristomii basi liberis.* Didymodon,
Cynontodium. *Hedw.*

11. TRICHOSTOMUM foliis lanceolato-subulâ-*trifari-*
*tis trifariis carinatis integerimis, capsulâ ova-*um.
â, caule ramoso.

Cynontodium trifarium. *Hedw. Sp. Musc. 57.*
Swartzia

Swartzia trifaria. *Hedw. Crypt.* v. 2. 76. t. 28.

Ehrb. Crypt. 174.

Didymodon trifarium. *Swartz. Musc. Suec.* 28.

Bryum trifarium. *Dicks. Crypt. fasc.* 3. 8.

With. 827. Hull. 261. Abbot. 238.

Angl. Three-ranked Fringe-moss.

In sterilibus montosis. *Dicks.*

On stones at Clapham springs, Bedfordshire.
Abbot.

Perennis. Fruct. fert Martio.

Caules densè cæspitosi, vix semiunciales, erecti, mox ramosi, foliosi. Folia carinata, integerrima, acuta, patula, tristè viridia; inferiora lanceolata; superiora subulata, trifaria. Flores terminales; ex auctoritate Hedwigii, hermaphroditici. Pedicellus solitarius, erectus, purpureus, subflexuosus, e caule post florescentiam ramoso, mox axillaris. Capula ovata, erecta, parva, fusca. Operculum conicum, rectum. Dentes peristomii 32, breves, incurvi, per paria subapproximati, sed basi discreti.

capilla- 2. **TRICHOSTOMUM** soliis capillaceis bifariis
ceum. vaginantibus, capsulâ elliptico-cylindraceâ, operculo conico, caulisbus densissimè cæspitosis.

Trichostomum capillaceum. *Engl. Bot.* t. 1152.

Cynodontium capillaceum. *Hedw. Sp. Musc.* 57.

Swartzia capillacea. *Hedw. Crypt.* v. 2. 72.
t. 26. *Ehrb. Crypt.* 164.

Didymodon capillaceum. *Schrad. Spicil.* 64.

Swartz. Musc. Suec. 28. *Roth. Germ.* v. 3.
199.

Bryum capillaceum. *Dicks. Crypt. fasc.* 1. 4.
t. 1. f. 6. *With. 831. Hull. 262. Fl. Dan.*

t. 1000. f. 1. Hoffm. Germ. v. 2. 42.

B. aestivum.

B. æstivum. *Huds.* 485, excluso Dillenii syn.
 B. n. 1806. *Hall. Hist.* v. 3. 44. t. 45. f. 1.
Angl. Capillary Fringe-moss.

In paludibus montosis.

About Llanberris. *Rev. H. Davies.* Plentiful on
 Ingleborough hill. *Huds.* On the Pentland
 hills near Edinburgh.

Perennis. Fruct. fert Aprili, Maio.

Cauls densissimè cæspitosi, altitudine variis, palmares,
 quandoque spithamæi, filiformes, infernè tomento
 ferrugineo, denso, connexi, supernè foliosi. Folia
 disticha, luteo-viridia, capillacea, acuta, integerrima,
 basi dilatata et vaginantia. Pedicelli terminales, erecti,
 tenuissimi, fulvi. Capsula erecta, parva, fere cylindracea,
 dilutè fusca, tenerima. Operculum conicum,
 breve, rectum, rufum. Peristomii dentes 32, breves,
 erecti, per paria approximati.

3. TRICHOSTOMUM foliis capillaceis basi *pusil-*
concavis, capsulâ ovato-cylindracea, operculo *lum.*
subulato, caule simplicissimo.

Trichostomum pusillum. *Hedw. Crypt.* v. 1.
 74. t. 28.

Didymodon pusillum. *Hedw. Sp. Musc.* 104.
Swartz. Musc. Suec. 29. *Roth. Germ.* v. 3.
 197.

Bryum pusillum. *Dicks. Crypt. fasc.* 2. 6. *Wittb.*
 812. *Hull.* 257.

B. Didymodon. *Hoffm. Germ.* v. 2. 43.
Angl. Dwarf Fringe-moss.

In arenosis. *Dicks.*

Annua. Fruct. fert vere.

Cauls breves, vix trilineares, erecti, simplicissimi, foli-
 osi. Folia pallidè viridia, undique imbricata, quan-
 doque secunda, capillacea, acuta, integerrima; basi
 VOL. III. Y dilatata,

dilatata, concava, vaginantia, uninervia. Pedicellus terminalis, erectus, fulvus vel roseus, nitidus. Capsula erecta, ovato-cylindracea, fusca, parva. Operculum subulatum, obliquum, basi rubrum. Peristomii dentes 32, erecti, rubri, per paria approximati.

- rigidu-* 4. **TRICHOSTOMUM** foliis lanceolatis acuminatis, capsulâ ellipticâ, operculo subulato, caule ramoso.

Didymodon rigidulum. *Hedw. Sp. Musc.* 104.

Crypt. v. 3. 8. t. 4. Roth. Germ. v. 3. 198.

Bryum rigidulum. *Dicks. Crypt. fasc.* 4. 12.

Hoffm. Germ. v. 2. 40.

Angl. Little Rigid Fringe-moss.

In muris.

In Northamptonshire, and near Dublin. *Mr. R. Brown.*

Pereennis. Fruct. fert vere.

Caules unciales, cæspitosi, ramosi, foliosi. Folia undique imbricata, patentia, latè lanceolata, carinata, integrerrima, acuminata, rigidula, siccitate torta, fuscescentia. Pedicelli mox laterales, erecti, rubri, vix unciales. Capsula erecta, elliptica, fusca. Operculum subulatum, gracile, arcuatum. Peristomium pallidè fuscum, erectum, dentibus 32, tenuissimis, per paria approximatis.

- papillo-* 5. **TRICHOSTOMUM** foliis subulatis carinatis, sum. capsulâ ellipticâ erectiusculâ basi hinc gibbosâ, operculo conico, caule ramoso.

Bryum papillosum. *Dicks. Crypt. fasc.* 4. 12.

t. 11. f. 5.

Angl. Papillary Fringe-moss.

In turfosis alpium Scoticarum. *Dicks.*

Perennis. Fl. . . .

Caules cæspitosi, ramosi, fastigiati, semiunciales, foliosi. Folia saturatè viridia, undique imbricata, erecto-patientia, subulata, carinata, integerrima. Pedicelli terminales, unciales, erecti, rubri, basi vaginati. Capsula erectiuscula, elliptica, castanea, basi hinc gibbosa, vel subcalcarata. Operculum conicum, colore capfulæ. Peristomii dentes 32, erecti; monente D. Turnero, per paria approximati.

** *Dentibus peristomii basi connexis.* Trichostomum. Hedw.

16. TRICHOSTOMUM foliis subulatis carinatis *cirrato*-apicem versùs ferratis siccitate tortilibus, *capitum*. fulâ ovatâ erectâ.

Angl. Curled Fringe-moss.

In sylvis subalpinis.

In woods near Stirling. Dr. Buchanan. North Wales. Rev. H. Davies.

Perennis. Fruct. fert Aprili.

Habitus Dicrani polyphylli, at major. Caules ramosi, cæspitosi, fastigiati, dense foliosi. Folia undique imbricata, linearis-subulata, acuta, mutica, carinata, apicem versùs laxè ferrata, lætè viridia, demùm nigricantia, arefactione involuta et crispa. Pedicelli terminales, erecti, vix unciales, fulvi, saepius gemini vel ternati. Capsula crecta, ovato-elliptica, pallida, substriata, ore rubro, coarctata. Operculum convexum, longius rostratum, rectum, fulvum. Calyptra fusco-rutilans, sulcata, recta, apice subulata, basi campanulata, margine crenata et multifida. Peristomium pulcherrimum, rectum, longitudine dimidii ferè capsulae, dentibus 32, basi connatis, purpureis, apice capillaribus, carneis.

17. TRICHOSTOMUM foliis lanceolatis: mu-hetero-crone *stichum.*

crone diaphano ferrato, capsulâ cylindraceâ erectâ, caule cæspitoso.

Trichostomum heterostichum. *Hedw. Sp. Musc.* 109. *Crypt. v. 2. 70. t. 25. Roth. Germ.* v. 3. 193. *Swartz. Musc. Suec.* 30. *Schrad. Spicil. 64. Ehrb. Crypt.* 153.

Bryum heterostichum. *Dicks. Crypt. fasc.* 4. 14. *Hoffm. Germ. v. 2. 40.*

Angl. Serrated Hoary Fringe-moss.

In saxis et ericetis montosis.

Perennis. Fruct. fert Maio.

Caulis cæspitosi, ramosi, unciales vel sesquiunciales, foliosi. Folia tristè viridia, imbricata, magis minùs secunda, lanceolata, carinata, unincrvia, integerima, in mucronem album, membranaceum, diaphanum, ferratum, producta. Pedicelli terminales, subsolitarii, erecti, fusi, tortiles, haud unciales. Capsula erecta, ovato-cylindracea, lœvis, fusca, ore purpureo. Peristomium breve, rubrum. Operculum, ex auctoritate Hedwigii, subulatum, obliquum.

lanuginosum. 8. *TRICHOSTOMUM* foliis lanceolatis: mucrone diaphano dentato, capsulâ ovatâ, caule procumbente pinnatim ramuloso.

Trichostomum lanuginosum. *Hedw. Sp. Musc.* 109. *Crypt. v. 3. 3. t. 2. Roth. Germ.* v. 3. 188. *Swartz. Musc. Suec.* 29.

T. serratum. *Ehrb. Crypt.* 94.

Bryum hypnoides. *Sp. Pl.* 1584, α . *Huds.* 480, ζ . *With.* 819, var. 1. *Hull.* 259. *Abbot.* 242. *Fl. Dan.* t. 476.

B. lanuginosum. *Hoffm. Germ. v. 2. 41.*

B. hypnoides polycephalon *lanuginosum*, montanum. *Dill. Musc.* 372. t. 47. f. 32.

B. hypnoides, capitulis plurimis erectis, *lanuginosum*. *Raii Syn.* 97.

Hypnum

*Hypnum n. 1780. Hall. Hist. v. 3. 37. t. 46. f. 4.
Angl. Toothed Hoary Fringe-moss.*

In sylvis siccis, rupibus, et arenosis montosis.

Perennis. Fruct. fert autumno.

Caules procumbentes, sèpè spithamæi, subdivisi, flexuosi, foliosi, ramulis numerosis, brevibus, alternis, incurvis. Folia luteo ac tristè viridia, imbricata, subsecunda, lanceolata, sive lincari-lanceolata, carinata, uninervia, integerrima, mucrone longissimo, tenuissimo, albo, diaphano, membranaceo, crebrè dentato, nec ferrato. Pedicelli in apicibus ramulorum, solitarii, erectiusculi, semiunciales, torti, fulvi. Capsula ovata, brevis, erecta, lævis, luteo-fusca. Operculum subulatum, rectiusculum, fulvum. Peristomium rubrum, præcedente longius. Calyptre conica, fusca, margine dentata.

9. **TRICHOSTOMUM** foliis lanceolatis recur-*ericoides*, vis uninervibus: mucrone diaphano serrulato, capsulâ ovatâ, caule erecto.

Trichostomum ericoides. Schrad. *Spicil.* 62.

Roth. Germ. v. 3. 191.

T. elongatum. Ehrb. *Crypt.* 233.

Bryum ericoides. Dicks. *Crypt. fasc.* 4. 14.

B. elongatum. Hoffm. *Germ. v. 2.* 41.

B. hypnoides. Sp. Pl. 1585, γ. Huds. 480, ε.
With. 820, var. 4. Hull. 259, var. 4.

B. hypnoides, ericæ facie, capsulis barbatis, alpinum. Dill. *Musc.* 371. t. 47. f. 31.

B. hypnoides, capitulis plurimis erectis, non lanuginosum. Dill. in Raii *Syn.* 478.

Angl. Heath-like Hoary Fringe-moss.

In scopolis udis subalpinis.

On the higher part of Snowdon. Richardson. Dill.

Perennis. Fruct. fert primâ æstate.

Caules erecti, breviores quam in præcedente, minusque ramosi, ramulis brevibus, subrecurvis. Folia luteo-viridia, recurvata, subsecunda, lanceolata, carinata, uninervia, integerima, mucrone diaphano, minute ferrulato, nec dentato. Pedicelli subterminales, erecti, unciales, torti, saturatè purpurei. Capsula ovato-cylindracea, elongata, erecta, lœvis, fusca. Operculum subulatum, rectum. Peristomium longitudine ferè capsulæ, rectum.

canescens. 10. **TRICHOSTOMUM** foliis ovato-lanceolatis enervibus canaliculatis: mucrone diaphano ferrulato, capsulâ ovatâ, caule erecto.

Trichostomum canescens. *Hedw.* *Sp. Musc.* 111. *Crypt.* v. 3. 5. t. 3. *Roth. Germ.* v. 3. 192. *Sibth.* 283.

Bryum hypnoides. *Huds.* 480, α .

B. canescens. *Haffm. Germ.* v. 2. 41.

B. hypnoides, hirsutie canescens, *vulgare*. *Dill. Musc.* 368. *t. 47. f. 27?* Folia haud in omni ejus exemplari enervia mihi scrupulum injecere.

B. trichoides, erectis capitulis, lanuginosum.

Raii Syn. 97.

Angl. Common Hoary Fringe-moss.

In apricis montosis, arenosis et sterilibus.

Perennis. Fruct. fert Februario, Martio, sed rariùs.

Caules cæspitosi, erectiusculi, ramosi, unciales seu bunciales. Folia imbricata, subrecurva, luteo-viridia, ovata vel ovato-lanceolata, integerima, subrevoluta, carinata et canaliculata, at prorsus enervia, mucrone diaphano, ferrulato, divaricato. Pedicelli laterales, erecti, unciales. Capsula ovata, erecta, fusca. Operculum subulatum, rectum. Peristomium longitudine ferè capsulæ, rectum, rubrum.

Hæc species, omnium hujusce tribus frequentissima, ab auctoribus nostris omnibus ferè neglecta, foliis longitudinaliter

tudinaliter canaliculatis, at enervibus, monente Hedwigio, certo certius dignoscenda est.

11. **TRICHOSTOMUM** foliis lanceolatis acu-fasciculatis minatis revolutis carinatis, capsulâ ovato-ob-lare. longâ, caule ramoso diffuso.

Trichostomum fasciculare. Schrad. *Spicil.* 61.

Hedw. Sp. Musc. 110. *Roth. Germ.* v. 3. 190.

Swartz. Musc. Suec. 29.

Bryum fasciculare. Hoffm. *Germ.* v. 2. 42.

B. hypnoides, ß. *Sp. Pl.* 1585. *Huds.* 480.

Wittb. 820, var. 7. *Hull.* 259.

B. lutescens. Dicks. *Crypt. fasc.* 4. 14.

B. hypnoides, hirsutie virescens, fasciculare alpinum. *Dill. Musc.* 370. t. 47. f. 28.

Angl. Beardless Hoary Fringe-moss.

In saxis montosis frequens. *Dicks.*

On Snowdon. *Dill.* North Wales. *Mr. Griffith,* and *Rev. H. Davies.* Ireland. *Dr. Scott.* On the Pentland hills near Edinburgh, 1782.

Perennis. Fruct. fert vere.

Caules cæspitosi, diffusi, ramosi, ramis brevibus. Folia imbricata, erecta, lanceolata, carinata, uninervia, integerrima, revoluta, plūs minus acuminata, nec apice diaphana; juniora luteo-viridia, pallida; reliqua fuscescentia, persistentia. Pedicelli numerosi, subterminales, solitarii, semiunciales, torti et flexuosi, fusci. Capsula ovato-cylindracea, erecta, fusca. Operculum subulatum, rectiusculum, fulvum, longitudine capsulae, apice tenuissimo. Peristomium rufum, capsulâ duplò vel triplò brevius.

12. **TRICHOSTOMUM** foliis lanceolatis carinatis acutis apice subdiaphanis, capsulâ ovatâ, carpon. caule ramoso diffuso.

- Trichostomum microcarpon. *Hedw. Sp. Musc.*
 112. t. 23. f. 1—5.
Bryum microcarpon. Hoffm. Germ. v. 2. 42.
B. hypnoides, ♂. Linn. Fl. Suec. 392. Fl. Dan.
t. 475.
B. hypnoides, γ. Huds. 480.
B. macrocarpon. With. 822. Hull. 260.
B. n. 396. Linn. Fl. Lapp. ed. 1. 314. ed. 2.
326. Herb. Linn.
B. hypnoides alpinum, setis et capsulis exiguis.
Dill. Musc. 370. t. 47. f. 29.
Hypnum n. 1782. Hall. Hist. v. 3. 37.
Angl. Small-fruited Hoary Fringe-moss.

In saxis alpinis.

On Snowdon. *Dill. and Mr. Griffith.*

Perennis. Fruct. fert Augusto.

Habitus præcedentis sed minor. Folia undique imbri-
 cata, erecta, lanceolata, acuta, carinata, uninervia,
 integerriama, vix revoluta, pilo cano, brevi, sæpius
 terminata; juniora luteo-viridia, reliqua nigricantia,
 persistentia. Pedicelli haud semiunciales, pallidi, torti.
 Capsula ovata, erecta, dilutè fusca, brevis, ore paten-
 tiuculo. Operculum subulatum.

obtusum. 13. TRICHOSTOMUM foliis lanceolatis cari-
 natis acuminatis revolutis, caule ramoso dif-
 fuso, operculo hemisphærico mutico.

Trichostomum obtusum. *Bridel. Muscol. v. 1.*
 125.

Bryum hypnoides, ♂. Huds. 480. With. 820,
var. 6.

B. hypnoides alpinum, operculis obtusis. Dill.
Musc. 371. t. 47. f. 30.

B. aciculariforme. Hoffm. Germ. v. 2. addend.
p. 42?

Angl. Blunt-fruited Hoary Fringe-moss.

In rupibus alpinis.

On Snowdon. *Dill.*

Perennis. Fruct. fert Augusto.

Habitus T. microcarpi, at colore vix lutescente. Folia siccitate arctè adpresfa, lanceolata, carinata, uninervia, integerrima, revoluta, acuminata, nec apice diaphana; omnia tristè virentia. Pedicelli fulvi, mox fusi, terminales, solitarii, haud unciales. Capsula erecta, elliptico-cylindracea, fusca. Operculum convexum, muticum, quâ notâ ab affinibus certè dignoscitur.
Herb. *Dill.*

14. TRICHOSTOMUM foliis linear-lanceolatis *glaucens-* acutis glaucescentibus, capsulâ ovatâ, caule *dens.* erecto subramoso.

Trichostomum glaucescens. *Hedw. Sp. Musc.*

112. *Crypt. v. 3. 103. t. 37, B. Swartz.*

Musc. Suec. 30.

Bryum glaucescens. *Dicks. Crypt. fasc. 4. 10.*

Angl. Glaucous Fringe-moss.

In umbrosis subalpinis Scotiæ. *Dicks.*

Perennis. Fl. . . .

(Caules cæspitosi, erecti, vix semiunciales, sæpius ramosi.

Folia erecto-patula, glauco-viridia, linear-lanceolata, acuminata, nec aristata, integerrima, carinata, uninervia. Pedicelli terminales, erecti, graciles, pallidè lutescentes. Capsula erecta, ovata, dilutè fusca, tenuis, obsoletè fulcata. Operculum subulatum, oblique. Peristomium purpureum, breve.

145. TRICHOSTOMUM foliis ovatis concavius-*pilifer-* culis acutis piliferis, capsulâ ovatâ, caule bre-*rum,* viissimo.

Bryum piliferum. *Dicks. Crypt. fasc. 4. 10.*

Angl.

Angl. Hairy Fringe-moss.

Ad aggeres umbrosos, prope Aberfeldy. *Dicks.*

Perennis? Fl. . . .

Caules cæspitosi, brevissimi, ut ferè nulli. Folia pauca, congesta, patula, ovata, concaviuscula, carinata, uninervia, integerrima, tristè ac pallidè virentia, pilo proprii ferè longitudinis aristata. Pedicellus solitarius, terminalis, erectus, capillaris, tortus, fuscus, semiuncialis et ultrà. Capsula erecta, ovata, brevis, fusca, lævis, ore patulo. Operculum subulatum, rectum. Peristomium breve.

flexifolium. 16. TRICHOSTOMUM? foliis oblongis vaginantibus flexuosis apice serratis, capsulâ ovato-cylindraceâ, caule basi ramoso.

Bryum flexifolium. *Dicks. Crypt. fasc. 3. 5. t. 7. f. 9. With. 815. Hill. 258.*

Angl. Wave-leaved Fringe-moss.

In declivibus sterilibus.

On barren sloping ground near Croydon, Surrey.

Dicks.

Perennis? Fl. . . .

Caules e basi ramosi, ramis patulis; foliosis, vix semiuncialibus. Folia dilutissimè ac lätè viridia, imbricata, patula, ovato-oblonga, concava, flexuosa, carinata, uninervia; basi vaginantia, integerrima; apice ferrata, mutica. Pedicelli ex apice ramorum, solitarii, erecti, straminei, capillares, haud unciales. Capsula ovato-cylindraceâ, erecta. Operculum conicum, obtusum. Peristomium nondùm vidi, hinc genus incertum.

lineare. 17. TRICHOSTOMUM foliis linear-lanceolatis imbricatis acutis, capsulâ ellipticâ, operculo conico oblique, caule erecto.

Bryum.

Bryum lineare. *Dicks. Crypt. fasc. 3. 6. t. 8.*
f. 2. With. 837. Hull. 265.
Angl. Linear-leaved Fringe-moss.

In udis fabulosis vel calcareis.

In the south of Scotland. *Dicks.* In Anglesea,
 where there is a constant dripping of water. *Rev.*
H. Davies.

Perennis. Fruct. fert Aprili.

Faciem quodammodo gerit Dicrani pellucidi, sed distinc-
 tissima est, et revera Trichostomi species. Confer
Schrad. Journ. v. 2. 177.

Caules densè cæspitosi, sesquiunciales, erecti, supernè
 ramosi, fastigiati. Folia laxè imbricata, flavo-virentia,
 erecto-patula, linear-lanceolata, integerima, carinata,
 uninervia, acuta, mutica, siccitate paululum incurva;
 inferiora marcescentia, ferrugineo-fusca, terrâ calcareâ
 subindè incrustata. Pedicelli apicem versus ramorum,
 erecti, rubri, ferè unciales. Capsula erecta, elliptica,
 brevis, fusca. Operculum conicum, obtusè acumi-
 natum, rectiusculum, rufum, capsulâ brevius. Pe-
 ristomium breve, fuscum, tenuissimum.

18. TRICHOSTOMUM foliis lanceolatis obtu- *linoides.*
 siusculis mucronulatis, capsulâ ovatâ, calyprâ
 subulatâ longissimâ, caule subsimplici.

Bryum linoides. *Dicks. Crypt. fasc. 3. 8. t. 8.*
f. 3. With. 824. Hull. 260. Hedw. Sp.
Musc. 114.

Angl. Flax-leaved Fringe-moss.

In pascais Scotiæ. *Dicks.*

Perennis. Fl. . . .

Caules erecti, trilincares, simplices et ramosi. Folia
 laxè imbricato-patula, fusco-viridia, lanceolata, inte-
 gerrima, obtusiuscula, nervo valido in mucronulum
 producto. Pedicellus terminalis, semiuncialis, erec-
 tus, tortus, fuscus. Capsula ovata, erecta, fusca, ob-
 soletè

foletè striata, ore patentiusculo. Operculum subulatum, fulvum, capsulâ duplò ferè longius. Peristomium tenuissimum, rufum, longitudine capsulæ, certè, in exemplaribus ab amicissimo Dicksono communicatis, rectum, nec contortum. Confer *Hedw. Sp. Musc.* 114.

fontinalis, 19. **TRICHOSTOMUM** foliis lanceolatis, capsuloides. capsulis ovatis subsessilibus in apicibus ramulorum, caule ramosissimo natante.

Trichostomum fontinaloides. *Hedw. Sp. Musc.*

114. *Crypt. v. 3. 36. t. 14. Roth. Germ.*
v. 3. 195. Sibth. 283.

Fontinalis minor. *Huds. 468. With. 788.*
Hull. 275. Abbot. 231. Lightf. 695. Engl.
Bot. t. 557. Dicks. Dr. Pl. 17.

• F. triangularis minor carinata, e cymis capsulifera. *Dill. Musc. 257. t. 33. f. 2. Herb. Dill.*

F. minor, foliis triangularibus minus complicatis, capitulis in summis ramulis sessilibus. *Dill. in Raii Syn. 79.*

Hypnum n. 1796. *Hall. Hist. v. 3. 41*, exclusis synonymis Dillenii et Linnæi.

H. fontinaloides. *Hoffm. Germ. v. 2. 79.*

β. *Fontinalis alpina*. *Dicks. Crypt. fasc. 2. 2. t. 4.*
f. 1. With. 789. Hull. 275.

Hypnum nigricans. *Villars. Dauph. v. 3. 905. B. Angl. River Fringe-moss.*

In fluviis : β in rivulis subalpinis.

About the shores of the Thames at Lambeth. *Dill.* and *Mr. Sowerby*. At Battersea. *Mr. Dickson*. In the Isis at Oxford. *Dill.* and *Sibth.* At Oakley mill, Bedfordshire. *Abbot*.

β. In the alpine rivulets of Scotland. *Dicks.* North. Wales.

Wales. *Mr. Griffith.* In the river at Matlock bath, Derbyshire. *Mr. Teesdale.* On rocks in the river at Kirkby Lonsdale, Westmoreland, above the bridge, June 1783.

Perennis. Fruct. fert Maio—Augusto.

Caules densè cæspitosi, fluitantes, ramosissimi, foliosi, nigri, longitudine varii. Folia atro-viridia, laxè imbricata, lanceolata, integrerrima, acuta, carinata, nervo centrali valido; siccitate undulata et spiraliter torta. Capsulæ solitariæ in apicibus ramlorum lateralium, subsessiles, erectæ, ovatæ, fuscæ, læves. Opculum conicum, subarcuatum, fuscum, capsula duplo brevius. Peristonii dentes rubri, basi variè connexi et perforati, siccitate subtorti, mox decidui.

Varietas subalpina β differt tantum colore nigriore, foliis arefactione magis tortuosis, paululumque angustioribus.

450. TORTULA. *Hedw. Fund.* v. 2. 92. t. 8.
f. 38, 39. *Schreb. Gen.* 760. *Engl. Bot.* t. 1101.

Barbula. *Hedw. Fund.* v. 2. 92. v. 1. t. 6.
f. 29.

Caps. oblonga. *Perist.* dentibus numerosis, filiformibus, spiraliter ac multipliciter convolutis.

1. TORTULA caule brevissimo, foliis patentibus *rigida*. rigidis involutis obtusis enervibus, capsula cylindracea, operculo conico.

Tortula rigida. *Swartz. Musc. Suec.* 40. *Roth.* *Germ.* v. 3. 211. *Schrad. Spicil.* 66. *Sibth.* 286.

Barbula rigida. *Hedw. Sp. Musc.* 115. *Crypt.* v. 1. 65. t. 25.

Bryum rigidum. *Huds.* 477. *With.* 813. *Hull.* 257. *Rehb.* 426. *Abbot.* 241. *Smith's Tour,* v. I.

v. 1. 191. *Dicks. Dr. Pl.* 47. *Ehrb. Crypt.*
34. *Engl. Bot.* t. 180.

B. stellatum. *Schreb. Lips.* 80. *Hoffm. Germ.*
v. 2. 45.

B. acaulon, ericæ tenuifoliæ Gerardi folio. *Dill.*
Musc. 388. t. 49. f. 55.

Angl. Rigid Screw-moss.

In rupibus, aggeribus, muris, et clivis cretaceis.

Near Wigmore, Herefordshire; L. Brown. *Dill.*

On Ingleborough hill, Yorkshire. *Huds.* On a bank about three miles from Norwich in the Yarmouth road, September 1781. It has since been found by the Rev. Mr. Bryant on Heydon church, Norfolk; by Mr. T. F. Forster on Banstead downs, and by other botanists in various places.

Perennis. Fruct. fert hyeme et vere.

Radix ramosissima, tomentosa, densa, pallida. Caulis brevissimus, simplex. Folia patentia, saturatè viridia, rigida, oblonga, obtusa, enervia, integrerrima, margine involuta; subtùs convexa; arefactione incurvata. Pedicelli plerumque solitarii, terminales, unciales, erecti, nitidi, rubri, apice slavecentes. Capitula erectiuscula, cylindracea, recta, fusca, lœvis, ore truncata. Operculum subulato-conicum, obtusiusculum, subrecurvum, longitudine vix dimidii capsulæ, basi rubrum. Peristomium rufum, operculo brevius. Calyptra cylindracea, gracilis, longitudine capsulæ, mox decidua.

Operculum in exemplaribus suecicis, ab amicissimo Professore Swartzio missis, quam in nostris duplò brevius est.

mucron-
nulata. 2. TORTULA caule subdiviso, foliis patulis lanceolatis mucronatis carinatis, capsulâ ovato-cylindraccâ, operculo subulato.

Tortula

- Tortula mucronulata.* *Swartz. Musc. Suec.* 40.
T. unguiculata. *Roth. Germ.* v. 3. 206.
T. decipiens. *Sibth.* 285.
Barbula unguiculata. *Hedw. Sp. Musc.* 118.
Crypt. v. 1. 59. t. 23, exclusis synonymis.
Bryum mucronulatum. *Dicks. Crypt. fasc.* 3. 3.
With. 817. *Hull.* 258. *Hoffm. Germ.* v. 2.
 44.
B. tenue barbatum, foliis angustioribus et rari-
 oribus. *Dill. Musc.* 385. t. 48. f. 49.
Angl. Pointed-leaved Screw-moss.

In ericetis et aggeribus arenosis, murisve.

IPerennis. Fruct. fert Februario—April.

Facies Dicrani purpurei. Caules erecti, cæspitosi, semi-unciales, simplices vel divisi, foliosi. Folia pallidè viridia, laxè imbricata, erecto-patula, lanceolata, mucronulata, integerrima, carinata, nervo valido prominente: in ramlulis junioribus duplò minora et remotiora. Pedicellus terminalis, erectus, haud uncialis, elegantè purpureus, apice viridi-flavescens, quandoque incrassatus. Capsula erecta, ovato-cylindracea, fusca, lœvis, ore purpureo. Operculum subulatum, obliquum, longitudine ferè capsulæ, purpureo-fulvum. Peristomium rubrum, longitudine operculi. Calyptra subulata, pellucida, operculo paulò major.

Variat pedicello sub capsulâ incrassato; an *Tortula unguiculata Sibthorpii*?

3. *TORTULA* caule ramoso, foliis patentibus *unguicu-*
lanceolatis carinatis acuminatis, capsulâ ovatâ, lata.
 operculo arcuato.

Tortula unguiculata. *Sibth.* 285?

Bryum unguiculatum. *Huds.* 482. *With.* 830.
Hull. 262. *Rehb.* 426? *Abbot.* 242. *Dicks.*
H. Sicc. fasc. 6. 21.

B. nervosum. *Hoffm. Germ.* v. 2. 44?

B. un-

B. unguiculatum et barbatum, surculis in summitate crassioribus. *Dill. Musc. 383. t. 48. f. 47.*

B. angustis viridibus foliis, capitulis erectis, brevibus pediculis insidentibus, calyprâ falcatâ, vel avium unguiculas referente. *Dill. in Raij Syn. 96.*

Angl. Birds-claw Screw-moss.

In muris et arenosis.

Perennis. Fruet. fert Martio.

Magnitudine præcedentis. Caules densè cæspitosi, plurumque divisi et fastigiati. Folia lutescentia, lanceolata sive linear-lanceolata, integerrima, apice attenuata et elongata, patentia, carinata, nervo valido; siccitatem inflexa. Pedicellus terminalis, erectusculus, vix uncialis, ruber. Capsula erecta, ovata, seu elliptica, fusca, laevis. Opereulum subulatum, incurvum, rufum, capsulâ duplò brevius.

fallax. 4. **TORTULA** caule ramoso, foliis linear-lanceolatis carinatis recurvato-patulis, capsulâ elliptico-cylindraceâ, opérculo obliquo.

Tortula fallax. Swartz. *Musc. Suec.* 40. Roth. Germ. v. 3. 212.

Barbula fallax. Hedw. *Sp. Musc.* 120. *Crypt.* v. 1. 62. t. 24.

Bryum fallax. Dicks. *Crypt. fasc.* 3. 5. With. 833. Hull. 264. Hoffm. Germ. v. 2. 44.

Angl. Fallacious Screw-moss.

In muris et arenosis.

Perennis. Fruet. fert primo vere.

Caules unciales, ramosi. Folia luteo-viridia, linear-lanceolata, acuta, subrevoluta, recurvato-patentia, carinata, nervo crasso: siccitatem parùm inflexa. Pedicellus erectus, uncialis, ruber, terminalis; statim, elongatis ramulis,

ramulis, lateralis fit. Capsula erecta; ferè cylindracea; dilutè fusca, lævis. Operculum longitudine capsulæ, subulatum, basi infractum vel obliquum, nec arcuatum. Peristomium rubrum, deciduum.

5. TORTULÀ caule ramuloſo; foliis cäulinis lan-convolu-
ceolatis carinatis; perichætialibus obtusis con- ta:
volutis enervibus, operculo obliquo:

Tortula convoluta. Swartz. Musc. Suec. 41.
Roth. Germ. v. 3. 208. Schrad. Spicil. 66.
Sibth. 285.

Barbula convoluta. Hedw. Sp. Musc. 120. Crypt.
v. 1 86. t: 32, excluso syn. Linnæi:
Bryum convolutum. Dicks. Crypt. fusc. 2. 6.
H. Sicc. fasc. 1. 20. With. 816. Hull. 254.
Relb. 427. Abböt. 237. Hoffm. Germ. v. 2.
43.

B. setaceum. Huds. 481. Lightf. 729:
B. stellare nitidum pallidum, capsulis tenuissi-
mis. Dill. Musc. 381: t. 48. f. 44.
B. trichoides exile pallidum; erectis capitulis e
surculis annotinis egredientibus. Raii Syn. 96:
Angl. Convoluted Screw-moss:

In ericetis, aggeribus et muris.

Perennis: Fruct. fert hyeme et vere.

Caules exigui; ramosi, latè cæspitosi; foliosi. Folia
lætè viridia, imbricata, patula, lanceolata, vel subovata,
brevia, obtusiuscula, integerrima, carinata, uninervia;
siccitatem adpressa et contorta. Perichætia e basi ra-
morum, solitaria, erecta, ovato-oblonga, pallida; fo-
liolis convolutis, obtusis, enervibus. Pedicelli soli-
tarii, erecti, unciales, capillares; citrini vel fulvi; nitidi.
Capsula ovato-cylindracea, gracilis, plūs minus incur-
va, fusca, lævis. Operculum capsulâ duplo brevius,
subulatum, obliquum, fuscum, nitidum. Peristomium
rubrum, longitudine operculi.

A Mnio setaceo Linnæi tota cælo diversa. Variat altitudine caulis, et operculi longitudine.

stellata. 6. **TORTULA** acaulis, foliis ovatis carinatis incurvis, capsulâ erectâ ovato-cylindraceâ sulcatâ, operculo obliquo.

Bryum stellatum. *Dicks. Crypt. fasc. 2. 6*, ex clisis synonymis. *With. 812.* *Hull. 257.*

Angl. Starry Screw-moss.

Ad aggeres et rivulorum margines Scotiæ. *Dicks.*

Perennis? *Fl. . . .*

Plantæ exiguæ, cæspitosæ, acaules. Folia numerosa, congesta, saturatè viridia, ovata, obtusiuscula, mutica, integerima, carinata, nervo valido, rubicundo; siccitate incurva. Pedicellus solitarius, terminalis, erectus, semiuncialis, rufus, perichaetio orbatus. Capsula erecta, ovato-cylindracea, brevis, fusca, longitudinaliter sulcata. Operculum longitudine capsulæ, subulatum, obliquum. Peristomium operculo æquale, fulvum. Calyptra cylindracea, operculo duplò longior.

Ex dictis patet quam multis et insignibus notis a præcedente discrepat; quid igitur existimandum de Hedwigii sententiâ, *Sp. Musc. 117?*

ruralis. 7. **TORTULA** caule ramoso, foliis obtusis recurvis apice piliferis: summis stellatis, capsulâ ovato-cylindraceâ.

Tortula ruralis. *Ehrb. Crypt. 184.* *Swartz.*
Musc. Suec. 39. *Roth. Germ. v. 3. 205.*
Sibth. 285.

Barbula ruralis. *Hedw. Sp. Musc. 121.*

Bryum rurale. *Sp. Pl. 1581.* *Huds. 476.* *With. 836.* *Hull. 265.* *Rehb. 426.* *Abbot. 241.*
Dicks. H. Sicc. fasc. 15. 18. *Hoffm. Germ. v. 2. 45.*

B. rurale

B. rurale unguiculatum hirsutum, elatius et ramosius. *Dill. Musc.* 352. t. 45. f. 12.

B. majus, erectis falcatis capitulis, foliis latiusculis extantibus, in pilum cānescentem desinentibus. *Raii Syn.* 94.

Muscus capillaris tectorum, densis cespitibus, capitulis oblongis, foliis in pilum oblongum desinentibus. *Vaill. Paris.* 133. t. 25. f. 3, benè.

Angl. Great Hairy Screw-moss.

In muris, tectis rusticis, et arborum truncis, frequens.

iPerennis. Fruet. fcrt Februārio—Aprili.

Caules dense cæspitosi, unciales vel biunciales, elastici, ramosi, fastigiati, foliosi. Folia saturatè viridia, ovato-oblonga, obtusissima, ferè obovata, integerrima, uninervia, recurvato-patentia, pilo laxo, flexuoso, sæpe fulvo, aristata; summa stellata, pallidiora. Pedicelli terminales, mox laterales, solitarii, erecti, firmi, rubri, unciales vel sesquiuunciales. Capsula erecta, ovato-cylindracea, lævis, castanea. Operculum subulatum, arcuatum, capsulâ paulò brevius. Peristomium operculo duplò brevius, conicum, obtusum, arctissimè convolutum, apice fulvum, basi albidum, pulcherrimum. Calyptra longa, subulata, fusca, basi laxa et subinflata.

3. TORTULA caule subsimplici brevi, foliis *subulata*, ovato-lanceolatis acuminatis, capsulâ cylindraceâ, operculo subulato recto.

Tortula subulata. *Hedw. Sp. Musc.* 122. t. 27.

Swartz. Musc. Suec. 39. *Roth. Germ.* v. 3.

201. *Sibth. 284. Engl. Bot.* t. 1101.

Bryum subulatum. *Sp. Pl.* 1581. *Huds.* 476.

With. 832. Hull. 263. Relb. 423. Abbot.

242. *Ligstf. 719. Curt. Lond. fasc. 3, t. 66.*

*Dicks. H. Sicc. fasc. 14. 19. Hoffm. Germ.
v. 2. 46. Ehrb. Crypt. 24.*

*B. capsulis longis subulatis. Dill. Musc. 350.
t. 45. f. 10.*

*B. erectis longis et acutis falcatis capitulis, ca-
lyptra subfuscâ, foliis serpylli pellucidis. Raii
Syn. 92.*

*β. Tortula pilosa. Schrad. Spicil. 66.
Angl. Awl-shaped Screw-moss.*

In umbrosis humidis frequens; rarius in platis borealibus.

β. On old oaks at Bruisyard, Suffolk. Mr. Turner.

Perennis. Fruct. fert hyeme et vere.

Caules cæspitosi, simplices vel subramosi, sæpius brevissimi. Folia stellatim patentia, pulchrè viridia, latè lanceolata, quandoque obovato-lanceolata, brevius acuminata, integerriina, unincrvia; siccitate inflexa et tortuosa. Pedicelli solitarii, terminales, rectiusculi, unciales, vel altiores, fulvi vel rubri. Capsula erectiuscula, longissima, cylindracea, castanea, lævis. Operculum capsulâ ter quaterve plerumque brevius, conico-subulatum, obtusum, rectum, ferrugineum. Peristomium ferè prioris, at dentibus apice evolutis et patentibus. Calyptra subulata, basi mox fissa.

β dignoscitur foliis apice piliferis; fortè species distincta.

*muralis. 9. TORTULA caule subsimplici brevissimo, fo-
liis ovatis acutis piliferis, capsulâ elliptico-cy-
lindracea, operculo conico.*

*Tortula muralis. Hedw. Sp. Musc. 123. Swartz.
Musc. Suec. 39. Roth. Germ. v. 3. 203. Sibth.
284. Ehrb. Crypt. 194.*

*Bryum murale. Sp. Pl. 1581. Huds. 477. With.
811. Hull. 256. Relb. 423. Abbot. 240.
Hoffm. Germ. v. 2. 45.*

B. tegu-

B. tegulare humile, pilosum et incanum. *Dill.*

Musc. 355. t. 45. f. 14.

B. minus, erectis minūs falcatis capitulis, foliis latiusculis congestis, in pilum canescentem desinentibus. *Raii Syn. 94.*

Muscus capillaris minor, capitulis erectis, vulgatissimus, foliis in pilum desinentibus. *Vaill.*

Paris. 133. t. 24. f. 15.

Angl. Wall Screw-moss.

In muris et tectis, omnium vulgatissima.

Perennis. Fruct. fert hyeme et vere.

Præcedente longè minor. Caules cæspitosi, pulvinati, brevissimi, simplices vel divisi. Folia congesta, patula, tristè virentia, ovata, seu obovato-oblonga, uninervia, integerrima, acuta, pilo albo, proprii sæpiùs longitudine, aristata. Pedicelli numerosi, solitarii, terminales, recti, ferruginei, semiunciales, vix unciales. Capsula fusca, erectiuscula, elliptico-cylindracea, demùm subarcuata. Operculum conico-fubulatum, capsulâ triplò plerumque brevius, obliquum. Peristomium conicum, breve, rufum. Calyptora fubulata, longitudine capsulæ.

Variat foliis breviùs aristatis, vix piliferis, quæ Tortula æstiva Bridelii, ex auctoritate Hedwigii; conf. *Dill.* loc. citato, f. F, G.

10. **TORTULA** caule brevissimo subsimplici, *cuneifolia*—foliis obovatis reticulatis pellucidis mucronulatis, capsulâ cylindraceâ, operculo conico.

Tortula cuneifolia. Roth. Germ. v. 3. 213.

*Bryum cuneifolium. Dicks. Crypt. fasc. 3. 7.
With. 811. Hull. 256. Hoffm. Germ. v. 2.*

45.

B. murale β. *Huds. 477.*

B. humile, pilis carens, viride et pellucidum.

Dill. Musc. 356. t. 45. f. 15.

Angl. Wedge-shaped Screw-moss.

In aggeribus et arenosis.

On banks at Streatham, Surrey; plentiful about Oxford. *Dill.*

Biennis. Fruct. fert vere.

A priore discrepat foliis latioribus, cuneiformibus seu obovatis, oculo armato pulchrè reticulatis, colore lætioribus, apice vel prorsùs muticis, vel mucronulo brevi tantummodo aristatis. Pedicellus semiuncialis, fulvus. Capsula erecta, fusca, cylindracea, retusa. Operculum breviusculum, conicum, leviter inflexum.

ericeto- 11. **TORTULA** caule ramoso, foliis lanceolatis acutis siccitate contortis: summis stellatis, capsulâ ovato-cylindracea.

Bryum ericetorum. *Dicks.* *Crypt. fasc.* 2. 5.
Witt. 826. *Hull.* 261.

B. ruralis facie, foliis minoribus et non pilosis,
Dill. Musc. 354. t. 45. f. 13.

Angl. Curling Heath Screw-moss.

In ericetis turfosis Scotiæ. *Dicks.*

Perennis. Fl. . . .

Caules erecti, ramosi. Folia densa, patula, angustè lanceolata, integerrima, acuta, mutica, nervo valido, lutescente; exsiccatione variè inflexa et contorta. Pedicellus vix uncialis, terminalis, atro-purpureus. Capsula erecta, ovato-cylindracea, retusa, fusca, nitidula.

tortuosa. 12. **TORTULA** caule ramoso fastigiato, foliis lanceolato-linearibus carinatis siccitate tortilibus, capsulâ ovato-cylindracea.

Tortula tortuosa. *Hedw.* *Sp. Musc.* 124. *Ehrb.*
Crypt. 204. *Swartz. Musc. Succ.* 39. *Roth.*
Germ. v. 3. 202. *Schrad. Spicil.* 64.

Bryum tortuosum. *Sp. Pl.* 1583. *Huds.* 486.
Witt.

With. 830. *Hull.* 255. *Lightf.* 727. *Relb.* 423. *Dicks. Dr. Pl.* 48. *Fl. Dan.* t. 880. f. 1, nec 2. *Hoffm. Germ.* v. 2. 46.

B. *cirratum*, setis et capsulis longioribus. *Dill. Musc.* 377. t. 48. f. 40.

B. *trichoides longifolium*, *crassiusculis caulinis*, *capitulis erectis aduncis acutis*. *Dill. in Raii Syn.* 98.

Hypnum n. 1787. *Hall. Hist.* v. 3. 38. t. 45. f. 2.

Angl. Frizzled Mountain Screw-moss.

In montibus Cambriæ, Scotiæ, et Angliæ borealis.

Perennis. Fruct. fert Julio—Octobri.

Caules uneiales vel biumiales, densè cæspitosi, ramosi, fastigiati. Folia patentia, lætè ae luteo-viridia, lanceolato-linearia, angusta, acuta, carinata, canaliculata, uninervia, apud nos certè integerrima, nequamquam ferrata; (confer Schrad. et Hoffm. loc. cit.) ; arefactione incurvato-erispa et undulosa. Pedicellus uneialis vel sesquiuncialis, terminalis, erectus, purpureus, nitidus, apice citrinus. Capsula erecta, ovato-cylindracea, lævis, ferruginea, pallida, ore rubro. Operculum cylindraceo-conicum, obtusum, reetiusculum, capsulâ duplò brevius. Peristomium purpureum, longitudine operculi, laxè tortum, nee, ut in reliquis, arctè in cylindrum convolutum.

Foliorum habitu, etiam peristomii indole, Trichostomis, potius quam reliquis Tortulis, videtur affinis.

13. **TORTULA** caule ramoso, foliis ovato-lan**-brevifo-**ceolatis carinatis revolutis, pedicellis laterali**-lia-**bus, capsulâ cylindraceâ.

Bryum brevifolium, *Dicks. Crypt. fasc.* 2. 4.

H. Sicc. fasc. 11. 19. *With.* 831. *Hull.* 262.

B. *palustre brevifolium*, capsulis nigricantibus. *Dill. Musc.* 377. t. 47. f. 39. *Herb. Dill.*

B. trichoides, erectis capitulis fusco-nigris. *Dill.*
in Raii Syn. 96.

Angl. Short-leaved Screw-moss.

In paludibus.

Sent from Yorkshire by Dr. Richardson. *Dill.* On bogs in Scotland. *Dicks.*

Perennis. Fl. . . .

Caules ferè unciales, erecti, ramosi. Folia dilutè viridia, ovato-lanceolata (nec linearia), acutiuscula, mutica, integerima, revoluta, nervo valido, rubro; siccitate subincurva. Pedicelli laterales, unciales, erecti, nigrantes. Capsula erecta, cylindracea, lævis, atro-fusca. Cætera in exemplaribus desunt.

barbatum. 14. *TORTULA* caule basi ramoso, foliis elliptico-lanceolatis patulis subrevolutis, pedicellis lateralibus, capsulâ ovatâ.

Bryum barbatum. *Curt. Lond. fasc.* 4. t. 65.
With. 829. *Hull.* 254. *Relb.* 426. *Abbot.* 237. *Hoffm. Germ.* v. 2. 45.

B. decipiens. *Abbot.* 239, ex ipso auctore.

B. unguiculatum et *barbatum*, tenuius et stelatum. *Dill. Musc.* 384. t. 48. f. 48.

Angl. Bearded Lateral Screw-moss.

In muris et cricetis sterilibus.

Not unfrequent about London, especially near Charlton. *Curtis.*

Perennis. Fruct. fert hyeme et primo vere.

Caules semiunciales, cæspitosi, fastigiati, basi ramosi. Folia patula, saturatè viridia, lanceolata, quandoque elliptica, integerima, mutica, carinata, subrevoluta; siccitate recurva: summa stellata. Pedicelli numerosi, laterales, rubri, haud unciales. Capsula erecta, ovata, brevis, ferruginea, nitida, lævis. Operculum subulatum, obliquum, longitudine ferè capsulæ. Peristomium rubrum. Calyptra subulata, acuminata.

Hanc speciem cum T. convolutâ confudit Hudsonus.
D. Davies.

Nomina trivialia hujus et sequentis reverâ inepta sunt.

15. **TORTULA** caule ramoso, foliis subulatis *imberbis*, patulis: basi ovatis, pedicellis subterminalibus, capsulâ elliptico-cylindraceâ.

Bryum imberbe. Linn. Mant. 309. Huds. 481.
Wittb. 829. Hull. 254.

B. tenue imberbe et pallidum, foliis crebrioribus.
Dill. Musc. 382. t. 48. f. 46.

B. perangustis foliis et cauliculis, foliis crebrioribus et circa extremitates magis congestis, capitulis erectis ad summitatem magis egredientibus. Rajii Syn. 99.

Angl. Deciduous Screw-moss.

In aggeribus graminosis et ad muros.

Perennis. Fruct. fert primo vere.

(Caules sesquiunciales, ramosissimi, fastigiati, graciles.

Folia luteo-viridia, patula, linearis-subulata, elongata, integerrima; basi ovata, subrevoluta, carinata; inferiora minora; summa stellata: omnia siccitate inflexa. Pedicelli apicem versus ramorum, erecti, unciales, capillares, purpurei. Capsula erecta, elliptico-cylindracea, fusca, lâvis. Operculum conico-subulatum, rectiusculum, capsulâ duplo brevius. Peristomium ferrugineum, mox deciduum; hinc character et nomen triviale apud Dillenium, Linnæum et Hudsonum; e quorum errore multi sefellerunt.

16. **TORTULA** caule ramoso fastigiato, foliis ob-*aristata*, longis obtusis mucronulatis siccitate incurvato-tortis, capsulâ cylindraceâ.

Bryum aristatum. Dicks. Crypt. fasc. 4. 12.
t. 11. f. 7.

Angl. Short-pointed Screw-moss.

In muris.

Near Croydon. *Mr. Dickson.* At Cullen's wood-bridge near Dublin. *Dr. W. Stokes.*

Perennis. Fruct. fert vere.

Caulis unciales, ramosi, fastigiati. Folia pallidè viridia, omnia ferè magnitudine et distantiâ æqualia, ovato-oblonga, obtusa, integerrima, subrevoluta, carinata: nervo validiore; mucronulo brevi terminata: siccitate inflexa et tortuosa. Pedicelli terminales, haud unciales, pùrpurei. Capsula erecta, cylindracea, gracilis, pallidè fusca, subdiaphana, lèvis, nitida. Operculum subulatum, arcuatum, rubrum, vix longitudine capsulae. Peristomium deciduum.

451. ORTHOTRICHUM. *Hedw. Crypt. v. 2.*

96. *Swartz. Act. Holm. ann. 1795. 247.*

Weissia. *Ehrh. Beitr. v. 1. 33. Schreb.*

Gen. 760.

Cap. oblonga. *Perist.* exterius dentibus sedecim: interius dentibus octo vel sedecim, filiformibus: quandoque nullum. *Calyptra* pilis erectis sèpiùs hispida, angulata,

* *Peristomio dupli-*

*stria-
num.*

1. ORTHOTRICHUM caule ramoso, foliis lanceolatis carinatis revolutis patulis, calyptrà integrâ, peristomio interno sedecimdentato.

Orthotrichum striatum. *Hedw. Sp. Musc. 163.*
Crypt. v. 2. 99. t. 36. Swartz. Musc. Suec.

42. Dicks. Crypt. fasc. 4. 5.

Bryum striatum. *Sp. Pl. 1579, a. With. 810.*
Abbot. 236. Relb. 425.

Polytrichum striatum. *Huds. 471, a. Hull. 248.*
P. capsulis subrotundis, pediculis brevissimis in-

sidentibus,

sidentibus, calyptrâ striatâ, arboreum ramosum majus. *Raii Syn.* 91.

P. Bryi ruralis facie, capsulis sessilibus, majus. *Dill. Musc.* 430. *t.* 55. *f.* 8. *Herb. Dill.*

Weissia striata. *Sibth.* 287. *Roth. Germ.* v. 3, 220.

W. ithyphylla major. *Ehrb. Beitr.* v. 1. 191. *Crypt.* 292.

Angl. Common Bristle-moss.

In arborum truncis.

IPerennis. Fruct. fert Februario—Maio.

Caulcs unciales vel sesquiunciales, erecti, fastigiati, varie ramosi, undique foliosi. Folia imbricato-patula, lanceolata, acuta, mutica, integerrima, revoluta, carinata, avenia, glaberrima, fulco-viridia; summa pallidiora, patentiora, quandoque apice erosa. Capsulæ laterales, breviùs pedicellatae, erectæ, folia parùm superantes, cylindraceo-ovatae, dilutè fuscæ, demùm apice sulcatae, vix tortæ. Calyptra conica, acuta, luteo-fusca, subrutilans, longitudinaliter sulcata, apice fulva, basi integerrima, undique pilis erectis luteis hispida. Operculum depresso, mucrone cylindraceo, obtuso. Peristomium externum e dentibus sedecim, æqualibus, complanatis, subfuscis, erectis, vix reflexis: internum e ciliis totidem albis, inflexis, articulatis.

ORTHOTRICHUM caule ramoso, foliis lan- *affine.* ceolatis carinatis revolutis patulis, calyptrâ subcrenatâ, peristomio interno octodentato.

Orthotrichum affine. *Schrad. Spicil.* 67. *Hoffm.* *Germ.* v. 2. 26.

Polytrichum capsulis sessilibus, foliis brevibus, rectis, carinatis. *Dill. Musc.* 432. *t.* 55. *f.* 10?

Weissia affinis. *Roth. Germ.* v. 3. 217.

Angl.

Angl. Pale straight-leaved Bristle-moss.

In rupibus, forsitan haud infrequens.

Near Edinburgh, 1782.

Perennis. Fruct. fert æstate et autumno.

Habitus prioris speciei. Folia pallidiora, lutescentia, erecto-patula, linear-lanceolata, acutiuscula, mutica, integerrima, revoluta, carinata, avenia. Capsulæ subfessiles, cylindraceæ, a primo ortu sulcatæ, demum subtortæ. Calyptra minus pilosa, margine saepius crenata et lacera. Operculi mucro gracilis, margo ruberrimus. Peristomium externum sedecimdentatum, dentibus per paria arctè reflexis: internum e ciliis octo, capillaribus, albis, inflexis, vix, nisi ætate proiectiori, articulatis.

Hanc a præcedente non distinxit Ehrhartus; ut e specie mine in Herbario Linnæano patet.

*pumi-
lum.*

3. *ORTHOTRICHUM* caule subramoso, foliis ovato-lanceolatis revolutis patulis, calyprâ nudâ, peristomio interno octodentato.

Orthotrichum pumilum. Swartz. *Musc. Suec.* 42. et 92. t. 4. f. 9. Dicks. *Crypt. fasc.* 4. 5. *Angl.* Dwarf Bristle-moss.

In cortice arborum. *Dicks.*

Near Kilcock in Ireland. *Mr. R. Brown.* Near Dublin. *Dr. Scott.*

Perennis. Fruct. fert vere.

Caules vix semiunciales, basi ramosi. Folia ovato-lanceolata, acutiuscula, mutica, carinata, exsiccatione imprimis revoluta, integerrima, luteo-viridia. Capsulæ terminales, subfessiles, cylindraceo-ovatæ, sulcatæ. Calyptra striata, tota glabra, albida, apice fusca. Operculi mucro brevis, rectus, pallidus. Peristomii externi dentes octo, nec sedecim, patuli; interni ciliæ octo, inflexæ, albæ.

4. *ORTHO-*

4. ORTHOTRICHUM 'caule subramoso, foliis *diaphanae* lanceolatis acuminatis apice diaphanis, calypnum: trâ nudiusculâ subcrenatâ.

Orthotrichum diaphanum. Schrad. *Spicil.* 69.

Weissia diaphana. Roth. *Germ.* v. 3. 223.

Angl. Sharp hairy-leaved Bristle-moss.

In arborum truncis et ad palos.

At Henham, Suffolk, and on trees in Norfolk.

Mr. D. Turner. Near Forfar, but rare. *Mr.*

G. Donn. Ireland. *Mr. Templeton.*

Perennis. Fruct. fert Aprili.

Caules cæspitosi, parùm ramosi, vix semiunciales. Folia olivaceo-viridia, vel nigricantia, erecto-patula, lanceolata, acuta, integerrima, subrevoluta, apice pallida, acuminata, acumine setaceo, diaphano, albo. Capulæ terminales, brévius pedicellatae, ovato-cylindraceæ, octofulcatæ. Calyptre pallida, campanulata, plicata, parùm pilosa, quandoque omnino nuda, margine inæqualiter crenata, apice rufo-fusca. Operculum rubrum, convexiusculum, mucrone brevi. Peristomii externi dentes sedecim, erectiusculi, lineari-lanceolati, angusti: interni ciliæ totidem, inflexæ, albæ.

5. ORTHOTRICHUM caule subsimplici, foliis *aristatae* obovato-oblongis obtusis acumine diaphano, tum. calyprâ dentatâ nudâ.

Orthotrichum diaphanum. Dick. *Crypt. fasc.*

4. 5. t. 10. f. 12.

Angl. Blunt hairy-leaved Bristle-moss.

In arborum truncis.

On a willow tree in the botanic garden at Edinburgh. *Mr. G. Donn.* Norfolk. *Mr. D. Turner.*

Perennis. Fruct. fert Decembri, Januario.

A præcedente differt colore atro-viridi, foliis plûs minus obtusis,

obtusis, aeumine albo, setaceo, diaphano. Capsulæ terminales, breviūs pedicellatæ, ovato-cylindraceæ, octosuleatæ. Calyptra pallida, plicata, prorsùs nuda, margine erenato-dentata, apice nigricans. Operculum convexiusculum, rubrum, muerone brevi. Peristomium externum sedecimdentatum, patens; internum e ciliis totidem, inflexis, albis.

Anne præcedentis varietas?

rivulare. 6. ORTHOTRICHUM caule ramoso, foliis ova-tis obtusis revolutis muticis, calyprâ dentatâ nudâ.

Angl. River Bristle-moss.

In saxis et rupibus rivulorum Hiberniæ.

On rocks in a rivulet near Balbriggen. *Dr. Scott.*

On a stone in a rivulet near Denyaghy. *Mr. Templeton.*

Perennis. Fruct. fert . . .

Caules cæspitosi, biunciales, ramosi, fastigiati, unicolor foliosi. Folia atro-viridia, recurvato-patula, ovata, obtusa, revoluta, carinata, mutica; siccitate adpressa. Capsulæ terminales, subsessiles, foliis obvallatæ, ova-tæ, profundè octosuleatæ, luteo-fuscæ. Calyptra pallidè virens, campanulata, octocostata, omnino nuda, margine dentato-lacera, apice acuta, fuscescens. Operculum convexum, pallidè fuscum, margine rubro, acumine recto, paululum elongato. Peristomium flavidum: exterius e dentibus sedecim, per paria connatis: interius e ciliis totidem inflexis.

crispum. 7. ORTHOTRICHUM caule ramoso, foliis linearibus arefactione recurvis undulatis, pedicellis elongatis apice incrassatis.

Orthotrichum crispum. *Hedw. Sp. Musc.* 162.

Crypt. v. 2. 96. t. 35. *Swärtz. Musc. Suec.*

42. Dicks. Crypt. fasc. 4. 6. *Engl. Bot.*
t. 996.

Bryum

Bryum crispum. *With.* 827. *Abbot.* 236. *Rehb.*
425.

B. striatum, ♂. *Sp. Pl.* 1580.

Polytrichum striatum, ♂. *Huds.* 471.

P. capillaceum. *Hull.* 249.

P. capillaceum crispum, calyptis acutis pilosissimis. *Dill. Musc.* 433. t. 55. f. 11.

P. capsulis oblongo-rotundis, calyptis pilosissimis. *Raii Syn.* 91.

Weissia ulophylla. *Ehrb. Beitr.* v. 1. 191. *Crypt.*
175.

W. crispa. *Sibth.* 287. *Roth. Germ.* v. 3. 222.

Angl. Curled Bristle-moss.

In arboribus.

Perennis. Fruct. fert Martio, Aprili.

Caules ferè unciales, basi ramosi. Folia aretè imbricata, linearia, acuta, integerrima, carinata, dilutè ac luteo-viridia, exsiccatione recurva et undulato-crispa. Pedicelli terminales, elongati, semiunciales, apice incrassati, torti, sulcati. Capsulæ cylindraceæ, octofulcatæ, pallidè fuscæ. Calyptra undique pilosa, pilis flavis, apice fusca. Operculum luteolum, mucrone elongato, attenuato, rectiusculo. Peristomium extermum sedecimdentatum, dentibus per paria approximatis, recurvatis: internum e ciliis octo, nec apud nos pluribus, albis, incurvis.

Illi calyptram tegentes, in hâc, uti in aliis speciebus, puleherrimè articulati sunt.

** *Peristomio simplici.*

ORTHOTRICHUM caule ramoso, foliis lan-

ceolatis carinatis revolutis, peristomio sim-

plici, calyprâ pilosâ dentatâ.

Orthotrichum anomalum. *Hedw. Sp. Musc.* 162.

Crypt. v. 2. 102. t. 37. *Dicks. Crypt. fasc.*
4. 6.

Bryum

- Bryum striatum*, $\beta.$ *Sp. Pl.* 1580.
B. tectorum. *With.* 810.
Polytrichum striatum, $\beta.$ *Huds.* 471. *Relb.*
ed. 1. 397.
P. tectorum. *Hull.* 248.
P. Bryi ruralis facie, capsulis sessilibus, minus.
Dill. Musc. 431. *t. 55. f. 9.*
P. capsulis subrotundis, pediculis brevissimis in-
sidentibus, calyptrâ striatâ, arboreum et ter-
reste, minus ramosum et breve. *Raii Syn.*
91.
Weissia anomala. *Sibth.* 287. *Roth. Germ.* v. 3.
215.
W. minor. *Ehrb. Crypt.* 282.
W. ithyphylla, $\beta.$ *Ehrb. Beitr.* v. 1. 191.
Angl. Rough Single-fringed Bristle-moss.

In tegulis, muris antiquis, et saxis.

Perennis. Fruct. fert Februario, Martio, etiam æstate.
 Caules ramosi, densè cæspitosi, sesquiunciales. Folia
 atro-viridia, lanceolata, acuta, revoluta, carinata, mu-
 tica, recta. Pedicelli parùm plerumque elongati, torti,
 sulcati, rubicundi, terminales. Capsulæ cylindraceæ,
 luteo-rutilantes, octosulcatæ, rectæ. Calyptre fusca,
 undique pilosa, margine acutè dentata. Operculum
 depresso-planum, mucrone recto, obtuso. Peristomium
 simplex, dentibus sedecim, brevibus, erecto-patulis,
 vix, nisi in ætate teneriori, per paria connexis.

- nudum.* 9. *ORTHOTRICHUM* caule ramoso, foliis ova-
 to-lanceolatis carinatis, peristomio simplici,
 calyptrâ nudâ subintegrâ.
Orthotrichum nudum. *Dicks. Crypt. fasc.* 4. 6.
t. 10. f. 13.
Bryum sessile. *With.* 810.
Angl. Smooth Single-fringed Bristle-moss.

In umbrosis ad ripas fluviorum.

On posts by the Thames at Wandsworth. Mr.
Sowerby.

Perennis: Fruct. fert April.

Caules ramosi, ferè unciales. Folia atro-viridia, ovato-lanceolata, acutiuscula, mutica, carinata, parùm, nisi exsiccata, revoluta. Capsulæ terminales, breviùs pedicellatæ, ovatæ, apice imprimis sulcatæ. Calyptra pallida, omnino nuda, inæqualiter sulcata, margine vix crenata. Opérulum demùm planiusculum, mucrone brevi, obtuso. Peristomium simplex, dentibus sedecim, erectis, per paria connatis, demùm distinc-tis:

10. ORTHOTRICHUM acaule, foliis radicali-Browniæ linearibus planis; perichætialibus ovatis *anum*. concavis, peristomio simplici, calyprâ nudâ.

Bryum Brownianum. Dicks. *Crypt. fasc.* 4. 7.
t. 10. f. 16.

Angl. Stemless Bristle-moss.

In rupibus et saxis in sylvis umbrosis.

By the river side at Roslin near Edinburgh. Mr.
R. Brown.

Anmia? Fruct. fert

In cæspitibus latis, planis, amorphis; rupes umbrosas instar Byssi investit. D. Brown.

Radix fibrosa: Folia radicalia pauca, crecta; oculo nudo capillaria, ad lentem linearia, angustissima, obtusa; integerrima, enervia, dilutè olivacea. Perichætium solitarium, imbricatum, olivaceum; foliis brevius, foliolis ovatis, acutis, concavis, carinatis, integerrimis. Pedicellus erectus, foliis duplò longior, ruber, gracilis. Capsula ovato-cylindracea, fusca, lævis. Calyptra rufa, acuminata, rectiuscula, octosulcata, glaberrima, margine integra. Peristomium simplex, sedecimdentatum, dentibus erectis, per paria connatis ut in præcedente.

Muscus valdè paradoxus, stirpe Phascum, pedicelli lon-gitudine

gitudine et facie Bryum, at peristomii, ut et calypræ, structurâ atque indole omnino Orthotrichum est.

452. PTEROGONIUM. Swartz. Musc. Suec. 26.
Engl. Bot. t. 1085.

Pterigynandrum. Hedw. Sp. Musc. 80.

Capf. oblonga, e perichaetio laterali. *Perist.* simplex, dentibus sedecim, linearibus, erectis.
Calyptra saepius pilosa.

gracile. 1. PTEROGONIUM caule repente, ramis fasciculatis teretibus recurvis, foliis ovatis ferrulatis.

Pterogonium gracile. Swartz. Musc. Suec. 26.
Engl. Bot. t. 1085.

Pterigynandrum gracile. Hedw. Sp. Musc. 80.
Crypt. v. 4. 16. t. 6.

Hypnum gracile. Linn. Mant. 310. Lightf.
766. Willd. 860. Hull. 272.

H. ornithopoides. Hudf. 508.

H. gracile ornithopodioides. Dill. Musc. 320.
t. 41. f. 55.

Encalypta gracilis. Roth. Germ. v. 3. 154.
Angl. Slender Wing moss.

In arboribus et rupibus montosis.

On Enfield chace and Marlborough downs. Dill.

In Ireland. Mr. D. Turner. North Wales. Rev.
H. Davies. Plentiful about Edinburgh.

Perennis. Fruet. sert vere et estate, sed rarius.

Caules siliformes, repentes, implexi, ramulis erecto-recurvatis, fasciculatis, cæspitosis, cylindraceis, obtusiusculis, undique foliosis. Folia saturatè viridia, in apricis lutescentia, aretè imbricata, latè ovata, enervia, minutè ferrulata sive denticulata. Pedicelli e perichaetio exiguo laterali, solitarii, erecti, haud unciales, capillares,

capillares, rubri. Capsula erectiuscula, cylindracea. Operculum conicum, obtusiusculum, subarcuatum. Peristomium pallidum. Calyptra cylindracea, albida, hirta, pilis erecto-patentibus, deciduis.

2. PTEROGONIUM caule repente ramosissimo; *Smithii.*
ramis pinnatis a refactione incurvis, foliis in-
tegerrimis, perichætiis pedicello subæquali-
bus.

Pterogonium Smithii. *Swartz.* in *Schrad. Journ.*

v. 2. 173.

Hypnum Smithii. *Dicks. Crypt. fasc.* 2. 10.
t. 5. f. 4. *Dr. Pl.* 19. *With.* 851. *Hull.*
270. *Hedw. Sp. Musc.* 264. t. 68. f. 5—7.
Sm. Tour, v. 3. 234. *Dillwyn Tr. of Linn.*
Soc. v. 6. 183.

Polytrichum Smithii. *Hull.* 249.

Muscus squamosus filicinus repens atro-virens
crispus, ramulos Abietis maris, conis onustos,
perbellè referens, foliis minimis subrotundis
creberrimis, ex unâ parte caulis appressis, al-
terâ expansis, capitulis oblongis erectis, pedi-
culis quasi orbatis, calyprâ subhirsutâ. *Mich.*
Gen. 114. n. 98.

Angl. Curling Wing-moss.

In truncis arborum rarissimè.

(On trees about Barham downs, Kent.

Perennis. Fruct. fert vere.

(Caules cæspitosi, depresso, repentes, rigiduli, ramosissimi, ramis tripinnatis, undique foliosis, siccitate arctè et elasticè incurvis. Folia saturatè viridia, opaca, imbricata, vaginantia, subrotunda, obtusa, concaviuscula, integerrima, basi uninervia. Perichætia lateralia, subulata, pallida, longitudine ferè pedicellorum, foliolis interioribus apice setaceis. Pedicelli solitarii, erexit, lutescentes, perichætio ferè tecti, apice torti. Cap-

ſula erecta, ovato-cylindracea, ferruginea, lævis, ore
subcoarctato. Peristomii dentes 16, pallidi, tenuissimi,
decidui. Operculum conicum, obliquum. Calyptra
flava, nitida, pilis erectis undique hispida.

453. NECKERA. Schreb. Gen. 762. Hedw.
Fund. v. 2. 93. t. 8. f. 47--50. Engl. Bot.
t. 1180.

Capſ. oblonga, e perichætio lateralī. *Perist.*
exterius dentibus ſedecim, acutis: *interius*
dentibus ſedecim, filiformibus, intermediis.
Calyptra lævis.

**Foliis bifariis.*

pumila. I. NECKERA ramis pinnatis, foliis distichis ova-
tis ſubundulatis, pedicello vix perichætium
superante.

Neckera pumila. Hedw. Sp. Musc. 205. Crypt.
v. 3. 49. t. 20.

Hypnum pennatum. Dicks. Crypt. fasc. 1. 5.
t. 1. f. 8. With. 852. Hull. 268.

Fontinalis pennata. Huds. 468, exclusis synony-
mis.

Angl. Small Feathery Neckera.

In arborum truncis rariūs.

On trees between Troutbeck and Ambleside, as
mentioned by Mr. Hudson, where I found it in
1782. It has been gathered in woods in Sussex
by Mr. Dickson; and in Scotland by the Rev.
Dr. Walker.

Peregrinis: Fl. . . .

Caules depreſſi, ramosi, foliosi, ramis pinnatis, ramulis
crebris, ſubalternis. Folia distichè imbricata, pallide
virentia, pellucida, membranacea, ovata, mucronu-
lata, integerima, compreſſa, ſæpius transversè cor-
rugata

rugata vel undulata, quandoque plana. Perichætia infrequentia, axillaria, solitaria, subulata, e foliolis pluribus, arctè imbricatis, quorum interiora in acumine gracili elongantur. Pedicellus vix perichætio longior, rectus, fulvus. Capsula erecta, ovata, fulva, lœvis. Operculum subulatum, subarcuatum.

Synonymi Hudsoniani veritas e loco natali constat.

2. NECKERA ramis pinnatis, foliis distichis ob-*crispa*, longis transversè rugosis, pedicello perichætio duplò longiore.

Neckera crispa. Hedw. Sp. Musc. 206. Swartz.

Musc. Suec. 70. Roth. Germ. v. 3. 336.

Hypnum crispum. Sp. Pl. 1589. Huds. 496.

With. 847. Hull. 269. Lightf. 745. Dicks.

H. Sicc. fasc. 2. 23. Ehrb. Phytoph. 79.

Engl. Bot. t. 617.

H. pennatum, undulatèm crispum, setis et capsulis brevibus. Dill. Musc. 273. t. 36. f. 12.

H. repens crispum, ramulis compressis, filicinorum more dispositis. Raii Syn. 89.

H. n. 1769. Hall. Hist. v. 3. 35. t. 46. f. 5.

Muscus vulgaris pennatus major. Baub. Pin.

369, ex Halleri auctoritate,

Angl. Crisped Neckera.

In collibus cretaceis, et nemorosis montosis siccis solo calcareo.

Perennis, Fruct. fert Aprili.

Præcedente duplò vel triplò major. Caules laxè imbricati et coacervati, complanati, duplicatè pinnati, undique foliosi. Folia distichè imbricata, lœtè viridia, nitida, pellucida, oblonga, integerima, obtusiuscula, elegantissimè corrugata. Perichætia subulata, pedicello duplò vel triplò breviora. Capsula ovaṭa, erecta, rufa, ore subcoartato. Operculum subulatum, rec-

tiusculum, gracile, basi conicum. Calyptra albida, tenuis.

** *Foliis undique imbricatis.*

heteromalla. 3. NECKERA caule ramoso diffuso, foliis ovatis acutis concavis undique imbricatis, capsulis sessilibus secundis.

Neckera heteromalla. *Hedw. Sp. Musc.* 202. *Crypt. v. 3. 38. t. 15. Sibth. 304. Engl. Bot. t. 1180.*

Sphagnum arboreum. Sp. Pl. 1570. Huds. 465. Abbot. 229.

S. heteromallum polycephalum. Dill. Musc. 248. t. 32. f. 6.

S. cauliferum et ramosum minus hirsutum, capitulis crebris pilosis per ramulorum longitudinem adnascientibus. Raii Syn. 105.

Fontinalis secunda. Linn. Meth. Musc. 30. With. 790. Hull. 275. Dicks. Dr. Pl. 18.

Phascum repens. With. 785. Hull. 251. Angl. Lateral Neckera.

In truncis arborum.

At Badminton, Gloucestershire. *W. Sherard.* On Enfield chace. *Dickson.* In Clapham wood, Bedfordshire. *Abbot.* Not rare in Norfolk. *Mr. Turner.*

Perennis. Fruct. fert Februario, Martio.

Caules sesquiunciales, ramosi, diffusi, rigidi, foliosi, ramis pinnatis. Folia undique patentia, imbricata, luteo-viridia, ovata, acuta, concava, integerima, carinata, uninervia. Capsulae in apicebus ramulorum laterarium, subsessiles, solitariæ, secundæ, ellipticæ, lœvæ. Perichaetia ovata, e foliis imbricatis, acuminatis, capsulam superantibus. Operculum conicum, acutum,

acutum, rectum. Calyptra fusca, parva, margine lacera.

4. NECKERA caule ramoso diffuso, foliis ovato-*curtipendula*-lanceolatis acuminatis planis patulis, capsulis *dula*. pedicellatis cernuis.

Neckera curtipendula. *Hedw. Sp. Musc.* 209.

Roth. Germ. v. 3. 341. *Swartz. Musc. Suec.*

71. *Ehrb. Crypt.* 76.

Hypnum curtipendulum. *Sp. Pl.* 1594. *Huds.* 504. *With.* 864. *Hull.* 274. *Dicks. H. Sicc.* *fusca.* 4. 19.

H. dentatum *curtipendulum*, viticulis rigidis. *Dill. Musc.* 333. t. 43. f. 69.

H. arboreum repens, capitulis reflexis, brevibus pediculis insidentibus. *Raii Syn.* 89.

Angl. Pendulous Neckera.

In rupibus et arborum truncis, haud vulgaris.

On Enfield chace, Marlborough downs, and about Snowdon. *Dill.* In Yorkshire. *Richardson.*

Perennis. Fruct. fert vere.

Caules decumbentes, implexi, ramosissimi, foliosi, ramis sœpè pinnatis, ramulis subindè elongatis, apice denu-datis, radicantibus. Folia undique imbricato-patula, luteo-viridia, plana, ovato-lanceolata, acuminata, integririma, uninervia, margine incrassata. Perichætia subulata, elongata, at pedicello duplò breviora. Capsula ovata, pendula, rufa. Operculum conicum, acutum.

5. NECKERA caule repente, ramis erectis cy-lindraceis, foliis lanceolatis patulis subsecun-dis, capsulis erectis.

Neckera viticulosoa. *Hedw. Sp. Musc.* 209. t. 48.

f. 4, 5. *Roth. Germ.* v. 3. 339. *Swartz.*

A a 4

Musc.

Musc. Suec. 71. *Sibth.* 304. *Erbh.* *Crypt.* 185.

Hypnum viticulosum. *Sp. Pl.* 1592. *Huds.* 501. *With.* 848. *Hull.* 269. *Lightf.* 754. *Relb.* 429. *Abbot.* 249. *Dicks.* *H. Sicc.* fasc. 8. 22. *Engl. Bot.* t. 265.

H. subhirsutum, viticulis gracilibus erectis, capsulis teretibus. *Dill. Musc.* 307. t. 39. f. 43.

H. repens trichoides arboreum majus, capitulis et surculis erectis, minus ramosis. *Raii Syn.* 85.

Muscus squamosus, viticulis longioribus glabris.

Vaill. Paris. 137. t. 23. f. 1.

Angl. Cylindrical Neckera.

Ad arborum radices in sylvis montosis siccis, vix in collibus cretaceis umbriosis.

Peregrinis. Fruct. fert vere,

Caules filiformes, repentes, ramosi. Rami numerosi, erecti, bi- vel tri-unciales, simplices vel ramosi, cylindracei, dense foliosi, obtusi, sursum incrassati. Folia saturatè viridia, opaca, sèpè lutescentia, undique imbricata, arcuato-subsecunda, lanceolata, acutiuscula, integrerrima, uninervia, subcarinata, siccitate crispa. Perichaetia exigua. Pedicelli erecti, capillares, unciales, nudi. Capsula ovato-cylindracea, erecta, fusca. Operculum subulatum, rectum. Calyptra pallida, mucronulata;

454. *HYPNUM.* *G. Pl.* 564. *Juss.* 11. *Schreb.* *Gen.* 762. *Hedw.* *Fund.* v. 2. 94. *Engl. Bot.* t. 1038.

Leskia. *Schreb. Gen.* 762. *Hedw. Fund.* v. 2. 93. t. 10. f. 62—65.

Caps. ovato-oblonga, e perichaetio squamofo, laterali.

lateralis. *Perist. exterius* dentibus sedecim, basi dilatatis; *interius* membranaceum, variè dentatum et laciniatum. *Calyptra* lævis,

* *Capulis erectis, Surculis teretibus.*

1. HYPNUM caule repente ramoso, ramis fili-*subtile*, formibus, foliis lanceolatis acutis enervibus laxiusculis, operculo conico.

Hypnum subtile, Hoffm. Germ. v. 2, 70. Dicks. Crypt. fasc. 4. 17.

Leskea subtilis. Hedw. Sp. Musc. 221. Crypt. v. 4. 23, t. 9. Swartz. Musc. Suec. 69. Roth. Germ. v. 3. 335.

Angl. Slender Feather-moss.

Ad radices arborum in humidioribus.

In Anglesea. Rev. H. Davies. Near Ormesby, Norfolk. Mr. D. Turner, In Ireland. Dr. Scott and Mr. Templeton.

Perennis. Fruct. fert Martio, Aprili.

Caules repentes, implexi, cæspitosi, filiformes, ramis numerosis, erectis, brevibus, subsimplicibus, teretibus, gracillimis, undique foliosis. Folia laxè imbricata, luteo-viridia, opaca, lanceolata, acutissima, integerima, prorsus enervia. Pedicelli erecti, capillares, rubri, vix unciales. Perichaetium exiguum, pallidum. Capsula ovato-cylindracea, æqualis, erecta, rufa, ore patenti; rarius maturitate subincurva. Operculum conicum, breve.

2. HYPNUM caulibus erectis cæspitosis, ramis *pulchell*-simplicibus, foliis ovato-lanceolatis acumina-*lum*, tis enervibus subsecundis, operculo conico.

Hypnum pulchellum. Dicks. Crypt. fasc. 2. 13. t. 5. f. 6. H. Sicc. fasc. 9. 22. With. 847. Hull. 269.

Leskea pulchella. *Hedw. Sp. Musc.* 220. t. 55.
f. 7—12.

Angl. Beautiful Tufted Feather-moss.

In sylvis umbrosis alpinis.

In Scotland. *Dickf.* and *Mr. G. Donn.* Ireland.

Dr. Scott and *Mr. Templeton.* North Wales.

Mr. Griffith.

Pérennis. Fl. . . .

Caules dense cæspitosi, erecti, sesquiunciales, ramosi; ramis confertis, simplicibus, erectis. Folia luteo-viridia, sericeo-nitida, imbricata, subsecunda, ovato-lanceolata, integerrima, concava, longius acuminata, enervia, sed plicâ longitudinali, nervum simulante, quandoque notata. Perichætium exiguum, acutiusculum, e ramorum basi. Pedicellus erectus, caulem superans, capillaris, ruber, apice incrassatus. Capsula erectiuscula, urceolato-oblonga, pallidè fusca, oculo armato punctulata, basi attenuata. Operculum conicum, mucronulatum. Peristomium exterius breve, flavescens; interius album, duplò longius, dentibus uniformibus.

Amicissimus D. Turner, qui de hoc genere observationes plurimas utilissimas mecum communicavit, errorem detexit in iconе Dicksonianâ, ubi folium nervo instructum malè prostat.

polyanthos. 3. *HYPNUM* caule repente ramosissimo, foliis lanceolatis acuminatis enervibus imbricatis erectis, operculo mucronulato.

Hypnum polyanthos. *Schreb. Lips.* 97.

H. filiforme. *Huds.* 497.

H. filiciforme. *Huds. ed. 1.* 421.

H. filicifolium. *Linn. Mant.* 2. 310.

H. filifolium. *With.* 862. *Hull.* 273.

H. sericeum ramosius et tenuius, capsulis acuminatis. *Dill. Musc.* 327. t. 42. f. 62. *Herb. Dill.*

H. repens trichoides terrestre viridius minus, capitulis cernuis, minus tumidis. Raii Syn. 84.

Leskea polyantha. Hedw. Sp. Musc. 229. Crypt. v. 4. 4. t. 2. Swartz. Musc. Suec. 68. Roth. Germ. v. 3. 330.

L. filiformis. Sibth. 303.

Angl. Thread-shaped Feather-moss.

Ad truncos arborum, lignaque putrescentia, haud infrequens.

IPerennis. Fruct. fert Februario, Martio.

(Caules repentes, arcte implexi, ramis sparsis, erectis, quandoque subpinnatis, foliosis, filiformibus. Folia sericeo-nitida, lutescentia, undique imbricata, erecta, rarius subsecunda, lanceolata, acuminata, integerrima, concaviuscula, enervia. Perichaetia elongata, pallida, numerosa. Pedicellus erectus, uncialis, vel semiuncialis, ruber, siccitate spiralis. Capsula erecta, elliptico-cylindracea, ferruginea, ore subcoarctato. Operculum convexum, mucronulatum.

Quot nomina inepta ex errore typographico in editione primâ Hudsoni, *filiciforme* pro *filiforme*!

44. *HYPNUM caule ramosissimo, ramis incurvis, attenuata* .
foliis ovatis nervosis imbricatis : summis sub- *tum.*
secundis, operculo conico.

Hypnum attenuatum. Schreb. Lips. 100. Dicks.
Crypt. fasc. 2. 13. With. 860. Hull. 272.
Hoffm. Germ. v. 2. 67.

*H. repens trichodes arboreum majus, caulinis
ramosis.* Dill. Musc. 331. t. 42. f. 66, *ex-*
clusa fig. d. Herb. Dill.

Leskea attenuata. Hedw. Sp. Musc. 230. Crypt.
v. 1. 33. t. 12. Roth. Germ. v. 3. 333. Timm.
Megap. 222.

Angl.

Angl. Taper-branched Feather-moss.

In truncis arborum in sylvis Scotiæ. *Dicks.*

Perennis. Fl. . . .

Caules ramosissimi, adscendentes, implexi, ramis ramulisque incurvis, teretibus, foliosis, apice incrassatis, quandoque attenuatis. Folia lætè viridia, opaca, undique imbricata, ovata, obtusiuscula, brevia, integerim, uninervia, superiora præcipue secunda. Geminæ masculinæ sparsæ, albidae, foliolis acuminatis, patulis, Perichætia ejusdem ferè figuræ, at longè minus frequentia. Pedicellus uncialis, flexuosus, atro-fuscus. Capsula elliptico-oblonga, erecta, fusca. Operculum conicum, obtusum.

cylindri- 5. **HYPNUM** caule ramoso, ramis incurvis cylindraceis acutis, foliis arctè imbricatis ovatis acutis enervibus.

Hypnum cylindricum. *Dicks. Crypt. fasc. 2. 12.*
Wittb. 861. Hull. 273.

H. julaceum sericeum repens, capsulis cylindraceis. *Dill. Musc. 322. t. 41. f. 57?*

Angl. Cylindrical Feather-moss.

Ad arborum truncos in sylvis Scotiæ, *Dicks.*

Perennis. Fl. . . .

Caules repentes, filiformes, foliosi. Rami adscendentes, cylindracei, incurvi, apice attenuati, undique foliosi. Folia arctè imbricata, luteo-viridia, opaca, parva, ovata, acutiuscula, enervia. Perichætium albidum, strictum. Pedicellus rectus, ferrugineus, haud uncialis. Capsula erecta, ovato-cylindracea, recta, ferruginea, Hypno filiformi similis.

Folia in *Herb. Dill.* majora, rariisque crassiiores, quam in speciminibus Dicksonianis.

medium. 6. **HYPNUM** caule repente, ramis sparsis, foliis ovato-

ovato-lanceolatis acutis uninervibus imbricatis, capsulâ elliptico-cylindraceâ.

Hypnum medium. *Dicks. Crypt. fasc. 2.* 12.
With. 847. *Hull.* 269.

H. trichodes, capsulis oblongis in setis brevioribus. *Dill. Musc.* 331. *t. 42.* *f. 65.*

Leskea polycarpa. *Erb. Crypt.* 96. *Hedw.*
Sp. Musc. 225. *Roth. Germ.* v. 3. 334.

Angl. Long-headed Feather-moss.

In truncis arborum.

Woods in Scotland. *Dicks.* Ireland, on stones.

Mr. Templeton. Near the roots of young oaks in Kempston wood, Bedfordshire. *Rev. Dr. Abbott.*

Perennis. Fruct. fert Februario.

(Caules repentes, implexi, ramis numerosis, procumbentibus, filiformibus, confertis, ramosis, apice saepè incurvis, quandoque incrassatis. Folia undique arctè imbricata, lætè at saturatè viridia, opaca, parva, ovato-lanceolata, acuta, integerrima, concaviuscula, nervo valido, prominente. Pedicelli copiosi, erecti, capillares, rubri, haud unciales. Capsula erecta, luteo-ferruginea, elliptico-cylindracea, elongata, apice quandoque incurva. Peristomium interius duplo longius, dentibus tenuissimis, conniventibus, flavescentibus. Operculum conicum.

77. *HYPNUM* caule repente, ramis erectis, foliis *inundatō*-
ovatis concavis obsoletè uninervibus subse-tum.
cundis, capsulâ cylindraceâ.

Hypnum inundatum. *Dicks. Crypt. fasc. 4.* 17.

H. heterophyllum aquaticum, *polycephalum*,
repens. *Dill. Musc.* 293. *t. 37.* *f. 27,* B.
Herb. Dill.

Leskea paludosa. *Hedw.* *Sp. Musc.* 221. *Crypt.*
v. 4. 1. *t. 1,* excluso syn. *Linnæi.*

Angl.

Angl. Marsh Feather-moss.

Ad radices arborum in inundatis.

In Norfolk. *Rev. H. Bryant.* On the trunks of willows, chiefly about the roots, near Beverley, also in Wiltshire, and by the Thames at Chiswick. *Mr. Teesdale.*

Perennis. Fruct. fert Junio—Septembri.

Caulis repentes, implexi, ramis erectis, apice sæpè attenuatis. Folia parva, ovata, concava, obtusiuscula, fusco-viridia, patula, subsecunda, nervo tenuissimo, obsoleto. Pedicelli ferè unciales, fulvi. Capsula erecta, cylindracea, gracilis, ferrugineo-flavescens. Operculum conicum, obtusiusculum, breve, coccineum.

sericeum. 8. HYPNUM caule repente, ramis adscendentibus confertis, foliis ovatis acuminatis erectis trinervibus, operculo conico.

Hypnum sericeum. Sp. Pl. 1595. Hudf. 505. With. 864. Hull. 274. Lightf. 762. Relb. 430. Abbot. 251. Curt. Lond. fasc. 2. t. 69.

H. vulgare sericeum recurvum, capsulis erectis cuspidatis. Dill. Musc. 323. t. 42. f. 59.

H. repens trichoides terrestre luteo-virens vulgare majus, capitulis erectis. Raii Syn. 84.

Leskea sericea. Hedw. Sp. Musc. 228. Crypt. v. 4. 43. t. 17. Swartz. Musc. Suec. 69. Roth. Germ. v. 3. 329.

Neckera sericea. Sibth. 305.

Muscus arboreus splendens sericeus. Vaill. Paris. t. 27. f. 3.

Angl. Silky Feather-moss.

In muris, tegulis, collibus, arborum yœ, truncis, vulgaris.

Perennis.

IPerennis. Fruct. fert hyeme et primo vere.

IPrioribus majus. Caules arctè repentes, ramulis numerosissimis, adscendentibus, confertis, brevibus, teretibus, acutis, undique foliosis. Folia imbricata, erecta, luteo-viridia, nitidissima, ovata, acuminata, integerrima, trinervia, nervis lateralibus medium versus evanescētibus. Perichætium oblongum, foliolis interioribus longissimè acuminatis. Pedicellus atropurpureus. Capsula erecta, ovata, fusca, ore coarctata, quandoque subincurva. Operculum conicum, obtusiusculum, sèpè aduncum.

1. *HYPNUM caule erecto simplici, ramis fastigiatissimis terminalibus, foliis ovatis uninervibus serrulatis, operculo acuminato.*

Hypnum dendroides. Sp. Pl. 1593. Huds. 503.

With. 859. Hull. 272. Relh. 430. Abbot.

250. Dicks. Dr. Pl. 96. H. Sicc. fasc. 13.

21. Ehrb. Crypt. 16. Fl. Dan. t. 823. f. 2.

H. dendroides sericeum, setis et capsulis longioribus erectis. Dill. Musc. 313. t. 40. f. 48.

H. erectum, arbusculam referens, ramulis subrotundis confertim nascentibus. Raii Syn. 81.

H. n. 1747. Hall. Hist. v. 3. 29.

Leskea dendroides. Hedw. Sp. Musc. 228.

Neckera dendroides. Swartz. Musc. Suec. 71. Roth. Germ. v. 3. 340.

Angl. Tree-shaped Feather-moss.

in sylvis et pascuis uidis.

Perennis. Fruct. fert Februario, Martio, sed rariùs.

radices repentes, ferrugineo-tomentosæ. Caules solitarii, erecti, stricti, 2—4-unciales, simplices, foliis arctè adpressis squamosi, apice ramosi, ramis numerosis, cylindraceis, acutis, fastigiatis, undique foliosis. Folia imbricata, lutescentia, nitida, ovata, acutiuscula, uninervia, apicem versus serrulata, utrinque plicâ longitudinali post exsiccationem notata. Perichætium cylindraceum.

cylindraceum. Pedicellus sesquiuncialis; ruber, nitidus. Capsula erecta, elliptico-cylindracea, ferruginea. Operculum conicum, acuminatum, rectiusculum, rufum. Peristomium interius; monente Hedwigio, ad basin ferè divisum.

- curva-*
tum. 10. HYPNUM caule repente, ramis fasciculatis incurvis, foliis ellipticis concavis integerrimis uninervibus; operculo subulato.

Hypnum curvatum. Swartz. *Musc.* Suec. 64.
Dicks. *Crypt. fasc.* 4. 18.

H. myosuroides. Hedw. *Sp. Musc.* 266. *Crypt.* v. 4. 20. t. 8. Roth. *Germ.* v. 3. 300. Huds. 568, β.

H. myosuron. With. 865. Hull. 274.

H. myosuroides crassius, capsulis erectis. Dill. *Musc.* 316. t. 41. f. 50. *Herb.* Dill.

H. repens, triangularibus angustis foliis, ramulis subrotundis. Dill. in Raii *Syn.* 83.

H. viviparum. Necker. *Gallobelg.* 475. Ehrb. *Crypt.* 57.

H. n. 1745. Hall. *Hist.* v. 3. 29.

Muscus denticulatus minor sericeus nostras, capitulo adianthi. Vauill. *Paris.* t. 28. f. 4.

Angl. Curve-branched Bushy Feather-moss.

In truncis arborum, et in rupibus calcareis.

Perennis. Fruct. fert vere et aestate.

Caules repentes, implexi. Rami numerosi, simplices et subdivisi, plus minus fasciculati, cylindracei, acuti, incurvi, elastici, magnitudine ferè praecedentis. Folia undique imbricata, pallide viridia, nitidiuscula, subsaccrosa, elliptica, acuta, integerrima, inflato-concava, medio tenus obsoletè uninervia. Pedicelli frequentes, unciales, nitidi, fulvi. Capsula ovata, ferruginea, semper erecta. Operculum conicum, mucronatum, subincurvum.

II. HYPNUM

11. HYPNUM caule repente, ramis fasciculatis *myosu-*
incurvis, foliis lanceolatis acuminatis ferru-
latis uninervibus, capsulâ demùm cernuâ.

Hypnum myosuroides. *Sp. Pl.* 1596. *Huds.*
 507. *With.* 865. *Hull.* 274. *Rehb.* 431.
Abbot. 251. *Swartz.* *Musc. Suec.* 64.
Necker. *Gallobelg.* 474.

II. *myosurum.* *Schrad. Krypt.* 17. *Roth.*
Germ. v. 3. 301.

H. myosuroides tenuius, capsulis nutantibus.
Dill. Musc. 317. *t. 41. f. 51.* *Herb. Dill.*

H. polyanthon, triangularibus angustis foliis.
Dill. in Raii Syn. 83.

H. n. 1752. *Hall. Hist.* *v. 3. 31.*

Leskia myosuroides. *Sibth.* 303.

Angl. Mouse-tail Feather-moss.

In rupibus calcareis copiosè.

Perennis. Fruct. fert vere.

Præcedenti simile, sed duplo minus, ramulis longè te-
 nuioribus, foliis laxioribus, angustioribus, minùsque
 concavis, colore magis lutescentibus, oculo armato
 ferratis. Capsula subcylindracea, demùm inclinata,
 vel subcernua. Operculum breve, mucronulatum,
 aduncum.

12. HYPNUM caule procumbente pinnato, fo- *nigro-*
*liis ovatis acuminatis concavis enervibus sub- *viride.**
secundis, operculo conico acuminato.

Hypnum nigro-viride. *Dicks. Crypt. fasc.* 4. 18.
H. myosuroides ♂. *Huds.* 508. *var. 2.* *With.*
 865. *Hull.* 274.

H. myosuroides sericeum tenuius, capsulis erec-
tis. *Dill. Musc.* 318. *t. 41. f. 53.*
Angl. Dark-green Silky Feather-moss.

In muris vetustis, rupibusve, et in sylvis ad terram.

Perennis. Fruct. fert vere.

Caules pollicares, bipollicares et ultra, depresso, repentes, pinnatum ramosi, ramis brevibus, simplicibus, erectis, cylindraceis, subcurvatis. Folia undique dense imbricata, subsecunda, fusco-viridia, quandoque ferrugineo-lutescentia, ovata, inflato-concava, enervia, integrerima, apice longius acuminata, plus minus falcata; juniora late viridia, nitida. Pedicellus uncialis, cretus, ruber. Capsula erecta, ovata, obtusa, rufa, deinceps subarcuata. Operculum (ex Dillenio) magnum, conicum, acutum.

Donnia- 13. HYPNUM caulis simplicibus cæspitosis num. basi pedunculiferis, foliis undique imbricatis ovatis enervibus, capsula cylindracea.

Angl. Green Simple Feather-moss.

E Scotiâ misit D. G. Donn. *Herb. D. Turner.*

Perennis? Fl. . . .

Caules cæspitosi, ut videtur decumbentes, scsquiunciales vel biunciales, simplicissimi, basi attenuati, flagelliformes, foliosi. Folia undique laxè imbricata, late viridia, ovata, obtusiuscula, quandoque mucronulata, integrerima, enervia, oculo armato reticulata. Pedicelli radicales, solitarii, caule breviores, flexuosi, carnei, basi coccinei, perichaetio exiguo. Capsula erecta, urceolato-cylindracea, laevis, fusca, ore patentiusculo. Peristomium flavum. Operculum convexum, brevius acuminatum, subaduncum.

** *Capulis erectis. Surculis planis; foliis bifariis.*

compla- 14. HYPNUM caule pinnato, foliis distichis natum. obtusis mucronulatis compressis enervibus, operculo rostrato obliquo.

Hypnum

Hypnum complanatum. *Sp. Pl.* 1588. *Huds.* 495. *With.* 844. *Hull.* 268. *Relb.* 431. *Abbot.* 244. *Lightf.* 742.

H. pennatum compressum et splendens, capsulis ovatis. *Dill. Musc.* 268. *t. 34. f. 7.*

H. repens filicifolium ramosum, ramulis appressis et magis complanatis. *Raii Syn.* 87.

Leskea complanata. *Hedw.* *Sp. Musc.* 231. *Ehrb.* *Crypt.* 224. *Timm.* *Megap.* 222. *Swartz.* *Musc. Suec.* 68. *Roth.* *Germ.* v. 3. 328. *Sibtb.* 302.

Muscus trichomanoides filicifolius splendens. *Vaill.* *Paris:* *t. 23. f. 4.*

Angl. Flat Feather-moss.

in truncis arborum.

Perennis. Fruct. fert Martio, Aprili.

Caules palmares, spithamei, et ultra, vel cæspitosi erecti, vel arborum cortici arctè adhærentes depresso, plani, inæqualiter pinnatim ramosi, foliosi, ramis apice sèpè attenuatis et elongatis, foliolosis. Folia dilutè ac lætè viridia, nitida, membranacea, distichè imbricata, compressa, ovato-oblonga, obtusa, integerrima, cnervia, apice mucronulata. Perichaëtium subulatum, elongatum, coarctatum. Pedicellus brevis. Capsula erecta, elliptica, brevis, fusca. Operculum obliquè rostratum.

5. *HYPNUM* caule subpinnato, ramis omnibus *trichomanoides.* obtusis, foliis distichis, enervibus acinaciformibus muticis, operculo subulato.

Hypnum trichomanoides. *Schreb.* *Lips.* 88. *With.* 844. *Hull.* 268. *Relb.* 431. *Hoffm.* *Germ.* v. 2. 56.

H. complanatum β. *Huds.* 495.

H. pennatum *trichomanoides* *splendens ramosum.* *Dill. Musc.* 269. *t. 34. f. 8.*

B b 2

H. erectum

H. erectum filicifolium ramosum, pinnulis obtusis. *Dill. in Raii Syn.* 87.

Leskea trichomanoides. *Ehrb. Crypt.* 234. *Hedw. Sp. Musc.* 231. *Swartz. Musc. Suec.* 68. *Roth. Germ. v. 3.* 326. *Wilden. Berol.* 318. *Sibth. 303.*

Angl. Blunt Fern-like Feather-moss.

In arborum cortice, quandoque in terrâ.

About London, *Dill.* Near Newcastle, *Mr. Winch.*

In Ireland, *Mr. Templeton.*

Perennis. Fruct. fert Februario, Martio.

A praeccidente differt caulibus brevioribus, ramis minus concinnae pinnatis, oninibus obtusis, nunquam apice attenuatis, foliis recurvato-acinaciformibus, convexisculis, muticis, nec mucronulatis, sub lente, ex observationibus D. Turneri, tenuerrimè ferrulatis, operculo denique longiore subulato.

*** *Capulis cernuis.* *Surculis planis;* *foliis bifariis.*

denticu- 16. HYPNUM foliis bifariis ovato-lanceolatis
latum. acutis integerrimis enervibus, capsulâ cylindraceâ, operculo conico.

Hypnum denticulatum. *Sp. Pl.* 1588. *Hedw.*
Sp. Musc. 237. *Crypt. v. 4.* 81. *t. 31.* *Huds.*
494. *With.* 843. *Hull.* 267. *Swartz.*
Musc. Suec. 52. *Roth. Germ. v. 3.* 264.
Engl. Bot. t. 1260.

H. denticulatîm pennatum, pinnulis duplicatis recurvis. *Dill. Musc.* 266. *t. 34. f. 5.*

H. repens filicifolium non ramosum, pediculis et capitulis longioribus ad radicem egrediéntibus, foliolis utrinque duplicatis. *Raii Syn.* 88.

Muscus

*Muscus squamosus non ramosus major et minor,
capitulis incurvis.* *Vaill. Paris. t. 29. f. 8.*

M. pennatus, capitulis adianthi. *Moris. Hist.
v. 3. 626. sect. 15. t. 6. f. 36.*

3. *Hypnum sylvaticum.* *Mant. 2. 310. Huds.
494. With. 844. Hull. 268. Swartz.
Mus. Suec. 52. Roth. Germ. v. 3. 263.*

*H. denticulatim pennatum, pinnulis simplicibus
rectioribus.* *Dill. Musc. 267. t. 34. f. 6.*

*H. repens filicifolium ramosum, pediculis et ca-
pitulis longioribus e foliorum alis egredien-
tibus, foliolis utrinque simplicibus.* *Dill. in
Raii Syn. 88.*

Angl. Sharp Fern-like Feather-moss.

In udis umbrosis.

Perennis. Fruct. fert Martio, Aprili.

Caulis longitudine varii, erecti vel patentes, simplices et subdivisi. Folia pallide viridia, nitidiuscula, bifariam imbricata, patentia, ovato-lanceolata, acuta, integrerima, enervia, oculo armato reticulata, basi amplexicaulia et compresas. Pedicelli erecti, e basi caulium simplicium, at ramosorum medium versus, unciales vel sesquiunciales, rubri, nitidi. Perichætium parvum, laxum. Capsula subcernua, cylindracea, ore patente, saepè incurva, demum fusca. Peristomium flavum. Operculum conicum, plus minus acuminatum.

β suadentibus summo Hedwigio et amicissimo D. Turnero huc reduxi. Differt præcipue caule, ætate for- sitan, ramoso, medium versus fructifero.

17. *HYPNUM* caule repente, ramis simpliciis- *crenula-*
culis, foliis subbisariis ovatis acutis crenulatis tum.
uninervibus, operculo conico.

Hypnum crenulatum. *Engl. Bot. t. 1261.*

Angl. Crenate Feather-moss.

In arboribus annosioribus, vel ligno putrefacente.

B b

Near

Near Newcastle. *Mr. Winch.*

Perennis. Fl. . . .

Præccidente majus, colore saturatiore. Caules repentes, ferrugineo-tomentosi, ramis crebris, patentibus, uncialibus, subsimplicibus. Folia subdistichè imbricata, lætè at saturatè viridia, nitida, ovata, acutissima, parùm concava, haud reticulata, minutè crenulata, nervo centrali solitario, apicem versùs evanescente. Pedicellus uncialis, ruber, perichætio pallido. Capsula arcuata, elliptico-cylindracea, fusca, basi saturatiore. Operculum conicum, obtusum. Peristomium ferrugineum.

serrula- 18. **HYPNUM** caule ramosissimo repente, foliis *tum.* ovato-lanceolatis acuminatis ferratis basi uninervibus, operculo subulato.

*Hypnum serrulatum. Hedw. Sp. Musc. 238.
t. 60. Engl. Bot. t. 1262.*

Angl. Serrulated Feather-moss.

Ad terram in Hiberniâ.

Sent from Ireland by Dr. Scott to Mr. Turner.

Perennis. Fl. . . .

Diffrert a duobus proximè præccidentibus magnitudine minori, caule multifariam ramoso. Folia subdisticha, ovato-lanceolata, saturatè viridia, parùm nitida, acuminata, inæqualiter ferrulata, obsoletè reticulata, medio tenus uninervia. Pedicellus uncialis, ruber, nitidus. Capsula cernua, ovato-oblonga. Operculum subulatum. Peristomium rufum.

leptè-vi- 19. **HYPNUM** caule filiformi, ramis sparsis simplicibus, foliis ovatis acutis ferrulatis uninervibus, operculo planiusculo.

*Hypnum rotundifolium. Dicks. Crypt. fasc. 2.
10, excluso synonymo. Wiltb. 8⁴⁵. Hull.
268.*

Angl. Bright-green Feather-mofs.

Ad arborum radices.

In woods in Scotland. *Dicks.* Ireland. *Dr. Scott.*

IPerennis. Fl. . . .

Caulis filiformis, serpens, squamulosus, ramis adscendentibus, simplicibus, brevibus, scè lanceolatis. Folia distichè imbricata, lætè ac dilutè viridia, ovata, acuta, concaviuscula, ferrulata, nervo solitario, concolore, pellucido, itaque obsoleto. Pedicellus vix uncialis, pallidè eroceus. Capsula oblonga, arcuata. Operculum planiusculum, umbo natum, nec acuminatum.

Ex exemplari Dicksoniano descriptionem confeci. A H. rotundifolio Scopolii differt toto habitu, facie et magnitudine, foliis imbricatis nec sparsis, deinde operculo brevissimo nec subulato.

210. *HYPNUM* caule repente filiformi ramoso, Teesdale-
foliis lanceolatis acutis integrerrimis uniner- lii.
vibus laxis, operculo subulato.

Hypnum intricatum. *Dicks. Crypt. fasc. 2. 10.*
With. 849. Huit. 269. Engl. Bot. t. 202;
nec Schreberi.

Angl. Teesdalian Feather-moss.

In sylvis umbrosis.

In woods on the south-east side of the river at
Matlock bath. *Mr. Teesdale.* In Colin glen,
near Belfast. *Mr. Templeton.*

IPerennis. Fruct. fert Junio, Julio.

Caules repentes, implexi, filiformes, ramosi, ramis adscendentibus, brevibus, subsimplicibus. Folia laxè imbricata, disticha, alterna, lanceolata, aeuta, integrerrima, carinata, uninervia, atro-viridia. Pedicelli haud unciales, ruſi, basi insigniter bulbosi. Perichaetia strictiuscula, foliolis ovatis, acuminatis, uninervibus. Capsula inclinata, ovato-urceolata. Operculum subulatum, breviusculum, sèpè curvatum.

H. Teesdalii, *Dicks. fasc. 4. 16*, est verum H. intricatum Schreberi, Hedwigii aliorumque; hinc nomen inventoris huic speciei transferre mihi visum est.

riparium. 21. HYPNUM caule patulo ramoso, foliis ovatis acuminatis integerrimis uninervibus laxis, operculo conico.

Hypnum riparium. Sp. Pl. 1595. Hedw. Sp. Musc. 241. Crypt. v. 4. 7. t. 3. Hud. 505. With. 861. Hull. 273. Relb. 436. Sibth. 301. Abbot. 250. Swartz. Musc. Suec. 63. Roth. Germ. v. 3. 320. Fl. Dan. t. 649. f. 1.

H. aquaticum, flagellis et teretibus et pennatis. *Dill. Musc. 308. t. 40. f. 44. B, C, D.*

H. ramosum fluitans pennatum. *Raii Syn. 81. Angl. Short-beaked Water Feather-moss.*

In rivulis, et ad ripas fluviorum.

Perennis. Fruet. fert toto ferè anno, imprimis autumno.

Caules decumbentes vel fluitantes, palmates et ultra, irregulariter ramosi, foliosi. Folia laxè patentia, remotiuscula, disticha, saturatè et fusco-viridia, ovata, plus minus elongata et acuminata, integerrima, nervo valido. Pedicelli rufi, unciales, vel biunciales. Capsula cernua, ovato-cylindraea, brevis, obtusa, arcuato-gibba. Operculum conicum, breve, acutum. Peristomium rubrum.

ruscifo- 22. HYPNUM caule ramoso, foliis ovatis acutis ferrulatis uninervibus imbricatis, operculo subulato.

Hypnum ruscifolium. Necker. Musc. 181. Roth. Germ. v. 1. 466. et v. 3. 322. Dicks. Crypt. fasc. 3. 10. With. 848. Hull. 269. Relb. 433. Sibth. 297. Abbot. 246.

II. ruscif-

H. rusciforme. *Necker. Gallobelg.* 481. *Weis.*
Crypt. 225.

H. rutabulum ♂. *Huds.* 497.

H. riparioides. *Hedw. Sp. Musc.* 242. *Crypt:*
fasc. 4. 10. *t.* 4. *Roth. Germ.* v. 3. 321.

H. foliis rusciformibus, capsulis subrotundis.
Dill. Musc. 298. *t.* 38. *f.* 31; *etiam t.* 40.
f. 44. *A. Herb. Dill.*

H. repens, triangularibus minoribus foliis, pe-
diculis et capitulis brevioribus et tumidio-
ribus, majus. *Dill. in Raii Syn.* 80.

β. H. prolixum. *Dicks. Crypt. fasc.* 2. 13. *With.*
852. Hull. 270.

H. rivulare. *Ehrb. Crypt.* 252.

H. riparium β. *Weber. Goett.* 81.

H. aquaticum prolixum, foliis ovatis. *Dill.*
Musc. 298. *t.* 38. *f.* 32. *et t.* 85. *f.* 20.
Herb. Dill.

γ. H. rotundifolium. *Abbot.* 245, nec *Dicksoni.*
Angl. Long-beaked Water Feather-moss.

In rivulis, et ad saxa lignaque in fluviis.

Perennis. Fruct. fert Martio—Septembri.

Præcedenti perquām affine sed distinctum. Caules ra-
mosi, natantes; in β elongati, et basi denudati; in γ
abbreviati. Folia atro-viridia, undique ferè at laxè
imbricata, parùm disticha, latè ovata, acuta, nec acu-
minata, oculo armato plūs minūs ferrulata, uninervia.
Pedicelli vix unciales, purpurei. Capsula ovata, ob-
tusa, brevis, subcernua. Operculum convexum, tu-
midum, acumine subulato, curvato.

23. HYPNUM caule procumbente ramosissimo, *Swartz-*
foliis subdistichis remotis ovatis ferrulatis ii.
uninervibus, operculo subulato incurvo.

Hypnum Swartzii. *Turn. Musc. Hibern.* MSS.
H. atro-

H. atro-virens. *Swartz. Musc. Suec.* 65, ex-
clusis synon.

Angl. Swartzian Feather-moss.

In arboribus.

Sent from Ireland to Mr. Turner by Dr. Scott
and Dr. Whittley Stokes.

Perennis. Fl. . . .

Caules procumbentes, repentes, ramosissimi, filiformes,
foliosi, divaricati, laxi. Folia colore varia, plus minus
atro-viridia, subdisticha, remota, haud imbricata, ova-
ta, acuta, ferrulata, concaviuscula, uninervia. Pedi-
celli vix unciales, purpurei, seabri. Capsula ovata,
cernua. Operculum subulatum, incurvum, longitu-
dine capsulae. Peristomium fulvum.

undula- 24. *HYPNUM* caule procumbente ramoso, fo-
tum. lijs lanceolatis compressis undulatis ener-
vibus, capsula sulcatâ, operculo subulato.

Hypnum undulatum. Sp. Pl. 1589. Hedw.
Sp. Musc. 242. Huds. 495. With. 849.
Hull. 269. Lightf. 744. Dicks. H. Sicc.
fasc. 6. 22. Engl. Bot. t. 1181. Swartz.
Musc. Suec. 53. Roth. Germ. v. 3. 267.
Ehrh. Crypt. 26.

H. pennatum undulatum, lycopodii instar spar-
sum. Dill. Musc. 271. t. 36. f. 11.

H. repens crispum, caulinulis compressis, lycop-
odii in morem per terram sparsis. Raii
Syn. 88.

Angl. Waved Feather-moss.

In sylvis montosis siccis.

Perennis. Fruct. fert Aprili—Augusto, haud frequenter.
Muscus formosus, foliis distichis, compressis, albidis,
nitidis, transversim pulcherrime undulatis, valde con-
spicuus. *Habitus* est ferè *Neckeræ* crispæ. Caules
ramosi,

ramosi, per terram sparsi, undique foliosi, basi attenuati. Folia basi binervia, absque nervo centrali. Pedicelli biunciales, pallidè purpurei, supernè subarcuati. Capsula cernua, elliptico-cylindracea, arcuata, longitudinaliter fulcatâ ut in Mnisiis; quod singulare. Operculum subulatum, recurvum.

25. HYPNUM caule erectiusculo ramofo, foliis *lucens.*
ovatis obtusis reticulatis enervibus, capsulâ
ovatâ, operculo subulato.

Hypnum lucens. Sp. Pl. 1589. Hedw. Sp.
Mus. 243. Huaf. 495. With. 845. Hull.
268. Lightf. 743. Dickf. H. Sicc. fasc. 2.
22. Roth. Germ. v. 3. 266. Ehrb. Crypt. 5.

H. pennatum aquaticum lucens, longis latisque
foliis. Dill. Musc. 270. t. 34. f. 10.

H. repens filicifolium ramosum, foliolis majo-
ribus magisque crebris. Dill. in Ruii Syn. 88.

Angl. Shining Membranaceous Feather-moss.

In udis umbrosis et palustribus.

Perennis. Fruct. fert Aprili—Julio.

Caules unciales vel biunciales, cæspitosi, erectiusculi,
undique foliosi, simplices vel ramosi. Folia sub-
disticha, laxè imbricata, plana; dilutè viridia, pellu-
cida, pulcherrimè reticulata ut in Jungermanniis, ova-
ta, sive obovata, obtusa, integerima, prorsùs ener-
via. Pedicelli solitarii vel gemini, unciales, rufi, apice
arcuati. Capsula ovata, brevis, turgida, lævis, demùm
pendula. Operculum subulatum, rectiusculum. Pe-
ristomium exterius rubrum: interius membrana-
ceum, plicatum, rectum, integrum, margine denti-
culatum.

Ad normam Hedwigii novum genus ex hâc specie for-
mari potuerit, quod habitus confirmaverit.

**** *Capulis cernuis.* *Surculis planis;* *foliis*
undique imbricatis.

26. HYPNUM caule bipinnato, foliis ovatis *con-splen-*
cavis dens.

cavis mucronulatis enervibus nitidis adpresso, operculo subulato recurvo.

Hypnum splendens. Hedw. Sp. Musc. 262. t. 67. f. 6—9. Sibth. 295. Host. Syn. 591.

H. parietinum. Huds. 499. Wilt. 853. Hull. 271. Lightf. 751. Relb. 434. Abbot. 247. Swartz. Musc. Suec. 53. Roth. Germ. v. 3. 276. Hoffm. Germ. v. 2. 60. Necker. Musc. 157. Weis. Crypt. 232.

H. proliferum. Ehrh. Crypt. 95. Willden. Berol. 321. Reich. Syst. v. 4. 489. Weber. Goett. 60. Pollich. v. 3. 136.

H. filicinum, tamarisci foliis majoribus splendentibus. Dill. Musc. 274. t. 35. f. 13.

H. repens filicinum, veluti spicatum. Raii Syn. 86.

H. n. 1765. Hall. Hist. v. 3. 33.

Muscus filicinus major slavecens ramosus. Vail. Paris. 140. t. 28. f. 1. t. 29. f. 1.

Angl. Shining Tamarisk Feather-moss.

In sylvis siccioribus, et ericetis.

Perennis. Fruct. fert Martio, Aprili, sed rarius, nisi in plagiis borealibus.

Caules spithamæi vel pedales, decumbentes, arcuati, proliferi, bipinnati, purpurei, nitidi, undique foliosi, ramis complanatis, apice attenuatis. Folia pallidè lutescentia, nitida, arctè imbricata, adpresso, parva, ovata, concava, integririma, encvia, mucronulata. Perichaetia saepius aggregata, subulata, coarctata. Pedicelli plerumque solitarii, sesquiunciales, rubri, nitidi. Capsula cernua, ferruginea, ovata, basi attenuata. Operculum basi conicum, apice subulatum, recurvum.

Hanc speciem cum *H. proliferum* apud Linnæum confundi videtur. *H. parietinum* ejus diversum est. Halterus Linnæum aliosque in errorem duxit de harum synonymis; confer *Linn. Mant.* 2: 507.

27. *HYPNUM* caule tripinnato, foliis cordatis *proliferis* acutis adpresso uninervibus trinervibus *rum*. opacis integerrimis, operculo subulato.

Hypnum proliferum. *Sp. Pl.* 1590. *Fl. Suec.*

395. *Huds.* 499. *With.* 853. *Hull.* 270.

Relb. 434. *Sibth.* 294. *Abbot.* 248. *Lightf.*

750. *Curt. Lond. fasc.* 1. t. 72. *Dicks. H.*

Sicc. fasc. 14. 21. *Swartz. Musc. Suec.* 53.

Roth. Germ. v. 3. 273. *Hoffm. Germ. v. 2.*

60. *Weis. Crypt.* 230.

H. tamariscinum. *Hedw. Sp. Musc.* 261. t. 67.

f. 1—5.

H. tamariscifolium. *Necker. Musc.* 158.

H. delicatum. *Ehrb. Crypt.* 301.

H. delicatulum. *Schrad. Spicil.* 73.

H. parietinum. *Willden. Berol.* 322. *Reich.*

Syst. v. 4. 490. *Weber. Goett.* 61. *Pollich.*

v. 3. 134.

H. filicinum, *tamarisci* foliis minoribus non splendentibus. *Dill. Musc.* 276. t. 35. f. 14.

H. repens filicinum minus, luteo-virens. *Raii Syn.* 86.

H. n. 1764. *Hall. Hist. v. 3.* 33.

Muscus filicinus major. *Vaill. Paris.* 140.

t. 25. f. 1. *Moris. v. 3.* 625. *sect. 15.* t. 5.

f. 20.

Angl. Opaque Proliferous Feather-moss.

In sylvis et ericetis frequens.

Perennis. Fruct. fert Martio, Aprili, at rarissime.

Præcedente plurimumque majus, ramis elegantè tripinnatis, filicinis, tomentosis, undique foliosis. Folia opaca, nec nitida, pulchre viridia, quandoque lutecentia, etiam fulva, arctè imbricata, cordata, vel cordato-ovata, integerrima, concaviuscula, acuta; maja trinervia; reliqua uninervia. Perichaetia sparsa, subulata,

ſubulata, nitida. Pedicelli ſepiūs aggregati, biunciales, purpurei. Capsula arcuato-cernua, cylindracea, fuſco-rubra. Operculum basi conicum, apice ſubulatum, curvatum.

recogni- 28. HYPNUM caule bipinnato, foliis cordatis
tum. acutis adpreſſis trinervibus opacis ferrulatis,
operculo conico acuto.

Hypnum recognitum. *Hedw. Sp. Musc.* 261.
Crypt. v. 4. 92. t. 35. *Roth. Germ.* v. 3.
279.

H. delicatulum. *Willden. Berol.* 321.

H. parietinum β. *Pollich. v.* 3. 135.

Angl. Lesser Opaque Feather-moss.

In pascuis et nemorosis, solo calcareo, rariūs.
Gathered at Matlock in 1790.

Perennis. Fruct. fert Junio, Julio.

Præcedenti perquām affine, at, ni fallor, distinctum.
Convenit colore luteteente, vel fulvo, opaco; foliorum
formā ac dispositione; denique toto habitu. Discrepat
magnitudine duplō vel triplō minore; caule vix proli-
fero, bipinnato nec tripinnato; foliis caulinis plūs
minūs ferrulatis, omnibus ferè trinervibus, nec, nisi
in ramulis tenuioribus, enervibus; pedicellis solitariis;
operculo breviūs et obtusiūs mucronulato.

In Britanniā haud anteā detectum fuit.

umbra- 29. HYPNUM caule bipinnato, foliis cordatis
tum. acuminatis ferrulatis plicato-striatis ener-
vibus, operculo conico inucronulato.

Hypnum umbratum. *Ehrb. Crypt.* 66. *Hedw.*
Sp. Musc. 263. t. 67. f. 10—12. *Hoffm.*
Germ. v. 2. 60. *Roth. Germ.* v. 3. 278.

Turn. Musc. Hibern. MSS.

Angl. Striated Proliferous Feather-moss.

In

In umbrosis.

In Ireland; Dr. Scott. *Mr. Turner.*

Perennis. Fl. . . .

Præcedente majus. Caulis interruptè pinnatus, quasi prolifer, ramis simplicibus pinnatisque. Folia pulchrè viridia, sive lutescentia, nitidiuscula nec opaca, patula, cordata, planiuscula, acuminata, ferrulata, enervia, sed multipliciter et longitudinaliter plicata, quasi sulcata. Perichætia laxiuscula, sèpè aggregata. Pedicelli solitarii, purpurei, ferè biunciales. Capsula ovata, brevis, arcuato-cernua. Operculum exactè conicum, breve, obtusum cuni mucronulo.

Fructificationem non nisi in speciminiibus exoticis vidi.

30. HYPNUM caule pinnato repente, foliis ova prælongo-lanceolatis acuminatis serratis uninervibus ^{gum.} patentibus, operculo subulato.

Hypnum prælongum. Sp. Pl. 1591. Hedw.
Sp. Musc. 258. Crypt. v. 4. 76. t. 29. Hud-
499. With. 855. Hull. 271. Relb. 434.
Sibth. 295. Abbot. 248. Swartz. Musc.
Suec. 54. Roth. Germ. v. 3. 280.

H. repens filicinum, triangularibus parvis foliis,
prælongum. Raii Syn. 80. Dill. Musc. 278.
t. 35. f. 15. A, C.

H. n. 1758. Hall. Hist. v. 3. 32.

Muscus filicinus minor. Vaill. Parif. t. 23.
f. 9.

Abbreviatum, caule erecto, vix pollicari, foliis con-
fertis. Turn. Musc. Hibern. MSS. Dill. t. 35.
f. 15. B.

Angl. Long Trailing Feather-moss.

In truncis arborum emortuorum, lignisve pu-
tridis.

B in

β in rupibus umbrosis.

Perennis. Fruct. fert Dcccmbri—Martio.

Caules graciles, arctè repentes, clongati, ramosi, pinnati, ramis subsimplicibus, tenuissimis, haud plerumque uncialibus. Folia pallidè lutescentia, quandoque saturatè viridia, laxè imbricato-patentia, lanceolata seu ovato-lanceolata, acuminata, acutè ferrata, nervo foliario, concolor. Pedicelli solitarii, purpurei, unciales vel paulò longiores. Perichaëtia laxa, foliolis recurvis. Capsula ovata, vel ovato-cylindracca, arcuata, cernua. Operculum subulatum, incurvum, longitudine ferè capsulæ.

Stokesii. 31. HYPNUM caule pinnato, foliis cordatis acuminatis serrulatis uninervibus patentibus, capsulâ ovatâ, operculo . . .

Angl. Alpine Trailing Feather-moss.

In rupibus Hiberniæ.

In the hollows of rocks under the eagles nest at Lough Bray. Dr. Stokes.

Perennis. Fl.

Habitu ad H. umbratum quodammodo accedit, characteribus verò cum præcedenti potius convenit. Caules laxè repentes, pinnati, vel subbipinnati. Folia imbricato-patentia, saturatè sed lætè viridia, nitidiuscula, cordata, acuminata, uninervia, crebrè serrulata, nec, ut in H. prælongo, acutè et remotè ferrata; siccitate planiuscula. Pedicelli unciales, atro-rusi. Perichaëtia laxa, albida. Capsula ovata, pendula. Operculum exemplario deest.

abietinum. 32. HYPNUM caule pinnato villoso, foliis cordatis acuminatis integerrimis uninervibus bisulcatis adpressis, operculo conico.

Hypnum abietinum. Sp. Pl. 1591. Hedw.
Sp. Mus. 353. Crypt. v. 4. 84. t. 32. Hud. 500.

500. *With.* 855. *Hull.* 271. *Rehb.* 435.
Sibth. 296. *Abbot.* 246. *Swartz.* *Musc.*
Suec. 54. *Roth.* *Germ.* v. 3. 272. *Necker.*
Musc. 163.

H. lutescens, alis subulatis tenacibus. *Dill.*
Musc. 280. t. 35. f. 17. *Herb.* *Dill.*

H. repens filicinum trichoides montanum, ramulis teretibus lutescentibus non divisis.
Dill. in *Raii Syn.* 86.

H. n. 1762. *Hall. Hist.* v. 3. 33.

Muscus terrestris, surculis filamentosis tenacibus abietinis, semel tantum divisis. *Moris. Hist.*
v. 3. 626. *sect.* 15. t. 5. f. 22.

M. pennatus minor, caulinis ramosis, in summitate veluti spicatus. *Vaill. Paris.* t. 29.
f. 12; nec tamen, monente Dillenio, Loefelii.

Angl. Fir-branched Feather-moss.

In collibus cretaceis siccis.

Perennis. Fruct. fert æstate, sed omnium rarissimè.
Caulis digitilongitudine, decumbentes, villosi, erubrè, ac plerumque simpliciter, pinnati, ramulis filiformibus, acutis, vix uncialibus, patentibus. *Folia* fulvo-viridia, opaca, undique imbricata, adpressa, cordata, acuminata, integerrima, paululum revoluta, uninervia, utrinque medio tenuis sulco longitudinali, nervum simulante, exarata. *Pedicelli* c speciminibus tantum Suecanis, in Herbario Linnaeano conservatis, vel ab amicissimo Swartzio missis, mihi obvii fuere; hi bunciales, purpurei, nitidi, flexuosi, capsulis ovato-cylindraceis, arcuato-cernuis. *Operculum*, monente Hedwigio, conicum.

Neckerus, ad hypothesis suam quo jure stabilendam, fructum ferri in hac specie, contra Linnaei assertionem, pernegavit.

* * * * * *Capſ. cernuis. Surculis teretibus; foliis undique imbricatis.*

alopecu- 33. HYPNUM caule erecto: infernè denudato: rum. supernè ramosissimo, foliis ovatis serratis uninnervibus, operculo subulato.

Hypnum alopecurum. *Sp. Pl.* 1594. *Hedw.*
Sp. Musc. 267. *Huds.* 503. *With.* 859.
Hull. 272. *Relb.* 436. *Sibth.* 299. *Abbot.*
249. *Swartz. Musc. Suec.* 60. *Roth. Germ.*
v. 3. 299. *Dicks. H. Sicc. fasc.* 13. 22.
Ebrh. Crypt. 37. *Engl. Bot. t.* 1182.

H. dendroides obscurius, setis et capsulis brevioribus subnutantibus. *Dill. Musc.* 315. t. 41.
f. 49.

H. palustre erectum, arbusculam referens, ramulis subrotundis. *Raii Syn.* 81.

Muscus dendroides elatior, radice repente.
Vaill. Paris. t. 23. f. 5.

Angl. Fox-tail Feather-moss.

In sylvis, ad rupes umbrosas vel speluncas humidas.

Perennis. Fruct. fert vere et primâ aëstate, sed rariis.

Radix repens, filiformis, horizontalis. Caules palmates vel spithamei, erecti; infernè simplicissimi, nudi, ferè aphylli; supernè ramosissimi, subpaniculati, ramis alternis fasciculatisve, simplicibus et divisis, arcuato-secundis, teretibus, acutis. Folia saturatè viridia, opaca, rariùs pallida vel subfulva, undique imbricato-patula, latè ovata, acutiuscula, concava, uninnervia, apicem versus ferrata. Pedicelli unguiculares, rubri, errecti. Perichaetia laxa. Capsula ovata, cernua. Operculum convexum, mucrone filiformi, recurvato, obtusiusculo.

Variat

Variat caulis maximè elongatis, ramosissimis, capillaribus, in torrentibus alpinis Hiberniæ. *D. Turner.*

4. **HYPNUM** caule subsimplici filiformi erecto, stramifoliis elliptico-lanceolatis enervibus concavis, neum. operculo convexo mucronulato.

Hypnum stramineum. *Dicks.* *Crypt. fasc.* 1. 6.

t. i. f. 9. *H. Sicc. fasc.* 11. 21. *With.* 860.

Hull. 273. *Hoffm. Germ.* v. 2. *addend. p. 77.*

Roth. Germ. v. 3. 325. *Swartz. Musc.*

Suec. 62.

Angl. Straw-like Feather-moss:

in aquosis gramineis:

in a bog on the west side of Hampstead heath.

Dicks. Near Yarmouth. *Mr. D. Turner.*

Perennis. Fl. . . .

caules palmatae et ultra, erecti, graciles, teretes, simplices vel subramosi, undique foliosi. Folia lutescentia, nitida, laxè imbricata; adpressa, elliptico-lanceolata, obtusiuscula, concava; integerriima, enervia, at, monente D. Turnero, sàpè longitudinaliter plicata. Pedicellus medium versus caulis; plerumque solitarius, erectus, biuncialis, capillaris, roseus, nitidus, perichaetio cylindraceo, stricto. Capsula ovata, arcuato-cernua, brevis, ferruginea. Operculum conico-convexum, mucronulatum. Calyptra apice atrata:

5. **HYPNUM** caule repente, ramis subsimpli-adnaticibus, foliis ovatis acuminatis integerrimis tum. enervibus concavis, operculo conico.

Hypnum adnatum. *Hedw.* *Sp. Musc.* 248.

t. 64. f. 5—10. *Turn. Musc. Hibern. MSS.*

Angl. Depressed Feather-moss.

ad terram in Hiberniâ.

at Rathmines near Dublin. *Dr. W. Stokes.*

Perennis. Fl. . . .

Caules repentes, depresso, vix unguiculares. Rami erecti scutuli, breves, subsimplices, undique foliosi. Folia fulvo-viridia, nitidula, imbricato-patula, parva, ovata, acuminata, concava, integerrima, enervia. Pedicelli unciales, rubri. Capsula cernua, ovata, ferruginea, ore patulo. Operculum conicum, breve, obliquum.

Differt ab *H. conferto* foliis integerimis, enervibus; ab abbreviato *Hedwigii* foliis acuminatis; ab utroque operculo brevi. *D. Turner.*

confertum. 36. HYPNUM caule repente, foliis ovatis acuminatis subserratis patulis mediotenus uninervibus, operculo subulato arcuato.

Hypnum confertum. *Dicks. Crypt. fasc. 4. 17.*
t. 11. f. 14.

H. clavellatum. *Huds. 508.* *With. 866.* *Hull. 274;*
excluso *Dillenii* synonymo.

Angl. Clustered Feather-moss.

In truncis arborum.

Perennis. Fruct. fert Februario.

Caules implexi; repentes, unciales. Rami copiosi, adscendentibus, breves, subsimplices. Folia late viridia, imbricato-patula, ovata, sive ovato-lanceolata, concava, acuminata, oculo armato serrulata, ad medium usque uninervia, nervo tenui, obsoleto. Pedicelli haud unciales, erecti, fulvi. Perichaetia e foliolis brevibus, angustatis. Capsula parva, ovata, obliqua. Operculum subulatum, arcuatum, longitudine ferè capsulæ.

An diversum a serrulato, n. 18?

murale. 37. HYPNUM caule repente, foliis ovatis integerimis arcte imbricatis mediotenus uninervibus, operculo subulato arcuato.

Hypnum murale. *Necker. Galio Belg. 479.* *Hedw.*
Sp. Musc. 240. *Crypt. v. 4. 79. t. 30.*
Dicks. Crypt. fasc. 3. 10. *With. 861.* *Hull.*
273.

273. *Roth. Germ.* v. 3. 318. *Hoffm. Germ.*
v. 2. 76.

H. confertum. *Engl. Bot.* t. 1038, monente
D. Turnero.

H. myosuroides γ. *Huds.* 508.

H. velutinum β. *Necker. Meth.* 172.

H. myosuroides brevius et crassius, capsulis cer-
nuis. *Dill. Musc.* 318. t. 41. f. 52. *Herb.*
Dill.

Angl. Wall Feather-moss.

in muris.

Perennis. Fruct. fert Januario, Februario.

Præcedenti simillimum, differt vero ramis concinnè ter-
retibus, e foliis arctè imbricatis, adpressis, integerri-
mis, breviùs mucronulatis; nec patulis, ferrulatis,
acuminatis. Capsula et operculum ferè prioris.

88. HYPNUM caule repente, foliis ovatis acu-*veluti-*
minatis ferratis mediotenùs uninervibus, cap- num.
fulâ ovatâ, operculo conico.

Hypnum velutinum. *Sp. Pl.* 1595. *Hedw.*

Sp. Musc. 272. *Crypt.* v. 4. 70. t. 27. *Huds.*

506. *With.* 866. *Hull.* 274. *Rehb.* 437.

Sibth. 301. *Abbot.* 251. *Swartz. Musc.*

Suec. 66. *Roth. Germ.* v. 3. 308.

H. velutinum, capsulis ovatis cernuis. *Dill.*

Musc. 326. t. 42. f. 61.

H. repens trichoides terrestre viridius minus,
capitulis tumidioribus cernuis. *Raii Syn.* 84.

H. repens trichoides terrestre minimum et
breve, capitulis tumidioribus cernuis. *Raii*
Syn. 84.

Angl. Velvet Feather-moss.

in pascuis, sylvis, et ad radices arborum.

Perennis. Fruct. fert Februario, Martio.

Caules repentes, densissimè implexi, pulvinati. Rami erecti, conferti, subsimplices, breves, foliosi. Folia luteo-viridia, imbricata, erecta, parva; basi ovata, uninervia; apice elongata, acuminata, sub lente serrata, nervo evanescente. Pedicelli unguiculares, rubri, apicem versus sèpè seabriuscùli. Capsula ovata, parva, ferruginea, arcuato-cernua. Operculum exactè conicum, acutum, rufum.

intricatum. 39. HYPNUM caule repente, foliis lanceolato-subulatis serratis mediotenuis uninervibus, capsulâ ventricosâ, operculo conico.

Hypnum intricatum. Schreb. Lips. 99. Hedw.

Sp. Musc. 271. Crypt. v. 4. 73. t. 28.

H. Teesdalii. Dicks. Crypt. fasc. 4. 16. t. 11. f. 13.

Angl. Matted Feather-moss.

In sylvis.

In Yorkshire. Mr. Teesdale. Ireland. Dr. Scott.

Perennis. Fruct. fert Martio.

A præcedente differt magnitudine parùm minore, foliis laxioribus, angustioribus, longius acuminatis; capsulâ paululùm ventricosiore, sub ore valde constrictâ.

Pedicellus etiam in hâc specie seabriusculus.

Annulus capsularum, annuente Hedwigio, duplice serie simbriatus, nec, ut in præcedente, ex ordine squamu-laruin simplici formatus.

serpens. 40. HYPNUM caule repente, foliis ovatis acuminatis integerrimis basi uninervibus, capsulâ cylindraccâ, operculo umbonato.

Hypnum serpens. Sp. Pl. 1596. Hedw. Sp.

Musc. 268. Crypt. v. 4. 45. t. 18. Hud. 507.

With. 865. Hull. 274. Lightf. 763.

Reh. 432. Sibth. 302. Abbot. 251. Swartz.

Musc.

Musc. Suec. 65. *Roth. Germ.* v. 3. 304.

Engl. Bot. t. 1037.

H. trichodes serpens, setis et capsulis longis erectis. *Dill. Musc.* 329. t. 42. f. 64. *Herb. Dill.*

H. repens trichoides terrestre minimum, capsulis majusculis oblongis erectis. *Raii Syn.* 85.

β. H. spinulosum. *Hedw. Sp. Musc.* 269. t. 69. f. 5—10.

Angl. Creeping White-veiled Feather-moss.

In umbrosis humidis, in truncis arborum, lignoque putrido, frequens.

Perennis. Fruct. fert imprimis Martio, Aprili.

Habitus duorum præcedentium. Caules prostrati, repentes, dense implexi, filiformes. Rami brevissimi, erecti, tenuissimi. Folia lætè viridia, exigua, laxè imbricata, erectiœcula, oyata, integerrima, concava, apice longius ac tenuius acuminata, basi, monente Bridelio, uninervia. Pedicelli copiosi, capillares, rubri, nitidi, apice pallidi. Capsula cylindracea, incurva, ore dilatata. Operculum plano-convexum, mucronulo brevi, subaduncō. Calyptre tenella, conico-cylindracea, alba, nitida, valdè conspicua.

β vix differt nisi pedicellis duplo circiter brevioribus, nervoque foliorum magis conspicuo, ultrà medium producto.

41. HYPNUM caule diffuso ramosissimo, foliis atro-virovatis concavis integerrimis carinatis, capsulârens. obovatâ, operculo subulato.

Hypnum atro-virens. *Dickf. Crypt. fasc.* 2. 10. *With.* 864. *Hull.* 274.

H. denticulatum exiguum pendulum atro-vires. *Dill. Musc.* 332. t. 43. f. 67.

Angl. Dark-green Round-branched Feather-moss.

Ad arbores in sylvis Scotiæ. *Dicks.*

Perennis. Fl. . . .

Caules diffusi, variè ramosissimi, undique foliosi, teretes. Folia fusco-viridia, opaca, undique imbricata, ovata, acuta, concava, integerrima, ad apicem usque uninervia et acutè carinata. Pedicelli unguiculares, atro-rufi. Perichætia albida, arctè imbricata, foliolis ovatis, concavis, subcarinatis. Capsula subcernua, ob-ovata, parva, fusca, ore dilatata. Operculum, ex Dilienio, subulatum, gracile, arcuatum.

filamen- 42. *HYPNUM caule ramosissimo implexo, foliis*
tosum. *ovatis integerrimis carinatis adpressis, capsulâ*
ovatâ, operculo conico.

Hypnum filamentosum. Dicks. Crypt. fasc. 2. 11.
With. 863. Hull. 273.

H. cæspitosum filamentosum minimum subfla-
vum. Dill. Musc. 282. t. 36. f. 18.

Muscus terrestris Virginianus minimus subfla-
vus. Moris. Hist. v. 3. 625. sect. 15. t. 5.
f. 13.

Angl. Thread-branched Yellow Feather-moss.

In sylvis Scotiæ. *Dicks.*

Perennis. Fl. . . .

Caules implexi, cæspitosi, ramosissimi, ramis elongatis, filiformibus. Folia undique arctè imbricata, adpressa, luteo-viridia, in fulvum vergentia, parva, ovata, acuta, concaviuscula, integerrima, carinata, uninervia. Pedicelli ferè unciales, capillares, fusi. Perichætia albida. Capsula cernua, ovata. Operculum conicum. Foliis cum præcedente arctè convenit, discrepat verò ramis elongatis filiformibus, capsulâ ovatâ, operculo conico, nec subulato.

tenel-
lum.

43. *HYPNUM caule repente ramosissimo, foliis*
subulatis enervibus integerrimis, capsulâ ova-
tâ, operculo subulato.

Hypnum

Hypnum tenellum. *Dicks. Crypt. fasc. 4. 16.*
t. 11. f. 12.

Angl. Tender Awl-leaved Feather-moss.

In truncis arborum.

Near Dover and Gravesend. *Mr. Dillwyn.* In Ireland. *Mr. Templeton.* On the under side of projecting stones and roots on Acton Burnell Hill, Shropshire. *Rev. E. Williams.*

Pereennis. Fruct. fert Februarij—Aprili.

Caules dense implexi, repentes, ramis brevibus, erectis, copiosis. Folia lætè at saturatè viridia, imbricata, erecta, subulata, acuta, tenuissima, integerrima, enervia, basi paululum dilatata. Pedicelli semiunguieulares, ramos superantes, erecti, apice demùm incurvi. Capsula pendula, ovata, brevis, obtusa, fusca. Operculum subulatum, longitudine ferè capsulae.

44. HYPNUM caule ramoso adscendente, foliis *albicans*.
 ovatis acuminatis trinervibus, capsulâ ovatâ,
 operculo conico mucronulato.

Hypnum albicans. *Necker. Musc. 180.* *Hedw.*
Sp. Musc. 251. *Crypt. v. 4. 13. t. 5.* *Dicks.*
Crypt. fasc. 1. 6. *With. 850.* *Hull. 269.*
Dicks. H. Sicc. fasc. 5. 21.

H. sericeum gracile albicans, capsulis subrotundis. *Dill. Musc. 328. t. 42. f. 63.*

H. terestre erectum humilius albicans, ramulis teretibus. *Dill. in Raii Syn. 83.*

Angl. Whitish Silky Feather-moss.

In arenosis et apricis.

Perennis. Fruct. fert Martio, Aprili, sed rariùs.

Caules adscendentes, unciales vel biunciales, divisi et ramosi, undique foliosi, ramis erecto-confertis, teretibus. Folia arctè imbricata, albo-virescentia, nitidula, ovata, acuminata, eoncava, integerrima, trinervia,

nervia, nervis apicem versùs plerumque evanescéntibus.
Pedicelli unciales, purpurei, propè basin ramorum.
Capsula ovata, brevis, ferruginea, arcuato-cernua.
Operculum conicum, mucronulo exiguo, acuto, nigro.

plumo- 45. **HYPNUM** caule repente, ramis confertis
sum. erectis, foliis ovatis acuminatis mediotenùs
uninervibus, operculo conico.

Hypnum plumosum. *Huds. ed. 1. 423. Sp.*
Pl. 1592. Wth. 852. Hull. 270. Sibib.
296. Abbot. 247. Hoffm. Germ. v. 2. 73.
Dicks. H. Sicc. fasc. 5. 22.

H. sericeum β . *Huds. 506.*

H. pseudo-plumosum. *Bridel. Musc.*

H. repens silicinum plumosum. *Dill. Musc. 280.*
t. 35. f. 16. Raii Syn. 86.

Angl. Rusty Silky Feather-moss.

Ad saxa, et arborum radices, tum in muris late-
ritiis.

Perennis. Fruct. fert hyeme et vere.

Caules prostrati, repentes, ferrugineo-tomentosi, ramis
numerosis, erectis, confertis, teretibus, subsimpli-
cibus. Folia imbricata, densa, ovata seu ovato-lan-
ceolata, acuminata, integerrima, concava, uninervia,
nervo vix medium attingente; inferiora ferruginea;
superiora luteo-viridia; omnia nitida. Pedicelli un-
guiculares, laeves, atro-rufi. Perichaetia apice patula.
Capsula inclinata, fusca, ovata, supra basin quodam-
modo constricta. Operculum conicum.

H. salebrosum *Hoffm. Germ. v. 2. 74,* *plumosum Hed-*
wigii, *distinctissima species est,* foliis lanceolatis tri-
nervibus laxè patentibus dilutè ac luteo-viridibus.

implex- 46. **HYPNUM** caule repente, ramis confertis
um. erectis, foliis lanceolatis acuminatis serrulatis
uninervibus, operculo conico.

Hypnum

Hypnum implexum. Swartz. *Act. Holm. ann.*,
1795. 264. *Turn. Musc. Hibern. MSS.*
H. plumosum. Swartz. *Musc. Suec.* 66.
Angl. Green Matted Feather-moss.

Ad saxa in montosis.

On a large stone near the foot of Ben Lawers, also
in Angus-shire. Mr. G. Donn. Near Dublin.
Dr. Scott and Dr. W. Stokes. Near Newcastle.
Mr. Winch.

Perennis. Fruct, fert vere.

Priore duplo tenuius. Caules repentes, ramis crebris,
erectis, plerumque simplicibus. Folia lætè viridia,
lutescentia, nitidiuscula, densè imbricata, patentius-
cula, lanceolata, concava, acuminata, e basi ad apicem
usque uninervia, summitatem versus, ex observatio-
nibus amicissimi Turneri, tenerrimè ferrulata. Pedi-
celli unguiculares, quandoque unciales, læves, rubri.
Perichaetium laxum, pallidum. Capsula ovata, cernua.
Operculum conicum.

47. *HYPNUM* caule procumbente ramosissimo, lutes-
foliis lanceolatis acuminatis trinervibus, pe- cens.
dicellis scabriusculis, operculo conico sub-
rostrato.

Hypnum lutescens. Huds. ed. 1. 421. Schreb.
Lips. 98. Hedw. Sp. Musc. 274. Crypt.
v. 4. 40. t. 16. With. 848. Hull. 269. Relh.
432. Sibth. 298. Abbot. 245. Dicks. H.
Sicc. fasc. 9. 21. Swartz. Musc. Suec. 65.
Roth. Germ. v. 3. 310. Hoffm. Germ. v. 2.
75.

H. sericeum γ. Huds. 506.

H. n. 1753. Hall. Hist. v. 3. 31.

H. sericeum, surculis longioribus et rectioribus,
capsulis

capsulis incurvis. *Dill. Musc.* 325. t. 42.
f. 60.

H. n. 23, varietas. *Raii Syn.* 84.

Neckera lutescens. *Willden. Berol.* 317.

Leskea longifolia. *Ehrb. Crypt.* 281.

Muscus cristam castrensem repræsentans, fla-
vescens, nemorosus, cassubicus. *Vaill. Paris.*
t. 27. f. 1; optimè monente Cel. Bridelio,
ex Herbarii Vaillantiani auctoritate.

Angl. Yellowish Feather-moss.

In ericetis et montosis siccis, præcipue calcareis.

Perennis. Fruct. fert Januario, Februario.

Caules palmares vel spithamæi, prostrati, ramosissimi,
divaricati, ramis teretibus, plus minus elongatis. Folia
lutescentia, nitida, imbricata, lanceolata, integerrima,
longius acuminata, trinervia, nervo centrali ad apicem
usque extenso, lateralibus medio evanescentibus. Pe-
dicelli ferè sesquiunciales, flexuosi, copiosi, rubri,
apicem versus tuberculis minutissimis scabri. Peri-
chætium coarctatum, foliolis interioribus apice capil-
laribus. Capsula subarcuato-crenua, ferruginea, ni-
tida. Operculum conicum, cum rostello brevi obtu-
siusculo.

Hujus affinitatem summam cum *H. salebroso Hoff-*
manni primus persentiit amicissimus D. Turnerus;
at me judice operculis, pedicellis, foliorumque nervis,
certè diserepant, nec confundenda sunt.

molle. 48. **HYPNUM** caule ramosissimo laxo filiformi,
foliis ovatis integerrimis concavis enervibus
acutiusculis, capsulâ ovatâ.

Hypnum molle. *Dicks. Crypt. fasc.* 2. 11. t. 5..
f. 8. *Hedw. Sp. Musc.* 273. t. 70. f. 7—10.

With. 862. *Hull.* 273.

Angl. Soft Water Feather-moss.

Ad rivulorum ripas in alpibus Scoticis. *Dicks.*

Perennis. Fl. . . .

Caules fasciculati, fluitantes, ramosissimi, teretes, flexuosi, foliosi, ramis obtusis. Folia luteo-viridia, parùm nitida, undique laxè imbricata, patentiuscula, ovata, concava, ventricosa, enervia, integrerrima, lævia; apice acutiuscula, mutica, complanata. Pedicelli rariores, haud unguiculares, incurvi. Perichaëtia e foliolis lanceolatis, acutis. Capsula cernua, ovato-subrotunda, ferruginea. Operculum adhuc ignotum.

Ex exemplaribus amicissimi Dicksoni, qui horum plurimorum mihi copiam fecit, descriptionem elaboravi.

49. HYPNUM caule pinnato, foliis ellipticis *purum.* ventricosis obtusis mucronulatis basi uninervibus, operculo conico adunco.

Hypnum purum. Sp. Pl. 1594. Hedw. Sp. Musc. 253. t. 66. f. 3—6. Hudsf. 504. With. 863. Hull. 273. Relb. 436. Sibth. 300. Abbot. 250. Curt. Lond. fasc. 3. t. 65. Fl. Dan. t. 706. f. 2. Roth. Germ. v. 3. 269. Swartz. Musc. Suec. 60.

H. elegans. Ehrb. Crypt. 242.

H. cypressiforme vulgare, foliis obtusis. Dill. Musc. 309. t. 40. f. 45.

H. terrestre erectum, ramulis teretibus, foliis inter rotunda et acuta medio modo se habentibus. Raii Syn. 81.

Muscus squamosus cypressiformis. Vaill. Paris. t. 28. f. 3.

Angl. Neat Meadow Feather-moss.

In pratis, pasuis, et nemorosis humidioribus, copiosè.

Perennis. Fruct. fert Novembre.

Caules

Caules erectiusculi, palmares, pinnato-ramosi, foliosi, ramis distichis, crebris, simplicibus, teretibus, acutis, patentibus vel arcuato-deflexis, undique foliosis. Folia pallidè sed pulcherumè viridia, mox lutescentia vel straminei coloris, pellucida, nitidissima, levissima, arctè imbricata, latè elliptica, integerriuma, ventricosa, apice obtusissima cum mucronulo plano, basi plus minus evidenter uninervia. Pedicelli biunciales, rubri: Perichaetium cylindraceum. Capsula cernua, ovata, arcuata, brevis. Operculum conicum, cum rostello adunco.

illecebrum. 50. **HYPNUM** caule vagè ramoso, ramis obtusis, foliis ellipticis ventricosis obtusis mucronulatis obfoletè uninervibus.

Hypnum illecebrium. *Hedw. Sp. Musc.* 252. t. 66. f. 1, 2. *Huds.* 504. *With.* 862. *Hull.* 273. *Sibth.* 300. *Schreb. Lips.* 95.

H. cupressiforme rotundius, vel *Illecebræ æmulum*. *Dill. Musc.* 311. t. 40. f. 46.

H. terreste erectum, ramulis teretibus, foliis subrotundis albo-virentibus cinetis. *Dill. Giss.* 220. *Raii Syn.* 81, sub n. 7.

H. n. 1744. *Hall. Hist.* v. 3. 29.

Muscus terrestris, surculis kali aut illecebræ æmulis, foliolis subrotundis, squamatim incumbentibus. *Raii Syn. ed.* 2. 37. *Vaill. Paris.* t. 25. f. 7.

Angl. Glasswort Feather-moss.

In ericetis, pascuis et nemorosis, rariùs.

In a wood at Heydon, Norfolk. *Rev. Mr. Bryant.*

Oxfordshire. *Sibth.* Near Eton. *Dr. Noebden.*

At Beddgelart, North Wales. *Mr. D. Turner.*

Shropshire. *Rev. E. Williams.*

Perennis. Fruct. apud nos haud profert.

A præ-

A præcedente vix differt nisi caulis brevioribus, inconditè ramosis, diffusis, ramulis obtusioribus, foliisque parùm latioribus. Variat foliis muticis.

51. **HYPNUM** caule pinnato compressiusculo, Schreberi. foliis ellipticis concavis mucronulatis enervi-
bus, operculo conico mucronulo recto.
Hypnum Schreberi. *Willden. Berolin.* 325.
Dicks. Crypt. fasc. 4. 18. *Roth. Germ. v. 3.*
270.
H. purum. *Ebrh. Crypt.* 311. *Huds.* 504, β.
With. 863, var. 2. *Hull.* 273, var. 2.
H. compressum. *Schreb. Lips.* 96. *Necker. Gallob.*
477. *Musc.* 159. *Hoffm. Germ. v. 2.* 58.
H. muticum. *Swartz. Musc. Suec.* 60.
H. cuspidatum β. *Weis. Crypt.* 253. *Web.*
Goett. 59.
H. parietinum. *Herb. Linn.* in operibus ejus
cum *H.* splendente nostro confusum.
H. n. 1742. *Hall. Hist. v. 3.* 28.
H. cypressiforme tenuius et compressius. *Dill.*
Musc. 312. t. 40. f. 47.
H. longum erectum, foliis angustis caulis appressis. *Raii Syn.* 83.
Muscus erectus, foliis angustis caulis appressis.
Raii Syn. ed. 2. 337. *Vaill. Paris.* t. 29. f. 10.
Angl. Schreberian Feather-moss.

In sylvis et paucis humidis, paludibusve.

In Anglesea. *Rev. H. Davies.* Glens and woody places in Ireland. *Mr. Templeton.* Near Yarmouth in very moist places. *Mr. D. Turner.*

Perennis. Fruct. fert hyeme et vere, rariūs.

H. puro simillimum, sed tenuius, ramis compressis, gracilioribus. Folia laxius imbricata, lanceolato-elliptica, adpressa, mucronulata, concava et obtusiuscula, sed longè

longè minus ventricosa, prorsus enervia. Pedicelli et capsulæ ferè H. puri, at operculum conicum, mucronulo breviore, recto; nec adunco.

nitens. 52. **HYPNUM** caule erecto, ramis abbreviatis, foliis lanceolatis acuminatis striatis, capsulâ oblongâ arcuato-cernuâ.

Hypnum nitens, Schreb. Lips. 92. Linn. fil. Meth. Musc. 34. Hedw. Sp. Musc. 255. Dicks. Crypt. fasc. 2. 12. H. Sicc. fasc. 1. 24. Willd. 852. Hull. 270. Swartz. Musc. Suec. 62. Roth. Germ. v. 3. 269. Hoffm. Germ. v. 2. 59.

H. trichodes. Neck. Musc. 161.

H. palustre erectum trichodes, ramulis crebris, luteo- et rufo-virentibus, glabris. Dill. Musc. 303. t. 39. f. 37.

Angl. Shining Yellow Feather-moss.

In paludosis.

Bogs in Scotland. Dicks. Near Acle, Norfolk.
Mr. D. Turner.

Perennis. Fruct. fert vere.

Caules erecti, palmares aut spithamei, foliosi, tomentosi, ramosi, ramis numerosis, sparsis, patentibus, brevibus, simplicibus. Folia, flava nitida, imbricata, lanceolata, acuminata, integerrima, striis tribus aut quinque post exsiccationem præcipue notata, nec, nisi omnia me fallunt, nervosa. Perichaetia subulata, foliolis plicato-striatis. Pedicelli capillares, biunciales, flexuosi, purpurei, læves. Capsula arcuato-cernua, oblonga, ferruginea. Operculum conicum, acutum, breve.

rufescens. 53. **HYPNUM** caule erecto, ramis elongatis, foliis lanceolatis acuminatis striatis, capsulâ ovatâ obliquâ.

Hypnum rufescens (ex errore refuscens). *Dicks.*

Crypt. fasc. 3. 9. t. 8. f. 4. With. 846. Hull. 269. Hoffm. Germ. v. 2. 59; sub n. 16.

H. caulinibus vix ramosis, foliis subulatis aristatis, capsulis gracilibus. *Hall. in Act. Helvet. v. 5. 12. n. 34.**

H. n. 1784. Hall. Hist. v. 3. 38; excluso syn. Dill.

Angl. Shining Reddish Feather-moss.

In rupibus uidis in alpibus Scoticis. *Dicks.*

IPerennis. Fruct: fert rarissimè:

(Caules erecti, sesquiunciales vel palinares, undique foliosi, teretiuseculi, ramosi, ramis erectiuseculis, elongatis, quandoque biuncialibus. Folia ferrugineo-lutescentia, nitida, imbricato-patentia, lanceolata, concava, integerrima, acuminata, interdum pilifera, enervia, siccitate verò, præcedentis more, longitudinaliter sulcata, ut multinervia videntur. Fructum non nisi in Herbario Linnæano vidi. Perichætium exiguum. Pedicellus ruber, uncialis, vel sesquiuncialis. Capsula suberecta, ovata, fusca, apice obliquè curvata.

54. *HYPNUM* catile pinnato, ramis cuspidatis, *cuspida-*
foliis oblongo-lanceolatis enervibus lœvibus, *tum.*
operculo conico acuto.

Hypnum cuspidatum. *Sp. Pl. 1595. Hedw.*
Sp. Musc. 254. Huds. 505. With. 862.
Hull. 273. Relb. 437. Sibth. 300. Abbot.
250. Dicks. H. Sicc. fasc. 7. 20. Swartz.
Musc. Suec. 61. Roth. Germ. v. 3. 271.
Hoffm. Gerin. v. 2. 58.

H. palustre, extremitatibus cuspidatis et pun-
gentibus. *Dill. Musc. 300. t. 39. f. 34.*

H. repens palustre, foliis triangularibus per cau-
les expansis, extremitatibus convolutis et acu-
minatis. *Raii Syn. 82.*

Angl. Pointed Bog Feather-moss.
In pascuis udis irriguis, et paludibus.

Perennis. Fruct. fert Martio, Aprili.

Caules erecti, longitudine varii, plerumque spithamæ; pinnatim ramosi, foliosi, ramis distichis, crebris, patentibus, brevibus, simplieibus, subcompressis; superioribus, e foliis convolutis, acutè cuspidatis. Folia lutescentia, nitida, imbricato-patentia, oblongo-lanceolata, acuta; caulinæ ovato-cordata; omnia integerrima, eoncava, lævia, enervia; summa arctè in cuspidem convoluta. Pedicelli 2—3-uneiales, rubri, subflexuosi. Perichætium subulatum, elongatum, strictum, foliolis striatis. Capsula ovato-oblonga, areuato-cernua, fulva. Operculum conicum, sæpius rectum, apice inuertonulatum.

***** *Caps. cernuis. Foliis squarrosis.*

cordifo- 55. *HYPNUM* caule erecto, ramis abbreviatis,
rim. foliis cordatis obtusiusculis uninervibus re-
motiusculis, operculo conico.

Hypnum cordifolium. *Hedw.* *Sp. Musc.* 254.
Crypt. v. 4. 97. t. 37. *Swärtz.* *Musc. Suec.*
62. *Roth.* *Germ.* v. 3. 319.

Angl. Heart-leaved Water Feather-moss.

In aquosis rariùs.

Near Newcastle; Mr. Winch. *Mr. D. Turner.*

Perennis. Fruct. fert vere.

Caules subæspitosi, erecti, debiles, foliosi, ramis paucis, sparsis, brevibus. Folia lætè viridia, nitida, interdùm lutescentia, remotiuscula, laxè patentia, cordato-ovata, amplexicaulia, integerrima, ad apieem usque uninervia, plus minus obtusa, basi pallida; superiora angustata; summa in cuspidem convoluta. Pedicelli (e specimini-
nibus exoticis) triunciales, capillares, flexuosi, rubri, basi atrati. Perichætium strictum, triplò quam in
præcedente minus. Capsula ovato-eylindracea, arcuato-cernua. Operculum conicum, acutum.

56. *HYPNUM*

§6. HYPNUM caule subpinnato, foliis lancicolatis *fluitans*.
acuminatis laxiusculis uninervibus, operculo
convexo mucronulato.

Hypnum fluitans. Linn. *Fl. Suec.* ed. 2. 399.

Hedw. *Sp. Musc.* 296. *Crypt.* v. 4. 94. t. 36.

With. 851. Hull. 270. Relb. 432. Sibth. 298.

Abbot. 245. Swartz. *Musc.* Suec. 57.

Roth. *Germ.* v. 3. 323. Hoffm. *Germ.* v. 2. 78.

Weis. *Crypt.* 226. Dickf. *H. Sicc. fasc.* 16. 22.

H. erectum aut *fluitans*, foliis oblongis perangustis acutis. Dill. *Musc.* 229. t. 38. f. 33.
Raii Syn. 82.

H. n. 1761. Hall. *Hist.* v. 3. 33.

Muscus fluitans, foliis et flagellis longis tenuibusque. Vaill. *Paris.* t. 33. f. 6.

Angl. Floating Feather-moss.

In aquis purioribus:

Perennis. Fruct. fert aestate, at rarissime.

Caules fluitantes, spithamæ vel pedales, filiformes, foliosi; subpinnati, rami sparsis, breviusculis, horizontalibus, apice deflexis. Folia dilutè fuscescentia, laxè imbricata, divaricato-patula, integerrima; basi ovato-lanceolata, concava, uninervia; apice subulata, tenuissimæ, elongata, enervia: inferiora sæpiù angustissimæ, scèrè setacea: superiora latiora, confertiora, falcato-secunda. Flores masculini numerosi, axillares, subglobosi. Pedicelli capillares, rufi; in Herb. Linn. triunciales; in Hort. Sicc. Dickfoni unciales. Perichaëtium oblongum, foliolis acuminatis. Capsula ovato-oblonga, brevis, arcuato-cernua. Operculum ex Hedwigio, convexo-conicum, mucronulatum.

§7. HYPNUM ramis vagis, foliis ovatis concavis *piliferum*, integerrimis piliferis inmediotenuis uninervibus, *rum.*
operculo subulato.

Hypnum piliferum. Schreb. *Lips.* 91. Hedw. *Sp.*

D d 2

Musc.

Musc. 275. *Crypt.* v. 4. 35. t. 14. *Swartz.*
Musc. Suec. 65. *Hoffm.* *Germ.* v. 2. 75.
Roth. *Germ.* v. 3. 312. *Turn.* *Musc.* *Hibern.*
MSS.

Angl. Hair-pointed Feather-moss.

In nemorosis udis.

In North Wales. *Rev. H. Davies.* Ireland. *Dr. Scott.*

Perennis. Fl. . . .

Caules procumbentes, biuneiales, palmares, spithamæi et ultra, vagè ramosi, ramis patulis, longitudine variis, teretibus, acutiusculis. Folia undique laxè imbrieata, pallidè viridia, nitidiuseula, ovata, obtusa, eoncava, integrerrima, lævia, apice pilifera, mediotenùs plerumque, (rariùs ad apicem usque) uninervia. Pedicelli rigiduli, rubri, sesquiunneales. Perichætium laxum, foliolis apice setaceis. Capsula ovata, ventricosa, brevis, eernua. Opereulum basi eonieum, mucrone filiformi, obtusiusculo, vix longitudine capfulæ. Hanc speciem cum *H. striato* eonfudit Cel. Dicksonus sub nomine *H. cassubici*. *Crypt. fasc.* 3. 10.

Rutabu- 58. *HYPNUM* ramis vagis, foliis ovatis acuminaliis ferrulatis mediotenùs uninervibus, operculo conico, pedicello scabro.

Hypnum rutabulum. Sp. *Pl.* 1590. *Hedw.*
Sp. *Musc.* 276. *Crypt.* v. 4. 29. t. 12. *Huds.*
 496. *With.* 851. *Hull.* 270. *Rehb.* 432.
Sibth. 297. *Abbot.* 245. *Swartz.* *Musc.*
Suec. 67. *Roth.* *Germ.* v. 3. 312. *Hoffm.*
Germ. v. 2. 77.

H. brevirostrum. *Ehrb.* *Crypt.* 85.

H. dentatum vulgatissimum, operculis obtusis.
Dill. *Musc.* 295. t. 38. f. 29.

H. repens triangularibus minoribus foliis. *Rau*
Syn. 80.

Muscus

Muscus myosuroides rutabulo fructu. *Vaill.*

Paris. t. 27. f. 8.

Angl. Common Rough-stalked Feather-moss.

Ubique in truncis arborum, dumetis, et ad sepes.

Perennis. Fruct. fert Martio.

Caules procumbentes, palmares aut spithamæi, vagè ramosi, elastici, ramis arcuatis, teretibus, apice attenuatis, undique foliosis. Folia pallidè viridia vel lutescentia, nitidiuscula, patula, ovata, basi lata, apice acuminata, oculo armato laxè ferrulata, ultrà medium uninervia. Pedicelli rubri, unciales vel sesquiunciales, undique tuberculis minutis scabri. Perichaetium laxum. Capsula ovata, arcuato-cernua. Operculum conicum, breve, obtusum cum exiguo mucrone.

59. **HYPNUM** ramis vagis, foliis ovato-deltoideis striatuminacuminatis subserrulatis uninervibus striatis, *tum.* operculo subulato.

Hypnum striatum. *Schreb. Lips. 91. Hedw.*

Sp. Musc. 275. Crypt. v. 4. 32. t. 13.

With. 850. Hull. 270. Sibth. 415. Swartz.

Musc. Suec. 60. Hoffm. Germ. v. 2. 75.

Roth. Germ. v. 3. 316.

H. longirostrum. *Ehrb. Crypt. 75.*

H. rutabulum γ. *Huds. 497. B. Relb. 433.*

H. vulgare dentatum, operculis cuspidatis. *Dill.*

Musc. 297. t. 38. f. 30. Herb. Dill.

H. repens triangularibus minoribus foliis, pediculis et capitulis brevioribus et tumidioribus, minus. *Raii Syn. 80.*

Angl. Common Striated Feather-moss.

In nemorosis uidis vel siccis, frequens.

Perennis. Fruct. fert Martio.

Species a præcedente, si quæ alia, distincta. Convenit cum eâ habitu et colore, sed plurimis notis discrepat.

Folia deltoidea, mediotenuis longitudinaliter multiplicatae, sulcatae. Pedicelli omnino laeves atque glaberrimi. Operculum basi convexum, acuminatum subulato, saepè arcuato, longitudine totius capsulae.

Varietatem ex Hibernia mecum communicavit amissimus D. Turnerus, cui folia manifestius ferrata, rami capsulaeque paululum elongati.

- flagellare.* 60. HYPNUM ramis fasciculatis, foliis deltoideis acuminatis serrulatis laevibus enervibus.

Hypnum flagellare. *Dicks. Crypt. fasc.* 2. 12. *exclusis synonymis. With.* 856. *Hull.* 271. *Angl.* Many-branched Water Feather-moss.

Ad saxa juxta rivulos in alpibus Scoticis. *Dicks.*

Perennis. Fl. . . .

Caules palmatae, ramosissimi; basi denudati, capillares, rigiduli. Rami numerosi, fasciculati, teretes, undique dense foliosi. Folia tristè viridia, sublutescentia, opaca, squarroso-patula, ovato-deltoides, acuminata, serrulata, prorsus non fallor encvia. E speciminibus Dicksonianis in Herbario D. Turneri, quibus fructus deest.

Diversum omnino ab *H. flagellari* *Hedw. Sp. Musc.* 282. t. 73; confer *H. alpinum*, n. 72.

- stellatum.* 61. HYPNUM caule erecto ramoso, foliis ovatis acuminatis integerrimis enervibus patentissimis, operculo conico.

Hypnum stellatum. *Schreb. Lips.* 92. *Hedw. Sp. Musc.* 280. *Dicks. Crypt. fasc.* 1. 5. t. 1. f. 7. *With.* 846. *Hull.* 268. *Rehb.* 437. *Sibth.* 301. *Abbot.* 246. *Swärtz. Musc. Suec.* 61. *Hoffm. Germ.* v. 2. 65. *Roth. Germ.* v. 3. 303. *Bridel. Musc.* t. 6. f. 2. *Dicks. H. Sicc. fasc.* 6. 23.

H. comata lutescente, extremitatibus stellatis. *Dill. Musc.* 302. t. 39. f. 35.

H. n.

H. n. 1732. *Hall. Hist. v. 3. 27.*

Muscus palustris, foliis et flagellis rigidiusculis,
seminibus in foliorum alis. *Vaill. Paris. t. 28.*
f. 10.

Angl. Yellow Starry Feather-moss.

In paludibus turfosis.

Perennis. Fruct. fert vere, haud frequentèr.

Caules erecti, plūs minus ramosi, ramis adscendentibus,
undique foliosis. Folia aureo-nitida, horizontaliter
patentia, basi latè ovata, integerima, lèvia, planius-
cula, enervia; apice longè ac tenuissimè acuminata;
inferiora ferruginea vel atrata. Pedicelli biunciales et
ultra, capillares, flexuosi, rubri. Perichaetia stricta,
foliolis lanceolatis, acuminatis, striatis. Capsula
ovato-cylindracea, arcuato-cernua. Operculum co-
nicum, acutum, breve.

62. HYPNUM caule ramoso subpinnato, foliis squarro-
ovatis acuminatis carinatis recurvis enervibus, *sum.*
operculo conico.

Hypnum squarrosum. *Sp. Pl. 1593. Hedw.*
Sp. Musc. 281. Hud. 502. With. 857.
Hull. 271. Relb. 435. Sibth. 299. Abbot.
249. Hoffm. Germ. v. 2. 65. Roth. Germ.
v. 3. 297.

H. repens, triangularibus reflexis foliis, majus.
Dill. Musc. 303. t. 39. f. 38. Raii Syn. 82.

G. H. repens, triangularibus reflexis foliis, minus.
Dill. Musc. 304. t. 39. f. 39. Raii Syn. 82.

H. squarrosum. *Swartz. Musc. Suec. 59.*
Ehrb. Crypt. 36.

H. n. 1733. Hall. Hist. v. 3. 27.

Muscus erectus, foliis reflexis. *Vaill. Paris.*
t. 27. f. 5.

Angl. Drooping-leaved Feather-moss.

In udis, nemorosis, et ad lēpes, vulgaris.

Perennis. Fruct. fert autumno et vere, sed rariūs.

Caulē adscendentēs, pinnatō-ramosī, ramis arcuatō-deflexis, undique foliosis, apice attenuatis. Folia luteo-viridia, densē imbricata, subquinquefaria, ovata, amplexicaulia, longē acuminata, integrērima, enervia, arcuato-deflexa, compresso-carinata, lāvia. Pedicelli unciales vel sesquiunciales, rubri, lāves. Capsula cernua, ovata, turgida, brevis. Operculūm conicūm, acutiusculum, breve.

β dignoscitur caule graciliore, ramis remotioribus, brevibus, foliis pallidioribus, vix lūtescentibus, operculo denique paulō producto.

loreum. 63. **HYPNUM** caule procumbente subpinnato, foliis subulatis striatis recurvis subsecundis, operculo convexo inueronulato.

Hypnum loreum. Sp. Pl. 1593. Hedw. Sp. Musc. 294. Huds. 502. With. 857. Hull. 272. Relb. 436. Sibth. 297. Dicks. H. Sicc. fasc. 7. 21. Ehrb. Crypt. 27. Swartz. Musc. Suec. 59. Hoffm. Germ. v. 2. 66. Roth. Germ. v. 3. 298.

H. lorcum montanum, capsulis subrotundis. Dill. Musc. 305. t. 39. f. 40.

H. repens, surculis magis erectis, foliis reflexis longioribus cinctis, operculo capituli magno. Rau Syn. 82.

Angl. Rambling Mountain Feather-moss.

In montosis siccioribus.

Perennis. Fruct. fert Martio, Aprili.

Præcedente duplō majus. Caulē diffusi, spithamæi aut pedales, pinnato-ramosissimi, ramis adscendentibus, teretibus, apice recurvis, acutis. Folia luteo-viridia, densē imbricata, arcuato-recurva, subsecunda; basi ovata, concava, integrērima, tri- aut quinque-sulcata, haud

haud nervosa; apice subulata, carinata, vix ferrulata. Pedicelli unciales vel sesquiunciales, purpurei, læves. Capsula cernua, ovato-subglobosa. Operculum convexum, rubicundum, mucronulatum.

64. HYPNUM caule adscendente subpinnato, *trique-*
foliis deltoideis acuminatis striatis undique trum,
patentibus, operculo conico.

Hypnum triquetrum. Sp. Pl. 1589. Hedw.

Sp. Musc. 256. Huds. 496. With. 850.

Hull. 270. Relb. 433. Sibth. 296. Abbot.

246. Swartz. Musc. Suec. 59. Hoffm. Germ.

v. 2. 59. Roth. Germ. v. 3. 267. Ehrb.

Crypt. 15. Dicks. H. Sicc. fasc. 16. 21.

H. vulgare triangulum, maximum et pallidum.

Dill. Musc. 293. t. 38. f. 28.

H. repens, triangularibus majoribus et pallidi-
oribus foliis. Raii Syn. 80.

Muscus squamosus major, sive vulgaris. Vail.

Paris. t. 28. f. 9.

Angl. Great Triangular Feather-moss.

In sylvis et ericetis, vulgaris.

Perennis. Fruct. fert hyeme et vere.

Caules erecti vel adscendentes, spithamæi aut pedales,
validi, ramosissimi, undique foliosi, ramis patentibus,
apice sæpè attenuatis et deflexis. Folia pallidè viridia,
nitidiuscula, conferta, undique squarroso-patentia, del-
toidea, multisulcata, obsoletè nervosa, plana nec cari-
nata, apice acuminata, minutè ferrulata. Pedicelli
biunciales, rubri, nitidi, læves: Perichaetium strictum,
foliolis apice setaceis, divaricatis. Capsula cernua,
ovata, brevis, ferruginea, demùm inæqualitè sulcata.
Operculum conicum, breve.

***** *Caps. cernuis. Foliis circinalibus.*

65. HYPNUM caule procumbente subpinnato, *rugo-*
foliis srtm.

foliis compresso-carinatis falcatis sulcatis uninervibus acuminatis, operculo conico.

Hypnum rugosum. *Linn. Mant.* 131. *Dicks.*
Crypt. fasc. 2. 12. *H. Sicc. fasc.* 1. 23.
With. 856. *Hull.* 271. *Swartz. Musc.*
Suec. 57.

H. lutescens crispum, lycopodii facie. *Dill.*
Musc. 289. *t. 37. f. 24.*

Angl. Yellow Wolf's-claw Feather-moss.

In paludibus Scotiæ.

Near Loch Rannoch, Perthshire. *Dicks.* Near
 Forfar. *Mr. G. Donn.*

Perennis. Fruct. fert vere.

Caules procumbentes, pinnatim-ramosi, undique foliosi,
 ramis eretis, obtusis, subaduncis. Folia luteo- vel
 fulvo-viridia, mox nigro-ferruginea, densè imbricata,
 faleato-secunda, lanceolata, acuminata, integerrima,
 compresso-carinata, ad apicem usque uninervia, sicei-
 tate longitudinaliter sulcata, nec transversè rugosa.
 Pedicelli atro-purpurei, biunciales, ereti. Capsula
 cylindracea, areuato-cernua, ore patula. Opereulum
 conium, acutum.

H. rugosum Hedwigii aliorumque ab hac specie prorsùs
 alienum est, folijsque gaudet transversè corrugatis.

scorpioides- 66. *HYPNUM* caule prostrato pinnato, foliis
 ventricosis falcatis adpressis encrvibus mu-
 ticis, operculo convexo mucronulato.

Hypnum scorpioides. *Sp. Pl.* 1592. *Hedw.*
Sp. Musc. 295. *Huds.* 501. *With.* 856.
Hull. 271. *Rehb.* 412. *Abbot.* 249. *Swartz.*
Musc. *Suec.* 58. *Hoffm.* *Germ.* v. 2. 63.
Dicks. *H. Sicc. fasc.* 3. 22. *Engl. Bot.*
t. 1039.

H. scorpioides palustre magnum, lycopodii in-
 star sparsum. *Dill. Musc.* 290. *t. 37. f. 25.*
Angl.

Angl. Scorpion Feather-moss.

In aquosis et paludibus turfosis.

Perennis. Fruct. fert hyeme et vere, sed rariūs.

Maxima hujus sectionis species. Caules prostrati, irregulariter pinnatim ramosi, undique foliosi, ramis medio incrassatis, apice aduncis. Folia aurco-viridia, mox fusca, nitida, arcte imbricata, ovata, concava, encervia, laevia, acutiuscula nec acuminata, superiora falcato-secunda. Pedicelli biunciales, rubri. Perichaetia subulata, stricta. Capsula arcuato-cernua, ovato-oblonga, fulva. Operculum convexum, mucronulatum.

Icon apud *Engl. Bot.* quamvis non optima, certe hujus speciei est.

67. HYPNUM caule erectiusculo pinnato, foliis revolutis subulatis compressis falcato-tortilibus, univens. nervibus striatis, operculo conico.

Hypnum revolvens. Swartz. *Musc. Suec.* 58.

et 101. t. 7. f. 14. *Turn. Musc. Hibern.* MSS.

Angl. Curling-leaved Feather-moss.

In udis Hiberniae; D. Scott. *Herb. Turn.*

Perennis. Fruct. fert. . . .

Priora duplo minus. Caules conferti, erectiusculi, vagè pinnati. Folia saepius fusco-atrata, rariūs fulva, nitidiuscula, arcte imbricata et conferta, secunda, lanceolato-subulata, acuminata, compresso-carinata, incurvato-tortilia, uninervia, siccitate striata. Pedicelli ultrà biunciales, rubri, flexuosi. Perichaetia parva, stricta. Capsula cylindracea, arcuato-cernua, ferruginea, subfusculata. Operculum conicum, breve, mucronulatum.

68. HYPNUM caule erecto pinnato, foliis ovatis aduncis acuminatis falcatis concavis uninervibus, operculo conico.

Hypnum aduncum. Sp. Pl. 1592. Hedw. Sp. Musc.

Musc. 295. *Crypt.* v. 4. 62. t. 24. *Huds.*
 501. *With.* 858. *Hull.* 272. *Lightf.* 753.
Relb. 430. *Abbot.* 248. *Swartz.* *Musc.*
Suec. 56. *Hoffm.* *Germ.* v. 2. 64. *Roth.*
Germ. v. 3. 289.

H. palustre erectum, summitatibus aduncis.
Dill. Musc. 292. t. 37. f. 26. *Raii Syn.* 82.
H. n. 1736. *Hall. Hist.* v. 3. 27.
Angl. Upright Hooked Feather-moss.

In uliginosis et aquosis, imprimis in montosis.

Perennis. Fruct. fert aestate.

Magnitudo fere praecedentis. Caules erecti, pinnati, altitudine variis, undique foliosi. Folia luteo-viridia, aurea, vel fulva, nitida, dense imbricata, secunda, ovata, acuminata, uninervia, concava, nec compresso-carinata, insigniter falcata, subtortilia, siccitate striata; summa arcte convoluta, adunca. Pedicelli bunciales, rubri. Perichaetia laxiuscula. Capsula priori simillima, ovato-cylindracea, arcuato-cernua, ferrugineo-fulva, laevis. Operculum convexo-conicum, breve, mucronulatum.

Varietatem laxiorem, magis elongatam, *H.* fluitans quodammodo referentem, e Sueciâ accepit optimus D. Turnerus.

uncina- 69. *HYPNUM* caule procumbente pinnato, folium subulatis falcatis uninervibus striatis, operculo conico, capsula constrictâ.

Hypnum uncinatum. *Hedw.* *Sp. Musc.* 289.
Crypt. v. 4. 65. t. 25. *Dicks.* *Crypt. fasc.*
 4. 19. *Swartz.* *Musc.* *Suec.* 56. *Hoffm.*
Germ. v. 2. 62. *Roth.* *Germ.* v. 3. 288.
Angl. Procumbent Hooked Feather-moss.

In udis montosis.

On stones by a rivulet a little to the north of
 Harrowgate,

Harrowgate, Yorkshire. *Mr. Teesdale.* Near Newcastle. *Mr. Winch.* By Lough Bray, Ireland. *Mr. D. Turner.*

Perennis. Fruct. fert autumno.

Præcedente parùm minus. Caules palmares, procumbentes, crebrè pinnati. Folia pallidè luteo viridia, nitidiuscula, densè imbricato-secunda, subulata, elongata, valdè falcata, etiam in circulum producta, compresso-carinata, uninervia, striata; summa convoluta, adunca. Pedicelli vix sesquiunciales, rubri. Perichætia subulata, elongata, foliolis setaceis. Capsula cernua, rufa, cylindracea, arcuata, gracillima, sub apice valdè constricta, ore patulo. Operculum conicum, acutum.

70. *HYPNUM caule repente, ramis erectis, foliis palustre,*
lanceolatis concavis enervibus secundis patulisque, operculo conico.

Hypnum palustre. Sp. Pl. 1593. Hud. 502.
With. 857. Hull. 272. Roth. Germ. v. 3.
295.

H. luridum. Hedw. Sp. Musc. 291. Crypt.
v. 4. 99. t. 38. Dicks. Crypt. fasc. 4. 18.
H. heterophyllum aquaticum polycephalum re-
pens. Dill. Musc. 293. t. 37. f. 27, A.
Herb. Dill.; nec B.

H. n. 1738. Hall. Hist. v. 3. 28.

Angl. Creeping Marsh Feather-moss.

In palustribus, et ad ligna putrescentia in aquosis.

Perennis. Fruct. fert æstate.

Caules repentes, tortuosi, implexi, unciales, vix biunciales, foliosi, sœpè denudati. Rami sparsi, erecti, simplices, unguiculares vel unciales. Folia tristè fusco-viridia, undique imbricata, in ramis masculinis (teste Hedwigio) erecto-patula, in fœmininis falcato-secunda, omnia ovata, concava, integerrima, plūs minùs acuta, enervia, lævia. Pedicelli unciales, rubri, nitidi.

nitidi. Perichætia subulata, pallida, foliolis obsoletè uninervibus apice acutis, patulis. Capsula ovato-cylindracea, brevis, arcuato-cernua, fulva. Operculum convexo-conicum, breve mucronulo acuto.

- fluvia-* 71. **HYPNUM** caule répente, ramis elongatis
tile. laxis, foliis ovatis concavis uninervibus fal-
catis, operculo convexo.

Hypnum fluviatile. Swartz. *Musc. Suec.* 63:
Hedw. Sp. Musc. 277. t. 71. f. 4. *Turn.*
Musc. Hibern. MSS.

H. luridum. Swartz. *Musc. Suec.* 58; ipso
auctore monente.

Angl. Creeping River Feather-moss.

Ad saxa in flaviis.

In Ireland. *Dr. Scott and Mr. R. Browne.*

Perennis. Fl. . . .

Caules prostrati, densè implexi, capillares, palmares aut spithamæi, ramosissimi, foliosi, demùm denudati, nigri. Rami subsimplices, adscendentes, elongati, laxi, undique foliosi. Folia imbricata, nitidiuscula, parva, ovata, concava, integerima, acuta, lævia, ad apicem usque uninervia; inferiora nigricantia; superiore pulchrè viridia; omnia magis minùsve falcato-secunda. Pedicelli unguiculares, carnei. Perichætia striata. Capsula cylindracea, demùm ovata, brevis, areuato-cernua. Operculum convexum, subconicum, mucronulo brevi, obtuso.

- alpinum.* 72. **HYPNUM** caule repente, ramis fasciculatis, foliis ovatis concavis acuminatis mediotenùs uninervibus: superioribus falcatis.

Hypnum flagellare. Hedw. *Sp. Musc.* 282:
t. 73. f. 1—3; exclusis synonymis.

Angl. Alpine-torrent Feather-moss.

Ad saxa in rivulis alpinis.

In the upper regions of Snowdon washed by
Llyn Fynnon Lâs. *Mr. D. Turner.* In Ireland.
Dr. Scott. In Scotland. *Mr. Dickson.*

Perennis. Fruct. fert Junio.

Caules palmares, ramis fasciculatis, densè foliosis, duplo
vel triplò quam in præcedente crassioribus, apice e fo-
liis falcatis et convolutis, subaduncis. Folia ovata,
sive ovato-lanceolata, acuminata, subaristata, concava,
integerrima, lævia, ad medium usque uninervia; cau-
lina ferrugineo-fusca, erecto-patula, vix secunda; ra-
mea pallida, luteo-viridia, nitida, falcato-secunda.
Pedicelli vix unciales, rubri. Perichætium laxum,
breve. Capsula ovata, brevis, subcernua. Operculum
conicum, medio constrictum.

Hujus exemplaria fructifera cum H. flagellari suo, cap-
sulis orbato, legit et confudit Cel. Dicksonus; hinc
proculdubio hallucinatus est Hedwigius, qui verum H.
flagellare nunquam vidit. Synonymon Dillenii de-
terminari non potest, ob exemplaris defectum in ejus
Herbario. Figura sanè, t. 39. f. 42, prorsùs aliena
videtur.

73. *HYPNUM* caule prostrato pinnato, foliis *cupressi-*
lanceolato-subulatis concavis enervibus fal-*forme*-
cato-secundis, operculo cuspidato.

Hypnum cupressiforme. Sp. Pl. 1592. *Hedw.*
Sp. Musc. 291. Crypt. v. 4. 59. t. 23. *Huds.*
500. *With.* 858. *Hull.* 272. *Rehb.* 435.
Sibth. 298. *Abbot.* 248. *Dicks.* *H. Sicc.*
fasc. 8. 21. *Swartz.* *Musc. Suec.* 57. *Hoffm.*
Germ. v. 2. 63. *Roth.* *Germ.* v. 3. 291.

H. crispum cupressiforme, foliis aduncis. *Dill.*
Musc. 287. t. 37. f. 23.

H. repens crispum cupressiforme. *Raii Syn.* 89.

H. n. 1737. *Hall. Hist.* v. 3. 27.

Neckera cupressiformis. *Willden. Berol.* 316.

Muscus squamosus ramosus minor et crispus.

Vaill. Paris. t. 27. f. 13.

β. Hypnum compressum. *Linn. Mant. 2. 310.*
Huds. 498.

H. filicinum sericeum, molle et pallidum, mucronibus aduncis. *Dill. Musc. 286. t. 36.*
f. 22.

Angl. Cypress-branched Feather-moss.

Ad radices arborum, rupes, vulgaris.

β in a wood at Tottenham. *Dill. Near Yarmouth. Mr. D. Turner.*

Perennis. Fruct. fert Aprili, Maio.

Caules prostrati, implexi, cæspitosi, pinnati, foliosi, ramis apice uncinatis, superne depresso-planis. Folia nitidiuscula, lætè viridia, sæpè lutescentia, quandoque fulva, arctè imbricata, subdisticha, terrain versus spectantia, falcato-secunda, ovato-lanceolata, concava, integrerima, enervia, apice setacea. Pedicelli sesquunciales, rubri, capillares, flexuosi. Perichaetium laxum. Capsula cylindracea, arcuato-cernua. Operculum convexum, longius cuspidatum, ferè subulatum.

β differt colore pallido, ramis tenuioribus, capsulâ subovatâ, erectiusculâ. Ab H. mollusco distinctissimum est.

dubium. 74. HYNUM caule prostrato pinnato, foliis ovatis acuminatis uninervibus secundis: superioribus falcatis, operculo conico.

Hypnum dubium. *Necker. Musc. 161. Dicks.*
Crypt. fasc. 3. 10. H. Sicc. fasc. 13. 23.

With. 854. Hull. 271. Hoffm. Germ. v. 2. 71.

H. filicinum β. *Huds. 498.*

H. repens filicinum trichodes palustre. *Dill. Musc. 286. t. 36. f. 21. Raii Syn. 85. n. 31, etiam n. 28.*

Leskeæ

Leskea incurvata. *Hedw. Sp. Musc.* 216. t. 53.

f. 8—14; optimè monente D. Turnero.

Angl. Fine Curve-leaved Feather-moss.

In palustribus. *Dicks.*

In a rivulet at Luttrell's town, Ireland. *Dr. Scott:*

Perennis. Fruct. fert Martio.

Caulis biunciales, prostrati, vagè pinnati, foliosi, ramis brevibus, subarcuatis. Folia luteo-fuscescentia, ovata, acuminata, integerrima, subrevoluta, nervo valido ad apicem usque producto; causina arctè imbricata, erecto-adpresca; ramca falcato-secunda, longius acuminata; omnia sèpè bistriata. Pedicelli sesquiunciales, flexuosi, purpurci. Perichætium albidum, ovatum, foliis striatis, et, ex auctoritate accuratissimi Turneri, sèpius denticulatis. Capsula ovato-cylindracea, ferruginea, demùm incurva, sub ore constricta. Operculum, ex Hedwigio et Dillenio, conicum, breve.

75. *HYPNUM* caule procumbente pectinato, *commun-*
foliis ovatis acuminatis tortilibus mediotenuis *tatum.*
uninervibus, operculo conico acuto.

Hypnum commutatum. *Hedw. Sp. Musc.* 284.

Crypt. v. 4. 68. t. 26. *Swartz. Musc. Suec.*

56. *Hoffm. Germ.* v. 2. 60. *Roth. Germ.*

v. 3. 284. *Turn. Musc. Hibern. MSS.*

H. repentis filicini crispi varietas B, C, D. *Dill.*

Musc. 283. t. 36. f. 19, B—D. *Herb. Dill.*

Angl. Larger Curled Fern Feather-moss.

In udis et palustribus, præcipue solo calcareo.

(On rocks by the river side near Bangor. *Mr. D.*

Turner. In Ireland. *Dr. Scott and Mr. Templeton.* Near Edinburgh, 1782.

Perennis. Fruct. fert æstatae.

Caulis palmates aut. spithamei, simplices vel divisi,

VOL. III.

E e

foliosi,

foliosi, ferrugineo-tomentosi, ramis adscendentibus, pectinato-pinnatis, ramiulis apice aduncis. Folia fulvo-viridia, parum nitida, parva, dense imbricato-patentia, secunda, vel sicca vel humeetata semper tortilia et crispa, ovata, integerrima, haud concava, longius acuminata, nervo centrali valido ultrà medium producto et in aeumine evanescente, cum pluribus exiguis brevibus ad basin ipsius folii dilatatam, ut in icona Hedwigianâ videri est. Pedicelli bi- vel triunciales, e ramis orti, atro-purpurei. Perichaëtia luteo-albida, ovato-subulata, foliolis interioribus acuminatis, striatis. Capsula arcuato-cernua, ovato-cylindracea, fusca, ore patulo. Operulum conicum, breve, cum exiguo mucrone.

A sequente dignoscitur foliis semper tortilibus. *D. Turner.*

*filici-
num.*

76. HYPNUM caule procumbente subpectinato foliis ovatis acuminatis uninervibus falcato-secundis, operculo convexo umbonato.

Hypnum filicinum. *Sp. Pl.* 1590. *Hedw. Sp. Musc.* 285. *t. 76. f. 5—10.* *Crypt.* v. 4. *t. 26. f. A, B.* *Huds.* 498. *With.* 854. *Hull.* 271. *Relh.* 433. *Sibth.* 294. *Abbot.* 247. *Swartz. Musc. Suec.* 55.

H. repens filicinum crispum. *Dill. Musc.* 282. *t. 36. f. 19, A, E, F.* *Raii Syn.* 85.

Angl. Lesser Fern Feather-moss.

In palustribus et rivulis montosis.

Perennis. Fruct. fert Maio.

Præcedente aliquantulum minus. Rami minùs exquisitè pectinati. Caulis densius tomentoso-ferrugineus. Folia aureo-vel fulvo-viridia, falcato-secunda, vix, etiam siecitate, crispa vel tortilia, nervo ad apicem usque producto. Pedicelli eaulini, sèpè approximati, biunciales, purpurei. Capsula cylindracea, arcuato-cernua. Operulum convexus, umbone brevi, reeto, acuto, nec subulatum.

77. HYPNUM

77. HYPNUM caule procumbente pectinato, foliis ovato-subulatis falcatis enervibus, capsula cum operatâ, operculo conico.

Hypnum molluscum. *Hedw. Sp. Musc.* 289.
Crypt. v. 4. 56. t. 22. *Swartz. Musc. Suec.*
55.

H. Crista-castrensis. *Huds.* 498. *With.* 854.
Hull. 271. *Lightf.* 749. *Relb.* 434. *Sibth.* 295.
Abbot. 247. *Dickf. H. Sicc. fasc.* 11. 20.

H. filicinum, cristam castrensem repræsentans.
Dill. Musc. 284. *t. 36. f. 20, A, vix B.* *Herb. Dill.*

H. n. 1767. *Hall. Hist. v. 3. 34.*

Muscus filicifolius luteus, folio crasso et undulato. *Raii Syn.* 86, *sub n. 32.*

M. terrestris repens subflavus, foliolis crispis minoribus, ramulisque densius confertis.
Moris. v. 3. 625. sect. 15. t. 5. f. 12. *Vaill.*
Paris. t. 27. f. 14.

Angl. Plumy Crested Feather-moss.

In montosis siccis calcareis.

Perennis. Fruct. fert Maio, Junio.

Pulcherrima species, proximis praecedentibus longè minor. Caules dense coacervati, pennæ instar plumosi, molles, ramosi, ramis pectinatis, ramulis incurvatis. Folia dilutè flavo-viridia, vix nitida, arctè imbricata, falcato-secunda, subulata, basi ovata, concava, prorsùs enervia. Pedicelli unciales, rubri, nitidi. Perichaetia flava, laxa, foliolis apice setaceis. Capsula cernua, ovata, brevis, turgida, ferruginea. Peristomium citrinum. Operculum conicum, acutissimum.

H. Crista-castrensis verum Linnæi atque Hedwigii,
Hall. n. 1768, a Dillenio cum hâc specie confusum, nondum me sciente in Britanniâ detectum fuit. Confusio haud extricabilis adest apud botanicos Germanicos de *H. filicino*, *mollusco*, *compresso* et *Crista-castrensi*.

455. FONTINALIS. G. Pl. 562. Juss. 11.
Hedw. Fund. v. 2. 96. t. 9. f. 53—55.

Caps. oblonga, lateralis, perichætio squamoso inclusa. Perist. exterius dentibus sedecim, basi dilatatis: interius reticulatum.

antipyretica. 1. FONTINALIS foliis complicato-carinatis trifariis acutis, perichætii foliolis obtusis, operculo subulato.

Fontinalis antipyretica. Sp. Pl. 1571. Hedw.
 Sp. Musc. 298. Huds. 467. Wiltb. 787.
 Hull. 275. Relb. 414. Sibth. 293. Abbot.
 231. Lightf. 694. Engl. Bot. t. 359. Roth.
 Germ. v. 3. 260. Swartz. Musc. Suec. 72.

F. triangularis major complicata, e foliorum alis capsulifera. Dill. Musc. 254. t. 33. f. 1.

F. major, foliis triangularibus complicatis, capitulis in foliorum alis sessilibus. Raii Syn. 79.

Hypnum n. 1794. Hall. Hist. v. 3. 40.

Muscus squamosus, foliis acutissimis, in aquis nascentes. Vaill. Paris. t. 33. f. 5.

Angl. Great Water-moss.

In fluviis lentè fluentibus, stagnisve.

Perennis. Fruct. fert æstate, sed rariùs.

Caules sinitantes, ramosi, spithamei vel pedales, foliosi. Folia atro-viridia, trifariam imbricata, ovato-lanceolata, acuta, integerrima, enervia, complicato-carinata. Capsulae laterales, solitariae, subsessiles, ellipticæ, perichaetio vestitæ, foliolis arcte imbricatis, obtusis, capsulam non superantibus. Operculum conico-subulatum, rectiusculum. Calyptra subulata. Peristomium exterius fuscum, deciduum; interius rubrum, pulcherrimè reticulatum.

squamosa. 2. FONTINALIS foliis imbricatis lanceolatis acuminatis

acuminatis concavis, perichætii foliolis operculisque obtusis.

Fontinalis squamosa. *Sp. Pl.* 1571. *Hedw.*

Sp. Musc. 299. *Crypt.* v. 3. 32. t. 12.

Huds. 467. *With.* 788. *Hull.* 275. *Lightf.*

696. *Dicks.* *H. Sicc. fasc.* 5. 25. *Roth.*

Germ. v. 3. 262. *Swartz. Musc. Suec.* 72.

F. squamosa tenuis sericea atro-virens. *Dill.*

Musc. 258. t. 33. f. 3.

Angl. Shining Scaly Water-moss.

In rivulis, præcipue alpinis.

In Wales, Scotland, and the north of England.

Dill. *Huds.* *Dicks.*

Perennis. Fruct. fert Augusto, Septembri.

Præcedente longè minor. Folia saturatè viridia, nitida,

arctè imbricata, plerumque trifaria, lanceolata, plus

minus angusta, concava, nec complicata, obsoletè ca-

rinata, enervia, apice subulata, nequaquam secunda.

Capsulæ laterales, elliptico-cylindraceæ. Perichætii

foliola latè ovata, obtusa, arctè imbricata, capsulâ bre-

viora. Operculum conicum, breve, obtusum. Ca-

lyptra subulata. Peristomium exterius fuscum, re-

curvum; interius atro-sanguineum.

Distinctissima a *F. antipyreticâ* foliis non complicatis,

vix carinatis, operculoque brevi, obtuso.

3. *FONTINALIS* foliis linear-setaceis patulis, *capilla-*
perichætii pédicellisque elongato-filiformi-*cea-*
bus: foliolis acutis capsulam superantibus.

Fontinalis capillacea. *Dicks.* *Crypt. fasc.* 2. 1.

With. 789. *Hull.* 275.

F. capillacea, calycibus stili instar cuspidatis.

Dill. Musc. 260. t. 33. f. 5.

Angl. Bristly Water-moss.

In rivulis alpinis Scottiæ. *Dicks.*

Perennis. Fl. . . .

Fusco-virens. Caulis palmaris vel spithamæus, pinnatim-ramosus. Folia imbricata, patentia, vix secunda, linearis-subulata, sive setacea, carinata. Capsulæ e ramorum axillis, longius pedicellatæ, ovatæ, breviusculæ. Perichaetium filiforme, contortum, capsulam superans, foliolis lanceolatis, acutis, interioribus majoribus et obtusioribus. E specimine ab amicissimo Dicksono mihi dono dato et cum Dillenianis collato.

Differt a *F. falcata* Swartzii et Hedwigii, quæ *F. capillacea* Linn. Fl. Suec., Ehrh. Crypt. 205, foliis laxioribus, haud falcato-secundis, pedicelloque cum capsulâ perichaetii foliolis superato, nec a medio ad apicem nudo, exerto.

456. FUNARIA. Schreb. Gen. 760. Hedw. Sp. Musc. 172.

Koelreutera. Hedw. Fund. v. 2. 95. t. 10.
f. 58—60. v. 1. t. 5. f. 21—26.

Caps. obovata. *Perist.* exterius dentibus sedecim, obliquis, apice cohærentibus: interius dentibus sedecim, planis. *Flores* terminales.
Calyptra quadrangula.

hygro-
metrica. 1. FUNARIA foliis concavis, capsula ventricosâ. Funaria hygrometrica. Hedw. Sp. Musc. 172.

Roth. Germ. v. 3. 225. Sibth. 288.

Mnium hygrometricum. Sp. Pl. 1575. With.
804. Relb. 417. Abbot. 233. Lightf. 709.
Ehrb. Crypt. 46. Fl. Dan. t. 648. f. 2.
Hoffm. Germ. v. 2. 40.

Bryum hygrometricum. Huds. 488. Hull. 265.
Engl. Bot. t. 342.

B. buibiforme aureum, calyprâ quadrangulari,
capsulis piriformibus nutantibus. Dill. Musc.
407. t. 52. f. 75.

B. aurcum,

B. aureum, capitulis reflexis piriformibus, calyptrâ quadrangulari, foliis in bulbi formam congestis. *Raii Syn.* 101.

Muscus foliis scutellatis, capitulo lypiformi nutante. *Vaill. Paris.* 135. t. 26. f. 16.
Angl. Twisting Cord-moss.

In ericetis arenosis humidiusculis, hortis et ambulacris neglectis, frequens.

Annua. Fructum fert Aprili, Maio.

Radix ferrugineo-tomentosa. Caules cæspitosi, brevissimi, ferè nulli. Folia congesta, sessilia, ovata, acuta, concava, ventricosa, integerrima, uninervia, inflexa, dilutè viridia, pellucida, sæpiùs matutino rore intùs detento pulchrè micantia, oculo armato reticulata. Flores dioici. Pedicellus erectus, altitudine maximè varians, sèpè biuncialis, flavus, basi purpureus, siccitate multiplicitè tortilis, apice incurvus. Capsula luteo-ferruginea, nutans, obovata, ventricosa, incutivâ, sulcata, fragilis, ore cœarctato. Operculum convexum, submuticum, margine coccineo. Peristomium exteriorius fulvum, dentibus obliquè tortis, conniventibus: interiorius dentibus rectis, transversis, planis. Calyptre albida, pellucida, apice subulata, basi plerumque quadrangula, subventricosa, mox, tuimente fructu, obliquè flexa.

Aliam speciem ex Hispaniolâ habemus, foliis arctè vaginantibus, capsulâ basi angustatâ.

457. BARTRAMIA. *Hedw. Crypt.* v. 2. 111.
Schreb. Gen. 761. *Engl. Bot. t.* 997.

Caps. sphærica, mox sulcata. Perist. exteriorius dentibus sedecim, basi dilatatis: interiorius membranaceum, plicatum, variè laciniatum. Calyptre lævis. Operculum depresso.

1. BARTRAMIA pedicellis lateralibus arcuatis *Halleri-*
E e 4 *folio ana.*

folio brevioribus, foliis linearis-subulatis uninervibus laevibus.

Bartramia Halleriana. *Hedw. Sp. Musc.* 164.

Crypt. v. 2. 111. t. 40. *Swartz. Musc.*

Suec. 73. *Roth. Germ. v. 3. 259.* *Engl.*

Bot. t. 997.

Bryum laterale. *Huds. 483.* *Lightf. 727.*

With. 823. *Hull. 254.* *Dicks. H. 3ic.*

fusc. 5. 20. *Ebrh. Crypt. 33.*

B. recurvum. *Wulf. in Jacq. Coll. v. 2. 224.*

B. n. 1802. *Hall. Hist. v. 3. 43. t. 46. f. 8.*

B. capillaceum, crebris per caulem capitulis.

Hall. It. Helvet. sect. 55. t. 1.

B. pomiforme, var. 1. *Scop. Carn. v. 2. 323.*

Mnium laterale. *Hoffm. Germ. v. 2. 54.*

Angl. Lateral Bartramia.

Ad rivulos in umbrosis alpinis.

By the sides of rivulets about Hart-fell near

Moffat. *Rev. Dr. Walker.* North of England.

Mr. Dickson. North Wales. *Mr. Griffith.*

Perennis. Fruct. fert Julio.

Caules cæspitosi, ercti, bi- vel triunciales, ramosi, foliosi, ferrugineo-tomentosi. Folia patula, densa, lætè viridia, subsecunda, mollia, tenuissima, linearis-subulata, carinata, uninervia, laevia nec sulcata; apicem versùs serrulata; basi dilatata. Pedicelli axillares, solitarii, incurvati, foliis breviores. Capsula globofa, mox paululum angustata, et longitudinaliter sulcata, sulcis sedecim, subcernua, ferruginca. Peristomium ferrugineum: interius plicatum, ferè integrum, ore tantummodo erosum. Operculum convexum, breve, Calyptra conica, decidua.

pomifor- 2. BARTRAMIA pedicellis erectis caulem superantibus, foliis subulatis uninervibus.

Bartramia

- Bartramia pomiformis. *Hedw. Sp. Musc.* 164.
Swartz. Musc. Suec. 73. *Roth. Germ.* v. 3.
 258. *Engl. Bot. t.* 998.
- Bryum pomiforme. *Sp. Pl.* 1580. *Huds.* 474.
Wittb. 822. *Hull.* 259. *Rehb.* 421. *Abbot.*
 240. *Dicks. H. Sicc. fasc.* 10. 20. *Ehrb.*
Crypt. 23. var. 2. *Scop. Carn.* v. 2. 323.
- B. capillaceum, capsulis sphæricis. *Dill. Musc.*
 339. t. 44. f. 1.
- B. trichoides virens, erectis majusculis ca-
 pitulis maliformibus. *Raii Syn.* 97.
- B. n. 1803. *Hall. Hist.* v. 3. 43. t. 46. f. 7.
- Muscus trichoides minimus sericeus capilla-
 ceus, capitulis sphæricis. *Morif.* v. 3. 628.
sect. 15. t. 6. f. 6. *Vaill. Paris.* 129. t. 24.
 f. 9.
- Angl.* Apple Bartramia.

In umbrosis et rupium fissuris.

Perennis. Fruct. fert Aprili, Maio.

(Caulēs densē cæspitosi, ramosi, fastigiati, foliosi, altitu-
 dine varii, sœpius unciales, infernè tomentoso-ferru-
 ginei. Folia pulchrè viridia, patentia, linearis-subulata,
 nervo valido, apice ferrulata, basi minus dilatata, sic-
 citate striata. Pedicelli solitarii, terminales, mox la-
 terales, caulem longè superantes, unciales vel sesqui-
 unciales, erecti, rubri. Capsula erecta, globosa, virens,
 demum fusca, arcuato-cernua, longitudinaliter sulcata.
 Peristomium breve; exterius rubrum; interius tenuis-
 simum. Operculum convexum, brevissimum. Ca-
 lyptra conico-subulata, decidua.

3. BARTRAMIA pedicellis erectis caulem su-*gracilis*.
 perantibus, foliis lanceolatis revolutis uniner-
 vibus, caule elongato.

Bartramia gracilis. *Flörk. in Schrad. Journ.*
Turn. Musc. Hibern. MSS. cum iconie.

B. Oede-

B. Oederiana. *Swartz. MSS. Mohr. Ind.* 5.
Bryum Oederi. Retz. Prod. 261. *Fl. Dan.*
t. 478.
Angl. Tall Slender Bartramia.

Iu montosis.

In Scotland. *Mr. Dickson.* 1788. Ireland.
Dr. Scott.

Perennis. Fl. . . .

Præcedenti arctè affinis, differt verò caulis altioribus,
 ferè palmaribus, gracilibus, parùm ramosis; foliis latioribus,
 lanceolatis; capsulâ minore, evidenter ar-
 cuato-cernuâ. Folia reverà lanceolata sunt, acuminata,
 margine revoluta, apicem versus ferrata.

fontana. 4. BARTRAMIA pedicellis erectis caulem su-
 perantibus, foliis ovatis integerrimis, ramis
 fasciculatis erectis filiformibus.

Mnium fontanum. *Sp. Pl.* 1574. *Hedw. Sp.*
Musc. 195. *With.* 799. *Lightf.* 708.
 ~ *Reh.* 419. *Abbot.* 232. *Hoffm. Germ.* v. 2,
 54. *Fl. Dan.* t. 298.

Bryum fontanum. *Huds.* 475. *Sibth.* 289.
Swartz. Musc. Suec. 48. *Rotb. Germ.* v. 3,
 234. *Engl. Bot.* t. 390.

B. palustre, scapis teretibus stellatis, capsulis
 magnis subrotundis. *Dill. Musc.* 340. t. 44.
 f. 2.

Muscus palustris, Adianto aurco affinis, scapis
 tenuibus, foliolis brevibus. *Raii Syn.* 98.

M. trichoides palustris erectus, coronatus capi-
 tulis sphæricis amplioribus. *Moris. v.* 3. 628.
 sect. 15. t. 6. f. 8.

Angl. Fountain Bartramia.

Ad rivulos in paludibus spongiosis copiosè.

Perennis.

Perennis. Fruct. fert Junio, Julio.

Caules densè cæspitosi, sæpè spithamæi, determinatè ramosi, foliosi, infernè tonientofo-ferruginei; ramis fasciculatis, erectis, fastigiatis, filiformibus. Folia luteo-viridia, imbricata, haud secunda, brevia, latè ovata, integrerrima, acuminata, carinata, uninervia. Flores dioici, terminales: masculi disciformes, foliis majoribus, rainulisque novellis, circumvallati. Pedicelli solitarii, erecti, triunciales et ultrà, caulem longè superantes, rubri. Capsula obliquè cernua, ovato-sphærica, fusca, multisulca. Operculum convexum, acuminé brevi, acuto, adunco. Peristomium breve, fuscum; interius duplicatò fimbriatum.

5. BARTRAMIA pedicellis recurvatis, foliis lan-*arcuata*.
ceolatis uninervibus sulcatis ferrulatis, ramis
sparsis patentibus.

Bartramia arcuata. *Engl. Bot.* t. 1237.

Mnium arcuatum. *Dicks. Crypt. fasc.* 3. 2.
t. 7. f. 3. *H. Sicc. fasc.* 6. 19. *With.* 803.

M. chrysocomum. *Hedw. Sp. Musc.* 74.

Hypnum chrysocomum. *Dicks. Crypt. fasc.* 2.
12.

H. palustre erectum, comâ luteâ, basi nigricante. *Dill. Musc.* 302. t. 39. f. 36. *Raii Syn.* 83.

Bryum arcuatum. *Hull.* 255.

Angl. Curve stalked Bartramia.

In paludibus subalpinis rariùs.

In the north of Yorkshire. *Richardson.* Scotland.

Dicks. Mount Glyder, Wales. *Dill.* Yorkshire and Cheshire. *Mr. Bradbury.* Near Llanberis. *Mr. Griffith* and *Mr. D. Turner.*

Perennis. Fruct. fert autumno, sed rariùs.

Caules erecti, ramosi, foliosi, dense ferrugineo-tomentosi, ramis sparsis, numerosis, patentibus. Folia imbricato-patula, luteo-viridia, nitida, lanceolata, acuminata,

nata, uninervia, fulcata, certè ferrulata. Pedicelli solitarii, terminales, mox laterales, breves, arcuati, rufi, perichaetio parvo, squamoso. Capsula cernua, globofa, virens, nitida, sulcata. Operculum convexisulcum, obtuse mucronulatum, coccineum. Peristomium exterius fulvum, breve, dentibus obtusis: interius membranaceum, album, laciniis sedecim, exterioris dentibus conforme.

Flores masculi terminales, stellati. *Dicks.*

458. MNIUM. *G. Pl. 563. Juss. 11.*

Cypf. cylindracea, mox suleata. *Perist.* *exterius* dentibus sedecim, basi dilatatis: *interius* membranaceum, laciniatum. *Calyptra* lævis. *Pedicellus* terminalis.

andro- 1. *MNIUM monoicum*, capsula rectâ, operculo conico, foliis undique imbricato-patulis apice denticulatis.

Mnium androgynum. *Sp. Pl. 1574. Huds. 472.*
Hoffm. Germ. v. 2. 46. Fl. Dan. t. 299.
Hedw. Theor. 149. t. 12. f. 48—50. Ebrh.
Crypt. 63. Schmidel. Ic. 216. t. 56. f. 1.
Engl. Bot. t. 1238.

M: perangustis et brevibus foliis. *Dill. Musc.*
230. t. 31. f. 1. Raii Syn. 78.

M. n. 1854. *Hall. Hist. v. 3. 56.*

Bryum androgynum. *Hedw. Sp. Musc. 178.*
With. 836. Hull. 262. Abbot. 238. Roth.

Germ. v. 3. 227. Swartz. Musc. Suec. 46.

Muscus capillaceus minimus, capitulo minimo pulvрulento. *Vaill. Paris. t. 29. f. 6.*

Angl. Narrow-leaved Spring-moss.

In uliginosis et umbrosis humidis.

Perennis. Fl. Martio: fructum fert, sed rarissime,
 Aprili, Maio.

Caulis

(Caules densè cæspitosi, erecti, unciales vel biunciales, ramosi, foliosi, ferrugineo-tomentosi. Folia luteo-viridia, pallida, pellucida, punctata, undique imbricata, erecto-patentia, lanceolata, acuta, subrevoluta, carinata, uninervia, apice denticulata; siccitate inflexa, parùm undulosa. Flores monoici, terminales; masculi pedunculati, globosi, nudi: fœniuei sessiles. Pedicellus fructûs uncialis vel ultrà, solitarius, capillaris, nitidus, roseus, apice flavescens. Capsula creticiuscula, cylindracea, gracilis, recta, mox longitudinaliter sulcata, ferruginea, ore patentiusculo. Operculum conicum, breve, obtusum, subaduncum, striatum. Peristomium exterius flavum, dentibus apice setaceis, longissimis, erectis, siccitate involutis: interius tenuissimum, membranaceum, album, laciñiis capillaribus.

2. *MNIUM* monoicum? capsulâ obovatâ rectâ, *conoide-*
operculo subulato, foliis undique imbricato-um.
patulis integerrimis.

Mnium conoideum. *Engl. Bot.* t. 1239.

Bryum conoideum. *Dicks. Crypt. fusc.* 4. 9.
t. 11. f. 2.

Angl. Club-fruited Spring-moss.

In truncis arborum Scotiæ et Hiberniæ.

(On the trunks of Beeches near Inverary. *Dickson.*

On pear trees at Orange grove near Belfast.

Mr. Templeton.

Perennis. Fl. . . .

Planta exigua. Caules subsimplices, densè foliosi, erecti. Folia tristè virentia, præcedentis inore pellucido-punctata, undique imbricato-patula, ovato-lanceolata, carinata, integerrima, obsoletè mucronulata. Pedicellus terminalis, brevis, rectus, flavescens. Capsula creticiuscula, obovata, obtusa, pallidè fusca, sulcata. Operculum subulatum, acutum, longitudine ferè capsulæ, sæpè curvatum. Peristomium exterius breviusculum: interius, ex D. Templeton, tenuissimè laciniatum.

3. *MNIUM*

palustre. 3. **MNIUM** dioicum, capsulâ obliquâ, operculo conico, foliis acutis: superioribus arcuatosecundis, caule erecto.

Mnium palustre. *Sp. Pl.* 1574. *Hedw. Sp. Musc.* 188. *Huds.* 472. *With.* 802. *Lightf.* 708. *Rehb.* 419. *Abbot.* 233. *Hoffm. Germ.* v. 2. 47. *Ehrh. Crypt.* 55. *Schmidel. Ic.* 218. *t. 56. f. 2.*

M. majus, ramis longioribus bifurcatis. *Dill. Musc.* 233. *t. 31. f. 3.* *Raii Syn.* 78.

M. n. 1855. *Hall. Hist.* v. 3. 56.

Bryum palustre. *Roth. Germ.* v. 3. 233. *Swartz. Musc. Suec.* 46. *Engl. Bot.* t. 391.

B. ventricosum. *Sibth.* 289, excluso syn. *Dicksoni*.

Angl. Greater Forked Spring-moss.

In palustribus spongiosis et irriguis.

Perennis. Fl. Aprili, Maio: fructum fert Junio, Julio.

Planta palmaris. Caules erecti, dichotomi, undique foliosi, ferruginco-tomentosi. Folia luteo ac pallidè viridia, imbricata, lanceolata, acuta, integerrima, canaliculata, carinata, uninervia, siccitate undulosa; superiora, quandoque omnia, falcato-secunda. Flores terminales, dioici: masculi disciformes, foliis majoribus patentibus circumvallati, at sàpè proliferi, pedicellos exferentes nudos, capitellatos, primæ speciei simillimos: fœminei tessiles. Pedicellus solitarius e dichotomiâ caulis, erectus, biuncialis, subflexuosus, pulcherrimè ruber, apice luteolus. Capsula subcernua, elliptico-cylindracea, obliquè incurvata, sulcata, fusca. Operculum conicum, obtusum, striatum, subaduncum, basi rubrum. Peristomium exterius sulphureum, dentibus erectis, apice setaceis: interius album, laci-niis capillaribus.

459. BRYUM. *G. Pl.* 564. *Juss.* 11. *Schreb.*
Gen. 761. *Hedw. Fund.* v. 2. 94.
 Webera. *Hedw. Fund.* v. 2. 95.
 Pohlia. *Schreb. Gen.* 760. *Hedw. Crypt.* v. 1.
 96.
 Meesia. *Schreb. Gen.* 761. *Hedw. Fund.* v. 2.
 97. t. 9. f. 56, 57.

Caps. ovato-oblonga, lævis. *Perist.* exterius dentibus sedecim, basi dilatatis: interius membranaceum, variè dentatum et laciniatum. *Calyptra* lævis. *Pedicellus* terminalis.

- II. BRYUM caule subsimplici, foliis lanceolatis *nutans*. acutis carinatis, capsulâ obovatâ penduiâ retusâ.

Bryum nutans. *Schreb. Lips.* 81. *Swartz.*
Musc. Suec. 46. *Röth. Germ.* v. 3. 252.
Ebrh. Crypt. 25. *Engl. Bot. t.* 1240.

B. sericeum. *With.* 839. *Hull.* 265. *Sibth.* 292.

B. trichodes lætè virens capitulis cernuis oblongis. *Dill. Musc.* 391. t. 50. f. 61. *Raii Syn.* 100.

Mnium nutans. *Hoffm. Germ.* v. 2. 49.

Webera nutans. *Hedw. Sp. Musc.* 168. *Crypt.* v. 1. 9. t. 4.

Angl. Silky Pendulous Thread-moss.

In ericetis montosis.

Near Woolwich. *Dill.* In Apsley wood, Bedfordshire. *Rev. Dr. Abbot.* On Cromford moor near Matlock, Derbyshire.

Perennis. Fruct. fert Julio, Augusto.

Caules cæspitosi, erecti, breves, sæpiùs simplices, apice foliosi. Folia pallidè viridia, nitida, lancicolata, acuta, carinata, uninervia, apice ferrulata. Pedicellus solitarius, terminalis, erectus, longissimus, bi-triuncialis, nitidus, purpureo-fulvus, apice pallidior. Capsula pendula, obovata, retusa, basi parùm attenuata vel elongata, luteo-ferruginea, opaca nec nitida. Operculum convexum, umbonatum. Peristomium flavescens, breve: interius varie laciniatum.

Flores hermaphroditici. *Hedw.*

aureum. 2. BRYUM caule simplicissimo, foliis setaceis: superioribus longissimis, capsulâ pendulâ pyriformi medio constrictâ.

Bryum aureum. *Schreb. Lips.* 81. *Huds.* 487. *Hull.* 265. *Dicks. H. Sicc. fasc.* 8. 19. *Engl. Bot. t.* 389.

B. mnioides. *With.* 838.

B. pyriforme. *Swartz. Musc. Suec.* 45. *Roth. Germ. v.* 3. 251.

B. pendulum. *Ehrb. Crypt.* 165.

B. trichodes aureum, capsulis pyriformibus nutantibus. *Dill. Musc.* 391. *t.* 50. *f.* 60. *Deering. Nottingh. append.* 7.

B. n. 1813. *Hall. Hist. v.* 3. 46. *t.* 45. *f.* 7.

Mnium pyriforme. *Sp. Pl.* 1576. *Hoffm. Germ. v.* 2. 50.

Webera pyriformis. *Hedw. Sp. Musc.* 169. *Crypt. v.* 1. 5. *t.* 3.

Angl. Golden Thread-moss.

In rupibus calcareis umbrosis, et speluncis siccis.

On the roofs of caverns in the rocks of Nottingham park. *Deering.* In Surrey. *Mr. Crowe* and *Mr. Dickson.* North Wales. *Rev. Mr. Wood.* On the inside of the frames in Messrs. Lee and Kennedy's garden at Hammersmith.

Annu.

Annua. Fruct. fert Maio—Julio.

Caules cæspitosi, crecti, simplices, semiunciales, filiformes. Folia luteo- ac pallidè-viridia, capillaria, laxa, carinata, integerrima; summa longissima, stellatim patentia. Flores hermaphroditici, terminales. Pedicellus solitarius, erectus, uncialis, capillaris, aureus, nitidus, subflexuosus. Capsula pulcherrima, aureofusca, nitidissima, pendula, pyriformis, subretusa, a basi ad medium usque attenuata. Operculum subconicum, breve, mucronulatum. Calyptra subulata, alba. Peristomium exterius flavum: interius album, tenuissimum, dupli scie dentatum.

3. BRYUM caule subsimplici, foliis linear-lan-*elongato-*
ceolatis margine incrassatis, capsulâ subcer-tum.
 nuâ cylindraceâ basi elongatâ.

Bryum elongatum. *Dicks. Crypt. fasc. 2. 8.*
With. 832. Hull. 263.

Pohlia elongata. *Hedw. Sp. Musc. 171. Crypt.*
v. 1. 96. t. 36. Swartz. Musc. Suec. 44.
Roth. Germ. v. 3. 224. Ehrb. Crypt. 64.

Mnium Pohlia. *Hoffm. Germ. v. 2. 48.*
Angl. Elongated Thread-moss.

In sylvis et rupibus alpinis.

On Clagwyn y garnedd, North Wales. *Herb.*
Dill. Near Loch Ness. *Dicks.* North Wales.
Mr. Griffith. At Luggy Lough, Ireland.
Dr. W. Stokes.

Perennis. Fruct. fert Junio.

Caulis erectus, brevis, simplex, demùm subramosus.
 Folia saturatè viridia, patula, lanceolata, vel linear-lanceolata, acuta, uninervia, margine paululùm incrassata, integerrima; siccatè flexuosa. Pedicellus sesquiuncialis, solitarius, erectiusculus, purpurcus, apice flavescens. Capsula subcernua, nec pendula, cylindracea, arcuata, fusca, ore subcoartata, basi in apophyse obconicam, propriæ ferè longitudinis, elongata.

Operculum conicum, acutum, breve. Peristomium exterius breviusculum, fuscum; interius album, simplici serie dentatum.

Hanc speciem cum Bryo nutante confudit Dillenius.

- trichodes.* 4. **BRYUM** caule subsimplici, foliis linearibus obtusis, capsulâ obovatâ incurvâ subcernuâ, pedicello longissimo.

Bryum trichodes. *Sp. Pl.* 1585. *Dicks. Crypt. fasc.* 2. 7. *H. Sicc. fasc.* 3. 21. *Hull.* 263.

B. trichodes aurcum, capsulis incurvis obtusis in setis longis. *Dill. Musc.* 389. t. 49. f. 58.

Mnium uliginosum. *With.* 800.

M. trichodes. *Hoffm. Germ.* v. 2. 47.

Meesia uliginosa. *Hedw. Sp. Musc.* 173. *Crypt. v. 1. 1. t. 1, 2.* *Swartz. Musc. Suec.* 44. *Roth. Germ.* v. 3. 256.

Angl. Long Capillary Thread-moss.

In uliginosis.

On bogs in Scotland. *Dicks.*

Perennis. Fruct. fert Junio.

Caules cæspitosi, vix unciales, quandoque longè breviiores, erecti, simplices et ramosi, nigri, basi ferruginco-tomentosi. Folia erecto-patentia, saturatè sed luteo-viridia, nitida, linear-lanceolata, obtusa, integriflora, carinata, nervo validiore, lutescente. Flores herinaphroditici. Pedicellus terminalis, solitarius, erectus, longissimus, ferè triuncialis, capillaris, ruber, basi fuscus, apice luteolus. Capsula inclinata, obovata, obtusa, incurva, fusca, apophysi brevi insidens. Operculum conicum, obtusum, brevissimum. Peristomium exterius breve: interius reticulatum, duplo longius. Calyptra subulata, recta.

- dealbatum.* 5. **BRYUM** caule brevissimo simplici, foliis lanceolatis

ceolatis reticulatis apice subserratis, capsulâ obovatâ cernuâ.

Bryum dealbatum. Dicks. *Crypt. fasc.* 8. t. 5.
f. 3. *With.* 816. *Hull.* 258.

Meesia dealbata. Swartz. *Musc. Suec.* 44 et
94. t. 5. f. 10. *Hedw. Sp. Musc.* 174. t. 41.
f. 6—9.

Angl. Pale-leaved Thread-moss.

In ericetis turfosis humidis montosis.

On Ben Lawers. Dicks. On St. Faith's bogs near
Norwich. Mr. Crowe.

Perennis? Fruct. fert æstate.

(Caules cæspitosi, bilineares, simplices. Folia albo-viridia, pellucida, reticulata, lanceolata, acuta, uninervia, apice obsoletè serrata. Pedicellus solitarius, terminalis, erectus, sesquiuncialis, ruber, subflexuosus. Capsula subcernua, obovata, vel subrotunda, arcuata, obtusa, ferruginea, opaca, apophysi valde constrictâ. Operculum convexum, mucronulatum, rubrum. Peristomium exterius breve, ferrugineum: interius reticulatum, duplò longius.

6. *BRYUM* caule simplici, foliis lanceolato-*cylindri-*
linearibus apicem versùs serratis, capsulâ *cum*
erectiusculâ cylindraceâ.

Bryum cylindricum. Dicks. *Crypt. fasc.* 4. 12.
t. 11. f. 4.

Angl. Cylindrical Thread-moss.

In montosis.

Near Halifax, Yorkshire. Dicks.

Annua? Fl. . . .

(Caulis haud semiuncialis, erectus, simplex. Folia lætè viridia, erecto-patentia, lanceolato-linearia, acuta, carinata, uninervia, basi integerrima, apicem versùs ferrata. Pedicellus erectus, terminalis, uncialis, rufus.

Capsula erectiuscula, pallidè fusca, tenuis, cylindracea, gracilis, recta, apophysi propriæ magnitudinis, obconicâ. Peristomium pallidum.

carneum. 7. BRYUM caule subsimplici, foliis lanceolatis acutis integerrimis reticulatis remotiusculis, capsulâ pendulâ turbinatâ.

Bryum carneum. *Sp. Pl.* 1587. *Huds.* 489. *With.* 837. *Hull.* 265. *Rehb.* 428. *Engl. Bot.* t. 360. *Roth. Germ.* v. 3. 231. *Swartz. Musc. Suec.* 47. *Schrad. Spicil.* 70.

B. delicatulum. *Hedw. Sp. Musc.* 179. *Crypt.* v. 1. 52. t. 20.

B. lanceolatum pellucidum, capsulis rotundis pendulis carnis. *Dill. Musc.* 400. t. 50. f. 69.

B. nitidum, foliis serpylli pellucidis angustioribus, reflexis capitulis subrotundis, carni coloris, in pediculis brevioribus. *Dill. in Raii Syn.* 102.

B. n. 1834. *Hall. Hist.* v. 3. 50. t. 45. f. 6.

Mnium carneum. *Hoffm. Germ.* v. 2. 51.

β. Bryum nitidum, perangustis serpyllinis pellucidis foliis, reflexis capitulis subrotundis virulentibus, longioribus pediculis nascentibus. *Dill. in Raii Syn.* 102; confer *Dill. Musc. loc. cit. f. G.*

Angl. Flesh-coloured Thread-moss.

In pastuis udis, irriguis.

On Black-heath, and near Dorking, Surrey. *Dill.* At Ditchingham, Norfolk, in wet drains near the bath. *Mr. Stone.* Cambridgeshire. *Rev. Mr. Hemsted.* On Marham fen, Norfolk. *Mr. Crowe.*

Perennis. Fruct. fert Aprili.

(Caules haud semiunciales, cæspitosi, eretici, simplices, apice demum quandoque ramosi, rubri, laxè foliosi. Folia alterna, pallidè viridia, pellucida, reticulata, laneeolata, acuta, integerrima, plana, uninervia. Pedicellus terminalis, solitarius, semiuncialis, subflexuofus, pallidè roseus. Capsula pendula, turbinata, subcampanulata, earnea, opaca, glauescens. Operculum conicum, breve, obtusum. Peristomium exterius ferrugineo-fuseum: interius flaveolens, duplii serie dentatum.

S. *BRYUM* caule ramoso erecto, foliis ovato-*alpinum*. lanceolatis acutis imbricatis, capsulâ oblongâ pendulâ.

Bryum alpinum. *Linn. Mant.* 309. *Huds.* 490. *With.* 841. *Hull.* 254. *Lightf.* 738. *Swartz.* *Musc. Suec.* 47.

B. hypnoides pendulum sericeum, comâ insigni atro-rubente. *Dill. Musc.* 394. t. 50. f. 64.
Angl. Red Alpine Thread-moss.

In rupibus alpinis.

On rocks covered with little earth near the ruined castle at Llanberis. *Dill.* In the north of England. *Huds.* On Ben Lomond. *Lightf.* North Wales. *Rev. H. Davies.* Isle of Shant. *Dr. De Ramm.* Luggy Lough, Ireland. *Dr. W. Stokes.*

Perennis. Fruct. fert Maio, Junio.

Caules cæspitosi, unciales vel biunciales, ramosi; ramis erectis, strictis, teretibus, densè foliosis, acutis. Folia undique imbrieata, viridi purpureoque puleherimè variata, nitida, ovato-laneeolata, acuta, integerrima, pellucida, nervo sèpiùs ruberriño; siccitate adpressa. Pedieelli laterales, solitarii, uneiales, rubri, reeti, eaulem superantes. Capsula pendula, obovato-cylindracea, demum rubra. Operculum hemisphæricum, umbonatum,

natum, fulvum vel rubrum. Peristomium albidum, vel roseum.

cæspiti- 9. BRYUM caule basi ramoso cæspitoso, foliis
tum. ovato-lanceolatis planis piliferis, capsulâ ob-
ovatâ pendulâ.

Bryum cæspiticium. Sp. Pl. 1586. Hedw.
Sp. Musc. 180. Huds. 488. Hull. 256.
Sibth. 290. Lightf. 736. Roth. Germ. v. 3.
239. Swartz. Musc. Suec. 50. Curt. Lond.
fasc. 3. t. 67. f. 1.

B. pendulum ovatum cæspiticium et pilosum,
setâ bicolori. Dill. Musc. 396. t. 50. f. 66,
exclusâ varietate F. G.

B. trichoides capitulis reflexis, pediculis imâ
mediatae rubris, summâ luteo-virentibus.
Raii Syn. 100.

Mnium cæspiticium. With. 807. Relb. 421.
Abbot. 235. Hoffm. Germ. v. 2. 50.

Hypnum n. 1790. Hall. Hist. v. 3. 39.

Muscus capillaceus minimus, capitulo nutante,
pediculo purpureo. Vaill. Paris. 134. t. 29.
f. 7.

Angl. Lesser Matted Thread-moss.

In muris, tectis, aggeribus et arenosis.

Perennis. Fruet. fert Martio, Aprili.

Caules densè cæspitosi, e basi ramosi, ramis trilinearibus,
simplicibus, patentibus, foliosis, obtusis. Folia lætè
viridia, ovato-lanceolata, plana, integerrima, uninervia,
pilo terminata; summa majora, stellatim patentia.
Pedicelli e basi ramorum, scèrè sesquiunciales, capillares,
erecti, basi purpurei, apice virides. Capsula pendula,
obovato-cylindracea. Operculum convexum, unih-
natum, fulvum. Peristomium flavum, dentibus acu-
minatis.

110. BRYUM caule brevissimo basi ramoso, foliis *bicolor*, ovato-lanceolatis acuminatis, capsulâ ovatâ obtusâ pendulâ.

Bryum bicolor. *Dicks. Crypt. fasc.* 4. 16.

B. præcedentis varietas parva. *Dill. Musc.* 397.
t. 50. f. 66, F, G.

Angl. Two-coloured Thread-moss.

In arenosis sterilibus siccis. *Dill. Dicks.*

Perennis. Fruct. fert Martio.

Caulis sparsi, haud bilineares, e basi ramosi, radicibus elongatis, fuscis. Folia luteo-viridia, parva, patula, ovato-lanceolata, plana, integerrima, mucronulata, nec pilifera. Pedicellus e basi ramorum, semiuncialis, erectus, ruberrimus, nitidus. Capsula pendula, majuscula, ovata, crassa, basi rubra, apice fulva, sèpè tota sanguinea. Operculum hemisphæricum, umbo-natum, nitidum, fulvum sive coccineum. Peristomium scèrè ut in *B. carneo*.

111. BRYUM caule basi ramoso cæspitoso, foliis *argenteo*-ovatis concavis piliferis imbricatis, capsulâ *um*. oblongâ pendulâ.

Bryum argenteum. *Sp. Pl.* 1586. *Hedw. Sp. Musc.* 181. *Huds.* 489. *With.* 838. *Hull.* 256. *Relb.* 427. *Sibth.* 293. *Abbot.* 243. *Lightf.* 735. *Swartz. Musc. Suec.* 47. *Roth.* *Germ. v. 3.* 229. *Ehrb. Crypt.* 262. *Dicks. H. Sicc. fasc.* 15. 19. *Curt. Lond. fasc.* 3. *t. 67. f. 2.*

B. pendulum julaceum, *argenteum et sericeum*. *Dill. Musc.* 392. *t. 50. f. 62.*

B. capitulis subrotundis reflexis, cauliculis tere-tibus argenteis. *Raii Syn.* 100.

B. n. 1821. *Hall. Hist. v. 3.* 47.

Mnium argenteum. *Hoffm. Germ. v. 2.* 51.

Muscus squamosus argenteus, ericæ folio.
Vaill. Paris. 134. t. 26. f. 3.
Angl. Silvery Thread-moss.

In muris, tectis, rupibus et apricis, ubique.

Perennis. Fruct. fert hyeme et vere.

Caules cæspitosi, pulvinati, semiunciales, e basi ramosi, ramis filiformibus erectis. Folia argenteo-nitida, arctè imbricata, ovata, integerrima, concava, uninervia, pilo brevi aristata. Pedicellus e basi ramorum, haud uncialis, capillaris, nitidus, basi purpureus, apice flavus. Capsula pendula, ovato-oblonga, brevis, rubicunda. Operculum conicum, brevissimum, fulvum. Peristomium luteum.

Zierii. 12. **BRYUM caule erecto ramoso, foliis ovatis concavis imbricatis acutis muticis, capsula clavata cernua.**

Bryum Zierii. *Dicks. Crypt. fasc. 2. 8. t. 4. f. 10.* *H. Sicc. fasc. 1. 19.* *Hedw. Sp. Musc. 182. t. 44. f. 1—4.* *With. 839. Hull. 256.* *Engl. Bot. t. 1021.*
Angl. Zierian Thread-moss.

Ad aggeres humidos in alpibus Scoticis et Cambro-britannicis.

Near Aberfeldy. *Dicks.* Carnarvonshire. *Mr. Griffith.*

Perennis. Fruct. fert autumno.

Caules cæspitosi, ramosi, crecti, ferè unciales, ramis teretibus, flexuosis, obtusis. Folia mox albida vel incarnata, nitida, arctè imbricata, ovata, integerrima, concava, mutica, uninervia. Pedicellus e basi ramorum, haud uncialis, erectiusculus, rubicundus. Capsula cernua, vix pendula, ante maturitatem clavato-cylindracca, viridis; demùm fusca, obovata, sive ovata, apophysi obconicæ insidens. Operculum conicum,

nicum, breve. Peristomium rubrum. Calyptra brevis, subulata, acuminata.

13. *BRYUM* caule basi ramoso, foliis ovatis con-julacea-
cavis imbricatis obtusis muticis, capsulâ cla-um.
vatâ pendulâ.

Bryum julaceum. Schrad. Spicil. 70. Roth.
Germ. v. 3. 230.

B. filiforme. Dicks. Crypt. fasc. 4. 16.

B. argenteum β. Sp. Pl. 1586. Huds. 489.
With. 839. Hull. 256.

B. n. 1821, β. Hall. Hist. v. 3. 47.

B. pendulum, surculis teretibus viridibus. Dill.
Mus. 394. t. 50. f. 63.

B. trichodes, capitulis reflexis, viridissimum
holosericeum. Dill. Giss. 226. app. 86. t. 2.
f. F.

Mnium holosericeum. Hoffm. Germ. v. 2. 51.
sub n. 18.

Angl. Slender-branched Thread-moss.

In glareosis et montosis.

In the walks of the botanic garden at Oxford. Dill.
In moist rather mountainous parts of Scotland.
Dicks.

Perennis. Fruct. fert Januario et Februario.

B. argenteo majus. Rami filiformes, graciles, erecti,
acuti. Folia nitida, luteo-viridia, (nec argentea,)
arctè imbricata, parva, ovata, integerrima, concavif-
cula, obtusa, mutica. Pedicellus e basi ramorum,
uncialis vel altior, subflexuosus, ruber. Capsula pen-
dula, obovato-cylindracea, fusca. Operculum con-
vexum, umbonatum, nitidum, pallidè fuscum.

4. *BRYUM* caule basi ramoso cæspitoso, foliis capilla-
obovatis re.

obovatis piliferis siccitate tortilibus, capsulâ clavatâ pendulâ.

Bryum capillare. Sp. Pl. 1586. Hedw. Sp. Musc. 182. Huds. 488. Hull. 255. Sibth. 290. Lightf. 737. Swartz. Musc. Suec. 50. Roth. Germ: v. 3. 238.

B. foliis latiusculis congestis, capsulis longis nutantibus. Dill. Musc. 398. t. 50. f. 67.

B. capitulis reflexis, foliolis latiusculis congestis. Raii Syn. 100.

Mnium capillare. Linn. Fl. Suec. 385. With. 805. Relb. 420. Abbot. 234. Hoffm. Germ. v. 2. 50.

Hypnum n. 1791. Hall. Hist. v. 3. 40.

Muscus capillaceus major, capitulis crassioribus cylindraceis nutantibus. Vaill. Paris. 134. t. 24. f. 6.

Angl. Greater Matted Thread-moss.

In sylvis, ericetis et aggeribus.

Perennis. Fruct. fert Martio, Aprili.

B. cæspititio simile, sed majus. Rami ferè unciales, erecti, apice incrassati atque stellati. Folia lætè viridia, obovata, obtusa, integerrima, uninervia, pilo terminata, siccitate tortuosa. Pedicelli sesquiunciales, firmi, rubri. Capsula pendula, cylindracea, obtusa, ferè clavata, duplò quam in B. cæspititio major. Operculum ferrugineum, nitidum, hemisphæricum, acumine brevi, recto. Peristomium exterius fulvum, dentibus acuminatis.

annoti- 15. *BRYUM* caule basi ramoso, foliis ovato-lan-

um. ceolatis pellucidis carinatis subserratis, capsulâ obovatâ pendulâ.

Bryum annotinum. Hedw. Sp. Musc. 183. t. 43. Huds.

Huds. 490. *Hull.* 255. *Sibth.* 291. *Swartz.*

Musc. *Suec.* 50. *Roth.* *Germ.* v. 3. 232.

B. *annotinum* lanceolatum pellucidum, capsulis oblongis pendulis. *Dill. Musc.* 399. t. 50. f. 68.

Mnium annotinum. *Sp. Pl.* 1576. *With.* 805.

Hoffm. *Germ.* v. 2. 49. *Leers.* *Herborn.* 231.

Muscus. *Fl. Dan.* t. 215.

Trentepohlia erecta. *Roth.* *Catal.* v. 1. 139.

Hoffm. *Germ.* v. 2. 17. t. 14.

Angl. Summer Thread-moss.

In pratis arenosis vel turfosis humidis, irriguis.

Perennis. Fruct. fert Maio—Augusto.

Caules rubri, ercti, foliosi: masculus simplex, flexuosus; sèpè elongatus, gemmiparus, sterilis: fœminæ basi ramosus. Folia lætè viridia, pellucida, oculo armato reticulata, ovato-lanceolata, carinata, uninervia, acuta, apicem versùs obsoletè ferrata. Pedicellus terminalis, solitarius, sesquiuncialis, ruber. Capsula pendula, subpyriformis. Operculum conicum, breve, acutum, carneum. Peristomium pallidum.

16. *BRYUM* caule basi diviso, foliis lanceolatis *compactum* carinatis apicem versùs ferratis, capsulâ ovatâ tum. pendulâ.

Bryum compactum. *Dicks.* *Crypt. fasc.* 4. 15.
t. 11. f. 11.

Angl. Compact Thread-moss.

In paludibus. *Dicks.*

Perennis. Fl. . .

Caules densè cæspitosi, semiunciales, stricti, vel simplices vel basi divisi, infernè denudati, rubri. Folia luteo-viridia, nitida, lanceolata, acuta, carinata, subrevoluta, c medio ad apicem ferrata; inferiora breviora, et ferè cordata. Pedicellus terminalis, haud uncialis, subflexuosus, ferrugineus. Capsula pendula, ferru-

ferruginina, ovata, brevis, margine dilatata, ferè turbinata. Operculum convexum, umbonatum.

- attenu-* 17. BRYUM caule simplici prolifero, foliis stellatis ovatis integerrimis reticulatis, capsulâ clavato-cylindraceâ erectâ.

Bryum attenuatum. *Dicks. Crypt. fasc.* 4. 8. t. 10. f. 8.

Angl. Slender Proliferous Thread-moss.

In paludibus montosis Scotiæ. *Dicks.*

Perennis. Fl. . .

Caules adscendentes, semiunciales, filiformes, indivisi, saepius annotino-proliferi, radicantes. Folia terminalia, stellatim-patentia, congesta, ovata, acuta, integriformia, uninervia, pellucida, reticulata. Pedicellus terminalis, erectus, semiuncialis, flaveolens. Capsula erecta, obovato-cylindracea, gracilis, pallidè fusca, ore subcoarctato. Operculum convexum, muticum. Peristomium, ex auctoritate amicissimi D. Turneri, est omnino Bryi.

- hornum.* 18. BRYUM caule simplici, foliis lanceolatis denticulato-serratis, pedicello arcuato, capsulâ ovatâ cernuâ.

Bryum hornum. *Huds.* 491. *Hull.* 266. *Sibth.* 290. *Curt. Lond. fasc.* 1. t. 71. *Dicks. H. Sicc.* *fusc.* 4. 23. *Roth. Germ.* v. 3. 237.

Swartz. Musc. Suec. 48. *Weber. Goelt.* 115.

B. stellare hornum sylvarum, capsulis magnis nutantibus. *Dill. Musc.* 402. t. 51. f. 71.

B. nitidum, capitulis majoribus reflexis, calyptrâ imum vergente, pediculis oblongis e caulinis novis egredientibus. *Raii Syn.* 102.

Mnium hornum. *Sp. Pl.* 1576. *With.* 804. *Lightf.* 711. *Rehb.* 419. *Abbot.* 233. *Hoffm. Germ.* v. 2. 48. *Ehrb. Crypt.* 3.

M. n.

M. n. 1846. *Hall. Hist.* v. 3. 54.

Muscus roseus, polygoni folio. *Vaill. Paris.* 134.
t. 24. f. 5.

Angl. Swan's neck Thread-moss.

In nemorosis udis.

Perennis. Fruct. fert Martio.

Praecendentibus majus. Caules cæspitosi, altitudine varii, simplices, foliosi, rubri, basi ferruginco-tomentosi. Folia lætè viridia, pellucida, membranacca, patentia, lanceolata, acuta, mutica, denticulato-serrata, nervo valido, rubro; siccitate undulosa. Flos masculus magnus, terminalis, discoideus, foliis majoribus, obtusis, circumvallatus: fœmineus in diversâ stirpe. Pedicellus terminalis, solitarius, plerumque biuncialis vel ultrâ; basi rectus, ruber; apice arcuatus, flavescens, sub fructu incrassatus. Capsula magna, cernua, ovato-elliptica, turgida, retusa, pallidè ferruginea, ore coccineo. Operculum hemisphæricum, mucronulatum. Peristomium exterius citrinum, dentibus acutis; interius album.

19. BRYUM caule simplici, foliis lanceolatis acu-crudum.
tis integerrimis: summis angustatis denticu-
latis, capsulâ oblongâ cernuâ.

Bryum crudum. *Huds.* 491. *Hull.* 266. *Roth.*

Germ. v. 3. 242. *Swartz. Musc. Suec.* 49.

B. sericeum. *Huds.* 487, ex D. Davies.

B. pendulum hornum molle, foliis et lanceolatis
et gramineis. *Dill. Musc.* 401. t. 51. f. 70.
Herb. Dill.

Mnium crudum. *Sp. Pl.* 1576. *Hedw. Sp.*

Musc. 189. *Crypt.* v. 1. 99. t. 37. *With.* 807.

Hoffm. Germ. v. 2. 48. *Ehrb. Crypt.* 152.

Angl. Transparent-green Thread-moss.

In udis alpinis.

In Wales. *Dill. and Rev. H. Davies.* On Crib
y Ddescil

y Ddescil and Cader Idris; Mr. Griffith.
With. In woods about Rydall, Westmoreland. *Huds.*

Perennis. Fruct. fert æstate.

Caules cæspitosi, simplices, erecti, vix unciales; basi denudati, atro-purpurei; apice densè foliosi. Folia nitidissima, lætè viridia, pellucida, patula, lanceolata, acuta, plana, uninervia, integerrima; inferiora breviora; summa angustiora, et apice plerumque obsoletè ferrata. Pedicellus terminalis, erectus, uncialis, ruber. Capsula cernua, elliptico-oblonga, ferruginea, ore vix rubicundo. Operculum hemisphæricum, apiculatum, luteolum. Peristomium citrinum. Calyptora recurva, decidua.

Mnium crudum Lightf. est *B. marginatum*.

margina 20. **BRYUM** caule simplici, foliis lanceolatis acutum. tis denticulatis margine incrassatis, capsulâ ovatâ, operculo conico.

Bryum marginatum. *Dicks. Crypt. fasc. 2. 9. t. 5. f. 1. With. 841. Hull. 267.*

Mnium crudum. *Lightf. 712,* cum descrip-
tione præcedentis, more suo e Dillenii pagi-
nis, haud felicitè, compilatâ.

Angl. Thick-edged Thread-moss.

In Scotiæ pasuis. *Dicks.*

In woods and moist shady places in the Lowlands.
Rev. Dr. Burgess.

Perennis. Fruct. fert Maio.

Caules unciales, simplices, erecti, rubri. Folia elliptico-
lanceolata, acuta, repanda, apicem versùs denticulata,
nervo margineque rubro, incrassato. Pedicellus fo-
litarius, erectus, uncialis, carneus. Capsula pendula,
ovata, retusa. Operculum conico-subulatum. Calypp-
tra subulata, rubicunda, recurva.

E speciminiibus a venerando amico Burgessio olim dono
datis

datis hanc speciem adumbravi. Nomen Dicksoni communicavit D. Turnerus.

21. BRYUM caule ramoso interruptè folioso, fo-*interrup-*
liis lanceolatis acutis undique ferrulatis, cap-*tum,*
sulâ pyriformi erectiusculâ.

Bryum interruptum. *Dicks. Crypt. fasc. 4. 13.*
t. 11. f. 6.

Angl. Interrupted Thread-moss.

In rupibus alpinis Scotiæ.

On Ben Nevis. *Dicks.*

Perennis. Fl. . .

Caules cæspitosi, sesquiuncialcs, determinatè-ramosi, in-
terruptè foliosi, rubri. Folia fasciculata, luteo-viridia,
nitida, reticulata, lanceolata, acuta, uninervia, æqua-
litèr ferrulata; inferiora remota, exigua, brevia. Pe-
dicellus terminalis, erectus, ruber, vix uncialis. Cap-
sula erectiuscula, pyriformis, fusca, ore dilatato. Oper-
culum hemisphæricum, umbone obtuso.

22. BRYUM caule basi ramoso, foliis lanceolatis *nigri-*
*strictis apicem versùs ferratis, capsulâ turbi-*cans.**
natâ pendulâ.

Bryum nigricans. *Dicks. Crypt. fasc. 4. 15.*
t. 11. f. 10.

Angl. Dark-green Thread-moss.

In paludibus turfosis Scotiæ. *Dicks.*

Perennis. Fl. . .

Caules cæspitosi, sæpiùs basi ramosi, foliosi. Folia
tristè virentia, imbricata, stricta, lanceolata, acuta,
carinata; apicem versùs ferrata; basi dilatata et inte-
gerrima. Pedicellus terminalis, ferrugineus, haud un-
cialis, apice arcuatus. Capsula pendula, turbinata,
ferruginea, ore patulo. Operculum convexum, ob-
tusè umbonatum. Peristomium flavum.

incli-
nans. 23 BRYUM? caule ramoso, foliis setaceis in-
tegerrimis siccitate crispis, capsulâ ovatâ in-
clinatâ, apophysi minutâ.

Bryum inclinans. *Dicks. Crypt. fasc. 4. 11. t. 11.*

f. 9.

Angl. Inclining Curling Thread-moss.

In alpibus Scoticis. *Dicks.*

On the summit of Craig Cailleach. *Mr. R. Brown.*

Perennis. Fl. . .

Caulis erectus, ramosus. Folia imbricato-patula, setacea,
carinata, integerrima, basi dilatata. Pedicelli latera-
les, erecti, semiunciales. Capsula inclinata, ovata,
apophysi minutâ.

Ex iconē et descriptione Dicksoni.

cubitale. 24. BRYUM caule adscendente subramoso, foliis
lanceolatis integerrimis decurrentibus, pedi-
cello geniculato, capsulâ clavatâ pendulâ.

Bryum cubitale. *Dicks. Crypt. fasc. 2. 9. t. 5.*
f. 2. *With. 840.* *Hull. 267.*

Angl. Elbow-shaped Thread-moss.

Ad aggeres humidos in alpibus Scoticis.

Near Abersfeldy, along with Bryum Zierii. *Dicks.*

Perennis. Fruct. fert autumno.

Caules adscendentēs, bi- vel triunciales, foliosi, ferrugi-
neo-tomentosi. Folia patula, lätè viridia, pellucida,
lanceolata, acuta, integerrima, margine incrassata, basi
plus minus sagittato-decurrentia, nervo valido, rubro.
Pedicellus terminalis, solitarius, tri- vel quadriuncialis,
aureo-fuscus, nitidus, basin versus geniculatus. Cap-
sula pendula, ferruginea, clavata, sive obovata, basi
attenuata.

“ Maximum inter omnia Brya hucusque nota.” *Dicks.*
A B. alpino, si quid aliud, distinctum. Conf. *With.*
loc. cit.

25. BRYUM caule determinatè ramoso, ramis *ventricosum* adscendentibus, foliis alternis lanceolatis apice *serratis*, capsulâ *ventricosa* pendulâ.

Bryum ventricosum. *Dicks. Crypt. fasc. 1. 4.*
Swartz. Musc. Suec. 49. *Reh. 427.*

B. *triquetrum*. *Huds. 490.* & *Hull. 256.*
B. *pseudo-triquetrum*. *Roth. Germ. v. 3. 243.*
B. *annotinum* *palustre*, *capsulis ventricosis pendulis*. *Dill. Musc. 454. t. 51. f. 72.*
B. *nitidum*, *foliis serpilli angustioribus*, *majus*. *Dill. in Raii Syn. 102.*

Mnium pseudo-triquetrum. *Hedw. Sp. Musc. 190.* *Crypt. v. 3. 19. t. 7.* *Hoffm. Germ. v. 2. 49.* *With. 808.*

M. triquetrum. *Abbot. 235.* *Lightf. 715.*
Angl. Swelling Bog Thread-moss.

In palūdibūs et ad mārgines rivulorum:

Perennis. Fruct. fert Maio.

Caules cæspitosi, determinatè ramosi, longius radicati, ferrugineo-tomentosi; ramis numerosis, adscendentibus, rubris, laxè foliosis. Folia pallidè viridia, alterna, vix trifaria, erecto-patula, lanceolata, acuta; uninervia, apice evidentè serrata, siccitate torta. Flores terminales, saepius (monente Hedwigio) hermaphroditici. Pedicellus ex apice caulis, seu rami præcedentis anni, sesquiuncialis vel biuncialis, erectus, basi purpureus, apice flavescens. Capsula pendula, ovata, brevis, basi ventricosa, sub ore paululum constricta. Operculum convexum, mueronulatum. Peristomium flavum.

26. BRYUM caule ramoso, foliis alternis linearis *bimum*. lanceolatis apice subserratis, capsulâ obovato-clavatâ pendulâ.

Bryum bimum. *Schreb. Lips. 83.* *Roth. Germ. v. 3. 243.* *Swartz. Musc. Suec. 50.*

B. *triquetrum* *B. Huds. 490.* *Hull. 256.*

- B. lanceolatum bimum, setis et capsulis longis pendulis. *Dill. Musc.* 405. t. 51. f. 73.
 B. nitidum, foliis serpilli pellucidis angustis, capitulis tumidis nutantibus, præaltis pediculis e surculis annotinis egredientibus. *Dill. in Raii Syn.* 102.
 Mnium bimum. *Hoffm. Germ.* v. 2. 48.
 M. pseudo-triquetrum, var. 2. *With.* 808.
Angl. Long-stalked Bog Thread-moss.

In uliginosis et palustribus montosis.

Perennis. Fruet. fert æstate.

Præcedente majus. Caulis bi- vel triuncialis, inconditè ramosus, atro-ferrugineus. Folia elongata, luteo-viridia, apice obsoletè serrata. Pedicelli ex apice ramorum bimorum vel trimorum, optimè notante Dillenio, bi- vel triunciales, atro-purpurei, nitidi, basin versùs subinde geniculati. Capsula pendula, ferruginea, ob-ovato-cylindracea, basi angustata, sub ore constricta, duplò ferè quàm in præcedente longior. Peristomium flavum.

- turbina-* 27. BRYUM caule ramoso, foliis ovatis aristatis *tum.* integerrimis, capsulâ pyriformi pendulâ.
Bryum turbinatum. *Swartz. Musc. Suec.* 49.
Roth. Germ. v. 3. 245.
 B. triquetrum γ. *Huds.* 491.
 B. palustre complicatum rubens, capsulis turbinatis pendulis. *Dill. Musc.* 406. t. 51. f. 74.
 B. nitidum rubens, capitulis reflexis, foliis angustis pellucidis, cauliculis proliferis. *Dill. in Raii Syn.* 102.
 Mnium turbinatum. *Hedw. Sp. Musc.* 191.
Crypt. v. 3. 22. t. 8. *Hoffm. Germ.* v. 2. 49.
 M. nutans. *With.* 803. *Hull.* 255.
Angl. Pear-shaped Bog Thread-moss.

In palustribus arenosis.

On Shooter's hill, and in Wales. *Dill.*

Perennis. Fruct. fert vere et æstate.

Caules vix unciales, cæspitosi, ramosi. Folia lætè viridia, patula, alterna, approximata, ovata, integerrima, uninervia, apice aristata. Pedicellus uncialis, purpleo-fuscus, subflexuosus. Capsula pendula, ferruginea, subrutilans, pyriformis, basi attenuata. Operculum convexum, muticum. Peristomium ferrugineum.

28. *BRYUM* caule ramoso, foliis ovatis acuminata *stellare*.
tis orbiculato-reticulatis: summis serratis stel-
latis, capsulâ oblongâ cernuâ.

Bryum stellare. *Roth. Germ.* v. 3. 240.

B. proliferum. var. 2. *Hull.* 266.

B. serpyllifolium ε. *Huds.* 492.

B. stellare roseum pendulum minus, capsulis et
setis longioribus. *Dill. Musc.* 412. t. 52.
f. 78.

B. roseum minus, foliis subrotundis. *Raii Syn.*
92.

Mnium stellare. *Hedw. Sp. Musc.* 191. *Hoffm.*
Germ. v. 2. 51.

M. proliferum, var. 2. *With.* 807.

Angl. Starry-leaved Thread-moss.

In sylvis et ericetis udis.

Perennis. Fl. . .

Caulis determinatè ramosus, uncialis vel biuncialis. Fo-
lia lætè viridia, ovata, breviùs acuminata, oculo ar-
matò pulcherrimè reticulata, interstitiis reticuli orbiculatis, nervo valido, rubro, apice ferè evanescente;
inferiora integerrima; summa apicem versùs ferrata,
stellatim patentia. Pedicellus terminalis, ferè biuncia-
lis, subarctatus, apice incrassatus. Capsula cernua,
elliptico-cylindracea. Operculum hemisphæricum.

- tuspidata*- 29. BRYUM caule simplici: sterili procumbente subramoso, foliis ovatis ferratis, capsulâ ovatâ, operculo hemisphærico.
- Bryum cuspidatum. *Swartz. Musc. Suec.* 51.
Roth. Germ. v. 3. 246. *Hull.* 266. *Sibtb.* 291.
- B. serpyllifolium γ. *Huds.* 492.
- B. pendulum, foliis variis pellucidis, capsulis ovatis. *Dill. Musc.* 413. t. 53. f. 79.
- Mnium cuspidatum. *Hedw. Sp. Musc.* 192.
t. 45. f. 5—8. Hoffm. Germ. v. 2. 52.
With. 804. *Rehb.* 420. *Abbot.* 234. *Ehrb.*
Crypt. 73.
- M. serpyllifolium β. *Sp. Pl.* 1577. *Weis.*
Crypt. 155.
- M. n. 1841. *Hall. Hist.* v. 3. 52.
Angl. Pointed-leaved Thyme Thread-moss.

In umbrosis humidiusculis.

Perennis. Fruct. fert Aprili, Maio.

Caulis foliosus, biformis; steriles procumbentes, radicantes, elongati, triunciales vel ultrâ, interdùm subramosi; fertiles erecti, simplices, unciales. Folia lètè viridia, pellucida, oculo armato minutè reticulata, alterna, latè ovata, subundulata, acutè ferrata, mucronulata, nervo valido, pallescente. Pedicellus terminalis, uncialis, flexuosus, roseus. Capsula pendula, ovata, brevis, ventricosa, retusa, lutea, hinc fulva, ore coccineo. Operculum hemisphærico-conicum, muticum. Peristomium citrinum.

- punctata*- 30. BRYUM caule subsimplici, foliis obovatis integrum. punctato-reticulatis, capsulâ ellipticâ, operculo subulato.
- Bryum punctatum. *Roth. Germ.* v. 3. 245.
Hull. 266. *Sibtb.* 291. *Engl. Bot.* t. 1183.
B. ser-

- B. serpyllifolium. *Swartz. Musc. Suec.* 51.
Huds. 492, α.
- B. pendulum, serpilli folio rotundiore pellucido, capsulis ovatis. *Dill. Musc. 416. t. 53. f. 81.*
- B. nitidum, serpilli rotundis et latioribus foliis pellucidis. *Raii Syn. 103.*
- Mnium punctatum. *Hedw. Sp. Musc. 193.*
With. 805. Relb. 420. Abbot. 234.
- M. serpyllifolium α . *Sp. Pl. 1577.*
Angl. Dotted-leaved Thyme Thread-moss.

In umbrosis irriguis.

Perennis. Fruct. fert Martio, Aprili.

Habitus præcedentis. Caules sesquiunciales, foliosi, rubri, sæpè flexuosi. Folia latè obovata, obtusissima, mucronulata, integerima, aut vix ferrulata, pulchrè ac evidentissimè reticulato-punctata, margine incrassato, rubro, repando, nervo valido, rubro; summa stellata. Pedicellus terminalis, uncialis vel sesquiuncialis, purpurascens, apice arcuatus. Capsula latè elliptica, pendula, virens, ore purpureo, demùm fusca. Operculum flavum, basi convexum, acumine subulato, obtusiusculo, recurvo, capsulâ duplò breviore.

Foliis reticulato-punctatis primâ facie dignoscitur.

31. BRYUM caule simplici, foliis obovato-spatulatis ferrulatis, capsulâ ovatâ, operculo *tum.* subulato.

Bryum rostratum. *Schrad. Spicil. 72,* ex ipso auctore.

B. serpyllifolium β . *Huds. 492.*

B. punctatum, var. 2. *Hull. 266.*

B. pendulum, serpilli folio longiore pellucido, capsulis oblongis cuspidatis. *Dill. Musc. 416. t. 53. f. 80. Herb. Dill.*

Mnium ellipticum. Hoffm. Germ. v. 2. 52.
M. punctatum, var. 2. With. 806.
M. n. 1840. Häll. Hist. v. 3. 52.
Angl. Long-beaked Thyme Thread-moss.

In palustribus montosis.

In the western part of Yorkshire. Richardson.

Perennis. Fruct. fert vere et æstate.

A præcedente differt caule breviori; foliis minoribus et angustioribus, haud reticulatis, in ipso specimine Dilleniano ferrulatis, nec integerrimis. Pedicelli solitarii vel aggregati, terminales, sesquiunciales, carnei, basi purpurei. Capsula ovata vel ovato-elliptica, pendula, virens, ore carneo. Operculum flavum, basi convexum, acumine subulato, obtuso, recurvo, capsulâ paululùm breviore.

roseum. 32. **BRYUM** caule subprolifero, foliis confertis stellatis ovatis acutis punctulatis margine scabris, operculo conico.

Bryum roseum. Schreb. Lips. 84. Swartz. Musc. Suec. 51. Roth. Germ. v. 3. 247.

B. serpyllifolium δ. Hud. 492.

B. proliferum. Sibth. 292.

B. stellare roseum majus, capsulis ovatis pendulis. Dill. Musc. 411. t. 52. f. 77.

B. roseum majus, foliis oblongis. Raii Syn. 92.

Mnium roseum. Hedw. Sp. Musc. 194. Weis. Crypt. 157.

M. proliferum. With. 806. Hull. 250. Abbot. 235.

M. serpyllifolium γ. Sp. Pl. 1578.

Muscus stellaris roseus. Baub. Pin. 361. Prodr. 151. Buxb. Cent. 2. t. 1. f. 3.

Angl. Rosaceous Thyme Thread-moss.

In sylvis et ericetis udis, haud vulgaris.

Perennis. Fruct. fert hyeme, sed rariūs.

Caulis steriles elongati, procumbentes, foliis minoribus, obtusis, alternis, tecti; fertiles erecti, unciales, sæpè proliferi, infernè squamosi, aphylli, apice foliosi, foliis confertis, stellatis, subdecurrentibus, ovatis, acutis, uninervibus, punctulatis, planis, margine scabris sive tenuissimè crenulatis. Pedicelli terminales, sæpiùs aggregati, sesquiunciales vel biunciales, rigidi, rubri, nitidi. Capsula pendula, ovato-oblonga, ferruginea, nitida. Operculum, ex auctoritate Dillenii, conicum, breve.

33. **BRYUM** caulis simplicibus apice demùm *ligularia*-
ramosis, foliis oblongis undulatis ferratis, pe-tum.
dicellis aggregatis.

Bryum ligulatum. Schreb. Lips. 84. Ehrb.

Crypt. 13. Swartz. Musc. Suec. 51?

B. undulatum. Sibth. 292. Roth. Germ. v. 3.
249.

B. dendroides. Web. Goett. 117.

B. dendroides polycephalon, phyllitidis folio
undulato pellucido, capsulis ovatis pendulis.
Dill. Musc. 410. t. 52. f. 76.

B. nitidum, foliis oblongis undatis, capitulis
cernuis, arbusculam referens. Raii Syn. 103.

Mnium undulatum. Hedw. Sp. Musc. 195.
With. 806. Hull. 250. Relb. 420. Abbot.
235. Weis. Crypt. 158.

M. serpyllifolium ð. Sp. Pl. 1578.

M. n. 1844. Hall. Hist. v. 3. 53.

Muscus polygoni folio. Vaill. Paris. t. 24. f. 3.
Mich. Gen. 108. t. 59. f. 5.

Angl. Long-leaved Thyme Thread-moss.

In sylvis humidis ad radices arborum.

Pereennis. Fruct. fert Aprili, haud frequentèr.
Hujus generis maxima et formosissima species. Radix
 repens. Caules erecti, suffruticosi, foliosi; steriles
 elongati; fertiles demùm determinatè ramosi, ramis
 stellatis, adscendentibus. Folia lätè viridia, approx-
 imata, alterna, decurrentia, patentia, linearis-oblonga,
 obtusa, uninervia, undulata, acutè ferrulata, siccitate
 mox flaccescentia. Pedicelli aggregati, biunciales,
 fulvi, nitidi. Capsula pendula, ovata, brevis, ferru-
 ginea. Operculum conicum, breve.

460. POLYTRICHUM. G. Pl. 563. Juss. 11.
 Schreb. Gen. 762. Hedw. Fund. v. 2. 90.
 t. 7. f. 37.

Caps. ventricosa. *Perist. exterius* dentibus tri-
 ginta-duo vel sexaginta-quatuor, brevibus,
 planis, inflexis: *interius* membrana orbicu-
 lata, transversa, apicibus dentium exterioris
 affixa. *Calyptra* sæpiùs duplex; *exterior* pi-
 losa.

* *Calyptrâ dupliči.*

commu- 1. POLYTRICHUM caule simplici, foliis li-
 ne, neari-lanceolatis ferrulatis, capsulâ erectâ
 quadrangulari, apophysi subrotundâ.

Polytrichum commune. Sp. Pl. 1573. Hedw.
 Sp. Musc. 88. Menz. Tr. of Linn. Soc. v. 4.

74. Huds. 469. With. 795. Hull. 247.
 Relh. 415. Sibth. 303. Abbot. 232. Dicks.
 H. Sicc. fasc. 17. 19? Swartz. Musc. Suec.
 75. Hoffm. Germ. v. 2. 23. Roth. Germ.
 v. 3. 346. Engl. Bot. t. 1197.

P. juccæfolium. Ehrb. Crypt. 214. Pl. Off.
 430.

P. quadrangulare vulgare, juccæ foliis serratis.
 Dill. Musc. 420. t. 54. f. 1.

P. vulgare

P. vulgare et majus, capsulâ quadrangulari.

Raii Syn. 90.

Muscus. Plum. Fil. t. B. f. 6. Mick. Gen.

t. 59. f. 1.

Angl. Common Hair-moss.

In sylvis uliginosis vel turfosis.

Perennis. Fruct. fert Maio, Junio.

Plantæ cæspitosæ. Radix longissima, simplex, filiformis, repens, villosa. Caulis plerumque simplicissimus, erectus, strictus, palmaris, spithameus aut pedalis, undique foliosus. Folia undique patentia, linear-lanceolata, acuta, ferrulata, saturatè viridia, nitida, glaberrima, ecarinata; basi membranacea, vaginantia. Flores dioici, terminales: masculi disciformes, pallidi, foliis numerosis, margine membranaceis, dilatatis, circumvallati, saepius e medio proliferi. Pedicellus solitarius, rectiusculus, firmus, bi- vel trunicalis, nitidus, ruber, apice pallidior, basi foliis pluribus dilutè fuscis, membranaceis, aristatis, vaginatus. Apophysis depreesa, torulosa, virens. Capsula erecta, tetraëdra, brevis, crassa, viridis, ore rubro. Operculum depresso, flavescentia, margine rubro, subcrenato, acuminè brevi, recto, gracili. Peristomium exterius dentibus 64, basi connatis, margine albis: intcrius membranaceum, planum. Calyptra duplex: exterior e pilis numerosissimis, aureo-fuscis, basi divaricatis, apice in mucronem coartatis, conflata: interior membranacea, brevior, tubulosa, basi fissa.

2. **POLYTRICHUM** caule simplici, foliis li-attenuatae lanceolatis patentibus ferratis, capsulatum. cernuâ quadrangulari basi constrictâ.

Polytrichum attenuatum. Menz. Tr. of Linn.

Soc. v. 4. 72. t. 6. f. 2. Engl. Bot. t. 1198.

*P. formosum. Hedw. Sp. Musc. 92. t. 19.
f. 1. a.*

P. Apuleii majus. Fuchs. Hist. 629?

Muscus

Muscus capillaris sive *Adiantum aureum majus*.

Ger. em. 1559?

Angl. Slender Hair-moss.

In ericetis turfosis.

On Sydenham Common, Kent. *Mr. Sowerby*.

Near Tintern Abbey, Monmouthshire. *Miss Hudson*.

Perennis. Fruct. fert Aprili.

Pulchra species, et forsitan haud infrequens, sed cum priore diù, a Dillenio aliisque ut videtur, confusa. Caule sæpiùs humiliore, foliis carinatis, evidentiùs ferratis et patentibus, paululùmque longioribus, capsulâ cernuâ, angustiore, longiore, basi constrictâ, at præcipue defectu apophysis, a *P. communi* differt.

gracile. 3. **POLYTRICHUM** caule simplici, foliis linear-lanceolatis erectiusculis ferratis, capsulâ cernuâ ovatâ subangulatâ basi constrictâ.

Polytrichum gracile. *Menz. Tr. of Linn. Soc.* v. 4. 73. t. 6. f. 3. *Hull.* 247.

P. longisetum. *Swartz. Musc. Suec.* 76, et 103. t. 8. f. 16.

Angl. Long-stalked Hair-moss.

In alpibus Scoticis. *Dickson*.

Perennis. Fruct. fert Maio..

Præcedenti affinis, discrepat verò caule breviori, foliis erectioribus, capsulâ minùs exactè tetraëdrâ, sed ovatâ, ventricosâ, basi parùm constrictâ. Apophysis nulla. Folia basin versus integerrima. Peristomii dentes 32, duplò quam in præcedente pauciores.

piliferum.

4. **POLYTRICHUM** foliis lanceolatis integerimis revolutis apice piliferis, capsulâ quadangulari, apophysi depresso.

- Polytrichum piliferum. *Schreb. Lips.* 74. *Menz.*
Tr. of Linn. Soc. v. 4. 75. *Hedw. Sp. Musc.* 90. *Hull.* 247. *Sibth.* 306. *Ehrb. Crypt.* 92. *Roth. Germ. v. 3.* 348. *Engl. Bot. t. 1199.*
- P. commune γ. *Sp. Pl.* 1573. *Huds.* 470.
With. 796. *Lightf.* 700.
- P. quadrangulare minus, juniperi foliis pilosis.
Dill. Musc. 426. *t. 54. f. 3.*
Angl. Bristle-leaved Hair-moss.

In ericetis arenosis aridis, et apricis glareosis.

Perennis. Fruct. fert Aprili, Maio.

Caules simplices, vix unciales, basi denudati, rubri; quandoque, ex observationibus Cl. Schwaegrichenii, ramosi. Folia congesta, patentia, atro-viridia, lanceolata, integerima, revoluta, pilo albo, in superioribus longiore magisque flexuoso, aristata. Flores masculi rubro variati. Pedicellus vix sesquiuncialis, saturatè ruber, rigidus. Capsula erecta, ovato-tetragona, viridis, apophysi depressâ rubrâ. Operculum conicum, rubrum. Calyptra exterior rufo-fusca, aureo nitens.

5. POLYTRICHUM foliis linear-lanceolatis *juniperi*-marginibus inflexis integerimis, capsulâ num. quadrangulari, apophysi depressâ, caule simplicissimo.

Polytrichum juniperinum. *Menz. Tr. of Linn. Soc. v. 4.* 76. *t. 6. f. 4.* *Hedw. Sp. Musc.* 89. *t. 18. f. 6—10.* *Relb.* 415. *Sibth.* 306. *Abbot.* 232. *Roth. Germ. v. 3.* 348. *Ehrb. Crypt.* 82. *Pl. Off.* 180. *Engl. Bot. t. 1200.*

P. commune β. *Sp. Pl.* 1573. *Huds.* 469.
With. 795. *Lightf.* 700.

P. quadrangulare, juniperi foliis brevioribus et rigidioribus.

rigidioribus. *Dill. Musc.* 424. t. 54. f. 2.
Herb. Dill.

P. montanum et minus, capsulâ quadrangulari.
Raii Syn. 90.

Angl. Juniper-leaved Hair-moss.

In ericetis montosis.

Perennis. Fruct. fert Aprili.

Præcedente major. Caules unciales vel sesquiunciales, undique foliosi, simplices. Folia atro-viridia, patentia, linear-lanceolata, acuta, mutica, integerima, margine arctè inflexo, nec revoluto. Pedicellus bi- vel triuncialis, rigidus, elegantè rubens. Capsula erecta, ovato-tetragona, virens, demùm cernua, fuscescens. Apophysis rubra, valdè depressa. Calyptre exterior magna, acumine fusco.

Dillenius folia ferrulata, minùs accuratè, delinavit.

strictum. 6. **POLYTRICHUM** foliis lanceolatis marginibus inflexis integerimis, capsulâ quadrangulari, apophysi depressâ, caule ramoso.

Polytrichum strictum. *Menz. Tr. of Linn. Soc.*
v. 4. 77. t. 7. f. 1. *Hull.* 247.

Angl. Close-leaved Hair-moss.

In apricis sterilioribus montosis.

On the tops of park walls about a mile east of Taymouth, and in other parts of Scotland.

Mr. Menzies. In North Wales. *Rev. H. Davies.* At Meavis bank near Edinburgh, and on Dersingham Moor, Norfolk, 1781.

Perennis. Fruct. fert Martio.

Caules cæspitosi, basi ramosi, tomento albido, molli, spongioso, ut et radices, densè vestiti. Folia erectissima, lætè viridia, lanceolata, acuta, submucronulata, integerima, margine arctè inflexo ut in priore; superiora, sive interiora, filo aristata. Pedicellus biuncialis,

uncialis, ruber. Capsula erecta, acutè quadrangula, viridis, apophysi depresso-planâ, rubicundâ. Operculum conicum, apice sèpè aduncum. Calyptra præcedentis.

Præcedentis varietatem censet D. Turnerus, nec forsitan immerito.

7. POLYTRICHUM foliis lanceolato-subulatis *alpinum*,
ferrulatis, capsulâ ovatâ cernuâ, apophysi
turbanatâ, caule ramoso.

Polytrichum alpinum. *Sp. Pl.* 1573. *Hedw.*
Sp. Musc. 92. *Menz. Tr. of Linn. Soc.* v. 4.
83. *Huds.* 470. *With.* 797. *Hull.* 248.
Lightf. 703. *Dicks. H. Sicc. fasc.* 12. 19.
Roth. Germ. v. 3. 349. *Swartz. Musc.*
Suec. 76. *Hoffm. Germ.* v. 2. 24. *Ehrh.*
Crypt. 2.

P. alpinum ramosum, capsulis e summitate el-
lipticis. *Dill. Musc.* 427. t. 55. f. 4.

P. n. 1800. *Hall. Hist.* v. 3. 43. t. 46. f. 6.
Angl. Alpine Hair-moss.

In ericetis et campis alpinis.

On Snowdon. *Dillenius.* On the top of Ben Lo-
mond.

Perennis. Fruct. fert Augusto.

Caules palmares; basi gracillimi, simplices, denudati;
apice ramosi, fastigiati, foliosi. Folia fusco-viridia,
vaginantia, recurvato-patula, lancolata, ecarinata,
denticulato-ferrata, apice subulata. Pedicelli solitarii,
sesquiunciales, lutescentes, subflexuosi. Capsula ovata,
fusca, mox cernua. Apophysis turbinata, capsulâ an-
gustior. Operculum planum, rostro subulato. Ca-
lypta fusco-rutilans. Peristomii dentes circiter 32,
albidi.

8. POLYTRICHUM foliis lanceolatis ferrulatis, *urnige-*
capsulârum.

capsulâ cylindraceâ erectâ apophysi destitutâ,
caule ramoso fastigiato.

Polytrichum urnigerum. *Sp. Pl.* 1573. *Hedw.*
Sp. Musc. 100. *t. 22. f. 5—7.* *Menz. Tr.*
of Linn. Soc. v. 4. 81. *With.* 797. *Hull.*
248. *Lightf.* 703. *Dicks. H. Sicc. fasc.*
12. 20. *Roth. Germ.* v. 3. 350. *Swartz.*
Musc. Suec. 77. *Hoffm. Germ.* v. 2. 24.
Ehrb. Crypt. 44. *Engl. Bot.* t. 1218.

P. alpinum β. *Huds.* 471.

P. ramosum, setis ex alis urnigeris. *Dill. Musc.*
427. *t. 55. f. 5.*

P. n. 1801. *Hall. Hist.* v. 3. 43.

Muscus erectus, *juniperi folio ramosus*. *Vaill.*
Paris. t. 28. f. 13.

Angl. Urn-headed Hair-moss.

In ericetis montosis Cambrobritannicis et Scoticis.

Perennis. Fruct. fert Augusto.

Habitus præcedentis, at caule breviori, ramis densius
fastigiatis. Folia lanceolata, planiuscula, ferrulata,
acuta, nec apice subulata. Pedicelli laterales, capil-
lares, purpurascentes. Capsula erecta, ovato-cylin-
dracea, gracilis, sub apice constricta. Apophysis
nulla. Operculum convexum, margine rubro, acu-
mine brevi, adunco. Calyptra pallida.

subro- 9. *POLYTRICHUM* foliis lanceolatis obtusis
tundum. *integerrimis rigidis*, capsulâ turbinatâ erec-
tiusculâ, caule brevissimo.

Polytrichum subrotundum. *Menz. Tr. of Linn.*
Soc. v. 4. 68. *Huds. ed.* 1. 400. *Hull.* 247.

P. pumilum. *Swartz. Musc. Suec.* 77, et 108.
t. 9. f. 19. *Hedw. Sp. Musc.* 97. *t. 21.*
f. 7—9.

P. nanum,

P. nanum, capsulis subrotundis galeritis, aloes folio non ferrato. *Dill. Musc.* 428. t. 55. f. 6. A—F.

Angl. Dwarf Round-headed Hair-moss.

In ericetis et aggeribus arenosis.

Annua? Fruct. fert hyeme et primo vere.

Radix densissimè tomentosa. Caulis brevissimus ut ferè nullus, simplicissimus. Folia numerosa, atro-viridia, rigida, crassiuscula, canaliculata, obtusa, integerrima, subtùs convexa, arefactione inflexa; inferiora quandoque apice dentato-erosa. Pedicellus haud uncialis, solitarius, rectiusculus, ruber, vel fusco-purpureus, apice incrassatus. Capsula erectiuscula, turbinata, parùm obliqua, fusca, ore dilatato. Apophysis nulla, nisi apex pedicelli incrassati. Peristomii dentes 32, tenuissimi, rufi, mox nigrantes. Operculum convexum, rubrum, mucronulo adunco. Calyptra luteo-fusca.

10. **POLYTRICHUM** foliis linearibus obtusis *nanum*. obsoletè serrulatis, capsulâ turbinatâ cernuâ, pedicello flexuoso, caule brevissimo.

Polytrichum nanum. *Hedw. Sp. Musc.* 95. *Crypt. v. 1. 35. t. 13.* *Menz. Tr. of Linn. Soc. v. 4. 69.* *Huds. 470, β.* *Hull. 247.* *Rehb. 416.* *Sibth. 306.* *Swartz. Musc. Suec. 77.* *Roth. Germ. v. 3. 351.* *Ebrh. Crypt. 54.*

P. subrotundum. *Curt. Lond. fasc. 2. t. 68.* *With. 796.* *Abbot. 232.*

P. ericoides. *Hoffm. Germ. v. 2. 24.*

P. nanum, capsulis subrotundis galeritis, aloes folio non ferrato. *Dill. Musc.* 428. t. 55. f. 6. G—L.

P. minus, capsulis subrotundis, calyprâ quasi lacerâ coronatis. *Raii Syn. 91.*

Mnium

Mnium polytrichoides α . *Sp. Pl.* 1576.
Angl. Slender Round-headed Hair-moss.

In ericetis et aridis.

Annua? Fruct. fert ineunte vere.

Præcedenti proxima at distinctissima species. Caulis vix ullus. Folia angustiora, minusque rigida, frequentius ferrulata. Pedicellus sesquiuncialis, gracilis, flexuosus, tortilis. Capsula subcernua, magnitudine ferè, et formâ, præcedentis.

alooides. II. **POLYTRICHUM** foliis lanceolatis obtusis rigidis apice ferratis, capsulâ cylindracea obliquâ, caule simplicissimo.

Polytrichum aloides. *Hedw.* *Sp. Musc.* 96.
Crypt. v. 1. 37. t. 14. *Menz. Tr. of Linn.*
Soc. v. 4. 70. *With.* 796. *Hull.* 247.
Sibth. 307. *Roth. Germ.* v. 3. 352. *Ehrh.*
Crypt. 45. *Hoffm. Germ.* v. 2. 24.

P. nanum. *Huds.* 470, α . *Lightf.* 701.

P. parvum, aloes folio ferrato, capsulis oblongis.
Dill. Musc. 429. t. 55. f. 7.

Mnium polytrichoides β . *Sp. Pl.* 1577.

Adiantum aureum medium in ericetis prove-
niens. *Vaill. Paris.* 131. t. 29. f. 11.

Angl. Dwarf Long-headed Hair-moss.

In ericetis humidiusculis.

Annua? Fruct. fert vere.

Habitus præcedentis, at magis caulescens, folia rigidiora, latiora, lanceolata, apicem versus evidenter ferrata. Pedicellus ferè sesquiuncialis, gracilis, parum flexuosus, ruber. Capsula erectiuscula, cylindracea, subobliqua, ore patentiusculo. Peristomii dentes 32. Operculum convexum, rostelllo adunco, brevi. Apophysis nulla. Carina foliorum laevis ut in præce-
dente.

I2. POLY-

12. POLYTRICHUM foliis lanceolatis obtusi-*rubell-*
usculis margine carinâque serratis, capsulâ *lum.*
cylindraceâ, caule ramoso.

Polytrichum rubellum. *Menz. Tr. of Linn.*

Soc. v. 2. 79. t. 7. f. 3. Hull. 248.

Angl. Reddish Dwarf Hair-moss.

In ericetis et locis sterilibus Angliæ et Scotiæ.
Menzies.

Perennis? Fruct. fert vere.

A præcedente differt caule plerumque ramoso, imprimis
verò foliis undique ferratis, dorsoque dentatis. Pedicelli
sesquiunciales, basi vaginati. Capsula erectiuscula.
Operculum convexum, mucrone subulato. Apophysis
nulla.

** *Calyptrâ simplici, glabriusculâ.*

13. POLYTRICHUM foliis linearibus subinte-*hercyni-*
gerimis arefactione incurvis, capsulâ *ovato-*
cum.
cylindraceâ errectâ, calyptre immaturâ pilosâ.

Polytrichum hercynicum. *Hedw. Sp. Musc. 94.*

Crypt. v. 1. 40. t. 15. With. 797. Hull.
248. Roth. Germ. v. 3. 353. Engl. Bot.
t. 1219.

Orthotrichum hercynicum. *Hoffm. Germ. v. 2.*
25.

Catharinea hercynica. *Ehrb. Crypt. 12.*

Bryum incurvum. *Huds. 479.*

Angl. Incurved Hair-moss.

In alpibus Cambrobritannicis et Scotticis.

Near Llanberris. *Rev. H. Davies.* In the High-
lands of Scotland. *Mr. Dickson.*

Perennis. Fruct. fert Julio.

Radix densè tomentosa. Caules cæspitosi, haud un-
ciales, subsimplices, foliosi. Folia rigida, linearia,
acutiuscula, canaliculata, carinata, sœpius integerrima;
VOL. III. H h interiora

interiora quandoque ferrata; omnia siccitate incurva. Pedicellus sesquiuncialis, rectiusculus, firmus, ruber. Capsula ovato-cylindracea, erecta, ore vix patulo. Peristomii dentes 32. Operculum conicum, mucronulo adunco. Calyptra cylindracea, simplex, primo ortu pilis erectis hispida, mox glabrata.

undula- 14. **POLYTRICHUM** foliis lanceolatis serratis
tum. arefactione crispis, capsulâ cylindraceâ cer-
nuâ, calyprâ apice exasperatâ.

Polytrichum undulatum. *Hedw. Sp. Musc.* 98.
Crypt. v. 1. 43. t. 16, 17. *Swartz. Musc.*
Suec. 78. *Roth. Germ.* v. 3. 354. *Rehb.*
416. *Sibth.* 307.

Catharinea Callibryon. *Ehrb. Crypt.* 83.
Bryum undulatum. *Sp. Pl.* 1582. *Huds.* 479.
With. 836. *Hull.* 265. *Abbot.* 243. *Curt.*
Lond. fasc. 1. t. 70. *Dicks. H. Sicc. fasc.* 10.
19. *Hoffm. Germ.* v. 2. 40. t. 1. *Fl. Dan.*
t. 477.

B. phyllitidis folio rugoso acuto, capsulis in-
curvis. *Dill. Musc.* 360. t. 46. f. 18.
B. capitulis oblongis rubentibus, foliis oblongis
angustis pellucidis rugosis. *Raii Syn.* 95.
Muscus erectus, linariæ folio, major. *Vaill.*
Paris. 132. t. 26. f. 17.
Angl. Undulated Hair-moss.

In umbrosis et ad saepes frequens.

Perennis. Fruct. fert Februario, Martio.
Radix ramosa, densè tomentosa. Caules sparsi, subsim-
plices, erecti, unciales vel sesquiunciales, foliosi. Folia
tenuia, lætè viridia, lanceolata, acuta; ferrata, undu-
lata, uninervia, arefactione mox crispa et tortuosa.
Pedicellus uncialis vel sesquiuncialis, rectus, ruber.
Capsula cylindracea, arcuata, cernua, fusca, apophysi
destituta, ore incrassato rubro. Peristomii dentes 32,
angusti.

angusti. Operculum hemisphæricum, mucrone filiformi, arcuato, longitudine ferè capsulæ. Calyptora cylindracea, simplex, nitida, apice quasi pilorum rudimentis denticulata seu exasperata.

ADDENDA

ADDENDA ET CORRIGENDA.

VOL. I.

p. 1. lin. 10. Post *Valerianam rubram* inferenda
Alchemilla arvensis.

5. *Chara hispida*. *Fl. Dan.* t. 154.
6. *C. flexilis*.
β. *C. translucens major flexilis*. *Vail.*
Act. Paris. Ann. 1719. t. 3. f. 8.
In Berrington pool, Shropshire. *Rev. E. Williams.*
14. *Circæa alpina* β. *C. intermedia*. *Ehrh.*
Herb. 101.
24. *Veronica hederifolia*. Winter weed, Nor-
folciensisibus. *D. Crowe*. Vix ultra
Maium flores profert.
27. *Pinguicula lusitanica*. Very common near
Newtown Barry, Ireland. *Rev. Mr. Butt.*
- 34, 75 et 77. *Characteres Milii et Agrostidis*
emendandi.
MILIUM. Cal. bivalvis, uniflorus, tumidus.
Semen corollâ persistente, cartilagineâ,
bivalvi, corticatum.

AGROSTIS.

AGROSTIS. *Cal.* bivalvis, uniflorus: valvulis acutis. *Cor.* bivalvis, inæqualis, membranacea. *Stigmata* plumosa.

39. Post Valerianam *Locustam*.

5. **VALERIANA** floribns triandris, caule di-*dentata*. chotomo, foliis linearis-lingulatis, fructu ovato acuminato inæqualitè tridentato.

Valeriana dentata. *Ehrb.* *Herb.* 122. *Willden. Sp. Pl.* v. 1. 183; exclusis minimùm synonymis Linnæi, Raii et Rivini.

Valerianella altera, nudo umbilicato et lævi semine, *Column. Ecphr.* 208. t. 209. f. 2.

Angl. Oval-fruited Corn Sallad.

In arvis.

Found in Cornwall by Mr. E. Forster, jun. in 1799.

Annua. Fl. autumno.

Habitus omnino V. *Locustæ*. Discrepat verò paniculâ iterum iterumque dichotomâ, cum flore ad omnem dichotomiam sessili; fructu ovato, hinc gibbo, lævigato, illinc umbilicato, apice tridentato, dentibus maximè inæqualibus,

41. lin. 20. pro *petalis* lege *laciniis*.

43. *Schœnus Mariscus.* Flores diandri. Stigma bi- tri- seu quadrifidum. *D. Sowerby.* Fructus drupa, semine elliptico, triangulare. Confer *Engl. Bot.* v. 14. t. 950.

44. *S. nigricans.* Semina alba, nitida. *Ehrhart.* *Beitr.* v. 4. 48, confirmante D. Wigg. *S. compressus.* Radix subrepens.

45. *S. rufus.* On the coast to the east of H h 3 Dunbar.

Dunbar. *Mr. J. Mackay.* In Fifeshire and Angus-shire. *Mr. G. Donn.* Near the bridge between Bootle and Crosby rabbit-warren between Liverpool and Ince. *Mr. J. Shepherd.*

47. *Cyperus longus.* At Walton in Gordano, Somersetshire. *Mr. Dyer.*

49. *Scirpus multicaulis.*

S. palustris. *Ebrh. Calam.* 61.

Not uncommon in various parts of Britain.

57. *S. sylvaticus.* Pro *Relb.* lege *Sibth.* 24.

61. *Nardus stricta.* *Cavan.* *Ic.* v. 3. 224. t. 204. f. 2.

62. *lin.* 18, pro *foliacei* lege *foliosi.*

64. *Panicum verticillatum.* Involucris retrorsum scabris, optimè monente Curtisio, a *P. viridi* certè dīgnoseitur.

In a field out of St. Giles's gates, Norwich.

67. *P. dactylon.* In charactere et descriptione dele *calycibus corollæ alternis.*

73. *Ante Alopecurum agrestem;*

alpinus. 2. *ALOPECURUS culmo erecto lœvi, spicâ ovatâ, glumis villosis muticis aristæ eocollinæ subæqualibus.*

Alopecurus alpinus. *Engl. Bot.* t. 1126.

Angl. Alpine Fox-tail-grass.

In alpibus Scoticis.

On mountains about Loch Nagore, Aberdeenshire. *Mr. G. Donn.*

Perennis. Fl. . . .

Radix

Radix repens, nigricans. Culmi pedales, erecti, basi geniculati, simplices, foliosi, glaberrimi. Folia radicalia linearia, angusta; caulina lata, acuta, abbreviata, glabra, vaginis elongatis, parùm ventricosis. Stipula brevissima. Spica haud uncialis, ovata, obtusa, in lobos fissilis. Calyx ovatus, purpurascens, undique villosus, glumis subretusis, basi connatis. Corolla elliptica, nuda, carinâ tantùm scabra, calyce parùm minor, basi aristata. Arista subtortilis, scabra, purpurea, vix calyce longior.

75. *Milium effusum*; planta perennis, nec annua.
78. *Agrostis littoralis*. On the Essex coast.
Dicks. H. Sicç. fasc. 16. 1. Near the powder magazine about 4 miles from Woolwich. *Mr. George Jackson. Engl. Bot. t. 1251.*
 Synonymon *Witheringii* excludendum.
82. *A. minima*, *KNAPPIA agrostidea*. *Engl. Bot. t. 1127.*
CHAR. GENERIS. *Cal.* bivalvis, uniflorus, obtusus, muticus. *Cor.* univalvis, truncata, hirsuta, mutica. *Flores* in rachi flexuosâ, alterni.
101. *Poa flexuosa*. *Engl. Bot. t. 1123.*
102. *P. bulbosa*. *Engl. Bot. t. 1071.* On sandy ground near the sea at Yarmouth. *R. Stone Esq.* Also at Lowestoft.
103. *P. cæsia*. Syn. *Witheringii* delendum.
105. Post *Poam pratensem*.
14. *POA paniculâ diffusâ*, spiculis ovatishumilis, subtrifloris, glumis acutis villo connexis, stipulâ brevissimâ obtusâ.

Poa humilis. *Ehrb. Calam.* 115. *Beitr.* v. 6. 84.

P. subcærulea. *Engl. Bot. t.* 1004; excl.
syn. *Witheringii.*

P. pratensis β , *alpina.* *Huds.* 39.

P. alpina γ . *Willden. Sp. Pl. v.* 1. 386.

P. n. 1456, γ . *Hall. Hist. v.* 2. 221.

Gramen alpinum paniculatum elegans, pa-
niculis ex spadiceo viridi et argenteo va-
riegatis. *Scheuchz. Prod.* 20. t. 3. *It.*
v. 2. 457. f. 17.

Angl. Short Blueish Meadow-grass.

In pascuis montosis.

In Westmoreland and Cumberland. *Huds.*
Anglesea. *Rev. H. Davies.*

Perennis. Fl. Junio.

Radix repens, fibris villosis. Herba glauca, pur-
purascens. Culmi palmates aut spithamei, ad-
scendentes, simplices, foliosi, supernè nudi. Fo-
lia brevia, patentia, recta, obtusiuscula, margine
scabra. Vaginæ elongatæ, compressæ, striatæ,
glabræ. Stipulæ omnes brevissimæ, obtusæ.
Panicula diffusa, pedicellis capillaribus, scabri-
usculis. Spiculæ erectæ, latè ovatæ, sæpiùs tri-
floræ. Glumæ calycinæ inæquales, acuminatæ,
trinerves, basi virides, marginem versùs purpu-
reæ. Flosculi acuti, nervosi, basi virides, medio
purpurei et spadicei, margine albi, scariosi, om-
nes villis copiosis, longissimis, connexi.

A Poâ alpinâ distinctissima flosculis villo connexis,
stipulis omnibus brevissimis, radice repente.

106. Ante Poam nemoralem.

glauca. 16. POA paniculâ attenuatâ erectâ, spiculis
ovatis subtrifloris, glumis subretusis basi
villosis, stipulâ brevissimâ.

Poa glauca. Fl. Dan. t. 964. *With.* 148.
P. n.

P. n. 1468. *Hall. Hist.* v. 2. 224; ex auctoritate D. Lachenalii.

Gramen paniculatum angustifolium montanum, paniculâ densâ, lccustis parvis muticis. *Scheuchz. Agr.* 180.

Angl. Slender Glaucous Meadow-grass.

In montibus Cambricis.

On Snowdon. *Mr. Griffith.*

Perennis. Fl. Junio, Julio.

Radix fibrosa. Herba glauca. Culmi pedales, stricti, graciles, foliosi, striati, glabri. Folia angusta, erecto-patentia, acuta, margine scabra. Vaginæ longitudine ferè foliorum, subcomprese, striatæ. Stipula brevissima, retusa, crenata. Panicula erecta, coarctata, pedunculis angulatis, scabris, subflexuosis. Spiculæ erectæ, ovatæ, seu ovato-lanceolatæ, glaucæ, demùm purpureo-spadiceæ. Glumæ calycinæ parùm inæquales, acutæ, carinatæ, læves; major trinervis. Flosculi bini vel terni, lanceolati, subretusi, vix nervosi, basin versùs villosi, haud villis complicatis connexi, margine purpureo alboque variati, scariosi. E speciminibus ab amicissimo D. Griffithio benevolè communiciatis.

A Poâ nemorali differt habitu minùs gracili, colore glauco, paniculâ strictâ, flosculis obtusioribus, coloratis, vix nervosis.

121. *Festuca calamaria.* *Engl. Bot.* t. 1005.

Poa trinervata. *Schrad. Spicil.* 3. *Willden.*

Sp. Pl. v. 1. 389.

In Dumbartonshire and Breadalbane, also near Fort Augustus. *Mr. G. Donn.* In Shrawley wood near Glashampton, Worcestershire. *Mr. Moseley.* In a wood near Newtown Barry, Ireland. *Rev. Mr. Butt.*

Obs.

Obs. Glumæ corollinæ interiores potius bicarinatæ sunt, carinis approximatis.

140. Dele lin. 15 et 16, et infere—

strigosa. 2. AVENA paniculâ oblongâ secundâ, calycibus bifloris, flosculis apice biaristatis.

Avena strigosa. *Engl. Bot. t. 1266. Schreb.*

Lips. 52. Retz. Obs. fasc. 1. 11. With.

166. *Hull. 26. Willden. Sp. Pl. v. 1.*

446. *Host. Gram. v. 2. 41. t. 56.*

Angl. Bristle-pointed Oat.

Inter segetes.

Annua. Fl. Julio.

Magnitudo Avenæ sativæ. Panicula oblonga, secunda, subcoarctata. Glumæ calycinæ subæquales, longitudine floscularum, nervosæ, scarifosæ, muticæ. Flosculi bini, paralleli, magnitudine æquales, pilosi, dorso aristati, aristâ geniculatâ, apice biaristati, aristis triplò ferè brevioribus, rectis. Glumæ interiores muticæ.

150. Lolium arvense. Aristæ apud *Engl. Bot. t. 1125* nimis validæ.

157. Triticum junceum. Dele syn. Relhani.

193. Potamogeton natans, foliis omnibus ellipticis petiolatis natantibus, petiolis inferioribus submersis aphyllis.

hetero-
phyllum:

2. POTAMOGETON foliis submersis membranaceis lineari-lanceolatis sessilibus; natantibus coriaceis ellipticis petiolatis, pedunculis incrassatis.

Potamogeton heterophyllum. *Schreb. Lips.*

21. *Hoffm. Germ. ann. 1800. 81.*

Willden. Sp. Pl. v. 1. 713. Engl. Bot. t. 1285.

P. pa-

P. palustre. *Teesdale Tr. of Linn. Soc.*
v. 5. 43. *Relb.* 64.

P. graminéum. *Fl. Dan.* t. 222: caret
foliis natantibus.

Angl. Various-leaved Pond-weed.

In fossis et stagnis.

In ditches near Beverley, Yorkshire. *Mr.*
Teesdale. Cambridgeshire. *Relb.* and *Rev.*
Mr. Hemsted. Berrington pool, Shrop-
shire. *Rev. Mr. Williams.* At Old Buck-
enham, Norfolk. *Mr. D. Turner.* Near
Forfar, Scotland. *Mr. G. Donn.*

Perennis. Fl. Julio—Septembri.

Radix repens, stolonifera, alba. Caules ramosi-
simi, divaricati, teretes, undique foliosi. Folia
integerrima, lætè viridia, tenuè nervosa; sub-
mersa numerosa, alterna, patentia, linear-lan-
ceolata, acuta, membranacea, sessilia; summa
natantia, ferotina, petiolata, elliptica, mucronu-
lata, latitudine varia. Stipulae intrafoliaceæ, ob-
longæ, obtusiusculæ. Pedunculi apicem versùs
caulis, sæpè conferti, erecti, sursùm incrassati.
Bracteæ stipulis latiores, carinatæ, concavæ.
Spica simplex, virens.

3. POTAMOGETON foliis inferioribus lan-*fluitans.*
ceolatis acuminatis membranaceis; su-
perioribus elliptico-lanceolatis subcoria-
ceis: omnibus petiolatis.

Potamogeton fluitans. *Roth. Germ.* v. 1.
72. v. 2. 202. *Beitr. fasc.* 2. 126. n. 4.
Hoffm. Germ. ann. 1800. 81. *Willden.*
Sp. Pl. v. 1. 713. *Engl. Bot.* t. 1286.
Angl. Long-leaved Floating Pond-weed.

In rivulis lentè fluentibus.

In ditches in marshy ground near Beverley.
Mr. Teesdale. Lilleshall mill-pool, Shrop-
shire.

shire. *Rev. Mr. Williams.* At Scole, Norfolk: *Mr. Woodward and Mr. D. Turner.*

Perennis. Fl. Julio, Augusto.

Herba tota ferè immerfa, foliis superioribus tantum subnatantibus, spicisque emersis. Radix repens. Caules fluitantes, longissimi, ramosi, foliosi. Folia rufo-viridia, integerrima, tenuè nervosa, alterna, plùs minùs petiolata; inferiora membranacea, lanceolata, acuminata, elongata, sèpè pedalia; superiora breviora, latiora, subelliptica, longius petiolata, magisque coriacea. Stipulæ latè lanceolatæ. Pedunculi axillares, solitarii, longitudine ferè foliorum, parùm incrassati. Spica densa, rufescens.

201. *Tillæa muscosa.* *Engl. Bot.* t. 116.

205. lin. 15. et 242. lin. 8. *Lobelia.* *Stigma capitatum,* nec simplex.

lin. 19. dele *bilocularis.*

213. lin. 27. *Lege calycem.*

218. *Pulmonaria maritima.* Herba quandoque viridis, nec glauca. *D. J. Mackay.*

226. lin. 4. *Lege exsertus.*

235. lin. 6. *Sibth.* 80, nec 248.

242. *Lobelia Dortmanna.* In the lake of Killarney, Ireland. *Rev. Mr. Butt.*

243. *Impatiens Noli me tangere.* In ordine naturali forsitan Corchoro affinis. Foliorum ferraturæ inferiores setaceæ. Folia in horto nostro noctu patentia, nequaquam flaccida, vel dependentia.

257. *Chironia Centaurium.*

Hujus varietates duas valde notabiles, in arenosis maritimis prope *Liverpool* cum vulgari vigentes, invenerunt D. Bostock M. D. et D. Shepherd, Hort. Botanici Liverpoolensis curator indefessus.

Harum prior cum vulgari certat magnitudine, florum formâ, et totâ facie; discrepat verò foliis angustis, spatulato-linearibus, laciniisque calycinis elongatis, angustatis, corollæ tubum superantibus. Altera longè humilior est, foliis latissimis, ferè orbiculatis, floribus parvis, congestis, laciniis calycinis etiam tubo longioribus, at minus quam in priore angustatis. Anne varietates vel species distinctæ?

261. Post *Lonicera Periclymenum*.

3. LONICERA pedunculis bifloris; baccis *Xylosteum*. distinctis, foliis integerrimis pubescens.

Lonicera Xylosteum. *Sp. Pl.* 248. *Willden.* *Sp. Pl.* v. 1. 986. *Børk. Outlin.* ed. 1. v. 2. 60. ed. 2. v. 2. 73. *With.* 247. *Sym. Syn.* 59. *Hull.* 53. *Engl. Bot.* t. 916. *Fl. Dan.* t. 808.

Xylosteum. *Riv. Monop. Irr.* t. 120.

Periclymenum rectum germanicum. *Ger.* *em.* 1294.

Angl. Upright Honeysuckle.

In dumetis siccioribus.

In the fissures of rocks under the Roman wall near Shewing-Sheels, Northumberland. *Wallis.* Edgbaston park near Birmingham. *With.* Plentifully, and certainly wild, in a coppice called the Hacketts, to the east of Houghton bridge, four miles from Arundel, Sussex. *Mr. W. Borrer.*

Frutex.

Frutex. Fl. Julio.

Frutex rectus, humilis, ramosissimus. Rami oppositi, teretes; juniores pubescentes, foliosi. Folia opposita, petiolata, ovata, acutiuscula, integerrima, mollia, pubescentia, decidua. Pedunculi axillares, oppositi, biflori, foliis breviores. Bracteæ exteriore geminæ, lanceolatæ, pilosæ; interiores minores, concavæ, obtusæ, solitariæ sub singulo flore. Flores lactei vel rubicundi, pubescentes, tubo brevi. Bacca rubra, subrotunda, discreta, succosa, unilocularis, subhexasperma.

282. *Cuscuta europæa.* Dele syn. Dicksoni.
295. *Tordylium maximum.* Engl. Bot. t. 1173.
Near Eton. Mr. Gotobed.
307. *Heracleum angustifolium* excludendum.
Planta Teesdalii est H. Sphondylii varietas, ab H. angustifolio Linnæi distincta.
325. Ante Scandicem *Anthriscum.*
2. SCANDIX seminibus nitidis ovato-subulatis, umbellis sessilibus lateralibus.
Scandix Ceresfolium. Sp. Pl. 368. Willden.
Sp. Pl. v. 1. 1450. With. 307. Hull.
63. Jacq. Austr. v. 4. 47. t. 390. Engl.
Bot. t. 1268.
Ceresfolium. Riv. Pent. Irr. t. 43.
C. vulgare sativum. Ger. em. 1038.
Gingidium. Fuchs. Hist. 216.
Angl. Garden Chervil.

Ad sepes.

About Worcester. Dr. Stokes. On a bank near Halesworth, Suffolk, plentifully. Mr. L. Wigg.

Annua.

Ceresfo-
lium.

Annua. Fl. Junio.

Habitus S. Anthrisci, sed foliis paululùm latoribus, glabrioribus, seminibusque lèvibus differt. Radix fusiformis. Herba tenera, nitida, pallida, dulcis et gràte aromatica. Caulis pedalis vel bipedalis, teres, ramosus, striatus, geniculis supernè pubescentibus, cæterùm glaber. Folia patentia, bipinnata, incisa; inferiora tripinnata, et longius pectiolata. Umbellæ laterales et terminales, sessiles, subquadriradiatæ, basi lanatæ. Umbellulæ ferè decemfloræ. Involucrum nullum. Involucella dimidiata, acuta. Flores regulares, albi. Semina lanceolata, glaberrima, striata, rostro brevissimo.

344. *Linum angustifolium.* In a field at Alerton hall, the seat of Mr. Roscoe, near Liverpool. *Mr. Shepherd.*

349. lin. 12. Peplis. Lege Cal. duodecimfidus.

384. Post *Juncum pilosum.*

17. *JUNCUS* foliis planis pilosis, paniculâ *Forsteri.* cymosâ erectâ, floribus solitariis, capulis acutis.

Juncus Forsteri. *Engl. Bot.* t. 1293.
Angl. Narrow-leaved Hairy Rush.

In nemorosis.

Under trees between Hoghill and Collier row in Hainhault forest, Essex, along with *J. pilosus*, first observed in 1795. *Mr. Edw. Forster, jun.* In a wood opposite the south front of Hafod house, Cardiganshire.

Perennis. Fl. Maio.

Radix fibrosa, nigra, vix stolonifera. Culmi spithamei vel pedales, erecti, simplices, graciles, teretes, glabri, foliosi. Folia linear-lanceolata, acuta, duplò vel triplò angustiora quam in *J. pilosus*, margine basin versùs pilosa; superiora magiora;

jora; radicaria minima. Pânieula terminalis, cymosa, ramosa, ramis omnibus erectis, subpatentibus, nunquam divaricatis vel refractis. Flores J. pilosi similes, sed lacinii ealycinis longius acuminatis. Capsula ovata, acuta, mucronulata, nec obtusa.

Species certè nova et distinctissima; in horto immutata manet.

422. lin. 29. Lege Socio.

433. Delenda omnino est Elatine *Alsinastrum*; specimen in Herb. Dill. est Centunculus *minimus*, monente Sibthorpio in manuscriptis! In loco ejus inserenda est

1. ELATINE foliis oppositis, caule radicante ramosissimo.

Elatine Hydropiper. Sp. Pl. 527. Willden. Sp. Pl. v. 2. 472. Fl. Dan. t. 156. Roth. Germ. v. 1. 175. v. 2. 458. Engl. Bot. t. 955.

Hydropiper. Buxb. Cent. 2. 36. t. 37. f. 3.

Alsinastrum serpyllifolium, flore roseo tripetalo, et flore albo tetrapetalo. Vail.

Paris. 5. t. 2. f. 1, 2.

Angl. Small Waterwort.

In lacuum marginibus arenosis inundatis.

About the eastern shore of Bomere pool near Condover, Shropshire. Rev. Mr. Williams.

Annua. Fl. Augusto.

Habitus Montiae fontanæ. Radix fibrosa, alba. Herba glaberrima, saepius immersa. Caules ramosissimi, diffusi, radicantes, foliosi, biunciales. Folia opposita, spatulata, integerrima. Flores axillares, solitarii, pedunculati, albi vel rosei, plerumque clausi, et, apud nos, trifidi, hexandri, trigyni; rarius, in eadem stirpe, quadrifidi, octandri, tetragyni. Semina numerosa, exigua,

Hydro-
piper.

exigua, oblonga, curvata, angulata; transversim pulchrè striata.

VOL. II.

455. lin. 16. *Saxifraga cæspitosa*. Syn. Scopolii excludendum. Confer *Engl. Bot.* t. 794.

467. *Silene paradoxa*.

Millerus, pro Lychnide majore noctiflorâ Dubrensi perenni Raianâ, in horto Chelseano coluit, et cum Linnæo communicavit, Silenem paradoxam; ut ex Herb. Linn. et Mus. Brit. (*H. Sicc.* n. 294. 30.) patet. At in Herbariis Buddlei et Petiveri, in eodem museo (*H. Sicc.* n. 124. 5, et n. 152. 86.) repositis, specimina Silenis cuiusdam, prorsus diversæ et mihi ignotæ, in ipso agro Dubrensi, ut videtur, lecta, nupèr vidi. Hæc verò, male conservata, ad novam generis tam difficilis speciem describendam, haud optimè valent; et in eorum loco natali Silene nutans tantum hodie observatur. Quid igitur faciam? Confusio pristina manebit, donèc planta Raiana botanicis de novo sese offerat.

470. *Silene conica*. Dele synonyma Lobelii et Gerardi, et hoc insere.

Lychnis caliculis striatis, secunda Clusii.

Ger. em. 470.

475. Post *Stellariam gramineam*.

5. STELLARIA caule pedunculis breviore, *scapigera*-
foliis linear-lanceolatis margine scabris, ra-
calyce trinervi petalis subæquali.

Stellaria scapigera. *Willden. Sp. Pl.* v. 2.

716. *Engl. Bot.* t. 1269.

Angl. Many-stalked Stitchwort.

Ad rivulos Scotiæ.

In Perthshire, and about Loch Nevs, Inverness-shire. *Mr. G. Donn.*

Perennis. Fl. æstate.

Caules brevissimi, cæspitosi, densè foliosi. Folia haud glauca, opposita, linear-lanceolata, acuta, glabra, uninervia, (nec trinervia,) nervo basi crassissimo, apicem versùs evanescente, margine crenulato-scabra. Pedunculi numerosissimi, axillares, erecti, caulem longè superantes, plerumque biunciales, tetragoni, glabri, sæpiùs simplices. Flores albi, inconspicui. Calycis foliola trinervia, acuta, margine membranacea, longitudine scè petalorum. Antheræ rubræ.

477. *Stellaria cerastoides.*

Caules lincâ pilosâ, longitudinali, notati, ut in *Stellariâ mediâ*. Pedunculi undique pubescentes. Confer *Engl. Bot.* t. 911.

480. lin. 3. *Lege subfusiformis.*

483. lin. 6. *Lege Segu. Veron. Suppl.* 180. t. 4. f. 3.

534. *Pyrus hybrida.*

In rocky places on Cairn na Callich and other mountains at the north end of the isle of Arran. *Mr. J. Mackay.*

538. Post *Rosam spinosissimam.*

involuta. 2. *ROSA* fructibus globosis pedunculisque aculeatissimis, aculeis caulinis numerosissimis rectiusculis, petalis involuto-clausis.

Angl. Prickly Unexpanded Rose.

In insulis occidentalibus Scotiæ. *D. Walker and D. J. Mackay.*

Frutex. Fl. Julio.

Habitus

Habitus R. spinosissimæ. Caulis undique aculeatus, aculcis magnitudine variis, quandoque subaduncis. Foliola 7 vel 9, elliptico-subrotunda, obtusa, ferrata, glabra, nervo subtus hirsuto. Petioli aculeati. Pedunculi terminales, solitarii, recti, uniflori, aculeatissimi. Germen globosum, minimè depresso, aculeis densis, divaricatis, rigidis, undique armatum. Calycis foliola acuminata, simplicia, extus muricata. Petala albidæ, rubro variata, involuto-clausa, nec patula.

568. Papaver *sommiferum*. Petala sœpè dilutè violacea, cum maculâ saturatiore.

Certainly wild on the banks of all the fen ditches, where the soil is sandy, in the parish of Hockwold cum Wilton, Norfolk. Rev. Mr. White.

- 604: Post Ajugam *reptantem*.

2. AJUGA pilosa, verticillis pyramidato-pyramidalis confertis multifloris, soliis radicalibus maximis obovatis obtusis crenatis.
- Ajuga pyramidalis. Sp. Pl. 785. Willden. Sp. Pl. v. 3. 8. Lightf. 302. With. 516. Hull. 131. Fl. Dan. t. 185. Engl. Bot. t. 1270.

Angl. Pyramidal Bugle.

In montibus Scoticis.

On Tore Aichaltie, a hill near Brahan castle, the seat of Lord Seaforth. Mr. Wm. Gibb.

Perennis. Fl. Maio, Junio.

Radix tuberosa, fibris longis. Herba undique plus minus pilosa aut villosa. Caulis solitarius, erectus, tetragonus, 2-5-uncialis, stolonibus destitutus. Folia numerosa, omnia ferè radicalia, petiolata, magna, obovata, obtusa, subcrenata, rubicunda. Verticilli plurimi, approximati, multiflori. Bracteæ magnæ, obtuse, incisæ, pallidè

purpureæ. Corolla purpurea, lineis saturationibus, palato flavo.

621. *Mentha gentilis*. About several villages in Shropshire. *Rev. Mr. Williams.*
622. *M. gracilis* β. *M. pratensis*, Sole. Est varietas *Menthæ rubræ*. Conf. *Tr. of Linn. Soc. v. 5. 275.*
624. *M. arvensis* δ. At Prestwick car, Northumberland. *Mr. Winch.*
648. *Bartsia viscosa*. *Engl. Bot. t. 1045.* At Allerton near Liverpool. *Mr. Robert Roscoe.*
651. *Melampyrum cristatum*. In Maple-bush lane, Gresenhall, Norfolk. *Mr. Crowe.*
677. Ante *Drabam vernam*.
- aizoides.* 1. *DRABA* scapo nudo, petalis emarginatis calyce duplò longioribus, foliis lanceolatis rigidis carinatis ciliatis.
Draba aizoides. *Linn. Mant. 91. Willden. Sp. Pl. v. 3. 424. Jacq. Austr. v. 2. 55. t. 192. Curt. Mag. t. 170. Engl. Bot. t. 1271.*
Moenchia aizoides. *Roth. Germ. v. 1. 273. v. 2. p. 2. 72.*
Sedum petræum. *Ger. em. 514.*
Angl. Yellow Alpine Whitlow-grass.
 In muris et rupibus maritimis Cambriæ australis.
 Abundantly on walls and inaccessible rocks about Fennard castle, 8 miles west of Swansea. *Dr. W. Turton.*
Perennis. Fl. Martio.

Radix

Radix fibrosa, multiceps. Folia omnia radicalia, cæspitosa, numerosa, densa, rosacea, persistentia, linear-lanceolata, rigida, nitida, carinata, uninervia, ciliata, ciliis albis, rigidis. Scapi solitarii, terminales, simplices, nudi, rigidi, teretes, glabri, sesquiunciales, corymboso-racemosi. Calycis foliola latè elliptica, concava, patentiuscula. Petala aurea, obovata, obtusa, subemarginata, calyce duplè longiora. Silicula elliptica, plana, acuta, glabra, polysperma, stylo elongato, persistente.

679. *Draba muralis.* *Scop. Insubr. fasc. 2. t. 15.*
Engl. Bot. t. 912.

Fl. Aprili, Maio.

684. *Thlaspi hirtum.* Dele signum †.

In a field below the garden at the Earl of Kinnoul's seat near Perth. *Mr. Miller.*

696. *Dentaria bulbifera.* On the High Rocks at Tunbridge wells. *Rev. Mr. Bale and Mr. Dickson.*

700. lin. 10. *Lege Stylus*, monente Curtisio, obliquus.

704. *Ante Sisymbrium monense.*

6. *SISYMBRIUM* siliquis erectis, foliis lan-

murale. caelatis inciso-ferratis, caulis paten-

tibus scabriusculis.

Sisymbrium murale. *Sp. Pl. 918. Willden.*

Sp. Pl. v. 3. 496. Dicks. Dr. Pl. 12.

Engl. Bot. t. 1090.

Eruca viminea, iberidis folio, luteo flore.

Barrel. Ic. t. 131.

Angl. Sand Rocket.

In arenosis et ruderatis maritimis.

Common throughout the isle of Thanet, particularly about Rainsgate. *Mr. Dillwyn.*
Below Bristol. *Mr. Edw. Forster.*

Annua. Fl. autumno.

Radix ramosa. Caules patentes, subramosi, foliosi, teretes, seabriuseuli, fetis deflexis; quandoque brevissimi, et ferè nulli. Folia alterna, petiolata, latè lanceolata, inæqualitèr sinuato-serrata, læviuseula, viridia, nec glauca. Peduncles terminales, longissimæ, corymboso-racemosi. Flores lutei, duplò quam in S. tenuifolio minores. Calyx glaber. Siliqua erecta, græcilis. Pedunculi et pedicelli sæpè hispidiusculi.

707. *Erysimum præcox.* *Engl. Bot. t. 1129.*

About Teignmouth, Dawlish, Kingsteinton, and Honiton, Devonshire. *Rev. Dr. Beeke.*

708. *E. cheiranthoides.* Not rare in Norfolk. *Mr. Crowe.*

710. *Cheiranthes fruticulosus* à Ch. Cheiri differt etiam petalis recurvis, rigidulis, nec flaccidis, laxè dependentibus. *D. Crowe.*

724. *Raphani Raphanistri* varietatem γ jampridem e Scotiâ accepi. Fortè species distincta.

725. lin. 9. Inscratur *Lysimachia vulgaris.*

734. Ad calcem, post *Vaill. Par. t. 15. f. 1.* *Raii Syn. 358. Hist. v. 2. 1059.*

736. *Geranium rotundifolium.* Dele Syn. Raii.

751. *Fumaria capreolata.* *Engl. Bot. t. 943.*
A very common weed every where about Liverpool.

762. *Orobus sylvaticus.*

Hujuscce plantæ varietatem, ut videtur, quam maximè singularē et inauditam, foliis simpli-cibus magnis gaudentem, at nunquam florentem, prope *Hafod*, Cambriæ, invenit D. Todd; mecumque communicavit Domina Mariamne Johnes.

764. *Lathyrus Nissolia.* At Wormleybury, Herts. *Miss Hume.*788. *Trifolium scabrum.* Not at Caister, but on Snettisham beach, and in an old chalk-pit near Wells, Norfolk. *Mr. Crowe.*792. lin. 1. *Lege deflexis.*

15. *TRIFOLIUM* spicis ovalibus multiflo-*procum-
ris*, vexillis sulcatis, caulis*procumbens*.
bentibus, petiolo communi basi elongato.
Synonymis addend. *Engl. Bot. t. 945.*

16. *TRIFOLIUM* spicis capitatis hemi-*minus*, sphæricis, pedunculis strictis, vexillis læ-
viusculis, caulis prostratis, petiolo
communi basi brevissimo.

Trifolium minus. *Relb. 290. Engl. Bot.
t. 1256.*

T. filiforme β. *Fl. Brit. 793;* cum sy-
nonymis.

Angl. Lesser Yellow Trefoil.

In pascuis glareosis.

Annua. Fl. Junio, Julio.

Radix fibrosa, parva, sæpè nodulosa. Caules pro-
cumbentes, subsimplices, teretes, pilosiusculi,
longitudine variis, spithamæ aut bipedales; "in
varietate majori fragiles et succosi," *D. Beeke.*
Petioli communes basi brevissimi, supernè elon-
gati.

gati. Foliola obcordata, dentata, striata; lateralia ferè sessilia. Stipulæ semiovatae. Pedunculi axillares, recti, sirmi, rigidi. Flores 12-15, capitati, subsessiles, flavi, demum fuscescentes. Dentes calycini valde inæquales, apice pilosi. Vexillum obsoletè fuleatum. Legumen obovato-ellipticum, monospermum, rariùs dispermum.

filiforme.

17. *TRIFOLIUM* spicis paucifloris laxis pedunculis flexuosis, vexillis lœvibus, caulinibus prostratis, foliolis omnibus subsessilibus.

Trifolium filiforme. *Rell.* 290. *Engl. Bot.* t. 1257. *Fl. Brit.* 792, α ; cum synonymis.

Angl. Slender Yellow Trefoil.

In pascuis glarcosis, graminosis.

Annua. Fl. Junio, Julio.

Præcedente pleniusque minus. Radix parva, sœpè nodulosa. Caules procumbentes, filiformes, gracilissimi, sœpiùs palmares, rariùs spithamei, glabri, vel apicem versus subpilosí, petioli communes breves. Foliola omnia subsessilia, parva, glabra. Stipulæ ovatae, parvae. Pedunculi capillares, flexuosi, laxi, pilosuscili. Flores 3—8, laxè penduli, secundi, pedicellis tenuissimis. Corolla flava, demum albida, vexillo lœvi. Dentes calycini inæquales, glaberrimi. Legumen obovatum, monospermum vel dispermum.

802. *Hypericum dubium.* At North Mimms, Hertfordshire. *Mr. Sabine.*

807. lin. 16. *Lege substipitatus.*

819. lin. 22. *Lege Pet. f. 1.*

830. *Hieracium mirorum.* Post varietatem γ infere

δ. Broad dented Hawk-lung. *Pet. H. Brit.* t. 13. f. 3; excluso synonymo Raii.

On Ben Gloc, and other mountains of Scotland. *Mr. J. Mackay.*

Folia dentato-incisa, immaculata. Caules spithamei, pilosi, subaphylli. Flores duplò quam in reliquis varietatibus majores, calyce villoso, atrō. Anne species distincta?

831. *Hieracium sylvaticum*, prorsùs excludendum. Planta apud *Ger. em.* 304 est *Cineraria integrifolia*, ut è speciminiibus à D. Lamb et D. Bicheno in loco natali lectis, et mecum benevolè communiciatis, satis patuit.
833. *Hieracium villosum*.
Hujus synonymon est *H. ovatum*, *Hull.* 175, ex auctore anonymo in *Month. Mag. v.* 16, 532.
848. *Carduus nutans*. *Raii Syn.* 193.
868. *Gnaphalium margaritaceum*. By a rivulet in the heart of Wire forest, Worcestershire. *Rev. Mr. Butt.* Found by Dr. Salt in a meadow at Longdon near Litchfield. *Lord Visc. Valentia.*
Senecio aquaticus.
Jacobaea latifolia palustris sive *aquatica*. *Raii Syn.* 178.
900. *Pyrethrum Parthenium*. Flores lutei, radio albo.

VOL. III.

935. *Ophrys Loeselii*.
Cymbidium Loeselii. *Swartz. in Nov. Act. Upsal. v. 6.* 76.
965. *Carex*

965: *Carex Davalliana*. On Lansdown hill near Bath, found there in 1800 by Mr. Groult in company with Mr. Lambert.

1005. *C. lœvigata*.

C. æthiopica. Schkubr. Car. 107. t. Z. f. 83.

In a boggy thicket to the west of Warley Common, Essex; also in Great Shrub-bush, a wood near Wanstead. Mr. E. Forster.

1011. *Littorella lacustris*. Variat foliis hirtis. D. E. Williams.

1053. *Salix malifolia*. Varietas videtur *S. hastata* Linnæi.

1063. *S. cinerea*. In a wood by the road side near Bromley, Kent. Mr. E. Forster.

1150. *Phascum patens*. Fruct. fert Augusto.

1190. *Grimmia cirrata*. Syn. Abbot. 238 est præcedentis.

1228. Post *Dicranum strumiferum*.

virens. 41. *DICRANUM* caule ramoso, foliis lineari-lanceolatis siccitate tortilibus, capsulâ arcuatâ lævi basi strumosâ.

Dicranum virens. Swartz. Musc. Suec. 33.

Hedw. Sp. Musc. 142. Crypt. v. 3. 77. t. 32.

Bryum virens. Swartz. Act. Nov. Upsal.

v. 4. 241; exclusis omnino synonymis.

Angl. Green Spur-fruited Fork-moss.

In montibus micaceis Hiberniæ et Scotiæ.

In Ireland. Dr. Scott. On Ben High and

Ben Lawers, also on the mountains of Corum Gorum in Inverness-shire. *Mr. G. Donn.*

Perennis. Fruct. fert æstate.

Caules biunciales vel palniares, ercti, dñsè cæspitosi, undique foliosi, infernè nigrantes, subtomentosi. Folia pulchrè et lætè virentia, linearilancolata, integerrima, subrevoluta, carinata, uninervia; basi dilatata, amplexicaulia; apice subulata, clongata, patula; siccitate incurvato-tortuosa. Pedicelli terminales, solitarii, vix unciales, graciles, virentes, demùm fusci; basi foliis valdè dilatatis vaginati. Capsula ferruginea, cylindracea, arcuato-incurvata, lævis, nec fulcata, basi attenuata, apophysi acutâ, hinc porrectâ, quasi calcarata. Calyptra pallidè vircns, basi constricta, apice nigra. Operculum subulato-conicum, capsulâ brevius. Peristomium rubrum.

A. D. pellucido differt capsulâ cylindracea, elongatâ basi, strumosâ; a polycarpo et strumifero capsulâ lævi.

1300. lin. 29. Lege *exemplari.*

1341. Bartramia *gracilis* omnino delendum, ex observationibus amicissimi D. Turneri nuper communicatis.

END OF THE THIRD VOLUME.


INDEX GENERUM


VOL. III.

A	DIANTUM	-	Pag.		Pag.		
Ajuga	-	-	1138	Elatine	-	-	1396
Alopecurus	-	-	1399	Empetrum	-	-	1072
Amaranthus	-	-	1386	Encalypta	-	-	1180
Andräa	-	-	1018	Equisetum	-	-	1102
Aristolochia	-	-	1178	Eriocaulon	-	-	1009
Arum	-	-	947	Fagus	-	-	1027
Aspidium	-	-	1023	Fontinalis	-	-	1336
Asplenium	-	-	1118	Funaria	-	-	1338
Atriplex	-	-	1126	Grimmia	-	-	1188
Avena	-	-	1090	Gymnostomum	-	-	1158
Bartramia	-	-	1390	Hippophae	-	-	1075
Betulá	-	-	1339	Humulus	-	-	1077
Blechnum	-	-	1011	Hydrocharis	-	-	1084
Bryonia	-	-	1135	Hymenophyllum	-	-	1141
Bryum	-	-	1018	Hypnum	-	-	1276
Buxbaumia	-	-	1347	Isoetes	-	-	1144
Buxus	-	-	1148	Juncus	-	-	1395
Carex	-	-	1013	Juniperus	-	-	1085
Carpinus	-	-	963	Knappia	-	-	1387
Ceratophyllum	-	-	1029	Lemna	-	-	956
Corylus	-	-	1020	Littorella	-	-	1011
Cyathea	-	-	1030	Lonicera	-	-	1393
Cypripedium	-	-	1138	Lycopodium	-	-	1108
Dicranum	-	-	1201, 1406	Malaxis			
Draba	-	-	1400				

INDEX GENERUM VOL. III.

	Pag.			Pag.
Malaxis	-	940	Ruscus	-
Mercurialis	-	1083	Sagittaria	-
Mnium	-	1344	Salix	-
Myrica	-	1076	Satyrium	-
Myriophyllum	-	1021	Scandix	-
Neckera	-	1272	Scolopendrium	-
Ophioglossum	-	1106	Serapias	-
Ophrys	-	931	Sisymbrium	-
Orchis	-	918	Sparganium	-
Orthotrichum	-	1262	Sphagnum	-
Osmunda	-	1107	Splachnum	-
Phascum	-	1149	Stellaria	-
Pilularia	-	1143	Tamus	-
Pinus	-	1031	Taxus	-
Poa	-	1387	Tetraphis	-
Polypodium	-	1113	Tortula	-
Polytrichum	-	1372	Trichostomum	-
Populus	-	1079	Trifolium	-
Potamogeton	-	1390	Typha	-
Poterium	-	1024	Urtica	-
Pteris	-	1136	Valeriana	-
Pterogonium	-	1270	Viscum	-
Quercus	-	1025	Xanthium	-
Rhodiola	-	1082	Zannichellia	-
Rosa	-	1398		955










RQ 7.12.1979



24ColorCard CameraColor.com

