

577 50
A TABULATED ARRANGEMENT

OF THE

PRINCIPAL DISEASES

CLASSIFICATIONS OF SKIN DISEASES.

TO ILLUSTRATE A PAPER BY

MR. TILBURY FOX.

READ BEFORE THE MEDICAL SOCIETY,

ON MONDAY, MARCH 7th 1884.

LONDON: GEORGE BARNWICK, 109, FLEET STREET.

HARDY'S ARRANGEMENT.

I.—MACULÆ and DEFORMITIES.

(A.)—DEFORMITIES OF PIGMENTATION.

1. *Augmentation*.—(a) Pigmentary naevi; (b) Lentigo; (c) Ephelis; (d) Nigrities.
2. *Decoloration*.—(a) Albinism; (b) Vitiligo.

(B.)—DEFORMITIES OF VASCULAR STRUCTURE.

- (a) Port-wine marks; (b) Naevi; (c) Sanguineous tumours.

(C.)—DEFORMITIES OF SEBACEOUS FOLLICLES.

- (a) Acne miliaris; (b) Molluscum.

(D.)—DEFORMITIES OF PAPILLÆ.

- (a) Warts.

(E.)—DEFORMITIES OF EPIDERMIS.

- (a) Corns; (b) Ichthyosis.

(F.)—DEFORMITIES OF DERMA.

Keloid.

II.—LOCAL INFLAMMATIONS.

(A.)—ERYTHEMA.

1. *Cause local*.—(a) simplex; (b) vesico-pustular; (c) intertrigo.
2. *General symptoms*.—(a) papulatum; (b) nodosum; (c) scarlatiniforme; (d) mammillatum; (e) copaliquè.
3. *Symptomatic and Secondary*.—(a) læve; (b) paratrimma; (c) pernio.

(B.)—URTICARIA.

(C.)—ECTHYMA.

1. *Acute*.—(a) simplex; (b) gangrenosum.
2. *Chronic*.—(a) infantilis; (b) cachecticum.

(D.)—ZONA (H. zoster).

(E.)—STROPHULUS.

- (a) Simplex (usual varieties); (b) pruriginosus (a mixed disease).

(F.)—PRURIGO.

1. *According to intensity*.—(a) mitis; (b) formicans.
2. *According to cause*.—(a) scabiosa; (b) pedicularis; (c) strophulus; (d) nervosa.
3. *According to seat*.—(a) podicis; (b) scroti; (c) pudendi.

2. *Inflammatory*.—(a) simplex; (b) indurata; (c) rosacea; (d) hypertrophica.

(H.)—PEMPHIGUS.

1. *Acute*.—(a) neonatorum; (b) of adults.

2. *Chronic*.—(a) diutinus; (b) foliaceus; (c) pruriginosus.

III.—PARASITIC.

IV.—ERUPTIVE FEVERS.

V.—SYMPTOMATIC ERUPTIONS.

Ex. Sudamina, Herpes, Purpura, Typhoid spots.

VI.—DARTRES.

(A.)—ECZEMA.

1. *According to aspect*.—(a) simplex; (b) rubra; (c) rimosum; (d) impetigo.

2. *According to configuration*.—(a) figuratum; (b) nummularis; (c) diffusum; (d) I. sparsa.

3. *According to seat*.—(a) pilaris; (b) capitis; (c) genitalis; (d) palmaris, &c.

(B.)—LICHEN.

(a) simplex; (b) circumscriptus; (c) agrius; (d) inveterata; (e) urticatus; (f) gyratus; (g) lividus;
(h) tropicus.

(C.)—PSORIASIS.

1. *Form*.—(a) guttata; (b) circinata; (c) gyrata; (d) diffusa.

2. *Seat*.—(a) communis; (b) capitis; (c) faciei; (d) palmaris; (e) palpebralis; (f) præputialis;
(g) genitalis, &c.

(D.)—PITYRIASIS.

(a) alba; (b) rubra; (c) nigra; (d) pilaris.

VII.—SCROFULOUS.

(a) Erythematous (Lupus); (b) Pustular (Impetigo rodens); (c) Verrucous; (d) Tubercular (with or without ulceration); (e) Phlegmonous; (f) Corneous.

VIII.—SYPHILITIC.

(a) Pigmentary; (b) Vesicular; (c) Papular; (d) Squamous; (e) Exanthems; (f) Pustular; (g) Bullous;
(h) Vegetating; (i) Tubercular.

IX.—CANCERS.

X.—EXOTICS (not seen in France).

Ex. Yaws, Elephantiasis.



Digitized by the Internet Archive
in 2019 with funding from
Wellcome Library

<https://archive.org/details/b30798760>

B. SPECIAL STRUCTURES AFFECTED.

E. Glands...

- Sweat.....
 - Excess of activity.....
 - Diminution of ditto.....
 - Alteration in character.....
 - Excessive secretion.....
 - Retention with or without diminished secretion.....
 - Alteration of secretion.....
 - Inflammation, generally with more or less retention.....
- Sebacaceous....
 - Duct open.....
 - Hypertrophy of Gland Sac.....
 - Duct shut.....
 - Sebaceous, Calcareous, and Serous Cysts.
 - Steatoma.
 - Stearrhœa havescens, nigricans.
 - Hordeolum.
 - Acne.
 - Orychia.
 - Psoriasis, &c.
 - Canties, Felting.
 - Calvities, Abnormal Direction.
 - Idrosis.
 - Anidrosis.
 - Chromidrosis, Osmidrosis, Hæmidrosis, &c.
 - Stearrhœa or Seborrhœa.
 - Ichthyosis spuria vel sebacea.
 - Comedones, Acne punctata, Xeroderma.
 - Cornua, Molluscum.
 - Lichen pilaris.

F. Nails

G. Hairs

- Scabies.....
- Grubs.....
- Prurigo.....
- Pruritus and Erythema.....
- Impetigo, &c.....
- A carus Scabiei et autumnalis.
- Steatozoon folliculorum.
- Petionulus corporis.
- Leptus.
- Pulex irritans (Flea).
- Cimex lectularius (bug).
- Filaria Medicinensis.
- Pediculus pubis.
- capitis.

3. Parasitic.

Diseases due to the presence of Animal Parasites, or Dermatozoa.....

Diseases due to the presence of Vegetable Parasites or Dermatophyta (Epiphytic Diseases).....

- Of body generally.....
 - Scabies.....
 - Grubs.....
 - Prurigo.....
 - Pruritus and Erythema.....
- Of Pubis.....
 - Impetigo, &c.....
- Of Scalp.....
 - Favus.....
 - Tinea tonsurans.....
 - decalvans.....
 - Polonica.....
 - Case of Dr. Martin.....
 - Herpes circinatus.....
 - Sycosis.....
 - Tinea tarsi.....
 - Chloasma.....
 - Madura Foot.....
- Of Scalp.....
 - Tinea tonsurans.....
 - decalvans.....
 - Polonica.....
 - Case of Dr. Martin.....
 - Herpes circinatus.....
 - Sycosis.....
 - Tinea tarsi.....
 - Chloasma.....
 - Madura Foot.....
- Of Body.....
 - Herpes circinatus.....
 - Sycosis.....
 - Tinea tarsi.....
 - Chloasma.....
 - Madura Foot.....
- Of Chin.....
 - Sycosis.....
 - Tinea tarsi.....
 - Chloasma.....
 - Madura Foot.....
- Eyelid.....
 - Tinea tarsi.....
 - Chloasma.....
 - Madura Foot.....
- Chest.....
 - Tinea tarsi.....
 - Chloasma.....
 - Madura Foot.....
- Foot.....
 - Madura Foot.....

Due to the presence of

Due to the presence of

- Achorion Schönleini.
- Trichophyton tonsurans.
- Microsporion Andouini.
- Trichophyton sporuloides.
- Zoogloea capillorum.
- T. tonsurans.
- Microsporion mentagrophytes.
- T. tonsurans.
- M. Furtur.
- Chionryphe Carteri.

Syphilis may modify most of the forms of skin disease, and is looked upon as a modifying agency, and not as deserving the formation of a special class. Syphilitic eruptions may be Pigmentary, Erythematous, Vesicular, Pustular, Papular, Scaly, Tubercular, and Vegetating.

THE MIXED MODE OF CLASSIFYING SKIN DISEASES.

BY W. TILBURY FOX.

1. Epidermal

A. Specific Eruptions of acute specific diseases

- { Pityriasis.
- { Ichthyosis vera.
- { Xeroderma.
- { Typhus, Typhoid.
- { Rubella, Scarlatina.
- { Variola, Vaccinia.
- { Erysipelas, Varicella.

Chiefly from some blood alteration of stomach origin.....

Direct poisoning

Local irritation

- { Erythema, Roseola,
- { Rosalia, Urticaria.
- { Insect Bites and Stings.
- { Pernio.
- { Burns, Scalds.
- { Eczema, Pemphigus.
- { Herpes, Miliaria.

B. Non-specific

Vesicular

Papular

Pustular

Sub order Furunculi

Squamous

- { Lichen, Strophulus, Prurigo.
- { Rupia, Impetigo, Ecthyma.
- { Furunculus, Pustula maligna, Anthrax.
- { Lepra and Psoriasis.
- { Elephantiasis.
- { Framboesia.
- { Keloid.

C. Degenerations

Homologous products

Heterologous

Strumous

Malignant

- { Lupus.
- { Epithelioma.
- { Cancer.
- { Canceroid.
- { Rodent Ulcer.

A. Diseases of the Papillæ.....

B. " Vascular Structures.....

C. " Nervous Supply.....

- { Warts, Corns, Callosities, Naevi.
- { Naevi (and Purpura?).
- { Pruritus, Hyperæsthesia, Anæsthesia.
- { Vitiligo.
- { Albinismus.*
- { Melasma, Naevi.

- { Leucopathia,* Melanopathia,* Chlorosis.
- { Itfersus Brown'd Skin * Syphilis *

Congenital

Diminution of activity ..

Increase of ditto

D. Of Chromatic function ..

Secondary to blood disease

A. STRUCTURES GENERALLY INVOLVED.

2. Dermal.

USUALLY

