



Cl. 5.12.02 Bk. 19639 EP
Vol. 1

THE ETHEL CARR PEACOCK
MEMORIAL COLLECTION


Matris amori monumentum



TRINITY COLLEGE LIBRARY
DURHAM, N. C.
1903

Gift of Dr. and Mrs. Dred Peacock

THI
R
No



Digitized by the Internet Archive
in 2016 with funding from
Duke University Libraries

ELEMENTARY AND PRACTICAL

ALGEBRA:

IN WHICH HAVE BEEN ATTEMPTED

IMPROVEMENTS

IN GENERAL ARRANGEMENT AND EXPOSITION:

AND IN THE MEANS OF THOROUGH DISCIPLINE IN THE PRINCIPLES AND
APPLICATIONS OF THE SCIENCE.

BY JAMES B. DODD, A. M.,

MORRISON PROFESSOR OF MATHEMATICS AND NATURAL PHILOSOPHY IN TRANSYLVANIA
UNIVERSITY.

FOURTEENTH EDITION.

NEW YORK:
PRATT, OAKLEY & COMPANY,
21 MURRAY STREET.

1860.

23457

Entered, according to Act of Congress, in the year 1852,
By JAMES B. DODD,
In the Clerk's Office of the District Court of Kentucky.

J. P. JONES & CO., STEREOTYPERS.
183 WILLIAM STREET

512.02

D639F2

P

copy 1

PREFACE.

THE following work is designed to furnish a practicable course of Algebra for the younger classes of students, without the omission of any thing important to a thorough education in the subjects which it embraces.

It aims at the most methodical *arrangement*, the clearest *expositions*, the best elementary *exercises*, and the most varied and useful *applications*:—in all these respects presenting some *new features*, which have been adopted as improvements in the method of teaching this science.

All that is appropriate to an Algebraic treatise in a general course of mathematical studies, or necessary in preparation for the higher works of the course, has been introduced, with the exception of a few subjects which are more exclusively preliminary to the Differential and Integral Calculus. These, with whatever else may be considered useful in a larger work, will shortly be added to the present treatise.

Between this work and the author's Arithmetic there will be found a mutual correspondence in many respects, though each is entirely complete in itself. The two are commended to the consideration of Teachers of Mathematics, and the Guardians and Friends of Education, as containing a practicable, progressive, and thorough course of study, for Schools, on these connected and important branches of science.

TRANSYLVANIA UNIVERSITY, }

July 20th, 1852.

23457

R E M A R K S

ON THE METHOD OF USING THIS WORK, AND CONDUCTING EXAMINATIONS IN ALGEBRA.

The following remarks may be useful to the less experienced Teacher using this work, who would make it fully efficient for the purposes intended.

1. The definitions and propositions numbered (1), (2), (3), &c., and the Rules I, II, III, &c., should be accurately memorized and recited by the Student.

2. The accompanying examples, illustrations, or demonstrations, should be required of the Student, and discussed with him on the part of the Teacher, with reference to the *principles* involved in them.


3. The *oral* exercises in the earlier parts of the work, should be exacted; and the Student should often be examined on the exercises under the Rules, with his book closed.

4. In the solution of Equations and Problems, he should explain each part of the operation, as exemplified in different parts of the work.

5. The Analysis of Contents (see the next page) will be convenient for reviews on the theory of the science; and such reviews should be frequent. The Student will thus become familiar with the phraseology, principles, and order of the science.

6. The Student's acquisitions will depend very much on the *exactness*, as well as on the frequency, with which he is examined. The requisitions made on him should be adapted to his capabilities,—which, it should be remembered, are liable to be sometimes *overrated*, and sometimes *underrated*, by Authors and Teachers.

ANALYSIS OF CONTENTS.

 This Analysis is designed to be used in oral examinations, in reviews. The Teacher will name the *topic* as presented in this table; the Learner will respond according to his knowledge of the subject.

For example: the Teacher will say, "Science and Art;" the Learner will respond, "Science is knowledge reduced to a system; Art is knowledge applied to practical purposes."

CHAPTER I.

PRELIMINARY DEFINITIONS AND EXERCISES.—PAGE 1 . . . 8.

SCIENCE and Art, (1).—A Unit—Numbers, (2).—Quantity—Whether Numbers are *quantities*, (3).—MATHEMATICS—Its most general Divisions—Arithmetic—Geometry, (4).—ALGEBRA, (5).—Symbols of Quantities, (6).—Symbols of Operations, the Sign + plus, (7).—The Sign — *minus*, (8).—The Sign \times *into*—A *point* (.) between Quantities—Quantities in *juxtaposition*, (9).—The sign \div *by*—Division otherwise denoted—An Integral Quantity, (10).—Use of the Parenthesis or Vinculum, (11).—Factors and Constant Product, (12).—Powers of Quantities, (13).—Roots of Quantities, (14).—The Coefficient of a Quantity, (15).—The Exponent of a Quantity, (16).—What an Integral Coefficient indicates—an Integral Exponent, (17).—Similar and Dissimilar Quantities, (18).—An Algebraic Monomial, (19).—An Algebraic Polynomial—A Binomial—A Trinomial, (20).—Whether the Value of a Polynomial is affected by changing the Order of its Terms, (21).—A Polynomial arranged by *powers*, (22).—A Homogeneous Polynomial—Dimensions and Degrees of the Terms of a Polynomial, (23).—Positive and Negative Quantities, (24).—Effect of a Negative Quantity in an Expression or a Calculation—Occasional Use of the Positive and Negative Signs, (25)

CHAPTER II.

ADDITION.—SUBTRACTION.—MULTIPLICATION.—DIVISION.—9 . . . 26.

Algebraic ADDITION, (26).—How to Add Similar Terms with *like signs*, (27).—Sum of Two Equal Similar Terms with *contrary signs*, (28).—How to Add unequal Similar Terms with *contrary signs*, (29) —Sum of Two or more Dissimilar Terms, (30).

RULE I. *For the Addition of Algebraic Quantities*, (31).

Calculations on the same Polynomial in two or more ()'s, (32).

Algebraic SUBTRACTION, (33).—How to Subtract a Monomial from another Quantity, (34).—How to Subtract a Monomial from a Dissimilar Quantity, (35).

RULE II. *For the Subtraction of Algebraic Quantities*, (36).

How to denote the Subtraction of a *negative* Monomial—of a Polynomial, (37).—Change of *signs* in a Polynomial without affecting its Value, (38).

Algebraic MULTIPLICATION—When the Multiplier is *positive*—When the Multiplier is *negative*, (39).—Product of Two Monomials, (40).—Exponent, in the Product, of a Letter occurring in both the Monomials multiplied together, (41).—*Sign of the Product*—*Reason* for this when both the Quantities are *negative*—When One of them is *positive* and the Other *negative* (42).—Product when Either of the Two Factors is 0, (43).

RULE III. *To Multiply a Monomial into a Polynomial*, (44).

RULE IV. *To Multiply a Polynomial into a Polynomial*, (45).

Algebraic DIVISION, (46).—How to find the Quotient of Two Monomials, (47).—Value of any Quantity with *exponent* 0, (48).—*Sign of the Quotient*, and Principle which determines it, (49)—Quotient of 0 divided by any Quantity, and of any Quantity divided by 0, (50).

RULE V. *To Divide a Monomial into a Polynomial*, (51).

RULE VI. *To Divide a Polynomial into a Polynomial*, (52).

How the *indicated Product* of Two or more Factors may be divided, (53).

CHAPTER III.

COMPOSITE QUANTITIES.—COMMON MEASURE.—COMMON MULTIPLE.—
27 . . . 40.

A Composite and a Prime Quantity, (54).—Decomposition of a Quantity --Divisor and Quotient as Factors of a Quantity, (55).—What the Difference of Two Quantities will divide, (56).—What the sum of two Quantities will divide, (57).—Product of the Sum and Difference of two Quantities, (58).—Square of the Sum of two Quantities, (59).—Square of the Difference of two Quantities, (60).—Case in which a Trinomial may be resolved into Two unequal Binomial Factors, (61).—One Quantity a MEASURE of another —A Common Measure of Two or more Quantities, (62).—Greatest Common Measure of Two or More Quantities, (63).—Composition of the Greatest Common Measure, (64).—Principle on which depends the Rule for the Greatest Common Measure, (65).

RULE VII.—*To Find the Greatest Common Measure of Two Quantities*, (66).

Whether the *Signs* in a Common Measure may be changed, (67).—Of a Factor which is common to all the Terms of the Dividend, (68).—Of a Factor which is contained in all the Terms of the two Polynomials, (69).—One Quantity a MULTIPLE of another.—A Common Multiple of Two or more Quantities, (70).—Least Common Multiple of Two or More Quantities, (71).—Composition of the Least Common Multiple, (72).—Relation of Least Common Multiple to Product and Greatest Common Measure, (73).

RULE VIII. *To Find the Least Common Multiple of Two or More Quantities*, (74).

CHAPTER IV.

FRACTIONS.—41 . . . 66.

An Algebraic FRACTION, (75).—Of a Quantity with a *negative* Exponent, (76).—Two Methods of representing the Quotient when the Divisor is not a Factor of the Dividend, (77).—How any Factor may be transferred from the

Numerator to the Denominator, and *vice versa*, (78).—*Reciprocal* of a Quantity, (79).—*Reciprocal* of a Fraction, (80).—Constant Value of a Fraction, (81).—When a Fraction is *positive*, and when *negative*, (82).—Signification of $+$ or $-$ prefixed to a Fraction, (83).—Changes of Signs without affecting the Value of a Fraction—How a Polynomial may be changed from Positive to Negative, (84).—A Fraction reduced to *lower Terms*—How a *binomial* Common Measure may often be discovered, (85).

RULE IX. *To Reduce a Fraction to its Lowest Terms*, (86).

When Two or more Fractions are said to have a Common Denominator.—How Fractions may be reduced, mentally, to a Common Denominator, (87).

RULE X. *To Reduce Two or More Fractions to a Common Denominator*, (88).

An Integral Quantity, (89).—A Mixed Quantity, (90)—An Improper Fraction, (91).

RULE XI. *To Reduce an Integral or a Mixed Quantity to an Improper Fraction*, (92).

RULE XII. *To Reduce an Improper Fraction to an Integral or a Mixed Quantity*, (93).

By what means the Sum of Two or More Fractions is found, (94).

RULE XIII. *For the Addition of Fractions*, (95).

By what means the Difference of Two Fractions is found, (96).

RULE XIV. *For the Subtraction of Fractions*, (97).

Product of Two or more Fractions, (98).—Effect of Multiplying by a Fraction, (99).—Compound Fractions, (100).—Equivalent of Multiplying Two or more Fractions together, (101).

RULE XV. *For the Multiplication of Fractions*, (102).

How a Fraction is Multiplied by its own Denominator, and what Cancellations may be made in the Multiplication of Fractions, (103).—Quotient of Two Fractions, (104)—Complex or Mixed Fractions, (105).

RULE XVI. *For the Division of Fractions*, (106).

CHAPTER V.

SIMPLE EQUATIONS.—67 . . . 90.

AN EQUATION—The First Member—The Second Member, (107).—For what purposes Equations are employed—How applied to the Solution of Questions, (108).—The Solution of an Equation—Verification of the Value found for the Unknown Quantity, (109).—A *Simple* Equation—A Quadratic Equation—A Cubic Equation, (110).—A Numerical Equation—A Literal Equation—An Identical Equation, (111).—Transformation of an Equation, (112).—An *Axiom*—*Axiom first, second, &c.*, (113).—How the Value of the Unknown Quantity is found—Transformations necessary, (114).—How to clear an equation of Fractions—How by means of Least Common Multiple—Advantage of this Method, (115).—How any term may be Transposed from one Side of an Equation to the other, (116).—Change of the Signs in an Equation, (117).

RULE XVII. *For the Solution of a Simple Equation containing but one unknown quantity, (118.)*

A Problem, and in what its Solution consists—General Method of forming the Equation of a Problem, (119).—Solution of Problems with Two or more Unknown Quantities—Independent Equations, (120).—General Method of solving Two Equations, (121).—Elimination by Addition or Subtraction, (122).—Elimination by Substitution, (123).—Elimination by Comparison, (124).—Solution of Three Equations—Of Four Equations, (125).—Of Problems in which there are Three or more Required Quantities, (126).

CHAPTER VI.

RATIO.—PROPORTION.—VARIATION.—91 . . . 110.

Ratio of one Quantity to another, (127).—*Sign* of Ratio, (128).—How the value of a Ratio may be represented, (129).—Direct and *Inverse* Ratio, (130).—Compound Ratio, (131) —Ratio of the *first* to the *last* of any Number of Quantities, (132).—Duplicate and Triplicate Ratios, (133).—Equimultiples and Equisubmultiples, (134).—Ratio of Equimultiples and Equisubmultiples, (135).—PROPORTION, (136).—Four Quantities in Proportion, (137).—Three quantities in Proportion, (138).—Direct and *Inverse* Propor-

tion, (139).—*Sign* of Proportion, (140).—Inverse Converted into Direct Proportion, (141).—VARIATION—Variation *direct*—Variation *inverse*, (142).—Product of Two Quantities varying *inversely* with each other, (143).—Variation, an Abbreviated Proportion, (144).—A *Theorem*—A Corollary, (145).—Ratio of two Fractions having a *common* Term, (146).—How the value of a Fraction *varies*, (147).—Corresponding Equalities and Inequalities between the Antecedents and Consequents of a Proportion, (148).—A Proportion converted into an Equation, (149).—A Fourth Proportional, how found, (150).—Product of the Extremes when Three Quantities are in Proportion, (151).—Mean Proportional, how found, (152).—An Equation converted into a Proportion, (153).—Ratio of the *first* to the *third* of Three Proportional Quantities, (154).—Proportion by Inversion, (155).—Proportion by Alternation, (156). What Multiplications may be made in a Proportion, (157).—What Divisions may be made in a Proportion, (158).—Proportion by Composition, (159).—Proportion by Division, (160).—Proportion between the Sum of two or more Antecedents and that of their Consequents, (161).—A Proportion derived from Two other Proportions in which there are *common* Terms, (162).—Proportion between the Sums and Differences of the Antecedents and Consequents, (163).—Products of the Corresponding Terms of Two or more Proportions, (164).—Proportion between the Powers or Roots of Proportional Quantities, (165).—Substitution of Factors in a Proportion, (166).—*General Solution* of a Problem, (167).—Two Numbers found from their Sum and Difference, (168).—An Algebraic *Formula*, (169).—Of a Proportion occurring in the Solution of a Problem, (170).—*Percentage*—Ratio of Percentage—Basis of Percentage, (171).—Amount of Percentage, how found, (172).—*Interest*—The Principal—The Amount, (173).—Amount of Interest, how found, (174).

CHAPTER VII.

ARITHMETICAL, HARMONICAL, AND GEOMETRICAL PROGRESSION.—111... 120.

An Arithmetical PROGRESSION, (175).—Last Term of an Arithmetical Progression, equal to what, (176).—Common Difference of the Terms, (177).—Sum of the two Extremes, (178).—Arithmetical Mean, (179).—Sum of all the Terms, (180).—*Formulas* in Arithmetical Progression, (181).—An Harmonical PROGRESSION, (182).—An Harmonical converted into an Arith-

metrical Progression, (183).—Harmonical Mean, (184).—A Geometrical PROGRESSION, (185).—Last Term of a Geometrical Progression, (186).—Power of the *ratio* found from the First and last Terms, (187).—Product of the two Extremes, (188).—Geometrical Mean, (189).—Sum of all the Terms, (190).—Sum of an *infinite* number of Decreasing Terms, (191).—*Formulas* in Geometrical Progression, (192).

CHAPTER VIII.

PERMUTATIONS AND COMBINATIONS.—INVOLUTION.—BINOMIAL.—THEOREM.—EVOLUTION.—121. . . . 146.

PERMUTATIONS, (193).—Number of Permutations, how found, (194).—COMBINATIONS, (195).—Number of Combinations, how found, (196).—INVOLUTION, (197).—A Higher Power found from lower Powers of the same Quantity, (198).—Powers of *unity*, (199).—Powers of Monomials, how found, (200).—Powers of Fractions, how found—Powers of a *mixed* Quantity, (201).—*Sign* to be Prefixed to a Power, (202).—Powers of Binomials, or of any Polynomials, (203).—BINOMIAL THEOREM—*Exponents* in any Power of $(a \pm b)$ —*Coefficients*—*Signs*, (204).—Formula for the development of $(a + b^n)$ —At what Term the development will *terminate*, (205).—EVOLUTION—Extracting the Square Root, in what it consists—The Cube Root, (206).—Roots of *unity*, (207).—Roots of Monomials, how found, (208).—Roots of Fractions, how found—Roots of a *mixed* Quantity, (209).—Of a Root whose Exponent is resolvable into two Factors, (210).—What denoted by the Numerator and Denominator of a Fractional Exponent, (211).—Root of a power of a Quantity, (212).—Equivalent Exponents, (213).—*Sign* to be Prefixed to an *odd* Root of a Quantity, (214).—*Sign* to be Prefixed to an *even* Root, (215).—Of an *even* Root of a *negative* Quantity, (216).—How the Roots of Polynomials may be discovered, (217).

RULE XVIII. *To Extract the Square Root of a Polynomial*, (218.)

Principle for determining the Number of Figures in the Square Root of a Number, (219).—Square of any two Parts into which a number may be divided, (220).—Periods to be formed in Extracting the Square Root of a Decimal Fraction—Why the last Period must be *complete*—Number of Decimal Figures to be made in the Root, (221).

RULE XIX. *To Extract the Cube Root of a Polynomial, (222.)*

Principle for determining the Number of Figures in the Cube Root of a Number, (223).—Cube of any two Parts into which a Number may be divided, (224).

RULE XX. *To Extract the Cube Root of a Number, (225).*

Periods to be formed in extracting the Cube Root of a Decimal Fraction.—Why the last period must be *Complete*—Number of decimal Figures to be made in the Root, (226).—How any Root whatever of a Polynomial might be extracted, (227).

RULE XXI. *To Extract any Root of a Polynomial, (228).*

CHAPTER IX.

IRRATIONAL OR SURD QUANTITIES.—IMAGINARY QUANTITIES.—147...166.

Perfect and Imperfect Powers, (229).—A Rational Quantity—An Irrational or *Surd* Quantity—Radical Quantities, (230).—Radical Sign—How this Sign may always be superseded, (231).—Similar and Dissimilar Surds, (232). How a Rational Quantity may be expressed under the Form of a Surd, (233). Transfer of an Exponent between Factors and Product, (234).—To what the Exponent of a Quantity may be changed, (235).—Product of a Rational and an Irrational Factor, (236).—How a Surd may be *simplified*, (237).—How a Fractional may be reduced to an *Integral* Surd, (238).—Surds of Different Roots reduced to the Same Root, (239).—How to find the *Sum* or Difference of Similar Surds—of *Dissimilar* Surds, (240).—How to find the *Product* or Quotient of Surds of the *same* root—of *different* Roots—of any two Roots of the *Same* Quantity, (241).—Expediency of *rationalizing* a Surd Divisor or Denominator, (242).—How a Monomial Surd may be made to produce a Rational Quantity—a *binomial* Surd—a *trinomial* Surd, (243).—Of the Powers and Roots of Irrational Quantities, (244).—Of the Square Root of a Numerical Binomial of the form $a \pm \sqrt{b}$ (245).—Of *Imaginary* Quantities—From what an Imaginary Quantity results, (246).—Of the Calculus of Imaginary Quantities—By what means the Sign affecting the Product of two Imaginaries may be determined, (247).—Resolution of an Imaginary Quantity, (248).—Product of two Imaginary Square Roots, (249)

CHAPTER X.

QUADRATIC AND OTHER EQUATIONS.—167 . . . 204.

A Quadratic Equation—A Cubic Equation—A Biquadratic Equation, (250).—A Pure Equation—An Affected and a Complete Equation, (251).—A Root of an Equation, (252).—Principle for determining a Divisor of an Equation, (253).—Principle for determining a Root of an Equation, (254). Number of Roots of an Equation—Whether the several Roots of an Equation are necessarily *unequal*, (255).

RULE XXII. *For the Solution of a Pure Equation*, (256).

How a *Surd* Quantity in an Equation may be *rationalized*, (257).—How two *Surds* in an Equation may be rationalized, (258).—Of an Equation containing a Fraction whose terms are both *irrational*, (259).

RULE XXIII. *For the Solution of a Complete Equation of the Second Degree*, (260).

How any Binomial of the form ax^2+bx may be made a *Perfect Square* (261).

RULE XXIV. *To Reduce an Equation of the form ax^2+bx to a Simple Equation*, (262).

Of Equations having a Quadratic Form with reference to a Power or *Root* of the Unknown Quantity, (263).—Of the Solution of two Equations, one or both of the Second or a Higher Degree, Containing two Unknown Quantities, (264).—Solutions by means of an Auxiliary Unknown Quantity, (265).—Solutions by means of two Auxiliary Unknown Quantities, (266).—General Law of the Coefficients of Equations, (267).—Determination of the Integral Roots of Equations, (268).—Solution of Equations by *Approximation*, (269).—General Method of Elimination, (270).

CHAPTER XI.

GENERAL DESCRIPTION OF PROBLEMS.—MISCELLANEOUS PROBLEMS.—205 . . . 217.

A Determinate Problem—How a Determinate Problem is represented, (270).—An Indeterminate Problem—How represented, (271).—An Impossible Problem—How represented, (272).—Greatest Product of any two Parts into which any Quantity may be divided, (273).—Signification of the Different Forms under which the value of the Unknown Quantity may be found in an Equation, (274).—*Inequalities*, (275).—For what purpose Inequalities are employed, (276).—Inequalities in the *same*, and in a *contrary sense*, (277).—Inequalities between *negative* quantities, (278).—Transformation of Inequalities, (279).



A L G E B R A .

CHAPTER I.

PRELIMINARY DEFINITIONS AND EXERCISES

Science and Art.

(1.) SCIENCE is knowledge reduced to a *system*.—ART is knowledge applied to practical purposes.

The Rules of Art are founded on the Principles of Science.

Numbers.—Quantity.

(2.) A UNIT is any thing regarded simply as *one*; and *numbers* are repetitions of a *unit*.

Thus the numbers *two, three, &c.*, are repetitions of the unit *one*.

(3.) QUANTITY is any thing which admits of being measured.

Thus a *line* is a quantity, and we express its measure in saying it is so many *feet* or *inches* long. *Time, weight, and distance* are also quantities.

Numbers are quantities; for every number expresses the measure of itself in *units*; and numbers are used to express the measures of all other quantities. Thus we express the measure of Time by a number of *days, hours, &c*

MATHEMATICS.

(4.) MATHEMATICS is the science of *quantity*. Its most general divisions are Arithmetic and Geometry.

ARITHMETIC is the science of *numbers*; or, when practically applied, the art of Calculation.

GEOMETRY is the science which treats of Extension—in *length*, *breadth*, and *height*, depth, or thickness.

(5.) ALGEBRA is a method of investigating the relations of both Arithmetical and Geometrical quantities, by *signs* or *symbols*.

Symbols of Quantities.

(6.) Quantities are represented in Algebra by *letters*,—*known* quantities usually by the *first*, and *unknown* or required quantities by the *last* letters of the Alphabet.

Thus the quantity *a* or *b*, that is, the quantity *represented* by *a* or *b*, will generally be understood as *known in value*; while the quantity *x* or *y* will be *unknown* or required.

Quantities represented by *letters* are called *literal* quantities, in contradistinction to *numbers* or numerical quantities.

Symbols of Operations.

(7.) The sign $+$ *plus* prefixed to a quantity, denotes that the quantity is to be *added*, or taken *additively*.

Thus $a+b$, *a plus b*, denotes that the quantity *b* is to be added to the quantity *a*.

(8.) The sign $-$ *minus* prefixed to a quantity, denotes that the quantity is to be *subtracted*, or taken *subtractively*.

Thus $a-b$, *a minus b*, denotes that the quantity *b* is to be subtracted from the quantity *a*.

(9.) The sign \times *into* between two quantities, denotes that the two quantities are to be *multiplied* together.

Thus $a \times b$, *a into b*, denotes that the two quantities *a* and *b* are to be multiplied together.

A *point* (.) between two literal quantities, or a numerical and a literal quantity, also denotes that the two quantities are to be multiplied together.

$a.b$ denotes *a into b*, and $3.a$ denotes *3 into a*, or *3 times a*.

The Product of *several numbers* may be denoted by points between them; thus $1.2.3$ denotes *1 into 2 into 3*, the same as $1 \times 2 \times 3$.

Quantities in juxtaposition, without any sign between them, are to be multiplied together. Thus ab denotes a and b multiplied together; and axy denotes a , x , and y multiplied together.

(10.) The sign \div by between two quantities, denotes that the quantity before the sign is to be *divided* by the one after it.

Thus $a \div b$, a by b , denotes that a is to be divided by b .

Division is also denoted by placing the *dividend* over the *divisor*, with a line between them, after the manner of a Fraction;

$\frac{a}{b}$ denotes a divided by b , the same as $a \div b$.

An *integral quantity*, in Algebra, is one which does not express any operation in *division*, whatever may be the *numerical values* which the letters represent.

(11.) A *parenthesis* () enclosing an algebraic expression, or a *vinculum* $\overline{\hspace{1cm}}$ drawn over it, connects the *value of that expression* with the sign which immediately precedes or follows it.

Thus $(a+b)c$, or $\overline{(a+b)}c$, a plus b in a parenthesis *into* c , denotes that the *sum* of a and b is to be multiplied into c .

The same thing would be denoted by $\overline{a+b} \times c$, a plus b under a vinculum *into* c .—In $a+bc$, only b would be multiplied into c .

The *vinculum*, and the expression affected by it, are sometimes set *vertically*.

Thus $\begin{array}{l} a \\ +b \\ -c \end{array} \Bigg| x$ is equivalent to $(a+b-c)x$
or $\overline{a+b-c} \times x$.

PRELIMINARY EXERCISES.

In the elementary *oral* Exercises which are occasionally inserted, the Student should write down the *quantities* as they are read to him.

☞ Suppose the letters a, b, c to represent the numbers 3, 4 5, respectively; then what is the *numerical* value of $a+b-c$?

What is the value of $ab+c$? Of $abc-bc+a$?

What is the value of $ac \div b$? Of $ab + b - ac$?

What is the value of $bc \div a$? Of $abc+ac-bc$?

What is the value of $(a+b)c$, a plus b in a *parenthesis* into c ?

What is the value of $(ab+c)a$? Of $(a+b+c)c$?

What is the value of $(a+b) \div c$? Of $(ab+b-c)a$?

What is the value of $(bc-a) \div b$? Of $(bc-a+b)b$?

The Teacher may propose other Exercises of the same nature, should he deem it necessary.

Factors.—Constant Product.

(12.) Two or more quantities multiplied together, are called the *factors* of their product; and the Product is the *same in value*, in whatever order its factors are taken.

Thus a and x are the *factors* of the product ax ; and this product is the same in value as xa . So a , b , and c are the factors of the product abc , or acb , or bca , &c.

It is most convenient to set *literal* factors according to the order of the same letters in the Alphabet; thus ax ; abc .

To understand why the product ax is equal to xa , consider a and x as representing *numbers*, and that the product of two numbers is the same, when either of them is made the *multiplier*.

For example, 25 times 7 is equal to 7 times 25. For 25 times 7 must be 7 times as many as 25 times 1, which is 25; that is, 25 times 7 is equal to 7 times 25.

☞ Prove that 14 times 9 is equal to 9 times 14.

Prove that 31 times 11 is equal to 11 times 31.

Prove that 23 times 15 is equal to 15 times 23.

Prove that 47 times 18 is equal to 18 times 47.

Powers and Roots.

(13.) The *first power* of a quantity is the *quantity itself*; thus the first power of 5 is 5, and the first power of a is a .

The *second power*, or *square*, of a quantity, is the product of the quantity *multiplied into itself*. Thus the second power, or *square*, of 5 is 5×5 , which is 25; and the second power of a is aa .

The *third power*, or *cube*, of a quantity is the product of the quantity *multiplied into its second power*, or square. Thus the third power, or cube of 5 is $5 \times 5 \times 5$, which is 125; and the third power of a is aaa .

☞ What is meant by the *fourth power* of a quantity? What is meant by the *fifth power* of a quantity? By the *seventh power* of a quantity?

What is the *square* of 3? The *cube* of 4? The fourth power of 2? The *square* of 7? The *cube* of 6? The fourth power of 10?

(14.) The *second root*, or square root, of a quantity, is that quantity whose *square is equal to the given quantity*. Thus the square root of 9 is 3; and the square root of aa is a .

The *third root*, or cube root, of a quantity, is that quantity whose *third power, or cube, is equal to the given quantity*. Thus the cube root of 8 is 2; and the cube root of aaa is a .

☞ What is meant by the *fourth root* of a quantity? What is meant by the *fifth root* of a quantity? By the *ninth root* of a quantity?

What is the square root of 16? The cube root of 27? The fourth root of 16? The square root of 81? The cube root of 1000?

What is the *square* of the square root of 4? The cube of the cube root of 125? The square of the cube root of 64? The cube of the square root of 16?

Coefficients and Exponents.

(15.) The *coefficient* of a quantity is any *multiplier* prefixed to that quantity.—In a more general sense, the coefficient of a quantity is any *factor* forming a product with that quantity.

Thus, in $3a$, 3 is the coefficient of a , and denotes 3 times a . In $5ax$, 5 is the coefficient, denoting 5 times ax . In $\frac{1}{2}x$, $\frac{1}{2}$ is the coefficient of x , and denotes *one-half* of x .

When *no numerical* coefficient is prefixed, a *unit* is always to be understood. Thus a is $1a$, once a , and ax is $1ax$, once ax .

(16.) The *exponent* of a quantity is an *integer* annexed to it, to denote a *power*, or a fraction annexed to denote a *root*, of that quantity.

Thus a^2 , a with exponent 2, denotes the second *power*, or square of a ; x^3 denotes the third *power*, or cube, of x ; and so on.

The *fractional exponents* $\frac{1}{2}$, $\frac{1}{3}$, $\frac{1}{4}$, and so on, denote, respectively the *square root*, cube root, &c., of the quantity to which they are annexed.

$a^{\frac{1}{2}}$, a with exponent $\frac{1}{2}$, denotes the square root of a ; $x^{\frac{1}{3}}$ denotes the cube root of x ; and so on.

When *no exponent* is annexed to a quantity, a *unit* is always to be understood. Thus a is a^1 , the first power of a . (13.)

An exponent is assigned to the product of two or more factors, by affecting such product with a parenthesis, or a vinculum, and the exponent.

Thus $(ax)^2$ or $\overline{ax^2}$, ax in a parenthesis, or under a vinculum, with exponent 2, denotes the square of the product ax ; whereas ax^2 denotes a into the square of x .

(17.) An *integral coefficient* indicates the repeated addition of a quantity to itself; while an *integral exponent* indicates the repeated multiplication of a quantity into itself.

Thus $3a$, 3 times a , is equivalent to $a+a+a$; while a^3 , the third power of a , is equivalent to aaa .

Coefficients and exponents are thus employed to abbreviate the language of Algebra.

☞ What is the *equivalent*, in Addition, of $2x$? Of $3ax$? Of $2abc$? Of $5a^2$? Of $4ay^2$? Of $3axy$?

What is the *equivalent*, in Multiplication, of a^2 ? Of x^3 ? Of ax^2 ? Of a^2x ? Of abx^3 ? Of ac^2x^2 ?

Allowing the value of a to be 4, what is the value of a^2 ? Of $a^{\frac{1}{2}}$? Of a^3 ? Of $5a^2$? Of $10a^{\frac{1}{2}}$? Of $a^{\frac{1}{2}}a^3$? Of $(4a)^{\frac{1}{2}}$?

Allowing a to be 4, and b 9, what is the value of $ab^{\frac{1}{2}}$? Of $a^{\frac{1}{2}}b$? Of $a^2b^{\frac{1}{2}}$? Of $(ab)^{\frac{1}{2}}$? Of $a^{\frac{1}{2}}b^{\frac{1}{2}}$? Of $\frac{1}{2}ab$? Of $\frac{1}{3}(ab)^{\frac{1}{2}}$?

Similar and Dissimilar Quantities.

(18.) *Similar* quantities are such as have all the *literal factors*, with their respective *exponents*, the same in each; otherwise, the quantities are *dissimilar*.

Thus the two quantities $2ax^2$ and $5ax^2$ are *similar*; while $3ax^2$ and $4a^2x$ are *dissimilar*, the literal factors not having the same exponents in each.

☞ Are the two quantities ab and $3ab$ similar, or *dissimilar*? Are $4a$ and $3x$ similar, or *dissimilar*? Are ay and $3ya$ similar, or *dissimilar*? $5ax^{\frac{1}{2}}$ and $7a^{\frac{1}{2}}x$? abc^2 and $5bc^2$? $2ab$ and $2ab^2$? $3axy^2$ and ay^2x ? $2bc^3$ and $3bc^{\frac{1}{2}}$?

Give an example of *three similar quantities*.—Give an example of *three dissimilar quantities*.—Another example of *three similar quantities*.—Another example of *three dissimilar quantities*.

Monomials and Polynomials.

(19.) An algebraic *monomial* is a symbol of quantity not composed of parts connected by the sign $+$ or $-$.

Thus $3a$, $5ax$, $2x^2$, and $\frac{1}{2}abc^3$ are monomials.

(20.) An algebraic *polynomial* consists of two or more monomials connected by the sign $+$ or $-$; and such monomials are called the *terms* of the polynomial.

Thus $5a^2 + bx$, and $ax^2 + 3b - 5c^2$ are polynomials.

A polynomial composed of *two terms* is called, more definitely, a *binomial*, and one composed of three terms, a *trinomial*.

(21.) The *value* of a polynomial is not affected by changing the *order of its terms*, without changing the *sign* prefixed to any term.

Thus $a + b - c$ is equivalent to $a - c + b$; for the result will evidently be the same, whether b be first added to a , and c then subtracted; or c be first subtracted, and b afterwards added.

(22.) A polynomial is arranged according to the *powers* of one of its letters, when the *exponents* of that letter increase, or *decrease*, continually in the successive terms.

Thus the polynomial $3a^3 + 4a^2x - 5ab$, is arranged according to the *descending powers* of a , since the exponents of a decrease continually in the successive terms.

The letter,—as a in this example,—according to which the polynomial is arranged, is called the *letter of arrangement*.

(23.) A polynomial is said to be *homogeneous*, when the *sum of the exponents* of the literal factors, is the same in each of its terms.

Thus $a^3 + 2ax^2 - bcy$ is homogeneous, since the sum of the exponents of the literal factors is the same, namely 3, in each term.

Each one of the literal factors composing a term, is called a *dimension* of that term; and the *degree* of any term is the ordinal of the number of its literal factors or *dimensions*.

Thus $4a^2x$ contains three literal factors, aax , and is therefore of three *dimensions*, or of the *third degree*.

☞ Arrange the polynomial $2x + 3a^2x^2 - 4ax^3 + a^3y^2$ according to the *descending powers* of a .—Arrange it according to the *ascending powers* of a .—Which of the terms of this polynomial are *homogeneous*? Tell the number of *dimensions* in each term.—Of what *degree* is each term?

Positive and Negative Quantities.

(24.) A *positive* quantity is one which enters *additively*, and a *negative* quantity is one which enters *subtractively*, into a calculation.

A positive quantity is denoted by the sign $+$, and a negative quantity by the sign $-$, prefixed to it.

In the polynomial $a+b-c$, the quantity b is *positive*, while c is *negative*.

A quantity with neither $+$ nor $-$ prefixed to it, is understood to be *positive*.

Thus in the preceding polynomial, the first term a is understood to be $+a$; for it may be regarded as $0+a$.

(25.) A *negative* quantity has an effect *contrary to that of an equal positive* quantity, in the expression, or calculation, into which it enters.

This is evident with regard to the *addition* and *subtraction* of the same quantity; the effect of *subtracting* it is contrary to that of *adding* it.

The effect of *multiplying* by a *negative* quantity is contrary to that of multiplying by an equal *positive* quantity.

For example; $3a \times 2$, $3a$ multiplied by *positive* 2, denotes that $3a$ is to be repeated *additively*, and is equivalent to $3a+3a$.

But $3a \times -2$, $3a$ multiplied by *negative* 2, denotes that $3a$ is to be repeated *subtractively*, and is equivalent to $-3a-3a$.

The same thing is true with respect to *dividing* by a negative, and an equal positive quantity.

The positive and negative signs are sometimes used to distinguish quantities as estimated in *contrary directions* from a given point or line.

Thus if Motion in one direction be denoted by the *sign* $+$, then motion in the contrary direction will be denoted by the *sign* $-$. If North Latitude be considered as *positive*, South Latitude will be *negative*.

CHAPTER II.

ADDITION.—SUBTRACTION.—MULTIPLICATION.—DIVISION.

ADDITION.

(26.) Algebraic ADDITION consists in finding the *simplest expression* for the value of two or more quantities connected together by the sign $+$ or $-$, and this equivalent expression is called the *sum of the quantities*.

The *simplest expression* for the value of $5a + 3a$, 5 times a plus 3 times a , is $8a$; then $8a$ is the *sum* of $5a$ and $3a$.

The simplest expression for the value of $5a - 3a$, 5 times a minus 3 times a , is $2a$; then $2a$ is the sum of $5a$ and $-3a$.

In the second example, observe that *adding* $-3a$ is equivalent to *subtracting* $3a$; the Adding of a *negative* quantity being the same as the Subtracting of an equal *positive* quantity.

Addition of Monomials.

(27.) *Similar* terms with *like signs*, are added together by taking the sum of their *coefficients*, annexing the common *literal factor*, and prefixing the common *sign*.

Thus $3ax + 2ax$ is $5ax$; just as 3 cents + 2 cents is 5 cents.

And $-3a - 2a$ is $-5a$; for $3a$ to be subtracted and $2a$ to be subtracted make $5a$ to be subtracted.

☞ What is the Sum of $4a$ and $3a$? Of ax and $5ax$? Of -2 and $-7x$? Of $5a^2$, $2a^2$, and $4a^2$? Of $-2b$, $-3b$, and $-5b$?

(28.) Two *equal similar* terms with contrary signs, when added together, mutually *cancel each other*; that is, their sum is 0.

Thus $3a - 3a$ is 0; that is, 0 is the simplest expression for the value of $3a - 3a$, and is therefore the *sum* of the two terms.

So $5a + x - 5a$ is equal to x ; for $5a$ and $-5a$ *cancel each other*, and x is therefore the sum of the three given terms.

☞ What is the Sum of $2ax$ and $-2ax$, that is, the simplest expression for the value of $2ax-2ax$? What is the Sum of $7b$ and $-7b$? Of a , $3x$, and $-a$? Of $3y^2$, $-3y^2$, and $5b^2$? Of ax^2 , 5 , and -5 ? Of $-abc$, $+a^2x$, and abc ?

(29.) Two *unequal similar* terms with contrary signs, are added together by taking the *difference of their coefficients*, annexing the common literal factor, and prefixing the *sign of the greater term*.

Thus $7a-3a$ is $4a$; just as $7\text{ cents} - 3\text{ cents}$ is 4 cents ; that is, the sum of $7a$ and $-3a$ is $4a$.

And $3a-7a$ is $-4a$. For $-7a$ is equal to $-3a-4a$ (27); then $3a-7a$ is equal to $3a-3a-4a$; $3a$ and $-3a$ cancel each other (28), and leave $-4a$.

☞ What is the Sum of $5ab$ and $-3ab$? Of $3a^2c$ and $-a^2c$? Of $9ax^2$ and $-4ax^2$? Of $-a^2x$ and $3a^2x$? Of $5ac$ and $-5ac$?

What is the simplest expression for the value of $2b-5b$, that is, the Sum of $2b$ and $-5b$? *How do you reason in finding that sum?*

What is the Sum of $3a^2$ and $-8a^2$? and how do you reason in finding it? What is the Sum of ax^2 and $-2ax^2$? and how do you reason in finding it? What is the Sum of $-11a^2$ and $5a^2$? and how do you reason in finding it?

(30.) The Sum of two or more *dissimilar* terms can only be indicated by arranging them in a Polynomial, so that each term shall have its *given sign prefixed to it*.

Thus the Sum of ax and bc^2 can only be indicated as $ax+bc^2$; and the Sum of ax and $-bc^2$ as $ax-bc^2$.

What is the Sum of $3a$ and $5b$? Of $5a$ and $-b$? Of $3a^2$ and $2x^2$? Of ax and by ? Of $-3b$ and $-4y$?

What is the Sum of $3b$, $5b$, and $6c$? Of a^2 , bx , and $3a^2$? Of $4c$ and $-3y$? Of $-ab$, $3xy$, and $-2c$? Of $3a^2$, $4b$, and $2c$? Of $5a$, -5 , and $5b$?

The preceding principles, and the following Rule, provide for all the cases in Algebraic Addition.

RULE I.

(31) For the Addition of Algebraic Quantities.

1. Find the *positive* and the *negative* Sum of similar terms, separately, (27), and then add together the *similar sums*, (28) and (29).

2. Connect the results thus found, and the *dissimilar* terms, in a Polynomial, prefixing to each term its proper sign. (30).

EXAMPLE.

To add together $4a^2 + 2bc - xy$, $2a^2 - 3bc + 5y$, $bc - a^2 + 3xy$, and $cy - 2a^2 - 5bc$.

$$\begin{array}{r}
 4a^2 + 2bc - xy \\
 2a^2 - 3bc + 5y \\
 -a^2 + bc + 3xy \\
 -2a^2 - 5bc + cy \\
 \hline
 3a^2 - 5bc + 2xy + cy + 5y
 \end{array}$$

The sum of the positive terms $2a^2$ and $4a^2$ is $6a^2$, and the sum of the similar *negative* terms $-2a^2$ and $-a^2$ is $-3a^2$, (27). Adding together these two similar sums, $6a^2$ and $-3a^2$, the result is $3a^2$, (29).

The sum of the positive terms bc and $2bc$ is $3bc$, and the sum of the similar *negative* terms $-5bc$ and $-3bc$ is $-8bc$. Adding together these two similar sums, $3bc$ and $-8bc$, the result is $-5bc$.

The sum of the similar terms $3xy$ and $-xy$ is $2xy$.

The results thus found, namely, $3a^2$, $-5bc$, and $2xy$, and the *dissimilar* terms cy and $5y$, are connected in a Polynomial.

EXERCISES.

1. Add together $ab + 3c^2 - 2x$, $3ab - c^2 + 5x$, $5c^2 - 2ab + y$, and $4ab - c^2 + x$.

Ans. $6ab + 6c^2 + 4x + y$.

2. Add together $4a^2 - 6b + 3$, $3a^2 + 6b - 7$, $5b + a^2 - bc$, and $a^2 - bc + 10$.

Ans. $9a^2 + 5b + 6 - 2bc$.

3. Add together $2b^2 + bc + x$, $3b^2 - 2bc + y$, $bc - 3b^2 + 5x$, and $b^2 + bc + 3y$.

Ans. $3b^2 + bc + 6x + 4y$.

4. Add together $5a - 6b^3 + 3$, $2b^3 - 4a - 7$, $7a - b^3 + c$, and $a + 3b^2 + 2c + 4$.

Ans. $9a - 5b^3 + 3c + 3b^2$.

5. Add together $a^2b - 5c + xy$, $4a^2b + 8c - xy$, $a^2b + c + 3xy$, and $a^2 + 3c - 13$.

Ans. $6a^2b + 7c + 3xy + a^2 - 13$.

6. Add together $a^3 + 2a^2 - 3bc$, $2a^3 - a^2 + bc$, $-a^3 - a^2 + b^2$, and $3a^3 + 3a^2 + 3b^2$. *Ans.* $5a^3 + 3a^2 - 2bc + 4b^2$.

7. Add together $2x + ab^2 - 3y^2$, $2ab^2 - 3x + y^2$, $-ab^2 + 5x - y^2$, and $5x + ab^2 - y$. *Ans.* $9x + 3ab^2 - 3y^2 - y$.

8. Add together $4a - 3b^2 + cx^2$, $b^2 - 3a + 3$, $2a + 3b^2 - cx^2$, and $2b^2 - 2a - 9$. *Ans.* $a + 3b^2 - 6$.

9. Add together $7b - 2a^2 - xy$, $5a^2 - 6b + 3xy$, $b - 3a^2 + 4$, and $a^2 - b - 3xy$. *Ans.* $b + a^2 - xy + 4$.

10. Add together $3b^2 - 2a^3 + 13$, $3a^3 - 2b^2 - 5$, $4ab + 7a^3 - 3$, and $2b^2 - a^3 + ab$. *Ans.* $3b^2 + 7a^3 + 5 + 5ab$.

11. Add together $ax^2 - 2y + b$, $2y + 2ax^2 - 3b$, $4ax^2 - y - b$, and $2b - 3ax^2 + b$. *Ans.* $4ax^2 - y$.

12. Add together $2c^2 + a^2 + 3bc$, $5c^2 - 3a^2 - 2bc$, $c^2 + 2a^2 - bc$, and $b + bc - 3$. *Ans.* $8c^2 + bc + b - 3$.

13. Add together $ab + a^2c - 5$, $3ab - 3a^2c + 7$, $2a^2c - 2ab - 3$, and $ab + a^2c + 5$. *Ans.* $3ab + a^2c + 4$.

14. Add together $3b^2 - 2a^2x + b$, $-b^2 + 3a^2x - 3b$, $b^2 - a^2x + c$, and $3b^2 + b - 3c$. *Ans.* $6b^2 - b - 2c$.

15. Add together $2a^2 + 3 - ac$, $3a^2 - 7 + ac$, $3ac - 5a^2 + 9$, and $3ac + 4 - a^2$. *Ans.* $-a^2 + 9 + 6ac$.

16. Add together $b^2c + 2 - y^2$, $y^2 - 3b^2c - 10$, $2b^2c - 3 + 2y^2$, and $b^2 - y^2 + 5$. *Ans.* $-6 + y^2 + b^2$.

17. Add together $a^3 + bc - \frac{1}{2}c$, $2a^3 - bc + \frac{3}{4}c$, $3a^3 + 3bc - \frac{1}{4}c$, and $a^3 + bc + c$. *Ans.* $7a^3 + 4bc + c$.

18. Add together $b^3 - 3a^2 + \frac{2}{3}c^2$, $2b^2 + 5a^2 + \frac{1}{2}c^2$, $b^2 - \frac{1}{3}c + 2$, and $5b^2 + 3a^2 - 2c + 3y$. *Ans.* $b^3 + 5a^2 + 1\frac{1}{6}c^2 + 8b^2 - 2\frac{1}{3}c + 2 + 3y$.

(32.) When the same Polynomial is enclosed in two or more *parentheses*, (11), this polynomial enters into a calculation in the same manner as the *common factor* of similar monomials.

Thus the Sum of $3(a+b)$ and $5(a+b)$, is evidently $8(a+b)$.

19. Add together $2(a-3b)+c$, $3(a-3b)-3c$, $-4(a-3b)+c$, and $5(a-3b)+3c$. *Ans.* $6(a-3b)+2c$.

20. Add together $3a^2+2(a-c^2)$, $a^2-3(a-c^2)$, $4a^2+5(a-c^2)$ and $-7a^2+(a-c^2)$. *Ans.* $a^2+5(a-c^2)$.

21. Add together $5(a+b-c)+3b^2$, $3(a+b-c)-2b^2$, $2(a+b-c)$ and $3b^2+2y+b^2$. *Ans.* $10(a+b-c)+5b^2+2y$.

22. Add together $2ab^2+3a(b+c^2)$, $ab^2-2a(b+c^2)$, $a(b+c^2)$ and $-ab^2-3a(b+c^2)$. *Ans.* $2ab^2-a(b+c^2)$.

23. Add together $5+\frac{1}{2}(c-d+m)+2b$, $1+\frac{3}{4}(c-d+m)-b+2$, and $2(c-d+m)+\frac{1}{2}b$. *Ans.* $8+3\frac{1}{4}(c-d+m)+1\frac{1}{2}b$.

SUBTRACTION.

(33.) Algebraic SUBTRACTION consists in finding the *difference* between the two quantities; that is, in finding *what quantity added to the quantity subtracted*, will produce that from which the subtraction is made.

Thus the difference found by subtracting $3a$ from $5a$ is $2a$, because $2a$ added to $3a$ produces $5a$, (27).

But the difference found by subtracting $-3a$ from $5a$ is $8a$; because $8a$ must be added to $-3a$ to produce $5a$, (29).

In the second example, observe that *subtracting* $-3a$ has an effect contrary to that of subtracting $3a$ in the first example, (25); the Subtracting of a *negative* quantity being the same as the Adding of an equal *positive* quantity.

Subtraction of Monomials.

(34.) A *monomial* is subtracted from another quantity, by changing the *sign* of the monomial, and then *adding it*, algebraically, to the other quantity.

Thus $5a$ subtracted from $2a$ gives $-3a$, because $-3a$ must be added to $5a$, to produce $2a$, (29); and this difference, $-3a$, will be found by changing $5a$ to $-5a$, and adding the $-5a$ to the $2a$.

☞ What difference will be found by subtracting $5a$ from $9a$? and why is that difference *true*? By subtracting $9a$ from $5a$?

What difference will be found by subtracting $3ab$ from $4ab$? and why is that difference true? By subtracting $-5x$ from $-10x$? and why is that difference true? By subtracting $7ax$ from $-3ax$.

(35.) When the two quantities are *dissimilar*, a monomial is subtracted by changing its *sign*, and then connecting it with the other quantity, in a Polynomial.

Thus $2x$ subtracted from $3a$ gives $3a-2x$; and $-2x$ subtracted from $3a$ gives $3a+2x$, (34), (30).

☞ What difference will be found by subtracting $2a$ from $3x$? By subtracting $-3b$ from $5ax$? By subtracting ax from $10b$? By subtracting $4y$ from $-9x^2$?

From the preceding we derive

RULE II.

(36.) *For the Subtraction of Algebraic Quantities.*

Change the sign of each term *to be subtracted*, + to - and - to +, or conceive these signs to be changed, and then proceed as in Algebraic Addition.

EXAMPLE.

To subtract $a^2 + 2ab - 3x - z$ from $3a^2 - 5ab + 5x - y$.

$$\begin{array}{r} 3a^2 - 5ab + 5x - y \\ a^2 + 2ab - 3x - z \\ \hline 2a^2 - 7ab + 8x - y + z \end{array}$$

Having set the polynomial which is to be subtracted under the one from which the subtraction is to be made, we conceive each term in the second line to have its *sign changed*; then $-a^2$ added to $3a^2$ makes $2a^2$; $-2ab$ added to $-5ab$ makes $-7ab$; and $3x$ added to $5x$ makes $8x$. These results, together with the $-y$ and $+z$, are connected in a polynomial.

EXERCISES.

- From $4ab + 3c^2 - xy$, subtract $2ab - c^2 + 3xy - 2z$.
Ans. $2ab + 4c^2 - 4xy + 2z$.
- From $5a^2 + 3bc - cd + 3x$, subtract $2a^2 - bc + 3cd$.
Ans. $3a^2 + 4bc - 4cd + 3x$.
- From $7a^2b - abc + 4xy$, subtract $a^2b + 5abc - xy + mz$.
Ans. $6a^2b - 6abc + 5xy - mz$.
- From $8a^3 - 4a^2b - 2bc + 10$, subtract $3a^3 + a^2b - 5$.
Ans. $5a^3 - 5a^2b - 2bc + 15$.
- From $3b^2c + abx + 2xy^2$, subtract $b^2c - 3abx + 2xy^2 - 3$.
Ans. $2b^2c + 4abx + 3$.
- From $4a^2c - 3c^2b + 6y + 20$, subtract $3c^2b + 6y + 3by$.
Ans. $4a^2c - 6c^2b + 20 - 3by$.
- From $-3a + 5b^2 - 7xy$, subtract $3b^2 + 2a - xy + bc$.
Ans. $-5a + 2b^2 - 6xy - bc$.
- From $2ab + 3bc - 2ax^2 + 15$, subtract $7ab + 3bc - 10$.
Ans. $-5ab - 2ax^2 + 25$.
- From $5ax - 3bc + 2b^2$, subtract $6ax - bc - 3b^2 + y^2$.
Ans. $-ax - 2bc + 5b^2 - y^2$.
- From $a^2b + 3abc + b^2c - 7$, subtract $\frac{1}{2}a^2b - abc + \frac{1}{3}b^2c$.
Ans. $\frac{1}{2}a^2b + 4abc + \frac{1}{3}b^2c - 7$

When one quantity is to be subtracted from the Sum of two or more quantities, the operation will be expedited by changing the signs of the *subtractive* quantity, and then *adding all the quantities together*.

11. From the Sum of $2a+3bc-1$, and $3a-4bc+3$, subtract $4a-bc-2+y$.
Ans. $a+4-y$.

12. From the Sum of $ax^2-6b^2+3ay^2$, and $2ax^2+6b^2-ay^2$, subtract $-3ax^2+b^2+2ay^2-5$.
Ans. $6ax^2-b^2+5$.

13. From the Sum of $3ab+bc-5x^2$, and $ab-3bc+x^2-3$, subtract $5bc-3x^2-4+y^2$.
Ans. $4ab-7bc-x^2+1-y^2$.

14. From the Sum of $5x-3xy+7y^2$, and $5xy-3x-4y^2-9$, subtract $-x+xy-3y^2+7$.
Ans. $3x+xy+6y^2-16$.

15. From the Sum of $2a-b+3cd$, $5a+3b-cd-3$, and $a+2b-cd$, subtract $4a-5b-cd$.
Ans. $4a+9b+2cd-3$.

16. From the Sum of $3x^2+y-5xy$, $2x^2-3y+xy$, and $4y-3x^2-xy+5$, subtract $x^2+5y-xy-3$.
Ans. $x^2-3y-4xy+8$.

17. From the Sum of $2ay^2-2b+3ax$, $2ay^2+b-ax$, and $3b-ay^2+3-y$, subtract $3ay^2+4b-ax+y$.
Ans. $-2b+3ax+3-2y$.

18. From the Sum of $3a^2+xy^2-2by$, $5a^2+3xy^2-3by$, and $3xy^2+4a^2+by$, subtract $2a^2-xy^2-5by+5$.
Ans. $10a^2+8xy^2+by-5$.

19. From the Sum of $a^2b^2-3y^2+5xy$, $3a^2b^2+3y^2-2xy$, and $5y^2+3a^2b^2-xy+6$, subtract $a^2b^2+xy-y^2-3$.
Ans. $6a^2b^2+6y^2+xy+9$.

20. From the Sum of $5y^2-3ax-2bc$, $4ax-2y^2+5bc$, and $3ax+y^2-bc+5$, subtract $ax-bc+3y^2-1$.
Ans. $y^2+3ax+3bc+6$.

21. From the Sum of a^2y-x^2+bc-d , $3x^2-3bc+5d$, and $a^2y+3x^2-4bc-d$, subtract $5x^2-2bc+7d$.
Ans. $2a^2y-4bc-4d$.

22. From the Sum of $5b^2+3ax+2y$, $3b^2-ax-y+1$, and $4ax-b^2-5y+2$, subtract $3b^2+2ax-y+10$.
Ans. $4b^2+4ax-3y-7$.

23. From the Sum of $7a+6x^2-y^2+1$, x^2+3y^2-c-3 , and $7x^2+y^2-3c+5$, subtract $5a+2x^2-2y^2+5c+2$.
Ans. $2a+12x^2+5y^2-9c+1$.

24. From the Sum of $2(a-x)+y$, $3(a-x)+2y$, and $5(a-x)-y$, subtract $4(a-x)+y-2$, (32).
Ans. $6(a-x)+y+2$.

25. From the Sum of $2b(a+c)^2+3$, $b(a+c)^2-1$, and $3b(a+c)^2-2$, subtract the sum of $b(a+c)^2+b$, and $5b(a+c)^2-b$.
Ans. 0.

Indicated Subtraction.

(37.) It is sometimes expedient merely to *indicate* the subtraction of a quantity, without performing the operation.

To denote the subtraction of a positive monomial, nothing more is necessary than to place the *sign* $-$ before it; thus $a-b$ denotes that b , that is, *positive* b , is to be subtracted from a .

The subtraction of a *negative* monomial, will be denoted by enclosing the quantity, with its negative sign, in a (), and prefixing the sign $-$ to the parenthesis.

Thus $a-(-b)$ denotes that *negative* b is to be subtracted from a . When the subtraction is performed, the expression becomes $a+b$.

The subtraction of a *polynomial* will be denoted by enclosing the polynomial in a (); and prefixing the sign $-$ to the parenthesis.

Thus $a-(b+c+d)$; in which the *sign* $-$ denotes that the *enclosed polynomial* is to be subtracted. When the subtraction is performed, the expression becomes $a-b-c-d$.

Change of Signs in a Polynomial.

(38.) The *signs of the value* of a Polynomial, will evidently be changed, by changing the signs of all the terms of the Polynomial.

The value of a Polynomial is therefore not affected by *changing the signs* of any or all of its terms, enclosing those terms in a (), and prefixing the *sign* $-$ to the parenthesis.

Thus $a+b-c$ is equivalent to $a-(-b+c)$, or $a-(c-b)$; for if $(c-b)$ be subtracted, as is required by the *sign* $-$ prefixed to it, the result will be $a+b-c$.

26. Under what different forms may the value of the polynomial $ab+2cd-3x+5$ be expressed?

27. Under what different forms may the value of the polynomial $ax-3y+4b^2-5c-7$ be expressed?

MULTIPLICATION.

(39.) Algebraic MULTIPLICATION consists in finding the Product of one quantity taken as many times, *additively*, or *subtractively*, as there are units in another quantity.

The quantity to be multiplied is called the *multiplicand*, and the multiplying quantity the *multiplier*: both together are called the *factors* of the product, (12).

When the Multiplier is *positive*, the multiplicand is repeated positively, or is repeatedly *added*.

Thus $5a \times 3$, $5a$ multiplied by *positive* 3, is $5a + 5a + 5a$, equal to $15a$; the multiplicand $5a$ being repeatedly *added*.

When the Multiplier is *negative*, the multiplicand is repeated *negatively*, or is repeatedly *subtracted*.

Thus $5a \times -3$, $5a$ multiplied by *negative* 3, is $-5a - 5a - 5a$, which is equal to $-15a$; the multiplicand $5a$ being repeatedly *subtracted*.

Multiplication of Monomials.

(40.) The Product of two *monomials* consists of the product of their *coefficients* multiplied into all their *literal factors*.

For example, $3ac \times 2x$ is equal to $6acx$; for the factors may be taken in the order $3 \times 2acx$, which becomes $6acx$ by substituting 6 for 3×2 .

☞ What is the Product of $3a \times 4b$? and how do you reason in finding that product? What is the Product of $5a^2b \times 2x$? and how do you reason in finding it? Of $7ac^2 \times xy$? Of $xy^2 \times 5$? Of $3 \times 7a^2b^2$?

(41.) When the *same letter* occurs in both the monomials multiplied together, its *exponent* in the Product will be the *sum* of its exponents in the two factors.

Thus $3a^2x \times 2ax$, or $3a^2x^1 \times 2a^1x^1$, is equal to $6a^2axx$, (40), and this product becomes $6a^3x^2$ by substituting a^3 and x^2 for their equivalents a^2a and xx , (13) and (16).

Observe that the exponents of a and x in the product $6a^3x^2$, are the *sums* of the exponents of the same letters, respectively, in the two monomials which are multiplied together.

☞ What is the Product of $4a^3 \times 2a^2$? Of $3ax \times 5ax^2$? Of $7a \times 2a^3b$? Of $4ac \times acb$? Of $a^2b^3 \times 5b$? Of $8a^2cb \times c$? Of $a^2b \times a^3b^2$?

Sign of the Product.

(42.) The Product of two quantities is *positive* when they both have the *same sign*, and *negative* when they have contrary signs;—in other words, *like signs* produce +, and *unlike signs* produce —, in multiplying.

When both the quantities are *positive*, their product is evidently *positive*,—as in common Arithmetic. Thus $3a \times 2$ is $3a + 3a$, equal to $6a$.

When both the quantities are *negative*, their product is *positive*, because such product results from repeatedly *subtracting a negative quantity*, (36).

For example, in $-3a \times -2$ the multiplier -2 denotes that $-3a$ is to be taken twice *subtractively*; and since the *sign* of a quantity is changed in subtracting it, the product is $3a + 3a$, or $6a$.

When one of the quantities is *positive* and the other *negative*, their product is *negative* because it results from repeatedly *subtracting a positive*, or adding a *negative* quantity.

Thus in $3a \times -2$, the $3a$ is to be taken twice *subtractively*, and the product is therefore $-3a - 3a$, equal to $-6a$.

☞ What is the Product of $-2x \times -3$? and how do you reason on the *sign* of the product? Of $-3ab \times -2$? Of $-5a^2 \times -3$? Of $-x \times -5$?

What is the Product of $5ax \times -4$? and how do you reason on the *sign* of the product? Of $4a^2c \times -3$? Of $7x^2 \times -5$? Of $b^2 \times -9$.

What is the Product of $-2ab \times 3$? and how do you reason on the *sign* of the product? Of $-3a^2x \times 4$? Of $-3a^2 \times 2$? Of $-c^3 \times 5\frac{1}{2}$?

What is the Product of $4x^2 \times -7$? and how do you reason on the *sign* of the product? Of $5ay \times -3$? Of $-3a \times -4a$? Of $5b \times -5$?

When the Multiplier or the Multiplicand is 0.

(43.) When *either* of the two factors multiplied together is 0, the product will be 0; for it is evident that 0 repeated any number of times, produces only 0; and the product is the same when the multiplicand and the multiplier are taken *the one for the other*.

Thus 5×0 is 0; $a \times 0$ is 0; $(a+b-c) \times 0$ is 0.

The preceding principles and the two following rules, embrace the subject of Algebraic Multiplication.

RULE III.

(44.) *To Multiply a Monomial into a Polynomial.*

Find the product of the monomial into *each term* of the polynomial, separately, and connect these products in a polynomial with the proper *sign prefixed to each*, (40), (41), (42).

EXAMPLE.

To multiply $3ab+bc^2-2xy-5$ by $2a^2x$

$$\begin{array}{r} 3ab+bc^2-2xy-5 \\ 2a^2x \\ \hline 6a^3bx+2a^2bc^2x-4a^2x^2y-10a^2x \end{array}$$

It will be most convenient to multiply from *left to right*. The first term of the multiplicand is *positive*, and the multiplier being also *positive*, the first term, $6a^3bx$, of the product is *positive*. In like manner the second term, $2a^2bc^2$, of the product is positive, &c.

EXERCISES.

1. Multiply $2a^2-3b+c-2y$ by 3. *Ans.* $6a^2-9b+3c-6y$.
2. Multiply $3x^2-xy-y^2+4$ by 2. *Ans.* $6x^2-2xy-2y^2+8$.
3. Multiply $4b+c^2-3y+1$ by $-y$. *Ans.* $-4by-c^2y+3y^2-y$.
4. Multiply $-a+3b-2x^2-3$ by $-2x$. *Ans.* $2ax-6bx+4x^3+6x$.
5. Multiply ab^2-b^3+2c-y by -5 . *Ans.* $-5ab^2+5b^3-10c+5y$.
6. Multiply $-3+ax-1+by$ by $-a$. *Ans.* $3a-a^2x+a-aby$.
7. Multiply $7x+by^2-4c+d^2$ by 4. *Ans.* $28x+4by^2-16c+4d^2$.
8. Multiply a^2b-c^2+x-2 by $3a$. *Ans.* $3a^3b-3ac^2+3ax-6a$.

RULE IV.

(45.) *To Multiply a Polynomial into a Polynomial.*

Multiply one of the polynomials by *each term*, separately, of the *other polynomial*, and add together the several products.

EXAMPLE.

To multiply $2a^2 + 4ac - c^2$ by $3a - 5c$

$$\begin{array}{r}
 2a^2 + 4ac - c^2 \\
 3a - 5c \\
 \hline
 6a^3 + 12a^2c - 3ac^2 \\
 \quad - 10a^2c - 20ac^2 + 5c^3 \\
 \hline
 6a^3 + 2a^2c - 23ac^2 + 5c^3.
 \end{array}$$

We multiply the first polynomial by $3a$, and then by $-5c$, according to the preceding Rule. The two products thus obtained are set with similar terms one under another, and added together.

The correctness of this Rule, as well as of the preceding one, will be evident upon considering, *first*, that if *each part* of a quantity be multiplied, the *whole* will be multiplied; and, *secondly*, that if one quantity be multiplied by *each part of another quantity*, the former will be multiplied by the *whole of the latter*.

EXERCISES.

- | | |
|---|---|
| 9. Multiply $a^2 - 2a + 5$ by $a + 3$. | <i>Ans.</i> $a^3 + a^2 - a + 15$ |
| 10. Multiply $2x^2 + 3x - 1$ by $x - 5$. | <i>Ans.</i> $2x^3 - 7x^2 - 16x + 5$. |
| 11. Multiply $a^2 - 2ax + x^2$ by $a + x$. | <i>Ans.</i> $a^3 - a^2x - ax^2 + x^3$. |
| 12. Multiply $x^2 + 3xy + y^2$ by $x - y$. | <i>Ans.</i> $x^3 + 2x^2y - 2xy^2 - y^3$. |
| 13. Multiply $b^2 - 3bc - c^2$ by $2b + c$. | <i>Ans.</i> $2b^3 - 5b^2c - 5bc^2 - c^3$. |
| 14. Multiply $2a^2 + 3ac - c^2$ by $a - 2c$. | <i>Ans.</i> $2a^3 - a^2c - 7ac^2 + 2c^3$. |
| 15. Multiply $2b^2 - 2bx + x^2$ by $2b - x$. | <i>Ans.</i> $4b^3 - 6b^2x + 4bx^2 - x^3$. |
| 16. Multiply $b^3 + b^2 + b$ by $b^3 + b^2$. | <i>Ans.</i> $b^6 + 2b^5 + 2b^4 + b^3$. |
| 17. Multiply $x^2 - 2c^2 + 2y$ by $c^2 + y$. | <i>Ans.</i> $c^2x^2 - 2c^4 + x^2y + 2y^2$. |
| 18. Multiply $a^2 + bx + y$ by $a^2 - bx$. | <i>Ans.</i> $a^4 + a^2y - b^2x^2 - bxy$. |

- 19 Find the Product of $(2a^2 - 4ab + 2b^2)(3a - 2b)$.
Ans. $6a^3 - 16a^2b + 14ab^2 - 4b^3$.
- 20 Find the Product of $(3x^2 - 2xy + 2y^2)(2x^2 + 3xy)$.
Ans. $6x^4 + 5x^3y - 2x^2y^2 + 6xy^3$.
- 21 Find the Product of $(2a^2 + 3ab - b^2)(a^2 - ab + b^2)$.
Ans. $2a^4 + a^3b - 2a^2b^2 + 4ab^3 - b^4$.
- 22 Find the Product of $(3x^2 - 2xy + 5)(x^2 + 2xy - 3)$.
Ans. $3x^4 + 4x^3y - 4x^2 - 4x^2y^2 + 16xy - 15$.
23. Find the Product of the three polynomials $(a+b)$, $(a-b)$, and $(a^2 + ab + b^2)$.
Ans. $a^4 + a^3b - ab^3 - b^4$.

DIVISION.

(46.) Algebraic DIVISION consists in finding a factor or Quotient, which, multiplied into a given *divisor*, will produce a given *dividend*.

Thus $6a^2x \div 2a$, $6a^2x$ divided by $2a$, gives the *quotient* $3ax$, because this factor, multiplied into the divisor $2a$, produces the dividend $6a^2x$.

Division of Monomials.

(47.) The Quotient of two *monomials* will be found by dividing the *coefficient* of the divisor into that of the dividend, and *subtracting the exponents* in the divisor from those of the same letters, respectively, in the dividend.

Thus $10a^3x^2y \div 5a^2x$ gives the quotient $2axy$, because this quotient, *multiplied into the divisor* $5a^2x$, produces the dividend $10a^3x^2y$; and $2axy$ is found by dividing 5 into 10, and diminishing the exponents of a and x in the dividend by the exponents of a and x in the divisor.

The dividend being the *product* of the divisor and quotient, the *exponents* in the dividend are the *sums* of the exponents of the same letters in the divisor and quotient, (41); hence the exponents in the quotient will always be found by subtracting as above.

☞ What is the Quotient of $6a^3x^2y \div 3ax$? and how do you prove that quotient true? What is the Quotient of $4ab^2c \div b$? and how do you prove that quotient true? What is the Quotient of $9a^2bx \div 2a$? and how do you prove that quotient true?

When any Exponent is reduced to 0.

(48.) Any quantity whatever with *exponent* 0 is equivalent to *unity*; and a *factor* with this exponent may therefore be canceled.

For example, $a^2 \div a^2$ gives the quotient a^0 , by subtracting the exponent of the divisor from that of the dividend.

But any quantity contains itself *once*, and therefore $a^2 \div a^2$ also gives the quotient 1; and these two quotients, a^0 and 1, must be *equivalent to each other*.

Since a in the preceding illustration may represent any quantity we please, any quantity whatever with *exponent* 0, is equivalent to 1, or is a *symbol of unity*.

To divide $10a^3b$ by $5a^3$.

Subtracting the exponent in the divisor from the exponent of a in the dividend, we find the quotient $2a^0b$. The factor a^0 is equivalent to 1, and may therefore be canceled without affecting the value of the quotient.

The quotient of $10a^3b \div 5a^3$ may therefore be given under the two different forms $2a^0b$ and $2b$.

☞ Under what *two different forms* may the Quotient of $6a^2x \div 2a^2bc$ be represented? Of $8ab^2 \div 2b^2$? Of $10abc \div 5a$? Of $9ax^2 \div x^2$? Of $7abc^2 \div 7c^2$? Of $5ax \div ax$? Of $3aby^2 \div ay^2$? Of $10x^2 \div x^2$?

Sign of the Quotient.

(49.) The Quotient of two quantities is *positive* when they both have the *same sign*, and *negative* when they have contrary signs;— in other words, *like signs* produce +, and *unlike signs* produce —, in dividing.

Thus $+ax \div +a$ gives $+x$, because $+a \times +x$ is $+ax$;
 $-ax \div -a$ gives $+x$, because $-a \times +x$ is $-ax$;
 $+ax \div -a$ gives $-x$, because $-a \times -x$ is $+ax$;
 and $-ax \div +a$ gives $-x$, because $+a \times -x$ is $-ax$; (42.)

From these examples, it will be seen that the *sign of the quotient* must be such that the quotient multiplied into the *divisor* shall produce the *dividend*.

It thus appears that the rule for the sign of the *quotient*, is the same as that for the sign of the *product*, of two quantities. The learner must be careful not to apply this rule in finding the *sum*, or *difference*, of two quantities.

☞ What is the Quotient of $4a^2x \div 2$? and how do you reason on the *sign* of the quotient? Of $6ac^2 \div 2c$? Of $9x^3y \div 3xy$? Of $a^3 \div a$?

What is the Quotient of $-7a^3 \div -a^2$? and how do you reason on the *sign* of the quotient? Of $-12c^2 \div -4$? Of $-a^3b \div -ab$? Of $-b^4 \div b^2$?

What is the Quotient of $8a^2 \div -a^2$? and how do you reason on the *sign* of the quotient? Of $10ab \div -2b$? Of $4a^2c \div -a$? Of $5x^2 \div -x^2$?

What is the Quotient of $-7a^3 \div 7a$? and how do you reason on the *sign* of the quotient? Of $-12ac \div 3a$? Of $-9a^3b \div 3$? Of $-20a^4 \div 4$?

What is the Quotient of $10a^3 \div 5a$? and why is that quotient true? What is the Quotient of $12x^2 \div -3$? and why is that quotient true? What is the Quotient of $-9a^3by \div 3a^3$? and why is that quotient true? What is the Quotient of $-20a^4x^3 \div -5ax$? and why is that quotient true? What is the Quotient of $-100axy \div -xy$? and why is that quotient true?

When the Dividend or the Divisor is 0.

(50.) The quotient of 0 divided by any quantity, is 0; but the quotient of any quantity divided by 0, is *infinitely great*.

First, let a represent any quantity we please; then $0 \div a$ gives the quotient 0, because this quotient multiplied into the *divisor* a , produces the *dividend* 0, (43).

Secondly, a divisor may be taken so small as to be contained a *great number of times* in any given dividend, represented by a ; if the divisor be still diminished, the quotient will be increased; and if the divisor be diminished *without limit*, the quotient will be *increased without limit*.

If therefore the divisor were diminished to 0, the quotient would be *unlimited*, that is, *infinitely great*.

The character ∞ is the sign of *infinity*; we have then $0 \div a$ equal to 0; and $a \div 0$ equal to ∞ , *infinity*

The preceding principles, and the two following Rules, embrace the subject of Algebraic Division, when the Quotient is an *integral* quantity

RULE V.

(51.) *To Divide a Monomial into a Polynomial.*

Find the quotient of the monomial divided into each term of the polynomial, separately, and connect these quotients in a polynomial, with the proper *sign prefixed to each*. (47), (48), (49).

EXAMPLE.

To divide $20a^2x - 15ax^2 + 30axy^2 - 5ax$ by $5ax$.

$$\begin{array}{r} 5ax \overline{) 20a^2x - 15ax^2 + 30axy^2 - 5ax} \\ \underline{4a \quad - 3x \quad + 6y^2 \quad - 1} \end{array}$$

Dividing into the first term of the dividend, we find the *quotient term* $4a$; dividing into the second term of the dividend, we find the quotient $-3x$; and so on, through the dividend.

EXERCISES.

1. Divide $3x^3 + 6x^2 + 3ax - 15x$ by $3x$. *Ans.* $x^2 + 2x + a - 5$.
2. Divide $2ab - 6a^2x + 8a^3y - 2a$ by $2a$. *Ans.* $b - 3ax + 4a^2y - 1$
3. Divide $14a^2 - 7ab + 21ax - 21a$ by $7a$. *Ans.* $2a - b + 3x - 3$.
4. Divide $a^3x + 3a^2x^2 - 6ax^3 + 3ax$ by a . *Ans.* $a^2x + 3ax^2 - 6x^3 + 3x$.
5. Divide $5b^2 - 10b^3 + 5b^4y - 15b^5$ by $5b$. *Ans.* $b - 2b^2 + b^3y - 3b^4$.
6. Divide $bx^2 + 2x^3 - 8cx^4 + 7x^5$ by x^2 . *Ans.* $b + 2x - 8cx^2 + 7x^3$.
7. Divide $4a^4 - 8a^3 - 4a^2b + 8a$ by $2a$. *Ans.* $2a^3 - 4a^2 - 2ab + 4$
8. Divide $ay^4 + a^2y^3 - a^3y^2 - ay$ by ay . *Ans.* $y^3 + ay^2 - a^2y - 1$.
9. Divide $-y + by^2 - 5y^3 + 3y^4$ by $-y$. *Ans.* $1 - by + 5y^2 - 3y^3$
10. Divide $3b^3 - 9b^2 + 12b - 15$ by $+ 3$. *Ans.* $b^3 - 3b^2 + 4b - 5$
11. Divide $-c^5 + 3c^4 - 6c^3 + c^2$ by $-c^2$. *Ans.* $c^3 - 3c^2 + 6c - 1$
12. Divide $8y^3 - 4ay + a^2y^2 - 3y$ by y . *Ans.* $8y^2 - 4a + a^2y - 3$
13. Divide $-10 + 20a - 15a^2 + 20$ by -5 . *Ans.* $2 - 4a + 3a^2 - 4$.
14. Divide $a^3b - a^2b^2 + a^3b^2 - a^4b$ by ab . *Ans.* $a^2 - ab + a^2b - a^3$.
15. Divide $x^4y^4 + x^3y^3 - x^2y^2 - xy$ by xy . *Ans.* $x^3y^3 + x^2y^2 - xy - 1$

RULE VI.

(52.) *To Divide a Polynomial into a Polynomial.*

1. Arrange the divisor and dividend according to the *ascending* or the *descending powers* of the same letter, (22).

2. Divide the *first term* of the divisor into the *first term* of the dividend; multiply the whole divisor by the *quotient* term; subtract the product from the dividend; divide into the remainder, as before, and so on,—observing to connect the several quotient terms in a polynomial, with the proper *sign* prefixed to each.

EXAMPLE.

To divide $6a^4x + 3a^3x^2 - 4a^2x^4 + x^6$ by $2ax + x^2$

$$\begin{array}{r}
 2ax + x^2 \) \ 6a^4x^2 + 3a^3x^3 - 4a^2x^4 + x^6 \ (\ 3a^3x - 2ax^3 + x^4 \\
 \underline{6a^4x^2 + 3a^3x^3} \\
 -4a^2x^4 + x^6 \\
 \underline{-4a^2x^4 - 2ax^5} \\
 2ax^5 + x^6 \\
 \underline{2ax^5 + x^6} \\

 \end{array}$$

The divisor and dividend are here arranged according to the *ascending* powers of a , or the *ascending* powers of x .

The first term $2ax$ of the divisor, divided into the first term $6a^4x^2$ of the dividend, gives the quotient term $3a^3x$. Multiplying the whole divisor by this quotient term, and subtracting the product from the dividend, the remainder is $-4a^2x^4 + x^6$.

We next divide the first term of the divisor into the first term, $-4a^2x^4$, of the remainder, and find the quotient term $-2ax^3$; &c.

By this Rule the divisor is multiplied by each *part of the quotient*, and the successive products subtracted from the dividend. When the dividend is thus *exhausted*, the product of the divisor and quotient is *equal to the dividend*; and the quotient is thus proved to be correct.

The Divisor, when a polynomial, is sometimes set on the left of the dividend, and the quotient under the divisor. This arrangement is thought to afford greater facility in multiplying the divisor by the quotient term.

EXERCISES.

16. Divide $a^2 - 2ax + x^2$ by $a - x$. *Ans.* $a - x$.
17. Divide $x^3 - 3ax^2 + 3a^2x - a^3$ by $x - a$. *Ans.* $x^2 - 2ax + a^2$.
18. Divide $6a^4 + 9a^2 - 15a$ by $3a^2 - 3a$. *Ans.* $2a^2 + 2a + 5$
19. Divide $a^4 + a^2x^2 + x^4$ by $a^2 - ax + x^2$. *Ans.* $a^2 + ax + x^2$.
20. Divide $2x^3 - 19x^2 + 26x - 16$ by $x - 8$. *Ans.* $2x^2 - 3x + 2$.
21. Divide $4x^4 - 5x^2c^2 + c^4$ by $2x^2 - 3xc + c^2$. *Ans.* $2x^2 + 3xc + c^2$.
22. Divide $a^4 - 2a^2x^2 + x^4$ by $a^2 + 2ax + x^2$. *Ans.* $a^2 - 2ax + x^2$.
23. Divide $12 - 4a - 3a^2 + a^3$ by $4 - a^2$. *Ans.* $3 - a$.
24. Divide $4y^4 - 9y^2 + 6y - 1$ by $2y^2 + 3y - 1$. *Ans.* $2y^2 - 3y + 1$.
25. Divide $2x^4 - 32$ by $x - 2$. *Ans.* $2x^3 + 4x^2 + 8x + 16$.
26. Divide $4a^5 - 64a$ by $2a - 4$. *Ans.* $2a^4 + 4a^3 + 8a^2 + 16a$.
27. Divide $6y^6 - 96y^2$ by $3y - 6$. *Ans.* $2y^5 + 4y^4 + 8y^3 + 16y^2$.
28. Divide $a^4 + 4x^4$ by $a^2 - 2ax + 2x^2$. *Ans.* $a^2 + 2ax + 2x^2$.
29. Divide $a^5 - x^5$ by $a^4 + a^3x + a^2x^2 + ax^3 + x^4$. *Ans.* $a - x$.
30. Divide $y^4 + 4y^2z^2 - 32z^4$ by $y + 2z$. *Ans.* $y^3 - 2y^2z + 8yz^2 - 16z^3$.

(53.) The *indicated Product* of two or more factors is divided by any quantity, when *either of the factors* is divided by that quantity.

Thus to divide $3a^2x(x^2 + y^2)$ by $3x$, we divide the factor $3a^2x$, and find the quotient $a^2(x^2 + y^2)$.

In the following exercises, the first of the given *factors* may be divided by the given divisor.

31. Divide $(a^2 + 2ay + y^2)(b^3 - cd^2 + 3)$ by $a + y$. *Ans.* $(a + y)(b^3 - cd^2 + 3)$.
32. Divide $(2x^3 - 6ax^2 + 6a^2x - 2a^3)(c^2 + 3cy - y^2)$ by $x - a$. *Ans.* $(2x^2 - 4ax + 2a^2)(c^2 + 3cy - y^2)$.
33. Divide $(a^4 + 4y^4)(3xy^2 - 5y^3 + 3y^4 - 4)$ by $a^2 - 2ay + 2y^2$. *Ans.* $(a^2 + 2ay + 2y^2)(3xy^2 - 5y^3 + 3y^4 - 4)$.
34. Divide $(8a^4 - 2a^3x - 13a^2x^2 - 3ax^3)(y^2 + 2xy)(y^3 + 3x^2y^2 - x^3)$ by $4a^2 + 5ax + x^2$. *Ans.* $(2a^2 - 3ax)(y^2 + 2xy)(y^3 + 3x^2y^2 - x^3)$.

CHAPTER III.

COMPOSITE QUANTITIES.—COMMON MEASURE.—COMMON MULTIPLE.

Composite and Prime Quantities.

(54.) A *composite* quantity is one which is the *product* of two factors, each differing from *unity*; and a *prime* quantity is one which is not the product of such factors.

Thus $3a^2$ is a *composite*, while $a+b$ is a *prime* quantity.

☞ Is ab a composite or a prime quantity? Is $a+5$ a composite or a prime quantity? a^2 ? $7x^2$? $2a+3b$? $5a-5c^2$?

Decomposition of Quantities.

(55.) *Decomposing* a composite quantity consists in resolving it into its *factors*. Any *divisor* of the quantity, and the corresponding *quotient*, are two factors into which it may be resolved, (46.)

Thus if the binomial $3x+6ax$ be divided by $3x$, the quotient will be $1+2a$; the binomial may therefore be resolved into the factors

$$3x(1+2a), \quad 3x \text{ into the binomial } 1+2a.$$

Resolve $2a+4ax-6a^2x^2$ into component factors.

Resolve $a^2x-3ax^2+8ay^2$ into component factors.

Resolve $4a^2+a^2b-5a^2y$ into component factors.

Resolve $2a^3-3ax+7a^4y$ into component factors.

Resolve $5ax+5a^2x-10a^3x^3$ into component factors.

Resolve $7ab-14ab^2-14abx$ into component factors.

A *monomial factor* may generally be found by mere inspection, and a composite polynomial be resolved into a monomial and a polynomial factor, as above.

No general Rule can be given for resolving a polynomial into the *polynomial factors* of which it may be composed. But there are particular Binomials which have well known *binomial divisors*, by means of which such Binomials may be resolved into two factors, (55).

The following divisions will be found, on trial, to terminate without remainders.

(56.) The Difference of two quantities will divide the difference of any *powers of the same degree* of those quantities.

Thus $a-b$ will divide a^2-b^2 , or a^3-b^3 , or a^4-b^4 , &c.

(57.) The Sum of two quantities will divide the *sum* of any *odd powers* or the *difference* of any *even powers*, of the same degree, of those quantities.

Thus $a+b$ will divide a^3+b^3 , or a^5+b^5 , or a^7+b^7 , &c.,
also $a+b$ will divide a^2-b^2 , or a^4-b^4 , or a^6-b^6 , &c.;

The factors of certain Binomials and Trinomials may also be known from the manner in which the *products* and *squares* of binomials are formed. This will be seen in the following propositions.

(58.) The Product of the *sum* and *difference* of two quantities is equal to the difference of the *squares* of those quantities.

Thus $(a+b)(a-b)$ is equal to a^2-b^2 ; and this last binomial may therefore be divided by either $a+b$ or $a-b$.

This proposition, it will be observed, is included in the two preceding ones, (56.) (57.)

☞ What is the Product of $(a+x)(a-x)$? What is the Product of $(a+5)(a-5)$? Of $(3+y)(3-y)$? Of $(x-1)(x+1)$?

(59.) The Square of the *sum* of two quantities is equal to the sum of the *squares* plus *twice* the product of the two quantities.

Thus $(a+b)(a+b)$, that is, the *square* of $a+b$, is equal to
 a^2+b^2+2ab or $a^2+2ab+b^2$.

This trinomial may therefore be divided by $a+b$.

☞ What is the Product of $(a+x)(a+x)$, or the *square* of $a+x$? What is the Square of $a+y$? Of $x+3$? Of $a+c$? Of $y+5$?

(60.) The Square of the *difference* of two quantities is equal to the sum of the *squares* minus *twice* the product of the two quantities.

Thus $(a-b)(a-b)$, that is, the square of $a-b$, is equal to
 a^2+b^2-2ab or $a^2-2ab+b^2$.

This trinomial may therefore be divided by $a-b$.

☞ What is the Product of $(a-x)(a-x)$, or the square of $a-x$? What is the Square of $a-y$? Of $y-2$? Of $b-x$? Of $1-y$?

The preceding principles will be found applicable to the following

EXERCISES.

1. Resolve $a^2 - x^2$ into component factors.
Ans. $(a-x)(a+x)$
2. Resolve $a^3 + y^3$ into component factors.
Ans. $(a+y)(a^2 - ay + y^2)$
3. Resolve $a^3 - x^3$ into component factors.
Ans. $(a-x)(a^2 + ax + x^2)$
4. Resolve $a^4 - y^4$ into component factors.
Ans. $(a^2 + y^2)(a+y)(a-y)$.
5. Resolve $a^3 - 8x^3$ into component factors.
Ans. $(a-2x)(a^2 + 2ax + 4x^2)$.
6. Resolve $1 - 8y^3$ into component factors.
Ans. $(1-2y)(1+2y+4y^2)$
7. Resolve $1 + 27x^3$ into component factors.
Ans. $(1+3x)(1-3x+9x^2)$.
8. Resolve $8a^3 - 27y^3$ into component factors.
Ans. $(2a-3y)(4a^2 + 6ay + 9y^2)$.
9. Resolve $a^3x^3 + c^3y^3$ into component factors.
Ans. $(ax+cy)(a^2x^2 - axcy + c^2y^2)$.
10. Resolve $a^4 - 16x^4$ into component factors.
Ans. $(a^2 + 4x^2)(a+2x)(a-2x)$.
11. Resolve $a^2 + 2ax + x^2$ into component factors.
Ans. $(a+x)(a+x)$.
12. Resolve $a^2 - 2ay + y^2$ into component factors.
Ans. $(a-y)(a-y)$.
13. Resolve $a^2 + 4ax + 4x^2$ into component factors.
Ans. $(a+2x)(a+2x)$.
14. Resolve $9a^2 - 6ay + y^2$ into component factors.
Ans. $(3a-y)(3a-y)$.
15. Resolve $4a^2 + 12ax + 9x^2$ into component factors.
Ans. $(2a+3x)(2a+3x)$
16. Resolve $a^2x^2 - 2ax + 1$ into component factors.
Ans. $(ax-1)(ax-1)$
17. Resolve $1 + 4xy + 4x^2y^2$ into component factors.
Ans. $(1+2xy)(1+2xy)$.
18. Resolve $4a^2x^2 - 12abxy + 9b^2y^2$ into component factors.
Ans. $(2ax-3by)(2ax-3by)$
19. Resolve $9a^4x^4 + 24a^2cx^2y + 16c^2y^2$ into component factors.
Ans. $(3a^2x^2 + 4cy)(3a^2x^2 + 4cy)$.
20. Resolve $16a^4x^2 - 32a^2cxy^2 + 16c^2y^4$ into component factors.
Ans. $(4a^2x - 4cy^2)(4a^2x - 4cy^2)$.

(61.) A Trinomial may be resolved into two *binomial* factors, whenever one of its three terms is a *square*, another the *sum of the products* of the square root of that term multiplied into any two quantities, and the remaining term the *product of the same two quantities*

For example, let it be required to decompose the trinomial

$$a^2 - a - 12.$$

The first term is the *square* of a ; the second term is the *sum of the products* of a multiplied into 3 and -4 ; and the third term is the product of 3 and -4 .

Now, from the manner in which the product of two binomials is formed, it is evident that

$$a^2 - a - 12 \text{ is equal to } (a+3)(a-4).$$

In like manner the following Trinomials may be decomposed.

21. Resolve $a^2 + 7a + 12$ into component factors.

$$\text{Ans. } (a+3)(a+4).$$

22. Resolve $x^2 + x - 30$ into component factors.

$$\text{Ans. } (x+6)(x-5).$$

23. Resolve $a^2 - 8a - 20$ into component factors.

$$\text{Ans. } (a-10)(a+2).$$

24. Resolve $y^2 - 10y - 39$ into component factors.

$$\text{Ans. } (y-13)(y+3).$$

25. Resolve $a^2x^2 + 9ax + 18$ into component factors.

$$\text{Ans. } (ax+6)(ax+3)$$

26. Resolve $a^2y^2 - 11ay + 28$ into component factors.

$$\text{Ans. } (ay-7)(ay-4)$$

27. Resolve $3ab^2 + 21ab + 36a$ into component factors

The given quantity may be resolved first into

$$3a(b^2 + 7b + 12);$$

and the *trinomial factor* thus obtained may be resolved into $(b+3)$
 $(b+4)$

$$\text{Ans. } 3a(b+3)(b+4).$$

28. Resolve $5x^2 - 5x - 60$ into component factors.

$$\text{Ans. } 5(x-4)(x+3).$$

29. Resolve $2ab^2 + 18ab + 36a$ into component factors.

$$\text{Ans. } 2a(b+6)(b+3).$$

30. Resolve $4ax^3 + 10ax^2 + 6ax$ into component factors.

$$\text{Ans. } ax(2x+3)(2x+2).$$

COMMON MEASURE.

(62.) One quantity is called a *measure* of another, or is said to *measure* another, if it will divide the other, without a remainder; and

A *common measure* of two or more quantities is any quantity that will *divide each of them*, without a remainder.

Thus $5a$ is a common measure of $10ab$ and $5a^2 + 15ab^2$.

☞ Name a Common measure of $4a^2b$ and $6ab^2$. Of $9a^2$ and $3a^2 + a^2c$. Of $5abc + 3a^2c$ and $4ac$. Of a^2x and $a^2x + a^3y$.

, *Greatest Common Measure.*

(63.) The *greatest common measure* of two or more quantities, is the greatest quantity that will divide each of them, without a remainder.

Thus $3a^2$ is the greatest common measure of $6a^2x + 3a^3y$ and $3a^2b - 6a^3cd$, since it is the greatest quantity that will divide each of these binomials, without a remainder.

Here it may be remarked, that when we speak of algebraic quantities as being *greater* or *less*, we have reference to the *coefficients* and *exponents* of the same letters, and not to any particular values which the letters may be supposed to represent.

Thus a^2 is, algebraically, always *greater* than a ; though its *numerical* value would be less than that of a , if the latter should be taken to represent a *proper fraction*, as $\frac{1}{2}$, $\frac{2}{3}$, &c.

☞ What is the Greatest Common Measure of $3a^2x + 4a^3x$ and $a^3y - 5a^2y^3$? Of $5a^3c + 10a^3c^2$ and $10a^2b - 5a^3bx$? Of $2ax^2 - 6a^2y$ and $4a^2x + 8ay^2 - 2a$?

When two quantities have no common measure but *unity*, they are said to be *prime to each other*.

(64.) The greatest common measure of two or more quantities is composed of all the factors which are *common to those quantities*, that is, all the factors which are found in *each quantity*.

For example, the common factors in $3axy^2$ and $6a^2y^2z$ are 3 , a , and y^2 ; and $3ay^2$ is the greatest common measure of the two quantities.

But as no general Rule can be given for resolving a Polynomial into the *polynomial* factors of which it may be composed, we make the Rule for finding the greatest common measure depend on the following proposition ;—

(65.) The greatest common measure of two or more quantities, is the same as that of the *least* of those quantities and the *remainder*, or remainders, if any, after dividing the least into the other, or each of the others.

We may suppose any two quantities to be represented by

$$a \text{ and } na+b;$$

n being some *integral* number, and b less than a .

It is plain that *any measure* of a will also measure na , n times a ; and, measuring na , if it measure $na+b$, it must also measure b . Hence there can be no common measure of a and $na+b$ which is not a common measure of a and b ; the greatest common measure, therefore, of a and b , will be the greatest common measure of a and $na+b$.

But b is the *remainder* after dividing a into $na+b$; hence the proposition is true for two quantities; and in like manner it may be proved for three or more quantities.

R U L E V I I.

(66.) *To find the Greatest Common Measure of two Quantities.*

1. Divide one of the quantities into the other, and the *remainder* into the divisor, and so on, until there is no remainder: the last *divisor* will be the greatest common measure required.—But,

2. When any factor is contained in *each term of the divisor*, without being contained in each term of the *corresponding dividend*,—such factor must be *anceled*, before dividing.—And

3. When the first term of the divisor is not a *measure* of the first term of the dividend—multiply the several terms of the latter by some quantity which will thus render its first term divisible, without a remainder.

NOTE—This Rule might be readily extended to three or more quantities; but we seldom or never have occasion to find the greatest common measure of more than two algebraic quantities.

EXAMPLE.

To find the greatest common measure of the polynomials

$$4a^2x - 5ax^2 + x^3 \text{ and } 3a^3 - 3a^2x + ax^2 - x^3.$$

$$\begin{array}{r}
 4a^2x - 5ax^2 + x^3 \\
 \hline
 4a^2 - 5ax + x^2
 \end{array}
 \left. \begin{array}{l}
 3a^3 - 3a^2x + ax^2 - x^3 \\
 \hline
 4 \\
 \hline
 12a^3 - 12a^2x + 4ax^2 - 4x^3 \quad (3a \\
 12a^3 - 15a^2x + 3ax^2 \\
 \hline
 3a^2x + ax^2 - 4x^3 \\
 \hline
 4 \\
 \hline
 12a^2x + 4ax^2 - 16x^3 \quad (3x \\
 12a^2x - 15ax^2 + 3x^3 \\
 \hline
 19ax^2 - 19x^3
 \end{array} \right)$$

$$\begin{array}{r}
 19ax^2 - 19x^3 \\
 \hline
 a - x
 \end{array}
 \left. \begin{array}{l}
 4a^2 - 5ax + x^2 \quad (4a - x \\
 4a^2 - 4ax \\
 \hline
 -ax + x^2 \\
 -ax + x^2
 \end{array} \right)$$

In this example the factor x is contained in each term of the divisor $4a^2x - 5ax^2 + x^3$, without being contained in each term of the dividend; we therefore cancel this factor, and take $4a^2 - 5ax + x^2$ for a divisor.

Then, the first term $4a^2$ of this divisor not being a *measure* of the first term $3a^3$ of the dividend, we multiply the dividend by 4, which renders the first term $12a^3$ divisible, without a remainder.

We also multiply the *remainder* $3a^2x + ax^2 - 4x^3$, still regarded as a dividend, by 4, to make the first term divisible, without a remainder.

In the next remainder, $19ax^2 - 19x^3$, the exponent of a , the *letter of arrangement*, being less than in the first term of the divisor,—we divide this remainder, after canceling the factor $19x^2$, into the former divisor, which now becomes the dividend.

The divisor $a - x$ completes the operation, and is the *greatest common measure* required.

It remains to elucidate the Rule.

The greatest common measure of the two polynomials, is the same as that of the divisor $4a^2x - 5ax^2 + x^3$ and the *remainder* after the first division, (65). On the same principle the greatest common measure of the first divisor and remainder, is the same as that of the first remainder and the second remainder; and so on, to the last remainder, which becomes the last divisor.

Hence, the last remainder, or divisor—being the greatest common measure of *itself* and the preceding divisor, and so on to the first remainder and divisor—is the greatest common measure of the given polynomials.

Again; since the greatest common measure is composed of all the factors found in *each* of the two polynomials, (64), it is not affected by canceling a factor,—as x in the first divisor,—which is found in only one of them; nor by introducing a *new factor*,—as in multiplying the first dividend by 4,—into only one of them,—since these expedients do not interfere with the original common factors.

The suppression of a factor, as x in this example, which is peculiar to the divisor, is necessary; for, otherwise, the dividend must be multiplied by this factor to render its first term divisible,—and this would introduce a *new common factor* into the two polynomials, and thus increase the greatest common measure, (64).

In the preceding Example, we might have taken the first remainder, $3a^2x + ax^2 - 4x^3$, for a divisor, and the first divisor for the dividend. The operation from that point would be as follows;—

$$\begin{array}{r}
 3a^2x + ax^2 - 4x^3 \quad 4a^2 - 5ax + x^2 \\
 \hline
 3a^2 + ax - 4x^2 \quad \left. \begin{array}{l} 3 \\ \hline 12a^2 - 15ax + 3x^2 \quad (4 \\ 12a^2 + 4ax - 16x^2 \\ \hline -19ax + 19x^2 \end{array} \right) \\
 \\
 -19ax + 19x^2 \\
 \hline
 -a + x \quad \left. \begin{array}{l} 3a^2 + ax - 4x^2 \quad (-3a - 4x \\ 3a^2 - 3ax \\ \hline 4ax - 4x^2 \\ 4ax - 4x^2 \end{array} \right)
 \end{array}$$

The greatest common measure is here found to be $-a+x$, which is the one before determined, with its *signs changed*.

The effect of changing the signs in the divisor, would only be to change the signs in the quotient resulting from the division. Hence

(67.) A Common measure of two or more quantities may have all its *signs changed*, without ceasing to be a common measure of those quantities.

The Rule which has been given, directs that one of the two quantities be divided by the other, &c., without distinguishing the divisor from the dividend. We therefore remark here, that the same common measure will be found by taking *either* of the two quantities for the divisor, and the other for the dividend.

If the divisor and dividend in the preceding Example, be interchanged with each other, the operation will be as follows ;—

$$\begin{array}{r} 4a^2x - 5ax^2 + x^3 \\ 3a \end{array} \Bigg) \begin{array}{r} 12a^3x - 15a^2x^2 + 3ax^3 \\ 12a^3x - 12a^2x^2 + 4ax^3 - 4x^4 \\ \hline -3a^2x^2 - ax^3 + 4x^4 \end{array} \quad (4x$$

$$\frac{-3a^2x^2 - ax^3 + 4x^4}{-3a^2 - ax + 4x^2}$$

$$\begin{array}{r} 3a^3 - 3a^2x + ax^2 - x^3 \\ 3a^3 + a^2x - 4ax^2 \end{array} \Bigg) \begin{array}{r} 3a^3 - 3a^2x + ax^2 - x^3 \\ \hline -4a^2x + 5ax^2 - x^3 \end{array} \quad (-a$$

$$\frac{-4a^2x + 5ax^2 - x^3}{3}$$

$$\begin{array}{r} -12a^2x + 15ax^2 - 3x^3 \\ -12a^2x - 4ax^2 + 16x^3 \end{array} \quad (4x$$

$$\frac{19ax^2 - 19x^3}{9ax^2 - 19x^3}$$

$$\frac{9ax^2 - 19x^3}{a-x}$$

$$a-x$$

$$\begin{array}{r} -3a^2 - ax + 4x^2 \\ -3a^2 + 3ax \end{array} \Bigg) \begin{array}{r} -3a^2 - ax + 4x^2 \\ \hline -4ax + 4x^2 \\ -4ax + 4x^2 \end{array} \quad (-3a+4x$$

$$-4ax + 4x^2$$

$$-4ax + 4x^2$$

In this case the factor x is contained in each term of the dividend, without being contained in each term of the divisor. This factor, therefore, does not enter into the composition of the greatest common measure, (64), and might be canceled before dividing; and this would simplify the operation. Hence

(68.) When any factor is common to all the terms of the *dividend*, and not to those of the divisor, such factor may be suppressed, without affecting the greatest common measure of the two quantities

It follows also from proposition (64), that

(69.) When the same factor is contained in all the terms of two polynomials, their greatest common measure may be found by multiplying this factor into the greatest common measure found for the polynomials *without this factor*.

For example, to find the greatest common measure of

$$a^2x - x^3 \text{ and } a^2x + ax^2 - 2x^3.$$

The factor x is contained in all the terms, and the two polynomials *without this factor* are $a^2 - x^2$ and $a^2 + ax - 2x^2$.

The greatest common measure of these will be found to be $a - x$; then, multiplying x into $a - x$, we find $ax - x^2$ for the greatest common measure of the given polynomials.

EXERCISES.

1. Find the greatest common measure of
 $a^3 - x^3$ and $a^4 - x^4$.

Ans. $a - x$

2. Find the greatest common measure of
 $a^2 - ax - 2x^2$ and $a^2 - 3ax + 2x^2$.

Ans. $a - 2x$

3. Find the greatest common measure of
 $3a^2 - 2a - 1$ and $4a^3 - 2a^2 - 3a + 1$.

Ans. $a - 1$

4. Find the greatest common measure of
 $x^2 + 2ax + a^2$ and $x^3 - a^2x$.

Ans. $x + a$

5. Find the greatest common measure of
 $2x^3 - 16x - 6$ and $3x^3 - 24x - 9$.

Ans. $x^3 - 8x - 3$

6. Find the greatest common measure of
 $a^2 - 5ax + 4x^2$ and $a^3 - a^2x + 3ax^2 - 3x^3$.
Ans. $a - x$.
7. Find the greatest common measure of
 $a^2 - 2ax + x^2$ and $a^3 - a^2x - ax^2 + x^3$.
Ans. $a^2 - 2ax + x^2$.
8. Find the greatest common measure of
 $6a^2 + 7ax - 3x^2$ and $6a^2 + 11ax + 3x^2$.
Ans. $2a + 3x$.
9. Find the greatest common measure of
 $7x^2 - 23xy + 6y^2$ and $5x^3 - 18x^2y + 11xy^2 - 6y^3$.
Ans. $x - 3y$.
10. Find the greatest common measure of
 $a^4 + a^2y^2 + y^4$ and $a^3 - 2a^2y + 2ay^2 - y^3$.
Ans. $a^2 - ay + y^2$.
11. Find the greatest common measure of
 $3a^5 + 6a^4x + 3a^3x^2$ and $a^3x + 3a^2x^2 + 3ax^3 + x^4$.
Ans. $a^2 + 2ax + x^2$.
12. Find the greatest common measure of
 $x^3 - ax^2 - 8a^2x + 6a^3$ and $x^4 - 3ax^3 - 8a^2x^2 + 18a^3x - 8a^4$.
Ans. $x^2 + 2ax - 2a^2$.

COMMON MULTIPLE.

(70.) One quantity is called a *multiple* of another, if it can be measured by the other, that is, divided by it, without a remainder; and

A *common multiple* of two or more quantities is any quantity that can be divided by each of them, without a remainder.

Thus $8a^3x^2$ is a common multiple of $2a^2x$ and $4x^2$.

☞ Name a Common Multiple of $3ab$ and $5c^2$. Of ax^2 and $2ay$. Of $3a^2$ and $4x^2$. Of $5bc$ and x^2 . Of y^3 and $5x^3$.

Least Common Multiple.

(71.) The *least common multiple* of two or more quantities, is the smallest quantity that can be divided by each of them, without a remainder.

Thus $6ax^2$ is the least common multiple of $6a$ and $6x^2$.

☞ What is the Least Common Multiple of $2ab$ and $3b^2$? Of $4x$ and $4y^2$? Of $5a$ and $3a^2b$? Of ab and $2bc^2$? Of $7ax$ and ax^2 ? Of $3a^2$ and $6ay^2$? Of axy and $5x^2y^2$?

(72.) The least common multiple of two or more quantities is composed of the *smallest selection of factors* that includes the factors of each given quantity.

For example, take the quantities $3ab^2c$ and $6a^2bxy$,
Resolving these two quantities into their *prime factors*, we have
 $3abbc$ and $3 \times 2 aabxy$.

If we take $3 \times 2 aabbcxy$, we shall have the smallest selection of factors that includes the factors of each of the two given quantities.

Then the product $6a^2b^2cxy$ is the least common multiple, because it is the smallest quantity that each of the given quantities will divide, without a remainder.

(73.) The least common multiple of *two quantities*, is equal to their *product* divided by their greatest common measure.

For since the greatest common measure of two quantities, is composed of all the factors which are *common to those quantities*, (64), these factors will enter *twice* in the *product* of the quantities.

If, therefore, the product be divided by the greatest common measure, the quotient will contain only those factors which are *common* to the two quantities, and those which are *peculiar to each* of them; and these are the factors of the least common multiple, (72).

RULE VIII.

(74.) *To find the Least Common Multiple of Two or more Quantities.*

1. Set the quantities in a line, from left to right, and divide any *two* or more of them by any *prime* quantity, greater than unity, that will divide them, without a remainder,—placing the quotients and the *undivided* quantities in a line below.

2. Divide any two or more of the quantities in the lower line, as before; and so on, until no two quantities in the lowest line can be divided. The product of the *divisors* and quantities in the *lowest line*, will be the least common multiple required.

3. If no two of the given quantities can be divided as above, the *product of all the quantities* will be their least common multiple.

EXAMPLE.

To find the least common multiple of

$$3ax^2, \quad 6a^2y, \quad \text{and} \quad 3y-4y^3$$

$$\begin{array}{r} 3) \ 3ax^2 \qquad 6a^2y \qquad 3y-4y^3 \\ \hline a) \ ax^2 \qquad 2a^2y \qquad 3y-4y^3 \\ \hline y) \ x^2 \qquad 2ay \qquad 3y-4y^3 \\ \hline \qquad x^2 \qquad 2a \qquad 3-4y^2 \end{array}$$

In the first operation we divide $3ax^2$ and $6a^2y$ by 3, and set down $3y-4y^3$ without dividing it; and in like manner in the second operation. In the third operation we set down x^2 without dividing it.

Then $3ayx^2 \times 2a \times (3-4y^2)$, equal to $18a^2x^2y-24a^2x^2y^3$, is the least common multiple of the three given quantities.

This Rule depends on proposition (72): the divisors and quantities in the lowest line, are the smallest selection of factors that includes the factors of each given quantity.

EXERCISES.

- Find the least common multiple of
 $ax^2, 2a^2y, 4y+y^2$, and ax^2+4x^2 ,
Ans. $8a^3x^2y + 2a^3x^2y^2 + 32a^2x^2y + 8a^2x^2y^2$.
- Find the least common multiple of
 $2ay^2, 4ay^2, 2x-4x^2$, and x^3+ax^2 .
Ans. $4ax^3y^2 - 8ax^4y^2 + 4a^2x^2y^2 - 8a^2x^3y^2$.
- Find the least common multiple of
 $3a^2, ax^2, 3a+6a^2$, and x^3-3x^2 .
Ans. $3a^2x^3 + 6a^3x^3 - 9a^2x^2 - 18a^3x^2$.
- Find the least common multiple of
 $4y^3, 2ay^2, 5a-5ab$, and $10a-5$.
Ans. $40a^2y^3 - 40a^2by^3 - 20ay^3 + 20aby^3$.
- Find the least common multiple of
 $5a, 10ab, 3y+3y^2$, and $6y^3+3y^2$.
Ans. $90aby^3 + 60aby^4 + 30aby^2$.

6. Find the least common multiple of

$$4, 4x^2, 8y-8, \text{ and } 2x^3-ax^3.$$

$$\text{Ans. } 16x^3y-16x^3-8ax^3y+8ax^3$$

7. Find the least common multiple of

$$10, 5ax, 4a-8y^2, \text{ and } 2x+6x^2.$$

$$\text{Ans. } 20a^2x-40axy^2+60a^2x^2-120ax^2y^2$$

8. Find the least common multiple of

$$3y^2, 12y, 5x^2-10, \text{ and } 4y-8y^3.$$

$$\text{Ans. } 60x^2y^2-120y^2-120x^2y^4+240y^4$$

9. Find the least common multiple of

$$14, 7a, 2y^2-4, \text{ and } 28+7ay.$$

$$\text{Ans. } 56ay^2+112a+14a^2y^3-28a^2y.$$

10. Find the least common multiple of

$$a^4-x^4, \text{ and } a^3-a^2x-ax^2+x^3, (73).$$

$$\text{Ans. } a^5-a^4x-ax^4+x^5.$$

11. Find the least common multiple of

$$x^2+2bx+b^2 \text{ and } x^3-b^2x. (73).$$

$$\text{Ans. } x^4+bx^3-b^2x^2-b^3x.$$

12. Find the least common multiple of

$$a^2-3ab+2b^2 \text{ and } a^2-ab-2b^2. (73).$$

$$\text{Ans. } a^3-2a^2b-ab^2+2b^3.$$

13. Find the least common multiple of

$$4a^3-2a^2-3a+1 \text{ and } 3a^2-2a-1. (73).$$

$$\text{Ans. } 12a^4-2a^3-11a^2+1.$$

CHAPTER IV.

FRACTIONS.

(75.) An algebraic FRACTION represents the Quotient of its *numerator* divided by its *denominator*.

Thus $\frac{a}{3x}$ represents the Quotient of a divided by $3x$.

In reading an algebraic Fraction, it will often be necessary to use the terms *numerator* and *denominator*, to avoid ambiguity in reference to the division which is expressed.

Thus if the Fraction

$$\frac{a+x}{y}$$

be read, $a+x$ divided by y , it might be understood that only x is to be divided by y . But the true sense would be conveyed by saying, *numerator* $a+x$, *denominator* y .

A Fraction is thus employed to represent the Quotient, when the divisor is not a factor of the dividend. The quotient in this case may also be represented by means of

Negative Exponents.

(76.) Any quantity with a *negative exponent* is equivalent to a *unit* divided by the same quantity with *the sign of its exponent changed*.

Thus a^{-2} , a with exponent -2 , is equivalent to $\frac{1}{a^2}$.

For $a^2 a^{-2}$ is equal to a^0 , (41);

and by *dividing each of these equals* by a^2 ,

we find a^{-2} equal to $\frac{a^0}{a^2}$ or $\frac{1}{a^2}$, (48).

☞ What is the fractional *equivalent* of a^{-3} ? and how is it proved? Of x^{-4} ? and how is it proved? Of y^{-5} ? and how is it proved? Of $(a+x)^{-2}$? Of $(x-y)^{-3}$?

(77.) When the Divisor is not a factor of the dividend, the Quotient may be represented by a *fraction* (75), or by the dividend *multiplied* into the divisor with the *sign of its exponent changed*.

$a \div x^2$ will give the quotient ax^{-2} , because this quotient multiplied into the *divisor* x^2 , produces ax^0 , (41), which is equal to a , the dividend, (48).

We have, therefore, ax^{-2} for an *integral*, and $\frac{a}{x^2}$ for a *fractional* form of the quotient of $a \div x^2$.

☞ What is the *integral* form of the Quotient of $a \div b^3$? and how is it proved? Of $x \div y^4$? and how is it proved? Of $1 \div a^2$? and how is it proved? Of $a^2 \div x$? Of $b \div ac^2$? Of $a \div 5$?

Transfer of Factors.

(78.) Any factor may be transferred from the *denominator* to the numerator, and *vice versa*, by changing the *sign of its exponent*.

For example, if we divide a by x^2y , fractionally,

$$\text{we have } a \div x^2y \text{ equal to } \frac{a}{x^2y}.$$

If we divide a by the factors x^2 and y , separately, we shall find

$$a \div x^2 \text{ equal to } ax^{-2}, (77), \text{ and } ax^{-2} \div y \text{ equal to } \frac{ax^{-2}}{y}.$$

Hence a divided by x^2y gives

$$\frac{a}{x^2y} \text{ or } \frac{ax^{-2}}{y},$$

These two quotients being necessarily equal to each other, we see that x^2 may be transferred from the denominator to the numerator, by changing the sign of its exponent.

If we also transfer the factor y , we shall have

$$\frac{a}{x^2y} \text{ equal to } \frac{ax^{-2}y^{-1}}{1} \text{ or } ax^{-2}y^{-1}, (77).$$

If we transfer the factor a from the numerator a , or $1a$, to the denominator, we find

$$\frac{a}{x^2y} \text{ equal to } \frac{1}{a^{-1}x^2y}.$$

☞ Under what *different* forms may the Quotient of $a \div bx^2$ be represented? Of $3a \div x^3$? Of $2c \div 3a^4$? Of $ab \div x^2y^2$?

Reciprocals of Quantities.

(79.) The *reciprocal* of a quantity is a *unit divided* by the quantity,

Thus the reciprocal of a^3 is $\frac{1}{a^3}$ or a^{-3} (76).

(80.) The reciprocal of a Fraction is equivalent to the fraction *inverted*; that is, with its numerator and denominator taken *the one for the other*.

Thus the reciprocal of $\frac{a}{x^2}$, or ax^{-2} (77), is $\frac{1}{ax^{-2}}$, which becomes $\frac{x^2}{a}$ by transferring x^{-2} from the denominator to the numerator, (78).

☞ Under what *different forms* may the Reciprocal of $\frac{a^2}{x}$ be expressed? The reciprocal of $\frac{a}{3c^2}$? The reciprocal of $\frac{ab}{y^3}$? Of $\frac{ax}{cy^2}$?

Constant Value of a Fraction.

(81.) The value of a Fraction *remains the same* when its numerator and denominator are both *multiplied*, or both *divided*, by the same quantity.

For example, if we multiply both terms of $\frac{a}{x^2}$ by y ,

we shall have $\frac{a}{x^2}$ equal to $\frac{ay}{x^2y}$.

• For the *first* of these fractions is equal to ax^{-2} ; and the second is equal to $ax^{-2}yy^{-1}$, (77).

But yy^{-1} is equal to y^0 , (41), and by cancelling this factor, (48), we find the *second* fraction also equal to ax^{-2} . Hence the two fractions are equal to each other.

☞ Prove that $\frac{a^2}{b}$ is equal to $\frac{a^2x}{bx}$. That $\frac{c^2}{x^3}$ is equal to $\frac{c^2m^2}{x^3m^2}$.

Signs of Fractions.

(82.) Since a Fraction represents the *quotient* of its numerator divided by its denominator, a Fraction is *positive* when its numerator and denominator have the same sign, and *negative* when they have contrary signs, (49).

☞ Say whether the Fraction $\frac{a}{b}$ is *positive* or *negative*. Say whether $\frac{a}{-b}$ is *positive* or *negative*. Say whether $\frac{-a}{-b}$ is *positive* or *negative*. Say whether $\frac{-ax}{by}$ is *positive* or *negative*.

(83.) The sign $+$ or $-$ *prefixed to a Fraction*,—not the sign of either the numerator or denominator,—shows whether the fraction enters *additively* or *subtractively* into a calculation.

Thus $+\frac{-a}{b}$ denotes that the fraction $\frac{-a}{b}$, which is in itself *negative*, (82), is to be *added*, in the calculation into which it enters.

When no sign is prefixed to a fraction, $+$ is always understood.

(84.) The signs of both the numerator and denominator *may be changed*, or the sign of either of them with the *sign prefixed to the Fraction*,—without affecting the value of the fraction.

Thus $\frac{a}{b}$ is equivalent to $\frac{-a}{-b}$, since both these fractions are *positive*, (82).

Also $\frac{a}{b}$ is equivalent to $-\frac{-a}{b}$. For the fraction $\frac{-a}{b}$ is *negative*, its numerator and denominator having contrary signs; this *negative* fraction becomes *positive*, when subtracted, as required by the sign $-$ *prefixed to it*, and it is then equivalent to the first fraction.

A Polynomial is changed from *positive* to *negative*, or from *negative* to *positive*, by changing the sign of *each of its terms*, (38).

For example, if $a-b$ is *positive*, a must be *greater* than b ; then, changing the signs, $b-a$ will be *negative*.

The Fraction $\frac{a-b}{x-y}$ is therefore equivalent to $\frac{b-a}{y-x}$.

☞ What other changes may be made in the *signs*, without affecting the value, of this fraction?

FRACTIONS REDUCED TO THEIR LOWEST TERMS.

(85.) A Fraction is reduced to *lower terms* by dividing its numerator and denominator by the same common *measure*. This simplifies the fraction, without altering its value, (81).

A *monomial* common measure may usually be known by inspection. Thus to reduce the Fraction

$$\frac{4a^2b}{6ac-8a^2x}$$

It is obvious that we have only to divide its numerator and denominator by $2a$. This gives us the equivalent Fraction,

$$\frac{2ab}{3c-4ax}$$

A *binomial* common measure may often be discovered from the principles which have been established for the decomposition of Polynomials.

Thus to reduce the Fraction

$$\frac{a^2-b^2}{a^2+2ab+b^2}$$

By proposition (57) we can divide the numerator by $a+b$; and by (59), we can divide the denominator by the same quantity

Thus dividing we find the equivalent Fraction

$$\frac{a-b}{a+b}$$

In all cases in which a Fraction admits of being reduced, we may apply

RULE IX.

(86.) *To Reduce a Fraction to its Lowest Terms.*

Divide the numerator and denominator by their greatest common measure, (66): the quotients will be the lowest terms of the given fraction.

EXERCISES.

1. Reduce $\frac{3ax^2}{3a^2b-9a^3}$ to its lowest terms.

$$\text{Ans. } \frac{x^2}{ab-3a^2}$$

2. Reduce $\frac{2a^2y+4ay^2}{8a}$ to its lowest terms.

$$\text{Ans. } \frac{ay+2y^2}{4}$$

3 Reduce $\frac{a^2 - b^2}{a^3 + b^3}$ to its lowest terms. (57).

$$\text{Ans. } \frac{a - b}{a^2 - ab + b^2}.$$

4. Reduce $\frac{a^4 - x^4}{a^5 - a^3x^2}$ to its lowest terms.

$$\text{Ans. } \frac{a^2 + x^2}{a^3}$$

5. Reduce $\frac{x^2 - a^2}{x^2 + 2ax + a^2}$ to its lowest terms.

$$\text{Ans. } \frac{x - a}{x + a}.$$

6. Reduce $\frac{a^3 - ay^2}{a^2 + 2ay + y^2}$ to its lowest terms.

$$\text{Ans. } \frac{a^2 - ay}{a + y}.$$

7. Reduce $\frac{a^2 - x^2}{a^3 - 3a^2x + 3ax^2 - x^3}$ to its lowest terms.

$$\text{Ans. } \frac{a + x}{a^2 - 2ax + x^2}.$$

8. Reduce $\frac{2ax^2 - a^2x - a^3}{2x^2 + 3ax + a^2}$ to its lowest terms.

$$\text{Ans. } \frac{ax - a^2}{x + a}$$

9. Reduce $\frac{a^3 + 2a^2x + 3a^2x^2}{2a^4 - 3a^3x - 5a^2x^2}$ to its lowest terms.

$$\text{Ans. } \frac{a + 2x + 3x^2}{2a^2 - 3ax - 5x^2}.$$

10. Reduce $\frac{6a^2 + 7ax - 3x^2}{6a^2 + 11ax + 3x^2}$ to its lowest terms.

$$\text{Ans. } \frac{3a - x}{3a + x}.$$

11. Reduce $\frac{a^4 + a^2x^2 + x^4}{a^4 + a^3x - ax^3 - x^4}$ to its lowest terms.

$$\text{Ans. } \frac{a^2 - ax + x^2}{a^2 - x^2}.$$

12. Reduce $\frac{5a^5 + 10a^4y + 5a^3y^2}{a^3y + 2a^2y^2 + 2ay^3 + y^4}$ to its lowest terms.

$$\text{Ans. } \frac{5a^4 + 5a^3y}{a^2y + ay^2 + y^3}.$$

FRACTIONS REDUCED TO A COMMON DENOMINATOR.

(87.) Two or more Fractions are said to have a *common denominator*, when they have the same quantity for a denominator.

Thus $\frac{ax}{a+b}$ and $\frac{a-x}{a+b}$ have a common denominator.

Two or more Fractions may often be reduced, very readily, to a common denominator, by multiplying both the numerator and denominator of one or more of them, so as to make the denominator the *same for each*.

For example, to reduce to a common denominator the Fractions

$$\frac{a}{2a-2x} \text{ and } \frac{bc}{3a-3x},$$

We have only to multiply the terms of the first fraction by 3, and those of the second by 2. This gives the equivalent Fractions

$$\frac{3a}{6a-6x} \text{ and } \frac{2bc}{6a-6x}. \quad (81).$$

Observe that this reduction does not alter the values of the given Fractions.

When this method cannot be obviously applied, we adopt

R U L E X.

(88.) *To Reduce two or more Fractions to a Common Denominator.*

1. Multiply each numerator by all the denominators *except its own*, for new *numerators*; and multiply all the denominators together, for a *common denominator*.

2. *If the Least Common Denominator* be required,—Find the least common *multiple* of the given denominators, for the Common denominator. Divide this Multiple by the denominator of each given Fraction, and multiply the quotient by the numerator, for the new numerators.

E X A M P L E S.

1. To reduce $\frac{a}{3x}$, $\frac{b}{6}$, and $\frac{c}{y+2}$ to a common denominator.

For the new numerators, we have

$$\begin{array}{l} a \cdot 6 \cdot (y+2), \text{ equal to } 6ay+12a; \\ b \cdot 3x \cdot (y+2), \quad \text{“} \quad 3bxy+6bx; \\ \text{and } c \cdot 3x \cdot 6 \quad \text{“} \quad 18cx. \end{array}$$

And the common denominator is

$$3x \cdot 6 \cdot (y+2), \text{ equal to } 18xy+36x.$$

The given Fractions are thus reduced to

$$\frac{6ay+12a}{18xy+36x}, \frac{3bxy+6bx}{18xy+36x}, \frac{18cx}{18xy+36x}, \text{ respectively.}$$

2. To reduce the same Fractions to the *least common denominator*.

The least common *multiple* of the denominators $3x$, 6 , and $y+2$ will be found to be $6xy+12x$, (74), which is the required *denominator*.

Dividing this Multiple by each given denominator, and multiplying the quotients by the given numerators, respectively, we find the new numerators,

$$2ay+4a, \quad bxy+2bx, \quad \text{and} \quad 6cx.$$

The Fractions reduced to their least common denominator, are then

$$\frac{2ay+4a}{6xy+12x}, \frac{bxy+2bx}{6xy+12x}, \frac{6cx}{6xy+12x}, \text{ respectively.}$$

Each of the given Fractions should be in its *lowest terms*, before proceeding to find their least common denominator; otherwise, the denominator found will not, in all cases, be the smallest by means of which the values of the several Fractions may be expressed.

In finding a Common Denominator as above, the numerator and denominator of each given fraction are multiplied by the *same quantity*.

Thus in the first example, $\frac{a}{3x}$ has both its terms multiplied by 6 and $y+2$,—producing the new terms $6ay+12a$ and $18xy+36x$.

Hence the values of the given fractions are not altered in reducing them to a common denominator, (81).

EXERCISES.

1. Reduce $\frac{a}{3}$, $\frac{b}{2x}$, and $\frac{c}{x-4}$ to a common denominator.

$$\text{Ans. } \frac{2ax^2-8ax}{6x^2-24x}, \frac{3bx-12b}{6x^2-24x}, \frac{6cx}{6x^2-24x}$$

2. Reduce $\frac{2}{a^2}$, $\frac{a}{y^2}$, and $\frac{a+c}{1-y}$ to a common denominator.

$$\text{Ans. } \frac{2y^2-2y^3}{a^2y^2-a^2y^3}, \frac{a^3-a^3y}{a^2y^2-a^2y^3}, \frac{a^3y^2+a^2cy^2}{a^2y^2-a^2y^3}$$

3. Reduce $\frac{a^2}{x}$, $\frac{c}{3y}$, and $\frac{a+5}{y^2}$ to a common denominator.

$$\text{Ans. } \frac{3a^2y^3}{3xy^3}, \frac{cxy^2}{3xy^3}, \frac{3axy+15xy}{3xy^3}.$$

4. Reduce $\frac{ax}{2}$, $\frac{a}{3x}$, and $\frac{b-1}{1+b}$ to a common denominator.

$$\text{Ans. } \frac{3ax^2+3abx^2}{6x+6bx}, \frac{2a+2ab}{6x+6bx}, \frac{6bx-6x}{6x+6bx}$$

5. Reduce $\frac{a^3}{y}$, $\frac{b}{2x^2}$, and $\frac{a+b}{xy}$ to a common denominator.

$$\text{Ans. } \frac{2a^3x^3y}{2x^3y^2}, \frac{bxy^2}{2x^3y^2}, \frac{2ax^2y+2bx^2y}{2x^3y^2}$$

6. Reduce $\frac{a}{2x^2}$, $\frac{b}{y}$, and $\frac{ab}{x^2+x^3}$ to the least common denominator.

$$\text{Ans. } \frac{ay+axy}{2x^2y+2x^3y}, \frac{2bx^2+2bx^3}{2x^2y+2x^3y}, \frac{2aby}{2x^2y+2x^3y}.$$

7. Reduce $\frac{x^2}{4}$, $\frac{a}{3y}$, and $\frac{a-x}{6y^2}$ to the least common denominator.

$$\text{Ans. } \frac{3x^2y^2}{12y^2}, \frac{4ay}{12y^2}, \frac{2a-2x}{12y^2}.$$

8. Reduce $\frac{a}{y^2}$, $\frac{b}{5}$, and $\frac{c}{y^3-y^2}$ to the least common denominator.

$$\text{Ans. } \frac{5ay-5a}{5y^3-5y^2}, \frac{by^3-by^2}{5y^3-5y^2}, \frac{5c}{5y^3-5y^2}.$$

9. Reduce $\frac{y}{4}$, $\frac{a}{2x^2}$, and $\frac{a^2}{2+x}$ to the least common denominator.

$$\text{Ans. } \frac{2x^2y+x^3y}{8x^2+4x^3}, \frac{4a+2ax}{8x^2+4x^3}, \frac{4a^2x^2}{8x^2+4x^3}.$$

10. Reduce $\frac{x-1}{3}$, $\frac{a^2}{3y}$, and $\frac{c}{6y-3y^2}$ to the least common denominator.

$$\text{Ans. } \frac{2xy-xy^2-2y+y^2}{6y-3y^2}, \frac{2a^2-a^2y}{6y-3y^2}, \frac{c}{6y-3y^2}$$

INTEGRAL AND MIXED QUANTITIES REDUCED TO IMPROPER FRACTIONS.

(89.) An *integral* quantity is one which does not contain any *fractional* expression; as $3ax^2$, or $2ab - 5xy$.

(90.) A *mixed* quantity is partly *integral* and partly *fractional*; as $3ax^2 + \frac{a}{5c}$, or $2ab - \frac{x}{1+y}$.

(91.) An *improper* Fraction is a fraction whose value may be expressed by an integral or a mixed quantity.

Thus $\frac{6a^2+5}{3a}$ is an improper fraction whose value is $2a + \frac{5}{3a}$.

RULE XI.

(92.) *To Reduce an Integral or a Mixed Quantity to an Improper Fraction.*

1. Under an *integral* quantity, regarded as a numerator, set 1 for a *denominator*. Or multiply the integral quantity by any proposed denominator; the product will be the numerator.

2. In a *mixed* quantity, multiply the integral part by the denominator annexed; *add the numerator* to the product when the sign before the fractional part is +, but *subtract the numerator* when this sign is -; and place the result over said denominator.

EXAMPLES.

1. To reduce $3a^2$ to a Fraction whose denominator shall be $a - 2x$.
 $3a^2$ is the same as $\frac{3a^2}{1}$; and by multiplying both terms of this fraction by the proposed denominator, we have

$$3a^2 \text{ equal to } \frac{3a^3 - 6a^2x}{a - 2x}, \quad (81).$$

2. To reduce $3a^2 - \frac{a - x^2}{2}$ to an improper Fraction

Multiplying $3a^2$ by the denominator 2, and *subtracting* the numerator $a - x^2$ from the product,—observing to *change the signs* of the numerator, (36),—we have

$$3a^2 - \frac{a - x^2}{2} \text{ equal to } \frac{6a^2 - a + x^2}{2}.$$

The reason of this operation will be evident, if we consider that, by multiplying the $3a^2$ by 2, we reduce the integral part to a *common denominator* with the fraction annexed to it, according to the first part of the Rule. The operation then consists in subtracting numerator from numerator, and placing their difference over the common denominator.

Another view to be taken of the preceding operations, is, that the integral term $3a^2$ is *multiplied*, and the product is then taken to be *divided*, by the same quantity, namely, the denominator. The value therefore remains the same.

EXERCISES.

1. Reduce $4ax^2$ to a Fraction whose denominator shall be $y+2$.

$$\text{Ans. } \frac{4ax^2y+8ax^2}{y+2}.$$

2. Reduce a^2+3x to a Fraction whose denominator shall be $2y^2$.

$$\text{Ans. } \frac{2a^2y^2+6xy^2}{2y^2}.$$

3. Reduce $5xy$ to a Fraction whose denominator shall be $3+x^2$.

$$\text{Ans. } \frac{15xy+5x^3y}{3+x^2}.$$

4. Reduce y^2-5 to a Fraction whose denominator shall be $3ax^2$.

$$\text{Ans. } \frac{3ax^2y^2-15ax^2}{3ax^2}.$$

5. Reduce $a+b$ to a Fraction whose denominator shall be $a-b$.

$$\text{Ans. } \frac{a^2-b^2}{a-b}.$$

6. Reduce $5ay^2$ to a Fraction whose denominator shall be a^2-b^2 .

$$\text{Ans. } \frac{5a^3y^2-5ab^2y^2}{a^2-b^2}.$$

7. Reduce x^2+1 to a Fraction whose denominator shall be $3a^2y$.

$$\text{Ans. } \frac{3a^2x^2y+3a^2}{3a^2y}.$$

8. Reduce ab^2-x to a Fraction whose denominator shall be $1-y$.

$$\text{Ans. } \frac{ab^2-x-ab^2y+xy}{1-y}.$$

9. Reduce $a^2 + x + \frac{a}{y}$ to an improper Fraction.

$$\text{Ans. } \frac{a^2y + xy + a}{y}.$$

10. Reduce $ax^2 + \frac{b}{y+1}$ to an improper Fraction.

$$\text{Ans. } \frac{ax^2y + ax^2 + b}{y+1}$$

11. Reduce $5a + 3b - \frac{a+x}{3}$ to an improper Fraction.

$$\text{Ans. } \frac{14a + 9b - x}{3}$$

12. Reduce $2a - 4c + \frac{a-y^2}{5}$ to an improper Fraction.

$$\text{Ans. } \frac{11a - 20c - y^2}{5}$$

13. Reduce $2 + y^2 - \frac{1-y^2}{4}$ to an improper Fraction.

$$\text{Ans. } \frac{7 + 5y^2}{4}.$$

14. Reduce $a^2 + x^2 - \frac{a^3 - x^3}{a-x}$ to an improper Fraction.

$$\text{Ans. } \frac{ax^2 - a^2x}{a-x}.$$

15. Reduce $a - x + \frac{a^2 - ax}{x}$ to an improper Fraction.

$$\text{Ans. } \frac{a^2 - x^2}{x}.$$

16. Reduce $1 + 2x - \frac{4x-4}{5x}$ to an improper Fraction.

$$\text{Ans. } \frac{x + 10x^2 - 4}{5x}.$$

17. Reduce $2x + 2y + \frac{5y^2}{2x-3y}$ to an improper Fraction.

$$\text{Ans. } \frac{4x^2 - 2xy - y^2}{2x-3y}.$$

18. Reduce $a + x - \frac{a^3 - x^3}{a^2 - ax + x^2}$ to an improper Fraction.

$$\text{Ans. } \frac{2x^3}{a^2 - ax + x^2}.$$

IMPROPER FRACTIONS REDUCED TO INTEGRAL OR MIXED
QUANTITIES.

By reversing the preceding Rule, we have

RULE XII.

(93.) *To Reduce an Improper Fraction to an Integral or a Mixed Quantity.*

Divide the numerator by the denominator for the *integral* part, and set the denominator under the remainder, if any, for the *fractional* part, of the result. Connect the fractional to the integral part by the sign +; or *change the sign* of the numerator or denominator, and connect it by the sign -.

EXAMPLE.

To reduce to an integral or a mixed quantity the Fraction

$$\frac{9a^2c + 3ab - 2b + y}{3a}$$

Dividing the numerator by the denominator, we find the integral quotient to be $3ac + b$, and the *remainder* $-2b + y$.

Setting the denominator under this remainder, and connecting the fraction so formed to the integral quotient, by the sign +, the result is

$$3ac + b + \frac{-2b + y}{3a}.$$

Or, changing the signs in the numerator $-2b + y$, and connecting the fraction by the sign -, the result, under a somewhat simpler form,

$$\text{is } 3ac + b - \frac{2b - y}{3a}.$$

This form is simpler than the preceding, as it dispenses with one sign in the numerator of the fractional part.

The reason of the preceding operation is evident from the consideration that every Fraction is equal to its numerator divided by its denominator, (75). After obtaining the quotient $3ac + b$,—the divisor $3a$ not being contained in the remainder $-2b + y$, the division of these terms is indicated by setting the divisor under them.

The fractional part of the result must evidently be *added* to the integral part; and this addition is indicated by placing the sign + before the fraction. But the value of the fraction annexed, will not be affected by changing the sign + before it to -, if at the same time we change the signs in the numerator, (84).

The Fraction formed of the divisor and remainder, will be in its *lowest terms*, or not, according as the improper fraction reduced, is, or is not, in its lowest terms.

For, if the dividend and divisor have any common *measure*, the divisor and remainder will have the same common measure, (65).

EXERCISES.

1. Reduce $\frac{a^3-x^3}{a-x}$ to an integral or a mixed quantity.

Ans. a^2+ax+x^2 .

2. Reduce $\frac{10x^2-5x+3}{5x}$ to an integral or a mixed quantity.

Ans. $2x-1+\frac{3}{5x}$.

3. Reduce $\frac{a^4-y^4}{a-y}$ to an integral or a mixed quantity.

Ans. $a^3+a^2y+ay^2+y^3$.

4. Reduce $\frac{x^4-3x^2y^2+ax}{x^2-3y^2}$ to an integral or a mixed quantity.

Ans. $x^2+\frac{ax}{x^2-3y^2}$.

5. Reduce $\frac{4a^2x^2-3ay-2b}{2a^2}$ to an integral or a mixed quantity.

Ans. $2x^2-\frac{3ay+2b}{2a^2}$.

6. Reduce $\frac{9ax^2-2x+3}{3a}$ to an integral or a mixed quantity.

Ans. $3x^2-\frac{2x-3}{3a}$.

7. Reduce $\frac{a^3+y^3}{a+y}$ to an integral or a mixed quantity.

Ans. a^2-ay+y^2 .

8. Reduce $\frac{x^2-y^2+4}{x+y}$ to an integral or a mixed quantity.

Ans. $x-y+\frac{4}{x+y}$

9. Reduce $\frac{4ay-2y^2+a^3}{2a-y}$ to an integral or a mixed quantity.

Ans. $2y+\frac{a^3}{2a-y}$.

ADDITION OF FRACTIONS.

(94.) The Sum of two or more Fractions is found by means of a *common denominator*.

$$\text{Thus the Sum of } \frac{a}{x} \text{ and } \frac{b}{x} \text{ is } \frac{a+b}{x}.$$

For it is evident that a divided by x , added to b divided by x , makes the sum of a and b divided by x .

In other words, if each of the *parts* a and b be divided by x , the *whole* $a+b$ will be divided by x .

Hence we have

RULE XIII.

(95.) *For the Addition of Fractions.*

1. If the fractions have not a *common denominator*, reduce them to a common denominator.

2. Add the numerators together, and place the Sum, as a numerator, over their common denominator.

3. *Mixed quantities* may be added under the form of *improper fractions*; or the integral and the fractional parts may be added separately.

EXAMPLE.

To add together $2a + \frac{b}{x}$ and $3a - \frac{c}{y}$.

Reducing these mixed quantities to improper fractions, they become

$$\frac{2ax+b}{x} \text{ and } \frac{3ay-c}{y}.$$

Reducing these fractions to a common denominator, we have

$$\frac{2axy+by}{xy} \text{ and } \frac{3axy-cx}{xy}.$$

Placing the sum of these numerators over the common denominator, the result is,

$$\frac{5axy+by-cx}{xy}, \text{ equal to } 5a + \frac{by-cx}{xy}.$$

Otherwise, by adding the *integral* and the *fractional* parts separately.—There will be less liability to error, if we change the *sign* before the fraction in the second quantity to +, and change the sign of its numerator; thus

$$+ \frac{-c}{y}, (84).$$

Then, reducing $\frac{b}{x}$ and $\frac{-c}{y}$ to a common denominator, they become

$$\frac{by}{xy} \text{ and } \frac{-cx}{xy}.$$

Adding these fractions together, and adding together the integral parts $2a$ and $3a$, we obtain

$$5a + \frac{by - cx}{xy}, \text{ as before.}$$

Improper fractions in the results obtained by this Rule, should be reduced to integral or mixed quantities; and proper fractions, to their lowest terms.

EXERCISES.

1. Add together $\frac{3a}{4}$, $\frac{2a}{3}$, and $\frac{3a+2}{3}$.

$$\text{Ans. } 2a + \frac{5a+8}{12}.$$

2. Add together $2x + \frac{a^2}{4}$, and $4x + \frac{2a^2}{5}$.

$$\text{Ans. } 6x + \frac{13a^2}{20}.$$

3. Add together $y^2 + \frac{2x}{3}$, and $3y^2 - \frac{x}{4}$.

$$\text{Ans. } 4y^2 + \frac{5x}{12}.$$

4. Add together $\frac{a^2}{2x}$, $2a^2$, and $\frac{2b-3}{5}$.

$$\text{Ans. } 2a^2 + \frac{5a^2 + 4bx - 6x}{10x}.$$

5. Add together $3x^2 + \frac{a+1}{2}$ and $\frac{2a+3}{4}$.

$$\text{Ans. } 3x^2 + a + \frac{5}{4}.$$

6. Add together x^2y , $\frac{a^2}{3}$, and $x^2y + \frac{1}{2}$.

$$\text{Ans. } 2x^2y + \frac{2a^2+3}{6}.$$

7 Add together $2a^2 - \frac{x}{4}$ and $\frac{a^2-3x}{2}$.

$$\text{Ans. } 2a^2 + \frac{2a^2-7x}{4}.$$

8. Add together $\frac{a^2}{y}$, $\frac{3a}{y}$, and $\frac{a^2-4}{y^2}$.

$$\text{Ans } \frac{a^2y+3ay+a^2-4}{y^2}.$$

9 Add together $\frac{x-1}{3}$, 2, and $\frac{3x+4}{5}$.

$$\text{Ans. } 2 + \frac{14x+7}{15}$$

10. Add together $2y + \frac{2a^3}{3}$ and $y - \frac{3a^3}{4}$.

$$\text{Ans. } 3y - \frac{a^3}{12}.$$

11. Add together $5a^2$, $\frac{3x}{c}$, and $\frac{a^2+1}{c^2}$.

$$\text{Ans. } 5a^2 + \frac{3cx+a^2+1}{c^2}.$$

12. Add together 10, $\frac{-x^2}{3}$, and $3 - \frac{x^2}{5}$.

$$\text{Ans. } 13 - \frac{8x^2}{15}.$$

13. Add together $\frac{a^2+x}{3}$ and $\frac{a^2-x+1}{4}$.

$$\text{Ans. } \frac{7a^2+x+3}{12}.$$

14. Add together $2y^2 - \frac{3}{a}$ and $\frac{3+y^2-x^2}{5}$.

$$\text{Ans. } 2y^2 - \frac{15-3a-ay^2+ax^2}{5a}$$

SUBTRACTION OF FRACTIONS.

(96.) The Difference of two Fractions is found by means of a *common denominator*.

Thus $\frac{a}{x}$ subtracted from $\frac{b}{x}$ leaves $\frac{b-a}{x}$.

For the last fraction, $\frac{b-a}{x}$, added to $\frac{a}{x}$ the one subtracted, produces $\frac{b}{x}$ the one from which the subtraction is made.

We have therefore

RULE XIV.

(97.) *For the subtraction of Fractions.*

1. If the fractions have not a *common denominator*, reduce them to a common denominator.

2. Subtract the numerator of the fraction to be subtracted from the other numerator, and place the Difference, as a numerator, over the common denominator.

3. A *mixed quantity* may be taken in subtraction under the form of an *improper fraction*; or the integral and the fractional part may be taken separately in subtracting.

EXAMPLE.

From $8x + \frac{a}{y}$ to subtract $3x - \frac{b}{w}$.

Reducing these mixed quantities to improper fractions, they become

$$\frac{8xy+a}{y} \text{ and } \frac{3xw-b}{w}.$$

Reducing these fractions to a common denominator,—subtracting the second of the resulting numerators from the first,—and placing the difference over the common denominator, we find

$$\frac{5xyw+aw+by}{yw}, \text{ equal to } 5x + \frac{aw+by}{yw}.$$

Otherwise, by taking the integral and the fractional parts separately.—To diminish the liability to error in adjusting the signs, we change the fraction in the second quantity to $+\frac{-b}{w}$, (84).

Then, reducing $\frac{a}{y}$ and $\frac{-b}{w}$ to a common denominator, they become

$$\frac{aw}{yw} \text{ and } \frac{-by}{yw}.$$

Subtracting the second of these fractions from the first, and $3x$ from $5x$, we find the difference of the given quantities to be

$$5x + \frac{aw + by}{yw}, \text{ as before.}$$

In all subsequent exercises, improper fractions in the results should be reduced to integral or mixed quantities; and proper fractions, to their lowest terms.

EXERCISES.

1. From $\frac{a^2 + 2}{3}$ subtract $\frac{b - 3}{2a}$.

$$\text{Ans. } \frac{2a^3 + 4a - 3b + 9}{6a}.$$

2. From $\frac{3a + x}{y}$ subtract $\frac{a - x}{y^2}$.

$$\text{Ans. } \frac{3ay + xy - a + x}{y^2}.$$

3. From $\frac{a^2}{1 + x}$ subtract $\frac{b}{1 - x}$.

$$\text{Ans. } \frac{a^2 - a^2x - b - bx}{1 - x^2}.$$

4. From $3a + \frac{b}{x}$ subtract $2a - \frac{c}{y}$.

$$\text{Ans. } a + \frac{by + cx}{xy}.$$

5. From $2x^2 - \frac{a}{x}$ subtract $x^2 - \frac{b}{y}$.

$$\text{Ans. } x^2 - \frac{ay - bx}{xy}.$$

6. From $\frac{1}{x-y}$ subtract $\frac{1}{x^2-y^2}$.

Ans. $\frac{x+y-1}{x^2-y^2}$.

7. From $3y^2 - \frac{a}{y}$ subtract $2y^2 + \frac{b}{x}$.

Ans. $y^2 - \frac{ax+by}{xy}$.

8. From $2a+b + \frac{x}{2}$ subtract $3b - \frac{y}{2}$.

Ans. $2a-2b + \frac{x+y}{2}$.

9. From $4x^2 - y - \frac{a^3}{3}$ subtract $y + \frac{c^3}{2}$.

Ans. $4x^2 - 2y - \frac{2a^3+3c^3}{6}$.

10. From $2b^2 + \frac{a+1}{2}$ subtract $b^2 - \frac{a-2}{3}$.

Ans. $b^2 + \frac{5a-1}{6}$.

11. From $5a^3 + \frac{x}{y+1}$ subtract $3a^3 + \frac{x}{y-1}$.

Ans. $2a^3 - \frac{2x}{y^2-1}$.

12. From $3x^2 - \frac{a+b}{2}$ subtract $-x^2 + \frac{a^2}{3}$.

Ans. $4x^2 - \frac{3a+3b+2a^2}{6}$.

13. From $2b^3 - \frac{1+a}{3}$ subtract $b^2 - \frac{1-a}{4}$.

Ans. $2b^3 - b^2 - \frac{1+7a}{12}$.

14. From $5 + \frac{a^2+x^2}{y}$ subtract $1 + \frac{a^2-x^2}{y^2}$.

Ans. $4 + \frac{a^2y+x^2y-a^2+x^2}{y^2}$.

MULTIPLICATION OF FRACTIONS.

(98.) The Product of two or more Fractions is equal to the product of their numerators divided by the *product of their denominators*

$$\text{Thus } \frac{a}{x^2} \text{ multiplied by } \frac{b}{y^3} \text{ produces } \frac{ab}{x^2y^3}.$$

For the first of these fractions is equivalent to ax^{-2} , and the second to by^{-3} , (77); and the product of these *two equivalents* is

$$abx^{-2}y^{-3}, \text{ (40).}$$

And by transferring the factors x^{-2} and y^{-3} to the *denominator*, we have

$$\frac{ab}{x^2y^3}, \text{ } ab \text{ divided by } x^2y^3, \text{ (78).}$$

From the preceding it follows, that

(99.) Multiplying by a Fraction finds *such a part of the multiplicand* as is expressed by the multiplier.

$$\text{Thus } a \times \frac{1}{2} \text{ or } \frac{a}{1} \times \frac{1}{2} \text{ produces } \frac{a}{2}, \frac{1}{2} \text{ of } a.$$

Compound Fractions.

(100.) A Fraction multiplied by a *fraction*, or divided by an *integer*, may be expressed by a *Compound fraction*, that is, a fraction of a fraction.

$$\text{For example, } \frac{a}{x} \times \frac{1}{2} \text{ is } \frac{1}{2} \text{ of } \frac{a}{x}, \text{ (99); also } \frac{a}{x} \div 2 \text{ is } \frac{1}{2} \text{ of } \frac{a}{x}.$$

Hence

(101.) Multiplying two or more Fractions together is equivalent to reducing a *compound* to a *simple* fraction.

$$\text{Thus } \frac{a}{x} \times \frac{1}{2} \text{ is } \frac{1}{2} \text{ of } \frac{a}{x}, \text{ equal to } \frac{a}{2x}.$$

From proposition (98), we have the following Rule.

RULE XV

(102.) *For the Multiplication of Fractions*

1. Multiply the numerators together for a *numerator*, and the denominators together for a *denominator*.

2. An *integral* quantity and a *fraction* are multiplied together, by multiplying the *numerator*, or dividing the denominator, by the integer.

3. A *mixed* quantity may be taken in multiplication under the form of an *improper fraction*; or the integral and the fractional part may be taken separately in multiplying.

It may also be remarked that

(103.) A Fraction is multiplied by *its own denominator*, by merely canceling the denominator.—And *equal factors may be canceled* in a numerator and its own or the other *denominator*, without altering the Product of the two fractions, (81).

EXAMPLE.

To multiply $\frac{a^2-x^2}{3y^2}$ by $\frac{2y}{a^2+2ax+x^2}$.

The first numerator is equal to $(a+x)(a-x)$, (58),

and the second denominator, to $(a+x)(a+x)$, (59),

By canceling the factor $a+x$ from these terms, and the factor y from the other numerator and denominator, the operation is reduced to

$$\frac{a-x}{3y} \times \frac{2}{a+x}; \text{ which produces } \frac{2a-2x}{3ay+3xy}.$$

EXERCISES.

1. Multiply together a^2-x^2 and $\frac{3}{a-x}$. Ans. $3a+3x$.

2. Multiply together a^2y and $x+\frac{3}{4}$. Ans. $a^2xy + \frac{3a^2y}{4}$.

3. Multiply together $\frac{a+x}{3}$ and $\frac{2}{a-x}$. Ans. $\frac{2a+2x}{3a-3x}$.

4. Multiply together $5y^2$ and $y^2-\frac{a}{3}$. Ans. $5y^4 - \frac{5ay^2}{3}$.

- 5 Multiply together $\frac{a+b}{3x^2}$ and $\frac{a-b}{a+b}$. Ans. $\frac{a-b}{3x^2}$.
- 6 Multiply together $2a^2-3x$ and $\frac{3b}{y-1}$. Ans. $\frac{6a^2b-9bx}{y-1}$.
7. Multiply together $a + \frac{ax}{a-x}$ and $x - \frac{ax}{a+x}$. Ans. $\frac{a^2x^2}{a^2-x^2}$.
8. Multiply together $\frac{a^4-b^4}{a+b}$ and $\frac{a^2}{ab-b^2}$. Ans. $\frac{a^4+a^2b^2}{b}$.
9. Multiply together $5y^2 - \frac{a}{x^2}$ and $5y^2 + \frac{a}{x^2}$. Ans. $25y^4 - \frac{a^2}{x^4}$.
10. Multiply together $\frac{3x^2-5x}{14}$ and $\frac{7a}{2x^3-3x}$. Ans. $\frac{3ax-5a}{4x^2-6}$.
-

DIVISION OF FRACTIONS.

(104.) The Quotient of two Fractions is equal to the dividend multiplied by the *reciprocal of the divisor*, (80).

Thus the Quotient of $\frac{a}{b}$ divided by $\frac{x}{y}$ is equal to $\frac{a}{b} \times \frac{y}{x}$.

For since the value of the dividend is not altered by multiplying each of its terms by both terms of the divisor, (81), the quotient is equal to

$$\frac{axy}{bxy} \div \frac{x}{y}$$

And by dividing the numerator of this dividend by the numerator of the divisor, and the denominator by the denominator, we have the

quotient $\frac{ay}{bx}$, which is equal to $\frac{a}{b} \times \frac{y}{x}$.

Observe that the Quotient *multiplied into the divisor*, produces the dividend, (46.)

Thus $\frac{ay}{bx} \times \frac{x}{y}$ produces $\frac{axy}{bxy}$, equal to $\frac{a}{b}$.

Complex or Mixed Fractions.

(105.) When the dividend or the divisor is a Fraction or a *mixed* quantity, the dividend over the divisor, with a line between them, forms a *complex* or *mixed* fraction.

Thus $\frac{a}{b} \div (x+y)$ equals $\frac{\frac{a}{b}}{x+y}$, numerator $\frac{a}{b}$, denominator $x+y$.

From the nature of Division, and the proposition before demonstrated, (104), we have

R U L E X V I.

(106.) *For the Division of Fractions.*

1. Divide the numerator of the divisor into the numerator of the dividend, and the denominator into the denominator; or *multiply the dividend by the reciprocal of the divisor.*

2. A *fraction is divided by an integral* quantity, by dividing the numerator, or multiplying the denominator, by the integer.

3. An *integral quantity is divided by a fraction*, by dividing the integer by the numerator, and multiplying by the denominator; or by multiplying the integer by the *reciprocal of the fraction.*

4. A *mixed* quantity may be taken in division under the form of an *improper fraction*: or the integral and the fractional part may be divided separately

E X A M P L E.

To divide $\frac{10c^2}{a^2-2ax+x^2}$ by $\frac{5c^3}{a^2-x^2}$.

The dividend must be multiplied by $\frac{a^2-x^2}{5c^3}$, which is the reciprocal of the divisor.

If we cancel the factor $a-x$ from $a^2-2ax+x^2$, (60), and from a^2-x^2 , (58); and also cancel the factor $5c^2$, the operation will be, (103),

$$\frac{2}{a-x} \times \frac{a+x}{c}; \text{ which produces } \frac{2a+2x}{ac-cx}.$$

By thus canceling common factors, we find the Quotient in its *lowest terms.*

EXERCISES.

1. Divide $2a^2 - 2y^2$ by $\frac{a+y}{5cx}$.

Ans. $10acx - 10cxy$

2. Divide $\frac{3a}{a^2 - x^2}$ by $\frac{3}{a-x}$.

Ans. $\frac{a}{a+x}$.

3. Divide $\frac{ax+b}{a}$ by $\frac{bx-a}{b}$.

Ans. $\frac{abx+b^2}{abx-a^2}$.

4. Divide $\frac{a^2+2ab+b^2}{4ax^2}$ by $\frac{a+b}{2x}$.

Ans. $\frac{a+b}{2ax}$

5. Divide $\frac{6a^2b}{x^2-2xy+y^2}$ by $\frac{3a^2}{x-y}$.

Ans. $\frac{2b}{x-y}$

6. Divide $3a^2 + \frac{c+x^2}{2}$ by $a^2 - \frac{x^2}{3}$.

Ans. $3 + \frac{9x^2+3c}{6a^2-2x^2}$

7. Divide $\frac{2x^2}{a^3+x^3}$ by $\frac{x}{x+a}$.

Ans. $\frac{2x}{x^2-ax+a^2}$

8. Divide $\frac{5a^2y-10ay^2+5a}{ax^2}$ by $\frac{5a}{x^2}$.

Ans. $y - \frac{2y^2-1}{a}$.

9. Divide $\frac{a^3+b^3}{2+3x}$ by $ab+b^2$.

Ans. $\frac{a^2-ab+b^2}{2b+3bx}$

10. Divide $2a^2+4ay+2y^2$ by $\frac{a+y}{2}$.

Ans. $4a+4y$

11. Add $\frac{a}{3}$ to $\frac{x}{4}$, and divide the Sum by $\frac{3y}{4}$. *Ans.* $\frac{4a+3x}{9y}$.
12. Add $\frac{a-y}{5}$ to $\frac{a+y}{4}$, and divide the Sum by $\frac{x^2}{10}$. *Ans.* $\frac{9a+y}{2x^2}$.
13. Add $-\frac{x^2}{2}$ to $\frac{2x^2-1}{3}$, and divide the Sum by $\frac{x^4-4}{2}$. *Ans.* $\frac{1}{3(x^2+2)}$.
14. Add $-\frac{ax}{2}$ to $\frac{2ax+1}{3}$, and divide the Sum by $\frac{(ax+2)^2}{3}$. *Ans.* $\frac{1}{2(ax+2)}$.
15. Subtract $\frac{y}{3}$ from $\frac{ay}{2}$, and divide the Remainder by $\frac{2x}{3}$. *Ans.* $\frac{3ay-2y}{4x}$.
16. Subtract $\frac{y^2}{4}$ from $\frac{a^2}{4}$, and divide the Remainder by $\frac{a+y}{2}$. *Ans.* $\frac{a-y}{2}$.
17. Subtract $\frac{x^2}{5}$ from $\frac{a^2}{5}$, and divide the Remainder by $\frac{a+x}{3}$. *Ans.* $\frac{3(a-x)}{5}$.
18. Multiply $\frac{2ab-3}{x}$ by $\frac{5a}{x}$, and divide the Product by $\frac{5}{x}$. *Ans.* $\frac{2a^2b-3a}{x}$.
19. Multiply $\frac{ax^2}{y^2}$ by $\frac{ay^2}{x^2}$, and divide the Product by $\frac{a+b}{3}$. *Ans.* $\frac{3a^2}{a+b}$.
20. Multiply $\frac{a-x}{6}$ by $\frac{a+x}{3}$, and divide the Product by $\frac{y}{9}$. *Ans.* $\frac{a^2-x^2}{2y}$.
21. Multiply $\frac{9+y^2}{2}$ by $\frac{9-y^2}{5}$, and divide the Product by $\frac{3-y}{10}$. *Ans.* $27+9y+3y^2+y^3$.

CHAPTER V.

SIMPLE EQUATIONS.

(107.) AN EQUATION is an expression denoting the *equality of two quantities* by means of the sign =, *equal to*, placed between them.

The quantity on the left of the sign = is called the *first member*, or *side*, and that on the right the *second member*, or *side*, of the Equation.

Thus $3x + ab = 5x + 8d - 9$ is an equation, in which $3x + ab$ is the first member, and $5x + 8d - 9$ is the second.

(108.) Equations are employed in the solution of particular mathematical questions, or in the investigation of general mathematical principles.

In the solution of questions, the *unknown* or required quantity is represented by a letter, usually x , or y , &c., and an Equation is then formed which expresses the relation between this and the *known* or given quantities.

To give a simple example;—Suppose we wish to find a number the *third* and *fourth* of which shall together make 35.

Let x represent the number to be found, and the Equation

$$\text{will be } \frac{x}{3} + \frac{x}{4} = 35.$$

(109.) The *solution* of an Equation consists in finding the value of the *unknown quantity* in the equation.

The value found for the unknown quantity is *verified*, or the Equation *satisfied*, when this value, substituted for its symbol in the equation, makes the *first member the same as the second*.

The value of x in the preceding equation is 60, since this number, substituted for x , satisfies the equation; thus

$$\frac{60}{3} + \frac{60}{4} = 35.$$

The mode of solution will vary with the

Different Degrees of Equations.

(110.) A *simple* Equation, or an equation of the *first degree*, is one which contains no power of the unknown quantity but its *first power*.

$3x+ax-4=20$ is a simple equation.

A *quadratic* Equation, or an equation of the *second degree*, is one in which the highest power of the unknown quantity is its second power or *square*.

$2x^2+3ax+5=30$ is a quadratic equation.

A *cubic* Equation, or an equation of the *third degree*, is one in which the highest power of the unknown quantity is its third power, or *cube*; and so on.

Equations are also distinguished as

Numerical, Literal, and Identical Equations.

(111.) A *numerical* Equation is one in which all the *known* quantities are expressed by *numbers*.

$2x+5x=25-3$ is a numerical equation.

A *literal* Equation is one in which some or all of the *known* quantities are represented by *letters*.

$2x+ax=25-3b$ is a literal equation,

in which a and b are supposed to represent quantities whose values are *known*.

An *identical* Equation is one in which the two members *are the same*, or become the same by performing the operations which are indicated in them.

Thus $3x-3ab=3(x-ab)$ is an identical equation.

Transformation of Equations.

(112.) The *transformation* of an Equation consists in changing its *form*, without destroying the *equality* of the two members,—for the purpose of finding the value of the *unknown* quantity, or of discovering some general truth or principle.

These transformations depend, for the most part, on the following

Axioms.

(113.) An *Axiom* is a truth which is *self-evident*,—neither admitting nor requiring any *demonstration*; such as,

1. Things which are equal to the same thing, *are equal to each other*.

2. If equals be added to equals, the *sums will be equal*.

3. If equals be taken from equals, the *remainders will be equal*.

4. If equals be multiplied by equals, the *products will be equal*.

5. If equals be divided by equals, the *quotients will be equal*.

6. Any like *powers* or *roots* of equal quantities, *are equal*

SOLUTION OF SIMPLE EQUATIONS CONTAINING BUT ONE
UNKNOWN QUANTITY.

(114.) The *value of the unknown* quantity is found by making its symbol stand alone on one side of the Equation, so as to be equal to known quantities on the other side.

In order to this, the following transformations may be necessary, or at least may be expedient.

1. Clearing the Equation of Fractions.
2. The Transposition and Addition of Terms.
3. Changing the Signs of all the Terms in the Equation.
4. Dividing the Equation by the Coefficient of the Unknown quantity.

We shall apply each of these transformations to the solution of the same Equation.

Clearing an Equation of Fractions.

(115.) An Equation is cleared of fractions by multiplying each numerator into all the denominators *except its own*—regarding each *integral* term as a numerator,—and omitting the given denominators

Let the Equation be

$$\frac{3x}{4} - 7 = x - \frac{28}{3} + \frac{x}{3}$$

Multiplying the numerator of each fraction by the denominators of the other two, and the integral terms 7 and x by all the denominators, we obtain

$$27x - 252 = 36x - 336 + 12x.$$

The equality of the two members is not destroyed in thus clearing the Equation of fractions, because each of the terms connected by the signs $+$ and $-$ in the two members, is thus multiplied by all the denominators. (103) (113...4).

An Equation may also be *cleared of fractions* by multiplying its two members by the *least common multiple* of the denominators;—observing that a fractional term will be multiplied by multiplying its numerator into the *quotient* of said multiple \div the denominator.

In the given Equation the least common multiple of the denominators is 12. Multiplying by 12, we find

$$9x - 84 = 12x - 112 + 4x.$$

The advantage of this method is, that the new equation is found in its *lowest terms*.

Transposition and Addition of Terms.

(116.) Any term may be transposed from one side of an Equation to the other by *changing its sign*.—All the similar terms may thus be placed on the same side, and then added together.

In the last Equation $9x - 84 = 12x - 112 + 4x$, by transposing -84 to the second member, and $12x$ and $4x$ to the first, we have

$$9x - 12x - 4x = -112 + 84;$$

And by adding together the similar terms,

$$-7x = -28.$$

The equality of the two members is not destroyed by transposing a term with *its sign changed* from one side to the other, because this is equivalent to adding the term with *its sign changed* to both sides.

Thus by adding 84 to both members of the equation

$$9x - 84 = 12x - 112 + 4x,$$

the term -84 is canceled in the first member (28). In like manner by adding $-12x$ to both members, $12x$ is canceled in the second member; so also with $4x$. (113...2).

From the preceding principles it follows, that

Two equal terms with like signs on opposite sides of the sign $=$, may be at once suppressed from the Equation.

Change of the Signs in an Equation.

(117.) All the signs in an Equation may be changed, $+$ to $-$ and $-$ to $+$, without affecting the equality of its two members.

This follows from the principle of Transposition, (116), since in transposing *all the terms*, the signs would all be changed, but the two members would still be equal.

In the Equation already found

$$-7x = -28,$$

we shall have, by Transposition,

$$28 = 7x, \text{ or } 7x = 28.$$

The only Transformation which remains towards finding the *value* of x in the equation at first assumed, is that of dividing by the *coefficient* of x , the unknown quantity. (113...5).

Dividing both members of the preceding Equation by the coefficient of x , we find

$$x = \frac{-28}{-7} = 4; \text{ or } x = \frac{28}{7} = 4.$$

We have thus found the value of x to be 4. This value may be verified by substituting it for x in the original equation.

We may now give

RULE XVII.

(118.) *For the Solution of a Simple Equation containing but one unknown quantity.*

1. Clear the Equation of *fractions*, if it contains any.
2. Transpose all the terms containing the *unknown* quantity to one side, and all the known terms to the other side, of the equation.
3. Add together all the *similar terms* in each member.
4. Divide both members by the *coefficient* of the unknown quantity;—observing that when the unknown quantity is found in two or more *dissimilar* terms, its coefficient will be the sum of its coefficients in those terms.

NOTE.—When the sum of the terms containing the unknown quantity, after transposition, is *negative*, it will generally be expedient, though it is never necessary, to make it *positive* by changing all the signs in the equation.

EXAMPLE.

Given $\frac{x-3}{2} + \frac{x}{3} = 20 - \frac{x+19}{2}$, to find the value of x .

Clearing the equation of fractions, by multiplying it by the least common *multiple* of the denominator, which is 6, we have

$$3x - 9 + 2x = 120 - 3x - 57.$$

By transposition,

$$3x + 2x + 3x = 120 - 57 + 9.$$

Adding similar terms,

$$8x = 72.$$

Dividing by the coefficient of x ,

$$x = \frac{72}{8} = 9.$$

Remark.—The student is apt to err in Clearing an Equation of its fractions, when, as in this Example, a fraction preceded by the negative sign has a *polynomial numerator*.

The sign $-$ before the fraction in the second member above, denotes that the fraction is to be *subtracted*. When this fraction is multiplied by the 6, the product $3x + 57$ is subtracted by *changing its signs*. This gives the terms $-3x - 57$ in the new equation.

EXERCISES.

Numerical Equations.

1. Given $4x - 8 = 13 - 3x$ to find the value of x .
Ans. $x = 3$
 2. Given $7x + 17 = 10x - 19$ to find the value of x .
Ans. $x = 12$.
 3. Given $8x + 6 = 36 - 7x$ to find the value of x .
Ans. $x = 2$.
 4. Given $59 - 7x = 4x + 26$ to find the value of x .
Ans. $x = 3$
 5. Given $20 - 4x - 12 = 92 - 10x$ to find the value of x .
Ans. $x = 14$
 6. Given $8 - 3x + 12 = 30 - 5x + 4$ to find the value of x .
Ans. $x = 7$
-
7. Given $\frac{x}{4} + 24 = \frac{3x}{2}$ to find the value of x .
Ans. $x = 19\frac{1}{2}$.
 8. Given $\frac{x}{2} + \frac{x}{3} = 13 - \frac{x}{4}$ to find the value of x .
Ans. $x = 12$.
 9. Given $\frac{x-5}{4} + 6x = \frac{284-x}{5}$ to find the value of x .
Ans. $x = 9$
 10. Given $\frac{x+1}{2} + \frac{x+2}{3} = 16 - \frac{x+3}{4}$ to find the value of x .
Ans. $x = 13$.
 11. Given $3x - \frac{2x-5}{3} = x + \frac{x}{6} + 13\frac{1}{3}$ to find the value of x .
Ans. $x = 10$.
 12. Given $\frac{12x+26}{5} - 2x = 15 - \frac{x+3}{3}$ to find the value of x .
Ans. $x = 12$
 13. Given $\frac{19-x}{2} = x + \frac{11-x}{3}$ to find the value of x .
Ans. $x = 5$
 14. Given $\frac{6x-10}{3} = \frac{18-4x}{3} + x$ to find the value of x .
Ans. $x = 4$.
 15. Given $\frac{3x+4}{5} - \frac{7x-3}{2} = \frac{x-16}{4}$ to find the value of x .
Ans. $x = 2$

16. Given $\frac{x+1}{3} - 16 = \frac{x+4}{5} - \frac{x+3}{4}$ to find the value of x .

Ans. $x=41$

17. Given $\frac{x-1}{7} = x-3 - \frac{x+4}{3}$ to find the value of x .

Ans. $x=8$.

18. Given $\frac{4(x+2)}{3} = 1 + \frac{3x+1}{2}$ to find the value of x .

Ans. $x=7$.

19. Given $x - \frac{4x+2}{5} = 3x - 1\frac{1}{3}$ to find the value of x .

Ans. $x = \frac{1}{3}$.

20. Given $\frac{x}{2} - \frac{x}{3} + 5 = \frac{6(x+2)}{8}$ to find the value of x .

Ans. $x=6$.

21. Given $\frac{x}{3} + 6x = \frac{4x-2}{5}$ to find the value of x .

Ans. $x = -\frac{6}{83}$.

22. Given $x - \frac{3x-5}{2} = 12 - \frac{2x-4}{3}$ to find the value of x .

Ans. $x=65$

23. Given $\frac{21-3x}{3} - \frac{2(2x+3)}{9} = 6 - \frac{5x+1}{4}$ to find the value of x .

Ans. $x=3$.

An Equation in which the unknown quantity is found in every term, with different *exponents* in different terms, may often be reduced to a *simple* Equation by dividing it by some *power of the unknown quantity*. (113...5).

Thus if $2x^3 = 10x^2$,—
by dividing by x^2 we have $2x=10$; hence $x=5$.

24. Given $\frac{x^2}{5} + 3x = 7x - \frac{3x^2}{5}$ to find the value of x .

Ans. $x=5$.

25. Given $\frac{3x}{4} - \frac{2x}{5} = \frac{x^2-10x}{2}$ to find the value of x .

Ans. $x = 10\frac{7}{16}$.

26. Given $\frac{4}{3x} + \frac{5}{2x} = \frac{2}{x^2-10x}$ to find the value of x .

Ans. $x = 10\frac{1}{2}$

Literal Equations.

27. Given $ax - c = \frac{x - b}{a + c}$ to find the value of x .

Clearing the equation of fractions, by multiplying it by the denominator $a + c$, we have

$$a^2x - ac + acx - c^2 = x - b.$$

By transposition

$$a^2x + acx - x = ac + c^2 - b.$$

Dividing by the coefficient of x ,

$$x = \frac{ac + c^2 - b}{a^2 + ac - 1}.$$

28. Given $x + \frac{ax}{c} = b - \frac{dx}{c}$ to find the value of x .

$$\text{Ans. } x = \frac{bc}{a + c + d}.$$

29. Given $bc - \frac{ab}{x} = -d - \frac{1}{x}$ to find the value of x .

$$\text{Ans. } x = \frac{ab - 1}{bc + d}.$$

30. Given $3x - a = x - \frac{bx - d}{3}$ to find the value of x .

$$\text{Ans. } x = \frac{3a + d}{6 + b}.$$

31. Given $4abx^2 = \frac{3ax^2 - 2bx + ax}{3}$ to find the value of x .

$$\text{Ans. } x = \frac{a - 2b}{12ab - 3a}.$$

32. Given $\frac{x(a - b)}{2} = a + \frac{ab}{4} - \frac{x}{3}$ to find the value of x .

$$\text{Ans. } x = \frac{3a(b + 4)}{6(a - b) + 4}.$$

Remark.—In an *identical* Equation the unknown quantity has no *determinate value*, since any quantity whatever may be substituted for it, and the equation will be *satisfied*

Thus in the equation

$$3x - 5 = 3x - 5,$$

the two members will be equal whatever be the value of x . (113...3)

PROBLEMS

In Simple Equations of one unknown Quantity.

(119.) A *Problem* is a question proposed for solution; and the solution of a problem by Algebra consists in *forming an Equation* which shall express the *conditions* of the problem, and then solving the equation.

The general method of *forming the Equation* of a problem, is, to represent a *required* quantity by x , or y , &c., and then to perform or indicate the same operations that would be necessary to *verify* the value of x or y , supposing that *value to have been found*.

EXAMPLES AND EXERCISES.

1. What number is that to the double of which if 13 be added, the sum will be 75?

Let x represent the required number;
then $2x$ will represent *twice the number*;
and, by the conditions of the problem, the equation will be

$$2x + 13 = 75.$$

The value of x in this equation is the number required.

Ans. 31.

2. Find a number such that if it be multiplied by 5, and 24 be subtracted from the product, the remainder will be 36.

Ans. 12.

3. What number is that to $\frac{1}{3}$ of which if 25 be added, the sum obtained will be equal to the number itself *minus* 39?

Ans. 96.

4. Find a number such that if $\frac{1}{4}$ of it be subtracted from three times the number, the remainder will be 77.

Ans. 28.

5. Find what number added to the sum of one half, one third, and one fourth of itself will equal 4 added to twice the number.

Ans. 48.

6. Divide the number 165 into two such parts that the less may be equal to $\frac{1}{10}$ of the greater.

Let x represent the *less part*;
then $165 - x$ will represent the *greater*;
and the equation will be

$$x = \frac{165 - x}{10}.$$

Ans. 15 and 150.

7. Divide the number 100 into two such parts that six times the less may be equal to twice the greater.

Ans. 25 and 75.

8. It is required to divide 75 into two such parts that 3 times the greater may exceed 7 times the less by 15. *Ans.* 21 and 54.

9. What sum of money is that to which if \$100 be added, $\frac{2}{3}$ of the amount will be \$400? *Ans.* \$500.

10. A prize of \$100 is to be divided between two persons,—the share of the first being $\frac{7}{9}$ of that of the other. What are the shares? *Ans.* $\$43\frac{3}{4}$; $\$56\frac{1}{4}$.

11. A post is $\frac{1}{4}$ of its length in the mud, $\frac{1}{3}$ of it in the water, and 15 feet above the water. What is the length of the post? *Ans.* 36 feet.

12. Find a number such that if it be divided by 12, the divisor dividend and quotient together shall make 64. *Ans.* 48.

13. In a mixture of wine and cider, $\frac{1}{2}$ of the whole *plus* 25 gallons was wine, and $\frac{1}{3}$ part minus 5 gallons was cider. What was the whole number of gallons in the mixture? *Ans.* 120.

14. After a person had expended \$10 more than $\frac{1}{3}$ of his money, he had \$15 more than $\frac{1}{2}$ of it remaining. What sum had he at first? *Ans.* \$150.

15. Divide the number 91 into two such parts that if the greater be divided by their difference, the quotient may be 7. *Ans.* 49 and 42.

16. A and B had equal sums of money; the first paid away \$25, and the second \$60, when it appeared that A had twice as much left as B. What sum had each? *Ans.* \$95.

17. After paying away $\frac{1}{4}$ of my money, and then $\frac{1}{3}$ of what was left, I had \$180. What sum had I at first? *Ans.* \$300.

18. A line 37 feet in length is to be divided into 3 parts, so that the first may be 3 feet less than the second, and the second 5 more than the third; what are the parts? *Ans.* 12, 15, and 10 feet.

19. A can perform a piece of work in 12 days, and B can perform the same in 15 days. In what time could both together do the work?

Let x represent the number of days. Then since A could do $\frac{1}{12}$ of the work, and B $\frac{1}{15}$ of it, in 1 day,

$\frac{x}{12}$ will represent the part of the work A could do in x days,

$\frac{x}{15}$ will represent the part of the work B could do in x days

The equation is $\frac{x}{12} + \frac{x}{15} = 1$, the *entire work*. *Ans.* $6\frac{2}{3}$ days.

20. If A could mow a certain meadow in 6 days, B in 8 days, and C in 5 days, in what time could the three together do it?

Ans. $2\frac{2}{59}$ days.

21. Out of a cask of wine, which had leaked away a third part, 20 gallons were afterwards drawn, and the cask was then found to be but half full; how much did it hold?

Ans. 120 gallons.

22. It is required to divide \$300 between A, B, and C, so that A may have twice as much as B, and C as much as the other two together.

Ans. A \$100, B \$50, C \$150.

23. A gentleman spends $\frac{2}{3}$ of his yearly income in board and lodging, $\frac{2}{3}$ of the remainder in clothes, and then has \$20 left. What is the amount of his income?

Ans. \$180.

24. A person at the time he was married, was 3 times as old as his wife, but 15 years afterwards he was only twice as old. What were their ages on their wedding day?

Ans. 45 and 15 years.

25. Two persons, A and B, lay out equal sums of money in trade; the first gains \$126, and the second loses \$87, and A's money is now double of B's; what did each lay out?

Ans. \$300.

26. A courier, who travels 60 miles a day, had been dispatched 5 days, when a second is sent to overtake him, who goes 75 miles a day, in what time will he overtake him?

Ans. 20 days.

27. An island is 60 miles in circumference, and two persons start together to travel the same way around it: A goes 15 miles a day; and B 20; in what time would the two come together again?

Ans. 12 days.

28. A man and his wife usually drank out a cask of beer in 12 days, but when the man was from home it lasted the woman 30 days; how many days would the man alone be in drinking it?

Let x be the number of days;

then $\frac{1}{x}$ is the part that he would drink in 1 day;

and since the woman would drink $\frac{1}{30}$ of it in 1 day, the equation will be

$$\frac{1}{x} + \frac{1}{30} = \frac{1}{12}, \text{ the part both would drink in 1 day.}$$

Ans. 20 days.

29. If A and B together can do a piece of work in 9 days, and A alone could do it in 15 days, in what time ought B alone to accomplish the work?

Ans. $22\frac{1}{2}$ days.

30. The hour and the minute hand of a clock or watch are exactly together at 12 o'clock; when are they next together?

Ans. $5\frac{5}{11}$ minutes past one

31. It is required to divide \$1000 between A, B, and C, so that A shall have half as much as B, and C half as much as A and B together.

Ans. \$222 $\frac{2}{3}$; \$444 $\frac{4}{9}$; \$333 $\frac{1}{3}$.

32. A person being asked the hour, answered that the time past noon was $\frac{2}{3}$ of the time till midnight; what was the hour?

Ans. 48 min. past 4.

33. It is required to divide the number 60 into two such parts that their product shall be equal to 3 times the square of the less; what are the parts?

Ans. 15 and 45.

34. How much wine at 90 cents a gallon, and how much at \$1.50 a gallon, will be required to form a mixture of 20 gallons which shall be worth \$1.25 a gallon?

Ans. 8 $\frac{1}{3}$ gal.; 11 $\frac{2}{3}$ gal.

35. A cistern is supplied with water by three pipes which would severally fill it in 4, 5, and 6 hours. In what time would three pipes running together fill the cistern?

Ans. 1 $\frac{2}{3}$ $\frac{3}{7}$ hours.

36. If \$1000 be divided between A, B, and C, so that B shall have as much as A and half as much more, and C as much as B and half as much more, what will be the portion of each?

Ans. \$210 $\frac{10}{19}$; \$315 $\frac{15}{19}$; \$473 $\frac{13}{19}$.

37. A person has a lease for 99 years, and $\frac{2}{3}$ of the time which has expired on it is equal to $\frac{4}{5}$ of that which remains. Required the time which remains on the lease.

Ans. 45 years.

38. A merchant bought cloth at the rate of \$7 for 5 yards, which he sold again at the rate of \$11 for 7 yards, and gained \$100. How many yards were thus bought and sold?

Ans. 583 $\frac{1}{3}$ yards.

39. A and B together possess the sum of \$9800; and five-sixths of the sum owned by A is the same as four-fifths of that owned by B. What is the sum owned by each?

Ans. \$4800; \$5000.

40. The assets of a bankrupt amounting to \$5600 are to be divided among his creditors A, B, and C, according to their respective claims. A's claim is $\frac{1}{2}$ of B's, and C's is $\frac{2}{3}$ of B's; what sum must each of the creditors receive?

Ans. \$1292 $\frac{4}{13}$; \$2584 $\frac{8}{13}$; \$1723 $\frac{1}{13}$.

SIMPLE EQUATIONS CONTAINING TWO OR MORE UNKNOWN QUANTITIES.

(120.) It is sometimes necessary to employ two or more unknown quantities in the solution of a Problem; and in this case there must be formed as many *independent* Equations as there are unknown quantities employed.

Two equations are said to be *independent* of each other when they express essentially *different conditions*, so that one of the equations is not a mere transformation of the other.

Equations which thus express different conditions of the same Problem, are sometimes called *simultaneous equations*.

Solution of Two Simple Equations Containing Two Unknown Quantities.

(121.) From two Equations containing two unknown quantities we may derive a *new equation* from which one of those quantities shall be *eliminated*, or made to *disappear*. The value of the remaining unknown quantity may be found from the new equation; and this value *put for its symbol* in one of the given equations, will determine the other unknown quantity.

Elimination by Addition or Subtraction.

(122.) The two terms which contain the *same letter* in two Equations, may be made *equal* by multiplying or dividing the equations by proper quantities. That letter will then be *eliminated* in the *sum*, or else in the *difference*, of the new equations.

. EXAMPLE .

Given the equations $2x + 3y = 23$
and $5x - 2y = 10$,

to find the values of x and y .

Multiplying the first equation by 2, and the second by 3, we have

$$4x + 6y = 46,$$

$$\text{and } 15x - 6y = 30. \quad (113\dots4).$$

Adding together the corresponding sides of these equations, we find

$$19x = 76. \quad (113\dots2).$$

which gives $x = 4$.

Putting 4 for x in the first of the two given equations, we obtain

$$8 + 3y = 23,$$

which gives $y = 5$.

Observe that if $6y$ had the *same sign* in the two equations which were *added together*, this term would have been eliminated by taking the *difference*, not the *sum*, of these equations.

Elimination by Substitution.

(123.) The value of one of the unknown quantities in an Equation, may be found in *terms of all the other quantities* in the equation. If this value be then substituted for its representative letter in another equation, that letter will be *eliminated*.

EXAMPLE.

Given, as before, $2x + 3y = 23$,
and $5x - 2y = 10$,
to find the values of x and y .

We will find the value of x in the first equation, as if the value of y were *known*.

By transposition, $2x = 23 - 3y$;
dividing both members by the coefficient of x ,

$$x = \frac{23 - 3y}{2}.$$

We now substitute this value of x for x in the second equation. In doing this, we must multiply this fraction by the coefficient 5 in the first term of that equation.

$$\text{Then } \frac{115 - 15y}{2} - 2y = 10.$$

Clearing this equation of fractions, transposing, &c., we shall find

$$y = 5.$$

Putting 5 for y in the first equation,

$$2x + 15 = 23;$$

which gives $x = 4$.

The values of x and y are thus found to be the same as before.

This method of Elimination depends on the evident principle, that equivalent algebraic expressions may be taken, *the one for the other*; that is, *equal quantities may be substituted for each other*.

Elimination by Comparison.

(124.) If the value of the same letter be found in each of two Equations, in terms of the other quantities in the equations, that letter will be *eliminated* by putting one of these values *equal to the other*.

EXAMPLE.

Given, as before, $2x+3y=23$,

and $5x-2y=10$,

to find the values of x and y .

We will find the value of x in each equation, as if the value of y were known.

Transposing and dividing in the first equation,

$$x = \frac{23-3y}{2}.$$

Transposing and dividing in the second equation,

$$x = \frac{10+2y}{5}.$$

Putting the first of these values of x equal to the second,

$$\frac{23-3y}{2} = \frac{10+2y}{5}, (113....1).$$

which will give $y=5$.

By substituting 5 for y in any of the preceding equations, the value of x will be found to be 4, as in the two preceding solutions.

Before applying any of the preceding methods of Elimination, the Equations should generally be cleared of *fractions*, if they contain any; the necessary *transpositions* must be made; and similar terms must be added together.

Elimination by Addition or Subtraction will generally be found the simplest method, since it is free from *fractional expressions*, which are likely to occur in the application of the other two methods.

Let the student apply each of the three methods to the first ten of the following Exercises.

Equations may be marked, for reference, by the numbers (1), (2), (3), &c., or the capitals (A), (B), (C), &c.

Thus (A) $2x+y=10$, would be called equation (A).

EXERCISES.

1. Given $2x+3y=29$, and $3x-2y=11$, to find the values of x and y .
Ans. $x=7$, and $y=5$.
2. Given $5x-3y=9$, and $2x+5y=16$, to find the values of x and y .
Ans. $x=3$, and $y=2$.
3. Given $x+2y=17$, and $3x+y=16$, to find the values of x and y .
Ans. $x=3$, and $y=7$.
4. Given $4x+y=34$, and $10y-x=12$, to find the values of x and y .
Ans. $x=8$, and $y=2$.
5. Given $3x+4y=88$, and $6x+5y=128$, to find the values of x and y .
Ans. $x=8$, and $y=16$.
6. Given $7x+3y=42$, and $8y-2x=50$, to find the values of x and y .
Ans. $x=3$, and $y=7$.
7. Given $8y-3x=29$, and $6y-4x=20$, to find the values of x and y .
Ans. $x=1$, and $y=4$.
8. Given $6x-5y=39$, and $7x-3y=54$, to find the values of x and y .
Ans. $x=9$, and $y=3$.
9. Given $12x-9y=3$, and $12x+16y=228$, to find the values of x and y .
Ans. $x=7$, and $y=9$.
10. Given $5x+7y=201$, and $8x-3y=137$, to find the values of x and y .
Ans. $x=22$, and $y=13$.
11. Given $7y + \frac{x}{7} = 99$, and $7x + \frac{y}{7} = 51$, to find the values of x and y .
Ans. $x=7$, and $y=14$.
12. Given $\frac{x}{2} + \frac{y}{3} = 7$, and $\frac{x}{3} + \frac{y}{2} = 8$, to find the values of x and y .
Ans. $x=6$, and $y=12$.
13. Given $64 - \frac{2x}{3} = \frac{4y}{5}$, and $77 - \frac{5x}{6} = \frac{9y}{10}$, to find the values of x and y .
Ans. $x=60$, and $y=30$.
14. Given $21 - 6y = \frac{x+8}{4}$, and $23 - 5x = \frac{y+6}{3}$, to find the values of x and y .
Ans. $x=4$, and $y=3$.
15. Given $\frac{3x+4y}{5} = 10 - \frac{x}{4}$, and $\frac{6x-2y}{3} = 14 - \frac{y}{6}$, to find the values of x and y .
Ans. $x=8$, and $y=4$.
16. Given $\frac{2x}{3} + \frac{3y}{5} = \frac{22}{5}$, and $\frac{3x}{5} + \frac{2y}{3} = 4\frac{7}{15}$, to find the values of x and y .
Ans. $x=3$, and $y=4$.
17. Given $\frac{x-2}{5} - \frac{10-x}{3} = \frac{y-10}{4}$, and $\frac{2y+4}{3} - \frac{2x+y}{8} = \frac{x+13}{4}$

to find the values of x and y .

$$\text{Ans. } x=7, \text{ and } y=10$$

18. Find the values of x and y in the equations

$$10x+55=10y+\frac{6x-35}{5}, \text{ and } \frac{80+3x}{15}, =18\frac{1}{3}-\frac{4x+3y-8}{7}.$$

$$\text{Ans. } x=10, \text{ and } y=15.$$

19. Find the values of x and y in the equations

$$x-\frac{y-a}{b}=c, \text{ and } y-\frac{a-x}{b}=d.$$

Clearing the equations of fractions, we have

$$(A) \quad bx-y+a=bc, \text{ and } (B) \quad by-a+x=bd.$$

By transposition in these two equations, we find

$$(C) \quad bx-y=bc-a, \text{ and } (D) \quad by+x=bd+a.$$

Multiplying equation C by b , in order to eliminate y ,

$$(E) \quad b^2x-by=b^2c-ab.$$

Adding together equations D and E, (122.)

$$b^2x+x=b^2c+bd-ab+a.$$

Dividing both sides of this equation by the co-efficient of x ,

$$x=\frac{b^2c+bd-ab+a}{b^2+1}.$$

The value of y will be found if we multiply equation B by b , subtract equation A, &c.

$$y=\frac{b^2d+ab-bc+a}{b^2+1}.$$

20. Given $\frac{x}{a}+\frac{y}{a}=1$, and $bx+cy=dc$, to find the values of x and y .

$$\text{Ans. } x=\frac{ac-de}{c-b}, \text{ and } y=\frac{de-ab}{c-b}$$

21. Given $2cx-4c=-3dy$, and $7x=\frac{4b}{a}$, to find the values of x

$$\text{and } y. \quad \text{Ans. } x=\frac{4b}{7a}, \text{ and } y=\frac{28ac-8bc}{21ad}$$

22. Given $x+\frac{by}{a}=\frac{c}{a}$, and $m-\frac{ny}{x}=\frac{d}{x}$, to find the values of x and y

$$\text{Ans. } x=\frac{nc+bd}{na+mb}, \text{ and } y=\frac{mc-ad}{na+mb}.$$

SOLUTION OF THREE OR MORE SIMPLE EQUATIONS CONTAINING
AS MANY UNKNOWN QUANTITIES.

(125.) From three Equations containing three unknown quantities, we may derive two *new equations* containing but *two of the unknown quantities*—by eliminating one of the unknown quantities from the first and second equations, and the same unknown quantity from either of these and the third equation.

The two equations thus obtained may be solved as already exemplified, (121). Two of the unknown quantities will thus be determined; and by substituting their values in one of the given equations, the value of the other unknown quantity may be readily found.

In a similar manner four equations containing *four unknown quantities*, may be reduced to three equations containing but three unknown quantities; and these three may then be reduced to two. Five equations may be reduced to *four*, these four to three, and these three to two, &c.

EXAMPLE.

Given the equations $x + y + z = 29$,
 $x + 2y + 3z = 62$,
 $6x + 4y + 3z = 120$, to find x , y , and z .

Multiplying the first equation by 3, in order to eliminate z ,
 $3x + 3y + 3z = 87$.

Subtracting this equation from the third equation,

$$3x + y = 33;$$

Subtracting the second equation from the third,

$$5x + 2y = 58. \quad (122).$$

The last two equations contain but two unknown quantities, x and y , from which we may find $x = 8$, and $y = 9$.

Substituting these values of x and y in the first equation,

$$8 + 9 + z = 29,$$

which gives $z = 12$.

If the value of x were found in the first equation, and substituted for x in the second and third equations, we should then have two new equations containing y and z , (123).

We might also have found the two *new equations* by finding the value of the same letter, as x , in each of the three given equations then putting the first of these values equal to the second, and either the first or the second equal to the third, (124)

The best methods of elimination to be adopted in particular cases, can be learned only from experience. Regard should be had to simplicity and brevity in the operations employed.

EXERCISES.

23. Given $x+y+z=9$, $x+2y+3z=16$, and $x+3y+4z=21$, to find the values of x , y , and z . *Ans.* $x=4$, $y=3$, $z=2$.

24. Given $x+y+z=18$, $x+3y+2z=38$, and $x+\frac{1}{3}y+\frac{1}{2}z=10$, to find the values of x , y , and z . *Ans.* $x=4$, $y=6$, $z=8$.

25. Given $3x-9y=33$, $4y+2z=5x-20$, and $11x-7y=37+6z$, to find the values of x , y , and z . *Ans.* $x=2$, $y=-3$, $z=1$.

26. Given $2z=21-\frac{1}{3}(x+y)$, $3x=72$, and $38=\frac{1}{2}(3x+y-z)$ to find the values of x , y , and z . *Ans.* $x=24$, $y=9$, $z=5$.

PROBLEMS

In Simple Equations of one, two, &c., Unknown Quantities

(126.) When two or more required quantities are so related that, when one of them is found, the others may be conveniently *derived from that one*, the Problem may be most readily solved by a single Equation. In some questions, however, it is necessary to represent each of the required quantities by an appropriate symbol, and then to form as many Equations as there are unknown quantities to be found.

EXAMPLES AND EXERCISES.

1. Find two numbers such that $\frac{1}{2}$ of the first with $\frac{1}{3}$ of the second shall be equal to 9, and $\frac{1}{4}$ of the first with $\frac{1}{5}$ of the second shall be equal to 5.

Let x represent the *first*, and y the *second* number, then by the conditions of the problem, we have

$$\frac{x}{2} + \frac{y}{3} = 9, \text{ and } \frac{x}{4} + \frac{y}{5} = 5.$$

The values of x and y in these equations are the numbers required. *Ans.* 8 and 15.

2. Divide the number 100 into two such parts that $\frac{1}{3}$ of the first and $\frac{1}{4}$ of the second part shall together make 30.

Ans. 60 and 40

3. Find two numbers such that their sum shall be 60, and the less number $\frac{1}{3}$ of the greater. *Ans.* 15 and 45.

4. At a certain election 946 men voted for two candidates, and the successful one had a majority of 558. How many votes were given for each candidate? *Ans.* 752 and 194.

5. Divide the number 48 into two such parts that the quotient of the greater part divided by 4, may be equal to 4 times the quotient of the greater part divided by the less. *Ans.* 32 and 16.

6. A, B, and C make a joint contribution which in the whole amounts to \$400; B contributes twice as much as A and \$20 more, and C as much as the other two together. What sum did each contribute? *Ans.* A \$60, B \$140, C \$200.

7. Find three numbers such that the sum of the 1st and 2d shall be 35, the sum of the 1st and 3d 40, and the sum of the 2d and 3d 45. *Ans.* 15, 20, and 25.

8. A sum of money was divided between A and B, so that B's share was $\frac{2}{5}$ of A's, and A's share exceeded $\frac{5}{9}$ of the whole sum by \$50. What was the share of each? *Ans.* A's \$450, B's \$270.

9. The stock of three traders amounted to \$760. The shares of the 1st and 2d together exceeded the share of the 3d by \$240; and the share of the 1st was \$360 less than the sum of the shares of the other two; what was the share of each?

Ans. \$200; \$300; \$260.

10. A man being asked the age of himself and son, replied, "If I were $\frac{1}{4}$ as old as I am +3 times the age of my son, I should be 45; and if he were $\frac{1}{4}$ his present age +3 times mine, he would be 111." Required their ages. *Ans.* 36 and 12.

11. A and B together have \$340, B and C together \$384, and A and C together \$356; what sum has each?

Ans. \$156; \$184; \$200.

12. A number which is expressed by two digits is equal to 4 times the sum of its digits, and if 18 be added to the number, its digits will be interchanged with each other; what is the number?

Let x represent the *tens'*, and y the *units'* figure of the number, then $10x+y$ will represent the number.

By the conditions of the problem the equations will be

$$10x+y=4(x+y),$$

$$10x+y+18=10y+x.$$

Ans. 24.

13. It is required to divide the number 35 into three such parts, that $\frac{1}{2}$ of the first, $\frac{1}{3}$ of the second, and $\frac{1}{4}$ of the third, shall all be equal to each other; what are the parts? *Ans.* 8, 12, and 16.

14. A and B have both the same income; A saves $\frac{1}{5}$ of his annually, but B, by spending \$50 per annum more than A, at the end of 4 years, finds himself \$100 in debt; what is their income?

Ans. \$125.

15. A gentleman purchased a chaise, horse, and harness for \$180; the horse cost twice as much as the harness, and the chaise twice as much as the horse and harness together; what was the price of each?

Ans. \$120; \$40; \$20.

16. A farmer purchased 100 acres of land for \$2450; for a part of the land he paid \$20 an acre, and for the other part \$30 an acre. How many acres were there in each part?

Ans. 55, and 45 acres.

17. What fraction is that to the numerator of which if 1 be added, the value will be $\frac{1}{2}$; but if 1 be added to the denominator, the value of the fraction will be $\frac{1}{3}$?

Ans. $\frac{3}{8}$.

18. A and B together possess an income of \$570; if A's income were 3 times, and B's 5 times as much as each really is, they would together have \$2350. What is the income of each?

Ans. \$250; and \$320.

19. How old are we? said a person to his father: 6 years ago, replied the latter, I was a third more than 3 times as old as you were; but in 3 years, if I multiply your age by $2\frac{1}{8}$, it will then be equal to mine. What were their ages?

Ans. 15 and 36.

20. Find a number such that if we subtract it from 4980, divide the remainder by 8, and subtract 123 from the quotient, we shall find a remainder equal to the number itself.

Ans. 444.

21. A laborer engaged for 40 days on these conditions; that for every day he worked he should receive 80 cents, but for every day he was idle he should forfeit 32 cents. At the end of the time he was entitled to \$15.20; how many days did he work, and how many was he idle?

Ans. 25, and 15 days.

22. A cistern containing 820 gallons is filled in 20 minutes by 3 pipes, the first of which conveys 10 gallons more, and the second 5 gallons less than the third, per minute. How much flows through each pipe in a minute?

Ans. 22, 7, and 12 gallons.

23. A trader maintained himself for 3 years at an expense of £50 a year, and each year augmented that part of his stock which was not thus expended by $\frac{1}{3}$ thereof. At the end of the third year his original stock was doubled; what was that stock?

Ans. £740.

24. A and B began to trade with equal sums of money. The first year A gained \$40, and B lost \$40; the second year A lost $\frac{1}{3}$ of what he had at the end of the first, and B gained \$40 less than twice what A lost; when it appeared that B had twice as much money as A. What sum did each begin with? *Ans.* \$320.

25. What fraction is that, whose numerator being doubled, and denominator increased by 7, the value becomes $\frac{2}{3}$; but the denominator being doubled, and the numerator increased by 2, the value becomes $\frac{3}{5}$? *Ans.* $\frac{4}{5}$.

26. A and B together can perform a piece of work in 8 days, A and C in 9 days, and B and C in 10 days. How many days would it take each person to perform the same work alone?

Let x , y , and z represent the number of days required for A, B, and C respectively;

Then $\frac{1}{x}$ is the part of the work that A could do in 1 day, &c., and, by the conditions of the problem, the equations will be

$$\frac{1}{x} + \frac{1}{y} = \frac{1}{8}, \quad \frac{1}{x} + \frac{1}{z} = \frac{1}{9}, \quad \frac{1}{y} + \frac{1}{z} = \frac{1}{10}.$$

By subtracting the second equation from the first, we shall eliminate x , and then by adding the third equation we shall eliminate z

$$\text{Ans. A } 14\frac{3}{4}\frac{4}{9} \text{ days, B } 17\frac{2}{4}\frac{3}{1}, \text{ C } 23\frac{7}{3}1.$$

27. From two places, which are 154 miles apart, two persons set out at the same time to meet each other, one traveling at the rate of 3 miles in 2 hours, and the other at the rate of 5 miles in 4 hours; in how many hours will they meet? *Ans.* 56 hours.

28. In a naval engagement, the number of ships captured was 7 more, and the number burned was 2 less, than the number sunk. Fifteen escaped, and the fleet consisted of 8 times the number sunk; in how many ships did the fleet consist? *Ans.* 32.

29. A and B together could have completed a piece of work in 15 days, but after laboring together 6 days, A was left to finish it alone, which he did in 30 days. In how many days could each have performed the work alone? *Ans.* 50, and $21\frac{3}{7}$ days.

30. On comparing two sums of money it is found, that $\frac{2}{3}$ of the first is \$96 less than $\frac{3}{4}$ of the second, and that $\frac{5}{8}$ of the second is as much as $\frac{4}{9}$ of the first. What are the sums?

$$\text{Ans. } \$720, \text{ and } \$512.$$

31. A privateer, running at the rate of 10 miles an hour, discovers a ship 18 miles off, making way at the rate of 8 miles an hour. In how many hours will the ship be overtaken? *Ans.* 9 hours.

32. In a composition of copper, tin, and lead, $\frac{1}{2}$ of the whole *minus* 16 pounds was copper, $\frac{1}{3}$ of the whole *minus* 12 pounds was tin, and $\frac{1}{4}$ of the whole *plus* 4 pounds was lead; what quantity of each was there in the composition? *Ans.* Copper 128, tin 84, lead 76 pounds.

33. The sum of \$660 was raised for a certain purpose by four persons, the first giving $\frac{1}{2}$ as much as the second, the third as much as the first and second, and the fourth as much as the second and third. What were the several sums contributed?

Ans. \$60, \$120, \$180, \$300.

34. Two pedestrians start from the same point, and go in the same direction; the first steps twice as far as the second, but the second makes 3 steps while the first is making 2. How far has each one gone when the first is 300 feet in advance of the second?

Ans. 1200, and 900 feet.

35. A merchant has cloth at \$3 a yard, and another kind at \$5 a yard. How many yards of each kind must he sell, to make 100 yards which shall bring him \$450?

Ans. 25, and 75 yards.

36. In the composition of a quantity of gunpowder, the nitre was 10 pounds more than $\frac{2}{3}$ of the whole, the sulphur $4\frac{1}{2}$ pounds less than $\frac{1}{6}$ of the whole, and the charcoal 2 pounds less than $\frac{1}{7}$ of the nitre. What was the amount of gunpowder?

Ans. 69 pounds.

37. Four places are situated in the order of the letters A, B, C, D. The distance from A to D is 34 miles; the distance from A to B is $\frac{2}{3}$ of the distance from C to D; and $\frac{1}{4}$ of the distance from A to B, *plus* $\frac{1}{2}$ of the distance from C to D, is 3 times the distance from B to C. What are the distances between A and B, B and C, C and D?

Ans. 12, 4, and 18 miles.

38. A vintner sold at one time 20 dozen of port wine, and 30 of sherry, for \$120; and at another time 30 dozen of port, and 25 of sherry, at the same prices as before, for \$140. What was the price of a dozen of each sort of wine?

Ans. \$3, and \$2.

39. A person pays, at one time, to two creditors, \$53, giving to one of them $\frac{4}{11}$ of the sum due to him, and to the other \$3 more than $\frac{1}{6}$ of his debt to him. At another time he pays them \$42, giving to the first $\frac{3}{7}$ of what remains due to him, and to the other $\frac{1}{3}$ of what remains due to him. What were the debts?

Ans. \$121, and \$36.

40. A farmer has 86 bushels of wheat at 4s. 6d. per bushel, with which he wishes to mix rye at 3s. 6d. per bushel, and barley at 3s. per bushel, so as to make 136 bushels, that shall be worth 4s. a bushel. What quantity of rye and of barley must he take?

Ans. 14, and 36 bushels.

41. A composition of copper and tin, containing 100 cubic inches, weighs 505 ounces. How many ounces of each metal does it contain, supposing the weight of a cubic inch of copper to be $5\frac{1}{4}$ ounces, and of a cubic inch of tin $4\frac{1}{4}$ ounces? *Ans.* 420, and 85 ounces.

42. A General having lost a battle, found that he had only one-half of his army *plus* 3600 men left, fit for action; $\frac{1}{8}$ of his men *plus* 600 being wounded, and the rest, who were $\frac{1}{5}$ of the whole army, either slain, taken prisoners, or missing. Of how many men did his army consist? *Ans.* 24000.

43. Two pipes, one of them running 5 hours, and the other 4, filled a cistern containing 330 gallons; and the same two pipes, the first running 2 hours, and the second 3, filled another cistern containing 195 gallons. How many gallons did each pipe discharge per hour? *Ans.* 30 and 45 gallons.

44. After A and B had been employed on a piece of work for 14 days, they called in C, by whose aid it was completed in 28 days. Had C worked with them from the beginning, the work would have been accomplished in 21 days. In how many days would C alone have accomplished the work? *Ans.* 42 days.

45. Some smugglers discovered a cave which would exactly hold their cargo, viz., 13 bales of cotton and 33 casks of wine. A revenue cutter coming in sight while they were unloading, they sailed away with 9 casks and 5 bales, leaving the cave two-thirds full. How many bales or casks would it contain? *Ans.* 24 bales or 72 casks.

46. A gentleman left a sum of money to be divided among four servants, so that the share of the first was $\frac{1}{2}$ the sum of the shares of the other three; the share of the second $\frac{1}{3}$ of the sum of the other three; and the share of the third $\frac{1}{4}$ of the sum of the other three; and it was also found that the share of the first exceeded that of the last by \$14. What was the whole sum? and the share of each?

Ans. Whole sum \$120; shares \$40; \$30; \$24; \$26.

CHAPTER VI.

RATIO—PROPORTION—VARIATION.

RATIO.

(127.) The *RATIO* of one quantity called the *antecedent* to another of the same kind called the *consequent*, is the *quotient* of the former divided by the latter.

Thus the *ratio* of 12 to 4 is 3, since 12 is 3 *times* 4;

and the ratio of 5 to 13 is $\frac{5}{13}$, since 5 is five *thirteenth*s of 13.

The antecedent and consequent together are called the *terms* of the ratio.

Sign of Ratio.

(128.) A *colon* (:) between two quantities denotes that the two quantities are taken as the *antecedent* and *consequent* of a ratio.

Thus 3 : 5, the ratio of 3 to 5; $a : b$, the ratio of a to b .

(129.) The *value* of a ratio may always be represented by making the antecedent the *numerator*, and the consequent the *denominator* of a Fraction.

Thus 3 : 5 is equal to $\frac{3}{5}$; and $a : b$ is equal to $\frac{a}{b}$, (75).

Direct and Inverse Ratio.

(130.) The *direct ratio* of the first of two quantities to the second, is the quotient of the *first divided by the second*; thus the direct ratio of 3 to 5 is $\frac{3}{5}$.

The *inverse ratio* of the first quantity to the second, is the direct ratio of the *second to the first*;—in other words, it is the direct ratio of the *reciprocals of the two quantities*.

Thus the *inverse ratio* of 3 to 5 is $\frac{5}{3}$;—

or it is the ratio of $\frac{1}{3}$ to $\frac{1}{5}$, equal to $\frac{1}{3} \div \frac{1}{5}$, equal to $\frac{5}{3}$, (127).

Hence *inverse* is often called *reciprocal* ratio. The term *ratio* used alone always means *direct* ratio.

Compound Ratio.

(131.) A *compound ratio* is the ratio of the *product* of two or more antecedents to the product of their consequents; and is equal to the product of all the *simple ratios*.

The compound ratio of a and b to x and y is $\frac{ab}{xy}$;

= the product of the simple ratios $\frac{a}{x}$ and $\frac{b}{y}$, or $\frac{a}{y}$ and $\frac{b}{x}$.

(132.) The ratio of the *first to the last* of any number of quantities, is equal to the product of the ratios of the first to the second, the second to the third, and so on to the last; that is, it is *compounded of all the intervening ratios*.

For example, take the quantities a, b, x, y . The ratios of the first to the second, the second to the third, &c., are

$\frac{a}{b}, \frac{b}{x}, \frac{x}{y}$; and their product is $\frac{abx}{bxy}, = \frac{a}{y}$, which is $a : y$.

Duplicate and Triplicate Ratios.

(133.) The *duplicate ratio* of two quantities is the ratio of their *squares*, and the *triplicate ratio* is the ratio of their *cubes*.

Thus the duplicate ratio of a to b is the ratio of a^2 to b^2 ;
and the triplicate ratio of a to b is the ratio of a^3 to b^3 .

The *subduplicate* ratio of quantities is the ratio of their *square roots*, and the *subtriplicate* ratio is the ratio of their *cube roots*.

Equimultiples and Equisubmultiples.

(134.) *Equimultiples* of two quantities are the products which arise from multiplying the quantities by the same *integer*, and *equisubmultiples* are the *quotients* which arise from dividing the quantities by the same *integer*.

Thus $3a$ and $3b$ are equimultiples of a and b , while, conversely, a and b are equisubmultiples of $3a$ and $3b$.

(135.) *Equimultiples*, or *equisubmultiples*, of two quantities have the same ratio as the quantities themselves, (81).

PROPORTION.

(136.) PROPORTION consists in an equality of the *ratios* of two or more antecedents to their respective consequents—but is usually confined to *four terms*.

(137.) *Four quantities* are in Proportion when the ratio of the *first* to the *second* is equal to the ratio of the third to the fourth; that is, when the first *divided by the second* is equal to the third divided by the fourth.

Thus the numbers 6, 3, 8, 4 are in proportion,
since the ratio $\frac{6}{3}$ equals the ratio $\frac{8}{4}$.

And the quantities a, b, x, y are in proportion,

when the ratio $\frac{a}{b}$ equals the ratio $\frac{x}{y}$, (129).

The first and third terms are the *antecedents* of the ratios; the second and fourth are the *consequents*. The first and fourth are the two *extremes*; the second and third are the two *means*.

The *fourth term* is called a fourth *proportional* to the other three taken in order; thus 4 is a fourth proportional to 6, 3, and 8.

(138.) *Three quantities* are in Proportion when the ratio of the first to the second is equal to the ratio of the *second to the third*,—the second term being called a *mean proportional* between the other two.

Thus the numbers 8, 4, 2 are in proportion,
since the ratio $\frac{8}{4}$ equals the ratio $\frac{4}{2}$;
and 4 is a *mean proportional* between 8 and 2.

Direct and Inverse Proportion.

(139.) A *direct* Proportion consists in an equality between two *direct* ratios, and an *inverse* or reciprocal Proportion in an equality between a direct and an inverse ratio.

Thus the numbers 6, 3, 8, 4 are in *direct* proportion; (137).

The same numbers in the order 6, 3, 4, 8 are in *inverse* proportion, since the direct ratio $\frac{6}{3}$ is equal to the *inverse* ratio $\frac{8}{4}$, (130).

The term *proportion* used alone always means *direct* proportion

Sign of Proportion.

(140.) A Proportion is denoted by a *double colon* ($::$), or the sign $=$ between the equal ratios of the proportion.

Thus $6 : 3 :: 8 : 4$, or $6 : 3 = 8 : 4$, or $\frac{6}{3} = \frac{8}{4}$ denotes that these numbers are in proportion, and is read
6 is to 3 as 8 is to 4.

To denote an *inverse* Proportion we employ the sign \neq between the two ratios of such proportion.

Thus $6 : 3 \neq 4 : 8$, denotes that 6 is to 3 *inversely* as 4 is to 8.

Inverse Converted Into Direct Proportion.

(141.) An *inverse* is converted into a *direct* Proportion by interchanging either antecedent and its consequent; or by substituting the *reciprocals* of either antecedent and its consequent.

Thus from the *inverse* proportion $6 : 3 \neq 4 : 8$, we get the *direct* proportion $3 : 6 = 4 : 8$, by interchanging 6 and 3,
or $\frac{1}{6} : \frac{1}{3} = 4 : 8$, by substituting $\frac{1}{6}$ and $\frac{1}{3}$.

The reason of this is evident from the nature of inverse ratio, (130).

VARIATION.

(142.) VARIATION is such a dependence of one term or quantity on another, that any *new value* of one of them will produce a new value of the other, in a constant ratio of increase or diminution.

1. One quantity *varies directly* as another when their dependence is such that if one of them be *multiplied*, the other must be multiplied, by the same quantity.

For example, the Interest on money, for a given *time* and *rate* per cent., *varies directly* as the Principal, since the Interest will be doubled, or tripled, &c., if the Principal be doubled, or tripled, &c.

2. One quantity *varies inversely* as another when their dependence is such that if one of them be multiplied, the other must be *divided*, by the same quantity.

For example, the Time in which a given amount of *interest* will accrue on a given *principal*, *varies inversely* as the Rate per cent., since the Time will be doubled, &c. if the Rate be *halved*, &c.

(143.) When one quantity *varies inversely* as another, the *product* of the two is always the same *constant quantity*.

For as one of the two quantities is *multiplied*, the other is *divided* by the same number; the product of the two will therefore be multiplied and divided by the same number; hence its value will remain *unchanged*.

Variation—an Abbreviated Proportion.

(144.) The two terms of a variation are the two *antecedents* in a Proportion in which the two consequents are not expressed, but may be understood, to complete the proportion.

Thus when we say that the Interest varies as the Principal, for a given *time* and *rate* per cent., it is understood, that

The Interest on any principal *is to* the Interest on any other Principal, for the same time and rate, *as* the *first* Principal *is to* the second.

Instead of saying "the Interest varies as the Principal," we may say, the Interest is *proportional* to the Principal; which is a brief method of expressing a Proportion by means of its *antecedents*,—the consequents being understood.

Sign of Variation.

The character \sim placed between two terms, denotes that one of them *varies as the other*. Thus $x \sim y$, x varies as y , or x is proportional to y .

$x \sim \frac{1}{y}$ denotes that x varies as the *reciprocal* of y ; or x varies reciprocally or *inversely* as y .

$x \sim \frac{y}{z}$ denotes that x varies *directly* as y , and *inversely* as z ; that is, x varies as the *quotient* of y divided by z .

THEOREMS IN PROPORTION.

(145.) A *Theorem* is a proposition to be demonstrated or proved.—

A *Corollary* is an inference drawn from a preceding proposition or demonstration.

THEOREM I.

(146.) Two Fractions having a *common denominator*, are to each other as *their numerators*; and two fractions having a common numerator are to each other *inversely as their denominators*

First. Let d be the common denominator;

then the ratio of $\frac{a}{d}$ to $\frac{c}{d}$ is

$$\frac{a}{d} \div \frac{c}{d} = \frac{ad}{cd} = \frac{a}{c}, \quad (127),$$

and $\frac{a}{c}$ is the ratio of the numerator a to the numerator c .

Secondly. Let n be the common numerator;

then the ratio of $\frac{n}{a}$ to $\frac{n}{c}$ is

$$\frac{n}{a} \div \frac{n}{c} = \frac{cn}{an} = \frac{c}{a};$$

and $\frac{c}{a}$ is the *inverse ratio* of the denominator a to the denominator c , (130).

Therefore, two fractions having a common denominator, &c.

(147.) *Corollary.* The value of a Fraction varies *directly as its numerator*, and *inversely as its denominator*.

THEOREM II.

(148.) In any Proportion, if one antecedent be greater than its consequent, the other antecedent will be *greater than its consequent*; if equal, equal; and if less, less.

Let $a : b :: x : y$;

$$\text{then } \frac{a}{b} = \frac{x}{y}, \quad (137).$$

Now if a be greater than b the first ratio will be greater than a unit, and consequently the second ratio will be greater than a unit, and therefore x will be greater than y . In like manner if a be equal to b , x will be equal to y , &c.

Hence, in any Proportion, if one antecedent, &c.

THEOREM III.

(149.) When four quantities are in Proportion, the product of the two extremes is equal to the product of the two means.

Let $a : b :: x : y$;

then is $ay = bx$.

For since the quantities are in proportion,

$$\frac{a}{b} = \frac{x}{y}, \quad (137).$$

Clearing the Equation of fractions,

$$ay = bx. \quad (115).$$

Therefore, when four quantities are in Proportion, &c.

(150.) *Cor. 1.* A fourth proportional to three given quantities, is found by dividing the product of the second and third by the first.

Thus from the equation $ay = bx$, we find $y = \frac{bx}{a}$.

(151.) *Cor. 2.* When three quantities are in Proportion, the product of the two extremes is equal to the square of the mean.

For let $a : b :: b : x$; then $ax = bb = b^2$.

(152.) *Cor. 3.* A mean proportional between two given quantities, is equal to the square root of their product.

Thus from the equation $ax = b^2$, we find $b = (ax)^{\frac{1}{2}}$.

THEOREM IV.

(153.) When the product of two quantities is equal to the product of two other quantities, either pair of factors may be made the extremes, and the other the means, of a Proportion.

Let $ab = xy$;

then will $a : x :: y : b$.

Dividing both sides of the given equation by b ,

$$a = \frac{xy}{b}.$$

Dividing both sides of this last equation by x ,

$$\frac{a}{x} = \frac{y}{b}.$$

Hence $a : x :: y : b$, (137).

In like manner a and b may be taken for the means, and x and y for the extremes. Therefore, when the product of two quantities, &c

THEOREM V.

(154.) If three quantities are in Proportion, the first will be to the *third* as the *square of the first* to that of the second, or the square of the second to that of the third.

$$\text{Let } a : b :: b : x ;$$

then will $a : x :: a^2 : b^2$; or $a : x :: b^2 : x^2$.

From the given proportion, we find

$$ax = b^2, \quad (151).$$

Multiplying both sides of this equation by a ,

$$a^2x = ab^2.$$

Converting this equation into a Proportion,

$$a : x :: a^2 : b^2, \quad (153).$$

And by multiplying both sides of the first equation by x , we shall, in like manner, find $a : x :: b^2 : x^2$.

Therefore, if three quantities are in proportion, &c.

THEOREM VI.

(155.) Four quantities in Proportion are also in proportion by *inversion*,—that is, when each antecedent and its consequent are *interchanged with each other*.

$$\text{Let } a : b :: x : y ;$$

then is $b : a :: y : x$.

From the given proportion we find

$$ay = bx, \quad (149).$$

Making b and x the extremes, and a and y the means,

$$b : a :: y : x, \quad (153).$$

Hence, four quantities in proportion are also in proportion, &c.

THEOREM VII.

(156.) Four quantities in Proportion are also in proportion by *alternation*,—that is, when the two *means*, or the two *extremes*, are *interchanged with each other*.

$$\text{Let } a : b :: x : y ;$$

then is $a : x :: b : y$; or $y : b :: x : a$.

From the given proportion we find

$$ay = bx, \quad (149).$$

This equation may be converted into the proportion

$$a : x :: b : y, \text{ or } y : b :: x : a, \quad (153).$$

Therefore, four quantities in proportion, &c.

THEOREM VIII.

(157.) In any Proportion, if the *two antecedents*, or the two consequents, or an antecedent and its consequent, be *multiplied by the same quantity*, the products and the remaining terms will be in proportion.

Let $a : b :: x : y$;
and let n be any numerical quantity ;
then will $an : b :: nx : y$, &c.

From the given proportion we have

$$ay = bx.$$

Multiplying both sides of this equation by n ,

$$any = bnx.$$

Converting this equation into a Proportion,

$$an : b :: nx : y ; \text{ or } a : bn :: x : ny ; \text{ or } a : b :: nx : ny, \quad (153).$$

Therefore, in any proportion, if the two antecedents, &c.

(158.) *Cor.* If the two *antecedents*, or the two consequents, or an antecedent and its consequent, be *divided by the same quantity*, the quotients and the remaining terms will be in proportion.

For dividing by a quantity is equivalent to *multiplying by its reciprocal*.

THEOREM IX.

(159.) Four quantities in Proportion are also in proportion by *composition*,—that is, the *sum of the first and second terms* is to the *first or second*, as the *sum of the third and fourth* is to the *third or fourth*

$$\text{Let } a : b :: x : y$$

then is $a + b : a :: x + y : x$.

From the given proportion we have

$$ay = bx.$$

Adding both sides of this equation to ax ,

$$ax + ay = ax + bx.$$

Resolving each member of this equation into its factors,

$$a(x + y) = x(a + b).$$

Converting this last equation into a Proportion,

$$a + b : a :: x + y : x, \quad (153).$$

By adding both sides of the first equation to by , it may be proved, in like manner, that $a + b : b :: x + y : y$.

Therefore, four quantities in proportion are also in proportion, &c

THEOREM X.

(160.) Four quantities in Proportion are also in proportion by *division*,—that is, the *difference of the first* and second terms is to the *first* or second, as the *difference of the third* and fourth is to the *third* or fourth.

Let $a : b :: x : y$;
then is $a - b : a :: x - y : x$.

From the given proportion we find

$$ay = bx.$$

Subtracting both sides of this equation from ax ,

$$ax - ay = ax - bx.$$

Resolving each member of this equation into its factors,

$$a(x - y) = x(a - b).$$

Converting this last equation into a Proportion,

$$a - b : a :: x - y : x.$$

By subtracting both sides of the first equation from by , it may be proved, in like manner, that $a - b : b :: x - y : y$.

Hence, four quantities in proportion are also in proportion, &c.

THEOREM XI.

(161.) When *any number of quantities* are in Proportion, the sum of any *two or more of the antecedents* is to the sum of their consequents, as any one antecedent is to its consequent.

Let $a : b :: c : d :: x : y$, &c. ;
then is $a + c : b + d :: x : y$.

From the given proportion we shall find

$$ay = bx, \text{ and } cy = dx, \quad (149).$$

Adding together the corresponding members of these two equations,

$$ay + cy = bx + dx.$$

Resolving each member of this equation into its factors,

$$(a + c)y = (b + d)x,$$

Converting this equation into a Proportion,

$$a + c : b + d :: x : y.$$

By adding xy to both sides of the *third* equation, we shall, in like manner, find that $a + c + x : b + d + y :: x : y$.

Therefore, when any number of quantities are in proportion, &c.

THEOREM XII.

(162.) If two Proportions have an *antecedent* and *its consequent*, or the two antecedents, or the two consequents, the *same in both*, the remaining terms will be in proportion.

$$\begin{aligned} &\text{Let } a : b :: x : y, \\ &\text{and } a : b :: w : z; \\ &\text{then will } x : y :: w : z. \end{aligned}$$

For the ratio of x to y is equal to the ratio of w to z , since each of these ratios is equal to the ratio of a to b ; hence

$$x : y :: w : z.$$

The two given proportions have an *antecedent* and its consequent *the same in both*. If the two antecedents were the same in both, the demonstration would be the same, after *interchanging the means*; and if the two consequents were the same, after interchanging the extremes, (156).

Hence, if two proportions have an antecedent and its consequent, &c.

Two or more Proportions having an antecedent and its consequent *the same in each*, form one *continued proportion*.

$$\begin{aligned} &\text{Thus, } a : b :: u : v, \\ &\quad a : b :: w : x, \\ &\quad \text{and } a : b :: y : z, \end{aligned}$$

form the continued proportion, $a : b :: u : v :: w : x : y : z$.

THEOREM XIII.

(163.) The sum of the *first* and *second* terms in any Proportion, is to their *difference*, as the sum of the third and fourth is to their difference.

$$\begin{aligned} &\text{Let } a : b :: x : y, \\ &\text{then is } a+b : a-b :: x+y : x-y. \end{aligned}$$

By Composition and Division, in the given proportion.

$$\begin{aligned} a+b : a &:: x+y : x; & (159); \\ a-b : a &:: x-y : x. & (160). \end{aligned}$$

These two proportions have the consequents a and x the same in both; hence $a+b : a-b :: x+y : x-y$, (162).

Therefore, the sum of the first and second terms in any proportion, &c.

THEOREM XIV.

(164.) The *products of the corresponding terms* of two or more Proportions, are in proportion.

$$\begin{aligned} \text{Let } a : b :: x : y, \\ \text{and } c : d :: w : z; \end{aligned}$$

then is $ac : bd :: xw : yz$.

From the two given proportions, we have

$$ay = bx, \text{ and } cz = dw.$$

Multiplying together the corresponding members of these equations.

$$ac \cdot yz = bd \cdot xw.$$

Converting this equation into a Proportion,

$$ac : bd :: xw : yz, \quad (153).$$

In like manner the demonstration may be extended to three or more proportions. Hence, the products of the corresponding terms, &c.

(165.) *Cor.* Like *powers or roots* of proportional quantities, are in proportion.

For if $a : b :: x : y$, by multiplying each term by itself, we shall have, according to the Theorem, $a^2 : b^2 :: x^2 : y^2$; and multiplying these by the given terms, we shall have $a^3 : b^3 :: x^3 : y^3$, &c.

THEOREM XV.

(166.) For *any factors* in an antecedent and its consequent, or the two antecedents, or the two consequents, in a Proportion, may be substituted any other quantities which have the *same ratio to each other*

$$\begin{aligned} \text{Let } a : b :: nx : py, \\ \text{and } n : p :: r : s; \end{aligned}$$

then will $a : b :: rx : sy$.

From the two given proportions, we find

$$an : bp :: nrx : psy, \quad (164).$$

Dividing the antecedents in this proportion by n , and the consequents by p ,

$$a : b :: rx : sy, \quad (158).$$

This proves the *first affirmation* in the Theorem. By interchanging the *extremes* in the two given proportions, and afterwards the *means*, (156), the other two affirmations in the theorem may be demonstrated.

Therefore, for any factors in an antecedent and its consequent, &c.

GENERAL SOLUTIONS OF PROBLEMS.—FORMULAS. APPLICATIONS
OF PROPORTION, &c.

(167.) In the *general solution* of a Problem, all the quantities are represented by *letters*; and the *unknown* being thus found in terms of *all the known* quantities, the result discloses a *rule* for the numerical computation in any given case of the problem.

EXAMPLE.

To find two numbers whose sum shall be s , and difference d .

Let x represent the *greater*, and y the *less* number.

From the conditions of the problem, we have

$$\begin{aligned}x + y &= s, \\ \text{and } x - y &= d.\end{aligned}$$

By adding the second equation to the first; and also subtracting the second from the first, we have

$$2x = s + d, \text{ and } 2y = s - d,$$

which give
$$x = \frac{s + d}{2}, \text{ and } y = \frac{s - d}{2}.$$

From these general values of x and y , we learn that the *greater* of two numbers is equal to $\frac{1}{2}$ of (the *sum* + the *difference*), and that the *less* is equal to $\frac{1}{2}$ of (the *sum* - the *difference*), of the two numbers; hence the following *rule*:

(168.) *To find two numbers from their sum and difference*,—Add the difference to the sum, and divide by 2, for the *greater* of the two numbers; subtract the difference from the sum, and divide by 2, for the *less* number.

For example, if the sum of two numbers be 500, and their difference 146,

$$\text{The greater number is } \frac{500 + 146}{2} = 323;$$

$$\text{and the less number is } \frac{500 - 146}{2} = 177.$$

(169.) An algebraic *Formula* is an equation between the *symbols* of certain quantities—resulting from the general solution of a Problem, or the investigation of some general principle.

Thus the equations which express the *values* of x and y in the preceding Example, are *formulas* for finding two numbers from the *sum* and *difference* of the numbers.

PROBLEMS

In Proportion, Percentage, Interest, &c

(170.) A Proportion occurring in the solution of a Problem, may be converted into an Equation by putting the *product of the two extremes* equal to the *product of the two means*, (149).

EXAMPLE.

To divide \$1000 between three persons, in the proportions of 2, 3, and 5; that is, so that A's share shall be to B's as 2 to 3, and B's to C's as 3 to 5.

Let x represent A's share; y , B's share; and z , C's share. Then by the conditions of the problem, we have

$$x + y + z = 1000;$$

$$x : y :: 2 : 3;$$

$$\text{and } y : z :: 3 : 5.$$

By converting the two Proportions into Equations, we find

$$3x = 2y, \text{ and } 5y = 3z.$$

We have now *three equations* from which to find the values of the *three unknown quantities*, (125).

The solution of the Problem may also be effected with *one unknown quantity*, by finding *fourth proportionals* for the shares of B and C.

Let x represent A's share;

$$\text{then } 2 : 3 :: x : \frac{3x}{2}; \text{ and } 2 : 5 :: x : \frac{5x}{2}, \quad (150).$$

Hence B's share is represented by $\frac{3x}{2}$, and C's by $\frac{5x}{2}$; and the equation of the problem is

$$x + \frac{3x}{2} + \frac{5x}{2} = \$1000.$$

Ans. The several shares are \$200, \$300, \$500

The *general* Problem of which the preceding is a particular case, may be stated thus;—To divide the sum s between three persons in the proportions of a , b , and c .

The *Formulas* which would be found for the several shares, are

$$\frac{as}{a+b+c}; \quad \frac{bs}{a+b+c}; \quad \frac{cs}{a+b+c}.$$

These Formulas translated into arithmetical language, would furnish a Proposition, or a Rule, which might be applied to any given case of the general problem.

It may be here remarked that algebraic Formulas express general principles and methods of solution with the utmost distinctness and brevity,—but that it is not always possible to translate them into concise and perspicuous phraseology.

EXERCISES.

1. Divide \$950 between two persons so that their shares shall be to each other as 3 to 5.

This problem might be solved without employing *proportion*, by observing that the first share will be $\frac{3}{8}$ of the second.

Ans. \$356 $\frac{1}{4}$; \$593 $\frac{3}{4}$.

2. Find the Formulas for dividing any given sum s between two persons so that the shares shall be to each other as any two numbers a and b .

Ans. $\frac{as}{a+b}$ and $\frac{bs}{a+b}$.

3. Divide the sum of \$3000 between A, B, and C, in the proportions of 1, 2, and 3.

Ans. \$500; \$1000; \$1500.

4. Divide the sum of \$7600 between three persons, in the proportions of $\frac{1}{2}$, $\frac{1}{4}$, and $\frac{1}{8}$.

Ans. \$4000; \$2000; \$1600.

5. A bankrupt is indebted to A \$400, and to B \$700. He is able to pay to both \$900; what sum should each of the two creditors receive?

The \$900 should be divided in the proportion of 400 and 700, or, of 4 and 7, (158).

Ans. \$327 $\frac{3}{11}$; \$572 $\frac{8}{11}$.

6. Three persons engaged in a speculation towards which they contributed, respectively, \$300, \$400, and \$500. The profit amounted to \$550; what are the respective shares of profit?

Ans. \$137 $\frac{1}{2}$; \$183 $\frac{1}{3}$; \$229 $\frac{1}{6}$.

7. A, B, and C in a joint mercantile adventure lost \$742. A's part of the capital employed was to B's as 4 to 3, and B's was to C's as 5 to 6; what amount of loss should be borne by each?

Ans. \$280; \$210; \$252.

8. Four persons rented a pasture, in which the first kept 8 oxen, the second 6, the third 10, and the fourth 12. The sum paid was \$40; what amount should have been paid by each person?

Ans. \$8 $\frac{8}{9}$; \$6 $\frac{2}{3}$; \$11 $\frac{1}{9}$; \$13 $\frac{1}{3}$.

9. A testator bequeathed his estate, amounting to \$7830, to his three children, in such a manner that the share of the first was to that of the second as $2\frac{1}{2}$ to 2, and the share of the second to that of the third as $3\frac{1}{2}$ to 3. What were the shares?

Ans. \$3150; \$2520; \$2160.

10. A, B, C, and D together have \$3000; A's part is to B's as 2 to 3, B and C together have \$1500, and C's part is to D's as 3 to 4. What is the sum possessed by each person?

Ans. \$500; \$750; \$750; \$1000

11. Three persons contributed funds in a joint speculation as follows: A \$200 for 5 months, B \$400 for 3 months, and C \$500 for 4 months. The profit amounted to \$600; what are the several shares of profit?

Each Dollar contributed produced a Profit *proportional to the Time* it was in the business. Each person's share of profit is therefore proportional to his amount of *capital* \times its *time*; in other words, the respective shares are to each other in the *compound ratio* of *capital* and *time*, (131),

Hence, A's share of profit is to B's as 200×5 to 400×3 ;
and A's is to C's as 200×5 to 500×4 .

Dividing the antecedent and consequent by 100, these ratios become 2×5 to 4×3 and 2×5 to 5×4 , (158).

By still further reductions, on the same principle, the ratios become 5 to 6, and 1 to 2.

Ans. $\$142\frac{6}{7}$; $\$171\frac{3}{7}$; $\$285\frac{5}{7}$.

12. Two persons rented a pasture for \$43. The first put into it 100 sheep for 15 days, and the second 120 sheep for 9 days; what amount of rent should be paid by each person?

Ans. \$25 and \$18.

13. A, B, and C trade together; A ventures \$1000 for 5 months, B \$1200 for 4 months, and C \$800 for 7 months. The profits of the partnership amount to \$2310; what share of profit should be assigned to each?

Ans. \$750; \$720; \$840.

14. An estate consisting of 1000 acres of land is to be divided between three persons, so that the first share shall be to the second as 2 to 3, and the first to the third as 1 to 2. What are the shares?

Ans. $222\frac{2}{3}$; $333\frac{1}{3}$; $444\frac{1}{3}$, acres.

15. Two men contracted to do a certain work for \$5000. In accomplishing the work, the first employed 100 laborers for 50 days; and the second 125 laborers for 60 days;—to what shares of the stipulated sum are the two men respectively entitled?

Ans. \$2000; and \$3000.

16. A gentleman bequeathed \$18000 to his widow and his three sons, in the proportions of 2, $2\frac{1}{2}$, 3, and $3\frac{1}{2}$, respectively. His widow dying before the division was effected, the whole is to be divided proportionably among the three sons. What are their several shares?

Ans. \$5000; \$6000; \$7000.

17. Find the Formulas for dividing, between two partners, the profits of a joint adventure, in which the first had the Capital a for the Time b , and the second the Capital c for the Time d .

Ans. $\frac{abs}{ab+cd}$ and $\frac{cbs}{ab+cd}$.

Problems in Percentage.

(171.) *Percentage* is an allowance at a certain rate per *hundred*; and this rate is called the *rate per cent.*, from the Latin *centum*, which means a *hundred*.

The *ratio* of percentage is the *rate per cent.* \div 100, and is therefore equal to the *rate per unit*.

The *basis* of percentage is the sum or number on which an amount of percentage is computed.

From these definitions it follows, that

(172.) The *basis* of percentage \times the *ratio* of percentage produces the *amount* of percentage; and, conversely, that the *amount* \div the *ratio* produces the *basis* of percentage.

☞ The Student may be required to write the *Rules* of Percentage which are indicated by the Formulas among the following problems.

18. A merchant finds that his capital, which is now \$12000, has increased in one year at the rate of 20 per cent.; what was his capital at the beginning of the year?

Let x represent his capital at the beginning of the year;

$$\text{then } 100 : 120 :: x : 12000. \quad \text{Ans. } \$10000.$$

19. What is the Formula for finding a sum of money which, increased at the rate of r per cent., shall amount to the sum a .

$$\text{Ans. } \frac{100a}{100+r}.$$

20. An agent receives \$500 to be laid out in merchandise, after deducting his commission of $1\frac{1}{2}$ per cent. on the amount of the purchase. What will be the amount of the purchase?

$$\text{Ans. } \$492\ 61'.$$

21. A merchant obtains an insurance at 2 per cent. on a stock of goods valued at \$7500, which includes this amount and the *premium* or the insurance. What is the sum insured? *Ans.* \$7653.06'.

22. What is the Formula for finding a sum of money which, diminished at the rate of r per cent., shall leave the sum a ?

$$\text{Ans. } \frac{100a}{100-r}.$$

23. The profits of a manufacturing company this year amount to \$3096, which is $3\frac{1}{4}$ per cent. less than their profits last year. What was the amount of profits last year? *Ans.* \$3200.

24. What must be the percentum of profit at which a quantity of merchandise, bought for \$3750, must be sold, that the whole amount of profit shall be \$1500? *Ans.* 40 per cent.

25. What is the Formula for finding the *rate per cent.* at which the sum s must be increased to produce the sum a ?

$$\text{Ans. } \frac{100(a-s)}{s}.$$

26. A quantity of silk was purchased for \$220, and, on account of its having become damaged, was sold for \$176. What was the per centum of the loss sustained?

Ans. 20 per cent.

27. What is the Formula for finding the *rate per cent.* at which the sum s must be diminished to leave the sum a ?

$$\text{Ans. } \frac{100(s-a)}{s}.$$

28. What amount of stock in an Insurance Office, at a discount of 5 per cent., could be purchased for \$3800?

Ans. \$4000.

29. A merchant finds that his capital, which is now \$4350, has decreased in one year at the rate of $12\frac{1}{2}$ per cent. What was his capital at the beginning of the year?

Ans. \$4971.42'.

30. What amount of stock in a manufacturing establishment, at an advance of $6\frac{1}{2}$ per cent., could be purchased for \$1200?

Ans. \$1126.76'.

31. A quantity of damaged cloth was sold for \$250,—which was at a loss of $16\frac{2}{3}$ per cent. For what sum was the cloth purchased?

Ans. \$300.

32. The population of a city increased from 7850 to 11775 inhabitants, in one year. What was the per centum of increase during the year?

Ans. 50 per cent.

33. An agent receives \$2030 to invest in merchandise—himself to retain a commission of $1\frac{1}{2}$ per cent. on the amount of the purchase. What is the sum to be invested?

Ans. \$2000.

34. A merchant wishes to effect an insurance on a stock of goods, amounting to \$3573, which shall cover both the value of the goods and the premium of insurance. What is the sum to be insured, allowing the rate to be $\frac{3}{4}$ per cent?

Ans. \$3600.

35. What amount of stock in a Savings Bank, at an advance of 5 per cent., could be purchased for \$4200? and what amount in another at a discount of 5 per cent., could be purchased for \$1995?

Ans. \$4000, and \$2100.

Problems in Interest, &c.

(173.) *Interest* is the price or premium paid for the *use of money*, and is reckoned at a certain percentum, *annually*, on the sum for which it is paid.

The *Principal* is the sum for which Interest is paid—the *Amount* is the sum of the Principal and Interest.

From the principles of Percentage, (172), it is evident that,

(174.) The Principal \times the *ratio* of percentage produces the *interest for one year*. For the ratio of percentage, in this case, is the interest of \$1 for one year.

☞ The Student may be required to write the *Rules* of Interest which are indicated by the Formulas among the following problems.

36. What Principal would amount to \$1000 in 5 years, allowing the rate of interest to be 6 per cent ?

Let x represent the Principal required ;

then $x \times \frac{6}{100}$ or $\frac{6x}{100}$ is the Interest for one year ;

and by adding 5 years' interest to the Principal,

$$\text{we have } x + \frac{30x}{100} = \$1000.$$

Ans. $\$769\frac{3}{13}$.

37. What is the Formula for finding the Principal which, at interest at r per cent., would amount to the sum a in t years ?

$$\text{Ans. } \frac{100a}{100 + rt}.$$

38. What Principal would amount to \$2500, in 10 years, allowing the rate of interest to be 7 per cent. ?

Ans. \$1470.588'.

39. At what Rate per cent. must \$1000 be put on interest, to amount to \$1150 in 2 years and 6 months ?

Ans. 6 per cent.

40. What is the Formula for finding the Rate per cent. of interest at which the sum s would amount to the sum a in t years ?

$$\text{Ans. } \frac{100(a-s)}{st}.$$

41. In how many years would \$6000 amount to \$7470, allowing the rate of interest to be 7 per cent ?

Ans. $3\frac{1}{2}$ years

42. What is the Formula for finding the Time in which the sum s would amount to the sum a , if the interest be at r per cent?

$$\text{Ans. } \frac{100(a-s)}{sr}.$$

43. What Principal would produce as much interest in $3\frac{1}{2}$ years, as \$500 would in 4 years, the rate of interest in both cases being 6 per cent?

$$\text{Ans. } \$571\frac{3}{7}.$$

44. At what Rate per cent. of interest would \$525 produce the same amount of interest in 5 years, that \$700 would produce at 5 per cent. in 3 years?

$$\text{Ans. } 4 \text{ per cent.}$$

45. A person who possessed a capital of \$70000, put the greater part of it at interest at 5 per cent., and the other part at 4 per cent. The interest on the whole was \$3250 per annum; required the two parts.

$$\text{Ans. } \$45000, \text{ and } \$25000.$$

46. The sum of \$200 is to be applied in part towards the payment of a debt of \$300, and in part to paying the Interest, at 6 per cent., *in advance*, for 12 months, on the remainder of the debt? What is the amount of the payment that can be made on the debt?

Let x represent the payment;

then $(300-x) \times \frac{6}{100}$ is the Interest on the remainder of the debt; and we have therefore the Equation,

$$x + (300-x) \times \frac{6}{100} = 200.$$

$$\text{Ans. } \$193.61'.$$

47. A is indebted to B \$1000, and is able to raise but \$600. With this sum A proposes to pay a part of the debt, and the Interest, at 8 per cent., in advance, on his Note at 2 years for the remainder. For what sum should the note be drawn?

$$\text{Ans. } \$476.19'.$$

48. Find the Formulas for dividing the sum s into two parts, one of which is to be applied towards the payment of a debt of n dollars, and the other to paying the interest, in advance, on the remainder of the debt, for t years, at r per cent. per annum.

$$\text{Ans. } \frac{100s-nrt}{100-rt}, \text{ and } \frac{rt(n-s)}{100-rt}.$$

What would be the RULE for finding the amount of *payment* that could be made on the debt?

CHAPTER VII.

ARITHMETICAL, HARMONICAL, AND GEOMETRICAL PROGRESSION.

ARITHMETICAL PROGRESSION.

(175.) An ARITHMETICAL PROGRESSION is a series of quantities which continually increase or decrease by a *common difference*.

Thus 1, 3, 5, 7, 9, is a Progression in which the quantities increase by the continual *addition* of the common difference 2.

And 15, 12, 9, 6, 3, is a progression in which the quantities decrease by the continual *subtraction* of the common difference 3.

The *first* and *last terms* of the Progression are called the two *extremes*, and all the intermediate terms the *means*.

The theory of Arithmetical Progression is contained in the following propositions.

The Last Term.

(176.) The *last term* of an *increasing* Arithmetical Progression, is equal to the first term + the product of the common difference \times the number of terms *less one*; and in a *decreasing* Progression it is equal to the first term — the same product.

Let a be the first term, and d the common difference; then in an increasing progression the series will be,

$$a, a+d, a+2d, a+3d, a+4d, \&c.;$$

and in a decreasing progression the series will be,

$$a, a-d, a-2d, a-3d, a-4d, \&c.$$

In these series the *fifth* or *last term* $a \pm 4d$, a plus or minus $4d$, is the first term a plus or minus 4 *times* the common difference d . And the proposition is evidently true for any number of terms.

(177.) *Cor.* The *common difference* of the terms in an Arithmetical Progression, is equal to the difference between the *two extremes* \div the number of terms *less one*.

The Sum of the Two Extremes.

(178.) The *sum of the two extremes* in an Arithmetical Progression, is equal to the sum of any two terms *equidistant from them*, or to twice the middle term when the number of terms is *odd*.

Let a be the first term, and d the common difference; then in an increasing progression the series will be,

$$a, a+d, a+2d, a+3d, a+4d, \&c.$$

Of these five terms the sum of the *first* and the *last*, is

$$a+(a+4d)=2a+4d.$$

The sum of the *second* and the *fourth*, which are equidistant from the extremes, is $(a+d)+(a+3d)$, also $=2a+4d$.

We see moreover, that the sum of the two extremes is equal to *twice the middle term*, $a+2d$.

In like manner the proposition will be found true for any number of terms; as also when the Progression is a *decreasing* one.

(179.) *Cor.* An *arithmetical mean* between two given terms, is equal to *half the sum* of those terms.

For the sum of the two given terms, considered as the two extremes of an Arithmetical Progression, is equal to twice the mean or middle term.

The Sum of all the Terms.

(180.) The *sum of all the terms* of an Arithmetical Progression, is equal to half the sum of the two extremes \times the number of terms.

To prove this proposition we add the several terms of an Arithmetical Progression to those of the same progression *reversed*; thus

$$\begin{array}{cccc} a, & a+d, & a+2d, & a+3d, \\ a+3d & a+2d & a+d & a \\ \hline 2a+3d & 2a+3d & 2a+3d & 2a+3d \end{array}$$

The sum $(2a+3d)+(2a+3d)$ &c. of the *two series*, is the sum of the two extremes in either series \times the number of terms; hence the sum of either series is equal to *half the sum* of the two extremes \times the number of terms. The demonstration will evidently apply to any number of terms.

FORMULAS IN ARITHMETICAL PROGRESSION.

(181.) In an Arithmetical Progression, let a be the first term, d the common difference, n the number of terms, l the last term, and S the sum of all the terms. Then

$$A \dots l = a \pm d(n-1) \quad (176);$$

$$B \dots S = \frac{1}{2}n(a+l) \quad (180).$$

The sign $+$ is to be prefixed to $d(n-1)$ when the progression is an increasing one, and $-$ when decreasing.

In these two Formulas we have the *five quantities*, a, d, n, l, s ; hence if any *three* of these quantities be given, the values of the other *two* may be found from the two Equations, (121).

HARMONICAL PROGRESSION.

(182.) An HARMONICAL PROGRESSION is a series of quantities such that, of any *three consecutive terms*, the first : the *third* :: the *difference* between the first and second : the *difference* between the second and third.

Thus the numbers 3, 4, 6, 12, are in harmonical progression,

$$\text{since } 3 : 6 :: 4-3 : 6-4,$$

$$\text{and } 4 : 12 :: 6-4 : 12-6.$$

An Harmonical *Proportion* consists of four terms such that the first *is to* the fourth *as* the difference between the first and second *is to* the difference between the third and fourth.

Thus a, b, c, d , are in Harmonical Proportion, if

$$a : d :: a-b : c-d;$$

$$\text{or } a : d :: b-a : d-c.$$

The numbers 16, 8, 3, 2, are in Harmonical Proportion, since

$$16 : 2 :: 16-8 : 3-2 .$$

The *first* and *last* terms of the Progression or proportion are called the two *extremes*, and all the intermediate terms the *means*.

An Harmonical converted into an Arithmetical Progression.

(183.) The *reciprocals* of the terms of an Harmonical progression are in Arithmetical progression.

Let a, b, c , be three consecutive terms of a decreasing harmonical progression.

$$\text{Then } a : c :: a - b : b - c, \quad (182);$$

Converting this proportion into an equation we have

$$ab - ac = ac - bc.$$

Dividing each term in this equation by abc , and reducing the several quotients to their *lowest terms*, we find

$$\frac{1}{c} - \frac{1}{b} = \frac{1}{b} - \frac{1}{a}.$$

Transposing the first and the last term of this equation

$$\frac{1}{a} - \frac{1}{c} = \frac{1}{b} - \frac{1}{b}.$$

We thus find that the difference between the reciprocals of a and c , is equal to the difference between the reciprocals of b and b ; hence these reciprocals are in Arithmetical Progression, (175).

The numbers 3, 4, 6, 12, form an Harmonical Progression, (182): by taking the *reciprocals* of the several terms we have the Arithmetical Progression

$$\frac{1}{3}, \frac{1}{4}, \frac{1}{6}, \frac{1}{12};$$

in which the common *difference* of the terms is $\frac{1}{12}$.

(184.) An *harmonical mean* between two given terms, is equal to *twice their product* divided by their *sum*.

From the first of the preceding equations, namely,

$$ab - ac = ac - bc,$$

$$\text{we shall find } b = \frac{2ac}{a+c};$$

and b is the harmonical mean between a and c .

The harmonical mean between 3 and 6, is

$$\frac{3 \times 6 \times 2}{3+6} = \frac{36}{9} = 4.$$

GEOMETRICAL PROGRESSION.

(185.) A GEOMETRICAL PROGRESSION is a series of quantities in which each succeeding term has the *same ratio* to the term which immediately precedes it.

Thus 1, 2, 4, 8, 16, is an increasing Progression in which each succeeding term is *double* the one which immediately precedes it; that is, the *ratio of the progression* is 2.

And 27, 9, 3, 1, $\frac{1}{3}$, is a decreasing progression in which each succeeding term is *one-third* of the one which immediately precedes it; and the ratio of the progression is consequently $\frac{1}{3}$.

Hence the *successive terms* of a Geometrical Progression consist of the *first term multiplied* continually into the *ratio*, that is, multiplied into the *successive powers of the ratio*.

The theory of Geometrical Progression is contained in the following propositions.

The Last Term.

(186.) The *last term* of a Geometrical Progression, is equal to the first term \times that *power of the ratio* which is expressed by the number of terms *less one*.

Let a be the first term, and r the *ratio* of the progression; then, multiplying a continually into r , the series will be

$$a, ar, ar^2, ar^3, ar^4, \&c.$$

Since the ratio r begins in the second term, with *exponent 1*, its exponent in the last term will always be *one less* than the number of terms; hence the last term consists of the first \times into that power of r which is expressed by the number of terms *minus 1*.

(187.) *Cor.* The *last term* of a Geometrical Progression \div the first term, gives that *power of the ratio* which is expressed by the number of terms *less one*.

Thus $ar^4 \div a = r^4$; the number of terms being *five*.

Product of the two Extremes.

(188.) The *product of the two extremes* in a Geometrical Progression, is equal to the product of any two terms *equidistant from them*, or to the *square of the middle term* when the number of terms is *odd*.

Let a be the first term, and r the ratio of the progression; then, multiplying a into the successive powers of r , the series is

$$a, ar, ar^2, ar^3, ar^4, \&c.$$

Of these five terms the product of the *first* and the *last*, is

$$a \times ar^4 = a^2 r^4.$$

The product of the *second* and the *fourth*, which are equidistant from the extremes, is $ar \times ar^3$, also $= a^2 r^4$.

We perceive moreover that the product of the two extremes is equal to the *square of the middle term* ar^2 .

In like manner the proposition will be found true for any number of terms.

(189.) *Cor.* A *geometrical mean*, or a mean proportional, between two given terms, is equal to the *square root of the product* of those terms.

For the product of the two given terms, considered as the two extremes of a Geometrical Progression, is equal to the square of the mean or middle term.

The Sum of all the Terms.

(190.) The *sum of all the terms* of a Geometrical Progression, is equal to the difference between the *first term* and the *product* of the *last term* \times the *ratio*, \div the difference between the *ratio* and a *unit*.

Let S represent the sum of the terms, and we shall have

$$S = a + ar + ar^2 + ar^3 + ar^4, \&c.$$

Multiplying both sides of this equation by the ratio r ,

$$Sr = ar + ar^2 + ar^3 + ar^4 + ar^5.$$

Subtracting the first equation from the second, we find

$$Sr - S = ar^5 - a;$$

$$\text{which gives } S = \frac{ar^5 - a}{r - 1}.$$

In the numerator of this value of S , observe that ar^5 is the last term ar^4 of the progression \times the ratio r . The demonstration will evidently apply to any number of terms.

(191.) The sum of an *infinite number of terms* in a decreasing Geometrical Progression, is equal to the *first term* divided by the difference between the *ratio* and a *unit*.

In a decreasing progression the terms continually diminish in a *constant ratio*; and if the number of terms be *infinite*, the last term will be 0. The last term \times the *ratio* will then be 0, and the expression for the sum of the terms, found above, will become

$$S = \frac{a}{1-r}.$$

The divisor in this case is $1-r$, because the ratio of the progression being a *proper fraction*, is less than a *unit*.

FORMULAS IN GEOMETRICAL PROGRESSION.

(192.) In a Geometrical Progression, let a be the first term, r the ratio, n the number of terms, l the last term, and s the sum of all the terms. Then

$$C \dots l = ar^{n-1}, \quad (186),$$

$$D \dots S = \frac{lr-a}{r-1}, \quad (190).$$

When the progression is a decreasing one, and the number of terms is infinite,

$$E \dots S = \frac{a}{1-r}, \quad (191).$$

In the Formulas C and D we have the five quantities, a , r , n , l , s ; hence if any *three* of these quantities be given, the values of the other *two* may be found from the two Equations, (121).

The principles which have been established in this Chapter may be applied to the solution of the following

Problems in Progressions.

1. The first term of an increasing Arithmetical progression is 3, the common difference of the terms is 2, and the number of terms 20. What is the *last term*? and the sum of all the terms?

Ans. 41, and 440.

2. The first term of a decreasing Arithmetical progression is 100, the common difference of the terms is 3, and the number of terms 34. What is the last term? and the sum of all the terms?

Ans. 1, and 1717.

3. What is the sum of the numbers 1, 2, 3, 4, 5, &c., continued to 1000 terms? *Ans.* 500500.

4. What is the *common difference* of the terms in an Arithmetical progression whose first term is 10, last term 150, and number of terms 21? *Ans.* 7.

5. If the *third* term of an Arithmetical progression be 40, and the *fifth* term 70, what will the *fourth* term be? *Ans.* 55.

6. If the first term of an Arithmetical progression be 5, and the fifth term 30, what will the *second*, *third*, and *fourth* terms be?

Find the *common difference*, (177), and thence the three intermediate terms. *Ans.* $11\frac{1}{4}$; $17\frac{1}{2}$; $23\frac{3}{4}$.

7. If the fourth term of an Arithmetical progression be 37, and the eighth term 60, what are the intermediate terms?

Ans. $42\frac{2}{3}$; $48\frac{1}{2}$, $54\frac{1}{3}$.

8. What is the sum of 25 terms of an increasing Arithmetical progression in which the first term is $\frac{1}{2}$, and the common difference of the terms also $\frac{1}{2}$? (176). *Ans.* $162\frac{1}{2}$.

9. The first term of an increasing Arithmetical progression, is 1, and the number of terms 23. What must be the *common difference*, that the sum of all the terms may be 100?

Let x represent the common difference;

then $1+22x$ is the last term, (176);

and $\frac{2+22x}{2} \times 23$ is the sum of the terms, (180).

Hence an Equation may be formed from which the value of x will be found.—Or we might substitute the numbers 1, 23, and 100, for a , n , and s in Formulas A and B, (181), and find the value of d , as one of the *two unknown quantities*. *Ans.* $\frac{7}{23}$.

10. If the first term of a decreasing Arithmetical progression is 100, and the number of terms 21, what must the common difference be, that the sum of the series may be 1260? *Ans.* 4.

11. A and B start together, and travel in the same direction; A goes 40 miles per day; B goes 20 miles the first day, and increases his rate of travel $\frac{3}{4}$ of a mile per day. How far will they be apart at the end of 40 days? *Ans.* 215 miles.

12. One Hundred stones being placed on the ground in a straight line, at the distance of 2 yards from each other; how far will a person travel who shall bring them, one by one, to a basket which is placed 2 yards from the first stone? *Ans.* 11 miles 840 yards.

13. Find the *third term* of an Harmonical progression whose first and second terms are 12 and 15 respectively.

If x represent the third term, we shall have

$$12 : x :: 15 - 12 : x - 15. \quad \text{Ans. } 20.$$

14. What is the first term of an Harmonical progression whose second and third terms are 30 and 20 respectively? *Ans.* 60.

15. What is the *fourth term* of an Harmonical *proportion* whose first, second, and third terms are 2, 3, and 8 respectively? *Ans.* 16.

16. If the first and third terms of an Harmonical progression be 25 and 40 respectively, what will the second term be? *Ans.* $30\frac{10}{13}$.

17. The first and fourth terms of an Harmonical progression, are 10 and 20 respectively. What are the two intermediate terms?

This problem may be solved by finding two *arithmetical means* between $\frac{1}{10}$ and $\frac{1}{20}$, and then taking the *reciprocals* of the terms thus found, (183). *Ans.* 12, and 15.

18. The fifth and eighth terms of an Harmonical progression are 20 and 40 respectively. What are the two intermediate terms? *Ans.* 24, and 30.

19. The first term of a Geometrical progression is 2, the ratio of the progression is 3, and the number of terms 4. What is the *last term*? and the sum of all the terms? *Ans.* 54, and 80.

20. The first term of a Geometrical progression is $\frac{1}{2}$, the ratio of the progression is $\frac{1}{3}$, and the number of terms 4. What is the last term? and the sum of all the terms? *Ans.* $\frac{1}{54}$, and $\frac{20}{27}$.

21. What is the sum of an infinite number of terms in the Geometrical progression whose first term is 100, and ratio $\frac{1}{4}$? *Ans.* $133\frac{1}{3}$.

22. What is the sum of an infinite number of terms in the Geometrical progression whose first term is 300, and ratio $\frac{1}{3}$? *Ans.* 450.

23. If the first and third terms of a Geometrical progression are 8 and 72 respectively, what is the *second term*?

The second term is equal to the *square root* of 8×72 , (189).

Or, considering the third as the *last term* of the progression, $72 \div 8 = 9$ is the *square of the ratio*, (187); then 3 is the *ratio of the progression*; and the second term is now readily obtained. *Ans.* 24.

24. If the third and fifth terms of a Geometrical progression be 75 and 300 respectively, what will the fourth term be? *Ans.* 150

25. If the first and fourth terms of a Geometrical progression are 3 and 24 respectively, what are the two intermediate terms?

Ans. 6 and 12.

26. If the seventh and tenth terms of a Geometrical progression are 6 and 750 respectively, what are the intermediate terms?

Ans. 30 and 150.

27. What is the sum of an infinite number of terms in the series $1, \frac{1}{2}, \frac{1}{4}, \&c.$, in which the ratio of the progression is evidently $\frac{1}{2}$?

Ans. 2.

28. If a body move forever at the rate of 2000 feet the first second 1000 the second, 500 the third, and so on, what is the utmost distance it can reach?

Ans. 4000.

29. If 10 yards of cloth be sold at the rate of \$1 for the first yard \$2 for the second, \$4 for the third, and so on, what would be the price of the last yard? and what would the whole amount to?

Ans. \$512, and \$1023.

30. If 13 acres of land were purchased at the rate of \$2 for the first acre, \$6 for the second, \$18 for the third, and so on, what would the last acre amount to?

Ans. \$1062882.

31. Allowing the interest of a sum of money to be \$500 the first year, \$400 the second, \$320 the third, and so on, forever, what would be the whole amount of interest?

Ans. \$2500.

32. Two bodies move at the same time, from the same point, in opposite directions. One goes 2 miles the first hour, 4 the second, 6 the third, and so on; the other goes 2 miles the first hour, 4 the second, 8 the third, &c.; how far will they be apart at the end of 12 hours?

Ans. 8346 miles.

33. A and B set out at the same time to meet each other. A travels 3, 4, 5, &c. miles on successive days, and B 3, $4\frac{1}{2}$, $6\frac{3}{4}$, &c. miles on successive days. They meet in 10 days; what is the distance between the two places from which they traveled?

Ans. $414\frac{5}{9}\frac{7}{2}$ miles.

CHAPTER VIII.

PERMUTATIONS AND COMBINATIONS.—INVOLUTION.—BINOMIAL THEOREM.—EVOLUTION.

PERMUTATIONS.

(193.) PERMUTATIONS are the different *orders of succession* in which a given number of things may be taken—either the *whole number together*, or the whole number taken *two and two*, or *three and three*, &c.

Thus the different Permutations of the three letters *a*, *b*, and *c*, when *all are taken together*, are

abc, acb, bac, cab, bca, cba.

And the different Permutations of the same letters when taken *two and two*, are

ab, ba, ac, ca, bc, cb.

Number of Permutations.

(194.) If *n* represent a given number of things, the number of *permutations* that can be formed of them, will be equal to

$n(n-1)(n-2)(n-3)(n-4)$, and so on,

until the number of *factors* multiplied together is equal to the number of things taken in *each permutation*.

To demonstrate this proposition,—suppose that we have *n* letters, *a*, *b*, *c*, *d*, &c., to be subjected to Permutations.

If we reserve one of the letters, as *a*, there will remain *n-1* letters; and by writing *a* before each of the remaining letters, we have *n-1* permutations of *n* letters, taken *two and two*, in which *a* stands first.

In like manner we should find *n-1* permutations of *n* letters, taken *two and two*, in which *b* stands first; and so for each of the *n* letters. Hence we shall have

n(n-1) permutations of n letters taken two and two.

Suppose now that the *n* letters are to be taken *three and three*.

By reserving *a*, and proceeding with *n-1* letters as before, we should find $(n-1)(n-2)$ permutations of *n-1* letters taken *two and two*; and by writing *a* before each of these permutations, we have

$(n-1)(n-2)$ permutations of n letters, taken *three* and *three*, in which a stands first.

We should find the same number of permutations of n letters, taken *three* and *three*, in which b stands first; and so for each of the n letters. Hence we shall have

$n(n-1)(n-2)$ permutations of n letters taken *three* and *three*.

Suppose now that the n letters are to be taken *four* and *four*.

By reserving a , and pursuing the operation in the same manner as before, we should find

$n(n-1)(n-2)(n-3)$ permutations of n letters taken *four* and *four*.

Thus the demonstration proceeds; the number of *factors* multiplied together being found always equal to the number of letters taken in *each permutation*.

As an Example of the application of the principle above demonstrated,—suppose it were required to determine the number of Permutations, or different *orders of succession*, that could be formed in a class composed of *six* pupils, by taking the *whole number* in each permutation.

Since the *six* pupils are to be taken in each Permutation, the number of *factors* to be employed is *six*; hence the number of permutations is

$$6 \times 5 \times 4 \times 3 \times 2 \times 1 = 720.$$

If the whole *six* were to be subjected to Permutations by taking *five* at a time, the number of permutations would be

$$6 \times 5 \times 4 \times 3 \times 2 = 720.$$

If the whole *six* were to be subjected to Permutations by taking *four* at a time, the number of permutations would be

$$6 \times 5 \times 4 \times 3 = 360.$$

☞ It will be observed above, that the number of Permutations will be the same, whether the whole number of things, or one less than the whole number, be taken in each permutation.

COMBINATIONS.

(195.) COMBINATIONS are the different *collections* which may be formed out of a given number of things, by taking the same number in each collection—without regard to the order of succession.

Thus the different Combinations which may be formed out of the three letters a , b , and c , by taking two at a time, are

$$ab, bc, ac.$$

Observe that ab and ba are not different *combinations*, but different *permutations*, of the letters a and b .

In Permutations we have regard to the *order of succession*, and may therefore have two permutations of *two things*. In Combinations we do not consider the order of succession; so that the combination of two or more things is the same, in whatever order they are taken.

Number of Combinations.

(196.) If n represent a given number of things, the number of *combinations* that can be formed out of them, will be equal to

$$\frac{n(n-1)(n-2)(n-3)}{1.2.3.4}, \text{ and so on,}$$

until the number of *factors* in the dividend, and also in the divisor, is equal to the number of things taken in *each combination*.

To demonstrate this proposition, we observe that the *numerator* in the preceding expression, is the number of *permutations* of n things taken *four and four*, (194).

On the same principle, the *denominator* $4 \times 3 \times 2 \times 1$ is the number of permutations of *four things taken all together*.

Now since there can be but *one combination* of 4 things taken *all together*, the number of *permutations* of n things taken *four and four* is $1.2.3.4$ times, that is, 24 times, the number of *combinations*.

Hence the number of Combinations of n things taken *four and four*, is $\frac{1}{24}$ of the number of Permutations; and will therefore be found by dividing the numerator by the denominator.

This mode of demonstration may evidently be applied whatever be the number of things taken in each combination.

PROBLEMS

In Permutations and Combinations.

1. In how many different ways might a company of 10 persons seat themselves around a table? (194). *Ans.* 3628800.

2. How many different numbers might be expressed by the 10 numeral figures, if 5 figures be used in each number? *Ans.* 30240.

3. In how many different ways may the names of the 12 months of the year be arranged one after another? *Ans.* 479001600.

4. How many different permutations of 8 men could be formed out of a company consisting of 15 men? *Ans.* 259459200.

5. In how many different ways might the seven prismatic colors, *red, orange, yellow, green, blue, indigo, and violet*, have been arranged in the solar spectrum? *Ans.* 5040.

6. How many different combinations of two colors could be formed out of the 7 prismatic colors? (195). *Ans.* 21.

7. How many different combinations of 5 letters may be formed out of the 26 letters of the Alphabet? *Ans.* 65780.

8. How many different combinations of 2 elements might be formed out of the 56 elements described in Chemistry? *Ans.* 1540.

9. In how many different ways might a company of 20 men be arranged, in single file, in a procession?

Ans. 2432902008176640000.

10. A farmer wishes to select a team of 6 horses out of a drove containing 10 horses. How many different choices for the team will he be able to make? *Ans.* 210.

11. In how many different ways might the planets Mercury, Venus, the Earth, Mars, Jupiter, Saturn, Uranus, and Neptune succeed one another in the solar system? *Ans.* 40320.

12. A company of 20 persons engaged to remain together so long as they might be able to combine in different couples in their evening walks. What time will be required to fulfil the engagement?

Ans. 190 days.

13. How many different permutations of 7 letters might be formed out of the 26 letters of the Alphabet? *Ans.* 3315312000.

14. In an exhibition of a Public School, 5 speakers are to be taken from a class of 15 students. How many different selections of the five might be made? and in how many different ways might the 5 succeed one another in the delivery of their speeches?

Ans. 3003 and 120.

15. Out of a Company consisting of 100 soldiers Six are to be taken for a particular service. How many different selections of the 6 might be made? and in how many different ways might the 6 chosen be disposed with regard to the *order of succession*?

Ans. 1192052400; and 720

INVOLUTION.

(197.) INVOLUTION consists in raising a given quantity to any required *power*. This may always be effected by multiplying the quantity into itself as many times *less one* as there are units in the *exponent* of the power.

Thus aa is a^2 , the second *power* or *square* of a ;
 aaa is a^3 , the third *power* or *cube* of a ; and so on.

Observe that one multiplication of a into itself produces the *second* power of a ; two multiplications produce the *third* power, and so on ; also that the number of times the quantity becomes a *factor* in raising a Power, is equal to the *exponent* of the Power.

(198.) A *higher power* of a quantity may also be found by multiplying together two or more lower powers (of the same quantity) the *sum of whose exponents* is equal to the exponent of the required power.

Thus $a^2 \times a^2$ produces a^4 ; $a^2 \times a^3$ produces a^5 , &c.

Powers of Unity, Monomials, Fractions, &c.

(199.) Every power of *unity* is *unity*, since any number of 1s multiplied together produce only 1 ; thus $1 \times 1 \times 1$ &c. = 1.

(200.) A *monomial* is raised to any required Power by raising its numerical *coefficient* to that power, and multiplying the *exponents* of its other factors by the exponent of the *power*.

Thus to find the *third* power of $4ax^2$, we raise 4 to its *third* power, which is $4 \times 4 \times 4 = 64$, and multiply the exponents of a and x by 3 : we thus obtain $64a^3x^6$.

Observe that multiplying the given *exponents* by the exponent of the required Power, is only a brief method of performing the requisite multiplications in the Involution of Monomials.

(201.) A *fraction* is raised to any required Power, by raising its numerator and denominator, separately, to that power.

Thus the *third* power of $\frac{a^2}{2x}$ is $\frac{a^6}{8x^3}$. (98).

A *mixed quantity* may be raised to any required Power by involving the equivalent *improper fraction*.

The Sign to be Prefixed to a Power.

(202.) Every *even power* of a quantity is *positive*; while every *odd power* has the same sign as the *quantity from which it is derived*.

Thus if a be *positive*, all its powers, as aa , aaa , and so on, will evidently be *positive*; but if a be *negative* we shall have

$$-a. -a = a^2; a^2. -a = -a^3; -a^3. -a = a^4; a^4. -a = -a^5, \&c.$$

from which it is plain that all the *even powers*, as the 2d, 4th, and so on, will be *positive*, while all the *odd powers* will be *negative*.

EXERCISES.

On the Powers of Monomials.

- | | |
|---|-----------------------------|
| 1. Find the square of $3ax^2$. | Ans. $9a^2x^4$ |
| 2. Find the cube of $-2a^2x$. | Ans. $-8a^6x^3$ |
| 3. Find the square of $-4ax^2$. | Ans. $16a^2x^4$ |
| 4. Find the cube of $3a^3x$. | Ans. $27a^9x^3$ |
| 5. Find the square of ab^2c^3 . | Ans. $a^2b^4c^6$ |
| 6. Find the cube of $-a^2x^2y$. | Ans. $-a^6x^6y^3$ |
| 7. Find the square of $\frac{1}{2}ax^2$. | Ans. $\frac{1}{4}a^2x^4$ |
| 8. Find the cube of $\frac{1}{2}ay^3$. | Ans. $\frac{1}{8}a^3y^9$ |
| 9. Find the square of $-\frac{2}{3}ab^n$. | Ans. $\frac{4}{9}a^2b^{2n}$ |
| 10. Find the cube of $-\frac{2}{3}ax^2$. | Ans. $-\frac{8}{27}a^3x^6$ |
| 11. Find the 4th power of $2a^n$. | Ans. $16a^{4n}$ |
| 12. Find the 4th power of $-\frac{1}{2}a^2$. | Ans. $\frac{1}{16}a^8$ |
| 13. Find the 4th power of $-3x^2$. | Ans. $81x^8$ |
| 14. Find the 5th power of $\frac{1}{2}x^2$. | Ans. $\frac{1}{32}x^{10}$ |
| 15. Find the 6th power of $-ax^2$. | Ans. a^6x^{12} |
| 16. Find the 6th power of $2y^2$. | Ans. $64y^{12}$ |
| 17. Find the 7th power of $-a^2y^n$. | Ans. $-a^{14}y^{7n}$ |
| 18. Find the 7th power of $\frac{1}{2}y$. | Ans. $\frac{1}{128}y^7$ |
| 19. Find the 8th power of a^2b^n . | Ans. $a^{16}b^{8n}$ |
| 20. Find the 8th power of $-\frac{1}{2}x^2$. | Ans. $\frac{1}{256}x^{16}$ |

POWERS OF POLYNOMIALS.

(203.) The Powers of a *binomial*, or of any *polynomial*, may be obtained by successive multiplications of the quantity into itself, (197).

Thus the 2d power or *square* of $a+b$, is

$$(a+b)(a+b)=a^2+2ab+b^2.$$

And the 3d power or *cube* of $a+b$, is

$$(a+b)(a+b)(a+b)=a^3+3a^2b+3ab^2+b^3.$$

The Involution of Binomials, however, and thence of Polynomials, is greatly facilitated by the application of NEWTON'S

BINOMIAL THEOREM.

(204.) This Theorem explains a general method of *developing* a Binomial according to any *exponent* with which the Binomial may be affected.

In developing $(a+b)^2$ by the Binomial Theorem we should obtain the 2d *power* or *square* of $(a+b)$.

In developing $(a+b)^{\frac{1}{2}}$ we should obtain the *square root* of $(a+b)$; and so for other exponents.

Development of $(a \pm b)^n$;

n representing any exponent.

1. The *Exponent* of a in the *first term* of the development, is the same as the *exponent of the Binomial*, and *decreases* by the subtraction of 1, continually, in the succeeding terms.

2. The *Exponent* of b entering as a *factor*, commences with 1 in the *second term* of the development, and *increases* by the addition of 1, continually, in the succeeding terms.

3. The *Coefficient* of the *first term* is 1; that of the *second term* is the same as the *exponent of the Binomial*;—and the coefficient of the second, or of any term, \times the exponent of a in that term, and \div by the *exponent of b* increased by 1, gives the coefficient of the succeeding term.

4. The *Signs* of the terms in the development of $(a+b)$ will all be $+$; while for $(a-b)$ they will be alternately $+$ and $-$.

From these principles we have the following

(205.) *Formula for the Development of the Binomial* $(a+b)^n$.

$$(a+b)^n = a^n + na^{n-1}b + \frac{n(n-1)}{2}a^{n-2}b^2 + \frac{n(n-1)(n-2)}{2 \times 3}a^{n-3}b^3, \text{ \&c}$$

When the exponent n is a *positive integer*, as 2, 3, or 4, &c., the development will *terminate* at the term in which the exponent of b becomes equal to the exponent of the Binomial.

For the exponent of a in that term will be 0, and 0 will thus become a *factor* in finding the *coefficient* of the next term; hence the next term will be 0, (43).

EXAMPLE I.

To find the 4th *power* of $(a+b)$, by the Binomial Theorem; that is, to *develop* $(a+b)^4$.

The *literal* factors without the *coefficients*, will be

$$a^4 \quad a^3b \quad a^2b^2 \quad ab^3 \quad b^4$$

By *computing* and introducing the *coefficients*, with the *signs*, we have

$$a^4 + 4a^3b + 6a^2b^2 + 4ab^3 + b^4.$$

The student will readily perceive the application of the preceding principles to the several terms of this Power.

Demonstration of the Binomial Theorem.

The principles which have been given for the development of $(a \pm b)^n$, will be demonstrated under the supposition that the exponent n is a *positive integer*.—A general demonstration would be equally applicable to *negative* or *fractional* exponents; such demonstration is unnecessary here, and is too abstruse for the present stage of our subject.

If we multiply together the *binomial factors*
 $a+b, a+c, a+d, a+e,$
 and decompose the product terms which contain the *lower powers* of a , the final Product may be represented as follows;

$$\begin{array}{cccc}
 a^4 & +b & | & a^3 & +bc & | & a^2 & +bcd & | & a & +bcde \\
 & +c & | & & +bd & | & & +bce & | & & \\
 & +d & | & & +be & | & & +bde & | & & \\
 & +e & | & & +cd & | & & +cde & | & & \\
 & & & & +ce & | & & & & & \\
 & & & & +de & | & & & & &
 \end{array}$$

In this Product observe that the *coefficients* of the powers of a in the successive terms, are as follows ;

The Coefficient of a^4 is 1 ; the coefficient of a^3 is the sum of the second terms, $b, c, \&c.$, of the binomial factors composing the Product ; the coefficient of a^2 is the sum of the products $bc, bd, \&c.$, of the second terms of the binomial factors *combined* two and two, (196) ; and the coefficient of a is the sum of the products $bcd, bce, \&c.$, of the second terms of the binomial factors *combined* three and three.

Suppose now that $c, d, e, \&c.$ are each equal to b , and the number of binomials equal to n . The Product of these factors will be the n th power of $(a+b)$; that is, it will be
the *development* of $(a+b)^n$.

The Exponent of a in the first term of the Product will evidently be n ; and, as exemplified in the preceding multiplication, this exponent will decrease by 1, continually, in the succeeding terms.

Hence we shall have

$$a^n, a^{n-1}, a^{n-2}, \&c. \text{ in the consecutive terms.}$$

The Coefficient of a^n will evidently be 1. The coefficient of a^{n-1} in the second term, will be n times b .

Hence the second term will be

$$nba^{n-1} \text{ or } na^{n-1}b.$$

The Coefficient of a^{n-2} in the third term, will be b^2 taken *as many times* as there are combinations of *two* letters in n letters, (196).

Hence the third term will be

$$\frac{n(n-1)}{1 \times 2} b^2 a^{n-2} \text{ or } \frac{n(n-1)}{1 \times 2} a^{n-2} b^2.$$

The Coefficient of a^{n-3} in the fourth term, will be b^3 taken *as many times* as there are combinations of *three* letters in n letters.

Hence the fourth term will be

$$\frac{n(n-1)(n-2)}{1 \times 2 \times 3} b^3 a^{n-3} \text{ or } \frac{n(n-1)(n-2)}{1 \times 2 \times 3} a^{n-3} b^3.$$

The several terms thus obtained agree with *Formula* (205) ; and the *law of development* thus indicated, will, in like manner, be found applicable to any number of terms.

With regard to the *Signs*,—it is evident that when a and b are *positive*, all the terms in any power of $(a+b)$ will be *positive*, (42).

When b is *negative*, all the *odd* powers of b will be *negative*, (202) ; and these negative powers multiplied by the positive powers of a , will cause the 2d, 4th, 6th, &c. terms to become *negative*.

EXAMPLE II.

To find the 5th power of the binomial $a-x$.

The literal factors without the *coefficients* are

$$a^5, a^4x, a^3x^2, a^2x^3, ax^4, x^5.$$

By computing and inserting the coefficients, with the *signs*, we have

$$a^5 - 5a^4x + 10a^3x^2 - 10a^2x^3 + 5ax^4 - x^5.$$

EXERCISES

On the Powers of Polynomials.

1. Find the square of $a-x$. *Ans.* $a^2 - 2ax + x^2$.
2. Find the cube of $a+y$. *Ans.* $a^3 + 3a^2y + 3ay^2 + y^3$.
3. Find the cube of $a+2b$.

By applying the principles of the Binomial Theorem, we obtain

$$a^3 + 3a^2 \cdot 2b + 3a(2b)^2 + (2b)^3;$$

which may be developed into *Ans.* $a^3 + 6a^2b + 12ab^2 + 8b^3$.

4. Find the square of $3a+y$. *Ans.* $9a^2 + 6ay + y^2$.
5. Find the cube of $2a-3x$. *Ans.* $8a^3 - 36a^2x + 54ax^2 - 27x^3$.
6. Find the square of $a+x-y$,

By operating on $(x-y)$ as if it were a *monomial*, and applying the Binomial Theorem to $a+(x-y)$, we obtain

$$a^2 + 2a(x-y) + (x-y)^2;$$

which may be developed into *Ans.* $a^2 + 2ax - 2ay + x^2 - 2xy + y^2$.

7. Find the square of $a-b+y$. *Ans.* $a^2 - 2ab + 2ay + b^2 - 2by + y^2$.
8. Find the square of $a-x-y$. *Ans.* $a^2 - 2ax + x^2 - 2ay + 2xy + y^2$.
9. Find the square of $a+b-2x$. *Ans.* $a^2 + 2ab + b^2 - 4ax - 4bx + 4x^2$.
10. Find the square of a^2+x^3 .

This square, according to the Binomial Theorem, may be indicated thus :

$$(a^2)^2 + 2a^2x^3 + (x^3)^2;$$

which may be developed into *Ans.* $a^4 + 2a^2x^3 + x^6$

11. Find the cube of a^2-y^3 . *Ans.* $a^6 - 3a^4y^3 + 3a^2y^6 - y^9$
12. Find the cube of $1+2x^2$. *Ans.* $1 + 6x^2 + 12x^4 + 8x^6$.
13. Find the fourth power of $a+x$. *Ans.* $a^4 + 4a^3x + 6a^2x^2 + 4ax^3 + x^4$
14. Find the fifth power of $a-y$. *Ans.* $a^5 - 5a^4y + 10a^3y^2 - 10a^2y^3 + 5ay^4 - y^5$

EVOLUTION.

(206.) EVOLUTION consists in extracting any required *root* of a given quantity, regarded as the corresponding power of the root to be found.

Extracting the square root consists in finding a quantity whose square is equal to a given quantity; extracting the *cube root* consists in finding a quantity whose cube is equal to a given quantity; and so on.

Roots of Unity, Monomials, Fractions, &c.

(207.) Every root of *unity* is *unity*, since the *square*, or *cube*, &c. of 1 is 1; thus $1 \times 1 = 1$; $1 \times 1 \times 1 = 1$; and so on.

In general terms, any *power* or *root* of a unit is a unit.

(208.) Any required Root of a *monomial* will be found by extracting the root of its numerical *coefficient*, and dividing the *exponents* of its literal factors by the *integer* corresponding to the root.

Thus the *square root* of $25 a^2 x$ is $5ax^{\frac{1}{2}}$, found by extracting the square root of 25, and dividing the exponents of a and x by 2.

And the *cube root* of $27a^6x^{\frac{1}{2}}$ is $3a^2x^{\frac{1}{6}}$, found by extracting the cube root of 27, and dividing the exponents of a and x by 3.

The correctness of this method will appear from considering that the Extracting of a Root is the reverse of raising the corresponding Power, (200).

Hence also the propriety of denoting roots by *fractional* exponents.

$a^{\frac{1}{2}}$ denotes the square root of a , because $a^{\frac{1}{2}} \times a^{\frac{1}{2}} = a$.

(209.) Any required Root of a *fraction* will be found by extracting the root of its numerator and denominator, separately.

Thus the square root of $\frac{4x^2}{9a^2y^4}$ is $\frac{2x}{3ay^2}$, found by extracting the square root of the numerator, and the square root of the denominator.

A Root of a *mixed quantity* would be found by extracting the root of the equivalent *improper fraction*.

(210.) A Root whose exponent is *resolvable into two factors*, may be found by extracting *in succession* the roots denoted by those factors.

The 4th root may be found by extracting the *square root* of the *square root*,—the exponent $\frac{1}{4}$ which denotes the 4th root, being $\frac{1}{2} \times \frac{1}{2}$.

Thus the 4th root of x^4 is the square root of x^2 ; the 4th root of 81 is the square root of 9; and the 4th root of 10000 is the square root of 100.

The 6th root may be found by extracting the square root of the *cube root*, or the cube root of the square root,— $\frac{1}{6}$ being equal to $\frac{1}{2} \times \frac{1}{3}$.

The correctness of this method of extracting Roots, is evident from considering, that, by raising in succession the *powers* denoted by two or more factors, we shall obtain the power denoted by the *product* of those factors.

Thus the *square* of the *cube* of a is $(a^3)^2 = a^6$, (200), and therefore, conversely, the *square root* of the *cube root* is the *sixth* root

Roots of Powers or Powers of Roots.

(211.) The Numerator of a fractional exponent denotes a *power* of the quantity affected, and the Denominator a *root of that power*.

Thus $a^{\frac{1}{2}}$ denotes the square root of the *first power* of a , or simply the square root of a . In like manner $a^{\frac{2}{3}}$ denotes the *cube root* of the square of a ; $a^{\frac{3}{4}}$ denotes the 4th root of the cube of a ; and so on. But, observe that

(212.) A Root of any *power* of a quantity, is equal to the same *power* of the same *root* of the quantity.

To illustrate this principle it may be shown that the *cube root* of the 6th *power* of a , is equal to the 6th *power* of the cube root of a .

The *cube root* of a^6 is a^2 since $a^2 \cdot a^2 \cdot a^2 = a^6$; and the 6th *power* of $a^{\frac{1}{3}}$ is also a^2 , since the *cube* of $a^{\frac{1}{3}}$ is a , and the *square* of *this cube*,—which gives the 6th *power* of $a^{\frac{1}{3}}$,—is a^2 , (200).

To give an application of this principle to *numbers*,—the cube root of the square of 8, is 4, and the square of the cube root of 8 is 4

Equivalent Exponents.

(213.) Two Exponents which are numerically equivalent, are also equivalent *exponentially*; and may therefore be substituted, the *one for the other*.

Thus $a^{\frac{2}{4}} = a^{\frac{1}{2}}$; the *fourth root* of a^2 is equivalent to the *square root* of a . The 4th root of a^2 is $a^{\frac{1}{2}}$, since $a^{\frac{1}{2}}$ raised to the 4th *power* produces a^2 .

In like manner $x^{\frac{3}{6}} = x^{\frac{1}{2}}$; $y^{\frac{4}{6}} = y^{\frac{2}{3}}$; $a^{\frac{4}{2}} = a^2$; &c.

To give an example in *numbers*,—the *fourth root* of the square of 9, that is, of 81, is 3, and the square root of 9 is also 3.

On this principle a fractional exponent may always be taken in its *lowest terms*.

The Sign to be Prefixed to a Root.

(214.) Every *odd Root* of a quantity has the same sign as the *quantity itself*.

For the quantity itself is an *odd power* with reference to such *root*, (206), and an *odd power* has the same sign as the quantity of which it is a power, (202).

(215.) Every *even Root* of a *positive* quantity is *ambiguous*, that is, the quantity itself does not determine whether the root is *positive* or *negative*.

This follows from the principle that every *even power* of a *negative*, as well as of a *positive* quantity, is *positive*, (202).

The square of $-a$, as well as of $+a$, is $+a^2$; hence the square root of a^2 is $+a$ or else $-a$; and when it is not known whether a^2 was derived from a or $-a$, it is uncertain which of these two is the square root.

This uncertainty as to the *sign* of the root, is expressed by the *ambiguous sign* \pm *plus* or *minus*; thus

the square root of a^2 is $\pm a$.

For a like reason the *fourth root* of a^4 is $\pm a$.

The square root of 9 is ± 3 ; the fourth root of 16 is ± 2 ; &c.

(216.) An *even Root* of a *negative* quantity is *impossible*, since there is no quantity which can be raised to an *even negative power*, (202).

Thus no quantity multiplied into itself will produce $-a^2$; this quantity therefore has *no square root*. For a like reason $-a^4$ has no square root or *fourth root*.

In like manner -9 has no square root; -16 has no square root or fourth root; $-25a^2$ has no square root, &c.

EXERCISES

On the Roots of Monomials.

- | | |
|---|---|
| 1. Find the square root of $4a^2x^4$. | <i>Ans.</i> $\pm 2ax^2$. |
| 2. Find the cube root of $8a^3y^6$. | <i>Ans.</i> $2ay^2$. |
| 3. Find the square root of $9a^4x$. | <i>Ans.</i> $\pm 3a^2x^{\frac{1}{2}}$. |
| 4. Find the cube root of $-27y^2$. | <i>Ans.</i> $-3y^{\frac{2}{3}}$. |
| 5. Find the square root of $16x^{\frac{1}{2}}$. | <i>Ans.</i> $\pm 4x^{\frac{1}{4}}$. |
| 6. Find the cube root of a^3xy^2 . | <i>Ans.</i> $ax^{\frac{1}{3}}y^{\frac{2}{3}}$. |
| 7. Find the square root of $25a^3$. | <i>Ans.</i> $\pm 5a^{\frac{3}{2}}$. |
| 8. Find the cube root of $-64y^2$. | <i>Ans.</i> $-4y^{\frac{2}{3}}$. |
| 9. Find the square root of $36x^6$. | <i>Ans.</i> $\pm 6x^3$. |
| 10. Find the cube root of $8ay^n$. | <i>Ans.</i> $2a^{\frac{1}{3}}y^{\frac{n}{3}}$. |
| 11. Find the square root of $\frac{1}{4}a^4y^2$. | <i>Ans.</i> $\pm \frac{1}{2}a^2y$. |
| 12. Find the cube root of $-8a^3x^{3n}$. | <i>Ans.</i> $-2ax^n$. |
| 13. Find the square root of $\frac{4}{9}xy^4$. | <i>Ans.</i> $\pm \frac{2}{3}x^{\frac{1}{2}}y^2$. |
| 14. Find the cube root of $-\frac{1}{8}ax^3$. | <i>Ans.</i> $-\frac{1}{2}a^{\frac{1}{3}}x$. |
| 15. Find the fourth root of $16a^4y^2$. | <i>Ans.</i> $\pm 2ay^{\frac{1}{2}}$. |
| 16. Find the cube root of $-\frac{8}{27}x^2$. | <i>Ans.</i> $-\frac{2}{3}x^{\frac{2}{3}}$. |
| 17. Find the fourth root of $\frac{1}{16}ay^2$. | <i>Ans.</i> $\pm \frac{1}{2}a^{\frac{1}{4}}y^{\frac{1}{2}}$. |
| 18. Find the fifth root of $-a^{10}x^2y$. | <i>Ans.</i> $-a^2x^{\frac{2}{5}}y^{\frac{1}{5}}$. |
| 19. Find the sixth root of a^3x^2y . | <i>Ans.</i> $\pm a^{\frac{1}{2}}x^{\frac{1}{3}}y^{\frac{1}{6}}$. |
| 20. Find the sixth root of ba^2y^{3n} . | <i>Ans.</i> $\pm b^{\frac{1}{6}}a^{\frac{1}{3}}y^{\frac{n}{2}}$. |

ROOTS OF POLYNOMIALS.

(217.) The method of extracting any required Root of a Polynomial, may be discovered from the manner in which the corresponding *power* of a polynomial is formed. This subject will be elucidated under the appropriate Rules.

RULE XVIII.

(218.) *To Extract the Square Root of a Polynomial.*

1. Arrange the Polynomial according to the powers of one of its letters, and take the square root of the *left hand term*, for the first term of the root.

2. Subtract the square of the root thus found from the given Polynomial; divide the remainder by *twice the root* already found, and annex the *quotient* to both the *root* and the *divisor*.

3. Multiply the divisor thus formed by the *last term in the root*; subtract the product from the dividend; divide the remainder by *twice the root* now found; and so on, as before.

EXAMPLE.

To extract the Square Root of

$$a^2 + 2ab + b^2.$$

The required Root, we already know, is $a + b$, since the *square* of this binomial is the given trinomial:—our object is to show that the root $a + b$ would be found by the foregoing Rule.

$$\begin{array}{r} a^2 + 2ab + b^2 \quad (a + b \\ a^2 \\ \hline 2a + b \quad 2ab + b^2 \\ \quad 2ab + b^2 \\ \hline \end{array}$$

The first term a of the root, is the square root of a^2 , the left hand term of the given polynomial. Subtracting the square of a , we have the remainder $2ab + b^2$.

We have now to discover a *divisor* of this remainder, which will give, for a quotient, the next term b of the root.

$2a$, that is, *twice the root* already found, divided into $2ab$, gives b ; and b annexed to $2a$ makes the divisor $2a + b$. This divisor multiplied by b , produces $2ab + b^2$,—which completes the operation.

The Rule is framed in accordance with the process thus discovered.

EXERCISES

On the Square Root of Polynomials.

1. Find the square root of the polynomial
 $a^4 - 4a^3 + 6a^2 - 4a + 1.$ *Ans.* $a^2 - 2a + 1.$
2. Find the square root of the polynomial
 $a^4 + 4a^3x + 6a^2x^2 + 4ax^3 + x^4.$ *Ans.* $a^2 + 2ax + x^2.$
3. Find the square root of the polynomial
 $1 - 6y + 13y^2 - 12y^3 + 4y^4.$ *Ans.* $1 - 3y + 2y^2.$
4. Find the square root of the polynomial
 $4x^6 - 4x^4 + 12x^3 + x^2 - 6x + 9.$ *Ans.* $2x^3 - x + 3.$
5. Find the square root of the polynomial
 $4a^4 + 12a^3x + 13a^2x^2 + 6ax^3 + x^4.$ *Ans.* $2a^2 + 3ax + x^2.$
6. Find the square root of the polynomial
 $1 - 6a + 13a^2 - 12a^3 + 4a^4.$ *Ans.* $1 - 3a + 2a^2.$
7. Find the square root of the polynomial
 $x^4 + 2ax^3 + 3a^2x^2 + 2a^3x + a^4.$ *Ans.* $x^2 + ax + a^2.$
8. Find the square root of the polynomial
 $9a^4 - 12a^3y + 28a^2y^2 - 16ay^3 + 16y^4.$ *Ans.* $3a^2 - 2ay + 4y^2.$
9. Find the square root of the polynomial
 $25 + 34x^2 + 12x^3 + 20x + 9x^4.$ *Ans.* $5 + 2x + 3x^2.$
10. Find the square root of the polynomial
 $a^4 + 4a^3y - 12ay + 4a^2y^2 + 9 - 6a^2.$ *Ans.* $a^2 + 2ay - 3.$
11. Find the square root of the polynomial
 $4 + 24x^2 - 16x + 4x^4 - 16x^3.$ *Ans.* $2 - 4x + 2x^2.$
12. Find the square root of the polynomial
 $1 - 2y + 7y^2 - 2y^3 + 5y^4 + 12y^5 + 4y^6.$ *Ans.* $1 - y + 3y^2 + 2y^3.$

SQUARE ROOT OF NUMBERS.

The preceding Rule is substantially the same as the Arithmetical Rule for extracting the square root of Numbers. To show its application, however, to the latter purpose, we must premise the following principles.

(219.) If a Number be separated into *periods of two figures* each,—from right to left,—these periods will correspond, respectively, to the *units, tens, hundreds, &c.*, in the Square Root of the number.

For since the square of *ten* is 100, the square of the *tens figure* in the root will leave *two vacant places* in the right of the given number; these two places must therefore correspond to the *units* in the root.

And since the square of a *hundred* is 10000, the square of the *hundreds* in the root leaves *four vacant places* in the right of the number; and the first two corresponding to the *units*, the *next two* must correspond to the *tens* in the root.

In like manner it is shown that the *third period* of two figures corresponds to the *hundreds* in the square root of the given number; and so on.

(220.) If a Number be divided into any two parts, the Square of the number will be equal to the *square of the first part + twice the first \times the second + the square of the second part.*

For, *a* representing the *first part* of the number, and *b* the *second*, *a + b* will be equal to the number; and

$$(a+b)^2 = a^2 + 2ab + b^2.$$

With these principles established we may proceed to the following

EXAMPLES.

1. To extract the Square Root of 529

$$\begin{array}{r} 5'29 \quad (\quad 23 \\ 4 \\ \hline 43) \overline{129} \\ \underline{129} \\ \hline \end{array}$$

The first period 29 corresponds to the *units figure*, and the 5 to the *tens figure* in the root, (219).

The greatest *integral square root* of the 5 is 2 *tens*, the square of which is 4 *hundreds*, and this subtracted leaves 129.

Since the root of the given number consists of two parts, *tens* and *units*, and the *square of the 2 tens* has been subtracted, the remainder
 $129 = \text{twice } 2 \text{ tens} \times \text{the units} + \text{the units}^2, (220).$

Doubling the root, 2 tens, already found, we take the product, 4 tens, for a divisor. The 0 in 4 tens, or 40, may be omitted in finding the next figure in the root, if we omit the 9 in the corresponding place in the 129.

We therefore say 4 in 12, 3 times. This 3 annexed to the 4 makes the 4 now become 4 tens. Then 43, or 4 tens + 3 units, \times 3 units, produces the remainder 129; and thus completes the operation.

2. To extract the Square Root of 36 84 49

$$\begin{array}{r} 36'84'49 \quad (\quad 607 \\ 36 \\ \hline 1207 \) \quad 84 \ 49 \\ \quad \quad 84 \ 49 \\ \hline \end{array}$$

Pointing off periods of two figures, from right to left, we find that the root will contain *units*, *tens*, and *hundreds*, (219).

The square root of the left hand period 36, is 6, the square of which subtracted leaves no remainder in that period.

We take the next period 84 to find the next figure in the root. Doubling the root 6, for a divisor, and omitting the 4, we say 12 in 8, 3 time.

Including the next period 49, and doubling the root 60 now found, for a divisor, we say 120 in 844, 7 times.

Annexing 7 to 120, the divisor becomes 1207, which multiplied by 7 produces 8449; and thus the operation is completed.

Square Root of Decimals.

(221.) In extracting the Square Root of a Decimal Fraction, the *periods* must be taken from the decimal point *towards the right*; and a 0 must be annexed, if necessary, to complete the last period.

The last period must be complete, because, by the principles of decimal multiplication, the square of a decimal Fraction must contain *twice as many* decimal figures as are in the root.

The number of decimal figures to be made in the root, is therefore the same as the number of *decimal periods*.

When an exact root cannot be found, decimal periods of 00 each may be annexed, and the root continued in *decimals* to any required exactness.

EXERCISES

On the Square Root of Numbers.

1. Find the square root of 11236, and of $\frac{256}{289}$. *Ans.* 106 ; and $\frac{16}{17}$.
2. Find the square root of 38809, and of $\frac{1156}{5041}$. *Ans.* 197 ; and $\frac{34}{71}$.
3. Find the square root of 75076, and of $\frac{6561}{9025}$. *Ans.* 274 ; and $\frac{81}{95}$.
4. Find the square root of 22801, and of $\frac{5929}{7056}$. *Ans.* 151 ; and $\frac{77}{84}$.
5. Find the square root of .582 169, and of $127\frac{13}{25}$.

Reduce the $\frac{13}{25}$ to an equivalent *decimal*, before proceeding with the evolution.

Ans. .763 ; and 11.292'.

6. Find the square root of 475.125, and of $346\frac{17}{30}$. *Ans.* 21.797' ; and 18.616'.
7. Find the square root of 37.4780, and of $470\frac{21}{100}$. *Ans.* 6.121' ; and 21.684'

RULE XIX.

(222.) *To Extract the cube Root of a Polynomial.*

1. Arrange the Polynomial according to the powers of one of its letters, and take the cube root of the *left hand term*, for the first term of the root.

2. Subtract the cube of the root thus found from the given Polynomial, and divide the remainder by 3 *times the square* of the root already found, and annex the *quotient* to the root.

3. *Complete the divisor* by adding to it the product which arises from annexing the last term in the root to 3 times the *other part* of the root, and *multiplying the result* by the last term.

4. Multiply the completed divisor by the *last term* in the root, and subtract the product from the dividend.

5. Find the next *incomplete divisor* by adding to the last complete divisor the *product which completed it*, and the *square of the last term* in the root ; divide, and complete the divisor, as before ; and so on.

EXAMPLE.

To extract the Cube Root of

$$a^3 + 3a^2b + 3ab^2 + b^3.$$

The required Root, we already know, is $a+b$, (203), and our object is to show that this root would be found by the Rule.

$$\begin{array}{r} a^3 + 3a^2b + 3ab^2 + b^3 \quad (a + b \\ a^3 \\ \hline 3a^2 + 3ab + b^2 \quad) \quad \begin{array}{l} 3a^2b + 3ab^2 + b^3 \\ \underline{3a^2b + 3ab^2 + b^3} \end{array} \end{array}$$

The first term a of the root, is the cube root of a^3 , the left hand term of the given polynomial. Subtracting a^3 , we have the remainder $3a^2b + 3ab^2 + b^3$.

We have now to find a *divisor* of this remainder, which will give, for a quotient, the next term b of the root.

$3a^2$, that is, 3 *times the square* of the root already found, divided into $3a^2b$, gives b . Adding $3ab$, and also b^2 , to $3a^2$, the completed divisor is $3a^2 + 3ab + b^2$.

This divisor multiplied by b , produces the last dividend, and thus shows that the operation is completed.

EXERCISES

On the Cube Root of Polynomials.

- Find the cube root of the polynomial
 $a^6 - 6a^5 + 15a^4 - 20a^3 + 15a^2 - 6a + 1.$ *Ans.* $a^2 - 2a + 1.$
- Find the cube root of the polynomial
 $1 - 6x + 21x^2 - 44x^3 + 63x^4 - 54x^5 + 27x^6.$ *Ans.* $1 - 2x + 3x^2.$
- Find the cube root of the polynomial
 $8 - 12x + 30x^2 - 25x^3 + 30x^4 - 12x^5 + 8x^6.$ *Ans.* $2 - x + 2x^2.$
- Find the cube root of the polynomial
 $8 + 36y^2 + 24y + 32y^3 + 6y^5 + y^6 + 18y^4.$ *Ans.* $2 + 2y + y^2$
- Find the cube root of the polynomial
 $9a^4 - 3a^5 - 13a^3 + a^6 - 12a + 8 + 18a^2.$ *Ans.* $a^2 - a + 2.$
- Find the cube root of the polynomial
 $21x^4 - 44x^3 - 54x + 63x^2 - 6x^5 + x^6 + 27.$ *Ans.* $x^2 - 2x + 3.$
- Find the cube root of the polynomial
 $y^6 - 9y^5 + 39y^4 - 99y^3 + 156y^2 - 144y + 64.$ *Ans.* $y^2 - 3y + 4$

CUBE ROOT OF NUMBERS.

The method of extracting the Cube Root of Numbers involves the following principles.

(223.) If a Number be separated into *periods of three figures* each,—from right to left,—these periods will correspond, respectively, to the *units, tens, hundreds, &c.*, in the Cube Root of the number.

For since the cube of *ten* is 1000, the cube of the *tens figure* in the root, will leave three vacant places in the right of the given number; these three places must therefore correspond to the *units* in the root.

And since the cube of *a hundred* is 1000000, the cube of the *hundreds* in the root leaves *six vacant* places in the right of the number; and the first three corresponding to the *units*, the *next three* must correspond to the *tens* in the root.

In like manner it is shown that the *third period* of three figures corresponds to the *hundreds* in the cube root of the given number, and so on.

(224.) If a Number be divided into any two parts, the Cube of the number will be equal to the *cube of the first part* + 3 *times the square of the first* × the *second* + 3 *times the first* × the *square of the second* + the *cube of the second*.

For *a* representing the *first part* of the number, and *b* the *second*, *a + b* will be equal to the number; and

$$(a + b)^3 = a^3 + 3a^2b + 3ab^2 + b^3.$$

As an example of the application of this principle to numbers, let 275 be divided into the two parts 200 and 75; then

$$(200 + 75)^3 = 200^3 + 3 \times 200^2 \times 75 + 3 \times 200 \times 75^2 + 75^3.$$

If the values of the several terms on the right be computed, and added together, we shall obtain the cube of 275.

We might now proceed to extract the Cube Root of Numbers in a manner corresponding to the Rule already given for the cube root of Polynomials; but the method by the following Rule is more direct and simple.

RULE XX.

(225.) *To Extract the Cube Root of a Number.*

1. Separate the number into *periods of three figures* each, from right to left, observing that the last period will sometimes have but one or two figures.

2. Take the greatest integral cube root of the left hand period, for the first figure of the root required,—subtract *its cube* from said period, —and to the remainder affix the next period for a *dividend*.

3. Take *3 times the square* of the root already found, for an incomplete divisor; divide it into the dividend *exclusive of its two right hand figures*, and annex the quotient to the root.

4. *Complete the divisor* by annexing to it 00, and adding the product which arises from annexing the last figure in the root to *3 times the other part* of the root, and multiplying the *result by the last figure*.

5. Multiply the completed divisor by the last figure in the root; subtract the product from the dividend; and to the remainder affix the next period for a new dividend.

6. Find the *next incomplete divisor* by adding to the last complete divisor the *product which completed it*, and the *square of the last figure* in the root; divide, and complete the divisor, as before; and so on.

In applying this Rule it will be convenient to have the following

Table of Roots and Cubes.

Roots, 1, 2, 3, 4, 5, 6, 7, 8, 9.

Cubes, 1, 8, 27, 64, 125, 216, 343, 512, 729.

We will first apply, and then demonstrate this Rule.

EXAMPLE.

To extract the Cube Root of 95 44 39 93.

	95'44 3'9 93 (457
	64
48	31 44 3
4800 + 625 = 54 25	27 12 5
5425 + 625 + 25 = 60 75	43 1 8 993
607500 + 9499 = 6169 99	43 1 8 993

The greatest integral cube root of the left hand period 95, is 4, the *cube* of which subtracted leaves the remainder 31. Affixing the next period, we obtain

the dividend 31443.

Three times the square of 4, the root already found, is 48. We divide 48 into 314, excluding the 43. The quotient would appear to be 6; but the divisor being as yet *too small*, we take 5 for the quotient.

To complete the divisor, we annex 00 to it, and add 625;—the 625 being obtained by annexing the 5 in the root to 3 times the 4, and multiplying the result 125 by 5.

Multiplying the completed divisor 5425 by 5, subtracting, and affixing the next period to the remainder, we find

the dividend 43 18 99 3.

For the next incomplete divisor, we add to the last complete divisor the product 625 which completed it, and the square 25 of the 5 in the root. Dividing by 6075, the quotient is 7.

To complete this divisor, we annex 00, and add 9499,—this last number being obtained by annexing the 7 to 3 times the 45, and multiplying the result 1357 by 7. The divisor now found multiplied by 7 produces the last dividend.

Demonstration of the Rule for the Cube Root of Numbers.

To facilitate the Demonstration, we will take a number containing but *two periods*,—whose root will therefore consist of *tens* and *units*, (223).

$$\begin{array}{r}
 91'125 \text{ (} 45 \\
 \hline
 64 \\
 \hline
 27125 \\
 \hline
 27125 \\
 \hline
 4800 + 625 = 5125
 \end{array}$$

The 4 in the root, found according to the Rule, we know is 4 *tens*, (223); and the second figure in the root will be *units*.

Let t represent the *tens*, and u the *units* of the root, then the given number $= t^3 + 3t^2u + 3tu^2 + u^3$, (224).

Hence the cube 64,—which is 64 *thousand*,—of the 4 *tens*, subtracted, leaves $27125 = 3t^2u + 3tu^2 + u^3$.

By the Rule for the cube root of a Polynomial, the *incomplete divisor* of this remainder—to be used for finding the next figure or term u of the root, is $3t^2 = 3$ times $(4 \text{ tens})^2 = 4800$.

In dividing we omit the 00, and at the same time exclude the 25 in the corresponding places of the dividend 27125.

By the Rule just referred to, the quantity to be added to complete the divisor $3t^2$, is

$$(3t+u)u=(3 \times 4 \text{ tens} + 5) \times 5 = 625;$$

that is, the product which arises from annexing the last figure in the root to 3 *times the other part* of the root, and *multiplying the result* by the last figure.

Suppose now that the given number contains *three periods*, and its root, consequently, three figures.

Regarding the two figures already found as constituting the *first part* of the root, the next *incomplete* divisor, by Rule XIX, would be

$$3t^2 + (3t+u)u + (3t+u)u + u^2;$$

which consists, first, of the last complete divisor $3t^2 + (3t+u)u$; secondly, of the product $(3t+u)u$ which was added to $3t^2$ to complete that divisor; thirdly, of the square of the last figure u in the root.

In like manner the Rule may be shown to be applicable, whatever be the number of figures in the required root.

Cube Root of Decimals.

(226.) In extracting the Cube Root of a Decimal Fraction, the *periods* must be taken from the decimal point *towards the right*, and 0 or 00 must be annexed, if necessary, to complete the last period.

The last period must be complete, because, by the principles of decimal multiplication, the cube of a decimal Fraction must contain 3 *times as many* decimal figures as are in the root.

The number of decimal figures to be made in the root, is therefore the same as the number of *decimal periods*.

When an exact root cannot be found, decimal periods of 000 each may be annexed, and the root *continued in decimals* to any required exactness.

EXERCISES

On the Cube Root of Numbers.

1. Find the cube root of 262144, and of $\frac{19683}{389017}$.
Ans. 64; and $\frac{27}{73}$.
2. Find the cube root of 2406104, and of $\frac{24389}{438976}$.
Ans. 134; and $\frac{29}{76}$.
3. Find the cube root of 22906304, and of $\frac{29791}{681472}$.
Ans. 284; and $\frac{31}{88}$.
4. Find the cube root of 479.2735, and of $8377\frac{23}{150}$.
Ans. 7.825'; and 20.309'.
5. Find the cube root of 5371.3745, and of $3059\frac{31}{200}$.
Ans. 17.513'; and 14.51'
6. Find the cube root of 403.73331, and of $.71200\frac{40}{123}$.
Ans. 7.390'; and .892'.
7. Find the cube root of 4370.666, and of $\frac{4913}{5832}$.
Ans. 16.34'; and $\frac{17}{18}$.
8. Find the cube root of 20796875, and of $3511\frac{101}{125}$.
Ans. 275; and 15.20.
9. Find the cube root of .202262003, and of $\frac{2744}{6859}$.
Ans. .587; and $\frac{14}{19}$.
10. Find the cube root of 103823, and of $2460\frac{3}{8}$.
Ans. 47; and 13.5.

EXTRACTION OF THE *n*th ROOT.

(227.) Any Root whatever of a Polynomial might be extracted,—by taking the root of its left hand term,—with this root forming an *incomplete* divisor,—with the quotient term, and the root already found, *completing* the divisor,—and so on, in a manner depending on the *order of the root* to be extracted.

But this method, which is preferable for the *square* and *cube*, becomes too complicated when applied to the higher roots.

By dispensing with the completed divisors, the operation may be simplified, and the process of Evolution *generalized*, as follows.

RULE XXI.

(228.) *To Extract any Root of a Polynomial.*

1. Understanding the *order of the root* to be denoted by n ,—arrange the Polynomial according to the powers of one of its letters, and take the n th root of its left hand term, for the first term of the root.

2. Subtract the n th power of the root found from the given Polynomial; and divide the remainder by n times the $(n-1)$ power of this root, for the second term of the root.

3. Subtract the n th power of the root now found from the given Polynomial, and, using the same divisor as before, proceed in the same manner till the n th power of the root becomes equal to the given Polynomial.

This Rule may also be applied to Numbers, by taking n figures in each period, from right to left, for *integers*, and from the decimal point towards the right, for *decimals*; and there will be less liability to error in finding the *quotient figure*, if new *divisors* be found for the second and subsequent remainders.

EXAMPLE.

To extract the 4th root of 30 49 800 625.

$$\begin{array}{r}
 30'4980'0625 \quad (\quad 235 \\
 \underline{2^4 = 16} \\
 4 \times 2^3 = 32 \quad) \quad 144980 \quad . \quad . \quad . \quad \text{Exclude } 980 \text{ in dividing.} \\
 \underline{23^4 = 279841} \\
 4 \times 23^3 = 48668 \quad) \quad 251390625 \quad . \quad . \quad \text{Exclude } 625 \text{ in dividing.} \\
 \underline{235^4 = 3049800625}
 \end{array}$$

*The preceding root might also be found by two extractions of the *square root*, (210); thus

The square root of 3049800625 is 55225; and the square root of the latter number is 235.

CHAPTER IX.

IRRATIONAL OR SURD QUANTITIES.—IMAGINARY QUANTITIES

Perfect and Imperfect Powers.

(229.) A Perfect Power, of any degree, is a quantity which has an exact *root* of the same degree;—otherwise, the quantity is called an *Imperfect Power*.

Thus 4 and $9a^2$ are perfect *squares*, having the exact square *roots* 2 and $3a$; while $2a$ and $8x^3$ are *imperfect squares*, since they have no exact square roots.

In like manner 8 and $27a^3$ are perfect *cubes*; while 9 and $25x^2$ are imperfect cubes, since they have no exact cube roots.

The Polynomial $a^2 + 2ab + b^2$ is a perfect square, whose root is $a + b$; (218); and $a^3 - 3a^2x + 3ax^2 - x^3$ is a perfect cube, whose root is $a - x$.

IRRATIONAL OR SURD QUANTITIES.

(230.) A *Rational* quantity is one which can be accurately expressed without any *indicated root*; as 2, $3a$, or $\frac{2}{3}x$.

An *Irrational* or *Surd* quantity is one which can be accurately expressed only under the form of a *root*,—being the indicated *root* of an *imperfect power*.

Thus $2^{\frac{1}{2}}$ is an *irrational* quantity, since the exact square root of 2 cannot be determined. Also $a^{\frac{1}{2}}$ and $x^{\frac{1}{3}}$ are irrational quantities.

By the common Rule for the square root of numbers, we should find the square root of $2 = 1.414213'$; but other decimal figures would succeed without end.

The term *irrational* when applied to a quantity, implies that such quantity has no determinable *ratio* to unity.

Irrational or Surd quantities—being expressed under the form of *roots*—are also called *Radical* quantities, from the Latin *Radix*, a *root*.

Radical Sign.

(231.) The *radical sign* $\sqrt{\quad}$ prefixed to a quantity, denotes the *square root* of the quantity; and is therefore equivalent to the *exponent* $\frac{1}{2}$.

Thus \sqrt{a} is the square root of a ; equivalent to $a^{\frac{1}{2}}$.

With the *index* 3 affixed to it, this sign denotes the *cube root*, and is then equivalent to the *exponent* $\frac{1}{3}$. With the index 4, it denotes the 4th root; and so on.

Thus $\sqrt[3]{a}$ is equivalent to $a^{\frac{1}{3}}$; and $\sqrt[4]{a}$ to $a^{\frac{1}{4}}$.

The radical sign may always be thus superseded by a *fractional exponent*; and this should be done whenever any obscurity would arise in calculation from using this sign.

For example, in extracting the square root of $\sqrt[3]{a}$, we substitute the exponent $\frac{1}{3}$, and find the required root to be

$$a^{\frac{1}{3} \div 2} = a^{\frac{1}{6}}; \text{ (208).}$$

A quantity preceding the $\sqrt{\quad}$, without + or - interposed, is a *coefficient* or multiplier of the Surd; and when no coefficient is expressed, a *unit* is understood.

Thus $5\sqrt{a}$ is 5 times the square root of a ; and \sqrt{ax} is the same as $1\sqrt{ax}$.

Similar and Dissimilar Surds.

(232.) *Similar Surds* are such as express the *same root* of the same quantity; otherwise, the Surds are *dissimilar*.

Thus $\sqrt{3}$ and $5\sqrt{3}$ are *similar*; while $\sqrt[3]{3}$ and $5\sqrt{3}$ are *dissimilar*. So $(a+b)^{\frac{1}{2}}$ and $3\sqrt{a+b}$ are similar Surds.

A Rational Quantity under the Form of a Surd.

(233.) A rational quantity may be expressed under the form of the *square root*, or *cube root*, &c., by placing the corresponding *power* of the quantity under the *exponent* or *sign* of the root.

To express $3a$ under the form of the *square root*, we place the square of $3a$, which is $9a^2$, under the *exponent* or *sign* of the square root, thus

$$3a = (9a^2)^{\frac{1}{2}} \text{ or } \sqrt{9a^2}.$$

TRANSFORMATION OR REDUCTION OF SURDS.

Certain Transformations of radical quantities are sometimes necessary, to adapt them to the purposes of calculation. These depend chiefly on the two following principles.

(234.) When two or more Factors have the same *exponent*, that exponent may be transferred to the *product* of those factors; and,

Conversely, the exponent of a Product may be transferred to *each of the factors* which compose that product.

Thus a^2x^2 is equal to $(ax)^2$; the product of the *squares* of a and x , is equal to the square of the *product* of a and x .

And $a^{\frac{1}{2}}x^{\frac{1}{2}}$ is equal to $(ax)^{\frac{1}{2}}$; the product of the *square roots* of a and x , is equal to the square root of the *product* of a and x .

These principles result from the methods of finding the *powers* and *roots* of quantities; thus by squaring ax ,—and extracting the square root of ax ,—we have

$$(ax)^2 = a^2x^2, (200); \text{ and } (ax)^{\frac{1}{2}} = a^{\frac{1}{2}}x^{\frac{1}{2}}, (208).$$

These methods of illustration may be applied to any other *power* or *root* as well as to the square and square root. Thus the product of the *cube roots* of two or more factors is equal to the cube root of the *product* of those factors; and so for any power or root.

To give examples in numbers;—

$$\sqrt{4} \times \sqrt{25} = \sqrt{100}; \text{ or } 2 \times 5 = 10.$$

$$\sqrt[3]{8} \times \sqrt[3]{27} = \sqrt[3]{216}; \text{ or } 2 \times 3 = 6.$$

(235.) The Exponent of a quantity may be changed to any *equivalent expression*, without altering the value of the power or root denoted.

This proposition is substantially the same as one already demonstrated, (213).

As examples, $a^{\frac{1}{2}} = a^{\frac{2}{4}} = a^{\frac{3}{6}} = a^{\frac{4}{8}};$

$$4^{\frac{1}{2}} = 4^{\frac{2}{4}} = 16^{\frac{1}{4}} = 2.$$

The following are the principal Transformations required in the calculation of Irrational or Surd quantities.

Product of a Rational and an Irrational Factor.

(236.) A *rational* multiplier or *coefficient* may be put under the form of the Surd annexed, and the product of the two radicals be then affected with the same fractional exponent or radical sign.

Thus $a\sqrt{3} = \sqrt{a^2}\sqrt{3} = \sqrt{3a^2}$, by putting the coefficient a under the form of the *square root*, (233), and transferring the sign $\sqrt{}$, which is equivalent to the *exponent* $\frac{1}{2}$, from the two factors to the product $3a^2$, (234).

In like manner $2\sqrt{5} = \sqrt{4} \times \sqrt{5} = \sqrt{20}$.

As an example of the *utility* of this Transformation,—suppose it were required to find an *approximate value* of $2\sqrt{5}$.

By the common method of extracting the square root, we might find an approximate value of $\sqrt{5}$, and such value multiplied by 2 would be an approximate value of $2\sqrt{5}$.

It is evident, however, that this process would *double*, in the product, the *deficiency* in the value found of $\sqrt{5}$; and the error thus arising will be greater as the coefficient of the Surd is greater.

By taking $\sqrt{20}$, instead of $2\sqrt{5}$, the approximate value will be found independently of the preceding *source of error*.

Surds Reduced to Simpler Forms.

(237.) A Surd is *simplified* by resolving the quantity under the $\sqrt{}$, or exponent, into two factors—one of which is a *perfect power* of like degree—and putting the extracted *root* of this factor as a coefficient to the indicated root of the other.

Thus $\sqrt{50} = \sqrt{25} \times \sqrt{2} = 5\sqrt{2}$;

or, $50^{\frac{1}{2}} = 25^{\frac{1}{2}} \times 2^{\frac{1}{2}} = 5 \times 2^{\frac{1}{2}}$, (234).

In like manner, $\sqrt{4a^2 + 8a^3} = \sqrt{4a^2} \sqrt{1 + 2a} = 2a\sqrt{1 + 2a}$.

In these examples the given Surds are *square roots*, and the factors 25 and $4a^2$ are accordingly *perfect squares*.

As an example of the *cube root*, we have

$\sqrt[3]{54} = \sqrt[3]{27} \times \sqrt[3]{2} = 3\sqrt[3]{2}$; 27 being a *perfect cube*.

When the given Surd is preceded by a Coefficient, this coefficient must be multiplied into the one found by the process of simplifying.

Thus $3\sqrt{32a^3} = 3\sqrt{16a^2} \sqrt{2a} = 3 \times 4a \sqrt{2a} = 12a\sqrt{2a}$.

If, in the preceding process, the square or cube factor be the *greatest* square or cube thus entering into the composition of the quantity under the $\sqrt{\quad}$, the Surd will be reduced to its *simplest form*.

Thus in each of the examples above given, the Surd is reduced to its simplest form. The second example would also give

$$\sqrt{4a^2 + 8a^3} = \sqrt{4} \sqrt{a^2 + 2a^3} = 2\sqrt{a^2 + 2a^3};$$

which is not the simplest form, since a^2 is still under the $\sqrt{\quad}$.

(238.) A Fractional Surd may be reduced to an *integral surd*, by multiplying both its terms, if necessary, to make the denominator a *perfect power*, and then resolving into factors, and proceeding as before, (237).

For example, $\sqrt{\frac{3}{8}} = \sqrt{\frac{6}{16}} = \sqrt{\frac{1}{16}} \sqrt{6} = \frac{1}{4} \sqrt{6};$

We multiply both terms of the first Fraction by 2, to make the denominator 16 a perfect *square*. The reciprocal of this denominator is the square factor, and the numerator 6 is the other factor.

Surds which are apparently *dissimilar*, often become *similar* when reduced to their simplest forms. They are thus prepared for Addition or Subtraction, as will be seen hereafter.

Surds of Different Roots reduced to the Same Root.

(239.) Two or more Surds of *different roots*, may be reduced to equivalent ones of the *same root*,—by reducing their fractional exponents to a *common denominator*,—raising each quantity under the $\sqrt{\quad}$ to the power denoted by the *numerator* of its new exponent,—and taking the root denoted by the common denominator.

Thus to reduce $\sqrt{5}$ and $\sqrt[3]{4}$ to the *same root*.

Reducing the exponents $\frac{1}{2}$ and $\frac{1}{3}$ to a common denominator,

we find $\sqrt{5} = 5^{\frac{3}{6}}$, and $\sqrt[3]{4} = 4^{\frac{2}{6}}$, (235).

Cubing the 5, and *squaring* the 4, according to the numerators of their new exponents, (211), we find

$$\sqrt{5} = 125^{\frac{1}{6}}, \text{ and } \sqrt[3]{4} = 16^{\frac{1}{6}}.$$

The *square* root and the *cube* root have thus been both reduced to the *sixth* root.—This kind of reduction is sometimes necessary in the Multiplication and Division of Surds,—as will be seen hereafter.

EXERCISES

On the Transformation or Reduction of Surds.

- | | |
|--|----------------------------------|
| 1. Find the Product of $3\sqrt{4a}$. (236). | <i>Ans.</i> $\sqrt{36a}$ |
| 2. Find the Product of $2\sqrt[3]{3x}$. | <i>Ans.</i> $\sqrt[3]{24x}$ |
| 3. Find the Product of $4\sqrt{ab}$. | <i>Ans.</i> $\sqrt{16ab}$ |
| 4. Find the Product of $a\sqrt[3]{5x}$. | <i>Ans.</i> $\sqrt[3]{5a^3x}$ |
| 5. Find the Product of $3x\sqrt{7}$. | <i>Ans.</i> $\sqrt{63x^2}$ |
| 6. Find the Product of $ax\sqrt[3]{10}$. | <i>Ans.</i> $\sqrt[3]{10a^3x^3}$ |
| 7. Find the Product of $7\sqrt{a+1}$. | <i>Ans.</i> $\sqrt{49a+49}$ |
| 8. Find the Product of $2\sqrt[3]{1-x}$. | <i>Ans.</i> $\sqrt[3]{8-8x}$ |
| 9. Find the Product of $(a+b)\sqrt{2}$. | <i>Ans.</i> $\sqrt{2(a+b)^2}$ |
| 10. Find the Product of $(a-x)\sqrt[3]{3}$. | <i>Ans.</i> $\sqrt[3]{3(a-x)^3}$ |
| 11. Find the Product of $(a+1)\sqrt{x}$. | <i>Ans.</i> $\sqrt{x(a+1)^2}$ |
| 12. Find the Product of $(a-1)\sqrt[3]{y}$. | <i>Ans.</i> $\sqrt[3]{y(a-1)^3}$ |

13. Reduce $\sqrt{245ay^5}$ to its *simplest form*. (237).

To discover the greatest *square factor* of 245, we will divide this number successively by the *square numbers*,

4, 9, 16, 25, 36, 49, 64, &c.,

and take the largest divisor that leaves no *remainder*.

Such divisor will be found to be 49;— $49)245(5$.

Then $\sqrt{245ay^5} = \sqrt{49y^4} \sqrt{5ay} = 7y^2 \sqrt{5ay}$.

- | | |
|---|-----------------------------|
| 14. Reduce $\sqrt{4a^2b}$ to its simplest form. | <i>Ans.</i> $2a\sqrt{b}$ |
| 15. Reduce $3\sqrt{a^3x}$ to its simplest form. | <i>Ans.</i> $3a\sqrt{ax}$ |
| 16. Reduce $\sqrt[3]{8y}$ to its simplest form. | <i>Ans.</i> $2\sqrt[3]{y}$ |
| 17. Reduce $2\sqrt{8ax^2}$ to its simplest form. | <i>Ans.</i> $4x\sqrt{2a}$ |
| 18. Reduce $\sqrt[3]{27a^4}$ to its simplest form. | <i>Ans.</i> $3a\sqrt[3]{a}$ |
| 19. Reduce $5\sqrt{48a^3y}$ to its simplest form. | <i>Ans.</i> $20a\sqrt{3ay}$ |
| 20. Reduce $2\sqrt[3]{64a^4}$ to its simplest form. | <i>Ans.</i> $8a\sqrt[3]{a}$ |

21. Reduce $3\sqrt{128x^3}$ to its simplest form. *Ans.* $24x\sqrt{2x}$
22. Reduce $a^3\sqrt[3]{250y}$ to its simplest form. *Ans.* $5a^3\sqrt[3]{2y}$
23. Reduce $2x\sqrt{432}$ to its simplest form. *Ans.* $24x\sqrt{3}$.
24. Reduce $3y^3\sqrt[3]{135}$ to its simplest form. *Ans.* $9y^3\sqrt[3]{5}$.
25. Reduce $\sqrt{8+12a^3}$ to its simplest form. *Ans.* $2\sqrt{2+3a^3}$.
26. Reduce $(a+b)^3\sqrt[3]{81x^3}$ to its simplest form. *Ans.* $3x(a+b)^3\sqrt[3]{3}$
27. Reduce $3\sqrt{a^2x-2a^3}$ to its simplest form. *Ans.* $3a\sqrt{x-2a}$.
28. Reduce $(a-x)^3\sqrt[3]{192a^3x}$ to its simplest form. *Ans.* $4a(a-x)^3\sqrt[3]{3x}$.
29. Reduce $4\sqrt{4x^2+8x^3}$ to its simplest form. *Ans.* $8x\sqrt{1+2x}$.

30. Reduce $3\sqrt{\frac{27}{50}}$ to an *integral* Surd in its *simplest form*. (238).

$$3\sqrt{\frac{27}{50}} = 3\sqrt{\frac{27 \times 2}{50 \times 2}} = 3\sqrt{\frac{54}{100}} = 3\sqrt{\frac{9}{100}} \sqrt{6} = \frac{9}{10} \sqrt{6}.$$

31. Reduce $2\sqrt{\frac{14}{25}}$ to an *integral* Surd in its simplest form. *Ans.* $\frac{2}{5} \sqrt{14}$
32. Reduce $4\sqrt{\frac{4ax^3}{11}}$ to an *integral* Surd in its simplest form. *Ans.* $\frac{8x}{11} \sqrt{11ax}$.
33. Reduce $a^2\sqrt[3]{\frac{16a}{3y^3}}$ to an *integral* Surd in its simplest form. *Ans.* $\frac{2a^2}{3y} \sqrt[3]{18a}$.
34. Reduce $5x\sqrt{\frac{1+y^2}{36}}$ to an *integral* Surd in its simplest form. *Ans.* $\frac{5x}{6} \sqrt{1+y^2}$.
35. Reduce $\frac{2}{5}\sqrt{\frac{18}{a+x}}$ to an *integral* Surd in its simplest form. *Ans.* $\frac{6}{5(a+x)} \sqrt{2(a+x)}$.

36. Reduce $5\sqrt{2}$ and $3\sqrt[3]{4}$ to Surds of the same root. (239).

By reducing the *exponents* of the Surd factor to the common denominator, we find

$$2^{\frac{1}{2}} = 2^{\frac{3}{6}} = 8^{\frac{1}{6}}; \text{ and } 4^{\frac{1}{3}} = 4^{\frac{2}{6}} = 16^{\frac{1}{6}}.$$

$$\text{Ans. } 5\sqrt[6]{8}, \text{ and } 3\sqrt[6]{16}.$$

Prefixing the rational *co-efficients*, we have

$$5\sqrt{2} = 5\sqrt[6]{8}, \text{ and } 3\sqrt[3]{4} = 3\sqrt[6]{16}.$$

37. Reduce $2\sqrt[3]{5}$ and $5\sqrt{3}$ to Surds of the same root.

$$\text{Ans. } 2\sqrt[6]{25}, \text{ and } 5\sqrt[6]{27}.$$

38. Reduce $a\sqrt{5}$ and $x\sqrt[3]{2}$ to Surds of the same root.

$$\text{Ans. } a\sqrt[6]{125}, \text{ and } x\sqrt[6]{4}.$$

39. Reduce $10\sqrt{10}$ and $2\sqrt[4]{3x}$ to Surds of the same root.

$$\text{Ans. } 10\sqrt[4]{100}, \text{ and } 2\sqrt[4]{3x}$$

40. Reduce $7\sqrt[3]{3y}$ and $2y\sqrt[4]{xy^2}$ to Surds of the same root.

$$\text{Ans. } 7\sqrt[12]{81y^4}, \text{ and } 2y\sqrt[12]{x^3y^6}$$

41. Reduce $a^2x\sqrt[4]{\frac{3}{4}}$ and $\frac{3}{4}\sqrt{a^2x}$ to Surds of the same root.

$$\text{Ans. } a^2x\sqrt[4]{\frac{3}{4}}, \text{ and } \frac{3}{4}\sqrt[4]{a^4x^2}.$$

42. Reduce $\frac{a}{2x}\sqrt[3]{2}$ and $5y\sqrt{\frac{1}{2x}}$ to Surds of the same root.

$$\text{Ans. } \frac{a}{2x}\sqrt[6]{4}, \text{ and } 5y\sqrt[6]{\frac{1}{8x^3}}.$$

43. Reduce $2x\sqrt{a}$ and $\frac{1}{y}\sqrt[4]{\frac{1}{2}}$ to Surds of the same root.

$$\text{Ans. } 2x\sqrt[4]{a^2}, \text{ and } \frac{1}{y}\sqrt[4]{\frac{1}{2}}.$$

44. Reduce $\frac{2}{3}\sqrt[3]{10}$ and $31\sqrt{\frac{a}{y}}$ to Surds of the same root.

$$\text{Ans. } \frac{2}{3}\sqrt[6]{100}, \text{ and } 31\sqrt[6]{\frac{a^3}{y^3}}.$$

45. Reduce $\frac{1}{2y}\sqrt[3]{\frac{1}{a}}$ and $\frac{1}{2x}\sqrt[4]{\frac{1}{y^2}}$ to Surds of the same root.

$$\text{Ans. } \frac{1}{2y}\sqrt[12]{\frac{1}{a^4}}, \text{ and } \frac{1}{2x}\sqrt[12]{\frac{1}{y^6}}.$$

ADDITION AND SUBTRACTION OF SURDS.

(240.) 1. The Sum, or Difference, of *similar surds* is obtained by prefixing the sum, or difference, of their *coefficients* as a coefficient to the common *radical factor*.

2. *Dissimilar surds* can be added together, or subtracted the one from the other, only by the proper *sign*; but Surds apparently dissimilar often become similar when reduced to their simplest forms, (237).

EXAMPLE.

To Add together $5\sqrt{80a}$ and $3\sqrt{125a}$.

Reducing the surds to their simplest forms, we find

$$5\sqrt{80a} = 5\sqrt{16}\sqrt{5a} = 20\sqrt{5a};$$

$$\text{and } 3\sqrt{125a} = 3\sqrt{25}\sqrt{5a} = 15\sqrt{5a}.$$

The two given Surds have thus become *similar*, (232.) Adding together the coefficients 20 and 15, we find the *Sum* $35\sqrt{5a}$.

The *Difference* of the two given Surds, is $(20 - 15)\sqrt{5a} = 5\sqrt{5a}$.

The Addition or Subtraction of similar Surds is evidently nothing more than the addition or subtraction of similar *monomials*; thus

$$20 \text{ times } \sqrt{5a} + 15 \text{ times } \sqrt{5a} \text{ is } 35 \text{ times } \sqrt{5a};$$

$$\text{just as } 20a + 15a \text{ is } 35a.$$

EXERCISES

On the Addition and Subtraction of Surds.

1. Find the Sum of $3\sqrt{27}$ and $2\sqrt{48}$. *Ans.* $17\sqrt{3}$.
2. Find the Difference between $\sqrt{50}$ and $\sqrt{72}$. *Ans.* $\sqrt{2}$.
3. Find the Sum of $7\sqrt{28}$ and $6\sqrt{63}$. *Ans.* $32\sqrt{7}$.
4. Find the Difference between $21\sqrt{2}$ and $5\sqrt{18}$. *Ans.* $6\sqrt{2}$.
5. Find the Sum of $\sqrt{180}$ and $\sqrt{405}$. *Ans.* $15\sqrt{5}$.

6. Find the Difference between $\sqrt{18}$ and $2\sqrt{50}$. *Ans.* $7\sqrt{2}$.
7. Find the Sum of $\sqrt{12a^2}$ and $\sqrt{27a^2}$. *Ans.* $5a\sqrt{3}$.
8. Find the Difference between $3\sqrt{24x^2}$ and $\sqrt{54x^2}$. *Ans.* $3x\sqrt{6}$
9. Find the Sum of $4\sqrt{3a}$ and $\sqrt{48a}$. *Ans.* $8\sqrt{3a}$
10. Find the Difference between $\sqrt{4a^3}$ and $\sqrt{9a^3}$. *Ans.* $a\sqrt{a}$.
11. Find the Sum of $3\sqrt[3]{4}$ and $7\sqrt[3]{4}$. *Ans.* $10\sqrt[3]{4}$
12. Find the Difference between $9\sqrt{200}$ and $\sqrt{288}$. *Ans.* $78\sqrt{2}$.
13. Find the Sum of $4\sqrt[3]{54}$ and $2\sqrt[3]{250}$. *Ans.* $22\sqrt[3]{2}$.
14. Find the Difference between $5\sqrt{9x^3}$ and $3\sqrt{x^3}$. *Ans.* $12x\sqrt{x}$.
15. Find the Sum of $2\sqrt[3]{16a}$ and $\sqrt[3]{54a}$. *Ans.* $7\sqrt[3]{2a}$.
16. Find the Difference between $3\sqrt[4]{10}$ and $5\sqrt[4]{10}$. *Ans.* $2\sqrt[4]{10}$.
17. Find the Sum of $5\sqrt{98x}$ and $10\sqrt{2x}$. *Ans.* $45\sqrt{2x}$
18. Find the Difference between $3a\sqrt[3]{5}$ and $a\sqrt[3]{5}$. *Ans.* $2a\sqrt[3]{5}$.
19. Find the Sum of $a\sqrt[3]{y^2}$ and $3b\sqrt[3]{y^2}$. *Ans.* $(a+3b)\sqrt[3]{y^2}$.
20. Find the Difference between $5\sqrt[4]{5}$ and $2a\sqrt[4]{5}$. *Ans.* $(5-2a)\sqrt[4]{5}$

21. Find the Sum of $(a+1)^{\frac{1}{2}}$ and $\sqrt{4a+4}$. *Ans.* $3(a+1)^{\frac{1}{2}}$
22. Find the Difference between $\sqrt{1+x}$ and $3(1+x)^{\frac{1}{2}}$. *Ans.* $2(1+x)^{\frac{1}{2}}$
23. Find the Sum of $2(a-x)^{\frac{1}{2}}$ and $\sqrt{9a-9x}$. *Ans.* $5(a-x)^{\frac{1}{2}}$
24. Find the Difference between $\sqrt[3]{2+y}$ and $4(y+2)^{\frac{1}{3}}$. *Ans.* $3(2+y)^{\frac{1}{3}}$
25. Find the Difference between $4(1+x^2)^{\frac{1}{2}}$ and $4\sqrt[3]{x^2+1}$. *Ans.* $4\left(\frac{1}{1+x^2} - \frac{1}{x^2+1}\right)^{\frac{1}{3}}$.

MULTIPLICATION AND DIVISION OF SURDS.

(241.) 1. The Product, or Quotient, of two Surds of the *same root*, is obtained by prefixing the product, or quotient, of their *coefficients* as a coefficient to the product, or quotient, of the *radical factors*,—the latter being affected with the same fractional *exponent* or radical sign.

2. Surds of *different roots* may be reduced to equivalent ones of the *same root*, (239), and then multiplied, or divided, as above. But

3. *Any two roots of the same quantity* may be multiplied into each other, by adding together their fractional *exponents*; or divided, the one into the other, by subtracting the exponent of the *divisor* from that of the *dividend*.

EXAMPLE.

To find the Product of $2\sqrt{10} \times 3\sqrt{2}$.

Since it is immaterial in what *order* the four factors are taken, we may take them in the order,

$2 \times 3 \sqrt{10} \times \sqrt{2}$; which gives the Product $6\sqrt{20}$, (234), $= 12\sqrt{5}$, (237)

The *Quotient* of $2\sqrt{10} \div 3\sqrt{2}$ is $\frac{2}{3}\sqrt{5}$, since this quotient multiplied by the *divisor* produces the *dividend*.

EXERCISES

On the Multiplication and Division of Surds.

- | | |
|---|-------------------------------|
| 1. Find the Product of $5\sqrt{8} \times 3\sqrt{5}$. | <i>Ans.</i> $30\sqrt{10}$. |
| 2. Find the Quotient of $6\sqrt{54} \div 3\sqrt{2}$. | <i>Ans.</i> $6\sqrt{3}$. |
| 3. Find the Product of $\sqrt{108} \times 2\sqrt{6}$. | <i>Ans.</i> $36\sqrt{2}$. |
| 4. Find the Quotient of $2\sqrt{96} \div \sqrt{54}$. | <i>Ans.</i> $2\frac{2}{3}$. |
| 5. Find the Product of $3\sqrt{5ax} \times 4\sqrt{20a}$. | <i>Ans.</i> $120a\sqrt{x}$. |
| 6. Find the Quotient of $4\sqrt{12a} \div 2\sqrt{6}$. | <i>Ans.</i> $2\sqrt{2a}$. |
| 7. Find the Product of $\sqrt{3ax} \times 3\sqrt{ax}$. | <i>Ans.</i> $3ax\sqrt{3}$. |
| 8. Find the Quotient of $6\sqrt{12x^2} \div 3\sqrt{4}$. | <i>Ans.</i> $2x\sqrt{3}$. |
| 9. Find the Product of $\sqrt[3]{18} \times 5\sqrt[3]{4}$. | <i>Ans.</i> $10\sqrt[3]{9}$. |
| 10. Find the Quotient of $4\sqrt[3]{72} \div 2\sqrt[3]{18}$. | <i>Ans.</i> $2\sqrt[3]{4}$. |

11. Find the Product of $5\sqrt{a} \times 3\sqrt[3]{a}$.

By reducing the surds to the *same root*, we obtain

$$5\sqrt[6]{a^3} \text{ and } 3\sqrt[6]{a^2}, (239).$$

These are to be multiplied together as before. Ans. $15\sqrt[6]{a^5}$.

But the given roots of the *same quantity a*, may also be multiplied into each other, by adding together their fractional *exponents* $\frac{1}{2}$ and $\frac{1}{3}$. Thus,

$$5a^{\frac{1}{2}} \times 3a^{\frac{1}{3}} = 15a^{\frac{5}{6}} = 15\sqrt[6]{a^5}, \text{ as before.}$$

Either of these two methods may be applied to roots of the same quantity. The first only is applicable to *different roots of different quantities*.

12. Find the Product of $3\sqrt[3]{x} \times 2\sqrt{x}$. Ans. $6\sqrt[6]{x^5}$.

13. Find the Quotient of $2\sqrt{3} \div 5\sqrt[3]{3}$. Ans. $\frac{2}{5}\sqrt[6]{3}$.

14. Find the Product of $4\sqrt{3} \times 2\sqrt[3]{2}$. Ans. $8\sqrt[6]{108}$.

15. Find the Quotient of $8\sqrt[3]{a} \div 4\sqrt[4]{a}$. Ans. $2\sqrt[12]{a}$.

16. Find the Product of $\sqrt{ax} \times 3\sqrt[4]{ax}$. Ans. $3\sqrt[4]{a^3x^3}$.

17. Find the Quotient of $4\sqrt{x^3} \div 2\sqrt[4]{x}$. Ans. $2x^4\sqrt{x}$.

18. Find the Product of $\frac{1}{2}\sqrt{5} \times \frac{4}{3}\sqrt[4]{10}$. Ans. $\frac{1}{6}\sqrt[4]{250}$.

19. Find the Quotient of $a^3\sqrt{\frac{1}{2}} \div a^3\sqrt[3]{\frac{1}{2}}$. Ans. $a^2\sqrt[6]{\frac{1}{2}}$.

20. Find the Product of $(3+2\sqrt{5}) \times (2-\sqrt{5})$.

In cases of this kind, in which a Surd is connected with another quantity by the sign $+$ or $-$, the Multiplication, or Division, must be performed as on *polynomials*.

$$\begin{array}{r} 3+2\sqrt{5} \\ 2-\sqrt{5} \\ \hline 6+4\sqrt{5} \\ -3\sqrt{5}-10 \\ \hline 6+\sqrt{5}-10=\sqrt{5}-4. \end{array}$$

Each term of the *multiplcand* is multiplied by each term of the *multiplier*, and the partial products are then added together.

Observe that $2\sqrt{5} \times -\sqrt{5} = -2\sqrt{25} = -10$.

21. Find the Product of $(2+3\sqrt{2}) \times (1+5\sqrt{2})$.

Ans. $13\sqrt{2}+32$.

22. Find the Product of $(4 - \sqrt{3}) \times (2 + 3\sqrt{3})$.

Ans. $10\sqrt{3} - 1$.

23. Find the Product of $(5 + 2\sqrt{6}) \times (1 + 2\sqrt{6})$.

Ans. $12\sqrt{6} + 29$.

24. Find the Product of $(1 - 4\sqrt{7}) \times (3 - 3\sqrt{7})$.

Ans. $87 - 15\sqrt{7}$.

25. Find the Quotient of $(\sqrt{20} + \sqrt{12}) \div (\sqrt{5} + \sqrt{3})$.

Ans. 2.

RATIONALIZATION OF SURD DIVISORS.

(242.) In computing an approximate value of an *irrational* numerical expression, it is expedient that a *surd divisor* or denominator be made *rational*.

For example, suppose we wish to compute an approximate value of

$$\frac{2}{\sqrt{3}}, \text{ 2 divided by the square root of 3.}$$

If we extract the square root of 3, for a divisor, a regard to *accuracy* will require that the root be continued to *several figures*, and hence will arise the inconvenience of *dividing by a large number*.

By multiplying both terms by the denominator, we have

$$\frac{2}{\sqrt{3}} = \frac{2\sqrt{3}}{\sqrt{9}} = \frac{\sqrt{12}}{3}, \text{ in which the divisor is } \textit{rational}.$$

The value will therefore be found by taking $\frac{1}{3}$ of the square root of 12; and by this method the computation is much simplified.

In pursuance of the object at present in view, it is necessary

To find a Multiplier of a given Surd which will cause the Product to be Rational.

(243.) 1. A *monomial* Surd will produce a rational quantity by being multiplied into itself with its *exponent subtracted from a unit*.

Thus $a^{\frac{1}{3}}$ multiplied by $a^{1-\frac{1}{3}}$, or $a^{\frac{1}{3}} \times a^{\frac{2}{3}} = a$, (241...3).

2. A *binomial* in which one or both terms contain an irrational *square root*, will produce a rational quantity by being multiplied into itself with a *sign changed*.

$$\text{Thus } (\sqrt{3} + \sqrt{2}) \times (\sqrt{3} - \sqrt{2}) = 3 - 2 = 1.$$

The product in this case is readily found on the principle, that the Product of the *sum* and *difference* of two quantities is equal to the *difference of the squares* of the two quantities.

3. A *trinomial* containing irrational *square roots* will produce a *binomial* Surd by being multiplied into itself with a *sign changed*; and this binomial may be rationalized as above.

These principles provide for the most useful cases of the subject under consideration.—In applying them to the rationalization of surd denominators, both terms of the given Fraction must be multiplied by the same quantity, (81).

EXERCISES

On the Rationalization of Surd Denominators.

1. Reduce $\frac{2+\sqrt[3]{3}}{\sqrt[3]{3}}$ to a Fraction having a *rational* denominator.
Ans. $\frac{2\sqrt[3]{9+3}}{3}$.
2. Reduce $\frac{5}{3+\sqrt{5}}$ to a Fraction having a rational denominator.
Ans. $\frac{15-5\sqrt{5}}{4}$.
3. Reduce $\frac{2a}{2-\sqrt{7}}$ to a Fraction having a rational denominator.
Ans. $\frac{4a+2a\sqrt{7}}{-3}$.
4. Reduce $\frac{3x}{\sqrt{a}-\sqrt{x}}$ to a Fraction having a rational denominator.
Ans. $\frac{3x\sqrt{a}+3x\sqrt{x}}{a-x}$.
5. Reduce $\frac{10}{a+\sqrt{10}}$ to a Fraction having a rational denominator.
Ans. $\frac{10a-10\sqrt{10}}{a^2-10}$.
6. Reduce $\frac{a-\sqrt[4]{2}}{\sqrt[4]{2}}$ to a Fraction having a rational denominator
Ans. $\frac{a\sqrt[4]{8}-2}{2}$.
7. Reduce $\frac{8}{\sqrt{3}+\sqrt{2+1}}$ to a Fraction having a rational denominator
Ans. $4+2\sqrt{2}-2\sqrt{6}$.

INVOLUTION AND EVOLUTION OF SURDS.

(244.) The Powers and Roots of *irrational* quantities are obtained, or *indicated*, according to the general principles of Involution and Evolution which have been established in the preceding Chapter.

We present here however a particular case of the

SQUARE ROOT OF BINOMIAL SURDS.

(245.) A Numerical Binomial of the form $a \pm \sqrt{b}$ admits of a square root in a *rational* and an *irrational* term, or two irrational terms, whenever $a^2 - b$ is a *perfect square*.

To determine the method to be pursued in this case of evolution, we must find

Formulas for the Square Root of $a \pm \sqrt{b}$.

The *square of the sum* of any two quantities, is equal to the *sum of their squares + twice their product*, (59). The binomial $a + \sqrt{b}$ may therefore represent the square of the sum of a *rational* and an *irrational* numerical term, or of two irrational terms, in the square root; a representing the *sum of the squares* of the two terms,—which sum will necessarily be *rational*,—and \sqrt{b} representing *twice the product* of the two terms.

In like manner $a - \sqrt{b}$ may represent the square of the difference of a rational and an irrational numerical term, or of two irrational numerical terms, in the square root, (60).

If therefore we take x and y to represent the two terms of the square root of $a \pm \sqrt{b}$, we shall have,

$$(1) \quad x^2 + y^2 = a;$$

$$(2) \quad x + y = \sqrt{a + \sqrt{b}};$$

$$(3) \quad x - y = \sqrt{a - \sqrt{b}};$$

Multiplying together equations (2) and (3), we have,

$$(4) \quad x^2 - y^2 = \sqrt{a^2 - b}. \quad (58.)$$

Adding together the (4) and (1), and also subtracting the (4) from the (1), and dividing by 2, we shall find,

$$(5) \quad x^2 = \frac{a + \sqrt{a^2 - b}}{2};$$

$$(6) \quad y^2 = \frac{a - \sqrt{a^2 - b}}{2}.$$

Extracting the square root of each of these equations,

$$\text{we find } x = \sqrt{\frac{a + \sqrt{a^2 - b}}{2}};$$

$$\text{and } y = \sqrt{\frac{a - \sqrt{a^2 - b}}{2}}.$$

Substituting these values of x and y in equations (2) and (3),—and interchanging the first and second members,—we have,

$$(A) \quad \sqrt{a + \sqrt{b}} = \sqrt{\frac{a + \sqrt{a^2 - b}}{2}} + \sqrt{\frac{a - \sqrt{a^2 - b}}{2}};$$

$$(B) \quad \sqrt{a - \sqrt{b}} = \sqrt{\frac{a + \sqrt{a^2 - b}}{2}} - \sqrt{\frac{a - \sqrt{a^2 - b}}{2}}.$$

These are the *Formulas* required. The right hand member of each will contain at most but two *irrational* terms, when $(a^2 - b)$ is a *perfect square*, that is, has an exact *square root*.

EXAMPLE.

To find the Square Root of $6 + 2\sqrt{5}$, or $6 + \sqrt{20}$, (236.)

Substituting 6 for a , and $\sqrt{20}$ for \sqrt{b} , in Formula (A),

$$\sqrt{6 + \sqrt{20}} = \sqrt{\frac{6 + \sqrt{36 - 20}}{2}} + \sqrt{\frac{6 - \sqrt{36 - 20}}{2}}.$$

And since $\sqrt{36 - 20} = \sqrt{16} = 4$, the second member reduces to

$$\sqrt{\frac{6+4}{2}} + \sqrt{\frac{6-4}{2}} = \sqrt{5+1}, \text{ the Root required.}$$

The root $\sqrt{5+1}$ may be verified by *squaring* it.

$$\text{Thus } (\sqrt{5+1})^2 = 5 + 2\sqrt{5} + 1 = 6 + 2\sqrt{5}.$$

Formula (B) would give the square root of $6 - 2\sqrt{5}$.

EXERCISES

On the Powers and Roots of Surds.

It will be readily perceived that by *cancelling the exponent or sign of a root, that root is raised to the corresponding power.*

Thus the *square* of \sqrt{a} is a ; the *cube* of $\sqrt[3]{a}$ is a , &c.

1. Find the *Square*, and also the *square root*, of $5\sqrt[3]{4}$.

The *square* of $5 \times 4^{\frac{1}{3}}$ is $25 \times 4^{\frac{2}{3}} = 25\sqrt[3]{16}$, (200).

Reducing this result to its *simplest form*, we have

$$25\sqrt[3]{16} = 25\sqrt[3]{8}\sqrt[3]{2} = 50\sqrt[3]{2}, \text{ (237).}$$

The *square root* of $5 \times 4^{\frac{1}{3}}$ is $5^{\frac{1}{2}} \times 4^{\frac{1}{6}}$, or $\sqrt{5}\sqrt[6]{4}$, (208).

Multiplying together the *two factors* of this root, we have

$$\sqrt{5}\sqrt[6]{4} = \sqrt[6]{125}\sqrt[6]{4} = \sqrt[6]{500}, \text{ (241 \dots 2).}$$

2. Find the *Square*, and also the *square root*, of $9\sqrt[3]{3}$.

Ans. $81\sqrt[3]{9}$, and $3\sqrt[6]{3}$.

3. Find the *Cube*, and also the *cube root*, of $3\sqrt{2}$.

Ans. $54\sqrt{2}$, and $\sqrt[6]{18}$.

4. Find the *Square*, and also the *square root*, of $2\sqrt[3]{3}$.

Ans. $4\sqrt[3]{9}$, and $\sqrt[6]{24}$.

5. Find the *Cube*, and also the *cube root*, of $3a\sqrt{3}$.

Ans. $81a^3\sqrt{3}$, and $\sqrt[6]{27a^2}$.

6. Find the *Square*, and also the *square root*, of $x^2\sqrt[3]{5}$.

Ans. $x^4\sqrt[3]{25}$, and $x\sqrt[6]{5}$.

7. Find the *Square*, and also the *square root*, of $3\sqrt{a+1}$.

Ans. $9(a+1)$, and $\sqrt[4]{9(a+1)}$.

8. Find the *Square*, and also the *square root*, of $4\sqrt[3]{1-x}$.

Ans. $16\sqrt[3]{(1-x)^2}$, and $2\sqrt[6]{1-x}$.

9. Find the *Cube*, and also the *cube root*, of $2\sqrt{a^2-x}$.

Ans. $8\sqrt{(a^2-x)^3}$, and $\sqrt[6]{4(a^2-x)}$.

10. Find the *Square*, and also the *square root*, of $3\sqrt[3]{2-a}$.

Ans. $9(2-a)$, and $\sqrt[4]{9(2-a)}$.

11. Find the Square of $2 + \sqrt{3}$, and of $2 - \sqrt{3}$.

These squares may be found by multiplying each binomial into itself, or, more readily, by applying the propositions relating to the squares of the *sum* and the *difference* of two quantities.

$$\text{Ans. } 7 + 4\sqrt{3}, \text{ and } 7 - 4\sqrt{3}.$$

12. Find the square of $3 + 2\sqrt{5}$, and of $5 - 3\sqrt{2}$.

$$\text{Ans. } 12\sqrt{5} + 29, \text{ and } 43 - 30\sqrt{2}.$$

13. Find the Square of $\sqrt{5 + 2\sqrt{a}}$, and of $\sqrt{3 - 5\sqrt{x}}$.

$$\text{Ans. } 5 + 2\sqrt{a}, \text{ and } 3 - 5\sqrt{x}.$$

14. Find the Square of $\sqrt{2 + 3\sqrt{5}}$, and of $\sqrt{a - 2x\sqrt{x}}$.

$$\text{Ans. } 6\sqrt{10} + 47, \text{ and } a - 4x\sqrt{ax} + 4x^3.$$

15. Find the Cube of $\sqrt[3]{1 + 5\sqrt{4}}$, and of $\sqrt[3]{a - 3\sqrt{a}}$.

$$\text{Ans. } 1 + 5\sqrt{4}, \text{ and } a - 3\sqrt{a}.$$

16. Find the Square of $\sqrt{3 + a\sqrt{3}}$, and of $\sqrt{3 - 3\sqrt{a}}$.

$$\text{Ans. } 3 + 6a + 3a^2, \text{ and } 3 - 6\sqrt{3a} + 9a.$$

17. Find the Cube of $2 + 2\sqrt{a}$, and of $2 - a\sqrt{2}$.

$$\text{Ans. } 8 + 24a + (24 + 8a)\sqrt{a}, \text{ and } 8 + 12a^2 - (12a + 2a^3)\sqrt{2}.$$

18. Find the Cube of $(\sqrt{a + \sqrt{2}})^{\frac{1}{3}}$ and of $(\sqrt[3]{1 - \sqrt{y}})^{\frac{1}{3}}$

$$\text{Ans. } \sqrt{a + \sqrt{2}}, \text{ and } \sqrt[3]{1 - \sqrt{y}}.$$

19. Find the Square root of $a + 2\sqrt{ax} + x$, (218).

$$\text{Ans. } \sqrt{a} + \sqrt{x}.$$

20. Find the Square root of $a + 2\sqrt{a} + 1$, and of $3 + 2a\sqrt{3} + a^2$.

$$\text{Ans. } \sqrt{a} + 1, \text{ and } \sqrt{3} + a.$$

21. Find the Square root of $x - 2\sqrt{x} + 1$, and of $5 - 2x\sqrt{5} + x^2$.

$$\text{Ans. } \sqrt{x} - 1, \text{ and } \sqrt{5} - x.$$

22. Find the Square root of $23 + 8\sqrt{7}$, (245). *Formula A.*

$$\text{Ans. } 4 + \sqrt{7}.$$

23. Find the Square root of $19 + 8\sqrt{3}$, and of $7 - 2\sqrt{10}$.

$$\text{Ans. } 4 + \sqrt{3}, \text{ and } \sqrt{5} - \sqrt{2}.$$

IMAGINARY QUANTITIES.

(246.) An *even root* of a *negative* quantity is *impossible*, and the symbol of such a root is therefore called an *imaginary* or *impossible* quantity, in contradistinction to *real* quantities.

Thus $\sqrt{-4}$, the square root of -4 , is *imaginary*, since there is no quantity whose *square* is -4 ; and for a like reason $\sqrt{-16}$ is *imaginary*.

An *Imaginary Quantity* results in calculation from some *impossibility* in the conditions of a *Problem*, and may therefore be regarded as a sign of such impossibility.

As an instance of this, suppose it were required *To find a number whose square subtracted from 5 shall leave 9.*

If x represent the required number, the *Equation* of the problem will be

$$5 - x^2 = 9.$$

From this equation, $x^2 = -4$; and hence $x = \sqrt{-4}$

The value of x being *imaginary* or impossible, shows that the *problem is impossible*; that is, that there is no number whose square subtracted from 5 will leave 9.

Imaginary Quantities may become the subjects of calculation like *real quantities*. Thus we may wish to *verify* an *imaginary* value of the unknown quantity in an *Equation*, by subjecting this value to all the operations which are performed on its *symbol* in the equation.

Calculus of Imaginary Quantities.

(247.) All the principles which have been established for the calculus of *radical* quantities are applicable to *imaginary* quantities, except that the ambiguous sign \pm does not precede the *product* of two *imaginary* quantities, as is the case, in general, with *even roots of real quantities*.

The *sign* which affects the *Product* of two *imaginaries* may always be determined by means of *imaginary* and *real factors* into which such quantities may be resolved.

(248.) An Imaginary Quantity may always be resolved into the *like root* of (-1) multiplied into the like root of a *positive* quantity which is equal to the negative quantity in the given imaginary.

Thus -4 being equal to -1×4 , we have $\sqrt{-4} = \sqrt{-1} \times \sqrt{4}$;
and $-a$ being equal to $-1 \times a$, we have $\sqrt{-a} = \sqrt{-1} \times \sqrt{a}$. (234.)

By means of this transformation it may be shown, that

(249.) The Product of two imaginary *square roots* is the *negative* square root of the product of the two quantities under the $\sqrt{}$, if the given roots are preceded by like signs, $+$ or $-$; otherwise, it is the *positive* square root of that product.

$$\begin{aligned} \text{For example, } \sqrt{-a} \cdot \sqrt{-b} &= -\sqrt{ab}; \\ \text{and, } \sqrt{-a} \cdot -\sqrt{-b} &= +\sqrt{ab}. \end{aligned}$$

On general principles of calculation, the Product of the square roots of $-a$ and $-b$, is equal to the square root of the product ab , which would be $+\sqrt{ab}$, or else $-\sqrt{ab}$, (215).

But the ambiguity in the *sign* to be prefixed to an *even root* in general, is removed when we know the factors which entered into the composition of the quantity whose root is considered. When a^2 , for example, is known to have been derived from $a \times a$, the square root of a^2 is a , and not $-a$.

To determine the *sign* of \sqrt{ab} in the first example, we have

$$\begin{aligned} \sqrt{-a} \cdot \sqrt{-b} &= \sqrt{-1} \cdot \sqrt{a} \cdot \sqrt{-1} \cdot \sqrt{b}, \quad (248) \\ &= \sqrt{-1} \cdot \sqrt{-1} \cdot \sqrt{a} \sqrt{b} \\ &= (\sqrt{-1})^2 \sqrt{ab} = -1 \sqrt{ab}, \text{ or } -\sqrt{ab}. \end{aligned}$$

In the second example, we have

$$\begin{aligned} \sqrt{-a} \cdot -\sqrt{-b} &= \sqrt{-1} \cdot \sqrt{a} \cdot -\sqrt{-1} \cdot \sqrt{b}, \quad (248), \\ &= \sqrt{-1} \cdot -\sqrt{-1} \cdot \sqrt{a} \sqrt{b} \\ &= -(\sqrt{-1})^2 \sqrt{ab} = -(-1) \sqrt{ab} = +\sqrt{ab}. \end{aligned}$$

In this example the square $(\sqrt{-1})^2$ has the sign $-$ before it, because it results from multiplying $\sqrt{-1}$ by $-\sqrt{-1}$, (42); and $-(-1)$ becomes $+1$ by changing the sign in subtracting.

CHAPTER X.

QUADRATIC AND OTHER EQUATIONS.

(250.) A *QUADRATIC EQUATION*, or an equation of the *second degree*, is one in which the highest power of the unknown quantity is its *square*, or second power, as

$$3x^2=12; \text{ or } 3x^2+4x=20.$$

A *Cubic Equation*, or an equation of the *third degree*, is one in which the highest power of the unknown quantity is its *cube*, or third power; and in like manner is defined a *Biquadratic Equation*, or an equation of the fourth degree, and so on.

An Equation containing two or more unknown quantities is of the *degree* which corresponds to the greatest number of *unknown factors* in any of its terms.

Thus $3xy+y=a$ is an Equation of the *second degree*, its first term containing the unknown factors xy .

And $x^2y-y=b$ is an equation of the third degree, its first term containing three unknown factors xy .

Pure and Affected Equations.

(251.) A *Pure Equation* is an equation which contains but *one power* of the unknown quantity; and is a Simple Equation, a Pure Quadratic, or a Pure Cubic, &c., according to its degree.

Thus $3x^2$ is a pure *quadratic*; $x^3=64$ is a pure *cubic*.

An *Affected Equation* is one which contains different powers of the unknown quantity. When these powers are in regular *ascension*, beginning with the first power, the Equation is also called a *complete* equation.

Thus $x^2+3x=10$ is an *affected*, and also a *complete* quadratic, $x^3-2x^2=75$, or $x^3-3x=110$, is an *affected cubic*; and $x^3+3x^2+4x=28$ is a complete cubic.

Roots of Equations.

(252.) A *Root* of an Equation is a value of the unknown quantity in the equation. It will presently be shown that an equation of the 2d degree has *two roots*, of the 3d degree *three roots*, and so on.

In the Simple Equation $3x=15$, the value of the unknown quantity x is 5; then 5 is the root of the equation.

It is evident that in a Simple Equation there can be but *one* value of the unknown quantity that will *satisfy the equation*. An equation of the 1st degree has therefore but *one root*.

GENERAL PROPERTIES OF EQUATIONS.

1. *Divisors of an Equation.*

(253.) If a be a *root* of an Equation of any degree, containing but one unknown quantity, x ; the equation—with all its terms *transposed to one side*—will be divisible by $x-a$.

Let a be a root of the Equation

$$x^2 + mx = s;$$

then will the equation $x^2 + mx - s = 0$ be divisible by $x - a$

For let r be the *remainder*, if any, after the quotient q has been obtained; then will

$$x^2 + mx - s = (x - a)q + r = 0;$$

the dividend being equal to the remainder added to the product of the *divisor* and *quotient*.

But a being a value of x , (252), we have $x - a = 0$; then the $(x - a) \times q$ is 0, (43); and consequently $r = 0$; that is, the division will leave *no remainder*.

The preceding demonstration is applicable to an equation of the third, fourth, or any higher degree.

(254.) Conversely, If an Equation of any degree, containing but one unknown quantity, x ,—with all its terms transposed to one side—be divisible by $x - a$; then a will be a *root* of the equation.

This is evident from considering that the divisibility of the Equation, as shown above, depends on the condition that $x - a = 0$, or that a is a value of x .

2. *Number of Roots of an Equation.*

(255.) Every Equation containing but one unknown quantity, has just as many *roots* as there are units in the exponent of the highest power of the unknown quantity in the equation.

Let a represent a root of the *cubic* Equation

$$x^3 + mx^2 + nx = s.$$

Transposing s to the first side of the equation, we have

$$x^3 + mx^2 + nx - s = 0.$$

Dividing this equation by $x - a$, (253), we shall obtain an equation of the *second degree*, which may be represented by

$$x^2 + px - q = 0.$$

Let b be a root of this equation. Dividing the equation by $x - b$ we shall obtain an equation of the *first degree*, represented by

$$x - u = 0.$$

The binomials $x - a$, $x - b$, and $x - u$, which represent the two *divisors* and the last *quotient*, are the factors of the dividend.

$$x^3 + mx^2 + nx - s.$$

The original equation is thus resolved, representatively, into

$$(x - a)(x - b)(x - u) = 0.$$

Since this equation is divisible by each of these *three binomial factors*, it follows that a , b , and u are *three roots* of the equation, (254).

And since the given equation—being of the 3d degree—cannot be resolved into more than *three* binomial factors, each containing the *first* power of x , it cannot have more than *three roots*.

The same method of demonstration will show the proposition to be true for an Equation of any other degree.

The several *roots* of an Equation are not necessarily *unequal*, though such will usually be found to be the case. The preceding Proposition shows that an Equation may be resolved into binomial factors, each containing a *root* of the equation. Two or more of these roots may be equal to each other.

In the Equation $x^2 - 5x + 6 = (x - 2)(x - 3) = 0$,

the two roots are 2 and 3.

In the Equation $x^3 - 7x^2 + 16x - 12 = (x - 2)(x - 2)(x - 3) = 0$

the three roots are 2, 2, and 3.

SOLUTION OF PURE EQUATIONS OF THE SECOND AND
HIGHER DEGREES.

The Equations belonging to this class are those which, in their simplest forms, contain but one power of the unknown quantity

R U L E X X I I .

(256.) *For the Solution of a Pure Equation.*

1. Reduce the Equation to the form

$$x^n = s; \text{ in which } x^n \text{ must be positive.}$$

2. Extract that *root* of both sides of the equation which corresponds to the *power* of the unknown quantity

E X A M P L E .

To find the value of x in the Equation

$$\frac{x^2}{4} + 5 = 2x^2 - 23.$$

Clearing the equation of its Fraction, we have

$$x^2 + 20 = 8x^2 - 92.$$

Transposing, and adding similar terms, we find

$$-7x^2 = -112.$$

Dividing both sides of this equation by the coefficient -7 ,

$$x^2 = 16.$$

Extracting the *square root* of both sides,

$$x = \pm 4, (215).$$

The *two values* of x are thus found to be 4 and -4 , (255), either of which will satisfy the given Equation.

It is evident that, in a Pure Equation of the *second degree*, the two values of the unknown quantity will always be *equal*, with *contrary signs*.

- The unknown quantity may enter an Equation in a *surd* expression, which it will be necessary to *rationalize* in the solution of the equation.

Thus in the equation $\sqrt{x} + \sqrt{1+x} = a$, it would be necessary to *rationalize* x , that is, to clear it of the *radical sign* before the value of x could be determined.

The following observations will assist the student in the

Rationalization of Surd Quantities in an Equation.

(257.) A Surd quantity in an Equation will be *rationalized* by transposing all the other terms to the other side of the equation, and raising both sides to the *power* corresponding to the indicated *root*.

To rationalize x in the Equation

$$\sqrt{x+1}-a=b.$$

By transposition, $\sqrt{x+1}=a+b$

Squaring both sides, $x+1=a^2+2ab+b^2.$

(258.) Two Surds in an Equation may be rationalized by successive involutions,—in the first of which it will generally be expedient to make one of the Surds stand alone on one side of the equation.

To rationalize x in the Equation

$$\sqrt{x+\sqrt{1+x}}=a.$$

By transposition, $\sqrt{x}=a-\sqrt{1+x}.$

Squaring both sides, $x=a^2-2a\sqrt{1+x}+1+x.$

By transposition, $2a\sqrt{1+x}=a^2+1.$

Squaring both sides, $4a^2(1+x)=a^4+2a^2+1.$

(259.) When an Equation contains a Fraction whose terms are both *irrational*, it will sometimes be expedient to rationalize its *denominator* before clearing the equation of the fraction.

To rationalize x in the Equation

$$\frac{\sqrt{x}}{1+\sqrt{x}}=a.$$

Multiplying both terms of the Fraction by $1-\sqrt{x}$, (243...2),

$$\frac{\sqrt{x-x}}{1-x}=a.$$

Clearing this equation of its Fraction, and transposing, we have

$$\sqrt{x}=a-ax+x.$$

The Surd in this equation will be rationalized by *squaring both sides*, as in the preceding examples.

By the preceding methods we may generally rationalize one or more Surds containing the unknown quantity in an Equation. Other expedients, however, such as the *extraction of roots* in possible cases, in the course of the operation, will sometimes be requisite; but these must be left to the care and skill of the computer.

EXERCISES

On Rationalization, and Pure Equations.

1. Find the value of x in the equation

$$24 - \sqrt{2x^2 + 9} = 15. \quad \text{Ans. } x = \pm 6$$
2. Find the value of x in the equation

$$13 - \sqrt{3x^2 + 16} = 5. \quad \text{Ans. } x = \pm 4$$
3. Find the value of x in the equation

$$35 + \sqrt[3]{x-5} = 40. \quad \text{Ans. } x = 130$$
4. Find the value of x in the equation

$$1 + 2\sqrt{x} = \sqrt{4x+21}. \quad \text{Ans. } x = 25$$
5. Find the value of x in the equation

$$\sqrt{x-32} = \sqrt{x-\frac{1}{2}}\sqrt{32}. \quad \text{Ans. } x = 50$$
6. Find the value of x in the equation

$$3 + \sqrt{x+4} \times \sqrt{x-4} = 10. \quad \text{Ans. } x = \pm \sqrt{65}$$
7. Find the value of x in the equation

$$a + \sqrt{x-3} \times \sqrt{x+3} = 4a. \quad \text{Ans. } x = \pm 3\sqrt{a^2+1}.$$
8. Find the value of x in the equation

$$\sqrt{x} + \sqrt{5+x} = \frac{10}{\sqrt{5+x}}. \quad \text{Ans. } x = 1\frac{2}{3}$$
9. Find the value of x in the equation

$$\frac{4\sqrt{6x-9}}{4\sqrt{6x+6}} = \frac{\sqrt{6x-2}}{\sqrt{6x+2}}. \quad \text{Ans. } x = 6$$
10. Find the value of x in the equation

$$\frac{\sqrt{x+c} + \sqrt{x-c}}{\sqrt{x+c} - \sqrt{x-c}} = b. \quad \text{Ans. } x = \frac{cb^2+c}{2b}.$$

SOLUTION OF COMPLETE EQUATIONS OF THE
SECOND DEGREE.

These Equations in their simplest forms contain no other power of the unknown quantity than its *square* and first power.

The value of the unknown quantity will be found by the following

R U L E X X I I I .

(260.) *For the Solution of a Complete Equation of the Second Degree.*

1. Reduce the Equation to the form

$$x^2 + bx = s; \text{ in which } x^2 \text{ must be positive.}$$

2. Add the *square* of half the coefficient of x , in the second term, to each side of the equation:—the first side will then be a *perfect square*.

3. Extract the square root of each side, and the result will be a Simple Equation,—from which the value of x may readily be found.

E X A M P L E .

To find the value of x in the equation

$$2x + \frac{3x-5}{2} = 5x - \frac{5x-5}{x-3}.$$

Clearing the equation of its fractions, transposing, and adding similar terms, we shall find

$$-3x^2 + 14x = -5.$$

Dividing both sides of this equation by -3 ,

$$x^2 - \frac{14x}{3} = \frac{5}{3}.$$

Adding the *square of half the coefficient* of x , in the second term, to each side,—which is called *completing the square*,

$$x^2 - \frac{14x}{3} + \frac{49}{9} = \frac{5}{3} + \frac{49}{9} = \frac{64}{9}.$$

Extracting the *square root* of each side,

$$x - \frac{7}{3} = \pm \frac{8}{3}.$$

Whence $x = \frac{7}{3} + \frac{8}{3} = 5$; or $x = \frac{7}{3} - \frac{8}{3} = -\frac{1}{3}$. (255).

Either of these two values of x , 5 or $-\frac{1}{3}$, will satisfy the given equation; and each of the binomials $x-5$ and $x+\frac{1}{3}$ will divide the equation

$$x^2 - \frac{14x}{3} - \frac{5}{3} = 0. \quad (253).$$

By performing the division it will be found that the left hand side of this equation, is the *product of the two binomials*; that is,

$$(x-5)(x+\frac{1}{3}) = x^2 - \frac{14x}{3} - \frac{5}{3} = 0.$$

On the preceding Rule we remark,

1. The method of *completing the square* in the first member, results from the composition of the *square* of a Binomial.

Thus the square of the binomial $a+b$ is $a^2+2ba+b^2$, in which b is half the coefficient of a in the second term.

2. If in reducing the equation to the required form, the first term should become $-x^2$, the signs of all the terms in the equation must be changed (117), before completing the square,—otherwise the root of the first side would be *imaginary*, (246).

3. The *square root* of the first side of the equation—after the completion of the square—will always be the square root of the *first term*, $+$ or $-$ *half the coefficient* of x in the second term, according as the second term is $+$ or $-$.

Hence, *without completing the square on the first side*, we may shorten the solution, by observing that

4. In an equation of the form $x^2+bx=s$, the value of x is half the co-efficient of x in the second term, taken with a *contrary sign*, $+$ the square root of (the *second member* of the equation $+$ the square of said half co-efficient).

Thus from the equation $x^2 - \frac{14x}{3} = \frac{5}{3}$, we immediately find

$$x = \frac{7}{3} + \sqrt{\frac{5}{3} + \frac{49}{9}} = \frac{7}{3} + \sqrt{\frac{64}{9}} = \frac{7}{3} + \frac{8}{3}$$

EXERCISES

On Quadratic Equations with One Unknown Quantity.

1. Find the value of x in the equation

$$x^2 - 15 = 45 - 4x. \quad \text{Ans. } x = 6 \text{ or } -10.$$
2. Find the value of x in the equation

$$x^2 + 10 = 65 + 6x. \quad \text{Ans. } x = 11 \text{ or } -5.$$
3. Find the value of x in the equation

$$2x^2 + 8x - 30 = 60. \quad \text{Ans. } x = 5 \text{ or } -9.$$
4. Find the value of x in the equation

$$3x^2 - 3x + 9 = 8\frac{1}{3}. \quad \text{Ans. } x = \frac{1}{3} \text{ or } \frac{2}{3}.$$
5. Find the value of x in the equation

$$5x^2 + 4x - 90 = 114. \quad \text{Ans. } x = 6 \text{ or } -6\frac{4}{5}.$$
6. Find the value of x in the equation

$$\frac{1}{2}x^2 - \frac{1}{4}x + 2 = 9. \quad \text{Ans. } x = 4 \text{ or } -3\frac{1}{2}.$$
7. Find the value of x in the equation

$$2x^2 + 12x + 36 = 356. \quad \text{Ans. } x = 10 \text{ or } -16.$$
8. Find the value of x in the equation

$$4x - 46 = \frac{36 - x}{x}. \quad \text{Ans. } x = 12 \text{ or } -\frac{3}{4}.$$
9. Find the value of x in the equation

$$8x^2 + 6 = 7x + 171. \quad \text{Ans. } x = 5 \text{ or } -4\frac{1}{3}.$$
10. Find the value of x in the equation

$$5x - 23 = \frac{25 - 3x}{x}. \quad \text{Ans. } x = 5 \text{ or } -1.$$
11. Find the value of x in the equation

$$3x^2 + 6 = 3x + 5\frac{1}{3}. \quad \text{Ans. } x = \frac{1}{3} \text{ or } \frac{2}{3}.$$
12. Find the value of x in the equation

$$\frac{x^2}{2} - \frac{x}{3} + 20\frac{1}{2} = 42\frac{2}{3}. \quad \text{Ans. } x = 7 \text{ or } -6\frac{1}{3}.$$
13. Find the value of x in the equation

$$\frac{8 - x}{2} = \frac{x - 2}{6} + \frac{2x - 11}{x - 3}. \quad \text{Ans. } x = 6 \text{ or } \frac{1}{2}.$$
14. Find the value of x in the equation

$$x + 4 = 13 - \frac{7x - 8}{x}. \quad \text{Ans. } x = 4 \text{ or } -2.$$
15. Find the value of x in the equation

$$\frac{x - 3a}{b} = \frac{9(b - a)}{x}. \quad \text{Ans. } x = 3b, \text{ or } 3(a - b)$$

Another Method of Solving Quadratic Equations.

(261.) A Binomial of the form ax^2+bx will be made a *perfect square* by multiplying it by 4 *times* the coefficient of x^2 , and adding the square of the given coefficient of x .

$$\text{Thus } (ax^2+bx)4a+b^2=4a^2x^2+4abx+b^2=(2ax+b)^2, \quad (59).$$

To apply this principle to solution of the Equation

$$3x^2-2x=65.$$

Multiplying both sides of the equation by 4 *times* the coefficient 3, and adding the square of the coefficient 2 to both sides, we have

$$36x^2-24x+4=780+4=784.$$

Extracting the square root of both sides, we find

$$6x-2=\pm 28; \text{ which gives } x=\frac{\pm 28+2}{6}=5 \text{ or } -4\frac{1}{3}.$$

The *square root of the first side* of the Equation prepared as above, will be x multiplied into *twice* the given coefficient of x^2 , + or - the given coefficient of x in the second term, according as this term is + or -. This is evident from the preceding illustration, (261.)

From these principles we derive

R U L E X X I V .

(262.) *To Reduce an Equation of the form $ax^2+bx=s$ to a Simple Equation.*

1. Double the coefficient of x^2 , and divide the first member by x .
2. Multiply the second member by 4 *times* the given coefficient of x^2 , add the square of the given coefficient of x , and extract the square root of the sum.

Applying this Rule to the numerical equation

$$3x^2-5x=50,$$

we immediately obtain $6x-5=\pm\sqrt{50\times 12+25}=\pm\sqrt{625}.$

When the coefficient of x^2 is *unity*, the multiplier of the second member will be simply 4; and when the coefficient of x is *unity*, the square to be added after multiplying the second member is 1.

This method of solving a Quadratic is preferable to the one first given, whenever the coefficient of x would give rise to a *fraction* in dividing the Equation by the coefficient of x^2 , or in *completing the square*, according to that method.

16. Find the value of x in the equation
 $3x^2 + 2x - 9 = 76.$ *Ans.* $x = 5$, or $-5\frac{2}{3}.$
17. Find the value of x in the equation
 $2x^2 - 14x + 2 = 18.$ *Ans.* $x = 8$, or $-1.$
18. Find the value of x in the equation
 $x^2 - 12x + 50 = 0.$ *Ans.* $x = 6 \pm \sqrt{-14}.$
19. Find the value of x in the equation
 $\frac{1}{2}x^2 - 12\frac{1}{2} = \frac{1}{2}x + 15\frac{1}{2}.$ *Ans.* $x = 8$, or $-7.$
20. Find the value of x in the equation
 $3x^2 - \frac{3}{4} = 5\frac{1}{4} + 2x.$ *Ans.* $x = \frac{1}{3} \pm \frac{1}{3}\sqrt{19}.$
21. Find the value of x in the equation
 $x^2 - \frac{5x}{6} + \frac{1}{6} = 0.$ *Ans.* $x = \frac{1}{2}$, or $\frac{1}{3}.$
22. Find the value of x in the equation
 $x + 4 = 13 - \frac{7x - 8}{x}.$ *Ans.* $x = 4$, or $-2.$
23. Find the value of x in the equation
 $\frac{x^2}{3} + \frac{4x}{5} - 34\frac{1}{5} = 0.$ *Ans.* $x = 9$, or $-11\frac{2}{5}.$
24. Find the value of x in the equation
 $x^2 - \frac{x}{2} = \frac{2}{3} - \frac{x}{3}.$ *Ans.* $x = \frac{1}{12} \pm \frac{1}{12}\sqrt{97}.$
25. Find the value of x in the equation
 $\frac{10}{x} - \frac{14 - 2x}{x^2} = 2\frac{4}{9}.$ *Ans.* $x = 3$, or $1\frac{10}{11}.$
26. Find the value of x in the equation
 $-\frac{x^2}{4} + 3x + 3 = 13.$ *Ans.* $x = 6 \pm \sqrt{-4}.$
27. Find the value of x in the equation
 $x^2 + \frac{17x}{4} = -\frac{17x}{4} - 4.$ *Ans.* $x = -\frac{1}{2}$, or -8
28. Find the value of x in the equation,
 $5x - \frac{3x - 3}{x - 3} = 2x + \frac{3x - 6}{2}.$ *Ans.* $x = 4$, or $-1.$
29. Find the value of x in the equation
 $\frac{x + 4}{3} - \frac{7 - x}{x - 3} = -1 + \frac{4x + 7}{9}.$ *Ans.* $x = 21$, or 5
30. Find the value of x in the equation
 $\frac{x + a}{x} - \frac{3x}{x - a} = 0.$ *Ans.* $x = \pm a\sqrt{-\frac{1}{3}}.$

*Equations in which the Unknown Quantity is contained
in a Surd Expression.*

31. Find the value of x in the equation

$$5 + \sqrt{x^3 + 36} = 15.$$

By transposition, $\sqrt{x^3 + 36} = 15 - 5 = 10.$

Squaring both sides, $x^3 + 36 = 100$; (257).

from which, $x^3 = 64.$

We have now a *pure cubic* equation, in which x has necessarily *three* values or *roots*, (255).

Extracting the *cube root* of each side of the last equation, we find
 $x = 4$, which is one value of x .

To find the other two values of x , we must reduce the *cubic* equation to a *quadratic* by division, (253).

Dividing each side of the equation $x^3 - 64 = 0$ by $x - 4$, we find

$$x^2 + 4x + 16 = 0, \text{ or } x^2 + 4x = -16; \text{ hence } x = -2 \pm \sqrt{-12}.$$

We have thus found $x = 4$, or $-2 + \sqrt{-12}$, or $-2 - \sqrt{-12}$; the first value being *real*, the other two *imaginary*.

Each of these imaginary values of x , as well as the *real* value 4, will satisfy the equation $x^3 = 64$.

Thus x being equal to $-2 + \sqrt{-12}$, we have, (247), (249),

$$x^2 = (-2 + \sqrt{-12})^2 = 4 - 4\sqrt{-12} - 12 = -8 - 4\sqrt{-12}.$$

and $x^3 = (-8 - 4\sqrt{-12})(-2 + \sqrt{-12}) = 16 + 48 = 64,$

In like manner the other imaginary value of x may be verified.

32. Find the value of x in the equation

$$6 + \sqrt{3x + 4} = 11. \qquad \text{Ans. } x = 7.$$

33. Find the value of x in the equation

$$24 - \sqrt{2x^2 + 9} = 15. \qquad \text{Ans. } x = \pm 6.$$

34. Find the value of x in the equation

$$20 - \sqrt{x^3 + 40} = 4$$

Ans. $x = 6$ or $-3 + \sqrt{-27}$, or $-3 - \sqrt{-27}.$

35. Find the value of x in the equation

$$x + \sqrt{x} = 20.$$

From this equation we shall find $x=25$ or 16 . The value 25 will not satisfy the original equation if the square root of x be restricted to its *positive* value; but this root is $\pm\sqrt{x}$, and 25 satisfies the equation for the *negative* root.

Two values of x may be found in each of the next three Exercises; but only that value is given in the *Ans.* which corresponds to *positive* roots in the given Equation.

36. Find the value of x in the equation

$$2\sqrt{x} + \sqrt{x+9} = 13. \quad \text{Ans. } x=16.$$

37. Find the value of x in the equation

$$2\sqrt{x} + \sqrt{2x+1} = \frac{21}{\sqrt{(2x+1)}}. \quad \text{Ans. } x=4.$$

38. Find the value of x in the equation

$$4\sqrt{x+16} = 7\sqrt{x+16} - x - 6. \quad \text{Ans. } x=9$$

39. Find the value of x in the equation

$$\frac{x+4}{2} = \sqrt{4 + \sqrt{2x^3 + x^2}}. \quad \text{Ans. } x=12, \text{ or } 4.$$

40. Find the value of x in the equation

$$(x-2)^2 = \frac{x + \sqrt{(x^2-9)}}{x - \sqrt{(x^2-9)}}.$$

By *rationalizing* the denominator, (243...2), we shall find

$$(x-2)^2 = \frac{(x + \sqrt{(x^2-9)})^2}{9}.$$

Extracting the square root of each side of this equation, we have

$$x-2 = \frac{x + \sqrt{(x^2-9)}}{3}. \quad \text{Ans. } x=5, \text{ or } 3$$

41. Find the value of x in the equation

$$x^2 - 6x + 9 = \frac{x + \sqrt{(x^2-16)}}{x - \sqrt{(x^2-16)}}. \quad \text{Ans. } x=5, \text{ or } 4$$

42. Find the value of x in the equation

$$\sqrt[3]{x^3+37} \times (x^3+37)^{\frac{2}{3}} = 64. \quad \text{Ans. } x=3, \text{ or } -\frac{3}{2} \pm \frac{3}{2}\sqrt{-3}.$$

*Equations of a Quadratic Form with reference to a Power
or Root of the Unknown Quantity.*

(263.) Any Equation containing the unknown quantity x in but two terms—with its exponent in one *double* its exponent in the other—is a Quadratic with reference to the *lower power* of x ; and the value of such power may be found accordingly.

To find the value of x in the equation

$$x^{\frac{1}{2}} + 4x^{\frac{1}{4}} = 21.$$

The higher *fractional* power $x^{\frac{1}{2}}$ is the *square* of the lower $x^{\frac{1}{4}}$; and the equation is therefore *quadratic* with reference to $x^{\frac{1}{4}}$.

Completing the square, we have

$$x^{\frac{1}{2}} + 4x^{\frac{1}{4}} + 4 = 21 + 4 = 25.$$

Extracting the square root of each side,

$$x^{\frac{1}{4}} + 2 = \pm 5;$$

$$\text{from which } x^{\frac{1}{4}} = 3, \text{ or } -7.$$

By raising each of these values to the *4th power*, we find

$$x = 81, \text{ or } 2401.$$

The first of these two values of x is easily verified. In verifying the value 2401 it must be observed that its 4th root is -7 , and that $4x^{\frac{1}{4}}$ is therefore -28 .

When x is in a *fractional* power, in the following Exercises, only that value will be given in the *Ans.* which satisfies the given form of the Equation.—Imaginary values of x are also omitted.

43. Find the value of x in the equation

$$x^4 - 2x^2 + 6 = 230.$$

$$\text{Ans. } x = \pm 4.$$

44. Find the value of x in the equation

$$x^6 + 20x^3 - 10 = 59.$$

$$\text{Ans. } x = \sqrt[3]{3}.$$

45. Find the value of x in the equation

$$2x^4 - x^2 + 20 = 23.$$

$$\text{Ans. } x = \frac{1}{2}\sqrt{6}.$$

46. Find the value of x in the equation

$$3x^{\frac{1}{2}} - 5x^{\frac{1}{4}} = -1\frac{1}{3}.$$

$$\text{Ans. } x = 3\frac{13}{81}.$$

47. Find the value of x in the equation

$$6x^{\frac{4}{3}} - 5x^{\frac{8}{3}} + 1184 = 0.$$

$$\text{Ans. } x = 8$$

48. Find the value of x in the equation

$$(x+12)^{\frac{1}{2}} + (x+12)^{\frac{1}{4}} = 6.$$

In this equation we must regard the binomial $(x+12)$ as the unknown quantity; and to simplify the operation we may represent this binomial by y .

$$\begin{aligned} \text{Then } y^{\frac{1}{2}} + y^{\frac{1}{4}} &= 6 \\ \text{which gives } y^{\frac{1}{4}} &= 2 \text{ or } -3. \end{aligned}$$

By restoring the binomial value of y , we have

$$(x+12)^{\frac{1}{4}} = 2, \text{ or } -3. \quad \text{Ans. } x=4.$$

49. Find the value of x in the equation

$$(2x+6)^{\frac{1}{2}} - 6 = -(2x+6)^{\frac{1}{4}}. \quad \text{Ans. } x=5.$$

50. Find the value of x in the equation

$$x^2 + 11 + \sqrt{x^2 + 11} = 42. \quad \text{Ans. } x = \pm 5$$

51. Find the value of x in the equation

$$\frac{1}{(2x-4)^2} = \frac{1}{8} + \frac{2}{(2x-4)^4}. \quad \text{Ans. } x=3, \text{ or } 1.$$

52. Find the value of x in the equation

$$4x^4 - 4x^3 + \frac{x}{2} = 33.$$

This equation may be reduced to the form of a *quadratic*, thus.—The first two terms of the *square root* of the first side will be found to be

$$2x^2 - x; \text{ and the remainder will be } -x^2 + \frac{x}{2}, \text{ which may be put under the form } -\frac{1}{2}(2x^2 - x).$$

Now the *square* of the root found, and the *remainder*, are together equivalent to the first side of the equation; hence we have

$$(2x^2 - x)^2 - \frac{1}{2}(2x^2 - x) = 33. \quad \text{Ans. } x=2, \text{ or } -1\frac{1}{2}.$$

A *Biquadratic* equation may be reduced to the form of a *Quadratic*, as above, whenever the remainder—after having found the first two terms of the square root of the first side—can be resolved into two factors, one of which is the same as *the part of the root* thus found

53. Find the value of x in the equation

$$x^3 - 8x^2 + 19x = 12.$$

Transposing the 12, and multiplying by x , we have

$$x^4 - 8x^3 + 19x^2 - 12x = 0.$$

We have now a Biquadratic equation which may be reduced to the form of a Quadratic by the method just exemplified.

We find the first two terms of the square root of the first side of the Equation, by the common Rule.

$$\begin{array}{r} x^4 - 8x^3 + 19x^2 - 12x(x^2 - 4x) \\ x^4 \\ \hline 2x^2 - 4x \sqrt{-8x^3 + 19x^2} \\ \phantom{\sqrt{-8x^3 + 19x^2}} - 8x^3 + 16x^2 \\ \hline \phantom{\sqrt{-8x^3 + 19x^2}} 3x^2 - 12x. \end{array}$$

The remainder, $3x^2 - 12x$, may be resolved into $3(x^2 - 4x)$, in which the binomial factor is the same as the part of the square root above found.

$$\text{Then } (x^2 - 4x)^2 + 3(x^2 - 4x) = 0.$$

Ans. $x = 4, 3, \text{ or } 1.$

54. Find the value of x in the equation

$$x^4 - 2x^3 + x = 5112.$$

Ans. $x = 9$

55. Find the value of x in the equation

$$x^4 + 2x^3 - 7x^2 - 8x = -12.$$

Ans. $x = 2, -3, 1, \text{ or } -2$

56. Find the value of x in the equation

$$x^4 - 10x^3 + 35x^2 - 50x + 24 = 0.$$

Ans. $x = 1, 2, 3, \text{ or } 4.$

57. Find the value of x in the equation

$$x^4 - 12x^3 + 44x^2 - 48x = 9009.$$

Ans. $x = 13$

58. Find the value of x in the equation

$$x^4 - 8ax^3 + 8a^2x^2 + 32a^3x = s.$$

Ans. $x = 2a \pm \sqrt{8a^2 \pm \sqrt{s + 16a^4}}.$

PROBLEMS

In Pure Equations and Affected Quadratics containing but One Unknown Quantity.

1. Find two numbers such that their product shall be 750, and the quotient of the greater divided by the less, $3\frac{1}{3}$.

Let x represent the greater of the two numbers ;
then will $\frac{750}{x}$ represent the less ; and the Equation will be

$$x \div \frac{750}{x} \text{ or } \frac{x^2}{750} = 3\frac{1}{3}.$$

From this equation we shall find $x=50$, or -50 . Each of these values will satisfy the Equation of the problem ; but only the *positive* one can be taken to answer the conditions of the problem itself, in which the required numbers are understood to be *positive*, as in the problems of common Arithmetic. *Ans.* 50, and 15.

2. Find a number such that if $\frac{1}{7}$ and $\frac{1}{8}$ of it be multiplied together, and the product divided by 3, the quotient will be $298\frac{2}{3}$. *Ans.* 224.

3. A mercer bought a piece of silk for £16 4s. ; and the number of shillings that he paid per yard, was to the number of yards, as 4 to 9. How many yards did he buy ? and what was the price per yard ?

Let x represent the number of shillings he paid per yard ;

then $4 : 9 :: x : \frac{9x}{4}$, the number of yards.

But without forming a Proportion, the number of yards is readily known to be $\frac{9}{4}$ of the price per yard.

Ans. 27 yards, at 12s. per yard.

4. Find two numbers which shall be to each other as 2 to 3, and the sum of whose squares shall be 208. *Ans.* 8 and 12.

5. A person bought a quantity of cloth for \$120 ; and if he had bought 6 yards more for the same sum, the price per yard would have been \$1 less. What was the number of yards ? and the price per yard ?
Ans. 24 yards, at \$5 per yard.

6. Divide the number 20 into two such parts that the squares of these parts may be in the proportion of 4 to 9. *Ans.* 8, and 12.

7. A merchant bought a quantity of flour for \$100, which he sold again at $\$5\frac{1}{4}$ per barrel, and in so doing gained as much as each barrel cost him. What was the number of barrels? *Ans.* 20.

8. Divide the number 800 into two such parts that the less divided by the greater, may be to the greater divided by the less, as 9 to 25.

Let x represent the less number :—we shall then have the Proportion

$$\frac{x}{800-x} : \frac{800-x}{x} :: 9 : 25;$$

which will be converted into an Equation by putting the product of the two *extremes* equal to the product of the two *means*.

Ans. 300, and 500

9. Two fields which differ in quantity by 10 acres, were each sold for \$2800, and one of them was valued at \$5 an acre more than the other. What was the number of acres in each? *Ans.* 70, and 80.

10. A and B started together on a journey of 150 miles. A traveled 3 miles an hour more than B, and completed the journey $8\frac{1}{3}$ hours before him. At what rate did each travel per hour?

Ans. 9, and 6 miles.

11. A man traveled 105 miles, and then found that if he had gone 2 miles less per hour, he would have been 6 hours longer on his journey. At what rate did he travel per hour? *Ans.* 7 miles.

12. A person has two pieces of silk which together contain 14 yards. Each piece is worth as many shillings per yard as there are yards in the piece, and their whole values are in the proportion of 9 to 16; how many yards are there in each piece? *Ans.* 6, and 8 yards.

13. A merchant sold a piece of linen for \$39, and in so doing gained as much per cent. as it cost him. What was the cost of the linen? *Ans.* \$30.

14. A grazier bought as many sheep as cost him \$100. After reserving 5 of the number, he sold the remainder for \$135, and gained \$1 a head on them; how many sheep did he buy? *Ans.* 50.

15. Find two numbers which shall be in the proportion of 7 to 9, and have the difference of their squares equal to 128.

Ans. 14, and 18.

16. An officer would arrange 2400 men in a solid body, so that each rank may exceed each file by 43 men. How many must be placed in rank and file? *Ans.* 75, and 32.

17. Two partners gained £18 by trade. A's money was employed in the business 12 months; and B's, which was £30, 16 months. A received for his capital and gain £26; what was the amount of his capital?

Let x represent A's capital; then $26 - x$ will be his gain; and since the gain is in the *compound ratio* of the *capital* and the *time* it was employed, we have

$$12x + 16 \times 30 : 12x :: 18 : 26 - x.$$

The first ratio in this proportion may be simplified by dividing the antecedent and the consequent by 12, (158). *Ans.* £20.

18. A detachment from an army was marching in regular column, with 5 men more in depth than in front; but upon the enemy's coming in sight, the front was increased by 845 men; and by this movement the detachment was drawn up in five lines. What was the number of men? *Ans.* 4550.

19. A company at a tavern had £8 15s. to pay, but before their bill was settled, two of them went away, when those who remained had 10s. apiece more to pay than before. How many were there in the company at first? *Ans.* 7.

20. Some gentlemen made an excursion, and each one took the same sum. Each gentleman had as many servants as there were gentlemen, and the number of dollars which each had was double the whole number of servants; also the whole sum taken with them was \$3456. What was the number of gentlemen? *Ans.* 12.

21. Divide the number 20 into two such parts, that the product of the whole number and one of the parts shall be equal to the square of the other. *Ans.* $10\sqrt{5} - 10$, and $30 - 10\sqrt{5}$.

22. A laborer dug two trenches, one of which was 6 yards longer than the other, for £17 16s, and the digging of each cost as many shillings per yard as there were yards in its length. What was the length of each? *Ans.* 10, and 16 yards.

23. There are two numbers whose product is 120, and if 2 be added to the less, and 3 subtracted from the greater, the product of the sum and the remainder will also be 120. What are the two numbers? *Ans.* 8 and 15.

24. Two persons lay out some money on speculation. A disposes of his bargain for £11, and gains as much per cent. as B lays out; B's gain is £36, and it appears that A gains 4 times as much per cent. as B. What sum did each lay out? *Ans.* A £5, B £120.

25. A set out from C towards D, and traveled 7 miles an hour. After he had gone 32 miles, B set out from D towards C, and went each hour $\frac{1}{19}$ of the whole distance; and after he had traveled as many hours as he went miles in one hour, he met A. Required the distance between the two places. *Ans.* 152, or 76 miles.

SOLUTION OF TWO EQUATIONS—ONE OR BOTH OF THE SECOND OR A HIGHER DEGREE—CONTAINING TWO UNKNOWN QUANTITIES.

(264) For the solution of two Equations, containing two unknown quantities, the method which naturally occurs is,

1. By elimination between the given equations to derive a *new equation* containing but one of the unknown quantities, and thence to find the *value* of that quantity.

2. By substituting this quantity for its symbol in one of the equations containing the other unknown quantity, to determine thence the value of that quantity.

There are, however, some facilitating expedients to be applied, in certain cases, to equations of the second and higher degrees: these will be exemplified as we proceed.

But the solution of two Equations—one or both of the second or a higher degree—containing two unknown quantities, may be impossible by the *method of quadratics*,—from the impossibility of deriving from them a new equation containing but one unknown quantity, which will admit of a quadratic solution

EXAMPLES AND EXERCISES.

1. Find the values of x and y in the equations

$$2x + y = 10, \text{ and } 2x^2 - xy + 3y^2 = 54.$$

From the first equation, we have

$$x = \frac{10 - y}{2}$$

$$\text{Then } 2x^2 = \frac{2(100 - 20y + y^2)}{4}, \text{ and } xy = \frac{10y - y^2}{2}.$$

By substituting these values in the second equation, we find

$$\frac{2(100 - 20y + y^2)}{4} - \frac{10y - y^2}{2} + 3y^2 = 54.$$

The value of y may be found from this equation; and by substituting the value of y for y in the first equation, the value of x may readily be determined.

Observe that the left hand fraction in the last equation may be reduced to *lower terms*; and the solution of the equation be thus somewhat simplified.

$$\text{Ans. } x = 3, \text{ or } \frac{41}{8}; y = 4, \text{ or } -\frac{1}{4}.$$

2. Find the values of
- x
- and
- y
- in the equations

$$x+y=9, \text{ and } x^2+y^2=45.$$

$$\text{Ans. } x=3, \text{ or } 6; y=6, \text{ or } 3$$

☞ Whenever x and y may be interchanged with each other, without changing the *form* of the given equations—as in the preceding example—the two values of one of these letters may be taken, in reverse order, for the two values of the other.

3. Find the values of
- x
- and
- y
- in the equations

$$xy=28, \text{ and } x^2+y^2=65.$$

$$\text{Ans. } x=\pm 7, \text{ or } \pm 4; y=\pm 4, \text{ or } \pm 7.$$

4. Find the values of
- x
- and
- y
- in the equations

$$x+4y=14, \text{ and } 4x-2y+y^2=11.$$

$$\text{Ans. } x=2, \text{ or } -46; y=3, \text{ or } 15.$$

5. Find the values of
- x
- and
- y
- in the equations

$$x+2y=7, \text{ and } x^2+3xy-y^2=23.$$

$$\text{Ans. } x=3, \text{ or } 15\frac{2}{3}; y=2, \text{ or } -4.$$

6. Find the values of
- x
- and
- y
- in the equations

$$x+\frac{y}{2}=11, \text{ and } xy+2y^2=120.$$

$$\text{Ans. } x=8, \text{ or } 17\frac{2}{3}; y=6, \text{ or } -13\frac{1}{3}.$$

7. Find the values of
- x
- and
- y
- in the equations

$$x-y=-2, \text{ and } x+\frac{y}{10}=\frac{3xy}{10}$$

$$\text{Ans. } x=2, \text{ or } -\frac{1}{3}; y=4, \text{ or } 1\frac{2}{3}.$$

8. Find the values of
- x
- and
- y
- in the equations

$$x-\frac{x-y}{2}=4, \text{ and } y-\frac{x+3y}{x+2}=1.$$

$$\text{Ans. } x=2, \text{ or } 5; y=6, \text{ or } 3.$$

9. Find the values of
- x
- and
- y
- in the equations

$$5(x+y)=13(x-y), \text{ and } x+y^2=25.$$

$$\text{Ans. } x=9, \text{ or } -14\frac{1}{16}; y=4, \text{ or } -6\frac{1}{4}.$$

Solutions by Means of an Auxiliary Unknown Quantity.

(265.) When the number of *unknown factors* is the same in every unknown term of the two Equations, the solution will often be facilitated by substituting for one of the unknown quantities the *product of the other into a third unknown quantity*.

10. Find the values of x and y in the equations

$$x^2 + xy = 54, \text{ and } 2xy + y^2 = 45.$$

The number of unknown *factors* in each of the unknown terms in these equations, is *two*.

If we assume x to be equal to vy , and substitute this product for x , the given equations will become

$$v^2y^2 + vy^2 = 54,$$

$$\text{and } 2vy^2 + y^2 = 45.$$

From the first of these equations we have

$$y^2 = \frac{54}{v^2 + v} \text{ and from the second } y^2 = \frac{45}{2v + 1}.$$

Putting these two values of y^2 equal to each other,

$$\frac{54}{v^2 + v} = \frac{45}{2v + 1}.$$

By solving this equation in the usual manner we shall find $v = 2$, or $-\frac{3}{5}$; then by substituting these values, successively, in either of the expressions for y^2 , we shall find the values of y ; and since $x = vy$, we may also readily obtain the value of x .

$$\text{Ans. } x = \pm 6, \text{ or } \mp 9\sqrt{-1}; \quad y = \pm 3, \text{ or } \pm 15\sqrt{-1}.$$

If in the preceding example an expression for the value of x or y were obtained from either equation, and substituted in the other, the resulting equation, when cleared of radical signs, would be of the *fourth degree*; and we have accordingly found *four values* for each of the unknown quantities, (255.)

11. Find the values of x and y in the equations

$$xy = 28, \text{ and } x^2 + y^2 = 65.$$

$$\text{Ans. } x = \pm 7, \text{ or } \pm 4; \quad y = \pm 4, \text{ or } \pm 7.$$

12. Find the values of x and y in the equations

$$x^2 + xy = 12, \text{ and } xy - 2y^2 = 1.$$

$$\text{Ans. } x = \pm 3, \text{ or } \pm \frac{8}{\sqrt{6}}; \quad y = \pm 1, \text{ or } \pm \frac{1}{\sqrt{6}}$$

13. Find the values of x and y in the equations

$$4x^2 - 2xy = 12, \text{ and } 2y^2 + 3xy = 8.$$

$$\text{Ans. } x = \pm 2, \text{ or } \mp 3\sqrt{\frac{1}{7}}; y = \pm 1, \text{ or } \pm 8\sqrt{\frac{1}{7}}.$$

14. Find the values of x and y in the equations

$$3y^2 - x^2 = 39, \text{ and } x^2 + 4xy = 256 - 4y^2.$$

$$\text{Ans. } x = \pm 6, \text{ or } \pm 102; y = \pm 5, \text{ or } \mp 59.$$

Solutions by Means of Two Auxiliary Unknown Quantities.

(266.) When the unknown quantities are *similarly involved* in each of the two Equations, the solution will sometimes be facilitated by substituting for the two unknown quantities the *sum and difference of two other unknown quantities*.

15. Find the values of x and y in the equations

$$x + y = 12, \text{ and } \frac{x^2}{y} + \frac{y^2}{x} = 18.$$

If we assume x equal to $v + z$, and y equal to $v - z$, we shall have

$$x + y = 2v = 12; \text{ and hence } v = 6.$$

Then $x = 6 + z$, and $y = 6 - z$.

Substituting these values of x and y in the second equation, we have

$$\frac{(6+z)^2}{6-z} + \frac{(6-z)^2}{6+z} = 18.$$

Clearing this equation of its fractions,

$$(6+z)^3 + (6-z)^3 = 18(36 - z^2).$$

By developing both sides of this last equation, and proceeding with the solution in the usual manner, we shall find $z = \pm 2$.

Having now found the values of both v and z , the values of x and y are easily obtained.

$$\text{Ans. } x = 8 \text{ or } 4; y = 4 \text{ or } 8.$$

16. Find the values of x and y in the equations

$$x + y = 10, \text{ and } x^3 + y^3 = 280.$$

$$\text{Ans. } x = 4 \text{ or } 6; y = 6 \text{ or } 4.$$

17. Find the values of x and y in the equations

$$x + y = 11, \text{ and } x^4 + y^4 = 2657.$$

$$\text{Ans. } x = 4 \text{ or } 7, y = 7 \text{ or } 4$$

18. Find the values of x and y in the equations

$$x + y = 10, \text{ and } x^5 + y^5 = 17,050.$$

$$\text{Ans. } x = 3 \text{ or } 7, y = 7 \text{ or } 3$$

Miscellaneous Solutions and Exercises.

19. Find the values of a and y in the equations

$$xy=6, \text{ and } x^2+x=18-y^2-y.$$

From the second equation,

$$x^2+y^2+x+y=18.$$

Adding *twice* the first,

$$x^2+2xy+y^2+x+y=30.$$

This last equation may be put under the form

$$(x+y)^2+(x+y)=30;$$

which is *quadratic* with reference to $x+y$, and from which we may therefore find the value of $x+y$.

$$\text{Ans. } x=2, \text{ or } 3; \text{ or } -3 \mp \sqrt{3}; y=3, \text{ or } 2; \text{ or } -3 \pm \sqrt{3}.$$

20. Find the values of x and y in the equations

$$x+y=6, \text{ and } x^2y^2+4xy=96.$$

$$\text{Ans. } x=2, \text{ or } 4; \text{ or } 3 \mp \sqrt{21}; y=4, \text{ or } 2; \text{ or } 3 \pm \sqrt{21}$$

21. Find the values of x and y in the equations

$$x^{\frac{2}{3}}y^{\frac{3}{2}}=2y^2, \text{ and } 8x^{\frac{1}{3}}-y^{\frac{1}{2}}=14.$$

Dividing the first equation by $y^{\frac{3}{2}}$, we find $x^{\frac{2}{3}}=2y^{\frac{1}{2}}$, or $y^{\frac{1}{2}}=\frac{1}{2}x^{\frac{2}{3}}$, and by substituting this value of $y^{\frac{1}{2}}$ in the second equation, that equation will become *quadratic* with reference to $x^{\frac{1}{3}}$.

$$\text{Ans. } x=2744, \text{ or } 8; y=9604, \text{ or } 4.$$

22. Find the values of x and y in the equations

$$x^2y-xy=6, \text{ and } x^3y-y=21.$$

Dividing the first equation by the second, we have

$$\frac{x^2y-xy}{x^3y-y}=\frac{6}{21}.$$

This equation will be simplified by reducing each of its two fractional members to its *lowest terms*.

$$\text{Ans. } x=2, \text{ or } \frac{1}{2}; y=3, \text{ or } -24.$$

23. Find the values of x and y in the equations

$$x^3+xy^2=39, \text{ and } x^2y+y^3=26.$$

$$\text{Ans. } x=3; y=2$$

24. Find the values of x and y in the equations
 $x - y = 4$, and $x^3 - y^3 = 316$.

Dividing each side of the second equation by the corresponding side of the first, we shall find, $x^2 + xy + y^2 = 79$.

Squaring the first equation, and subtracting, we have $3xy = 63$

$$\text{Ans } x = 7, \text{ or } -3; y = 3, \text{ or } -7$$

25. Find the values of x and y in the equations
 $x^2 - y^2 = 5$, and $x^4 - y^4 = 65$.

$$\text{Ans. } x = \pm 3, y = \pm 2$$

26. Find the values of x and y in the equations
 $x + y = 60$, and $2(x^2 + y^2) = 5xy$.

The second equation may be put under the form $x^2 + y^2 - 2\frac{1}{2}xy = 0$, and the solution will be facilitated by subtracting this from the square of the first equation.

$$\text{Ans. } x = 40, \text{ or } 20; y = 20, \text{ or } 40.$$

27. Find the values of x and y in the equations

$$xy = 8, \text{ and } \frac{x^2}{y} + \frac{y^2}{x} = 9.$$

Multiplying the two equations together, we find $x^3 + y^3 = 72$; and, multiplying this equation by x^3 , we have $x^6 + x^3y^3 = 72x^3$.

From the first equation, $x^3y^3 = 8^3 = 512$; and if this number be substituted in the preceding equation we shall have a *quadratic* with reference to x^3 .

$$\text{Ans. } x = 4, \text{ or } 2; y = 2, \text{ or } 4.$$

28. Find the values of x and y in the equations

$$xy = 25, \text{ and } x^3 + y^3 = 10xy.$$

The second equation will be reduced to the same form as the second in the preceding example, by dividing it by xy .

$$\text{Ans. } x = 5, y = 5.$$

29. Find the values of x and y in the equations

$$x^2 - y^2 - (x + y) = 8, \text{ and } (x - y)^2(x + y) = 32.$$

Dividing each equation by $x + y$, we have

$$x - y - 1 = \frac{8}{x + y}, \text{ and } (x - y)^2 = \frac{32}{x + y}.$$

Transposing -1 , and squaring, we obtain

$$(x - y)^2 = \left(\frac{8}{x + y} + 1 \right)^2 = \frac{32}{x + y};$$

from which will result a *quadratic* with reference to $x + y$.

$$\text{Ans. } x = 5; y = 3.$$

30. Find the values of x and y in the equations

$$x + \sqrt{xy} + y = 7, \text{ and } x^2 + xy + y^2 = 21.$$

Dividing the second equation by the first, we have

$$x - \sqrt{xy} + y = 3.$$

We shall now obtain two equations of simpler forms by adding the third equation to the first, and subtracting it from the first.

$$\text{Ans. } x=1, \text{ or } 4; y=4, \text{ or } 1$$

31. Find the values of x and y in the equations

$$\frac{x+y}{2} = \sqrt{xy} + 4, \text{ and } \sqrt{xy} = \frac{2xy}{x+y} + \frac{12}{5}.$$

From the first equation, by transposition,

$$\sqrt{xy} = \frac{x+y}{2} - 4.$$

Clearing this equation of its fraction, and squaring,

$$4xy = (x+y)^2 - 16(x+y) + 64.$$

By equating the two values of \sqrt{xy} , from the second and third equations,

$$\frac{2xy}{x+y} + \frac{12}{5} = \frac{x+y}{2} - 4;$$

$$\text{or } \frac{2xy}{x+y} = \frac{x+y}{2} - \frac{32}{5}.$$

Clearing this equation of its fractions,

$$4xy = (x+y)^2 - \frac{64}{5}(x+y).$$

We shall now obtain a simple equation by subtracting this last equation from the fourth equation.

$$\text{Ans. } x=2, \text{ or } 18; y=18, \text{ or } 2.$$

32. Find the values of x and y from the equation and proportion

$$\begin{aligned} xy^2 - x &= 3; \\ x^2 y^4 - x^2 : x^2 + x^2 y^2 + x^2 y^4 &:: 5 : 7. \end{aligned}$$

In any proportion the difference of the first and second terms is to the first, as the difference of the third and fourth is to the third, (160).

Hence from the given proportion, we shall find

$$2x^2 + x^2 y^2 : x^2 y^4 - x^2 :: 2 : 5.$$

Dividing the first antecedent and consequent by x^2 , (158)

$$2 + y^2 : y^4 - 1 :: 2 : 5.$$

$$\text{Ans. } x=1, \text{ or } -\frac{6}{5}; y=2, \text{ or } \sqrt{-\frac{3}{2}}$$

33. Find the values of x and y in the equations

$$x^2 + xy + y^2 = 13, \text{ and } x^4 + x^2y^2 + y^4 = 91.$$

Dividing the second equation by the first,

$$x^2 - xy + y^2 = 7.$$

Adding the third equation to the first, and dividing the result by 2: and also subtracting the third from the first,

$$x^2 + y^2 = 10;$$

$$\text{and } 2xy = 6.$$

The solution now proceeds by adding the latter of these two equations to the former, and also subtracting the latter from the former, and extracting the *square roots* of the resulting equations.

$$\text{Ans. } x = \pm 3; y = \pm 1.$$

34. Find the values of x and y in the equations

$$x^2y = x^2y^2 - x^2, \text{ and } x^2y^2 + x^2 = x^3y^3 - x^3$$

Dividing each equation by x^2 , we have

$$y = y^2 - 1, \text{ and } y^2 + 1 = xy^3 - x.$$

The value of y is to be found from the first of these equations, and substituted in the second.

$$\text{Ans. } x = \frac{1}{2}\sqrt{5}; y = \frac{1}{2} \pm \frac{1}{2}\sqrt{5}.$$

35. Find the values of x and y in the equations

$$xy = x^2 - y^2, \text{ and } x^2 + y^2 = x^3 - y^3.$$

If we assume y to be equal to xv , and substitute this product for y in the two equations, (264,) we shall have

$$x^2v = x^2 - x^2v^2, \text{ and } x^2 + x^2v^2 = x^3 - x^3v^3.$$

These equations may be solved in the same manner as those in the preceding example. Or the value of x may be found, in terms of y , from the first of the given equations, and substituted in the second.

$$\text{Thus } x = \frac{y}{2} \pm \sqrt{\frac{5y^2}{4}} = \frac{y}{2} \pm \frac{y}{2}\sqrt{5}.$$

Then for the second equation,

$$\left(\frac{y}{2} \pm \frac{y}{2}\sqrt{5}\right)^2 + y^2 = \left(\frac{y}{2} \pm \frac{y}{2}\sqrt{5}\right)^3 - y^3.$$

$$\text{Ans. } x = \frac{1}{4}(5 \pm \sqrt{5}); y = \pm \frac{1}{2}\sqrt{5}.$$

36. Find the values of x and y in the equations

$$x^3 - x^2y - xy^2 + y^3 = 576, \text{ and } x^3 + x^2y + xy^2 + y^3 = 2336$$

Subtracting the first equation from the second,

$$2x^2y + 2xy^2 = 1760.$$

Adding this to the second equation, we obtain

$$x^3 + 3x^2y + 3xy^2 + y^3 = 4096.$$

By extracting the *cube root* of this equation, we shall find the value of $x+y$, which may be substituted for $x+y$ in the third equation.

$$\text{Ans. } x=11; y=5$$

37. Find the values of x and y in the equations

$$xy=320, \text{ and } x^3 - y^3 = 61(x-y)^3.$$

Dividing the second equation by $x-y$, we have

$$x^2 + xy + y^2 = 61(x-y)^2.$$

By converting this equation into a *proportion*, (153,)

$$x^2 + xy + y^2 : (x-y)^2 :: 61 : 1.$$

The solution now proceeds by developing the term $(x-y)^2$,—subtracting each consequent from its antecedent, and forming a proportion of the *consequents* and *remainders*, &c, (160).

$$\text{Ans. } x=20; y=16.$$

38. Find the values of x , y , and z , in the equations,

$$x^2 + y^2 + xy = 37; x^2 + z^2 + xz = 49; y^2 + z^2 + yz = 61.$$

Subtracting the first equation from the second, and *decomposing*,

$$(z-y)(z+y) + (z-y)x, \text{ or } (z+y+x)(z-y) = 12;$$

$$\text{from which } z+y+x = \frac{12}{z-y}.$$

Proceeding in like manner with the second and third equations,

$$\text{we shall find } y+x+z = \frac{12}{y-x}.$$

Hence the right hand members of the last two equations are equal to each other. (113 1); and since the numerators are the same, we have

$$z-y = y-x, \text{ from which } 2y = x+z.$$

By substituting $2y$ for $x+z$ in the sixth equation, we shall find

$$y^2 - yx = 4.$$

The value of x from this equation, is to be substituted in the first equation.

$$\text{Ans. } x=3; y=4; z=5.$$

PROBLEMS

In Quadratic Equations of One or More Unknown Quantities.

1. The *area* of a rectangular lot of ground is 384 square rods, and its length is to its breadth as 3 is to 2. Required the length and breadth of the lot.

The *area*, in square measure, of a rectangle, is expressed by the product of the number of *linear* units in its *length* \times the number of linear units in its *breadth*.

The length and breadth must be taken in the same denomination in multiplying: the *area* will be found in the corresponding denomination of *square measure*.

Let x represent the *length*, and y the *breadth* of the lot; then by the conditions of the problem,

$$xy = 384,$$

$$\text{and } x : y :: 3 : 2.$$

Or, if x represent the length, $\frac{2x}{3}$ will represent the breadth, and we shall then have

$$\frac{2x^2}{3} = 384. \quad \text{Ans. 24, and 16 rods.}$$

2. The length of a rectangular garden exceeds its breadth by 6 rods, and its area is 216 square rods. What are the length and breadth of the garden? Ans. 18, and 12 rods.

3. Find two numbers whose sum shall be 24, and whose product shall be equal to 35 times their difference. Ans. 14 and 10.

4. Divide a line 20 inches in length into two such parts that the rectangle or product of the whole line and one of the parts shall be equal to the square of the other part.

$$\text{Ans. } 10\sqrt{5} - 10, \text{ and } 30 - 10\sqrt{5}.$$

5. Find the dimensions of a rectangular field, so that its length shall be equal to twice its breadth, and its area 800 square rods.

$$\text{Ans. 40, and 20 rods.}$$

6. The sum of the two digits of a certain number is 10, and if their product be increased by 40, the digits will be reversed. What is the number? Ans. 46.

7. The sum of two fractions is $1\frac{1}{6}$, and the sum of their *reciprocals* is $3\frac{1}{2}$; what are the two fractions? *Ans.* $\frac{1}{2}$ and $\frac{2}{3}$.

8. The *perimeter*, or sum of the four sides of a rectangle, is 112 rods, and its area is 720 square rods. What are the length and breadth of the rectangle? *Ans.* 36, and 20 rods.

9. Divide the number 60 into two such parts that their product shall be to the sum of their squares as 2 to 5. *Ans.* 20 and 40

10. A merchant bought a piece of cloth for \$120, and after cutting off 4 yards, sold the remainder for what the whole cost him—by which he made \$1 a yard on what he sold. How many yards did the piece contain? *Ans.* 24.

11. Divide the number 100 into two such parts that the difference of their square roots shall be 2. *Ans.* 64, and 36.

12. A garden which is 20 rods square is surrounded by a walk whose area is equal to $\frac{1}{4}$ of the area of the garden itself. What is the breadth of the walk? *Ans.* $5\sqrt{5}-10$ rods.

13. The sum of the squares of two numbers is 325, and the difference of their squares is 125. What are the numbers? *Ans.* 15 and 10.

14. The area of a rectangular court-yard is 875 square rods, and if its length and breadth were each increased by 5 rods, its area would then be 1200 square rods. What are the dimensions of the yard? *Ans.* 35 and 25 rods.

15. The difference of two numbers is 4, and the difference of their cubes is 448. What are the two numbers? *Ans.* 8 and 4.

16. A grocer sold 80 pounds of mace and 100 pounds of cloves for £65, and finds that he has sold 60 more of cloves for £20 than of mace for £10. What was the price of each per pound. *Ans.* 10s. and 5s.

17. The fore-wheel of a carriage makes 6 revolutions more than the hind wheel in going 120 yards; but if the circumference of each be increased 1 yard, it will make only four revolutions more in going the same distance. What is the circumference of each wheel? *Ans.* 4, and 5 yards.

18. Find four numbers in arithmetical progression, such, that the product of the two *extremes* shall be 45, and the product of the two *means* 77.

Let x be the first term, and y the common difference of the terms: then the numbers will be

$$x, x+y, x+2y, x+3y; \quad (175).$$

and by the conditions of the problem we shall have

$$x^2+3xy=45; \text{ and } x^2+3xy+2y^2=77.$$

Ans. 3, 7, 11, and 15.

19. A farmer has a field 16 rods long and 12 rods wide, which he wishes to enlarge so that it may contain just twice as much area, without altering the proportion of the sides. What will be the dimensions of the field when thus enlarged? *Ans.* $16\sqrt{2}$; and $12\sqrt{2}$.

20. Find three numbers, such, that the difference of the first and second shall be two less than the difference of the second and third, their sum 17, and the sum of their squares 115.

The solution will be facilitated by assuming x to represent the second number, and y the difference of the first and second.

Ans. 3, 5, and 9.

21. There are two square gardens which together contain 1025 square rods, and a side of the one exceeds a side of the other by 5 rods. What are the sides of the two gardens? *Ans.* 20, and 25 rods.

22. Find two numbers, such, that their sum, their product, and the difference of their squares shall all be equal to one another.

Take $x+y$ to represent the greater, and $x-y$ the less number.

Ans. $\frac{3}{2} \pm \sqrt{\frac{5}{4}}$, and $\frac{1}{2} \pm \sqrt{\frac{5}{4}}$.

23. A merchant received \$12 for a quantity of linen, and an equal sum, at 50 cents less per yard, for a quantity of calico, which exceeded the quantity of linen by 32 yards. What was the quantity of each? *Ans.* 16, and 48 yards.

24. Find two numbers whose sum multiplied by the greater shall be equal to 192, and whose difference multiplied by the less shall be equal to 32.

The solution will be facilitated by taking x to represent one of the required numbers, and xy the other.

Ans. 12, and 4.

25. Three merchants gained \$1444; of which their respective shares were such that B's, added to the square root of A's, made \$920; but if added to the square root of C's it made \$912. What was the share of each? *Ans.* \$400, \$900, and \$144.

26. The sum of three numbers in harmonical progression is 13, and the product of the two extremes is 18. What are the numbers?

If x and y represent the two extremes, the mean term will be

$$\frac{2xy}{x+y} \quad (184).$$

Ans. 6, 4 and 3.

27. There is a rectangular field whose length is to its breadth as 4 to 3. A part of this field, which is equal to $\frac{1}{4}$ of the whole, being in meadow, there remain for ploughing 1296 square rods. What are the dimensions of the field? *Ans.* 48, and 36 rods.

28. The sum of three numbers in geometrical progression is 21, and the sum of their squares is 189. What are the numbers?

If x and y represent the two extremes, the mean term will be

$$\sqrt{xy}, \quad (189). \quad \text{Ans. } 3, 6, \text{ and } 12.$$

29. A and B set out from two places which are distant 110 miles, and traveled towards each other. A went five miles an hour; and the number of hours in which they met was greater, by four, than the number of miles B went per hour. What was B's rate of traveling?

Ans. 6 miles per hour.

30. The arithmetical mean between two numbers exceeds the geometrical mean by 13, and the geometrical mean exceeds the harmonical mean by 12. What are the numbers? *Ans.* 234 and 104.

31. Three merchants made a joint stock, by which they gained a sum less than that stock by \$80. A's share of the gain was \$60, and his contribution to the stock was \$17 more than B's; also B and C together contributed \$325. How much did each contribute?

Ans. \$75, \$58, and \$267

32. Of three numbers in geometrical progression the greatest exceeds the least by 15, and the difference of the squares of the greatest and the least is to the sum of the squares of the three numbers as 5 to 7. What are the numbers?

Assume x to represent the first term, and y the *ratio* of the progression. *Ans.* 5, 10, and 20.

33. Two persons set out from different places, and traveled towards each other. On meeting, it appeared that A had traveled 24 miles more than B, and that A could have gone B's journey in 8 days, while B would have been 18 days in performing A's journey. What distance was traveled by each? *Ans.* 72, and 48 miles.

34. The joint stock of two partners was \$416. A's money was in the business 9 months, and B's 6 months. When they shared stock and gain, the first received \$228, and the second \$252; what was each man's amount of stock? *Ans.* A's \$192, B's \$224.

35. The sum of \$700 was divided among four persons, A, B, C, and D, whose shares were in geometrical progression; and the difference between the greatest and the least was to the difference between the two means as 37 to 12. What were the several shares?

Ans. \$108, \$144, \$192, and \$256.

SOLUTION OF AFFECTED CUBIC AND HIGHER EQUATIONS.

Various methods have been devised for the solution of Affected Equations of the third and higher degrees. Some of these methods are very prolix,—while others are of limited application; we shall explain those which are the most useful in a practical point of view, without attempting a full exposition of this subject.

Under the head of General Properties of Equations, we have already noticed the *divisors*, (253), and the number of *roots*, (255) of equations; we here present the

General Law of the Coefficients of Equations.

(267) When the terms of any *complete* Equation containing but one unknown quantity x , are all arranged, according to the descending powers of x , in the first member—with the known or absolute term for the *last term*—and the coefficient of the first term is *unity*; then

1. The coefficient of the second term is equal to the sum of all the *roots* of the equation, with their *signs* changed.

2. The coefficient of the third term is equal to the sum of the *products* of all the roots *combined two and two*, with their signs changed, &c.

3. The known or absolute term is equal to the product of all the roots, with their signs changed.

To demonstrate these principles with reference to a Cubic Equation, let the three roots be denoted by a, b , and $-c$; then $x-a, x-b$, and $x+c$ are the *divisors* of the equation, and the equation may be accordingly resolved into

$$(x-a)(x-b)(x+c)=0, \quad (253).$$

By performing the multiplication which is here indicated, and decomposing the terms containing the like powers of x in the product, we find

$$x^3+(c-a-b)x^2+(ab-ac-bc)x+abc=0.$$

In this equation the coefficient of x^2 is the sum of the roots a, b , and $-c$, with their signs changed; the coefficient of x is the sum of the products of the roots *combined two and two*, with their signs changed; and the known or absolute term abc is the product of all the roots, with their signs changed.

The same principles may be demonstrated, in like manner, in reference to an Equation of the second, or of any of the higher degrees.

An application of these principles may be made to the equation $x^2 + 3x - 10 = (x-2)(x+5) = 0$, whose roots are 2, and -5 ; or to $x^3 - 19x + 30 = (x-2)(x+5)(x-3) = 0$, whose roots are 2, -5 , & 3

In the second equation it will be observed that the second term containing x^2 , is wanting, since its coefficient, that is, the sum of the roots 2, -5 , and 3, is 0; and that $19x$ therefore corresponds to the third term, (267...2).

Determination of the Integral Roots of Equations.

(268.) If an equation containing but one unknown quantity x , with all its terms transposed to one side, be divisible by $x +$ or $-$ any number, that number, with a contrary sign, will be a root of the equation (254).

The trial numbers to be used in this division, are the factors or divisors of the known term of the equation, since that term is equal to the product of all the roots of the equation, (267...3).

When an equation has any integral roots, such roots may be readily determined by an application of these principles.

EXAMPLE.

To find the values of x in the equation

$$x^3 + 3x^2 - 4x = 12.$$

The divisors of the known term 12 are 1, 2, 3, 4, 6, and 12; and it will be found, on trial, that the equation

$$x^3 + 3x^2 - 4x - 12 = 0,$$

is divisible by $x-2$, $x+2$, and $x+3$; hence the values of x , or roots of the equation, are 2, -2 , and -3 .

After any one of the three roots has been determined, the two remaining ones may be obtained directly from the quadratic equation which results from dividing the given equation by $x +$ or $-$ the root already found.

$$(x-2)x^3 + 3x^2 - 4x - 12 = (x^2 + 5x + 6).$$

By dividing the given equation by $x-2$, we thus find

$$x^2 + 5x + 6 = 0,$$

or $x^2 + 5x = -6$, which gives $x = -2$ or -3

By this method the last two roots are found the same as before.

SOLUTION OF EQUATIONS BY APPROXIMATION.

(269.) The following method of solution may be applied to an Equation of any degree—even to one in which the unknown quantity is left, without rationalization, in a *surd* expression.

1. By trial find two numbers—differing by a unit or less—which being substituted for x in the given Equation, will produce results, the one less and the other greater than the *known term* of the equation; then,

The difference between the *two results*,
Is to the difference between the *two assumed numbers*,
As the difference between *either result* and the *known term*,
Is to the *correction*, nearly, required in the corresponding assumed number.

2. Take the corrected root thus obtained for one of two numbers to be substituted for x , and find, and apply, a *correction* as before.

We shall thus obtain a *nearer* value of the unknown quantity; and the approximation may be carried, in like manner, to any required exactness.

EXAMPLE.

To find an approximate value of x in the equation.

$$x^3 + x^2 + x = 100.$$

First, It will be found that x is more than 4, and less than 5. Substituting these numbers for x , we have

$$\begin{array}{r} 64 \dots\dots x^3 \dots\dots 125 \\ 16 \dots\dots x^2 \dots\dots 25 \\ 4 \dots\dots x \dots\dots 5 \\ \hline 84 \qquad \qquad \qquad 155 \end{array}$$

The difference between the two results is $155 - 84 = 71$; and the difference between the *less result* and the known term 100 is 16.

Then $71 : 1 :: 16 : \text{the correction} . 225$.

This correction, added to the *less* assumed number, gives 4.225 for an *approximate* value of x .

Secondly, By substituting 4.2 and 4.3 for x , we have

$$\begin{array}{r} 74.088 \dots\dots x^3 \dots\dots 79.507 \\ 17.64 \dots\dots x^2 \dots\dots 18.49 \\ 4.2 \dots\dots x \dots\dots 4.3 \\ \hline 95.928 \qquad \qquad \qquad 102.297 \end{array}$$

Forming a proportion between the difference of these two results, and the difference between the *greater result* and the known term 100,
 $6.369 : .1 :: 2.297 : \text{the correction } .036.$

By subtracting this correction from the *greater* assumed number 4 3, we have 4.264 for a *nearer* value of x .

For the next approximation we should take 4.264 and 4,265 to be substituted for x in the given Equation. The value of x would then be found to be 4.2644299 *very nearly*.

In the Proportion for finding the *correction*, it is best to employ the *less error* in the results of the substitution.

Thus in the first substitution, in this Example, the error in the less result 84 is $(100 - 84) = 16$, and this being less than the error in the 155, we employ 16 in the first proportion.

But in the second substitution, the error in the greater result 102.297 is less than the error in the 95.928, and we accordingly use $(102.297 - 100) = 2.297$ in the second proportion.

Each approximative solution will generally *double the number of true figures in the root*. Thus in the preceding Example we found by trial that 4 is the first figure in the root, and the first solution gives 4.2 for the first two correct figures; the next solution gives 4.264; and the number of figures will again be doubled by a third solution. This property determines the number of figures which need be found in the successive *corrections* of the assumed numbers.

To find the other Roots of the given Equation, we would divide

$$x^3 + x^2 + x - 100 = 0 \text{ by } x - 4,2644, \text{ \&c.} = 0, (253).$$

We should thus obtain a *quadratic* equation, from which the other two values of x might be determined, according to the usual method.

☞ When all the Roots of an Equation have been found, we may *verify* them by the property that, with their signs changed, their sum must be equal to the *coefficient of the second term of the equation*, (267...1).

Thus the sum of the *three roots* of the equation in the preceding Example, with their signs changed, would be *unity*, which is the coefficient of the second term x^2 .

EXERCISES

On Affected Cubic and Biquadratic Equations.

1. Find the values of x in the equation

$$x^3 - 6x^2 + 11x = 6, \quad (268).$$

Ans. $x = 1, 2, \text{ or } 3.$
2. Find the values of x in the equation

$$x^3 - 9x^2 + 26x = 24.$$

Ans. $x = 2, 3, \text{ or } 4.$
3. Find the values of x in the equation

$$x^3 - 3x^2 - 6x = -8.$$

Ans. $x = 1, 4, \text{ or } -2.$
4. Find the values of x in the equation

$$2x^3 - 6x^2 - 8x = -24.$$

Ans. $x = 2, 3, \text{ or } -2.$
5. Find the values of x in the equation

$$x^4 + 2x^3 - 13x^2 - 14x + 24 = 0.$$

Ans. $1, -2, 3, \text{ or } -4$

6. Find an approximate value of x in the equation

$$x^3 + 10x^2 + 5x = 260, \quad (269).$$

Ans. $x = 4.117'.$
7. Find an approximate value of x in the equation

$$x^3 - 15x^2 + 63x = 50.$$

Ans. $x = 1.028'.$
8. Find an approximate value of x in the equation

$$x^3 - 17x^2 + 54x = 350.$$

Ans. $x = 14.95'.$
9. Find an approximate value of x in the equation

$$x^4 - 3x^2 - 75x = 10000.$$

Ans. $x = 10.23'.$
0. Find an approximate value of x in the equation

$$2x^4 - 16x^3 + 40x^2 - 30x + 1 = 0.$$

Ans. $x = 1.284'.$
1. Find an approximate value of x in the equation

$$\left(\frac{1}{5}x^2 - 15\right)^2 + x\sqrt{x} = 90.$$

Ans. $x = 1^{\circ} 5'.$

ELIMINATION BY THE METHOD OF COMMON DIVISOR.

(269.) The following is a general method of Elimination, and, for Equations of the *higher degrees*, it will sometimes be found preferable to any other.

Transpose all the terms of the two Equations to one side; then divide one into the other, and the *remainder* into the *divisor*, and so on, as in finding the Greatest Common Measure, (66,) until one of the two unknown quantities is *eliminated from the remainder*; and put this remainder = 0.

To eliminate x from the equations

$$x^2 + xy = 10, \text{ and } xy + 2y^2 = 24.$$

$$\begin{array}{r} xy + 2y^2 - 24 \\ x \\ \hline x^2 + xy - 10 \end{array} \left) \begin{array}{l} x^2y + 2xy^2 - 24x \\ \underline{x^2y + xy^2 - 10y} \\ xy^2 - 24x + 10y, \text{ or } x(y^2 - 24) + 10y. \end{array}$$

$$\begin{array}{r} x^2 + xy - 10 \\ y^2 - 24 \\ \hline x(y^2 - 24) + 10y \end{array} \left) \begin{array}{l} x^2(y^2 - 24) + x(y^3 - 24y) - 10y^2 + 240 \\ \underline{x^2(y^2 - 24) + 10xy} \\ x(y^3 - 24y) - 10xy - 10y^2 + 240 \\ \underline{x(y^3 - 24y) + 10y^2} \\ -10xy - 20y^2 + 240 \end{array}$$

$$\begin{array}{r} x(y^2 - 24) + 10y \\ y \\ \hline xy + 2y^2 - 24 \end{array} \left) \begin{array}{l} x(y^3 - 24y) + 10y^2(y^2 - 24) \\ \underline{x(y^3 - 24y) + 2y^4 - 48y^2 - 24y^2 + 576} \\ -2y^4 + 82y^2 - 576 = 0. \end{array}$$

In the remainder $-10xy - 20y^2 + 240$, we cancel the factor 10 and change the *signs*, for the next divisor. In dividing into this divisor, we take the binomial $y^2 - 24$ for the quotient, and multiply the divisor by this binomial.

The first remainder is equal to 0, because the divisor and dividend are each equal to 0; and it follows hence that *each subsequent remainder is equal to 0*.

The operation will be much more simple if we divide the first equation by the second: the result will be the same.

CHAPTER XI.

GENERAL DESCRIPTION OF PROBLEMS.—INEQUATIONS.

MISCELLANEOUS PROBLEMS.

1. *Determinate Problems.*

(270.) A *Determinate Problem* is one in which the given conditions determine the values of the unknown or required quantities.

A *Determinate Problem* is represented by as many *independent* equations as there are different conditions to be expressed, or unknown quantities to be determined, (120.)

All the Problems which have hitherto been proposed in this work, are *determinate*; and no example of this kind need be here given.

2. *Indeterminate Problems.*

(271.) An *Indeterminate Problem* is one in which the given conditions do not determine the values of the required quantities,—admitting either of an *unlimited* number of values to those quantities, or else of a variety of values, within certain limits.

An *Indeterminate Problem* is represented either by a less number of independent Equations than there are unknown quantities to be determined, or by an *identical* equation.

We give an example of each of these forms of indeterminateness.

EXAMPLE I.

To find three numbers such that the first shall be 5 less than the second, and the sum of the second and third shall be 12.

This Problem contains but two conditions; and if we represent the three required numbers by x , y , and z , we shall have only the two Equations

$$\begin{aligned}y - x &= 5; \\ y + z &= 12.\end{aligned}$$

By subtracting the first equation from the second, we have

$$x+z=7.$$

This equation will admit of an *unlimited* number of values of x and z ; for we may assume any value whatever for one of the letters, as x , and determine thence the corresponding value of z .

Thus if $x=\frac{1}{4}$, $z=6\frac{3}{4}$; if $x=\frac{1}{2}$, $z=6\frac{1}{2}$; if $x=\frac{3}{4}$, $z=6\frac{1}{4}$, &c.; and from the values of x or z , we might obtain the corresponding values of y from one of the given equations.

If, however, the required numbers were limited to *integral* values. the third equation would be satisfied only by

$$x=1, 2, 3, 4, 5, \text{ or } 6, \text{ and } z=6, 5, 4, 3, 2, \text{ or } 1.$$

In the first equation $y=5+x$, which would give

$$y=6, 7, 8, 9, 10, \text{ or } 11.$$

EXAMPLE II.

To find a number such, that $\frac{3}{4}$ of it, diminished by $\frac{1}{3}$ of it, and by 5, shall be equal to $\frac{1}{12}$ of the excess of 5 times the number above 60

The equation of this problem will be

$$\frac{3x}{4} - \frac{x}{3} - 5 = \frac{5x-60}{12}.$$

Clearing the equation of its fractions,

$$9x-4x-60=5x-60;$$

$$\text{or } 5x-60=5x-60.$$

This last is an *identical* Equation, which will be satisfied by attributing to x any numerical value whatever. The problem is therefore entirely *indeterminate*.

We may obtain an expression for the value of x from the last equation. Thus, by transposition,

$$5x-5x=60-60.$$

By adding similar terms, and retaining x as a symbol in the first member, we have

$$0x=0;$$

$$\text{which gives } x=\frac{0}{0}.$$

Hence $\frac{0}{0}$ is a symbol of an *indeterminate* quantity.

The same thing will appear from considering that the *quotient* of $0 \div 0$ is *any quantity whatever*; inasmuch as the *divisor* $0 \times$ *any quantity* will produce the *dividend* 0, (43).

3. Impossible Problems.

(272.) An *Impossible Problem* is one in which there is some condition, expressed or implied, which *cannot be fulfilled*.

An Impossible Problem is represented by a greater number of *independent* equations than there are unknown quantities to be determined; or by an equation in which the value of the unknown quantity is *negative—zero—infinite—or imaginary*.

We subjoin an example of each of these forms of impossibility

EXAMPLE I.

To find two numbers whose sum shall be 10, difference 2, and product 20.

Representing the two numbers by x and y , we shall have

$$x+y=10; \quad x-y=2; \quad xy=20.$$

From the first and second equations the values of x and y will be found to be $x=6$, and $y=4$. The third equation cannot, therefore, be fulfilled; that is, the problem is *impossible*.

If the third equation were $xy=24$, the problem would be possible, but this would not be an *independent* equation, since it may be derived from the other two.

Thus, squaring the first and second equations, and subtracting, we find

$$4xy=96, \text{ or } xy=24.$$

EXAMPLE II.

To find a number which, added to 17 and to 53, will make the first sum equal to $\frac{1}{4}$ of the second.

If x represent the number, the equation will be

$$17+x=\frac{53+x}{4}.$$

From this equation we shall find $x=-5$. This number, added to 17 and 53, gives 12 and 48, and $12=\frac{1}{4}$ of 48.

The problem is *impossible* in an arithmetical sense, according to which addition always implies *augmentation*; and it is in this sense only that the problem would be considered.

To make it arithmetically consistent, it should be stated thus:

To find a number which, *subtracted* from 17 and from 53, will make the first *remainder* equal to $\frac{1}{4}$ of the second.

EXAMPLE III. •

To find a number such, that if 82 be increased by 3 times that number, $\frac{1}{6}$ of the sum will be equal to $13\frac{2}{3}$.

If x represent the number, the equation will be

$$\frac{82+3x}{6}=13\frac{2}{3}.$$

Clearing the equation of its fractions, we find

$$82+3x=82;$$

$$\text{which gives } 3x=82-82=0;$$

$$\text{and } x=\frac{0}{3}=0, (50).$$

Hence no number can be found that will fulfil the conditions of the problem; that is, the problem is *impossible*. The result shows that $\frac{1}{6}$ of 82 itself is equal to $13\frac{2}{3}$.

EXAMPLE IV.

To find a number such, that the sum of $\frac{1}{4}$ of it and $\frac{2}{3}$ of it, diminished by 2, shall be equal to $\frac{1}{12}$ of it increased by 3.

If x represent the number, the equation will be

$$\frac{x}{4}+\frac{2x}{3}-2=\frac{11x}{12}+3.$$

From this equation we shall find

$$3x+8x-24=11x+36; \text{ or } 11x-11x=0x=60;$$

$$\text{and } x=\frac{60}{0}=\infty, \text{ infinity; } (50).$$

The result shows that it would require a number *infinitely great*, to fulfil the conditions of the problem. The problem is therefore *impossible*.

EXAMPLE V.

To divide the number 24 into two such parts, that their product shall be 150.

If x represent one of the two parts, $24-x$ will represent the other and the equation will be

$$24x-x^2=150$$

$$\text{or } x^2-24x=-150, (117);$$

$$\text{which gives } x=12\pm\sqrt{144-150},$$

$$=12\pm\sqrt{-6}.$$

In this value of x , the part $\sqrt{-6}$ is *imaginary*, that is, it is an *impossible* quantity, (246); hence the problem is impossible.

That the preceding problem is impossible, will also appear from the following general proposition; viz:

(273.) The *square of one half* of any quantity is greater than the product of any *two unequal parts* of the quantity.

Let s represent any number, and d the difference between any two parts of the number; then the respective parts are

$$\frac{s+d}{2} \text{ and } \frac{s-d}{2} \text{ (168), and their product is } \frac{s^2-d^2}{4}.$$

This product will vary directly as its numerator s^2-d^2 (147); and will therefore have its *greatest value* when $d=0$. In that case the product becomes $\frac{1}{4}s^2$, which is the square of $\frac{1}{2}s$, half the given number.

It may also be remarked here, that *the sum of the two equal factors* of any quantity, *is less than the sum of any two unequal factors* into which the quantity can be resolved.

For, s representing the *sum*, and d the difference, of the two factors of a quantity, those factors are

$$\frac{s+d}{2} \text{ and } \frac{s-d}{2} \text{ (168), and their product is } \frac{s^2-d^2}{4}.$$

This product will retain a *constant value*, if s^2 and d^2 be *equally diminished*, for then the value of s^2-d^2 will remain *constant*. In this diminution, s^2 will have its *least value* when d^2 is made 0; so that s , the sum of the two factors, will have its least value when $a=0$, and the two factors will then be *equal to each other*.

Signification of the Different Forms under which the Value of the Unknown Quantity may be found in an Equation.

(274.) 1. *Positive* values of the unknown or required quantities, fulfil the conditions of problems in the sense in which they are proposed.

2. A value of the unknown quantity of the form $\frac{0}{0}$, shows that the problem from which the equation was derived is *indeterminate*.

3. A *negative* value of the unknown quantity, in an equation of the *first degree*, indicates an *impossibility* in the problem, produced by taking this quantity *additively*, instead of *subtractively*, or *vice versa*.

4. When the value of the unknown quantity in an equation is *zero*, *infinite*, or *imaginary*, the problem from which the equation was derived is *impossible*.

MISCELLANEOUS PROBLEMS.

1. A, B, and C together have \$2000. B has \$100 less than twice as much as A, and C \$400 less than twice as much as the other two together. What sum has each?

Ans. A, \$300 ; B, \$500 ; C, \$1200.

2. A gentleman has three plantations. The first contains 250 acres, the second as much as the first and $\frac{1}{6}$ of the third, and the third as much as the first and second. What is the whole number of acres?

Ans. 1200 acres.

3. A company of workmen had been employed on a piece of work for 24 days, and had half finished it, when, by calling in the assistance of 16 more men, the remaining half was completed in 16 days. What was the original number of men?

Ans. 32 men

4. A's money was equal to $\frac{3}{4}$ of B's. A paid away \$50 less than $\frac{2}{3}$ of his, and B \$50 more than $\frac{3}{4}$ of his, when it was found that the latter had remaining only $\frac{1}{3}$ as much as the former. What sum had each at first?

Ans. A, \$300 ; B, \$400.

5. A person wishing to enclose a piece of ground with palisades, found, that if he set them one foot asunder, he would not have enough by 150, but if he set them one yard asunder, he would have too many by 70. What was the number of his palisades?

Ans. 180 palisades.

6. From two tracts of land of equal size, were sold quantities in the proportion of 3 to 5. If 150 acres less had been sold from the one which is now the smaller of the two, only $\frac{2}{3}$ as much would have been taken from it as from the other ; how many acres were sold from each?

Ans. 150, and 250 acres.

7. A and B had adjoining farms, which, in quantity, were in the ratio of 4 to 5. A sold to B 50 acres, and afterwards purchased from B one-third of his entire tract, when it was found that the original ratio of their quantities of land had been reversed. How many acres had each at first?

Ans. A, 200 ; B, 250 acres.

8. A waterman can row down the middle of the stream, on a certain river, 5 miles in $\frac{3}{4}$ of an hour ; but it takes him $1\frac{1}{2}$ hours to return, though he keeps along shore, where the current is but half as strong as in the middle. What is the velocity of the middle of the stream?

Ans. $2\frac{2}{3}$ miles per hour.

9. A farmer has three flocks of sheep, whose numbers are in the proportion of 2, 3, and 5. If he sell 20 from each flock, the whole number will be diminished in the proportion of 4 to 3 ; how many has he in each flock?

Ans. 48, 72, and 120 sheep

10. The sum of \$1170 is to be divided between three persons, A, B, and C, in proportion to their ages. Now A's age is to B's as 1 to $1\frac{1}{3}$, and to C's as 1 to 2; what are the respective shares?

Ans. \$270; \$360; \$540.

11. A hare is 50 leaps before a greyhound, and takes 4 leaps to the greyhound's 3; but 2 of the greyhound's leaps are as much as 3 of the hare's. How many leaps must the greyhound take to catch the hare?

Ans. 300.

12. A vintner has two casks of wine, the contents of which are in the proportion of 5 to 6, and if $\frac{1}{6}$ of the quantity in the second were to be drawn off, the contents of the two casks would be equal. How many gallons are there in each?

Ans. This problem is *indeterminate*; how is its indeterminateness indicated?

13. A person looking at his watch, and being asked what o'clock it was, replied that it was between *eight* and *nine*, and that the hour and minute hands were exactly together. What was the time?

Ans. 43m. $38\frac{2}{11}$ s. past *eight*.

14. A criminal having escaped from prison, traveled 10 hours before his escape was known. He was then pursued, and gained upon 3 miles an hour. When his pursuers had been 8 hours on the way they met an express going at the same rate as themselves, who had met the criminal 2 hours and 24 minutes before. In what time from the commencement of the pursuit will the criminal be overtaken?

Ans. 20 hours.

15. A regiment of militia containing 875 men is to be raised from three counties, A, B, and C. The quotas of A and B are in the proportion of 2 and 3, and of B and C in the proportion of 4 to 5. What is the number to be raised by each?

Ans. 200, 300, and 375 men.

16. If 19 pounds of gold, in air, weighs 18 pounds in water; 10 pounds of silver, in air, weighs 9 in water; and a mass of 106 pounds, composed of gold and silver, weighs 99 pounds in water; what are the respective quantities of gold and silver in the mass?

Ans. 76, and 30 pounds.

17. A farmer having mixed a certain quantity of corn and oats, found that if he had taken 6 bushels more of each, there would have been 7 bushels of corn to 6 of oats; but if he had taken 6 bushels less of each, there would have been 6 bushels of corn to 5 of oats. How many bushels of each were mixed?

Ans. 78, and 66 bushels.

18. Two persons, A and B, can perform a piece of work in 16 days. They work together for 4 days, when A being called off, B is left to finish it, which he does in 36 days more. In what time could each do it separately?

Ans. 24, and 48 days.

19. A merchant has two casks containing unequal quantities of wine. Wishing to have the same quantity in each, he pours from the first into the second as much as the second contained at first; then he pours from the second into the first as much as was left in the first; and then again from the first into the second as much as was left in the second, when there are found to be 16 gallons in each cask. How many gallons did each cask contain at first? *Ans.* 22, and 10 gallons.

20. A fisherman being asked how many fish he had caught, replied, If 5 be added to one-third of the number that I caught yesterday, it will make half the number I have caught to-day; or if 5 be subtracted from three times this half, it will leave the number I caught yesterday. How many were caught each day?

Ans. This problem is *impossible*; how is its impossibility indicated?

21. A laborer engaged for n days, on condition that he should receive p pence for each day that he worked, and forfeit q pence for each day that he idled. At the end of the time he received s pence; how many days did he work? and how many was he idle?

Ans. Worked $\frac{nq+s}{p+q}$; was idle $\frac{np-s}{p+q}$ days.

22. A, B, and C engage in a joint speculation. A invests \$2000 for 5 months, B \$2400 for 4 months, and C \$1600 for 7 months. The profits amount to \$4620; what is each man's share of profit?

Ans. \$1500, \$1440, \$1680.

23. A farmer wishes to mix rye worth 40 cents a bushel, and oats worth $26\frac{2}{3}$ cents a bushel, in such quantities as to produce 100 bushels which shall be worth 30 cents a bushel. What quantity of each must be taken?

Ans. 25, and 75 bushels.

24. Three persons engage in a joint mercantile adventure, in which the first has the capital a for the time b , the second the capital c for the time d , and the third the capital e for the time f . Their profits amount to s ; what is each partner's share of profit?

Ans. $\frac{abs}{ab+cd+ef}$; $\frac{cds}{ab+cd+ef}$; $\frac{efs}{ab+cd+ef}$.

25. A church which cost \$40,000 is insured, annually, at $1\frac{1}{2}$ per cent., for such an amount, that, in case of its being destroyed by fire, the Insurance Company shall be liable for the cost of the edifice, and the *premium* of insurance. What is the sum insured?

Ans. 40609.13'.

26. Four towns are situated in the order of the first four letters of the alphabet. The distance from A to D is 34 miles; the distance from A to B is to the distance from C to D as 2 to 3; and $\frac{1}{4}$ of the distance from A to B added to half the distance from C to D, is 3 times the distance from B to C. What are the respective distances?

Ans. 12 4, and 18 miles

27. A commission merchant receives the sum of s dollars to invest in merchandise—himself to retain a commission of r per cent. on the amount of the purchase. What is the sum to be invested?

$$\text{Ans. } \frac{100s}{100+r}.$$

28. A sum of money was equally divided among a number of persons, by first giving to A \$100 and $\frac{1}{6}$ of the remainder, then to B \$200 and $\frac{1}{6}$ of the remainder, then to C \$300 and $\frac{1}{6}$ of the remainder; and so on. What was the sum divided? and the number of persons?

$$\text{Ans. } \$2500, \text{ and } 5 \text{ persons.}$$

29. The sum of \$500 is to be applied in part towards the payment of a debt of \$900, now due, and in part to paying the interest, at 7 per cent., in advance, on the remainder of the debt, on which a credit of 12 months is to be allowed. What is the amount of payment that can be made on the debt?

$$\text{Ans. } 469.89'$$

30. A besieged garrison had such a quantity of bread as would, if distributed to each man at 10 ounces a day, last 6 weeks; but having lost 1200 men in a sally, the governor was enabled to increase the allowance to 12 ounces per day. What was the original number of men?

$$\text{Ans. } 7200 \text{ men.}$$

31. A is indebted to B the sum of s dollars, and is able to raise but a dollars. With this latter sum A proposes to pay a part of the debt, and the interest, at r per cent., in advance, on his note at n years, for the remainder. For what sum should the note be drawn

$$\text{Ans. } \frac{100(s-a)}{100-rn}.$$

32. The crew of a ship consisted of her complement of sailors and a number of soldiers. Now there were 22 sailors to every three guns, and 10 over. Also the whole number of men was 5 times the number of soldiers and guns together. But after an engagement, in which the slain were $\frac{1}{4}$ of the survivors, there wanted 5 of being 13 men to every 2 guns. Required the number of guns, soldiers, and sailors.

$$\text{Ans. } 90 \text{ guns, } 55 \text{ soldiers, } 670 \text{ sailors.}$$

33. Two sums of money, amounting together to \$600, were put at interest—the smaller at 2 per cent. more than the other. The interest of the larger sum was afterwards increased, and that of the smaller diminished, 1 per cent. By this the interest of the whole was augmented *one-twentieth*. But if the interest of the greater sum had been so increased, without any diminution of the other, the interest of the whole would have been increased *one-tenth*. What were the two sums? and the two rates of interest?

$$\text{Ans. } \$400 \text{ at } 6 \text{ per cent. ; } \$200 \text{ at } 8 \text{ per cent.}$$

34. Two persons purchase 300 acres of land at \$2 per acre, each one paying \$300; but the first takes the more fertile portion of the

tract, at 25 cents above the mean price per acre, and the second the remainder at 25 cents below the mean price per acre. How many acres has each?

Ans. This problem is *impossible*; how is its impossibility indicated?

35. The *area* of a field is 432 square rods, and the sum of its length and breadth is equal to twice their difference. Required the length and breadth of the field.

Ans. 36, and 12 rods.

36. A and B lay out some money on speculation. A disposes of his interest in the business for £11, and gains as much per cent. as B lays out; B gains £36, and it appears that A gains 4 times as much per cent. as B. What was the capital of each?

Ans. £5, and £120

37. A garden which is 12 rods in length, and 8 rods in breadth, is surrounded by a walk whose area is equal to $\frac{1}{6}$ of the area of the garden itself. Required the breadth of the walk.

Ans. $-5 + \sqrt{29}$.

38. A and B hired a pasture, into which A put 4 horses, and B as many as cost him 18 shillings a week. Afterwards B put in two additional horses, and found that he must pay 20 shillings a week. At what rate was the pasture hired?

Ans. 30s. per week.

39. A gentleman bought a rectangular piece of ground, at \$10 for every rod in its *perimeter*. If the same area had been in the form of a square, and had been purchased in the same way, it would have cost \$20 less; and a square piece of the same perimeter would have contained $12\frac{1}{4}$ square rods more. What were the length and breadth of the lot?

Ans. 16, and 9 rods.

40. A person being asked the ages of himself and his wife, replied, that the product of their ages added to the square of his age, would make 1560. but added to the square of hers would make 1144. What were their ages?

Ans. 30, and 22.

41. A and B purchased a farm containing 900 acres for which they paid \$900 each. On dividing the land, it was agreed that A should have his choice of situation, and pay 45 cents per acre more than B. How many acres should each have taken? and at what price per acre?

Ans. A 400 acres at \$2.25; B 500 acres at \$1.80.

42. A capital of \$13,000 was divided into two parts, which were put at interest in such a manner that the income was the same from each. If the first part had been at the same rate of interest as the second, it would have produced an income of \$360; and if the second part had been at the same rate as the first, it would have produced an income of \$490. What were the two rates of interest?

Ans. 7 and 6 per cent.

43. A departs from London towards Lincoln at the same time at which B leaves Lincoln for London. When they met, A had traveled 20 miles more than B, having gone as far in $6\frac{2}{3}$ days as B had in all the time; and it appeared that B would not reach London under 15

days. What is the distance between the two places? and how far had each man traveled?

Ans. Distance, 100 miles; A had gone 60, B 40 miles.

44. The product of the two dimensions of a rectangular piece of land, subtracted from the square of the greater dimension, leaves 300 square rods, and subtracted from the square of the less, leaves 200 square rods. What are the dimensions of the piece?

Ans. This problem is *impossible*; how is its impossibility indicated? and in what does this impossibility consist?

45. A person being asked the ages of his two children, replied, that the difference of their ages was 3 years, and the product multiplied by the sum of their ages was 308. What were their ages?

This problem will result in an Affected Cubic Equation.

Ans. 7 and 4 years.

46. A gentleman who had a square lot of ground, reserved 10 square rods out of it, and sold the remainder for \$432, which was as many dollars per square rod as there were rods in a side of the whole square? What was the length of its sides?

Ans. 8 rods.

47. A and B set out together from the same place, and travel in the same direction. A goes the first day 28 miles, the second 26, and so on, in arithmetical progression; while B goes uniformly 20 miles per day. In how many days will the two be together again?

Ans. 9 days.

48. A farmer wishes to build a crib whose capacity shall be 1620 cubic feet, and whose length, breadth, and height shall be in an arithmetical progression decreasing by the common difference 3. What must be the dimensions of the crib?

It may be well to remind the student here, that *cubic measure*, or measure of capacity, is found by multiplying together *length*, *breadth*, and *height* or depth.

Ans. 15, 12, and 9 feet.

49. One traveler sets out to go from A to B, at the same time at which another sets out from B to A. They both travel uniformly, and at such rates, that the former, 4 hours after their meeting, arrives at B, and the latter at A, in 9 hours after. In how many hours did each one perform the journey?

Ans. 10. and 15 hours.

50. A lady on being asked the ages of her three little boys, answered that they were in harmonical progression; and the sum of their ages was 22 years; and that if the ages of the two elder were each increased by $\frac{1}{3}$ of itself, the three would then be in geometrical progression. What were the respective ages?

Ans. 4, 6, and 12 years.

51. A person wishes to construct two cubical reservoirs which shall differ in their linear dimensions by 4 feet, and which shall together contain 5824 cubic feet. What must be the dimensions of the two reservoirs?

Ans. 12, and 16 feet.

52. Two partners, A and B, divided their gain, which was \$60, when B's share was found to be \$20. A's capital was in trade 4 months; and if the number 50 be divided by A's capital, the quotient will be the number of months that B's capital, which was \$100, continued in trade. What was A's capital? and the time B's was in trade?
Ans. A's Capital \$50; B's 1 month in trade.

53. Let there be a square whose side is 110 inches; it is required to assign the length and breadth of a rectangle whose perimeter shall be greater than that of the square by 4 inches, but whose area shall be less than the area of the square by 4 square inches.

Ans. 126, and 96 inches.

54. There is a number consisting of three digits which increase from left to right by the common difference 2; and the product of the three digits is 105. Required the number.
Ans. 357.

55. A person bought 2 pieces of cloth for \$63. For the first piece he paid as many dollars per yard as there were yards in both pieces, and for the second as many dollars per yard as there were yards in the first more than in the second; also the first piece cost six times as much as the second. What was the number of yards in each piece?

Ans. 6, and 3 yards.

56. There is a number consisting of 4 digits which decrease from left to right by the common difference 2; and the product of the four digits is 945. Required the number.

Ans. 9753.

57. A gentleman purchased two square lots of ground for \$300; each of them cost as many cents per square rod as there were rods in a side of the other, and the greater contained 500 square rods more than the less. What was the cost of each lot?

Ans. \$180, and \$120.

58. A merchant bought a number of bales of cloth. The number of pieces in each bale was 10 more than the number of bales, and the number of yards in each piece was 5 more than the number of pieces in each bale; and the whole quantity was 1500 yards. What was the number of bales?

Ans. 5 bales.

59. A person dies, leaving children, and a fortune of \$46800, which, by his will, is to be divided equally amongst them. Immediately after the death of the father, two of the children also die, in consequence of which each surviving one receives \$1950 more than he was entitled to by the will. How many children did the father leave?

Ans. 8 children.

60. A coach set out from Cambridge for London with 4 more *outside* than *inside* passengers. Seven outside passengers went at 2 shillings less than 4 inside ones, and the fare of the whole amounted to £9. At the end of half the journey, 3 more outside and one more inside passenger were taken up, in consequence of which the fare of

the whole was increased in the proportion of 19 to 15. Required the number of passengers at first, and the fare of each.

Ans. 5 inside, and 9 outside passengers; fares 18 and 10 shillings.

61. There is a fraction which, inverted and increased by $\frac{1}{3}$, will be less than 2; and if its numerator be increased by 2, the value of the fraction will be greater than $\frac{1}{2}$. What is the fraction?

Ans. Between $\frac{3}{5}$ and $\frac{1}{6}$.

62. In a purse which contains 24 coins of silver and copper, each silver coin is worth as many pence as there are copper coins, each copper coin is worth as many pence as there are silver coins, and the whole is worth 18 shillings. How many were there of each kind of coins?

Ans. 6, and 18.

63. The square root of a certain number plus 4, is less than 9; and ten times the square root of the number minus 2, is greater than eight times the square root of the number plus 4. What is the number?

Ans. Between 9 and 25.

64. A and B traveled on the same road, and at the same rate, from Huntingdon to London. At the 50th mile stone from London, A overtook a drove of geese, which were proceeding at the rate of 3 miles in 2 hours; and 2 hours afterwards met a wagon which was moving at the rate of 9 miles in 4 hours. B overtook the same drove of geese at the 45th mile stone, and met the same wagon 40 minutes before he came to the 31st mile stone. Where was B when A reached London?

Ans. 25 miles from London.

TEXT BOOKS

FOR SCHOOLS AND COLLEGES,

INCLUDING THE PRIMARY, ENGLISH, AND CLASSICAL DEPARTMENT, NATURAL
SCIENCE, GEOGRAPHY, MATHEMATICS, BOOK-KEEPING, ETC.

PUBLISHED BY

PRATT, OAKLEY AND COMPANY,

NO. 21 MURRAY STREET, NEW YORK.

* It will be noticed that most of these works were written by Teachers of the
highest eminence.

ELEMENTS OF ASTRONOMY; with explanatory Notes and elegant Illustrations. By John Brocklesby, A. M., Professor in Trinity College
\$1 25.

From the Connecticut Common School Journal.

We take pleasure in calling the attention of teachers and students to this truly excellent book. It is not a milk-and-water compilation, without principles and without demonstration. It contains the elements of the science in their proper integrity and proportions. Its author is a learned man and a practical instructor, as the author of every school-book should be. The style is a model for a text-book, combining in a high degree perspicuity, precision, and vivacity. In a word, it is the very best elementary work on Astronomy with which we are acquainted.

This notice is echoed by a large number of academies, who are promptly introducing the book.

ELEMENTS OF METEOROLOGY; designed for Schools and Academies. By John Brocklesby, A. M., Professor of Mathematics and Natural Philosophy in Trinity College, Hartford 84 cents.

The subject of Meteorology is of the deepest interest to all. Its phenomena everywhere surround us, and ought to be as familiarly known to the scholar as his arithmetic or philosophy. This work treats of Winds in general, Hurricanes, Tornadoes, Water-spouts, Rain, Fogs, Clouds, Dew, Snow, Hail, Thunder-storms, Rainbows, Haloes, Meteorites, Northern Lights, &c.

It has proved highly satisfactory in the school-room, and is now the established text-book in a very large number of our best high schools and academies, where the natural sciences are taught.

It is highly commended by Prof. Olmsted, Prof. Silliman, Dr. J. L. Corstock of Pa., Prof. Love, of Mo., and a host of eminent instructors.

VIEWS OF THE MICROSCOPIC WORLD ; designed for General Reading, and as a Hand-book for Classes in Natural Sciences. By Prof Broekles by \$1 12.

By the aid of a powerful microscope, the author has given us highly instructive accounts of Infusorial Animalcules, Fossil Infusoria, Minute Aquatic Animals, Structure of Wood and Herbs, Crystallization, Parts of Insects, &c. &c.

To those who are necessarily deprived of the aid of a microscope, and even to those who have it, this is a most valuable work. It is clearly and pleasantly written. The sections on the Animalcules, Infusoria, and Crystallization, are very beautifully illustrated with large and expensive plates. The descriptions of the different kinds of these wonderful little animals, many of which multiply by billions in a few hours, are really very instructive. There is no better school library book in the world. It should be read by every man, woman and child.

HUMAN PHYSIOLOGY ; designed for Colleges and the Higher Classes in Schools, and for General Reading. By Worthington Hooker, M. D. Professor of the Theory and Practice of Medicine in Yale College. Illustrated with nearly 200 engravings. \$1 25.

This is an original work, and not a compilation. It presents the subject in a new light, and at the same time embraces all that is valuable for its purpose that could be drawn from the most eminent sources. The highest encomiums are received from all quarters ; a few are subjoined.

From CALEB J. HALLOWELL, *Alexandria High School, Va.*

Hooker's Physiology was duly received. We propose to adopt it as a text book and shall order in the course of a fortnight.

From the Boston Medical and Surgical Journal.

We can truly say that we believe this volume is of great value, and we hope that the rare merits of the diligent author will be both appreciated and patronized

From B. F. TEWKSBURY, *Lenoxville, Pa.*

I am ready to pronounce it unqualifiedly the most admirable book or work on the human system that has fallen under my notice, and they have not been few. If any one desires a complete and thorough elucidation of the great science discussed, they can nowhere be better satisfied than in the perusal of Dr. Hooker's most excellent work.

AN INTRODUCTORY WORK ON HUMAN PHYSIOLOGY, by Prof Hooker, has just been published, designed for all persons commencing the study. Dr. Hooker's works seem to have taken their place decidedly at the head of all treatises on the subject of Physiology. They are rapidly going into seminaries and normal schools in all parts of the country, and the best institutions express their "delight at the result." 60 cents.

A COMPARATIVE ENGLISH-GERMAN GRAMMAR ; based on the affinity of the two languages. By Prof. Elias Peissner, late of the University of Munich, now of Union College, Schenectady. \$1.00

From the New York Churchman.

Of all the German Grammars we have ever examined, this is the most modest and unpretending, and yet it contains a system and a principle which is the life of it, as clear, as practical, as effective for learning grammar as any thing we have ever seen put forth, with so much more pretense of originality and show of philosophy. It will be found, too, we think, that the author has not only presented a new idea of much interest in itself, but has admirably carried it out in the practical lessons and exercises of his work.

From PROF. J. FOSTER, *of Schenectady.*

I have examined Prof. Peissner's German Grammar with some attention, have marked with interest the rapid advancement of students here using it as a text book and have myself carefully tested it in the instruction of a daughter eleven years of age. The result is a conviction that it is most admirably adapted to secure easy, pleasant, and real progress, and that from no other work which has come under my notice can so satisfactory a knowledge of the language be obtained in a given time.

WHITLOCK'S GEOMETRY AND SURVEYING, is a work for advanced students, possessing the highest claims upon the attention of Mathematical Teachers. \$1 50.

In comparison with other works of the kind, it presents the following advantages:

1. A better connected and more progressive method of geometrizing, calculated to enable the student to go alone.
2. A fuller, more varied, and available practice, by the introduction of more than four hundred exercises, arithmetical, demonstrative, and algebraical, so chosen as to be serviceable rather than amusing, and so arranged as greatly to aid in the acquisition of the theory.
3. The bringing together of such a body of geometrical knowledge, theoretical and practical, as every individual on entering into active life demands.
4. A system of surveying which saves two-thirds of the labor required by the ordinary process.

This work is well spoken of universally, and is already in use in some of the best institutions of this country. It is recommended by Prof. Pierce, of Cambridge, Prof. Smith, of Middletown, Prof. Dodd, of Lexington, and many other eminent mathematicians.

From E. M. MOASE, Esq.

I consider that I have obtained more mathematical knowledge from Whitlock's Geometry than from all other text-books combined. Unlike too many treatises of a similar nature, it is eminently calculated to make mathematicians

PROF. J. B. DODD'S MATHEMATICAL SERIES

COMPRISES

AN ELEMENTARY AND PRACTICAL ARITHMETIC.....	\$0 40
HIGH SCHOOL ARITHMETIC.....	0 84
ELEMENTS OF ALGEBRA.....	0 84
HIGHER ALGEBRA.....	1 50
KEY TO ALGEBRA.....	0 84
ELEMENTS OF GEOMETRY.....	1 00

These books are believed to be unrivaled in the following particulars:

1. The philosophical accurateness with which their topics are arranged, so as to show the mutual dependence and relationship of their subjects.
2. The scientific correctness and practical convenience of their greatly improved nomenclature.
3. The clear and concise manner in which principles are stated and explanations are given.
4. Brevity and completeness of rules.
5. The distinctness with which the true connection between Arithmetic and its cognate branches is developed.
6. The excellent and thorough intellectual discipline superinduced.

RECOMMENDATIONS.

From R. T. P. ALLEN, Superintendent of Kentucky Military Institute.

Upon a careful examination of a manuscript Treatise on Arithmetic, by Prof. Dodd, I find it greatly superior to all others which have come under my notice, in system, completeness, and nomenclature. The arrangement is natural, the system complete, and the nomenclature greatly improved. These improvements are not slight; they are fundamental—eminently worthy the attention of the mathematical teacher, and give a character of unity to the work which at once distinguishes it from all others on this subject.

From C. M. WRIGHT, Associate Principal of Mount Palatine Academy.

I have examined Dodd's Arithmetic, and am fully persuaded that it is superior to any other with which I am acquainted. I could speak in detail were it necessary, but all that is required to establish its reputation and introduction, is to have it known by teachers

From M. S. LITTLEFIELD, *Grand Rapids, Mich*

I have Dodd's Higher Arithmetic, and unhesitatingly pronounce it the best work for advanced classes I have ever seen.

From E. HINDS, Esq., *of Newtown Academy.*

I have recently adopted Dodd's High School Arithmetic, and like it much. Having seen that Prof. Dodd is also author of an Algebra, I should like to see that work before forming a new class.

From H. ELIAS, Esq., *Palmyra, Mo.*

I have fairly tested Dodd's Algebra, and am much pleased with it. If I like its geometry as well as the Algebra, I shall forthwith introduce it into my school.

From PROF. W. H. DE PUY.

We have introduced Dodd's Algebra into the Genesee Wesleyan Seminary as a permanent text-book.

From R. H. MOORE, *Ill.*

Dodd's Algebra possesses excellencies pertaining to no other work

From REV J. A. McCANLEY, *Va.*

I am much pleased with Dodd's Algebra, and will introduce it.

From OSCAR HARRIS, *N. J*

I use Professor Dodd's Algebra, and shall continue it as our regular text-book

From PROF. A. L. HAMILTON, *President of Andrew College.*

I have examined with some care Prof. Dodd's Elements of Geometry, and, so far as I am capable of judging, I conceive it to be in many respects decidedly the best work of the kind extant. For simplicity, exactness, and completeness, it can have no superior. Like his Arithmetic and Algebra, in many important particulars, his Geometry stands pre-eminent and alone.

A NEW COMMON-SCHOOL ARITHMETIC, by Prof. Dodd, is in press.

The Department of Public Instruction in Canada has repeatedly ordered Prof Dodd's books, as well as many of F. B. & Co.'s other publications, for use in schools

SCHELL'S INTRODUCTORY LESSONS IN ARITHMETIC; designed as an Introduction to the study of any Mental or Written Arithmetic. It contains a large amount of mental questions together with a large number of questions to be performed on the slate, thus combining mental and written exercises for young beginners. This is a very attractive little book, superior to any of its class. It leads the pupil on by the easiest steps possible, and yet insures constant progress. 20 cents.

From GEO. PAYNE QUACKENBOS, *Rector of Henry street Grammar School, N. Y*

It is unnecessary to do more than to ask the attention of teachers to this work, they cannot examine it impartially without being convinced of its superior merits. It will, no doubt, become one of the most popular of school-books.

From J. MARKHAM, *Ohio.*

I wish to introduce Schell's little Arithmetic. It is just the thing for beginners. Send six dozen

From G. C. MERRIFIELD, *Ind.*

I am highly pleased with Schell's little book, and shall use it.

From D. F. DEWOLF *Ohio.*

Schell's little book for children is a *beau-ideal* of my own, and of course it suits

From D. G. HEFFRON, *Sup't. Schools, Utica.*

The School Committee have adopted Schell's Arithmetic for our public schools. Send us three hundred.

AN INTELLECTUAL AND PRACTICAL ARITHMETIC; or, First Lessons in Arithmetical Analysis. By T. L. ENOS, Graduate of the New York State Normal Schools. 25 cents.

The same *clearness* and *conciseness* characterize this admirable book that belong to the works of Prof. Dodd. The natural arrangements of the text, and the logical mode of solving the questions, is a peculiar and important feature belonging to this book alone.

From PROF. C. M. WRIGHT.

I have examined with care and interest Enos' Mental Arithmetic, and shall introduce it at once into the Academy.

From PROFS. D. I. PINCKNEY, S. M. FELLOWS, S. SEARLE, *Rock River Seminary*

We have examined an intellectual Arithmetic, by J. L. Enos, and like it much. We shall immediately use it in our school.

PROF. PALMER'S BOOK-KEEPING; Key and Blanks. 67 cents

This excellent book is superior to the books generally used, because:

1. It contains a large number of business blanks to be filled by the learner, such as deeds, mortgages, agreements, assignments, &c., &c.

2. Explanations from page to page, from article to article, and to settle principles of law in relation to deeds, mortgages, &c., &c.

3. The exercises are to be written out, *after being calculated*. In other works, the pupil is expected to copy, merely.

Palmer's Book-Keeping is used in the New York Public Schools, and extensively in Academies. It is recommended by Horace Webster, LL. D., G. B. Docharty, LL. D., and a large number of *accountants* and teachers.

REV. P. BULLIONS' ENGLISH AND CLASSICAL SERIES,

COMPRISING

PRACTICAL LESSONS IN ENGLISH GRAMMAR AND COMPOSITION.....	\$0 25
PRINCIPLES OF ENGLISH GRAMMAR.....	0 50
PROGRESSIVE EXERCISES IN ANALYSIS AND PARSING.....	0 15
INTRODUCTION TO ANALYTICAL GRAMMAR.....	0 30
NEW, OR ANALYTICAL AND PRACTICAL ENGLISH GRAMMAR.....	0 63
LATIN LESSONS, WITH EXERCISES IN PARSING. By Geo. Spencer, A. M. Half cloth, enlarged.....	0 63
BULLIONS' PRINCIPLES OF LATIN GRAMMAR.....	1 00
BULLIONS' LATIN READER. With an Introduction on the Idioms of the Latin Language. An improved Vocabulary.....	1 00
BULLIONS' CÆSAR'S COMMENTARIES.....	1 00
BULLIONS' CICERO'S ORATIONS. With reference both to Bullions', and Andrew's, and Stoddard's Latin Grammar.....	1 15
BULLIONS' SALLUST.....	1 00
BULLIONS' GREEK LESSONS FOR BEGINNERS.....	0 75
BULLIONS' PRINCIPLES OF GREEK GRAMMAR.....	1 13
BULLIONS' GREEK READER With Introduction on the Idioms of the Greek Language, and Improved Lexicon.....	1 75
BULLIONS' LATIN EXERCISES.....	1 25
COOPER'S VIRGIL.....	2 00

In this series of books, the three Grammars, English, Latin, and Greek, are all on the same plan. The general arrangement, definitions, rules, &c., are the same, and expressed in the same language, as nearly as the nature of the case would admit. To those who study Latin and Greek, much time and labor, it is believed, will be saved by this method, both to teacher and pupil. The analogy and peculiarities of the different languages being kept in view, will show what is common to all. or pecu

war to each; the confusion and difficulty unnecessarily occasioned by the use of elementary works differing widely from each other in language and structure, will be avoided, and the progress of the student rendered much more rapid, easy, and satisfactory.

No series of Grammars, having this object in view, has heretofore been prepared, and the advantages which they offer cannot be obtained in an equal degree by the study of any other Grammars now in use. They form a complete course of elementary books, in which the substance of the *latest* and *best* Grammars in each language has been compressed into a volume of convenient size, beautifully printed on superior paper, neatly and strongly bound, and are put at the lowest prices at which they can be afforded.

The elementary works intended to follow the Grammars—namely, the Latin Reader and the Greek Reader—are also on the SAME PLAN; are prepared with special references to these works, and contain a course of elementary instruction so unique and simple as to furnish great facilities to the student in these languages.

NOTICES.

From PROF. C. S. PENNEL, Antioch College, Ohio.

Bullions' books, by their superior arrangement and accuracy, their completeness as a series, and the references from one to the other, supply a want more perfectly than any other books have done. They bear the marks of the *instructor* as well as the scholar. It requires more than learning to make a good school-book.

From J. B. THOMPSON, A. M., late Rector of the Somerville Classical Institute, N. J.

I use Bullions' works—all of them—and consider them the best of the kind that have been issued in this or any other language. If they were *universally* used we would not have so many superficial scholars, and the study of the classics would be more likely to serve the end for which it was designed—the strengthening and adorning of the mind.

From A. C. RICHARDS, Esq., Clay Co., Ga.

We think Bullions' Latin Grammar, in the arrangement of its syntax and the correctness of its rules, the manner of treating prosody, and the conjugations of the verbs, superior to any other. If his Greek Reader is as good as the Latin Reader, we shall introduce it.

It is almost superfluous to publish notices of books so extensively used.

Within the last few months Dr. Bullions' English Grammar has been introduced into the Public, and many of the Private Schools, the Latin School, the English High School, the City Normal School, of the city of Boston; Normal Schools of Bridgewater and Westfield, Marlborough Academy; cities Salem, Newburyport, &c., Mass.; Portsmouth, Concord, and several academies in New Hampshire; and re-adopted in Albany and Troy, New York. They are used in over seventy academies in New York, and in many of the most flourishing institutions in every State of the Union. Also, in the Public Schools of Washington, D. C., and of Canada, in Oregon and Australia. The classical Series has been introduced into several colleges, and it is not too much to say that Bullions' Grammars bid fair to become the Standard Grammars of the country.

THE STUDENTS' SERIES

BY J. S. DENMAN, A. M.

	Cents.
THE STUDENTS' PRIMER.....	7
“ “ SPELLING-BOOK.....	13
“ “ FIRST READER.....	13
“ “ SECOND “.....	23
“ “ THIRD “.....	44
“ “ FOURTH “.....	73
“ “ FIFTH “.....	94
“ “ SPEAKER.....	9

The Publishers feel justified in claiming that the Students' Series is decidedly *the best* for teaching reading, and spelling that has yet appeared. The plan of teaching includes, in the first steps, an ingenious and original mode of repetition which is very pleasing and encouraging to the pupil. The first books of the series are very instructive, and the later portions consist of fine selections, which are not hackneyed. Prof. Page, late Principal of the New York State Normal School, said of this system: "*It is the best I ever saw for teaching the first principles of Reading.*" Such testimony is of the highest value, and none need be afraid to use the books on such a recommendation.

The numerous notices from all parts of the country where these books have been used, cannot be introduced here. They have just gone into the schools of Seneca County, N. Y., without solicitation; and the same is true of many important schools where they have been examined.

From C. B. CRUMB, N. Y.

The Students' Series is, in my opinion, *the best in use*. I believe a class of young students will learn *twice as much*, with the same labor, as they would from any other system. The books of this Series excel in the purity and attraction of their style I have introduced them.

DR. COMSTOCK'S SERIES OF BOOKS ON THE SCIENCES, viz:

INTRODUCTION TO NATURAL PHILOSOPHY. For Children.....	\$0 41
SYSTEM OF NATURAL PHILOSOPHY, newly revised and enlarged, including late discoveries..	1 04
ELEMENTS OF CHEMISTRY. Adapted to the present state of the Science.....	1 00
THE YOUNG BOTANIST. New edition.....	50
ELEMENTS OF BOTANY. Including Vegetable Physiology, and a Description of Common Plants. With Cuts.....	1 25
OUTLINES OF PHYSIOLOGY, both Comparative and Human. To which is added OUTLINES OF ANATOMY, excellent for the general scholar and ladies' schools.	80
NEW ELEMENTS OF GEOLOGY. Highly Illustrated.....	1 25
ELEMENTS OF MINERALOGY. Illustrated with numerous Cuts.....	75
NATURAL HISTORY OF BIRDS. Showing their Comparative Size. A new and valuable feature.....	50
NATURAL HISTORY OF BEASTS. Ditto.....	50
NATURAL HISTORY OF FERES AND BEASTS. Do. Cloth.....	1 00
QUESTIONS AND ILLUSTRATIONS TO THE PHILOSOPHY.....	30

All the above works are fully illustrated by elegant cuts.

The Philosophy has been republished in Scotland, and translated for the use of schools in Prussia. The many valuable additions to the work by its transatlantic editors, Prof. Lees, of Edinburgh, and Prof. Hoblyn, of Oxford, have been embraced by the author in his last revision. The Chemistry has been entirely revised, and contains all the late discoveries, together with methods of analyzing minerals and metals. Portions of the series are in course of publication in London. Such testimony, in addition to the general good testimony of teachers in this country, is sufficient to warrant us in saying that no works on similar subjects can equal them, or have ever been so extensively used. Continual applications are made to the publishers to replace the Philosophy in schools where, for a time, it has given way to other books. The style of Dr. Comstock is so clear, and his arrangement is so excellent, that no writer can be found to excel him for school purposes, and he takes constant pains to include new discoveries, and to consult eminently scientific men.

HON. J. OLNEY'S GEOGRAPHICAL SERIES.

PRIMARY GEOGRAPHY; with Colored Maps. 25 cents.

QUARTO GEOGRAPHY; with elegant Cuts, Physical Geography Tables, Map of the Atlantic Ocean &c. 75 cents

OLNEY'S SCHOOL GEOGRAPHY AND ATLAS. Containing Ancient Geography, Physical Geography, Tables, an entirely new Chart of the World, to show its physical conformation, as adapted to purposes of commerce, and also for the purpose of reviewing classes; also a Chronological Table of Discoveries. \$1 12.

OLNEY'S OUTLINE MAPS. Of the World, United States, Europe, Asia, Africa, America, and Canada, with Portfolio and Book of Exercises \$6.

All the recent improvements are included in Olney's Quarto and School Geographies. They are not obsolete or out of date, but fully "up to the times." In elegance or completeness they are not surpassed.

Mr. Olney commenced the plan of simplifying the first lesson, and teaching a child by what is familiar, to the exclusion of astronomy. He commenced the plan of having only those things represented on the maps which the pupil was required to learn. He originated the system of classification, and of showing the government, religion, &c., by symbols. He first adopted the system of carrying the pupil over the earth by means of the Atlas. His works first contained cuts, in which the dress, architecture, animals, internal improvements, &c., of each country are grouped, so as to be seen at one view. His works first contained the world as known to the Ancients, as an aid to Ancient History, and a Synopsis of Physical Geography, with maps. In short, we have seen no valuable feature in any geography which has not originally appeared in these works; and we think it not too much to claim that, in many respects, most other works are copies of these. We think that a *fair and candid examination* will show that Olney's Atlas is the largest, most systematic, and complete of any yet published, and that the Quarto and Modern School Geographies contain more matter, and that better arranged, than any similar works; and they are desired to test the claims here asserted.

It is impossible to give here more than a fractional part of the recommendations, of the first order, which the publishers have received for the foregoing list of books. Enough has been given to show the claims of the books to examination and use.

All these works are made in very neat, durable style, and are sold as low as a moderate remuneration will allow. Copies supplied to teachers for their own use at one-fifth off from the retail price, and postage paid. Large institutions are furnished sample copies without charge.

PRATT, OAKLEY & CO.

21 MURRAY STREET, NEW YORK.



512.02 D639E2 23457
Dodd P cop.1

Dodd's Algebra

DATE DUE	ISSUED TO
EB 20 '51	<i>Karl Gane</i>

512.02 D639E2 cop.1 23457
P

D031140870



Duke University Libraries