

DUDLEY TOWN LIBRARY
BAY STATE SCHOOL
MORRIS

NAVAL POSTGRADUATE SCHOOL Monterey, California



THESIS

B1251

MODELS FOR TARGET DETECTION TIMES

by

Bae, Deok Hwan

September 1989

Thesis Advisor

Patricia A. Jacobs

Approved for public release; distribution is unlimited.

Unclassified
Security classification of this page

REPORT DOCUMENTATION PAGE

Report Security Classification Unclassified		1b Restrictive Markings			
Security Classification Authority		3 Distribution Availability of Report Approved for public release; distribution is unlimited.			
4 Declassification/Downgrading Schedule		5 Monitoring Organization Report Number(s)			
6a Name of Performing Organization Naval Postgraduate School		6b Office Symbol (if applicable) 55jc	7a Name of Monitoring Organization Naval Postgraduate School		
6c Address (city, state, and ZIP code) Monterey, CA 93943-5000		7b Address (city, state, and ZIP code) Monterey, CA 93943-5000			
8a Name of Funding Sponsoring Organization		8b Office Symbol (if applicable)	9 Procurement Instrument Identification Number		
6c Address (city, state, and ZIP code)		10 Source of Funding Numbers			
		Program Element No	Project No	Task No	Work Unit Accession No
1 Title (Include security classification) MODELS FOR TARGET DETECTION TIMES					
2 Personal Author(s) Bae, Deok Hwan					
3a Type of Report Master's Thesis		13b Time Covered From To	14 Date of Report (year, month, day) September 1989	15 Page Count 102	
6 Supplementary Notation The views expressed in this thesis are those of the author and do not reflect the official policy or position of the Department of Defense or the U.S. Government.					
7 Cosat: Codes		18 Subject Terms (continue on reverse if necessary and identify by block number)			
Field	Group	Subgroup	Weibull Regression, hierarchical models, maximum likelihood estimates, generalized linear models.		
9 Abstract (continue on reverse if necessary and identify by block number) Some battlefield models have a component in them which models the time it takes for an observer to detect a target. Different observers may have different mean detection times due to various factors such as the type of sensor used, environmental conditions, fatigue of the observer, etc. Two parametric models for the distribution of time to target detection are considered which can incorporate these factors. Maximum likelihood estimation procedures for the parameters are described. Results of simulation experiments to study the small sample behavior of the estimators are presented.					
22a Name of Responsible Individual Patricia A. Jacobs			22b Telephone (Include Area code) (408) 646-2258	22c Office Symbol 62Pw	

FORM 1473, 84 MAR

83 APR edition may be used until exhausted
All other editions are obsolete

security classification of this page

Unclassified

T245453

Approved for public release; distribution is unlimited.

Models for Target Detection Times

by

Bae, Deok Hwan
Major, Korean Air Force
B.S., Korean Air Force Academy, 1979

Submitted in partial fulfillment of the
requirements for the degree of

MASTER OF SCIENCE IN OPERATIONS RESEARCH

from the

NAVAL POSTGRADUATE SCHOOL
September 1989

ABSTRACT

Some battlefield models have a component in them which models the time it takes for an observer to detect a target. Different observers may have different mean detection times due to various factors such as the type of sensor used, environmental conditions, fatigue of the observer, etc. Two parametric models for the distribution of time to target detection are considered which can incorporate these factors. Maximum likelihood estimation procedures for the parameters are described. Results of simulation experiments to study the small sample behavior of the estimators are presented.

TABLE OF CONTENTS

I. INTRODUCTION	1
II. A HIERARCHICAL MODEL FOR TIMES TO DETECTION	2
A. MODEL	2
B. ESTIMATION	3
1. The Likelihood Equation and Maximum Likelihood Estimates	3
2. Initial Condition	7
3. Implementation in Simulation	11
C. SIMULATION PROCEDURES AND RESULTS	12
1. Simulation	12
2. Results	13
III. A WEIBULL REGRESSION MODEL	21
A. MODEL	21
B. ESTIMATION	22
1. The likelihood Equation and Maximum Likelihood Estimators.	22
a. Newton's Procedure to solve for β_i	23
b. Newton Procedure to solve for ξ_i	24
2. Initial Condition	25
3. Recursive Procedure to find $\{\beta_i\}$ and ξ_i	26
4. Implementation in Simulation	28
C. SIMULATION PROCEDURE AND RESULTS	29
1. Simulation	29
2. Results	29
IV. CONCLUSION	37
V. LIST OF REFERENCES	38
APPENDIX A. HISTOGRAMS FOR THE ESTIMATED GAMMA PARAME- TERS	39

APPENDIX B. HISTOGRAM FOR THE WEIBULL REGRESSION PARAMETERS	57
APPENDIX C. SIMULATION PROGRAM FOR THE ESTIMATION OF GAMMA PARAMETERS	84
APPENDIX D. SIMULATION PROGRAM FOR THE ESTIMATION OF REGRESSION PARAMETERS	86
INITIAL DISTRIBUTION LIST	88

LIST OF TABLES

Table 1.	MEAN BIAS, MEAN SQUARE ERROR AND STANDARD ERROR AT $O=10$ AND $UC=37\%$	15
Table 2.	MEAN BIAS AND MEAN SQUARE ERROR AT $O=10$ AND $UC=37\%$	16
Table 3.	MEAN BIAS AND MEAN SQUARE ERROR AT $O=10$ AND $UC=37\%$	16
Table 4.	MEAN BIAS, MEAN SQUARE ERROR AND STANDARD ERROR AT $O=25$ AND $UC=62\%$	17
Table 5.	MEAN BIAS AND MEAN SQUARE ERROR AT $O=25$ AND $UC=62\%$	18
Table 6.	MEAN BIAS AND MEAN SQUARE ERROR AT $O=25$ AND $UC=62\%$	18
Table 7.	MEAN BIAS, MEAN SQUARE ERROR AND STANDARD ERROR AT $O=40$ AND $UC=75\%$	19
Table 8.	MEAN BIAS AND MEAN SQUARE ERROR AT $O=40$ AND $UC=75\%$	20
Table 9.	MEAN BIAS AND MEAN SQUARE ERROR AT $O=40$ AND $UC=75\%$	20
Table 10.	MEAN BIAS, MEAN SQUARE ERROR AND STANDARD ERROR AT $O=1.3$ AND $UC=45\%$	31
Table 11.	MEAN BIAS AND MEAN SQUARE ERROR AT $O=1.3$ AND $UC=45\%$	32
Table 12.	MEAN BIAS AND MEAN SQUARE ERROR AT $O=1.3$ AND $UC=45\%$	32
Table 13.	MEAN BIAS AND MEAN SQUARE ERROR AT $O=1.3$ AND $UC=45\%$	32
Table 14.	MEAN BIAS, MEAN SQUARE ERROR AND STANDARD ERROR AT $O=2.5$ AND $UC=75\%$	33
Table 15.	MEAN BIAS AND MEAN SQUARE ERROR AT $O=2.5$ AND $UC=75\%$	34
Table 16.	MEAN BIAS AND MEAN SQUARE ERROR AT $O=2.5$ AND	

UC = 75%	34
Table 17. MEAN BIAS AND MEAN SQUARE ERROR AT O=2.5 AND UC = 75%	34
Table 18. MEAN BIAS, MEAN SQUARE ERROR AND STANDARD ERROR AT O=4.0 AND UC=89%	35
Table 19. MEAN BIAS AND MEAN SQUARE ERROR AT O=4.0 AND UC = 89%	36
Table 20. MEAN BIAS AND MEAN SQUARE ERROR AT O=4.0 AND UC = 89%	36
Table 21. MEAN BIAS AND MEAN SQUARE ERROR AT O=4.0 AND UC = 89%	36

LIST OF FIGURES

Figure 1.	COMPARISON BETWEEN DIFFERENT ESTIMATES (O = 10, UC = 37%)	39
Figure 2.	COMPARISON BETWEEN DIFFERENT ESTIMATES (O = 25, UC = 62%)	40
Figure 3.	COMPARISON BETWEEN DIFFERENT ESTIMATES (O = 40, UC = 75%)	41
Figure 4.	COMPARISON BETWEEN DIFFERENT ESTIMATES (O = 10, UC = 37%)	42
Figure 5.	COMPARISON BETWEEN DIFFERENT ESTIMATES (O = 25, UC = 62%)	43
Figure 6.	COMPARISON BETWEEN DIFFERENT ESTIMATES (O = 40, UC = 75%)	44
Figure 7.	COMPARISON BETWEEN DIFFERENT ESTIMATES (O = 10, UC = 37%)	45
Figure 8.	COMPARISON BETWEEN DIFFERENT ESTIMATES (O = 25, UC = 62%)	46
Figure 9.	COMPARISON BETWEEN DIFFERENT ESTIMATES (O = 40, UC = 75%)	47
Figure 10.	COMPARISON BETWEEN DIFFERENT ESTIMATES (O = 10, UC = 37%)	48
Figure 11.	COMPARISON BETWEEN DIFFERENT ESTIMATES (O = 25, UC = 62%)	49
Figure 12.	COMPARISON BETWEEN DIFFERENT ESTIMATES (O = 40, UC = 75%)	50
Figure 13.	COMPARISON BETWEEN DIFFERENT ESTIMATES (O = 10, UC = 37%)	51
Figure 14.	COMPARISON BETWEEN DIFFERENT ESTIMATES (O = 25, UC = 62%)	52
Figure 15.	COMPARISON BETWEEN DIFFERENT ESTIMATES (O = 40, UC = 75%)	53
Figure 16.	COMPARISON BETWEEN DIFFERENT ESTIMATES (O = 10,	

UC = 37%)	54
Figure 17. COMPARISON BETWEEN DIFFERENT ESTIMATES (O = 25, UC = 62%)	55
Figure 18. COMPARISON BETWEEN DIFFERENT ESTIMATES (O = 40, UC = 75%)	56
Figure 19. COMPARISON BETWEEN DIFFERENT ESTIMATES (O = 1.3, UC = 45%)	57
Figure 20. COMPARISON BETWEEN DIFFERENT ESTIMATES (O = 2.5, UC = 75%)	58
Figure 21. COMPARISON BETWEEN DIFFERENT ESTIMATES (O = 4.0, UC = 89%)	59
Figure 22. COMPARISON BETWEEN DIFFERENT ESTIMATES (O = 1.3, UC = 45%)	60
Figure 23. COMPARISON BETWEEN DIFFERENT ESTIMATES (O = 2.5, UC = 75%)	61
Figure 24. COMPARISON BETWEEN DIFFERENT ESTIMATES (O = 4.0, UC = 89%)	62
Figure 25. COMPARISON BETWEEN DIFFERENT ESTIMATES (O = 1.3, UC = 45%)	63
Figure 26. COMPARISON BETWEEN DIFFERENT ESTIMATES (O = 2.3, UC = 75%)	64
Figure 27. COMPARISON BETWEEN DIFFERENT ESTIMATES (O = 4.0, UC = 89%)	65
Figure 28. COMPARISON BETWEEN DIFFERENT ESTIMATES (O = 1.3, UC = 45%)	66
Figure 29. COMPARISON BETWEEN DIFFERENT ESTIMATES (O = 2.5, UC = 75%)	67
Figure 30. COMPARISON BETWEEN DIFFERENT ESTIMATES (O = 4.0, UC = 89%)	68
Figure 31. COMPARISON BETWEEN DIFFERENT ESTIMATES (O = 1.3, UC = 45%)	69
Figure 32. COMPARISON BETWEEN DIFFERENT ESTIMATES (O = 2.5, UC = 75%)	70
Figure 33. COMPARISON BETWEEN DIFFERENT ESTIMATES (O = 4.0, UC = 89%)	71

Figure 34. COMPARISON BETWEEN DIFFERENT ESTIMATES (O= 1.3, UC= 45%)	72
Figure 35. COMPARISON BETWEEN DIFFERENT ESTIMATES (O= 2.5, UC= 75%)	73
Figure 36. COMPARISON BETWEEN DIFFERENT ESTIMATES (O= 4.0, UC= 89%)	74
Figure 37. COMPARISON BETWEEN DIFFERENT ESTIMATES (O= 1.3, UC= 45%)	75
Figure 38. COMPARISON BETWEEN DIFFERENT ESTIMATES (O= 2.5, UC= 75%)	76
Figure 39. COMPARISON BETWEEN DIFFERENT ESTIMATES (O= 4.0, UC= 89%)	77
Figure 40. COMPARISON BETWEEN DIFFERENT ESTIMATES (O= 1.3, UC= 45%)	78
Figure 41. COMPARISON BETWEEN DIFFERENT ESTIMATES (O= 2.5, UC= 75%)	79
Figure 42. COMPARISON BETWEEN DIFFERENT ESTIMATES (O= 4.0, UC= 89%)	80
Figure 43. COMPARISON BETWEEN DIFFERENT ESTIMATES (O= 1.3, UC= 45%)	81
Figure 44. COMPARISON BETWEEN DIFFERENT ESTIMATES (O= 2.3, UC= 75%)	82
Figure 45. COMPARISON BETWEEN DIFFERENT ESTIMATES (O= 4.0, UC= 89%)	83

ACKNOWLEDGEMENTS

For their significant contributions toward this thesis, I would like to thank the following people :

- Professor P.A.Jacobs for her tolerance, professional assistance, and thorough review.
- Professor D.P.Gaver for his encouragement and guidance.
- Most especially, my wife, In-Sook, my daughters, Yun-Jeong and Yun-Hae for their prayers, patience, and earnest support.

I. INTRODUCTION

Some battlefield models have a component in them which models the time it takes for an observer to detect a target. A common model is that the time to detect a target is a random variable having a distribution $P_{\infty}(1 - e^{-\lambda t})$, [Ref. 1]. However different observers may have different mean detection times due to various factors such as the type of sensor used, environmental condition, fatigue of the observer, etc. In this thesis we will consider two parametric models for the distribution of the time to detection which can incorporate these factors.

Suppose there are M observers, such as tank crews. Observer i is presented with N_i targets. In Chapter 2, observer i has a parameter A_i , which reflects the ability of the observer to detect a target. The parameters of the M observers, $A_1, A_2, A_3, \dots, A_M$, are assumed independent, having a common gamma distribution. Given A_i , the detection times of observer i are conditionally independent, having Weibull distributions. In chapter 2 the parameters of the conditional Weibull distributions are assumed known and interest is in estimating the parameters of the gamma distribution.

In chapter 3 observer i has explanatory variables $x_{i1}, x_{i2}, x_{i3}, \dots, x_{ip}$ relating to his j^{th} target representing factors which influence his time to detection. The detection times for the observers are independent random variables having Weibull distributions. The location parameter of the Weibull distribution for the detection time of the j^{th} target by the i^{th} observer is of the form $\mu_{ij} = e^{\xi_i} \underline{\mu}$ where $\underline{\mu} = \sum_{k=1}^p x_{ijk} \beta_k$; the shape parameter is of the form e^{ξ_i} . Interest is in estimation of $\{\beta_k\}$ and $\{\xi_i\}$.

In both chapters the data for the observers can be censored. The i^{th} observer has an opportunity time O_{ij} to detect his j^{th} target. If the j^{th} target is not detected in time O_{ij} , the time to detection is censored. Data for the i^{th} observer consist of (possibly censored) times to detect the N_i targets.

Chapter 2 and 3 present iterative procedures based on maximum likelihood to estimate the parameters. Results from simulation studies of small sample size behavior of the estimators are given.

These models and estimation procedures should be of eventual use in the Army MANPRINT program, an objective of which is to better understand the human contribution to battlefield performance [Ref. 2].

II. A HIERARCHICAL MODEL FOR TIMES TO DETECTION

A. MODEL

Suppose there are M observers who attempt to detect targets on a battlefield. Observer i has a parameter A_i , $i = 1, 2, 3, \dots, M$, which reflects the ability of the observer to detect a target. Observer i is presented with N_i targets. The targets are presented one at a time. Let U_{ij} be the time it takes for observer i to detect target j . Assume, given $A_i = \theta$, the U_{ij} , $j = 1, 2, 3, \dots, N_i$, are conditionally independent random variables with Weibull distributions

$$P\{U_{ij} \leq t \mid A_i = \theta\} = 1 - \exp\left\{-\theta(t/\mu_{ij})^{\epsilon}\right\} \quad t \geq 0 \quad (2.1)$$

independent of other observers.

Further, assume the parameters A_i , $i = 1, 2, 3, \dots, M$, are independent identically distributed having a Gamma distribution with density function as follows:

$$g(\theta) = \frac{\alpha(\alpha\theta)^{\gamma-1}}{\Gamma(\gamma)} e^{-\alpha\theta} \quad (2.2)$$

where $\theta > 0$. The A_i variations are introduced to represent individual observer differences. Now let

$$Z_{ij} = \ln U_{ij} .$$

It follows that

$$\begin{aligned} P\{Z_{ij} \leq z \mid A_i = \theta\} &= P\{U_{ij} \leq e^z \mid A_i = \theta\} \\ &= 1 - \exp\left\{-\theta[\exp(z - \ln \mu_{ij})]^{\epsilon}\right\} . \end{aligned} \quad (2.3)$$

Hence, the conditional density function of Z_{ij} , given $A_i = \theta$, is

$$f_{Z_{ij} \mid A_i}(z \mid \theta) = \theta \exp\{(z - \ln \mu_{ij})e^{-\frac{z}{\epsilon}}\} e^{-\frac{z}{\epsilon}} \exp\{-\theta(\exp\{(z - \ln \mu_{ij})e^{-\frac{z}{\epsilon}}\})\} . \quad (2.4)$$

When the i^{th} observer is presented with his j^{th} target, he gets a length of time called opportunity time O_{ij} to detect it. An observer either successfully detects the target within this time or is unsuccessful. Data for the i^{th} observer consist of times of detections for

the successes and the lengths of opportunity times for the failures. For each $i = 1, 2, 3, \dots, M$, $j = 1, 2, 3, \dots, N_i$, let

$$Y_{ij} = \min(\ln U_{ij}, \ln O_{ij}) \quad (2.5)$$

and

$$\Delta_{ij} = \begin{cases} 1 & \text{if } U_{ij} \leq O_{ij} \\ 0 & \text{otherwise} \end{cases} \quad (2.6)$$

The Y_{ij} are the censored ln-detection times and Δ_{ij} is an indicator of whether or not the ln-time to detect the j^{th} target by the i^{th} observer is censored. Let

$$C_i = \sum_{j=1}^{N_i} \Delta_{ij} \quad (2.7)$$

be the number of targets detected by observer i .

In this Chapter we will assume $\{\mu_{ij}\}$ and $\{\xi_i\}$ are known constants. We are interested in estimating the parameters η and β_0 with $\alpha \equiv e^\eta$ and $\frac{\gamma}{\alpha} \equiv e^{\beta_0}$ using maximum likelihood. These parameter estimates can be used to predict future times to detection for an observer given his past performance. In the next section we describe a Newton - Raphson procedure for solving the likelihood equations, [Ref. 3], [Ref. 4]. In the final section of this chapter we describe a simulation experiment to study the small sample properties of the estimators. Results of the simulation are also presented.

B. ESTIMATION

1. The Likelihood Equation and Maximum Likelihood Estimates

Given $A_i = \theta$, it follows from equation (2.3) and (2.4) that the conditional likelihood function for observer i using the censored ln-times y_{ij} is

$$L_i(\underline{\mu}, \underline{\xi}; \theta) = \prod_{j=1}^{N_i} \left[\theta e^{(\theta y_{ij} - \ln \mu_{ij})e^{-\xi_i}} e^{-\xi_i} \right]^{\Delta_{ij}} \exp \left[-\theta e^{(\theta y_{ij} - \ln \mu_{ij})e^{-\xi_i}} \right] \quad (2.8)$$

Let

$$S_i = \sum_{j=1}^{N_i} \exp[(y_{ij} - \ln \mu_{ij})e^{-\xi_i}] . \quad (2.9)$$

Rewriting equation (2.8)

$$\begin{aligned} L_i(\underline{\mu}, \underline{\xi}; \theta) &= \theta^{C_i} \exp\left\{\sum_{j=1}^{N_i} \Delta_{ij}[(y_{ij} - \ln \mu_{ij})e^{-\xi_i} - \xi_i]\right\} \exp(-\theta S_i) \\ &= \theta^{C_i} K_i \exp(-\theta S_i) \end{aligned} \quad (2.10)$$

where

$$K_i = \exp\left\{\sum_{j=1}^{N_i} \Delta_{ij}[(y_{ij} - \ln \mu_{ij})e^{-\xi_i} - \xi_i]\right\}. \quad (2.11)$$

The unconditional likelihood for observer i is

$$\begin{aligned} L_i(\underline{\mu}, \underline{\xi}, \alpha, \gamma) &= \int_0^{\infty} L_i(\underline{\mu}, \underline{\xi}; \theta) g(\theta) d\theta \\ &= K_i \int_0^{\infty} \theta^{C_i} e^{-\theta S_i} \frac{\alpha(\alpha\theta)^{\gamma-1}}{\Gamma(\gamma)} e^{-\alpha\theta} d\theta \\ &= K_i \alpha^{\gamma} \left(\frac{1}{S_i + \alpha}\right)^{C_i + \gamma} \prod_{k=0}^{C_i-1} (\gamma + k) . \end{aligned} \quad (2.12)$$

If $C_i = 0$, then $\prod_{k=0}^{C_i-1} (\gamma + k) = 1$. Recall the parametrization $\alpha = e^{\eta}$ and $\frac{\gamma}{\alpha} = e^{\beta_0}$. The unconditional ln-likelihood function for observer i can be rewritten as

$$\begin{aligned} \ln L_i(\underline{\mu}, \underline{\xi}, \eta, \beta_0) &= \ln K_i + e^{\eta+\beta_0}\eta - (C_i + e^{\eta+\beta_0}) \ln(S_i + e^{\eta}) \\ &\quad + \sum_{k=0}^{C_i-1} \ln(e^{\eta+\beta_0} + k) \end{aligned} \quad (2.13)$$

where if $C_i = 0$, then $\sum_{k=0}^{C_i-1} \ln(e^{\eta+\beta_0} + k) = 0$. Since the observers are independent, the unconditional ln-likelihood for all M observers is

$$\ln L = \sum_{i=1}^M \left\{ \ln K_i + e^{\eta+\beta_0}\eta - (C_i + e^{\eta+\beta_0}) \ln(S_i + e^{\eta}) + \sum_{k=0}^{C_i-1} \ln(e^{\eta+\beta_0} + k) \right\} .$$

The derivative of the ln-likelihood with respect to η is

$$\begin{aligned} \frac{\partial \ln L}{\partial \eta} &= \sum_{i=1}^M \left\{ e^{\eta + \beta_0 \eta} + e^{\eta + \beta_0} - e^{\eta + \beta_0} \ln(S_i + e^\eta) - \left(\frac{C_i + e^{\eta + \beta_0}}{S_i + e^\eta} \right) e^\eta \right\} \\ &+ \sum_{i=1}^M \sum_{k=0}^{C_i-1} \left(\frac{e^{\eta + \beta_0}}{e^{\eta + \beta_0} + k} \right). \end{aligned} \quad (2.15)$$

The derivative with respect to β_0 is

$$\frac{\partial \ln L}{\partial \beta_0} = \sum_{i=1}^M \left\{ e^{\eta + \beta_0 \eta} - e^{\eta + \beta_0} \ln(S_i + e^\eta) + \sum_{k=0}^{C_i-1} \frac{e^{\eta + \beta_0}}{e^{\eta + \beta_0} + k} \right\}. \quad (2.16)$$

We are assuming $\{\mu_i\}$ and $\{\xi_i\}$ are known. The problem is to find the maximum likelihood estimates of η and β_0 ; that is, find η and β_0 such that $\frac{\partial \ln L}{\partial \eta} = 0$ and $\frac{\partial \ln L}{\partial \beta_0} = 0$. Note that

$$\frac{\partial \ln L}{\partial \eta} = \frac{\partial \ln L}{\partial \beta_0} + \sum_{i=1}^M \left[e^{\eta + \beta_0} - \left(\frac{C_i + e^{\eta + \beta_0}}{S_i + e^\eta} \right) e^\eta \right]. \quad (2.17)$$

Thus, if $\frac{\partial \ln L}{\partial \beta_0} = 0$, then to solve the equation $\frac{\partial \ln L}{\partial \eta} = 0$, we need to solve the equation

$$\begin{aligned}
0 = f(\eta, \beta_0) &= \sum_{i=1}^M \left[e^{\eta + \beta_0} - \left(\frac{C_i + e^{\eta + \beta_0}}{S_i + e^{\eta}} \right) e^{\eta} \right] \\
&= \sum_{i=1}^M \frac{e^{\eta + \beta_0}(S_i + e^{\eta}) - C_i e^{\eta} - e^{2\eta + \beta_0}}{S_i + e^{\eta}} \\
&= \sum_{i=1}^M e^{\eta} \frac{(e^{\beta_0} S_i - C_i)}{S_i + e^{\eta}} .
\end{aligned} \tag{2.18}$$

The derivatives of $f(\eta, \beta_0)$ with respect to η and β_0 are as follows :

$$\begin{aligned}
\frac{\partial f}{\partial \eta} &= - \sum_{i=1}^M \frac{(e^{\beta_0} S_i - C_i) e^{\eta}}{(S_i + e^{\eta})^2} \\
&= e^{\eta} f(\eta, \beta_0) ;
\end{aligned} \tag{2.19}$$

$$\frac{\partial f}{\partial \beta_0} = \sum_{i=1}^M \frac{e^{\beta_0}}{S_i + e^{\eta}} . \tag{2.20}$$

Solving the equation $\frac{\partial \ln L}{\partial \beta_0} = 0$ is equivalent to solving the equation

$$0 = g(\eta, \beta_0) \equiv \sum_{i=1}^M \left[\eta - \ln(S_i + e^{\eta}) + \sum_{k=0}^{C_i-1} \frac{1}{e^{\eta + \beta_0} + k} \right] . \tag{2.21}$$

The derivatives of $g(\eta, \beta_0)$ with respect to η and β_0 are

$$\frac{\partial g}{\partial \eta} = \sum_{i=1}^M \left[1 - \frac{e^{\eta}}{S_i + e^{\eta}} - \sum_{k=0}^{C_i-1} \frac{e^{\eta + \beta_0}}{(e^{\eta + \beta_0} + k)^2} \right] ; \tag{2.22}$$

$$\frac{\partial g}{\partial \beta_0} = - \sum_{i=1}^M \sum_{k=0}^{C_i-1} \frac{e^{\eta + \beta_0}}{(e^{\eta + \beta_0} + k)^2} \quad (2.23)$$

where if $C_i = 0$, then the sum involving $C_i - 1$ is zero. A Newton procedure to solve the equations $\frac{\partial L}{\partial \eta} = 0$ and $\frac{\partial L}{\partial \beta_0} = 0$ would use the following equations

$$0 = f(\eta^0, \beta_0^0) + \frac{\partial f(\eta^0, \beta_0^0)}{\partial \eta} (\eta - \eta^0) + \frac{\partial f(\eta^0, \beta_0^0)}{\partial \beta_0} (\beta_0 - \beta_0^0) \quad (2.24)$$

$$0 = g(\eta^0, \beta_0^0) + \frac{\partial g(\eta^0, \beta_0^0)}{\partial \eta} (\eta - \eta^0) + \frac{\partial g(\eta^0, \beta_0^0)}{\partial \beta_0} (\beta_0 - \beta_0^0) \quad (2.25)$$

where η^0 and β_0^0 are current values for η and β_0 . However note that if $f(\eta, \beta_0) = 0$, then $\frac{\partial f}{\partial \eta} = 0$. Hence, a Newton procedure that is more stable numerically would use the equations

$$0 = f(\eta^0, \beta_0^0) + \frac{\partial f(\eta^0, \beta_0^0)}{\partial \beta_0} (\beta_0 - \beta_0^0) \quad (2.26)$$

$$0 = g(\eta^0, \beta_0^0) + \frac{\partial g(\eta^0, \beta_0^0)}{\partial \eta} (\eta - \eta^0) \quad (2.27)$$

which results in

$$\beta_0 - \beta_0^0 = \frac{-f(\eta^0, \beta_0^0)}{\frac{\partial f}{\partial \beta_0}(\eta^0, \beta_0^0)} \quad (2.28)$$

$$\eta_0 - \eta_0^0 = \frac{-g(\eta^0, \beta_0^0)}{\frac{\partial g}{\partial \eta}(\eta^0, \beta_0^0)} \quad (2.29)$$

2. Initial Condition

In this subsection, we describe a rough way to provide initial estimates of η and

β_0 to start the iterative Newton procedure of the previous section.

The uncensored random variable U_{ij} has the same distribution as

$$U_{ij} = \mu_{ij} \left(\frac{W_{ij}}{A_i} \right) e^{t_i} \quad (2.30)$$

where W_{ij} is unit exponential random variable. Let

$$\begin{aligned} Z_{ij} &= \ln U_{ij} \\ &= \ln \mu_{ij} + e^{t_i} \ln W_{ij} - e^{t_i} \ln A_i ; \end{aligned} \quad (2.31)$$

then

$$E[\ln W_{ij}] = -0.5772 \quad (2.32)$$

$$VAR[\ln W_{ij}] = \frac{\pi^2}{6} . \quad (2.33)$$

[Ref. 5: p.943]. An approximation to the moments of $\ln A_i$ is given by the first two terms of a Taylor expansion of $\ln(A_i)$ about the mean $E[A_i] = \frac{\gamma}{\alpha}$

$$\ln(A_i) \cong \ln\left(\frac{\gamma}{\alpha}\right) + \frac{1}{\left(\frac{\gamma}{\alpha}\right)} \left(A_i - \frac{\gamma}{\alpha}\right); \quad (2.34)$$

thus

$$E[\ln A_i] \cong \ln \gamma - \ln \alpha ; \quad (2.35)$$

$$\begin{aligned} VAR[\ln A_i] &\cong \left(\frac{\alpha}{\gamma}\right)^2 VAR[A_i] \\ &= \left(\frac{\alpha}{\gamma}\right)^2 \frac{\gamma}{\alpha^2} \\ &= \frac{1}{\gamma} . \end{aligned} \quad (2.36)$$

Let

$$\begin{aligned} V_{ij} &= (Z_{ij} - \ln \mu_{ij}) e^{-t_i} \\ &= \ln W_{ij} - \ln A_i . \end{aligned} \quad (2.37)$$

Hence,

$$E[V_{ij}] = -0.5772 - (\ln \gamma - \ln \alpha) \quad (2.38)$$

$$\begin{aligned} VAR[V_{ij}] &= VAR_2[AR[\ln W_{ij}]] + VAR[\ln A_{ij}] - 2cov(\ln W_{ij}, \ln A_{ij}) \\ &= \frac{\pi^2}{6} + \frac{1}{\gamma} . \end{aligned} \quad (2.39)$$

Rough estimates of the first and second moments of V_{ij} are given by

$$\hat{M}_1 = \frac{\sum_{i=1}^M \sum_{j=1}^{N_i} \Delta_{ij} (Z_{ij} - \ln \mu_{ij}) e^{-Z_{ij}}}{\sum_{i=1}^M \sum_{j=1}^{N_i} \Delta_{ij}} \quad (2.40)$$

and

$$\hat{M}_2 = \frac{\sum_{i=1}^M \sum_{j=1}^{N_i} \Delta_{ij} [(Z_{ij} - \ln \mu_{ij}) e^{-Z_{ij}}]^2}{\sum_{i=1}^M \sum_{j=1}^{N_i} \Delta_{ij}} . \quad (2.41)$$

Recall the parametrization

$$\alpha = e^\eta ; \quad (2.42)$$

$$\frac{\gamma}{\alpha} = e^{\beta_0} . \quad (2.43)$$

Thus

$$\gamma = e^\eta e^{\beta_0} \quad (2.44)$$

and

$$\begin{aligned} E[V_{ij}] &= -0.5772 - \ln \gamma + \ln \alpha \\ &= -0.5772 - [\eta + \beta_0] + \eta \\ &= -0.5772 - \beta_0 ; \end{aligned} \quad (2.45)$$

$$\begin{aligned}
 VAR[V_{ij}] &= \frac{\pi^2}{6} + \frac{1}{\gamma} \\
 &= \frac{\pi^2}{6} + e^{-(\eta + \beta_0)} .
 \end{aligned}
 \tag{2.46}$$

Equating the rough estimates of the moments of V_{ij} with the mean and variance of V_{ij} we obtain initial estimates


$$\hat{\beta}_0 = -\hat{M}_1 - 0.5772 ; \tag{2.47}$$

$$\hat{\eta} = -\ln[\hat{M}_2 - (\hat{M}_1)^2 - \frac{\pi^2}{6}] - \hat{\beta}_0 . \tag{2.48}$$

If $\hat{M}_2 - (\hat{M}_1)^2 - \frac{\pi^2}{6} < 0$, we set $\hat{\eta} = -\hat{\beta}_0$. The latter condition usually occurs in the simulations indicating that a better initial condition might be found.

3. Implementation in Simulation

The following flow diagram describes one replication of a simulation experiment.



C. SIMULATION PROCEDURES AND RESULTS

1. Simulation

All simulations were carried out on an IBM 3179 G computer at the Naval Postgraduate School using the APL GRAFSTAT random number package [Ref. 6]. Histograms of simulated estimates were produced by an experimental APL package GRAFSTAT which the Naval Postgraduate School is using under a test agreement with IBM Watson Research center, Yorktown, Height, NY. The simulation is replicated for $R = 100$ replications. Each simulation experiment of 100 replications starts with the same random number seed. The mean bias (M.B) and its standard error (S.E(M.B)), mean square error (M.S.E) and its standard error (S.E(M.S.E)) from $R = 100$ replications are computed as follows :

$$M.B = \frac{1}{R} \sum_{i=1}^R (\hat{\beta}_i - \beta) \quad (2.49a)$$

$$S.E(M.B) = \sqrt{\frac{1}{R(R-2)} \sum_{i=1}^R ((\hat{\beta}_i - \beta) - M.B)^2} \quad (2.49b)$$

$$M.S.E = \frac{1}{R} \sum_{i=1}^R (\hat{\beta}_i - \beta)^2 \quad (2.50a)$$

$$S.E(M.S.E) = \sqrt{\frac{1}{R(R-2)} \sum_{i=1}^R ((\hat{\beta}_i - \beta)^2 - M.S.E)^2} \quad (2.50b)$$

where $\hat{\beta}_i$ is the point estimate of the true value β for the i th replication and R is the number of replications.

The simulation experiment to study the sampling properties of the estimators of η and β_0 is as follows.

1. Give arbitrary constants for μ_j and ξ_i ($\mu_j = 4.2$, $\xi_i = 0$)
2. Set the same opportunity time, O_{ij} , for M observers, $i = 1, 2, 3, \dots, M$, $j = 1, 2, 3, \dots, N_j$. The constant opportunity time is changed to give different censoring levels. The different values for O_{ij} are 10, 25, 40.

3. Generate gamma random variables for M observers ($A_1, A_2, A_3, \dots, A_M$ with shape parameter $\gamma = 1.5$, scale parameter $\alpha = 6.5$).
4. Generate independent exponential random variables with mean 1 ($W_{ij}, i = 1, 2, 3, \dots, M, j = 1, 2, 3, \dots, N_i$) .
5. Compute the detection time U_{ij} for observer i to detect target j

$$U_{ij} = \mu_{ij} \left(\frac{W_{ij}}{A_i} \right) e^{\xi_i}$$

6. Compare $\ln U_{ij}$ with $\ln O_{ij}$, then choose the smaller one for the data

$$Y_{ij} = \min(\ln U_{ij}, \ln O_{ij})$$

and compute the censoring indicator Δ_{ij} as in equation (2.6)

7. Compute the moments \hat{M}_i and \hat{M}_j for the observations that are not censored as in equation (2.40) and equation (2.41).
8. Compute the initial value for η and β_0 as in equation (2.47) and equation (2.48)
9. Use equation (2.28) and (2.29) to find new values for η and β_0
10. Iterate the procedure until the differences between successive values of η and β_0 are small, (less than 10^{-4}).

2. Results

In this section results from the simulation experiments will be reported. Simulation experiments were done for various numbers of observers and targets and values of the opportunity time. The numbers of observers considered are $M = 5, 15$, and 30 . The numbers of targets considered are $N_i = 5, 15, 30$, and 50 for each observer. For all simulation experiments, the $\mu_{ij} = 4.2$, $\xi_i = 0$, $\gamma = 1.5$ and $\alpha = 6.5$. From equations (2.42) and (2.44) the true values of η and β_0 are found to be 1.8718 and -1.4663 . Histograms of the difference between the estimates and the true values, $\hat{\eta}_i - \eta$ and $\hat{\beta}_0 - \beta_0$ are shown in Figures 1 through 18 of Appendix A. For Figures 1 through 6, $O_{ij} = 10$; for Figures 7 through 12, $O_{ij} = 25$; for Figures 13 through 18, $O_{ij} = 40$. The fraction of detected level or uncensoring level (UC) for R replications of each simulation is determined by following equation

$$UC = \frac{\sum_{r=1}^R \sum_{i=1}^M C_i(r)}{\sum_{r=1}^R \sum_{i=1}^M N_i(r)} \quad (2.51)$$

where $C_i(r)$ is numbers of targets detected by observer i in the r^{th} replication, $N_i(r)$ is the number of targets presented to observer i in the r^{th} replication. The UC's for the simulations using the same opportunity time are then averaged to obtain the mean UC for that opportunity time. For each Figure of Appendix A and B, the mean uncensoring level for the opportunity time is given in parenthesis. Mean bias and mean square error are recorded in tables 1 through 9 for each of the estimators; their standard errors appear in parenthesis below. Tables 1, 4 and 7 present all of the means and standard errors for different opportunity times. The other tables present the same results in a more convenient fashion; the mean biases are displayed with the mean square error in parenthesis. For each Table, the opportunity time(O) and average uncensoring level(UC) for that opportunity time are given at the top of table.

- Results for $\hat{\eta}$

The histograms of $\hat{\eta} - \eta$ are centered about 0 with a slight amount of skewness to the right. Increasing the opportunity time, which results in less censoring, has very little effect on mean bias and mean square error. Increasing the number of targets for a fixed number of observers has some tendency to decrease the mean square error and bias. Increasing the number of observers for a fixed number of targets has the greatest effect on decreasing the mean bias and mean square error.

- Results for $\hat{\beta}_0$

The histograms of $\hat{\beta}_0 - \beta_0$ are centered around 0 with some skewness to the left. Once again changing the opportunity time has little effect on mean bias and mean square error. Changing the number of targets for a fixed number of observers also has little effect. Increasing the number of observers for a fixed number of targets has the greatest effect on decreasing the mean bias and mean square error.

Table 1. MEAN BIAS, MEAN SQUARE ERROR AND STANDARD ERROR AT
O = 10 AND UC = 37%

Number of Observers	Number of Targets	$\hat{\eta}$		$\hat{\beta}_0$	
		M.B (S.E)	M.S.E (S.E)	M.B (S.E)	M.S.E (S.E)
5	5	0.79 (0.12)	2.04 (0.17)	-0.07 (0.06)	0.36 (0.09)
	15	0.58 (0.11)	1.33 (0.15)	-0.08 (0.05)	0.25 (0.05)
	30	0.42 (0.09)	0.97 (0.12)	-0.12 (0.04)	0.19 (0.03)
	50	0.43 (0.07)	0.68 (0.08)	-0.05 (0.04)	0.14 (0.03)
15	5	0.19 (0.09)	0.81 (0.10)	-0.02 (0.03)	0.12 (0.02)
	15	0.19 (0.08)	0.72 (0.01)	-0.01 (0.03)	0.07 (0.01)
	30	0.09 (0.04)	0.20 (0.03)	-0.07 (0.03)	0.07 (0.01)
	50	0.14 (0.05)	0.23 (0.04)	-0.02 (0.03)	0.06 (0.01)
30	5	0.29 (0.07)	0.54 (0.10)	-0.03 (0.02)	0.05 (0.01)
	15	0.13 (0.05)	0.21 (0.04)	-0.01 (0.02)	0.03 (0.01)
	30	0.08 (0.03)	0.10 (0.01)	-0.04 (0.02)	0.03 (0.01)
	50	0.01 (0.03)	0.09 (0.01)	+0.004 (0.02)	0.02 (0.003)

Table 2. MEAN BIAS AND MEAN SQUARE ERROR AT $O=10$ AND $UC=37\%$

	<i>M.B(M.S.E) for $\hat{\eta}$</i>			
	5 TGT	15 TGT	30 TGT	50 TGT
5 OBS	0.79(2.04)	0.58(1.3)	0.42(0.97)	0.43(1.68)
15 OBS	0.19(0.81)	0.19(0.72)	0.09(0.20)	0.14(0.23)
30 OBS	0.29(0.54)	0.13(0.21)	0.08(0.09)	0.01(0.09)

Table 3. MEAN BIAS AND MEAN SQUARE ERROR AT $O=10$ AND $UC=37\%$

	<i>M.B(M.S.E) for $\hat{\beta}_0$</i>			
	5 TGT	15 TGT	30 TGT	50 TGT
5 OBS	-0.07(0.36)	-0.08(0.25)	-0.12(0.19)	-0.05(0.14)
15 OBS	-0.02(0.12)	-0.009(0.07)	-0.07(0.07)	-0.02(0.06)
30 OBS	-0.03(0.05)	-0.01(0.03)	-0.04(0.03)	-0.004(0.02)

Table 4. MEAN BIAS, MEAN SQUARE ERROR AND STANDARD ERROR AT
O = 25 AND UC = 62%

Number of Observers	Number of Targets	$\hat{\eta}$		$\hat{\beta}_0$	
		M.B (S.E)	M.S.E (S.E)	M.B (S.E)	M.S.E (S.E)
5	5	0.83 (0.13)	2.23 (0.23)	-0.06 (0.05)	0.25 (0.07)
	15	0.52 (0.08)	0.97 (0.14)	-0.08 (0.05)	0.20 (0.04)
	30	0.47 (0.01)	1.00 (0.14)	-0.12 (0.04)	0.18 (0.03)
	50	0.47 (0.07)	0.76 (0.10)	-0.05 (0.04)	0.13 (0.02)
15	5	0.20 (0.07)	0.55 (0.08)	-0.01 (0.03)	0.09 (0.01)
	15	0.15 (0.05)	0.29 (0.05)	-0.01 (0.03)	0.07 (0.01)
	30	0.10 (0.04)	0.20 (0.03)	-0.06 (0.03)	0.06 (0.01)
	50	0.16 (0.05)	0.23 (0.04)	-0.12 (0.03)	0.06 (0.01)
30	5	0.17 (0.06)	0.37 (0.12)	-0.03 (0.02)	0.04 (0.01)
	15	0.13 (0.04)	0.18 (0.03)	-0.005 (0.02)	0.03 (0.01)
	30	0.09 (0.03)	0.09 (0.01)	-0.03 (0.02)	0.03 (0.01)
	50	0.01 (0.03)	0.10 (0.01)	0.02 (0.02)	0.02 (0.01)

Table 5. MEAN BIAS AND MEAN SQUARE ERROR AT $O=25$ AND $UC=62\%$

	<i>M.B(M.S.E) for $\hat{\eta}$</i>			
	5 TGT	15 TGT	30 TGT	50 TGT
5 OBS	0.83(2.20)	0.52(0.97)	0.47(1.00)	0.47(0.76)
15 OBS	0.20(0.55)	0.15(0.29)	0.10(0.20)	0.16(0.22)
30 OBS	0.17(0.37)	0.13(0.18)	0.09(0.08)	0.008(0.10)

Table 6. MEAN BIAS AND MEAN SQUARE ERROR AT $O=25$ AND $UC=62\%$

	<i>M.B(M.S.E) for $\hat{\beta}_0$</i>			
	5 TGT	15 TGT	30 TGT	50 TGT
5 OBS	-0.06(0.25)	-0.08(0.20)	-0.12(0.18)	-0.05(0.13)
15 OBS	-0.01(0.09)	-0.007(0.07)	-0.06(0.06)	-0.02(0.01)
30 OBS	-0.03(0.04)	-0.005(0.03)	-0.03(0.03)	0.02(0.02)

Table 7. MEAN BIAS, MEAN SQUARE ERROR AND STANDARD ERROR AT
O = 40 AND UC = 75%

Number of Observers	Number of Targets	$\hat{\eta}$		$\hat{\beta}_0$	
		M.B (S.E)	M.S.E (S.E)	M.B (S.E)	M.S.E (S.E)
5	5	0.89 (0.12)	2.22 (0.28)	-0.06 (0.05)	0.20 (0.04)
	15	0.55 (0.09)	1.08 (0.16)	-0.06 (0.04)	0.18 (0.03)
	30	0.49 (0.09)	1.03 (0.16)	-0.12 (0.04)	0.18 (0.03)
	50	0.61 (0.12)	1.71 (0.17)	-0.05 (0.04)	0.13 (0.02)
15	5	0.24 (0.08)	0.70 (0.18)	-0.01 (0.03)	0.09 (0.01)
	15	0.17 (0.05)	0.29 (0.05)	-0.01 (0.03)	0.07 (0.01)
	30	0.12 (0.04)	0.20 (0.03)	-0.06 (0.03)	0.06 (0.01)
	50	0.35 (0.04)	0.19 (0.03)	-0.03 (0.03)	0.06 (0.01)
30	5	0.13 (0.05)	0.24 (0.03)	-0.02 (0.02)	0.04 (0.01)
	15	0.20 (0.07)	0.55 (0.03)	-0.01 (0.02)	0.04 (0.01)
	30	0.16 (0.08)	0.68 (0.03)	-0.04 (0.02)	0.03 (0.01)
	50	0.02 (0.03)	0.09 (0.01)	+0.01 (0.02)	0.02 (0.01)

Table 8. MEAN BIAS AND MEAN SQUARE ERROR AT $O=40$ AND $UC=75\%$

	<i>M.B(M.S.E) for $\hat{\eta}$</i>			
	5 TGT	15 TGT	30 TGT	50 TGT
5 OBS	0.89(2.20)	0.55(1.1)	0.49(1.00)	0.61(1.70)
15 OBS	0.24(0.70)	0.17(0.29)	0.12(0.20)	0.35(2.00)
30 OBS	0.13(0.24)	0.20(0.55)	0.16(0.68)	0.02(0.09)

Table 9. MEAN BIAS AND MEAN SQUARE ERROR AT $O=40$ AND $UC=75\%$

	<i>M.B(M.S.E) for $\hat{\beta}_0$</i>			
	5 TGT	15 TGT	30 TGT	50 TGT
5 OBS	-0.06(0.20)	-0.06(0.18)	-0.12(0.18)	-0.05(0.13)
15 OBS	-0.007(0.09)	-0.009(0.07)	-0.06(0.06)	-0.03(0.06)
30 OBS	-0.02(0.04)	-0.01(0.04)	-0.04(0.03)	-0.01(0.02)

III. A WEIBULL REGRESSION MODEL

A. MODEL

Suppose there are M observers. Each observer is presented with N_j targets. Let U_{ij} be the time it takes for observer i to detect target j . Let $x_{i1}, x_{i2}, x_{i3}, \dots, x_{ip}$ be the values of explanatory variables which may effect U_{ij} (e.g., terrain, atmospheric conditions, fatigue etc.)

Assume that U_{ij} are independent random variables with Weibull distributions

$$\begin{aligned} P\{U_{ij} \leq t\} &= 1 - \exp\left[-(te^{-\underline{x}_i \underline{\beta}})^{\xi_{ij}}\right] \\ &= 1 - \exp\left[-(t/\mu_{ij})^{\xi_{ij}}\right] \end{aligned} \quad (3.1)$$

where $\mu_{ij} = e^{\underline{x}_i \underline{\beta}}$ with

$$\underline{x}_i \underline{\beta} = \sum_{k=1}^p x_{ijk} \beta_k \quad (3.2)$$

It follows from equation (3.1) that the distribution of the ln-detection time is

$$\begin{aligned} P\{\ln U_{ij} \leq t\} &= P\{U_{ij} \leq e^t\} \\ &= 1 - \exp\left\{-\left(e^t e^{-\underline{x}_i \underline{\beta}}\right)^{\xi_{ij}}\right\} \\ &= 1 - \exp\left[-\exp\left[(t - \underline{x}_i \underline{\beta})e^{-\xi_{ij}}\right]\right] \end{aligned} \quad (3.3)$$

The derivative of equation (3.3) with respect to t is following

$$\begin{aligned} P\{\ln U_{ij} \in dt\} &= \exp\left[-\exp\left[(t - \underline{x}_i \underline{\beta})e^{-\xi_{ij}}\right]\right] \exp\left\{(t - \underline{x}_i \underline{\beta})e^{-\xi_{ij}}\right\} e^{-\xi_{ij}} \\ &= \exp\left[-\exp\left[(t - \underline{x}_i \underline{\beta})e^{-\xi_{ij}}\right]\right] \exp\left\{(t - \underline{x}_i \underline{\beta})e^{-\xi_{ij}} - \xi_{ij}\right\} \end{aligned} \quad (3.4)$$

When the i^{th} observer is presented with his j^{th} target, he gets a length of time called an opportunity time O_{ij} to detect it. The observer either successfully detects the target within the opportunity time or is unsuccessful. Data for the i^{th} observer consist of times of detections for the successes and the lengths of opportunity times for the failures. For each $i = 1, 2, 3, \dots, M$, $j = 1, 2, 3, \dots, N_j$, let

$$Y_{ij} = \min(\ln U_{ij}, \ln O_{ij}) \quad (3.5)$$

and

$$\Delta_{ij} = \begin{cases} 1 & \text{if } U_{ij} \leq O_{ij} \\ 0 & \text{otherwise} . \end{cases} \quad (3.6)$$

The Y_{ij} are the censored In-detection times and Δ_{ij} is an indicator of whether or not the In-time to detect the j^{th} target by the i^{th} observer is censored. Let

$$C_i = \sum_{j=1}^{N_i} \Delta_{ij} \quad (3.7)$$

be the number of targets detected by observer i . The next section of this chapter will discuss estimation of the parameters $\{\beta_k\}$ [Ref. 4]. These parameter estimates might be used to predict future times to detection for an observer given his past performance. In the last section of the chapter results of a simulation study of the estimation procedure will be given.

B. ESTIMATION

1. The likelihood Equation and Maximum Likelihood Estimators.

The likelihood for the i^{th} observer is

$$L_i(\underline{\beta}, \underline{\xi}) = \prod_{j=1}^{N_i} \exp\{\Delta_{ij}[(y_{ij} - x_{ij}\underline{\beta})e^{-\xi_i} - \xi_i]\} \exp\{-\exp[(y_{ij} - x_{ij}\underline{\beta})e^{-\xi_i}]\} . \quad (3.8)$$

The ln likelihood for the i^{th} observer is

$$\ln L_i(\underline{\beta}, \underline{\xi}) = \sum_{j=1}^{N_i} \{\Delta_{ij}[(y_{ij} - x_{ij}\underline{\beta})e^{-\xi_i} - \xi_i] - \exp[(y_{ij} - x_{ij}\underline{\beta})e^{-\xi_i}]\} . \quad (3.9)$$

Since the observers are assumed to be independent, the ln likelihood for all observers is

$$\begin{aligned}
\ln L(\underline{\beta}, \underline{\xi}) &= \sum_{i=1}^M \ln L_i(\underline{\beta}, \underline{\xi}) \\
&= \sum_{i=1}^M \sum_{j=1}^{N_i} \{ \Delta_{ij} [(y_{ij} - x_{ij}\underline{\beta})e^{-\xi_i} - \xi_i] - \exp[(y_{ij} - x_{ij}\underline{\beta})e^{-\xi_i}] \} .
\end{aligned} \tag{3.10}$$

a. Newton's Procedure to solve for β_k

The partial derivative of equation (3.10) with respect to β_k is

$$\begin{aligned}
\frac{\partial}{\partial \beta_k} \ln L_i(\underline{\beta}, \underline{\xi}) &= \sum_{j=1}^{N_i} \{ \Delta_{ij} [(-x_{ijk})e^{-\xi_i}] - \exp[(y_{ij} - x_{ij}\underline{\beta})e^{-\xi_i}] (-x_{ijk}e^{-\xi_i}) \} \\
&= \sum_{j=1}^{N_i} \{ -\Delta_{ij} + \exp[(y_{ij} - x_{ij}\underline{\beta})e^{-\xi_i}] x_{ijk}e^{-\xi_i} \} .
\end{aligned} \tag{3.11}$$

A Newton procedure to solve equation (3.11) for $\underline{\beta}$ assuming $\{\xi_i\}$ known uses the second derivatives

$$\frac{\partial^2 L}{\partial \beta_k \partial \beta_h} = \sum_{i=1}^M \sum_{j=1}^{N_i} -\exp[(y_{ij} - x_{ij}\underline{\beta})e^{-\xi_i}] x_{ijk}e^{-\xi_i} x_{ijh}e^{-\xi_i} . \tag{3.12}$$

Let

$$w_{ij} = \sqrt{\exp[(y_{ij} - x_{ij}\underline{\beta})e^{-\xi_i}]} ; \tag{3.13}$$

$$u_{ijh} = x_{ijk}e^{-\xi_i} w_{ij} . \tag{3.14}$$

The Newton procedure to solve equations (3.11) for $\underline{\beta}$ can be written as

$$\begin{aligned}
0 = \frac{\partial}{\partial \beta_k} \ln L_i(\underline{\beta}, \underline{\xi}) &= \sum_{i=1}^M \sum_{j=1}^{N_i} \frac{[-\Delta_{ij} + w_{ij}^2]}{w_{ij}} u_{ijk} \\
&\quad - \sum_{i=1}^M \sum_{j=1}^{N_i} \sum_{h=1}^P w_{ij}^2 x_{ijh} e^{-\xi_i} x_{ijk} e^{-\xi_i} [\beta_k - \beta_k^0] \\
&= \sum_{i=1}^M \sum_{j=1}^{N_i} \left\{ \frac{[-\Delta_{ij} + w_{ij}^2]}{w_{ij}} + \sum_{h=1}^P u_{ijh} \beta_h^0 - \sum_{h=1}^P u_{ijh} \beta_h \right\}
\end{aligned} \tag{3.15}$$

which are of the form of the normal equations for Least Squares regression with dependent variables

$$z_{ij} = \frac{[-\Delta_{ij} + w_{ij}^2]}{w_{ij}} + \sum_{h=1}^P u_{ijh} \beta_h^0 \tag{3.16}$$

and independent variables

$$u_{ijk} = w_{ij} x_{ijk} e^{-\xi_i} . \tag{3.17}$$

b. Newton Procedure to solve for ξ_i

The partial derivative of equation (3.10) with respect to ξ_i is

$$\begin{aligned}
\frac{\partial}{\partial \xi_i} \ln L_i(\underline{\beta}, \underline{\xi}_i) &= \sum_{j=1}^{N_i} \left[\{ \Delta_{ij} [(y_{ij} - x_{ij} \underline{\beta}) e^{-\xi_i} (-1) - 1] \} \right. \\
&\quad \left. - \{ \exp[(y_{ij} - x_{ij} \underline{\beta}) e^{-\xi_i}] (y_{ij} - x_{ij} \underline{\beta}) e^{-\xi_i} (-1) \} \right] \\
&= -C_i + \sum_{j=1}^{N_i} (y_{ij} - x_{ij} \underline{\beta}) e^{-\xi_i} \{ -\Delta_{ij} + \exp[(y_{ij} - x_{ij} \underline{\beta}) e^{-\xi_i}] \}
\end{aligned} \tag{3.18}$$

A Newton procedure to solve equation (3.15) for ξ_i , assuming $\underline{\beta}$ known uses the second derivative

$$\begin{aligned}
\frac{\hat{c}^2}{\hat{\partial} \xi_i^2} \ln L_i(\underline{\beta}, \underline{\xi}_i) &= - \sum_{j=1}^{N_i} (y_{ij} - \mathbf{x}_{ij} \underline{\beta}) e^{-\xi_i} \{-\Delta_{ij} + \exp[(y_{ij} - \mathbf{x}_{ij} \underline{\beta}) e^{-\xi_i}]\} \\
&\quad + \sum_{j=1}^{N_i} (y_{ij} - \mathbf{x}_{ij} \underline{\beta}) e^{-\xi_i} \{ \exp[(y_{ij} - \mathbf{x}_{ij} \underline{\beta}) e^{-\xi_i}] (y_{ij} - \mathbf{x}_{ij} \underline{\beta}) e^{-\xi_i} (-1) \} \\
&= -C_i - \sum_{j=1}^{N_i} (y_{ij} - \mathbf{x}_{ij} \underline{\beta})^2 e^{-2\xi_i} \exp[(y_{ij} - \mathbf{x}_{ij} \underline{\beta}) e^{-\xi_i}]
\end{aligned} \tag{3.19}$$

if $\frac{\hat{c}}{\hat{\partial} \xi_i} \ln L_i(\underline{\beta}, \underline{\xi}_i) = 0$. Therefore the Newton equation for ξ_i is

$$\begin{aligned}
0 = \frac{\hat{c}}{\hat{\partial} \xi_i} \ln L_i(\underline{\beta}, \underline{\xi}_i) &= -C_i + \sum_{j=1}^{N_i} \Delta_{ij} (y_{ij} - \mathbf{x}_{ij} \underline{\beta}) e^{-\xi_i} \{-\Delta_{ij} + \exp[(y_{ij} - \mathbf{x}_{ij} \underline{\beta}) e^{-\xi_i}]\} \\
&\quad + \left(-C_i - \sum_{j=1}^{N_i} \{[(y_{ij} - \mathbf{x}_{ij} \underline{\beta}) e^{-\xi_i}]^2 \exp[(y_{ij} - \mathbf{x}_{ij} \underline{\beta}) e^{-\xi_i}]\} \right) [\xi_i - \xi_i^0]
\end{aligned} \tag{3.20}$$

where ξ_i^0 is the current value of ξ_i . Solving results in the equation

$$\begin{aligned}
\xi_i - \xi_i^0 &= \frac{\left\{ +C_i - \sum_{j=1}^{N_i} (y_{ij} - \mathbf{x}_{ij} \underline{\beta}) e^{-\xi_i} [-\Delta_{ij} + \exp\{(y_{ij} - \mathbf{x}_{ij} \underline{\beta}) e^{-\xi_i}\}] \right\}}{-C_i - \sum_{j=1}^{N_i} \{[(y_{ij} - \mathbf{x}_{ij} \underline{\beta}) e^{-\xi_i}]^2 \exp[(y_{ij} - \mathbf{x}_{ij} \underline{\beta}) e^{-\xi_i}]\}}
\end{aligned} \tag{3.21}$$

2. Initial Condition

In this subsection, we describe a rough way to provide initial estimates of $\mathbf{x}_{ij} \underline{\beta}$ to start the iterative Newton procedure of the previous section. It follows from equation (3.1) that

$$\begin{aligned}
P\{(U_{ij})e^{-\xi_i} \leq t\} &= P\{U_{ij} \leq t e^{\xi_i}\} \\
&= 1 - \exp\left[-\left(t e^{\xi_i} e^{-\underline{x}_{ij}\underline{\beta}}\right)e^{-\xi_i}\right] \\
&= 1 - \exp\left[-e^{-\underline{x}_{ij}\underline{\beta}}e^{-\xi_i} t\right]
\end{aligned} \tag{3.22}$$

Hence,

$$\begin{aligned}
E[(U_{ij})e^{-\xi_i}] &= \left(e^{-\underline{x}_{ij}\underline{\beta}}e^{-\xi_i}\right)^{-1} \\
&= e^{\underline{x}_{ij}\underline{\beta}}e^{-\xi_i}
\end{aligned} \tag{3.23}$$

and

$$\ln E[(U_{ij})e^{-\xi_i}] = \underline{x}_{ij}\underline{\beta} - \xi_i \tag{3.24}$$

As a result, for all the observed U_{ij} censored or uncensored, we will put

$$e^{\xi_i} \ln U_{ij} = \underline{x}_{ij}\underline{\beta}^0 \tag{3.25}$$

We will take $\xi_i^0 = 0$ for convenience. Thus the initial value of $\underline{x}_{ij}\underline{\beta}^0$ for each observation u_{ij} is

$$\underline{x}_{ij}\underline{\beta}^0 = \ln u_{ij} \tag{3.26}$$

3. Recursive Procedure to find $\{\beta_k\}$ and ξ_i

1. Put $\xi_i = 0$ for $i = 1, 2, 3, \dots, M$
2. Compute the initial value

$$\underline{x}_{ij}\underline{\beta}^0 = e^{\xi_i} \ln U_{ij} \tag{3.27}$$

3. Iteration

- a. Compute dependent and independent variables for regression

$$W_{ij} = \sqrt{\exp\{(y_{ij} - \underline{x}_{ij}\underline{\beta}^0)e^{-\xi_i}\}} \tag{3.28}$$

$$u_{ijk} = W_{ij} x_{ijk} e^{-\xi_i} \quad (\text{dependent variable})$$

$$z_{ij} = \frac{-\Delta_{ij} + W_{ij}^2}{W_{ij}} + W_{ij}\underline{x}_{ij}\underline{\beta}^0 \quad (\text{independent variable})$$

Let

$$Z = \begin{bmatrix} z_{11} \\ \vdots \\ z_{1N_1} \\ z_{21} \\ \vdots \\ z_{2N_2} \\ \vdots \\ z_{M1} \\ \vdots \\ z_{MN_M} \end{bmatrix} \quad U = \begin{bmatrix} u_{111} & u_{112} & \cdots & u_{11p} \\ u_{121} & u_{122} & \cdots & u_{12p} \\ \vdots & \vdots & \vdots & \vdots \\ \vdots & \vdots & \vdots & \vdots \\ u_{1N_11} & u_{1N_12} & \cdots & u_{1N_1p} \\ u_{211} & u_{212} & \cdots & u_{21p} \\ \vdots & \vdots & \vdots & \vdots \\ \vdots & \vdots & \vdots & \vdots \\ u_{2N_21} & u_{2N_22} & \cdots & u_{2N_2p} \\ \vdots & \vdots & \vdots & \vdots \\ \vdots & \vdots & \vdots & \vdots \\ u_{M11} & u_{M12} & \cdots & u_{M1p} \\ \vdots & \vdots & \vdots & \vdots \\ \vdots & \vdots & \vdots & \vdots \\ u_{MN_M1} & u_{MN_M2} & \cdots & u_{MN_Mp} \end{bmatrix}$$

b. Compute regression estimates

$$\underline{\beta} = (\underline{U}^T \underline{U})^{-1} \underline{U}^T \underline{Z} \quad (3.29)$$

c. Recompute the initial value

$$x_{ij}\underline{\beta} = \sum_{k=1}^p x_{ijk}\beta_k \quad (3.30)$$

d. Update ξ_i as in equation (3.21)

$$\xi_i - \xi_i^0 = \frac{\left\{ + C_i - \sum_{j=1}^{N_i} (y_{ij} - x_{ij}\underline{\beta}) e^{-\xi_i} [-\Delta_{ij} + \exp\{(y_{ij} - x_{ij}\underline{\beta}) e^{-\xi_i}\}] \right\}}{- C_i - \sum_{j=1}^{N_i} \{[(y_{ij} - x_{ij}\underline{\beta}) e^{-\xi_i}]^2 \exp[(y_{ij} - x_{ij}\underline{\beta}) e^{-\xi_i}]\}}$$

e. Put

$$\xi_i^0 = \xi_i$$

$$\beta_k^0 = \beta_k$$

f. Return to step 3

4. Iterate until the following conditions are satisfied


$$\max \left(\frac{|\beta_k - \beta_k^0|}{|\beta_k^0|} \right) \leq 1E-4, \quad k = 1, 2, 3, \dots, p$$

and

$$\max \left(\frac{|\hat{\xi}_i - \xi_i^0|}{|\xi_i^0|} \right) \leq 1E-4, \quad i = 1, 2, 3, \dots, M.$$

4. Implementation in Simulation

The following flow diagram describes one replication of a simulation experiment.



C. SIMULATION PROCEDURE AND RESULTS

1. Simulation

The number of replications for each simulation experiment is 100. Each simulation experiment of 100 replications starts with the same random number seed. The mean bias, its standard error and mean square error, standard error of mean square error from $R = 100$ replications are computed as in equations (2.49a) through (2.50b). In each of the experiments the number of covariates for each observer is 2.

The simulation experiment to study the sampling properties of the estimators of $\{\beta_k\}$ and ξ , is as follows.

1. Give arbitrary constant values for the true values of β_k and ξ . In all the simulations $\beta_1 = 0.2$, $\beta_2 = 0.3$ and $\xi = 0$.
2. The same opportunity time, O_{ij} , $i = 1, 2, 3, \dots, M$, and $j = 1, 2, 3, \dots, N_j$, is used for M observers. The values of the constant opportunity time are 1.3, 2.5, 4.0.
3. Generate random numbers
 - a. Generate independent normal random numbers with mean 1 and variance 0.5, x_{ij1} , $i = 1, 2, 3, \dots, M$, $j = 1, 2, 3, \dots, N_j$.
 - b. Generate independent normal random numbers with mean 2 and variance 1, x_{ij2} , $i = 1, 2, 3, \dots, M$, $j = 1, 2, 3, \dots, N_j$.
 - c. Generate exponential random numbers with mean 1, W_{ij} , $i = 1, 2, 3, \dots, M$, $j = 1, 2, 3, \dots, N_j$.
4. Compute μ_{ij} as follows :

$$\mu_{ij} = \exp(\beta_1 x_{ij1} + \beta_2 x_{ij2}) . \quad (3.31)$$

5. Compute the detection time U_{ij} that it takes for observer i to detect target j as

$$U_{ij} = \mu_{ij} (W_{ij})^{\epsilon_{ij}} . \quad (3.32)$$

6. Compare $\ln U_{ij}$ with $\ln O_{ij}$, then choose smaller one for the data

$$Y_{ij} = \min(\ln U_{ij}, \ln O_{ij})$$

and compute Δ_{ij} as equation (3.7)

7. Compute the initial value as in equation (3.27)
8. Compute the values of the regression variables, z_{ij} and u_{ij} as in equation (3.16) and (3.17)
9. Compute regression estimates as in equation (3.29)
10. Recompute the initial value with new $\hat{\beta}$ value as in equation (3.30)
11. Compute ξ , as in equation (3.21)
12. Iterate the procedure until the differences between successive values of β_1 , β_2 and ξ , are small, (less than 10^{-4}).

2. Results

In this section results from the simulation experiments will be reported. For R replications, the uncensoring level (UC) for an experiment is computed as follows :

$$UC = \frac{\sum_{r=1}^R \sum_{i=1}^M C_i(r)}{\sum_{r=1}^R \sum_{i=1}^M N_i(r)}$$

where $C_i(r)$ is the number of the targets detected by observer i in replication r and $N_i(r)$ is the number of targets presented to observer i in replication r . The UC's for the simulations using the same opportunity time are then averaged to obtain the mean UC for that opportunity time. Figures 19 through 45 show histograms of $(\hat{\beta}_k - \beta_k)$, $k = 1, 2$ and $(\hat{\xi}_i - \xi_i)$. Tables 10 through 21 show mean square errors and mean biases for each of the estimates, their standard errors appear in parenthesis below. Tables 10, 14 and 18 present all the simulation results for different values of opportunity times. The other tables present the same results in a more convenient fashion. The mean biases are displayed with the mean square error in parenthesis; for each table, the opportunity time(O) and average uncensoring(UC) for that are given at the top of table.

- Results for $\hat{\beta}_1$ and $\hat{\beta}_2$

The histograms for $\hat{\beta}_1 - \beta_1$ and $\hat{\beta}_2 - \beta_2$ tend to be somewhat centered around 0. For small numbers of targets and observers the histograms tend to be slightly skewed to the left. Increasing the observation time has little effect on the mean bias and mean square error. Increasing the number of observers for a fixed number of targets tends to decrease the mean square error but has less effect on the mean bias. Increasing the number of targets for a fixed number of observers decreases the mean bias and mean square error. The standard errors of the mean biases mean square errors are large.

- Results for $\hat{\xi}_i$

The histograms of $\hat{\xi}_i - \xi_i$ tend to be centered about 0. There is a tendency for slight skewness to the right for small numbers of observers and targets. Changing the opportunity time has little effect on the mean bias and mean square error. The more targets there are for observer i , the smaller the mean bias and mean square error for $\hat{\xi}_i$. The standard errors of the mean biases and mean square errors are large.

Table 10. MEAN BIAS, MEAN SQUARE ERROR AND STANDARD ERROR AT
 $O = 1.3$ AND $UC = 45\%$

Number of Observers	Number of Targets	$\hat{\beta}_1$		$\hat{\beta}_2$		$\hat{\xi}_i$	
		M.B (S.E)	M.S.E (S.E)	M.B (S.E)	M.S.E (S.E)	M.B (S.E)	M.S.E (S.E)
5	5	-0.06 (0.03)	0.07 (0.01)	-0.02 (0.02)	0.03 (0.01)	-0.05 (0.01)	0.02 (0.01)
	15	-0.02 (0.02)	0.02 (0.01)	-0.02 (0.01)	0.01 (0.001)	-0.02 (0.01)	0.01 (0.001)
	30	-0.003 (0.01)	0.02 (0.01)	-0.02 (0.01)	0.01 (0.002)	-0.03 (0.01)	0.01 (0.001)
	50	-0.01 (0.001)	0.01 (0.001)	-0.01 (0.01)	0.003 (0.001)	-0.01 (0.01)	0.004 (0.001)
15	5	-0.05 (0.01)	0.02 (0.003)	-0.04 (0.01)	0.01 (0.001)	-0.02 (0.01)	0.01 (0.001)
	15	-0.03 (0.01)	0.01 (0.001)	-0.02 (0.01)	0.003 (0.001)	-0.01 (0.01)	0.002 (0.0003)
	30	-0.01 (0.01)	0.01 (0.001)	-0.01 (0.004)	0.002 (0.001)	-0.01 (0.004)	0.002 (0.0002)
	50	-0.01 (0.001)	0.003 (0.0004)	-0.01 (0.003)	0.001 (0.0004)	-0.01 (0.004)	0.001 (0.0002)
30	5	-0.05 (0.01)	0.01 (0.002)	-0.04 (0.01)	0.005 (0.001)	-0.02 (0.005)	0.003 (0.001)
	15	-0.02 (0.01)	0.003 (0.0004)	-0.02 (0.003)	0.001 (0.0002)	-0.01 (0.003)	0.001 (0.0002)
	30	-0.01 (0.01)	0.003 (0.0004)	-0.01 (0.003)	0.001 (0.0001)	-0.004 (0.003)	0.001 (0.0001)
	50	-0.01 (0.004)	0.002 (0.0002)	-0.01 (0.003)	0.001 (0.0001)	-0.01 (0.002)	0.001 (0.0001)

Table 11. MEAN BIAS AND MEAN SQUARE ERROR AT $\theta = 1.3$ AND UC = 45%

	<i>M.B(M.S.E) for β_1</i>			
	5 TGT	15 TGT	30 TGT	50 TGT
5OBS	-0.04(0.06)	-0.02(0.02)	-0.004(0.02)	-0.003(0.01)
15OBS	-0.05(0.02)	-0.02(0.007)	-0.01(0.005)	-0.01(0.003)
30OBS	-0.008(0.01)	-0.05(0.003)	-0.006(0.003)	-0.01(0.002)

Table 12. MEAN BIAS AND MEAN SQUARE ERROR AT $\theta = 1.3$ AND UC = 45%

	<i>M.B(M.S.E) for β_2</i>			
	5 TGT	15 TGT	30 TGT	50 TGT
5 OBS	-0.03(0.02)	-0.02(0.007)	-0.02(0.005)	-0.01(0.002)
15 OBS	-0.04(0.007)	-0.009(0.003)	-0.01(0.002)	-0.01(0.001)
30 OBS	-0.04(0.004)	-0.01(0.001)	-0.01(0.001)	-0.009(0.0006)

Table 13. MEAN BIAS AND MEAN SQUARE ERROR AT $\theta = 1.3$ AND UC = 45%

	<i>M.B(M.S.E) for ξ</i>			
	5 TGT	15 TGT	30 TGT	50 TGT
5 OBS	-0.05(0.14)	-0.02(0.007)	-0.02(0.005)	-0.006(0.004)
15 OBS	-0.01(0.005)	-0.009(0.002)	-0.005(0.002)	-0.008(0.001)
30 OBS	-0.02(0.002)	-0.01(0.001)	-0.004(0.0007)	0.008(0.0006)

Table 14. MEAN BIAS, MEAN SQUARE ERROR AND STANDARD ERROR AT O = 2.5 AND UC = 75%

Number of Observers	Number of Targets	$\hat{\beta}_1$		$\hat{\beta}_2$		$\hat{\xi}_r$	
		M.B (S.E)	M.S.E (S.E)	M.B (S.E)	M.S.E (S.E)	M.B (S.E)	M.S.E (S.E)
5	5	-0.04 (0.02)	0.06 (0.01)	-0.03 (0.01)	0.02 (0.003)	-0.05 (0.01)	0.14 (0.01)
	15	-0.02 (0.02)	0.02 (0.003)	-0.02 (0.01)	0.01 (0.001)	-0.02 (0.01)	0.01 (0.001)
	30	-0.004 (0.01)	0.02 (0.003)	-0.02 (0.01)	0.005 (0.001)	-0.02 (0.01)	0.01 (0.001)
	50	-0.003 (0.01)	0.01 (0.001)	-0.01 (0.01)	0.002 (0.0003)	-0.01 (0.01)	0.004 (0.001)
15	5	-0.05 (0.01)	0.02 (0.003)	-0.04 (0.01)	0.01 (0.001)	-0.01 (0.01)	0.005 (0.001)
	15	-0.02 (0.01)	0.01 (0.001)	-0.02 (0.01)	0.003 (0.0004)	-0.01 (0.01)	0.002 (0.0003)
	30	-0.01 (0.01)	0.01 (0.001)	-0.01 (0.004)	0.002 (0.0002)	-0.01 (0.004)	0.002 (0.0002)
	50	-0.01 (0.01)	0.003 (0.0004)	-0.01 (0.003)	0.001 (0.0002)	-0.01 (0.003)	0.001 (0.0002)
30	5	-0.01 (0.01)	0.01 (0.001)	-0.04 (0.01)	0.004 (0.001)	-0.02 (0.01)	0.003 (0.0004)
	15	-0.05 (0.01)	0.003 (0.0004)	-0.02 (0.003)	0.001 (0.0002)	-0.01 (0.003)	0.001 (0.0002)
	30	-0.01 (0.01)	0.003 (0.0003)	-0.01 (0.003)	0.001 (0.0001)	-0.004 (0.003)	0.001 (0.0001)
	50	-0.01 (0.004)	0.0012 (0.0002)	-0.01 (0.002)	0.001 (0.0001)	-0.01 (0.002)	0.001 (0.0001)

Table 15. MEAN BIAS AND MEAN SQUARE ERROR AT $\theta = 2.5$ AND UC = 75%

	<i>M.B(M.S.E) for $\hat{\beta}_1$</i>			
	5 TGT	15 TGT	30 TGT	50 TGT
5 OBS	-0.06(0.07)	-0.02(0.20)	-0.003(0.02)	-0.005(0.01)
15 OBS	-0.05(0.02)	-0.03(0.007)	-0.01(0.006)	-0.008(0.003)
30 OBS	-0.05(0.01)	-0.02(0.003)	-0.006(0.003)	-0.01(0.002)

Table 16. MEAN BIAS AND MEAN SQUARE ERROR AT $\theta = 2.5$ AND UC = 75%

	<i>M.B(M.S.E) for $\hat{\beta}_2$</i>			
	5 TGT	15 TGT	30 TGT	50 TGT
5 OBS	-0.02(0.03)	-0.02(0.007)	-0.02(0.005)	-0.009(0.003)
15 OBS	-0.04(0.007)	-0.02(0.003)	-0.01(0.002)	-0.01(0.001)
30 OBS	-0.04(0.004)	-0.02(0.001)	-0.01(0.001)	-0.009(0.001)

Table 17. MEAN BIAS AND MEAN SQUARE ERROR AT $\theta = 2.5$ AND UC = 75%

	<i>M.B(M.S.E) for $\hat{\xi}_1$</i>			
	5 TGT	15 TGT	30 TGT	50 TGT
5 OBS	-0.046(0.02)	-0.03(0.008)	-0.03(0.004)	-0.006(0.004)
15 OBS	-0.02(0.01)	-0.01(0.002)	-0.006(0.002)	-0.009(0.001)
30 OBS	-0.02(0.003)	-0.01(0.001)	-0.004(0.0008)	-0.009(0.0006)

Table 18. MEAN BIAS, MEAN SQUARE ERROR AND STANDARD ERROR AT
 $O = 4.0$ AND $UC = 89\%$

Number of Observers	Number of Targets	$\hat{\beta}_1$		$\hat{\beta}_2$		$\hat{\xi}_i$	
		M.B (S.E)	M.S.E (S.E)	M.B (S.E)	M.S.E (S.E)	M.B (S.E)	M.S.E (S.E)
5	5	-0.03 (0.01)	0.02 (0.003)	-0.03 (0.02)	0.03 (0.01)	-0.07 (0.01)	0.02 (0.003)
	15	-0.01 (0.01)	0.01 (0.001)	-0.02 (0.01)	0.01 (0.001)	-0.02 (0.01)	0.01 (0.001)
	30	-0.01 (0.01)	0.01 (0.001)	-0.02 (0.01)	0.01 (0.001)	-0.03 (0.01)	0.004 (0.001)
	50	-0.002 (0.01)	0.01 (0.001)	-0.01 (0.01)	0.004 (0.001)	-0.01 (0.01)	0.003 (0.0004)
15	5	-0.03 (0.01)	0.01 (0.001)	-0.04 (0.01)	0.01 (0.001)	-0.04 (0.01)	0.01 (0.001)
	15	-0.02 (0.01)	0.003 (0.0004)	-0.02 (0.001)	0.004 (0.0001)	-0.02 (0.0004)	0.002 (0.0003)
	30	-0.01 (0.001)	0.002 (0.0003)	-0.01 (0.001)	0.002 (0.0003)	-0.01 (0.004)	0.001 (0.0002)
	50	-0.01 (0.004)	0.002 (0.0002)	-0.01 (0.04)	0.002 (0.0002)	-0.01 (0.003)	0.001 (0.0002)
30	5	-0.03 (0.01)	0.004 (0.001)	-0.04 (0.01)	0.006 (0.001)	-0.04 (0.01)	0.004 (0.001)
	15	-0.01 (0.004)	0.002 (0.0002)	-0.02 (0.004)	0.002 (0.0003)	-0.02 (0.003)	0.001 (0.0002)
	30	-0.03 (0.0003)	0.001 (0.00)	-0.01 (0.0004)	0.001 (0.00)	-0.01 (0.0003)	0.001 (0.00)
	50	-0.01 (0.0003)	0.001 (0.00)	-0.01 (0.003)	0.001 (0.0001)	-0.01 (0.002)	0.001 (0.0001)

Table 19. MEAN BIAS AND MEAN SQUARE ERROR AT $O=4.0$ AND $UC=89\%$

	<i>M.B(M.S.E) for $\hat{\beta}_1$</i>			
	5 TGT	15 TGT	30 TGT	50 TGT
5 OBS	-0.03(0.02)	-0.009(0.008)	-0.01(0.008)	-0.002(0.005)
15 OBS	-0.03(0.01)	-0.02(0.003)	-0.008(0.002)	-0.006(0.002)
30 OBS	-0.03(0.004)	-0.01(0.002)	-0.003(0.001)	-0.008(0.0008)

Table 20. MEAN BIAS AND MEAN SQUARE ERROR AT $O=4.0$ AND $UC=89\%$

	<i>M.B(M.S.E) for $\hat{\beta}_2$</i>			
	5 TGT	15 TGT	30 TGT	50 TGT
5 OBS	-0.03(0.03)	-0.02(0.01)	-0.02(0.008)	-0.008(0.004)
15 OBS	-0.04(0.009)	-0.02(0.004)	-0.01(0.002)	-0.0074(0.002)
30 OBS	-0.04(0.004)	-0.02(0.002)	-0.0097(0.001)	-0.008(0.0007)

Table 21. MEAN BIAS AND MEAN SQUARE ERROR AT $O=4.0$ AND $UC=89\%$

	<i>M.B(M.S.E) for $\hat{\xi}$</i>			
	5 TGT	15 TGT	30 TGT	50 TGT
5 OBS	-0.07(0.02)	-0.02(0.008)	-0.03(0.004)	-0.0005(0.003)
15 OBS	-0.04(0.004)	-0.02(0.002)	-0.01(0.001)	-0.011(0.001)
30 OBS	-0.04(0.004)	-0.02(0.001)	-0.008(0.0006)	-0.011(0.0006)

IV. CONCLUSION

This thesis considers two models for the times until detection of targets. Each model has M observers. The i^{th} observer is presented with N_i targets. In the model of Chapter 2, observer i has a random variable A_i which reflects his ability to detect a target. The random variables $\{A_i\}$ are independent identically distributed having a gamma distribution. Given A_i , the times to target detections for observer i are conditionally independent Weibull random variables with known parameters. Simulation experiments indicate that increasing the number of observers for a fixed number of targets provides the greatest decrease in the mean bias and mean square error of the estimates of the parameters of the gamma distribution that describes the variation between individuals. This is not surprising, since observing more individuals sampled from a fixed population should better estimate properties of that population. The model of Chapter 3 is a Weibull regression model. In this case the simulation experiments indicate that increasing the number of targets for a fixed number of observers provides the greatest decrease in the mean bias and mean square error of the estimators.

A topic for future investigation is to combine the two models and estimation procedures to provide estimates for a hierarchical gamma Weibull regression model. Another topic is to investigate using the fitted hierarchical model to predict future performance of the observers.

V. LIST OF REFERENCES

1. Mackey, D. C., and others, *CASTFOREM - UPDATE (combined arms and support task force evaluation model)*, Department of the Army, US Army TRADOC Analysis Command, White Sand Missile Range, New Mexico 88002 - 5502.
2. Army Regulation 602-2, *Manpower and Personnel Integration(MANPRINT) in the Material Acquisition Process*.
3. Johnson, L. W. and Riess, R. D., *Numerical Analysis*, Addison-Wesley, Inc, 1977.
4. Gaver, D. P., Jacobs, P. A., and O'Muircheartaigh, I., *Hierarchical Models for Target Detection Times*, Naval Postgraduate School Technical Report. To appear.
5. Abramowitz, M. and Stegun, I., *Handbook of Mathematical Functions*, National Bureau of Standards, 7th corrected printing, 1968.
6. Lewis, P. A. W. and Uribe, L., *The New Naval Postgraduate School Random Number Package - LLRANDOM II*, Naval Postgraduate School Technical Report, NPS55-81-005, Monterey, CA, 1981.

APPENDIX A. HISTOGRAMS FOR THE ESTIMATED GAMMA PARAMETERS

5 OBSERVERS (UC : 37 PERCENT)




Figure 1. COMPARISON BETWEEN DIFFERENT ESTIMATES ($O=10$,
UC = 37%): 5,15,30,50 targets for 5 observers ($\hat{\eta} - \eta$)

5 OBSERVERS (UC : 62 PERCENT)






Figure 2. COMPARISON BETWEEN DIFFERENT ESTIMATES ($O=25$, $UC=62\%$): 5,15,30,50 targets for 5 observers ($\hat{\eta} - \eta$)

5 OBSERVERS (UC : 75 PERCENT)


5 TARGETS



15 TARGETS



30 TARGETS



50 TARGETS




Figure 3. COMPARISON BETWEEN DIFFERENT ESTIMATES (O = 40, UC = 75%): 5,15,30,50 targets for 5 observers ($\hat{\eta} - \eta$)

15 OBSERVERS (UC : 37 PERCENT)




Figure 4. COMPARISON BETWEEN DIFFERENT ESTIMATES ($O=10$, UC = 37%): 5,15,30,50 targets for 15 observers ($\hat{\eta} - \eta$)

15 OBSERVERS (UC : 62 PERCENT)




Figure 5. COMPARISON BETWEEN DIFFERENT ESTIMATES ($O=25$, $UC=62\%$): 5,15,30,50 targets for 15 observers ($\hat{\eta} - \eta$)

15 OBSERVERS (UC : 75 PERCENT)






Figure 6. COMPARISON BETWEEN DIFFERENT ESTIMATES ($O=40$, $UC=75\%$): 5,15,30,50 targets for 15 observers ($\hat{\eta} - \eta$)

30 OBSERVERS (UC : 37 PERCENT)


5 TARGETS



15 TARGETS



30 TARGETS



50 TARGETS




Figure 7. COMPARISON BETWEEN DIFFERENT ESTIMATES (O=10, UC=37%): 5,15,30,50 targets for 30 observers ($\hat{\eta} - \eta$)

30 OBSERVERS (UC : 62 PERCENT)






Figure 8. COMPARISON BETWEEN DIFFERENT ESTIMATES ($O=25$, UC = 62%): 5,15,30,50 targets for 30 observers ($\hat{\eta} - \eta$)

30 OBSERVERS (UC : 75 PERCENT)


5 TARGETS



15 TARGETS



30 TARGETS



50 TARGETS




Figure 9. COMPARISON BETWEEN DIFFERENT ESTIMATES (O = 40, UC = 75%): 5,15,30,50 targets for 30 observers ($\hat{\eta} - \eta$)

5 OBSERVERS (UC : 37 PERCENT)




Figure 10. COMPARISON BETWEEN DIFFERENT ESTIMATES (O = 10, UC = 37%): 5,15,30,50 targets for 5 observers ($\hat{\beta}_0 - \beta_0$)

5 OBSERVERS (UC : 62 PERCENT)




Figure 11. COMPARISON BETWEEN DIFFERENT ESTIMATES (O = 25, UC = 62%): 5,15,30,50 targets for 5 observers ($\hat{\beta}_0 - \beta_0$)

5 OBSERVERS (UC : 75 PERCENT)






Figure 12. COMPARISON BETWEEN DIFFERENT ESTIMATES ($O = 40$, UC = 75%): 5,15,30,50 targets for 5 observers ($\hat{\beta}_0 - \beta_0$)

15 OBSERVERS (UC : 37 PERCENT)


5 TARGETS



15 TARGETS



30 TARGETS



50 TARGETS




Figure 13. COMPARISON BETWEEN DIFFERENT ESTIMATES (O = 10, UC = 37%): 5,15,30,50 targets for 15 observers ($\hat{\beta}_0 - \beta_0$)

15 OBSERVERS (UC : 62 PERCENT)




Figure 14. COMPARISON BETWEEN DIFFERENT ESTIMATES (O = 25, UC = 62%): 5,15,30,50 targets for 15 observers ($\hat{\beta}_0 - \beta_0$)

15 OBSERVERS (UC : 75 PERCENT)




Figure 15. COMPARISON BETWEEN DIFFERENT ESTIMATES ($O = 40$, UC = 75%): 5,15,30,50 targets for 15 observers ($\hat{\beta}_0 - \beta_0$)

30 OBSERVERS (UC : 37 PERCENT)




Figure 16. COMPARISON BETWEEN DIFFERENT ESTIMATES ($\sigma = 10$, UC = 37%): 5,15,30,50 targets for 30 observers ($\hat{\beta}_0 - \beta_0$)

30 OBSERVERS (UC : 62 PERCENT)




Figure 17. COMPARISON BETWEEN DIFFERENT ESTIMATES ($O = 25$, $UC = 62\%$): 5,15,30,50 targets for 30 observers ($\hat{\beta}_0 - \beta_0$)

30 OBSERVERS (UC : 75 PERCENT)





Figure 18. COMPARISON BETWEEN DIFFERENT ESTIMATES (O = 40, UC = 75%); 5,15,30,50 targets for 30 observers ($\hat{\beta}_0 - \beta_0$)


APPENDIX B. HISTOGRAM FOR THE WEIBULL REGRESSION PARAMETERS

5 OBSERVERS (UC : 45 PERCENT)


5 TARGETS



15 TARGETS



30 TARGETS



50 TARGETS




Figure 19. COMPARISON BETWEEN DIFFERENT ESTIMATES ($O = 1.3$,
UC = 45%): 5, 15, 30, 50 targets for 5 observers ($\hat{\beta}_1 - \beta_1$)

5 OBSERVERS (UC : 75 PERCENT)






Figure 20. COMPARISON BETWEEN DIFFERENT ESTIMATES ($\sigma = 2.5$, UC = 75%): 5, 15, 30, 50 targets for 5 observers ($\hat{\beta}_1 - \beta_1$)

5 OBSERVERS (UC : 89 PERCENT)


5 TARGETS



15 TARGETS



30 TARGETS



50 TARGETS




Figure 21. COMPARISON BETWEEN DIFFERENT ESTIMATES ($O=4.0$,
 $UC=89\%$): 5, 15, 30, 50 targets for 5 observers ($\hat{\beta}_1 - \beta_1$)

15 OBSERVERS (UC : 45 PERCENT)




Figure 22. COMPARISON BETWEEN DIFFERENT ESTIMATES ($O = 1.3$, UC = 45%): 5, 15, 30, 50 targets for 15 observers ($\hat{\beta}_1 - \beta_1$)

15 OBSERVERS (UC : 75 PERCENT)




Figure 23. COMPARISON BETWEEN DIFFERENT ESTIMATES ($O = 2.5$, UC = 75%): 5, 15, 30, 50 targets for 15 observers ($\hat{\beta}_1 - \beta_1$)

15 OBSERVERS (UC : 89 PERCENT)





Figure 24. COMPARISON BETWEEN DIFFERENT ESTIMATES ($\sigma = 4.0$, UC = 89%): 5, 15, 30, 50 targets for 15 observers ($\hat{\beta}_1 - \beta_1$)

30 OBSERVERS (UC : 45 PERCENT)

5 TARGETS



15 TARGETS






Figure 25. COMPARISON BETWEEN DIFFERENT ESTIMATES ($O = 1.3$, $UC = 45\%$): 5, 15, 30, 50 targets for 30 observers ($\hat{\beta}_1 - \beta_1$)

30 OBSERVERS (UC : 75 PERCENT)






Figure 26. COMPARISON BETWEEN DIFFERENT ESTIMATES ($O = 2.3$,
 UC = 75%): 5, 15, 30, 50 targets for 30 observers ($\hat{\beta}_1 - \beta_1$)

30 OBSERVERS (UC : 89 PERCENT)


5 TARGETS



15 TARGETS



30 TARGETS



50 TARGETS




Figure 27. COMPARISON BETWEEN DIFFERENT ESTIMATES ($\sigma = 4.0$,
UC = 89%): 5, 15, 30, 50 targets for 30 observers ($\hat{\beta}_1 - \beta_1$)

5 OBSERVERS (UC : 45 PERCENT)




Figure 28. COMPARISON BETWEEN DIFFERENT ESTIMATES ($O = 1.3$, UC = 45%): 5, 15, 30, 50 targets for 5 observers ($\hat{\beta}_2 - \beta_2$)

5 OBSERVERS (UC : 75 PERCENT)




Figure 29. COMPARISON BETWEEN DIFFERENT ESTIMATES ($\sigma = 2.5$, UC = 75%): 5, 15, 30, 50 targets for 5 observers ($\hat{\beta}_2 - \beta_2$)

5 OBSERVERS (UC : 89 PERCENT)





Figure 30. COMPARISON BETWEEN DIFFERENT ESTIMATES (O = 4.0, UC = 89%): 5, 15, 30, 50 targets for 5 observers ($\hat{\beta}_2 - \beta_2$)


15 OBSERVERS (UC : 45 PERCENT)

5 TARGETS




BETA2 - BETA2 TRUE

30 TARGETS




BETA2 - BETA2 TRUE

15 TARGETS



BETA2 - BETA2 TRUE

50 TARGETS



BETA2 - BETA2 TRUE

Figure 31. COMPARISON BETWEEN DIFFERENT ESTIMATES ($O = 1.3$,
 UC = 45%): 5, 15, 30, 50 targets for 15 observers ($\hat{\beta}_2 - \beta_2$)

15 OBSERVERS (UC : 75 PERCENT)




Figure 32. COMPARISON BETWEEN DIFFERENT ESTIMATES ($\sigma = 2.5$, UC = 75%); 5, 15, 30, 50 targets for 15 observers ($\hat{\beta}_2 - \beta_2$)

15 OBSERVERS (UC : 89 PERCENT)




Figure 33. COMPARISON BETWEEN DIFFERENT ESTIMATES ($O = 4.0$, $UC = 89\%$): 5, 15, 30, 50 targets for 15 observers ($\hat{\beta}_2 - \beta_2$)

30 OBSERVERS (UC : 45 PERCENT)




Figure 34. COMPARISON BETWEEN DIFFERENT ESTIMATES ($O = 1.3$, UC = 45%): 5, 15, 30, 50 targets for 30 observers ($\hat{\beta}_2 - \beta_2$)

30 OBSERVERS (UC : 75 PERCENT)




Figure 35. COMPARISON BETWEEN DIFFERENT ESTIMATES ($O = 2.5$,
 UC = 75%): 5, 15, 30, 50 targets for 30 observers ($\hat{\beta}_2 - \beta_2$)

30 OBSERVERS (UC : 89 PERCENT)




Figure 36. COMPARISON BETWEEN DIFFERENT ESTIMATES ($O = 4.0$, $UC = 89\%$): 5, 15, 30, 50 targets for 30 observers ($\hat{\beta}_2 - \beta_2$)

5 OBSERVERS (UC : 45 PERCENT)




Figure 37. COMPARISON BETWEEN DIFFERENT ESTIMATES ($O = 1.3$, UC = 45%): 5, 15, 30, 50 targets for 5 observers ($\hat{\xi}_i - \xi_i$)

5 OBSERVERS (UC : 75 PERCENT)




Figure 38. COMPARISON BETWEEN DIFFERENT ESTIMATES ($\sigma = 2.5$, UC = 75%): 5, 15, 30, 50 targets for 5 observers ($\hat{\xi}_i - \xi_i$)

5 OBSERVERS (UC : 89 PERCENT)




Figure 39. COMPARISON BETWEEN DIFFERENT ESTIMATES ($O = 4.0$, UC = 89%): 5, 15, 30, 50 targets for 5 observers ($\hat{\xi}_i - \xi_i$)

15 OBSERVERS (UC : 45 PERCENT)




Figure 40. COMPARISON BETWEEN DIFFERENT ESTIMATES ($O = 1.3$, $UC = 45\%$): 5, 15, 30, 50 targets for 15 observers ($\hat{\xi}_i - \xi_i$)

15 OBSERVERS (UC : 75 PERCENT)




Figure 41. COMPARISON BETWEEN DIFFERENT ESTIMATES ($O = 2.5$, UC = 75%): 5, 15, 30, 50 targets for 15 observers ($\hat{\xi}_i - \xi_i$)

15 OBSERVERS (UC : 89 PERCENT)




Figure 42. COMPARISON BETWEEN DIFFERENT ESTIMATES ($O=4.0$, UC=89%): 5, 15, 30, 50 targets for 15 observers ($\hat{\xi}_i - \xi_i$)

30 OBSERVERS (UC : 45 PERCENT)




Figure 43. COMPARISON BETWEEN DIFFERENT ESTIMATES ($O = 1.3$, $UC = 45\%$): 5, 15, 30, 50 targets for 30 observers ($\hat{\xi}_i - \xi_i$)

30 OBSERVERS (UC : 75 PERCENT)




Figure 44. COMPARISON BETWEEN DIFFERENT ESTIMATES ($O = 2.3$, $UC = 75\%$): 5, 15, 30, 50 targets for 30 observers ($\hat{\xi}_i - \xi_i$)

30 OBSERVERS (UC : 89 PERCENT)




Figure 45. COMPARISON BETWEEN DIFFERENT ESTIMATES ($O = 4.0$, $UC = 89\%$): 5, 15, 30, 50 targets for 30 observers ($\hat{\xi}_i - \xi_i$)

**APPENDIX C. SIMULATION PROGRAM FOR THE ESTIMATION
OF GAMMA PARAMETERS**

```

VBAE[ ]∇
∇ R←M BAE N;MU;TA;H;E;L;OBS;W;G1;G;Y;C;DEL;S;
M1;M2;NIN;L;V;F;A;B;IN;JN;C1
[1] MU←4.2
[2] TA←0
[3] IN←NIN+JN←2ρ0
[4] OBS←40
[5] W←(M,N)ρ(M×N) EXPRAND 1
[6] G1←Φ(N,M)ρG←M GAMRAND 1.5 0.154
[7] Y←ΦMU×(W+G1)×(TA)
[8] C1←+/C←+/DEL←Y≤OBS
[9] Y←Y|@OBS
[10] S←+/S1←*H←(Y-ΦMU)×*-1×TA
[11] M1←(+/+/DEL×H)÷(L←+/C)
[12] M2←(+/+/DEL×(H*2))÷L
[13] V←(M2-(M1*2))-((01)*2)÷6
[14] IN[2]←(*-1×M1)-0.5772
[15] →A2×1(V>0)
[16] V←1
[17] A2:IN[1]←(*-1×V)-IN[2]
[18] →L7
[19] L6:IN←NIN
[20] L7:NIN[2]←IN[2]+(*-1×+/((A-C)+B)÷(+(A←(*IN[2])×S)+
(B←S+*IN[1]))
[21] JN←IN[1],NIN[2]
[22] F←C SUM JN
[23] NIN[1]←IN[1]+(*-1×+/((IN[1]-ΦB))+F[;1])÷
(+(1-*IN[1]÷B)-F[;2])
[24] →L6×1((|IN[1]-NIN[1])÷IN[1])>0.0001)^
((|IN[2]-NIN[2])÷IN[2])>0.0001)
[25] □←NIN
[26] R←(NIN+(*-1.872 1.4665)),(C1+M×N)
∇
∇SUM[ ]∇
∇ R←C SUM IN;DD;BB;AA;D1;D2;I
[1] DD←10
[2] I←1
[3] L3:→L4×1(C[I]=0)
[4] D1←+/+(AA←*IN[1]+IN[2])+BB←(1C[I])-1
[5] D2←+/AA÷(AA+BB)*2
[6] →L5
[7] L4:D1+D2←0
[8] L5:DD←DD,D1,D2
[9] →L3×1(ρC)≥I←I+1
[10] R←DD←((ρC),2)ρDD

```

```

∇
∇THESIS[ ]
∇ R←K THESIS J;I;RES;M;N
[1] □RL←466801747
[2] M←1↑J
[3] N←1↑J
[4] I←1
[5]
[6] RES←(K,3)ρ0
[7] L0:RES[I;]←M BAE N
[8] →L0×1K≥I←I+1
[9] R←RES
∇

```

APPENDIX D. SIMULATION PROGRAM FOR THE ESTIMATION
OF REGRESSION PARAMETERS

```

VBAE2[ ]V
V R←M BAE2 N;OBS;BO;BN;W;U1;M1;M2;MU;ZZ;RR;T0;T1;S;
Y;C;IV;W1;X1;X2;DEL;T;A;T2;C1
OBS←4.0
[1]
[2] J←1
[3] T1←T2+Q(N,M)P T←T0+MPO
[4] BO←BN←2PO
[5] X1←(M,N)P(M×N) NORRAND 1 0.5
[6] X2←(M,N)P(M×N) NORRAND 2 1
[7] W←(M,N)P(M×N) EXPRAND 1
[8] U1←*((0.2×X1)+0.3×X2)
[9] Y←@U1×W*(T2)
[10] DEL←Y≤@OBS
[11] C1←+/C←+/DEL
[12] Y←Y\@OBS
[13] IV←(T2)×Y
[14] LL:BO←BN
[15] T0←T
[16] T2←T1
[17] W1←*((Y-IV)×*(-1×T2))*0.5
[18] M1←W1×X1×*(-1×T2)
[19] M2←W1×X2×*(-1×T2)
[20] MU←Q(2,(M×N))P(,M1),(,M2)
[21] Z←Q,Z←(W1×IV)+( (-1×DEL)+W1*2)÷W1
[22] BN←(E(QMU)+.×MU)+.×((QMU)+.×Z)
[23] IV←(X1×BN[1])+(X2×BN[2])
[24] RR←(Y-IV)×*(-1×T2)
[25] T1←Q(N,M)PT1←T←,T←T+(C-+/RR×((-1×DEL)+*RR))+
((-1×C)-+/RR*2)×RR)
[26] A←(Γ/(BN-BO)+BN)>0.0001
[27] S←|(T0-T)÷T
[28] →LE×1300≤J←J+1
[29] →LL×1(AV(Γ/S)>0.0001)
[30] LE:R←(BN←BN- 0.2 0.3),(C1+(M×N)),((+/T)+M)
V
VTHESIS[ ]
V R←K THESIS J;I;RES;M;N
[1] QRL←466801747
[2] M←1↑J
[3] N←-1↑J
[4] I←1
[5] RES←(K,4)PO
[6] L0:RES[I;]←M BAE2 N
[7] →L0×1K≥I←I+1
[8] R←RES

```



```

▽
▽STAT[ ]
▽ R←STAT K;DF;ANS;II;JJ;KK;TR;MS
[1] II←JJ+KK+1
[2]
[3] DF← 5 10 20 30 40 50
[4] TR← 10 20 50
[5] ANS←(((ρDF)×(ρTR)),K,4)ρ0
[6] MSE←(((ρDF)×(ρTR)),7)ρ0
[7] LO:ANS[II;;]←MS+K THESIS(DF[JJ],TR[KK])
[8] MSE[II;]←((+MS)+K),((+MS[ 1 2 4]*2)+K)
[9] II←II+1
[10] →LO×1 (ρTR)≥KK+KK+1
[11] →LO×1 (ρDF)≥JJ+JJ+KK+1
[12] R←ANS
▽

```

INITIAL DISTRIBUTION LIST

	No. Copies
1. Defense Technical Information Center Cameron Station Alexandria, VA 22304-6145	2
2. Library, Code 0142 Naval Postgraduate School Monterey, CA 93943-5002	2
3. Professor P. A. Jacobs, Code 55JC Department of Operations Research Naval Postgraduate School Monterey, California 93943-5000	1
4. Professor D. P. Gaver, Code 55GV Department of Operations Research Naval Postgraduate School Monterey, California 93943-5000	1
5. Professor Peter. Purdue, Code 55PD Department of Operations Research Naval Postgraduate School Monterey, California 93943-5000	1
6. Professor Samuel Parry, Code 55PY Department of Operations Research Naval Postgraduate School Monterey, California 93943-5000	1
7. Lt. Col. Bard Mansager TRADOC Monterey, California 93945	1
8. Professor Kang, Keebom, Code 55KA Department of Operations Research Naval Postgraduate School Monterey, California 93943-5000	1
9. Personnel Management Office Air Force Headquarters Chungcheon Nam Do Republic of Korea	1
10. Air Force Central Libray Chongwon Gun, Chungcheong Bug Do Republic of Korea	2
11. 3rd Department of Air Force College Chongwon Gun, Chungcheong Bug Do Republic of Korea	1

- | | | |
|-----|---|----|
| 12. | Library of Air Force Academy
Chongwon Gun, Chungcheong Bug Do
Republic of Korea | 2 |
| 13. | Bae, Deok Hwan
Postal Code 180-11
Anjung Ri 153
Pang Seong-Moeyn, Pyeong Tack-kun, Kyeong ki-Do
Republic of Korea | 10 |
| 14. | Choi, Moon Soo
SMC 1840
Naval Postgraduate School
Monterey, California 93943 | 1 |

614-583

Thesis

B1251 Bae

c.1 Models for target de-
tection times.

22 FEB 93

00504

Thesis

B1251 Bae

c.1 Models for target de-
tection times.

thesB1251

Models for target detection times.



3 2768 000 86366 6

DUDLEY KNOX LIBRARY