

Historic, archived document

Do not assume content reflects current scientific knowledge, policies, or practices.

United States Department of Agriculture
Bureau of Biological Survey

Wildlife Research and Management Leaflet BS-32

Washington, D. C.

*

January 1936

RAISING BADGERS IN CAPTIVITY

Prepared in Section of Fur Resources, Division of Wildlife Research

The active demand for serviceable pelts of all kinds and the good prices received for badger skins have resulted in many attempts to raise badgers in captivity. Badger raising, however, is still in its infancy, and little as yet can be predicted as to its future. Fur farmers engaged in the experiments say that the animals tolerate pen conditions very well.

General Description

The badger is one of the larger members of the weasel family. It has a broad, heavy body and a short, muscular neck. Its head is rather small and pointed, and the ears are small and rounded. The forelegs are short, powerful, and well developed, with claws more than an inch in length and well adapted to burrowing, while the hind legs are small and the claws short.

The fur on the back varies from light to dark silvery buff-gray. This blends to a yellowish white on the under side of the body. The head and neck are marked with a white stripe on top, and white patches appear also below the eyes and on the ears. The hair is longest on the sides of the body. Primeness of the pelt varies somewhat, but it is usually in best condition from the latter part of December to the middle of March.

Pens and Dens

A great variety of pens and dens can be used in raising badgers in captivity. While no definite plan of construction has been devised, there are certain fundamental rules that should be observed. Sufficient space should be given for exercise, and the enclosure should be so made as to prevent escape of the animals. The floor should be either of concrete or of earth covered with heavy woven wire, such as that used for the sides, so as to prevent the animals from burrowing out. The den should be well ventilated, and may be either above or below the surface of the ground. The pens should be on well-drained soil, with both shade and sunlight.

Feeding

Badgers in the wild subsist largely on ground squirrels and other small animals. In captivity their food may consist of a mixture of cereals, milk, eggs, and meat. The definite quantity of food required

has not been determined, but it is usually safe to give as much as the animals will readily clean up. The ration used on most fox farms appears to be suitable. In cold climates badgers usually require more food prior to hibernation than at other times. The practice of feeding once daily is followed on some fur farms. On others, two feedings a day are given with good results.

Breeding

There is a conflict of testimony as to the time of breeding. Some have asserted that it is in the fall season, others that it occurs during March and April. The breeding season and the length of the gestation period will remain in doubt until further series of experiments and observations have been made. It is known, however, that only one litter a year is produced, usually in May or June. The number in the litter may vary from one to seven, though it probably averages about four.