
Historic, archived document

Do not assume content reflects current scientific knowledge, policies, or practices.



SEVERAL BREEDS of cattle in the United States are recognized as dairy breeds. Although much alike in what is known as general dairy conformation, these breeds differ to some extent in certain characteristics. What these characteristics are, the factors to consider in selecting a breed, and the history of the origin and development of the breeds are questions of interest to both the beginner and the established breeder of dairy cattle. These are the topics discussed in this bulletin.

This bulletin supersedes Farmers' Bulletin 893, Breeds of Dairy Cattle.

Washington, D.C.

Issued February, 1925; revised May, 1927.

n.

DAIRY CATTLE BREEDS

By AMER B. NYSTROM, Dairy Husbandman, Bureau of Dairy Industry

CONTENTS

Pa	ge		Page
Dairy cattle in the United States What is a dairy breed? Registration Which breed to select The score card	$3 \\ 4 \\ 6$	Dutch Belted Guernsey Holstein-Friesian Jersey	- 14 - 27 - 22 - 26

DAIRY CATTLE IN THE UNITED STATES

A CCORDING to estimates made by the United States Department of Agriculture there were about 33,000,000 dairy cattle of all ages in the United States on January 1, 1926. Three per cent of these cattle, or about 1,000,000, are purebred, and represent six breeds, namely, Ayrshire, Brown Swiss, Dutch Belted, Guernsey, Holstein-Friesian, and Jersey. The improvement that must be made in the 97 per cent that are not purebred must come largely from the 3 per cent that are purebred. Likewise, any increase in the number of our purebred dairy cows, and even the maintenance of our grade dairy herds at their present number and efficiency, will be accomplished principally through the use of purebred bulls. For these reasons purebred dairy cattle have played in the past and will play in the future a very important rôle in the dairy industry of the Nation.

NUMBER AND DISTRIBUTION OF BREEDS

Tables 1 and 2 show the number and distribution of the various breeds in the United States, by sections and by States. The information in Table 1, showing total number of each breed on January 1, 1920, was brought out by an inquiry sent to 14,000 special livestock reporters of the branch then known as the Bureau of Markets and Crop Estimates of the United States Department of Agriculture. Grades and scrubs were listed with the respective breeds to which they seemed to belong.

Table 2 shows the number of registered purebreds of the dairy breeds on January 1, 1920, as ascertained by the census.

Table 3 gives the average annual production of milk and butterfat of the cows having official yearly records in the breed associations.

TABLE 1.—Estimated	number and perc	centage of cattle of	dairy breeds, including
purebreds and gra	des, in the Unite	d States, January	1, 1920, by sections
-	,	, 0	, , ,

Breed	Total	United States	North Atlan- tic States	North Cen- tral, East	North Cen- tral, West	South Atlan- tic States	South Cen- tral States	Far West
A yrshire Brown Swiss Dutch Belted Guernsey Holstein-Friesian Jersey Total	Number 412,000 170,000 1,57,000 1,993,000 11,069,000 9,554,000 23,355,000	$\begin{array}{c} Per \ cent \\ 1.8 \\ .7 \\ .7 \\ 8.5 \\ 47.4 \\ 40.9 \\ \hline 100.0 \end{array}$	$Per \ cent \\ 5.8 \\ .5 \\ 10.6 \\ 65.3 \\ 17.3 \\ \hline 100.0$	Per cent 0.8 1.3 .2 12.1 56.2 29.4 100.0	$Per \ cent \\ 1. 9 \\ 1. 3 \\ . 6 \\ 11. 9 \\ 54. 3 \\ 30. 0 \\ 100. 0$	Per cent 0.7 .3 .3 8.4 21.1 69.2 100.0	Per cent 2.3 1.3 17.3 79.1 100.0	Per cent 0.8 .4 5.1 58.9 34.8 100.0

1

TABLE 2.—Purebred	(registered)	cattle of	the ?	dairy	breeds	on	farms	in	1920,	by
S	states and sec	ctions, as	shoi	wn by	the cens	us				,

Division and State	Total	Ayr- shire	Brown Swiss	Guern- sey	Holstein- Friesian	Jersey	All other breeds ¹
United States	916, 602	30, 509	8, 283	79, 446	528, 621	231, 834	37, 909
Geographic divisions:							
New England Middle Atlantic	85, 724 239, 764	9,780 12,883	349 1, 904	10, 311 21, 114	42, 721 171, 124	20,305 25,815	2, 258 6, 924
East North Central	289,859	12, 883 3, 735	4, 199	29.640	1 185.475	25, 815 57, 167	9,643
West North Central	106, 967	· 1,859	1, 386	8,250	62,055	23,697	9,720
South Atlantic East South Central	49, 119 34, 651	519 27	43	5, 949 298	15, 445 5, 902	25,245 27,024	1, 918 1, 397
West South Central	43, 268	60	13	263	9,724	30,650	2, 558
Mountain Pacific	22,534	324	146	838	12,689	6, 926 15, 005	1,611
Pacific	44, 716	1,322	240	2, 783	23, 486	15,005	1,880
New England:							
Maine New Hampshire	15, 683	1, 134	62	1, 836	7, 206	4, 999	446
New Hampshire	10, 750	1, 214	75 59	1, 151 2, 193	6,695	1,348	267 630
Vermont Massachusetts Rhode Island	28, 549 18, 807	3, 808 1, 880	80	3, 348	13, 413 10, 006	8, 446 2, 904	589
Rhode Island	1,651	494		217	542	351	47
Connecticut	10, 284	1, 250	73	1, 566	4,859	2, 257	279
Middle Atlantic: New York	153, 037	9, 521	1, 347	9, 749	114,662	13, 411	4, 347
New Jersey	11, 538	265	23	1, 747	7,810	1, 368	325
New Jersey Pennsylvania East North Central:	75, 189	3, 097	534	9, 618	7, 810 48, 652	1, 368 11, 036	2, 252
East North Central:	TO 000	1 001	324	4, 960	00 007		. 0.400
Ohio Indiana	70, 882 21, 115	1, 021 509	324 131	4,900	38, 327 8, 477	23, 842 9, 921	2,408 862
Illinois	36, 412	202	1.385	1,369	25, 124	7, 317	1,015
Illinois Michigan	46,533	291	429	3,369 18,727	32,702	8.296	1,446
Wisconsin West North Central:	114, 917	1, 712	1,930	18,727	80, 845	7, 791	3, 912
Minnesota	32, 668	399	483	4,468	22, 830	2, 508	1, 980
Minnesota Iowa	20,286	271	447	1,716	10.916	3, 629	3, 307
Miscouri North Dakota South Dakota	19, 037	110	135	760	5, 569	10, 708	1,755
North Dakota	4, 797 5, 248	226 85	23 119	346 135	2, 937 4, 027	481 312	784 570
Nebraska	7,873	74	38	348	5,368	1,275	770
Nebraska Kansas	17, 058	694	141	477	10,408	4, 784	554
outh Atlantic:	1,691	. 1		246	1, 245	172	27
Delaware Maryland	1, 691	113	9	1, 867	4,073	2,323	283
District of Columbia	186			. 1	175	10	
Virginia	9, 586	$25 \\ 272$		1,696	4,160	3, 223	482
West Virginia	4, 450 7, 697	272 44	$32 \\ 1$	333 789	1, 134 1, 613	2, 546 4, 978	133 272
North Carolina	5, 184	4		644	1,008	3, 389	139
Georgia Florida	8,727	14	1	305	1,700	6, 224 2, 380	483
East South Central:	2, 930	46		68	337	2, 380	99
Kentucky	8, 829	20		40	2,046	6,421	302
Kentucky Tennessee	11, 347	2	1	111	1,383	9,424	426
Alabama	6,108	4	2	46	1,142	4,608	306
Mississippi Vest South Central:	8, 367	1		101	1, 331	6, 571	363
Arkansas	6,950	3	1	19	2,001	4,627	299
Louisiana	3, 415	1	2	94	1,009	2,201	108
Oklahoma	9, 539	37	3 7	91 59	3, 741 2, 973	5,104	563
Texas Aountain:	23, 364	19	•	. 59	2, 813	18, 718	1, 588
Montana	3, 451	13	54	176	2,453	560	195
Idaho	4, 138	46	48	197	2, 049	1, 579	219
Idaho Wyoming Colorado New Mexico	1,071	21 114	14 30	31 241	747	$\begin{array}{c}167\\1,605\end{array}$	91 401
New Mexico	6,448 1,327	114	30	₹41 42.	4, 057 438	1,005	401
Arizona	2,772	109		48	1, 778	669	168
Utah	2, 772 2, 922			93	970	1,706	153
Nevada	405	21		10	197	133	44
Pacific: Washington	12, 720	404	69	941	7.673	3, 402	231
Oregon	12,852	323	135	697	3,624	7, 771	302
California	19, 144	595	36	1,145	12, 189	3, 832	1, 347

¹ Including animals reported as purebred, but with breed not specified.

	Number of cows	To what date		Milk	Butterfat		
Breed	and heifers			MIIK	Quantity	Test	
A yshire Brown Swiss Dutch Belted Guernsey Holstein-Friesian Jersey	6, 368 476 46 20, 931 23, 622 22, 355	Jan. Jan. Jan. Jan. Jan. Jan.	1, 1925 1, 1926 1, 1926 1, 1926 1, 1926 1, 1925	Pounds 10, 256 12, 534 10, 141 9, 615 15, 697 8, 287	Pounds 407.48 501.60 384.64 480.10 532.44 444.37	Per cent 3. 97 4. 00 3. 80 4. 99 3. 39 5. 36	

 TABLE 3.—Average yearly production of milk and butterfat of the cows of different

 breeds that have official yearly records

WHAT IS A DAIRY BREED?

The term "dairy breed" has been accepted by stockmen and investigators as referring to the breeds of cattle that are especially well fitted for the production of milk and butterfat. Such breeds represent the efforts made by breeders of many generations toward improving the milking capacity of certain classes of cows. Because of this fact the inherent tendency of purebred dairy cattle to produce milk is greater than that of a native or unimproved cow. This inherent capacity is transmitted to the offspring. As a result, the mating of a purebred dairy animal with a native or scrub produces a grade animal which is superior to the scrub in production and in other dairy characteristics.

A purebred dairy animal is one that meets the requirements for registration laid down by the association for that breed in the United States. A grade is the offspring resulting from mating a purebred with a scrub, or from mating animals not purebred but having close purebred ancestors. The offspring of a purebred and a grade is also a grade, and through progressive improvement such animals become high grade. The names of the breeds (Ayrshire, Brown Swiss, etc.) may refer to either purebreds or grades; but to prevent misunderstanding it is desirable to precede the breed name with the word "purebred" or "grade."

In addition to the breeds of dairy cattle mentioned, certain other breeds having good milking qualities are kept for dairy purposes. Such cattle, which are often referred to as dual-purpose animals because of their ability to produce satisfactory carcasses as well as a good milk flow, include the Shorthorn, Red Polled, and Devon. The qualities of these are discussed in Farmers' Bulletin 612, "Breeds of Beef Cattle."

REGISTRATION

A purebred dairy animal is one whose sire and dam are eligible to be recorded by name and number in a register of the breed, commonly called the herdbook. An animal thus qualified may itself be recorded in the same herd register, provided the sire and dam are registered, and provided it also qualifies with regard to color. Additional rules and requirements for registration are laid down by the various breed organizations. Copies of these rules may be obtained by writing to the association concerned, as listed on page 30 of this bulletin. In addition to the herd register, there is for each breed another register in which are entered the names of purebred cows that have completed records meeting specified requirements of milk and butterfat production under definite regulations. Bulls that have a certain number of tested daughters are also recorded in this register. This record of tested cows and proved bulls is called by various names— Advanced Registry for the Ayrshires and Dutch Belted, Register of Production for the Brown Swiss, Advanced Register for the Guernseys and Holsteins, and Register of Merit for the Jerseys.

The requirements for admission to this special register of production and the rules under which the records are made vary somewhat with different breeds. Detailed information on this point may be obtained from the breed associations concerned.

WHICH BREED TO SELECT

Sometimes too much emphasis is given to the question which breed to choose, and too little to the matter of getting good individuals that is, those that are well bred and high producers. There are three points, however, that should be considered in deciding which breed to select. These are: (1) The breed that predominates in the locality where the new herd is to be located, (2) personal preference, and (3) market requirements for the product.

THE BREED THAT PREDOMINATES

A dairyman just starting with purebreds may feel that since all his neighbors have one breed of cattle, he should get another breed so as to have a monopoly in the business of selling breeding stock. There is no question about the monopoly, but there would probably be no business to monopolize. It is difficult for an isolated small breeder to dispose of his surplus stock to advantage, while if there are many breeders with the same breed buyers are attracted to the locality because of the better chance to get the desired animals from one or more of the several breeders.

There are other advantages to a dairyman in having the same breed as his neighbor, such as the possibility of exchanging bulls, and of owning good purebred bulls cooperatively. These advantages are obtained by those having grade herds as well as by those with purebreds. Then there is also the opportunity for taking advantage of special breed sales of surplus stock, and, lastly, the advantage of bringing the community together in other endeavors which usually result where there is but one breed.

PERSONAL PREFERENCE

In a district where no breed is established, or in sections where all the different breeds are about equally represented, the prospective breeder must be guided largely by his personal preference. A person usually takes a liking to one breed, for reasons not easily explainable. Naturally he would take more interest in caring for animals of that breed than for those of a breed that he does not like so well.

Personal preference, however, must not overshadow the matter of quality of individual animals. If high-producing individuals of the breed not so well liked are available at reasonable cost, and individuals of the same quality of the breed well liked are not available except at a much higher cost, it may be wiser to select the former, for usually a dairyman soon begins to like a breed with which he is doing well.

MARKET REQUIREMENTS FOR PRODUCT

Market requirements for the product should not be overemphasized in selecting the breed. For a time a dairyman may sell his product in a market where low-testing milk has the advantage, while later the conditions may be changed, and a high-testing milk will sell to better advantage. Obviously, a breeder can not shift from one breed to another to meet the fluctuations in market demands.

When selling to a city milk plant, however, the price paid for the extra fat over the basic test, or deducted from the standard price when




FIG. 1.-Diagram of cow showing names and location of parts

the milk is below basic test, may well be considered in selecting the breed. The point here is that sometimes in some whole-milk markets the differential may favor high-testing milk, and at other times or in other markets it may favor low-testing milk.

In summing up the matter of which breed to select this point should be kept in mind—there are good cows and poor cows in all breeds, and, other things being equal, the breeder or dairyman who gets good individuals to begin with will have a good chance for success, no matter what breed he selects.

THE SCORE CARD

Each breed association has a scale of points, or score card, for bulls and cows of that breed. The card gives definite values for the various characteristics of conformation, and emphasizes points requiring special attention from breeders. The purpose of the score card is to teach beginners the art of judging, and also to encourage the formation of what is considered by breeders, through their associations, as the ideal type. It tends to make the breed uniform in appearance. The scale of points for a cow is given in this bulletin with the description of each breed.

In order to make the score cards more useful a diagram is given in Figure 1, which names and locates the various parts referred to on the score cards.

AYRSHIRE

ORIGIN AND HISTORY

The Ayrshire breed originated in southwestern Scotland, in the county of Ayr, in the latter part of the eighteenth century. Doubtless cattle from several neighboring countries were used in the formation of the breed, though there is no record of direct foreign importations to the county of Ayr at that time. While this foreign blood probably had a good effect on the ultimate value of the breed, the substantial and efficient development of the breed seems to have come about mostly through subsequent judicious selection and mating.

IMPORTATION AND DISTRIBUTION

The first importations into the United States occurred in 1822. Since then Ayrshires have been imported almost every year, either from Scotland or Canada. As indicated by Tables 1 and 2, there are about 30,000 registered Ayrshires in the United States and about 400,000 grades carrying more or less Ayrshire blood. These cattle are found scattered through 46 of the 48 States of the Union, though by far the largest numbers are to be found in the Northeastern States.

GENERAL CHARACTERISTICS

The Ayrshire has a well-built, stocky body, not heavily covered with flesh, but giving the appearance of great vigor and vitality. The calves weigh from 60 to 80 pounds at birth. The weight of mature bulls varies from 1,500 to 2,000 pounds, with an average of about 1,650 pounds, while mature cows range in weight from 850 to 1,250 pounds, and average about 1,050 pounds.

The color varies from almost pure white to nearly all cherry red or brown, with any combination of these colors. Usually the tail is white. The horns are large, and turn gracefully outward, then forward and back, giving a distinctive appearance to the head.

Ayrshire cows are noted for their symmetrical udders, which extend well forward and back, with no tendency to be pendent. The quarters are generally even, the teats medium in size and well placed.

Dairy Callle Breeds





FIG. 2.-Ayrshire bull, Strathglass Roamer 28189. Grand champion, National Dairy Show, 1925



F16.3.—Ayrshire bull, Beuchan Peter Pan 12971. Fifty-seven of his daughters arc in the Advanced Registry 45282°-27-2




FIG. 4 .- Ayrshire cow, Lily of Willowmoor 22269. Champion butterfat producer of the breed




FIG. 5.-Ayrshire cow, Garclaugh May Mischief 27944. Champion milk producer of the breed

Scale of points for Ayrshire cow

Head		10
Forehead, broad and clearly defined	1	10
Horns, wide set on and inclining upward	ĩ	,
Face, of medium length, slightly dished; clean cut, showing veins	2	
Muzzle, broad and strong without coarseness, nostrils large	1	
Jaws, wide at the base and strong	1	
Eves, full and bright with placid expression	3	
Ears, of medium size and fine, carried alert	1	
Neck, fine throughout, throat clean, neatly joined to head and shoul-		
ders, of good length, moderately thin, nearly free from loose skin,		
elegant in bearing	~	3
Fore quartersShoulders, light, good distance through from point to point but		10
shown at withow, amosthly blanding into hady	9	
sharp at withers, smoothly blending into body Chest, low, deep and full between back and forelegs	$\frac{2}{6}$	
	1	
Brisket, light Legs and feet, legs straight and short, well apart, shanks fine and	· *	
smooth, joints firm, feet of medium size, round, solid and deep	1	
Body	1	13
Back, short and straight, chine lean, sharp and open jointed	4	-0
Loin, broad, strong, and level	$\bar{2}$	
Ribs, long, broad, wide apart, and well sprung	3	
Abdomen, capacious, deep, firmly held up with strong muscular		
development Flank, thin and arching	3	
Flank, thin and arching	- 1	
Hind quarters		11
Hind quarters Rump, wide, level, long from hooks to pin bones, a*reasonable		
pelvic arch allowed Hooks, wide apart and not projecting above back nor unduly over-	3	
Hooks, wide apart and not projecting above back nor unduly over-	2	
laid with fat Pin bones, high, wide apart	· 1	
Thighs, thin, long, and wide apart	$\frac{1}{2}$	
Tail fine long and set on level with back	1	
Tail, fine, long, and set on level with back Legs and feet, legs strong, short, straight, when viewed from be-	-	
hind and set well apart; shanks fine and smooth, joints firm,		
feet medium size, round, solid, and deep	2	
Udder, long, wide, deep but not pendulous nor fleshy: firmly attached		
to the body, extending well up benind and lar forward; quarters		
even; sole nearly level and not indented between teats, udder veins		4 - 19 -
well developed and plainly visible	'	22
Teats, evenly placed, distance apart from side to side equal to half the breadth of udder, from back to front equal to one-third the	1	5.2
the breadth of udder, from back to front equal to one-third the		
length; length $2\frac{1}{2}$ to $3\frac{1}{2}$ inches, thickness in keeping with length,		8
hanging perpendicularly and not tapering Mammary veins, large, long, tortuous, branching, and entering large		0
		5
orifices		U
well upward		2
well upward Color, red of any shade, brown, or these with white; mahogany and white, or white, each color distinctly defined. (Brindle markings		
white, or white, each color distinctly defined. (Brindle markings		
allowed but not desired)		2
Covering Skin, medium thickness, mellow, and elastic		6
Skin, medium thickness, mellow, and elastic	3	
Hair, soft and fine	2	
Secretions, oily, of rich brown or yellow color	1-	
Style, active, vigorous, showing strong character, temperament inclined		
to nervousness but still docile		4
Weight, at maturity not less than 1,000 pounds		4
Total		100
		_ • •

9

PRODUCTION

Ayrshire milk contains a percentage of butterfat that is about the average of all the dairy breeds. The 6,368 cows and heifers that completed official records up to January 1, 1925, produced an average of 10,256 pounds of milk per cow, containing 407.48 pounds of butterfat, or 3.97 per cent butterfat.

The 10 highest producers of butterfat and milk among the Ayrshires are listed below:

Ten highest Ayrshire yearly butterfat and milk production records in the United States

Butterfat production	Milk production		
Cow	Butter- fat	Cow	Milk
Lily of Willowmoor 22269 Vis Bountiful Lassie 58096 Auchenbrain Brown Kate 4th 27943 Garclaugh Ma y Mischief 27944 Auchenbrain Yellow Kate 3d 36910 Agawam Bess Howie 43781 Harperland Spicy Lass 40652 Jean Armour 3d 32219 Nancy Whitehall 47810 Bloomer's Queen 30119	866. 21	Garclaugh May Mischief 27944 Vis Bountiful Lassie 58096 Mistress Thistle of South Farm 49818 Auchenbrain Brown Kate 4th 27943 Lily of Willowmoor 22269 Garclaugh Spottie 27950 Nancy Whitehall 47810 Jean Armour 3d 32219 Bloomer's Queen 39119 Wil'owmoor May Mischief 2d A 34173	Pounds 25, 32 24, 55 23, 02 23, 02 22, 53 22, 53 21, 53 21, 53 21, 82 21, 16

BULLS

The 10 Ayrshire sires having the largest number of daughters with official yearly records, up to January 1, 1926, are listed below.

Ten Ayrshire sires having the largest number of Advanced-Registry daughters

Sire	Number cf daugh- ters	Sire	Number of daugh- ters
Beuchan Peter Pan 12971	57	Nox'email 7312	35
Finlayston 8882.	55	White Cloud of Hickory Island 10377	32
Baron's Best Bargenoch 12858.	47	Netherton Statesman 11643	31
Earl's Choice of Spring Hill 8289	43	Kate's Good Gift 15426.	30
Kate's Champion of Penhurst 18782	38	Morton Mains Queechy 11537	27

BROWN SWISS

ORIGIN AND HISTORY

The original home of the Brown Swiss breed is in Switzerland, where it has been developed during many centuries. It is probably one of the oldest breeds in existence, and it is thought that no outside blood has been introduced since records began.

IMPORTATION AND DISTRIBUTION

The first importation of Brown Swiss into the United States was made in Massachusetts in 1869 and another in 1882. A number of importations have been made since, but only in small numbers. After 1906 there were only a few importations because of regulations due to the prevalence of the foot-and-mouth disease in Europe. As shown in Tables 1 and 2, there are in the United States about 8,000 registered animals, and about 162,000 grades carrying more or less Brown Swiss blood. These are to be found scattered in 37 States, the largest numbers being in Wisconsin, Illinois, New York, Minnesota, Michigan, Iowa, Pennsylvania, and Ohio. Of late years the breed has made notable increase in popularity.

GENERAL CHARACTERISTICS

The large frame of the Brown Swiss cattle indicates that they have been developed for service as draft animals as well as for milk. They are substantial in appearance, well proportioned, with the body well covered with flesh. The calves weigh from 65 to 90 pounds at birth.




FIG. 6.-Brown Swiss bull, Reuben 2927. Twenty-three of his daughters are in the Advanced Registry

The heifers are slow in maturing. When full grown the cows weigh from 1,100 to 1,500 pounds, averaging about 1,250 pounds; and the bulls range in weight from 1,500 to 2,200 pounds, averaging about 1,750 pounds.

The color of the Brown Swiss varies from dark to light brown, and at some seasons of the year approaches gray. There is usually a light stripe of gray along the back. White splashes near the udder are found on some animals, but white splashes on the sides of the body or on the back are objectionable. The hair between the horns is usually a lighter shade than the body. The nose, switch, tongue, and horn tips are always black, and there is usually a light or mealy ring around the muzzle.




FIG. 7.-Brown Swiss bull, Ursula Stasis 10719. Grand champion, National Dairy Show, 1925




FIG. 8 .- Brown Swiss cow, June's College Girl 11427. Champion butterfat producer of the breed




FIG. 9.-Brown Swiss cow, Believe 4245. World's champion of her breed in production of milk

Scale of points for Brown Swiss cow or heifer

	2
	2
Ears, large, fringed inside with light-colored hair, skin inside of ear a deep	P
orange color	2
Nose, black, large, and square, with mouth surrounded by mealy colored band, tongue black	2
Eves, moderately large, full, and bright	2
Horns, short, regularly set with black tips	2
Neck, straight, throat clean, neatly joined to head, shoulders of good length,	
	1
Chest, low, deep, and full between and back of forelegs	ŝ
Baek, level to setting of tail and broad across the loin	6
Ribs, long and broad, wide apart and well sprung, with thin, arching flanks_	2
	5
	1
Thighs, wide, quarters not thin	î
Legs, short and straight, with good hoofs	ĵ
Tail, slender, well set on, with good switch	2
Hide of medium thickness, mellow and elastic	3
Color-shades from dark to light brown, at some seasons of the year gray;	í
white splashes near udder not objectionable, light stripe along back.	
White splashes on body or sides objectionable. Hair between horns usually	
	1
Fore udder, wide, deep, well rounded but not pendulous nor fleshy, extending	×,
far forward on the abdomen12	,
Rear udder, wide, deep, but not pendulous nor fleshy, extending well up	7
behind 12	9
	8

 Milk veins large, long, tortuous, elastic, and entering good wells______
 6

 Disposition, quiet______
 2

 Size, evidence of constitution, and stamina______
 5

Total_____ 100

PRODUCTION

The Brown Swiss produces milk of average quality compared with the other breeds of dairy cattle. The 476 cows and heifers that completed yearly records and were admitted to the Register of Production up to January 1, 1926, have an average yearly production of 12,534 pounds of milk and 501.6 pounds of butterfat per cow, with an average butterfat test of 4 per cent. The 10 highest butterfat and milk producers among the Brown Swiss are listed below.

Ten highest Brown Swiss yearly butterfat and milk production records in the United States

Butterfat production	Milk production		
Cow	Butter- fat	Cow	Milk
June's College Girl 11427 Believe 4245. Hawthorn Dairy Maid 6753 Alice Lee 2d 8777. Clothilda Werder's Aggie 10691 Bodlie Merrillan 12103. Darling D. 8845. Swiss Valley Girl 8th 6574. Hawthorn Dairy Maid 2d 8550. Millicent of Walhalla 11178	927. 23 914. 38 880. 22 873. 51 860. 52	Believe 4245 Alice Lee 2d 8777 Une's College Girl 11427 Clothilda Werder's Aggie 10691 Hawthorn Dairy Maid 6753 Sterling Pride of Lake View 9530 Goldie Merrillan 12103 Belle Windsor of Tarpleywick 9000 Hawthorn Dairy Maid 2d 8550 Swiss Valley Girl 8th 6574	Pounds 25, 848 24, 845 24, 571, 6 22, 809 22, 765 22, 597 22, 093 21, 211 21, 179 20, 812

BULLS

The 10 Brown Swiss sires having the largest number of daughters with official yearly records, up to January 1, 1926, are listed below.

Ten Brown Swiss sires having the largest number of daughters in Register of Production

Sire	Number of daugh- ters	Sire	Number of daugh- ters
Reuben 2927	23	Vogel's College Boy 5638	8
Junker 2365.	10	Casper C. 1999.	7
Collier 2075.	10	Ben Hanson 2373	7
College Master 2986.	10	Sunny Hill Bob 6170	7
Tom Phylis 1769.	9	Merney's Second Son 3280	7

DUTCH BELTED

ORIGIN AND HISTORY

The Dutch Belted breed originated in Holland about two centuries ago. The breed gets its name both from the original home and from the distinctive color marking. This breed has probably been developed from the same cattle as the Holstein-Friesian. The early records show that the Dutch Belted were bred by the nobility of Holland, and while the unusual color marking was perhaps the chief basis of selection, the qualities of milk production and dairy refinement were not lost sight of.

14

IMPORTATION AND DISTRIBUTION

The first importation of Dutch Belted cattle into the United States was made probably in 1838. The first importation of importance, however, was made in 1840 by P. T. Barnum for show purposes. These cattle later were placed on a farm, and this seems to be the beginning of the Dutch Belted cattle in America. A number were imported from that time on until 1885, and some in 1906 and 1907. Since then no importations have been made on account of the prevalence of foot-and-mouth disease in Europe. As shown in Table 1, there are about 6,000 purebred Dutch Belted cattle in the United States, and about 150,000 grades.

GENERAL CHARACTERISTICS

Dutch Belted cattle have the general dairy conformation, which includes fineness of bone and freedom from beefiness. The aim of the breeders of these cattle is to breed animals that have no white other than that of the standard belt around the body. This belt begins back of the shoulder and may extend to the front of the hips but must not be narrower than 6 inches at the narrowest point. There must be no black spots in the belt on females. The width of the belt on each animal tends to be uniform around the body. The remainder of the animal is coal black except that females may have not to exceed 3 inches of white on hind feet above the hoof, and males may have not to exceed $2\frac{1}{2}$ inches of white on one hind foot above the hoof.

Calves at birth range in weight from 60 to 90 pounds. Welldeveloped mature cows weigh from 1,000 to 1,500 pounds, averaging about 1,200 pounds; and bulls from 1,500 to 2,000 pounds, averaging about 1,700 pounds.

Scale of points for Dutch Belted cow

Body color, black, with a clearly defined continuous white belt. The belt to be of medium width, beginning behind the shoulder and extending	
nearly to the hips	8
Head, comparatively long and somewhat dishing; broad between the eyes. Poll, prominent; muzzle, fine; dark tongue	6
Eyes, black, full and mild. Horns long compared with their diameter Neck, fine and moderately thin and should harmonize in symmetry with the	4
head and shoulders	6
Shoulders fine at top, becoming deep and broad as they extend backward	-
and downward, with a low chest	4
Barrel large and deep with well-developed abdomen; ribs well rounded and	2.0
free from fat	10
Hips broad, and chine level, with full loin	10
Rump high, long, and broad	6
Hind quarters long and deep, rear line incurving; tail long, slim, tapering to a full switch	8
Legs short, clean, standing well apart	3
Udder large, well-developed front and rear; teats of convenient size and	0
apart: mammary veins large, long, and crooked, entering large orifices	20
Escutcheon	$\overline{2}$
Hair fine and soft; skin of moderate thickness of a rich, dark, or yellow color-	3
Quiet disposition and free from excessive fat	4
General condition and apparent constitution	6
Perfection	100




Fig. 10.-Dutch Belted cow, Angelina 2641. Leading butterfat producer of the breed




FIG. 11.-Dutch Belted cow, Gem of Columbia 2033. Leading milk producer of the breed

PRODUCTION

By referring to Table 3 it will be seen that the Dutch Belted cow ranks between the Holstein and the Ayrshire in the percentage of butterfat contained in her milk. The Dutch Belted cows and heifers that finished yearly official records up to January 1, 1926, show an average production of 10,141 pounds of milk and 384.64 pounds of butterfat, with an average test of 3.8 per cent. The 10 highest producers of butterfat and milk among Dutch

Belted cows are listed below.

Ten highest Dutch Belted yearly butterfat and milk production records in the United States

Butterfat production	-	Milk production	
Cow	Butterfat	Cow	Milk
Angelina 2641	531, 19 516, 07 504, 99 501, 09 484, 31 447, 64	Gem of Columbia 2038. Angelina 2641. Ferndell 1961. Priscilla of Lakeview 2937. Glenbeulah's Beauty 2172. Peapack Anna 1487. Peapack Dutchess 1390, imported. Orange Lilly 3126. Eunice 1597. Fritzi 1832.	13, 478 13, 385 13, 296 13, 159 13, 065 11, 575

BULLS

The 10 Dutch Belted sires having the largest number of daughters, with official yearly records, up to January 1, 1926, are listed below.

Dutch Belted sires having two or more daughters with Advanced-Register official yearly records

Defendant 1185. Keith 934. Sutton's Gay Lad 494. Michigans Brings 1959	Monarch of Lakeview 1219. Bruce W. 729. Samoset 1134.	Glenbeulah's Duke 1092. Columbia King 1015. Ka No No of Council Grove 1184.
Michigans Prince 1258.) <u> </u>	

GUERNSEY

ORIGIN AND HISTORY

The Guernsey breed originated in the Channel Islands, near the north coast of France. It is thought that this breed has been developed from a cross between the large red and brindle cattle of Normandy and the small red cattle of Brittany, in France. The exact date of origin is unknown, but it was probably in the latter part of the seventeenth century or before.

All the cattle in the Channel Islands were at one time known as After laws had been enacted forbidding the importation Aldernevs. of cattle from the Continent or between the islands of Guernsey and Jersey, two distinct breeds came to be recognized. The one on the islands of Alderney, Sark, and Guernsey became known as the Guernsey breed and the one on Jersey Island as the Jersey breed.




FIG. 12,—Guernsey bull, Yeoman's King of the May 17053. He has 103 daughters in the Advanced Registry




FIG. 13.-Guernsey bull, May Rose Cherub 70934. Grand champion, National Dairy Show, 1925




FIG. 14.—Guernsey cow, Anesthesia Faith of Hill Stead 114354. Champion butterfat producer of the breed




FIG. 15.-Guernsey cow, Murne Cowan 19597. Champion milk producer of the breed

IMPORTATION AND DISTRIBUTION

The first cattle from the Channel Islands brought to America were called Alderneys. They were imported in the latter part of the eighteenth century and may have been either Guernsey or Jersey cattle. The first animals recorded in the herdbook of the American Guernsey Cattle Club were brought over in 1830. A few more were imported in the next two decades, but not until about 1870 were extensive importations made. Since that time importations have been made nearly every year.

According to Tables 1 and 2, there are about 80,000 purebred Guernsey cattle in the United States, and about 1,900,000 grades. They are to be found in every State of the Union, but by far the largest numbers are found in the North Atlantic and North Central States.

GENERAL CHARACTERISTICS

In size the Guernseys are about equal to the Ayrshires and slightly smaller than the Brown Swiss. The calves weigh from 55 to 85 pounds at birth, and reach maturity early. When mature the cows weigh from 800 to 1,400 pounds, averaging about 1,050; and the bulls from 1,200 to 2,200 pounds, averaging about 1,600 pounds.

The color of the Guernseys is fawn and white, with fawn predominating. A light cherry red with white is also found. Sometimes white may be entirely lacking except on the legs. The switch is usually white and the tongue light in color. The horns are of moderate size and amber in color. The skin is yellow.

Scale of points for Guernsey cow

Style and symmetry, attractive individuality revealing vigor, femininity and breed character; a harmonious blending and correlation of parts; an active well-balanced walk	5
Head, moderately long, clean-cut, showing femininity and breed character; a lean face; wide mouth and broad muzzle with open nostrils; strong jaws; full bright eyes with gentle expression; forehead broad between the eyes and moderately dishing; bridge of nose straight	5
Horns, yellow, small at base; of medium length; inclining forward; not too spreading	1
Neck, long and thin; clean throat, smoothly blending into shoulders Withers, chine rising above shoulder blades, with open vertebrae	$\frac{2}{2}$
Shoulders, shoulder blades set smoothly against chine and chest wall, form- ing neat junction with the body	2
Chest, wide, and deep at heart with least possible depression back of the shoulders	4
Back, appearing straight from withers to hips Loin, strong, broad, and nearly level laterally; width carried forward to junction with the ribs	$\frac{5}{3}$
Hips, wide apart, approximately level with the back; free from excess tissue_ Rump, long, continuing with level of the back; approximately level between hip bone and pin bones. Pin bones well apart	$\frac{3}{2}$
Thurls, wide apart and high	$\hat{2}$
Barrel; deep and long with well-sprung ribs. Individual ribs: Long, flat, wide apart, and free from excess tissue	10
Thighs, incurving when viewed from side, thin and wide apart when viewed from rear; well cut up between the thighs	2
Legs, flat flinty bone, tendons clearly defined; front legs straight; hind legs	-
nearly upright from hock to pastern, set wide apart and nearly straight when viewed from behind. Pastern: Strong and springy	2
Hide, loose and pliable, and not thick, with oily feelings; hair, fine and silky Tail, long, tapering with neat, strong, level attachment, neatly set between	3
pin bones; fine bone and hair; nicely balanced switch	2

Udder, uniformly fine in texture; free from meatiness; covered with pliable	
velvety skin	3
Veins prominent	1
Attachment to body: Strong, long, and wide	. 4
Extending well forward; extending well up behind	4
Sole: Level between teats	2
Teats: Of even convenient size; cylindrical in shape; well apart and squarely placed, plumb	3
Mammary veins, long, tortuous, prominent and branching with large numer- ous wells	3
Secretions indicating color of product, indicated by the pigment secretion of skin which should be a deep yellow inclining toward orange in color; especially discernible in the ear, at the end of bone of tail, around the eyes and nose, on the udder and teats, and at the base of horns; hoofs and	
horns amber colored	20
Color markings, a shade of fawn with white markings	2
Size, mature cows, about 1,100 pounds in milking condition	2
Total	100

PRODUCTION

Guernsey milk has a high per cent of butterfat and a yellow color. The 20,931 cows and heifers that completed official records up to January 1, 1926, produced an average per cow of 9,615.2 pounds of milk and 480.1 pounds of butterfat, the average fat test being 4.99 per cent.

The 10 highest butterfat and milk producers among the Guernseys are listed below.

Ten highest Guernsey yearly butterfat and milk production records in the United States

Butterfat production		Milk production	
Cow Anesthesia Faith of Hill Stead 114354 Countess Prue 43785 Murne Cowan 19697 May Rilma 22761. Gertrude Claire 99550	Butterfat Pounds 1, 112. 50 1, 103. 28 1, 098. 18 1, 073. 41 1, 020. 00	Cow Murne Cowan 19597 Katherine's Trixie 100396 Gertrude Claire 99550 Nella Jay 4th 38233 Molly's Lassie 86472	Milk Pounds 24,008 21,071 20,738 20,710 20,592
Nella Jay 4th 38233 Langwater Nancy 27943 Dairy Maid Queen of Spring Hill 74067 Langwater Hope 27946 Lillia of Iowa 2d 61500	1,019.25 1,011.66 1,011.18 1,003.17 998.40	Miss Daisy of Maple Hill 72610 Tom's Daisy 54541 Jean Du Luth Coronet 52816 Langwater Hope 27946 Yeksa's Top's of Gold's Fannie 22362	20, 315 20, 298 20, 004 19, 882 19, 795

BULLS

The 10 Guernsey sires having the largest number of daughters with official yearly records, up to January 1, 1926, are listed below.

Ten Guernsey sires having the largest number of Advanced-Register daughters

Sire	Number of daugh- ters	Sire	Number of daugh- ters
Governor of the Chêne (R. G. A. S. 1297 P. S.)	114 103 71 70 58	Galaxy's Sequel 16904 (Imp.) Ne Plus Ultra 15265 Cora's Governor of Chilmark 8971 (Imp.). Beda's May King 11893 Florham Laddie 20431	53 48 47 45 42

HOLSTEIN-FRIESIAN

ORIGIN AND HISTORY

The cattle from which our present Holstein-Friesian breed has descended were developed in northern Holland, especially in the Province of Friesland, and in the neighboring Provinces of northern Germany. The time of their origin as a recognized distinct breed is unknown, but it is probable that they have been selected for their dairy qualities for about 2,000 years.

Before 1885 there were two associations furthering the interests of this breed in the United States. One maintained a Holstein herdbook, and the other a Dutch-Friesian herdbook. In 1885 the two associations were combined into the Holstein-Friesian Association of America, and from that time on only one herd register has been maintained. This is known as the Holstein-Friesian herdbook. While the official name of the breed is Holstein-Friesian, the single word Holstein is more common in ordinary use.

IMPORTATION AND DISTRIBUTION

The first importations of Holsteins into the United States were made in 1795, and afterwards a few were brought in from time to time up to 1879, following which heavy importations were made each year until 1887. Thereafter only a few were imported up to 1905, and since then, because of the prevalence of foot-and-mouth disease in Europe, very few have been imported. According to Tables 1 and 2 there are about 528,000 purebred

According to Tables 1 and 2 there are about 528,000 purebred Holsteins in the United States, and about 10,500,000 grades. Holstein cattle are found throughout all the 48 States of the Union, though by far the largest numbers are in New York, Wisconsin, Pennsylvania, Ohio, Michigan, and Illinois, in the order named. These six States contain more than 60 per cent of the purebred Holstein cattle in the United States.

GENERAL CHARACTERISTICS

The Holsteins are the largest of the dairy breeds. They have large frames, not heavily covered with flesh. The calves weigh from 70 to 105 pounds at birth. The mature bulls weigh from 1,600 to 2,200, and average about 1,900 pounds; and the mature cows weigh from 1,100 to 1,750, and average about 1,250 pounds. The color is black and white, with the colors sharply defined rather than blended. They may be nearly all white or black, but no solid-color animal can be registered.

Scale of points for Holstein-Friesian cow

Forehead, broad between the eyes; dishing	2
Face, of medium length; clean cut; feminine; the bridge of the nose straight_	1
Muzzle, broad, with strong lips; nostrils, large and open; jaws strong	3
Ears, of medium size; of fine texture; well carried	1
Eyes, large; full; mild; bright	2
Horns, small; tapering finely toward the tips; set moderately narrow at base;	
inclining forward; well curved inward	1
Neck, long; fine and clean at junction with the head; evenly and smoothly	
joined to shoulder	3
Shoulders, slightly lower than the hips; smooth and rounding over tops;	
moderately broad and full at sides	3

Crops, full; level with the shoulders	
Chine, straight; strong; broadly developed, with open vertebra	
Loin and hips, broad; level or nearly level between the hip bones; level and strong laterally; spreading from chine broadly and nearly level; hip bones	
fairly prominent Rump, long; broad with roomy pelvis; nearly level laterally; full above the thurls; carried out straight to tail head	:
Pin bones, wide between; nearly level with hips	
Thurls, high; broad through Tail head and tail, strong at base without coarseness; the setting well back; tail long, tapering finely to a full switch	
Chest, deep; wide; well filled and smooth in the brisket; broad between the forearms; full in the foreflanks	
Barrel, long; deep; well rounded; strongly and trimly held up Flanks, deep; full	
Thighs, wide; deep; straight behind; wide and moderately full at the out- sides; twist well cut out and filled with development of udder; escutcheon well defined	
Mammary veins, large, tortuous, entering large orifices or double extension; with additional developments, such as branches and connections entering numerous orifices	
Udder, capacious; flexible; quarters even and of uniform texture, filling the space in the rear below the twist, extending well forward; broad and well	
attached Teats, well formed; plumb; of convenient size; properly placed	1
Legs, medium length; clean; nearly straight; wide apart; firmly and squarely set under the body; arms wide, strong, and tapering	
Hair and hide, hair healthy in appearance; fine and soft; hide of medium thickness; mellow and loose	
	10

PRODUCTION

The Holsteins produce a larger quantity of milk, with a lower butterfat content, than any other dairy breed. The milk is not so highly colored as that from the Guernseys and Jerseys. The 23,622 official records of Holstein cows and heifers that were

completed up to January 1, 1926, show an average yearly production of 15,697 pounds of milk and 532.44 pounds of butterfat, the average test being 3.39 per cent. The 10 highest butterfat and milk producers among the Holsteins

are listed below.

Ten	highest	Holstein	y early	butterfat	and	milk	production	records	in	the	United
					State	88					

Butterfat production		. Milk production	
Cow	Butter- fat	Cow	Milk
De Kol Pius Segis Dixie 295135 ¹ May Walker O.lie Homestead 300043 Hollywcod Lilith Palmyra Abbekerk 400491 Duchess Skylark Ormsby 124514 Bess Johanna Ormsby 263431 Redfield Segis Johanna 735550 Finderne Pride Johanna Rue 121083 Segis Pietertje Prospect 221846 Dutchland Pietertje Creamelle 394861 Dutchland Pietertje Creamelle 394861 Orahamholm Colantha Pauline Segis 405465	1, 206. 79 1, 205. 09 1, 198. 09 1, 182. 93 1, 176. 47 1, 158. 95 1, 148. 54	Segis Pietertje Prospect 221846 Helm Veeman Woodcrest 486877 Kolrain Marion Finderne 317396 Nooksack Lunde Oregon De Kol 301119. Queen Carlotta De Kol 311674 Adirondac Wietske Dairy Maid 204072 Grahamholm Colantha Pauline Segis 405465. Princess Aaggie Polkadot De Kol 372024. Grahamholm Colantha Segis Maid 405466	Pounds 37, 381 36, 218 35, 340 35, 085 34, 511 34, 430 34, 402 34, 071 33, 812

¹ Canadian cow.




Fig. 16.—Holstein bull, Sir Pietertje Ormsby Mercedes 44931. He has 65 yearly-record daughters in the Advanced Register




FIG. 17.—Holstein bull, Johanna Rag Apple Pabst 346005. Grand champion, National Dairy Show, 1925




FIG. 18.--Holstein cow, Segis Pietertje Prospect 221846. Has highest yearly milk record of all the breeds




FIG. 19.—De Kol Plus Segis Dixie 295135. Has highest yearly butterfat record of all the breeds (Canadian cow)

BULLS

The 10 Holstein-Friesian sires having the largest number of daughters with yearly records, up to January 1, 1926, are listed below.

Ten Holstein sires with the largest number of yearly-record daughters

Sire	Number of daugh- ters	Sire	Number of daugh- ters
Sir Inka Prilly Segis 80914 Judge Segis 80912 Sir Johanna Fayne 42147 King Pontiac Champion 53418 Matador Segis Walker 148839	67	Sir Pietertje Ormsby Mercedes 44931 Woodcrest Tehee 74219 Colantha Sir Walker Korndyke 95460 King of the Pontiacs 39037 Ormsby Korndyke Lad 102469	

JERSEY

ORIGIN AND HISTORY

The Jersey breed originated in the Island of Jersey, one of the group of Channel Islands, between England and France. In 1789 a law was passed prohibiting the importation of cattle into Jersey Island except for immediate slaughter. Shortly afterwards the cattle on that island became known by the name of Jersey instead of Alderney. No outside blood has been introduced since that time.

IMPORTATION AND DISTRIBUTION

The first importation of Jerseys into the United States was made in 1850. A few more were brought over about 20 years later, and from 1870 to 1890 there were numerous importations. Since 1890 many Jerseys have been imported every year.

The Jerseys are more evenly distributed in the United States than any other breed. There were about 232,000 purebred and about 9,300,000 grade Jerseys in this country in 1920, well scattered throughout all the 48 States.

GENERAL CHARACTERISTICS

The Jersey is the smallest of the breeds discussed in this bulletin. The calves weigh from 40 to 75 pounds at birth. The heifers develop rapidly and mature sufficiently to drop the first calf at 24 months of age. The mature cows weigh from 700 to 1,200 pounds, averaging about 900 pounds, and the bulls weigh from 1,200 to 1,800, averaging about 1,500 pounds.

The color of Jerseys is usually some shade of fawn or cream color, though different shades of mouse color, gray, and brown are common, and some individuals approach black. They may be solid color of any of these shades, or spotted with white. The muzzles and tongues are usually black or lead colored, but light-colored tongues are not uncommon, and around the muzzle is a white or mealy ring.




FIG. 20.-Jersey bull, Pogis 99th of Hood Farm 94502. He has 111 daughters in the Register of Merit




FIG. 21.-Jersey hull, Fern's Rochette Nohle 193011. Grand champion, National Dairy Show, 1925





FIG. 22 .- Jersey eow, Madeline of Hillside 389336. Champion milk producer of the breed



F1G. 23.-Jersey cow, Darling's Jolly Lassie 435948. Champion butterfat producer of the breed

Scale of points for Jersey cow

DAIRY TEMPERAMENT AND CONSTITUTION

Head, 7:	
A. Medium size, lean; face dished; broad between eyes; horns medium	
size, incurving	3
B. Eyes full and placid; ears medium size, fine, carried alert; muzzle	·
broad, with wide-open nostrils and muscular lips; jaw strong	- 4
Neck, 4:	
Thin, rather long, with clean throat, neatly joined to head and shoulders_	4
Body, 37: A. Shoulders light, good distance through from point to point, but thin	
at withers; chest deep and full between and just back of forelegs	5
B. Ribs amply sprung and wide apart, giving wedge shape, with deep,	
large abdomen, firmly held up, with strong, muscular development	10
C. Back straight and strong, with prominent spinal processes; loins	
broad and strong	5
D. Rump long to tail setting, and level from hip bones to rump bones	6
E. Hip bones high and wide apart	33
F. Thighs flat and wide apart, giving ample room for udder	3
G. Legs proportionate to size and of fine quality, well apart, with good feet, and not to weave or cross in walking	2
H. Hide loose and mellow	$\frac{2}{2}$
I. Tail thin, long, with good switch, not coarse at setting on	์ 1
1. Tail tinin, long, with good switch, not coarse at setting on	् ै
MAMMARY DEVELOPMENT	
Udder, 26:	C
A. Large size, flexible, and not fleshy B. Broad, level or spherical, not deeply cut between teats	- 6 - 4
C. Fore udder full and well rounded, running well forward of front teats_	10
D. Rear udder well rounded, and well out and up behind	6
Teats, 8:	Ū,
Of good and uniform length and size, regularly and squarely placed	8
Milk veins, 4:	0
Large, long, tortuous and elastic, entering large and numerous orifices	4
Sino At	

a high-class animal, with capacity for feed and productiveness at pail_____10 Total score_______100

PRODUCTION

Jersey milk is yellow and rich in butterfat. The 22,355 cows and heifers that had Register-of-Merit records up to January 1, 1925, averaged 8,287 pounds of milk and 444.37 pounds of butterfat a year, with an average test of 5.36 per cent.

The 10 highest butterfat and milk producers among the Jerseys are listed below.

۲

Ten highest Jersey yearly butterfat and milk production records in the United States

Butterfat production		Milk production		
Cow	Butter- fat	Cow	Milk	
Darling's Jolly Lassie 435948 Groff's Constance 367292 Prince's Emma of H. S. F. 359390 Lad's lota 350672 Fauvic Ruth 385463 Madeline of Hillside 389336 Plain Mary 268206 Vive La France 319616 Lady's Siken Glow 313311 St. Mawes Lad's Lady 451568	1, 130. 09 1, 109. 99 1, 048. 07 1, 047. 28 1, 044. 05 1, 040. 08 1, 039. 29 1, 038. 70	Madeline of Hillside 389336 Fauvic's Star 313018 Abigail of Hillside 457241. Passport 219742 Red Lady 396118 Lad's Likeness 338246. Eminent's Jimp's Owl 297471 St. Mawes Blossom Rosaire 355800 Miss Incz 377204 Raleigh's Sybil 372752.	20, 359 19, 695 19, 608 19, 223 19, 099 19, 061	

BULLS

The 10 Jersey sires having the largest number of daughters with official yearly records, up to January 1, 1926, are listed below.

Ten Jersey sires with largest number of daughters in Register of Merit

Sire	Number of daugh- ters	Sire	Number of daugh- ters
Pogis 99th of Hood Farm 94502	113	Sophie 19th's Tormentor 113302	75
	83	Hood Farm Torono 60326	74
	83	Imported Golden Fern's Noble 145762	65
	81	Irene's King Pogis 73182	62
	79	Sayda's Heir 3d 74817	62

BREED ASSOCIATIONS

The various breed associations and clubs maintain offices and forces whose duty it is (1) to keep the herdbooks for their respective breeds; (2) to keep a record of the animals that have qualified for the additional registration because of meritorious performance; and (3) to further the interest of the breed in other ways. The official names of these organizations and their addresses are as follows:

American Guernsey Cattle Club, Peterboro, N. H. American Jersey Cattle Club, 324 West Twenty-third Street, New York

City.

Aryshire Breeders' Association, Brandon, Vt. Brown Swiss Cattle Breeders' Association, Beloit, Wis. Dutch Belted Cattle Association of America, Rockville, Conn. Holstein-Friesian Association of America, Brattleboro, Vt.

U.S. GOVERNMENT PRINTING OFFICE: 1927