

1971
A99.9
F763245
Cop. 3

2455

FOREST SERVICE
U.S. DEPARTMENT OF AGRICULTURE

ROCKY MOUNTAIN FOREST AND RANGE EXPERIMENT STATION

Basal Area Growth of Arizona Mixed Conifer Species

Robert S. Embry and Gerald J. Gottfried¹

Growth data were collected from four small mixed conifer watersheds totaling 1,800 acres in east-central Arizona. Annual gross basal area increment was estimated to be 4.027 square feet per acre. This represents a 2.3 percent annual increase.
KEY WORDS: Stand increment estimates, basal area measurement, mixed conifer forests.

Mixed conifer stands occupy about 6 percent of the commercial forest land in Arizona. The areas they occupy, however, are some of the most productive lands in the State. These mixed conifer forests are mostly uncut, all-aged stands of Engelmann spruce (Picea engelmannii), blue spruce (Picea pungens), Douglas-fir (Pseudotsuga menziesii), white fir (Abies concolor), corkbark fir (Abies lasiocarpa var. arizonica),² ponderosa pine (Pinus ponderosa), southwestern white pine (Pinus strobiformis), and quaking aspen (Populus tremuloides) in a wide variety of mixtures.

Large-scale harvesting of these mixed stands began in 1966. Optimum management practices have not yet been fully established. Growth information needed to prescribe satisfactory timber and watershed management practices is not available for these mixed species.

This Note presents data that will provide a means of estimating gross basal area and volume growth in mixed conifer stands.

¹Embry is Associate Silviculturist, located at Flagstaff in cooperation with Northern Arizona University; Gottfried is Associate Hydrologist, located at Tempe in cooperation with Arizona State University. Station's central headquarters is maintained at Fort Collins in cooperation with Colorado State University.

²On the Kaibab Plateau, corkbark fir is replaced by subalpine fir (Abies lasiocarpa var. lasiocarpa (Hook.) Nutt.).

Procedures

Growth data were obtained during an over-story inventory of about 1,800 acres of virgin mixed conifer forests on the four Willow-Thomas Creek watersheds on the Apache National Forest in east-central Arizona. This inventory was made according to methods developed by Ffolliott and Worley.³ Increment borings were taken from one tree out of every four selected with a 25 BAF angle gage for diameter measurement. Growth information was collected from 991 trees on 556 sample points. Average annual gross basal area increment (BAI) for each d.b.h. class was computed from the increment core measurements. The distribution of sample trees by species is shown below:

Species	Number of trees
Douglas-fir	277
Ponderosa pine	172
White fir	155
Quaking aspen	121
Engelmann spruce	118
Southwestern white pine	100
Blue spruce	24
Corkbark fir	24
Total	991

³Ffolliott, Peter F., and David P. Worley. An inventory system for multiple use evaluations. U. S. Forest Serv. Res. Pap. RM-17, 15 p. 1965. Rocky Mt. Forest and Range Exp. Sta., Ft. Collins, Colo.

LIBRARY

A basal area stand table was constructed from the basic inventory data by conventional point-sampling techniques (table 1).

Results

Basal area increment.—The sapling-small pole size class accounts for almost half the total gross BAI of 4.027 square feet per acre per year (table 2). One-third of the total is produced by Douglas-fir.

The large standard errors associated with the sapling-small pole class are the result of converting basal area per acre to number of trees per acre in the double-sampling procedure to compute BAI. The smaller the diameter, the larger the number of trees it represents on a per-acre basis.

Basal area growth percentage.—The total gross annual BAI of 4.027 square feet per acre represents a 2.3 percent annual increase. The sapling-small pole class shows the greatest class increase with 6.7 percent (table 3). Corkbark fir has the greatest species increase, with 3.7 percent.

Application of Results

The prediction of growth is necessary for the formulation of any management plan.

Growth percentage provides a quick, simple, and relatively accurate method of predicting growth.

Volume growth is a function of basal area and height increment plus any changes in form. Volume growth may be a function of basal area increment alone in mature trees, in which height and form change very slowly. Such a condition would be restricted to the larger size classes in the mixed conifers.

If a short prediction period of 5 to 10 years is used and a precise estimate is not required, the growth percentages in table 3 may be used for all size classes. The volume in each species-size class multiplied by the respective growth percentage gives a useful and quick estimate of gross annual volume growth. At present we have total cubic-foot volume tables only for ponderosa pine. When such information becomes available for the other species, these growth percentages may be used.

The growth percentage method must be used with caution. Davis⁴ states that “. . . the main difficulty with growth percent as a management tool is that it is not a quantity, but a relationship with all the shifty characteristics of percentages in general.” Growth percentages vary by species, diameter class, and prediction period and are not constant for any stand, especially a natural, mixed stand.

⁴Davis, K. P. *American Forest Management*. p. 98. N. Y.: McGraw-Hill Book Co., Inc. 1954.

Table 1.--Basal area per acre on an 1,800-acre mixed conifer stand in east-central Arizona

Size class	Douglas-	Quaking	White	Ponderosa	Engelmann	White	Corkbark	Blue	Total	Distri- bution
	fir	aspen	fir	pine	spruce	pine	fir	spruce		
	Square feet									Percent
Sapling-small poles (0.1-6.9 inches)	8.45	6.92	3.55	2.43	3.51	1.57	1.03	0.45	27.91	16
Poles (7.0-10.9 inches)	6.56	12.41	2.74	1.26	5.44	1.71	1.89	.63	32.64	18
Small sawtimber (11.0-16.9 inches)	11.38	7.15	3.69	3.87	9.22	2.43	2.16	1.03	40.93	23
Medium sawtimber (17.0-22.9 inches)	10.39	1.66	4.50	7.15	4.41	1.80	.63	.76	31.30	18
Large sawtimber (23.0 inches plus)	18.97	.04	11.38	10.30	1.44	2.43	.22	.18	44.96	25
Total	55.75	28.18	25.86	25.01	24.02	9.94	5.93	3.05	177.74	
	Percent									
Distribution	31	16	14	14	14	6	3	2		

Table 2.--Gross annual basal area increment per acre on an 1,800-acre mixed conifer stand in east-central Arizona (Confidence intervals are at the 95 percent level)

Size class	Douglas-fir	Quaking aspen	White fir	Ponderosa pine	Engelmann spruce	White pine	Corkbark fir	Blue spruce	Total	Distribution
	----- Square feet -----									----- Percent -----
Sapling-small poles (0.1-6.9 inches)	0.687±.214	0.262±.077	0.203±.084	0.188±.146	0.266±.140	0.105	0.125	0.043	1.879	47
Poles (7.0-10.9 inches)	.147±.041	.340±.095	.061±.023	.027±.012	.133±.045	.061±.024	.047±.031	.016	.832	21
Small sawtimber (11.0-16.9 inches)	.193±.038	.126±.035	.053±.023	.060±.021	.185±.046	.035±.013	.040±.020	.016±.011	.708	17
Medium sawtimber (17.0-22.9 inches)	.094±.021	.019	.049±.014	.066±.017	.054±.017	.020±.006	.006	.009±.006	.317	8
Large sawtimber (23.0 inches plus)	.120±.018	<.001	.080±.017	.059±.011	.012±.006	.015±.006	.001	.004	.291	7
Total	1.241	.747	.446	.400	.650	.236	.219	.088	4.027	
	----- Percent -----									
Distribution	31	19	11	10	16	6	5	2		

Blocked-in values are means of four or fewer sample trees expanded to a per-acre value. Intervals could not be calculated because of insufficient data.

Table 3.--Annual gross basal area growth percentage for mixed conifers in 1,800-acre stand in east-central Arizona

Size class	Douglas-fir	Quaking aspen	White fir	Ponderosa pine	Engelmann spruce	White pine	Corkbark fir	Blue spruce	Average ¹
	----- Percent -----								
Sapling-small poles (0.1-6.9 inches)	8.1	3.8	5.7	7.8	7.6	6.7	12.1	9.4	6.7
Poles (7.0-10.9 inches)	2.2	2.7	2.2	2.1	2.4	3.6	2.5	2.6	2.5
Small sawtimber (11.0-16.9 inches)	1.7	1.8	1.4	1.5	2.0	1.4	1.8	1.6	1.7
Medium sawtimber (17.0-22.9 inches)	.9	1.1	1.1	.9	1.2	1.1	1.0	1.2	1.0
Large sawtimber (23.0 inches plus)	.6	1.0	.7	.6	.8	.6	.6	2.0	.6
Average ¹	2.2	2.6	1.7	1.6	2.7	2.4	3.7	2.9	¹ 2.3

¹Averages are derived from tables 1 and 2; i.e., $\frac{4.027 \text{ (table 2)}}{177.74 \text{ (table 1)}} = 2.3 \text{ percent.}$

