Eadē dicit & for Eadē

Sum J: Kenns. pred. 1.

יְהוָה

# PARADISI IN SOLE
## Paradisus Terrestris.

or

A Garden of all sorts of pleasant, flowers which our English ayre will permitt to be noursed vp:

with

A Kitchen garden of all manner of herbes, rootes, & fruites, for meate or sause vsed with vs,

and

An Orchard of all sorte of fruitbearing Trees and Shrubbes fit for our Land

together

With the right orderinge planting & preserving of them and their vses & vertues

Collected by John Parkinson Apothecary of London.

1629

Qui veut parangonner l'artifice a Nature,
Et nos parcs a l'Eden, indiscret il mesure.

Le pas de l'Elephant par le pas du ciron,
Et de l'Migle le vol par cil du mouscheron,

Mo. Bot. Garden,
1893

# TO THE QVEENES MOST EXCELLENT MAIESTIE.

*Madame,*

KNowing your Maieſtie ſo much delighted with all the faire Flowers of a Garden, and furniſhed with them as farre beyond others, as you are eminent before them; this my VVorke of a Garden, long before this intended to be publiſhed, and but now only finiſhed, ſeemed as it were deſtined, to bee firſt offered into your Highneſſe hands, as of right challenging the proprietie of Patronage from all others. Accept, I beſeech your Maieſtie, this ſpeaking Garden, that may informe you in all the particulars of your ſtore, as well as wants, when you cannot ſee any of them freſh vpon the ground: and it ſhall further encourage him to accompliſh the remainder; who, in praying that your Highneſſe may enioy the heauenly Paradiſe after the many yeares fruition of this earthly, ſubmitteth to be

*Your Maieſties*

*in all*

*humble deuotion,*

IOHN PARKINSON.

# TO THE COVRTEOVS READER.

ALthough the ancient Heathens did appropriate the first inuention of the knowledge of Herbes, and so consequently of Physicke, some vnto Chiron the Centaure, and others vnto Apollo or Æculapius his sonne; yet wee that are Christians haue out of a better Schoole learned, that God, the Creator of Heauen and Earth, at the beginning when he created Adam, inspired him with the knowledge of all naturall things (which successiuely descended to Noah afterwardes, and to his Posterity): for, as he was able to giue names to all the liuing Creatures, according to their seuerall natures; so no doubt but hee had also the knowledge, both what Herbes and Fruits were fit, eyther for Meate or Medicine, for Vse or for Delight. And that Adam might exercise this knowledge, God planted a Garden for him to liue in, (wherein euen in his innocency he was to labour and spend his time) which hee stored with the best and choysest Herbes and Fruits the earth could produce, that he might haue not onely for necessitie whereon to feede, but for pleasure also; the place or garden called Paradise importing as much, and more plainly the words set downe in Genesis the second, which are these; Out of the ground the Lord God made to grow euerie tree pleasant to the sight and good for meate; and in the 24. of Numbers, the Parable of Balaam, mentioning the Aloe trees that God planted; and in other places if there were neede to recite them. But my purpose is onely to shew you, that Paradise was a place (whether you will call it a Garden, or Orchard, or both, no doubt of some large extent) wherein Adam was first placed to abide; that God was the Planter thereof, hauing furnished it with trees and herbes, as well pleasant to the sight, as good for meate, and that hee being to dresse and keepe this place, must of necessity know all the things that grew therein, and to what vses they serued, or else his labour about them, and knowledge in them, had been in vaine. And although Adam lost the place for his transgression, yet he lost not the naturall knowledge, nor vse of them: but that, as God made the whole world, and all the Creatures therein for Man, so hee may vse all things as well of pleasure as of necessitie, to bee helpes vnto him to serue his God. Let men therefore, according to their first institution, so vse their seruice, that they also in them may remember their seruice to God, and not (like our Grand-mother Eve) set their affections so strongly on the pleasure in them, as to deserue the losse of them in this Paradise, yea and of Heauen also. For truly from all sorts of Herbes and Flowers we may draw matter at all times not only to magnifie the Creator that hath giuen them such diuersities of formes, sents and colours, that the most cunning

Worke-man cannot imitate, and such vertues and properties, that although wee know many, yet many more lye hidden and vnknowne, but many good instructions also to our selues: That as many herbes and flowers with their fragrant sweete smels doe comfort, and as it were reuiue the spirits, and perfume a whole house; euen so such men as liue vertuously, labouring to doe good, and profit the Church of God and the Common wealth by their paines or penne, doe as it were send forth a pleasing sauour of sweet instructions, not only to that time wherein they liue, and are fresh, but being drye, withered and dead, cease not in all after ages to doe as much or more. Many herbes and flowers that haue small beautie or sauour to commend them, haue much more good vse and vertue: so many men of excellent rare parts and good qualities doe lye hid vnknown and not respected, vntill time and vse of them doe set forth their properties. Againe, many flowers haue a glorious shew of beauty and brauery, yet stinking in smell, or else of no other vse: so many doe make a glorious ostentation, and flouriſh in the world, when as if they stinke not horribly before God, and all good men, yet surely they haue no other vertue then their outside to commend them, or leaue behind them. Some also rise vp and appear like a Lilly among Thornes, or as a goodly Flower among many Weedes or Grasse, eyther by their honourable authoritie, or eminence of learning or riches, whereby they excell others, and thereby may doe good to many. The frailty also of Mans life is learned by the soone fading of them before their flowring, or in their pride, or soone after, being either cropt by the hand of the spectator, or by a sudden blast withered and parched, or by the reuolution of time decaying of it owne nature: as also that the faireſt flowers or fruits first ripe, are sooneſt and first gathered. The mutabilitie also of states and persons, by this, that as where many goodly flowers & fruits did grow this yeare and age, in another they are quite pulled or digged vp, and eyther weedes and grasse grow in their place, or some building erected thereon, and their place is no more known. The Ciuill respects to be learned from them are many also: for the delight of the varieties both of formes, colours and properties of Herbes and Flowers, hath euer beene powerfull ouer dull, vnnurtured, rusticke and sauage people, led only by Natures instinct; how much more powerfull is it, or should be in the mindes of generous persons? for it may well bee said, he is not humane, that is not allured with this obiect. The study, knowledge, and trauel in them, as they haue been entertained of great Kings, Princes and Potentates, without disparagement to their Greatnesse, or hinderance to their more serious and weighty Affaires: so no douht vnto all that are capable thereof, it is not onely pleasant, but profitable, by comforting the minde, spirits and senses with an harmelesse delight, and by enabling the iudgement to conferre and apply helpe to many dangerous diseases. It is also an Instructer in the verity of the genuine Plants of the Ancients, and a Corrector of the many errours whereunto the world by continuance hath bin diuerted, and almost therein fixed, by eradicating in time, and by degrees, the pertinacious wilfulnesse of many, who because they were brought vp in their errours, are most vnwilling to leaue them without consideration of the good or euill, the right or wrong, they draw on therewith. And for my selfe I may well say, that had not mine owne paines and studies by a naturall inclination beene more powerfull in mee then any others helpe (although some through an euill disposition and ignorance haue so far traduced me as to say this was rather another mans worke then mine owne, but I leaue them to their folly) I had neuer done so much as I here publish; nor been fit or prepared for a larger, as time may suddenly (by Gods permission) bring to light, if the maleuolent dispositions of degenerate spirits doe not hinder the accomplishment.

But

*But perswading my selfe there is no showre that produceth not some fruit, or no word but worketh some effect, eyther of good to perswade, or of reproofe to euince, I could not but declare my minde herein, let others iudge or say what they please. For I haue alwaies held it a thing vnfit, to conceale or bury that knowledge God hath giuen, and not to impart it, and further others therewith as much as is conuenient, yet without ostentation, which I haue euer hated. Now further to informe the courteous Reader, both of the occasion that led me on to this worke, and the other occurrences to it. First, hauing perused many Herbals in Latine, I obserued that most of them haue eyther neglected or not knowne the many diuersities of the flower Plants, and rare fruits are known to vs at this time, and (except* Clusius) *haue made mention but of a very few. In English likewise we haue some extant, as* Turner *and* Dodonæus *translated, who haue haue said little of Flowers,* Gerard *who is last, hath no doubt giuen vs the knowledge of as many as he attained vnto in his time, but since his daies we haue had many more varieties, then he or they euer heard of, as may be perceiued by the store I haue here produced. And none of them haue particularly seuered those that are beautifull flower plants, fit to store a garden of delight and pleasure, from the wilde and vnfit: but haue enterlaced many, one among another, whereby many that haue desired to haue faire flowers, haue not known either what to choose, or what to desire. Diuers Bookes of Flowers also haue been set forth, some in our owne Countrey, and more in others, all which are as it were but handfuls snatched from the plentifull Treasury of Nature, none of them being willing or able to open all sorts, and declare them fully; but the greatest hinderance of all mens delight was, that none of them had giuen any description of them, but the bare name only. To satisfie therefore their desires that are louers of such Delights, I took vpon me this labour and charge, and haue here selected and set forth a Garden of all the chiefest for choyce, and faireſt for shew, from among all the seuerall Tribes and Kindreds of Natures beauty, and haue ranked them as neere as I could, or as the worke would permit, in affinity one vnto another. Secondly, and for their sakes that are studious in Authors, I haue set down the names haue bin formerly giuen vnto them, with some of their errours, not intending to cumber this worke with all that might bee said of them, because the deciding of the many controuersies, doubts, and questions that concerne them, pertaine more fitly to a generall History: yet I haue beene in some places more copious and ample then at the first I had intended, the occasion drawing on my desire to informe others with what I thought was fit to be known, reseruing what else might be said to another time & worke; wherein (God willing) I will inlarge my selfe, the subiect matter requiring it at my hands, in what my small ability can effect. Thirdly, I haue also to embellish this Worke set forth the figures of all such plants and flowers as are materiall and different one from another: but not as some others haue done, that is, a number of the figures of one sort of plant that haue nothing to distinguish them but the colour, for that I hold to be superfluous and waste. Fourthly, I haue also set down the Vertues and Properties of them in a briefe manner, rather desiring to giue you the knowledge of a few certaine and true, then to relate, as others haue done, a needlesse and false multiplicitie, that so there might as well profit as pleasure be taken from them, and that nothing might be wanting to accomplish it fully. And so much for this first part, my Garden of pleasant and delightfull Flowers. My next Garden consisteth of Herbes and Rootes, fit to be eaten of the rich and poor as nourishment and food, as sauce or condiment, as sallet or refreshing, for pleasure or profit; where I doe as well play the Gardiner, to shew you (in briefe, but not at large) the times*

*and*

*and manner of sowing, setting, planting, replanting, and the like (although all these things, and many more then are true, are set down very largely in the seuerall bookes that others haue written of this subiect) as also to shew some of the Kitchen vses (because they are Kitchen herbes &c.) although I confesse but very sparingly, not intending a treatise of cookery, but briefly to giue a touch thereof; and also the Physicall properties, to shew somewhat that others haue not set forth; yet not to play the Empericke, and giue you receipts of medicines for all diseases, but only to shew in some sort the qualities of Herbes, to quicken the minds of the studious. And lastly an Orchard of all sorts of domesticke or forraine, rare and good fruits, fit for this our Land and Countrey, which is at this time better stored and furnished then euer in any age before. I haue herein endeauoured, as in the other Gardens, to set forth the varieties of euery sort in as briefe a manner as possibly could be, without superfluous repetitions of descriptions, and onely with especiall notes of difference in leaues, flowers and fruits. Some few properties also are set downe, rather the chiefest then the most, as the worke did require. And moreouer before euery of these parts I haue giuen Treatises of the ordering, preparing and keeping the seuerall Gardens and Orchard, with whatsoeuer I thought was conuenient to be known for euery of them.*

*Thus haue I shewed you both the occasion and scope of this Worke, and herein haue spent my time, paines and charge, which if well accepted, I shall thinke well employed, and may the sooner hasten the fourth Part, A Garden of Simples; which will be quiet no longer at home, then that it can bring his Master newes of faire weather for the iourney.*

Thine in what he may,

IOHN PARKINSON.

Ioanni

Ioanni Parkinsono *Pharmacopoeo Londinensi solertissimo Botanico consummatissimo*
*T.D.M. S.P.D.*

POema panegyricum Opus tuum indefessi laboris, vtilitatis eximiæ postulat, & meriti iure à me extorqueret (mi Parkinsone) si fauentibus Musis, & secundo Apolline in bicipiti somniare Parnasso, & repentè Poetæ mihi prodire liceret. In fœtus tui bonis auibus in lucem editi, & prolixiorem nepotum seriem promittentis laudes, alii Deopleni Enthousiastæ carmine suos pangant elenchos; quos sub figmentis ampullata hyperbolicarum vocum mulcedine, vates ferè auribus mentibusue insinuant. Veritas nuditatis amans, fuco natiuum candorem obumbranti non illustranti perpetuum indixit bellum: In simplicitate, quam assertionum neruosa breuitas exprimit, exultat. Audi quid de te sentiam, Tu mihi sis in posterum Crateuas Brittannus; inter omnes, quotquot mihi hic innotuerunt, peritissimus, exercitatissimus, oculatissimus, & emunctissimæ naris Botanicus: Cuius opera in fortunata hac Insula rem herbariam tractari, emendari, augeri, & popularibus tuis vernaculo sermone ad amussim tradi, non decentiæ modo, sed etiam necessitatis est. Macte tua sedulitate (Vir optime) neque te laborum tam arduis lucubrationibus datorum hactenus pœniteat, vel deinceps impendendorum pigeat. Difficilia quæ pulchra. Leniet debitæ laudis dulcedo vigiliarum acerbitatem, & Olympicum stadium cito pede, à carceribus ad metas alacriter decurrentem nobile manet βραβεῖον. Sed memento Artem longam, Vitam esse breuem. Μηδὲν ἀναβαλλόμενος. Vide quid ad antiquum illum, cuius si non animam, saltem genium induisti, Crateuam scribat Hippocrates, Τέχνης πάσης ἀλλότριον ἀναβολὴ ἐπιτηδειοτάτη δὲ καὶ πάνυ; ἐν ᾗ ψυχῆς κίνδυνος ἢ ὑπέρθεσις. Nobilissimam Medicinæ partem Botanicam esse reputa. Floræ nunc litasti & Pomonæ, Apollini vt audio propediem HORTO MEDICO facturus. Amabò integræ Vestæ sacra conficito, eiusque variegatum multis simplicium morbifugorum myriadibus sinum absolutè pandito, quem sine velo nobis exhibeas. Nulla dies abeat sine linea. Sic tandem fructus gloriæ referes vberrimos, quos iustè sudoribus partos, vt in cruda & viridi senectute decerpas diu, iisque longum fruaris opto. *Vale. Datum Londini Calendas Octobris anno salutis* 1629.

Theodorus de Mayerne *Eques aurat. in Aula Regum Magnæ Britanniæ Iacobi & Caroli P. & F. Archiatrorum Comes.*

## Ad eximium arte & vſu Pharmacopæum & Botanographum *I. Parkinſonum.*

Gu. Turnerus. M. D.

Io. Gerardus Chirurgus.

*HErbarum vires, primus te (magne Britannæ)*
*Edocuit medicas, inclytus arte ſophus.*
*Atque cluens herbis alter, Chironis alumnus,*
*Deſcripſit plantas, neu cadat vlla ſalus.*
*Fortunate ſenex, ſis tu nunc tertius Heros*
*Hortos qui reſeras, deliciaſque ſoli,*
*Et flores Veneris lætos, herbaſque virentes,*
*Arboreos fætus, pharmacum & arte potens.*
*Poſteritas iuſtos poſthac tibi ſolvet honores,*
*Laudabitque tuæ dexteritatis opus.*

Ottuellus Meuerell. D.M. & Collegiæ Med. Lond. ſocius.

---

## Amico ſuo *Ioanni Parkinſono.*

*EXtollunt alij quos* (Parkinſone) *labores*
*Da mihi iam veniam comminuiſſe tuos.*
*Extremos poteris credi migraſſe per Indos:*
*Cum liber haud aliud quam tuus hortus hic eſt:*
*Ipſe habitare Indos tecum facis, haud petis Indos*
*I nunc, & tua me comminuiſſe refer.*
*Eſt liber Effigies, tuus hic qui pingitur hortus,*
*Digna manu facies hæc, facieque manus!*
*Vidi ego ſplendentem varigatis vndique gemmis*
*Vna fuit Salomon, turba quid ergo fuit?*
*Vt vario ſplendent Pallacia regia ſumptu,*
*Et Procerum turbis Atria tota nitent:*
*Tunc cum feſta dies veniam dedit eſſe ſuperbis*
*Quoſque ficus texit, nunc tria rura tegunt:*
*Plena tuo pariter ſpectatur Curia in Horto,*
*Hic Princeps, Dux hic, Sponſaque pulchra Ducis.*
*Quæque dies eſt feſta dies, nec parcius vnquam*
*Luxuriant, lauta hæc; Quotidiana tamen.*
*Ecce velut Patriæ Paradiſi haud immemor Exul,*
*Hunc naturali pingit amore ſibi.*
*Pingit & ad vivum ſub eodem nomine, & hic eſt*
*Fronticuli ſudor quem cerebrique dedit:*
*Aſtat Adam medius Paradiſo noſter in iſto*
*Et ſpecies nomen cuique dat ipſe ſuum.*
*Hos cape pro meritis, qui florem nomine donas*
*Æternum florens tu tibi Nomen habe.*

Guilielmus Atkins.

## Ad Amicum *Ioannem Parkinſonum* Pharmacopæum, & Archibotanicum Londinenſem.

*AFrica quas profert Plantas, quas India mittit,*
*Quas tua dat tellus, has tuus hortus habet:*
*Atque harum Species, florendi tempora, vires,*
*Et varias formas iſte libellus habet:*
*Neſcio plus librum talem mirabor, an hortum*
*Totus ineſt horto mundus; at iſte libro.*
Parkinſone *tuus liber, & labor, & tua ſit laus,*
*Herbas dum nobis das; datur herba tibi.*

Guilielmus Brodus Pharmacopæus
ac Philobotanicus Londinenſis.

---

## Ad Amicum *Ioannem Parkinſonum* Pharmacopæum & Botanicum inſignem. Carmen.

*QVam magno pandis Floræ penetralia nixu*
*Atque facis cœlo liberiore frui?*
*Omnibus vt placeas, ô quam propenſa voluntas,*
*Solicituſque labor nocte dieque premit?*
*Quam magno cultum ſtudio conquirere in hortum*
*Herbarum quicquid mundus in orbe tenet,*
*Immenſus ſumptus, multoſque extenſus in annos*
*Te labor afficiunt? & data nulla quies.*
*Talia quærenti, ſurgit novus ardor habendi,*
*Nec tibi tot ſoli munera magna petis;*
*Deſcriptos vivâ profers ſub imagine flores,*
*Tum profers menſæ quicquid & hortus alit,*
*Laudatos nobis fructus & promis honores,*
*Profers, quas celebrant nullibi ſcripta virum,*
*Herbarum ſpecies, quibus eſt quoque grata venuſtas;*
*Sic nos multiplici munere, Amice, beas.*
*Hoc cape pro meritis, florum dum gratia floret,*
*Suntque herbis vires; en tibi Nomen erit.*
*In ſerum ſemper tua gloria floreat ævum,*
*Gloria quæ in longum non peritura diem.*

Thomas Iohnſon vtriuſque
Societatis conſors.

ANNO AETATIS SVAE LXII EFFIGIES ANNO CIↃDCXXIX
THOMAS IOHNSON VIRTVTIS SOCIETATIS CONIVRA

# THE ORDERING OF THE GARDEN OF PLEASVRE.

## CHAP. I.

*The ſituation of a Garden of pleaſure, with the nature of ſoyles, and how to amend the defects that are in many ſorts of ſituations and grounds.*

THE ſeuerall ſituations of mens dwellings, are for the moſt part vnauoideable and vnremoueable; for moſt men cannot appoint forth ſuch a manner of ſituation for their dwelling, as is moſt fit to auoide all the inconueniences of winde and weather, but muſt bee content with ſuch as the place will afford them; yet all men doe well know, that ſome ſituations are more excellent than others: according therfore to the ſeuerall ſituation of mens dwellings, ſo are the ſituations of their gardens alſo for the moſt part. And although diuers doe diuerſly preferre their owne ſeuerall places which they haue choſen, or wherein they dwell; As ſome thoſe places that are neare vnto a riuer or brooke to be beſt for the pleaſantneſſe of the water, the eaſe of tranſportation of themſelues, their friends and goods, as alſo for the fertility of the ſoyle, which is ſeldome bad neare vnto a riuers ſide; And others extoll the ſide or top of an hill, bee it ſmall or great, for the proſpects ſake; And againe, ſome the plaine or champian ground, for the euen leuell thereof: euery one of which, as they haue their commodities accompanying them, ſo haue they alſo their diſcommodities belonging vnto them, according to the Latine Prouerbe, *Omne commodum fert ſuum incommodum.* Yet to ſhew you for euerie of theſe ſituations which is the fitteſt place to plant your garden in, and how to defend it from the iniuries of the cold windes and froſts that may annoy it, will, I hope, be well accepted. And firſt, for the water ſide, I ſuppoſe the North ſide of the water to be the beſt ſide for your garden, that it may haue the comfort of the South Sunne to lye vpon it and face it, and the dwelling houſe to bee aboue it, to defend the cold windes and froſts both from your herbes, and flowers, and early fruits. And ſo likewiſe I iudge for the hill ſide, that it may lye full open to the South Sunne, and the houſe aboue it, both for the comfort the ground ſhall receiue of the water and raine deſcending into it, and of defence from winter and colds. Now for the plaine leuell ground, the buildings of the houſe ſhould be on the North ſide of the garden, that ſo they might bee a defence of much ſufficiency to ſafeguard it from many iniurious cold nights and dayes, which elſe might ſpoyle the pride thereof in the bud. But becauſe euery one cannot ſo appoint his dwelling, as I here appoint the fitteſt place for it to be, euery ones pleaſure thereof ſhall be according to the ſite, coſt, and endeauours they beſtow, to cauſe it come neareſt to this proportion, by ſuch helpes of bricke or ſtone wals to defend it, or by the helpe of high growne and well ſpread trees, planted on the North ſide thereof, to keepe it the warmer. And euery of theſe three ſituations, hauing the faireſt buildings of the houſe facing the garden in this manner before ſpecified, beſides the benefit of ſhelter it ſhall haue from them, the buildings and roomes abutting thereon, ſhall haue reciprocally the beautifull proſpect into it, and haue both ſight and ſent of whatſoeuer is excellent, and worthy to giue content out from it, which is one of the greateſt pleaſures a garden can yeeld his Maſter. Now hauing ſhewed you the beſt place where this your

garden ſhould be, let me likewiſe aduiſe you where it ſhould not be, at leaſt that it is the worſt place wherein it may be, if it be either on the Weſt or Eaſt ſide of your houſe, or that it ſtand in a mooriſh ground, or other vnwholſome ayre (for many, both fruits, herbes, and flowers that are tender, participate with the ayre, taking in a manner their chiefeſt thriuing from thence) or neare any common Lay-ſtalles, or common Sewers, or elſe neare any great Brew-houſe, Dye-houſe, or any other place where there is much ſmoake, whether it be of ſtraw, wood, or eſpecially of ſea-coales, which of all other is the worſt, as our Citie of London can giue proofe ſufficient, wherein neither herbe nor tree will long proſper, nor hath done euer ſince the vſe of ſea-coales beganne to bee frequent therein. And likewiſe that it is much the worſe, if it bee neare vnto any Barnes or Stackes of corne or hey, becauſe that from thence will continually with the winde bee brought into the garden the ſtrawe and chaffe of the corne, the duſt and ſeede of the hey to choake or peſter it. Next vnto the place or ſituation, let mee ſhew you the grounds or ſoyles for it, eyther naturall or artificiall. No man will deny, but the naturall blacke mould is not only the fatteſt and richeſt, but farre exceedeth any other either naturall or artificiall, as well in goodneſſe as durability. And next thereunto, I hold the ſandy loame (which is light and yet firme, but not looſe as ſand, nor ſtiffe like vnto clay) to be little inferiour for this our Garden of pleaſure; for that it doth cauſe all bulbous and tuberous rooted plants to thriue ſufficiently therein, as likewiſe all other flower-plants, Roſes, Trees, &c. which if it ſhall decay by much turning and working out the heart of it, may ſoone be helped with old ſtable manure of horſes, being well turned in, when it is old and almoſt conuerted to mould. Other grounds, as chalke, ſand, grauell, or clay, are euery of them one more or leſſe fertill or barren than other; and therefore doe require ſuch helpes as is moſt fit for them. And thoſe grounds that are ouer dry, looſe, and duſtie, the manure of ſtall fedde beaſts and cattell being buried or trenched into the earth, and when it is thorough rotten (which will require twice the time that the ſtable ſoyle of horſes will) well turned and mixed with the earth, is the beſt ſoyle to temper both the heate and drineſſe of them. So contrariwiſe the ſtable dung of horſes is the beſt for cold grounds, to giue them heate and life. But of all other ſorts of grounds, the ſtiffe clay is the very worſt for this purpoſe; for that although you ſhould digge out the whole compaſſe of your Garden, carry it away, and bring other good mould in the ſtead thereof, and fill vp the place, yet the nature of that clay is ſo predominant, that in a ſmall time it will eate out the heart of the good mould, and conuert it to its owne nature, or very neare vnto it: ſo that to bring it to any good, there muſt bee continuall labour beſtowed thereon, by bringing into it good ſtore of chalke, lime, or ſand, or elſe aſhes eyther of wood or of ſea-coales (which is the beſt for this ground) well mixed and turned in with it. And as this ſtiffe clay is the worſt, ſo what ground ſoeuer commeth neareſt vnto the nature thereof, is neareſt vnto it in badneſſe, the ſignes whereof are the ouermuch moyſture thereof in Winter, and the much cleauing and chapping thereof in Summer, when the heate of the yeare hath conſumed the moyſture, which tyed and bound it faſt together, as alſo the ſtiffe and hard working therein: but if the nature of the clay bee not too ſtiffe, but as it were tempered and mixed with ſand or other earths, your old ſtable ſoyle of horſes will helpe well the ſmall rifting or chapping thereof, to be plentifully beſtowed therin in a fit ſeaſon. Some alſo do commend the caſting of ponds and ditches, to helpe to manure theſe ſtiffe chapping grounds. Other grounds, that are ouermoiſt by ſprings, that lye too neare the vpper face of the earth, beſides that the beds thereof had need to be laid vp higher, and the allies, as trenches and furrowes, to lye lower, the ground it ſelfe had neede to haue ſome good ſtore of chalke-ſtones beſtowed thereon, ſome certaine yeares, if it may be, before it be laid into a Garden, that the Winter froſts may breake the chalke ſmall, and the Raine diſſolue it into mould, that ſo they may bee well mixed together; than which, there is not any better manure to ſoyle ſuch a moiſt ground, to helpe to dry vp the moyſture, and to giue heate and life to the coldneſſe thereof, which doth alwayes accompany theſe moiſt grounds, and alſo to cauſe it abide longer in heart than any other. For the ſandy and grauelly grounds, although I know the well mollified manure of beaſts and cattell to be excellent good, yet I know alſo, that ſome commend a white Marle, and ſome a clay to be well ſpread thereon, and after turned thereinto: and for the chalkie ground, *è conuerſo*, I commend fatte clay to helpe it. You muſt vnderſtand, that the leſſe rich or more barren that your ground is, there nee-

deth

deth the more care, labour, and coſt to bee beſtowed thereon, both to order it rightly, & ſo to preſerue it from time to time: for no artificiall or forc't ground can endure good any long time, but that within a few yeares it muſt be refreſhed more or leſſe, according as it doth require. Yet you ſhall likewiſe vnderſtand, that this Garden of pleaſure ſtored with theſe Out-landiſh flowers; that is, bulbous and tuberous rooted plants, and other fine flowers, that I haue hereafter deſcribed, and aſſigned vnto it, needeth not ſo much or ſo often manuring with ſoyle, &c. as another Garden planted with the other ſorts of Engliſh flowers, or a Garden of ordinary Kitchin herbes doth. Your ground likewiſe for this Garden had neede to bee well cleanſed from all annoyances (that may hinder the well doing or proſpering of the flowers therein) as ſtones, weedes, rootes of trees, buſhes, &c. and all other things cumberſome or hurtfull; and therefore the earth being not naturally fine enough of it ſelfe, is vſed to bee ſifted to make it the finer, and that either through a hurdle made of ſticks, or lathes, or through ſquare or round ſieues platted with fine and ſtrong thin ſtickes, or with wyers in the bottome. Or elſe the whole earth of the Garden being courſe, may be caſt in the ſame manner that men vſe to try or fine ſand from grauell, that is, againſt a wall; whereby the courſer and more ſtony, falling downe from the fine, is to be taken away from the foote of the heape, the finer ſand and ground remaining ſtill aboue, and on the heape. Or elſe in the want of a wall to caſt it againſt, I haue ſeene earth fined by it ſelfe in this manner: Hauing made the floore or vpper part of a large plat of ground cleane from ſtones, &c. let there a reaſonable round heape of fine earth be ſet in the midſt thereof, or in ſtead thereof a large Garden flower-pot, or other great pot, the bottome turned vpwards, and then poure your courſe earth on the top or head thereof, one ſhouell full after another ſomewhat gently, and thereby all the courſe ſtuffe and ſtones will fall downe to the bottome round about the heape, which muſt continually be carefully taken away, and thus you may make your earth as fine as if it were caſt againſt a wall, the heape being growne great, ſeruing in ſtead thereof. Thoſe that will not prepare their grounds in ſome of theſe manners aforeſaid, ſhall ſoone finde to their loſſe the neglect thereof: for the traſh and ſtones ſhall ſo hinder the encreaſe of their roots, that they will be halfe loſt in the earth among the ſtones, which elſe might be ſaued to ſerue to plant whereſoeuer they pleaſe.

## Chap. II.

### *The frame or forme of a Garden of delight and pleaſure, with the ſeuerall varieties thereof.*

ALthough many men muſt be content with any plat of ground, of what forme or quantity ſoeuer it bee, more or leſſe, for their Garden, becauſe a more large or conuenient cannot bee had to their habitation: Yet I perſwade my ſelfe, that Gentlemen of the better ſort and quality, will prouide ſuch a parcell of ground to bee laid out for their Garden, and in ſuch conuenient manner, as may be fit and anſwerable to the degree they hold. To preſcribe one forme for euery man to follow, were too great preſumption and folly: for euery man will pleaſe his owne fancie, according to the extent he deſigneth out for that purpoſe, be it orbicular or round, triangular or three ſquare, quadrangular or foure ſquare, or more long than broad. I will onely ſhew you here the ſeuerall formes that many men haue taken and delighted in, let euery man chuſe which him liketh beſt, or may moſt fitly agree to that proportion of ground hee hath ſet out for that purpoſe. The orbicular or round forme is held in it owne proper exiſtence to be the moſt abſolute forme, containing within it all other formes whatſoeuer; but few I thinke will chuſe ſuch a proportion to be ioyned to their habitation, being not accepted any where I think, but for the generall Garden to the Vniuerſity at Padoa. The triangular or three ſquare is ſuch a forme alſo, as is ſeldome choſen by any that may make another choiſe, and as I thinke is onely had where another forme cannot be had, neceſſitie conſtraining them to be therewith content. The foure ſquare forme is the moſt vſually accepted with all, and doth beſt agree to any mans dwelling, being (as I ſaid before) behinde the houſe, all the backe windowes thereof opening into it. Yet if it bee longer than the breadth, or broader than the length, the proportion of walkes, ſquares, and knots may be ſoon brought to the ſquare forme, and be ſo caſt, as the beauty thereof may



bee no lesse than the foure square proportion, or any other better forme, if any be. To forme it therfore with walks, crosse the middle both waies, and round about it also with hedges, with squares, knors and trayles, or any other worke within the foure square parts, is according as euery mans conceit alloweth of it, and they will be at the charge: For there may be therein walkes eyther open or close, eyther publike or priuate, a maze or wildernesse, a rocke or mount, with a fountaine in the midst thereof to conuey water to euery part of the Garden, eyther in pipes vnder the ground, or brought by hand, and emptied into large Cisternes or great Turkie Iarres, placed in conuenient places, to serue as an ease to water the neerest parts thereunto. Arbours also being both gracefull and necessary, may be appointed in such conuenient places, as the corners, or else where, as may be most fit, to serue both for shadow and rest after walking. And because many are desirous to see the formes of trayles, knots, and other compartiments, and because the open knots are more proper for these Out-landish flowers; I haue here caused some to be drawne, to satisfie their desires, not intending to cumber this worke with ouer manie; in that it would be almost endlesse, to expresse so many as might bee conceiued and set downe, for that euery man may inuent others farre differing from these, or any other can be set forth. Let euery man therefore, if hee like of these, take what may please his mind, or out of these or his own conceit, frame any other to his fancy, or cause others to be done as he liketh best, obseruing this *decorum*, that according to his ground he do cast out his knots, with conuenient roome for allies and walkes; for the fairer and larger your allies and walkes be, the more grace your Garden shall haue, the lesse harme the herbes and flowers shall receiue, by passing by them that grow next vnto the allies sides, and the better shall your Weeders cleanse both the beds and the allies.

## Chap. III.

*The many sorts of herbes and other things, wherewith the beds and parts of knots are bordered to set out the forme of them, with their commodities and discommodities.*

IT is necessary also, that I shew you the seuerall materials, wherewith these knots and trayles are set forth and bordered; which are of two sorts: The one are liuing herbes, and the other are dead materials; as leade, boords, bones, tyles, &c. Of herbes, there are many sorts wherewith the knots and beds in a Garden are vsed to bee set, to shew forth the forme of them, and to preserue them the longer in their forme, as also to be as greene, and sweete herbes, while they grow, to be cut to perfume the house, keeping them in such order and proportion, as may be most conuenient for their seuerall natures, and euery mans pleasure and fancy: Of all which, I intend to giue you the knowledge here in this place; and first, to begin with that which hath beene most anciently receiued, which is Thrift. This is an euerliuing greene herbe, which many take to border their beds, and set their knots and trayles, and therein much delight, because it will grow thicke and bushie, and may be kept, being cut with a paire of Garden sheeres, in some good handsome manner and proportion for a time, and besides, in the Summer time send forth many short stalkes of pleasant flowers, to decke vp an house among other sweete herbes: Yet these inconueniences doe accompany it; it will not onely in a small time ouergrow the knot or trayle in many places, by growing so thicke and bushie, that it will put out the forme of a knot in many places: but also much thereof will dye with the frosts and snowes in Winter, and with the drought in Summer, whereby many voide places will be seene in the knot, which doth much deforme it, and must therefore bee yearely refreshed: the thickenesse also and bushing thereof doth hide and shelter snayles and other small noysome wormes so plentifully, that Gilloflowers, and other fine herbes and flowers being planted therein, are much spoyled by them, and cannot be helped without much industry, and very great and daily attendance to destroy them. Germander is another herbe, in former times also much vsed, and yet also in many places; and because it will grow thicke, and may be kept also in some forme and proportion with cutting, and that the cuttings are much vsed as a strawing herbe for houses, being pretty and sweete, is also much affected by diuers: but this also will often dye and grow out of forme, and besides that, the stalkes will grow too great, hard and stubby, the rootes doe so farre shoote vnder ground, that vpon a little continuance thereof, will

ſpread into many places within the knot, which if continually they be not plucked vp, they will ſpoile the whole knot it ſelfe; and therefore once in three or foure yeares at the moſt, it muſt be taken vp and new ſet, or elſe it will grow too royniſh and cumberſome. Hyſſope hath alſo been vſed to be ſet about a knot, and being ſweet, will ſerue for ſtrewings, as Germander: But this, although the rootes doe not runne or creep like it, yet the ſtalkes doe quickly grow great aboue ground, and dye often after the firſt yeares ſetting, whereby the grace of the knot will be much loſt. Marierome, Sauorie, and Thyme, in the like manner being ſweete herbes, are vſed to border vp beds and knors, and will be kept for a little while, with cutting, into ſome conformity; but all and euery of them ſerue moſt commonly but for one yeares vſe, and will ſoone decay and periſh: and therefore none of theſe, no more than any of the former, doe I commend for a good bordering herbe for this purpoſe. Lauander Cotton alſo being finely ſlipped and ſet, is of many, and thoſe of the higheſt reſpect of late daies, accepted, both for the beauty and forme of the herbe, being of a whitiſh greene mealy colour, for his ſent ſmelling ſomewhat ſtrong, and being euerliuing and abiding greene all the Winter, will, by cutting, be kept in as euen proportion as any other herbe may be. This will likewiſe ſoone grow great and ſtubbed, notwithſtanding the cutting, and beſides will now and then periſh in ſome places, eſpecially if you doe not ſtrike or put off the ſnow, before the Sunne lying vpon it diſſolue it: The rarity & nouelty of this herbe, being for the moſt part but in the Gardens of great perſons, doth cauſe it to be of the greater regard, it muſt therfore be renewed wholly euery ſecond or third yeare at the moſt, becauſe of the great growing therof. Slips of Iuniper or Yew are alſo receiued of ſome & planted, becauſe they are alwayes green, and that the Iuniper eſpecially hath not that ill ſent that Boxe hath, which I will preſently commend vnto you, yet both Iuniper and Yew will ſoon grow too great and ſtubbed, and force you to take vp your knot ſooner, than if it were planted with Boxe. Which laſtly, I chiefly and aboue all other herbes commend vnto you, and being a ſmall, lowe, or dwarfe kinde, is called French or Dutch Boxe, and ſerueth very well to ſet out any knot, or border out any beds: for beſides that it is euer greene, it being reaſonable thicke ſet, will eaſily be cut and formed into any faſhion one will, according to the nature thereof, which is to grow very ſlowly, and will not in a long time riſe to be of any height, but ſhooting forth many ſmall branches from the roote, will grow very thicke, and yet not require ſo great tending, nor ſo much periſh as any of the former, and is onely receiued into the Gardens of thoſe that are curious. This (as I before ſaid) I commend and hold to bee the beſt and ſureſt herbe to abide faire and greene in all the bitter ſtormes of the ſharpeſt Winter, and all the great heates and droughts of Summer, and doth recompence the want of a good ſweet ſent with his freſh verdure, euen proportion, and long laſting continuance. Yet theſe inconueniences it hath, that beſides the vnpleaſing ſent which many miſlike, and yet is but ſmall, the rootes of this Boxe do ſo much ſpread themſelues into the ground of the knot, and doe draw from thence ſo much nouriſhment, that it robbeth all the herbes that grow neare it of their ſap and ſubſtance, thereby making all the earth about it barren, or at leaſt leſſe fertile. Wherefore to ſhew you the remedy of this inconuenience of ſpreading, without either taking vp the Boxe of the border, or the herbes and flowers in the knot, is I thinke a ſecret knowne but vnto a few, which is this: You ſhall take a broad pointed Iron like vnto a Sliſe or Cheſſill, which thruſt downe right into the ground a good depth all along the inſide of the border of Boxe ſomewhat cloſe thereunto, you may thereby cut away the ſpreading rootes thereof, which draw ſo much moiſture from the other herbes on the inſide, and by this meanes both preſerue your herbes and flowers in the knot, and your Boxe alſo, for that the Boxe will be nouriſhed ſufficiently from the reſt of the rootes it ſhooteth on all the other ſides. And thus much for the liuing herbes, that ſerue to ſet or border vp any knot. Now for the dead materials, they are alſo, as I ſaid before diuers: as firſt, Leade, which ſome that are curious doe border their knots withall, cauſing it to be cut of the breadth of foure fingers, bowing the lower edge a little outward, that it may lye vnder the vpper cruſt of the ground, and that it may ſtand the faſter, and making the vpper edge either plain, or cut out like vnto the battlements of a Church: this faſhion hath delighted ſome, who haue accounted it ſtately (at the leaſt coſtly) and fit for their degree, and the rather, becauſe it will be bowed and bended into any round ſquare, angular, or other proportion as one liſteth, and is not much to be miſliked, in that the Leade

doth

doth not easily breake or spoile without much iniury, and keepeth vp a knot for a very long time in his due proportion: but in my opinion, the Leade is ouer-hot for Summer, and ouer-cold for Winter. Others doe take Oaken inch boords, and sawing them foure or fiue inches broad, do hold vp their knot therewith: but in that these boordes cannot bee drawne compasse into any small scantling, they must serue rather for long outright beds, or such knots as haue no rounds, halfe rounds, or compassings in them. And besides, these boordes are not long lasting, because they stand continually in the weather, especially the ends where they are fastned together will sooneſt rot and perish, and so the whole forme will be spoyled. To preuent that fault, some others haue chosen the shanke bones of Sheep, which after they haue beene well cleansed and boyled, to take out the fat from them, are stucke into the ground the small end downewards, and the knockle head vpwards, and thus being set side to side, or end to end close together, they set out the whole knot therewith, which heads of bones although they looke not white the first yeare, yet after they haue abiden some frosts and heates will become white, and prettily grace out the ground: but this inconuenience is incident to them, that the Winter frosts will raise them out of the ground oftentimes, and if by chance the knockle head of any doe breake, or be strucke off with any ones foot, &c. going by, from your store, that lyeth by you of the same sort, set another in the place, hauing first taken away the broken peece: although these will last long in forme and order, yet because they are but bones many mislike them, and indeed I know but few that vse them. Tyles are also vsed by some, which by reason they may bee brought compasse into any fashion many are pleased with them, who doe not take the whole Tyle at length, but halfe Tyles, and other broken peeces set somewhat deepe into the ground, that they may stand fast, and these take vp but little roome, and keepe vp the edge of the beds and knots in a pretty comely manner, but they are often out of frame, in that many of them are broken and spoiled, both with mens feete passing by, the weather and weight of the earth beating them downe and breaking them, but especially the frosts in Winter doe so cracke off their edges, both at the toppes and sides that stand close one vnto another, that they must bee continually tended and repaired, with fresh and sound ones put in the place of them that are broken or decayed. And lastly (for it is the latest inuention) round whitish or blewish pebble stones, of some reasonable proportion and bigneſſe, neither too great nor too little, haue beene vsed by some to be set, or rather in a manner but laide vpon the ground to fashion out the traile or knot, or all along by the large grauelly walke sides to set out the walke, and maketh a pretty handsome shew, and because the stones will not decay with the iniuries of any time or weather, and will be placed in their places againe, if any should be thrust out by any accident, as also that their sight is so conspicuous vpon the ground, especially if they be not hid with the store of herbes growing in the knot; is accounted both for durability, beauty of the sight, handsomneſſe in the worke, and ease in the working and charge, to be of all other dead materials the chiefest. And thus, Gentlemen, I haue shewed you all the varieties that I know are vsed by any in our Countrey, that are worth the reciting (but as for the fashion of Iawe-bones, vsed by some in the Low-Countries, and other places beyond the Seas, being too grosse and base, I make no mention of them) among which euery one may take what pleaseth him best, or may most fitly be had, or may best agree with the ground or knot. Moreouer, all these herbes that serue for borderings, doe serue as well to be set vpon the ground of a leuelled knot; that is, where the allies and foot-pathes are of the same leuell with the knot, as they may serue also for the raised knot, that is, where the beds of the knot are raised higher than the allies: but both Leade, Boordes, Bones, and Tyles, are only for the raised ground, be it knot or beds. The pebble stones againe are onely for the leuelled ground, because they are so shallow, that as I said before, they rather lye vpon the earth than are thrust any way into it. All this that I haue here set downe, you must vnderstand is proper for the knots alone of a Garden. But for to border the whole square or knot about, to serue as a hedge thereunto, euery one taketh what liketh him best; as either Priuet alone, or sweete Bryer, and white Thorne enterlaced together, and Roses of one, or two, or more sorts placed here and there amongst them. Some also take Lauander, Rosemary, Sage, Southernwood, Lauander Cotton, or some such other thing. Some againe plant Cornell Trees, and plash them, or keepe them lowe, to

forme

forme them into an hedge. And some againe take a lowe prickly shrubbe, that abideth alwayes greene, described in the end of this Booke, called in Latine *Pyracantha*, which in time will make an euer greene hedge or border, and when it beareth fruit, which are red berries like vnto Hawthorne berries, make a glorious shew among the greene leaues in the Winter time, when no other shrubbes haue fruit or leaues.

## Chap. IV.

### *The nature and names of diuers Out-landish flowers, that for their pride, beauty, and earlinesse, are to be planted in Gardens of pleasure for delight.*

HAuing thus formed out a Garden, and diuided it into his fit and due proportion, with all the gracefull knots, arbours, walkes, &c. likewise what is fit to keepe it in the same comely order, is appointed vnto it, both for the borders of the squares, and for the knots and beds themselues; let vs now come and furnish the inward parts, and beds with those fine flowers that (being strangers vnto vs, and giuing the beauty and brauery of their colours so early before many of our owne bred flowers, the more to entice vs to their delight) are most beseeming it: and namely, with Daffodils, Fritillarias, Iacinthes, Saffron-flowers, Lillies, Flowerdeluces, Tulipas, Anemones, French Cowslips, or Beares eares, and a number of such other flowers, very beautifull, delightfull, and pleasant, hereafter described at full, whereof although many haue little sweete sent to commend them, yet their earlinesse and exceeding great beautie and varietie doth so farre counteruaile that defect (and yet I must tell you with all, that there is among the many sorts of them some, and that not a few, that doe excell in sweetnesse, being so strong and heady, that they rather offend by too much than by too little sent, and some againe are of so milde and moderate temper, that they scarce come short of your most delicate and dantiest flowers) that they are almost in all places with all persons, especially with the better sort of the Gentry of the Land, as greatly desired and accepted as any other the most choisest, and the rather, for that the most part of these Out-landish flowers, do shew forth their beauty and colours so early in the yeare, that they seeme to make a Garden of delight euen in the Winter time, and doe so giue their flowers one after another, that all their brauery is not fully spent, vntil that Gilliflowers, the pride of our English Gardens, do shew themselues: So that whosoeuer would haue of euery sort of these flowers, may haue for euery moneth seuerall colours and varieties, euen from Christmas vntill Midsommer, or after; and then, after some little respite, vntill Christmas againe, and that in some plenty, with great content and without forcing; so that euery man may haue them in euery place, if they will take any care of them. And because there bee many Gentlewomen and others, that would gladly haue some fine flowers to furnish their Gardens, but know not what the names of those things are that they desire, nor what are the times of their flowring, nor the skill and knowledge of their right ordering, planting, displanting, transplanting, and replanting; I haue here for their sakes set downe the nature, names, times, and manner of ordering in a briefe manner, referring the more ample declaration of them to the worke following. And first of their names and natures: Of Daffodils there are almost an hundred sorts, as they are seuerally described hereafter, euery one to be distinguished from other, both in their times, formes, and colours, some being eyther white, or yellow, or mixt, or else being small or great, single or double, and some hauing but one flower vpon a stalke, others many, whereof many are so exceeding sweete, that a very few are sufficient to perfume a whole chamber, and besides, many of them be so faire and double, eyther one vpon a stalke, or many vpon a stalke, that one or two stalkes of flowers are in stead of a whole nose-gay, or bundell of flowers tyed together. This I doe affirme vpon good kuowledge and certaine experience, and not as a great many others doe, tell of the wonders of another world, which themselues neuer saw nor euer heard of, except some superficiall relation, which themselues haue augmented according to their owne fansie and conceit. Againe, let me here also by the way tell you, that many idle and ignorant Gardiners and others, who get names by stealth, as they doe many other things, doe call

some

some of these Daffodils Narcisses, when as all know that know any Latine, that Narcissus is the Latine name, and Daffodill the English of one and the same thing; and therefore alone without any other Epithite cannot properly distinguish seuerall things. I would willingly therefore that all would grow iudicious, and call euery thing by his proper English name in speaking English, or else by such Latine name as euery thing hath that hath not a proper English name, that thereby they may distinguish the seuerall varieties of things and not confound them, as also to take away all excuses of mistaking; as for example: The single English bastard Daffodill (which groweth wilde in many Woods, Groues, and Orchards in England.) The double English bastard Daffodill. The French single white Daffodill many vpon a stalke. The French double yellow Daffodill. The great, or the little, or the least Spanish yellow bastard Daffodill, or the great or little Spanish white Daffodill. The Turkie single white Daffodill, or, The Turkie single or double white Daffodill many vpon a stalke, &c. Of Fritillaria, or the checkerd Daffodill, there are halfe a score seuerall sorts, both white and red, both yellow and blacke, which are a wonderfull grace and ornament to a Garden in regard of the Checker like spots are in the flowers. Of Iacinthes there are aboue halfe an hundred sorts, as they are specified hereafter; some like vnto little bells or starres, others like vnto little bottles or pearles, both white and blew, sky-coloured and blush, and some starlike of many pretty various formes, and all to giue delight to them that will be curious to obserue them. Of Crocus or Saffron flowers, there are also twenty sorts; some of the Spring time, others flowring onely in the Autume or Fall, earlier or later than another, some whereof abide but a while, others indure aboue a moneth in their glorious beauty. The Colchicum or Medowe Saffron, which some call the sonne before the father, but not properly, is of many sorts also; some flowring in the Spring of the yeare, but the most in Autume, whereof some haue faire double flowers very delightfull to behold, and some party coloured both single and double so variable, that it would make any one admire the worke of the Creatour in the various spots and stripes of these flowers. Then haue wee of Lillies twenty seuerall sorts and colours, among whom I must reckon the Crowne Imperiall, that for his stately forme deserueth some speciall place in this Garden, as also the Martagons, both white and red, both blush and yellow, that require to be set by themselues apart, as it were in a small round or square of a knot, without many other, or tall flowers growing neare them. But to tell you of all the sorts of Tulipas (which are the pride of delight) they are so many, and as I may say, almost infinite, doth both passe my ability, and as I beleeue the skill of any other. They are of two especiall sorts, some flowring earlier, and others later than their fellowes, and that naturally in all grounds, wherein there is such a wonderfull variety and mixture of colours, that it is almost impossible for the wit of man to descipher them thoroughly, and to giue names that may be true & seuerall distinctions to euery flower, threescore seuerall sorts of colours simple and mixed of each kind I can reckon vp that I haue, and of especiall note, and yet I doubt not, but for euery one of them there are ten others differing from them, which may be seen at seuerall times, and in seuerall places: & besides this glory of variety in colors that these flowers haue, they carry so stately & delightfull a forme, & do abide so long in their brauery (enduring aboue three whole moneths from the first vnto the last) that there is no Lady or Gentlewoman of any worth that is not caught with this delight, or not delighted with these flowers. The Anemones likewise or Windeflowers are so full of variety and so dainty, so pleasant and so delightsome flowers, that the sight of them doth enforce an earnest longing desire in the minde of any one to be a possessour of some of them at the least: For without all doubt, this one kinde of flower, so variable in colours, so differing in forme (being almost as many sorts of them double as single) so plentifull in bearing flowers, and so durable in lasting, and also so easie both to preserue and to encrease, is of it selfe alone almost sufficient to furnish a garden with their flowers for almost halfe the yeare, as I shall shew you in a fit and conuenient place. The Beares eares or French Cowslips must not want their deserued commendations, seeing that their flowers, being many set together vpon a stalke, doe seeme euery one of them to bee a Nosegay alone of it selfe; and besides the many differing colours that are to be seene in them, as white, yellow, blush, purple, red, tawney, murrey, haire colour, &c. which encrease much delight in all sorts of the Gentry of the Land, they are not vnfurnished with a pretty sweete sent;

CHAP.

which

which doth adde an encreaſe of pleaſure in thoſe that make them an ornament for their wearing. Flowerdeluces alſo are of many ſorts, but diuided into two eſpeciall kindes; the one bearing a leafe like a flagge, whoſe rootes are tuberous, thicke and ſhort (one kinde of them being the Orris rootes that are ſold at the Apothecaries, whereof ſweete powders are made to lye among garments) the other hauing round rootes like vnto Onions, and narrow long leaues ſomewhat like graſſe: Of both theſe kindes there is much variety, eſpecially in their colours. The greater Flagge kinde is frequent enough and diſperſed in this Land, and well doth ſerue to decke vp both a Garden and Houſe with natures beauties: But the chiefe of all is your Sable flower, ſo fit for a mourning habit, that I thinke in the whole compaſſe of natures ſtore, there is not a more patheticall, or of greater correſpondency, nor yet among all the flowers I know any one comming neare vnto the colour of it. The other kinde which hath bulbous or Onion like rootes, diuerſifieth it ſelfe alſo into ſo many fine colours, being of a more neate ſhape and ſuccinct forme than the former, that it muſt not bee wanting to furniſh this Garden. The Hepatica or Noble Liuerwoort is another flower of account, whereof ſome are white, others red, or blew, or purple, ſomewhat reſembling Violets, but that there are white threads in the middeſt of their flowers, which adde the more grace vnto them; and one kinde of them is ſo double, that it reſembleth a double thicke Daſie or Marigold, but being ſmall and of an excellent blew colour, is like vnto a Button: but that which commendeth the flower as much as the beauty, is the earlineſſe in flowring, for that it is one of the very firſt flowers that open themſelues after Chriſtmas, euen in the midſt of Winter. The Cyclamen or Sowebread is a flower of rare receipt, becauſe it is naturally hard to encreaſe, and that the flowers are like vnto red or bluſh coloured Violets, flowring in the end of Summer or beginning of Autumne: the leaues likewiſe hereof haue no ſmall delight in their pleaſant colour, being ſpotted and circled white vpon greene, and that which moſt prefereth it, is the Phyſicall properties thereof for women, which I will declare when I ſhall ſhew you the ſeuerall deſcriptions of the varieties in his proper place. Many other ſorts of flowers there are fit to furniſh this Garden, as Leucoium or Bulbous Violet, both early and late flowring. Muſcari or Muske Grape flower. Starre flowers of diuers ſorts. Phalangium or Spiderwort, the chiefe of many is that ſort whoſe flowers are like vnto a white Lilly. Winter Crowfoote or Wolfes bane. The Chriſtmas flower like vnto a ſingle white Roſe. Bell flowers of many kindes. Yellow Larkes ſpurre, the prettieſt flower of a ſcore in a Garden. Flower-gentle or Floramour. Flower of the Sunne. The Maruaile of Peru or of the world. Double Marſh Marigold or double yellow Buttons, much differing and farre exceeding your double yellow Crowfoote, which ſome call ~~Batchelours Buttons.~~ Double French Marigolds that ſmell well, and is a greater kinde than the ordinary, and farre ſurpaſſeth it. The double red Ranunculus or Crowfoote (farre excelling the moſt glorious double Anemone) and is like vnto our great yellow double Crowfoote. Thus hauing giuen you the knowledge of ſome of the choiſeſt flowers for the beds of this Garden, let me alſo ſhew you what are fitteſt for your borders, and for your arbours. The Iaſmine white and yellow. The double Honyſockle. The Ladies Bower, both white, and red, and purple ſingle and double, are the fitteſt of Outlandiſh plants to ſet by arbours and banqueting houſes, that are open, both before and aboue to helpe to couer them, and to giue both ſight, ſmell, and delight. The ſorts of Roſes are fitteſt for ſtandards in the hedges or borders. The Cherry Bay or Lauroceraſus. The Roſe Bay or Oleander. The white and the blew Syringa or Pipe tree, are all gracefull and delightfull to ſet at ſeuerall diſtances in the borders of knots; for ſome of them giue beautifull and ſweete flowers. The Pyracantha or Prickly Corall tree doth remaine with greene leaues all the yeare, and may be plaſhed, or laid downe, or tyed to make a fine hedge to border the whole knot, as is ſaid before. The Wilde Bay or Laurus Tinus, doth chiefly deſire to be ſheltered vnder a wall, where it will beſt thriue, and giue you his beautifull flowers in Winter for your delight, in recompence of his fenced dwelling. The Dwarfe Bay or Meſereon, is moſt commonly either placed in the midſt of a knot, or at the corners thereof, and ſometimes all along a walke for the more grace. And thus to fit euery ones fancy, I haue ſhewed you the variety of natures ſtore in ſome part for you to diſpoſe of them to your beſt content.

CHAP.

## Chap. V.

### *The nature and names of those that are called vsually English flowers.*

THose flowers that haue beene vsually planted in former times in Gardens of this Kingdome (when as our forefathers knew few or none of those that are recited before) haue by time and custome attained the name of English flowers, although the most of them were neuer naturall of this our Land, but brought in from other Countries at one time or other, by those that tooke pleasure in them where they first saw them: and I doubt not, but many other sorts than here are set downe, or now knowne to vs, haue beene brought, which either haue perished by their negligence or want of skill that brought them, or else because they could not abide our cold Winters; those onely remaining with vs that haue endured of themselues, and by their encreasing haue beene distributed ouer the whole Land. If I should make any large discourse of them, being so well knowne to all, I doubt I should make a long tale to small purpose: I will therefore but briefly recite them, that you may haue them together in one place, with some little declaration of the nature and quality of them, and so passe to other matters. And first of Primroses and Cowslips, whereof there are many prettie varieties; some better knowne in the West parts of this Kingdome, others in the North, than in any other, vntill of late being obserued by some curious louers of varieties, they haue been transplanted diuersly, and so made more common: for although we haue had formerly in these parts about London greene Primroses vsually, yet we neuer saw or heard of greene Cowslips both single and double but of late dayes, and so likewise for Primroses to be both single and double from one roote, and diuers vpon one stalke of diuers fashions, I am sure is not vsuall: all which desire rather to bee planted vnder some hedge, or fence, or in the shade, than in the Sunne. Single Rose Campions, both white, red, and blush, and the double red Rose Campion also is knowne sufficiently, and will abide moderate Sunne as well as the shade. The flower of Bristow or None-such is likewise another kinde of Campion, whereof there is both white flowring plants and blush as well as Orange colour, all of them being single flowers require a moderate Sunne and not the shadow: But the Orange colour None-such with double flowers, as it is rare and not common, so for his brauery doth well deserue a Master of account that will take care to keepe and preserue it. Batchelours Buttons both white and red, are kindes of wilde Campions of a very double forme, and will reasonably well like the Sunne but not the shade. Wall-flowers are common in euery Garden, as well the ordinary double as the single, and the double kinde desireth no more shade than the single, but the greater kindes both double and single must haue the Sunne. Stock-Gilloflowers likewise are almost as common as Wall-flowers, especially the single kindes in euery womans Garden, but the double kindes are much more rare, and possessed but of a few, and those onely that will bee carefull to preserue them in Winter; for besides that the most of them are more tender, they yeeld no seede as the single kindes doe to preserue them, although one kinde from the sowing of the seed yeeld double flowers: They will all require the comfort of the Sunne, especially the double kindes, and to be defended from cold, yet so as in the Summer they doe not want water wherein they much ioy, and which is as it were their life. Queenes Gilloflowers (which some call Dames Violets, and some Winter Gilloflowers, are a kinde of Stock-Gilloflower) planted in Gardens to serue to fill vp the parts thereof for want of better things, hauing in mine opinion neither sight nor sent much to commend them. Violets are the Springs chiefe flowers for beauty, smell, and vse, both single and double, the more shadie and moist they stand the better. Snapdragon are flowers of much more delight, and in that they are more tender to keep, and will hardly endure the sharpe Winters, vnlesse they stand well defended, are scarce seene in many Gardens. Columbines single and double, of many sorts, fashions, and colours, very variable both speckled and party coloured, are flowers of that respect, as that no Garden would willingly bee without them, that could tell how to haue them, yet the rarer the flowers are, the more trouble to keepe; the ordinary sorts on the contrary

trary part will not be loſt, doe what one will. Larkes heeles, or ſpurres, or toes, as in ſeuerall Countries they are called, exceed in the varietie of colours, both ſingle and double, any of the former times; for vntill of late dayes none of the moſt pleaſant colours were ſeene or heard of: but now the ſingle kindes are reaſonable well diſperſt ouer the Land, yet the double kindes of all thoſe pleaſant colours (and ſome other alſo as beautifull) which ſtand like little double Roſes, are enioyed but of a few: all of them riſe from ſeed, and muſt be ſowne euery yeare, the double as well as the ſingle. Panſyes or Hartes eaſes of diuers colours, and although without ſent, yet not without ſome reſpect and delight. Double Poppies are flowers of a great and goodly proportion, adorning a Garden with their variable colours to the delight of the beholders, wherein there is ſome ſpeciall care to be taken, leſt they turne ſingle; and that is, if you ſee them grow vp too thicke, that you muſt pull them vp, and not ſuffer them to grow within leſſe than halfe a yard diſtance, or more one from another. Double Daiſies are flowers not to be forgotten, although they be common enough in euery Garden, being both white and red, both bluſh and ſpeckled, or party coloured, beſides that which is called Iacke an Apes on horſebacke, they require a moiſt and ſhadowie place; for they are ſcorched away, if they ſtand in the Sunne in any dry place. Double Marigolds alſo are the moſt common in all Gardens. And ſo are the French Marigolds that haue a ſtrong heady ſent, both ſingle and double, whoſe glorious ſhew for colour would cauſe any to beleeue there were ſome rare goodneſſe or vertue in them. Theſe all are ſometimes preſerued in the Winter, if they bee well defended from the cold. But what ſhall I ſay to the Queene of delight and of flowers, Carnations and Gilloflowers, whoſe brauery, variety, and ſweete ſmell ioyned together, tyeth euery ones affection with great earneſtneſſe, both to like and to haue them? Thoſe that were knowne, and enioyed in former times with much acceptation, are now for the moſt part leſſe accounted of, except a very few: for now there are ſo many other varieties of later inuention, that troubleth the other both in number, beauty, and worth: The names of them doe differ very variably, in that names are impoſed and altered as euerie ones fancy will haue them, that carryed or ſent them into the ſeuerall Countries from London, where their trueſt name is to be had, in mine opinion. I will here but giue you the names of ſome, and referre you to the worke enſuing for your further knowledge. The red and the gray Hulo. The old Carnation, differing from them both. The Gran Pere. The Camberſiue. The Sauadge. The Chriſtall. The Prince. The white Carnation, or Delicate. The ground Carnation. The French Carnation. The Douer. The Oxford. The Briſtow. The Weſtminſter. The Daintie. The Granado, and many other Gilloflowers too tedious to recite in this place, becauſe I haue amply declared them in the booke following. But there is another ſort of great delight and varietie, called the Orange tawny Gilloflower, which for the moſt part hath riſen from ſeed, and doth giue ſeed in a more plentifull manner than any of the former ſorts, and likewiſe by the ſowing of the ſeed there hath been gained ſo many varieties of that excellent worth and reſpect, that it can hardly be expreſſed or beleeued, and called by diuers names according to the marking of the flowers; as The Infanta. The Stript Tawny. The Speckled Tawny. The Flackt Tawny. The Griſeld Tawny, and many others, euery one to bee diſtinguiſhed from others: Some alſo haue their flowers more double and large than others, and ſome from the ſame ſeed haue ſingle flowers like broad ſingle Pinkes: the further relation of them, *viz.* their order to ſowe, encreaſe, and preſerue them, you ſhall haue in the ſubſequent diſcourſe in a place by it ſelfe. Pinkes likewiſe both ſingle and double are of much variety, all of them very ſweete, comming neare the Gilloflowers. Sweete Williams and Sweete Iohns, both ſingle and double, both white, red, and ſpotted, as they are kindes of wilde Pinkes, ſo for their grace and beauty helpe to furniſh a Garden, yet deſire not to ſtand ſo open to the Sunne as the former. Double and ſingle Peonies are fit flowers to furniſh a Garden, and by reaſon of their durability, giue out freſh pleaſure euery yeare without any further trouble of ſowing. And laſtly, Hollihocks both ſingle and double, of many and ſundry colours, yeeld out their flowers like Roſes on their tall branches, like Trees, to ſute you with flowers, when almoſt you haue no other to grace out your Garden: the ſingle and double doe both yeeld ſeed, and yet doe after their ſeeding abide many yeares. Thus haue I ſhewed you moſt of the Engliſh, as well as (I did before) the Outlandiſh

landish flowers, that are fit to furnish the knots, trailes, beds, and borders of this Garden. Roses onely, as I said before, I reserue to circle or encompasse all the rest, because that for the most part they are planted in the outer borders of the quarters, and sometimes by themselues in the middle of long beds, the sorts or kindes whereof are many, as they are declared in their proper place: but the White Rose, the Red, and the Damaske, are the most ancient Standards in England, and therefore accounted naturall.

## CHAP. VI.

### *The order and manner to plant and replant all the sorts of Out-landish flowers spoken of before, as well those with bulbous rootes, as others with stringie rootes.*

WHereas it is the vsuall custome of most in this Land, to turne vp their Gardens, and to plant them againe in the Spring of the yeare, which is the best time that may bee chosen for all English flowers, yet it is not so for your Out-landish flowers. And herein indeede hath beene not onely the errour of a great many to hinder their rootes from bearing out their flowers as they should, but also to hinder many to take delight in them, because as they say they will not thriue and prosper with them, when as the whole fault is in the want of knowledge of the fit and conuenient time wherein they should bee planted. And because our English Gardiners are all or the most of them vtterly ignorant in the ordering of these Out-landish flowers, as not being trained vp to know them, I haue here taken vpon mee the forme of a new Gardiner, to giue instructions to those that will take pleasure in them, that they may be the better enabled with these helpes I shall shew them, both to know how they should be ordered, and to direct their Gardiners that are ignorant thereof, rightly to dispose them according to their naturall qualities. And I doe wish all Gentlemen and Gentlewomen, whom it may concerne for their owne good, to bee as carefull whom they trust with the planting and replanting of these fine flowers, as they would be with so many Iewels; for the rootes of many of them being small, and of great value, may be soone conueyed away, and a cleanly tale faire told, that such a roote is rotten, or perished in the ground if none be seene where it should be, or else that the flower hath changed his colour, when it hath been taken away, or a counterfeit one hath beene put in the place thereof; and thus many haue been deceiued of their daintiest flowers, without remedy or true knowledge of the defect. You shall therefore, if you will take the right course that is proper for these kindes of flowers, not set or plant them among your English flowers; for that when the one may be remoued, the other may not be stirred: but plant those rootes that are bulbous, or round like Onions, eyther in knots or beds by themselues which is the best, or with but very few English or Out-landish flower plants that haue stringie rootes: For you must take this for a generall rule, that all those rootes that are like Lillies or Onions, are to bee planted in the moneths of Iuly or August, or vnto the middle or end of September at the furthest, if you will haue them to prosper as they should; and not in the Spring of the yeare, when other gardening is vsed. Yet I must likewise giue you to vnderstand, that if Tulipas, and Daffodils, and some other that are firme and hard rootes, and not limber or spongie, being taken vp out of the ground in their fit season, that is, in Iune, Iuly, and August, and likewise kept well and dry, may bee reserued out of the ground vntill Christmas or after, and then (if they could not be set sooner) being set, will thriue reasonable well, but not altogether so well as the former, being set long before: but if you shall remoue these bulbous rootes againe, either presently after their planting hauing shot their small fibres vnder the round rootes, and sprung likewise vpwards, or before they be in flower at the soonest (yet Tulipas, Daffodils, and many other bulbous, may be safely remoued being in flower, and transplanted into other places, so as they be not kept too long out of the ground) you shall much endanger them either vtterly to perish, or to be hindered from bearing out their flowers they then would haue

borne, and for two or three years after from bearing flowers againe. For the order of their planting there are diuers wayes, some whereof I will shew you in this place: Your knot or beds being prepared fitly, as before is declared, you may place and order your rootes therein thus, Eyther many rootes of one kind set together in a round or cluster, or longwise crosse a bed one by another, whereby the beauty of many flowers of one kinde being together, may make a faire shew well pleasing to many; Or else you may plant one or two in a place dispersedly ouer the whole knot, or in a proportion or diameter one place answering another of the knot, as your store will suffer you, or your knot permit: Or you may also mingle these rootes in their planting many of diuers sorts together, that they may giue the more glorious shew when they are in flower; and that you may so doe, you must first obserue the seuerall kindes of them, which doe flower at one and the same time, and then to place them in such order and so neare one vnto another, that their flowers appearing together of seuerall colours, will cause the more admiration in the beholders: as thus, The Vernall Crocus or Saffron flowers of the Spring, white, purple, yellow, and stript, with some Vernall Colchicum or Medow Saffron among them, some Deus Caninus or Dogges teeth, and some of the small early Leucoium or Bulbous Violet, all planted in some proportion as neare one vnto another as is fit for them, will giue such a grace to the Garden, that the place will seeme like a peece of tapestry of many glorious colours, to encrease euery ones delight: Or else many of one sort together, as the blew, white and blush Grape flowers in the same manner intermingled, doe make a maruellous delectable shew, especially because all of them rise almost to an equall height, which causeth the greater grace, as well neare hand as farre of. The like order may be kept with many other things, as the Hepatica, white, blew, purple, and red set or sowne together, will make many to beleeue that one roote doth beare all those colours: But aboue and beyond all others, the Tulipas may be so matched, one colour answering and setting of another, that the place where they stand may resemble a peece of curious needle-worke, or peece of painting: and I haue knowne in a Garden, the Master as much commended for this artificiall forme in placing the colours of Tulipas, as for the goodnesse of his flowers, or any other thing. The diuers sorts and colours of Anemones or Winde-flowers may be so ordered likewise, which are very beautifull, to haue the seuerall varieties planted one neare vnto another, that their seuerall colours appearing in one place will be a very great grace in a Garden, or if they be dispersed among the other sorts of flowers, they will make a glorious shew. Another order in planting you may obserue; which is this, That those plants that grow low, as the Aconitum Hyemale or Winter-wolues bane, the Vernall Crocus or Saffron-flowers of diuers sorts, the little early Leucoium or Bulbous Violet, and some such other as rise not vp high, as also some Anemones may be very well placed somewhat neare or about your Martagons, Lillies, or Crownes Imperiall, both because these little plants will flower earlier than they, and so will bee gone and past, before the other greater plants will rise vp to any height to hinder them; which is a way may well be admitted in those Gardens that are small, to saue roome, and to place things to the most aduantage. Thus hauing shewed you diuers wayes and orders how to plant your rootes, that your flowers may giue the greater grace in the Garden, let mee shew you likewise how to set these kindes of rootes into the ground; for many know not well eyther which end to set vpwards or downewards, nor yet to what depth they should be placed in the ground. Daffodils if they be great rootes, will require (as must bee obserued in all other great plants) to bee planted somewhat deeper then the smaller of the same kinde, as also that the tops or heads of the rootes be about two or three fingers breadth hid vnder ground. The Tulipas likewise if you set them deepe, they will be the safer from frosts if your ground be cold, which will also cause them to be a little later before they be in flower, yet vsually if the mould be good, they are to be set a good hand breadth deep within the ground, so that there may be three or foure inches of earth at the least aboue the head, which is the smaller end of the roote: for if they shall lye too neare the vpper face or crust of the earth, the colds & frosts will pierce and pinch them the sooner. After the same order and manner must Hyacinthes, whether great or small, and other such great rootes be planted. Your greater rootes, as Martagons, Lillies, and Crownes Imperiall, must be set much deeper then any other bulbous roote, because they are greater rootes then others, and by themselues also, as

is

is most vsuall either in some square, round, triangle, or other small part in the Garden, because they spread and take vp a very great deale of ground. All of them likewise are to be set with the broad end of the roote downewards, and the small end vpwards, that is, both Lillies, Daffodils, Hyacinthes, and Tulipas, and all other sorts of round rootes, which shew one end to bee smaller than another. But the Colchicum or Medow-Saffron onely requireth an exception to this generall rule, in regard the roote thereof hath a small eminence or part on the one side thereof, which must bee set or planted downeward, and not vpward; for you shall obserue, if the roote lye a little moist out of the ground, that it will shoote fibres out at the small long end thereof, although you may perceiue when you take it vp, that the fibres were at the other broad end or side of the roote. As for the Crowne Imperiall, which is a broad round roote and flat withall, hauing a hole in the middle, for the most part quite thorow, when it is taken vp in his due time out of the ground, you shall perceiue the scales or cloues of the rootes to bee a little open on the vpperside, and close and flat on the vnderside, which will direct you which part to set vpward, as also that the hole is bigger aboue then it is below. The Persian Lilly is almost like vnto the Crowne Imperiall, but that the roote thereof is not so flat, and that it hath a smaller head at the one part, whereby it may be discerned the plainer how to be set. The Fritillaria is a small white root diuided as it were into two parts, so that many haue doubted, as formerly in the Crowne Imperiall, what part to set vppermost; you shall therefore marke, that the two parts of the roote are ioyned together at the bottome, where it shooteth out fibres or small stringie rootes, as all other sorts of bulbous rootes doe, and withall you shall see, that betweene the two parts of the roote a small head will appeare, which is the burgeon that will spring vp to beare leaues and flowers. In the rootes of Anemones there are small round swelling heads, easie enough to be obserued if you marke it, which must be set vpwards. All other sorts of stringie rooted plants (and not bulbous or tuberous rooted) that lose their greene leaues in Winter, will shew a head from whence the leaues and flowers will spring, and all others that keepe their greene leaues, are to bee planted in the same manner that other herbes and flower-plants are accustomed to be. But yet for the better thriuing of the stringie rooted plants, when you will plant them, let me informe you of the best way of planting, and the most sure to cause any plant to comprehend in the ground without failing, and is no common way with any Gardiner in this Kingdome, that euer I heard or knew, which is thus: Presuming that the stringie rooted plant is fresh and not old gathered, and a plant that being remoued will grow againe, make a hole in the ground large enough where you meane to set this roote, and raise the earth within the hole a little higher in the middle then on the sides, and set the roote thereon, spreading the strings all abroad about the middle, that they may as it were couer the middle, and then put the earth gently round about it, pressing it a little close, and afterwards water it well, if it be in Summer, or in a dry time, or otherwise moderately: thus shall euery seuerall string of the roote haue earth enough to cause it to shoote forth, and thereby to encrease farre better than by the vsuall way, which is without any great care and respect to thrust the rootes together into the ground. Diuers other flower plants are but annuall, to bee new sowne euery yeare; as the Maruaile of the world, the Indian Cresses, or yellow Larkes heeles, the Flower of the Sunne, and diuers other: they therefore that will take pleasure in them, that they may enioy their flowers the earlier in the yeare, and thereby haue ripe seede of them while warme weather lasteth, must nurse vp their seedes in a bed of hot dung, as Melons and Cowcumbers are, but your bed must be prouided earlier for these seeds, than for Melons, &c. that they may haue the more comfort of the Summer, which are to be carefully tended after they are transplanted from the hot bed, and couered with straw from colds, whereby you shall not faile to gaine ripe seed euery yeare, which otherwise if you should misse of a very kindly & hot Summer, you should neuer haue. Some of these seedes neede likewise to be transplanted from the bed of dung vnder a warme wall, as the Flower of the Sunne, and the Maruaile of the world, and some others, and that for a while after their transplanting, as also in the heate of Summer, you water them at the roote with water that hath stood a day or two in the Sunne, hauing first laid a round wispe of hay or such other thing round about the roote, that so all helpes may further their giuing of ripe seede. One or two rules more I will giue you concerning

these dainty flowers, the first whereof is this, That you shall not bee carefull to water any of your bulbous or tuberous rooted plants at any time; for they all of them do better prosper in a dry ground than in a wet, onely all sorts of tuberous rooted Flower deluces vpon their remouall had neede of a little water, and some will doe so also to such Tulipas and other bulbous rootes as they transplant, when they are in flower, and this is I grant in some sort tolerable, if it bee not too much, and done onely to cause the stalke and flower to abide sometime the longer before they wither, but else in no other case to be permitted. The second rule is, That I would aduise you to water none of your dainty flowers or herbes, with any water that hath presently before been drawne out of a well or pumpe, but onely with such water that hath stood open in the Sunne in some cisterne, tubbe, or pot for a day at the least, if more the better: for that water which is presently drawne out of a well, &c. is so cold, that it presently chilleth & killeth any dainty plant be it younger or elder grown, wherof I haue had sufficient proofe: and therfore I giue you this caution by mine own experience. Thus haue I directed you from point to point, in all the particulars of preparing & planting that belong to this Garden, sauing only that yet I would further enforme you, of the time of the flowring of these Out-landish plants, according to the seuerall moneths in the yeare, that euery one may know what flowers euery moneth yeeldeth, and may chuse what them liketh best, in that they may see that there is no moneth, but glorieth in some peculiar sorts of rare flowers. I would likewise rather in this place shew you, the true and best manner & order to encrease and preserue all sorts of Gilloflowers & Carnations, then ioyne it with the Chapter of Gilloflowers in the worke following, because it would in that place take vp too much roome. And lastly, I must of necessity oppose three sundry errours, that haue possessed the mindes of many both in former and later times, which are, that any flower may be made to grow double by art, that was but single before by nature: And that one may by art cause any flower to grow of what colour they will: And that any plants may be forced to flower out of their due seasons, either earlier or later, by an art which some can vse. All which being declared, I then suppose enough is spoken for an introduction to this worke, referring many other things to the seuerall directions in the Chapters of the booke.

## Chap. VII.

### *The seuerall times of the flowring of these Out-landish flowers, according to the seuerall moneths of the yeare.*

I Intend in this place onely to giue you briefly, the names of some of the chiefest of these Out-landish flowers, according to the seuerall moneths of the yeare wherein they flower, that euery one seeing what sorts of flowers euery moneth yeeldeth, may take of them which they like best. I begin with Ianuary, as the first moneth of the yeare, wherein if the frosts be not extreme, you shall haue these flowers of plants; the Christmas flower or Helleborus niger verus, Winter wolues bane or Aconitum hyemale, Hepatica or Noble Liuer wort blew and red, and of shrubbes, the Laurus Tinus or Wilde Bay tree, and Mesereon or the dwarfe Bay: but because Ianuarie is oftentimes too deepe in frosts and snow, I therefore referre the Hepaticas vnto the moneth following, which is February, wherein the weather beginneth to be a little milder, and then they will flower much better, as also diuers sorts of Crocus or Saffron flower will appeare, the little early Summer foole or Leucoium bulbosum, and towards the latter end thereof the Vernall Colchicum, the Dogges tooth Violet or Deus Caninus, and some Anemones, both single and double, which in some places will flower all the Winter long. March will yeeld more varieties; for besides that it holdeth some of the flowers of the former moneth, it will yeeld you both the double blew Hepatica, and the white and the blush single: then also you shall haue diuers other sorts of Crocus or Saffron flowers, Double yellow Daffodils, Orientall Iacinths and others, the Crowne Imperiall, diuers sorts of early Tulipas, some sorts of French Cowslips, both tawney, murry, yellow, and blush, the early Fritillaria or checkerd Daffo-

dill,

dill, and some other sorts of early Daffodils, and many sorts of Anemones. In Aprill commeth on the pride of these strangers; for herein you may behold all the sorts of Auricula Vrsi or Beares Eares, many sorts of Anemones, both single and double, both the sorts of Tulipas, the earlier vntill the middle of the moneth, and the later then beginning; which are of so many different colours, that it is almost impossible to expresse them, the white, red, blacke, and yellow Fritillarias, the Muscari or Muske Grape flower, both ash colour and yellow. Diuers other sorts of Iacinths and Daffodils, both single and double, the smaller sorts of Flowerdeluces, the Veluet Flowerdeluce and double Honysuckles, with diuers others. May likewise at the beginning seemeth as glorious as Aprill, although toward the end it doth decline, in regard the heate of the Sunne hath by this time drawne forth all the store of natures tenderest dainties, which are vsually spent by the end of this moneth, and then those of stronger constitution come forward. Herein are to bee seene at the beginning the middle flowring Tulipas, and at the end the later sort: some kindes of Daffodils, the Day Lillies, the great white Starre flower, the Flowerdeluce of Constantinople or the mourning Sable flower, the other sorts of Flowerdeluces. Single and double white Crowfoote, and single and double red Crowfoot, the glory of a Garden: the early red Martagon, the Persian Lilly, the yellow Martagon, the Gladiolus or Corne flagge, both white, red, and blush: the double yellow Rose, and some other sorts of Roses. In Iune doe flower the white and the blush Martagon, the Martagon Imperiall, the mountaine Lillies, and the other sorts of white and red Lillies, the bulbous Flowerdeluces of diuers sorts, the red flowred Ladies bower, the single and double purple flowred Ladies bower, the white Syringa or Pipetree, for the blew Pipe tree flowreth earlier, the white and the yellow Iasmin. Iuly holdeth in flower some of the Ladies bowers and Iasmines, and besides doth glory in the Female Balsame apple, the Indian Cresses or yellow Larkes spurres, the purple Flower-gentle and the Rose Bay. In August begin some of the Autumne bulbous flowers to appeare, as the white and the purple Colchicum or Medow Saffron, the purple mountaine Crocus or Saffron flower, the little Autumne Leucoium and Autumne Iacinth, the Italian Starrewort, called of some the purple Marigold, the Meruaile of Peru or of the world, the Flower of the Sunne, the great blew Bell-flower, the great double French Marigold. September flourisheth with the Flower of the Sunne, the Meruaile of the world, the purple Marigold, and blew Bell-flower spoken of before, and likewise the other sorts of Medow Saffron, and the double kinde likewise, the siluer Crocus, the Autumne yellow Daffodill, Cyclamen also or Sowbread shew their flowers in the end of this moneth. October also will shew the flowers of Cyclamen, and some of the Medow Saffrons. In Nouember, as also sometimes in the moneth before, the party coloured Medow Saffron may bee seene, that will longest hold his flower, because it is the latest that sheweth it selfe, and the ash coloured mountaine Crocus. And euen December it selfe will not want the true blacke Hellebor or Christmas flower, and the glorious shew of the Laurus Tinus or wilde Bay tree. Thus haue I shewed you some of the flowers for euery moneth, but I referre you to the more ample declarion of them and all the others, vnto the work following.

## Chap. VIII.

### *The true manner and order to encrease and preserue all sorts of Gilloflowers, as well by slippes as seedes.*

BEcause that Carnations and Gilloflowers bee the chiefest flowers of account in all our English Gardens, I haue thought good to entreate somewhat amply of them, and that a part by it selfe, as I said a little before, in regard there is so much to be said concerning them, and that if all the matters to be entreated of should haue beene inserted in the Chapter of Gilloflowers, it would haue made it too tedious and large, and taken vp too much roome. The particular matters whereof I mean in this place to entreate are these: How to encrease Gilloflowers by planting and by sowing,

sowing, and how to preserue them being encreased, both in Summer from noysome and hurtfull vermine that destroy them, and in Winter from frosts, snowes, and windes, that spoile them. There are two wayes of planting, whereby to encrease these faire flowers; the one is by slipping, which is the old and ready vsuall way, best knowne in this Kingdome; the other is more sure, perfect, ready, and of later inuention, *videlicet*, by laying downe the branches. The way to encrease Gilloflowers by slipping, is so common with all that euer kept any of them, that I thinke most persons may thinke me idle, to spend time to set downe in writing that which is so well known vnto all: Yet giue me leaue to tell them that so might imagine, that (when they haue heard or read what I haue written thereof, if they did know fully as much before) what I here write, was not to informe them, but such as did not know the best, or so good a way as I teach them: For I am assured, the greatest number doe vse, and follow the most vsuall way, and that is not alwaies the best, especially when by good experience a better way is found, and may be learned; and therefore if some can doe a thing better than others, I thinke it is no shame to learne it of them. You shall not then (to take the surest course) take any long spindled branches, nor those branches that haue any young shootes from the ioynts on them, nor yet sliue or teare any slippe or branch from the roote; for all these waies are vsuall and common with most, which causeth so many good rootes to rot and perish, and also so many slippes to be lost, when as for the most part, not the one halfe, or with some, not a third part doth grow and thriue of those slippes they set. And although many that haue store of plants, doe not so much care what hauocke they make to gaine some, yet to saue both labour and plants, I doe wish them to obserue these orders: Take from those rootes from whence you intend to make your encrease, those shootes onely that are reasonable strong, but yet young, and not either too small and slender, or hauing any shootes from the ioynts vpon them; cut these slippes or shootes off from the stemme or roote with a knife, as conueniently as the shoote or branch will permit, that is, either close vnto the maine branch, if it be short, or leauing a ioynt or two behinde you, if it be long enough, at which it may shoote anew: When you haue cut off your slippes, you may either set them by and by, or else as the best Gardiners vse to doe, cast them into a tubbe or pot with water for a day or two, and then hauing prepared a place conuenient to set them in, which had neede to bee of the finest, richest, and best mould you can prouide, that they may thriue therein the better, cut off your slippe close at the ioynt, and hauing cut away the lowest leaues close to the stalke, and the vppermost euen at the top, with a little sticke make a little hole in the earth, and put your slippe therein so deep, as that the vpper leaues may be wholly aboue the ground, (some vse to cleaue the stalke in the middle, and ~~put a little earth or clay within~~ the cleft, but many good and skilfull Gardiners doe not vse it); put the earth a little close to the slippe with your finger and thumbe, and there let it rest, and in this manner doe with as many slippes as you haue, setting them somewhat close together, and not too farre in sunder, both to saue ground and cost thereon, in that a small compasse will serue for the first planting, and also the better to giue them shadow: For you must remember in any case, that these slippes new set, haue no sight of the Sunne, vntill they be well taken in the ground, and shot aboue ground, and also that they want not water, both vpon the new planting and after. When these slippes are well growne vp, they must be transplanted into such other places as you thinke meete; that is, either into the ground in beds, or otherwise, or into pots, which that you may the more safely doe, after you haue well watered the ground, for halfe a day before you intend to transplant them, you shall separate them seuerally, by putting down a broad pointed knife on each side of the slippe, so cutting it out, take euery one by it selfe, with the earth cleauing close vnto the root, which by reason of the moisture it had formerly, and that which you gaue presently before, will be sufficient with any care had, to cause it to hold fast vnto the roote for the transplanting of it: for if the earth were dry, and that it should fall away from the roote in the transplanting, it would hazzard and endanger the roote very much, if it did thriue at all. You must remember also, that vpon the remouing of these slips, you shadow them from the heate of the Sunne for a while with some straw or other thing, vntill they haue taken hold in their new place. Thus although it bee a little more labour and care than the ordinary way is, yet it is surer, and will giue you plants that

will be ſo ſtrongly growne before Winter, that with the care hereafter ſpecified, you ſhall haue them beare flowers the next yeare after, and yeeld you encreaſe of ſlippes alſo. To giue you any ſet time, wherein theſe ſlippes will take roote, and begin to ſhoote aboue ground, is very hard to doe; for that euery ſlip, or yet euery kinde of Gilloflower is not alike apt to grow; nor is euery earth in like manner fit to produce and bring forward the ſlippes that are ſet therein: but if both the ſlippe be apt to grow, and the earth of the beſt, fit to produce, I thinke within a fortnight or three weekes, you ſhall ſee them begin to put forth young leaues in the middle, or elſe it may be a moneth and more before you ſhall ſee any ſpringing. The beſt time likewiſe when to plant, is a ſpeciall thing to be knowne, and of as great conſequence as any thing elſe: For if you ſlippe and ſet in September, as many vſe to doe, or yet in Auguſt, as ſome may thinke will doe well, yet (vnleſſe they be the moſt ordinary ſorts, which are likely to grow at any time, and in any place) the moſt of them, if not all, will either aſſuredly periſh, or neuer proſper well: for the more excellent and dainty the Gilloflower is, the more tender for the moſt part, and hard to nurſe vp will the ſlippes be. The beſt time therefore is, that you cut off ſuch ſlippes as are likely, and ſuch as your rootes may ſpare, from the beginning of May vntill the middle of Iune at the furtheſt, and order them as I haue ſhewed you before, that ſo you may haue faire plants, plenty of flowers, and encreaſe ſufficient for new ſupply, without offence or loſſe of your ſtore. For the enriching likewiſe of your earth, wherein you ſhall plant your ſlippes, that they may the better thriue and proſper, diuers haue vſed diuers ſorts of manure; as ſtable ſoyle of horſe, beaſts or kine, of ſheepe, and pigeons, all which are very good when they are thoroughly turned to mould, to mixe with your other earth, or being ſteeped in water, may ſerue to water the earth at times, and turned in with it. And ſome haue likewiſe proued Tanners earth, that is, their barke, which after they haue vſed, doth lye on heapes and rot in their yards, or the like mould from wood-ſtackes or yards; but eſpecially, and beyond all other is commended the Willow earth, that is, that mould which is found in the hollow of old Willow trees, to be the moſt principall to mixe with other good earth for this purpoſe. And as I haue now giuen you directions for the firſt way to encreaſe them by ſlipping, ſo before I come to the other way, let mee giue you a caueat or two for the preſeruing of them, when they are beginning to runne vtterly to decay and periſh: The one is, that whereas many are ouer greedy to haue their plants to giue them flowers, and therefore let them runne all to flower, ſo farre ſpending themſelues thereby, that after they haue done flowring, they grow ſo weake, hauing out ſpent themſelues, that they cannot poſſibly be preſerued from the iniuries of the ſucceeding Winter; you ſhall therefore keepe the kinde of any ſort you are delighted withall, if you carefully looke that too many branches doe not runne vp and ſpindle for flowers, but rather either cut ſome of them downe, before they are run vp too high, within two or three ioynts of the rootes; or elſe plucke away the innermoſt leaues where it ſpringeth forwards, which you ſee in the middle of euery branch, before it be runne vp too high, which will cauſe them to breake out the faſter into ſlips and ſuckers at the ioynts, to hinder their forward luxurie, and to preſerue them the longer: The other is, If you ſhall perceiue any of your Gilloflower leaues to change their naturall freſh verdure, and turne yellowiſh, or begin to wither in anie part or branch thereof, it is a ſure ſigne that the roote is infected with ſome cancker or rottenneſſe, and will ſoone ſhew it ſelfe in all the reſt of the branches, whereby the plant will quickly be loſt: to preſerue it therefore, you ſhall betime, before it be runne too farre, (for otherwiſe it is impoſſible to ſaue it) either couer all or moſt of the branches with freſh earth, or elſe take the faireſt ſlippes from it, as many as you can poſſibly, and caſt them into a pot or tubbe with water, and let them there abide for two or three daies at the leaſt: the firſt way hath recouered many, being taken in time. Thus you ſhall ſee them recouer their former ſtiffeneſſe and colour, and then you may plant them as you haue beene heretofore directed; and although many of them may periſh, yet ſhall you haue ſome of them that will grow to continue the kinde againe. The other or ſecond way to encreaſe Gilloflowers by planting, is, as I ſaid before, by in-laying or laying downe the branches of them, and is a way of later inuention, and as frequently vſed, not onely for the tawney or yellow Gilloflower, and all the varieties therof, but with the other kinds of Gilloflowers, whereof experience hath ſhewed

that

that they will likewise take if they be so vsed; the manner whereof is thus: You must choose out the youngest, likeliest, and lowest branches that are neerest the ground (for the vpper branches will sooner breake at the ioynt, than bend downe so low into the earth, without some pot with earth raised vp vnto them) and cut it on the vnderside thereof vpwards at the second ioynt next vnto the roote, to the middle of the branch, and no more, and not quite thorough in any case, and then from that second ioynt vnto the third, slit or cut the branch in the middle longwise, that so it may be the more easily bended into the ground, the cut ioynt seeming like the end of a slippe, when you haue bended downe the branch where it is cut into the ground (which must bee done very gently for feare of breaking) with a little sticke or two thrust slopewise, crosse ouer it, keepe it downe within the earth, and raise vp sufficient earth ouer it, that there it may lye and take roote, which commonly will be effected within sixe weekes or two moneths in the Summer time, and then (or longer if you doubt the time too short for it to take sufficient roote) you may take or cut it away, and transplant it where you thinke good, yet so as in any case you shadow it from the heate of the Sunne, vntill it haue taken good hold in the ground. The other way to encrease Gilloflowers, is by sowing the seede: It is not vsuall with all sorts of Gilloflowers to giue seede, but such of them as doe yeeld seede may be encreased thereby, in the same manner as is here set downe. The Orange tawney Gilloflower and the varieties thereof is the most vsuall kinde, (and it is a kinde by it selfe, how various soeuer the plants be that rise from the seede) that doth giue seede, and is sowne, and from thence ariseth so many varieties of colours, both plaine and mixt, both single and double, that one can hardly set them downe in writing: yet such as I haue obserued and marked, you shall finde expressed in the Chapter of Gilloflowers in the worke following. First therefore make choise of your seede that you intend to sowe (if you doe not desire to haue as many more single flowers as double) that it bee taken from double flowers, and not from single, and from the best colours, howsoeuer some may boast to haue had double and stript flowers from the seede of a single one; which if it were so, yet one Swallow (as we say) maketh no Summer, nor a thing comming by chance cannot bee reckoned for a certaine and constant rule; you may be assured they will not vsually doe so: but the best, fairest, and most double flowers come alwaies, or for the most part, from the seede of those flowers that were best, fairest, and most double; and I doe aduise you to take the best and most double: for euen from them you shall haue single ones enow, you needе not to sowe any worser sort. And againe, see that your seede bee new, of the last yeares gathering, and also that it was full ripe before it was gathered, lest you lose your labour, or misse of your purpose, which is, to haue faire and double flowers. Hauing ~~now made choise of your seede,~~ and prepared you a bedde to sowe them on, the earth whereof must be rich and good, and likewise sifted to make it the finer; for the better it is, the better shall your profit and pleasure bee: hereon, being first made leuell, plaine, and smooth, sowe your seede somewhat thinne, and not too thicke in any case, and as euenly as you can, that they be not too many in one place, and too few in another, which afterwards couer with fine sifted earth ouer them about one fingers thickneſſe; let this be done in the middle of Aprill, if the time of the yeare be temperate, and not too cold, or else stay vntill the end of the moneth: after they are sprung vp and growne to be somewhat bigge, let them bee drawne forth that are too close and neare one vnto another, and plant them in such place where they shall continue, so that they stand halfe a yard of ground distance asunder, which after the planting, let be shadowed for a time, as is before specified; and this may bee done in the end of Iuly, or sooner if there be cause. I haue not set downe in all this discourse of planting, transplanting, sowing, setting, &c. any mention of watering those slips or plants, not doubting but that euery ones reason will induce them to thinke, that they cannot prosper without watering: But let this Caueat be a sufficient remembrance vnto you, that you neuer water any of these Gilloflowers, nor yet indeede any other fine herbe or plant with cold water, such as you haue presently before drawne out from a pumpe or Well, &c. but with such water as hath stood open in the aire in a cisterne, tubbe, or pot, for one whole day at the least; if it be two or three daies it will be neuer the worse, but rather the better, as I haue related before: yet take especiall heede that you doe not giue them too much to ouer-glut them at any time, but temperately to irrorate

rorate, bedew or sprinkle them often. From the seedes of these Gilloflowers hath risen both white, red, blush, stamell, tawny lighter and sadder, marbled, speckled, striped, flaked, and that in diuers manners, both single and double flowers, as you shall see them set downe in a more ample manner in the Chapter of Gilloflowers. And thus much for their encrease by the two wayes of planting and sowing: For as for a third way, by grafting one into or vpon another, I know none such to be true, nor to be of any more worth than an old Wiues tale, both nature, reason, and experience, all contesting against such an idle fancy, let men make what ostentation they please. It now resteth, that we also shew you the manner how to preserue them, as well in Summer from all noysome and hurtfull things, as in the Winter and Spring from the sharp and chilling colds, and the sharpe and bitter killing windes in March. The hurtfull things in the Summer are especially these, too much heate of the Sunne which scorcheth them, which you must be carefull to preuent, by placing boughes, boords, clothes or mats, &c. before them, if they bee in the ground; or else if they bee in pots, to remoue into them into the shadow, to giue them refreshing from the heate, and giue them water also for their life: too much water, or too little is another annoyance, which you must order as you see there is iust cause, by withholding or giuing them water gently out of a watering pot, and not cast on by dishfuls: Some also to water their Gilloflowers, vse to set their pots into tubbes or pots halfe full of water, that so the water may soake in at the lower holes in each flower pot, to giue moisture to the roots of the Gilloflowers onely, without casting any water vpon the leaues, and assuredly it is an excellent way to moisten the rootes so sufficiently at one time, that it doth saue a great deale of paines many other times. Earwickes are a most infestuous vermine, to spoyle the whole beauty of your flowers, and that in one night or day; for these creatures delighting to creepe into any hollow or shadowie place, doe creepe into the long greene pods of the Gilloflowers, and doe eate away the white bottomes of their leaues, which are sweete, whereby the leaues of the flowers being loose, doe either fall away of themselues before, or when they are gathered, or handled, or presently wither within the pods before they are gathered, and blowne away with the winde. To auoide which inconuenience, many haue deuised many waies and inuentions to destroy them, as pots with double verges or brimmes, containing a hollow gutter betweene them, which being filled with water, will not suffer these small vermine to passe ouer it to the Gilloflowers to spoile them. Others haue vsed old shooes, and such like hollow things to bee set by them to take them in: but the best and most vsuall things now vsed, are eyther long hollow canes, or else beasts hoofes, ~~which~~ being turned downe vpon sticks ends set into the ground, or into the pots of earth, will soone draw into them many Earwickes, lying hid therein from sunne, winde, and raine, and by care and diligence may soone bee destroyed, if euery morning and euening one take the hoofes gently off from the sticks, and knocking them against the ground in a plain allie, shake out all the Earwicks that are crept into them, which quickly with ones foot may be trode to peeces. For sodain blasting with thunder and lightening, or fierce sharpe windes, &c. I know no other remedy, vnlesse you can couer them therefrom when you first foresee the danger, but patiently to abide the losse, whatsoeuer some haue aduised, to lay litter about them to auoide blasting; for if any shall make tryall thereof, I am in doubt, he shall more endanger his rootes thereby, being the Summer time, when any such feare of blasting is, than any wise saue them from it, or doe them any good. For the Winter preseruation of them, some haue aduised to couer them with Bee-hiues, or else with small Willow sticks, prickt crossewise into the ground ouer your flowers, and bowed archwise, and with litter laid thereon, to couer the Gilloflowers quite ouer, after they haue beene sprinkled with sope ashes and lyme mixt together: and this way is commended by some that haue written thereof, to be such an admirable defence vnto them in Winter, that neither Ants, nor Snailes, nor Earwickes shall touch them, because of the sope ashes and lyme, and neyther frosts nor stormes shall hurt them, because of the litter which so well will defend them; and hereby also your Gilloflowers will bee ready to flower, not onely in the Spring very early, but euen all the Winter. But whosoeuer shall follow these directions, may peraduenture finde them in some part true, as they are there set downe for the Winter time, and while they are kept close and couered; but let them bee assured,

that

that all such plants, or the most part of them, will certainely perish and dye before the Summer be at an end: for the sope ashes and lyme will burne vp and spoile any herbe; and againe, it is impossible for any plant that is kept so warme in Winter, to abide eyther the cold or the winde in the Spring following, or any heate of the Sun, but that both of them will scorch them, and carry them quite away. One great hurt vnto them, and to all other herbes that wee preserue in Winter, is to suffer the snow to lye vpon them any time after it is fallen, for that it doth so chill them, that the Sunne afterward, although in Winter, doth scorch them and burne them vp: looke therefore vnto your Gilloflowers in those times, and shake or strike off the snow gently off from them, not suffering it to abide on them any day or night if you can; for assure your selfe, if it doth not abide on them, the better they will be. The frosts likewise is another great annoyance vnto them, to corrupt the rootes, and to cause them to swell, rot, and breake: to preuent which inconuenience, I would aduise you to take the straw or litter of your horse stable, and lay some thereof about euery roote of your Gilloflowers (especially those of the best account) close vnto them vpon the ground, but be as carefull as you can, that none thereof lye vpon the greene leaues, or as little as may be, and by this onely way haue they been better defended from the frosts that spoile them in Winter, then by any other that I haue seen or knowne. The windes in March, and Sunneshine dayes then, are one of the greatest inconueniences that happeneth vnto them: for they that haue had hundreds of plants, that haue kept faire and greene all the Winter vntill the beginning or middle of March, before the end thereof, haue had scarce one of many, that either hath not vtterly perished, or been so tainted, that quickly after haue not been lost; which hath happened chiefly by the neglect of these cautions before specified, or in not defending them from the bitter sharpe windes and sunne in this moneth of March. You shall therefore for their better preseruation, besides the litter laid about the rootes, which I aduise you not to remoue as yet, shelter them somewhat from the windes, with eyther bottomlesse pots, pales, or such like things, to keep away the violent force both of windes and sun for that moneth, and for some time before & after it also: yet so, that they be not couered close aboue, but open to receiue ayre & raine. Some also vse to wind withes of hey or straw about the rootes of their Gilloflowers, and fasten them with stickes thrust into the ground, which serue very well in the stead of the other. Thus haue I shewed you the whole preseruation of these worthy and dainty flowers, with the whole manner of ordering them for their encrease: if any one haue any other better way, I shall be as willing to learne it of them, as I haue beene to giue them or any others the knowledge of that I haue here set downe.

## Chap. IX.

*That there is not any art whereby any flower may be made to grow double, that was naturally single, nor of any other sent or colour than it first had by nature; nor that the sowing or planting of herbes one deeper than other, will cause them to be in flower one after another, euery moneth in the yeare.*

THe wonderfull desire that many haue to see faire, double, and sweete flowers, hath transported them beyond both reason and nature, feigning and boasting often of what they would haue, as if they had it. And I thinke, from this desire and boasting hath risen all the false tales and reports, of making flowers double as they list, and of giuing them colour and sent as they please, and to flower likewise at what time they will, I doubt not, but that some of these errours are ancient, and continued long by tradition, and others are of later inuention: and therefore the more to be condemned, that men of wit and iudgement in these dayes should expose themselues in their writings, to be rather laughed at, then beleeued for such idle tales. And although in the contradiction of them, I know I shall vndergoe many calumnies, yet notwithstanding, I will endeauour to set downe and declare so much, as I hope may by reason

perswade

perswade many in the truth, although I cannot hope of all, some being so strongly wedded to their owne will, and the errours they haue beene bred in, that no reason may alter them. First therefore I say, that if there were any art to make some flowers to grow double, that naturally were single, by the same art, all sorts of flowers that are single by nature, may be made to grow double: but the sorts of flowers that are single by nature, whereof some are double, were neuer made double by art; for many sorts abide still single, whereof there was neuer seene double: and therefore there is no such art in any mans knowledge to bring it to passe. If any man shall say, that because there are many flowers double, whereof there are single also of the same kinde, as for example, Violets, Marigolds, Daisyes, Daffodils, Anemones, and many other, that therefore those double flowers were so made by the art of man: *viz.* by the obseruation of the change of the Moone, the constellations or coniunctions of Planets, or some other Starres or celestiall bodies. Although I doe confesse and acknowledge, that I thinke some constellations, and peraduenture changes of the Moone, &c. were appointed by the God of nature, as conducing and helping to the making of those flowers double, that nature hath so produced; yet I doe deny, that any man hath or shall euer be able to proue, that it was done by any art of man, or that any man can tell the true causes and seasons, what changes of the Moone, or constellations of the Planets, wrought together for the producing of those double flowers, or can imitate nature, or rather the God of nature, to doe the like. If it shall bee demanded, From whence then came these double flowers that we haue, if they were not so made by art? I answer, that assuredly all such flowers did first grow wilde, and were so found double, as they doe now grow in Gardens, but for how long before they were found they became double, no man can tell; we onely haue them as nature hath produced them, and so they remaine. Againe, if any shall say, that it is likely that these double flowers were forced so to be, by the often planting and transplanting of them, because it is obserued in most of them, that if they stand long in any one place, and not be often remoued, they will grow still lesse double, and in the end turne single. I doe confesse, that *Facilior est descensus quàm ascensus*, and that the vnfruitfulnesse of the ground they are planted in, or the neglect or little care had of them, or the growing of them too thicke or too long, are oftentimes a cause of the diminishing of the flowers doublenesse; but withall you shall obserue, that the same rootes that did beare double flowers (and not any other that neuer were double before) haue returned to their former doublenesse againe, by good ordering and looking vnto: single flowers haue only beene made somewhat fairer or larger, by being planted in the richer ~~and more fruit~~-full ground of the Garden, than they were found wilde by nature; but neuer made to grow double, as that which is naturally so found of it selfe: For I will shew you mine owne experience in the matter. I haue been as inquisitiue as any man might be, with euery one I knew, that made any such report, or that I thought could say any thing therein, but I neuer could finde any one, that could assuredly resolue me, that he knew certainly any such thing to be done: all that they could say was but report, for the obseruation of the Moone, to remoue plants before the change, that is, as some say, the full of the Moone, others the new Moone, whereupon I haue made tryall at many times, and in many sorts of plants, accordingly, and as I thought fit, by planting & transplanting them, but I could neuer see the effect desired, but rather in many of them the losse of my plants. And were there indeed such a certaine art, to make single flowers to grow double, it would haue beene knowne certainly to some that would practise it, and there are so many single flowers, whereof there were neuer any of the kinde seene double, that to produce such of them to be double, would procure both credit and coyne enough to him that should vse it; but *Vltra posse non est esse*: and therefore let no man beleeue any such reports, bee they neuer so ancient; for they are but meere tales and fables. Concerning colours and sents, the many rules and directions extant in manie mens writings, to cause flowers to grow yellow, red, greene, or white, that neuer were so naturally, as also to be of the sent of Cinamon, Muske, &c. would almost perswade any, that the matters thus set downe by such persons, and with some shew of probability, were constant and assured proofes thereof: but when they come to the triall, they all vanish away like smoake. I will in a few words shew you the matters and manners of their proceedings to effect this purpose: First (they say) if you shall steepe

your

your seedes in the lees of red Wine, you shall haue the flowers of those plants to be of a purple colour. If you will haue Lillies or Gilloflowers to be of a Scarlet red colour, you shall put Vermillion or Cynaber betweene the rinde and the small heads growing about the roote: if you will haue them blew, you shall dissolue Azur or Byse between the rinde and the heads: if yellow, Orpiment: if greene, Vardigrease, and thus of any other colour. Others doe aduise to open the head of the roote, and poure into it any colour dissolued, so that there be no fretting or corroding thing therein for feare of hurting the roote, and looke what colour you put in, iust such or neare vnto it shall the colour of the flower bee. Some againe doe aduise to water the plants you would haue changed, with such coloured liquor as you desire the flower to be of, and they shall grow to be so. Also to make Roses to bee yellow, that you should graft a white Rose (some say a Damaske) vpon a Broome stalke, and the flower will be yellow, supposing because the Broome flower is yellow, therefore the Rose will be yellow. Some affirme the like, if a Rose be grafted on a Barbery bush, because both the blossome and the barke of the Barbery is yellow, &c. In the like manner for sents, they haue set downe in their writings, that by putting Cloues, Muske, Cinamon, Benzoin, or any other such sweete thing, bruised with Rose water, between the barke and the body of trees, the fruit of them will smell and taste of the same that is put vnto them; and if they bee put vnto the toppe of the rootes, or else bound vnto the head of the roote, they will cause the flowers to smell of that sent the matter put vnto them is of: as also to steep the seeds of Roses, and other plants in the water of such like sweet things, and then to sowe them, and water them morning and euening with such like liquor, vntill they be growne vp; besides a number of such like rules and directions set downe in bookes, so confidently, as if the matters were without all doubt or question: whenas without all doubt and question I will assure you, that they are all but meere idle tales & fancies, without all reason or truth, or shadow of reason or truth: For sents and colours are both such qualities as follow the essence of plants, euen as formes are also; and one may as well make any plant to grow of what forme you will, as to make it of what sent or colour you will; and if any man can forme plants at his will and pleasure, he can doe as much as God himselfe that created them. For the things they would adde vnto the plants to giue them colour, are all corporeall, or of a bodily substance, and whatsoeuer should giue any colour vnto a liuing and growing plant, must be spirituall: for no solide corporeall substance can ioyne it selfe with the life and essence of an herbe or tree, and the spirituall part of the colour thereof is not the same with the bodily substance, but is a meere vapour that riseth from the substance, and feedeth the plant, whereby it groweth, so that there is no ground or colour of reason, that a substantiall colour ~~should giue colour to a~~ growing herbe or tree: but for sent (which is a meere vapour) you will say there is more probability. Yet consider also, that what sweete sent soeuer you binde or put vnto the rootes of herbes or trees, must be either buried, or as good as buried in the earth, or barke of the tree, whereby the substance will in a small time corrupt and rot, and before it can ioyne it selfe with the life, spirit, and essence of the plant, the sent also will perish with the substance: For no heterogeneall things can bee mixed naturally together, as Iron and Clay; and no other thing but homogeneall, can be nourishment or conuertible into the substance of man or beast: And as the stomach of man or beast altereth both formes, sents, and colours of all digestible things; so whatsoeuer sent or colour is wholsome, and not poysonfull to nature, being receiued into the body of man or beast, doth neither change the bloud or skinne into that colour or sent was receiued: no more doth any colour or sent to any plant; for the plants are onely nourished by the moisture they draw naturally vnto them, be it of wine or any other liquor is put vnto them, and not by any corporeall substance, or heterogeneall vapour or sent, because the earth like vnto the stomach doth soone alter them, before they are conuerted into the nature and substance of the plant. Now for the last part I vndertooke to confute, that no man can by art make all flowers to spring at what time of the yeare hee will; although, as I haue here before shewed, there are flowers for euery moneth of the yeare, yet I hope there is not any one, that hath any knowledge in flowers and gardening, but knoweth that the flowers that appeare and shew themselues in the seuerall moneths of the yeare, are not one and the same, and so made to flower by art; but that they are seuerall sorts of plants, which

will

will flower naturally and constantly in the same moneths one yeare, that they vse to doe in another, or with but little alteration, if the yeares proue not alike kindly: As for example, those plants that doe flower in Ianuary and February, will by no art or industry of man be caused to flower in Summer or in Autumne; and those that flower in Aprill and May, will not flower in Ianuary or February; or those in Iuly, August, &c. either in the Winter or Spring: but euery one knoweth their owne appointed naturall times, which they constantly obserue and keepe, according to the temperature of the yeare, or the temper of the climate, being further North or South, to bring them on earlier or later, as it doth with all other fruits, flowers, and growing greene herbes, &c. except that by chance, some one or other extraordinarily may be hindered in their due season of flowring, and so giue their flowers out of time, or else to giue their flowers twice in the yeare, by the superaboundance of nourishment, or the mildnesse of the season, by moderate showers of raine, &c. as it sometimes also happeneth with fruits, which chance, as it is seldome, and not constant, so we then terme it but *Lusus naturæ*: or else by forcing them in hot stoues, which then will perish, when they haue giuen their flowers or fruits. It is not then, as some haue written, the sowing of the seedes of Lillies, or any other plants a foote deepe, or halfe a foote deepe, or two inches deepe, that will cause them to be in flower one after another, as they are sowne euery moneth of the yeare; for it were too grosse to thinke, that any man of reason and iudgement would so beleeue. Nor is it likewise in the power of any man, to make the same plants to abide a moneth, two, or three, or longer in their beauty of flowring, then naturally they vse to doe; for I thinke that were no humane art, but a supernaturall worke. For nature still bendeth and tendeth to perfection, that is, after flowring to giue fruit or seede; nor can it bee hindered in the course thereof without manifest danger of destruction, euen as it is in all other fruit-bearing creatures, which stay no longer, then their appointed time is naturall vnto them, without apparent damage. Some things I grant may be so ordered in the planting, that according to that order and time which is obserued in their planting, they shall shew forth their faire flowers, and they are Anemones, which will in that manner, that I haue shewed in the worke following, flower in seuerall moneths of the yeare; which thing as it is incident to none or very few other plants, and is found out but of late, so likewise is it knowne but vnto a very few. Thus haue I shewed you the true solution of these doubts: And although they haue not beene amplified with such Philosophicall arguments and reasons, as one of greater learning might haue done, yet are they truely and sincerely set downe, that they may serue *tanquam galeatum*, against all the calumnies and obiections of wilfull and obdurate persons, that will not be reformed. As first, that all double flowers were so found wilde, being the worke of nature alone, and not the art of any man, by planting or transplanting, at or before the new or full Moone, or any other obseruation of time, that hath caused the flower to grow double, that naturally was single: Secondly, that the rules and directions, to cause flowers to bee of contrary or different colours or sents, from that they were or would be naturally, are meere fancies of men, without any ground of reason or truth. And thirdly, that there is no power or art in man, to cause flowers to shew their beauty diuers moneths before their naturall time, nor to abide in their beauty longer then the appointed naturall time for euery one of them.

# THE GARDEN OF PLEASANT FLOWERS.

## CHAP. I.

### *Corona Imperialis.* The Crowne Imperiall.

BEcauſe the Lilly is the more ſtately flower among manie: and amongſt the wonderfull varietie of Lillies, knowne to vs in theſe daies, much more then in former times, whereof ſome are white, others bluſh, ſome purple, others red or yellow, ſome ſpotted, others without ſpots, ſome ſtanding vpright, others hanging or turning downewards, The Crowne Imperiall for his ſtately beautifulneſs, deſerueth the firſt place in this our Garden of delight, to be here entreated of before all other Lillies: but becauſe it is ſo well knowne to moſt perſons, being in a manner euery where common, I ſhall neede onely to giue you a relation of the chiefe parts thereof (as I intend in ſuch other things) which are theſe: The roote is yellowiſh on the outſide, compoſed of fewer, but much thicker ſcales, then any other Lilly but the Perſian, and doth grow ſometimes to be as great as a pretty bigge childes head, but ſomewhat flat withall, from the ſides whereof, and not from the bottome, it ſhooteth forth thicke long fibres, which periſh euery yeare, hauing a hole in the midſt thereof, at the end of the yeare, when the old ſtalke is dry and withered, and out of the which a new ſtalke doth ſpring againe (from a bud or head to be ſeen within the hollowneſſe on the one ſide) the yeare following: the ſtalke then filling vp the hollowneſſe, riſeth vp three or foure foote high, being great, round, and of a purpliſh colour at the bottome, but greene aboue, beſet from thence to the middle thereof with many long and broad greene leaues, very like to the leaues of our ordinary white Lilly, but ſomewhat ſhorter and narrower, confuſedly without order, and from the middle is bare or naked without leaues, for a certaine ſpace vpwards, and then beareth foure, ſixe, or tenne flowers, more or leſſe, according to the age of the plant, and the fertility of the ſoyle where it groweth; The buddes at the firſt appearing are whitiſh, ſtanding vpright among a buſh or tuft of greene leaues, ſmaller then thoſe below, and ſtanding aboue the flowers, after a while they turne themſelues, and hang downewards euerie one vpon his owne footeſtalke, round about the great ſtemme or ſtalke, ſometimes of an euen depth, and other while one lower or higher than another, which flowers are neare the forme of an ordinary Lilly, yet ſomewhat leſſer and cloſer, conſiſting of ſixe leaues of an Orange colour, ſtriped with purpliſh lines and veines, which adde a great grace to the flowers: At the bottome of the flower next vnto the ſtalke, euery

leafe thereof hath on the outſide a certaine bunch or eminence, of a darke purpliſh colour, and on the inſide there lyeth in thoſe hollow bunched places, certaine cleare drops of water like vnto pearles, of a very ſweete taſte almoſt like ſugar: in the midſt of each flower is a long white ſtile or pointell, forked or diuided at the end, and ſixe white chiues tipt with yellowiſh pendents, ſtanding cloſe about it: after the flowers are paſt, appeare ſixe ſquare ſeede veſſels ſtanding vpright, winged as it were or welted on the edges, yet ſeeming but three ſquare, becauſe each couple of thoſe welted edges are ioyned cloſer together, wherein are contained broad, flat, and thinne ſeedes, of a pale browniſh colour, like vnto other Lillies, but much greater and thicker alſo. The ſtalke of this plant doth oftentimes grow flat, two, three, or foure fingers broad, and then beareth many more flowers, but for the moſt part ſmaller then when it beareth round ſtalkes. And ſometimes it happeneth the ſtalke to be diuided at the top, carrying two or three tufts of greene leaues, without any flowers on them. And ſometimes likewiſe, to beare two or three rowes or crownes of flowers one aboue another vpon one ſtalke, which is ſeldome and ſcarce ſeene, and beſides, is but meere accidentall: the whole plant and euery part thereof, as well rootes, as leaues and flowers, doe ſmell ſomewhat ſtrong as it were the ſauour of a Foxe, ſo that if any doe but come neare it, he cannot but ſmell it, which yet is not vnwholſome.

I haue not obſerued any variety in the colour of this flower, more then that it will be fairer in a cleare open ayre, and paler, or as it were blaſted in a muddy or ſmoakie ayre. And although ſome haue boaſted of one with white flowers, yet I could neuer heare that any ſuch hath endured in one vniforme colour.

### The Place.

This plant was firſt brought from Conſtantinople into theſe Chriſtian Countries, and by the relation of ſome that ſent it, groweth naturally in Perſia.

### The Time.

It flowreth moſt commonly in the end of March, if the weather be milde, and ſpringeth not out of the ground vntill the end of February, or beginning of March, ſo quicke it is in the ſpringing: the heads with ſeed are ripe in the end of May.

### The Names.

It is of ſome called *Lilium Perſicum*, the Perſian Lilly: but becauſe wee haue another, which is more vſually called by that name, as ſhall be ſhewed in the next Chapter, I had rather with Alphonſus Pancius the Duke of Florence his Phyſitian, (who firſt ſent the figure thereof vnto M^ſ^. Iohn de Brancion) call it *Corona Imperialis*, The Crowne Imperiall, then by any other name, as alſo for that this name is now more generally receiued. It hath been ſent alſo by the name *Tuſai*, and *Tuſchai*, and *Turfani*, or *Turfanda*, being, as it is like, the Turkiſh names.

### The Vertues.

For any Phyſicall Vertues that are in it, I know of none, nor haue heard that any hath been found out: notwithſtanding the ſtrong ſent would perſwade it might be applyed to good purpoſe.

## Chap. II.

### *Lilium Perſicum.* The Perſian Lilly.

THe roote of the Perſian Lilly is very like vnto the root of the Crowne Imperiall, and loſing his fibres in like maner euery yeare, hauing a hole therin likewiſe where the old ſtalke grew, but whiter, rounder, and a little longer, ſmaller, and not ſtinking at all like it, from whence ſpringeth vp a round whitiſh greene ſtalke, not much

1 *Corona Imperialis.* The Crowne Imperiall. 2 *Lilium Persicum.* The Persian Lilly.
3 *Martagon Imperiale.* The Martagon Imperiall.

much lower than the Crowne Imperiall, but much ſmaller, beſet from the bottome to the middle thereof, with many long and narrow leaues, of a whitiſh or blewiſh greene colour, almoſt like to the leafe of a Tulipa: from the middle vpwards, to the toppe of the ſtalke, ſtand many flowers one aboue another round about it, with leaues at the foote of euery one of them, each whereof is pendulous or hanging downe the head, like vnto the Crowne Imperiall, and not turning vp any of the flowers againe, but ſmaller than in any other kinde of Lilly, yea not ſo bigge as the flower of a Fritillaria, conſiſting of ſixe leaues a peece, of a dead or ouerworne purpliſh colour, hauing in the midſt a ſmall long pointell, with certaine chiues tipt with yellow pendents: after the flowers are paſt (which abide open a long time, and for the moſt part flower by degrees, the loweſt firſt, and ſo vpwards) if the weather be temperate, come ſixe ſquare heads or ſeede veſſels, ſeeming to be but three ſquare, by reaſon of the wings, very like to the heads of the Crowne Imperiall, but ſmaller and ſhorter, wherein are contained ſuch like flat ſeed, but ſmaller alſo, and of a darker colour.

### The Place.

This was, as it is thought, firſt brought from Perſia vnto Conſtantinople, and from thence, ſent vnto vs by the meanes of diuers Turkie Merchants, and in eſpeciall, by the procurement of M^r. Nicholas Lete, a worthy Merchant, and a louer of all faire flowers.

### The Time.

It ſpringeth out of the ground very neare a moneth before the Crowne Imperiall, but doth not flower till it bee quite paſt (that is to ſay) not vntill the latter end of Aprill, or beginning of May: the ſeed (when it doth come to perfection, as it ſeldome doth) is not ripe vntill Iuly.

### The Names.

It hath been ſent by the name of *Pennachio Perſiano*, and wee thereupon doe moſt vſually call it *Lilium Perſicum*, The Perſian Lilly. Cluſius ſaith it hath been ſent into the Low-Countries vnder the name of Suſam giul, and he thereupon thinking it came from Suſis in Perſia, called it *Lilium Suſianum*, The Lilly of Suſis.

### The Vertues.

Wee haue not yet heard, that this hath beene applyed for any Phyſicall reſpect.

## CHAP. III.

### *Martagon Imperiale, ſiue Lilium Montanum maius,* The Martagon Imperiall.

VNder this title of *Lilium Montanum*, or *Lilium Silueſtre*, I do comprehend only thoſe kindes of Lillies, which carry diuers circles of greene leaues ſet together at certaine diſtances, round about the ſtalke, and not ſparſedly as the two former, and as other kindes that follow, doe. And although there bee many of this ſort, yet becauſe their chiefeſt difference is in the colour of the flower, wee will containe them all in one Chapter, and begin with the moſt ſtately of them all, becauſe of the number of flowers it beareth vpon one ſtalke. The Imperiall Lilly hath a ſcaly roote, like vnto all the reſt of the Lillies, but of a paler yellow colour, cloſely compact or ſet together, being ſhort and ſmall oftentimes, in compariſon of the greatneſſe of the

ſtemme growing from it. The ſtalke is browniſh and round at the bottome, and ſometimes flat from the middle vpwards, three foote high or more, beſet at certaine diſtances with rondles or circles of many broad leaues, larger and broader for the moſt part than any other of this kinde, and of a darke green colour: It hath two or three, and ſometimes foure of theſe rondles or circles of leaues, and bare without any leafe betweene; but aboue toward the tops of the ſtalkes, it hath here and there ſome leaues vpon it, but ſmaller than any of the other leaues: at the toppe of the ſtalke come forth many flowers, ſometime three or foure ſcore, thicke thruſt, or confuſedly ſet together, and not thinne or ſparſedly one aboue another, as in the leſſer of this kinde of Mountaine Lilly. It hath been ſometimes alſo obſerued in this kinde, that it hath borne manie flowers at three ſeuerall ſpaces of the ſtalke, one aboue another, which hath made a goodly ſhew; each flower whereof is pendulous, hanging downe, and each leafe of the flower turning vp againe, being thicke or fleſhy, of a fine delayed purple colour, ſpotted with many blackiſh or browniſh ſpots, of a very pleaſant ſweet ſent, which maketh it the more acceptable: in the middle of the flower hangeth downe a ſtile or pointell, knobbed or buttoned at the end with ſixe yellow chiues, tipt with looſe pendents of an Orient red or Vermillion colour, which will eaſily ſticke like duſt vpon any thing that toucheth them: the heads or ſeede veſſels are ſmall and round, with ſmall edges about them, wherein is contained flat browne ſeede like other Lillies, but leſſer. The root is very apt to encreaſe or ſet of, as we call it, wherby the plant ſeldome commeth to ſo great a head of flowers, but riſeth vp with many ſtalkes, and then carry fewer flowers.

Of this kinde there is ſometimes one found, that beareth flowers without any ſpots: the leaues whereof and ſtalke likewiſe are paler, but not elſe differing. *Martagon Imperiale flore non punctato.*

*Martagon flore albo.* The White Martagon.

We haue alſo ſome other of this kind, the firſt wherof hath his ſtalke & leafe greener than the former, the ſtalke is a little higher, but not bearing ſo thicke a head of flowers, although much more plentifull than the leſſer Mountaine Lilly, being altogether of a fine white colour, without any ſpots, or but very few, and that but ſometimes alſo: the pendents in the middle of this flower are not red, as the former, but yellow; the roote of this, and of the other two that follow, are of a pale yellow colour, the cloues or ſcales of them being brittle, and not cloſely compact, yet ſo as if two, and ſometimes three ſcales or cloues grew one vpon the head or vpperpart of another; which difference is a ſpeciall note to know theſe three kindes, from any other kinde of Mountaine Lilly, as in all old rootes that I haue ſeene, I haue obſerued, as alſo in them that are reaſonably well growne, but in the young rootes it is not yet ſo manifeſt.

*Martagon flore albo maculato.* The White ſpotted Martagon.

The ſecond is like vnto the firſt in all things, ſaue in this, that the flowers hereof are not altogether ſo white, and beſides hath many reddiſh ſpots on the inſide of the leaues of the flower, and the ſtalke alſo is not ſo greene but browniſh.

*Martagon flore carneo.* The bluſh Martagon.

A third ſort there is of this kinde, whoſe flowers are wholly of a delayed fleſh colour, with many ſpots on the flowers, and this is the difference hereof from the former.

*Lilium Montanum ſiue ſilueſtre minus.* The leſſer Mountaine Lilly.

The leſſer Mountaine Lilly is ſo like in root vnto the greater that is firſt deſcribed, that it is hard to diſtinguiſh them aſunder; but when this is ſprung vp out of the ground, which is a moneth after the firſt: it alſo carrieth his leaues in rondles about the ſtalke, although not altogether ſo great nor ſo many. The flowers are more thinly ſet on the ſtalkes one aboue another, with more diſtance betweene each flower than the former, and are of a little deeper fleſh colour or purple, ſpotted in the ſame manner. The buds

1 *Martagon flore albo.* The white Martagon. 2 *Martagon sive Lilium Canadense maculatum.* The spotted Martagon, or Lilly of Canada. 3 *Martagon Pomponium.* The Martagon Pompony, or early red Martagon.

or

or heads of flowers, in some of these before they be blowne, are hoary white, or hairie, whereas in others, there is no hoarinesse at all, but the buddes are smooth and purplish: in other things this differeth not from the former.

Of this sort also there is one that hath but few spots on the flowers, whose colour is somewhat paler than the other. *Lilium Montanum non maculatum.*

*Martagon Canadense maculatum.* The spotted Martagon of Canada.

Although this strange Lilly hath not his flowers hanging downe, and turning vp again, as the former kinds set forth in this Chapter; yet because the green leaues stand at seuerall ioynts as they do, I must needs insert it here, not knowing where more fitly to place it. It hath a small scaly roote, with many small long fibres thereat, from whence riseth vp a reasonable great stalke, almost as high as any of the former, bearing at three or foure distances many long and narrow greene leaues, but not so many or so broad as the former, with diuers ribbes in them: from among the vppermost rundle of leaues breake forth foure or fiue flowers together, euery one standing on a long slender foote stalke, being almost as large as a red Lilly, but a little bending downewards, and of a faire yellow colour, spotted on the inside with diuers blackish purple spots or strakes, hauing a middle pointell, and sixe chiues, with pendents on them.

## The Place.

All these Lillies haue been found in the diuers Countries of Germany, as Austria, Hungaria, Pannonia, Stiria, &c. and are all made Denisons in our London Gardens, where they flourish as in their owne naturall places. The last was was brought into France from Canada by the French Colonie, and from thence vnto vs.

## The Time.

They flower about the later end of Iune for the most part, yet the first springeth out of the ground a moneth at the least before the other, which are most vsually in flower before it, like vnto the Serotine Tulipas, all of them being early vp, and neuer the neere.

## The Names.

The first is vsually called *Martagon Imperiale*, the Imperiall Martagon, and is *Lilium Montanum maius*, the greatest Mountaine Lilly; for so it deserueth the name, because of the number of flowers vpon a head or stalke. Some haue called it *Lilium Saracenicum*, and some *Hemerocallis*, but neither of them doth so fitly agree vnto it.

The second is *Lilium Montanum maius flore albo*, and of some *Martagon Imperiale flore albo*, but most vsually *Martagon flore albo*, the white Martagon. The second sort of this second kinde, is called *Martagon flore albo maculato*, the spotted white Martagon. And the third, *Martagon flore carneo*, the blush Martagon.

The third kinde is called *Lilium Montanum*, the Mountaine Lilly, and some adde the title *minus*, the lesser, to know it more distinctly from the other. Some also *Lilium Siluestre*, as Clusius, and some others, and of Matthiolus *Martagon*. Of diuers women here in England, from the Dutch name, Lilly of Nazareth. The last hath his title *Americanum & Canadense*, and in English accordingly.

## Chap. IV.

### 1. *Martagon Pomponeum sive Lilium rubrum præcox, vel Lilium Macedonicum.*
The early red Martagon, or Martagon Pompony.

AS in the former Chapter we described vnto you such Lillies, whose flowers being pendulous, turne their leaues backe againe, and haue their greene leaues, set by spaces about the stalke: so in this wee will set downe those sorts, which carry their greene leaues more sparsedly, and all along the stalke, their flowers hanging downe, and turning vp againe as the former, and begin with that which is of greatest beauty, or at least of most rarity.

*1. Martagon Pomponeum angusti folium præcox.*

1. This rare Martagon hath a scaly root closely compact, with broader and thinner scales than others, in time growing very great, and of a more deepe yellow colour then the former, from whence doth spring vp a round greene stalke in some plants, and flat in others, two or three foote high, bearing a number of small, long, and narrow greene leaues, very like vnto the leaues of Pinkes, but greener, set very thicke together, and without order about the stalke, vp almost vnto the toppe, and lesser by degrees vpwards, where stand many flowers, according to the age of the plant, and thriuing in the place where it groweth; in those that are young, but a few, and more sparsedly, and in others that are old many more, and thicker set: for I haue reckoned threescore flowers and more, growing thicke together on one plant with mee, and an hundred flowers on another: these flowers are of a pale or yellowish red colour, and not so deep red as the red Martagon of Constantinople, hereafter set down, nor fully so large: yet of the same fashion, that is, euery flower hanging downe, and turning vp his leaues againe. It is not so plentifull in bearing of seede as the other Lillies, but when it doth, it differeth not but in being lesse.

*2. Martagon angusti folium magis serotinum.*

There is another, whose greene leaues are not so thicke set on the stalke, but else differeth not but in flowring a fortnight later.

*3. Martagon Pomponeum latifolium præcox.*

There is another also of this kind, so like vnto the former in root, stalk, flower, & maner of growing, that the difference is hardly discerned; but consisteth chiefly in these two points: First, that the leaues of this are a little broader and shorter then the former; and secondly, that it beareth his flowers a fortnight earlier than the first. In the colour or forme of the flower, there can no difference bee discerned, nor (as I said) in any other thing. All these Lillies doe spring very late out of the ground, euen as the yellow Martagons doe, but are sooner in flower then any others.

*4. Martagon flore phœniceo.*

A fourth kinde hereof hath of late been knowne to vs, whose leaues are broader and shorter then the last, and the flowers of a paler red, tending to yellow, of some called a golden red colour: but flowreth not so early as they.

### 2. *Lilium rubrum Byzantinum, sive Martagon Constantinopolitanum.*
The red Martagon of Constantinople.

1. The red Martagon of Constantinople is become so common euery where, and so well knowne to all louers of these delights, that I shall seeme vnto them to lose time, to bestow many lines vpon it; yet because it is so faire a flower, and was at the first so highly esteemed, it deserueth his place and commendations, howsoeuer encreasing the plenty hath not made it dainty. It riseth out of the ground early in the spring, before many other Lillies, from a great thicke yellow scaly root, bearing a round brownish stalke, beset with many faire greene leaues confusedly thereon, but not so broad as the common white Lilly, vpon the toppe whereof stand one, two, or three, or more flowers, vpon long foote stalkes, which hang downe their heads, and turne vp their leaues againe, of an excellent red crimson colour, and sometimes paler, hauing a long pointell in the middle, compassed with sixe whitish chiues, tipt with loose yellow pendents, of a reasonable good sent, but somewhat faint. It likewise beareth seede in heads, like vnto the other, but greater.

*Martagon Constantinopolitanum maculatum.*
The red spotted Martagon of Constantinople.

We haue another of this kinde, that groweth somewhat greater and higher, with a larger flower, and of a deeper colour, spotted with diuers blacke spots, or strakes and lines, as is to be seene in the Mountaine Lillies, and in some other hereafter to be described; but is not so in the former of this kinde, which hath no shew of spots at all. The whole plant as it is rare, so it is of much more beauty then the former.

2. *Martagon Pannonicum, sive Exoticum flore spadicee.*
The bright red Martagon of Hungarie.

Although this Martagon or Lilly bee of another Countrey, yet by reason of the neerenesse both in leafe and flower vnto the former, may more fitly be placed next vnto them, then in any other place. It hath his roote very like the other, but the leaues are somewhat larger, and more sparsedly set vpon the stalke, else not much vnlike: the flowers bend downe, and turne vp their leaues againe, but somewhat larger, and of a bright red, tending to an Orenge colour, that is, somewhat yellowish, and not crimson, like the other.

3. *Martagon Luteum punctatum.* The Yellow spotted Martagon.

1. This Yellow Martagon hath a great scaly or cloued roote, and yellow, like vnto all these sorts of turning Lillies, from whence springeth vp a round greene strong stalke, three foote high at the least, confusedly set with narrow long greene leaues, white on the edges vp to the very toppe thereof almost, hauing diuers flowers on the head, turning vp againe as the former doe, of a faint yellowish, or greenish yellow colour, with many blacke spots or strakes about the middle of the leafe of euery flower, and a forked pointell, with sixe chines about it, tipt with reddish pendents, of a heauie strong smell, not very pleasant to many. It beareth seede very plentifully, in great heads, like vnto the other former Lillies, but a little paler.

2. *Martagon Luteum non maculatum.* The Yellow Martagon without spots.

The other yellow Martagon differeth in no other thing from the former, but onely that it hath no spots at all vpon any of the leaues of the flowers; agreeing with the former, in colour, forme, height, and all things else.

3. *Martagon Luteum serotinum.* The late flowring Yellow Martagon.

There is yet another yellow Martagon, that hath no other difference then the time of his flowring, which is not vntill Iuly, vnlesse in this, that the flower is of a deeper yellow colour.

The Place.

The knowledge of the first kindes of these early Martagons hath come from Italy, from whence they haue bin sent into the Low-Countries, and to vs, and, as it seemeth by the name, whereby they haue bin sent by some into these parts, his originall should be from the mountaines in Macedonia.

The second sort is sufficiently knowne by his name, being first brought from Constantinople, his naturall place being not farre from thence, as it is likely. But the next sort of this second kinde, doth plainly tell vs his place of birth to be the mountaines of Pannonia or Hungarie.

The third kindes grow on the Pyrenæan mountaines, where they haue been searched out, and found by diuers louers of plants, as also in the Kingdome of Naples.

The

### The Time.

The first early Martagons flower in the end of May, or beginning of Iune, and that is a moneth at the least before those that come from Constantinople, which is the second kinde. The two first yellow Martagons flower somewhat more early, then the early red Martagons, and sometimes at the same time with them. But the third yellow Martagon, as is said, flowreth a moneth later or more, and is in flower when the red Martagon of Constantinople flowreth. And although the early red and yellow Martagons, spring later then the other Martagons or Lillies, yet they are in flower before them.

### The Names.

The first early red Lillies or Martagons haue beene sent vnto vs by seuerall names, as *Martagon Pomponeum*, and thereafter are called Martagon of Pompony, and also *Lilium* or *Martagon Macedonicum*, the Lilly or Martagon of Macedonia. They are also called by Clusius *Lilium rubrum præcox*, the one *angustiore folio*, the other *latiore folio*. And the last of this kinde hath the title *flore phœnicco* added or giuen vnto it, that is, the Martagon or Lilly of Macedonia with gold red flowers.

The Martagons of Constantinople haue beene sent by the Turkish name *Zufiniare*, and is called *Martagon*, or *Lilium Byzantinum* by some, and *Hemerocallis Chalcedonica* by others; but by the name of the Martagon of Constantinople they are most commonly receiued with vs, with the distinction of *maculatum* to the one, to distinguish the sorts. The last kinde in this *classis*, hath his name in his title, as it hath been sent vnto vs.

The Yellow Martagons are distinguished in their seuerall titles, as much as is conuenient for them.

## Chap. V.

*Lilium Aureum & Lilium Rubrum.* The Gold and Red Lillies.

There are yet some other kindes of red Lillies to bee described, which differ from all the former, and remaine to be spoken of in this place. Some of them grow high, and some lowe, some haue small knots, which wee call bulbes, growing vpon the stalkes, at the ioynts of the leaues or flowers, and some haue none: all which shall be intreated of in their seuerall orders.

*Lilium pumilum cruentum.* The dwarfe red Lilly.

The dwarfered Lilly hath a scaly roote, somewhat like vnto other Lillies, but white, and not yellow at all, and the cloues or scales thicker, shorter, and fewer in number, then in most of the former: the stalke hereof is not aboue a foote and a halfe high, round and greene, set confusedly with many faire and short greene leaues, on the toppe of which doe stand sometimes but a few flowers, and sometimes many, of a faire purplish red colour, and a little paler in the middle, euery flower standing vpright, and not hanging downe, as in the former, on the leaues whereof here and there are some blacke spots, lines or markes, and in the middle of the flower a long pointell, with some chiues about it, as is in the rest of these Lillies.

*Lilium rubrum multiplici flore.* This kinde is sometimes found to yeeld double flowers, as if all the single flowers should grow into one, and so make it consist of many leaues, which notwithstanding his

1 *Martagon rubrum siue luteum.* The red or the yellow Martagon. 2 *Lilium Bulbiferum.* The red bulbed Lilly. 3 *Lilium aureum.* The gold red Lilly. 4 *Lilium album.* The white Lilly.

his so continuing sundry yeares, vpon transplanting, will *redire ad ingenium*, that is, quickly come againe to his old byas or forme.

*Lilium Aureum.* The Gold red Lilly.

The second red Lilly without bulbes groweth much higher then the first, and almost as high as any other Lilly: the roote hereof is white and scaly, the leaues are somewhat longer, and of a darke or sad greene colour; the flowers are many and large, standing vpright as all these sorts of red Lillies doe, of a paler red colour tending to an Orenge on the inside, with many blacke spots, and lines on them, as in the former, and more yellow on the outside: the seede vessels are like vnto the roundish heads of other Lillies, and so are the seedes in them likewise.

1. *Lilium minus bulbiferum.* The dwarfe bulbed Lilly.

The first of the Lillies that carrieth bulbes on the stalke, hath a white scaly roote like the former; from whence riseth vp a small round stalke, not much higher then the first dwarfe Lilly, seeming to be edged, hauing many leaues thereon of a sad green colour set about it, close thrust together: the greene heads for flowers, will haue a kind of woollinesse on them, before the flowers begin to open, and betweene these heads of flowers, as also vnder them, and among the vppermost leaues, appeare small bulbes or heads, which being ripe if they be put into the ground, or if they fall of themselues, will shoote forth leaues, and beare flowers within two or three yeares like the mother plant, and so will the bulbes of the other hereafter described: the flowers of this Lilly are of a faire gold yellow colour, shadowed ouer with a shew of purple, but not so red as the first, or the next to bee described. This Lilly will shoote strings vnder ground, like as the last red Lilly will doe also, whereat will grow white bulbed roots, like the rootes of the mother plant, thereby quickly encreasing it selfe.

2. *Lilium Cruentum bulbiferum.* The Fierie red bulbed Lilly.

The second bulbed Lilly riseth vp with his stalke as high as any of these Lillies, carrying many long and narrow darke greene leaues about it, and at the toppe many faire red flowers, as large or larger then any of the former, and of a deeper red colour, with spots on them likewise, hauing greater bulbes growing about the toppe of the stalke and among the flowers, then any else.

*Lilium Cruentum flore pleno.* The Fierie red double Lilly.

The difference of this doth chiefly consist in the flower, which is composed of manie leaues, as if many flowers went to make one, spotted with black spots, and without any bulbes when it thus beareth, which is but accidentall, as the former double Lilly is said to be.

3. *Lilium maius bulbiferum.* The greater bulbed red Lilly.

The third red Lilly with bulbes, riseth vp almost as high as the last, and is the most common kinde we haue bearing bulbes. It hath many leaues about the stalke, but not of so sad a greene colour as the former: the flowers are of as pale a reddish yellow colour as any of the former, and comming neerest vnto the colour of the Gold red Lilly. This is more plentifull in bulbes, and in shooting strings, to encrease rootes vnder ground, then the others.

The Place.

These Lillies doe all grow in Gardens, but their naturall places of growing is the Mountaines and the Vallies neere them in Italy, as Matthiolus saith:

saith: and in many Countries of Germany, as Hungarie, Austria, Stiria, and Bohemia, as Clusius and other doe report.

### The Time.

They flower for the most part in Iune, yet the first of these is the earliest of all the rest.

### The Names.

All these Lillies are called *Lilia Rubra*, Red Lillies: Some call them *Lilium Aureum, Lilium Purpureum, Lilium Puniceum, & Lilium Cruentum.* Some also call them *Martagon Chimistarum.* Clusius calleth these bulbed Lillies *Martagon Bulbiferum.* It is thought to be *Hyacinthus Poetarum*, but I referre the discussing thereof to a fitter time. Wee haue, to distinguish them most fitly (as I take it) giuen their proper names in their seuerall titles.

## CHAP. VI.

### *Lilium Album.* The White Lilly.

NOw remaineth onely the White Lilly, of all the whole family or stocke of the Lillies, to bee spoken of, which is of two sorts. The one is our common or vulgar White Lilly; and the other, that which was brought from Constantinople.

### *Lilium Album vulgare.* The ordinary White Lilly.

The ordinary White Lilly scarce needeth any description, it is so well knowne, and so frequent in euery Garden; but to say somewhat thereof, as I vse to doe of euery thing, be it neuer so common and knowne; it hath a cloued or scaly roote, yellower and bigger then any of the red Lillies: the stalke is of a blackish greene colour, and riseth as high as most of the Lillies, hauing many faire, broad, and long greene leaues thereon, larger and longer beneath, and smaller vpon the stalke vpwards; the flowers are many or few, according to the age of the plant, fertility of the soile, and time of standing where it groweth: and stand vpon long greene foote stalkes, of a faire white colour, with a long pointell in the middle, and white chiues tipt with yellow pendents about it; the smell is somewhat heady and strong.

### *Lilium Album Byzantinum.* The White Lilly of Constantinople.

The other White Lilly, differeth but little from the former White Lilly, either in roote, leafe, or flower, but only that this vsually groweth with more number of flowers, then euer we saw in our ordinary White Lilly: for I haue seene the stalke of this Lilly turne flat, of the breadth of an hand, bearing neere two hundred flowers vpon a head, yet most commonly it beareth not aboue a dozen, or twenty flowers, but smaller then the ordinary, as the greene leaues are likewise.

### The Place.

The first groweth onely in Gardens, and hath not beene declared where it is found wilde, by any that I can heare of. The other hath beene sent from Constantinople, among other rootes, and therefore is likely to grow in some parts neere thereunto.

### The Time.

They flower in Iune or thereabouts, but shoote forth greene leaues in

Autumne, which abide greene all the Winter, the stalke springing vp betweene the lower leaues in the Spring.

The Names.

It is called *Lilium Album*, the White Lilly, by most Writers; but by Poets *Rosa Iunonis*, Iuno's Rose. The other hath his name in his title.

The Vertues.

This Lilly aboue all the rest, yea, and I thinke this onely, and none of the rest is vsed in medicines now adayes, although in former times Empericks vsed the red; and therefore I haue spoken nothing of them in the end of their Chapters, reseruing what is to be said in this. This hath a mollifying, digesting, and cleansing quality, helping to suppurate tumours, and to digest them, for which purpose the roote is much vsed. The water of the flowers distilled, is of excellent vertue for women in trauell of childe bearing, to procure an easie deliuery, as Matthiolus and Camerarius report. It is vsed also of diuers women outwardly, for their faces to cleanse the skin, and make it white and fresh. Diuers other properties there are in these Lillies, which my purpose is not to declare in this place. Nor is it the scope of this worke; this that hath been said is sufficient: for were it not, that I would giue you some taste of the qualities of plants (as I said in my Preface) as I goe along with them, a generall worke were fitter to declare them then this.

## Chap. VII.

### *Fritillaria.* The checkerd Daffodill.

ALthough diuers learned men do by the name giuen vnto this delightfull plant, thinke it doth in some things partake with a Tulipa or Daffodill, and haue therefore placed it betweene them; yet I, finding it most like vnto a little Lilly, both in roote, stalke, leafe, flower, and seede, haue (as you see here) placed it next vnto the Lillies, and before them. Hereof there are many sorts found out of late, as white, red, blacke, and yellow, besides the purple, which was first knowne; and of each of them there are also diuers sorts: and first of that which is most frequent, and then of the rest, euery one in his place and order.

#### 1. *Fritillaria vulgaris.* The common checkerd Daffodill.

The ordinary checkerd Daffodill (as it is vsually called, but might more properly be called the small checkerd Lilly) hath a small round white roote, and somewhat flat, made as it were of two cloues, and diuided in a maner into two parts, yet ioyning together at the bottome or seate of the roote, which holdeth them both together: from betweene this cleft or diuision, the budde for the stalke &c. appeareth, which in time riseth vp a foote, or a foote and a halfe high, being round and of a browniſh greene colour, especially neere vnto the ground, whereon there standeth dispersedly foure or fiue narrow long and greene leaues, being a little hollow: at the toppe of the stalke, betweene the vpper leaues (which are smaller then the lowest) the flower sheweth it selfe, hanging or turning downe the head, but not turning vp againe any of his leaues, as some of the Lillies before described doe; (sometimes this stalke beareth two flowers, and very seldome three) consisting of fixe leaues, of a reddish purple colour, spotted diuersly with great spots, appearing like vnto square checkers, of a deeper colour; the inside of the flower is of a brighter colour then the outside, which hath some greennesse at the bottome of euery leafe: within the flower there appeare

fixe

1 *Fritillaria vulgaris.* The common Fritillaria. 2 *Fritillaria flore atrorubente.* The darke red Fritillaria. 4 *Fritillaria alba.* The white Fritillaria. 7 *Fritillaria lutea punctata.* The yellow checkerd Fritillaria. 8 *Fritillaria lutea Italica.* The great yellow Italian Fritillaria. 10 *Fritillaria lutea Lusitanica.* The small yellow Fritillaria of Portugall. 11 *Fritillaria Pyrenea.* The blacke Fritillaria. 12 *Fritillaria umbellifera.* The Spanish blacke Fritillaria.

ſixe chiues tipt with yellow pendents, and a three-forked ſtile or pointell compaſſing a greene head, which when the flower is paſt, riſeth vpright againe, and becommeth the ſeede veſſell, being ſomewhat long and round, yet hauing a ſmall ſhew of edges, flat at the head, like the head of a Lilly, and without any crowne as the Tulipa hath, wherein is contained pale coloured flat ſeede, like vnto a Lilly, but ſmaller.

*Fritillaria vulgaris pallidior, præcox, & ſerotina.*

There is ſome variety to be ſeene in this flower; for in ſome the colour is paler, and in others againe of a very high or deepe colour: ſometimes alſo they haue eight leaues, and ſometimes ten or twelue, as if two flowers were made one, which ſome thereupon haue called a Double Fritillaria. Some of them likewiſe doe flower very early, euen with or before the early flowring Tulipas; and ſome againe flower not vntill a moneth or more after the former.

2. *Fritillaria flore atrorubente.* The bloud red Fritillaria.

The roote of this Fritillaria is ſomewhat rounder and cloſer then the former, from whence the ſtalke riſeth vp, being ſhorter and lower then in any other of theſe kindes, hauing one or two leaues thereon, and at the top thereof two or three more ſet cloſer together, which are broader, ſhorter, and whiter then any of them before, almoſt like vnto the leaues of the yellow Fritillaria, from among which toppe leaues commeth forth the flower, ſomewhat bending downe, or rather ſtanding forth, being larger then any of the former, and almoſt equall in bigneſſe vnto the yellow Fritillaria, of a duskie gray colour all ouer on the outſide, and of a very darke red colour on the inſide, diuerſly ſpotted or ſtraked: this very hardly encreaſeth by the roote, and as ſeldome giueth ripe ſeede, but flowreth with the other firſt ſorts, and before the blacke, and abideth leſſe time in flower then any.

3. *Fritillaria maxima purpurea ſiue rubra.* The great purple or red Fritillaria.

This great Fritillaria hath his roote equall to the bigneſſe of the reſt of his parts, from whence riſeth vp one, & oftentimes two ſtalks, hauing one, two or three flowers a peece on them, as nature and the ſeaſons are fitting: euery one of theſe flowers are larger and greater then any of the former deſcribed, and pendulous as they are, of a ſad red or purpliſh colour, with many thwart lines on them, and ſmall long markes, which hardly ſeeme checkerwiſe, nor are ſo eminent or conſpicuous as in the former: the ſtalke is ſtrong and high, whereon are ſet diuers long whitiſh greene leaues, larger and broader then thoſe of the former.

4. *Fritillaria alba.* The white Fritillaria.

The white Fritillaria is ſo like vnto the firſt, that I ſhall not neede to make another deſcription of this: it ſhall (I hope) be ſufficient to ſhew the chiefe differences, and ſo proceed to the reſt. The ſtalke and leaues of this are wholly greene, whereby it may eaſily be knowne from the former, which, as is ſaid, is browniſh at the bottome. The flower is white, without almoſt any ſhew of ſpot or marke in it, yet in ſome the markes are ſomewhat more plainly to be ſeene, and in ſome againe there is a ſhew of a faint kinde of bluſh colour to be ſeene in the flower, eſpecially in the inſide, the bottomes of the leaues of euery flower ſometimes are greeniſh, hauing alſo a ſmall liſt of greene, comming downe towards the middle of each leafe: the head or ſeede veſſell, as alſo the ſeede and the roote, are ſo like vnto the former, that the moſt cunning cannot diſtinguiſh them.

5. *Fritillaria flore duplici albicante.* The double bluſh Fritillaria.

This Fritillaria hath a round flattiſh white roote, very like vnto the laſt Fritillaria, bearing a ſtalke with long greene leaues thereon, little differing from it, or the firſt ordinary Fritillaria: the flower is ſaid to be conſtant, compoſed of many leaues, being ten at the leaſt, and moſt vſually twelue, of a pale whitiſh purple colour, ſpotted like vnto the paler ordinary Fritillaria that is early, ſo that one would verily thinke it were



but

but an accidentall kinde thereof, whereas it is (as is ſaid before) held to bee conſtant, continuing in this manner.

6. *Fritillaria flore luteo paro.* The pure yellow Fritillaria.

The pure yellow Fritillaria hath a more round, and not ſo flat a whitiſh roote as the former kindes, and of a meane bigneſſe; from the middle riſeth vp a ſtalke a foote and a halfe high, and ſometimes higher, whereon are ſet without order diuers long and ſomewhat broad leaues of a whitiſh greene colour, like vnto the leaues of the blacke Fritillaria, but not aboue halfe ſo broad: the flower is ſomewhat ſmall and long, not much vnlike to the blacke for ſhape and faſhion, but that the leaues are ſmaller and rounder pointed, of a faint yellowiſh colour, without any ſhew of ſpots or checkers at all, eyther within or without the flower, hauing ſome chiues and yellow pendents in the middle, as is to be ſeene in all of them: the ſeede is like the firſt kinde.

7. *Fritillaria flore luteo vario ſiue punctato.* The checkerd yellow Fritillaria.

This Fritillaria groweth not much lower then the former, and browniſh at the riſing vp, hauing his leaues whiter, broader, and ſhorter then it, and almoſt round pointed. The flower is greater, and larger ſpread then any other before, of a faire pale yellow colour, ſpotted in very good order, with fine ſmall checkers, which adde a wonderfull pleaſing beauty thereunto: it hath alſo ſome liſts of greene running downe the backe of euery leafe. It ſeldome giueth ſeede; the roote alſo is like the other, but not ſo flat.

8. *Fritillaria lutea maxima Italica.* The great yellow Italian Fritillaria.

This kinde of Fritillaria riſeth vp with a round and browne greene ſtalke, whereon are ſet diuers leaues ſomewhat broad and ſhort, which compaſſe the ſtalke at the bottome of them, of a darke greene colour; at the toppe of the ſtalke, which bendeth a little downewards, doe moſt vſually ſtand three or foure leaues, betweene which commeth forth moſt vſually but one flower, which is longer then the laſt, hanging downe the head as all the others doe, conſiſting of ſixe leaues, of a darke yellowiſh purple colour, ſpotted with ſome ſmall red checkers. This kinde flowreth late, and not vntill all the reſt are paſt.

9. *Fritillaria Italorum polyanthos flore paruo.* The ſmall Italian Fritillaria.

This ſmall Italian Fritillaria carrieth more ſtore of flowers on the ſtalke, but they are much ſmaller, and of a yellowiſh greene colour, ſpotted with long and ſmall darke red checkers or markes: the ſtalke hath diuers ſmall ſhort greene leaues thereon, vnto the very toppe.

10. *Fritillaria lutea Iuncifolia Luſitanica.* The ſmall yellow Fritillaria of Portugall.

The leaues of this Fritillaria are ſo ſmall, narrow and long, that it hath cauſed them to take the name of ruſhes, as if you ſhould call it, The ruſh leafed Fritillaria, which ſtand on a long weake round ſtalke, ſet without order: the flower is ſmall and yellow, but thicker checkerd with red ſpots then any of the other yellow Fritillaria's; the ſtalk of the flower, at the head thereof, being alſo of a yellowiſh colour.

11. *Fritillaria Pyrenæa ſiue Apenninea.* The blacke Fritillaria.

The roote of this kinde doth often grow ſo great, that it ſeemeth like vnto the roote of a ſmall Crowne Imperiall: the ſtalke is ſtrong, round, and high, ſet without order, with broader and whiter greene leaues then any of the former, bearing one, two, or three flowers; ſometimes at the toppe, being not ſo large as thoſe of the ordinary purple Fritillaria, but ſmaller, longer, and rounder, ſometimes a little turning vp the brims or edges of the leaues againe, and are of a yellowiſh ſhining greene colour on

the

the inside, sometimes spotted with red spots almost through the whole inside of the flower, vnto the very edge, which abideth of a pale yellow colour, and sometimes there are very few spots to be seene, and those from the middle onely on the inside (for on the outside there neuer appeareth any spots at all in this kinde) and sometimes with no shew of spots at all, sometimes also of a more pale greene, and sometime of a more yellow colour: the outside of the flowers doe likewise vary, for in some the outside of the leaues are of a darke sullen yellow, &c. else more pale yellow, and in other of a darke purplish yellow colour, which in some is so deepe, and so much, that it rather seemeth blacke then purple or yellow, and this especially about the bottome of the flower, next vnto the stalke, but the edges are still of a yellowish greene: the head of seede, and the seede likewise is like vnto the former, but bigger in all respects.

12. *Fritillaria Hispanica vmbellifera.* The Spanish blacke Fritillaria.

This Fritillaria is no doubt of kindred to the last recited, it is so like, but greater in all parts thereof, as if growing in a more fruitfull soile, it were the stronger and lustier to beare more store of flowers: the flowers grow foure or fiue from the head together, hanging downe round about the stalke, like vnto a Crowne Imperiall, and are of a yellowish greene colour on the inside, spotted with a few red spots, the outside being blackish as the former.

The Place.

The first of these plants was first brought to our knowledge from France, where it groweth plentifully about Orleance; the other sorts grow in diuers other Countries, as some in Portugall, Spaine, Italy, &c. as their names doe import, and as in time they haue been obserued by those that were curious searchers of these rarities, haue been sent to vs.

The Time.

The early kindes doe flower in the beginning of Aprill or thereabouts, according to the mildenesse or sharpenesse of the precedent Winter. The other doe flower after the first are past, for a moneths space one after another, and the great yellow is very late, not flowring vntill about the middle or end of May.

The Names.

This hath receiued diuers names: some calling it *Flos Meleagridis*, the Ginny Hen Flower, of the variety of the colours in the flower, agreeing with the feathers of that Bird. Some call it *Narcissus Caparonius*, of the name of the first inuentor or finder thereof, called Noel Caperon, an Apothecary dwelling in Orleance, at the time he first found it, and was shortly after the finding thereof taken away in the Massacre in France. It is now generally called *Fritillaria*, of the word *Fritillus*, which diuers doe take for the Chesse borde or table whereon they play, whereunto, by reason of the resemblance of the great squares or spots so like it, they did presently referre it. It is called by Lobel *Lilionarcissus purpureus variegatus, & tessulatus*, making it a kinde of Tulipa; but as I said in the beginniug of the Chapter, it doth most neerely resemble a small pendulous Lilly, and might therefore rightly hold the name of *Lilium variegatum*, or in English, the checkerd Lilly. But because the errour which first referred it to a Daffodill, is growne strong by custome of continuance, I leaue to euery one their owne will, to call it in English eyther Fritillaria, as it is called of most, or the checkerd Daffodill, or the Ginnie Hen flower, or, as I doe, the checkerd Lilly. I shall not neede in this place further to explaine the seuerall names of euery of them, hauing giuen you them in their titles.

The

### The Vertues.

I haue not found or heard by any others of any property peculiar in this plant, to be applied either inwardly or outwardly for any disease: the chiefe or onely vse thereof is, to be an ornament for the Gardens of the curious louers of these delights, and to be worne of them abroad, which for the gallant beauty of many of them, deserueth their courteous entertainment, among many other the like pleasures.

## Chap. VIII.

### *Tulipa.* The Turkes Cap.

NExt vnto the Lillies, and before the Narcissi or Daffodils, the discourse of Tulipas deserueth his place, for that it partaketh of both their natures; agreeing with the Lillies in leaues, flowers, and seede, and somewhat with the Daffodils in rootes. There are not onely diuers kindes of Tulipas, but sundry diuersities of colours in them, found out in these later dayes by many the searchers of natures varieties, which haue not formerly been obserued: our age being more delighted in the search, curiosity, and rarities of these pleasant delights, then any age I thinke before. But indeede, this flower, aboue many other, deserueth his true commendations and acceptance with all louers of these beauties, both for the stately aspect, and for the admirable varietie of colours, that daily doe arise in them, farre beyond all other plants that grow, in so much, that I doubt, although I shall in this Chapter set downe the varieties of a great many, I shall leaue more vnspoken of, then I shall describe; for I may well say, there is in this one plant no end of diuersity to be expected, euery yeare yeelding a mixture and variety that hath not before been obserued, and all this arising from the sowing of the seede. The chiefe diuision of Tulipas, is into two sorts: *Præcoces*, early flowring Tulipas, and *Serotinæ*, late flowring Tulipas. For that sort which is called *Mediæ* or *Dubiæ*, that is, which flower in the middle time betweene them both, and may be thought to be a kinde or sort by it selfe, as well as any of the other two: yet because they doe neerer participate with the *Serotinæ* then with the *Præcoces*, not onely in the colour of the leafe, being of the same greennesse with the *Serotinæ*, and most vsually also, for that it beareth his stalke and flower, high and large like as the *Serotinæ* doe; but especially, for that the seede of a *Media Tulipa* did neuer bring forth a *Præcox* flower (although I know Clusius, an industrious, learned, and painfull searcher and publisher of these rarities, saith otherwise) so farre as euer I could, by mine owne care or knowledge, in sowing their seede apart, or the assurance of any others, the louers and sowers of Tulipa seede, obserue, learne, or know: and because also that the seede of the *Serotinæ* bringeth forth *Medias*, and the seede of *Medias Serotinæ*, they may well bee comprehended vnder the generall title of *Serotinæ*: But because they haue generally receiued the name of *Mediæ*, or middle flowring Tulipas, to distinguish betweene them, and those that vsually doe flower after them; I am content to set them downe, and speake of them seuerally, as of three sorts. Vnto the place and ranke likewise of the *Præcoces*, or early flowring Tulipas, there are some other seuerall kinds of Tulipas to be added, which are notably differing, not onely from the former *Præcox Tulipa*, but euery one of them, one from another, in some speciall note or other: as the *Tulipa Bolonienſis flore rubro*, the red Bolonia Tulipa. *Tulipa Bolonienſis flore luteo*, the yellow Bolonia Tulipa. *Tulipa Perſica*, the Persian Tulipa. *Tulipa Cretica*, the Candie Tulipa, and others: all which shall bee described and entreated of, euery one apart by it selfe, in the end of the ranke of the *Præcoces*, because all of them flower much about their time. To begin then with the *Præcox*, or early flowring Tulipas, and after them with the *Medias* and *Serotines*, I shall for the better method, diuide their flowers into foure primary or principall colours, that is to say, White, Purple, Red, and Yellow, and vnder euery one of these colours, set downe the seuerall varie-

ties

ties of mixtures we haue seene and obserued in them, that so they may be both the better described by me, and the better conceiued by others, and euery one placed in their proper ranke. Yet I shall in this, as I intend to doe in diuers other plants that are variable, giue but one description in generall of the plant, and then set downe the varietie of forme or colour afterwards briefly by themselues.

*Tulipa præcox.* The early flowring Tulipa.

The early Tulipa (and so all other Tulipas) springeth out of the ground with his leaues folded one within another, the first or lowest leafe riseth vp first, sharpe pointed, and folded round together, vntill it be an inch or two aboue the ground, which then openeth it selfe, shewing another leafe folded also in the bosome or belly of the first, which in time likewise opening it selfe, sheweth forth a third, and sometimes a fourth and a fifth: the lower leaues are larger then the vpper, and are faire, thicke, broad, long, and hollow like a gutter, and sometimes crumpled on the edges, which will hold water that falleth thereon a long time, of a pale or whitish greene colour, (and the *Mediæ* and *Serotinæ* more greene) couered ouer as it were with a mealinesse or hoarinesse, with an eye or shew of rednesse towards the bottome of the leaues, and the edges in this kinde being more notable white, which are two principall notes to know a *Præcox Tulipa* from a *Media* or *Serotina*: the stalke with the flower riseth vp in the middle, as it were through these leaues, which in time stand one aboue another, compassing it at certaine vnequall distances, and is often obserued to bend it selfe crookedly downe to the ground, as if it would thrust his head thereinto, but turning vp his head (which will be the flower) againe, afterwards standeth vpright, sometimes but three or foure fingers or inches high, but more often halfe a foote, and a foot high, but the *Mediæ*, and *Serotinas* much higher, carrying (for the most part) but one flower on the toppe thereof, like vnto a Lilly for the forme, consisting of sixe leaues, greene at the first, and afterwards changing into diuers and sundry seuerall colours and varieties, the bottomes likewise of the leaues of these sometimes, but most especially of the *Mediæ*, being as variable as the flower, which are in some yellow, or green, or blacke, in others white, blew, purple, or tawnie; and sometimes one colour circling another: some of them haue little or no sent at all, and some haue a better then others. After it hath been blowne open three or foure dayes or more, it will in the heate of the Sunne spread it selfe open, and lay it selfe almost flat to the stalke: in the middle of the flower standeth a greene long head (which will be the seed vessell) compassed about with sixe chiues, which doe much vary, in being sometimes of one, and sometimes of another colour, tipt with pendents diuersly varied likewise: the head in the middle of the flower groweth after the flower is fallen, to be long, round, and edged, as it were three square, the edges meeting at the toppe, where it is smallest, and making as it were a crowne (which is not seen in the head of any Lilly) and when it is ripe, diuideth it selfe on the inside into sixe rowes, of flat, thinne, browniſh, gristly seede, very like vnto the seede of the Lillies, but brighter, stiffer, and more transparent: the roote being well growne is round, and somewhat great, small and pointed at the toppe, and broader, yet roundish at the bottome, with a certaine eminence or seate on the one side, as the roote of the Colchicum hath; but not so long, or great, it hath also an hollownesse on the one side (if it haue borne a flower) where the stalke grew, (for although in the time of the first springing vp, vntill it shew the budde for flower, the stalke with the leaues thereon rise vp out of the middle of the roote; yet when the stalke is risen vp, and sheweth the budde for flower, it commeth to one side, making an impression therein) couered ouer with a browniſh thin coate or skin, like an Onion, hauing a little woollinesse at the bottome; but white within, and firme, yet composed of many coates, one folding within another, as the roote of the Daffodils be, of a reasonable good taste, neyther very sweete, nor yet vnpleasant. This description may well serue for the other Tulipas, being *Mediæ* or *Serotinas*, concerning their springing and bearing, which haue not any other great variety therein worth the note, which is not expressed here; the chiefe difference resting in the variety of the colours of the flower, and their seuerall mixtures and markes, as I said before: sauing onely, that the flowers of some are great and large, and of others smaller, and the leaues of some long

and

1 *Tulipa præcox alba sine rubra, &c. unius coloris.* The early white or red Tulipa, &c. being of one colour. 2 *Tulipa præcox purpurea oris albis.* The early purple Tulipa with white edges, or the Prince. 3 *Tulipa præcox variegata.* The early stript Tulipa. 4 *Tulipa præcox rubra oris luteis.* The early red Tulipa with yellow edges, or the Duke.

and pointed, and of others broad and round, or bluntly pointed, as shall bee shewed in the end of the Chapter: I shall therefore onely expresse the colours, with the mixture or composure of them, and giue you withall the names of some of them, (for it is impossible I thinke to any man, to giue seuerall names to all varieties) as they are called by those that chiefly delight in them with vs.

*Tulipa præcox Alba.*

1 *Niuea tota interdum purpureis staminibus, vel saltem luteis, fundo puro haud luteo.*
2 *Alba siue niuea fundo luteo.*
3 *Albida.*
4 *Alba, venis cæruleis in dorso.*
5 *Alba purpureis oris.* } *Harum flores vel*
6 *Alba carneis oris.* } *constantes, vel*
7 *Alba sanguineis oris.* } *dispergentes.*
8 *Alba oris magnis carneis, & venis intro respicientibus.*
9 *Alba extra, carnei vero coloris intus, oras habens carneas saturatiores.*
10 *Albida, oris rubris, vel oris purpureis.*
11 *Albida purpurascentibus maculis extra, intus vero carnei vinacissimi.*
12 *Alba, purpureis maculis aspersa extra, intus vero alba purpurantibus oris.*
13 *Dux Alba, i. e. coccineis & albis variata flammis, à medio ad oras intercursantibus.*
14 *Princessa, i.e. argentei coloris maculis purpurascentibus.*
15 *Regina pulcherrima, albis & sanguineis aspersa radijs & punctis.*

The early White Tulipa.

1 The flower whereof is either pure snow white, with purple sometimes, or at least with yellow chiues, without any yellow bottome.
2 Or pure white with a yellow bottome.
3 Or milk white that is not so pure white.
4 White with blew veines on the outside.
5 White with purple edges. } Some of these abiding constant, & others spreading or running.
6 White with blush edges. }
7 White with red edges. }
8 White with great blush edges, and some strakes running from the edge inward.
9 White without, and somewhat blush within, with edges of a deeper blush.
10 Whitish, or pale white with red or purple edges.
11 Whitish without, with some purplish veins & spots, & of a liuely blush within.
12 White without, spotted with small purple spots, and white within with purple edges.
13 A white Duke, that is, parted with white & crimson flames, from the middle of each leafe to the edge.
14 The Princesse, that is, a siluer colour spotted with fine deepe blush spots.
15 The Queen, that is, a fine white sprinkled with bloud red spots, and greater strakes.

*Tulipa præcox purpurea.*

1 *Purpurea satura rubescens, vel violacea.*
2 *Purpurea pallida, Columbina dicta.*
3 *Persici coloris saturi.*
4 *Persici coloris pallidioris.*
5 *Paoniæ floris coloris.*
6 *Rosea.*
7 *Chermesina peramena.*
8 *Chermesina parum striata.*

The early purple Tulipa.

1 A deep reddish purple, or more violet.
2 A pale purple, called a Doue colour.
3 A deep Peach colour.
4 A paler Peach colour.
5 A Peony flower colour.
6 A Rose colour.
7 A Crimson very bright.
8 A Crimson stript with a little white.

9 *Princep-*

9 *Princeps, i.e. purpurea saturatior vel dilutior, oris albis magnis vel paruis, fundo luteo, vel albo orbe, quæ multum variatur, & colore, & oris, ita vt purpurea elegans oris magnis albis, dicta est, Princeps excellens, &*

10 *Princeps Columbina, purpurea dilutior.*

11 *Purpurea Chermesina, rubicundioris coloris, albidis vel albis oris.*

12 *Purpurea, vel obsoleta albidis oris Princeps Brancion.*

13 *Purpurea diluta, oris dilutioris purpurei coloris.*

14 *Purpurea in exterioribus, carnei verò ad medium intus, oris albis, fundo luteo.*

15 *Purpurea albo plumata extra, oris albis, purpurascens intus, fundo luteo, vel orbe albo.*

16 *Alia, minus elegans plumata, minoribúsq; oris albidis.*

9 A Prince or Bracklar, that is, a deepe or pale purple, with white edges, greater or smaller, and a yellow bottome, or circled with white, which varieth much, both in the purple & edges, so that a faire deep purple, with great white edges, is called, The best or chiefe Prince, and

10 A paler purple with white edges, called a Doue coloured Prince.

11 A Crimson Prince or Bracklar.

12 A Brancion Prince, or purple Brancion.

13 A purple with more pale purple edges.

14 Purple without, and blush halfe way within, with white edges, and a yellow bottome.

15 Purple feathered with white on the out side, with white edges, and pale purple within, the ground being a little yellow, or circled with white.

16 Another very neere vnto it, but not so fairely feathered, being more obscure, and the edges not so great or whitish.

## *Tulipa præcox rubra.*

## The early red Tulipa.

1 *Rubra vulgaris fundo luteo, & aliquando nigro.*

2 *Rubra saturo oris luteis paruis, dicta Roan.*

3 *Baro, i.e. rubra magis intensa, oris luteis paruis.*

4 *Dux maior & minor, i.e. rubra magis aut minus elegans saturo, oris luteis maximis vel minoribus, & fundo luteo magno. Alia aliis est magis amœna, in aliis etiam fundo nigro vel obscuro viridi.*

5 *Ducissa, i.e. Duci similis, at plus lutei quàm rubri, oris magnis luteis, & rubore magis aut minus intus in gyrum acto, fundo item luteo magno.*

6 *Testamentum Brancion, i.e. rubra sanguinea saturo, aut minus rubra, oris pallidis, magnis vel paruis: alia aliis magis aut minus elegans diuersimodo.*

1 An ordinary red, with a yellow, & sometimes a blacke bottome.

2 A deep red, with a small edge of yellow, called a Roane.

3 A Baron, that is, a faire red with a small yellow edge.

4 A Duke, a greater and a lesser, that is, a more or less faire deep red, with greater or lesser yellow edges, and a great yellow bottome. Some of this sort are much more or lesse faire then others, some also haue a blacke or darke greene bottome.

5 A Dutchesse, that is like vnto the Duke, but more yellow then red, with greater yellow edges, and the red more or lesse circling the middle of the flower on the inside, with a large yellow bottome.

6 A Testament Brancion, or a Brancion



Mo. Bot. Garden. 1893

7 *Flambans, ex rubore & flauedine radiata, vel striata fundo luteo.*
8 *Mali Aurantij coloris, ex rubore, & flauedine integrè, non separatim mixta, oris luteis paruis, vel absq; oris.*
9 *Minij, siue Cinabaris coloris, i.e. ex purpurea, rubedine, & flauedine radiata, vnguibus luteis, & aliquando oris.*
10 *Rex Tuliparum, i.e. ex sanguineo & aureo radiatim mixta, à flammea diuersa, fundo luteo, orbe rubro.*
11 *Tunica Morionis, i.e. ex rubore & aureo separatim diuisa.*

Duke, that is, a faire deepe red, or lesse red, with a pale yellow or butter coloured edge, some larger others smaller: and some more pleasing then others, in a very variable manner.
7 A Flambant, differing from the Dutchesse; for this hath no such great yellow edge, but streaks of yellow through the leafe vnto the very edge.
8 An Orenge colour, that is, a reddish yellow, or a red and yellow equally mixed, with small yellow edges, and sometimes without.
9 A Vermillion, that is, a purplish red, streamed with yellow, the bottome yellow, and sometimes the edges.
10 The Kings flower, that is, a crimson or bloud red, streamed with a gold yellow, differing from the Flambant, the bottome yellow, circled with red.
11 A Fooles coate, parted with red and yellow guardes.

## *Tulipa præcox lutea.*

1 *Lutea siue flaua.*
2 *Pallida lutea siue straminea.*
3 *Aurea, oris rubicundis.*
4 *Straminea, oris rubris.*
5 *Aurea, rubore persusa extra.*
6 *Aurea, vel magis pallida, rubore in gyrum acta simillima Ducissæ, nisi minus rubedinis habet.*
7 *Aurea, extremitatibus rubris, dici potest, Morianis Pileus præcox.*

## The early yellow Tulipa.

1 A faire gold yellow without mixture.
2 A strawe colour.
3 A faire yellow with reddish edges.
4 A strawe colour, with red edges.
5 A faire yellow, reddish on the out side onely.
6 A gold or paler yellow, circled on the inside a little with red, very like the Dutchesse, but that it hath lesse red therein.
7 A gold yellow with red toppes, and may be called, The early Fooles Cap.

*Tulipa de Caffa.* The Tulipa of Caffa.

There is another sort or kinde of early Tulipa, differing from the former, whose pale greene leaues being as broad and large as they, and sometimes crumpled or waued at the edges, in some haue the edges onely of the said leaues for a good breadth, of a whitish or whitish yellow colour, and in others, the leaues are listed or parted with whitish yellow and greene: the stalke riseth not vp so high as the former, and beareth a flower at the toppe like vnto the former, in some of a reddish yellow colour, with a russet coloured ground or bottome, and in others, of other seuerall colours: the seede and roote is so like vnto others of this kinde, that they cannot be distinguished.

There is (as I doe heare) of this kinde, both *Præcoces*, and *Serotinæ*, early flowring, and late flowring, whereof although wee haue not so exact knowledge, as of the rest, yet I thought good to speake so much, as I could hitherto vnderstand of them, and giue others leaue (if I doe not) hereafter to amplifie it.

*Tulipa Bolonicnsis, siue Bombycina flore rubro major.*
The greater red Bolonia Tulipa.

There are likewise other kindes of early Tulipas to bee spoken of, and first of the red Bolonia Tulipa; the roote whereof is plainly discerned, to be differing from all others: for that it is longer, and not hauing so plaine an eminence at the bottome thereof, as the former and later Tulipas, but more especially because the toppe is plentifully stored with a yellowish silke-like woollinesse: the outside likewise or skinne is of a brighter or paler red, not so easie to be pilled away, and runneth vnder ground both downeright and sidewise (especially in the Countrey ground and ayre, where it will encrease aboundantly, but not either in our London ayre, or forc't grounds) somewhat like vnto the yellow Bolonia Tulipa next following. It shooteth out of the ground with broad and long leaues, like the former; but neither so broad, nor of so white or mealy a greene colour as the former, but more darke then the late flowring Tulipa, so that this may bee easily discerned by his leafe from any other Tulipa aboue the ground, by one that is skilfull. It beareth likewise three or foure leaues vpon the stalke, like the former, and a flower also at the toppe of the same fashion, but that the leaues hereof are alwayes long, and somewhat narrow, hauing a large blacke bottome, made like vnto a cheuerne, the point whereof riseth vp vnto the middle of the leafe, higher then any other Tulipa; the flower is of a pale red colour, nothing so liuely as in the early or late red Tulipas, yet sweeter for the most part then any of them, and neerest vnto the yellow Bolonia Tulipa, which is much about the same sent.

*Tulipa pumilio rubra, siue Bergomensis rubra media & minor.*
The dwarfe red Bergomo Tulipa, a bigger and a lesser.

There are two other sorts hereof, and because they were found about Bergomo, do carry that name, the one bigger or lesser then another, yet neither so great as the former, hauing very little other difference to bee obserued in them, then that they are smaller in all parts of them.

*Tulipa Bolonicnsis flore luteo.* The yellow Bolonia Tulipa.

The roote of this Tulipa may likewise bee knowne from the former red (or any other Tulipa) in that it seldome commeth to bee so bigge, and is not so woolly at the toppe, and the skinne or outside is somewhat paler, harder, and sharper pointed: but the bottome is like the former red, and not so eminent as the early or late Tulipas. This beareth much longer and narrower leaues then any (except the Persian & dwarfe yellow Tulipas) and of a whitish greene colour: it beareth sometimes but one flower on a stalke, and sometimes two or three wholly yellow, but smaller, & more open then the other kinds, and (as I said) smelleth sweete, the head for seede is smaller then in others, and hath not that crowne at the head thereof, yet the seed is like, but smaller.

*Tulipa Narbonensis, siue Monspelliensis vel pumilio.*
The French or dwarfe yellow Tulipa.

This Tulipa is very like vnto the yellow Bolonia Tulipa, both in roote, leafe, and flower, as also in the colour thereof, being yellow: the onely difference is, that it is in all things lesser and lower, and is not so apt to beare, nor so plentifull to encrease by the roote.

*Tulipa Italica maior & minor.* The Italian Tulipa the greater and the lesser.

Both these kindes of Tulipas doe so neere resemble the last kinde, that I might almost say they were the same, but that some difference which I saw in them, maketh mee set them apart; and consisteth in these things, the stalkes of neither of both these rise so high, as of the first yellow Bolonia Tulipa: the leaues of both sorts are writhed in and out at the edges, or made like a waue of the sea, lying neerer the ground, and the flower being yellow within, is browniſh or reddish on the backe, in the middle of the three outer leaues the edges appearing yellow. Both these kindes doe differ one from the other in nothing, but in that one is bigger, and the other smaller then the other which I saw with Iohn Tradescante, my very good friend often remembred.

*Tulipa Lusitanica, siue pumilio versicolor.* The dwarfe stript Tulipa.

This dwarfe Tulipa is also of the same kindred with the three last described; for there is no other difference in this from them, then that the flower hath some red veins running in the leaues thereof.

There are two other sorts of dwarfe Tulipas with white flowers, whereof Lobel hath made mention in the Appendix to his *Aduersaria*; the one whereof is the same that Clusius setteth forth, vnder the title of *Pumilio altera*: but because I haue not seen either of them both, I speake no further of them.

*Tulipa pumilio alba.* The white dwarfe Tulipa.

But that white flower that Iohn Tradescante shewed me, and as hee saith, was deliuered him for a white Pumilio, had a stalke longer then they set out theirs to haue, and the flower also larger, but yet had narrower leaues then other sorts of white Tulipas haue.

*Tulipa Bicolor.* The small party coloured Tulipa.

Vnto these kindes, I may well adde this kinde of Tulipa also, which was sent out of Italy, whose leaues are small, long, and narrow, and of a darke greene colour, somewhat like vnto the leaues of an Hyacinth: the flower is small also, consisting of sixe leaues, as all other Tulipas doe, three whereof are wholly of a red colour, and the other three wholly of a yellow.

*Tulipa Persica.* The Persian Tulipa.

This rare Tulipa, wherewith we haue beene but lately acquainted, doth most fitly deserue to be described in this place, because it doth so neerely participate with the Bolonia and Italian Tulipas, in roote, leafe, and flower: the roote hereof is small, couered with a thicke hard blackish shell or skinne, with a yellowish woollinesse both at the toppe, and vnder the shell. It riseth out of the ground at the first, with one very long and small round leafe, which when it is three or foure inches high, doth open it selfe, and shew forth another small leafe (as long almost as the former) breaking out of the one side thereat, and after it a third, and sometimes a fourth, and a fift; but each shorter then other, which afterwards be of the breadth of the dwarfe yellow Tulipa, or somewhat broader, but much longer then any other, and abiding more hollow, and of the colour of the early Tulipas on the inside: the stalke riseth vp a foot and a halfe

1 *Tulipa Bombycina flore rubro.* The red Bolonia Tulipa. 2 *Tulipa Bononiensis flore luteo.* The yellow Bolonia Tulipa. 3 *Tulipa pumilio rubra sive lutea.* The red or yellow dwarfe Tulipa. 4 *Folium Tulipa de Cassa per totum striatum.* The leafe of the Tulipa of Cassa striped throughout the whole leafe. 5 *Folium Tulipa de Cassa per oras striatum.* The leafe of the Tulipa of Cassa striped at the edges onely. 6 *Tulipa Persica.* The Persian Tulipa. 7 *Tulipa Cretica.* The Tulipa of Candie. 8 *Tulipa Armeniaca.* The Tulipa of Armenia.

high sometimes, bearing one flower thereon, composed of sixe long and pointed leaues of the forme of other small Tulipas, and not shewing much bigger then the yellow Italian Tulipa, and is wholly white, both inside and outside of all the leaues, except the three outtermost, which haue on the backe of them, from the middle toward the edges, a shew of a browniſh blush, or pale red colour, yet deeper in the midst, and the edges remaining wholly white: the bottomes of all these leaues are of a darke or dun tawnie colour, and the chiues and tippes of a darkish purple or tawnie also. This doth beare seed but seldome in our Country, that euer I could vnderstand, but when it doth, it is small like vnto the Bolonia or dwarfe yellow Tulipas, being not so plentifull also in parting, or setting of by the roote as they, and neuer groweth nor abideth so great as it is brought vnto vs, and seldome likewise flowreth after the first yeare: for the rootes for the most part with euery one grow lesse and lesse, decaying euery yeare, and so perish for the most part by reason of the frosts and cold, and yet they haue been set deepe to defend them, although of their owne nature they will runne downe deep into the ground.

*Tulipa Byzantina duobus floribus Clusij.* The small Tulipa of Constantinople.

The small Tulipa of Constantinople, beareth for the most part but two leaues on the stalke, which are faire and broad, almost like vnto the Candy Tulipa, next hereunto to be described: the stalke it selfe riseth not aboue a foote high, bearing sometimes but one flower, but most commonly two thereon, one below another, and are no bigger then the flowers of the yellow Bolonia Tulipa, but differing in colour; for this is on the outside of a purpliſh colour, mixed with white and greene, and on the inside of a faire blush colour, the bottome and chiues being yellow, and the tippes or pendents blackish: the roote is very like the yellow Bolonia Tulipa.

*Tulipa Cretica.* The Tulipa of Candie.

This Tulipa is of later knowledge with vs then the Persian, but doth more hardly thriue, in regard of our cold climate; the description whereof, for so much as wee haue knowledge, by the sight of the roote and leafe, and relation from others of the flower, (for I haue not yet heard that it hath very often flowred in our Country) is as followeth. It beareth faire broad leaues, resembling the leaues of a Lilly, of a greeniſh colour, and not very whitiſh: the stalke beareth thereon one flower, larger and more open then many other, which is eyther wholly white, or of a deepe red colour, or else is variably mixed, white with a fine reddiſh purple, the bottomes being yellow, with purpliſh chiues tipt with blackiſh pendents: the roote is small, and somewhat like the dwarfe yellow Tulipa, but somewhat bigger.

*Tulipa Armeniaca.* The Tulipa of Armenia.

This small Tulipa is much differing from all the former (except the small or dwarfe white Tulipas remembred by Lobel and Clusius, as is before set downe) in that it beareth three or foure small, long, and somewhat narrow greene leaues, altogether at one ioynt or place; the stalke being not high, and naked or without leaues from them to the toppe, where it beareth one small flower likevnto an ordinary red Tulipa, but somewhat more yellow, tending to an Orenge colour with a blacke bottome: the roote is not much bigger then the ordinary yellow Bolonia Tulipa, before set downe.

And these are the sorts of this first *Classis* of early Tulipas.

*Tulipa media.* The meaner or middle flowring Tulipa.

For any other, or further description of this kinde of Tulipa, it shall not neede, hauing giuen it sufficiently in the former early Tulipa, the maine difference consisting first in the time of flowring, which is about a moneth after the early Tulipas, yet some more some lesse: for euen in the *Præcoces*, or early ones, some flower a little earlier, and later then others, and then in the colours of the flowers; for wee haue obserued many

colours, and mixtures, or varieties of colours in the *Medias*, which we could neuer see in the *Præcoces*, and so also some in the *Præcoces*, which are not in the *Medias*: yet there is farre greater varieties of mixture of colours in these *Medias*, then hath been obserued in all the *Præcoces*, (although Clusius saith otherwise) eyther by my selfe, or by any other that I haue conuersed with about this matter, and all this hath happened by the sowing of the seede, as I said before. I will therefore in this place not trouble you with any further circumstance, then to distinguish them, as I haue done in the former early Tulipas, into their foure primary colours, and vnder them, giue you their seuerall varieties and names, for so much as hath come to my knowledge, not doubting, but that many that haue trauelled in the sowing of the seed of Tulipas many yeares, may obserue each of them to haue some variety that others haue not: and therefore I thinke no one man can come to the knowledge of all particular distinctions.

| *Tulipa media alba.* | The white meane flowring Tulipa. |
|---|---|
| 1 *Niuea, fundo albo vel luteo.* | 1 A snow white, with a white or yellow bottome. |
| 2 *Argentea, quasi alba cineracea fundo lutescente, purpureis staminibus.* | 2 A siluer colour, that is, a very pale or whitish ashe colour, with a yellowish bottome and purple chiues. |
| 3 *Margaritina alba, carneo dilutissima.* | 3 A Pearle colour, that is, white, with a wash or shew of blush. |
| 4 *Alba, fundo caruleo vel nigro.* | 4 A white, with a blew or black bottome. |
| 5 *Albida.* | 5 A Creame colour. |
| 6 *Alba, oris rubris.* | 6 A white, with red edges. |
| 7 *Alba, purpureis oris.* | 7 A white, with purple edges. |
| 8 *Alba, oris coccineis.* | 8 A white, with crimson edges. |
| *Hæc tria genera in aliquibus constanter tenent oras, in alijs dispergunt.* | *These three sorts doe hold their edges constant in some, but well spread in others.* |
| 9 *Albida primum, deinde albidior, oris purpureis, & venis intrò respicientibus, dicta nobis Hackquenay.* | 9 A pale or whitish yellow, which after a few dayes groweth more white, with purplish red edges, and some streakes running inward from the edge, which we call an Hackney. |
| 10 *Alba, sanguineo colore variata, fundo vel albissimo, vel alio.* | 10 A white mixed with a bloud red very variably, and with a pure white, or other coloured bottome. |
| 11 *Alba, radiatim disposita flammis, & maculis coccineis.* | 11 A white, streamed with crimson flames, and spots through the whole flower. |
| 12 *Alba, purpurea rubedine plumata, diuersarum specierum, quæ cum superiore, vel albo, vel luteo, vel paruo cæruleo constant fundo, quæ constanter tenent punctatos colores, & non dispergunt, sed post trium aut quatuor dierum spatium pulchriores apparent.* | 12 A white, speckled with a reddish purple, more or lesse, of diuers sorts, with white, yellow, or blew bottomes, all which doe hold their markes constant, and doe not spread their colours, but shew fairer after they haue stood blown three or foure dayes. |
| 13 *Panni argentei coloris, i.e. alba, plumata, punctata, striata, vel diuersimodè variata, rubedine dilutiore, vel saturatiore purpurea, interius vel exterius, vel vtrinq;, diuersarum specierum.* | 13 A cloth of siluer of diuers sorts, that is, a white spotted, striped, or otherwise marked with red or purple, in some paler, in some deeper, either on the inside, or on the outside, or on both. |
| 14 *Tunica morionis alba varia, i.e. ex albo & purpureo striata diuersimodè, fundo albo vel alio.* | 14 A white Fooles coate of diuers sorts, that is, purple or pale crimson, and white, as it were empaled together, eyther with a white ground or other, whereof there is great variety. |
| 15 *Holias alba vel albida, absq; fundo, vel fundo purpureo caruleo, vel caruleo albo circundato, diuersè signata, vel variata intus ad medietatem foliorum, sursum in orbem vt plurimum, vel ad oras pertingens amplas & albas. Hæ species tantoperè multiplicantur, vt vix sint explicabiles.* | 15 A white Holias, that is, a faire white, or paler white, eyther without a bottome, or with a blewish purple bottome, or blew and white circling the bottome, and |

*Tanta est huius varietas, vel multitudine, vel striarum paucitate & distinctione, vel fundis variantibus, vt ad tedium esset perscribere.*

and from the middle vpwards, speckled and straked on the inside for the most part, with bloud red or purplish spots and lines vnto the very edges, which abide large and white. Of this kinde there are found very great varieties, not to be expressed.

Of this sort there is so much variety, some being larger or fairer marked then others, their bottomes also varying, that it is almost impossible to express them.

*Tulipa media purpurea.*

The meane flowring purple Tulipa.

1 *Purpurea saturo.*
1 A faire deep purple.

2 *Purpurea dilutior, diuersarum specierum, quarum Rosea vna, Carnea sit altera.*
2 A paler purple, of many sorts, whereof a Rose colour is one, a Blush another.

3 *Persici coloris, duarum aut trium specierum.*
3 A Peach colour of two or three sorts.

4 *Chermesina, obscura, aut pallida.*
4 A Crimson, deepe, or pale.

5 *Stamela, intensior aut remissior.*
5 A Stamell, darke or light.

6 *Xerampelina.*
6 A Murrey.

7 *Purpurea, striata.*
7 A purple, stript and sported.

8 *Persici saturi, vel diluti coloris, vndulata, vel radiata.*
8 A Peach colour, higher or paler, waued or stript.

9 *Columbina, oris & radijs albis.*
9 A Doue colour, edged and straked with white.

10 *Purpurea rubra, oris albis, simili Præcoci, dicta Princeps.*
10 A faire red purple, with white edges, like vnto the early Tulipa, called a Prince

11 *Chermesina, vel Heluola, lineis albis in medio, & versus oras, fundo cæruleo, vel albo, itemq; albo orbe.*
11 A faire Crimson, or Claret wine colour, with white lines both in the middle, and towards the edges, most haue a blew bottome, yet some are white, or circled with white.

12 *Purpurea remissior, aut intensior, oris albis, paruis aut magnis, vt in Principe præcoci, fundo vel cæruleo orbe albo, vel albo orbe cæruleo ample.*
12 A light or deepe purple, with white edges, greater or smaller, like the early Prince, the bottomes eyther blew circled with white, or white circled with a large blew.

13 *Holias Heluola, sanguineis guttis intus à medio sursum in orbem, fundo cæruleo.*
13 A purple Holias, the colour of a pale Claret wine, marked and sported with bloud red spots, round about the middle of each leafe vpward on the inside only, the bottome being blew.

14 *Tunica Morionis purpurea rubra saturo, albido striata, quam in alba saturatior, fundo ex cæruleo & albo.*
14 A Crimson Fooles Coate, a darke crimson, and pale white empaled together, differing from the white Fooles Coate, the bottome blew and white.

15 *Purpurea rubra saturo vel dilutò, albo vel albedine, punctata vel striata diuersimodè, dicta Cariophyllata.*
15 A deeper or paler reddish purple, spotted or striped with a paler or purer white, of diuers sorts, called the Gilloflower Tulipa.

*Tulipa*

1 *Tulipa rubra & lutea varia* The Fooles Coate red and yellow. 2 *Tulipa Holeas alba absq; fundo.* The white Holeas without a bottome. 3 *Tulipa argentea, vel punctata, &c.* The cloth of siluer, or other spotted Tulipa. 4 *Tulipa alba flammis coccineis.* The white Fooles Coate. 5 *Tulipa Holeas alba, &c. fundo purpureo, &c.* A white Holeas, &c. with a purple bottome, &c. 6 *Tulipa rubra & lutea flammea, &c.* A red and yellow flamed Tulipa, &c. 7 *Tulipa alba striata & punctata.* A white striped and spotted Tulipa. 8 *Tulipa altera variata, &c.* Another variable Tulipa.

*Tulipa mediæ rubra.*

1 *Rubra communis, fundo luteo, vel nigro.*
2 *Mali Aurantij coloris.*
3 *Cinabaris coloris.*
4 *Laterityj coloris.*
5 *Rubra, luteo aspersa.*
6 *Rubra, oris luteis.*
7 *Testamentum Brancion rubra satura, oris pallidis, diuersarum specierum, rubore variantium, & orarum amplitudine.*

8 *Cinabaris radiata, magis aut minus serotina.*
9 *Rubra purpurascens obsoleta, exterioribus foliis, perfusa luteo intus, oris pallidis luteis.*
10 *Rubra purpurascens elegans extra, & intus lutescens, oris pallidis luteis, fundo luteo vel viridi.*

11 *Rubra flambans coccinea, crebris maculis luteis absq; fundo.*
12 *Flambans elegantior rubra, i.e. radijs luteis intercurrentibus ruborem*

13 *Flambans remissior vtroq; colore.*
14 *Panni aurei coloris.*
15 *Tunica Morionis verior, seu Palto du Sot. optima, tenijs amplis amœnis & crebris, ex rubro & flauo separatim diuisis & excurrentibus, flos constans.*

16 *Tunica Morionis altera, tenijs minoribus & minus frequentibus, magis aut minus alia alijs inconstans.*

17 *Tunica Morionis pallida, i.e. tenijs vel strijs frequentioribus in vtroq; colore pallidis, flos est constans & elegans.*
18 *Pileus Morionis, radijs luteis, in medio foliorum latis, per ruborem excurrentibus, fundo luteo, apicibus luteis, & tribus exterioribus foliis luteis oris rubris, vel absq; oris.*

## The meane flowring red Tulipa.

1 A faire red which is ordinary, with a yellow or blacke bottome.
2 A deepe Orenge colour.
3 A Vermillion.
4 A pale red, or Bricke colour.
5 A Gingeline colour.
6 A red with small yellow edges.
7 A Testament Brancion of diuers sorts, differing both in the deepnesse of the red, and largenesse of the pale coloured edges.
8 A Vermillion flamed, flowring later or earlier.
9 A dead purplish red without, and of a yellowish red within, with pale yellow edges.
10 A bright Crimson red on the outside, more yellowish on the inside, with pale yellow edges, and a bottome yellow or greene.
11 A red Flambant, spotted thicke with yellow spots without any bottome.
12 A more excellent red Flambant, with flames of yellow running through the red.
13 A pale coloured Flambant.
14 A cloth of gold colour.
15 A true Fooles Coate, the best is a faire red & a faire yellow, parted into guards euery one apart, varied through euery leafe to the very edge, yet in most abiding constant.
16 Another Fooles Coate, not so fairely marked, nor so much, some of these are more or lesse constant in their marks, & some more variable then others.
17 A pale Fooles Coate, that is, with pale red, and pale yellow guardes or stripes very faire and constant.
18 A Fooles Cappe, that is, with lists or stripes of yellow running through the middle of euery leafe of the red, broader at the bottome then aboue, the bottome being yellow, the three outer leaues being yellow with red edges, or without.

1 *Tulipa tricolor.* A Tulipa of three colours. 2 *Tulipa Macedonica, siue de Cæsa varia.* The Tulipa of Cæsa purple, with pale white stripes. 3 *Tulipa Heluola charmesina versicolor.* A pure Claret wine colour variable. 4 *Tulipa Caryophyllata Wilmeri.* Mr. Wilmers Gilloflower Tulipa. 5 *Tulipa Chermesina flammis albis.* A Crimson with white flames. 6 *Tulipa Goliah.* A kind of Zwisler called Goliah. 7 *Tulipa le Zwisse.* A Tulipa called the Zwisler. 8 *Tulipa alba flammis coccineis.* Another white Flambant or Fooles Coate. 9 *Tulipa Cinnabarina albo flammata.* The Vermillion flamed. 10 *Tulipa plumata rubra & lutea.* The feathered Tulipa red and yellow.

19 *Le Suisse, tenijs radiata magnis ex rubore & pallore.*

20 *Altera dicta Goliah à floris magnitudine, tenijs radiata simillima le Suisse, nisi rubor & albedo sint elegantiores.*

21 *Holias rubra, i.e. sanguinea argenteis radijs, & guttis in orbem dispositis, præsertim interiùs, fundo viridi saturo.*

22 *Holias coccinea, rubra coccinea, alboradiata in orbem, circa medium foliorum interiùs, fundo albo.*

23 *Alia huic similis, fundo albo & cæruleo.*

19 A Swisse, paned with a faire red and pale white or strawe colour.

20 A Goliah, so called of the bignesse of the flower, most like to the Swisse in the marks and guardes, but that the red and white is more liuely.

21 A red Holias. A bloud red stript with siluer white veines and spots, with a darke green bottome.

22 A Crimson red Holias, that is, a faire purplish red, spotted with white circlewise about the middle of the inner leaues, and a white bottome.

23 Another like thereunto, with a blew and white bottome.

*Tulipa media lutea.*

1 *Lutea, sive Aurea vulgaris.*
2 *Straminea.*
3 *Sulphurea.*

4 *Mali Aurantij pallidi coloris.*
5 *Lutea dilutè purpurea striata, aurei panni pallidi instar,*
6 *Pallidè lutea fuscedine adumbrata.*
7 *Flaua, oris rubris magnis, aut paruis.*

8 *Straminea oris rubris magnis intensis, vel paruis remissis.*
9 *Obscura & fuliginosa lutea, instar Folij decidui, ideoq; Folium mortuum appellatur.*
10 *Flaua, rubore perfusa, etiamque striata per totum, dorso coccineo, oris pallidis.*

11 *Pallidè lutea, perfusa & magis aut minus rubore striata, fundo vel luteo, vel viridi.*
12 *Testamentum Clusij, i.e. lutea pallida fuligine obfusca, exteriùs & interiùs adoras vsq; pallidas, per totum verò floris medium, maculis interiùs asperfa instar omnium aliarum Holias, dorso obscuriore, fundo viridi.*

The meane flowring yellow Tulipa.

1 A faire gold yellow.
2 A Strawe colour.
3 A Brimstone colour pale yellowish greene.
4 A pale Orenge colour.
5 A pale cloth of gold colour.
6 A Custard colour a pale yellow shadowed ouer with a browne.
7 A gold yellow with red edges, greater or smaller.
8 A Strawe colour with red edges, deeper or paler, greater or smaller.
9 A sullen or smoakie yellow, like a dead leafe that is fallen, and therefore called, *Fueille mort.*
10 A yellow shadowed with red, and striped also through all the leaues, the backside of them being of a red crimson, and the edges pale.
11 A pale yellow, shadowed and striped with red, in some more in some lesse, the bottomes being either yellow or green.
12 A *Testamentum Clusij*, that is, a shadowed pale yellow, both within & without, spotted round about the middle on the inside, as all other Holias are, the backe of the leaues being more obscure or shadowed with pale yellow edges, and a greene bottome.

13 *Flam-*

13 *Flambans lutea, diuersimodè intus magis aut minus striata, vel in alijs extra maculata rubore, fundo vt plurimum nigro, vel in alijs luteo.*

13 A yellow Flambant of diuers sorts, that is, the whole flower more or lesse streamed or spotted on the inside, and in some on the outside with red, the bottome in most being blacke, yet in some yellow.

14 *Flambans pallidior & elegantior.*

14 A paler yellow Flambant more beautifull.

15 *Holias lutea intensior vel remissior diuersimodè, in orbem radiata interius, rubris maculis ad supremas vsq; oras, aliquoties crebrè, aliàs parcè, fundo viridi, vel tanetto obscuro.*

15 A yellow Holias, paler or deeper yellow very variable, spotted on the inside round about the middle, with red sometimes plentifully, or else sparingly, with a green or dark tawny bottome.

16 *Holias straminea rubore striata & punctata, instar alba Holias.*

16 A strawe coloured Holias, spotted and streamed with red, as is to bee seene in the white Holias.

17 *Tunica Morionis lutea, alijs dicta Flammea, in qua color flauus magis & conspicuus rubore, diuersimodè radiata.*

17 A yellow Fooles coate, of some called a flame colour, wherein the yellow is more then the red, diuersly streamed.

*Huc reddenda esset viridium Tuliparum classis, quæ diuersarum etiam constat specierum. Vna viridis intensior, cuius flos semper ferè semiclausus manet staminibus fimbriatis. Altera remissior, instar Psittaci pennarum viridium, luteo variata oris albis. Tertia adhuc dilutiori viriditate oris purpureis. Quarta, cujus folia æqualiter purpura diluta, & viriditate diuisa sunt. Quinta, foliis longissimis stellamodo expansis, ex rubore & viriditate coacta.*

Vnto these may be added the greene Tullipa, which is also of diuers sorts. One hauing a great flower of a deepe green colour, seldome opening it selfe, but abiding alwaies as it were halfe shut vp and closed, the chiues being as it were feathered. Another of a paler or yellowish green, paned with yellow, and is called, The Parret, &c. with white edges. A third of a more yellowish green, with red or purplish edges. A fourth, hath the leaues of the flower equally almost parted, with greene and a light purple colour, which abiding a long time in flower, groweth in time to be fairer marked: for at the first it doth not shew it selfe so plainely diuided. Some call this a greene Swisser. A fifth hath the longest leaues standing like a starre, consisting of greene and purple.

*Tulipa Serotina.* The late flowring Tulipa.

The late flowring Tulipa hath had his description expressed in the precedent discourse, so that I shall not neede to make a repetition of what hath already beene set downe. The greatest matter of knowledge in this kinde is this, That it hath no such plentifull variety of colours or mixtures in his flowers, as are in the two former sorts, but is confined within these limits here expressed, as farre as hath come to our knowledge.

*Tulipa Serotina.* The late flowring Tulipa.

*Rosea intensior, aut remissior.*
A Rose colour deeper or paler.

*Rubra vulgaris, aut saturatior, & quasi nigricans, fundo luteo vel nigro, vel nigro orbe, aureo incluso, dicta Oculus Solis.*
An ordinary red, or else a deeper red like blacke bloud, with a blacke or yellow bottome, or blacke circled with yellow, called the Suns eye.

*Lutea communis.*
An ordinary yellow.

*Lutea oris rubris.*
A yellow with red edges.

*Lutea guttis sanguineis, fundo nigro vel varie.*
A yellow with red spots and veines, the bottome blacke or discoloured.

There yet remaine many obſeruations, concerning theſe beautifull flowers, fit to be knowne, which could not, without too much prolixity, be comprehended within the body of the deſcription of them; but are reſerued to bee intreated of a part by themſelues.

All ſorts of Tulipas beare vſually but one ſtalke, and that without any branches: but ſometimes nature is ſo plentifull in bearing, that it hath two or three ſtalkes, and ſometimes two, or more branches out of one ſtalke (euery ſtalke or branch bearing one flower at the toppe) but this is but ſeldome ſeene; and when it doth happen once, it is hardly ſeene againe in the ſame roote, but is a great ſigne, that the roote that doth thus, being an old roote, will the ſame yeare part into diuers rootes, whereof euery one, being of a reaſonable greatneſſe, will beare both his ſtalke and flower the next yeare, agreeing with the mother plant in colour, as all the of-ſets of Tulipas doe for the moſt part: for although the young of-ſets of ſome doe vary from the maine roote, euen while it groweth with them, yet being ſeparated, it will bee of the ſame colour with the mother plant.

There groweth oftentimes in the *Medias*, and ſometimes alſo in the *Præcoces*, but more ſeldome, a ſmall bulbe or roote, hard aboue the ground, at the bottome of the ſtalke, and betweene it and the lower leafe, which when the ſtalke is dry, and it ripe, being put into the ground, will bring forth in time a flower like vnto the mother plant, from whence it was taken.

The flowers alſo of Tulipas conſiſt moſt commonly of ſixe leaues, but ſometimes they are ſeene to haue eight or tenne, or more leaues; but vſually, thoſe rootes beare but their ordinary number of ſixe leaues the next yeare: the head for ſeede then, is for the moſt part foure ſquare, which at all other times is but three ſquare, or when the flower wanteth a leafe or two, as ſometimes alſo it doth, it then is flat, hauing but two ſides.

The forme of the flower is alſo very variable; for the leaues of ſome Tulipas are all ſharpe pointed, or all blunt and round pointed, and many haue the three outer leaues ſharpe pointed, and the three inner round or pointed, and ſome contrariwiſe, the three outermoſt round pointed, and the three inner ſharpe pointed. Againe, ſome haue all the leaues of the flowers long and narrow, and ſome haue them broader and ſhorter. Some *Præcoces* alſo haue their flowers very large and great, equall vnto eyther the *Media*, or *Serotina*, which moſt commonly are the largeſt, and others haue them as ſmall as the Bolonia Tulipa.

The bottomes of the leaues of the flowers are alſo variably diuerſified, and ſo are both the chiues or threeds that ſtand vp about the head, and the tips or pendents that are hanging looſe on the toppes of them; and by the difference of the bottomes or chiues, many flowers are diſtinguiſhed, which elſe are very like in colour, and alike alſo marked.

For the ſmell alſo there is ſome diuerſity; for that the flowers of ſome are very ſweete, of others nothing at all, and ſome betweene both, of a ſmall ſent, but not offenſiue: and yet ſome I haue obſerued haue had a ſtrong ill ſent; but how to ſhew you to diſtinguiſh them, more then by your owne ſenſe, I cannot: for the ſeedes of ſweete ſmelling Tulipas doe not follow their mother plant, no more then they doe in the colour.

And laſtly, take this, which is not the leaſt obſeruation, worth the noting, that I haue obſerued in many: When they haue beene of one entire colour for diuers yeares, yet in ſome yeare they haue altered very much, as if it had not beene the ſame, *viz.* from a purple or ſtamell, it hath beene variably either parted, or mixed, or ſtriped with white, eyther in part, or through the whole flower, and ſo in a red or yellow flower, that it hath had eyther red or yellow edges, or yellow or red ſpots, lines, veines, or flames, running through the red or yellow colour, and ſometimes it hath happened, that three leaues haue been equally parted in the middle with red and yellow, the other three abiding of one colour, and in ſome the red had ſome yellow in it, and the yellow ſome red ſpots in it alſo; whereof I haue obſerued, that all ſuch flowers, not hauing their originall in that manner, (for ſome that haue ſuch or the like markes from the beginning, that is, from the firſt and ſecond yeares flowring, are conſtant, and doe not change) but as I ſaid, were of one colour at the firſt, doe ſhew the weakneſſe

weakneſſe and decay of the roote, and that this extraordinary beauty in the flower, is but as the brightneſſe of a light, vpon the very extinguiſhing thereof, and doth plainly declare, that it can doe his Maſter no more ſeruice, and therefore with this iollity doth bid him good night. I know there is a common opinion among many (and very confidently maintained) that a Tulipa with a white flower, hath changed to beare a red or yellow, and ſo of the red or yellow, and other colours, that they are likewiſe inconſtant, as though no flowers were certaine: but I could neuer either ſee or heare for certaine any ſuch alteration, nor any other variation, but what is formerly expreſſed. Let not therefore any iudicious be carried away with any ſuch idle conceit, but rather ſuſpect ſome deceit in their Gardeners or others, by taking vp one, and putting in another in the place, or elſe their owne miſtaking.

Now for the ſowing, planting, tranſplanting, choiſe, and ordering of Tulipas, which is not the leaſt of regard, concerning this ſubiect in hand, but (as I think) would be willingly entertained; What I haue by my beſt endeauours learned, by mine owne paines in almoſt forty yeares trauell, or from others informations, I am willing here to ſet downe; not doubting, but that ſome may adde what hath not come to my knowledge.

Firſt, in the ſowing of ſeedes of Tulipas, I haue not obſerued (whatſoeuer others haue written) nor could of certainty learne of others, that there doth ariſe from the ſeedes of *Præcoces* any *Medias* or *Serotine* Tulipas, (or but very ſeldome) nor am certainly aſſured of any: but that the ſeedes of all *Præcoces* (ſo they be not doubtfull, or of the laſt flowring ſorts) will bring *Præcoces*: And I am out of doubt, that I neuer ſaw, nor could learne, that euer the ſeede of the *Medias* or *Serotines* haue giuen *Præcoces*, but *Medias* or *Serotines*, according to their naturall kinde. But if there ſhould bee any degeneration, I rather incline to thinke, that it ſooner commeth to paſſe (*à meliore ad peius*, for *facilis eſt deſcenſus*, that is) that *Præcoces* may giue *Medias*, then that *Medias* or *Serotines* ſhould giue *Præcoces*.

For the choiſe of your ſeede to ſowe. Firſt, for the *Præcoces*, Cluſius ſaith, that the *Præcox Tulipa*, that beareth a white flower, is the beſt to giue the greateſt variety of colours. Some among vs haue reported, that they haue found great variety riſe from the ſeede of the red *Præcox*, which I can more hardly beleeue: but Cluſius his experience hath the greater probability, but eſpecially if it haue ſome mixture of red or purple in it. The purple I haue found to be the beſt, next thereunto is the purple with white edges, and ſo likewiſe the red with yellow edges, each of them will bring moſt of their owne colours. Then the choiſe of the beſt *Medias*, is to take thoſe colours that are light, rather white then yellow, and purple then red; yea white, not yellow, purple, not red: but theſe againe to be ſpotted is the beſt, and the more the better; but withall, or aboue all in theſe, reſpect the ground or bottome of the flower, (which in the *Præcox Tulipa* cannot, becauſe you ſhall ſeldome ſee any other ground in them but yellow) for if the flower be white, or whitiſh, ſpotted, or edged, and ſtraked, and the bottome blew or purple (ſuch as is found in the Holias, and in the Cloth of ſiluer, this is beyond all other the moſt excellent, and out of queſtion the choiſeſt of an hundred, to haue the greateſt and moſt pleaſant variety and rarity. And ſo in degree, the meaner in beauty you ſowe, the leſſer ſhall your pleaſure in rarities be. Beſtowe not your time in ſowing red or yellow Tulipa ſeede, or the diuers mixtures of them; for they will (as I haue found by experience) ſeldome be worth your paines. The *Serotina*, or late flowring Tulipa, becauſe it is ſeldome ſeene, with any eſpeciall beautifull variety, you may eaſily your ſelues gheſſe that it can bring forth (euen as I haue alſo learned) no raritie, and little or no diuerſity at all.

The time and manner to ſowe theſe ſeedes is next to be conſidered. You may not ſowe them in the ſpring of the yeare, if you hope to haue any good of them; but in the Autumne, or preſently after they be thorough ripe and dry: yet if you ſowe them not vntill the end of October, they will come forward neuer the worſe, but rather the better; for it is often ſeene, that ouer early ſowing cauſeth them to ſpring out of the ground ouer early, ſo that if a ſharpe ſpring chance to follow, it may goe neere to ſpoile all, or the moſt of your ſeede. Wee vſually ſowe the ſame yeares ſeede, yet if you chance to keepe of your owne, or haue from others ſuch ſeed, as is two years old, they will thriue and doe well enough, eſpecially if they were ripe and well gathered:

You must not sowe them too thicke, for so doing hath lost many a pecke of good seede, as I can tell; for if the seede lye one vpon another, that it hath not roome vpon the sprouting, to enter and take roote in the earth, it perisheth by and by. Some vse to tread downe the ground, where they meane to sowe their seede, and hauing sowne them thereon, doe couer them ouer the thickenesse of a mans thumbe with fine sifted earth, and they thinke they doe well, and haue good reason for it: for considering the nature of the young Tulipa rootes, is to runne downe deeper into the ground, euery yeare more then other, they thinke to hinder their quicke descent by the fastnesse of the ground, that so they may encrease the better. This way may please some, but I doe not vse it, nor can finde the reason sufficient; for they doe not consider, that the stiffenesse of the earth, doth cause the rootes of the young Tulipas to bee long before they grow great, in that a stiffe ground doth more hinder the well thriuing of the rootes, then a loose doth, and although the rootes doe runne downe deeper in a loose earth, yet they may easily by transplanting be holpen, and raised vp high enough. I haue also seene some Tulipas not once remoued from their sowing to their flowring; but if you will not lose them, you must take them vp while their leafe or stalke is fresh, and not withered: for if you doe not follow the stalke downe to the roote, be it neuer so deepe, you will leaue them behinde you. The ground also must be respected; for the finer, softer, and richer the mould is, wherein you sowe your seede, the greater shall be your encrease and varietie: Sift it therefore from all stones and rubbish, and let it be either fat naturall ground of it selfe, or being muckt, that it bee thoroughly rotten: but some I know, to mend their ground, doe make such a mixture of grounds, that they marre it in the making.

After the seede is thus sowne, the first yeares springing bringeth forth leaues, little bigger then the ordinary grasse leaues; the second yeare bigger, and so by degrees euery yeare bigger then other. The leaues of the *Præcoces* while they are young, may be discerned from the *Medias* by this note, which I haue obserued. The leaues of them doe wholly stand vp aboue the ground, shewing the small footstalkes, whereby euerie leafe doth stand, but the leaues of the *Medias* or *Serotines* doe neuer wholly appeare out of the ground, but the lower part which is broad, abideth vnder the vpper face of the earth. Those Tulipas now growing to bee three yeares old, (yet some at the second, if the ground and ayre be correspondent) are to bee taken vp out of the ground, wherein yee shall finde they haue runne deepe, and to be anew planted, after they haue been a little dryed and cleansed, eyther in the same, or another ground againe, placing them reasonable neare one vnto another, according to their greatnesse, which being planted and couered ouer with earth againe, of about an inch or two thickenesse, may be left vntaken vp againe for two yeare longer, if you will, or else remoued euery yeare after, as you please; and thus by transplanting them in their due season (which is still in the end of Iuly, or beginning of August, or thereabouts) you shall according to your seede and soyle, haue some come to bearing, in the fifth yeare after the flowring, (and some haue had them in the fourth, but that hath beene but few, and none of the best, or in a rich ground) some in the sixth and seuenth, and some peraduenture, not vntill the eighth or tenth yeare: but still remember, that as your rootes growe greater, that in re-planting you giue them the more roome to be distant one from another, or else the one will hinder, if not rot the other.

The seede of the *Præcoces*, doe not thriue and come forward so fast as the *Medias* or *Serotines*, nor doe giue any of-sets in their running downe as the *Medias* doe, which vsually leaue a small roote at the head of the other that is runne downe euery yeare; and besides, are more tender, and require more care and attendance then the *Medias*, and therefore they are the more respected.

This is a generall and certaine rule in all Tulipas, that all the while they beare but one leafe, they will not beare flower, whether they bee seedlings, or the of-sets of elder rootes, or the rootes themselues, that haue heretofore borne flowers; but when they shew a second leafe, breaking out of the first, it is a certaine signe, that it will then beare a flower, vnlesse some casualty hinder it, as frost or raine, to nip or spoile the bud, or other vntimely accident befall it.

To set or plant your best and bearing Tulipas somewhat deeper then other rootes, I hold it the best way; for if the ground bee either cold, or lye too open to the cold

Northerne

Northerne ayre, they will be the better defended therein, and not suffer the frosts or cold to pierce them so soone: for the deepe frosts and snowes doe pinch the *Præcoces* chiefly, if they bee too neare the vppermost crust of the earth; and therefore many, with good successe, couer ouer their ground before Winter, with either fresh or old rotten dung, and that will maruellously preserue them. The like course you may hold with seedlings, to cause them to come on the forwarder, so it bee after the first yeares sowing, and not till then.

To remoue Tulipas after they haue shot forth their fibres or small strings, which grow vnder the great round roores, (that is, from September vntill they bee in flower) is very dangerous; for by remouing them when they haue taken fast hold in the ground, you doe both hinder them in the bearing out their flower, and besides, put them in hazzard to perish, at least to bee put backe from bearing for a while after, as oftentimes I haue proued by experience: But when they are now risen to flower, and so for any time after, you may safely take them vp if you will, and remoue them without danger, if you haue any good regard vnto them, vnlesse it be a young bearing roote, which you shall in so doing much hinder, because it is yet tender, by reason it now beareth his first flower. But all Tulipa roots when their stalke and leaues are dry, may most safely then be taken vp out of the ground, and be so kept (so that they lye in a dry, and not in a moist place) for sixe moneths, without any great harme: yea I haue knowne them that haue had them nine moneths out of the ground, and haue done reasonable well, but this you must vnderstand withall, that they haue not been young but elder rootes, and they haue been orderly taken vp and preserued. The dryer you keep a Tulipa roote the better, so as you let it not lye in the sunne or winde, which will pierce it and spoile it.

Thus Gentlewomen for your delights, (for these pleasures are the delights of leasure, which hath bred your loue & liking to them, and although you are herein predominant, yet cannot they be barred from your beloued, who I doubt not, wil share with you in the delight as much as is fit) haue I taken this paines, to set downe, and bring to your knowledge such rules of art, as my small skill hath enabled mee withall concerning this subiect, which of all other, seemed fittest in this manner to be enlarged, both for the varietie of matter, and excellency of beautie herein; and also that these rules set forth together in one place, might saue many repetitions in other places, so that for the planting and ordering of all other bulbous rootes, and the sowing the seedes of them, you may haue recourse vnto these rules, *(tanquam ad normam & examen)* which may serue in generall for all other, little diuersitie of particulars needing exception.

## The Place.

The greater Tulipas haue first beene sent vs from Constantinople, and other parts of Turkie, where it is said they grow naturally wilde in the Fields, Woods, and Mountaines; as Thracia, Macedonia, Pontus about the Euxine Sea, Cappadocia, Bithynia, and about Tripolis and Aleppo in Syria also: the lesser haue come from other seuerall places, as their names doe descipher it out vnto vs; as Armenia, Persia, Candye, Portugall, Spaine, Italy, and France. They are all now made Denizens in our Gardens, where they yeeld vs more delight, and more encrease for their proportion, by reason of the culture, then they did vnto their owne naturals.

## The Time.

These doe flower some earlier, some later, for three whole moneths together at the least, therein adorning out a Garden most gloriously, in that being but one kinde of flower, it is so full of variety, as no other (except the Daffodils, which yet are not comparable, in that they yeeld not that alluring pleasant variety) doe the like besides. Some of the *Præcoces* haue beene in flower with vs, (for I speake not of their owne naturall places, where the Winters are milder, and the Spring earlier then ours) in the moneth of Ianuary, when the Winter before hath beene milde, but many in February,

and all the *Præcoces*, from the beginning to the end of March, if the yeare be kindly: at what time the *Medias* doe begin, and abide all Aprill, and part of May, when the *Serotines* flower and fade; but this, as I said, if the yeare be kindly, or else each kinde will be a moneth later. The seede is ripe in Iune and Iuly, according to their early or late flowring.

## The Names.

There haue beene diuers opinions among our moderne Writers, by what name this plant was knowne to the ancient Authors. Some would haue it be *Cosmosandalos*, of the Ancient. Dodonæus referreth it to πυπῶν of Theophrastus, in his seuenth Booke and thirteenth Chapter: but thereof he is so briefe, that besides the bare name, wee cannot finde him to make any further relation of forme, or quality. And Bauhinus, vpon Matthiolus Commentaries of Dioscorides, and in his Pinax also, followeth his opinion. Camerarius in his Hortus Medicus is of opinion, it may be referred to the Helychrysuni of Crateua. Gesner, as I thinke, first of all, and after him Lobel, Camerarius, Clusius and many others, referre it to the Satyrium of Dioscorides: and surely this opinion is the most probable for many reasons. First, for that this plant doth grow very frequent in many places of Greece, and the lesser Asia, which were no doubt sufficiently knowne both to Theophrastus, and Dioscorides, and was accounted among bulbous rootes, although by sundry names. And secondly, as Dioscorides setteth forth his Satyrium, so this most commonly beareth three leaues vpon a stalke (although sometimes with vs it hath foure or fiue) like vnto a Lilly, whereof some are often seen to be both red, in the first springing, and also vpon the decaying, especially in a dry time, and in a dry ground: the flower likewise of some is white, and like a Lilly; the roote is round, and as white within as the white of an egge, couered with a browne coate, hauing a sweetish, but not vnpleasant taste, as any man without danger many try. This description doth so liuely set forth this plant, that I thinke wee shall not neede to be any longer in doubt, where to finde Dioscorides his Satyrium Triphyllum, seeing wee haue such plenty growing with vs. And thirdly, there is no doubt, but that it hath the same qualities, as you shall hereafter heare further. And lastly, that plant likewise that beareth a red flower, may very well agree with his Erythronium; for the descriptions in Dioscorides are both alike, as are their qualities, the greatest doubt may be in the seede, which yet may agree vnto Lin or Flaxe as fitly, or rather more then many other plants doe, in many of his comparisons, which yet wee receiue for currant. For the seede of Tulipas are flat, hard, and shining as the seede of *Linum* or Flaxe, although of another colour, and bigger, as Dioscorides himselfe setteth it downe. But if there should be a mistaking in the writing of λίνω for χίονι, in the Greeke Text, as the slippe is both easie and likely, it were then out of all question the same: for the seede is very like vnto the seede of Lillies, as any man may easily discerne that know them, or will compare them. It is generally called by all the late Writers, *Tulipa*, which is deriued from the name *Tulpan*, whereby the Turkes of *Dalmatia* doe entitle their head Tyres, or Caps; and this flower being blowne, laide open, and inuerted, doth very well resemble them. We haue receiued the early kinde from Constantinople, by the name of *Cafa lale*, and the other by the name of *Cauala lale*. Lobel and others doe call it *Lilio-narcissus*, because it doth resemble a Lilly in the leafe, flower, and seede, and a Daffodill in the roote. We call it in English the Turkes Cap, but most vsually Tulipa, as most other Christian Countries that delight therein doe. Daleschampius calleth it Oulada.

## The Vertues.

Dioscorides writeth, that his first Satyrium is profitable for them that



haue

haue a convulsion in their necke, (which wee call a cricke in the necke) if it be drunke in harsh (which we call red) wine.

That the roots of Tulipas are nourishing, there is no doubt, the pleasant, or at least the no vnpleasant taste, may hereunto perswade; for diuers haue had them sent by their friends from beyond Sea, and mistaking them to bee Onions, haue vsed them as Onions in their pottage or broth, and neuer found any cause of mislike, or any sense of euill quality produced by them, but accounted them sweete Onions.

Further, I haue made tryall of them my selfe in this manner. I haue preserued the rootes of these Tulipas in Sugar, as I haue done the rootes of Eringus, Orchis, or any other such like, and haue found them to be almost as pleasant as the Eringus rootes, being firme and sound, fit to be presented to the curious; but for force of Venereous quality, I cannot say, either from my selfe, not hauing eaten many, or from any other, on whom I haue bestowed them: but surely, if there be any speciall propertie in the rootes of Orchis, or some other tending to that purpose, I thinke this may as well haue it as they. It should seeme, that Dioscorides doth attribute a great Venereous faculty to the seede, whereof I know not any hath made any especiall experiment with vs as yet.

## Chap. IX.

### *Narcissus.* The Daffodill.

THere hath beene great confusion among many of our moderne Writers of plants, in not distinguishing the manifold varieties of Daffodils; for euery one almost, without consideration of kinde or forme, or other speciall note, giueth names so diuersly one from another, that if any one shall receiue from seuerall places the Catalogues of their names (as I haue had many) as they set them down, and compare the one Catalogue with the other, he shall scarce haue three names in a dozen to agree together, one calling that by one name, which another calleth by another, that very few can tell what they meane. And this their confusion, in not distinguishing the name of *Narcissus* from *Pseudonarcissus*, is of all other in this kinde the greatest and grossest errour. To auoide therefore that gulfe, whereof I complaine that so manie haue bin endrenched; and to reduce the Daffodils into such a methodicall order, that euery one may know, to what *Classis* or forme any one doth appertaine, I will first diuide them into two principall or primary kindes: that is, into *Narcissos*, true Daffodils, and *Pseudonarcissos*, bastard Daffodils: which distinction I hold to be most necessarie to be set downe first of all, that euery one may be named without confusion vnder his owne primary kind, and then to let the other parts of the subdiuision follow, as is proper to them, and fittest to expresse them. Now to cause you to vnderstand the difference betweene a true Daffodill and a false, is this; it consisteth onely in the flower, (when as in all other parts they cannot bee distinguished) and chiefly in the middle cup or chalice; for that we doe in a manner onely account those to bee *Pseudonarcissos*, bastard Daffodils, whose middle cup is altogether as long, and sometime a little longer then the outter leaues that doe encompasse it, so that it seemeth rather like a trunke or long nose, then a cup or chalice, such as almost all the *Narcissi*, or true Daffodils haue; I say almost, because I know that some of them haue their middle cup so small, that we rather call it a crowne then a cup; and againe, some of them haue them so long, that they may seem to be of the number of the *Pseudonarcissi*, or bastard Daffodils: but yet may easily be knowne from them, in that, although the cup of some of the true Daffodils be great, yet it is wider open at the brim or edge, and not so long and narrow all alike as the bastard kindes are; and this is the chiefe and onely way to know how to seuer these kindes, which rule holdeth certaine in all, except that kinde which is called *Narcissus Iuncifolius reflexo flore*, whose cup is narrow, and as long as the leaues that turne vp againe.

Secondly,

Secondly, I will subdiuide each of these again apart by themselues, into foure sorts; and first the *Narcissos*, or true Daffodils into,

*Latifolios*, broad leafed Daffodils.

*Angustifolios*, narrow leafed Daffodils.

*Iuncifolios*, Rushe Daffodils, and

*Marinos*, Sea Daffodils.

These sorts againe doe comprehend vnder them some other diuisions, whereby they may the better be distinguished, and yet still bee referred to one of those foure former sorts: as

*Monanthos*, that is, Daffodils that beare but one flower, or two at the most vpon a stalke, and

*Polyanthos*, those that beare many flowers together vpon a stalke: as also

*Simplici flore*, those that beare single flowers, and

*Multiplici flore*, or *flore pleno*, that is, haue double flowers.

*Vernales*, those that flower in the Spring, and among them some that are earlier; and therefore called

*Præcoces*, early flowring Daffodils, and

*Autumnales*, those that flower in Autumne onely.

And lastly, with the *Pseudonarcissos*, or bastard Daffodils; I will keepe the same order, to distinguish them likewise into their foure seuerall sorts; and as with the true Daffodils, so with these false, describe vnder euery sort: first, those that beare single flowers, whether one or many vpon a stalke; and then those that beare double flowers, one or many also. As for the distinctions of *maior* and *minor*, greater and lesser, and of *maximus* and *minimus*, greatest and least, they doe not onely belong to these Daffodils; and therefore must be vsed as occasion permitteth, but vnto all other sort of plants. To begin therefore, I thinke fittest with that stately Daffodill, which for his excellency carrieth the name of None such.

### 1. *Narcissus latifolius omnium maximus, amplo calice flauo, siue Nompareille.*
### The great None such Daffodill, or Incomparable Daffodill.

This *Narcissus Nompareille* hath three or foure long and broad leaues, of a grayish greene colour, among which riseth vp a stalke two foote high at the least, at the toppe whereof, out of a thinne skinnie huske, as all Daffodils haue, commeth forth one large single flower, and no more vsually, consisting of sixe very pale yellow large leaues, almost round at the point, with a large cuppe in the middle, somewhat yellower then the leaues, the bottome whereof next vnto the stalke is narrow and round, rising wider to the mouth, which is very large and open, and vneuenly cut in or indented about the edges. The cup doth very well resemble the chalice, that in former dayes with vs, and beyond the Seas is still vsed to hold the Sacramentall Wine, that is with a narrower bottome, and a wide mouth. After the flower is past, sometimes there commeth (for it doth not often) a round greene head, and blacke round seede therein, like vnto other Daffodils, but greater. The roote is great, as other Daffodils that beare large flowers, and is couered ouer with a browniſh coate or skinne. The flower hath little or no sent at all.

*Flore geminato* This doth sometimes bring forth a flower with ten or twelue leaues, and a cup much larger, as if it would be two, euen as the flower seemeth.

### 2. *Narcissus omnium maximus flore & calice flauo.*
### The great yellow Incomparable Daffodill.

This other kinde differeth neither in forme, nor bigneſſe of leafe or flower from the former, but in the colour of the circling leaues of the flower, which are of the same yellow colour with the cup.

*Flore geminato.* This doth sometimes degenerate and grow luxurious also, bringing forth two flowers vpon a stalke, each distinct from other, and sometimes two flowers thrust together, as if they were but one, although it be but seldome; for it is not a peculiar kinde that is constant, yearly abiding in the same forme.

3. *Narcissus*

3. *Narcissus maximus griseus calice flauo.* The gray Peerlesse Daffodill.

This Peerlesse Daffodill well deserueth his place among these kindes, for that it doth much resemble them, and peraduenture is but a difference raised from the seede of the former, it is so like in leafe and flower, but that the leaues seeme to be somewhat greater, and the fixe outer leaues of the flower to be of a glistering whitish gray colour, and the cup yellow, as the former, but larger.

4. *Narcissus latifolius flauo flore amplo calice, siue Mattenesse.*
The lesser yellow Nompareille, or the Lady Mattenesses Daffodill.

The leaues of this Daffodill, are somewhat like vnto the leaues of the first kind, but not altogether so long or broad: the stalke likewise riseth not vp fully so high, and beareth one flower like the former, but lesser, and both the cuppe and the leaues are of one colour, that is, of a pale yellow, yet more yellow then in the former: the cup of this also is lesser, and a little differing; for it is neither fully so small in the bottome, nor so large at the edges, nor so crumpled at the brimmes, so that all these differences doe plainly shew it to be another kinde, quite from the former.

The Place.

The places of none of these are certainly knowne to vs where they grow naturally, but we haue them onely in our Gardens, and haue beene sent, and procured from diuers places.

The Time.

They flower sometimes in the end of March, but chiefly in Aprill.

The Names.

The first and second haue been sent vs by the name of *Narcisse Nompareille*, as it is called in French; and in Latine, *Narcissus omnium maximus amplo calice flauo*, and *Narcissus Incomparabilis*, that is, the Incomparable Daffodill, or the greatest Daffodill of all other, with a large yellow cuppe: but assuredly, although this Daffodill doth exceed many other, both in length and bigneſſe, yet the great Spanish bastard Daffodill, which shall be spoken of hereafter, is in my perswasion oftentimes a farre higher and larger flower; and therefore this name was giuen but relatiuely, we may call it in English, The great None such Daffodill, or the Incomparable Daffodill, or the great Peerlesse Daffodill, or the Nompareille Daffodill, which you will: for they all doe answer either the French or the Latine name; and because this name *Nompareille* is growne currant by custome, I know not well how to alter it. The third kinde may passe with the title giuen it, without controule. The last is very well knowne beyond the Seas, especially in the Low Countries, and those parts, by the Lady Matteneſſe Daffodill, because Clusius receiued it from her. We may call it in English, for the correspondency with the former, The lesser yellow Nompareille, or Peerlesse Daffodill, or the Lady Matteneſſe Daffodill, which you will.

*Narcissus Indicus flore rubro, dictus Iacobæus.*
The Indian Daffodill with a red flower.

This Indian Daffodill is so differing, both in forme, not hauing a cuppe, and in colour, being red, from the whole Family of the Daffodils (except the next that followeth, and the Autumne Daffodils) that some might iustly question the fitneſſe of his place here. But because as all the plants, whether bulbous or other, that come from the

the Indies, either East or West (although they differ very notably, from those that grow in these parts of the world) must in a generall survey and muster be ranked euery one, as neere as the surueiours wit will direct him, vnder some other growing with vs, that is of neereſt likeneſſe; Euen so vntill some other can direct his place more fitly, I shall require you to accept of him in this, with this descriptiõ that followeth, which I must tell you also, is more by relation then knowledge, or sight of the plant it selfe. This Daffodill hath diuers broad leaues, somewhat like vnto the common or ordinary white Daffodill, of a grayish greene colour; from the sides whereof, as also from the middle of them, rise vp sometimes two stalkes together, but most vsually one after another (for very often it flowreth twice in a Summer) and often also but one stalke alone, which is of a faint reddish colour, about a foote high or more, at the toppe whereof, out of a deepe red skinne or huske, commeth forth one flower bending downewards, consisting of sixe long leaues without any cup in the middle, of an excellent red colour, tending to a crimson; three of these leaues that turne vpwards, are somewhat larger then those three that hang downewards, hauing sixe threads or chiues in the middle, tipt with yellow pendents, and a three forked stile longer then the rest, and turning vp the end thereof againe: the roote is round and bigge, of a browniſh colour on the outside, and white within. This is set forth by Aldinus, Cardinall Farnesius his Physitian, that at Rome it rose vp with stalkes of flowers, before any leaues appeared.

### The Place, Time, and Names.

This naturally groweth in the West Indies, from whence it was brought into Spaine, where it bore both in Iune and Iuly, and by the Indians in their tongue named AZCAL XOCHITL, and hath beene sent from Spaine, vnto diuers louers of plants, into seuerall parts of Christendome, but haue not thriued long in these transalpine colder Countries, so far as I can heare.

## *Narcissus Trapezunticus flore luteo præcocissimus.*
## The early Daffodill of Trebizond.

Because this Daffodill is so like in flower vnto the former, although differing in colour, I thought it the fittest place to ioyne it the next thereunto. This early Daffodill hath three or foure short very greene leaues, so like vnto the leaues of the Autumne Daffodill, that many may easily bee deceiued in mistaking one for another, the difference consisting chiefly in this, that the leaues of this are not so broad or so long, nor rise vp in Autumne: in the midst of these leaues riseth vp a short green stalke, an handfull high, or not much higher vsually, (I speake of it as it hath often flowred with mee, whether the cause be the coldnesse of the time wherein it flowreth, or the nature of the plant, or of our climate, I am in some doubt; but I doe well remember, that the stalkes of some plants, that haue flowred later with me then the first, haue by the greater strength, and comfort of the Sunne, risen a good deale higher then the first) bearing at the top, out of a whitish thinne skinne striped with greene, one flower a little bending downewards, consisting of sixe leaues, laid open almost in the same manner with the former Indian Daffodill, whereof some doe a little turne vp their points againe, of a faire pale yellow colour, hauing sixe white chiues within it, tipt with yellow pendents, and a longer pointell: the roote is not very great, but blackish on the outside, so like vnto the Autumne Daffodill, but that it is yellow vnder the first or outermost coate, that one may easily mistake one for another.

### The Place.

It was sent vs from Constantinople among other rootes, but as wee may ghesse by the name, it should come thither from Trapezunte or Trebizond.

### The Time.

It flowreth sometimes in December, if the former part of the Winter haue

1 *Narcissus Nonpareille.* The incomparable Daffodill. 2 *Narcissus Mattenese.* The lesser yellow Nomparelle Daffodill. 3 *Narcissus Iacobæus flore rubro.* The red Indian Daffodill. 4 *Narcissus Trapezunticus,* The early Daffodill of Trabesond. 5 *Narcissus Montanus albus apophysous traditus.* The white winged Daffodill. 6 *Narcissus Montanus, sive Nonpareille totus albus.* The white Nompareille, or Peerlesse Daffodill. 7 *Narcissus albus oblongo calice.* The white Daffodill with a long cup.

haue been milde; but most vsually about the end of Ianuary, or else in Februarie the beginning or the end.

The Names.

Wee doe vsually call it from the Turkish name, *Narcissus Trapezunticus*, and some also call it *Narcissus vernus præcox*, as Clusius doth, in English, The early Daffodill of Trebizond.

*Narcissus Montanus albus apophysibus præditus.*
The white Mountaine Daffodill with eares, or
The white winged Daffodill.

This Mountaine Daffodill riseth vp with three or foure broad leaues, somewhat long, of a whitish greene colour, among which riseth vp a stalke a foote and a halfe high, whereon standeth one large flower, and sometimes two, consisting of sixe white leaues a peece, not very broad, and without any shew of yellownesse in them, three whereof haue vsually each of them on the backe part, at the bottome vpon the one side of them, and not on both, a little small white peece of a leafe like an eare, the other three hauing none at all: the cup is almost as large, or not much lesse then the small Nompareille, small at the bottome, and very large, open at the brimme, of a faire yellow colour, and sometimes the edges or brimmes of the cup will haue a deeper yellow colour about it, like as if it were discoloured with Saffron: the flower is verie sweete, the roote is great and white, couered with a pale coate or skinne, not verie blacke, and is not very apt to encrease, seldome giuing of-sets; neither haue I euer gathered seede thereof, because it passeth away without bearing any with me.

*Narcissus Montanus, sive Nompareille totus albus amplo calice.*
The white Nompareille Daffodill.

This white Nompareille Daffodill, is in roote and leafe very like vnto the former mountain or winged Daffodill, but that they are a little larger: the stalke from among the leaues riseth vp not much higher then it, bearing at the top one large flower, composed of sixe long white leaues, each whereof is as it were folded halfe way together, in the middle whereof standeth forth a large white cup, broader at the mouth or brims then at the bottome, very like vnto the lesser Nompareille Daffodill before remembred, which hath caused it to be so entituled: the sent whereof is no lesse sweete then the former.

The Place.

The naturall places of these Daffodils are not certainly knowne to vs; but by the names they carry, they should seeme to bee bred in the Mountaines.

The Time.

These flower not so early as many other kindes doe, but rather are to bee accounted among the late flowring Daffodils; for they shew not their flowers vntill the beginning of May, or the latter end of Aprill, with the soonest.

The Names.

The names set downe ouer the heads of either of them be such, whereby they are knowne to vs: yet some doe call the first *Narcissus auriculatus*, that is to say, The Daffodill with eares: and the other, *Narcissus Nompareille totus albus*, that is to say, The white Nompareille, or Peerlesse Daffodill.

1. *Narciſſus albus oblongo calice luteo præcox minor.*
The ſmall early white Daffodill with a long cup.

The leaues of this early Daffodill are broad, very greene, and not whitiſh as others, three or foure ſtanding together, about a foote long or better, among which riſeth vp a greene ſtalke, not full ſo high as the leaues, bearing one flower at the toppe thereof of a reaſonable bigneſſe, but not ſo great as the later kindes that follow are, conſiſting of ſix whitiſh leaues, but not perfect white, hauing a ſhew of a Creame colour appearing in them; in the middle is a long round yellow cup, about halfe an inch long or better. The ſmell of this flower is reaſonable ſweete, the roote is of a reaſonable bigneſſe, yet leſſer then the rootes of the later kindes.

2. *Narciſſus pallidus oblongo calice flauo præcox.*
The early Strawe coloured Daffodill with a long cup.

The leaues of this Daffodill are as greene as the former, but much narrower; and the leaues of the flower are more enclining to yellow, but yet very pale, as if it were a light ſtrawe colour, and ſeeme to bee a little more narrow and pointed then the former: the cup of this, is as long and yellow as the precedent. The ſmell whereof is very like the former, yet neither of them being ſo ſweete as thoſe that follow.

3. *Narciſſus albus oblongo calice luteo ſerotinus maior.*
The great late flowring white Daffodill with a long cup.

This later flowring Daffodill hath his leaues ſomewhat narrow & long, of a grayiſh or whitiſh greene colour, among which the ſtalke riſeth vp a foote and a halfe high, bearing one flower at the toppe, made of ſix white leaues, hauing the cup in the middle thereof as long as the former, and of a deepe yellow: the edges of this cuppe are ſometimes plaine, and ſometimes a little crumpled; they are often alſo circled at the brimmes with a Saffron colour, and often alſo without it, the ſmell whereof is very pleaſant, and not heady: the roote hereof is reaſonable bigge, and couered ouer rather with a pale then blackiſh skinne. This flower doth ſometimes alter his forme into eight leaues, which being narrow and long, ſeeme like a white ſtarre, compaſſing a yellow trunke.

4. *Narciſſus totus pallidus oblongo calice ſerotinus minor.*
The late pale coloured Daffodill with a long cup.

There is another of this kinde, whoſe flower is wholly of a pale white, or yellowiſh colour, differing neither in leafe nor roote from the former.

5. *Narciſſus pallidus oblongo calice flauo ſerotinus.*
The Strawe coloured late flowring Daffodill with a long yellow cup.

The chiefe difference of this Daffodill from the former, conſiſteth in the colour of the top of the flower, which is of a more yellow colour, and a little larger then the former, and the brimmes or edges of the cup of a deeper yellow, or Saffron colour. The ſmell of this is no leſſe ſweete then in the former.

6. *Narciſſus albus oblongo calice flauo ſerotinus, duobus floribus in caule.*
The late white Daffodill with a long cup, and two flowers on a ſtalke.

This Daffodill is ſurely a kinde of it ſelfe, although it be ſo like the former, abiding conſtant in his forme and manner of flowring, vſually bearing without miſſing two flowers vpon a ſtalke, very like vnto the former great white kinde, that one cannot know any greater matter of difference betweene them, then that it beareth two flowers on a ſtalke: the cuppes whereof are ſeldome touched with any ſhew of Saffron colour on them at the brimmes or edges, as ſome of the former haue.

### The Place.

All these Daffodils doe grow on the Pyrenæan mountaines, and haue been sought out, and brought into these parts, by those curious or couetous searchers of these delights, that haue made vs partakers of them.

### The Time.

The former kindes flower earlier by a fortnight then the later, the one in the later end of March, and the other not vntill the middle of Aprill.

### The Names.

Their names are giuen to euery one of them in their seuerall titles, as fitly as may best agree with their natures; and therefore I shall not neede to speake any further of them.

*Narcissus mediolutens vulgaris.*

The common white Daffodill called Primrose Peerlesse.

This Daffodill is so common in euery Countrey Garden almost through England, that I doubt I shall but spend my time in vaine, to describe that which is so well knowne, yet for their sakes that know it not, I will set downe the description of it in this manner. It hath long limber and broad leaues, of a grayish greene colour, among which riseth vp a stalke, bearing at the toppe out of a skinnie huske sometimes but one flower, but most commonly two flowers, and seldome three or more, but larger for the most part, then any that beare many flowers vpon a stalke, of a pale whitish Creame colour, tending somewhat neare vnto the colour of a pale Primrose (which hath caused our Countrey Gentlewomen, I thinke, to entitle it Primrose Peerlesse) with a small round flat Crowne, rather then a cup in the middle, of a pale yellow colour, with some pale chiues standing therein, being of a sweete, but stuffing sent: the roote is reasonable great, and encreasing more then a better plant.

*Narcissus mediocroceus serotinus.* The late flowring white Daffodill.

This Daffodill hath much smaller leaues, and shorter then the last, the stalke also riseth not so high by much, and beareth but one flower thereon, of a pure white colour, made of six small leaues, and somewhat narrow, standing seuerally one from another, and not so close together as the former, but appearing like a starre: the cup is small and round, of a pale yellow colour, but saffrony about the brims, hauing six small pale chiues in the middle, the smell whereof is much sweeter then in the former.

### The Place.

The first is thought to grow naturally in England, but I could neuer heare of his naturall place. I am sure it is plentifull enough in all Country Gardens, so that wee scarce giue it place in our more curious parkes. The second liueth onely with them that delight in varieties.

### The Time.

The first Daffodill flowreth in the middle time, being neither of the earliest, nor of the latest; but about the middle, or end of Aprill. The other flowreth with the latest in May.

### The Names.

I shall not neede to trouble you with further repetitions of names, they hauing been set downe in their titles, which are proper to them.

1 *Narcissus vulgaris medio luteus.* The common white Daffodil, or Primrose Peerlesse. 2 *Narcissus medio purpureus maximus.* The great white purple ringed Daffodill. 3 *Narcissus medio purpureus præcox.* The early purple ringed Daffodill. 4 *Narcissus medio purpureus stellatus.* The starry purple ringed Daffodill. 5 *Narcissus Persicus.* The Persian Daffodill. 6 *Narcissus Autumnalis minor.* The lesser Winter Daffodill. 7 *Narcissus Autumnalis maior.* The greater Winter Daffodill.

1. *Narcissus medio purpureus præcox.* The early purple ringed Daffodill.

This early Daffodill hath many long grayish greene leaues, somewhat narrower and stiffer then the former common white Daffodill, among which riseth vp a long naked hollow stalke (as all other Daffodils haue) bearing at the toppe one flower, and seldome two, made of sixe long white leaues, standing close together about the stalke; the cup is yellow, and so flat, that it might rather bee called a crowne: for it standeth very close to the middle, and very open at the brimmes, circled with a reddish or purple coloured ring, hauing certaine chiues in the middle of it also. The smell hereof is very sweete, exceeding many other.

2. *Narcissus medio purpureus serotinus.* The late purple ringed Daffodill.

The leaues of this Daffodill are alwayes broader then the former early one, and some are very neare twice as broad: the flower is very like the former, being large, and his leaues standing close one to the side of another; the ring likewise that compasseth the yellow coronet, is sometimes of a paler reddish purple, and sometimes as deepe a red as the former: so that it differeth not in any other materiall point, then that it flowreth not vntill the other is past and gone. The sent of this is like the former, the roote hereof is greater, as well as the leafe and flower.

3. *Narcissus medio purpureus maximus.*
The great white purple ringed Daffodill.

There is another kinde, whose flower (as well as leaues and rootes) is larger then any other of this kinde, which onely maketh it a distinct sort from the other: it flowreth also with the later sort of these purple ringed Daffodils.

4. *Narcissus medio purpureus stellaris.* The starry purple ringed Daffodill.

This Daffodill hath his leaues a little narrower and greener then the former sorts, the flower also of this hath his sixe white leaues not so broad, but narrower, and seeming longer then they, not closing together, but standing apart one from another, making it seeme like a white starre: it hath also a yellow coronet in the middle, circled about with purple, like the former. This doth smell nothing so sweete as the first, but yet hath a good sent.

The Place.

The first, third, and fourth of these Daffodils, haue alwayes beene sent vs from Constantinople among other bulbous rootes, so that wee know no further of their naturall places.

The second groweth in many places of Europe, both in Germany, France, and Italy, as Clusius hath noted.

The Time.

The first flowreth very early in March, euen with the first Daffodils. The second, third, and fourth, about a moneth after.

The Names.

The early and starre Daffodils, haue been sent vs by the Turkish name of *Deuebohini*, and *Serincade*. But their names, they haue receiued since, to bee endenizond with vs, are set downe in their seuerall titles.

*Narcissus Persicus.* The Persian Daffodill.

This Persian Daffodill differeth from all other kindes of Daffodils in his manner of growing,

growing, for it neuer hath leaues and flowers at one time together, wherein it is like vnto a Colchicum, yet in roote and leafe it is a Daffodill. The roote is a little blackish on the outside, somewhat like the roote of the Autumne Daffodill, from whence riseth vp a naked foote stalke, bearing one pale yellow flower, breaking through a thinne skinne, which first enclosed it, composed of six leaues, the three outermost being a little larger then the rest, in the middle of the flower there are six small chiues, and a longer pointell. The whole flower is of an vnpleasant sent: After the flower is past, come vp the leaues, sometimes before Winter, but most vsually after the deepe of Winter is past with vs, in the beginning of the yeare, which are broad, long, and of a pale greene colour, like the leaues of other Daffodils, but not greene as the Autumne Daffodill is, and besides they doe a little twine themselues, as some of the Pancratium, or bastard Sea Daffodils doe.

*Narcissus Autumnalis maior.* The greater Autumne or Winter Daffodill.

The greater Autumne Daffodill riseth vp with three or foure faire broad and short leaues at the first, but afterwards grow longer, of a very deepe or darke greene colour, in the middle of which riseth vp a short, stiffe, round footestalke, bearing one faire yellow flower on the head thereof (inclosed at the first in a thinne skinne, or huske) and consisteth of six leaues as the former, with certaine chiues in the middle, as all or most other Daffodils haue, which passeth away without shew of any seed, or head for seed, although vnder the head there is a little greene knot, which peraduenture would beare seede, if our sharpe Winters did not hinder it. The roote is great and round, couered ouer with a blackish skinne or coate.

*Narcissus Autumnalis minor.* The lesser Autumne or Winter Daffodill.

Clusius setteth downe, that the manner of the flowring of this lesser Daffodill, is more like vnto the Persian Daffodill, then vnto the former greater Autumne kind; but I doe finde that it doth in the same sort, as the greater kinde, rise vp with his leaues first; and the flowers a while after: the flower of this is lesser, and a little paler then the flower of the greatef kinde, but consisting in like sort of six leaues, narrow and sharpe pointed; the greene leaues also are almost of as deepe a greene colour, as the greater kinde, but smaller and narrower, and a little hollow in the middle. The roote is also alike, but lesser, and couered with a blackish skinne as the former. This hath sometimes borne blacke round seede in three square heads.

## The Place.

The Persian Daffodill hath beene sent sometimes, but very seldome, among other rootes from Constantinople, and it is probable by the name whereby it was sent, that it should naturally grow in Persia.

The other two haue likewise beene sent from Constantinople, and as it is thought, grow in Thracia, or thereabouts.

## The Time.

They all doe flower much about one time, that is, about the end of September, and in October.

## The Names.

The first hath been sent by the name of *Serincade Persiana*, and thereupon is called *Narcissus Persicus*, The Persian Daffodill.

The other two haue been thought by diuers to be Colchica, and so haue they called them, vpon no other ground, but that their flower is in forme and time somewhat like Colchicum, when as if they had marked them better, they might plainly discerne, that in all other things they did resemble Daffodils; but now the names of *Colchicum luteum maius, & minus*, is quite lost,

loſt, time hauing worne them out, and they are called by moſt Herbariſts now adayes, *Narciſſus Autumnalis maior & minor*, The greater and the leſſer Autumne Daffodill.

Thus farre haue I proceeded with thoſe Daffodils, that hauing broad leaues, beare but one ſingle flower, or two at the moſt vpon a ſtalke: And now to proceed with the reſt, that haue broad leaues, and beare ſingle flowers, but many vpon a ſtalke.

*Narciſſus Africanus aureus maior.* The great yellow Daffodill of Africa.

This braue and ſtately Daffodill hath many very long and broad leaues, of a better greene colour, then many others that are grayiſh, among which appeareth a ſtalke, not riſing to the height of the leaues, bearing at the toppe out of a skinnie hoſe many faire, goodly, and large flowers, to the number of ten or twelue, if the roote bee well growne, and ſtand in a warme place, euery one being larger then any of the French, Spaniſh, or Turkie Daffodils, that beare many ſingle flowers vpon a ſtalke, and commeth neere vnto the bigneſſe of the Engliſh Daffodill, called Primroſe Peerleſſe, before deſcribed, or that French kinde hereafter deſcribed, that beareth the largeſt flowers, many vpon a ſtalke (which ſome would make to bee a kinde of that Engliſh Daffodill, but bearing more flowers) and of a faire ſhining yellow colour, hauing large, round, and open cups or boules, yellower then the outer leaues; and is of ſo exceeding ſweete a ſent, that it doth rather offend the ſenſes by the aboundance thereof: the roote is great, and couered with a blackiſh browne coate or skinne.

*Narciſſus Africanus aureus minor.* The leſſer Barbary Daffodill.

This leſſer kinde is very neere the ſame with the former, but that it lacketh ſomewhat of his ſtatelineſſe of height, largeneſſe of flower and cup (being of a paler yellow) and beauty of colour, for it beareth neither of theſe equall vnto the former, but is in them all inferiour. And thus by this priuatiue, you may vnderſtand his poſitiue, and that ſhall be ſufficient at this time.

*Narciſſus Byzantinus totus luteus.* The yellow Turkie Daffodill.

Whereas the laſt deſcribed, came ſhort of the beauty of the former, ſo this lacketh of that beauty is in the laſt; for this, although it haue very long leaues, and a high ſtalke, yet the flowers are neither ſo many, as not being aboue foure or fiue, nor ſo large, being not much greater then the ordinary French Daffodill hereafter deſcribed, nor the colour ſo faire, but much paler, and the cup alſo ſmaller; and herein conſiſteth the chiefeſt differences betweene this, and both the other, but that the ſent of this is alſo weaker.

The Place.

The firſt and the ſecond grow in Barbary, about Argier, and Fez, as by the relation of them, that haue brought them into theſe parts, wee haue been enformed.

The laſt hath been often brought from Conſtantinople among other varieties of Daffodils, but from whence they receiued them, I could neuer learne.

The Time.

Theſe Daffodils do flower very early, euen with the firſt ſort of Daffodils, I meane after they haue been accuſtomed vnto our climate: for oftentimes vpon their firſt bringing ouer, they flower in Ianuary or February, eſpecially if they be preſerued from the froſts, and kept in any warme place; for they are very tender, and will ſoone periſh, being left abroad.

The Names.

The firſt is called by diuers in French, *Narciſſe d'Algiers*, and in many places

places of the Low Countries, *Narcissen van Heck*, or *Narcissus Heckius*; by diuers others *Narcissus Africanus aureus maior*, we may call it in English, The great African Daffodill, or the great Barbary Daffodill, or the great yellow Daffodill of Argiers, which you please.

The second hath no other variation of name, then a diminutiue of the former, as is set downe in the title.

The third is no doubt the same, that Clusius setteth downe in the twelfth Chapter of his second Booke of the History of more rare plants, and maketh the fourth sort, which came from Constantinople, and may also be the same, which he maketh his fifth, which (as he saith) he receiued from Doctour *Simor Touar* of Seuill in Spaine. Wee call it, from the place from whence we receiued it, *Narcissus Byzantinus*, with the addition of *totus luteus*, to put a difference from other sorts that come from thence also: in English, The yellow single Daffodill of Turkie.

*Narcissus Sulphureus maior*. The greater Lemon coloured Daffodill.

The greater of these Daffodils, beareth three or foure greene and very long leaues, a foote and a halfe long at the least, among which riseth vp a round, yet crested stalke, not so high as the leaues, bearing fiue or sixe single flowers thereon, euery one of them being greater then the ordinary French or Italian Daffodils, with many flowers vpon a stalke; of a faint, but yet pleasant yellow colour at the first, which after they haue been in flower a fortnight or thereabouts, change into a deeper, or more sullen yellow colour: the cup in the middle is likewise larger, then in those formerly named, and of a deeper yellow colour then the outer leaues, hauing onely three chiues within it. The smell is very pleasant.

*Narcissus Sulphureus minor*. The lesser Lemon coloured Daffodill.

This lesser Daffodill hath broader and shorter leaues then the former, of the colour of other Daffodils, and not greene like the former: the stalke of this riseth vp higher then the leaues, bearing foure or fiue flowers vpon shorter footestalkes, and no bigger then the French Daffodill, of a pale yellow, which most doe call a Brimstone colour, the cup or rather crowne in the middle, is small, and broad open, of a little deeper yellow, hauing many chiues within it, and is as it were sprinkled ouer with a kinde of mealinesse. The smell of this is not full so pleasant as the former.

## The Place.

Both these haue been gathered on the Pyrenæan Mountaines, and both likewise haue been sent out of Italy.

## The Time.

They both flower in the middle time of the Daffodils flowring, that is, in Aprill.

## The Names.

They haue their Latine names expressed in their titles, and so are their English also, if you please so to let them passe; or else according to the Latine, you may call them, The greater and the lesser Brimstone coloured Daffodils; some haue called them *Narcissus Italicus*, but the Italians themselues haue sent them by the name of *Narcisso Solfarigno*.

*Narcissus totus albus polyanthos*. The milke white Daffodill many vpon a stalke.

The leaues of this Daffodill are of a meane size, both for length and breadth, yet somewhat greener then in the ordinary sorts, that haue some whiteneſſe in them: the flowers

flowers are many vpon the ſtalke, as ſmall for the moſt part, as any of theſe kindes that beare many together, being wholly of a milke, or rather ſnow white colour, both the cuppe, which is ſmall, and the outer leaues that compaſſe it; after which come ſmall heads, wherein is contained round blacke ſeede, as all other Daffodils doe, although ſome greater, and others leſſer, according to the proportion of the plants: the roote is couered ouer with a blackiſh skinne or coate; the ſmell is very ſweete.

There are two other ſorts more of this kinde, the differences whereof are, that the one hath his leaues ſomewhat broader, and the flowers greater then the former: And the other ſmaller leaues and flowers alſo, whoſe cups being ſmall, are neuer ſeene fully open, but as it were halfe cloſed at the brimmes.

*Narciſſus latifolius totus albus, mediocri calice reflexus.*
The milke white Daffodill with the great cup.

There is yet another ſort of theſe milke white Daffodils, whoſe leaues are as broad as any of the former, and whoſe cup in the middle of the flower, is ſomewhat larger then in any of the leſſer ſorts, and leſſer then in the greater kinde: but the leaues of the flowers doe a little turne themſelues vpwards, which maketh a chiefe difference.

## The Place.

Theſe Daffodils grow in Spaine, from whence I receiued many that flouriſhed a while, but periſhed by ſome fierce cold Winters: they likewiſe grow in France, from whence many alſo haue been brought vnto vs. They haue likewiſe been ſent from Conſtantinople to vs, among other kindes of Daffodils.

## The Time.

They that come from Conſtantinople, for the moſt part doe flower earlier then the other, euen after they are accuſtomed to our ayre. Some of them flower notwithſtanding in the end of March, the reſt in Aprill.

## The Names.

They are vſually called *Narciſſus totus albus polyanthos*, adding thereunto the differences of *maior*, *medius*, and *minor*, that is, The milke white Daffodill, the greater, the middle, and the leſſer; for ſo ſome doe diſtinguiſh them. The laſt, for diſtinction, hath his name in his title ſufficient to expreſſe him.

1. *Narciſſus Narbonenſis, ſiue medio luteus præcox,*
The early French Daffodill.

The leaues of this Daffodill, ſpring vp out of the ground a moneth or two ſometimes before the other of this kinde, that follow; being alſo ſhorter, and narrower: the ſtalke likewiſe is not very high, bearing diuers flowers at the top, breaking through a thinne skinne, as is vſuall with all the Daffodils, euery one whereof is ſmall, conſiſting of ſix white leaues, and a ſmall yellow cup in the middle, which is of a prettie ſmall ſent, nothing ſo ſtrong as many others: the roote is great and round, and ſeldome parteth into of-ſets, euen as all the other that follow, bearing many ſingle flowers, doe.

2. *Narciſſus Narbonenſis vulgaris.* The ordinary French Daffodill.

This Daffodill hath long and broad greene leaues, a little hollowiſh in the middle, and edged on both ſides; the ſtalke is a foote and a halfe high, bearing at the toppe diuers flowers, ſomewhat larger then the former, conſiſting of ſix white leaues, ſomewhat round; the cup is yellow in the middle, ſmall and round, like vnto an Acorne cuppe, or a little fuller in the middle: this is the forme of that ſort which was firſt

brought

1 *Narcissus Africanus aureus major.* The great yellow Daffodill of Africa. 2 *Narcissus Africanus luteus minor.* The lesser yellow Daffodill of Africa. 3 *Narcissus Narbonensis medio luteus.* The French Daffodill. 4 *Narcissus Pisanus, vel totus albus.* The Italian Daffodill, or the all white Daffodill. 5 *Narcissus Mussart.* Mussart his Daffodill. 6 *Narcissus Anglicus polyanthos.* The great English Daffodill.

brought vnto vs: But since there is found out some, whose cup is shorter, others flatter, some of a paler, others of a deeper yellow colour, and some that haue their cuppe longer then the rest. The rootes of them all are couered with a blackish skin or coate.

3. *Narcissus Narbonensis maior amplo flore.*
The French Daffodill with great flowers.

The leaues of this Daffodill are somewhat like vnto the last, but not so broad, yet full as long, and spring sooner out of the ground, yet not so early as the first of these kindes: the stalke hereof is flatter, and riseth higher, bearing foure or fiue flowers, much larger then any of this kinde; for euery one of them doth equall the English Daffodill, before described, but whiter then it, and the yellow cup larger, and more open then in any of the rest. The roote of this is not so great, or round, as the former, but is more plentifull in of-sets, then any other of these French, or Italian kindes.

4. *Narcissus Pisanus.* The Italian Daffodill.

This Italian Daffodill hath his leaues as large, or larger then the second French Daffodill, and his stalke somewhat higher, bearing many white flowers, very like vnto the common French Daffodill, but somewhat larger also; and the yellow cup in the middle likewise is larger, and rounder, then is vsually seen in any of the French kinds, except the last with the greatest flowers.

5. *Narcissus mediocrocens polyanthos.*
The French Daffodill with Saffron coloured cups.

This French Daffodill hath diuers leaues of a grayish greene colour, not so broad or long as the last recited Daffodill, but comming neerer vnto the second French kinde, the flowers likewise are white, and many vpon a stalke, like thereunto, but the yellow cup is somewhat large, and circled with a Saffron like brimme or edge, which maketh the chiefest difference.

6. *Narcissus mediocrocens alter, dictus Mussart.* Mussart his Daffodill.

The affinity between this & the last, (for it is not the same to be expressed vnder one title) hath made me ioyne it next vnto it, yet because it hath a notable difference, it deserueth a place by himselfe. The leaues are large and long, and the flowers, being white, are larger also then in any other, except the greatest, but the cup hereof is small and short, rather seeming a coronet then a cup, of a deepe Saffron colour all about the brimmes or edges.

7. *Narcissus Anglicus polyanthos.* The great English Daffodill.

This Daffodill hath his leaues not much broader or longer, then the French kinde with great flowers, before described, the stalke with flowers riseth not fully so high as it, bearing many flowers thereon, not altogether so white, yet whiter then the former English Daffodill, called Primrose Peerlesse, but nothing so large, and with short, broad, and almost round leaues, standing close one vnto another: the yellow cup in the middle is bowle fashion, being somewhat deeper then in any of the former kinds, but not much greater: the smell hereof is very sweete and pleasant.

8. *Narcissus Narbonensis, sive medio luteus serotinus maior.*
The greater late flowring French Daffodill.

The roote as well as the leaues of this Daffodill, are greater, larger, broader, and longer then in any other of the former French, or Italian kindes; the stalke is as high as any of them, bearing at the toppe fiue or sixe white flowers, standing open spread like a starre, and not close together, euery one whereof is large, and round pointed, the

the cup is yellow, small and short, yet not lying flat to the flower, but a little standing out with some threads in the middle, as all the former Daffodils haue. This is not so sweete as the earlier kindes.

*9. Narcissus medioluteus alter serotinus calice breui.*
The lesser late flowring French Daffodill.

This Daffodill is of the same kinde with the last described, the onely difference is, that it is lesser, and the yellow cuppe in the middle of the flower, is somewhat shorter then the former, although the former be shorter then many others, otherwise it differeth not, no not in time; for it flowreth late as the former doth.

## The Place.

These Daffodils haue been brought vs from diuers places: The first and second grow naturally in many places of Spaine, that are open to the Sea: they grow likewise about Mompelier, and those parts in France. They haue been likewise sent among many other sorts of Daffodils from Constantinople, so that I may thinke, they grow in some places neere thereunto.

The fourth groweth plentifully in Italy, about Pisa in Tuscane, from whence we haue had plants to furnish our Gardens.

The seuenth is accounted beyond Sea to be naturall of our Country, but I know not any with vs that haue it, but they haue had it from them.

The rest haue been brought at diuers times, but wee know no further of their naturall places.

## The Time.

The first flowreth earlier then any of the rest by a moneth, euen in the beginning of March, or earlier, if the weather be milde. The other in Aprill, some a little before or after another. The late kinds flower not vntill May.

## The Names.

There can be no more said of the names of any of them, then hath beene set out in their titles; for they distinguish euery sort as fitly as we can: onely some doe call the first two sorts, by the name of *Donax Narbonensis*.

After all these Daffodils, that hauing broad leaues beare single flowers, either one or many vpon a stalke, I shall now goe on to set forth those broad leafed Daffodils, that carry double flowers, either one or many vpon a stalke together, in the same order that we haue vsed before.

1. *Narcissus albus multiplex.* The double white Daffodill.

The leaues of this Daffodill are not very broad, but rather of a meane size, being of the same largeneffe with the leaues of the purple ringed Daffodill, the stalke riseth vp to be a foote and a halfe high, bearing out of a thinne white skinne or hose, one flower and no more, consisting of many leaues, of a faire white colour, the flower is larger then any other double white Daffodill, hauing euery leafe, especially the outermost, as large almost as any leafe of the single Daffodill with the yellow cup, or purple ring. Sometimes it happeneth, that the flower is very little double, and almost single, but that is either in a bad ground, or for that it hath stood long in a place without remouing; for then it hath such a great encrease of rootes about it, that it draweth away into many parts, the nourishment that should be for a few: but if you doe transplant it, taking away the of-sets, and set his rootes single, it will then thriue, and beare his flower as goodly and double, as I haue before described it: and is very sweete.

2. *Narcissus mediopurpureus multiplex.* The double purple ringed Daffodill.

There is little difference in the leaues of this kinde, from the leaues of the single purple

ple ringed Daffodill; for it is probable it is of the ſame kinde, but by natures gift (and not by any humane art) made more plentifull, which abideth conſtant, and hath not that dalliance, which oftentimes nature ſheweth, to recreate the ſenſes of men for the preſent, and appeareth not againe in the ſame forme: the chiefeſt difference is, that the flower (being but ſometimes one on a ſtalke, and ſometimes two) conſiſteth of ſix white outer leaues, as large as the leaues of the ſingle kinde, hauing many ſmall yellow peeces, edged with purple circles round about them, inſtead of a cup; and in the middle of theſe peeces, ſtand other ſix white leaues, leſſer then the former, and a yellow cup edged with a purple circle likewiſe, parted into peeces, and they comprehend a few other white leaues, ſmaller then any of the other, hauing among them ſome broken peeces of the cup, with a few chiues alſo in the middle of the flower. The flower is very ſweete.

There is of this kinde another, whoſe flower hath not ſo plaine a diſtinction, of a triple rowe of leaues in it: but the whole flower is confuſedly ſet together, the outer leaues being not ſo large, and the inner leaues larger then the former; the broken yellow cuppe, which is tipt with purple, running diuerſly among the leaues; ſo that it ſheweth a fairer, and more double flower then the former, as it is indeed.

### 3. *Narciſſus mediolateus corona duplici.*
### The Turkie Daffodill with a double crowne.

This Daffodill hath three or foure leaues, as large and long almoſt, as the great double Daffodill of Conſtantinople next following hath: the ſtalke likewiſe is very neere as great, but as high altogether, bearing at the toppe foure or fiue flowers, the leaues whereof are as large, as of the firſt or ſecond kinde of French Daffodils, before deſcribed, but not altogether of ſo pure a white colour; and being ſix in number, ſtand like the former ſingle French Daffodils, but that the yellow cup in the middle of this is thicke and double, or as it were crumpled together, not ſtanding very high to be conſpicuous, but abiding lowe and ſhort, ſo that it is not preſently marked, vnleſſe one looke vpon it preciſely; yet is exceeding ſweete. The roote is like vnto the roote of the purple ringed Daffodill, or ſomewhat bigger.

### 4. *Narciſſus Chalcedonicus flore pleno albo polyanthos.*
### The double white Daffodill of Conſtantinople.

This beautifull and goodly Daffodill (wherewith all Floriſts greatly deſire to bee acquainted, as well for the beauty of his double flowers, as alſo for his ſuperabounding ſweete ſmell, one ſtalke with flowers being inſtead of a noſegay) hath many very broad, and very long leaues, ſomewhat greener then gray, among which riſeth vp a ſtrong round ſtalke, being ſometimes almoſt flat, and ribbed, bearing foure or fiue, or more white flowers at the toppe, euery one being very great, large, and double, the leaues being confuſedly ſet together, hauing little peeces of a yellow cup running among them, without any ſhew of that purple ring that is in the former, and fall away without bearing ſeed, euen as all, or moſt other double flowers doe: the ſmell is ſo exceeding ſweet and ſtrong, that it will ſoone offend the ſenſes of any, that ſhall ſmell much vnto it: the roote is great and thicke, couered with a blackiſh coate.

### 5. *Narciſſus Chalcedonicus fimbriatus multiplex polyanthos.*
### The great double purple ringed Daffodill of Conſtantinople.

This Daffodill differeth very little or nothing in leafe from the former, the onely difference is in the flowers, which although they bee double, and beare many vpon a ſtalke, like vnto them, yet this hath the peeces of the yellow cuppes tipt with purple, as if they were ſhred or ſcattered among the white leaues, whereas the other hath only the yellow, without any ſhew of purple tips vpon them: the ſmell of this is as ſtrong as of the other.

1 *Narcissus albus multiplex.* The double white Daffodill. 2 *Narcissus medioluteus corona duplici.* The Turkie Daffodill with a double crowne. 3 *Narcissus mediopurpureus multiplex.* The double purple ringed Daffodill. 4 *Narcissus chalcedonicus flore pleno albo polyanthos.* The double white Daffodill of Constantinople.

*6. Narciſſus Cyprius flore pleno luteo polyanthos.*
The double yellow Daffodill of Cyprus.

The leaues of this Daffodill are almoſt as broad and long as the former, the ſtalke is a foot high and more, bearing foure or fiue flowers on the top, euery one very double, and of a fine pale yellow colour, of a ſtrong heady ſent. The root of this is alſo like the former.

The Place.

The firſt of theſe Daffodils, was firſt brought into England by Mr. Iohn de Franqueuille the elder, who gathered it in his owne Countrey of Cambray, where it groweth wilde, from whoſe ſonne, Mr. Iohn de Franqueuille, now liuing, we all haue had it. The reſt haue come from Conſtantinople at ſeuerall times; and the laſt is thought to come from Cyprus. Wee haue it credibly affirmed alſo, that it groweth in Barbary about Fez and Argiers. Some of the double white kindes grow in Candy, and about Aleppo alſo.

The Time.

The Turkie kindes doe for the moſt part all flower early, in the end of March, or beginning of Aprill at the furtheſt, and the firſt double, about the middle or end of Aprill.

The Names.

All theſe Daffodils, except the firſt, haue had diuers Turkiſh names ſet vpon the packets, wherein they haue been ſent, but there is ſmall regard of certainty to be expected from them; for that the name *Serincade*, without any more addition, which is a ſingle Daffodill, hath beene impoſed vpon that parcell of rootes, that haue borne moſt of them double flowers of diuers ſorts; and the name *Serincade Catamer lale*, which ſignifieth a double flowred Daffodill, hath had many ſingle white flowers, with yellow cups, and ſome whoſe flowers haue been wholly white, cuppe and all, and ſome purple ringed, and double alſo among them. Their names, whereby they are knowne and called with vs, are, as fitly as may be, impoſed in their titles: And this I hope ſhall ſuffice, to haue ſpoken of theſe ſorts of Daffodils.

Hauing finiſhed the diſcourſe of the former ſort of broad leafed Daffodils, it is fit to proceede to the next, which are *Angustifolios Narciſſos*, thoſe Daffodils that haue narrow leaues, and firſt to ſet downe thoſe that beare ſingle flowers, whether one or many flowers vpon a ſtalke, and then thoſe that beare double flowers in the ſame manner.

*Narciſſus Virgineus.* The Virginia Daffodill.

This plant I thought fitteſt to place here in the beginning of this *Claſsis*, not finding where better to ſhroud it. It hath two or three long, and very narrow leaues, as greene as the leaues of the great *Leucoium bulboſum*, and ſhining withall, which grow ſometimes reddiſh, eſpecially at the edges: the ſtalke riſeth vp a ſpanne high, bearing one flower and no more on the head thereof, ſtanding vpright like a little Lilly or Tulipa, made of ſix leaues, wholly white, both within and without, except that at the bottome next to the ſtalke, and a little on the backſide of the three outer leaues, it hath a ſmall daſh or ſhew of a reddiſh purple colour: it hath in the middle a few chiues, ſtanding about a ſmall head pointed; which head groweth to bee ſmall and long, containing ſmall blackiſh flat ſeede: the roote is ſmall, long, and round, a little blackiſh on the outſide, and white on the inſide.

The

The Place.

This bulbous plant was brought vs from Virginia, where they grow aboundantly; but they hardly thriue and abide in our Gardens to beare flowers.

The Time.

It flowreth in May, and seldome before.

The Names.

The Indians in Virginia do call it *Attamusco*, some among vs do call it *Lilionarcissus Virginianus*, of the likenesse of the flower to a Lilly, and the leaues and roote to a Daffodill. Wee for breuity doe call it *Narcissus Virgineus*, that is, The Daffodill of Virginia, or else you may call it according to the former Latine name, The Lilly Daffodill of Virginia, which you will; for both names may serue well to expresse the plant.

*Narcissus angustifolius albidus præcox oblongo calice.*
The early white narrow leafed Daffodill with a long cup.

This Daffodill hath three or foure narrow, long, and very greene leaues, a foote long for the most part: the stalke riseth not vp so high as the leaues, whereon standeth one flower, not altogether so great as the late flowring Daffodill, with a long cuppe, described before among the broad leafed ones, which consisteth of six pale coloured leaues, not pure white, but hauing a wash of light yellow among the white: the cuppe in the middle is round and long, yet not so long as to bee accounted a bastard Daffodill, within which is a middle pointell, compassed with six chiues, hauing yellow mealy pendents.

The Place.

This Daffodill groweth with the other sorts of broad leafed ones, on the Pyrenæan Mountaines, from whence they haue beene brought vnto vs, to furnish our Gardens.

The Time.

It flowreth early, a moneth before the other sorts of the same fashion, that is, in the beginning of March, if the time be milde, which the other before spoken of doe not.

The Names.

It hath no other name that I know, then is expressed in the title.

2. *Narcissus mediocroceus tenuifolius.* The small Daffodill with a Saffron crown.

This small Daffodill hath foure or fiue narrow leaues about a spanne long, among which riseth vp a stalke some nine inches high, bearing at the toppe one small white flower, made of six leaues, with a small yellow cup in the middle, shadowed ouer at the brimmes with a Saffron colour: the roote is small, round, and little long withall, couered with a blackish skinne or coate.

3. *Narcissus minimus mediopurpureus.* The least purple ringed Daffodill.

This little Daffodill hath small narrow leaues, shorter by much then any of the purple ringed Daffodils, before described: the stalke and flower keepe an equall proportion to the rest of the plant, being in forme and colour of the flower, like vnto the

Starre Daffodill before recited, but vnlike in the greatnesse: this also is to bee obserued, that the purple colour that circleth the brimmes of the cuppe, is so small, that sometimes it is not well perceiued.

4. *Narcissus minimus Iuncifolij flore.* The least Daffodill of all.

This least Daffodill hath two or three whitish greene leaues, narrower then the two last recited Daffodils, and shorter by halfe, being not aboue two or three inches long, the stalke likewise is not aboue three or foure inches high, bearing one single flower at the toppe, somewhat bigger then the smalnesse of the plant should seeme to beare, very like vnto the least Rush Daffodill, and of the same bignesse, or rather somewhat bigger, being of a faint yellow colour, both leaues, and cup, or crowne, (if you please so to call it); for the middle part is spread very much, euen to the middle of the leaues almost, and lyeth flat open vpon the flower: the roote is small, euen the smallest of any Daffodill, and couered with a blackish skinne or coate.

The Place.

The first of these Daffodils haue beene brought vs from the Pyrenæan Mountaines, among a number of other rare plants, and the last by a French man, called Francis le Veau, the honestest roote-gatherer that euer came ouer to vs. The second was sent to M^r. Iohn de Franqueuille, before remembred, who imparted it to mee, as hee hath done many other good things; but his naturall place wee know not.

The Time.

They all flower about the latter end of Aprill.

The Names.

Being brought without names, wee haue giuen them their names according to their face and fashion, as they are set downe in their titles.

*Narcissus Autumnalis minor albus.* The little white Autumne Daffodill.

This little Autumne Daffodill riseth with his flowers first out of the ground, without any leaues at all. It springeth vp with one or two stalkes about a finger long, euery one bearing out of a small huske one small white flower, laid open abroad like vnto the Starre white Daffodill, before spoken of: in the middle of the flower is a small yellow cup of a meane size, and after the flower is past, there commeth in the same place a small head, containing small, round, blacke seede, like vnto the Autumne Hyacinth: the leaues come vp after the seede is ripe and gone, being small and narrow, not much bigger then the Autumne Hyacinth: the roote is small and blackish on the outside.

The Place.

This Daffodill groweth in Spaine, where Clusius saw it, and brought it into these parts.

The Time.

It flowreth in the beginning of Autumne, and his seede is ripe in the end of October in those hot Countries, but in ours it will scarce abide to shew a flower.

The Names.

The Spaniards, as Clusius reporteth, call it *Tonada*, and he vpon the sight thereof,

1 *Narcissus Virgineus.* The Virginian Daffodill. 2 *Narcissus minimus luncifolij flore.* The least Daffodill of all. 3 *Narcissus Autumnalis minor albus.* The little white Autumne Daffodill. 4 *Narcissus albus Autumnalis medio obsoletus.* The white Autumne Daffodill with a sullen crown. 5 *Narcissus luncifolius maximus amplo calice.* The great Junquilla with the largest flower or cup. 6 *Narcissus totus albus flore plano Virginianus.* The double white Daffodill of Virginia.



5. Narcissus odorus. Spa. hist.

thereof, *Narcissus Autumnalis minor albus*, and wee in English thereafter, The little white Autumne Daffodill.

*Narcissus albus Autumnalis medio obsoletus.*
The white Autumne Daffodill with a sullen crowne.

This Autumne Daffodill hath two or three leaues at the most, and very narrow, so that some doe reckon it among the Rush Daffodils, being somewhat broad at the bottome, and more pointed at the toppe, betweene these leaues commeth vp the stalke, bearing vsually two flowers and no more at the toppe, made of sixe white leaues a peece, pointed and not round: the cup is small and round, like vnto the cup or crowne of the least Rush Daffodill, of a yellow colour at the bottome, but toward the edge of a dunne or sullen colour.

*Narcissus angustifolius luteus semper florens Caccini.*
The yellow Italian Daffodill of Caccini.

This Daffodill beareth a number of small, long, narrow, and very greene leaues, broader then the leaues of any Rush Daffodill, among which rise vp diuers stalkes, bearing at the head two or three flowers a peece, each of them being small and yellow, the cup or crowne is small also, of a deeper yellow then the flower. The Nobleman of Florence, who first sent this plant to Christian Porret at Leyden, after the death of Carolus Clusius, writeth that euery stalke doth beare with him more store of flowers, then are formerly set downe, and that it neuer ceaseth to beare flowers, but that after one or moe stalkes haue been in flower together, and are past, there succeed other in their places.

The Place.

The first is naturall of Spain, the naturall place of the other is not known to vs.

The Time.

The times of the flowring, are set downe both in the title and in the descriptions; the one to be in Autumne, the other to be all the Summer long.

The Names.

The Latine names are imposed on them, as are fittest for them, and the last by that honourable man that sent it, which is most fit to continue, and not to bee changed. But wee, to let it bee knowne by an English name to English people, haue entituled it, The yellow Italian Daffodill of Caccini: if any man can giue it a more proper name, I shall bee therewith right well content.

*Narcissus angustifolius, sive Iuncifolius maximus amplo calice.*
The great Iunquilia with the large flower or cup.

Although this Daffodill importeth by his name, not to be of this family, but of the next, considering it is so like vnto them, but bigger; yet I haue thought good to place it in the end of these narrow leafed Daffodils, as being indifferent, whether it should bee referred to this or to that. For this carrieth diuers long greene leaues, like vnto the other Rush Daffodils, but thicker and broader, so that it may without any great errour, bee reckoned among these narrow leafed Daffodils, bearing at the toppe two or three very faire large flowers, with a large and more open cuppe, then in any other of the Rush Daffodils, both of them of a faire yellow colour, yet the cuppe a little deeper then the flower, and a little crumpled about the edges, and hath a pretty sharpe sent: the roote is greater and longer then the other Rush Daffodill, and couered likewise with a blackish coate.

The

The Place.

We haue this in Gardens onely, and haue not heard of his naturall place.

The Time.

It flowreth in Aprill.

The Names.

I leaue it indifferent, as I said, whether you will call it *Narcissus angustifolius*, or *Iuncifolius magno calice*, or *maximus*, because it is the greatest of all the rest of that kinde.

*Narcissus totus albus flore pleno Virginianus.*

The double white Daffodill of Virginia.

The roote of this Daffodill, is very like vnto the former single Virginia Daffodill, set forth in the first place of this ranke of narrow leafed Daffodils, but that it is a little bigger and rounder, being a little long withall, and blackish also on the outside, as that is: from whence riseth vp two leaues, somewhat broader then the former: but of a like greennesse: the stalke riseth vp betweene these two leaues, about a span high, or not much higher, bearing one faire double snow white flower, very like in the fashion vnto the pale yellow double Daffodill, or bastard Daffodill of Robinus, hereafter described: For it is in the like manner laid open flat, and composed of six rowes of leaues, euery rowe lying in order iust opposite, or one before another, whereof those six leaues that make the first or outermost course, are the greatest, and all the rest lying, as I said, one vpon or before another, are euery rowe smaller then others from the middle of this flower, thrusteth forth a small long pointed forke or horne, white as the flower is.

The Place.

The place is named to be Virginia, but in what part it is not known to vs.

The Time.

It flowreth in the end of Aprill.

The Names.

It may be that this doth grow among the former single kinde, and called by the same name Attamusco, for that the plant is not much differing, yet hereof I am not certaine: But we, from the forme and countenance of the plant, doe call it *Narcissus Virginianus*, The Virginian Daffodill, and because it beareth a double flower, it hath the title of double added vnto it.

The third order of Daffodils, I said in the beginning, was of *Iuncifolios*, Rush Daffodils, which are now next to be entreated of, I shall herein keepe the same order I vsed in the former; but because I finde none of this order, that beare but one flower vpon a stalke, I must begin with those that beare many.

1. *Narcissus Iuncifolius albus.* The white Iunquilia.

This white Rush Daffodill hath small long leaues, a little broader, and of a whiter greene colour then the ordinary yellow Rush Daffodils: the stalke riseth vp halfe a foote high or more, bearing two or three small white flowers vpon a stalke, yet somewhat bigger then the common yellow Rush Daffodill, hauing a small round cuppe in the middle, white also as the leaues are. The seede is small, blacke,

blacke, and round, as other ſeedes of Daffodils are: the roote is ſmall and round, couered with a blackiſh coate.

*Narciſſus Iuncifolius albus magno calice.* The white Iunquilia with a great cup.

There is of this kinde another ſort, that hath the cup in the middle of the flower, a little larger then the other, but in all other things alike.

2. *Narciſſus Iuncifolius flore albo reflexo.*
The white turning Iunquilia, or Ruſh Daffodill.

This turning white Daffodill hath foure or fiue long greene leaues, yet ſhorter and broader then the ordinary yellow Iunquilia, and fully as greene alſo, from among which riſeth vp a ſlender greene ſtalke, a foote high, bearing out of a thinne skinnie huske, three or foure, or more ſnow white flowers, ſtanding vpon long greene foot-ſtalkes, euery flower hanging downe his head, and turning vp his ſix narrow and long leaues, euen to the very foot-ſtalke againe: from the middle of the flower hangeth downe a long round cuppe, as white as the leaues, within which are contained three ſmall white chiues, tipt with yellow, and a ſmall long pointell, thruſting out beyond the brimmes of the cup: after the flowers are paſt, there come vp in their places ſmall three ſquare heads, wherein is contained very ſmall, round, and blacke ſhining ſeede: the roote is ſmall, round, and a little long withall, couered with a blackiſh browne coate or skin. The flower is quite without any good ſent, or indeed rather none at all.

3. *Narciſſus Iuncifolius flore luteo reflexo.*
The yellow turning Iunquilia, or Ruſh Daffodill.

The leaues of this Ruſh Daffodill are greater and longer then the former, and of a paler greene colour: the ſtalke riſeth ſomewhat higher, bearing two or three flowers thereon wholly of a gold yellow colour, both the cuppe and the leaues that turne vp againe.

4. *Narciſſus Iuncifolius calice albo reflexis foliis luteis.*
The yellow turning Iunquilia with a white cup.

This Daffodill hath his long ruſh-like leaues ſtanding vpright as the former, betweene which riſeth vp a greene ſtalke, about a foote high or more, bearing two or three flowers thereon, whoſe turning leaues are of a faire pale yellow, and the cuppe pale white, and not ſo pure a white as the former.

5. *Narciſſus Iuncifolius calice luteo reflexis foliis albidis.*
The white turning Iunquilia with a yellow cup.

As the laſt had the leaues of the flower that turne vp againe yellow, and the cuppe whitiſh, ſo this hath contrariwiſe the turning leaues of a whitiſh yellow, and the long cup yellower, elſe in his long green leaues, or any other thing, there is ſmall difference.

6. *Narciſſus Iuncifolius luteus magno calice.*
The Iunquilia, or Ruſh Daffodill with a great cup.

This Ruſh Daffodill hath bigger leaues, and longer then the ordinary yellow Ruſh Daffodill, being a little flat on the one ſide, and round on the other, but of the ſame greenneſſe with all the reſt: the ſtalke riſeth vp two foote high, bearing two, and ſometimes three flowers thereon, being of a faire yellow colour, with a large open cup in the middle, of a little deeper yellow colour, like vnto the great Iunquilia with the large flower, before ſet downe, whereof this is a kinde, no doubt; but that is larger and greater then this, both in leafe, flower, cup, &c. and this onely ſomewhat leſſe in all parts then that.

7. *Narciſſus*

1 *Narcissus lunæfolius albus.* The white Iunquilia. 2 *Narcissus lunæfolius flore albo reflexo.* The white turning Iunquilia. 3 *Narcissus lunæfolius calice luteo reflexis foliis albis.* The yellow turning Iunquilia. 4 *Narcissus lunæfolius luteus magno calice.* The yellow Iunquilia with a great cuppe. 5 *Narcissus lunæfolius luteus maior vulgaris.* The ordinary yellow Iunquilia. 6 *Narcissus lunæfolius Autumnalis flore viridi.* The greene Autumne Iunquilia. 7 *Narcissus angustifolius aureus multiplex.* The golden double narrow leated Daffodill. 8 *Narcissus lunæfolius flore pleno.* The double Iunquilia.

6 Narcissus viridiflorus, Schmo. Spec. Inst.

7. *Narciſſus Iuncifolius luteus vulgaris maior.*
The ordinary Iunquilia, or Ruſh Daffodill.

This ordinary Ruſh Daffodill hath foure or fiue long greene round leaues, like vnto Ruſhes, whereof it tooke the name: among theſe leaues riſeth vp the ſtalke, round and greene, a foote and a halfe high very often, bearing at the toppe three or foure flowers all yellow, but much ſmaller then the laſt, and ſo is the cup alſo: the ſeede is ſmall and blacke, incloſed in ſmall cornered heads; the roote is blackiſh on the outſide. The ſmell of the flower is very ſweete in all theſe ſorts of Ruſh Daffodils.

8. *Narciſſus Iuncifolius luteus medius.* The ſmaller Iunquilia, or Ruſh Daffodill.

The leaues of this Daffodill are like vnto the former, but ſmaller and rounder, the ſtalke riſeth not vp ſo high, nor are the flowers ſo great, but the leaues of the flower are a little rounder, and not ſo pointed as in the former, in all things elſe alike, ſauing leſſer.

9. *Narciſſus Iuncifolius luteus minor.* The leaſt Iunquilia, or Ruſh Daffodill.

This leaſt Daffodill hath fiue or ſix ſmall greene leaues, a little broader, and not ſo long as the laſt, among which riſeth vp a ſtalke almoſt a foote high, bearing one or two ſmall flowers at the toppe, of a paler yellow colour then the former, with a yellow open cuppe, or crowne rather in the middle, bigger then in either of the laſt two: the roote is very ſmall and blacke, like vnto the laſt in roundneſſe and colour.

10. *Narciſſus Iuncifolius luteus albicantibus lineis diſtinctus.*
The yellow Iunquilia, or Ruſh Daffodill with white lines.

This Ruſh Daffodill hath round, greene, and long leaues, like vnto the ordinary Ruſh Daffodill, with a ſtalke bearing two or three yellow flowers, hauing leaues ſomewhat round at the point or end, with a line or ſtrake of white in the middle of euerie one of them, the cup is ſhort, and crowne faſhion, a little crumpled about the brims: the ſeede, roote, or any thing elſe differeth not.

11. *Narciſſus Iuncifolius Autumnalis flore viridi.*
The Autumne Ruſh Daffodill with a greene flower.

This ſtrange Ruſh Daffodill (I call it ſtrange, not onely becauſe it differeth from all others of this kinde, but alſo becauſe there are but few in theſe parts that haue had it, and fewer that doe ſtill enioy it, in that it is periſhed withall that had it) hath but one onely leafe, very long, round, and greene, in all that euer I ſaw growing, which beareth no flower while that greene leafe is freſh, and to bee ſeene: but afterwards the ſtalke riſeth vp, being like vnto the former greene leafe, round, naked, and greene vp to the toppe, where two or three flowers breake forth out of a ſmall thin skinne, euery one conſiſting of ſix ſmall and narrow greene leaues, very ſharpe pointed at the end, and as it were ending in a ſmall pricke or thorne: in the middle whereof is a ſmall round cup, or rather crowne, of the ſame colour with the leaues and ſtalke, which flower ſmelleth very ſweete, ſomewhat like vnto the reſt of the Ruſh Daffodils: this ſheweth not his flower vntill October, and the froſts quickly following after their flowring, cauſe them ſoone to periſh.

12. *Narciſſus anguſtifolius aureus multiplex.*
The golden double narrow leafed Daffodill.

The leaues of this Daffodill are very narrow, and of a whitiſh greene colour, not aboue foure or fiue inches long, from among which riſeth vp a ſtalke about a foote high, bearing at the top one flower, conſiſting of ſome outer leaues, which are of a yel-

low

low colour, and of many other leaues in the middle being smaller, and set thicke and round together of a more yellow gold colour, but with some whiter leaues among them, the middle part a little pointing forth: the flower standeth long before it doth perfect his colour, and abideth long in flower before the colour decay: the roote is in fashion almost like the ordinary Iunquilia, or Rush Daffodill. I acknowledge this Daffodill hath not his proper place; but because the figure is set in this table, let it thus passe at this time.

13. *Narcissus Iuncifolius luteus flore pleno.* The double Iunquilia, or Rush Daffodill.

The double Rush Daffodill hath his long greene leaues round, like the leaues of the common or ordinary Rush Daffodill, and of the same bigneſſe, among which riseth vp a long slender greene stalke, bearing two or three, seldome more small flowers, yellow and double, that is, with diuers rowes of leaues, hauing the yellow cup such as is in the single flower, broken into small shreads or peeces, running among the leaues of the flower, which peeces in some flowers are not so easily seene, being smaller then in others, this beareth no button or head vnder the flower for seede, his roote is round and blackish, brovvne on the outside, so like vnto the common Rush Daffodill, that it is almost impossible to know the one from the other.

There is another of this kinde, whose flowers are smaller, and not so double, one, two, or three at the most vpon a stalke, and of lesse beauty by much. *Alter minori flore.*

### The Place.

All these Rush Daffodils, doe for the most part grow in Spaine and France, and on the Pyrenæan Mountaines, which are betweene Spaine and France, which Mountains are the Nourseries of many of the finest flowers, that doe adorne the Gardens of these louers of natures pride, and gathered in part by industrious, learned, generous men, inhabiting neare thereunto, and in part by such as make a gaine of their labours, bestowed vpon these things. Onely that with the greene flower was gathered in Barbary, and imparted vnto vs from France.

### The Time.

They flower in the Spring, that is, in March and Aprill, except such whose time is set downe to be in Autumne.

### The Names.

Their names are specified in their titles, and therefore I shall not need to set downe any further repetitions.

To conclude therefore this discourse of true Daffodils, there remaineth to speake of the Sea Daffodils, which (as I said in the beginning) is but one, that is frequent, and doth abide with vs. But there bee some others found about the Cape of good Hope, and in the West Indies, and brought into these parts rather for ostentation, then continuance, where they haue flowred onely once (if peraduenture so often) so that being such strangers, of so remote Countries, and of so diuers natures, I shall but shew you some of them, rather cursorily then curiously; and but onely for your satisfaction, giue you knowledge of two or three of them, that there haue beene seene such in flower, and that they are scarce to bee seene againe, except they bee fetcht a new euery yeare that they be seene.

*Narcissus Marinus, siue tertius Matthioli.*

The great white Sea Daffodill, or Matthiolus his third Daffodill.

The roote of this Daffodill by long continuance, standing in one place without being remoued, groweth to be much greater and larger, then any other Daffodill whatsoeuer,

ſoeuer, and as bigge as any meane Squilla or Sea Onion roote, hauing many long, thicke, and white fibres, or long rootes, diuerſly branched, and ſpread vnder the vpper part of the earth, beſide ſome others that grow downward, and periſh not euery yeare, as the fibres of all, or moſt of the other Daffodils doe; and therefore this plant will not thriue, and beare flowers, if it be often tranſplanted, but rather deſire to abide in one place without remouing, as I ſaid, and that not to be ouerſhadowed, or couered with other herbes ſtanding too neare it, which then will flouriſh, and beare aboundantly: from this roote, which is couered with many blackiſh coates, ariſeth ſix or ſeuen, or more leaues, twice ſo broad almoſt, as any of the former Daffodils, but not ſo long by halfe as many of them, being but ſhort, in compariſon of the breadth, and of a white greene colour; from the middle of which leaues, as alſo from the ſides ſometimes, ſpringeth vp one or two, or more ſtalkes, roundiſh and thicke, and ſometimes a little flat and cornered, a foote high or ſomewhat more, bearing at the toppe, out of a skinnie huske, eight, ten, twelue, or more very large flowers, conſiſting of ſix white leaues a peece, ſpread or laid open, with a white ſhort cuppe or crowne in the middle, lying flat vpon the leaues, cut or diuided into ſix corners (and not whole, as the cuppe or crowne of any other ſingle Daffodill) from euery of which edges, or corners of this cup or crowne, ſtandeth one white long thread, a little crooked or turning vp at the end, tipt with a yellow pendent, and ſome other white threads tipt with yellow pendents, ſtanding alſo in the middle: after the flower is paſt, there come vp great three ſquare heads, wherein the ſeede is contained, which is great, blacke, and round, like vnto the ſeede of other Daffodils, but greater: the flower hath a reaſonable good ſent, but not very ſtrong.

## The Place.

It was firſt found by the Sea ſide, in the Iſle of Sardinia, and on the high Mountaines alſo of the ſame Iſle, where it hath borne by report, thirty fiue flowers vpon a ſtalke: it groweth likewiſe about Illyricum, and in diuers other places.

## The Time.

It ſpringeth later out of the ground then any other Daffodill, that is to ſay, not vntill the later end of March, or beginning of Aprill, and flowreth in the end of May, or the beginning of Iune: the ſeede is ripe in the end of Iuly, or beginning of Auguſt.

## The Names.

The firſt that hath made mention of this Daffodill, was Matthiolus, who placed it in the third place among his Daffodils, and is moſt vſually now adayes called, *Narciſſus tertius Matthioli*, Matthiolus his third Daffodill, the rather, becauſe Cluſius vpon a more mature deliberation, firſt referred it thereunto, but called it at the firſt, *Lilionarciſſus Hemerocallidis facie*, and, as hee ſaith, Iacobus Plateau (who firſt ſent him the figure hereof, with the deſcription) called it *Lilionarciſſus Orientalis*, but Cluſius vpon certaine information, that it grew in the places aforeſaid, miſliked the name of *Orientalis*, and added *Hemerocallis*, which yet is not fit, for that his *Hemerocallis Valentina*, is a plaine Pancration or Sea baſtard Daffodill, whoſe middle cup is longer then the cup of any true Daffodill, which (as I ſaid in the beginning of this Chapter) is the chiefeſt note of difference, betweene a true and a baſtard Daffodill. I receiued the ſeede of this Daffodill among many other ſeedes of rare plants, from the liberality of Mr. Doctor Flud, one of the Phyſitians of the Colledge in London, who gathered them in the Vniuerſity Garden at Piſa in Italy, and brought them with him, returning home from his trauailes into thoſe parts, by the name of *Martagon rariſsimum*, (and hauing ſowne them, expected fourteene yeares, before I ſaw them beare a flower, which the firſt yeare that it did flower, bore foure ſtalkes of flowers,

1 *Narcissus tertius Matthioli.* The great white Sea Daffodill. 2 *Narcissus Indicus Autumnalis.* The Indian Autumne Daffodill. 3 *Narcissus marinus Africanus.* The Sea Daffodill of Africa. 4 *Narcissus marinus exoticus.* The strange Sea Daffodill.

flowers, with euery one of them eight or ten flowers on them) which of all other names, doth least answer the forme or qualities of this plant. It may most fitly be called *Narcissus marinus maximus*, in English, The great Sea Daffodill, both because it is a true Daffodill, and the greatest of all other, and also because it hath not been found, but in Islands, or else in other places neare the Sea. Lobelius entituleth it *Pancratium Indicum alterum vernum, sive Narcissus Indicus alter facie Pancratij Monspeliaci*, but all this is wide from the matter, as may easily be known, by that that hath been said before. It is generally (as I said before) called of all *Narcissus tertius Matthioli*, Matthiolus his third Daffodill, which may either so passe with vs, or as I called it, The great Sea Daffodill, which you will, & so Clusius doth lastly entitle it.

1. *Pancratium Indicum, aut Narcissus Indicus Autumnalis quorundam Lobelij.*
The Indian Autumne Daffodill of Lobel.

This plant hath in my opinion, a farre nearer resemblance vnto an Hyacinthus, then vnto any Daffodill: But because Lobel hath so set it forth, I will so publish it vnto you, leauing it to iudgement. The roote is, as he saith, a span long, and of the thicknesse of a mans arme, couered with many white shells, whereof the outermost are of a darke red or Chesnut colour: the flowers rise vp in September, and October, being eight or ten in number, euery one by it selfe vpon a small footstalke, made of six leaues a peece, somewhat long, narrow, and pointed, like vnto the flowers of the English Colchicum, or Medowe Saffron, of a whitish yellow dunne colour, with six long threads in the middle: the greene leaues are long and broad, and broad pointed.

2. *Narcissus Marinus Africanus, sive Exoticus Lobelij.*
The Sea Daffodill of Africa.

The roote of this strange plant (which of some likenesse is called a Daffodill) is very great, made as it were of many scaly cloues, from whence riseth vp a small short stalke, bearing hard aboue the ground two faire broad greene pointed leaues, more long then broad, so compassing the stalke at the bottome, that it seemeth to run through them: the stalke is spotted with diuers discoloured spots, and is bare or naked from these two leaues vnto the toppe, where it beareth one faire double flower, like vnto a double Auemone, of a delayed reddish colour, tending to a blush, with many threads set about the middle head.

3. *Narcissus Marinus Exoticus.* The strange Sea Daffodill

This strange Sea Daffodill, hath fiue or six large and long leaues of a pale greene colour, from among which riseth vp a strong and bigge stalke, bearing at the toppe, out of a thinne hose or skinne, many very large flowers, made of six long and pointed leaues apeece, of a blewish purple colour, with a large round open cup in the middle, of a sadder colour then the leaues: the roote is very great, yet like vnto other great Daffodils, the outer skins whereof are of a darke browne colour.

The Place.

The Indian Daffodils grew in the vpper part of Hispaniola in the West Indies, and brought hither, where they all soone perished.
The other grew neare the Cape of good Hope, and was brought into the parts of Holland and thereabouts, from whence we had it, & perished also.
The last is vnknowne where it was gathered.

The Time.

The first flowred in Autumne, as it is said.
The other in the first Summer of their bringing.
And so did the last, but the same rootes will not flower with vs againe.

The

The Names.

So much hath been said of their names in their titles, as hath come to our knowledge; and therefore let that suffice.

Thus hauing gone through the whole Family of the true Daffodils, (for so much as hath come to our knowledge) and set them downe euery one by his name, and in his order; it is fit that we speake of their bastard brethren, and shew you them also, in the same order held with the former, as neare as the plenty of variety herein, which is not the like with the former, will giue leaue, that when you know them both by face and name, you may the better know to place or distinguish of others, that haue not passed vnder this rod.

*Pseudonarcissus aureus Hispanicus maximus.*

The great yellow Spanish bastard Daffodill.

The roote of this kinde of Daffodill is reasonable great, and blackish on the outside, desiring to be deepe in the ground; and therefore will runne downe, where it will then encrease into many of-sets, from whence rise vp many thicke, long, and stiffe leaues, of a grayish greene colour, among which riseth vp a round strong stalke, sometimes three foote high or better, bearing at the toppe one onely faire great yellow flower, standing forth right, and not pendulous, consisting of six short and somewhat broad leaues, with a very great, large, and long trunke, of an equall largenesse, but open at the mouth, and turning vp the brimmes a little, which are somewhat crumpled: after the flower is past, there commeth in the place a three square head, containing round blacke seede, like vnto other Daffodils.

*Pseudonarcissus Pyrenæus Hispanico & Anglico similis.*

The Mountaine bastard Daffodill of diuers kindes.

There is much variety in this kinde of bastard Daffodill: For one sort hath verie broad and whitish greene leaues, somewhat short in comparison of others, that are of that breadth: the flower is wholly yellow, but a little paler then the former Spanish kinde, hauing the leaues of his flower long, and somewhat narrow, standing like wings about the middle trunke, which is as long as the leaues, and smaller then in many other of this kinde, but a little yellower then the wings. Another sort hath narrower green leaues then this last, and longer, the flower is all yellow, but the trunke is larger, wider, and more open at the mouth then the former, and almost as large as the former Spanish, but not so high as the last. A third hath the wings of the flower of a Strawe colour, but the trunke is long and narrow, of a faire yellow. A fourth hath such like flowers, but that it is shorter, both the wings and the trunke: Some likewise haue the wings of the flower longer, then the long trunke, and some shorter. Some also are all yellow, and some haue their wings onely a little more pale or white, like the English kinde: Some againe haue their trunkes long and narrow, others haue them larger and wider open, and crumpled at the brimmes; so that it is needlesse, to spend a great deale of time and labour vpon such smally respected flowers, but that in the beholding of them, we may therein admire the worke of the Creatour, who can frame such diuersity in one thing: But this is beside the text, yet not impertinent.

*Pseudonarcissus pallidus præcox.* The early Strawe coloured bastard Daffodill.

The leaues of this Daffodill are of a meane size, betweene the broadest and the narrower kindes, of a grayish greene colour, and not very long: the stalke riseth vp a foot high or more, whereon standeth one large great flower, equalling the greatest Spanish bastard Daffodill, before described, in the largenesse of his trunke, and hauing the brimmes turned vp a little, which maketh it seeme the larger: the wings or outer leaues are in a maner as short, as they are in the greatest Spanish kinde, (and not long flagging down, like vnto the Mountain kinds) and stand straight outright: all the whole flower is

of one euen colour, that is, of a fine pale yellow, somewhat like vnto the colour of a Lemon peele or rinde, but somewhat whiter, which vsually we call a Strawe colour: the greatnesse of the flower, the earlinesse of the flowring, and the difference of colour from all the rest of this kinde, hath made me entreate of it apart by it selfe, as being no lesse worthy.

*Pseudonarcissus Hispanicus flore albo maior.*
The great white Spanish bastard Daffodill.

This bastard Daffodill hath diuers leaues rising vp together, long and broad, somewhat like vnto the first Spanish kinde, but a little broader, and of a whiter greene colour, yet not so white, as in the lesser Spanish white kindes, hereafter described: among these leaues riseth vp a round strong stalke, about two foote high, bearing one white flower at the toppe, bending downe the head, as all these white kindes doe, but is not of so pure a white, as the lesser kindes that follow, yet whiter then the greatest white Spanish kinde, next of all to be described: the whole flower, as well trunke as wings, is much larger then the lesser white kindes, and almost equalling the first Spanish yellow, but a little longer and narrower, a little crumpled and turning vp at the brimmes: the head and seede are like the first; the roote is greater and thicker then the first Spanish, and doth not encrease so much, nor is couered with a blacke, but rather with a whitish coate.

*Pseudonarcissus Hispanicus maximus albidus.*
The greatest Spanish white bastard Daffodill.

This kinde of bastard Daffodill is very like the last mentioned Daffodill, both in leaues and flowers, but larger in both: the flower of this is not full so white, but hath some shew of paleness therein, and more vpon the first opening of the flower then afterwards, and is as great altogether, as the great Spanish yellow, at the least with a longer, and somewhat narrower trunke: the seede is like vnto the former, and so is the roote also, but greater, being white on the outside, and not blacke.

*Pseudonarcissus Hispanicus flore albo medius & minor.*
The two lesser white Spanish bastard Daffodils.

There are two other of these kindes of white Spanish Daffodils, one greater or lesser then the other, but neither of them so great as the former. The leaues of both are of a whitish greene colour, one a little broader then the other: the flowers of both are pure white, and bending downe the heads, that they almost touch the stalke againe, the greater flower hath the longer and narrower trunke; and the lesser flower, the shorter and wider open, yet both a little crumpled at the edges or brimmes: the rootes of both are like one vnto another, but differ in the greatnesse. From the seede of these haue sprung much variety, few or none keeping either colour or height with the mother plants.

*Pseudonarcissus Anglicus vulgaris.* Our common English wilde bastard Daffodill.

This bastard Daffodill is so common in all England, both in Copses, Woods, and Orchards, that I might well forbeare the description thereof, and especially, in that growing wilde, it is of little respect in our Garden: but yet, lest I bee challenged of ignorance in common plants, and in regard of some variety therein worth the marking, I will set downe his description and variety as briefly as I may: It hath three or foure grayish greene leaues, long and somewhat narrow, among which riseth vp the stalke, about a span high or little higher, bearing at the toppe, out of a skinnie huske, as all other Daffodils haue, one flower (although sometimes I haue seene two together) somewhat large, hauing the six leaues that stand like wings, of a pale yellow colour, and the long trunke in the middle of a faire yellow, with the edges or brimmes a little crumpled or vneuen: after the flower is past, it beareth a round head, seeming three square, containing round blacke seede; the roote is somewhat blackish on the outside.

But

1 *Pseudonarcissus Hispanicus maximus aureus.* The great yellow Spanish bastard Daffodill. 2 *Pseudonarcissus Pyrenaus variformis.* The Mountaine bastard Daffodill of diuers kindes 3 *Pseudonarcissus Hispanicus maior albus.* The greater white Spanish bastard Daffodill. 4 *Pseudonarcissus Hispanicus minor albus.* The lesse Spanish white bastard Daffodill. 5 *Pseudonarcissus tubo sexangulari.* The six cornered bastard Daffodill. 6 *Pseudonarcissus maximus aureus, siue Roseus Tradescanti.* Iohn Tradescants great Rose Daffodill. 7 *Pseudonarcissus aureus Anglicus maximus.* Master Wilmers great double Daffodill. 8 *Pseudonarcissus Hispanicus aureus flore pleno.* The double Spanish Daffodill, or Parkinsons double Daffodill. 9 *Pseudonarcissus Gallicus maior flore pleno.* The greater double French Daffodill. 10 *Pseudonarcissus Anglicus flore pleno.* The double English Daffodill, or Gerrards double Daffodill.

But there is another of this kinde like vnto the former, whose further description you haue here before; the wings of which flower are much more white then the former, and in a manner of a milke white colour, the trunke remaining almost as yellow as the former, and not differing in any thing else.

*Pseudonarcissus tubo sexangulari.* The six cornered bastard Daffodill.

This kinde of Daffodill hath two or three long, and somewhat broader leaues then the last, between which commeth forth a stalke, bearing one flower somewhat large, hauing the six outer leaues of a pale yellow colour, and the long trunke plaited or cornered all along vnto the very edge into six parts, of a little deeper yellow then the wings.

The Place.

The first great Spanish kinde was brought out of Spaine. The rest from the Pyrenæan Mountaines, onely the last sauing one is plentifull in our owne Countrey, but the white sort of that kinde came with the rest from the same Mountaines.

The Time.

The pale or third kinde, and the English bee the most early, all the rest flower in Aprill, and the greatest yellow somewhat earlier, then the other greater or lesser white.

The Names.

Their seuerall names are expressed in their titles sufficient to distinguish them, and therefore there needeth no more to be said of them.

1. *Pseudonarcissus aureus maximus flore pleno, sive Roseus Tradescanti.*
The greatest double yellow bastard Daffodill, or
Iohn Tradescant his great Rose Daffodill.

This Prince of Daffodils (belongeth primarily to Iohn Tradescant, as the first founder thereof, that we know, and may well bee entituled the Glory of Daffodils) hath a great round roote, like vnto other Daffodils, couered with a brownish outer skinne or peeling, from whence riseth vp foure or fiue somewhat large and broad leaues, of a grayish greene colour, yet not fully so long and large as the next following Daffodill: from the middle whereof riseth vp a stalke almost as high and great as it, bearing at the toppe (out of a skinnie huske) one faire large great flower (the budde, before it breake open, being shorter and thicker in the middle, and ending in a longer and sharper point then any of the other Daffodils) very much spread open, consisting of smaller and shorter leaues then the next, but more in number, and thicker and rounder set together, making it seeme as great and double as any Prouince Rose, and intermixt with diuers yellow and pale leaues, as it were in rowes one vnder another. It abideth long in flower, and spreadeth, by standing long, to be the broadest in compasse of any of the Daffodils, but falleth away at the last without giuing any seede, as all double Daffodils doe.

2. *Pseudonarcissus aureus Anglicus maximus.* Mr. Wilmers great double Daffodill.

The other great double Daffodill doth so neare resemble our ordinary English double kinde, that I doe not finde therein any greater difference, then the largenesse both of leaues and flowers, &c. and the statelinesse of growth. It beareth three or foure large, long, and broad leaues, somewhat longer and broader then the former, and of a whitish greene colour: the stalke riseth to bee two foote high, growing (in a fruitfull and fat soyle) strong, and somewhat round, bearing at the toppe, out of a thin skinne, one great and faire double flower, each leafe whereof is twice as large and broad

broad as the former, diuerſly intermixt with a rowe of paler, and a rowe of deeper yellow leaues, wholly diſperſed throughout the flower, the pale colour as well as the deeper yellow, in this as in the other ſmall Engliſh kinde, growing deeper by ſtanding: ſometimes the leaues hereof are ſcattered, and ſpread wholly, making it ſhew a faire, broad, open flower: and ſometimes the outer leaues ſtand ſeparate from the middle trunke, which is whole and vnbroken, and very thicke of leaues: and ſometimes the middle trunke will bee halfe broken, neither expreſſing a full open double flower, nor a cloſe double trunke, as it is likewiſe ſeene in the ſmall Engliſh kinde, as ſhall bee declared in his place: this beareth no ſeede; the roote hereof is thicke and great, and encreaſeth as well as any other Daffodill.

### 3. *Pſeudonarciſſus aureus Hiſpanicus flore pleno.*
The great double yellow Spaniſh baſtard Daffodill, or Parkinſons Daffodill.

This double Spaniſh Daffodill hath diuers leaues riſing from the roote, ſtiffer, narrower, and not of ſo whitiſh a greene colour as the former, but more ſullen or grayiſh, plainely reſembling the leaues of the ſingle great kinde, from whence this hath riſen: the ſtalke hereof likewiſe riſeth almoſt as high as it, and neare the height of the laſt recited double, bearing one double flower at the toppe, alwayes ſpread open, and neuer forming a double trunke like the former, yet not ſo faire and large as it, the outermoſt leaues whereof being of a greeniſh colour at the firſt, and afterward more yellow, doe a little turne themſelues backe againe to the ſtalke, the other leaues are ſome of a pale yellow, and others of a more gold yellow colour, thoſe that ſtand in the middle are ſmaller, and ſome of them ſhew as if they were hollow trunked, ſo that they ſeeme to be greeniſh, whitiſh, yellow, and gold yellow, all mixed one among another: the root is great, round, and whitiſh on the inſide, couered with darke coloured skinnes or peelings. I thinke none euer had this kinde before my ſelfe, nor did I my ſelfe euer ſee it before the yeare 1618. for it is of mine own raiſing and flowring firſt in my Garden.

### 4. *Pſeudonarciſſus Gallicus maior flore pleno.*
The greater double French baſtard Daffodill.

This greater double Daffodill, hath his whitiſh greene leaues longer and broader then the ſmaller French kinde, hereafter following, to bee deſcribed, and broader, longer, and more limber then the double Engliſh kinde: the ſtalke riſeth vp not much higher, then the ſmaller French kinde, but a little bigger, bearing at the top one great double flower, which when it is fully and perfectly blowne open (which is but ſeldome; for that it is very tender, the leaues being much thinner, and thereby continually ſubiect, vpon any little diſtemperature of the time, to cleaue ſo faſt one vnto another, that the flower cannot blow open faire) is a faire and a goodly flower, larger by halfe then the ſmaller kinde, and fuller of leaues, of the ſame pale whitiſh yellow, or Lemon colour, with the leſſer, or rather a little whiter, and not ſet in the ſame order of rowes as it is, but more confuſedly together, and turning backe the ends of the outermoſt leaues to the ſtalke againe, and hauing the bottome of the flower on the backſide ſomewhat greene, neither of which is found in the leſſer kinde: the roote is very like vnto the leſſer kinde, but a little bigger and longer.

### 5. *Pſeudonarciſſus Anglicus flore pleno.*
The double Engliſh baſtard Daffodill, or Gerrards double Daffodill.

The leaues of this double Daffodill are very like vnto the ſingle kinde, being of a whitiſh greene colour, and ſomewhat broad, a little ſhorter and narrower, yet ſtiffer then the former French kinde: the ſtalke riſeth vp about a foote high, bearing at the toppe one very double flower, the outermoſt leaues being of the ſame pale colour, that is to bee ſeene in the wings of the ſingle kinde; thoſe that ſtand next them, are ſome as deepe a yellow as the trunke of the ſingle, and others of the ſame pale colour, with ſome greene ſtripes on the backe of diuers of the leaues: thus is the whole flower variably intermixt with pale and deepe yellow, and ſome greene ſtripes among them,

when

when it is fully open, and the leaues disperſed and broken. For ſometimes the flower ſheweth a cloſe and round yellow trunke in the middle, ſeparate from the pale outer wings, which trunke is very double, ſhewing ſome pale leaues within it, diſperſed among the yellow: And ſometimes the trunke is more open, or in part broken, ſhewing forth the ſame colours intermixt within it: the flower paſſeth away without giuing any ſeede, as all other bulbous rootes doe that beare double flowers: the roote is ſmall, very like vnto the French double kindes, eſpecially the leſſer, that it is verie hard to know the one from the other.

### The Place.

The firſt and greateſt kinde, we had firſt from Iohn Tradeſcante (as I ſaid before) whether raiſed from ſeed, or gained from beyond Sea, I know not.

The ſecond we firſt had from Vincent Sion, borne in Flanders, dwelling on the Banke ſide, in his liues time, but now dead; an induſtrious and worthy louer of faire flowers, who cheriſhed it in his Garden for many yeares, without bearing of any flowers vntill the yeare 1620. that hauing flowred with him, (and hee not knowing of whom hee receiued it, nor hauing euer ſeene the like flower before) he ſheweth it to Mr. Iohn de Franqueuille, of whom he ſuppoſed he had receiued it, (for from beyond Sea he neuer receiued any) who finding it to bee a kinde neuer ſeene or knowne to vs before, cauſed him to reſpect it the more, as it is well worthy. And Mr. George Wilmer of Stratford Bowe Eſquire, in his liues time hauing likewiſe receiued it of him (as my ſelfe did alſo) would needes appropriate it to himſelfe, as if he were the firſt founder thereof, and call it by his owne name Wilmers double Daffodill, which ſince hath ſo continued.

The third is of mine owne foſtering or raiſing, as I ſaid before; for aſſuredly, it is riſen from the ſeede of the great Spaniſh ſingle kinde, which I ſowed in mine owne Garden, and cheriſhed it, vntill it gaue ſuch a flower as is deſcribed.

The fourth is not certainly knowne where his originall ſhould be: Some thinke it to be of France, and others of Germany.

The laſt is aſſuredly firſt naturall of our owne Countrey, for Mr. Gerrard firſt diſcouered it to the world, finding it in a poore womans Garden in the Weſt parts of England, where it grew before the woman came to dwell there, and, as I haue heard ſince, is naturall of the Iſle of Wight.

### The Time.

They doe all flower much about one time, that is, from the middle or end of March, as the yeare is forward, vnto the middle of Aprill.

### The Names.

Vpon the three firſt I haue impoſed the names in Latine, as they are expreſſed in their titles: and for the Engliſh names, if you pleaſe, you may let them paſſe likewiſe as they are expreſſed there alſo, that thereby euery one may be truely diſtinguiſhed, and not confounded. The fourth, beſides the name in the title, is called of ſome *Narciſſus Germanicus*, which whether it be of Germany, or no, I know not; but that the name ſhould import ſo much. The laſt doth vſually carry Mr. Gerrards name, and called Gerrards double Daffodill.

1. *Pſeudonarciſſus anguſtifolius flore flaueſcente tubo quaſi abſciſſo.*
The narrow leafed baſtard Daffodill with the clipt trunke.

This kinde of Daffodill hath long and narrow grayiſh greene leaues, bearing one ſingle flower at the toppe of his ſtalke, like vnto the former ſingle baſtard kindes, before

fore specified, hauing his outer leaues of a pale yellow colour, and his trunke of a deeper yellow: the chiefe differences in this from the former, is in the leaues, being narrow, and then in the trunke of the flower, which is not crumpled or turned vp, as most of the other are; and that the brimmes or edges of the flower is as if it had beene clipt off, or cut euen.

2. *Pseudonarcissus Hispanicus medius & minor luteus.*
The two lesser Spanish yellow bastard Daffodils.

These two lesser kindes of Spanish Daffodils, doe but differ in greatnesse the one from the other, and not in any thing else; so that in declaring the one, you may vnderstand the other to bee a little greater. The lesser then hath three or foure narrow short whitish greene leaues, from among which commeth forth a short stalke, not aboue an hand breadth, or halfe a foote high, bearing one single flower, not fully standing outright, but a little bending downe, consisting of six small leaues, standing as wings about a small, but long trunke, a little crumpled at the brimmes: the whole flower, as well leaues as trunke, are of one deepe yellow colour, like vnto the great Spanish kinde: the roote is but small, and couered with a darkish coate. The other is in all parts greater, and (as I said) differeth not else.

3. *Pseudonarcissus Hispanicus luteus minimus.*
The least Spanish yellow bastard Daffodill.

The leaues of this small kinde are smaller and shorter then the former, seldome exceeding the length of three inches, and very narrow withall, but of the same grayish greene colour with the former: euery flower standeth vpon a small and short foote-stalke, scarce rising aboue the ground, so that his nose, for the most part, doth lye or touch the ground, and is made after the same fashion, and of the same colour with the former, but much smaller, as his roote is so likewise.

4. *Pseudonarcissus Gallicus minor flore pleno.*
The lesser French double bastard Daffodill.

The rootes of this lesser French kinde (if I may lawfully call it, or the greater kinde before specified, a bastard Daffodill; for I somewhat doubt thereof, in that the flower of either is not made after the fashion of any of the other bastard Daffodils, but doth more nearely resemble the forme of the double white Daffodill, expressed before among the true Daffodils) are like vnto the double English kinde, as also to the former double greater French kinde, and the leaues are of the same whitish greene colour also, but narrower and not longer: the stalke riseth a little higher then the English, and not fully so high as the greater French, bearing one faire double flower thereon, of a pale yellow or Lemon colour, consisting of six rowes of leaues, euery rowe growing smaller then other vnto the middle, and so set and placed, that euery leafe of the flower doth stand directly almost in all, one vpon or before another vnto the middle, where the leaues are smallest, the outermost being the greatest, which maketh the flower seeme the more beautifull: this and the greater kinde hath no trunke, or shew of any other thing in the middle, as all or most of the other former double bastard Daffodils haue, but are flowers wholly composed of leaues, standing double euen to the middle.

The Place.

The first is vndoubtedly a naturall of the Pyrenæan Mountaines.
The Spanish kindes grew in Spaine, and
The French double kinde about Orleance in France, where it is said to grow plentifully.

The Time.

The first flowreth at the end of March.

The

The Spanish kindes are the most early, flowring betimes in March.
The French double doth flower presently after.

The Names.

More cannot bee said or added, concerning the names of any of these Daffodils, then hath been set downe in their titles: onely the French kinde is most vsually called Robinus his Daffodill.

*Pseudonarcissus Iuncifolius albus.* The white bastard Rush Daffodill, or Iunquilia.

This bastard Rush Daffodill hath two or three long and very greene leaues, very like vnto the small yellow Rush Daffodill, formerly described, but not altogether so round, among which riseth vp a short stalke, seldome halfe a foote high, bearing at the toppe, out of a small skinnie huske, one small white flower, sometime declining to a pale colour, hauing six small and short leaues, standing about the middle of the trunke, which is long, and much wider open at the mouth, then at the bottome: the small outer leaues or wings are a little tending to greene, and the trunke (as I said) is either white, or whitish, hauing the brimmes a little vneuen: the seede is small, blacke, and round, like vnto other Rush Daffodils, but smaller.

*Pseudonarcissus Iuncifolius luteus maior.*
The greater yellow Iunquilia, or bastard Daffodill.

The leaues of this greater kinde are longer, greater, and a little broader then the former; the stalke also is higher, and the flower larger, more open at the mouth and crumpled, then the white, but wholly of a yellow colour: the seede and the roots are bigger, according to the proportion of the plant.

*Pseudonarcissus Iuncifolius luteus minor.* The lesser yellow bastard Iunquilia.

This is so like vnto the last in all things, that I shall not neede to trouble you with repetitions of the same things formerly spoken; the chiefest difference is the smalnesse of the plant in all parts.

*Pseudonarcissus Iuncifolius luteus serotinus.* The late yellow bastard Iunquilia.

There is likewise a third kinde, as great as the greater yellow, and in all his parts expressing and equalling it, but is accounted the fairer, and flowreth somewhat later.

The Place.

The Pyrenæan Hils haue afforded vs all these varieties, and wee preserue them carefully; for they are all tender.

The Time.

All these flower in Aprill, except the last, which is a moneth later.

The Names.

The French and Lowe-Countrey men call them *Trompettes*, that is, Trumpets, from the forme of the trunke; wee sometimes call them also by that name, but more vsually bastard Iunquilia's.

*Pseudonarcissus marinus albus, Pancratium vulgo.*
The white Sea bastard Daffodill.

The Sea bastard Daffodill (to conclude this Chapter, and the discourse of Daffodils)

1 *Pseudonarcissus tubo quasi abscisso.* The bastard Daffodill with the clipt trunke. 2 *Pseudonarcissus Hispanicus minor.* The lesser Spanish bastard Daffodill. 3 *Pseudonarcissus Hispanicus minimus.* The least Spanish bastard Daffodill. 4 *Pseudonarcissus Gallicus minor flore pleno.* The lesser double French bastard Daffodil. 5 *Pancratium flore albo.* The white Sea bastard Daffodil. 6 *Pseudonarcissus Iunciformis luteus maior.* The greater yellow bastard Iunquilla. 7 *Pseudonarcissus Iunciformis luteus minor.* The lesser yellow bastard Iunquilla. 8 *Pseudonarcissus Iunciformis luteus serotinus.* The late yellow bastard Iunquilla. 9 *Leucoium bulbosum præcox maius.* The great early bulbous Violet. † *Leucoium bulbosum præcox minus.* The lesser early bulbous Violet. 10 *Leucoium bulbosum autumnale.* The small Autumne bulbous Violet. 11 *Leucoium bulbosum maius serotinum.* The great late flowring bulbous Violet.

dils) hath diuers broad whitish greene leaues, but not very long, among which riseth vp a stiffe round stalke, at the top whereof breaketh out of a great round skinny huske, fiue or six flowers, euery one made somewhat of the fashion of the great bastard Rush Daffodill, but greater, and wholly white; the six leaues, being larger and longer then in the Rush kinde, and extending beyond the trunke, are tipt with greene at the point of each leafe, and downe the middle likewise on the backside. The trunke is longer, larger, and wider open at the mouth, cut in or indented at the brims or edges, and small at the bottome, with diuers white threeds in the middle, and is very sweet: vnder the flower is a round greene head, which groweth very great, hauing within it, when it is ripe, flat and blacke seede: the roote is great and white.

*Flore luteo, & flore rubro.*

It is reported, that there are found other sorts; some that beare yellow flowers, and others that beare red: but we haue seene none such, and therefore I can say no more of them.

## The Place.

This kinde groweth neare the Sea side, both in Spaine, Italy, and France, within the Straights, and for the most part, vpon all the Leuant shoare and Islands also, but will seldome either flower, or abide with vs in these colder Countries, as I haue both seene by those that I receiued from a friend, and heard by others.

## The Time.

It flowreth in the end of Summer, that is, in August and September.

## The Name.

Diuers doe call it *Pancratium*, as the learned of Mompeher, and others, with the addition of *flore Lilij*, after they had left their old errour, in taking it to be *Scylla*, and vsing it for *Scylla*, in the *Trochisces* that go into Andromachus Treakle. The learned of Valentia in Spaine, as Clusius saith, doe call it *Hemerocallis*, thinking it to be a Lilly; and Clusius doth thereupon call it, *Hemerocallis Valentina*: but in my opinion, all these are deceiued in this plant; for it is neither a Lilly, to haue the name of *Hemerocallis* giuen vnto it, nor *Scylla*, nor *Pancratium*, as many doe yet call it: for certainly this is a kinde of Daffodill; the forme both of roote, leafe, and flower, doth assure me that haue seene it, and not *Pancratium*, which (as Dioscorides testifieth) is a kinde of *Scylla*, and in his time called *Scylla*, with a red roote, and a leafe like a Lilly, but longer, and was vsed both with the same preparation and quantity, and for the same diseases that *Scylla* was vsed, but that his force was weaker: all which doth plainly shew the errours that many learned men haue been conuersant in, and that all may see how necessary the knowledge of Herbarisme is to the practice of Physicke; And lest the roote of this Sea bastard Daffodill bee vsed in the stead of an wholsome remedy, which (as Clusius maketh mention) was deadly to him that did but cut his meate with that knife, which had immediately before cut this roote, and done in malice by him, that knew the force thereof, to kill his fellow, it working the more forceably by the euill attracting quality of the iron.

## The Vertues of Daffodils in generall.

Howsoeuer Dioscorides and others, doe giue vnto some of them speciall properties, both for inward and outward diseases, yet know I not any in these dayes with vs, that apply any of them as a remedy for any griefe, whatsoeuer Gerrard or others haue written.

## Chap. X.

### *Leucoium bulbosum.* The bulbous Violet.

HAuing thus set downe the whole family, both of the true and bastard Daffodils, I should next set in hand with the Hyacinths; but because *Leucoium bulbosum*, The bulbous Violet is a plant that doth challenge a place next vnto the Daffodils, as most nearly partaking with them, and a little with the Hyacinthes, I must of necessity interpose them, and shew their descriptions and differences, whereof some are early, of the first Spring, others later, and some of the Autumne.

### *Leucoium bulbosum præcox maius.* The greater early bulbous Violet.

This bulbous Violet hath three or foure very greene, broad, flat, and short leaues, among which riseth vp a naked greene stalke, bearing out of a small skinny hose (as the former Daffodils doe) one white flower, hanging downe his head by a very small foot-stalke, made of six leaues, of an equall length, euery one whereof is tipt at the end with a small greenish yellow spot: after the flower is past, the head or seed-vessell groweth to be reasonable great, somewhat long and round, wherein is contained hard round seede, which being dry, is cleare, and of a whitish yellow colour: the roote is somewhat like a Daffodill roote, and couered with a blackish outside or skinne.

### *Leucoium bulbosum præcox minus.* The lesser early bulbous Violet.

This lesser kinde riseth vp with two narrow grayish greene leaues, between which commeth forth the stalke, fiue or six inches high, bearing one small pendulous flower, consisting of three white leaues, which are small and pointed, standing on the outside, and hauing three other shorter leaues, which seeme like a cup in the middle, being each of them round at the ends, and cut in the middle, making the forme of an heart, with a greene tippe or spot at the broad end or edge: the seede is whitish, inclosed in long and round heads, like the former, but lesser: the roote is like a small Daffodill, with a blackish gray coate, and quickly diuideth into many of-sets.

There is another of this kinde, that came among other bulbous rootes from Constantinople, and differeth in nothing from it, but that it is a little greater, both in root, leafe, and flower. *Minus Byzantinum.*

#### The Place.

The two first are found in many places of Germany, and Hungary. The third, as I said, was brought from Constantinople.

#### The Time.

The two lesser sorts doe most commonly flower in February, if the weather be any thing milde, or at the furthest in the beginning of March, but the first is seldome in flower, before the other be well neare past, or altogether.

#### The Names.

Lobel and Dodonæus call the lesser kinde *Leucoium triphyllum*, and *Leuconarcissolirion triphyllum*, of the three leaues in the flower. Some doe call it *Viola bulbosa alba.* The first or greater kinde is called by Lobel, *Leuconarcissolirion pauciioribus floribus*; and by Dodonæus, *Leucoium bulbosum hexaphyllum.* We doe most vsually call them, *Leucoium bulbosum præcox maius, & minus*, The greater, or the lesser early bulbous Violet. In Dutch, *Somer Sottekens*, and not *Druifkens*, which are Grape-flowers, as some haue thought.

1. *Leucoium bulbosum Vernum minimum.*
The small bulbous Violet of the Spring.

This small *Leucoium* sendeth forth his small and long greene leaues, like haires in Autumne, and before Winter, which abide greene vntill Aprill, and then wither away quite, and about May there ariseth vp a naked slender stalke, at the toppe whereof breake forth two small white flowers, made of six leaues a peece, hanging downe their heads, the three inner leaues being a little larger then the three outward, a little reddish neare the stalke, and very sweet: the root is small and round, and couered with a darke coate.

2. *Leucoium bulbosum Autumnale.* The small Autumne bulbous Violet.

As the former small *Leucoium* sprang vp with his leaues without flowers in Autumne, so this contrariwise, riseth vp with his slender browniſh stalke of flowers in Autumne, before any greene leaues appeare, whereon stand two or three very small snow white pendulous flowers, consisting of six leaues a peece, and a little reddish at the bottome of the flower next vnto the stalke, so like vnto the former, that one would take them to be both one: after which, there grow small browne heads, containing small, blacke, round seed; after the flower is past, and the seede is ripening, and sometimes after the heads are ripe, the leaues begin to spring vp, which when they are full growne, are long, greene, and as small, or smaller then the leaues of the Autumne Hyacinth, which abide all the Winter, and Spring following, and wither away in the beginning of Summer: the roote is small, long, and white.

3. *Leucoium maius bulbosum serotinum.*
The great late flowring bulbous Violet.

The late bulbous Violet hath three or foure broad flat greene leaues, very like vnto the first, but longer, among which riseth vp a flattish stalke, being thicker in the middle then at both edges, on the toppe whereof stand three or foure flowers, hanging downe their heads, consisting of six leaues a peece, all of an equall length and bigneſſe, wholly white, except that each leafe hath a greene tippe at the end of them: the seede hereof is blacke and round; the roote is reasonable great and white.

The Place.

The two former small ones were first found in Spaine, and Portugall, and sent to me by Guillaume Boel; but the first was so tender, that scarce one of a score sprang with me, or would abide. The greatest haue beene found wilde in Germany and Austria.

The Time.

The small ones haue their times expressed in their titles and descriptions, the last flowreth not vntill May.

The Names.

These names that are set downe in their titles, doe passe with all Herbarists in these daies.

The Vertues.

Wee haue not knowne these plants vsed Physically, either inwardly or outwardly, to any purposes in these dayes.

## Chap. XI.

### *Hyacinthus.* The Hyacinth or Iacinth.

THe Iacinths are next to be entreated of, whereof there are many more kindes found out in these later times, then formerly were knowne, which for order and method sake, I will digest vnder seuerall sorts, as neare as I can, that auoiding confusion, by enterlacing one among another, I may the better put euery sort vnder his owne kinde.

#### *Hyacinthus Indicus maior tuberosa radice.*
#### The greater Indian knobbed Iacinth.

I haue thought fittest to begin with this Iacinth, both because it is the greatest and highest, and also because the flowers hereof are in some likenesse neare vnto a Daffodill, although his roote be tuberous, and not bulbous as all the rest are. This Indian Iacinth hath a thicke knobbed roote (yet formed into seuerall heads, somewhat like vnto bulbous rootes) with many thicke fibres at the bottome of them; from the diuers heads of this roote arise diuers strong and very tall stalkes, beset with diuers faire, long, and broad leaues, ioyned at the bottome close vnto the stalke, where they are greatest, and grow smaller to the very end, and those that grow higher to the toppe, being smaller and smaller, which being broken, there appeare many threeds like wooll in them: the toppes of the stalkes are garnished with many faire large white flowers, each whereof is composed of six leaues, lying spread open, as the flowers of the white Daffodill, with some short threeds in the middle, and of a very sweete sent, or rather strong and headie.

#### *Hyacinthus Indicus minor tuberosa radice.*
#### The smaller Indian knobbed Iacinth.

The roote of this Iacinth is knobbed, like the roote of Arum or Wake Robin, from whence doe spring many leaues, lying vpon the ground, and compassing one another at the bottome, being long and narrow, and hollow guttered to the end, which is small and pointed, no lesse woolly, or full of threeds then the former: from the middle of these leaues riseth vp the stalke, being very long and slender, three or foure foot long, so that without it be propped vp, it will bend downe, and lye vpon the ground, whereon are set at certaine distances many short leaues, being broad at the bottome, where they doe almost compasse the stalke, and are smaller toward the end where it is sharpe pointed: at the top of the stalke stand many flowers, with a small peece of a green leafe at the bottome of euery foot-stalke, which seeme to bee like so many white Orientall Iacinths, being composed of six leaues, which are much thicker then the former, with six chiues or threeds in the middle, tipt with pale yellow pendents.

### The Place.

They both grow naturally in the West Indies, from whence being first brought into Spaine, haue from thence been dispersed vnto diuers louers of plants.

### The Time.

They flower not in these cold Countries vntill the middle of August, or not at all, if they bee not carefully preserued from the iniury of our cold Winters; and then if the precedent Summer be hot, it may be flower a moneth sooner.

### The Names.

Clusius calleth the lesser (for I thinke hee neuer saw the first) *Hyacinthus Indicus*

*Indicus tuberosa radice*, that is in English, The Indian Iacinth with a tuberous roote: Some would call these *Hyacinthus Eriophorus Indicus*, that is, The Indian woolly Iacinth, because they haue much wooll in them when they are broken; yet some doe doubt that they are not two plants seuerall, as of greater and lesser, but that the greatnesse is caused by the fertility of the soyle wherein it grew.

### 1. *Hyacinthus Botroides maior Moschatus, siue Muscari flore flauo.*
### The great yellow Muske Grape-flower, or yellow Muscari.

This Muske Iacinth or Grape-flower, hath fiue or six leaues spread vpon the ground in two or three heads, which at the first budding or shooting forth out of the ground, are of a reddish purple colour, and after become long, thicke, hollow, or guttered on the vpperside, of a whitish greene colour, and round and darke coloured vnderneath: in the middle of these heads of leaues, rise vp one or two hollow weake browniſh stalkes, sometimes lying on the ground with the weight of the flowers, (but especially of the seede) yet for the most part standing vpright, when they are laden towards the toppe, with many bottle-like flowers, which at their first appearing, and vntill the flowers begin to blow open, are of a browne red colour, and when they are blowne, of a faire yellow colour, flowring first below, and so vpwards by degrees, euery one of these flowers is made like vnto a little pitcher or bottle, being bigge in the belly, and small at the mouth, which is round, and a little turned vp, very sweete in smell, like vnto Muske, whereof it tooke the name *Muscari*; after the flowers are past, there come three square thicke heads, puffed vp as if it were bladders, made of a spongie substance, wherein are here and there placed blacke round seed: the roote is long, round, and very thicke, and white on the outside, with a little woollinesse on them, being broken, and full of a slimie iuice, whereunto are annexed thicke, fat, and long fibres, which perish not as most of the other Iacinths; and therefore desireth not to bee often remoued, as the other sorts may.

### 2. *Hyacinthus Botroides maior Moschatus, seu Muscari flore cineritio.*
### The Ashcoloured Muske Grape-flower, or Muscari.

This Muscari differeth not in rootes, or forme of leaues or flowers from the former, the chiefe differences are these: the leaues hereof do not appeare so red at the first budding out of the ground, nor are so darke when they are fully growne; the stalke also most vsually hath more store of flowers thereon, the colour whereof at the first budding is a little duskie, and when they are full blowne, are of a bleake, yet bright ashcolour, with a little shew of purple in them, and by long standing change a little more gray; being as sweete, or as some thinke, more sweete then the former: the roote (as I said) is like the former, yet yeeldeth more encrease, and will better endure our cold clymate, although it doth more seldome giue ripe seede.

### 3. *Hyacinthus Botroides maior Moschatus, siue Muscari flore rubro.*
### The red Muske Grape-flower.

This kinde (if there be any such, for I am in some doubt thereof) doth chiefly differ in the colour of the flower from the first, in that this should beare flowers when they are blowne, of a red colour tending to yellownesse.

### 4. *Hyacinthus Botroides maior Moschatus, siue Muscari flore albo.*
### The white Muske Grape-flower.

This also is said to haue (if there bee such an one) his leaues like vnto the second kinde, but of a little whiter greene, and the flowers pale, tending to a white: the roots of these two last are said vsually not to grow to be so great as of the former two.

### The Place.

The rootes of the two first sorts, haue been often sent from Constantino-

ple,

1 *Hyacinthus Indicus maior tuberosa radice.* The greater Indian knobbed Iacinth. 2 *Hyacinthus Indicus minor tuberosa radice.* The lesser Indian knobbed Iacinth. 3 *Muscari flore flauo.* The yellow Muscari. 4 *Muscari flore cineritio.* The ascheoloured Muscari. 5 *Hyacinthus Botroides caruleus amœnus.* The skie coloured Grape-flower. 6 *Hyacinthus Botroides flore albo.* The white Grape-flower. † *Hyacinthus Botroides ramosus.* The branched Grape-flower.

ple, among many other ſorts of rootes, and it may be come thither from beyond the Boſphorus in Aſia; we haue them in our Gardens.

The other two ſorts are ſprung (it is probable, if they be *in rerum natura*) from the ſeede of the two former; for we could neuer get ſuch from Conſtantinople, as if the Turkes had neuer knowledge of any ſuch.

## The Time.

They flower in March or Aprill, as the yeare is temperate, but the firſt is ſooneſt vp out of the ground.

## The Names.

The two former haue beene ſent from Turkie by the name of *Muſchoromi* and *Dipcadi*. Matthiolus calleth it *Bulbus vomitorius*, ſaying that no root doth more prouoke vomit then it. Caſpar Bauhinus doth moſt properly call it *Hyacinthus Moſchatus*. It is moſt generally called *Muſcari*, by all Herbariſts and Floriſts, yet becauſe it doth ſo neerely reſemble the Grape-flower, I haue named it *Hyacinthus Botroides maior Muſchatus*, to put a difference from the leſſer Grape-flowers that follow; in Engliſh, The great Muske Grape-flower, or Muſcari.

*Hyacinthus Botroides minor cæruleus obſcurus.*
The darke blew Grape-flower.

This Grape-flower hath many ſmall, fat, and weake leaues lying vpon the ground, which are ſomewhat browniſh at their firſt comming vp, and of a ſad greene afterwards, hollow on the vpperſide, and round vnderneath, among which riſe vp round, ſmooth, weake ſtalkes, bearing at the toppe many ſmall heauie bottle-like flowers, in ſhape like the former Muſcari, but very thicke thruſt together, ſmaller, and of a very darke or blackiſh blew colour, of a very ſtrong ſmell, like vnto Starch when it is new made, and hot: the root is round, and blackiſh without, being compaſſed with a number of ſmall rootes, or of-ſets round about it, ſo that it will quickly choke a ground, if it be ſuffered long in it. For which cauſe, moſt men doe caſt it into ſome by-corner, if they meane to preſerue it, or caſt it out of the Garden quite.

*Alter maior.* There is another of this kinde that is greater, both in leafe and flower, and differeth not in colour or any thing elſe.

*Hyacinthus Botroides cæruleus amœnus.* The skie coloured Grape-flower.

This Iacinth ſpringeth vp with fewer leaues then the firſt, and not reddiſh, but green at his firſt appearing; the leaues, when they are full growne, are long and hollow, like the former, but greener, ſhorter, and broader, ſtanding vpright, and not lying along vpon the ground as they doe: the flowers grow at the toppe of the ſtalke, more ſparſedly ſet thereon, and not ſo thicke together, but like a thinne bunch of grapes, and bottle-like as the former, of a perfect blew or skie-colour, euery flower hauing ſome white ſpots about the brimmes of them: this hath a very ſweet ſmell, nothing like the former: this roote is whiter, and doth not ſo much encreaſe as the former, yet plentifull enough.

*Hyacinthus Botroides ramoſus.* The branched Grape-flower.

Of this kinde, there is another found to grow with many branches of flowers, breaking out from the ſides of the greater ſtalkes or branches: the leaues as all the reſt of the plant is greater then the former.

*Hyacinthus Botroides flore albo.* The white Grape-flower.

The white Grape-flower hath his greene leaues a little whiter, then the blew or skie

skie coloured Grape-flower, his flowers are very pure white, alike sparsedly set on the stalkes, but a little lower and smaller then it, in all other things there is no difference.

*Hyacinthus Botroides flore albo rubente.* The blush Grape-flower.

The roote of this Grape-flower groweth greater, then either the skie coloured, or white Grape-flower, and seldome hath any small rootes or of-sets, as the other haue: his leaues also are larger, and somewhat broader; the flowers are of a pale, or bleake blush colour out of a white, and are a little larger, and grow a little higher and fuller of flowers then the white.

**The Place.**

They naturally grow in many places both of Germany and Hungary; in Spaine likewise, and on Mount Baldus in Italy, and Narbone in France, about the borders of the fields: we haue them in our Gardens for delight.

**The Time.**

These flower from the beginning of March, or sooner sometimes, vntill the beginning of May.

**The Names.**

They are most commonly called *Botroides*, but more truely *Botryodes*, of *Βότρυς* the Greeke word, which signifieth a bunch or cluster of grapes: Lobelius calleth the white one, *Dipcadi flore albo*, transferring the name *Dipcadi*, whereby the *Muscari* is called to this Iacinth, as if they were both one. Their seuerall names, whereby they are knowne and called, are set downe in their titles. The Dutchmen call them *Druekens*, as I said before. Some English Gentlewomen call the white Grape-flower Pearles of Spaine.

1. *Hyacinthus Comosus albus.* The white haired Iacinth.

This Iacinth doth more neerly resemble the Grape-flowers, then the faire haired Iacinths that follow, whereof it beareth the name, in that it hath no haire or threeds at the toppe of the stalke or sides, as they: and therefore I haue placed it next vnto them, and the other to follow it, as being of another kinde. The root hereof is blackish, a little long and round, from whence rise vp three or foure leaues, being smooth and whitish, long, narrow, and hollow, like a trough or gutter on the vpperside: among which the stalke riseth vp a foote high or more, bearing at the toppe diuers small flowers, somewhat like the former, but not so thicke set together, being a little longer, and larger, and wider at the mouth, and as it were diuided into six edges, of a darke whitish colour, with some blacker spots about the brimmes on the inside: the heads or seede-vessels are three square, and somewhat larger, then the heads of any of the former lesser Grape-flowers, wherein is contained round blacke seede.

2. *Hyacinthus Comosus Byzantinus.* The Turkie faire haired Iacinth.

This other Iacinth which came from Constantinople, is somewhat like the former, but that it is bigger, both in roote, and leafe, and flower, and bearing greater store of flowers on the head of the stalke: the lower flowers, although they haue short stalkes at their first flowring, yet afterwards the stalkes grow longer, and those that are lower, stand out further then those that are higheſt, whose foot-stalkes are short, and almost close to the stemme, and of a more perfect purple then any below, which are of a duskie greenish purple colour: the whole stalke of flowers seem like a Pyramis, broad belowe, and small aboue, or as other compare it, to a water sprinkle; yet neither of both these Iacinths haue any threeds at the tops of the stalkes, as the other following haue.

3. *Hyacinthus*

### 3. *Hyacinthus Comosus maior purpureus.*
### The great purple faire haired Iacinth.

This faire haired Iacinth hath his leaues softer, longer, broader, and lesse hollow then the former, lying for the most part vpon the ground: the stalke riseth vp in the midst of the leaues, being stronger, higher, and bearing a greater and longer head of flowers also then they: the flowers of this stand not vpon such long foote-stalkes, but are shorter below, and close almost to the stalke aboue, hauing many bright purplish blew threeds, growing highest aboue the flowers, as it were in a bush together, euery one of these threeds hauing a little head at the end of them, somewhat like vnto one of the flowers, but much smaller: the rest of the flowers below this bush, are of a sadder or deader purple, and not so bright a colour, and the lowest worst of all, rather enclining to a greene, like vnto the last Turkie kinde: the whole stalke with the flowers vpon it, doth somewhat resemble a long Purse tassell, and thereupon diuers Gentlewomen haue so named it: the heads and seede are like vnto the former, but greater: the roote is great and white, with some rednesse on the outside.

### 4. *Hyacinthus Comosus ramosus purpureus.*
### The faire haired branched Iacinth.

The leaues of this Iacinth are broader, shorter, and greener then of the last, not lying so weakly on the ground, but standing somewhat more vpright: the stalke riseth vp as high as the former, but branched out on euery side into many tufts of threeds, with knappes, as it were heads of flowers, at the ends of them, like vnto the head of threeds at the toppe of the former Iacinth, but of a little darker, and not so faire a blewish purple colour: this Iacinth doth somewhat resemble the next Curld haire Iacinth, but that the branches are not so fairely composed altogether of curled threeds, nor of so excellent a faire purple or Doue colour, but more duskie by much: the roote is greater and shorter then of the next, and encreaseth faster.

### 5. *Hyacinthus Pennatus, siue Comosus ramosus elegantior.*
### The faire Curld-haire Iacinth.

This admirable Iacinth riseth vp with three or foure leaues, somewhat like vnto the leaues of the Muske Grape-flower, but lesser; betweene which riseth vp the stalke about a foote high, or somewhat more, bearing at the toppe a bush or tuft of flowers, which at the first appearing, is like vnto a Cone or Pineapple, and afterwards opening it selfe, spreadeth into many branches, yet still retaining the forme of a Pyramis, being broad spread below, and narrow vp aboue: each of these branches is againe diuided into many tufts of threeds or strings, twisted or curled at the ends, and of an excellent purple or Doue colour, both stalkes and haires. This abideth a great while in his beauty, but afterwards all these flowers (if you will so call them) do fall away without any seede at all, spending it selfe as it should seeme in the aboundance of the flowers: the roote is not so great as the last, but white on the outside.

### The Place.

The two first haue been sent diuers times from Constantinople, the third is found wilde in many places of Europe, and as well in Germany, as in Italy. The two last are onely with vs in Gardens, and their naturall places are not knowne vnto vs.

### The Time.

The three former kindes doe flower in Aprill, the two last in May.

### The Names.

The first and second haue no other names then are expressed in their titles.

1 *Hyacinthus Comosus albus.* The white haired Iacinth. 2 *Hyacinthus Comosus Byzantinus.* The Turkie faire haired Iacinth. 3 *Hyacinthus Comosus maior purpureus.* The purple faire haired Iacinth, or Purse tassels. 4 *Hyacinthus Comosus ramosus, sive Calamistratus.* The faire haired branched Iacinth. 5 *Hyacinthus Pannatus, sive Comosus elegantior.* The faire curld haire Iacinth.

tles. The third is called of ſome onely *Hyacinthus maior*, and of others *Hyacinthus comoſus maior*: We call it in Engliſh, The purple faire haired Iacinth, becauſe of his tuft of purple threeds, like haires at the toppe, and (as I ſaid) of diuers Gentlewomen, purple taſſels. The fourth is called by ſome as it is in the title, *Hyacinthus comoſus ramoſus*, and of others *Hyacinthus Calamiſtratus*. And the laſt or fifth is diuerſly called by diuers, Fabius Columna in his *Phytobaſanos* the ſecond part, calleth it *Hyacinthus Sanneſius*, becauſe hee firſt ſaw it in that Cardinals Garden at Rome. Robin of Paris ſent to vs the former of the two laſt, by the name of *Hyacinthus Pennatus*, and *Hyacinthus Calamiſtratus*, when as others ſent the laſt by the name *Pennatus*, and the other by the name of *Calamiſtratus*; but I thinke the name *Cincinnatus* is more fit and proper for it, in that the curled threeds which ſeeme like haires, are better expreſſed by the word *Cincinnus*, then *Calamiſtrum*, this ſignifying but the bodkin or inſtrument wherewith they vſe to friſle or curle the haire, and that the buſh of haire it ſelfe being curled. Some alſo haue giuen to both theſe laſt the names of *Hyacinthus Comoſus Parnaſsi*, the one fairer then the other. Of all theſe names you may vſe which you pleaſe; but for the laſt kinde, the name *Cincinnatus*, as I ſaid, is the more proper, but *Pennatus* is the more common, and *Calamiſtratus* for the former of the two laſt.

### 1. *Hyacinthus Orientalis Brumalis, ſiue præcox flore albo.*
The white Winter Orientall Iacinth.

This early Iacinth riſeth vp with his greene leaues (which are in all reſpects like to the ordinary Orientall Iacinths, but ſomewhat narrower) before Winter, and ſometimes it is in flower alſo before Winter, and is in forme and colour a plaine white Orientall Iacinth, but ſomewhat leſſer, differing onely in no other thing, then the time of his flowring, which is alwayes certaine to be long before the other ſorts.

### 2. *Hyacinthus Orientalis Brumalis, ſiue præcox flore purpureo.*
The purple Winter Orientall Iacinth.

The difference of colour in this flower cauſeth it to bee diſtinguiſhed, for elſe it is of the kindred of the Orientall Iacinths, and is, as the former, more early then the reſt that follow: Vnderſtand then, that this is the ſame with the former, but hauing fine blewiſh purple flowers.

### 3. *Hyacinthus Orientalis maior præcox, dictus Zumbul Indi.*
The greateſt Orientall Iacinth, or Zumbul Indi.

The roote of this Orientall Iacinth, is vſually greater then any other of his kinde, and moſt commonly white on the outſide, from whence riſe vp one or two great round ſtalkes, ſpotted from within the ground, with the lower part of the leaues alſo vpward to the middle of the ſtalkes, or rather higher, like vnto the ſtalkes of Dragons, but darker; being ſet among a number of broad, long, and ſomewhat hollow greene leaues, almoſt as large as the leaues of the white Lilly: at the toppe of the ſtalkes ſtand more ſtore of flowers, then in any other of this kinde, euery flower being as great as the greateſt ſort of Orientall Iacinths, ending in ſix leaues, which turne at the points, of a faire blewiſh purple colour, and all ſtanding many times on one ſide of the ſtalkes, and many times on both ſides.

### 4. *Hyacinthus Orientalis vulgaris diuerſorum colorum.*
The ordinary Orientall Iacinth.

The common Orientall Iacinth (I call it common, becauſe it is now ſo plentifull in all Gardens, that it is almoſt not eſteemed) hath many greene leaues, long, ſomewhat broad and hollow, among which riſeth vp a long greene round ſtalke, beſet from the middle thereof almoſt, with diuers flowers, ſtanding on both ſides

of

of the ſtalkes, one aboue another vnto the toppe, each whereof next vnto the foote-ſtalke is long, hollow, round, and cloſe, ending in ſix ſmall leaues laid open, and a little turning at the points, of a very ſweete ſmell: the colours of theſe flowers are diuers, for ſome are pure white, without any ſhew of other colour in them: another is almoſt white, but hauing a ſhew of blewneſſe, eſpecially at the brims and bottomes of the flowers. Others againe are of a very faint bluſh, tending towards a white: Some are of as deepe a purple as a Violet; others of a purple tending to rednesse, and ſome of a paler purple. Some againe are of a faire blew, others more watchet, and ſome ſo pale a blew, as if it were more white then blew: after the flowers are paſt, there riſe vp great three ſquare heads, bearing round blacke ſeede, great and ſhining: the roote is great, and white on the outſide, and oftentimes purpliſh alſo, flat at the bottome, and ſmall at the head.

There is a kinde of theſe Iacinths, whoſe flowers are of a deepe purpliſh Violet colour, hauing whitiſh lines downe the backe of euery leafe of the flower, which turne themſelues a little backwards at the points. *Flore purpureo violaceo lineis albicantis in dorſo.*

There is another, whoſe flowers ſtand all opening one way, and not on all ſides, but are herein like the great Zumbul Indi, before ſet out. *Floribus antrorſum reſpicientibus.*

There is againe another kinde which flowreth later then all the reſt, and the flowers are ſmaller, ſtanding more vpright, which are either white or blew, or mixt with white and purple. *Serotinus erectis floribus diuerſorum colorum.*

5. *Hyacinthus Orientalis folioſo caule.* The buſhy ſtalked Orientall Iacinth.

This ſtrange Iacinth hath his rootes, leaues, and flowers, like vnto the former Orientall Iacinths: the onely difference in this is, that his ſtalke is not bare or naked, but hath very narrow long leaues, growing diſperſedly, and without order, with the flowers thereon, which are blew, and hauing for the moſt part one leafe, and ſometimes two at the foote, or ſetting on of euery flower, yet ſometimes it happeneth, ſome flowers to be without any leafe at the bottome, as nature, that is very variable in this plant, liſteth to play: the heads and ſeede are blacke and round, like the other alſo.

6. *Hyacinthus Orientalis flore duplici.* The bleake Orientall Iacinth once double.

This double Iacinth hath diuers long leaues, like vnto the other Orientall Iacinths, almoſt ſtanding vpright, among which riſeth vp a ſtalke, browniſh at the firſt, but growing greene afterwards, bearing many flowers at the toppe, made like the flowers of the former Iacinths, and ending in ſix leaues, greene at the firſt, and of a blewiſh white when they are open, yet retaining ſome ſhew of greenneſſe in them, the brims of the leaues being white; from the middle of each flower ſtandeth forth another ſmall flower, conſiſting of three leaues, of the ſame colour with the other flower, but with a greene line on the backe of each of theſe inner leaues: in the middle of this little flower, there ſtand ſome threeds tipt with blacke: the ſmell of this flower is not ſo ſweete as of the forme; the heads, ſeede, and rootes are like the former.

7. *Hyacinthus Orientalis flore pleno caruleo, vel purpuro violaceo.* The faire double blew, or purple Orientall Iacinth.

The leaues of theſe Iacinths are ſmaller, then the leaues of moſt of the other former ſorts; the ſtalkes are ſhorter, and ſmaller, bearing but three or foure flowers on the heads of them for the moſt part, which are not compoſed like the laſt, but are more faire, full, and double of leaues, where they ſhew out their full beauties, and of a faire blew colour in ſome, and purple in others, ſmelling pretty ſweete; but theſe doe ſeldome beare out their flowers faire; and beſides, haue diuers other flowers that will be either ſingle, or very little double vpon the ſame ſtalke.

8. *Hyacinthus Orientalis candidiſsimus flore pleno.* The pure white double Orientall Iacinth.

This double white Iacinth hath his leaues like vnto the ſingle white Orientall Iacinth;

cinth; his stalke is likewise long, slender, and greene, bearing at the toppe two or three flowers at the most, very double and full of leaues, of a pure white colour, without any other mixture therein, hanging downe their heads a little, and are reasonable sweete. I haue this but by relation, not by sight, and therefore I can giue no further assurance as yet.

The Place.

All these Orientall Iacinths, except the last, haue beene brought out of Turkie, and from Constantinople: but where their true originall place is, is not as yet vnderstood.

The Time.

The two first (as is said) flower the earliest, sometimes before Christmas, but more vsually after, and abide a great while in flower, in great beauty, especially if the weather be milde, when as few or no other flowers at that time are able to match them. The other greatest kinde flowreth also earlier then the rest that follow, for the most part. The ordinary kindes flower some in March, and some in Aprill, and some sooner also; and so doe the double ones likewise. The bushy stalked Iacinth flowreth much about the same time.

The Names.

The former two sorts are called *Hyacinthus Orientalis Brumalis*, and *Hyacinthus Orientalis præcox flore albo*, or *cæruleo*. The third is called of many *Zumbul Indicum*, or *Zumbul Indi*, and corruptly *Simboline*; of others, and that more properly, *Hyacinthus Orientalis maior præcox*. The Turkes doe call all Iacinths *Zumbul*, and by adding the name of *Indi*, or *Arabi*, do shew from what place they are receiued. In English, The greatest Orientall Iacinth; yet some doe call it after the Turkish name *Zumbul Indi*, or *Simboline*, as is said before. The rest haue their names set downe in their titles, which are most fit for them.

*Hyacinthus Hispanicus minor Orientalis facie.*

The little Summer Orientall Iacinth.

This little Iacinth hath foure or fiue long narrow greene leaues, lying vpon the ground, among which riseth vp a slender smooth stalke, about a spanne high or more,
*Flore cæruleo.* bearing at the toppe many slender bleake blew flowers, with some white stripes and edges to be seene in most of them, fashioned very like vnto the flowers of the Orientall Iacinth, but much smaller: the flower hath no sent at all; the seede is like the seede of the English Iacinth, or Harefbels: the roote is small and white.

*Flore albo.* There is another of this kinde, differing in nothing but in the colour of the flower, which is pure white.

*Flore rubente.* There is also another, whose flowers are of a fine delayed red colour, with some deeper coloured veines, running along the three outer leaues of the flower, differing in no other thing from the former.

The Place.

These plants haue been gathered on the Pyrenæan Mountaines, which are next vnto Spaine, from whence, as is often said, many rare plants haue likewise been gathered.

The Time.

They flower very late, euen after all or most of the Iacinths, in May for the most part.

The




1 *Hyacinthus Orientalis brumalis.* The Winter Orientall Iacinth. 2 *Zumbul Indi.* The greateſt Orientall Iacinth. 3 *Hyacinthus Orientalis vulgaris.* The ordinary Orientall Iacinth. 4 *Hyacinthus Orientalis foliosa caule.* The buſhy ſtalked Orientall Iacinth. 5 *Hyacinthus Orientalis flore duplici.* The Orientall Iacinth once double. 6 *Hyacinthus Orientalis flore pleno cæruleo.* The faire double blew Orientall Iacinth.

The Names.

They are called eyther *Hyacinthus Hiſpanicus minor Orientalis facie*, as it is in the title, or *Hyacinthus Orientalis facie*, that is to ſay, The leſſer Spaniſh Iacinth, like vnto the Orientall: yet ſome haue called them, *Hyacinthus Orientalis ſerotinus minor*, The leſſer late Orientall Iacinth, that thereby they may be knowne from the reſt.

*Hyacinthus Hiſpanicus obſoletus.* The Spaniſh dunne coloured Iacinth.

This Spaniſh Iacinth ſpringeth very late out of the ground, bearing foure or fiue ſhort, hollow, aud ſoft whitiſh greene leaues, with a white line in the middle of euery one of them, among which riſe vp one or more ſtalkes, bearing diuers flowers at the toppes of them, all looking one way, or ſtanding on the one ſide, hanging downe their heads, conſiſting of ſix leaues, three whereof being the outermoſt, lay open their leaues, and turne back the ends a little again: the other three which are innermoſt, do as it were cloſe together in the middle of the flower, without laying themſelues open at all, being a little whitiſh at the edges: the whole flower is of a purpliſh yellow colour, with ſome white and green as it were mixed among it, of no ſent at all: it beareth blacke and flat ſeede in three ſquare, great, and bunched out heads: the roote is reaſonable great, and white on the outſide, with many ſtrong white fibres at it, which periſh not yearely, as the fibres of many other Iacinths doe, and as it ſpringeth late, ſo it holdeth his greene leaues almoſt vntill Winter.

*Mauritanicus.* There hath been another hereof brought from about Fez and Marocco in Barbary, which in all reſpects was greater, but elſe differed little.

*Maximus Æthiopicus.* There was another alſo brought from the Cape of good Hope, whoſe leaues were ſtronger and greener then the former, the ſtalke alſo thicker, bearing diuers flowers, confuſedly ſtanding vpon longer foote-ſtalkes, yet made after the ſame faſhion, but that the three inner leaues were whitiſh, and dented about the edges, otherwiſe the flowers were yellow and greeniſh on the inſide.

The Place.

Theſe plants grow in Spaine, Barbary, and Ethiopia, according as their names and deſcriptions doe declare.

The Time.

The firſt flowreth not vntill Iune; for, as I ſaid, it is very late before it ſpringeth vp out of the ground, and holdeth his leaues as is ſaid, vntill September, in the meane time the ſeede thereof ripeneth.

The Names.

They haue their names according to the place of their growing; for one is called *Hyacinthus Hiſpanicus obſoletioris coloris*. The other is called alſo *Hyacinthus Mauritanicus*. And the laſt, *Hyacinthus Æthiopicus obſoletus*. In Engliſh, The Spaniſh, Barbary, or Ethiopian Iacinth, of a dunne or duskie colour.

*Hyacinthus Anglicus Belgicus, vel Hiſpanicus.*
Engliſh Hareſ-bels, or Spaniſh Iacinth.

Our Engliſh Iacinth or Hareſ-bels is ſo common euery where, that it ſcarce needeth any deſcription. It beareth diuers long and narrow greene leaues, not ſtanding vpright, nor yet fully lying vpon the ground, among which ſpringeth vp the ſtalke, bearing at the toppe many long and hollow flowers, hanging downe their heads all forwards

forwards for the most part, parted at the brimmes into six parts, turning vp their points a little againe, of a sweetish, but heady sent; somewhat like vnto the Grape-flower: the heads for seede are long and square, wherein is much blacke seede: the colour of the flowers are in some of a deeper blew, tending to a purple; in others of a paler blew, or of a bleake blew, tending to an ash colour: Some are pure white, and some are party coloured, blew and white; and some are of a fine delayed purplish red or blush colour, which some call a peach colour. The rootes of all sorts agree, and are alike, being white and very slimie; some whereof will be great and round, others long and slender, and those that lye neare the toppe of the earth bare, will be greene.

*Hyacinthus Hispanicus maior flore campanulæ instar.*
The greater Spanish bell-flowred Iacinth.

This Spanish bell-flowred Iacinth, is very like the former English or Spanish Iacinth, but greater in all parts, as well of leaues as flowers, many growing together at the toppe of the stalke, with many short greene leaues among them, hanging downe their heads, with larger, greater, and wider open mouths, like vnto bels, of a darke blew colour, and no good sent.

The Place.

The first groweth in many places of England, the Lowe-Countries, as we call them, and Spaine, but the last chiefly in Spaine.

The Time.

They flower in Aprill for the most part, and sometimes in May.

The Names.

Because the first is more frequent in England, then in Spain, or the Lowe-Countries, it is called with vs *Hyacinthus Anglicus*, The English Iacinth; but it is also called as well *Belgicus*, as *Hispanicus*: yet Dodonæus calleth it *Hyacinthus non scriptus*, because it was not written of by any Authour before himselfe. It is generally knowne in England by the name of Harebels. The other Spanish Iacinth beareth his name in his title.

*Hyacinthus Eriophorus.* The Woolly Iacinth.

This Woolly Iacinth hath many broad, long, and faire greene leaues, very like vnto some of the Iacinths, but stiffer, or standing more vpright, which being broken, doe yeeld many threeds, as if a little fine cotton wooll were drawne out: among these leaues riseth vp a long greene round stalke, a foote and a halfe high or more, whereon is set a great long bush of flowers, which blowing open by degrees, first below, and so vpwards, are very long in flowring: the toppe of the stalke, with the flowers, and their little footstalkes, are all blew, euery flower standing outright with his stalke, and spreading like a starre, diuided into six leaues, hauing many small blew threeds, standing about the middle head, which neuer gaue ripe seede, as farre as I can heare of: the root is white, somewhat like the root of a Muscari, but as full of wooll or threeds, or rather more, then the leaues, or any other part of it.

The Place.

This hath been sent diuers times out of Turkie into England, where it continued a long time as well in my Garden as in others, but some hard frosty Winters caused it to perish with me, and diuers others, yet I haue had it againe from a friend, and doth abide fresh and greene euery yeare in my Garden.

### The Time.

This flowred in the Garden of Mr. Richard Barnesley at Lambeth, onely once in the moneth of May, in the yeare 1606. after hee had there preserued it a long time: but neither he, nor any elfe in England that I know, but those that saw it at that time, euer saw it beare flower, either before or since.

### The Names.

It is called by diuers *Bulbus Eriophorus*, or *Laniferus*, that is, Woolly Bulbous; but because it is a Iacinth, both in roote, leafe, and flower, and not a *Narcissus*, or Daffodill, it is called *Hyacinthus Eriophorus*, or *Laniferus*, The Woolly Iacinth. It is very likely, that Theophrastus in his seuenth Book & thirteenth Chapter, did meane this plant, where hee declareth, that garments were made of the woolly substance of a bulbous roote, that was taken from between the core or heart of the roote (which, as hee saith, was vsed to be eaten) and the outermost shels or peelings; yet Clusius seemeth to fasten this woolly bulbous of Theophrastus, vpon the next Iacinth of Spaine.

*Hyacinthus Stellatus Baticus maior, vulgò Perüanus.*

The great Spanish Starry Iacinth, or of Peru.

This Iacinth (the greatest of those, whose flowers are spread like a starre, except the two first Indians) hath fiue or six, or more, very broad, and long greene leaues, spread vpon the ground, round about the roote, which being broken are woolly, or full of threeds, like the former: in the middle of these leaues riseth vp a round short stalke, in comparison of the greatnesse of the plant (for the stalke of the Orientall Iacinth is sometimes twice so high, whose roote is not so great) bearing at the toppe a great head or bush of flowers, fashioned in the beginning, before they bee blowne or separated, very like to a Cone or Pineapple, and begin to flower belowe, and so vpwards by degrees, euery flower standing vpon a long blackish blew foote-stalke, which when they are blowne open, are of a perfect blew colour, tending to a Violet, and made of six small leaues, laid open like a starre; the threeds likewise are blewish, tipt with yellow pendents, standing about the middle head, which is of a deeper blew, not hauing any good sent to be perceiued in it, but commendable only for the beauty of the flowers: after the flowers are past, there come three square heads, containing round blacke seede: the roote is great, and somewhat yellowish on the outside, with a knobbe or bunch at the lower end of the roote, (which is called the seate of the roote) like vnto the Muscari, Scylla, and many other bulbous rootes, at which hang diuers white, thicke, and long fibres, whereby it is fastened in the ground, which perish not euery yeare, but abide continually, and therefore doth not desire much remouing.

*Hyacinthus Stellatus Baticus, sive Perüanus flore albo.*

The great white Spanish starry Iacinth.

This other Spanish Iacinth is in most parts like vnto the former, but that his leaues are not so large, nor so deep a greene: the stalks of flowers likewise hath not so thicke a head, or bush on it, but fewer and thinner set: the flowers themselues also are whitish, yet hauing a small dash of blush in them: the threeds are whitish, tipt with yellow pendents: the seede and rootes are like vnto the former, and herein consisteth the difference betweene this and the other sorts.

*Hyacinthus Stellatus Baticus, sive Perüanus flore carneo.*

The great blush coloured Spanish Starry Iacinth.

This likewise differeth little from the two former, but onely in the colour of the flowers;

1 *Hyacinthus Orientalis facie.* The little Summer Orientall Iacinth. 2 *Hyacinthus Mauritanicus.* The Barbary Iacinth. 3 *Hyacinthus obsoletus Hispanicus.* The Spanish duskie Iacinth. 4 *Hyacinthus Hispanicus flore campanula.* The greater Spanish bel-flowred Iacinth. 5 *Hyacinthus Anglicus.* The English Iacinth or Harebels. 6 *Hyacinthus Eriophorus.* The Woolly Iacinth. 7 *Hyacinthus Stellaris Baticus maior, sive Peruanus.* The great Spanish Starry Iacinth, or of Peru.

flowers; for this being found growing among both the other, hath his head of flowers as great and large as the first, but the buds of his flowers, before they are open, are of a deepe blush colour, which being open, are more delayed, and of a pleasant pale purple, or blush colour, standing vpon purplish stalkes: the heads in the middle are whitish, and so are the threeds compassing it, tipt with yellow.

The Place.

These doe naturally grow in Spaine, in the Medowes a little off from the Sea, as well in the Island Gades, vsually called Cales, as likewise in other parts along the Sea side, as one goeth from thence to Porto Santa Maria, which when they be in flower, growing so thicke together, seeme to couer the ground, like vnto a tapistry of diuers colours, as I haue beene credibly enformed by Guillaume Boel, a Freeze-lander borne, often before and hereafter remembred, who being in search of rare plants in Spaine, in the yeare of our Lord 1607. after that most violent frosty Winter, which perished both the rootes of this, and many other fine plants with vs, sent mee ouer some of these rootes for my Garden, and affirmed this for a truth, which is here formerly set downe, and that himselfe gathered those he sent mee, and many others in the places named, with his owne hands; but hee saith, that both that with the white, and with the blush flowers, are farre more rare then the other.

The Time.

They flower in May, the seede is ripe in Iuly.

The Names.

This hath beene formerly named *Eriophorus Peruanus*, and *Hyacinthus Stellatus Peruanus*, The Starry Iacinth of Peru, being thought to haue grown in Peru, a Prouince of the West Indies; but he that gaue that name first vnto it, eyther knew not his naturall place, or willingly imposed that name, to conceale it, or to make it the better esteemed. It is most generally receiued by the name *Hyacinthus Peruanus*, from the first imposer thereof, that is, the Iacinth of Peru: but I had rather giue the name agreeing most fitly vnto it, and call it as it is indeede *Hyacinthus Stellatus Baticus*, The Spanish Starry Iacinth; and becausse it is the greatest that I know hath come from thence, I call it, The great Starry Iacinth of Spaine, or Spanish Iacinth.

*Hyacinthus Stellatus vulgaris, siue Bifolius Fuchsij.*
The common blew Starry Iacinth.

This Starry Iacinth (being longest knowne, and therefore most common) riseth out of the ground, vsually but with two browne leaues, yet sometimes with three, inclosing within them the stalke of flowers, the buds appearing of a darke whitish colour, as soone as the leaues open themselues, which leaues being growne, are long, and hollow, of a whitish greene on the vpper side, and browne on the vnder side, and halfe round, the browne stalke rising vp higher, beareth fiue or sixe small starre-like flowers thereon, consisting of six leaues, of a faire deepe blew, tending to a purple. The seede is yellowish, and round, contained in round pointed heads, which by reason of their heauinesse, and the weaknesse of the stalke, lye vpon the ground, and often perish with wet and frosts, &c. The roote is somewhat long, and couered with a yellowish coate.

*Hyacinthus stellatus flore albo.* The white Starry Iacinth.

The white Starry Iacinth hath his leaues like the former, but greene and fresh, not browne, and a little narrower also: the buddes for flowers at the first appeare a little blush, which when they are blowne, are white, but yet retaine in them a small shew of that blush colour.

We

We haue another, whose flowers are pure white, and smaller then the other, the leaues whereof are of a pale fresh greene, and somewhat narrower. *Flore niueo.*

*Hyacinthus Stellatus flore rubente.* The blush coloured Starry Iacinth.

The difference in this from the former, is onely in the flowers, which are of a faire blush colour, much more eminent then in the others, in all things else alike.

*Hyacinthus Stellatus Martius, sive præcox cæruleus.* The early blew Starry Iacinth.

This Iacinth hath his leaues a little broader, of a fresher greene, and not browne at all, as the first blew Iacinth of Fuchsius last remembred: the buds of the flowers, while they are enclosed within the leaues, and after, when the stalke is gowne vp, doe remaine more blew then the buds of the former: the flowers, when they are blowne open, are like the former, but somewhat larger, and of a more liuely blew colour: the roote also is a little whiter on the outside. This doth more seldome beare seede then the former.

*Hyacinthus Stellatus præcox flore albo.* The white early Starry Iacinth.

There is also one other of this kinde, that beareth pure white flowers, the green leafe thereof being a little narrower then the former, and no other difference.

*Hyacinthus Stellatus præcox flore suaue rubente.* The early blush coloured Starry Iacinth.

This blush coloured Iacinth is very rare, but very pleasant, his flowers being as large as the first of this last kinde, and somewhat larger then the blush of the other kinde: the leaues and rootes differ not from the last recited Iacinth.

## The Place.

All these Iacinths haue beene found in the Woods and Mountaines of Germany, Bohemia, and Austria, as Fuchsius and Gesner doe report, and in Naples, as Imperatus and others doe testifie. Wee cherish them all with great care in our Gardens, but especially the white and the blush of both kindes, for that they are more tender, and often perish for want of due regard.

## The Time.

The common kindes, which are first expressed, flower about the middle of February, if the weather bee milde, and the other kindes sometimes a fortnight after, that is, in March, but ordinarily much about the same time with the former.

## The Names.

The first is called in Latine *Hyacinthus Stellatus vulgaris*, and *Hyacinthus Stellatus bifolius*, and *Hyacinthus Stellaris Fuchsij*, and of some *Hyacinthus Stellatus Germanicus*; wee might very well call the other kinde, *Hyacinthus Stellatus vulgaris alter*, but diuers call it *Præcox*, and some *Martius*, as it is in the title. In English they may bee seuerally called: the first, The common; and the other, The early Starry Iacinth (notwithstanding the first flowreth before the other) for distinction sake.

The *Hyacinthus* seemeth to be called *Vacinium* of Virgil in his Eclogues; for hee alwayes reckoneth it among the flowers that were vsed to decke Garlands, and neuer among fruits, as some would haue it. But in that hee calleth it *Vacinium nigrum*, in seuerall places, that doth very fitly answer the common

common receiued custome of those times, that called all deepe blew colours, such as are purples, and the like, blacke; for the Violet it selfe is likewise called blacke in the same place, where he calleth the *Vacinium* blacke; so that it seemeth thereby, that he reckoned them to be both of one colour, and we know the colour of the Violet is not blacke, as we doe distinguish of blacke in these dayes. But the colour of this Starry Iacinth, being both of so deepe a purple sometimes, so neare vnto a Violet colour, and also more frequent, then any other Iacinth with them, in those places where Virgil liued, perswadeth me to thinke, that Virgil vnderstood this Starry Iacinth by *Vacinium*: Let others iudge otherwise, if they can shew greater probabilitie.

### 1. *Hyacinthus Stellatus Byzantinus nigra radice.*
### The Starry Iacinth of Turkie with the blacke roote.

This Starry Iacinth of Constantinople hath three or foure fresh greene, thinne, and long leaues, of the bignesse of the English Iacinth, but not so long, betweene which riseth vp a slender lowe stalke, bearing fiue or six small flowers, dispersedly set thereon, spreading open like a starre, of a pale or bleake blew colour: the leaues of the flowers are somewhat long, and stand as it were somewhat loosly, one off from another, and not so compactly together, as the flowers of other kindes: it seldome beareth ripe seede with vs, because the heads are so heauie, that lying vpon the ground, they rotte with the wet, or are bitten with the frosts, or both, so that they seldome come to good: the roote is small in some, and reasonable bigge in others, round, and long, white within, but couered with deepe reddish or purplish peelings, next vnto it, and darker and blacker purple on the outside, with some long and thicke white fibres, like fingers hanging at the bottome of them, as is to be seene in many other Iacinths: the roote it selfe for the most part doth runne downewards, somewhat deep into the ground.

### 2. *Hyacinthus Stellatus Byzantinus maior.*
### The greater Starry Iacinth of Constantinople.

This Iacinth may rightly be referred to the former Iacinth of Constantinople, and called the greater, it is so like thereunto, that any one that knoweth that, will soone say, that this is another of that sort, but greater as it is in all his parts, bearing larger leaues by much, and more store, lying vpon the ground round about the roote: it beareth many lowe stalkes of flowers, as bleake, and standing as loosly as the former: onely the roote of this, is not black on the outside, as the other, but three times bigger.

### 3. *Hyacinthus Stellatus Byzantinus alter, sine flore boraginis.*
### The other Starry Iacinth of Constantinople.

This other Iacinth hath for the most part onely foure leaues, broader and greener then the first, but not so large or long as the second: the stalke hath fiue or six flowers vpon it, bigger and rounder set, like other starry Iacinths, of a more perfect or deeper blew then either of the former, hauing a whitish greene head or vmbone in the middle, beset with six blew chiues or threeds, tipt with blacke, so closly compassing the vmbone, that the threeds seeme so many prickes stucke into a clubbe or head; some therefore haue likened it to the flower of Borage, and so haue called it: after the flowers are past, come vp round white heads, wherein is contained round and white seede: the roote is of a darke whitish colour on the outside, and sometimes a little reddish withall.

### The Place.

The first and the last haue beene brought from Constantinople; the first among many other rootes, and the last by the Lord Zouch, as Lobel witnesseth. The second hath been sent vs out of the Lowe-Countries, but from whence they had it, we do not certainly know. They growe with vs in our Gardens sufficiently.

The

The Time.

These flower in Aprill, but the first is the earliest of the rest, and is in flower presently after the early Starry Iacinth, before described.

The Names.

The former haue their names in their titles, and are not knowne vnto vs by any other names that I know; but as I said before, the last is called by some, *Hyacinthus Boraginis flore.* The first was sent out of Turkie, by the name of *Susam giul*, by which name likewise diuers other things haue beene sent, so barren and barbarous is the Turkish tongue.

*Hyacinthus Stellatus Æstivus maior.* The greater Summer Starry Iacinth.

This late Iacinth hath diuers narrow greene leaues, lying vpon the ground, somewhat like the leaues of the English Iacinth, but stiffer and stronger; among which riseth vp a round stiffe stalke, bearing many flowers at the toppe thereof, and at euery foote-stalke of the flowers a small short leafe, of a purplish colour: the flowers are starre-like, of a fine delayed purplish colour, tending to a pale blew or ash colour, striped on the backe of euery leafe, and hauing a pointed vmbone in the middle, with some whitish purple threeds about it, tipt with blew: the seede is blacke, round, and shining, like vnto the seede of the English Iacinth, but not so bigge: the roote is round and white, hauing some long thicke rootes vnder it, besides the fibres, as is vsuall in many other Iacinths.

*Hyacinthus Stellatus Æstivus minor.* The lesser Summer Starry Iacinth.

This lesser Iacinth hath diuers very long, narrow, and shining greene leaues, spread vpon the ground round about the roote, among which riseth vp a very short round stalke, not aboue two inches high, carrying six or seuen small flowers thereon, on each side of the stalke, like both in forme and colour vnto the greater before described, but lesser by farre: the seede is blacke, contained in three square heads: the roote is small and white, couered with a browne coate, and hauing some such thicke rootes among the fibres, as are among the other.

The Place.

Both these Iacinths grow naturally in Portugall, and from thence haue been brought, by such as seeke out for rare plants, to make a gaine and profit by them.

The Time.

They both flower in May, and not before: and their seed is ripe in Iuly.

The Names.

Some doe call these *Hyacinthus Lusitanicus*, The Portugall Iacinth. Clusius, who first set out the descriptions of them, called them as is expressed in their titles; and therefore we haue after the Latine name giuen their English, according as is set downe. Or if you please, you may call them, The greater and the lesser Portugall Iacinth.

*Hyacinthus Stellaris flore cinereo.* The ash coloured Starry Iacinth.

This ash coloured Iacinth, hath his leaues very like vnto the leaues of the English Iacinth, and spreading vpon the ground in the same manner, among which rise vp one or two stalkes, set at the toppe with a number of small starre-like flowers, bushing bigger

ger below then aboue, of a very pale or white blew, tending to an aſh colour, and very ſweete in ſmell: the ſeede is blacke and round, like vnto the ſeede of the Engliſh Iacinth, and ſo is the roote, being great, round, and white; ſo like, I ſay, that it is hard to know the one from the other.

The Place.

The certaine originall place of growing thereof, is not knowne to vs.

The Time.

It flowreth in Aprill.

The Names.

Some doe call this *Hyacinthus Someri*, Somers Iacinth, becauſe as Lobel ſaith, he brought it firſt into the Lowe-Countries, eyther from Conſtantinople, or out of Italy.

*Hyacinthus Stellatus Lilifolio & radice cæruleo.*
The blew Lilly leafed Starre Iacinth.

This Iacinth hath ſix or ſeuen broad greene leaues, ſomewhat like vnto Lilly leaues, but ſhorter (whereof it tooke his name as well as from the roote) ſpread vpon the ground, and lying cloſe and round: before the ſtalke riſeth out from the middle of theſe leaues, there doth appeare a deepe hollow place, like a hole, to bee ſeene a good while, which at length is filled vp with the ſtalke, riſing thence vnto a foote or more high, bearing many ſtarre-like flowers at the toppe, of a perfect blew colour, neare vnto a Violet, and ſometimes of paler or bleake blew colour, hauing as it were a ſmall cuppe in the middle, diuided into ſix peeces, without any threeds therein: the ſeede is blacke and round, but not ſhining: the roote is ſomewhat long, bigge belowe, and ſmall aboue, like vnto the ſmall roote of a Lilly, and compoſed of yellow ſcales, as a Lilly, but the ſcales are greater, and fewer in number.

*Hyacinthus Stellatus Lilifolius albus.* The white Lilly leafed Starre Iacinth.

The likeneſſe of this Iacinth with the former, cauſeth me to be briefe, and not to repeate the ſame things againe, that haue already been expreſſed: You may therefore vnderſtand, that except in the colour of the flower, which in this is white, there is no difference betweene them.

*Flore carneo.* I heare of one that ſhould beare bluſh coloured flowers, but I haue not yet ſeene any ſuch.

The Place.

Theſe Iacinths haue been gathered on the Pyrenæan Hils, in that part of France that is called Aquitaine, and in ſome other places.

The Time.

Theſe flower in Aprill, and ſometimes later.

The Names.

Becauſe the roote is ſo like vnto a Lilly, as the leafe is alſo, it hath moſt properly beene called *Hyacinthus Stellatus Lilifolio & radice*, or for breuity *Lilifolius*, that is, The Starry Lilly leafed Iacinth. It is called *Sarabug* by the Inhabitants where it groweth, as Cluſius maketh the report from Venerius, who further ſaith, that by experience they haue found the cattell to ſwell and dye, that haue eaten of the leaues thereof.

*Hyacinthus*

1 *Hyacinthus stellatus præcox cæruleus.* The early blew starry Iacinth. 2 *Hyacinthus stellatus præcox albus.* The early white starry Iacinth. 3 *Hyacinthus stellatus Byzantinus nigra radice.* The Turkie starry Iacinth with a blacke roote. 4 *Hyacinthus Byzantinus alter siue flore Boraginis.* The other starry Iacinth of Constantinople. 5 *Hyacinthus æstiuus maior.* The greater Summer starry Iacinth. 6 *Hyacinthus stellatus floredimetco.* The ash coloured starry Iacinth. 7 *Hyacinthus stellatus Liliofolius.* The Lilly leafed starre Iacinth. 8 *Hyacinthus Autumnalis.* The Autumne Iacinth. 9 *Scilla alba siue Hyacinthus marinus.* The Sea Onion or Squill.

*Hyacinthus Autumnalis maior.* The greater Autumne Iacinth.

The greater Autumne Iacinth hath fiue or six very long and narrow greene leaues, lying vpon the ground; the stalkes are set at the toppe with many starre-like flowers, of a pale blewish purple colour, with some pale coloured threeds, tipt with blew, standing about the head in the middle, which in time growing ripe, containeth therein small blacke seede, and roundish: the roote is great and white on the outside.

*Hyacinthus Autumnalis minor.* The lesser Autumne Iacinth.

This lesser Iacinth hath such like long and small leaues, but narrower then the former: the stalke is not full so high, but beareth as many flowers on it as the other, which are of a pale or bleake purple colour, very like vnto it also: the roote and seed are like the former, but smaller. These both for the most part, beare their flowers and seede before the greene leaues rise vp much aboue the ground.

*Flore albo.* There is a kinde hereof found that beareth white flowers, not differing in any other thing from the smaller purple kinde last mentioned.

The Place.

The first and last are onely kept in Gardens, and not knowne to vs where their naturall place of growing wilde may be.

The second groweth wilde in many places of England. I gathered diuers rootes for my Garden, from the foote of a high banke by the Thames side, at the hither end of Chelsey, before you come at the Kings Barge-house.

The Time.

The greatest flowreth in the end of Iuly, and in August.

The other in August and September, you shall seldome see this plant with flowers and greene leaues at one time together.

The Names.

They haue their names giuen them, as they are expressed in their titles, by all former Writers, except Daleschampius, or hee that set forth that great worke printed at Lyons; for hee contendeth with many words, that these plants can bee no Iacinths, because their flowers appeare before their leaues in Autumne, contrary to the true Iacinth, as he saith: and therefore he would faine haue it referred to *Theophrastus bulbus in libro primo cap.* 12. and calleth it his *Tiphyum* mentioned in that place, as also *Bulbus æstiuus Daleschampij.* Howsoeuer these things may carry some probability in them, yet the likenesse both of rootes, and flowers especially, hath caused very learned Writers to entitle them as is set downe, and therefore I may not but let them passe in the like manner.

The Vertues.

Both the rootes and the leaues of the Iacinths are somewhat cold and drying, but the seede much more. It stayeth the loosnesse of the belly. It is likewise said to hinder young persons from growing ripe too soone, the roote being drunke in wine. It helpeth them also whose vrine is stopt, and is auaileable for the yellow Iaundise; but as you heare some are deadly to cattell, I therefore wish all to bee well aduised which of these they will vse in any inward physicke.

*Scilla alba.* The Sea Onion or Squill.

As I ended the discourse of both the true and the bastard Daffodils, with the Sea kindes

kindes of both sorts; so I thinke it not amisse, to finish this of the Iacinths with the description of a Sea Iacinth, which (as you see) I take to be the *Scilla*, or Sea Onion, all his parts so nearely resembling a Iacinth, that I know not where to ranke him better then in this place, or rather not any where but here. You shall haue the description thereof, and then let the iudicious passe their sentence, as they thinke meetest.

The Squill or Sea Onion (as many doe call it) hath diuers thicke leaues, broad, long, greene, and hollowish in the middle, and with an eminent or swelling ribbe all along the backe of the leafe, (I relate it as I haue seene it, hauing shot forth his leaues in the ship by the way, as the Mariners that brought diuers rootes from out of the Straights, did sell them to mee and others for our vse) lying vpon the ground, somewhat like vnto the leaues of a Lilly: these spring vp after the flowers are past, and the seed ripe, they abiding all the Winter, and the next Spring, vntill the heate of the Summer hath spent and consumed them, and then about the end of August, or beginning of September, the stalke with flowers ariseth out of the ground a foote and a halfe high, bearing many starre-like flowers on the toppe, in a long spike one aboue another, flowring by degrees, the lowest first, and so vpwards, whereby it is long in flowring: very like, as well in forme as bignesse, to the flowers of the great Starre of Bethlehem (these flowers I haue likewise seene shooting out of some of the rootes, that haue been brought in the like manner:) after the flowers are past, there come vp in their places thicke and three square heads, wherin is contained such like flat, black, and round seed, as the Spanish duskie Iacinth before described did beare, but greater: the root is great & white, couered with many peelings or couerings, as is plainly enough seen to any that know them, and that sometimes wee haue had rootes, that haue beene as bigge as a pretty childes head, and sometimes two growing together, each whereof was no lesse then is said of the other.

*Scilla rubra sive Pancratium verum.* The red Sea Onion.

The roote of this Squill, is greater oftentimes then of the former, the outer coates or peelings being reddish, bearing greater, longer, stiffer, and more hollow leaues, in a manner vpright: this bringeth such a like stalke and flowers, as the former doth, as Fabianus Ilges, Apothecary to the Duke of Briga, did signifie by the figure thereof drawne and sent to Clusius.

The Place.

They grow alwayes neare the Sea, and neuer farre off from it, but often on the very baich of the Sea, where it washeth ouer them all along the coasts of Spaine, Portugal, and Italy, and within the Straights in many places: it will not abide in any Garden farre from the Sea, no not in Italy, as it is related.

The Time.

The time wherein they flower, is expressed to be in August and September: the seede to be ripe in October and Nouember, and the greene leaues to spring vp in Nouember and December.

The Names.

These are certainly the true kindes of *Scilla* that should bee vsed in medicines, although (as Clusius reporteth) the Spaniards forbade him to taste of the red Squill, as of a most strong and present poison. Pliny hath made more sorts then can be found out yet to this day with vs: that *Scilla* that is called *Epimenidia*, because it might be eaten, is thought to be the great *Ornithogalum*, or Starre of Bethlehem. *Pancratium* is, I know, and as I said before, referred to that kinde of bastard Sea Daffodill, which is set forth before in the end of the history of the bastard Daffodils; and diuers also would make the *Narcissus tertius Matthioli*, which I call the true Sea Daffodill, to be a *Pancratium*, but seeing Dioscorides (and no other is against him)

maketh *Pancratium* to be a kinde of Squill with reddish rootes, I dare not vphold their opinion against such manifest truth.

### The Vertues.

The Squill or Sea Onion is wholly vsed physically with vs, because wee can receiue no pleasure from the sight of the flowers. Pliny writeth, that Pithagoras wrote a volume or booke of the properties thereof, for the singular effects it wrought; which booke is loft, yet the diuers vertues it hath is recorded by others, to be effectuall for the spleene, lungs, stomach, liuer, head and heart; and for dropsies, old coughs, laundise, and the wormes; that it cleareth the sight, helpeth the tooth-ache, cleanseth the head of scurfe, and running sores; and is an especiall Antidote against poison: and therefore is vsed as a principall ingredient into the *Theriaca Andromachi*, which we vsually call Venice Treakle. The Apothecaries prepare hereof, both Wine, Vinegar, and Oxymel or Syrupe, which is singular to extenuate and expectorate tough flegme, which is the cause of much disquiet in the body, and an hinderer of concoction, or disgestion in the stomach, besides diuers other wayes, wherein the scales of the rootes, being dryed, are vsed. And Galen hath sufficiently explained the qualities and properties thereof, in his eight Booke of Simples.

## Chap. XII.

### *Ornithogalum.* Starre of Bethlehem.

After the Family of the Iacinths, must needes follow the kindes of Starre-flowers, or Starres of Bethlehem, as they are called, for that they doe so nearely resemble them, that diuers haue named some of them Iacinths, and referred them to that kindred: all of them, both in roote, leafe, and flower, come nearer vnto the Iacinths, then vnto any other plant. They shall therefore bee next described, euery one in their order, the greatest first, and the rest following.

### *Ornithogalum Arabicum.* The great Starre-flower of Arabia.

This Arabian Starre-flower hath many broad, and long greene leaues, very like vnto the leaues of the Orientall Iacinth, but lying for the most part vpon the ground, among which riseth vp a round greene stalke, almost two foote high, bearing at the toppe diuers large flowers, standing vpon long foote-stalkes, and at the bottome of euery one of them a small short pointed greene leafe: these flowers are made of six pure white leaues a peece, laid open as large as an ordinary Daffodill, but of the forme of a Starre Iacinth, or Starre of Bethlehem, which close as they doe euery night, and open themselues in the day time, especially in the Sunne, the smell whereof is pretty sweete, but weake: in the middle of the flower is a blackish head, composed with six white threeds, tipt with yellow pendents: the seede hath not beene obserued with vs: the roote is great and white, with a flat bottome, very impatient of our cold Winters, so that it seldome prospereth or abideth with vs; for although sometimes it doe abide a Winter in the ground, yet it often lyeth without springing blade, or any thing else a whole yeare, and then perisheth: or if it doe spring, yet many doe not beare, and most after their first bearing doe decay and perish. But if any be desirous, to know how to preserue the roote of this plant, or of many other bulbous rootes that are tender, such as the great double white Daffodill of Constantinople, and other fine Daffodils, that come from hot Countries; let them keepe this rule: Let either the roote be planted in a large pot, or tubbe of earth, and housed all the Winter, that so it may bee defended from the frosts; Or else (which is the easier way) keepe the roote out of the ground euery yeare, from September, after the leaues and stalkes are past, vntill February, in

some dry, but not hot or windy place, and then plant it in the ground vnder a South wall, or such like defended place, which will spring, and no doubt prosper well there, in regard the greatest and deepest frosts are past after February, so that seldome any great frosts come after, to pierce so deepe as the roote is to be set, or thereby to doe any great harme to it in such a place.

### The Place.

This hath been often sent out of Turkie, and likewise out of Italy; I had likewise two rootes sent mee out of Spaine by Guillaume Boel before remembred, which (as hee said) hee gathered there, but they prospered not with me, for want of the knowledge of the former rule. It may be likely that Arabia is the place, from whence they of Constantinople receiue it.

### The Time.

It flowreth in May, if it be of the first yeares bringing; or in Iune, if it haue been ordered after the manner before set downe.

### The Names.

It hath been sent out of Italy by the name of *Lilium Alexandrinum*, The Lilly of Alexandria, but it hath no affinity with any Lilly. Others call it *Hyacinthus Arabicus*; and the Italians, *Iacintho del pater nostro*: but it is no Iacinth neither, although the flowers be like some of them. Some also would referre it to a *Narcissus* or Daffodill, and it doth as little agree with it, as with a Lilly, although his flowers in largenesse and whitenesse resemble a Daffodill. Clusius hath most fitly referred it to the stocke or kindred of *Ornithogala*, or Starres of Bethlehem, as wee call them in English, and from the Turkish name, *Zumbul Arabi*, entituled it *Ornithogalum Arabicum*, although *Zumbul*, as I haue before declared, is with them, a Iacinth, wee may call it in English, The Arabian Starre-flower, or Starre of Bethlehem, or the great Starre-flower of Arabia.

### 1. *Ornithogalum maximum album.*

The greatest white Starre-flower, or Starre of Bethlehem.

This great Starre-flower hath many faire, broad, long, and very fresh green leaues, rising vp very early, and are greater, longer, and greener then the leaues of any Orientall Iacinth, which doe abide greene, from the beginning or middle of Ianuary, or before sometimes, vntill the end of May, at which time they begin to fade, and the stalke with the head of flowers beginneth to rise, so that it will haue either few or no leaues at all, when the flowers are blowne: the stalke is strong, round, and firme, rising two foote high or more, bearing at the toppe a great bush of flowers, seeming at the first to be a great greene eare of corne, for it is made spike-fashion, which when the flowers are blowne, doth rise to be very high, slender or small at the head aboue, and broad spread and bushing below, so that it is long in flowring; for they flower below first, and so vpwards by degrees: these flowers are snow white, without any line on the backside, and is therein like vnto the former, as also in whitenesse, but nothing so large, with a white vmbone or head in the middle, beset with many white threeds, tipt with yellow: the seede is blacke and round, contained in three square heads: the roote is great, thicke, and short, and somewhat yellowish on the outside, with a flat bottome, both like the former, and the next that followeth.

### 2. *Ornithogalum maius spicatum album.*

The great white spiked Starre-flower.

This spiked Starre-flower in his growing, is somewhat like vnto the last described,

but ſpringeth not vp ſo early, nor hath his leaues ſo greene, or large, but hath broad, long, whitiſh greene hollow leaues, pointed at the end, among which riſeth vp the ſtalke, which is ſtrong and high, as the former, hauing a great buſh of flowers at the toppe, ſtanding ſpike-faſhion, ſomewhat like the former, flowring in the ſame maner by degrees, firſt below, and ſo vpwards; but it is not ſo thicke ſet with flowers, nor ſo farre ſpread at the bottome as it, the flowers alſo are not ſo white, and each of the leaues of them haue a greene line downe the backe, leauing the edges on both ſides white: after the flowers are paſt, the heads for ſeede grow three ſquare, like the other, bearing ſuch like blacke ſeede therein: the roote hereof is vſually bigger then the laſt, and whiter on the outſide.

### 3. *Ornithogalum Pannonicum.* The Hungarian Starre-flower.

This Hungarian Starre-flower ſhooteth out diuers narrow, long, whitiſh greene leaues, ſpread vpon the ground before Winter, which are very like vnto the leaues of Gilloflowers, and ſo abide aboue ground, hauing a ſtalke riſing in the middle of them the next Spring, about halfe a foote high or thereabouts, bearing many white flowers at the toppe, with greene lines downe the backe of them, very like vnto the ordinary Starres of Bethlehem: the roote is greater, thicker, and longer then the ordinary Starres, and for the moſt part, two ioyned together, ſomewhat grayiſh on the out ſide.

### 4. *Ornithogalum vulgare.* The Starre of Bethlehem.

The ordinary Starre of Bethlehem is ſo common, and well knowne in all countries and places, that it is almoſt needleſſe to deſcribe it, hauing many greene leaues with white lines therein, and a few white flowers ſet about the toppe of the ſtalke, with greeniſh lines downe the backe: the roote is whitiſh, and encreaſeth aboundantly.

### 5. *Aſphodelus bulboſus Galeni, ſiue Ornithogalum maius flore ſubnireſcente.* The bulbous Aſphodill, or greene Starre-flower.

Diuers haue referred this plant vnto the Aſphodils, becauſe (as I thinke) the flowers hereof are ſtraked on the backe, and the leaues long and narrow, like vnto the Aſphodils; but the roote of this being bulbous, I rather (as ſome others doe) ioyne it with the *Ornithogala*, for they alſo haue ſtrakes on the backe of the flowers. It hath many whitiſh greene leaues, long and narrow, ſpread vpon the ground, which ſpring vp in the beginning of the yeare, and abide vntill May, and then they withering, the ſtalke ſpringeth vp almoſt as high as the firſt, hauing many pale yellowiſh greene flowers, but ſmaller, and growing more ſparſedly about the ſtalke vpon ſhort foot-ſtalkes, but in a reaſonable long head ſpike-faſhion: the ſeede is like vnto the ſecond kinde, but ſmaller: the roote is ſomewhat yellowiſh, like the firſt great white kinde.

#### The Place.

The firſt is onely nurſed in Gardens, his originall being not well knowne, yet ſome attribute it vnto *Pannonia* or Hungary. The ſecond hath been found neare vnto Barcinone, and Toledo in Spaine. The third was found in Hungary by Cluſius. Our ordinary euery where in the fields of Italy and France, and (as it is ſaid) in England alſo. And the laſt groweth likewiſe by the corne fields in the vpper Hungary.

#### The Time.

They flower in Aprill and May, and ſometimes in Iune.

#### The Names.

The firſt is called by Cluſius *Ornithogalum maximum album*, becauſe it is greater

1 *Ornithogalum Arabicum.* The great starre-flower of Arabia. 2 *Ornithogalum maximum album* The greatest white starre flower. 3 *Ornithogalum maius spicatum album.* The great white spiked starre flower. 4 *Ornithogalum Pannonicum album.* The Hungarian starre flower. 5 *Asphodelus bulbosus Galeni, sive Ornithogalum maius subviridescente flore.* The bulbed Asphodill, or greene starre flower. 6 *Ornithogalum Hispanicum minus.* The little starre-flower of Spaine. 7 *Ornithogalum luteum.* The yellow starre-flower of Bethlehem. 8 *Ornithogalum Neapolitanum.* The starre-flower of Naples.

greater then the next, which hee tooke formerly for the greateſt: but it might more fitly, in my iudgement, bee called *Aſphodelus bulboſus albus* (if there be any *Aſphodelus bulboſus* at all) becauſe this doth ſo nearly reſemble that, both in the early ſpringing, and the decay of the greene leaues, when the ſtalkes of flowers doe riſe vp. Diuers alſo doe call it *Ornithogalum Pannonicum maximum album.*

The ſecond hath his name in his title, as moſt authors doe ſet it downe, yet in the great Herball referred to Dalechampius, it is called *Ornithogalum magnum Myconi.*

The third hath his name from the place of his birth, and the other from his popularity, yet Dodonæus calleth it *Bulbus Leucanthemos.*

The laſt is called by diuers *Aſphodelo-hyacinthinus*, and *Hyacintho-aſphodelus Galeni.* Dodonæus calleth it *Aſphodelus famina*, and *Aſphodelus bulboſus.* But Lobel, and Gerrard from him, and Dodonæus, doe make this to haue white flowers, whereas all that I haue ſeene, both in mine owne, and in others Gardens, bore greeniſh flowers, as Cluſius ſetteth it truely downe. Lobel ſeemeth in the deſcription of this, to confound the *Ornithogalum* of Mompelier with it, and calleth it *Aſphodelus hyacinthinus forte Galeni*, and ſaith that ſome would call it *Pancratium Monſpelienſe*, and *Aſphodelus Galeni.* But as I haue ſhewed, the *Ornithogalum ſpicatum* and this, doe plainly differ the one from the other, and are not both to be called by one name, nor to be reckoned one, but two diſtinct plants.

*Ornithogalum Æthiopicum.* The Starre-flower of Æthiopia.

The leaues of this plant are a foote long, and at the leaſt an inch broad, which being broken, are no leſſe woolly then the woolly Iacinth: the ſtalke is a cubit high, ſtrong and greene; from the middle whereof vnto the toppe, ſtand large ſnow white flowers, vpon long, greene, thicke foot-ſtalkes, and yellowiſh at the bottome of the flower; in the middle whereof ſtand ſix white threeds, tipt with yellow chiues, compaſſing the head, which is three ſquare, and long containing the ſeede: the roote is thicke and round, ſomewhat like the *Aſphodelus Galeni.*

The Place.

This plant was gathered by ſome Hollanders, on the Weſt ſide of the Cape of good Hope.

The Time.

It flowred about the end of Auguſt with thoſe that had it.

The Names.

Becauſe it came from that part of the continent beyond the line, which is reckoned a part of Æthiopia, it is thereupon ſo called as it is ſet downe.

*Ornithogalum Neopolitanum.* The Starre-flower of Naples.

This beautifull plant riſeth out of the ground very early, with foure or fiue hollow pointed leaues, ſtanding round together, of a whitiſh greene colour, with a white line downe the middle of euery leafe on the inſide, ſomewhat narrow, but long, (Fabius Columna ſaith, three foot long in Italy, but it is not ſo with vs) in the middle of theſe leaues riſeth vp the ſtalke, a foote and a halfe high, bearing diuers flowers at the toppe, euery one ſtanding in a little cuppe or huske, which is diuided into three or foure parts, hanging downe very long about the heads for ſeede: after the flower is paſt, theſe flowers doe all hang downe their heads, and open one way, although their little foot-ſtalkes come forth on all ſides of the greater ſtalke, being large, and compoſed of ſix long leaues, of a pure white on the inſide, and of a blewiſh or whitiſh greene colour

on

on the outside, leauing the edges of euery leafe white on both sides: in the middle of these flowers stand other small flowers, each of them also made of six small white leaues a peece, which meeting together, seeme to make the shew of a cuppe, within which are contained six white threeds, tipt with yellow, and a long white pointell in the middle of them, being without any sent at all: after the flowers are past, come vp great round heads, which are too heauie for the stalke to beare; and therefore lye downe vpon the leaues or ground, hauing certaine lines or stripes on the outside, wherein is contained round, blacke, rough seede: the roote is great and white, and somewhat flat at the bottome, as diuers of these kindes are, and doe multiply as plentifully into small bulbes as the common or any other.

## The Place.

This Starre-flower groweth in the Medowes in diuers places of Naples, as Fabius Columna, and Ferrantes Imperatus doe testifie, from whence they haue been sent. And Matthiolus, who setteth out the figure thereof among his Daffodils, had (it should seeme) seene it grow with him.

## The Time.

It flowreth in May, although it begin to spring out of the ground oftentimes in Nouember, but most vsually in Ianuary: the seede is ripe in Iuly.

## The Names.

Matthiolus reckoneth this (as is said) among the Daffodils, for no other respect, as I conceiue, then that he accounted the middle flower to bee the cuppe or trunke of a Daffodill, which it doth somewhat resemble, and setteth it forth in the fourth place, whereupon many doe call it *Narcissus quartus Matthioli*, The fourth Daffodill of Matthiolus. Fabius Columna calleth it *Hyacinthus aruorum Ornithogali flore.* Clusius (to whom Imperatus sent it, in stead of the Arabian which hee desired) calleth it of the place from whence he receiued it, *Ornithogalum Neopolitanum*, and we thereafter call it in English, The Starre-flower of Naples.

*Ornithogalum Hispanicum minus.* The little Starre-flower of Spaine.

Clusius hath set forth this plant among his *Ornithogala* or Starre-flowers, and although it doth in my minde come nearer to a *Hyacinthus*, then to *Ornithogalum*, yet pardon it, and let it passe as he doth. From a little round whitish roote, springeth vp in the beginning of the yeare, fiue or six small long green leaues, without any white line in the middle of them, among which rise vp one or two small stalkes, an hand length high or better, bearing seuen or eight, or more flowers, growing as it were in a tuft or vmbell, with small long leaues at the foote of euery stalke, the lower flowers being equall in length with the vppermost, of a pale whitish blew or ash colour, with a strake or line downe the backe of euery leafe of them, with some white threeds standing about a blewish head in the middle: these flowers passe away quickly, and giue no seed, so that it is not knowne what seede it beareth.

## The Place.

This groweth in Spaine, and from thence hath been brought to vs.

## The Time.

It flowreth in May.

## The Names.

It hath no other name then is set down in the title, being but lately found out.

I. Ornithogalum

1. *Ornithogalum album vnifolium.* The white starre-flower with one blade.

This little starre-flower I bring into this place, as the fittest in my opinion where to place it, vntill my minde change to alter it. It hath a very small round white roote, from whence springeth vp one very long and round greene leafe, like vnto a rush, but that for about two or three inches aboue the ground, it is a little flat, and from thence springeth forth a small stalke not aboue three or foure inches high, bearing at the top thereof three or foure small white flowers, consisting of six leaues a peece, within which are six white chiues, tipt with yellow pendents, standing about a small three square head, that hath a white pointell sticking as it were in the middest thereof: the flower is pretty and sweete, but not heady.

*Ornithogalum luteum.* The yellow Starre of Bethlehem.

This yellow Starre-flower riseth vp at the first, with one long, round, greenish leafe, which openeth it selfe somewhat aboue the ground, and giueth out another small leafe, lesser and shorter then the first, and afterward the stalke riseth from thence also, being foure or fiue inches high, bearing at the toppe three or foure small green leaues, and among them foure or fiue small yellow starre-like flowers, with a greenish line or streake downe the backe of euery leafe, and some small reddish yellow threeds in the middle: it seldome giueth seede: the roote is round, whitish, and somewhat cleare, very apt to perish, if it bee any little while kept dry out of the ground, as I haue twice tryed to my losse.

The Place.

The first grew in Portugall, and Clusius first of all others desciphers it. The other is found in many places both of Germany and Hungary, in the moister grounds.

The Time.

The first flowreth in May: the other in Aprill, and sometimes in March.

The Names.

Carolus Clusius calleth the first *Bulbus vnifolius*, or *Bolbine*, but referreth it not to the stocke or kindred of any plant; but (as you see) I haue ranked it with the small sorts of *Ornithogalum*, and giue it the name accordingly.

The other is referred for likenesse of forme, and not for colour, vnto the *Ornithogala*, or Starres of Bethlehem. It is called by Tragus and Fuchsius *Bulbus siluestris*, because of the obuiousnesse. Cordus taketh it to be *Sisyrinchium*. Lacuna calleth it *Bulbus esculentus*. Lobel and others in these dayes generally, *Ornithogalum luteum*, and wee thereafter in English, The yellow Starre-flower, or Starre of Bethlehem.

The Vertues.

The first kinde being but lately found out, is not knowne to be vsed. The rootes of the common or vulgar, are (as Matthiolus saith) much eaten by poore people in Italy, either rawe or roasted, being sweeter in taste then any Chesnut, and seruing as well for a necessary food as for delight. It is doubtfull whether any of the rest may be so vsed; for I know not any in our Land hath made any experience.

There are many other sorts of Starre-flowers, which are fitter for a generall then this History; and therefore I referre them thereunto.

## CHAP. XIII.

### *Moly.* Wilde Garlicke.

VNto the former Starre-flowers, must needes bee ioyned another tribe or kindred, which carry their straked flowers Starre-fashion, not spikewise, but in a tuft or vmbell thicke thrust or set together. And although diuers of them smell not as the former, but most of their first Grandfathers house, yet all doe not so; for some of them are of an excellent sent. Of the whole Family, there are a great many which I must leaue, I will onely select out a few for this our Garden, whose flowers for their beauty of statelinesse, forme, or colour, are fit to bee entertained, and take place therein, euery one according to his worth, and are accepted of with the louers of these delights.

1. *Moly Homericum, vel potius Theophrasti.* The greatest Moly of Homer.

Homers Moly (for so it is most vsually called with vs) riseth vp most commonly with two, and sometimes with three great, thicke, long, and hollow guttured leaues, of a whitish greene colour, very neare the colour of the Tulipa leafe, hauing sometimes at the end of some of the leaues, and sometimes apart by it selfe, a whitish round small button, like vnto a small bulbe, the like whereof also, but greater, doth grow betweene the bottome of the leaues and the stalke neare the ground, which being planted when it is ripe, will grow into a roote of the same kinde: among these leaues riseth vp a round, strong, and tall stalke, a yard high or better, bare or naked vnto the toppe, where it beareth a great tuft or vmbell of pale purplish flowers, all of them almost standing vpon equall foot-stalkes, or not one much higher then another, consisting of fiue leaues a peece, striped downe the backe with a small pale line, hauing a round head or vmbone with some threeds about it in the midst: These flowers doe abide a great while blowne before they vade, which smell not very strong, like any Onion or Garlicke, but of a faint smell: and after they are past come the seede, which is blacke, wrapped in white close huskes: the roote groweth very great, sometimes bigger then any mans closed fist, smelling strong like Garlicke, whitish on the outside, and greene at the toppe, if it be but a while bare from the earth about it.

2. *Moly Indicum siue Cancasson.* The Indian Moly.

The Indian Moly hath such like thicke large leaues, as the Homers Moly hath, but shorter and broader, in the middle whereof riseth vp a short weake stalke, almost flat, not hauing any flowers vpon it, but a head or cluster of greenish scaly bulbes, inclosed at the first in a large thinne skinne, which being open, euery bulbe sheweth it selfe, standing close one vnto another vpon his foot-stalke, of the bigneſſe of an Acorne, which being planted, will grow to bee a plant of his owne kinde: the roote is white and great, couered with a darke coate or skinne, which encreaseth but little vnder ground; but besides that head, it beareth small bulbes aboue the ground, at the bottome of the leaues next vnto the stalke, like vnto the former.

The Place.

Both these doe grow in diuers places of Spaine, Italy, and Greece; for the last hath been sent out of Turkie among other rootes. Ferrantes Imperatus a learned Apothecary of Naples, sent it to diuers of his friends in these parts, and hath described it in his naturall history among other plants, printed in the Italian tongue. It grew also with Iohn Tradescante at Canterbury, who sent me the head of bulbes to see, and afterwards a roote, to plant it in my Garden.

The

### The Time.

The first flowreth in the end of May, and abideth vnto the midst of Iuly, and sometimes longer. The other beareth his head of bulbes in Iune and Iuly.

### The Names.

We haue receiued them by their names expressed in their titles, yet the last hath also been sent by the name of *Ornithogalum Italicum*, but as all may easily see, it is not of that kindred.

1. *Moly montanum Pannonicum bulbiferum primum.*
The first bulbed Moly of Hungary.

This first Hungarian Moly hath three or foure broad and long greene leaues, folded together at the first, which after open themselues, and are carried vp with the stalke, standing thereon one aboue another, which is a foote high; at the toppe whereof doe grow a few sad reddish bulbes, and betweene them long footstalkes, bearing flowers of a pale purplish colour; after which followeth blacke seede, inclosed in roundish heads: the roote is not great, but white on the outside, very like vnto the roote of Serpents Moly, hereafter described, encreasing much vnder ground, & smelling strong.

2. *Moly montanum Pannonicum bulbiferum secundum.*
The second bulbed Moly of Hungary.

The second Moly hath narrower greene leaues then the former: the stalke is about the same height, and beareth at the toppe a great cluster of small greene bulbes, which after turne of a darker colour; from among which come forth long foot-stalks, whereon stand purplish flowers: the roote is couered with a blackish purple coate or skinne.

3. *Moly Serpentinum.* Serpents Moly.

This Moly must also be ioyned vnto the bulbous Molyes, as of kindred with them, yet of greater beauty and delight, because the bulbes on the heads of the small stalkes are redder, and more pleasant to behold: the stalke is lower, and his grassie winding leaues, which turne themselues (whereof it tooke the name) are smaller, and of a whiter greene colour: it beareth among the bulbes purplish flowers also, but more beautifull, the sent whereof is nothing so strong: the roote is small, round, and whitish, encreasing into a number of small rootes, no bigger then pease round about the greater roote.

4. *Moly caule & folijs triangularibus.* The three cornered Moly.

This three square Moly hath foure or fiue long, and somewhat broad pale greene leaues, flat on the vpper side, and with a ridge downe the backe of the leafe, which maketh it seeme three square: the stalke which riseth vp a foote and a halfe high or better, is three square or three cornered also, bearing at the toppe out of a skinnie huske diuers white flowers, somewhat large and long, almost bell-fashion, with stripes of greene downe the middle of euery leafe, and a few chiues tipt with yellow in the middle about the head, wherein when it is ripe, is inclosed small blacke seede: the roote is white on the outside, and very like the yellow Moly; both roote, leafe, and flower hath a smacke, but not very strong of Garlicke.

5. *Moly Narcissinis folijs.* Daffodill leafed Moly.

This Moly hath many long, narrow, and flat greene leaues, very like vnto the leaues of a Daffodill, from whence it tooke his name (or rather of the early greater *Leucoium bulbosum*,

1 *Moly Homericum vel potius Theophrasti.* The greater Moly of Homer. 2 *Moly Indicum siue Caucasen.* The Indian Moly. 3 *Moly Pannonicum bulbiferum.* The bulbed Moly of Hungary. 4 *Moly Serpentinum.* Serpents Moly. 5 *Moly purpureum Neapolitanum.* The purplish Moly of Naples. 6 *Moly caule & folijs triangularibus.* The three cornered Moly. 7 *Moly latifolium flore luteo.* The yellow Moly. 8 *Moly Dioscoridem Hispanicum.* The Spanish Moly of Dioscorides. 9 *Moly Zibettinum vel Moschatum.* The sweete smelling Moly of Mompelier. 10 *Moly serotinum Coniferum.* The late Pineapple Moly.

6. Allium triquetrum. Spa. hist.

*bulboſum*, or bulbed Violet before deſcribed, ioyned next vnto the Daffodils, becauſe it is ſo like them) among which riſeth vp two or three ſtalkes ſometimes, each of a foot and a halfe high, bearing at the toppe, incloſed in a skinny hoſe, as all the Molyes haue, a number of ſmall purpliſh flowers, which doe not long abide, but quickly fade: the ſeede is blacke as others are; the roote is ſometimes knobbed, and more often bulbed, hauing in the knobs ſome markes of the old ſtalkes to be ſeene in them, and ſmelleth ſomewhat like Garlicke, whereby it may be knowne.

6. *Moly montanum latifolium luteo flore.* The yellow Moly.

The yellow Moly hath but one long and broad leafe when it doth not beare flower, but when it will beare flower, it hath two long and broad leaues, yet one alwaies longer and broader then the other, which are both of the ſame colour, and neare the bigneſſe of a reaſonable Tulipa leafe: betweene theſe leaues groweth a ſlender ſtalke, bearing at the toppe a tuft or vmbell of yellow flowers out of a skinnie hoſe, which parteth three wayes, made of ſix leaues a peece, laid open like a Starre, with a greeniſh backe or outſide, and with ſome yellow threeds in the middle: the ſeede is blacke, like vnto others: the roote is whitiſh, two for the moſt part ioyned together, which encreaſeth quickly, and ſmelleth very ſtrong of Garlicke, as both flowers and leaues doe alſo.

7. *Moly Pyrenæum purpureum.* The purple mountaine Moly.

This purple Moly hath two or three leaues, ſomewhat like the former yellow Moly, but not ſo broad, nor ſo white: the ſtalke hath not ſo many flowers thereon, but more ſparingly, and of an vnpleaſant purple colour: the roote is whitiſh, ſmelling ſomewhat ſtrongly of Garlicke, but quickly periſheth with the extremity of our cold Winters, which it will not abide vnleſſe it be defended.

8. *Moly montanum latiſolium purpureum Hiſpanicum.*
The purple Spaniſh Moly.

This Moly hath two broad and very long greene leaues, like vnto the yellow Moly, in this, that they doe compaſſe one another at the bottome of them, between which riſeth vp a ſtrong round ſtalke, two foote high or more, bearing at the toppe, out of a thinne huske, a number of faire large flowers vpon long foot-ſtalkes, conſiſting of ſix leaues a peece, ſpread open like a Starre, of a fine delayed purple or bluſh colour, with diuers threeds of the ſame colour, tipt with yellow, ſtanding about the middle head: betweene the ſtalke and the bottome of the leaues it hath ſome ſmall bulbes growing, which being planted, will ſoone ſpring and encreaſe: the roote alſo being ſmall and round, with many fibres thereat, hath many ſmall bulbes ſhooting from them; but neither roote, leafe, nor flower, hath any ill ſent of Garlicke at all.

9. *Moly purpureum Neapolitanum.* The purple Moly of Naples.

The Neapolitane Moly hath three or foure ſmall long greene leaues ſet vpon the ſtalke after it is riſen vp, which beareth a round head of very fine purple flowers, made of ſix leaues a peece, but ſo cloſing together at the edge, that they ſeeme like vnto ſmall cuppes, neuer laying themſelues open, as the other doe; this hath ſome ſent of his originall, but the roote more then any part elſe, which is white and round, quickly encreaſing as moſt of the Molyes doe.

10. *Moly pyxidatum argenteum Hiſpanicum.*
The Spaniſh ſiluer cupped Moly.

This Spaniſh Moly hath two or three very long ruſh like leaues, which riſe vp with the ſtalke, or rather vaniſh away when the ſtalke is riſen vp to bee three foote high or more, bearing a great head of flowers, ſtanding cloſe at the firſt, but afterwards ſpreading much one from another, euery flower vpon a long foote-ſtalke, being of a white

siluer colour, with stripes or lines on euery side, and fashioned small and hollow, like a cuppe or boxe: the seede I could neuer obserue, because it flowreth so late, that the Winter hindereth it from bearing seede with vs: the roote is small and round, white, and in a manner transparent, at least so shining, as if it were so, and encreaseth nothing so much, as many of the other sorts: this hath no ill sent at all, but rather a pretty smell, not to bee misliked.

### 11. *Moly serotinum Coniferum.* The late Pineapple Moly.

This late Moly that was sent me with the last described, and others also from Spain, riseth vp with one long greene leafe, hollow and round vnto the end, towards this end on the one side, breaketh out a head of flowers, enclosed in a thinne skinne, which after it hath so stood a good while, (the leafe in the meane time rising higher, and growing harder, becommeth the stalke) breaketh, and sheweth a great bush or head of buds for flowers, thicke thrust together, fashioned very like vnto the forme of a Pineapple (from whence I gaue it the name) of the bignes of a Walnut: after this head hath stood in this manner a moneth or thereabouts, the flowers shew themselues to bee of a fine delayed or whitish purple colour, with diuers stripes in euery of them, of the same cup-fashion with the former, but not opening so plainly, so that they cannot bee discerned to bee open, without good heede and obseruation. It flowreth so late in Autumne, that the early frosts doe quickly spoile the beauty of it, and soone cause it to rotte: the roote is small and round, and shining like the last, very tender also, as not able to abide our sharpe Winters, which hath caused it vtterly to perish with me.

### 12. *Moly Dioscorideum.* Dioscorides his Moly.

The roote of this small Moly is transparent within, but couered with a thicke yellowish skinne, of the bigneſſe of an Hasell Nut, or somewhat bigger, which sendeth forth three or foure narrow grassie leaues, long and hollow, and a little bending downwards, of a whitish greene colour, among which riseth vp a slender weake stalke, a foot and a halfe high, bearing at the toppe, out of a thinne skinne, a tuft of milke white flowers, very like vnto those of Ramsons, which stand a pretty while in their beauty, and then passe away for the most part without giuing any seede: this hath little or no sent of Garlicke.

We haue another of this sort that is lesser, and the flowers rounder pointed.

### 13. *Moly Dioscorideum Hispanicum.* The Spanish Moly of Dioscorides.

This Moly came vnto me among other Molyes from Spaine, and is in all things like vnto the last described, but fairer, larger, and of much more beauty, as hauing his white flowers twice as great as the former; but (as it seemeth) very impatient of our Winters, which it could not at any hand endure, but quickly perished, as some others that came with it also.

### 14. *Moly Moschatum vel Zibettinum Monspelienſe.* The sweete smelling Moly of Mompelier.

This sweete Moly, which I haue kept for the last, to close vp your senses, is the smallest, and the fineſt of all the rest, hauing foure or fiue small greene leaues, almost as fine as haires, or like the leaues of the Feather-grasse: the stalke is about a foote high, bearing fiue or six or more small white flowers, laid open like Starres, made of six leaues a peece, of an excellent sweete sent, resembling Muske or Ciuet; for diuers haue diuersly censured of it. It flowreth late in the yeare, so that if the precedent Summer bee either ouer moist, or the Autumne ouer early cold, this will not haue that sweete sent, that it will haue in a hot drie time, and besides must be carefully respected: for it will hardly abide the extremity of our sharpe Winters.

### The Place.

The places of these Molyes, are for the most part expressed in their titles, or in their descriptions.

### The Time.

The time is set downe, for the most part to bee in Iune and Iuly, the rest later.

### The Names.

To make further relation of names then are expressed in their tiles, were needlesse; let these therefore suffice.

### The Vertues.

All these sorts of Molyes are small kindes of wilde Garlicke, and are to be vsed for the same purposes that the great Garden Garlicke is, although much weaker in their effects. For any other especiall property is in any of these, more than to furnish a Garden of variety, I haue not heard at all.

And thus much may suffice of these kindes for our Garden, reseruing manie others that might be spoken of, to a generall worke, or to my Garden of Simples, which as God shall enable me, and time giue leaue, may shew it selfe to the world, to abide the iudicious and criticke censures of all.

## Chap. XIIII.

### *Asphodelus.* The Asphodill.

THere remaine some other flowers, like vnto the last described, to be specified, which although they haue no bulbous rootes, yet I thinke them fittest to bee here mentioned, that so I may ioyne those of neerest similitude together, vntill I haue finished the rest that are to follow.

1. *Asphodelus maior albus ramosus.* The great white branched Asphodill.

The great white Asphodill hath many long, and narrow, hollow three square leaues, sharpe pointed, lying vpon the ground round about the roote: the stalke is smooth, round, and naked without leaues, which riseth from the midst of them, diuided at the toppe into diuers branches, if the plant bee of any long continuance, or else but into two or three small branches, from the sides of the maine great one, whereon doe stand many large flowers Starre-fashion, made of six leaues a peece, whitish on the inside, and straked with a purplish line downe the backside of euery leafe, hauing in the middle of the flowers some small yellow threeds: the seede is blacke, and three square, greater then the seede of Bucke wheate, contained in roundish heads, which open into three parts: the roote is composed of many tuberous long clogges, thickest in the middle, and smaller at both ends, fastened together at the head, of a darke grayish colour on the outside, and yellow within.

2. *Asphodelus albus non ramosus.* The white vnbranched Aspodill.

The vnbranched Asphodill is like vnto the former, both in leaues and flowers, but that the flowers of this are whiter, and without any line or strake on the backe side,

and

and the stalkes are without branches: the rootes likewise are smaller, and fewer, but made after the same fashion.

3. *Asphodelus maior flore carneo.* The blush coloured Asphodill.

This Asphodill is like to the last in forme of leaues and branches, and differeth in this, that his leaues are marked with some spots, and the flowers are of a blush or flesh colour, in all other things alike.

4. *Asphodelus minimus albus.* The least white Asphodill.

This least Asphodill hath foure or fiue very narrow long leaues, yet seeming three square like the greatest, bearing a small stalke, of about a foote high among them, without any branches, and at the toppe a few white flowers, straked both within and without, with a purplish line in the middle of euery leafe. The rootes are such like tuberous clogges as are in the former, but much lesser.

5. *Asphodelus albus minor siue Fistulosus.* The little hollow white Aspnodill.

This little white Asphodill hath a number of leaues growing thicke together, thicker and greener then those of the small yellow Asphodill, or Kings Speare next following, among which riseth vp diuers round stalkes, bearing flowers from the middle to the toppe, Starre-fashion, with small greene leaues among them, which are white on the inside, and striped on the backe with purple lines, like vnto the first described: the seede, and heads containing them, are three square, like the seede of the little yellow Asphodill: the rootes of this kinde are not glandulous, as the former, but stringie, long and white: the whole plant is very impatient of our cold Winters, and quickly perisheth, if it be not carefully preserued, both from the cold, and much wet in the Winter, by housing it; and then it will abide many yeares: for it is not an annuall plant, as many haue thought.

6. *Asphodelus luteus minor, siue Hastula regia.*
The small yellow Asphodill, or Kings speare.

This small yellow Asphodill, which is vsually called the Kings speare, hath many long narrow edged leaues, which make them seeme three square, of a blewish or whitish greene colour: the stalke riseth vp three foote high oftentimes, beset with small long leaues vp vnto the very flowers, which grow thicke together spike-fashion one aboue another, for a great length, and wholly yellow, laid open like a Starre, somewhat greater then the last white Asphodill, and smaller then the first, which when they are past yeeld round heads, containing blacke cornered seede, almost three square: the rootes are many long yellow strings, which spreading in the ground, doe much encrease.

## The Place.

All these Asphodils doe grow naturally in Spaine and France, and from thence were first brought vnto vs, to furnish our Gardens.

## The Time.

All the glandulous rooted Asphodils doe flower some in May, and some in Iune; but the two last doe flower, the yellow or last of them in Iuly, and the former white one in August and September, and vntill the cold and winter hinder it.

## The Names.

Their seuerall names are giuen them in their titles, as much as is fit for this

this diſcourſe. For to ſhew you that the Greekes doe call the ſtalke of the great Aſphodill Ἀσφοδελὸς, and the Latines *Albucum*, or what elſe belongeth to them, is fitter for another worke, vnto which I leaue them.

The baſtard Aſphodils ſhould follow next in place, if this worke were fit for them; but becauſe I haue tyed my ſelfe to expreſſe onely thoſe flowers and plants, that for their beauty, or ſent, or both, doe furniſh a Garden of Pleaſure, and they haue none, I leaue them to a generall Hiſtory of plants, or that Garden of Simples before ſpoken of, and will deſcribe the Lilly Aſphodils, and the *Phalangia* or Spider-worts, which are remaining of thoſe, that ioyne in name or faſhion, and are to be here inſerted, before I paſſe to the reſt of the bulbous rootes.

1. *Liliaſphodelus phœniceus.* The gold red Day Lilly.

Becauſe the rootes of this and the next, doe ſo nearely agree with the two laſt recited Aſphodils, I haue ſet them in this place, although ſome doe place them next after the Lillies, becauſe their flowers doe come neareſt in forme vnto Lillies; but whether you will call them Aſphodils with Lilly flowers, as I thinke it fitteſt, or Lillies with Aſphodill rootes, or Lillies without bulbous rootes, as others doe, I will not contend.

The red Day Lilly hath diuers broad and long freſh greene leaues, folded at the firſt as it were double, which after open, and remaine a little hollow in the middle; among which riſeth vp a naked ſtalke three foot high, bearing at the toppe many flowers, one not much diſtant from another, and flowring one after another, not hauing lightly aboue one flower blown open in a day, & that but for a day, not laſting longer, but cloſing at night, and not opening againe; whereupon it had his Engliſh name, The Lilly for a day: theſe flowers are almoſt as large as the flowers of the white Lilly, and made after the ſame faſhion, but of a faire gold red, or Orange tawny colour. I could neuer obſerue any ſeede to follow theſe flowers; for they ſeeme the next day after they haue flowred, (except the time be faire and dry) to bee ſo rotten, as if they had lyen in wet to rotte them, whereby I thinke no ſeede can follow: the rootes are many thicke and long yellow knobbed ſtrings, like vnto the ſmall yellow Aſphodill rootes, but ſomewhat greater, running vnder ground in like ſort, and ſhooting young heads round about.

2. *Liliaſphodelus luteus.* The yellow Day Lilly.

I ſhall not neede to make a repetition of the deſcription of this Day Lilly, hauing giuen you one ſo amply before, becauſe this doth agree thereunto ſo nearely, as that it might ſeeme the ſame; theſe differences onely it hath, the leaues are not fully ſo large, nor the flower ſo great or ſpread open, and the colour thereof is of a faire yellow wholly, and very ſweet, which abideth blowne many daies before it fade, and hath giuen blacke round ſeede, growing in round heads, like the heads of the ſmall yellow Aſphodill, but not ſo great.

Cluſius hath ſet downe, that it was reported, that there ſhould be another Liliaſphodill with a white flower, but we can heare of none ſuch as yet; but I rather thinke, that they that gaue that report might be miſtaken, in thinking the Sauoye Spider-wort to be a white Liliaſphodill, which indeede is ſo like, that one not well experienced, or not well regarding it, may ſoone take one for another.

The Place.

Their originall is many moiſt places in Germany.

The Time.

They flower in May and Iune.

The Names.

They are called by ſome *Liliago*, and *Lilium non bulboſum*, and *Liliaſphodelus*,

1 *Asphodelus maior albus ramosus.* The great white branched Asphodill. 2 *Asphodelus minor albus seu fistulosus.* The little hollow white Asphodill. 3 *Asphodelus minor luteus, siue Hastula regia.* The small yellow Asphodill, or Kings speare. 4 *Liliasphodelus luteus.* The yellow Day Lilly. 5 *Liliasphodelus phœniceus.* The gold red Day Lilly.

*delus*. In Englisſh we call them both Day Lillies, but the name doth not ſo well agree with the laſt, as with the firſt, for the cauſes aboue ſpecified.

### The Vertues.

The rootes of Aſphodill hath formerly beene had in great account, but now is vtterly neglected; yet by reaſon of their ſharpeneſſe they open and cleanſe, and therefore ſome haue of late vſed them for the yellow Iaundiſe. The Day Lillies haue no phyſicall vſe that I know, or haue heard.

## Chap. XV.

### *Phalangium*. Spider-wort.

THeſe plants doe ſo nearely reſemble thoſe that are laſt ſet forth, that I thinke none that knowes them, will doubt, but that they muſt follow next vnto them, being ſo like vnto them, and therefore of the faireſt of this kinde firſt.

#### 1. *Phalangium Alſobrogicum*. The Sauoye Spider-wort.

The Sauoye Spider-wort ſpringeth vp with foure or fiue greene leaues, long and narrow, yet broader at the bottome, narrower pointed at the end, and a little hollow in the middle; among which riſeth vp a round ſtiffe ſtalke, a foote and a halfe high, bearing at the toppe one aboue another, ſeuen or eight, or more flowers, euery one as large almoſt as the yellow Day Lilly laſt deſcribed, but much greater then in any other of the Spider-worts, of a pure white colour, with ſome threeds in the middle, tipt with yellow, and a ſmall forked pointell: after the flowers are paſt, the heads or ſeede veſſels grow almoſt three ſquare, yet ſomewhat round, wherein is contained blackiſh ſeede: the rootes are many white, round, thicke, brittle ſtrings, ioyned together at the head, but are nothing ſo long, as the rootes of the other *Phalangia* or Spider-worts.

#### 2. *Phalangium maius Italicum album*. The great Italian Spider-wort.

This great Spider-wort hath diuers long and narrow leaues ſpread vpon the ground, and not riſing vp as the former, and not ſo broad alſo as the former, but ſomewhat larger then thoſe that follow: the ſtalke is bigger, but ſeldome riſeth vp ſo high as the next, whereof this is a larger kinde, hauing a long vnbranched ſtalke of white flowers, laid open like ſtarres as it hath, but ſomewhat greater: the rootes are long and white, like the next, but ſomewhat larger.

#### 3. *Phalangium non ramoſum vulgare*. Vnbranched Spider-wort.

The leaues of this Spider-wort doe ſeeme to bee little bigger or longer then the leaues of graſſe, but of a more grayiſh green colour, riſing immediately from the head or tuft of rootes; among which riſe vp one or two ſtalkes, ſometimes two or three foote long, beſet toward the toppe with many white Starre-like flowers, which after they are paſt turne into ſmall round heads, containing blacke ſeede, like vnto the ſeed of the little yellow Aſphodill, but leſſer: the rootes are long white ſtrings, running vnder ground.

#### 4. *Phalangium ramoſum*. Branched Spider-wort.

The branched Spider-wort hath his leaues ſomewhat broader then the former, and of a more yellowiſh greene colour: the ſtalke hereof is diuerſly branched at the top, bearing many white flowers, like vnto the former, but ſmaller: the ſeedes and rootes are like the former in all things.

1 *Phalangium Alpobrogicum.* The Sauoye Spider-wort. 2 *Phalangium non ramosum.* Vnbranched Spider-wort. 3 *Phalangium ramosum.* Branched Spider-wort. 4 *Phalangium Ephemerum Virginianum.* Iohn Tradescante's Spider-wort.

### The Place.

The first groweth on the Hils neare vnto Sauoye, from whence diuers, allured with the beauty of the flower, haue brought it into these parts.

The second came vp in my Garden, from the seede receiued out of Italy. The others grow in Spaine, France, &c.

### The Time.

The vnbranched Spider-wort most commonly flowreth before all the other, and the branched a moneth after it: the other two about one time, that is, towards the end of May, and not much after the vnbranched kinde.

### The Names.

The first (as I said before) hath beene taken to be a white Lilliasphodill, and called *Liliasphodelus flore albo*; but Clusius hath more properly entituled it a *Phalangium*, and from the place of his originall, gaue him his other denomination, and so is called of most, as is set downe in the title.

The other haue no other names then are expressed in their titles, but only that Cordus calleth them *Liliago*; and Dodonæus, *lib. 4. hist. plant.* would make the branched kinde to bee *Moly alterum Plinij*, but without any good ground.

### The Vertues.

The names *Phalangium* and *Phalangites* were imposed on these plants, because they were found effectuall, to cure the poyson of that kinde of Spider, called *Phalangium*, as also of Scorpions and other Serpents. Wee doe not know, that any Physitian hath vsed them to any such, or any other purpose in our dayes.

## 5. *Phalangium Ephemerum Virginianum Ioannis Tradescant.*

The soon fading Spider-wort of Virginia, or Tradescant his Spider-wort.

This Spider-wort is of late knowledge, and for it the Christian world is indebted vnto that painfull industrious searcher, and louer of all natures varieties, Iohn Tradescant (sometimes belonging to the right Honourable Lord Robert Earle of Salisbury, Lord Treasurer of England in his time, and then vnto the right Honourable the Lord Wotton at Canterbury in Kent, and lastly vnto the late Duke of Buckingham) who first receiued it of a friend, that brought it out of Virginia, thinking it to bee the Silke Grasse that groweth there, and hath imparted hereof, as of many other things, both to me and others; the description whereof is as followeth:

From a stringie roote, creeping farre vnder ground, and rising vp againe in many places, springeth vp diuers heads of long folded leaues, of a grayish ouer-worne greene colour, two or three for the most part together, and not aboue, compassing one another at the bottome, and abiding greene in many places all the Winter; otherwhere perishing, and rising anew in the Spring, which leaues rise vp with the great round stalke, being set thereon at the ioynts, vsually but one at a ioynt, broad at the bottome where they compasse the stalke, and smaller and smaller to the end: at the vpper ioynt, which is the toppe of the stalke, there stand two or three such like leaues, but smaller, from among which breaketh out a dozen, sixteene, or twenty, or more round green heads, hanging downe their heads by little foot-stalkes, which when the flower beginneth to blow open, groweth longer, and standeth vpright, hauing three small pale greene leaues for a huske, and three other leaues within them for the flower, which lay themselues open flat, of a deepe blew purple colour, hauing an vmbone or small head in the middle, closely set about with six reddish, hairy, or feathered threeds, tipt with yellow pendents: this flower openeth it selfe in the day, & shutteth vsually at night,

night, and neuer openeth againe, but perisheth, and then hangeth downe his head againe; the greene huske of three leaues, closing it selfe againe into the forme of a head, but greater, as it was before, the middle vmbone growing to bee the seede vessell, wherein is contained small, blackish, long seede: Seldome shall any man see aboue one, or two at the most of these flowers blowne open at one time vpon the stalke, whereby it standeth in flowring a long time, before all the heads haue giuen out their flowers.

### The Place.

This plant groweth in some parts of Virginia, and was deliuered to Iohn Tradescant.

### The Time.

It flowreth from the end of May vntill Iuly, if it haue had greene leaues all the Winter, or otherwise, vntill the Winter checke his luxuriousnesse.

### The Names.

Vnto this plant I confesse I first imposed the name, by considering duely all the parts thereof, which vntill some can finde a more proper, I desire may still continue, and to call it *Ephemerum Virginianum Tradescanti*, Iohn Tradescante's Spider-wort of Virginia, or *Phalangium Ephemerum Virginianum*, The soone fading or Day Spider-wort of Virginia.

### The Vertues.

There hath not beene any tryall made of the properties since wee had it, nor doe we know whether the Indians haue any vse thereof.

## Chap. XVI.

### *Colchicum.* Medowe Saffron.

TO returne to the rest of the bulbous and tuberous rooted plants, that remaine to bee entreated of, the *Colchica* or Medowe Saffrons are first to bee handled, whereof these later dayes haue found out more varieties, then formerly were knowne; some flowring in the Spring, but the most in Autumne, and some bearing double, but the greatest part single flowers: whereof euery one in their order, and first of our owne Country kindes.

### 1. *Colchicum Anglicum album.* The white English Medowe Saffron.

It is common to all the Medowe Saffrons, except that of the Spring, and one other, to beare their flowers alone in Autumne or later, without any green leaues with them, and afterwards in February, their greene leaues: So that I shall not neede to make manie descriptions, but to shew you the differences that consist in the leaues, and colours of the flowers; and briefly to passe (after I haue giuen you a full description of the first) from one vnto another, touching onely those things that are note worthy. The white English Medowe Saffron then doth beare in Autumne three or foure flowers at the most, standing seuerally vpon weake foote-stalkes, a fingers length or more aboue the ground, made of six white leaues, somewhat long and narrow, and not so large as most of the other kindes, with some threeds or chiues in the middle, like vnto the Saffron flowers of the Spring, wherein there is no colour of Saffron, or vertue to that effect: after the flowers are past and gone, the leaues doe not presently follow, but the roote remaineth in the ground without shew of leafe aboue ground, most part of the Winter, and then in February there spring vp three or foure large and long greene

leaues,

leaues, when they are fully growne vp, standing on the toppe of a round, weake, green, and short foote-stalke, somewhat like the leaues of white Lillies, but not so large, and in the middest of these leaues, after they haue been vp some time, appeare two or three loose skinny heads, standing in the middle of the leaues vpon short, thicke, greene stalkes, and being ripe, conteine in them round small browniſh seede, that lye as it were loose therein, and when the head is dry, may bee heard to rattle being shaken: the roote is white within, but couered with a thicke blackish skinne or coate, hauing one side thereof at the bottome longer then the other, with an hollownesse also on the one side of that long eminence, where the flowers rise from the bottome, and shooting downe from thence a number of white fibres, whereby it is fastened in the ground: the greene leaues afterwards rising from the top or head of the roote.

2. *Colchicum Anglicum purpureum.* The purple Englisſh Medowe Saffron.

There is no difference at all in this Medowe Saffron from the former, but only in the colour of the flowers, which as they were wholly white in the former, so in this they are of a delayed purple colour, with a small shew of veines therein.

3. *Colchicum Pannonicum album.* The white Hungary Medowe Saffron.

The greatest difference in this *Colchicum* from the former Englisſh white one, is, that it is larger both in roote, leafe, and flower, and besides, hath more store of flowers together, and continuing longer in beauty, without fading so soone as the former, and are also somewhat of a fairer white colour.

4. *Colchicum Pannonicum purpureum.* The purple Hungary Medowe Saffron.

This purple Medowe Saffron is somewhat like vnto the white of this kinde, but that it beareth not so plentifully as the white, nor doth the roote grow so great; but the flowers are in a manner as large as they, and of the like pale delayed purple colour, or somewhat deeper, as is in the purple Englisſh, with some veines or markes vpon the flowers, making some shew of a checker on the out side, but not so conspicuous, as in the true checkerd kindes. Wee haue a kinde hereof is party coloured with white streakes and edges, which abide constant, and hath been raised from the seede of the former.

5. *Colchicum Byzantinum.* Medowe Saffron of Constantinople.

This Medowe Saffron of Constantinople hath his leaues so broad and large, that hardly could any that neuer saw it before, iudge it to be a *Colchicum*; for they are much larger then any Lilly leaues, and of a darke greene colour: the flowers are correspondent to the leaues, larger and more in number then in any of the former purple kindes, of the same colour with the last purple kinde, but of a little deeper purple on the inside, with diuers markes running through the flowers, like vnto it, or vnto checkers, but yet somewhat more apparantly: the roote is in the middle greater and rounder then the others, with a longer eminence, whereby it may easily bee knowne from all other sorts.

6. *Colchicum Lusitanicum Fritillaricum.* The checkerd Medowe Saffron of Portugall.

The flowers of this Medowe Saffron are larger and longer then the flowers of either the Englisſh or Hungarian, and almost as large as the last before mentioned, and of the same colour, but a little deeper, the spots and markes whereof are somewhat more easie to be seene euen a farre off, like vnto the flower of a Fritillaria, from whence it tooke his significatiue name: the leaues of this Medowe Saffron doe rise vp sooner then in any other of the Autumne kindes; for they are alwayes vp before Winter, and are foure or fiue in number, short rather then long, broad belowe, and pointed at the end, canaled or hollow, and standing round aboue the ground, one encompassing another at the bottome, like the great Spanish Starre Iacinth, called the Iacinth of Peru,

but

1 *Colchicum Pannonicum.* The Hungarian Medow Saffron. 2 *Colchicum Byzantinum.* Medowe Saffron of Conſtantinople. 3 *Colchicum Luſitanicum Fritillaricum.* The checkerd Medowe Saffron of Portugall. 4 *Colchicum Neapolitanum Fritillaricum.* The checkerd Medowe Saffron of Naples. 5 *Colchicum Fritillaricum Chienſe.* The checkerd Medowe Saffron of Chio or Sio. 6 *Colchicum Hermodactylum.* Phyſicall Medowe Saffron.

but shorter, and of a pale or grayish greene colour, differing from the colour of all the other Medowe Saffrons: the roote is like the roote of the English or Hungarian without any difference, but that it groweth somewhat greater. It is one of the first Medowe Saffrons that flower in the Autumne.

### 7. *Colchicum Neapolitanum Fritillaricum.*
### The checkerd Medowe Saffron of Naples.

This checkerd Medowe Saffron of Naples, is very like vnto the last recited checkerd Saffron of Portugall, but that the flower is somewhat larger, yet sometimes very little, or not at all: the greatest marke to distinguish them is, that the flowers of this are of a deeper colour, and so are the spots on the flowers likewise, which are so conspicuous, that they are discerned a great way off, more like vnto the flowers of a deepe Fritillaria, then the former, and make a goodlier and a more glorious shew: the leaues of this doe rise vp early after the flowers, and are somewhat longer, of a darker greene colour, yet bending to a grayish colour as the other, not lying so neatly or round, but stand vp one by another, being as it were folded together: neither of both these last named checkerd Medowe Saffrons haue giuen any seede in this Countrey, that euer I could learne or heare of, but are encreased by the roote, which in this is like the former, but a little bigger.

### 8. *Colchicum Fritillaricum Chiense.*
### The checkerd Medowe Saffron of Chio or Sio.

This most beautifull Saffron flower riseth vp with his flowers in the Autumne, as the others before specified doe, although not of so large a size, yet farre more pleasant and delightfull in the thicke, deepe blew, or purple coloured beautifull spots therein, which make it excell all others whatsoeuer: the leaues rise vp in the Spring, being smaller then the former, for the most part three in number, and of a paler or fresher greene colour, lying close vpon the ground, broad at the bottome, a little pointed at the end, and twining or folding themselues in and out at the edges, as if they were indented. I haue not seene any seede it hath borne: the roote is like vnto the others of this kinde, but small and long, and not so great: it flowreth later for the most part then any of the other, euen not vntill Nouember, and is very hard to be preserued with vs, in that for the most part the roote waxeth lesse and lesse euery yeare, our cold Country being so contrary vnto his naturall, that it will scarce shew his flower; yet when it flowreth any thing early, that it may haue any comfort of a warme Sunne, it is the glorie of all these kindes.

### 9. *Colchicum versicolor.* The party coloured Medowe Saffron.

The flowers of this Medowe Saffron most vsually doe not appeare, vntill most of the other Autumne sorts are past, except the last, which are very lowe, scarce rising with their stalkes three fingers breadth aboue the ground, but oftentimes halfe hid within the ground: the leaues whereof are smaller, shorter, and rounder, then in any of the other before specified, some being altogether white, and others wholly of a very pale purple, or flesh colour; and some againe parted, the one halfe of a leafe white, and the other halfe of the same purple, and sometimes striped purple and white, in diuers leaues of one and the same flower: and againe, some will be the most part of the leafe white, and the bottome purple, thus varying as nature list, that many times from one roote may be seene to arise all these varieties before mentioned: these flowers doe stand long before they fade and passe away; for I haue obserued in my Garden some that haue kept their flower faire vntill the beginning of Ianuary, vntill the extremitie of the Winter frosts and snowes haue made them hide their heads: the leaues therefore accordingly doe rise vp after all other, and are of a browniſh or darke greene colour at their first springing vp, which after grow to be of a deepe greene colour: the roote is like the former English or Hungarian kindes, but thicker and greater for the most part, and shorter also.

10. *Colchicum variegatum alterum.* Another party coloured Medowe Saffron.

There is another, whose party coloured flowers rise a little higher, diuersly striped and marked, with a deeper purple colour, and a pale or whitish blush throughout all the leaues of the flower.

11. *Colchicum montanum Hispanicum minus.* The little Spanish Medowe Saffron.

The flowers of this little Medowe Saffron are narrower and smaller then any of the former, and of a deeper reddish purple colour then either the English or Hungarian kindes: the greene leaues also are smaller then any other, lying on the ground, of a deepe or sad greene colour, rising vp within a while after the flowers are past, and doe abide greene all the Winter long: the roote is small and long, according to the rest of the plant, and like in forme to the others.

12. *Colchicum montanum minus versicolore flore.*
The small party coloured Medowe Saffron.

This little kinde differeth not from the Spanish kinde last set forth, but in the varietie of the flower, which is as small as the former, the three inner leaues being almost all white, and the three outer leaues some of them pale or blush, and some party coloured, with a little greene on the backe of some of them.

13. *Colchicum Hermodactilum.* Physicall Medowe Saffron.

This Physicall Medowe Saffron springeth vp with his leaues in Autumne, before his flowers appeare beyond the nature of all the former kindes, yet the flower doth, after they are vp, shew it selfe in the middle of the greene leaues, consisting of six white leaues, with diuers chiues in the middle, and passeth away without giuing any seede that euer I could obserue: the greene leaues abide all the Winter and Spring following, decaying about May, and appeare not vntill September, when (as I said) the flowers shew themselues presently after the leaues are sprung vp.

14. *Colchicum atropurpureum.* The darke purple Medowe Saffron.

The greatest difference in this kinde consisteth in the flower, which at the first appearing is as pale a purple, as the flower of the former Hungarian kinde: but after it hath stood in flower two or three dayes, it beginneth to change, and will after a while become to bee of a very deepe reddish purple colour, as also the little foote-stalke whereon it doth stand: the flower is of the bignesse of the Hungarian purple, and so is the greene leafe: the seede and roote is like the English purple kinde.

15. *Colchicum atropurpureum variegatum.*
The party coloured darke purple Medowe Saffron.

We haue of late gained another sort of this kinde, differing chiefly in the flower, which is diuersly striped thorough euery leafe of the flower, with a paler purple colour, whereby the flower is of great beauty: this might seeme to bee a degeneration from the former, yet it hath abiden constant with me diuers yeares, and giueth seede as plentifully as the former.

16. *Colchicum flore pleno.* Double flowred Medowe Saffron.

The double Medowe Saffron is in roote and leafe very like vnto the English kinde: the flowers are of a fine pale or delayed purple colour, consisting of many leaues set thicke together, which are somewhat smaller, as in the English flower, being narrow and long, and as it were round at the points, which make a very double flower, hauing

some chiues with their yellow tips, dispersed as it were among the leaues in the middle: it flowreth in September, a little after the first shew of the earlier Medowe Saffrons are past.

17. *Colchicum variegatum pleno flore.*
The party coloured double Medowe Saffron.

We haue another of these double kinds (if it be not the very same with the former, varying in the flower as nature pleaseth oftentimes; for I haue this flower in my garden, as I here set it forth, euery yeare) whose flowers are diuersified in the partition of the colours, as is to be seene in the single party coloured Medowe Saffron before described, hauing some leaues white, and others pale purple, and some leaues halfe white and halfe purple, diuersly set or placed in the double flower, which doth consist of as many leaues as the former, yet sometime this party coloured flower doth not shew it selfe double like the former, but hath two flowers, one rising out of another, making each of them to be almost but single flowers, consisting of eight or ten leaues a peece: but this diuersity is not constant; for the same roote that this yeare appeareth in that manner, the next yeare will returne to his former kinde of double flowers againe.

18. *Colchicum Vernum.* Medowe Saffron of the Spring.

This Medowe Saffron riseth vp very early in the yeare, that is, in the end of Ianuarie sometimes, or beginning, or at the furthest the middle of February, presently after the deepe Frosts and Snowes are past, with his flowers inclosed within three greene leaues, which opening themselues as soone almost as they are out of the ground, shew their buds for flowers within them very white oftentimes, before they open farre, and sometimes also purplish at their first appearing, which neuer shew aboue two at the most vpon one roote, and neuer rise aboue the leaues, nor the leaues much higher then they, while they last: the flower consisteth of six leaues, long and narrow, euery leafe being diuided, both at the bottome and toppe, each from other, and ioyned together onely in the middle, hauing also six chiues, tipt with yellow in the middle, euery chiue being ioyned to a leafe, of a pale red or deepe blush colour, when it hath stood a while blowne, and is a smaller flower then any Medowe Saffron, except the small Spanish kindes onely, but continueth in his beauty a good while, if the extremity of sharpe Frosts and Windes doe not spoile it: the leaues wherein these flowers are enclosed, at their first comming vp, are of a browni sh greene colour, which so abide for a while, especially on the outside, but on the inside they are hollow, and of a whitish or grayish greene colour, which after the flowers are past, grow to bee of the length of a mans longest finger, and narrow withall: there riseth vp likewise in the middle of them the head or seede vessell, which is smaller and shorter, and harder then any of the former, wherein is contained small round browne seede: the roote is small, somewhat like vnto the roores of the former, but shorter, and not hauing so long an eminence on the one side of the bottome.

19. *Colchicum Vernum atropurpureum.* Purple Medowe Saffron of the Spring.

The flower of this Medowe Saffron, is in the rising vp of his leaues and flowers together, and in all things else, like vnto the former, onely the flowers of this sort are at their first appearing of a deeper purple colour, and when they are blowne also are much deeper then the former, diuided in like manner, both at the bottome and toppe as the other, so that they seeme, like as if six loose leaues were ioyned in the middle part, to make one flower, and hath his small chiues tipt with yellow, cleauing in like manner to euery leafe.

The Place.

All these Medowe Saffrons, or the most part of them, haue their places expressed in their titles; for some grow in the fields and medowes of the champion grounds, others on the mountaines and hilly grounds. The English kindes grow in the West parts, as about Bathe, Bristow, Warmister,

1 *Colchicum montanum Hispanicum.* The little Spanish Medowe Saffron. 2 *Colchicum montanum minus versicolore flore.* The small party coloured Medowe Saffron. 3 *Colchicum versicolor.* The party coloured Medowe Saffron. 4 *Colchicum variegatum alterum.* Another party coloured Medowe Saffron. 5 *Colchicum atropurpureum.* The darke purple Medowe Saffron. 6 *Colchicum atropurpureum variegatum.* The variable darke purple Medowe Saffron. 7 *Colchicum vernum,* Medowe Saffron of the spring. 8 *Colchicum flore pleno.* Double Medowe Saffron.

ster, and other places also. The double kindes are thought to come out of Germany.

### The Time.

Their times likewise are declared in their seuerall descriptions: those that are earliest in Autumne, flower in August and September, the later in October, and the latest in the end of October, and in Nouember. The other are said to bee of the Spring, in regard they come after the deepe of Winter (which is most vsually in December and Ianuary) is past.

### The Names.

The generall name to all these plants is *Colchicum*, whereunto some haue added *Ephemerum*, because it killeth within one dayes space; and some *Strangulatorium*. Some haue called them also *Bulbus agrestis*, and *Filius ante Patrem*, The Sonne before the Father, because (as they thinke) it giueth seede before the flower: but that is without due consideration; for the root of this (as of most other bulbous plants) after the stalke of leaues and seede are dry, and past, may be transplanted, and then it beginneth to spring and giue flowers before leaues, (and therein onely it is differing from other plants) but the leaues and seede follow successiuely after the flowers, before it may be remoued againe; so that here is not seede before flowers, but contrarily flowers vpon the first planting or springing, and seede after, as in all other plants, though in a diuers manner.

The *Colchicum Hermodactilum* may seeme very likely to bee the *Colchicum Orientale* of Matthiolus, or the *Colchicum Alexandrinum* of Lobelius: And some thinke it to be the true *Hermodactilus*, and so call it, but it is not so. We doe generally call them all in English, Medowe Saffrons, or *Colchicum*, according to the Latine, giuing to euery one his other adiunct to know it by.

### The Vertues.

None of these are vsed for any Physicall respect, being generally held to be deadly, or dangerous at the least. Only the true Hermodactile (if it be of this tribe, and not this which is here expressed) is of great vse, for paines in the ioynts, and of the hippes, as the *Sciatica*, and the like, to be taken inwardly. Costæus in his Booke of the nature of plants, saith, that the rootes of our common kindes are very bitter in the Spring of the yeare, and sweet in Autumne, which Camerarius contradicteth, saying, that he found them bitter in Autumne, which were (as he saith) giuen by some impostors to diuers, as an antidote against the Plague.

## Chap. XVII.

### *Crocus.* Saffron.

THere are diuers sorts of Saffrons, whereof many doe flower in the Spring time, and some in Autumne, among whom there is but one onely kinde, that is called tame or of the Garden, which yeeldeth those blades or chiues that are vsed in meates and medicines, and many wayes profitable for other respects, none of the rest, which are all wilde kindes, giuing any blade equall vnto those of the tame kinde, or for any other vse, then in regard of their beautifull flowers of seuerall varieties, and as they haue been carefully sought out, and preserued by diuers, to furnish a Garden of dainty curiosity. To entreate therefore of these, I must, to obserue an orderly declaration, diuide them into two primary families: the former shall be of those that yeeld their pleasant flowers in the Spring of the yeare, and the other that send out their

their colours in the Autumne, among whom that *Rex pomarij* (as I may so call it) the tame or manured kinde, properly called of the Garden, is to be comprehended, for that it giueth his pleasant flowers at that time among others. I shall againe distribute those of the Spring time into three chiefe colours, that is, into white, purple, and yellow, and vnder euery one of them, comprehend the seuerall varieties that doe belong vnto them; which course I will also hold with those of the Autumne, that thus being rightly ranked, they may the more orderly be described.

### 1. *Crocus Vernus albus purus minor.*
The smaller pure white Saffron flower of the Spring.

This small Saffron flower springeth vp in the beginning of the yeare, with three or foure small greene leaues, somewhat broader, but much shorter then the true Saffron leaues, with a white line downe the middle of euery leafe: betweene these leaues, out of a white skinne, riseth vp one or two small flowers, made of six leaues a peece, as all the rest in generall are, of a pure white colour, without any mixture in it, which abide not in flower aboue a weeke, or rather lesse, so sodainly is the pleasure of this, and the purple lost: it flowreth not for the most part, vntill a moneth after the yellow Crocus appeareth in flower, and the ordinary stript Crocus is past: the seede is small, round, and reddish, yet not so red as the seede of the yellow, contained in three square heads, yet seldome beareth, but encreaseth by the roote plentifully enough, which is small, round, and flat at the bottome, somewhat white on the outside, but whiter within, shooting out small sprouts on euery side of the roote, which is the best note to know this kinde and the lesser purple, which are both alike, from all other rootes of Saffron flowers.

### 2. *Crocus albus maior multiflorus.* The great snow white Crocus.

This greater Saffron flower riseth vp vsually with three or foure greene leaues, larger then the former, with a white line in euery one of them: the flowers are greater, and more in number, rising together, but flowring one after another, of a pure snow white colour, and abiding but little longer in flower then the former.

### 3. *Crocus albus maior alter dictus Mæsiacus.*
The great white Saffron flower or Crocus of Mesia.

This great white Crocus of Mesia, riseth vp out of the ground, almost as early as the first sort of the yellow, with foure or fiue leaues, being very like vnto the leaues of the yellow Crocus, and as large, with white lines in them: the flowers also are as large as the flowers of the yellow, and many also rising one after another like vnto it, but not of so pure a white colour, as the former or last described, but rather tending to a Milky or Creame colour: the roote is not couered with any reddish, but rather pale skinnes or coates.

### 4. *Crocus albus Mæsiacus fundo violaceo.*
The great white Crocus of Mesia with a blew bottome.

There is another of this kinde, like vnto the former in all things, sauing that the bottomes of the flowers of this kinde, with some part of the stalke next the flower, are of a pale shining purple colour, and rising vp a pretty way into the flower; whereas another also of this kind, hath a little shew or marke of blew, and not purple, at the bottome of the flower onely, which maketh a difference.

### 5. *Crocus albus fundo purpureo.* The white Crocus with a purple bottome.

This Saffron flower is of the same kinde with the first, both in roote, leafe, and flower, in none of them differing from it, but in that the bottome of this flower, with that part of the short foote-stalke next vnto it, is of a violet or purple colour, and sometimes hauing here and there some purple small lines, or spots on the white leaues: it flowreth also with the first white, or somewhat later.

*6. Crocus vernus albus ſtriatus.* The white ſtript Crocus.

This ſtript Saffron flower is likewiſe neare the ſame firſt kind, or firſt white Crocus, hauing the like leaues and flowers, ſomewhat larger, but as ſoone fading almoſt as it: but herein this flower differeth, that it hath pale blewiſh lines and ſpots in all the leaues thereof, and more principally in the three outer leaues: the root is alſo white on the outſide, like the firſt white, but greater, with young ones growing round about it.

*7. Crocus vernus albus polyanthos verſicolor.*
The greater party coloured white Crocus.

The greater party coloured Saffron flower, hath his greene leaues like vnto the ſecond great white Crocus before mentioned, hauing more flowers then any of the former, except the firſt great white, the leaues whereof haue greater ſtripes then the laſt recited Crocus, but of a purple Violet colour, making each leafe ſeeme oftentimes to haue as much purple as white in them: the roote hereof is ſomewhat like the ſecond white, but of a little more duskie colour on the outſide, and not budding out on the ſides at all, or very little.

*8. Crocus vernus albus verſicolor.* The leſſer party coloured white Crocus.

The leaues and flowers of this other party coloured Crocus, are for bigneſſe in a manner equall with the laſt, but hath not ſo many flowers riſing together from the roote: the flower is finely marked with blew ſtrakes on the white flower, but nothing ſo much as in the former: the roote alſo is like the laſt.

*9. Crocus Epiſcopalis.* The Biſhops Crocus.

This party coloured or Biſhops Saffron flower, is very like both in leaues and rootes vnto the Neapolitane blew Crocus, but ſomewhat greater: the flowers doe abide not ſo long time blowne, and hath all the leaues either wholly white, with blew ſtripes on both ſides of them, or wholly of a fine delayed blew Violet colour, and the three innermoſt more blew and finely ſtriped, both on the inſide and outſide of them, and ſometimes it hath been ſeen to haue three leaues white, and three leaues of a pale blew.

10. *Crocus vernus ſtriatus vulgaris.* The ordinary ſtript Crocus.

There is another ſort of ſtript Saffron flower, which is moſt common and plentifull in moſt Gardens, which I muſt needes bring vnder the ranke of theſe white kinds, although it differre very notably, both in roote, leafe, and flower, from all of them: the leaues of this riſe vp ſooner then the yellow or white Crocus, lying ſpread vpon the ground for the moſt part, but narrower then any of the former: among theſe leaues ſpring vp diuers flowers, almoſt as large as the former great white Crocus, of a very bleake or pale purple colour, tending to white on the inſide, and in many almoſt white; with ſome ſmall whitiſh chiues tipt with yellow in the middle: the three outer leaues are of a yellowiſh white colour on the backe ſide of them, ſtript euery one of them with three broad ſtripes, of a darke murrey or purple colour, and a little ſprinkled with ſome ſmall purple lines, on both ſides of thoſe ſtripes; but on the inſide, of the ſame pale purple or white colour with the reſt: the ſeede hereof is ſomewhat darker coloured then of the white, and is more liberall in bearing: the roote is differing from all the former, being rounder and bigger then any of them, except the kindes of Miſia, yet ſomewhat flat withall, not hauing any ſhootes from the ſides, but ſetting off into rootes plentifully, hauing a round circle compaſſing the bottome of the roote, which eaſily falleth away, when it is taken vp out of the ground, and couered with a browne coate, ſomewhat neare the colour of the yellow Crocus, but not altogether ſo bright: it flowreth vſually the firſt of all theſe ſorts, or with the firſt of the early yellowes.



11. *Crocus*

1 *Crocus vernus albus minor.* The small white Saffron flower of the spring. 2 *Crocus vernus Mæsiacus albus.* The great white Crocus of Misia. 3 *Crocus vernus albus striatus.* The white stript Crocus. 4 *Crocus vernus albus polyanthos versicolor* The party coloured white Crocus. 5 *Crocus albus fundo purpureo.* The white Crocus with a purple bottome. 6 *Crocus vernus Neapolitanus.* The great blew Crocus of Naples. 7 *Crocus vernus purpureus maximus.* The great purple Crocus. 8 *Crocus vernus purpureus striatus.* The purple stript Crocus. 9 *Crocus vernus purpureus Capillarifolio.* The purple Crocus with small leaues. 10 *Crocus vernus flavus striatus.* The yellow stript Crocus. 11 *Crocus vernus intus versicolor.* The cloth of gold Crocus.

11. *Crocus vernus striatus Turcicus.* The Turkie stript Crocus.

There is another of this kinde, whose flower is a little larger, and of a deeper purple colour, both on the inside and outside; the greene leafe also is bigger, and of a more whitish colour.

12. *Crocus vernus Capillarifolio albus.* The white Crocus with small leaues.

This white Crocus is in all things like vnto the purple of the same kinde, but that the flower of this is wholly white: the full description therefore hereof, you shall haue in that purple with small leaues, of this kinde hereafter set downe, whereunto I referre you.

13. *Crocus vernus purpureus minor.* The smaller purple Crocus.

The smaller purple Saffron flower of the Spring, hath his greene leaues so like vnto the first white flowred Saffron, that they can hardly be distinguished, onely they seem to bee a little narrower: the flower is also much about the same bignesse, or a little bigger, and seldome beareth aboue one flower from a roote, euen as the first doth, of a deepe purple Violet colour, the bottome of the flower, with the vpper part of the stalke next thereunto, being of a deeper or blacker purple; in the middle of the flower are some pale chiues tipt with yellow pendents, and a longer pointell, diuided or forked at the toppe: the roote of this is in all things so like vnto the first white, that it is impossible for the most cunning and conuersant in them, to know the one from the other. This beareth seede very sparingly, as the white doth, and is reddish like vnto it, but recompenseth that defect with a plentifull encrease by the roote: it likewise flowreth at the very same time with the white, and endureth as small a time.

14. *Crocus vernus purpureus maximus.* The greatest purple Crocus.

This great purple Crocus is of the same kinde with the next described, as well in roote as leafe, but greater; for the greene leaues hereof are the greatest and broadest of all other Crocus, with a large white line in the middle of euery one: it springeth vp much later then the former, and doth not shew his flower vntill the other bee past a good while: the flowers also are the largest of all these Crocus of the Spring time, and equalling, if not surpassing that purple kinde that flowreth in Autumne, hereafter set forth, of a very faire and deepe Violet colour, almost as deepe as the former: the seed vessels are large also and white, wherein is contained pale reddish seede, like vnto the next blew kinde, but somewhat greater: the roote is (as I said before) like vnto the next, that is, flat and round, with a duskie coloured outside, whose head for springing in it is as hardly discerned.

*Alter Apicibus albidis.* We haue one of this kinde, the toppes onely of whose purple flower are whitish, for the breadth of halfe the naile of a mans hand, which abideth constant euery yeare in that manner, and therefore is a difference fit to be remembred.

15. *Crocus vernus Neapolitanus siue caeruleus maior.*
The greater blew Crocus of Naples.

This great blew Crocus riseth vp with diuers greene leaues, broader then any of the former (except the last) with a white line running downe the inside of euery leafe, as in the former, among which riseth vp, out of diuers great long white skinnes, diuers large flowers, but not fully so great as the former, consisting of six leaues, of a paler blew or Violet colour then in the former, hauing in the middle of the flowers a few pale threeds, tipt with yellow, and a longer pointell of a gold yellow colour, forked or diuided at the toppe, smelling sweeter then in the former, and abiding a great while longer, being in flower vsually euen with the stript yellow Crocus', or before the former purple, and yeelding more plenty of seede: the roote hereof is not very great, but a little darke on the outside, being round and flat withall, that one can hardly know which is the vpperside thereof.

This

This kinde differeth very little from the former, either in roote, leafe, or flower, for the bigneſſe or colour, but that it ſeemeth to be a little bleaker or paler blew, becauſe it flowreth a little earlier. *Crocus Neapolitanus praecocior.*

16. *Crocus vernus purpureus ſtriatus.* The ſtript purple Crocus.

The leaues of this ſtript purple Saffron flower, are as large and broad as the laſt, or rather a little longer: the flowers alſo are as plentifull, and as large, of a fine delayed purple colour on the outſide, with three broad ſtrakes or lines downe the backe of the three outer leaues, and of a little deeper purple on the inſide, as the other three leaues are alſo of a deeper purple colour, and are ſtriped with the ſame deepe purple about the ground, or bottome of the leaues: this ſometimes yeeldeth three ſquare heads, containing in them browniſh ſeede: the roote is like vnto the laſt, and flowreth much about the time of the former.

17. *Crocus vernus purpurens verſicolor.* The ſiluer ſtript purple Crocus.

This ſtript Saffron flower, is in leaues and flowers ſomewhat like vnto the laſt ſtript purple, but a little ſmaller: the flowers are of a little deeper purple through the whole leaues, ſtriped with white lines, both on the leaues, and towards the edges, which maketh a peculiar difference from all the reſt: the roote of this is not ſo flat, though like it, and couered with a darke aſh coloured skinne: it flowreth about the ſame time.

18. *Crocus purpureus flammeus maior.* The greater purple flame coloured Crocus.

The greene leaues of this Crocus or Saffron flower, are of a reaſonable breadth and length, and of a pleaſant freſh greenneſſe, with a faire broad white line downe the middle of them, but riſing not out of the ground ſo early as the next deſcribed Crocus: the flowers are likewiſe of a meane bigneſſe, of a pale purple on the outſide, ſomewhat whitiſh, eſpecially the three outer leaues; but on the inſide of a deeper purple, and ſtriped with great ſtripes like flames, hauing ſome chiues in the middle, and a longer one alſo feathered a little at the toppe: the roote is white on the outſide, ſomewhat flat and round, but not ſo flat as the Neapolitane Crocus before deſcribed.

19. *Crocus purpureus flammeus minor.* The leſſer purple flame coloured Crocus.

This Crocus hath almoſt as broad and long greene leaues as the former, and of the ſame verdure, which riſe vp earlier then it, and is in flower likewiſe ſomewhat before it, being ſmaller for ſize by a little, but of as deepe a purple on the outſide, as on the inſide, flamed with faire broad ſtripes from the middle of the leaues, or ſomewhat lower vnto the edges: each of theſe giue ſeed that is of a pale reddiſh colour: the roote is very like vnto the former, but a little leſſer.

20. *Crocus vernus purpureus Capillariſolio.* The purple Crocus with ſmall leaues.

This ſmall kinde of Saffron flower riſeth out of the ground, with two or three long and ſmall green leaues, very like vnto the leaues of the fine Fether-Graſſe hereafter deſcribed, ſtanding vpright at the firſt, but afterwards lying vpon the ground; among which come the flowers, ſometimes three, but moſt vſually two vpon one ſtalke, if the roote be not young, which then will beare but one on a ſtalke, which is very ſhort, ſo that the flowers ſcarce ariſe aboue the ground, yet laying themſelues open in the day time, if it be faire, and the Sunne doe ſhine, otherwiſe they keepe cloſe, and doe not open at all: and after one flower is paſt, which doth not laſt abcue three or foure dayes at the moſt, the others follow, which are of a bleake blewiſh purple in the middle of the flower, and of a deeper purple towards the ends or points of the leaues, but of a more ſullen or darke purple on the outſide of them, and yellowiſh at the bottome, with ſome yellow chiues in the middle: the ſeede is ſmall and darker coloured then any of the former Crocus, contained alſo in ſmaller heads, ſtanding one by another

vpon

vpon the ſame ſhort foote-ſtalke, which then riſeth vp a little higher, ſhewing the maner of the ſtanding of the flowers, which in their flowring time could not ſo eaſily bee diſcerned: the roote is very ſmall and round, hauing one ſide at the bottome lower then the other, very like the roote of a *Colchicum* or Medowe Saffron, and ſomewhat neare reſembling alſo the hoofe of an horſe foote, couered with a very thicke skinne, of a darke or blackiſh browne colour: this flowreth the laſt of all the former ſorts of Saffron flowers, euen when they are all paſt.

### 21. *Crocus vernus purpureus ſtriatus Capillarifolio.* The ſtript purple Crocus with ſmall leaues.

This ſmall ſtript purple Saffron flower hath ſuch like leaues, as the laſt deſcribed hath, betweene which riſeth the flower vpon as ſhort a foote-ſtalke, conſiſting of ſix leaues like the former, of a faire purple colour on the outſide of the three outer leaues, with three lines or ſtrakes downe euery leafe, of a deeper purple colour, and on the inſide of a paler purple, as the other three leaues are alſo, with ſome chiues tipt with yellow pendents, and a forked pointell in the middle: the roote of this is ſomewhat bigger then the former, and rounder, but couered with as thicke and as browne a skinne: it flowreth about the ſame time with the former.

### 22. *Crocus vernus luteus ſiue Maſiacus.* The yellow Crocus.

The yellow Crocus or Saffron flower, riſeth vp with three or foure leaues out of the ground, being ſomewhat neare the breadth of the great purple kindes, with a white line in them, as in moſt of the reſt: the flowers ſtand in the middle of theſe leaues, and are very large, of a gold yellow colour, with ſome chiues, and a forked point in the middle: the ſeede hereof is of a brighter colour then in any of the other: the roote is great and round, as great or greater then a Wall Nut ſometimes, and couered with reddiſh skinnes or coates, yeelding more ſtore of flowers then moſt of the former, and beginning to blowe with the firſt ſorts, or preſently after, but outlaſt many of them, and are of a pleaſant good ſent.

*Flore aureo.* Of this kinde we haue ſome, whoſe flowers are of a deeper gold yellow colour then others, ſo that they appeare reddiſh withall.

*Flore pallido.* And we haue alſo another ſort, whoſe flowers are very pale, betweene a white and a yellow, not differing in any thing elſe.

*Flore viridante luteo.* And another ſmaller, whoſe flower hath a ſhew of greenneſſe in the yellow, and more greene at the bottome.

### 23. *Crocus vernus flauus ſtriatus.* The yellow ſtript Crocus.

This kinde of yellow ſtript Crocus or Saffron flower, riſeth vp with more ſtore of narrower and greener leaues then the former, and after the leaues are ſpread, there riſe vp many yellow flowers from among them, which are not of ſo faire and bright a yellow colour, but more dead and ſullen, hauing on the backſide of each of the three outermoſt leaues, three ſmall ſtripes, of an ouer-worne or dull purple colour, with ſome chiues and a pointell in the middle: the roote of this kinde, is very like the roote of the former yellow, but ſomewhat ſmaller and ſhorter, and couered with the like reddiſh skinnes, but a little ſadder: it flowreth not ſo early as the former yellow, but abideth almoſt as long as it.

### 24. *Crocus vernus luteus verſicolor primus.* The beſt cloth of gold Crocus.

The faireſt cloth of gold Crocus or Saffron flower, riſeth vp very early, euen with the firſt, or the firſt of all other Crocus, with three or foure very narrow and ſhort leaues, of a whiter colour then any of the former, which by and by after doe ſhew forth the flowers, riſing from among them out of the ſame white skinne, which includeth the leaues, but are not ſo plentifull as the former yellow, being but two or three at the moſt, of a faire gold yellow colour, yet ſomewhat paler then the firſt, hauing

uing on the backe of euery of the three outer leaues, three faire and great stripes, of a faire deepe purple colour, with some small lines at the sides or edges of those purple stripes; on the inside of these flowers, there is no signe or shew of any line or spot, but wholly of a faire gold yellow, with chiues and a fethertopt pointell in the middle: the seede hereof is like the former, but not so red: the roote of this kinde is easily knowne from the roote of any other Saffron flower, because the outer peelings or shels being hard, are as it were netted on the outside, hauing certaine ribbes, rising vp higher then the rest of the skinnes, diuided in the forme of a net-worke, of a darke browne colour, and is smaller and rounder then the former yellow, and not encreasing so plentifully by the roote.

25. *Crocus vernus luteus versicolor alter.*
The second cloth of gold, or Duke Crocus.

There is no difference either in roote, leafe, or colour of flower, or time of flowring in this sort from the last before mentioned; for the flower of this is of the same bignesse and colour, the only note of difference is in the marking of the three outer leaues, which haue not three stripes like the former, but are wholly of the same deepe purple colour on the backe of them, sauing that the edges of them are yellow, which is the forme of a Duke Tulipa, and from thence it tooke the name of a Duke Crocus.

26. *Crocus vernus versicolor pallidelutens.* The pale cloth of gold Crocus.

We haue a third sort of this kinde of cloth of gold Crocus, which hath leaues and flowers like the former, but differeth in this, that the colour of the flower is of a paler yellow by much, but stript in the same manner as the first, but with a fainter purple colour: the roote also is netted like them, to shew that this is but a variation of the same kinde.

27. *Crocus vernus versicolor albidoluteus.* The cloth of siluer Crocus.

The chiefest note of difference in this Saffron flower is, that being as large a flower as any of the former of this kinde, it is of so pale a yellowish white, that it is more white then yellow, which some doe call a butter colour: the three outer leaues are striped on the backe of them, with a paler purple blew shining colour, the bottome of the flower, and the vpper part of the stalke, being of the same purple blew colour: the roote of this is also netted as the other, to shew it is a variety of the same kinde.

And thus much for those Saffron flowers that come in the Spring time; now to those that flower in Autumne onely: and first of the true Saffron.

1. *Crocus verus sativus Autumnalis.* The true Saffron.

The true Saffron that is vsed in meates and medicines, shooteth out his narrow long greene leaues first, and after a while the flowers in the middle of them appeare about the end of August, in September and October, according to the soile, and climate where they growe; these flowers are as large as any of the other former or later sorts, composed of six leaues a peece, of a murrey or reddish purple colour, hauing a shew of blew in them: in the middle of these flowers there are some small yellow chiues standing vpright, which are as vnprofitable, as the chiues in any other of the wilde Saffrons, before or hereafter specified; but besides these, each flower hath two, three, or foure greater and longer chiues, hanging downe vpon or betweene the leaues, which are of a fierie red colour, and are the true blades of Saffron, which are vsed physically or otherwise, and no other: All these blades being pickt from the seuerall flowers, are laid and prest together into cakes, and afterwards dryed very warily on a Kill to preserue them; as they are to be seene in the shops where they are sold. I neuer heard that euer it gaue seede with any: the roote groweth often to be as great, or greater then a green Wall Nut, with the outer shell on it, couered with a grayish or ash-coloured skin, which breaketh into long hairie threeds, otherwise then in any other roote of Crocus.

2. *Crocus Byzantinus argenteus.* The siluer coloured Autumne Crocus.

This Saffron flower springeth vp in October, and seldome before, with three or foure short greene leaues at the first, but growing longer afterwards, and in the midst of them, presently after they haue appeared, one flower for the most part, and seldome two, consisting of six leaues, the three outermost whereof are somewhat larger then the other three within, and are of a pale bleake blew colour, almost white, which many call a siluer colour, the three innermost being of a purer white, with some yellow chiues in the middle, and a longer pointell ragged or fethered at the toppe: this very seldome beareth seede, but when the yeare falleth out to bee very milde; it is small, round, and of a darke colour: the roote is pretty bigge, and rounder then any other Crocus, without any flat bottome, and couered with a darke russet skinne.

3. *Crocus Pyrenæus purpureus.* The purple mountaine Crocus.

This purple Saffron flower of the Autumne, riseth vp but with one flower vsually, yet sometimes with two one after another, without any leaues at all, in September, or sometimes in August, standing vpon a longer foote-stalke then any kinde of Saffron flower, either of the Spring or Autumne, and is as large as the flower of the greatest purple Saffron flower of the Spring, of a very deepe Violet purple colour, which decayeth after it hath stood blowne three or foure dayes, and becommeth more pale, hauing in the middle some yellow chiues, and a long fether topt pointell, branched, and rising sometimes aboue the edges of the flowers: about a moneth after the flowers are past, and sometimes not vntill the first of the Spring, there riseth vp three or foure long and broad greene leaues, with a white line in euery one of them, like vnto the first purple Vernall kindes, which abide vntill the end of May or Iune: the roote is small and white on the outside, so like vnto the roote of the lesser Vernall purple or white Crocus, that it cannot be distinguished, vntill about the end of August, when it doth begin to shoot, and then by the early shooting vp a long white sprout for flower, it may be knowne. I neuer could obserue it to giue any seede, the Winter (as I thinke) comming on it so quickly after the flowring, being the cause to hinder it.

4. *Crocus montanus Autumnalis.* The Autumne mountaine Crocus.

The mountaine Saffron flower springeth vp later then any of the former, and doth not appeare vntill the middle or end of October, when all the flowers of the former are past, appearing first with three or foure short greene leaues, like vnto the Byzantine Crocus, and afterwards the flowers betweene them, which are of a pale or bleake blew tending to a purple, the foote-stalkes of them being so short, that they scarce appeare aboue ground at the first, but after two or three dayes they grow a little higher: the roote is very great and flat bottomed, couered with a grayish duskie coate or skinne, and encreaseth very little or seldome.

## The Place.

The seuerall places of these Saffron flowers, are in part set downe in their titles; the others haue beene found out, some in one Countrey, and some in another, as the small purple and white, and stript white in Spaine: the yellow in Mesia about Belgrade, the great purple in Italy; and now by such friends helpes as haue sent them, they prosper as well in our Gardens, as in their naturall places. Yet I must giue you this to vnderstand, that some of these formerly expressed, haue been raised vp vnto vs by the sowing of their seede.

## The Time.

Their seuerall times are likewise expressed in their descriptions; for some shew forth their pleasant flowers in the Spring, wherein for the three first moneths,

1 *Crocus vernus luteus vulgaris.* The common yellow spring Crocus. 2 *Crocus vernus sativus Autumnalis.* The true Saffron. 3 *Crocus Byzantinus argenteus.* The siluer coloured Autumne Crocus. 4 *Crocus Pyrenæus purpureus.* The purple mountaine Crocus. 5 *Crocus montanus Autumnalis,* The Autumne mountaine Crocus. 6 *Sisyrinchium maius,* The greater Spanish Nut.

moneths, our Gardens are furnished with the varietie of one sort or another: the rest in Autumne, that so they might procure the more delight, in yeelding their beauty both early and late, when scarce any other flowers are found to adorne them.

### The Names.

I shall not neede to trouble you with an idle tale of the name of Crocus, which were to little purpose, nor to reiterate the former names imposed vpon them; let it suffice that the fitteſt names are giuen them, that may distinguish them one from another; onely this I must giue you to vnderstand, that the gold yellow *Crocus* or Saffron flower, is the true *Crocus Mæsiacus*, as I shewed before; and that neither the yellow stript, or cloth of gold (which wee so call after the Dutch name *Gaud Laken*) is the true *Mæsiacus*, as some suppose; and that the great white Saffron flower, by reason of his likenesse vnto the gold yellow, is called *Crocus albus Mæsiaci facie*, or *facie lutei*, that is, The white Saffron flower that is like the *Mæsiacus* or yellow.

### The Vertues.

The true Saffron (for the others are of no vse) which wee call Englisſh Saffron, is of very great vse both for inward and outward diseases, and is very cordiall, vsed to expell any hurtfull or venemous vapours from the heart, both in the small Pockes, Measels, Plague, Iaundise, and many other diseases, as also to strengthen and comfort any cold or weake members.

## Chap. XVIII.

### *Sisyrinchium.* The Spanish Nut.

I Can doe no otherwise then make a peculiar Chapter of this plant, because it is neither a *Crocus*, although in the roote it come somewhat neare vnto that kinde that is netted; but in no other part agreeing with any the delineaments of a Saffron flower, and therefore could not be thrust into the Chapter amongst them: neither can I place it in the forefront of the Chapter of the *Iris bulbosa*, or bulbous Flowerdeluces, because it doth not belong to that Family: and although the flower thereof doth most resemble a Flowerdeluce, yet in that no other parts thereof doe fitly agree thereunto, I haue rather chosen to seate it by it selfe betweene them both, as partaking of both natures, and so may serue in stead of a bridge, to passe from the one to the other, that is, from the *Crocus* or Saffron flower, to the *Iris bulbosa* or bulbous rooted Flowerdeluce, which shall follow in the next Chapter by themselues.

The Spanish Nut hath two long and narrow, soft and smooth greene leaues, lying for the most part vpon the ground, and sometimes standing vp, yet bending downewards; betweene these leaues riseth vp a small stalke, halfe a foote high, hauing diuers smooth soft greene leaues vpon it, as if they were skinnes, through which the stalke passeth; at the toppe whereof stand diuers flowers, rising one after another, and not all flowring at once: for seldome shall you haue aboue one flower blowne at a time, each whereof doth so quickly passe and fade away, that one may well say, that it is but one dayes flower, or rather the flower of a few houres: the flower it selfe hath nine leaues, like vnto a Flowerdeluce, whereof the three that fall downe, haue in each of them a yellow spot: the other three, which in the Flowerdeluces are hollow and ridged, couering the other three that fall downe, in this stand vpright, and are parted at the ends: the three that stand vp in the middle are small and short: the whole flower is smaller then any Flowerdeluce, but of sundry colours; for some are of an excellent skie colour blew, others of a Violet purple, others of a darker purple colour, and some white, and many others mixed, either pale blew and deepe purple, or white and blew

mixed

mixed or striped together very variably, quickly fading as I said before: the seede is enclosed in small cods, so thinne and transparent, that one may easily see, and tell the seeds as they lye, which are of a brownish red colour: the roote is small, blackish and round, wrapped in a thicke skinne or huske, made like vnto a net, or somewhat like vnto the roote of the cloth of gold Crocus: when the plant is in flower, it is found to haue two rootes one aboue another, whereof the vppermost is firme and sound, and the vndermost loose and spongie, in like manner as is found in the rootes of diuers Orchides or Satyrions, Bee-flowers and the like, and without any good taste, or sweetnesse at all, although Clusius saith otherwise.

*Sisyrinchium Mauritanicum.* The Barbary Nut.

There is another of this kinde, not differing from the former in any other notable part, but in the flower, which in this is of a delayed purplish red colour, hauing in each of the three lower leaues a white spot, in stead of the yellow in the former, but are as soone fading as they.

The Place.

The former doe grow very plentifully in many parts both of Spaine and Portugall, where Guillaume Boel, a Dutch man heretofore remembred often in this Booke, found them; of the sundry colours specified, whereas Clusius maketh mention but of one colour that he found.

The other was found in that part of Barbary, where Fez and Morocco do stand, and brought first into the Lowe-Countries: but they are both very tender, and will hardly abide the hard Winters of these colder regions.

The Time.

The first flowreth in May and Iune, the last not vntill August.

The Names.

The name *Sisyrinchium* is generally imposed vpon this plant, by all authors that haue written thereof, thinking it to bee the right *Sisyrinchium* of Theophrastus: but concerning the Spanish name *Nozelha*, which Clusius saith it is called by in Spaine, I haue beene credibly enformed by the aforenamed Boel, that this roote is not so called in those parts; but that the small or common stript Crocus is called *Nozelha*, which is sweete in taste, and desired very greedily by the Shepheards and Children, and that the roote of this *Sisyrinchium* or Spanish Nut, is without any taste, and is not eaten. And againe, that there is not two kindes, although it grow greater, and with more flowers, in those places that are neare the Sea, where both the washing of the Sea water, and the moisture and ayre of the Sea, causeth the ground to bee more fertile. This I thought good, from the true relation of a friend, to giue the world to vnderstand, that truth might expell errour.

The Vertues.

These haue not been knowne to bee vsed to any Physicall purpose, but wholly neglected, vnlesse some may eate them, as Clusius reporteth.

## Chap. XIX.

### *Iris bulbosa.* The bulbous Flowerdeluce.

THe Flowerdeluces that haue bulbous rootes are of two sorts, the one greater then the other: the greater bearing larger and broader leaues and flowers, and the lesser narrower. But before I giue you the descriptions of the vsuall greater kindes, I must needes place one or two in the fore-front that haue no fellowes; the one is called of Clusius, his broad leafed Flowerdeluce, and the other a Persian, somewhat like vnto it, which although they differ notably from the rest, yet they haue the neareſt resemblance vnto those greater kindes, that come next after them.

### *Iris bulbosa prima latifolia Clusij.*
Clusius his first great bulbous Flowerdeluce.

This Flowerdeluce hath diuers long and broad leaues, not stiffe, like all the other, but soft and greenish on the vpperside, and whitish vnderneath; among which rise vp sometimes seuerall small, short, slender stalkes, and sometimes but one, not aboue halfe a foote high, bearing at the top one flower a peece, somewhat like vnto a Flowerdeluce, consisting of nine leaues, whereof those three that stand vpright, are shorter and more closed together, then in other sorts of Flowerdeluces; the other three that fall downe, turne vp their ends a little, and those three, that in other Flowerdeluces doe couer them at the bottome, stand like the vpright leaues of other Flowerdeluces, but are parted into two ends, like vnto two small eares: the whole flower is of a faire blew, or pale skie colour in most, with a long stripe in the middle of each of the three falling leaues, and in some white, but more seldome: the roote is reasonable great, round and white, vnder the blackish coates wherewith it is couered, hauing many long thicke white rootes in stead of fibres, which make them seeme to be Asphodill rootes. The flower is very sweete.

### *Iris bulbosa Persica.* The Persian bulbous Flowerdeluce.

This Persian Flowerdeluce is somewhat like vnto the former, both in roote and in leafe, but that the leaues are shorter and narrower, and the flower being much about the same fashion, is of a pale blew russetish colour, each of the three lower falling leaues are almost wholly of a browne purple colour, with a yellow spot in the middle of them: this as it is very rare, so it seldome beareth flowers with vs.

#### The Place.

The first groweth in many places of Spaine and Portugall, from whence I and others haue often had it for our Gardens, but by reason of the tendernesse thereof, it doth hardly endure the sharpnesse of our cold Winters, vnlesse it be carefully preserued.

The other is said to come from Persia, and therefore it is so entituled, and is as tender to be kept as the other.

#### The Time.

The first flowreth most vsually not vntill May with vs, yet many times sooner: but in Ianuary and February, as Clusius saith, in the naturall places thereof.

The other is as early oftentimes when it doth flower with vs.

#### The Names.

Because Clusius by good iudgement referreth the first to the greater kindes

kindes of Flowerdeluces, and placeth it in the fore ranke, calling it *Iris bulbosa latifolia prima*, that is, The first broad leafed Flowerdeluce, and all others doe the like, I haue (as you see) in the like manner put it before all the other, and keepe the same name. The Spaniards, as he saith, called it *Lirio espadanal*, and they of Corduba, *Lirios azules*.

The other hath no other name then as it is in the title.

1. *Iris bulbosa maior sive Anglica cærulea.*

The blew Englisn bulbous Flowerdeluce.

This bulbous Flowerdeluce riseth vp early, euen in Ianuary oftentimes, with fiue or six long and (narrow, in comparison of any great breadth, but in regard of the other kinde) broad whitish green leaues, crested or straked on the backside, and halfe round, the inside being hollow like a trough or gutter, white all along the inside of the leafe, and blunt at the end; among which riseth vp a stiffe round stalke, a cubit or two foot high, at the toppe whereof, out of a skinnie huske, commeth forth one or two flowers, consisting of nine leaues a peece, three whereof that are turned downewards, are larger and broader then the other, hauing in each of them a yellow spot, about the middle of the leafe, other three are small, hollow, ridged or arched, couering the lower part next the stalke of those falling leaues, turning vp their ends, which are diuided into two parts, other three stand vpright, and are very small at the bottome of them, and broader toward the toppe: the whole flowet is of a faire blew colour; after the flowers are past, come vp three square heads, somewhat long, and lanke, or loose, containing in them round yellowish seede, which when it is ripe, will rattle by the shaking of the winde in the dry huskes: the roote of this kinde is greater and longer then any of the smaller kindes with narrow leaues, couered with diuers browne skinnes, which seeme to be fraught with long threeds like haires, especially at the small or vpper end of the roote, which thing you shall not finde in any of the smaller kindes.

2. *Iris bulbosa maior purpurea & purpuro violacea.*

The paler or deeper purple great bulbous Flowerdeluce.

These purple Flowerdeluces differ not from the last described, either in roote or leafe: the chiefest difference consisteth in the flowers, which in these are somewhat larger then in the former, and in the one of a deepe blew or Violet purple colour, and in the other of a deepe purple colour, in all other things alike.

There is also another, in all other things like vnto the former, but only in the flower, which is of a pale or bleake blew, which we call an ash-colour. *Flore cinereo.*

3. *Iris bulbosa maior purpurea variegata sive striata.*

The great purple stript bulbous Flowerdeluce.

There is another of the purple kinde, whose flower is purple, but with some veines or stripes of a deeper Violet colour, diuersly running through the whole leaues of the flower.

And another of that bleake blew or ash-colour, with lines and veines of purple in the leaues of the flowers, some more or lesse then other. *Flore cinereo striata purpureo.*

And againe another, whose flower is of a purple colour like vnto the second, but that round about that yellow spot, in the middle of each of the three falling leaues (as is vsuall in all the bulbous Flowerdeluces) there is a circle of a pale blew or ash-colour, the rest of the leafe remaining purple, as the other parts of the flower is. *Flore purpureo orbe cinereo.*

4. *Iris bulbosa maior flore rubente.*

The great peach coloured bulbous Flowerdeluce.

There is another of these greater kindes, more rare then any of the former, not differing in roote, leafe, or flower, from the former, but onely that the flower in this is of a pale reddish purple colour, comming somewhat neare vnto the colour of a peach blossome.

### 5. *Iris bulbosa maior sive latifolia alba.*
### The great white bulbous Flowerdeluce.

The great white bulbous Flowerdeluce, riseth not vp so early out of the ground as the blew or purple doth, but about a moneth or more after, whose leaues are somewhat larger, and broader then of the others: the stalke is thicker and shorter, bearing vsually two very large and great flowers, one flowring a little before the other, yet oftentimes both in flower together in the end, of a bleake blewish white colour, which wee call a siluer colour, while they are in the budde, and before they be blowne open, but then of a purer white, yet with an eye or shew of that siluer colour remaining in them, the three falling leaues being very large, and hauing that yellow spot in the middle of each of them: the seedes are likewise included in heads, like vnto the blew or purple kindes, but larger, and are of a reddish yellow colour like them: the roote likewise is not differing, but greater.

### 6. *Iris bulbosa maior alba variegata.*
### The great white stript bulbous Flowerdeluce.

This white stript Flowerdeluce, is in roote, leafe, and flower, and in manner of growing, like vnto the former white Flowerdeluce; the onely difference is in the marking of the flower, being diuers from it: for this hath in the white flower great veines, stripes, or markes, of a Violet blew colour, dispersed through the leaues of the flower very variably, which addeth a superexcellent beauty to the flower.

### 7. *Iris bulbosa maior sive latifolia versicolor.*
### The great party coloured bulbous Flowerdeluce.

There is no difference in this from the former, but in the flower, which is of a whitish colour in the three falling leaues, hauing a circle of ash-colour about the yellow spot, the three rigged leaues being likewise whitish, but ridged and edged with that ash-colour, and the three vpright leaues of a pale blewish white colour, with some veines therein of a blewish purple.

*Varietas.* There hath beene brought vnto vs diuers rootes of these kindes, with the dryed flowers remaining on them, wherein there hath beene seene more varieties, then I can well remember to expresse, which variety it is very probable, hath risen by the sowing of the seeds, as is truely obserued in the narrower leafed kinde of Flowerdeluce, in the Tulipa, and in some other plants.

*Flore luteo.* Wee haue heard of one of this kinde of broad leafed Flowerdeluces, that should beare a yellow flower, in the like manner as is to be seene in the narrow leafed ones: but I haue not seene any such, and therefore I dare report no further of it, vntill time hath discouered the truth or falshood of the report.

### The Place.

Lobelius is the first reporter, that the blew Flowerdeluce or first kinde of these broad leafed Flowerdeluces, groweth naturally in the West parts of England; but I am in some doubt of the truth of that report: for I rather thinke, that some in their trauels through Spaine, or other parts where it groweth, being delighted with the beauty of the flower, did gather the rootes, and bring them ouer with them, and dwelling in some of the West parts of England, planted them, and there encreasing so plentifully as they doe, they were imparted to many, thereby in time growing common in all Countrey folkes Gardens thereabouts. They grow also, and all the other, and many more varieties, about Thoulouse, from whence Plantinianus Gassanus both sent and brought vs them, with many other bulbous rootes, and rare plants gathered thereabouts.

1 *Iris bulbosa latifolia prima Clusii.* Clusius his first great bulbous Flowerdeluce. 2 *Iris bulbosa maior cærulea sive Anglica.* The great blew or English bulbous Flowerdeluce. 3 *Iris bulbosa maior purpureo variegata.* The great purple striped bulbous Flowerdeluce. 4 *Iris bulbosa angustifolia maior alba.* The greater white narrow leafed bulbous Flowerdeluce. 5 *Iris bulbosa angustifolia versicolor.* The party coloured narrow leafed bulbous Flowerdeluce. 6 *Iris bulbosa angustifolia Africana.* The purple African bulbous narrow leafed Flowerdeluce.

**The Time.**

These doe flower vsually in the end of May, or beginning of Iune, and their seede is ripe in the end of Iuly or August.

**The Names.**

Lobel calleth the first English blew Flowerdeluce, *Hyacinthus Poetarum flore Iridis, & propter Hyacinthinum colorem, id est violaceum dictus*: but I know not any great good ground for it, more then the very colour; for it is neither of the forme of a Lilly, neither hath it those mourning markes imprinted in it, which the Poet faineth to bee in his Hyacinth. It is most truely called an *Iris*, or Flowerdeluce (and there is great difference betweene a Lilly and a Flowerdeluce, for the formes of their flowers) because it answereth thereunto very exactly, for the flower, and is therefore called vsually by most, either *Iris bulbosa Anglica*, or *Iris bulbosa maior sive latifolia*, for a difference betweene it, and the lesser with narrow leaues: In English, eyther The great English bulbous Flowerdeluce, or the great broad leafed bulbous Flowerdeluce, which you will, adding the other name, according to the colour.

And thus much for these broad leafed bulbous Flowerdeluces, so much as hath come to our knowledge. Now to the seuerall varieties of the narrow leafed bulbous Flowerdeluces, so much likewise as we haue been acquainted with.

*Iris bulbosa minor sive angustifolia alba.*
The smaller white or narrow leafed bulbous Flowerdeluce.

This first Flowerdeluce, which beareth the smaller flower of the two white ones, that are here to bee described, springeth out of the ground alwaies before Winter, which after breaketh forth into foure or fiue small and narrow leaues, a foote long or more, of a whitish greene on the inside, which is hollow and chanalled, and of a blewish greene colour on the outside, and round withall: the stalke of this kinde is longer and slenderer then the former, with some shorter leaues vpon it, at the toppe whereof, out of short skinny leaues, stand one or two flowers, smaller, shorter, and rounder then the flowers of the former broad leafed Flowerdeluces, but made after the same proportion with nine leaues, three falling downewards, with a yellow spot in the middle, other three are made like a long arch, which couer the lower part next the stalke of those falling leaues, and turne vp at the ends of them, where they are diuided into two parts: the other three stand vpright, betweene each of the three falling leaues, being somewhat long and narrow: the flower is wholly (sauing the yellow spot) of a pure white colour, yet in some hauing a shew of some blew throughout, and in others towards the bottome of the three vpright leaues: after the flowers are past, there rise vp so many long cods or seede vessels, as there were flowers, which are longer and smaller then in the former, and a little bending like a Cornet, with three round squares, and round pointed also, which diuiding it selfe when the seede is ripe into three parts, doe shew six seuerall cells or places, wherein is contained such like round reddish yellow seedes, but smaller then the former: the roote is smaller and shorter then the former, and without any haires or threeds, couered with browne thin skinnes, and more plentifull in giuing encrease.

*Iris bulbosa angustifolia alba flore maiore.*
The greater white narrow leafed bulbous Flowerdeluce.

I shall not neede to make a seuerall description to euery one of these Flowerdeluces that follow, for that were but to make often repetition of one thing, which being once done, as it is, may well serue to expresse all the rest, and but onely to adde the especiall

ciall differences, either in leafe or flower, for bigneſſe, colour, or forme, as is expedient to expreſſe and diſtinguiſh them ſeuerally. This greater white bulbous Flowerdeluce is like vnto the laſt deſcribed in all parts, ſauing that it is a little larger and higher, both in leafe, ſtalke, and flower, and much whiter then any of theſe mixed ſorts that follow, yet not ſo white as the former: the roote hereof is likewiſe a little bigger and rounder in the middle.

*Albeſcens.* Milke white.

There is another, whoſe falling leaues haue a little ſhew of yellowneſſe in them, and ſo are the middle ridges of the arched leaues, but the vpright leaues are more white, not differing in roote or leafe from the firſt white.

*Argentea.* Siluer colour.

And another, whoſe falls are of a yellowiſh white, like the laſt, the arched leaues are whiter, and the vpright leaues of a blewiſh white, which we call a ſiluer colour.

*Albida.* Whitiſh.

Another hath the fals yellowiſh, and ſometimes with a little edge of white about them, and ſometimes without; the vpright leaues are whitiſh, as the arched leaues are, yet the ridge yellower.

*Albida labris luteis.* White with yellow fals.

Another hath his fals yellow, and the vpright leaues white, all theſe flowers are about the ſame bigneſſe with the firſt.

*Albida angustior.* The narrow white.

But we haue another, whoſe flower is ſmaller, and almoſt as white as the ſecond, the lower leaues are ſmall, and doe as it were ſtand outright, not hauing almoſt any fal at all, ſo that the yellow ſpot ſeemeth to be the whole leafe, the arched leaues are not halfe ſo large as in the former, and the vpright leaues bowe themſelues in the middle, ſo that the tops doe as it were meete together.

And another of the ſame, whoſe falling leaues are a little more eminent and yellow, with a yellower ſpot.

*Aurea ſiue lutea Hiſpanica.* The Spaniſh yellow.

We haue another kinde that is called the Spaniſh yellow, which riſeth not vp ſo high, as ordinarily moſt of the reſt doe, and is wholly of a gold yellow colour.

*Pallide lutea.* Straw colour.

There is another, that vſually riſeth higher then the former yellow, and is wholly of a pale yellow, but deeper at the ſpot.

*Albida lutea.* Pale Straw colour.

There is alſo another like vnto the pale yellow, but that the falling leaues are whiter then all the reſt of the flower.

*Mauritanica flaua ſirotina minor.* The ſmall Barbary yellow.

There is a ſmaller or dwarfe kinde, brought from the backe parts of Barbary, neare the Sea, like vnto the yellow, but ſmaller and lower, and in ſtead of vpright leaues, hath ſmall ſhort leaues like haires: it flowreth very late, after all others haue almoſt giuen their ſeede.

*Verſicolor Hiſpanica cærulea labris albis.* The party coloured Spaniſh

We haue another ſort is called the party coloured Spaniſh bulbous Flowerdeluce, whoſe falling leaues are white, the arched leaues of a whitiſh ſiluer colour, and the vpright leaues of a fine blewiſh purple.

*Diuerſitas.* The diuerſity or variation of this flower.

Yet ſometimes this doth vary; for the falling leaues will haue either an edge of blew, circling the white leaues, the arched leaues being a little blewer, and the vpright leaues more purple.

Or the fals will be almoſt wholly blew, edged with a blewer colour, the arched leaues pale blew, and the vpright leaues of a purpliſh blew Violet colour.

Or the fals white, the arched leaues pale white, as the vpright leaues are.

Or not of ſo faire a blewiſh purple, as the firſt ſort is.

Some of them alſo will haue larger flowers then others, and be more liberall in bearing flowers: for the firſt ſort, which is the moſt ordinary, ſeldome beareth aboue one flower on a ſtalke, yet ſometimes two. And of the others there are ſome that wil beare vſually two and three flowers, yet ſome againe will beare but one. All theſe kindes ſmell ſweeter then many of the other, although the moſt part be without ſent.

*cærulea ſiue purpurea minor Luſitanica præcox.* The ſmall early purple Portugall.

There is another kinde, that is ſmaller in all the parts thereof then the former, the ſtalke is ſlender, and not ſo high, bearing at the toppe one or two ſmall flowers, all wholly of a faire blewiſh purple, with a yellow ſpot in

in euery one of the three falling leaues, this vsually flowreth early, euen with the first bulbous Flowerdeluces.

*Purpurea maior.* The greater purple,

We haue another purple, whose flower is larger, and stalke higher, and is of a very reddish purple colour, a little aboue the ground, at the foote or bottome of the leaues and stalke: this flowreth with the later sort of Flowerdeluces.

*Purpurea serotina* The late purple.

There is another, whose flower is wholly purple, except the yellow spot, and flowreth later then any of the other purples.

*Purpura rubescens labrio caruleo.* A reddish purple with blew fals.

There is yet another purple, whose vpright leaues are of a reddish purple, and the falling leaues of a blew colour.

*Purpura rubescens labrio albido caruleo.* A reddish purple with whitish blew fals.

And another of a reddish purple, whose falling leaues are of a whitish blew colour, in nothing else differing from the last.

*Purpurea labrio luteo.* Party coloured purple & yellow

Another hath his falling leaues of a faire gold yellow, without any stripe, yet in some there are veines running through the yellow leaues, and some haue an edge of a sullen darke colour about them: the vpright leaues in euery of these, are of a Violet purple.

*Purpurea labrio ex albido caruleo & luteo mixtis.* Party coloured purple with stript yellow fals.

Another is altogether like this last, but that the falling leaues are of a pale blew and yellow, trauersing one the other, and the arched leaues of a pale purplish colour.

*Subpurpurea labrio luteo.* Pale purple with yellow fals.

Another hath his vpright leaues of a paler purple, and the falling leaues yellow.

A paler purple.

And another little differing from it, but that the arched leaues are whitish.

*Subcaerulea labrio luteo.* Party coloured blew and yellow

Another whose vpright leaues are of a pale blew, and the falling leaues yellow.

*Crinio coloris elegantioris.* A faire haire colour.

And another of the same sort, but of a little paler blew.

*Altera obsoletior.* A dull haire colour.

We haue another sort, whose vpright leaues are of a faire browniish yellow colour, which some call a *Fuille mort*, and others an haire colour; the falling leaues yellow.

And another of the same colour, but somewhat deader.

*Iris bulbosa Africana serpentariæ caule.*

## The purple or murrey bulbous Barbary Flowerdeluce.

This Flowerdeluce as it is more strange (that is, but lately knowne and possessed by a few) so it is both more desired, and of more beauty then others. It is in all respects, of roote, leafe, and flower, for the forme like vnto the middle sort of these Flowerdeluces, onely the lowest part of the leaues and stalke, for an inch or thereabouts, next vnto the ground, are of a reddish colour, spotted with many spots, and the flower, being of a meane size, is of a deepe purplish red or murrey colour the whole flower throughout, except the yellow spot in the middle of the three lower or falling leaues, as is in all others.

*Purpura cærulea obsoleta labris fuscis.* The duskie party coloured purple,

And lastly, there is another sort, which is the greatest of all these narrow leafed Flowerdeluces, in all the parts of it; for the roote is greater then any of the other, being thicke and short: the leaues are broader and longer, but of the same colour: the stalke is stronger and higher then any of them, bearing two or three flowers, larger also then any of the rest, whose falling leaues are of a duskie yellow, and sometimes with veines and borders about the brimmes, of another dunne colour, yet hauing that yellow spot that is in all: the arched leaues are of a sullen pale purplish yellow, and the vpright leaues of a dull or duskie blewish purple colour: the heads or hornes for seede are likewise greater, and so is the seede also a little.

## The Place.

These Flowerdeluces haue had their originall out of Spaine and Portugall, as it is thought, except those that haue risen by the sowing, and those which are named of Africa.

The

### The Time.

These flower in Iune, and sometimes abide vnto Iuly, but vsually not so early as the former broad leafed kindes, and are soone spoiled with wet in their flowring.

### The Names.

The seuerall names, both in Latine and English, are sufficient for them as they are set downe; for we know no better.

### The Vertues.

There is not any thing extant or to be heard, that any of these kindes of Flowerdeluces hath been vsed to any Physicall purposes, and serue onely to decke vp the Gardens of the curious.

And thus much for these sorts of bulbous Flowerdeluces, and yet I doubt not, but that there are many differences, which haue risen by the sowing of the seede, as many may obserue from their owne labours, for that euery yeare doth shew forth some variety that is not seene before. And now I will conuert my discourse a while likewise, to passe through the seuerall rankes of the other kindes of tuberous rooted Flowerdeluces, called Flagges.

## Chap. XX.

*Iris latifolia tuberosa.* The Flagge or Flowerdeluce.

THere are two principall kindes of tuberous or knobby rooted Flowerdeluces, that is, the tall and the dwarfe, or the greater and the lesser; the former called *Iris maior* or *latifolia*, and the other *Iris minor*, or rather *Chamæiris*; and each of these haue their lesser or narrow leafed kindes to bee comprehended vnder them: Of all which in their order. And first of that Flowerdeluce, which for his excellent beautie and raritie, deserueth the first place.

*Iris Chalcedonica sive Susiana maior.* The great Turkie Flowerdeluce.

The great Turkie Flowerdeluce, hath diuers heads of long and broad fresh greene leaues, yet not so broad as many other of those that follow, one folded within another at the bottome, as all other of these Flowerdeluces are: from the middle of some one of those heads (for euery head of leaues beareth not a flower) riseth vp a round stiffe stalke, two foote high, at the toppe whereof standeth one flower (for I neuer obserued it to beare two) the largest almost, but rarest of all the rest, consisting of nine leaues, like the others that follow, but of the colour almost of a Snakes skinne, it is so diuersly spotted; for the three lower falling leaues are very large, of a deepe or darke purple colour, almost blacke, full of grayish spots, strakes, and lines through the whole leaues, with a blacke thrume or freeze in the middle of each of them: the three arched leaues that couer them, are of the same darke purple colour, yet a little paler at the sides, the three vpper leaues are very large also, and of the same colour with the lower leaues, but a little more liuely and fresh, being speckled and straked with whiter spots and lines; which leaues being laid in water, will colour the water into a Violet colour, but if a little Allome be put therein, and then wrung or pressed, and the iuice of these leaues dryed in the shadow, will giue a colour almost as deepe as Indico, and may serue for shadowes in limming excellent well: the flower hath no sent that can be perceiued, but is onely commendable for the beauty and rarity thereof: it seldome beareth seedes in these cold Countries, but when it doth, it is contained in great heads,

being

being browniſh and round, but not ſo flat as in other ſorts, the roots are more browne on the outſide, and growing tuberous thicke, as all other that are kept in Gardens.

*Iris Chalcedonica ſiue Suſiana minor.* The leſſer Turkie Flowerdeluce.

There is another hereof little differing, but that the leafe is of a more yellowiſh greene colour, and the flower neither ſo large or faire, nor of ſo perſpicuous markes and ſpots, nor the colour of that liuely (though darke) luſtre.

The Place.

Theſe haue been ſent out of Turkie diuers times among other things, and it ſhould ſeeme, that they haue had their originall from about Sufis, a chiefe Citie of Perſia.

The Time.

They flower in May moſt vſually, before any of the other kindes.

The Names.

They haue been ſent vnto vs, and vnto diuers other in other parts, from Conſtantinople vnder the name of *Alaia Suſiana*, and thereupon it hath been called, both of them and vs, either *Iris Chalcedonica*, or *Suſiana*, and for diſtinction *maior* or *minor*: In Engliſh, The Turkie Flowerdeluce, or the Ginnie Hen Flowerdeluce, the greater or the leſſer.

*Iris alba Florentina.* The white Flowerdeluce.

The great white Flowerdeluce, hath many heads of very broad and flat long leaues, encloſing or folding one within another at the bottome, and after a little diuided one from another toward the top, thin edged, like a ſword on both ſides, and thicker in the middle: from the middle of ſome of theſe heads of leaues, riſeth vp a round ſtiffe ſtalk, two or three foot high, bearing at the top one, two, or three large flowers, out of ſeuerall huskes or skins, conſiſting or nine leaues, as all the other do, of a faire white colour, hauing in the middle of each of the three falling leaues, a ſmall long yellow frize or thrume, as is moſt vſuall in all the ſorts of the following Flowerdeluces, both of the greater and ſmaller kindes: after the flowers are paſt, come the ſeed, incloſed in thicke ſhort pods, full traught or ſtored with red roundiſh and flat ſeede, lying cloſe one vpon another: the roote is tuberous or knobby, ſhooting out from euery ſide ſuch like tuberous heads, lying for the moſt part vpon or aboue the ground, and faſtened within the ground with long white ſtrings or fibres, which hold them ſtrongly, and encreaſeth faſt. There is another like vnto this laſt in all things, ſauing that the colour of the flower is of a more yellowiſh white, which we vſually call a Straw colour.

*Flore pallido.*

*Iris alba maior Verſicolor.* The white party coloured Flowerdeluce.

This variable Flowerdeluce is like vnto the former, but that the leaues are not ſo large and broad, the flower hereof is as large almoſt, and as white as the former, but it hath a faire liſt or line of a blewiſh purple downe the backe of euery one of the three vpright leaues, and likewiſe round about the edges, both of the vpper and lower leaues, and alſo a little more purpliſh vpon the ridge of the arched leaues, that couer the falling leaues: the roote hereof is not ſo great as of the former white, but a little ſlenderer and browner.

*Iris Dalmatica maior.* The great Dalmatian Flowerdeluce.

This greater Flowerdeluce of Dalmatia, hath his leaues as large and broad as any of the Flowerdeluces whatſoeuer, his ſtalke and flower doe equall his other proportion,

tion, onely the colour of the flower is differing, being of a faire watchet or bleake blew colour wholly, with the yellow frize or thrum downe the middle of the lower or falling leaues, as before is said to be common to all these sorts of Flowerdeluces, in all other parts it little differeth, sauing onely this is obserued to haue a small shew of a purplish red about the bottome of the greene leaues.

*Iris purpurea siue vulgaris.* The common purple Flowerdeluce.

This Flowerdeluce, which is most common in Gardens, differeth nothing at all from those that are formerly described, either in roote, leafe, or flower for the forme of them, but onely that the leaues of this are not so large as the last, and the flower it selfe is of a deep purple or Violet colour, and sometimes a little declining to rednesse, especially in some places.

Sometimes this kinde of Flowerdeluce will haue flowers of a paler purple colour, comming neare vnto a blew, and sometimes it will haue veines or stripes of a deeper blew, or purple, or ash colour, running through all the vpper and lower leaues. *Purpurea pallidior versicolor.*

There is another like vnto this, but more purple in the fals, and more pale in the vpright leaues. *Cærulea labris purpureis.*

*Iris Asiatica cærulea.* The blew Flowerdeluce of Asia.

This Flowerdeluce of Asia, is in largenesse of leaues like vnto the Dalmatian, but beareth more store of flowers on seuerall branches, which are of a deeper blew colour, and the arched leaues whitish on the side, and purplish on the ridges, but in other things like vnto it.

There is another neare vnto this, but that his leaues are a little narrower, and his flowers a little more purple, especially the vpper leaues. *Purpurea.*

*Iris Damascena.* The Flowerdeluce of Damasco.

This is likewise altogether like the Flowerdeluce of Asia, but that it hath some white veines in the vpright leaues.

*Iris Lusitanica biflora.* The Portugall Flowerdeluce.

This Portugall Flowerdeluce is very like the common purple Flowerdeluce, but that this is not so large in leaues, or flowers, and that it doth often flower twice in a yeare, that is, both in the Spring, and in the Autumne againe, and besides, the flowers haue a better or sweeter sent, but of the like purple or Violet colour as it is, and comming forth out of purplish skins or huskes.

*Iris Camerarij siue purpurea versicolor maior.*
The greater variable coloured purple Flowerdeluce.

The greater of the variable purple Flowerdeluces, hath very broad leaues, like vnto the leaues of the common purple Flowerdeluce, and so is the flower also, but differing in colour, for the three lower leaues are of a deepe purple colour tending to rednesse, the three arched leaues are of the colour with the vpper leaues, which are of a pale or bleake colour tending to yellownesse, shadowed ouer with a smoakie purplish colour, except the ridges of the arched leaues, which are of a more liuely purple colour.

*Iris purpurea versicolor minor.* The lesser variable purple Flowerdeluce.

This Flowerdeluce differeth not in any thing from the last, but onely that it hath narrower greene leaues, and smaller and narrower flowers, else if they be both conferred together, the colours will not seeme to varie the one from the other any whit at all.

There is another somewhat neare vnto these two last kindes, whose huskes from whence *Altera minus fuliginea.*

whence the flowers doe shoote forth, haue purple veines in them, and so haue the falling purplish leaues, and the three vpright leaues are not so smoakie, yet of a dun purple colour.

*Iris cærulea versicolor.* The blew party coloured Flowerdeluce.

This party coloured Flowerdeluce hath his leaues of the same largenesse, with the lesser variable purple Flowerdeluce last described, and his flowers diuersly marked: for some haue the tals blew at the edges, and whitish at the bottome, the arched leaues of a yellowish white, and the vpright leaues of a whitish blew, with yellowish edges. Some againe are of a darker blew, with browniſh spots in them. And some are so pale a blew, that we may well call it an ash-colour: And lastly, there is another of this sort, whose vpright leaues are of a faire pale blew, with yellowish edges, and the falling leaues parted into two colours, sometimes equally in the halfe, each side suitable to the other in colour: And sometimes hauing the one leafe in that manner: And sometimes but with a diuers coloured list in them; in the other parts both of flower and leafe, like vnto the other.

*Iris lutea variegata.* The yellow variable Flowerdeluce.

This yellow variable Flowerdeluce loseth his leaues in Winter, contrary to all the former Flowerdeluces, so that his roote remaineth vnder ground without any shew of leafe vpon it: but in the beginning of the Spring it shooteth out faire broad leaues, falling downwards at the points or ends, but shorter many times then any of the former, and so is the stalke likewise, not rising much aboue a foote high, whereon are set two or three large flowers, whose falling leaues are of a reddish purple colour, the three that stand vpright of a smoakie yellow, the arched leaues hauing their ridges of a bleake colour tending to purple, the sides being of the former smoakie yellow colour, with some purplish veines at the foote or bottome of all the leaues: the roote groweth somewhat more slender and long vnder ground, and of a darker colour then manie of the other.

Another sort hath the vpright leaues of a reasonable faire yellow, and stand more vpright, not bowing downe as most of the other, and the purple fals haue pale edges.

*Varietas.* Some haue their greene leaues party coloured, white and greene, more or lesse, and so are the huskes of the flowers, the arched leaues yellow, as the vpright leaues are, with purplish veines at the bottome. And some haue both the arched and vpright leaues of so pale a yellow, that we may almost call it a straw colour, but yellower at the bottome, with purple veines, and the falling leaues purple, with two purple spots in them.

And these are the sorts of the greater tuberous or Flagge Flowerdeluces that haue come to our knowledge: the next hereunto are the lesser or narrow leafed kindes to be described; and first of the greatest of them.

1. *Iris angustifolia Tripolitana aurea.* The yellow Flowerdeluce of Tripoly.

This Flowerdeluce I place in the forefront of the narrow leafed Flowerdeluces, for the length of the leaues, compared with the breadth of them; it may fitly bee called a narrow leafed Flowerdeluce, although they be an inch broad, which is broader then any of them that follow, or some of those are set downe before, but as I said, the length make them seem narrow, and therfore let it take vp his roome in this place, with the description that followeth. It beareth leaues a yard long, or not much lesse, and an inch broad, as is said before, or more, of a sad greene colour, but not shining: the stalke riseth vp to be foure or fiue foote high, being strong and round, but not very great, bearing at the toppe two or three long and narrow gold yellow flowers, of the fashion of the bulbous Flowerdeluces, as the next to bee described is, without any mixture or variation therein: the heads for seede are three square, containing within them many flat cornered seedes: the roote is long and blackish, like vnto the rest that follow, but greater and fuller.

1 *Iris Chalcedonica sive Susiana maior.* The great Turkie Flowerdeluce. 2 *Iris alba Florentina.* The white Flowerdeluce. 3 *Iris latifolia variegata.* The variable Flowerdeluce. 4 *Chamæiris latifolia major.* The greater dwarfe Flowerdeluce.

2. *Iris anguſtifolia maior cærulea.*
The greater blew Flowerdeluce with narrow leaues.

This kinde of Flowerdeluce hath his leaues very long and narrow, of a whitiſh greene colour, but neither ſo long or broad as the laſt, yet broader, thicker and ſtiffer then any of the reſt with narrow leaues that follow: the ſtalke riſeth ſometimes no higher then the leaues, and ſometimes a little higher, bearing diuers flowers at the top, ſucceſſiuely flowring one after another, and are like vnto the flowers of the bulbous Flowerdeluces, but of a light blew colour, and ſometimes deeper: after the flowers are paſt, riſe vp ſix cornered heads, which open into three parts, wherein is contained browne ſeede, almoſt round: the roote is ſmall, blackiſh and hard, ſpreading into many long heads, and more cloſely growing or matting together.

3. *Iris anguſtifolia purpurea marina.* The purple narrow leafed Sea Flowerdeluce.

This Sea Flowerdeluce hath many narrow hard leaues as long as the former, and of a darke greene colour, which doe ſmell a little ſtrong: the ſtalke beareth two or three flowers like the former, but ſomewhat leſſe, and of a darke purple or Violet colour: in ſeede and roote it is like the former.

4. *Iris anguſtifolia purpurea verſicolor.*
The variable purple narrow leafed Flowerdeluce.

The leaues of this Flowerdeluce are very like the former Sea Flowerdeluce, and do a little ſtinke like them; the flowers are differing, in that the vpper leaues are wholly purple or violet, and the lower leaues haue white veines, and purple running one among another: the ſeede and rootes differ not from the former purple Sea kinde.

5. *Iris anguſtifolia minor Pannonica ſiue verſicolor Cluſij.*
The ſmall variable Hungarian Flowerdeluce of Cluſius.

This Hungarian Flowerdeluce (firſt found out by Cluſius, by him deſcribed, and of him tooke the name) riſeth vp with diuers ſmall tufts of leaues, very long, narrow, and greene, growing thicke together, eſpecially if it abide any time in a place; among which riſeth vp many long round ſtalkes, higher then the leaues, bearing two or three, or foure ſmall flowers, one aboue another, like the former, but ſmaller and of greater beauty: for the lower leaues are variably ſtriped with white and purple, without any thrume or fringe at all; the vpper leaues are of a blewiſh fine purple or Violet colour, & ſo are the arched leaues, yet hauing the edges a little paler: the heads for ſeede are ſmaller, and not ſo cornered as the other, containing ſeedes much like the former, but ſmaller: the roote is blacke and ſmall, growing thicker and cloſer together then any other, and ſtrongly faſtened in the ground, with a number of hard ſtringie rootes: the flowers are of a reaſonable good ſent.

6. *Iris anguſtifolia maior flore duplici.* The greater double blew Flowerdeluce.

This Flowerdeluce, differeth not either in roote or leafe from the firſt great blew Flowerdeluce of Cluſius, but onely in that the leaues grow thicker together, and that the flowers of this kinde are as it were double with many leaues confuſedly ſet together, without any diſtinct parts of a Flowerdeluce, and of a faire blew colour with many white veines and lines running in the leaues; yet oftentimes the ſtalke of flowers hath but two or three ſmall flowers diſtinctly ſet together, riſing as it were out of one huske.

7. *Iris anguſtifolia minor alba Cluſij.*
The ſmall white Flowerdeluce of Hungary.

This likewiſe differeth little from the former Hungarian Flowerdeluce of Cluſius, but

1 *Iris angustifolia Tripolitana.* The yellow Flowerdeluce of Tripoli. 2 *Iris angustifolia maior carulea.* The greater blew Flowerdeluce with narrow leaues. 3 *Iris angustifolia minor Pannonica siue versicolor Clusii.* The small variable Hungarian Flowerdeluce of Clusius. 4 *Iris angustifolia maior flore aupliçi* The greater double blew Flowerdeluce. 5 *Chamairis angustifolia minor.* The lesser Grasse Flowerdeluce. 6 *Iris tuberosa.* The veluet Flowerdeluce.

but that the leafe is of a little paler greene colour, and the flower is of a faire whitish colour, with some purple at the bottome of the leaues.

Next after these narrow leafed Flowerdeluces, are the greater and smaller sorts of dwarfe kindes to follow; and lastly, the narrow or grasse leafed dwarfe kindes, which will finish this Chapter of Flowerdeluces.

1. *Chamæiris latifolia maior alba.* The greater white dwarfe Flowerdeluce.

This dwarfe Flowerdeluce hath his leaues as broad as some of the lesser kindes last mentioned, but not shorter; the stalke is very short, not aboue halfe a foote high or thereabouts, bearing most commonly but one flower, seldome two, which are in some of a pure white, in others paler, or somewhat yellowish through the whole flower, except the yellow frize or thrume in the middle of euery one of the falling leaues: after the flowers are past, come forth great heads, containing within them round pale seed: the roote is small, according to the proportion of the plant aboue ground, but made after the fashion of the greater kindes, with tuberous peeces spreading from the sides, and strong fibres or strings, whereby they are fastened in the ground.

2. *Chamæiris latifolia maior purpurea.* The greater purple dwarfe Flowerdeluce.

There is no difference either in roote, leafe, or forme of flower in this from the former dwarfe kinde, but onely in the colour of the flower, which in some is of a very deepe or blacke Violet purple, both the toppes and the fals: in others the Violet purple is more liuely, and in some the vpper leaues are blew, and the lower leaues purple, yet all of them haue that yellow frize or thrume in the middle of the falling leaues, that the other kindes haue.

*Altera.* There is another that beareth purple flowers, that might be reckoned, for the smalnesse and shortnesse of his stalke, to the next kinde, but that the flowers and leaues of this are as large as any of the former kindes of the smaller Flowerdeluces.

3. *Chamæiris latifolia minor alba.* The lesser white dwarfe Flowerdeluce.

There is also another sort of these Flowerdeluces, whose leaues and flowers are lesse, and wherein there is much variety. The leaues of this kinde, are all for the most part somewhat smaller, narrower, and shorter then the former: the stalke with the flower vpon it scarce riseth aboue the leaues, so that in most of them it may be rather called a foote-stalke, such as the Saffron flowers haue, and are therefore called of manie ἀκαυλοι, without stalkes; the flowers are like vnto the first described of the dwarfe kindes, and of a whitish colour, with a few purplish lines at the bottome of the vpper leaues, and a list of greene in the falling leaues.

*Straminea.* Another hath the flowers of a pale yellow, called a Straw colour, with whitish stripes and veines in the fals, and purplish lines at the bottome of the vpper leaues.

4. *Chamæiris latifolia minor purpurea.* The lesser purple dwarfe Flowerdeluce.

The difference of this from the former, consisteth more in the colour then forme of the flower, which is of a deep Violet purple, sometimes paler, and sometimes so deep, that it almost seemeth blacke: And sometimes the fals purplish, and the vpper leaues blew. Some of these haue a sweete sent, and some none.

*Cærulea.* There is another of a fine pale or delayed blew colour throughout the whole flower.

5. *Chamæiris latifolia minor suauerubens.*
The lesser blush coloured dwarfe Flowerdeluce.

This Flowerdeluce hath the falling leaues of the flower of a reddish colour, and the thrumes blew: the vpper and arched leaues of a fine pale red or flesh colour, called a blush colour; in all other things it differeth not, and smelleth little or nothing at all.

6. *Chamæiris*

*6. Chamæiris latifolia minor lutea versicolor.*
The lesser yellow variable dwarfe Flowerdeluce.

The falling leaues of this Flowerdeluce are yellowish, with purple lines from the middle downewards, sometimes of a deeper, and sometimes of a paler colour, and white thrumes in the middle, the vpper leaues are likewise of a yellowish colour, with purple lines in them: And sometimes the yellow colour is paler, and the lines both in the vpper and lower leaues of a dull or dead purple colour.

*3. Chamæiris latifolia minor cærulea versicolor.*
The lesser blew variable dwarfe Flowerdeluce.

The vpper leaues of this flower are of a blewish yellow colour, spotted with purple in the broad part, and at the bottome very narrow: the falling leaues are spread ouer with pale purplish lines, and a small shew of blew about the brimmes: the thrume is yellow at the bottome, and blewish aboue: the arched leaues are of a blewish white, being a little deeper on the ridge.

And sometimes the vpper leaues are of a paler blew rather whitish, with the yellow: both these haue no sent at all.

*8. Chamæiris marina purpurea.* The purple dwarfe Sea Flowerdeluce.

This small Flowerdeluce is like vnto the narrow leafed Sea Flowerdeluce before described, both in roote, leafe, and flower, hauing no other difference, but in the smalnesse and lownesse of the growing, being of the same purple colour with it.

*9. Chamæiris angustifolia maior.* The greater Grasse Flowerdeluce.

This Grasse Flowerdeluce hath many long and narrow darke greene leaues, not so stiffe as the former, but lither, and bending their ends downe againe, among which rise vp diuers stalkes, bearing at the toppe two or three sweete flowers, as small as any of them set downe before, of a reddish purple colour, with whitish yellow and purple strakes downe the middle of the falling leaues: the arched leaues are of a horse flesh colour all along the edges, and purple vpon the ridges and tips that turne vp againe: vnder these appeare three browne aglets, like vnto birds tongues: the three vpper leaues are small and narrow, of a perfect purple or Violet colour: the heads for seede haue sharper and harder cornered edges then the former: the seedes are somewhat grayish like the former, and so are the rootes, being small, blacke, and hard, growing thicke together, fastened in the ground with small blackish hard strings, which hardly shoote againe if the roote be remoued.

10. *Chamæiris angustifolia minor.* The lesser Grasse Flowerdeluce.

This Flowerdeluce is in leaues, flowers, and rootes so like the last described, that but onely it is smaller and lower, it is not to be distinguished from the other. And this may suffice for these sorts of Flowerdeluces, that furnish the Gardens of the curious louers of these varieties of nature, so farre forth as hath passed vnder our knowledge. There are some other that may be referred hereunto, but they belong to another history; and therefore I make no mention of them in this place.

The Place.

The places of most of these are set downe in their seuerall titles; for some are out of Turkie, others out of Hungaria, Dalmatia, Illyria, &c. as their names doe import. Those that grow by the Sea, are found in Spaine and France.

The

### The Time.

Some of these do flower in Aprill, some in May, and some not vntill Iune.

### The Names.

The names expressed are the fittest agreeing vnto them, and therefore it is needlesse againe to repeate them. Many of the rootes of the former or greater kindes, being dryed are sweete, yet some more then other, and some haue no sent at all: but aboue all the rest, that with the white flower, called of Florence, is accounted of all to be the sweetest root, fit to be vsed to make sweete powders, &c. calling it by the name of *Orris* rootes.

## *Iris tuberosa.* The Veluet Flowerdeluce.

Vnto the Family of Flowerdeluces, I must needes ioyne this peculiar kinde, because of the neare resemblance of the flower, although it differ both in roote and leafe; lest therefore it should haue no place, let it take vp a roome here in the end of the Flowerdeluces, with this description following. It hath many small and foure square leaues, two foote long and aboue sometimes, of a grayish greene colour, stiffe at the first, but afterwards growing to their full length, they are weak and bend downe to the ground: out of the middle, as it were of one of these leaues, breaketh out the stalke, a foot high and better, with some leaues thereon, at the toppe whereof, out of a huske riseth one flower, (I neuer saw more on a stalke) consisting of nine leaues, whereof the three that fall downe are of a yellowish greene colour round about the edges, and in the middle of so deepe a purple, that it seemeth to be blacke, resembling blacke Veluet: the three arched leaues, that couer the lower leaues to the halfe, are of the same greenish colour that the edges and backside of the lower leaues are: the three vppermost leaues, if they may be called leaues, or rather short peeces like eares, are green also, but wherein a glimpse of purple may be seene in them: after the flower is past, there followeth a round knob or whitish seede vessell, hanging downe by a small foote-stalke, from betweene the huske, which is diuided as it were into two leaues, wherein is contained round white seede. The roote is bunched or knobbed out into long round rootes, like vnto fingers, two or three from one peece, one distant from another, and one longer then another, for the most part of a darkish gray colour, and reddish withall on the outside, and somewhat yellowish within.

### The Place.

It hath beene sent out of Turkie oftentimes (as growing naturally thereabouts) and not knowne to grow naturally any where else.

### The Time.

It flowreth in Aprill or May, sometimes earlier or later, as the Spring falleth out to be milde or sharpe.

### The Names.

Matthiolus contendeth to make it the true *Hermodactylus*, rather from the shew of the rootes, which (as is said) are like vnto fingers, then from any other good reason: for the rootes hereof eyther dry or greene, do nothing resemble the true *Hermodactyli* that are vsed in Physicke, as any that knoweth them may easily perceiue, either in forme or vertue. It is more truely referred to the Flowerdeluces, and because of the tuberous rootes, called *Iris tuberosa*, although all the Flowerdeluces in this Chapter haue tuberous rootes,

rootes, yet this much differing from them all. In English it is vsually called, The Veluet Flowerdeluce, because the three falling leaues seeme to be like smooth blacke Veluet.

The Vertues.

Both the rootes and the flowers of the great Flowerdeluces, are of great vse for the purging and cleansing of many inward, as well as outward diseases, as all Authors in Physicke doe record. Some haue vsed also the greene rootes to cleanse the skinne, but they had neede to be carefull that vse them, lest they take more harme then good by the vse of them. The dryed rootes called *Orris* (as is said) is of much vse to make sweete powders, or other things to perfume apparrell or linnen. The iuice or decoction of the green roots doth procure both neezing to be snuft vp into the nostrils, and vomiting very strongly being taken inwardly.

## Chap. XXI.

### *Gladiolus*. Corne Flagge.

NExt vnto the Flagges or Flowerdeluces, come the *Gladioli* or Corne Flagges to bee entreated of, for some resemblance of the leaues with them. There are hereof diuers sorts, some bigger and some lesser, but the chiefest difference is in the colour of the flowers, and one in the order of the flowers. Of them all in their seuerall orders.

### *Gladiolus Narbonensis*. The French Corne Flagge.

The French Corne Flagge riseth vp with three or foure broad, long, and stiffe greene leaues, one as it were out of the side of another, being ioyned together at the bottome, somewhat like vnto the leaues of Flowerdeluces, but stiffer, more full of ribbes, and longer then many of them, and sharper pointed: the stalke riseth vp from among the leaues, bearing them on it as it riseth, hauing at the toppe diuers huskes, out of which come the flowers one aboue another, all of them turning and opening themselues one way, which are long and gaping, like vnto the flowers of Foxegloue, a little arched or bunching vp in the middle, of a faire reddish purple colour, with two white spots within the mouth thereof, one on each side, made like vnto a Lozenge that is square and long pointed: after the flowers are past, come vp round heads or seede vessels, wherein is contained reddish flat seede, like vnto the seede of the Fritillaria, but thicker and fuller: the roote is somewhat great, round, flat, and hard, with a shew as if it were netted, hauing another short spongie one vnder it, which when it hath done bearing, and the stalke dry, that the roote may be taken vp, sticketh close to the bottome, but may be easily taken away, hauing vsually a number of small rootes encreased about it, the least whereof will quickly grow, so that if it be suffered any long time in a Garden, it will rather choake and pester it, then be an ornament vnto it.

### *Gladiolus Italicus binis floribus ordinibus*. The Italian Corne Flagge.

The Italian Corne Flagge is like vnto the French in roote, leafe, and flower, without any other difference, then that the roote is smaller and browner, the leafe and stalke of a darker colour, and the flowers (being of a little darker colour like the former, and somewhat smaller) stand out on both sides of the stalke.

### *Gladiolus Byzantinus*. Corne Flagge of Constantinople.

This Corne Flagge that came first from Constantinople, is in all things like vnto the French Corne Flagge last described, but that it is larger, both in rootes, leaues, and flowers,

flowers, and likewise that the Flowers of this, which stand not on both sides, are of a deeper red colour, and flower later, after all the rest are past: the roote hereof being netted as plainly as any of the former, is as plentifull also to giue encrease, but is more tender and lesse able to abide our sharpe cold Winters.

*Gladiolus flore rubente.* Blush Corne Flagge.

This blush kinde is like vnto the French Corne Flagge in all respects, sauing onely that the flowers are of a pale red colour, tending to whitenesse, which wee vsually call a blush colour.

*Gladiolus flore albo.* White Corne Flagge.

This white Corne Flagge also differeth not from the last, but onely that the rootes are whiter on the outside, the leaues are greener, without any brownnesse or darknesse as in the former, and the flowers are snow white.

*Gladiolus purpureus minor.* The small purple Corne Flagge.

This also differeth not from any of the former, but onely in the smalnesse both of leafe, stalke, and flowers, which stand all on the one side, like vnto the French kinde, and of the same colour: the roote of this kinde is netted more then any other.

## The Place.

They grow in France and Italy, the least in Spaine, and the Byzantine, as it is thought, about Constantinople, being (as is said) first sent from thence. Iohn Tradescante assured mee, that hee saw many acres of ground in Barbary spread ouer with them.

## The Time.

They all flower in Iune and Iuly, and the Byzantine latest, as is said before.

## The Names.

It hath diuers names; for the Latines call it *Gladiolus*, of the forme of a sword, which the leafe doth resemble. The Romanes *Segetalis*, because it groweth in the Corne fields. Some call it *Victorialis rotunda*, to put a difference between it, and the *longa*, which is a kinde of Garlicke. Plinie saith, that *Gladiolus* is *Cyperus*, but to decide that controuersie, and many others, belongeth to another discourse, this being intended only for pleasure. Gerard mistaketh the French kinde for the Italian.

## The Vertues.

The roote being bruised, and applyed with Frankinsense (and often of it selfe without it) in the manner of a pultis or plaister, is held of diuers to be singular good to draw out splinters, thornes, and broken bones out of the flesh. Some take it to be effectuall to stirre vp Venerie, but I somewhat doubt thereof: For Galen in his eighth Booke of Simples, giueth vnto it a drawing, digesting, and drying faculty.

CHAP.

1 *Gladiolus Narbonensis*, The French Corne Flagge. 2 *Gladiolus Italicus*. The Italian Corne Flagge. 3 *Gladiolus Byzantinus*. Corne Flagge of Constantinople. 4 *Palma Christi mas*. The great male handed Satyrion. 5 *Orchis Hermaphroditica candida*, The white Butterflie Orchis. 6 *Orchis Melitias flue apifera*. The Bee flower or Bee Orchis. 7 *Dens Caninus flore purpurante*, Dogges tooth Violet with a pale purplish flower. 8 *Dens Caninus flore albo*. Dogges tooth Violet with a white flower.

## Chap. XXII.

### *Orchis siue Satyrium.* Bee flowers.

ALthough it is not my purpose in this place, to giue a generall history of all the sorrs of Orchides, Satyrions, and the rest of that kinde; yet because many of them are very pleasant to behold, and, if they be planted in a conuenient place, will abide some time in Gardens, so that there is much pleasure taken in them: I shall intrude some of them for curiosities sake, to make vp the prospect of natures beautifull variety, and only entreate of a few, leauing the rest to a more ample declaration.

1. *Satyrium Basilicum siue Palma Christi mas.*
The greater male handed Satyrion.

This handed Satyrion hath for the most part but three faire large greene leaues, neare vnto the ground, spotted with small blackish markes: from among which riseth vp a stalke, with some smaller leaues thereon, bearing at the toppe a bush or spike of flowers, thicke set together, euery one whereof is made like a body, with the belly broader belowe then aboue, where it hath small peeces adioyned vnto it: the flower is of a faire purple colour, spotted with deeper purple spots, and hauing small peeces like hornes hanging at the backes of the flowers, and a small leafe at the bottome of the foote-stalke of euery flower: the rootes are not round, like the other Orchides, but somewhat long and flat, like a hand, with small diuisions belowe, hanging downe like the fingers of a hand, cut short off by the knockles, two alwayes growing together, with some small fibres or strings aboue the heads of these rootes, at the bottome of the stalke.

2. *Satyrium Basilicum siue Palma Christi femina.*
The female handed Satyrion.

This female Satyrion hath longer and narrower leaues then the former, and spotted with more and greater spots, compassing the stalke at the bottome like the other: this beareth likewise a bush of flowers, like vnto the other, but that each of these haue heads like hoods, whereas the former haue none: in some they are white with purple spots, and in others of a reddish purple, with deep or darke coloured spots: the roots are alike.

3. *Orchis Hermaphroditica candida.* The white Butterflie Orchis.

The rootes of this kinde take part with both the sorts of *Orchis* and *Satyrium*, being neither altogether round, nor fully handed, and thereupon it tooke the name, to signifie both kindes: the leaues are two in number, seldome more, being faire and broad, like vnto the leaues of Lillies, without any spot at all in them: at the toppe of the stalke stand many white flowers, not so thicke set as the first or second, euery one being fashioned like vnto a white Butterflie, with the wings spread abroad.

4. *Orchis Melitias siue apifera.* The Bee flower or Bee Orchis.

This is a small and lowe plant for the most part, with three or foure small narrow leaues at the bottome: the stalke is seldome aboue halfe a foote high, with foure or fiue flowers thereon one aboue another, hauing round bodies, and somewhat flat, of a kind of yellowish colour, with purple wings aboue them, so like vnto an honey Bee, that it might soone deceiue one that neuer had seene such a flower before: the roots are two together, round and white, hauing a certaine *muccilagineſſe* or clammineſſe within them, without any taste almost at all, as all or the most part of these kindes haue.

5. *Orchis Sphegodes.* Gnats Satyrion.

The leaues of this Orchis are somewhat larger then of the Bee flower, the stalke also somewha:

somewhat higher: the flowers are fewer on the toppe, but somewhat larger then of the Bee flowers, made to the resemblance of a Gnat or great long Flie: the rootes are two round bulbes, as the other are.

*6. Orchis Myodes.* Flie Orchis.

The Flie Orchis is like vnto the last described, both in leafe and roote, the difference is in the flower, which is neither so long as the Gnat Satyrion, nor so great as the Bee Orchis, but the neather part of the Flie is blacke, with a list of ash-colour crossing the backe, with a shew of legges hanging at it: the naturall Flie seemeth so to bee in loue with it, that you shall seldome come in the heate of the day, but you shall finde one sitting close thereon.

The Place.

These grow in many places of England, some in the Woods, as the Butterflie, and the two former handed Satyrions: others on dry bankes and barren balkes in Kent, and many other places.

The Time.

They flower for the most part in the beginning or middle of May, or thereabouts.

The Names.

Their seuerall names are expressed in their titles, so much as may suffice for this discourse.

The Vertues.

All the kindes of Orchis are accounted to procure bodily lust, as well the flowers distilled, as the rootes prepared.

The rootes boyled in red Wine, and afterwards dryed, are held to bee a singular good remedie against the bloody Flixe.

## Chap. XXIII.

*Dens Caninus.* Dogs tooth Violet.

VNto the kindes of Orchides, may fitly be ioyned another plant, which by many is reckoned to be a *Satyrium*, both from the forme of roote and leafe, and from the efficacy or vertue correspondent thereunto. And although it cannot be the *Satyrium Erythronium* of Dioscorides, as some would entitle it, for that as I haue shewed before, his *Satyrium tryphillum* is the Tulipa without all doubt; yet because it differeth very notably, and carrieth more beauty and respect in his flower then they, I shall entreate thereof in a Chapter by it selfe, and set it next vnto them.

*Dens Caninus flore albo.* Dogs tooth Violet with a white flower.

The white Dogs tooth hath for his roote a white bulbe, long and small, yet vsually greater then either of the other that follow, bigger belowe then aboue, with a small peece adioyning to the bottome of it, from whence rise vp in the beginning of the Spring, after the Winter frosts are past, two leaues for the most part (when it will flower, or else but one, and neuer three together that euer I saw) closed together when they first come vp out of the ground, which inclose the flower betweene them: the leaues when they are opened do lay themselues flat on the ground, or not much aboue it, one opposite vnto the other, with the stalke and the flower on it standing betweene them, which leaues are of a whitish greene colour, long and narrow, yet broader in the

middle then at both ends, growing leſſe by degrees each way, ſpotted and ſtriped all ouer the leaues with white lines and ſpots: the ſtalke riſeth vp halfe a foote high or more, bearing at the toppe one flower and no more, hanging downe the head, larger then any of the other of this kinde that follow, made or conſiſting of ſix white long and narrow leaues, turning themſelues vp againe, after it hath felt the comfort of the Sunne, that they doe almoſt touch the ſtalke againe, very like vnto the flowers of *Cyclamen* or Sowebread: it hath in the middle of the flower ſix white chiues, tipt with darke purple pendents, and a white three forked ſtile in the middle of them: the flower hath no ſent at all, but commendable onely for the beauty and forme thereof: after the flower is paſt, commeth in the place a round head ſeeming three ſquare, containing therein ſmall and yellowiſh ſeede.

*Dens Caninus flore purpuraſcente.* Dogs tooth with a pale purple flower.

This other Dogs tooth is like vnto the former, but leſſer in all parts, the leafe whereof is not ſo long, but broad and ſhort, ſpotted with darker lines and ſpots: the flower is like the other, but ſmaller, and of a delayed purple colour, very pale ſometimes, and ſometimes a little deeper, turning it ſelfe as the other, with a circle round about the vmbone or middle, the chiues hereof are not white, but declining to purple: the roote is white, and like vnto the former, but leſſer, as is ſaid before.

*Dens Caninus flore rubro.* Dogs tooth with a red flower.

This is in all things like vnto the laſt, both for forme and bigneſſe of flower and leafe: the chiefe difference conſiſteth in this, that the leaues hereof are of a yellowiſh mealy greene colour, ſpotted and ſtreaked with redder ſpots and ſtripes, and the flower of a deeper reddiſh purple colour, and the chiues alſo more purpliſh then the laſt, in all other things it is alike.

## The Place.

The ſorts of *Dens Caninus* doe growe in diuers places; ſome in Italy on the Euganean Hils, others on the Apenine, and ſome about Gratz, the chiefe Citie of Stiria, and alſo about Bayonne, and in other places.

## The Time.

They flower in March moſt vſually, and many times in Aprill, according to the ſeaſonableneſſe of the yeare.

## The Names.

Cluſius did call it firſt *Dentali*, and Lobel, and from him ſome others *Satyrium*, and *Erythronium*, but I haue ſaid enough hereof in the beginning of the Chapter. It is moſt commonly called *Dens Caninus*, and we in Engliſh, either Dogs tooth, or Dogs tooth Violet. Geſner called it *Hermodactylus*, and Matthiolus *Pſeudohermodactylus*.

## The Vertues.

The roote hereof is held to bee of more efficacy for venereous effects, then any of the Orchides and Satyrions.

They of Stiria vſe the rootes for the falling ſickneſſe.

Wee haue had from Virginia a roote ſent vnto vs, that wee might well iudge, by the forme and colour thereof being dry, to be either the roote of this, or of an Orchis, which the naturall people hold not onely to be ſingular to procure luſt, but hold it as a ſecret, loth to reueale it.

## Chap. XXIIII.

### *Cyclamen.* Sowebread.

THe likeneſſe of the flowers, and the ſpotting of the leaues of the *Dens Caninus*; with theſe of the *Cyclamen* or Sowebread, maketh mee ioyne it next thereunto: as alſo that after the bulbous rooted plants I might begin with the tuberous that remaine, and make this plant the beginning of them. Of this kinde there are diuers ſorts, differing both in forme of leaues and time of flowring: for ſome doe flower in the Spring of the yeare, others afterwards in the beginning of Summer: but the moſt number in the end of Summer, or beginning of Autumne or Harueſt, whereof ſome haue round leaues, others cornered like vnto Iuie, longer or ſhorter, greater or ſmaller. Of them all in order, and firſt of thoſe that come in the Spring.

1. *Cyclamen Vernum flore purpureo.* Purple flowred Sowebread of the Spring.

This Sowebread hath a ſmaller roote then moſt of the others, yet round and blackiſh on the outſide, as all or moſt of the reſt are (I ſpeake of them that I haue ſeene; for Cluſius and others doe report to haue had very great ones) from whence riſe vp diuers round, yet pointed leaues, and ſomewhat cornered withall, greene aboue, and ſpotted with white ſpots circlewiſe about the leafe, and reddiſh vnderneath, which at their firſt comming vp are folded together; among which come the flowers, of a reddiſh purple colour and very ſweete, euery one vpon a ſmall, long, and ſlender reddiſh foote-ſtalke, which hanging downe their heads, turne vp their leaues againe: after the flowers are paſt, the head or ſeede veſſell ſhrinketh downe, winding his footeſtalke, and coyling it ſelfe like a cable, which when it toucheth the ground, there abideth hid among the leaues, till it be growne great and ripe, wherein are contained a few ſmall round ſeedes, which being preſently ſowne, will growe firſt into round rootes, and afterwards from them ſhoote forth leaues.

2. *Cyclamen Vernum flore albo.* White flowred Sowebread of the Spring.

The white flowring Sowebread hath his leaues like the former, but not fully ſo much cornered, bearing ſmall ſnow white flowers, as ſweete as the other: and herein conſiſteth the chiefeſt difference, in all other things it is alike.

3. *Cyclamen Vernum Creticum flore albo.* White Candy Sowebread of the Spring.

This Sowebread is ſomewhat like the former white kinde, but that the leaues grow much larger and longer, with more corners at the edges, and more eminent ſpots on them: the flowers alſo ſomewhat longer and larger, and herein conſiſteth the whole difference.

4. *Cyclamen Æſtivum.* Summer Sowebread.

Summer Sowebread hath round leaues like vnto the Romane Sowebread, but ſomewhat cornered, yet with ſhorter corners then the Iuie leafed Sowebread, full of white ſpots on the vpperſide of the leaues, and very purple vnderneath, ſometimes they haue fewer ſpots, and little or no purple vnderneath: the flowers hereof are as ſmall, as purple, and as ſweete, as the purple Sowebread of the Spring time: the roote hereof is likewiſe ſmall, blacke, and round.

5. *Cyclamen Romanum rotundifolium.* Romane Sowebread with round leaues.

The Romane Sowebread hath round leaues, ſomewhat like vnto the common Sowebread, but not fully ſo round pointed at the ends, a little cornered ſometimes alſo, or as it were indented, with white ſpots round about the middle of the leaues,

and very conspicuous, which make it seeme the more beautifull: the flowers appeare in Autumne, and are shorter, and of a deeper purplish red colour then the Iuie Sowebread, rising vp before the leaues for the most part, or at least with them, and little or nothing sweete: the roote is round and blacke, vsually not so flat as it, but growing sometimes to bee greater then any other kinde of Sowebread. There is sometimes some variety to be seene, both in the leaues and flowers of this kinde; for that sometime the leaues haue more corners, and either more or lesse spotted with white: the
*Varietas.* flowers likewise of some are larger or lesser, longer or rounder, paler or deeper coloured one then another. This happeneth most likely from the sowing of the seede, causing the like variety as is seene in the Iuie leafed Sowebread. It doth also many times happen from the diuersity of soyles and countries where they grow: the seed of this, as of all the rest, is small and round, contained in such like heads as the former, standing almost like the head of a Snake that is twined or folded within the body thereof. This and the other Autumnall kindes, presently after their sowing in Autumne, shoote forth leaues, and so abide all the Winter, according to their kinde.

### 6. *Cyclamen folio hederæ autumnale.* Iuie leafed Sowebread.

The Iuie leafed Sowebread groweth in the same manner that the former doth, that is, bringeth forth flowers with the leaues sometimes, or most commonly before them, whose flowers are greater then the common round leafed Sowebread, somewhat longer then the former Romane or Italian Sowebreads, and of a paler purple colour, almost blush, without that sweete sent as is in the first kinde of the Spring: the greene leaues hereof are more long then round, pointed at the ends, and hauing also one or two corners on each side, sometimes much spotted on the vpperside with white spots and marks, and sometimes but a little or not at all; and so likewise sometimes more or lesse purple vnderneath: all the leaues and flowers doe stand vsually euery one seuerally by themselues, vpon their owne slender foote-stalkes, as most of all the other
*Varietas.* kindes doe: but sometimes it happeneth, that both leaues and flowers are found growing from one and the same stalke, which I rather take to be accidentall, then naturall so to continue: the seede hereof is like the former kindes, which being sowne produceth variety, both in the forme of the leaues, and colour and smell of the flowers: some being paler or deeper, and some more or lesse sweete then others: the leaues also, some more or lesse cornered then others: the root groweth to be great, being round and flat, and of a blackish browne colour on the outside.

### 7. *Cyclamen autumnale hederæfolio flore albo.* Iuie leafed Sowebread with white flowers.

There is one of this kinde, whose leaues are rounder, and not so much cornered as the former, flowring in Autumne as the last doth, and whose flowers are wholly white, not hauing any other notable difference therein.

### 8. *Cyclamen autumnale angustifolium.* Long leafed Sowebread.

This kinde of Sowebread may easily be knowne from all the other kindes, because his leafe is longer and narrower then others, fashioned at the bottome thereof with points, somewhat like vnto *Arum* or Wake Robin leaues: the flowers are like the former sorts for forme, but of a purple colour. There is also another of this kinde in all things like the former, but that the flowers are white.

### 9. *Cyclamen Antiochenum Autumnale flore purpureo duplici.* Double flowred Sowebread of Antioch.

This Sowebread of Antioch with double flowers, hath his leaues somewhat round, like vnto the leaues of the Summer Sowbread, but with lesse notches or corners, & full of white spots on them: it beareth flowers on stalks, like vnto others, & likewise some stalks that haue two or three flowers on them, which are very large, with ten or twelue

leaues

1 *Cyclamen Vernum flore purpureo* Purple flowred Sowebread of the Spring. 2 *Cyclamen æstivum.* Summer Sowebread. 3 *Folium Cyclaminu Cretici vernalis flore candido.* A leafe of Candie Sowebread. 4 *Cyclamen Romanum Autumnale.* Romane Sowebread of the Autumne. 5 *Cyclamen hederæfolio Autumnale.* Ivie leafed Autumne Sowebread. 6 *Folium Cyclaminis Autumnalis flore albo.* A leafe of the Autumne Sowebread with a white flower. 7 *Folium Cyclaminis angustifolij Autumnalis.* A leafe of the long leafed Sowebread. 8 *Cyclamen Antiochenum Autumnale flore ample purpureo duplici.* The double flowred Sowebread of Antioch. 9 *Cyclamen vulgare folio rotundo.* The common round leafed Sowebread.

leaues a peece, of a faire Peach colour, like vnto the flowers of purple Sowebread of the Spring, and deeper at the bottome.

There are of this kinde some, whose flowers appeare in the Spring, and are as large and double as the former, but of a pure white colour.

There are of these Sowebreads of Antioch, that haue but single flowers, some appearing in the Spring, and others in Autumne.

## 10. *Cyclamen vulgare folio rotundo.* The common Sowebread.

The common Sowebread (which is most vsed in the Apothecaries Shops) hath many leaues spread vpon the ground, rising from certaine small long heads, that are on the greater round rootes, as vsually most of the former sorts doe, being in the like manner folded together, and after spread themselues into round greene leaues, somewhat like vnto the leaues of *Asarum*, but not shining, without any white spots on the vpperside for the most part, or but very seldome, and reddish or purplish vnderneath, and very seldome greener: the flowers stand vpon small foot-stalkes, and shew themselues open for the most part, before any leaues doe appeare, being smaller and shorter then those with Iuie leaues, and of a pale purple colour, yet sometimes deeper, hanging downe their heads, and turning vp their leaues againe, as all others doe, but more sweete then many other of the Autumne flowers: after the flowers are past, come the heads turning or winding themselues downe in like manner as the other do, hauing such like seede, but somewhat larger, and more vneuen, or not so round at the least: the roote is round, and not flat, of a browner colour, and not so blacke on the outside as many of the others.

### The Place.

The Sowebreads of the Spring doe both grow on the Pyrenæan Mountaines in Italy, and in Candy, and about Mompelier in France; Antioch in Syria also hath yeelded some both of the Spring and Autumne. Those with round and Iuie leaues grow in diuers places both of France and Italy: and the common in Germany, and the Lowe-Countries. But that Autumne Sowebread with white flowers, is reported to grow in the Kingdome of Naples. I haue very curiously enquired of many, if euer they found them in any parts of England, neare or farther off from the places where they dwell: but they haue all affirmed, that they neuer found, or euer heard of any that haue found of any of them. This onely they haue assured, that there groweth none in the places, where some haue reported them to grow.

### The Time.

Those of the Spring doe flower about the end of Aprill, or beginning of May. The other of the Summer, about the end of Iune or in Iuly. The rest some in August, and September, others in October.

### The Names.

The Common Sowebread is called by most Writers in Latine, *Panis Porcinus*, and by that name it is knowne in the Apothecaries shops, as also by the name *Arthanita*, according to which name, they haue an ointment so called, which is to be made with the iuice hereof. It is also called by diuers other names, not pertinent for this discourse. The most vsuall name, whereby it is knowne to most Herbarists, is *Cyclamen* (which is the Greeke word) or as some call it *Cyclaminus*, adding thereunto their other seuerall titles. In English, Sowebread.

### The Vertues.

The leaues and rootes are very effectuall for the spleene, as the Ointment before remembred plainly proueth, being vsed for the same purpose, and

and that to good effect. It is vsed also for women in long and hard trauels, where there is danger, to accelerate the birth, either the roote or the leafe being applyed. But for any amorous effects, I hold it meere fabulous.

## CHAP. XXV.

### *Anemone.* Windeflower and his kindes.

THe next tuberous rooted plants that are to follow (of right in my opinion) are the *Anemones* or Windeflowers, and although some tuberous rooted plants, that is, the Asphodils, Spiderworts, and Flowerdeluces haue beene before inserted, it was, both because they were in name or forme of flowers sutable to them whom they were ioyned vnto, and also that they should not be seuered and entreated of in two seuerall places: the rest are now to follow, at the least so many of them as be beautifull flowers, fit to furnish a Florists Garden, for natures delightsome varieties and excellencies. To distinguish the Family of *Anemones* I may, that is, into the wilde kindes, and into the tame or mannured, as they are called, and both of them noursed vp in Gardens; and of them into those that haue broader leaues, and into those that haue thinner or more iagged leaues: and of each of them, into those that beare single flowers, and those that beare double flowers. But to describe the infinite (as I may so say) variety of the colours of the flowers, and to giue to each his true distinction and denomination, *Hic labor, hoc opus est*, it farre passeth my ability I confesse, and I thinke would grauell the best experienced this day in Europe (and the like I laid concerning Tulipas, it being as contingent to this plant, as is before said of the Tulipa, to be without end in yeelding varieties:) for who can see all the varieties that haue sprung from the sowing of the seede in all places, seeing the variety of colours risen from thence, is according to the variety of ayres & grounds wherein they are sowne, skill also helping nature in ordering them aright. For the seede of one and the same plant sowne in diuers ayres and grounds, doe produce that variety of colours that is much differing one from another; who then can display all the mixtures of colours in them, to set them downe in so small a roome as this Booke? Yet as I haue done (in the former part of this Treatise) my good will, to expresse as many of each kinde haue come to my knowledge, so if I endeauour the like in this, I hope the courteous wil accept it, and hold me excused for the rest: otherwise, if I were or could be absolute, I should take from my self and others the hope of future augmentation, or addition of any new, which neuer will be wanting. To begin therefore with the wilde kinds (as they are so accounted) I shall first entreate of the *Pulsatillas* or Pasque flowers, which are certainly kindes of wilde *Anemones*, both in leafe and flower, as may well be discerned by them that are iudicious (although some learned men haue not so thought, as appeareth by their writings) the rootes of them making one speciall note of difference, from the other sorts of wilde *Anemones*.

#### 1. *Pulsatilla Anglica purpurea.* The purple Pasque flower.

The Pasque or Passe flower which is of our owne Country, hath many leaues lying on the ground, somewhat rough or hairie, hard in feeling, and finely cut into many small leaues, of a darke greene colour, almost like the leaues of Carrets, but finer and smaller, from among which rise vp naked stalkes, rough or hairie also, set about the middle thereof with some small diuided leaues compassing them, and rising aboue these leaues about a spanne, bearing euery one of them one pendulous flower, made of six leaues, of a fine Violet purple colour, but somewhat deepe withall, in the middle whereof stand many yellow threeds, set about a middle purple pointell: after the flower is past, there commeth vp in the stead thereof a bushie head of long seedes, which are small and hoarie, hauing at the end of euery one a small haire, which is gray likewise: the roote is small and long, growing downewards into the ground, with a tuft of haire at the head thereof, and not lying or running vnder the vpper crust thereof, as the other wilde *Anemones* doe.

2. *Pulsa-*

2. *Pulsatilla Danica.* The Passe flower of Denmarke.

There is another that was brought out of Denmarke, very like vnto the former, but that it is larger both in roote and leafe, and flower also, which is of a fairer purple colour, not so deepe, and besides, will better abide to bee mannured then our English kinde will, as my selfe haue often proued.

*Vtriusque flore albo & flore duplici.*

Of both these sorts it is said, that some plants haue bin found, that haue borne white flowers. And likewise one that bore double flowers, that is, with two rowes of leaues.

3. *Pulsatilla flore rubro.* The red Passe flower.

Lobel, as I take it, did first set forth this kinde, being brought him from Syria, the leaues whereof are finer cut, the flower smaller, and with longer leaues, and of a red colour.

4. *Pulsatilla flore luteo.* The yellow Passe flower.

The yellow Passe flower hath his leaues cut and diuided, very like vnto the leaues of the first kinde, but somewhat more hairie, greene on the vpperside, and hairie vnderneath: the stalke is round and hoary, the middle whereof is beset with some small leaues, as in the other, from among which riseth vp the stalke of the flower, consisting of six leaues of a very faire yellow colour on the inside, and of a hoary pale yellow on the outside; after which followeth such an head of hairie thrummes as in the former: the roote is of the bignesse of a mans finger.

5. *Pulsatilla flore albo.* The white Passe flower.

The white Passe flower (which Clusius maketh a kinde of *Anemone*, and yet as hee saith himselfe, doth more nearely resemble the *Pulsatilla*) hath, from amongst a tuft or head of haires, which grow at the toppe of a long blacke roote, many leaues standing vpon long stalkes, which are diuided as it were into three wings or parts, and each part finely cut and diuided, like vnto the Passe flower of Denmarke, but somewhat harder in handling, greenish on the vpperside, and somewhat gray vnderneath, and very hairie all ouer: among these leaues rise vp the stalkes, beset at the middle of them with three leaues, as finely cut and diuided as those belowe, from aboue which standeth the flower, being smaller, and not so pendulous as the former, but in the like manner consisting of six leaues, of a snow white colour on the inside, and a little browner on the outside, with many yellow thrums in the middle: after the flower is past, riseth vp such a like hoary head, composed as it were of many haires, each whereof hath a small seede fastened vnto it, like as the former Passe flowers haue.

The Place.

The first is found in many places of England, vpon dry bankes that lye open to the Sunne.

The second was first brought, as I take it, by Doctor Lobel from Denmarke, & is one of the two kinds, that Clusius saith are common in Germanie, this bearing a paler purple flower, and more early then the other, which is the same with our English, whose flower is so darke, that it almost seemeth blacke.

The red kinde, as Lobel saith, came from Syria.

The yellow Passe flower, which Clusius maketh his third wilde *Anemone*, was found very plentifully growing at the foote of St. Bernards Hill, neare vnto the Cantons of the Switzers.

The white one groweth on the Alpes neare Austria, in France likewise, and other places.

The

1 *Pulsatilla purpurea cum folio, semine, & radice.* The purple Pasque flower with leafe, seed, and root. 2 *Pulsatilla luteo flore.* The yellow Pasque flower. 3 *Pulsatilla rubra Syriaca Lobelij.* Red Pasque flower of Lobel. 4 *Pulsatilla rubra Swertij.* Swertz his red Pasque flower. 5 *Pulsatilla flore albo.* White Pasque flower. 6 *Anemone siluestris alba Matthioli.* The wilde white broad leafed Windflower. 7 *Anemone siluestris tenuifolia alba.* The wilde single white Windflower. 8 *Anemone siluestris tenuifolia lutea.* The yellow wilde thin leafed Windflower. 9 *Anemone siluestris trifolia Dodonæi.* The three leafed wilde Windflower. 10 *Anemone siluestris flore pleno albo.* The double white wilde Windflower. 11 *Anemone siluestris flore pleno purpureo.* The double purple wilde Windflower. * *Semen separatim divulsum.* The seed separated. † *Radix cum folio inferiore.* The roote with a lower leafe.

### The Time.

All of them doe flower early in the yeare, that is, in the beginning of Aprill, about which time most commonly Easter doth fall.

### The Names.

Their proper names are giuen to each in their seuerall titles, being all of them kindes of wilde *Anemones*, as I said in the beginning of the Chapter, and so for the most part all Authors doe acknowledge them. We call them in English, because they flower about Easter, Pasque Flower, which is the French name for Easter, or *Euphoniæ gratia*, Passe Flower, which may passe currant, without any further descant on the name, or else *Pulsatilla*, if you will, being growne old by custome.

### The Vertues.

The sharpe biting and exulcerating quality of this plant, causeth it to be of little vse, notwithstanding Ioachimus Camerarius saith in his *Hortus Medicus*, that in Borussia, which is a place in Italy, as I take it, the distilled water hereof is vsed with good successe, to be giuen to them that are troubled with a Tertian Ague; for he saith that it is *medicamentum* ἐνδεγματικὸν, that is, a medicine of force to helpe obstructions.

*Anemone siluestris latifolia alba siue tertia Matthioli.*
The white wilde broad leafed Windflower.

This Windflower hath diuers broad greene leaues, cut into diuisions, and dented about, very like vnto a broad leafed Crowfoote, among which riseth vp a stalke, hauing some such like cut leaues in the middle thereof, as growe below, but smaller; on the toppe whereof standeth one large white flower, consisting of fiue leaues for the most part, with some yellow threads in the middle, standing about such a greene head as is in the tame or garden *Anemones*, which growing greater after the flower is past, is composed of many small seedes, wrapped in white wooll, which as soone as they are ripe, raise themselues vp from the bottome of the head, and flye away with the winde, as the other tame or garden kindes doe: the roote is made of a number of long blacke strings, encreasing very much by running vnder ground, and shooting vp in diuers places.

*Anemone siluestris tenuifolia lutea.* The yellow wilde thin leafed Windflower.

The yellow wilde *Anemone* riseth vp with one or two small round naked stalkes, bearing about the middle of them, small, soft, and tender iagged leaues, deeply cut in and indented on the edges about, from aboue which doth grow the stalke, bearing small yellow flowers, standing vpon weake foote-stalkes, like vnto a small Crowfoot, with some threads in the middle: the roote is long and small, somewhat like vnto the roote of Pollipodie, creeping vnder the vpper crust of the earth: this kinde is lower, and springeth somewhat earlier then the other wilde kindes that follow.

*Anemone siluestris tenuifolia alba simplex.*
The single white thin leafed wilde Windflower.

This white wilde *Anemone* riseth vp with diuers leaues vpon seuerall long stalkes; which are somewhat like vnto the former, but that they are somewhat harder, and not so long, nor the diuisions of the leaues so finely snipt about the edges, but a little broader, and deeper cut in on euery side: the flowers hereof are larger and broader then the former, white on the inside, and a little purplish on the outside, especially at the

the bottome of the flower next vnto the stalke: the roote of this is very like vnto the last.

There is another of this kinde, whose flowers are purple, in all other things it is like vnto the white. *Purpurea.*

And likewise another, with a blush or carnation coloured flower. *Coccinea siue suaue rubens.*

There is one that is onely nursed vp with vs in Gardens, that is somewhat like vnto these former wilde *Anemones* in roote and leafe, but that the flower of this, being pure white within, and a little purplish without, consisting of eight or nine small round pointed leaues, hath sometimes some leaues vnder the flower, party coloured white and greene: the flower hath likewise a greene head, like a Strawberry, compassed about with white threads, tipt with yellow pendents. *Peregrina alba.*

And another of the same kinde with the last, whose flower consisting of eight or nine leaues, is of a greenish colour, except the foure outermost leaues, which are a little purplish, and diuided at the points into three parts; the middle part is of a greenish white colour, with a greene head in the middle as the other. *Peregrina viridis.*

*Anemone siluestris trifolia Dodonæi.* The three leafed wilde Windflower.

This wilde *Anemone* hath his rootes very like vnto the former kindes; the leaues are alwaies three set together at the toppe of slender stalkes, being small and indented about, very like vnto a three leafed Grasse, but smaller: the flower consisteth of eight small leaues, somewhat like vnto a Crowfoote, but of a whitish purple or blush colour, with some white threads, and a greene rough head in the middle.

*Anemone siluestris flore pleno albo.* The double white wilde Windflower.

This double kinde is very like vnto the single white kinde before described, both in his long running rootes, and thin leaues, but somewhat larger: the flowers hereof are very thicke and double, although they be small, and of a faint sweete sent, very white after it is full blowne for fiue or six dayes, but afterwards it becommeth a little purplish on the inside, but more on the outside: this neuer giueth seede (although it haue a small head in the middle) like as many other double flowers doe.

*Anemone siluestris flore pleno purpureo.* The double purple wilde Windflower.

This double purple kinde hath such like iagged leaues as the last described hath, but more hoarie vnderneath: the flower is of a fine light purple toward the points of the leaues, the bottomes being of a deeper purple, but as thicke, and full of leaues as the former, with a greene head in the middle, like vnto the former: this kinde hath small greene leaues on the stalkes vnder the flowers, cut and diuided like the lower leaues.

## The Place.

The first broad leafed *Anemone* groweth in diuers places of Austria and Hungary. The yellow in diuers woods in Germany, but not in this Countrey that euer I could learne. The other single wilde kindes, some of them are very frequent throughout the most places of England, in Woods, Groues, and Orchards. The double kindes were found, as Clusius saith, in the Lowe-Countries, in a Wood neare Louaine.

## The Time.

They flower from the end of March (that is the earliest) and the beginning of Aprill, vntill May, and the double kindes begin within a while after the single kinds are past.

## The Names.

They are called *Ranunculi siluarum*, and *Ranunculi nemorum*, and as Clusius

sius would haue them, *Leimonia* of Theophrastus; they are generally called of most Herbarists *Anemones siluestres*, Wilde *Anemones* or Windflowers. The Italians call them *Gengeuo salnatico*, that is, Wilde Ginger, because the rootes are, besides the forme, being somewhat like small Ginger, of a biting hot and sharpe taste.

*Anemone Lusitanica siue hortensis latifolia flore simplici luteo.*
The single Garden yellow Windflower or Anemone.

This single yellow Anemone or Windflower hath diuers broad round leaues, somewhat diuided and endented withall on the edges, browniſh at the first rising vp out of the ground, and almost folded together, and after of a sad greene on the vpperside, and reddish vnderneath; among which rise vp small slender stalkes, beset at the middle of them with two or three leaues, more cut and diuided then those belowe, with small yellow flowers at the toppe of them, consisting of ten or twelue leaues a peece, hauing a few yellow threads in the middle of them, standing about a small greene head, which in time growing ripe hath small flat seede, inclosed within a soft wooll or downe, which is easily blowne away with the winde: the roote groweth downeward into the ground, diuersly spread with branches here and there, of a browniſh yellow on the outside, and whitiſh within, so brittle, that it can hardly bee touched without breaking.

*Anemone latifolia flore luteo duplici.* The double yellow Anemone or Windflower.

This double yellow Anemone hath such broad round leaues as the single kinde hath, but somewhat larger or ranker: the stalkes are beset with larger leaues, more deeply cut in on the edges: the flowers are of a more pale yellow, with some purpliſh veines on the outside, and a little round pointed; but they are all on the inside of a faire yellow colour, consisting of two rowes of leaues, whereof the innermost is the narrower, with a small greene head in the middle, compassed with yellow threads as in the former: the roote is like the roote of the single; neither of these haue any good sent, and this springeth vp and flowreth later then the single kinde.

*Anemone latifolia purpurea stellata siue papaveracea.*
The purple Starre Anemone or Windflower.

The first leaues of this purple Anemone, which alwayes spring vp before Winter, (if the roote be not kept too long out of the ground,) are somewhat like the leaues of *Sanicle* or Selfe-heale, but the rest that follow are more deeply cut in and iagged; among which rise vp diuers round stalkes, beset with iagged leaues as all other Anemones are, aboue which leaues, the stalkes rising two or three inches high, beare one flower a peece, composed of twelue leaues or more, narrow and pointed, of a bleake purple or whitiſh ash-colour, somewhat shining on the outside, and of a fine purple colour tending to a murrey on the inside, with many blackiſh blew threads or thrummes in the middle of the flower, set about a head, whereon groweth the seede, which is small and blacke, inclosed in soft wooll or downe, which flieth away with the winde, carrying the seede with it, if it be not carefully gathered: the roote is blackiſh on the outside, and white within, tuberous or knobby, with many fibres growing at it.

*Anemone purpurea Stellata altera.* Another purple Starre Anemone.

There is so great diuersity in the colours of the flowers of these broad leafed kinds of Anemones or Windflowers, that they can very hardly be expressed, although in their leaues there is but little or no difference. I shall not neede therefore to make seuerall descriptions of euery one that shall be set downe; but it will be sufficient, I thinke, to giue you the distinctions of the flowers: for as I said, therein is the greatest and chiefest difference. This other Starre Anemone differeth not from the former in leafe or flower, but onely that this is of a more pale sullen colour on the outside, and of a paler purple colour on the inside.

There

1 *Anemone latifolia flore luteo simplici.* The single yellow Anemone. 2 *Anemone latifolia flore luteo duplici.* The double yellow Anemone. 3 *Anemone latifolia flore purpureo Stellato.* The purple Starre Anemone. 4 *Anemone latifolia purpurea dilutior.* The pale purple Starre Anemone. 5 *Anemone latifolia flore miniato dilute.* The pale red Anemone. 6 *Anemone latifolia coccinea Cardinalis dicta.* The Cardinall Anemone. 7 *Anemone latifolia incarnata Hispanica.* The Spanish incarnate Anemone. 8 *Anemone latifolia Pæne simplex dicta.* The lesser Orenge tawney Anemone. 9 *Anemone latifolia flore carneo.* The carnation Anemone. 10 *Anemone latifolia Arantiaca sive Pæne maior.* The double Orenge tawney Anemone. 11 *Anemone Superitica sive Cyparissia.* The double Anemone of Cyprus. 12 *Anemone latifolia flore pleno albicante.* The double pale blush Anemone. 13 *Anemone Chalcedonica maxima.* The great Spanish Marigold Anemone. 14 *Anemone Coronaria sive Persica.* The double Persian Anemone. ✝ *Anemone latifolia radix.* The roote of a great Anemone.

*Viola purpurea* There is another, whose flower hath eight leaues, as many of them that follow haue (although diuers sorts haue but six leaues in a flower) and is of a Violet purple, and therefore is called, The Violet purple Anemone.

*Varietas.* Of all these three sorts last described, there be other that differ only in hauing white bottomes, some smaller and some larger.

*Purpurea striata.* There is also another of the same Violet purple colour with the former, but a little paler, tending more to rednesse, whose flowers haue many white lines and stripes through the leaues, and is called, The purple stript Anemone.

*Carnea vivacissima simplex.* There is another, whose greene leaues are somewhat larger, and so is the flower likewise, consisting of eight leaues, and sometimes of more, of the colour of Carnation silke, sometimes pale and sometimes deeper, with a whitish circle about the bottome of the leaues, which circle in some is larger, and more to be seene then in others, when the flower layeth it selfe open with the heate of the Sunne, hauing blewish threads in the middle. This may be called, the Carnation Anemone.

*Perficiviolacea.* We haue another, whose flower is betweene a Peach colour and a Violet, which is vsually called a Gredeline colour.

*Cochenille.* And another of a fine reddish Violet or purple, which we call, The Cochenille Anemone.

*Cardinalis.* And another of a rich crimson red colour, and may be called, The Cardinall Anemone.

*Sanguinea.* Another of a deeper, but not so liuely a red, called, The bloud red Anemone.

*Cramesina.* Another of an ordinary crimson colour, called, The crimson Anemone.

*Coccinea.* Another of a Stamell colour, neere vnto a Scarlet.

*Incarnata.* Another of a fine delayed red or flesh colour, and may bee called, The Incarnadine Anemone.

*Incarnata Hispanica.* Another whose flower is of a liuely flesh colour, shadowed with yellow, and may be called, The Spanish Incarnate Anemone.

*Rubescens.* Another of a faire whitish red, which we call, The Blush Anemone.

*Moschutella.* Another whose flower consisteth of eight leaues, of a darke whitish colour, stript all ouer with veines of a fine blush colour, the bottomes being white, this may be called, The Nutmegge Anemone.

*Enfumata.* Another whose flower is of a pale whitish colour, tending to a gray, such as the Monkes and Friers were wont to weare with vs, and is called, A Monkes gray.

*Pauo maior simplici flore.* There is another, whose leafe is somewhat broader then many or most of the Anemones, comming neare vnto the leafe of the great double Orenge coloured Anemone; the flower whereof is single, consisting of eight large or broad leaues, very neare vnto the same Orenge colour, that is in the double flower hereafter described, but somewhat deeper. This is vsually called in Latine, *Pauo maior simplici flore*, and we in English, The great single Orenge tawnie Anemone.

*Pauo minor.* There is likewise of this kinde another, whose flower is lesser, and called, The lesser Orenge tawnie Anemone.

*Varietas magna ex semine.* There is besides these expressed, so great a variety of mixt colours in the flowers of this kinde of Anemone with broad leaues, arising euery yeare from the sowing of the seede of some of the choisest and fittest for that purpose, that it is wonderfull to obserue, not onely the variety of single colours, but the mixture of two or three colours in one flower, besides the diuersity of the bottomes of the flowers, some hauing white or yellowish bottomes, and some none, and yet both of the same colour; and likewise in the thrums or threads in the middle: But the greatest wonder of beauty is in variety of double flowers, that arise from among the other single ones, some hauing two or three rowes of leaues in the flowers, and some so thicke of leaues as a double Marigold, or double Crowfoote, and of the same seuerall colours that are in the single flowers, that it is almost impossible to expresse them seuerally, and (as is said before) some falling out to bee double in one yeare, which will proue single or lesse double in another,

other, yet very many abiding constant double as at the first; and therefore let this briefe recitall be sufficient in stead of a particular of all the colours.

*Anemone Chalcedonica maxima versicolor.*
The great double Windflower of Constantinople.

This great Anemone of Constantinople hath broader and greener leaues then any of the former kindes, and not so much diuided or cut in at the edges, among which rise vp one or two stalkes, (seldome more from one roote) hauing some leaues about the middle of the stalke, as other Anemones haue, and bearing at the toppes of the stalkes one large flower a peece, very double, whose outermost leaues being broadest, are greenish at the first, but afterwards red, hauing sometimes some greene abiding still in the leaues, and the red striped through it: the other leaues which are within these are smaller, and of a perfect red colour; the innermost being smallest, are of the same red colour, but turned somewhat inward, hauing no thrummes or threads in the middle, as the former haue, and bearing no seede: the roote is blackish on the outside, and white within, thicke and tuberous as the other kindes, but thicker set and close together, not shooting any long slender rootes as others doe. Some Gentlewomen call this Anemone, The Spanish Marigold.

*Anemone Chalcedonica altera siue Pano maior flore duplici.*
The great double Orenge tawney Anemone.

This other great Anemone of Constantinople hath his large leaues so like vnto the last, that one can hardly distinguish them asunder; the stalke hath also such like leaues set vpon it, bearing at the toppe a faire large flower, consisting of many leaues set in two or three rowes at the most, but not so thicke or double as the last, yet seeming to be but one thicke rowe of many small and long leaues, of an excellent red or crimson colour, wherein some yellow is mixed, which maketh that colour is called an Orenge tawney; the bottomes of the leaues are red, compassed with a whitish circle, the thrummie head in the middle being beset with many darke blackish threads: the roote is like the former.

*Anemone Superitica siue Cyparissia.* The double Anemone of Cyprus.

This Anemone (which the Dutchmen call Superitz, and as I haue beene enformed, came from the Isle of Cyprus) hath leaues very like the last double Anemone, but not altogether so large: the flower consisteth of smaller leaues, of colour very neare vnto the last double Orenge coloured Anemone, but more thicke of leaues, and as double as the first, although not so great a flower, without any head in the middle, or thrums about it as is in the last, and differeth not in the roote from either of them both.

Somewhat like vnto this kinde, or as it were betweene this and the first kinde of these great double Anemones, we haue diuers other sorts, bearing flowers very thicke and double; some of them being white, or whitish, or purple, deeper or paler, and some of a reddish colour tending to Scarlet or a Carnation colour, and some also of a blush or flesh colour, and diuers other colours, and all of them continue constant in their colours.

*Anemone Cacumeni Maringi siue Persica.* The double Persian Anemone.

This rare Anemone, which is said to come out of Persia to Constantinople, and from thence to vs, is in leafe and roote very like vnto the former double Anemones before described; onely the flower hereof is rather like vnto the second great double Orenge coloured Anemone, vsually called *Pano maior flore pleno*, being composed of three rowes of leaues, the outtermost rowe consisting of ten or twelue larger leaues, and those more inward lesser and more in number, but all of them variably mixed with white, red, and yellow, hauing the bottomes of the leaues white: but instead of a middle head with thrums about it, as the other hath, this hath a few narrow leaues, of a deepe yellow colour in the middle of the flower, standing vpright,

Hauing thus farre proceeded in the two parts of the kindes of Anemones or Windflowers, it remaineth to entreate of the reſt, which is thoſe Anemones which haue thin cut leaues, whereof ſome haue reckoned vp thirty ſorts with ſingle flowers, which I confeſſe I haue not ſeene; but ſo many as haue come to my knowledge, I ſhall here ſet downe.

*Anemone tenuifolia ſiue Geranifolia cærulea.*
The Watchet Anemone or Storkes bill leafed Windflower.

This firſt Windflower with thin cut leaues, riſeth not out of the ground vntil the great Winter froſts be paſt, that is, about the middle or end of February, and are ſomewhat browniſh at their firſt appearing, but afterwards ſpread into wings of greene leaues, ſomewhat broader then the reſt that follow, diuided into three parts, & each part into three leaues, euery one cut in about the edges, one ſtanding againſt another vpon a long ſlender foote-ſtalke, and the end leafe by it ſelfe: among theſe riſeth vp two or three greene ſtalkes, garniſhed with ſuch like thin leaues as are at the bottome, from aboue which riſe the flowers, but one vpon a ſtalke, conſiſting of fourteene or fifteene ſmall pale blew or watchet leaues, leſſer then any of the ſingle kindes that follow, compaſſing many whitiſh threads, and a ſmall greene head in the middle, ſomewhat like the head of the wilde Crowfoote, wherein is contained ſuch like ſeede: the roote is blackiſh without, thruſting out into long tuberous peeces, ſomewhat like vnto ſome of the broad leafed Anemones.

*Alba.* Of this kinde there is another, whoſe leaues are not browne at their firſt riſing, but greene, and the flowers are white, in other things not differing.

*Anemone tenuifolia purpurea vulgaris.*
The ordinary purple Anemone with thin leaues.

This purple Anemone which is moſt common, and therefore the leſſe regarded, hath many winged leaues ſtanding vpon ſeuerall ſtalkes, cut and diuided into diuers leaues, much like vnto the leaues of a Carrot; among which riſe vp ſtalkes with ſome leaues thereon (as is vſuall to the whole Family of Anemones, both wilde and tame, as is before ſaid;) at the toppes whereof ſtand the flowers, made of ſix leaues moſt vſually, but ſometimes they will haue ſeuen or eight, being very large, and of a perfect purple Violet colour, very faire and liuely: the middle head hath many blackiſh thrums or threads about it, which I could neuer obſerue in my Gardens to beare ſeed: the roote is ſmaller, and more ſpreading euery way into ſmall long flat tuberous parts, then any other kindes of ſingle or double Anemones.

*Carnea pallida.* There is another very like in leafe and roote vnto the former, but the flower is nothing ſo large, and is whitiſh, tending to a bluſh colour, and of a deeper bluſh colour toward the bottome of the flower, with blackiſh blew thrums in the middle, and giueth no ſeede that I could euer obſerue.

*Carnea viuida unguibus albis.* There is likewiſewiſe another like vnto the laſt in leafe and flower, but that the flower is larger then it, and is of a liuely bluſh colour, the leaues hauing white bottomes.

*Alba venis purpureis.* And another, whoſe flower is white, with purple coloured veines and ſtripes through euery leafe, and is a leſſer flower then the other.

*Anemone tenuifolia coccinea ſimplex.* The ſingle Scarlet Anemone with thin leaues.

The leaues of this Scarlet Windflower are ſomewhat like vnto the former, but a little broader, and not ſo finely cut and diuided: the flower conſiſteth of ſix reaſonable large leaues, of an excellent red colour, which we call a Scarlet; the bottomes of the leaues are large and white, and the thrums or threads in the middle of a blackiſh purple colour: the roote is tuberous, but conſiſting of thicker peeces, ſomewhat like vnto the rootes of the broad leafed Anemones, but ſomewhat browne, and not ſo blacke, and moſt like vnto the roote of the double Scarlet Anemone.

*Coccinea abſq; unguibus.* There is another of this kinde, whoſe flower is neare vnto the ſame colour, but this hath no white bottomes at all in his leaues.

We

*Flore holosericeo.* We haue another which hath as large a flower as any single, and is of an Orient deepe red crimson Veluet colour.

*Sanguinea.* There is another of a deeper red colour, and is called, The bloud red single Anemone.

*Rubra fundo luteo.* And another, whose flower is red with the bottomes yellow.

*Coccinea dilutior.* Another of a perfect crimson colour, whereof some haue round pointed leaues, and others sharpe pointed, and some a little lighter or deeper then others.

*Alba staminibus purpureis.* There is also one, whose flower is pure white with blewish purple thrums in the middle.

*Carnea Hispanica.* And another, whose flower is very great, of a kinde of sullen blush colour, but yet pleasant, with blewish threads in the middle.

*Alba carneis venis.* And another with blush veines in euery leafe of the white flower.

*Alba purpureis unguibus.* And another, the flower whereof is white, the bottomes of the leaues being purple.

*Purpurascens.* Another whose flower consisteth of many small narrow leaues, of a pale purple or blush colour on the outside, and somewhat deeper within.

*Facie florum pomi simplex.* There is another like in leafe and roote vnto the first Scarlet Anemone, but the flower hereof consisteth of seuen large leaues without any bottomes, of a white colour, hauing edges, and some large stripes also of a carnation or flesh colour to bee seene in them, marked somewhat like an Apple blossome, and thereupon it is called in Latine, *Anemone tenuifolia simplex alba instar florum pomi*, or *facie florum pomi*, that is to say in English, The single thin leafed Anemone with Apple blossome flowers.

*Multiplex.* I haue heard that there is one of this kinde with double flowers.

1. *Anemone tenuifolia flore coccineo pleno vulgaris.*
The common double red or Scarlet Anemone.

The leaues of this double Anemone are very like vnto the leaues of the single Scarlet Anemone, but not so thin cut and diuided as that with the purple flower: the flower hereof when it first openeth it selfe, consisteth of six and sometimes of seuen or eight broad leaues, of a deepe red, or excellent Scarlet colour, the middle head being thick closed, and of a greenish colour, which after the flower hath stood blowne some time, doth gather colour, and openeth it selfe into many small leaues, very thicke, of a more pale red colour, and more Stamell like then the outer leaues: the root of this is thicke and tuberous, very like vnto the root of the single Scarlet Anemone.

2. *Anemone tenuifolia flore coccineo pleno variegata.*
The party coloured double Crimson Anemone.

We haue a kinde hereof, varying neither in roote, leafe, or forme of flower from the former, but in the colour, in that this will haue sometimes the outer broad leaues party coloured, with whitish or blush coloured great streakes in the red leaues both inside and outside; as also diuers of the middle or inner leaues striped in the same manner: the roote hereof giueth fairer flowers in some yeares then in others, and sometimes giue flowers all red againe.

3. *Anemone tenuifolia flore coccineo saturo pleno.*
The double crimson Veluet Anemone.

Wee haue another also, whose flower is of a deepe Orenge tawny crimson colour, neare vnto the colour of the outer leaues, of the lesser French Marigold, and not differing from the former in any thing else.

4. *Anemone tenuifolia flore pleno suauerubente.* The greater double blush Anemone.

There is small difference to be discerned, either in the roote or leaues of this from the

the former double Scarlet Anemone, sauing that the leaues hereof are a little broader, and seeme to bee of a little fresher greene colour: the flower of this is as large almost, and as double as the former, and the inner leaues likewise almost as large as they, being of a whitish or flesh colour at the first opening of them, but afterwards become of a most liuely blush colour; the bottomes of the leaues abiding of a deeper blush, and with long standing, the tops of the leaues will turne almost wholly white againe.

5. *Anemone tenuifolia flore albo pleno.* The double white Anemone.

This double white Anemone differeth little from the former blush Anemone, but in that it is smaller in all the parts thereof, and also that the flower hereof being wholly of a pure white colour, without any shew of blush therein, hath the middle thrummes much smaller and shorter then it, and not rising vp so high, but seeme as if they were chipped off euen at the toppes.

6. *Anemone tenuifolia flore pleno albicante.* The lesser double blush Anemone.

This small double blush Anemone differeth very little from the double white last recited, but onely in the colour of the flower: for they are both much about the bignesse one of another, the middle thrums likewise being as small and short, and as euen aboue, onely the flower at the first opening is almost white, but afterwards the outer leaues haue a more shew of blush in them, and the middle part a little deeper then they.

7. *Anemone tenuifolia flore pleno purpureo violaceo.* The double purple Anemone.

This double purple Anemone is also of the same kindred with the first double red or Scarlet Anemone for the form or doubleneſſe of the flower, consisting but of six or seuen leaues at the most in this our Country, although in the hotter it hath ten or twelue, or more as large leaues for the outer border, and as large small leaues for the inner middle also, and almost as double, but of a deepe purple tending toward a Violet colour, the outer leaues being not so deepe as the inner: the roote and leafe commeth neare vnto the single purple Anemone before described, but that the roote spreadeth not so small and so much.

8. *Anemone tenuifolia flore pleno purpureo cæruleo.*
The double blew Anemone.

This Anemone differeth not in any thing from the former double purple, but onely that the flower is paler, and more tending to a blew colour.

9. *Anemone tennifolia flore pleno roseo.* The double Rose coloured Anemone.

The double Rose coloured Anemone differeth also in nothing from the former double purple, but onely in the flower, which is somewhat smaller, and not so thicke and double, and that it is of a reddish colour, neare vnto the colour of a pale red Rose, or of a deepe coloured Damaske.

10. *Anemone tennifolia flore pleno carneo viuacissimo.*
The double Carnation Anemone.

This Anemone, both in roote, leafe, and flower, commeth neerest vnto the former double white Anemone, for the largeneſſe and doubleneſſe of the flower, and in the smalneſſe of the middle thrums, and euenneſſe at the toppes of them, being not so large and great a flower as the double purple, either in the inner or outter leaues, but yet is very faire, thicke and double, and of a most liuely Carnation silke colour, very deepe, both the outer leaues and middle thrums also so bright, that it doth as it were amaze, and yet delight the minde of the beholder, but by long standing in the Sun, waxe a little paler, and so passe away as all the most beautifull flowers doe.

1 *Anemone tenuifolia simplex purpurea.* The single purple Anemone with thin cut leaues. 2 *Anemone tenuifolia simplex alba pura.* The single pure white Anemone. 3 *Anemone tenuifolia simplex chermesina.* The single bright Crimson Anemone. 4 *Anemone tenuifolia simplex sanguinea.* The single bloudred Anemone. 5 *Anemone tenuifolia simplex facie florum pomi.* The single Apple bloome Anemone. 6 *Anemone tenuifolia simplex purpurascens.* The single purplish blush Anemone. 7 *Anemone tenuifolia simplex alba unguibus carnea.* The single white Anemone with blush bottomes. 8 *Anemone tenuifolia flore pleno coccineo.* The double red or ordinary Scarlet Anemone. 9 *Anemone tenuifolia flore pleno rubrofusca coma Amarantina.* The double purple Velues Anemone. 10 *Anemone tenuifolia flore pleno purpureo violaceo.* The double blewish purple Anemone. 11 *Anemone tenuifolia flore pleno incarnedini colorn sericei vivacissimi.* The double Carnation Anemone, or of a liuely Carnation silke colour.

11. *Anemone tenuifolia flore rubrofusco pleno coma Amarantina.*
The double purple Veluet Anemone.

This double Veluet Anemone is in all things like the last described Carnation Anemone, but somewhat larger, the difference consisteth in the colour of the flower, which in this is of a deep or sad crimson red colour for the outer leaues, and of a deep purple Veluet colour in the middle thrums, resembling the colour of the lesser *Amaranthus purpureus*, or Purple flower gentle hereafter described, whereof it tooke the name, which middle thrums are as fine and small, and as euen at the toppes as the white or last Carnation Anemones.

12. *Anemone tenuifolia flore pleno tricolor.*
The double purple Veluet Anemone of three colours.

This double Anemone also is very like the last described Anemone, but that in the middle of the purple thrums, there thrusteth forth a tuft of threads or leaues of a more light crimson colour.

And thus much for the kindes of Anemones or Windflowers, so farre forth as haue hitherto come to our knowledge; yet I doubt not, but that more varieties haue beene elsewhere collected, and will be also in our Countrey daily and yearly obserued by diuers, that raise them vp from sowing the seede, wherein lyeth a pretty art, not yet familiarly knowne to our Nation, although it be very frequent in the Lowe-Countries, where their industry hath bred and nourished vp such diuersities and varieties, that they haue valued some Anemones at such high rates, as most would wonder at, and none of our Nation would purchase, as I thinke. And I doubt not, if wee would be as curious as they, but that both our ayre and soyle would produce as great variety, as euer hath been seene in the Lowe-Countries; which to procure, if any of our Nation will take so much paines in sowing the seedes of Anemones, as diuers haue done of Tulipas: I will set them downe the best directions for that purpose that I haue learned, or could by much search and tryall attaine vnto; yet I must let them vnderstand thus much also, that there is not so great variety of double flowers raised from the seede of the thin leafed Anemones, as from the broad leafed ones.

First therefore (as I said before) concerning Tulipas, there is some speciall choice to be made of such flowers, whose seed is fittest to be taken. Of the *Latifolias*, the double Orenge tawney seede being sowne, yeeldeth pretty varieties, but the purples, and reds, or crimsons, either *Latifolias* or *Tenuifolias*, yeeld small variety, but such as draw neerest to their originall, although some be a little deeper or lighter then others. But the light colours be they which are the chiefe for choice, as white, ash-colour, blush or carnation, light orenge, simple or party coloured, single or double, if they beare seede, which must bee carefully gathered, and that not before it bee thorough ripe, which you shall know by the head; for when the seede with the wollinesse beginneth to rise a little of it selfe at the lower end, it must bee then quickly gathered, lest the winde carry it all away. After it is thus carefully gathered, it must be laid to dry for a weeke or more, which then being gently rubbed with a little dry sand or earth, will cause the seede to be somewhat better separated, although not thoroughly from the woollinesse or downe that compasseth it.

Within a moneth at the most after the seede is thus gathered and prepared, it must be sowne; for by that meanes you shall gaine a yeare in the growing, ouer that you should doe if you sowed it in the next Spring.

If there remaine any woollinesse in the seede, pull it in sunder as well as you can, and then sowe your seede reasonable thin, and not too thicke, vpon a plaine smooth bed of fine earth, or rather in pots or tubbes, and after the sowing, sift or gently straw ouer them some fine good fresh mould, about one fingers thickenesse at the most for the first time: And about a moneth after their first springing vp, sift or straw ouer them in like manner another fingers thickenesse of fine earth, and in the meane time if the weather proue dry, you must water them gently and often, but not to ouerglut them with moisture; and thus doing, you shall haue them spring vp before Winter, and

grow

grow pretty strong, able to abide the sharpe Winter in their nonage, in vsing some little care to couer them loosely with some fearne, or furse, or beane hame, or straw, or any such, which yet must not lye close vpon them, nor too farre from them neither.

The next Spring after the sowing, if you will, but it is better if you stay vntill August, you may then remoue them, and set them in order by rowes, with sufficient distance one from another, where they may abide, vntill you see what manner of flower each plant will beare, which you may dispose of according to your minde.

Many of them being thus ordered (if your mould be fine, loose, and fresh, not stonie, clayish, or from a middin) will beare flowers the second yeare after the sowing, and most or all of them the third yeare, if the place where you sowe them, be not annoyed with the smoake of Brewers, Dyers, or Maultkils, which if it be, then will they neuer thriue well.

Thus much haue I thought good to set downe, to incite some of our owne Nation to be industrious; and to helpe them forward, haue giuen such rules of directions, that I doubt not, but they will vpon the tryall and view of the variety, proceede as well in the sowing of Anemones as of Tulipas.

I cannot (Gentlewomen) withhold one other secret from you, which is to informe you how you may so order Anemones, that after all others ordinarily are past, you may haue them in flower for two or three moneths longer then are to be seene with any other, that vseth not this course I direct you.

The ordinary time to plant Anemones, is most commonly in August, which will beare flower some peraduenture before Winter, but most vsually in February, March, and Aprill, few or none of them abiding vntill May; but if you will keepe some roots out of the ground vnplanted, vntill February, March, and Aprill, and plant some at one time, and some at another, you shall haue them beare flower according to their planting, those that shall be planted in February, will flower about the middle or end of May, and so the rest accordingly after that manner: And thus may you haue the pleasure of these plants out of their naturall seasons, which is not permitted to be enioyed in any other that I know, Nature being not so prone to bee furthered by art in other things as in this. Yet regard, that in keeping your Anemonie rootes out of the ground for this purpose, you neither keep them too dry, nor yet too moist, for sprouting or rotting; and in planting them, that you set them not in too open a sunny place, but where they may be somewhat shadowed.

## The Place.

I shall not need to spend much time in relating the seuerall places of these Anemones, but onely to declare that the most of them that haue not beene raised from seed, haue come from Constantinople to vs; yet the first broad leafed or yellow Anemone, was first found in Portugall, and from thence brought into these parts. And the first purple Starre Anemone in Germanie, yet was the same sent among others from Constantinople also. And the first thin cut leafed Anemone came first out of Italy, although many of that sort haue come likewise from Constantinople. And so haue the double red or Scarlet Anemones, and the great double blush, which I first had by the gift of Mr. Humfrey Packington of Worcestershire Esquire, at Haruington.

## The Time.

The times of their flowring are sufficiently expressed in the descriptions, or in the rules for planting.

## The Names.

The Turkish names whereby the great double broad leafed kindes haue beene sent vnto vs, were *Giul Catamer*, and *Giul Catamer lale*; And *Binizade*, *Binizante*, and *Galipoli lale* for the thinne cut leafed Anemones. All Authors haue called them *Anemones*, and are the true *Herba venti*.

We

Wee call them in Englisſh eyther Anemones, after the Greeke name, or Windflowers, after the Latine.

### The Vertues.

There is little vſe of theſe in Phyſicke in our dayes, eyther for inward or outward diſeaſes; onely the leaues are vſed in the Ointment called *Marciatum*, which is compoſed of many other hot herbes, and is vſed in cold griefes, to warme and comfort the parts. The roote, by reaſon of the ſharpeneſſe, is apt to drawe downe rheume, if it be taſted or chewed in the mouth.

## CHAP. XXVI.

### *Aconitum.* Wolfebane.

THere be diuers ſorts of Wolfebanes which are not fit for this booke, but are reſerued for a generall Hiſtory or Garden of Simples, yet among them there are ſome, that notwithſtanding their euill quality, may for the beauty of their flowers take vp a roome in this Garden, of whom I meane to entreate in this place: And firſt of the Winter Wolfesbane, which for the beauty, as well as the earlineſſe of his flowers, being the firſt of all other, that ſhew themſelues after Chriſtmas, deſerueth a prime place; and therefore for the likeneſſe of the rootes vnto the Anemones, I ioyne it next vnto them.

#### 1. *Aconitum Hyemale.* The Winters Wolfesbane.

This little plant thruſteth vp diuers leaues out of the ground, in the deepe of Winter oftentimes, if there be any milde weather in Ianuary, but moſt commonly after the deepe froſts, bearing vp many times the ſnow vpon the heads of the leaues, which like vnto the Anemone, doe euery leafe riſe from the roote vpon ſeuerall ſhort foote-ſtalkes, not aboue foure fingers high, ſome hauing flowers in the middle of them, (which come vp firſt moſt vſually) and ſome none, which leaues ſtand as it were round, the ſtalke riſing vp vnder the middle of the leafe, deeply cut in and gaſhed to the middle ſtalke almoſt, of a very faire deepe greene colour, in the middle whereof, cloſe vnto the leafe, ſtanderh a ſmall yellow flower, made of ſix leaues, very like a Crowfoote, with yellow threads in the middle: after the flower is fallen, there riſe vp diuers ſmall hornes or cods ſet together, wherein are contained whitiſh yellow round ſeede. The roote is tuberous, ſo like both for ſhape and colour vnto the rootes of Anemones, that they will eaſily deceiue one not well experienced, but that it is browner and ſmoother without, and yellow within, if it be broken.

#### 2. *Aconitum flore albido, ſiue Aconitum luteum Ponticum.* The whitiſh yellow Wolfesbane.

This Wolfesbane ſhooteth not out of the ground vntill the Spring be well begun, and then it ſendeth forth great broad greene leaues, deeply cut in about the edges, not much vnlike the leaues of the great wilde Crowfoote, but much greater; from among which leaues riſeth vp a ſtrong ſtiffe ſtalke, three foote high, hauing here and there leaues ſet vpon it, like vnto the loweſt, but ſmaller; the toppe of the ſtalke is diuided into three or foure branches, whereon are ſet diuers pale yellow flowers, which turne at the laſt to be almoſt white, in faſhion like almoſt vnto the flowers of the Helmet flower, but much ſmaller, and not gaping ſo wide open: after the flowers are paſt come vp diuers ſhort poddes, wherein is contained blacke ſeede: the roote is made of a number of darke browne ſtrings, which ſpread and faſten themſelues ſtrongly in the ground.

3. *Napellus*

3. *Napellus verus flore cæruleo.* Blew Helmet flower or Monkes hood.

The Helmet flower hath diuers leaues of a fresh greene colour on the vpperside, and grayish vnderneath, much spread abroad and cut into many slits and notches, more then any of the Wolfebanes; the stalke riseth vp two or three foot high, beset to the top with the like leaues, but smaller: the toppe is sometimes diuided into two or three branches, but more vsually without, whereon stand many large flowers one aboue another, in forme very like vnto a Hood or open Helmet, being composed of fiue leaues, the vppermost of which and the greatest, is hollow, like vnto an Helmet or Headpeece, two other small leaues are at the sides of the Helmet, closing it like cheekes, and come somewhat vnder, and two other which are the smallest hang down like labels, or as if a close Helmet were opened, and some peeces hung by, of a perfect or faire blew colour, (but grow darker, hauing stood long) which causeth it be so nourished vp in Gardens, that their flowers, as was vsuall in former times, and yet is in many Countrey places, may be laid among greene herbes in windowes and roomes for the Summer time: but although their beauty may be entertained for the vses aforesaid, yet beware they come not neare your tongue or lippes, lest they tell you to your cost, they are not so good as they seeme to be: in the middest of the flower, when it is open and gapeth wide, are seene certaine small threads like beards, standing about a middle head, which when the flower is past, groweth into three or foure, or more small blackish pods, containing in them blacke seede: the rootes are browniish on the outside, and white within, somewhat bigge and round aboue, and small downewards, somewhat like vnto a small short Carrot roote, sometimes two being ioyned at the head together. But the name *Napellus* anciently giuen vnto it, doth shew they referred the forme of the roote vnto a small Turnep.

*Anthora.* The wholsome Helmet flower, or counterpoison Monkes hood.

This wholsome plant I thought good to insert, not onely for the forme of the flower, but also for the excellent properties thereof, as you shall haue them related hereafter. The rootes hereof are small and tuberous, round and somewhat long, ending for the most part in a long fibre, and with some other small threads from the head downeward: from the head whereof riseth vp diuers greene leaues, euery one seuerally vpon a stalke, very much diuided, as finely almost as the leaues of Larkes heeles or spurres: among which riseth vp a hard round stalke, a foote high and better, with some such leaues thereon as grow belowe, at the toppe whereof stand many small yellowish flowers, formed very like vnto the former whitish Wolfesbane, bearing many blacke seedes in pods afterwards in the like manner.

Many more sorts of varieties of these kindes there are, but these onely, as the most specious, are noursed vp in Florists Gardens for pleasure; the other are kept by such as are Catholicke obseruers of all natures store.

The Place.

All these grow naturally on Mountaines, in many shadowie places of the Alpes, in Germany, and elsewhere.

The Time.

The first flowreth (as is said) in Ianuary, and February, and sometimes vntill March be well spent, and the seede is soone ripe after.

The other three flower not vntill Iune and Iuly.

The Names.

The first is vsually called *Aconitum hyemale Belgarum.* Lobelius calleth it *Bulbosum*

*Bulbosus vnifolius Batrachoides*, *Aconitum Elleboraceum*, and *Ranunculus Monophyllos*, and some by other names. Most Herbarists call it *Aconitum hyemale*, and we in English thereafter, Winters Wolfesbane; and of some, Yellow Aconite.

The second is called by most Writers, *Aconitum luteum Ponticum*: Some also *Lupicida*, *Luparia*, and *Canicida*, of the effect in killing Wolues and Dogs: And some, because the flower is more white then yellow, doe call it *Aconitum flore albido*, we call it in English, The whitish yellow Aconite, or Wolfesbane, but some afrer the Latine name, The yellow Wolfesbane.

The third is called generally *Napellus*, and *Verus*, because it is the true *Napellus* of the ancient Writers, which they so termed from the forme of a Turnep, called *Napus* in Latine.

The fourth is called *Aconitum Salutiferum*, *Napellus Moysis*, *Antora* and *Anthora*, *quasi Antithora*, that is, the remedy against the poisonfull herbe *Thora*, in English according to the title, eyther wholsome Helmet flower, or counterpoison Monkes hood.

### The Vertues.

Although the first three sorts of plants be very poisonfull and deadly, yet there may bee very good vse made of them for sore eyes (being carefully applyed, yet not to all sorts of sore eyes neither without discretion) if the distilled water be dropped therein.

The rootes of the counterpoison Monkes hood are effectuall not onely against the poison of the poisonfull Helmet flower, and all others of that kinde, but also against the poison of all venemous beasts, the plague or pestilence, and other infectious diseases, which raise spots, pockes, or markes in the outward skinne, by expelling the poison from within, and defending the heart as a most soueraigne Cordiall. It is vsed also with good successe against the wormes of the belly, and against the paines of the Wind collick.

## Chap. XXVII.

### *Ranunculus*. The Crowfoote.

NExt vnto the Aconites, of right are to follow the *Ranunculi*, or Crowfeete, for the neareneſſe both of forme, of leaues, and nature of the plants, although lesse hurtfull, yet all of them for the most part being sharpe and exulcerating, and not without some danger, if any would be too bold with them. The whole Family of the *Ranunculi* is of a very large extent, and I am constrained within the limits of a Garden of Pleasure; I must therefore select out onely such as are fit for this purpose, and set them here downe for your knowledge, leauing the rest for that other generall worke, which time may perfect and bring to light, if the couetous mindes of some that should be most affected towards it, doe not hinder it: or if the helpe of generous spirits would forward it.

#### 1. *Ranunculus montanus albus humilior*. The lowe white mountaine Crowfoot.

This lowe Crowfoote hath three or foure broad and thicke leaues, almost round, yet a little cut in and notched about the edges, of a fine greene and shining colour on the vpperside, and not so green vnderneath, among which riseth a small short stalke, bearing one snow white flower on the toppe, made of fiue round pointed leaues, with diuers yellow threads in the middle, standing about a greene head, which in time groweth to be full of seede, in forme like vnto a small greene Strawberry: the roote is composed of many white strings.

*Duplici flore.* There is another of this lowe kinde, whose leaues are somewhat more deeply cut in on the edges, and the flower larger, and sometimes a little double, as it were with two rowes of leaues, in other things not differing from the former.

2. *Ranunculus*

2. *Ranunculus montanus albus maior vel elatior.*
The great single white mountaine Crowfoote.

The leaues of this Crowfoote are large and greene, cut into three, and sometimes into fiue speciall diuisions, and each of them besides cut or notched about the edges, somewhat resembling the leaues of the Globe Crowfoote, but larger: the stalke is two foote and a halfe high, hauing three small leaues set at the ioynt of the stalke, where it brancheth out into flowers, which stand foure or fiue together vpon long foote-stalkes, made of fiue white leaues a peece, very sweete, and somewhat larger then the next white Crowfoote, with some yellow threads in the middle compassing a greene head, which bringeth seede like vnto other wilde Crowfeete: the roote hath many long thicke whitish strings, comming from a thicke head.

3. *Ranunculus montanus albus minor.* The lesser single white Crowfoote.

This Crowefoote hath faire large spread leaues, cut into fiue diuisions, and somewhat notched about the edges, greene on the vpperside, and paler vnderneath, hauing many veines running through the leaues: the stalke of this riseth not so high as the former, although this be reasonable tall, as being neare two foote high, spread into many branches, bearing such like white flowers, as in the former, but smaller: the seede of this is like the former, and so are the rootes likewise.

4. *Ranunculus albus flore pleno.* The double white Crowfoot.

The double white Crowfoote is of the same kinde with the last single white Crowfoote, hauing such like leaues in all respects: the onely difference is in the flowers, which in this are very thicke and double. Some doe make mention of two sorts of double white Crowfeete, one somewhat lower then another, and the lower likewise bearing more store of flowers, and more double then the higher: but I confesse, I haue neuer seene but one sort of double, which is the same here expressed, not growing very high, and reasonably well stored with flowers.

5. *Ranunculus præcox Rutæfolio siue Coriandrifolio.*
The early Coriander leafed Crowfoote.

This Crowfoote hath three or foure very greene leaues, cut and diuided into many small peeces, like vnto the wing of leaues of Rue, or rather like the lower leaues of the Coriander (for they well resemble either of them) euery of them standing vpon a long purplish stalke, at the toppe whereof groweth the flower alone, being composed or made of twelue small white leaues, broad pointed, and a little endented at the ends, somewhat purplish on the outside, and white on the inside, sustained by diuers small greene leaues, which are in stead of a cup or huske: in the middle of the flower are many small white threads, tipt with yellow pendents, standing about a small greene head, which after groweth to bee full of seedes like a Strawberry, which knobs giue small blackish seede: the roote is white and fibrous.

6. *Ranunculus Thalictrifolio maior.* The great colombine leafed Crowfoot.

The lower leaues of this Crowfoote haue long stalkes, and are very like vnto the smaller leaues of Colombines, or the great Spanish *Thalictrum*, which hath his leaues very like vnto a Colombine, foure or fiue rising from the roote: the stalke riseth about a foote and a halfe high, somewhat reddish, beset here and there with the like leaues, at the toppe whereof stand diuers small white flowers, made of fiue leaues a peece, with some pale white threads in the middle: the seede is round and reddish, contained in small huskes or hornes: the roote is made of a bush or tuft of white strings.

7. *Ranunculus Thalictrifolio minor Asphodeli radice.*
The small white Colombine leafed Crowfoote.

This small Crowfoote hath three or foure winged leaues spread vpon the ground, standing vpon long stalkes, and consisting of many small leaues set together, spreading from the middle ribbe, euery leafe somewhat resembling both in shape and colour the smallest and youngest leaues of Colombines: the flowers are white, standing at the toppe of the stalkes, made of fiue round leaues: the root hath three or foure thick, short, and round yellowish clogs hanging at the head, like vnto the Asphodill roote. The great Herball of Lyons, that goeth vnder the name of *Daleschampius*, saith, that Dr. Myconus found it in Spaine, and sent it vnder the name of Oenanthe; and therefore Ioannes Molineus who is thought to haue composed that booke, set it among the vmbelliferous plants, because the Oenanthes beare vmbels of flowers and seede, and haue tuberous or cloggy rootes; but with what iudgement, let others say, when they haue compared the vmbels of flowers and seede of the Oenanthes, with the flowers and seede of this plant, and whether I haue not more properly placed it among the *Ranunculi* or Crowfeete, and giuen it a denomination agreeable to his forme.

8. *Ranunculus Globosus.* The Globe Crowfoot.

This Crowfoote (which in the Northerne countries of England where it groweth plentifully, is called Locker goulous) hath many faire, broad, darke greene leaues next the ground, cut into fiue, sixe, or seuen diuisions, and iagged besides at the edges; among which riseth vp a stalke, whereon are set such like leaues as are belowe, but smaller, diuided toward the toppe into some branches, on the which stand seuerall large yellow flowers, alwayes folded inward, or as a close flower neuer blowing open, as other flowers doe, consisting of eleuen leaues for the most part, set or placed in three rowes, with many yellow threads in the middle, standing about a greene rough head, which in time groweth to be small knops, wherein are contained blacke seede: the roote is composed of many blackish strings.

9. *Ranunculus pratensis flore multiplici.* The double yellow field Crowfoot.

There is little or no difference in the leaues of this double Crowfoot, from those of the single kindes that growe in euery medowe, being large and diuided into foure or fiue parts, and indented about the edges, but they are somewhat smaller, and of a fresher greene: the flowers stand on many branches, much diuided or separated, being not very great, but very thicke and double: the roote runneth and creepeth vnder ground like as the single doth.

10. *Ranunculus Anglicus maximus multiplex.*
The Garden double yellow Crowfoot or Batchelours buttons.

This great double Crowfoote, which is common in euery Garden through England, hath many great blackish greene leaues, iagged and cut into three diuisions, each to the middle ribbe: the stalkes haue some smaller leaues on them, and those next vnder the branches long and narrow: the flowers are of a greenish yellow colour, very thicke and double of leaues, in the middle whereof riseth vp a small stalke, bearing another double flower, like to the other, but smaller: the roote is round, like vnto a small white Turnep, with diuers other fibres annexed vnto it.

11. *Ranunculus Gramineus.* Grasse leafed Crowfoot.

The leaues of this Crowfoote are long and narrow, somewhat like vnto Grasse, or rather like the leaues of single Gilloflowers or Pinckes, being small and sharpe pointed, a little hollow, and of a whitish greene colour: among these leaues rise vp diuers slender stalkes, bearing one small flower at the toppe of each, consisting of fiue yellow leaues,

1 *Aconitum Hyemale.* Winter Wolfesbane. 2 *Aconitum flore albido sive luteum Ponticum.* The whitish yellow Wolfesbane. 3 *Napellus verus.* Blew Helmets or Monkes hood. 4 *Anthora.* The counterpoison Monkes hood. 5 *Ranunculus humilis albus simplex.* The single white low Crowfoot. 6 *Ranunculus humilis albus duplici flore.* The double lowe white Crowfoot. 7 *Ranunculus Coriandrifolio* The early Coriander leaved Crowfoot. 8 *Ranunculus montanus elatior albus.* The great single white mountain Crowfoot. 9 *Ranunculus montanus albus flore pleno* The double white mountain Crowfoot. 10 *Ranunculus Thalictrifolio minor.* The lesser Colombine leaved Crowfoot. 11 *Ranunculus globosus.* The globe Crowfoot.



2. Aconitum Lycoctonum, L. mill. dict.

leaues, with some threads in the middle: the roote is composed of many thicke, long, round white strings.

There is another of this kinde that beareth flowers with two rowes of leaues, as if it were double, differing in nothing else.

12. *Ranunculus Lusitanicus Autumnalis.* The Portugall Autumne Crowfoot.

This Autumne Crowfoote hath diuers broad round leaues lying on the ground, set vpon short foote-stalkes, of a faire greene colour aboue, and grayish vnderneath, snipt all about the edges, hauing many veines in them, and sometimes swelling as with blisters or bladders on them; from among which rise vp two or three slender and hairy stalkes, bearing but one small yellow flower a peece, consisting of fiue and sometimes of six leaues, and sometimes of seuen or eight, hauing a few threads in the middle, set about a small greene head, like vnto many of the former Crowfeete, which bringeth small blacke seede: the roote is made of many thicke short white strings, which seeme to be grumous or kernelly rootes, but that they are somewhat smaller, and longer then any other of that kinde.

13. *Ranunculus Creticus latifolius.* The broad leafed Candy Crowfoot.

This Crowfoote of Candy, hath the greatest and broadest leaues of all the sorts of Crowfeete, being almost round, and without any great diuisions, but onely a few notches about the edges here and there, as large or larger sometimes then the palme of a mans hand; among which riseth vp the stalke, not very high when it doth first flower, but afterwards, as the other flowers doe open themselues, the stalke groweth to be a foote and a halfe high, or thereabouts, hauing some leaues on it, deeply cut in or diuided, and bearing many faire yellow flowers, consisting of fiue leaues a peece, being somewhat whitish in the middle, when the flower hath stood blowne a little time: the roote is composed of a number of small kernelly knobs, or long graines, set thicke together. This flowreth very early, being vsually in flower before the end of March, and oftentimes about the middle thereof.

14. *Ranunculus Creticus albus.* The white Candy Crowfoote.

The leaues of this Crowfoote are very like vnto the leaues of the red Crowfoote of Tripoli or Asia, hereafter set downe, being somewhat broad and indented about the edges, some of the leaues being also cut in or gashed, thereby making it as it were three diuisions, of a pale greene colour, with many white spots in them: the stalke riseth vp a foote high, with some leaues on it, more diuided then the lower, and diuided at the toppe into two and sometimes into three branches, each of them bearing a faire snow white flower, somewhat large, included at the first in a browniish huske or cup of leaues, which afterwards stand vnder the flowers, consisting of fiue white large round pointed leaues, in the middle whereof is set many blackish purple thrums, compassing a small long greene head, composed of many scales or chaffie whitish huskes, when they are ripe, which are the seede, but vnprofitable in all that euer I could obserue: the rootes are many small graines or kernels, set together as in the former, and much about the same colour, that is, of a darke or duskie grayish colour, but much smaller.

*Alba purpureis oris & venis.* There is another of this kinde, whose flowers haue purple edges, and sometimes some veines of the same purple in the leaues of the flowers, not differing in any other thing from the former.

*Alba oris rubris.* And another, whose edges of the flowers are of a bright red colour.

15. *Ranunculus Creticus flore argenteo.* The Argentine, or cloth of siluer Crowfoot.

The greene leaues of this Crowfoote are as small and thinne, cut in or diuided on the edges, as the last two sorts; the stalke riseth vp somewhat higher, and diuided into some branches, bearing at the toppe of euery of them one flower, somewhat smaller then the former, composed of six, seuen, and sometimes of eight small round pointed

1 *Ranunculus gramineus flore simplici & duplici.* The single and the double grasse Crowfoot. 2 *Ranunculus Lusitanicus Autumnalis.* The Portugall Autumne Crowfoot. 3 *Ranunculus Creticus latifolius.* The broad leafed Candy Crowfoot. 4 *Ranunculus Anglicus maximus multiplex.* The double English Crowfoot. 5 *Ranunculus pratensis flore multiplici.* The double yellow field Crowfoot. 6 *Ranunculus Creticus albus.* The white Candy Crowfoot. 7 *Ranunculus Asiaticus flore albo vel pallido vario.* The white or the straw coloured Crowfoot with red tops & edges. 8 *Ranunculus Tripolitanus flore rubro simplici.* The single red Crowfoot of Tripoli. 9 *Ranunculus Asiaticus flore rubro amplo.* The large single red Crowfoot of Asia. 10 *Ranunculus Asiaticus flore rubro pleno.* The double red Crowfoot of Asia. 11 *Caltha palustris flore pleno.* Double Marsh Marigold or Batchelours buttons.

leaues, of a whitish yellow blush colour on the inside wholly, except sometimes a little stript about the edges: but the outside of euery leafe is finely stript with crimson stripes, very thicke, somewhat like vnto a Gilloflower: in the middle riseth vp a small blacke head, compassed about with blackish blew threads or thrums, which head is as vnfruitfull for seede in our Countrey as the former. This flower hath no such greene leaues vnder it, or to enclose it before it be blowne open as the former: the roots are in all things like the former.

16. *Ranunculus Asiaticus siue Tripolitanus flore rubro.*
The single red Crowfoote of Asia or Tripoli.

The lower leaues of this red Crowfoote are alwayes whole without diuisions, being onely somewhat deeply indented about the edges, but the other that rise after them are more cut in, sometimes into three, and sometimes into fiue diuisions, and notched also about the edges: the stalke riseth higher then any of the former, and hath on it two or three smaller leaues, more cut in and diuided then those belowe: at the toppe whereof standeth one large flower, made of fiue leaues, euery one being narrower at the bottome then at the toppe, and not standing close and round one to another, but with a certaine distance betweene, of a duskie yellowish red colour on the outside, and of a deepe red on the inside, the middle being set with many thrums of a darke purple colour: the head for seede is long, and scaly or chaffie, and idle in like manner as the rest: the roote is made of many graines or small kernels set together, and closing at the head, but spreading it selfe, if it like the ground, vnder the vpper crust of the earth into many roots, encreasing from long strings, that runne from the middle of the small head of graines, as well as at the head it selfe.

17. *Ranunculus Asiaticus flore amplo rubro.* The large single red Crowfoot of Asia.

There hath come to vs out of Turkie, together with the former, among many other roots, vnder the same title, a differing sort of this Crowfoote, whose leaues weare broader, and much goaler; the flower also larger, and the leaues thereof broader, sometimes eight in a flower, standing round and close one to another, which maketh the fairer shew: in all other things it is like the former.

18. *Ranunculus Asiaticus flore rubro vario simplici.*
The red stript single Crowfoote of Asia.

This party coloured Crowfoote differeth not eyther in roote or leafe from the former, the chiefest difference is in the flower, which being red, somewhat like the former, hath yet some yellow stripes or veines through euery leafe, sometimes but little, and sometimes so much, that it seemeth to bee party coloured red and yellow: this sort is very tender; for we haue twice had it, and yet perished with vs.

19. *Ranunculus Asiaticus flore luteo vario simplici.*
The yellow stript single Crowfoote of Asia.

There is little difference in the roote of this Crowfoote from the last described, but the leaues are much different, being very much diuided, and the flower is large, of a fine pale greenish yellow colour, consisting of six and seuen, and sometimes of eight or nine round leaues; the toppes whereof haue reddish spots, and the edges sometimes also, with such purplish thrums in the middle that the other haue. None of these former Crowfeete with kernelly roots, haue euer beene found to haue giuen so good seed in England, as that being sowne, any of them would spring vp; for hereof tryall hath been often made, but all they haue lost their labour, that haue bestowed their paines therein, as farre as I know.

20. *Ranunculus Asiaticus flore rubro pleno.*
The double red Crowfoote of Asia.

The double red Crowfoote hath his roots and leaues so like vnto the single red kinde, that none can perceiue any differ ice, or know the one from the other, vntill the budde of the flower doe appeare, which after it is any thing forward, may be perceiued to be greater and fuller then the budde of the single kinde. This kinde beareth most vsually but one faire large double flower on the toppe of the stalke, composed of many leaues, set close together in three or foure rowes, of an excellent crimson colour, declining to Scarlet, the outter leaues being larger then the inner; and in stead of thrummes, hath many small leaues set together: it hath likewise six small narrow greene leaues on the backside of the flower, where the stalke is fastened to the flower.

There is of this double kinde another sort, whose flower is of the same colour with the former, but out of the middle of the flower ariseth another double flower, but smaller. *Polifero flore.*

The Place.

These plants grow naturally in diuers Countries; some in France, and Germany, and some in England, some in Spaine, Portugall, and Italy, and some haue been sent out of Turkie from Constantinople, and some from other parts, their titles for the most part descrying their Countries.

The Time.

Some of them flower early, as is set downe in their descriptions, or titles. The others in Aprill and May. The white Candy Crowfoote, and the other single and double sorts of Asia, about the same time, or somewhat later, and one in Autumne, as it is set downe.

The Names.

The names that are giuen seuerally to them may well serue this worke, that thereby they may bee distinguished one from another: For to set downe any further controuersie of names, how fitly or vnfitly they haue beene called, and how variably by diuers former Writers, is fitter for a generall History, vnto which I leaue what may be said, both concerning these and the rest: Onely this I would giue you to vnderstand, that the Turkie kindes haue been sent to vs vnder the names of *Terebolos* for the single, and *Terebolos Catamer lale* for the double, and yet oftentimes, those that haue been sent for double, haue proued single, so little fidelity is to bee found among them.

The Vertues.

All or most of these plants are very sharpe and exulcerating, yet the care and industry of diuers learned men haue found many good effects in many of them. For the roots and leaues both of the wilde kindes, and of some of these of the Garden, stamped and applyed to the wrists, haue driuen away the fits in Feuers. The roote likewise of the double English kinde is applyed for pestilent sores, to helpe to breake them, by drawing the venome to the place. They helpe likewise to take away scarres and markes in diuers places of the body.

## Chap. XXVIII.

### *Caltha palustris flore pleno.* Double Marsh Marigold.

AS an appendix to the Crowfeete, I must needes adde this plant, yet seuerally by it selfe, because both it and his single kinde are by most adioyned thereunto, for the neare resemblance both in shape and sharpenesse of quality. The single kinde I leaue to the Ditch sides, and moist grounds about them, as the fittest places for it, and onely bring the double kinde into my Garden, as fittest for his goodly proportion and beauty to be entertained, and haue place therein.

The double Marsh Marigold hath many broad and round greene leaues, a little endented about the edges, like vnto the single kinde, but not altogether so large, especially in a Garden where it standeth not very moist: the stalkes are weake, round, hollow, and greene, diuided into three or foure branches at the toppe, with leaues at the seuerall ioynts, whereon stand very double flowers, of a gold yellow colour: the fiue outer leaues being larger then any of the rest that are encompassed by them, which fall away after they haue stood blowne a great while (for it endureth in flower a moneth or more, especially if it stand in a shadowie place) without bearing any seed: the roots are composed of many thicke, long, and round whitish strings, which runne downe deep into the ground, and there are fastened very strongly.

#### The Place.

This plant groweth naturally in diuers Marshes, and moist grounds in Germany, yet in some more double then in others; it hath long agoe beene cherished in our Gardens.

#### The Time.

It flowreth in Aprill or May, as the yeare proueth earlier or later: all his leaues doe in a manner quite perish in Winter, and spring anew in the end of February, or thereabouts.

#### The Names.

There is great controuersie among the learned about the single kinde, but thereof I shall not neede to speake in this place; if God permit I may in a fitter. This is called generally in Latine, *Caltha palustris multiplex*, or *flore pleno*. And wee in English (after the Latine, which take *Caltha* to be that which wee vsually call *Calendula*, a Marigold) The double Marsh Marigold.

#### The Vertues.

The roote hereof is sharpe, comming neare vnto the quality of the Crowfeete, but for any speciall property, I haue not heard or found any.

## Chap. XXIX.

*Hepatica nobilis siue trifolia.* Noble Liuerwort.

NExt vnto the Crowfeete are to follow the Hepaticas, because of the likenesse with them, seeming to be small Crowfeete in all their parts, but of another and more wholsome kinde. Their diuersity among themselues consisteth chiefly in the colour of the flowers, all of them being single, except one which is very thicke and double.

1. *Hepatica flore caeruleo simplici maior.*

The great single blew Hepatica or noble Liuerwort.

The flowers of this Hepatica doe spring vp, blow open, and sometimes shed and fall away, before any leaues appeare or spread open. The rootes are composed of a bush of blackish strings, from the seuerall heads or buttons whereof, after the flowers are risen and blowne, arise many fresh greene leaues, each seuerally standing vpon his foot-stalke, folded together, and somewhat browne and hairy at their first comming, which after are broad, and diuided at the edges into three parts: the flowers likewise stand euery one vpon his owne seuerall foote-stalke, of the same height with the leaues for the most part, which is about foure or fiue fingers breadth high, made of six leaues most vsually, but sometimes it will haue seuen or eight, of a faire blew colour, with many white chiues or threads in the middle, standing about a middle green head or vmbone, which after the flower is fallen groweth greater, and sheweth many small graines or seede set close together (with three small greene leaues compassing them vnderneath, as they did the flower at the bottome) very like the head of seed of manie Crowfeete.

2. *Hepatica minor flore pallido caeruleo.* The small blew Hepatica.

The leaues of this Hepatica are smaller by the halfe then the former, and grow more aboundantly, or bushing thicke together: the flowers (when it sheweth them, for I haue had the plant halfe a score yeares, and yet neuer saw it beare flower aboue once or twice) are of a pale or bleake blew colour, not so large as the flowers of the former.

3. *Hepatica flore purpureo.* Purple Hepatica or noble Liuerwort.

This Hepatica is in all things like vnto the first, but onely the flowers are of a deeper blew tending to a Violet purple: and therefore I shall not neede to reiterate the former description.

4. *Hepatica flore albo minor.* The lesser white Hepatica.

The flowers of this Hepatica are wholly white, of the bignesse of the red or purple, and the leaues somewhat smaller, and of a little whiter or paler greene colour, else in all other things agreeing with the former.

5. *Hepatica alba magno flore.* The great white Hepatica.

There is no other difference herein from the last, but that the flower being as white, is as large as the next.

6. *Hepatica albida siue argentea.* Ash-coloured or Argentine Hepatica.

Both the leaues and the flowers of this Hepatica are larger then any of the former, except the last: the flowers hereof at the first opening seeme to bee a of blush ash-colour, which doe so abide three or foure dayes, decaying still vntill it turne almost white,

white, hauing yet still a shew of that blush ash-colour in them, till the very last.

7. *Hepatica alba straminibus rubris.* White Hepatica with red threads.

There is no difference between this Hepatica and the first white one, sauing that the threads in the middle of the flower, being white, as in the former, are tipt at the ends with a pale reddish colour, which adde a great beauty to the flowers.

8. *Hepatica flore rubro.* Red Hepatica or noble Liuerwort.

The leaues of this Hepatica are of a little browner red colour, both at their first comming vp, and afterwards, especially in the middle of the leafe more then any of the former: the flowers are in forme like vnto the rest, but of a bright blush, or pale red colour, very pleasant to behold, with white threads or chiues in the middle of them.

9. *Hepatica flore purpureo multiplici siue pleno.*
The double purple Hepatica.

The double Hepatica is in all things like vnto the single purple kinde, sauing onely that the leaues are larger, and stand vpon longer foote-stalkes, and that the flowers are small buttons, but very thicke of leaues, and as double as a flower can be, like vnto the double white Crowfoote before described, but not so bigge, of a deepe blew or purple colour, without any threads or head in the middle, which fall away without giuing any seede.

10. *Hepatica flore caruleo pleno.* The double blew Hepatica.

In the colour of this flower, consisteth the chiefest difference from the last, except one may say it is a little lesse in the bignesse of the flower, but not in doublenesse of leaues.

## The Place.

All these plants with single flowers grow naturally in the Woods, and shadowie Mountaines of Germany in many places, and some of them in Italy also. The double kinde likewise hath been sent from Alphonsus Pantius out of Italy, as Clusius reporteth, and was also found in the Woods, neare the Castle of Starnbeg in Austria, the Lady Heusenstains possession, as the same Clusius reporteth also.

## The Time.

These plants doe flower very early, and are of the first flowers that shew themselues presently after the deepe frosts in Ianuary, so that next vnto the Winter Wolfesbane, these making their pride appeare in Winter, are the more welcome early guests. The double kinde flowreth not altogether so early, but sheweth his flower, and abideth when the others are past.

## The Names.

They haue obtained diuers names; some calling them *Hepatica*, *Hepatica nobilis*, *Hepaticum trifolium*, *Trifolium nobile*, *Trifolium aureum*, and some *Trinitas*, and *Herba Trinitatis*. In English you may call them either Hepatica, after the Latine name, as most doe, or Noble Liuerwort, which you please.

## The Vertues.

These are thought to coole and strengthen the liuer, the name importing as much; but I neuer saw any great vse of them by any the Physitians of our London Colledge, or effect by them that haue vsed them in Physicke in our Country.

CHAP.

1 *Hepatica flore albo amplo simplici.* The large white Hepatica. 2 *Hepatica flore rubro simplici.* The red Hepatica. 3 *Hepatica flore purpureo pleno.* The double purple Hepatica. 4 *Geranium tuberosum.* Knobbed Cranes bill. 5 *Geranium Batrachoides flore albo vel cæruleo.* The blew or white Crowfoote Cranes bill. 6 *Geranium Hematodes.* The red Rose Cranes bill. 7 *Geranium Romanum striatum.* The variable stript Cranes bill. 8 *Geranium Creticum.* Candy Cranes bill.

7. Geranium striatum: Spr. hist.

## CHAP. XXX.

*Geranium.* Storkes bill or Cranes bill.

AS was ſaid before concerning the Crowfeet, of their large extent and reſtraint, the like may be ſaid of the Storkes bils or Cranes bils; for euen of theſe as of them, I muſt for this worke ſet forth the deſcriptions but of a few, and leaue the reſt to a generall worke.

1. *Geranium tuberoſum vel bulboſum.* Bulbous or knobbed Cranes bill.

The knobbed Cranes hath three or foure large leaues ſpread vpon the ground, of a grayiſh or rather duſty greene colour, euery one of them being as it were of a round forme, but diuided or cut into ſix or ſeuen long parts or diuiſions, euen vnto the middle, which maketh it ſeeme to be ſo many leaues, each of the cuts or diuiſions being deeply notched or indented on both ſides; among which riſeth vp a ſtalke a foote high or better, bearing thereon diuers pale but bright purple flowers, made of fiue leaues a peece, after which come ſmall heads with long pointed beakes, reſembling the long bill of a Storke or Crane, or ſuch like bird, which after it is ripe, parteth at the bottome where it is biggeſt, into foure or fiue ſeedes, euery one whereof hath a peece of the beake head faſtened vnto it, and falleth away if it bee not gathered: the roote is tuberous and round, like vnto the roote of the *Cyclamen* or ordinary Sowbread almoſt, but ſmaller, and of a darke ruſſet colour on the outſide, and white within, which doth encreaſe vnder ground, by certaine ſtrings running from the mother root into ſmall round bulbes, like vnto the rootes of the earth Cheſnut, and will preſently ſhoote leaues, and quickly grow to beare flowers, but will not abide to be kept long dry out of the ground, without danger to be vtterly ſpoiled.

*Geranium Batrachoides flore cæruleo.* The blew Crowfoote Cranes bill.

This Crowfoote Cranes bill hath many large leaues, cut into fiue or ſix parts or diuiſions, euen to the bottome, and iagged beſides on the edges, ſet vpon very long ſlender foote-ſtalkes, very like the leaues of the wilde Crowfoot; from among which riſe vp diuers ſtalkes with great ioynts, ſomewhat reddiſh, ſet with leaues like the former: the toppes of the ſtalkes are ſpread into many branches, whereon ſtand diuers flowers, made of fiue leaues a peece, as large as any of the wilde or field Crowfeete, round pointed, of a faire blew or watchet colour, which being paſt, there doe ariſe ſuch heads or bils, as other of the Cranes bils haue: the roote is compoſed of many reddiſh ſtrings, ſpreading in the ground, from a head made of diuers red heads, which lye oftentimes eminent aboue the ground.

*Geranium Batrachoides flore albo.* The white Crowfoote Cranes bill.

This Cranes bill is in leafe and flower altogether like the former, the onely difference betweene them conſiſteth in the colour of the flower, which in this is wholly white, and as large as the former: but the roote of this hath not ſuch red heads as the other hath.

*Geranium Batrachoides flore albo & cæruleo vario.*
The party coloured Crowfoote Cranes bill.

The flowers of this Cranes bill are variably ſtriped and ſpotted, and ſometimes diuided, the one halfe of euery leafe being white, and the other halfe blew, ſometimes with leſſer or greater ſpots of blew in the white leafe, very variably, and more in ſome years then in others, that it is very hard to expreſſe all the varieties that may be obſerued in the flowers, that blow at one time. In all other parts of the plant, it is ſo like vnto the former, that vntill it be in flower, the one cannot be knowne from the other.

*Geranium*

5. *Geranium Batrachoides alterum flore purpureo.*
Purple Crowfoote Cranes bill.

This purple Cranes bill hath many leaues rising from the roote, set vpon long foote stalkes, somewhat like vnto the other, yet not so broad, but more diuided or cut, that is, into seuen or more slits, euen to the middle, each whereof is likewise cut in on the edges more deeply then the former; the stalkes are somewhat knobbed at the ioynts, set with leaues like vnto the lower, and bearing a great tuft of buds at the toppes of the branches, which breake out into faire large flowers, made of fiue purple leaues, which doe somewhat resemble the flower of a Mallow, before it be too full blowne, each whereof hath a reddish pointell in the middle, and many small threads compassing it, this vmbell or tuft of buds doe flower by degrees, and not all at once, and euery flower abideth open little more then one day, and then shedderh, so that euery day yeeldeth fresh flowers, which because they are so many, are a long while before they are all past or spent: after the flowers are past, there arise small beake heads or bils, like vnto the other Cranes bils, with small turning seede: the roote is composed of a great tuft of strings, fastened to a knobby head.

6. *Geranium Romanum versicolor siue striatum.* The variable stript Cranes bill.

This beautifull Cranes bill hath many broad yellowish greene leaues arising from the roote, diuided into fiue or six parts, but not vnto the middle as the first kindes are: each of these leaues hath a blackish spot at the bottome corners of the diuisions, the whole leafe as well in forme as colour and spots, is very like vnto the leafe of the *Geranium fuscum*, or spotted Cranes bill, next following to be described, but that the leaues of this are not so large as the other: from among these leaues spring vp sundry stalkes a foote high and better, ioynted and knobbed here and there, bearing at the tops two or three small white flowers, consisting of fiue leaues a peece, so thickly & variably striped with fine small reddish veines, that no green leafe that is of that bigneſſe can shew so many veines in it, nor so thick running as euery leafe of this flower doth: in the middle of the flower standeth a small pointell, which when the flower is past doth grow to be the seed vessell, wherein is set diuers small seeds, like vnto the small seedes of other Cranes bils: the root is made of many small yellow threads or strings.

7. *Geranium fuscum siue maculatum.* Swart tawny or spotted Cranes bill.

The leaues of this Cranes bill are in all points like the last described, as well in the forme and diuisions as colour of the leaues, being of a yellowish greene colour, but larger and stronger by much: the stalkes of this rise much higher, and are ioynted or knobbed with reddish knees or ioynts, on the tops whereof stand not many although large flowers, consisting of fiue leaues a peece, each whereof is round at the end, and a little snipt round about, and doe bend or turne themselues backe to the stalkewards, making the middle to be higheſt or most eminent; the colour of the flower is of a darke or deepe blackish purple, the bottome of euery leafe being whiter then the rest; it hath also a middle pointell standing out, which afterwards bring forth seede like vnto others of his kinde: the roote consisteth of diuers great strings, ioyned to a knobby head.

8. *Geranium Hematodes.* The red Rose Cranes bill.

This Cranes bill hath diuers leaues spread vpon the ground, very much cut in or diuided into many parts, and each of them againe slit or cut into two or three peeces, standing vpon slender long foote-stalkes, of a faire greene colour all the Spring and Summer, but reddish in Autumne: among these leaues spring vp slender and weake stalkes, beset at euery ioynt (which is somewhat reddish) with two leaues for the most part, like vnto the lower: the flowers grow seuerally on the toppe of the stalkes, and not many together in bunches or branches, as in all other of the Cranes bils, euery flower being as large as a single Rose Campion flower, consisting of fiue large leaues,

of a deeper red colour then in any other Cranes bill at the first opening, and will change more blewish afterwards: when the flower is past, there doth arise such like beakes as are in others of the same kinde, but small: the roote is hard, long, and thicke, with diuers branches spreading from it, of a reddish yellow colour on the outside, and whitish within, which abideth and perisheth not, but shooteth forth some new greene leaues, which abide all the Winter, although those that turne red doe fall away.

### *Geranium Creticum.* Candy Cranes bill.

Candy Cranes bill beareth long and tender stalkes, whereon growe diuers broad and long leaues, cut in or ragged on the edges: the toppes of the stalkes are branched into many flowers, made of fiue leaues of a reasonable bignesse, and of a faire blew or watchet colour, with a purplish pointell in the middle, which being past, there follow beake heads like other Cranes bils, but greater, containing larger, greater, and sharper pointed seede, able to pierce the skinne, if one be not warie of it: the roote is white and long, with some fibres at it, and perisheth when it hath perfected his seede, and will spring of it owne sowing many times, if the Winter be not too sharpe, otherwise (being annuall) it must be sowne in the Spring of the yeare.

#### The Place.

Most of these Cranes bils are strangers vnto vs by nature, but endenizond in our English Gardens. It hath beene reported vnto mee by some of good credit, that the second or Crowfoot Cranes bill hath been found naturally growing in England, but yet I neuer saw it, although I haue seen many sorts of wilde kindes in many places. Matthiolus saith that the first groweth in Dalmatia and Illyria very plentifully. Camerarius, Clusius, and others, that most of the rest grow in Germany, Bohemia, Austria, &c. The last hath his place recorded in his title.

#### The Time.

All these Cranes bils doe for the most part flower in Aprill, and May, and vntill the middle of Iune. The variable or stript Cranes bill is vsually the latest of all the rest.

#### The Names.

The first is vsually called *Geranium tuberosum*, of some *Geranium bulbosum*, of the likenesse of the roote vnto a bulbe: It is without controuersie *Geranium primum* of Dioscorides. The second is called *Geranium Gratia Dei*, of others, *Geranium cæruleum*. The blew Cranes bill Lobel calleth it *Batrachoides*, because both leafe and flower are like vnto a Crowfoote; and the affinity with the Cranes bils in the seede causeth it rather to be referred to them then to the Crowfeete. The stript Cranes bill is called by some *Geranium Romanum*. The last saiuing one is called *Geranium Hæmatodes*, or *Sanguineum*, of Lobel *Geranium Gruinale Hæmatodes supinum radice repente*. In English it may be called after the Greek and Latine, The bloudy Cranes bill, but I rather call it, The Rose Cranes bill, because the flowers are as large as single Roses, or as the Rose Campion. Some of them are called in many places of England Bassinets.

#### The Vertues.

All the kindes of Cranes bils are accounted great wound herbes, and effectuall to stay bleedings, yet some more then others. The Emperickes of Germanie, as Camerarius saith, extoll it wonderfully, for a singular remedie against the Stone, both in the reines and bladder.

## CHAP. XXXI.

### *Sanicula guttata maior.* Spotted Sanicle.

HAuing long debated with my selfe, where to place this & the other plants that follow in the two next Chapters, I haue thought it not amisse for this worke to set them downe here, both before the Beares eares, which are kindes of Sanicle, as the best Authors doe hold, and after the Cranes bils, both for some qualities somewhat resembling them, and for some affinity of the flowers with the former.

The spotted Sanicle hath many small round leaues, bluntly endented about the edges, somewhat like vnto the leaues of our white Saxifrage, of a full greene colour aboue, and whitish hairy, and somewhat reddish withall vnderneath: the stalkes are set here and there with the like leaues, rising a foote and a halfe high or more, very much diuided at the toppe into sundry small branches, bearing many very small white flowers, consisting of fiue small leaues, wherein are many small red spots to be seene, as small as pins points, of a pretty sweete sent, almost like Hawthorne flowers, in the middle whereof are many small threads compassing a head, which when it is ripe containeth small blacke seede: the roote is scaly, or couered with a chaffie matter, hauing many small white fibres vnderneath, whereby it is fastened in the ground.

There is another of this kinde, like both in roote, leafe, and flower to the former, the onely difference is, that this is lesser then the former, and hath no spots in the flower, as the other hath. *Minor non guttata.*

We haue also another smaller kinde then the last, both in leafe and flower, the leaues whereof are smaller, but rounder, and more finely snipt or indented about the edges, like the teeth of a fine sawe: the stalke is little aboue a span high, hauing many small white flowers spotted as the first, but with fewer spots. *Minus guttata.*

#### The Place.

These growe in the shadowie Woods of the Alpes, in diuers places, and with vs they more delight in the shade then the sunne.

#### The Time.

All these Sanicles doe flower in May, and continue flowring vntill Iune, and the seede soone ripeneth after: the rootes abide all the Winter, with some leaues on them, springing a fresh in the beginning of the yeare.

#### The Names.

The former two are called by Clusius *Sanicula montana*, and by others *Sanicula guttata*: by Lobel *Geum Alpinum*. The third or last hath been sent vs vnder the name of *Sanicula montana altera minor*.

#### The Vertues.

The name imposed on these plants doe certainly assure vs of their vertues, from the first founders, that they are great healers, and from their taste, that they are great binders.

## Chap. XXXII.

### *Cotyledon altera Matthioli.* Spotted Nauelwort.

THis spotted Nauelwort, as many doe call it, hath many thicke small leaues, not so broad as long, of a whitish greene colour, lying on the ground in circles, after the manner of the heads of Houseleeke, and dented about the edges; from the middle whereof sometimes (for it doth not flower euery yeare in many places) ariseth vp a stalke, scarce a foote high, beset with such like leaues as are belowe, but somewhat longer: from the middle of the stalke vp to the top it brancheth forth diuersly, with a leafe at euery ioynt, bearing three or foure flowers on euery branch, consisting of fiue white leaues, spotted with small red spots, like vnto the spotted Sanicle, but with fewer and greater spots, hauing a yellowish circle or eye in the bottome of euery flower, and many whitish threads with yellowish tips in it: the seede is small and blacke, contained in small round heads: the roote is small, long, and threadie, shooting out such heads of leaues, which abide all the Winter, those that beare flower perishing.

### *Cotyledon altera minor.* Small dented Nauelwort.

There is another like vnto that before described in most things, the differences be these: It hath shorter leaues then the former, and dented about the edges in the like manner: the flowers hereof are white, but greater, made of six leaues, and most vsually without any spots at all in them, some are seene to haue spots also: the heads or seede vessels are more cornered then the former.

### *Cotyledon altera flore rubro stellato.* Small red flowred Nauelwort.

This hath also many heads of leaues, but more open, which are longer, greener, and sharper pointed then eyther of the former, somewhat reddish also, and not dented about the edges, but yet a little rough in handling: the stalke ariseth from among the leaues, being somewhat reddish, and the leaues thereon are reddish pointed, diuided at the toppe into many branches, with diuers flowers thereon, made of twelue small long leaues, standing like a starre, of a reddish purple colour, with many threads therein, set about the middle head, which is diuided at the toppe into many small ends, like pods or hornes, containing therein very small seede: the root is small like the former.

### *Sedum serratum flore rubente maculato.* The Princes Feather.

This kinde of Sengreene is composed of heads of larger, broader, and thinner leaues then any of the former, of a sadder greene colour, somewhat vneuenly endented about the edges, and not so close set together, but spreading forth into seuerall heads like as the former sorts doe, although not so plentifully; from the middle of diuers of which heads rise vp browniſh or reddish stalkes, set with smaller leaues thereon to the middle thereof, and then brancheth forth into seuerall sprigs, set with diuers small reddish flowers consisting of fiue leaues a peece, the innerside of which are of a pale red, somewhat whitish, spotted with many small bloud red spots, as small almost as pins points, with some small threads in the middle, standing about a small greene head, which turneth into the seede vessell, parted foure wayes at the head, wherein is contained small blackish seede: the rootes are small threads, which spread vnder the ground, and shoote vp seuerall heads round about it.

### The Place.

All these growe in Germany, Hungarie, Austria, the Alpes, and other such like places, where they cleaue to the rocke it selfe, that hath but a crust of earth on it to nourish them. They will abide in Gardens reasonable well, if they be planted in shadowie places, and not in the sun.

The

### The Time.

They flower for the most part in the end of May, and sometimes sooner or later, as the yeare falleth out.

### The Names.

The first is called by Matthiolus, *Cotyledon altera Dioscoridis*, and *Vmbilicus alter*, but it is not the true *Cotyledon altera* of Dioscorides; for *Sedum vulgare maius*, Our common Houseleeke, by the consent of the best moderne Writers, is the true *Cotyledon altera* of Dioscorides, or *Vmbilicus Veneris alter*. I hold it rather to bee a kinde of small Houseleeke, as the other two likewise are. The second is called by some *Aizoum* or *Sedum minus serratum*. The third hath his name in his title. Wee doe call them Nauelworts in English rather then Houseleekes, *Euphoniæ gratia*. The last may be called dented Sengreene with reddish spotted flowers, but some of our English Gentlewomen haue called it, The Princes Feather, which although it be but a by-name, may well serue for this plant to distinguish it, and whereby to be knowne.

### The Vertues.

They are all held to be cold and moist, like vnto other Houseleekes.

---

## Chap. XXXIII.

*Soldanella Alpina.* Mountaine Soldanella or blew Moonewort.

THis beautifull plant hath many round and hard leaues, set vpon long foote-stalkes, a little vneuenly cut about the edges, greene on the vpperside, and of a grayish greene vnderneath, and sometime reddish like the leaues of Sowbread, which because they doe somewhat resemble the leaues of *Soldanella marina*, which is the Sea Bindweede, tooke the name thereof: the stalkes are slender, small, round, and reddish, about a span high, bearing foure or fiue flowers at the toppe, euery one hanging downe their heads, like vnto a Bell flower, consisting but of one leafe (as most of the Bindweeds doe) plated into fiue folds, each of them ending in a long point, which maketh the flower seem to haue fiue leaues, each whereof is deeply cut in on the edges, and hauing a round greene head in the middle, with a pricke or pointell at the end thereof: the flower is of a faire blew colour, sometimes deeper or paler, or white, as nature listeth without any smell at all: the middle head, after the flower is fallen, riseth to be a long round pod, bearing that pricke it had at the end thereof, wherein is contained small greenish seede: the roote hath many fibres shooting from a long round head or roote.

### The Place.

This groweth on the Alpes, which are couered with snow the greatest part of the yeare, and will hardly abide transplanting.

### The Time.

In the naturall places it flowreth not vntill the Summer moneths, Iune, Iuly, and August, after the snow is melted from the Hils, but being brought into Gardens, it flowreth in the beginning of Aprill, or thereabouts.

### The Names.

This plant, by reason of the likenesse of leaues with *Soldanella*, as was before said, is called by many *Soldanella*, but yet is no Bindweede; and therefore I rather call it in English a Mountaine Soldanella, then as Gerrard doth, Mountaine Bindweede. It is likewise called by some, *Lunaria minor cærulea*, The lesser blew Lunary or Moonwort, and so I would rather haue it called.

### The Vertues.

They that imposed the name of *Lunaria* vpon this plant, seeme to referre it to the wound or consolidating herbes, but because I haue no further relation or experience, I can say no more thereof vntill tryall hath taught it. Some also from the name *Soldanella*, which is giuen it, because of the likenesse of the leaues, haue vsed it to help the Dropsie, for which the Sea plant is thought to be effectuall.

## Chap. XXXIIII.

### *Auricula Vrsi.* Beares eares.

THere are so many sundry and seuerall sorts of Beares eares, the variety consisting as well in the differing colours of the flowers, as the forme and colour of the leaues, that I shall not comprehend and set downe vnto you all the diuersities by many, that are risen vpto those that haue beene industrious in the sowing of the seedes of the seuerall sorts of them; yet if you accept of these that I doe here offer vnto you, I shall giue you the knowledge of others, as time, occasion, and the view of them shall enable me. And because they are without all question kinds of Cowslips, I haue set them downe before them in the first place, as being of more beautie and greater respect, or at the least of more rarity vnto vs. To dispose them therefore into order, I shall ranke them vnder three principall colours, that is to say, Red or Purple, White, and Yellow, and shew you the varieties of each of them (for so many as are come to my knowledge) apart by themselues, and not promiscuously as many others haue done.

#### 1. *Auricula Vrsi flore purpureo.* Purple Beares eare, or The Murrey Cowslip.

This purple Beares eare or Cowslip hath many greene leaues, somewhat long and smooth, narrow from the bottome of the leafe to the middle, and broad from thence to the end, being round pointed, and somewhat snipt or endented about the edges; in the middle of these leaues, and sometimes at the sides also, doe spring round greene stalkes foure or fiue fingers high, bearing at the top many flowers, the buds whereof, before they are blowne, are of a very deepe purple colour, and being open, are of a bright, but deepe purple, vsually called a Murrey colour, consisting of fiue leaues a peece, cut in at the end as it were into two, with a whitish ring or circle at the bottome of each flower, standing in small greene cups, wherein after the flowers are fallen, are contained very small heads, not rising to the height of the cups, bearing a small pricke or pointell at the toppe of them, wherein is little blackish seede: the roote hath many whitish strings fastened to the maine long roote, which is very like vnto a Primrose or Cowslip roote, as it is in all other parts besides.

#### 2. *Auricula Vrsi purpurea absq; orbe.* The murrey Cowslip without eyes.

There is another of this kinde, whose leafe is somewhat lesse, as the flower is also, but

but of the same colour, and sometimes somewhat redder, tending to a Scarlet, without any circle at the bottome of the flower, in no other things differing from it.

3. *Auricula Vrsi minor flore tannetto.* Tawney Beares eares.

The leaues of this kinde haue a greater shew of mealinesse to be seene in them, and not much smaller then the former, yet snipt or endented about the ends like vnto them: the flowers are many, of the same fashion with the former, but smaller, each whereof is of as deepe a murrey or tawnie colour when it is blowne, as the buds of the former are before they are blowne, hauing a white circle at the bottome of the flower, and yellowish in the middle belowe the circle.

4. *Auricula Vrsi flore rubro saturo orbe luteo.*
Deepe or bloud red Beares eares with eyes.

This kinde hath small and long greene leaues, nothing mealy, but snipt about the edges, from the middle of the leaues forwards to the ends: the flowers hereof are of a deepe red colour, tending to a bloud red, with a deepe yellow circle, or rather bottome in the middle.

*Auricula Vrsi flore rubro saturo absque orbe.*

There is another of this kinde, whose leaues are somewhat mealy, and smaller then any (that I haue seene) that haue mealy leaues: the flowers are of the same deepe red colour with the last described, yet hath no circle or bottome of any other colour at all.

5. *Auricula Vrsi flore purpureo caruleo.* The Violet coloured Beares eare.

We haue another, whose leaues are somewhat mealy and large; the flowers whereof are of a paler purple then the first, somewhat tending to a blew.

6. *Auricula Vrsi flore obsoleto magno.* The Spaniards blush Beares eare.

This great Beares eare hath as large leaues as any other of this kindred whatsoeuer, and whitish or mealy withall, somewhat snipt about the edges, as many other of them are: the flowers stand at the toppe of a strong and tall stalke, larger then any of the other that I haue seene, being of a duskie blush colour, resembling the blush of a Spaniard, whose tawney skinne cannot declare so pure a blush as the English can; and therefore I haue called it the Spaniards blush.

7. *Auricula Vrsi flore rubello.* Scarlet or light red Beares eares.

The leaues of this kinde are very like the leaues of the first purple kinde, but that they are not so thicke; of a little paler greene colour, and little or nothing snipt about the edges: the flowers are of a bright, but pale reddish colour, not halfe so deepe as the two last with white circles in the bottomes of them, in other things this differeth not from others.

8. *Auricula Vrsi Roseo colore.* The Rose coloured Beares eare.

We haue another, whose leafe is a little mealy, almost as large as any of the former, whose flowers are of a light red colour, very neare the colour of an ordinary Damaske Rose, with a white eye at the bottome.

9. *Auricula Vrsi flore caruleo folio Boraginis.*
Blew Beares eares with Borage leaues.

This plant is referred to the kindred or family of the Beares eares, onely for the forme of the flower sake, which euen therein it doth not assimilate to the halfe; but because it hath passed others with that title, I am content to insert it here, to giue you the

1 *Auricula Vrſi flore purpureo.* Purple Cowſlips or Beares eares. 2 *Auricula Vrſi flore tannetto.* Tawney Beares eare. 3 *Auricula Vrſi flore & folio Boraginis.* Blew Beares eares with Borage leaues. 4 *Auricula Vrſi flore carneo* Bluſh Beares eare. 5 *Auricula Vrſi maxima lutea flore elegauti.* The greateſt faire yellow Beares eares with eyes. 6 *Auricula Vrſi altera flore luteo.* The yellow Beares eare. 7 *Auricula Vrſi crinis coloris ſiue flore fuſco.* The haire coloured Beares eare. 8 *Corniſa Matthioli.* Beares eare Sanicle.

the knowledge thereof, and rather to ſatisfie others then my ſelfe with the place thereof: the deſcription whereof is as followeth: It hath diuers broad rough hairy leaues ſpread vpon the ground, ſomewhat like vnto the leaues of Borage for the roughneſſe, but not for the largeneſſe; the leaues hereof being ſomewhat rent in ſome places at the edges: from among theſe leaues riſe vp one, or two, or more browniſh, round, and hairy ſtalkes, a ſpan high or thereabouts, bearing at the toppes three or foure flowers a peece, conſiſting of fiue large pointed leaues, of a faire blew or light azur colour, with ſome ſmall yellow threads in the middle, ſtanding in ſmall greene cups: the roote is long and browniſh, hauing many ſmall fibres annexed vnto it.

10. *Auricula Vrſi maior flore albo.* The great white Beares eare.

This white Beares eare hath many faire whitiſh greene leaues, ſomewhat paler then the leaues of any of the kindes of Beares eares, and a little ſnipt about the ends, as manie other are: among theſe leaues riſe vp ſtalkes foure or fiue inches high, bearing at the toppe many flowers like vnto the ſmall yellow Beares eare hereafter ſet downe, of a pale whitiſh colour, tending to yellow at the firſt opening of the flower, which after two or three dayes change into a faire white colour, and ſo continue all the while it flowreth: the roote is like the purple kinde, as all or moſt of the reſt are, or very little differing.

11. *Auricula Vrſi minor flore albo.* The leſſer white Beares eare.

The leſſer Beares eare hath ſmaller leaues, of a little darker green colour: the ſtalke and flowers are likewiſe leſſer then the former, and haue no ſhew of yellowneſſe at all, eyther in budde or flower, but is pure white, differing not in other things from the reſt.

12. *Auricula Vrſi maxima lutea flore eleganti.*
The greateſt faire yellow Beares eare with eyes.

This yellow Beares eare hath many faire large thicke leaues, ſomewhat mealy or hoary vpon the greenneſſe, being larger then any other kinde, except the ſixth, and the next yellow that followeth, ſmooth about the edges, and without any endenting at all: the ſtalke is great, round, and not higher then in other of the former, but bearing manie more flowers thereon then in any other kinde, to the number of thirty many times, ſtanding ſo round and cloſe together, that they ſeeme to be a Noſegay alone, of the ſame faſhion with the former, but that the leaues are ſhorter and rounder, yet with a notch in the middle like the reſt, of a faire yellow colour, neither very pale nor deepe, with a white eye or circle in the bottome, about the middle of euery flower, which giueth it the greater grace: the ſeede is of a blackiſh browne colour, like vnto others, but contained in greater round heads then any other, with a ſmall pointell ſticking in the middle: the roote is greater and thicker then any other, with long ſtrings or fibres like vnto the other ſorts, but greater.

13. *Auricula Vrſi maior lutea folio in cano.* The greater yellow Beares eare.

This greater yellow Beares eare hath his leaues larger, and more mealy or hoarie then the laſt, or any other of theſe kindes: the flowers are not ſo many, but longer, and not ſo thicke thruſting together as the firſt, but of a deeper yellow colour, without any eye or circle in the middle.

14. *Auricula Vrſi maior flore pallido.* The great Straw coloured Beares eare.

This hath almoſt as mealy leaues as the laſt, but nothing ſo large; the flowers are of a faire ſtrawe colour, with a white circle at the bottome of them, theſe three laſt haue no ſhew or ſhadow of any other colour in any part of the edge, as ſome others that follow haue.

15. *Auricula*

15. *Auricula Vrſi minor flore pallente.* The leſſer ſtraw coloured Beares eare.

We haue another, whoſe leafe is leſſe mealy, or rather pale green, and a little mealy withall; the flowers whereof are of a paler yellow colour then the laſt, and beareth almoſt as many vpon a ſtalke as the firſt great yellow.

16. *Auricula Vrſi minor lutea.* The leſſer yellow Beares eares.

The leaues of this Beares eare are nothing ſo large as either of the three former yellow kindes, but rather of the bigneſſe of the firſt white kinde, but yet a little larger, thicker, and longer then it, hauing vnder the greenneſſe a ſmall ſhew of mealineſſe, and ſomewhat ſnipt about the edges: the flowers are of a pale yellow colour, with a little white bottome in them: the ſeed and rootes are like vnto the other kindes.

17. *Auricula Vrſi flore flauo.* The deepe yellow or Cowſlip Beares eare.

This kinde hath ſomewhat larger leaues then the laſt, of a yellowiſh greene colour, without any mealineſſe on them, or endenting about the edges, but ſmooth and whole: the flowers are not larger but longer, and not laide open ſo fully as the former, but of as deepe a yellow colour as any Cowſlip almoſt, without any circle in the bottome: neither of theſe two laſt haue any ſhew of other colour then yellow in them, ſauing the white in the eye.

18. *Auricula Vrſi verſicolor prima ſiue flore rubeſcente.* The bluſh Beares eare.

The bluſh Beares eare hath his leaues as large, and as hoary or mealy as the third greater yellow, or ſtraw coloured Beares eare; among which riſeth vp a ſtalke about foure inches high, bearing from ſix to twelue, or more faire flowers, ſomewhat larger then the ſmaller yellow Beares eare before deſcribed, hauing the ground of the flower of a darke or dunne yellow colour, ſhadowed ouer a little with a ſhew of light purple, which therefore we call a bluſh colour, the edges of the flower being tipt with a little deeper ſhew of that purple colour, the bottome of the flower abiding wholly yellow, without any circle, and is of very great beauty, which hath cauſed me to place it in the forefront of the variable coloured Beares eares. And although ſome might thinke it ſhould be placed among the firſt ranke of Beares eares, becauſe it is of a bluſh colour, yet ſeeing it is aſſuredly gained from ſome of the yellow kindes by ſowing the ſeede, as many other ſorts are, as may be ſeene plainly in the ground of the flower, which is yellow, and but ſhadowed ouer with purple, yet more then any of the reſt that follow; I thinke I haue giuen it his right place: let others of skill & experience be iudges herein.

19. *Auricula Vrſi crinis coloris.* Haire coloured Beares eares.

The leaues of this kinde are more mealy like then the laſt bluſh kinde, but ſomewhat longer and larger, and ſnipt about the edges in the ſame manner, from the middle of the leafe forwards: the flower is vſually of a fine light browne yellow colour, which wee doe vſually call an Haire colour, and ſometimes browner, the edges of the flower haue a ſhew or ſhadow of a light purple or bluſh about them, but more on the outſide then on the inſide.

20. *Auricula Vrſi verſicolor lutea.* The yellow variable Beares eare.

This variable Beares eare hath his greene leaues ſomewhat like vnto the deepe yellow, or Cowſlip Beares eare before deſcribed, but ſomewhat of a freſher greene, more ſhining and ſmaller, and ſnipt about the edges towards the ends, as many of thoſe before are: the flowers are of a faire yellow colour, much laid open when it is full blowne, that it ſeemeth almoſt flat, dafht about the edges onely with purple, being more yellow in the bottome of the flower, then in any other part.

21. *Auricula Vrſi verſicolor luteſcente viridi flore.* The variable green Beares eare.

This kinde of Beares eare hath greene leaues, very like vnto the laſt deſcribed, and ſnipt in the like manner about the edges, but in this it differeth, that his leaues do turne or fold themſelues a little backwards: the flowers are of a yellowiſh greene colour, more cloſed then the former, hauing purpliſh edges, eſpecially after they haue ſtood blowne ſome time, and haue little or none at the firſt opening: theſe haue no circles at all in them.

Many other varieties are to be found, with thoſe that are curious conſeruers of theſe delights of nature, either naturally growing on the mountaines in ſeuerall places, from whence they (being ſearched out by diuers) haue been taken and brought, or elſe raiſed from the ſeede of ſome of them, as it is more probable: for ſeuerall varieties haue beene obſerued (and no doubt many of theſe before ſpecified) to bee gotten by ſowing of the ſeedes, euery yeare lightly ſhewing a diuerſity, not obſerued before, either in the leafe, diuers from that from whence it was taken, or in the flowers. I haue onely ſet downe thoſe that haue come vnder mine owne view, and not any by relation, euen as I doe with all or moſt of the things contained in this worke.

## The Place.

Many of theſe goodly plants growe naturally on mountaines, eſpecially the Alpes, in diuers places; for ſome kindes that growe in ſome places, doe not in others, but farre diſtant one from the other. There hath likewiſe ſome beene found on the Pyrenæan mountaines, but that kinde with the blew flower and Borage leafe, hath beene gathered on the mountaines in Spaine, and on the Pyrenæans next vnto Spaine.

## The Time.

They all flower in Aprill and May, and the ſeede is ripe in the end of Iune, or beginning of Iuly, and ſometimes they will flower againe in the end of Summer, or in Autumne, if the yeare proue temperate, moiſt, and rainie.

## The Names.

It is very probable, that none of theſe plants were euer knowne vnto the ancient Writers, becauſe we cannot be aſſured, that they may be truely referred vnto any plant that they name, vnleſſe we beleeue Fabius Columna, that it ſhould be *Aliſma* of Dioſcorides, for thereunto hee doth referre it. Diuers of the later Writers haue giuen vnto them diuers names, euery one according to his owne conceit. For Geſner calleth it *Lunaria arthritica*, and *Paralytica Alpina*. Matthiolus accounteth it to bee of the kindred of the Sanicles, and ſaith, that in his time it was called by diuers Herbariſts, *Auricula Vrſi*, which name hath ſince bin receiued as moſt vſuall. We in Engliſh call them Beares eares, according to the Latine, or as they are called by diuers women, French Cowſlips; they may be called Mountaine Cowſlips, if you will, for to diſtinguiſh betweene them and other Cowſlips, whereof theſe are ſeuerall kindes.

*Sanicula Alpina ſiue Cortuſa Matthioli.* Beares eare Sanicle.

I cannot chuſe but inſert this delicate plant in the end of the Beares eares, for that it is of ſo neare affinity, although it differ much in the forme of the leaues, the deſcription whereof is in this manner: The leaues that ſpring vp firſt are much crumpled, and as it were folded together, which afterwards open themſelues into faire, broad, and roundiſh leaues, ſomewhat rough or hairy, not onely cut into fine diuiſions, but ſomewhat notched alſo about the edges, of a darke greene colour on the vpperſide, and

more

more whitish greene vnderneath; amongst these leaues riseth vp one or two naked round stalkes, fiue or six inches high, bearing at the toppes diuers small flowers, somewhat sweete, like vnto the first purple Beares eare, hanging downe their heads, consisting of fiue small pointed leaues a peece, of a darke reddish purple colour, with a white circle or bottome in the middle, and some small threads therein: after the flowers are past, there come small round heads, somewhat longer then any of the Beares eares, standing vpright vpon their small foot-stalkes, wherein is contained small round and blackish seede: the roote consisteth of a thicke tuft of small whitish threads, rather then rootes, much enterlaced one among another: the leaues of this plant dye downe euery yeare, and spring vp a new in the beginning of the yeare, whereas all the Beares eares doe hold their leaues greene all the Winter, especially the middlemost, which stand like a close head, the outermost for the most part perishing after seed time.

## The Place.

This groweth in many shadowie Woods both of Italy and Germany; for both Clusius hath described it, finding it in the Woods of Austria and Stiria; and Matthiolus setteth it downe, hauing receiued it from Anthonius Cortusus, who was President of the Garden at Padua, and found it in the woody mountaines of Vicenza, neare vnto Villestagna, whereon (as Matthiolus saith) there is found both with white flowers as well as with blew, but such with white flowers or blew we neuer could see or heare further of.

## The Time.

It flowreth much about the time of the Beares eares, or rather a little later, and the seede is ripe with them.

## The Names.

Clusius calleth it *Sanicula montana*, and *Sanicula Alpina*, and referreth it to the *Auricula Vrsi*, or Beares eare, which it doth most nearly resemble: but Matthiolus referreth it to the *Cariophyllata* or *Auens*, making it to be of that tribe or family, and calleth it *Cortusa* of him that first sent it him. Wee may call it eyther *Cortusa*, as for the most part all Herbarists doe, or Beares eare Sanicle as Gerrard doth.

## The Vertues.

All the sorts of Beares eares are Cephalicall, that is, conducing helpe for the paines in the head, and for the giddinesse thereof, which may happen, eyther by the sight of steepe places subiect to danger, or otherwise. They are accounted also to be helping for the Palsey, and shaking of the ioynts; and also as a Sanicle or wound-herbe. The leaues of the *Cortusa* taste a little hot, and if one of them bee laide whole, without bruising, on the cheeke of any tender skind woman, it will raise an orient red colour, as if some *fucus* had beene laide thereon, which will passe away without any manner of harme, or marke where it lay: This is Cortusus his obseruation. Camerarius in his *Hortus Medicus* saith, that an oyle is made thereof, that is admirable for to cure wounds.

## Chap. XXXV.

### *Primula veris & Paralysis.* Primroses and Cowslips.

WE haue so great variety of Primroses and Cowslips of our owne Country breeding, that strangers being much delighted with them, haue beene often furnished into diuers Countries, to their good content: And that I may set them downe in some methodicall manner, as I haue done other things, I will first set downe all the sorts of those we call Primroses, both single and double, and afterwards the Cowslips with their diuersities, in as ample manner as my knowledge can direct me. And yet I know, that the name of *Primula veris* or Primrose, is indifferently conferred vpon those that I distinguish for *Paralyses* or Cowslips. I doe therefore for your better vnderstanding of my distinction betweene Primroses and Cowslips, call those onely Primroses that cariy but one flower vpon a stalke, be they single or double, except that of Master Hesket, and that with double flowers many vpon a stalke, set out in Gerards Herball, which is his onely, not found (as I thinke) *in rerum natura*, I am sure, such a one I could neuer heare of: And those Cowslips, that beare many flowers vpon a stalke together constantly, be they single or double also. I might otherwise distinguish them also by the leafe; that all the Primroses beare their long and large broad yellowish greene leaues, without stalkes most vsually; and all the Cowslips haue small stalkes vnder the leaues, which are smaller, and of a darker greene, as vsually, but that this distinction is neither so certaine and generall, nor so well knowne.

### 1. *Primula veris flore albo.* The single white Primrose.

The Primrose that groweth vnder euery bush or hedge, in all or most of the Woods, Groues, and Orchards of this Kingdome, I may well leaue to his wilde habitation, being not so fit for a Garden, and so well knowne, that I meane not to giue you any further relation thereof: But we haue a kinde hereof which is somewhat smaller, and beareth milke white flowers, without any shew of yellownesse in them, and is more vsually brought into Gardens for the rarity, and differeth not from the wilde or ordinary kinde, either in roote or leafe, or any thing else, yet hauing those yellow spots, but smaller, and not so deepe, as are in the other wilde kinde.

### 2. *Primula veris flore viridi simplici.* The single greene Primrose.

The single greene Primrose hath his leaues very like vnto the greater double Primrose, but smaller, and of a sadder greene colour: the flowers stand seuerally vpon long foot-stalkes, as the first single kinde doth, but larger then they, and more laide open, of the same, or very neare the same yellowish greene colour that the huske is of, so that at the first opening, the huske and the flower seeme to make one double greene flower, which afterwards separating themselues, the single flower groweth aboue the huske, and spreadeth it selfe open much more then any other single Primrose doth, growing in the end to be of a paler greene colour.

### 3. *Primula veris flore viridante & albo simplici.* The single greene and white Primrose.

The leaues of this differ in a manner nothing from the former, neither doth the flower but only in this, that out of the large yellowish green huskes, which contain the flowers of the former, there commeth forth out of the middle of each of them either a small peece of a whitish flower, or else a larger, sometimes making vp a whole flower, like an ordinary Primrose.

### 4. *Primula veris flore viridi duplici.* The double greene Primrose.

This double Primrose is in his leaues so like the former single greene kindes, that the

1 *Primula veris flore albo.* The white Primrose. 2 *Primula veris flore viridi & albo simplici.* The green and white Primrose. 3 *Primula veris flore viridi duplici.* The double green Primrose. 4 *Primula veris Heskeri.* Master Heskets double Primrose. 5 *Primula veris flore pleno vulgaris.* The ordinary double Primrose. 6 *Paralysis veris flore viridante simplici.* The single green Cowslip. 7 *Paralysis flore geminato coccineo.* Double Cowslips or hose in hose. 8 *Paralysis inodora flore geminato.* Double Oxelips hose in hose. 9 *Paralysis flore & calice crispo.* Curld Cowslips or Gaskins. 10 *Paralysis minor angustifolia flore rubro.* Red Birds eyen. 11 *Paralysis hortensis flore pleno vulgaris.* Double Paigles. 12 *Paralysis fatua.* The foolish Cowslip, or laske an Apes or horse backe. 13 *Paralysis flore viridi roseo calamistrato.* The double greene feathered Cowslip.

the one cannot be knowne from the other vntill it come to flower, and then it beareth vpon euery ſtalke a double green flower, of a little deeper green colour then the flower of the former ſingle kinde conſiſting but of two rowes of ſhort leaues moſt vſually, and both of an equall height aboue the huske, abiding a pretty time in flower, eſpecially if it ſtand in any ſhadowed place, or where the Sun may come but a while vnto it.

5. *Primula veris Hesketi flore multiplici ſeparatim diuiſo.*
Maſter Heskets double Primroſe.

Maſter Heskets double Primroſe is very like vnto the ſmall double Primroſe, both in leafe, roote, and heigth of growing, the ſtalke not riſing much higher then it, but bearing flowers in a farre different manner; for this beareth not only ſingle flowers vpon ſeuerall ſtalkes, but ſomtimes two or three ſingle flowers vpon one ſtalk, and alſo at the ſame time a bigger ſtalke, and ſomewhat higher, hauing one greene huske at the toppe thereof, ſometimes broken on the one ſide, and ſometimes whole, in the middle whereof ſtandeth ſometimes diuers ſingle flowers, thruſt together, euery flower to be ſeene in his proper forme, and ſometimes there appeare with ſome whole flowers others that are but parts of flowers, as if the flowers were broken in peeces, and thruſt into one huske, the leaues of the flowers (being of a white or pale Primroſe colour, but a little deeper) ſeldome riſing aboue the height of the very huske it ſelfe; and ſometimes, as I haue obſerued in this plant, it will haue vpon the ſame ſtalke, that beareth ſuch flowers as I haue here deſcribed vnto you, a ſmall flower or two, making the ſtalke ſeeme branched into many flowers, whereby you may perceiue, that it will vary into many formes, not abiding conſtant in any yeare, as all the other ſorts doe.

6. *Primula hortenſis flore pleno vulgaris.* The ordinary double Primroſe.

The leaues of this Primroſe are very large, and like vnto the ſingle kind, but ſomewhat larger, becauſe it groweth in gardens: the flowers doe ſtand euery one ſeuerally vpon ſlender long footeſtalkes, as the ſingle kinde doth, in greeniſh huskes of a pale yellow colour, like vnto the field Primroſe, but very thicke and double, and of the ſame ſweete ſent with them.

7. *Primula veris flore duplici.* The ſmall double Primroſe.

This Primroſe is both in leafe, roote, and flower, altogether like vnto the laſt double Primroſe, but that it is ſmaller in all things; for the flower riſeth not aboue two or three fingers high, and but twice double, that is, with two rowes of leaues, yet of the very ſame Primroſe colour that the former is of.

8. *Paralyſis vulgaris pratenſis flore flauo ſimplici odorato.*
The Common field Cowſlip.

The common fielde Cowſlip I might well forbeare to ſet downe, being ſo plentifull in the fields: but becauſe many take delight in it, and plant it in their gardens, I will giue you the deſcription of it here. It hath diuers green leaues, very like vnto the wilde Primroſe, but ſhorter, rounder, ſtiffer, rougher, more crumpled about the edges, and of a ſadder greene colour, euery one ſtanding vpon his ſtalke, which is an inch or two long: among the leaues riſe vp diuers round ſtalkes, a foote or more high, bearing at the toppe many faire yellow ſingle flowers, with ſpots of a deeper yellow, at the buttome of each leafe, ſmelling very ſweete. The rootes are like to the other Primroſes, hauing many fibres annexed to the great roote.

9. *Paralyſis altera odorata flore pallido polyanthos.* The Primroſe Cowſlip.

The leaues of this Cowſlip are larger then the ordinary fielde Cowſlip, and of a darke yellowiſh greene colour: the flowers are many ſtanding together, vpon the toppes of the ſtalkes, to the number of thirty ſometimes vpon one ſtalke, as I haue counted them in mine owne Garden, and ſometimes more, euery one hauing a longer

foote

foote stalke then the former, and of as pale a yellowish colour almost as the fielde Primrose, with yellow spots at the bottome of the leaues, as the ordinary hath, and of as sweet a sent.

10. *Paralysis flore viridante simplici.* The single greene Cowslip.

There is little difference in leafe or roote of this from the first Cowslip, the chiefest varietie in this kinde is this, that the leaues are somewhat greener, and the flowers being in all respects like in forme vnto the first kinde, but somewhat larger, are of the same colour with the greene huskes, or rather a little yellower, and of a very small sent; in all other things I finde no diuersitie, but that it standeth much longer in flower before it fadeth, especially if it stand out of the Sunne.

11. *Paralysis flore & calice crispo.* Curl'd Cowslips or Gallegaskins.

There is another kinde, whose flowers are folded or crumpled at the edges, and the huskes of the flowers bigger than any of the former, more swelling out in the middle, as it were ribbes, and crumpled on the sides of the huskes, which doe somewhat resemble mens hose that they did weare, and tooke the name of Gallegaskins from thence.

12. *Paralysis flore geminato odorato.*
Double Cowslips one within another, or Hose in Hose.

The only difference of this kinde from the ordinary field Cowslip is, that it beareth one single flower out of another, which is as a greene huske, of the like sent that the first hath, or somewhat weaker.

13. *Paralysis flore flauo simplici inodoro absque calicibus.* Single Oxe lippes.

This kinde of Cowslip hath leaues much like the ordinary kinde, but somewhat smaller: the flowers are yellow like the Cowslip, but smaller, standing many vpon a stalke, but bare or naked, that is, without any huske to containe them, hauing but little or no sent at all; not differing in any thing else from the ordinary Cowslip.

14. *Paralysis flore geminato inodora.* Double Oxelips Hose in Hose.

As the former double Cowslip had his flowers one within another, in the very like manner hath this kinde of Cowslip or Oxelippe, sauing that this hath no huske to containe them, no more then the former single Oxelippe hath, standing bare or naked, of the very same bignesse each of them, and of the same deepe yellow colour with it, hauing as small a sent as the former likewise.

Wee haue another of this kinde, whose leaues are somewhat larger, and so are the *Flore pallidiore* flowers also, but of a paler yellow colour.

15. *Paralysis inodora calicibus dissectis.* Oxelips with iagged huskes.

This kinde differeth not from the first Oxelip in the smalnesse of the greene leaues, but in the flower, which standing many together on a reasonable high stalke, and being very small and yellow, scarce opening themselues or layde abroade as it, hath a greene huske vnder each flower, but diuided into sixe seuerall small long peeces.

16. *Paralysis flore fatuo.* The Frantick, or Foolish Cowslip:
Or Iacke an apes on horse backe.

Wee haue in our gardens another kinde, not much differing in leaues from the former Cowslip, and is called Fantasticke or Foolish, because it beareth at the toppe of the stalke a bush or tuft of small long greene leaues, with some yellow leaues, as it were peeces of flowers broken, and standing among the greene leaues. And sometimes

some stalkes among those greene leaues at the toppe (which are a little larger then when it hath but broken peeces of flowers) doe carry whole flowers in huskes like the single kinde.

17. *Paralysis minor flore rubro.* Red Birds eyes.

This little Cowslippe (which will hardly endure in our gardens, for all the care and industrie we can vse to keepe it) hath all the Winter long, and vntill the Spring begin to come on, his leaues so closed together, that it seemeth a small white head of leaues, which afterwards opening it selfe, spreadeth round vpon the ground, and hath small long and narrow leaues, snipt about the edges, of a pale greene colour on the vpper-side, & very white or mealy vnderneath, among these leaues rise vp one or two stalks, small & hoary, halfe a foot high, bearing at the top a bush or tuft of much smaller flowers, standing vpon short foot stalkes, somewhat like vnto Cowslips, but more like vnto the Beares eares, of a fine reddish purple colour, in some deeper, in others paler, with a yellowish circle in the bottomes of the flowers, like vnto many of the Beares eares, of a faint or small sent: the seede is smaller than in any of the former kindes, and so are the rootes likewise, being small, white and threddy.

18. *Paralysis minor flore albo.* White Birds eyes.

This kinde differeth very little or nothing from the former, sauing that it seemeth a little larger both in leafe and flower, and that the flowers hereof are wholly white, without any great appearance of any circle in the bottome of them, vnlesse it be well obserued, or at least being nothing so conspicuous, as in the former.

*Flore geminato.* These two kindes haue sometimes, but very seldome, from among the middle of the flowers on the stalke, sent out another small stalke, bearing flowers theron likewise.

19. *Paralysis hortensis flore pleno.* Double Paigles or Cowslips.

The double Paigle or Cowslip hath smaller and darker greene leaues then the single kinde hath, and longer stalkes also whereon the leaues doe stand: it beareth diuers flowers vpon a stalke, but not so many as the single kinde, euery one whereof is of a deeper and fairer yellow colour then any of the former, standing not much aboue the brimmes of the huskes that hold them, consisting of two or three rowes of leaues set round together, which maketh it shew very thicke and double, of a prettie small sent, but not heady.

20. *Paralysis flore viridante pleno.* Double greene Cowslips.

This double greene Cowslip is so like vnto the single greene kinde formerly expressed, that vntill they be neare flowring, they can hardly be distinguished: but when it is in flower, it hath large double flowers, of the same yellowish greene colour with the single, and more laid open then the former double Paigle.

21. *Paralysis flore viridante siue calamistrato.*
The greene Rose Cowslip, or double greene feathered Cowslip.

There is small difference in the leaues of this double kinde from the last, but that they are not of so darke a greene: the chiefest difference consisteth in the flowers, which are many, standing together at the toppes of the stalkes, but farre differing from all other of these kindes: for euery flower standing vpon his owne stalke, is composed of many very small and narrow leaues, without any huske to containe them, but spreading open like a little Rose, of a pale yellowish greene colour, and without any sent at all, abiding in flower, especially if it stand in a shadowie place out of the sunne, aboue two moneths, almost in as perfect beauty, as in the first weeke.

The Place.

All these kindes as they haue been found wilde, growing in diuers places

in

in England, so they haue been transplanted into Gardens, to be there nourished for the delight of their louers, where they all abide, and grow fairer then in their naturall places, except the small Birds eyes, which will (as I said) hardly abide any culture, but groweth plentifully in all the North Countries, in their squally or wet grounds.

## The Time.

These doe all flower in the Spring of the yeare, some earlier and some later, and some in the midst of Winter, as they are defended from the colds and frosts, and the mildnesse of the time will permit: yet the Cowslips doe alwayes flower later then the Primroses, and both the single and double greene Cowslips latest, as I said in their descriptions, and abide much after all the rest.

## The Names.

All these plants are called most vsually in Latine, *Primulæ veris*, *Primulæ pratenses*, and *Primulæ siluarum*, because they shew by their flowring the new Spring to bee comming on, they being as it were the first Embassadours thereof. They haue also diuers other names, as *Herba Paralysis*, *Arthritica*, *Herba Sancti Petri*, *Claues Sancti Petri*, *Verbasculum odoratum*, *Lunaria arthritica*, *Phlomis*, *Alisma siluarum*, and *Alismatis alterum genus*, as Fabius Columna calleth them. The Birds eyes are called of Lobel in Latine, *Paralytica Alpina*, *Sanicula angustifolia*, making a greater and a lesser. Others call them *Sanicula anguſtifolia*, but generally they are called *Primula veris minor*. I haue (as you see) placed them with the Cowslips, putting a difference betweene Primroses and Cowslips. And some haue distinguished them, by calling the Cowslips, *Primula veris Elatior*, that is, the Taller Primrose, and the other *Humilis*, Lowe or Dwarfe Primroses. In English they haue in like manner diuers names, according to seuerall Countries, as Primroses, Cowslips, Oxelips, Palsieworts, and Petty Mulleins. The first kindes, which are lower then the rest, are generally called by the name of Primroses (as I thinke) throughout England. The other are diuersly named; for in some Countries they call them Paigles, or Palsieworts, or Petty Mulleins, which are called Cowslips in others. Those are vsually called Oxelips, whose flowers are naked, or bare without huskes to containe them, being not so sweete as the Cowslip, yet haue they some little sent, although the Latine name doth make them to haue none. The Franticke, Fantasticke, or Foolish Cowslip, in some places is called by Country people, Iacke an Apes on horse-backe, which is an vsuall name with them, giuen to many other plants, as Daisies, Marigolds, &c. if they be strange or fantasticall, differing in the forme from the ordinary kinde of the single ones. The smallest are vsually called through all the North Country, Birds eyen, because of the small yellow circle in the bottomes of the flowers, resembling the eye of a bird.

## The Vertues.

Primroses and Cowslips are in a manner wholly vsed in Cephalicall diseases, either among other herbes or flowers, or of themselues alone, to ease paines in the head, and is accounted next vnto Betony, the best for that purpose. Experience likewise hath shewed, that they are profitable both for the Palsie, and paines of the ioynts, euen as the Beares eares are, which hath caused the names of *Arthritica*, *Paralysis*, and *Paralytica*, to bee giuen them. The iuice of the flowers is commended to cleanse the spots or marks of the face, whereof some Gentlewomen haue found good experience.

## Chap. XXXVI.

### *Pulmonaria.* Lungwort, or Cowslips of Ierusalem.

ALthough these plants are generally more vsed as Pot-herbes for the Kitchen, then as flowers for delight, yet because they are both called Cowslips, and are of like forme, but of much lesse beauty, I haue ioyned them next vnto them, in a distinct Chapter by themselues, and so may passe at this time.

1. *Pulmonaria maculosa.* Common spotted Cowslips of Ierusalem.

The Cowslip of Ierusalem hath many rough, large, and round leaues, but pointed at the ends, standing vpon long foot-stalkes, spotted with many round white spots on the vppersides of the sad greene or browne leaues, and of a grayer greene vnderneath: among the leaues spring vp diuers browne stalkes, a foote high, bearing many flowers at the toppe, very neare resembling the flowers of Cowslips, being of a purple or reddish colour while they are buds, and of a darke blewish colour when they are blowne, standing in browniſh greene huskes, and sometimes it hath beene found with white flowers: when the flowers are past, there come vp small round heads, containing blacke seed: the roote is composed of many long and thicke blacke strings.

2. *Pulmonaria altera non maculosa.* Vnspotted Cowslips of Ierusalem.

The leaues of this other kinde are not much vnlike the former, being rough as they are, but smaller, of a fairer greene colour aboue, and of a whiter greene vnderneath, without any spots at all vpon the leaues: the flowers also are like the former, and of the same colour, but a little more branched vpon the stalkes then the former: the rootes also are blacke like vnto them.

3. *Pulmonaria angustifolia.* Narrow leafed Cowslips of Ierusalem.

The leaues hereof are somewhat longer, but not so broad, and spotted with whitish spots also as the former: the stalke hereof is set with the like long hairy leaues, but smaller, being a foote high or better, bearing at the toppe many flowers, standing in huskes like the first, being somewhat reddish in the bud, and of a darke purplish blew colour when they are blowne open: the seede is like the former, all of them doe well resemble Bugloſſe and Comfrey in most parts, except the roote, which is not like them, but stringie, like vnto Cowslips, yet blacke.

### The Place.

The Cowslips of Ierusalem grow naturally in the Woods of Germany, in diuers places, and the first kinde in England also, found out by Iohn Goodier, a great searcher and louer of plants, dwelling at Maple-durham in Hampshire.

### The Time.

They flower for the most part very early, that is, in the beginning of Aprill.

### The Names.

They are generally called in Latine, *Pulmonaria*, and *maculosa*, or *non maculosa*, is added for distinctions sake. Of some it is called *Symphitum maculosum*, that is, spotted Comfrey. In English it is diuersly called; as spotted Cowslips of Ierusalem, Sage of Ierusalem, Sage of Bethlehem, Lungwort, and

and spotted Comfrey, and it might bee as fitly called spotted Bugloſſe, whereunto it is as like as vnto Comfrey, as I said before.

### The Vertues.

It is much commended of some, to bee singular good for vlcered lungs, that are full of rotten matter. As also for them that spit bloud, being boyled and drunke. It is of greatest vse for the pot, being generally held to be good, both for the lungs and the heart.

## Chap. XXXVII.

### 1. *Bugloſſum & Borrago.* Bugloſſe and Borage.

Although Borage and Bugloſſe might as fitly haue been placed, I confesse, in the Kitchen Garden, in regard they are wholly in a manner spent for Physicall properties, or for the Pot, yet because anciently they haue been entertained into Gardens of pleasure, their flowers hauing been in some respect, in that they haue alwaies been enterposed among the flowers of womens needle-worke, I am more willing to giue them place here, then thrust them into obscurity, and take such of their tribe with them also as may fit for this place, either for beauty or rarity.

The Garden Bugloſſe and Borage are so well knowne vnto all, that I shall (I doubt) but spend time in waste to describe them; yet not vsing to passe ouer any thing I name and appropriate to this Garden so sleightly, they are thus to bee knowne: Bugloſſe hath many long, narrow, hairy, or rough sad greene leaues, among which rise vp two or three very high stalks, branched at the top, whereon stand many blew flowers, consisting of fiue small round pointed leaues, with a small pointell in the middle, which are very smooth, shining, and of a reddish purple while they are buds, and not blowne open, which being fallen, there groweth in the greene huske, wherein the flower stood, three or foure roundish blacke seedes, hauing that thread or pointell standing still in the middle of them: the roote is blacke without, and whitish within, long, thicke, and full of slimie iuice (as the leaues are also) and perisheth not euery yeare, as the roote of Borage doth.

### 2. *Borrago.* Borage.

Borage hath broader, shorter, greener, and rougher leaues then Bugloſſe, the stalkes hereof are not so high, but branched into many parts, whereon stand larger flowers, and more pointed at the end then Bugloſſe, and of a paler blew colour for the most part (yet sometimes the flowers are reddish, and sometimes pure white) each of the flowers consisting of fiue leaues, standing in a round hairy whitish huske, diuided into fiue parts, and haue a small vmbone of fiue blackish threads in the middle, standing out pointed at the end, and broad at the bottome: the seed is like the other: the root is thicker and shorter then the roote of Bugloſſe, somewhat blackish without also, and whitish within, and perisheth after seede time, but riseth of it owne seede fallen, and springeth in the beginning of the yeare.

### 3. *Borrago ſemper virens.* Euerliuing Borage.

Euerliuing Borage hath many broad greene leaues, and somewhat rough, more resembling Comfrey then Borage, yet not so large as either; the stalkes are not so high as Borage, and haue many small blew flowers on them, very like to the flowers of Bugloſſe for the forme, and Borage for the colour: the rootes are blacke, thicker then either of them, somewhat more spreading, and not perishing, hauing greene leaues all the Winter long, and thereupon tooke his name.

### 4. *Anchuſa.*

### 4. *Anchusa*. Sea Bugloſſe or Alkanet.

The Sea Bugloſſe or Alkanet hath many long, rough, narrow, and darke greene leaues, ſpread vpon the ground (yet ſome that growe by the Sea ſide are rather hoarie and whitiſh) among theſe leaues riſeth vp a ſtalke, ſpread at the toppe into many branches, whereon ſtand the flowers in tufts, like vnto the Garden Bugloſſe, or rather Comfrey, but leſſer; in ſome plants of a reddiſh blew colour, and in others more red or purpliſh, and in others of a yellowiſh colour: after which come the ſeedes, very like vnto Bugloſſe, but ſomewhat longer and paler: the roote of moſt of them being tranſplanted, are ſomewhat blackiſh on the outſide, vntill the later end of Summer, and then become more red: for thoſe that grow wilde, will be then ſo red, that they will giue a very deepe red colour to thoſe that handle them, which being dryed keepe that red colour, which is vſed to many purpoſes; the roote within being white, and hauing no red colour at all.

### 5. *Limonium Rauwolfij*. Marſh Bugloſſe.

This Limonium (which I referre here to the kindes of Bugloſſe, as preſuming it is the fitteſt place where to inſert it) hath many long, narrow, and ſomewhat rough leaues lying vpon the ground, waued or cut in on both ſides, like an Indenture, ſomewhat like the leaues of Ceterach or Miltwaſt, among which riſe vp two or three ſtalkes, ſomewhat rough alſo, and with thin skinnes like wings, indented on both ſides thereof alſo, like the leaues, hauing three ſmall, long, rough, and three ſquare leaues at euery ioynt where it brancheth forth; at the toppe whereof ſtand many flowers vpon their foote ſtalkes, in ſuch a manner, as is not ſeene in any other plant, that I know: for although that ſome of the ſmall winged foot ſtalkes are ſhorter, and ſome longer, ſtanding as it were flatwiſe, or all on one ſide, and not round like an vmbell, yet are they euen at the toppe, and not one higher than another; each of which ſmall foote ſtalkes doe beare foure or fiue greeniſh heads or huskes, ioyned together, out of each of which doe ariſe other pale or bleake blew ſtiffe huskes, as if they were flowers, made as it were of parchment, which hold their colour after they are dry a long time; and out of theſe huskes likewiſe, doe come (at ſeuerall times one after another, and not all at one time or together) white flowers, conſiſting of fiue ſmall round leaues, with ſome white threds in the middle: after theſe flowers are paſt, there come in their places ſmall long ſeede, incloſed in many huskes, many of thoſe heads being idle, not yeelding any good ſeede, but chaffe, eſpecially in our Countrey, for the want of ſufficient heate of the Sunne, as I take it: the roote is ſmall, long, and blackiſh on the outſide, and periſheth at the firſt approach of Winter.

#### The Place.

Borage and Bugloſſe grow onely in Gardens with vs, and ſo doth the *Semper virens*, his originall being vnknowne vnto vs. Alkanet or Sea Bugloſſe groweth neare the Sea, in many places of France, and Spaine, and ſome of the kindes alſo in England. But the Limonium or Marſhe Bugloſſe groweth in Cales, and Malacca in Spaine, and is found alſo in Syria, as Rauwolfius relateth: and in other places alſo no doubt; for it hath beene ſent vs out of Italie, many yeares before eyther Guillaume Boel found it in Cales, or Cluſius in Malacca.

#### The Time.

Borage and Bugloſſe doe flower in Iune, and Iuly, and ſometimes ſooner, and ſo doth the euer-liuing or neuer dying Borage, but not as Gerrard ſaith, flowring Winter and Summer, whereupon it ſhould take his name, but leaueth flowring in Autumne, and abideth greene with his leaues all the Winter,

1 *Pulmonaria latifolia maculosa.* Cowslips of Ierusalem. 2 *Pulmonaria angustifolia.* Narrow leafed Cowslips of Ierusalem. 3 *Borrago.* Borage. 4 *Borrago semper virens.* Euerliuing Borage. 5 *Anchusa.* Sea Bugloſſe or Alkanet. 6 *Limonium Raunolfij.* Marſh Bugloſſe.

ter, flowring the next Spring following. The other flower not vntill Iuly, and so continue, especially the Marshe Bugloſſe vntill September bee well spent, and then giueth seede, if early frosts ouertake it not; for it seldome commeth to be ripe.

The Names.

Our ordinary Borage by the consent of all the best moderne Writers, is the true *Bugloſſum* of Dioscorides, and that our Bugloſſe was vnknowne to the ancients. The *Borago semper virens*, Lobel calleth *Bugloſſum semper virens*, that is, Euer-liuing, or greene Bugloſſe: but it more resembleth Borage then Bugloſſe; yet becauſe Bugloſſe abideth greene, to auoyde that there should not be two *Bugloſſa semper virentia*, I had rather call it Borage then Bugloſſe. Anchusa hath diuers names, as Dioscorides setteth downe. And some doe call it *Fucus herba*, from the Greeke word, becauſe the roote giuing so deepe a colour, was vsed to dye or paint the skinne. Others call it *Bugloſſum Hiſpanicum*, in Engliſh Alkanet, and of some Orchanet, after the French. Limonium was found by Leonhartus Rauwolfius, neere vnto Ioppa, which he setteth downe in the second Chapter of the third booke of his trauayles, and from him first knowne to theſe parts: I haue, as you see, referred it to the kindes of Bugloſſe, for that the flowers haue some resemblance vnto them, although I know that *Limonium genuinum* is referred to the Beetes. Let it therefore here finde a place of residence, vntill you or I can finde a fitter; and call it as you thinke beſt, eyther Limonium as Rauwolfius doth, or Marshe Bugloſſe as I doe, or if you can adde a more proper name, I shall not be offended.

The Vertues.

Borage and Bugloſſe are held to bee both temperate herbes, beeing vsed both in the pot and in drinkes that are cordiall, especially the flowers, which of Gentlewomen are candid for comfitts. The Alkanet is drying, and held to be good for wounds, and if a peece of the roote be put into a little oyle of Peter or Petroleum, it giueth as deepe a colour to the oyle, as the Hypericon doth or can to his oyle, and accounted to be singular good for a cut or greene wound.

The Limonium hath no vse that wee know, more then for a Garden; yet as Rauwolfius saith, the Syrians vse the leaues as sallats at the Table.

## Chap. XXXVIII.

### *Lychnis.* Campions.

THere bee diuers sorts of Campions, as well tame as wilde, and although some of them that I shall here entreate of, may peraduenture be found wilde in our owne Countrey, yet in regard of their beautifull flowers, they are to bee respected, and nourſed vp with the reſt, to furniſh a garden of pleasure; as for the wilde kindes, I will leaue them for another discourse.

1. *Lychnis Coronaria rubra ſimplex.* The single red Roſe Campion.

The single red Roſe Campion hath diuers thicke, hoary, or woolly long greene leaues, abiding greene all the winter, and in the end of the spring or beginning of summer, shooteth forth two or three hard round woolly stalkes, with some ioynts thereon, and at euery ioynt two such like hoary greene leaues as those below, but smaller, diuerſly branched at the toppe, hauing one flower vpon each seuerall long foot stalke, consisting

consisting of fiue leaues, somewhat broade and round pointed, of a perfect red crimson colour, standing out of a hard long round huske, ridged or crested in foure or fiue places; after the flowers are fallen there come vp round hard heads, wherein is contained small blackish seed: the roote is small, long and wooddy, with many fibres annexed vnto it, and shooteth forth anew oftentimes, yet perisheth often also.

2. *Lychnis Coronaria alba simplex.* The white Rose Campion.

The white Rose Campion is in all things like the red, but in the colour of the flower, which in this is of a pure white colour.

3. *Lichnis Coronaria albescens sine incarnata maculata & non maculata.*
The blush Rose Campion spotted and not spotted.

Like vnto the former also are these other sorts, hauing no other difference to distinguish them, but the flowers, which are of a pale or bleake whitish blush colour, especially about the brims, as if a very little red were mixed with a great deale of white, the middle of the flower being more white; the one being spotted all ouer the flower, with small spots and streakes, the other not hauing any spot at all.

4. *Lychnis Coronaria rubra multiplex.* The double red Rose Campion.

The double red Rose Campion is in all respects like vnto the single red kinde, but that this beareth double flowers, consisting of two or three rowes of leaues at the most, which are not so large as the single, and the whole plant is more tender, that is, more apt to perish, then any of the single kindes.

5. *Lychnis Chalcedonica flore simplici miniato.* Single Nonesuch, or Flower of Bristow, or Constantinople.

This Campion of Constantinople hath many broad and long greene leaues, among which rise vp sundry stiffe round hairy ioynted stalks three foot high, with two leaues euery ioynt: the flowers stand at the toppes of them, very many together, in a large tuft or vmbell, consisting of fiue small long leaues, broade pointed, and notched-in in the middle, of a bright red orenge colour, which being past, there come in their places small hard whitish heads or seede vessels, containing blacke seede, like vnto the seede of sweet Williams, and hauing but a small sent; the roote is very stringie, fastening it selfe very strongly in the ground, whereby it is much encreased.

Of the single kinde there is also two or three other sorts, differing chiefly in the colour of the flowers. The one is pure white. Another is of a blush colour wholly, without variation. And a third is very variable; for at the first it is of a pale red, and after a while groweth paler, vntill in the end it become almost fully white; and all these diuersities of the flowers are sometimes to bee seene on one stalke at one and the same time. *Flore albo. Et carneo. Verficolor.*

6. *Lychnis Chalcedonica flore miniato pleno.*
Double Flower of Bristow, or Nonesuch.

This glorious flower being as rare as it is beautifull, is for rootes, beeing stringie, for leaues and stalkes being hairy and high, and for the flowers growing in tufts, altogether like the first single kinde: but herein consisteth the chiefest difference, that this beareth a larger vmbell or tuft of flowers at the toppe of the stalke, euery flower consisting of three or foure rowes of leaues, of a deeper orenge colour then it, which addeth the more grace vnto it, but passeth away without bearing seede, as most other double flowers doe, yet recompenceth that defect with encrease from the roote.

7. *Lychnis plumaria siluestris simplex & multiplex.*
The featherd wilde Campion single and double.

The leaues of this wilde Campion are somewhat like the ordinary white wilde

Campion, but not so large, or rather resembling the leaues of sweete Williams, but that they grow not so close, nor so many together: the stalkes haue smaller leaues at the ioynts then those belowe, and branched at the toppe, with many pale, but bright red flowers, iagged or cut in on the edges, like the feathered Pinke, whereof some haue taken it to be a kinde, and some for a kinde of wilde William, but yet is but a wilde Campion, as may be obserued, both by his huske that beareth the flowers, and by the grayish roundish seede, being not of the Family of Pinkes and Gillowers, but (as I said) of the Campions: the roote is full of strings or fibres.

*Flore pleno.* The double kinde is very like vnto the single kinde, but that it is lower and smaller, and the flowers very double.

8. *Lychnis siluestris flore pleno rubro.* Red Batchelours buttons.

The double wilde Campion (which of our Countrey Gentlewomen is called Batchelours buttons) is very like both in rootes, leaues, stalkes, and flowers vnto the ordinary wilde red Campion, but somewhat lesser, his flowers are not iagged, but smooth, and very thicke and double, so that most commonly it breaketh his short huske, wherein the flower standeth on the one side, seldome hauing a whole huske, and are of a reddish colour.

9. *Lychnis siluestris flore albo pleno.* White Batchelours buttons.

As the leaues of the former double Campion was like vnto the single kinde that had red flowers, so this hath his leaues like vnto the single white kinde, differing in no other thing from it, but in the doubleness of the flowers, which by reason of the multiplicity of leaues in them thrusting forth together, breaketh his huskes wherein the flowers doe stand, as the other doth, and hath scarce one flower in many that is whole.

10. *Ocymoides arborea semper virens.* Strange Bassil Campion.

This Strange Campion (for thereunto it must bee referred) shooteth forth many round, whitish, wooddy, but brittle stalkes, whereon stand diuers long, and somewhat thicke leaues, set by couples, narrow at the bottome, and broader toward the point, of a very faire greene and shining colour, so that there is more beauty in the greene leaues, which doe so alwaies abide, then in the flowers, which are of a pale red or blush colour, consisting of fiue small long broad pointed leaues, notched in the middle, which doe not lye close, but loosly as it were hanging ouer the huskes: after the flowers are past, there come heads that containe blackish seede: the roote is small, hard, white, and threadie.

11. *Muscipula Lobelij siue Ben rubrum Monspelienſium.* Lobels Catch Flie.

I must needes insert this small plant, to finish this part of the Campions, whereunto it belongeth, being a pretty toye to furnish and decke out a Garden. It springeth vp (if it haue beene once sowne and suffered to shed) in the later end of the yeare most commonly, or else in the Spring with fiue or six small leaues, very like vnto the leaues of Pinkes, and of the same grayish colour, but a little broader and shorter, and when it beginneth to shoote vp for flower, it beareth smaller leaues on the clammy or viscous stalkes (fit to hold any small thing that lighteth on it) being broad at the bottome compassing them, and standing two at a ioynt one against another: the toppes of the stalkes are diuersly branched into seuerall parts, euery branch hauing diuers small red flowers, not notched, but smooth, standing out of small, long, round, stript huskes, which after the flowers are past, containe small grayish seede: the roote is small, and perisheth after it hath giuen seede; but riseth (as is before said) of its owne seede, if it be suffered to shed.

The Place.

The Rose Campions, Flowers of Bristow, or None such, the Bassil Campion,

1 *Lychnis Coronaria simplex.* Single Rose Campion. 2 *Lychnis Coronaria rubra multiplex.* The double red Rose Campion. 3 *Lychnis Chalcedonica simplex.* Single None such, or flower of Bristow. 4 *Lychnis Chalcedonica flore pleno* Double None such, or flower of Bristow. 5 *Lychnis plumaria multiplex.* Pleasant in sight. 6 *Lychnis sylvestris flore pleno rubro.* Red Batchelours Buttons. 7 *Lychnis sylvestris flore pleno albo.* White Batchelours Buttons. 8 *Muscipula Lobelij.* Lobels Catch Flie.

1, 2. Agrostemma coronaria, L. mod. sat.

pion, and the Catch Flie, haue been sent vs from beyond the Seas, and are onely noursed vp in Gardens with vs; the other Campions that are double, haue been naturally so found double wilde (for no art or industry of man, that euer I could be assured of to be true, be it by neuer so many repetitions of transplantations, and planeticall obseruations (as I haue said in he beginning of this worke) could bring any flower, single by nature, to become double, notwithstanding many affirmations to that purpose, but whatsoeuer hath been found wilde to be double, nature her selfe, and not art hath so produced it) and being brought into Gardens, are there encreased by slipping, and parting the roote, because they giue no seede.

## The Time.

All of them doe flower in the Summer, yet none before May.

## The Names.

The first kindes are called *Lychnides sativæ*, and *coronariæ*, in English generally Rose Campions. The next is called *Lychnis Chalcedonica*, and *Byzantina*; in English, of some Nonesuch, and of others Flower of Bristow, and after the Latine, Flower of Constantinople, because it is thought the seede was first brought from thence; but from whence the double of this kinde came, we cannot tell. The names of the others of this kinde, both single and double, are set downe with their descriptions. The feathered Campions are called *Armoraria pratensis*, and *Flos Cuculi*, and of Clusius and others thought to be *Odontitis Plinij*. Some call them in English Crowflowers, and Cuckowe-Flowers; and some call the double hereof, The faire Maide of France. The Bassil Campions were sent ouer among many other seedes out of Italy, by the name of *Ocimoides arborea semper virens*. *Arborea*, because the stalke is more wooddy and durable then other Campions: And *semper virens*, because the leaues abide greene Winter and Summer. Clusius calleth it *Lychnis semper virens*, because it is certainly a Campion. The last is diuersly called of Authors; Lobel calleth it *Muscipula*: Others *Armoraria altera*: Dodonæus *Armerius flos quartus*. Clusius *Lychnis siluestris altera*, in his Spanish obseruations, and *prima* in his History of plants, and saith, the learned of Salmantica in Spaine called it, *Ben rubrum*, as Lobel saith, they of Mompelier doe also: and by that name I receiued it firft out of Italy. It hath the name of Catch Flie, of *Muscipula* the Latine word, because the stalkes in the hot Summer dayes haue a certaine viscous or clammy humour vpon them, whereby it easily holdeth (as I said before) whatsoeuer small thing, as Flies, &c. lighteth vpon it.

## The Vertues.

We know none in these dayes, that putteth any of these to any Physicall vse, although some haue in former times.

---

# Chap. XXXIX.

*Keiri sive Leucoium luteum.* Wall-flowers, or Wall Gilloflowers.

THere are two sorts of Wall-flowers, the one single, the other double, and of each of them there is likewise some differences, as shall be shewed in their descriptions.

1. *Keiri ſiue Leucoium luteum ſimplex vulgare.* Common ſingle Wall-flowers.

The common ſingle Wall-flower which groweth wilde abroad, and yet is brought into Gardens, hath ſundry ſmall, narrow, long, and darke greene leaues, ſet without order vpon ſmall round whitiſh wooddy ſtalkes, which beare at the tops diuers ſingle yellow flowers one aboue another, euery one hauing foure leaues a peece, and of a very ſweete ſent: after which come long pods, containing reddiſh ſeede: the roote is white, hard and thready.

2. *Keiri ſiue Leucoium luteum ſimplex maius.* The great ſingle Wall-flower.

There is another ſort of ſingle Wall-flower, whoſe leaues as well as flowers are much larger then the former: the leaues being of a darker and ſhining greene colour, and the flowers of a very deepe gold yellow colour, and vſually broader then a twentie ſhilling peece of gold can couer: the ſpike or toppe of flowers alſo much longer, and abiding longer in flower, and much ſweeter likewiſe in ſent: the pods for ſeede are thicker and ſhorter, with a ſmall point at the end: this is ſlower to encreaſe into branches, as alſo to be encreaſed by the branches, and more tender to be preſerued; for the hard froſts doe cauſe it to periſh, if it be not defended from them

3. *Keiri ſimplex flore albo.* White Wall-flower.

This Wall-flower hath his leaues as greene as the great kinde, but nothing ſo large: the flowers ſtand at the toppe, but not in ſo long a ſpike, and conſiſteth of foure leaues, of a very white colour, not much larger then the common kinde, and of a faint or weaker ſent: the pods are nothing ſo great as the former great one: this is more eaſie to be propagated and encreaſed alſo, but yet will require ſome care in defending it from the colds of the Winter.

4. *Keiri ſiue Leucoium luteum vulgare flore pleno.* Common double Wall-flowers.

This ordinary double Wall-flower is in leaues and ſtalke very like vnto the firſt ſingle kinde, but that the leaues hereof are not of ſo deepe a greene colour: the flowers ſtand at the top of the ſtalkes one aboue another, as it were a long ſpike, which flower by degrees, the loweſt firſt, and ſo vpwards, by which it is a long time in flowring, and is very double, of a gold yellow colour, and very ſweete.

5. *Keiri ſiue Leucoium luteum alterum flore pleno.* Pale double Wall-flowers.

Wee haue another ſort of this kinde of double Wall-flower, whoſe double flowers ſtand not ſpike-faſhion as the former, but more open ſpread, and doe all of them blowe open at one time almoſt, and not by degrees as the other doth, and is of a paler yellow colour, not differing in any thing elſe, except that the greene leaues hereof are of a little paler greene then it.

6. *Keiri ſiue Leucoium luteum maius flore pleno ferrugineo.*
Double red Wall-flowers.

We haue alſo another ſort of double Wall-flower, whoſe leaues are as greene, and almoſt as large as the great ſingle yellow kinde, or full as bigge as the leaues of the white Wall-flower: the flowers hereof are not much larger then the ordinary, but are of a darker yellow colour then the great ſingle kinde, and of a more browniſh or red colour on the vnderſide of the leaues, and is as it were ſtriped.

7. *Keiri ſiue Leucoium maximum luteum flore pleno.*
The greateſt double yellow Wall-flower.

This great double Wall-flower is as yet a ſtranger in England, and therefore what I here

here write is more vpon relation (which yet I beleeue to be most true) then vpon sight and speculation. The leaues of this Wall flower are as greene and as large, if not larger then the great single kinde: the flowers also are of the same deepe gold yellow colour with it, but much larger then any of the former double kindes, and of as sweet a sent as any, which addeth delight vnto beauty.

### The Place.

The first single kind is often found growing vpon old wals of Churches, and other houses in many places of England, and also among rubbish and stones. The single white and great yellow, as well as all the other double kindes, are noursed vp in Gardens onely with vs.

### The Time.

All the single kindes doe flower many times in the end of Autumne, and if the Winter be milde all the Winter long, but especially in the moneths of February, March, and Aprill, and vntill the heate of the Spring doe spend them: but the other double kindes doe not continue flowring in that manner the yeare throughout, although very early sometimes, and very late also in some places.

### The Names.

They are called by diuers names, as *Viola lutea*, *Leucoium luteum*, and *Keiri*, or *Cheiri*, by which name it is chiefly knowne in our Apothecaries shops, because there is an oyle made thereof called *Cheirinum*: In English they are vsually called in these parts, Wall-flowers: Others doe call them Bee-flowers; others Wall-Gilloflowers, Winter-Gilloflowers, and yellow Srocke-Gilloflowers; but we haue a kinde of Stocke-Gilloflower that more fitly deserueth that name, as shall be shewed in the Chapter following

### The Vertues.

The sweetnesse of the flowers causeth them to be generally vsed in Nosegayes, and to decke vp houses; but physically they are vsed in diuers manners: As a Conserue made of the flowers, is vsed for a remedy both for the Appoplexie and Palsie. The distilled water helpeth well in the like manner. The oyle made of the flowers is heating and resoluing, good to ease paines of strained and pained sinewes.

## Chap. XL.

### *Leucoium.* Stocke-Gilloflower.

THere are very many sorts of Stocke-Gilloflowers both single and double, some of the fields and mountaines, others of the Sea marshes and medowes; and some noursed vp in Gardens, and there preserued by seede or slippe, as each kinde is aptest to bee ordered. But because some of these are fitter for a generall History then for this our Garden of Pleasure, both for that diuers haue no good sent, others little or no beauty, and to be entreated of onely for the variety, I shall spare so many of them as are not fit for this worke, and onely set downe the rest.

#### 1. *Leucoium simplex sativum diuersorum colorum.*
Garden Stocke-Gilloflowers single of diuers colours.

These single Stocke-Gilloflowers, although they differ in the colour of their flowers,

1 *Keiri siue Leucoium luteum vulgare.* Common Wall-flowers. 2 *Keiri siue Leucoium luteum maius simplex.* The great single Wall-flower. 3 *Keiri siue Leucoium luteum flore pleno vulgare.* Ordinary double Wall-flowers. 4 *Keiri maius flore pleno ferruginio.* The great double red Wall-flower. 5 *Leucoium sativum simplex.* Single Stocke-Gilloflowers. 6 *Leucoium sativum simplex flore striato.* Single stript Stocke-Gilloflowers.

ers, yet are in leafe and manner of growing, one so like vnto another, that vntill they come to flower, the one cannot be well knowne that beareth red flowers, from another that beareth purple; and therfore one description of the plant shall serue, with a declaration of the sundry colours of the flowers. It riseth vp with round whitish woody stalkes, two, three, or foure foot high, whereon are set many long, and not very broad, soft, and whitish or grayish greene leaues, somewhat round pointed, and parted into diuers branches, at the toppes whereof grow many flowers, one aboue another, smelling very sweet, consisting of foure small, long, and round pointed leaues, standing in small long huskes, which turne into long and flat pods, sometimes halfe a foote long, wherein is contained flat, round, reddish seedes, with grayish ringes or circles about them, lying flat all along the middle rib of the pod on both sides: the roote is long, white, and woody, spreading diuers wayes. There is great variety in the colours of the flowers: for some are wholly of a pure white colour, others of a most excellent crimson red colour, others againe of a faire red colour, but not so bright or liuely as the other, some also of a purplish or violet colour, without any spot, marke, or line in them at all. There are againe of all these colours, mixed very variably, as white mixed with small or great spottes, strakes or lines of pure or bright red, or darke red, and white with purple spots and lines; and of eyther of them whose flowers are almost halfe white, and halfe red, or halfe white, and halfe purple. The red of both sorts, and the purple also, in the like manner spotted, striped, and marked with white, differing neyther in forme, nor substance, in any other point.

2. *Leucoium satiuum albido luteum simplex.*
The single pale yellow Stocke-Gilloflower.

There is very little difference in this kind from the former, for the manner of growing, or forme of leaues or flower. Only this hath greener leaues, and pale yellow almost white flowers, in all other things alike: this is of no great regard, but only for rarity, and diuersity from the rest.

3. *Leucoium Melancholicum.* The Melancholick Gentleman.

This wilde kinde of stocke gilloflower hath larger, longer and greener leaues then any of the former kindes, vneuenly gashed or sinuated on both edges lying on the ground, and a little rough or hairy withall: from among which rise vp the stalks, a yard high or more, and hairy likewise, bearing theron here and there some such like leaues as are below, but smaller, and at the top a great number of flowers, as large or larger then any of the former single kindes, made of 4. large leaues a peece also, standing in such like long huskes, but of a darke or sullen yellowish colour: after which come long roundish pods, wherein lye somewhat long but rounder and greater seede then any stocke gilloflower, and nearer both in pod and seede vnto the *Hesperis* or Dames Violet: this perisheth not vsually after seede bearing, although sometimes it doth.

4. *Leucoium marinum Syriacum.* Leuant stocke gilloflowers.

This kind of stocke gilloflower riseth vp at the first with diuers long and somewhat broad leaues, a little vneuenly dented or waued on the edges, which so continue the first yeare after the sowing: the stalke riseth vp the next yeare to bee two foot high or more, bearing all those leaues on it that it first had, which then do grow lesse sinuated or waued then before: at the top whereof stand many flowers, made of foure leaues a peece, of a delayed purple colour, but of a small sent which turne into very long and narrow flat pods, wherein are contained flat seed like the ordinary stocke gilloflowers, but much larger and of a darke or blackish browne colour: the root is white, and groweth deepe, spreading in the ground, but growing woody when it is in seede, and perisheth afterwards.

5. *Leuco-*

*5. Leucoÿ alterum genus, flore tam multiplici quam simplici ex seminio oriundum.*

Another sort of Stocke gilloflowers bearing as well double as single flowers from seede.

This kinde of Stocke gilloflower differeth neyther in forme of leaues, stalkes, nor flowers from the former, but that it oftentimes groweth much larger and taller; so that whosoeuer shall see both these growing together, shall scarce discerne the difference, onely it beareth flowers, eyther white, red or purple, wholly or entire, that is, of one colour, without mixture of other colour in them (for so much as euer I haue obserued, or could vnderstand by others) which are eyther single, like vnto the former, or very thicke and double, like vnto the next that followeth; but larger, and growing with more store of flowers on the long stalke. But this you must vnderstand withall, that those plants that beare double flowers, doe beare no seede at all, and is very seldome encreased by slipping or cutting, as the next kinde of double is: but the onely way to haue double flowers any yeare, (for this kinde dyeth euery winter, for the most part, after it hath borne flowers, and seldome is preserued) is to saue the seedes of those plants of this kinde that beare single flowers, for from that seede will rise, some that will beare single, and some double flowers, which cannot bee distinguished one from another, I meane which will be single and which double, vntill you see them in flower, or budde at the least. And this is the only way to preserue this kinde: but of the seed of the former kinde was neuer known any double flowers to arise, and therefore you must be carefull to marke this kinde from the former.

*6. Leucoium flore pleno diuersorum colorum.*

Double Stocke Gillowflowers of diuers colours.

This other kinde of Stock gilloflower that beareth onely double flowers, groweth not so great, nor spreadeth his branches so farre, nor are his leaues so large, but is in all things smaller, and lower, and yet is woody, or shrubby, like the former, bearing his flowers in the like manner, many vpon a long stalke, one aboue another, and very double, but not so large as the former double, although it grow in fertile soyle, which are eyther white, or red, or purple wholly, without any mixture, or else mixed with spots and stripes, as the single flowers of the first kinde, but more variably, and not in all places alike, neuer bearing seede, but must be encreased, only by the cutting of the young sproutes or branches, taken in a fit season: this kinde perisheth not, as the former double kinde doth, so as it bee defended in the winter from the extreame frosts, but especially from the snow falling, or at the least remaining vpon it.

*7. Leucoium satiuum luteum flore pleno.*

The double yellow Stocke Gilloflower.

This double yellow Stock gilloflower is a stranger in England, as far as I can learne, neyther haue I any further familiaritie with him, then by relation from Germany, where it is affirmed to grow only in some of their gardens, that are curious louers of these delights, bearing long leaues somewhat hoary or white, (and not greene like vnto the Wallflower, whereunto else it might be thought to be referred) like vnto the Stock gilloflowers, as the stalkes and branches also are, and bearing faire double flowers, of a faire, but pale yellow colour. The whole plant is tender, as the double Stock gilloflowers are, and must be carefully preserued in the winter from the coldes, or rather more then the last double, lest it perish.

The Place.

The single kindes, especially some of them, grow in Italie, and some in Greece, Candy, and the Isles adiacent, as may be gathered out of the verses in Plutarches Booke *De Amore fraterno:*

*Inter*

*Inter Echinopodas velut, aſperam & inter Ononim,*
*Interdum creſcunt mollia Leucoia.*

Which ſheweth, that the ſoft or gentle ſtocke gilloflowers doe ſometimes grow among rough or prickely Furſe and Cammocke. The other ſorts are only to be found in gardens.

### The Time.

They flower in a manner all the yeare throughout in ſome places, eſpecially ſome of the ſingle kindes, if they ſtand warme, and defended from the windes and cold : the double kindes flower ſometimes in Aprill, and more plentifully in May, and Iune; but the double of ſeed, flowreth vſually late, and keepeth flowring vnto the winter, that the froſtes and colde miſtes doe pull it downe.

### The Names.

It is called *Leucoium, & Viola alba :* but the name Leucoium (which is in Engliſh the white Violet) is referred to diuers plants; we call it in Engliſh generally, Stocke gilloflower, (or as others doe, Stocke gillouer) to put a difference betweene them, and the Gilloflowers and Carnations, which are quite of another kindred, as ſhall be ſhewne in place conuenient.

### The Vertues.

Theſe haue no great vſe in Phyſick that I know: only ſome haue vſed the leaues of the ſingle white flowred kinde with ſalt, to be laid to the wreſts of them that haue agues, but with what good ſucceſſe I cannot ſay, if it happen well I thinke in one (as many ſuch things elſe will) it will fayle in a number.

## Chap. XLI.

### 1. *Heſperis, ſiue Viola Matronalis.* Dames Violets, or Queenes Gilloflowers.

THe ordinary Dames Violets, or Queene Gilloflowers, hath his leaues broader, greener, and ſharper pointed, then the Stock gilloflowers, and a little endented about the edges: the ſtalkes grow two foot high, bearing many greene leaues vpon them, ſmaller then thoſe at the bottome, and branched at the toppe, bearing many flowers, in faſhion much like the flowers of ſtocke gilloflowers, conſiſting of foure leaues in like manner, but not ſo large, of a faint purpliſh colour in ſome, and in others white, and of a pretty ſweet ſent, eſpecially towards night, but in the day time little or none at all: after the flowers are paſt, there doe come ſmall long and round pods, wherein is contained, in two rowes, ſmall and long blacke ſeede: the roote is wholly compoſed of ſtringes or fibres, which abide many yeares, and ſpringeth freſh ſtalks euery yeare, the leaues abiding all the Winter.

### 2. *Heſperis Pannonica.* Dames Violets of Hungary.

The leaues of this Violet are very like the former, but ſmoother and thicker, and not at all indented, or cut in on the edges: the flowers are like the former, but of a ſullen pale colour, turning themſelues, and ſeldome lying plaine open, hauing many purple veines, and ſtreakes running through the leaues of the flowers, of little or no ſent in the day time, but of a very ſweete ſent in the euening and morning; the ſeedes are alike alſo, but a little browner.

1 *Leucoium Melancholicum.* Sullen Stocke-Gilloflowers. 2 *Leucoium sativum flore pleno.* Double Stocke-Gilloflowers. 3 *Leucoium sativum flore pleno vario.* Party coloured Stocke-Gilloflowers. 4 *Leucoium marinum Syriacum.* Leuant Stocke-Gilloflowers. 5 *Hesperis vulgaris.* Dames Violets or Winter Gilloflowers. 6 *Lysimachia lutea siliquosa Virginiana.* The tree Primrose of Virginia. 7 *Viola Lunaris sive Bolbonach.* The white Sattin flower.

3. *Lyſimachia lutea ſiliquoſa Virgiana.* The tree Primroſe of Virginia.

Vnto what tribe or kindred I might referre this plant, I haue ſtood long in ſuſpence, in regard I make no mention of any other *Lyſimachia* in this work: leſt therfore it ſhould loſe all place, let me ranke it here next vnto the Dames Violets, although I confeſſe it hath little affinity with them. The firſt yeare of the ſowing the ſeede it abideth without any ſtalke or flowers lying vpon the ground, with diuers long and narrow pale greene leaues, ſpread oftentimes round almoſt like a Roſe, the largeſt leaues being outermoſt, and very ſmall in the middle: about May the next yeare the ſtalke riſeth, which will be in Summer of the height of a man, and of a ſtrong bigge ſize almoſt to a mans thumbe, round from the bottome to the middle, where it groweth creſted vp to the toppe, into as many parts as there are branches of flowers, euery one hauing a ſmall leafe at the foote thereof: the flowers ſtand in order one aboue another, round about the tops of the ſtalks, euery one vpon a ſhort foot-ſtalke, conſiſting of foure pale yellow leaues, ſmelling ſomewhat like vnto a Primroſe, as the colour is alſo (which hath cauſed the name) and ſtanding in a greene huske, which parteth it ſelfe at the toppe into foure parts or leaues, and turne themſelues downewards, lying cloſe to the ſtalke: the flower hath ſome chiues in the middle, which being paſt, there come in their places long and cornered pods, ſharpe pointed at the vpper end, and round below, opening at the toppe when it is ripe into fiue parts, wherein is contained ſmall browniſh ſeed: the roote is ſomewhat great at the head, and wooddy, and branched forth diuerſly, which periſheth after it hath borne ſeede.

## The Place.

The two firſt grow for the moſt part on Hils and in Woods, but with vs in Gardens onely.

The laſt, as may be well vnderſtood by the title, came out of Virginia.

## The Time.

They flower in May, Iune, and Iuly.

## The Names.

The name of *Heſperis* is impoſed by moſt Herbariſts vpon the two firſt plants, although it is not certainly knowne to be the ſame that Theophraſtus doth make mention of, in his ſixth Booke and twenty fiue Chapter *de cauſis plantarum*: but becauſe this hath the like effects to ſmell beſt in the euening, it is (as I ſaid) impoſed vpon it. It is alſo called *Viola Marina Matronalis, Hyemalis, Damaſcena* and *Muſchatella*: In Engliſh, Dames Violets, Queens Gilloflowers, and Winter Gilloflowers.

The laſt hath his Latine name in the title as is beſt agreeing with it, and for the Engliſh, although it be too fooliſh I confeſſe, yet it may paſſe for this time till a fitter be giuen, vnleſſe you pleaſe to follow the Latine, and call it Virginia Looſe-ſtrife.

## The Vertues.

I neuer knew any among vs to vſe theſe kindes of Violets in Phyſicke, although by reaſon of the ſharpe biting taſte, Dodonæus accounteth the ordinary ſort to be a kinde of Rocket, and ſaith it prouoketh ſweating, and vrine: and others affirme it to cut, digeſt, and cleanſe tough flegme. The Virginian hath not beene vſed by any that I know, either inwardly or outwardly.

## CHAP. XLII.

*Viola Lunaris siue Bolbonach.* The Sattin flower.

VNto the kindes of Stocke-Gilloflowers I thinke fitteſt to adioyne theſe kindes of Sattin-flowers, whereof there are two ſorts, one frequent enough in all our Countrie, the other is not ſo common.

1. *Viola Lunaris vulgaris.* The common white Sattin flower.

The firſt of theſe Sattin flowers, which is the moſt common, hath his leaues broad belowe, and pointed at the end, ſnipt about the edges, and of a darke greene colour: the ſtalkes are round and hard, two foot high, or higher, diuided into many branches, ſet with the like leaues, but ſmaller: the tops of the branches are beſet with many purpliſh flowers, like vnto Dames Violets, or Stocke-Gilloflowers, but larger, being of little ſent: after the flowers are paſt, there come in their places round flat thin cods, of a darke colour on the outſide, but hauing a thinne middle skinne, that is white and cleare ſhining, like vnto very pure white Sattin it ſelfe, whereon lye flat and round browniſh ſeede, ſomewhat thicke and great: the rootes periſh when they haue giuen their ſeede, and are ſomewhat round, long, and thicke, reſembling the rootes of *Lilium non bulboſum*, or Day Lilly, which are eaten (as diuers other rootes are) for Sallets, both in our owne Country, and in many places beſide.

2. *Viola Lunaris altera ſeu peregrina.* Long liuing Sattin flower.

This ſecond kinde hath broader and longer leaues then the former, the ſtalkes alſo are greener and higher, branching into flowers, of a paler purple colour, almoſt white, conſiſting of foure leaues in like manner, and ſmelling pretty ſweete, bearing ſuch like pods, but longer and ſlenderer then they: the rootes are compoſed of many long ſtrings, which dye not as the former, but abide, and ſhoot out new ſtalkes euery yeare.

### The Place.

The firſt is (as is ſaid) frequent enough in Gardens, and is found wilde in ſome places of our owne Country, as Maſter Gerard reporteth, whereof I neuer could be certainly aſſured, but I haue had it often ſent mee among other ſeedes from Italy, and other places. The other is not ſo common in Gardens, but found about Watford, as he ſaith alſo.

### The Time.

They flower in Aprill or May, and ſometimes more early.

### The Names.

It hath diuers names, as well in Engliſh as in Latine; for it is called moſt vſually *Bolbonach*, and *Viola Lunaris*: Of ſome *Viola latifolia*, and of others *Viola Peregrina*, and *Lunaria Græca*, *Lunaria maior*, and *Lunaria odorata*, and is thought to be *Thlaſpi Cretenſe*: In Engliſh, White Satten, or Satten flower: Of ſome it is called Honeſty, and Penny-flower.

### The Vertues.

Some doe vſe to eate the young rootes hereof, before they runne vp to flower, as Rampions are eaten with Vinegar and Oyle; but wee know no Phyſicall vſe they haue.

## Chap. XLIII.

### *Linum silvestre & Linaria.* Wilde Flaxe and Tode Flaxe.

ALthough neither the manured Line or Flaxe is a plant fit for our Garden, nor many of the wilde sorts, yet there are some, whose pleasant and delightfull aspect doth entertaine the beholders eyes with good content, and those I will set downe here for varietie, and adioyne vnto them some of the *Linarias*, or Tode Flaxe, for the neare affinity with them.

1. *Linum silvestre flore albo.* Wilde Flaxe with a white flower.

This kinde of wilde Flaxe riseth vp with diuers slender branches, a foote high or better, full of leaues, standing without order, being broader and longer then the manured Flaxe: the tops of the branches haue diuers faire white flowers on them, composed of fiue large leaues a peece, with many purple lines or strikes in them: the seede vessell as well as the seede, is like vnto the heads and seede of the manured Flaxe: the rootes are white strings, and abide diuers yeares, springing fresh branches and leaues euery yeare, but not vntill the Spring of the yeare.

2. *Linum silvestre luteum.* Wilde Flaxe with a yellow flower.

This wilde Flaxe doth so well resemble a kinde of St. Iohns wort, that it will soone deceiue one that doth not aduisedly regard it: For it hath many reddish stalkes, and small leaues on them, broader then the former wilde Flaxe, but not so long, which are well stored with yellow flowers, as large as the former, made of fiue leaues a peece, which being past, there come small flattish heads, containing blackish seede, but not shining like the former: the rootes hereof dye not euery yeare, as many other of the wilde kindes doe, but abide and shoote out euery yeare.

3. *Linaria purpurea.* Purple Tode Flaxe.

This purple Tode Flaxe hath diuers thicke, small, long, and somewhat narrowish leaues, snipt about the edges, of a whitish greene colour, from among which rise vp diuers stalkes, replenished at the tops with many small flowers, standing together one aboue another spike-fashion, which are small and somewhat sweete, while they are fresh, fashioned somewhat like the common Tode flaxe that groweth wilde abroad almost euery where, but much smaller, with a gaping mouth, but without any crooked spurre behinde, like vnto them, sometimes of a sad purple neare vnto a Violet, and sometimes of a paler blew colour, hauing a yellow spot in the middle or gaping place: after the flowers are past, there come small, hard, round heads, wherein are contained small, flat, and grayish seede: the roote is small, and perisheth for the most part euery yeare, and will spring againe of it owne sowing, if it be suffered to shed it selfe, yet some hard Winters haue killed the seede it should seeme, in that sometimes it faileth to spring againe, and therefore had neede to be sowne anew in the Spring.

4. *Linaria purpurea odorata.* Sweete purple Tode Flaxe.

The lower leaues of this purple Tode Flaxe are nothing like any of the rest, but are long and broad, endented about the edges, somewhat resembling the leaues of the greater wilde white Daisie: the stalke is set at the bottome with such like leaues, but a little more diuided and cut in, and still smaller and smaller vpward, so that the vppermost leaues are very like the common Tode Flaxe, the toppe whereof is branched, hauing diuers small flowers growing along vpon them, in fashion and colour almost like the last described Tode Flaxe, but not altogether so deepe a purple: the heads and seedes are very like the former, but that the seede of this is reddish: the flowers

in

1 *Linum siluestre flore albo.* Wilde Flaxe with a white flower. 2 *Linaria purpurea siue cærulea.* Purple Tode Flaxe. 3 *Linaria purpurea odorata.* Sweete purple Tode Flaxe. 4 *Scoparia siue Beluidere Italorum* Broome Tode Flaxe. 5 *Antirrhinum maius.* The greater Snapdragon. 6 *Chamænerium flore delphinij.* The willowe flower.

in their naturall hot Countries haue a fine sent, but in these colder, little or none at all: the rootes are small and threadie, and perish after they haue flowred and seeded.

5. *Linaria Valentina.* Tode Flaxe of Valentia.

This Spanish Tode Flaxe hath three or fonre thicker and bigger stalkes then the former, bearing small broad leaues, like vnto the small Centory, two or three together at a ioynt, round about the lower end of the stalkes, but without any order vpwards, at the toppes whereof stand many flowers, in fashion like vnto the common kinde, and almost as large, of a faire yellow colour, but the gaping mouth is downie, and the spurre behinde of a purplish colour.

6. *Scoparia siue Beluidere Italorum.* Broome Tode Flaxe.

Although this plant haue no beautifull flowers, yet because the greene plant full of leaues is so delightfull to behold, being in Italy and other places planted not onely in their Gardens, but set likewise in pots to furnish their Windowes, and euen with vs also hath growne to be so dainty a greene bush, that I haue thought it worthy to be among the delights of my Garden; the description whereof is as followeth: This pleasant Broome Flaxe riseth vp most vsually with one straight vpright square stalke, three foote and a halfe high or better in our Gardens, branching it selfe out diuers waies, bearing thereon many long narrow leaues, like the Garden Line or Flaxe, very thicke set together, like vnto a bush, or rather like vnto a faire greene Cypresse tree, growing broad belowe, and spire-fashion vpwards, of a very faire greene colour: at the seuerall ioynts of the branches, towards the tops, and among the leaues, there come forth small reddish flowers, not easily seene nor much regarded, being of no beauty, which turne into small round blackish gray seede: the rootes are a number of blackish strings set together, and the whole plant perisheth euery yeare at the first approach of any cold ayre, as if it neuer had beene so faire a greene bush.

The Place.

These kindes of wilde Flaxe doe growe naturally in diuers places, some in Germany, some in Spaine, and some in Italy. Those that delight in the beauty of natures variety, doe preserue them, to furnish vp the number of pleasant aspects.

The Time.

They all flower in the Summer moneths, and soone after perfect their seede.

The Names.

Their names are sufficiently expressed in their titles, yet I must giue you to vnderstand, that the last is called of some *Linaria magna*, and of others *Osyris*.

The Vertues.

The wilde Flaxe hath no medecinable vertue appropriate vnto it that is knowne. The Tode Flaxe is accounted to be good, to cause one to make water.

## Chap. XLIIII.

### *Antirrhinum.* Snapdragon.

THere is some diuersity in the Snapdragons, some being of a larger, and others of a lesser stature and bignesse; and of the larger, some of one, and some of another colour, but because the small kindes are of no beautie, I shall at this time onely entreate of the greater sorts.

#### 1. *Antirrhinum album.* White Snapdragon.

The leaues of these Snapdragons (for I doe vnder one description comprehend the rest) are broader, longer, and greener then the leaues of the Garden Flaxe, or of the wilde Flaxe set confusedly vpon the tender greene branches, which are spread on all sides, from the very bottome, bearing at the toppes many flowers, somewhat resembling the former Tode Flaxe, but much larger, and without any heele or spurre, of a faire white colour, with a yellow spot in the mouth or gaping place: after the flowers are past, there come vp in their places hard round seede vessels, fashioned somewhat like vnto a Calues head, the snout being cut off, wherein is contained small blacke seede: the rootes are many white strings, which perish in most places after they haue giuen seede, notwithstanding any care or paines taken with them to preserue them aliue, and yet they will abide in some places where they are defended in the Winter.

#### 2. *Antirrhinum purpureum siue roseum.* Purple Snapdragon.

The purple Snapdragon is in stalkes, leaues, and flowers altogether like the former, and as large and great in euery part, or greater; the only difference is, that this beareth pale Stammell or Rose coloured flowers, with a yellow spot in the mouth, and sometimes of a paler colour, almost blush.

#### 3. *Antirrhinum variegatum.* Variable Snapdragon.

This variable kinde is somewhat lesse, and tenderer then the last described, hauing also a reddish or blush coloured flower, lesser then the former, but much bigger then the middle kinde of Snapdragon (which is not set downe in this worke) the yellow spot in the mouth of it hath some white about it, and extending to both sides of the spot: the heads and seede are like the former: the rootes are smaller, but neuer will abide after they haue giuen flowers and seede.

#### 4. *Antirrhinum luteum.* Yellow Snapdragon.

There is likewise another of these kindes, that beareth leaues as large as any of the former, & very faire yellow flowers, as large likewise as they, not differing in any thing else from the first; let not any therefore imagine this to be a *Linaria* or Tode Flaxe: for all parts are answerable vnto the Snapdragons.

### The Place.

All these are nourished with vs in our Gardens, although in Spaine and Italy they are found growing wilde.

### The Time.

They flower for the most part the second yeare after the sowing, from Aprill vntill Iuly, and the seede is quickly ripe after.

### The Names.

The name *Antirrhinum* is vsually giuen to this plant, although it fully agreeth not eyther with the description of Dioscorides, or Theophrastus: It hath also diuers other names in Latine, as *Orontium*, *Canis cerebrum Os Leonis*, *Leo herba*, *&c.* In English Calues snout, from the forme of the seede vessels, and Snapdragon, or Lyons mouth, from the forme of the flowers.

### The Vertues.

They are seldome or neuer vsed in Phyficke by any in our dayes.

## Chap. XLV.

### *Chamænerium flore delphinÿ.* The Willowe flower.

THis plant riseth vp with many strong, woddy, round, browniſh great ſtalkes, three or foure foote high, beset here and there without order, with one broad and long whitish greene leafe at a ioynt, somewhat like vnto a *Lysimachia*, or Willow herbe, as also vnto a Peach leafe, but larger and longer: at the toppe of the branches stand many flowers one aboue another, of a pale reddish purple colour, consisting of fiue leaues, spread open with an heele or spurre behinde them, with many yellow threads in the middle, much larger then any flower of the Larkes spurres, and smelling somewhat sweete withall; it beareth a shew of long pods with seede, but I could neuer obserue the seede: the rootes are like the rootes of *Lysimachia*, or the ordinary yellow Looſe-ſtrife, or Willowe herbe, but greater: running and spreading vnder ground, and shooting vp in many places, whereby it filleth a ground that it likes quickly: the stalkes dye downe euery yeare, and spring againe in many places farre asunder.

### The Place.

Wee haue not knowne where this Willowe flower groweth naturally, but we haue it standing in an out corner of our Gardens, to fill vp the number of delightfull flowers.

### The Time.

It flowreth not vntill May, and abideth a long while flowring.

### The Names.

It may seeme to diuers, that this is that plant that Dodonæus called *Pseudolysimachium purpureum minus*, and Lobel seemeth by the name of *Delphinium buccinum* to aime at this plant, but withall calleth it *Chamænerium Gesneri*, and giueth the same figure that Dodonæus hath for his *Pseudolysimachium*: But that is one kinde of plant (which hath smaller and shorter stalkes, and very narrow long leaues, whose flowers stand vpon long slender cods, full of downe, with reddish seede, like vnto the *Lysimachia siliquosa siluestris*, and rootes that abide many yeares, but creepe not) and this is another, much greater, whose true figure is not extant in any Author that I know. It is vsually called *Chamænerium flore delphinÿ*; but the name of *Delphinium buccinum* in my minde may not so conueniently be applyed vnto it. It is called in English, The Willowe flower, for the likenesse of the leaues, and the beauty and respect of the flowers.

The

The Vertues.

There is no vse hereof in Physicke that euer I could learne, but is onely cherished among other sorts of flowers, that serue to decke and set forth a Garden of varieties.

## Chap. XLVI.

### *Aquilegia.* Colombines.

THere are many sorts of Colombines, as well differing in forme as colour of the flowers, and of them both single and double carefully noursed vp in our Gardens, for the delight both of their forme and colours.

### 1. *Aquilegia vulgaris flore simplici.* Single Colombines.

Because the whole difference of these Colombines standeth in the varieties of the forme, and colour of the flowers, and little in the leaues, I shall not neede to make anie repetitions of the description of them, seeing one onely shall suffice for each peculiar kinde. The Colombine hath diuers large spread leaues, standing on long stalkes: euery one diuided in seuerall partitions, and roundly endented about the edges, in colour somewhat like the leaues of Celondine, that is, of a darke blewish greene colonr: the stalkes rise vp sometimes two or three foote high, diuided vsually into many branches, bearing one long diuided leafe at the lower ioynt, aboue which the flowers growe, euery one standing on a long stalke, consisting of fiue hollow leaues, crooked or horned at the ends, turning backward, the open flower shewing almost like vnto a Cinquefoile, but more hollow: after the flowers are past, there arise small long cods, foure or fiue together, wherein are contained blacke shining seede: the rootes are thicke and round, for a little space within the ground, and then diuided into branches, ending in many small fibres, abiding many yeares, and shooting a fresh euery Spring from the round heads, that abide all the Winter. The variety of the colours of these flowers are very much, for some are wholly white, some of a blew or violet colour, others of a blush or flesh colour, or deepe or pale red, or of a dead purple, or dead murrey colour, as nature listeth to shew it selfe.

### 2. *Aquilegia vulgaris flore pleno.* Double Colombines.

The double Colombines differ not in leafe or manner of growing from the single, so that vntill they come to flower, they cannot bee discerned one from another; the onely difference is, it beareth very thicke and double flowers, that is, many horned or crooked hollow leaues set together, and are not so large as the leaues of the single flowers. The variety of colours in this double kinde is as plentifull, or rather more then in the single; for of these there is party coloured, blew and white, and spotted very variably, which are not in the single kinde, and also a very deepe red, very thicke and double, but a smaller flower, and lesse plentifull in bearing then many of the other double sorts. These double kindes doe giue as good seede as the single kindes doe, which is not obserued in many other plants.

### 3. *Aquilegia inuersis corniculis.* Double inuerted Colombines.

These Colombines are not to be distinguished eyther in roote, leaues, or seed from the former, the flowers onely make the difference, which are as double as the former, but that the heeles or hornes of these are turned inward, and stand out in the middle of the flowers together: there is not that plentifull variety of colours in this kinde, as there is in the former: for I neuer saw aboue three or foure seuerall colours in this kinde,

kinde, that is, white, purpliſh, reddiſh, and a dun or darke ouerworne purpliſh colour. Theſe double flowers doe likewiſe turne into pods, bearing ſeede, continuing his kind, and not varying into the former.

4. *Aquilegia Roſea.* Roſe Colombines.

The leaues and other parts of this kinde of Colombine, differ little or nothing from the former, the diuerſitie conſiſteth likewiſe in the flowers, which although they ſtand in the ſame manner ſeuerally vpon their ſmall ſtalkes, ſomewhat more ſparingly then the former doe, yet they haue no heeles or hornes, eyther inward or outward, or very ſeldome, but ſtand ſometimes but with eight or tenne ſmooth ſmall plaine leaues, ſet in order one by one in a compaſſe, in a double rowe, and ſometimes with foure or fiue rowes of them, euery one directly before the other, like vnto a ſmall thick double Roſe layd open, or a ſpread Marigold: yet ſometimes it happeneth, that ſome of theſe flowers will haue two or three of the firſt rowes of leaues without any heele, and the reſt that are inward with each of them a peece of a ſmall horne at them, as the former haue: the colours of theſe flowers are almoſt as variable, and as variably mixed as the former double kindes. This likewiſe giueth ſeede, preſeruing his owne kinde for the moſt part.

5. *Aquilegia degener.* Degenerate Colombines.

This kinde of Colombine might ſeeme to ſome, to bee but a caſuall degeneration, and no true naturall kinde, happening by ſome cauſe of tranſplanting, or otherwiſe by the art of man: but I haue not ſo found it, in that it keepeth, and holdeth his own proper forme, which is like vnto the double Roſe Colombine, but that the outermoſt row of leaues are larger then any of the reſt inwardes, and is of a greeniſh, or elſe of a purpliſh greene colour, and is not altogether ſo apt to giue good ſeed like the former.

## The Place.

The ſingle kindes haue beene often found in ſome of the wooddy mountaines of Germany, as Cluſius ſaith, but the double kindes are chiefly cheriſhed in gardens.

## The Time.

They flower not vntill May, and abide not for the moſt part when Iune is paſt, and in the meane time perfecteth their ſeede.

## The Names.

Coſtæus doth call this plant *Pothos* of Theophraſtus, which Gaza tranſlateth *Deſiderium.* Dalechampius vpon Athenæus, calleth it *Dioſanthos*, or *Iouis flos* of Theophraſtus, who in his ſixth Booke and ſeuenth Chapter reckoneth them both, that is, *Dioſanthos* and *Pathos*, to be Summer flowers, but ſeuerally. Dodonæus *Leoherba*, and Geſner *Leontoſtomium.* Fabius Columna in his Phytobaſanos, vnto whom Cluſius giueth the greateſt approbation, referreth it to the *Iſopyrum* of Dioſcorides. All later Writers doe generally call it, eyther *Aquileia*, *Aquilina*, or *Aquilegia*; and we in Engliſh, generally (I thinke) through the whole Countrey, Colombines. Some doe call the *Aquilegia roſea*, *Aquilegia ſtellata*, The ſtarre Colombine; becauſe the leaues of the flowers doe ſtand ſo directly one by another, beſides the doubleneſſe, that they ſomewhat repreſent eyther a Roſe or a Starre, and thereupon they giue it the name eyther of a Starre or Roſe.

## The Vertues.

Some in Spaine, as Camerarius ſaith, vſe to eate a peece of the roote here-

of

1 *Aquilegia simplex.* The single Colombine. 2 *Aquilegia flore multiplici.* The double Colombine. 3 *Aquilegia versicolor.* The party coloured Colombine. 4 *Aquilegia inuersis corniculis.* The double inuerted Colombine. 5 *Aquilegia Rosea sive Stellata.* The Rose or the Starre Colombine. 6 *Thalictrum Hispanicum album.* White Spanish tufts.

of fasting, many dayes together, to helpe them that are troubled with the stone in the kidneyes. Others vse the decoction, of both herbe and roote in wine, with a little Ambargrise, against those kinds of swounings, which the Greekes call ἀδυναμία. The seede is vsed for the iaundise, and other obstructions of the liuer. Clusius writeth from the experience of Franciscus Rapard, a chiefe Physician of Bruges in Flanders, that the seede beaten and drunke is effectuall to women in trauell of childe, to procure a speedy deliuerie, and aduiseth a second draught thereof should be taken if the first succeede not sufficiently.

## Chap. XLVII.

*Thalictrum Hispanicum.* Spanish Tufts, or Tufted Colombines.

From among the diuersities of this plant, I haue selected out two sorts for this my garden, as hauing more beautie then all the rest; leauing the other to be entreated of, where all in generall may be included. I haue in this place inserted them, for the likenesse of the leaues only, being in no other part correspondent, and in a Chapter by themselues, as it is most fit.

*Thalictrum Hispanicum album.* White Spanish tufted Colombines.

These plants haue both one forme, in roote, leafe and flower, and therefore neede but one description. The leaues are both for colour and forme so like vnto Colombines leaues (although lesser and darker, yet more spread, and on larger stalkes) that they may easily deceiue one, that doth not marke them aduisedly; for the leaues are much more diuided, and in smaller parts, and not so round at the ends: the stalkes are round, strong, and three foote high at the least, branching out into two or three parts, with leaues at the seuerall ioynts of them, at the toppes whereof stand many flowers, which are nothing but a number of threads, made like vnto a small round tuft, breaking out of a white skinne, or leafe, which incloseth them, and being vnblowne, shew like vnto little buttons: the colour of these threds or tufts in this are whitish with yellow tips on them, and somewhat purplish at the bottome, hauing a strong but no good sent, and abiding in their beautie (especially if they grow in the shade, and not too hot in the sun) a great while, and then fall away, like short downe or threds: the seed vessels are three square, containing small, long, and round seede; the rootes are many long yellow stringes, which endure and encrease much.

*Thalictrum Montanum purpureum.* Purple tufted Colombines.

This purple tufted Colombine differeth onely from the former, in that it is not so high nor so large, and that the colour of the flower or tuft is of a blewish purple colour with yellow tips, and is much more rare then the other.

### The Place.

These grow both in Spaine and Italie.

### The Time.

They flower in the end of May, or in Iune, and sometime later.

### The Names.

Some doe call them *Thalictrum*, and some *Thalictrum*. Others *Ruta paistris*, and *Ruta pratensis*, and some *Rhabarbarum Monachorum*, or *Pseudorhabarbarum*,

*rhabarbarum*, by reason that the rootes being yellow, haue an opening qualitie, and drying as Rubarbe. In English what other fit Names to giue these then I haue expressed in the titles, I know not.

### The Vertues.

The are a little hot and drying withall, good for old Vlcers, as Dioscorides saith, to bring them to cicatrising: in Italy they are vsed against the Plague, and in Saxonye against the Iaundise, as Camerarius saith.

## CHAP. XLVIII.

### *Radix caua.* Hollow roote.

THe likenesse of the leaues likewise of this plant with Colombines, hath caused mee to insert it next the other, and although some of this kinde bee of small respect, being accounted but foolish, yet let it fill vp a waste corner, that so no place be vnfurnished.

1. *Radix Caua maior flore albo.* The white Hollow roote.

The leaues of this hollow roote breake not out of the ground, vntill the end of March, or seldome before, and are both for proportion and colour somewhat like vnto the leaues of Colombines, diuided into fiue parts, indented about the edges, standing on small long footestalkes of a whitish greene colour, among which rise vp the stalkes, without any leaues from the bottome to the middle, where the flowers shoote forth one aboue another, with euery one a small short leafe at the foote thereof, which are long and hollow, with a spurre behinde it, somewhat like vnto the flowers of Larckes spurres, but hauing their bellies somwhat bigger, and the mouth not so open, being all of a pure white colour: after the flowers are past, arise small long and round cods, wherein are contained round blackish seede: the roote is round and great, of a yellowish browne colour on the outside, and more yellow within, and hollow vnderneath, so that it seemeth but a shell: yet being broken, euery part will grow: it abideth greene aboue ground but a small time.

2. *Radix Caua maior flore carneo.* Blush colourd Hollow roote.

The blush Hollow roote is in all things like vnto the former, but onely that the flowers hereof are of a delayed red or purple colour, which we call blush: and sometimes of a very deepe red or purple colour; but very rare to meete with.

3. *Radix Caua minor, seu Capnos fabacea radice.* Small hollow roote.

This small kinde hath his leaues of a blewish greene colour, yet greener and smaller then the former, growing more thicke together: the flowers are like in proportion vnto the former in all respects, but lesser, hauing purplish backes, and white bellyes: standing closer and thicker together vpon the short stalkes: the roote is solid or firme, round and a little long withall, two being vsually ioyned together, yellowish both within and without: but I haue seene the dry roots that came from beyond Sea hither, that haue beene as small as hasell nuts, and somewhat flat with the roundnesse, differing from those that growe with vs, whether the nature thereof is to alter by manuring, I know not.

### The Place.

The greater kindes Clusius reporteth he found in many places of Hungarie,

rie, and the other parts neere thereunto: the lesser in the lower Germany, or Low Countries, as we call them.

### The Time.

These are most truely to bee reckoned Vernall plants, for that they rise not out of the ground vntill the Spring bee come in, and are gone likewise before it be past, remaining vnder ground all the rest of the yeare, yet the lesser abideth longer aboue ground then the greater.

### The Names.

Concerning the former of these, there is a controuersie among diuers, whether it should be *Thesium* of Theophrastus, or *Eriphium* of Galen, but here is no fit place to trauerse those opinions. Some would haue it to bee *Corydalis*, and some referre it to Plinie his *Capnos Cheledonia*, for the likenesse it hath both with Fumeterie and Celandine. It is generally called of all moderne Writers, *Radix Cana*, and we in English thereafter, Hollow roote. The lesser for the firmenesse of his round roote, is vsually called, *Capnos fabacea radice*, and the Dutch men thereafter, **Boonkens Hollwortell:** we of the likenesse with the former, doe call it the lesse Hollow roote.

### The Vertues.

Some by the bitternesse doe coniecture (for little proofe hath beene had thereof, but in outward cases) that it clenseth, purgeth, and dryeth withall.

## CHAP. XLIX.

### *Delphinium.* Larkes heeles.

OF Larkes heeles there are two principall kindes, the wilde kinde, and the tame or garden; the wilde kinde is of two sorts, one which is with vs noursed vp chiefly in gardens, and is the greatest; the other which is smaller and lower, often found in our plowed landes, and elsewhere: of the former of these wilde sorts, there are double as well as single: and of the tame or more vpright, double also and single: and of each of diuers colours, as shall be set downe.

#### 1. *Delphinium maius sive vulgare.* The ordinary Larkes heeles.

The common Larkes heele spreadeth with many branches much more ground then the other, rather leaning or bending downe to the ground, then standing vpright, whereon are set many small long greene leaues, finely cut, almost like Fennell leaues: the branches end in a long spike of hollow flowers, with a long spurre behinde them, very like vnto the flowers of the Hollow roote last described, and are of diuers seuerall colours, as of a blewish purple colour, or white, or ash colour or red, paler or deeper, as also party coloured of two colours in a flower: after the flowers are past, (which in this kinde abide longer then in the other) there come long round cods, containing very blacke seede: the root is hard after it groweth vp to seede, spreading both abroad and deepe, and perisheth euery yeare, vsually raising it selfe from it own sowing, as well as from the seede sowen in the spring time.

*Varietas.*

#### 2. *Delphinium vulgare flore pleno.* Double common Larkes heeles.

Of this vulgar kinde there is some difference in the flower, although in nothing else: the flowers stand many vpon a stalke like the former, but euery one of them are as if three

three or foure small flowers were ioyned together, with euery one his spurre behinde; the greatest flower being outermost, and as it were containing the rest, which are of a pale red, or deepe blush colour : Another of this kinde will beare his flowers with three or foure rowes of leaues in the middle, making a double flower with one spurre behinde onely : and of this kinde there is both with purple, blew, blush, and white flowers, and party coloured also; these doe all beare feed like the single, wherby it is encreased euery yeare.

3. *Delphinium aruense.* Wilde Larkes spurres.

This wilde Larkes spurre hath smaller and shorter leaues, smaller and lower branches, and more thinly or sparsedly growing vpon them, then any of the former : the flowers likewise are neyther so large as any of the former, nor so many growing together, the cods likewise haue smaller seede, and is harder to grow in gardens then any of the former; the most vsuall colour hereof is a pale reddish or blush colour, yet sometimes they are found both white and blew, and sometimes mixt of blew and blush, variably disposed, as nature can when she listeth ; but are much more rare.

4. *Diphinium elatius flore simplici diuersorum colorum.*
Single vpright bearing Larkes heeles of many colours.

The difference betweene this and the last is, that the leaues of this are not fully so greene, nor so large; the stalkes grow vpright, to the height of a man, and sometimes higher, hauing some branches thereon, but fewer then the former, and standing likewise vpright, and not leaning downe as the former : the toppes of the stalkes are better stored with flowers then the other, being sometimes two foote long and aboue, of the same fashion, but not altogether so large, but of more diuers and seueral colours, as white, pale, blush, redde deeper or paler, ashcoloured, purple or violet, and of an ouerworne blewish purple, or iron colour : for of all these we haue simple, without any mixture or spot : but we haue other sorts, among the simple colours, that rise from the same seede, and will haue flowers that wil be halfe white, and halfe blush or purple, or one leafe white, and another blush or purple, or else variably mixed and spotted: the seede and seede vessels are like the former but larger and harder.

5. *Delphinium elatius flore pleno diuersorum colorum.*
Double vpright Larkes heeles of many colours.

These double Larkes heeles cannot bee knowne from the single of the same kinde, vntill they come towards flowring; for there appeare many flowers vpon the stalkes, in the same manner, and of as many colours almost as of the single, except the party coloured, which stand like little double Roses, layd or spread broade open, as the Rose Colombine without any heeles behinde them, very delightfull to behold, consisting of many small leaues growing together, and after they are fallen there come vp in their places three or foure small cods set together, wherein is contained here and there (for all are not full of seede, as the single kindes) blacke seede, like vnto all the rest, but smaller, which being sowen will bring plants that will beare both single and double flowers againe, and it often happeneth, that it variably altereth in colours from it owne sowing : for none of them hold constantly his owne colour, (so farre as euer I could obserue) but fall into others as nature pleaseth.

6. *Delphinium Hispanicum paruum.* Spanish wilde Larkes spurres.

This small Larkes spurre of Spaine, hath diuers long and broad leaues next the ground, cut-in on both sides, somewhat like vnto the leafe of a Scabious, or rather that kinde of Stœbe, which Lobel calleth *Crupina*, for it doth somewhat neerly resemble the same, but that this is smooth on the edges, and not indented besides the cuts, as the *Crupina* is, being of a whitish greene colour, and somewhat smooth and soft in handling : among the leaues riseth vp a whitish greene stalke, hauing many smaller

leaues vpon it that grow belowe, but not diuided, branching out into many ſmall ſtalkes, bearing flowers like vnto the wilde Larkes heeles, but ſmaller, and of a bleake blewiſh colour, which being paſt, there come vp two or three ſmall cods ioyned together, wherein is blacke ſeede, ſmaller and rounder then any of the former: the roote is ſmall and thready, quickly periſhing with the firſt cold that ouertaketh the plant.

### The Place.

The greateſt or firſt wilde kindes growe among corne in many countries beyond the Seas, and where corne hath beene ſowne, and for his beauty brought and nouriſhed in our Gardens: the leſſer wilde kinde in ſome fields of our owne Country. The Spaniſh kinde likewiſe in the like places, which I had among many ſeedes that Guillaume Boel brought mee out of Spaine. The firſt double and ſingle haue been common for many yeares in all countries of this Land, but the tall or vpright ſingle kindes haue been entertained but of late yeares. The double kindes are more rare.

### The Time.

Theſe flower in the Summer onely, but the Spaniſh wilde kinde flowreth very late, ſo that oftentimes in our Country, the Winter taketh it before it can giue ripe ſeede: the double kindes, as well the vpright as the ordinary or wilde, are very choiſe and dainty many times, not yeelding good ſeede.

### The Names.

They are called diuerſly by diuers Writers, as *Conſolida regalis*, *Calcaris flos*; *Flos regius*, *Buccinum Romanorum*, and of Matthiolus, *Cuminum ſilueſtre alterum Dioſcoridis*: but the moſt vſuall name with vs is *Delphinium*: but whether it be the true *Delphinium* of Dioſcorides, or the Poets Hyacinth, or the flower of Aiax, another place is fitter to diſcuſſe then this. Wee call them in Engliſh Larkes heeles, Larkes ſpurres, Larkes toes or clawes, and Monkes hoods. The laſt or Spaniſh kinde came to mee vnder the name of *Delphinium latifolium trigonum*, ſo ſtiled eyther from the diuiſion of the leaues, or from the pods, which come vſually three together. Bauhinus vpon Matthiolus calleth it, *Conſolida regalis peregrina paruo flore*.

### The Vertues.

There is no vſe of any of theſe in Phyſicke in theſe dayes that I know, but are wholly ſpent for their flowers ſake.

## Chap. L.

*Balſamina fœmina.* The Female Balſam Apple.

I Haue ſet this plant in this place, for ſome likeneſſe of the flower, rather then for any other compariſon, euen as I muſt alſo with the next that followeth. This plant riſeth vp with a thicke round reddiſh ſtalke, with great and bunched ioynts, being tender and full of iuice, much like to the ſtalke of Purſlane, but much greater, which brancheth it ſelfe forth from the very ground, into many ſtalkes, bearing thereon manie long greene leaues, ſnipt about the edges, very like vnto the Almond or Peach tree leaues; among which from the middle of the ſtalkes vpwards round about them, come forth vpon ſeuerall ſmall ſhort foot-ſtalkes many faire purpliſh flowers, of two or three colours in them, faſhioned ſomewhat like the former Larkes heeles, or Monks hoods, but that they are larger open at the mouth, and the ſpurres behinde crooke or bend downewards: after the flowers are paſt, there come in their places round rough heads,

1 *Radix Cana maior flore albo.* The white flowred Hollow roote. 2 *Capnos fabacea radice.* The small Hollow roote. 3 *Delphinium flore simplici.* Single Larkes spurs. 4 *Delphinium vulgare flore medio duplici.* Larkes spurs double in the middle. 5 *Delphinium vulgare flore pleno.* Common Larks spurs double. 6 *Delphinium elatius flore pleno.* Double vpright Larkes spurs. 7 *Delphinium Hispanicum parvum.* Small Spanish Larkes spurs. 8 *Balsamina fœmina.* The Female Balsam apple. 9 *Nasturtium Indicum.* Indian Cresses, or yellow Larkes spurs.

heads, pointed at the end, greene at the first, and a little yellower when they bee ripe, containing within them small round blackish seede, which will soone skippe out of the heads, if they be but a little hardly pressed betweene the fingers: the rootes spread themselues vnder ground very much from the toppe, with a number of small fibres annexed thereunto: this is a very tender plant, dying euery yeare, and must bee sowne carefully in a pot of earth, and tended and watered in the heate of Summer, and all little enough to bring it to perfection.

### The Place.

Wee haue alwaies had the seede of this plant sent vs out of Italy, not knowing his originall place.

### The Time.

It flowreth from the middle of Iuly, to the end of Auguſt: the seed doth seldome ripen with vs, especially if the Summer be backward, so that wee are oftentimes to seeke for new and good seede from our friends againe.

### The Names.

Some vse to call it *Charantia fœmina*, *Balsamina fœmina*, *Balsamella*, and *Anguillara*, *Herba Sancta Katharina*. We haue no other English name to call it by, then the Female Balsame Apple, or *Balsamina*.

### The Vertues.

Some by reason of the name, would attribute the property of Balme vnto this plant, but it is not sufficiently knowne to haue any such; yet I am well perswaded, there may bee some extradinary quality in so beautifull a plant, which yet lyeth hid from vs.

## CHAP. LI.

*Nasturtium Indicum.* Indian Cresses, or yellow Larkes heeles.

THe likenesse (as I said before) of this flower likewise, hauing spurres or heeles maketh me ioyne it with the rest, which is of so great beauty and sweetnesse withall, that my Garden of delight cannot bee vnfurnished of it. This faire plant spreadeth it selfe into very many long trayling branches, enterlaced one within another very confusedly (yet doth it not winde it selfe with any claspers about either pole or any other thing, but if you will haue it abide close thereunto, you must tye it, or else it will lye vpon the ground) foure or fiue foot in length at the least, wherby it taketh vp a great deale of ground: the leaues are smooth, greene, and as round as the Penniwort that groweth on the ground, without any cut or incisure therein at all in any part, the stalkes whereof stand in the middle of each leafe, and stand at euery ioynt of the stalke, where they are a little reddish, and knobbed or bunched out: the flowers are of an excellent gold yellow colour, and grow all along these stalkes, almost at euery ioynt with the leaues, vpon pretty long foote-stalkes, which are composed of fiue leaues, not hollow or gaping, but standing open each leafe apart by it selfe, two of them, that be larger and longer then the other, stand aboue, and the other two that are lesser belowe, which are a little iagged or bearded on both sides, and the fift lowest: in the middle of each of the three lower leaues (yet sometimes it is but in two of them) there is a little long spot or streake, of an excellent crimson colour, with a long heele or spurre behinde hanging downe: the whole flower hath a fine small sent, very pleasing, which being placed in the middle of some Carnations or Gillo-

flowers

flowers (for they are in flower at the same time) make a delicate Tussimussie, as they call it, or Nosegay, both for sight and sent: After the flower is past, come the seede, which are rough or vneuen, round, greenish yellow heads, sometimes but one, and sometimes two or three standing together vpon one stalke, bare or naked of themselues, without any huske, containing a white pulpy kernell: the rootes are small, and spreading vnder ground, which perish with the first frosts, and must be sowne a new euery yeare; yet there needeth no bed of horse-dung for the matter: the naturall ground will be sufficient, so as you defend it a little from those frosts, that may spoile it when it is newly sprung vp, or being yet tender.

### The Place.

This goodly plant was first found in the West Indies, and from thence sent into Spaine vnto Monardus and others, from whence all other parts haue receiued it. It is now very familiar in most Gardens of any curiosity, where it yearly giueth ripe seed, except the yeare be very vnkindly.

### The Time.

It flowreth sometimes in Iune, but vsually in Iuly (if it be well defended and in any good ground) and so continueth flowring, vntill the cold frosts and mistes in the middle or end of October, doe checke the luxurious nature thereof, and in the meane time the seede is ripe, which will quickly fall downe on the ground, where for the most part the best is gathered.

### The Names.

Some doe reckon this plant among the *Clematides* or *Conuolvuli*, the Clamberers or Bindweedes; but (as I said) it hath no claspers, neither doth it winde it selfe: but by reason of the number of his branches, that run one within another, it may seeme to climbe vp by a pole or sticke, which yet doth but onely close it, as hauing something whereon to leane or rest his branches. Monardus and others call it *Flos sanguineus*, of the red spots in the flowers, as also *Mastuerzo de las Indias*, which is *Nasturtium Indicum*, by which name it is now generally knowne and called, and wee thereafter in English, Indian Cresses, yet it may bee called from the forme of the flowers onely, Yellow Larkes heeles.

### The Vertues.

The Spaniards and others vse the leaues hereof in stead of ordinary Cresses, because the taste is somewhat sharpe agreeing thereunto, but other Physicall properties I haue heard of none attributed to it.

## Chap. LII.

### *Viola.* Violets.

THe Garden Violets (for the Wilde I leaue to their owne place) are so well knowne vnto all, that either keepe a Garden, or hath but once come into it, that I shall (I thinke) but lose labour and time to describe that which is so common. Yet because it is not onely a choise flower of delight, notwithstanding the popularity, and that I let not passe any thing without his particular description, I must also doe so by this. And hereunto I must adde that kinde of Violet, which, although it want that smell of the other, goeth beyond it in variety of dainty colours, called *Viola tricolor & flammea*, or Harts eases.

1. *Viola simplex Martia.* Single March Violets.

The single Garden Violet hath many round greene leaues, finely snipt or dented about the edges, standing vpon seuerall small stalkes, set at diuers places of the many creeping branches, which as they runne, doe here and there take roote in the ground, bearing thereon many flowers seuerally at the ioynts of the leaues, which consist of fiue small leaues, with a short round tayle or spurre behinde, of a perfect blew purple colour, and of a very sweete sent, it bringeth forth round seede vessels, standing likewise vpon their seuerall small stalkes, wherein is contained round white seede: but these heads rise not from where the flowers grew, as in all other plants that I know, but apart by themselues, and being sowne, will produce others like vnto it selfe, whereby there may be made a more speedy encrease to plant a Garden (as I haue done) or any other place, then by slipping, as is the vsuall manner: the rootes spread both deepe and wide, taking strong hold in the ground.

*Flore albo.* Of this kinde there is another that beareth white flowers, not differing in smell or any thing else from the former.

*Flore obsoleto.* And also another, that beareth flowers of a dead or sad reddish colour, in all other things alike, sauing that this hath not altogether so good a sent as the other.

2. *Viola Martia flore multiplici.* Double March Violets.

There is no difference betweene this Violet and the former, in any other thing then in the doubleneſſe of the flowers, which haue so many leaues set and thrust together, that they are like vnto hard buttons. There is of this double kinde both white and purple, as in the single; but the white sort is seldome so thicke and double as the purple: but of the red colour to be double I neuer heard.

3. *Viola flammea siue tricolor.* Harts eases or Pansies.

The Harts ease hath his leaues longer, and more endented or cut in on the edges then the Violet hath, and somewhat round withall: the stalkes are vpright, yet weake, and ready to fall downe, and lye vpon the ground, set here and there with the like leaues, from whence come forth the flowers, of little or no sent at all, made like vnto a Violet, yet more open, and with larger leaues; but so variably mixed with blew or purple, white and yellow, that it is hard to set downe all the varieties: For some flowers will be more white, and but some spots of purple or blew in the two vpper leaues, and the lower leaues with some stripes of yellow in the middle: others will haue more purple in them then any other colour, both in the vpper and lower leaues, the side leaues blew, and the middle yellow, and others white and blew with yellow stripes, as nature listeth to distribute their colours: the seede is small, whitish, and round, contained in small round heads: the roote perisheth euery yeare, and raiseth it selfe vp plentifully by it owne sowing, if it be suffered.

4. *Viola tricolor flore duplici.* Double Harts ease.

We haue in our Gardens another sort, that beareth flowers with more leaues then the former, making it seeme to be twice double, and that onely in Autumne; for the first flowers are single that come in Summer: This is of that sort that beareth purple flowers: And it is to be obserued, that the seed of this kinde will not all bring double flowers, but only some, if the ground be fit and liking, so that if you haue once had of this double kinde, you shall seldome misse to haue double flowers againe euery yeare of it owne growing or sowing.

5. *Viola flammea lutea maxima.* The great yellow Pansie.

There is one other kinde of Harts ease, that decketh vp our Gardens not to be forgotten, whose leaues and flowers are like the former, but more plentifull in stalkes and branches, and better abideth our Winters: the flowers are larger then any of the former,

former, of a faire pale yellow colour, with some yellower stripes now and then about the middle; for it is sometimes without any stripes, and also of a little deeper yellow colour: this is to bee encreased by slips, which will soone comprehend in a moist or moistened ground, for that I neuer could obserue that it bore seede.

### The Place.

These plants were first wilde, and by manuring brought to be both fairer in colour, and peraduenture of a better sent then when they grew wilde.

### The Time.

The Violets flower in March, and sometimes earlier, and if the yeare be temperate and milde, in Autumne againe. The double Violets, as they are later before they flower then the single, so they hold their flowers longer. The Harts ease flowreth seldome vntill May; but then some will abide to flower vntill the end of Autumne almost, especially if the frosts be not early.

### The Names.

The Violet is called *Viola nigra*, *purpurea*, and *Martia*: In English, Violets, March Violets, and purple Violets. The Harts ease is called *Viola flammea*, *Viola tricolor*, *Viola multicolor*, and of some, *Iacea*, *Flos trinitatis*, and *Herba clauellata*: In English, Harts ease, and Pansies, of the French name *Pensees*. Some giue it foolish names, as Loue in idlenesse, Cull mee to you, and Three faces in a hood. The great yellow Harts ease is so called, because it is like in forme, and is the greatest of all other, although it haue not that diuersity of colours in it that the other haue.

### The Vertues.

The properties of Violets are sufficiently knowne to all, to coole and moisten: I shall forbeare to recite the many vertues that may be set downe, and onely let you know, that they haue in them an opening or purging quality, being taken either fresh and greene, or dryed, and made into powder, especially the flowers; the dryed leaues will doe the like, but in greater quantity. Costæus in his booke of the nature of all plants saith, that the distilled water of Harts ease, is commended in the French disease, to be profitable, being taken for nine dayes or more, and sweating vpon it, which how true it is, I know not, and wish some better experience were made of it, before we put any great confidence in that assertion.

## Chap. LIII.

### *Epimedium.* Barrenwort.

THis pretty plant riseth vp out of the ground with vpright, hard, round, small stalkes, a foote and a halfe high, or not two foote high at the highest, diuided into three branches for the most part, each branch whereof is againe diuided for the most part into three other branches, and each of them beare three leaues (seldome either more or lesse) set together, yet each vpon his owne foote-stalke, each leafe being broad, round, and pointed at the end, somewhat hard or dry in feeling, hayrie, or as it were prickly about the edges, but very tenderly, without harme, of a light greene colour on the vpperside, and a little whiter vnderneath: from the middle of the stemme or stalke of leaues doth likewise come forth another long stalke, not much higher then those with the leaues on them, diuided into other branches, each

whereof

whereof hath likewise three flowers, each vpon his owne footestalke, consisting of eight small leaues a peece, yet seeming to be but of foure leaus spread or layd open flat, for that the foure vppermost, which are the smaller and being yellow, doe lye so close on the foure vndermost, wch are a little broader and red, that they shew as if they were yellow flowers with red edges, hauing yellow threds tipt with greene, standing in the middle of the flowers: the vnderside of the lower leaues are of a pale yellowish red, striped with white lines: after the flowers are past, there come small long pods, wherin are contained flat reddish seede: the rootes are small, reddish and hard, spreading, branching and enterlacing themselues very much, and is fit to be placed on some shady side of a garden: the whole plant is rather of a strong then any good sent, yet is cherished for the pleasant varietie of the flowers.

### The Place.

Cæsalpinus saith it groweth on the mountaines of Liguria, that is nigh vnto Ligorne, in the Florentine Dominion. Camerarius saith, nigh vnto Vicenzo in Italie. Bauhinus on the Euganian hils, nigh vnto Padoa, and in Romania in shadowie wet grounds.

### The Time.

It flowreth from Iune vntill the end of Iuly, and to the middle of August, if it stand, as I said it is fittest, in a shadowie place.

### The Names.

It is of most Writers accepted for the true *Epimedium* of Dioscorides, though he saith it is without flower or seede, being therein eyther mistaken, or mis-informed, as he was also in *Dictamnus* of Candy, and diuers other plants. From the triple triplicitie of the standing of the stalkes and leaues, and quadriplicitie of the flowers, it might receiue another name in English then is already imposed vpon it: but lest I might be thought to be singular or full of noueltie, let it passe with the name Barrenwort, as it is in the title.

### The Vertues.

It is thought of diuers to agree in the propertie of causing barrennesse, as the ancients doe record of *Epimedium*.

## Chap. LIIII.

### *Papaver satiuum.* Garden Poppies.

OF Poppies there are a great many sorts, both wilde and tame, but because our Garden doth entertaine none, but those of beautie and respect, I wil onely giue you here a few double ones, and leaue the rest to a general survey.

#### 1. *Papaver multiplex album.* Double white Poppies.

The double white Poppy hath diuers broade, and long whitish greene leaues, giuing milke (as all the rest of the plant aboue ground doth, wheresoeuer it is broken) very much rent or torne in on the sides, and notched or indented besides, compassing at the bottome of them a hard round brittle whitish greene stalke, branched towards the toppe, bearing one faire large great flower on the head of euery branch, which before it breaketh out, is contained within a thin skinne, and being blowne open is very thick of leaues, and double, somewhat iagged at the ends, and of a white colour; in the middle

1 *Viola Martia simplex.* Single March Violets. 2 *Viola Martia multiplex.* Double March Violets. 3 *Viola flammea siue tricolor.* Ordinary garden Pansies or Harts ease. 4 *Viola flammea lutea magna.* Great yellow Pansies. 5 *Viola trieglor duplex* Double Pansies or Harts eases. 6 *Epimedium.* Barrenwort. 7 *Papauer sativum flore pleno.* Double garden Poppies. 8 *Papauer sativum flore pleno laciniato.* Double feathered Poppies. 9 *Nigella Hispanica flore ample.* Spanish Nigella or Fenell flower. 10 *Nigella multiplex cærulea* Double blew Nigella or Fenell flower. 11 *Nigella duplex flore albo.* Double white Nigella. 12 *Tormina flore pleno.* Double wilde Pellitory.

middle whereof ſtandeth a round head or bowle, with a ſtriped crowne on the heade of it, very like a ſtarre, compaſſed about with ſome threds, wherein when it is ripe, is contained ſmall, round, white ſeede, diſpoſed into ſeuerall cels : the roote is hard, wooddy, and long, periſhing euery yeare, and muſt bee new ſowne euery Spring, if they doe not ſpring of their own ſowing, which if it doe, the flowers are ſeldome ſo faire and double as they that are ſowne in the Spring : the whole plant is of a ſtrong heady ſmell.

2. *Papauer multiplex rubeſcens.* Double red or bluſh Poppies.

This other kind of double Poppy differeth not in any other thing from the former, but only in the colour of the flowers, which are of a bright red, tending to a bluſh colour, parted, paned or ſtriped in many places with white, and exceedingly more iagged then the former, almoſt like a feather at the ends, the bottomes of all the leaues being white : the ſeede hereof is white as the former, which is not ſo in any other Poppie, that beareth not a full white flower.

3. *Papauer multiplex nigrum ſiue purpureum.*
Double purple or murry Poppies.

This kinde varyeth both in flowers and ſeede, although neyther in leaues or any other thing from the firſt : the flowers are thicke and double, and ſomewhat iagged at the ends, in ſome more, in ſome leſſe, eyther red or bluſh, or purpliſh red, more or leſſe, or of a ſad murrey or tawney, with browne, or blacke, or tawny bottomes : the ſeede is eyther of a grayiſh blew colour, or in others more blackiſh.

4. *Papauer Rhœas flore multiplici.* The double red field Poppie.

This double Poppie is like the wilde or fielde Poppie, which is well knowne to all to haue longer, narrower, and more iagged greene leaues then the former, the ſtalkes more hairy, and the flower of a deepe yellowiſh red colour, knowne to all. Now this differeth in nothing from it, but in the doubleneſſe of the flower, which is very thicke and double, but not ſo large as the former. This riſeth of ſeede in the like manner as they doe, and ſo to bee preſerued.

## The Place.

From what place they haue beene firſt gathered naturally I cannot aſſure you, but we haue had them often and long time in our gardens, being ſent from Italie and other places. The double wilde kindes came from Conſtantinople, which whether it groweth neere vnto it or further off, we cannot tell as yet.

## The Time.

They flower in the beginning or middle of Iune at the furtheſt, the ſeede is ripe within a ſmall while after.

## The Names.

The generall knowne name to all, is *Papauer*, Poppie : the ſeuerall diſtinctions are according to their colours. Yet our Engliſh Gentlewomen in ſome places, call it by a by-name, Ione ſiluer pinne : *ſubauditur*, Faire without and fowle within.

## The Vertues.

It is not vnknowne, I ſuppoſe to any, that Poppie procureth ſleepe, for which cauſe it is wholly and onely vſed, as I thinke : but the water of the wilde

wilde Poppies, besides that it is of great vse in Pleuresies, and Rheumatick-or thinne Distillations, is found by daily experience, to bee a soueraigne remedy against surfeits ; yet some doe attribute this propertie to the water of the wilde Poppies.

## CHAP. LV.

*Nigella.* The Fenell flower, or Nigella.

AMong the many sorts of Nigella, both wilde and tame, both single and double, I will onely set downe three sorts, to be nourled vp in this garden, referring the rest to a Physicke garden, or a generall Historie, which may comprehend all.

1. *Nigella Hispanica flore simplici.* The great Spanish Nigella.

Spanish Nigella riseth vp with diuers greene leaues, so finely cut, and into so many parts, that they are finer then Fenell, and diuided somewhat like the leaues of Larkes heeles, among which rise vp stalkes, with many such like leaues vpon them, branched into three or foure parts, at the toppe of each whereof standeth one faire large flower, like vnto other single Nigella's, consisting of fiue or six leaues sometimes, of a bleake blew, or of a purplish blew colour, with a greene head in the middle, compassed about with seuen or eight small blewish greene flowers, or peeces of flowers rather, made like gaping hoodes, with euery of them a yellowish line thwart or crosse the middle of them, with some threds also standing by them : after the flower is past the head groweth greater, hauing fiue, seuen or eight hornes as it were at the toppe, greater and longer, and standing closer together then any other Nigella, spreading very like a starre, or the crowne of the Poppy head, but larger and longer, each whereof being folded together, openeth a little when the head is ripe, which is greater aboue, and smaller below, and not so round as the others are, containing within them small yellowish greene seede, or not so blacke as the other sorts : the rootes are small and yellow, perishing euery yeare as the others likewise doe.

2. *Nigella Damascena flore multiplici.*
Double blew Nigella, or The Fenell flower.

The double Nigella is in leaues, stalkes and rootes, very like vnto the former Nigella, so that the one can very hardly bee discerned from the other before this rise vp to flower, except it be that the leaues hereof are not fully so large as they : the flower consisteth of three or foure rowes of leaues, layde one vpon another, of a pale blew colour, with a greene round head compassed with diuers short threads in the middle, and hauing fiue or sixe such small greene Fenell-like leaues vnder the flower, to beare it vp (as it were) below, which adde a greater grace to the flowers, which at the first sheweth sometimes white, but changeth quickely after : the horned heads hereof are like vnto the heads of the other wilde kinde, which are somewhat rounder and greater, hauing within them blacke vneuen seedes, but without any sent.

3. *Nigella Catrina flore albo multiplici.* Double white Nigella.

This double white Nigella hath such like leaues as the last hath, but somewhat larger, of a yellower greene colour, and not so finely cut and iagged : the flowers are somewhat lesse, and lesser double then the former, and in colour white, hauing no greene leaues vnder the flower, as the former hath, the head whereof in the middle is very like the head of the last double kinde, but not so great, wherein is contained black seede for the most part, and sweete like the Romane Nigella, which only is sweet besides this : yet sometimes it is not so blacke, but rather a little more white or yellowish: the roote is yellow, and perisheth as the others euery yeare.

The

### The Place.

All these, and the rest be found wilde in diuers Countreyes, as France, Spaine, Italie, &c. but wee onely cherish them in our Gardens for our delight.

### The Time.

They flower in the end of Iune, and in Iuly, or thereabouts.

### The Names.

They are called *Melanthium, Gith*, and *Nigella*, and of some *Flos Diuæ Catherinæ*. We may either call them *Nigella* according to the Latine name, or the Fenell flower, as some doe, because the double blew Nigella hath small Fenell-like leaues bearing vp the flower, as I shewed before in the description.

### The Vertues.

These Nigella's are nothing so hot in qualitie as the single Romane kind is, as may well be knowne by the smell of the seede thereof, and therefore are not fit to be vsed in the steed of it, as many ignorant persons vse to doe: for the single Romane seede is vsed to helpe paines, and cold distillations in the head, and to dry vp the rheume. Pena saith, that the presſed oyle of the seede as well taken inwardly as vsed outwardly is an excellent remedy for the hardnesse and swelling of the spleene.

## Chap. LVI.

*Ptarmica siluestris flore pleno.* Double wilde Pelletory.

THe double wilde Pelletorie hath straight and slender stalkes, beset with long and narrow leaues, snipt round about the edges, in all points like vnto the single wilde kinde, that groweth common with vs almost euery where: on the toppes of the stalkes stand foure or fiue, or more white flowers, one aboue another, with a greene leafe at the bottome of the footestalke of euery one of them, beeing small, thicke, and very double, with a little yellowishnesse in the middle of euery flower, like both for forme and colour vnto the flower of the double Featherfew, but smaller: the rootes are many long strings, running here and there in the ground: this hath no smell at all, but is delightsome only for the double white flowers.

### The Place.

It is only cherished in some few Gardens, for it is very rare.

### The Time.

It flowreth in the end of Iune or thereabouts.

### The Names.

It is called of most *Ptarmica*, or *Sternutamentoria*, of his qualitie to prouoke neesing; and of some *Pyrethrum*, of the hot biting taste. We vsually call it Double wilde Pelletorie, and some Sneesewort, but *Elleborus albus* is vsually so called, and I would not two things should be called by one name, for the mistaking and mis-using of them.

The

### The Vertues.

The properties hereof, no doubt, may well bee referred to the single kinde, beeing of the same qualitie, yet as I take it, a little more milde and temperate.

## Chap. LVII.

*Parthenium flore pleno.* Double Featherfew.

Featherfew that beareth double flowers is so like vnto the single kinde, that the one cannot be discerned from the other, vntill it come to flower, bearing broad, pale or fresh greene leaues, much cut in on the sides : the stalkes haue such like leaues on them as grow below, from the toppes whereof come forth many double white flowers, like vnto the flowers of the former wilde Pelletory, but larger, and like also vnto the flowers of the double Camomill : the sent whereof is as strong as of the single.

### The Place.

We haue this kinde only in Gardens, and as it is thought by others, is peculiar only to our owne Countrey.

### The Time.

It flowreth in the end of May, and in Iune and Iuly.

### The Names.

It is called diuersly by diuers : Some thinke it to be *Parthenium* of Dioscorides, but not of Galen; for his *Parthenium* is a sweet herbe, and is thought to bee *Amaracus*, that is Marierome : others call it *Matricaria*; and some *Amarella*. Gaza translateth it *Muraleum*, *Theoph. lib. 7. cap. 7.* It is generally in these parts of our Country called Double Feaverfew, or Featherfew.

### The Vertues.

It is answerable to all the properties of the single kinde which is vsed for womens diseases, to procure their monthly courses chiefly. It is held to bee a speciall remedy to helpe those that haue taken *Opium* too liberally. In Italy some vse to eate the single kinde among other greene herbes, as Camerarius saith, but especially fryed with egges, and so it wholly loseth his strong and bitter taste.

## Chap. LVIII.

*Chamæmelum.* Camomill.

Our ordinary Camomill is well knowne to all, to haue many smal trayling branches, set with very fine smal leaues, bushing and spreading thicke ouer the ground, taking roote still as it spreadeth : the toppes of the branches haue white flowers, with yellow thrummes in the middle, very like vnto the Featherfew, before described, but somewhat greater, not so hard, but more soft and gentle in handling, and the whole herbe to be of a very sweet sent.

1. *Chamæmelum nudum.* Naked Camomill.

We haue another sort of Camomill in some Gardens, but very rare, like vnto the former, but that it is whiter, finer, and smaller, and raiseth it selfe vp a little higher, and beareth naked flowers; that is, without that border of white leaues that is in the former, and consisteth onely of a yellow round thrummie head, smelling almost as sweete as the former.

2. *Chamæmelum flore pleno.* Double flowred Camomill.

The double Camomill groweth with his leaues vpon the ground, as the other single kinde doth, but of a little fresher greene colour, and larger withall: the stalkes with the flowers on them, doe raise themselues vp a little higher then the ordinary, and bearing one or two flowers vpon a stalk, which are composed of many white leaues set together in diuers rowes, which make a fine double flower, with a little yellow spot in the middle for the most part of euery one, and are much larger then any single kinde, smelling better, and more pleasing then the ordinary: this doth creepe vpon the ground as the other, but is more tender to be kept in the Winter. Yet if you saue the flowers hereof (and so will the double Featherfew also) when they haue stood long, and ready to fade, and keepe them dry vntill the Spring, and then breaking them or pulling them to peeces, sowe them, there will spring vp from them Camomill, and also Featherfew, that will againe beare double flowers.

## The Place.

Our ordinary Camomill groweth wilde in many places of our Country, and as well neare London as in other places. The others are onely found in our Gardens, where they are cherished. Bauhinus saith, that the double flowred Camomill is found wilde about Orleance in France.

## The Time.

The double kinde is vsually in flower in Iune, before the ordinary kinde, and most commonly past before it flowreth, which is not vntill Iuly or August. The naked Camomill flowreth betweene them both, or later.

## The Names.

Camomill is called *Anthemis, Leucanthemis*, and *Leucanthemum*, of the whiteneſſe of the flowers; and *Chamæmelum* of the corrupted Italian name *Camomilla*. Some call the naked Camomill, *Chrysanthemum odoratum*. The double Camomill is called by some *Chamæmelum Romanum flore multiplici*.

## The Vertues.

Camomill is put to diuers and sundry vses, both for pleasure and profit, both for inward and outward diseases, both for the sicke and the sound, in bathings to comfort and strengthen the sound, and to ease paines in the diseased, as also in many other formes applyed outwardly. The flowers boyled in Posset drinke prouoketh sweat, and helpeth to expell colds, aches, and other griefes. A Syrupe made of the iuice of the double Camomill, with the flowers and white wine, as Bauhinus saith, is vsed by some against the Iaundise and Dropsie, caused by the euill disposition of the splene.

1 *Parthenium flore pleno.* Double Featherfew. 2 *Chamæmelum nudum.* Naked Camomill. 3 *Chamæmelum flore pleno.* Double Camomill. 4 *Pyrethrum officinarum.* Pelletory of Spaine. 5 *Flos Adonis flore rubro & flore luteo.* Adonis flower both red & yellow. 6 *Helleborus niger ferulaceus siue Buphthalmum.* The great Oxe eye or the great yellow Anemone. 7 *Buphthalmum vulgare.* The common yellow Oxe eye.

## Chap. LIX.

### *Pyrethrum officinarum.* Pelletory of Spaine.

I Must needes adioyne vnto the Camomils this fine and tender plant, for some neare resemblance it hath with them in face, though not in quality. It is a small and lowe plant, bearing many fine greene leaues vpon his slender branches, which leane or lye down vpon the ground, diuided into many parts, yet somewhat larger and broader then Camomill, the stalkes whereof are bigger, and more iuicie then it: the flowers that stand at the toppes of the stalkes are single, but much larger then any Camomill flower, hauing a pale or border of many leaues, white on the vpperside, and reddish vnderneath, set about the yellow middle thrumme; but not standing so close together ioyning at the bottome, as the Camomill flowers doe, but more seuered one from another: it beareth small whitish seede, which is hardly found and discerned from the chaffe: the roote is long, and growing downe right, of the bignesse of a mans finger or thumbe in our Countrey, but not halfe so great where it groweth naturally, with some fibres and branches from the sides thereof, of a very hot, sharpe, and biting taste, drawing much water into the mouth, after it hath been chewed a while: the plant with vs is very tender, and will hardly or not at all endure the hardnesse and extremities of our Winters, vnlesse it be very carefully preserued.

#### The Place.

It groweth in Spaine wilde in many places, and in other hot Countries, where it may feele no frosts to cause it perish.

#### The Time.

It flowreth so late with vs, that it is not vntill August, that oftentimes we cannot gather ripe seedes from it, before it perish.

#### The Names.

The name *Pyrethrum* (taken from πυρ, that is, *ignis*, fire) is giuen to this plant, because of the heate thereof, and that the roote is somewhat like in shew, but specially in property vnto the true *Pyrethrum* of Dioscorides, which is an vmbelliferous plant, whose rootes are greater, and more feruent a great deale, and haue a hayrie bush or toppe as *Meum*, and many other vmbelliferous plants haue. It is also called in Latine, *Salinaris*, of the effect in drawing much moisture into the mouth, to be spit out. We doe vsually call it Pelletory of Spaine.

#### The Vertues.

It is in a manner wholly spent to draw rheume from the teeth, by chewing it in the mouth, thereby to ease the tooth-ach, and likewise from the head, in the paines thereof.

## Chap. LX.

### *Flos Adonis flore rubro.* Red Adonis flower.

ADonis flower may well be accounted a kinde of Camomill, although it hath some especiall differences, hauing many long branches of leaues lying vpon the ground, and some rising vp with the stalke, so finely cut and iagged, that they much resemble the leaues of Mayweed, or of the former *Nigella*: at the top of the stalkes, which rise a foote high or better, stand small red flowers, consisting of six or eight round leaues, hauing a greene head in the middle, set about with many blackish threads, without any smell at all: after the flowers are past, there grow vp heads with many roundish white seedes at the toppes of them, set close together, very like vnto the heads of seede of the great Oxe eye, set downe in the next Chapter, but smaller: the rootes are small and thready, perishing euery yeare, but rising of his owne seede againe, many times before Winter, which will abide vntill the next yeare.

Yellow Adonis flower is like vnto the red, but that the flower is somewhat larger, and of a faire yellow colour. *Flore luteo.*

#### The Place.

The first groweth wilde in the corn fields in many places of our own country, as well as in others, and is brought into Gardens for the beauties sake of the flower. The yellow is a stranger, but noursed in our Gardens with other rarities.

#### The Time.

They flower in May or Iune, as the yeare falleth out to be early or late: the seed is soone ripe after, and will quickly fall away, if it be not gathered.

#### The Names.

Some haue taken the red kinde to be a kinde of Anemone; other to be *Eranthemum* of Dioscorides: the most vsuall name now with vs is *Flos Adonis*, and *Flos Adonidis*: In English, where it groweth wilde, they call it red Maythes, as they call the Mayweede, white Maythes; and some of our English Gentlewomen call it Rosarubie: we vsually call it Adonis flower.

#### The Vertues.

It hath been certainly tryed by experience, that the seed of red Adonis flower drunke in wine, is good to ease the paines of the Collicke and Stone.

## Chap. LXI.

### *Buphthalmum.* Oxe eye.

VNder the name *Buphthalmum*, or Oxe eye, are comprehended two or three seuerall plants, each differing from other, both in face and property, yet because they all beare one generall name, I thinke fittest to comprise them all in one Chapter, and first of that which in leafe & seed commeth neerest to the Adonis flower.

#### 1. *Buphthalmum maius siue Helleborus niger ferulaceus.*
Great Oxe eye, or the yellow Anemone.

This great Oxe eye is a beautifull plant, hauing many branches of greene leaues

leaning or lying vpon the ground for the most part, yet some standing vpright, which are as fine, but shorter then Fenell; some of them ending in a small tuft of green leaues, and some hauing at the toppes of them one large flower apeece, somewhat reddish or browniſh on the outside, while they are in bud, and a while after, and being open, shew themselues to consist of twelue or fourteene long leaues, of a faire shining yellow colour, set in order round about a greene head, with yellow thrums in the middle, laying themselues open in the sunne, or a faire day, but else remaining close: after the flower is past, the head growing greater, sheweth it selfe compact of many round whitish seede, very like vnto the head of seede of the Adonis flower last described, but much greater: the rootes are many long blackish fibres or strings, set together at the head, very like vnto the rootes of the lesser blacke Hellebor or Bearefoote, but somewhat harder, stiffer, or more brittle, and seeming without moisture in them, which abide and encrease euery yeare.

2. *Buphthalmum minus, seu Anthemis flore luteo.* Small Oxe eye.

This plant might seeme to be referred to the Camomils, but that it is not sweete, or to the Corne-Marigolds, but that the stalkes and leaues are not edible: it is therefore put vnder the Oxe eyes, and so we will describe it; hauing many weake branches lying vpon the ground, beset with winged leaues, very finely cut and iagged, somewhat like vnto Mayweede, but a little larger: the flowers are like vnto the Corne Marigold, and larger then any Camomill, being wholly yellow, as well the pale or border of leaues, as the middle thrummes: the rootes are somewhat tough and long.

3. *Buphthalmum vulgare.* Common Oxe eye.

This Oxe eye riseth vp with hard round stalkes, a foote and a halfe high, hauing many winged leaues vpon them, made of diuers long and something broad leaues, snipt about the edges, set together somewhat like vnto Tansie, but smaller, and not so much winged: the flowers stand at the toppes of the stalkes, of a full yellow colour, both the outer leaues and the middle thrum, and not altogether so large as the last: the rootes of this kinde perish euery yeare, and require a new sowing againe.

## The Place.

The first groweth in diuers places of Austria, Bohemia, and those parts, it hath beene likewise brought out of Spaine. The second in Prouence, a country in France. The last in diuers places, as well of Austria as Morauia, and about Mentz and Norimberg, as Clusius setteth downe. We haue them in our Gardens, but the first is of the greatest respect and beauty.

## The Time.

The first flowreth betimes, oftentimes in March, or at the furthest in Apill; the seede is ripe in May, and must be quickly gathered, left it bee lost. The other two flower not vntill Iune.

## The Names.

The first is called *Buphthalmum* of Dodonæus, *Pseudohelleborus* of Matthiolus, *Helleborus niger ferulaceus Theophrasti* by Lobel, of some others *Ellebor us niger verus*, vsing it for the true blacke Ellebor, but it is much differing, as well in face as properties. Of others *Sesamoides minus*. Some haue thought it to be a yellow Anemone, that haue looked on it without further iudgement, and by that name is most vsually knowne to most of our English Gentlewomen that know it. But it may most fitly be called a *Buphthalmum*, as Dodonæus doth, and *Hispanicum* or *Austriacum*, for distinctions sake. We doe most vsually call it *Helleborus niger ferulaceus*, as Lobel doth: Bauhinus calleth

calleth it *Helleborus niger tenuifolius Buphthalmi flore.* The second is called *Buphthalmum Narbonense*: In English, The French, or lesser Oxe eye, as the first is called, The great Oxe eye. The last, The common Oxe eye.

### The Vertues.

The first hath been vsed in diuers places for the true blacke Ellebor, but now is sufficiently knowne to haue been an errour; but what Physicall property it hath, other then Matthiolus hath expressed, to be vsed as Setterwort for cattell, when they rowell them, to put or draw the rootes hereof through the hole they make in the dewe lappe, or other places, for their coughes or other diseases, I know not, or haue heard or read of any. The others likewise haue little or no vse in Physicke now a dayes that I know.

## Chap. LXII.

### *Chrysanthemum.* Corne Marigold.

Although the sorts of Corne Marigolds, which are many, are fitter for another then this worke, and for a Catholicke Garden of Simples, then this of Pleasure and Delight for faire Flowers; yet giue me leaue to bring in a couple: the one for a corner or by-place, the other for your choisest, or vnder a defenced wall, in regard of his statelinesse.

1. *Chrysanthemum Creticum.* Corne Marigold of Candy.

This faire Corne Marigold hath for the most part one vpright stalke, two foote high, whereon are set many winged leaues, at euery ioynt one, diuided and cut into diuers parts, and they againe parted into seuerall peeces or leaues: the flowers growe at the toppes of the stalkes, rising out of a scaly head, composed of ten or twelue large leaues, of a faire, but pale yellow colour, and more pale almost white at the bottome of the leaues, round about the yellow thrumme in the middle, being both larger and sweeter then any of the other Corne Marigolds: the seede is whitish and chaffie: the roote perisheth euery yeare.

2. *Chrysanthemum Perüuianum, sive Flos Solis.*
The golden flower of Peru, or the Flower of the Sunne.

This goodly and stately plant, wherewith euery one is now adayes familiar, being of many sorts, both higher and lower (with one stalke, without branches, or with many branches, with a blacke, or with a white seede, yet differing not in forme of leaues or flowers one from another, but in the greatnesse or smalnesse) riseth vp at the first like vnto a Pompion with two leaues, and after two, or foure more leaues are come forth, it riseth vp into a great stalke, bearing the leaues on it at seuerall distances on all sides thereof, one aboue another vnto the very toppe, being sometimes, and in some places, seuen, eight, or ten foote high, which leaues standing out from the stemme or stalke vpon their seuerall great ribbed foote-stalkes, are very large, broad belowe, and pointed at the end, round, hard, rough, of a sad greene colour, and bending downewards: at the toppe of the stalke standeth one great, large, and broad flower, bowing downe the head vnto the Sunne, and breaking forth from a great head, made of scaly greene leaues, like vnto a great single Marigold, hauing a border of manie long yellow leaues, set about a great round yellow thrumme, as it were in the middle, which are very like vnto short heads of flowers, vnder euery one whereof there is a seede, larger then any seede of the Thistles, yet somewhat like, and lesser, and rounder then any Gourd seede, set in so close and curious a manner, that when the seede is taken out, the head with the hollow places or cels thereof, seemeth very like vnto an hony combe; which seede is in some plants very blacke, in the hotter countries, or very white,

white, and great, or large, but with vs is neither so large, blacke, or white; but sometimes blackish or grayish. Some sort riseth not vp halfe the height that others doe, and some againe beare but one stemme or stalke, with a flower at the toppe thereof; and others two or three, or more small branches, with euery one his flower at the end; and some so full of branches from the very ground almost, that I haue accounted threescore branches round about the middle stalke of one plant, the lowest neare two yards long, others aboue them a yard and a halfe, or a yard long, with euery one his flower thereon; but all smaller then those that beare but one or two flowers, and lesser also for the most part then the flower on the middle stalke it selfe. The whole plant, and euery part thereof aboue ground hath a strong resinous sent of Turpentine, and the heads and middle parts of the flowers doe oftentimes (and sometimes the ioynts of the stalke where the leaues stand) sweat out a most fine thin & cleare Rosin or Turpentine, but in small quantity, and as it were in drops, in the heate and dry time of the year, so like both in colour, smell, and taste vnto cleare Venice Turpentine, that it cannot be knowne from it: the roote is strongly fastened in the ground by some greater roots branching out, and a number of small strings, which growe not deepe, but keepe vnder the vpper crust of the earth, and desireth much moisture, yet dyeth euery yeare with the first frosts, and must be new sowne in the beginning of the Spring.

### The Place.

Their places are set downe in their titles, the one to come out of Candy, the other out of Peru, a Prouince in the West Indies.

### The Time.

The first flowreth in Iune, the other later, as not vntill August, and sometimes so late, that the early frosts taking it, neuer suffer it to come to ripenes.

### The Names.

The first hath his name in his title. The second, besides the names set downe, is called of some *Planta maxima*, *Flos maximus*, *Sol Indianus*, but the most vsuall with vs is, *Flos Solis*: In English, The Sunne Flower, or Flower of the Sunne.

### The Vertues.

There is no vse of either in Physicke with vs, but that sometimes the heads of the Sunne Flower are dressed, and eaten as Hartichokes are, and are accounted of some to be good meate, but they are too strong for my taste.

## Chap. LXIII.

### *Calendula.* Marigolds.

Some haue reckoned vp many sorts of Marigolds, I had rather make but two, the single and the double; for doubtlesse, those that be most double, rise from the best seede, which are the middlemost of the great double, and some will be lesse double, whose seede is greater then the rest, according to the ground where it groweth; as also those that be of a paler colour, doe come of the seed of the yellower sort.

1. *Calendula maxima.* The great Garden Marigold.

The Garden Marigold hath round greene stalkes, branching out from the ground into many parts, whereon are set long flat greene leaues, broader and rounder at the

1 *Chrysanthemum Creticum.* Corne Marigolds of Candy. 2 *Flos Solis.* The Flower of the Sunne. 3 *Calendula.* Marigolds. 4 *Aster Atticus sive Italorum.* The purple Marigold. 5 *Pilosella maior,* Golden Mouse-eare. 6 *Scorsonera Hispanica.* Spanish Vipers grasse. 7 *Tragopogon.* Goates beard, or goe to bed at noone.

point then any where else, and smaller also at the setting to of the stalke, where it compasseth it about: the flowers are sometimes very thicke and double (breaking out of a scaly clammy greene head) composed of many rowes of leaues, set so close together one within another, that no middle thrume can bee seene, and sometimes lesse double, hauing a small browne spot of a thrume in the middle: and sometimes but of two or three rowes of leaues, with a large browne thrume in the middle; euery one whereof is somewhat broader at the point, and nicked into two or three corners, of an excellent faire deepe gold yellow colour in some, and paler in others, and of a pretty strong and resinous sweete sent: after the flowers are past, there succeede heads of crooked seede, turning inward, the outermost biggest, and the innermost least: the roote is white, and spreadeth in the ground, and in some places will abide after the seeding, but for the most part perisheth, and riseth againe of his owne seede. Sometimes this Marigold doth degenerate, and beareth many small flowers vpon short stalkes, compassing the middle flower: but this happeneth but seldome, and therefore accounted but *lusus naturæ*, a play of nature, which she worketh in diuers other plants besides.

2. *Calendula simplex.* The single Marigold.

There is no difference betweene this and the former, but that the flowers are single, consisting of one rowe of leaues, of the same colour; eyther paler or deeper yellow, standing about a great browne thrumme in the middle: the seed likewise is alike, but for the most part greater then in the double kindes.

## The Place.

Our Gardens are the chiefe places for the double flowers to grow in; for we know not of any other naturall place: but the single kinde hath beene found wilde in Spaine, from whence I receiued seede, gathered by Guillaume Boel, in his time a very curious, and cunning searcher of simples.

## The Time.

They flower all the Summer long, and sometimes euen in winter, if it be milde, and chiefly at the beginning of those monethes, as it is thought.

## The Names.

They are called *Caltha* of diuers, and taken to bethat *Caltha*, wherof both Virgil and Columella haue written. Others doe call them *Calendula*, of the Kalendes, that is the first day of the monthes, wherein they are thought chiefly to flower; and thereupon the Italians call them, *Fiori di ogni mese*, that is, The Flowers of euery moneth: We cal them in English generally, eyther Golds, or Marigolds.

## The Vertues.

The herbe and flowers are of great vse with vs among other pot-herbes; and the flowers eyther greene or dryed, are often vsed in posiets, broths, and drinkes, as a comforter of the heart and spirits, and to expel any malignant or pestilential quality, gathered neere thereunto. The Syrupe and Conserue made of the fresh flowers, are vsed for the same purposes to good effect.

## Chap. LXIIII.

### *Aster.* Starre-wort.

Dioscorides and other of the ancient Writers, haue set forth but one kinde of Starre-wort, which they call *Aster Atticus*, of the place no doubt, where the greatest plentie was found, which was the Countrey of Athens: the later Writers haue found out many other plants which they referre to this kinde, calling them by the same name. It is not my purpose to entreate of them all, neyther doth this garden fitly agree with them: I shall therefore select out one or two from the rest, and giue you the knowledge of them, leauing the rest to their proper place.

### 1. *Aster Atticus flore luteo.* Yellow Starre-wort.

This Starre-wort riseth vp with two or three rough hairy stalkes, a foote and a halfe high, with long, rough or hairie, browniſh, darke greene leaues on them, diuided into two or three branches: at the toppe of euery one whereof standeth a flat scaly head, compassed vnderneath with fiue or sixe long, browne, rough greene leaues, standing like a Starre, the flower it selfe standing in the middle thereof, made as a border of narrow, long, pale yellow leaues, set with a browniſh yellow thrume: the roote dyeth euery yeare, hauing giuen his flower.

### 2. *Aster Atticus Italorum flore purpureo.* Purple Italian Starre-wort.

This Italian Starre-wort hath many wooddy, round brittle stalkes, rising from the roote, somewhat higher then the former, sometimes standing vpright, and otherwhiles leaning downewards, whereon are set many somewhat hard, and rough long leaues, round pointed, without order vp to the toppe, where it is diuided into seuerall branches, whereon stand the flowers, made like vnto a single Marigold, with a border of blewish purple leaues, set about a browne middle thrume, the heads sustaining the flowers, are composed of diuers scaly greene leaues, as is to be seene in the Knapweedes or Matfelons, which after the flowers are past yeelde a certaine downe, wherein lye small blacke and flat seedes, somewhat like vnto Lettice seede, which are carried away with the winde: the roote is composed of many white strings, which perisheth not as the former, but abideth, and springeth afresh euery yeare.

### The Place.

The first is found in Spaine, as Clusius, and in France, as Lobel say. The other hath beene found in many places in Germany, and Austria: in Italie also, and other places; we haue it plentifully in our Gardens.

### The Time.

The first flowreth in Summer. And the other not vntill August or September.

### The Names.

The first is called *Aster Atticus flore luteo, Bubonium, & Inguinalis*, and of many is taken to be the true *Aster Atticus* of Dioscorides: yet Matthiolus thinketh not so, for diuers good reasons, which hee setteth downe in the Chapter of *Aster Atticus*, as any man may vnderstand, if they will but reade the place, which is too long to bee inserted here. The other is thought by Matthiolus, to bee the truer *Aster Atticus*, (vnto whom I must also consent) and constantly also affirmed to be the *Amellus Virgilij*, as may be seene in the same place: but it is vsually called at this day, *Aster Italorum flore cæruleo* or *purpureo*,

*purpureo.* Their Englisſh names are ſufficiently expreſſed in their titles, yet ſome call the laſt, The purple Marigold, becauſe it is ſo like vnto one in form.

### The Vertues.

They are held, if they bee the right, to bee good for the biting of a mad dogge, the greene herbe being beaten with old hogs greaſe, and applyed; as alſo for ſwolne throats : It is likewiſe vſed for botches that happen in the groine, as the name doth import.

## Chap. LXV.

*Piloſella maior.* Golden Mouſe-eare.

SOme reſemblance that the flowers of this plant hath with the former Golds, maketh me to inſert it in this place, although I know it agreeth not in any other part, yet for the pleaſant aſpect thereof, it muſt bee in this my garden, whoſe deſcription is as followeth: It hath many broade greene leaues ſpread vpon the ground, ſpotted with pale ſpots, yet more conſpicuous at ſometimes then at other; ſomewhat hairy both on the vpper and vnderſide, in the middle of theſe leaues riſe vp one, two or more blackiſh hairy ſtalkes, two foote high at the leaſt, bare or naked vp to the top, where it beareth an vmbell, or ſhort tuft of flowers, ſet cloſe together vpon ſhort ſtalkes, of the forme or faſhion of the Haukeweedes, or common Mouſe-eare, but ſomewhat ſmaller, of a deep gold yellow, or orenge tawney colour, with ſome yellow threds in the middle, of little or no ſent at all: after the flowers are paſt, the heads carry ſmall, ſhort, blacke ſeede, with a light downie matter on them, ready to bee carried away with the winde, as many other plants are, when they be ripe : the rootes ſpread vnder ground, and ſhoote vp in diuers other places, whereby it much encreaſeth, eſpecially if it be ſet in any moiſt or ſhadowie place.

### The Place.

It groweth in the ſhadowie woods of France, by Lions, and Mompelier, as Lobell teſtifieth: we keepe it in our gardens, and rather in a ſhadowie then ſunnie place.

### The Time.

It flowreth in Somer, and ſometimes againe in September.

### The Names.

It is called by Lobell, *Pulmonaria Gallorum Hieratÿ facie :* and the Herbariſts of France take it to be the true *Pulmonaria* of Tragus. Others call it *Hieratium flore aureo.* Pelleterius *Hieratium Indicum.* Some *Piloſella*, or *Auricula muris maior flore aureo.* And ſome *Chondrilla flore aureo.* Dalechamptus would haue it to bee *Corchorus*, but farre vnfitly. The fitteſt Engliſh name we can giue it, is Golden Mouſe-eare, which may endure vntill a fitter bee impoſed on it : for the name of Grim the Collier, whereby it is called of many, is both idle and fooliſh.

### The Vertues.

The French according to the name vſe it for the defects of the lunges, but with what good ſucceſſe I know not.

## Chap. LXVI.

### *Scorsonera.* Vipers grasse.

ALthough there be foure or fiue sorts of *Scorsonera*, yet I shall here desire you to be content with the knowledge only of a couple.

1. *Scorsonera Hispanica maior.* The greater Spanish Vipers grasse.

This Spanish Vipers grasse hath diuers long, and somewhat broad leaues, hard and crumpled on the edges, and sometimes vneuenly cut in or indented also, of a blewish greene colour: among which riseth vp one stalke, and no more for the most part, two foote high or thereabouts, hauing here and there some narrower long leaues thereon then those below: the toppe of the stalke brancheth it selfe forth into other parts, euery one bearing a long scaly head, from out of the toppe whereof riseth a faire large double flower, of a pale yellow colour, much like vnto the flower of yellow Goates beard, but a little lesser, which being past, the seede succeedeth, being long, whitish and rough, inclosed with much downe, and among them many other long smooth seedes, which are limber and idle, and are carryed away at the will of the winde: the roote is long, thicke and round, brittle and blacke, with a certaine roughnesse on the outside: but very white within, yeelding a milkie liquor being broken, as euery other part of the plant doth besides, yet the roote more then any other part, and abideth many yeares without perishing.

2. *Scorsonera Pannonica purpurea.* Purple flowred Vipers grasse.

This purple flowred Vipers grasse hath long and narrow leaues, of the same blewish greene colour with the former: the stalke riseth vp a foote and a halfe high, with a few such like leaues, but shorter thereon, breaking at the toppe into two or three parts, bearing on each of them one flower, fashioned like the former, and standing in the like scaly knoppe or head, but of a blewish purple colour, not fully so large, of the sweetest sent of any of this kinde, comming neerest vnto the smell of a delicate perfume.

#### The Place.

The first is of Spaine. The other of Hungarie and Austrich: which now furnish our gardens.

#### The Time.

They flower in the beginning of May: the seede is soone ripe after, and then perishing downe to the roote for that yeare, springeth afresh before Winter againe.

#### The Names.

They are called after the Spanish name *Scorsonera*, which is in Latine *Viperaria*, of some *Viperina*, and *Serpentina*: Wee call them in English Vipers grasse, or *Scorsonera*.

#### The Vertues.

Manardus as I thinke first wrote hereof, and saith that it hath been found to cure them that are bitten of a Viper, or other such like venemous Creature. The rootes hereof being preserued with sugar, as I haue done often, doe eate almost as delicate as the Eringus roote, and no doubt is good to comforr and strengthen the heart, and vitall spirits. Some that haue vsed the preserued roote haue found it effectuall to expelling winde out of the stomacke, and to helpe swounings and faintnesse of the heart.

## Chap. LXVII.

### *Tragopogon.* Goates beard.

I Must in this place set downe but two sorts of Goates beards; the one blew or ash-colour, the other red or purple, and leaue the other kindes: some to bee spoken of in the Kitchin Garden, and others in a Physicall Garden.

#### 1. *Tragopogon flore caruleo.* Blew Goates beard.

All the Goates beards haue long, narrow, and somewhat hollow whitish greene leaues, with a white line downe the middle of euery one on the vpperside: the stalke riseth vp greater and stronger then the Vipers grasse, bearing at the toppe a great long head or huske, composed of nine or ten long narrow leaues, the sharpe points or ends whereof rise vp aboue the flower in the middle, which is thicke and double, somewhat broad and large spread, of a blewish ash-colour, with some whitish threads among them, shutting or closing it selfe within the greene huske euery day, that it abideth blowing, vntill about noone, and opening not it selfe againe vntill the next morning: the head or huske, after the flower is past, and the seede neare ripe, openeth it selfe; the long leaues thereof, which closed not before now, falling downe round about the stalke, and shewing the seede, standing at the first close together, and the doune at the toppe of them: but after they haue stood a while, it spreadeth it selfe round, and is ready to be carried away with the winde, if it be not gathered: the seede it selfe is long, round, and rough, like the seede of the Vipers grasse, but greater and blacker: the roote is long, and not very great, but perisheth as soone as it hath borne seede, and springeth of the fallen seede, that yeare remaining greene all Winter, and flowring the next yeare following: the whole yeeldeth milke as the former, but somewhat more bitter and binding.

#### 2. *Tragopogon purpureum.* Purple Goates beard.

There is little difference in this kind from the former, but that it is a little larger, both in the leafe, and head that beareth the seed: the flowers also are a little larger, and spread more, of a darke reddish purple colour, with some yellow dust as it were cast vpon it, especially about the ends: the roote perisheth in the like manner as the other.

### The Place.

Both these haue been sent vs from the parts beyond the Seas, I haue had them from Italy, where no doubt they grow naturally wilde, as the yellow doth with vs: they are kept in our Gardens for their pleasant flowers.

### The Time.

They flower in May and Iune: the seede is ripe in Iuly.

### The Names.

Their generall name is after the Greeke word *Tragopogon*, which is in Latine, *Barba hirci*: In English, Goates beard; the head of seede when it is readie to bee carried away with the winde, causing that name for the resemblance: and because the flower doth euery day close it selfe at noone (as I said before) and openeth not againe vntill the next Sunne, some haue fitly called it, Goe to bed at noone.

### The Vertues.

The rootes of these kindes are a little more bitter and more binding also then

then the yellow kinde expreſſed in the Kitchin Garden; and therefore fitter for medicine then for meate, but yet is vſed as the yellow kinde is, which is more fit for meate then medicine. The diſtilled water is good to waſh old ſores and wounds.

## CHAP. LXVIII.

### *Flos Africanus.* The French Marigold.

OF the French or African Marigolds there are three kindes as principall, and of each of them both with ſingle and double flowers: of theſe, ſome diuerſity is obſerued in the colour of the flowers, as well as in the forme or largeneſſe, ſo that as you may here ſee, I haue expreſſed eight differences, and Fabius Columna nine or ten, in regard hee maketh a diuerſity of the paler and deeper yellow colour: and although the leſſer kinde, becauſe of its euill ſent, is held dangerous, yet for the beauty of the flower it findeth roome in Gardens.

### 1. *Flos Africanus maior ſiue maximus multiplex.* The great double French Marigold.

This goodly double flower, which is the grace and glory of a Garden in the time of his beauty, riſeth vp with a ſtraight and hard round greene ſtalke, hauing ſome creſts or edges all along the ſtalke, beſet with long winged leaues, euery one whereof is like vnto the leafe of an Aſh, being compoſed of many long and narrow leaues, ſnipt about the edges, ſtanding by couples one againſt another, with an odde one at the end, of a darke or full greene colour: the ſtalke riſeth to be three or foure foote high, and diuideth it ſelfe from the middle thereof into many branches, ſet with ſuch like leaues to the toppes of them, euery one bearing one great double flower, of a gold yellow colour aboue, and paler vnderneath, yet ſome are of a pale yellow, and ſome betweene both, and all theſe riſing from one and the ſame ſeede: the flower, before it be blowne open, hath all the leaues hollow; but when it is full blowne open, it ſpreadeth it ſelfe larger then any Prouince Roſe, or equall vnto it at the leaſt, if it be in good earth, and riſeth out of a long greene huske, ſtriped or furrowed, wherein after the flower is paſt, (which ſtandeth in his full beauty a moneth, and oftentimes more, and being gathered, may be preſerued in his full beauty for two moneths after, if it be ſet in water) ſtandeth the ſeede, ſet thicke and cloſe together vpright, which is blacke, ſomewhat flat and long: the roote is full of ſmall ſtrings, whereby it ſtrongly comprehendeth in the ground: the flower of this, as well as the ſingle, is of the very ſmell of new waxe, or of an honie combe, and not of that poiſonfull ſent of the ſmaller kindes.

### 2. *Flos Africanus maior ſimplex.* The great ſingle French Marigold.

This ſingle Marigold is in all things ſo like vnto the former, that it is hard to diſcerne it from the double, but by the flowers, onely the ſtalke will be browner then the double; and to my beſt obſeruation, hath and doth euery yeare riſe from the ſeede of the double flower: ſo that when they are in flower, you may ſee the difference (or not much before, when they are in bud) this ſingle flower euer appearing with thrums in the middle, and the leaues, which are the border or pale ſtanding about them, ſhewing hollow or fiſtulous, which after lay themſelues flat and open (and the double flower appearing with all his leaues folded cloſe together, without any thrum at all) and are of a deeper or paler colour, as in the double.

### 3. *Flos Africanus fiſtuloſo flore ſimplex & multiplex.* Single and double French Marigolds with hollow leafed flowers.

As the former two greateſt ſorts haue riſen from the ſeede of one and the ſame (I

meane the pod of double flowers) so doe these also, not differing from it in any thing, but that they are lower, and haue smaller greene leaues, and that the flower also being smaller, hath euery leafe abiding hollow, like vnto an hollow pipe, broad open at the mouth, and is of as deepe a yellow colour for the most part as the deepest of the former, yet sometimes pale also.

### 4. *Flos Africanus minor multiplex.* The lesser double French Marigold.

The lesser double French Marigold hath his leaues in all things like vnto the former, but somewhat lesser, which are set vpon round browne stalkes, not so stiffe or vpright, but bowing and bending diuers wayes, and sometimes leaning or lying vpon the ground: the stalkes are branched out diuersly, whereon are set very faire double flowers like the former, and in the like greene huskes, but smaller, and in some the outermost leaues will be larger then any of the rest, and of a deeper Orenge colour, almost crimson, the innermost being of a deepe gold yellow colour, tending to crimson: the whole flower is smaller, and of a stronger and more vnpleasant sauour, so that but for the beautifull colour, and doubleneſſe of the flower, pleasant to the eye, and not to any other sense, this kinde would finde roome in few Gardens: the rootes and seedes are like the former, but lesser.

### 5. *Flos Africanus minor simplex.* The small single French Marigold.

This single kinde doth follow after the last in all manner of proportion, both of stalkes, leaues, seedes, and rootes: the flowers onely of this are single, hauing fiue or six broad leaues, of a deepe yellow crimson colour, with deepe yellow thrummes in the middle, and of as strong a stinking sent, or more then the last.

#### The Place.

They growe naturally in Africa, and especially in the parts about Tunis, and where old Carthage stood, from whence long agoe they were brought into Europe, where they are onely kept in Gardens, being sowne for the most part euery yeare, vnlesse in some milde Winters. The last single and double kindes (as being more hardy) haue sometimes endured: but that kinde with hollow leafed flowers, as Fabius Columna setteth it downe, is accounted to come from Mexico in America.

#### The Time.

They flower not vntill the end of Summer, especially the greater kindes: but the lesser, if they abide all the Winter, doe flower more early.

#### The Names.

They haue been diuersly named by diuers men: Some calling them *Caryophyllus Indicus*, that is, Indian Gilloflowers, and *Tanacetum Peruianum*, Tansie of Peru, as if it grew in Peru, a Prouince of America; and *Flos Indicus*, as a flower of the Indies; but it hath not beene knowne to haue beene brought from thence. Others would haue it to be *Othonna* of Plinie, and others; some to be *Lycopersicum* of Galen. It is called, and that more truely, *Flos Tunetensis*, *Flos Africanus*, and *Caltha Africana*, that is, the flower of Tunis, the flower of Africa, the Marigold of Africa, and peraduenture *Pedna Panerum*. We in English most vsually call them, French Marigolds, with their seuerall distinctions of greater or smaller, double or single. To that with hollow leafed flowers, Fabius Columna giueth the name of *Fistulose flore*, and I so continue it.

The

1 *Flos Africanus maximus multiplex.* The greateſt double French Marigold. 2 *Flos Africanus maior multiplex.* The greater double French Marigold
3 *Flos Africanus maximus ſimplex* The greateſt ſingle French Marigold. 4 *Flos Africanus multiplex fiſtuloſus.* The doule hollow French Marigold.
5 *Flos Africanus ſimplex fiſtuloſus.* The ſingle hollow French Marigold. 6 *Flos Africanus minor multiplex.* Theſmaller double French Marigold. 7 *Flos Africanus minor multiplex alter.* Another ſort of the leſſer double French Marigold. 8 *Flos Africanus minor ſimplex.* The leſſer ſingle French Marigold.

The Vertues.

We know no vſe they haue in Phyſicke, but are cheriſhed in Gardens for their beautifull flowers ſake.

## Chap. LXIX.

*Caryophyllus hortenſis.* Carnations and Gilloflowers.

TO auoide confuſion, I muſt diuide Gilloflowers from Pinkes, and intreate of them in ſeuerall Chapters. Of thoſe that are called Carnations or Gilloflowers, as of the greater kinde, in this Chapter; and of Pinkes, as well double as ſingle, in the next. But the number of them is ſo great, that to giue ſeuerall deſcriptions to them all were endleſſe, at the leaſt needleſſe: I will therefore ſet downe onely the deſcriptions of three (for vnto theſe three may be referred all the other ſorts) for their faſhion and manner of growing, and giue you the ſeuerall names (as they are vſually called with vs) of the reſt, with their variety and mixture of colours in the flowers, wherein conſiſteth a chiefe difference. I account thoſe that are called Carnations to be the greateſt, both for leafe and flower, and Gilloflowers for the moſt part to bee leſſer in both; and therefore will giue you each deſcription apart, and the Orenge tawnie or yellow Gilloflower likewiſe by it ſelfe, as differing very notably from all the reſt.

1. *Caryophyllus maximus Harwicenſis ſiue Anglicus.*
The great Harwich or old Engliſh Carnation.

I take this goodly great old Engliſh Carnation, as a preſident for the deſcription of all the reſt of the greateſt ſorts, which for his beauty and ſtatelineſſe is worthy of a prime place, hauing beene alwayes very hardly preſerued in the Winter; and therefore not ſo frequent as the other Carnations or Gilloflowers. It riſeth vp with a great thicke round ſtalke, diuided into ſeuerall branches, ſomewhat thickly ſet with ioynts, and at euery ioynt two long greene rather then whitiſh leaues, ſomewhat broader then Gilloflower leaues, turning or winding two or three times round (in ſome other ſorts of Carnations they are plaine, but bending the points downewards, and in ſome alſo of a darke reddiſh greene colour, and in others not ſo darke, but rather of a whitiſh greene colour:) the flowers ſtand at the toppes of the ſtalkes in long, great, and round greene huskes, which are diuided into fiue points, out of which riſe many long and broad pointed leaues, deeply iagged at the ends, ſet in order round and comely, making a gallant great double flower, of a deepe Carnation colour, almoſt red, ſpotted with many bluſh ſpots and ſtrakes, ſome greater and ſome leſſer, of an excellent ſoft ſweete ſent, neither too quicke as many others of theſe kinds are, nor yet too dull, and with two whitiſh crooked threads like hornes in the middle: this kinde neuer beareth many flowers, but as it is ſlow in growing, ſo in bearing, not to be often handled, which ſheweth a kinde of ſtatelineſſe, fit to preſerue the opinion of magnificence: the roote is branched into diuers great, long, wooddy rootes, with many ſmall fibres annexed vnto them.

2. *Caryophyllus hortenſis flore pleno rubro.* The red or Cloue Gilloflower.

The red Cloue Gilloflower, which I take as a preſident for the ſecond ſort, which are Gilloflowers, grow like vnto the Carnations, but not ſo thicke ſet with ioynts and leaues: the ſtalkes are more, the leaues are narrower and whiter for the moſt part, and in ſome doe as well a little turne: the flowers are ſmaller, yet very thicke and double in moſt, and the greene huskes wherein they ſtand are ſmaller likewiſe then the former: the ends of the leaues in this flower, as in all the reſt, are dented or iagged, yet in ſome more then in others; ſome alſo hauing two ſmall white threads, crooked at the ends like hornes, in the middle of the flower, when as diuers other haue none. Theſe

kindes,

1 *Caryophyllus maximus rubro varius.* The great old Carnation or gray Hulo. 2 *Caryophyllus maior rubro & albo varius.* The white Carnation. 3 *Caryophyllus albo rubens.* The Camberſine or the Pooleflower. 4 *Caryophyllus Cantij ſtriatus.* The faire made of Kent. 5 *Caryophyllus Sabaudicus carneus.* The blush Sauadge. 6 *Caryophyllus Xerampelinus.* The Gredeline Carnation. 7 *Caryophyllus dictus Grimelo.* The Grimelo or Prince. 8 *Caryophyllus albus maior,* The great white Gilloflower. 9 *Elegans Heroina Bradſhawij.* Maſter Bradſhawes dainty Lady.

kindes, and especially this that hath a deepe red crimson coloured flower, doe endure the cold of our winters, and with lesse care is preserued : these sorts as well as the former doe very seldome giue any seede, as far as I could euer obserue or learne.

3. *Caryophyllus Silesiacus flore pleno miniato.*
The yellow or Orenge tawny Gilloflower.

This Gilloflower hath his stalkes next vnto the ground, thicker set, and with smaller or narrower leaues then the former for the most part : the flowers are like vnto the Cloue Gilloflowers, and about the same bignesse and doublenesse most vsually, yet in some much greater then in others ; but of a pale yellowish Carnation colour, tending to an Orenge, with two small white threds, crooked at the ends in the middle, yet some haue none, of a weaker sent then the Cloue Gilloflower : this kinde is more apt to beare seede then any other, which is small, black, flat, and long, and being sowen, yeelde wonderfull varieties both of single and double flowers : some being of a lighter or deeper colour then the mother plants : some with stripes in most of the leaues : Others are striped or spotted, like a speckled Carnation or Gilloflower, in diuers sorts, both single and double : Some againe are wholly of the same colour, like the mother plant, and are eyther more or lesse double then it, or else are single with one row of leaues, like vnto a Pinck ; and some of these likewise eyther wholly of a crimson red, deeper or lighter, or variably spotted, double or single as a Pinck, or bluish eyther single or double, and but very seldome white : yet all of them in their greene leaues little or nothing varying or differing.

*Cariophylli maximi.*

## CARNATIONS.

*Caryophyllus maximus dictus Hulo rubro-variui.*

THe gray *Hulo* hath as large leaues as the former old Carnation, and as deepely iagged on the edges : it hath a great high stalke, whereon stand the flowers, of a deepe red colour, striped and speckled very close together with a darkish white colour.

*Caryophyllus maximus dictus Hulo ruber non variatus.*

The red *Hulo* is also a faire great flower, of a stamell colour, deeply iagged as the former, and groweth very comely without any spot at all in it, so that it seemeth to bee but a stamell Gilloflower, saue that it is much greater.

*Caryophyllus maximus dictus Hulo caruleo purpureus.*

The blew *Hulo* is a goodly faire flower, being of a faire purplish murrey colour, curiously marbled with white, but so smally to be discerned, that it seemeth only purple, it hath so much the Mastrie in it ; it resembleth the Brassill, but that it is much bigger.

*Caryophyllus maximus dictus Grimelo siue Princeps.*

The *Grimelo* or Prince is a faire flower also, as large as any Chrystall or larger, being of a faire crimson colour, equally for the most part striped with white, or rather more white then red, thorough euery leafe from the bottome, and standeth comely.

*Caryophyllus maximus Incarnadinus albus.*

The white Carnation or Delicate, is a goodly delightfull fair flower in his pride and perfection, that is, when it is both marbled and flaked, or striped and speckled with white vpon an incarnate crimson colour, beeing a very comely flower, but abideth not constant, changing oftentimes to haue no flakes or strakes of white, but marbled or speckled wholly.

*Caryophyllus maximus Incarnadinus Gallicus.*

The French Carnation is very like vnto the white Carnation, but that it hath more specks, and fewer stripes or flakes of white in the red, which hath the mastrie of the white.

*Caryophyllus maximus Incarnadinus grandis.*

The ground Carnation (if it be not the same with the graund or great old Carnation first set downe, as the alteration but of one letter giueth the coniecture) is a thicke flower, but spreadeth

not

not his leaues abroade as others doe, hauing the middle standing higher then the outer leaues, and turning vp their brimmes or edges; it is a sad flower, with few stripes or spots in it: it is very subiect to breake the pod, that the flower seldome commeth faire and right; the greene leaues are as great as the *Hulo* or Lombard red.

*Caryophyllus maximus Chrystallinus.*

The Chrystall or Chrystalline (for they are both one, howsoeuer some would make them differ) is a very delicate flower when it is well marked, but it is inconstant in the markes, being sometimes more striped with white and crimson red, and sometimes lesse or little or nothing at all, and changing also sometimes to be wholly red, or wholly blush.

*Caryophyllus maximus flore rubro.*

The red Chrystall, which is the red hereof changed, is the most orient flower of all other red Gilloflowers, because it is both the greatest, as comming from the Chrystall, as also that the red hereof is a most excellent crimson.

*Caryophyllus maximus dictus Fragrans.*

The Fragrant is a faire flower, and thought to come from the Chrystall, being as large, but of a blush red colour, spotted with small speckes, no bigger then pinnes points, but not so thicke as in the Pageant.

*Caryophyllus maximus Sabaudicus varius.*

The stript Sauadge is for forme and bignesse equall with the Chrystall or White Carnation, but as inconstant as eyther of them, changing into red or blush; so that few branches with flowers containe their true mixtures, which are a whitish blush, fairely striped with a crimson red colour, thicke and short, with some spots also among.

*Caryophyllus maximus Sabaudicus carneus.*

The blush Sauadge is the same with the former, the same root of the stript Sauadge, as I said before, yeelding one side or part whose flowers will be eyther wholly blush, or hauing some small spots, or sometimes few or none in them.

*Caryophyllus maximus Sabaudicus ruber.*

The red Sauadge is as the blush, when the colour of the flower is wholly red without any stripes or spots in them, and so abideth long; yet it is sometimes seene, that the same side, or part, or roote being separate from the first or mother plant, will giue striped and well marked flowers againe.

*Caryophyllus maximus Oxoniensis.*

The Oxeford Carnation is very like vnto the French Carnation, both for forme, largenesse and colour: but that this is of a sadder red colour, so finely marbled with white thereon, that the red hauing the maistry, sheweth a very sad flower, not hauing any flakes or stripes at all in it.

*Caryophyllus maximus Regius, siue Bristoliensis maior.*

The Kings Carnation or ordinary Bristow, is a reasonable great flower, deepely iagged, of a sad red, very smally striped and speckled with white: some of the leaues of the flower on the one side will turne vp their brimmes or edges: the greene leafe is very large.

*Caryophyllus maximus Granatensis.*

The greatest *Granado* is a very faire large flower, bigger then the Chrystall, and almost as bigge as the blew *Hulo*: it is almost equally diuided and stript with purple and white, but the purple is sadder then in the ordinary *Granado* Gilloflower, else it might bee said it were the same, but greater. Diuers haue taken this flower to bee the *Gran Pere*, but you shall haue the difference shewed you in the next ensuing flower.

*Caryophyllus maximus Gran Pere dictus.*

The *Gran Pere* is a fair great flower, and comely for the forme, but of no great beautie for colour, because although it be stript red and white like the Queenes Gilloflower, yet the red is so sad that it taketh away all the delight to the flower.

*Caryophyllus maximus Camberfine dictus.*

The Camberfine is a great flower and a faire, beeing a redde flower, well marked or striped with white, somewhat like vnto a

Sauadge

Sauadge, say some, but that the red is not crimson as the Sauadge; others say the Daintie, but not so comely : the leaues of the flowers are many, and thrust together, without any due forme of spreading.

*Caryophyllus maximus Longobardicus ruber.*

The great Lombard red is a great sad red flower, so double and thick of leaues, that it most vsually breaketh the pod, and seldome sheweth one flower among twenty perfect: the blades or greene leaues are as large as the *Hulo*.

## *Caryophylli majores.* GILLOFLOWERS.

*Caryophyllus maior Westmonasterienſis.*

THe lustie Gallant or Westminster (some make them to be one flower, and others to bee two, one bigger then the other) at the first blowing open of the flower sheweth to be of a reasonable size and comelinesse, but after it hath stood blowen some time it sheweth smaller and thinner : it is of a bright red colour, much striped and speckled with white.

*Caryophyllus maior Briſtolienſis purpureus.*

The Bristow blew hath greene leaues, so large, that it would seeme to bring a greater flower then it doth, yet the flower is of a reasonable size, and very like vnto the ordinary *Granado* Gilloflower, striped and flaked in the same manner, but that the white of this is purer then that, and the purple is more light, and tending to a blew: this doth not abide constant, but changeth into purple or blush.

*Caryophyllus maior Briſtolienſis carneus.*

The Bristow blush is very like the last both in leafe and flower, the colour only sheweth the difference, which seldome varyeth to be spotted, or change colour.

*Caryophyllus maior Dorobernienſis ruber.*

The red Douer is a reasonable great Gilloflower and constant, being of a faire red thicke poudered with white spots, and seemeth somewhat like vnto the ground Carnation.

*Caryophyllus maior Dorobernienſis dilutus ſiue albus.*

The light or white Douer is for forme and all other things more comely then the former, the colour of the flower is blush, thicke spotted with very small spots, that it seemeth all gray, and is very delightfull.

*Caryophyllus maior Cantii.*

The faire maide of Kent, or Ruffling Robin is a very beautiful flower, and as large as the white Carnation almost: the flower is white, thicke poudered with purple, wherein the white hath the mastrie by much, which maketh it the more pleasant.

*Caryophyllus maior Regineus.*

The Queenes Gilloflower is a reasonable faire Gilloflower although very common, striped red and white, some great and some small with long stripes.

*Caryophyllus maior elegans.*

The Daintie is a comely fine flower, although it be not great, and for the smallnesse and thinnesse of the flower being red so finely marked, striped and speckled, that for the liuelinesse of the colours it is much desired, beeing inferiour to very few Gilloflowers.

*Caryophyllus maior Braſilienſis.*

The Brassill Gilloflower is but of a meane size, being of a sad purple colour, thicke poudered and speckled with white, the purple herein hath the mastrie, which maketh it shew the sadder, it is vnconstant, varying much and often to bee all purple : the greene leaues lye matting on the ground.

*Caryophyllus maior Granatenſis.*

The *Granado* Gilloflower is purple and white, flaked and striped very much: this is also much subiect to change purple. There is a greater and a lesser of this kinde, besides the greatest that is formerly described.

The

*Caryophyllus Turcicus.* The Turkie Gilloflower is but a small flower, but of great delight, by reason of the well marking of the flower, being most vsually equally striped with red and white.

*Caryophyllus Cambrensis Poole.* The Poole flower, growing naturally vpon the rockes neare Cogshot Castle in the Isle of Wight, is a small flower, but very pleasant to the eye, by reason of the comely proportion thereof; it is of a bright pale red, thicke speckled, and very small with white, that it seemeth to bee but one colour, the leaues of the flower are but smally iagged about: it is constant.

*Caryophyllus Pegma dilutior.* The light or pale Pageant is a flower of a middle size, very pleasant to behold, and is both constant and comely, and but that it is so common, would be of much more respect then it is: the flower is of a pale bright purple, thicke poudered, and very euenly with white, which hath the mastery, and maketh it the more gracefull.

*Caryophyllus Pegma saturatior.* The sad Pageant is the same with the former in forme and bignesse, the difference in colour is, that the purple hath the mastery, which maketh it so sad, that it doth resemble the Brassill for colour, but is not so bigge by halfe.

*Caryophyllus Heroina dictus elegans Magistri Bradshawij.* Master Bradshawe his dainty Lady may bee well reckoned among these sorts of Gilloflowers, and compare for neatenesse with most of them: the flower is very neate, though small, with a fine small iagge, and of a fine white colour on the vnderside of all the leaues, as also all the whole iagge for a pretty compasse, and the bottome or middle part of the flower on the vpperside also: but each leafe is of a fine bright pale red colour on the vpperside, from the edge to the middle, which mixture is of wonderfull great delight.

*Caryophyllus albus optimus maior Londinensis & alius.* The best white Gilloflower groweth vpright, and very double, the blades growe vpright also, and crawle not on the ground.

The London white is greater and whiter then the other ordinary white, being wholly of one colour.

*caryophyllus maior rubens & minor.* The stamell Gilloflower is well knowne to all, not to differ from the ordinary red or cloue Gilloflower, but only in being of a brighter or light red colour: there is both a greater and a lesser of this kinde.

*caryophyllus purpureus maior & minor.* The purple Gilloflower a greater and a lesse: the stalke is so slender, and the leaues vpon them so many and thicke, that they lye and traile on the ground: the greatest is almost as bigge as a Chrystall, but not so double: the lesse hath a smaller flower.

*caryophyllus Persico violaceus.* The Gredeline Gilloflower is a very neate and handsome flower, of the bignesse of the Cloue red Gilloflower, of a fine pale reddish purple or peach colour, enclining to a blew or violet, which is that colour is vsually called a gredeline colour: it hath no affinity with eyther Purple, Granado, or Pageant.

*Caryophyllus purpuro cæruleus.* The blew Gilloflower is neither very double nor great, yet round and handsome, with a deepe iagge at the edge, and is of an exceeding deepe purple colour, tending to a tawnie: this differeth from all other sorts, in that the leafe is as greene as grasse, and the stalkes many times red or purple: by the greene leaues it may be knowne in the Winter, as well as in the Summer.

*Caryophyllus carneus.* The blush Gilloflower differeth not from the red or stamell, but only in the colour of the flower, which is blush.

*Caryophyllus Silesiacus maximus Wittie.* Iohn Wittie his great tawny Gilloflower is for forme of growing, in leafe and flower altogether like vnto the ordinary tawny, the flower onely, because it is the faireſt and greatest that any o-

ther

ther hath nourled vp, maketh the difference, as also that it is of a faire deepe scarlet colour.

There are also diuers other Tawnies, either lighter or sadder, either lesse or more double, that they cannot be numbered, and all rising (as I said before) from sowing the seede of some of them: besides the diuersities of other colours both simple and mixed, euery yeare and place yeelding some variety was not seen with them before: I shall neede but onely to giue you the names of some of them we haue abiding with vs, I meane such as haue receiued names, and leaue the rest to euery ones particular denomination.

Of Blushes there are many sorts, as the deepe blush, the pale blush, the Infanta blush, a blush enclining to a red, a great blush, the fairest and most double of all the other blushes, and many others both single and double.

Of Reds likewise there are some varieties, but not so many as of the other colours; for they are most dead or deepe reds, and few of a bright red or stamell colour; and they are single like Pinkes, either striped or speckled, or more double striped and speckled variably, or else

There are neither purple nor white that rise from this seede that I haue obserued, except one white in one place.

*Caryophyllus Silesiacus striatus.*

The striped Tawny are either greater or lesser, deeper or lighter flowers twenty sorts and aboue, and all striped with smaller or larger stripes, or equally diuided, of a deeper or lighter colour: and some also for the very shape or forme will bee more neate, close, and round; others more loose, vnequall, and sparsed.

*Caryophyllus Silesiacus marmor-amulus.*

The marbled Tawny hath not so many varieties as the striped, but is of as great beauty and delight as it, or more: the flowers are greater or smaller, deeper or lighter coloured one then another, and the veines or markes more conspicuous, or more frequent in some then in others: but the most beautifull that euer I did see was with Master Ralph Truggie, which I must needes therefore call

*Heroina Rodolphi florum Imperatoris.*

Master Tuggies Princesse, which is the greatest and fairest of all these sorts of variable tawnies, or seed flowers, being as large fully as the Prince or Chrystall, or something greater, standing comely and round, not loose or shaken, or breaking the pod as some other sorts will; the marking of the flower is in this manner: It is of a stamell colour, striped and marbled with white stripes and veines quite through euery leafe, which are as deeply iagged as the Hulo: sometimes it hath more red then white, and sometimes more white then red, and sometimes so equally marked, that you cannot discerne which hath the mastery; yet which of these hath the predominance, still the flower is very beautifull, and exceeding delightsome.

*Caryophyllus Silesiacus assulosus*

The Flaked Tawny is another diuersity of these variable or mixt coloured flowers, being of a pale reddish colour, flaked with white, not alwaies downeright, but often thwart the leaues, some more or lesse then others; the marking of them is much like vnto the Chrystall: these also as well as others will be greater or smaller, and of greater or lesse beauty then others.

*Caryophyllus Silesiacus plumatus.*

The Feathered Tawny is more rare to meete with then many of the other; for most vsually it is a faire large flower and double, equalling the Lumbard red in his perfection: the colour hereof is vsually a scarlet, little deeper or paler, most curiously feathered and streamed with white through the whole leafe.

*Caryophyllus Silesiacus punctatus.*

The Speckled Tawny is of diuers sorts, some bigger, some lesse,

1 *Heroina Radolphi florum Imperatoris Princessa dicta.* Master Tuggie his Princesse. 2 *Caryophyllus Oxoniensis.* The French or Oxford Carnation. 3 *Caryophyllus Westmonasteriensis.* The Gallant or Westminster Gilloflower. 4 *Caryophyllus Bristoliensis.* The Bristow. 5 *Caryophyllus Chrystallinus.* The Chrystall or Chrystalline. 6 *Caryophyllus Sabaudicus striatus.* The stript Sauadge. 7 *Caryophyllus Granatensis maximus.* The Granpere or greatest Granado. 8 *Caryophyllus peramœnus.* The Dainty. 9 *Caryophyllus Silesiacus maximus Iuvenij Ioannis* Iohn Witty his great tawny Gilloflower. 10 *Caryophyllus Silesiacus striatus.* The stript Tawny. 11 *Caryophyllus mariner similis.* The wrinkled Tawny. 12 *Caryophyllus rosinus rotundus magistri Tuggie.* Master Tuggie his Rose Gilloflower.



*Caryophyllus roseus rotundus Magistri Tuggie.*

lesse, some more, and some lesse spotted then others: Vsually it is a deepe scarlet, speckled or spotted with white, hauing also some stripes among the leaues.

Master Tuggie his Rose Gilloflower is of the kindred of these Tawnies, being raised from the seede of some of them, and onely possessed by him that is the most industrious pr eseruer of all natures beauties, being a different sort from all other, in that it hath round leaues, without any iagge at all on the edges, of a fine stamell full colour, without any spot or strake therin, very like vnto a small Rose, or rather much like vnto the red Rose Campion, both for forme, colour, and roundnesse, but larger for size.

## The Place.

All these are nourished with vs in Gardens, none of their naturall places being knowne, except one before recited, and the yellow which is *Silesia*; many of them being hardly preserued and encreased.

## The Time.

They flower not vntill the heate of the yeare, which is in Iuly (vnlesse it be an extraordinary occasion) and continue flowring, vntill the colds of the Autumne checke them, or vntill they haue wholly out spent themselues, and are vsually encreased by the slips.

## The Names.

Most of our later Writers doe call them by one generall name, *Caryophyllus sativus*, and *flos Caryophylleus*, adding thereunto *maximus*, when wee meane Carnations, and *maior* when we would expresse Gilloflowers, which name is taken from Cloues, in that the sent of the ordinary red Gilloflower especially doth resemble them. Diuers other seuerall names haue beene formerly giuen them, as *Vetonica*, or *Betonica altera*, or *Vetonica altilis*, and *coronaria*. *Herba Tunica*, *Viola Damascena*, *Ocellus Damascenus*, and *Barbaricus*. Of some *Cantabrica Plinij*. Some thinke they were vnknowne to the Ancients, and some would haue them to be *Iphium* of Theophrastus, whereof he maketh mention in his fixth and seuenth Chapters of his sixth booke, among Garland and Summer flowers; others to be his *Dios anthos*, or *Iouis flos*, mentioned in the former, and in other places. We call them in English (as I said before) the greatest kindes, Carnations, and the others Gilloflowers (*quasi* Iuly flowers) as they are seuerally expressed.

## The Vertues.

The red or Cloue Gilloflower is most vsed in Physicke in our Apothecaries shops, none of the other being accepted of or vsed (and yet I doubt not, but all of them might serue, and to good purpose, although not to giue so gallant a tincture to a Syrupe as the ordinary red will doe) and is accounted to be very Cordiall.

# CHAP. LXX.

## *Caryophylli siluestres.* Pinkes.

THere remaine diuers sorts of wilde or small Gilloflowers (which wee vsually call Pinkes) to be entreated of, some bearing single, and some double flowers, some smooth, almost without any deepe dents on the edges, and some iagged, or as it were feathered. Some growing vpright like vnto Gilloflowers, others creeping

1 *Caryophyllus silvestris simplex.* The vsuall Engle Pinke. 2 *Caryophyllus multiplex silvestris.* Double Pinkes. 3 *Cariophyllus silvestris plumarius.* Feathered or iagged Pinkes. 4 *Caryophyllus Stellatus.* Starre Pinkes. 5 *Caryophyllus retens.* Matted Pinkes. 6 *Caryophyllus mediterraneus.* The great Thistle or Sea Gilloflower. 7 *Caryophyllus marinus.* The ordinary Thistle or Sea Cushion.

ping or spreading vnder the toppe or cruſt of the ground, ſome of one colour, ſome of another, and many of diuers colours: As I haue formerly done with the Gilloflowers, ſo muſt I doe with theſe that are entertained in our Gardens, onely giue you the deſcriptions of ſome three or foure of them, according to their variety, and the names of the reſt, with their diſtinctions.

### 1. *Caryophyllus minor ſilueſtris multiplex & ſimplex.*
Double and ſingle Pinkes.

The ſingle and double Pinkes are for forme and manner of growing, in all parts like vnto the Gilloflowers before deſcribed, ſauing onely that their leaues are ſmaller and ſhorter, in ſome more or leſſe then in others, and ſo are the flowers alſo: the ſingle kindes conſiſting of fiue leaues vſually (ſeldome ſix) round pointed, and a little ſnipt for the moſt part about the edges, with ſome threads in the middle, either crooked or ſtraight: the double kindes being leſſer, and leſſe double then the Gilloflowers, hauing their leaues a little ſnipt or endented about the edges, and of diuers ſeuerall colours, as ſhall hereafter be ſet downe, and of as fragrant a ſent, eſpecially ſome of them as they: the rootes are long and ſpreading, ſomewhat hard and wooddy.

### 2. *Caryophyllus plumarius.* Feathered or iagged Pinkes.

The iagged Pinkes haue ſuch like ſtalkes and leaues as the former haue, but ſomewhat ſhorter and ſmaller, or graſſe-like, and of a whitiſh or grayiſh greene colour likewiſe: the flowers ſtand in the like manner at the toppes of the ſtalkes, in long, round, ſlender, greene huskes, conſiſting of fiue leaues, very much cut in on the edges, and iagged almoſt like a feather, of a light red, or bright purple colour, with two white threads ſtanding in the middle, crooked like a horne at the end, and are of a very good ſent. Some of theſe haue not thoſe two crooked threads or hornes in the middle, but haue in their ſtead many ſmall threads, not crooked at all: the ſeedes of them all are like vnto the ſeedes of Gilloflowers, or the other Pinkes, that is, ſmall, blacke, long, and flat: the rootes are ſmall and wooddy likewiſe.

### 3. *Caryophyllus plumarius albus orbe rubro ſiue Stellatus.* Starre Pinkes.

Of this kinde there is another ſort, bearing flowers almoſt as deeply cut or iagged as the former, of a faire white colour, hauing a ring or circle of red about the bottome or lower part of the leaues, and are as ſweete as the former: this being ſowne of ſeede doth not giue the ſtarre of ſo bright a red colour, but becommeth more dunne.

### 4. *Caryophyllus plumarius Auſtriacus ſiue Superba Auſtriaca.*
The feathered Pinke of Auſtria.

This kinde of Pinke hath his firſt or lower leaues, ſomewhat broader and greener then any of the former Pinkes, being both for breadth and greenneſſe more like vnto the Sweete Iohns, which ſhall bee deſcribed in the next Chapter: the leaues on the ſtalkes are ſmaller, ſtanding by couples at euery ioynt, at the toppes whereof ſtand ſuch like iagged flowers as the laſt deſcribed, and as large, but more deeply cut in or iagged round about, ſome of them of a purpliſh colour, but the moſt ordinary with vs are pure white, and of a moſt fragrant ſent, comforting the ſpirits and ſenſes a farre off: the ſeedes and rootes are like vnto the former. Some haue miſtaken a kinde of wilde Campion, growing in our Woods, and by the paths ſides in Hornſie Parke, and other places, to be this feathered Pinke: but the flowers declare the difference ſufficiently.

### 5. *Caryophyllus minor repens ſimplex & multiplex.*
Single and double matted Pinkes.

The matted Pinke is the ſmalleſt, both for leafe and flower of all other Pinkes that are nouriſhed in Gardens, hauing many ſhort and ſmall graſſie greene leaues vpon the ſtalkes,

ſtalkes, which as they grow and lye vpon the ground (and not ſtanding ſo vpright as the former) doe take roote againe, whereby it quickly ſpreadeth, and couereth a great deale of ground in a little ſpace: the flowers are ſmall and round, a little ſnipt about the edges, whereof ſome are white, and ſome red, and ſome are white ſpotted with red, and ſome red ſpotted with white, all of them being ſingle flowers. But there is another of this kinde, not differing in leafe, but in flower: for that the firſt flowers are but once double, or of two rowes of leaues, of a fine reddiſh colour, ſpotted with ſiluer ſpots: but thoſe that follow, are ſo thicke and double, that they oftentimes doe breake the pod or huske; being not altogether of ſo deepe a red colour, but more pale.

*6. Caryophyllus Mediterraneus ſiue Marinus maior.*
Great Sea Gilloflower or Great Thrift.

Vnto theſe kindes of Pinkes I muſt needs adde, not only our ordinary Thrift (which is more frequent in gardens, to empale or border a knot, becauſe it abideth greene Winter and Summer, and that by cutting, it may grow thick, and be kept in what form one liſt, rather then for any beautie of the flowers) but another greater kinde, which is of as great beautie and delight almoſt as any of the former Pinkes, as well for that the leaues are like vnto Gilloflowers, being longer and larger then any Pinkes, and of a whitiſh greene colour like vnto them, not growing long or by couples vpon the ſtalkes as Pinkes and Gilloflowers doe, but tufting cloſe vpon the ground, like vnto the common Thrift: as alſo that the ſtalkes, riſing from among the leaues (being ſometimes two foote high (as I haue obſerued in my garden) are yet ſo ſlender and weake, that they are ſcarce able to beare the heads of flowers, naked or bare, both of leaues and ioynts, ſauing only in one place, where at the ioynt each ſtalke hath two ſmall and very ſhort leaues, not riſing vpwards as in all other Gilloflowers, Pinkes, and other herbes, but growing downewards) and doe beare each of them a tuft or vmbell of ſmall purpliſh, or bluſh coloured flowers, at the toppes of them ſtanding ſomewhat like vnto ſweete Williams, but more roundly together, each flower conſiſting of fiue ſmall, round, ſtiffe or hardiſh leaues, as if they were made of paper, the bottome or middle being hollow, not blowing all at once as the ordinary Thrift, but for the moſt part one after another, not ſhewing vſually aboue foure or fiue flowers open at one time (ſo farre as I could obſerue in the plants that I kept) ſo that it was long before the whole tuft of flowers were paſt; but yet the hoter and dryer the time was, the ſooner it would be gone: the ſeede I haue not perfectly obſerued, but as I remember, it was ſomewhat like vnto the ſeede of Scabious; I am ſure nothing like vnto Gilloflowers or Pinkes: the roote is ſomewhat great, long and hard, and not ſo much ſpreading in the ground as Gilloflowers or Pinkes.

*Caryophyllus Marinus.* Thrift, or Sea Cuſhion.

Our common Thrift is well knowne vnto all, to haue many ſhort and hard greene leaues, ſmaller then many of the graſſes, growing thicke together, and ſpreading vpon the ground: the ſtalkes are naked of leaues a ſpanne high, bearing a ſmall tuft of light purple, or bluſh coloured flowers, ſtanding round and cloſe thruſting together.

## Double Pinkes.

THe double white Pinke is onely with more leaues in it then the ſingle, which maketh the difference.

The double red Pinke is in the ſame manner double, differing from the ſingle of the ſame colour.

The double purple Pinke differeth not

## Single Pinkes.

THe ſingle white ordinary Pinke hath a ſingle white flower of fiue leaues, finely iagged about the edges.

The ſingle red Pink is like the white, but that the leaues are not ſo much iagged, and the flower is of a pale purpliſh red colour.

from the single purple for colour, but only in the doubleneſſe of the flower.

The *Granado* Pinke differeth not from the Gilloflower of the ſame name, but in the ſmalneſſe both of leaues and flower.

The double Matted Pinke is before deſcribed.

The double bluſh Pinke is almoſt as great as the ordinary bluſh Gilloflower, and ſome haue taken it for one, but the greene leaues are almoſt as ſmall as Pinks, and therefore I referre it to them.

The ſingle purple Pinke is of a faire purple colour, like almoſt vnto the purple Gilloflower.

The great bluſh Pinke hath broader and larger leaues in the flower then any other Pinke, and of a faire bluſh colour.

The white Featherd Pinke hath the edges of the flower more finely and deeply cut in then the former.

The red or light purple featherd Pinke is like the former featherd Pinke, but only differeth in colour.

The Starre Pinke is a faire flower, finely iagged on the edges, with a faire red circle at the lower end of the leaues on the inſide.

The white featherd Pinke of Auſtria is deſcribed before. The purple featherd Pinke of Auſtria is ſo likewiſe. The ſingle matted Pinke is before deſcribed. The ſpeckled Pinke is a ſmall flower hauing ſmall ſpotsof red here and there diſperſed ouer the white flower.

Thoſe ſingle flowers being like vnto Pinkes that riſe from the ſowing of the orenge tawney, I bring not into this *claſsis*, hauing already ſpoken of them in the precedent Chapter.

## The Place.

Theſe are all like as the former, nouriſhed in Gardens with vs, although many of them are found wilde in many places of Auſtria, Hungarie, and Germany, on the mountaines, and in many other places, as Cluſius recordeth. The ordinary Thrift groweth in the ſalt Marſhes at Chattam by Rocheſter, and in many other places in England : but the great kinde was gathered in Spaine, by Guillaume Boel that painefull ſearcher of ſimples, and the ſeede thereof imparted to me, from whence I had diuers plants, but one yeare after another they all periſhed,

## The Time.

Many of theſe Pinkes both ſingle and double, doe flower before any Gilloflower, and ſo continue vntill Auguſt, and ſome, moſt of the Summer and Autumne.

## The Names.

The ſeuerall titles that are giuen to theſe Pinkes, may ſuffice for their particular names: and for their generall they haue beene expreſſed in the former Chapter, beeing of the ſame kindred, but that they are ſmaller, and more frequently found wilde. The two ſorts of Thrift are called *Caryophyllus Marinus*. The greater, *Maior & Mediterraneus*; In Engliſh, The greater or Leuant Thrift, or Sea Gilloflower. The leſſer *Minimus*, and is accounted of ſome to be a graſſe, and therefore called *Gramen Marinum & Polyanthemum*; In Engliſh, Thrift, Sea graſſe, and our Ladies Cuſhion, or Sea Cuſhion.

## The Vertues.

It is thought by diuers, that their vertues are anſwerable to the Gilloflowers, yet as they are of little vſe with vs, ſo I thinke of as ſmall effect.

## Chap. LXXI.

### *Armerius.* Sweet Iohns, and sweet Williams.

THese kindes of flowers as they come neereſt vnto Pinkes and Gilloflowers, though manifeſtly differing, ſo it is fitteſt to place them next vnto them in a peculiar Chapter.

1. *Armerius anguſtifolius rubens ſimplex.* Single red ſweete Iohns.

The ſweete Iohn hath his leaues broader, ſhorter and greener then any of the former Gilloflowers, but narrower then ſweete Williams, ſet by couples, at the ioynts of the ſtalkes, which are ſhorter then moſt of the former, and not aboue a foote and a halfe high, at the tops whereof ſtand many ſmall flowers, like vnto ſmall Pinkes, but ſtanding cloſer together, and in ſhorter huskes, made of fiue leaues, ſmaller then moſt of them, and more deeeply iagged then the Williams, of a red colour in the middle, and white at the edges, but of a ſmall or ſoft ſent, and not all flowring at once, but by degrees: the ſeede is blacke, ſomewhat like vnto the ſeede of Pinkes, the roote is diſperſed diuerſly, with many ſmall fibres annexed vnto it.

2. *Armerius anguſtifolius albus ſimplex.* Single white ſweet Iohns.

This white Iohn differeth not in any thing from the former, but onely that the leafe doth neuer change browniſh, and that the flower is of a faire white colour, without any mixture.

3. *Armerius anguſtifolius duplex.* Double ſweet Iohns.

There is of both thoſe former kindes, ſome whoſe flowers are once double, that is, conſiſting of two or three rowes of leaues, and the edges not ſo deepely iagged; not differing in any thing elſe.

4. *Armerius latifolius ſimplex flore rubro.*
Single red ſweet Williams.

The ſweet Williams doe all of them ſpread into many very long trayling branches, with leaues lying on the ground, in the very like manner that the ſweete Iohns doe: the chiefe differences betweene them are, that theſe haue broader, and darker greene leaues, ſomewhat browniſh, eſpecially towards the points, and that the flowers ſtand thicker and cloſer, and more in number together, in the head or tuft, hauing many ſmall pointed leaues among them, but harmleſſe, as all men know; the colour of the flower is of a deep red, without any mixture or ſpot at all.

5. *Armerius latifolius flore rubro multiplici.*
Double red ſweete Williams.

The double kinde differeth not from the ſingle kinde of the ſame colour, but only in the doubleneſſe of the flowers, which are with two rowes of leaues in euery flower.

6. *Armerius latifolius variegatus ſiue verſicolor.*
Speckled ſweete Williams, or London pride.

Theſe ſpotted Williams are very like the firſt red Williams, in the forme or maner of growing, hauing leaues as broade, and browne ſometimes as they, the flowers ſtand as thicke or thicker, cluſtring together, but of very variable colours: for ſome flowers will be of a fine delayed red, with few markes or ſpots vpon them, and others will

will bee full peckled or sprinkled with white or siluer spots, circlewise about the middle of the flowers, and some will haue many specks or spots vpon them dispersed: all these flowers are not blowne at one time, but some are flowring, when others are decaying, so that abiding long in their pride, they become of the more respect: The seede is blacke, as all the rest, and not to be distinguished one from another: the roots are some long, and some small and threddy, running vnder the vpper crust of the earth.

7. *Armerius latifolius flore rubro saturo holoserico.*
Sweet Williams of a deepered or murrey colour.

The leaues of this kinde seeme to be a little larger, and the ioints a little redder then the former, but in the flower consisteth the chiefest difference, which is of a deepe red, or murrey purple colour, like vnto veluet of that colour, without any spots, but smooth, and as it were soft in handling, hauing an eye or circle in the middle, at the bottome of the leaues.

8. *Armerius latifolius simplex flore albo.*
Single white sweete Williams.

The white kinde differeth not in forme, but in colour from the former, the leaues are not browne at all, but of a fresh greene colour, and the flowers are wholly white, or else they are all one.

## The Place.

These for the most part grow wilde in Italie, and other places: we haue them in our Gardens, where they are cherished for their beautifull varietie.

## The Time.

They all generally doe flower before the Gilloflowers or Pinkes, or with the first of them: their seede is ripe in Iune and Iuly, and doe all well abide the extremitie of our coldest winters.

## The Names.

They all generally are called *Armerius*, or *Armeria*, as some doe write, and distinguished as they are in their titles: Yet some haue called them *Vetonica agrestis*, and others *Herba Tunica, Scarlatea, & Caryophyllus siluestris*: Wee doe in English in most places, call the first or narrower leafed kindes, Sweet Iohns, and all the rest Sweete Williams; yet in some places they call the broader leafed kindes that are not spotted, Tolmeiners, and London tufts: but the speckled kinde is termed by our English Gentlewomen, for the most part, London pride.

## The Vertues.

We haue not knowne any of these vsed in Physicke.

# Chap. LXXII.

## *Bellis.* Daisie.

THere be diuers sorts of Daisies, both great and small, both single and double, both wilde growing abroade in the fieldes, and elsewhere, and manured growing only in Gardens: of all which I intend not to entreate, but of those that are of most beautie and respect, and leaue the rest to their proper place.

1 *Armerius angustifolius simplex.* Single sweete Iohns. 2 *Armerius angustifolius multiplex* Double sweet Iohns. 3 *Armerius latifolius simplex* Single sweete Williams. 4 *Armerius latifolius versicolor.* Spotted sweet Williams or pride of London. 5 *Armerius latifolius multiplex* Double sweet Williams. 6 *Bellis hortensis minor multiplex.* Double Garden Daisies. 7 *Bellis minor hortensis flore varie.* Double red Daisies strict 8 *Bellis minor hortensis prolifera.* Double fruitfull Daisies or lacke an Apes on horseback. 9 *Bellis caerulea siue Globularia.* Double blew Daisies or blew Globeflower. 10 *Bellis lutea montana siue Globularia lutea montana,* Double yellow Daisies or yellow Globeflower.

1. *Bellis maior flore albo pleno.* The great double white Daiſie.

The great Daiſie with the double white flower, is in all things ſo like vnto the great ſingle kinde, that groweth by the high wayes, and in diuers medowes and fields, that there is no difference but in the flower, which is double. It hath many long, and ſomewhat broad leaues lying vpon the ground, deepely cut in on both ſides, ſomewhat like vnto an oaken leafe; but thoſe that are on the ſtalkes are ſhorter, narrower, and not ſo deeply cut in, but onely notched on the edges: the flowers at the toppe are (as I ſaid) white and double, conſiſting of diuers rowes of leaues, being greater in compaſſe then any of the double Daiſies that follow, but nothing ſo double of leaues.

2. *Bellis minor flore rubro ſimplici.* Single red Daiſies.

This ſingle Daiſie (like as all the reſt of the ſmall Daiſies) hath many ſmooth, greene, round pointed leaues lying on the ground, a little ſnipt about the edges; from among which riſe many ſlender round foote-ſtalkes, rather then ſtalkes or ſtems, about an hand breadth high at the moſt, and oftentimes not halfe ſo high, bearing one flower a peece, conſiſting of many ſmall leaues, as a pale or border ſet about a middle thrumme: the leaues of this kinde are almoſt wholly red, whereas in the wilde they are white or whitiſh, enclining to red on the edges, the middle being yellow in both ſorts: the rootes are many ſmall white threads or ſtrings.

3. *Bellis minor hortenſis flore pleno variorum colorum.*
Double Garden Daiſies of diuers colours.

The leaues of all the double Daiſies are in forme like vnto the ſingle ones, but that they are ſmaller, and little or nothing ſnipt or notched about the edges: the ſmall ſtalkes likewiſe are ſmaller and lower, but bearing as double flowers as any that growe on the ground, being compoſed of many ſmall leaues, thicke thruſt together, of diuers colours; for ſome are wholly of a pure white, others haue a little red, either diſperſed vpon the white leaues, or on the edges, and ſometimes on the backes of the leaues: ſome againe ſeeme to be of a whitiſh red, or more red then white, when as indeede they are white leaues diſperſed among the red; others of a deepe or darke red colour, and ſome are ſpeckled or ſtriped with white and red through the whole flower: and ſome the leaues will bee red on the vpperſide, and white vnderneath; and ſome alſo (but thoſe are very rare) are of a greeniſh colour.

4. *Bellis minor hortenſis prolifera.* Double double Daiſies or childing Daiſies.

There is no difference either in leafe or roote in this kinde from the former double Daiſies: the chiefeſt variety conſiſteth in this, that it beareth many ſmall double flowers, ſtanding vpon very ſhort ſtalkes round about the middle flower, which is vſually as great and double as any of the other double kindes, and is either wholly of a deepe red colour, or ſpeckled white and red as in ſome of the former kindes, or elſe greeniſh, all the ſmall flowers about it being of the ſame colour with the middlemoſt.

5. *Bellis cærulea ſiue Globularia.* Blew Daiſies.

The likeneſſe and affinity that this plant hath with the former, both in the forme of leafe and flower, as alſo in the name, hath cauſed me to inſert it, and another rare plant of the ſame kinde, in this place, although they be very rare to be met with in our Engliſh Gardens. This beareth many narrower, ſhorter, and blacker greene leaues then the former, lying round about vpon the ground; among which riſe vp ſlender, but ſtiffe and hard ſtalks, halfe a foot high or more, ſet here and there with ſmall leaues, and at the top a ſmall round head, compoſed of many ſmall blew leaues, ſomewhat like vnto the head of a Scabious: It hath bin found likewiſe with a white head of flowers: the roote is hard and ſtringie: the whole plant is of a bitter taſte.

### 6. *Globularia lutea montana.* Yellow Daisies.

This mountaine yellow Daisie or Globe-flower hath many thicke, smooth, round pointed leaues, spread vpon the gronnd like the former; among which spring diuers small round rushie stalkes, a foote high, bearing about the middle of them two small leaues at the ioynts, and at the toppes round heads of flowers thrust thicke together, standing in purplish huskes, euery of which flowers do blow or spread into fiue leaues, starre-fashion, and of a faire yellow colour, smelling like vnto broome flowers, with many small threads in the middle compassing a flat pointell, horned or bended two wayes: after the flowers are past rise vp the seede vessels, which are round, swelling out in the middle, and diuided into foure parts at the toppes, containing within them round, flat, blacke seede, with a small cut or notch in them: the roote is a finger long, round and hard, with a thicke barke, and a woddy pith in the middle, of a sharpe drying taste and strong sent: the leaues are also sharpe, but bitter.

#### The Place.

The small Daisies are all planted, and found onely in Gardens, and will require to be replanted often, lest they degenerate into single flowers, or at least into lesse double. The blew Daisie is naturall of Mompelier in France, and on the mountaines in many places of Italy, as also the yellow kinde in the Kingdome of Naples.

#### The Time.

The Daisies flower betimes in the Spring, and last vntill May, but the last two flower not vntill August or September.

#### The Names.

They are vsually called in Latine *Bellides*, and in English Daisies. Some call them *Herba Margarita*, and *Primula veris*, as it is likely after the Italian names, of *Marguerite*, and *Fior di prima vera gentile*. The French call them *Pasquettes*, and *Marguerites*, and the Fruitfull sort, or those that beare small flowers about the middle one, *Margueritons*: our English women call them Iacke an Apes on horse-backe, as they doe Marigolds before recited, or childing Daisies: but the Physitians and Apothecaries doe in generall call them, especially the single or Field kindes, *Consolida minor*. The blew Daisie is called *Bellis carulea*, and *Globularia*, of some *Scabiosæ pumilum genus*. The Italians call it *Botanaria*, because the heads are found like buttons. The yellow, *Globularia montana*, is onely described by Fabius Columna, in his last part of *Phytobasanos*, and by him referred vnto the former *Globularia*, although it differ in some notable points from it.

#### The Vertues.

The properties of Daisies are certainly to binde, and the roote especially being dryed, they are vsed in medicines to that purpose. They are also of speciall account among those herbes, that are vsed for wounds in the head.

## CHAP. LXXIII.

### *Scabiosa.* Scabious.

THe sorts of Scabious being many, yeeld not flowers of beauty or respect, fit to bee cherished in this our Garden of delight; and therefore I leaue them to the Fields and Woods, there to abide. I haue onely two or three strangers to bring to your acquaintance, which are worthy this place.

1. *Scabiosa flore albo.* White flowred Scabious.

This white Scabious hath many long leaues, very much iagged or gasht in on both sides, of a meane bigneſſe, being neither so large as many of the field, nor so small as any of the small kindes: the stalkes rise about a foote and a halfe high, or somewhat higher, at the tops whereof grow round heads, thicke set with flowers, like in all points vnto the field Scabious, but of a milke white colour.

2. *Scabiosa rubra Austriaca.* Red Scabious of Austria.

This red Scabious hath many leaues lying vpon the ground, very like vnto Deuils bit, but not so large, being shorter and snipt, not gashed about the edges, of a light greene colour; yet (there is another of a darker greene colour, whose flower is of a deeper red) the stalkes haue diuers such leaues on them, set by couples at the ioynts as grow belowe, and at the tops small heads of flowers, each consisting of fiue leaues, the biggest flowers standing round about in the outer compasse, as is vsuall almost in all the kinds of Scabious, of a fine light purple or red colour: after the flowers are past, come the seede, which is somewhat long and round, set with certaine haires at the head thereof, like vnto a Starre: the roote is composed of a number of slender strings, fastened at the head.

3. *Scabiosa rubra Indica.* Red flowred Indian Scabious.

This (reputed Indian) Scabious hath many large faire greene leaues lying on the ground, iagged or cut in on both sides to the middle ribbe, euery peece whereof is narrower then that at the end, which is the broadest: among these leaues rise vp sundry slender and weake stalkes, yet standing vpright for the most part, set with smaller and more iagged leaues at certaine distances, two or three at euery ioynt, branching forth at the toppe into other smaller branches, bearing euery one head of flowers, like in forme vnto other Scabiouses, but of an excellent deepe red crimson colour (and sometimes more pale or delayed) of no sent at all: after which doe come small roundish seede, like vnto the field Scabious: the roote is long and round, compassed with a great many small strings, and perisheth vsually as soone as it hath borne out his flowers and seede: otherwise if it doe not flower the first yeare of the sowing, if it be carefully defended from the extremity of Winter, it will flower the sooner the next yeare, as I my selfe haue often found by experience.

### The Place.

The first is sometimes found wilde in our owne Countrey, but it is very geason, and hath been sent among other rare seedes from Italy.

The second was first found and written of by Clusius, in Pannonia and Austria, where it is very plentifull.

The third hath been sent both from Spaine & Italy, and is verily thought to grow naturally in both those parts.

The

1 *Scabiosa flore albo.* White flowred Scabious 2 *Scabiosa rubra Austriaca.* Red Scabious of Austria. 3 *Scabiosa rubra Indica,* Red flowred Indian Scabious. 4 *Cyanus vulgaris minor.* Corn-flower of diuers colours. 5 *Cyanus Baticus.* Spanish Corn-flower. 6 *Cyanus floridus Turcicus.* The braue Sultans flower. 7 *Carthamus sativus.* Spanish Saffron.

### The Time.

The first and second flower earlier then the last, for that it flowreth not vntill September or October, (vnlesse it be not apt to beare the first yeare as I before said) so that many times (if none be more forward) it perisheth without bearing ripe seede, whereby we are oftentimes to seeke new seede from our friends in other parts.

### The Names.

They haue all one generall name of Scabious, distinguished eyther by their flower, or place of growing, as in their titles: yet the last is called of diuers *Scabiosa exotica*, because they thinke the name *Indica*, is not truely imposed vpon it.

### The Vertues.

Whether these kindes haue any of the vertues of the other wilde kinds, I know none haue made any experience, and therefore I can say no more of them.

## Chap. LXXIIII.

### *Cyanus*. Corne flower, or blew Bottles.

VNder the name of *Cyanus* are comprehended, not onely those plants which from the excellent blew colour of their flowers (furnishing or rather pestering the Corne fieldes) haue peculiarly obtained that name, and which doth much vary also, in the colour of the flowers, as shallbe shewed; but some other plants also for their neere resemblance, but with seuerall distinctions. The *Cyanus maior*, *Ptarmica Austriaca*, *Ptarmica Imperati*, and many others which may be adioyned vnto them, do more fitly belong to the Garden of Simples, whereunto I leaue them, and will here only entreate of those that may most please the delight of our Gentle Florists, in that I labour and striue, to furnish this our garden, with the chiefest choyse of natures beauties and delights.

#### 1. *Cyanus vulgaris diuersorum colorum*. Corne flower of diuers colours.

All these sorts of Corne flowers are for the most part alike, both in leaues and flowers one vnto another for the forme: the difference betweene them consisteth in the varying colour of the flowers: For the leaues are long, and of a whitish greene colour, deeply cut in on the edges in some places, somewhat like vnto the leaues of a Scabious: the stalkes are two foote high or better, beset with such like leaues but smaller, and little or nothing slit on the edges: the toppes are branched, bearing many smal greene scaly heads, out of which rise flowers, consisting of fiue or sixe, or more long and hollow leaues, small at the bottome, and opening wider and greater at the brims, notched or cut in on the edges, and standing round about many small threds in the middle: the colours of these flowers are diuers, and very variable; for some are wholly blew, or white, or blush, or of a sad, or light purple, or of a light or dead red, or of an ouerworne purple colour, or else mixed of these colours, as some, the edges white, and the rest blew or purple, or the edges blew or purple, and the rest of the flower white, or striped, spotted, or halfed, the one part of one colour, and the other of another, the threds likewise in the middle varying in many of them; for some will haue the middle thrume of a deeper purple then the outer leaues, and some haue white or blush leaues, the middle thrume being reddish, deeper or paler: After the flowers are past, there come small, hard, white and shining seede in those heads, wrapped

ped or set among a deale of flockie matter, as is most vsuall, in all plants that beare scaly heads: the rootes are long and hard, perishing euery yeare when it hath giuen seede.

2. *Cyanus floridus Turcicus.* The Sultans flower.

As a kinde of these Corne flowers, I must needes adioyne another stranger, of much beautie, and but lately obtained from Constantinople, where, because (as it is said) the great Turke, as we call him, saw it abroade, liked it, and wore it himselfe; all his vassals haue had it in great regard, and hath been obtained from them, by some that haue sent it into these parts. The leaues whereof are greener, and not only gashed, but finely snipt on the edges: the stalkes are three foote high, garnished with the like leaues as are below, and branched as the former, bearing large scaly heads, and such like flowers but larger, hauing eight or nine of those hollow gaping leaues in euery flower, standing about the middle threds (if it be planted in good and fertile ground and be well watered, for it soone starueth and perisheth with drought) the circling leaues are of a fine delayed purple or blush colour, very beautifull to behold; the seede of this is smaller and blacker, and not enclosed in so much dounie substance, as the former (yet in our Countrey the seede is not so blacke, as it came vnto vs, but more gray) the roote perisheth likewise euery yeere.

3. *Cyanus Bæticus supinus.* The Spanish Corne-flower.

This Spanish kinde hath many square low bending or creeping stalkes, not standing so vpright as the former, but branching out more diuersly; so that one plant will take vp a great deale of ground: the leaues are broader then any of the rest, softer also, of a pale or whitish greene colour, and not much gashed on the edges: the flowers stand in bigger heads, with foure or fiue leaues vnder euery head, and are of a light pale purple or blush colour; after which come seede, but not so plentifully, yet wrapped in a great deale of flockie matter, more then any: the roote groweth downe deepe into the ground, but perisheth euery yeare as they doe.

The Place.

The first or former kindes, grow many times in the Corne fields of our own Countrey, as well as of others, especially that sort with a blew flower: but the other sorts or colours are not so frequent, but are nourished in gardens, where they will vary wonderfully.

The second as is before set downe, groweth in Turkie: and the last in Spaine, found out and first sent to vs by that industrious searcher of simples, Guillaume Boel before remembred.

The Time.

The first doe flower in the end of Iune, and in Iuly, and somtimes sooner. The other two later, and not vntill August most commonly, and the seede is soone ripe after.

The Names.

The first is generally called *Cyanus*, and some following the Ditch name, call it *Flos frumenti*. The olde Writers gaue it the name of *Bapti sæcuba*, which is almost worne out. We doe call them in English, Blew Bottles, and in some places, Corne flowers, after the Ditch names. The second hath beene sent by the name of *Ambreboi*, which whether it be a Turkie or Arabian name, I know not. I haue called it from the place, from whence we had it, *Turcicus*, and for his beauty, *Floridus*. The Turkes themselues as I vnderstand, doe call it The Sultans flower, and I haue done so likewise, that it may bee distinguished from all the other kindes, or else you may call it,

The Turkey blush Corne flower, which you please. The last was sent by the name of *Iacea Betica*, but I had rather to referre it to the *Cyanus*, or Corne flowers, because the flowers are like vnto the Corne flowers, and not vnto the Iaceas or Knapweedes.

### The Vertues.

These had no vse in Physicke in Galen and Dioscorides time, in that (as it is thought) they haue made no mention of them: We in these dayes doe chiefly vse the first kindes (as also the greater sort) as a cooling Cordiall, and commended by some to be a remedy, not onely against the plague and pestilentiall diseases, but against the poison of Scorpions and Spiders.

## Chap. LXXV.

### *Iacea Marina Betica.* Spanish Sea Knapweede.

THere are a great many sorts of Knapweedes, yet none of them all fit for this our Garden, but this only stranger, which I haue beene bold to thrust in here, for that it hath such like gaping or open flowers, as the former Corne flowers haue, but notably differing, and therefore deserueth a peculiar Chapter, as partaking both with *Cyanus* and *Iacea*. It hath many long and narrow leaues vneuenly dented or waued on both edges (and not notched, gashed or indented, as many other herbes are) being thicke, fleshie and brittle, a little hairy, and of an ouerworne darke greene colour, among which rise lowe weake stalkes, with such like leaues as grow at the bottome, but smaller, bearing but here and there a flower, of a bright reddish purple colour, like in forme vnto the Corne flowers, but much larger, with many threds or thrumes in the middle of the same colour, standing vp higher then any of the former: this flower riseth out of a large scaly head, all set ouer with small sharpe (but harmelesse) white prickles: the seedes are blackish, like vnto the Knapweedes, and larger then any of the former Corne flowers: the roote is great and thicke, growing deepe into the ground, fleshie and full of a slimie or clammy iuice, and easie to bee broken, blackish on the outside, and whitish within, enduring many yeares, like as the other Knapweedes, or Matfelons doe, growing in time to be very thicke and great.

### The Place.

It groweth naturally by the Sea side in Spaine, from whence I receiued the seedes of Guillaume Boel, and did abide well in my garden a long time, but is now perished.

### The Time.

It flowreth in the beginning of Iuly, or thereabouts, and continueth not long in flower: but the head abideth a great while, and is of some beauty after the flower is past; yet seldome giueth good seed with vs.

### The Names.

It hath no other name then is set down in the title, being altogether a Noueliſt, and not now to be seene with any saue my selfe.

### The Vertues.

We haue not yet kuown any vse hereof in Physick.

## Chap. LXXVI.

### *Cnicus siue Carthamus sativus.* Bastard or Spanish Saffron.

THere are two or three sorts of *Cnicus* or bastard Saffrons which I passe ouer, as not fit for this Garden, and onely set downe this kinde, whose flowers are of a fairer and more liuely colour in our Country, then any hath come ouer from Spaine, where they manure it for the profit they make thereof, seruing for the dying of Silke especially, and transporting great quantities to diuers Countries. It hath large broad leaues, without any prickes at all vpon them in our Country, growing vpon the stalke, which is strong, hard, and round, with shorter leaues thereon vp to the toppe, where they are a little sharpe pointed, and prickly about the edges sometimes, which stalke riseth three or foure foote high, and brancheth it selfe toward the toppe, bearing at the end of euery branch one great open scaly head, out of which thrusteth out many gold yellow threads, of a most orient shining colour, which being gathered in a dry time, and kept dry, will abide in the same delicate colour that it bare when it was fresh, for a very long time after: when the flowers are past, the seede when it is come to maturity, which is very seldome with vs, is white and hard, somewhat long, round, and a little cornered: the roote is long, great, and wooddy, and perisheth quickly with the first frosts.

#### The Place.

It groweth in Spaine, and other hot Countries, but not wilde, for that it is accounted of the old Writers, Theophrastus and Dioscorides, to be a manured plant.

#### The Time.

It flowreth with vs not vntill August, or September sometimes, so that it hardly giueth ripe seede (as I said) neither is it of that force to purge, which groweth in these colder Countries, as that which commeth from Spaine, and other places.

#### The Names.

The name *Cnicus* is deriued from the Greekes, and *Carthamus* from the Arabians, yet still *sativus* is added vnto it, to shew it is no wilde, but a manured plant, and sowne euery where that wee know. Of some it is called *Crocus hortensis*, and *Saracenicus*, from the Italians which so call it. We call it in English Bastard Saffron, Spanish Saffron, and Catalonia Saffron.

#### The Vertues.

The flowers are vsed in colouring meates, where it groweth beyond Sea, and also for the dying of Silkes: the kernels of the seede are onely vsed in Physicke with vs, and serueth well to purge melancholicke humours.

## Chap. LXXVII.

### *Cardus.* Thistles.

YOu may somewhat maruaile, to see mee curious to plant Thistles in my Garden, when as you might well say, they are rather plagues then pleasures, and more trouble to weede them out, then to cherish them vp, if I made therein no distinction or choise; but when you haue viewed them well which I bring in, I will

then abide your censure, if they be not worthy of some place, although it be but a corner of the Garden, where something must needes be to fill vp roome. Some of them are smooth, and without prickes at all, some at the heads onely, and some all ouer; but yet not without some especiall note or marke worthy of respect: Out of this discourse I leaue the Artichoke, with all his kindes, and reserue them for our Kitchin Garden, because (as all know) they are for the pleasure of the taste, and not of the smell or sight.

1. *Acanthus sativus.* Garden Beares breech.

The leaues of this kinde of smooth thistle (as it is accounted) are almost as large as the leaues of the Artichoke, but not so sharp pointed, very deeply cut in and gashed on both edges, of a sad green & shining colour on the vpperside, and of a yellowish green vnderneath, with a great thicke rib in the middle, which spread themselues about the root, taking vp a great deale of ground. After this plant hath stood long in one place, and well defended from the iniury of the cold, it sendeth forth from among the leaues one or more great and strong stalkes, three or foure foote high, without any branch at all, bearing from the middle to the top many flowers one aboue another, spike-fashion round about the stalke, with smaller but not diuided greene leaues at euery flower, which is white, and fashioned somewhat like vnto a gaping mouth; after which come broad, flat, thicke, round, browniſh yellow seede (as I haue well obserued by them haue beene sent me out of Spaine, and which haue sprung vp, and doe grow with me; for in our Countrey I could neuer obserue any seede to haue growne ripe) the rootes are composed of many great and thicke long strings, which spread farre in and vnder the ground, somewhat darkish on the outside, and whitish within, full of a clammy moisture (whereby it sheweth to haue much life) and doe endure our Winters, if they be not too much exposed to the sharpe violence thereof, which then it will not endure, as I haue often found by experience.

2. *Acanthus siluestris.* Wilde or prickly Beares breech.

This prickly Thistle hath diuers long greenish leaues lying on the ground, much narrower then the former, but cut in on both sides, thicke set with many white prickes and thornes on the edges: the stalke riseth not vp so high, bearing diuers such like thornie leaues on them, with such a like head of flowers on it as the former hath: but the seede hereof (as it hath come to vs from Italy and other places, for I neuer saw it beare seed here in this Country) is blacke and round, of the bignesse of a small pease: the roote abideth reasonable well, if it be defended somewhat from the extremity of our Winters, or else it will perish.

3. *Eringium Pannonicum siue Montanum.* Hungary Sea Holly.

The lower leaues of this Thistle that lye on the ground, are somewhat large, round, and broad, hard in handling, and a little snipt about the edges, euery one standing vpon a long foote-stalke: but those that growe vpon the stalke, which is stiffe, two or three foote high, haue no foote-stalke, but encompasse it, two being set at euery ioynt, the toppe whereof is diuided into diuers branches, bearing small round rough heads, with smaller and more prickly leaues vnder them, and more cut in on the sides then those belowe: out of these heads rise many blew flowers, the foote-stalkes of the flowers, together with the toppes of the branches, are likewise blew and transparent, or shining.

*Flore albo.* We haue another of this kinde, the whole toppes of the stalkes, with the heads and branches, are more white then blew: the seede contained in these heads are white, flat, and as it were chaffie: the roote is great and whitish, spreading farre into many branches, and somewhat sweete in taste, like the ordinary Sea Holly rootes.

4. *Carduus mollis.* The gentle Thistle.

The leaues of this soft and gentle Thistle that are next vnto the ground, are greene:

1 *Acanthus sativus*, Garden Beares breech. 2 *Acanthus silvestris*. Wilde Beares breech. 3 *Eringium Pannonicum* Mountaine Sea Holly. 4 *Carlina humilis*. The lowe Carline Thistle. 5 *Carduus sphærocephalus maior*. The greater Globe-Thistle. 6 *Carduus sphærocephalus minor*. The lesser Globe-Thistle. 7 *Carduus Eriocæphalus*. The Friers crowne. 8 *Fraxinella*, Bastard Dittanie.

1. Acanthus mollis, L. mid. dict.
2. A. spinosus, L. mid. dict.

on the vpperside, and hoary vnderneath, broad at the bottome, somewhat long pointed, and vneuenly notched about the edges, with some soft hairie prickles, not hurting the handler, euery one standing vpon a short foote-stalke; those that growe about the middle stalke are like the former, but smaller and narrower, and those next the toppe smallest, where it diuideth it selfe into small branches, bearing long and scaly heads, out of which breake many reddish purple threads: the seede is whitish and hard, almost as great as the seede of the greater Centory: the roote is blackish, spreading vnder the ground, with many small fibres fastened vnto it, and abideth a great while.

### 5. *Carlina humilis.* The lowe Carline Thistle.

This lowe Thistle hath many iagged leaues, of a whitish greene colour, armed with small sharp white prickles round about the edges, lying round about the root vpon the ground, in the middle whereof riseth vp a large head, without any stalke vnder it, compassed about with many small and long prickly leaues, from among which the flower sheweth it selfe, composed of many thin, long, whitish, hard shining leaues, standing about the middle, which is flat and yellow, made of many thrums or threads like small flowers, wherein lye small long seede, of a whitish or siluer colour: the roote is somewhat aromaticall, blackish on the outside, small and long, growing downewards into the ground. There is another of this kinde that beareth a higher stalke, and a redder flower, but there is a manifest difference betweene them.

### 6. *Carduus Spherocephalus sive Globosus maior.* The greater Globe Thistle.

The greatest of these beautifull Thistles, hath at the first many large and long leaues lying on the ground, very much cut in and diuided in many places, euen to the middle ribbe, set with small sharpe (but not very strong) thornes or prickles at euery corner of the edges, greene on the vpperside, and whitish vnderneath: from the middle of these leaues riseth vp a round stiffe stalke, three foote and a halfe high, or more, set without order with such like leaues, bearing at the toppe of euery branch a round hard great head, consisting of a number of sharpe bearded huskes, compact or set close together, of a blewish greene colour, out of euery one of which huskes start small whitish blew flowers, with white threads in the middle of them, and rising aboue them, so that the heads when they are in full flower, make a fine shew, much delighting the spectators: after the flowers are past, the seede encreaseth in euery one, or the most part of the bearded huskes, which doe still hold their round forme, vntill that being ripe it openeth it selfe, and the huskes easily fall away one from another, containing within them a long whitish kernell: the roote is great and long, blackish on the outside, and dyeth euery yeare when it hath borne seede.

### 7. *Carduus Globosus minor.* The lesser Globe Thistle.

The lesser kinde hath long narrow leaues, whiter then the former, but cut in and gashed on the edges very much with some small prickes on them; the stalke is not halfe so long, nor the heads halfe so great, but as round, and with as blew flowers as the greater: this seldome giueth ripe seede, but recompenseth that fault, in that the roote perisheth not as the former, but abideth many yeares.

### 8. *Carduus Eriocephalus sive Tomentosus.* The Friers Crowne.

This woolly Thistle hath many large and long leaues lying on the ground, cut in on both sides into many diuisions, which are likewise somewhat vnequally cut in or diuided againe, hauing sharpe white prickles at euery corner of the diuisions, of a dead or sad greene colour on the vpperside, and somewhat woolly withall, and grayish vnderneath: the stalke is strong and tall, foure or fiue foote high at the least, branching out into diuers parts, euery where beset with such like leaues as growe below; at the toppe of euery branch there breaketh out a great whitish round prickly head, flattish at the toppe, so thicke set with wooll, that the prickles seeme but small spots or haires, and

and doth so well resemble the bald crowne of a Frier, not onely before it be in flower, but especially after it hath done flowring, that thereupon it deseruedly receiued the name of the Friers Crowne Thistle: out of these heads riseth forth a purple thrumme, such as is to be seene in many other wilde Thistles, which when they are ripe, are full of a flockie or woolly substance, which breake at the toppe shedding it, and the seede which is blackish, flat, and smooth: the roote is great and thicke, enduring for some yeares, yet sometimes perishing, if it be too much exposed to the violence of the frosts in Winter.

### The Place.

The first groweth naturally in Spaine, Italy, and France, and in many other hot Countries, and growe onely in Gardens in these colder climates, and there cherished for the beautifull aspect both of the greene plants, and of the stalkes when they are in flower. The Carline Thistle is found both in Germany and Italy in many places, and as it is reported, in some places of the West parts in England. The others are found some in France, some in Hungary, and on the Alpes, and the last in Spaine.

### The Time.

They doe all flower in the Summer moneths, some a little earlier or later then others.

### The Names.

The first is called *Acanthus sativus* (because the other that is prickly, is called *siluestris* or *spinosus*) and *Branca vrsina*; In English, Branck vrsine, and Beares breech. The third is called *Eringium montanum, Alpinum*, and *Pannonicum latifolium*: In English, Mountaine or Hungary Sea Holly. The fourth is called *Carduus mollis*, The gentle Thistle, because it hath no harmfull prickles, although it seeme at the first shew to be a Thistle. The fifth is called of diuers *Chamaleo albus*, and *Carlina*, as if they were both but one plant; but Fabius Columna hath in my iudgement very learnedly descided that controuersie, making *Carlina* to be *Ixine* of Theophrastus, and *Chamaleo* another differing Thistle, which Gaza translateth *Vernilago*. We call it in English, The Carline Thistle. The other haue their names in their titles, as much as is conuenient for this discourse.

### The Vertues.

The first hath alwaies been vsed Physically, as a mollifying herbe among others of the like slimie matter in Glisters, to open the body; yet Lobel seemeth to make no difference in the vse of them both (that is, the prickly as well as the smooth.) The Carline Thistle is thought to bee good against poysons and infection. The rest are not vsed by any that I know.

## CHAP. LXXVIII.

### *Fraxinella.* Bastard Dittany.

HAuing finished those pleasing Thistles, I come to other plants of more gentle handling, and first bring to your consideration this bastard Dittany, whereof there are found out two especiall kindes, the one with a reddish, the other with a whitish flower, and each of these hath his diuersity, as shall be presently declared.

1. *Fraxinella flore rubente.* Bastard Dittany with a reddish flower.

This goodly plant riseth vp with diuers round, hard, browniſh stalkes, neare two foote

foote high, the lower parts whereof are furnished with many winged leaues, somewhat like vnto Liquerice, or a small young Ashe tree, consisting of seuen, nine, or eleuen leaues set together, which are somewhat large and long, hard and rough in handling, of a darkish greene colour, and of an vnpleasant strong resinous sent: the vpper parts of the stalkes are furnished with many flowers, growing spike fashion, at certaine distances one aboue another, consisting of fiue long leaues a peece, whereof foure that stand on the two sides, are somewhat bending vpwards, and the fift hanging downe, but turning vp the end of the leafe a little againe, of a faint or pale red colour, striped through euery leafe with a deeper red colour, and hauing in the middle a tassell of fiue or six long purplish threds, that bowe downe with the lower leafe, and turne vp also the ends againe, with a little freese or thrume at the ends of euery one: after the flowers are past, arise hard, stiffe, rough, clammy huskes, horned or pointed at the end, foure or fiue standing together, somewhat like the seede vessels of the Wolfes-banes, or Colombines, but greater, thicker and harder, wherein is contained round shining blacke seede, greater then any Colombine seede by much, and smaller then Peony seede: the roote is white, large, and spreading many wayes vnder ground, if it stand long: the whole plant, as well roots as leaues and flowers, are of a strong sent, not so pleasing for the smell, as the flowers are beautifull to the sight.

2. *Fraxinella flore rubro.* Bastard Dittaine with a red flower.

This differeth not from the former eyther in roote, leafe or flower for the forme, but that the stalkes and leaues are of a darker greene colour, and that the flowers are of a deeper red colour, (and growing in a little longer spike) wherein the difference chiefly consisteth, which is sufficient to distinguish them.

3. *Fraxinella flore albo.* Bastard Dittanie with a white flower.

The white flowred *Fraxinella* hath his leaues and stalkes of a fresher greene colour then any of the former; and the flowers are of a pure white colour, in forme differing nothing at all from the other.

4. *Fraxinella flore albo cæruleo.*
Bastard Dittanie with an ash coloured flower.

The colour of the flower of this *Fraxinella* onely putteth the difference betweene this, and the last recited with a white flower: for this beareth a very pale, or whitish blew flower, tending to an ash colour.

## The Place.

All these kindes are found growing naturally, in many places both of Germany, and Italie: and that with the white flower, about Franckford, which being sent me, perished by the way by long and euill carriage.

## The Time.

They flower in Iune and Inly, and the seede is ripe in August.

## The Names.

The name *Fraxinella* is most generally imposed on those plants, because of the resemblance of them vnto young Ashes, in their winged leaues. Yet some doe call them *Dictamus albus*, or *Dictamnus albus*, and *Diptamus albus*, as a difference from the *Dictamnus Creticus*, which is a farre differing plant. Some would haue it to be *Tragium* of Dioscorides, but beside other things wherein this differeth from *Tragium*, this yeeldeth no milkie iuice, as Dioscorides saith *Tragium* doth: We in English doe eyther call it *Fraxinella*, or after the other corrupted name of *Dictamus*, Bastard Dittanie.

The

### The Vertues.

It is held to be profitable against the stingings of Serpents, against contagious and pestilent diseases, to bring downe the feminine courses, for the paines of the belly and the stone, and in Epilepticall diseases, and other cold paines of the braines: the roote is the most effectuall for all these, yet the seede is sometimes vsed.

## CHAP. LXXIX.

### *Legumina.* Pulse.

IF I should describe vnto you all the kindes of Pulse, I should vnfold a little world of varieties therein, more knowne and found out in these dayes, then at any time before, but that must bee a part of a greater worke, which will abide a longer time before it see the light. I shall only select those that are fit for this garden, and set them downe for your consideration. All sorts of Pulse may be reduced vnder two generall heads, that is, of Beanes and Pease, of each whereof there is both tame and wilde: Of Beanes, besides the tame or vsuall garden Beane, and the French or Kidney Beane, (whereof I meane to entreate in my Kitchen garden, as pertinent thereto) there is the Lupine or flat Beane, whereof I meane to entreate here, and the blacke Beane and others which must bee reserued for the Physicke Garden. And of the kindes of Pease some are fit for this Garden, (whereunto I will adioyne two or three other plants as neerest of affinitie, the flowers of some, and the fruit of others being delightfull to many, and therefore fit for this garden) some for the Kitchen, the rest for the Physicke garden. And first of Lupines or flat Beanes, accepted as delightfull to many, and therefore fit for this garden.

### 1. *Lupinus sativus albus.* The white garden Lupine.

The garden Lupine riseth vp with a great round stalke, hollow and somewhat woolly, with diuers branches, whereon grow vpon long footestalkes many broade leaues, diuided into seuen or nine parts, or smaller leaues, equally standing round about, as it were in a circle, of a whitish greene colour on the vpperside, and more woolly vnderneath: the flowers stand many together at seuerall ioynts, both of the greater stalke, and the branches, like vnto beanes, and of a white colour in some places, and in others of a very bleake blew tending to white: after the flowers are past, there come in their places, long, broade, and flat rough cods, wherein are contained round and flat seede, yellowish on the inside, and couered with a tough white skin, and very bitter in taste: the rootes are not very great, but full of small fibres, whereby it fasteneth it selfe strongly in the ground, yet perisheth euery yeare, as all the rest of these kindes doe.

### 2. *Lupinus caruleus maximus.* The greater blew Lupine.

The Stemme or stalke of this Lupine is greater then the last before recited, as also the leaues more soft and woolly, and the flowers are of a most perfect blew colour, with some white spots in the middle: the long rough greenish cods are very great and large, wherein are contained hard, flat and round seede, not so white on the outside as the former, but somewhat yellower, greater also, and more rough or hard in handling.

### 3. *Lupinus caruleus minor.* The lesser blew Lupine.

This kinde of wilde Lupine differeth not in the forme of leafe or flower from the former, but only that it is much smaller, the leaues are greener, and haue fewer diuisions in them: the flower is of as deepe a blew colour as the last; the cods likewise are small and long, containing small round seede, not so flat as the former, but more

Minimus. discoloured or spotted on the outside, then the greater kinde is. There is a lesser kind then this, not differing in any thing from this, but that it is lesser.

### 4. *Lupinus flore luteo.* The yellow Lupine.

The yellow Lupine groweth not vsually so high, but with larger leaues then the small blew Lupine; the flowers grow in two or three rundles or tufts, round about the stalke and the branches at the ioynts, of a delicate fine yellow colour, like in fashion vnto the other kindes, being larger then the last, but nothing so large as the greater kindes, and of a fine small sent: the seede is round, and not flat, but much about the forme and bignesse of the small blew, or somewhat bigger, of a whitish colour on the outside, spotted with many spots.

#### The Place.

The first groweth in many places of Greece, and the Easterne Countries beyond it, where it hath beene anciently cherished for their foode, being often watered to take away the bitternesse. It groweth also in these Western parts, but still where it is planted. The great blew Lupine is thought to come from beyond the parts of Persia, in Caramania. The lesser blew is found very plentifully wilde, in many places both of Spaine and Italy. The last hath beene brought vs likewise out of Spaine, whereas it is thought it groweth naturally. They all grow now in the gardens of those, that are curious louers of these delights.

#### The Time.

They flower in Summer, and their seede is ripe quickly after.

#### The Names.

They are generally called *Lupini*. Plautus in his time saith, they were vsed in Comedies in stead of money, when in any Scene thereof there was any shew of payment, and therefore he calleth them *Aurum Comicum*. And Horace hath this Verse,

*Nec tamen ignorant, quid distent æra Lupinis,*

to shew that counterfeit money (such as counters are with vs, or as these Lupines were vsed in those times) was easily knowne from true and currant coine. In English wee vsually call them after the Latine name, Lupines; and some after the Dutch name, Figge-beanes, because they are flat and round as a Figge that is pressed; and some Flat-beanes for the same reason. Some haue called the yellow Lupine, Spanish Violets: but other foolish names haue beene giuen it, as Virginia Roses, and the like, by knauish Gardiners and others, to deceiue men, and make them beleeue they were the finders out, or great preseruers of rarities, of no other purpose, but to cheate men of their money: as you would therefore auoyde knaues and deceiuers, beware of these manner of people, whereof the skirts of our towne are too pitifully pestered.

#### The Vertues.

The first or ordinary Lupine doth scoure and cleanse the skin from spots, morphew, blew markes, and other discolourings thereof, beeing vsed eyther in a decoction or ponther. Wee seldome vse it in inward medicines, not that it is dangerous, but of neglect, for formerly it hath beene much vsed for the wormes, &c.

1 *Lupinus maior.* The great Lupine. 2 *Lupinus luteus.* The yellow Lupine. 3 *Lathyrus latifolius seu Pisum perenne.* Peaſe euerlaſting. 4 *Piſum quadratum.* The crimſon bloſſomd or ſquare Peaſe. 5 *Medica cochleata vulgaris* Snailes or Barbary buttons. 6 *Medica ſpinoſa.* Prickly Snailes. 7 *Medica ſpinoſa altera.* Another ſort of prickly Snailes. 8 *Medica ſelliculo lato.* Broad buttons or Snailes. 9 *Medica Lunata.* Halfe Moons. 10 *Hedyſarum clypeatum.* The red Sattinflower, or French Honyſuckle. 11 *Scorpioides minus.* The leſſer Caterpiller. 12 *Scorpioides maius.* The greater Caterpiller. 13 *Orobus Venetus.* Blew vpright Peaſe euerlaſting.



5. Medicago cochleata Spr. hist

1. *Lathyrus latifolius, ſiue Piſum perenne.* Peaſe euerlaſting.

This kinde of wilde Peaſe that abideth long, and groweth euery yeare greater then other, ſpringeth vp with many broade trayling branches, winged as it were on both the ſides, diuerſly diuided into other ſmaller branches, at the ſeuerall ioynts whereof ſtand two hard, not broad, but ſomewhat long greene leaues, and diuers twining claſpers, in ſundry places with the leaues, from betweene the branches and the leaues, at the ioynts towards the toppes, come forth diuers purpliſh peaſe like bloſſomes, ſtanding on a long ſtemme or ſtalke, very beautifull to behold, and of a pretty ſent or ſmell: after which come ſmall, long, thin, flat, hard skind cods, containing ſmall round blackiſh ſeede: the roote is great and thicke, growing downe deepe into the ground, of the thickneſſe ſometimes of a mans arme, blackiſh on the outſide, and whitiſh within, with ſome branches and a few fibres annexed thereunto.

2. *Orobus Venetus.* Blew vpright euerlaſting Peaſe.

This pretty kinde of Peaſe bloſſome beareth diuers ſlender, but vpright greene branches ſomewhat cornered, two foote high or thereabouts, hauing at ſeuerall diſtances on both ſides of them certaine winged leaues, ſet together vpon long foote-ſtalkes one againſt another, conſiſting of ſix or eight leaues, ſomewhat broade and pointed, and without any odde one at the end: at the ioynts toward the toppes, betweene the leaues and the ſtalkes, come forth many flowers ſet together at the end of a pretty long footeſtalke, of the faſhion of the former Peaſe bloſſome, but ſomewhat ſmaller, and of a purpliſh violet colour: after which come ſlender and long pointed pods rounder then they, wherein is contained ſmall round grayiſh peaſe: the roote is blacke, hard or woody, abiding after ſeede bearing as the former doth, and ſhooting afreſh euery yeare.

3. *Lathyrus annuus ſiliquis orobi.* Partie coloured Cichelings.

This ſmall Pulſe or wild Peaſe, hath two or three long ſlender winged branches, with ſmaller leaues theron then the former, and without any claſpers at all on them: the flowers ſtand ſingle, euery one by it ſelfe, or two at the moſt together, the middle leaues whereof that cloſe together are white, and the vpper leaues of a reddiſh purple colour: after which come long round flattiſh cods, bunched out in the ſeuerall places where the ſeedes lye, like vnto the cods of *Orobus* or the bitter Vetch, but greater: the roote is ſmall and dyeth euery yeare.

4. *Piſum quadratum.* The crimſon bloſſomd or ſquare codded Peaſe.

This pretty kinde of Pulſe might very well for the forme of the leaues, be referred to the kindes of *Lotus* or Trefoiles: but becauſe I haue none of that kindred to entreate of in this Worke, I haue thought fitteſt to place it here before the Medica's, becauſe both pods and ſeedes are like alſo. It hath three or foure ſmall weake ſtalkes, diuided into many branches, hauing two ſtalkes of leaues at euery ioynt, and three ſmall ſoft leaues ſtanding on a very ſmall ſtalke, comming from the ioynts: the flowers ſtand for the moſt part two together, of a perfect red or crimſon colour, like in forme almoſt vnto a Peaſe bloſſome; after which come long thicke and round cods, with two skinnes or filmes, running all along the cod at the backe or vpperſide, and two other ſuch like filmes, all along the belly or vnder ſide, which make it ſeeme foure ſquare, wherein there lye round diſcoloured Peaſe, ſomewhat ſmaller and harder then ordinary Peaſe: the roote is ſmall and periſheth euery yeare.

5. *Medica Cochleata vulgaris.* Snailes or Barbary buttons.

The plant that beareth theſe pretty toyes for Gentlewomen, is ſomewhat like vnto a Threeleafed graſſe or Trefoile, hauing many long trayling branches lying vpon the ground, whereon at diuers places are three ſmall greene leaues, ſet together at the end of a little footeſtalke, each of them a little ſnipt about the edges: at ſeuerall diſtances,

from

from the middle of these branches to the ends of them, come forth the flowers, two for the most part standing together vpon a little foot stalke, which are of a pale yellow colour, very small, and of the forme of a Pease blossome: after which come smooth heads, which are turned or writhen round, almost like a Snaile, hard and greene at the first, somewhat like a greene button (from the formes of both which came their names) but afterwards growing whiter, more soft and open, wherein lyeth yellowish round and flat seede, somewhat like vnto the Kidney beane: the roote is small and stringie, dying downe euery yeare, and must be new sowne in the spring, if you desire to haue it.

6. *Medica spinosa minor.* Prickly or thorny Snailes, or Buttons.

This kinde of *Medica* is in all things very like vnto the former, both in the long trayling branches, & three leaues alwaies growing together, but a little greater pale yellow flowers, and crooked or winding heads: but herein chiefly consisteth the difference, that this kinde hath his heads or buttons harder, a little greater, more closed together, and set with short and somewhat hard prickles, all the head ouer, which being pulled open, haue those prickles standing on each side of the filme or skinne, whereof the head consisteth, somewhat like vnto a fish bone, and in this kinde goeth all one way; in which are contained such like seedes for the forme, as are in the former, but great and blacke, and shining withall.

7. *Medica spinosa altera.* Small thorney Buttons, or Snailes.

This other kinde is also like vnto the last described in all other things, except in the heads or buttons, which are a little smaller, but set with longer and softer prickes vpon the filmes, and may easily bee discerned to goe both forwards and backewards, one enterlacing within another, wherein are contained such like flat and blacke shining seede, made after the fashion of a kidney, as are in the former, but somewhat smaller: the roote perisheth in like manner euery yeare.

8. *Medica lata.* Broade Buttons.

This kinde differeth not from the first in leafe or flower, the fruite onely hereof is broade and flat, and not so much twined as it.

9. *Medica Lunata.* Halfe Moones.

This is also a kinde of these Medicke fodders, hauing a trefoyle leafe and yellow flowers like the former sorts, but both somewhat larger, the chiefest difference consisteth in the head or fruite, which is broade and flat, and not twined like the rest, but abideth halfe closed, resembling a halfe Moone (and thereupon hath assumed both the Latine and English name) wherein is contained flat seede, kidney fashion like the former.

10. *Hedysarum clypeatum.* The red Sattin flower.

This red flowred Fitchling, hath many stalkes of winged faire greene leaues, that is, of many set on both sides a middle ribbe, whereof that at the end is the greatest of the rest: from the ioynts where the leaues stand, come forth pretty long small stalkes, bearing on them very many flowers, vp to the toppe one aboue another, of an excellent shining red or crimson colour, very like vnto Sattin of that colour, and sometimes of a white colour, (as Master William Coys, a Gentleman of good respect in Essex, a great and ancient louer and cherisher of these delights, and of all other rare plants, in his life time assured me, he had growing in his garden at Stubbers by North Okenden) which are somewhat large, and more closed together, almost flat and not open, as in most of the other sorts: after the flowers are past, there come rough, flat, round huskes, somewhat like vnto the old fashioned round bucklers without pikes, three or foure standing one vpon or aboue another, wherein are contained small

small browniſh ſeede: the roote periſheth the ſame yeare it beareth ſeede, for oftentimes it flowreth not the firſt yeare it is ſowne.

## 11. *Scorpioides maius & minus.*

### Great and ſmall Caterpillers.

Vnder one deſcription I comprehend both theſe ſorts of Scorpions graſſe, or Caterpillers, or Wormes, as they are called by many, whereof the greater hath been known but of late yeares; and ioyne them to theſe pulſes, not hauing a fitter place where to inſert them. It is but a ſmall low plant, with branches lying vpon the ground, and ſomewhat long, broad, and hard leaues theron, among which come forth ſmall ſtalkes, bearing at the end for the moſt part, two ſmall pale yellowiſh flowers, like vnto Tares or Vetches, but ſmaller, which turne into writhed or crooked tough cods; in the greater ſort they are much thicker, rounder and whiter, and leſſer wound or turned together then in the ſmaller, which are ſlenderer, more winding, yet not cloſing like vnto the Snailes, and blacker more like vnto a Caterpiller then the other, wherein are contained browniſh yellow ſeede, much like vnto a *Medica*: the rootes of both are ſmall and fibrous, periſhing euery yeare.

### The Place.

Theſe are found ſeuerally in diuers and ſeuerall places, but wee ſow and plant them vſually to furniſh our gardens.

### The Time.

They doe all flower about the moneths of Iune and Iuly, and their ſeede is ripe ſoone after: but the ſecond is earlier then the reſt.

### The Names.

The firſt is called *Clymenum* of Matthiolus, and *Lathyris* of Lobel and others: but *Lathyris* in Greeke is *Cataputia* in Latine, which is our Spurge, farre differing from this Pulſe; and therefore *Lathyrus* is more proper to diſtinguiſh them aſunder, that two plants ſo farre vnlike ſhould not bee called by one name: this is alſo called *Lathyrus latifolius*, becauſe there is another called *auguſtifolius*, that differeth from it alſo: It is moſt vſually called with vs, *Piſum perenne*, and in Engliſh Peaſe bloſſome, or Peaſe euerlaſting. The ſecond is called by Cluſius, *Orobus venetus*, becauſe it was ſent him from Venice, with another of the ſame kinde that bore white flowers; yet differeth but little or nothing from that kinde he found in Hungary, that I thinke the ſeuerall places of their growing only cauſe them to beare ſeuerall names, and to be the ſame in deede. Although I yeeld vnto Cluſius the Latine name which doth not ſufficiently content mee; yet I haue thought good to giue it a differing Engliſh name, according as it is in the title. The third, becauſe I firſt receiued it among other ſeeds from Spaine, I haue giuen it the name, as it is entituled. The fourth is called of ſome *Sandalida Cretica, & Lotus ſiliquoſus flore rubello, Lotus tetragonolobus, Piſum rubrum, & Piſum quadratum*: We vſually call it in Engliſh, Crimſon Peaſe, or ſquare Peaſe. The *Medica Cochleata* is called of Dodonæus *Trifolium Cochleatum*, but not iudged to be the true *Medica*. Wee call it in Engliſh, Medick fodder, Snailes Clauer, or as it is in the title, and ſo the reſt of the Medica's accordingly. The *Hedyſarum clypeatum* or *Securidaca* is called of Dodonæus *Onobrichis altera*, and we in Engliſh for the likeneſſe, The red Sattin flower, although ſome fooliſhly call it, the red or French Honyſuckle. The laſt is called by Lobel, *Scorpioides bupleurifolio*, I haue called it *minus*, becauſe the greateſt ſort which came to me out of Spaine was not knowne vnto him: in Engliſh they are generally called Caterpillers.

The

The Vertues.

The Medica's are generally thought to feede cattell fat much more then the Medow Trefoile, or Clauer grasse, and therefore I haue known diuers Gentlemen that haue plowed vp some of their pasture grounds, and sowen them with the seedes of some Medica's to make the experience. All the other sorts are pleasures to delight the curious, and not any way profitable in Physicke that I know.

## CHAP. LXXX.

### *Peonia.* Peonie.

THere are two principall kindes of Peonie, that is to say, the Male and the Female. Of the male kinde, I haue onely known one sort, but of the Female a great many; which are thus to be distinguished. The Male his leafe is whole, without any particular diuision, notch or dent on the edge, & his rootes long & round, diuided into many branches, somewhat like to the rootes of Gentian or Elecampane, and not tuberous at all. The Female of all sorts hath the leaues diuided or cut in on the edges, more or lesse, and hath alwaies tuberous rootes, that is, like clogs or Asphodill rootes, with many great thick round peeces hanging, or growing at the end of smaller strings, and all ioyned to the toppe of the maine roote.

1. *Peonia mas.* The Male Peonie.

The Male Peonie riseth vp with many browniſh stalkes, whereon doe grow winged leaues, that is, many faire greene, and sometimes reddish leaues, one set against another vpon a stalke, without any particular diuision in the leafe at all: the flowers stand at the toppes of the stalkes, consisting of fiue or six broade leaues, of a faire purplish red colour, with many yellow threds in the middle, standing about the head, which after riseth to be the seede vessels, diuided into two, three or foure rough crooked pods like hornes, which when they are ful ripe, open and turn themselues down one edge to another backeward, shewing within them diuers round black shining seede, which are the true seede, being full and good, and hauing also many red or crimson graines, which are lancke and idle, intermixed among the blacke, as if they were good seede, whereby it maketh a very pretty shew: the roots are great, thick and long, spreading in the ground, and running downe reasonable deepe.

2. *Peonia fœmina vulgaris flore simplici.*
The ordinary single Female Peonie.

This ordinary Female Peonie hath many stalkes, with more store of leaues on them then the Male kinde hath, the leaues also are not so large, but diuided or nicked diuersly on the edges, some with great and deepe, and others with smaller cuts or diuisions, and of a darke or dead greene colour: the flowers are of a strong heady sent, most vsually smaller then the male, and of a more purple tending to a murrey colour, with yellow thrumes about the head in the middle, as the male kinde hath: the heads or hornes with seed are like also but smaller, the seede also is blacke, but lesse shining: the rootes consist, as I said, of many thicke and short tuberous clogs, fastened at the ends of long strings, and all from the head of the roote, which is thicke and short, and tuberous also, of the same or the like sent with the male.

3. *Peonia fœmina vulgaris flore pleno rubro.*
The double red Peonie.

This double Peonie as well as the former single, is so frequent in euerie Garden of note, through euery Countrey, that it is almost labour in vaine

to deſcribe it: but yet becauſe I vſe not to paſſe ouer any plant ſo ſlightly, I will ſet down the deſcription briefly, in regard it is ſo common. It is very like vnto the former ſingle female Peony, both in ſtalkes and leaues, but that it groweth ſomewhat higher, and the leaues are of a freſher greene colour: the flowers at the tops of the ſtalkes are very large, thicke, and double (no flower that I know ſo faire, great, and double; but not abiding blowne aboue eight or ten daies) of a more reddiſh purple colour then the former female kinde, and of a ſweeter ſent: after theſe flowers are paſt, ſometimes come good ſeed, which being ſowne, bring forth ſome ſingle flowers, and ſome double: the rootes are tuberous, like vnto the former female.

4. *Paonia famina flore carneo ſimplici.* The ſingle bluſh Peony.

The ſingle bluſh Peony hath his ſtalkes higher, and his leaues of a paler or whiter greene colour then the double bluſh, and more white vnderneath (ſo that it is very probable it is of another kinde, and not riſen from the ſeede of the double bluſh, as ſome might thinke) with many veines, that are ſomewhat diſcoloured from the colour of the leafe running through them: the flowers are very large and ſingle, conſiſting of fiue leaues for the moſt part, of a pale fleſh or bluſh colour, with an eye of yellow diſperſed or mixed therewith, hauing many whitiſh threads, tipt with yellow pendents ſtanding about the middle head: the rootes are like the other female Peonies.

5. *Paonia famina flore pleno albicante.* The double bluſh Peony.

The double bluſh Peony hath not his ſtalkes ſo high as the double red, but ſomewhat lower and ſtiffer, bearing ſuch like winged leaues, cut in or diuided here and there in the edges, as all theſe female kindes are, but not ſo large as the laſt: the flowers are ſmaller, and leſſe double by a good deale then the former double red, of a faint ſhining crimſon colour at the firſt opening, but decaying or waxing paler euery day: ſo that after it hath ſtood long (for this flower ſheddeth not his leaues in a great while) it will change ſomewhat whitiſh; and therefore diuers haue ignorantly called it, the double white Peony: the ſeedes, which ſometimes it beareth, and rootes, are like vnto the former female kindes, but ſomewhat longer, and of a brighter colour on the outſide.

6. *Paonia famina Byzantina.* The ſingle red Peony of Conſtantinople.

This red Peony of Conſtantinople is very like in all things vnto the double red Peonie, but that the flowers hereof are ſingle, and as large as the laſt, and that is larger then either the ſingle female, or the male kinde, conſiſting of eight leaues, of a deeper red colour then either the ſingle or double Peonies, and not purpliſh at all, but rather of the colour of an ordinary red Tulipa, ſtanding cloſe and round together: the roots of this kinde haue longer clogs, and not ſo ſhort as of the ordinary female kinde, and of a paler colour on the outſide.

## The Place.

All theſe Peonies haue beene ſent or brought from diuers parts beyond the Seas; they are endenized in our Gardens, where wee cheriſh them for the beauty and delight of their goodly flowers, as well as for their Phyſicall vertues.

## The Time.

They all flower in May, but ſome (as I ſaid) abide a ſmall time, and others many weekes.

## The Names.

The name *Paonia* is of all the later Writers generally giuen to theſe plants, although they haue had diuers other names giuen by the elder Writers, as *Roſa ſatuina*, *Idæus dactylus*, *Aglaophotis*, and others, whereof to ſet downe

1 *Pæonia mas cum semine.* The male Peony & the seed. 2 *Pæonia fœmina Byzantina.* The female red Peony of Constantinople. 3 *Pæonia fœmina flore pleno vulgaris.* The ordinary double Peony. 4 *Pæonia flore pleno albicante.* The double white Peony. 5 *Helleborus vernus atrorubente flore.* The early white Ellebor with a darke red flower. 6 *Helleborus niger vernus.* The Christmas flower. 7 *Calceolus Mariæ.* Our Ladies Slipper.

downe the causes, reasons, and errours, were to spend more time then I intend for this worke. Wee call them in English, Peonie, and distinguish them according to their titles.

### The Vertues.

The male Peony roote is farre aboue all the rest a most singular approued remedy for all Epilepticall diseases, in English, The falling sickenesse (and more especially the greene roote then the dry) if the disease be not too inueterate, to be boyled and drunke, as also to hang about the neckes of the younger sort that are troubled herewith, as I haue found it sufficiently experimented on many by diuers. The seede likewise is of especiall vse for women, for the rising of the mother. The seede of the female kinde, as well as the rootes, are most vsually sold, and may in want of the other be (and so are generally) vsed.

## Chap. LXXXI.

### *Helleborus niger.* Beares foote.

There are three sorts of blacke Hellebor or Beares foote, one that is the true and right kinde, whose flowers haue the most beautifull aspect, and the time of his flowring most rare, that is, in the deepe of Winter about Christmas, when no other can bee seene vpon the ground: and two other that are wilde or bastard kindes, brought into many Gardens for their Physicall properties; but I will only ioyne one of them with the true kinde in this worke, and leaue the other for another.

### 1. *Helleborus niger verus.* The true blacke Hellebor, or Christmas flower.

The true blacke Hellebor (or Beare foote as some would call it, but that name doth more fitly agree with the other two bastard kindes) hath many faire greene leaues rising from the roote, each of them standing on a thicke round fleshly stiffe green stalke, about an hand breadth high from the ground, diuided into seuen, eight, or nine parts or leaues, and each of them nicked or dented, from the middle of the leafe to the pointward on both sides, abiding all the Winter, at which time the flowers rise vp on such short thicke stalkes as the leaues stand on, euery one by it selfe, without any leafe thereon for the most part, or very seldome hauing one small short leafe not much vnder the flower, and very little higher then the leaues themselues, consisting of fiue broad white leaues, like vnto a great white single Rose (which sometimes change to be either lesse or more purple about the edges, as the weather or time of continuance doth effect) with many pale yellow thrummes in the middle, standing about a greene head, which after groweth to haue diuers cods set together, pointed at the ends like hornes, somewhat like the seede vessels of the *Aconitum hyemale*, but greater & thicker, wherein is contained long, round, and blackish seede, like the seede of the bastard kindes: the rootes are a number of browniſh strings running downe deepe into the ground, and fastened to a thicke head, of the bignesse of a finger at the toppe manie times, and smaller still downewards.

### 2. *Helleboraster minor.* The lesser bastard blacke Hellebor, or Beare foote.

The smaller Beare foote is in most things like vnto the former true blacke Hellebor; for it beareth also many leaues vpon short stalkes, diuided into many leaues also, but each of them are long and narrow, of a blacker greene colour, snipt or dented on both edges, which feele somewhat hard or sharpe like prickes, and perish euery yeare, but rise againe the next Spring: the flowers hereof stand on higher stalkes, with some leaues on them also, although but very few, and are of a pale greene colour, like in

forme vnto the flowers of the former, but smaller, hauing also many greenish yellow threads or thrums in the middle, and such like heads or seede vessels, and blackish seed: the rootes are stringie and blackish like the former.

### The Place.

The first groweth onely in the Gardens of those that are curious, and delight in all sorts of beautifull flowers in our Countrey, but wilde in many places of Germany, Italy, Greece, &c.

The other groweth wilde in many places of England, as well as the other greater sort, which is not here described; for besides diuers places within eight or ten miles from London, I haue seen it in the Woods of Northamptonshire, and in other places.

### The Time.

The first of these plants doth flower in the end of December, and beginning of Ianuary most vsually, and the other a moneth or two after, and sometime more.

### The Names.

The first is called *Helleborus*, or *Elleborus niger verus*, and is the same that both Theophrastus and Dioscorides haue written of, and which was called *Melampodion*, of Melampus the Goateheard, that purged and cured the mad or melancholicke daughters of Prætus with the rootes thereof. Dodonæus calleth it *Veratrum nigrum primum*, and the other *secundum*: Wee call it in English, The true blacke Hellebor, or the Christmas flower, because (as I said) it is most commonly in flower at or before Christmas. The second is a bastard or wilde kinde thereof, it so nearely resembleth the true, and is called of most of the later Writers, *Pseudoelleborus niger minor*, or *Helleboraster minor*, for a distinction betweene it and the greater, which is not here described: and is called in English, The smaller or lesser Beare foote, and most vsed in Physicke, because it is more plentifull, yet is more churlish and strong in operation then the true or former kinde.

### The Vertues.

The rootes of both these kindes are safe medecines, being rightly prepared, to be vsed for all Melancholicke diseases, whatsoeuer others may feare or write, and may be without danger applied, so as care and skill, and not temerary rashnesse doe order and dispose of them.

The powder of the dryed leaues, especially of the bastard kinde, is a sure remedy to kill the wormes in children, moderately taken.

## Chap. LXXXII.

### *Elleborus albus.* White Ellebor or Neesewort.

THere are two sorts of great white Ellebors or Neeseworts, whereas there was but one kinde knowne to the Ancients; the other being found out of later dayes: And although neither of both these haue any beauty in their flowers, yet because their leaues, being faire and large, haue a goodly prospect, I haue inserted them in this place, that this Garden should not be vnfurnished of them, and you not vnacquainted with them.

1. *Elleborus albus vulgaris.* White Ellebor or Neesing roote.

The first great white Ellebor riseth at the first out of the ground, with a whitish greene great round head, which growing vp, openeth it selfe into many goodly faire large greene leaues, plaited or ribbed with eminent ribbes all along the leaues, compassing one another at the bottome, in the middle whereof riseth vp a stalke three foot high or better, with diuers such like leaues thereon, but smaller to the middle thereof; from whence to the toppe it is diuided into many branches, hauing many small yellowish, or whitish greene starre-like flowers all along vpon them, which after turne into small, long, three square whitish seede, standing naked, without any huske to containe them, although some haue written otherwise: the roote is thicke and reasonable great at the head, hauing a number of great white strings running downe deepe into the ground, whereby it is strongly fastened.

2. *Elleborus albus præcox sive atrorubente flore.*
The early white Ellebor with reddish flowers.

This other Ellebor is very like the former, but that it springeth vp a moneth at the least before it, and that the leaues are not fully so thicke or so much plaited, but as large or larger, and doe sooner perish and fall away from the plant: the stalke hereof is as high as the former, bearing such like starry flowers, but of a darke or blackish red colour: the seede is like the other: the roote hath no such head as the other (so farre as I haue obserued, both by mine own and others plants) but hath many long white strings fastened to the top, which is as it were a long bulbous scaly head, out of which spring the leaues.

## The Place.

The first groweth in many places of Germany, as also in some parts of Russia, in that aboundance, by the relation of that worthy, curious, and diligent searcher and preseruer of all natures rarities and varieties, my very good friend, Iohn Tradescante, often heretofore remembred, that, as hee said, a good ship might be loaden with the rootes hereof, which hee saw in an Island there.

The other likewise groweth in the vpland wooddy grounds of Germanie, and other the parts thereabouts.

## The Time.

The first springeth vp in the end or middle of March, and flowreth in Iune. The second springeth in February, but flowreth not vntill Iune.

## The Names.

The first is called *Elleborus albus*, or *Helleborus albus*, the letter *H*, as all Schollers know, being but *aspirationis nota*: and *Veratrum album flore viridante*, of some *Sanguis Herculis*. The other is called *Elleborus albus præcox*, and *flore atrorubente*, or *atropurpurante*. We call the first in English, White Ellebor, Neesewort, or Neesing roote, because the powder of the roote is vsed to procure neesing; and I call it the greater, in regard of those in the next Chapter. The other hath his name according to the Latine title, most proper for it.

## The Vertues.

The force of purging is farre greater in the roote of this Ellebor, then in the former; and therefore is not carelesly to bee vsed, without extreame danger; yet in contumatious and stubborne diseases it may bee vsed with good

good caution and aduice. There is a Syrupe or Oxymel made hereof in the Apothecaries shops, which as it is dangerous for gentle and tender bodies, so it may be very effectuall in stronger constitutions. Pausanias *in Phocicis*, recordeth a notable stratagem that Solon vsed in besieging the Citie of Cirrheus, *viz.* That hauing cut off the riuer Plistus from running into the Citie, he caused a great many of these rootes to be put into a quantity thereof, which after they had steeped long enough therein, and was sufficiently infected therewith, he let passe into the Citie againe: whereof when they had greedily drunke, they grew so weake and feeble by the superpurgation thereof, that they were forced to leaue their wals vnmand, and not guarded, whereby the Amphyctions their enemies became masters of their Citie. The like stratagems are set downe by diuers other Authors, performed by the helpe of other herbes.

## CHAP. LXXXIII.

### *Elleborine.* Small or wilde white Ellebor.

THe likenesse of the leaues of these plants, rather then any other faculty with the former white Ellebor, hath caused them to be called *Elleborine*, as if they were smaller white Ellebors. And I for the same cause haue ioyned them next, whereof there are found many sorts: One which is the greater kinde, is of greatest beauty; the other which are lesser differ not much one from another, more then in the colour of the flowers, whereof I will onely take three, being of the most beautie, and leaue the rest to another worke.

#### 1. *Helleborine vel Elleborine maior, sive Calceolus Mariæ.*
Our Ladies Slipper.

This most beautifull plant of all these kindes, riseth vp with diuers stalkes, a foote and a halfe high at the most, bearing on each side of them broad greene leaues, somewhat like in forme vnto the leaues of the white Ellebor, but smaller and not so ribbed, compassing the stalke at the lower end; at the tops of the stalkes come forth one, or two, or three flowers at the most, one aboue another, vpon small short foote-stalkes, with a small leafe at the foote of euery stalke: each of these flowers are of a long ouall forme, that is, more long then round, and hollow withall, especially at the vpper part, the lower being round and swelling like a belly: at the hollow part there are two small peeces like eares or flippets, that at the first doe couer the hollow part, and after stand apart one from another, all which are of a fine pale yellow colour, in all that I haue seene (yet it is said there are some found, that are more browne or tending to purple) there are likewise foure long, narrow, darke coloured leaues at the setting on of the flower vnto the stalke, wherein as it were the flower at the first standeth: the whole flower is of a pretty small sent: the seede is very small, very like vnto the seede of the *Orchides* or Satyrions, and contained in such like long pods, but bigger: the roots are composed of a number of strings enterlacing themselues one within another, lying within the vpper crust of the earth, & not spreading deep, of a darke browniſh colour.

#### 2. *Elleborine minor flore albo.*
The small or wilde white Ellebor with a white flower.

This smaller wilde white Ellebor riseth vp in the like manner vnto the former, and not much lower, bearing such like leaues, but smaller, and of a whiter greene colour, almost of the colour and fashion of the leaues of Lilly Conually; the top of the stalke hath many more flowers, but lesser, growing together, spike-fashion, with small short leaues at the stalke of euery flower, which consisteth of fiue small white leaues, with a small close hood in the middle, without any sent at all: the seede and seede vessels are like

like vnto the former, but smaller: the roots are many small strings, dispersing themselues in the ground.

3. *Elleborine minor flore purpurante.*

The small or wilde white Ellebor with blush flowers.

The leaues of this kinde are like vnto the last described, but somewhat narrower: the stalkes and flowers are alike, but smaller also, and of a pale purplish or blush colour, which causeth the difference.

## The Place.

The first groweth in very many places of Germany, and in other Countries also. It groweth likewise in Lancashire, neare vpon the border of Yorkeshire, in a wood or place called the Helkes, which is three miles from Ingleborough, the highest Hill in England, and not farre from Ingleton, as I am enformed by a courteous Gentlewoman, a great louer of these delights, called Mistris Thomasin Tunstall, who dwelleth at Bull-banke, neare Hornby Castle in those parts, and who hath often sent mee vp the roots to London, which haue borne faire flowers in my Garden. The second groweth in many places of England, and with the same Gentlewoman also before remembred, who sent me one plant of this kinde with the other. The last I haue not yet knowne to growe in England; but no doubt many things doe lye hid, and not obserued, which in time may bee discouered, if our Country Gentlemen and women, and others, in their seuerall places where they dwell, would be more carefull and diligent, and be aduertised either by themselues, or by others capable and fit to be imployed, as occasion and time might serue, to finde out such plants as growe in any the circuits or limits of their habitations, or in their trauels, as their pleasures or affaires leade them. And becausе ignorance is the chiefe cause of neglect of many rare things, which happen to their view at sometimes, which are not to be seene againe peraduenture, or not in many yeares after, I would heartily aduise all men of meanes, to be stirred vp to bend their mindes, and spend a little more time and trauell in these delights of herbes and flowers, then they haue formerly done, which are not onely harmlesse, but pleasurable in their time, and profitable in their vse. And if any would be better enformed, and certified of such things they know not, I would be willing and ready to my best skill to aduertise them, that shall send any thing vp to me where I dwell in London. Thus farre I haue digressed from the matter in hand, and yet not without some good vse I hope, that others may make of it.

## The Time.

The two first flower earlier then the last, and both the first about one time, that is, in the end of Aprill, or beginning of May. The last in the end of May, or in Iune.

## The Names.

The first is called *Elleborine recentiorum maior*, and *Calceolus Marie*: Of some thought to be *Cosmosandalos*, because it is *Sandali forma*. In English we call it our Ladies Slipper, after the Dutch name. The other two lesser kinds haue their names in their titles: I haue thought it fit to adde the title of small white Ellebors vnto these, for the forme sake, as is before said.

## The Vertues.

There is no vse of these in Physicke in our dayes that I know.

CHAP.

## Chap. LXXXIIII.

### *Lilium Conuallium.* Lilly Conually.

THe remembrance of the Conuall Lilly, spoken of in the precedent Chapter, hath caused me to insert these plants among the rest, although differing both in face and properties; but lest it should lose all place, let it keepe this. It is of two sorts, differing chiefly in the colour of the flowers, the one being white, and the other reddish, as shall be shewed in their descriptions following.

#### 1. *Lilium Conuallium flore albo.* The white Lilly Conually.

The white Conuall or May Lilly, hath three or foure leaues rising together from the roote, one enclosed within another; each whereof when it is open is long and broad, of a grayish shining greene colour, somewhat resembling the leaues of the former wilde Neesewort, at the side whereof, and sometime from the middle of them, riseth vp a small short naked foote-stalke, an hand breadth high or somewhat more, bearing at the toppe one aboue another many small white flowers, like little hollow bottles with open mouths, nicked or cut into fiue or six notches, turning all downewards one way, or on one side of the stalke, of a very strong sweete sent, and comfortable for the memory and senses, which turne into small red berries, like vnto Asparagus, wherein is contained hard white seede: the rootes runne vnder ground, creeping euery way, consisting of many small white strings.

#### 2. *Lilium Conuallium flore rubente.* May Lillies with red flowers.

This other May Lilly differeth neither in roote, leafe, nor forme of flower from that before, but onely in the colour of the flower, which is of a fine pale red colour, being in my iudgement not altogether so sweet as the former.

### The Place.

The first groweth aboundantly in many places of England. The other is a stranger, and groweth only in the Gardens of those that are curious louers of rarities.

### The Time.

They both flower in May, and the berries are ripe in August.

### The Names.

The Latines haue no other name for this plant but *Lilium Conuallium*, although some would haue it to be *Lilium vernum* of Theophrastus, and others *Oenanthe* of the same Author. Gesner thinketh it to be *Callionymus*. Lonicerus to be *Cacalia*, and Fuchsius to be *Ephemerum non lethale*: but they are all for the most part mistaken. We call it in English Lilly Conually, May Lilly, and of some Liriconfancie.

### The Vertues.

The flowers of the white kinde are often vsed with those things that help to strengthen the memory, and to procure ease to Apoplecticke persons. Camerarius setteth downe the manner of making an oyle of the flowers hereof, which he saith is very effectuall to ease the paines of the Goute, and such like diseases, to be vsed outwardly, which is thus: Hauing filled a glasse with the flowers, and being well stopped, set it for a moneths space in an Ants hill, and after being drayned cleare, set it by to vse.

## Chap. LXXXV.

### *Gentiana.* Gentian or Fell-wort.

THere are diuers sorts of Gentians or Fell-wortes, some greater, others lesser, and some very small; many of them haue very beautifull flowers, but because some are very suddenly past, before one would thinke they were blowne open, and others will abide no culture and manuring, I will onely set forth vnto you two of the greater sorts, and three of the lesser kindes, as fittest, and more familiarly furnishing our gardens, leauing the rest to their wilde habitations, and to bee comprehended in a generall Worke.

#### 1. *Gentiana maior flore flauo.* The great Gentian.

The great Gentian riseth vp at the first, with a long, round and pointed head of leaues, closing one another, which after opening themselues, lye vpon the ground, and are faire, long and broad, somewhat plaited or ribbed like vnto the leaues of white Ellebor or Neesewort, but not so fairely or eminently plaited, neyther so stiffe, but rather resembling the leaues of a great Plantane: from among which riseth vp a stiffe round stalke, three foote high or better, full of ioynts, hauing two such leaues, but narrower and smaller at euery ioynt, so compassing about the stalke at the lower end of them, that they will almost hold water that falleth into them: from the middle of the stalke to the toppe, it is garnished with many coronets or rundles of flowers, with two such greene leaues likewise at euery ioynt, and wherein the flowers doe stand, which are yellow, layd open like starres, and rising out of small greenish huskes, with some threds in the middle of them, but of no sent at all, yet stately to behold, both for the order, height and proportion of the plant: the seede is browne and flat, contained in round heads, somewhat like vnto the seede of the *Fritillaria*, or checkerd Daffodill, but browner: the rootes are great, thicke and long, yellow, and exceeding bitter.

#### 2. *Gentiana maior folio Asclepiadis.* Swallow-wort Gentian.

This kinde of Gentian hath many stalkes rising from the roote, neere two foote high, whereon grow many faire pale greene leaues, set by couples, with three ribs in euery one of them, and doe somewhat resemble the leaues of *Asclepias* or Swallow-wort, that is, broade at the bottome, and sharpe at the point: the flowers grow at the seuerall ioynts of the stalkes, from the middle vpwards, two or three together, which are long and hollow, like vnto a bell flower, ending in fiue corners, or pointed leaues, and folded before they are open, as the flowers of the Bindeweedes are, of a faire blew colour, sometimes deeper, and sometimes paler: the heads or seede vessels haue two points or hornes at the toppes, and containe within them flat grayish seed, like vnto the former, but lesse: the rootes hereof are nothing so great as the former, but are yellow, small and long, of the bignesse of a mans thumbe.

#### 3. *Gentiana minor Cruciata.* Crosse-wort Gentian.

This small Gentian hath many branches lying vpon the ground, scarce lifting themselues vpright, and full of ioynts, whereat grow vsually foure leaues, one opposite vnto another, in manner of a Crosse, from whence it tooke his name, in shape very like vnto *Saponaria* or Sopewort, but shorter, and of a darker greene colour: at the tops of the stalkes stand many flowers, thick thrusting together, and likewise at the next ioynt vnderneath, euery one of them standing in a darke blewish greene huske, and consisting of fiue small leaues, the points or ends whereof only appeare aboue the huskes wherein they stand, and are hardly to be seene, but that they are of a fine pale blew colour, and that many grow together: the seed is small and brown, hard, and somewhat

like

1 *Lilium Conuallium.* Liriconfancy or Lilly Conually. 2 *Gentiana maior.* The great Gentian. 3 *Gentianella verna.* Small Gentian of the Spring. 4 *Gentiana cruciata.* Crosslewort Gentian. 5 *Pneumonanthe seu Gentiana Autumnalis.* Autumne Gentian. 6 *Saponaria flore duplici.* Double flowred Sopewort. 7 *Plantago Rosea.* Rose Platane.

like vnto the seed of the Marian Violets, or Couentry bels: the roots are small and whitish, dispersing themselues diuersly in the ground, of as bitter a taste almost as the rest.

4. *Gentianella Verna.* Small Gentian of the Spring.

The small Gentian of the Spring hath diuers small hard greene leaues, lying vpon the ground, as it were in heads or tufts, somewhat broade below, and pointed at the end, with fiue ribs or veines therein, as conspicuous as in the former Gentians, among which riseth vp a small short stalke, with some smaller leaues thereon, at the toppe whereof standeth one faire, large, hollow flower, made bell fashion, with wide open brimmes, ending in fiue corners or diuisions, of the most excellent deepe blew colour that can be seene in any flower, with some white spots in the bottome on the inside: after the flower is past, there appeare long and round pods, wherein are contained small blackish seede: the rootes are small, long, pale yellow strings, which shoot forth here and there diuers heads of leaues, and thereby encrease reasonable well, if it finde a fit place, and ground to grow, or else will not be noursed vp, with all the care and diligence can be vsed: the whole plant is bitter, but not so strong as the former.

5. *Gentiana Autumnalis siue Pneumonanthe.*
Calathian Violet or Autumne Gentian.

This Gentian that flowreth in Autumne, hath in some places higher stalkes then in others, with many leaues thereon, set by couples as in other Gentians, but long and narrow, yet shewing the three ribbes or veines that are in each of them: the toppes of the stalkes are furnished euery one with a flower or two, of an excellent blew purple colour, ending in fiue corners, and standing in long huskes: the rootes are somewhat great at the top, and spreading into many small yellow strings, bitter as the rest are.

6. *Saponaria flore duplici.* Double flowred Sopeworte.

Vnto these kindes of Gentians, I must needes adde these following plants, for that the former is of some neere resemblance in leafe with some of the former. And because the ordinary Sopeworte or Bruiseworte with single flowers is often planted in Gardens, and the flowers serue to decke both the garden and the house; I may vnder the one describe them both: for this with double flowers is farre more rare, and of greater beautie. It hath many long and slender round stalkes, scarce able to sustaine themselues, and stand vpright, being ful of ioynts and ribbed leaues at them, euery one somewhat like a small Gentian or Plantane leafe: at the toppes of the stalkes stand many flowers, consisting of two or three rowes of leaues, of a whitish or pale purple colour, and of a strong sweet sent, somewhat like the smell of Iasmin flowers, standing in long and thicke pale greene huskes, which fall away without giuing any seede, as most other double flowers doe that encrease by the roote, which spreadeth within the ground, and riseth vp in sundry distant places like the single.

7 *Plantago Rosea.* Rose Plantane.

This other plant is in all things like vnto the ordinary Plantane or Ribworte, that groweth wilde abroade in many places, whose leaues are very large: but in stead of the long slender spike, or eare that the ordinary hath, this hath eyther a thicke long spike of small greene leaues vpon short stalkes, or else a number of such small greene leaues layd round-wise like vnto a Rose, and sometimes both these may be seene vpon one and the same roote, at one and the same time, which abide a great while fresh vpon the roote, and sometimes also giueth seede, especially from the more long and slender spikes.

The Place.

Some of these Gentians grow on the toppes of hils, and some on the sides and foote of them in Germany and other Countreyes: some of them also vpon barren heaths in those places, as also in our owne Countrey, especially

ly the Autumne Gentian, and as it is reported, the Vernall likewise. The single or ordinary Sopeworte is found wilde in many places with vs, but the double came to vs from beyond the Sea, and is scarce known or heard of in England. The Rose Plantaine hath beene long in England, but whether naturall thereof or no, I am not assured.

### The Time.

They flower for the most part in Iune and Iuly, but the small Gentian of the Spring flowreth somewhat earlier, and that of the Autumne in August and September.

### The Names.

*Gentiana* is the generall name giuen to the Gentians. We call them in English Gentian, Fellworte, Bitterwort, and Baldmoney. *Saponaria* taketh his name from the scouring qualitie it hath: Wee call it in English Sopewort, and in some places Bruisewort. Some haue thought it to bee *Struthium* of Dioscorides, or at least haue vsed it for the same causes, but therein they are greatly deceiued, as Matthiolus hath very well obserued thereon, and so is Dodonæus, that thought it to be *Alisma*. The Rose Plantaine is so called of the double spikes it carrieth.

### The Vertues.

The wonderfull wholsomnesse of Gentian cannot bee easily knowne to vs, by reason our daintie tastes refuse to take thereof, for the bitternesse sake: but otherwise it would vndoubtedly worke admirable cures, both for the liuer, stomacke and lunges. It is also a speciall counterpoison against any infection, as also against the violence of a mad dogges tooth: wilde Sopewort is vsed in many places, to scoure the countrey womens treen, and pewter vessels, and physically some make great boast to performe admirable cures in Hydropicall diseases, because it is diuretically, and in *Lue Venerea*, when other Mercuriall medicines haue failed. The Rose Plantaine no doubt hath the same qualities that the ordinary hath.

## CHAP. LXXXVI.

### *Campanula.* Bell-flowers.

VNder the title of Bell-flowers are to bee comprehended in this Chapter, not only those that are ordinarily called *Campanula*, but *Viola Mariana*, and *Trachelium* also, whereof the one is called Couentry, the other Canterbury Bells.

#### 1. *Campanula Persicifolio alba, vel cærulea.*
Peach-leafed Bell-flowers white or blew.

The Peach-leafed Bell-flower hath many tufts, or branches of leaues lying vpon the ground, which are long and narrow, somewhat like vnto the leafe of an Almond or Peach tree, being finely nicked about the edges, and of a sad greene colour, from among which rise vp diuers stalkes, two foote high or more, set with leaues to the middle, and from thence vpwards, with many flowers standing on seuerall small foote-stalkes, one aboue another, with a small leafe at the foote of euery one: the flowers stand in small greene huskes, being small and round at the bottome, but wider open at the brimme, and ending in fiue corners, with a three forked clapper in the middle, set about with some small threds tipt with yellow, which flowers in some plants are pure white

white, and in others of a pale blew or watchet colour, hauing little or no ſent at all: the ſeede is ſmall, and contained in round flat heads, or ſeede veſſels: the roote is very ſmall, white and threddy, creeping vnder the vpper cruſt of the ground, ſo that oftentimes the heat and drought of the Summer wil goe near to parch and wither it vtterly: it requireth therefore to be planted in ſome ſhadowie place.

### 2. *Campanula maior, ſiue Pyramidalis.*
The great or ſteeple Bell-flower.

This great Bell-flower hath diuers ſtalkes, three foote high or better, whereon grow diuers ſmooth, darke, greene leaues, broade at the bottome, and ſmall at the point, ſomewhat vneuenly notched about the edges, and ſtanding vpon longer footeſtalkes below then thoſe aboue: the flowers are blew, and in ſome white, not ſo great or large as the former, but neare of the ſame faſhion, growing thicker and more plentifully together, with ſmaller leaues among them, buſhing thicke below, and riſing ſmaller and thinner vp to the toppe, in faſhion of a *Pyramis* or ſpeere Steeple: the roote is thicke and whitiſh, yeelding more ſtore of milke being broken (as the leaues and ſtalks alſo doe) then any other of the Bell-flowers, euery one whereof doe yeelde milke, ſome more and ſome leſſe.

### 3. *Viola Mariana flore albido vel purpureo.*
Couentry Bels white or purple.

The leaues of Couentry Bels are of a pale or freſh greene colour, long, and narrow next vnto the bottome, and broader from the middle to the end, and ſomewhat round pointed, a little hairy all ouer, and ſnipt about the edges: the ſtalkes riſe vp the yeare after the ſowing, being ſomewhat hairy alſo, and branching forth from the roote, into diuers parts, whereon ſtand diuers leaues, ſmaller then the former, and of a darker greene colour: at the end of euery branch ſtand the flowers, in greene huskes, from whence come large, round, hollow Bels, ſwelling out in the middle, and riſing ſomewhat aboue it, like the necke of a pot, and then ending in fiue corners, which are either of a faire or faint white, or of a pale blew purpliſh colour, and ſometimes of a deeper purple or violet: after the flowers are paſt, there riſe vp great ſquare, or cornered ſeede veſſels, wherein is contained in diuers diuiſions, ſmall, hard, ſhining, browne, flat ſeeds: the roote is white, and being young as in the firſt yeares ſowing, is tender, and often eaten as other Rampions are; but the next yeare, when it runneth vp to ſeede, it groweth hard, and periſheth: ſo that it is to be continued by euery other yeares ſowing.

### 4. *Trachelium maius flore albo vel purpureo.*
Great Canterbury Bels white or purple.

The greater Canterbury Bels, or Throateworte, hath many large rough leaues, ſomewhat like vnto Nettle leaues, being broad and round at the bottome, and pointed at the end, notched or dented on the edges, and euery one ſtanding on a long footſtalk: among theſe leaues riſe vp diuers ſquare rough ſtalkes, diuided at the toppe into diuers branches, whereon grow the like leaues as grow below, but leſſer; toward the ends of the branches ſtand the flowers, mixed with ſome longer leaues, euery one in his ſeuerall huske, which are hollow, long and round, like a bell or cup, wide open at the mouth, and cut at the brimme into fiue corners, or diuiſions, ſomewhat leſſer then the Couentry Bels, in ſome of a pure white, and others of a faire deepe purple violet colour, and ſometimes paler: after the flowers are paſt, come ſmaller and rounder heades then in the former, containing flat ſeede, but blacker, and not ſo redde as the laſt: the roote is hard and white, diſperſing it ſelfe into many branches vnder ground, not periſhing euery yeare as the former (although it loſeth all the leaues in winter) but abiding many yeares, and encreaſing into diuers heades or knobs, from whence ſpring new leaues and branches.

5. *Trache-*

1 *Campanula persicifolia.* Peach leafed Bell-flower. 2 *Trachelium maius simplex.* Canterbury Bels. * *Trachelium flore duplici.* Double Canterbury Bels. 3 *Viola Mariana.* Couentry Bels. 4 *Trachelium Giganteum.* Giants Throatewort. 5 *Trachelium minus.* The lesser Throatewort. 6 *Trachelium Americanum siue Cardinalis planta.* The rich crimson Cardinals flower.

5. *Trachelium maius flore duplici albo & cæruleo.*
Canterbury Bels with double flowers both white and blew.

Of this kinde of Throateworte or Canterbury Bels, there is another sort, not differing in any thing from the former, but in the doubleneſſe of the flower: For there is of both the kindes, one that beareth double white flowers, and the other blew: Of each whereof I receiued plants from friends beyond the Sea, which grow well with me.

6. *Trachelium Giganteum flore purpurante.*
Pale purple Giants Throateworte.

This Bell-flower, although it hath a Gigantine name, yet did I neuer perceiue it in my Garden, to riſe vp higher then the former, the epithite beeing in my perſwaſion, only giuen for difference ſake: the leaues whereof are not ſo rough, but as large, and dented about the edges, ſomewhat larger pointed, and of a freſher greene colour: the ſtalkes beare ſuch like leaues on them, but more thinly or diſperſedly ſet, hauing a flower at the ſetting on of euery one of the leaues, from the middle vpwards, and are ſomewhat like the great Throateworte in forme, but of a pale or bleake reddiſh purple colour, turning the brims or corners a little backwards, with a forked clapper in the middle, ſufficient eminent and yellow: the ſeede hereof is white, and plentifull in the heads, which will abide all the winter vpon the ſtalkes, vntill all the ſeede being ſhed, the heads remaining ſeeme like torne rags, or like thin peeces of skin, eaten with wormes: the roote is great, thicke and white, abiding long without periſhing.

*Flore albo.* There is another which differeth not any thing but in the flower, which is white.

7. *Trachelium minus flore albo & purpureo.*
Small Throateworte or Canterbury Bells both white and purple.

The leſſer Throateworte hath ſmaller leaues, nothing ſo broade or hard as the former great kinde, but long, and little or nothing dented about the edges: the ſtalkes are ſquare and browniſh, if it beare purple flowers, and greene if it beare white flowers, which in forme are alike, and grow in a buſh or tuft, thicke ſet together, more then any of the former, and ſmaller alſo, being not much bigger then the flowers of the fielde, or garden Rampions: the roote is laſting, and ſhooteth afreſh euery yeare.

8. *Trachelium Americarum flore ruberrimo, ſiue Planta Cardinalis.*
The rich crimſon Cardinals flower.

This braue plant, from a white roote ſpreading diuers wayes vnder ground, ſendeth forth many greene leaues, ſpread round about the head thereof, each whereof is ſomewhat broade and long, and pointed at the end, finely alſo ſnipt about the edges: from the middle whereof ariſeth vp a round hollow ſtalke, two foote high at the leaſt, beſet with diuers ſuch leaues as grow below, but longer below then aboue, and branching out at the toppe aboundantly, euery branch bearing diuers greene leaues on them, and one at the foote of euery of them alſo, the toppes whereof doe end in a great large tuft of flowers, with a ſmall greene leafe at the foote of the ſtalke of euery flower, each footeſtalke being about an inch long, bearing a round greene huske, diuided into fiue long leaues or points turned downwards, and in the midſt of euery of them a moſt rich crimſon coloured flower, ending in fiue long narrow leaues, ſtanding all of them foreright, but three of them falling downe, with a long vmbone ſet as it were at the backe of them, bigger below, and ſmaller aboue, and at the toppe a ſmall head, being of a little paler colour then the flower, but of no ſent or ſmell at all, commendable only for the great buſh of ſo orient red crimſon flowers: after the flowers are paſt, the ſeede commeth in ſmall heads, cloſed within thoſe greene husks that held the flowers, which is very like vnto the ſeede veſſels of the *Viola Mariana*, or Couentry Bels, and is ſmall and browniſh.

The

### The Place.

All these Bell-flowers do grow in our Gardens, where they are cherished for the beautie of their flowers. The Couentry Bels doe not grow wilde in any of the parts about Couentry, as I am credibly informed by a faithfull Apothecary dwelling there, called Master Brian Ball, but are noursed in Gardens with them, as they are in other places. The last groweth neere the riuer of Canada, where the French plantation in America is seated.

### The Time.

They flower from May vntill the end of Iuly or August, and in the mean time the seed is ripe: But the Peache-leafed Bell-flowers, for the most part, flower earlier then the other.

### The Names.

The first is generally called *Campanula Persicifolia*, in English Peach-leafed Bell-flower. The second is called *Campanula maior*, *Campanula lactescens Pyramidalis*, and *Pyramidalis Lutetiana* of Lobel, in English, Great or Steeple Bell-flower. The third is vsually called *Viola Mariana*, and of some *Viola Marina*. Lobel putteth a doubt whether it be not *Medium* of Dioscorides, as Matthiolus and others doe thinke; but in my opinion the thickneſse of the roote, as the text hath it, contradicteth all the rest. We call it generally in English Couentry Bels. Some call it Marian, and some Mercuries Violets. The fourth and fift are called *Trachelium* of *Ceruicaria*, of some *Vvularia*, because many haue vsed it to good purpose, for the paines of the *Vvula*, or Throate: Yet there is another plant, called also by some *Vvularia*, which is *Hippoglossum*, Horse tongue, or Double tongue. The sixt hath his title to descipher it out sufficiently, as is declared. The seuenth is called *Trachelium minus*, and *Cerviaria minor*, of some *Saponaria altera*; in English, Small Throateworte, or Small Canterbury Bels. The last hath his name in the title, as it is called in France, from whence I receiued plants for my Garden with the Latine name: but I haue giuen it in English.

### The Vertues.

The Peach-Bels as well as the others may safely bee vsed in gargles and lotions for the mouth, throate, or other parts, as occasion serueth. The rootes of many of them, while they are young, are often eaten in sallets by diuers beyond the Seas.

## CHAP. LXXXVII.

### *Campana Cærulea sive Convolvulus Cæruleus.*
### Blew Bell flowers, or blew Bindeweede.

THere are two other kindes of Bell-flowers, much differing from the Tribe or Familie of the former, because of their climbing or winding qualitie, which I must needes place next them, for the likenesse of the flowers, although otherwise they might haue beene placed with the other clamberers that follow. Of these there is a greater, and a lesser, and of each likewise some difference, as shall be declared.

1. *Convolvulus cæruleus maior rotundifolius.*
The greater blew Bindweede, or Bell-flower with round leaues.

This goodly plant riseth vp with many long and winding branches, whereby it climbeth and windeth vpon any poles, herbes, or trees, that stand neare it within a great compasse, alwaies winding it selfe contrary to the course of the Sunne: on these branches doe growe many faire great round leaues, and pointed at the end, like vnto a Violet leafe in shape, but much greater, of a sad greene colour: at the ioynts of the branches, where the leaues are set, come forth flowers on pretty long stalkes, two or three together at a place, which are long, and pointed almost like a finger, while they are buds, and not blowne open, and of a pale whitish blew colour, but being blowne open, are great and large bels, with broad open mouths or brims ending in fiue corners, and small at the bottome, standing in small greene huskes of fine leaues: these flowers are of a very deepe azure or blew colour, tending to a purple, very glorious to behold, opening for the most part in the euening, abiding so all the night and the next morning, vntill the Sunne begin to growe somewhat hot vpon them, and then doe close, neuer opening more: the plant carrieth so many flowers, if it stand in a warme place, that it will be replenished plentifully, vntill the cold ayres and euenings stay the luxury thereof: after the flowers are past, the stalkes whereon the flowers did stand, bend downwards, and beare within the huskes three or foure blacke seedes, of the bignesse of a Tare or thereabouts: the rootes are stringy, and perish euery yeare.

2. *Convolvulus trifolius sive hederaceus purpureus.*
The greater purple Bindeweede, or Bell-flower with cornered leaues.

The growing and forme of this Bindeweede or Bell-flower, is all one with the former, the chiefest differences consisting in the forme of the leafe, which in this is three cornered, like vnto an Iuie leafe with corners; and in the flower, which is of a deeper blew, tending more to a deepe purple Violet, and somewhat more reddish in the fiue plaites of each flower, as also in the bottomes of the flowers.

3. *Convolvulus tennifolius Americanus.* The red Bell-flower of America.

Although this rare plant (because wee seldome haue it, and can as hardly keepe it) be scarce knowne in these cold Countries, yet I could not but make mention of it, to incite those that haue conueniencie to keepe it, to be furnished of it. It springeth vp at the first from the seede with two leaues, with two long forked ends, which abide a long time before they perish, betweene which riseth vp the stalke or stemme, branching forth diuers waies, being of a browniish colour, which windeth it selfe as the former great Bell-flower doth, whereon are set at seuerall ioynts diuers winged leaues, that is to say, many small narrow and long leaues set on both sides of the middle ribbe, and one at the end: from these ioynts arise long stalkes, at the ends whereof stand two or three small, long, hollow flowers, fashioned very like vnto the flowers of a Bindeweede, or the flowers of Tabacco, and ending in the like manner in fiue points, but not so much laide open, being of a bright red colour, plaited as the Bindeweedes or Bell-flowers before they be open, with some few threads in the middle, which turne into long pointed cods, wherein is contained long and blacke seede, tasting hot like Pepper: the roote is small and stringy, perishing euery yeare, and with vs will seldome come to flower, because our cold nights and frosts come so soone, before it cannot haue comfort enough of the Sun to ripen it.

4. *Convolvulus cæruleus minor Hispanicus.*
The Spanish small blew Bindeweede.

This small Bindeweede hath small long leaues, somewhat broader then the next that followeth, and not so broad as the common small Bindeweede (that groweth

euery

euery where wilde on the bankes of fields abroad) set vpon the small trayling branches, which growe aboue two or three foote high: from the middle of these branches, and so vnto the toppes of them, come forth the flowers at the ioynts with the leaues, folded together at the first into fiue plaites, which open into so many corners, of a most excellent faire skie coloured blew (so pleasant to behold, that often it amazeth the spectator) with white bottomes, and yellowish in the middle, which turne into small round white heads, wherein are contained small blackish cornered seede, somewhat like the former, but smaller: the roote is small and threddy, perishing as the former euery yeare: this neuer windeth it selfe about any thing, but leaneth by reason of the weakenesse of the branches, and dyeth euery yeare after seede time, and not to be sowne againe vntill the next Spring.

5. *Convolvulus purpureus Spicæfolius.* Lauander leafed Bindeweede.

This small purple Bindeweede, where it naturally groweth, is rather a plague then a pleasure, to whatsoeuer groweth with it in the fields; yet the beauty of the flower hath caused it to be receiued into Gardens, bearing longer and smaller leaues then the last, and such like small Bell-flowers, but of a sad purple colour: the roote is liuing, as the common kinds are, and springeth againe where it hath been once sowne, without feare of perishing.

## The Place.

The first two greater kindes haue beene sent vs out of Italy, but whether they had them from the East Indies, or from some of the Easterne Countries on this side, wee know not: but they thriue reasonable well in our Country, if the yeare be any thing kindly. The next came out of America, as his name testifieth. The lesser blew kinde groweth naturally in many places both of Spaine and Portugall (from whence I first receiued seedes from Guillaume Boel, heretofore remembred.) The last groweth wilde in the fields, about Dunmowe in Essex, and in many other places of our owne Countrey likewise.

## The Time.

The three first greater kindes flower not vntill the end of August, or thereabouts, and the seede ripeneth in September, if the colds and frosts come not on too speedily. The lesser kindes flower in Iune and Iuly.

## The Names.

The first is called of some *Campana Lazura*, as the Italians doe call it, or *Campana cærulea*, of others *Convolvulus cæruleus maior, siue Indicus*, and *Flos noctis*. Of some *Nil Auicennæ*. The second is called *Convolvulus trifolius*, or *hederaceus*, for the distinction of the leaues. In English wee call them eyther Great blew Bell-flowers, or more vsually, Great blew Bindeweedes. That of America is diuersly called by diuers. It is called *Quamoclit* of the Indians, and by that name it was sent to Ioachinus Camerarius out of Italy, where it is so called still, as Fabius Columna setteth it downe, and as my selfe also can witnesse it, from thence being so sent vnto mee: but Andræas Cæsalpinus calleth it, *Iasminum folio Millefolij*, supposing it to be a Iasmine. Camerarius saith, it may not vnfitly be called *Convolvulus tenuifolius*, accounting it a kinde of Bindeweede. Columna entituleth it *Convolvulus pennatus exoticus rarior*, and saith it cannot bee referred to any other kinde of plant then to the Bindeweedes. Hee that published the *Curæ posteriores* of Clusius, giueth it the name of *Iasminum Americanum*, which I would doe also, if I thought it might belong to that Family; but seeing the face and forme of the plant better agreeing with the Bindeweedes or Bell-flowers,

I haue

I haue (as you see) inserted it among them, and giuen it that name may bee most fit for it, especially because it is but an annuall plant. The lesser kindes haue their names sufficiently expressed in their titles.

The Vertues.

We know of no vse these haue in Physicke with vs, although if the first be *Nil* of Auicen, both he and Serapio say it purgeth strongly.

## Chap. LXXXVIII.

### *Stramonium.* Thorne-Apple.

VNto the Bell-flowers, I must adioyne three other plants, in the three seuerall Chapters following, for some affinity of the flowers: and first of the Thorne-Apples, whereof there are two especiall kindes, that is, a greater and a lesser, and of each some diuersity, as shall be set downe.

### 1. *Stramonium maius album.* The great white flowred Thorne-Apple.

The greater Thorne-Apple hath a great, strong, round greene stalke, as high as any man, if it be planted in good ground, and of the bignesse of a mans wrest almost at the bottome, spreading out at the toppe into many branches, whereon stand many very large and broad darke greene leaues, cut in very deeply on the edges, and hauing manie points or corners therein: the flowers come forth at the ioynts, betweene two branches towards the toppe of them, being very large, long, and wide open, ending in fiue points or corners, longer and larger then any other Bell-flowers whatsoeuer: after the flowers are past, come the fruit, which are thorny long heads, more prickly and greene then the lesser kindes, which being ripe openeth it selfe into three or foure parts, hauing a number of flat blackish seede within them: the roote is aboundant in fibres, whereby it strongly taketh hold in the ground, but perisheth with the first frosts; yet the seede that is shed when the fruit is ripe, commeth vp the next yeare.

### 2. *Stramonium maius purpureum.* The great purple flowred Thorne-Apple.

This purple Thorne-Apple is in largenesse of leaues, thicknesse and height of stalke, greatnesse and forme of flowers and fruit, euery way equall and correspondent vnto the former, the chiefe differences be these: the stalke is of a darke purple colour; the leaues are of a darker greene, somewhat purplish, and the flowers are of light purple or pale Doue colour, enclining to white, and whiter at the bottome.

### 3. *Stramonium minus seu Nux Metel flore albo.* The smaller Thorne-Apple with a white flower.

The smaller Thorne-Apple riseth vp with one round stalke, of the bignesse of a mans finger, and neuer much aboue two foote high with vs, bearing a few large, broad, smooth leaues thereon, without any branches at all, which are vneuenly rent or torne about the edges, with many ribs, and smaller veines running through them, yet lesser by much then the greater kinde: at the ioynts where the leaues stand, come forth long and large white flowers, with broad or wide open brims, folded together before their opening, as the other former Bell-flowers or Bindeweedes, but hauing their fiue corners more pointed or horned then either they, or the former Thorne-Apples: after the flowers are past, succeed small fruit, rounder and harder, set with harder, but blunt prickes then the former, wherein is contained brownish yellow flat seede,

sticking

1 *Convolvulus major cæruleus.* The greater blew Bindweed or Bell-flower. 2 *Convolvulus trifolius seu hederaceus.* The great purple Bindeweed. 3 *Convolvulus minor cæruleus Hispanicus.* The Spanish small blew Bindweed. 4 *Stramonium maius seu Pomum spinosum.* The great Thorne Apple. 5 *Datura seu Stramonium cæruleum.* The small Thorne-Apple. 6 *Stramonium flore duplici.* The double flowred Thorne-Apple. 7 *Stramonium flore geminato.* Double Thorne-Apple one out of another 8 *Tabacco latifolium.* Broad leafed Tabacco. 9 *Mirabilis Peruana.* The Meruaile of the world.

sticking to the inward pulpe: the roote is not very great, but full of strings, and quickly perisheth with the first frosts.

4. *Stramonium minus flore geminato purpurante.*
The small double flowred purple Thorne-Apple.

In the flower of this plant, consisteth the chiefest difference from the former, which is as large as the last, pointed into more hornes or corners, and beareth two flowers, standing in one huske, one of them rising out from the middle of the other, like vnto those kindes of Cowslips and Oxelips, called double, or Hose in hose, before described, which are of a pale purplish colour on the outside, and almost white within: the fruit is round like the last, and beareth such like seede, so that vntill it bee in flower, their difference can hardly bee discerned: this is more tender then the last, although euen it is so tender, that it seldome beareth ripe seede with vs.

*Flore duplici.* Sometimes (for I think it is not another kind) the flower will haue as it were double rowes of leaues, close set together, and not consisting of two, rising so distinctly one aboue another.

## The Place.

All these kindes haue been brought or sent vs out of Turkie and Egypt; but Garcias, and Christopherus Acosta, with others, affirme that they grow in the East Indies. The lesser kindes are very rare with vs, because they seldome come to maturity; and therefore we are still to seeke of new seede to sowe. The greater kindes are plentifull enough in our Gardens, and will well abide, and giue ripe fruit.

## The Time.

The smaller kindes flower later then the greater; and therefore their fruit are the sooner spoiled with the cold ayres, dewes, and frosts, that come at the latter end of the yeare: but the greater kinds neuer misse lightly to ripen.

## The Names.

Both the greater and smaller kindes are generally called *Stramonium*, *Stramonia*, *Pomum spinosum*, and *Datura*. Bauhinus vpon Matthiolus his Comentaries on Dioscorides, calleth it *Solanum fœtidum spinosum*. Some learned men haue referred it to *Nux Metel*, of the Arabian Authors. Wee call them generally in English, Thorne-Apples, and distinguish them by their titles of greater and lesser, single and double.

## The Vertues.

The East Indian lasciuious women performe strange acts with the seed (of the smaller kinde, as I suppose, or it may be of either) giuing it their husbands to drinke. The whole plant, but especially the seed, is of a very cold and soporiferous quality, procuring sleep and distraction of senses. A few of the seeds steeped and giuen in drinke, will cause them that take it to seem starke drunke or dead drunke, which fit will within a few houres weare away, and they recouer their senses againe, as a drunken man raysed after sleep from his wine. It may therefore (in my opinion) be of safe and good vse to one, that is to haue a legge or an arme cut off, or to be cut for the stone, or some other such like cure to be performed, to take away the sense of paine for the time of doing it; otherwise I hold it not fit to be vsed without great caution. But the greene leaues of the greater kindes (as also of the lesser, but that with vs they are not so plentifull) are by tryed experience, found to be excellent good for any scalded or burned part, as also to take away any hot inflammations, being made vp into a salue or ointment with suet, waxe, and rosin, &c. or with *Axungia*, that is, Hogs larde.

CHAP.

## CHAP. LXXXIX.

### *Tabacco.* Indian Henbane, or Tabacco.

THere hath beene formerly but three kindes of Tabacco knowne vnto vs, *two* of them called Indian, and the third Englisth Tabacco. In these later yeares, we haue had in our gardens about London (before the suppressing of the planting) three or foure other sorts at the least, and all of the Indian kinde, hauing some especiall difference, eyther in leafe, or flower, or both: And in regard the flowers of some of these carry a pretty shew, I shall only entreate of them, and not of the Englisth kind.

### *Tabacco latifolium.* Broade leafed Tabacco.

The great Indian Tabacco hath many very large, long, thicke, fat and faire greene leaues, standing foreright for the most part, and compassing the stalkes at the bottome of them, being somewhat pointed at the end: the stalke is greene and round, sixe or seuen foote high at sometimes, and in some places, in others not past three or foure foote high, diuided towards the toppe into many branches, with leaues at euery ioynt, and at the toppes of the branches many flowers, the bottomes hereof are long and hollow, and the toppes plaited or folded before they are open, but being open, are diuided sometimes into foure, or more vsually into fiue corners, somewhat like vnto other of the Bell-flowers, but lying a little flatter open, of a light carnation colour. The seede is very small and browne, contained in round heads, that are clammy while they are greene, and pointed at the end: the roote is great, whitish, and woody at the head, dispersing many long branches, and small fibres vnder the ground, whereby it is strongly fastened, but perisheth with our violent frosts in the winter, if it be left abroad in the garden, but if it be housed, or safely prouided for against the frostes, the rootes will liue, and spring afresh the next yeare.

There is of this kinde another sort, whose leaues are as large and long as the former, but thicker, and of a more dead greene colour, hanging downe to the ground-ward, and scarce any standing forth-right, as the former, vnlesse they bee very young: the flowers of this kinde are almost whole, without any great shew of corners at the brims or edges, in all other things there is no difference.

There is another, whose large and thicke flat leaues doe compasse the stalke at the bottome, and are as it were folded together one side vnto another: the flowers are of a deeper blush, or carnation colour, and with longer points and corners then in any of the former; and in these two things consisteth the difference from the others, and is called Verines Tabacco.

Another hath his leaues not so large and long as the first, and these haue short foote-stalkes, whereon they stand, and doe not compasse the stalke as the other doe: the flower hereof is like the first, but smaller, and of a little paler colour.

### *Tabacco angustifolium.* Narrow leafed Tabacco.

This kinde of Tabacco hath somewhat lower, and smaller stalkes, then any of the former: the leaues hereof are smaller and narrower, and not altogether so thicke, but more pointed, and euery one standing vpon a footstalke, an inch and a halfe long at the least: the flowers hereof stand thicker together, vpon the small branches, somewhat larger, of a deeper blush colour, and more eminent corners then in any the former: the seed and roots are alike, and perish in like manner, vnlesse it be brought into a cellar, or other such couert, to defend it from the extremitie of the Winter.

### The Place.

America or the West Indies is the place where all these kindes doe grow naturally, some in one place, and some in another, as in Peru, Trinidado,

Hispaniola, and almost in euery Iland and Countrey of the continent thereof: with vs they are cherished in gardens, as well for the medicinable qualities, as for the beauty of the flowers.

### The Time.

It flowreth in August, seldome before, and the seede is ripe quickly after. If it once sowe it selfe in a Garden, it will giue next year after young plants: but for the most part they will spring vp late, and therefore they that would haue them more early, haue sowen the seede vpon a bed of dung, and transplanted them afterwards.

### The Names.

This plant hath gotten many names. The Indians call it in some places *Petum*, in others *Picielt*, and *Perebecennu*, as Ouiedus and others doe relate. The Spaniards in the Indies first called it *Tabacco*, of an Iland where plenty of it grew. It hath in Christendome receiued diuers other names, as *Nicotiana*, of one Nicot a French man, who seeing it in Portugall, sent it to the French Queene, from whom it receiued the name of *Herba Regina*. Lobel calleth it *Sancta herba, & Sana sancta Indorum*. Some haue adiudged it to be an *Hiosciamus*, and therefore call it *Peruvianus*. The most vsuall name whereby we call it in English, is Tabacco.

### The Vertues.

The herbe is, out of question, an excellent helpe and remedy for diuers diseases, if it were rightly ordered and applyed, but the continuall abuse thereof in so many, doth almost abolish all good vse in any. Notwithstanding if men would apply their wits to the finding out of the vertues, I make no doubt but many strange cures would bee performed by it, both inward and outward. For outward application, a Salue made hereof (as is before recited of the Thorne apple leaues) cureth vlcers, and wounds of hard curation: And for inward helpes, a Syrupe made of the iuice and sugar, or honey, procureth a gentle vomit (but the dryed leafe infused in wine much more) and is effectuall in astmaticall diseases, if it bee carefully giuen. And likewise cleanseth cankers and fistulaes admirably, as hath beene found by late experience. The ashes of Tabacco is often vsed, and with good successe, for cuts in the hands, or other places, and for other small greene wounds.

## Chap. XC.

### *Mirabilia Peruviana.* The Meruaile of Peru.

THis plant yeeldeth in our Gardens fiue or sixe seuerall varieties of beautifull flowers, as pure white, pure yellow, pure red, white and red spotted, and red and yellow spotted. But besides these, I haue had some other sorts, among which was one, of a pale purple or peach colour: all which, comming vnto mee out of Spaine with many other, seedes in an vnkindly yeare (an early winter following a cold summer) perished with mee; yet I plainely might discerne by their leaues, and manner of growing, to be diuers from them that we now haue and keepe. I shall need therefore (because the chiefest difference consisteth almost in the flowers) to giue only one description of the plant, and therein shew the varieties as is before declared.

### *Admirabilis.* The Meruaile of the World.

The stalke of this meruellous plant is great and thick, bigger then any mans thumbe, bunched

bunched out or swelling at euery ioynt, in some the stalkes will bee of a faire greene colour, and those will bring white, or white and red flowers : in others they will bee reddish, and more at the ioynts, and those giue red flowers; and in some of a darker greene colour, which giue yellow flowers; the stalkes and ioynts of those that will giue red and yellow flowers spotted, are somewhat browniſh, but not so red as those that giue wholly red flowers: vpon these stalkes that spread into many branches, doe grow at the ioynts vpon seuerall footestalkes, faire greene leaues, broad at the stalke, and pointed at the end: at the ioynts likewise toward the vpper part of the branches, at the foote of the leaues, come forth seuerall flowers vpon short footestalkes, euery one being small, long and hollow from the bottome to the brimme, which is broade spread open, and round, and consist but of one leafe without diuision, like vnto a Bell flower, but not cornered at all: which flowers, as I said, are of diuers colours, and diuersly marked and spotted, some being wholly white, without any spot in them for the most part, through all the flowers of the plant; so likewise some being yellow, and some wholly red; some plants againe being mixed and spotted, so variably either white and red, or purple, (except here and there some may chance to be wholly white, or red or purple among the rest) or red and yellow through the whole plant, (except as before some may chance in this kinde to be eyther wholly red, or wholly yellow) that you shall hardly finde two or three flowers in a hundred, that will bee alike spotted and marked, without some diuersitie, and so likewise euery day, as long as they blow, which is vntill the winters, or rather autumnes cold blastes do stay their willing proneneſſe to flower: And I haue often also obserued, that one side of a plant will giue fairer varieties then another, which is most commonly the Easterne, as the more temperate and shadowie side. All these flowers doe open for the most part, in the euening, or in the night time, and so stand blowne open, vntill the next mornings sun beginne to grow warme vpon them, which then close themselues together, all the brims of the flowers shrinking into the middle of the long necke, much like vnto the blew Bindeweede, which in a manner doth so close vp at the sunnes warme heate: or else if the day be temperate and milde, without any sunne shining vpon them, the flowers will not close vp for the most part of that day, or vntill toward night: after the flowers are past, come seuerall seedes, that is, but one at a place as the flowers stood before, of the bigneſſe (sometimes) of small peaſe, but not so round, standing within the greene huskes, wherein the flowers stood before, being a little flat at the toppe, like a crowne or head, and round where it is fastened in the cup, of a blacke colour when it is ripe, but else greene all the while it groweth on the stalke, and being ripe is soone shaken downe with the wind, or any other light shaking: the roote is long and round, greater at the head, and smaller downwards to the end, like vnto a Reddish, spreading into two or three, or more branches, blackish on the outside and whitish within. These rootes I haue often preserued by art a winter, two or three (for they will perish if they be left out in the garden, vnleſſe it be vnder a house side) because many times, the yeare not falling out kindely, the plants giue not ripe seede, and so we should be to seeke both of seede to sow, and of rootes to set, if this or the like art to keep them, were not vsed; which is in this manner: Within a while after the first frosts haue taken the plants, that the leaues wither and fall, digge vp the rootes whole, and lay them in a dry place for three or foure dayes, that the superfluous moysture on the outside, may be spent and dryed, which done, wrap them vp seuerally in two or three browne papers, and lay them by in a boxe, chest or tub, in some conuenient place of the house all the winter time, where no winde or moist ayre may come vnto them; and thus you shall haue these rootes to spring a fresh the next yeare, if you plant them in the beginning of March, as I haue sufficiently tryed. But some haue tryed to put them vp into a barrell or firkin of sand, or ashes, which is also good if the sand and ashes be thorough dry, but if it bee any thing moist, or if they giue againe in the winter, as it is vsuall, they haue found the moisture of the rootes, or of the sand, or both, to putrefie the rootes, that they haue beene nothing worth, when they haue taken them forth. Take this note also for the sowing of your seede, that if you would haue variable flowers, and not all of one colour, you must choose out such flowers as be variable while they grow, that you may haue the seede of them: for if the flowers bee of one entire colour, you shall haue for the most part from those seedes, plants that will bring flowers all of that colour, whether it be white, red or yellow.

### The Place.

These plants grow naturally in the West Indies, where there is a perpetuall summer, or at the least no cold frosty winters, from whence the seede hath been sent into these parts of Europe, and are dispersed into euery garden almost of note.

### The Time.

These plants flower from the end of Iuly sometimes, or August, vntill the frosts, and cold ayres of the euenings in October, pull them down, and in the meane time the seed is ripe.

### The Names.

Wee haue not receiued the seedes of this plant vnder any other name, then *Mirabilia Peruviana*, or *Admirabilis planta*. In English wee call them, The meruaile of Peru, or the meruaile of the world: yet some Authors haue called it *Gelseminum*, or *Iasminum rubrum, & Indicum*: and Bauhinus *Solanum Mexicanum flore magno*.

### The Vertues.

We haue not knowne any vse hereof in Physicke.

## CHAP. XCI.

*Malua.* Mallowes.

OF the kindred of Mallowes there are a great number, some of the gardens, others wilde, some with single flowers, others with double, some with whole leaues, others with cut or diuided: to entreate of them all is not my purpose, nor the scope of this worke, but onely of such whose flowers, hauing beautie and respect, are fit to furnish this garden, as ornaments thereunto. And first of those single kindes, whose flowers come neerest vnto the fashion of the former Bell-flowers, and after to the double ones, which for their brauery, are entertained euery where into euery Countrey womans garden.

1. *Malua Hispanica flore carneo ample.*
The Spanish blush Mallow.

The Spanish Mallow is in forme and manner of growing, very like vnto our common fielde Mallow, hauing vpright stalkes two or three foote high, spread into diuers branches, and from the bottome to the toppe, beset with round leaues, like vnto our Mallowes, but somewhat smaller, rounder, and lesse diuided, yet larger below then aboue: the flowers are plentifully growing vpon the small branches, folding or writhing their leaues one about another before they bee blowne, and being open consist of fiue leaues, with a long forked clapper therein, of the same colour with the flower: the chiefest difference from the common consisteth in this, that the leaues of these flowers are longer, and more wide open at the brimmes (almost like a Bell-flower) and of a faire blush or light carnation colour, closing at night, and opening all the day: after the flowers are past, there come such like round heads, with small blacke seede, like vnto the common kinde, but somewhat smaller: the roote is small and long, and perisheth euery yeare.

2. *Alcea vulgaris flore carneo.* Vervaine Mallow with blush flowers.

There is a Mallow that hath long stalkes, and flowers like vnto the common wilde Mal-

1 *Malua Hiſpanica flore carneo ample.* The Spaniſh Mallow. 2 *Alcea Veneta.* The Venice Mallow. 3 *Alcea Americana.* Thorney Mallow. 4 *Alcea Ægyptia.* The Mallow of Egypt. 5 *Althea frutex.* The ſhrubbe Mallow. 6 *Malua hortenſis ſimplex.* Single Hollihockes. 7 *Malua roſea multiplex.* Double Hollihockes.

Mallow, and of the same deepe colour with it, so that you can hardly know it from the ordinary kinde, which is found growing wilde together with it, but onely by the leafe, which is as round and as large as the former, but cut into many fine diuisions, euen to the stalke that vpholdeth it, that it seemeth to consist onely of ragges, or peeces of leaues: Of this kinde I take a plante for this garden, growing in all respects like vnto it, but differing onely in the colour of the flowers, which are of the same blush or light carnation colour, or not much differing from the former Spanish kinde, with some veines therein of a deeper colour: the root hereof liueth, as the root of the common wilde kinde doth.

### 3. *Alcea peregrina sive vesicaria.*
### Venice Mallow, or Good night at noone.

The Venice Mallow hath long and weake stalkes, most vsually lying or leaning vpon the ground, hauing here and there vpon them long leaues and somewhat broad, cut in or gashed very deepely on both edges, that it seemeth as if they were diuers leaues set together, euery one standing on a long footestalke: at the ioynts of these stalkes, where the leaues are set, come forth seuerall flowers, standing vpon long foot-stalkes, which are somewhat larger then any of the former flowers, consisting of fiue leaues, small at the bottome, and wide at the brimmes, of a whitish colour tending to a blush, and sometimes all white, with spots at the bottomes of the leaues on the inside, of a very deepe purple or murrey colour, which addeth a great grace to the flower, and hauing also a long pestle or clapper in the middle, as yellow as gold: these flowers are so quickly faded and gone, that you shall hardly see any of them blowne open, vnlesse it bee betimes in the morning before the Sunne doe grow warme vpon them, for as soone as it feeleth the Sunnes warme heate, it closeth vp and neuer openeth againe, so that you shall very seldome see a flower blowne open in the day time, after nine a clocke in the morning: after these flowers are past, there rise vp in their places thinne, round, shining or transparent bladders, pointed at the toppe, and ribbed down all along, wherein are contained small, round, blackish seede: the roote is long and small, and perisheth euery yeare.

### 4. *Alcea fruticosa pentaphyllea.* Cinquefoile Mallow.

The stalkes of this Mallow are very long, hard or wooddy, more then of any of the other Mallowes: at the lower part whereof, and vp to the middle, stand diuers leaues vpon long footestalkes, parted or diuided into fiue parts or leaues, and dented about the edges; but vpwards from the middle to the toppe, the leaues haue but three diuisions: among these leaues stand large wide open flowers, of the colour of the common Mallow: the seede is smaller then in any other Mallow, but the rootes are great and long, spreading in the ground like vnto the roots of Marsh Mallowes, springing vp afresh euery yeare from the roote.

### 5. *Sabdariffa seu Alcea Americana.* Thorney Mallowe.

This Thorney Mallowe hath greene leaues next vnto the ground, that are almost round, but pointed at the end, and dented very much about the edges; the other leaues that growe vpon the stalke are diuided into three parts, like vnto a trefoile, and some of them into fiue diuisions, all of them dented about the edges: the stalke is reddish, with some harmelesse prickles in sundry places thereon, and riseth vp three or foure foote high in a good ground, a fit place, and a kindly yeare, bearing plenty of flowers vpon the stalkes, one at the foote of euery leafe, the toppe it selfe ending in a long spike, as it were of buddes and leaues together: the flowers are of a very pale yellow, tending to a white colour, spotted in the bottome of each of the fiue leaues, with a deepe purple spot, broad at the lower part, and ending in a point about the middle of the leafe, which are quickly fading, and not abiding aboue one day, with a long pestle in the middle diuided at the toppe: after the flower is past, commeth vp a short prickly podde, set within a small greene huske or cup that bore the flower, wherein is contai-

ned whitish, or rather browniſh yellow ſeede, flat and ſomewhat round, like vnto the ſeedes of Hollyhocke: the roote is ſtringie, and quickly periſheth; for it will hardly endure in our cold Country to giue flowers, much leſſe ſeede, vnleſſe (as I ſaid before) it happen in a kindly yeare, and be well planted and tended.

### 6. *Bamia ſeu Alcea Ægyptia.* The Mallow of Egypt.

This Mallow is alſo as tender to nourſe vp as the laſt, hauing the lower leaues broad like a Marſh Mallow, and of a freſh greene colour; but thoſe that growe vpon the ſtalke, and vp to the toppe, are diuided into fiue parts or points, but are not cut in to the middle ribbe, like the former Thorney Mallow, yet dented about the edges like vnto them: the flowers growe at the ſetting to of the leaues, like vnto a Mallow for forme, but of a whitiſh colour; after which come long fiue ſquare pointed pods, with hard ſhels, wherein are contained round blackiſh gray ſeede, as bigge as a Vetch or bigger: the roote periſheth quickly with vs, euen with the firſt froſts.

### 7. *Althæa frutex flore albo vel purpureo.* Shrubbe Mallow with a white or purple flower.

There are diuers ſorts of ſhrubbe Mallowes, whereof ſome that haue their ſtemmes or ſtalkes leſſe wooddy, dye downe to the ground euery yeare, and others that abide alwayes, are more wooddy: Of the former ſorts I intend not to ſpeake, referring them to a fitter place; and of the other, I will onely giue you the knowledge of one or two in this place, although I doe acknowledge their fitteſt place had been to be among the ſhrubbes; but becauſe they are Mallowes, I pray let them paſſe with the reſt of their kindred, and their deſcriptions in this manner: Theſe wooddy kindes of ſhrub Mallowes haue ſomewhat large, long, and diuided leaues, of a whitiſh greene colour, ſoft alſo, and as it were woolly in handling, ſet diſperſedly on the whitiſh hard or wooddy ſtalkes: their flowers are large, like vnto a ſingle Roſe or Hollyhocke, in the one being white with purple ſpots in the bottome; in the other either of a deepe red colour, or elſe of a paler purple, with a deeper bottome, and with veines running in euery leafe: they are ſomewhat tender, and would not be ſuffered to be vncouered in the Winter time, or yet abroad in the Garden, but kept in a large pot or tubbe, in the houſe or in a warme cellar, if you would haue them to thriue.

### 8. *Malua hortenſis roſea ſimplex & multiplex diuerſorum colorum.* Hollihockes ſingle and double of ſeuerall colours.

I ſhall not neede to make many deſcriptions of Hollihockes, in regard the greateſt difference conſiſteth in the flowers, which are in ſome ſingle, in ſome double, in ſome of one colour, and in others of other colours: for the loweſt leaues of Hollihockes are all round, and ſomewhat large, with many corners, but not cut in or diuided, ſoft in handling; but thoſe that growe vp higher are much more diuided into many corners: the ſtalkes ſometimes growe like a tree, at the leaſt higher then any man, with diuers ſuch diuided leaues on them, and flowers from the middle to the toppe, where they ſtand as it were a long ſpike of leaues and buds for flowers together: the flowers are of diuers colours, both ſingle and double, as pure white, and pale bluſh, almoſt like a white, and more bluſh, freſh and liuely, of a Roſe colour, Scarlet, and a deeper red like a crimſon, and of a darke red like blacke bloud; theſe are the moſt eſpeciall colours both of ſingle and double flowers that I haue ſeene: the ſingle flowers conſiſt of fiue broad and round leaues, ſtanding round like vnto ſingle Roſes, with a middle long ſtile, and ſome chiues aboue them: the double flowers are like vnto double Roſes, very thicke, ſo that no ſtile or vmbone is ſeene in the middle, and the outermoſt rowe of leaues in the flowers are largeſt, the innermoſt being ſmaller and thicke ſet together: after the flowers are paſt, there come vp as well in the double as ſingle, flat round heads, like flat cakes, round about the bottomes whereof growe flat whitiſh ſeede: the roote is long and great at the head, white and tough, like the roote of the common Mallowes, but greater, and will reaſonably well abide the Winter.

The

### The Place.

The firſt groweth wilde in Spaine. The ſecond in our owne Countrey. The third is thought to growe in Italy and Venice; but Lobel denieth it, ſaying, that it is there onely in Gardens, and is more plentifull in theſe parts then with them. The fourth Cluſius ſaith he found in many places of Germany. The fifth is ſuppoſed to be firſt brought out of the Weſt Indies, but an Arabicke name being giuen it, maketh me ſomewhat doubtfull how to beleeue it. The ſixth groweth in Egypt, where it is of great vſe, as Proſper Alpinus hath ſet downe in his Booke of Egyptian plants. The ſeuenth groweth in ſome parts both of Spaine and France. The laſt is not found but in Gardens euery where.

### The Time.

The firſt, ſecond, third, fourth, and laſt, doe flower from Iune vntill the end of Iuly and Auguſt. The reſt flower very late, many times not vntill September or October.

### The Names.

The firſt and ſecond haue their names ſufficiently expreſſed in their titles. The third is diuerſly called, as *Malua horaria*, *Alcea veſicaria*, *Alcea Veneta*, *Alcea Peregrina*, and of Matthiolus, *Hypecoum*. The moſt vſuall Engliſh name is Venice Mallow. The fourth is called *Alcea fruticoſa pentaphyllea*, and *Cannabinifolio*, or *Pentaphyllifolio*: In Engliſh, Cinquefoile Mallow. The fifth hath been ſent vnder the name of *Sabdariſa*, and *Sabdariffa*, and (as I ſaid) is thought to be brought from America, and therefore it beareth the name of that Country. The ſixth is called in Egypt, *Bamia*, or *Bammia*, and by that name ſent with the addition *del Cayro* vnto it: In Engliſh, Egyptian Mallow, or Mallow of Egypt. The ſeuenth is called *Althæa frutex*, and of ſome *Althæa arborea*: In Engliſh, Shrubbe Mallow, becauſe his ſtemme is woodie, and abideth as ſhrubbes and trees doe. The eight and laſt is called *Malua hortenſis*, *Malua Roſea*, and of ſome *Roſa vltra marina*: In Engliſh, of ſome Hockes, and vſually Hollihockes.

### The Vertues.

All ſorts of Mallowes, by reaſon of their viſcous or ſlimie quality, doe helpe to make the body ſoluble, being vſed inwardly, and thereby helpe alſo to eaſe the paines of the ſtone and grauell, cauſing them to be the more eaſily voided: being outwardly applyed, they mollifie hard tumors, and helpe to eaſe paines in diuers parts of the body; yet thoſe that are of moſt vſe, are moſt common. The reſt are but taken vpon credit.

## Chap. XCII.

### *Amaranthus.* Flower-gentle.

WE haue foure or fiue ſorts of Flower-gentle to trimme vp this our Garden withall, which doe differ very notably one from another, as ſhall be declared in their ſeuerall deſcriptions; ſome of which are very tender, and muſt be carefully regarded, and all little enough to cauſe them beare ſeede with vs, or elſe wee ſhall bee to ſeeke euery yeare: others are hardy enough, and will hardly be loſt out of the Garden.

1. *Amaranthus purpureus minor.* The small purple Flower-gentle.

This gallant purple Veluet flower, or Flower-gentle, hath a crested stalke two foote high or more, purplish at the bottome, but greene to the toppe, whereout groweth many small branches, the leaues on the stalkes and branches are somewhat broad at the bottome, and sharpe pointed, of a full greene colour, and often somewhat reddish withall, like in forme vnto the leaues of Blites (whereof this and the rest are accounted *species*, or sorts) or small Beetes: the flowers are long, spikie, soft, and gentle tufts of haires, many as it were growing together, broad at the bottome, and small vp at the toppe, pyramis or steeple-fashion, of so excellent a shining deepe purple colour, tending to a murrey, that in the most excellent coloured Veluet, cannot be seene a more orient colour, (and I thinke from this respect, the French call it *Passe velours*, that is to say, passing Veluet in colour) without any smell at all, which being bruised giueth the same excellent purple colour on paper, and being gathered in his full strength and beauty, will abide a great time (if it be kept out of the winde and sunne in a dry place) in the same grace and colour: among these tufts lye the seede scattered, which is small, very blacke, and shining: the rootes are a few threddy strings, which quickly perish, as the whole plant doth, at the first approach of Winter weather.

2. *Amaranthus Coccineus.* Scarlet Flower-gentle.

The leaues of this Flower-gentle are longer, and somewhat narrower then the former; the stalke groweth somewhat higher, bearing his long tufts at seuerall leaues, as also at the toppe of the stalkes, many being set together, but separate one from another, and each bowing or bending downe his head, like vnto a Feather, such as is worn in our Gallants and Gentlewomens heads, of an excellent bloudy Scarlet colour: the seede is blacke, like vnto the former: the roote perisheth quicklier, because it is more tender.

3. *Amaranthus tricolor.* Spotted or variable Flower-gentle.

The chiefest beauty of this plant consisteth in the leaues, and not in the flowers; for they are small tufts growing all along the stalke, which is nothing so high as the former, especially with vs, and at the ioynts with the leaues: the leaues hereof are of the same fashion that the former are, and pointed also; but euery leafe is to be seene parted into greene, red, and yellow, very orient and fresh (especially if it come to his full perfection, which is in hot and dry weather) diuided not all alike, but in some leaues, where the red or yellow is, there will be greene, and so varying, that it is very pleasant to behold: the seede hereof is blacke and shining, not to bee knowne from the former.

4. *Amaranthus Carnea spica.* Carnation Flower-gentle.

There is another more rare then all the rest, whose leaues are somewhat longer, and narrower then the first, and like vnto the second kinde: the spikes are short, many set together, like branches full of heads or eares of corne, euery one whereof hath some long haires sticking out from them, of a deep blush, tending to a carnation colour.

5. *Amaranthus purpureus maior panniculis sparsis.*
Great Floramour, or purple Flower-gentle.

The great Floramour hath one thicke, tall, crested, browne red stalke, fiue or six foote high, from whence spring many great broad leaues, like vnto the former for the forme, but much larger & redder for the most part, especially the lowest, which brancheth forth into diuers parts, & from between these leaues, & the stalks or branches, as also at the tops of them, stand long, spikie, round, & somewhat flat tufts, of a more reddish purple colour then the first, and diuided also into seuerall parts, wherin when they

are

are full ripe, are to be seen an innumerable company of white seed, standing out among the short thrums, and do then easily fall away with a little touching; euery one of these white seed hath as it were an hole halfe bored through therin: the root is a great bush of strings, spreading in the ground, whereby it is strongly fastened, yet perisheth euery yeare, after it hath giuen his seede.

### The Place.

All these plants growe in the Easterne Countries, as Persia, Syria, Arabia, &c. except the greatest, which hath been brought out of the West Indies, where it is much vsed, especially the seede: they are all, except it, noursed vp with much care in our Gardens, and yet in a backward or cold yeare they will not thriue, for that they desire much heate: but the greatest doth alwayes giue ripe seede euery yeare.

### The Time.

They beare their gallant tufts or spikes for the most part in August, and some not vntill September.

### The Names.

The name *Amaranthus* is giuen to all these plants, taken from the Greeke word ἀμαραντὸς, *non marcescens*, or *non senescens*, that is, neuer waxing old, and is often also imposed on other plants, who haue the same property, that is, that their flowers being gathered in a fit season, will retaine their natiue colour a long time, as shall be shewed in the Chapter following. Diuers do thinke the first to be *Phlox*, or *Flamma* of Theophrastus. The third is called *Gelosia*, or *Celosia* of Tragus. Spigelius in his *Isagoges* saith, it is generally taken to be *Sophonia*, whereof Plinie maketh mention; and Lobel, to bee the Persians *Theombroton* of Plinie. The Italians, from whom I had it (by the meanes of Mr. Doctor Iohn More, as I haue had many other rare simples) call it, *Blito di tre colori*, A three coloured Blite. The fifth, which is the greatest, hath been sent from the West Indies by the name of *Quiniua*, as Clusius reporteth. The name Flower-gentle in English, and *Flor amour*, which is the French, of *Flos amoris*, and *Passe velours*, as is before said, or Veluet flower, according to the Italian, *Fior veluto*, are equally giuen to all these plants, with their seuerall distinctions, as they are expressed in their titles.

### The Vertues.

Diuers suppose the flowers of these plants doe helpe to stay the fluxe of bloud in man or woman, because that other things that are red or purple doe performe the same. But Galen disproueth that opinion very notably, *in lib. 2. & 4. de simpl. medicament. facultatibus.*

## CHAP. XCIII.

### *Helichrysum, siue Amaranthus luteus.*
### Golden Flower-gentle, Goldilockes, or Gold-flower.

THe propinquity of property (as I before said) hath caused the affinity in name, and so in neighbourhood in these plants, wherein there are some diuersity; and although they differ from them before in many notable points, yet they all agree with themselues in the golden, or siluer heads or tufts they beare; and therefore I haue

1 *Amaranthus purpureus maior.* The great purple Flower-gentle. 2 *Amaranthus purpureus minor.* The lesser purple Flower-gentle. 3 *Amaranthus tricolor.* Variable Flower-gentle. 4 *Amaranthus Coccineus.* Scarlet Flower gentle. 5 *Amaranthus carnea spica.* Carnation Flower-gentle. 6 *Chrysocoma* Golden Cassidonie. 7 *Helichrysum Creticum.* Candy Goldilockes. 8 *Gnaphalium Americanum.* Long or Life everlasting. 9 *Gnaphalium montanum,* Cats foote. 10 *Gnaphalium rosæum.* The Cotton Rose.

haue compriſed them in one Chapter, and will begin with that which commeth neareſt vnto the *Helichryſum* of Dioſcorides, or *Aurelia* (as Gaza tranſlateth it) of Theophraſtus.

1. *Heliochryſum.* The Golden flower of life.

This firſt Golden tuft riſeth vp with many hard, round, white ſtalkes, a foote and a halfe high, whereon at certaine diſtances ſtand many fine cut leaues, or rather one leafe cut into many ſmall fine pàrts, almoſt as ſmall as Fenell, but grayiſh, like vnto the Cudweedes or Cotton-weedes (whereof certainly theſe are ſpeciall kindes) at the toppes of the ſtalkes ſtand many round flowers, of a pale gold colour, in an vmbell cloſe together, yet euery flower vpon his owne ſtalke, and all of an euen height, which will keepe the colour, being gathered, and kept dry for a long time after, and are of a hot and quicke ſent: the roote is ſmall and wooddy, ſpreading vnder the vpper cruſt of the earth, and liueth long in his owne naturall place, but very hardly endureth the cold of our Winters, vnleſſe they be milde, or it be well defended.

2. *Helichryſum Creticum.* Candy Goldilockes.

Candy Goldilockes hath two or three ſmall ſlender white branches, ſet here and there very ſcatteringly, with ſmall, long, and narrow hoary leaues, hauing yellow heads of flowers at the tops made into vmbels or tufts, not ſo round and euen as the former, but longwiſe one aboue another, the heads being made as it were of ſcales, looſly, and not ſo cloſely ſet together, as in the next following, which when they are full ripe, doe paſſe into doune, and are blowne away with the winde, hauing a ſmall reddiſh ſeede at the end; but will abide a long time, as the other in his beauty, being gathered in time, as the reſt will doe.

3. *Helichryſum Orientale ſiue Amaranthus luteus.*
Golden Flower-gentle.

This moſt beautifull plant is very like vnto the former Candy Goldilockes laſt deſcribed, but growing vp higher, with many more branches, and more hoary, white, and woolly, hauing alſo long and narrow white leaues, but ſomewhat broader, and thicker ſet on the branches: the tufts of flowers or vmbels likewiſe doe conſiſt of longer and larger heads, more ſcaly, and cloſer compact together, of an excellent pale gold yellow colour, and ſhining, with ſome yellow threads or thrummes in the middle: the roote dyeth not euery yeare, but liueth long, eſpecially in the South and Eaſt Countries, where no colds or froſts are felt; but will require extraordinary care and keeping, and yet ſcarce ſufficient to preſerue it in theſe cold Countries.

4. *Chryſocome ſiue Stœchas Citrina.* Golden tufts or Golden Caſſidony.

This Golden flower is ſomewhat like the former of theſe two laſt deſcribed, hauing hoary ſtalkes and leaues, ſtanding confuſedly on them, being long, and narrower then any of the former: the tops of the ſtalkes are diuided into many parts, each bearing a ſmall long yellow head or flower at the toppe, with ſome yellow thrummes in them, which heads being many, are diffuſedly ſet together, like a looſe or ſparſed vmbell, keeping their colour long before they wither, and when they are ripe, haue thinne ſmall reddiſh ſeede, like Mariorome ſeede, but ſmaller; the roote is ſmall and blacke: the whole plant, as well leaues and flowers, as rootes, are of a ſtrong ſharpe ſent, yet pleaſant.

5. *Argyrocome ſiue Gnaphalium Americanum.*
Liue long or Life euerlaſting.

This ſiluer tuft or Indian Cotton weede, hath many white heads of leafes at their firſt ſpringing out of the ground, couered with a hoary woollineſſe like cotton, which riſing into hard, thicke round ſtalkes, containe ſtill the ſame hoarineſſe vpon them, as alſo vpon the long and narrow leaues which are ſet thereon, eſpecially on the vnder

ſide,

side, for the vpper sides are of a darke shining greene colour: the stalkes are diuided at the toppe into many small branches, each whereof haue many scaly tufted heads set together, couered ouer with cotton before their opening, and then disseuering one from another, abiding very white on the outside, when they are fully growne, but with a small yellow thrume in the midde of euery flower, which in time turne into yellow doune, apt to be blowne away with euery winde: the roots are long and black on the outside, creeping vnder ground very much.

### 6. *Gnaphalium montanum flore albo & flore purpureo.* White and purple Cats foote.

This small Cudweede or Cottonweede, hath many small white woolly leaues growing from the roote, which is composed of a few small blackish threds, and lying vpon the ground somewhat like vnto the leaues of a small Mouse-eare, but smaller; from among which riseth vp a small stalke of halfe a foote high or thereabouts, beset here and there with some few leaues, at the top whereof commeth forth a tuft of small flowers, set close together, in some of a pure white, in others of a purple or reddish colour, in some of a pale red or blush, and in others of a white and purple mixt together, which for the beauty is much commended and desired, but will hardly abide to be kept in Gardens, so vnwilling they are to leaue their naturall abiding.

### 7. *Gnaphalium Roseum.* The Cotton Rose.

This little rose Cotton weede hath many such like woolly leaues, growing as the former from the roote vpon small short branches, not full an hand breadth high, in fashion somewhat like vnto Daysie leaues, but lesser, and round pointed: at the toppe of euery stalke or branch, standeth one flower, composed of two rowes of small white leaues, layd open like a Starre or a Rose, as it beareth the name, hauing a round head in the middle made of many yellow threds or thrumes, which falling away, there riseth vp a small round head, full of small seedes: the root is small, long and threddy.

## The Place.

The foure first plants doe grow naturally in many of the hot Countries of Europe, as Spaine, Italie, and Prouince in France; as also in Candy, Barbary, and other places, and must be carefully kept with vs in the winter time. The Liue long was brought out of the West Indies, and groweth plentifully in our gardens. The two last doe grow as well in the colder Countries of Germany, as in France and other places.

## The Time.

They all flower in the end of September, if they will shew out their beauty at all with vs, for sometimes it is so late, that they haue no faire colour at all, especially the foure first sorts.

## The Names.

Variable and many are the names that seuerall Writers do call these foure first sorts of plants, as *Helichrysum*, *Heliochrysum*, or *Elichrysum*. *Eliochrysum*, *Chrysocome*, *Coma aurea*, *Amaranthus luteus*, *Stoechas Citrina*, and *Aurelia*, with others, needlesse here to be recited: it is sufficient for this worke, to giue you knowledge that their names are sufficient as they are expressed in their titles: The fift is called *Gnaphalium* by Carolus Clusius, from the likenesse of the vmbels or tufts of heads, though greater and white: for as I said before, the Cotton weedes are of kindred with the golden tufts: It hath been called by our English Gentlewomen, Liue long, and Life euerlasting, because of the durabilitie of the flowers in their beautie. The two last are cal-

led *Gnaphalium*, according to their titles; and in English they may passe vnder those names are set downe with them.

### The Vertues.

The foure first are accounted to bee hot and dry, and the three last to bee cold and dry: yet all of them may to some good purpose bee applyed to rheumaticke heads. The former foure are likewise vsed to cause vrine, and in baths to comfort and heate cold parts. They are also layd in chests and wardrobes, to keepe garments from moths; and are worne in the heads and armes of Gentiles and others, for their beautifull aspect.

## CHAP. XCIIII.

### *Canna Indica.* The Indian flowring Reede.

THere are two kindes or sorts of this beautifull plant, the one with a red flower, the other with a yellow, spotted with reddish spots, both which in some kindly yeares haue borne their braue flowers, but neuer any ripe seede, and doth not abide the extremities of our winters, eyther abroade or vnder couert, vnlesse it meete with a stoue or hot-house, such as are vsed in Germany, or such other like place: For neyther house nor cellar will preserue it, for want of heate.

### *Canna Indica flore rubro.* Red flowred Indian Reede.

This beautifull plant riseth vp with faire greene, large, broade leaues, euery one rising out of the middle of the other, and are folded together, or writhed like vnto a paper Coffin (as they call it) such as Comfitmakers and Grocers vse, to put in their Comfits and Spices, and being spread open, another riseth from the bottome thereof, folded in the same manner, which are set at the ioynts of the stalke when it is risen vp, like vnto our water Reede, and growing (if it runne vp for flower) to be three or foure foote high, as I haue obserued in mine owne garden: the flowers grow at the toppe of the stalke one aboue another, which before their opening are long, small, round, and pointed at the end, very like vnto the claw of a Crauise or Sea-Crab, and of the same red or crimson colour, but being open, are very like vnto the flower of *Gladiolus* or Corne-flagge, but of a more orient colour then at the first, and standing in a rough huske, wherein afterwards standeth a three square head, containing therein round blacke seede, of the bignesse of a pease: the roote is white and tuberous, growing into many knobs, from whence arise such other leaues and stalkes, whereby it encreaseth very much, if it be righrly kept and defended.

### *Canna Indica flore flauo punctato,* Yellow spotted Indian Reede.

This Reede groweth vp with leaues and flowers, in all points so like vnto the former, that it cannot bee knowne from it, vntill it come to flower, which is of a yellow colour, spotted with reddish spots, without any other difference.

### The Place.

These plants grow naturally in the West Indies, from whence they were first sent into Spaine, and Portugall, where Clusius saith he saw them planted by the houses sides, flowring in winter, which might be in those warme Countreyes. We preserue them with great care in our gardens, for the beautifull aspect of their flowers.

### The Time.

They flower not with vs vntill the end, or middle of Auguſt, at the ſooneſt.

### The Names.

They are called of ſome *Canna Indica*, and *Arundo Indica*, of others *Cannacorus*, and of ſome *Flos Cancri*, becauſe the colour of the flowers, as well as the forme of the buds, are ſo like vnto a Sea-Crabs cle, or claw.

### The Vertues.

There is not any vſe of theſe in Phyſicke that I know.

## CHAP. XCV.

## *Mandragoras.* Mandrake.

THe Mandrake is diſtinguiſhed into two kindes, the male and the female; the male hath two ſorts, the one differing from the other, as ſhall be ſhewed; but of the female I know but one: The male is frequent in many gardens, but the female, in that it is more tender and rare, is nourſed vp but in a few.

### *Manadrgoras mas.* The male Mandrake.

The male Mandrake thruſteth vp many leaues together out of the ground, which being full growne, are faire, large and greene, lying round about the roote, and are larger and longer then the greateſt leaues of any Lettice, whereunto it is likened by Dioſcorides and others: from the middle, among theſe leaues, riſe vp many flowers, euery one vpon a long ſlender ſtalke, ſtanding in a whitiſh greene huske, conſiſting of fiue pretty large round pointed leaues, of a greeniſh white colour, which turne into ſmall round apples, greene at the firſt, and of a pale red colour when they are ripe, very ſmooth and ſhining on the outſide, and of a heady or ſtrong ſtuffing ſmell, wherein is contained round whitiſh flat ſeede: the roote is long and thicke, blackiſh on the outſide, and white within, conſiſting many times but of one long roote, and ſometimes diuided into two branches a little below the head, and ſometimes into three or more, as nature liſteth to beſtow vpon it, as my ſelfe haue often ſeene, by the tranſplanting of many, as alſo by breaking and cutting off of many parts of the rootes, but neuer found harme by ſo doing, as many idle tales haue beene ſet downe in writing, and deliuered alſo by report, of much danger to happen to ſuch, as ſhould digge them vp or breake them; neyther haue I euer ſeene any forme of man like or woman-like parts, in the rootes of any: but as I ſaid, it hath oftentimes two maine rootes running down-right into the ground, and ſometimes three, and ſometimes but one, as it likewiſe often happeneth to Parſneps, Carrots, or the like. But many cunning counterfeit rootes haue bin ſhaped to ſuch formes, and publickly expoſed to the view of all that would ſee them, and haue been tolerated by the chiefe Magiſtrates of the Citie, notwithſtanding that they haue beene informed that ſuch practices were meere deceit, and vnſufferable; whether this happened through their ouer-credulitie of the thing, or of the perſons, or through an opinion that the information of the truth roſe vpon enuy, I know not, I leaue that to the ſearcher of all hearts: But this you may bee bold to reſt vpon, and aſſure your ſelues, that ſuch formes as haue bin publickly expoſed to be ſeene, were neuer ſo formed by nature, but only by the art and cunning of knaues and deceiuers. and let this be your *Galeatum* againſt all ſuch vaine, idle and ridiculous toyes of mens inuentions.

There is likewiſe another ſort of theſe male Mandrakes, which I firſt ſaw at Canterbury, with my very louing and kinde friende Iohn Tradeſcante, in the garden of the Lord Wotton, whoſe gardiner he was at that time ; the leaues whereof were of a more grayiſh greene colour, and ſomewhat folded together, when as the former kind that grew hard by it, was of the ſame forme that is before deſcribed, and ordinary in all others : but whether the apples were differing from the other, I know not, nor did they remember that euer it had borne any.

*Mandragoras famina.* The female Mandrake.

The female Mandrake doth likewiſe put vp many leaues together, from the head of the roote, but they are nothing ſo large, and are of a darker greene colour, narrower alſo and ſhining, more crumpled, and of a ſtronger ſent : the flowers are many, riſing vp in the middle of the leaues, vpon ſlender ſtalkes, as in the male kind, but of a blewiſh purple colour, which turne into ſmall round fruite or apples, and not long like a peare (as Cluſius reporteth that ſaw them naturally growing in Spaine) greene at the firſt, and of a pale yellowiſh colour, when they are full ripe; of a more pleaſing, or if you will, of a leſſe heady ſent then the apples of the male, wherein is contained ſuch like ſeede, but ſmaller and blacker : the rootes are like the former, blacke without and white within, and diuided in the ſame manner as the male is, ſometimes with more, and ſometimes with fewer parts or branches.

## The Place.

They grow in many places of Italie, as Matthiolus reporteth, but eſpecially on Mount Garganus in Apulia. Cluſius ſaith hee found the female in many wet grounds of Spaine, as alſo in the borders of thoſe medowes that lye neere vnto riuers and water courſes. The male is cheriſhed in many Gardens, for pleaſure as well as for vſe : but the female as is ſaid, is both very rare, and farre more tender.

## The Time.

The Male flowreth in March, and the fruit is ripe in Iuly. The Female, if it be well preſerued, flowreth not vntill Auguſt, or September; ſo that without extraordinary care, we neuer ſee the fruite thereof in our gardens.

## The Names.

*Mandragoras mas* is called *albus*, as the *Famina* is called *niger*, which titles of blacke and white, are referred vnto the colour of the leaues : the female is called alſo *Thridacias*, from the likeneſſe of Lettice, whereunto they ſay in forme it doth carry ſome ſimilitude. Dioſcorides ſaith, that in his time the male was called *Morion*, and both of them *Antimelum*, and *Circæa*. Wee call them in Engliſh, The male, and the female Mandrake.

## The Vertues.

The leaues haue a cooling and drying qualitie, fit for the oyntment *Populeon*, wherein it is put. But the Apples haue a ſoporiferous propertie, as Leuinus Lemnius maketh mention in his Herball to the Bible, of an experiment of his owne. Beſides, as Dioſcorides firſt, and then Serapio, Auicen, Paulus Ægineta, and others alſo do declare, they conduce much to the cooling and cleanſing of an hot *matrix*. And it is probable, that Rachel knowing that they might be profitable for her hot and dry body, was the more earneſt with Leah for her Sonne Rubens Apples, as it is ſet downe *Geneſis* 30. *verſe* 14. The ſtrong ſent of theſe apples is remembred alſo, *Cant.* 7. 13. although ſome would diuert the ſignification of the Hebrew word, דודאים,

vnto

vnto Violets, or some other sweet flowers, in the former place of *Genesis*, and the fruit of *Musa*, or Adams Apples in this place of the *Canticles*. Hamilcar the Carthaginian Captaine is said to haue infected the wine of the Lybians (his enemies against whom he fought) with the apples of Mandrake, whereby they being made exceeding drowsie, he obtained a famous victory ouer them.

## CHAP. XCVI.

### *Pomum Amoris.* Loue Apples.

ALthough the beautie of this plant consisteth not in the flower, but fruit, yet giue me leaue to insert it here, lest otherwise it haue no place: whereof there are two especiall sorts, which wee comprehend in one Chapter, and distinguish them by *maius* and *minus*, greater and smaller: yet of the greater kinde, we haue noursed vp in our Gardens two sorts, that differ only in the colour of the fruite, and in nothing else.

### *Pomum Amoris maius fructu rubro.*
### Great Apple of Loue the ordinary red sort.

This greater kinde of Loue Apples, which hath beene most frequently cherished with vs, hath diuers long and trayling branches, leaning or spreading vpon the ground, not able to sustaine themselues, whereon doe grow many long winged leaues, that is, many leaues set on both sides, and all along a middle ribbe, some being greater, and others lesse, iagged also and dented about the edges, of a grayish ouer-worne greene colour, somewhat rough or hairy in handling; from among the leaues and the branches come forth long stalkes, with diuers flowers set thereon, vpon seuerall short footstalks, consisting of sixe, and sometimes of eight small long yellow leaues, with a middle pricke or vmbone, which after the flowers are fallen, riseth to be the fruite, which are of the bignesse of a small or meane Pippin, vneuenly bunched out in diuers places, and scarce any full round without bunches, of a faire pale reddish colour, or somewhat deeper, like vnto an Orenge, full of a slimie iuice and watery pulpe, wherein the seede lyeth, which is white, flat and somewhat rough: the roote shooteth with many small strings and bigger branches vnder ground, but perisheth at the first feeling of our winter weather. The fruite hereof by often sowing it in our Land, is become much smaller then I haue here described it: but was at the first, and so for two or three yeares after, as bigge as I haue related it.

### *Pomum Amoris maius fructu luteo.* Yellow Amorous Apples.

Of the same kinde is this other sort of Amorous Apples, differing in nothing but the colour of the fruite, which is of a pale yellow colour, hauing bunches or lobes in the same manner, and feede also like the former.

### *Pomum Amoris minus, siue Mala Ethiopica parua.*
### Small Loue Apples.

The small Apples of Loue in the very like manner, haue long weake trayling branches, beset with such like leaues as the greater kinde hath, but smaller in euery part: the flowers also stand many together on a long stalke, and yellow as the former, but much smaller: the fruite are small, round, yellowish red berries, not much bigger then great grapes, wherein are contained white flat seede, like the other, but smaller: the roote perisheth in like manner euery yeare, and therefore must bee new sowen euery spring, if you will haue the pleasure of their sight in the garden; yet some yeares I haue known them rise of their owne sowing in my garden.

The

### The Place.

They growe naturally in the hot Countries of Barbary, and Ethiopia; yet some report them to be first brought from Peru, a Prouince of the West Indies. Wee onely haue them for curiosity in our Gardens, and for the amorous aspect or beauty of the fruit.

### The Time.

They flower in Iuly and August, and their fruit is ripe in the middle or end of September for the most part.

### The Names.

The first is named diuersly by diuers Authors; for Lobel, Camerarius, and others, call them *Poma amoris*. Dodonæus *Aurea Mala*. Gesnerus first, and Bauhinus after him, make it to be a kinde of *Solanum Pomiferum*. Anguillara taketh it to be *Lycopersicum* of Galen. Others thinke it to bee *Glaucium* of Dioscorides. The last is called *Mala Æthiopica parua*, and by that title was first sent vnto vs, as if the former were of the same kinde and country. We call them in English, Apples of Loue, Loue-Apples, Golden Apples, or Amorous Apples, and all as much to one purpose as another, more then for their beautifull aspect.

### The Vertues.

In the hot Countries where they naturally growe, they are much eaten of the people, to coole and quench the heate and thirst of their hot stomaches. The Apples also boyled, or infused in oyle in the sunne, is thought to be good to cure the itch, assuredly it will allay the heate thereof.

## Chap. XCVII.

### *Digitalis.* Foxegloue.

THere are three principall sorts of Foxegloues; a greater, a middle or meane sort, and a lesser, and of them, three especiall colours, that is, purple, white, and yellow; the common purple kinde that groweth abroad in the fields, I leaue to his wilde habitation: and of the rest as followeth.

#### 1. *Digitalis maxima ferruginea.* Dun coloured Foxegloues.

The leaues of this Foxegloue are long and large, of a grayish green colour, finely cut or dented about the edges, like the teeth of a fine sawe; among which commeth vp a strong tall stalke, which when it was full growne, and with ripe seede thereon, I haue measured to be seuen foot high at the least, whereon grow an innumerable company (as I may so say, in respect of the aboundance) of flowers, nothing so large as the common purple kinde, that groweth wilde euery where in our owne Countrey, and of a kinde of browne or yellowish dunne colour, with a long lippe at euery flower; after them come seede, like the common kinde, but in smaller heads: the rootes are stringie like the ordinary, but doe vsually perish, or seldome abide after it hath giuen seed.

#### 2. *Digitalis maior flore carneo.* Blush coloured Foxegloues.

This kinde of Foxegloues hath reasonable large leaues, yet not altogether so large

as

1 *Canna Indica.* The Indian Reed. 2 *Mandragoras mas.* The male Mandrake. 3 *Pomum amoris maius,* Great Apples of loue. 4 *Digitalis maior flore luteo amplo.* The great yellow Foxegloue. 5 *Digitalis media flore luteo rubente.* Orenge tawny Foxegloues. 6 *Digitalis maxima ferruginea.* Dun coloured Foxegloues.

as the common field kinde : the flowers are also smaller then the common sort, but of a blush colour.

3. *Digitalis media flore luteo rubente.* Orenge tawnie Foxegloue.

As this Foxegloue is none of the greatest, so also is it none of the smallest; but a sort betweene both, hauing leaues in some proportion correspondent to the lesser yellow Foxegloue, but not so large as the lesser white: the flowers are long and narrow, almost as large as the last white, but nothing so large as the first white, of a faire yellowish browne colour, as if the yellow were ouershadowed with a reddish colour, and is that colour wee vsually call an Orenge tawnie colour : the seede is like the former: the rootes perish euery yeare that they beare seede, which is vsually the second yeare of the springing.

4. *Digitalis maior alba.* The greater white Foxegloue.

This white Foxegloue is in all things so like vnto the purple wilde kinde, that it can hardly be distinguished from it, vnlesse it be in the fresher greennesse and largenesse of the leaues: the flowers are as great in a manner as the purple, but wholly white, without any spot in them: the seed and other things agree in all points.

5. *Digitalis alba altera seu minor.* The lesser white Foxegloue.

We haue in our Gardens another sort of white Foxegloue, whose leaues are like vnto the last described, but not altogether so long or large, and of a darker greene colour: the stalke groweth not so high, as not full three foote : the flowers are pure white, fashioned like vnto the former, but not so great or large, in all other things alike: the rootes hereof did abide sometime in our Gardens, but since perished, and the seede also, since when we neuer could obtaine from any our friends of that kinde againe.

6. *Digitalis maior lutea flore amplo.*
The great yellow Foxegloue.

The leaues of this greater yellow Foxegloue, are in forme somewhat like vnto the common purple kinde, but not altogether so large: the stalke groweth to bee three or foure foote high, whereon stand many long hollow pendulous flowers, in shape like the ordinary purple: but somewhat shorter, and more large and open at the brimmes, of a faire yellow colour, wherein are long threads, like as in the others: the roote hereof is greater at the head, and more wooddy then any of the reft, with many smaller fibres, spreading themselues in the ground, and abideth almost as well as our common purple kinde.

7. *Digitalis minor lutea sive pallida.*
The small pale yellow Foxegloue.

This small pale yellow Foxegloue hath somewhat short, broad, smooth and darke greene leaues, snipt or dented about the edges very finely: the stalke is two foot high, beset with such like leaues, but lesser: the flowers are moe in number then in any of the rest, except the first and greatest, and growe along the vpper part of the stalke, being long and hollow, like the other, but very small, and of a pale yellow colour almost white: the seede vessels are small like the former, wherein are contained seede like the rest, but smaller: the rootes are stringy, but durable, and seldome perish with any iniury of the extreamest frosts.

The Place.

The great white kinde hath been often, and in many places found wilde in our owne Country, among or hard by the common purple kinde. All the rest are strangers, but cherished in our Gardens.

The

### The Time.

They flower in Iune and Iuly, and some in August, their seede becomming ripe quickly after.

### The Names.

Onely the name *Digitalis*, is of all Writers giuen vnto these plants; for it is not knowne to bee remembred of any of the old Authors. Wee call them generally in English, Foxegloue; but some (as thinking it to bee too foolish a name) doe call them Finger-flowers, because they are like vnto the fingers of a gloue, the ends cut off.

### The Vertues.

Foxegloues are not vsed in Physicke by any iudicious man that I know; yet some Italians of Bononia, as Camerarius saith, in his time vsed it as a wound herbe.

## Chap. XCVIII.

### *Verbascum.* Mullein.

THere be diuers kindes of Mullein, as white Mullein, blacke Mullein, wooddy Mullein, base Mullein, Moth Mullein, and Ethiopian Mullein, all which to distinguish or to describe, is neither my purpose, nor the intent of this worke, which is to store a Garden with flowers of delight, and sequester others not worthy of that honour. Those that are fit to bee brought to your consideration in this place, are first, the *Blattarias*, or Moth Mulleins, and then the wooddy Mullein, which otherwise is called French Sage, and lastly, the Ethiopian Mullein, whose beauty consisteth not in the flower, but in the whole plant; yet if it please you not, take it according to his Country for a Moore, an Infidell, a Slaue, and so vse it.

#### 1. *Blattaria lutea odorata.* Sweete yellow Moth Mullein.

The yellow Moth Mullein whose flower is sweete, hath many hard grayish greene leaues lying on the ground, somewhat long and broad, and pointed at the end: the stalks are two or three foot high, with some leaues on them, & branching out from the middle vpwards into many long branches, stored with many small pale yellow flowers, of a pretty sweete sent, somewhat stronger then in the other sorts, which seldome giueth seede, but abideth in the roote, liuing many yeares, which few or none of the others doe.

#### 2. *Blattaria lutea maior siue Hispanica.* The great yellow Moth Mullein.

This Spanish kinde hath larger and greener leaues then the former, and rounder and larger then the next that followeth: the stalke is higher then in any of the Moth Mulleins, being for the most part foure or fiue foote high, whereon toward the toppe growe many goodly yellow flowers, consisting of fiue leaues, as all the rest doe, not so thicke set as the former, but much larger, with some small purplish threads in the middle: the ends whereof are fashioned somewhat like as if a Flie were creeping vp the flower, which turne into round heads, sometimes two or three or more standing together, but vsually one, wherein lye small duskie seed: the roote is not great nor full of threads, and doth perish most vsually hauing giuen seede, except the Winter bee very milde.

3. *Blattaria*

3. *Blattaria lutea altera vulgatior.* The ordinary yellow Moth Mullein.

This yellow Moth Mullein (which is the most frequent in our Gardens) hath longer, and narrower leaues then any of the former, and roundly notched or dented on the edges, of a darke greene colour: the stalke is sometimes branched, but most vsually single, whereon stand many gold yellow flowers, not fully so large as the Spanish kinde, but with the like purple threads in the middle: the seede is small, and contained in the like round heads, but alwaies euery one single by it selfe: the roote perisheth euery yeare that it beareth seede.

4. *Blattaria flore luteo purpurascente.* Cloth of gold Moth Mullein.

The greatest point of difference betweene this and the last described, consisteth chiefly in the colour of the flower, which in this is of the colour of cloth of gold, that is, the ground yellow, and ouershadowed with a bright crimson colour, which is a fine colour of much delight: the threads in the middle are not so purple red as in the former, but much about the colour of the flower: this is not so willing to giue seede, and will as hardly abide in the roote, and hath out of question risen from the seede of the former.

5. *Blattaria flore albo.* White Moth Mullein.

The leaues of the white Moth Mullein are somewhat like vnto the yellow, yet not altogether so much roundly notched about the edges, but rather a little dented, with sharper notches: the stalke riseth as high as the yellow, and hath now and then some branches about it: the flowers hereof are pure white, as large and great as the ordinary yellow, or somewhat larger, with the like purple threads in the middle, as are in the yellow: the seed is like the other; the root perisheth in like maner, and will not endure.

6. *Blattaria flore purpureo.* Purple Moth Mullein.

The Purple Moth Mullein hath his leaues lying on the ground, broader and shorter then any of the other, of a more grayish greene colour, and without any denting for the most part about the edges, sharpe pointed also at the end of the leafe; among the leaues riseth vp the stalke, not so high as either the white or the yellow, and many times branched, bearing many flowers thereon, of the same fashion, and no whit smaller, of a faire deepe blewish colour tending to rednesse, the threads in the middle of the flowers being yellow: the seede vessels hereof are somewhat smaller then any of the former, except the first sweete yellow kinde: the roote hereof is long, thicke, and blackish on the outside, abiding very well from yeare to yeare, and riseth well also from the sowing of the seede.

7. *Blattaria flore cærulco.* Blew Moth Mullein.

This blew Moth Mullein is in all respects like vnto the former purple kinde, saiuing onely in the colour of the flower, which is of a blewish violet colour, and is not much inferiour either in greatnesse of the plant, or in the largenesse of the flower, vnto the former purple kinde, and endureth many yeares in the like manner. And these be all the sorts of this kinde of Moth Mullein, that I haue seene and noursed vp for this my Garden, without interposing any vnknowne, not seene, or vnworthy.

8. *Verbascum siluestre siue quartum Matthioli.*
Wooddy Mullein or French Sage.

Wooddy Mullein or French Sage, hath diuers wooddy branches two or three foot high, very hoary or white, whereon at seuerall ioynts stand diuers thicke leaues, white also and hoary, long, somewhat broad, round pointed, and rough, somewhat resembling the leaues of Sage in the forme and roughnesse, but not in the sent, whereof our people

people gaue it the name of Sage, calling it French Sage (when as it is as great a stranger in France as in England, yet they doe with this as with many other things, calling them French, which come from beyond the Seas; as for example, all or most of our bulbous flowers, they call French flowers, &c.) at the toppes of the stalkes and branches, at certaine distances, are placed round about them many gaping flowers, like vnto the flowers of Sage, but yellow: after which now and then come seede, somewhat bigger then the Moth Mulleins, and lesse then the next Mullein of Ethiopia: the roote is wooddy at the toppe, with diuers blackish strings growing from it, and endureth as well aboue ground with his leaues, as vnder it with his rootes.

### 9. *Æthiopis.* Ethiopian Mullein.

This Mullein of Ethiopia hath many great, broad, and large leaues lying on the ground, rent or torne in diuers of them very much on the sides, of so hoary a white greene colour, that it farre passeth any of the white Mulleins, that growe wilde abroad in our owne Country; for they are of a yellowish white hoarinesse, nothing so pleasant to looke on as this: in the middle of these leaues riseth vp a square strong stalke, foure or fiue foote high, set full of such like leaues as growe belowe, but much lesser, and lesser still vp to the toppe, all hoary and woolly, as the rest, and diuided into manie branches, spreading farre, and taking vp a great compasse of ground, more then any one roote of Garden Clary, or other such like plant: at each of the stalkes and branches are set two small leaues, and with them, round about the stalkes, stand many small gaping flowers, of a pale bleake blew colour: the seede is almost as large as Garden Clary seede, and of the same forme and colour: the roote is wooddy, and perisheth as soone as it hath borne seede, which is vsually the second yeare after the sowing; for the first yeare it seldome runneth vp to flower.

### 10. *Lamium Pannonicum siue Galeopsis Pannonica.* Hungary dead Nettle or the Dragon flower.

Let mee thrust this plant into this place, rather then make a peculiar Chapter, because I haue no other of the same stocke or kindred to be ioyned with it, and is a pretty ornament in a Garden. The leaues whereof are very large, round, and great, rough or full of veines, which make it seeme crumpled, dented or deepely notched about the edges, and of a very darke greene colour, and sometimes browniſh, or of a darke reddish colour withall, euery one standing on a long foote-stalke, very like in forme vnto the great white Arch-Angell leaues, but farre larger and blacker: the stalkes are great and foure square, hauing leaues and flowers standing round about them at the ioynts like coronets, which flowers are very great, long, and wide gaping open, of a darke red or purple colour, with some whitenesse or spots in the iawes, and some hairinesse also on the sides, which stand in full flower two or three moneths most vsually, and sometimes longer, after which come browniſh seede: the roote is a great tuft or bush of long whitish strings, and encreaseth euery yeare, not fearing the greatest iniuries of our coldest and extreamest Winters.

### The Place.

All these plants are strangers in our Countrey, and onely preserued in Gardens, to furnish them with variety; but (as I said) the cloth of gold Moth Mullein hath been raised from seed in our owne Country.

### The Time.

The last flowreth first, before all the rest, beginning in Aprill. The Moth Mulleins in May and Iune. The French Sage in Iuly.

### The Names.

All the sorts of *Blattaria* may bee comprehended vnder the kindes of

*Verbascum nigrum*, as any one but meanely exercised in the knowledge of plants, may discerne. And although Plinie saith, that Moths doe most frequently haunt where *Blattaria* either groweth, or is laid, yet it is not obserued sufficiently in our Country so to doe, notwithstanding the name of Moth Mullein is generally giuen them. The last is generally called with vs *Lamium Pannonicum*, but certainely it is the *Galeosis maxima Pannonica* of Clusius.

The Vertues.

Other qualities I haue not found hath been allotted vnto the *Blattaria* or Moth Mullein, then those of Plinie, to engender Moths. Weevse none of these plants in Physicke in these daies.

## Chap. XCIX.

### *Valeriana.* Valerian.

THe many sorts of Valerian (or Set-wall as many doe call them) are fitter for a generall worke, or a generall Physicall Garden of Simples, then this of delightfull flowers. I will therefore select out a few, worthy of the place, and offer them to your considerations.

#### 1. *Valeriana rubra Dodonæi.* Red Valerian.

This Valerian hath diuers hard, but brittle whitish greene stalkes, rising from the roote, full of tuberous or swelling ioynts, whereat stand two leaues, on each side one, and now and then some small leaues from betweene them, which are somewhat long and narrow, broadest in the middle, and small at both ends, without either diuision or incisure on the edges, of a pale greene colour: the stalkes are branched at the top into diuers parts, at the ends whereof stand many flowers together, as it were in an vmbell or tuft, somewhat like vnto the flowers of our ordinary Valerian, but with longer neckes, and of a fine red colour, very pleasant to behold, but of no sent of any Valerian: after these flowers haue stood blowne a very great while, they sodainely fall away, and the seede is ripe very quickly after, which is whitish, standing vpon the branches naked, as the Valerians doe, and very like vnto them, with a little white doune at the end of euery one of them, whereby they are soone carried away with the winde: the roote is great, thicke, and white, continuing long, and shooting out new branches euery yeare, and smelling somewhat like a Valerian.

#### 2. *Nardus Montana tuberosa.* Knobbed Mountaine Valerian.

This kinde of Valerian or Spiknard, if you will so call it, hath his first leaues lying on the ground, without any diuision in them at all, being smooth, and of a dark greene colour, which so abide all the winter; but those that spring vp after, and when it runneth vp to flower, are cut in on the edges, very like vnto the iagged leaues of the great garden Valerian, and so the elder they grow, the more cut and iagged they are: the stalke and flowers are very like the stalke with flowers of the garden Valerian, but of a darke or deepe red colour, and more store of them thrust together, by double the number almost: the seede is like the seede of the great Valerian: the root is tuberous, or knobbed in many parts, round about, aboue and below also, with some fibres shooting from them, whereby it is encreased, and smelleth very like the roote of the garden Setwall, or not altogether so strong.

3. *Valeriana*

1 *Elatteria flore albo.* Moth Mullein with a white flower. 2 *Blattaria flore purpureo* Moth Mullein with a purple flower. 3 *Verbascum quartum Matthioli.* French Sage. 4 *Æthiopis.* Ethiopian Mullein. 5 *Valeriana rubra Dodonæi.* Red Valerian. 6 *Valeriana Græca.* Greek Valerian. 7 *Laminm Pannonicum.* Hungary dead Nettle. 8 *Cardamine flore pleno.* Double Cuckowe flower or Ladies smocks.

### 3. *Valeriana Græca.* Greeke Valerian.

The Greek Valerian hath many winged leaues lying vpon the ground, that is, many small leaues set on both sides of a middle ribbe, very like vnto the wilde Valerian, that groweth by the ditch sides, but much smaller and tenderer, among which rise vp one or two round brittle stalkes, two foote high or thereabouts, whereon are set at the ioynts, such like leaues as grow below, but smaller : the toppes of the stalkes are diuided into many small branches, thicke set together, full with flowers, consisting of fiue small round leaues a peece, layd open like vnto the Cinquefoile flower, with some white threds in the middle, tipt with yellow pendents : the colour of these flowers in some plants, is of a faire bleake blew colour, and in others pure white : And I doe heare of one beyond the Seas (if the report bee true, for I haue not seene such a one) which should beare red flowers : after the flowers are past, there come vp in their places small hard huskes or heads, containing small blackish seedes : the roote is composed of a number of small long blackish threds, fastened together at the head, without any sent at all of a Valerian, eyther in roote or leafe ; and why it should bee called a Valerian I see no great reason, for it agreeth with none of them, in flower or seede, and but onely with the wilde Valerian in leafe, as I said before : but as it is, we so giue it you, and for the flowers sake is receiued into our gardens, to helpe to fill vp the number of natures rarities and varieties.

#### The Place.

All these Valerians are strangers, but endenizond for their beauties sake in our Gardens. The Mountaine Valerian I had of the liberalitie of my louing friend Iohn Tradescante, who in his trauaile, and search of natures varieties, met with it, and imparted thereof vnto me.

#### The Time.

They flower in the Summer moneths, and seed quickly after.

#### The Names.

The first is generally called of most, *Valeriana rubra Dodonæi*, who saith also that some would haue it to be *Behen rubrum*. Some call it *Valerianthon*, others make it a kinde of *Ocimastrum*, and some *Saponaria altera*, with other names, which are to no great purpose to set downe in this place, it beeing fitter for a generall worke to discusse of names, wherein both reading, knowledge and iudgement must bee shewen, to correct errours, and set downe the truth, that one may rest thereon. The others haue their names in their titles sufficient to distinguish them.

#### The Vertues.

The Mountaine Valerian is of all the the rest here set downe of most vse in Physicke, the rest hauing little or none that I know, although it be much weaker then the great garden kinde, or the Indian Nardus, in whose steed anciently it was vsed, in oyles, oyntments, &c.

## CHAP. C.

### *Cardamine.* Cuckow flowers, or Ladies smockes.

OF the common sorts of Cuckow flowers that grow by ditch-sides, or in moist medowes, & wet grounds, it is not my purpose here to write, but of one or two other, the most specious or faire of all the tribe, that doe best befit this garden.

1. *Cardamine flore pleno.* Double Cuckow flowers.

The double *Cardamine* hath a few winged leaues, weake and tender, lying on the ground, very like vnto the single medow kinde; from among which riseth vp a round greene stalke, set here and there, with the like leaues that grow below, the top wherof hath a few branches, whereon stand diuers flowers, euery one vpon a small footestalk, consisting of many small whitish round leaues, a little dasht ouer with a shew of blush, set round together, which make a double flower: the roote creepeth vnder ground, sending forth small white fibres, and shooteth vp in diuers places.

2. *Cardamine trifolia.* Trefoile Ladies smockes.

This small plant hath diuers hard, darke round greene leaues, somewhat vneuen about the edges, alwayes three set together on a blackish small footstalke, among which rise vp small round blackish stalkes, halfe a foote high, with three small leaues at the ioynts, where they branch forth; at the toppes whereof stand many flowers, consisting of foure leaues a peece, of a whitish or blush colour very pale: after which come vp small, thicke and long pods, wherein is contained small round seede: the root is composed of many white threds, from the heads whereof runne out small strings, of a dark purple colour, whereby it encreaseth.

## The Place.

The first with the double flower is found in diuers places of our owne Countrey, as neere Micham about eight miles from London; also in Lancashire, from whence I receiued a plant, which perished, but was found by the industrie of a worthy Gentlewoman, dwelling in those parts heretofore remembred, called Mistresse Thomasin Tunstall, a great louer of these delights. The other was sent me by my especial good friend Iohn Tradescante, who brought it among other dainty plants from beyond the Seas, and imparted thereof a roote to me.

## The Time.

The last most vsually flowreth before the former, yet not much differing, that is, in the end of Aprill or in May.

## The Names.

The first is a double kinde of that plant, that growing wilde abroade, is vsually called *Cardamine altera*, and *Sisymbrium alterum* of Dioscorides, and of some *Flos cuculi*, but not fitly; for that name is more vsually giuen vnto the wilde featherd Campions, both single and double, as is before expressed: yet for want of a fitter name, wee may call it in English, eyther Cuckowe flower, or Ladyes smockes, which you will. The second hath beene sent vnder the name of *Sanicula trifolia*, but the most frequent name now receiued, is *Cardamine trifolia*, and in English Trefoile Ladies spockes.

## The Vertues.

The double Ladies smockes are of the same qualitie with the single, and is thought to be as effectuall as Watercresses. The propertie of the other I thinke is not much knowne, although some would make it a wound herbe.

## Chap. CI.

### *Thlaspi Creticum.* Candy Tufts.

OF the many sorts of *Thlaspi* it is not the scope of this worke to relate, I will select but onely two or three, which for their beautie are fit to bee inserted into this garden.

*Thlaspi Creticum vmbellatum flore albo & purpureo.*
Candy Tufts white and purple.

This small plant riseth seldome aboue a foote and a halfe high, hauing small, narrow, long and whitish greene leaues, notched or dented with three or foure notches on each side, from the middle to the point-wards; from among which rise vp the stalkes, branched from the bottome almost into diuers small branches, at the toppes whereof stand many small flowers, thick thrust together in an vmbell or tuft, making them seeme to be small, round, double flowers of many leaues, when as euery flower is single, and standeth a part by it selfe, of a faire white colour in some plants, without any spot, and in others with a purplish spot in the centre or middle, as if some of the middle leaues were purple; in others againe the whole flower is purplish all ouer, which make a pretty shew in a garden: the seede is contained in many small and flat seed vessels, which stand together in an vmbell, as the flowers did, in which are contained somewhat reddish seede, like vnto some other sorts of *Thlaspi*, called Treakle Mustards: the roote is small and hard, and perisheth euery yeare hauing giuen seede.

*Thlaspi Marinum Baticum.* We haue another sort, whose leaues before it sendeth forth any stalke, are a little toothed, or finely dented about the edges, and brancheth not so much out, but carryeth an vmbell of purplish flowers like vnto the former, and paler yellow seede.

#### The Place.

These doe grow in Spaine and Candie, not farre from the Sea side.

#### The Time.

These *Thlaspi* giue not their flowers vntill the end of Iune, or beginning of Iuly, and the seed is ripe soone after.

#### The Names.

The first is named by some, *Draba*, or *Arabis*, as Dodonæus, but *Draba* is another plant differing much from this. We call one sort, *Thlaspi Creticum*, and the other *Thlaspi Baticum marinum*, because the one came from Spaine, and the other from Candy; we giue it in English, the name of Tufts, because it doth fit the forme of the flowers best, although ordinarily all the *Thlaspi* are Englished Wilde Mustardes.

#### The Vertues.

Candy, or Spanish Tufts, is not so sharpe biting in taste, as some other of the Thlaspies are, and therefore is not to be vsed in medicines, where *Thlaspi* should be in the stead thereof.

## Chap. CII.

### *Clematis.* Clamberers, or Creepers.

HAuing shewed you all my store of herbes bearing fine flowers, let mee now bring to your consideration the rest of those plants, be they Shrubs or Trees, that are cherished in our garden, for the beauty of their flowers chiefly, or for some other beautifull respect: and first I will begin with such as creepe on the ground, without climing, and then such as clime vp by poles, or other things, that are set or grow neere them, fit to make Bowers, and Arbours, or else are like them in forme, in name, or some other such qualitie or propertie.

#### 1. *Clematis Daphnoides, sive Vinca peruinca simplex minor diuersorum colorum.*
Single Perwinkle of diuers colours.

The smaller Perwinkle which not onely groweth wilde in many places, but is most vsuall in our Gardens, hath diuers creeping branches, trayling or running vpon the ground, shooting out small fibres at the ioynts, as it creepeth, taking thereby hold in the ground, and rooteth in diuers places: at the ioynts of these branches stand two small darke greene shining leaues, somewhat like vnto small Baye leaues, but smaller, and at the ioynts likewise with the leaues, come forth the flowers, one at a ioynt, standing vpon a tender footestalke, being somewhat long and hollow, parted at the brims, sometimes into foure leaues, and sometimes into fiue, the most ordinary sort is of a pale or bleake blew colour, but some are pure white, and some of a darke reddish purple colour: the root is in the body of it, little bigger then a rush, bushing in the ground, and creeping with his branches farre about, taking roote in many places, whereby it quickely possesseth a great compasse; and is therefore most vsually planted vnder hedges, or where it may haue roome to runne.

#### 2. *Vinca peruinca flore duplici purpurea.*
Double purple Perwinkle.

The double Perwinkle is like vnto the former single kinde, in all things except in the flower, which is of that darke reddish purple colour that is in one of the single kindes; but this hath another row of leaues within the flower, so that the two rowes of leaues causeth it to be called double, but the leaues of these are lesser then the single. I haue heard of one with a double white flower, but I haue not yet seene it.

#### 3. *Clematis Daphnoides siue Peruinca maior.*
The greater Perwinkle.

This greater Perwinkle is somewhat like the former, but greater, yet his branches creepe not in that manner, but stand more vpright, or lesse creeping at the least: the leaues also hereof stand by couples at the ioynts, but they are broader and larger by the halfe: the flowers are larger, consisting of fiue leaues that are blew, a little deeper then the former blew: this plant is farre tenderer to keepe then the other, and therefore would stand warme, as well as in a moist shadowie place.

#### 4. *Clematis altera siue vrens flore albo.*
Burning Clamberer, or Virgins Bower.

This Caustick or burning Climer, hath very long and climing tender branches, yet somewhat woody below, which winde about those things that stand neere it, couered with a browniish greene barke, from the ioynts whereof shoote forth many winged leaues, consisting for the most part of fiue single leaues, that is, two and two together, and one at the end, which are a little cut in or notched on the edges here and there,

there, but euery part of them is lesser then the leaues of the next following Climer, without any clasping tendrels to winde about any thing at all: towards the vpper part of the branches, with the said leaues, come forth long stalks, wheron stand many white flowers clustering together, opening the brims into sixe or eight small leaues, spreading like a starre, very sweet of smell, or rather of a strong heady sent, which after turne into flattish and blackish seede, plumed at the head, which plume or feather flyeth away with the winde after it hath stood long, and leaueth the seede naked or bare: the roote is white and thicke, fleshie and tender, or easie to be broken, as my selfe can well testifie, in that desiring to take a sucker from the roote, I could not handle it so tenderly, but that it broke notwithstanding all my care. Master Gerard in his Herball maketh mention of one of this kinde with double white flowers, which hee saith he recouered from the seede was sent him from Argentine, that is Strasborough, whereof hee setteth forth the figure with double flowers; but I neuer saw any such with him, neither did I euer heare of any of this kinde with double flowers. Clusius indeed saith, that hee receiued from a friend some seede vnder the name of *Clematis flore albo pleno*: but he doubteth whether there bee any such: the plants that sprang with him from that seede, were like vnto the vpright kinde called *Flammila Matthioli*, or *Iouis cresta*, as he there saith: but assuredly I haue beene informed from some of my especiall friends beyond Sea, that they haue a double white *Clematis*, and haue promised to send it; but whether it will be of the climing or vpright sort, I cannot tell vntill I see it: but surely I doe much doubt whether the double will giue any good seede.

### 5. *Clematis altera sine peregrina flore rubro.* Red Ladies Bower.

This Climer hath many limber and weake climing branches like the former, couered with a browne thin outer barke, and greene vnderneath: the leaues stand at the ioynts, consisting but of three leaues or parts, whereof some are notched on one side, and some on both, without any clasping tendrels also, but winding with his branches about any thing standeth next vnto it: the flowers in like manner come from the same ioynts with the leaues, but not so many together as the former vpon long footstalkes, consisting of foure leaues a peece, standing like a crosse, of a darke red colour; the seed is flat and round, and pointed at the end, three or foure or more standing close together vpon one stalk, without any doune vpon them at all, as in the former: the roots are a bundell of browniish yellow strong strings, running down deep into the ground, from a bigge head aboue.

### 6. *Clematis peregrina flore purpureo simplici.* Single purple Ladies Bower.

This Ladies Bower differeth in nothing from the last described, but onely in the colour of the flower, which is of a sad blewish purple colour; so that the one is not possible to be known from the other, vntill they be in flower.

### 7. *Clematis peregrina flore purpureo pleno.* Double flowred purple Ladies Bower.

This double *Clematis* hath branches and leaues so neere resembling the single kinds, that there can be knowne no difference, vnlesse it be, that this groweth more goale and great, and yeeldeth both more store of branches from the ground, and more spreading aboue: the chiefest marke to distinguish it is the flower, which in this is very thicke and double, consisting of a number of smaller leaues, set close together in order in the middle, the foure outermost leaues that encompasse them, being much broader and larger then any of the inward, but all of a dull or sad blewish purple colour, the points or ends of the leaues seeming a little darker then the middle of them: this beareth no seede that euer I could see, heare of, or learne by any of credit, that haue noursed it a great while; and therefore the tales of false deceitfull gardiners, and others, that diliuer such for truth, to deceiue persons ignorant thereof, must not bee credulously entertained.

In

In the great booke of the Garden of the Bishop of Eystot (which place is neere vnto Noremberg) in Germany, I reade of a *Clematis* of this former kinde, whose figure is thereto also annexed, with double flowers of an incarnate, or pale purple tending to a blush colour, whereof I haue not heard from any other place. *clematis peregrina flore carneo caruleo.*

8. *Flammula Iouis erecta.* Vpright Virgins Bower.

This kinde of *Clematis* hath diuers more vpright stalkes then any of the foure last described, sometimes foure or fiue foote high, or more; yet leaning or bending a little, so that it had some neede of sustaining, couered with a browniſh barke; from whence come forth on all sides diuers winged leaues, consisting of fiue or seuen leaues, set on both sides of a middle ribbe, whereof one is at the end: the tops of the stalkes are diuided into many branches, bearing many white sweet smelling flowers on them, like in fashion vnto the white Virgins Bower; after which come such like feather topt seede, which remaine and shew themselues, being flat like the other, when the plumes are blowne abroad: the roote spreadeth in the ground from a thicke head, into many long strings, and fasteneth it selfe strongly in the earth; but all the stalkes dye downe euery yeare, and spring afresh in the beginning of the next.

9. *Clematis caerulea Pannonica.* The Hungarian Climer.

The stalks of this plant stand vpright, & are foure square, bearing at euery ioynt two leaues, which at the first are closed together, and after they are open, are somewhat like vnto the leaues of *Asclepias*, or Swallow-wort: from the tops of the stalks, and sometimes also from the sides by the leaues commeth forth one flower, bending the head downward, consisting of foure leaues, somewhat long & narrow, standing like a crosse, and turning vp their ends a little againe, of a faire blew or skie colour, with a thicke pale yellow short thrumme, made like a head in the middle: after the flower is past, the head turneth into such a like round feather topt ball, as is to be seene in the Trauellers ioy, or *Viorna* (as it is called) that groweth plentifully in Kent, and in other places by the way sides, and in the hedges, wherein is included such like flat seede. These stalkes (like as the last) dye downe to the ground euery yeare, and rise againe in the Spring following, shooting out new branches, and therby encreaseth in the root.

10. *Maracoc siue Clematis Virginiana.* The Virginia Climer.

Because this braue and too much desired plant doth in some things resemble the former Climers, so that vnto what other family or kindred I might better conioyne it I know not; let me I pray insert it in the end of their Chapter, with this description. It riseth out of the ground (very late in the yeare, about the beginning of May, if it be a plant hath risen from the seed of our owne sowing, and if it be an old one, such as hath been brought to vs from Virginia, not till the end thereof) with a round stalke, not aboue a yard and a halfe high (in any that I haue seene) but in hotter Countries, as some Authors haue set it downe, much higher, bearing one leafe at euery ioynt, which from the ground to the middle thereof hath no claspers, but from thence vpwards hath at the same ioynt with the leafe both a small twining clasper, like vnto a Vine, and a flower also: euery leafe is broad at the stalke thereof, and diuided about the middle on both sides, making it somewhat resemble a Figge leafe, ending in three points, whereof the middlemost is longest: the bud of the flower, before it doe open, is very like vnto the head or seede vessell of the ordinary single *Nigella*, hauing at the head or top fiue small crooked hornes, which when this bud openeth, are the ends or points of fiue leaues, that are white on the inside, and lay themselues flat, like vnto an Anemone, and are a little hollow like a scoope at the end, with fiue other smaller leaues, and whiter then they lying betweene them, which were hid in the bud before it opened, so that this flower being full blowne open, consisteth of ten white leaues, laide in order round one by another: from the bottome of these leaues on the inside, rise diuers twined threads, which spread and lay themselues all ouer these white leaues, reaching beyond the points of them a little, and are of a reddish peach colour: towards the bot-

tomes

tomes likewise of these white leaues there are two red circles, about the breadth of an Oten strawe, one distant from another (and in some flowers there is but one circle seen) which adde a great grace vnto the flower; for the white leaues shew their colour through the peach coloured threads, and these red circles or rings vpon them being also perspicuous, make a tripartite shew of colours most delightfull: the middle part of this flower is hollow, and yellowish; in the bottome whereof riseth vp an vmbone, or round stile, somewhat bigge, of a whitish greene colour, spotted with reddish spots like the stalkes of Dragons, with fiue round threads or chiues, spotted in the like manner, and tipt at the ends with yellow pendents, standing about the middle part of the said vmbone, and from thence rising higher, endeth in three long crooked hornes most vsually (but sometimes in foure, as hath beene obserued in Rome by Dr. Aldine, that set forth some principall things of Cardinall Farnesius his Garden) spotted like the rest, hauing three round greene buttons at their ends: these flowers are of a comfortable sweete sent, very acceptable, which perish without yeelding fruit with vs, because it flowreth so late: but in the naturall place, and in hot Countries, it beareth a small round whitish fruit, with a crowne at the toppe thereof, wherein is contained (while it is fresh, and before it be ouer dried) a sweet liquor, but when it is dry, the seede within it, which is small, flat, somewhat rough and blacke, will make a ratling noise: the rootes are composed of a number of exceeding long and round yellowish browne strings, spreading farre abroad vnder the ground (I haue seene some rootes that haue beene brought ouer, that were as long as any rootes of *Sarsa parilla*, and a great deale bigger, which to be handsomely laid into the ground, were faine to be coyled like a cable) and shooting vp in seuerall places a good distance one from another, whereby it may be well encreased.

The Iesuites Figure of the Maracoc.

GRANADILLVS FRVTEX INDICVS CHRISTI PASSIONIS IMAGO.

## The Place.

The first blew Perwinkle groweth in many Woods and Orchards, by the hedge sides in England, and so doth the white here and there, but the other single and double purple are in our Gardens onely. The great Perwinkle groweth in Prouence of France, in Spaine, and Italy, and other hot Countries, where also growe all the twining Clamberers, as well single as double: but both the vpright ones doe growe in Hungary and thereabouts. The surpassing delight of all flowers came from Virginia. Wee preserue them all in our Gardens.

## The Time.

The Perwinkles doe flower in March and Aprill. The Climers not vntill the end of Iune, or in Iuly, and sometimes in August. The Virginian somewhat later in August; yet sometimes I haue knowne the flower to shew it selfe in Iuly.

## The Names.

The first is out of question the first *Clematis* of Dioscorides, and called of many

1 *Thalspi creticum.* Candy tufts. 2 *Vinca pervinca flore simplici.* Single Perwinkle. 3 *Vinca pervinca flore duplici.* Double Perwinkle. 4 *Flammula Matthioli.* Vpright Virgins Bower. 5 *Clematis peregrina flore simplici.* The single Ladies Bower. 6 *clematis peregrina flore pleno purpureo.* Double flowred Ladies Bower. 7 *Maracoc sive Clematis Virginiana.* The Virginian Climer.

many *Clematis Daphnoides* (but not that plant that is simply called *Daphnoides*, for that is *Laureola*) and is vsually called *Vinca pervinca*: but it is not *Chamædaphne*, for that is another plant, as shall be shewed in his place; some call it *Centunculus*: In English wee call it Perwinkle. The other is *Clematis altera* of Dioscorides, and is called also *Clematis peregrina*, whose distinctions are set downe in their titles: In English, Ladies Bower, or Virgins Bower, because they are fit to growe by Arbours, to couer them. The first vpright Clamberer is called, and that rightly of some, *Clematis erecta*, or *surrecta*. Of others, *Flammula frutex*, and *Flammula Iouis*, or *surrecta*: In English, Vpright Virgins Bower. The next is called by Clusius, *Clematis Pannonica cærulea*, who thought it to be *Climeni species*, by the relation of others, at the first, but after entituled it, *Clematis*: In English, the Hungarian Climer. The last may be called in Latine, *Clematis Virginiana*: In English, The Virgin or Virginian Climer; of the Virginians, *Maracoc*: of the Spaniards in the West Indies *Granadilla*, because the fruit (as is before said) is in some fashion like a small Pomegranate on the outside; yet the seede within is flattish, round, and blackish. Some superstitious Iesu-ite would faine make men beleeue, that in the flower of this plant are to be seene all the markes of our Sauiours Passion; and therefore call it *Flos Passionis*: and to that end haue caused figures to be drawne, and printed, with all the parts proportioned out, as thornes, nailes, speare, whippe, pillar, &c. in it, and all as true as the Sea burnes, which you may well perceiue by the true figure, taken to the life of the plant, compared with the figures set forth by the Iesuites, which I haue placed here likewise for euery one to see: but these bee their aduantagious lies (which with them are tolerable, or rather pious and meritorious) wherewith they vse to instruct their people; but I dare say, God neuer willed his Priests to instruct his people with lyes: for they come from the Diuell, the author of them. But you may say I am beside my Text, and I am in doubt you will thinke, I am in this besides my selfe, and so nothing to be beleeued herein that I say. For, for the most part, it is an inherent errour in all of that side, to beleeue nothing, be it neuer so true, that any of our side shall affirme, that contrarieth the assertions of any of their Fathers, as they call them: but I must referre them to God, and hee knoweth the truth, and will reforme or deforme them in his time. In regard whereof I could not but speake (the occasion being thus offered) against such an erroneous opinion (which euen Dr. Aldine at Rome, before remembred, disproued, and contraried both the said figures and name) and seek to disproue it, as doth (I say not almost, but I am affraid altogether) leade many to adore the very picture of such things, as are but the fictions of superstitious brains: for the flower it selfe is farre differing from their figure, as both Aldine in the aforesaid booke, and Robinus at Paris in his *Theatrum Floræ*, doe set forth; the flowers and leaues being drawne to the life, and there exhibited, which I hope may satisfie all men, that will not be perpetually obstinate and contentious.

## The Vertues.

Costæus saith hee hath often seene, that the leaues of Perwinkle held in the mouth, hath stayed the bleeding at the nose. The French doe vse it to stay the menstruall fluxes. The other are causticke plants, that is, fiery hot, and blistering the skinne; and therefore (as Dioscorides saith) is profitable to take away the scurfe, leprye, or such like deformities of the skin. What property that of Virginia hath, is not knowne to any with vs I thinke, more then that the liquor in the greene fruit is pleasant in taste; but assuredly it cannot be without some speciall properties, if they were knowne.

## CHAP. CIII.

### *Chamælea.* Dwarfe Spurge Oliue, or Dwarfe Baye.

I Haue three sorts of *Chamælea* to bring to your consideration, euery one differing notably from other; two of them of great beauty in their flowers, as well as in the whole plant: the third abiding with greene leaues, although it haue no beauty in the flower, yet worthy of the place it holds. And vnto these I must adioyne another plant, as comming neerest vnto them in the brauery of the flowers.

#### 1. *Chamælea Germanica siue Mezereon floribus dilutioris coloris & saturatioris.* Dwarfe Baye, or flowring Spurge Oliue.

We haue two sorts of this Spurge Oliue or Dwafe Baye, differing onely in the colour of the flowers. They both rise vp with a thicke wooddy stemme, fiue or six foot high sometimes, or more, and of the thickneſſe (if they be very old) of a mans wrest at the ground, spreading into many flexible long branches, couered with a tough grayish barke, beset with small long leaues, somewhat like vnto Priuet leaues, but smaller and paler, and in a manner round pointed: the flowers are small, consisting of foure leaues, many growing together sometimes, and breaking out of the branches by themselues: in the one sort of a pale red at the first blowing, and more white afterwards; the other of a deeper red in the blossome, and continuing of a deeper red colour all the time of the flowring, both of them very sweete in smell: after the flowers are past, come the berries, which are greene at the first, and very red afterwards, turning blackish red, if they stand too long vpon the branches: the rootes spread into many tough long branches, couered with a yellowish barke.

#### 2. *Chamælea Alpina.* Mountaine Spurge Oliue.

This Mountaine Laurell riseth vp with a small wooddy stemme, three or foure foot high, or more, branching forth towards the vpper parts into many slender and tough branches, couered with a rough hoary greene barke, beset at the ends thereof with flatter, fuller, and smaller round pointed leaues then the former, of a grayish greene colour on the vpperside, and hoary vnderneath, which abide on the branches in Winter, and fall not away as the former: the flowers are many set together at the ends of the branches, greater then the former, and consisting of foure leaues a peece, of a light blush colour, standing in small grayish huskes, of little or no sent at all: the fruit followeth, which are small long graines or berries, of an excellent red colour, which afterwards turne blacke: the roote is long, and spreadeth about vnder the vpper part of the earth.

#### 3. *Chamælea tricocces.* Widowe Wayle.

This three berried Spurge Oliue hath no great stemme at all, but the whole plant spreadeth from the ground into many flexible tough greene branches, whereon are set diuers narrow, long, darke greene leaues all along the branches, which abide greene all the Winter: the flowers are very small, scarce to be seene, and come forth between the leaues and the stalke, of a pale yellow colour, made of three leaues; after which come small blackish berries, three vsually set together: the roote spreadeth it selfe in the ground not very farre, being hard and wooddy, and often dyeth, if it bee not well defended from the extremity of our sharpe Winters.

#### 4. *Cneorum Matthioli.* Small Rocke Roses.

I was long in doubt in what place I should dispose of this plant, whether among the Campions, as Bauhinus, or among these, as Clusius doth; but lest my Gorden should want it wholly, let it take vp roome for this time here. This gallant plant hath diuers

long, weake, slender, but yet tough branches lying vpon the ground, diuided vsually into other smaller branches, whereon growe many, small, long, and somewhat thicke leaues, somewhat like vnto the leaues of the former *Mezereon*, set without any order to the very tops, from whence doe come forth a tuft of many small flowers together, made or consisting of foure leaues a peece, of a bright red or carnation colour, and very sweete withall, which turne into small round whitish berries, wherein is contained small round seede, couered with a grayish coate or skinne: the roote is long and yellowish, spreading diuers wayes vnder the ground, and abideth many yeares shooting forth new branches.

*Flore albo.* It hath beene obserued in some of these plants, to bring forth white flowers, not differing in any thing else.

## The Place.

The first sorts growe plentifully in many places of Germany. The second in the mountaines by Sauoye. The third in Prouence and Spaine. The last in diuers parts of Germany, Bohemia, and Austria, and about Franckford.

## The Time.

The two first sorts are most vsually in flower about Christmas, or in Ianuary, if the weather be not violent, and sometimes not vntill February. The second flowreth not vntill Aprill. The third in May. The berries of them ripen some in Iune and Iuly; some in August and September, as their flowring is earlier or later. The last flowreth as well in the Spring as in Autumne, so apt and plentifull it is in bearing, and the seede at both times doth ripen soone after.

## The Names.

The first is called of some *Chamelea*, with this addition *Germanica*, that it may differ from the third, which is the true *Chamelea* of Dioscorides, as all the best Authors doe agree, and is also called *Piper montanum* of the Italians. It is generally called *Mezereon*, and is indeede the true *Mezereon* of the Arabians, and so vsed in our Apothecaries shops, whersoeuer the Arabians *Mezereon* is appointed, although the Arabians are so intricate and vncertaine in the descriptions of their plants, confounding *Chamelea* and *Thymelea* together. Matthiolus maketh it to be *Daphnoides* of Dioscorides; but in my opinion he is therein mistaken: for all our best moderne Writers doe account our *Laureola*, which hath blacke berries, to bee the true *Daphnoides*: the errour of his Countrey might peraduenture drawe him thereunto; but if hee had better considered the text of Dioscorides, that giueth black berries to *Daphnoides*, and red to *Chamædaphne*, he would not so haue written; and truly, I should thinke (as Lobel doth) with better reason, that this *Chamelea* were Dioscorides *Chamædaphne*, then hee to say it were *Daphnoides*: for the description of *Chamædaphne*, may in all parts be very fitly applyed to this *Chamelea*: and euen these words, *Semen annexum folijs*, wherein may be the greatest doubt in the description, may not vnfitly bee construed, that as is seene in the plant, the berries growe at the foote of the leaues, about the branches: the faculties indeede that Dioscorides giueth to *Chamædaphne*, are (if any repugnancie be) the greatest let or hinderance, that this *Chamelea* should not be it: but I leaue the discussing of these and others of the like nature, to our learned Physitians; for I deale not so much with vertues as with descriptions. The second is called of Lobel *Chamælea Alpina incana*, of Clusius *Chamælea secunda*, and saith hee had it out of Italy. Wee may call it in English, Mountaine Spurge Oliue, as it is in the description, or Mountaine Laurell, which you will. The last hath the name of *Cneorum*, first giuen it by Matthiolus, which since is continued by all others. Bauhinus (as I said) referreth it to the Mountaine Campions, but Clusius

(as

1 *Chamelæa Germanica seu Mezereon.* Mezereon or Dwarfe Bay. 2 *Chamelæa Alpina.* Mountain Spurge Oliue. 3 *Cneorum Matthioli.* Small Rocke Roses. 4 *Laurus Tinus sive siluestris.* The wild Bay tree. 5 *Oleander sive Laurus Rosea.* The Rose Bay tree. 6 *Laurocerasus.* The Bay Cherrie tree.

(as I doe) to the kindes of *Chamælæa* or *Thymælæa*. For want of an Engliſh name I haue (as you ſee, and that is according to the name the Germane women, as Cluſius ſaith, doe call it) entituled it the Small Rocke Roſe; which may abide vntill a fitter may be conferred vpon it.

### The Vertues.

All theſe plants except the laſt, as well leaues as berries, are violent purgers, and therefore great caution is to bee had in the vſe of them. The laſt hath not beene applyed for any diſeaſe that I know.

## Chap. CIII.

### *Laurus*. The Bay Tree.

MY meaning is not to make any deſcription of our ordinary Bayes in this place (for as all may very well know, they may be for an Orchard or Courtyard, and not for this Garden) but of two or three other kindes, whoſe beautifull aſpect haue cauſed them to be worthy of a place therein: the one is called *Laurus Tinus*, The wilde Baye: the other *Laurus Roſea* or *Oleander*, The Roſe Bay: and a third is *Lauroceraſus*, The Cherry Bay; which may haue not onely ſome reſpect for his long buſh of ſweet ſmelling flowers, but eſpecially for the comely ſtatelineſſe of his gallant euer freſh greene leaues; and the rather, becauſe with vs in moſt places, it doth but *fruteſcere*, vſe to bee Shrub high, not *arboreſcere*, Tree high, which is the more fit for this Garden.

#### 1. *Laurus Tinus ſiue ſilueſtris*. The wilde Bay tree.

This wilde Baye groweth ſeldome to bee a tree of any height, but abideth for the moſt part low, ſhooting forth diuers ſlender branches, whereon at euery ioynt ſtand two leaues, long, ſmooth, and of a darke greene colour, ſomewhat like vnto the leaues of the Female Cornell tree, or between that and Baye leaues: at the toppes of the branches ſtand many ſmall white ſweete ſmelling flowers, thruſting together, as it were in an vmbell or tuft, conſiſting of fiue leaues a peece, the edges whereof haue a ſhew of a waſh purple, or light bluſh in them, which for the moſt part fall away without bearing any perfect ripe fruit in our Countrey: Yet ſometimes it hath ſmall black berries, as if they were good, but are not. In his naturall place it beareth ſmall, round, hard and pointed berries, of a ſhining blacke colour, for ſuch haue come often to my hands (yet Cluſius writeth they are blew); but I could neuer ſee any ſpring that I put into the ground. This that I here deſcribe, ſeemeth to me to be neither of both thoſe that Cluſius ſaw growing in Spain and Potugall, but that other, that (as he ſaith) ſprang in the low Countreyes of Italian ſeede.

#### 2. *Laurus Roſea ſiue Oleander*. The Roſe Bay.

Of the Roſe Bay there are two ſorts, one bearing crimſon coloured flowers, which is more frequent, and the other white, which is more rare. They are ſo like in all other things, that they neede but one deſcription for both. The ſtemme or trunke is many times with vs as bigge at the bottome as a good mans thumbe, but growing vp ſmaller, it diuideth it ſelfe into branches, three for the moſt part comming from one ioynt or place, and thoſe branches againe doe likewiſe diuide themſelues into three other, and ſo by degrees from three to three, as long as it groweth: the loweſt of theſe are bare of leaues, hauing ſhed or loſt them by the cold of winters, keeping onely leaues on the vppermoſt branches, which are long, and ſomewhat narrow, like in forme vnto Peach leaues, but thicker, harder, and of a darke greene colour on the vpperſide, and yellowiſh

yellowish greene vnderneath: at the tops of the young branches come forth the flowers, which in the one sort before they are open, are of an excellent bright crimson colour, and being blowen, consist of foure long and narrow leaues, round pointed, somewhat twining themselues, of a paler red colour, almost tending to blush, and in the other are white, the greene leaues also being of a little fresher colour: after the flowers are past, in the hot countries, but neuer in ours, there come vp long bending or crooked flat pods, whose outward shell is hard, almost woody, and of a browne colour, wherein is contained small flat browniſh seede, wrapped in a great deale of a browniſh yellow doune, as fine almost as silke, somewhat like vnto the huskes of *Asclepias*, or *Periploca*, but larger, flatter and harder; as my selfe can testifie, who had some of the pods of this Rose bay, brought mee out of Spaine, by Master Doctor Iohn More, the seedes whereof I sowed, and had diuers plants that I raised vp vnto a reasonable height, but they require, as well old as young, to bee defended from the colde of our winters.

### 3. *Laurocerasus.* The Bay Cherry.

This beautifull Bay in his naturall place of growing, groweth to bee a tree of a reasonable bigneſse and height, and oftentimes with vs also if it bee pruined from the lower branches; but more vsually in these colder Countries, it groweth as a shrub or hedge bush, shooting forth many branches, whereof the greater and lower are couered with a darke grayish greene barke, but the young ones are very greene, whereon are set many goodly, faire, large, thicke and long leaues, a little dented about the edges, of a more excellent fresh shining greene colour, and farre larger then any Bay leafe, and compared by many to the leaues of the *Pomeritron* tree (which because wee haue none in our Countrey, cannot be so well known) both for colour and largeneſse, which yeeld a most gracefull aspect: it beareth long stalkes of whitish flowers, at the ioynts of the leaues both along the branches and towards the ends of them also, like vnto the Birds Cherry or *Padus Theophrasti*, which the French men call *Putier & Cerisier blanc*, but larger and greater, consisting of fiue leaues with many threds in the middle: after which commeth the fruite or berries, as large or great as Flanders Cherries, many growing together one by another on a long stalke, as the flowers did, which are very blacke and shining on the outside, with a little point at the end, and reasonable sweete in taste, wherein is contained a hard round stone, very like vnto a Cherry stone, as I haue obserued as well by those I receiued out of Italie, as by them I had of Master Iames Cole a Merchant of London lately deceased, which grew at his house in Highgate, where there is a faire tree which hee defended from the bitterneſse of the weather in winter by casting a blanket ouer the toppe thereof euery yeare, thereby the better to preserue it.

#### The Place.

The first is not certainly knowne from whence it came, and is communicated by the suckers it yeeldeth. The second groweth in Spaine, Italie, Greece, and many other places: that with white flowers is recorded by Bellonius, to grow in Candy. The last, as Matthiolus, and after him Clusius report, came first from Constantinople: I had a plant hereof by the friendly gift of Master Iames Cole, the Merchant before remembred, a great louer of all rarities, who had it growing with him at his countrey house in Highgate aforesaid, where it hath flowred diuers times, and borne ripe fruit also.

#### The Time.

The first flowreth many times in the end of the yeare before Christmas, and often also in Ianuary, but the most kindly time is in March and Aprill, when the flowers are sweetest. The second flowreth not vntill Iuly. The last in May, and the fruit is ripe in August and September.

### The Names.

The first is called *Laurus silvestris*, and *Laurus Tinus* : in English Wilde Bay, or Sweete flowring Bay. The second is called *Laurus Rosea, Oleander, Nerium*, and *Rhododendros* : in English The Rose Bay, and Oleander. The last was sent by the name of *Trebezon Curmasi*, that is to say, *Dactylus Trapezuntina*, but not hauing any affinitie with any kinde of Date, Bellonius as I thinke first named it *Laurocerasus*, and *Cerasus Trapezuntina*. Dalechampius thinketh it to bee *Lotus Aphricana*, but Clusius refuteth it. Those stones or kernels that were sent me out of Italie, came by the name of *Laurus Regia*, The Kings Bay. Wee may most properly call it according to the Latine name in the title, The Cherry bay, or Bay Cherry, because his leaues are like vnto Bay-leaues, and both flowers and fruit like vnto the Birdes Cherry or Cluster Cherry, for the manner of the growing; and therfore I might more fitly I confesse haue placed it in my Orchard among the sorts of Cherries : but the beautifulnesse of the plant caused mee rather to insert it here.

### The Vertues.

The wilde Bay hath no propertie allotted vnto it in Physicke, for that it is not to be endured, the berries being chewed declare it to be so violent hot and choking. The Rose Bay is said by Dioscorides, to be death to all foure footed beasts, but contrariwise to man it is a remedie against the poison of Serpents, but especially if Rue bee added vnto it. The Cherry Bay is not knowne with vs to what physicke vse it may be applyed.

---

## Chap. CIIII.

*Cerasus flore multiplici.* The Rose or double blossomd Cherry.
*Malus flore multiplici.* The double blossomd Apple tree. And
*Malus Persica flore multiplici.* The double blossomd Peach tree.

THe beautifull shew of these three sorts of flowers, hath made me to insert them into this garden, in that for their worthinesse I am vnwilling to bee without them, although the rest of their kindes I haue transferred into the Orchard, where among other fruit trees, they shall be remembred : for all these here set downe, seldome or neuer beare any fruite, and therefore more fit for a Garden of flowers, then an Orchard of fruite.

### *Cerasus flore pleno vel multiplici.*
### The Rose Cherry, or double blossomd Cherry.

The double blossomed Cherry tree is of two sorts for the flower, but not differing in any other part, from the ordinary English or Flanders Cherry tree, growing in the very like manner : the difference consisteth in this, that the one of these two sorts hath white flowers lesse double, that is, of two rowes or more of leaues, and the other more double, or with more rowes of leaues, and besides I haue obserued in this greater double blossomd Cherry, that some yeares most of the flowers haue had another smaller and double flower, rising vp out of the middle of the other, like as is to bee seene in the double English Crow-foote, and double redde *Ranunculus* or Crowfoote, before described : this I say doth not happen euery yeare, but sometimes. Sometimes also these trees will giue a few berries, here and there scattered, and that with lesse double flowers more often, which are like vnto our English Cherries both for taste and bignesse. These be very fit to be set by Arbours.

*Malus*

1 *Cerasus flore pleno.* The double blossomd Cherry tree. 2 *Malus flore multiplici* The double blossomd Apple tree. 3 *Malus Persica flore pleno.* The double blossomd Peach tree. 4 *Periclymenum perfoliatum.* Double Honisuckle. 5 *Periclymenum rectum.* Vpright Honisuckle.

*Malus flore multiplici.* The double bloſſomd Apple tree.

This double bloſſomd Apple tree is altogether like vnto our ordinary Pippin tree in body, branch and leafe, the only difference is in the flower, which is altogether whitiſh, ſauing that the inner leaues towards the middle are more reddiſh, but as double and thicke as our double Damaske Roſes, which fall away without bearing fruit.

*Malus Perſica flore multiplici.* The double bloſſomd Peach tree.

This Peach tree for the manner of growing, is ſo like vnto an ordinary Peach tree, that vntill you ſee it in bloſſome you can perceiue no difference : the flower is of the ſame colour with the bloſſomes of the Peach, but conſiſting of three or foure, or more rowes of leaues, which fall often away likewiſe without bearing any fruite; but after it hath abiden ſome yeares in a place doth forme into fruite, eſpecially being planted againſt a wall.

The Place.

Both the Cherry trees are frequent in many places of England, nourſed for their pleaſant flowers. The Apple is as yet a ſtranger. And the Peach hath not been ſeen or knowne, long before the writing hereof.

The Time.

They all flower in April & May, which are the times of their other kinds.

The Names.

Their names are alſo ſufficiently expreſſed to know them by.

The Vertues.

Cherries, Peaches and Apples, are recorded in our Orchard, and there you ſhall finde the properties of their fruit: for in that theſe beare none or very few, their bloſſomes are of moſt vſe to grace and decke the perſons of thoſe that will weare or beare them.

## Chap. CV.

*Periclymenum.* Honyſuckles.

THe Honiſuckle that groweth wilde in euery hedge, although it be very ſweete, yet doe I not bring into my garden, but let it reſt in his owne place, to ſerue their ſenſes that trauell by it, or haue no garden. I haue three other that furniſh my Garden, one that is called double, whoſe branches ſpreade far, and being very fit for an arbour will ſoone couer it : the other two ſtand vpright, and ſpreade not any way far, yet their flowers declaring them to be Honiſuckles, but of leſſe delight, I conſort them with the other.

*Periclymenum perfoliatum ſive Italicum.* The double Honiſuckle.

The truncke or body of the double Honiſuckle, is oftentimes of the bigneſſe of a good ſtaffe, running out into many long ſpreading branches, couered with a whitiſh barke, which had neede of ſome thing to ſuſtaine them, or elſe they will fall down to the ground (and therefore it is vſually planted at an arbour, that it may run thereon,

or

or against a house wall, and fastened thereto in diuers places with nailes) from whence spring forth at seuerall distances, and at the ioynts, two leaues, being like in forme vnto the wilde Honisuckles, and round pointed for the most part; these branches diuiding themselues diuers wayes, haue at the toppes of them many flowers, set at certaine distances one aboue another, with two greene leaues at euery place, where the flowers doe stand, ioyned so close at the bottome, and so round and hollow in the middle, that it seemeth like a hollow cuppe or sawcer of flowers: the flowers stand round about the middle of these cuppes or sawcers, being long, hollow, and of a whitish yellow colour, with open mouthes dasht ouer with a light shew of purple, and some threds within them, very sweet in smell, like both in forme and colour vnto the common Honisuckles, but that these cuppes with the flowers in them are two or three standing one aboue another (which make a far better shew then the common, which come forth all at the heade of the branches, without any greene leaues or cuppes vnder them) and therefore these were called double Honisuckles.

*Periclymenum rectum fructu rubro.* Red Honisuckles.

This vpright Woodbinde hath a straight woody stemme, diuided into seuerall branches, about three or foure foote high, couered with a very thinne whitish barke, whereon stand two leaues together at the ioynts, being lesser then the former, smooth and plaine, and a little pointed: the flowers come forth vpon slender long footstalks at the ioynts where the leaues stand, alwayes two set together, and neuer more, but seldome one alone, which are much smaller then the former, but of the same fashion, with a little button at the foote of the flower; the buds of the flowers before they are open are very reddish, but being open are not so red, but tending to a kinde of yellowish blush colour: after which come in their places two small red berries, the one withered for the most part, or at least smaller then the other, but (as Clusius saith) in their naturall places they are both full and of one bignesse.

*Periclymenum rectum fructu cæruleo.* Blew berried Honisuckles.

This other vpright Woodbinde groweth vp as high as the former, or rather somewhat higher, couered with a blackish rugged barke, chapping in diuers places, the younger branches whereof are somewhat reddish, and couered with an hoary doune: the leaues stand two together at the ioints, somewhat larger then the former, and more whitish vnderneath: the flowers are likewise two standing together, at the end of a slender footestalke, of a pale yellowish colour when they are blowne, but more reddish in the bud: the berries stand two together as the former, of a darke blewish colour when they are fully ripe, and full of a red liquour or iuice, of a pleasant taste, which doth not only dye the hands of them that gather them, but serueth for a dying colour to the inhabitants where they grow plentifully, wherein are contained many flat seede: The roote is woody as the former is.

## The Place.

The firw groweth in Italie, Spaine, and Prouence of France, but not in the colder countreyes, vnlesse it be there planted, as is most frequent in our countrey. The others grow in Austria, and Stiria, as Clusius saith, and are entertained into their gardens onely that are curious.

## The Time.

The first flowreth vsually in Aprill, the rest in May.

## The Names.

The first is called *Periclymenum*, *Caprifolium perfoliatum*, and *Italicum*, as a difference from the common kinde: In English Double Woodbinde,

or

or double Honisuckles. The others, as they are rare, and little knowne, so are their names also: yet according to their Latine, I haue giuen them English names.

The Vertues.

The double Honisuckle is as effectuall in all things, as the single wilde kinde, and besides, is an especiall good wound herbe for the head or other parts. I haue not knowne the vpright kindes vsed in Physicke.

## Chap. CVI.

*Iasminum siue Gelsemium.* Iasmine or Gesmine.

WE haue but one sort of true Iasmine ordinarily in our Gardens throughout the whole Land; but there is another greater sort, which is farre more tender, brought out of Spaine, and will hardly endure any long time with vs, vnlesse it be very carefully preserued. Wee haue a third kinde called a yellow Iasmine, but differeth much from their tribe in many notable points: but because the flowers haue some likenesse with the flowers of the true Iasmine, it hath been vsually called a Iasmine; and therefore I am content for this Garden to conioyne them in one Chapter.

1. *Iasminum album.* The white Iasmine.

The white Iasmine hath many twiggy flexible greene branches, comming forth of the sundry bigger boughes or stems, that rise from the roote, which are couered with a grayish darke coloured barke, hauing a white pith within it like the Elder, but not so much: the winged leaues stand alwaies two together at the ioynts, being made of manie small and pointed leaues, set on each side of a middle ribbe, six most vsually on both sides, with one at the end, which is larger, more pointed then any of the rest, and of a darke greene colour: at the toppes of the young branches stand diuers flowers together, as it were in an vmbell or tuft, each whereof standeth on a long greene stalke, comming out of a small huske, being small, long, and hollow belowe, opening into fiue white small, pointed leaues, of a very strong sweete smell, which fall away without bearing any fruit at all, that euer I could learne in our Country; but in the hot Countries where it is naturall, it is said to beare flat fruit, like Lupines: the rootes spread farre and deepe, and are long and hard to growe, vntill they haue taken strong hold in the ground.

2. *Iasminum Catalonicum.* The Spanish Iasmine.

This Catalonia Iasmine groweth lower then the former, neuer rising halfe so high, and hath slender long greene branches, rising from the toppe of the wooddy stemme, with such like leaues set on them as the former, but somewhat shorter and larger: the flowers also are like vnto the former, and stand in the same manner at the end of the branches, but are much larger, being of a blush colour before they are blowne, and white with blush edges when they are open, exceeding sweete of smell, more strong then the former.

3. *Iasminum luteum, siue Trifolium fruticans alijs Polemonium.* The yellow Iasmine.

This that is called the yellow Iasmine, hath many long slender twiggy branches rising from the roote, greene at the first, and couered with a darke grayish barke afterwards, whereon are set at certaine distances, three small darke greene leaues together, the end leafe being alwaies the biggest: at the ioynts where the leaues come forth, stand

ſtand long ſtalkes, bearing long hollow flowers, ending in fiue, and ſome in ſix leaues, very like vnto the flowers of the firſt Iaſmine, but yellow, whereupon it is vſually called the Yellow Iaſmine: after the flowers are paſt, there come in their places round blacke ſhining berries, of the bigneſſe of a great Peaſe, or bigger, full of a purpliſh iuyce, which will dye ones fingers that bruiſe them but a little: the roote is tough, and white, creeping farre about vnder the ground, ſhooting forth plentifully, whereby it greatly encreaſeth.

### The Place.

The firſt is verily thought to haue been firſt brought to Spaine out of Syria, or thereabouts, and from Spaine to vs, and is to be ſeene very often, and in many of our Country Gardens. The ſecond hath his breeding in Spaine alſo, but whether it be his originall place we know not, and is ſcarce yet made well acquainted with our Engliſh ayre. The third groweth plentifully about Mompelier, and will well abide in our London Gardens, and any where elſe.

### The Time.

The firſt flowreth not vntill the end of Iuly. The ſecond ſomewhat earlier. The third in Iuly alſo.

### The Names.

The firſt is generally called *Iaſminum album*, and *Gelſeminum album*: In Engliſh, The white Iaſmine. The ſecond hath his name in his title, as much as may be ſaid of it. The third hath been taken of ſome to be a *Cytiſus*, others iudge it to be *Polemonium*, but the trueſt name is *Trifolium fruticans*, although many call it *Iaſminum luteum*: In Engliſh moſt vſually, The yellow Iaſmine, for the reaſons aforeſaid; or elſe after the Latine name, Shrubbie Trefoile, or Make-bate.

### The Vertues.

The white Iaſmines haue beene in all times accepted into outward medicines, eyther for the pleaſure of the ſweete ſent, or profit of the warming properties. And is in theſe dayes onely vſed as an ornament in Gardens, or for ſent of the flowers in the houſe, &c. The yellow Iaſmine, although ſome haue adiudged it to be the *Polemonium* of Dioſcorides, yet it is not vſed to thoſe purpoſes by any that I know.

## Chap. CVII.

### *Syringa.* The Pipe tree.

VNder the name of *Syringa*, is contained two ſpeciall kinds of Shrubs or Trees, differing one from another; namely, the *Lilac* of Matthiolus, which is called *Syringa cærulea*, and is of two or three ſorts: And the *Syringa alba*, which alſo is of two ſorts, as ſhall bee declared.

### 1. *Lilac ſiue Syringa cærulea.* The blew Pipe tree.

The blew Pipe tree riſeth ſometimes to be a great tree, as high and bigge in the bodie as a reaſonable Apple tree (as I haue in ſome places ſeene and obſerued) but moſt vſually groweth lower, with many twigs or branches riſing from the roote, hauing as much pith in the middle of them as the Elder hath, couered with a grayiſh greene barke,

barke, but darker in the elder branches, with ioynts set at a good distance one from another, and two leaues at euery ioynt, which are large, broad, and pointed at the ends, many of them turning or folding both the sides inward, and standing on long foote stalkes: at the toppes of the branches come forth many flowers, growing spike-fashion, that is, a long branch of flowers vpon a stalke, each of these flowers are small, long, and hollow belowe, ending aboue in a pale blewish flower, consisting of foure small leaues, of a pretty small sent: after the flowers are past, there come sometimes (but it is not often in our Country, vnlesse the tree haue stood long, and is grown great, the suckers being continually taken away, that it may growe the better) long and flat cods, consisting as it were of two sides, a thin skinne being in the midst, wherein are contained two long flattish red seede: the rootes are strong, and growe deepe in the ground.

2. *Syringa flore lacteo siue argenteo.* The siluer coloured Pipe tree.

This Pipe tree differeth not from the former blew Pipe tree, either in stemme or branches, either in leaues or flowers, or manner of growing, but onely in the colour of the flower, which in this is of a milke, or siluer colour, which is a kinde of white, wherein there is a thinne wash, or light shew of blew shed therein, comming somewhat neare vnto an ash-colour.

3. *Lilac laciniatis foliis.* The blew Pipe tree with cut leaues.

This Pipe tree should not differ from the first in any other thing then in the leaues, which are said to be cut in on the edges into seuerall parts, as the relation is giuen *à viris fide dignis*; for as yet I neuer saw any such; but I here am bold to set it downe, to induce and prouoke some louer of plants to obtaine it for his pleasure, and others also.

4. *Syringa flore albo simplici.* The single white Pipe tree.

The single white Pipe tree or bush, neuer commeth to that height of the former, but abideth alwaies like a hedge tree or bush, full of shootes or suckers from the roote, much more then the former: the young shootes hereof are reddish on the outside, and afterward reddish at the ioynts, and grayish all the rest ouer: the young as well as the old branches, haue some pith in the middle of them, like as the Elder hath: the leaues stand two at a ioynt, somewhat like the former, but more rugged or crumpled, as also a little pointed, and dented about the edges: the flowers growe at the toppes of the branches, diuers standing together, consisting of foure white leaues, like vnto small Muske Roses, and of the same creame colour, as I may call it, with many small yellowish threads in the middle, and are of a strong, full, or heady sent, not pleasing to a great many, by reason of the strange quickenesse of the sent: the fruit followeth, being flat at the head, with many leafie shels or scales compassing it, wherein is enclosed small long seede: the rootes runne not deepe, but spread vnder the ground, with many fibres annexed vnto them.

5. *Syringa Arabica flore albo duplici.*
The double white Pipe tree.

This Pipe tree hath diuers long and slender branches, whereon growe large leaues, somewhat like vnto the leaues of the former single white kinde, but not so rough or hard, and not at all dented about the edges, two alwaies standing one against another at euery ioynt of the stalke, but set or disposed on contrary sides, and not all vpon one side; at the ends whereof come forth diuers flowers, euery one standing on his owne foote-stalke, the hose or huske being long and hollow, like vnto the white Iasmine, and the flowers therin consisting of a double rowe of white and round pointed leaues, fiue or six in a rowe, with some yellownesse in the middle, which is hollow, of a very strong and heady sweet sent, and abiding a long time flowring, especially in the hotter Countries, but is very tender, and not able to abide any the least cold weather with vs;

for

1 *Iasminum album vulgare.* The ordinary white Iasmine. 2 *Iasminum Americanum siue Conuolulus Americanus.* The Iasmine or Bindweed of America. 3 *Iasminum luteum vulgare.* The yellow Iasmine. 4 *Lilac seu Syringa caerulea.* The blew Pipe tree. 5 *Syringa alba vulgaris.* The single white Syringa or Pipe tree. 6 *Syringa flore albo duplici.* The double white Syringa. 7 *Sambucus rosea.* The Elder or Gelder Rose.

for the cold windes will (as I vnderstand) greatly molest it : and therefore must as charily be kept as Orenge trees with vs, if wee will haue it to abide.

## The Place.

The first groweth in Arabia (as Matthiolus thinketh, that had it from Constantinople.) We haue it plentifully in our Gardens. The second and third are strangers with vs as yet. The fourth is as frequent as the first, or rather more, but his originall is not knowne. The last hath his originall from Arabia, as his name importeth.

## The Time.

The first, second, and third flower in Aprill, the other two not vntill May.

## The Names.

The first is called of Matthiolus *Lilac*, and by that name is most vsually called in all parts. It is also called *Syringa cærulea*, because it commeth neerest vnto those woods, which for their pithy substance, were made hollow into pipes. It is called of all in English, The blew Pipe tree. It seemeth likely, that Petrus Bellonius in his third Booke and fiftieth Chapter of his obseruations (making mention of a shrubbe that the Turkes haue, with Iuie leaues alwaies greene, bearing blew or violet coloured flowers on a long stalke, of the bignesse and fashion of a Foxe taile, and thereupon called in their language a Foxe taile) doth vnderstand this plant here expressed. The certainty whereof might easily be knowne, if any of our Merchants there residing, would but call for such a shrubbe, by the name of a Foxe taile in the Turkish tongue, and take care to send a young roote, in a small tubbe or basket with earth by Sea, vnto vs here at London, which would be performed with a very little paines and cost. The second and third, as kindes thereof, haue their names in their titles. The fourth is called by Clusius and others, *Frutex Coronarius*; some doe call it *Lilac flore albo*, but that name is not proper, in that it doth confound both kindes together. Lobel calleth it *Syringa Italica*. It is now generally called of all *Syringa alba*, that is in English, The white Pipe tree. Some would haue it to bee *Ostrys* of Theophrastus, but Clusius hath sufficiently cleared that doubt. Of others *Ligustrum Orientale*, which it cannot be neither; for the *Cyprus* of Plinie is Dioscorides his *Ligustrum*, which may be called *Orientale*, in that it is most proper to the Easterne Countries, and is very sweete, whose seede is like vnto Coriander seede. The last is called by diuers *Syringa Arabica flore albo duplici*, as most fitly agreeing thereunto. Of Basilius Besslerus that set forth the great booke of the Bishop of Eystot in Germany his Garden, *Syringa Italica flore albo pleno*, because, as it is likely, hee had it from Italy. It is very likely, that Prosper Alpinus in his booke of Egyptian plants, doth meane this plant, which hee there calleth *Sambach, siue Iasminum Arabicum*. Matthæus Caccini of Florence in his letter to Clusius entituleth it *Syringa Arabica, siue Iasminum Arabicum, siue Iasminum ex Gine*, whereby hee declareth that it may not vnfitly be referred to either of them both. We may call it in English as it is in the title, The double white Pipe tree.

## The Vertues.

We haue no vse of these in Physicke that I know, although Prosper Alpinus saith, the double white Pipe tree is much vsed in Egypt, to help women in their trauailes of childbirth.

## CHAP. CVIII.

### *Sambucus Rosea.* The Elder or Gelder Rose.

ALthough there be diuers kindes of Elders, yet there is but one kinde of Elder Rose, whereof I meane to intreate in this Chapter, being of neare affinity in some things vnto the former Pipe trees, and which for the beauty of it deserueth to be remembred among the delights of a Garden.

### *Sambucus Rosea.* The Gelder Rose.

The Gelder Rose (as it is called) groweth to a reasonable height, standing like a tree, with a trunke as bigge as any mans arme, couered with a darke grayish barke, somewhat rugged and very knotty: the younger branches are smooth and white, with a pithy substance in the middle, as the Elders haue, to shew that it is a kind thereof, whereon are set broad leaues, diuided into three parts or diuisions, somewhat like vnto a Vine leafe, but smaller, and more rugged or crumpled, iagged or cut also about the edges: at the toppes of euery one of the young branches, most vsually commeth forth a great tuft, or ball as it were, of many white flowers, set so close together, that there can be no distinction of any seuerall flower seene, nor doth it seeme like the double flower of any other plant, that hath many rowes of leaues set together, but is a cluster of white leaued flowers set together vpon the stalke that vpholdeth them, of a small sent, which fall away without bearing any fruit in our Country, that euer I could obserue or learne: The roote spreadeth neither farre nor deepe, but shooteth many small rootes and fibres, whereby it is fastened in the ground, and draweth nourishment to it, and sometimes yeeldeth suckers from it.

### The Place.

It should seeme, that the naturall place of this Elder is wet and moist grounds, because it is so like vnto the Marsh Elder, which is the single kind hereof. It is onely noursed vp in Gardens in all our Country.

### The Time.

It flowreth in May, much about the time of the double Peony flower, both which being set together, make a pleasant variety, to decke vp the windowes of a house.

### The Names.

It is generally called *Sambucus Rosea*: In English, The Elder Rose, and more commonly after the Dutch name, the Gelder Rose. Dalechampius seemeth to make it *Thraupalus* of Theophrastus, or rather the single Marsh Elder; for I thinke this double kinde was not knowne in Theophrastus his time.

### The Vertues.

It is not applyed to any Physicall vse that I know.

## CHAP. CIX.

### *Rosa.* The Rose tree or bush.

THe great varietie of Roses is much to be admired, beeing more then is to bee seene in any other shrubby plant that I know, both for colour, forme and smell. I haue to furnish this garden thirty sorts at the least, euery one notably differing from the other, and all fit to be here entertained: for there are some other, that being wilde and of no beautie or smell, we forbeare, and leaue to their wilde habitations. To distinguish them by their colours, as white, red, incarnate, and yellow, were a way that many might take, but I hold it not so conuenient for diuers respects: for so I should confound those of diuers sorts one among another, and I should not keepe that methode which to me seemeth most conuenient, which is, to place and ranke euery kinde, whether single or double, one next vnto the other, that so you may the better vnderstand their varieties and differences: I will therefore beginne with the most ancient, and knowne Roses to our Countrey, whether naturall or no I know not, but assumed by our precedent Kings of all others, to bee cognisances of their dignitie, the white Rose and the red, whom shall follow the damaske, of the finest sent, and most vse of all the other sorts, and the rest in their order.

### 1. *Rosa Anglica alba.* The English white Rose.

The white Rose is of two kindes, the one more thicke and double then the other: The one riseth vp in some shadowie places, vnto eight or ten foote high, with a stocke of a great bignesse for a Rose. The other growing seldome higher then a Damaske Rose. Some doe iudge both these to be but one kinde, the diuersitie happening by the ayre, or ground, or both. Both these Roses haue somewhat smaller and whiter greene leaues then in many other Roses, fiue most vsually set on a stalke, and more white vnderneath, as also a whiter greene barke, armed with sharpe thornes or prickles, whereby they are soone known from other Roses, although the one not so easily from the other: the flowers in the one are whitish, with an eye or shew of a blush, especially towards the ground or bottome of the flower, very thicke double, and close set together, and for the most part not opening it selfe so largely and fully as eyther the Red or Damaske Rose. The other more white, lesse thicke and double, and opening it selfe more, and some so little double as but of two or three rowes, that they might be held to be single, yet all of little or no smell at all. To describe you all the seuerall parts of the Rose, as the bud, the beards, the threds &c. were needlesse, they are so conuersant in euery ones hand, that I shall not neede but to touch the most speciall parts of the varieties of them, and leaue a more exact relation of all things incident vnto them, vnto a generall worke.

### 2. *Rosa Incarnata.* The Carnation Rose.

The Carnation Rose is in most things like vnto the lesser white rose, both for the growing of the stocke, and bignesse of the flower, but that it is more spreade abroade when it is blown then the white is, and is of a pale blush colour all the flower thorough-out, of as small a sent as the white one is almost.

*Rosa Belgica siue Vitrea.*

This kinde of Rose is not very great, but very thicke and double, and is very variable in the flowers, in that they will be so different one from another: some being paler then others, and some as it were blasted, which commeth not casually, but naturally to this rose: but the best flowers (whereof there will bee still some) will be of a bright pale murrey colour, neere vnto the Veluet rose, but nothing so darke a colour.

### 3. *Rosa Anglica rubra.* The English red Rose.

The red Rose (which I call English, not only for the reason before expressed, but be-

cause

cause (as I take it) this Rose is more frequent and vsed in England, then in other places) neuer groweth so high as the damaske Rose bush, but most vsually abideth low, and shooteth forth many branches from the roote (and is but seldome suffered to grow vp as the damaske Rose into standards) with a greene barke, thinner set with prickles, and larger and greener leaues on the vpperside then in the white, yet with an eye of white vpon them, fiue likewise most vsually set vpon a stalke, and grayish or whitish vnderneath. The Roses or Flowers doe very much vary, according to their site and abiding; for some are of an orient, red or deepe crimson colour, and very double (although neuer so double as the white) which when it is full blowne hath the largest leaues of any other Rose; some of them againe are paler, tending somewhat to a damaske; and some are of so pale a red, as that it is rather of the colour of the canker Rose, yet all for the most part with larger leaues then the damaske, and with many more yellow threds in the middle: the sent hereof is much better then in the white, but not comparable to the excellencie of the damaske Rose, yet this Rose being well dryed and well kept, will hold both colour and sent longer then the damaske, bee it neuer so well kept.

### 4. *Rosa Damascena.* The Damaske Rose.

The Damaske Rose bush is more vsually noursed vp to a competent height to stand alone, (which we call Standards) then any other Rose: the barke both of the stocke and branches, is not fully so greene as the red or white Rose: the leaues are greene with an eye of white vpon them, so like vnto the red Rose, that there is no great difference betweene them, but that the leaues of the red Rose seeme to bee of a darker greene. The flowers are of a fine deepe blush colour, as all know, with some pale yellow threds in the middle, and are not so thicke and double as the white, nor being blowne, with so large and great leaues as the red, but of the most excellent sweet pleasant sent, far surpassing all other Roses or Flowers, being neyther heady nor too strong, nor stuffing or vnpleasant sweet, as many other flowers.

### 5. *Rosa Prouincialis sive Hollandica Damascena.* The great double Damaske Prouince or Holland Rose.

This Rose (that some call *Centifolia Batauica incarnata*) hath his barke of a reddish or browne colour, whereby it is soone discerned from other Roses. The leaues are likewise more reddish then in others, and somewhat larger, it vsually groweth very like the Damaske rose, and much to the same height: the flowers or roses are of the same deepe blush colour that the damaske roses are, or rather somewhat deeper, but much thicker, broader, and more double, or fuller of leaues by three parts almost, the outer leaues turning themselues backe, when the flower hath stood long blowne, the middle part it selfe (which in all other roses almost haue some yellow threds in them to be seene) being folded hard with small leaues, without any yellow almost at all to be seene, the sent whereof commeth neerest vnto the damaske rose, but yet is short of it by much, howsoeuer many doe thinke it as good as the damask, and to that end I haue known some Gentlewomen haue caused all their damaske stockes to bee grafted with prouince Roses, hoping to haue as good water, and more store of them then of damask Roses; but in my opinion it is not of halfe so good a sent as the water of damaske Roses: let euery one follow their own fancie.

### 6. *Rosa Prouincialis rubra.* The red Prouince Rose.

As the former was called *incarnata*, so this is called *Batauica centifolia rubra*, the difference being not very great: the stemme or stocke, and the branches also in this, seeming not to be so great but greener, the barke being not so red; the leaues of the same largenesse with the former damaske Prouince. The flowers are not altogether so large, thicke and double, and of a little deeper damaske or blush colour, turning to a red Rose, but not comming neere the full colour of the best red Rose, of a sent not so sweete as the damaske Prouince, but comming somewhat neere the sent of the ordinary

dinary red rose, yet exceeding it. This rose is not so plentifull in bearing as the damaske Prouince.

7. *Rosa Prouincialis alba.* The white Prouince Rose.

It is said of diuers, that there is a white Prouince Rose, whereof I am not *oculatus testis*, and therfore I dare not giue it you for a certaintie, and indeed I haue some doubt, that it is the greater and more double white rose, whereof I gaue you the knowledge in the beginning: when I am my selfe better satisfied, I shall bee ready to satisfie others.

8. *Rosa versicolor.* The party coloured Rose, of some Yorke and Lancaster.

This Rose in the forme and order of the growing, is neerest vnto the ordinary damaske rose, both for stemme, branch, leafe and flower: the difference consisting in this, that the flower (being of the same largenesse and doublenesse as the damask rose) hath the one halfe of it, sometimes of a pale whitish colour, and the other halfe, of a paler damaske colour then the ordinary; this happeneth so many times, and sometimes also the flower hath diuers stripes, and markes in it, as one leafe white, or striped with white, and the other halfe blush, or striped with blush, sometimes also all striped, or spotted ouer, and other times little or no stripes or markes at all, as nature listeth to play with varieties, in this as in other flowers: yet this I haue obserued, that the longer it abideth blowen open in the sun, the paler and the fewer stripes, markes or spots will be seene in it: the smell whereof is of a weake damaske rose sent.

9. *Rosa Chrystallina.* The Chrystall Rose.

This Rose is very like vnto the last described, both for stocke, branch and leafe: the flower hereof is not much different from it, being no great large or double Rose, but of a meane size, striped and marked with a deeper blush or red, vpon the pale coloured leafe, that it seemeth in the marking and beauty thereof, to bee of as much delight as the Chrystall Gilloflower: this, euen like the former, soone fadeth and passeth away, not yeelding any great store of flowers any yeare.

10. *Rosa rubra humilis siue pumilio.*
The dwarfe red Rose, or Gilloflower Rose.

This Rose groweth alwayes low and small, otherwise in most respects like vnto the ordinary redde Rose, and with few or no thornes vpon it: the Flowers or Roses are double, thicke, small and close, not so much spread open as the ordinary red, but somewhat like vnto the first double white Rose before expressed; yet in some places I haue seene them more layde open then these, as they grew in my garden, being so euen at the toppes of the leaues, as if they had been clipt off with a paire of sheeres, and are not fully of so red a colour as the red Prouince Rose, and of as small or weak sent as the ordinary red Rose, or not so much.

11. *Rosa Francofurtensis.* The Franckford Rose.

The young shootes of this Rose are couered with a pale purplish barke, set with a number of small prickes like haires, and the elder haue but very few thornes: the flower or rose it selfe hath a very great bud or button vnder it, more then in any other rose, and is thicke and double as a red rose, but so strongly swelling in the bud, that many of them breake before they can be full blowen, and then they are of a pale red rose colour, that is, betweene a red and a damaske, with a very thicke broade and hard vmbone of short yellow threds or thrumes in the middle, the huske of the flower hauing long ends, which are called the beards of the rose, which in all other are iagged in some of them, in this hath no iagge at all: the smell is neerest vnto a red Rose.

12. *Rosa*

1 *Roſa Damaſcena.* The Damaske Roſe. 2 *Roſa Prouincialis ſiue Hollandica.* The great Prouince Roſe. 3 *Roſa Francofurtenſis.* The Franckford Roſe. 4 *Roſa rubra humilis.* The dwarfered Roſe. 5 *Roſa Hungarica.* The Hungarian Roſe. 6 *Roſa lutea multiplex.* The great double yellow Roſe.

12. *Roſa Hungarica.* The Hungarian Roſe.

The Hungarian Roſe hath greene ſhootes ſlenderly ſet with prickes, and ſeldome groweth higher then ordinarily the red Roſe doth; the ſtemme or ſtocke being much about that bigneſſe: the flower or roſe is as great, thicke and double, as the ordinary red Roſe, and of the ſame faſhion, of a paler red colour, and beeing neerely looked vpon is finely ſpotted with faint ſpots, as it were ſpreade ouer the red; the ſmell wherof is ſomewhat better then the ſmell of the ordinary red Roſe of the beſt kinde.

13. *Roſa Holoſerica ſimplex & multiplex.*
The Veluet Roſe ſingle and double.

The old ſtemme or ſtock of the veluet Roſe is couered with a dark coloured barke, and the young ſhootes of a ſad greene with very few or no thornes at all vpon them: the leaues are of a ſadder greene colour then in moſt ſorts of Roſes, and very often ſeuen on a ſtalke, many of the reſt hauing but fiue: the Roſe is eyther ſingle or double: the ſingle is a broade ſpread flower, conſiſting of fiue or ſixe broade leaues with many yellow threds in the middle: the double hath two rowes of leaues, the one large, which are outermoſt, the other ſmaller within, of a very deepe red crimſon colour like vnto crimſon veluet, with many yellow threds alſo in the middle; and yet for all the double rowe of leaues, theſe Roſes ſtand but like ſingle flowers: but there is another double kinde that is more double then this laſt, conſiſting oftentimes of ſixteene leaues or more in a flower, and moſt of them of an equall bigneſſe, of the colour of the firſt ſingle roſe of this kinde, or ſomewhat freſher; but all of them of a ſmaller ſent then the ordinary red Roſe.

14. *Roſa ſine ſpinis ſimplex & multiplex.*
The Roſe without thornes ſingle and double.

The Roſe without thornes hath diuers greene ſmooth ſhootes, riſing from the root, without any pricke or thorne at all vpon them, eyther young or old: the leaues are not fully ſo large as of the red roſe: the flowers or roſes are not much bigger then thoſe of the double Cinamon Roſe, thicke ſet together and ſhort, of a pale red Roſe colour, with diuers pale coloured veines through euery leafe of the flower, which hath cauſed ſome to call it The marbled Roſe, and is of a ſmall ſent, not fully equall to the red Roſe. The ſingle of this kinde differeth not in any other thing from the former, then in the doubleneſſe or ſingleneſſe of the flowers, which in this are not halfe ſo double, nor yet fully ſingle, and are of a paler red colour.

*Roſa ſine ſpina flore albo.* I haue heard likewiſe of a white Roſe of this kinde, but I haue ſeene none ſuch as yet, and therefore I can ſay no more thereof.

15. *Roſa Cinamomea ſimplex & multiplex.*
The Cinamon Roſe ſingle and double.

The ſingle Cinamon Roſe hath his ſhootes ſomewhat red, yet not ſo red as the double kinde, armed with great thornes, like almoſt vnto the Eglantine buſh, thereby ſhowing, as well by the multiplicitie of his ſhootes, as the quickneſſe and height of his ſhooting, his wilde nature: On the ſtemme and branches ſtand winged leaues, ſometimes ſeuen or more together, which are ſmall and greene, yet like vnto other Roſes. The Roſes are ſingle, of fiue leaues a peece, ſomewhat large, and of a pale red colour, like vnto the double kinde, which is in ſhootes redder, and in all other things like vnto the ſingle, but bearing ſmall, ſhort, thicke and double Roſes, ſomewhat like vnto the Roſe without thornes, but a little leſſer, of a paler red colour at the end of the leaues, and ſomewhat redder and brighter toward the middle of them, with many yellow ſhort thrumes; the ſmall ſent of Cinamon that is found in the flowers hath cauſed it to beare the name.

16. *Roſa*

16. *Rosa lutea simplex.* The single yellow Rose.

This single yellow Rose is planted rather for variety then any other good vse. It often groweth to a good height, his stemme being great and wooddy, with few or no prickes vpon the old wood, but with a numher of small prickes like haires, thicke set, vpon the younger branches, of a darke colour somewhat reddish, the barke of the young shootes being of a sad greene reddish colour: the leaues of this Rose bush are smaller, rounder pointed, of a paler greene colour, yet finely snipt about the edges, and more in number, that is, seuen or nine on a stalke or ribbe, then in any other Garden kinde, except the double of the same kinde that followeth next: the flower is a small single Rose, consisting of fiue leaues, not so large as the single Spanish Muske Rose, but somewhat bigger then the Eglantine or sweete Briar Rose, of a fine pale yellow colour, without any great sent at all while it is fresh, but a little more, yet small and weake when it is dryed.

17. *Rosa lutea multiplex siue flore pleno.*
The double yellow Rose.

The double yellow Rose is of great account, both for the rarity, and doubleness of the flower, and had it sent to the rest, would of all other be of highest esteeme. The stemme or stocke, the young shoots or branches, the small hairy prickes, and the small winged leaues, are in all parts like vnto the former single kinde; the chiefest difference consisteth in the doubleness of the flower or Rose, which is so thicke and double, that very often it breaketh out on one side or another, and but a few of them abiding whole and faire in our Countrey, the cause whereof wee doe imagine to bee the much moisture of our Countrey, and the time of flowring being subiect to much raine and showers; many therefore doe either plant it against a wall, or other wayes defend it by couering: againe, it is so plentifull in young shootes or branches, as also in flowers at the toppe of euery branch, which are small and weake for the most part, that they are not able to bring all the flowers to ripeneſſe; and therefore most of them fall or wither away without comming to perfection (the remedy that many doe vse for this inconuenience last recited is, that they nippe away most of the buds, leauing but some few vpon it, that so the vigour of the plant may be collected into a few flowers, whereby they may the better come to perfection, and yet euen thus it is hardly effected) which are of a yellowish greene colour in the bud, and before they be blowne open, but then are of a faire yellow colour, very full of leaues, with many short haires rather then leaues in the middle, and hauing short, round, greene, smooth buttons, almost flat vnder them: the flower being faire blowne open, doth scarce giue place for largeness, thickness, and doubleness, vnto the great Prouence or Holland Rose. This Rose bush or plant is very tender with vs here about London, and will require some more care and keeping then the single of this kinde, which is hardy enough; for I haue lost many my selfe, and I know but a few about this towne that can nourse it vp kindly, to beare or scarce to abide without perishing; but abideth well in euery free aire of all or the most parts of this Kingdome: but (as I heare) not so well in the North.

18. *Rosa Moschata simplex & multiplex.*
The Muske Rose single and double.

The Muske Rose both single and double, rise vp oftentimes to a very great height, that it ouergroweth any arbour in a Garden, or being set by an house side, to bee ten or twelue foote high, or more, but more especially the single kinde, with many green farre spread branches, armed with a few sharpe great thornes, as the wilder sorts of Roses are, whereof these are accounted to be kindes, hauing small darke greene leaues on them, not much bigger then the leaues of Eglantine: the flowers come forth at the toppes of the branches, many together as it were in an vmbell or tuft, which for the most part doe flower all at a time, or not long one after another, euery one standing on a pretty long stalke, and are of a pale whitish or creame colour, both the single and

the

the double; the single being small flowers, consisting of fiue leaues, with many yellow threads in the middle: and the double bearing more double flowers, as if they were once or twice more double then the single, with yellow thrummes also in the middle, both of them of a very sweete and pleasing smell, resembling Muske: some there be that haue auouched, that the chiefest sent of these Roses consisteth not in the leaues, but in the threads of the flowers.

19. *Rosa Moschata multiplex altera: alijs Damascena alba, vel verisimilior Cinamomea flore pleno albo.*
The double white Damaske Muske Rose.

This other kinde of Muske Rose (which with some is called the white Damaske Muske, but more truely the double white Cinamon Rose) hath his stemme and branches also shorter then the former, but as greene: the leaues are somewhat larger, and of a whiter greene colour; the flowers also are somewhat larger then the former double kinde, but standing in vmbels after the same manner, or somewhat thicker, and of the same whitish colour, or a little whiter, and somewhat, although but a little, neare the smell of the other, but nothing so strong. This flowreth at the time of other Roses, or somewhat later, yet much before the former two sorts of Muske Roses, which flower not vntill the end of Summer, and in Autumne; both which things, that is, the time of the flowring, and the sent being both different, shew plainly it cannot be of the tribe of Muske Roses.

20. *Rosa Hispanica Moschata simplex.* The Spanish Muske Rose.

This Spanish Rose riseth to the height of the Eglantine, and sometimes higher, with diuers great greene braneches, the leaues whereof are larger and greener then of the former kindes: the flowers are single Roses, consisting of fiue whiter leaues then in any of the former Muske Roses, and much larger, hauing sometimes an eye of a blush in the white, of a very sweete smell, comming neareſt vnto the last recited Muske Rose, as also for the time of the flowring.

21. *Rosa Pomifera maior.* The great Apple Rose.

The stemme or stocke of this Rose is great, couered with a darke grayish barke, but the younger branches are somewhat reddish, armed here and there with great and sharpe thornes, but nothing so great or plentifull as in the Eglantine, although it be a wilde kinde: the leaues are of a whitish greene colour, almost like vnto the first white Rose, and fiue alwaies set together, but seldome seuen: the flowers are small and single, consisting of fiue leaues, without any sent, or very little, and little bigger then those of the Eglantine bush, and of the very same deepe blush colour, euery one standing vpon a rough or prickly button, bearded in the manner of other Roses, which when the flowers are fallen growe great, somewhat long and round, peare-fashion, bearing the beards on the tops of them; and being full ripe are very red, keeping the small prickles still on them, wherein are many white, hard, and roundish seedes, very like vnto the seede of the Heppes or Eglantine berries, lying in a soft pulpe, like vnto the Hawthorne berries or Hawes: the whole beauty of this plant consisteth more in the gracefull aspect of the red apples or fruit hanging vpon the bushes, then in the flowers, or any other thing. It seemeth to be the same that Clusius calleth *Rosa Pumila*, but that with me it groweth much higher and greater then he saith his doth.

22. *Rosa siluestris odora siue Eglenteria simplex.*
The single Eglantine or sweete Briar bush.

The sweete Briar or Eglantine Rose is so well knowne, being not onely planted in Gardens, for the sweetenesse of the leaues, but growing wilde in many woods and hedges, that I thinke it lost time to describe it; for that all know it hath exceeding long greene shootes, armed with the cruellest sharpe and strong thornes, and thicker set

then

1 *Rosa sine spinis multiplex.* The double Rose without thorns. 2 *Rosa Cinamomea flore pleno.* The double Cinamon Rose. 3 *Rosa Holoserica simplex.* The single Veluet Rose. 4 *Rosa Holoserica duplex.* The double Veluet Rose. 5 *Rosa Moschata multiplex.* The double Muske Rose. 6 *Rosa Moschata Hispanica simplex.* The single Spanish Muske Rose. 7 *Rosa Pomifera maior.* The great Apple Rose. 8 *Rosa siluestris sive Eglanteria duplex.* The double Eglantine Rose.

then is in any Rose either wilde or tame: the leaues are smaller then in most of those that are noursed vp in Gardens, seuen or nine most vsually set together on a ribbe or stalke, very greene and sweete in smell, aboue the leaues of any other kinde of Rose: the flowers are small single blush Roses, of little or no sent at all, which turne into reddish berries, stuffed within with a dounie or flocky matter or substance, wherein doth lye white hard seede.

23. *Rosa silnestris odora sine Eglenteria flore duplici.*
The double Eglantine.

The double Eglantine is in all the places that I haue seene it a grafted Rose, (but I doubt not, but that his originall was naturall, and that it may be made naturall againe, as diuers other Roses are.) It groweth and spreadeth very well, and with a great head of branches, whereon stand such like leaues as are in the single kinde, but a little larger, not smelling fully so sweete as it: the flowers are somewhat bigger then the single, but not much, hauing but one other rowe of leaues onely more then the former, which are smaller, and the outer leaues larger, but of the same pale reddish purple colour, and smelleth somewhat better then the single.

24. *Rosa semper virens.* The euer greene Rose bush.

This Rose or bush is very like vnto a wilde single Eglantine bush in many respects, hauing many very long greene branches, but more slender and weake, so that many times they bend downe againe, not able to sustaine themselues without some helpe, and armed with hooked thornes as other Roses be; the winged leaues consist of seuen for the most part, whereof those two that are lowest and opposite, are smallest, the next two bigger then they, the third couple bigger then any of the rest belowe, and the end leafe biggest of all: this proportion generally it holdeth in euery winged leafe through the whole plant, which at the first comming forth are somewhat reddish, with the young branch that shooteth out with them, but being full growne, are of a deepe greene colour, and somewhat shining, dented about the edges, and fall not away from the branches as other Roses doe, but abide thereon for the most part all the Winter: the flowers stand foure or fiue together at the tops of the branches, being single Roses, made of fiue leaues a peece, of a pure white colour, much larger then the ordinary Muske Rose, and of a fine sent, comming nearest thereunto, with many yellow chiues or threads in the middle.

25 Eglenteria Austriaca. vid Theat. Botanic.

The Place.

Some of these Roses had their originall, as is thought in England, as the first and second; for these dryed red Roses that come ouer to vs from beyond the Seas, are not of the kinde of our red Rose, as may well be perceiued by them that will compare our English dryed leaues with those. Some in Germany, Spaine, and Italy. Some againe in Turkie, as the double yellow Rose, which first was procured to be brought into England, by Master Nicholas Lete, a worthy Merchant of London, and a great louer of flowers, from Constantinople, which (as wee heare) was first brought thither from Syria; but perished quickly both with him, and with all other to whom hee imparted it: yet afterwards it was sent to Master Iohn de Franqueuille, a Merchant also of London, and a great louer of all rare plants, as well as flowers, from which is sprung the greatest store, that is now flourishing in this Kingdome.

The Time.

The Cinamon Rose is the earliest for the most part, which flowreth with vs about the middle of May, and sometimes in the beginning. The ordinary Muske Roses both single and double flower lateste, as is said. All the other flower much about one time, in the beginning of Iune, or thereabouts, and continue flowring all that moneth, and the next throughout for the most part, and the red vntill August be halfe past.

The

### The Names.

The seuerall names, whereby they are most commonly knowne vnto vs in this Countrey, are expressed in their titles; but they are much differing from what they are called in other Countries neare vnto vs, which to compare, conferre, and agree together, were a worke of more paines then vse: But to proportion them vnto the names set downe by Theophrastus, Pliny, and the rest of the ancient Authors, were a worke, wherein I might be sure not to escape without falling into errour, as I verily beleeue many others haue done, that haue vndertaken to doe it: I will therefore for this worke desire that you will rest contented, with so much as hath already been deliuered, and expect an exact definition and complete satisfaction by such a methodicall course as a generall History will require, to be performed by them that shall publish it.

### The Vertues.

The Rose is of exceeding great vse with vs; for the Damaske Rose (besides the superexcellent sweete water it yeeldeth being distilled, or the perfume of the leaues being dryed, seruing to fill sweete bags) serueth to cause solublenesse of the body, made into a Syrupe, or preserued with Sugar moist or dry candid. The Damaske Prouince Rose, is not onely for sent neerest of all other Roses vnto the Damaske, but in the operation of solubility also. The red Rose hath many Physicall vses much more then any other, seruing for many sorts of compositions, both cordiall and cooling, both binding and loosing. The white Rose is much vsed for the cooling of heate in the eyes: diuers doe make an excellent yellow colour of the iuyce of white Roses, wherein some Allome is dissolued, to paint or colour flowers or pictures, or any other such things. There is little vse of any other sort of Roses; yet some affirme, that the Muske Roses are as strong in operation to open or loosen the belly as the Damaske Rose or Prouince.

## Chap. CXI.

### *Cistus.* The Holly Rose or Sage Rose.

THere are three principall kindes of *Cistus*, the male, the female, and the gumme or sweete smelling *Cistus* bearing *Ladanum*, called *Ledon*. Of each of these three there are also diuers sorts: Of them all to intreate in this worke is not my minde, I will onely select out of the multitude some few that are fit for this our Garden, and leaue the rest to a greater.

### 1. *Cistus mas.* The male Holly Rose or Sage Rose.

The male *Cistus* that is most familiar vnto our Countrey, I meane that will best abide, is a small shrubby plant, growing seldome aboue three or foure foote high with vs, hauing many slender brittle wooddy branches, couered with a whitish barke, whereon are set many whitish greene leaues, long and somewhat narrow, crumpled or wrinckled as it were with veines, and somewhat hard in handling, especially the old ones; for the young ones are softer, somewhat like vnto Sage leaues for the forme and colour, but much smaller, two alwaies set together at a ioynt: the flowers stand at the toppe of the branches, three or foure together vpon seuerall slender foot-stalkes, consisting of fiue small round leaues a peece, somewhat like vnto a small single Rose, of a fine reddish purple colour, with many yellow threads in the middle, with-

out any sent at all, and quickly fading or falling away, abiding seldome one whole day blowne at the most: after the flowers are past, there come vp round hard hairie heads in their places, containing small browniſh seede: the roote is wooddy, and will abide some yeares with vs, if there be some care had to keepe it from the extreamity of our Winters frostes, which both this, and many of the other sorts and kinds, will not abide doe what we can.

### 2. *Cistus fœmina.* The female Holly Rose.

The female Holly Rose groweth lower, and smaller then the former male kinde, hauing blackish branches, lesse woody, but not lesse brittle then it: the leaues are somewhat rounder and greener, but a little hard or rough withall, growing in the same manner vpon the branches by couples: the flowers grow at the toppes of the branches, like vnto the former, consisting of fiue leaues, but somewhat lesser, and wholly white, with yellow threds in the middle, as quickly fading, and of as little sent as the former: the heads and seede are somewhat bigger then in the former.

### 3. *Chamæcistus Friſicus.* The dwarfe Holly Rose of Friseland.

This dwarfe Cistus is a small low plant, hauing diuers shootes from the rootes, full of leaues that are long and narrow, very like vnto the leaues of the French Spikenard or *Spica Celtica*; from among which leaues shoote forth short stalkes, not aboue a span high, with a few smaller leaues thereon; and at the toppes diuers small flowers one aboue another, consisting of six small round leaues, of a yellow colour, hauing two circles of reddish spots round about the bottome of the leaues, a little distant one from another, which adde much grace to the flower: after the flowers are past, there come in their places small round heads, being two forked at the end, containing within them small browniſh chaffie seede: the roote is small and slender, with many fibres thereat creeping vnder ground, and shooting forth in diuers places, whereby it much encreaseth: the whole plant, and euery part of it, smelleth strong without any pleasant sent.

### 4. *Cistus annuus.* The Holly Rose of a yeare.

This small Cistus that endureth but a year (and will require to be sowne euery year, if ye will haue it) riseth vp with straight, but slender hard stalkes, set here and there confusedly with long and narrow greenish leaues, very like vnto the leaues of the Gum Cistus or Ledon, being a little clammy withall: at the toppe of the stalkes, and at the ioynts with the leaues, stand two or three pale yellow flowers, consisting of fiue leaues a peece, with a reddish spot neere the bottome of euery leafe of the flower, as quickely fading as any of the former: after which follow small three square heades, containing small seede, like vnto the first female kinde, but somewhat paler or yellower: the root is small and woody, and perisheth as soone as it hath borne seede.

### 5. *Cistus Ledon.* The Gum Cistus, or Sweete Holly Rose.

This sweete Holly Rose or Gum Cistus, riseth higher, and spreadeth larger then the former male kind doth, with many blackish woody branches, whereon are set diuers long and narrow darke greene leaues, but whitish vnderneath, two alwayes standing together at a ioint, both stalks and leaues bedeawed as it were continually with a clammy sweete moisture (which in the hot Countries is both more plentifull, and more sweet then in ours) almost transparent, and which being gathered by the inhabitants, with certaine instruments for that purpose (which in some places are leather thongs, drawne ouer the bushes, and after scraped off from the thongs againe, and put together) is that kind of blacke sweet gum, which is called *Ladanum* in the Apothecaries shops: at the tops of the branches stand single white flowers, like vnto single Roses, being larger then in any of the former kindes, consisting of fiue leaues, whereof euery one hath at the bottome a dark purplish spot, broad below, and small pointed vpwards, with some yellow threds in the middle: after which are past, there arise cornered

heads,

1 *Cistus mas* The male Holly Rose. 2 *Chamæcistus Frisucus.* The dwarfe Holly Rose of Frisia. 3 *Cistus Ledon.* The sweet Holly Rose or gumme Cistus. 4 *Ledum Alpinum.* The mountaine Holly Rose. 5 *Ledum Silesiacum.* The sweet Mary Rose of Silesia. 6 *Rosmarinum aureum,* Gilded Rosemary.

heads, containing such small brownish seede as is in the former male kinde: the roote is woody, and spreadeth vnder ground, abiding some yeares, if it be placed vnder a wall, where it may bee defended from the windes that often breake it, and from the extremitie of our winters, and especially the snow, if it lye vpon it, which quickly causeth it to perish.

### 6. *Ledum Alpinum seu Rosa Alpina.* The Mountaine sweet Holly Rose.

The fragrant smell with properties correspondent of two other plants, causeth me to insert them in this Chapter, and to bring them to your knowledge, as well worthy a fit place in our Garden. The first of them hath diuers slender woody branches, two foote high or thereabouts, couered with a grayish coloured barke, and many times leaning downe to the ground, whereby it taketh roote againe: vpon these branches grow many thicke, short, hard greene leaues, thicke set together, confusedly without order, sometimes whitish vnderneath, and sometimes yellowish: the toppes of the branches are loden with many flowers, which cause them to bend downwards, being long, hollow and reddish, opening into fiue corners, spotted on the outside with many white spots, and of a paler red colour on the inside, of a fine sweet sent: after the flowers are past, there follow small heads, containing small browniſh seede: the root is long, hard and woody, abiding better if it comprehend in the ground, then some of the former, because his originall is out of a colder country.

### 7. *Ledum Silesiacum.* The sweete Mary Rose, or Rosemary of Silesia.

This other sweete plante riseth vp with woody ash-coloured branches two foote high or more, which shoote forth other branches, of a reddish or purplish colour, couered with a browniſh yellow hoarinesse, on which are set many narrow long greene leaues, like vnto Rosemary leaues, but couered with the like hoarinesse as the stalks are (especially in the naturall places, but not so much being transplanted) and folding the sides of the leaues so close together, that they seeme nothing but ribbes, or stalkes, of an excellent sweet and pleasant sent; at the ends of the branches there grow certaine browniſh scaly heads, made of many small leaues set thicke together, out of which breake forth many flowers, standing in a tuft together, yet seuerally euery one vpon his owne footstalke, consisting of fiue white leaues, with certaine white threds in the middle, smelling very sweete: after which rise small greene heads, spotted with browniſh spots, wherein is contained very small, long, yellowish seede: the roote is hard and woodie.

#### The Place.

The first, second, fourth and fifth, grow in the hot Countries, as Italie, Spaine, &c. The third, and the two last in the colder Countries, as Friseland, Germanie, Bohemia.

#### The Time.

They do all flower in the Summer moneths of Iune, Iuly and August, and their seede is ripe quickly after.

#### The Names.

The first, second, fourth and fift, haue their names sufficiently expressed in their desctiptions. The third was sent vnto Clusius, vnder the name of *Herculus Friscus*, because of the strong sent: but he referreth it to the kinds of *Chamæcistus*, that is, dwarfe or low *Cistus*, both for the low growth, and for the flowers and seede sake. The fixt is diuersly called; for Clusius calleth it *Ledum Alpinum*: others, *Nerium Alpinum*, making it to bee a Rose Bay.

Gesner

Gesner according to the Countrey peoples name, *Rosa Alpina*, and *Rosa Montana*. Lobel calleth it *Balsamum Alpinum*, of the fragrant smell it hath, and *Chamærhododendros Chamalææfolio*. And some haue called it *Euonymus*, without all manner of iudgement. In English wee may call it, The Mountaine Rose, vntill a fitter name be giuen it. The last is called of Matthiolus, *Rosmarinum siluestre*, but of Clusius *Ledum*, referring it to their kindred; and *Silesiacum*, because he found it in that Countrey; or for distinction sake, as he saith, it may bee called, *Ledum foliis Rosmarini*, or *Ledum Bohemicum*. Cordus, as it seemeth in his History of Plants, calleth it *Chamapeuce*, as though he did account it a kinde of low Pine, or Pitch tree.

### The Vertues.

The first, second, and fift, are very astringent, effectuall for all sorts of fluxes of humours. The sweet Gum called *Ladanum*, made artificially into oyle, is of singular vse for *Alopecia*, or falling of the haire. The seed of the fourth is much commended against the stone of the Kidneyes. The sweete Rosemary of Silesia is vsed of the inhabitants, where it naturally groweth, against the shrinking of sinewes, crampes, or other such like diseases, wherof their daily experience makes it familiar, being vsed in bathing or otherwise.

## Chap. CXII.

### *Rosmarinum.* Rosemary.

THere hath beene vsually knowne but one sort of Rosemary, which is frequent through all this Country: but there are some other sorts not so well known; the one is called Gilded Rosemary; the other broade leafed Rosemary; a third I will adioyne, as more rare then all the other, called Double flowred Rosmary, because few haue heard thereof, much lesse seene it, and my selfe am not well acquainted with it, but am bold to deliuer it vpon credit.

### 1. *Libanotis Coronaria sive Rosmarinum vulgare.* Our Common Rosmary.

This common Rosemary is so well knowne through all our Land, being in euery womans garden, that it were sufficient but to name it as an ornament among other sweete herbes and flowers in our Garden, seeing euery one can describe it: but that I may say something of it, It is well obserued, as well in this our Land (where it hath been planted in Noblemens, and great mens gardens against bricke wals, and there continued long) as beyond the Seas, in the naturall places where it groweth, that it riseth vp in time vnto a very great height, with a great and woody stemme (of that compasse, that (being clouen out into thin boards) it hath serued to make lutes, or such like instruments, and here with vs Carpenters rules, and to diuers other purposes) branching out into diuers and sundry armes that extend a great way, and from them againe into many other smaller branches, wheron are set at seueral distances, at the ioynts, many very narrow long leaues, greene aboue, and whitish vnderneath; among which come forth towards the toppes of the stalkes, diuers sweet gaping flowers, of a pale or bleake blewish colour, many set together, standing in whitish huskes; the seed is small and red, but thereof seldome doth any plants arise that will abide without extraordinary care; for although it will spring of the seede reasonable well, yet it is so small and tender the first yeare, that a sharpe winter killeth it quickly, vnlesse it be very well defended: the whole plant as well leaues as flowers, smelleth exceeding sweete.

### 2. *Rosmarinum striatum, sive aureum.* Gilded Rosemary.

This Rosemary differeth not from the former, in forme or manner of growing, nor

in the forme or colour of the flower, but only in the leaues, which are edged, or stri-ped, or pointed with a faire gold yellow colour, which so continueth all the yeare throughout, yet fresher and fairer in Summer then in Winter; for then it will looke of a deader colour, yet so, that it may be discerned to be of two colours, green & yellow.

3. *Rosmarinum latifolium.* Broade leafed Rosemary.

This broad leafed Rosemary groweth in the same manner that the former doth, but that we haue not seene it in our Countrey since we had it to grow so great, or with such woody stemmes: the leaues stand together vpon the long branches after the same fa-shion, but larger, broader and greener then the other, and little or nothing whitish vnderneath: the flowers likewise are of the same forme and colour with the ordinary, but larger, and herein consisteth the difference.

4. *Rosmarinum flore duplici.* Double flowred Rosmary.

The double flowred Rosmary thus far differeth from the former, that it hath stron-ger stalkes, not so easie to breake, fairer, bigger and larger leaues, of a faire greene co-lour, and the flowers are double, as the Larkes heele or spurre: This I haue onely by relation, which I pray you accept, vntill I may by sight better enforme you.

## The Place.

Our ordinary Rosmary groweth in Spaine, and Prouence of France, and in others of those hot Countryes, neere the Sea side. It will not abide (vnlesse kept in stoues) in many places of Germany, Denmarke, and those colder Countries. And in some extreame hard winters, it hath well neere perished here in England with vs, at the least in many places: but by slipping it is vsually, and yearly encreased, to repleniish any garden.

## The Time.

It flowreth oftentimes twice in the yeare; in the Spring first, from April vntill the end of May or Iune, and in August and September after, if the yeare before haue been temperate.

## The Names.

Rosmary is called of the ancient Writers, *Libanotis*, but with this distinction, *Stephanomatica*, that is, *Coronaria*, because there were other plants called *Libanotis*, that were for other vses, as this for garlands, where flowers and sweete herbes were put together. The Latines call it *Rosmarinum*. Some would make it to be *Cneorum nigrum* of Theophrastus, as they would make Lauander to bee his *Cneorum album*, but Matthiolus hath sufficiently confuted that errour.

## The Vertues.

Rosmary is almost of as great vse as Bayes, or any other herbe both for inward and outward remedies, and as well for ciuill as physicall purposes. Inwardly for the head and heart; outwardly for the sinewes and ioynts: for ciuill vses, as all doe know, at weddings, funerals, &c. to bestow among friends: and the physicall are so many, that you might bee as well tyred in the reading, as I in the writing, if I should set down all that might be said of it. I will therefore onely giue you a taste of some, desiring you will be con-tent therewith. There is an excellent oyle drawne from the flowers alone by the heate of the Sunne, auaileable for many diseases both inward and outward, and accounted a soueraigne Balsame: it is also good to helpe dim-

nesse

nesse of sight, and to take away spots, markes and scarres from the skin; and is made in this manner. Take a quantitie of the flowers of Rosemary, according to your owne will eyther more or lesse, put them into a strong glasse close stopped, set them in hot horse dung to digest for fourteene dayes, which then being taken forth of the dung, and vnstopped, tye a fine linnen cloth ouer the mouth, and turne downe the mouth thereof into the mouth of another strong glasse, which being set in the hot Sun, an oyle will distill downe into the lower glasse; which preserue as precious for the vses before recited, and many more, as experience by practice may enforme diuers.

There is another oyle Chymically drawne, auaileable in the like manner for many the same inward and outward diseases, *viz.* for the heart, rheumaticke braines, and to strengthen the memory, outwardly to warme and comfort cold benummed sinewes, whereof many of good iudgement haue had much experience.

## Chap. CXIII.

### *Myrtus.* The Mirtle tree or bush.

IN the hot Countreyes, there haue been many sorts of Mirtles found out, naturally growing there, which will not fructifie in this of ours, nor yet abide without extraordinary care, and conueniencie withall, to preserue them from the sharpenesse of our winters. I shall only bring you to view three sorts in this my Garden, the one with a greater, the other two with lesser leaues, as the remainder of others which wee haue had, and which are preserued from time to time, not without much paine and trouble.

### 1. *Myrtus latifolia.* The greater leafed Mirtle.

The broader leafed Mirtle riseth vp to the height of foure or fiue foote at the most with vs, full of branches and leaues growing like a small bush, the stemme and elder branches whereof are couered with a dark coloured bark, but the young with a green, and some with a red, especially vpon the first shooting forth, whereon are set many fresh greene leaues, very sweet in smell, and very pleasant to behold, so neer resembling the leaues of the Pomegranate tree that groweth with vs, that they soone deceiue many that are not expert therein, being somewhat broade and long, and pointed at the ends, abiding alwaies green: at the ioynts of the branches where the leaues stand, come forth the flowers vpon small footestalkes, euery one by it selfe consisting of fiue small white leaues, with white threds in the middle, smelling also very sweet: after the flowers are past, there doe arise in the hot Countries, where they are naturall, round blacke berries, when they are ripe, wherein are contained many hard white crooked seedes, but neuer in this Countrey, as I said before: the roote disperseth it selfe into many branches, with many fibres annexed thereto.

### 2. *Myrtus minor, seu minore folio.* The smaller leafed Mirtle.

The smaller leafed Mirtle is a low shrub or bush, like vnto the former, but scarce rising so high, with branches spreading about the stemme, much thicker set with leaues then the former, smaller also, and pointed at the ends, of a little deeper greene colour, abiding greene also winter and summer, and very sweete likewise: the flowers are white like vnto the former, and as sweete, but shew not themselues so plentifull on the branches: the fruit is blacke in his naturall places, with seedes therein as the former.

### 3. *Myrtus minor rotundiore folio.* Boxe Mirtle.

Wee haue another sort of this small kinde of Mirtle, so like vnto the former both for smalnesse, deepe greene colour of the leaues, and thicke growing of the branches, that

that it will be thought of moſt, without good heede, and comparing the one with the other, to be the very ſame with the former: but if it bee well viewed, it will ſhew, by the roundneſſe at the ends of the leaues very like vnto the ſmall Boxe leaues, to be another differing kinde, although in nothing elſe. Wee nourſe them with great care, for the beautifull aſpect, ſweete ſent and raritie, as delights and ornaments for a garden of pleaſure, wherein nothing ſhould be wanting that art, care and coſt might produce and preſerue: as alſo to ſet among other euer greene plants to ſort with them.

### The Place.

Theſe, and many other ſorts of Mirtles grow in Spaine, Portugall, Italie, and other hot Countries in great aboundance, where they make their hedges of them: wee (as I ſaid) keepe them in this Countrey, with very great care and diligence.

### The Time.

The Mirtles doe flower very late with vs, not vntill Auguſt at the ſooneſt, which is the cauſe of their not fructifying.

### The Names.

They are called in Latine *Myrtus*, and in Engliſh Mirtle tree, without any other diuerſitie of names, for the generall title. Yet the ſeuerall kindes haue had ſeuerall denominations, in Plinies time, and others, as *Romana, Coniugala, Terentina, Egyptia, alba, nigra, &c.* which haue noted the differences, euen then well obſerued.

### The Vertues.

The Mirtle is of an aſtringent qualitie, and wholly vſed for ſuch purpoſes.

## Chap. CXIIII.

*Malus Punica ſiue Granata.* The Pomegranet tree.

THere are two kindes of Pomegranet trees, The one tame or manured, bearing fruit, which is diſtinguiſhed of ſome into two ſorts, of others into three, that is, into ſower, and ſweet, and into ſower ſweete. The other wilde, which beareth no fruite, becauſe it beareth double flowers, like as the Cherry, Apple, and Peach tree with double bloſſomes, before deſcribed, and is alſo diſtinguiſhed into two ſorts, the one bearing larger, the other leſſer flowers. Of the manured kinde wee haue onely one ſort (ſo farre as we know) for it neuer beareth ripe fruit in this our Countrey) which for the beautifull aſpect, both of the greene verdure of the leaues, and faire proportion and colour of the flowers, as alſo for the raritie, are nourſed in ſome few of their gardens that delight in ſuch raritie s: for in regard of the tenderneſſe, there is neede of diligent care; that is, to plant it againſt a brick wall, and defend it conueniently from the ſharpeneſſe of our winters, to giue his Maſter ſome pleaſure in ſeeing it beare flowers: And of the double kinde we haue as yet obtained but one ſort, although I ſhall giue you the knowledge and deſcription of another.

### 1. *Malus Punica ſatiua.* The tame Pomegranet tree.

This Pomegranet tree groweth not very high in his naturall places, and with vs ſomtimes it ſhooteth forth from the roote many browniſh twigges or branches, or if it bee pruned from them, and ſuffered to grow vp, it riſeth to bee ſeuen or eight foote high, ſpreading

1 *Myrtus latifolia maior.* The great leafed Myrtle. 2 *Myrtus angustifolia minor.* The small leafed Myrtle. 3 *Myrtus buxifolia minor.* The Boxe leafed Myrtle. 4 *Malus Granatus simplici flore.* The ordinary Pomegranet tree. 5 *Balaustium Romanum seu minus.* The lesser double flowred Pomegranet tree. 6 *Balaustium maius sive Cyprium.* The greater double flowred Pomegranet. 7 *Pseudocapsicum sive Amomum Plinij.* The Winter Cherry tree. 8 *Ficus Indica cum suo fructu.* The Indian Figgetree and his fruit.

ſpreading into many ſmall and ſlender branches, here and there ſet with thornes, and with many very faire greene ſhining leaues, like in forme and bigneſſe vnto the leaues of the larger Myrtle before deſcribed, euery one hauing a ſmall reddiſh foote-ſtalke vpon theſe branches: among the leaues come forth here and there, long, hard, and hollow reddiſh cups, diuided at the brimmes, wherein doe ſtand large ſingle flowers, euery one conſiſting of one whole leafe, ſmaller at the bottome then at the brimme, like bels, diuided as it were at the edges into fiue or ſix parts, of an orient red or crimſon colour in the hotter Countries; but in this it is much more delayed, and tendeth neare vnto a bluſh, with diuers threads in the middle. The fruit is great and round, hauing as it were a crowne on the head of it, with a thicke tough hard skinne or rinde, of a browniſh red colour on the outſide, and yellow within, ſtuffed or packt full of ſmall graines, euery one encompaſt with a thin skin, wherein is contained a cleare red iuyce or liquor, either of a ſweet (as I ſaid before) or ſower taſte, or betweene them both of a winie taſte: the roote diſperſeth it ſelfe very much vnder ground.

2. *Balauſtium maius ſiue Malus Punica ſilueſtris maior.*
The greater wilde or double bloſſomd Pomegranet tree.

The wilde Pomegranet is like vnto the tame in the number of purpliſh branches, hauing thornes, and ſhining faire greene leaues, ſomewhat larger then the former: from the branches likewiſe ſhoote forth flowers, farre more beautifull then thoſe of the tame or manured ſort, becauſe they are double, and as large as a double Prouince Roſe, or rather more double, of an excellent bright crimſon colour, tending to a ſilken carnation, ſtanding in browniſh cups or huskes, diuided at the brims vſually into foure or fiue ſeuerall points, like vnto the former, but that in this kinde there neuer followeth any fruit, no not in the Country, where it is naturally wilde.

3. *Balauſtium minus.* The ſmaller wilde Pomegranet tree.

This ſmaller kinde differeth from the former in his leaues, being of a darker greene colour, but not in the height of the ſtemme, or purpliſhneſſe of his branches, or thorns vpon them; for this doth ſhew it ſelfe more like vnto a wilde kind then it: the flowers hereof are much ſmaller, and not ſo thicke and double, of a deeper or ſadder red Orenge tawny colour, ſet alſo in ſuch like cups or huskes.

## The Place.

The tame or manured kinde groweth plentifully in Spaine, Portugall, and Italy, and other in other warme and hot countries. Wee (as I ſaid before) preſerue it with great care. The wilde I thinke was neuer ſeene in England, before Iohn Tradeſcante my very louing good friend brought it from the parts beyond the Seas, and planted it in his Lords Garden at Canterbury.

## The Time.

They flower very late with vs, that is, not vntill the middle or end of Auguſt, and the cold euenings or froſts comming ſo ſoone vpon it, doth not onely hinder it from bearing, but many times the ſharpe winters ſo pinch it, that it withereth it downe to the ground, ſo that oftentimes it hardly ſpringeth againe.

## The Names.

The name *Malus Punica* for the tree, and *Malum Punicum* for the fruit, or *Malus Granata*, and *Malum Granatum*, is the common name giuen vnto this tree, which is called in Engliſh the Pomegarnet or Pomegranet tree. The flowers of the tame kinde are called *Cytini*, as Dioſcorides ſaith, although Plinie ſeemeth either to make *Cytinus* to be the flower of the wilde kinde, or

*Balauſtium*

*Balaustium* to be the flower of both tame and wilde kinde: but properly, as I take it, *Cytinus* is the cup wherein the flower as well of the tame as wilde kinde doth stand; for vnto the similitude of them, both the flowers of *Asarum*, and the seede vessels of *Hyosciamus* are compared and resembled, and not vnto the whole flower: the barke or rinde of the fruit is called of diuers *Sidion*, and in the Apothecaries shops *Psidium*, and *cortex Granatorum*. The wilde kinde is called *Malus Punica silvestris*: In English, The wilde Pomegranet tree; the flower thereof is properly called *Balaustium*. The lesser kind is vsually called *Balaustium Romanum*, as the greater is called *Creticum* and *Cypxinum*, because they growe in Candy and Cyprus.

### The Vertues.

The vse of all these Pomegranets is very much in Physicke, to coole and binde all fluxibility both of body and humours: they are also of singular effect in all vlcers of the mouth, and other parts of the body, both of man and woman. There is no part of them but is applyed for some of these respects. The rinde also of the Pomegranet is vsed of diuers in stead of Gaules, to make the best sort of writing Inke, which is durable to the worlds end.

## Chap. CXV.

### *Amomum Plinij seu Pseudocapsicum.*
### Tree Night shade or the Winter Cherry tree.

I Haue adioyned this plant, for the pleasurable beauty of the greene leaues, and red berries. It groweth vp to be a yard or foure foote high at the most, hauing a small wooddy stemme or stocke, as bigge as ones finger or thumbe, couered with a whitish greene barke, set full of greene branches, and faire greene leaues, somewhat vneuen sometimes on the edges, narrower then any Night shade leaues, and very neare resembling the leaues of the *Capsicum*, or Ginny pepper, but smaller and narrower, falling away in the Winter, and shooting fresh in the Spring of the yeare: the flowers growe often two or three together, at the ioynts of the branches with the leaues, being white, opening starre-fashion, and sometimes turning themselues backe, with a yellow pointell in the middle, very like vnto the flowers of Night shade: after the flowers are past, come forth in their stead small greene buttons, which after turne to be pleasant round red berries, of the bignesse of small Cherries when they are ripe, which with vs vsually ripen not vntill the Winter, or about Christmas, wherein are contained many small whitish seede that are flat: all the whole plant, as well leaues and flowers as seede, are without either smell or taste: the roote hath many yellowish strings and fibres annexed vnto it.

### The Place.

The originall place hereof is not well knowne, but is thought to bee the West Indies. It hath been planted of long time in most of these Countries, where it abideth reasonable well, so that some care bee had thereof in the extreamity of the Winter.

### The Time.

It flowreth sometimes in Iune, but vsually in Iuly and August, and the fruit is not ripe (as is said) vntill the Winter.

The

### The Names.

This plant hath diuers names; for it is thought to be that kinde of *Amomum* that Plinie ſetteth downe. Dodonæus calleth it *Pſeudocapſicum*, for ſome likeneſſe in the leafe and fruit vnto the ſmall *Capſicum* or Ginnie Pepper, although much vnlike in the taſte and property. Others doe call it *Strichnodendron*, that is, *Solanum arboreſcens*, and wee in Engliſh according thereunto, Tree Night ſhade. But ſome Latine aſſes corrupting the Latine word *Amomum*, doe call it the Mumme tree. Dalechampius calleth it *Solanum Americum, ſeu Indicum*, and ſaith the Spaniards call it in their tongue, *Guindas de las Indias*, that is, *Ceraſa Indiana*, Indian Cherries, which if any would follow, I would not bee much againſt it: but many Gentlewomen doe call them Winter Cherries, becauſe the fruit is not throughly ripe vntill Winter.

### The Vertues.

I finde no Phyſicall property allotted vnto it, more then that by reaſon of the inſipidity, it is held to be cooling.

## Chap. CXVI.

### *Ficus Indica minor.* The ſmaller Indian Figgetree.

THis Indian Figgetree, if you will call it a tree (becauſe in our Country it is not ſo, although it groweth in the naturall hot Countries from a wooddy ſtemme or body into leaues) is a plant conſiſting only of leaues, one ſpringing out of another, into many branches of leaues, and all of them growing out of one leafe, put into the ground halfe way, which taking roote, all the reſt riſe out thereof, thoſe belowe for the moſt part being larger then thoſe aboue; yet all of them ſomewhat long, flat, and round pointed, of the thickneſſe of a finger vſually, and ſmalleſt at the lower end, where they are ioyned or ſpring out of the other leaues, hauing at their firſt breaking out a ſhew of ſmall, red, or browne prickes, thicke ſet ouer all the vpper ſide of the leaues, but with vs falling away quickly, leauing onely the markes where they ſtood: but they haue beſides this ſhew of great prickes, a few very fine, and ſmall, hard, white, and ſharpe, almoſt inſenſible prickes, being not ſo bigge as haires on the vnderſide, which will often ſticke in their fingers that handle them vnaduiſedly, neither are they to be diſcerned vnleſſe one look preciſely for them: the leaues on the vnderſide hauing none of thoſe other great pricks or marks at all, being of a faire freſh pale green colour: out of the vppermoſt leaues breake forth certaine greene heads, very like vnto leaues (ſo that many are deceiued, thinking them to be leaues, vntill they marke them better, and be better experienced in them) but that they growe round and not flat, and are broad at the toppe; for that out of the tops of euery of them ſhooteth out a pale yellow flower, conſiſting of two rowes of leaues, each containing fiue leaues a peece, laid open with certaine yellow threads, tipt with red in the middle: this greene head, vntill the flower be paſt, is not of halfe that bigneſſe that it attaineth vnto after, yet ſeldome or neuer commeth vnto perfection with vs, being long and round, like vnto a Figge, ſmall belowe, and greater aboue, bearing vpon the flat or broad head the marke of the flower; ſome holding ſtill on them the dryed leaues, and others hauing loſt them, ſhew the hollowneſſe which they haue in the toppe or middle of the head, the ſides round about being raiſed or ſtanding vp higher: this head or figge in our Country abideth greene on the outſide, and little or nothing reddiſh within (although it abide all the Winter, and the Summer following, as ſometimes it doth) for want of that heate and comfort of the Sunne it hath in his naturall place, where it groweth

reddiſh

reddish on the outside, and containing within it a bloudy red clammy iuyce, making the vrine of them that eate of them as red as bloud, which many seeing, were in doubt of themselues, lest their vrine were not very bloud; of what sweetnesse, like a figge, in the naturall places, I am not well assured, yet affirmed: but those that haue beene brought vnto me, whose colour on the outside was greenish, were of a reddish purple within, and contained within them round, small, hard seede, the taste was flat, watery, or insipide: the roote is neither great, nor disperseth it selfe very deepe or farre, but shooteth many small rootes vnder the vpper crust of the earth.

There is a greater kinde hereof, whose leaues are twice or thrice as bigge, which hauing been often brought vs, will seldome abide more then one Summer with vs, our Winters alwaies rotting the leaues, that it could not be longer kept.

## The Place.

This Indian Figge tree groweth dispersedly in many places of America, generally called the West Indies: The greater kinde in the more remote and hot Countries, as Mexico, Florida, &c. and in the Bermudas or Summer Islands, from whence wee haue often had it. The lesser in Virginia, and those other Countries that are nearer vnto vs, which better endureth with vs.

## The Time.

It flowreth with vs sometimes in May, or Iune; but (as I said) the fruit neuer commeth to perfection in this Country.

## The Names.

Diuers doe take it to bee *Opuntia Plinij*, whereof hee speaketh in the 21. Booke and 17. Chapter of his Naturall History: but he there saith, *Opuntia* is an herbe, sweete and pleasant to be eaten, and that it is a wonder that the roote should come from the leafe, and so to growe; which words although they descipher out the manner of the growing of this plant, yet because this is a kinde of tree, and not an herbe, nor to be eaten, it cannot bee the same: but especially because there is an herbe which groweth in the same manner, or very neare vnto it, one leafe standing on the toppe or side of another, being a Sea plant, fit to be eaten with vinegar and oyle (as many other herbes are that growe in the salt marshes, or neare the Sea, whereof Sea Purslane is one) which Clusius calleth *Lychen Marinus*, and (as Clusius saith) Cortusus very fitly called *Opuntia marina*, and out of doubt is the verie same *Opuntia* that Theophrastus maketh mention of, and Plinie out of him. Our English people in Virginia, and the Bermuda Island, where it groweth plentifully, because of the form of the fruit, which is somewhat like to a Peare, & not being so familiarly acquainted with the growing of Figs, sent it vnto vs by the name of the prickly Peare, from which name many haue supposed it to be a Peare indeede, but were therein deceiued.

## The Vertues.

There is no other especiall property giuen hereunto, by any that haue written of the West Indies, then of the colouring of the vrine, as is before said.

## CHAP. CXVII.

### *Yucca sive Iucca.* The supposed Indian Iucca.

THis rare Indian plant hath a great thicke tuberous roote (spreading in time into many tuberous heads) from the head whereof shooteth forth many long, hard, and guttured leaues, very sharpe pointed, compassing as it were one another at the bottome, of a grayish greene colour, which doe not fall away, but abide euer greene on the plant; from the middle whereof springeth forth (now and then, but not euery yeare) a strong round stalke, diuided into diuers branches, whereon stand diuers white, and somewhat large flowers, hanging downe their heads, consisting of six leaues, with diuers veines, of a weake reddish or blush colour, spread on the backe of the three outer leaues, especially from the middle of the leaues to the bottome, and not rising to the edge of the leafe of any flower, which fall away without bearing any seede in our Country, as farre as euer could be obserued either in the plant that Master Gerard kept a long time by him, or by Robinus at Paris his plant, which Master Gerard sent vnto him, or yet by that plant, that Vespasian Robin the sonne of old Robin sent vnto Master Iohn de Franqueuille, and now abideth and flourisheth in my Garden.

#### The Place.

It was first brought into England (as Master Gerard saith) from the West Indies, by a seruant of Master Thomas Edwards, an Apothecary of Exeter, and imparted to him, who kept it vnto his death: but perished with him that got it from his widow, intending to send it to his Country house.

#### The Time.

It flowreth not vntill Iuly, and the flowers fall away sodainely, after they haue beene blowne open a while.

#### The Names.

Master Gerard first as I thinke called it *Iucca*, supposing it to bee the true *Yuca* of *Theuet*, wherewith the Indians make bread, called *Cassana*: but the true *Iucca* is described to haue a leafe diuided into seuen or nine parts, which this hath not: Yet not knowing by what better name to call it, let it hold still his first imposition, vntill a fitter may be giuen it.

#### The Vertues.

Wee haue not heard of any, that hath either read, heard, or experimented the faculties hereof, nor yet whether it hath good or euill taste; for being rare, and possessed but by a few, they that haue it are loth to cut any thereof, for feare of spoiling and losing the whole roote.

Some haue affirmed, that in some parts of Turkie, where as they say this plant groweth, they make a kinde of cloth from the threads are found running through the leaues; but I finde the threads are so strong and hard, that this cannot be that plant the relators meane is vsed in that manner.

1 *Yuca sive Iucca.* The Indian Iucca. 2 *Arbor vitæ.* The tree of life. 3 *Arbor Iudæ.* Iudas tree. 4 *Laburnum.* Beane Trefoile. 5 *Cytisus.* Tree Trefoile.

## CHAP. CXVIII.

### *Arbor vitæ.* The tree of life.

THe tree of life riseth vp in some places where it hath stood long, to be a tree of a reasonable great bignesse and height, couered with a redder barke then any other tree in our Country that I know, the wood whereof is firme and hard, and spreadeth abroad many armes and branches, which againe send forth many smaller twigges, bending downewards; from which twiggy or slender branches, being flat themselues like the leaues, come forth on both sides many flat winged leaues, somewhat like vnto Sauine, being short and small, but not pricking, seeming as if they were brayded or folded like vnto a lace or point, of a darke yellowish greene colour, abiding greene on the branches Winter and Summer, of a strong resinous taste, not pleasing to most, but in some ready to procure casting, yet very cordiall and pectorall also to them that can endure it: at the toppes of the branches stand small yellowish dounie flowers, set in small scaly heads, wherein lye small, long, browniſh seede, which ripen well in many places, and being sowne, doe spring and bring forth plants, which with some small care will abide the extreamest Winters we haue.

### The Place.

The first or originall place where it naturally groweth, as farre as I can learne or vnderstand, is that part of America which the French doe inhabite, about the riuer of Canada, which is at the backe of Virginia Northward, and as it seemeth, first brought by them from thence into Europe, in the time of Francis the first French King, where it hath so plentifully encreased, and so largely beene distributed, that now few Gardens of respect, either in France, Germany, the Lowe-Countries, or England, are without it.

### The Time.

It flowreth in the end of May, and in Iune; the fruit is ripe in the end of August and Sptember.

### The Names.

All the Writers that haue written of it, since it was first knowne, haue made it to be *Thuyæ genus*, a kinde of Thuya, which Theophrastus compareth vnto a Cypresse tree, in his fifth Book and fifth Chapter: but *Omne simile non est idem*, and although it haue some likenesse, yet I verily beleeue it is *proprium sui genus*, a proper kinde of it owne, not to bee paralleld with any other. For wee finde but very few trees, herbes, or plants in America, like vnto those that growe in Europe, the hither part of Africa, or in the lesser Asia, as experience testifieth. Some would make it to be *Cedrus Lycia*, but so it cannot be. The French that first brought it, called it *Arbor vitæ*, with what reason or vpon what ground I know not: but euer since it hath continued vnder the title of the Tree of life.

### The Vertues.

It hath beene found by often experience, that the leaues hereof chewed in the morning fasting, for some few dayes together, haue done much good to diuers, that haue beene troubled with shortnesse of breath, and to helpe to expectorate thinne purulentous matter stuffing the lungs. Other properties I haue not heard that it hath; but doubtlesse, the hot resinous smell and taste

taste it hath, both while it is fresh, and after it hath beene long kept dry, doth euidently declare his tenuity of parts, a digesting and cleansing quality it is possessed with, which if any industrious would make tryall, hee should finde the effects.

## CHAP. CXIX.

### *Arbor Iudæ.* Iudas tree.

IVdas tree riseth vp in some places, where it standeth open from a wall, and alone free from other trees (as in a Garden at Battherſey, which sometimes agoe belonged to Master Morgan, Apothecary to the late Queene Elizabeth of famous memory) to be a very great and tall tree, exceeding any Apple tree in height, and equall in bignesse of body thereunto (as my selfe can testifie, being an eye witnesse thereof) when as it had many stalkes of flowers, being in the bud, breaking out of the body of the tree through the barke in diuers places, when as there was no bough or branch near them by a yard at the least, or yet any leafe vpon the tree, which they gathered to put among other flowers, for Nosegayes) and in other places it groweth to bee but an hedge bush, or plant, with many suckers and shootes from belowe, couered with a darke reddish barke, the young branches being more red or purplish: the flowers on the branches come forth before any shew or budding of leaues, three or foure standing together vpon a small foote-stalke, which are in fashion like vnto Pease blossomes, but of an excellent deepe purplish crimson colour: after which come in their places so many long, flat, large, and thinne cods, of a browniſh colour, wherein are contained small, blackish browne, flat, and hard seede: the roote is great, and runneth both deepe, and farre spreading in the earth: the leaues come forth by themselues, euery one standing on a long stalke, being hard & very round, like vnto the leafe of the largest *Asarum*, but not so thick, of a whitish green on the vpper side, and grayish vnderneath, which fall away euery yeare, and spring a fresh after the Spring is well come in, and the buds of flowers are sprung.

There is another of this kinde, growing in some places very high, somewhat like the former, and in other places also full of twiggy branches, which are greener then the former, as the leaues are likewise: the flowers of this kinde are wholly white, and the cods nothing so red or browne, in all other things agreeing together. *Flore albo.*

### The Place.

The former groweth plentifully in many places of Spaine, Italy, Prouence in France, and in many other places. The other hath beene sent vs out of Italy many times, and the seede hath sprung very well with vs, but it is somewhat tender to keepe in the Winter.

### The Time.

The flowers (as I said) appeare before the leaues, and come forth in Aprill and May, and often sooner also, the leaues following shortly after; but neither of them beareth perfect seede in our Country, that euer I could learne, or know by mine owne or others experience.

### The Names.

Some would referre this to *Cercis*, whereof Theophrastus maketh mention in his first Booke and eighteenth Chapter, among those trees that beare their fruit in cods, like as Pulse doe: and hee remembreth it againe in the fourteenth Chapter of his third Booke, and maketh it not vnlike the white

Poplar tree, both in greatnesse and whitenesse of the branches, with the leafe of an Iuie, without corners on the one part, cornered on the other, and sharpe pointed, greene on both sides almost alike, hauing so slender long footestalkes that the leaues cannot stand forthright, but bend downwards, with a more rugged barke then the white Poplar tree. Clusius thinketh this large description is but an ample description of the third kinde of Poplar, called *Lybica*, the Aspen tree, which Gaza translateth *Alpina*: but who so will well consider it, shall finde it neyther answerable to any Poplar tree, in that it beareth not cods as *Cercis* doth; nor vnto this *Arbor Iudæ*, because it beareth not white branches. Clusius saith also, that the learned of Mompelier in his time, referred it to *Colytea* of Theophrastus in his third booke and seuenteenth chapter, where he doth liken it to the leaues of the broadest leafed Bay tree, but larger and rounder, green on the vpperside, and whitish vnderneath, and whereunto (as he saith) Theophrastus giueth cods in the fourteenth chapter of the same third booke: and by the contracting of their descriptions both together, saith, they agree vnto this Iudas tree. But I find some doubts and differences in these places: for the *Colutæa* that Dioscorides mentioneth in the said fourteenth chapter of his third booke, hath (as he saith there) a leafe like vnto the Willow, and therefore cannot bee the same *Colutæa* mentioned in the seuenteenth chapter of the same third book, which hath a broade Bay leafe: indeede hee giueth seede in cods: but that with broade Bay leaues is (as he saith) without eyther flower or fruite; and besides all this, he saith the rootes are very yellow, which is not to bee found in this *Arbor Iudæ*, or Iudas tree: let others now iudge if these things can bee well reconciled together. Some haue for the likenesse of the cods vnto Beane cods, called it *Fabago*. And Clusius called it *Siliqua siluestris*. It is generally in these dayes called *Arbor Iudæ*, and in English after the Latine name, vntill a fitter may be had, Iudas tree.

### The Vertues.

There is nothing extant in any Author of any Physicall vse it hath, neyther hath any later experience found out any.

## CHAP. CXX.

### *Laburnum.* Beane Trefoile.

THere be three sorts of these codded trees or plants, one neere resembling another, whereof *Anagyris* of Dioscorides is one. The other two are called *Laburnum*; the larger whereof Matthiolus calleth *Anagyris altera*, and so doe some others also: the third is of the same kinde with the second, but smaller. I shall not for this our Garden trouble you or my selfe with any more of them then one, which is the lesser of the two *Liburnum*, in that it is more frequent, and that it will far better abide then the *Anagyris*, which is so tender, that it will hardly endure the winters of our Countrey: and the greater *Laburnum* is not so easily to be had.

### *Laburnum.* Beane Trefoile.

This codded tree riseth vp with vs like vnto a tall tree, with a reasonable great body, if it abide any long time in a place, couered with a smooth greene barke; the branches are very long, greene, pliant, and bending any way, whereon are set here and there diuers leaues, three alwaies standing together vpon a long stalk, being somwhat long, and not very narrow, pointed at the ends, greene on the vpperside, and of a siluer shining colour vnderneath, without any smell at all: at the ioynts of these branches, where the leaues stand, come forth many flowers, much like vnto broome flowers, but not so large

large or open, growing about a very long branch or ſtalke, ſometimes a good ſpan or more in length, and of a faire yellow colour, but not very deepe ; after which come flat thin cods, not very long or broade, but as tough and hard as the cods of Broome; wherein are contained blackiſh ſeede, like, but much leſſe then the ſeede of *Anagyris vera* (which are as big as a kidney beane, purpliſh and ſpotted) : the roote thruſteth down deepe into the ground, ſpreading alſo farre, and is of a yellowiſh colour.

### The Place.

This tree groweth naturally in many of the woods of Italie, and vpon the Alpes alſo, and is therefore ſtill accounted to be that *Laburnum* that Plinie calleth *Arbor Alpina*. It groweth in many gardens with vs.

### The Time.

It flowreth in May, the fruit or cods, and the ſeedes therein are ripe in the end of Auguſt, or in September.

### The Names.

This tree (as I ſaid before) is called of Matthiolus *Anagyris altera, ſiue ſecunda*, of Cordus, Geſner and others, eſpecially of moſt now adayes, *Laburnum*. It is probable in my opinion, that this ſhould bee that *Coluteа* of Theophraſtus, mentioned in the fourteenth Chapter of his third book with the leafe of a Willow ; for if you take any one leafe by it ſelfe, it may well reſemble a Willow leafe both for forme and colour, and beareth ſmall ſeed in cods like vnto pulſe as that doth. Of ſome it hath beene taken for a kinde of *Cytiſus*, but not truely. We call it in Engliſh, Beane Trefoile, in regard of his cods and ſeede therein, ſomewhat like vnto Kidney Beanes, and of the leaues, three alwayes ſtanding together, vntill a more proper name may bee giuen it.

### The Vertues.

There is no vſe hereof in Phyſicke with vs, nor in the naturall place of the growing, ſaue only to prouoke a vomit, which it will doe very ſtrongly.

## Chap. CXXI.

## *Cytiſus*. Tree Trefoile.

THere are ſo many ſorts of *Cytiſus* or Tree trefoiles, that if I ſhould relate them all, I ſhould weary the Reader to ouerlooke them, whereof the moſt part pertaine rather to a generall worke then to this abſtract. I ſhall not therefore trouble you with any ſuperfluous, but only with two, which we haue nourſed vp to furniſh waſte places in a garden.

### *Cytiſus Maranthæ*. Horned Tree Trefoile.

This Tree Trefoile which is held of moſt Herbariſts to bee the true *Cytiſus* of Dioſcorides, riſeth vp to the height of a man at the moſt, with a body of the bigneſſe of a mans thumbe, couered with a whitiſh bark, breaking forth into many whitiſh branches ſpreading farre, beſet in many places with ſmall leaues, three alwayes ſet together vpon a ſmall ſhort footeſtalke, which are rounder, and whiter then the leaues of Beane Trefoile : at the ends of the branches for the moſt part, come forth the flowers three or foure togethers, of a fine gold colour, and of the faſhion of Broome flowers, but not

not so large : after the flowers are past, there come in their places crooked flat thinne cods, of the fashion of a halfe moone, or crooked horne, whitish when they are ripe, wherein are contained blackish seede : the roote is hard and woody, spreading diuers wayes vnder the ground : the whole plant hath a pretty small hot sent.

*Cytisus vulgatior.* The common Tree Trefoile.

This *Cytisus* is the most common in this Land, of any the other sorts of tree trefoiles, hauing a blackish colourd barke, the stemme or body whereof is larger then the former, both for height and spreading, bearing also three leaues together, but smaller and greener then the former : the flowers are smaller, but of the same fashion and colour: the cods blackish and thin, and not very long, or great, but lesser then Broome cods, wherein there lyeth small blackish hard seede : the roote is diuersly disperfed in the ground.

The Place.

The first groweth in the kingdome of Naples, and no doubt in many other places of Italie, as Matthiolus saith. The other groweth in diuers places of France.

The Time.

They flower for the most part in May or Iune : the seede is ripe in August or September.

The Names.

The first (as I said) is thought of most to be the true *Cytisus* of Dioscorides, and as is thought, was in these later dayes first found by Bartholomæus Maranta of Naples, who sent it first to Matthiolus, and thereupon hath euer since beene called after his name, *Cytisus Maranthæ*. Some doe call it *Cytisus Lunatus*, because the cods are made somewhat like vnto an halfe Moone. We call it in English, Horned Tree Trefoile. The other is called *Cytisus vulgaris* or *vulgatior*; in English, The common Tree Trefoile, because we haue not any other so common.

The Vertues.

The chiefest vertues that are appropriate to these plants, are to procure milke in womens breasts, to fatten pullen, sheep &c. and to be good for bees.

## Chap. CXXII.

*Colutea.* The Bastard Sena Tree.

WEe haue in our Gardens two or three sorts of the Bastard Sena tree; a greater as I may so call it, and two lesser : the one with round thin transparent skins like bladders, wherein are the seede : the others with long round cods, the one bunched out or swelling in diuers places, like vnto a Scorpions tale, wherein is the seede, and the other very like vnto it, but smaller.

1. *Colutea Vesicaria.* The greater Bastard Sena with bladders.

This shrub or tree, or shrubby tree, which you please to call it, riseth vp to the height of a pretty tree, the stemme or stock being sometimes of the bignesse of a mans arme, couered with a blackish greene rugged barke, the wood whereof is harder then of an Elder,

Elder, but with an hollownesse like a pith in the heart or middle of the branches, which are diuided many wayes, and whereon are set at seuerall distances, diuers winged leaues, composed of many small round pointed, or rather flat pointed leaues, one set against another, like vnto Licoris, or the Hatchet Fitch; among these leaues come forth the flowers, in fashion like vnto Broome flowers, and as large, of a very yellow colour: after which appeare cleare thinne swelling cods like vnto thinne transparent bladders, wherein are contained blacke seede, set vpon a middle ribbe or sinew in the middle of the bladder, which if it be a little crushed betweene the fingers, will giue a cracke, like as a bladder full of winde. The roote groweth branched and woody.

2. *Colutea Scorpioides maior*. The greater Scorpion podded Bastard Sena.

This Bastard Sena groweth nothing so great or tall, but shooteth out diuersly, like vnto a shrub, with many shoots springing from the root: the branches are greener, but more rugged, hauing a white barke on the best part of the elder growne branches; for the young are greene, and haue such like winged leaues set on them as are to be seen in the former, but smaller, greener, and more pointed: the flowers are yellow, but much smaller, fashioned somewhat like vnto the former, with a reddish stripe downe the backe of the vppermost leafe: the long cods that follow are small, long and round, distinguished into many diuisions or dents, like vnto a Scorpions tayle, from whence hath risen the name: in these seuerall diuisions lye seuerall blacke seede, like vnto the seede of Fenigrecke: the roote is white and long, but not so woody as the former.

3. *Colutea Scorpioides minor*. The lesser Scorpion Bastard Sena.

This lesser Bastard Sena is in all things like the former, but somewhat lower, and smaller both in leafe, flower, and cods of seede, which haue not such eminent bunches on the cods to be seene as the former.

The Place.

They grow as Matthiolus saith about Trent in Italie, and in other places: the former is frequent enough through all our Countrey, but the others are more rare.

The Time.

They flower about the middle or end of May, and their seede is ripe in August. The bladders of the first will abide a great while on the tree, if they be suffered, and vntill the winde cause them to rattle, and afterwards the skins opening, the seed will fall away.

The Names.

The name *Colutea* is imposed on them, and by the iudgement of most writers, the first is taken to bee that *Colutea* of *Lipara* that Theophrastus maketh mention of, in the seuenteenth chapter of his third booke. But I should rather thinke that the *Scorpioides* were the truer *Colutea* of Theophrastus, because the long pods thereof are more properly to bee accounted *siliquæ*, then the former which are *vesicæ tumentes*, windy bladders, and not *siliquæ*: and no doubt but Theophrastus would haue giuen some peculiar note of difference if he had meant those bladders, and not these cods. Let others of iudgement be vmpeeres in this case; although I know the currant of writers since Matthiolus, doe all hold the former *Colutea vesicaria* to be the true *Colutea Liparæ* of Theophrastus. Wee call it in English, Bastard Sena, from Ruellius, who as I thinke first called it Sena, from the forme of the leaues. The second and third (as I said before) from the forme of the cods receiued their names, as it is in the titles and descriptions; yet they may as properly be called *Siliquosæ*, for that their fruite are long cods.

The

### The Vertues.

Theophrastus saith it doth wonderfully helpe to fatten sheepe: But sure it is found by experience, that if it be giuen to man it causeth strong castings both vpwards and downwards; and therefore let euery one beware that they vse not this in steede of good Sena, lest they feele to their cost the force thereof.

## CHAP. CXII.

*Spartum Hispanicum frutex.* Spanish Broome.

ALthough Clusius and others haue found diuers sorts of this shrubby Spartum or Spanish Broome, yet because our Climate will nourse vp none of them, and euen this very hardly, I shall leaue all others, and describe vnto you this one only in this manner: Spanish Broome groweth to bee fiue or sixe foote high, with a woody stemme below, couered with a darke gray, or ash-coloured barke, and hauing aboue many pliant, long and slender greene twigs, whereon in the beginning of the yeare are set many small long greene leaues, which fall away quickly, not abiding long on; towards the tops of these branches grow the flowers, fashioned like vnto Broom flowers, but larger, as yellow as they, and smelling very well; after which come small long cods, crested at the backe, wherein is contained blackish flat seede, fashioned very like vnto the Kidney beanes: the roote is woody, dispersing it selfe diuers waies.

### The Place.

This groweth naturally in many places of France, Spaine and Italie, wee haue it as an ornament in our Gardens, among other delightfull plants, to please the senses of sight and smelling.

### The Time.

It flowreth in the end of May, or beginning of Iune, and beareth seede, which ripeneth not with vs vntill it be late.

### The Names.

It is called *Spartium Graecorum*, and *Spartum frutex*, to distinguish it from the sedge or rush, that is so called also. Of some it is called *Genista*, and thought not to differ from the other *Genista*, but they are much deceiued; for euen in Spaine and Italie, the ordinary *Ganista* or Broome groweth with it, which is not pliant, and fit to binde Vines, or such like things withall as this is.

### The Vertues.

There is little vse hereof in Physicke, by reason of the dangerous qualitie of vomiting, which it doth procure to them that take it inwardly: but being applyed outwardly, it is found to helpe the *Sciatica*, or paine of the hippes.

CHAP.

The

1 *Colutæa vulgaris.* Ordinary bastard Sene. 2 *Periploca recta Virginiana.* Virginian Silke. 3 *Colutæa Scorpioides.* Scorpion bastard Sene. 4 *Spartum Hispanicum.* Spanish Broome. 5 *Ligustrum.* Priuet. 6 *Salvia variegata.* Party coloured Sage. 7 *Maiorana aurea.* Guilded Marierome.

2. Asclepias Syriaca spu. bast.

## Chap. CXXIIII.

*Periploca recta Virginiana.* Virginian Silke.

LEst this stranger should finde no hospitality with vs, being so beautifull a plant, or not finde place in this Garden, let him be here receiued, although with the last, rather then not at all. It riseth vp with one or more strong and round stalkes, three or foure foote high, whereon are set at the seuerall ioynts thereof two faire, long, and broad leaues, round pointed, with many veines therein, growing close to the stemme, without any foote-stalke: at the tops of the stalkes, and sometimes at the ioynts of the leaues, groweth forth a great bush of flowers out of a thinne skinne, to the number of twenty, and sometimes thirty or forty, euery one with a long foote-stalke, hanging downe their heads for the most part, especially those that are outermost, euery one standing within a small huske of greene leaues, turned to the stalkeward, like vnto the Lysimachia flower of Virginia before described, and each of them consisting of fiue small leaues a peece, of a pale purplish colour on the vpperside, and of a pale yellowish purple vnderneath, both sides of each leafe being as it were folded together, making them seeme hollow and pointed, with a few short chiues in the middle: after which come long and crooked pointed cods standing vpright, wherein are contained flat browniſh seede, dispersedly lying within a great deale of fine, soft, and whitish browne silke, very like vnto the cods, seede, and silke of *Asclepias*, or Swallow-wort, but that the cods are greater and more crooked, and harder also in the outer shell: the roote is long and white, of the bignesse of a mans thumbe, running vnder ground very far, and shooting vp in diuers places, the heads being set full of small white grumes or knots, yeelding forth many branches, if it stand any time in a place: the whole plant, as well leaues as stalkes, being broken, yeeld a pale milke.

### The Place.

It came to me from Virginia, where it groweth aboundantly, being raised vp from the seede I receiued.

### The Time.

It flowreth in Iuly, and the seede is ripe in August.

### The Names.

It may seeme very probable to many, that this plant is the same that Prosper Alpinus in the twenty fift Chapter of his Booke of Egyptian plants, nameth *Beidelsar*; and Honorius Bellus in his third and fourth Epistles vnto Clusius (which are at the end of his History of plants) calleth *Ossar frutex*: And Clusius himselfe in the same Booke calleth *Apocynum Syriacum, Palæstinum*, and *Ægyptiacum*, because this agreeth with theirs in very many and notable parts; yet verily I thinke this plant is not the same, but rather another kinde of it selfe: First, because it is not *frutex*, a shrub or wooddy plant, nor keepeth his leaues all the yeare, but loseth both leaues and stalks, dying down to the ground euery yeare: Secondly, the milke is not caustick or violent, as Alpinus and Bellus say *Ossar* is: Thirdly, the cods are more crooked then those of Clusius, or of Alpinus, which Honorius Bellus acknowledgeth to be right, although greater then those he had out of Egypt: And lastly, the rootes of these doe runne, whereof none of them make any mention. Gerard in his Herball giueth a rude figure of the plant, but a very true figure of the cods with seede, and saith the Virginians call it *wisanck*, and referreth it to the *Asclepias*, for the likenesse of the cods stuffed with silken

silken doune. But what reason Caspar Bauhinus in his *Pinax Theatri Botanici* had, to call it (for it is Clusius his *Apocynum Syriacum*) by the name of *Lapathum Ægyptiacum lactescens siliqua Asclepiadis*, I know none in the world: for but that he would shew an extreame singularity in giuing names to plants, contrary to all others (which is very frequent with him) how could he thinke, that this plant could haue any likenesse or correspondencie, with any of the kindes of Dockes, that euer he had seene, read, or heard of, in face, or shew of leaues, flowers, or seede; but especially in giuing milke. I haue you see (and that not without iust and euident cause) giuen it a differing Latine name from Gerard, because the *Asclepias* giueth no milke, but the *Periploca* or *Apocynum* doth; and therefore fitter to be referred to this then to that. And because it should not want an English name answerable to some peculiar property thereof, I haue from the silken doune called it Virginian Silke: but I know there is another plant growing in Virginia, called Silke Grasse, which is much differing from this.

### The Vertues.

I know not of any in our Land hath made any tryall of the properties hereof. Captaine Iohn Smith in his booke of the discouery and description of Virginia, saith, that the Virginians vse the rootes hereof (if his be the same with this) being bruised and applyed to cure their hurts & diseases.

## Chap. CXXV.

### *Ligustrum.* Primme or Priuet.

BEcause the vse of this plant is so much, and so frequent throughout all this Land, although for no other purpose but to make hedges or arbours in Gardens, &c. whereunto it is so apt, that no other can be like vnto it, to bee cut, lead, and drawne into what forme one will, either of beasts, birds, or men armed, or otherwise: I could not forget it, although it be so well knowne vnto all, to be an hedge bush growing from a woodly white roote, spreading much within the ground, and bearing manie long, tough, and plyant sprigs and branches, whereon are set long, narrow, and pointed sad greene leaues by couples at euery ioynt: at the tops whereof breake forth great tufts of sweete smelling white flowers, which when they are fallen, turne into small blacke berries, hauing a purple iuyce within them, and small seede, flat on the one side, with an hole or dent therein: this is seene in those branches that are not cut, but suffered to beare out their flowers and fruit.

### The Place.

This bush groweth as plentifully in the Woods of our owne Countrey, as in any other beyond the Seas.

### The Time.

It flowreth sometimes in Iune, and in Iuly; the fruit is ripe in August and September.

### The Names.

There is great controuersie among the moderne Writers concerning this plant, some taking it to be ῥάμνος of Dioscorides, other to be *Phillyrea* of Dioscorides, which followeth next after *Cyprus*. Plinie maketh mention of *Cyprus* in two places; in the one he saith, *Cyprus* hath the leafe of *Zizyphus*,

or the Iuiube tree: in the other he saith, that certain do affirme, that the *Cyprus* of the East Country, and the *Ligustrum* of Italy is one and the same plant: whereby you may plainly see, that our Priuet which is *Ligustrum*, cannot be that *Cyprus* of Plinie with Iuiube leaues: Besides, both Dioscorides & Plinie say, that *Cyprus* is a tree; but all know that *Ligustrum*, Priuet, is but an hedge bush: Againe, Dioscorides saith, that the leaues of *Cyprus* giue a red colour, but Priuet giueth none. Bellonius and Prosper Alpinus haue both recorded, that the true *Cyprus* of Dioscorides groweth plentifully in Egypt, Syria, and those Easterne Countries, and noursed vp also in Constantinople, and other parts of Greece, being a merchandise of much worth, in that they transport the leaues, and young branches dryed, which laid in water giue a yellow colour, wherewith the Turkish women colour the nailes of their hands, and some other parts of their bodies likewise, delighting much therein: and that it is not our *Ligustrum*, or Priuet, because *Cyprus* beareth round white seede, like Coriander seede, and the leaues abide greene alwaies vpon the tree, which groweth (if it bee not cut or pruined) to the height of the Pomegranet tree. I haue (I confesse) beyond the limits I set for this worke spoken concerning our Priuet, because I haue had the seede of the true *Cyprus* of Dioscorides sent mee, which was much differing from our Priuet, and although it sprang vp, yet would not abide any time, whereas if it had beene our Priuet, it would haue beene familiar enough to our Countrey.

### The Vertues.

It is of small vse in physicke, yet some doe vse the leaues in Lotions, that serue to coole and dry fluxes or sores in diuers parts.

## Chap. CXXVI.

*Saluia variegata.* Party coloured Sage. And
*Maiorana verficolor siue aurea.* Yellow or golden Marierome.

VNto all these flowers of beauty and rarity, I must adioyne two other plants, whose beauty consisteth in their leaues, and not in their flowers: as also to separate them from the others of their tribe, to place them here in one Chapter, before the sweete herbes that shall follow, as is fittest to furnish this our Garden of pleasure. This kinde of Sage groweth with branches and leaues, very like the ordinary Sage, but somewhat smaller, the chiefest difference consisteth in the colour of the leaues, being diuersly marked and spotted with white and red among the greene: for vpon one branch you shall haue the leaues seuerally marked one from another, as the one halfe of the leafe white, and the other halfe greene, with red shadowed ouer them both, or more white then greene, with some red in it, either parted or shadowed, or dasht here and there, or more greene then white, and red therein, eyther in the middle or end of the leafe, or more or lesse parted or striped with white and red in the greene, or else sometimes wholly greene the whole branch together, as nature listeth to play with such varieties: which manner of growing rising from one and the same plant, because it is the more variable, is the more delightfull and much respected.

There is another speckled Sage parted with white and greene, but it is nothing of that beauty to this, because this hath three colours euidently to bee discerned in euery leafe almost, the red adding a superaboundant grace to the rest.

*Maiorana aurea siue verficolor.* Yellow or golden Marierome.

This kinde of Marierome belongeth to that sort is called in Latine *Maiorana latifolia,*

*lia*, which Lobel setteth forth for *Hyssopus Græcorum genuina:* In English Winter Marierome, or pot Marierome: for it hath broader and greater leaues then the sweete Marierome, and a different vmbell or tuft of flowers. The difference of this from that set forth in the Kitchin Garden, consisteth chiefly in the leaues, which are in Summer wholly yellow in some, or but a little greene, or parted with yellow and greene more or lesse, as nature listeth to play: but in Winter they are of a darke or dead greene colour, yet recouering it selfe againe: the sent hereof is all one with the pot Marierome.

Wee haue another parted with white and greene, much after the manner with the former.

The Place, Time, Names, and Vertues of both these plants, shall be declared where the others of their kindes are specified hereafter, and in the Kitchen Garden, for they differ not in properties.

## CHAP. CXXVII.

### *Lauendula.* Lauender Spike.

AFter all these faire and sweete flowers before specified, I must needes adde a few sweete herbes, both to accomplish this Garden, and to please your senses, by placing them in your Nosegayes, or else where, as you list. And although I bring them in the end or last place, yet are they not of the least account.

#### 1. *Lauendula maior.* Garden Lauender.

Our ordinary Garden Lauender riseth vp with a hard wooddy stemme aboue the ground, parted into many small branches, whereon are set whitish, long, and narrow leaues, by couples one against another; from among which riseth vp naked square stalkes, with two leaues at a ioynt, and at the toppe diuers small huskes standing round about them, formed in long and round heads or spikes, with purple gaping flowers springing out of each of them: the roote is wooddy, and spreadeth in the ground: The whole plant is of a strong sweete sent, but the heads of flowers much more, and more piercing the senses, which are much vsed to bee put among linnen and apparrell.

There is a kinde hereof that beareth white flowers, and somewhat broader leaues, but it is very rare, and seene but in few places with vs, because it is more tender, and will not so well endure our cold Winters. *Flore albo.*

#### 2. *Lauendula minor seu Spica.* Small Lauender or Spike.

The Spike or small Lauender is very like vnto the former, but groweth not so high, neither is the head or spike so great and long, but shorter and smaller, and of a more purplish colour in the flower: the leaues also are a little harder, whiter, and shorter then the former; the sent also is somewhat sharper and stronger. This is not so frequent as the first, and is nourished but in some places that are warme, and where they delight in rare herbes and plants.

### The Place.

Lauender groweth in Spaine aboundantly, in many places so wilde, and little regarded, that many haue gone, and abiden there to distill the oyle thereof whereof great quantity now commeth ouer from thence vnto vs: and also in Lanquedocke, and Prouence in France.

### The Time.

It flowreth early in those hot Countries, but with vs not vntill Iune and Iuly.

### The Names.

It is called of ſome *Nardus Italica*, and *Lauendula*, the greater is called *Fœmina*, and the leſſer *Mas*. We doe call them generally Lauender, or Lauender Spike, and the leſſer Spike, without any other addition.

### The Vertues.

Lauender is little vſed in inward phyſicke, but outwardly; the oyle for cold and benummed parts, and is almoſt wholly ſpent with vs, for to perfume linnen, apparrell, gloues, leather, &c. and the dryed flowers to comfort and dry vp the moiſture of a cold braine.

## CHAP. CXXVIII.

### *Stœchas*. Sticadoue, Caſſidony, or French Lauender.

CAſſidony that groweth in the Gardens of our Countrey, may peraduenture ſomewhat differ in colour, as well as in ſtrength, from that which groweth in hotter Countries; but as it is with vs, it is more tender a great deale then Lauender, and groweth rather like an herbe then a buſh or ſhrub, not aboue a foote and a halfe high, or thereabouts, hauing many narrow long greene leaues like Lauender, but ſofter and ſmaller, ſet at ſeuerall diſtances together about the ſtalkes, which ſpread abroad into branches: at the tops whereof ſtand long and round, and ſometimes foure ſquare heads, of a darke greeniſh purple colour, compact of many ſcales ſet together; from among which come forth the flowers, of a blewiſh purple colour, after which follow ſeede veſſels, which are ſomewhat whitiſh when they are ripe, containing blackiſh browne ſeede within them: the roote is ſomewhat wooddy, and will hardly abide the iniuries of our cold Winters, except in ſome places onely, or before it haue flowred: The whole plant is ſomewhat ſweete, but nothing ſo much as Lauender.

### The Place.

Caſſidony groweth in the Iſlands Stœchades, which are ouer againſt Marſelles, and in Arabia alſo: we keep it with great care in our Gardens.

### The Time.

It flowreth the next yeare after it is ſowne, in the end of May, which is a moneth before any Lauender.

### The Names.

It is called of ſome *Lauendula ſilueſtris*, but moſt vſually *Stœchas*: in Engliſh, of ſome Stickadoue, or French Lauender; and in many parts of England, Caſſidony.

### The Vertues.

It is of much more vſe in phyſicke then Lauender, and is much vſed for old paines in the head. It is alſo held to be good for to open obſtructions, to expell melancholy, to cleanſe and ſtrengthen the liuer, and other inward parts, and to be a Pectorall alſo.

CHAP.

## Chap. CXXIX.

*Abrotanum femina sive Santolina.* Lauender Cotton.

THis Lauender Cotton hath many wooddy, but brittle branches, hoary or of a whitish colour, whereon are set many leaues, which are little, long, and foure square, dented or notched on all edges, and whitish also: at the tops of these branches stand naked stalkes, bearing on euery one of them a larger yellow head or flower, then eyther Tansie or Maudeline, whereunto they are somewhat like, wherein is contained small darke coloured seede: the roote is hard, and spreadeth abroad with many fibres: the whole plant is of a strong sweete sent, but not vnpleasant, and is in many places planted in Gardens, to border knots with, for which it will abide to be cut into what forme you thinke best; for it groweth thicke and bushy, very fit for such workes, besides the comely shew the plant it selfe thus wrought doth yeeld, being alwayes greene, and of a sweet sent; but because it quickly groweth great, and will soon runne out of forme, it must be euery second or third yeare taken vp, and new planted.

### The Place.

It is onely planted in Gardens with vs, for the vses aforesaid especially.

### The Time.

It flowreth in Iuly, and standeth long in the hot time of the yeare in his colour, and so will doe, if it be gathered before it haue stood ouer long.

### The Names.

Diuers doe call it as Matthiolus doth, *Abrotanum femina*, and *Santolina*; and some call it *Chamaecyparissus*, because the leaues thereof, are somewhat like the leaues of the Cypresse tree: Wee call it in English generally Lauender Cotton.

### The Vertues.

This is vsually put among other hot herbes, eyther into bathes, ointments, or other things, that are vsed for cold causes. The seede also is much vsed for the wormes.

## Chap. CXXX.

*Ocimum.* Bassill.

BAssill is of two sorts (besides other kindes) for this our Garden, the one whereof is greater, the other lesse in euery part thereof, as shall be shewed.

1. *Ocimum Citratum.* Common Bassill.

Our ordinary Garden Bassill hath one stalke rising from the root, diuersly branched out, whereon are set two leaues alwayes at a ioynt, which are broad, somewhat round, and pointed, of a pale greene colour, but fresh, a little snipt or dented about the edges, and of a strong or heady sent, somewhat like a Pomecitron, as many haue compared it, and thereof call it *Citratum*: the flowers are small and white, standing at the tops of the branches, with two smal leaues at euery ioynt vnder them, in some plants green, in o-

theirs browne vnder them : after which commeth blackiſh ſeede : the roote periſheth at the firſt approach of winter weather, and is to be new ſowen euery yeare.

2. *Ocimum minimum ſiue Gariophyllatum.* Buſh Baſill.

The buſh Baſill groweth not altogether ſo high, but is thicker ſpreade out into branches, whereon grow ſmaller leaues, and thicker ſet then the former, but of a more excellent and pleaſant ſmell by much : the flowers are white like the former, and the ſeede blacke alſo like it, and periſheth as ſuddenly, or rather ſooner then it, ſo that it requireth more paines to get it, and more care to nourſe it, becauſe we ſeldome or neuer haue any ſeede of it.

*Ocimum Indicum.* Indian Baſill.

The Indian Baſill hath a ſquare reddiſh greene ſtalke, a foote high or better, from the ioynts whereof ſpreade out many branches, with broade fat leaues ſet thereon, two alwayes together at the ioynt, one againſt another, as other Baſils haue, but ſomewhat deepely cut in on the edges, and oftentimes a little crumpled, ſtanding vpon long reddiſh footeſtalkes, of a darke purple colour, ſpotted with deeper purple ſpots, in ſome greater, in others leſſer: the flowers ſtand at the tops of the ſtalkes ſpike-faſhion, which are of a white colour, with reddiſh ſtripes and veines running through them, ſet or placed in darke purple coloured huskes : the ſeede is greater and rounder then the former, and ſomewhat long withall : the roote periſheth in like manner as the other former doe. The whole plant ſmelleth ſtrong, like vnto the other Baſils.

## The Place.

The two laſt ſorts of Baſils are greater ſtrangers in our Country then the firſt which is frequent, and only ſowen and planted in curious gardens. The laſt came firſt out of the Weſt Indies.

## The Time.

They all flower in Auguſt, or Iuly at the ſooneſt, and that but by degrees, and not all at once.

## The Names.

The firſt is vſually called *Ocimum vulgare*, or *vulgatius*, and *Ocimum Citratum*. In Engliſh, Common or Garden Baſill. The other is called *Ocimum minimum*, or *Gariophyllatum*, Cloue Baſill, or Buſh Baſill. The laſt eyther of his place, or forme of his leaues, being ſpotted and curled, or all, is called *Ocimum Indicum maculatum, latifolium & criſpum.* In Engliſh according to the Latine, Indian Baſill, broade leafed Baſill, ſpotted or curled Baſill, which you pleaſe.

## The Vertues.

The ordinary Baſill is in a manner wholly ſpent to make ſweet, or waſhing waters, among other ſweet herbes, yet ſometimes it is put into noſegayes. The Phyſicall properties are, to procure a cheerefull and merry heart, whervnto the ſeede is chiefly vſed in pouder, &c. and is moſt vſed to that, and to no other purpoſe.

CAPH.

1 *Santolina.* Lauender Cotton. 2 *Lauendula.* Lauender Spike. 3 *Stœchas.* Cassidony. 4 *Chamedrys.* Germander. 5 *Ocimum minus.* Fine Bassill. 6 *Marum.* Herbe Masticke. 7 *Maiorana.* Sweete Marieromé.

## Chap. CXXXI.

### *Maiorana.* Sweete Marierome.

WEe haue many sorts of Marierome; some that are sweete, and but Summer plants; others that are greater and not so sweet; and some also that are wilde. Of all these I will onely select some of the choisest that are fit for this place, and leaue the other for the next garden, and the garden of simples, or a generall worke: yet hereunto I will adioyne another sweete plant called Mastick, as participating neerer with them then with Time, whereunto many doe referre it.

#### 1. *Maiorana maior æstiua.* Common sweet Marierome.

The sweet Marierome that is most frequently sowen in our Country, is a low herbe little aboue a foote high when it is at the highest, full of branches, and small whitish soft roundish leaues, smelling very sweet: at the toppes of the branches stand diuers small scaly heads, like vnto knots, (and therefore of some called knotted Marierome) of a whitish greene colour, out of which come here and there small white flowers, and afterwards small reddish seede: the roote is composed of many small threds or strings, which perish with the whole plant euery yeare.

#### 2. *Maiorana tennifolia.* Marierome gentle.

This Marierome hath likewise diuers small branches, growing low, and not higher then the former, but hauing finer and smaller leaues, hoary and soft, but much sweeter: the heads are like vnto the former, and so are the flowers and seede, and the whole plant abiding but a Summer in the like manner.

#### 3. *Marum.* Herbe Mastick.

The neerer resemblance that this herbe hath with Marierome then with Tyme (as I said before) hath made me place it next vnto the small sweet Marierome. It riseth vp with a greater, and a more woody stalke then Marierome, two foote high or better in some places, where it liketh the ground and ayre, branching out on all sides towards the vpper part, leauing the stemme bare below, if it bee old, otherwise being young, thinly furnishing the branches from the bottome with small greene leaues, bigger then the leaues of any Tyme, and comming neere vnto the bignesse and forme of the last recited finer Marierome, but of a greener colour: at the toppes of the branches stand small white flowers on a head, which afterwards turne into a loose tuft of a long white hoary matter, like vnto soft doune, with some leaues vnderneath and about it, which abide not long on the stalkes, but are blowne away with the winde: the seede is so small if it haue any, that I haue not obserued it: the roote is threddy: the whole plant is of a sweete resinous sent, stronger then the Marierome, and abideth our winters, if it be carefully planted and regarded.

### The Place.

The sweete Marieromes grow naturally in hot Countreyes: the first in Spaine &c. the second is thought to come out of Syria, or Persia first into Italie, where they much esteeme it, and plant it curiously and carefully in pots, and set them in their windowes, beeing much delighted therewith for the sweet sent it hath. The first is vsually sowen euery yeare in most gardens with vs: but the second is very rare and daintie, and must as daintely be preserued, being more tender then the former. The herbe Mastick is thought to be first brought out of Candie, Clusius saith he found it in Spaine: It is planted by slippes, (and not sowen) in many gardens, and is much replanted

for

for increase, but prospereth onely, or more frequently, in loamie or clay grounds then in any other soyle.

### The Time.

The sweete Marieromes beare their knots or scaly heads in the end of Iuly, or in August. Herbe Masticke in Iune many times, or in the beginning of Iuly.

### The Names.

The first of the two sweet Marieromes called *Maiorana* in Latine *à maiore cura*, is taken of most writers to be the *Amaracus* or *Sampsuchum* of Dioscorides, Theophrastus and Plinie, although Galen doth seem a little to dissent therefrom. The other sweet Marierome hath his name in his title as much as can be said of it. The next is thought by the best of the moderne Writers to be the true *Marum* that Galen preferreth for the excellent sweetnesse, before the former Marierome in making the *Oleum*, or *vnguentum Amaricinum*, and seemeth to incline to their opinion that thought *Amaracus* was deriued from *Marum*. It is the same also that Galen and others of the ancient Writers make mention of, to go into the composition of the *Trochisci Hedychroi*, as well as *Amaracus* among the ingredients of the *Theriaca Andromachi*. In English we call it Masticke simply, or Herbe Mastick, both to distinguish it from that Tyme that is called Masticke Tyme, and from the Masticke Tree, or Gum, so called. Some of later times, and Clusius with them, haue thought this to be Dioscorides his *Tragoriganum*, which doth somewhat resemble it: but there is another plant that Matthiolus setteth forth for *Marum*, that in Lobels opinion and mine is the trueſt *Tragoriganum*, and this the trueſt *Marum*.

### The Vertues.

The sweete Marieromes are not onely much vsed to please the outward senses in nosegayes, and in the windowes of houses, as also in sweete pouders, sweete bags, and sweete washing waters, but are also of much vse in Physicke, both to comfort the outward members, or parts of the body, and the inward also: to prouoke vrine being stopped, and to ease the paines thereof, and to cause the feminine courses. Herbe Masticke is of greater force to helpe the stopping of vrine, then the Marierome, and is put into Antidotes, as a remedie against the poyson of venemous Beasts.

---

## CHAP. CXXXI.

### *Thymum.* Tyme.

THere are many kindes of Tyme, as they are vsually called with vs, some are called of the garden, and others wilde, which yet for their sweetnesse are brought into gardens, as Muske Tyme, and Lemon Tyme; and some for their beauty, as embroidered or gold yellow Tyme, and white Tyme. But the true Tyme of the ancient Writers, called *Capitatum*, as a speciall note of distinction from all other kindes of Tyme, is very rare to be seene with vs here in England, by reason of the tenderneſſe, that it will not abide our Winters. And all the other sorts that with vs are called garden Tymes, are indeede but kindes of wilde Tyme, although in the defect or want of the true Tyme, they are vsed in the stead of it. With the Tymes I must doe as I did with the Marieromes in the Chapter before, that is, reserue the most common in vse, for the common vse of the Kitchen, and shew you only those here, that are not put to that vse: and first with the true Tyme, because it is knowne but to a few.

1. *Thymum legitimum capitatum.* The true Tyme.

The true Tyme is a very tender plant, hauing hard and hoary brittle branches, ſpreading from a ſmall wooddy ſtemme, about a foote and a halfe high, whereon are ſet at ſeuerall ioynts, and by ſpaces, many ſmall, long, whitiſh, or hoary-greene leaues, of a quicke ſent and taſte: at the tops of the branches ſtand ſmall long whitiſh greene heads, ſomewhat like vnto the heads of *Stœchas*, made as it were of many leaues or ſcales, out of which ſtart forth ſmall purpliſh flowers (and in ſome white, as Bellonius ſaith) after which commeth ſmall ſeede, that ſoone falleth out, and if it be not carefully gathered, is ſoone loſt, which made (I thinke) Theophraſtus to write, that this Tyme was to be ſowne of the flowers, as not hauing any other ſeede: the root is ſmall and wooddy. This holdeth not his leaues in Winter, no not about Seuill in Spaine, where it groweth aboundantly, as Cluſius recordeth, finding it there naked or ſpoiled of leaues. And will not abide our Winters, but periſheth wholly, roote and all.

2. *Serpillum hortenſe ſiue maius.* Garden wilde Tyme.

The wilde Tyme that is cheriſhed in gardens groweth vpright, but yet is lowe, with diuers ſlender branches, and ſmall round greene leaues, ſomewhat like vnto ſmall fine Marierome, and ſmelling ſomewhat like vnto it: the flowers growe in roundels at the toppes of the branches, of a purpliſh colour: And in another of this kinde they are of a pure white colour.

There is another alſo like hereunto, that ſmelleth ſomewhat like vnto Muske; and therefore called Muske Tyme, whoſe greene leaues are not ſo ſmall as the former, but larger and longer.

3. *Serpillum Citratum.* Lemon Tyme.

The wilde Tyme that ſmelleth like vnto a Pomecitron or Lemon, hath many weake branches trayling on the ground, like vnto the firſt deſcribed wilde Tyme, with ſmall darke greene leaues, thinly or ſparſedly ſet on them, and ſmelling like vnto a Lemon, with whitiſh flowers at the toppes in roundels or ſpikes.

4. *Serpillum aureum ſiue verſicolor.* Guilded or embroidered Tyme.

This kinde of wilde Tyme hath ſmall hard branches lying or leaning to the ground, with ſmall party coloured leaues vpon them, diuided into ſtripes or edges, of a gold yellow colour, the reſt of the leafe abiding greene, which for the variable mixture or placing of the yellow, hath cauſed it to be called embroidered or guilded Tyme.

The Place.

The firſt groweth as is ſaid before, about Seuill in Spaine, in very great aboundance as Cluſius ſaith; and as Bellonius ſaith, very plentifully on the mountaines through all Greece. The others growe ſome in this Country, and ſome in others: but wee preſerue them with all the care wee can in our gardens, for the ſweete and pleaſant ſents and varieties they yeeld.

The Time.

The firſt flowreth not vntill Auguſt; the reſt in Iune and Iuly.

The Names.

Their names are ſeuerally ſet downe in their titles, as is ſufficient to diſtinguiſh them; and therefore I ſhall not neede to trouble you any further with them.

The

The Vertues.

The true Tyme is a speciall helpe to melancholicke and splenetick diseases, as also to flatulent humours, either in the vpper or lower parts of the body. The oyle that is Chimically drawne out of ordinary Tyme, is vsed (as the whole herbe is, in the stead of the true) in pils for the head and stomach. It is also much vsed for the toothach, as many other such like hot oyles are.

## Chap. CXXXII.

### *Hyssopus.* Hyssope.

THere are many varieties of Hyssope, beside the common or ordinary, which I reserue for the Kitchen garden, and intend onely in this place to giue you the knowledge of some more rare: *viz.* of such as are noursed vp by those that are curious, and fit for this garden: for there are some other, that must be remembred in the Physicke garden, or garden of Simples, or else in a generall worke.

### 1. *Hyssopus foliis niueis.* White Hyssope.

This white Hyssope is of the same kinde and smell with the common Hyssope; but differeth, in that this many times hath diuers leaues, that are wholly of a white colour, with part of the stalke also: others are parted, the one halfe white, the other halfe greene, and some are wholly greene, or with some spots or stripes of white within the greene, which makes it delightfull to most Gentlewomen.

### 2. *Hyssopus foliis cinereis.* Russet Hyssope.

As the last hath party coloured leaues, white and greene, so this hath his leaues of an ash-colour, which of some is called russet; and hath no other difference either in forme or smell.

### 3. *Hyssopus aureus.* Yellow or golden Hyssope.

All the leaues of this Hyssope are wholly yellow, or but a little greene in them, and are of so pleasant a colour, especially in Summer, that they prouoke many Gentlewomen to weare them in their heads, and on their armes, with as much delight as many fine flowers can giue: but in Winter their beautifull colour is much decayed, being of a whitish greene, yet recouer themselues againe the next Summer.

### 4. *Hyssopus surculis densis.* Double Hyssope.

As this kinde of Hyssope groweth lower then the former or ordinary kinde, so it hath more branches, slenderer, and not so wooddy, leaning somewhat downe toward the ground, so wonderfully thicke set with leaues, that are like vnto the other, but of a darker greene colour, and somewhat thicker withall, that it is the onely fine sweete herbe, that I know fittest (if any be minded to plant herbes) to set or border a knot of herbes or flowers, because it will well abide, and not growe too wooddy or great, nor be thinne of leaues in one part, when it is thicke in another, so that it may be kept with cutting as smooth and plaine as a table. If it be suffered to growe vp of it selfe alone, it riseth with leaues as before is specified, and flowreth as the common doth, and of the same sent also, not differing in any thing, but in the thicknesse of the leaues on the stalkes and branches, and the aptnesse to be ordered as the keeper pleaseth.

*Chamædrys.* Germander.

Left Germander should be vtterly forgotten, as not worthy of our Garden, seeing many (as I said in my treatise or introduction to this Garden) doe border knots therewith: let me at the least giue it a place, although the last, being more vsed as a strewing herbe for the house, then for any other vse. It is (I thinke) sufficiently knowne to haue many branches, with small and somewhat round endented leaues on them, and purplish gaping flowers: the rootes spreading far abroad, and rising vp againe in many places.

The Place.

These Hyssopes haue beene most of them noursed vp of long time in our English Gardens, but from whence their first originall should be, is not well knowne. The Germander also is onely in Gardens, and not wilde.

The Time.

They flower in Iune and Iuly.

The Names.

The seuerall names whereby they are knowne to vs, are set forth in their titles; and therefore I neede not here say more of them then onely this, that neyther they here set downe, nor the common or ordinary sort, nor any of the rest not here expressed, are any of them the true Hyssope of the ancient Greeke Writers, but *supposititiæ*, vsed in the stead thereof. The Germander, from the forme of the leaues like vnto small oaken leaues, had the name *Chamædrys* giuen it, which signifieth a dwarfe Oake.

The Vertues.

The common Hyssope is much vsed in all pectorall medicines, to cut fleagme, and to cause it easily to be auoided. It is vsed of many people in the Country, to be laid vnto cuts or fresh wounds, being bruised, and applyed eyther alone, or with a little Sugar. It is much vsed as a sweet herbe, to be in the windowes of an house. I finde it much commended against the Falling Sickenesse, especially being made into Pils after this manner: Of Hyssope, Horhound, and Castor, of each halfe a dramme, of Peony rootes (the male kinde is onely fit to be vsed for this purpose) two drams, of *Assa fœtida* one scruple: Let them be beaten, and made into pils with the iuyce of Hyssope; which being taken for seuen dayes together at night going to bed, is held to be effectual to giue much ease, if not thoroughly to cure those that are troubled with that disease. The vse of Germander ordinarily is as Tyme, Hyssope, and other such herbes, to border a knot, whereunto it is often appropriate, and the rather, that it might be cut to serue (as I said) for a strewing herbe for the house among others. For the physicall vse it serueth in diseases of the splene, and the stopping of vrine, and to procure womens courses.

Thus haue I led you through all my Garden of Pleasure, and shewed you all the varieties of nature noursed therein, pointing vnto them, and describing them one after another. And now lastly (according to the vse of our old ancient Fathers) I bring you to rest on the Grasse, which yet shall not be without some delight, and that not the least of all the rest.

CHAP.

1 *Thymum legitimum.* The true Tyme. 2 *Serpillum maius hortense.* Garden wilde Tyme. 3 *Serpillum Citratum.* Lemon Tyme. 4 *Hyssopus versicolor sive aureus.* Golden Hyssope. 5 *Chamædrys.* Germander. 6 *Spartum Austriacum sive Gramen plumarium minus.* The lesser feather Grasse. 7 *Gramen striatum vel sulcatum.* Painted Grasse or Ladies Laces.

## Chap. CXXXIII.

### *Gramina.* Grasses.

THere are among an infinite number (as I may so say) of Grasses, a few onely which I thinke fit to be planted in this Garden, both for the rarity of them, and also for your delight, and the excellent beauty that is in them aboue many other plants. One of them hath long agoe bin respected, and cherished in the country gardens of many Gentlewomen, and others. The others are knowne but vnto a few.

### 1. *Gramen striatum.* Painted Grasse or Ladies laces.

This kinde of Grasse hath many stiffe, hard, round stalkes, full of ioynts, whereon are set at euery ioynt one long leafe, somewhat broad at the bottome, where it compasseth the stalke, and smaller to the end, where it is sharpe pointed, hard or rough in handling, and striped all the length of the leafe with white streakes or lines, that they seeme party coloured laces of white and greene: the tops of the stalkes are furnished with long spikie tufts, like vnto the tufts of Couch Grasse: the rootes are small, white, and threddy, like the rootes of other Grasses.

### 2. *Gramen Plumarium minus.* The lesser Feather-Grasse.

This lesser Feather-Grasse hath many small, round, and very long leaues or blades, growing in tufts, much finer and smaller then any other Grasse that I know, being almost like vnto haires, and of a fresh greene colour in Summer, but changing into gray, like old hay in Winter, being indeede all dead, and neuer reuiuing; yet hardly to be plucked away vntill the Spring, and then other greene leaues or rushes rise vp by them, and in their stead, and are aboue a foote in length: from the middle of these tufts come forth rounder and bigger rushes, which are the stalkes, and which haue a chaffie round eare about the middle thereof, which when it is full growne, is somewhat higher then the toppes of the leaues or rushes, opening it selfe (being before close) at the top, and shewing forth three or foure long ayles or beards, one aboue another, which bend themselues a little downewards (if they stand ouer long before they are gathered, and will fall off, and be blowne away with the winde) being so finely feathered on both sides, all the length of the beard, and of a pale or grayish colour, that no feather in the taile of the Bird of Paradise can be finer, or to be compared with them, hauing sticking at the end of euery one of them, within the eare, a small, long, whitish, round, hard, and very sharpe pointed graine, like vnto an oaten graine, that part of the stalke of the feather that is next vnder it, and aboue the seede for some two or three inches, being stiffe and hard, and twining or curling it selfe, if it be suffered to stand too long, or to fall away, otherwise being straight as the feather it selfe: the roote is composed of many long, hard, small threddy strings, which runne deepe and far, and will not willingly be remoued, in that it gaineth strength euery yeare by standing.

### 3. *Gramen Plumarium maius.* The greater Feather-Grasse.

The greater Feather-Grasse is like vnto the lesser, but that both the leaues and the feathers are greater, and nothing so fine, grosser also, and of lesse beauty and respect, though whiter then it; and therefore is not so much regarded: for I haue knowne, that many Gentlewomen haue vsed the former lesser kinde, being tyed in tufts, to set them in stead of feathers about their beds, where they haue lyen after childe-bearing, and at other times also, when as they haue been much admired of the Ladies and Gentles that haue come to visit them.

### The Place.

The first of these Grasses, as Lobel saith, groweth naturally in the woods and hils of Sauoy. It hath long agoe beene receiued into our English gardens.

dens. The second, as Clusius saith, in Austria, from whence also (as I take it) the greater came, and are both in the gardens of those, that are curious obseruers of these delights.

## The Time.

The first is in its pride for the leaues all the Spring and Summer, yeelding his bush in Iune. The other giue their feather-like sprigs in Iuly and August, and quickly (as I said) are shed, if they be not carefully gathered.

## The Names.

The first is called by Lobel *Gramen sulcatum*, or *striatum album*; of others *Gramen pictum*. The French call it *Aiguellettes d' armes*, of the fashion that their Ensignes, Pennons, or Streamers vsed in wars were of, that is, like vnto a party coloured curtaine. In English vsually Ladies laces, and Painted Grasse. The first of the other two is called *Gramen plumarium* or *plumosum*, and *minus* is added for the distinction of it. Clusius calleth it *Spartum Austriacum*, of the likenesse and place where he found it. The last is called *Gramen plumarium*, or *plumosum maius*, The greater Feather-Grasse.

## The Vertues.

These kindes of Grasses are not in any time or place that I doe heare of applyed to any Physicall vse; and therefore of them I will say no more; but here I will end the prime part of this worke.

## The Vertues.

The Single Vertues of Cradles are not in any time or place that I do here of applyed to any Particularity; and therefore of them I will say no more: but here I will end the prime part of this work.

# THE ORDERING OF THE KITCHEN GARDEN.

## Chap. I.

*The ſituation of a Kitchen Garden, or Garden of Herbes, and what ſort of manure is fitteſt to helpe the decaying of the ſoyle thereof.*

HAuing giuen you the beſt rules and inſtructions that I can for your flower Garden, and all the flowers that are fit to furniſh it, I now proceede to your herbe garden, which is not of the leaſt reſpect belonging to any mans houſe, nor vtterly to bee neglected for the many vtilities are to be had from it, both for the Maſters profit and pleaſure, and the meynies content and nouriſhment: all which if I ſhould here ſet down, I had a large field to wander in, and matter ſufficient to entreat of, but this worke permitteth not that libertie: and I thinke there are but few but eyther know it already, or conceiue it ſufficiently in their minds. Paſſing therefore no further in ſuch diſcourſes, I come to the matter in hand, which is to ſhew you where the fitteſt place is for an herbe garden. As before I ſhewed you that the beautie of any worthy houſe is much the more commended for the pleaſant ſituation of the garden of flowers, or of pleaſure, to be in the ſight and full proſpect of all the chiefe and choiſeſt roomes of the houſe; ſo contrariwiſe, your herbe garden ſhould bee on the one or other ſide of the houſe, and thoſe beſt and choyſe roomes: for the many different ſents that ariſe from the herbes, as Cabbages, Onions, &c. are ſcarce well pleaſing to perfume the lodgings of any houſe; and the many ouer-tures and breaches as it were of many of the beds thereof, which muſt neceſſarily bee, are alſo as little pleaſant to the ſight. But for priuate mens houſes, who muſt like their habitations as they fall vnto them, and cannot haue time or meanes to alter them, they muſt make a vertue of neceſſity, and conuert their places to their beſt aduantage, by making their profit their chiefeſt pleaſure, and making one place ſerue for all vſes. The choyce of ground for this Garden, is (as I ſaid before) where it is fat, fertill and good, there needeth the leſſe labour and coſt: and contrariwiſe, where it is cold, wet, dry or barren, there muſt bee the more helpes ſtill added to keepe it in heart. For this Garden by reaſon of the much and continuall ſtirring therein, the herbes and rootes drawing out the ſubſtance of the fertilitie thereof more aboundantly then in the former, muſt be continually holpen with ſoyle, or elſe few things of goodneſſe or worth will come forward therein. The ſtable ſoyle of horſes is beſt and more proper for any colde grounds, for being the hotteſt, it will cauſe any the ſeedes for this Garden to proſper well, and be more forward then in any other ground that is not ſo holpen. The ſtable ſoyle of Cattell is of a colder and moiſter nature, and is therefore more proper for the

the hot ſandy or grauelly grounds, and although it bee longer before it bee brought to mould then that of horſes, yet it will outlaſt it more then twice ſo long. Let euery one therefore take according to the nature of the ground ſuch helpes as are moſt fit and conuenient, as I haue here and before ſhewed. But I doe here ingenuouſly confeſſe my opinion of theſe forcings and helpings of ground, that howſoeuer it doth much good to ſome particular things, which becauſe they delight in heate, and cannot be brought to perfection without it in this our Countrey, which is colder then their naturall from whence they are brought, muſt therfore haue artificiall helpes to forward them; yet for many other things the compoſt doth much alter and abate the naturall vigour, and quickeneſſe of taſte, that is perceiued in them that grow in a naturall fat or ſandy ſoile that is not ſo holpen.

## Chap. II.

### *The forme of a Garden of herbes for neceſſary vſes, with the ordering thereof.*

AS our former Garden of pleaſure is wholly formable in euery part with ſquares, trayles, and knots, and to bee ſtill maintained in their due forme and beautie: ſo on the contrary ſide this Garden cannot long conſerue any forme, for that euery part thereof is ſubiect to mutation and alteration. For although it is conuenient that many herbes doe grow by themſelues on beds, caſt out into ſome proportion fit for them, as Tyme, Hiſſope, Sage, &c. yet many others may bee ſowen together on a plot of ground of that largeneſſe that may ſerue euery mans particular vſe as he ſhall haue occaſion to employ it, as Reddiſh, Lettice and Onions, which after they are growne vp together may be drawne vp and taken away, as there is occaſion to ſpend them: but Carrots or Parſneps being ſowen with others muſt bee ſuffered to grow laſt, becauſe they require a longer time before they be fit to be taken vp. Other herbes require ſome large compaſſe of ground whereon they may grow of themſelues without any other herbes growing among them, as Artichokes, Cowcumbers, Melons, Pompions. And ſome will doe ſo with their Cabbages alſo, but the beſt and moſt frugall way now vſed, is to plant them round about the border of your plot or ground whereon you plant Cowcumbers, Pompions, or other things, in that by this meanes ſo much ground will be well ſaued, and the other things be no whit hindered thereby, which elſe a great deale of ground muſt be employed for them apart. So that by this that I haue here ſaid, you may perceiue the forme of this Garden is for the moſt part, to bee ſtill out of forme and order, in that the continuall taking vp of the herbes and rootes that are ſowen and planted, cauſeth the beds or parts of this Garden to lye broken, diſmembered, and out of the order that at the firſt it was put into. Remember herewithall that (as I ſaid before) this Garden requireth the continuall helpe of ſoyle to be brought into it, in that the plenty of theſe manner of herbes and rootes doe ſo much waſte the fertilitie and fatneſſe of the ground, that without continuall refreſhing it would quickly become ſo poore and barren, that it would not yeelde the worth of the ſeede. The ordinary time to ſoyle a Garden, is to bring in manure or dung before Chriſtmas, and eyther bury it ſome ſmall depth, not too deepe, or elſe to lay it vpon the ground that the winter froſtes may pierce it, and then turne it ſhallow into the ground to ſow your ſeeds in the Spring.

## CHAP. III.

### *How to order diuers Garden herbes, both for their sowing, spending, and gathering of the seede.*

OVr chiefest and greatest Gardiners now adaies, doe so prouide for themselues euery yeare, that from their owne grounds they gather the seede of many herbes that they sowe againe: for hauing gained the best kind of diuers herbes, they will be still furnished with the same, and be not to seeke euery yeare for new that oftentimes will not yeelde them halfe the profit that their choyce seede will: I say of many herbes, but not of all; for the best of them all hath not ground sufficient for all sorts, nor will our climate bring some to that perfection that other forraine doth, and therefore the seede of some things are continually brought from beyond Sea vnto vs. And againe although our chiefe Gardiners doe still prouide their owne seede of diuers things from their owne ground, because as I said it is of the best kinde, yet you must vnderstand also, that good store of the same sortes of seeds are brought from beyond the Seas, for that which is gathered in this Land is not sufficient to serue euery mans vse in the whole Kingdome by many parts; yet still it is true, that our English seede of many things is better then any that commeth from beyond the Seas: as for example, Reddish, Lettice, Carrots, Parsneps, Turneps, Cabbages, and Leekes, of all which I intend to write in this place; for these are by them so husbanded, that they doe not sow their owne grounds with any other seede of these sorts but their owne: which that you may know the manner how to doe, I will here set it downe, that euery one may haue the best directions if they will follow them. Of Reddish there are two sorts, one more early then the other: they vse therfore to sow their early Reddish first, that they may haue the earliest profit of them, which is more worth in one fortnight, then in a moneth after. And to effect this they haue some artificiall helps also; which are these: They vse to digge vp a large plot of ground where they intend to sow their seede a little before or after Christmas, casting it into high balkes or ridges fiue or sixe foote asunder, which they suffer to lye and take all the extreame frosts in Ianuary to mellow the earth, and when the frostes are past, they then beginne to bring into it good store of fresh stable dung, which they laye neyther too deepe nor too thicke, and couer it with the mould a hand breadth thickenesse aboue the dung, which doth giue such warmth and comfort to whatsoeuer is sowen thereon, that it forceth it forward much sooner then any other way can doe: And to preuent both the frostes, and the cold bitter windes which often spoyle their seede new sprung vp, they vse to set great high and large mattes made of reedes, tyed together, and fastened vnto strong stakes, thrust into the ground to keepe them vp from falling, or being blowne down with the winde; which mattes they place on the North and East side to breake the force of these winds, and are so sure and safe a defence, that a bricke wall cannot better defend any thing vnder it, then this fence will. In this manner they doe euery yeare to bring forward their seede to gaine the more by them, and they that will haue Reddish early, must take the same course. The other sort of Reddish for the most part is sowen in Februarie, a fortnight after the other at the least, and likewise euery moneth after vnto September, that they may haue young continually. For the blacke Reddish, although many in many places doe sowe it in the same time, and in the same manner that the ordinary is sowen, yet the nature thereof is to runne vp to seede more speedily then the other, if it haue so rich ground to grow vpon, and therefore the best time to sow it is in August, that so it may abide all winter, wherein is the chiefest time for the spending thereof, and to keepe it vntill the beginning of the next yeare from running vp to seede: the gathering whereof, as also of the other sort, is all after one manner, that is, to be pulled vp when the pods change whitish, and then hanged vpon bushes, pales, or such other thing, vntill they bee thorough dry, and then beaten or thrashed out vpon a smooth plancher, or vpon clothes, as euery ones store is, and their conueniencie. Lettice is sowen oftentimes with the early Reddish, in the same manner before said, that they may haue Lettice likewise as early as the time of the year will permit them, which

they

they pull vp where they grow too thicke, ſpending them firſt, and ſo taking vp from time to time, vntill they ſtand two foote in ſunder one from another, and beginne to ſpindle and ſhoote vp for ſeede. In this is vſed ſome arte to make the plants ſtrong to giue the better ſeede without danger of rotting or ſpoyling with the wet, which often happeneth to thoſe about whom this caution is not obſerued: Before your Lettice is ſhot vp, marke out the choyſeſt and ſtrongeſt plantes which are fitteſt to grow for ſeede, and from thoſe when they are a foote high, ſtrippe away with your hand the leaues that grow loweſt vpon the ſtalke next the ground, which might rot, ſpoyle or hinder them from bearing ſo good ſeede; which when it is neere to be ripe, the ſtalkes muſt be cut off about the middle, and layde vpon mats or clothes in the Sunne, that it may there fully ripen and be gathered; for it would be blowne away with the winde if it ſhould be ſuffered to abide on the ſtalkes long. Parſneps muſt be ſowen on a deep trenched mellow ground, otherwiſe they may run to ſeede the firſt yeare, which then are nothing worth: or elſe the rootes will be ſmall ſtaruelings and ſhort, and runne into many ſpires or branches, whereby they will not bee of halfe the worth. Some vſe to ſow them in Auguſt and September, that ſo they may bee well growne to ſerue to ſpend in Lent following, but their beſt time is in February, that the Summers growth may make them the fairer and greater. When they runne vp to ſeede, you ſhall take the principall or middle heades, for thoſe carry the Maſter ſeede, which is the beſt, and will produce the faireſt rootes againe. You ſhall hardly haue all the ſeede ripe at one inſtant, for vſually the chiefeſt heads will be fallen before the other are ripe: you muſt therefore ſtill looke them ouer, and cut them as they ripen. Carrots are vſually ſowen in March and Aprill, and if it chance that ſome of them doe runne vp for ſeede the ſame year, they are to be weeded out, for neyther the ſeed nor roots of them are good: You muſt likewiſe pull them vp when they are too thicke, if you will haue them grow fair, or for ſeed, that they may grow at the leaſt three or foure foot in ſunder: the ſtalkes of Carrots are limber, and fall downe to the ground; they muſt therefore be ſuſtained by poles layde acroſſe on ſtalkes thruſt into the ground, and tyed to the poles and ſtalkes to keepe them vp from rotting or ſpoyling vpon the ground: the ſeed hereof is not all ripe at once, but muſt be tended and gathered as it ripeneth, and layd to dry in ſome dry chamber or floore, and then beaten out with a ſtick, and winnowed from the refuſe. Turneps are ſowne by themſelues vpon a good ground in the end of Iuly, and beginning of Auguſt, to haue their rootes beſt to ſpend in winter; for it often happeneth that thoſe ſeedes of Turneps that are ſowen in the Spring, runne vp to ſeede the ſame yeare, and then it is not accounted good. Many doe vſe to ſow Turneps on thoſe grounds from whence the ſame yeare they haue taken off Reddiſh and Lettice, to make the greater profit of the ground, by hauing two crops of increaſe in one yeare. The ſtalkes of Turneps will bend downe to the ground, as Carrots doe, but yet muſt not be bound or ordered in that manner, but ſuffered to grow without ſtaking or binding, ſo as they grow of ſome good diſtance in ſunder: when the ſeede beginneth to grow ripe, be very carefull to preſerue it from the birds, which will be moſt buſie to deuour them. You ſhall vnderſtand likewiſe that many doe account the beſt way to haue the faireſt and moſt principall ſeede from all theſe fore-recited herbes, that after they are ſowen, and riſen to a reaſonable growth, they be tranſplanted into freſh ground. Cabbages alſo are not only ſowen for the vſe of their heads to ſpend for meat, but to gather their ſeede likewiſe, which howſoeuer ſome haue endeauoured to doe, yet few haue gained good ſeede, becauſe our ſharpe hard froſtes in winter haue ſpoyled and rotted their ſtockes they preſerued for the purpoſe; but others haue found out a better and a more ſure way, which is, to take vp your ſtocks that are fitteſt to be preſerued, and bring them into the houſe, and there wrap them eyther in clothes, or other things to defend them from the cold, and hang them vp in a dry place, vntill the beginning of March following, then planting them in the ground, and a little defend them at the firſt with ſtraw caſt ouer them from the cold nights, thereby you may be ſure to haue perfect good ſeede, if your kinde be of the beſt: Sowe your ſeed in the moneths of February or March, and tranſplant them in May where they may ſtand to grow for your vſe, but be carefull to kill the wormes or Caterpillers that elſe will deuoure all your leaues, and be carefull alſo that none of the leaues bee broken in the planting, or otherwiſe rubbed, for that oftentimes hindereth the well cloſing of them. Leekes are

for

for the moſt part wholly nourſed vp from the ſeede that is here gathered; and becauſe there is not ſo much ſtore of them either ſowne or ſpent, as there is of Onions by the twentieth part, we are ſtill the more carefull to be prouided from our owne labours; yet there be diuers Gardiners in this Kingdome, that doe gather ſome ſmall quantity of Onion ſeede alſo for their owne or their priuate friends ſpending. The ſowing of them both is much about one time and manner, yet moſt vſually Leeks are ſowne later then Onions, and both before the end of March at the furtheſt; yet ſome ſowe Onions from the end of Iuly to the beginning of September, for their Winter prouiſion. Thoſe that are ſowne in the Spring, are to be taken vp and tranſplanted on a freſh bed prepared for the purpoſe, or elſe they will hardly abide a Winter; but hauing taken roote before Winter, they will beare good ſeede in the Summer following: You muſt ſtake both your Leekes and your Onion beds, and with poles laid a croſſe, binde your lopple headed ſtalkes vnto them, on high as well as belowe, or elſe the winde and their owne weight will beare them downe to the ground, and ſpoile your ſeede. You muſt thinne them, that is, pull vp continually after they are firſt ſprung vp thoſe that growe too thicke, as you doe with all the other herbes before ſpoken of, that they may haue the more roome to thriue. Of all theſe herbes and rootes before ſpoken of, you muſt take the likelieſt and faireſt to keepe for your ſeede; for if you ſhould not take the beſt, what hope of good ſeede can you expect? The time for the ſpending of theſe herbes and rootes, not particularly mentioned, is vntill they begin to runne vp for ſeede, or vntill they are to be tranſplanted for ſeede, or elſe vntill Winter, while they are good, as euery one ſhall ſee cauſe.

## Chap. IIII.

### *How to order Artichokes, Melons, Cowcumbers, and Pompions.*

THere are certaine other herbes to be ſpoken of, which are wholly nourſed vp for their fruit ſake, of whom I ſhall not need to ſay much, being they are ſo frequent in euery place. Artichokes being planted of faire and large ſlips, taken from the roote in September and October (yet not too late) will moſt of them beare fruit the next yeare, ſo that they be planted in well dunged ground, and the earth raiſed vp like vnto an Anthill round about each roote, to defend them the better from the extreame froſts in Winter. Others plant ſlips in March and Aprill, or ſooner, but although ſome of them will beare fruit the ſame yeare, yet all will not. And indeede many doe rather chooſe to plant in the ſpring then in the fall, for that oftentimes an extreame hard Winter following the new ſetting of ſlips, when they haue not taken ſufficient heart and roote in the ground, doth vtterly pierce and periſh them, when as they that are ſet in the Spring haue the whole Summers growth, to make them ſtrong before they feele any ſharpe froſts, which by that time they are the better able to beare. Muske Melons haue beene begun to bee nourſed vp but of late dayes in this Land, wherein although many haue tryed and endeauoured to bring them to perfection, yet few haue attained vnto it: but thoſe rules and orders which the beſt and skilfulleſt haue vſed, I will here ſet downe, that who ſo will, may haue as good and ripe Melons as any other in this Land. The firſt thing you are to looke vnto, is to prouide you a peece of ground fit for the purpoſe, which is either a ſloping or ſheluing banke, lying open and oppoſite to the South Sunne, or ſome other fit place not ſheluing, and this ground alſo you muſt ſo prepare, that all the art you can vſe about it to make it rich is little enough; and therefore you muſt raiſe it with meere ſtable ſoyle, thorough rotten & well turned vp, that it may be at the leaſt three foote deepe thereof, which you muſt caſt alſo into high beds or balkes, with deepe trenches or furrowes betweene, ſo as the ridges may be at the leaſt a foot and a halfe higher then the furrowes; for otherwiſe it is not poſſible to haue good Melons growe ripe. The choiſe of your ſeede alſo is another thing of eſpeciall regard, and the beſt is held to be Spaniſh, and not French, which hauing once gained, be ſure to haue ſtill of the ſame while they laſt

good,

good, that you may haue the ſeede of your owne ripe Melons from them that haue eaten them, or ſaue ſome of the beſt your ſelfe for the purpoſe. I ſay while they laſt good; for many are of opinion, that no ſeede of Muske Melons gathered in England, will endure good to ſowe againe here aboue the third yeare, but ſtill they muſt be renewed from whence you had your choiſeſt before. Then hauing prepared a hot bed of dung in Aprill, ſet your ſeedes therein to raiſe them vp, and couer them, and order them with as great care or greater then Cowcumbers, &c. are vſed, that when they are ready, they may be tranſplanted vpon the beds or balkes of that ground you had before prepared for them, and ſet them at the leaſt two yards in ſunder, euery one as it were in a hole, with a circle of dung about them, which vpon the ſetting being watered with water that hath ſtood in the Sunne a day or two, and ſo as often as neede is to water, couer them with ſtrawe (ſome vſe great hollow glaſſes like vnto bell heads) or ſome ſuch other things, to defend them both from the cold euenings or dayes, and the heate of the Sunne, while they are young and new planted. There are ſome that take vpon them great skill, that miſlike of the raiſing vp of Melons, as they doe alſo of Cowcumbers, on a hot bed of horſe dung, but will put two or three ſeedes in a place in the very ground where they ſhall ſtand and growe, and thinke without that former manner of forcing them forwards, that this their manner of planting will bring them on faſt and ſure enough, in that they will plucke away ſome of the worſt and weakeſt, if too many riſe vp together in a place; but let them know for certaine, that howſoeuer for Cowcumbers their purpoſe and order may doe reaſonable well, where the ground is rich and good, and where they ſtriue not to haue them ſo early, as they that vſe the other way, for Muske Melons, which are a more tender fruit, requiring greater care and trouble in the nourſing, and greater and ſtronger heate for the ripening, they muſt in our cold climate haue all the art vſed vnto them that may be, to bring them on the more early, and haue the more comfort of the Sunne to ripen them kindly, or elſe they will not bee worth the labour and ground. After you haue planted them as aforeſaid, ſome of good skill doe aduiſe, that you be carefull in any dry ſeaſon, to giue them water twice or thrice euery weeke while they are young, but more afterward when they are more growne, and that in the morning eſpecially, yea and when the fruit is growne ſomewhat great, to water the fruit it ſelfe with a watering pot in the heate of the day, is of ſo good effect, that it ripeneth them much faſter, and will giue them the better taſte and ſmell, as they ſay. To take likewiſe the fruit, and gather it at the full time of his ripeneſſe is no ſmall art; for if it be gathered before his due time to be preſently eaten, it will be hard and greene, and not eate kindly; and likewiſe if it be ſuffered too long, the whole goodneſſe will be loſt: You ſhall therefore know, that it is full time to gather them to ſpend preſently, when they begin to looke a little yellowiſh on the outſide, and doe ſmell full and ſtrong; but if you be to ſend them farre off, or keepe them long vpon any occaſion, you ſhall then gather them ſo much the earlier, that according to the time of the carriage and ſpending, they may ripen in the lying, being kept dry, and couered with woollen clothes: When you cut one to eate, you ſhall know it to be ripe and good, if the ſeede and pulpe about them in the middle be very wateriſh, and will eaſily be ſeparated from the meate, and likewiſe if the meate looke yellow, and be mellow, and not hard or greene, and taſte full and pleaſant, and not wateriſh: The vſuall manner to eate them is with pepper and ſalt, being pared and ſliced, and to drowne them in wine, for feare of doing more harme. Cowcumbers and Pompions, after they are nourſed vp in the bed of hot dung, are to be ſeuerally tranſplanted, each of them on a large plot of ground, a good diſtance in ſunder: but the Pompions more, becauſe their branches take vp a great deale more ground, & beſides, will require a great deale more watering, becauſe the fruit is greater. And thus haue you the ordering of thoſe fruits which are of much eſteeme, eſpecially the two former, with all the better ſort of perſons; and the third kinde is not wholly refuſed of any, although it ſerueth moſt vually for the meaner and poorer ſort of people, after the firſt early ripe are ſpent.

## CHAP. V.

### *The ordering of diuers ſorts of herbes for the pot, for meate, and for the table.*

TYme, Sauory, and Hyſſope, are vſually ſowne in the Spring on beds by themſelues, euerie one a part; but they that make a gaine by ſelling to others the young rootes, to ſet the knots or borders of Gardens, doe for the moſt part ſowe them in Iuly and Auguſt, that ſo being ſprung vp before Winter, they will be the fitter to be taken vp in the Spring following, to ſerue any mans vſe that would haue them. Sage, Lauender, and Roſemary, are altogether ſet in the Spring, by ſlipping the old ſtalkes, and taking the youngeſt and likelieſt of them, thruſting them either twined or otherwiſe halfe a foote deepe into the ground, and well watered vpon the ſetting; if any ſeaſonable weather doe follow, there is no doubt of their well thriuing: the hot Sunne and piercing drying Windes are the greateſt hinderances to them; and therefore I doe aduiſe none to ſet too ſoone in the Spring, nor yet in Autumne, as many doe practiſe: for I could neuer ſee ſuch come to good, for the extremity of the Winter comming vpon them ſo ſoone after their ſetting, will not ſuffer their young ſhootes to abide, not hauing taken ſufficient ſtrength in the ground, to maintain themſelues againſt ſuch violence, which doth often pierce the ſtrongeſt plants. Marierome and Baſſill are ſowne in the Spring, yet not too early; for they are tender plants, and doe not ſpring vntill the weather bee ſomewhat warme: but Baſſill would bee ſowne dry, and not haue any water of two or three daies after the ſowing, elſe the ſeede will turne to a gelly in the ground. Some vſe to ſowe the ſeed of Roſemary, but it ſeldome abideth the firſt Winter, becauſe the young plants being ſmall, and not of ſufficient ſtrength, cannot abide the ſharpneſſe of ſome Winters, notwithſtanding the couering of them, which killeth many old plants; but the vſuall way is to ſlippe and ſet, and ſo they thriue well. Many doe vſe to ſowe all or the moſt ſorts of Pot-herbes together on one plot of ground, that they neede not to goe farre to gather all the ſorts they would vſe. There are many ſorts of them well knowne vnto all, yet few or none doe vſe all ſorts, but as euery one liketh; ſome vſe thoſe that others refuſe, and ſome eſteem thoſe not to bee wholeſome and of a good reliſh, which others make no ſcruple of. The names of them are as followeth, and a ſhort relation of their ſowing or planting.

Roſemary, Tyme, and Sauorie are ſpoken of before, and Onions and Leekes.

Mints are to bee ſet with their rootes in ſome by-place, for that their rootes doe creepe ſo farre vnder ground, that they quickly fill vp the places neare adioyning, if they be not puld vp.

Clarie is to be ſowne, and ſeedeth and dyeth the next yeare, the herbe is ſtrong, and therefore a little thereof is ſufficient.

Nep is ſowne, and dyeth often after ſeeding, few doe vſe it, and that but a little at a time: both it and Clarie are more vſed in Tanſies then in Broths.

Coſtmarie is to be ſet of rootes, the leaues are vſed with ſome in their Broths, but with more in their Ale.

Pot Marierome is ſet of rootes, being ſeparated in ſunder.

Penniroyall is to be ſet of the ſmall heads that haue rootes, it creepeth and ſpreadeth quickly.

Alliſanders are to be ſowne of ſeede, the tops of the rootes with the greene leaues are vſed in Lent eſpecially.

Parſley is a common herbe, and is ſowne of ſeede, it ſeedeth the next yeare and dyeth: the rootes are more vſed in broths then the leaues, and the leaues almoſt with all ſorts of meates.

Fennell is ſowne of ſeede, and abideth many yeares yeelding ſeede: the rootes alſo are vſed in broths, and the leaues more ſeldome, yet ſerue to trimme vp many fiſh meates.

Borage is ſowne of ſeede, and dyeth the next yeare after, yet once being ſuffered to ſeede in a Garden, will ſtill come of it owne ſhedding.

Bugloſſe commeth of ſeede, but abideth many yeares after it hath giuen ſeede, if it ſtand not in the coldeſt place of the Garden.

Marigolds are ſowne of ſeede, and may be after tranſplanted, they abide two or three yeares, if they be not ſet in too cold a place: the leaues and flowers are both vſed.

Langedebeefe is ſowne of ſeede, which ſhedding it ſelfe will hardly be deſtroyed in a Garden.

Arrach is to be ſowne of ſeede, this likewiſe will riſe euery yeare of it owne ſeed, if it be ſuffered to ſhed it ſelfe.

Beetes are ſowne of ſeede, and abideth ſome yeares after, ſtill giuing ſeede.

Blites are vſed but in ſome places; for there is a generall opinion held of them, that they are naught for the eyes: they are ſowne euery yeare of ſeede.

Bloodwort once ſowne abideth many yeares, if the extremity of the froſts kill it not, and ſeedeth plentifully.

Patience is of the ſame nature, and vſed in the ſame manner.

French Mallowes are to be ſowne of ſeede, and will come of it owne ſowing, if it be ſuffered to ſhed it ſelfe.

Ciues are planted onely by parting the rootes; for it neuer giueth any ſeede at all.

Garlicke is ordered in the ſame manner, by parting and planting the rootes euerie yeare.

Theſe be all the ſorts are vſed with vs for that purpoſe, whereas I ſaid before, none vſeth all, but euery one will vſe thoſe they like beſt: and ſo much ſhall ſuffice for pot-herbes.

## Chap. VI.

### *The manner and ordering of many ſorts of herbes and rootes for Sallets.*

IF I ſhould ſet downe all the ſorts of herbes that are vſually gathered for Sallets, I ſhould not onely ſpeake of Garden herbes, but of many herbes, &c. that growe wilde in the fields, or elſe be but weedes in a Garden; for the vſuall manner with many, is to take the young buds and leaues of euery thing almoſt that groweth, as well in the Garden as in the Fields, and put them all together, that the taſte of the one may amend the relliſh of the other: But I will only ſhew you thoſe that are ſown or planted in gardens for that purpoſe. Aſparagus is a principall & delectable Sallet herbe, whoſe young ſhootes when they are a good handfull high aboue the ground, are cut an inch within the ground, which being boyled, are eaten with a little vinegar and butter, as a Sallet of great delight. Their ordering with the beſt Gardiners is on this wiſe: When you haue prouided ſeede of the beſt kinde, you muſt ſowe it either before Chriſtmas, as moſt doe, or before the end of February; the later you ſowe, the later and the more hardly will they ſpring: after they are growne vp, they are to be tranſplanted in Autumne on a bed well trenched in with dung; for elſe they will not bee worth your labour, and ſet about a foote diſtance in ſunder, and looke that the more carefull you are in the replanting of them, the better they will thriue, and the ſooner growe great: after fiue or ſix yeares ſtanding they vſually doe decay; and therefore they that ſtriue to haue continually faire and great heads, doe from ſeede raiſe vp young for their ſtore. You muſt likewiſe ſee that you cut not your heads or young ſhoote too nigh, or too much, that is, to take away too many heads from a roote, but to leaue a ſufficient number vncut, otherwiſe it will kill the heart of your rootes the ſooner, cauſing them to dye, or to giue very ſmall heads or ſhootes; for you may well conſider with your ſelfe, that if the roote haue not head enough left it aboue the ground to ſhoote greene this yeare, it will not, nor cannot proſper vnder ground to giue encreaſe the next yeare. The ordering of Lettice I haue ſpoken of before, and ſhall not neede here to repeate what hath beene already ſaid, but referre you thereunto for the ſowing, planting, &c. onely I will here ſhew you the manner of ordering them for Sallets. There are ſome ſorts of Lettice that growe very great, and cloſe their heads, which are called Cab-

bage

bage Lettice, both ordinary and extraordinary, and there are other ſorts of great Lettice that are open, and cloſe not, or cabbage not at all, which yet are of an excellent kinde, if they be vſed after that eſpeciall manner is fit for them, which is, That when they are planted (for after they are ſowne, they muſt be tranſplanted) of a reaſonable diſtance in ſunder, and growne to be of ſome bigneſſe, euery one of them muſt bee tyed together with baſt or thread toward the toppes of the leaues, that by this meanes all the inner leaues may growe whitiſh, which then are to be cut vp and vſed: for the keeping of the leaues cloſe doth make them taſte delicately, and to bee very tender. And theſe ſorts of Lettice for the moſt part are ſpent after Summer is paſt, when other Lettice are not to be had. Lambes Lettice or Corne Sallet is an herbe, which abiding all Winter, is the firſt Sallet herbe of the yeare that is vſed before any ordinarie Lettice is ready; it is therefore vſually ſowne in Auguſt, when the ſeede thereof is ripe. Purſlane is a Summer Sallet herbe, and is to be ſowne in the Spring, yet ſomewhat late, becauſe it is tender, and ioyeth in warmth; and therefore diuers haue ſowne it vpon thoſe beddes of dung, whereon they nourſed vp their Cowcumbers, &c. after they are taken away, which being well and often watered, hath yelded Sallet vntill the end of the yeare. Spinach is ſowne in the Spring, of all for the moſt part that vſe it, but yet if it be ſowne in Summer it will abide greene all the Winter, and then ſeedeth quickly: it is a Sallet that hath little or no taſte at all therein, like as Lettice and Purſlane, and therefore Cookes know how to make many a good diſh of meate with it, by putting Sugar and Spice thereto. Coleworts are of diuers kinds, and although ſome of them are wholly ſpent among the poorer ſort of people, yet ſome kindes of them may be dreſſed and ordered as may delight a curious palate, which is, that being boyled tender, the middle ribs are taken cold, and laid in diſhes, and vinegar and oyle poured thereon, and ſo eaten. Coleflowers are to be had in this Country but very ſeldome, for that it is hard to meete with good ſeede: it muſt bee ſowne on beds of dung to force it forward, or elſe it would periſh with the froſt before it had giuen his head of flowers, and tranſplanted into verie good and rich ground, leſt you loſe the benefit of your labours. Endiue is of two ſorts, the ordinary, and another that hath the edges of the leaues curld or crumpled; it is to be whited, to make it the more dainty Sallet, which is vſually done in this manner: After they are grown to ſome reaſonable greatneſſe (but in any caſe before they ſhoote forth a ſtalke in the midſt for ſeede) they are to be taken vp, and the rootes being cut away, lay them to dry or wither for three or foure houres, and then bury them in ſand, ſo as none of them lye one vpon another, or if you can, one to touch another, which by this meanes will change whitiſh, and thereby become verie tender, aud is a Sallet both for Autumne and Winter. Succorie is vſed by ſome in the ſame manner, but becauſe it is more bitter then Endiue, it is not ſo generally vſed, or rather vſed but of a verie few: and whereas Endiue will ſeede the ſame yeare it is ſowne, and then dye, Succorie abideth manie yeares, the bitterneſſe thereof cauſing it to be more Phyſicall to open obſtructions; and therefore the flowers pickled vp, as diuers other flowers are vſed to be now adaies, make a delicate Sallet at all times when there is occaſion to vſe them. Of red Beetes, the rootes are onely vſed both boyled and eaten cold with vinegar and oyle, and is alſo vſed to trimme vp or garniſh forth manie ſorts of diſhes of meate: the ſeede of the beſt kinde will not abide good with vs aboue three yeares, but will degenerate and growe worſe; and therefore thoſe that delight therein muſt be curious, to be prouided from beyond Sea, that they may haue ſuch as will giue delight. Sorrell is an herbe ſo common, and the vſe ſo well knowne, both for ſawce, and to ſeaſon broths and meates for the ſound as well as ſicke perſons, that I ſhall not neede to ſay anie more thereof. Cheruill is a Sallet herbe of much vſe, both with French and Dutch, who doe much more delight in herbes of ſtronger taſte then the Engliſh doe: it is ſowne early, and vſed but a while, becauſe it quickly runneth vp to ſeede. Sweete Cheruill, or as ſome call it, Sweete Cis, is ſo like in taſte vnto Aniſe ſeede, that it much delighteth the taſte among other herbes in a Sallet: the ſeede is long, thicke, blacke, and cornered, and muſt be ſowne in the end of Autumne, that it may lye in the ground all the Winter, and then it will ſhoote out in the Spring, or elſe if it be ſowne in the Spring, it will not ſpring vp that yeare vntill the next: the leaues (as I ſaid before) are vſed among other herbes: the rootes likewiſe are not onely cordiall, but alſo held to be preſeruatiue againſt the Plague, either greene, dryed, or preſerued

with ſugar. Rampion roores are a kinde of Sallet with a great many, being boyled tender, and eaten cold with vinegar and pepper. Creſſes is an herbe of eaſie and quick growth, and while it is young eaten eyther alone, or with parſley and other herbes: it is of a ſtrong taſte to them that are not accuſtomed thereunto, but it is much vſed of ſtrangers. Rocket is of the ſame nature and qualitie, but ſomewhat ſtronger in taſte: they are both ſowen in the Spring, and riſe, ſeede and dye the ſame yeare. Tarragon is an herbe of as ſtrong a taſte as eyther Rocket or Creſſes, it abideth and dyeth not euery yeare, nor yet giueth ripe ſeede (as far as euer could bee found with vs) any yeare, but maketh ſufficient increaſe within the ground, ſpreading his roots all abroad a great way off. Muſtard is a common ſawce both with fiſh and fleſh, and the ſeed thereof (and no part of the plant beſides) is well knowne how to be vſed being grownded, as euery one I thinke knoweth. The roores of horſe Radiſh likewiſe beeing grownd like Muſtard, is vſed both of ſtrangers and our owne nation, as ſawce for fiſh. Tanſie is of great vſe, almoſt with all manner of perſons in the Spring of the yeare: it is more vſually planted of the roores then otherwiſe; for in that the roores ſpread far and neere they may be eaſily taken away, without any hurt to the reſt of the roores. Burnet, although it be more vſed in wine in the Summer time then any way elſe, yet it is likewiſe made a ſallet herbe with many, to amend the harſh or weak relliſh of ſome other herbs. Skirrets are better to be ſowen of the ſeed then planted from the roots, and will come on more ſpeedily, and be fairer roores: they are as often eaten cold as a Sallet, being boyled and the pith taken out, as ſtewed with butter and eaten warme. Let not Parſley and Fenell be forgotten among your other Sallet herbes, wherof I haue ſpoken before, and therefore need ſay no more of them. The flowers of Marigolds pickt cleane from the heads, and pickled vp againſt winter, make an excellent Sallet when no flowers are to be had in a garden. Cloue Gilloflowers likewiſe preſerued or pickled vp in the ſame manner (which is *ſtratum ſuper ſtratum*, a lay of flowers, and then ſtrawed ouer with fine dry and poudered Sugar, and ſo lay after lay ſtrawed ouer, vntill the pot bee full you meane to keepe them in, and after filled vp or couered ouer with vinegar) make a Sallet now adayes in the higheſt eſteeme with Gentles and Ladies of the greateſt note: the planting and ordering of them both is ſpoken of ſeuerally in their proper places. Goates bearbe that groweth in Gardens only, as well as that which groweth wilde in Medowes, &c. bearing a yellow flower, are vſed as a Sallet, the roores beeing boyled and pared are eaten cold with vinegar, oyle and pepper; or elſe ſtewed with butter and eaten warme as Skirrets, Parſneps &c. And thus haue you here ſet downe all thoſe moſt vſuall Sallets are vſed in this Kingdome: I ſay the moſt vſuall, or that are nourſed vp in Gardens; for I know there are ſome other wilde herbes and roores, as Dandelion &c. but they are vſed onely of ſtrangers, and of thoſe whoſe curioſitie ſearcheth out the whole worke of nature to ſatisfie their deſires.

## Chap. VII.

### *Of diuers Phyſicall herbes fit to be planted in Gardens, to ſerue for the eſpeciall vſes of a familie.*

HAuing thus ſhewed you all the herbes that are moſt vſually planted in Kitchen Gardens for ordinary vſes, let mee alſo adde a few other that are alſo nourſed vp by many in their Gardens, to preſerue health, and helpe to cure ſuch ſmall diſeaſes as are often within the compaſſe of the Gentlewomens skils; who, to helpe their owne family, and their poore neighbours that are farre remote from Phyſitians and Chirurgions, take much paines both to doe good vnto them, and to plant thoſe herbes that are conducing to their deſires. And although I doe recite ſome that are mentioned in other places, yet I thought it meete to remember them altogether in one place. Angelica, the garden kinde, is ſo good an herbe, that there is no part thereof but is of much vſe, and all cordiall and preſeruatiue from infectious or contagious diſeaſes, whether you will diſtill the water of the herbe, or preſerue or candie the roores or the greene ſtalkes, or vſe the ſeede in pouder or in diſtillations, or decoctions with other things: it is ſowen of ſeede, and will abide vntill

it giue ſeede, and then dyeth. Rue or Herbe grace is a ſtrong herbe, yet vſed inwardly againſt the plague as an Antidote with Figs and Wall-nuts, and helpeth much againſt windy bodies: outwardly it is vſed to bee layde to the wreſtes of the hands, to driue away agues: it is more vſually planted of ſlips then raiſed from ſeede, and abideth long if ſharpe froſtes kill it not. Dragons being diſtilled are held to be good to expell any euill thing from the heart: they are altogether planted of the rootes. Setwall, Valerian, or Capons tayle, the herbe often, but the roote much better, is vſed to prouoke ſweating, thereby to expell euill vapours that might annoy the heart: it is only planted of the rootes when they are taken vp, and the young replanted. Aſarabacca, the leaues are often vſed to procure vomiting being ſtamped, and the ſtrained iuice to a little quantitie, put into a draught of ale and drunke, thereby to eaſe the ſtomacke of many euill and groſſe humours that there lye and offend it; diuers alſo take the leaues and rootes a little boyled in wine, with a little ſpice added thereunto, to expell both tertian and quartan agues: the rootes of our Engliſh growing is more auaileable for theſe purpoſes then any outlandiſh: it is planted by the roote; for I could neuer ſee it ſpring of ſeede. Maſterwort commeth ſomewhat neere in propertie vnto Angelica, and beſides very effectuall to diſperſe winde in the bodie, whether of the collicke or otherwiſe; as alſo very profitable to comfort in all cold cauſes: it yeeldeth ſeede, but yet is more vſually planted from the rootes being parted. Balme is a cordiall herbe both in ſmell and taſte, and is wholly vſed for thoſe purpoſes, that is, to comfort the heart being diſtilled into water either ſimple or compound, or the herbe dryed and vſed: it is ſet of the rootes being parted, becauſe it giueth no ſeede that euer I could obſerue. Camomill is a common herbe well knowne, and is planted of the rootes in alleyes, in walkes, and on bankes to ſit on, for that the more it is troden on, and preſſed downe in dry weather, the cloſer it groweth, and the better it will thriue: the vſe thereof is very much, both to warme and comfort, and to eaſe paines being applyed outwardly after many faſhions: the decoction alſo of the flowers prouoketh ſweat, and they are much vſed againſt agues. Featherfew is an herbe of greater vſe for women then for men, to diſſolue flatulent or windy humours, which cauſeth the paines of the mother: ſome vſe to take the iuice thereof in drinke for agues: it is as well ſowen of the ſeede as planted of the rootes. Coſtmary is vſed among thoſe herbes that are put in ale to cauſe it haue a good relliſh, and to be ſomewhat phyſicall in the moneth of May, and doth helpe to prouoke vrine: it is ſet of the rootes being parted. Maudlin is held to be a principall good herbe to open and cleanſe the liuer, and for that purpoſe is vſed many wayes, as in ale, in tanſies, and in broths &c. the ſeed alſo is vſed, and ſo is the herbe alſo ſometimes, to kill the wormes in children: it is ſowen of the ſeede, and planted alſo of the ſeparated rootes. Caſſidonie is a ſmall kinde of Lauender, but differing both in forme and qualitie: it is much vſed for the head to eaſe paines thereof, as alſo put among other things to purge melancholicke diſeaſes: it is ſowen of ſeede, and abideth not a winter vnleſſe it bee well defended, and yet hardly giueth ripe ſeede againe with vs. Smallage is a great opening herbe, and much more then eyther Parſley or Fenell, and the rootes of them all are often vſed together in medicines: it is ſowen of ſeede, and will not bee wanting in a Garden if once you ſuffer it to ſow it ſelfe. Cardus Benedictus, or the Bleſſed Thiſtle, is much vſed in the time of any infection or plague, as alſo to expell any euill ſymptome from the heart at all other times. It is vſed likewiſe to be boyled in poſſet drink, & giuen to them that haue an ague, to help to cure it by ſweating or otherwiſe. It is vſually ſowen of ſeed, and dyeth when it hath giuen ſeed. Winter Cherries are likewiſe nurſed vp in diuers gardens, for that their propertie is to giue helpe to them that are troubled eyther with the ſtopping or heate of their vrine: the herbe and berries are often diſtilled, but the berries alone are more often vſed: after it is once planted in a garden it will runne vnder ground, & abide well enough. Celondine is held to bee good for the iaundiſe, it is much vſed for to cleere dim eyes, eyther the iuice or the water dropped into them: it is ſowen of ſeede, and being once brought into a garden, will hardly be weeded out; the ſeede that ſheddeth will ſo ſow it ſelfe, and therefore ſome corner in a garden is the fitteſt place for it. Tabacco is of two ſorts, and both vſed to be planted in Gardens, yet the Engliſh kinde (as it is called) is more to be found in our Countrey Gardens then the Indian ſort: the leaues of both ſorts indifferently, that is, of eyther of which is next at hand, being ſtamped and boy-

led eyther by it selfe, or with other herbes in oyle or hogs suet, doe make an excellent salue for greene wounds, and also to clense old vlcers or sores; the iuice of the greene leaues drunke in ale, or a dryed leafe steeped in wine or ale for a night, and the wine or ale drunke in the morning, prouoketh to cast, but the dryed leafe much stronger then the greene: they are sowen of seede, but the Indian kinde is more tender, and will not abide a winter with vs abroade. Spurge that vsually groweth in Gardens, is a violent purger, and therefore it is needfull to be very carefull how it is vsed: the seede is more ordinarily vsed then any other part of the plant, which purgeth by vomiting in some, and both vpwards and downwards in many; the iuice of the herbe, but especially the milke thereof, is vsed to kill wartes: it is sowen of seede, and when it doth once shed it selfe, it will still continue springing of the fallen seede. Bearefoote is sowen of seed, and will hardly abide transplanting vnlesse it bee while it is young; yet abideth diuers yeares, if it stand not in too cold a place. This I speake of the greater kinde; for the lower small wilde kind (which is the most ordinary in this land) will neuer decay: the leaues are sometimes vsed greene, but most vsually dryed and poudered, and giuen in drinke to them that haue the wormes: it purgeth melancholy, but especially the roots. In many Countries of this Land, and elsewhere, they vse to thrust the stalk of the great kinde through the eare or dewlap of Kine and Cattell, to cure them of many diseases. Salomons Seale, or (as some call it) Ladder to heauen, although it doth grow wilde in many places of this Land, yet is planted in Gardens: it is accounted an excellent wound herbe to consolidate, and binde, insomuch that many vse it with good successe to cure ruptures, and to stay both the white and the red fluxe in women: it is planted altogether of the rootes, for I could neuer finde it spring from the seede, it is so strong. Comfry likewise is found growing wilde in many places by ditch sides, and in moist places, and therefore requireth some moist places of the garden: it is wholly vsed for knitting, binding, and consolidating fluxes and wounds, to be applyed either inwardly or outwardly: The rootes are stronger for those purposes then any other parts of the plant. Licoris is much vsed now adaies to bee planted in great quantitie, euen to fill many acres of ground, whereof riseth a great deale of profit to those that know how to order it, and haue fit grounds for it to thriue in; for euery ground will not be aduantagious: It will require a very rich, deepe and mellow ground, eyther naturall or artificiall; but for a priuate house where a small quantitie will serue, there needeth not so much curiositie: it is vsually planted of the top heads, when the lower rootes (which are the Licoris that is vsed) and the runners are cut from them. Some vse to make an ordinary drinke or beuerage of Licoris, boyled in water as our vsuall ale or beere is with malt, which fermented with barme in the same manner, and tunned vp, serueth in stead thereof, as I am credibly informed: It is otherwise in a manner wholly spent for colds, coughes and rheumes, to expectorate flegme, but vsed in diuers formes, as in iuice, in decoctions, syrrups, roules, trochisces, and the greene or dryed roote of it selfe.

And these are the most ordinary Physicall herbes that are vsed to be planted in gardens for the vse of any Country familie, that is (as I said before) farre remote from Physitians or Chirurgions abidings, that they may vse as occasion serueth for themselues or their neighbours, and by a little care and paines in the applying may doe a great deale of good, and sometimes to them that haue not wherewith to spend on themselues, much lesse on Physitians or Chirurgions, or if they haue, may oftentimes receiue lesse good at their hands then at others that are taught by experience in their owne families, to be the more able to giue helpe to others.

The

# THE KITCHEN GARDEN.

## THE SECOND PART,

CHAP. I.

Ontaining as well all sorts of herbes, as rootes and fruits, that are vsually planted in Gardens, to serue for the vse of the Table whether of the poore or rich of our Countrey: but herein I intend not to bring any fruite bearing trees, shrubbes, or bushes; for I reserue them for my Orchard, wherin they shal be set forth. So that in these three parts, I suppose the exquisite ornament of any worthy house is consummate for the exteriour bounds, the benefit of their riches extending also to the furnishing of the most worthy inward parts thereof: but because many take pleasure in the sight and knowledge of other herbes that are Physicall, and much more in their properties and vertues, if vnto these three I should adde a Physicke Garden, or Garden of Simples, there would be a quadripartite complement, of whatsoeuer arte or nature, necessitie or delight could affect: which to effect (as many my friends haue intreated it at my hands) will require more paines and time then all this worke together: yet to satisfie their desires and all others herein, that would bee enformed in the truth, and reformed of the many errours and slips set forth and published heretofore of plants by diuers, I shall (God assisting and granting life) labour to performe, that it may shew it selfe to the light in due conueniencie, if these bee well and gratefully accepted. And because I ended with some sweete herbes in the former part, I will in this part beginne with the rest, which I reserued for this place, as fitter for the pot and kitchen then for the hand or bosome, and so descend to other herbes that are for meat or sallets: and after them to those rootes that are to be eaten, as meate or as sallets: and lastly the fruits that grow neere, or vpon the ground, or not much aboue it; as the Artichoke, &c. in which I make a shorter description then I did in the former, rather endeauouring to shew what they are, and whereunto they are vsed, then the whole varietie or any exact declaration: which methode, although in some sort it may bee fitting for this purpose, yet it is not for an history or herball: I shall therefore require their good acceptance for whose sake I doe it, not doubting, but that I, or others, if they write againe of this subiect, may polish and amende what formerly hath beene eyther mis set, or not so thoroughly expressed, besides some additions of new conceits; seeing I treade out a new path, and therefore those that follow may the easilier see the Meanders, and so goe on in a direct line.

## Chap. I.

*Maiorana latifolia, sive maior Anglica.* Winter, or pot Marierome.

WInter Marierome is a small bushie herbe like vnto sweete Marierome, being parted or diuided into many branches, whereon doe grow broader and greener leaues, set by couples, with some small leaues likewise at the seuerall ioynts all along the branches: at the tops whereof grow a number of small purplish white flowers set together in a tuft, which turne into small and round seed, bigger then sweet Marierome seede: the whole plant is of a small and fine sent, but much inferiour to the other, and is nothing so bitter as the sweete Marierome, and thereby both the fitter and more willingly vsed for meates: the roote is white and threddy, and perisheth not as the former, but abideth many yeares.

### The Vse of winter Marierome.

The vse of this Marierome is more frequent in our Land then in others, being put among other pot-herbes and farsing (or faseting herbes as they are called) and may to good profit bee applyed in inward as well as outward griefes for to comfort the parts, although weaker in effect then sweete Marieromes.

## Chap. II.

*Thymum vulgatius sive durius.* Ordinary Garden Tyme.

THe ordinary Garden Tyme is a small low wooddy plant with brittle branches, and small hard greene leaues, as euery one knoweth, hauing small white purplish flowers, standing round about the tops of the stalkes: the seed is small and browne, darker then Marierome seed: the root is woody, and abideth well diuers Winters.

*Thymum latifolium.* Masticke Tyme.

This Tyme hath neyther so wooddy branches, nor so hard leaues, but groweth lower, more spreading, and with somewhat broader leaues: the flowers are of a purplish white colour, standing in roundles round about the stalkes, at the ioynts with leaues at them likewise. This Tyme endureth better and longer then the former, and by spreading it selfe more then the former, is the more apt to bee propagated by slipping, because it hath beene seldome seene to giue seede: It is not so quicke in sent or taste as the former, but is fitter to set any border or knot in a garden, and is for the most part wholly employed to such vses.

### The Vse of Tyme.

To set downe all the particular vses whereunto Tyme is applyed, were to weary both the Writer and Reader; I will but only note out a few: for besides the physicall vses to many purposes, for the head, stomacke, spleene, &c. there is no herbe almost of more vse, in the houses both of high and low, rich and poore, both for inward and outward occasions; outwardly for bathings among other hot herbes, and among other sweete herbes for strewings: inwardly in most sorts of broths, with Rosmary, as also with other faseting (or rather farsing) herbes, and to make sawce for diuers sorts both fish and flesh, as to stuffe the belly of a Goose to bee rosted, and after put into the sawce, and the pouder with breade to strew on meate when it

is

1 *Maiorana maior Anglica.* Pot Marierome. 2 *Thymum vulgatius.* Garden Tyme. 3 *Satureia.* Sauorie. 4 *Hyssopus.* Hyssope. 5 *Pulegium.* Penniroyall. 6 *Salvia maior.* Common Sage. 7 *Salvia minor primata.* Sage of vertue.

is roſted, and ſo likewiſe on roſted or fryed fiſh. It is held by diuers to bee a ſpeedy remedy againſt the ſting of a Bee, being bruiſed and layd thereon.

## Chap. III.

### *Satureia ſiue Thymbra.* Sauorie.

THere are two ſorts of of Sauory, the one called Summer, and the other Winter Sauorie: The Summer Sauory is a ſmall tender herbe, growing not aboue a foote and a halfe high, or thereabouts, riſing vp with diuers brittle branches, ſlenderly or ſparſedly ſet with ſmall long leaues, ſoft in handling, at euery ioynt a couple, one againſt another, of a pleaſant ſtrong and quicke ſent and taſte: the flowers are ſmall and purpliſh, growing at the toppes of the ſtalkes, with two ſmall long leaues at the ioynts vnder them: the ſeede is ſmall, and of a darke colour, bigger then Tyme ſeede by the halfe: the roote is wooddy, and hath many ſtrings, periſhing euery yeare wholly, and muſt bee new ſowen againe, if any will haue it.

The Winter Sauorie is a ſmall low buſhie herbe, very like vnto Hyſſope, but not aboue a foote high, with diuers ſmall hard branches, and hard darke green leaues thereon, thicker ſet together then the former by much, and as thicke as common Hyſſope, ſometimes with foure leaues or more at a ioynt, of a reaſonable ſtrong ſent, yet not ſo ſtrong or quicke as the former: the flowers are of a pale purpliſh colour, ſet at ſeuerall diſtances at the toppes of the ſtalkes, with leaues at the ioynts alſo with them, like the former: the roote is woody, with diuers ſmall ſtrings thereat, and abideth all the winter with his greene leaues: it is more vſually encreaſed by ſlipping or diuiding the roote, and new ſetting it ſeuerally againe in the Spring, then by ſowing the ſeed.

### The Vſe of Sauorie.

The Summer Sauorie is vſed in other Countryes much more then with vs in their ordinary diets, as condiment or ſawce to their meates, ſometimes of it ſelfe, and ſometimes with other herbes, and ſometimes ſtrewed or layde vpon the diſhes as we doe Parſley, as alſo with beanes and peaſe, riſe and wheate; and ſometimes the dryed herbe boyled among peaſe to make pottage.

The Winter Sauorie is one of the (farſing) faſeting herbes as they call them, and ſo is the Summer Sauorie alſo ſometimes. This is vſed alſo in the ſame manner that the Summer Sauorie is, ſet downe before, and to the ſame purpoſes: as alſo to put into puddings, ſawſages, and ſuch like kindes of meates. Some doe vſe the ponder of the herbe dryed (as I ſayd before of Tyme) to mixe with grated bread, to breade their meate, be it fiſh or fleſh, to giue it the quicker relliſh. They are both effectuall to expell winde.

## Chap. IIII.

### *Hyſſopus.* Hyſſope.

GArden Hyſſope is ſo well knowne to all that haue beene in a Garden, that I ſhall but *actum agere*, to beſtow any time thereon, being a ſmall buſhie plant, not riſing aboue two foote high, with many branches, woody below, and tender aboue, whereon are ſet at certaine diſtances, ſundry ſmall, long and narrow greene leaues: at the toppe of euery ſtalke ſtand blewiſh purple gaping flowers, one aboue another in a long ſpike or eare: after which followeth the ſeede, which is ſmall and blackiſh: the rootes are compoſed of many threddy ſtrings, the whole plant is of a ſtrong ſweet ſent.

The

### The Vſe of Hyſſope.

Hyſſope is much vſed in Ptiſans and other drinkes, to help to expectorate flegme. It is many Countrey peoples medicine for a cut or greene wound, being bruiſed with ſugar and applyed. I finde it is alſo much commended againſt the falling ſickeneſſe, eſpecially being made into pils after the manner before rehearſed. It is accounted a ſpeciall remedy againſt the ſting or biting of an Adder, if the place be rubbed with Hyſſope, bruiſed and mixed with honey, ſalt and cummin ſeede. A decoction thereof with oyle, and annointed, taketh away the itching and tingling of the head, and vermine alſo breeding therein. An oyle made of the herbe and flowers, being annointed, doth comfort benummed ſinewes and ioynts.

## Chap. V.

*Pulegium.* Pennyroyall.

PEnnyroyall alſo is an herbe ſo well knowne, that I ſhall not neede to ſpend much time in the deſcription of it: hauing many weake round ſtalkes, diuided into ſundry branches, rather leaning or lying vpon the ground then ſtanding vpright; whereon are ſet at ſeuerall ioynts, ſmall roundiſh darke greene leaues: the flowers are purpliſh that grow in gardens, yet ſome that grow wilde are white, or more white then purple, ſet in roundles about the tops of the branches; the ſtalkes ſhoote forth ſmall fibres or rootes at the ioynts, as it lyeth vpon the ground, thereby faſtening it ſelfe therein, and quickly increaſeth, and ouer-runneth any ground, eſpecially in the ſhade or any moiſt place, and is replanted by breaking the ſprouted ſtalkes, and ſo quickely groweth.

Other ſorts of Pennyroyall are fit for the Phyſicke Garden, or Garden of Simples.

### The Vſe of Pennyroyall.

It is very good and wholeſome for the lunges, to expell cold thin flegme, and afterwards to warme and dry it vp: and is alſo of the like propertie as Mintes, to comfort the ſtomacke, and ſtay vomiting. It is alſo vſed in womens baths and waſhings: and in mens alſo to comfort the ſinewes. It is yet to this day, as it hath beene in former times, vſed to bee put into puddings, and ſuch like meates of all ſorts, and therefore in diuers places they know it by no other name then Pudding-graſſe.

The former age of our great Grandfathers, had all theſe hot herbes in much and familiar vſe, both for their meates and medicines, and therewith preſerued themſelues in long life and much health: but this delicate age of ours, which is not pleaſed with any thing almoſt, be it meat or medicine, that is not pleaſant to the palate, doth wholly refuſe theſe almoſt, and therefore cannot be partaker of the benefit of them.

## Chap. VI.

*Salvia.* Sage.

THere are two eſpeciall kindes of Sage nourſed vp in our Gardens, for our ordinary vſe, whereof I intend to write in this place, leauing the reſt to his fitter place. Our ordinary Sage is reckoned to bee of two ſorts, white and red, both

both of them bearing many foure ſquare wooddy ſtalkes, in ſome whiter, in others redder, as the leaues are alſo, ſtanding by couples at the ioynts, being long, rough, and wrinkled, of a ſtrong ſweete ſent: at the tops of the ſtalkes come forth the flowers, ſet at certaine ſpaces one aboue another, which are long and gaping, like vnto the flowers of Clary, or dead Nettles, but of a blewiſh purple colour; after which come ſmall round ſeede in the huske that bore the flower: the roote is wooddy, with diuers ſtrings at it: It is more vſually planted of the ſlips, pricked in the Spring time into the ground, then of the ſeed.

*Saluia minor ſiue pinnata.* Small Sage or Sage of vertue.

The leſſer Sage is in all things like vnto the former white Sage, but that his branches are long and ſlender, and the leaues much ſmaller, hauing for the moſt part at the bottome of each ſide of the leafe a peece of a leafe, which maketh it ſhew like finns or eares: the flowers alſo are of a blewiſh purple colour, but leſſer. Of this kinde there is one that beareth white flowers.

## The Vſe of Sage.

Sage is much vſed of many in the moneth of May faſting, with butter and Parſley, and is held of moſt much to conduce to the health of mans body.

It is alſo much vſed among other good herbes to bee tund vp with Ale, which thereupon is termed Sage Ale, whereof many barrels full are made, and drunke in the ſaid moneth chiefly for the purpoſe afore recited: and alſo for teeming women, to helpe them the better forward in their childebearing, if there be feare of abortion or miſcarrying.

It is alſo vſed to be boyled among other herbes, to make Gargles or waters to waſh ſore mouths and throates: As alſo among other herbes, that ſerue as bathings, to waſh mens legs or bodies in the Summer time, to comfort nature, and warme and ſtrengthen aged cold ſinewes, and lengthen the ſtrength of the younger.

The Kitchen vſe is either to boyle it with a Calues head, and being minced, to be put with the braines, vinegar and pepper, to ſerue as an ordinary ſawce thereunto: Or being beaten and iuyced (rather then minced as manie doe) is put to a roſted Pigges braines, with Currans for ſawce thereunto. It is in ſmall quantity (in regard of the ſtrong taſte thereof) put among other faſting herbes, to ſerue as ſawce for peeces of Veale, when they are farſed or ſtuffed therewith, and roſted, which they call Olliues.

For all the purpoſes aforeſaid, the ſmall Sage is accounted to be of the more force and vertue.

## Chap. VII.

*Horminum ſatiuum.* Garden Clary.

THere is but one ſort of Garden Clary, though many wilde, which hath foure ſquares ſtalks, with broad rough wrinkled whitiſh leaues, ſomewhat vneuenly cut in on the edges, and of a ſtrong ſweete ſent, growing ſome next the ground, & ſome by couples vpon the ſtalkes: the flowers growe at certaine diſtances, with two ſmall leaues at the ioynts vnder them, ſomewhat like vnto the flowers of Sage, but leſſer, and of a very whitiſh or bleake blew colour: the ſeede is of a blackiſh browne colour, ſomewhat flat, and not ſo round as the wilde: the rootes ſpread not farre, and periſh euery yeare that they beare flowers and ſeede. It is altogether to bee ſowne of ſeed in the Spring time, yet ſometimes it will riſe of it owne ſowing.

The

### The Vſe of Clary.

The moſt frequent and common vſe of Clary, is for men or women that haue weake backes, to helpe to comfort and ſtrengthen the raines, being made into Tanſies and eaten, or otherwiſe. The ſeede is vſed of ſome to be put into the corner of the eye, if any mote or other thing haue happened into it: but aſſuredly although this may peraduenture doe ſome good, yet the ſeede of the wilde will doe much more. The leaues taken dry, and dipped into a batter made of the yolkes of egges, flower, and a little milke, and then fryed with butter vntill they be criſpe, ſerue for a diſh of meate accepted with manie, vnpleaſant to none.

## Chap. VIII.

*Nepeta.* Nep.

Although thoſe that are Herbariſts do know three ſorts of Nep, a greater & two leſſer, yet becauſe the leſſer are not vſuall, but in the Gardens of thoſe that delight in natures varieties, I do not here ſhew you them. That which is vſuall (and called of manie Cat Mint) beareth ſquare ſtalkes, but not ſo great as Clarie, hauing two leaues at euery ioynt, ſomewhat like vnto Balme or Speare Mintes, but whiter, ſofter, and longer, and nicked about the edges, of a ſtrong ſent, but nothing ſo ſtrong as Clary: the flowers growe at the toppes of the ſtalkes, as it were in long ſpikes or heads, ſomewhat cloſe together, yet compaſſing the ſtalkes at certaine ioynts, of a whitiſh colour, for forme and bigneſſe like vnto Balme, or ſomewhat bigger: the rootes are compoſed of a number of ſtrings, which dye not, but keepe greene leaues vpon them all the Winter, and ſhoote anew in the Spring. It is propagated both by the ſeede, and by ſlipping the rootes.

### The Vſe of Nep.

Nep is much vſed of women either in baths or drinkes to procure their feminine courſes: as alſo with Clarie, being fryed into Tanſies, to ſtrengthen their backes. It is much commended of ſome, if the iuyce thereof be drunke with wine, to helpe thoſe that are bruiſed by ſome fall, or other accident. A decoction of Nep is auaileable to cure the ſcabbe in the head, or other places of the body.

## Chap. IX.

*Meliſſa.* Baulme.

The Garden Baulme which is of common knowne vſe, hath diuers ſquare blackiſh greene ſtalkes, and round, hard, darke, greene pointed leaues, growing thereon by couples, a little notched about the edges, of a pleaſant ſweete ſent, drawing neareſt to the ſent of a Lemon or Citron; and therefore of ſome called *Citrago*: the flowers growe about the toppes of the ſtalkes at certaine diſtances, being ſmall and gaping, of a pale carnation colour, almoſt white: the rootes faſten themſelues ſtrongly in the ground, and endure many yeares, and is encreaſed by diuiding the rootes; for the leaues dye downe to the ground euery yeare, leauing no ſhew of leafe or ſtalke in the Winter.

The

### The Vſe of Baulme.

Baulme is often vſed among other hot and ſweete herbes, to make baths and waſhings for mens bodies or legges, in the Summer time, to warme and comfort the veines and ſinewes, to very good purpoſe and effect, and hath in former ages beene of much more vſe then now adaies. It is alſo vſed by diuers to be ſtilled, being ſteeped in Ale, to make a Baulme water, after the manner they haue beene taught, which they keepe by them, to vſe in the ſtead of *Aqua vitæ*, when they haue any occaſion for their owne or their neighbours Families, in ſuddaine qualmes or paſſions of the heart: but if they had a little better direction (for this is ſomewhat too rude) it would doe them more good that take it: For the herbe without all queſtion is an excellent helpe to comfort the heart, as the very ſmell may induce any ſo to beleeue. It is alſo good to heale greene wounds, being made into ſalues: and I verily thinke, that our forefathers hearing of the healing and comfortable properties of the true naturall Baulme, and finding this herbe to be ſo effectuall, gaue it the name of Baulme, in imitation of his properties and vertues. It is alſo an herbe wherein Bees doe much delight, as hath beene found by experience of thoſe that haue kept great ſtore; if the Hiues bee rubbed on the inſide with ſome thereof, and as they thinke it draweth others by the ſmell thereof to reſort thither. Plinie ſaith, it is a preſent remedy againſt the ſtinging of Bees.

## Chap. X.

### *Mentha.* Mintes.

THere are diuers ſorts of Mints, both of the garden, and wilde, of the woods, mountaines, and ſtanding pooles or waters: but I will onely in this place bring to your remembrance two or three ſorts of the moſt vſuall that are kept in gardens, for the vſes whereunto they are proper.

Red Mint or browne Mint hath ſquare browniſh ſtalkes, with ſomewhat long and round pointed leaues, nicked about the edges, of a darke greene colour, ſet by couples at euery ioynt, and of a reaſonable good ſent: the flowers of this kinde are reddiſh, ſtanding about the toppes of the ſtalkes at diſtances: the rootes runne creeping in the ground, and as the reſt, will hardly be cleared out of a garden, being once therein, in that the ſmalleſt peece thereof will growe and encreaſe apace.

Speare Mint hath a ſquare greene ſtalke, with longer and greener leaues then the former, ſet by couples, of a better and more comfortable ſent, and therefore of much more vſe then any other: the flowers hereof growe in long eares or ſpikes, of a pale red or bluſh colour: the rootes creepe in the ground like the other.

Party coloured or white Mint hath ſquare greene ſtalkes and leaues, ſomewhat larger then Speare Mint, and more nicked in the edges, whereof many are parted, halfe white and halfe greene, and ſome more white then greene, or more green then white, as nature liſteth: the flowers ſtand in long heads cloſe ſet together, of a bluſh colour: the rootes creepe as the reſt doe.

### The Vſe of Mintes.

Mintes are oftentimes vſed in baths, with Baulme and other herbes, as a helpe to comfort and ſtrengthen the nerues and ſinewes.

It is much vſed either outwardly applyed, or inwardly drunke, to ſtrengthen and comfort weake ſtomackes, that are much giuen to caſting: as alſo for feminine fluxes. It is boyled in milke for thoſe whoſe ſtomackes are

1 *Herminum sativum.* Garden Clary. 2 *Nepeta.* Nep. 3 *Melissa.* Baulme. 4 *Mentha sativa.* Garden Mintes.
5 *Balsamita mas, seu Costus hortorum.* Costmary. 6 *Ageratum.* Maudeline.

apt to cauſe it to curdle. And applyed with ſalt, is a good helpe for the biting of a mad dogge.

It is vſed to be boyled with Mackarell, and other fiſh.

Being dryed, is often and much vſed with Penniroyall, to bee put into puddings: as alſo among peaſe that are boyled for pottage.

Where Dockes are not ready at hand, they vſe to bruiſe Mintes, and lay them vpon any place that is ſtung with Bees, Waſpes, or ſuch like, and that to good purpoſe.

## CHAP. XI.

*Balſamita mas & famina, ſeu Coſtus hortorum maior & minor.*
Coſtmary and Maudeline.

COſtmary or Alecoaſt is a ſweet herbe, bearing many broad and long pale green leaues, ſnipped about the edges, euery one vpon a long foote-ſtalke; among which riſe vp many round greene ſtalkes, with ſuch like leaues on them, but leſſer vp to the toppe, where it ſpreadeth it ſelfe into three or foure branches, euery one bearing an vmbell or tuft of gold yellow flowers, ſomewhat like vnto Tanſie flowers, but leſſer, which turne into ſmall heads, containing ſmall flat long ſeede: the roote is ſomewhat hard and ſtringy, and being diuided, is replanted in the Spring of the yeare for increaſe.

Maudeline hath ſomewhat long and narrow leaues, ſnipt about the edges: the ſtalks are two foot high, bearing many yellow flowers on the tops of the branches, in an vmbell or tuft like vnto Tanſie: the whole herbe is ſweete, and ſomewhat bitter, and is replanted by ſlipping.

### The Vſe of Coſtmary and Maudeline.

Coſtmary is of eſpeciall vſe in the Spring of the yeare, among other ſuch like herbes, to make Sage Ale, and thereupon I thinke it tooke the name of Alecoaſt.

It is alſo vſed to be put among other ſweete herbes, to make ſweete waſhing water, whereof there is great ſtore ſpent.

The leaues haue an eſpeciall vertue to comfort both the ſtomack and heart, and to warme and dry a moiſt braine. The ſeede is much vſed in the Country, to be giuen to children for the wormes, in the ſtead of wormſeed, and ſo is the ſeede of Maudeline alſo.

Maudeline is much vſed with Coſtmary and other ſweet herbes, to make ſweete waſhing water: the flowers alſo are tyed vp with ſmall bundels of Lauender toppes, theſe being put in the middle of them, to lye vpon the toppes of beds, preſſes, &c. for the ſweete ſent and ſauour it caſteth. It is generally accounted of our Apothecaries to be the true *Eupatorium* of Auicen, and the true *Ageratum* of Dioſcorides; but Dodonæus ſeemeth to contradict both.

## CHAP. XII.

*Tanacetum vulgare & criſpum.* Tanſie.

OVr Garden Tanſie hath many hard greene leaues, or rather wings of leaues; for they are many ſmall ones, ſet one againſt another all along a middle ribbe or ſtalke, and ſnipt about the edges: in ſome the leaues ſtand cloſer and thicker, and ſomewhat crumpled, which hath cauſed it to be called double or curld Tanſie,

ſie, in others thinner and more ſparſedly: It riſeth vp with many hard ſtalks, whereon growe at the tops vpon the ſeuerall ſmall branches gold yellow flowers like buttons, which being gathered in their prime, will hold the colour freſh a long time: the ſeede is ſmall, and as it were chaffie: the roote creepeth vnder ground, and ſhooteth vp againe in diuers places: the whole herbe, both leaues and flowers, are of a ſharpe, ſtrong, bitter ſmell and taſte, but yet pleaſant, and well to be endured.

### The Vſe of Tanſie.

The leaues of Tanſie are vſed while they are young, either ſhred ſmall with other herbes, or elſe the iuyce of it and other herbes fit for the purpoſe, beaten with egges, and fryed into cakes (in Lent and the Spring of the yeare) which are vſually called Tanſies, and are often eaten, being taken to be very good for the ſtomack, to helpe to digeſt from thence bad humours that cleaue thereunto: As alſo for weak raines and kidneyes, when the vrine paſſeth away by drops: This is thought to be of more vſe for men then for women. The ſeed is much commended againſt all ſorts of wormes in children.

## CHAP. XIII.

*Pimpinella ſine Sanguiſorba.* Burnet.

BVrnet hath many winged leaues lying vpon the ground, made of many ſmall, round, yet pointed greene leaues, finely nicked on the edges, one ſet againſt another all along a middle ribbe, and one at the end thereof; from among which riſe vp diuers round, and ſometimes creſted browne ſtalkes, with ſome few ſuch like leaues on them as growe belowe, but ſmaller: at the toppes of the ſtalkes growe ſmall browne heads or knaps, which ſhoote forth ſmall purpliſh flowers, turning into long and browniſh, but a little cornered ſeede: the roote groweth downe deepe, being ſmall and browniſh: the whole plant is of a ſtipticke or binding taſte or quality, but of a fine quicke ſent, almoſt like Baulme.

### The Vſe of Burnet.

The greateſt vſe that Burnet is commonly put vnto, is to put a few leaues into a cup with Claret wine, which is preſently to be drunke, and giueth a pleaſant quicke taſte thereunto, very delightfull to the palate, and is accounted a helpe to make the heart merrie. It is ſometimes alſo while it is young, put among other Sallet herbes, to giue a finer relliſh thereunto. It is alſo vſed in vulnerary drinkes, and to ſtay fluxes and bleedings, for which purpoſes it is much commended. It hath beene alſo much commended in contagious and peſtilentiall agues.

## CHAP. XIIII.

*Hippolapathum ſatiuum, ſiue Rhabarbarum Monachorum.*
Monkes Rubarbe or Patience.

GArden Patience is a kinde of Docke in all the parts thereof, but that it is larger and taller then many others, with large and long greene leaues, a great, ſtrong, and high ſtalke, with reddiſh or purpliſh flowers, and three ſquare ſeede, like as all other Dockes haue: the roote is great and yellow, not hauing any ſhew of fleſh coloured veines therein, no more then the other kinde with great round thin leaues,

commonly called *Hippolapathum rotundifolium*, Baſtard Rubarbe, or Monkes Rubarbe, the properties of both which are of very weake effect : but I haue a kinde of round leafed Dock growing in my Garden, which was ſent me from beyond Sea by a worthy Gentleman, M^r. D^r. Matth. Liſter, one of the Kings Phyſitians, with this title, *Rhapontícum verum*, and firſt grew with me, before it was euer ſeen or known elſewhere in England, w^ch by proof I haue found to be ſo like vnto the true Rubarbe, or the Rha of Pontus, both for forme and colour, that I dare ſay it is the very true Rubarbe, our climate only making it leſſe ſtrong in working, leſſe heauy, and leſſe bitter in taſte: For this hath great and thicke rootes, as diuerſly diſcoloured with fleſh coloured veines as the true Rubarbe, as I haue to ſhew to any that are deſirous to ſee and know it ; and alſo other ſmaller ſprayes or branches of rootes, ſpreading from the maine great roote, which ſmaller branches may well be compared to the *Rhapontícum* which the Merchants haue brought vs, which we haue ſeene to be longer and ſlenderer then Rubarbe, but of the very ſame colour : this beareth ſo goodly large leaues, that it is a great beauty in a garden to behold them : for I haue meaſured the ſtalke of the leafe at the bottome next the roote to bee of the bigneſſe of any mans thumbe ; and from the roote to the leafe it ſelfe, to bee two foote in length, and ſometimes more ; and likewiſe the leafe it ſelfe, from the lower end where it is ioyned to the ſtalke, to the end or point thereof, to bee alſo two foote in length, and ſometimes more ; and alſo in the broadeſt part of the leafe, to be two foote or more ouer in breadth : it beareth whitiſh flowers, contrary to all other Dockes, and three ſquare browniſh ſeede as other Dockes doe, but bigger, and therefore aſſuredly it is a Docke, and the true Rubarbe of the Arabians, or at the leaſt the true *Rhapontícum* of the Ancients. The figure of the whole plant I haue cauſed to be cut, with a dryed roote as it grew in my garden by it ſelfe, and haue inſerted it here, both becauſe Matthiolus giueth a falſe figure of the true Rubarbe, and that this hath not been expreſſed and ſet forth by any before.

### The Vſe of Patience, and of the Rubarbe.

The leaues of Patience are often, and of many vſed for a pot-herbe, and ſeldome to any other purpoſe : the roote is often vſed in Diet-beere, or ale, or in other drinkes made by decoction, to helpe to purge the liuer, and clenſe the blood. The other Rubarbe or *Rhapontícum*, wherof I make mention, and giue you here the figure, I haue tryed, and found by experience to purge gently, without that aſtriction that is in the true Rubarbe is brought vs from the Eaſt Indies, or China, and is alſo leſſe bitter in taſte ; whereby I coniecture it may bee vſed in hot and feaueriſh bodies more effectually, becauſe it doth not binde after the purging, as the Eaſt India Rubarbe doth : but this muſt bee giuen in double quantitie to the other, and then no doubt it will doe as well : The leaues haue a fine acide taſte : A ſyrrupe therefore made with the iuice and ſugar, cannot but be very effectuall in deiected appetites, and hot fits of agues ; as alſo to helpe to open obſtructions of the liuer, as diuers haue often tryed, and found auaileable by experience.

## Chap. XV.

### *Lapathum ſanguineum.* Blood-wort.

AMong the ſorts of pot-herbes Blood-worte hath alwayes beene accounted a principall one, although I doe not ſee any great reaſon therein, eſpecially ſeeing there is a greater efficacie of binding in this Docke, then in any of the other : but as common vſe hath receiued it, ſo I here ſet it downe. Blood-worte is one of the ſorts of Dockes, and hath long leaues like vnto the ſmaller yellow Docke, but ſtriped with red veines, and ouer-ſhadowed with red vpon the greene leafe, that it ſeemeth almoſt wholly red ſometimes : the ſtalke is reddiſh, bearing ſuch like leaues, but ſmaller

1 *Tanacetum.* Tanſie. 2 *Pimpinella.* Burnet. 3 *Rhaponticum verum ſeu potius Rhabarbarum verum.* True Raponticke or rather true Rubarbe. 4 *Lapathum ſativum ſeu Patientia.* Monkes Rubarbe or Patience. 5 *Lapathum ſanguineum.* Bloudwort. 6 *Acetoſa.* Sorrell.



3. Rheum rhaponticum, L. Fruit. distr.

ſmaller vp to the toppe, where it is diuided into diuers ſmall branches, whereon grow purpliſh flowers, and three ſquare darke red ſeede, like vnto others: the roots are not great, but ſomewhat long, and very red, abiding many yeares, yet ſometimes ſpoiled with the extremitie of winter.

### The Vſe of Blood-worte.

The whole and onely vſe of the herbe almoſt, ſerueth for the pot, among other herbes, and, as I ſaid before, is accounted a moſt eſpeciall one for that purpoſe. The ſeede therof is much commended for any fluxe in man or woman, to be inwardly taken, and ſo no doubt is the roote, being of a ſtipticke qualitie.

## Chap. XVI.

*Oxalis ſiue Acetoſa.* Sorrell.

SOrrell muſt needes bee reckoned with the Dockes, for that it is ſo like vnto them in all things, and is of many called the ſower Docke. Of Sorrels there are many ſorts, but I ſhall not trouble you with any other in this place, then the common Garden Sorrell, which is moſt knowne, and of greateſt vſe with vs; which hath tender greene long leaues full of iuice, broade, and bicorned as it were, next vnto the ſtalke, like as Arrach, Spinach, and our Engliſh Mercurie haue, of a ſharpe ſower taſte: the ſtalkes are ſlender, bearing purpliſh long heads, wherein lye three ſquare ſhining browne ſeede, like, but leſſer then the other: the root is ſmaller then any of the other Dockes, but browne, and full of ſtrings, and abideth without decaying, hauing greene leaues all the winter, except in the very extremitie thereof, which often taketh away all or moſt of his leaues.

### The Vſe of Sorrell.

Sorrell is much vſed in ſawces, both for the whole, and the ſicke, cooling the hot liuers, and ſtomackes of the ſicke, and procuring vnto them an appetite vnto meate, when their ſpirits are almoſt ſpent with the violence of their furious or fierie fits; and is alſo of a pleaſant relliſh for the whole, in quickning vp a dull ſtomacke that is ouer-loaden with euery daies plenty of diſhes. It is diuers waies dreſſed by Cooks, to pleaſe their Maſters ſtomacks.

## Chap. XVII.

*Bugloſſum luteum, ſiue Lingua Bouis.* Langdebeefe.

VNto this place may well bee referred our ordinary Borage and Bugloſſe, ſet forth in the former Booke, in regard of the properties whereunto they are much employed, that is, to ſerue the pot among other herbes, as is ſufficiently knowne vnto all. And yet I confeſſe, that this herbe (although it bee called *Bugloſſum luteum*, as if it were a kind of Bugloſſe) hath no correſpondency with Bugloſſe or Borage in any part, ſauing only a little in the leafe; & our Borage or Bugloſſe might more fitly, according to the Greeke name, bee called Oxe tongue or Langdebeefe; and this might in my iudgement more aptly be referred to the kinds of *Hieratium* Hawkeweed, whereunto it neereſt approacheth: but as it is commonly receiued, ſo take it in this place, vntill it come to receiue the place is proper for it. It hath diuers broad and long darke green leaues, lying vpon the ground, very rough in handling, full of ſmall haires or prickes, ready to enter into the hands of any that handle it; among which riſeth

vp

1 *Lingua bouis sive Buglossum luteum.* Langdebeefe. 2 *Atriplex sive Olus aureum.* Arrach. 3 *Blitum;* Blites.
4 *Beta.* Beetes. 5 *Hipposelinum sive Olus atrum.* Allisanders. 6 *Selinum dulce.* Sweete Parsley.

vp a round greene hairy or prickly ſtalk, bearing at the toppe, among a few ſmall green leaues, diuers ſmall yellow flowers in rough heads, which turne into doune, containing within them browne yellowiſh ſmall long ſeedes, ſomewhat like vnto the ſeede of Hawkeweede: the roote is wooddy, w hich periſheth quickly after it hath borne ſeed; but is tender while it is young.

### The Vſe of Langdebeefe.

The leaues are onely vſed in all places that I know, or euer could learne, for an herbe for the pot among others, and is thought to bee good to looſen the belly.

## Chap. XVIII.

### *Atriplex ſiue Olus Aureum.* Arrach.

THere be diuers kindes of Arrach, or Orach, as ſome doe call them; ſome of the Garden, whereof I meane to entreate in this place; others wilde of the Fieldes, &c. and others of the Sea, which are not to bee ſpoken of in this worke, but referred to a generall hiſtorie. The white garden Arrach, or Orach, hath diuers leaues, ſtanding vpon their ſeuerall footeſtalkes, broade at the bottome, ending in two points like an arrow, with two feathers at the head, and ſmall pointed at the end of the leafe, of a whitiſh yellow greene colour, and as it were ſtrewed ouer with flower or meale, eſpecially while they are young: the ſtalke likewiſe is mealy, bearing many branches with ſmall yellow flowers on them, which turne into ſmall leafie ſeeds: the rooote groweth ſomewhat deepe in the ground, with many ſmall threds faſtened thereto: it quickly ſpringeth vp of the ſeede, groweth great, and fadeth away as ſoon as it hath borne ſeede.

The purple Arrach is in all things like vnto the white, ſauing onely in the colour of the leafe, ſtalke, ſeede, &c. which are all of a mealy duſty purpliſh colour.

### The Vſe of Arrach.

Arrach is cold and moiſt, and of a lubricke or ſlippery qualitie, whereby it quickely paſſeth through the ſtomacke and belly, and maketh it ſoluble, and is of many vſed for that purpoſe, being boyled and buttered, or put among other herbes into the pot to make pottage.

There are many diſhes of meate made with them while they are young, for being almoſt without ſauour of themſelues, they are the more conuertible into what relliſh any one will make them with Sugar, Spice &c.

## Chap. XIX.

### *Blitum.* Blites.

THere be diuers ſorts of Blites, ſome whereof I haue entreated in the former part of this worke, vnder the title of *Amaranthus*, Flower gentle: others that are nourſed vp in Gardens, I will ſet forth in this place, which are onely two, that haue come to my knowledge, that is, the white and the red, and are of a qualitie as neere vnto Arrach as vnto Beetes, participating of both, and therefore I haue placed them betwixt them. The white Blite hath leaues ſomewhat like vnto Beetes, but ſmaller, rounder, and of a whitiſh greene colour, euery one ſtanding vpon a ſmall long footeſtalke: the ſtalke riſeth vp two or three foote high, with many ſuch like leaues thereon: the flowers grow at the top in long round tufts or cluſters, wherein are contained

tained ſmall round ſeede: the roote is very full of threds or ſtrings.

The red Blite is in all things like the white, but that his leaues and tufted heades are exceeding red at the firſt, and after turne more purpliſh.

### The Vſe of Blites.

Blites are vſed as Atrach, eyther boyled of it ſelfe or ſtewed, which they call Loblolly, or among other herbes to bee put into the pot; and yet ſome doe vtterly refuſe it, becauſe in diuers it prouoketh caſtings. It is altogether inſipide or without taſte, but yet by reaſon of the moiſt ſlipperie qualitie it hath, it helpeth to looſen the belly. The vnſauorineſſe whereof hath in many Countries growne into a prouerbe, or by-word, to call dull, ſlow, or lazie perſons by that name: They are accounted more hurtfull to the ſtomacke, and ſo to the head and eyes, then other herbes, and therefore they are the leſſe vſed.

## Chap. XX.

### *Beta.* Beetes.

THere are many diuerſities of Beetes, ſome growing naturally in our own Country, others brought from beyond Sea; whereof ſome are white, ſome greene, ſome yellow, ſome red: the leaues of ſome are of vſe only, and the root not vſed: others the roote is only vſed, and not the leaues: and ſome againe, both roote and leafe. The ancient Authors, as by their workes appeare, knew but two ſorts, the white and the blacke Beete, whereof the white is ſufficiently known, and was of them termed *Sicula*, of the later Phyſitians *Sicla*, becauſe it was thought firſt to be brought from Sicilie: the blacke abideth ſome controuerſie; ſome thinking that our common greene Beete, becauſe it is of a darke greene colour, was that they called the blacke Beete; others that our ſmall red Beete, which is of a darke red colour, was their black Beete, which in my opinion is the more likely: But to come to the matter in hand, and giue you the deſcriptions of them which are in vſe with vs, and leaue controuerſies to ſuch a worke as is fit for them, wherein all ſuch matters may be diſcuſſed at large.

The common white Beete hath many great leaues next the ground (in ſome hot Countries growing to be three foote long, and very broade, in our Countrey they are very large, but nothing neere that proportion) of a whitiſh greene colour; the ſtalke is great, ſtrong, and ribbed or creſted, bearing great ſtore of leaues vpon it vp to the very toppe almoſt: the flowers grow in very long tufts, ſmall at the ends, and turning down their heads, which are ſmall pale greeniſh yellow burres, giuing cornered prickly ſeede: the roote is great, long and hard, when it hath giuen ſeede, of no vſe at all, but abideth a former winter with his leaues vpon it, as all other ſorts following doe.

The common red Beet differeth not from the white Beete, but only that it is not ſo great, and both the leaues and rootes are ſomewhat red: the leaues bee in ſome more red then in others, which haue but red veines or ſtrakes in them, in ſome alſo of a freſh red, in others very darke red: the roote hereof is red, ſpongy, and not vſed to bee eaten.

The common greene Beete is alſo like vnto the white Beete, but of a darke greene colour. This hath beene found neere the ſalt Marſhes by Rocheſter, in the foote-way going from the Lady Leueſons houſe thither, by a worthy, diligent and painefull obſeruer and preſeruer both of plants and all other natures varieties, often remembred before in this worke, called Iohn Tradeſcante, who there finding it, gaue me the knowledge thereof, and I haue vpon his report ſet it here down in this manner:

The Romane red Beete, called *Beta rapoſa*, is both for leafe and roote the moſt excellent Beete of all others: his rootes bee as great as the greateſt Carrot, exceeding red both within and without, very ſweete and good, fit to bee eaten: this Beete groweth higher then the laſt red Beete, whoſe rootes are not vſed to bee eaten: the leaues likewiſe

wiſe are better of taſte, and of as red a colour as the former red Beete : the roote is ſometimes ſhort like a Turnep, whereof it took the name of *Rapa* or *rapoſa*; and ſometimes as I ſaid before, like a Carrot and long : the ſeede is all one with the leſſer red Beete.

The Italian Beete is of much reſpect, whoſe faire greene leaues are very large and great, with great white ribbes and veines therein: the ſtalke in the Summer time, when it is growen vp to any height, is ſix ſquare in ſhew, and yellowiſh withall, as the heades with ſeede vpon them ſeeme likewiſe.

The great red Beete that Maſter Lete a Merchant of London gaue vnto Maſter Gerrard, as he ſetteth it downe in his Herball, ſeemeth to bee the red kinde of the laſt remembred Beete, whoſe great ribbes as he ſaith, are as great as the middle ribbe of the Cabbage leafe, and as good to bee eaten; whoſe ſtalke roſe with him to the height of eight cubits, and bore plenty of ſeede.

### The Vſe of Beetes.

Beetes, both white, greene and red, are put into the pot among other herbes, to make pottage, as is commonly known vnto all, and are alſo boyled whole, both in France vſually with moſt of their boyled meates, and in our Countrey, with diuers that delight in eating of herbes.

The Italian Beete, and ſo likewiſe the laſt red Beete with great ribbes, are boyled, and the ribbes eaten in ſallets with oyle, vinegar and pepper, and is accounted a rare kinde of ſallet, and very delicate.

The roote of the common red Beete with ſome, but more eſpecially the Romane red Beete, is of much vſe among Cookes to trimme or ſet out their diſhes of meate, being cut out into diuers formes and faſhions, and is grown of late dayes into a great cuſtome of ſeruice, both for fiſh and fleſh.

The rootes of the Romane red Beete being boyled, are eaten of diuers while they are hot with a little oyle and vinegar, and is accounted a delicate ſallet for the winter; and being cold they are ſo vſed and eaten likewiſe.

The leaues are much vſed to mollifie and open the belly, being vſed in the decoction of Gliſters. The roote of the white kinde ſcraped, and made vp with a little honey and ſalt, rubbed on and layd on the belly, prouoketh to the ſtoole. The vſe of eating Beetes is likewiſe held to bee helpefull to ſplenetick perſons.

## Chap. XXI.

### *Hippoſelinum, ſiue Olus atrum.* Aliſanders.

ALiſanders hath beene in former times thought to be the true Macedonian Parſley, and in that errour many doe yet continue : but this place giueth not leaue to diſcuſſe that doubt : but I muſt here only ſhew you, what it is, and to what vſe it is put ordinarily for the Kitchen. The leaues of Aliſanders are winged or cut into many parts, ſomewhat reſembling Smallage, but greater, broader, and more cut in about the edges : the ſtalkes are round and great, two foote high or better, bearing diuers leaues on them, and at the toppe ſpokie roundles of white flowers on ſeuerall ſmall branches, which turne into blacke ſeede, ſomewhat cornered or creſted, of an aromaticall bitter taſte : the roote is blacke without, and white within, and abideth well the firſt year of the ſowing, periſhing after it hath borne ſeed.

### The Vſe of Aliſanders.

The tops of the rootes, with the lower part of the ſtalkes of Aliſanders, are vſed in Lent eſpecially, and Spring of the yeare, to make broth, which although it be a little bitter, yet it is both wholſome, and pleaſing to a great many,

many, by reaſon of the aromaticall or ſpicie taſte, warming and comforting the ſtomack, and helping it digeſt the many wateriſh and flegmaticke meates are in thoſe times much eaten. The rootes alſo either rawe or boyled are often eaten with oyle and vinegar. The ſeede is more vſed phyſically then theroote, or any other part, and is effectuall to prouoke plenty of vrine in them that piſſe by drops, or haue the Strangury: It helpeth womens courſes, and warmeth their benummed bodies or members, that haue endured fierce cold daies and nights, being boyled and drunke.

## Chap. XXII.

### *Selinum dulce.* Sweete Parſley or ſweete Smallage.

THis kinde of ſweete Parſley or Smallage, which ſoeuer you pleaſe to call it; for it reſembleth Smallage as well in the largeneſſe of the leaues, as in the taſte, yet ſweeter and pleaſanter, is (as I take it) in this like vnto ſweete Fennell (that hath his ſweetneſſe from his naturall foyle and clymate; for howſoeuer it bee reaſonable ſweete the firſt yeare it is ſowne with vs, yet it quickly doth degenerate, and becommeth no better then our ordinarie Fennell afterwards). The firſt yeare it is ſowne and planted with vs (and the firſt that euer I ſaw, was in a Venetian Ambaſſadours Garden in the Spittle yard, neare Biſhops gate ſtreete) is ſo ſweete and pleaſant, eſpecially while it is young, as if Sugar had beene mingled with it: but after it is growne vp high and large, it hath a ſtronger taſte of Smalladge, and ſo likewiſe much more the next yeare; that it groweth from the ſeed was gathered here: the leaues are many, ſpreading farre about the roote, broader and of a freſher greene colour then our ordinary Smalladge, and vpon longer ſtalkes: the ſeed is as plentifull as Parſley, being ſmall and very like vnto it, but darker of colour.

### The Vſe of ſweete Parſley.

The Venetians vſe to prepare it for meate many waies, both the herbe and the roote eaten rawe, as many other herbes and rootes are, or boyled or fryed to be eaten with meate, or the dryed herbe poudered and ſtrewed vpon meate; but moſt vſually either whited, and ſo eaten rawe with pepper and oyle, as a dainty Sallet of it ſelfe, or a little boyled or ſtewed: the taſte of the herbe being a little warming, but the ſeede much more, helpeth cold windy ſtomackes to digeſt their meate, and to expell winde.

## Chap. XXIII.

### *Petroſolinum & Apium.* Parſley and Smalledge.

WE haue three ſorts of Parſley in our Gardens, and but one of Smalladge; Our common Parſley, Curld Parſley, and Virginia Parſley; which laſt, although it be but of late knowne, yet it is now almoſt growne common, and of as good vſe as the other with diuers. Our common Parſley is ſo well knowne, that it is almoſt needleſſe to deſcribe it, hauing diuers freſh greene leaues, three alwaies placed together on a ſtalke, and ſnipt about the edges, and three ſtalkes of leaues for the moſt part growing together: the ſtalkes growe three or foure foote high or better, bearing ſpikie heads of white flowers, which turne into ſmall ſeede, ſomewhat ſharpe and hot in taſte: the roote is long and white.

Curld Parſley hath his leaues curled or crumpled on the edges, and therein is the onely difference from the former.

Virginia

Virginia Parſley is in his leafe altogether like vnto common Parſley for the forme, conſiſting of three leaues ſet together, but that the leaues are as large as Smallage leaues, but of a pale or whitiſh greene colour, and of the ſame taſte of our common Parſley: the ſeede hereof is as the leaues, twice if not thrice as bigge as the ordinary Parſley, and periſheth when it hath giuen ſeede, abiding vſually the firſt yeare of the ſowing.

Smallage is in forme ſomewhat like vnto Parſley, but greater and greener, and leſſe pleaſant, or rather more bitter in taſte: the ſeede is ſmaller, and the root more ſtringy.

### The Vſe of Parſley.

Parſley is much vſed in all ſorts of meates, both boyled, roaſted, fryed, ſtewed, &c. and being greene it ſerueth to lay vpon ſundry meates, as alſo to draw meate withall. It is alſo ſhred and ſtopped into poudered beefe, as alſo into legges of Mutton, with a little beefe ſuet among it, &c.

The rootes are often vſed to be put into broth, to helpe to open obſtructions of the liuer, reines, and other parts, helping much to procure vrine.

The rootes likewiſe boyled or ſtewed with a legge of Mutton, ſtopped with Parſley as aforeſaid, is very good meate, and of very good relliſh, as I haue proued by the taſte; but the rootes muſt bee young, and of the firſt yeares growth, and they will haue their operation to cauſe vrine.

The ſeed alſo is vſed for the ſame cauſe, when any are troubled with the ſtone, or grauell, to open the paſſages of vrine.

Although Smallage groweth in many places wilde in moiſt grounds, yet it is alſo much planted in Gardens, and although his euill taſte and ſauour doth cauſe it not to be accepted into meates as Parſley, yet it is not without many ſpeciall good properties, both for outward and inward diſeaſes, to helpe to open obſtructions, and prouoke vrine. The iuyce cleanſeth vlcers; and the leaues boyled with Hogs greaſe, healeth felons on the ioynts of the fingers.

## Chap. XXIIII.

### *Fœniculum.* Fenell.

THere are three ſorts of Fenell, whereof two are ſweete. The one of them is the ordinary ſweete Fenell, whoſe ſeedes are larger and yellower then the common, and which (as I ſaid before in the Chapter of ſweete Parſley) doth ſoone degenerate in this our Country into the common. The other ſweete Fenell is not much knowne, and called Cardus Fenell by thoſe that ſent it out of Italy, whoſe leaues are more thicke and buſhie then any of the other. Our common Fenell, whereof there is greene and red, hath many faire and large ſpread leaues, finely cut and diuided into many ſmall, long, greene, or reddiſh leaues, yet the thicker tufted the branches be, the ſhorter are the leaues: the ſtalkes are round, with diuers ioynts and leaues at them, growing fiue or ſix foot high, bearing at the top many ſpoakie rundels of yellow flowers: the Common, I meane, doth turne into a darke grayiſh flat ſeede, and the Sweete into larger and yellower: the roote is great, long, and white, and endureth diuers yeares.

### The Vſe of Fenell.

Fenell is of great vſe to trimme vp, and ſtrowe vpon fiſh, as alſo to boyle or put among fiſh of diuers ſorts, Cowcumbers pickled, and other fruits, &c. The rootes are vſed with Parſley rootes, to be boyled in broths and drinkes to open obſtructions. The ſeed is of much vſe with other things to expell winde. The ſeede alſo is much vſed to be put into Pippin pies, and diuers

other

1 *Petrosolinum.* Parsley. 2 *Apium.* Smallage. 3 *Fœniculum.* Fenell. 4 *Anethum.* Dill. 5 *Myrrhis siue Cerefolium magnum.* Sweete Cherrill. 6 *Cerefolium vulgare.* Common Chesrill.

other ſuch baked fruits, as alſo into bread, to giue it the better relliſh.

The ſweete Cardus Fenell being ſent by Sir Henry Wotton to Iohn Tradeſcante, had likewiſe a large direction with it how to dreſſe it; for they vſe to white it after it hath been tranſplanted for their vſes, which by reaſon of the ſweetneſſe by natnre, and the tenderneſſe by art, cauſeth it to be the more delightfull to the taſte, eſpecially with them that are accuſtomed to feede on greene herbes.

## Chap. XXV.

### *Anethum.* Dill.

Dill doth much growe wilde, but becauſe in many places it cannot be had, it is therefore ſowne in Gardens for the vſes whereunto it ſerueth. It is a ſmaller herbe then Fenell, but very like, hauing fine cut leaues, not ſo large, but ſhorter, ſmaller, and of a ſtronger and quicker taſte: the ſtalke is ſmaller alſo, and with few ioynts and leaues on them, bearing ſpoakie tufts of yellow flowers, which turne into thinne, ſmall, and flat ſeedes: the roote periſheth euery yeare, and riſeth againe for the moſt part of it owne ſowing.

### The Vſe of Dill.

The leaues of Dill are much vſed in ſome places with Fiſh, as they doe Fenell; but becauſe it is ſo ſtrong many doe refuſe it.

It is alſo put among pickled Cowcumbers, wherewith it doth very well agree, giuing vnto the cold fruit a pretty ſpicie taſte or relliſh.

It being ſtronger then Fenell, is of the more force to expell winde in the body. Some vſe to eate the ſeed to ſtay the Hickocke.

## Chap. XXVI.

### *Myrrhis ſiue Cerefolium maius & vulgare.*
### Sweet Cheruill and ordinary Cheruill.

The great or ſweete Cheruill (which of ſome is called Sweete Cicely) hath diuers great and faire ſpread winged leaues, conſiſting of many leaues ſet together, deeply cut in the edges, and euery one alſo dented about, very like, and reſembling the leaues of Hemlockes, but of ſo pleaſant a taſte, that one would verily thinke, he chewed the leaues or ſeedes of Aniſeedes in his mouth: The ſtalke is reaſonable great, and ſomewhat cornered or creſted about three or foure foote high, at the toppe whereof ſtand many white ſpoakie tufts of flowers, which change into browne long cornered great ſeede, two alwaies ioyned together: the roote is great, blackiſh on the outſide, and white within, with diuers fibres annexed vnto it, and periſheth not, but abideth many yeares, and is of a ſweete, pleaſant, and ſpicie hot taſte, delightfull vnto many.

The common Cheruill is a ſmall herbe, with ſlender leaues, finely cut into long peeces, at the firſt of a pale yellowiſh greene colour, but when the ſtalke is growne vp to ſeede, both ſtalkes and leaues become of a darke red colour: the flowers are white, ſtanding vpon ſcattered or thin ſpread tufts, which turne into ſmall, long, round, and ſharpe pointed ſeedes, of a browniſh blacke colour: the roote is ſmall, with diuers long ſlender white ſtrings, and periſheth euery yeare.

The

### The Vſes of theſe Cheruils.

The common Cheruill is much vſed of the French and Dutch people, to bee boyled or ſtewed in a pipkin, eyther by it ſelfe, or with other herbes, whereof they make a Loblolly, and ſo eate it. It is vſed as a pot-herbe with vs.

Sweete Cheruill, gathered while it is young, and put among other herbes for a ſallet, addeth a meruellous good reliſh to all the reſt. Some commend the greene ſeedes ſliced and put in a ſallet of herbes, and eaten with vinegar and oyle, to comfort the cold ſtomacke of the aged. The roots are vſed by diuers, being boyled, and after eaten with oyle and vinegar, as an excellent ſallet for the ſame purpoſe. The preſerued or candid rootes are of ſingular good vſe to warme and comfort a cold flegmaticke ſtomack, and is thought to be a good preſeruatiue in the time of the plague.

## Chap. XXVII.

### *Malua Criſpa.* French Mallowes.

THe curld or French Mallow groweth vp with an vpright greene round ſtalke, as high vſually as any man, whereon from all ſides grow forth round whitiſh greene leaues, curld or crumpled about the edges, like a ruffe, elſe very like vnto an ordinary great Mallow leafe: the flowers grow both vpon the ſtalke, and on the other branches that ſpring from them, being ſmall and white; after which come ſmall caſes with blacke ſeede like the other Mallowes: the roote periſheth when it hath borne ſeede, but abideth vſually the firſt yeare, and the ſecond runneth vp to flower and ſeede.

### The Vſe of French Mallowes.

It is much vſed as a pot-herbe, eſpecially when there is cauſe to moue the belly downward, which by his ſlippery qualitie it doth helpe forward. It hath beene in times paſt, and ſo is to this day in ſome places, vſed to be boyled or ſtewed, eyther by it ſelfe with butter, or with other herbes, and ſo eaten.

## Chap. XXVIII.

### *Intubum.* Succorie and Endiue.

I Put both Succorie and Endiue into one chapter and deſcription, becauſe they are both of one kindred; and although they differ a little the one from the other, yet they agree both in this, that they are eaten eyther greene or whited, of many.

Endiue, the ſmooth as well as the curld, beareth a longer and a larger leafe then Succorie, and abideth but one yeare, quickely running vp to ſtalke and ſeede, and then periſheth: whereas Succorie abideth many years, and hath long and narrower leaues, ſomewhat more cut in, or torne on the edges: both of them haue blew flowers, and the ſeede of the ſmooth or ordinary Endiue is ſo like vnto the Succorie, that it is very hard to diſtinguiſh them aſunder by ſight; but the curld Endiue giueth blackiſh and flat ſeede, very like vnto blacke Lettice ſeede: the rootes of the Endiue periſh, but the Succorie abideth.

### The Vſe of Succory and Endiue.

Although Succorie bee ſomewhat more bitter in taſte then the Endiues,

yet it is oftentimes, and of many eaten greene, but more vſually being buried a while in ſand, that it may grow white, which cauſeth it to loſe both ſome part of the bitterneſſe, as alſo to bee the more tender in the eating; and Horace ſheweth it to be vſed in his time, in the 32. Ode of his firſt Book, where he ſaith,

*Me paſcunt Oliuæ, me Cithoreа leueſq; Maluæ.*

Endiue being whited in the ſame, or any other manner, is much vſed in winter, as a ſallet herbe with great delight; but the curld Endiue is both farre the fairer, and the tenderer for that purpoſe.

## Chap. XXIX.

### *Spinachia, ſiue Olus Hiſpanicum.* Spinach.

SPinach or Spinage is of three ſorts (yet ſome doe reckon of foure, accounting that herbe that beareth no ſeede to be a ſort of it ſelfe, when it is but an accident of nature, as it falleth out in Hempe, Mercury, and diuers other herbes) two that bear prickly ſeed, the one much greater then the other: the third that beareth a ſmooth ſeede, which is more daintie, and nourſed vp but in few Gardens: The common Spinach which is the leſſer of the two prickly ſorts, hath long greene leaues, broad at the ſtalke, and rent, or torne as it were into foure corners, and ſharpe pointed at the ends: it quickly runneth vp to ſtalke, if it be ſowen in the Spring time; but elſe, if at the end of Summer, it will abide all the winter green, and then ſuddenly in the very beginning of the Spring, runne vp to ſtalke, bearing many leaues both below and at the toppe, where there doth appeare many ſmal greeniſh flowers in cluſters, and after them prickly ſeede: The other greater ſort that hath prickly ſeede, is in all things like the former, but larger both in ſtalke, leafe and ſeede. The ſmooth Spinach hath broader, and a little rounder pointed leaues then the firſt, eſpecially the lower leaues; for thoſe that grow vpwards vpon the ſtalke, are more pointed, and as it were three ſquare, of as darke a greene colour as the former: at the ſeuerall ioynts of the ſtalkes and branches, ſtand cluſtering many ſmall greeniſh flowers, which turne into cluſters of round whitiſh ſeede, without any prickles at all vpon them: the roote is long, white and ſmall, like vnto the other, with many fibres at it: If it be often cut, it will grow the thicker, or elſe ſpindle vp very thinly, and with but few leaues vpon the ſtalke.

### The Vſe of Spinage.

Spinage is an herbe fit for ſallets, and for diuers other purpoſes for the table only; for it is not knowne to bee vſed Phyſically at all. Many Engliſh that haue learned it of the Dutch people, doe ſtew the herbe in a pot or pipkin, without any other moiſture then it owne, and after the moiſture is a little preſſed from it, they put butter, and a little ſpice vnto it, and make therewith a diſh that many delight to eate of. It is vſed likewiſe to be made into Tartes, and many other varieties of diſhes, as Gentlewomen and their Cookes can better tell then my ſelfe; vnto whom I leaue the further ordering of theſe herbes, and all other fruits and rootes of this Garden: For I intend only to giue you the knowledge of them, with ſome briefe notes for their vſe, and no more.

1 *Malua crispa.* French Mallowes. 2 *Endiuia.* Endiue. 3 *Cicherium.* Succory. 4 *Spinachia.* Spinach. 5 *Lactuca crispa.* Curld Lettice. 6 *Lactuca*
*patula.* An open Lettice. 7 *Lactuca capitata vulgaris.* Ordinary cabbage Lettice. 8 *Lactuca capitata Romana.* The great Romane cabbage Lettice. 9 *Lactuca agnina.*
*cornina.* Corne Sallet or Lambes Lettice.

## Chap. XXX.

### *Lactuca.* Lettice.

THere are ſo many ſorts, and ſo great diuerſitie of Lettice, that I doubt I ſhall ſcarce be beleeued of a great many. For I doe in this Chapter reckon vp vnto you eleauen or twelue differing ſorts; ſome of little vſe, others of more, being more common and vulgar; and ſome that are of excellent vſe and ſeruice, which are more rare, and require more knowledge and care for the ordering of them, as alſo for their time of ſpending, as ſome in the ſpring, ſome in ſummer, others in autumne, and ſome being whited for the winter. For all theſe ſorts I ſhall not neede many deſcriptions, but only ſhew you which doe cabbage, and which are looſe, which of them are great or ſmall, white, greene or red, and which of them beare white ſeeds, and which of them blacke. And laſtly I haue thought good to adde another Sallet herbe, which becauſe it is called Lambes Lettice of many, or Corne Sallet of others, is put in only to fill vp a number in this Chapter, and that I muſt ſpeake of it, and not that I thinke it to be any of the kindes of Lettice.

All ſorts of Lettice, after a while that they haue cloſed themſelues, if they bee of the Cabbage kindes, or otherwiſe being looſe, and neuer cloſing, ſend forth from among the middle of their leaues a round ſtalke (in ſome greater, in others leſſer, according to their kinde) full of leaues like vnto the lower, branching at the toppe into ſundry parts, whereon grow diuers ſmall ſtar-like flowers, of a pale yellowiſh colour; after which come ſeede, eyther white or blackiſh, as the plant yeeldeth, whereat hangeth ſome ſmall peece of a cottony doune, wherewith the whole head is ſtored, and is carried away with the winde, if it be not gathered in time: the roote is ſomewhat long and white, with ſome fibres at it, and periſheth quickely after the ſeede is ripe.

The Romane red Lettice is the beſt and greateſt of all the reſt. For Iohn Tradeſcante that firſt, as I thinke, brought it into England, and ſowed it, did write vnto mee, that after one of them had been bound and whited, when the refuſe was cutaway, the reſt weighed ſeuenteene ounces: this hath blacke ſeede.

The white Romane Lettice is like vnto it, hauing long leaues like a Teaſell, it is in goodneſſe next vnto the red, but muſt be whited, that it may eate kindly: the ſeede hereof is white.

The Virginia Lettice hath ſingle and very broade reddiſh leaues, and is not of any great regard, and therefore is kept but of a few: it beareth blacke ſeede.

The common Lumbard Lettice that is looſe, and another kinde thereof that doth ſomewhat cabbage, haue both white ſeedes.

The Venice Lettice is an excellent Cabbage Lettice, and is beſt to bee ſowen after Midſummer for lateward Lettice; they be ſometimes as great as the crowne of a mans hatt: the ſeede hereof is white, and groweth to be of a meane height.

Our common Cabbage Lettice is well known, and beareth blacke ſeede.

The curld Lettice which is open, and differeth but little from Endiue, beareth black ſeede.

Another ſort of curld Lettice doth cabbage, and is called Flanders Cropers, or Cropers of Bruges; this groweth loweſt, and hath the ſmalleſt head, but very hard and round, and white while it groweth: the ſeed is blacke.

A kinde of Romane Lettice is of a darke green colour, growing as low as the Venice Lettice, and is an excellent kinde, bearing blacke ſeede.

And laſtly our winter Lettice is wonderfull hardy to endure our cold: It is but ſingle, and muſt be ſowen at Michaelmas, but will be very good, before any of the other good ſorts ſowen in the Spring, will be ready to be vſed, and beareth white ſeed.

To inſtruct a nouice (for I teach not a Gardiner of knowledge) how to gather his ſeede that it may be good, is in this manner: Let him marke out thoſe plants that hee meaneth ſhall run vp for ſeede, which muſt be the moſt likely; & after they haue begun to ſhoote forth ſtalkes, ſtrip away the lower leaues, for two or three hands breadth aboue the ground, that thereby in taking away the loweſt leaues, the ſtalke doe not rot, nor the ſeed be hindered in the ripening.

There

There are two manner of wayes to whiten Lettice to make them eate the more tender : the one is by rayſing vp earth like moale hils, round about the plants while they are growing, which will make them grow white : the other is by tying vp all the looſe leaues round together while it groweth, that ſo the cloſe tying may make it grow white, and thereby be the more tender.

Lambes Lettice or Corne Sallet is a ſmall plant while it is young, growing cloſe vpon the ground, with many whitiſh greene, long and narrow, round pointed leaues, all the winter, and in the beginning of the ſpring (if it bee ſowen in autumne, as it is vſuall to ſerue for an early ſallet) riſeth vp with ſmall round ſtalkes, with two leaues at euery ioynt, branching forth at the toppe, and bearing tufts of ſmall bleake blew flowers, which turne into ſmall round whitiſh ſeede : the roote is ſmall and long, with ſome ſmall threds hanging thereat : the whole plant is of a wateriſh taſte, almoſt inſipide.

### The Vſe of Lettice.

All ſorts of Lettice are ſpent in ſallets, with oyle and vinegar, or as euery one pleaſe, for the moſt part, while they are freſh and greene, or whited, as is declared of ſome of the ſorts before, to cauſe them to eate the more delicate and tender. They are alſo boyled, to ſerue for many ſorts of diſhes of meate, as the Cookes know beſt.

They all coole a hot and fainting ſtomacke.

The iuice of Lettice applyed with oyle of Roſes to the foreheads of the ſicke and weake wanting ſleepe, procureth reſt, and taketh away paines in the head : bound likewiſe to the cods, it helpeth thoſe that are troubled with the Colts euill. If a little camphire be added, it reſtraineth immoderate luſt : but it is hurtfull to ſuch as are troubled with the ſhortneſſe of breath.

Lambes Lettice is wholly ſpent for ſallets, in the beginning of the yeare, as I ſaid, before any almoſt of the other ſorts of Lettice are to be had.

## Chap. XXXI.

*Portulaca.* Purſlane.

PVrſlane hath many thicke round ſhining red ſtalkes, full of iuice, lying vpon the ground for the moſt part ; whereon are ſet diuers long, thicke, pale green leaues, ſometimes alone by themſelues, and ſometimes many ſmall ones together with them ; among which grow ſmall yellow flowers, which ſtand in little greene huskes, containing blacke ſeede : the roote is ſmall, and periſheth euery yeare, and muſt be new ſowen in Aprill, in the alleyes of the Garden betweene the beds, as ſome haue heretofore vſed, where it may haue the more moiſture, or, as I haue ſeene in ſome Gardens, vpon thoſe beds of dung that Gardiners haue vſed to nourſe vp their Cowcumbers, Melons, and Pompions, whereon after they haue beene taken away, they haue ſowen Purſlane, where if it be much watered, the warmth of the dung, and the water giuen it, the Purſlane hath grown great and large, and continued vntill winter.

### The Vſe of Purſlane.

It is vſed as Lettice in ſallets, to coole hot and faint ſtomackes in the hot time of the yeare, but afterwards if only for delight, it is not good to bee too prodigall in the vſe thereof.

The ſeede of Purſlane doth coole much any inflammation inward or outward, and doth a little binde withall.

## Chap. XXXII.

*Dracoherba siue Tarchon & Dracunculus hortensis.* Tarragon.

TArragon hath long and narrow darke greene leaues, growing on slender and brittle round stalkes, two or three foote high, at the tops whereof grow forth long slender spikes of small yellowish flowers, which seldome giue any good seede, but a dustie or chaffie matter, which flieth away with the winde: the roote is white, and creepeth about vnder ground, whereby it much encreaseth: the whole herbe is of a hot and biting taste.

### The Vse of Tarragon.

It is altogether vsed among other cold herbes, to temper their coldnesse, and they to temper its heate, so to giue the better relish vnto the Sallet; but many doe not like the taste thereof, and so refuse it.

There are some Authors that haue held Tarragon not to be an herbe of it owne kinde, but that it was first produced, by putting the seede of Lin or Flaxe into the roote of an Onion, being opened and so set into the ground, which when it hath sprung, hath brought forth this herbe Tarragon, which absurd and idle opinion, Matthiolus by certaine experience saith, hath been found false.

## Chap. XXXIII.

*Nasturtium hortense.* Garden Cresses.

GArden Cresses growe vp to the height of two foote or thereabouts, hauing many small, whitish, broad, endented, torne leaues, set together vpon a middle ribbe next the ground, but those that growe higher vpon the stalkes are smaller and longer: the tops of the stalkes are stored with white flowers, which turne into flat pods or pouches, like vnto Shepheard purse, wherein is contained flat reddish seede: the roote perisheth euery yeare: the taste both of leaues and seedes are somewhat strong, hot, and bitter.

### The Vse of Cresses.

The Dutchmen and others vse to eate Cresses familiarly with their butter and bread, as also stewed or boyled, either alone or with other herbes, whereof they make a Horch potch, and so eate it. Wee doe eate it mixed among Lettice or Purslane, and sometimes with Tarragon or Rocket, with oyle and vinegar and a little salt, and in that manner it is very sauoury to some mens stomackes.

The vse of Cresses physically is, it helpeth to expectorate tough flegme, as also for the paines of the breast, and as it is thought taketh away spots, being laid to with vinegar. The seede is giuen of many to children for the wormes.

Chap.

1 *Portulaca.* Purſlane. 2 *Dracho herba ſeu Tarchon.* Tarragon. 3 *Eruca ſatiua.* Garden Rocket. 4 *Naſturtium ſatiuum.* Garden Creſſes.
5 *Sinapi.* Muſtard. 6 *Aſparagus.* Aſparagus or Sperage.

## Chap. XXXIIII.

### *Eruca ſatiua.* Garden Rocket.

OVr Garden Rocket is but a wilde kinde brought into Gardens; for the true Romane Rocket hath larger leaues; this hath many long leaues, much torne or rent on the edges, ſmaller and narrower then the Romane kinde: the flowers hereof are of a pale yellowiſh colour, whereas the true is whitiſh, conſiſting of foure leaues: the ſeede of this is reddiſh, contained in ſmaller and longer pods then the true, which are ſhorter and thicker, and the ſeede of a whitiſh yellow colour: the rootes of both periſh as ſoone as they haue giuen ſeede. Some haue taken one ſort of the wilde kinde for Muſtard, and haue vſed the ſeede for the ſame purpoſe.

### The Vſe of Rocket.

It is for the moſt part eaten with Lettice, Purſlane, or ſuch cold herbes, and not alone, becauſe of its heate and ſtrength; but that with the white ſeede is milder. The ſeede of Rocket is good to prouoke vrine, and to ſtirre vp bodily luſt.

The ſeede bruiſed, and mixed with a little vinegar, and of the gall of an Oxe, cleanſeth the face of freckles, ſpots, and blew markes, that come by beatings, fals, or otherwaies.

Matthiolus ſaith, that the leanes boyled, and giuen with ſome Sugar to little children, cureth them of the cough.

The ſeede is held to be helpfull to ſplenetick perſons; as alſo to kill the wormes of the belly.

## Chap. XXXV.

### *Sinapi ſatiuum.* Garden Muſtard.

THe Muſtard that is moſt vſuall in this Country, howſoeuer diuers doe for their priuate vſes ſowe it in their Gardens or Orchards, in ſome conuenient corner, yet the ſame is found wilde alſo abroad in many places. It hath many rough long diuided leaues, of an ouerworne greene colour: the ſtalke is diuided at the toppe into diuers branches, whereon growe diuers pale yellow flowers, in a great length, which turne into ſmall long pods, wherein is contained blackiſh ſeede, inclining to redneſſe, of a fiery ſharpe taſte: the roote is tough and white, running deepe into the ground, with many ſmall fibres at it.

### The Vſe of Muſtard.

The ſeede hereof grownd between two ſtones, fitted for the purpoſe, and called a Querne, with ſome good vinegar added vnto it, to make it liquid and running, is that kinde of Muſtard that is vſually made of all ſorts, to ſerue as ſawce both for fiſh and fleſh.

The ſame liquid Muſtard is of good vſe, being freſh, for Epilepticke perſons, to warme and quicken thoſe dull ſpirits that are ſopite and ſcarce appeare, if it be applyed both inwardly and outwardly.

It is with good ſucceſſe alſo giuen to thoſe that haue ſhort breathes, and troubled with a cough in the lungs.

## Chap. XXXVI.

### *Asparagus.* Sperage or Asparagus.

ASparagus riseth vp at the first with diuers whitish greene scaly heads, very brittle or easie to breake while they are young, which afterwards rise vp into very long and slender greene stalkes, of the bignesse of an ordinary riding wand at the bottome of most, or bigger or lesser, as the rootes are of growth, on which are set diuers branches of greene leaues, shorter and smaller then Fennell vp to the toppe, at the ioynts whereof come forth small mossie yellowish flowers, which turne into round berries, greene at the first, and of an excellent red colour when they are ripe, shewing as if they were beades of Corrall, wherein are contained exceeding hard and blacke seede: the rootes are dispersed from a spongious head into many long, thicke, and round strings, whereby it sucketh much nourishment out of the ground, and encreaseth plentifully thereby.

We haue another kinde hereof that is of much greater account, because the shootes are larger, whiter, and being dressed taste more sweete and pleasant, without any other difference.

### The Vse of Asparagus.

The first shootes or heads of Asparagus are a Sallet of as much esteeme with all sorts of persons, as any other whatsoeuer, being boyled tender, and eaten with butter, vinegar, and pepper, or oyle and vinegar, or as euery ones manner doth please; and are almost wholly spent for the pleasure of the pallate. It is specially good to prouoke vrine, and for those that are troubled with the stone or grauell in the reines or kidneyes, because it doth a little open and cleanse those parts.

## Chap. XXXVII.

### *Brassica.* Cabbages and Coleworts.

THere is greater diuersity in the forme and colour of the leaues of this plant, then there is in any other that I know groweth vpon the ground. But this place requireth not the knowledge of all sorts which might be shewen, many of them being of no vse with vs for the table, but for delight, to behold the wonderfull variety of the workes of God herein. I will here therefore shew you onely those sorts that are ordinary in most Gardens, and some that are rare, receiued into some especiall Gardens: And first of Cabbages, and then of Coleworts.

Our ordinary Cabbage that closeth hard and round, hath at the first great large thicke leaues, of a grayish greene colour, with thicke great ribbes, and lye open most part of the Summer without closing, but toward the end of Summer, being growne to haue many leaues, it then beginneth to growe close and round in the middle, and as it closeth, the leaues growe white inward; yet there be some kindes that will neuer be so close as these, but will remaine halfe open, which wee doe not account to be so good as the other: in the middle of this head, the next yeare after the sowing, in other Countries especially, and sometimes in ours, if the Winter be milde, as may be seene in diuers Gardens (but to preuent the danger of our Winter frosts, our Gardiners now doe vse to take vp diuers Cabbages with their rootes, and tying a cloth or some such thing about the rootes, doe hang them vp in their houses, where they may be defended from cold, and then set them againe after the frosts are past) and then there shooteth out a great thicke stalke, diuided at the toppe into many branches, bearing thereon diuers small flowers, sometime white, but most commonly yellow, made of foure leaues, which turne into long, round, and pointed pods, containing therein small

round

round ſeede, like vnto Turnep ſeede : the roote ſpreadeth not farre nor deepe, and dyeth vſually in any great froſte ; for a ſmall froſt maketh the Cabbage eate the tenderer.

The red Cabbage is like vnto the white, laſt ſpoken of, but differing in colour and greatneſſe ; for it is ſeldome found ſo great as the white, and the colour of the leaues is very variable, as being in ſome ſtript with red, in others more red, or very deepe red or purple.

The ſugar loafe Cabbage, ſo called becauſe it is ſmaller at the toppe then it is at the bottome, and is of two ſorts, the one white, the other greene.

The Sauoy Cabbadge, one is of a deepe greene coloured leafe, and curld when it is to be gathered ; the other is yellowiſh : neyther of both theſe doe cloſe ſo well as the firſt, but yet are vſed of ſome, and accounted good.

The Cole flower is a kinde of Coleworte, whoſe leaues are large, and like the Cabbage leaues, but ſomewhat ſmaller, and endented about the edges, in the middle whereof, ſometimes in the beginning of Autumne, and ſometimes much ſooner, there appeareth a hard head of whitiſh yellow tufts of flowers, cloſely thruſt together, but neuer open, nor ſpreading much with vs, which then is fitteſt to be vſed, the green leaues being cut away cloſe to the head : this hath a much pleaſanter taſte then eyther the Coleworte, or Cabbage of any kinde, and is therefore of the more regard and reſpect at good mens tables.

The ordinary Coleworte is ſufficiently knowne not to cloſe or cabbage, and giueth ſeede plentifully enough.

The other Colewortes that are nourſed vp with thoſe that delight in curioſities, beſides the aforeſaid ordinary greene, which is much vſed of Dutchmen, and other ſtrangers, are theſe : The Curld Coleworte eyther wholly of a greene colour, or of diuers colours in one plant, as white, yellow, red, purple or crimſon, ſo variably mixed, the leaues being curld on the edges, like a ruffe band, that it is very beautifull to behold.

There is alſo another curld Colewort of leſſe beauty and reſpect, being but a little curld on the edges, whoſe leaues are white, edged with red, or green edged with white.

Two other there are, the one of a popingaye greene colour : the other of a fine deepe greene, like vnto the Sauoyes.

Then there is the Cole rape, which is alſo a kinde of Coleworte, that beareth a white heade, or headed ſtalke aboue the ground, as bigge as a reaſonable Turnep, but longer, and from the toppe thereof ſpringeth out diuers great leaues, like vnto Colewortes ; among which riſe diuers ſtalkes that beare yellow flowers, and ſeede in pods, almoſt as ſmall as Muſtard ſeede : the roote is ſomewhat long, and very buſhie with threds.

## The Vſe of Cabbages and Colewortes.

They are moſt vſually boyled in poudered beefe broth vntil they be tender, and then eaten with much fat put among them.

The great ribs of the Popingay, and deepe greene Colewortes, beeing boyled and layde into diſhes, are ſerued to the table with oyle and vinegar in the Lent time for very good ſallets.

In the cold Countries of Ruſſia and Muſcouia, they pouder vp a number of Cabbages, which ſerue them, eſpecially the poorer ſort, for their moſt ordinary foode in winter ; and although they ſtinke moſt grieuouſly, yet to them they are accounted good meate.

It is thought, that the vſe of them doth hinder the milke in Nurſes breaſts, cauſing it to dry vp quickely : but many women that haue giuen ſucke to my knowledge haue denyed that aſſertion, affirming that they haue often eaten them, and found no ſuch effect. How it might proue in more delicate bodies then theirs that thus ſaid, I cannot tell : but Matthiolus auerreth it to encreaſe milke in Nurſes breaſtes ; ſo differing are the opinions of many. The ſeede groſſely bruiſed and boyled a little in fleſh broth, is a preſent remedie for the Collicke ; the ſeede and the broth being taken together, eaſing them that are troubled therewith of all griping paines : as alſo for the ſtone in the kidneyes. A Lohoc or licking Electuary made of the pulpe of

the

1 *Brassica capitata.* Close Cabbage. 2 *Brassica patula.* Open Cabbage. 3 *Brassica Sabaudica crispa.* Curld Sauoye Co ewort. 4 *Caulis floride.* Cole flower. 5 *Caulis crispa.* Curld Co'ewort. 6 *Caulis crispa varieta.* Changeable curld Colewort. 7 *Rapocaulis.* Cole rape,

the boyled ſtalkes, and a little honey and Almond milke, is very profitable for ſhortneſſe of breath, and thoſe that are entring into a Conſumption of the lunges. It hath beene formerly held to be helpefull in all diſeaſes: for Criſippus, an ancient Phyſitian, wrote a whole Volume of the vertues, applying it to all the parts of the body: which thing neede not ſeeme wonderfull, in that it is recorded by writers, that the old Romanes hauing expelled Phyſitians out of their Common-wealth, did for many hundred of yeares maintaine their health by the vſe of Cabbages, taking them for euery diſeaſe.

## Chap. XXXVIII.

### *Siſarum.* Skirrets.

After all the herbes before rehearſed, fit for ſallets, or otherwiſe to bee eaten, there muſt follow ſuch rootes as are vſed to the ſame purpoſe: and firſt, Skirrets haue many leaues next the ground, compoſed of many ſmall ſmooth green leaues, ſet each againſt other vpon a middle ribbe, and euery one ſnipt about the edges: the ſtalke riſeth vp two or three foote high, ſet with the like leaues, hauing at the toppe ſpoakie tufts of white flowers, which turne into ſmall ſeede, ſomewhat bigger and darker then Parſley ſeede: the rootes be many growing together at one head, beeing long, ſlender, & rugged or vneuen, of a whitiſh colour on the outſide, and more white within, hauing in the middle of the roote a long ſmall hard pith or ſtring: theſe heads are vſually taken vp in February and March, or ſooner if any ſo pleaſe, the greater number of them being broken off to bee vſed, the reſt are planted againe after the heads are ſeparated, and hereby they are encreaſed euery yeare by many; but it is now adayes more ſowen of the ſeed, which come forwards well enough if the ground be fat and good.

### The Vſe of Skirrets.

The rootes being boyled, peeled and pithed, are ſtewed with butter, pepper and ſalt, and ſo eaten; or as others vſe them, to roule them in flower, and fry them with butter, after they haue beene boyled, peeled and pithed: each way, or any way that men pleaſe to vſe them, they may finde their taſte to be very pleaſant, far beyond any Parſnep, as all agree that taſte them.

Some doe vſe alſo to eate them as a ſallet, colde with vinegar, oyle, &c. being firſt boyled and dreſſed as before ſaid. They doe helpe to prouoke vrine, and as is thought, to procure bodily luſt, in that they are a little windy.

## Chap. XXXIX.

### *Paſtinaca ſatiua latifolia.* Parſneps.

The common garden Parſnep hath diuers large winged leaues lying vpon the ground, that is, many leaues ſet one by another on both ſides of a middle ſtalk, ſomewhat like as the Skirret hath, but much larger, and cloſer ſet: the ſtalke riſeth vp great and tall, fiue or ſix foot high ſomtimes, with many ſuch leaues thereon at ſeuerall ioynts; the top whereof is ſpread into diuers branches, whereon ſtand ſpoakie rundles of yellow flowers, which turne into browniſh flat ſeede: the root is long, great and white, very pleaſant to bee eaten, and the more pleaſant if it grow in a fat ſandy ſoyle.

There is another ſort of garden Parſnep, called the Pine Parſnep, that is not common in euery Garden, and differeth from the former in three notable parts. The root is not ſo long, but thicker at the head and ſmaller below; the ſtalke is neither ſo bigge,

nor

1 *Sisarum.* Skirrits. 2 *Pastinaca latifolia.* Parsneps. 3 *Pastinaca tenuifolia.* Carrets. 4 *Napum.* Turneps. 5 *Napus sativus.* Navewes.
6 *Raphanus niger.* Blacke Raddish. 7 *Raphanus vulgaris.* Common Raddish.

nor ſo high; and the ſeede is ſmaller: yet as Iohn Tradeſcante ſaith (who hath giuen me the relation of this, and many other of theſe garden plants, to whom euery one is a debtor) the roote hereof is not altogether ſo pleaſant as the other.

Moreouer the wilde kinde, which groweth in many places of England (and wherof in ſome places there might be gathered a quarter ſacke full of the ſeede) if it be ſowen in Gardens, and there well ordered, will proue as good as the former kinde of Garden Parſneps.

### The Vſe of Parſneps.

The Parſnep root is a great nouriſher, and is much more vſed in the time of Lent, being boyled and ſtewed with butter, then in any other time of the yeare; yet it is very good all the winter long. The ſeede helpeth to diſſolue winde, and to prouoke vrine.

## Chap. XL.

### *Paſtina ſatiua tenuifolia.* Carrots.

THe Carrot hath many winged leaues, riſing from the head of the roote, which are much cut and diuided into many other leaues, and they alſo cut and diuided into many parts, of a deepe greene colour, ſome whereof in Autumne will turne to be of a fine red or purple (the beautie whereof allureth many Gentlewomen oftentimes to gather the leaues, and ſticke them in their hats or heads, or pin them on their armes in ſtead of feathers): the ſtalke riſeth vp among the leaues, bearing many likewiſe vpon it, but nothing ſo high as the Parſnep, being about three foote high, bearing many ſpoakie tufts of white flowers, which turne into ſmall rough ſeede, as if it were hairy, ſmelling reaſonable well if it bee rubbed: the roote is round and long, thicke aboue and ſmall below, eyther red or yellow, eyther ſhorter or longer, according to his kinde; for there is one kinde, whoſe roote is wholly red quite thorough-out; another whoſe roote is red without for a pretty way inward, but the middle is yellow.

Then there is the yellow, which is of two ſorts, both long and ſhort: One of the long yellow ſorts, which is of a pale yellow, hath the greateſt and longeſt roote, and likewiſe the greateſt head of greene, and is for the moſt part the worſt, being ſpongy, and not firme.

The other is of a deepe gold yellow colour, and is the beſt, hauing a ſmaller head, or tuft of greene leaues vpon it.

The ſhorte rootes are likewiſe diſtinguiſhed, into pale and deepe yellow colours.

### The Vſe of Carrots.

All theſe ſorts being boyled in the broth of beefe, eyther freſh or ſalt, but more vſually of ſalted beefe, are eaten with great pleaſure, becauſe of the ſweeteneſſe of them: but they nouriſh leſſe then Parſneps or Skirrets.

I haue not often knowne the ſeede of this Garden kinde to bee vſed in Phyſicke: but the wilde kinde is often and much vſed to expell winde, &c.

## Chap. XLI.

### *Rapum hortenſe.* Turneps.

THere are diuers ſorts of Turneps, as white, yellow, and red: the white are the moſt common, and they are of two kinds, the one much ſweeter then the other.

The yellow and the red are more rare, and nourſed vp only by thoſe that are curious: as alſo the Nauewe, which is ſeene but with very few.

The

The ordinary Garden Turnep hath many large, and long rough greene leaues, with deepe and vneuen gashes on both sides of them: the stalke riseth vp among the leaues about two foote high, spread at the toppe into many branches, bearing theron yellow flowers, which turne into long pods, with blackish round seede in them: the roote is round and white, some greater, some smaller; the best kinde is knowne to be flat, with a small pigges tale-like roote vnderneath it; the worser kinde which is more common in many places of this land, both North and West, is round, and not flat, with a greater pigges tayle-like roote vnderneath.

The yellow kinde doth often grow very great, it is hardly discerned from the ordinary kinde while it groweth, but by the greatnesse and spreading of the leaues beeing boyled, the roote changeth more yellow, somewhat neare the colour of a Carrot.

The red Turnep groweth vsually greater then any of the other, especially in a good ground, being of a faire red colour on the outside, but being pared, as white as any other on the inside. This, as Matthiolus saith, doth grow in the Countrey of Anania, where hee hath seene an infinite number of them that haue waighed fifty pound a peece, and in some places hee saith, a hundred pound a peece, both which we would thinke to be incredible, but that we see the kind is greatly giuen to grow, and in warme Countries they may so thriue, that the bulke or bignesse of the roote may so farre passe the growth of our Countrey, as that it may rise to that quantity aboue specified.

The Navew gentle is of two kindes, a smaller and a greater; the smaller is vsually called in France, *Naveau de Caue*, the roote is somewhat long with the roundnesse; this kinde is twice as bigge as a mans thumbe, and many of them lesse: The other is long and great, almost as big as the short Carrot, but for the most part of an vneuen length, and roundnesse vnto the very end, where it spreadeth into diuers small long fibres: neyther of them doth differ much from the Turnep, in leafe, flower or seed.

### The Vse of Turneps.

Being boyled in salt broth, they all of them eate most kindly, and by reason of their sweetnesse are much esteemed, and often seene as a dish at good mens tables: but the greater quantitie of them are spent at poore mens feasts. They nourish much, and engender moist and loose flesh, and are very windy. The seede of the Navew gentle is (as I take it) called of Andromachus in the composition of his Treakle, *Bunias dulcis*: for Dioscorides and Plinie doe both say, that the seede of the tame Bunias or Napus is put into Antidotes, and not the seede of the wilde, which is more sharpe and bitter; neyther the seede of the Turnep, which is called in Greeke γογγύλη in Latine *Rapum*, because the seede is not sweete.

## Chap. XLII.

### *Raphanus.* Raddish.

THere are two principall kindes of Garden Raddish, the one is blackish on the outside, and the other white; and of both these there is some diuision againe, as shall be shewed. Dittander and horse Raddish be reckoned kinds thereof.

The ordinary Raddish hath long leaues, vneuenly gashed on both sides, the stalke riseth vp to the height of three or foure foote, bearing many purplish flowers at the top, made of foure leaues a peece, which turne into thicke and short pods, wherein are contained round seede, greater then Turnep or Coleworte seede, and of a pale reddish colour: the roote is long, white, and of a reddish purple colour on the outside toward the toppe of it, and of a sharpe biting taste.

There is a small kind of Raddish that commeth earlier then the former, that we haue had out of the low Countries, not differing in any thing else.

The blacke Raddish I haue had brought me out of the lowe Countries, where they sell them in some places by the pound, and is accounted with them a rare winter sallet:

the roote of the beſt kinde is blackiſh on the outſide (and yet the ſeede gathered from ſuch an one, hath after the ſowing againe, giuen rootes, whereof ſome haue beene blacke, but the moſt part white on the outſide) and white within, great and round at the head, almoſt like a Turnep, but ending ſhorter then a Raddiſh, and longer then a Turnep, almoſt peare-faſhion, of a firmer and harder ſubſtance then the ordinary Raddiſh, but no leſſe ſharpe and biting, and ſomewhat ſtrong withall; the leaues are ſomewhat ſmaller, and with deeper gaſhes, the flower and ſeede are like the former, but ſmaller.

Another ſort of blacke Raddiſh is like in leafe and ſeede to the former, but the flower is of a lighter purple colour: the roote is longer and ſmaller, and changeth alſo to bee white as the former doth, ſo that I thinke they haue both riſen from one kinde.

The Horſe Raddiſh is a kinde of wilde Raddiſh, but brought into Gardens for the vſe of it, and hath great large and long greene leaues, which are not ſo much diuided, but dented about the edges: the roote is long and great, much ſtronger in taſte then the former, and abideth diuers yeares, ſpreading with branches vnder ground.

Dittander is likewiſe a wilde kinde hereof, hauing long pointed blewiſh greene leaues, and a roote that creepeth much vnder ground: I confeſſe this might haue bin placed among the herbes, becauſe the leaues and not the rootes are vſed; but let it paſſe now with the kindes of Raddiſh.

### The Vſe of theſe Raddiſhes.

Raddiſhes doe ſerue vſually as a *ſtimulum* before meat, giuing an appetite thereunto; the poore eate them alone with bread and ſalt. Some that are early ſowen, are eaten in Aprill, or ſooner if the ſeaſon permit; others come later; and ſome are ſowen late to ſerue for the end of Summer: but (as of all things elſe) the earlier are the more accepted.

The blacke Raddiſhes are moſt vſed in the winter, (yet ſome in their naturall and not forc'd grounds, haue their rootes good moſt part of the Summer) and therefore muſt bee ſowen after Midſomer; for if they ſhould bee ſowen earlier, they would preſently runne vp to ſtalke and ſeed, and ſo loſe the benefit of the roote. The Phyſicall propertie is, it is often vſed in medicines that helpe to breake the ſtone, and to auoyde grauell.

The Horſe Raddiſh is vſed Phyſically, very much in Melancholicke, Spleneticke and Scorbuticke diſeaſes. And ſome vſe to make a kinde of Muſtard with the rootes, and eate it with fiſh.

Dittander or Pepperworte is vſed of ſome cold churliſh ſtomackes, as a ſawce or ſallet ſometimes to their meate, but it is too hot, bitter and ſtrong for weake and tender ſtomackes.

Our Gardiners about London vſe great fences of reede tyed together, which ſeemeth to bee a mat ſet vpright, and is as good as a wall to defend the cold from thoſe things that would be defended, and to bring them forwards the earlier.

## Chap. XLIII.

### *Cepæ.* Onions.

WEe haue diuers ſorts of Onions, both white and red, flat, round and long, as ſhall be preſently ſhewed: but I will doe with theſe as I doe with the reſt, only giue you one deſcription for them all, and afterwards their ſeuerall names and varieties, as they are to be known by.

Our common Garden Onion hath diuers long greene hollow leaues, ſeeming halfe flat; among which riſeth vp a great round hollow ſtalke, bigger in the middle then any where elſe, at the toppe whereof ſtandeth a cloſe round head, couered at the firſt with a thin skinne, which breaketh when the head is growne, and ſheweth forth a great vm-

bell

1 *Raphanus rusticanus.* Horse Raddish. 2 *Lepidium siue Piperitis.* Dittander. 3 *Cepæ rotundæ.* Round Onions. 4 *Cepæ longæ.* Long Onions. 5 *Porrum.* Leekes 6 *Allium.* Garlicke. 7 *Rapunculus.* Rampions. 8 *Tragopogon.* Goates beard.

bell of white flowers, which turne into blacke seede : but then the head is so heauie that the stalke cannot sustaine it, but must be vpheld from falling to the ground, lest it rot and perish : the roote as all know is round, in some greater, in others lesser, or flat, in some red on the outside only, in others quite thorough out, in some white, and very sharpe and strong, in others milder, and more pleasant, and some so pleasant that they may be eaten as an Apple : All these kindes of Onions, contrary to the nature of all other bulbous rootes, haue no off-set, or other roote growing to it, but are euery one alone single by themselues ; and therefore it seemeth, the Latines, as Columella recordeth, haue giuen it the name *Vnio*, and the French it should seeme following the Latine, and the English the French, do call it *Oignon* and *Onion*, as an vnite, or as if they were but one and one, and dye euery yeare after seed bearing.

The red flat kinde is most vsually with vs the strongest of them all, yet I haue had a great red Onion brought mee from beyond Sea, that was as great almost as two mens fistes, flat and red quite thoroughout, and very pleasant both to smell vnto, and to eate, but did quickly degenerate; so that we plainly see, that the soyle and climate doth giue great alteration to plants of all sorts.

The long kinde wee call St. Omers Onions, and corruptly among the vulgar, St. Thomas Onions.

The other red kinde we call Strasborough Onions, whose outside onely is red, and are very sharpe and fierce.

The white Onions both long and flat, are like vnto Chalke-stones lying vpon the ground, when they are ripe and fit to be gathered.

And lastly, there is the Spanish Onion, both long and flat, very sweete, and eaten by many like an apple, but as Iohn Tradescante saith, who hath beene in Spaine, that the Spaniards themselues doe not eate them so familiarly, as they doe those white Onions that come out of our owne Countrey, which they haue there more plentifully then their sweete Onions.

### The Vse of Onions.

Onions are vsed many wayes, as sliced and put into pottage, or boyled and peeled and layde in dishes for sallets at supper, or sliced and put into water, for a sawce for mutton or oysters, or into meate roasted being stuffed with Parsly, and so many waies that I cannot recount them, euery one pleasing themselues, according to their order, manner or delight.

The iuice of Onions is much vsed to be applyed to any burnings with fire, or with Gun-pouder, or to any scaldings with water or oyle, and is most familiar for the Country, where vpon such sudden occasions they haue not a more fit or speedy remedie at hand : The strong smell of Onions, and so also of Garlicke and Leekes, is quite taken away from offending the head or eyes, by the eating of Parsley leaues after them.

## Chap. XLIIII.

### *Porrum.* Leekes.

THere be likewise sundry sorts of Leekes, both great and small. Leekes are very like vnto Onions, hauing long green hollow-like leaues, flattish on the one side, and with a ridge or crest on the backe side : if they bee suffered to grow vncut, then in the second or third yeare after the sowing, they will send forth a round and slender stalke, euen quite thoroughout, and not swollen or bigger in the middle like the Onion, bearing at the toppe a head of purplish flowers, and blacke seede after them, very like vnto Onion seede, that it is hard to distinguish them : the root is long and white, with a great bush of fibres hanging at it, which they call the beards.

The vnset Leeke hath longer and slenderer roots then the other, which being transplanted, groweth thicker and greater.

The

The French Leeke, which is called the Vine Leeke, is the beſt of all others.
Our common kinde is of two ſorts, one greater then another.
Another ſort encreaſeth altogether by the roote, as Garlicke doth.
And then Ciues, which are the ſmalleſt, and encreaſe aboundantly only by the roor.

Some doe account Scalions to be rather a kinde of Onions then Leekes, and call them *Cepa Aſcalonica*, or *Aſcalonitides*, which will quickly ſpend it ſelfe, if it be ſuffered to be vncut; but all Authors affirme, that there is no wilde kinde of Onion, vnleſſe they would haue it to be *Gethyum*, whereof Theophraſtus maketh mention, ſaying, that it hath a long necke (and ſo theſe Scalions haue) and was alſo of ſome called *Gethyllides*, which antiquity accounted to be dedicated to Latona, the mother of Apollo, becauſe when ſhe was bigge with childe of Apollo, ſhe longed for theſe Leekes.

## The Vſe of Leekes.

The old World, as wee finde in Scripture, in the time of the children of Iſraels being in Egypt, and no doubt long before, fed much vpon Leekes, Onions, and Garlicke boyled with fleſh; and the antiquity of the Gentiles relate the ſame manner of feeding on them, to be in all Countries the like, which howſoeuer our dainty age now refuſeth wholly, in all ſorts except the pooreſt; yet Muſcouia and Ruſſia vſe them, and the Turkes to this day, (as Bellonius writeth) obſerue to haue them among their diſhes at their tables, yea although they be *Baſhas*, *Cades*, or *Vaiuodas*, that is to ſay, Lords, Iudges, or Gouernours of countries and places. They are vſed with vs alſo ſometimes in Lent to make pottage, and is a great and generall feeding in Wales with the vulgar Gentlemen.

Onions boyled or roſted vnder the embers, and mixed with ſugar and butter, are good for thoſe that are troubled with coughes, ſhortneſſe of breath, and wheeſing. An Onion made hollow at the bottome, and ſome good Treakle put into it, with a little iuyce of Cirrons (or Lemons in the ſtead thereof) being well baked together vnder the embers, after the hole is ſtopped againe, and then ſtrained forth, and giuen to one that hath the plague, is very helpefull, ſo as hee be laid to ſweate vpon it.

Ciues are vſed as well to be ſhred among other herbes for the pot, as to be put into a Sallet among other herbs, to giue it a quicker relliſh.

Leekes are held to free the cheſt and lungs from much corruption and rotten flegme, that ſticketh faſt therein, and hard to be auoided, as alſo for them that through hoarſeneſſe haue loſt their voice, if they be eyther taken rawe, or boyled with broth of barley, or ſome ſuch other ſupping, fit and conducing thereunto. And baked vnder hot embers is a remedy againſt a ſurfeit of Muſhromes.

The greene blades of Leekes being boyled and applyed warme to the *Hemorrhoides* or piles, when they are ſwolne and painfull, giue a great deale of eaſe.

# Chap. XLV.

## *Allium*. Garlicke.

IHaue ſpoken of diuers ſorts of Garlicke called Moly, in the former booke; I ſhall needе in this place to ſhew onely thoſe kindes, that this Garden nourſeth vp, and leaue the reſt to his fit time and place.

Garlicke hath many long greene leaues, like vnto Onions, but much larger, and not hollow at all as Onions are: the ſtalke riſeth vp to be about three foote high, bearing ſuch a head at the toppe thereof as Onions and Leekes doe, with purpliſh flowers, and blacke ſeede like Leekes: the roote is white within, couered ouer with many purpliſh

pliſh skins, and is diuided into many parts or cloues, which ſerue both to ſet againe for increaſe, and alſo to vſe as neede ſhall require, and is of a very ſtrong ſmell and taſte, as euery one knoweth, paſſing either Onions or Leekes, but exceeding wholſome withall for them that can take it.

*Allium Vrſinum.* Ramſons.

Ramſons are another kinde of Garlicke, and hath two or three faire broad leaues, of a freſh or light greene colour, pointed at the end: the ſtalke groweth about an hand length high, bearing many ſmall and pure white ſtarre-like flowers at the toppe, and afterwards ſmall, blacke, and ſmooth round ſeede: the roote is alſo diuided into many parts, whereby it is much encreaſed, and is much milder then the former, both in ſmell and taſte.

### The Vſe of Garlicke.

It being well boyled in ſalt broth, is often eaten of them that haue ſtrong ſtomackes, but will not brooke in a weake and tender ſtomacke.

It is accounted, and ſo called in diuers Countries, The poore mans Treakle, that is, a remedy for all diſeaſes. It is neuer eaten rawe of any man that I know, as other of the rootes aforeſaid, but ſodden alwaies and ſo taken.

Ramſons are oftentimes eaten with bread and butter, and otherwiſe alſo, as euery mans affection and courſe of life leadeth him to vſe.

## Chap. XLVI.

*Rapunculus ſiue Rapuntium.* Rampions.

Garden Rampions are of two ſorts, the one greater, the other leſſer: the leaues of Rampions are in the one ſomewhat broad like a Beete, in the other ſomewhat long and narrow, and a little broader at the end, of a light greene colour, lying flat vpon the ground all the firſt winter, or yeare of the ſpringing, and the next Spring ſhooteth forth ſtalkes two or three foote high, bearing at the toppe, in the bigger ſort, a long ſlender ſpike of ſmall horned or crooked flowers, which open their brimmes into foure leaues; in the leſſer many ſmall purpliſh bels, ſtanding vpon ſeuerall ſmall foote-ſtalkes, which turne into heads, bearing ſmall blackiſh ſeede: the root is white, branched into two or three rootes, of the bigneſſe and length of a mans finger or thumbe.

### The Vſe of Rampions.

The rootes of both are vſed for Sallets, being boyled, and then eaten with oyle and vinegar, a little ſalt and pepper.

## Chap. XLVII.

*Tragopogon.* Goates beard.

Goates beard hath many long and narrow leaues, broader at the bottome, and ſharper at the end, with a ridge downe the backe of the leafe, and of a pale greene colour; among which riſeth vp a ſtalke of two or three foote high, ſmooth and hollow, bearing thereon many ſuch like leaues, but ſmaller and ſhorter, and at the toppe thereof on euery branch a great double yellow flower, like almoſt vnto the flower of a Dandelion, which turneth into a head, ſtored with doune, and long whitiſh ſeede therein, hauing on the head of euery one ſome part of the doune,

and

and is carried away with the winde if it bee neglected: the roote is long and round, somewhat like vnto a Parsnep, but farre smaller, blackish on the outside, and white within, yeelding a milkie iuyce being broken, as all the rest of the plant doth, and of a very good and pleasant taste. This kinde, as also another with narrower leaues, almost like grasse, growe wilde abroad in many places, but are brought into diuers Gardens. The other two kindes formerly described in the first part, the one with a purple flower, and the other with an ash-coloured, haue such rootes as these here described, and may serue also to the same purpose, being of equall goodnesse, if any will vse them in the same manner; that is, while they are young, and of the first yeares sowing, else they all growe hard, in running vp to seede.

### The Vse of Goates beard.

If the rootes of any of these kindes being young, be boyled and dressed as a Parsnep, they make a pleasant dish of meate, farre passing the Parsnep in many mens iudgements, and that with yellow flowers to be the best.

They are of excellent vse being in this manner prepared, or after any other fit and conuenient way, to strengthen those that are macilent, or growing into any consumption.

## Chap. XLVIII.

### *Carum.* Carawayes.

CArawayes hath many very fine cut and diuided leaues lying on the ground, being alwaies greene, somewhat resembling the leaues of Carrots, but thinner, and more finely cut, of a quicke, hot, and spicie taste: the stalke riseth not much higher then the Carrot stalke, bearing some leaues at the ioynts along the stalke to the toppe, where it brancheth into three or foure parts, bearing spoakie vmbels of white flowers, which turne into small blackish seede, smaller then Aniseede, and of a hotter and quicker taste: the roote is whitish, like vnto a Parsnep, but much smaller, more spreading vnder ground, and a little quicke in taste, as all the rest of the plant is, and abideth long after it hath giuen seede.

### The Vse of Carawayes.

The rootes of Carawayes being boyled may be eaten as Carrots, and by reason of the spicie taste doth warme and comfort a cold weake stomacke, helping to dissolue winde (whereas Carrots engender it) and to prouoke vrine, and is a very welcome and delightfull dish to a great many, yet they are somewhat stronger in taste then Parsneps.

The seede is much vsed to bee put among baked fruit, or into bread, cakes, &c. to giue them a relish, and to helpe to digest winde in them are subiect thereunto.

It is also made into Comfits, and put into *Trageas*, or as we call them in English, Dredges, that are taken for the cold and winde in the body, as also are serued to the table with fruit.

Chap.

## Chap. XLIX.

### *Pappas sive Battatas.* Potatoes.

THree sorts of Potatoes are well knowne vnto vs, but the fourth I rest doubtfull of, and dare not affirme it vpon such termes as are giuen vnto it, vntill I may be better informed by mine owne sight.

The Spanish kinde hath (in the Islands where they growe, either naturally, or planted for increase, profit, and vse of the Spaniards that nourse them) many firme and verie sweete rootes, like in shape and forme vnto Asphodill rootes, but much greater and longer, of a pale browne on the outside, and white within, set together at one head; from whence rise vp many long branches, which by reason of their weight and weakenesse, cannot stand of themselues, but traile on the ground a yard and a halfe in length at the least (I relate it, as it hath growne with vs, but in what other forme, for flower or fruit, we know not) whereon are set at seuerall distances, broad and in a manner three square leaues, somewhat like triangled Iuie leaues, of a darke greene colour, the two sides whereof are broad and round, and the middle pointed at the end, standing reasonable close together: thus much we haue seene growe with vs, and no more: the roote rather decaying then increasing in our country.

The Potatoes of Virginia, which some foolishly call the Apples of youth, is another kinde of plant, differing much from the former, sauing in the colour and taste of the roote, hauing many weake and somewhat flexible branches, leaning a little downwards, or easily borne downe with the winde or other thing, beset with many winged leaues, of a darke grayish greene colour, whereof diuers are smaller, and some greater then others: the flowers growe many together vpon a long stalke, comming forth from betweene the leaues and the great stalkes, euery one seuerally vpon a short foot-stalke, somewhat like the flower of Tabacco for the forme, being one whole leafe six cornered at the brimmes, but somewhat larger, and of a pale blewish purple colour, or pale doue colour, and in some almost white, with some red threads in the middle, standing about a thicke gold yellow pointell, tipped with greene at the end: after the flowers are past, there come vp in their places small round fruit, as bigge as a Damson or Bullcis, greene at the first, and somewhat whitish afterwards, with many white seedes therein, like vnto Nightshade: the rootes are rounder and much smaller then the former, and some much greater then others, dispersed vnder ground by many small threads or strings from the rootes, of the same light browne colour on the outside, and white within, as they, and neare of the same taste, but not altogether so pleasant.

The Potatos of Canada, (which hath diuers names giuen it by diuers men, as Bauhinus vpon Matthiolus calleth it, *Solanum tuberosum esculentum*, Pelleterius of Middleborough in his *Plantarum Synonimia*, *Heliotropium Indicum tuberosum*, Fabius Columna in the second part of his *Phytobasanos*, *Flos Solis Farnesianus*, *sive Aster Peruanus tuberosus*: We in England, from some ignorant and idle head, haue called them Artichokes of Ierusalem, only because the roote, being boyled, is in taste like the bottome of an Artichoke head: but they may most fitly be called, Potatos of Canada, because their rootes are in forme, colour and taste, like vnto the Potatos of Virginia, but greater, and the French brought them first from Canada into these parts) riseth vp with diuers stiffe, round stalkes, eight or tenne foote high in our Country, where they haue scarce shewed their flowers, whereas the very head of flowers in other Countries, as Fabius Columna expresseth it, being of a Pyramis or Sugar loafe fashion, broade spreading below, and smaller pointed vpwards towards the toppe, is neere of the same length, whereon are set large and broade rough greene leaues, very like vnto the leaues of the flower of the Sunne, but smaller, yet growing in the very same manner, round about the stalkes: at the very later end of Summer, or the beginning of Autumne, if the roote bee well planted and defended, it will giue a shew of a few small yellow flowers at the top, like vnto the flowers of *Aster* or Starre-worte, and much smaller then any flower of the Sunne, which come to no perfection with vs: the roote, while the plant

is

1 *Carum.* Carawayes. 2 *Battatas Hispanorum.* Spanish Potatoes. 3 *Papas seu Battatas Virginianorum.* Virginia Potatoes. 4 *Battatas de Canada.* Potatoes of Canada, or Artichokes of Ierusalem.

is growing aboue ground, encreaſeth not to his full growth, but when the Summer is well ſpent, and the ſpringing of the ſtalk is paſt, which is about the end of Auguſt, or in September, then the root is perceiued to be encreaſed in the earth, and will before Autumne be ſpent, that is, in October, ſwell like a mound or hillocke, round about the foote of the ſtalkes, and will not haue his rootes fit to be taken vp, vntill the ſtalkes be halfe withered at the ſooneſt; but after they be withered, and ſo all the winter long vntill the Spring againe, they are good, and fit to bee taken vp and vſed, which are a number of tuberous round rootes, growing cloſe together; ſo that it hath beene obſerued, that from one roote, being ſet in the Spring, there hath been forty or more taken vp againe, and to haue ouer-filled a pecke meaſure, and are of a pleaſant good taſte as many haue tryed.

### The Vſe of all theſe Potato's.

The Spaniſh Potato's are roaſted vnder the embers, and being pared or peeled and ſliced, are put into ſacke with a little ſugar, or without, and is delicate to be eaten.

They are vſed to be baked with Marrow, Sugar, Spice, and other things in Pyes, which are a daintie and coſtly diſh for the table.

The Comfit-makers preſerue them, and candy them as diuers other things, and ſo ordered, is very delicate, fit to accompany ſuch other banquetting diſhes.

The Virginia Porato's being dreſſed after all theſe waies before ſpecified, maketh almoſt as delicate meate as the former.

The Potato's of Canada are by reaſon of their great increaſing, growne to be ſo common here with vs at London, that euen the moſt vulgar begin to deſpiſe them, whereas when they were firſt receiued among vs, they were dainties for a Queene.

Being put into ſeething water they are ſoone boyled tender, which after they bee peeled, ſliced and ſtewed with butter, and a little wine, was a diſh for a Queene, beeing as pleaſant as the bottome of an Artichoke: but the too frequent vſe, eſpecially being ſo plentifull and cheape, hath rather bred a loathing then a liking of them.

## Chap. L.

### *Cinara.* Artichokes.

THe fruits that grow vpon or neere the ground, are next to be entreated of, and firſt of Artichokes, whereof there be diuers kindes, ſome accounted tame and of the Garden, others wilde and of late planted in Gardens, Orchards or Fieldes, of purpoſe to be meate for men.

The Artichoke hath diuers great, large, and long hollowed leaues, much cut in or torne on both edges, without any great ſhew of prickles on them, of a kinde of whitiſh greene, like vnto an aſh colour, whereof it tooke the Latine name *Cinara*: the ſtalke is ſtrong, thicke and round, with ſomeskins as it were downe all the length of them, bearing at the toppe one ſcaly head, made at the firſt like a Pine-apple, but after growing greater, the ſcales are more ſeparate, yet in the beſt kindes lying cloſe, and not ſtaring, as ſome other kindes doe, which are eyther of a reddiſh browne, whitiſh, or greeniſh colour, and in ſome broade at the ends, in others ſharpe or prickly: after the head hath ſtood a great while, if it bee ſuffered, and the Summer proue hot and kindly, in ſome there will breake forth at the toppe thereof, a tuft of blewiſh purple thrumes or threds, vnder which grow the ſeede, wrapped in a great deale of dounie ſubſtance: but that roote that yeeldeth flowers will hardly abide the next winter; but elſe being cut off when it is well growne, that dounie matter abideth cloſe in the middle of the head, hauing the bottome thereof flat and round, which is that matter or ſubſtance that is vſed to be eaten: the roote ſpreadeth it ſelfe in the ground reaſonable

1 *Cinara satiua rubra.* The red Artichoke. 2 *Cinara satiua alba.* The white Artichoke. 3 *Cinara patula.* The French Artichoke. 4 *Cinara siluestris.* The Thistle Artichoke. 5 *Cardunus esculentus.* The Chardon.

ble well, yeelding diuers heads of leaues or ſuckers, whereby it is increaſed.

The white Artichoke is in all things like the red, but that the head is of a whitiſh aſhe colour, like the leaues, whereas the former is reddiſh.

We haue alſo another, whoſe head is greene, and very ſharpe vpwards, and is common in many places.

Wee haue had alſo another kinde in former times that grew as high as any man, and branched into diuers ſtalkes, euery one bearing a head thereon, almoſt as bigge as the firſt.

There is another kinde, called the Muske Artichoke, which groweth like the French kinde, but is much better in ſpending, although it haue a leſſer bottome.

The French Artichoke hath a white head, the ſcales whereof ſtand ſtaring far aſunder one from another at the ends, which are ſharpe: this is well known by this qualitie, that while it is hot after it is boyled, it ſwelleth ſo ſtrong, that one would verily thinke it had bin boyled in ſtinking water, which was brought ouer after a great froſte that had well nigh conſumed our beſt kindes, and are now almoſt cleane caſt out again, none being willing to haue it take vp the roome of better.

There is a lowe kinde that groweth much about Paris, which the French eſteeme more then any other, and is lower then the former French kinde, the head whereof as well as the leaues, is of a freſher greene colour, almoſt yellowiſh.

Then there is the Thiſtle Artichoke, which is almoſt a wilde kinde, and groweth ſmaller, with a more open and prickly head then any of the former.

And laſtly, the Chardon as they call it, becauſe it is almoſt of the forme and nature of a Thiſtle, or wilde Artichoke. This groweth high, and full of ſharpe prickles, of a grayiſh colour. Iohn Tradeſcante aſſured mee, hee ſaw three acres of Land about Bruſſels planted with this kinde, which the owner whited like Endiue, and then ſold them in the winter: Wee cannot yet finde the true manner of dreſſing them, that our Countrey may take delight therein.

All theſe kindes are encreaſed by ſlipping the young ſhootes from the root, which being replanted in February, March, or Aprill, haue the ſame yeare many times, but the next at the moſt, borne good heads.

We finde by dayly experience, that our Engliſh red Artichoke is in our Countrey the moſt delicate meate of any of the other, and therefore diuers thinking it to bee a ſeuerall kinde, haue ſent them into Italie, France, and the Lowe Countries, where they haue not abode in their goodneſſe aboue two yeare, but that they haue degenerated; ſo that it ſeemeth, that our ſoyle and climate hath the preheminence to nouriſh vp this plant to his higheſt excellencie.

## The Vſe of Artichokes.

The manner of preparing them for the Table is well knowne to the youngeſt Houſewife I thinke, to bee boyled in faire water, and a little ſalt, vntill they bee tender, and afterwardes a little vinegar and pepper, put to the butter, poured vpon them for the ſawce, and ſo are ſerued to the Table.

They vſe likewiſe to take the boyled bottomes to make Pyes, which is a delicate kinde of baked meate.

The Chardon is eaten rawe of diuers, with vinegar and oyle, pepper and ſalt, all of them, or ſome, as euery one liketh for their delight.

## Chap. LI.

### *Fabæ & Phaſeoli.* Garden and French Beanes.

THe Garden Beane is of two colours, red or blacke, and white, yet both riſe from one; the ſmall or fielde Beanes I make no mention of in this place; but the French or Kidney Beane is almoſt of infinite ſorts and colours: we doe not for all that intend to trouble you in this place, with the knowledge or relation of any more then is fit for a Garden of that nature, that I haue propounded it in the beginning.

Our ordinary Beanes, ſeruing for foode for the poorer ſort for the moſt part, are planted as well in fieldes as in gardens, becauſe the quantity of them that are ſpent taketh vp many acres of land to be planted in, and riſe vp with one, two or three ſtalks, according to the fertilitie of the ſoyle, being ſmooth and ſquare, higher then any man oftentimes, whereon are ſet at certaine diſtances, from the very bottome almoſt to the toppe, two long ſmooth fleſhy and thicke leaues almoſt round, one ſtanding by another at the end of a ſmall footeſtalke: betweene theſe leaues and the ſtalke, come forth diuers flowers, all of them looking one way for the moſt part, which are cloſe a little turned vp at the brimmes, white and ſpotted with a blackiſh ſpot in the middle of them, and ſomwhat purpliſh at the foot or bottome, of the forme almoſt of Broome or Peaſe flowers, many of which that grow vpward toward the toppe, doe ſeldome beare fruit, and therefore are gathered to diſtill, and the toppes of the ſtalkes cut off, to cauſe the reſt to thriue the better; after which grow vp long great ſmooth greene pods, greater then in any other kinde of Pulſe, which grow blacke when they are ripe, and containe within them two, three or foure Beanes, which are ſomewhat flat and round, eyther white or reddiſh, which being full ripe grow blackiſh: the roote hath diuers fibres annexed vnto the maine roote, which dyeth euery yeare.

The French or Kidney Beane riſeth vp at the firſt but with one ſtalke, which afterwards diuideth it ſelfe into many armes or branches, euery one of them being ſo weak, that without they be ſuſtained with ſtickes or poles, whereon with their winding and claſpers they take hold, they would lye fruitleſſe vpon the ground: vpon theſe branches grow forth at ſeuerall places long footeſtalkes, with euery of them three broade, round and pointed greene leaues at the end of them, towards the tops whereof come forth diuers flowers, made like vnto Peaſe bloſſomes, of the ſame colour for the moſt part that the fruit will be of, that is to ſay, eyther white, or yellow, or red, or blackiſh, or of a deepe purple &c. but white is moſt vſuall for our Garden; after which come long and ſlender flat pods, ſome crooked, and ſome ſtraight, with a ſtring as it were running downe the backe thereof, wherein are contained flattiſh round fruit, made to the faſhion of a kidney: the roote is long, and ſpreadeth with many fibres annexed vnto it, periſhing euery yeare.

### The Vſe of theſe Beanes.

The Garden Beanes ſerue (as I ſaid before) more for the vſe of the poore then of the rich: I ſhall therefore only ſhew you the order the poore take with them, and leaue curioſity to them that will beſtow time vpon them. They are only boyled in faire water and a little ſalt, and afterwards ſtewed with ſome butter, a little vinegar and pepper being put vnto them, and ſo eaten: or elſe eaten alone after they are boyled without any other ſawce. The water of the bloſſomes diſtilled, is vſed to take away ſpots, and to cleer the skin. The water of the greene huskes or cods is good for the ſtone.

The Kidney Beanes boyled in water huske and all, onely the ends cut off, and the ſtring taken away, and ſtewed with butter &c. are eſteemed more ſauory meate to many mens pallates, then the former, and are a diſh more oftentimes at rich mens Tables then at the poore.

## Chap. LII.

### *Pisum.* Peaſe.

THere is a very great variety of manured Peaſe known to vs, and I think more in our Country then in others, whereof ſome proſper better in one ground and country, and ſome in others: I ſhall giue you the deſcription of one alone for all the reſt, and recite vnto you the names of the reſt.

Garden Peaſe are for the moſt part the greateſt and ſweeteſt kinds, and are ſuſtained with ſtakes or buſhes. The Field Peaſe are not ſo vſed, but growe without any ſuch adoe. They ſpring vp with long, weake, hollow, and brittle (while they are young and greene) whitiſh greene ſtalkes, branched into diuers parts, and at euery ioynt where it parteth one broad round leafe compaſſing the ſtalke about, ſo that it commeth as it were thorough it: the leaues are winged, made of diuers ſmall leaues ſet to a middle ribbe, of a whitiſh greene colour, with claſpers at the ends of the leaues, whereby it taketh hold of whatſoeuer ſtandeth next vnto it: betweene the leaues and the ſtalkes come forth the flowers, ſtanding two or three together, euery one by it ſelfe on his owne ſeuerall ſtalke, which are either wholly white, or purple, or mixed white and purple, or purple and blew: the fruit are long, and ſomewhat round cods, whereof ſome are greater, others leſſer, ſome thicke and ſhort, ſome plaine and ſmooth, others a little crooked at the ends; wherein alſo are contained diuers formes of fruit or peaſe; ſome being round, others cornered, ſome ſmall, ſome great, ſome white, others gray, and ſome ſpotted: the roote is ſmall, and quickly periſheth.

The kindes of Peaſe are theſe:

| | |
|---|---|
| The Rounciuall. | The gray Peaſe. |
| The greene Haſting. | The white Haſting. |
| The Sugar Peaſe. | The Peaſe without skins. |
| The ſpotted Peaſe. | |

The Scottiſh or tufted Peaſe, which ſome call the Roſe Peaſe, is a good white Peaſe fit to be eaten.

The early or French Peaſe, which ſome call Fulham Peaſe, becauſe thoſe grounds thereabouts doe bring them ſooneſt forward for any quantity, although ſometimes they miſcarry by their haſte and earlineſſe.

### *Cicer Arietinum.* Rams Ciches.

This is a kinde of Pulſe, ſo much vſed in Spaine, that it is vſually one of their daintie diſhes at all their feaſts: They are of two ſorts, white and red; the white is onely vſed for meate, the other for medicine. It beareth many vpright branches with winged leaues, many ſet together, being ſmall, almoſt round, and dented about the edges: the flowers are either white or purple, according to the colour of the Peaſe which follow, and are ſomewhat round at the head, but cornered and pointed at the end, one or two at the moſt in a ſmall roundiſh cod.

### The Vſe of Peaſe.

Peaſe of all or the moſt of theſe ſorts, are either vſed when they are greene, and be a diſh of meate for the table of the rich as well as the poore, yet euery one obſeruing his time, and the kinde: the faireſt, ſweeteſt, youngeſt, and earlieſt for the better ſort, the later and meaner kindes for the meaner, who doe not giue the deereſt price: Or

Being dry, they ſerue to boyle into a kinde of broth or pottage, wherein many doe put Tyme, Mints, Sauory, or ſome other ſuch hot herbes, to giue it the better reliſh, and is much vſed in Towne and Countrey in the Lent time,

1 *Faba satiua.* Garden Beanes. 2 *Phaseoli satiui.* French Beanes. 3 *Pisum vulgare.* Garden Peaſe. 4 *Pisum vmbellatum ſiue Roſeum.* Roſe Peaſe or Scottiſh Peaſe. 5 *Pisum Saccharatum.* Sugar Peaſe. 6 *Pisum maculatum.* Sported Peaſe. 7 *Cicer Arietinum.* Rams Ciches or Cicers.

2. P. vulgaris, L. mei dict.
5. Pisum saccharum, L. E. mei dict.

time, especially of the poorer sort of people.

It is much vsed likewise at Sea for them that goe long voyages, and is for change, because it is fresh, a welcome diet to most persons therein.

The Rams Ciches the Spaniards call *Grauancos*, and *Garauancillos*, and eate them boyled and stewed as the most dainty kinde of Pease that are, they are of a very good relliſh, and doe nouriſh much, but yet are not without that windy quality that all sorts of Pulse are subiect vnto: they increase bodily lust much more then any other sorts, and as it is thought, doth helpe to encrease seede.

## Chap. LIII.

### *Cucumer.* The Cowcumber.

Of Cowcumbers there are diuers sorts, differing chiefly in the forme and colour of the fruit, and not in the forme of the plant; therefore one description shall serue in stead of all the rest.

The Cowcumber bringeth forth many trailing rough greene branches lying on the ground, all along whereof growe seuerall leaues, which are rough, broad, vneuen at the edges, and pointed at the ends, with long crooked tendrels comming forth at the same ioynt with the leafe, but on the other side therof: between the stalks & the leaues at the ioynts come forth the flowers seuerally, euery one standing on a short foot-stalke, opening it selfe into fiue leaues, of a yellowish colour, at the bottome whereof groweth the fruit, long and greene at the first, but when it is thorough ripe, a little yellowish, hauing many furrowes, and vneuen bunches all the length of it, wherein is a white firme substance next vnto the skin, and a cleare pulpe or watery substance, with white flat seede lying disperſed through it: the roote is long and white, with diuers fibres at it.

#### The kindes.

The first described is called, The long greene Cowcumber.

There is another is called, The short Cowcumber, being short, and of an equall bigneſſe in the body thereof, and of an vnequall bigneſſe at both ends.

The long Yellow, which is yellowish from the beginning, and more yellow when it is ripe, and hath beene measured to be thirteene inches long: but this is not that small long Cowcumber, called of the Latines, *Cucumis anguinus*.

Another kinde is early ripe, called The French kinde.

The Dantsicke kinde beareth but small fruit, growing on short branches or runners: the pickled Cowcumbers that are vsually sold are of this kind.

The Muscouie kinde is the smalleſt of all other, yet knowne, and beareth not aboue foure or fiue at the most on a roote, which are no bigger then small Lemons.

#### The Vse of Cowcumbers.

Some vse to cast a little salt on their sliced Cowcumbers, and let them stand halfe an houre or more in a dish, and then poure away the water that commeth from them by the salt, and after put vinegar, oyle, &c. thereon, as euery one liketh: this is done, to take away the ouermuch watcriſnneſſe and coldneſſe of the Cowcumbers.

In many countries they vse to eate Cowcumbers as wee doe Apples or Peares, paring and giuing slices of them, as we would to our friends of some dainty Apple or Peare.

The pickled Cowcumbers that come from beyond Sea, are much vsed

with

with vs for ſawce to meate all the Winter long. Some haue ſtriuen to equall them, by pickling vp our Cowcumbers at the later end of the yeare, when they are cheapeſt, taking the little ones and ſcalding them thoroughly well, which after they put in brine, with ſome Dill or Fenell leaues and ſtalkes: but theſe are nothing comparable to the former, wee either miſſing of the right and orderly pickling of them, or the kinde it ſelfe differing much from ours (as I ſaid of the Dantſicke kinde) for ours are neither ſo tender and firme, nor ſo ſauoury as the other.

The rawe or greene Cowcumbers are fitteſt for the hotter time of the yeare, and for hot ſtomackes, and not to be vſed in colder weather or cold ſtomackes, by reaſon of the coldneſſe, whereby many haue been ouertaken.

The ſeede is vſed phyſically in many medicines that ſerue to coole, and a little to make the paſſages of vrine ſlippery, and to giue eaſe to hot diſeaſes.

## Chap. LIIII.

### *Melo.* Milions or Muske Melons.

THere bee diuers ſorts of Melons found out at this day, differing much in the goodneſſe of taſte one from another. This Countrey hath not had vntill of late yeares the skill to nourſe them vp kindly, but now there are many that are ſo well experienced therein, and haue their ground ſo well prepared, as that they will not miſſe any yeare, if it be not too extreme vnkindly, to haue many ripe ones in a reaſonable time: yet ſome will be later then others alwayes.

The Melon is certainly a kinde of Cowcumber, it doth ſo neare reſemble it, both in the manner of his growing, hauing rough trailing branches, rough vneuen leaues, and yellow flowers: after which come the fruit, which is rounder, thicker, bigger, more rugged, and ſpotted on the outſide then the Cowcumber, of a ruſſet colour, and greene vnderneath, which when it groweth full ripe, will change a little yellowiſh, being as deepe furrowed and ribbed as they, and beſides hauing chaps or rifts in diuers places of the rinde: the inward hard ſubſtance is yellow, which onely is eaten: the ſeede which is bigger, and a little yellower then the Cowcumber, lying in the middle onely among the moiſter pulpe: the ſmell and changing of his colour, fore-ſhew their ripeneſſe to them that are experienced: the roote is long, with many fibres at it. The fruit requireth much watering in the hot time of the day, to cauſe them to ripen the ſooner, as I haue obſerued by diuers of the beſt skill therein.

### The Vſe of the kindes of Melons.

The beſt Melon ſeede doe come to vs out of Spaine, ſome haue come out of Turkie, but they haue been nothing ſo good and kindly.

Some are called Sugar Melons, others Peare Melons, and others Muske Melons.

They haue beene formerly only eaten by great perſonages, becauſe the fruit was not only delicate but rare; and therfore diuers were brought from France, and ſince were nourſed vp by the Kings or Noblemens Gardiners onely, to ſerue for their Maſters delight: but now diuers others that haue skill and conueniencie of ground for them, doe plant them and make them more common.

They paire away the outer rinde, and cut out the inward pulpe where the ſeede lyeth, ſlice the yellow firme inward rinde or ſubſtance, & ſo eate it with ſalt and pepper (and good ſtore of wine, or elſe it will hardly diſgeſt) for this is firmer, & hath not that moiſture in it that the Cowcumbers haue. It is alſo more delicate, and of more worth, which recompenſeth the paine.

The ſeed of theſe Melons are vſed as Cowcumbers phyſically, and together with them moſt vſually.

## Chap. LV.

### *Pepo.* Pompions.

WE haue but one kinde of Pompion (as I take it) in all our Gardens, notwithstanding the diuersities of bigneſſe and colour.

The Pompion or great Melon (or as ſome call it Milion) creepeth vpon the ground (if nothing bee by it whereon it may take hold and climbe) with very great, ribbed, rough, and prickly branches, whereon are ſet very large rough leaues, cut in on the edges with deepe gaſhes, and dented beſides, with many claſpers alſo, which winde about euery thing they meete withall: the flowers are great and large, hollow and yellow, diuided at the brims into fiue parts, at the bottome of which, as it is in the reſt, groweth the fruit, which is very great, ſometimes of the bigneſſe of a mans body, and oftentimes leſſe, in ſome ribbed or bunched, in others plaine, and either long or round, either green or yellow, or gray, as Nature liſteth to ſhew her ſelfe; for it is but waſte time, to recite all the formes and colours may be obſerued in them: the inner rinde next vnto the outer is yellowiſh and firme: the ſeede is great, flat, and white, lying in the middle of the watery pulpe: the roote is of the bigneſſe of a mans thumbe or greater, diſperſed vnder ground with many ſmall fibres ioyned thereunto.

Gourds are kindes of Melons; but becauſe wee haue no vſe of them, wee leaue them vnto their fit place.

#### The Vſe of Pompions.

They are boyled in faire water and ſalt, or in powdered beefe broth, or ſometimes in milke, and ſo eaten, or elſe buttered. They vſe likewiſe to take out the inner watery ſubſtance with the ſeedes, and fill vp the place with Pippins, and hauing laid on the couer which they cut off from the toppe, to take out the pulpe, they bake them together, and the poore of the Citie, as well as the Country people, doe eate thereof, as of a dainty diſh.

The ſeede hereof, as well as of Cowcumbers and Melons, are cooling, and ſerue for emulſions in the like manner for Almond milkes, &c. for thoſe are troubled with the ſtone.

## Chap. LVI.

### *Fragaria.* Strawberries.

THere be diuers ſorts of Strawberries, whereof thoſe that are nourſed vp in Gardens or Orchards I intend to giue you the knowledge in this place, and leaue the other to a fitter; yet I muſt needs ſhew you of one of the wilde ſorts, which for his ſtrangeneſſe is worthy of this Garden: And I muſt alſo enforme you, that the wilde Strawberry that groweth in the Woods is our Garden Strawberry, but bettered by the ſoyle and tranſplanting.

The Strawberry hath his leaues cloſed together at the firſt ſpringing vp, which afterwards ſpread themſelues into three diuided parts or leaues, euery one ſtanding vpon a ſmall long foote-ſtalke, greene on the vpperſide, grayiſh vnderneath, and ſnipped or dented about the edges; among which riſe vp diuers ſmall ſtalkes, bearing foure or fiue flowers at the tops, conſiſting of fiue white round pointed leaues, ſomewhat yellowiſh in the bottome, with ſome yellow threads therein; after which come the fruit, made of many ſmall graines ſet together, like vnto a ſmall Mulberry or Raſpis, reddiſh when it is ripe, and of a pleaſant winy taſte, wherein is encloſed diuers ſmall blackiſh ſeede: the roote is reddiſh and long, with diuers ſmall threads at it, and ſendeth forth

1 *Cucumis longus vulgaris.* The ordinary Cowcumber. 2 *Cucumis Hispanicus.* The long yellow Spanish Cowcumber. 3 *Melo vulgaris.* The ordinary Melon. 4 *Melo maximus optimus.* The greateſt Muske Melon. 5 *Pepo.* The Pompion. 6 *Fraga vulgaris.* Common Strawberries. 7 *Fraga Bohemica maxima.* The great Bohemia Strawberries. 8 *Fraga aculeata.* The prickly Strawberry.

forth from the head therof long reddiſh ſtrings running vpon the ground, which ſhoot forth leaues in many places, whereby it is much encreaſed.

The white Strawberry differeth not from the red, but in the colour of the fruite, which is whiter then the former when it is thorough ripe, enclining to redneſſe.

The greene Strawberry likewiſe differeth not, but that the fruit is green on all ſides when it is ripe, ſaue on that ſide the Sun lyeth vpon it, and there it is ſomewhat red.

The Virginia Strawberry carryeth the greateſt leafe of any other, except the Bohemian, but ſcarce can one Strawberry be ſeene ripe among a number of plants; I thinke the reaſon thereof to be the want of skill, or induſtry to order it aright. For the Bohemia, and all other Strawberries will not beare kindly, if you ſuffer them to grow with many ſtrings, and therefore they are ſtill cut away.

There is another very like vnto this, that Iohn Tradeſcante brought with him from Bruſſels long agoe, and in ſeuen yeares could neuer ſee one berry ripe on all ſides, but ſtill the better part rotten, although it would euery yeare flower abundantly, and beare very large leaues.

The Bohemia Strawberry hath beene with vs but of late dayes, but is the goodlieſt and greateſt, both for leafe next to the Virginian, and for beauty farre ſurpaſſing all; for ſome of the berries haue beene meaſured to bee neere fiue inches about. Maſter Queſter the Poſtmaſter firſt brought them ouer into our Country, as I vnderſtand, but I know no man ſo induſtrious in the carefull planting and bringing them to perfection in that plentifull maner, as Maſter Vincent Sion who dwelt on the Banck ſide, neer the old Paris garden ſtaires, who from ſeuen rootes, as hee affirmed to me, in one yeare and a halfe, planted halfe an acree of ground with the increaſe from them, beſides thoſe he gaue away to his friends, and with him I haue ſeene ſuch, and of that bigneſſe before mentioned.

One Strawberry more I promiſed to ſhew you, which although it be a wilde kinde, and of no vſe for meate, yet I would not let this diſcourſe paſſe, without giuing you the knowledge of it. It is in leafe much like vnto the ordinary, but differeth in that the flower, if it haue any, is greene, or rather it beareth a ſmall head of greene leaues, many ſet thicke together like vnto a double ruffe, in the midſt whereof ſtandeth the fruit, which when it is ripe, ſheweth to be ſoft and ſomwhat reddiſh, like vnto a Strawberry, but with many ſmall harmleſſe prickles on them, which may be eaten and chewed in the mouth without any maner of offence, and is ſomewhat pleaſant like a Strawberry: it is no great bearer, but thoſe it doth beare, are ſet at the toppes of the ſtalks cloſe together, pleaſant to behold, and fit for a Gentlewoman to weare on her arme, &c. as a raritie in ſtead of a flower.

## The Vſe of Strawberries.

The leaues of Strawberries are alwaies vſed among other herbes in cooling drinkes, as alſo in lotions, and gargles for the mouth and throate: the rootes are ſometimes added to make it the more effectuall, and withall ſomwhat the more binding.

The berries themſelues are often brought to the Table as a reare ſeruice, whereunto claret wine, creame or milke is added with ſugar, as euery one liketh; as alſo at other times, both with the better and meaner ſort, and are a good cooling and pleaſant diſh in the hot Summer ſeaſon.

The water diſtilled of the berries, is good for the paſſions of the heart, cauſed by the perturbation of the ſpirits, being eyther drunke alone, or in wine; and maketh the heart merry.

Some doe hold that the water helpeth to clenſe the face from ſpots, and to adde ſome cleereneſſe to the skinne.

## Chap. LVII.

*Angelica.* Garden Angelica.

HAuing thus furniſhed you out a Kitchen Garden with all ſorts of herbes, roots & fruits fit for it, and for any mans priuate vſe, as I did at the firſt appropriate it; let me a little tranſcend, and for the profit & vſe of Country Gentlewomen and others, furniſh them with ſome few other herbes, of the moſt eſpeciall vſe for thoſe ſhall need them, to be planted at hand in their Gardens, to ſpend as occaſion ſhall ſerue, and firſt of Angelica.

Angelica hath great and long winged leaues, made of many broade greene ones, diuided one from another vpon the ſtalk, which is three foot long or better ſomtimes, among which riſe vp great thicke and hollow ſtalkes with ſome few ioynts, whereat doth alwayes ſtand two long leaues compaſſing the ſtalke at the bottome, in ſome places at the ioynts ſpring out other ſtalkes or branches, bearing ſuch like leaues but ſmaller, and at the tops very large vmbels of white flowers, that turne into whitiſh ſeede ſomewhat thicke: the roote groweth great with many branches at it, but quickly periſheth after it hath borne ſeede: to preſerue the roote therefore the better, they vſe to cut it often in the yeare, thereby to hinder the running vp to ſeede: the whole plant, both leafe, roote and ſeede, is of an excellent comfortable ſent, ſauour and taſte.

### The Vſe of Angelica.

The diſtilled water of Angelica, eyther ſimple or compound, is of eſpeciall vſe *in deliquium animi, vel cordis tremores & paſsiones*, that is, ſwounings, when the ſpirits are ouercome and faint, or tremblings and paſſions of the heart, to expell any windy or noyſome vapours from it. The green ſtalkes or the young rootes being preſerued or candied, are very effectuall to comfort and warme a colde and weake ſtomacke: and in the time of infection is of excellent good vſe to preſerue the ſpirits and heart from infection. The dryed roote made into pouder, and taken in wine or other drinke, will abate the rage of luſt in young perſons, as I haue it related vnto me vpon credit: A Syrupe made thereof in this manner, is very profitable to expectorate flegme out of the cheſt and lunges, and to procure a ſweete breath. Into the greene ſtalke of Angelica as it ſtandeth growing, make a great gaſhe or inciſion, wherein put a quantitie of fine white Sugar, letting it there abide for three dayes, and after take it forth by cutting a hole at the next ioynt vnder the cut, where the Syrupe reſteth, or cut off the ſtalke, and turne it downe, that the Syrupe may drayne forth; which keepe for a moſt delicate medicine.

## Chap. LVIII.

*Dracunculus hortenſis ſiue Serpentaria.* Dragons.

DRagons riſeth out of the ground with a bare or naked round whitiſh ſtalke, ſpotted very much with purpliſh ſpots and ſtrakes, bearing at the toppe therof a few greene leaues very much diuided on all ſides, ſtanding vpon long foote-ſtalkes, in the middle whereof (if the roote be old enough) commeth forth a great long huske or hoſe, green on the outſide, and of a darke purpliſh colour on the inſide, with a ſlender long reddiſh peſtell or clapper in the middle: the roote is great, round, flat and whitiſh on the outſide, and whiter within, very like vnto the rootes of *Arum*, or Wakerobin, and taſting ſomewhat ſharpe like it.

### The Vſe of Dragons.

The chiefe vſe whereunto Dragons are applyed, is, that according to an old receiued cuſtome and tradition (and not the iudgement of any learned Author) the diſtilled water is giuen with Mithridatum or Treakle to expell noyſome and peſtilentiall vapours from the heart.

## Chap. LIX.

### *Ruta.* Garden Rue, or Herbe Grace.

GArden Rue or Herbe Grace groweth vp with hard whitiſh wooddy ſtalkes, whereon are ſet diuers branches of leaues, being diuided into many ſmall ones, which are ſomewhat thicke and round pointed, of a blewiſh greene colour: the flowers ſtand at the tops of the ſtalkes conſiſting of foure ſmall yellow leaues, with a greene button in the middle, and diuers ſmall yellow threds about it, which growing ripe, containe within them ſmall blacke ſeede: the roote is white and wooddy, ſpreading farre in the ground.

### The Vſe of Rue.

The many good properties whereunto Rue ſerueth, hath I thinke in former times cauſed the Engliſh name of Herbe Grace to be giuen vnto it. For without doubt it is a moſt wholeſome herbe, although bitter and ſtrong, and could our dainty ſtomackes brooke the vſe thereof, it would worke admirable effects being carefully and skilfully applyed, as time and occaſion did require: but not vndiſcreetly or hand ouer head, as many vſe to doe that haue no skill. Some doe rippe vp a beade rowle of the vertues of Rue, as Macer the Poet and others, in whom you ſhall finde them ſet downe, to bee good for the head, eyes, breaſt, liuer, heart, ſpleene, &c. In ſome places they vſe to boyle the leaues of Rue, and keep them in pickle, to eate them as Sampire for the helpe of weake eyes. It is very auaileable in gliſters or drinkes againſt the winde or the collicke, and to procure vrine that is ſtayed by the paines therof. The diſtilled water is often vſed for the ſame purpoſes aforeſaid: but beware of the too frequent or ouermuch vſe thereof, becauſe it heateth exceedingly, and waſteth nature mightily.

## Chap. LX.

### *Carduus Benedictus.* The Bleſſed Thiſtle.

CArduus benedictus or the bleſſed Thiſtle, hath many weake tender branches lying for the moſt part on the ground, whereon are ſet long and narrow leaues, much cut in or waued about the edges, hairy or rough in handling, yet without any hard or ſharpe thornes or prickles at all, that the tendereſt hand may touch them without harme: but thoſe that grow toward the toppes of the ſtalkes are ſomewhat more prickly, and the heads which grow on the tops of the ſeuerall branches are ſomewhat ſharpe, ſet with prickles like a Thiſtle: the flower is yellow, and the ſeede lying within the woolly or flocky doune like to all other thiſtles, are blackiſh, long and round, with a few haires on the head of them: the roote is white, and periſheth euery yeare after it hath giuen ſeede.

### The Vſe of the bleſſed Thiſtle.

The diſtilled water hereof is much vſed to be drunke againſt agues of all ſortes, eyther peſtilentiall or humorall, of long continuance or of leſſe:

but

1 *Angellica.* Angellica. 2 *Dracunculus hortensis.* Dragons. 3 *Rutahortensis.* Garden Rue, or Herbegrace. 4 *Cardnus benedictus.* The blessed Thistle. 5 *Alkakengi sive Solanum Halicacabum & Vesicarium.* Winter Cherries. 6 *Asarum.* Asarabacca. 7 *Liquiritia.* Licoris.

but the decoction of the herbe giuen in due time, hath the more forcible operation: it helpeth to expell wormes, becauſe of the bitterneſſe, and is thereby alſo a friend to the ſtomack ouercharged with chollar, and to clenſe the liuer: it prouoketh ſweate and vrine, is helpefull to them are troubled with the ſtone, and to eaſe paines in the ſides.

## Chap. LXI.

*Solanum veſicarium, ſiue Alkakengi.* Winter Cherries.

THe Winter Cherry hath a running or creeping roote in the ground, of the bigneſſe wany times of ones little finger, ſhooting forth at ſeuerall ioynts in ſeuerall places, whereby it quickly ſpreadeth a great compaſſe of ground: the ſtalke riſeth not aboue a yard high, whereon are ſet many broade and long greene leaues, ſomewhat like vnto the leaues of Nightſhade, but larger: at the ioynts whereof come forth whitiſh flowers made of fiue leaues a peece, which after turne into green berries, incloſed with thin skins or bladders, which change to bee reddiſh when they grow ripe, the berry likewiſe being reddiſh, and as large as a Cherry, wherein are contained many flat and yellowiſh ſeed lying within the pulpe: which being gathered and ſtrung vp, are kept all the yeare to be vſed vpon occaſion.

### The Vſe of Winter Cherries.

The diſtilled water of the herbe and fruit together, is often taken of them that are troubled with the ſharpneſſe or difficultie of vrine, and with the ſtone in the kidneyes, or grauel in the bladder: but the berries themſelues either greene or dryed boyled eyther in broth, in wine, or in water, is much more effectuall: It is likewiſe conducing to open obſtructions of the liuer, &c. and thereby to helpe the yellow Iaundiſe.

## Chap. LXII.

*Aſarum.* Aſarabacca.

ASarabacca, from a ſmall creeping roote ſet with many fibres, ſhooteth forth diuers heads, and from euery of them ſundry leaues, euery one ſtanding vpon a long greene ſtalke, which are round, thicke, and of a very ſad or darke greene colour, and ſhining withall: from the rootes likewiſe ſpring vp ſhort ſtalkes, not fully foure fingers high, at the toppe of euery one of which ſtandeth the flower, in faſhion very like the ſeede veſſell of Henbane ſeede, of a greeniſh purple colour, which changeth not his forme, but groweth in time to containe therein ſmall cornered ſeed: the greene leaues abide all the winter many times, but vſually ſheddeth them in winter, and recouereth freſh in the ſpring.

### The Vſe of Aſarabacca.

The leaues are much and often vſed to procure vomits, fiue or ſeuen of them bruiſed, and the iuice of them drunke in ale or wine. An extract made of the leaues with wine artificially performed, might bee kept all the yeare thorough, to bee vſed vpon any preſent occaſion, the quantitie to bee proportioned according to the conſtitution of the patient. The roote worketh not ſo ſtrongly by vomit, as the leaues, yet is often vſed for the ſame purpoſe, and beſides is held auaileable to prouoke vrine, to open obſtructions in the liuer and ſpleene, and is put among diuers other ſimples, both into Mithridatum and Andromachus Treakle, which is vſually called Venice Treakle. A dram of the dryed roots in pouder giuen in white wine a little before the fit of an ague, taketh away the ſhaking fit, & therby cauſe the hot fit to be the more remiſſe, and in twice taking expell it quite.

## Chap. LXIII.

### *Glycyrrhiza siue Liqueritia.* Licorice.

ALthough there are two sorts of Licorice set downe by diuers Authors, yet because this Land familiarly is acquainted but with one sort, I shall not neede for this Garden, to make any further relation of that is vnknowne, but onely of that sort which is sufficiently frequent with vs. It riseth vp with diuers wooddy stalks, whereon are set at seuerall distances many winged leaues, that is to say, many narrow long greene leaues set together on both sides of the stalke, and an odde one at the end, very well resembling a young Ashe tree sprung vp from the seede: this by many yeares continuance in a place without remouing, and not else, will bring forth flowers many standing together spike-fashion one aboue another vpon the stalkes, of the forme of Pease blossomes, but of a very pale or bleake blew colour, which turne into long somewhat flat and smooth cods, wherein is contained small round hard seede: the roote runneth downe exceeding deep into the ground, with diuers other smaller roots and fibres growing with them, and shoote out suckers from the maine rootes all about, whereby it is much encreased, of a browniſh colour on the outside, and yellow within, of a farre more weake sweete taste, yet far more pleasing to vs then that Licorice that is brought vs from beyond Sea; because that, being of a stronger sweet taste hath a bitternesse ioyned with it, which maketh it the lesse pleasing and acceptable to most.

### The Vse of Licorice.

Our Englisſh Licorice is now adaies of more familiar vse (as I said before) then the outlandisſh, and is wholly spent and vsed to helpe to digest and expectorate flegme out of the chest and lunges, and doth allay the sharpenesse or saltnesse thereof. It is good also for those are troubled with shortnesse of breath, and for all sorts of coughes. The iuice of Licorice artificially made with Hyssope water, serueth very well for all the purposes aforesaid. It being dissolued with Gum Tragacanth in Rose water, is an excellent Lohoc or licking medicine to breake flegme, and to expectorate it, as also to avoyde thin frothy matter, or thin salt flegme, which often fretteth the lunges. It doth also lenifie exulcerated kidneyes, or the bladder, and helpeth to heale them. It is held also good for those that cannot make their water but by drops, or a small deale at a time.

The dryed root finely minced, is a speciall ingredient into all Trageas or Dredges, seruing for the purposes aforesaid, but the vse of them is almost wholly left now adaies with all sorts.

Thus haue I shewed you not only the herbes, rootes and fruites, nourſed vp in this Garden, but such herbes as are of most necessary vses for the Country Gentlewomens houses: And now I will shew you the Orchard also.

# THE ORDERING OF THE ORCHARD.

## The third part, or ORCHARD.

### CHAP. I.

*The ſituation of an Orchard for fruit-bearing trees, and how to amend the defects of many grounds.*

AS I haue done in the two former parts of this Treatiſe, ſo I meane to proceede in this; firſt to ſet downe the ſituation of an Orchard, and then other things in order: And firſt, I hold that an Orchard which is, or ſhould bee of ſome reaſonable large extent, ſhould be ſo placed, that the houſe ſhould haue the Garden of flowers iuſt before it open vpon the South, and the Kitchen Garden on the one ſide thereof, ſhould alſo haue the Orchard on the other ſide of the Garden of Pleaſure, for many good reaſons: Firſt, for that the fruit trees being grown great and tall, will be a great ſhelter from the North and Eaſt windes, which may offend your chiefeſt Garden, and although that your Orchard ſtand a little bleake vpon the windes, yet trees rather endure theſe ſtrong bitter blaſts, then other ſmaller and more tender ſhrubs and herbes can doe. Secondly, if your Orchard ſhould ſtand behinde your Garden of flowers more Southward, it would ſhadow too much of the Garden, and beſides, would ſo binde in the North and Eaſt, and North and Weſt windes vpon the Garden, that it would ſpoile many tender things therein, and ſo much abate the edge of your pleaſure thereof, that you would willingly wiſh to haue no Orchard, rather then that it ſhould ſo much annoy you by the ſo ill ſtanding thereof. Thirdly, the falling leaues being ſtill blowne with the winde ſo aboundantly into the Garden, would either ſpoile many things, or haue one daily and continuall attending thereon, to cleanſe and ſweepe them away. Or elſe to auoide theſe great inconueniences, appoint out an Orchard the farther off, and ſet a greater diſtance of ground betweene. For the ground or ſoile of the Orchard, what I haue ſpoken concerning the former Garden for the bettering of the ſeuerall grounds, may very well ſerue and be applyed to this purpoſe. But obſerue this, that whereas your Gardens before ſpoken of may be turned vp, manured, and bettered with ſoile if they growe out of heart, your Orchard is not ſo eaſily done, but muſt abide many yeares without altering; and therefore if the ground be barren, or not good, it had the more neede to bee amended, or wholly made good, before you make an Orchard of it; yet ſome there be that

that doe appoint, that where euery tree ſhould bee ſet, you onely digge that place to make it good: but you muſt know, that the rootes of trees runne further after a little times ſtanding, then the firſt compaſſe they are ſet in; and therefore a little compaſſe of ground can maintaine them but a little while, and that when the rootes are runne beyond that ſmall compaſſe wherein they were firſt ſet, and that they are come to the barren or bad ground, they can thriue no better then if they had beene ſet in that ground at the firſt, and if you ſhould afterwards digge beyond that compaſſe, intending to make the ground better further off, you ſhould much hurt the ſpreading rootes, and put your trees in danger: the ſituation of hils in many places is grauelly or chalky, which is not good for trees, becauſe they are both too ſtonie, and lacke mellow earth, wherein a tree doth moſt ioy and proſper, and want moiſture alſo (which is the life of all trees) becauſe of the quicke deſcent of raine to the lower grounds: and beſides all theſe inconueniences there is one more; your trees planted either on hils or hill ſides, are more ſubiect to the fury and force of windes to be ouerturned, then thoſe that growe in the lower grounds; for the ſtrongeſt and moſt forcible windes come not vſually out of the North Eaſt parts, where you prouide beſt defence, but from the South and Weſt, whence you looke for the beſt comfort of the Sunne. To helpe therefore manie of the inconueniences of the hils ſides, it were fit to cauſe manie leauels to bee made thereon, by raiſing the lower grounds with good earth, and ſuſtaining them with bricke or ſtone wals, which although chargeable, will counteruaile your coſt, beſide the pleaſure of the walkes, and proſpect of ſo worthy a worke. The plaine or leuell grounds as they are the moſt frequent, ſo they are the moſt commendable for an Orchard, becauſe the moulds or earths are more rich, or may better and ſooner be made ſo; and therefore the profits are the more may be raiſed from them. A ſtiffe clay doth nouriſh trees well, by reaſon it containeth moiſture; but in regard of the coldneſſe thereof, it killeth for the moſt part all tender and early things therein: ſea-cole aſhes therefore, bucke aſhes, ſtreete ſoyle, chaulke after it hath lyen abroad and been broken with many yeares froſts and raine, and ſheepes dung, are the moſt proper and fitteſt manure to helpe this kinde of ſoyle. The dry ſandy ſoile, and grauelly ground are on the contrary ſide as bad, by reaſon of too much heate and lacke of moiſture: the dung of kine or cattell in good quantity beſtowed thereon, will much helpe them. The amending or bettering of other ſorts of grounds is ſet down toward the end of the firſt Chapter of the firſt part of this worke, whereunto I will referre you, not willing to repeate againe the ſame things there ſet downe. The beſt way to auoide and amend the inconueniences of high, boiſterous, and cold windes, is to plant Walnut trees, Elmes, Oakes or Aſhes, a good diſtance without the compaſſe of your Orchard, which after they are growne great, will bee a great ſafeguard thereunto, by breaking the violence of the windes from it. And if the ſoyle of your Orchard want moiſture, the conueying of the ſinke of the houſe, as alſo any other draine of water thereinto, if it may be, will much helpe it.

## Chap. II.

### *The forme of an Orchard, both ordinary, and of more grace and rarity.*

ACcording to the ſituation of mens grounds, ſo muſt the plantation of them of neceſſitie be alſo; and if the ground be in forme, you ſhall haue a formall Orchard: if otherwiſe, it can haue little grace or forme. And indeed in the elder ages there was ſmall care or heede taken for the formality; for euery tree for the moſt part was planted without order, euen where the maſter or keeper found a vacant place to plant them in, ſo that oftentimes the ill placing of trees without ſufficient ſpace betweene them, and negligence in not looking to vphold them, procured more waſte and ſpoile of fruit, then any accident of winde or weather could doe. Orchards in moſt places haue not bricke or ſtone wals to ſecure them, becauſe the extent thereof being larger

larger then of a Garden, would require more coſt, which euery one cannot vndergoe; and therefore mud wals, or at the beſt a quicke ſet hedge, is the ordinary and moſt vſuall defence it findeth almoſt in all places : but with thoſe that are of ability to compaſſe it with bricke or ſtone wals, the gaining of ground, and profit of the fruit trees planted there againſt, will in ſhort time recompenſe that charge. If you make a doubt how to be ſure that your Orchard wall ſhall haue ſufficient comfort of the Sunne to ripen the fruits, in regard the trees in the Orchard being ſo nigh thereunto, and ſo high withall, will ſo much ſhadow the wall, that nothing will ripen well, becauſe it will want the comfort of the Sunne: you may follow this rule and aduice, to remedy thoſe inconueniences. Hauing an Orchard containing one acre of ground, two, three, or more, or leſſe, walled about, you may ſo order it, by leauing a broad and large walke betweene the wall and it, containing twenty or twenty foure foote (or yards if you will) that the wall ſhall not be hindered of the Sun, but haue ſufficient comfort for your trees, notwithſtanding the height of them, the diſtance betweene them and the wall being a ſufficient ſpace for their ſhadow to fall into: and by compaſſing your Orchard on the inſide with a hedge (wherein may bee planted all ſorts of low ſhrubs or buſhes, as Roſes, Cornellian Cherry trees plaſhed lowe, Gooſeberries, Curran trees, or the like) you may encloſe your walke, and keepe both it and your Orchard in better forme and manner, then if it lay open. For the placing of your trees in this Orchard, firſt for the wals: Thoſe ſides that lye open to the South & Southweſt Sunne, are fitteſt to bee planted with your tendereſt and earlieſt fruits, as Apricockes, Peaches, Nectarius, and May or early Cherries: the Eaſt, North and Weſt, for Plums and Quinces, as you ſhall like beſt to place them. And for the Orchard it ſelfe, the ordinary manner is to place them without regard of meaſure or difference, as Peares among Apples, and Plums among Cherries promiſcuouſly; but ſome keepe both a diſtance and a diuiſion for euery ſort, without intermingling: yet the moſt gracefull Orchard containeth them all, with ſome others, ſo as they be placed that one doe not hinder or ſpoile another; and therefore to deſcribe you the modell of an Orchard, both rare for comelineſſe in the proportion, and pleaſing for the profitableneſſe in the vſe, and alſo durable for continuance, regard this figure is here placed for your direction, where you muſt obſerue, that your trees are here ſet in ſuch an equall diſtance one from another euery way, & as is fitteſt for them, that when they are grown great, the greater branches ſhall not gall or rubbe one againſt another; for which purpoſe twenty or ſixteene foot is the leaſt to be allowed for the diſtance euery way of your trees, & being ſet in rowes euery one in the middle diſtance, will be the moſt gracefull for the plantation, and beſides, giue you way ſufficient to paſſe through them, to pruine, loppe, or dreſſe them, as need ſhall require, and may alſo bee brought (if you pleaſe) to that gracefull delight, that euery alley or diſtance may be formed like an arch, the branches of either ſide meeting to be enterlaced together. Now for the ſeuerall ſorts of fruit trees that you ſhall place in this modell, your beſt direction is to ſet Damſons, Bulleis, and your taler growing Plums on the outſide, and your lower Plums, Cherries, and Apples on the inſide, hauing regard, that you place no Peare tree to the Sunward, of any other tree, leſt it ouer-



ſhadow

ſhadow them: Let your Peare trees therefore be placed behinde, or on the one ſide of your lower trees, that they may be as it were a ſhelter or defence on the North & Eaſt ſide. Thus may you alſo plant Apples among Plums and Cherries, ſo as you ſuffer not one to ouer-growe or ouer-toppe another; for by pruning, lopping, and ſhredding thoſe that growe too faſt for their fellowes, you may ſtill keepe your trees in ſuch a conformity, as may be both moſt comely for the ſight, and moſt profitable for the yeelding of greater and better ſtore of fruit. Other ſorts of fruit trees you may mixe among theſe, if you pleaſe, as Filberds, Cornellian Cherries in ſtanderds, and Medlers: but Seruice trees, Baye trees, and others of that high ſort, muſt be ſet to guard thereſt. Thus haue I giuen you the faireſt forme could as yet be deuiſed; and from this patterne, if you doe not follow it preciſely, yet by it you may proportion your Orchard, be it large or little, be it walled or hedged.

## Chap. III.

### *Of a nourſery for trees, both from ſowing the kernels, and planting ſtockes to graft vpon.*

ALthough I know the greater ſort (I meane the Nobility and better part of the Gentrie of this Land) doe not intend to keepe a Nurſery, to raiſe vp thoſe trees that they meane to plant their wals or Orchards withall, but to buy them already grafted to their hands of them that make their liuing of it: yet becauſe many Gentlemen and others are much delighted to beſtowe their paines in grafting themſelues, and eſteeme their owne labours and handie worke farre aboue other mens: for their incouragement and ſatisfaction, I will here ſet downe ſome conuenient directions, to enable them to raiſe an Orchard of all ſorts of fruits quickly, both by ſowing the kernels or ſtones of fruit, and by making choiſe of the beſt ſorts of ſtockes to graft on: Firſt therefore to begin with Cherries; If you will make a Nurſery, wherein you may bee ſtored with plenty of ſtockes in a little ſpace, take what quantitie you thinke good of ordinarie wilde blacke Cherrie ſtones, cleanſed from the berries, and ſowe them, or pricke them in one by one on a peece of ground well turned vp, and large enough for the quantitie of ſtones you will beſtowe thereon, from the midſt of Auguſt vnto the end of September, which when they are two or three yeares old, according to their growth, you may remoue them, and ſet them anew in ſome orderly rowes, hauing pruned their tops and their rootes, which at the next yeares growth after the new planting in any good ground, or at the ſecond, will be of ſufficient bigneſſe to graft vpon in the bud what ſorts of Cherries you thinke beſt: and it is fitteſt to graft them thus young, that pruning your ſtockes to raiſe them high, you may graft them at fiue or ſix foote high, or higher, or lower, as you ſhall ſee good, and being thus grafted in the bud, will both more ſpeedily and ſafely bring forward your grafts, and with leſſe danger of loſing your ſtockes, then by grafting them in the ſtocke: for if the bud take not by inoculating the firſt yeare, yet your tree is not loſt, nor put in any hazzard of loſſe; but may be grafted anew the yeare following, if you will, in another place thereof, whereas if you graft in the ſtocke, and it doe not take, it is a great chance if the ſtocke dye not wholly, or at leaſt be not ſo weakened both in ſtrength and height, that it will not bee fit to bee grafted a yeare or two after. In the ſame manner as you doe with the blacke, you may deale with the ordinary Engliſh red Cherrie ſtones, or kernels, but they are not ſo apt to growe ſo ſtraight and high, nor in ſo ſhort a time as the blacke Cherrie ſtones are, and beſides are ſubiect in time to bring out ſuckers from the rootes, to the hinderance of the ſtockes and grafts, or at the leaſt to the deformitie of your Orchard, and more trouble to the Gardiner, to pull or digge them away. Plumme ſtones may bee ordered in this manner likewiſe, but you muſt make choiſe of your Plums; for although euery Plumme is not ſo fit for this purpoſe, as the white Peare Plumme, becauſe it groweth the goaleſt and freeſt, the barke being ſmooth and apteſt to be raiſed, that they may be grafted vpon; yet diuers other Plummes may be taken, if they be not at hand, or to be had, as the blacke and red Peare Plumme, the

white

white and red Wheate Plumme, becauſe they are neareſt in goodneſſe vnto it. Peach ſtones will be ſoone raiſed vp to graft other ſorts of Peaches or Nectorins vpon, but the nature of the Peach roote being ſpongie, is not to abide long. As for Almonds, they will be raiſed from their ſtones to be trees of themſelues; but they will hardly abide the remouing, and leſſe to bee grafted vpon. Apricocke ſtones are the worſt to deale withall of any ſort of ſtone fruit; for although the Apricocke branches are the fitteſt ſtockes to graft Nectorins of the beſt ſorts vpon, yet thoſe that are raiſed from the kernels or ſtones will neuer thriue to be brought on for this purpoſe; but will ſtarue and dye, or hardly grow in a long time to be a ſtraight and fit ſtocke to be grafted, if it be once remoued. Your Cornelian Cherrie trees are wholly, or for the moſt part raiſed from the ſtones or kernels; yet I know diuers doe increaſe them, by laying in their loweſt branches to taks roote: and thus much for ſtone fruits. Now for Apples and Peares, to be dealt withall in the ſame manner as aforeſaid. They vſe to take the preſſing of Crabs whereas Veriuyce is made, as alſo of Cidar and Perry where they are made, and ſowing them, doe raiſe vp great ſtore of ſtockes; for although the beating of the fruit doth ſpoile many kernels, yet there will bee enough left that were neuer toucht, and that will ſpring: the Crabbe ſtockes ſome preferre for the fitteſt, but I am ſure, that the better Apple and Peare kernels will growe fairer, ſtraighter, quicklier, and better to be grafted on. You muſt remember, that after two or three yeares you take vp theſe ſtockes, and when you haue pruned both toppe and roote, to ſet them againe in a thinner and fitter order, to be afterwards grafted in the bud while they are young, as I ſhall ſhew you by and by, or in the ſtocke if you will ſuffer them to growe greater. Now likewiſe to know which are the fitteſt ſtockes of all ſorts to chooſe, thereon to graft euery of theſe ſorts of fruits, is a point of ſome skill indeede; and therefore obſerue them as I doe here ſet them downe: for bee you aſſured, that they are certaine rules, and knowne experiences, whereunto you may truſt without being deceiued. Your blacke Cherrie ſtockes (as I ſaid before) are the fitteſt and beſt for all ſorts of Cherries long to abide and proſper, and euen May or early Cherry will abide or liue longer, being grafted thereon, either in the budde or in the ſtocke, then on the ordinary red Cherry ſtocke; but the red Cherry ſtocke is in a manner the onely tree that moſt Nurſery men doe take to graft May Cherries on in the ſtocke (for it is but a late experience of many, to graft May Cherries in the bud) many alſo doe graft May Cherries on Gaſcoigne Cherry ſtockes, which doe not onely thriue well, but endure longer then vpon any ordinay Cherry ſtocke: For indeede the May Cherries that are grafted vpon ordinary red Cherrie ſtockes, will hardly hold aboue a dozen yeares bearing well, although they come forwarder at the firſt, that is, doe beare ſooner then thoſe that are grafted on Gaſcoigne or blacke Cherry ſtockes; but as they are earlier in bearing, ſo they are ſooner ſpent, and the Gaſcoigne and blacke Cherry ſtockes that are longer in comming forward, will laſt twice or thrice their time; but many more grafts will miſſe in grafting of theſe, then of thoſe red Cherry ſtockes, and beſides, the natures of the Gaſcoigne and blacke Cherry ſtockes are to riſe higher, and make a goodlier tree then the ordinary red ſtocke will, which for the moſt part ſpreadeth wide, but riſeth not very high. The Engliſh red Cherry ſtocke will ſerue very well to graft any other ſort of Cherry vpon, and is vſed in moſt places of this Land, and I know no other greater inconuenience in it, then that it ſhooteth out many ſuckers from the roote, which yet by looking vnto may ſoone bee remoued from doing any harme, and that it will not laſt ſo long as the Gaſcoigne or blacke Cherry ſtocke will. May Cherries thus grafted lowe, doe moſt vſually ſerue to be planted againſt a wall, to bring on the fruit the earlier; yet ſome graft them high vpon ſtandards, although not many, and it is, I thinke, rather curioſity (if they that doe it haue any wals) then anie other matter that cauſeth them thus to doe: for the fruit is naturally ſmall, though early, and the ſtandard Cherries are alwaies later then the wall Cherries, ſo that if they can ſpare any roome for them at their wals, they will not plant many in ſtandards. Now concerning Plummes (as I ſaid before) for the ſowing or ſetting of the ſtones, ſo I ſay here for their choiſe in grafting of them, either in the budde or ſtocke. The white Peare Plumme ſtocke, and the other there mentioned, but eſpecially the white Peare Plumme is the goodlieſt, freeſt, and fitteſt of all the reſt, as well to graft all ſort of Plummes vpon, as alſo to graft Apricockes, which can be handſomely, and to any

good

good purpoſe grafted vpon no other Plum ſtocke, to riſe to beeworth the labour and paine. All ſorts of Plums may be grafted in the ſtocke,and ſo may they alſo in the bud; for I know none of them that will refuſe to be grafted in the bud, if a cunning hand performe it well; that is, to take off your bud cleanely and well, when you haue made choice of a fit cyon: for,as I ſhal ſhew you anon,it is no ſmall peece of cunning to chuſe your cyon that it may yeeld fit buds to graft withall,for euery plum is not of a like aptnes to yeeld them: But Apricocks cannot be grafted in the ſtock for any thing that euer I could heare or learne, but only in the bud, and therefore let your Plum ſtocke bee of a reaſonable ſize for Apricockes eſpecially, and not too ſmall, that the graft ouergrow not the ſtocke, and that the ſtocke bee large enough to nouriſh the graft. As your Plum ſtockes ſerue to graft both Apricockes and Plummes, ſo doe they ſerue alſo very well to graft Peaches of all ſorts; and although Peach ſtockes will ſerue to be grafted with Peaches againe, yet the Peach ſtocke (as I ſaid before) will not endure ſo long as the Plumme ſtocke, and therefore ſerueth but for neceſſity if Plum ſtocks be not ready,or at hand, or for the preſent time, or that they afterwards may graft that ſort of Peach on a Plumme ſtocke: for many might loſe a good fruit, if when they meete with it, and haue not Plumme ſtockes ready to graft it on, they could not be aſſured that it would take vpon another Peach ſtocke or branch, or on the branch of an Apricocke eyther. Plumme ſtockes will ſerue likewiſe very well for ſome ſorts of Nectorins; I ſay, for ſome ſorts,and not for all: the greene and the yellow Nectorin will beſt thriue to be grafted immediately on a Plumme ſtocke; but the other two ſorts of red Nectorins muſt not be immediately grafted on the Plumme ſtocke,but vpon a branch of an Apricocke that hath beene formerly grafted on a Plumme ſtocke, the nature of theſe Nectorins being found by experience to be ſo contrary to the Plum ſtocke, that it will ſterue it, and both dye within a yeare,two or three at the moſt: Diuers haue tryed to graft theſe red Nectorins vpon Peach ſtockes, and they haue endured well a while; but ſeeing the Peach ſtocke will not laſt long it ſelfe, being ouerweake, how can it hold ſo ſtrong a nature as theſe red Nectorins, which will (as I ſaid before) ſterue a Plum ſtocke that is ſufficient durable for any other Plumme?

Apricocke ſtockes from the ſtones are hardly nurſed vp, and worſe to be remoued, and if a red Nectorin ſhould be grafted on an Apricock rayſed from the ſtone,and not remoued,I doubt it might happen with it as it doth with many other trees raiſed from ſtones or kernels, and not remoued, that they would hardly beare fruit: for the nature of moſt trees raiſed from ſtones or kernels, and not remoued, is to ſend great downe-right rootes,and not to ſpread many forwards; ſo that if they be not cut away that others may ſpreade abroad, I haue ſeldome ſeene or known any of them to beare in any reaſonable time; and therefore in remouing, theſe great downe-right rootes are alwayes ſhred away, and thereby made fit to ſhoote others forwards. Hereby you may perceiue, that theſe red Nectorins will not abide to bee grafted vpon any other ſtocke well, then vpon an Apricocke branch, although the green and the yellow (as I ſaid before) will well endure and thriue vpon Plums. The ſuckers or ſhootes both of Plums and Cherries that riſe from their rootes, eyther neare their ſtockes, or farther off, ſo that they bee taken with ſome ſmall rootes to them, will ſerue to bee ſtockes, and will come forward quickly; but if the ſuckers haue no ſmall roots whereby they may comprehend in the ground,it is almoſt impoſſible it ſhould hold or abide. There is another way to rayſe vp eyther ſtockes to graft on, or trees without grafting, which is, by circumciſing a faire and fit branch in this manner: About Midſomer, when the ſappe is thoroughly riſen (or before if the yeare be forward) they vſe to binde a good quantity of clay round about a faire and ſtraight branch, of a reaſonable good ſize or bigneſſe, with ſome conuenient bands,whether it be ropes of hey, or of any other thing,about an handfull aboue the ioynt,where the branch ſpreadeth from the tree,and cutting the barke thereof round about vnder the place where the clay is bound, the ſap is hereby hindered from riſing, or deſcending further then that place ſo circumciſed,whereby it will ſhoote out ſmall knubs and rootes into the clay, which they ſuffer ſo to abide vntill the beginning of winter, whenas with a fine Sawe they cut off that branch where it was circumciſed,and afterwardes place it in the ground where they would haue it to grow, and ſtake it, and binde it faſt, which will ſhoote forth rootes, and will become eyther a faire tree to beare fruite without grafting, or elſe a fit ſtocke to graft on according

ding

ding to the kinde: but oftentimes this kinde of propagation misseth, in that it sendeth not forth rootes sufficient to cause it to abide any long time. Let me yet before I leaue this narration of Plummes, giue you one admonition more, that vpon whatsoeuer Plumme stocke you doe graft, yet vpon a Damson stocke that you neuer striue to graft, for it (aboue all other sorts of Plumme stockes) will neuer giue you a tree worth your labour. It remaineth only of stone fruit, that I speake of Cornelles, which as yet I neuer saw grafted vpon any stocke, being as it should seeme vtterly repugnant to the nature thereof, to abide grafting, but is wholly raysed vp (as I said before) eyther from the stones, or from the suckers or layers. For Peares and Apples your vsuall stockes to graft on are (as I said before, speaking of the nursing vp of trees from the kernels) your Crabbe stockes, and they bee accepted in euery Countrey of this Land as they may conueniently be had, yet many doe take the stockes of better fruit, whether they bee suckers, or stockes raysed from the kernels (and the most common and knowne way of grafting, is in the stocke for all sorts of them, although some doe vse whipping, packing on, or incising, as euery one list to call it: but now we doe in many places begin to deale with Peares and Apples as with other stone fruit, that is, graft them all in the bud, which is found the most compendious and safest way both to preserue your stocke from perishing, and to bring them the sooner to couer the stock, as also to make the goodlier and straighter tree, being grafted at what height you please:) for those stockes that are raysed from the kernels of good fruit (which are for the most part easily knowne from others, in that they want those thornes or prickles the wilde kindes are armed withall:) I say for the most part; for I know that the kernels of some good fruite hath giuen stockes with prickles on them (which, as I thinke, was because that good fruite was taken from a wilde stocke that had not beene long enough grafted to alter his wilde nature; for the longer a tree is grafted, the more strength the fruite taketh from the graft, and the lesse still from the stocke) being smoother and fairer then the wilde kinds, must needes make a goodlier tree, and will not alter any whit the taste of your fruit that is grafted thereon, but rather adde some better relish thereunto; for the Crabbe stockes yeelding harsh fruite, must giue part of their nature to the grafts are set thereon, and therefore the taste or relish, as well as some other naturall properties of most fruits, are somewhat altered by the stocke. Another thing I would willingly giue you to vnderstand concerning your fruits and stockes, that whereas diuers for curiosity and to try experiments haue grafted Cherries vpon Plumme stockes, or Plums on Cherry stockes, Apples vpon Peare stockes, and Peares vpon Apple stockes, some of these haue held the graft a yeare, two or three peraduenture, but I neuer knew that euer they held long, or to beare fruite, much lesse to abide or doe well: bestow not therefore your paines and time on such contrary natures, vnlesse it be for curiositie, as others haue done: Yet I know that they that graft peares on a white thorne stocke haue had their grafts seeme to thriue well, and continue long, but I haue seldome seene the fruite thereof answerable to the naturall wilde Peare stocke; yet the Medlar is knowne to thriue best on a white thorne. And lastly, whereas diuers doe affirme that they may haue not only good stockes to graft vpon, but also faire trees to bear store of fruit from the kernels of Peares or Apples being prickt into the ground, and suffered to grow without remouing, and then eyther grafted or suffered to grow into great trees vngrafted; and for their bearing of fruite, assigne a dozen or twenty yeares from the first setting of the kernels, and abiding vngrafted, I haue not seene or heard that experience to hold certaine, or if it should be so, yet it is too long time lost, and too much fruit also, to waite twenty yeares for that profit may be gained in a great deale of lesse time, and with more certainty. Vnto these instructions let mee adde also one more, which is not much known and vsed, and that is, to haue fruit within foure or fiue years from the first sowing of your stones or kernels in this manner: After your stones or kernels are two or three yeares old, take the faireſt toppe or branch, and graft it as you would doe any other cyon taken from a bearing tree, and looke what rare fruite, eyther Peare or Apple, the kernell was of that you sowed, or Peach or Plum &c. the stone was set, such fruite shall you haue within two or three yeares at the most after the grafting, if it take, and the stocke be good. And thus may you see fruit in farre lesse time then to stay vntill the tree from a kernell or stone beareth fruit of it selfe.

## Chap. IIII.

### *The diuers manners of grafting all sorts of fruits vsed in our Land.*

THe most vsuall manner of grafting in the stocke is so common and well known in this Land to euery one that hath any thing to doe with trees or an Orchard, that I think I shall take vpon mee a needlesse worke to set downe that is so well knowne to most; yet how common soeuer it is, some directions may profit euery one, without which it is not easily learned. And I doe not so much spend my time and paines herein for their sakes that haue knowledge, but for such as not knowing would faine be taught priuately, I meane, to reade the rules of the arte set downe in priuate, when they would refuse to learne of a Gardiner, or other by sight: and yet I discommend not that way vnto them to learne by sight; for one may see more in an instant by sight, then he shall learn by his own practice in a great while, especially if he be a little practised before he see a cunning hand to doe it. There are many other kindes of grafting, which shall be spoken of hereafter, and peraduenture euen they that know it well, may learne something they knew not before.

1. The grafting in the stocke, is, to set the sprigge of a good fruit into the body or stocke of another tree, bee it wilde or other, bee it young or old, to cause that tree to bring forth such fruit as the tree bore from whence you took the sprigge, and not such as the stocke or tree would haue borne, if it had not beene grafted, and is performed in this manner: Looke what tree or stocke you will chuse to graft on, you must with a small fine sawe and very sharpe, whip off, or cut off the head or toppe thereof at what height you eyther thinke best for your purpose, or conuenient for the tree: for if you graft a great tree, you cannot without endangering the whole, cut it downe so low to the ground, as you may without danger doe a small tree, or one that is of a reasonable size; and yet the lower or neerer the ground you graft a young tree, the safer it is both for your stocke and graft, because the sappe shall not ascend high, but soone giue vigour to the graft to take and shoote quickly: After you haue cut off the toppe of your stocke, cut or smooth the head thereof with a sharpe knife, that it may be as plaine and smooth as you can, and then cleaue it with a hammer or mallet, and with a strong knife, cleauer or chessell, either in the middle of it if it be small, or of a reasonable size, or on the sides an inch or more within the barke, if it be great: into both sides of the cleft put your grafts, or into one if the stocke bee smaller; which grafts must bee made fit for the purpose on this fashion: Hauing made choise of your grafts from the toppe branches especially, or from the sides of that tree wherof you would haue the fruit, and that they be of a reasonable good size, not too small or too great for your stockes, and of one or the same yeares shoote; (and yet many doe cut an inch or more of the olde wood with the sprigge of the last yeares growth, and so graft the old and young together (but both are good, and the old wood no better then the young) cut your graft not too long, but with two, three or foure eyes or buds at the most, which at the lower or bigger end for an inch long or more (for the greater stockes, and an inch or lesse for the lesser sort) must be so cut, that it be very thin on the one side from the shoulders downward, and thicker on the other, and thin also at the end, that it may goe downe close into the cleft, and rest at the shoulders on the head of the stocke: but take heede that in cutting your grafts your knife bee very sharpe that you doe not rayse any of the barke, eyther at the sides or the end, for feare of losing both your paines and graft, and stocke too peraduenture; and let not your grafts bee made long before you set them, or else put the ends of them in water to keepe them fresh and cleane: when you set them you must open the cleft of your stocke with a wedge or chessell as most doe, that the graft may goe easily into it, and that the barke of both graft and stocke may ioyne close the one to the other, which without stirring or displacing must bee so left in the cleft, and the wedge or chessell gently pulled forth; but because in the doing hereof consisteth in a manner the whole losse or gaine of your paines, graft and stocke, to preuent which inconuenience I doe vse an iron Instrument, the forme whereof is shewne in the following

lowing page, marked with the letter A, crooked at both ends, and broade like vnto a cheſſell, the one bigger, and the other leſſer, to fit all ſorts of ſtockes, and the iron handle ſomewhat long betweene them both, that being thruſt or knocked downe into the cleft, you may with your left hand open it as wide as is fit to let in your graft, without ſtrayning, which being placed, this iron may bee pulled or knocked vp againe without any mouing of your graft: when you haue thus done, you muſt lay a good handfull or more (according to the bigneſſe of your ſtocke) of ſoft and well moiſtned clay or loame, well tempered together with ſhort cut hey or horſe dung, vpon the head of your ſtocke, as lowe or ſomewhat lower then the cleft, to keepe out all winde, raine or ayre from your graft vntill Midſomer at the leaſt, that the graft be ſhot forth ſomewhat ſtrongly, which then if you pleaſe may be remoued, and the cleft at the head only filled with a little clay to keepe out earewigs, or other things that may hurt your graft.

A. The Iron Inſtrument with cheſſels at each end, the one bigger and the other leſſer, to keepe the cleft of the Tree open vntill the graft bee placed in the ſtocke, which with a knock vpwards will be eaſily taken away.

B. The ſmall Penne-knife with a broad and thinne ended hafte, to raiſe the ſides both of the bud and the down-right ſlit in the body or arme of a Tree to be grafted in the bud.

C. A pen or quil cut halfe round to take off a bud from the branch.

D. An Iuory Inſtrument made to the ſame faſhion.

E. A ſhielde of braſſe made hollow before to be put into the ſlit, to keepe it open vntill the bud be put into its place.

F. The manner of grafting called inciſing or ſplicing.

G. A Ladder made with a ſtoole at the toppe, to ſerue both to graft higher or lower, and alſo to gather fruit without ſpoyling or hurting any buddes or branches of Trees.

1. The firſt ſlit in the body or arme of a Tree to be grafted in the bud with the croſſe cut at the head.
2. The ſame ſlit opened on both ſides, ready to receiue the budde ſhould be put therein: theſe ſmall peeces ſerue as well as trees to ſhew the manner and order of the grafting.
3. The branch of a Tree with one budde cut ready to be taken off, and another not yet touched.
4. The bud cleane taken off from the branch, both the foreſide and backſide.
5. The graft or bud now put into the ſtocke or tree you intend to be grafted: but the binding thereof is omitted.

12. Inarching is another manner of grafting in the ſtocke, and is more troubleſome, and more caſuall alſo then the former, and is rather a curioſity then any way of good ſpeede, certainety or profit, and therefore vſed but of a few. Yet to ſhew you, the man-

manner thereof, it is thus: Hauing a tree well growne, bee it high or low, yet the lower the better, with young branches well ſpread, they vſe to ſet ſtockes round about it, or on the one ſide as you pleaſe; into which ſtockes they ingraft the young branches of the well growne tree as they are growing (before they cut them from the tree) by bowing downe the branch they intend to graft, and putting it into the ſtocke, hauing firſt cut off the head thereof, and cut a notch in the middle of the head a little ſlope on both ſides, wherein the branch muſt be fitted: let the branch be cut thinne on the vnderſide, only of that length as may ſuffice to fit the notch in the ſtocke, leauing about halfe a yarde length of the branch, to riſe aboue or beyond the ſtocke, which beeing bound on, and clayed ouer or couered with red or greene ſoft waxe, they let ſo abide, that if it take in the ſtocke they cut off the branch a little below the grafting place in Nouember following, and remouing the ſtocke, they haue thus gained a grafted and growne tree the firſt yeare: but it is vſually ſeene, that where one branch taketh, three doe miſſe: yet this manner of grafting was much in vſe for May Cherries, when they were firſt known to vs, and the way thought to be a rare manner of grafting to encreaſe them, vntill a better way was found out, which now is ſo common and good alſo, that this is not now ſcarce thought vpon.

3. Another kinde of grafting in the ſtocke is called of ſome whipping, of ſome ſplicing, of others inciſing, and of others packing on (and as I heare, is much vſed in the Weſt parts eſpecially, and alſo in the North parts of this Land) and is performed in this manner: Take and ſlice the branch of a tree (ſo as the branch be not too bigge) or elſe a young tree of two, or three, or foure yeares growth at the moſt, quite off ſlope wiſe, about an inch and a halfe long or more, and cut a deep notch in the middle thereof, then fit into it a graft iuſt of that ſize or bigneſſe, cut on both ſides with ſhoulders, and thin at the end, that it may ioyne cloſe in the notch, and neyther bigger or leſſer, but that the barke of the one may bee fitted iuſt to the barke of the other, the figure wherof is expreſſed at the letters E.F. which ſhew the one to be with a ſhoulder & the other without; binde them gently together with baſt, and put clay or waxe ouer the place, vntill it be taken: this is much vſed of late dayes for ſuch young trees as are riſen of ſtones or kernels after the ſecond or third yeares growth, and thriue very well in that it not only ſaueth much time, but diuers checks by remouing and grafting.

4. Inoculating or grafting in the budde is another manner of grafting, which is the taking of a budde from one tree, and putting it into the barke of another tree, to the end, that thereby you may haue of the ſame kinde of fruit the tree bare from whence the budde was taken; and although it bee ſufficiently knowne in many places of this Land, yet as I vnderſtand, good Gardiners in the North parts, and likewiſe in ſome other places, can ſcarce tell what it meaneth, or at the leaſt how to doe it well. It is performed after a different faſhion from the former, although they all tend vnto one end, which is the propagating of trees. You muſt for this purpoſe obſerue, that for thoſe trees you would graft, either with, or vpon, you chooſe a fit time in Summer, when the ſappe is well riſen, and your graft well ſhot, that the barke will riſe eaſily and cleanly, both of ſtocke and graft, which time I cannot appoint, becauſe both the years doe differ in earlineſſe, and the ſeuerall parts or countries of this Land likewiſe one from another, but moſt vſually in theſe Southerne parts, from the beginning of Iune vnto the end of it, or to the middle of Iuly, or either ſomewhat before or after. Firſt (as I ſaid) hauing taken the fitteſt time of the yeare, you muſt take eſpeciall care, that your grafts be well growne, and of the ſame yeares ſhoote, and alſo that the buds or eyes haue but ſingle leaues at them, as neare as you can: for I would vtterly refuſe thoſe buds that haue aboue two leaues as vnprofitable, either in Peaches or any other fruit; and therefore ſee that your grafts or cyons bee taken from the chiefeſt place of the tree, that is, either from the toppe, or from a ſunnie ſide thereof, and not from the contrarie ſide if you may otherwiſe, not from any vnder-boughes; for ſeeing your graft is ſo ſmall a thing, you had neede take the more care that it be the beſt and faireſt. You muſt to take off this eye or budde from the ſprigge, haue a ſmall ſharpe pen-knife, the end of the haft being made flat and thinne, like a cheſſell or wedge, the figure whereof is ſet forth at the letter B, and a pen or gooſe quill cut, to be leſſe then halfe round, and to be broad at the end, but not ſharpe pointed like a penne, or elſe ſuch a peece of bone or Iuorie made in that faſhion as the quill is, to bee thinne, hollow, or

halfe

halfe round, the figures of both which are marked with the letters C, D. with your knife cut the barke of the bud (hauing firſt cut off the leafe, leauing onely the ſhort foote ſtalke thereof at the bud) about a ſtrawes breadth aboue the eye thereof halfe round, and then from that round or ouerthwart cut, with your knife cut it downe on both ſides of the eye, cloſe to the bud ſlopewiſe about an inch long or thereabouts, that it bee broad at the head aboue the eye, and pointing at the end like a ſheild or ſcutcheon; and then cutting away the reſt of the barke from about it, with the thinne flat end of the haft of your knife raiſe vp both ſides of your bud a little, and with your quill or bone put vnder the barke, raiſe your budde, and thruſt it quite off, beginning at the toppe or head of your eye; but ſee that you thruſt it off cloſe to the wood of the branch or ſprigge, and that you doe not leaue the eye of the budde behinde ſticking vpon the branch; for if that eye be left or loſt, your bud is worth nothing; you muſt caſt it away, and cut another that may haue that eye abiding within the budde on the inſide: you may perceiue if that eye be wanting, if you ſee an emptie hole in the place where the eye ſhould be, to fill it vp on the inſide thereof; thus hauing taken off your bud well and cleanly, which is ſet forth vnto you at the figures 3 and 4. preſently ſet it on the tree you would graft (for your ſmall bud can abide no delay, leſt by taking the ayre too long it become dry, and nothing worth) in this manner: Cut the barke of your tree you would graft in a ſmooth place, at what height you pleaſe, firſt aboue or ouerthwart, and then downe right in the middle thereof, more then an inch long, the figure whereof you ſhall haue at the figure 1. and then raiſe vp both ſides of the barke, firſt one, and then another, with the flat and thinne haft end of your knife, a prettie way inwards (for if the barke will not riſe eaſily, the ſtocke is not then fit to graft vpon) put in your budde into the cleft with the point downewards, holding the ſtalke of the leafe that is with the budde betweene your fingers of the one hand, and opening the cleft with the flat end of your knife with the other hand, that the head of your bud may be put cloſe vnder the ouerthwart cut in the ſtocke or tree (which muſt not be raiſed or ſtirred as the ſides are) & the eye of the bud ſtand iuſt in the middle of the ſlit that is downeright, and then cloſing the barke of the ſtocke or tree ſoftly vnto the bud thus put in with your fingers, let it be bound gently with a ſmall long peece of baſte, or other ſuch like ſoft thing, firſt aboue the eye, & then compaſſing it belowe as cloſe as you can, but not too hard in any caſe, vntil you haue bound it all ouer the ſlit you made, eſpecially the lower end, leſt any winde get in to dry and ſpoile it; and hauing tyed both ends thereof faſt, leaue it ſo for a fornight or ſomewhat more, in which ſpace it will take and hold, if it be well done, which you ſhall perceiue, if the bud abide green, and turne not blacke, when you haue vnlooſed the tying; for if it hold faſt to the tree, and be freſh and good, tye it vp gently againe, and ſo leaue it for a fortnight longer, or a moneth if you will, and then you may take away your binding cleane: this budde will (if no other miſchance happen vnto it) ſpring and ſhoote forth the next yeare, (and ſometimes the ſame yeare, but that is ſeldome) and therefore in the beginning of the yeare, cut off the head of the grafted tree about an handfull aboue the grafted place, vntill the graft be growne ſtrong, and then cut it off cloſe, that the head may be couered with the graft, and doe not ſuffer any buds to ſprout beſides the graft, either aboue or belowe it. If you graft diuers buds vpon one ſtocke (which is the beſt way) let that onely remaine and abide that ſhooteth beſt forth, and rubbe off, or take away the other: the ſeuerall parts of this grafting I haue cauſed to be expreſſed for your further information.

5. Grafting in the ſcutcheon is accounted another kinde of grafting, and differeth verie little from grafting in the budde: the difference chiefly conſiſteth in this, that in ſtead of the downe right ſlit, and that aboue ouerthwart, they take away iuſt ſo much barke of the great tree, as your bud is in bigneſſe, which vſually is a little larger then the former, and placing it therein, they binde it as formerly is ſaid: ſome vſe for this purpoſe a paire of compaſſes, to giue the true meaſure both of bud and ſtocke; this manner of grafting is moſt vſed vpon greater trees, whoſe young branches are too high to graft vpon in the former maner, and whoſe tops they cut off (for the moſt part) at the latter end of the next yeare after the bud is taken: both theſe waies were inuented to ſaue the loſſe of trees, which are more endangered by grafting in the ſtocke, then

then any of these waies; and besides, by these waies you may graft at a farre grea height without losse.

## Chap. V.

### *Of the manner of grafting and propagating all sorts of Roses.*

HAuing now spoken of the grafting of trees, let mee adioyne the properties of Roses, which although they better fit a Garden then an Orchard, yet I could not in a fitter place expresse them then here, both for the name and affinity of grafting, & because I do not expresse it in the first part. All sorts of Roses may be grafted (although all sorts are not, some seruing rather for stockes for others to be grafted on) as easily as any other tree, & is only performed, by inoculating in the same maner I haue set downe in the former Chapter of grafting trees in the bud; for both stocke and budde must bee dealt with after the same fashion. And although some haue boasted of grafting Roses by slicing or whipping, as they call it, or in the stocke, after the first manner, set downe in the former Chapter, yet I thinke it rather a bragge, not hauing seene or heard any true effect proceede from that relation. The sweete Briar or Eglantine, the white and the Damaske Roses, are the chiefest stockes to graft vpon. And if you graft lowe or neare the ground, you may by laying downe that graft within the ground, after it hath bin shot out well, and of a years growth, by pinning it fast downe with short stickes, a thwart or acrosse, cause that grafted branch, by taking roote, to become a naturall Rose, such as the graft was, which being separated and transplanted after it hath taken root wel, will prosper as well as any naturall sucker. And in this maner, by laying downe branchese at length into the ground, if they be full of spreading small branches, you may increase all sorts of Roses quickly and plentifully; for they will shoote forth rootes at the ioynt of euery branch: But as for the manner of grafting white Roses or Damaske vpon Broome stalkes or Barbary bushes, to cause them to bring forth double yellow Roses, or vpon a Willowe, to beare greene Roses, they are all idle conceits, as impossible to be effected, as other things, whereof I haue spoken in the ninth Chapter of my first part, concerning a Garden of flowers, vnto which I referre you to be satisfied with the reasons there alledged. And it is the more needlesse, because we haue a naturall double yellow Rose of it owne growing. The sowing of the seedes of Roses (which are sometimes found vpon most sorts of Roses, although not euery yeare, and in euerie place) hath bin formerly much vsed; but now the laying downe of the young shootes is a way for increase so much vsed, being safe and verie speedie to take, especially for those Roses that are not so apt to giue suckers, that it hath almost taken quite away the vse of sowing of the seedes of Roses, which yet if anie one bee disposed to make the triall, they must gather the seede out of the round heads, from amongst the doune, wherein they lye verie like vnto the berries of the Eglantine or sweete Briar bush, and especially of those Roses that bee of the more single kindes, which are more apt to giue berries for seed then the more double, although sometimes the double Roses yeeld the like heads or berries. Their time of sowing is in the end of September (yet some reserue them vntill February) and their manner of noursing is to bee transplanted, after the first or second yeares growth, and tended carefully, that while they are young they be not lost for want of moisture in the dry time of Summer.

## CHAP. VI.

### *Certaine rules and obſeruations in and after grafting, not remembred in the former Chapter.*

THe time of ſome manners of grafting being not mentioned before, muſt here be ſpoken of. For the grafting of all ſorts of trees in the ſtocke, the moſt vſuall time is from the middle of February vntill the middle of March, as the yeare and the countrie is more forward or backward, with vs about London wee neuer paſſe midde March: but becauſe the May Cherrie is firſt ripe, and therefore of a very forward nature, it doth require to be grafted ſomewhat ſooner then others. The time of gathering likewiſe, or cutting your grafts for grafting in the ſtocke, is to be obſerued, that they bee not long gathered before they bee grafted, for feare of being too dry, which I commend, howſoeuer diuers ſay, if they be long kept they are not the worſe; and therefore if you be forced to haue your grafts from farre, or by ſome other chance to keepe them long, be carefull to keepe them moiſt, by keeping their ends ſtucke in moiſt clay; but if neare hand, neglect no time I ſay after the cutting of them for their grafting, but either the ſame, or the next day, or verie ſpeedily after, in the meane time being put into the ground to keepe them freſh. The grafts taken from old trees, becauſe they are ſtronger, and ſhoote forth ſooner, are to bee ſooner grafted then thoſe that are taken from younger trees: of a good branch may bee made two, and ſometimes three grafts ſufficient for anie reaſonable ſtocke. For whipping, the time is ſomewhat later then grafting in the ſtocke, becauſe it is performed on younger trees, which (as I ſaid before) doe not ſo early bud or ſhoote forth as the elder. Inarching likewiſe is performed much about the later end of the grafting time in the ſtocke; for being both kindes thereof they require the ſame time of the yeare. The times of the other manners of graftings are before expreſſed, to bee when they haue ſhot forth young branches, from whence your buds muſt be taken; and therefore need not here againe to be repeated. If a graft in the ſtocke doth happen not to ſhoote forth when others do (ſo as it holdeth green) it may perchance ſhoot out a moneth or two after, & do well, or elſe after Midſummer, when a ſecond time of ſhooting, or the after Spring appeareth: but haue an eſpeciall care, that you take not ſuch a graft that ſhal haue nothing but buds for flowers vpon it, and not an eye or bud for leaues (which you muſt be carefull to diſtinguiſh) for ſuch a graft after it hath ſhot out the flowers muſt of neceſſitie dye, not hauing wherewith to maintaine it ſelfe. Alſo if your good graft doe miſſe, and not take, it doth hazzard your ſtocke at the firſt time, yet manie ſtockes doe recouer to be grafted the ſecond time; but twice to faile is deadly, which is not ſo in the inoculating of buds in the greene tree: for if you faile therein three, or three times three, yet euerie wound being ſmall, and the tree ſtill growing greene, will quickly recouer it, and not be afterwards ſeen. Some vſe to graft in the ſtocke the ſame yeare they remoue the ſtocke, to ſaue time, & a ſecond checke by grafting; but I like better both in grafting in the ſtocke, and in the bud alſo, that your trees might be planted in the places where you would haue them growe, for a yeare or two at the leaſt before you graft them, that after grafting there ſhould be no remouall, I neede not be tedious, nor yet I hope verie ſollicitous to remember many other triuiall, or at the leaſt common knowne things in this matter. Firſt, for the time to remoue trees, young or old, grafted or vngrafted, to be from a fortnight after Michaelmas vntill Candlemas, or if neede be, ſomewhat after, yet the ſooner your remoue is, the better your trees will thriue, except it be in a very moiſt ground. For the manner or way to ſet them: *viz.* in the high and dry grounds ſet them deeper, both to haue the more moiſture, and to be the better defended from windes; and in the lower and moiſter grounds ſhallower, and that the earth be mellow, well turned vp, and that the finer earth bee put among the ſmall rootes, wherein they may ſpread, and afterwards gently troden downe, that no hollowneſſe remaine among the rootes: as alſo that after ſetting (if the time be not ouermoiſt) there may be ſome water powred to the rootes, to moiſten and faſten them the better; and in the dry time of Summer, after the ſetting, let them not want moiſture, if you will

haue

haue them thriue and prosper; for the want thereof at that time, hath often killed manie a likely tree. To stake and fence them also if neede bee after they are new set, and so to continue for two or three yeares after, is verie expedient, lest windes or other casualties spoile your paines, and ouerthrow your hopes. And likewise to defend your grafts from birds lighting on them, to breake or displace them, to sticke some prickes or sharpe pointed stickes longer then your graft into your clay, that so they may be a sure defence of it: As also to tye some woollen cloathes about the lower end of your stockes, or thrust in some thornes into the ground about the roores, to defend them from hauing their barkes eaten by Conies, or hurt by some other noisome vermine.

## Chap. VII.

### *Obseruations for the dressing and well keeping of Trees and an Orchard in good order.*

THere are two manner of waies to dresse and keepe trees in good order, that they may bee both gracefull and fruitfull; the one is for wall-trees, the other is for standards: for as their formes are different, so is their keeping or ordering. Wall trees, because they are grafted lowe, and that their branches must be plasht or tackt vnto the wall to fasten them, are to be so kept, that all their branches may be suffered to growe, that shoote forth on either side of the bodie, and led either along the wall, or vpright, and one to lappe ouer or vnder another as is conuenient, and still with peeces of lists, parings of felt, peeces of soft leather, or other such like soft thing compassing the armes or branches, fastened with small or great nailes, as neede requireth, to the wals, onely those buds or branches are to be nipped or cut off, that shoot forward, and will not so handsomely be brought into conformity, as is fitting; yet if the branches growe too thicke, to hinder the good of the rest, or too high for the wall, they may, nay they must be cut away or lopped off: and if anie dead branches also happen to be on the trees, they must be cut away, that the rest may haue the more libertie to thriue. Diuers also by carefully nipping away the waste and superfluous buds, doe keepe their trees in conformity, without much cutting. The time to pruine or plash, or tye vp wall trees, is vsually from the fall of the leafe, to the beginning of the yeare, when they begin to blossome, and most especially a little before or after Christmas: but in any case not too late, for feare of rubbing off their buds. Some I know doe plash and tye vp their wall trees after bearing time, while the leaues are greene, and their reason is, the buds are not so easie or apt to bee rubbed from the branches at that time, as at Christmas, when they are more growne: but the leaues must needes be very cumbersome, to hinder much both the orderly placing, and close fastening of them to the wall. This labour you must performe euery yeare in its due time; for if you shall neglect and ouerslip it, you shall haue much more trouble, to bring them into a fit order againe, then at the first. The standard trees in an Orchard must be kept in another order; for whereas the former are suffered to spread at large, these must be pruined both from superfluous branches that ouerload the trees, & make them lesse fruitfull, as well as lesse sightly, and the vnder or water boughes likewise, that drawe much nourishment from the trees, and yet themselues little the better for it, I meane to giue fruit. If therefore your Orchard consist of young trees, with a little care and paines it may bee kept in that comely order and proportion it was first destined vnto; but if it consist of old growne trees, they will not without a great deale of care and paines be brought into such conformitie, as is befitting good and comely trees: for the marke of those boughes or branches that are cut off from young trees, will quickly be healed againe, the barke growing quickly ouer them, whereby they are not worse for their cutting; but an old tree if you cut off a bough, you must cut it close and cleanly, and lay a searcloth of tallow, waxe, and a little pitch melted together vpon the place, to keepe off both the winde, sunne, and raine, vntill the barke haue couered it ouer againe: and in this manner you must deale with all such short stumps of branches, as are either broken short off with the winde, or by careleslnesse or

want

want of skill, or elſe ſuch armes or branches as are broken off cloſe, or ſliued from the body of the tree: for the raine beating and falling into ſuch a place, will in ſhort time rotte your tree, or put it in danger, beſides the deformity. Some vſe to fill vp ſuch an hole with well tempered clay, and tacke a cloth or a peece of leather ouer it vntill it be recouered, and this is alſo not amiſſe. Your young trees, if they ſtand in anie good ground, will bee plentifull enough in ſhooting forth branches; bee carefull therefore if they growe too thicke, that you pruine away ſuch as growe too cloſe (and will, if they be ſuffered, ſpoile one another) as they may be beſt ſpared, that ſo the ſunne, ayre, and raine may haue free acceſſe to all your branches, which will make them beare the more plentifully, and ripen them the ſooner and the more kindly. If anie boughes growe at the toppe too high, cut them alſo away, that your trees may rather ſpread then growe too high. And ſo likewiſe for the vnder boughes, or anie other that by the weight of fruit fall or hang downe, cut them off at the halfe, and they will afterwards riſe and ſhoote vpwards. You ſhall obſerue, that at all thoſe places where anie branches haue been cut away, the ſappe will euer bee readie to put forth: if therefore you would haue no more branches riſe from that place, rubbe off or nippe off ſuch buddes as are not to your minde, when they are new ſhot: and thus you may keep your trees in good order with a little paines, after you haue thus pruined and dreſſed them. One other thing I would aduertiſe you of, and that is how to preſerue a fainting or decaying tree which is readie to periſh, if it be not gone too farre or paſt cure, take a good quantitie of oxe or horſe bloud, mixe therewith a reaſonable quantitie of ſheepe or pigeons dung, which being laid to the roote, will by the often raines and much watering recouer it ſelfe, if there bee anie poſſibilitie; but this muſt bee done in Ianuarie or Februarie at the furtheſt.

## CHAP. VIII.

### *Diuers other obſeruations to be remembred in the well keeping of an Orchard.*

THere be diuers other things to be mentioned, whereof care muſt be had, either to doe or auoide, which I thinke fit in this Chapter promiſcuouſly to ſet down, that there may be nothing wanting to furniſh you with ſufficient knowledge of the care, paines, and caſualties that befall an Orchard: for it hath many enemies, and euery one laboureth as much as in them lye, to ſpoile you of your pleaſure, or profit, or both, which muſt bee both ſpeedily and carefully preuented and helped; and they are theſe: Moſſe, Caterpillars, Ants, Earwigs, Snailes, Moales, and Birds. If Moſſe begin to ouergrowe your trees, looke to it betimes, leſt it make your trees barren: Some vſe to hacke, and croſſehacke, or cut the barke of the bodies of their trees, to cauſe it fall away; but I feare it may endanger your trees. Others do either rubbe it off with a haire cloth, or with a long peece of wood formed like a knife, at the end of a long ſticke or pole, which if it bee vſed cautelouſly without hurting the buds, I like better. Caterpillars, ſome ſmoake them with burning wet ſtrawe or hay, or ſuch like ſtuffe vnder the trees; but I doe not greatly like of that way: others cut off the boughes whereon they breed, and tread them vnder their feete, but that will ſpoile too manie branches; and ſome kill them with their hands: but ſome doe vſe a new deuiſed way, that is, a pompe made of lattin or tin, ſpout-faſhion, which being ſet in a tubbe of water vnder or neare your trees, they will cauſe the water to riſe through it with ſuch a force, and through the branches, that it will waſh them off quickly. To deſtroy Ants, that eate your fruit before and when it is ripe, ſome vſe to annoint the bodies of their trees with tarre, that they may not creepe vp on the branches; but if that doe not helpe, or you will not vſe it, you muſt be carefull to finde out their hill, and turne it vp, pouring in ſcalding water, either in Summer, but eſpecially if you can in Winter, and that will ſurely deſtroy them. I haue ſpoken of Earwigs in the firſt part of this worke, entreating of the annoyances of Gilloflowers, and therefore I referre you thereunto: yet one way more I will

will here relate which ſome doe vſe, and that is with hollow canes of halfe a yard long or more, open at both ends for them to creepe in, and ſtucke or laid among the branches of your trees, will ſoone drawe into them many Earwigs, which you may ſoone kill, by knocking the cane a little vpon the ground, and treading on them with your foote. Snailes muſt be taken with your hands, and that euerie day, eſpecially in the morning when they will be creeping abroad. Moales by running vnder your trees make them leſſe fruitfull, and alſo put them in danger to be blowne downe, by leauing the ground hollow, that thereby the rootes haue not that ſtrength in the ground, both to ſhoote and to hold, that otherwiſe they might haue. Some haue vſed to put Garlicke, and other ſuch like things into their holes, thinking thereby to driue them away, but to no purpoſe: others haue tryed manie other waies; but no way doth auaile anie thing, but killing them either with a Moale ſpade, or a trappe made for the purpoſe as manie doe know: and they muſt bee watched at their principall hill, and trenched round, and ſo to be caught. Birds are another enemie both to your trees and fruit; for the Bullfinch will deſtroy all your ſtone fruit in the budde, before they flower, if you ſuffer them, and Crowes, &c. when your Cherries are ripe: for the ſmaller birds, Lime twigs ſet either neare your trees, or at the next water where they drinke, will helpe to catch them and deſtroy them. And for the greater birds, a ſtone bowe, a birding or fowling peece will helpe to leſſen their number, and make the reſt more quiet: or a mill with a clacke to ſcarre them away, vntill your fruit be gathered. Some other annoyances there are, as ſuckers that riſe from the rootes of your trees, which muſt be taken away euerie yeare, and not ſuffered to growe anie thing great, for feare of robbing your trees of their liuelihood. Barke bound, is when a tree doth not ſhoote and encreaſe, by reaſon the barke is as it were drie, and will not ſuffer the ſappe to paſſe vnto the branches: take a knife therefore, and ſlit the barke downe almoſt all the length of the tree in two or three places, and it will remedy that euill, and the tree will thriue and come forward the better after. Barke pilled is another euill that happeneth to ſome trees, as well young as old, either by reaſon of caſuall hurts, or by the gnawing of beaſts, howſoeuer it bee, if it bee anie great hurt, lay a plaiſter thereon made of tallow, tarre, and a little pitch, and binde it thereto, letting it ſo abide vntill the wound bee healed: yet ſome doe only apply a little clay or loame bound on with ropes of hay. The Canker is a ſhrewd diſeaſe when it happeneth to a tree; for it will eate the barke round, and ſo kill the very heart in a little ſpace. It muſt be looked vnto in time before it hath runne too farre; moſt men doe wholly cut away as much as is fretted with the Canker, and then dreſſe it, or wet it with vinegar or Cowes piſſe, or Cowes dung and vrine, &c. vntill it be deſtroyed, and after healed againe with your ſalue before appointed. There are yet ſome other enemies to an Orchard: for if your fence be not of bricke or ſtone, but either a mudde wall, or a quicke ſet or dead hedge, then looke to it the more carefully, and preuent the comming in of either horſe, or kine, ſheepe, goates, or deere, hare, or conie; for ſome of them will breake through or ouer to barke your trees, and the leaſt hole almoſt in the hedge will giue admittance to hares and conies to doe the like. To preuent all which, your care muſt be continuall to watch them or auoide them, and to ſtoppe vp their entrance. A dogge is a good ſeruant for many ſuch purpoſes, and ſo is a ſtone bowe, and a peece to make vſe of as occaſion ſhall ſerue. But if you will take that medicine for a Canker ſpoken of before, which is Cowes dung and vrine mixed together, and with a bruſh waſh your trees often to a reaſonable height, will keepe hares and conies from eating or barking your trees. Great and cold windes doe often make a great ſpoile in an Orchard, but great trees planted without the compaſſe thereof, as Wall-nuts, Oakes, Elmes, Aſhes, and the like, will ſtand it in great ſtead, to defend it both early and late. Thus haue I ſhewed you moſt of the euils that may happen to an Orchard, and the meanes to helpe them, and becauſe the number is great and daily growing, the care and paines muſt be continuall, the more earneſt and diligent, leſt you loſe that in a moment that hath been growing many yeares; or at the leaſt the profit or beauty of ſome yeares fruit.

CHAP.

## CHAP. IX.

### *The manner and way how to plant, order, and keepe other trees that beare greene leaues continually.*

THe way to order thoſe trees that beare their leaues greene continually, is differing from all others that doe not ſo : for neyther are they to bee planted or remoued at the time that all other trees are ſet, nor doe they require that manner of dreſſing, pruining and keeping, that others doe. And although many ignorant perſons and Gardiners doe remoue Bay trees, and are ſo likewiſe perſwaded that all other trees of that nature, that is, that carry their greene leaues continually, may bee remoued in Autumne or Winter, as well as all other trees may bee; yet it is certaine it is a great chance if they doe thriue and proſper that are ſet at that time, or rather it is found by experience, that ſcarce one of ten proſpereth well that are ſo ordered. Now in regard that there be diuers trees and ſhrubs mentioned here in this booke that beare euer greene leaues, wherein there is very great beauty, and many take pleaſure in them ; as the ordinary Bay, the Roſe Bay, and the Cherry Bay trees, the Indian Figge, the Cypreſſe, the Pine tree, the Mirtle and dwarfe Boxe, and many others ; I will here ſhew you how to plant and order them, as is fitteſt for them. For in that they doe not ſhed their greene leaues in winter as other trees doe, you may in reaſon be perſwaded that they are of another nature; and ſo they are indeede: for ſeeing they all grow naturally in warme Countries, and are from thence brought vnto vs, we muſt both plant them in a warmer place, and tranſplant them in a warmer time then other trees be, or elſe it is a great hazzard if they doe not periſh and dye, the cold and froſts in the winter being able to pierce them through, if they ſhould bee tranſplanted in winter, before they haue taken roote. You muſt obſerue and take this therefore for a certaine rule, that you alwaies remoue ſuch trees or ſhrubbes as are euer greene in the ſpring of the yeare, and at no time elſe if you will doe well, that is, from the end of March, or beginning of Aprill, vnto the middle or end of May, eſpecially your more dainty and tender plants, ſhadowing them alſo for a while from the heate of the Sun, and giuing them a little water vpon their planting or tranſplanting ; but ſuch water as hath not preſently been drawn from a Well or Pumpe, for that will go neer to kill any plant, but ſuch water as hath ſtood in the open ayre for a day at the leaſt, if not two or three. Yet for dwarfe Boxe I confeſſe it may endure one moneth to be earlier planted then the reſt, becauſe it is both a more hardy and lowe plant, and thereby not ſo much ſubiect to the extremitie of the colde : but if you ſhould plant it before winter, the froſts would raiſe it out of the ground, becauſe it cannot ſo ſoone at that time of the yeare take roote, aud thereby put it in danger to be loſt. Moreouer all of them will not abide the extremitie of our winter froſts, and therefore you muſt of neceſſity houſe ſome of them, as the Roſe Bay, Mirtle, and ſome others, but the other ſorts being ſet where they may bee ſomewhat defended from the cold windes, froſtes, and ſnow in winter, with ſome couering or ſhelter for the time, will reaſonably well endure and beare their fruit, or the moſt of them. If any be deſirous to be furniſhed with ſtore of theſe kinds of trees that will be nourſed vp in our Country, he may by ſowing the ſeed of them in ſquare or long woodden boxes or cheſts made for that purpoſe, gaine plenty of them : but hee muſt be carefull to couer them in winter with ſome ſtraw or fearne, or beane hame, or ſuch like thing layd vpon croſſe ſticks to beare it vp from the plants, and after two or three yeares that they are growne ſomewhat great and ſtrong, they may bee tranſplanted into ſuch places you meane they ſhall abide : yet it is not amiſſe to defend them the firſt yeare after they are tranſplanted, for their more ſecuritie : the ſeedes that are moſt vſually ſowen with vs, are, the Cypreſſe tree, the Pine tree, the Baye, the Pyracantha or prickly Corall tree, and the Mirtle : the Roſe Bay I haue had alſo riſen from the ſeede that was freſh, and brought me from Spaine. But as for Orenge trees, becauſe they are ſo hardly preſerued in this our cold climate (vnleſſe it bee with ſome that doe beſtow the houſing of them, beſides a great deale more of care and reſpect vnto them) from the bitterneſſe of our cold long winter weather (although their

kernels

kernels being put into the ground in the Spring or Summer, and if care bee had of them and conuenient keeping, will abide, and by grafting the good fruite on the crab ſtocke they may bee in time nurſed vp) I doe not make any other eſpeciall account of them, nor giue you any further relation of their ordering. Now for the ordering of theſe trees after they are eyther planted of young ſets, or tranſplanted from the ſeede, it is thus : Firſt for Bay trees, the moſt vſuall way is to let them grow vp high to bee trees, and many plant them on the North or Eaſt ſide of their houſes that they may not bee ſcorched with the Sunne; but the bitter winters which we often haue, doe pinch them ſhrewdly, inſomuch that it killeth euen well growne trees ſometimes downe to the roote : but ſome doe make a hedge of them being planted in order, and keep them low by lopping of them continually, which will make them buſh and ſpread. The Cypreſſe tree is neuer lopped, but ſuffered to grow with all the branches from a foote aboue the ground, if it may be, ſtraight vpright; for that is his natiue grace and greateſt beautie, and therefore the more branches doe dye that they muſt bee cut away, the more you deforme his propertie. The Pine tree may be vſed in the ſame manner, but yet it wil better endure to ſuſtaine pruining then the Cypreſſe, without any ſuch deformitie. The Lauroceraſus or Cherry Bay may be diuerſly formed, that is, it may be either made to grow into a tall tree by ſhredding ſtill away the vnder branches, or elſe by ſuffering all the branches to grow to be a low or hedge buſh, & both by the ſuckers and by laying downe the lower branches into the earth, you may ſoone haue much increaſe ; but this way will cauſe it to bee the longer before it beare anie fruit. The Roſe Baye will verie hardlie bee encreaſed either by ſuckers or by layers, but muſt bee ſuffered to grow without lopping, topping or cutting. The Pyracantha or Prickly Corall tree may bee made to grow into a reaſonable tall tree by ſhredding away the lower branches, or it may be ſuffered to grow lowe into an hedge buſh, by ſuffering all the branches to grow continually, you may alſo propagate it by the ſuckers, or by laying downe the lower branches. The Myrtle of all ſorts abideth a low buſh ſpreading his branches full of ſweete leaues and flowers, without anie great encreaſe of it ſelfe, yet ſometimes it giueth ſuckers or ſhootes from the rootes : but for the more ſpeedie propagating of them, ſome doe put the cuttings of them into the earth, and thereby increaſe them. There are ſome other trees that are not of any great reſpect, as the Yew tree, and the Savine buſh, both which may be encreaſed by the cuttings, and therefore I need not make any further relation or amplification of them, and to ſay thus much of them all, is (I thinke) ſufficient for this Worke.

## Chap. X.

### *The ordering, curing, and propagating Vines of all ſorts.*

IN moſt places of this countrie there is ſmall care or paines taken about the ordering of Vines : it ſufficeth for the moſt part with them that haue anie, to make a frame for it to ſpread vpon aboue a mans height, or to tacke it to a wall or window, &c. and ſo to let it hang downe with the branches and fruit, vntill the weight thereof, and the force of windes doe teare it downe oftentimes, and ſpoile the grapes : and this way doth ſomewhat reſemble that courſe that the Vineyard keepers obſerue in the hot countries of Syria, Spaine, and Italy, and in the furtheſt parts of France as I hear likewiſe : for in moſt of theſe hot countries they vſe to plant an Oliue betweene two Vines, and let them runne thereupon. But manie of the other parts of France, &c. doe not ſuffer anie trees to growe among their Vines ; and therefore they plant them thicke, and pruine them much and often, and keepe them lowe in compariſon of the other way, faſtening them to pearches or poles to hold them vp. And according to that faſhion many haue aduentured to make Vineyards in England, not onely in theſe later daies, but in ancient times, as may wel witneſſe the ſundrie places in this Land, entituled by the name of Vineyards; and I haue read, that manie Monaſteries in this Kingdome hauing Vineyards, had as much wine made therefrom, as ſufficed their couents yeare by yeare : but long ſince they haue been deſtroyed, and the knowledge how to order a Vineyard is alſo vtterly periſhed with them. For although diuers, both Noble

bles and Gentlemen, haue in these later times endeauoured to plant and make Vineyards, and to that purpose haue caused Frenchmen, being skilfull in keeping and dressing of Vines, to be brought ouer to performe it, yet either their skill failed them, or their Vines were not good, or (the most likely) the soile was not fitting, for they could neuer make anie wine that was worth the drinking, being so small and heartlesse, that they soone gaue ouer their practice. And Indeede the soile is a maine matter to bee chiefly considered to seate a Vineyard vpon: for euen in France and other hot countries, according to the nature of the soile, so is the rellish, strength, and durabilitie of the wine. Now although I think it a fruitlesse labour for any man to striue in these daies to make a good Vineyard in England, in regard not only of the want of knowledge, to make choise of the fittest ground for such Vines as you would plant therupon, but also of the true maner of ordering them in our country; but most chiefly & aboue all others, that our years in these times do not fal out to be so kindly and hot, to ripen the grapes, to make anie good wine as formerly they haue done; yet I thinke it not amisse, to giue you instructions how to order such Vines as you may nourse vp for the pleasure of the fruit, to eate the grapes being ripe, or to preserue and keepe them to bee eaten almost all the winter following: And this may be done without any great or extraordinarie paines. Some doe make a lowe wall, and plant their Vines against it, and keepe them much about the height thereof, not suffering them to rise much higher: but if the high bricke or stone wals of your Garden or Orchard haue buttresses thereat, or if you cause such to bee made, that they bee somewhat broade forwards, you may the more conueniently plant Vines of diuers sorts at them, and by sticking down a couple of good stakes at euery buttresse, of eight or ten foot high aboue ground, tacking a few lathes acrosse vpon those stakes, you may therunto tye your Vines, & carry them theron at your pleasure: but you must be carefull to cut them euery year, but not too late, and so keepe them downe, and from farre spreading, that they neuer runne much beyond the frame which you set at the buttresses: as also in your cutting you neuer leaue too many ioynts, nor yet too few, but at the third or fourth ioint at the most cut them off. I doe aduise you to these frames made with stakes and lathes, for the better ripening of your grapes: for in the blooming time, if the branches of your vines bee too neare the wall, the reflection of the Sunne in the day time, and the colde in the night, doe oftentimes spoile a great deale of fruit, by piercing and withering the tender foot-stalkes of the grapes, before they are formed, whereas when the blossomes are past, and the fruit growing of some bignesse, then all the heate and reflection you can giue them is fit, and therefore cut away some of the branches with the leaues, to admit the more Sunne to ripen the fruit. For the diuers sorts of grapes I haue set them downe in the Booke following, with briefe notes vpon euerie of them, whether white or blacke, small or great, early or late ripe; so that I neede not here make the same relation again. There doth happen some diseases to Vines sometimes, which that you may helpe, I thinke it conuenient to informe you what they are, and how to remedy them when you shall be troubled with any such. The first is a luxurious spreading of branches and but little or no fruit: for remedie whereof, cut the branches somewhat more neere then vsuall, and bare the roote, but take heed of wounding or hurting it, and in the hole put either some good old rotten stable dung of Horses, or else some Oxe blood new taken from the beasts, and that in the middle of Ianuarie or beginning of Februarie, which being well tempered and turned in with the earth, let it so abide, which no doubt, when the comfort of the blood or dung is well soaked to the bottome by the raines that fall thereon, will cause your Vine to fructifie againe. Another fault is, when a Vine doth not bring the fruit to ripenesse, but either it withereth before it be growne of any bignesse, or presently after the blooming: the place or the earth where such a Vine standeth, assuredly is too cold, and therefore if the fault bee not in the place, which cannot bee helped without remouing to a better, digge out a good quantity of that earth, and put into the place thereof some good fresh ground well heartned with dung, and some sand mixed therewith (but not salt or salt water, as some doe aduise, nor yet vrine, as others would haue) and this will hearten and strengthen your Vine to beare out the frut vnto maturitie. When the leaues of a Vine in the end of Summer or in Autumne, vntimely doe turne either yellow or red, it is a great signe the earth is

too hot and drie ; you muſt therefore in ſtead of dung and ſand, as in the former defect is ſaid, put in ſome freſh loame or ſhort clay, well mixed together with ſome of the earth, and ſo let them abide, that the froſts may mellow them. And laſtly, a Vine ſometimes beareth ſome ſtore of grapes, but they are too many for it to bring to ripeneſſe ; you ſhall therefore helpe ſuch a Vine (which no doubt is of ſome excellent kinde, for they are moſt vſually ſubiect to this fault) by nipping away the bloſſomes from the branches, and leauing but one or two bunches at the moſt vpon a branch, vntill the Vine be growne older, and thereby ſtronger, and by this meane inured to beare out all the grapes to ripeneſſe. Theſe be all the diſeaſes I know doe happen to Vines: for the bleeding of a Vine it ſeldome happeneth of it ſelfe, but commeth either by cutting it vntimely, that is, too late in the yeare, (for after Ianuarie, if you will be well aduiſed, cut not any Vine) or by ſome caſuall or wilfull breaking of an arme or a branch. This bleeding in ſome is vnto death, in others it ſtayeth after a certaine ſpace of it ſelfe: To helpe this inconuenience, ſome haue ſeared the place where it bleedeth with an hot iron, which in many haue done but a little good ; others haue bound the barke cloſe with packe-thred to ſtay it ; and ſome haue tied ouer the place, being firſt dried as well as may bee, a plaiſter made with waxe roſſen and turpentine while it is warme. Now for the propagating of them: You muſt take the faireſt and goaleſt ſhot branches of one yeares growth, and cut them off with a peece of the old wood vnto it, and theſe being put into the ground before the end of Ianuarie at the furtheſt, will ſhoote forth, and take roote, and ſo become Vines of the ſame kinde from whence you tooke them. This is the moſt ſpeedy way to haue increaſe: for the laying downe of branches to take roote, doth not yeelde ſuch ſtore ſo plentifully, nor doe ſuckers riſe from the rootes ſo aboundantly ; yet both theſe waies doe yeelde Vines, that being taken from the old ſtockes will become young plants, fit to bee diſpoſed of as any ſhall thinke meete.

## Chap. XI.

### *The way to order and preſerue grapes, fit to be eaten almoſt all the Winter long, and ſometimes vnto the Spring.*

Although it bee common and vſuall in the parts beyond the Sea to dry their grapes in the Sunne, thereby to preſerue them all the year, as the Raiſins of the Sunne are, which cannot bee done in our Countrie for the want of ſufficient heate thereof at that time : or otherwiſe to ſcald them in hot water (as I heare) and afterwards to dry them, and ſo keepe them all the yeare, as our Malaga Raiſins are prepared that are packed vp into Frayles: yet I doe intend to ſhew you ſome other waies to preſerue the grapes of our Countrie freſh, that they may be eaten in the winter both before and after Chriſtmas with as much delight and pleaſure almoſt, as when they were new gathered. One way is, when you haue gathered your grapes you intend to keepe, which muſt be in a dry time, and that all the ſhrunke, dried, or euill grapes in euery bunch be picked away, and hauing prouided a veſſell to hold them, be it of wood or ſtone which you will, and a ſufficient quantitie of faire and cleane drie ſand ; make *ſtratum ſuper ſtratum* of your grapes and the ſand, that is, a lay of ſand in the bottome firſt, and a lay of grapes vpon them, and a lay or ſtrowing againe of ſand vpon thoſe grapes, ſo that the ſand may couer euery lay of grapes a fingers breadth in thickneſſe, which being done one vpon another vntill the veſſell be full, and a lay of ſand vppermoſt, let the veſſell be ſtopped cloſe, and ſet by vntill you pleaſe to ſpend them, being kept in ſome drie place and in no ſellar : let them bee waſhed cleane in faire water to take away the ſand from ſo many you will ſpend at a time. Another way is (which Camerarius ſetteth downe he was informed the Turkes vſe to keepe grapes all the winter vnto the next ſummer) to take ſo much meale of Muſtard ſeede, as will ſerue to ſtrow vpon grapes, vntill they haue filled their veſſels, whereon afterwards they poure new wine before it hath boiled, to fill vp their veſſels therwith, and being ſtopped vp cloſe, they keepe them a certaine time, and ſelling them with their liquour to them that will

vſe

vſe them, they doe waſh the ſeedes or meale from them when they vſe them. Another way is, that hauing gathered the faireſt ripe grapes, they are to be caſt vpon threds or ſtrings that are faſtened at both ends to the ſide walks of a chamber, neere vnto the ſeeling thereof, that no one bunch touch another, which will bee ſo kept a great while, yet the chamber muſt be well defended from the froſts, and cold windes that pierce in at the windowes, leſt they periſh the ſooner : and ſome will dippe the ends of the branches they hang vp firſt in molten pitch, thinking by ſearing vp the ends to keepe the bunches the better; but I doe not ſee any great likelihood therein. Your chamber or cloſet you appoint out for this purpoſe muſt alſo bee kept ſomewhat warme, but eſpecially in the more cold and froſtie time of the yeare, leſt it ſpoile all your coſt and paines, and fruſtrate you of all your hopes : but although the froſts ſhould pierce and ſpoile ſome of the grapes on a bunch, yet if you be carefull to keepe the place warme, the fewer will be ſpoiled. And thus haue I ſhewed you the beſt directions to order this Orchard rightly, and all the waies I know are vſed in our Countrie to keep grapes good anie long time after the gathering, in regard wee haue not that comfort of a hotter Sun to preſerue them by its heate.

The fruits themſelues ſhall follow euerie one in their order; the lower ſhrubbes or buſhes firſt, and the greater afterwards.

# THE THIRD PART CALLED THE ORCHARD,

CONtaining all sorts of trees bearing fruit for mans vse to eate, proper and fit for to plant an Orchard in our climate and countrie: I bound it with this limitation, because both Dates, Oliues, and other fruits, are planted in the Orchards of Spaine, Italy, and other hot countries, which will not abide in ours. Yet herein I will declare whatsoeuer Art, striuing with Nature, can cause to prosper with vs, that whosoeuer will, may see what can bee effected in our countrie. And first to begin with the lower shrubbes or bushes, and after ascend to the higher trees.

## CHAP. I.

### *Rubus Idæus.* Raspis.

THe Raspis berrie is of two sorts, white and red, not differing in the forme either of bush, leafe, or berry, but onely in the colour and taste of the fruit. The Raspis bush hath tender whitish stemmes, with reddish small prickes like haires set round about them, especially at the first when they are young; but when they grow old they become more wooddy and firme, without any shew of thornes or prickles vpon them, and hath onely a little hairinesse that couereth them: the leaues are somewhat rough or rugged, and wrinkled, standing three or fiue vpon a stalke, somewhat like vnto Roses, but greater, and of a grayer greene colour: the flowers are small, made of fine whitish round leaues, with a dash as it were of blush cast ouer them, many standing together, yet euery one vpon his owne stalke, at the tops of the branches; after which come vp small berries, somewhat bigger then Strawberries, and longer, either red or white, made of many graines, more eminent then in the Strawberry, with a kinde of douninesse cast ouer them, of a pleasant taste, yet somewhat sowre, and nothing so pleasant as the Strawberrie. The white Raspis is a little more pleasant then the red, wherein there is small seede inclosed: the rootes creepe vndergound verie farre, and shoote vp againe in many places, much encreasing thereby.

There is another whose stemme and branches are wholly without prickles: the fruit is red, and somewhat longer, and a little more sharpe.

### The Vse of Raspis.

The leaues of Raspis may be vsed for want of Bramble leaues in gargles, and other decoctions that are cooling and drying, although not fully to that effect.

The Conſerue or Syrupe made of the berries, is effectuall to coole an hot ſtomacke, helping to refreſh and quicken vp thoſe that are ouercome with faintneſſe.

The berries are eaten in the Summer time, as an afternoones diſh, to pleaſe the taſte of the ſicke as well as the ſound.

The iuyce and the diſtilled water of the berries are verie comfortable and cordiall.

It is generally held of many, but how true I know not, that the red wine that is vſually ſold at the Vintners, is made of the berries of Raſpis that grow in colder countries, which giueth it a kinde of harſhneſſe: And alſo that of the ſame berries growing in hotter climates, which giueth vnto the wine a more pleaſant ſweetneſſe, is made that wine which the Vintners call Alligant: but we haue a Vine or Grape come to vs vnder the name of the Alligant Grape, as you ſhall finde it ſet downe hereafter among the Grapes; and therefore it is likely to be but an opinion, and no truth in this, as it may be alſo in the other.

## Chap. II.

*Ribes rubra, alba, nigra.* Currans red, white, and blacke.

THe buſhes that beare thoſe berries, which are vſually called red Currans, are not thoſe Currans either blew or red, that are ſold at the Grocers, nor any kind thereof; for that they are the grapes of a certaine Vine, as ſhall be ſhewed by and by: but a farre differing kinde of berry, whereof there are three ſorts, red, white, and blacke.

The red Curran buſh is of two ſorts, and groweth to the height of a man, hauing ſometimes a ſtemme of two inches thickneſſe, and diuers armes and branches, couered with a ſmooth, darke, browniſh barke, without anie pricke or thorne at all vpon anie part thereof, whereon doe growe large cornered blackiſh greene leaues, cut in on the edges, ſeeming to be made of fiue parts, almoſt like a Vine leafe, the ends a little pointing out, and ſtanding one aboue another on both ſides of the branches: the flowers are little and hollow, comming forth at the ioynts of the leaues, growing many together on a long ſtalke, hanging downe aboue a fingers length, and of an herbie colour: after which come ſmall round fruit or berries, greene at the firſt, and red as a Cherry when they are ripe, of a pleaſant and tart taſte: the other differeth not in anie other thing then in the berries, being twice as bigge as the former: the roote is wooddy, and ſpreadeth diuerſly.

The white Curran buſh riſeth vſually both higher then the red, and ſtraighter or more vpright, bigger alſo in the ſtemme, and couered with a whiter barke: the leaues are cornered, ſomewhat like the former, but not ſo large: the flowers are ſmall and hollow like the other, hanging downe in the ſame manner on long ſtalkes, being of a whiter colour: the berries likewiſe growe on the long ſtalkes, ſomewhat thicker ſet together, and of a cleare white colour, with a little blacke head, ſo tranſparent that the ſeedes may be eaſily ſeene thorough them, and of a more pleaſant winie taſte then the red by much.

The blacke Curran buſh riſeth higher then the white, with more plentifull branches, and more pliant and twiggie: the ſtemme and the elder branches being couered with a browniſh barke, and the younger with a paler: the flowers are alſo like vnto little bottles as the others be, of a greeniſh purple colour, which turne into blacke berries, of the bigneſſe of the ſmaller red Currans: the leaues are ſomewhat like vnto the leaues of the red Currans, but not ſo large: both branches, leaues, and fruit haue a kind of ſtinking ſent with them, yet they are not vnwholſome, but the berries are eaten of many, without offending either taſte or ſmell.

### The Vſe of Currans.

The red Currans are vſually eaten when they are ripe, as a refreſhing to an hot

1 *Rubus Idæus.* The Raſpis. 2 *Ribes fructu rubro vel albo.* White or red Currans. 3 *Groſſularia vulgaris.* The ordinary Gooſeberry
4 *Groſſularia fructu rubro.* The great red Gooſeberry. 5 *Groſſularia aculeata.* The prickly Gooſeberry. 6 *Oxyacantha ſeu Berberis;* The
Barbary buſh. 7 *Auellana Byzantina.* The Filberd of Conſtantinople. 8 *Auellana rubra noſtras.* The beſt red Filberd.

hot ſtomacke in the heate of the yeare, which by the tartneſſe is much delighted. Some preſerue them, and conſerue them alſo as other fruits, and ſpend them at neede.

The white Currans, by reaſon of the more pleaſant winie taſte, are more accepted and deſired, as alſo becauſe they are more daintie, and leſſe common.

Some vſe both the leaues and berries of the blacke Currans in ſawces, and other meates, and are well pleaſed both with the ſauour and taſte thereof, although many miſlike it.

## Chap. III.

*Vva Criſpa ſiue Groſſularia.* Gooſeberries or Feaberries.

WEe haue diuers ſorts of Gooſeberries, beſides the common kinde, which is of three ſorts, ſmall, great, and long. For wee haue three red Gooſeberries, a blew and a greene.

The common Gooſeberrie, or Feaberrie buſh, as it is called in diuers Countries of England, hath oftentimes a great ſtemme, couered with a ſmooth darke coloured bark, without anie thorne thereon, but the elder branches haue here and there ſome on them; and the younger are whitiſh, armed with verie ſharpe and cruell crooked thorns, which no mans hand can well auoide that doth handle them, whereon are ſet verie greene and ſmall cornered leaues cut in, of the faſhion almoſt of Smallage, or Hawthorne leaues, but broad at the ſtalke: the flowers come forth ſingle, at euerie ioynt of the leafe one or two, of a purpliſh greene colour, hollow and turning vp the brims a little: the berries follow, bearing the flowers on the heads of them, which are of a pale greene at the firſt, and of a greeniſh yellow colour when they are ripe, ſtriped in diuers places, and cleare, almoſt tranſparent, in which the ſeede lyeth. In ſome theſe berries are ſmall and round; in others much greater; a third is great, but longer then the other: all of them haue a pleaſant winie taſte, acceptable to the ſtomacke of anie (but the long kinde hath both the thicker skin, and the worſer taſte of the other) and none haue been diſtempered by the eating of them, that euer I could heare of.

The firſt of the red Gooſeberries is better knowne I thinke then the reſt, and by reaſon of the ſmall bearing not much regarded; the ſtemme is ſomewhat bigge, and couered with a ſmooth darke coloured barke, the younger branches are whiter, and without anie thorne or pricke at all, ſo long, weake, ſmall, and ſlender, that they lye vpon the ground, and will there roote againe: the leaues are like vnto the former Gooſeberries, but larger: the flowers and berries ſtand ſingle, and not manie to bee found anie yeare vpon them, but are ſomewhat long, and are as great as the ordinarie Gooſeberry, of a darke browniſh red colour, almoſt blackiſh when they are ripe, and of a ſweetiſh taſte, but without any great delight.

The ſecond red Gooſeberry riſeth vp with a more ſtraight ſtemme, couered with a browniſh barke; the young branches are ſtraight likewiſe, and whitiſh, and grow not ſo thicke vpon it as the former red kinde, and without any thorne alſo vpon them: the leaues are like vnto the former red, but ſmaller: the berries ſtand ſingly at the leaues as Gooſeberries doe, and are of a fine red colour when they are ripe, but change with ſtanding to be of a darker red colour, of the bigneſſe of the ſmall ordinary Gooſeberry, of a pretty tart taſte, and ſomewhat ſweete withall.

The third red Gooſeberry which is the greateſt, and knowne but vnto few, is ſo like vnto the common great Gooſeberry, that it is hardly diſtinguiſhed: the fruit or berries grow as plentifully on the branches as the ordinary, and are as great & round as the great ordinary kinde, but reddiſh, and ſome of them paler, with red ſtripes.

The blew Gooſeberry riſeth vp to bee a buſh like vnto the red Curran, and of the ſame bigneſſe and height, with broader and redder leaues at the firſt ſhooting out, then the ſecond red Gooſeberry: the berries are more ſparingly ſet on the branches, then on the ſmall red, and much about the ſame bigneſſe, or rather leſſer, of the colour of a Damſon, with an ouerſhadowing of a blewiſh colour vpon them, as the Damſon hath, before it be handled or wiped away.

The greene prickly Gooſeberry is very like vnto the ordinary Gooſeberry in ſtemme and branches, but that they are not ſtored with ſo many ſharpe prickles; but the young ſhootes are more plentifull in ſmall prickles about, and the greene leafe is a little ſmaller: the flowers are alike, and ſo are the berries, being of a middle ſize, and not very great, greene when they are thorough ripe as well as before, but mellower, and hauing a few ſmall ſhort prickles, like ſmall ſhort haires vpon them, which are harmleſſe, and without danger to anie the moſt dainty and tender palate that is, and of a verie good pleaſant taſte. The ſeede hereof hath produced buſhes bearing berries, hauing few or no prickles vpon them.

### The Vſe of Gooſeberries.

The berries of the ordinary Gooſeberries, while they are ſmall, greene, and hard, are much vſed to bee boyled or ſcalded to make ſawce, both for fiſh and fleſh of diuers ſorts, for the ſicke ſometimes as well as the ſound, as alſo before they be neere ripe, to bake into tarts, or otherwiſe, after manie faſhions, as the cunning of the Cooke, or the pleaſure of his commanders will appoint. They are a fit diſh for women with childe to ſtay their longings, and to procure an appetite vnto meate.

The other ſorts are not vſed in Cookery that I know, but ſerue to bee eaten at pleaſure; but in regard they are not ſo tart before maturity as the former, they are not put to thoſe vſes they be.

## Chap. IIII.

### *Oxyacantha, ſed potius Berberis.* Barberries.

THe Barberry buſh groweth oftentimes with very high ſtemmes, almoſt two mens height, but vſually ſomewhat lower, with manie ſhootes from the roote, couered with a whitiſh rinde or barke, and yellow vnderneath, the wood being white and pithy in the middle: the leaues are ſmall, long, and very greene, nicked or finely dented about the edges, with three ſmall white ſharpe thornes, for the moſt part ſet together at the ſetting on of the leaues: the flowers doe growe vpon long cluſtering ſtalkes, ſmall, round, and yellow, ſweete in ſmell while they are freſh, which turne into ſmall, long, and round berries, white at the firſt, and very red when they are ripe, of a ſharpe ſowre taſte, fit to ſet their teeth on edge that eate them: the roote is yellow, ſpreading far vnder the vpper part of the ground, but not very deepe.

There is (as it is thought) another kinde, whoſe berries are thrice as bigge as the former, which I confeſſe I haue not ſeene, and know not whether it be true or no: for it may peraduenture be but the ſame, the goodneſſe of the ground and ayre where they growe, and the youngneſſe of the buſhes cauſing that largeneſſe, as I haue obſerued in the ſame kinde, to yeeld greater berries.

There is ſaid to be alſo another kinde, whoſe berries ſhould be without ſtones or ſeede within them, not differing elſe in anie thing from the former: but becauſe I haue long heard of it, and cannot vnderſtand by all the inquirie I haue made, that any hath ſeene ſuch a fruit, I reſt doubtfull of it.

### The Vſe of Barberries.

Some doe vſe the leaues of Barberries in the ſtead of Sorrell, to make ſawce for meate, and by reaſon of their ſowreneſſe are of the ſame quality.

The berries are vſed to be pickled, to ſerue to trimme or ſet out diſhes of fiſh and fleſh in broth, or otherwiſe, as alſo ſometime to bee boyled in the broth, to giue it a ſharpe relliſh, and many other wayes, as a Maſter Cooke can better tell then my ſelfe.

The

The berries are preſerued and conſerued to giue to ſicke bodies, to helpe to coole any heate in the ſtomacke or mouth, and quicken the appetite.

The depurate iuyce is a fine menſtrue to diſſolue many things, and to verie good purpoſe, if it be cunningly handled by an Artiſt.

The yellow inner barke of the branches, or of the rootes, are vſed to be boyled in Ale, or other drinkes, to be giuen to thoſe that haue the yellow iaundiſe: As alſo for them that haue anie fluxes of choller, to helpe to ſtay and binde.

Cluſius ſetteth downe a ſecret that hee had of a friend, of a cleane differing propertie, which was, that if the yellow barke were laid in ſteepe in white wine for the ſpace of three houres, and afterwards drunke, it would purge one very wonderfully.

## Chap. V.

### *Nux Auellana.* The Filberd.

THe Filberd tree that is planted in Orchards, is very like vnto the Haſell nut tree that groweth wilde in the woods, growing vpright, parted into many boughes and tough plyable twigges, without knots, couered with a browniſh, ſpeckled, ſmooth, thinne rinde, and greene vnderneath: the leaues are broad, large, wrinkled, and full of veines, cut in on the edges into deepe dents, but not into any gaſhes, of a darke greene colour on the vpperſide, and of a grayiſh aſh colour vnderneath: it hath ſmall and long catkins in ſtead of flowers, that come forth in the Winter, when as they are firme and cloſe, and in the Spring open themſelues ſomewhat more, growing longer, and of a browniſh yellow colour: the nuts come not vpon thoſe ſtalkes that bore thoſe catkins, but by themſelues, and are wholly incloſed in long, thicke, rough huskes, bearded as it were at the vpper ends, or cut into diuers long iagges, much more then the wood nut: the nut hath a thinne and ſomewhat hard ſhell, but not ſo thicke and hard as the wood nut, in ſome longer then in other, and in the long kinde, one hath the skinne white that couereth the kernels, and another red.

There is another ſort of the round kinde that came from Conſtantinople, whoſe huske is more cut, torne, or iagged, both aboue and belowe, then any of our country; the barke alſo is whiter, and more rugged then ours, and the leaues ſomewhat larger.

We haue had from Virginia Haſell nuts, that haue beene ſmaller, rounder, browner, thinner ſheld, and more pointed at the end then ours: I know not if any hath planted of them, or if they differ in leafe or any thing elſe.

### The Vſe of Filberds.

Filberds are eaten as the beſt kinde of Haſell nuts, at bankets among other dainty fruits, according to the ſeaſon of the yeare, or otherwiſe, as euery one pleaſe: But Macer hath a Verſe, expreſſing prettily the nature of theſe nuts, which is,

*Ex minimis nucibus nulli datur eſca ſalubris.*

that is, There is no wholſome food or nouriſhment had from theſe ſmall kinde of nuts.

Yet they are vſed ſometime phyſically to be roſted, and made into a Lohoc or Electuary, that is vſed for the cough or cold. And it is thought of ſome, that Mithridates meant the kernels of theſe nuts, to be vſed with Figs and Rue for his Antidote, and not of Walnuts.

## Chap. VI.

### *Vitis.* The Vine.

THere is so great diuersities of Grapes, and so consequently of Vines that bear them, that I cannot giue you names to all that here grow with vs : for Iohn Tradescante my verie good friend, so often before remembred, hath assured me, that he hath twentie sorts growing with him, that hee neuer knew how or by what name to call them. One description therefore shall serue (as I vse to doe in such varieties) for all the rest, with the names afterwards, of as many as we can giue, and the seuerall formes, colours and proportions of the grapes.

The manured Vine, in the places where it hath abiden long time, groweth to haue a great bodie, stemme or trunke, sometimes of the bignesse of a mans arme, sleeue and all, spreading branches if it bee suffered without end or measure, but vsually stored with many armes or branches, both old and new, but weake, and therefore must bee sustained; whereof the old are couered with a thin scaly rinde, which will often chap and peele off of it selfe; the youngest being of a reddish colour, smooth and firme, with a hollownesse or pith in the middle: from the ioints of the young branches, and sometimes from the bodie of the elder, breake out on euerie side broade greene leaues, cut on the edges into fiue diuisions for the most part, and besides notched or dented about: right against the leafe, and likewise at other ioynts of the branches, come forth long twining or clasping tendrels, winding themselues about any thing standeth next vnto them: at the bottome of these leaues come forth clusters of small greenish yellow bloomes or flowers, and after them the berries, growing in the same manner in clusters, but of diuers formes, colours, tastes and greatnesse. For some grapes are great, others lesse, some very small (as the Currans that the Grocers sell) some white, some red, blew, blacke, or partie-coloured, some are are as it were square, others round: some the clusters are close, others open, some are sweete, others sower or harsh, or of some other mixed taste; euerie one differing from others, verie notably either in taste, colour or forme; within euerie one of which grapes, (and yet there is a grape without stones) are contained one, two, or more kernels or stones, some of them being smal, others greater: the rootes spread far and deepe. They that keepe their Vines in the best order, doe cut them low, not suffering them to grow high, or with too many branches, whereby they grow the better, take vp the lesser roome, and bring their grapes fairer and sweeter.

#### The kinds of Vines and Grapes.

Our ordinarie Grape both white and red, which excelleth Crabs for verjuice, and is not fit for wine with vs.

The white Muscadine Grape is a verie great Grape, sweete and firme, some of the bunches haue weighed sixe pound, and some of the grapes halfe an ounce.

The redde Muscadine is as great as the white, and chiefly differeth in colour.

The Burlet is a very great white Grape, but fitter for veriuice then wine for the most part; yet when a hot yeare happeneth fit for it, the Grape is pleasant.

The little blacke Grape that is ripe very early.

The Raisin of the Sunne Grape is a very great Grape, and very great clusters, of a reddish colour when it is ripe with vs, yet in an extraordinarie hot yeare, it hath got a little blewnesse cast ouer it by the heat: but naturally verie blew.

The Curran Grape (or the Grape of Corinth) is the least Grape of all, and beareth both few, and verie seldome with vs, but in reasonable great clusters, and of a blackish blew colour, when they are ripe with vs, and very

sweet

ſweete. There is another ſort of them that are red or browne, and of a ſower taſte, nothing ſo ſweete.

The Greeke wine Grape is a blackiſh Grape, and very ſweete.

The Frontignack is a white Grape, of a verie ſweete and delicate taſte, as the wine declareth, that ſmelleth as it were of Muske.

The ſquare Grape is reported to bear a Grape not fully round, but ſided, or as it were ſquare, whereby it became ſo called.

The Damaſco Grape is a great white grape, very ſweete, and is the true *Vva Zibeba*, that the Apothecaries ſhould vſe in the *Trochiſci Cyphi:* and ſuch wee haue had in former times come ouer vnto vs in great, long and round white boxes, containing halfe an hundred weight a peece.

The Ruſſet Grape is a reaſonable faire grape, exceeding ſweet and whitiſh, with a thicke skinne, cruſted ouer with a ſhew of aſh colour.

The white long Grape is like vnto a Pigeons egge, or as it were pointed pendent like a Pearle.

The partie-coloured Grape is a reaſonable great Grape, and diſcoloured when it is ripe, ſometimes the whole bunches, and ſometimes but ſome of the grapes being parted whitiſh, and blacke halfe through, verie variably.

The Rheniſh wine Grape is a white Grape, and endureth the cold of winter when it commeth earely, more then the Muſcadine before ſet downe, and is nothing ſo ſweete.

The White wine Grape is verie like vnto the Rhine Grape, the ſoile only and climate adding more ſweetneſſe vnto the one then to the other.

The Claret wine Grape is altogether like the white Grape, but that it is not white, but of a reddiſh colour, which lying bruiſed vpon the skins before they are preſſed, giue that Claret tincture to the wine.

The Teint is a Grape of a deeper or darker colour, whoſe iuice is of ſo deepe a colour, that it ſerueth to colour other wine.

The Burſarobe is a faire ſweete white Grape of much eſteeme about Paris.

The Alligant is a verie ſweete Grape, giuing ſo deep and liuely a coloured red wine, that no other whatſoeuer is comparable to it, and therfore vſually called Spaniards blood.

The blew or blacke Grape of Orleans is another blacke Grape, giuing a darke coloured ſweete wine much commended in thoſe parts.

The Grape without ſtones is alſo a kinde by it ſelfe, and groweth naturally neere Aſcalon, as Brochard affirmeth, the wine whereof is redde, and of a good taſte.

The Virginia Vine, whereof I muſt needes make mention among other Vines, beareth ſmall Grapes without any great ſtore of iuice therein, and the ſtone within it bigger then in any other Grape: naturally it runneth on the ground, and beareth little.

## The Vſe of Vines, Grapes, and other parts that come of them.

The greene leaues of the Vine are cooling and binding, and therefore good to put among other herbes that make gargles and lotions for ſore mouthes.

And alſo to put into the broths and drinke of thoſe that haue hot burning feauers, or any other inflammation.

They ſtay (as it is held for true) womens longings, if they be either taken inwardly, or applyed outwardly.

Wine is vſually taken both for drinke and medicine, and is often put into ſawces, broths, cawdles, and gellies that are giuen to the ſicke. As alſo into diuers Phyſicall drinkes, to be as a *vehiculum* for the properties of the ingredients.

It is diſtilled likewiſe after diners manners, with diuers things, for diuers & ſundry waters to drinke, & for diuers purpoſes both inward and outward.

As

1 *Vuæ nigræ minores.* The ſmall blacke Grape. 2 *Vuæ cæruleæ maiores.* The great blew Grape. 3 *Vuæ Moſchatellinæ.* The Muſcadine Grape. 4 *Vuæ Buriotenſes.* The Burlet Grape. 5 *Vuæ inſolatæ.* The Rayſins of the ſunne Grape. 6 *Ficus.* The Figge tree.

Alſo diſtilled of it ſelfe, is called Spirit of wine, which ſerueth to diſſolue, and to draw out the tincture of diuers things, and for many other purpoſes.

The iuice or veriuice that is made of greene hard grapes, before they be ripe, is vſed of the Apothecaries to be made into a Syrupe, that is very good to coole and refreſh a faint ſtomacke.

And being made of the riper grapes is the beſt veriuice, farre exceeding that which is made of crabs, to be kept all the yeare, to be put both into meates and medicines.

The grapes of the beſt ſorts of Vines are preſſed into wine by ſome in theſe dayes with vs, and much more as I verily beleeue in times paſt, as by the name of Vineyard giuen to many places in this Kingdome, eſpecially where Abbies and Monaſteries ſtood, may bee coniectured: but the wine of late made hath beene but ſmall, and not durable, like that which commeth from beyond Sea, whether our vnkindly yeares, or the want of skill, or a conuenient place for a Vineyard be the cauſe, I cannot well tell you.

Grapes of all ſorts are familiarly eaten when they are ripe, of the ſicke ſometimes as well as the ſound.

The dryed grapes which we call great Rayſins, and the Currans which we call ſmall Rayſins, are much vſed both for meates, broths, and ſawces, in diuers manners, as this Countrey in generall aboue any other, wherein many thouſands of Frailes full, Pipes, Hogs-heads, and Buts full are ſpent yearly, that it breedeth a wonder in them of thoſe parts where they growe and prouide them, how we could ſpend ſo many.

The Rayſins of the Sunne are the beſt dryed grapes, next vnto the Damaſco, and are very wholſome to eate faſting, both to nouriſh, and to helpe to looſen the belly.

The dryed Lees of wine called Argoll or Tartar, is put to the vſe of the Goldſmith, Dyer, and Apothecary, who doe all vſe it in ſeuerall manners, euery one in his art.

Of it the Apothecaries make *Cremor Tartari*, a fine medicine to bee vſed, as the Phyſitian can beſt appoint, and doth helpe to purge humours by the ſtoole.

Thereof likewiſe they make a kinde of water or oyle, fit to bee vſed, to take away freckles, ſpots, or any ſuch deformities of the face or skinne, and to make it ſmooth. It cauſeth likewiſe haire to growe more aboundantly in thoſe places where it naturally ſhould growe.

The liquor of the Vine that runneth forth when it is cut, is commended to be good againſt the ſtone whereſoeuer it be; but that liquor that is taken from the end of the branches when they are burnt, is moſt effectuall to take away ſpots and markes, ring-wormes and tetters in any place.

## Chap. VII.

*Ficus*. The Figge tree.

THe Figge trees that are nourſed vp in our country are of three ſorts, whereof two are high; the one bearing againſt a wall goodly ſweete and delicate Figs, called Figs of Algarua, and is blewiſh when it is ripe: the other tall kinde is nothing ſo good, neither doth beare ripe Figges ſo kindly and well, and peraduenture may be the white ordinary kinde that commeth from Spaine. The third is a dwarfe kinde of Figge tree, not growing much higher then to a mans body or ſhoulders, bearing excellent good Figges and blew, but not ſo large as the firſt kinde.

The Figge trees of all theſe three kindes are in leaues and growing one like vnto another, ſauing for their height, colour, and ſweetneſſe of the fruit, hauing many armes or branches, hollow or pithy in the middle, bearing very large leaues, and ſomewhat thicke, diuided ſometimes into three, but vſually into fiue ſections, of a darke greene colour on the vpperſide, and whitiſh vnderneath, yeelding a milkie iuyce when it is broken,

broken, as the branches also or the figges when they are greene; the fruit breaketh out from the branches without anie blossome, contrary to all other trees of our Orchard, being round and long, fashioned very like vnto a small Peare, full of small white grains or kernels within it, of a very sweete taste when it is ripe, and very mellow or soft, that it can hardly be carried farre without bruising.

The other two sorts you may easily know and vnderstand, by so much as hath been said of them. Take onely this more of the Figge tree, That if you plant it not against a bricke wall, or the wall of an house, &c. it will not ripen so kindly. The dwarfe Figge tree is more tender, and is therefore planted in great square tubs, to be remoued into the sunne in the Summer time, and into the house in Winter.

### The Vse of Figges.

Figges are serued to the table with Raysins of the Sunne, and blanched Almonds, for a Lenten dish.

The Figs that growe with vs when they are ripe, and fresh gathered, are eaten of diuers with a little salt and pepper, as a dainty banquet to entertaine a freind, which seldome passeth without a cup of wine to wash them downe.

In Italy (as I haue beene enformed by diuers Gentlemen that haue liued there to study physicke) they eate them in the same manner, but dare not eate many for feare of a feuer to follow, they doe account them to be such breeders of bloud, and heaters of it likewise.

The Figges that are brought vs from Spaine, are vsed to make Ptisan drinkes, and diuers other things, that are giuen them that haue coughes or colds.

It is one of the ingredients also with Nuts and Rice, into Mithridates counterpoison.

The small Figges that growe with vs, and will not ripen, are preserued by the Comfitmakers, and candid also, to serue as other moist or candid banquetting stuffe.

## Chap. VIII.

### *Sorbus.* The Seruice tree.

THere are two kindes of Seruice trees that are planted in Orchards with vs, and there is also a wilde kinde like vnto the later of them, with Ashen leaues, found in the woods growing of it selfe, whose fruit is not gathered, nor vsed to bee eaten of any but birds. And there is another kinde also growing wilde abroad in many places, taken by the Country people where it groweth, to be a Seruice tree, and is called in Latine, *Aria Theophrasti*, whose leaues are large, somewhat like Nut tree leaues, but greene aboue, and grayish vnderneath: some doe vse the fruit as Seruices, and for the same purposes to good effect, yet both of these wilde kindes wee leaue for another worke, and here declare vnto you onely those two sorts are noursed vp in our Orchards.

The more common or ordinary Seruice tree with vs, is a reasonable great tree, couered with a smooth barke, spread into many great armes, whereon are set large leaues, very much cut in on the edges, almost like vnto a Vine leafe, or rather like vnto that kind of Maple, that is vsually called the Sycomore tree with vs: the flowers are white, and growe many clustering together, which after bring forth small browne berries when they are ripe, of the bignesse almost of Hasell nuts, with a small tuft, as if it were a crowne on the head, wherein are small blacke kernels.

The other kinde, which is more rare with vs, and brought into this Land by Iohn Tradescante, heretofore often remembred, hath diuers winged leaues, many set together like vnto an Ashen leafe, but smaller, and euery one endented about the edges: the flowers growe in long clusters, but nothing so many, or so close set as the wilde kinde: the fruit of this tree is in some round like an Apple, and in others a little longer

like a Peare, but of a more pleasant taste then the ordinarie kinde, when they are ripe and mellowed, as they vse to doe with both these kindes, and with Medlars.

### The Vse of Seruices.

They are gathered when they growe to be neare ripe (and that is neuer before they haue felt some frosts) and being tyed together, are either hung vp in some warme roome, to ripen them thoroughly, that they may bee eaten, or (as some vse to doe) lay them in strawe, chaffe, or branne, to ripen them.

They are binding, fit to be taken of them that haue any scouring or laske, to helpe to stay the fluxe; but take heed, lest if you binde too much, more paine and danger may come thereof then of the scouring.

## Chap. IX.

### *Mespilus.* The Medlar tree.

THere are three sorts of Medlers: The greater and the lesser English, and the Neapolitan.

The great and the small English Medlar differ not one from the other in any thing, but in the size of the fruit, except that the small kinde hath some prickes or thornes vpon it, which the great one hath not, bearing diuers boughes or armes, from whence breake forth diuers branches, whereon are set long and somewhat narrow leaues, many standing together; in the middle whereof, at the end of the branch, commeth the flower, which is great and white, made of fiue leaues, broad at the ends, with a nicke in the middle of euery one; after which commeth the fruit, being round, and of a pale browniſh colour, bearing a crowne of those small leaues at the toppe, which were the huske of the flower before, the middle thereof being somewhat hollow, and is harsh, able to choake any that shall eate it before it be made mellow, wherein there are certaine flat and hard kernels.

The Medlar of Naples groweth likewise to bee a reasonable great tree, spreading forth armes and branches, whereon are set many gashed leaues, somewhat like vnto Hawthorne leaues, but greater, and likewise diuers thornes in many places: the flowers are of an herbie greene colour, and small, which turne into smaller fruit then the former, and rounder also, but with a small head or crowne at the toppe like vnto it, and is of a more sweete and pleasant taste then the other, with three seeds only therein ordinarily.

### The Vse of Medlars.

Medlars are vsed in the same manner that Seruices are, that is, to be eaten when they are mellowed, and are for the same purposes to binde the body when there is a cause: yet they as well as the Seruices, are often eaten by them that haue no neede of binding, and but onely for the pleasant sweetnesse of them when they are made mellow, and sometimes come as a dish of ripe fruit at their fit season, to be serued with other sorts to the table.

## Chap. X.

### *Lotus.* The Lote or Nettle tree.

THe first kinde of Lote tree, whereof Dioscorides maketh mention, is but of one kinde; but there are some other trees spoken of by Theophrastus, that may be referred thereunto, which may bee accounted as bastard kindes thereof, of which I meane to entreate in this Chapter, hauing giuen you before the description of

1 *Sorbus legitima.* The true Seruice tree. 2 *Sorbus vulgaris siue Torminalis.* The ordinary Seruice tree. 3 *Mespilus vulgaris.* The common Medlar tree. 4 *Mespilus Aronia.* The Medlar of Naples. 5 *Lotus arbor.* The Nettle tree. 6 *Lotus Virginiana.* The Pishamin or Virginia Plumme. 7 *Cornus mas.* The Cornell Cherry tree.

of another kinde hereof (by the opinion of good Authors) vnder the name of *Lauro-cerasus*.

The first or true Lote tree groweth to be a tree of a great height, whose bodie and elder branches are couered with a smooth darke greene barke, the leaues are somewhat rough in handling, of a darke greene colour, long pointed, and somewhat deepe dented about the edges, somewhat like vnto a Nettle leafe, and oftentimes growe yellow toward Autumne: the flowers stand here and there scattered vpon the branches, after which come round berries like vnto Cherries, hanging downewards vpon long foot-stalkes, greene at the first, and whitish afterwards; but when they are ripe they become reddish, and if they be suffered to hang too long on the branches, they grow blackish, of a pleasant austere taste, not to be misliked, wherein is a hard round stone.

The second, which is a bastard kinde, and called *Guaiacum Patauinum*, groweth to bee a faire tree, with a smooth darke greene barke, shooting out many faire great boughes, and also slender greene branches, beset with faire broad greene leaues, almost like vnto the leaues of the Cornell tree, but larger: the flowers growe along the branches close vnto them, without any or with a very short foote-stalke consisting of foure greene leaues, which are as the huske, containing within it a purplish flower, made of foure leaues somewhat reddish: the fruit standeth in the middle of the green huske, greene at the first, and very harsh, but red and round when it is ripe, and somewhat like a Plumme, with a small point or pricke at the head thereof, and of a reasonable pleasant taste or relish, wherein are contained flat and thicke browne seeds or kernels, like vnto the kernels of *Cassia Fistula*, somewhat hard, and not so stonie, but that it may somewhat easily be cut with a knife.

The third is called in Virginia *Pishamin*, The Virginia Plumme (if it be not all one with the former Guaiacana, whereof I am more then halfe perswaded) hath growne with vs of the kernels that were sent out of Virginia, into great trees, whose wood is very hard and brittle, and somewhat white withall: the branches are many, and grow slender to the end, couered with a very thinne greenish bark, whereon doe grow many faire broad greene leaues, without dent or notch on the edges, and so like vnto the former *Guaiacum*, that I verily thinke it (as I before said) to bee the same. It hath not yet borne flower or fruit in our Countrey that I can vnderstand: but the fruit, as it was sent to vs, is in forme and bignesse like vnto a Date, couered with a blackish skinne, set in a huske of foure hard leaues, very firme like vnto a Date, and almost as sweete, with great flat and thicke kernels within them, very like vnto the former, but larger.

### The Vse of these Lote trees.

The first sort is eaten as an helper to coole and binde the body: the last, as Captaine Smith relateth in the discouery of Virginia, if the fruit be eaten while it is greene, and not ripe, is able by the harsh and binding taste and quality to draw ones mouth awry (euen as it is said of the former Guaiacana) but when it is thorough ripe it is pleasant, as I said before.

## Chap. XI.

### *Cornus mas.* The Cornell tree.

THe Cornell tree that is planted in Orchards, being the male (for the female is an hedge bush) is of two sorts, the one bearing red, the other whiter berries, which is very rare yet in our country, and not differing else.

It groweth to a reasonable bignesse and height, yet neuer to any great tree, the wood whereof is very hard, like vnto horne, and thereof it obtained the name: the body and branches are couered with a rugged barke, and spreadeth reasonable well, hauing somewhat smooth leaues, full of veines, plaine, and not dented on the edges: the flowers are many small yellow tufts, as it were of short haires or threads set together, which come forth before any leafe, and fall away likewise before any leafe bee much open: the fruit are long and round berries, of the bignesse of small Oliues, with an hard

hard round ſtone within them, like vnto an Oliue ſtone, and are of a yellowiſh red when they are ripe, of a reaſonable pleaſant taſte, yet ſomewhat auſtere withall.

The white (as I ſaid) is like vnto the red, but onely that his fruit is more white when it is ripe.

### The Vſe of the Cornelles.

They helpe to binde the body, and to ſtay laskes, and by reaſon of the pleaſantneſſe in them when they are ripe, they are much deſired.

They are alſo preſerued and eaten, both for rarity and delight, and for the purpoſe aforeſaid.

## Chap. XII.

### *Ceraſus.* The Cherry tree.

THere are ſo many varieties and differences of Cherries, that I know not well how to expreſſe them vnto you, without a large relation of their ſeuerall formes. I will therefore endeauour after one generall deſcription (as my cuſtome is in many other the like variable fruits) to giue as briefe and ſhort notes vpon all the reſt, as I can both for leafe and fruit, that ſo you may the better know what the fruit is, when you haue the name.

The Engliſh Cherrie tree groweth in time to be of a reaſonable bigneſſe and height, ſpreading great armes, and alſo ſmall twiggy branches plentifully; the leaues whereof are not verie large or long, but nicked or dented about the edges: the flowers come forth two or three or foure at the moſt together, at a knot or ioynt, euerie one by it ſelfe, vpon his owne ſmall and long footeſtalke, conſiſting of fiue white leaues, with ſome threds in the middle; after which come round berries, greene at the firſt, and red when they are through ripe, of a meane bigneſſe, and of a pleaſant ſweete taſte, ſomewhat tart withall, with a hard white ſtone within it, whoſe kernell is ſomewhat bitter, but not vnpleaſant.

The Flanders Cherrie differeth not from the Engliſh, but that it is ſomewhat larger, and the Cherry ſomewhat greater and ſweeter, and not ſo ſower.

The early Flanders Cherry is more rathe or early ripe, almoſt as ſoone as the May Cherry, eſpecially planted againſt a wall, and of many falſe knaues or Gardiners are ſold for May Cherrie trees.

The May Cherrie in a ſtandard beareth ripe fruite later then planted againſt a wall, where the berries will be red in the verie beginning of May ſometimes.

The Arch-Dukes Cherrie is one of the faireſt and beſt cherries wee haue, being of a very red colour when it is ripe, and a little long more then round, and ſomewhat pointed at the end, of the beſt relliſh of any Cherrie whatſoeuer, and of a firme ſubſtance; ſcarce one of twentie of our Nurſerie men doe ſell the right, but giue one for another: for it is an inherent qualitie almoſt hereditarie with moſt of them, to ſell any man an ordinary fruit for whatſoeuer rare fruit he ſhall aske for: ſo little they are to be truſted.

The ounce Cherrie hath the greateſt and broadeſt leafe of any other cherrie, but beareth the ſmalleſt ſtore of cherries euerie yeare that any doth, and yet bloſſometh well: the fruit alſo is nothing anſwerable to the name being not verie great, of a pale yellowiſh red, neere the colour of Amber, and therefore ſome haue called it, the Amber Cherrie.

The great leafed Cherrie is thought of diuers to bee the Ounce Cherrie, becauſe it hath almoſt as great a leafe as the former: but the fruit of this alſo doth not anſwer the expectation of ſo great a leafe, being but of a meane bigneſſe, and a ſmall bearer, yet of a pale reddiſh colour.

The true Gaſcoign Cherry is known but to a few, for our Nurſery men do ſo change the names of moſt fruits they ſell, that they deliuer but very few true names to any: In former times before our wilde blacke Cherrie was found to grow plentifully in our owne woods in many places of this Land, the French continually ſtored vs with wilde ſtockes to graft vpon, which then were called Gaſcoigne ſtocks, but ſince they haue ſo

termed

termed another red Cherrie, and obtruded it vpon their cuſtomers: but the true is one of our late ripe white Cherries, euen as Gerard ſaith, it is a great cherrie and ſpotted: and this is that Cherrie I ſo commend to be a fit ſtocke to graft May cherries vpon.

The Morello Cherrie is of a reaſonable bigneſſe, of a darke red colour when they are full ripe, and hang long on, of a ſweetiſh ſower taſte, the pulpe or ſubſtance is red, and ſomewhat firme: if they be dryed they will haue a fine ſharpe or ſower taſte very delectable.

The Hartlippe Cherrie is ſo called of the place where the beſt of this kinde is nourſed vp, being betweene Sittingbourne and Chattam in Kent, and is the biggeſt of our Engliſh kindes.

The ſmaller Lacure or Hart Cherrie is a reaſonable faire Cherrie, full aboue, and a little pointing downward, after the faſhion of an heart, as it is vſually painted, blackiſh when it is full ripe, and leſſer then the next.

The great Lacure or Hart Cherrie differeth not in forme, but in greatneſſe, being vſually twice as great as the former, and of a reddiſh blacke colour alſo: both of them are of a firme ſubſtance, and reaſonable ſweete. Some doe call the white cherrie, the White hart cherrie.

The Luke Wardes Cherrie hath a reaſonable large leafe, and a larger flower then many other: the cherries grow with long ſtalkes, and a ſtone of a meane ſize within them, of a darke reddiſh colour when they are full ripe, of a reaſonable good relliſh, and beareth well.

The Corone Cherrie hath a leafe little differing from the Luke Wardes cherrie; the fruit when it is ripe, is of a faire deepe red colour, of a good bigneſſe, and of a verie good taſte, neither verie ſweete or ſower: the pulpe or iuice will ſtaine the hands.

The Vrinall Cherrie in a moſt fruitfull yeare is a ſmall bearer, hauing many yeares none, and the beſt but a few; yet doth bloſſome plentifully euery yeare for the moſt part: the cherrie is long and round, like vnto an Vrinall, from whence it tooke his name; reddiſh when it is full ripe, and of an indifferent ſweete relliſh.

The Agriot Cherrie is but a ſmall Cherrie, of a deepe redde colour when it is ripe, which is late; of a fine ſharpe taſte, moſt pleaſant and wholſome to the ſtomacke of all other cherries, as well while they are freſh as being dryed, which manner they much vſe in France, and keepe them for the vſe both of the ſicke and ſound at all times.

The Biguarre Cherrie is a fair cherrie, much ſpotted with white ſpots vpon the pale red berry, and ſometimes diſcoloured halfe white and halfe reddiſh, of a reaſonable good relliſh.

The Morocco Cherrie hath a large white bloſſome, and an indifferent big berrie, long and round, with a long ſtalke of a darke reddiſh purple colour, a little tending to a blew when it is full ripe, of a firme ſubſtance: the iuice is of a blackiſh red, diſcolouring the hands or lips, and of a pleaſant taſte: Some doe thinke that this and the Morello be both one.

The Naples Cherrie is alſo thought to bee all one with the Morello or Morocco.

The white Spaniſh Cherrie is an indifferent good bearer, the leafe and bloſſome ſomewhat large, and like the Luke Wardes cherrie: the cherries are reaſonable faire berries, with long ſtalkes and great ſtones, white on the outſide, with ſome redneſſe, on the one ſide of a firme ſubſtance, and reaſonable ſweet, but with a little aciditie, and is one of the late ripe ones: But there is another late ripe white Cherry, which ſome call the Gaſcoigne, before remembred.

The Flanders cluſter Cherrie is of two ſorts, one greater then another: the greater kinde hath an indifferent large leafe; the bloſſomes haue many threds within them, ſhewing as it were many parts, which after turne into cluſters of berries, foure, fiue or ſixe together, and but with one ſtalke vnder them, as if they grew one out of another, and ſometimes they will beare but two or three, and moſt of them but one cherry on a ſtalke, which are red when they are ripe, very tender, and wateriſh ſweete in eating.

The leſſer is in all things like the greater, but ſmaller, which maketh the difference.

The wilde cluſter or birds cluſter Cherry beareth many bloſſomes ſet all along the ſtalkes, and cherries after them in the ſame maner, like a long thinne bunch of grapes, and therefore called of ſome the Grape cherry: there are of them both red and blacke.

The

1 *Cerasus præcox*. The May Cherry. 2 *Cerasus Batauica*. The Flanders Cherry. 3 *Cerasus Hispanica siue alba*. The white Cherry. 4 *Cerasus platephyllos*. The great leafed Cherry. 5 *Cerasus Luca Wardi*. Luke Wards Cherry. 6 *Cerasus Neapolitana*. The Naples Cherry. 7 *Cerasus Cordata*. The Heart Cherry. 8 *Cerasus maculata*. The bignarre or spotted Cherry. 9 *Cerasus autumnacemosa*. The wilde cluster Cherry. 10 *Cerasus Corymbifera*. The Flanders cluster Cherry. 11 *Cerasus Archiducis*. The Archdukes Cherry. 12 *Chamacerasus*. The dwarfe Cherry.

The ſoft ſheld Cherrie is a ſmall red cherrie when it is ripe, hauing the ſtone within it ſo ſoft and tender, that it may eaſily be broken in the eating of the cherrie.

Iohn Tradeſcantes Cherrie is moſt vſually ſold by our Nurſery Gardiners, for the Archdukes cherrie, becauſe they haue more plenty thereof, and will better be increaſed, and becauſe it is ſo faire and good a cherrie that it may be obtruded without much diſcontent: it is a reaſonable good bearer, a faire great berrie, deepe coloured, and a little pointed.

The Baccalaos or New-found-land Cherrie hath a ſhining long leafe, moſt like vnto a Peach leafe, the bloſſomes come very many together as it were in an vmbell, which is ſuch a cluſter as is neither like the Flanders cluſter, nor the wilde cluſter cherrie bloſſome: it bringeth forth berries ſtanding in the ſame manner euerie one vpon his own footeſtalke, being no bigger then the largeſt berrie of the red Curran tree or buſh, of a pale or wateriſh red colour when it is ripe.

The ſtrange long cluſter Cherrie, or *Padus Theophraſti Dalechampio* is reckoned by the Author of that great Herball that goeth vnder his name, among the ſorts of cherries; and ſo muſt I vntill a fitter place be found for it. It groweth in time to be a great tree, with a ſad coloured barke both on the bodie and branches, whereon doe grow many leaues, ſomewhat broade, ſhorter, harder, and a little more crumpled then any cherrie leafe: the bloſſomes are very ſmall, and of a pale or whitiſh colour, ſmelling very ſweete and ſtrong, or rather heady, like Orenge flowers, growing on ſmall long branches, very like the toppe of flowers vpon the Laburnum or Beane trefoile trees: after which come ſmall blacke berries, growing together all along the long ſtalke, like vnto the wilde cluſter or birds cherrie mentioned before, but not much bigger then tares, with ſmall ſtones within them, and little or no ſuſtance vpon them: the French call the tree *Putier*, becauſe the wood thereof ſtinketh, and make it to be wonderfull that the bloſſomes of the tree ſhould be ſo ſweete, and the wood ſo ſtinking.

The Cullen Cherrie is a darke red cherrie like the Agriot, which they of thoſe parts neere Cullen and Vtrecht &c. vſe to put into their drinke, to giue it the deeper colour.

The great Hungarian Cherrie of Zwerts is like both in leafe and fruite vnto the Morello cherrie, but much greater and fairer, and a far better bearer: for from a ſmall branch hath beene gathered a pound of cherries, and this is vſuall continually, and not accidentally, moſt of them foure inches in compaſſe about, and very many of them more of a faire deepe red colour, and very ſweete, excelling the Arch-Dukes cherry, or any other whatſoeuer.

The Cameleon or ſtrange changeable Cherry deſeruedly hath this name, although of mine owne impoſition, not only becauſe it beareth vſually both bloſſomes, greene and ripe fruit at one time thereupon, but that the fruit will be of many formes; ſome round, ſome as it were ſquare, and ſome bunched forth on one ſide or another, abiding conſtant in no faſhion, but for the moſt part ſhewing forth all theſe diuerſities euerie yeare growing vpon it: the fruit is of a very red colour, and good taſte.

The great Roſe Cherry, or double bloſſomd Cherry differeth not in any thing from the Engliſh Cherrie, but only in the bloſſomes, which are very thicke of white leaues, as great and double as the double white Crowfoote, before remembred, and ſomtimes out of the middle of them will ſpring another ſmaller flower, but double alſo; this ſeldome beareth fruit, but when it doth I ſuppoſe it commeth from thoſe bloſſomes are the leaſt double, and is red, no bigger then our ordinary Engliſh cherrie.

The leſſer Roſe or double bloſſomd Cherrie beareth double flowers alſo, but not ſo thicke and double as the former; but beareth fruit more plentifully, of the ſame colour and bigneſſe with the former.

The Dwarfe Cherrie is of two ſorts; one whoſe branches fall downe low, round about the body of it, with ſmall greene leaues, and fruit as ſmall, of a deep red colour.

The other, whoſe branches, although ſmall, grow more vpright, hauing greener ſhining leaues: the fruit is little bigger then the former, red alſo when it is ripe, with a little point at the end: both of them of a ſweetiſh relliſh, but more ſower.

The great bearing Cherry of Maſter Millen is a reaſonable great red cherry, bearing very plentifully, although it bee planted againſt a North wall, yet it will bee late ripe, but of an indifferent ſweet and good relliſh.

The long finger Cherry is another ſmall long red one, being long & round like a finger, wherof it took the name: this is not the Vrinall cherry before, but differing from it.

The

### The Vſe of Cherries.

All theſe ſorts of Cherries ſerue wholly to pleaſe the palate, and are eaten at all times, both before and after meales.

All Cherries are cold, yet the ſower more then the ſweete; and although the ſweete doe moſt pleaſe, yet the ſower are more wholſome, if there bee regard taken in the vſing.

The Agriot or ſower Cherries are in France much vſed to bee dryed (as is ſaid before) as Pruines are, and ſo ſerue to miniſtred to be the ſick in all hot diſeaſes, as feuers &c. being both boyled in their drinkes, and taken now and then of themſelues, which by reaſon of their tartneſſe, doe pleaſe the ſtomacke paſſing well.

The Gum of the Cherrie tree is commended to bee good for thoſe are troubled with the grauell or ſtone. It is alſo good for the cough being diſſolued in liquour, and ſtirreth vp an appetite. The diſtilled water of the blacke Cherries, the ſtones being broken among them, is vſed for the ſame purpoſe, for the grauell, ſtone, and winde.

## Chap. XIII.

### *Prunus.* The Plumme tree.

THere are many more varieties of Plummes then of Cherries, ſo that I muſt follow the ſame order with theſe that I did with them, euen giue you their names apart, with briefe notes vpon them, and one deſcription to ſerue for all the reſt. And in this recitall I ſhall leaue out the Apricockes which are certainly a kind of Plum, of an eſpeciall difference, and not of a Peach, as Galen and ſome others haue thought, and ſet them in a chapter by themſelues, and only in this ſet down thoſe fruits are vſually called Plums.

The Plum tree (eſpecially diuers of them) riſeth in time to bee a reaſonable tall and great tree, whoſe bodie and greater armes are couered with a more rugged barke, yet in ſome more or leſſe, the younger branches being ſmooth in all, the leaues are ſomewhat rounder then thoſe of the Cherrie tree, and much differing among themſelues, ſome being longer, or larger, or rounder then others, and many that are exerciſed herein, can tell by the leafe what Plum the tree beareth (I ſpeake this of many, not of all) as in many Cherries they can doe the like: the flowers are white, conſiſting of fiue leaues: the fruit is as variable in forme, as in taſte or colour, ſome being ouall, or Peare faſhion or Almond like, or ſphericall or round, ſome firme, ſome ſoft and wateriſh, ſome ſweete, ſome ſower or harſh, or differing from all theſe taſtes: and ſome white, others blacke, ſome red, others yellow, ſome purple, others blew, as they ſhall bee briefly ſet downe vnto you in the following lines, where I meane not to inſert any the wilde or hedge fruit, but thoſe only are fit for an Orchard, to be ſtored with good fruit: and of all which ſorts, the choyſeſt for goodneſſe, and rareſt for knowledge, are to be had of my very good friend Maſter Iohn Tradeſcante, who hath wonderfully laboured to obtaine all the rareſt fruits hee can heare off in any place of Chriſtendome, Turky, yea or the whole world; as alſo with Maſter Iohn Millen, dwelling in Olde ſtreete, who from Iohn Tradeſcante and all others that haue had good fruit, hath ſtored himſelfe with the beſt only, and he can ſufficiently furniſh any.

The Amber Primordian Plumme is an indifferent faire Plumme, early ripe, of a pale yellowiſh colour, and of a wateriſh taſte, not pleaſing.

The red Primordian Plumme is of a reaſonable ſize, long and round, reddiſh on the outſide, of a more dry taſte, and ripe with the firſt ſorts in the beginning of Auguſt.

The blew Primordian is a ſmall plumme, almoſt like the Damaſcene, and is ſubiect to drop off from the tree before it be ripe.

The white Date Plum is no very good plum.

The

The red Date plumme is a great long red pointed plumme, and late ripe, little better then the white.

The blacke Muſſell plumme is a good plumme, reaſonable drye, and taſteth well.

The red Muſſell Plumme is ſomewhat flat as well as round, of a very good taſte, and is ripe about the middle of Auguſt.

The white Muſſell plumme is like the redde, but ſomewhat ſmaller, and of a whitiſh greene colour, but not ſo well taſted.

The Imperiall plum is a great long reddiſh plum, very wateriſh, and ripeneth ſomewhat late.

The Gaunt plum is a great round reddiſh plum, ripe ſomewhat late, and eateth wateriſh.

The red Peſcod plum is a reaſonable good plum.

The white Peſcod plum is a reaſonable good relliſhed plumme, but ſomewhat wateriſh.

The greene Peſcod plum is a reaſonable big and long pointed plum, and ripe in the beginning of September.

The Orenge plum is a yellowiſh plum, moiſt, and ſomewhat ſweetiſh.

The Morocco plumme is blacke like a Damſon, well taſted, and ſomewhat drye in eating.

The Dine plum is a late ripe plum, great and whitiſh, ſpeckled all ouer.

The Turkie plum is a large long blackiſh plum, and ſomewhat flat like the Muſſell plum, a well relliſhed dry plum.

The Nutmeg plumme is no bigger then a Damſon, and is of a greeniſh yellow colour when it is ripe, which is with vs about Bartholmew tide, and is a good plum.

The Perdigon plumme is a dainty good plumme, early blackiſh, and well relliſhed.

The Verdoch plum is a great fine greene ſhining plum fit to preſerue.

The Ienua plum is the white Date plum, before remembred.

The Barberry plum is a great early blacke plum, and well taſted.

The Pruneola plum is a ſmall white plum, of a fine tart taſte: it was wont to bee vſually brought ouer in ſmall round boxes, and ſold moſt commonly at the Comfit-makers, (cut in twaine, the ſtone caſt away) at a very deere rate: the tree groweth and beareth well with vs.

The Shepway Bulleis is of a darke blewiſh brown colour, of a larger ſize then the ordinary, and of a ſharpe taſte, but not ſo good as the common.

The white and the blacke Bulleis are common in moſt Countries, being ſmall round plums, leſſer then Damſons, ſharper in taſte, and later ripe.

The Fluſhing Bulleis groweth with his fruite thicke cluſtring together like grapes.

The Winter Creke is the lateſt ripe plum of all ſorts, it groweth plentifully about Biſhops Hatfield.

The white Peare plum early ripe, is of a pale yellowiſh greene colour.

The late ripe white Peare plum is a greater and longer plum, greeniſh white, and is not ripe vntill it be neere the end of September, both wateriſh plums.

The blacke Peare plum is like vnto the white Peare plumme, but that the colour is blackiſh when it is ripe, and is of a very good relliſh, more firme and drye then the other.

The red Peare plumme is of the ſame faſhion and goodneſſe, but is the worſt of the three.

The white Wheate plum is a wateriſh fulſome plum.

The red Wheate plum is like the other for taſte.

The Bowle plum is flat and round, yet flatter on the one ſide then on the other, which cauſed the name, and is a very good relliſhed blacke plum.

The Friars plumme is a very good plum, well taſted, and comming cleane from the ſtone, being blacke when it is ripe, and ſome whitiſh ſpots vpon it.

The Catalonia plum is a very good plum.

The don Alteza is alſo a very good plum.

The Muſcadine plum, ſome call the Queene mother plumme, and ſome the Cherry plum, is a faire red plum, of a reaſonable bigneſſe, and ripe about Bartholmew tide.

The Chriſtian plum, called alſo the Nutmeg plum; the tree groweth very ſhrubby,

and

1 *Prunum Imperiale.* The Imperiall Plum. 2 *Prunum Turcicum.* The Turkey Plum. 3 *Prunum præcox rubrum.* The red Primordian Plum. 4 *Prunum Mytellinum.* The Mussell Plum. 5 *Prunum Ambarinum.* The Amber Plum. 6 *Prunum Regineum.* The Queen mother Plum. 7 *Prunum viride.* The green Oysterly Plum. 8 *Prunum Arantiacum.* The Orenge Plum. 9 *Prunum Myristicum.* The Nutmeg Plum. 10 *Prunum Siliquosum.* The Peacod Plum. 11 *Prunum Gandavense.* The Gaunt Plum. 12 *Prunum Dactylites.* The Date Plum. 13 *Prunum Pyrum præcox.* The early Peare Plum.

and will abide good for ſix weekes at the leaſt after it is gathered, and after all other plums are ſpent.

The Cherry plum remembred before, ſpeaking of the Muſcadine plum, is a very good plum, but ſmall.

The Amber plum is a round plum, as yellow on the outſide almoſt as yellow waxe, of a ſowre vnpleaſant taſte that which I taſted, but I thinke it was not the right; for I haue ſeene and taſted another of the ſame bigneſſe, of a paler colour, farre better relliſhed, and a firmer ſubſtance, comming cleane from the ſtone like an Apricocke.

The Apricocke plum is a good plum when it is in its perfection, but that is ſeldome; for it doth moſt vſually cracke, thereby diminiſhing much of its goodneſſe, and beſides yeeldeth gumme at the crackes.

The Eaſon plum is a little red plum, but very good in taſte.

The Violet plum is a ſmall and long blackiſh blew plum, ripe about Bartholmew tide, a very good dry eating fruit.

The Grape plum is the Fluſhing Bulleis before remembred

The Dennie plum is called alſo the Cheſton, or the Friars plum before remembred.

The Damaske Violet plum, or Queen mother plum ſpoken of before.

The blacke Damaſcene plum is a very good dry plum, and of a darke blew colour when it is ripe.

The white Damſon is nothing ſo well relliſhed as the other.

The great Damſon or Damaske plum is greater then the ordinary Damſon, and ſweeter in taſte.

The blew Damſon well knowne, a good fruit.

The Coferers plum is flat, like vnto a Peare plum, it is early ripe and blacke, of a very good relliſh.

The Margate plum the worſt of an hundred.

The green Oyſterly plum is a reaſonable great plum, of a whitiſh green colour when it is ripe, of a moiſt and ſweete taſte, reaſonable good.

The red Mirobalane plum groweth to be a great tree quickly, ſpreading very thicke and farre, very like the blacke Thorne or Sloe buſh: the fruit is red, earlier ripe, and of a better taſte then the white.

The white Mirobalane plum is in moſt things like the former red, but the fruit is of a whitiſh yellow colour, and very pleaſant, eſpecially if it be not ouer ripe: both theſe had need to be plaſhed againſt a wall, or elſe they will hardly beare ripe fruit.

The Oliue plum is very like a greene Oliue, both for colour and bigneſſe, and groweth lowe on a ſmall buſhing tree, and ripeneth late, but is the beſt of all the ſorts of greene plums.

The white diapred plum of Malta, ſcarce knowne to any in our Land but Iohn Tradeſcante, is a very good plum, and ſtriped all ouer like diaper, and thereby ſo called.

The blacke diapred plum is like the Damaſcene plum, being blacke with ſpots, as ſmall as pins points vpon it, of a very good relliſh.

The Peake plum is a long whitiſh plum, and very good.

The Piſhamin or Virginia plum is called a plum, but vtterly differeth from all ſorts of plums, the deſcription whereof may truely enforme you, as it is ſet downe in the tenth Chapter going before, whereunto I referre you.

## The Vſe of Plums.

The great Damaske or Damſon Plummes are dryed in France in great quantities, and brought ouer vnto vs in Hogs-heads, and other great veſſels, and are thoſe Prunes that are vſually ſold at the Grocers, vnder the name of Damaske Prunes: the blacke Bulleis alſo are thoſe (being dryed in the ſame manner) that they call French Prunes, and by their tartneſſe are thought to binde, as the other, being ſweet, to looſen the body.

The Bruneola Plumme, by reaſon of his pleaſant tartneſſe, is much accounted of, and being dryed, the ſtones taken from them, are brought ouer to vs in ſmall boxes, and ſold deere at the Comfitmakers, where they very often accompany all other ſorts of banquetting ſtuffes.

Some

Some of these Plums, because of their firmnesse, are vndoubtedly more wholsome then others that are sweete and watersh, and cause lesse offence in their stomackes that eate them; and therefore are preserued with Sugar, to be kept all the yeare. None of them all is vsed in medicines so much as the great Damson or Damaske Prune, although all of them for the most part doe coole, lenifie, and draw forth choller, and thereby are fittest to be vsed of such as haue chollericke Agues.

## Chap. XIIII.

### *Mala Armeniaca sive Præcocia.* Apricockes.

THe Apricocke (as I said) is without question a kinde of Plumme, rather then a Peach, both the flower being white, and the stone of the fruit smooth also, like a Plumme, and yet because of the excellencie of the fruit, and the difference therein from all other Plummes, I haue thought it meete to entreate thereof by it selfe, and shew you the varieties haue been obserued in these times.

The Apricocke tree riseth vp to a very great height, either standing by it selfe (where it beareth not so kindly, and very little in our country) or planted against a wall, as it is most vsuall, hauing a great stemme or body, and likewise many great armes or branches, couered with a smooth barke: the leaues are large, broad, and almost round, but pointed at the ends, and finely dented about the edges: the flowers are white, as the Plumme tree blossomes, but somewhat larger, and rounder set: the fruit is round, with a cleft on the one side, somewhat like vnto a Peach, being of a yellowish colour as well on the inside as outside, of a firme or fast substance, and dry, not ouermoist in the eating, and very pleasant in taste, containing within it a broad and flat stone, somewhat round and smooth, not rugged as the Peach stone, with a pleasant sweete kernell (yet some haue reported, that there is such as haue their kernels bitter, which I did neuer see or know) and is ripe almost with our first or earliest Plummes, and thereof it tooke the name of *Præcox*; and it may bee was the earliest of all others was then knowne, when that name was giuen.

The great Apricocke, which some call the long Apricocke, is the greatest and fairest of all the rest.

The smaller Apricocke, which some call the small round Apricocke, is thought to be small, because it first sprang from a stone: but that is not so; for the kinde it selfe being inoculated, will bee alwaies small, and neuer halfe so faire and great as the former.

The white Apricocke hath his leaues more folded together, as if it were halfe double: it beareth but seldome, and very few, which differ not from the ordinary, but in being more white, without any red when it is ripe.

The Mascoline Apricocke hath a finer greene leafe, and thinner then the former, and beareth very seldome any store of fruit, which differeth in nothing from the first, but that it is a little more delicate.

The long Mascoline Apricocke hath his fruit growing a little longer then the former, and differeth in nothing else.

The Argier Apricocke is a smaller fruit then any of the other, and yellow, but as sweete and delicate as any of them, hauing a blackish stone within it, little bigger then a Lacure Cherry stone: this with many other sorts Iohn Tradescante brought with him returning from the Argier voyage, whither hee went voluntary with the Fleete, that went against the Pyrates in the yeare 1620.

### The Vse of Apricockes.

Apricockes are eaten oftentimes in the same manner that other dainty Plummes are, betweene meales of themselues, or among other fruit at banquets.

They are alſo preſerued and candid, as it pleaſeth Gentlewomen to beſtowe their time and charge, or the Comfitmaker to ſort among other candid fruits.

Some likewiſe dry them, like vnto Peares, Apples, Damſons, and other Plummes.

Matthiolus doth wonderfully commend the oyle drawne from the kernels of the ſtones, to annoint the inflamed *hæmorrhoides* or piles, the ſwellings of vlcers, the roughneſſe of the tongue and throate, and likewiſe the paines of the eares.

## Chap. XV.

### *Mala Perſica.* Peaches.

AS I ordered the Cherries and Plummes, ſo I intend to deale with Peaches, becauſe their varieties are many, and more knowne in theſe dayes then in former times: but becauſe the Nectorin is a differing kinde of Peach, I muſt deale with it as I did with the Apricocke among the Plummes, that is, place it in a Chapter by it ſelfe.

The Peach tree of it ſelfe groweth not vſually altogether ſo great, or high as the Apricocke, becauſe it is leſſe durable, but yet ſpreadeth with faire great branches, from whence ſpring ſmaller and ſlenderer reddiſh twigges, whereon are ſet long narrow greene leaues, dented about the edges: the bloſſomes are greater then of any Plumme, of a deepe bluſh or light purple colour: after which commeth the fruit, which is round, and ſometimes as great as a reaſonable Apple or Pippin (I ſpeake of ſome ſorts; for there be ſome kindes that are much ſmaller) with a furrow or cleft on the one ſide, and couered with a freeſe or cotton on the outſide, of colour either ruſſet, or red, or yellow, or of a blackiſh red colour; of differing ſubſtances and taſtes alſo, ſome being firme, others wateriſh, ſome cleauing faſt to the ſtone on the inſide, others parting from it more or leſſe eaſily, one excelling another very farre, wherein is contained a rugged ſtone, with many chinkes or clefts in it, the kernell whereof is bitter: the roots growe neither deepe nor farre; and therefore are ſubiect to the winds, ſtanding alone, and not againſt a wall. It ſooner waxeth old and decayeth, being ſprung of a ſtone, then being inoculated on a Plumme ſtocke, whereby it is more durable.

The great white Peach is white on the outſide as the meate is alſo, and is a good well relliſhed fruit.

The ſmall white Peach is all one with the greater, but differeth in ſize.

The Carnation Peach is of three ſorts, two are round, and the third long; they are all of a whitiſh colour, ſhadowed ouer with red, and more red on the ſide is next the ſunne: the leſſer round is the more common, and the later ripe.

The grand Carnation Peach is like the former round Peach, but greater, and is as late ripe, that is, in the beginning of September.

The red Peach is an exceeding well relliſhed fruit.

The ruſſet Peach is one of the moſt ordinary Peaches in the Kingdome, being of a ruſſet colour on the outſide, and but of a reaſonable relliſh, farre meaner then many other.

The Iſland Peach is a faire Peach, and of a very good relliſh.

The Newington Peach is a very good Peach, and of an excellent good relliſh, being of a whitiſh greene colour on the outſide, yet halfe reddiſh, and is ripe about Bartholmew tide.

The yellow Peach is of a deepe yellow colour; there be hereof diuers ſorts, ſome good and ſome bad.

The St. Iames Peach is the ſame with the Queenes Peach, here belowe ſet downe, although ſome would make them differing.

The Melocotone Peach is a yellow faire Peach, but differing from the former yellow both in forme and taſte, in that this hath a ſmall crooked end or point for the moſt part, it is ripe before them, and better relliſhed then any of them.

The

1 *Malus Armeniaca sive Præcoxia.* The Apricoeke. 2 *Malus Persica Melocotonea.* The Melocotone Peach. 3 *Persica Moschatellina,* The Nutmeg Peach 4 *Persica nigra.* The blacke Peach. 5 *Persica Carnea longa.* The long Carnation Peach. 6 *Persica Reginea.* The Queenes Peach. 7 *Amygdalus,* The Almond. 8 *Persica de Troas.* The Peach du Troas. 9 *Nuciperfica rubra optima,* The best Romane red Nectorin. 10. *Nuciperfica rubra altera,* The bastard red Nectorin with a pincking blossome.

The Peach *du Troas* is a long and great whitiſh yellow Peach, red on the outſide, early ripe, and is another kinde of Nutmeg Peach.

The Queenes Peach is a faire great yellowiſh browne Peach, ſhadowed as it were ouer with deepe red, and is ripe at Bartholmew tide, of a very pleaſant good taſte.

The Romane Peach is a very good Peach, and well relliſhed.

The Duraſme or Spaniſh Peach is of a darke yellowiſh red colour on the outſide, and white within.

The blacke Peach is a great large Peach, of a very darke browne colour on the outſide, it is of a wateriſh taſte, and late ripe.

The Alberza Peach is late ripe, and of a reaſonable good taſte.

The Almond Peach, ſo called, becauſe the kernell of the ſtone is ſweete, like the Almond, and the fruit alſo ſomewhat pointed like the Almond in the huske; it is early ripe, and like the Newington Peach, but leſſer.

The Man Peach is of two ſorts, the one longer then the other, both of them are good Peaches, but the ſhorter is the better relliſhed.

The Cherry Peach is a ſmall Peach, but well taſted.

The Nutmeg Peach is of two ſorts, one that will be hard when it is ripe, and eateth not ſo pleaſantly as the other, which will bee ſoft and mellow; they are both ſmall Peaches, hauing very little or no reſemblance at all to a Nutmeg, except in being a little longer then round, and are early ripe.

Many other ſorts of Peaches there are, whereunto wee can giue no eſpeciall name; and therefore I paſſe them ouer in ſilence.

### The Vſe of Peaches.

Thoſe Peaches that are very moiſt and wateriſh (as many of them are) and not firme, doe ſoone putrefie in the ſtomacke, cauſing ſurfeits oftentimes; and therefore euery one had neede bee carefull, what and in what manner they eate them: yet they are much and often well accepted with all the Gentry of the Kingdome.

The leaues, becauſe of their bitterneſſe, ſerue well being boyled in Ale or Milke, to be giuen vnto children that haue wormes, to help to kill them, and doe gently open the belly, if there be a ſufficient quantity vſed.

The flowers haue the like operation, that is, to purge the body ſomewhat more forceably then Damaske Roſes; a Syrupe therefore made of the flowers is very good.

The kernels of the Peach ſtones are oftentimes vſed to be giuen to them that cannot well make water, or are troubled with the ſtone; for it openeth the ſtoppings of the vritory paſſages, whereby much eaſe enſueth.

## Chap. XVI.

### *Nuciperſica.* Nectorins.

I Preſume that the name *Nuciperſica* doth moſt rightly belong vnto that kinde of Peach, which we call Nectorins, and although they haue beene with vs not many yeares, yet haue they beene knowne both in Italy to Matthiolus, and others before him, who it ſeemeth knew no other then the yellow Nectorin, as Dalechampius alſo: But we at this day doe know fiue ſeuerall ſorts of Nectorins, as they ſhall be preſently ſet downe; and as in the former fruits, ſo in this, I will giue you the deſcription of one, and briefe notes of the reſt.

The Nectorin is a tree of no great bigneſſe, moſt vſually leſſer then the Peach tree, his body and elder boughes being whitiſh, the younger branches very red, whereon grow narrow long greene leaues, ſo like vnto Peach leaues, that none can well diſtinguiſh them, vnleſſe it be in this, that they are ſomewhat leſſer: the bloſſomes are all reddiſh, as the Peach, but one of a differing faſhion from all the other, as I ſhall ſhew you by and by: the fruit that followeth is ſmaller, rounder, and ſmoother then Peaches, without any cleft on the ſide, and without any douny cotton or freeze at all; and herein

herein is like vnto the outer greene rinde of the Wallnut, whereof as I am perſwaded it tooke the name, of a faſt and firme meate, and very delicate in taſte, eſpecially the beſt kindes, with a rugged ſtone within it, and a bitter kernell.

The Muske Nectorin, ſo called, becauſe it being a kinde of the beſt red Nectorins, both ſmelleth and eateth as if the fruit were ſteeped in Muske: ſome thinke that this and the next Romane Nectorin are all one.

The Romane red Nectorin, or cluſter Nectorin, hath a large or great purpliſh bloſſome, like vnto a Peach, reddiſh at the bottome on the outſide, and greeniſh within: the fruit is of a fine red colour on the outſide, and groweth in cluſters, two or three at a ioynt together, of an excellent good taſte.

The baſtard red Nectorin hath a ſmaller or pincking bloſſome, more like threads then leaues, neither ſo large nor open as the former, and yellowiſh within at the bottome: the fruit is red on the outſide, and groweth neuer but one at a ioynt; it is a good fruit, but eateth a little more rawiſh then the other, euen when it is full ripe.

The yellow Nectorin is of two ſorts, the one an excellent fruit, mellow, and of a very good relliſh; the other hard, and no way comparable to it.

The greene Nectorin, great and ſmall; for ſuch I haue ſeene abiding conſtant, although both planted in one ground: they are both of one goodneſſe, and accounted with moſt to be the beſt relliſhed Nectorin of all others.

The white Nectorin is ſaid to bee differing from the other, in that it will bee more white on the outſide when it is ripe, then either the yellow or greene: but I haue not yet ſeene it.

### The Vſe of Nectorins.

The fruit is more firme then the Peach, and more delectable in taſte; and is therefore of more eſteeme, and that worthily.

## Chap. XVII.

### *Amygdala.* Almonds.

THe Almond alſo may be reckoned vnto the ſtock or kindred of the Peaches, it is ſo like both in leafe and bloſſome, and ſomewhat alſo in the fruit, for the outward forme, although it hath onely a dry skinne, and no pulpe or meate to bee eaten: but the kernell of the ſtone or ſhell, which is called the Almond, maketh recompenſe of that defect, whereof ſome are ſweete, ſome bitter, ſome great, ſome ſmall, ſome long, and ſome ſhort.

The Almond tree groweth vpright, higher and greater then any Peach; and is therefore vſually planted by it ſelfe, and not againſt a wall, whoſe body ſometime exceedeth any mans fadome, whereby it ſheweth to be of longer continuance, bearing large armes, and ſmaller branches alſo, but brittle, whereon are ſet long and narrow leaues, like vnto the Peach tree: the bloſſomes are purpliſh, like vnto Peach bloſſoms, but paler: the fruit is ſomewhat like a Peach for the forme of the skinne or outſide, which is rough, but not with any ſuch cleft therein, or with any pulpe or meate fit to bee eaten, but is a thicke dry skinne when it is ripe, couering the ſtone or ſhell, which is ſmooth and not rugged, and is either long and great, or ſmall, or thicke and ſhort, according as the nut or kernell within it is, which is ſweete both in the greater and ſmaller, and onely one ſmaller kinde which is bitter: yet this I haue obſerued, that all the Almond trees that I haue ſeene growe in England, both of the ſweete and bitter kindes, beare Almonds thicke and ſhort, and not long, as that ſort which is called the Iorden Almond.

### The Vſe of Almonds.

They are vſed many wayes, and for many purpoſes, either eaten alone with Figges, or Rayſins of the Sunne, or made into paſte with Sugar and Roſewater for Marchpanes, or put among Floure, Egges, and Sugar, to make

make Mackerons, or cruſted ouer with Sugar, to make Comfits, or mixed with Roſewater and Sugar, to make Butter, or with Barley water, to make Milke, and many other waies, as euery one liſt, that hath skill in ſuch things.

The oyle alſo of Almonds is vſed many waies, both inwardly and outwardly, for many purpoſes; as the oyle of ſweete Almonds mixt with poudered white Sugar Candy, for coughes and hoarſeneſſe, and to be drunk alone, or with ſome other thing (as the Syrupe of Marſh Mallowes) for the ſtone, to open and lenifie the paſſages, and make them ſlipperie, that the ſtone may paſſe the eaſier. And alſo for women in Child bed after their ſore trauell. And outwardly either by it ſelfe, or with oyle of Tartar to make a creame, to lenifie the skin, parched with the winde or otherwiſe, or to annoint the ſtomacke either alone, or with other things to helpe a cold.

The oyle of bitter Almonds is much vſed to be dropped into their eares that are hard of hearing, to helpe to open them. And as it is thought, doth more ſcoure and cleanſe the skin then the ſweet oyle doth, and is therefore more vſed of many for that purpoſe, as the Almonds themſelues are.

## Chap. XVIII.

### *Mala Arantia.* Orenges.

I Bring here to your conſideration, as you ſee, the Orenge tree alone, without mentioning the Citron or Lemmon trees, in regard of the experience we haue ſeen made of them in diuers places: For the Orenge tree hath abiden with ſome extraordinary looking and tending of it, when as neither of the other would by any meanes be preſerued any long time. If therefore any be deſirous to keepe this tree, he muſt ſo prouide for it, that it be preſerued from any cold, either in the winter or ſpring, and expoſed to the comfort of the ſunne in ſummer. And for that purpoſe ſome keepe them in great ſquare boxes, and lift them to and fro by iron hooks on the ſides, or cauſe them to be rowled by trundels, or ſmall wheeles vnder them, to place them in an houſe, or cloſe gallerie for the winter time: others plant them againſt a bricke wall in the ground, and defend them by a ſhed of boardes, couered ouer with ſeare-cloth in the winter, and by the warmth of a ſtoue, or other ſuch thing, giue them ſome comfort in the colder times: but no tent or meane prouiſion will preſerue them.

The Orenge tree in the warme Countries groweth very high, but with vs (or elſe it is a dwarfe kinde thereof) riſeth not very high: the barke of the elder ſtemmes being of a darke colour, and the young branches very greene, whereon grow here and there ſome few thornes: the leaues are faire, large, and very greene, in forme almoſt like a Bay leafe, but that it hath a ſmall eare, or peece of a leafe, faſhioned like vnto an heart vnder euery one of them, with many ſmall holes to be ſeene in them, if you hold them vp betweene you and the light, of a ſweet but ſtrong ſmell, naturally not falling away, but alwaies abiding on, or vntill new be come vp, bearing greene leaues continually: the flowers are whitiſh, of a very ſtrong and heady ſent; after which come ſmall round fruit, greene at the firſt, while they are ſmall, and not neere maturitie, but being grown and ripe, are (as all men know) red on the out ſide, ſome more pale then others, and ſome kindes of a deeper yellowiſh red, according to the climate, and as it receiueth the heate of the ſunne, wherein is contained ſower or ſweete iuice, and thicke white kernels among it: it beareth in the warme Countries both bloſſomes and greene fruit continually vpon it, and ripe fruit alſo with them for the beſt part of the yeare, but eſpecially in Autumne and Winter.

### The Vſe of Orenges.

Orenges are vſed as ſawce for many ſorts of meates, in reſpect of their ſweete ſowerneſſe, giuing a relliſh of delight, whereinſoeuer they are vſed.

The inner pulpe or iuice doth ſerue in agues and hot diſeaſes, and in Summer to coole the heate of deiected ſtomackes, or fainting ſpirits.

The

1 *Malus Arantia.* The Orenge tree. 2 *Malus.* The Apple tree. 3 *Malum Carbonarium.* The Pomewater. 4 *Malum Curtipendulum.* The golden Pippin.
5 *Melapium.* The Pearemaine. 6 *Malum Reginæum.* The Queene Apple. 7 *Malum primo maturum.* The Gennering. 8 *Malum Regale.* The pound Roy-
all. 9 *Malum Kentij ad fervescendum.* The Kentish Codlin. 10 *Malum Reginæum spurium.* The Bardfield Quining.

The dryed rinde, by reaſon of the ſweete and ſtrong ſent, ſerueth to bee put among other things to make ſweet pouthers.

The outer rindes, when they are clenſed from all the inner pulpe and skins, are preſerued in Sugar, after the bitterneſſe by often ſteepings hath been taken away,& do ſerue either as Succots,and banquetting ſtuffes, or as ornaments to ſet out diſhes for the table, or to giue a relliſh vnto meats, whether baked or boyled: Phyſically they helpe to warme a cold ſtomack, and to digeſt or breake winde therein: or they are candid with Sugar, and ſerue with other dryed Iunquets.

The water of Orange flowers is oftentimes vſed as a great perfume for glones, to waſhe them,or in ſtead of Roſe-water to mixe with other things.

It is vſed to bee drunke by ſome, to preuent or to helpe any peſtilentiall feuer.

The oyntment that is made of the flowers, is very comfortable both for the ſtomache, againſt the could or cough,or for the head,for paines and diſineſſe.

The kernels or ſeede beeing caſt into the grounnd in the ſpring time, will quickely grow vp, (but will not abide the winter with vs, to bee kept for growing trees) and when they are of a finger length high,being pluckt vp, and put among ſallats, will giue them a maruellous fine aromaticke or ſpicy taſte, very acceptable.

The ſeed or kernels are a little cordiall, although nothing ſo much as the kernels of the Pomecitron.

## Chap. XIX.

### *Poma.* Apples.

THe ſorts of Applesare ſo many, and infinite almoſt as I may ſay, that I cannot giue you the names of all, though I haue endeauoured to giue a great many, and I thinke it almoſt impoſſible for any one,to attaine to the full perfection of knowledge herein, not onely in regard of the multiplicitie of faſhions, colours and taſtes, but in that ſome are more familiar to one Countrey then to another, being of a better or worſe taſte in one place then in another, and therefore diuerſly called: I will therefore as I haue done before,giue you the deſcription of the Tree in generall,as alſo of the Paradiſe or dwarfe Apple, becauſe of ſome eſpeciall difference, and afterwards the names of as many, with their faſhions, as haue come to my knowledge, either by ſight or relation: for I doe confeſſe I haue not ſeene all that I here ſet downe, but vſe the helpe of ſome friends, and therefore if it happen that the ſeuerall names doe not anſwer vnto ſeuerall ſorts, but that the ſame fruit may bee called by one name in one Country, that is called by another elſewhere, excuſe it I pray you; for in ſuch a number, ſuch a fault may eſcape vnknowne.

The Apple tree for the moſt part is neyther very high, great or ſtraight, but rather vſually boweth and ſpreadeth(although in ſome places it groweth fairer and ſtraighter then in others) hauing long and great armes or boughes, and from them ſmaller branches, whereon doe grow ſomewhat broade, and long greene leaues, nicked about the edges: the flowers are large and white, with bluſh coloured ſides, conſiſting of fiue leaues: the fruit (as I ſaid) is of diuers formes, colours and taſtes, and likewiſe of a very variable durabilitie; for ſome muſt be eaten preſently after they are gathered, and they are for the moſt part the earlieſt ripe; others will abide longer vpon the trees, before they bee fit to be gathered; ſome alſo will be ſo hard when others are gathered, that they will not be fit to be eaten, for one, two or three months after they bee gathered; and ſome will abide good but one,two or three moneths,and no more; and ſome will be beſt,after a quarter or halfe a yeares lying, vnto the end of that yeare or the next.

The Paradiſe or dwarfe Apple tree groweth nothing ſo high as the former,and many times not much higher then a man may reach, hauing leaues and flowers altogether like the other, the fruit is a faire yellow Apple,and reaſonable great,but very light and ſpongy or looſe, and of a bitteriſh ſweet taſte, nothing pleaſant. And theſe faults al-

ſo

ſo are incident vnto this tree, that both bodie and branches are much ſubiect vnto cancker, which will quickely eate it round, and kill it; beſides it will haue many bunches, or tuberous ſwellings in many places, which grow as it were ſcabby or rough, and will ſoone cauſe it to periſh : the roote ſendeth forth many ſhootes and ſuckers, whereby it may be much increaſed. But this benefit may be had of it, to recompence the former faults, That being a dwarfe Tree, whatſoeuer fruit ſhall bee grafted on it, will keepe the graft low like vnto it ſelfe, and yet beare fruit reaſonable well. And this is a pretty way to haue Pippins, Pomewaters, or any other ſort of Apples (as I haue had my ſelfe, and alſo ſeene with others) growing low, that if any will, they may make a hedge rowe of theſe low fruits, planted in an Orchard all along by a walke ſide: but take this Caueat, if you will auoide the danger of the cancker and knots, which ſpoile the tree, to graft it hard vnto the ground, that therby you may giue as little of the nature of the ſtock thereunto as poſſibly you can, which wil vndoubtedly help it very much.

## The kindes or ſorts of Apples.

The Summer pippin is a very good apple firſt ripe, and therefore to bee firſt ſpent, becauſe it will not abide ſo long as the other.

The French pippin is alſo a good fruit and yellow.

The Golding pippin is the greateſt and beſt of all ſorts of pippins.

The Ruſſet pippin is as good an apple as moſt of the other ſorts of pippins.

The ſpotted pippin is the moſt durable pippin of all the other ſorts.

The ordinary yellow pippin is like the other, and as good; for indeed I know no ſort of pippins but are excellent good well relliſhed fruites.

The great pearemaine differeth little either in taſte or durabilitie from the pippin, and therefore next vnto it is accounted the beſt of all apples.

The ſummer pearemaine is of equall goodneſſe with the former, or rather a little more pleaſing, eſpecially for the time of its eating, which will not bee ſo long laſting, but is ſpent and gone when the other beginneth to be good to eate.

The Ruſſetting is alſo a firme and a very good apple, not ſo wateriſh as the pippin or pearemaine, and will laſt the beſt part of the year, but will be very mellow at the laſt, or rather halfe dryed.

The Broading is a very good apple.

The Pomewater is an excellent good and great whitiſh apple, full of ſap or moiſture, ſomewhat pleaſant ſharpe, but a little bitter withall : it will not laſt long, the winter froſts ſoone cauſing it to rot, and periſh.

The Flower of Kent is a faire yellowiſh greene apple both good and great.

The Gilloflower apple is a fine apple, and finely ſpotted.

The Marligo is the ſame, that is called the Marigold apple, it is a middle ſized apple, very yellow on the outſide, ſhadowed ouer as it were with red, and more red on one ſide, a reaſonable well relliſhed fruit.

The Blandrill is a good apple.

The Dauie Gentle is a very good apple

The Gruntlin is ſomewhat a long apple, ſmaller at the crowne then at the ſtalke, and is a reaſonable good apple.

The gray Coſterd is a good great apple, ſomewhat whitiſh on the outſide, and abideth the winter.

The greene Coſterd is like the other, but greener on the outſide continually.

The Haruy apple is a faire great goodly apple, and very well relliſhed.

The Dowſe apple is a ſweetiſh apple not much accounted of.

The Pome-paris is a very good apple.

The Belle boon of two ſorts winter and ſummer, both of them good apples, and fair fruit to look on, being yellow and of a meane bigneſſe.

The pound Royall is a very great apple, of a very good and ſharpe taſte.

The Doues Bill a ſmall apple.

The Deuſan or apple Iohn is a delicate fine fruit, well relliſhed when it beginneth to be fit to be eaten, and endureth good longer then any other apple.

The Maſter William is greater then a pippin, but of no very good relliſh.

The Maſter Iohn is a better taſted apple then the other by much.

The

The Spicing is a well tasted fruite.

*Pome de Rambures*, *Pome de Capanda*, *Pome de Calual* } all faire and good apples brought from France.

The Queene apple is of two sorts, both of them great faire red apples, and well relished, but the greater is the best.

The Bastard Queene apple is like the other for forme and colour, but not so good in taste: some call this the bardfield Queening.

The Boughton or greening is a very good and well tasted apple.

The Leathercoate apple is a good winter apple, of no great bignesse, but of a very good and sharpe taste.

The Pot apple is a plaine Country apple.

The Cowsnout is no very good fruit.

The Gildiling apple is a yellow one, not much accounted.

The Cats head apple tooke the name of the likenesse, and is a reasonable good apple and great.

The Kentish Codlin is a faire great greenish apple, very good to eate when it is ripe; but the best to coddle of all other apples.

The Stoken apple is a reasonable good apple.

The Geneting apple is a very pleasant and good apple.

The Worcester apple is a very good apple, as bigge as a Pomewater.

*Donime Conadis* is a French apple, and of a good relliſh.

The French Goodwin is a very good apple.

The old wife is a very good, and well relliſhed apple.

The towne Crab is an hard apple, not so good to be eaten rawe as roasted, but excellent to make Cider.

The Virgilling apple is a reasonable good apple.

The Crowes egge is no good relliſhed fruit, but nourſed vp in some places of the common people.

The Sugar apple is so called of the sweetnesse.

Sops in wine is so named both of the pleasantnesse of the fruit, and beautie of the apple.

The womans breast apple is a great apple.

The blacke apple or pippin is a very good eating apple, and very like a Pearemaine, both for forme and bignesse, but of a blacke sooty colour.

Tweenty sorts of Sweetings and none good.

The Peare apple is a small fruit, but well relliſhed being ripe, and is for shape very like vnto a small short Peare, and greene.

The Paradise apple is a faire goodly yellow apple, but light and spongy, and of a bitterish sweet taste, not to be commended.

The apple without blossome, so called because although it haue a small shew of a blossome, yet they are but small threds rather than leaues, neuer shewing to bee like a flower, and therefore termed without blossome: the apple is neyther good eating nor baking fruit.

Wildings and Crabs are without number or vse in our Orchard, being to be had out of the woods, fields and hedges rather then any where else.

## The Vse of Apples.

The best sorts of Apples serue at the last course for the table, in most mens houses of account, where, if there grow any rare or excellent fruit, it is then set forth to be seene and tasted.

Diuers other sorts serue to bake, either for the Masters Table, or the meynes sustenance, either in pyes or pans, or else stewed in dishes with Rose-water and Sugar, and Cinamon or Ginger cast vpon.

Some kinds are fittest to roast in the winter time, to warme a cup of wine, ale or beere; or to be eaten alone, for the nature of some fruit is neuer so good, or worth the eating, as when they are roasted.

Some

Some ſorts are fitteſt to ſcald for Codlins, and are taken to coole the ſtomacke, as well as to pleaſe the taſte, hauing Roſewater and Sugar put to them.

Some ſorts are beſt to make Cider of, as in the Weſt Countrey of England great quantities, yea many Hogſheads and Tunnes full are made, eſpecially to bee carried to the Sea in long voyages, and is found by experience to bee of excellent vſe, to mixe with water for beuerage. It is vſually ſeene that thoſe fruits that are neither fit to eate raw, roaſted, nor baked, are fitteſt for Cider, and make the beſt.

The iuice of Apples likewiſe, as of pippins, and pearemaines, is of very good vſe in Melancholicke diſeaſes, helping to procure mirth, and to expell heauineſſe.

The diſtilled water of the ſame Apples is of the like effect.

There is a fine ſweet oyntment made of Apples called *Pomatum*, which is much vſed to helpe chapt lips, or hands, or for the face, or any other part of the skinne that is rough with winde, or any other accident, to ſupple them, and make them ſmooth.

## Chap. XX.

### *Cydonia.* Quinces.

WEe haue ſome diuerſities of Quinces, although not many, yet more then our elder times were acquainted with, which ſhall be here expreſſed.

The Quince tree groweth oftentimes to the height and bigneſſe of a good Apple tree, but more vſually lower, with crooked and ſpreading armes and branches farre abroad, the leaues are ſomewhat round, and like the leaues of the Apple tree, but thicker, harder, fuller of veines, and white on the vnderſide: the bloſſomes or flowers are white, now and then daſht ouer with bluſh, being large and open, like vnto a ſingle Roſe: the fruit followeth, which when it is ripe is yellow, and couered with a white cotton or freeze, which in the younger is thicker and more plentifull, but waxeth leſſe and leſſe, as the fruit ripeneth, being bunched out many times in ſeuerall places, and round, eſpecially about the head, ſome greater, others ſmaller, ſome round like an Apple, others long like a Peare, of a ſtrong heady ſent, accounted not wholſome or long to be endured, and of no durabilitie to keepe, in the middle whereof is a core, with many blackiſh ſeedes or kernels therein, lying cloſe together in cels, and compaſſed with a kinde of cleare gelly, which is eaſier ſeene in the ſcalded fruit, then in the raw.

The Engliſh Quince is the ordinarie Apple Quince, ſet downe before, and is of ſo harſh a taſte being greene, that no man can endure to eate it rawe, but eyther boyled, ſtewed, roaſted or baked; all which waies it is very good.

The Portingall Apple Quince is a great yellow Quince, ſeldome comming to bee whole and faire without chapping; this is ſo pleaſant being freſh gathered, that it may be eaten like vnto an Apple without offence.

The Portingall Peare Quince is not fit to be eaten rawe like the former, but muſt be vſed after ſome of the waies the Engliſh Quince is appointed, and ſo it will make more dainty diſhes then the Engliſh, becauſe it is leſſe harſh, will bee more tender, and take leſſe ſugar for the ordering then the Engliſh kinde.

The Barbary Quince is like in goodneſſe vnto the Portingall Quince laſt ſpoken of, but leſſer in bigneſſe.

The Lyons Quince.

The Brunſwicke Quince.

### The Vſe of Quinces.

There is no fruit growing in this Land that is of ſo many excellent vſes as this, ſeruing as well to make many diſhes of meate for the table, as for ban-

banquets, and much more for the Phyſicall vertues, whereof to write at large is neither conuenient for mee, nor for this worke: I will onely briefly recite ſome, as it were to giue you a taſte of that plenty remaineth therein, to bee conuerted into ſundry formes: as firſt for the table, while they are freſh (and all the yeare long after being pickled vp) to be baked, as a dainty diſh, being well and orderly cookt. And being preſerued whole in Sugar, either white or red, ſerue likewiſe, not onely as an after diſh to cloſe vp the ſtomacke, but is placed among other Preſerues by Ladies and Gentlewomen, and beſtowed on their friends to entertaine them, and among other ſorts of Preſerues at Banquets. Codiniacke alſo and Marmilade, Ielly and Paſte, are all made of Quinces, chiefly for delight and pleaſure, although they haue alſo with them ſome phyſicall properties.

We haue for the vſe of phyſicke, both Iuyce and Syrupe, both Conſerue and Condite, both binding and looſening medicines, both inward and outward, and all made of Quinces.

The Ielly or Muccilage of the ſeedes, is often vſed to be laid vpon womens breaſts, to heale them being ſore or rawe, by their childrens default giuing them ſucke.

Athenæus reciteth in his third booke, that one Philarchus found, that the ſmell of Quinces tooke away the ſtrength of a certaine poiſon, called *Phariacum*. And the Spaniards haue alſo found, that the ſtrength of the iuyce of white Ellebor (which the Hunters vſe as a poyſon to dippe their arrow heads in, that they ſhoote at wilde beaſts to kill them) is quite taken away, if it ſtand within the compaſſe of the ſmell of Quinces. And alſo that Grapes, being hung vp to bee kept, and ſpent in Winter, doe quickly rot with the ſmell of a Quince.

## Chap. XXI.

### *Pyra.* Peares.

THe variety of peares is as much or more then of apples, and I thinke it is as hard in this, as before in apples, for any to be ſo exquiſite, as that hee could number vp all the ſorts that are to be had: for wee haue in our country ſo manie, as I ſhall giue you the names of by and by, and are hitherto come to our knowledge: but I verily beleeue that there be many, both in our country, and in others, that we haue not yet knowne or heard of; for euery yeare almoſt wee attaine to the knowledge of ſome, we knew not of before. Take therefore, according to the manner before held, the deſcription of one, with the ſeuerall names of the reſt, vntill a more exact diſcourſe be had of them, euery one apart.

The Peare tree groweth more ſlowly, but higher, and more vpright then the apple tree, and not leſſe in the bulke of the body: his branches ſpread not ſo farre or wide, but growe vprighter and cloſer: the leaues are ſomewhat broader and rounder, greene aboue, and whiter vnderneath then thoſe of the apple tree: the flowers are whiter and greater: the fruit is longer then round for the moſt part, ſmaller at the ſtalke, and greater at the head, of ſo many differing formes, colours, and taſtes, that hardly can one diſtinguiſh rightly between them, the times alſo being as variable in the gathering and ſpending of them, as in apples: the roote groweth deeper then the apple tree, and therefore abideth longer, and giueth a faſter, cloſer, & ſmoother gentle wood, eaſie to be wrought vpon.

#### The kindes of Peares.

The Summer bon Chretien is ſomewhat a long peare, with a greene and yellow ruſſetiſh coate, and will haue ſometimes red ſides; it is ripe at Michaelmas: ſome vſe to dry them as they doe Prunes, and keepe them all the yeare after. I haue not ſeene or heard any more Summer kindes hereof then this one, and needeth no wall to nourſe it as the other.

The

1 *Malus Cotonea*, The Quince tree. 2 *Cydonium Lusitanicum*. The Portingall Quince 3 *Pyrus*. The Peare tree. 4 *Pyrum Pompeianum, sive Cucumerinum byemale*. The Winter Bon Chretien. 5 *Pyrum pictum vel striatum*. The painted or striped Peare of Ierusalem. 6 *Pyrum Palatinale*. The Burgomot Peare. 7 *Pyrum Cucumerinum sive Pompeianum æstivum*. The Summer Bon Chretien. 8 *Pyrum Volemum*. The best Warden. 9 *Pyrum Librale*. The pound Peare. 10 *Pyrum Windsorienum*. The Windsor Peare. 11 *Pyrum Cucumerinum*. The Gratiola Peare. 12 *Pyrum Caryophyllatum*. The Gilloflower Peare.

The Winter bon Chretien is of many ſorts, ſome greater, others leſſer, and all good, but the greateſt and beſt is that kinde that groweth at Syon: All the kinds of this Winter fruit muſt be planted againſt a wall, or elſe they will both ſeldome beare, and bring fewer alſo to ripeneſſe, comparable to the wall fruit: the kindes alſo are according to their laſting; for ſome will endure good much longer then others.

The Summer Bergomot is an excellent well relliſhed peare, flattiſh, & ſhort, not long like others, of a meane bigneſſe, and of a darke yellowiſh greene colour on the outſide.

The Winter Bergomot is of two or three ſorts, being all of them ſmall fruit, ſomewhat greener on the outſide then the Summer kindes; all of them very delicate and good in their due time: for ſome will not be fit to bee eaten when others are will-nigh ſpent, euery of them outlaſting another by a moneth or more.

The Diego peare is but a ſmall peare, but an excellent well relliſhed fruit, taſting as if Muske had been put among it; many of them growe together, as it were in cluſters.

The Duetete or double headed peare, ſo called of the forme, is a very good peare, not very great, of a ruſſettiſh browne colour on the outſide.

The Primating peare is a good moiſt peare, and early ripe.

The Geneting peare is a very good early ripe peare.

The greene Cheſill is a delicate mellow peare, euen melting as it were in the mouth of the eater, although greeniſh on the outſide.

The Catherine peare is knowne to all I thinke to be a yellow red ſided peare, of a full wateriſh ſweete taſte, and ripe with the foremoſt.

The King Catherine is greater then the other, and of the ſame goodneſſe, or rather better.

The Ruſſet Catherine is a very good middle ſized peare.

The Windſor peare is an excellent good peare, well knowne to moſt perſons, and of a reaſonable greatneſſe: it will beare fruit ſome times twice in a yeare (and as it is ſaid) three times in ſome places.

The Norwich peare is of two ſorts, Summer and Winter, both of them good fruit, each in their ſeaſon.

The Worſter peare is blackiſh, a farre better peare to bake (when as it will be like a Warden, and as good) then to eate rawe; yet ſo it is not to be miſliked.

The Muske peare is like vnto a Catherine peare for bigneſſe, colour, and forme, but farre more excellent in taſte, as the very name importeth.

The Roſewater peare is a goodly faire peare, and of a delicate taſte.

The Sugar peare is an early peare, very ſweete, but wateriſh.

The Summer Popperin } both of them are very good firme dry peares, ſomewhat
The Winter Popperin } ſpotted, and browniſh on the outſide.

The greene Popperin is a winter fruit, of equall goodneſſe with the former.

The Souerainge peare, that which I haue ſeene and taſted, and ſo termed vnto me, was a ſmall browniſh yellow peare, but of a moſt dainty taſte; but ſome doe take a kind of Bon Chretien, called the Elizabeth peare, to be the Soueraigne peare; how truely let others iudge.

The Kings peare is a very good and well taſted peare.

The peare Royall is a great peare, and of a good relliſh.

The Warwicke peare is a reaſonable faire and good peare.

The Greenfield peare is a very good peare, of a middle ſize.

The Lewes peare is a browniſh greene peare, ripe about the end of September, a reaſonable well relliſhed fruit, and very moiſt.

The Biſhop peare is a middle ſized peare, of a reaſonable good taſte, not very wateriſh; but this property is oftentimes ſeene in it, that before the fruit is gathered, (but more vſually thoſe that fall of themſelues, and the reſt within a while after they are gathered) it will be rotten at the core, when there wil not be a ſpot or blemiſh to be ſeene on the outſide, or in all the peare, vntill you come neare the core.

The Wilford peare is a good and a faire peare.

The Bell peare a very good greene peare.

The Portingall peare is a great peare, but more goodly in ſhew then good indeed.

The Gratiola peare is a kinde of Bon Chretien, called the Cowcumber peare, or Spinola's peare.

The Rowling peare is a good peare, but hard, and not good before it bee a little rowled or bruiſed, to make it eate the more mellow.

The

The Pimpe peare is as great as the Windſor peare, but rounder, and of a very good relliſh.

The Turnep peare is a hard winter peare, not ſo good to eate rawe, as it is to bake.

The Arundell peare is moſt plentifull in Suffolke, and there commended to be a verie good peare.

The Berry peare is a Summer peare, reaſonable faire and great, and of ſo good and wholſome a taſte, that few or none take harme by eating neuer ſo many of them.

The Sand peare is a reaſonable good peare, but ſmall.

The Morley peare is a very good peare, like in forme and colour vnto the Windſor, but ſomewhat grayer.

The peare pricke is very like vnto the Greenfield peare, being both faire, great, and good.

The good Rewell is a reaſonable great peare, as good to bake as to eate rawe, and both wayes it is a good fruit.

The Hawkes bill peare is of a middle ſize, ſomewhat like vnto the Rowling peare.

The Petworth peare is a winter peare, and is great, ſomewhat long, faire, and good.

The Slipper peare is a reaſonable good peare.

The Robert peare is a very good peare, plentifull in Suffolke and Norfolke.

The pound peare is a reaſonable good peare, both to eate rawe, and to bake.

The ten pound peare, or the hundred pound peare, the trueſt and beſt, is the beſt Bon Chretien of Syon, ſo called, becauſe the grafts coſt the Maſter ſo much the fetching by the meſſengers expences, when he brought nothing elſe.

The Gilloflower peare is a winter peare, faire in ſhew, but hard, and not fit to bee eaten rawe, but very good to bake.

The peare Couteau is neither good one way nor other.

The Binſce peare is a reaſonable good winter peare, of a ruſſetiſh colour, and a ſmall fruit: but will abide good a long while.

The Pucell is a greene peare, of an indifferent good taſte.

The blacke Sorrell is a reaſonable great long peare, of a darke red colour on the outſide.

The red Sorrell is of a redder colour, elſe like the other.

The Surrine is no very good peare.

The Summer Haſting is a little greene peare, of an indifferent good relliſh.

Peare Gergonell is an early peare, ſomewhat long, and of a very pleaſant taſte.

The white Genneting is a reaſonable good peare, yet not equall to the other.

The Sweater is ſomewhat like the Windſor for colour and bigneſſe, but nothing neare of ſo good a taſte.

The bloud red peare is of a darke red colour on the outſide, but piercing very little into the inner pulpe.

The Hony peare is a long greene Summer peare.

The Winter peare is of many ſorts, but this is onely ſo called, to bee diſtinguiſhed from all other Winter peares, which haue ſeuerall names giuen them, and is a very good peare.

The Warden or Luke Wards peare of two ſorts, both white and red, both great and ſmall.

The Spaniſh Warden is greater then either of both the former, and better alſo.

The peare of Ieruſalem, or the ſtript peare, whoſe barke while it is young, is as plainly ſeene to be ſtript with greene, red, and yellow, as the fruit it ſelfe is alſo, and is of a very good taſte: being baked alſo, it is as red as the beſt Warden, whereof Maſter William Ward of Eſſex hath aſſured mee, who is the chiefe keeper of the Kings Granary at Whitehall.

Hereof likewiſe there is a wilde kinde no bigger then ones thumbe, and ſtriped in the like manner, but much more.

The Choke peares, and other wilde peares, both great and ſmall, as they are not to furniſh our Orchard, but the Woods, Forreſts, Fields, and Hedges, ſo wee leaue them to their naturall places, and to them that keep them, and make good vſe of them.

## The Vſe of Peares.

The moſt excellent ſorts of Peares, ſerue (as I ſaid before of Apples) to

make an after-course for their masters table, where the goodnesse of his Orchard is tryed. They are dryed also, and so are an excellent repaste, if they be of the best kindes, fit for the purpose.

They are eaten familiarly of all sorts of people, of some for delight, and of others for nourishment, being baked, stewed, or scalded.

The red Warden and the Spanish Warden are reckoned among the most excellent of Peares, either to bake or to roast, for the sicke or for the sound: And indeede, the Quince and the Warden are the two onely fruits are permitted to the sicke, to eate at any time.

Perry, which is the iuyce of Peares pressed out, is a drinke much esteemed as well as Cyder, to be both drunke at home, and carried to the Sea, and found to be of good vse in long voyages.

The Perry made of Choke Peares, notwithstanding the harshnesse, and cuill taste, both of the fruit when it is greene, as also of the iuyce when it is new made, doth yet after a few moneths become as milde and pleasant as wine; and will hardly bee knowne by the sight or taste from it: this hath beene found true by often experience; and therefore wee may admire the goodnesse of God, that hath giuen such facility to so wilde fruits, altogether thought vseless, to become vsefull, and apply the benefit thereof both to the comfort of our soules and bodies.

For the Physicall properties, if we doe as Galen teacheth vs, *in secundo Alimentorum*, referre the qualities of Peares to their seuerall tastes, as before he had done in Apples, we shall not neede to make a new worke; those that are harsh and sowre doe coole and binde, sweet do nourish and warme, and those betweene these, to haue middle vertues, answerable to their temperatures, &c.

*Much* more might be said, both of this and the other kinds of fruits; but let this suffice for this place and worke, vntill a more exact be accomplished.

## Chap. XXII.

### *Nux Iuglans.* The Wallnut.

ALthough the Wallnut tree bee often planted in the middle of great Courtyards, where by reason of his great spreading armes it taketh vp a great deale of roome, his shadow reaching farre, so that scarce any thing can well grow neare it; yet because it is likewise planted in fit places or corners of Orchards, and that it beareth fruit or nuts, often brought to the table, especially while they are freshest, sweetest, and fittest to be eaten, let not my Orchard want his company, or you the knowledge of it. Some doe thinke that there are many sorts of them, because some are much greater then others, and some longer then others, and some haue a more frangible shell then others; but I am certainly perswaded, that the soyle and climate where they grow, are the whole and onely cause of the varieties and differences. Indeed Virginia hath sent vnto vs two sorts of Wallnuts, the one blacke, the other white, whereof as yet wee haue no further knowledge. And I know that Clusius reporteth, he tooke vp at a banquet a long Wallnut, differing in forme and tenderneſſe of shell from others, which being set, grew and bore farre tenderer leaues then the other, and a little snipt about the edges, which (as I said) might alter with the soyle and climate: and besides you may obserue, that many of Clusius differences are very nice, and so I leaue it.

The Wallnut tree groweth very high and great, with a large and thicke body or trunke, couered with a thicke clouen whitish greene barke, tending to an ash-colour; the armes are great, and spread farre, breaking out into smaller branches, whereon doe grow long & large leaues, fiue or seuen set together one against another, with an odde one at the end, somewhat like vnto Ashen leaues, but farre larger, and not so many on a stalke, smooth, and somewhat reddish at the first springing, and tender also, of a reasonable good sent, but more strong and headie when they growe old: the fruit or nut is great and round, growing close to the stalkes of the leaues, either by couples or by

three

three set together, couered with a double shell, that is to say, with a greene thicke and soft outer rinde, and an inner hard shell, within which the white kernell is contained, couered with a thinne yellow rinde or peeling, which is more easily peeled away while it is greene then afterwards, and is as it were parted into foure quarters, with a thinne wooddy peece parting it at the head, very sweete and pleasant while it is fresh, and for a while after the gathering; but the elder they growe, the harder and more oily: the catkins or blowings are long and yellow, made of many scaly leaues set close together, which come forth early in the Spring, and when they open and fall away, vpon their stalkes arise certaine small flowers, which turne into so many nuts.

### The Vse of Wallnuts.

They are often serued to the table with other fruits while they abide fresh and sweete; and therefore many to keepe them fresh a long time haue deuised many wayes, as to put them into great pots, and bury them in the ground, and so take them out as they spend them, which is a very good way, and will keepe them long.

The small young nuts while they are tender, being preserued or candid, are vsed among other sorts of candid fruits, that serue at banquets.

The iuyce of the outer greene huskes are held to be a soueraigne remedy against either poyson, or plague, or pestilentiall feuer.

The distilled water of the huskes drunke with a little vinegar, if the fits growe hot and tedious, is an approued remedy for the same.

The water distilled from the leaues, is effectuall to be applyed to fluent or running vlcers, to dry and binde the humours.

Some haue vsed the pouder of the catkins in white wine, for the suffocation or strangling of the mother.

The oyle of Wallnuts is vsed to varnish Ioyners workes. As also is accounted farre to excell Linseede oyle, to mixe a white colour withall, that the colour bee not dimmed. It is of excellent vse for the coldnesse, hardnesse and contracting of the sinewes and ioynts, to warme, supple, and to extend them.

## Chap. XXIII.

### *Castanea Equina.* The Horse Chesnut.

ALthough the ordinary Chesnut is not a tree planted in Orchards, but left to Woods, Parkes, and other such like places; yet wee haue another sort which wee haue noursed vp from the nuts sent vs from Turky, of a greater and more pleasant aspect for the faire leaues, and of as good vse for the fruit. It groweth in time to be a great tree, spreading with great armes and branches, whereon are set at seuerall distances goodly faire great greene leaues, diuided into six, seuen, or nine parts or leaues, euery one of them nicked about the edges, very like vnto the leaues of *Ricnus*, or *Palma Christi*, and almost as great: it beareth at the ends of the branches many flowers set together vpon a long stalke, consisting of foure white leaues a peece, with many threads in the middle, which afterwards turne into nuts, like vnto the ordinary Chesnuts, but set in rougher and more prickly huskes: the nuts themselues being rounder and blacker, with a white spot at the head of each, formed somewhat like an heart, and of a little sweeter taste.

### The Vse of this Chesnut.

It serueth to binde and stop any maner of fluxe, be it of bloud or humours, either of the belly or stomacke; as also the much spitting of bloud. They are roasted and eaten as the ordinary sort, to make them taste the better.

They are vsually in Turkie giuen to horses in their prouender, to cure them of coughes, and helpe them being broken winded.

Chap.

## Chap. XXIIII.

### *Morus.* The Mulberrie.

THere are two ſorts of Mulberries ſufficiently known to moſt, the blackiſh and the white : but wee haue had brought vs from Virginia another ſort, which is of greater reſpect then eyther of the other two, not onely in regard of the raritie, but of the vſe, as you ſhall preſently vnderſtand.

### 1. *Morus nigra.* The blacke Mulberrie.

The blacke Mulberrie tree groweth oftentimes tall and great, and oftentimes alſo crooked, and ſpreading abroade, rather then high; for it is ſubiect to abide what forme you will conforme it vnto : if by ſuffering it to grow, it will mount vp, and if you will binde it, or plaſh the boughes, they will ſo abide, and be carried ouer arbours, or other things as you will haue it. The bodie groweth in time to bee very great, couered with a rugged or thicke barke, the armes or branches being ſmoother, whereon doe grow round thicke leaues pointed at the ends, and nicked about the edges, and in ſome there are to be ſeene deep gaſhes, making it ſeeme ſomewhat like the Vine leafe: the flowers are certaine ſhort dounie carkings, which turne into greene berries at the firſt, afterwards red, and when they are full ripe blacke, made of many graines ſet together, like vnto the blacke berrie, but longer and greater : before they are ripe, they haue an auſtere and harſh taſte, but when they are full ripe, they are more ſweete and pleaſant; the iuice whereof is ſo red, that it will ſtaine the hands of them that handle and eate them.

### 2. *Morus alba.* The white Mulberrie.

The white Mulberrie tree groweth not with vs to that greatneſſe or bulke of bodie that the blacke doth, but runneth vp higher, ſlenderer, more knotty, hard and brittle, with thinner ſpreade armes and branches : the leaues are like the former, but not ſo thicke ſet on the branches, nor ſo hard in handling, a little paler alſo, hauing ſomewhat longer ſtalkes: the fruit is ſmaller and cloſer ſet together, greene, and ſomewhat harſh before they be ripe, but of a wonderfull ſweetneſſe, almoſt ready to procure loathing when they are thorough ripe, and white, with ſuch like ſeede in them as in the former, but ſmaller.

### 3. *Morus Virginiana.* The Virginia Mulberrie.

The Virginia Mulberry tree groweth quickely with vs to be a very great tree, ſpreading many armes and branches, whereon grow faire great leaues, very like vnto the leaues of the white Mulberrie tree : the berry or fruit is longer and redder then either of the other, and of a very pleaſant taſte.

### The Vſe of Mulberries.

The greateſt and moſt eſpeciall vſe of the planting of white Mulberries, is for the feeding of Silke wormes, for which purpoſe all the Eaſterne Countries, as Perſia, Syria, Armenia, Arabia &c. and alſo the hither part of Turkie, Spaine alſo and Italie, and many other hot Countries doe nouriſh them, becauſe it is beſt for that purpoſe, the wormes feeding thereon, giuing the fineſt and beſt ſilke ; yet ſome are confident that the leaues of the blacke will doe as much good as the white : but that reſpect muſt be had to change your ſeede, becauſe therein lyeth the greateſt myſterie. But there is a Booke or Tractate printed, declaring the whole vſe of whatſoeuer can belong vnto them : I will therefore referre them thereunto, that would

1 *Nux Inglans.* The Wallnut. 2 *Caſtanea equina.* The horſe Cheſnut. 3 *Morus nigra vel alba.* The Mulberry. 4 *Morus Virginiana.* The Virginia Mulberry. 5 *Laurus vulgaris.* The ordinary Bay tree. 6 *Laurea Ceraſus Virginiana.* The Virginia Cherry Bay.

would further vnderſtand of that matter.

Mulberries are not much deſired to be eaten, although they be ſomewhat pleaſant, both for that they ſtaine their fingers and lips that eate them, and doe quickly putrefie in the ſtomacke, if they bee not taken before meate.

They haue yet a Phyſicall vſe, which is by reaſon of the aſtringent quality while they are red, and before they bee ripe, for ſore mouthes and throats, or the like, whereunto alſo the Syrup, called Diamoron, is effectuall.

## *Corollarium.*

## A COROLLARIE To this Orchard.

THere are certaine other trees that beare no fruit fit to bee eaten, which yet are often ſeene planted in Orchards, and other fit and conuenient places bout an houſe, whereof ſome are of eſpeciall vſe, as the Bay tree &c. others for their beauty and ſhadow are fit for walkes or arbours; ſome being euer green are moſt fit for hedge-rowes; and ſome others more for their raritie then for any other great vſe, wherof I thought good to entreat apart by themſelues, and bring them after the fruit trees of this Orchard, as an ornament to accompliſh the ſame.

### 1. *Laurus.* The Bay tree.

THere are to bee reckoned vp fiue kindes of Bay trees, three whereof haue been entreated of in the firſt part, a fourth wee will only bring here to your conſideration, which is that kinde that is vſually planted in euery mans yard or orchard, for their vſe throughout the whole land, the other we will leaue to bee conſidered of in that place is fit for it.

The Bay tree riſeth vp oftentimes to carry the face of a tree of a meane bigneſſe in our Countrey (although much greater in the hoter) and oftentimes ſhooteth vp with many ſuckers from the roote, ſhewing it ſelfe more like to a tall ſhrubbe or hedge-buſh, then a tree, hauing many branches, the young ones whereof are ſometimes reddiſh, but moſt vſually of a light or freſh greene colour, when the ſtemme and elder boughes are couered with a darke greene barke: the leaues are ſomewhat broad, and long pointed as it were at both the ends, hard and ſometimes crumpled on the edges, of a darke greene colour aboue, and of a yellowiſh greene vnderneath, in ſmell ſweet, in taſte bitter, and abiding euer greene: the flowers are yellow and moſſie, which turne into berries that are a little long as well as round, whoſe ſhell or outermoſt peele is greene at the firſt, and blacke when it is ripe; wherein is contained an hard bitter kernell, which cleaueth in two parts.

#### The Vſe of Bayes.

The Bay leaues are of as neceſſary vſe as any other in Garden or Orchard; for they ſerue both for pleaſure and profit, both for ornament and for vſe, both for honeſt Ciuill vſes, and for Phyſicke, yea both for the ſicke and for the ſound, both for the liuing and for the dead: And ſo much might be ſaid of this one tree, that if it were all told, would as well weary the Reader, as the Relater: but to explaine my ſelfe; It ſerueth to adorne the houſe of God as well as of man: to procure warmth, comfort and ſtrength to the limmes of men and women, by bathings and annoyntings outward, and by drinkes &c. inward to the ſtomacke, and other parts: to ſeaſon veſſels &c. wherein are preſerued our meates, as well as our drinkes: to crowne or encircle

circle as with a garland, the heads of the liuing, and to sticke and decke forth the bodies of the dead: so that from the cradle to the graue we haue still vse of it, we haue still neede of it.

The berries likewise serue for stitches inward, and for paines outward, that come of cold eyther in the ioynts, sinewes, or other places.

## 2. *Laurea Cerasus, siue Laurus Virginiana.* The Virginian Bay, or Cherry Baye.

THis Virginian (whether you will call it a Baye, or a Cherrie, or a Cherrie Bay, I leaue it to euery ones free will and iudgement, but yet I thinke I may as well call it a Bay as others a Cherrie, neither of them being answerable to the tree, which neyther beareth such berries as are like Cherries, neither beareth euer greene leaues like the Bay: if it may therefore bee called the Virginia Cherry Bay, for a distinction from the former Bay Cherry that beareth faire blacke Cherries, it will more fitly agree thereunto, vntill a more proper may be imposed) riseth vp to be a tree of a reasonable height, the stemme or bodie thereof being almost as great as a mans legge, spreading forth into diuers armes or boughes, and they againe into diuers small branches, whereon are set without order diuers faire broade greene leaues, somewhat like vnto the former Bay leaues, but more limber and gentle, and not so hard in handling, broader also, and for the most part ending in a point, but in many somewhat round pointed, very finely notched or toothed about the edges, of a bitter taste, very neere resembling the taste of the Bay leafe, but of little or no sent at all, either greene or dryed, which fall away euery autumne, and spring afresh euery yeare: the blossomes are small and white, many growing together vpon a long stalke, somewhat like the Bird Cherry blossomes, but smaller, and come forth at the ends of the young branches, which after turne into small berries, euery one set in a small cup or huske, greene at the first, and blacke when they are ripe, of the bignesse of a small pease, of a strong bitter taste, and somewhat aromaticall withall, but without any fleshy substance like a Cherry at all vpon it; for it is altogether like a berry.

### The Vse of this Virginia Cherry Bay.

Being a stranger in our Land, and possessed but of a very few, I doe not heare that there hath beene any triall made thereof what properties are in it: let this therefore suffice for this present, to haue shewed you the description and forme thereof, vntill we can learne further of his vses.

## 3. *Pinus.* The Pine tree.

MY purpose in this place is not to shew you all the diuersities of Pine trees, or of the rest that follow, but of that one kinde is planted in many places of our Land for ornament and delight, and there doth reasonably well abide: take it therefore into this Orchard, for the raritie and beautie of it, though we haue little other vse of it.

The Pine tree groweth with vs, though slowely, to a very great height in many places, with a great straight bodie, couered with a grayish greene barke, the younger branches are set round about, with very narrow long whitish greene leaues, which fall away from the elder, but abide on the younger, being both winter and summer alwaies greene. It hath growing in sundry places on the branches, certaine great hard wooddy clogs (called of some apples, of others nuts) composed of many hard woody scales, or tuberous knobs, which abide for the most part alwaies greene in our Countrey, and hardly become browniish, as in other Countries, where they haue more heat and comfort of the Sun, and where the scales open themselues; wherein are contained white long and round kernels, very sweete while they are fresh, but quickely growing oylely and rancide.

The

### The Vſe of the Pine apples and kernels.

The Cones or Apples are vſed of diuers Vintners in this City, being painted, to expreſſe a bunch of grapes, whereunto they are very like, and are hung vp in their buſhes, as alſo to faſten keyes vnto them, as is ſeene in many places.

The kernels within the hard ſhels, while they are freſh or newly taken out, are vſed many waies, both with Apothecaries, Comfit-makers, and Cookes : for of them are made medicines, good to lenifie the pipes and paſſages of the lungs and throate, when it is hoarſe. Of them are made Comfits, Paſtes, Marchpanes, and diuers other ſuch like : And with them a cunning Cooke can make diuers Keck ſhoſes for his Maſters table.

Matthiolus commendeth the water of the greene apples diſtilled, to take away the wrinkles in the face, to abate the ouer-ſwelling breaſts of Maidens, by fomenting them after with linnen clothes, wet in the water; and to reſtore ſuch as are rauiſht into better termes.

## 4. *Abies.* The Firre tree.

THe Firre tree groweth naturally higher then any other tree in theſe parts of Chriſtendome where no Cedars grow, and euen equalling or ouer-topping the Pine : the ſtemme or bodie is bare without branches for a great height, if they bee elder trees, and then branching forth at one place of the bodie foure wayes in manner of a croſſe, thoſe boughes againe hauing two branches at euery ioynt, on which are ſet on all ſides very thicke together many ſmall narrow long hard whitiſh greene leaues, and while they are young tending to yellowneſſe, but nothing ſo long or hard or ſharpe pointed as the Pine tree leaues, growing ſmaller and ſhorter to the end of the branches : the bloomings are certaine ſmall long ſcaly catkins, of a yellowiſh colour, comming forth at the ioynts of the branches, which fall away : the cones are ſmaller and longer then of the Pine tree, wherein are ſmall three ſquare ſeede contained, not halfe ſo big as the Pine kernels.

### The Vſe of the Firre tree.

The vſe of this tree is growne with vs of late daies to bee more frequent for the building of houſes then euer before : for hereof (namely of Deale timber and Deale boords) are framed many houſes, and their floores, without the helpe of any other timber or boord of any other tree almoſt ; as alſo for many other workes and purpoſes. The yellow Roſſen that is vſed as well to make ſalues as for many other common vſes, is taken from this tree, as the Pitch is both from the Pitch and Pine trees, and is boyled to make it to bee hard, but was at the firſt a yellow thin cleere Turpentine, and is that beſt ſort of common Turpentine is altogether in vſe with vs, as alſo another more thicke, whitiſh, and troubled, both which are vſed in ſalues, both for man and beaſt (but not inwardly as the cleere white Venice Turpentine is) and ſerueth both to draw, cleanſe and heale. Dodonæus ſeemeth to ſay, that the cleere white Turpentine, called Venice Turpentine, is drawn from the Firre : but Matthiolus confuteth that opinion, which Fulſius alſo held before him.

## 5. *Ilex arbor.* The euer-greene Oake.

THe *Ilex* or euer-greene Oake riſeth in time to be a very great tree, but very long and ſlow in growing (as is to be ſeene in the Kings priuy Garden at Whitehall, growing iuſt againſt the backe gate that openeth into the way going to Weſtminſter, and in ſome other places) ſpreading many fair large great armes and branches, whereon are ſet ſmall and hard greene leaues, ſomewhat endented or cornered, and prickly

1 *Pinus* The Pine tree. 2 *Abies.* The Firre tree. 3 *Ilex.* The euer greene Oake. 4 *Cupressus.* The Cipresse tree. 5 *Arbutus.* The Strawberry tree. 6 *Alaternus.* The euer greene Priuet.

prickly on the edges, especially in the young trees, and sometimes on those branches that are young and newly sprung forth from the elder rootes, but else in a manner all smooth in the elder growne, abiding greene all the winter as well as summer, and are of a grayish greene on the vnderside. It beareth in the spring time certaine slender long branches (like as other Okes doe) with small yellowish mossie flowers on them, which fall away, and are vnprofitable, the acornes not growing from those places, but from others which are like vnto those of our ordinary Oake, but smaller and blacker, and set in a more rugged huske or cuppe. This and no other kinde of *Ilex* doe I know to grow in all our land in any Garden or Orchard: for that kind with long and narrower leaues, and not prickly, growing so plentifully as Matthiolus saith in Tuscane, I haue not seen: and it is very probable to bee the same that Plinie remembreth to haue the leafe of an Oliue; but not as some would haue it, that *Smilax* Theophrastus maketh mention of in his third Booke and sixteenth Chapter of his Historie of Plants, which the Arcadians so called, and had the leafe of the *Ilex*, but not prickly: for Theophrastus saith, the timber of *Smilax* is smooth and soft, and this of the *Ilex* is harder, and stronger then an Oake.

### The Vse of the *Ilex* or euer-greene Oake.

Seeing this is to be accounted among the kindes of Oake (and all Oakes by Dioscorides his opinion are binding) it is also of the same qualitie, but a little weaker, and may serue to strengthen weake members. The young tops and leaues are also vsed in gargles for the mouth and throate.

---

### 6 *Cupressus*. The Cypresse tree.

THe Cypresse tree that is noursed vp by vs, in our Country, doth grow in those places where it hath beene long planted, to a very great height, whose bodie and boughes are couered with a reddish ash-coloured bark; the branches grow not spreading, but vpright close vnto the bodie, bushing thicke below, and small vpwards, spire fashion, those below reaching neere halfe the way to them aboue, whereon doe grow euer greene leaues, small, long and flat, of a resinous sweete smell, and strong taste, somewhat bitter: the fruit, which are called nuts, grow here and there among the boughes, sticking close vnto them, which are small, and clouen into diuers parts, but close while they are young, of a russetish browne colour; wherein are contained small browne seede, but not so small as motes in the Sunne, as Matthiolus and others make them to be.

### The Vse of the Cypresse tree.

For the goodly proportion this tree beareth, as also for his euer-greene head, it is and hath beene of great account with all Princes, both beyond, and on this side of the Sea, to plant them in rowes on both sides of some spatious walke, which by reason of their high growing, and little spreading, must be planted the thicker together, and so they giue a goodly, pleasant and sweet shadow: or else alone, if they haue not many, in the middle of some quarter, or as they thinke meete. The wood thereof is firme and durable, or neuer decaying, of a brown yellow colour, and of a strong sweete smell, whereof Chests or Boxes are made to keepe apparell, linnen, furres, and other things, to preserue them from moths, and to giue them a good smell.

Many Physicall properties, both wood, leaues and nuts haue, which here is not my purpose to vnfold, but only to tell you, that the leaues being boyled in wine, and drunke, helpe the difficultie of making vrine, and that the nuts are binding, fit to bee vsed to stay fluxes or laskes, and good also for ruptures.

### 7. *Arbutus.* The Strawberry tree.

THe Strawberry tree groweth but ſlowly, and riſeth not to the height of any great tree, no not in France, Italy, or Spaine: and with vs the coldneſſe of our country doth the more abate his vigour, ſo that it ſeldome riſeth to the height of a man: the barke of the body is rough, and ſmooth in the younger branches: the leaues are faire and greene, very like vnto Baye leaues, finely dented or ſnipped about the edges, abiding alwayes greene thereon both Winter and Summer: the flowers come forth at the end of the branches vpon long ſtalkes, not cluſtering thicke together, but in long bunches, and are ſmall, white, and hollow, like a little bottle, or the flower of Lilly Conually, which after turne into rough or rugged berries, moſt like vnto Strawberries (which hath giuen the name to the tree) ſomewhat reddiſh when they are ripe, of a harſh taſte, nothing pleaſant, wherein are contained many ſmall ſeedes: It hardly bringeth his fruit to ripeneſſe in our countrey; for in their naturall places they ripen not vntill Winter, which there is much milder then with vs.

#### The Vſe of the Strawberry tree.

Amatus Luſitanus I thinke is the firſt that euer recorded, that the water diſtilled from the leaues and flowers hereof, ſhould bee very powerfull againſt the plague and poyſons: for all the ancient Writers doe report, that the fruit hereof being eaten, is an enemy to the ſtomacke and head. And Cluſius likewiſe ſetteth downe, that at Liſhbone, and other places in Portingall where they are frequent, they are chiefly eaten, but of the poorer ſort, women and boyes. They are ſomewhat aſtringent or binding, and therefore may well ſerue for fluxes. It is chiefly nourſed with vs for the beauty and rareneſſe of the tree, for that it beareth his leaues alwayes green.

### 8. *Alaternus.* The euer greene Priuet.

THe tree which we haue growing in our country called *Alaternus*, groweth not to be a tree of any height; but abiding lowe, ſpreadeth forth many branches, whereon are ſet diuers ſmall and hard greene leaues, ſomewhat round for the forme, and endented a little by the edges: it beareth many ſmall whitiſh greene flowers at the ioynts of the ſtalkes, and ſetting on of the lower leaues cluſtering thicke together, which after turne into ſmall blacke berries, wherein are contained many ſmall graines or ſeedes: the beauty and verdure of theſe leaues abiding ſo freſh all the yeare, doth cauſe it to be of the greater reſpect; and therefore findeth place in their Gardens onely, that are curious conſeruers of all natures beauties.

#### The Vſe of the euer greene Priuet.

It is ſeldome vſed for any Phyſicall property, neither with vs, nor in the places where it is naturall and plentifull: but as Cluſius reporteth, hee learned that the Portingall Fiſhermen do dye their nets red with the decoction of the barke hereof, and that the Dyers in thoſe parts doe vſe the ſmall peeces of the wood to ſtrike a blackiſh blew colour.

### 9. *Celaſtrus Theophraſti Cluſio.* Cluſius his Celaſtrus.

ALthough the Collector (who is thought to be Ioannes Molineus of the great Herball or Hiſtory of plants, and generally bearing Daleſchampius name, becauſe the finding and relation of diuers herbes therein expreſſed, is appropiate to him, and printed at Lyons) of all our moderne Writers doth firſt of all others appoint the *Celaſtrus*, whereof Theophraſtus onely among all the ancient Writers of

plants maketh mention, to be the firſt *Alaternus* that Cluſius hath ſet forth in his Hiſtory of rarer plants: yet I finde, that Cluſius himſelfe before his death doth appropiate that *Celaſtrus* of Theophraſtus to another plant, growing in the Garden at Leyden, which formerly of diuers had beene taken to be a kinde of *Laurus Tinus*, or the wilde Baye; but he impugning that opinion for diuers reſpects, decyphreth out that Leyden tree in the ſame manner that I doe: and becauſe it is not onely faire, in bearing his leaues alwayes greene, but rare alſo, being nourſed vp in our Land in very few places, but principally with a good old Lady, the widow of Sir Iohn Leuſon, dwelling neere Rocheſter in Kent; I thought it fit to commend it for an ornament, to adorne this our Garden and Orchard. It groweth vp to the height of a reaſonable tree, the body whereof is couered with a darke coloured barke, as the elder branches are in like manner; the younger branches being greene, whereon are ſet diuers leaues thicke together, two alwayes at a ioynt, one againſt another, of a ſad but faire greene colour on the vpperſide, and paler vnderneath, which are little or nothing at all ſnipped about the edges, as large as the leaues of the *Laurus Tinus*, or wilde Baye tree: at the end of the young branches breake forth between the leaues diuers ſmall ſtalkes, with foure or fiue flowers on each of them, of a yellowiſh greene colour, which turne into ſmall berries, of the bigneſſe of blacke Cherries, greene at the firſt, and red when they begin to be ripe, but growing blacke if they hang too long vpon the branches, wherein is contained a hard ſhell, and a white hard kernell within it, couered with a yellowiſh skin. This abideth (as I ſaid before) with greene leaues as well Winter as Summer; and therefore fitteſt to be planted among other of the ſame nature, to make an euer greene hedge.

### The Vſe of Cluſius his Celaſtrus.

Being ſo great a ſtranger in this part of the Chriſtian world, I know none hath made tryall of what property it is, but that the taſte of the leaues is ſomewhat bitter.

## 10. *Pyracantha.* The euer greene Hawthorne, or prickly Corall tree.

THis euer greene ſhrubbe is ſo fine an ornament to a Garden or Orchard, either to be nourſed vp into a ſmall tree by it ſelfe, by pruining and taking away the ſuckers and vnder branches, or by ſuffering it to grow with ſuckers, thicke and plaſhing the branches into a hedge, for that it is plyable to be ordered either way; that I could not but giue you the knowledge thereof, with the deſcription in this manner. The younger branches are couered with a ſmooth darke blewiſh greene barke, and the elder with a more aſh coloured, thicke ſet with leaues without order, ſome greater and others ſmaller, ſomewhat like both in forme and bigneſſe vnto the leaues of the Barberry tree, but ſomewhat larger, and more ſnipt about the edges, of a deeper green colour alſo, and with ſmall long thornes ſcattered here & there vpon the branches: the flowers come forth as well at the ends of the branches, as at diuers places at the ioynts of the leaues, ſtanding thicke together, of a pale whitiſh colour, a little daſht ouer with a ſhew of bluſh, conſiſting of fiue leaues a peece, with ſome ſmall threads in the middle, which turne into berries, very like vnto Hawthorne berries, but much redder and dryer, almoſt like poliſhed Corall, wherein are contained foure or fiue ſmall yellowiſh white three ſquare ſeede, ſomewhat ſhining. It is thought to be the *Oxyacantha* of Dioſcorides; but ſeeing Dioſcorides doth explaine the forme of the leafe in his Chapter of Medlars, which he concealed in the Chapter of *Oxyacantha*, it cannot be the ſame: for *Meſpilus Anthedon* of Theophraſtus, or *Aronia* of Dioſcorides, hath the leafe of *Oxyacantha*, as Dioſcorides ſaith, or of Smalladge, as Theophraſtus, which cannot agree to this Thorne; but doth moſt liuely delineate out our white Thorne or Hawthorne, that now there is no doubt, but that *Oxyacantha* of Dioſcorides is the Hawthorne tree or buſh.

### The Vſe of this Corall tree.

Although Lobel maketh mention of this tree to grow both in Italy, and

1 *Celastrus Theophrasti Clusio.* Clusius his Celastrus. 2 *Pyracantha.* The euer green prickly Corall tree. 3 *Taxus.* The Yewe tree. 4 *Buxus arbor.* The Boxe tree 5 *Buxus humilis.* The lowe or dwarfe Boxe. 6 *Sabina.* The Sauine tree. 7 *Paliurus.* Christs thorne. 8 *Larix.* The Larch tree.

Prouence in France, in ſome of their hedges, yet he ſaith it is neglected in the naturall places, and to be of no vſe with them: neither doe I heare, that it is applyed to any Phyſicall vſe with vs, but (as I before ſaid) it is preſerued with diuers as an ornament to a Garden or Orchard, by reaſon of his euer greene leaues, and red berries among them, being a pleaſant ſpectacle, and fit to be brought into the forme of an hedge, as one pleaſe to lead it.

## 11. *Taxus*. The Yewe tree.

THe Yewe tree groweth with vs in many places to bee a reaſonable great tree, but in hoter countries much bigger, couered with a reddiſh gray ſcaly barke; the younger branches are reddiſh likewiſe, whereon grow many winged leaues, that is, many narrow long darke greene leaues, ſet on both ſides of a long ſtalke or branch, neuer dying or falling away, but abiding on perpetually, except it be on the elder boughes: the flowers are ſmall, growing by the leaues, which turne into round red berries, like vnto red Aſparagus berries, in taſte ſweetiſh, with a little bitterneſſe, and cauſing no harme to them for any thing hath been knowne in our country,

### The Vſe of the Yewe tree.

It is found planted both in the corners of Orchards, and againſt the windowes of Houſes, to be both a ſhadow and an ornament, in being alwayes greene, and to decke vp Houſes in Winter: but ancient Writers haue euer reckoned it to be dangerous at the leaſt, if not deadly.

## 12. *Buxus*. The Boxe tree.

THe Boxe tree in ſome places is a reaſonable tall tree, yet growing ſlowly; the trunke or body whereof is of the bigneſſe of a mans thigh, which is the biggeſt that euer I ſaw: but ſometimes, and in other places it groweth much lower, vſually not aboue a yard, or a yard and a halfe high, on the backe ſides of many Houſes, and in the Orchards likewiſe: the leaues are ſmall, thicke and hard, and ſtill the greater or leſſer the tree is, the greater or leſſer are the leaues, round pointed, and of a freſh ſhining greene colour: the flowers are ſmall and greeniſh, which turne into heads or berries, with foure hornes, whittiſh on the outſide, and with reddiſh ſeede within them.

*Buxus aureus.* Gilded Boxe. There is another kinde hereof but lately come to our knowledge, which differeth not in any thing from the former, but onely that all the leaues haue a yellow liſt or gard about the edge of them on the vpperſide, and none on the lower, which maketh it ſeeme very beautifull; and is therefore called gilded Boxe.

*Buxus humilis.* Dwarfe Boxe. We haue yet another kinde of Boxe, growing ſmall and lowe, not aboue halfe a foote, or a foote high at the moſt, vnleſſe it be neglected, which then doth grow a little more ſhrubby, bearing the like leaues, but ſmaller, according to the growth, and of a deeper greene colour: I could neuer know that this kinde euer bore flower or ſeede, but is propagated by ſlipping the roote, which encreaſeth very much.

### The Vſe of Boxe.

The wood of the Boxe tree is vſed in many kindes of ſmall works among Turners, becauſe it is hard, cloſe, and firme, and as ſome haue ſaid, the roots much more, in regard of the diuers waues and crooked veines running through it. It hath no Phyſicall vſe among the moſt and beſt Phyſitians, although ſome haue reported it to ſtay fluxes, and to be as good as the wood of *Guaiacum*, or *Lignum vitæ* for the French diſeaſe. The leaues and branches ſerue both Summer and Winter to decke vp houſes; and are many times giuen to horſes for the bots.

The lowe or dwarfe Boxe is of excellent vſe to border vp a knot, or the

lon

long beds in a Garden, being a maruailous fine ornament thereunto, in regard it both groweth lowe, is euer greene, and by cutting may bee kept in what maner euery one pleaſe, as I haue before ſpoken more largely.

13. *Sabina.* The Sauine tree or buſh.

THe Sauine tree or buſh that is moſt vſuall in our country, is a ſmall lowe buſh, not ſo high as a man in any place, nor ſo bigge in the ſtemme or trunke as a mans arme, with many crooked bending boughes and branches, whereon are ſet many ſmall, ſhort, hard, and prickly leaues, of a darke green colour, freſh and green both Winter and Summer: it is reported, that in the naturall places it beareth ſmall blacke berries, like vnto Iuniper, but with vs it was neuer knowne to beare any.

The Vſe of Sauine.

It is planted in out-yards, backſides, or voide places of Orchards, as well to caſt clothes thereon to dry, as for medicines both for men and horſes: being made into an oyle, it is good to annoint childrens bellies for to kill the Wormes: and the powder thereof mixed with Hogs greaſe, to annoint the running ſores or ſcabs in their heads; but beware how you giue it inwardly to men, women, or children. It is often put into horſes drenches, to helpe to cure them of the bots, and other diſeaſes.

14. *Paliurus.* Chriſts thorne.

THis thorny ſhrubbe (wherewith as it is thought, our Sauiour Chriſt was crowned, becauſe as thoſe that haue trauelled through Paleſtina and Iudæa, doe report no other thorne doth grow therein ſo frequent, or ſo apt to be writhed) riſeth in ſome places to a reaſonable height, but in our country ſeldome exceedeth the height of a man, bearing many ſlender branches, full of leaues, ſet on either ſide thereof one by one, which are ſomewhat broad and round, yet pointed, and full of veines, thicke ſet alſo with ſmall thornes, euen at the foote of euery branch, and at the foote of euery leafe one or two, ſome ſtanding vpright, others a little bending downe: the flowers are ſmall and yellow, ſtanding for the moſt part at the end of the branches, many growing vpon a long ſtalke, which after turne into round, flat, and hard ſhelly fruit, yet couered with a ſoft fleſhy skinne, within which are included two or three hard, ſmall, and browne flat ſeeds, lying in ſeuerall partitions. The leaues hereof fall away euery yeare, and ſpring forth afreſh againe the next May following. The rarity and beauty of this ſhrubbe, but chiefly (as I thinke) the name hath cauſed this to be much accounted of with all louers of plants.

The Vſe of Chriſts thorne.

Wee haue ſo few of theſe ſhrubbes growing in our country, and thoſe that are, doe, for any thing I can vnderſtand, neuer beare fruit with vs; that there is no other vſe made hereof then to delight the owners: but this is certainly receiued for the *Paliurus* of Dioſcorides and Theophraſtus, and thought alſo by Matthiolus to be the very true *Rhamnus tertius* of Dioſcorides. Matthiolus alſo ſeemeth to contradict the opinion is held by the Phyſitians of Mompelier, and others, that it cannot be the *Paliurus* of Theophraſtus. It is held to be effectuall to helpe to breake the ſtone, both in the bladder, reines, and kidneyes: the leaues and young branches haue an aſtringent quality, and good againſt poyſons and the bitings of ſerpents.

15. *Larix.*

### 15. *Larix.* The Larch tree.

THe Larch tree, where it naturally groweth, riſeth vp to be as tall as the Pine or Firre tree, but in our Land being rare, and nourſed vp but with a few, and thoſe onely louers of rarities, it groweth both ſlowly, and becommeth not high: the barke hereof is very rugged and thicke, the boughes and branches grow one aboue another in a very comely order, hauing diuers ſmall yellowiſh knobs or bunches ſet thereon at ſeuerall diſtances; from whence doe yearely ſhoote forth many ſmall, long, and narrow ſmooth leaues together, both ſhorter and ſmaller, and not ſo hard or ſharpe pointed as either the Pine or Firre tree leaues, which doe not abide the Winter as they doe, but fall away euery yeare, as other trees which ſhed their leaues, and gaine freſh euery Spring: the bloſſomes are very beautifull and delectable, being of an excellent fine crimſon colour, which ſtanding among the greene leaues, allure the eyes of the beholders to regard it with the more deſire: it alſo beareth in the naturall places (but not in our Land that I could heare) ſmall ſoft cones or fruit, ſomewhat like vnto Cypreſſe nuts, when they are greene and cloſe.

#### The Vſe of the Larch tree.

The coles of the wood hereof (becauſe it is ſo hard and durable as none more) is held to be of moſt force being fired, to cauſe the Iron oare to melt, which none other would doe ſo well. Matthiolus conteſteth againſt Fuchſius, for deeming the Venice Turpentine to be the liquid Roſſen of the Firre tree, which he aſſureth ~~vpon his owne~~ experience and certaine knowledge, to be drawne from this Larch tree, and none other; which cleere Turpintine is altogether vſed inwardly, and no other, except that of the true Turpintine tree, and is very effectuall to cleanſe the reines, kidneyes, and bladder, both of grauell and the ſtone, and to prouoke vrine: it is alſo of eſpeciall property for the *gonorrhœa*, or running of the reines, as it is called, with ſome powder of white Amber mixed therewith, taken for certaine dayes together. Taken alſo in an Electuary, it is ſingular good for to expectorate rotten flegme, and to helpe the conſumption of the lungs. It is vſed in plaiſters and ſalues, as the beſt ſort of Turpintine. The Agaricke that is vſed in phyſicke, is taken from the bodies and armes of this tree. And Matthiolus doth much inſiſt againſt Braſauolus, that thought other trees had produced Agaricke, affirming them to be hard *Fungi*, or Muſhroms (ſuch as wee call Touch-wood) wherwith many vſe to take fire, ſtrooke thereinto from ſteele.

---

### 16. *Tilia.* The Line or Linden tree.

THere are two ſorts of Line trees, the male and the female; but becauſe the male is rare to be ſeene, and the female is more familiar, I will onely giue you the deſcription of the female, and leaue the other.

The female Line tree groweth exceeding high and great, like vnto an Elme, with many large ſpreading boughes, couered with a ſmooth barke, the innermoſt being very plyant and bending from whence come ſmaller branches, all of them ſo plyable, that they may bee led or carried into any forme you pleaſe: the leaues thereon are very faire, broad, and round, ſomewhat like vnto Elme leaues, but fairer, ſmoother, and of a freſher greene colour, dented finely about the edges, and ending in a ſharpe point: the flowers are white, and of a good ſmell, many ſtanding together at the top of a ſtalke, which runneth all along the middle ribbe of a ſmall long whitiſh leafe; after which come ſmall round berries, wherein is contained ſmall blackiſh ſeede: this tree is wholly neglected by thoſe that haue them, or dwell neere them, becauſe they ſuppoſe it to be fruitleſſe, in regard it beareth chaffie huskes, which in many places fall away, without giuing ripe ſeede.

The

1 *Tilia fœmina.* The Line or Linden tree. 2 *Tamariſcus.* The Tamariske tree. 3 *Acer maius latifolium.* The Sycomore tree. 4 *Staphylodendron.* The bladder nut. 5 *Rhus Myrtifolia.* The Mirtle leafed Sumach. 6 *Rhus Virginiana.* The Buck-horne tree. 7 *Vitis ſeu potius Hedera Virginenſis.* The Virginia Vine or rather Iuie.

### The Vſe of the Line tree.

It is planted both to make goodly Arbours, and Summer banquetting houſes, either belowe vpon the ground, the boughes ſeruing very handſomely to plaſh round about it, or vp higher, for a ſecond aboue it, and a third alſo: for the more it is depreſſed, the better it will grow. And I haue ſeene at Cobham in Kent, a tall or great bodied Line tree, bare without boughes for eight foote high, and then the branches were ſpread round about ſo orderly, as if it were done by art, and brought to compaſſe that middle Arbour: And from thoſe boughes the body was bare againe for eight or nine foote (wherein might bee placed halfe an hundred men at the leaſt, as there might be likewiſe in that vnderneath this) & then another rowe of branches to encompaſſe a third Arbour, with ſtayres made for the purpoſe to this and that vnderneath it: vpon the boughes were laid boards to tread vpon, which was the goodlieſt ſpectacle mine eyes euer beheld for one tree to carry.

The coles of the wood are the beſt to make Gunpowder. And being kindled, and quenched in vinegar, are good to diſſolue clotted bloud in thoſe that are bruiſed with a fall. The inner barke being ſteeped in water yeeldeth a ſlimie iuyce, which is found by experience, to be very profitable for them that haue been burnt with fire.

---

### 17. *Tamarix.* Tamariske tree.

THe Tamariske tree that is common in our country, although in ſome places it doth not grow great, yet I haue ſeene it in ſome other, to be as great as a great apple tree in the body, bearing great arms; from whoſe ſmaller branches ſpring forth young ſlender red ſhootes, ſet with many very fine, ſmall, and ſhort leaues, a little criſped, like vnto the leaues of Sauine, not hard or rough, but ſoft and greene: the flowers be white moſſie threads, which turne into dounie ſeede, that is carried away with the winde.

*Tamariſcus foliis ablidis.* White Tamariske.

There is another kinde hereof very beautifull and rare, not to be ſeene in this Land I thinke, but with Mr. William Ward, the Kings ſeruant in his Granary, before remembred, who brought me a ſmall twigge to ſee from his houſe at Boram in Eſſex, whoſe branches are all red while they are young, and all the leaues white, abiding ſo all the Summer long, without changing into any ſhew of greene like the other, and ſo abideth conſtant yeare after yeare, yet ſhedding the leaues in Winter like the other.

### The Vſe of Tamariske.

The greateſt vſe of Tamariske is for ſpleneticke diſeaſes, either the leaues or the barke made into drinkes; or the wood made into ſmall Cans or Cups to drinke in.

---

### 18. *Acer maius latifolium.* The great Maple or Sycomore tree.

THe Sycomore tree, as we vſually call it (and is the greateſt kind of Maple, cheriſhed in our Land onely in Orchards, or elſewhere for ſhade and walkes, both here in England, and in ſome other countries alſo) groweth quickly to bee a faire ſpreading great tree, with many boughes and branches, whoſe barke is ſomewhat ſmooth: the leaues are very great, large, and ſmooth, cut into foure or fiue diuiſions, and ending into ſo many corners, euery one ſtanding on a long reddiſh ſtalke: the bloomings are of a yellowiſh greene colour, growing many together on each ſide of a long ſtalke, which after turne into long and broad winged ſeede, two alwaies ſtanding together on a ſtalke, and bunched out in the middle, where the ſeed or kernell lyeth, very like vnto the common Maple growing wilde abroad, but many more together, and larger.

The

### The Vſe of the Sycomore tree.

It is altogether planted for ſhady walkes, and hath no other vſe with vs that I know.

## 19. *Nux Veſicaria.* The bladder Nut.

THis tree groweth not very high, but is of a meane ſtature, when it is preſerued and pruined to grow vpright, or elſe it ſhooteth forth many twigges from the rootes, and ſo is fit to plant in a hedge rowe, as it is vſed in ſome places: the body and armes are couered with a whitiſh greene barke: the branches and leaues on them are like vnto the Elder, hauing three or fiue leaues ſet one againſt another, with one of them at the end, each whereof is nicked or dented about the edges: the flowers are ſweete and white, many growing together on a long ſtalke, hanging downeward, in forme reſembling a ſmall Daffodill, hauing a ſmall round cup in the middle, and leaues about it: after which come the fruit, incloſed in ruſſetiſh greene bladders, containing one or two browniſh nuts, leſſer then Haſell nuts, whoſe outer ſhell is not hard and woody, like the ſhell of a nut, but tough, and hard withall, not eaſie to breake, within which is a greene kernell, ſweetiſh at the firſt, but lothſome afterwards, ready to procure caſting, and yet liked of ſome people, who can well endure to eate them.

### The Vſe of the Bladder Nut.

The greateſt vſe that I know the tree or his fruit is put vnto, is, that it is receiued into an Orchard, either for the rarity of the kinde, being ſuffered to grow into a tree, or (as I ſaid before) to make an hedge, being let grow into ſuckers.

Some Quackſaluers haue vſed theſe nuts as a medicine of rare vertue for the ſtone, but what good they haue done, I neuer yet could learne.

## 20. *Rhus Myrtifolia.* The Mirtle leafed Sumach.

THis lowe ſhrubbe groweth ſeldome to the height of a man, hauing many ſlender branches, and long winged leaues ſet thereon, euery one whereof is of the bigneſſe of the broad or large Mirtle leafe, and ſet by couples all the length of the ribbe, running through the middle of them. It beareth diuers flowers at the tops of the branches, made of many purple threads, which turne into ſmall blacke berries, wherein are contained ſmall, white, and rough ſeed, ſomewhat like vnto Grape kernels or ſtones. This vſeth to dye down to the ground in my Garden euery Winter, and riſe vp again euery Spring, whether the nature thereof were ſo, or the coldneſſe of our climate the cauſe therof, I am not well aſſured. It is alſo rare, and to be ſeen but with a few.

### The Vſe of this Sumach.

It is vſed to thicken or tanne leather or hides, in the ſame manner that the ordinary Sumach doth; as alſo to ſtay fluxes both in men and women.

## 21. *Rhus Virginiana.* The Virginia Sumach, or Buckes horne tree of Virginia.

THis ſtrange tree becommeth in ſome places to bee of a reaſonable height and bigneſſe, the wood whereof is white, ſoft, and pithy in the middle, like vnto an Elder, couered with a darke coloured barke, ſomewhat ſmooth: the young branches that are of the laſt yeares growth are ſomewhat reddiſh or browne, very ſoft and

and ſmooth in handling, and ſo like vnto the Veluet head of a Deere, that if one were cut off from the tree, and ſhewed by it ſelfe, it might ſoone deceiue a right good Woodman, and as they grow ſeeme moſt like thereunto, yeelding a yellowiſh milke when it is broken, which in a ſmall time becommeth thicke like a gumme: the leaues grow without order on the branches, but are themſelues ſet in a ſeemly order on each ſide of a middle ribbe, ſeuen, nine, ten, or more on a ſide, and one at the end, each whereof are ſomewhat broad and long, of a darke greene colour on the vpperſide, and paler greene vnderneath, finely ſnipped or toothed round about the edges: at the ends of the branches come forth long and thicke browne tufts, very ſoft, and as it were woolly in handling, made all of ſhort threads or thrums; from among which appeare many ſmall flowers, much more red or crimſon then the tufts, which turne into a very ſmall ſeede: the roote ſhooteth forth young ſuckers farre away, and round about, whereby it is mightily encreaſed.

### The Vſe of this Sumach.

It is onely kept as a rarity and ornament to a Garden or Orchard, no bodie, that I can heare of, hauing made any tryall of the Phyſicall properties.

---

### 22. *Vitis, ſeu potius Hedera Virginenſis.* The Virginia Vine, or rather Iuie.

THis ſlender, but tall climing Virginia Vine (as it was firſt called; but Iuie, as it doth better reſemble) riſeth out of the ground with diuers ſtems, none much bigger then a mans thumbe, many leſſe; from whence ſhoote forth many long weake branches, not able to ſtand vpright, vnleſſe they be ſuſtained: yet planted neere vnto a wall or pale, the branches at ſeuerall diſtances of the leaues will ſhoote forth ſmall ſhort tendrels, not twining themſelues about any thing, but ending into foure, fiue, or ſix, or more ſmall ſhort and ſomewhat broad clawes, which will faſten like a hand with fingers ſo cloſe thereunto, that it will bring part of the wall, morter, or board away with it, if it be pulled from it, and thereby ſtay it ſelfe, to climbe vp to the toppe of the higheſt chimney of a houſe, being planted thereat: the leaues are crumpled, or rather folded together at the firſt comming forth, and very red, which after growing forth, are very faire, large, and greene, diuided into foure, fiue, ſix, or ſeuen leaues, ſtanding together vpon a ſmall foote-ſtalke, ſet without order on the branches, at the ends whereof, as alſo at other places ſometime, come forth diuers ſhort tufts of buds for flowers; but we could neuer ſee them open themſelues, to ſhew what manner of flower it would be, or what fruit would follow in our country: the roote ſpreadeth here and there, and not very deepe.

### The Vſe of this Virginian.

We know of no other vſe, but to furniſh a Garden, and to encreaſe the number of rarities.

And thus haue I finiſhed this worke, and furniſhed it with whatſoeuer Art and Nature concurring, could effect to bring delight to thoſe that liue in our Climate, and take pleaſure in ſuch things; which how well or ill done, I muſt abide euery ones cenſure: the iudicious and courteous I onely reſpect, let Momus bite his lips, and eate his heart; and ſo Farewell.

*FINIS.*

# Index omnium ſtirpium

## quæ in hoc opere continentur.


A

*Abies* page 600
*Abrotanum fœminum, ſiue Santolina* 449
*Acanthus Aculeatus* 330
*Acanthus ſatiuus* ibid.
*Acer maius ſiue Sycomorus* 610
*Acetoſa* 486
*Aconitum baccifer um*, i. e. *Chriſtophoriana*
*Aconitum flore albido* 214
*Aconitum hyemale* ibid.
*Aconitum luteum Ponticum* ibid.
*Aconitum ſalutiferum* 216
*Admirabilis peruana* 364
*Æthiopis* 385
*Agluophotis*, i.e. *Paonia* 603
*Alaternus* 148
*Albucum* 369
*Alcea Ægyptia ſiue Bamia* 368
*Alcea Americana* ib.
*Alcea fruticoſa pentaphyllea* ib.
*Alcea peregrina ſiue veſicaria*
*Aliſma Dodonæi*, i.e. *Saponaria*
*Aliſma Dioſcoridis Fab. Columna*, i. e. *Auricula Vrſi*
*Aliſma ſyluarum*, i.e. *Paralyſis* 613
*Allium* 369
*Althæa frutex*
*Amaracus*, i.e. *Maiorana.*
*Amarella*, i.e. *Matricaria*
*Amaranthus panniculis ſparſis* 371
*Amaranthus purpureus* ib.
*Amaranthus tricolor* ib.
*Amaranthus luteus*, i.e. *Heliocryſum*
*Ambreboi*, i.e. *Cyanus Orientalis*
*Amellus Virgilij*, i.e. *Aſter Atticus Italorum*
*Anemum Plinij*, i.e. *Pſeudocapſicum Dodonei*
*Amygdalus* 583
*Anagyris altera*, i.e. *Laburnum.*
*Anchuſa* 251
*Anemone eiuſque ſpecies* 199 ad 214
*Anethum* 494.
*Angelica* 529
*Anthemis flore luteo* 294
*Anthemis Leucanthemis*, i.e. *Chamemelum*
*Anthericos* 148
*Anthora* 494
*Antimelum*, i.e. *Mandragoras*
*Antirrhinum* 269
*Apium* 491
*Apocynum Syriacum* 444
*Apocynum Virginianum* 445
*Aquilegia* 271
*Arbor Alpina Plinij*, i.e. *Laburnum*
*Arbor Iudæ* 437
*Arbor Vitæ* 438
*Arbutus* 603
*Argyrocome*, i.e. *Gnaphalium Americanum*
*Armerius* 319
*Armoraria pratenſis* 256
*Armoraria altera*, i.e. *Muſcipula Lobelij*
*Arthanita*, i.e. *Cyclamen*
*Arthritica*, i.e. *Paralyſis*
*Arundo Indica, ſiue Canna Indica* 376
*Aſarum* 532
*Aſparagus* 503
*Aſphodelus bulboſus albus* 138
*Aſphodelus bulboſus Galeni* ibid.
*Aſphodelus hyacinthinus*, i.e. *bulboſus*
*Aſphodelus maior albus* 146
*Aſphodelus minor luteus*, i.e. *Haſtula regia*
*Aſter Atticus Italorum* 299
*Aſter Peruanus Columnæ*, i.e. *Battatas de Canada*
*Attamuſco*, i.e. *Narciſſus Virginianus*
*Atriplex*, i.e. *Olus aureum* 488
*Auellanæ, & Byzantinæ*
*Aurelia*, i.e. *Chryſocome*
*Auricula muris maior*, i.e. *Pulmonaria Gallorum*
*Auricula Vrſi eiuſque ſpecies* 232

B

*Balauſtium* 439
*Balſamina fœmina & Balſamella* 278
*Balſamita mas & fœmina* 482
*Balſamum alpinum*, i.e. *Ledum alpinum* 424
*Bamia*, i.e. *Alcea Ægyptia*
*Baptiſecula*, i.e. *Cyanus*
*Barba hirci*, i.e. *Tragopogon*
*Battatas Hiſpanorum, Virginianum, & Canadenſe* 518
*Behen rubrum*, i. *Valeriana rubra Dodonæi*

Bellis cærulea sive Globularia 321
Bellis maior flore pleno 322
Bellis minor flore pleno eiusque species ibid.
Belvidere Italorum, i. Scoparia, sive Linaria magna 268
Ben rubrum Monspelliense, i. Muscipula Lobelij
Berberis 561
Beta 488
Binizade & Binizante, i. Anemone tennifolia
Blattaria 383
Blito di tre colori 372
Blitum 488
Bolbonach, i. Viola lunaris & latifolia 265
Borrago, Borrago semper virens 249
Botanaria, i. Globularia
Branca vrsina, i. Acanthus sativus
Brassica eiusque species 503
Bubonium sive Inguinalis, i. After Atticus Italorum
Bulbus agrestis, i. Colchicum
Buccinum Romanorum, i. Delphinium
Bulbus Eriophorus 124
Bulbus esculentus Lacunæ, i. Ornithogalum luteum 140
Bulbus Leucanthemos i. Ornithogalum album
Bulbus unifolius 140
Bulbus vomitorius Matthioli, i. Muscari
Buglossum 249
Buglossum Hispanicum, i. Anchusa
Buglossum luteum 486
Bunias dulcis, i. Napus 509
Buphthalmum 293
Buphthalmum maius i. Helleborus niger serulaceus
Buxus arbor. Buxus humilis, & Buxus verficoribus solijs 606


## C


CAcalia Loniceri. i. Lilium Convallium
Calcaris flos, i. Delphinium
Calceolus Mariæ 367
Calendula maxima & simplex 296. 298
Callionymus Gesneri, i. Lilium convallium
Caltha, i. Calendula
Caltha Africana, i. Flos Africanus
Caltha palustris flore pleno 224
Camomilla vulgaris, & flore pleno 290
Campanula maior pyramidalis 354
Campana lazura, i. Convolvulus cæruleus maior 359
Campanula persicifolia alba & cærulea 351
Canicida, i. Aconitum luteum Ponticum
Canis cerebrum, i. Antirrhinum
Cannacorus, i. Canna Indica
Canna Indica flore luteo punctato 376
Canna Indica flore rubro ibid.
Cantabrica Plinij, i. Caryophyllus
Capnos fabaceæ radice, i. Radix cana minor 275
Caprifolium perfoliatum sive Italicum 405
Cardamine flore pleno, & trifolia 389
Cardnus benedictus 530
Carduus Eriocephalus, i. tomentosus 332
Carduus mollis 330
Carlina humilis 332
Carthamus sive Cnicus sativus 329
Carum 515
Caryophylli maiores & maximi 316
Caryophylli sylvestres 314
Caryophyllus marinus & mediterraneus 317
Caryophyllus Indicus, i. Flos Africanus
Cassana, i. Iucca 434
Cavala lale, i. Tulipæ præcoces
Caffalale, i. Tulipæ mediæ
Caucafon, i. Moly Indicum
Caulis vulgaris, Crispa, Subandica 504
Caulis florida ibid
Caulo rapum ibid.
Cedrus Lycia 436
Celastrus 603
Cepa alba, rubra &c. 510
Cerasa Indiana 432
Cerasus flore pleno 402
Cerasorum diuersitas 571
Cerasus Trapezuntina, i. Laurocerasus
Cercis 437
Cerefolium maius & vulgare 494
Cervicaria, i. Trachelium
Chamæcistus Frisicus 424
Chamæcyparissus, i. Santolina
Chamædaphne 498
Chamedrys 456
Chamæiris angustifolia 187
Chamæiris latifolia 186
Chamælea Alpina 397
Chamælea Germanica, i. Mesereon 397
Chamælea tricoccos ibid
Chamæleo, vide Carlina
Chamæmalus, i. Malus Paradisens
Chamæmelum, i. Camomilla
Chamænerium flore Delphinij 270
Chamæpauce Cordi, i. Ledum Silesiacum
Chamærhododendros Chamalaæfolio Lobelij, i. Ledum Alpidum
Charantia fœmina, i. Balsamina fœmina
Cheiri sive Keiri, i. Leucoium luteum 256
Chondrilla aurea, i. Pilosella maior 300
Chrysanthemum odoratum, i. Chamæmelum nudum
Chrysanthemum Peruvianum, i. Flos Solis
Chrysanthemum Creticum 295
Chrysoceme sive Stœchas citrina 374
Cichorium 495
Cinara alba, rubra, moschata &c. 518
Circæa. i. Mandragoras
Cistus annuus 422
Cistus mas 421
Cistus famina 422
Cistus Ledon ibid
Claues sancti Petri, i. Paralysis
Clematis Daphnoides, i. Vinca pernica 391
Clematis altera sive vrens flore albo ibid
Clematis peregrina flore rubro 392
Clematis peregrina flore purpureo ibid
Clematis peregrina flore purpureo pleno ibid
Clematis peregrina flore carneo pleno 393
Clematis cærulea Pannonica ibid
Clematis surrecta sive Flammula Iouis ibid
Clematis flore albo pleno, vel surrecta flore duplici 392
Clematis Virginiana, i. Maracos 393
Clymenum Matthioli, i. Lathyrus latifolius sive Pisum perenne
Cnecrum Matthioli 397

*Cneorum nigrum & album Theophrasti* 426
*Cnicus siue Carthamus sativus* 329
*Colchicum Anglicum Byzantinum &c.* 154 *&c.*
*Colchicum vernum* 158
*Colytea Theophrasti* 438
*Colutea vulgaris, & Scorpioides* 440
*Coma aurea siue Heliochrysum*
*Condrilla aurea*, i. *Pilosella maior*
*Consolina minor*, i. *Bellis minor vulgaris*
*Consolida regalis*, i. *Delphinium*
*Convolvulus Americanus* 358
*Convolvulus caerulens maior* 357
*Convolvulus cæruleus minor* 358
*Convolvulus purpureus maior* ibid
*Convolvulus purpureus minor spicafolijs* 359
*Corchorus Dalechampij*, i. *Pilosella maior*
*Cornus mas fructu albo, rubro* 570
*Corona Imperialis* 28
*Cortusa Matthioli* 240
*Corydalis* 276
*Cosmosandalos*, i. *Calceolus Mariæ*
*Costus hortorum maior & minor* 482
*Cotonea malus* 589
*Cotyledon altera minor, & flore rubro* 232
*Crocus vernus albus, purpureus &c.* 160 ad 170
*Crocus Hispanicus*, i. *Carthami flores*
*Cucumis hortensis* 524
*Cuminum sylvestre alterum Dioscoridis Matthiolo*, i. *Delphinium*
*Cupressus* 601
*Cyanus Batitus supinus* 327
*Cyanus floridus Turcicus* ibid.
*Cyanus minor variorum colorum* 326
*Cyanus Orientalis*, i. *floridus Turcicus*
*Cyclamen Antiochenum* 196
*Cyclamen autumnale hederæfolio* ibid
*Cyclamen vernum. Veronense &c.* 195
*Cyclaminus idem*
*Cydonia malus* 589
*Cyprus Plinij*, i. *Ligustrum Orientale* 414. 445
*Cytisus vulgaris. Maranthæ &c* 440

D

*Dactylus Trapezuntinus*, i. *Lauroceraſus*
*Daphnoides*, i. *Laureola* 398
*Datura Turcarum*, i. *Stramonium*
*Delphinium* 276
*Delphinium buccinum*, i. *Chamænerium*
*Delphinium Hispanicum parvum* 277
*Dens caninus* 193
*Dentali*, i. *Dens caninus*
*Devebohini*, i. *Narcissus medio purpureus*
*Dictamus albus & Diptamus albus*, i. *Fraxinella* 380
*Digitalis*
*Dioscanthos Theophrasti Dalechampio*, i *Aquilegia*
*Dioscanthos, siue Iouis flos*, i. *Caryophyllus*
*Dipcadi*, i. *Muscari*
*Dracoherba, seu Tarchon* 500
*Draba siue Arabis Dodonæi*, i. *Thlaspi Creticum*
*Dracunculus maior* 529

E.

*Elleborine flore albo &c.* 347
*Elleborus albus vulgaris & præcox* 346
*Elleborus niger, vel Helleborus niger* 344
*Endiuia* 495
*Ephemerum lethale*, i *Colchicum*
*Ephemerum non lethale*, i. *Lilium conuallium*
*Ephemerum Virginianum* 152
*Epimedium* 283
*Eranthemum*, i. *Flos Adonidis*
*Eriphium Galeni*, i. *Radix cana*
*Eruca sativa* 502
*Eryngium Montanum siue Pannonicum* 330

F

*Faba vulgaris* 521
*Fabago*, i. *Arbor Inda*
*Ficus Arbor & humilis* 566
*Ficus Indica* 432
*Flamma Theophrasti* i. *Amaranthus minor purpureus*
*Flammula Iouis*, i. *Clematis surrecta*
*Flos Adonis* 293
*Flos Africanus* 303
*Flos Amoris*, i *Amaranthus*
*Flos Cancri*, i. *Canna Indica*
*Flos Cardinalis*, i. *Trachelium Americanum* 356
*Flos Caryophylleus, siue Caryophyllus* 314
*Flos Constantinopolitanus siue Lychnis Chalcedonica*
*Flos maximus*, i. *Flos Solis*
*Flos Cuculi*, i. *Cardamine*
*Flos Cuculi*, i. *Lychnis sylvestris plumaria*
*Flos frumenti*, i. *Cyanus minor*
*Flos noctis*, i. *Convolvulus maior purpureus*
*Flos Diuæ Katharinæ*, i. *Nigella*
*Flos Meleagridis*, i. *Fritillaria*
*Flos Regius*, i. *Delphinium*
*Flos Indicus vel Tunetensis*, i. *Africanus*
*Flos Solis Farnesianus Columnæ*, i. *Battatas de Canada*
*Flos Solis* 295
*Faniculum* 492
*Fragaria* 526
*Fraxinella* 333
*Fritillaria* 44
*Frutex Coronarius Clusij*, i. *Syringa alba* 414
*Fucus herba*, i. *Anchusa*

G.

*Gallipo lale*, i. *Anemone tenuifolia*
*Galeopsis Pannonica siue Lamium Pannonicum* 385
*Gelosia Tragi*, i. *Amaranthus tricolor*
*Gelseminum*, i. *Iasminum*
*Genista Hispanica siue Spartum Hispanicum* 442
*Gentiana maior* 350
*Gentiana Asclepiadis folio* ibid
*Gentiana cruciata* ibid
*Gentianella verna* 352
*Gentianella autumnalis siue Pneumonanthe* ibid
*Geranium tuberosum vel bulbosum* 228
*Geum Alpinum Lobelij*, i. *Sanicula guttata*
*Geranium batrachoides, fuscum &c.* 229
*Gith*, i. *Nigella*
*Giul catamer lale*, i. *Anemone latifolia multiplex*
*Gladiolus Byzantinus, Italicus &c.* 189
*Globularia cærulea* 322
*Globularia lutea montana* 323
*Glycyrrhiza siue Liqueritia* 533

*Gnaphalium Americanum* 374
*Gnaphalium Montanum, siue Pes Cati vel Pilosella minor Montana* 375
*Gnaphalium Roseum* ibid.
*Gramen striatum vel pictum* 458
*Gramen marinum*, i. *Caryophyllus marinus minor*
*Gramen plumarium vel plumosum* 458
*Grossularia siue Vua crispa vulgaris, baccis rubris, caeruleis, aculeatis &c.* 560
*Guaiacana siue Guaiacum Patauinum idem est cum Pishamin Virginianorum, Loti species*

H.

*Hastula regia*, i. *Asphodelus luteus minor*
*Hedera Virginiana* 612
*Hedysarum clypeatum* 339
*Heliotropium Indicum Pelleterij*, i *Battatas de Canada*
*Heliocrysum* 374
*Helleboraster siue Pseudohelleborus* 345
*Helleborus albus & vernus praecox* 346
*Helleborus niger verus* 344
*Helleborus niger ferulaceus*, i. *Buphthalmum maius* 293
*Helleborine* 347
*Hemerocallis*, i. *Martagon*
*Hemerocallis Valentina Clusij*, i. *Pseudonarcissus marinus, vel Pancratium vulgo*
*Hepatica nobilis siue trifolia* 225
*Herba clauellata*, i. *Viola tricolor*
*Herba sancti Petri*, i. *Paralysis*
*Herba Sanctae Catharinae*, i. *Balsamina famina*
*Herba Margarita*, i. *Bellis minor*
*Herba Regina*, i. *Tabacco*
*Herba Tunica*, i. *Caryophyllus & Armerius*
*Herba Trinitatis*, i. *Hepatica trifolia*
*Hermodactylus Matthioli* 188
*Gesneri* 194
*Hermodactylum Colchicum* 160
*Hesperis* 262
*Hippolapathum*, i. *Patientia siue Lapathum sativum, & Rhabarbarum Monachorum* 483
*Hippolapathum rotundifolium* 484
*Hipposelinum siue Olus atrum* 490
*Hirculus Friscus*, i. *Chamacistus Fricisca*
*Horminum sativum* 478
*Hyacinthus & eius genera* 111 ad 133
*Hyacynthus Poetarum*, i. *Iris bulbosa Anglicana*
*Hyssopus foliis aureis* 455
*Hyssopus vulgaris* 476
*Hypecoum Matthioli*, i. *Alcea vesicaria*

I.

*Iacea Batica* 328
*Iacea marina Batica* ib.
*Iasminum Americanum* 359
*Iasminum Arabicum* 410
*Iasminum album* 406
*Iasminum Catalonicum* ibid
*Idaeus dactylus*, i. *Paeonia*
*Ilex arbor* 600
*Intubum*, i. *Endiuia & Cichorium* 495
*Iouis flos*, i. *Caryophyllus*
*Iphium Theophrasti*, i. *Caryophyllus*
*Iris bulbosa & eius varietas* 171 ad 179
*Iris Chalcedonica* 179
*Iris Dalmatica Damascena &c.* 180 &c.
*Iris Persica* 172
*Iris Tripolitana* 182
*Iris tuberosa* 188
*Iucca* 434
*Ixine Theophrasti*, i. *Carlina*

K

*Keiri, siue Leucoium luteum* 256
*Keiri albo flore* 257

L

*Laburnum* 438
*Lactuca agnina. Capitata &c.* 498
*Lamium Pannonicum*, i. *Galeopsis Pannonica* 385
*Lapathum sanguineum* 484
*Larix* 608
*Lathyrus latifolius*, i. *Pisum perenne* 338
*Lathyrus siliquis orobi* ib.
*Lauendula mas & famina* 447
*Lauendula syluestris*, i. *Stachas*
*Laurus* 598
*Laurus regia*, i. *Lauroceransus* 401
*Laurus Rosea*, i. *Oleander*
*Laurus Tinus* 400
*Laurea Cerasus* 599
*Lauro cerasus* 401
*Ledum Alpinum. Silesiacum* 424
*Ledum id est Cistus Ledon*
*Leimonia Theophrasti Clusio*, i. *Anemone syluestris*
*Leontostomium*, i. *Aquilegia*
*Leucoium bulbosum* 109
*Leucoium hortense simplex & multiplex* 258
*Leucoium luteum multiplex* 257
*Leucoium marinum Syriacum* 260
*Leucoium melancholicum* ibid
*Leuconacissolirion*, i. *Leucoium bulbosum*
*Libanotis Coronaria*, i. *Rosmarinus*
*Ligustrum* 445
*Ligustrum Orientale*, i. *Cyprus Plinij* 410
*Lilac*, i. *Syringa caerulea* 407
*Lilac flore argenteo* 408
*Liliago*, i. *Phalangium: vel alijs Lilium non bulbosum siue Liliasphodelus*
*Liliasphodelus* 148
*Lilionarcissus*, i. *Tulipa*
*Lilium Alexandrinum*, i. *Ornithogalum Arabicum* 135
*Lilium album* 40
*Lilium aureum, rubrum &c* 39
*Lilium conuallium* 349
*Lilium Macedonicum* 36
*Lilium Montanum* 33
*Lilium non bulbosum*, i. *Liliasphodelus*
*Lilium Persicum vel Susianum* 28
*Lilium syluestre*, i. *Montanum*
*Lilium variegatum*, i. *Fritillaria*
*Limonium peregrinum Rauwolfij* 250
*Linaria magna siue Belvidere Italorum* 268
*Liquerutia* 583
*Linaria caerulea purpurea odorata* 66
*Lingua Bouis vel Buglossum luteum* 486
*Linum syluestre album, luteum* 266

*Lotus Africana Dalechampij*, i. *Laurocerasus*
*Lotus Arbor* 568
*Lotus tetragonolobus siue siliquosus flore rubello*, i. *Pisum quadratum*
*Lunaria Arthritica*, i. *Peralysis, etiamque & Auricula Vrsi*
*Lunaria Graca* { *maior* } i. *Bolbonach, seu Viola* { *odorata* } *latifolia* 265
*Lunaria carulea*, i. *Soldanella Alpina, siue Montana* 234
*Lupinus* 335
*Lychnis Chalcedonica flore simplici, & flore pleno* 253
*Lychnis Coronaria* 252
*Lychnis plumaria sylueſtris multiplex* 253
*Lychnis ſylueſtris flore albo pleno, & flore rubro pleno* 254
*Lycoperſicum Galeni*, i. *Flos Africanus, pomum Amoris Anguillare* 511

Lepidium.

M.

*Maiorana aurea* 446
*Maiorana tenuifolia* 452
*Maiorana vulgaris* ibid
*Maiorana latifolia vulgaris* 474
*Mala Æthyopica* 379
*Mala Arantia* 584
*Mala Armeniaca siue præcocia* 579
*Malus Cotonea vel Cydonia* 589
*Malus Granata seu Punica* 428
*Malorum Perſicorum diuerſitas* 580
*Malorum varia genera* 586
*Malus Punica ſatiua* 428
*Malus Punica ſylueſtris*, i. *Balauſtium* 495
*Maluacriſpa* 366
*Malua Hiſpanica flore carneo amplo* 370
*Malua horaria*, i. *Alcea peregrina* 369
*Malua hortenſis ſimplex & multiplex*
*Malua Roſea*, i. *hortenſis* 377
*Mandragoras mas & fœmina* 393
*Maracoc siue Clematis Virginiana*
*Marguerites & Margueritons*, i. *Bellis minor multiplex* 33
*Martagon album, flore carneo* 34
*Martagon Chymiſtarum*, i. *Lilium aureum ſiue rubrum* 35
*Martagon Byzantinum ſeu Conſtantinopolitanũ* 33
*Martagon Pannonicum flore ſpadiceo* 36
*Martagon Imperiale* 34
*Martagon Pomponeum*
*Martagon Phœniceo flore*
*Martagon rariſſimum*, i. *Narciſſus tertius Matthioli* 289
*Matricaria flore pleno* 338. &c.
*Medica Cochleata. Spinoſa*
*Medium Dioſcoridis*, i. *Viola Mariana*
*Melampodium*, i. *Helleborus niger* 479
*Melanthium*, i. *Nigella* 525
*Meliſſa* 480
*Melo Moſcatus &c.* 568
*Mentha*
*Meſpilus Aronia. Vulgaris*
*Mezereon*, i. *Chamælea Germanica* 394
*Mirabilia Peruviana* 364.
*Moly alterum genus Pliny Dodonæo*, i. *Phalangium* 141 ad 146
*Moly varia genera*
*Morion*, i. *Mandragoras*
*Morus vulgaris, alba, Virginiana* 599
*Muſchoromi*, i. *Muſcari*
*Moſchatella*, i. *Heſperis*
*Muſcipula Lobelij*, i. *Benrubrum Monſp.* 254
*Myrobalanus prunus* 577
*Myrrhis*, i. *Cerefolium maius*
*Myrtus maior & minor* 427

Marum. 452.

N.

*Napellus* 215
*Napellus Moiſis*, i. *Anthora* 216
*Napus maior & minor* 509
*Narciſſus Caperonius*, i. *Fritillaria*
*Narciſſorum varia genera* 67 ad 108
*Narciſſus marinus*, i. *tertius Matthioli*
*Narciſſus Iacobæus flore rubro* 169
*Narciſſus Matthioli*, i. *Ornithogalum Neapolitanum*
*Narciſſus Trapezunticus* 30
*Narciſſus Virgineus* 86 & 91
*Nardus Italica*, i. *Lauendula*
*Nardus montana* 386
*Naſturtium Indicum* 280
*Naſturtium hortenſe* 500
*Nepeta* 479
*Nerium*, i. *Oleander*
*Nerium Alpinum*, i. *Ledum Alpinum*
*Nicotiana*, i. *Tabacco*
*Nigella flore albo duplici* 287
*Nigella flore caruleo multiplici* ibid
*Nigella Hiſpanica* ibid
*Nil Auicennæ*, i. *Convolvulus cæruleus maior*
*Nozelha*, i. *Crocus Cluſio. Syſirinchium Boelio* 171
*Nuciperſica & eius varietas* 583
*Nux Auellana, Byzantina* 562
*Nux Iuglans* 595
*Nux Metel*, i. *Stramonium maius*
*Nux veſicaria ſeu Staphylodendron* 611

O.

*Ocellus Barbaricus, & Damaſcenus*, i. *Caryophyllus*
*Ocimaſtrum Valerianthon*, i. *Valeriana rubra Dodonæi*
*Ocimoides ſemper virens* 254
*Ocimum citratum, Indicum* 450
*Odontitis*, i. *Lychnis plumaria*
*Oenanthe Myconi*, i. *Ranunculus thalictri folio minor* 218

*Olus album*, i. *Lactuca agnina*
*Olus atrum*, i. *Hippoſelinum*
*Olus aureum*, i. *Atriplex* 488.
*Olus hiſpanicum*, i. *Spinachia*
*Opuntia*, i. *Lychen marinus* 433
*Opuntia ſeu Ficus Indica* ibid
*Orchis Melitias ſeu Apifera* 192
*Orchis hermaphroditica &c.* ibid.
*Ornithogalum Æthiopicum* 138
*Ornithogalum Arabicum* 134
*Ornithogalum luteum* 140
*Ornithogalum Neapolitanum* 138
*Ornithogalum Pannonicum* 136 & 138
*Orobus Venetus* 338

*Orontium*, i. *Antirrhinum*
*Os Leonis*, i. *Antirrhinum*
*Ostrys Theophrasti* 410
*Osyris*, i. *Scoparia vel Linaria magna*
*Oxalis siue Acetosa* 487
*Oxyacantha siue Berberis & sine acinis* 561

P.

*PAdus Theophrasti*, i. *Cerisier blanc Gallorum siue Cerasus racemosus*
*Paeonia mas & famina simplex & multiplex* 341 &c.
*Paliurus* 607
*Palma Christi*, i. *Ricinus*
*Palma Christi*, i. *Orchis siue Satyrium Basilicum*
*Pancratium*, i. *Scilla rubra, & non Pseudonarcissus marinus, vide pag.* 108 & 153
*Panis porcinus*, i. *Cyclamen.*
*Papaver sativum flore pleno* 286
*Papaver sylvestre flore pleno* ibid.
*Pappas siue Battatas Hispanorum Canadense Virginianum* 516
*Paralysis diuersarum specierum* 242 &c.
*Paralytica alpina maior & minor*, i. *Auricula Vrsi & Paralysis minor*
*Parthenium*, i. *Matricaria*
*Parthenium Galeni*, i. *Amaracus*
*Pastinaca latifolia* 506
*Pastinaca tenuifolia* 508
*Pedna Penorum*, i. *Flos Africanus*
*Pennachio Persiano*, i. *Lilium Persicum*
*Pepo* 526
*Peruinca siue Vinca peruinca* 391
*Periclymenum perfoliatum* 404
*Periclymenum rectum* 415
*Periploca Virginiana* 444
*Pes cati*, i. *Gnaphalium Montanum*
*Petroselinum* 491
*Petum & Picielt*, i. *Tabacco.*
*Perebecenuc*, i. *Tabacco*
*Phalangium Allobrogicum* 150
*Phalangium ephemerum Virginianum* 152
*Phalangium Italicum maius* 150
*Phalangium ramosum & non ramosum* ibid.
*Phaseolus vulgaris* 521
*Phillyrea* 445
*Phlomitis*, i. *Æthiopis*
*Phlox siue flamma Theophrasti*, i. *Viola flammea siue tricolor*
*Piloselle maior siue Chondrilla aurea* 300
*Piloselle minor montana*, i. *Gnophalium montanum* 375
*Pimpinella siue Sanguisorba* 483
*Pinus* 599
*Piper montanum*, i. *Chamalea frutex vel semen*
*Pisamin Virginianorum*, i. *Guaiacum Pataninum*
*Pisum perenne*, *Lathyrus latifolius* 340
*Pisum quadratum rubrum* 338
*Pisum vulgare, roseum, maculatum &c.* 522
*Planta Cardinalis*, i. *Trachelium Americanum* 356
*Planta maxima*, i, *Flos Solis*
*Plantago rosea* 352
*Pneumonanthe*, i. *Gentiana autumnalis* 352
*Poma amoris maiora & minora* 379
*Pomorum varietas* 586
*Pomum spinosum*, i. *Stramonium* 362
*Porrum* 512
*Portulaca* 499
*Pothos Theophrasti*, i. *Aquilegia*
*Primula veris simplex & multiplex* 242. &c.
*Prunorum magna varietas* 575
*Pseudocapsicum*, i. *Amomum Plinij* 431
*Pseudohelleborus Matthioli*, i. *Helleborus niger fernlaceus*
*Pseudohermodactylus Matthioli*, i. *Dens Caninus*
*Pseudonarcissus Anglicus, Germanicus, Hispanicus, &c.* 99. &c.
*Pseudolotus Matthioli*, i. *Laurocerasus, est & Guaiacum Patavinum & Pisamin Virginianorum*
*Pseudorhabarbarum est Rhabarbarum Monachorum* 274
*Psidium*, i. *Cortex Granatorum*
*Ptarmica flore pleno* 288
*Pulegium* 477
*Pulmonaria Tragi & Gallorum*, i. *Piloselle maior* 300
*Pulmonaria maculosa & non maculosa* 248
*Pulsatilla* 200
*Pyracantha* 604
*Pyramidalis Lutetiana*, i. *Campanula maior siue lactescens* 354
*Pyretrum officinarum* 292
*Pyretrum sylvestre* 288
*Pyrus & eius varietas* 590

piperitis. 

Q.

*QVamoclit Indorum*, i. *Convolvulus Americanus* 358
*Quinúa Indorum*, i. *Amaranthus maior siue paniculis sparsis*

R.

*RAdix Caua maior, & minor* 275
*Ranunculus Anglicus, Asiaticus, Creticus &c.* 276 ad 223
*Ranunculus monophyllos*, i. *Aconitum hyemale*
*Ranunculus nemorum & sylvarum*, i. *Anemones sylvestres*
*Raphanus vulgaris & nigra radice* 509
*Rapum hortense, luteum, rubrum* ibid.
*Rapunculus hortensis* 514
*Rhabarbarum Monachorum & Pseudorhabarbarum* 274
*Rhabarbarum & Rhaponticum verum* 483
*Rhododendron*, i. *Oleander*
*Rhus Virginiana* 611
*Rhus Myrtifolia* ibid.
*Ribes fructu albo, nigro, rubro* 558
*Rosa Alpina*, i. *Ledum alpidum* 424
*Rosa Iunonis*, i. *Lilium album*
*Rosa satuina*, i. *Paeonia*
*Rosa montana*, i. *Alpina*
*Rosa vltramarina*, i. *Malua rosea*
*Rosa Anglica, Cinamomea, Damascena, &c.* 412 ad 425
*Rosmarinum vulgare* 425
*Rosmarinum aureum* ibid
*Rosmarinum latifolium* 426
*Rosmarinum sylvestre Matthioli*, i. *Ledum Alpinum*
*Rubus Idæus* 557
*Rubus Idæus non spinosus* ibid

*Ruta hortensis* 530
*Ruta palustris vel pratensis*, i. *Thalictrum*

S

*Sabdarifa* i. *Alcea Americana*
*Sabina* 607
*Saliuaris*, i. *Pyretrum*
*Saluia maior & minor* 478
*Saluia variegata* 446
*Sambach Arabum*, i. *Syringa Arabica flore duplici*
*Sambacus Rosea* 411
*Sampsuchum*, i. *Maiorana*
*Sana sancta & Sancta herba*, i. *Tabacco*
*Sandalida Cretica*. i. *Pisum quadratum*
*Sanicula Alpina vel montana*, i. *Cortusa Matthioli*
*Sanicula trifolia*, i. *Cardamine trifolia*
*Sanicula guttata* 231
*Sanguisorba vel Pimpinella* 483
*Sanguis Herculis*, i. *Elleborus albus*
*Santolina*, i. *Abrotanum fœmina* 449
*Saponaria flore duplici* 352
*Saponaria altera*. i. *Trachelium minus, & Valeriana rubra Dodon.*
*Sarahug*, i. *Hyacinthus Lillifolius*
*Satureia vel Thymbra* 476
*Satyrium Erythronium & Triphyllum Dioscoridis* 66
i. *Tulipa*
*Satyrium Orchidis species* 192
*Scabiosa rubra Austriaca* 324
*Scabiosa rubra Indica* ibid
*Scarlatea*, i. *Armerius*
*Scoparia*, i. *Linaria magna* 268
*Scorpioides maius & minus* 340
*Scorsonera* 301
*Scylla alba, rubra* 133
*Sedum serratum* 232
*Segetalis*, i. *Gladiolus*
*Selinum dulce* 491
*Serincade*, i. *Narcissus medio purpureus*
*Serincade catamer lale*, i. *Narcissus flore pleno*
*Serpentaria*, i. *Dracunculus*
*Serpentina*, i. *Scorsonera*
*Serpillum aureum, Citratum &c.* 454
*Sesamoides minus, aliquibus est Helleborus niger fernulaceus*
*Sicla & Sicula*, i. *Beta*
*Sidium idem quod Psidium*
*Siliqua sylueſtris*, i. *Arbor Iudæ*
*Simboline idem quod Zumbul Indicum, Hyacinthi Orientalis species* 502
*Sinapi* 506
*Sisarum*
*Sisyrinchium alterum*, i, *Cardamine altera* 171
*Sisyrinchium Mauritanicum &c,*
*Sisyrinchium Cordi*, i. *Ornithogalum*
*Solanum arboreſcens*, i. *Amomum Plinij*
*Solanum fœtidum spinosum Bauhini*, i. *Stramonium*
*Solanum Mexiocanum Bauhini*, i. *Mirabilia*
*Solanum pomiferum Gesneri & Bauhini*, i. *Pomum amoris*
*Solanum vesicarium*, i. *Alkakengi*
*Solanum esculentum Bauhini*, i. *Battatas de Virginia*
*Sol Indianus*, i. *Flos solis* 234
*Soldanella alpina*
*Sophonia*, i. *Amaranthus tricolor*
*Sorbus legitima & Torminalis* 567
*Spartum Austriacum*, i. *Gramen plumosum*
*Spartum Hispanicum frutex*, i. *Genista Hispanica* 442
*Spinachia* 496
*Staphylodendron*, i. *Nux vesicaria*
*Sternutamentoria*, i. *Ptarmica*
*Stœchas* 448
*Stœchas Citrina*, i. *Chrysocome*
*Struthium non est Saponaria* 353
*Stramonium maius & minus* 360
*Sumach Virginense* 612
*Susamgul*, i. *Lilium Persicum, & Hyacinthus stellatus, Byzantinus alter*
*Sycomorus, siue Acer maius latifolium* 610
*Symphitum maculosum*, i *Pulmonaria*
*Syringa alba, cærulea* 408
*Syringa Arabica flore albo duplici* 408
*Syringa Italica Lobelij*, i. *Syringa flore albo simplici*
*Syringa Italica flore albo pleno Besleri* 410

T

*Tabacco* 363
*Tamarix vel Tamariscus* 610
*Tanacetum vulgare* 482
*Tanacetum Peruanum*, i. *Flos Africanus*
*Tarchon herba* 500
*Taxus arbor* 606
*Thalictrum vel Thalietrum Hispanicum* 274
*Thesium Theophrasti*, i. *Radix cana*
*Thlaspi Beticum marinum* 390
*Thraupalus Theophrasti Dalechampio*, i. *Sambucus rosea*
*Thridacias*, i. *Mandragoras*
*Thuya*, i. *Arbor vitæ* 436
*Thymbra*, i *Satureia* 476
*Thymum legitimum capitatum* 454
*Thymum durius & latifolium* 474
*Tilia fœmina* 608
*Trachellum maius & minus* 354
*Trachelium Americanum* 356
*Tragium Dioscoridis*, i. *Fraxinella*
*Tragopogon cæruleum, purpureum* 302
*Tragopogon luteum* 514
*Tragoriganum Matthioli* 453
*Trifolium fruticans*, i. *Iasminum luteum*
*Trifolium* {*aureum*, *nobile*} i *Hepatica*
*Trinitas, seu herba trinitatis*, i. *Hepatica*
*Tulipa Armeniaca Bolonienſis, Bombycina, Byzantina, Cretica &c.* 52 &c.
*Tulipæ mediæ* 55
*Tulipæ præcoces* 48
*Tulipæ serotinæ* 61
*Tusai & Turfana*, i. *Corona Imperialis*

V.

*Vaccinium Virgilij* 128
*Valeriana rubra Dodonæi* 386
*Valeriana Græca* 388
*Valerianthon*, i. *Valeriana rubra Dodonæi*
*Verbascum odoratum &* {i. *Paralysis*
*Verbasculum odoratum*}

*Veratrum*

*Veratrum album & nigrum*, i. Helleborus albus & niger
*Vernilago*, i. *Chamaleo albus*
*Vetonica altera, vel altilis, aut Coronaria*, i. *Caryophyllus hortensis*
*Vetonica agrestis*, i. *Armerius*
*Victorialis rotunda*, i. *Gladiolus*
*Vincaperuinca* 391
*Viola alba*, i. *Leucoium*
*Viola alba bulbosa*, i. *Leucoium bulbosum*
*Viola Damascena*, i. *Hesperis*
*Viola flammea*, i. *Tricolor*
*Viola hyemalis*, i. *Hesperis*
*Viola latifolia* & *Viola Lunaris* } Bolbonach 265
*Viola lutea*, i. *Leucoium luteum sive Keiri*
*Viola peregrina*, i. Bolbonach
*Viola mariana* 354
*Viola martia* 281
*Viola Matronalis*, i. *Hesperis*

T

*Viola tricolor simplex & duplex* 282
*Viperaria & Viperina*, i. *Scorsonera*
*Vitis Corinthiaca, Damascena &c.* 563
*Vitis Virginiana* 564
*Vitis Virginense seu potius* Hedera *Virginiana* 612
*Vmbilicus Veneris*, i, *Cotiledon* 234
*Vva crispa*, i. *Grossularia* 560
*Vvularia*, i. *Trachelium, est &* Hippoglossum

Y.

*Vcca, sive Iucca* 434

Z.

*Zambach Arabi*, i. *Iasminum Arabicum*
*Zuziniare*, i. *Martagon Constantinopolitanum*
*Zumbul Arabi*, i. *Ornithogalum Arabicum*
*Zumbul Indi*, i. *Orientalis maior praecox*

# A Table of the Engliſh names of ſuch Plants as are contained in this Booke.


A.

White *Aconite* 214
Yellow *Aconite*, or winter Welfes bane, ibid.
*Adonis* flower 293
*Alkanet*, or Sea Bugloſſe 250
*Anemone*, or winde-flower, & the kinds 199 to 214
Yellow *Anemone* 194
*Alliſanders* 490
*Almond*, and the kinds 583
*Angelica* 529
*Apricocks* 579
*Apples*, and the ſeueral ſorts 585
Double bloſſomd *Apple* tree 404
*Apples* of Loue 379
Thorne *Apples* 360
*Arrach* white and purple 488
*Aſarabacca* 532
*Aſparagus* 503
*Aſphodill* and his kindes 146
*Aſphodill* with Lilly flowers 148

B

*Baldmony* or Gentian 350
*Balme* 479
The *Balſame* apple 278
*Barberies* 561
*Barbery Buttons*, and Thorny *Buttons* 339
*Barrenwort* 283
*Batchelours Buttons* double, white and red 254
*Batchelours Buttons* yellow 218 & 224
The *Bay* tree 598
The Cherry *Bay* tree, or *Bay* Cherry 401
The dwarfe *Bay* 397
The Kings *Bay*, that is, the Cherry *Bay* 400
The Roſe *Bay* 599
The Virginia *Bay* Cherry 400
The wilde *Bay* 192 & 258
The *Bee-flower* 330
*Beares breech* 235 &c.
*Beares* eares, and the ſorts 240
*Beares* eare Sanicle 344
*Beares* foote 489
*Beetes*, and the kindes 521
Garden *Beanes* and French *Beanes* 353
*Bell* flowers, and the kinds 354 & 356
Canterburie *Bels* 354
Couentry *Bels* 488
*Blites* 484
*Bloodwort* 359
The great blew *Bindeweed* 360
The ſmall blew *Bindeweed* 326
Blew *Bottles* 249
*Borrage*, and euer liuing *Borage*
Ladies *Bower*, and Virgins *Bower* ſingle and double 393
Dwarfe *Boxe* and guilded *Boxe* 606
Flower of *Briſtow*, or None ſuch 253
Spaniſh *Broome* 442
Double flowred *Bruiſewort* or Sopewort 352
Garden *Bugloſſe* 249
Marſh *bugloſſe* and Sea *bugloſſe* 250
*Burnet* 483
*Butterflie* Orchis 192

C

*Cabbage*, and his kindes 503
*Calues ſnout* or Snapdragon 269
Double *Camomill* and naked *Camomill* 290
Roſe *Campion* 252
Feathed wilde *Campion* ſingle and double 253
The Crimſon *Cardinals* flower 356
*Carawayes* 515
*Carnations* and Gilloflowers 306
*Caterpillers* great and ſmall 340
*Caſſidonie* 443
Lobels *Catch flye* 254
Cluſius his *Celaſtrus* 604
Sweete *Cheruill* or great *Cheruill* 494
Garden *Cheruill* ib.
Party coloured *Cicheling* 338
The *Chriſtmas* flower 344
The *Cherry* tree, and the kindes thereof 571
The double bloſſomd *Cherry* tree 402
The *Cypreſſe* tree 602
The ſweet gum *Ciſtus* 422
Burning *Clamberer* or *Climer* 391
Hungarian *Climer* 393
Virginian *Climer* or Maracoc ibid
*Coleflower*, *Colewort*, *Colerape* 504
*Colombines* 271
Tufted *Colombines* 274
The prickly euer greene *Corall* tree 604
*Corneflower* 326
*Corne ſallet* or Lambes Lettice 498
The *Cornell* tree 570
*Coſtmary* 482
*Cotten weedes* 379

*Cowcumbers*, diuers 524
*Cowslips* of diuers sorts 242. &c.
French *Cowslips* or Beareseares 235
*Cowslips* of Ierusalem 248
*Cranes* bill, and the kinds 228
Garden *Cresses* 500
Indian *Cresses* 280
*Crow* flower 253
*Crow* foote of diuers kinds 216 to 223
*Crown* Imperial 28
The double *Cuckow* flower 253 and 389
*Currans*, white, red and blacke 558
The true *Curran* Vine and Grape 563

D

*Affodils*, and the diuersities thereof, from 67 to 108
Checkerd *Daffodill*, and the kinds 44
*Dittander* 5ii
Bastard *Dittanie* 333
*Dogs* tooth Violet 193
*Dragons* 529
The *Dragon* flower 385
Double *Dasie*, and blew *Dasies* 321

E.

White *Ellebor* of two sorts 346
Wilde white *Ellebor* 347
The true black *Ellebor* or Christmas flower 344
Garden *Endiue* 495

F

*Ellworte* or Gentian 350
*Fennell* 492
*Fennell* flower 287
Double *Fetherfew* 289
The Princes *Feather* 232
The *Fig* tree, and the kinds 566
The Indian *Figge* tree 433
The *Finger* flower 383
The *Firre* tree 600
The Corne *Flagge* 189
The *flagge* or *flowerdeluce* 79 &c.
The *flowerdeluce* of Constantinople 79
The *flowerdeluce* of Persia 172
The bulbous *flowerdeluces* 172 to 179
The veluet *flowerdeluce* 188
Wilde *flaxe* or Tode *flaxe* 266
*Foxe* gloues, and the kinds 380
*Fillbeards* ordinary, and of Constantinople 562
*Flower* of Bristow, or None such, single and double 253
Purple *flower* gentle, and the kinds 371
Golden *flower* gentle or golden *flower* of life 372
The *flower* of the Pasion, or Maracoc 393
The *flower* of the Sun 395
The Sultans *flower* 327
The *friars* Crowne 332
*Fritillaria* or checkerd Daffodil 44
Double *fritillaria* ibid.

G

*Garlicke* 513
*Gentian* great and smal 350
*Germander* 456
*Gilloflowers* and Carnations 306
Queenes *Gilloflowers* or Dames Violets 262
Stocke *Gilloflowers* single and double 258
The *Ginny* hen-flower, that is, Fritillaria 44
*Goats* beard blew and purple 302
*Goats* beard yellow 514
Candy *Goldilocks* 372
*Golds*, that is, Marigolds 296
*Gooseberries* of diuers sorts 560
Herbe *Grace* or Rue 530
*Grape* flower 114
Vipers *Grasse* 301
Feather *Grasse* 458
Painted *Grasse* ib.
The *Guaiacum* of Padoa 570

H

*Hares* bels 122
*Hearts* ease single and double 282
The blew *Helmet* flower or Monkes hood 215
The wholsome *Helmet* flower ibid
*Hollihockes* single and double 369
*Holewort* or *Hollow-roote* 275
*Hyssope* common (Hellebore. 346. 344) 476
Guilded *Hyssope* 455
Hungary or mountaine Sea *Holly* 330
*Honisockles* double 404
Red *Honysockles* or vpright *Honysockles* 405
The euer green *Hawthorne* tree, or the euer green prickly Coraltree 604

I

*Iacinths*, and the seueral sorts 111 to 133
White *Iasmine* and yellow *Iasmine* 406
Double white *Iasmine* 408
Sweet *Iohns* single and double 319
*Ione* filuer Pin, that is, Poppies double 286
*Iudas* tree 437
The supposed Indian *Iucca* 434
The Virginia *Ivye* 612

K

Spanish Sea *Knapweede* 328

L

*Ladies* laces or painted grasse 458
*Ladies* smockes double 389
*Lambes Lettice* 498
*Langedebeefe* 486
The *Larch* tree 608
*Larkes* heeles or spurs single and double 276
Yellow *Larkes* heele, that is, Indian Cresses 280
*Lauender* spike 447
*Lauender* cotton 449
French *Lauender* or Sticadoue 448
Mountaine *Laurell* 398
*Leekes* 512
*Lettice*, and the kinds thereof 498
*Licorice* 533
The tree of *Life* 436
The chekerd *Lilly*, that is, Fritillaria 349
The Conual *Lilly*, or *Lilleconvally*

The Day *Lilly* 148
The Persian *Lilly* 28. 30
The Mountaine *Lilly* 33
The red or gold *Lilly* 39
The white *Lilly* 40
The *Line* or *Linden* tree 608
*Line-long*, or *Life everlasting* 375
Noble *Liverwort* or Hepatica 225
*Lungwort* or Cowslips of Ierusalem 248
*Lupines* white, blew and yellow 335

M.

SWeet *Marierome* 452
Guilded or yellow *Marierome* 446
French *Mallowes* 495
Spanish *Mallowe* 366
Shrub *Mallow* 369
Thorny *Mallow* 368
Venice *Mallow* ibid.
Red *Maithes*, and white 293
*Mandrake* male and female 377
The great *Maple* or Sycomore tree 611
*Marigolds* 296
Corne *Marigolds* of Candy 295
French *Marigolds* 303
Double Marsh *Marigolds* 224
The Spanish *Marigold* is the greatest double broade leafed Anemone 207
The blew or purple *Marigold* 299
*Martagons* of diuers sorts 33
*Mastick* the herbe 452
*Medlars* 568
The *Melancholy* Gentleman 260
Muske *Melons* 525
The *Mernaile* of the world, *of Peru. ib.* 364
*Mirtles* 427
*Moly* or *Mountaine* Garlicke 144 &c.
*Monkes* hoode, or Larkes spurs 276
*Monkes* hood or helmet flower 215
Counterpoyson *Monkes* hood 216
Blew *Moone-wort* 234
Halfe *Moones* 339
Golden *Mouse-eare* 330
*Mulberries*, and Virginia *Mulberry* 599
*Moth-Mullein* 383
Woody *Mullein* or French Sage 384
Ethiopian *Mullein* 385
The *Mumme* tree 432
*Mustard* 502

N.

SPotted *Nauelwort* 232
*Navew* 509
The *Nectorin*, and the kinds thereof 582
*Neesewort*, or *Neesing roote* 346
*Neppe* 479
The *Nettle tree* 568
Hungarian dead *Nettle* 385
*Nigella*, or the Fenel-flower 287
Tree *Night shade*, that is, the Winter Cherry tree 432
*None-such*, or the flower of Bristow 253
The bladder *Nut* 611
The Filberd *Nut* of Constantinople 562
The Spanish or Barberry *Nut* 171
The Wall *Nut* 594

O

THe euer greene *Oake* 606
Spurge *Olive* 397
Mountaine Spurge *Olive* ibid
*Onions*, and the kinds 510
Sea *Onion* 133
*Orchis* of Virginia 194
*Orenges* 584
*Oxe* eye 293
*Oxe* lips 245

P

*Pansies* single and double 282
*Parsley*, and sweet *Parsley* 491
Virginia *Parsley* 492
*Parsneps* 506
*Pasque* flower, or *Passe* flower 200
*Patience*, or Monkes Rubarbe 483
*Peaches*, and the kinds 586
Double blossomd *Peach* tree 404
*Peares*, and the seuerall sorts 590
The prickly *Peare*, that is, the Indian Figge 115
*Pearles* of Spaine 522
Garden *Pease* of diuers sorts 338
Crimson *Pease* blossome ibid
*Pease* euerlasting ibid
Blew vpright euerlasting *Pease* 292
*Pelletory* of Spaine 288
Double wilde *Pelletory*
*Penny* flower, that is, white Sattin
*Peony* single and double 342
*Perwinkle* single and double 392
*Pinkes* single and double 314
The *Pine* tree 599
The blew and the white *Pipe* tree 408
The double white *Pipe* tree, or double Iasmine 410
Rose *Plantane* 352
*Plums*, and the kindes 575
The *Pomegranet* tree 428
The double blossomd *Pomegranet* tree 430
*Pompions* 526
Double garden *Poppies* 284
Double wilde *Poppy* 286
*Potato's* of Spaine, of Virginia, of Canada 516
*Pride* of London 310
*Primme* or *Privet* 445
The euer-greene *Privet* 603
*Primroses*, and the kinds 242 &c.
Tree *Primrose* of Virginia 264
*Purslane* 499
*Purse* tassels 116 & 118
*Pushamin* or *Pishamin*, the Virginia *Plum* 570

Q.

QVinces, and the kinds 589

R

RAmpions 514
*Raspis*, white and red 557
*Reddish*, blacke *Reddish*, horse *Reddish* 509
Red and yellow flowred Indian *Reede* 376

*Rosarubie*, that is, Adonis flower 293
*Rose* tree, and the seuerall kinds 412. to 425
Iuno's *Rose*, that is, the white Lilly
The Elder or Guelder *Rose* 411
The Holly *Rose* or Sage *Rose* 421
The Mountaine *Rose* 424
Rocke *Roses* 397
*Rosemarie* common and gilded &c. 425
The Marie *Rose* or *Rosemary* of Silesia 424
*Rocket* 502
Garden *Rue* or Herbe grace 530
True *Rubarbe*, Monkes *Rubarbe*, *Rubarbe* of Pontus 483

S

*Sage* great and small 478
Guilded *Sage* 446
*Sage* of Ierusalem 248
French *Sage* 384
*Saffron* flowers of diuers sorts of the spring time and of the fall 160 to 170
Medow *Saffrons* or Colchicum, that is, the Sonne before the Father, and the kindes 154
Spotted *Sanicle* 231
Beares eare *Sanicle* 240
*Satyrion* 192
The *Sauine* tree 607
Summer *Sauorie* and winter *Sauorie* 476
White *Sattin* flower 265
Red *Sattin* flower 339
*Scabious* white and red 324
*Scorsonera* or Vipers grasse 501
Bastard *Sena* tree 440
The true and the ordinary *Seruice* 567
Mountaine *Setwall* 386
Virginia *Silke* 444
*Skirrets* 506
Our Ladies Slipper 347
*Smallage* 491
Ladies *Smocks* double 388
*Snayles* 338
*Snapdragons* 269
Mountaine *Soldanella* 434
Double flowred *Sopewort* 352
*Sorrell* 486
*Sowbread*, and the kindes 195 to 199
The Kings *Speare* or yellow Asphodill 148
*Sperage* or Asparagus 503
*Spiderworte*, and the kindes 150 &c.
*Spinach* 496
*Starre* flowers of diuers sorts, or *Stars* of Bethlehem 130, to 140
The greene *Starre* flower or bulbed Asphodill of Galen 136
*Starwort* or *Sharewort*, and Italian *Starwort* 299
*Sticadoue* or Cassidonie 448
*Stock gilloflowers* single and double 258
*Storkes bils* of diuers sorts 228
*Strawberries* of many sorts 526
The *Strawberry* tree 603
*Succory* 495
The *Sultans* flower, or Turkie Corne flower 327
The *Sun* flower or flower of the Sun 295
The Virginia *Sumach* 611
The Myrtle leafed *Sumach* ibid
The *Sycomore* tree 610

T

INdian *Tabacco* of diuers sorts 363
The greene and the white *Tamariske* tree 610
*Tansie* single and double 482
The blessed *Thistle* 530
The gentle *Thistle*. Globe *Thistle* &c. 332.
Christs *Thorne* (thorne apple. 360) 607
*Thrift* ordinary, and the great Sea *Thrift* 317
*Throatwort*, and Giants *Throatewort* single and double 354
Beane *Trefoyle* 438
Shrub *Trefoile* 407
Tree *Trefoile* 439
Candie *Tufts* 390
Golden *Tufts* 375
Spanish *Tufts* 274 & 340
The early flowring *Tulipa* 46
The meane flowring *Tulipa* 54
The dwarfe *Tulipa* 52
The Persian *Tulipa* ibid
The Turkes Cap, that is, the *Tulipa*
The true *Time* 454
Guilded *Time*, Moske and Lemmon *Time* ibid
Garden *Time*, and Mastick *Time* 474
Tarragon 501

V

REd *Valerian* of Dodonæus 386
Greeke *Valerian* 388
Mountaine *Valerian* 386
*Violets* single and double 282
The bulbous *Violet* 169
The Dogs tooth *Violet* 193
Dames *Violets* 262
Mercuries *Violets* 357
*Vines*, and the seuerall kinds of grapes 564
*Vipers* grasse 301
*Virginia Vine* 564
*Virginia Vine*, or *Virginia Ivie* 612

W

THe *Walnut* tree 594
Single and double *Wall-flowers* of many sorts 257
*Widow Wayle* 397
The *Willow-flower* 270
The *Wind-flower* or Anemone, single and double of many sorts 199 to 214
*Wild Wind-flower* single and double 202
Sweet *Williams*, and the kinds 319
*Winter* Gilloflowers 258
*Winter Wolfes* bane, and yellow *Wolfes* bane 214
*Winter* Cherries 532
The *Winter* Cherrie tree 431
Double *Wood-bine* or Honisuckle 404

Y

THe *Yew* tree 606

THE

# *A Table of the Vertues and Properties of the Hearbes contained in this Booke.*


A.

IN feare of *Aborſment*, or miſ-carrying in women 478
Good againſt *Aches* 290
Good in hot *Agues* and to driue away their fits 223.484.532.536.538.564 575.579.584 595
From whence *Agaricke* is taken 608
A Syrupe of *Angelica* 529
Good for deiected appetites, 484.486. 499.561.562.578.575.584.
To perfume *Apparell*, Leather, &c. 421 448.586
For the *Apoplexie* 349
*Aſtringent* or to binde 428.431

B

FOr weake *Backes* and reines 479.483
A *Balſame* for green wounds 426
To cauſe *Barrenneſſe* 284
Good for *Bathing* 294.376.421.449 474.477.478.479.480
Good for *Bees* 440.480
Good to take away the ſting of *Bees* and waſpes 476.479.480.482
Good to open and mollifie the *Belly*, and make it ſoluble 333.370.421.488 489.490.495.566.578.582
To clenſe the *Blood* 484
To diſſolue clotted *Blood* 610
To encreaſe *Blood* 567
To helpe the ſpitting of *Blood* 595
To ſtay *Bleeding* 230.396.483
Hurtfull for a ſhort *Breath* 499
Good for the ſhortneſſe of *Breath* 364. 436.502.506.513.533
For a ſtinking *Breath* 529
To make a ſweet *Breeth* *ibid*
For cold and moiſt *Braines* 335.427.448 481.482
To abate the ouer-ſwelling of Maidens *Breaſts* 600
To helpe paines in the *Breaſt* 500
To heale womens ſore *Breaſts* 590
To draw out broken *Bones* &c. out of the fleſh 190
For *Bruiſes* by fals &c. 479.608.610
Good for *Burnings* or ſcaldings 362.512 610

C.

TO clenſe *Cankers* 364
To procure *Caſtings* or vomitings 189.434.442.489.532
To ſtay *Caſting* or vomiting 477.480
To feede or make Cattell fat 348.440 442
*Cauſticke* or burning plants 396
The beſt *Coales* for Gunpowder 610
The beſt *Coales* for laſting 608
For the Winde *Collicke* 216.293.453 455.476.504.508.515.530
To *Clenſe*, purge and dry 276
To ſet an orient red *Colour* on the cheeke of a woman 241
To make a deepe blew *Colour* 179.603
To make a red *Colour* ibid
To giue a luſtre to a white *Colour* 595
To make a yellow *Colour* 421
For the *Colts* euill, or immoderate luſt 499
To warme and comfort *Cold* griefes 214 376.434.477.478.480.491
*Cooling* and drying 323.353.378.421 438.486,499.528.562.557.564.576 562.568.576.578.590.594.598.602 603.607
*Cooling* and moiſtning 234.283.380.432 579
For a *Conſumption* 519.608
*Cordiall* to comfort the heart 170, 216 252

252.298.301.314.328.421. 427.436
450.480.482.483.526.528.530.558
586
For the *Cough* in young children 502
Good for *Coughes* and colds 134.295.513
533.562.567.575.584.586
Mithridates *Counterpoyson* 567
Good for *Crampes* and shrinking of sinews
425

D.

TO procure an easie and speedy *Deliuery* to women in trauell 40.199.274
410
To stay rheumaticke *Distillations* 288
376.386.427
To cure the biting of a mad *Dogge* 300
353.482
To helpe the *Dropsie* 235.290.353
To cause *Drowsinesse* like vnto drunkennes
362

E.

FOr paines in the *Eares* 580.584
Good for sore and weake *Eyes* 216
421.427.479.530
Hurtfull to the head and *Eyes* 489
For the *Epilepsie* or falling sickneſſe 194
335.344.456.477.502

F.

TO cleanse the *Face*, and other patts of the skinne, and make it fresh 40.189
247.336.396.500.502.521.528.566
584.589.600
For the *Falling* sickneſſe 194.335.344
456.477.502
*Farsing* or *fasseting* herbes 474.476.478
To heale *Felons* on the ioynts of the fingers 492
To procure the Feminine courses 289
335.453.456.477.479.491
To stay the Feminine or menstruall courses 372.396.480.486.608.611
To cleanse *Fistula's* 364
To expell thin *Flegme* 477
To extenuate & expectorate tough *Flegme*
134.264.436.448.456.477. 500, 529
530.533.608
For the bloody *Flixe* and all other *Fluxes*
193.231.372.425. 431.446, 483.486
568.595.602.605.608, 611
To take away *Freckles*, spots, &c. 500
566
For the *French* disease 283.353.606

G

TO stay the *Gonorrhea* or running of the reynes 608
Good for the *Gout* 349

H.

TO cause the *haire* to grow 566
For the falling of the *Haire* 425
Hurtfull to the *Head* and eyes 489
Good for the *Head* and *Heart* 134.170
298.426.455.474.530.586
Good for the *Head* and stomack 455.474
530.586
For paines in the *Head* 288.292.426
448.499
For the swimming and dissineſſe of the *Head* 241.247.586
For the passion of the *Heart*, and to make it merry 480.528.529,589
To expell venemous vapours from the *Heart* 170.301.529.530.586
To procure *Health* 477.478
To ease the *Hemorrhodes* or piles 513
580
For an *Hoarseneſſe* 580.584.600
For the bots in *Horses* 606.607
For the cough in *Horses* 595
To stay the *Hickock* 494
To purge flegmatick and watery *Humours*
329

I.

FOr the yellow *Iaundise* 132.134.150
170.275.290.531.562
To make excellent *Inke* 431
To asswage hot *Inflammations* 362.378
380
To helpe the *Itching* of the *Head* 477
To cure the *Itch* 380

K.

TO heale exulcerated *Kidneyes* 533

L.

TO stay a *Laske* or looseneſſe 132.323
421.425.446.483.562.568.570.571
578.595
For the *Lepry* and deformity of the skin
306
For chapt *Lips* and hands &c. 589
To tanne or thicken *Leather* 611
To clense the *Liuer* 484.532
To coole and strengthen the *Liuer* 226
448.486
Good for the *Lunges* and old coughes
134.249.300. 353.364.436. 448.456
477.502.506.513. 529. 530.533.600
608
To stay immoderate lust 499.529

M.

TO take away blew *Markes* 427.500
502.566
For the *Measels* and small pocks 170.216
To straw on *Meate* 474.476

For

For *Melancholicke* diſeaſes 345.450.448 455,510,589
To comfort & ſtrengthen cold and weake *Members*, 170,290,407,427,448,449 453,474,477,491,598
To ſtrengthen the *Memory* 427,428
To keep *Milke* from curdling in the ſtomacke 482
To increaſe *Milke* in womens breaſts 440 504
For the *Morphew* and other diſcolourings of the skin, 336,396,427,566
For the *Mother* in women 344,378
To keep garments from *Moths* 376,611
To engender *Moths* 386
Good to waſh and clenſe the *Mouth* 428 431, 446,528,557,564,598
To cure Vlcers in the *Mouth* 431
For a ſurfet of *Muſhroms* 513

N.

TO waſte *Nature* 530
For a cricke in the *Necke* 66
To procure *Neeſing* 189,289
To ſtay bleeding at the *Noſe* 396

O.

TO open *Obſtructions* 448,484,492. 532
To take away the offence of the ſmell of *Onions*, Garlicke, Leekes &c. 512
A remedie for them that haue taken *Opium* too liberally 289

P.

TO eaſe *Paines* 290,370
For the *Palſie* 241,247
For the *Plague* or peſtilentiall feuers 160, 170,216,275,298,328,333,335,353, 483,495,513,529,530,586,595,603.
From whence *Pitch* is made 600
For the ſmall *Pocks* 170,216
An eſpeciall Antidote againſt *Poyſon* 134,339,353
Againſt the *Poyſon* of the Helmet flower and other venemous herbes, and againſt all other infectious diſeaſes 216,333, 335,353,483,529
Againſt the *Poyſon* of the Spider Phalangium, Scorpions, Serpents, and other venemous beaſts 152,301,328,333, 335,402,453.477,607
To take away the ſtrength of certaine *Poyſons* 590
*Poyſon* to all fourefooted beaſts 402
For ſweet *Pouthers* and ſweet bags 189 421,450,453,586
To *Purge* gently 284,421,566
To *Purge* vehemently, and ſtubborne diſeaſes 189,346,360,400,562
To fatten *Pullen* 440,442

R.

TO reſtore *Raviſhment* 600
To hinder young perſons from growing *Ripe* too ſoone 132
Fences of *Reedes* as good as wals 510
To draw *Rheume* 214,288,292
To ſtay *Rheumaticke* diſtillations 287 288,376,427,482
From whence *Roſſen* is made 60
Good for *Ruptures* 602

S.

TO put into *Sawſages* &c. 476,477,478, 480,482
For *Scaldings* and burnings 362,512
To take away *Scars* and markes 223, 247 336,521,568
For the *Sciatica* or paine in the hippes and ioynts 160,442
To clenſe the head of *Scurfe* 134,396 477,479,607
For the *Scurvie* 389,510
To increaſe *Seede* 524
To eaſe paines in the *Sides* 532,599
To cleare the *Sight* 134,427,479,530
To comfort the *Sinewes* and ioynts 426 427,477,478,480,595,599
To procure *Sleepe* 286,362,378,499
To keep the body *Soluble* 333,376,421 480,488,495,566,578,582
To clenſe the head of running *Sores* 134 607
For the ſhrinking of *Sinewes* 425,426 599
To breake plague *Sores* 223,300
Good for *Sores* and wounds 303,446 600
Good for the *Spleene* 198,288.455.456 474.490.502.510.530.610
To draw out *Splinters* &c. out of the fleſh 190
Good for *Stitches* 599
Good for the *Stomacke* and Liuer 134, 353.455,473,483.491,532
Hurtfull to the head and *Stomacke* 489 607
For cold and windy *Stomacks* 301,455 476.491,495,586
To coole an hot *Stomacke* 380,486, 499.525,526,528,558,560,562,566, 575,584,589
To warme a cold *Stomacke* 474,477,480 482,495,500,515,529,586,598
For the *Stone* in the reynes and kidneyes 230,274,293,335,370,425,492,503. 504,510,521,526,532,533,566.575. 582,584.607.608.611
For the *Strangury* 491
Good againſt *Surfets* 287

To

To cause *Surfets* 582
To prouoke *Sweating* 264.290.532
To helpe *Swounings* 274.301.529

T.

TO draw out *Thornes* &c. out of the flesh 190
Good for swollen *Throats* 300
To wash the mouth and *Throate* 357.478 528.598
For the *Tooth-ach* 134.292
The poore mans *Treakle* 514
For *Trageas* or dredges 573
Common *Turpentine* from whence it is taken 600
Venice *Turpentine* from whence it is taken 600.608

V.

TO helpe *Venery* 66.190.193,194. 502.506.524
To take away *Vermine* and Lice in the head &c. 477
To recouer the *Voyce* being lost 513
To cicatrise old *Vlcers* 275.364
To clense *Vlcers* 492.580
For running *Vlcers* 364.595
To procure *Vomiting* 189.439.442. 532
To stay *Vomiting* 477.480
To cure the biting of a *Viper* or Adder 302
To cause the *Vrine* to seeme blood 433
For the stopping of *Vrine* 132.264.353 376.453.456.483.491.492.502.503 506.508.515.525.530.532.533.582 602.608

W.

TO purge *watery* humours 329
To make sweete *Waters* 421,450.453 482
To breake *Winde* 301.455.476.491, 494.508.515.529.530.575.586
To stay the longing of *Women* with childe 561.564
For the *Wormes* 134.216.336.345.449. 482.483.500.502.532.582.607
Good to cure *Wounds* 241.252.364.389 445.446.456.480.483.492
Good *Wound* herbes 230.231.235.241. 323.383.389.406.

*Faults escaped in some Copies.*

Folio 8. line 14. for own reade home. f.12.l.27. for trouble reade treble. f.42. l.5. reade, like vnto that of a Lilly f.66.l.42 for χήρε read χρίνε or λείεται. f.73. l.37. for top of the flower, read cup. f.134.l.36. for composed reade compassed. f.150.l.4. for hath, reade haue, and line 5. for is are. f.173.l.12. put out these Wordes, the inside, in the beginning of the line. f.189.l.38. reade *Binis florum ordinibus*. f.218.l.19. reade goulons, and line 28 pratensis. f.272. line 36. read Pothos. f.276 l.12. Chelidonia. f.281.l.37. for hath, haue. and l.28. Mastuerzo. f.284. l.15. Vicenza. f.287.l.39. Citrina. f.290 l.39. reade prouoke, and, helpe. f.329.l.37. for Melancholicke, reade Flegmaticke. f.330.331.333. reade Eryngium in all places f.336.l.8. reade, and not very flat. f.356.l.31. Americanum. f.357.l.26. Cervicaria. f.358. l.45. reade, before it can haue. f.372.l.9. blot out, except it. f.389. for spockes, reade smockes. f.393.l.3. in the margent for cæruleo, read pleno. f.397.l.10. reade dwarfe. f.424.l.45. Hirculus. f.428.l.20. Tarentina. f.431.l.10. Cyprium. l.19. Amomum. f.438.l.17. for Dioscorides, reade Theophrastus. f.442.l.3. for castings reade purgings. f.509.l.35. reade γογγύλη. f.513.l.24. transferre all that clause of Onions vnto the other side, vnder the vse of Onions. f.516.l.37. transferre these words, [Bauhinus vpon Matthiolus calleth it *Solanum tuberosum esculentum*] vnto the former Potatoes of Virginia. f.520 l.13. for swelleth, read smelleth. f.541.l.51. reade, after your stockes rayed from stones. f.566.l.20. for as, read and. and l.29, euery one. f.567.l.24. for Rice, read Rue. f.575 l.8. reade serue to be ministred to the sicke. f.588 l.3. Capandu. f.594.l.18. for facility, read faculty. f.595.l.39. reade Ricinus. f.600.l.4. Fuchsius

LONDON,
*Printed by* HVMFREY LOWNES *and* ROBERT YOVNG
*at the signe of the Starre on Bread-street hill.*
1629.