


504313

A MANUAL
OF
STRUCTURAL BOTANY:

FOR THE USE OF
CLASSES, SCHOOLS, & PRIVATE STUDENTS

BY M. C. COOKE.



WITH UPWARDS OF 200 ILLUSTRATIONS BY RUFFLE.

NEW EDITION.

LONDON:
ROBERT HARDWICKE, 192, PICCADILLY;
AND ALL BOOKSELLERS.

CONTENTS.

	PAGE
1. INTRODUCTORY	1
2. ULTIMATE ELEMENTS	3
3. CARBONACEOUS COMPOUNDS	8
4. NITROGENOUS COMPOUNDS	12
5. MINERAL COMPOUNDS	14
6. CELLS	18
7. VASCULAR TISSUE	21
8. CELL DEVELOPMENT	23
9. THE EPIDERMIS	25
10. THE DESCENDING AXIS	29
11. ASCENDING AXIS—EXOGENS	32
12. DITTO ENDOGENS AND ACHROGENS	36
13. SUBTERRANEAN STEMS	38
14. LEAF-BUDS	41
15. LEAF-STRUCTURE	43
16. LEAF-FORM	47
17. LEAF-ORDER	51
18. LEAF-APPENDAGES	53
19. FLOWER-BUDS	55
20. INFLORESCENCE	58
21. THE CALYX	61
22. THE COROLLA	63
23. STAMENS	67
24. THE PISTIL	72
25. THE OVULE, OR SEED-BUD	75
26. THE FRUIT	78
27. SEED	84
28. GERMINATION	87
29. REPRODUCTION OF ALGÆ	89
30. DITTO LICHENS	91
31. DITTO FUNGI	94
32. DITTO MOSSES	96
33. DITTO FERNS	98
34. VEGETATIVE PHENOMENA	100
35. EPIPHYTES AND PARASITES	103
36. GALLS AND EXCRESCENCES	104
37. VITAL ACTION	106
38. BOTANICAL DESCRIPTION	109
39. DESCRIPTIVE MODELS	111
40. SUMMARY	113
INDEX	139

P R E F A C E.

HAVING experienced the want of a cheap Manual to place in the hands of students in the Botanical Classes established for Operatives, in connection with the Department of Science and Art, I have made this effort to supply the want in the manner which, to my judgment, appeared best calculated to render it serviceable to them. I have purposely divided the Manual into Forty Sections, corresponding with the Forty Lessons which the Department requires should be given. Each of these sections contains an outline of the chief points necessary to be demonstrated to a Class at the respective lessons, unencumbered with such illustrations as a teacher would give, but which would needlessly have swelled this volume and augmented its price. In its present form, it is calculated to refresh the memory of the student concerning the past, rather than, of itself, instruct him in the future of his studies. No attempt has been made to popularize the subject;—this must be considered rather as a skeleton of dry details, to be filled up and attired according to the taste of the demonstrator. The Illustrations are not intended to supersede diagrams, drawings, and the liberal use of the black board, but

as helps for those who may not be able to use the pencil with sufficient facility in their note-books. The orthography and etymology of the technicalities of this branch of the science will, it is hoped, prove an assistance to the teacher, as there will be less necessity for him to write up every new term as it occurs, so as to familiarize his pupils with its orthography. No note has been taken of Economic Botany. Information of this kind is more readily obtainable, and the oral instruction which the teacher may give in the course of his demonstrations, will be more popular in its character and more easily remembered.

Independently of the primary object of the work, it is believed that it will be found useful to the young botanist and private student as a prelude to other and more expensive treatises.

M. C. C.

November, 1861.

NOTE TO THE PRESENT EDITION.

A NEW edition of this Manual being demanded, it has been thoroughly revised; the Chemical nomenclature corrected, to bring it into harmony with the requirements of advanced science, and several minor alterations adopted, which, it is hoped, will greatly improve the work, and ensure for it a continuance of public favour.

June, 1870.

ovule

orthotropal
campylotropal
anatropal
Pisum
Secundine
Tortile
Quartile
quintine or Embryosac
Hilum
Microphyte
Chalazal, Raphe

Fruit - Pericarp, Seed

Seed - Embryo, Albumen or perisperm.

embryo - Cotyledon
Radicle, Plumule.

Fruit

Alocarpous, Syncarpous, pericarp = Epicarp
mesocarp, endocarp

Septicidal
Loculicidal
Septitragal

A MANUAL OF STRUCTURAL BOTANY.

SECTION I.

INTRODUCTORY.

WE are accustomed to regard the whole of the material world as divisible into three great *kingdoms*, *i. e.*, mineral, vegetable, and animal. It is not so easy to determine the limits of the latter two, as to distinguish them from the former, so intimately do they blend into each other. No definition has yet been attempted to which exception might not be taken. The old distinction that animals live and move; that plants live, but do not move; and that minerals neither live nor move; will only hold good to a certain extent, for we have *fixed* animals and *moving* plants; although animals are generally possessed of, and plants deficient in, the power of locomotion. Another distinction, that plants respire oxygen under the influence of light, and animals carbonic acid gas, was accepted, until certain plants, of a low organization, were found to respire carbonic acid gas under all circumstances. Chlorophyll and starch were supposed to be exclusively vegetable products till the former was discovered in the fresh-water hydra, and the latter in the brain of man.

Dr. Lankester gives the following distinctions :—

“ I. Plants derive their nourishment from mineral or inorganic substances, animals from organized matters.

“ II. Plants absorb their nutriment to a greater or lesser extent by their whole external surface, whilst animals possess a mouth, and take up their nutriment from an internal bag or stomach.

“ III. Plants are mostly fixed, whilst animals have a power of moving about. To this distinction, however, there are numerous exceptions.

“ IV. The tissues of plants are distinguished by the tendency to absorb carbonic acid and throw off oxygen, whilst those of animals absorb oxygen and throw off carbonic acid gas.

“ V. Both plants and animals are developed from an impressible and motile protoplasm, but in animals this substance becomes differentiated into nerves and muscles.”—(Lectures on Botany,—Science and Art Department.)

The study of all that relates to the vegetable kingdom is included in the science of *Botany*.

I. That department of the science of Botany which has reference to the textures of which plants are composed, and the forms of their organs, is termed *Structural Botany*, *Vegetable Anatomy*, or *Organo-graphy*.

This department of the science may be regarded as again subdivided into *Vegetable Histology*, which confines itself to the tissues of plants, and *Vegetable Morphology*, which treats of the forms of the various organs.

II. The second department of Botanical Science is that in which plants are considered as living beings, and which takes especial cognizance of their vital functions. It is therefore termed *Physiological Botany*.

The aid of Chemistry and the allied sciences is

requisite for a thorough acquaintance with these departments.

III. Another department concerns itself with all that relates to the arrangement and classification of plants, and is termed *Systematic Botany*. The knowledge of this depends on an acquaintance with the former divisions of the science.

IV. The distribution of plants over the surface of the globe forms another division, which, from its intimate connection with geography, is termed *Geographical Botany*.

V. *Fossil Botany*, or a special study of the vegetable remains found deposited in geological formations, links Botany with Geology, and being based upon the results obtained in the former divisions of the science, applies them in elucidation of the plant-history of the past. It has also been called *Geological Botany*.

These departments have been named in the order in which they should be studied. The first and second only will be treated of in this volume, not separately but simultaneously, the functions of the organs with the descriptions of their forms.

The title selected must therefore be accepted in a broad and general sense as embracing these two subdivisions, and not as restricted to one.

SECTION II.

THE ULTIMATE ELEMENTS OF PLANTS.

An *Element*, or simple substance, contains, so far as we know, but one kind of matter. Sixty-five elements in various forms of combination make up the whole mass of the earth, air, and water.

Fifteen elements invariably occur in plants and animals. These elements are found combined together in the vegetable and the animal structures as *compounds*.

When a plant is burnt, some of its elements enter into new combinations, become *volatile*, and are dissipated in the air as gases or vapours. The volatile elements of plants are four—oxygen, hydrogen, nitrogen, and carbon.

The non-volatile or *fixed* elements of plants are eleven in number ; viz., phosphorus, sulphur, silicon, fluorine, chlorine, potassium, sodium, calcium, magnesium, iron, and manganese.

Phosphorus, sulphur, and fluorine are sometimes found in the volatile part of plants, while, on the other hand, in the fixed part, carbon and oxygen always occur, nitrogen rarely. Two other elements, iodine and bromine, are contained in seaweeds, while several additional metals, copper, aluminium, zinc, and rubidium, have been detected in particular plants.

ELEMENTS OF THE VOLATILE PART OF PLANTS.

O, Symbol ; atomic weight, 16. OXYGEN (*Oxys*, acid ; *gennao*, I produce) is only known as a gas. The atmosphere contains in 100 measures 21 of oxygen and 79 of nitrogen, another elementary gas. The majority of the compounds which form the food of plants contain much oxygen ; of these compounds, *carbon dioxide*, often called carbonic acid, contains 2 atoms of oxygen and 1 atom of carbon (CO_2), while another, *water*, contains 2 atoms of hydrogen and 1 atom of oxygen (H_2O). Plants have the power, under the solar influence, of decomposing carbon dioxide and other oxygen compounds. In the case of carbon dioxide the carbon is retained by the plant and re-combined with other elements, and the oxygen is

set free. Thus the atmosphere is purified and supplied afresh with oxygen, the great supporter of combustion and life. In the putrefaction of vegetable and animal matters oxygen is absorbed, and they thus once more return to the inorganic or mineral world.

H, Symbol ; 1, atomic weight. **HYDROGEN** (*Hydro*, water ; *gennao*, I produce) is only known as a gas. It is the lightest body in nature. Burnt in the air, it combines with oxygen, forming water. Water enters into the composition of plants, and through the agency of water they take up a large proportion of the materials of their structure.

N, Symbol ; 14, atomic weight. **NITROGEN** (*Nitron*, nitre ; *gennao*, I produce) is only known as a gas. It is chemically inactive. Combined with hydrogen it forms *ammonia* (NH_3), with hydrogen and oxygen *nitric acid* (HNO_3).

C, Symbol ; 12, atomic weight. **CARBON** (*Carbo*, coal) is a solid element existing in three different states,—the diamond, graphite or blacklead, and charcoal or lamp-black. When burnt, carbon yields carbon dioxide, a gas which is also evolved in the oxidation of the carbon of food which goes on in the animal body.

THE FIXED ELEMENTS OF PLANTS.

P, Symbol ; 31, atomic weight. **PHOSPHORUS** (*Phos*, light ; *phero*, I bring) is a wax-like solid in one of its forms. It does not naturally occur free, but only in combination with metals as phosphides, or with metals and oxygen as phosphates. It is extremely inflammable, and emits light by slow combustion at the ordinary temperature.

S, Symbol ; 32, atomic weight. **SULPHUR** (*Sal*, salt ; *pyr*, fire) is a brittle, yellow, combustible solid, which melts and vaporizes by heat. Sulphur and hydrogen unite to form an offensive-smelling gas, hydrosulphuric acid (H_2S) ; with oxygen it unites, burning with a blue flame and forming sulphur dioxide (SO_2), a pungent gas. Sulphates contain sulphur, a metal, and oxygen ; sulphuric acid, sulphur, hydrogen, and oxygen (H_2SO_4).

Si, Symbol ; 28, atomic weight. **SILICON** (*Silex*, flint). United with oxygen, silicon forms a dioxide called silica (SiO_2). This is common as flint, agate, sand, and quartz, and enters largely into the composition of the stems of meadow grasses and wheat.

F, Symbol ; 19, atomic weight. **FLUORINE** (*Fluo*, I flow) is found in minute quantity in many, perhaps in all, plants. Combined with calcium it forms the mineral fluorspar (CaF_2), and in this state occurs in bones and teeth.

Cl, Symbol ; 35.5, atomic weight. **CHLORINE** (*Chloros*, green) is known usually as a gas of a pale greenish-yellow colour and suffocating odour. Combined with sodium it constitutes common salt, sodium chloride ($NaCl$). A compound of chlorine, calcium, and oxygen ($CaCl_2O_2$) is used for disinfection and bleaching. This element occurs abundantly in marine and maritime plants.

I, Symbol ; 127, atomic weight. **IODINE** (*Iodes*, violet-coloured). A heavy, greyish-black, lustrous and crystallized solid, which turns on heating into a violet-coloured vapour. It stains the skin yellow, and gives with starch a rich blue colour : it occurs in seaweeds as metallic iodide.

Br, Symbol ; 80, atomic weight. **BROMINE** (*Bromos*, stench), a heavy, deep brownish-red liquid, which gives off a vapour of the same colour and of disagreeable odour.

K, Symbol ; 39, atomic weight. **POTASSIUM** (*Potashes*), a bluish-white soft metal, lighter than water, and decomposing at ordinary temperatures. Potassium is more abundant in cultivated than in wild plants. It occurs in them as chloride and sulphate, and in the salts of the vegetable acids.

Na, Symbol ; 23, atomic weight. **SODIUM** (*Natrium*, from *natron*, native sodium carbonate) is a silver-white metal, rather lighter than water. In its occurrence and uses in plants it closely resembles potassium.

Ca, Symbol ; 40, atomic weight. **CALCIUM** (*Calx*, lime). Its oxide is lime (CaO), its hydrate, slaked lime (CaH_2O_2), and its carbonate, chalk and whitening (CaCO_3). Calcium carbonate is dissolved by water holding carbon dioxide in solution. Many plants contain much calcium, as oxalate, citrate, &c.

Mg, Symbol ; 24, atomic weight. **MAGNESIUM** (*Magnesia*, in Asia Minor) is a silver-like metal, which burns with a brilliant white light when heated, and forms magnesia (MgO). It occurs abundantly in the grain of wheat.

Fe, Symbol ; 56, atomic weight. **IRON** (*Ferrum*, iron). The presence of iron in plants seems connected with their green colour.

Mn, Symbol ; 55, atomic weight. **MANGANESE** occurs in minute quantity in plants ; but it is a variable ingredient.

Of the other metals found in plants copper seems most generally diffused, though it occurs in very small traces only.

SECTION III.

CARBONACEOUS COMPOUNDS.

CELLULOSE ($C_6H_{10}O_5$) is a compound of carbon, hydrogen, and oxygen. It constitutes the basis of vegetable tissues. The elastic transparent membrane of cell-walls is composed of this substance, which is permeable by fluids. It is allied to starch, becomes yellow on the addition of iodine, and when acted upon by sulphuric acid and iodine, assumes a blue colour. *a*

LIGNINE is found deposited upon cellulose, forming the woody fibres of plants, and giving strength and solidity. When lignine is submitted to the action of sulphuric acid, it is converted into glucose. *Solub in H₂O, Soluble in Alkali*

STARCH ($C_6H_{10}O_5$) is a common product of plants. It exists in the form of granules, or minute cells, which become blue upon the addition of iodine. These granules possess such distinctive features in size and shape, that the mixture of the starch from one kind of plant with that of another may be detected by the microscope. They are ordinarily oblong, spheroidal bodies or flattish discs, marked with parallel striations. Some are of very irregular form. In almost all *marginal* instances there is a ~~central~~ spot or hilum, and the series of lines arrange themselves around it. There is also a cell-wall, but no central cavity. The markings

Insoluble in cold & boiling water, Alcohol, ether, dilute acids and almost in weak alkaline solutions. Convert to a treacly looking ~~substance~~^{light} by the action of strong Sulphuric acid, which after dilution & boiling is changed 1st into dextrine & then into grape sugar. Has been discovered in the tissues of molluscan animals. (N.S. 1: 111.)

6. It has been detected in an amorphous condition by Schlieden in the bark of Jamaica Sarsaparilla & in some seeds & underground stems.

Starch occurs in mature structures removed from the influence of light (as pith, root, seed

Chlorophyll - occurs in young structures exposed to light - When starch occurs in active vegetating parts it is invested by chlorophyll.

(I.O.)

^{dull white}
Arrowroot, Sago, Potato, Tous-le-mois, ^{Bananafruit} are most
abundant in starch. ^{banana}

Single grains distinct in Potato, wheat, peas
Aggregated - West Indian Arrowroot, Portland Sago (
Arum Maculatum)

Amorphous starch [Jamaica Sarsaparilla,
Cardamomum minus (Lud), Cassia
arenaria (Lendugrad stem), certain
Fungi, Ornithogalum nutans &
Lancedatum -

Sago from - Sagu, Metroxylon
Brazil - (Alvaula acuminata)
Portland Sago (itum maculatum)
Tapioca - (Cassava)
Tous-les-mois (Canna)

are supposed to be produced either by the foldings of the cell-walls, or by the successive layers of starch.

This substance is insoluble in cold water; when submitted to the action of boiling water, the cells swell, the cell-walls burst, their contents are discharged, and the mass becomes a tenacious paste. Starch is found in almost all the cellular parts of plants, but is most plentiful in roots and tubers, and in the grains of cereals.



Fig. 1.

Starch is convertible by heat into dextrine, or by the action of boiling weak sulphuric acid, into a mixture of dextrine and glucose (grape sugar), and ultimately into glucose. *When sold up by tubers.*

Plants contain more starch at certain periods than at others. During germination and at the time of flowering much of it is converted into sugar. It is this conversion of starch into sugar during germination which has been taken advantage of in the process of malting, to convert the starch of grain into sugar by forced germination.

Inuline is a variety of starch, obtained from dahlia tubers, elecampane, Jerusalem artichoke, and some few other plants.

Lichenine ($C_6H_{10}O_5$) is the peculiar starch of lichens. It is found in Iceland moss (*Cetraria Islandica*), Reindeer moss (*Cænomyce rangiferina*), and carrageen, or Irish moss (*Chondrus crispus*).

DEXTRINE ($C_6H_{13}O_5$). During the change of starch into sugar, an intermediate stage occurs in which the starch is converted into dextrine or soluble starch; for it is then soluble in water. Starch must first be converted into dextrine or sugar before it can afford nourishment to the growing parts of plants.

SUGAR may be said to exist in plants under three forms.

Cane sugar ($C_{12}H_{22}O_{11}$), which is found in the sugar-cane, maple, and beet-root. This is not a fermentable sugar, but must first be converted into glucose.

Grape sugar, or *glucose* ($C_6H_{12}O_6$), gives sweetness to most fruits, is a product of the metamorphoses of starch, cane sugar, and woody fibre. This is fermentable.

Mannite ($C_6H_{14}O_6$) is an unfermentable sugar found in the manna ash, mushrooms, celery, some sea-weeds and other plants.

GUM ($C_{12}H_{22}O_{11}$) is one of the substances produced abundantly in plants. It exists in many seeds, exudes from the stems and branches of trees, and is found in the juices of others. It may be considered as a fixed dextrine.

The following are its chief varieties :—

Arabine, soluble in cold water. Ex. Gum arabic. *fyr*

Bassorine, partly soluble. Ex. Gum tragacanth. *Al. Orch.*

Cerasine, insoluble part of cherry gum. *& plum*

Pectine, gelatinous. Ex. Pulpy fruits.

Mucilage, present in mallows and linseed.

RESINS contain much carbon and hydrogen, with little or no oxygen. They are found in plants, sometimes in a fluid state, called balsams and oleo-resins, and at other times solid. The most important solid resins are the copals, or those which partake of their character. These substances are insoluble in water, but soluble in spirits of turpentine.

ORGANIC ACIDS are produced by processes going on

Constituents of Plants

Elementary

~~of~~ O, H, N, C, P, S, Se, Cl, Br,
I, Fl, K, Na, Ca, Mg, Fe, Mn, Cu &c.

II Proximate Principles

1. ~~Non-nitrogenous~~ organic
~~Carbohydrates~~

Cellulose, lignine | Starch, inuline,
lechenine, dextrine,

Sugars - cane sugar, grape sugar
mannite

Gums - arabine, Basorine, Cerasine
Pectine, mucilage.

Resins - Balsams and oleo-resins
(liquid); Copals & (solid)

Acids - Acetic, Citric, Tartaric,
Malic, Tannic, Hydrocyanic
Oxalic, Gallic.

Volatile oils

~~Essences~~

Caoutchouc, Fixed oils

Wax

(See 6.13)

in living plants, and exist in vegetable juices, often combined with peculiar bases and alkaloids.

PECTOSE and other neutral compounds of carbon, hydrogen, and oxygen are also found in fruits and roots.

Acetic acid ($C_2H_4O_2$) occurs in plants as potassium acetate.

Citric acid ($C_6H_8O_7$) occurs in fruits of the orange tribe.

Tartaric acid ($C_4H_6O_6$) occurs in the juice of the grape.

Malic acid ($C_4H_6O_5$) occurs in the apple, gooseberry, &c.

Tannic acid ($C_{27}H_{12}O_{17}$) in oak and other barks.

Hydrocyanic acid (CyH where Cy is = CN) in cherry-laurel leaves.

Oxalic acid ($C_2H_2O_4$) with potash in sorrel, rhubarb, and other plants.

Gallic acid ($C_7H_6O_5$) in all kinds of galls by the conversion of tannic acid.

VOLATILE OILS. The chemical composition of these oils is extremely various. They are obtained by distillation from various aromatic plants. Oils from coniferous trees consist generally of carbon and hydrogen only. Otto of roses and oils from labiate plants contain oxygen in addition; whilst the oils of alliaceous plants contain sulphur also. Many volatile oils deposit *stearoptine*, which is nearly allied to camphor.

Camphor ($C_{15}H_{16}O$). This may be considered as a sort of solid volatile oil, less widely diffused, and only obtained commercially from two or three species of plants. It is light, inflammable, and evaporates on exposure to the air.

CAOUTCHOUC. A hydrocarbon, very widely diffused through plants, but more profusely developed in tropical regions. It is found associated with resin in the milky juices of many kinds of figs and euphorbias.

FIXED OILS contain carbon, hydrogen, and oxygen. When decomposed they yield fat—acids, and glycerine. Found most abundantly in the seeds of plants. They generally congeal at a lower temperature than the freezing-point of water, but some are solid at the ordinary temperature of the air. They all require a high temperature for their evaporation. Olive oil is found in the fruit.

WAX. A kind of fatty matter found in the fruits and stems of plants, sometimes covering the fruit with a "bloom," or appearing in scaly masses upon the stems. It is soluble in ether, insoluble in water, and combustible.

SECTION IV.

NITROGENOUS COMPOUNDS.

There are certain nitrogenous products found in plants, most commonly in greatest quantity in the seeds. These are termed flesh-formers or albuminoids, as they serve to build up the flesh of animals, and are closely related to *albumen*, the white of the egg. They consist of carbon, oxygen, hydrogen, and nitrogen, with some sulphur.

FIBRINE is the essential part of the gluten of wheat, the grey elastic mass which is left when dough is long washed in water.

Nitrogenous

Fibrine

Caseine or Legumine

Albumen

Emulsine

Diastase

The Alkaloids - viz

Quinine

Morphine

Solanine

Veratrine

Aconitine

Stychnine

Atropine

Piperine

Thine or Caffeine

(See P. 15)

CASEINE or *Legumine* is an essential part of the seeds of leguminous plants ; it is also the chief flesh-former in milk.

ALBUMEN occurs in a soluble form in the expressed juice of many plants : it coagulates or separates in flocks when liquids are heated to 60°—70° centigrade.

These three nitrogenous compounds are seldom found alone. In the juice of potatoes there is, besides fibrine and albumen, vegetable caseine ; and in leguminous seeds there is always, besides the caseine, a certain quantity of albumen.

EMULSINE is a nitrogenous compound found in oily seeds, as in almonds. In bitter almonds it is associated with *amygdaline*, and the two, when boiled with water, yield hydrocyanic acid, and oil of bitter almonds.

DIASTASE is a nitrogenized substance procured from malt, and developed during the germination of seeds. It is an albuminoid in an altered state, and has the power of converting starch into sugar.

The following table shows the proportions of digestible nitrogenous products and carbo-hydrates in 100 parts of the edible portions of certain plants :—

	Water.	Nitro- genous.	Carbo- hydrates.	Ashes.
Peas	14	23	52	2·5
Beans	14	25·5	46	3·5
Lentils	14	24	52	2·6
Oats	14	12	61	3
Barley	14	9	66	2·3
Potatoes	75	2	21	1
Turnips	89	1	9	1

ALKALOIDS are nitrogenous compounds found in plants, and generally constitute their active principles. They usually occur combined with organic acids.

- 1 *Quinine* ($C_{20}H_{24}O_2N_2$) exists in cinchona barks.
- 2 *Morphine* ($C_{17}H_{19}ON_3$) exists with others in opium.
- 3 *Solanine* ($C_{43}H_{71}NO_{16}$) is found in plants of the potato tribe.
- 4 *Veratrine* ($C_{32}H_{52}O_2N_2$) occurs in sebadilla.
- 5 *Aconitine* ($C_3H_{47}NO_7$) is found in monkshood and other species of aconite.
- 6 *Strychnine* ($C_{21}H_{22}N_2O_2$) is the active principle of *nux vomica*.
- 7 *Atropine* ($C_{17}H_{23}O_3N$) the active principle of belladonna.
- 8 *Piperine* ($C_{34}H_{58}O_6N_2$) is found in pepper.
- 9 *Theine* or *Caffeine* ($C_8H_{10}N_4O_2$), the principle of tea and coffee.

✓ **CHLOROPHYLL.** The green colouring matter of plants; generally occurs in a granular form, floating in the fluid contents of cells. It partakes somewhat of the character of wax, and is only developed under the influence of light, and in the presence of iron. *a.*

SECTION V.

MINERAL COMPOUNDS.

As a prelude to the mineral compounds, an important constituent of vegetable structures in their vital state must not be forgotten, *i. e.* water.

a. According to Frémy it is composed of two principle
(1) Phyllocyanin (blue) & (2) Phylloxanthin (yellow).
All ~~other~~ colouring matters of cells (except green,
are called Chromule. According to Frémy & Cloz
all the staining matters of flowers are ~~is~~ composed
of three principles (1) Cyanin (blue or rose) (2)
Xanthin (~~yellow~~) & (3) Xanthin (both yellow)

Mineral or Inorganic.

Water

Ammonia

Nitrates

Sulphates

Phosphates

Chlorides, Bromides &c

Metallic bases combined with
various organic Acids &c.

WATER (H_2O) is found in many aquatic plants to the extent of 90 per cent. Potatoes contain 75 per cent., turnips 90, and carrots 80 per cent. A sunflower consumes daily 22 ounces of water; consequently, if every plant occupies 4 square feet of soil, the plants of one acre would require 1,826,706 lb. during the four summer months. But the ground around the plants would also sustain grass and weeds, which require water to an equal extent at least. Therefore 3 million pounds of water would be required for an acre of sunflowers. It has been estimated that an acre of cabbages would require 5 million pounds, and an acre of hops 6 or 7 million pounds of water. Through the agency of this fluid, plants take up most of the elements of their composition. Much of this is absorbed by means of the roots.

The water and combustible portion of plants form by far the greatest proportion,—the mineral or incombustible being seldom present, even in their dried state, to the amount of 10 per cent.

The mineral matter, or ash left after incineration, varies in different plants, and in different parts of the same plant. In the dried leaves of the elm, more than 10 per cent. of mineral matter exists, while the wood affords less than 2 per cent.

In 1,000 lb. of the grain, and the same quantity of the straw of wheat, the following mineral substances are found :—

	Grain.	Straw.
Potash (K_2O)	5·5	4·9
Soda (Na_2O)	·6	1·2
Lime (CaO)	·6	2·6
Magnesia (MgO)	2·2	1·1
Silica (SiO_2)	·3	28·2
Sulphur trioxide	·4	1·2
Phosphorus pentoxide	8·2	2·3
	<hr/>	<hr/>
	17·8 lb.	41·5 lb.

These substances are variously combined, as sulphates, phosphates, and chlorides ; the silica, however, is mostly free.

The large proportion of silica in the above analyses shows its importance to plants of the grass tribe.

As all these fixed substances must be derived by plants from the soil in a state of solution, it becomes important to ascertain the quality of the soil, and its fitness to supply the requisite materials to the plants we would cultivate.

The importance of such a consideration is thus beautifully illustrated by Schleiden, the celebrated German botanist :—

“ The beautiful orchidaceous plant, the ladies’ slipper, grows over all parts of the Swiss Fore Alps, where the soil is formed of the alpine limestone ; it accompanies the whole Swabian muschelkalk, and disappears suddenly when we come to the sand of the Jura and Keuper formations on this side of the Danube. It next makes its appearance on the muschelkalk of Thuringia, and comes down with that on the Werra as far as the neighbourhood of Göttingen, then leaps over the Bunter sandstone of the Lower Eichsfeld, the granite of the Upper Hartz, and again gladdens the eye of the wanderer on the calcareous formations eastward of the Brocken. It is sought in vain over all the clay and sand formation of the northern German plains, till in the extreme north it again shows itself at Rügen, where the chalk rocks of Arkona and Stubbenkammer lift their heads.

“ On the western coast of France grow various insignificant-looking shore plants, species of *Salsola* and *Salicornia*, which the inhabitants there use to obtain soda from the ashes. When we travel from thence towards the east, we everywhere miss these little plants, even when searching most carefully, and

merely one or other of them makes its appearance in such places where the soil is moistened by some salt spring. At last we arrive at the Great Steppes of the south-east of Russia, which in summer are often covered with a thick crust of salt, showing themselves to be the ancient bottom of some dried-up sea, and here these plants are found growing in the same abundance and luxuriance as in the west of France. On the northern coast of Germany the little pale-red maiden pink grows upon the arid sand dunes, and is universally distributed over the sandy plains of northern Germany ; but these are succeeded by the granite, clay, slate, and gypsum of the Hartz, the porphyry and muschelkalk of Thuringia, and our little pink is not met with again till we arrive at the Keuper sand plains, on the further side of the Maine, surrounding the venerable city of Nuremberg. It extends farther south, through the Palatinate, till the muschelkalk of the Suabian Alps again sets a limit to it ; but it leaps over these and the whole alpine region, and at last appears on the sandy soils of northern Italy. How is it that these plants everywhere disdain the richest soils in their range of geographical distribution, and are confined to perfectly determinate geognostic formations ? Must not the lime, salt, and sand (or silex) have a most distinct influence in the matter ?”

Of the elements of plants, only oxygen and nitrogen occur free or uncombined in them. The other elements are variously combined together : some of these compounds are here given :—

Water, H_2O .

Carbon dioxide = carbonic acid gas, CO_2 .

Ammonia with various acids.

- Nitrates, such as potassium and calcium nitrates.
 Sulphates, such as potassium and calcium sulphates.
 Phosphates, such as calcium and magnesium phosphates.
 Chlorides, such as sodium and potassium chlorides.
 Oxalates,
 Citrates,
 Tartrates,
 Acetates, } as those of calcium and potassium.

Raphides (Gr. needles) are crystals, or clusters of crystals, commonly of calcium oxalate, but also sometimes of the phosphate, tartrate, sulphate, or carbonate. They are found generally in cells (Fig. 2), and occasionally in great numbers. Turkey rhubarb root



Fig. 2. contains above 30 per cent. *Hyalanth - Cactus dit*
2. Equisetum contains silica - dit
 C. *En crista h. c. g.*
 (1) *Acicular Raphides.*
 (2) *Conglomerate Raphides.*
 SECTION VI.
Cytoplasts = a mass of crystals hanging in a cell - dit.
 CELLS. *Inticacids, Monas etc.*

The simplest form of vegetable life, and the most elementary phase from which all others are developed, is that of a minute vesicle, or cell. *Ex. Botrococcus or redonnis, Oscillatoria*



CELLS are very small transparent bladders, or cysts, with walls, or boundaries, of cellulose, and generally contain fluids.

An egg may be accepted as an illustration of a cell, of which the shell represents the cell-wall.

The normal form of the cell is that of a sphere, or spheroid, having a length not much exceeding its

Carbonate of lime - calcareous.
 Sulphate of lime - siliceous.
 carbonate of Aluminobromine, Alkali - Alkaline
 Phosphate - cellular.

Cells

The ^{regular} form of cells is modified by (1) ^{irregular} pressure (2) ^{irregular} unequal stretching in different parts.

Forms (1) Spherical (2) Elliptical (3) Polygonal generally dodecahedral with ^(figs 5 or 6) hexagonal sections (4) Stellate (fig 7) (5) Cylindrical (fig 11) (6) Funiform (fig 12) (7) Fibrilliform

In the cellular plants (Algae) (2) pollen (3) spores
 the cells are Spherical or modified into cylindrical, oblong & c or show projections in their surface called papillae, or spines, of various kinds - The cells of hairs & c also are more or less unstrained in development.



Cellular Tissue or parenchyma Short cells thin walled long cells.

- (1) Round or elliptical. (Fig 3) Ex. Parenchyma. Ex. Succulent plants, pulpy part of leaves of fern.
- (2) Spongiform (Fig 7). Ex. Rush, under surface of leaves various
- (3) Regular or complete (Fig 8 & 9). Ex. Pith of Dicotyledons.
- (4) Elongated (Fig 12) (cylindrical or fusiform). Algae & stems of Monocots.
- (5) Tabular (t) (a) uniform (m) Epidermis (t)

(6) Fella Contorta or interlacing fibrilliform tissue Ex. Fungus (2)

(7) Fibro-cellular Tissue (Mistletoe, Orchid - Spine)
 Colenchyma - the cells thickened by starch deposits (Algae)
 Sclerenchyma - the cells thickened by lignin - Stones of fruit

Cell Contents

~~Mid-Brain~~

- Protoplasm or primordial utricle etc see page (28)

breadth (*Fig. 3*). An aggregation of these cells, or vesicles, constitutes the *cellular tissue* of plants.

Were the cell-walls to retain their original form when compacted together, there would be numerous spaces left unoccupied between the points of attachment; these might be termed *intercellular spaces* or *lacunæ* (*lacuna*, Lat. a little hole).

Slight pressure causes the cells to accommodate themselves to the space they have to occupy, and they assume that form which shall best fill up the interstices and occupy the least room, namely, a modification of the dodecahedral with an hexagonal section (*Fig. 5*). Hence, this is the most common form in the cellular portions of plants.

The simplest plants, as algæ and fungi (*Fig. 4*), consist entirely of cellular tissue, and are hence called *Cellulares*.

The pith of trees, rushes (*Fig. 7*), and the rice-paper of the Chinese, are cellular, as are also all plants in their earliest stages.

Parenchyma (Gr. for the substance of the lungs, liver, &c.) is the name generally given to cellular tissue ~~which exhibits hexagonal cells when cut across.~~



Fig. 4.




Fig. 6.




Fig. 5.




Fig. 7

When a spiral line is coiled up in the interior of

cells, it is called *Fibro-cellular tissue* (*Fig. 8*). This is a very common form.

The spiral is sometimes broken up, so that the thread appears in rings (*Fig. 9*), and, when still more broken, in bars, or portions of rings, or elongated dots (*Fig. 10*).



Figs. 8.

9.

10.

The spiral thread is deposited within upon the cell-wall subsequent to the formation of the cell itself. It sometimes becomes free, and at others remains attached.

The spiral threads and movable fibres which are found in the antheridia of mosses, and elsewhere in cryptogamic plants, and which exhibit peculiar movements, so as at one time to have been considered as animalculæ, are doubtless of this nature and origin.

The microscope is necessary for the examination of all cell-structures.

The membrane of the cell-wall permits the circulation of fluids through it, although pores are not often visible.

A kind of cellular tissue consisting of short cylindrical cells, is called *Pitted tissue*. These may originally have been spheroidal cells, the walls of which have become ruptured in one direction so as to assume a cylindrical character (*Fig. 11*).



Fig. 11.

The walls of these tubes are pitted or dotted, in consequence of the unequal deposit in their interior. It is a very common form of tissue in wood, and conveys fluids with facility in the direction of its length.

As all forms of tissue are derived from the simple cell, of which they are but subsequent modifications, intermediate forms will be met with of all the varieties, and the line of demar-

cation can scarcely be fixed, the division or

Modifications of cells by ligninous deposit

1. Dotted or pitted - Ex. Sphagnum, Mistletoe, (Fig 10)
2. Spiral - Orchid, ~~Mistletoe~~ (Fig 8) Seed of *Clonaria*, Cycas Rev. etc.
3. Annular - Ex. Cactus (Fig 9) 4. Reticulated - Ex. ~~Mistletoe~~

II Pseudo-sclerenchyma

When elongated cells have thick walls ~~thickened~~ by ligninous deposits their combination forms the following kind of tissues which are all called Pseudo-sclerenchyma

I. Pseudo-sclerenchyma (fusiform, uniform)

- (a) Woody tissue - Ex. Wood of trees
- (b) Woody tissue of the ~~stem~~ (very long) [vast bundles of Monocots & dicots outside the stem of Monocots & in stems of many dicots]
- * Liber or bast tissues - Ex. Commonly in the inner bark of dicots, hemp, flax, New Zealand flax, Pita flax, Seem, Jute, China grass

II Disc bearing woody tissues

The glandular tissue - Coniferous & other gymnosperms

Sperms ~~and~~ plants [discs opposite - Pine
- alternate - Spruce, Abies]

In oaks & fir the ~~cells~~ ^{cells} contain spirals as well in addition to ^{to} ~~the~~ ^{the} ~~discs~~ ^{discs} ~~in~~ ⁱⁿ ~~the~~ ^{the} ~~Star anise~~ ^{Star anise} & the discs have no central pits.

- * 1. ~~Hemp~~ Hemp - *Cannabis Sativa* [Cannabis]
- 2. New Zealand Flax - *Phormium Tenax* [Phormium]
- 3. Pita flax - *Agave Americana* [Agave]
- 4. Pine - *Pinus* [Pinus]

Less (from Loggia unearns)

Vascular Tissue

~~Angienchyma
Fusiform vessels or Vascular tissue~~

1. Composed of Fusiform vessels - Pluranchyma or Bothranchyma
2. Pitted or dotted vessels - Bothranchyma or Zaphranchyma
3. Spiral vessel - Zoanchyma

- 1) Pluranchyma - (fusiform & uniform) - S.Wood
- 2) Bothranchyma or Zaphranchyma } - (Pitted or dotted)
- 3) Zoanchyma (spiral)
- 4) Glandular or Dispersing
- 5) Laticiferous
- 6) Scalariform

Kind of vessels of Tissues formed by them -

1. Pitted or dotted (Bothranchyma or Zaphranchyma) - Wood of Dicotyledons (Spiral fibres in the dotted vessels of stem & xylem)
2. Spiral vessels (Zoanchyma) Medullary Sheath of Dicty. Vasc. bundles of Marantaceae (Pithecolobium) of some higher Polyp. Simple - only one fibre of Claves.
Compound - many fibres - ex. stem of Banana, young shoot of Achras, & in the Pithecolobium plant.

classification under different names being more artificial than natural, serving to mark phases of the same tissue, rather than to be designations of different elementary organs.

The terms *cellular* and *vascular* tissue are still employed, with this limitation of the meaning and value attached to them.

SECTION VII

VASCULAR TISSUE.

Under the general name of *Vascular tissue* is included the form of tissue which, with greater propriety, is termed *woody tissue*, or *pleurenchyma* (*pleura*, Gr. a rib), and *fibro-vascular* tissue.

WOODY TISSUE consists of elongated tubes of a fusiform or spindle shape, tapering at each end. It is cylindrical, long, fine, and tough, characteristics which distinguish it from cellular tissue.

It is found in the wood, inner bark, and veins of leaves.

The walls of this kind of tissue become thickened by successive layers of lignine, and when cut in section, the concentric lines or markings of the successive deposits may often be traced (*Fig. 12*).

The strength of this kind of tissue renders it extremely valuable in the arts, when divested by maceration of all extraneous matters. In this way hemp, flax, China grass, Manila hemp, &c. are prepared. The distinct forms of woody tissue in linen,



Fig. 12.

and cellular tissue in cotton, are easily detected under the microscope.

Glandular woody tissue is a peculiar variety of *pleurenchyma*, occurring in coniferous and some few other plants.

The sides of the tubes are furnished with circular disks, sometimes in single, and at others in double or triple rows. These disks or depressions have small orifices in their centre (*Figs. 13, 14*).



A recognition of this kind of tissue in coal determines its coniferous or cycadaceous origin.




Fig. 14.

VASCULAR, or Fibro-
Fig. 13. vascular tissue, consists of thin cylindrical tubes, with fibre spirally coiled up within them. It has also been called *Trachenchyma* (*trachea*, Lat. windpipe).

Spiral vessels are the type of this kind of tissue. These vessels, which are fusiform (spindle-shaped), overlap at the ends, where, the intervening membrane being absorbed, they communicate freely with each other. They are found in the medullary sheath, veins of leaves, &c.



Fig. 15.

The fibre in the interior may be unrolled, and is sometimes single; and at others several fibres are united together, side by side, and may be unrolled like a ribbon.

When the spiral fibre is broken up, or does not make a complete coil in the tube, a modification of the above kind of tissue results, called *ducts* or *false tracheæ*, to which special names are also given, according to the phases it assumes.

When the fibre occurs in rings, it is called *annular*.

When the tubes are prismatic, and the fibre is

Vessels

Form or kind of Vessels

1. Pitted or dotted ex. Wood of Dicotyledons
2. Spiral - (Stem of Banana (Cannaceae), young shoot of Asparagus, ^{ex. Mollusca, shell, Vasc. bundle of Monocots, of some other plants})
3. Annular -
4. Reticulated -
5. Scalariform -
6. Laticiferous -

Vessels & vascular tissue

1. Spiral vessels (Tracheidyma)

The spirals may be easily uncoiled by pulling asunder (1) the leaves of Banana, Hyacinth or (2) the young shoot of leaf buds of Geranium, Rose, Strawberry etc -

3. Annular vessels - Stems of soft rapidly growing Dicotyledons, of Monocotyledons, & in Pteridoms - especially in the Equisetaceae (Pteridium) Melon.

1. Pith plant (Comp) - Fibres seen - leaves of Banana, Hyacinth, Geranium - ~~can~~ can be uncoiled by pulling the leaves gently asunder.

(4) Reticulated vessels forming a network of spirals) similar situations with annular - Ex. Melon.

(5) Scaliform vessels - Ex. Vine, (Cylindrical) ferns frismatic, lycopods.

4, 5, are but modifications of true Sieve & Radial

(6) Laticiferous - Dandelion, Celadon, Castor (Galaxiandra vitile), India rubber (Ficus elastica), Urtica elastica, Siphonia Brasiliana, gutta percha (Sonchus oleraceus) Liber & pith of Xylem, Veins of leaves (the

is also a kind of milky juice) Euphorbia & Secundum of Monocotyledons & higher angiosperms.

These form tissues called Leucodermis on account of the movements that are observed in the milk - Diastole & Anastole by their walls & anastomosis.

broken and arranged at equal distances above each other, like the steps of a ladder, it is termed *scalariform* (*scale*, Lat. a ladder), as in ferns (*Fig. 16*).

Sometimes the arrangement is very irregular.

These ducts seem to convey fluids.

Laticiferous vessels, or *Oinenchyma* (*kineo*, Gr. I move), are branched tubes, which unite and run into each other in a very irregular manner, forming a complete network (*Fig. 17*).




Fig. 17.




Fig. 16.

The walls are not marked. It commonly occurs in the liber of Exogens.

These vessels convey a fluid called *latex*, which is either transparent or coloured. It has of late been suspected to be of a formative character.

SECTION VIII.

CELL DEVELOPMENT.

The history of the development of cells is still confused. Much as has been written, the entire question is one of considerable uncertainty.

Schleiden has referred the origin of cells to the formation of a central point amongst the formative fluid, around which a congregation of atoms takes place till a firm layer is formed; this becomes ultimately the cell-wall, which enlarges, assimilates fluid, and becomes a cell, containing within it the *nucleus*, or, as he terms it, *cytoblast* (*kutos*, Gr. a cell; *blastos*, a germ).

Cells have generally, when full grown, a thickened disk, or *nucleus*, of this kind, attached to their sides, and these sometimes contain other bodies, called *nucleoli* (Fig. 18).



Fig. 18.

New cells may be formed on the outside of old cells, as branches; this process has been termed *gemmation*, or budding. This fact has been proved in the growth of sea-weeds (Fig. 19).



Fig. 19.

New cells may also be produced from old cells by *fissuration*, or internal subdivision.

A division, or septum, separates the old cell into two portions, each of which becomes a complete cell. (This is shown in Fig. 20.)




Fig. 20.

Another theory assumes that the *nucleus* separates into parts, each forming the new nucleus, around each of which a new cell-wall is developed, and ultimately the old cell-wall is ruptured and disappears.

These methods may be characterized as :—

- 1st. *Isolated* formation in mucus.
- 2nd. *Gemmation*, or *Exogenous* formation.
- 3rd. *Fissuration*, or by division of cells.
- 4th. *Endogenous*, or by division of nucleus.

Cells increase with immense rapidity, especially in some fungi. Dr. Lindley estimates that in *Bovista gigantea*, the great puff-ball, cells are developed at the rate of 66,000,000 per minute.

Spiral threads owe their origin to internal deposits in cells. It is not improbable that gum, derived from the transformation of starch, is the chief agent.

Truncated or tubular cells appear to be derived from a congeries of cells, placed in apposition, and

Cell development

1. Gemmatem budding — ex. Sea weeds
2. Fission —

oil - In the rind of orange, lemon; leaves of orange,
Mercury, Myrtle etc

Wax - Wax palm.

~~Raphides~~ Air - In the cells of Pongrood

Raphides - Reticular - Hyacinth, Rumex
Conglomerate - Beetroot.

Epidermis

(1) Epidermis - Consists of a layer or layers of
compressed cells (tabular) of varying outline -

Hexagonal - Iris & many other Monocotyledons

Trigonal - Zigzag - chairs, Simon - chairs

Polygonal - Square Rhomboid etc.

under circumstances of pressure or circulation favourable to the final obliteration of their walls in the direction of their attachment.

Similarly may the laticiferous tubes have originated in part from such truncated cells, which have at the same time become branched and anastomosed, so that all trace of their origin is obliterated.

The *Contents* of cells are—

The *primordial utricle*, or *protoplasm*, which is the deposited lining of the cell-wall.

Deposits of *sclerogen* (makes hard), in layers, give great solidity to some cells. *Ex.* Vegetable ivory.

Water, generally containing other matters in solution.

Jelly, which is perhaps the same as *pectin*, the mucilage which abounds in some algæ.

Starch, *chlorophyll*, *chromule* (colouring matter other than green), *wax*, *oil*, *camphor*, *resin*, and *raphides*.

Other compounds are found deposited in the *vascular* system.

SECTION IX.

THE EPIDERMIS.

The external cellular covering of plants, or epidermal layers, may be readily detached from young leaves and stems for examination.

—The *Epidermis* (*epi*, Gr. upon; *derma*, skin) is extended over all parts of plants exposed to the atmosphere, except the stigma.

It usually consists of a layer or layers of compressed cells, having




Fig. 27.

their walls bounded by waved (*Fig. 22*) or straight (*Fig. 21*) lines, and without intercellular spaces. These cells are usually colourless, and with fluid contents. (a)

22 A superficial pellicle is spread over and upon the epidermis, and is considered by some as a secretion of the epidermal cells below it. *Very thick in the upper surface of the stem.* An imperfectly formed epidermis, called *Epiblema* (*epi*, Gr. upon; *blema*, a wound), supplies the place of the epidermis in submerged plants, and on the extremities of growing roots.

The name of *Epithelium* (*epi*, Gr. upon; *thēlus*, tender) has been given to a very thin cuticular covering found lining the ovary and other internal parts of the plant. *a canal of the style in many flowering plants.*

There are openings of a peculiar character between some of the cells of the epidermis, on parts exposed to the air. These are called *Stomata* (Gr. mouths). *Breathe*




Fig. 22.

They are usually formed by two semilunar cells, the ends of which unite around an oval orifice (*Fig. 22*). *The cells open & close.*

The *stomata* communicate with the intercellular passages, and perform some of the functions of mouths to the plant, from their fancied resemblance

to which the name arose. They are scattered on the surface of the epidermis, sometimes with regularity, but generally without any apparent order.

Stomata occur on the green parts of plants, on the leaves and their appendages principally. They are not found either on cellular or aquatic plants, *roots, ribs of*

On leaves they generally occupy the *under surface*.

(a) In the Thallophytes (Fungi, Algae, lichens etc) there's no Epidermis but they have a cuticle - There is Siliceous matter in the ~~cuticle~~^{epidermis} of Equisetum - Carbonate of lime in Chara

Epiblema - ~~outer~~ ^{submerged} plants (2) admixture of growing roots -

Epidermis

Spiral cells in the Epidermis of aerial roots of Orchids.
Thick - Oblander, American Aloe

Stomata

Usually oval or elliptical - Rounded in Rumex; ^{composed of 2 cells}
quadrangular in Amer. Aloe, Yucca glauca - In Nerium (Oblander) the small stomata are on the pits or depressions on the underside of the leaf (together with hairs)

of 4 cells either in 2 pairs (e.g. Ficus Elastica) or arranged in a circular manner (e.g. Proteaceae), of 3 or more tiers of cells (e.g. Marchantia - Hepaticae)

Stomata
Stomata found in reptiles of Cruciferae, on side of Walnut.

(Quadrangular Stomata in - American Aloe, Yucca glauca
Rounded Stomata in - Rumex Heliosa.)

Stomata are more abundant in woody than in herbaceous plants; ^{more} in leathery leaves than in those of thinner texture, least in succulent leaves - On the dull surface of a leaf more than on the glossy leaf

In clauder there are pits lined with hairs & Very small ~~pits~~ stomata (Hair-like stomata Balfour)



Stomata occur in the joint of the wall flower & in the seed of the Walnut.

Hairs

(a) ~~Simple~~ Unicellular Cabbage, Draba (pink
~~simple~~ - cabbage Alyssum (Stellate,
 Forked - Draba or Whitlow (Cruciferae)
 grass

Stellate - Alyssum (Cruciferae)

(b) ~~Unicellular~~ Unicellular
~~simple~~ Simple -

Urtica galepa 2 Moniliform -  Su. - Urtica of Urtica
 (Urtica galepa)
Tridacantha 3 Partitoid -  Tridacantha in Urtica
 (Virginia Urtica)

4 Capitate (Fig 26)

5. Incurvate or Hooked (27)

6 - Glochidiate or Barbed (Fig 28)

Long

7 Stellate - Draba (Fig 24)

Malpighia Urus 8 Peltate - Malpighia Urus - (Fig 29)

Spiral fibres in rows of:

(1) outer coat of Scin of Neanthidium

(2) Fruit of Salvia Homiumum

(7) Scabrous - In Chaetis (Elaeagnus), Boop
Ramenta or Ramentaceous hairs = 2 Callos of Sci
Scabrous - Ramenta

of such as are freely exposed to the air, with but few on the upper. In floating leaves the stomata occupy the *upper* surface; and on leaves which grow vertically, an equal number of stomata are found on *both* sides.

The leaves of the mistletoe have been found to possess 200 stomata to the square inch on both sides, the clove pink 38,500 on both sides, and the lilac 160,000 to the square inch on the under side and but few on the upper.

Stomata open in a moist state of the atmosphere, and close when dry.

Their functions appear to be the regulation of the transpiration, and through them the liberated gaseous bodies escape.

There are also conditions of, or appendages to, the epidermis, to which various terms are applied. It may be *naked*, *glabrous* (smooth), *silky*, *pubescent* (downy), *hairy*, *shaggy*, *velvety*, *woolly*, *glandular*, *punctated* (dotted), *scabrous* (rough), *rugose* (wrinkled), *reticulated* (netted), *verruose* (warted), and *papillose*.

Of the appendages the most important are *Hairs*. They are composed of one or more cells, and vary much both in position and structure. Sometimes they are simple (*Fig. 23*), forked, divided, or branched, and consist of a single cell. At others they are of the same forms, and consist of various cells placed in apposition.

Several hairs, radiating from a common centre, assume a starlike or *stellate* form in the mallow tribe and *Cruciferae* (*Fig. 24*).

The hairs investing the seeds of the cotton plant are the most valuable in a commercial sense of any of the appendages of the epidermis.



Fig. 23.



Fig. 24.

Silk cotton consists of hairs from the seed-vessels of several species of *Bombax*, *Eriodendron*, *Asclepias*, &c.

Clavate (*clava*, a club) hairs gradually expand from the base to the apex (Fig. 25).

Capitate (*caput*, a head) hairs have a distinct round head (Fig. 26).

Uncinate (*uncus*, a hook), are hooked hairs (Fig. 27).

Barbed hairs have one or more hooks around the apex (Fig. 28).

Peltate (*pelta*, a buckler) hairs are attached at the centre (Fig. 29).

The hairs of *Drosera*, or sundew, have spiral fibres coiled in the interior.

Prickles are hardened hairs, and are only attached to the epidermis. *Ex. Rose. Bramble.*

Setæ are bristles or stiff hairs formed of a-

Deciduous hairs, on falling away, often leave depressions, or openings, in the epidermis, which have been called *false stomata*.

Glandular hairs are such as contain secretions, either in distended cells, at their apex or their base. Of this character are the receptacles of the fragrant volatile oil which gives its odour to the sweet-briar, and also the stinging hairs of the nettle (Fig. 30).

Root-hairs are prolongations of the superficial cells of the newest portions of roots, which they bring into close contact with the particles of soil. They die away when the epiblema is formed by the thickening of the superficial cells.



Fig. 25.



Fig. 27.




Fig. 29.



Fig. 28.



Fig. 26.


G. Stoma
2.

Fig. 30.

Hairs may be trans formations of

- (1) Flower stalks - Plum cotinoid, Wig tree
- (2) Calyx - Dandelion & other composite plants.

Balls of hairs secrete colouring matter in Pris, Virginia spider wort.

Bristles - are rigid hairs formed of a single cell (Sclero). Sitosa or Setae - with bristles

Sclero - are hairs united by a membrane &c. oleose

• Glands Secretory are ex. oleaster, sunsp
Movements seen in Alant ap. -

- (1) oily - in flower stalks of Dillany rose
- (2) Waxy - in Rosulelia (of beautiful green colour)
- (3) Resinous
- (4) Gummy - stipules of Cinchona
- (5) Saccharine
- (6) saline
- (7) Acid fluid - Chickpea (flower stalks)
- (8) Irritant

Sensitive hairs in Drosera; Grossa

Honey glands - in petals of Fritellaria imperialis

Cellular glands - in the Cinchona tribo Althia, hairs - in Sampanda

Islands

(A) External
a. Sacked

1. Unicellular - Salvia, Antanthinum may
 2. Multicellular - ~~Antanthinum~~ no,
- b. Sepals - Helianthus, Begonia.
- Papillae (Soft) - Plant
water vascular (hard)
- Stings - Contain irritant

(B) Internal - Nettle, Malpighia laevis
no, orange
Nectaris - Nettle cups, Trillium, Impatiens (Cm
(Dennis) 1946)

Duckweed, Pistia contain a peculiar kind of cellular sheath called Plecochiza.

Hairs in the roots of Orchids,
aerial roots of Gony, grasses as mites



Dicotyledons are Exorhizal -
Monocotyledons - Endorhizal } Richer -
Acotyledons - Heterorhizal -

Roots of these avoid light & air & are
destitute of the green colour of leaves & stems (3)
bear no leaves (4) their growing points are
protected by sheaths called calyptrae

SECTION X.

THE DESCENDING AXIS.

From the surface of the soil the plant-axis ordinarily extends upwards and downwards; in the former direction being called the *stem*, in the latter the *root*.

As the ascending axis divides and branches, so does the descending axis, or root; but, whilst the stem produces its branches through the medium of a leaf-bud, the root is without leaf-buds; neither is there a true epidermis, stomata, or a distinct pith.

The extremities of the roots are covered with loose cellular layers, termed *spongioles*. (2)

Roots continue to grow by their extremities, which are ever removing further from the central axis and penetrating new soil, the less recent portion becoming part of the non-absorbing root, as the new extremity performs the function of absorption.

The internal structure of the root is similar to that of the stem of which it is an extension, except that the pith and medullary sheath are absent. 6.

The point of union between the ascending and descending axis is termed the *collum*, or neck. It is sometimes marked by a distinct swelling at the base of the stem.

One class of roots consists of a simple extension of the axis downwards, which may become afterwards more or less branched. Another class consists of a number of rootlets proceeding from one point of union or attachment to the stem.

When a simple or tap root is large and succulent, ^{primary} the roots of *Dicotyledons* consist of the direct prolongation of the radicle - these called *Erwahirab*.

and tapers gradually towards its extremity, it is called a *conical* root (*Fig. 31*). *Ex.* Carrot.

But if it first enlarges, and afterwards diminishes, so as to become spindle-shaped (*Fig. 32*), it is termed *fusiform* (*fusus*, Lat. a spindle). *Ex.* Radish.

When a root expands still more in a lateral direction, becoming globular, or more like a boy's top (*Fig. 33*), it is *napiform* (*napus*, Lat. a turnip). *Ex.* Turnip.

If instead of a single axis there are a number of rootlets (*Fig. 34*) proceeding from, or near, a common point of attachment, and these are all threadlike in form, it is called a *fibrous* root. *Ex.* Grasses.

When some of these rootlets become enlarged and succulent (*Fig. 35*), assuming an ovate form, a *tubercular* root is formed. (This must not be confounded with similar swellings in an underground stem called *tubers*.) *Ex.* Orchis.

When the swellings are numerous, and succeed each other so as to present the appearance of a string of beads (*Fig. 36*), it is called a *moniliform* root (*monile*, Lat. a necklace).

If, instead of being separated, the



Fig. 31.



Fig. 32.




Fig. 33.



Fig. 34.




Fig. 35.

All roots other than the mere prolongation of the radicle (which is called a tap root) are adventitious.

Aerial roots are a variety of adventitious roots
ex - Banana, Mangrove, Screw pine, Ivy etc.

Epiphytes - have aerial roots which absorb nourishment from the air - ex. ~~Mistletoe~~ orchids -

Parasites - penetrate their roots (adventitious) into the substance of their host - ex Mistletoe, Booroom
• rape, Rafflesia etc.

As to duration roots are (1) Annual (2) Bicennial or (3) perennial

Modifications of roots

(A) of Taproot -

- (1) Conical - Carrot, Monkshood
- (2) Fusiform - Radish -
- (3) Napi form - Turnip



(4) Placental form

(5) Contorted in position.

(6) Abruptor praemorse.

ex. Sambucus

(Devilbit Scabrous)

(B) of Adventitious roots

- (1) Fibrous - ex. grasses, Bulbous plants.
- (2) Tubercular - ex. Orchis or Orchis ^{palmated - tubercly divided} ~~excess~~ - ex. Orchis ^{maculata}.
- (3) Moheliform - ex. Sclerogomum ^{triste}.
- (4) Annulated - ex. Ipe ^{caluan}.
- (5) Coralline - Coralloid ^{inmate}.

(*Spiraea filifundula*)

1) Nodulose - Dropwort, succulent at extrem

2) Fasciculated - Dahlia. ... roots collected together.

Leontodon viduus-avis, *Dryopteris*

Pracnense, tuberc - ...

Twisted

Root like process of ... are supports of plant ... adventitious roots - Fig tree, pine tree.

Parasites - are plants which send roots into other plant.

Epiphytic - Do not send roots into the supporting plant it does not draw nourishment from it.

Parasitic device - nourishment - Dr. Dodd's ... *Rafflesia*.

orchid, Vine = Epiphyte.

Dodder, *Mistletoe* - stem parasite

Rafflesia - Root parasite.

Fully formed *Laqueum* root has no pith & medullary sheath (pith is developed in some *Dicranum* in the root of Walnut, hazel & chestnut).

The roots of *Phemelia*, *Vignea japonica*, *elloula*

Pracny & *Japan Pracnense* have the process forming adventitious buds.

The Dodder is at first parasitic, after it is *Mistletoe* (viscous album) complete parasite. It is parasitic the whole life.




Fig. 36.

swellings are compressed together so as to appear as rings around the root (Fig. 37), it is *annulated* (*annulus*, Lat. a ring). *Ex.* *Ipecacuanha*.

When some of the rootlets become succulent and enlarge towards their extremities (Fig. 38), a *nodulose* root is formed (*nodus*, Lat. a knot). *Ex.* Dropwort.



Fig. 37.



Fig. 38.

When all become succulent and enlarged, so as to resemble a bundle of fusiform rootlets (Fig. 39), the root is said to be *fasciculated* (*fasciculus*, Lat. a little bundle). *Ex.* *Dahlia*.




Fig. 39.

Roots that last only one year are *annual*; if two years, *biennial*; but if longer, *perennial*.

The functions of the root consist not only in fixing the plant firmly to the earth, but also in obtaining nutriment from the soil, and often in storing up organizable matter for its support.

When two fluids of different densities are separated by a permeable medium, two currents will be established—one outwards, called *exosmose* (*exo*, Gr. outwards; *osmos*, impulsion), and one inwards, called *endosmose* (*endon*, Gr. within), which will continue

until the two fluids are of equal density. By means of such currents vegetative action is carried on. The fluid contents of roots are of greater density than the moisture of the soil; an *endosmose* current is therefore established, for which the root-hairs and the newest part of the root protected by the *spongioles* are the medium, accompanied also by a much smaller outward or *exosmose* current. Evaporation continually going on in the exposed portions of the plant, the contained fluids are never reduced to the density of the medium in which they grow; hence the inward current, or absorption, does not cease while the vital action is unimpaired.

Adventitious roots are formed in some plants for the purpose of affording additional support to the stem.

In the *ivy* they serve the office of tendrils. From the branches of the *banyan* they are produced downwards to the soil, in which they fix themselves, and develop true roots.

In the *screw-pine* and *mangrove* they are developed in a similar manner from the lower parts of the stem, and become rooted in the soil.

SECTION XI.

ASCENDING AXIS.—EXOGENS.

A typical plant may be characterized as consisting of *stem* and *leaves*.

The *stem* is the axis of growth, and has, ordinarily, a portion above ground, which may be considered as the *ascending axis*, and a subterranean portion, which is the *descending axis*. All branches, whether of the ascending or descending axis, are portions or offsets of the stem.

Stem - (1) Ascends sucking light & air (2) bears
leaf buds (3) is coloured more or less green.

~~And produce~~

Trees - plants producing permanent woody stems
& not less than five times the height of a man.

Shrubs - less than five times the height of a man.

Shrubs of small stature are called bushes or under
shrubs.

Caulis = herbaceous stem

Culm = stalk of grasses.

Caudex & stipe = stems of palms & tree ferns -
anomalous stems

(1) Tortoise or elephant's foot

(2) globular - Melon cactus, jointed, columnar, or angular
in other cactuses -

(3) Pseudoball - oval or rounded stem of orchids -

Parts & Composition of Progenous stem

(1) Pith (or Medulla) - Not central - composed of
(a) hexagonal cells & (2) laticiferous vessels -

(2) Medullary sheath - composed of spiral vessels

(3) Wood -

All the parts of a plant are referable to one of these typical forms, *i. e.*, stem and leaf.

The *ascending axis*, or, as commonly called, the *stem*, in its earlier stages consists almost entirely of cellular tissue. Fibrous or woody portions are afterwards developed. Some plants are entirely devoid of this axis, and consist merely of cellular expansions, called a *thallus* (*thallos*, Gr. a frond). These plants are denominated *thallogens*.

Provision is made for a regular and systematic development of leaves and branches around the axis. The points from whence leaves and buds issue are termed *nodes* (*nodus*, Lat. a knot), and the spaces between them *internodes*.

Sometimes the internodes are so much shortened that the nodes are clustered together just above the surface of the ground, and the leaves appear in the form of a rosette, and without any evident stem. *Ex.* Primrose.

The internal structure and method of increase in stems has formed the basis of a division of them into three great groups or classes, *Exogens*, *Endogens*, and *Acrogens*.

Exogens (*exo*, Gr. outward; *gennao*, to produce), to which class our forest trees belong, are characterized by the possession of a separable bark (2) wood arranged in concentric layers, the youngest at the circumference, (3) a central pith, and (4) medullary rays proceeding from the pith to the bark (*Fig. 40*).

The *pith*, or *medulla* (Lat. marrow), is composed of cellular tissue, the cells being




Fig. 40.

often hexagonal. In the young plant it occupies a large proportion of the stem. In some plants it is more largely developed than in others. *Ex.* Elder.

At first the pith is greenish and succulent, but afterwards becomes pale and dry. It does not increase in growth after the first year.

The rice-paper of the Chinese and the shola pith of India are the medulla of certain plants.

Medullary sheath. This immediately surrounds the pith, and consists principally of spiral vessels. Prolongations from the pith pass between these vessels into, and intersect the woody layers (medullary rays).

Wood is deposited annually in layers, or circles, upon that of the previous year. A transverse section of an exogenous stem shows these concentric zones, or rings of wood, each of which is generally the produce of a single year. Hence the ages of trees have been calculated from the number of these concentric rings. In some trees the circles are marked much more distinctly than in others. The external layers are the youngest. The internal, or oldest, occasionally exhibit a distinct difference in colour. *Ex.* Lignum vitæ.

The concentric rings are not always developed equally all around the tree; so that the pith is not always in the centre. On that side of the stem which has, during growth, been most exposed to the light, the woody layers will be found thicker, and the entire rings of some years will give evidence of a more vigorous growth in a wider circle for that year.

Duramen (*durus*, Lat. hard). The interior, or *heart-wood*, called also *duramen*, from its greater hardness, is black in the ebony, and brown in coccus-wood, while the external layers of *alburnum* (*albus*, Lat. white) or *upwood*, are pale-coloured, soft, and permeable.

Cambium. Between the external layers of wood and the bark, a copious semifluid mucilage is found in

Part of an Injurious Stem & their composition

- (1) Pith (Medulla) Composed of (1) Hexagonal cells (2) of laticiferous vessels.
- (2) Medullary sheath - Composed of spiral vessels
- (3) Wood - Composed of pleuruchyma & Bostrychyma - ~~Some Annular & reticulated vessels~~ occasionally some Annular & reticulated vessels.
In Conifers no Bostrychyma, but composed of disobeying
- (4) Cambium Layer - Cells pleuruchyma only.
- (5) Medullary rays - Composed of muriform parenchyma (very long fibres - Arctostichia or Menispermata and small - fern, fish)
- (6) Bark
 - (a) Endophloem - Composed of tabular cells (2) laticiferous vessels | interspaces between libra cells - laquea lenticular shear
 - (b) Mesophloem or green layer - Composed of (1) polyhedral cells (2) laticiferous
 - (c) Epi-phloem or suberous or corky layer composed of tabular cells (tabular parenchyma)
- (7) Epidermis - Formed of compressed cells.
- (8) Article - Membranous secretion of the cells of the epidermis

Mesophloem - cells contain chlorophyll

Epiphloem - gives stem to young barks, green in colour owing to chlorophyll (Gray). Got very much deviated

~~different kinds of cells -~~ ~~the~~ ~~projections of the~~ ~~epithelium~~ ~~(cavities)~~ ~~which give it a rough~~ ~~or~~ ~~when~~ ~~cut~~ ~~the~~ ~~wood~~ ~~grows~~ ~~by~~ ~~addition~~ ~~to~~ ~~the~~ ~~outside~~ ~~in~~ ~~each~~ ~~part~~ ~~of~~ ~~the~~ ~~bark~~ ~~grows~~ ~~by~~ ~~addition~~ ~~to~~ ~~the~~ ~~inside~~ - each layer of the bark grows separately by means of cells placed between it & the next layer inside - The 2 or 3 layers cease to grow as they gradually ^{generally} become cracked from the pressure of the growing wood (cracked on Smooth - Birch, holly, etc.)

~~The part essential to the development of life is~~
The active part of an Englemann's cone is
in its development to give rise

- (1) New layers of wood
- (2) Cambium
- (3) Recently formed liber

the spring, to which the name of *cambium* (*cambio*, Lat. to change) has been given. Cells are afterwards formed in it, which become elongated and deposited, and originate the new layer of wood for the year.

Medullary rays may be seen radiating from the centre in almost any transverse section of an exogenous stem. They constitute the medium of communication between the pith and bark; and consist chiefly of flattened muriform cells. These intersections separate the wood into a series of wedges, more or less widely separated. These are sometimes very distinct. *Ex.* Clematis.

The *Bark* occupies the most external position in the plant. It consists of three layers, besides the epidermis, *i. e.*—

Epiphylœum (*epi*, Gr. upon; *phloios*, bark).

Mesophylœum (*mesos*, Gr. middle).

Endophylœum (*endon*, Gr. within), or *Liber*.

The *Epiphylœum*, outer or *suberous* layer of bark, consists of flat tabulated cells. In some instances it is very thin, and in others largely developed in a thick corky layer, which readily separates from the series below it. *Ex.* Cork oak.

The *Mesophylœum*, or middle layer, is generally developed to a less extent than the outer. It consists of cells differing in form from those of the epiphylœum, with thicker walls, and intercellular spaces. The laticiferous vessels occur in this series. The mesophylœum is occasionally much developed. *Ex.* Larch.

The *Endophylœum*, or *liber*, is the fibrous or vascular layer of the bark; it is the *bast-layer*, so useful for textile purposes. During growth it separates into a kind of network. *Ex.* Lace bark and Cuba bast.

Additions are made to the cortical layers *internally*.

The *cellular* system of the stem has been seen to

consist, in exogens, of the pith, medullary rays, and two outer layers of the bark.

Peculiar and anomalous forms occasionally occur, in which secondary woody masses surround the central or primary one.

SECTION XII.

ASCENDING AXIS—ENDOGENS AND ACGROGENS.

Endogens (*endon*, Gr. within; *gennao*, to produce) include the palms and grasses, as well as other natural orders of plants. The chief features in endogens consist in the cellular and vascular systems being mixed together, or rather, in the cellular portion being traversed throughout its extent by vascular fibres. They do not possess a distinct central pith, medullary rays, concentric rings, or a separable bark. The external portion of the stem is the hardest. The name originated with the belief that the stem increased by additions at the centre, so that the materials earlier accumulated were pushed outwards. The structure of a palm stem being accepted as a type, it has been shown that the new fibres proceeding from the new leaves downwards first direct themselves to




Fig. 41.

the centre, whence they soon diverge, crossing the existing fibres in their progress to the circumference, where they become at last the most external (*Fig. 42*). Exogens increase in diameter indefinitely, endogens do not thus increase beyond a certain period (*Fig. 41—42*).

At the upper portion of the

Ex. Endogenous Stem (1) Shows no distinction of pith, & bark (2) they have no medullary rays (3) or concentric circles or zones -

Endogenous Stem - Consist of vascular bundles dispersed among cells or tissues & interlacing with one another. No pith, separate bark, medullary sheath, medullary ray or cambium - ~~wood~~ & hard outside soft inside. the hard part being formed by the ends of the interlacing vascular bundles -

The vascular bundles have their origin in the primum Vegetationis of the stem & ~~downward~~ ~~in~~ in growth upward & outward in the leaves & downward & outward ~~and~~ toward the circumference -

In the yuccas & Dragon palms the vascular bundles after having reached the pith are continued downward as fibers layers between it & the original vascular bundles. Somewhat after the manner that layers of wood are produced by the cambium in Quercus. Thus there is no limit to their increase in girth.

Strobilium stem - Consists (1) of pith & pith compound of parenchyma & pleurochyma (2) central ~~to~~ ...

composed of plemenchyma & scalariform vessels arranged in a spiral manner)

The vascular bundles are composed of (1) woody tissue or plemenchyma, spiral vessels, fiber cells & laticiferous vessels towards the pith all these vessels disappear so that the bundle is composed only of plemenchyma

The outermost part of an Eudicots stem is the hardest, therefore twining plant produce no deep furrows upon their surface—

Q

The stem is

- (1) Dichotomous - In Date palm.
- (2) Very wide - Dragon trees; this is owing to the cortical integument remaining soft
- (3) In Xanthorrhoea the structure of the stem is like that of an exogen owing to the presence of horizontal fibres radiating from the centre & cutting the vascular bundles at right angles.
- (4) The stem of Sarcocolla present a circle of vascular wedges resembling in appearance the stems of certain exogens —

Aerogenous stem composed of (1) plemenchyma
(2) Scalariform vessels (3) dotted vessels.

palm stem, the vascular bundles proceeding from the leaves contain spiral, porous, and laticiferous vessels, mixed with threads resembling liber. These all successively disappear, except the last, or become converted into a complete network of fibrous tissue as they reach the circumference.

In grasses the stem is often hollow from node to node, at which point the fibres cross over to opposite sides, and by interlacing form a dissepiment or partition across the stem. In the rapid growth of this kind of stem the more cellular central portion is ruptured and disappears.

Circles are sometimes visible in a transverse section of a palm stem from the upper portion, but these bear no analogy to the concentric rings of exogens.

The loose thready character of palm stems renders them almost valueless as timber trees: this is often compensated for by other valuable products.

Acrogens (*akron*, Gr. summit). In this instance the additions to the stem are dependent upon the union of the bases of the leaf-stalks, and the extension by growth of fresh leaves at the summit.

To this class, tree ferns belong.

Acrogens have no other stem than that formed by the union of the bases of the leaf-stalks with the axis from



Fig. 42.




Fig. 43.

which they were developed. Scars are left on the surface of the stem where the upper portions of the leaves have fallen away (*Fig. 43*).

A section of a tree fern exhibits a circle of irregular vascular bundles near the circumference, which were originally each of them the vascular system of a leaf-stalk. The rest of the stem is cellular, which falls away eventually in the centre, and leaves the stem hollow.

The stem of a tree fern may always be distinguished from that of an endogen by the presence of the vascular bundles, and the rhomboidal scars on the exterior.

Acrogenous stems are seldom branched or furcate.

The leaves of acrogens are denominated *fronds* (*frons*, *frondis*, Lat. a leaf), and the petioles, or stems, are called *stipes* (Lat. a trunk of a tree).

Stems are said to be *procumbent* (*procumbo*, Lat. I lie flat) when they lie on the ground.

Scandent stems (*scando*, Lat. I climb) are such as climb by means of suckers or tendrils.

Voluble stems (*volubilis*, Lat. easily turned) are those which entwine around other plants, as the hop and honeysuckle.

SECTION XIII. (a)

SUBTERRANEAN STEMS.

Under the name of subterranean stems we have included also those which are prostrate and not strictly subterranean.

The majority of forms to be described in this section are popularly designated as *roots*.

It must be remembered that *roots* do not possess

unconscionable number of some certain Stem
 of ultrapinnate - spiral clearly
 of stale - chamber

As the position or posture stems are
1. Direction

- (1) Prostrate or procumbent
- (2) Decumbent - when a prostrate stem rises towards its extremity
- (3) Ascending - Rising obliquely from the base
- (4) Scandent or climbing - when they cling to other plants or walls & by tendrils or root like processes - Ex. Ivy
- (5) Voluble or twining (a) D. to left - Ex. Cucurbit
Dodder (b) left to right - Ex. Hop, Juncus
Richardsonii (c) to left & left to right alternately - Ex. Wisteria, Passiflora

Modifications of the Stem

2. Aerial

1. The runner or Flagellum (45) Ex: Strawberry
 2. The Offset (Shorter & thicker runners. Ex: Houseleek)
 3. The Stolon (A branch from the air curving down to the earth & then giving off roots & stem. Ex: Plantain, grass)
 4. The Sucker (First subterranean then Aerial, Stolon the Ex: Rose, mint)
 - (5) The Rhizome (46) Ex: ginger, Jamruic, Iris
- 1-4 - modified branch. 5. Modified branch or stem

3. Subterranean

1. The Soboles or creeping stem (46) Ex: Sanjambh, Branch [9]
2. The Tuber. (47) Ex: Potato, Jerusalem Artichoke
- x 3. The Bulb. (48) Have fleshy scales Ex: Onion, Tulip
4. The Corm. (49) are solid Ex: Saffron, Crocus
- Colchicum & R



runner - plant producing suckers.



stolon

Subterranean stems are distinguished from roots by (1) the power of forming regular buds (2) by having rudimentary leaves or scales on their surface -

Tubers occur in dicotyledons. Corms in monocotyledons

cloves = young bulbs developed in the axils of the scales of bulbs. stencil bulbs - In Helianthus

scales (modified leaves) or buds (rudimentary leaves) or nodes (whence buds are developed).

The *Rhizome* (*rhiza*, Gr. a root), or rootstock, is a thick, procumbent stem, partly, and sometimes entirely, under the surface of the soil. It develops roots from its under side, and leaves from its upper. Its surface generally bears the scars left by the falling away of old leaves (*Fig. 44*).

.. *Ex. Iris, Tugeric, ginger, Asparagus, Solomon*

The *Flagellum* (Lat. a young twig), or runner, is a long, slender, procumbent branch, which develops a leaf-bud from its upper surface, and roots from its under, at each node. Each vegetating node becomes a perfect plant (*Fig. 45*).

Ex. Strawberry.

The *Soboles* (Lat. a shoot, or young branch) is a creeping underground stem, or branch, which emits roots from its under surface and leaves from its upper. It thus




Fig. 44.




Fig. 45.




Fig. 46.

resembles a rhizome, but is much more slender, and subterranean in its character (Fig. 46). *Ex.* Couch grass. *Saxifraga* *caux* or *Sedge*



The *Tuber* (Lat. a knob) is a thickened portion of an underground stem or branch, which serves as a depository for starch, and other nutritious secretions for the service of the plant (Fig. 47). *Ex.* Potato. *Jerusalem Artichoke*

A scaly modification of the tuber exists in the species of *Maranta* which yields arrowroot.

Bulbs are scaly modifications of leaf-buds developed upon a flattish disk, from the under surface of which roots are directed downwards (Fig. 48). *Ex.* Onion *Lily*, *Jalap*



Fig. 48.

The *Pseudo-bulbs* of orchids are thickenings of the base of the stem by the deposit of bassorine; they are produced above the surface, and are of a green colour.

Corms (*kormos*, Gr. a stem) are also expansions of the base of ^{the stem,} the stem,



Fig. 49.

differing from bulbs in being solid and not scaly, and from pseudo-bulbs in being subterranean, and consequently not green. Corms also contain starch and other nutritious secretions. Some botanists consider them as buds (Fig. 49). *Ex.* Meadow saffron.

Bulbs, *pseudo-bulbs*, and *corms* are confined to endogenous plants.

(a) The corns give off buds from their apex in Crocus; from the side in Colchicum which grow at the expense of the old Corm.

Bulbs - (1) annual - The outermost scale has an axillary bud, which grows into a new bulb after the flowering season is over, and its terminal point grows into the flowering axis, while the old bulb with scales ~~withers~~ - Hence such bulbs have been called proventives because the bulb of any ^{given} year is formed before the time leaves & flowers. Ex. Garden Tulip.

(2) Biennial - The bud is seated in the innermost scale ~~is~~ at the base of the axis which has just flowered - when this bud flowers it is at first surrounded (by the decay of the axis) by the scales of the parent axis but after the flowering is over these quickly die out & the axis which has just flowered gives origin to a number of scales replacing the ~~most~~ ^{upper} of these giving rise to a new bud - Such bulbs are called proventives as the leaf scales of each axis are developed after it has flowered & biennial as it bears growths belonging to 2 seasons on the same axis ^{1st year} Ex. Crown Imperial (*Fritillaria imperialis*) *



I. Vernation

A. Of Single leaves

1. Folded or bent &

1. Reclinate (50) *Aspidodendron*, Tulip
2. Conduplicate (52) oak, Rose, cherry
3. Plicate (53) Palm, Vine, Currant, Sycamore
red, oak (Hicksville)

2. Rollled

1. Circinate (51) Fern, Sundew
2. Convolute (54) Banana, Fig   *Arauc.*
3. Involute (55) Waterbily, violet, apple,
4. Revolute (56) Dock, Rosemary, Azalea

- 2 -

(3) perennial bulbs - retain the products of the condensed axes of several years in a state of vegetative condition - (years usually 2 part of one future) - the bulbs are part-ventricles in the perennial but may be distinguished as being indefinite -

Cornus. In Cornus each new corn ends in a list of glumes which are the rudiments of the ~~stem~~ future axis. This axis grows ^{off from} its base develops later latently rootlets to produce a new corn, at the summit of the last & at the same time a bud is produced in the axil of the uppermost leaves of the growing axis.

SECTION XIV.

L E A F - B U D S .

Leaf-buds are either axillary or terminal, *i. e.*, they constitute the extremity or growing point of branches, or are produced in the *axils* of previously developed leaves.


Because of their capability of being removed and grafted or budded upon other plants, they have been called *fixed embryos*.

They are generally protected by scales of a firmer texture than the leaves themselves, which serve a temporary purpose, and then fall away.


These scales are lapped over each other, in accordance with a spiral arrangement, so that the bud resembles a small cone.

The arrangement of leaves in the bud is termed *vernation* (*ver*, Lat. spring), and follows different plans in different plants, according to regular fixed laws.

The following are the most common forms of *vernation*. First, of individual leaves.

 *Reclinate* (*reclino*, Lat. I bend back), folded down from the apex to the base (*Fig. 50*). *Ex.* Liriodendron.

Circinate (*circino*, Lat. I turn round), rolled from apex to base, as in the fronds of ferns (*Fig. 51*).

 *Conduplicate* (*conduplico*, I double), folded laterally (*Fig. 52*). *Ex.* Oak.

Plicate (Lat. knit together), several folds like a fan (*Fig. 53*). *Ex.* Vine. *Currant, Syca*

Convolute (*convolvo*, Lat. I entwine), rolled upon itself (*Fig. 54*). *Ex.* Apricot. *Banana*



Fig. 51.



Fig. 53.



Fig. 54.

Involute (*involvere*, Lat. I fold in), with the edges rolled inwards (Fig. 55). *Ex.* Violet. *Water lily*



Fig. 55.

Revolvute (*revolvere*, Lat. I roll back), with the edges rolled outwards (Fig. 56). *Ex.* Rosemary. *Dock*, *Azalea*



Fig. 56.

The arrangement of all the leaves in the bud gives rise to other names.



Fig. 57.

Valvate (*valvæ*, Lat. folding-doors), nearly in a circle, at the same level touching each other by the edges (Fig. 57).



Fig. 59.

Imbricated (*imbrex*, Lat. a roof-tile), applied over each other at different levels, as the tiles of a house (Fig. 58). *Ex.* Lilac.



Fig. 58.

Twisted or spiral. When the margin of one leaf overlaps that of another (Fig. 59).



Fig. 61.

Induplicate (Lat. folded in). The leaves are folded so as to touch at their edges, or the margins are folded inwards (Fig. 60).



Fig. 60.

Equitant (*equito*, Lat. I ride), conduplicate leaves covering each other in opposition (Fig. 61). *Ex.* Privet.



Fig. 62.

Obvolute (*obvolvō*, Lat. I wrap round). Half of one leaf covers half of another, so that it becomes half-equitant (Fig. 62). *Ex.* Sage.

The bud is often protected by a resinous exudation or a downy covering.

Bulbs (see Fig. 49) are subterranean buds, in the centre of which is a growing point covered with scales. New bulbs are formed like buds in the axils of the scales.

Bulbils are small rounded bodies resembling bulbs,

Vernation (Continued)

B. Of Several leaves

α. The leaves plane or slightly convex.

1. Valvate (57)
2. Imbricated (58) Lilac, white holly-
3. Spiral or Twisted (59)

β. The leaves rolled or bent

1. Induplicate (60) ^{involute} ~~and slightly convex, or~~ ~~to curl back of each other.~~
2. Equeitant (61) Iris
3. Ovolute or $\frac{1}{2}$ equitant (62) Sage, Dog wort.

(4) Supervolute - one convolute over another
convolute.  Apricot

The leaf buds or ovules are produced by the elongation of the parenchymatous system of the stem or its branches in the form of a conical parenchymatous mass connected with the pith; around this vascular bundles are developed connected with the wood of the stem. On the outside of this again is a spiriferous mass which ultimately becomes the bark little conical cellular projections and webbed which are the rudimentary leaves.

Organs protecting the buds in old countries are called Scabs or tegments.

Budding, Grafting, Layering consist in separating the leaf bud or similar part...

Plants raised by such means propagate the individual peculiarities of their plants, while those raised from seed have merely a specific identity.

consisting of a number of thickened scales, frequently consolidated together. They are found in the axils of the leaves of some lilies, are easily detached, and capable of producing young plants under favourable circumstances.

Adventitious buds in the form of woody nodules are found in the bark of some trees, and are called by Dutrochet embryo-buds, but more appropriately by Dr. Lankester, abortive branches. *Ex.* Cedar.

Leaf-buds may become arrested and transformed into *spines*, or thorns, as in the sloe and barberry.

Under other conditions, leaf-buds may be converted into *tendrils*, as in the passion-flower.




Fig. 63.

Branches are but lateral expansions of the stem, or developed leaf-buds proceeding from the cellular system of the stem; branches of the first year, directly from the pith and medullary sheath (*Fig. 63 a*), and those developed afterwards communi-

cating through the medullary rays (*Fig. 63 b*) with the central medulla.

SECTION XV.

LEAF-STRUCTURE.

Leaves are appendages to the stems of plants, and may be regarded as expansions of the bark.

A leaf generally consists of an expanded portion, called the *blade* or *limb*, and a stalk or *petiole*, which latter sometimes expands at the base into a *sheath* or *vagina* embracing the stem. or becomes developed into leaflets called *stipulæ*.

(a) Leaves consist of vascular tissue in the *veins* or *ribs*, with cellular tissue, or *parenchyma*, filling up the interstices, and an *epidermis* covering all. In what are called skeleton leaves, the cellular portion is removed by maceration, and the network of the fibro-vascular remains. The lattice-leaf plant of Madagascar has skeleton leaves with very little *parenchyma*.

The *Epidermis* (*epi*, Gr. upon; *derma*, skin) consists of more or less compressed cells, having a different structure on the upper and under side of the leaf. The cells of the epidermis are generally flat, with sinuous, wavy, and irregular margins fitting in to each other, and leaving no openings or intercellular spaces, excepting *stomata*.




Fig. 64.

These latter orifices occur principally in the epidermis of the under surface of aerial leaves, and the upper surface of floating leaves (*Fig. 64 s s*). The cells of the epidermis are transparent and colourless.

The *Parenchyma* is the cellular tissue of the leaf which fills up the interstices between the veins. It consists of two series of cells, each containing chlorophyll. Below the upper epidermis of the leaf the cells are arranged more closely together, perpendicularly to the surface, and with very small spaces between them. The cells beneath are arranged horizontally, are more irregular, with larger intercellular spaces. This arrangement varies in different plants; in some, where both sides of the leaf are equally exposed to the light, the cellular series are similar on both sides. 6

Submerged leaves have no true epidermis, no fibro-vascular system, and no stomata. They consist of closely-compacted cells on the surface, which become

When the petiole is absent the leaf is Sessile
When the stipule is wanting it is exstipulate
As to duration leaves are (1) fragacious or caduce
falling soon after its appearance (2) deciduous or
annual (3) persistent, evergreen or perennial
again they are (1) non-articulated when they separate
by decaying on the stem (2) articulated when they
separate by means of articulation. *

The remains of non-articulate leaves are called
religios or indurva (the stem being thin indurvat
The scars left by the articulated leaves are
called Scar or licatrix. *

b) (2) In floating leaves the compact cells
are the broad on the under side, the spongiform on
the upper side - (3) In Succulent leaves the paren-
chyma is everywhere compacted.

*. Leaves are generally placed horizontally -
In Acacia, Eucalyptus they are placed vertically
Anything placed on the axil (angle) of the leaf is
called axillary, above it supra-axillary, below
it infra-axillary.

Leaves are Aerial or Submerged

In *Theophrasta gussiae* there are three layers.

II Venation Vascular Venation

- I Reticulated Ex. Dicotyledons.
- II Parallel Ex. Monocotyledons.
- III Forked or Fuscate Ex. Acetyledons.

I Reticulated

- (a) Unicostate - [chestnut]
Feather-veined Ex. Beech, oak, Holly, Spruce
Curved-veined Ex. Deadnettle, Lilac
- (b) Multicostate
Ribs - Convergent Ex. Cinnamon, Cassia
Divergent Ex. Melon, Castor oil.

elongated and compressed so as to resemble veins. The internal cells are very irregularly disposed, with large intercellular spaces containing air, which serve to support the leaf. *i. e. to float*

The *vascular system* of leaves is also double. On the upper side there are spiral vessels, reticulated or porous vessels, and woody fibre, and on the under side laticiferous vessels and fibres. These ramify through the leaf in the form of either simple or branching veins. This vascular system is a continuation of that of the stem, the upper corresponds to the woody layers, and the under to the *liber*, or *endophlœura* (a)

The *venation*, or distribution of the veins of leaves, varies in different plants.

In thick or *succulent* leaves they cannot be distinguished, on account of the mass of cellular tissue in which they are imbedded.

In the lower orders of plants, such as lichens and sea-weeds, and in submerged leaves, they are absent altogether.

In an ordinary leaf, as in that of the oak, there is a central continuation of the petiole, or footstalk, which is larger than the other veins. This is called the *midrib*; it gives off other veins from either side, which are called *primary*,^{veins} and if these give rise again to others, they are called *secondary*, &c. Leaves of this kind, with one large rib, are called *unicostate* (*unus*, Lat. one; *costa*, rib).

In such leaves as those of the sycamore, in the place of one central rib, there are two or more diverging from the point of junction of the blade with the petiole. These are *multicostate* (Lat. many-ribbed).

In all these instances there is a complete network of veins produced; hence the term *reticulated* (*reticulum*, Lat. a net), or net-veined, has been applied (*Fig. 65*).




Fig. 65.

In some instances the veins end in curvatures within the margin of the leaf, in others the veins proceed directly to the margin, as in the oak. The latter, for the sake of distinction, have been termed *feather-veined*.

In plants of a certain kind, of which grasses may be taken as a familiar example, there is a midrib, and veins running parallel to it from the base to the apex of the leaf (Fig. 66); or, in other and less familiar instances, there exists a central midrib, with veins running from it to the margin of the leaf, in a straight, or nearly straight direction, as in the plantain and banana of the tropics. In both cases there are generally small veinlets uniting the veins together, but they do not form a regular network, and are insufficient to prevent the leaf being torn down in strips between every pair of veins. These are instances of *parallel-veined leaves*.




Fig. 66.



Fig. 67.

In the leaves of ferns a third kind of venation is distinguishable, in which the veins divide in a forked manner. This kind of venation is therefore termed *furcate* (*furca*, Lat. a fork) (Fig. 67).

As a general rule, exogenous plants are possessed of leaves with a reticulated or netted venation, endogenous plants of leaves with a parallel venation, and acrogens of furcate-veined leaves.

SECTION XVI.

L E A F - F O R M .

Many distinctive features are imparted to leaves by the relative development of their vascular or cellular tissue. When the vascular increases most rapidly, and the parenchyma is insufficient to occupy the intermediate spaces, the leaf becomes deeply divided; sometimes the parenchyma is reduced to mere threads along each side of the principal veins. This feature is exhibited by the leaves of the horseradish when grown in sand or very poor soil. The vascular tissue may be prolonged beyond the margin of the leaf, terminating in spines, as in the holly. Under favourable circumstances, the cellular portion is most developed, so that the spaces between the veins of the leaf are too small to allow of its lying level between them; in this case it forms prominences, giving to the leaf a curled, crumpled, or corrugated appearance, as in the Savoy cabbage.

The innumerable forms of leaves may be well enough characterized for ordinary purposes by a few prominent types.


They are *simple*, when the blade is undivided, or when the divisions do not proceed directly to the midrib or *petiole* (a)

When these conditions are not fulfilled, the leaf is said to be *compound*.

Simple leaves, when undivided, are—

1) **ACICULAR** (needle-shaped), when long and filament-like, as in firs (x) *Pine, Larochre.*

SUBULATE (awl-shaped) when narrow and tapering from base to apex, as in the yew

(Fig. 68). Furze (Ulex Europæus)  Fig. 68.

LINEAR, when long and narrow (*Fig. 69*), but of equal width throughout. *Grass, Marsh Marigold*

LANCEOLATE, shaped like a lance-head (*Fig. 70*). *White willow (Salix alba)*. Broad middle of tapering ends

OVATE, egg-shaped, broadest towards the base (*Fig. 71*). *Porwinkle (Viola major)*

OBOVATE is reversely ovate, broadest at the apex (*Fig. 72*).

OVAL, or *elliptical*, widest in the middle (*Fig. 73*).

Fig. 69.

ORBICULAR, or round (*Fig. 74*). This form of leaf is generally *peltate*.

long
a



Fig. 71.



Fig. 72.




Fig. 73.




Fig. 74.

RENIFORM, or kidney-shaped (*Fig. 75*). *Ascum Eutopium*.

CORDATE, heart-shaped (*Fig. 76*). *Black Bryony (Tamus communis)*

OBCORDATE, reversely heart-shaped (*Fig. 77*).

SAGITTATE (*sagitta*, Lat. an arrow), shaped like an arrow-head (*Fig. 78*).

SPATHULATE (*spatula*, Lat. a kind of spoon), resembling a spoon (*Fig. 79*).



Fig. 75.



Fig. 76.






Fig. 77.



Fig. 78.



Fig. 79.



III Leaf-form Summary.

- | | |
|-------------------------------------|--|
| 1. Oblique (p. 47 a. 2) | 11. Orbicular 74 |
| 2. Acicular (p. 47. x) | 12. Reniform 75 |
| 3. Sabulate 68 | 13. Cordate 76 hinc |
| 4. Linear 69 | 14. Obcordate 77 |
| 5. Lanceolate 70 | 15. Sagittate 78 Scamin |
| 6. ovate 71 | 16. Spatulate 79 Pinnate |
| 7. Obovate 72 or 48a. b | 17. Pauriform 80 fig 1 |
| 8. Cuneate or wedge shaped (48a: a) | 18. Hastate 82 hinc |
| 9. oval or elliptical 73 | 19. Auriculate 82a |
| 10. Oblong. 73. a | 20. Subrotund 74a |
| 23. Irregular | 21. Lunate (16) 22. Subrotund |

Stem - leaves

Attachments

Fissures - fid. Half way between the margin & midrib (leaf or petiole)

Articulations - partite. Nearly to the base or midrib (leaf - partite)

Segments - Saele - Down to the midrib & base (The Saele have abrad base & have no articulations, the leaflets have articulations & narrow base) Leaf dissected

- 1. Bifid, trifid, quadrifid, quinquefid, septemfid, multifid
- 2. Bipartite, tripartite, quinque-partite, Multipartite.
- 3. Birsuled, trisected, &c.

More usually. A Feather-veined or Linear

Pinnatifid, pinnatepartite, pinnatisected } If further

is, tri, or _____ } decompos

Special Modifications - (1) Pectinate - close & narrow divisions in

midrib (2) Lopate - terminal large, rounded, lobed ^{smaller & smaller}

a. The stalk of a leaflet is called a petiole ^{or partial petiole}

or partial petiole & that of the compound leaf Rachis or Common petiole

The leaflets of a compound leaf may be at once distinguished from the separate leaves of a branch by their being all Situated in the same plane.

The presence of more than one articulation is a positive proof that the leaf is Compound. The converse is not true. The leaf of an orange is compound because there are two articulations (1) between stem & petiole (2) between petiole & blade.

* Laciniate (Pinnate type) deep & irregular



Fig. 80.

PANDURIFORM (*pandus*, Lat. bent in the middle), somewhat of the form of a fiddle (Fig. 80). *Fiddle Leaf*

Simple leaves, when divided, may be—

BILOBATE, or two-lobed.

TRILOBATE, or three-lobed (Fig. 81).

HASTATE, halberd-shaped (Fig. 82). *Wax's Sorrel (Rumex acetosella)*

QUINQUANGULAR, five-angled (Fig. 83).

PALMATE, with five lobes, resembling a hand (Fig. 84).

PEDATE (*pes*, Lat.

a foot), from a supposed resemblance to the foot of a bird (Fig. 85). *Ex. Hellebore.*

A leaf is said to be *compound* when

it is divided down to the midrib, so that the subdivisions appear as distinct leaves, which are called, in such instances, *leaflets. or folioles. &*

TERNATE, when composed of three leaflets (Fig. 86).

When this division is repeated upon itself, the leaf is

bilobate; and when again repeated, *triternate*.

DIGITATE (*digitus*, Lat. a finger), when there are five leaflets, like the fingers of a hand (Fig. 87).



Fig. 81.



Fig. 83.



Fig. 82.



Fig. 84.



Fig. 85.



Fig. 86.



Fig. 87.



Fig. 89.



Fig. 90.



Fig. 91.

PINNATE (*pinnata*, Lat. a feather), when the leaflets are arranged along the sides of the petiole (Fig. 88). They may be *oppositely* or *alternately* pinnate.

When each leaflet is again divided in a pinnate manner, the leaf is *bipinnate* (Fig. 89); and when again subdivided, *tripinnate* (Fig. 90). When not referable to any one of these forms, by a more complicated subdivision, the leaf is said to be *supra-decompound*.

The venation of compound leaves agrees with that of simple leaves; the compound character resulting from deficient parenchyma.

When the base of a leaf embraces the stem, it is *amplexicaul* (*amplexus*, Lat. embracing, *caulis*, stem) (Fig. 91).

When it runs along and is united to the stem, as a winged appendage, it is *decurrent* (*decurro*, Lat. I run along) (Fig. 92). *Ex.* Thistle.



Fig. 88.



Fig. 92

V. Margins

1. Serrate, Biserrate, Serrulate, Runcinate
 2. Crenate, Acutely crenate, crenulated, Bicerate.
 3. Dentate, duplicatodentate,
 4. Ciliated 5. Simiate. 6 Wavy or undulated (7) Crisped
- Serrulate - When minutely Serrate; Ex. *Barosma Serratifolia*.

c. Crenulated - Minutely crenate.

4. Bicerate - Crenate teeth crenate again.

5. Duplicato-dentate - The teeth of a dentate leaf dentate again. (313. c)

d. ~~Wavy~~ Wavy or Undulated - Slightly Sinuous. Holly

Crisped or curled - Very irregular twisted or

Curled - Ex. Garden Endive, Curled Dalk

Curled Mint.

3)

II.

Apex.

^{Simple}

(1) Acute

(2) Obtuse or blunt

(3) Acuminate (long tapering)

(4) Cuspidate

(5) Mucronate

(6) Abrupt or truncate

Divided - (1) Emarginate (2) Cordate (3) Retuse.





Fig. 93.

When the bases of two leaves unite, they are *connate* (Fig. 93). *Ex.* Teasel.

The margins of leaves also present distinctive features. When smooth or without serratures, they are *entire*. *Ex.* *Orchistube*

SERRATE (*serra*, Lat. a saw), when indented like a saw (Fig. 94). *Ex.* Dead-nettle.




Fig. 94.

BISERRATE, when each of such indentations is again similarly indented (Figs. 95). *Ex.* Nettle-leaved Bell-flower.




Fig. 95.



Fig. 96.

RUNCINATE (*runcina*, Lat. a large saw), toothed like a pit-saw (Fig. 96).

CRENATE, with convex serratures (Fig. 97). *Ex.* *Viola tricolor*, ground-hy.



Fig. 97.

DENTATE (*dens*, Lat. a tooth), with concave serratures (Fig. 98). *Ex.* Melon.




Fig. 98.

ACUTELY CRENATE, when the serratures are rectangular (Fig. 99).




Fig. 99.

CILIATED (*cilium*, Lat. an eyelash), when fringed with hairs (Fig. 100).




Fig. 100.

SINUATE, or waved; often irregularly (Fig. 101). *oak*.




Fig. 101.

SECTION XVII

LEAF-ORDER.

Leaves are arranged symmetrically about the stem; this is called *Leaf-order*, or *Phyllotaxis* (*phullon*, Gr. a leaf, *taxis* order). There are regular modes of

which leaves are developed, in accordance with a definite plan. (a)

Each node is capable of developing a leaf, and when the arrangement is such as to produce single leaves at different levels on the axis, the leaves are then said to be *alternate*. (1) *Ex.* Lime.

When some of the internodes are shortened, so as to bring two leaves to the same level at opposite sides of the axis, they are called *opposite*. (2) *Ex.* Lilac.




Fig. 102.

If more than two leaves are developed at the same level, they are said to be *verticillate*; and the leaves thus arranged constitute a *verticel* (*verto*, Lat. I turn), or *whorl* (Fig. 102). *Ex.* Goosegrass.

When pairs of opposite leaves cross at right angles, so that the first and third, and second and fourth pair are perpendicular to each other, they are *decussate* (*decusso*, Lat. I cut across). *Ex.* Dead-nettle. *Their stem is generally square.*

Alternate leaves are arranged in varied order in different species of plants. When they form two rows on opposite sides of the stem, so that the third leaf is over the first, the arrangement is *distichous* (*dis*, Gr. twice; *stichos*, rank).

The *spiral* may be accepted as typical of leaf-order, and the different arrangements may be represented by fractions, in which the numerator shall represent the number of turns in the spiral, and the denominator the number of leaves encountered in a spiral proceeding right or left from any given leaf to the next one immediately above it.

Thus $\frac{1}{2}$ represents the *distichous* arrangement; two leaves being encountered in one turn of the spiral, the third being over the first. *Ex.* Lime. *grass, Iris, Elm*

- (a) When the direction of the spiral (right to left or left to right) is the same both in the stem & the branch it is called homodromous when different heterodromous.



- (c) $\frac{1}{2}, \frac{1}{3}, \frac{2}{5}, \frac{3}{8}, \frac{5}{13}, \frac{8}{21}, \frac{13}{34}, \frac{21}{55}$ etc.
 $\frac{1}{4}, \frac{1}{5}, \frac{2}{7}, \frac{3}{14}, \frac{5}{23}, \frac{8}{37}$ etc.
 $\frac{1}{2}, \frac{2}{3}, \frac{3}{5}, \frac{5}{8}, \frac{8}{13}, \frac{13}{21}$ etc.

When the angular divergence divides exactly 360° , the arrangement is called rectilinear strobilar curviserial. The arrangement of the leaves is a distinct characteristic of species.

The leaves of Dicotyledons are mostly articulated (not oak etc). Those of Monocotyledons & Acotyledons are non-articulated.

- (a) A similar swelling at the base of a petiole of a compound leaf is called a stigma. Both Palms & Strains may be seen in the Sensitive plant.

Stigma of petiole

1. Sheath - in grasses (The membranous termination is called a ligule)
2. Air bladders - in Water plants as Pontederia, Japa
3. Winged - having a leaf-like border. Orange, Venus fly-trap etc.
4. Decurrent (leaf) - The wing continued down along the stem. e.g. Strain, Strain, Strain

And $\frac{1}{3}$ the *tristichous*. *Ex.* Sedge. ^{*Quercus, Tulip, Albu.*} Most common arrangement represents the *tetrastichous*. In this instance two turns of the spiral must be traversed before a leaf is encountered, which is directly over the first, and which, in this instance, is the sixth. *Ex.* Oak. ^{*Most common arrangement.*} *Ex.* Laurel. ^{*Apple, cherry, Pear, Holl.*} $\frac{2}{5}$ is another common arrangement. $\frac{5}{8}$ is found exemplified in the Houseleek. ^{*Flax*} $\frac{8}{13}$ in the lanceolate Plantain. ^{*Ex. European, Cor.*} $\frac{13}{21}$ in another species of Plantain (*Plantago media*), and other plants. (a)

In some instances a certain number of leaves are *fascicled*, or collected in a bundle, surrounded by a sheath at the base. *Ex.* Fir. (b)

A theory has been propounded, based on accidental coincidences, that the angles at which the primary veins of the leaf diverge from the mid-rib, the principal branches from the stem and those from the root coincide; but the facts do not appear to us to be proven. (c)

SECTION XVIII.

LEAF-APPENDAGES.

The *petiole* (*petiolus*, Lat. a stalk), or stalk of a leaf, forms an articulation, more or less complete, at the point of union with the stem. In some instances a transverse articulation takes place, as in the orange. The cushion-like swelling sometimes occurring at the base is called a *pulvinus* (Lat. a cushion); in the gooseberry it is changed into spines (*Fig.* 103).





Fig. 103.



Occasionally, the petiole not only becomes leaf-like, but the margins are folded in so as to form a kind of urn or pitcher, called an *ascidium* (*askidion*, Gr. a small leather bottle), if closed, and an *ampulla* (Lat. an open vessel), if open (*Fig. 104*). *Ex.* *Pitcher-plant*.
When the lamina is abortive, the petiole may become transformed into a *tendrill*; or when a lamina is present, the petiole may be extended beyond it in a tendrill-like manner. The upper leaflets of a pinnate leaf are also sometimes converted into *tendrills*.

On either side of the petiole, at its base, expansions and modifications occur, which are denominated *stipules* (Lat. a straw).

These are extremely variable in form and appearance (*Fig. 105 s*).

When they embrace the stem, so as to inclose it in a kind of sheath, such form of stipule is called an *ochrea* (Lat. a boot). *Ex.* *Rhubarb*. τ σ σ .

When they resemble leaves in appearance, they are said to be *foliaceous*. In some instances they are very large and foliaceous. *Ex.* *Pea*.



Fig. 105.




Fig. 106.

is owing to the separation of the vascular bundles before they reach the laminae & their filling up of the intervals by parenchyma. A phyllodium or phyllode (which occurs only in Equisetum) is distinguished from true blades by

- (1) parallel venation (2) vertical position (3) The similarity of appearance of ~~these~~ ^{the} two surfaces - *

Most pitchers are formed by the folding of the phyllodium, but some are formed by the folding of the blade -

Indrials are modified

- (1) Midrib - in gloriosa superba (5) leaf - vine dry
 (2) leaflet - Lathyrus Sylvesteris (6) Branch - Papier
 (3) Petiole - Lathyrus Aphaca flower, vein, g
 (4) Stipule - Smilax, Melon. Branch

Stipules - very large in Pansy, Common Pea.

Indrial - Lathyrus ... Melon, Smilax


Stipels - Stipules of leaflets (called thin stipel
Stipellate) Bean, Bladder nut, Desmodium

1. Adnate - adhering to each side of base of petiole. Rose

(2) Caulinary - separate on each side of base of petiole

Ex: Pansy & many Willows

Intrapetioles (3) Synochrotrichous or opposite - two joining by their inner margins. Ex. Astragalus.

Plumose (4) Apithary - joining by the inner margins 

(5) Ochrea. (6) Interpetioles.

(1 - leaflets, spines, Corbice, Scaly are modifications of stipules)

Stipules perform the function of leaf
axils, the leafy part is called by a main part.
General view of the leaves in the
three great classes of plants

In Dicotyledons. Venation reticulated, leaves articulated
often compound & variously toothed or incised at their
margin.

In Monocotyledons - Venation parallel (with few
exceptions). Leaves not articulated, margins
usually entire, commonly simple, often sheathing
& seldom have stipules.

In Acotyledons - Venation Forked or furcate
leaves non-articulated, frequently toothed
- or incised, often Compound.

Decurrent leaves often become prostrate
Cylindrical - e.g. Helianthemum, Oniscus.

They may occur between the bases of the petioles of opposite leaves, and are then described as *interpetiolarly* (Fig. 106 s). *Ex.* Cinchona. *haca*.

If the stipules are attached through their entire length on either side of the petiole, they are *adnate*. *Ex.* Rose.

The fibrous material found at the base of palm-leaves is often considered as of a stipulate character, and is termed a *reticulum* (Lat. a net).

The sheathing petiole of grasses is of an allied character, and is called a *vagina*.

Stipules are sometimes converted into tendrils. *Ex.* *Smilax* sp.

Or they may be consolidated into thorns (Fig. 107). *Ex.* Acacia.

Regard must be paid more to their position than appearance, on account of their varied modifications.

When the leaflets of a compound leaf have at their bases small appendages of a similar character, these are called *stipels*. *Ex.* Bean.

Leaves having stipules are described as *stipulate*, but when absent, *exstipulate*.




Fig. 107.

SECTION XIX

FLOWER-BUDS.

Flowers in the bud may be described as fixed points surrounded by rudimentary modified leaves. They differ from leaf-buds in the central point being

fixed instead of growing. Sometimes this central point departs from its fixed character, and, by continuing its growth, assumes the form of a branch, with leaves but little differing from the ordinary leaves of the plant. Such a growth is an unnatural one, and is generally characterized as a monstrosity.

Flower-buds proceed from the axils of leaves. Such leaves are termed *floral leaves*, or *bracts* (*bractea*, Lat. a thin leaf).

Sometimes these bracts are coloured, as in *Bougainvillea*; at others they remain green, as in the Lime.

All intermediate leaves, of whatever size, form, or colour, intervening between the bract and the calyx, are called *bractlets*.

A single bract may become so enlarged and folded as to inclose many flowers, which are arranged on a *spadix*. Such form of bract is termed a *spathe*. It is either membranaceous, foliaceous, or woody (*Fig. 108*). *Ex. Arum*.

If several bracts are arranged in a *whorl* or *verticel* around the base of certain kinds of inflorescence, such as an umbel or capitulum, such a verticel of bracts is called an *involucre*. *Ex. Fool's parsley.*

Small bracts which overlap, or imbricate, like the tiles of a house, are often termed *scales*. When the bracts of an involucre cohere and form a cup around the base of the fruit, as in the acorn, it is called a *cupule* (*cupula*, Lat. a little cup (*Fig. 109*)).

The manner in which the *floral en-*



Fig. 108.




Fig. 109.

Bracts

1. Scales - Bracts of catkin or Amentum Ex. Willows, oak, Birch
2. Involucre - Whole of bracts around a flower or series of flower stalks - Dandelion, umbels, Daisy etc.
Compound umbel has (1) general involucre (2) partial involucre or involucel - The each bract of involucre of composite flowers is called a perianth.
3. cupule - Involucre forming a (cup like) part of the fruit
Ex. Acorn, Hazel nut.
4. Spathe - Large sheathing bract completely enclosing one or a number of flowers - Iris, Arum, Palms, Banana
5. Glumes, paleae, Squamulae are the three sets of bracts of grasses from without inwards.

Arduana of speciation

9) of single ~~two~~ parts of the floral envelopes

The same terms are used as in venation with the simple addition of the term

Crumpled or corrugated - Ex. petals of Poppy (Papaver) & Rock-rose (Helianthemum)

10) of several parts of the floral envelopes

(a) Circular.

- (110) (1) Valvate - Calyx of Lime (Components slightly thick overlapped & thin margins of the petals) (Sepals)
(111) - (2) Induplicate - Pet. of Quercus ulmifolia, Calyx of Quercus
(3) Reduplicate - Ex. Calyx of Hollyhock, Petals (in

(112) Contorted or twisted. Hollyhock (Crotia) Fls. (immature), Vi
(113) Imbricated. Mallow (Crotia)

- (1) Imbricate. Ex. Camellia japonica (Calyx)
(2) Cochlear - Frog's mouth, in wh. the 2 of the 5 leaves instead of being wholly outside as is the case of in quinquefidal is wholly inside.
Ex. Frog's mouth

(3) Vexillary - The superior vexillum overlaps the rest of the petals
Ex. Leguminosae.

velopes, i. e., the calyx and corolla, are arranged in the bud, is termed *æstivation* (*æstivus*, Lat. belonging to summer).

This arrangement often differs from that of leaf-buds considerably, and is commonly of one of the following kinds:—



Fig. 111.

VALVATE. This is the same as valvate vernation in leaf-buds (Fig. 110).

INDUPLICATE. When the margins are folded inwards (Fig. 111).



Fig. 110.

CONTORTED, or TWISTED. When twisted in the bud (Fig. 112).



Fig. 113.

IMBRICATE. When overlapping as in the tiles of a house (Fig. 113).



Fig. 112.

QUINCUNCIAL (*quincunx*, Lat. arranged as five in cards). Five sepals or petals, arranged in a spiral, so that two are exterior, two interior, and one half exterior and half interior (Fig. 114).



Fig. 115.



Fig. 114.



Fig. 117.

ALTERNATE. When the inner whorl alternates with the outer (Fig. 115).

PLICATIVE. When folded or plaited together (Fig. 116).



Fig. 116.

VEXILLARY (*vexillum*, Lat. a standard). A common form of æstivation in papilionaceous plants, where *v* is the vexillum, *a* the alæ, and *c* the carina (Fig. 117).

The different forms of æstivation are each perma-

ment in the species in which they are found, and are of some importance in systematic botany.

SECTION XX.

INFLORESCENCE.

The arrangement of flowers on the axis of growth is called the *Inflorescence* or *Anthotaxis* (*anthos*, Gr. a flower; *taxis*, order).

The axis, or primary stem supporting a flowering head, is termed the *rachis* (*rachis*, Gr. a backbone). The stalk supporting a flower is a *peduncle* (*pes*, Lat. a foot); and if this is branched, then the small branches which it gives off are *pedicels*.

A flower having no stalk is said to be *sessile*; but if stalked, *pedicellate*.

The following are the most common forms of inflorescence:—

SPIKE. When the peduncles are very short or wanting altogether (*Fig. 118*). *Ex. Plantago foliosa*

Amentum or *catkin*, a *deciduous* spike, bearing unisexual flowers.

Ex. Willow, oak, Birch, poplar

Spadix, a *succulent* spike, surrounded by a sheathing bract or *spathe* (*Fig. 118*). *Ex. Arum.*

Strobilus (Gr. a fir cone), a dense spike of female flowers, covered with membranaceous scales. *Ex. Hop.*



Fig. 118.




Fig. 119.

Pedicels

Peduncle - is the main axis of the flower stalk
Pedical - the branches of a peduncle bearing a flower.

1. Rachis or axis - elongated peduncle bearing sessile flowers or pedicels.
2. Receptacle - shortened horizontally dilated peduncle
Ex. Huckle (Hatter), Chamomile (Conical), Borstinia (Lvs. Fig (pear shaded)) [Prunosea, cowslip etc]
3. Scape - a radical peduncle. Ex. Fulph, Hyacinth,
In Cashed nut the peduncle becomes pulpy fruit.
4. Phyllod - leaflike Ex. Butcherbroom, Eupophylla.

Inertia - The peduncle is extraaxillary in lime & woody,
- Nightshade (Solanaceae)

Kind of Inflorescence

Indefinite, Opening - Centripetal (1) ^{leguminosae} Silicle, axilla (2) Whorl
(a) With elongated primary axis - development
or opening of flowers from base to apex. E. to Centre.
(b) With dilated primary axis, opening from circumference

- (a) Elongated primary axis (4) Strobilus (distillate)
(1) Spike - (118) (5) Cone (distillate) (119)
(2) amentum or catkin (unisexual) (6) housta or spikelet
(3) Spadix (7) Raceme. (120)

axillary cyme

(8) Corymb (122) (9) panicle (121) (10) Thyrus - a kind of panicle with the pedicels very short chestnut, lilac

(6) Dilated ^{or shortened} racemose axis

(1) Capitulum (125) or anthodium

(2) Hypanthodium (127) usually unilocular sunflower

(3) Umbel (128) Compound umbel (129) whorls in 2, 3, 4, 5, 6, 7, 8, 9, 10, 12, 15, 20, 30, 40, 50, 60, 70, 80, 90, 100, 120, 150, 200, 300, 400, 500, 600, 700, 800, 900, 1000

N.B. Spike is sepals raceme. Capitulum is sepals umbel

Definite

Solitary terminal - ex. Gentian, Wood Anemone (Nemorosa)

The order of development or opening is from apex base when the peduncle is elongated & from centre to circumference when it is shortened (proliterated).

Cyme - is applied to all definite inflorescence.

(1) True cyme (= Corymbae or umbellate cyme)
When a more or less branched definite inflorescence develops in a Corymbae or umbellate manner - These are Dichotomous, trichotomous & c according to the number of branches [

(2) Spiked cyme: a cyme with sepals flowers ex. Sedum

(3) Racemose cyme - cyme with pedicels of equal length. ex. Campy nula

(4) Panicled cyme - with branched pedicels - ex. Princh

Cone, a dense spike of female flowers covered with woody scales (*Fig. 119*).

Ex. Fir.

Locusta, a spikelet or flower-cluster of grasses.




Fig. 121.

RACEME (*racemus*, Lat. a bunch) differs from a spike, in the flowers being supported upon pedicels of equal length (*Fig. 120*). *Ex. Ribes.*

PANICLE (*panicula*, Lat. down of reeds) is a compound raceme, or a raceme with branched pedicels (*Fig. 121*). *Ex. Poa.*

Thyrus (*thursos*, Gr. a



Fig. 120.

Bacchanal's wand) is a compact panicle in which the pedicels are

short and often thickened, or succulent. Some botanists characterize it as a panicle having the middle peduncles longer than those above and below them. *Ex. Lilac and Horse-chestnut.*

CORYMB (*corymbus*, Lat. summit), a kind of raceme, in which the pedicels are gradually elongated, from the apex downwards, so that all the flowers are brought to the same level, or nearly so (*Fig. 122*). *Ex. Cerasus. Hawthorn &c.*



Fig. 122.

When these pedicels are at the same time branched, it is called a *compound corymb*.

CYME (*kuma*, Gr. a wave) is a panicle with the pedicels so developed or arrested, that the flowers



Fig. 123.

are brought to nearly the same level as in an umbel (Figs. 123, 124). *Ex.* Elder.



Fig. 124.

This kind of inflorescence should be studied rather in its development than its form. The order of expansion is *centrifugal*.

When the basal flowers expand first, the order of expansion is *centripetal* (*centrum*, Lat. centre; *peto*, I seek).

When the central, or uppermost flowers expand first, it is *centrifugal* (*centrum*, Lat. centre; *fugo*, I fly).

CAPITULUM (*caput*, Lat. a head).

When numerous flowers are arranged upon a nearly flat disk or receptacle (Fig. 125).

Ex. Daisy, *Thistle*, *Dandelion*

Glomerulus (like a ball). When the *capitulum* assumes a globular form (Fig. 126).

Ex. Teasel.

Hypanthodium (*upo*, Gr. under; *anthos*, a flower).

When the receptacle folds upwards, so as partially or entirely to inclose the flowers (Fig. 127). *Ex.* Fig.

UMBEL (*umbella*, Lat. a sun-shade.)

When all the flowers are supported upon pedicels of equal length which arise from a common point, and spread out like radii (Fig. 128). *Ex.* Ivy.

If each of the pedicels of an umbel gives rise to a secondary umbel, the entire inflorescence is



Fig. 125.




Fig. 126.




Fig. 127.



Fig. 128.

Helicoid or Spiral Cyme - (Sometimes Circinate or gyrate Cyme) Flowers developed only on one side of the upper extremity is more or less coiled ~~at~~ ~~of~~ ~~the~~ ~~leaf~~.

Contracted axillary or axilside - Cyme crowded with flowers placed on short pedicels of nearly equal length - Ex. Sweet-William ^{some this is} Pinks order

7. Globose - number of sessile flowers collected upon a rounded head or short spike -

8. Verticillaster - Whorl the central flowers opening first - Ex. ~~White~~ White deadnettle.

elliptical inflorescence

This is when one set of inflorescence is Centrifugal the other centripetal - Thus in the Compositae the florets are centripetal, but the general inflorescence is Centrifugal. In labiate plants (ex. White deadnettle) the partial inflorescence is Centrifugal but the general inflorescence is Centripetal.

The calyx has epidermis & stomata & hairs etc. Red
With venation like true leaves. Veins composed of spiral
vessels -

Sessile - almost without exception, generally entire (except
as in Peony & those divided at their margins) (toothed) in
many species of Dock, Passiflora foetida (divided)

Induction - Erect, Cornicent (turned inwards),
Divergent, reflexed or patulous.

(a) Polysepalous ~~7/8~~ ~~arranged~~

Regular - if the sepals are of equal size & like
form symmetrically arranged. Ex. Ranunculus,
or else Irregular - Ex. Monkshood.

termed a compound umbel (*Fig. 129*). *Ex.* Carrot, &c. *Trill. Bot. Alley, Newbrook*

Most of these forms, by repetition upon themselves, become *compound*.

The pedicels are but branches, and the various forms and developments of these flower-branches only modifications of the stem.




Fig. 129.

SECTION XXI.

THE CALYX.

The flower and its appendages consist ordinarily of four whorls of modified leaves; the outer two constituting the *floral envelopes*, and the inner two the *essential organs*.

The exterior whorl of the floral envelopes is the *calyx* (Gr. a cup). It consists of modifications of leaves, called *sepals* (*sepia*, Lat. I inclose), either united in part, or altogether or entirely free. When the *sepals* are free, the calyx is said to be *polysepalous* (*polus*, Gr. many); and when united, *gamosepalous* (*gamos*, Gr. union), or monosepalous (*monos*, Gr. one). These sepals present all the characters of leaves, and are sometimes, in cases of monstrosity, converted into ordinary leaves. They may be green, as in the rose, or coloured, as in the *fuchsia*.¹ When green they are *foliaceous* (leaf-like), and when coloured, *petaloid* (petal-like). The external envelope of the flower is always considered the calyx.

In some instances the flower appears to have a

double calyx, the outer being called the *epicalyx* (*epi*, Gr. upon). *Ex.* Mallow. *Strawberry* *any other of the Rubi.*

In a monosepalous calyx, the part formed by the union of the sepals is termed the *tube* of the calyx, and the free portion the *limb*, *the orifice*, *the throat of the*

Occasionally the calyx is *globular*, *campanulate* (bell-shaped), *turbinate* (top-shaped), or *inflated*.



Fig. 130.

When certain parts are enlarged and prolonged into a kind of spur, the calyx is *spurred*, or *calcarate* (*Fig.* 130). *Ex.* *Tropæolum*.

In composite plants the calyx is adherent to the ovary, and the limb becomes developed into a *pappus* (Lat. thistle-down), or hairy crown, by means of which the fruit, when ripe, is transported through the air (*Fig.* 131). *Ex.* Dandelion.



Fig. 131.

In the bright yellow *Eschscholtzia*, the upper portion of the calyx is ruptured from the lower, and carried upon the expanding corolla, leaving a permanent ring or fringe at the base. This form of calyx is termed *operculate* (*operculum*, Lat. a lid), or *calyptrate* (*kuluptra*, Gr. a covering) (*Fig.* 132).

When the calyx adheres to the ovary, it is said to be *adherent*, or *superior*; but when free from the ovary, it is *non-adherent*, or *inferior*. *Wall. flower. on the cup &c.*

Plants that are entirely destitute of floral envelopes are *achlamydeous* (*a*, Gr. without; *chlamys*, a tunic). *Ex.* Willows.



Fig. 132.

When but one floral envelope is present, they

Gamosepalous - Entire, ^{-dentate} Toothed (Campanium), ^{-fid} Cleft
 (Centauris) Pastate (primverul)

then entire the number of sepals is determined by the
 position the principal veins corresponding with the
 midribs of the sepals -

Regular & irregular

Special names (see the Corolla)

- (1) Hooded or galeate - Ex. Monkshood (one sep. forming
 as a hood)
 (2) Gibbous or saccate - little sacs at the base - Ex. Wallflower
 (3) Calcinate or spurred - with tubular prolongation downward
 Ex. Loxopur, Cathedraibine
 (4) Calyptrate or operculate - Ex calyptrae

Caducious - Poppy

Deciduous - Buttercup

Persistent - Mallow, Henbane

Forms part of fruit - Ex. Apple, pear, gooseberry, Melon,
 Cucumbers &c

When the calyx is adherent the modifications of the
 limb are (1) petaloid - orchis, Iris, Chocoma

(2) Thalaceous - ^{duyee} ~~Campanium~~ rim - Mastor

(3) Membranous - Scunflower absent - OX - eye }
 obsolete

(4) Pappose Compositae (101)
 (Plumae - Valerian
 Pilose - Dandelion)

The pappus is stalked (Dandelion) or sessile
 (Valerian)

Macroscent - When persistent and assumes a striated appearance - Ex. *Campanula* Heath

Acrocent - Persistent sometimes to grow after flowering forming a fleshy expansion round the fruit - Ex. *Winter cherry* & *Strophelia*

The Corolla

Distinguished from calyx by its coloured nature & more delicate structure.

Composed of parenchyma & veins of spiral vessels, no epidermis but no stomata as a rule.

Outline, incision, dentation &c described by the same terms as the leaves. But at its margin it is Embricated or fringed when they present long thread like processes. Ex. *Dianthus*.

Thick & fleshy - Ex. *Stapelia*

Dry & membranous. Ex. *Heath*

Stiff & hard - Ex. *Xylocopa*.

They are erect, connivent, divergent, patulous or reflexed as the Sepals

(A) Polypetalous Corolla

(a) Regular -

(1) Cruciform or cross - ^{no four petals} - Arranged in the form of a cross - Ex. *Wall flower* & other *Cruciferae*.

(2) Caryophyllaceous - Five petals with long lobes arched in the tube of the calyx & with the limbs commonly at rt angles to the claws.

are *monochlamydeous* (*monos*, Gr. one). *Ex.* Meze-
reum.

When both floral envelopes are present, *dichlamy-
deous* (*dis*, Gr. twice). *Ex.* Geranium.

SECTION XXII.

THE COROLLA.

The inner whorl of floral envelopes is termed the *corolla* (Lat. a little crown), and consists of more or less coloured leaves, and is generally the showy portion of the flower. The corolline leaves are termed *petals* (*petalon*, Gr. a leaf), and are generally placed alternately with the sepals of the calyx.

When the petals are more or less united, they form a *monopetalous* corolla; and when distinct, a *polypetalous* corolla. Petals are less leaf-like than sepals.

In endogens, the calyx and corolla are often similar in size, shape, and colour; hence the entire floral envelope is, in such instances, usually termed a *perianth* (*peri*, Gr. around; *anthos*, a flower). *Ex.* Tulip.

Petals are sometimes sessile; at others, the base is prolonged into an *unguis* or *claw*, when the petal is said to be *unguiculate*, and the upper or free portion is called the *lamina* or *limb* (*Fig.* 133). *Ex.* Pink.

The margins of petals, like those of true leaves, are either entire or toothed. When the teeth form a fringe round the margin, the petal becomes *fimbriated* (*fimbria*, Lat. a fringe). *Ex.* Pink.

Or a petal may be concave or hollowed like a boat, and is *navicular* (*navis*, Lat. a ship).

In other instances, one or more of the petals are



Fig. 133.

prolonged into a spur, and are called *calcarate* (*calcar*, Lat. a spur). *Ex.* Columbine.

The forms of petals are described generally by the same terms as simple leaves.

When all the petals in a flower are equal in size and shape, such a flower is said to be *regular*, and when otherwise, *irregular*.

We have, therefore,—

Regular monopetalous corolla. *Ex.* Bluebell.

Irregular monopetalous corolla. *Ex.* Calceolaria.

Regular polypetalous corolla. *Ex.* Buttercup.

Irregular polypetalous corolla. *Ex.* Pea.

Of the *regular monopetalous corolla* there are varieties of form, some of the more general of which have received distinct names; as—




Fig. 134.

CAMPANULATE, or bell-shaped (*campanula*, Lat. a little bell) (*Fig.* 134). *Ex.* *Campanula*. *Herb.*

INFUNDIBULIFORM, or funnel-shaped (*infundibulum*, Lat. a funnel). In this variety the corolla assumes the shape

of an inverted cone (*Fig.* 135). *Ex.* *Convolvulus*. *Tobacco*



Fig. 135.



Fig. 136.

HYOCRATERIFORM, or saucer shaped (*krater*, Gr. a cup), when the corolla consists of a straight tube surmounted by flat and spreading limbs (*Fig.* 136). *Ex.* Primrose.

TUBULAR, when it consists of a long cylindrical tube, as in the central florets of a daisy (*Fig.* 137).



Fig. 137.

Crossans - (sup. 66)

1) Irregular

- (1) Papilionaceus - mesepior, 2 lateral, 2 inferior (p.f.)
- Gamopetalous Corolla
(~~partly~~ ~~both~~ cleft, toothed, entire fls in calyx)

2) Regular -

(*) Campanulate

(1) Tubular (2) Campanulate (3) infundibuliform

(4) Hypocrateriform (5) Rotate or wheel shaped (6) Ureol-
ate or pitcher shaped or urn shaped

3) Irregular.

(1) Labiata or lipped. ex. Rosemary.

(2) Ringent or gaping (a modification of labiate)
ex. Dead nettle.

(3) Personate or masked

(4) Calceolate - ex. Calceolaria.
(slipper shaped)

(5) Ligulate or strap-shaped -

(6) Digittate or glove-shaped - Elongated bill-like
with resemblance to the fingers of a glove - ex. Fox-glove

(7) Irregularly rotate. ex. Speedwell.

(8) Irregularly Salver shaped or Hypocrateriform. ex. Red Valerian

Anomalous forms of appendages.

- 1) Saccate or gibbous - with a little bag or sac in the lower part. Ex. Snapdragon, Valerian.
- 2) Spurred or calcareate - The tube of a gamopetalous corolla prolonged downwards - Ex. Chelidonium.
- Simple - only one whorl.
- ~~Compound~~ - more than one whorl. Ex. White water lily.

Scales or hairlike processes in the inner surface

These situated at the junction of the claw & limb or at the base - Ex. Mignonette, Crowfoot, Lychnis, Ranunculus -

Near the throat of gamopetalous corolla - Ex. Forget me - not, Dadder &c.)

Form a cup-shaped process or corona in the Daffodil - (Corolla then called crowned)
Ex. Daffodil, Corolla of Pea flower.

Furcation

caducous, deciduous, persistent when it is usually marcescent (i.e. becomes dry & shrivelled) Ex. Heaths, Campanula.

ROTATE, or wheel-shaped (*rota*, Lat. a wheel), when the tube is very short and the limbs are spreading (Fig. 138).

Ex. Forget-me-not. *Bitter Sweet*




Fig. 138.



Fig. 139.

URCEOLATE, or urn-shaped (*urceolus*, Lat. a little pitcher), when the tube narrows at both ends and expands in the middle, as in some kinds of heath (*Erica*) (Fig. 139). *ros. lbony*

Irregular monopetalous corollas.

LABIATE, or lipped (*labium*, Lat. a lip), when the limb has two divisions opposed to each other, with a fancied resemblance to lips. *Ex.* Ground ivy.



Fig. 140.

RINGENT (*ringor*, Lat. to grin), when the lips are widely separated, and the upper is much arched (Fig. 140). *Ex.* Dead-nettle.

PERSONATE, or masked (*persona*, Lat. a mask), when the lips are closed, so that there is only a chink between them (Fig. 141). *Ex.* Snap-dragon.




Fig. 141.

CALCEOLATE (*calceolus*, Lat. a slipper), when the lips are hollowed out into a slipper-like form (Fig. 142). *Ex.* Calceolaria.

LIGULATE (*ligula*, Lat. a little tongue), when, as in the florets of many of the composite plants, the tube ends in a strap-shaped process on one side (Fig. 143). *Ex.* Dandelion.



Fig. 143.



Fig. 142.

Regular polypetalous corollas.

Fig. 144.

ROSACEOUS, when, as in the rose tribe, there are five spreading petals without claws (Fig. 144). *Ex.* Strawberry. *Single rose.*

CARYOPHYLLACEOUS, when, as in the pink tribe (*caryophyllaceæ*), there are five petals with long narrow claws, and spaces between them. *Ex.* Stitchwort. *Single pink, Campion, Snow*




Fig. 145.

CRUCIFORM (*crux*, Lat. a cross), when, as in crossworts (*cruciferae*), there are four petals arranged in the form of a cross (Fig. 145). *Ex.* Wallflower.

Irregular polypetalous corolla.

PAPILIONACEOUS (*papilio*, Lat. a butterfly), when there are five petals arranged as in the pea tribe; one superior, called the *vexillum* or standard (Fig. 146, *v*); two lateral, called *alæ* or wings (Fig. 146, *a*); two inferior, often partially united and called the *carina* or keel (Fig. 146, *c*). *Ex.* Broom.




Fig. 146.

What is ordinarily termed the *disk* is an enlargement of the receptacle, either as a cup, flat disk, or cushion be-

neath or around the base of the ovary, and sometimes between the stamens and petals, or with one or both of these organs inserted upon it. When the disk is lobed or divided, the parts are often denominated *glands*; they should rather be considered as *abortive stamens*.

Hermaprodit - male + female (bisexual)

Epilinaris - male or female (unisexual)

Neuter - having neither stamens nor pistils -

Monocious - staminate & pistillate flowers upon the same plant.

Dioecious - staminate & pistillate flowers in diff. plant.

Polygamous - staminate, pistillate & hermaproditic flowers on the same plant - Ex. Palms, Pellitory.

Stamen = Filament (stalk), Anther (blade) containing pollen
no Filament = Sepals. No pollen = abortive.

A. Filament

(a) Structure - spiral vessels in the centre surrounded by parenchyma & covered by epidermis with ornithoid stomata & hairs -

(b) Form - Filiform - tapering from base to apex. Ex. Rose

Capillary - very slender - Ex. Seeds & grasses

Blavate or club shaped - dilating from base to apex ^{in the lathyrus.}

Awl shaped - enlarged at the base & tapering upwards - Ex. Flowering Rush (Butomus umbellatus)

Petalloid - flattened - Ex. Water lily, Launa.

Forked - Allium

Forked - cranberry

Appendiculate (i.e. with various appendages, no scales etc. Ex. Boraginaceae)

Short - Ex. Boraginaceae, Primrose etc.

Young - Ex. Dahlia, Lily, Apricot etc.

Of various colours - Blue, black, white, yellow, red.

NECTARIES.—This term is confined to certain cavities, sometimes found at the base of petals, which secrete honey. Formerly the term was vaguely applied to other parts of the flower, but, except with the above limitation, may now be considered as obsolete. *Ex.* Ranunculus.

SECTION XXIII.

STAMENS.

The two inner whorls of the flower, called the *essential organs*, consist of stamens and pistil.

In some instances the stamens are present in one flower and the pistil in another. Such plants are said to be *diclinous* (*dis*, Gr. twice; *klinè*, bed) ^{or unisexual}.

When both ^{staminate} *stamiferous* and ^{pistillate} *pistilliferous* flowers are found on the same plant, it is said to be *monoecious* (*monos*, Gr. one; *ekion*, habitation). *Ex.* Hazel.

When the male and female flowers are found on different plants, the term *dioecious* (*dis*, Gr. twice) is applied. *Ex.* Hemp.

The stamens, or male organs, are represented, for brevity in writing, by the following sign ♂.

The pistil, or female organ, by the sign ♀.

The name *hermaphrodite* is applied to flowers in which both these organs are present, and the sign ♂♀, a combination of the two, is employed.

As the petals of the corolla are generally alternate with the sepals of the calyx, so the stamens are usually alternate with the petals, and consequently opposite to the sepals.

When the stamens are so placed as to spring apparently from beneath the pistil (*Fig.* 147), they are *hypogynous*.



Fig. 147.

gynous (*upo*, Gr. under ; *gunè*, female) *Ex.* Buttercup. *Poppay*.



When the stamens are inserted upon the calyx, and thus stand around the pistil, they are said to be *perigynous* (*peri*, Gr. around) (*Fig.* 148). *Ex.* Peach. *Cherry*. *Strawberry*.

When the filaments are united to the petals, so as apparently to spring from them, they are described as *epipetalous* (upon the petals) (*Fig.* *Prunrose* 149). *Ex.* Honeysuckle.



Fig. 150.

And when the stamens are inserted upon the ovary, they are *epigynous* (*epi*, Gr. upon) (*Fig.* 150). *Ex.* Umbellifers. *Joy*. *Lampbrush*.



Fig. 149.

In certain instances the filaments are closely united with the pistil into one column, and are *gynandrous* (*gunè*, Gr. female ; *aner*, *gen.* *andros*, a male). *Ex.* Orchis. *Bistort*.

When the number of stamens is less than twenty, they are said to be *definite* ; when more than twenty, *indefinite* (∞).

STAMENS consist of a stalk or support called the *filament* (*filum*, Lat. a thread) ; a head, or *anther* (*antheros*, Gr. belonging to a flower), and a powdery mass which the anthers contain, called *pollen* (fine flour) (*Fig.* 151).



Fig. 151.

When the anther is absent, the stamen is *abortive*.

The *filament* is usually thread-like, but sometimes

- Direction - erect, incurved, recurved, pendulous etc.
- Dedinate - filaments turned towards the same side of the flower - Ex. Horse Chestnut.
- Genucurvate - upper part forming an angle with the lower - Ex. Mahonia -

Sometimes there is a joint at the angle - Here the lower part is not filament but a flower stalk supporting a single ~~filament~~ stamen the other parts being abortive - Ex. Euphorbia

Spiral - Ex. Pellitory (Parietaria)

Duration - deciduous or persistent.

(B) Anther - At first consists of parenchyma soon after an epidermis; then four masses of large cells (two in each half) develop to form the pollen which gradually enlarge generally uniting into a single cavity in each half lobe.

• Quadricular (four cavities or chambers) Ex. Flowering Tobacco.

• Bilocular or dithecal - Ex. Barberry (most common)

• Unilocular - Ex. Mallow, Milkwort, Lady's Mantle.

• In Salvia ^{or Salvia} the connective becomes elongated into a kind of stalk each end of which bears an anther lobe.

One lobe in this case is sterile - It is then called the sterile lobe which the connective is attached to is called the back, the opposite the face. The face contains 1st a groove indicating the union of the lobes & 2nd two stamens (one on each lobe) on the line of opening

When the face is turned toward the pistil or the centre the anther is intorse ex. Mulberry
 When it is turned toward the petals or outward it is extorse - ex. Iris.

Structure - Two layers (1) Epithelium - outer & (2) Endothelium - inner (consisting of one or more rows of fibrous cells). At this stage there is no Endothelium & the epithelium is very thin.

Anthers having appendages formed by modified connectives are called appendiculate - ex. Magnolia (rounded upanin), Oleander (feathery process) Heart's ease (spur)

Forms of Anther lobes of anthers -

(1) Lobes

rounded - ex. Mercurialis annua

oval - ex. Almond

linear - ex. Acalypha

linear & sinuate - ex. Groundnut

(2) Anther

(1) oval, oblong &c

(2) Bitarcate - ex. Vaccinium uliginosum

quadrifurcate - ex. Gynalthena prostrata

(3) Sagittate - oleander

(4) Cordate - wallflower.

(5) Bifurcate at each end - grasses.

There may also be appendages from the lobes like those from the connectives - In this case also the anther is appendiculate.
 ex. G. ...

becomes *petaloid* (petal-like). In the common water-lily, all stages of transition, from the simple filament to the fully-developed petal, may be traced. In the Fuchsia, Tulip, and other cultivated plants, the same transition may be often observed.

The filaments, petals, and sepals, are regarded as metamorphosed leaves, and are sometimes characterized as *staminate*, *corolline*, and *calycine* leaves.

The length of the filament varies in different plants, its function being to support the anther in such position that its contents may be shed upon the stigma.

When, of four stamens, two have the filaments long and two short, they are said to be *didynamous* (*dis*, Gr. twice; *dunamis*, superiority.)

When, of six stamens, two are longer than the four others, *tetradynamous* (*tetra*, Gr. four).

Filaments are sometimes bearded, or *stipose* (*stupa*, Lat. tow), with delicate hairs. *Ex.* Mullein.

When all the filaments are united, so as to form a tube inclosing the pistil, they are *monadelphous* (*monos*, Gr. one; *adelphos*, brother). *Ex.* Mallow.

When united in *two* bundles, whether of equal or unequal number, they are *diadelphous* (*dis*, Gr. twice). *Ex.* Pea. *Fumitory* (equal)

When the filaments are united into *three* or more bundles, *polyadelphous* (*polus*, Gr. many). *Ex.* St. John's wort. *Castor oil orange* (more than 3) (2)

The **ANTHER** generally consists of two lobes united by a *connective*.

When the anthers appear to be united through their whole length to the filament on either side of it, so that the filament appears to pass between them, they are *adnate* (*Fig.* 151, c). *Water lily, Magnolia*

When the filament terminates at the base of the anthers, they are *innate* (*Fig.* 151, b). *Carex*

When the filament is attached in the middle of the anthers, so that they are movable, as in many grasses, it is *versatile* (*verto*, Lat. I turn) (*Fig. 151, a*). *introrse* (inwardly)

Anthers are commonly oval in shape; but are sometimes spherical, linear, curved, or flexuose.

The pollen is shed either by means of valves or pores. *Ex.* Lily and Heath.

The anthers are called *introrse* (*introrsum*, Lat. inwardly), when they open on the surface turned towards the centre; *extrorse* (outwardly), when they open on the outer surface.

The anther-lobe is *unilocular* or *bilocular*, as it contains one or two cells.

Even when the filaments are united, the anthers often remain separate and distinct; but in certain cases the anthers are themselves united together, and are then termed *syngenesious* (*sun*, Gr. together; *genesis*, origin). *Ex.* Dandelion. *All. Compositae, Adelia, violet &c.*

POLLEN consists of a number of vesicles or cases called grains containing a peculiar semi-fluid called *fovilla*.

The shape of these grains is very variable, but the spherical and polygonal are the most common forms.

The surface is sometimes smooth, and at others studded with points or prominences (*Figs. 152 & 153*).

Pollen grains occasionally adhere in masses, or are contained in bags.

Although yellow or brown is the ordinary colour of pollen grains, they are sometimes of other shades.

Their function is the impregnation of the ovules.



Fig. 152.

Fig. 153.

- Dehiscence — 1. Longitudinal — Wallflower, Dulif
 (usually in unilocular anthers) 2. Transverse — Alchemilla, Comua, Lavan
 3. Porous or apical — Saffron (apex) Heath (late
~~the position~~)
 4. Valvular — Ex. Barbary (?), Saffron

Relative length — (a) Relative to corda.

Included — Shorter than the tube of the corda — Ex. Forgetmen

Exserted — longer than the tube of the corda — Ex. Valerian.

(b) Among themselves

Pollen

Development & structure. The pollens correspond to the honey
stoves the upper & lower epidermis of the blade of the leaf
 The primordial utricle of the large cells developed in the young
 anther becomes infolded so as to divide them into four portions
 each of which substitutes a cellular wall on its outside —

There are thus three walls (1) that of the parent or mother
cell (2) that formed by the primordial utricle called that
 of the special parent cell (3) ~~the~~ cellular wall

Pollen cells — Generally the (1) & (2) are
 ruptured & the pollen cells are set free in
 the boches of the anther — Some times the wall
 of the special parent cell is not completely absorbed
 thus giving rise to a compound body consisting of 4 cells.

Ex. Penifloca graeca. If the membranes of two or more
 united parent cells are incompletely absorbed we get
 pollen masses of eight, twelve or any multiple of four.

Ex. Inga anomala 8. Acacia 12 ~~or more~~

Pollinia = pollen masses ~~of~~ waxy nature in orchids.
The pollinia of orchidaceae are subtended upon a stalk
(caudicle) adhering at the base to glandular mass
called (retinaculum) ~~for~~ in ~~the~~ ~~an~~ ~~place~~
upon a little projection (pedicellum)

a. Wall - Two membranes (1) Intine (internal) first
formed, smooth, ^{very} delicate, transparent & generally forms
a complete ~~lining~~ ^{lining} of the (2) Exine (external) hard
thick, resisting layer forming a kind of cuticle
over the intine. It varies in appearance

- (1) Smooth (2) with granular processes (3) with protuberances
(4) reticulations. The colour of the pollen cells resides in
the exine - red (verbascum) blue (in Epilobium)
black (Dulcis), green, white etc.
(5) pores or (6) slits or both (1 slit ^{or pore} mostly, 3 diately
rarely) ~~Some~~ Sometimes the pores have
lid like processes by. Peppin flower wh burst when the
pollens are discharged. The pollen cells are then sperulate
Gymnospermous plants have more than one cell in the
pollen -

(A) Content = granular protoplasm called ovule
which is semi fluid -

(V) Form - spherical, oval, polyhedral, triangular etc.

(S) Schisence - This takes place when the intine
is entered by water, sulphuric acid etc. The intine
send protuberances through the pores or slit of the
exine & burst by the ~~water~~ fluid which has entered,

The number of pollen grains produced is very great. It has been said that a single plant of *Wisteria sinensis* produced 6,750,000 stamens, and these, if perfect, would have contained 27,000,000,000 pollen grains.

In some cases the stamens approach the pistil in succession, and shed the pollen upon the stigma. In other cases, when the anthers are ripe, the elastic filaments are liberated from the corolla, and spring towards the pistil with a jerk, and scatter the pollen. Bees and other insects also assist in scattering the pollen from the anthers upon the stigma.

The Rosebay willow-herb, *Epilobium angustifolium*, has a four-lobed stigma, supported on a style rather longer than the filaments of the stamens. When the flower first opens, the lobes are closely applied together by their faces, and both style and stamens are pendulous (*Fig. 154*).



Fig. 154.

As the anthers approach maturity, the style becomes erect, and the stamens commence elevating themselves (*Fig. 155*).

As the anthers approach maturity, the style becomes erect, and the stamens commence elevating themselves (*Fig. 155*).




Fig. 155.



Fig. 156.

By the time that the anthers are fully matured, the lobes of the stigma have divided and curled outwards and downwards in a circinate manner, so that they may be reached by the anthers. the filaments then becoming



Fig. 157.

erect, and the pollen is discharge upon the lobes of the stigma (Fig. 156).

After discharging the contents of their anthers, the stamens droop and become pendulous again, whilst the style remains erect (Fig. 157).

SECTION XXIV.

THE PISTIL.

The centre of a complete flower is occupied by the pistil (*pistillum*, Lat. a pestle). It constitutes the inner whorl, and is the female organ of the plant. When mature, it is converted into fruit, and contains the seeds.

It is composed of modified leaves called *carpels* (*karpos*, Gr. fruit), or *carpellary leaves*. When it consists of but one carpel, it is *simple*, and when of more than one united, it is *compound* (Fig. 158).



Fig. 158.

The pistil includes the ovary, style, and stigma. The style is nonessential, and is sometimes absent). *When the stigma is sepale &*



Fig. 159.

When the carpels of a pistil remain distinct, the term *apocarpous* (*apo*, Gr. separate; *karpos*, fruit) is applied (Fig. 159). *Ex.* *Ranunculus*.

When the carpels are united a *syncarpous* (*syn*

endosmose & discharge the contents away - The pollen tube developed after the pollen cell is placed upon the stigma is also a prolongation of the antine through the pores or ~~sub~~ of the exine & filled with provida to be carried to the placental ovules.

Disk

Disk = All bodies between calyx & pistil or upon or in connection with these organs but which cannot be properly referred to ~~the~~ any of the floral parts (Bentley) (N.B. Between stamen & pistil Dalman). Generally such bodies are found between the stamens & pistil.

Examples - (1) ~~the~~ ^{ovary} rose (fleshy ring round pistil). Tree raony (dark red cup covering the greater part of the pistil). Rose cherry (waxy lining of the tube of the calyx). Umbelliferous plants (a swelling over the top of the ovary called a Stylidium). Crucifers (glend). Stone crop, Vine (seals). Columbine (petaloid expansion).
Disks are hypogynous perigynous (attached to calyx) & epigynous.

Pistil

The ventral suture corresponding to the junction of the edges of the lamina of the carpellary leaf is turned toward the axis, the ~~other~~ ^{other} corresponding to the midrib toward the circumference of the flower the dorsal suture.

2) Nature.

Leaf = modified leaf. (1) In Double Cherry they are transformed into an ordinary leaf (2) its development is exactly like that of an ordinary leaf.

3) Structure - It is a leaf folded with the upper surface in the outside & the upper inside. The ovary is the homologue of the blade & its structure consists of spiral vessels surrounded by parenchyma & enveloped by epidermis - The upper epidermis of the leaf forms the epi the lesion - The epidermis of the outside ~~is~~ that of the lower surface of the leaf & we frequently find stomata in it - The placenta is formed at the junction of the margins of the lamina by the development of a layer of cellular tissue - In this the vessels are attached & it is essentially double being formed by two margins meeting.

The style is formed by the apex of the leaf, the margins of which have been rolled inward & united - It is hollow, the canal being interrupted ~~by~~ usually by loose cells or filiform cells. At the period of formation these cells as well as those of the Stigma secrete a viscid fluid. These three form the conducting tissue (= loose tissue surrounding the canal (2) filiform cells (3) viscid fluid)

Stigma = an expansion externally of the conducting tissue. No epidermis. Divides sometimes prolonged into papillae, hairlike or feather processes.

Gr. together) ovary is formed (*Fig. 160*).
Ex. Pear.

Sometimes the ovary is *pedicellate*, or borne on a stalk or *gynophore* (*gynē*, Gr. pistil; *phoreo*, I bear). *Ex. Euphorbia. Dictamnus, Dianthus, Papaver.*

Sometimes the axis is borne beyond the ovaries to which the styles are united. This extension of the *thalamus* is called a *carpophore*. *Ex. Geranium.*

Along the united edges of the carpels, a thickening takes place, and a projection or process, called a *placenta*, is formed, to which the ovules are attached. *Single ovary has the placenta always*

When the carpels fold inwards, so that the placenta proceed to the centre and unite, they form a *central* or *axile* placenta bearing the ovules, which are then turned outwards (*Fig. 161*). *Ex. Iris.*

When the carpels are folded inwards, so that the placenta are only projections from the walls of the ovary, these are called *parietal* (*paries*, Lat. a wall) placenta (*Fig. 162*). *Ex. Poppy. Ononis.*

The united edges of the carpels, when they proceed to the centre, divide the ovary into cells, and each division is then called a *septum* (fence or inclosure), or a *dissepiment* (*dissepio*, Lat. I separate).

When the dissepiments formed by the inwardly-folded carpels are obliterated, the ovules are borne on a column standing free in the centre of the ovary, and called a free central placenta, as in Pink and Primrose.

When the ovary is divided into cells by means of the *dissepiments*, it is described as *bilocular*, *trilocular*, or *multilocular*, according to the number of its cells.



Fig. 160.



Fig. 161.



Fig. 162.

If undivided, and consequently but one-celled, it is said to be *unilocular*.

A cell or an ovary may contain one or more ovules. If the ovary contains but one ovule, it is *monospermous*; if it contains many ovules, it is *polyspermous*.

In the fruit of cruciferous plants, the *placentæ* prolong and form a partition, called a *replum* (Lat. leaf of a door), through the entire length of the fruit.

The ovary may be *free*, or unattached to the calyx, or any other portion of the flower, when it becomes *superior*. *Ex.* Lily.

When the calyx is united to the ovary, the calyx becomes superior and the ovary *inferior*. *Ex.* Gooseberry.



Fig. 163.

The **STYLE** (*stylus*, Lat. a pillar) is a prolongation of the carpels upwards, and is the support of the *stigma* (Fig. 163). Sometimes the stigma is sessile upon the ovary, and the style is absent.

When the style proceeds from the summit of the ovary, it is *apical* (*apex*, Lat. top); when from the side, it is *lateral*; and when from the base, it is *basilar*.

The style is commonly *cylindrical* and simple. It is sometimes *angular* or *petaloid*. It may be cleft or *bifurcate*.

After fertilization, if the style falls away, it is *deciduous*; but if it remains, it is *persistent*.

The **STIGMA** (Lat. a mark) consists of loose cellular tissue, and secretes a viscid substance upon its surface, which causes the pollen grains to adhere when shed upon it from the anthers.

The stigma is sometimes borne upon the summit of the style, sometimes laterally, and sometimes is sessile upon the ovary.

The divisions of the stigma generally correspond to

*Digynus = two carpels only

False Dissepiments are that of not formed by the union of the sides of the contiguous ovaries -

(1) Paragynata - horizontal - Ex. Cassia fistula

(2) Paragynata - i.e. Datura, Stramonium where one is true
(i.e. Thorn-apple)
some specimens

(3) Paragynata - Cruiferales, Gerard & flax (dorsal suture)
(i.e. Thorn-apple)
Astragalus (folding inwards of the dorsal suture), Oxytropis
& Phala (folding inward of the ventral suture)

ovary Stalked or Stipitate in Dianthus Caryophyllus;
Dictamnus fraxinella (Stalk called Gynophore)

The style is apical when it arises from the geometrical apex of the ovary - Ex. Primrose - natural (e.g. Strawberry) inflexed towards the side. Basilar (Althemia)

Persistent when it is a continuation of the ovary. Deciduous when it is articulated to the ovary (Ex. Sisyrinchium)

Gynobasic is when the basilar or lateral gynobasic style appears arising from the Thalamus (when the ovary is Gynobasic) Ex. Labiales. Cephalophore (See p. 73)

Compound style bifid, trifid &c or bipartite, tripartite &c ending with cleavage between two or more cham halfway between the apex & the base of the style -

Indusium = a ring of hairs below the stigma - Ex. Golden rod & Sobolewsce, style hairy in Compositae.

In orchis the Stigma is placed where the Gynobasic (i.e. the cham formed by the union of pistil & cham in Gynandria, flowers)


Stigma - is bifid, trifid or divided bifidate, lobate or
 trilobate when the divisions are thick ^{of filaments} in the compartments)

Lamellae = plate shaped like stigmae in. Dipsosaurus, allium
 lobes are (1) thick fleshy in. allium (2) feathery grasses (3)
 fringed or lacinate in. Rumex.

Thalamus

is the apex of the peduncle or pedicel or that part of
 the floral axis in axillary flower upon which the
 different parts of the flower are arranged.

1. Cylindrical - in Maguslia, Leiodendron.
2. Conical - Ranunculus, Raspberry.
3. Hemispherical - Strawberry
4. Tabular - in. Heliumbium
5. Concave - in Rose.
6. Placenta - in. Primula etc.
7. Carpophore - in. Geranium
8. Median proliferation (= a branch bearing true axes)
 in. antennated bee.
9. Hypophore - in Papier flower, Pink etc.



the number of carpels in the ovary. When these divisions are large, they are termed *lobes*. Hence we may have a *bilobate* or *trilobate* stigma.

Occasionally stigmas assume very extraordinary and peculiar forms. In *Sarracenia* they are surmounted by an umbrella-shaped appendage; in the poppy they are rags diverging upon a flat sessile shield. In some instances they are globular, in others hemispherical, or ovoid. They may be naked and smooth, or feathery.

The proper limitation of stigma should be to that portion of the pistil which receives the pollen grains. In *Iris* this is a cleft at the back of the style; in *Sarracenia* it is placed beneath the umbrella-like covering; and in many orchids it is of a hollow cup-like form.

The axis, or growing-point, sometimes extends between and even beyond the carpels, in the form of a beak, as in geraniums; or it becomes enlarged and fleshy, as in the strawberry; or conical, as in the raspberry. Whichever form it assumes, or if reduced to a mere thread, it still constitutes what is termed the *torus* (Lat. a couch), *thalamus* (Lat. a bed), or *receptacle* (*recipio*, Lat. I receive).

Instances occasionally occur in which the *thalamus* grows beyond the flower and produces leaves. *Ex.* Rose.

SECTION XXV.

THE OVULE, OR SEED-BUD.

The small bodies attached to the *placenta* in the ovary are the *ovules*, destined, after fertilization, to become the seeds.

Certain leaves, under favourable circumstances, de-

velop buds from their margins. The carpellary leaves bearing the ovules on their margins are of the same character, ovules being the buds produced by carpellary leaves.

Dr. Lindley has seen the ovules of the Columbine, and Professor Henslow the ovules of Mignonette, converted into true leaves.

Usually the ovules are contained in an ovary ; but in coniferous and cycadaceous plants the ovules are naked in the axils, or on the margins of metamorphosed leaves.

When attached immediately to the placentæ, the ovules are *sessile*; but when stalked, the prolongation by which they are attached is called a *funiculus* (a cord), or *podosperm* (seed-foot): the point of union of the *funiculus* with the ovule is termed the *hilum* (Lat. scar of a bean), or *base*.

The ovule consists of a *nucleus* and its *coverings*.




Fig. 164.

The integuments, or coverings of the nucleus, have a small aperture, which is termed the *foramen* (*Fig. 164, f*), or *micro-pyle* (*micros*, Gr. little ; *pylon*, gate).

The point of union between the base of the nucleus and the integuments is the *chalaza*.

The apex of the nucleus is turned towards this opening.

When the *foramen* is near the base of the ovule, the nucleus is inverted ; but a connection is kept up between the base of the nucleus and the base of the ovule by a process termed a *raphe* (*Fig. 164, r*).

There is a swelling at the base of the nucleus where the *raphe* is attached, which is the *chalaza* (*Fig. 164, c*), still at the base of the *nucleus*, but by inversion carried to the apex of the ovule.

In the matured ovule, or seed, the radicle is next the *foramen*.



Fig. 165.

The positions of the ovule in the ovary give rise to various terms.

When attached at the base of the ovary, it is *erect* (Fig. 165); if attached to the side, a little above the base, and with the ovule pointing upwards, it is *ascending* (Fig. 166).



Fig. 166.

If it is suspended from the top of the ovary, it is *inverted*, or *pendulous* (Fig. 167).



Fig. 167.

And if from the wall near the summit, it is, in that case, *suspended* (Fig. 168).



Fig. 168.

The curvatures of the ovule are also of importance.

It is *orthotropal* (*orthos*, Gr.



Fig. 170.

straight; *tropos*, mode) when erect (Fig. 169).

Camptotropal (*kamptos*, Gr. curved), when curved downwards towards the placenta (Fig. 170)



Fig. 169.

Anatropal (*anatrepo*, Gr. I subvert), when not only curved downwards but also grown to the lower half (Fig. 171).



Fig. 172.

Campylotropal (*kampulos*, Gr. curved), when attached at the middle (Fig. 172).



Fig. 171.



Fig. 174.

Lycotropal (*lycos*, Gr. a hollow disk), when horseshoe-shaped (Fig. 173).



Fig. 173.

Semianatropal (*semi*, Lat. half), when half free (Fig. 174).

In Cryptogamic plants, there being no ovules, the

functions of these bodies are performed by other minute analogues, called *spores*.

When the process of fertilization takes place, the mature pollen is discharged from the anthers upon the stigma, to which it is attached by means of the viscid secretion already alluded to. Whilst in this position, the pollen grains develop small tubular processes, which pass through the loose cellular structure of the stigma, down the style to the ovary. These tubes then penetrate the ovules, and discharge into them the contents of the pollen grains, and thus a union is effected between the fovilla, or fertilizing principle of the pollen, and the semi-fluid contents of the ovule. After this process, the style commonly falls away, and the ovary ripens into fruit, inclosing fertile seeds.

SECTION XXVI.

THE FRUIT.

The mature ovary constitutes the *fruit*, and the matured ovules the *seeds* of plants.

In the popular acceptance of the term *fruit*, other parts of the flower are also sometimes included; as the pulpy disk, or receptacle, in the strawberry, the united calyx and ovary in the apple and gooseberry, and the bracts and calyx united into a cupula, or cup, containing the acorn.

Important changes sometimes occur in the process of transformation of ovary into fruit. The ovary may be *unilocular*, and afterwards develop into a *multilocular* fruit, or the ovary may be *multilocular* and the fruit *unilocular*. In the ash, a two-celled ovary is re-

Fruit

Inferior - Plum, Apple, Pear, Quince etc.

Superior - Poppy, Pea.

Fruit = pericarp + Seed.

Pericarp - (1) Exocarpe = Epidermis of flower surface of Cot. leaf

(2) Mesocarpe (Sarcocarpe) = Parenchyma

(3) Endocarpe (Stone or pit) = Epidermis of upper part of cot. leaf

In Apple (inferior fruit). Exocarpe = Epidermis of under part.

Calyx. Mesocarpe = rest of calyx + oblong sac of inner lining

Endocarpe = inner lining of ovary.

Dehiscent pericarp in Bladder Senna.

In Orange - rind = Epi- & meso-carpe, partitions = Endocarpe

pulp = Saccharate parenchyma from inner lining of ovary
only the placentas only.

Dehiscent

1. Valvular
2. Transverse or Circumscissile
- (3) Porous.

Valvular Dehiscence

(a) Simple fruit - (1) Subequal. Ex. Hellebore, Collumbon
(ventral) Magnolia (dorsal) Pea, Bean, etc. (dorsal-ventral).

(2) Compound fruit with many cells

1. Septicidal 2. Loculicidal, 3. Septifragal

(b) with one cell (See text p 80)

Septicidal - valves placentiferous at their borders - Ex. Pea

Septifragal - with one cell in heart

valvular dehiscence either from above downwards or from below upwards (as in the Cruciferae)

II Transverse or circumscissile dehiscence (or poriculate)

Ex. Henbane (*Hyoscyamus*) periwinkle (*Cataglyphis*)
Monkeypot (*Lecythis* Maria?)

It is supposed that in these cases the fruits are formed by the folding of the compound leaves like that of orange or ~~lemon~~ that its lower part is formed by the unfolding out of the thalamus & the upper part by the Compellory leaf.

Other legumes separate transversely into a many pods as there are seeds - Ex.

Cordilla, Medicago, etc. etc.

III Porous - (1) Apical } Ex. Poppy, Antirrhinum, Campanula
(2) lateral }
(3) Basilar }

Kinds of fruits

1. Fruits formed by a single flower
 - a. Simple fruits
 - b. Apocarpous fruits
 - c. Syncarpous fruits
2. Fruits formed by the combination of several flowers.
 - a. Simple fruits -
 1. Legume or pod - Superior, one celled, seed on ventral suture, dehiscing by both sutures
Pea, Bean or other leguminosae.

placed by a one-celled fruit. In *Cassia fistula*, a one-celled ovary becomes a multilocular fruit.

The placentæ may also undergo a pulpy development, inclosing the seeds, as in the grape.

The envelope of the seeds is called the *pericarp* (*peri*, Gr. around ; *karpos*, fruit), and ordinarily consists of three layers.

The *epicarp* (*epi*, Gr. outside) is the external layer.

The *mesocarp* (*mesos*, Gr. middle), the intermediate, and—

The *endocarp* (*endon*, Gr. within), the internal layer.

Sometimes the three are blended, as in the nut ; at others separable, as in the peach. In the latter instance the skin is the *epicarp*, the pulp the *mesocarp*, and the hard shelly covering of the seed, or kernel, the *endocarp*.

The inner union of the carpellary leaves is called the *ventral* (*venter*, Lat. the belly) *suture* (*sutura*, Lat. a seam) ; the outer, corresponding to the rib of the leaf, the *dorsal* (*dorsum*, Lat. the back) *suture*.

The sutures may be so firmly united as not to give way on the ripening of the fruit : it is then said to be *indehiscent* ; when the sutures open to allow the seeds to escape, the fruit is termed *dehiscent* (*dehisco*, Lat. I open). By dehiscence the pericarp becomes divided into parts called *valves*.

Indehiscent fruits may be *dry*, as in the filbert, or *fleshy*, as the apple.

Dehiscent fruits may separate by one suture or both, and they may carry the placenta with them, or leave it standing in the centre. In the latter case it is called a *columella*.

The separation may proceed from the base to the apex, or from the apex to the base. Transverse, or operculate divisions, sometimes occur.

Dehiscence may also take place by means of orifices or *pores*, as in the poppy.

Fruit composed of a single carpel may dehisce by one suture only, or by both together; this kind of dehiscence is called *sutural*. *Ex.* Pea.



Fig. 175.

Fruit composed of several united carpels may dehisce through the dissepi-ments, so that it is again resolved into its original carpels. This mode of dehiscence is *septifragal* (*septum*, Lat. inclosure; *cædo*, I cut) (Fig. 175). *Ex.* Foxglove. *Adonis*.



Fig. 176.

If the carpels remain united at their edges and dehisce through the back of the loculaments, each valve being formed by the halves of contiguous cells, it is *loculicidal* (*loculus*, Lat. cell) (Fig. 176). *Ex.* Yellow Iris.



Fig. 177.

If the fruit opens by the dorsal suture and the septa remain attached to the centre, it is called *septifragal* dehiscence (*septum* and *frango*, Lat. I break) (Fig. 177). *Ex.* *Salvia*, *Adonis*.

Fruits may also be the result of one flower, as in ordinary instances; or of several united, as in the Pine-apple.

When the bracts and floral envelopes are united in the fruit, it is termed *anthocarpous* (*anthos*, Gr. flower; *karpos*, fruit).

Fruits formed out of one or more free carpels are said to be *apocarpous* (*apo*, Gr. separate). *Ex.* Bean.

When several carpels are united, they constitute a *syncarpous* (*syn*, Gr. together) fruit. *Ex.* Orange.

Indehiscent fruits may be classed in two divisions, containing *one* or *more* seeds respectively.

Indehiscent fruits containing *one seed* may be again subdivided into *dry* and *fleshy*.

- Stomatoc
indiscut of legume, contorted (In Lucuma medicap), Cyath
cap (Scorpiurus sulcata) Curved like avocado eg.
Spurious discut - ment - Astragalus, Phase (vertical from
Cassia fistula (transverse) [dorsal & ventral Sutures req,
Indiscut - Cassia, Arachis, Pterocarpus,
 1. Lomentum or lomentaceous legume - A modification
 of the legume which is contracted in a moniform
manner. Ex. Hedysarum, Leuca Sapoteae.
 2. Drupe - Superior, one celled, indeciscent, one or two seeded
pulpy mesocarp, stony endocarp. Ex. Plum, cherry,
Peach &c.

- Uticle - Superior, one celled, few seeded, usually in
indeciscent with their membranes inflating pericarp.
 Ex. Chenopodium -
 1. Aposporous fruits
 1. Follicle - Superior, one celled, one or many seeded,
dehiscing by ventral Suture only (or therefore
one valued) - (N.B. legume is distinguished
 from follicle by being two-valved i.e. dehiscip
by both Sutures); or one suture only as Magnolia
(dorsal). Ex. Chumbina, Hellebore, Katapu
Monett, (or more black); Pasony, periwinkle
(leaf) hairs dendron, Magnolia (numerous)
 2. Achaenium or Achene - Superior, one celled, one seeded
indeciscent. Ex. Urnalis, Arenaria, Labiatae
Pyragmata. (A chene rarely single). Stachys,
Stachys

3. Elaeis - A number of drupes aggregated together. In Raspberry. (By some a number of Achenes ~~forming~~ more than a single series are also called elaeis - In Ranunculus, Rose, Honeysuckle etc -

C. Synsarpous Fruits (ovary compound)

1. Superior or Inferior

(α) Indehiscent (dry)
(β) Dehiscent (dry)

2. Inferior. (γ) fleshy indehiscent

2. Inferior (α) Dry indehiscent

(β) Dry dehiscent

(γ) fleshy indehiscent

1. Superior

a. Dry indehiscent -

1. Caryopsis - Superior, one celled, one seeded, indehiscent with dry membranous pericarp - (In Achene the pericarp is not united to the seed but in Caryopsis it is - The latter is considered compound then being two or more stigmas) In Wheat, Barley etc.

Generally in grasses.

2. Samara - Superior, 2 or more celled, dry indehiscent, few-seeded, pericarp winged (i.e. a combination of Achenes with wings ~~in~~ united) In Maple, Ash, Elm etc. By some each winged portion

is called a Samara.

3. Carcerule - Superior, many celled, each cell dry indehiscent one or few seeded (This is as if a number of achenes without wings united)

Dry indehiscent fruits with one seed are :—

ACHENIUM (*a-chaino*, Gr. I open not).

With a pericarp separable from the seed (*Fig. 178*). *Ex.* Sunflower.



Fig. 178.

CARYOPSIS (*karua*, Gr. a nut; *opsis*, appearance). With a pericarp inseparable from the seed. *Ex.* Wheat.

UTRICLE (*uter*, Lat. a bladder). With an inflated pericarp. *Ex.* Chenopodium.

GLANS (Lat. an acorn). With a hardened pericarp, accompanied

by bracts at the base, or inclosed in an involucre (*Fig. 179*). *Exam.* Acorn, Chestnut.



Fig. 179.




Fig. 180.

Samara (Lat.

seed of an elm). With a winged pericarp (*Fig. 180*).

Ex. Sycamore, sometimes included under *Glans*.

Pulpy indehiscent fruits with one seed :—

DRUPE (*drupæ*, Lat. unripe olives). With a succulent, or fibrous mesocarp, and hard endocarp (*Fig. 181*).

Ex. Cherry.



Fig. 181.

Etærio (*etæi-*

ros, Gr. a companion). Several small drupes aggregated together (*Fig. 182*).

Ex. Raspberry.



Fig. 182.

Pulpy indehiscent fruits with more than one seed are :—

BERRY (*bacca*). A succulent fruit, in



Fig. 183.

which the seeds are immersed in a pulp mass (Figs. 183, 184). *Ex.* Gooseberry.

Pepo (*pepon*, Gr. a pumpkin). With three or more parietal placentæ bearing seeds, often with a central cavity. *Ex.* Melon.



Fig. 184.

Hesperidium (golden fruit of Hesperides) is a name given to the fruit of the orange tribe, in which the epicarp and mesocarp form a separable rind.

N.B.—The *Pepo* and *Hesperidium* are modifications of the Berry.




Fig. 185.

POME (*pomum*, Lat. an apple). The calyx, with the epicarp and mesocarp, form a fleshy mass, and the endocarp is scaly or horny, forming separate cells inclosing the seeds (Fig. 185). *Ex.* Apple.

In the annexed engraving, C represents the remains of the stamens, and free limbs of the calyx; T, the

calyx tube, become succulent and adherent to E, the epicarp; M indicates the mesocarp; N the endocarp, and S the seeds.

Dehiscent fruits are all dry.

Those which are apocarpous are:—

FOLLICLE (*folliculus*, Lat. a little bag). A carpel having no dorsal suture, and dehiscing by the ventral suture (Fig. 186).



Fig. 186.

LEGUME (*legumen*, Lat. pulse). A carpel having dorsal and ventral sutures, and dehisc-

4. Amphisarca - Superior, many celled, indehiscent
many seeded woody externally, pulpy internally -
Ophalocarpus, Adansonia, Crescentia.

(B) Dry dehiscent -

(1) Kapsule - Superior, one or more celled, many
seeded, dry dehiscent.

~~like~~ Colchicum, Dalura (Valerian) Poppy, Antirrhinum
(porous), humpreel, Henbane (Ornithoglossum - when
the fruit called Pypis) Lychnis, Mignonne (has
one celled - gentiana, Heart's Ease

Two or more celled - Colchicum, Dalura, Scrophularia
Twisted (Heliclytes), Stilted (Glicium anisatum)

Very common fruit - Papaveraceae, Caryophyllaceae
Portulacaceae, Scrophulariaceae, Liliaceae,
Gentianaceae &c.

Regma = capsule bursting with elasticity -
Picinus, Hura crepitans

2. Siliqua - Superior, one or two celled, many seeded,
long, narrow, dehiscent by 2 valves from
below upward, leaving the seeds attached to
the two parietal placentas. When several
connected by a replum. Wall flower &
many cruciferae (Stigmas opposite the
placentas).

3. Siliqua - Broad, short Siliqua - Shepherd's Purse
1 & 3 only found in the Cruciferae. Sometimes
m

9. Fleshy indehiscent

1. Idiosperidium. Superior, many celled, few seeded, indehiscent with separable rind from of Epic + mesocarp of pulpy seed embedded in the pulp - Ex. orange, lemon etc.
2. Tryma. Superior, one celled, one seeded, pulp indehiscent with separable rind (this mesocarp with hard endocarp from the inner lining of which spines or septines extend to divide the seed into deep lobes. Ex. Walnut (very similar to Drupe except that it is formed from a carp. ovary)
3. Nauculanium - Superior berry - Ex. grape

II Synsarpous - Inferior

- ~~1. Gynocarp - inferior, dry, indehiscent, 2 celled, 2 seeded~~
~~id. Dry indehiscent.~~
1. Ochrocarp - Synsarpous, inferior, dry, indehiscent, 2 or more celled + 2 or more seeded - Ex. Hemlock, Forsk. torrey & other umbelliferous plants. (ie two or more inferior achenes joined together)
 2. Cypselis - ~~Ham~~ inferior, Synsarpous achenes with hairs - Ex. all Compositae.
 3. Nat - inferior, dry, hard, indehiscent one celled, one or two seeded.

ing by both or either (*Fig.* 187).

Those which are syncarpous are:—



Fig. 187.

CAPSULE

(*capsula*, Lat. a little chest). A dry fruit opening by valves or pores.

Valvular capsule. *Ex.* Foxglove.

Porous capsule (*Figs.* 188, 189). *Ex.* Poppy.

Sometimes the capsule dehisces by a lid, in which case it is called

a *Pyxidium* (a little box) (*Fig.* 190). *Ex.* Henbane.

Siliqua (*siliqua*, Lat. a husk or pod). A variety of capsule

opening by two valves, detached from below upwards, leaving the seeds attached to both sides of a *replum* (*Fig.* 191).

Ex. Wallflower.

Silicle (*silicula*, dimin. of *siliqua*), a pouch or short pod. *Ex.* Candy-tuft.

Anthocarpous fruits:—

SOROSIS (*soros*, Gr. a cluster).

A succulent anthocarpous fruit. *Ex.* Pine-apple.

SYCONUS (*sucon*, Gr. a fig).

A pulpy hollowed axis, inclosing the achenia of numerous flowers (*Fig.* 192). *Ex.* Fig.

STROBILUS (*strobilus*, Gr. a fir-cone). A spike



Fig. 188.




Fig. 189.



Fig. 190.



Fig. 191.




Fig.




Fig. 193.

covered with scales or bracts, with seeds at their base (Fig. 193). *Ex.* Fir cone. Hop.

SECTION XXVII.

SEED.

The *seed* is the matured ovule.

When the seeds are inclosed in a seed-vessel, as in most flowering plants, such plants are termed *angiospermous* (*aggeion*, Gr. a vessel ; *sperma*, a seed).

In conifers the seeds are naked, or not provided with a seed-vessel ; hence such plants are termed *gymnospermous* (*gymnos*, Gr. naked).

In mignonette the seed-vessels open very early and expose the seeds.

The seed has also its own coverings, independent of the pericarp.

The general covering of the seed is called the *spermoderm* (*derma*, Gr. covering). It consists of an external membrane, the *testa* (shell) or *episperm*, and an internal lining or *tegmen*, the *endopleura* (*endon*, Gr. within ; *pleura*, side).

The *episperm* or *testa* is cellular, and is often coloured. In the Circassian bean and crab's-eye bean they are of a bright red, and in cotton they are covered with hairs.

The *endopleura* is cellular also. It is generally thin and transparent, and is often closely applied to the embryo.

An additional covering to the seed is present in a few instances, which is derived from an expansion of the *placenta*, after the fertilization of the ovule. This

constitutes the *mace* of the nutmeg, and is termed the *arillus*.

The attachment of the seeds, as of the ovule, is called a *funiculus* (Lat. a little cord). Occasionally it is of such length that the seeds are pendent from them as from a cord. *Ex.* Magnolia.

The scar left upon seeds after detaching them from the funiculus is the *hilum* (Lat. the scar of a bean).

The seeds sometimes contain, interposed between the embryo and the seed-coat, a substance called *albumen* ^{Fig. 2} (Lat. white of an egg), which consists of starch, oil, and nitrogenous compounds.

Seeds in which the albumen is present are called *albuminous*, and those from which it is absent, *ex-albuminous*.

The albumen forms the great bulk of the cocoa-nut; and the hard substance of the vegetable-ivory nut, and the horny substance of the coffee berry are alike albumen.

The EMBRYO (*embryon*, Gr. a fœtus), or young plant, consists of *radicle* (*radix*, root), or end of axis (*r*), *plumule* (*plumula*, feather), or young stem (*q*), and *cotyledons* (*c*), or seed-leaves (*Fig.* 194).

Sometimes the embryo occupies the entire seed, and at others only a portion. The seeds of some plants contain more than one embryo.

The plumule may generally be seen lying between the cotyledons.

The class of plants now associated under the name of *Endogens* correspond to another division formerly adopted, called *monocotyledons* (*monos*, Gr. one), from the fact of their having but one cotyledon.

Plants having two cotyledons, and which correspond to Exogens, were called *dicotyledons*.




Fig. 194.

Embryos containing more than two cotyledons are said to be *polycotyledonous*.

Embryos deficient in cotyledons, as in *Cuscuta*, which has no leaves, are *acotyledonous*.

Cotyledons are generally entire, but sometimes lobed; they are either flat or folded in various ways.

The cotyledons may be *conduplicate* (Lat. folded together) or folded laterally (Fig. 195).

Or *reclinate* (Lat. bent downwards), folded from apex to base (Fig. 196).

Or *convolute* (Lat. rolled over), rolled up laterally (Fig. 197).

Or *circinate* (*kirkos*, Gr. a circle), rolled up like the young fronds of ferns (Fig. 198).

Or *contortuplicate* (Lat. twisted into plaits), crumpled (Fig. 199).

When the cotyledons have their faces applied to each other, and the radicle is folded on their edges, so as to be *lateral*, the cotyledons are said to be *accumbent* (*accumbo*, I lie at the side) (Fig. 200).

When the cotyledons are applied in the same manner, and the radicle is folded on their back so as to be dorsal, the cotyledons are *incumbent* (*incumbo*, I lie on the back) (Fig. 201).

The two latter forms and terms are applicable to the seeds of cruciferous plants, in which the distinction becomes one of importance.

The seed contains the germ of a new plant. Its function is the reproduction of its species. When matured, the seeds are detached from the parent plant and dispersed in various ways. Some are expelled to



Fig. 195.



Fig. 196.



Fig. 197.



Fig. 198.



Fig. 199.



Fig. 200.



Fig. 201.

a considerable distance by the elastic bursting of the seed-vessel; others are supplied with appendages, by means of which they are readily wafted through the air; and others fall to the ground and germinate in the immediate neighbourhood of their parent.

Seeds are sometimes produced in immense numbers. One capsule of the poppy has been found to contain 40,000 seeds.

SECTION XXVIII.

GERMINATION.

The period during which seeds retain their vitality is very variable. Melon-seeds have been known to vegetate after forty years. Seeds capable of germinating have been stated to have been found in a Roman tomb fifteen or sixteen centuries old. Seeds have also germinated which were found in a sand-pit twenty-five feet deep, where they were supposed to have been deposited at least 2,000 years ago. Other seeds require planting soon after ripening to insure germination.

Certain conditions are essential to germination.

Moisture is of great importance. Dry seeds will not vegetate. Dampness is necessary to enable the young plant to burst its hard covering.

Temperature should be such as is suited to the character of the plant to be reared.

Air is also essential. Seeds buried deep in the soil do not germinate in that condition.

Darkness is also favourable. Seeds germinate best when excluded from the light.

Important internal changes take place in seeds during germination. Starch is converted into dextrine, or sugar, by oxidation, and diastase is developed

During these changes considerable heat is evolved; substances which were before insoluble become soluble. The seed is swollen, the embryo enlarges and ruptures the integument, protrudes, and the growth of a new plant commences. During its early stages the young plant is nourished by the cotyledons until the primary leaves are formed; the radicle invariably proceeding downwards, and the plumule upwards from the *collum* or neck, which lies intermediate between the two.

The germination is *exorhizal* (*exo*, Gr. outwards; *rhiza*, root) when the radicle is directly prolonged into the root. *Ex.* Maple.

It is *endorhizal* (*endon*, Gr. within) when the roots pierce through the integument at the extremity of the radicle. *Ex.* Wheat.

The ruptured integument surrounding the base of the young root as with a sheath, is termed the *colcorhiza*.

In acotyledons, there being no distinct embryo in the spore, it germinates from any part of its surface, and is termed *heterorhizal* (*eteros*, Gr. diverse).

Some seeds will germinate within twenty-four hours of planting. *Ex.* Cress.

Seeds with a stony pericarp will sometimes lie dormant in the soil for upwards of a year.

Some seeds will germinate while still attached to the parent, and young ferns commence their existence attached to the fronds of their parents.

The leaves of *Bryophyllum calycinum*, when placed upon damp soil, will produce buds, which develop into young plants, upon their edges.

In other plants buds are developed in the axils of the leaves, which are capable of germination under favourable circumstances.



SECTION XXIX.

REPRODUCTION OF ALGÆ.

Seaweeds are furnished with a double system of fructification. The majority of the red series have this double fructification, the *capsular* on one plant, and the *granular* on another. A great modification of shape exists, but there is a unity of principle, so that they are at once alike and unlike.

Dr. Lindley says, "I am very much inclined to adopt the opinion that the two sorts of fructification observable in algæ are the first attempts at the agency which, in higher plants, performs the office of sexes, without, however, having their qualities established, and each capable of producing a new plant without the aid of the other."

In the *capsular* fruit, we meet with capsules of different forms, according to species, containing tufts or bundles of spores. The capsules vary in size and form; one kind is termed a *ceramidium* (*keramos*, Gr. a little measure), another is called a *coccidium* (*kokkos*, Gr. a berry; *oidos*, resemblance), and others are termed *favellæ*.

In *Fig. 202 (1)* is represented a magnified portion of a frond of *Polydiphonia* bearing a *ceramidium*, and *Fig. 202 (2)*, the *ceramidium* only, still more magnified.

The *granular* fruit consists of *tetraspores* (*tetra*, Gr. four), or granules, which, on being examined by the microscope, are found to be composed of four spores or seeds, united together. These granules are imbedded in the terminal por-




Fig. 202.

tion of the frond or its branchlets, or occasionally supported on short pedicels, collected in *conceptacles* or *stichidia* (*stichos*, Gr. a row).




Fig. 203.

In Fig. 203 (1) is shown a portion of a frond of *Polysiphonia*, bearing tetraspores, and Fig. 203 (2), a tetraspore removed and more highly magnified.

Nemathecia (*nema*, Gr. a thread) are warts, concealed under leafy processes, and composed of thin moniliform filaments. This is a kind of fructification concerning which but little is known.

Antheridia (*anthos*, Gr. a flower) are not uncommon. They are little cells, in each of which is contained one or more *spermatozoids* (*sperma*, Gr. seed ; *zoon*, life).

SPERMATOZOIDS are thread-like bodies, varied in form, but generally spiral, and furnished with movable cilia. They are very small, and, after escaping from their parent cells, continue to move about for some time, until they have sought out and fecundated the spores.

The seed consists of a cell filled with a dark-coloured semi-fluid mass, which is called the *endochrome* (*endon*, Gr. within ; *chroma*, colour). This seed produces a perfect plant like its parent ; but the process of fecundation is still involved in mystery.

SECTION XXX.

REPRODUCTION OF LICHENS.

In Lichens (*lichenos*, Gr. a letter or wart) we have a double reproductive system.

1 The *primary* consists of organs called, respectively, *apothecia*, *spermagones*, and *pycnides*.

Apothecia (*apothèkè*, Gr. a repository) generate and protect the spores from which new plants are developed.

Spermagones (*sperma*, Gr. seed; *gonè*, generation) produce and discharge bodies called *spermatia*, whose function appears to be the fertilization of the spores.

Pycnides (*puknotès*, Gr. denseness), which give rise to ³spore-like bodies called *stylospores*, whose function is undetermined.

2 *APOTHECIA* appear like peltate disks on the surface of lichens (*Fig. 204, a*).

They are sometimes open (*Fig. 204, b*), as in a section of the lichen family, called *gymnocarpous* (*gumnos*, Gr. naked; *karpos*, fruit), and sometimes closed, as in the *angiocarpous* section (*aggos*, Gr. a vessel).

In *gymnocarpous* lichens, the *apothecium* consists of a more or less flattened disk called a *thalamium* (*thalamos*, Gr. a receptacle), which contains the spores, and an *exciple* (*excipio*, Lat. to receive), or envelope, which




Fig. 204.

surrounds and protects it. In this case the upper surface of the *thalamium* is exposed (*Fig.* 204, *b*).

In *angiocarpous* lichens, the *exciple* incloses the *thalamium* in a globular envelope, from which the spores issue, when ripe, by means of a pore or fissure. In these instances the organ is termed a *perithecium* (*peri*, Gr. around).

The *thalamium* consists of a series of elongated cells, placed on end, closely packed together and united by means of a tenacious mucus. These cells arise from the smaller, somewhat irregular cells, which constitute the base of the *thalamium*, and called the *hypothecium* (*upo*, Gr. under).

The elongated cells thus packed together upon the hypothecium, are of two kinds; *thecæ* (*thèkè*, Gr. a sac), and *paraphyses* (*paraphusis*, Gr. an offset).

Thecæ are the parent cells of the spores. They are elongated cells, containing, at first, a mucous matter which becomes granular, and ultimately consolidates into eight, four, or two spores. When the spores are mature, the *theca* ruptures, and the spores escape.

Paraphyses are elongated, club-shaped bodies, thinner and longer than *thecæ*. They consist of a series of six or eight cells, attached end to end, like a string of beads, and contain mucus; the terminal cell being larger, and, with its contents, coloured. They precede the *thecæ* in order of development, have a distinct structure, and, apparently, a distinct function not yet clearly defined.

The *spores* (*spora*, Gr. a seed) are generally ovoid in form, and consist of one or more cells. In their early stage they are generally colourless, acquiring a yellow tint as they ripen. The young spore contains mucus and granular matter, sometimes mixed with globules of oil. At maturity these become fused into a mass, partaking frequently of an oily nature.

Spermagones^{ia} are usually minute capsules, imbedded in the *thallus*, and opening on its surface by pores. From the inner wall a number of delicate filaments project, called *sterigmata* (*sterigma*, Gr. a support); from their ends or sides the *spermatia* are developed. These escape from the *spermagone*, when mature, which takes place long before the maturity of the *spores* (Fig. 204, c).

These *spermatia* are minute linear bodies, colourless and transparent; not possessed of cilia or other appendages, but often exhibiting molecular motion.

The *spermagones* only appear as black point-like bodies, scattered or clustered on the surface of lichens. No proof has yet been obtained of the fecundating influence of *spermatia*; it is only still matter of probability.

Pycnides^{ia} are rare bodies, differing from *spermagones* in being larger, and supporting, upon pedicels, bodies resembling spores, and called *stylospores* (*stulos*, Gr. a pillar). These are oblong and colourless, and contain granular matter.

The *secondary* system consists of *gonidia*.

Beneath the upper or cortical layer of the thallus of lichens, is a thin bright-green layer of globular cells, loosely aggregated, called the *gonidic* layer. These isolated cellules are the *gonidia*. They assume the offices, and partake of the character, of both vegetative and reproductive cells, and are peculiar to the thallus of lichens. They frequently burst through the outer layer of the lichen and appear as a powdery mass on the surface, termed *soredia*, giving it a powdery or frosted appearance. They appear to be cellular buds or reproductive cells, destined, without preparatory fertilization, to multiply the individual.

SECTION XXXI.

REPRODUCTION OF FUNGI.

The common mushroom is a well-known example of the group of cryptogamic plants included under the general name of Fungi. Let us employ this plant to illustrate our descriptions.

In this instance we have a number of individual plants, composed of thread-like filaments, which constitute what is called the *mycelium* (*mukès*, Gr. fungus), or *spawn*. This is either concealed beneath, or appears on the surface of the matrix, upon which the fungus vegetates.




Fig. 205.

From a community of the individual plants which constitute the *mycelium* (*Fig. 205, m*), a common pedicel, or *stipe* (Lat. a trunk), proceeds (*Fig. 205, s*), surmounted by a round knob or protuberance, which enlarges and assumes the form of a cap (*Fig. 205, p*), and is called the *pileus* (Lat. a cap). In the process of growth, whereby the

rounded summit of the stipe is converted into the cap-like form, the wrapper which envelops it ruptures, or breaks away from the stipe, leaving a ring, or *annulus* (Lat. a ring), showing the point of attachment, around the stipe.

The under portion of the *pileus*, in agarics, consists of a number of plates or *gills* radiating from the stipe. These are covered with a membrane called the *hymenium* (*umèn*, Gr. a membrane), which bears the spores, or reproductive organs.

SECTION XXXII.

†*Gay*

REPRODUCTION OF MOSSES.




Fig. 208.

The spores of mosses are borne in spore-cases, or *thecae*, which dehisce in the split-mosses by valves (Fig. 208, 1). In urn-mosses the dehiscence is operculate (Fig. 208, 3). The organs of fructification in mosses are of two kinds. The most conspicuous consists of a *sporangium*, or spore-case, an urn-shaped capsule, supported upon a stalk or *seta*, and, at first, covered with a *calyptra*, or hood (Fig. 208, 2), which is sometimes *dimidiate*, or split on one side; at others *mitriform*, or cleft at the base. When the *calyptra* is removed, the *sporangium* is found to possess a lid, or *operculum* (Lat. a cover), which, when it falls off, shows the mouth of the urn, or *sporangium*, to be sometimes crowned with a *peristome* (*peri*, Gr. around; *stoma*, mouth), or fringe, beneath which is an elastic *annulus*, or ring; and sometimes naked.

In the centre of the urn is a *columella*, and occasionally a transverse *septum* or *epiphragm* (*phragma*, Gr. partition) is interposed between the *operculum* and the contents of the urn. Within the urn are contained a number of spherical spores, each of which divides into four sporules. In some instances, spiral filaments (Fig. 208, 4), called *elaters* (Gr. impellers), are also present, *i. e.* in the moss-like plants called *Hepaticæ*.

When a fleshy protuberance occurs at the base of the theca, and on one side, it is called a *struma* (Lat. a swelling).

The urn usually grows from a fleshy tubercle called an *apophysis* (*apophysis*, Gr. excrescence).

To the other kind of organs, the name of *antheridia* or *staminidia* has been given. They are found in the axils of the leaves, and consist of membranous sacs, opening at their summit, and discharging a turbid fluid, and *phytozoæ* (*phuton*, Gr. a plant; *zoon*, an animal; the antheridia are generally in clusters mixed with *paraphyses* (Gr. an offset),

A *phytozoa* is a peculiar body, rolled up in a circular or spiral manner, which exhibits movements at a certain period of its existence, from which the name has originated (*Fig. 209, b*).

The Rev. M. J. Berkeley says: "Perennial mosses and *Hepaticæ* so far agree with phænogams, that there is a fresh crop of fruit every year; but then the result of impregnation is not a seed, but an organ, which produces the reproductive bodies or spores, much after the fashion in which pollen grains are generated. Ferns and club-mosses, though living to a hundred years, and producing a crop of spores each year, are impregnated but once, and that before the plant has assumed its true habit. The result of impregnation in these, is something more or less resembling the embryo of a phænogam, and, like that, produces, by progressive development, a perfect plant."



Fig. 209.

SECTION XXXIII.

REPRODUCTION OF FERNS




Fig. 210.

The substance of the fronds of ferns is traversed by veins which bear from certain determinate points the fructification (Fig. 210, 1). The part to which these organs are attached is called the *receptacle* (*recipio*, Lat. to receive): a correct appreciation of the nature and position of this receptacle is necessary in a determination of groups and species.

Each mass or cluster of seed-cases is called a *sorus*. This consists of a number of *spore-cases*, *sporangia* (*spora*, Gr. a seed; *aggos*, a vessel), or *thecæ* (*theke*, Gr. a sac), minute roundish bodies, often nearly surrounded by an elastic band or *ring* (Fig. 210, 2), which is sometimes wanting. When mature, the ring, by its elasticity, causes the *thecæ* to burst and discharge the *spores*, as a minute, almost impalpable dust (Fig. 210, 3, *magnified*).

When the ring is present, the *sporangia* are stalked; but when absent, they are sessile.

The *thecæ* sometimes arise from the surface of the frond, while at other times they spring from below, having a cuticular covering in the form of an *indusium* (Lat. a shirt), or *involute* (*involutum*, Lat. a cover).

The margin of the frond may be so folded as to cover the *thecæ*, and at times the whole frond is converted into clusters of *thecæ*.

The *indusium* is commonly a membranaceous scale of the form of the *sori*, inclosing the young *thecae*; but ultimately becoming ruptured and thrust back or cast off (*Fig. 211*). In some species the *indusium* is cup-shaped, containing the spore-cases; but this form is rare in British ferns.

The *sori* are said to be *marginal* when they project beyond the margin of the frond, and *dorsal* when placed on some part of the under surface of the frond.

Certain cellular *papillæ* (Lat. teats) on the margin, or upper surface of the fronds, have been considered by some as antheridia, each of the cells containing a spiral fibre. But this question cannot be considered as yet settled beyond dispute. There are some who contend for a duality of organs in ferns analogous to the anthers and ovules of flowering plants, and state that among the *thecae*, filaments are found which are equivalent to stamens.

The following is the most generally accepted history of the reproductive development of ferns. The spores at first germinate by the protrusion of a tubular process, which enlarges into a small leafy plate, or *prothallus*, with filamentous rootlets. When completely formed, this *prothallus* bears on its under surface bodies of two kinds, viz., *antheridia* and *archegonia*. The *antheridia* are unicellular structures, in which a second cell is formed, the interior of which becomes divided into a number of minute *sperm-cells*. When ripe, the sperm-cells escape, each containing a *spermatozoid*. The *archegonia* are formed of four tiers of cells, with a central canal leading to an *embryo-sac*:



Fig. 211.

in which lies a germinal body, having somewhat the character of an ovule, which ultimately becomes fertilized by the entrance of the spermatozoids liberated from the antheridia. The fecundated embryo, or ovule-like body which lies in the embryo sac, then develops as a bud, and produces true leaves, which mature into a plant resembling the original parent from which the spore first sprung.

SECTION XXXIV.

VEGETATIVE PHENOMENA.

ROTATION.—This is a movement of fluids confined to the cells themselves, and is the result of vital action. In young cells, and in the cells of aquatic plants of all ages, this movement has been observed. It proceeds in each cell independently of its neighbour. In some the movement will be slow, in others rapid, and is generally most evident near the circumference, passing up one side and down the other.

CIRCULATION is the general motion, in plants, of fluids absorbed by the roots upwards into the leaves and back again to the stem. This current enters by the extremities of the roots, passes up by the woody portions of the stems of exogenous plants, and the vascular bundles of endogens, into the leaves, where much of the water which the fluid, hitherto termed *ascending sap*, contained, is evaporated under the influence of light and heat, and the thickened fluid enriched by carbonic acid absorbed from air, now called the *descending sap*, returns by the liber of the bark to the stem, whence it passes horizontally by means of the medullary rays, and is diffused through the plant, depositing, in its progress, the materials essential to its growth and vigour.

ASSIMILATION.—In the upward course of the sap, its

water and carbonic acid are partly decomposed and deposited. Upon reaching the leaves exposed to light, water is given off, carbonic acid absorbed, and chlorophyll formed by the decomposition of carbonic acid and water; oxygen is returned to the atmosphere, and carbon, nitrogen, and hydrogen assimilated for the use of the plant. *Light* is the main cause of this assimilating process, which, therefore, does not take place during the night. The parts of plants excluded from the light by artificial means are blanched, or rather prevented forming the colouring and other secretions for which light is essential. *Ex.* Endive.

The average rate of perspiration in plants has been estimated as equal to seventeen times that of a man.

During the night, plants re-absorb oxygen from the air and part with a small quantity of carbonic acid. It is a vulgar error to suppose that the latter is given off in sufficient quantity by plants growing in a room to have any appreciable or detrimental effect upon the atmosphere.

HEAT is developed by plants during growth, sometimes to a considerable extent. Tubers and bulbs have been found to possess, during winter, a temperature 6° or 7° above the external air. In germination, much heat is evolved, and again during flowering. The spathes of araceous plants, which contain many flowers, have been subjected to experiment. In some of these the thermometer has risen at sunrise to 30° above the surrounding air. The bulb of a thermometer plunged 11 inches deep in the trunk of a walnut-tree rose 2° or 3° . This, however, was due to the absorptive power of the trunk for the sun's heat. A species of *Colocasia* was found to attain a heat of 10° above the atmosphere in the middle of the day. The exciting cause of this evolution of heat may be found in the oxidation going on in plants at all times, but with greater vigour during the flowering season.

IRRITABILITY.—Some of the most simple forms of Algae have an oscillatory motion, and the spores of others move about for some time, as it has been supposed, by means of vibratile ciliæ. Plants of a higher order exhibit movements incidental to peculiar states of the atmosphere. Some flowers are hygrometric, others close at stated periods. The sensitive plant (*Mimosa sensitiva*) closes up its leaves when touched. The leaves of *Desmodium gyrans* move about by rapid jerks. A species of *Dionea*, called Venus's fly-trap, folds up if an insect alights. The leaflets of the wood sorrel fall back at the approach of night. Many other plants exhibit staminal movements. In the Passion-flower the styles move towards the stamens. Thus, in some part or other, every plant exhibits irritability.

LUMINOSITY has been noticed chiefly in Fungi. A species of Agaric, in Western Australia, is said to afford a phosphorescent light of sufficient intensity to read by. Decaying wood, called *touchwood*, caused by the mycelium of fungi, is often phosphorescent; and in Brazil certain mushrooms give out a light like that of fire-flies. Luminosity has also been attributed to some phanerogamous plants; but the evidence of its existence is questionable.

ODOURS are given off from some plants only during the day, from others only at night. Certain plants, as the woodruff, are fragrant only when dried, others only whilst in the fresh state. From the investigations of certain German botanists, we learn that white flowers have the largest number of fragrant species, and that a large proportion of yellow and brown flowers possessing any odour, are disagreeable. Of 1,193 white flowers examined, 187 were odoriferous, whilst only 12 were disagreeable. Meteorological changes greatly influence the odours of plants.

SECTION XXXV.

EPIPHYTES AND PARASITES.

Plants which grow without attachment to the soil are called *Epiphytes* and *Parasites*.

Epiphytes (*epi*, Gr. upon; *phuton*, a plant), or air-plants, are generally Orchids, or their allies. They are found in tropical climates attached to the surface of trees, and are sometimes nearly covered with a mass of showy flowers. Some of these aerial plants develop long root-like processes, clad with a green epidermis containing stomata. These serve all the purposes of leaves, and would rather deserve to be considered as such did they not often possess a layer of delicate air-cells covering the epidermis, which has caused them to be termed by some botanists *coated roots*.

Epiphytes derive very little of their nourishment from the matrix on which they flourish, but chiefly from the medium in which they grow.

Parasites (*para*, Gr. beside; *sitos*, food) grow and prey upon other plants. They are more numerous than epiphytes, and are found in all climates. Some of these are flowering plants, and others without any such organs. Some are brownish, without true leaves, but provided with scales. Others are green, and possessed of leaves. To the former the dodder belongs, and to the latter the mistletoe.

A remarkable group of parasites are found in the Eastern Archipelago and South America, which produce immense flowers, but no leaves. These extraordinary plants are of very rapid growth. It is reported of them, that the *Rafflesia*, which, when full grown, is a yard across, and when unexpanded is

as large as a middle-sized cabbage, only takes about three months for its complete formation.

Some parasites attach themselves to the roots of plants by suckers. *Ex.* Broom-rape.

Others insinuate their extremities through the bark into the wood of trees, and flourish on their branches. *Ex.* Mistletoe.

A great many Fungi are true parasites, producing diseased states of corn, and dry-rot in wood, or are found growing upon, but without any great injury to, other plants; as *Polyporus*, &c.

SECTION XXXVI.

GALLS AND EXCRESCENCES.

Galls and excrescences of various kinds, forms, and sizes, are met with on different parts of plants, resulting from disease in some form or other, whether induced by the puncture of an insect or not.

The artichoke gall of the oak is a bud which has been checked in its development by the puncture of an insect, and which is then converted into a gall consisting of imbricated scales, and resembling a small cone.




Fig. 212.

The Chinese and Japanese galls (*Fig.* 212) of commerce are leaf-galls, or cases, which ultimately contain a colony of insects.

In some parts of Europe a gall is common on one or more species of oak, in which the insect, having attacked the *cupula*, or acorn-cup, it

becomes developed into a curiously-shaped gall, known commercially as the knoppern (*Fig. 213*).

Some galls are formed in the substance of leaves, and ultimately burst through, and appear on the surface like flat disks, with perforations in the centre. *Oak spangles* are of this kind.

At one of the meetings of the Linnæan Society, a curious gall was shown by Dr. Farre, found on the leaves of a species of oak from Mexico. It consisted of an aggregation of hollow cylindrical tubes, nearly an inch in length, furnished with a fringed orifice. Their colour was white, suffused with red, especially towards the apex.

Galls are of two kinds, called respectively *galls* and *cases*. The *galls* are of a more or less solid or ligneous character,



Fig. 214.

and form the temporary abode of a single insect. The *cases* are hollow, and of a horny texture, and contain a colony of insects. Of the former kind are the "galls" of commerce (*Fig. 214*); the British galls, first discovered in Devonshire, but now becoming so



Fig. 215.

universally diffused through England; the apple galls, or Sodom apples; and the galls found on many species of oak other than those enumerated. The *cases* are represented by the commercial galls of China and Japan, the Kakrasingheo



Fig. 213.

galls of the Himalayas (*Fig. 215*), the galls of the Pistachio, Terebinth, and other species of *Rhus*, *Pistacia*, and *Terminalia*.

The majority of the galls enumerated contain a large percentage of tannin, and are, therefore, much employed in the arts; but there are others which do not contain the astringent principle. Such, for instance, are the galls found on the leaves of the elm in Southern Europe, which are sometimes the size of the fist, and contain a sweet and viscid fluid; also the *Baisonge*, found on the leaves of a species of sage in the East, which are collected and sold for food: they are sweet, and, with sugar or honey, are affirmed to make an excellent sweetmeat.

Warts, or *verrucae* (Lat. warts), are enlarged and thickened cells, containing starch and other substances, which are developed in a globular form upon the stems of certain plants.

Monstrous developments of individual organs often assume the appearance of excrescences, and, indeed, those which we have enumerated seem to be but the result of concentrated efforts, on the part of the plant, to repair the injury it has sustained, or manifestations of disease, faintly resembling the tumorous and cancerous diseases to which the animal structure is subject.

SECTION XXXVII.

VITAL ACTION.

An answer to the inquiry, "How does a plant grow?" may be fairly attempted upon the basis of the foregoing pages. The organs of plants and their functions having been described as fully as the limits of the present work would allow, it only remains to show how the structure of the plant is built up by their action.

From the soil in which the plant grows, it absorbs all its inorganic constituents dissolved in the fluid which enters the roots under the influence of endosmose. This fluid passes upwards from the roots to the leaves through the medium of the woody layers, especially the newest and most external, there to be elaborated under the influence of light.

The ascending fluid having reached the leaves, exhales much of its water in the form of vapour, through their stomata, absorbs additional carbonic acid, and, being elaborated and thickened, returns downwards by the bark and transversely by the medullary rays, depositing in its course its various contributions to the growing structure of the plant.

It must be remembered that very important changes take place in the sap during its passage through the leaves. It enters them in a crude state, and issues from them in a favourable condition for supplying all the requirements of the plant. Not only has water, but also oxygen, in considerable quantities, been exhaled, and carbonic acid has been inhaled; and thus, by the union of carbon with the hydrogen and oxygen of water in the rising sap, the descending fluid returns charged with carbonaceous secretions.

From this survey, it would seem evident that the following conditions are necessary to plant life:—

I. A soil containing all the inorganic substances which enter into the structure of the plant to be grown therein. Because the plant is unable to obtain them from any other source.

II. Sufficiency of water at the root to supply the place of that continually evaporating from the leaves, as well as a medium for the introduction of inorganic matters. Because water can only enter by the spongioles in sufficient quantity to supply that deficiency, and inorganic substances can only enter in solution.

III. Heat sufficient to evaporate the water of the ascending sap, when it becomes diffused through the leaves. Because, otherwise, the fluids of the plant would not exceed in density the water of the soil, and no endosmotic current would be established.

IV. Leaf-surface equivalent to the size of the tree. Because, otherwise, there is insufficient provision for bringing the crude sap under the influence of heat, light, and air.

V. Plenty of light to elaborate the secretions. Because, in the absence of light, plants etiolate and die ; absorption of carbon increasing relatively with intensity of light.

VI. An atmosphere containing carbonic acid gas. Because the carbon is requisite for the formation of the carbonaceous secretions, and plants exhibit a more vigorous habit in an atmosphere more than ordinarily charged with carbonic acid.

Hence, during the winter, when some of these conditions are not fulfilled, there is a cessation of growth.

The autumnal tints of leaves are believed to be due to the oxidation of their green colouring matter, or chlorophyll.

The fall of leaves is assumed to result from the choking up of their tissues by continued deposits during evaporation. Being unable to perform their functions, they become sickly, dry up, die, and, weakened at their point of attachment or articulation, fall off by their own weight, or, contracting by loss of moisture, disconnect themselves and fall.

In plant life, as in other manifestations of the Great Author of Nature, man meets with many phenomena which he is compelled to admire, but is puzzled to explain.

SECTION XXXVIII.

BOTANICAL DESCRIPTION.

It is a good test of the progress of the student in Botany, that he shall be able to describe a plant correctly, naming the parts with precision, and describing its features with accuracy. It is also a test of careful and correct observation, and should be the aim of every student to accomplish it well.

To dissect and examine the flower and seed, a pocket lens will be necessary.

Nothing must be taken for granted ; but every part should be carefully examined, noticing the result.

The botanical description should be as terse and compact as possible ; omitting all superfluous expressions, at the same time guarding against the use of either a wrong or a vague term.

Should the plant to be described be in flower only, it will be necessary either to examine the plant again when in fruit, to complete the description, or, if this cannot be done, let the description end with the flower. The parts should be described in the order of development.

ROOT.—Its form, whether conical, fibrous, &c., and also whether annual or perennial.

Care must be taken not to call tubers, rhizomes, or other underground stems, by the name of *roots*.

STEM.—Whether erect or prostrate, solid or fistular, cylindrical or angular, naked or pubescent, &c.

LEAVES.—Whether alternate or opposite, stipulate or exstipulate, radical or cauline, simple or compound, entire or serrated ; and character of venation.

INFLORESCENCE.—Whether racemose or cymose, ter-

minial or axillary, erect or drooping, and the nature of the bracteal appendages.

FLOWERS.—Commencing with the *calyx*, whether regular or irregular, monosepalous or polysepalous, persistent or deciduous, superior or inferior. *Corolla*, whether regular or irregular, monopetalous or poly-petalous, sessile or unguiculate. *Stamens*, whether definite or indefinite, hypogynous or epigynous, monadelphous or polyadelphous. *Filaments*, whether short or long, filiform or petaloid. *Anthers*, whether one or more celled, introrse or extrorse, innate or adnate, and whether opening by pores or valves. *Ovary*, superior or inferior, one or many celled, one or many seeded, syncarpous or apocarpous. *Style*, terminal or lateral, filiform or petaloid, persistent or deciduous. *Stigma*, whether simple or bifid, terminal or lateral. *Ovules*, whether erect or otherwise, anatropal or orthotropal, definite or indefinite. *Placenta*, whether axile or parietal.

FRUIT.—Indehiscent or dehiscent. If dehiscent, whether by pores or valves, and if loculicidal or septicidal, also the arrangement of the placentæ. *Seeds*, whether spherical or angular, smooth or hairy, albuminous or exalbuminous, whether the *albumen* is horny, oily, or mealy. The *embryo*, monocotyledonous or dicotyledonous, straight, annular, or spiral; whether the *radicle* is directed to the apex or base, and whether the *cotyledons* are flat, spiral, accumbent, or incumbent.

Plants should be described in conformity with the following scheme, commencing with the root and ending with the seed:—

19

20

21

SCHEME.

Root	Character. (<i>Concell. fibrous &c, Ann</i>)
Stem	General character. (<i>Erect, prostrate; Sol. and glan; naked</i>)
Leaves.....	Arrangement. (<i>alternate opposite</i>)
	Stipulation. (<i>stipulate, instipulate</i>)
	Composition. (<i>Simple, Compound.</i>)
	Form.
	Margin and incision.
	Venation.
Inflorescence	General character.
	Bracteal appendages.
Flower.....	Peculiarities of Calyx.
	Corolla.
	Stamens. (<i>filaments, anthers</i>)
	Disk.
	Pistil (Ovary, Style, Stigma, Placentation).
Fruit.....	General character.
	Dehiscence.
	Placentation.
	Arrangement of embryo.
Habitat, or place	of growth.

SECTION XXXIX.

DESCRIPTIVE MODELS.

As it will be advisable that this lesson should be devoted to the practical description of plants, in conformity with the *Scheme* already given, a botanical description, by Dr. Lindley, is appended as a model for imitation.

CARDAMINE PRATENSIS.

Cuckoo flower.

N.O. Brassicacæ.

Roots fibrous, slightly branched; proceeding from the sides of a half-subterranean, perennial, green *tuber*, whose sides are marked by wide scars, and very short tooth-like branches

STEM erect, annual, smooth, terete, about a foot high.

LEAVES: **RADICAL**, on long stalks, thin, dark green, distantly pinnate; *leaflets* stalked, in two to four pairs, sometimes alternate, wavy, orbicular, entire, angular, or toothed; the terminal one much larger than the other; **CAULINE**, more closely pinnate, in from five to seven pairs, with an odd one; *leaflets* linear, obtuse, entire.

RACEMES terminal and axillary, two or three, nodding, longer than the leaves, somewhat corymbose, ebracteate.

SEPALS four. Oblong, obtuse, membranous at the edge: the two opposite slightly saccate.

PETALS four. Cruciate, pale lilac or white, veiny, with a slight tooth on one side of the unguis.

STAMENS tetradynamous, hypogynous, erect, longer than the calyx; *filaments* subulate, rigid, somewhat herbaceous, the two lateral much shorter than the others; *anthers* ovate, innate, erect, opening longitudinally: those of the longer filaments extrorse, of the shorter introrse.

OVARY superior, terete, as long as the longest filaments, two-celled; *placentæ*, two in each cell, next the dissepiment, parietal, polyspermous; *style* very short; *stigma* capitate.

DISK: four minute green glands; one surrounding the base of each shorter stamen, one free, between the bases of each longer stamens.

FRUIT (rarely produced). **SILIQUEA** linear, compressed, slightly torulose; *valves* thin, flat.

SEEDS oblong, lenticular, bright brown, arranged in a single line, suspended from very short funicles.

EMBRYO exalbuminous, with a superior radicle as long as the accumbent cotyledons.

SECTION XL.

SUMMARY.

The following Summary has been prepared in a tabular form, and, had it been found practicable to have printed it entire upon an open sheet, would have presented, at one view, the whole scope of the foregoing pages. Having carefully studied all that has preceded it, a slight effort will enable the student to comprehend this table, which will then serve as a mentor, pointing out centres around which memory will gather the results of our forty lessons.

Ultimate Elements.

}	Oxygen.
	Hydrogen.
	Nitrogen.
	Carbon.
	Phosphorus.
	Sulphur.
	Chlorine.
	Iodine.
	Bromine.
	Silicon.
	Potassium.
	Sodium.
	Calcium.
	Magnesium.
	Aluminium.
Iron.	
Manganese.	
Copper.	

Proximate Principles.

	}	Cellulose.		
		Lignine.		
		Starch	{ Inulins.	
			{ Lichenine.	
		Doxtrine.		
		Sugar	{ Cane Sugar.	
			{ Grape Sugar.	
			{ Mannite.	
<i>Carbonaceous or (Nonazotic)</i>			Gum.	
			Resin.	
		Organic Acids.		
		Volatile Oils.		
		Camphor.		
		Caoutchouc.		
		Fixed Oils.		
		Wax.		
	}	Fibrine.		
		Caseine.		
<i>Nitrogenous</i>			Albumen.	
			Chlorophyll—Chromale.	
			Emulsine.	
		Diastase.		
		Alkaloids.		
	}	Water.		
<i>Mineral</i>			Mineral constituents of the ash of plants.	

Elementary Tissues.

	}	Vesicles or Cells—Lacuna	
CELLULAR ..			Parenchyma.
			Fibro-cellular.
	}	Pitted.	
			Woody tissue.
	}	Glandular.	
			Fibro-vascular
FASCULAR ..			{ Spiral vessels.
		{ Ducts.	
		{ Annular vessels.	
		{ Scalariform.	
		Laticiferous.	

Nutritive Organs.

<p>EPIDERMAL.</p>		<p>Cuticle.</p>	<p>{ Stomata. Hairs. Prickles. Stings.</p>	<p>Epidermis</p>	<p>Epiderma. Epithelium.</p>	
<p>AXIAL.</p>		<p>Descending ..</p>	<p>{ Simple..... Compound .. Adventitious</p>	<p>{ Fibrils..... Spongioles .. Exogenous .. Endogenous .. Acrogenous.. Rhizome Flagellum .. Soboles Tuber</p>	<p>{ Pith or Medulla Sheath..... Medullary Rays Duramen..... Alburnum Liber</p>	<p>{ Medullary System. Ligneous System. Cortical System.</p>
<p>FOLLICULAR.</p>		<p>Tillus.</p>	<p>{ Petiole..... Lamina</p>	<p>{ Buds. Bulbs.</p>	<p>{ Ascidia. Phylodia.</p>	
<p>Stipule</p>		<p>Reticulum.</p>				

Reproductive Organs.

I.—*Phanerogamous.*

	Bracts.	Involucre.	Epicalyx.
FLORAL	{ Envelopes	{ Calyx (Sepals)	{ Perianth or Perigone.
	{ Essentials	{ Corolla (Petals).....	
	{ Columna	Stamens	{ Anther { Pollen- grains } Fovilla,
		Pistil.....	{ Filament, Stigma, Style.
	{ Pericarp	Ovary	{ Carpels, Ovules.
		{ Epicarp { Sarcocarp.	
POSTFLORAL	{ Seed.....	Mesocarp ..	{ Cotyledons, Plumule—Collar, Radicle.
		Endocarp, Spermoderm, Albumen. Embryo	

II.—*Cryptogamous.*

FILICAL	{ Thecæ Indusium. Annulus. }	{ Antheridia Archegonia }	{ Peristome. Operculum, Spores. Columella, Calyptra, Vaginule. }
		{ Antheridium Paraphyses }	
MUSCAL	{ Perigone Epigone }	{ Antheridium Archegonia }	{ Sporangium .. Seta Theca Paraphyses. }
		{ Thalamium Apothecium, Exciple. }	
LICHENAL	{ Pycnidia Spermagonia Hymenium }	{ Stylospores Spermatozoids Spores. }	{ Stylospores Spermatozoids Spores. }
		{ Basidia Paraphyses, Spermagonia Pycnidia Stylospores. }	
FUNGAL	{ Tetraspores..... }	{ Conceptacles .. Stichidia..... Ceramidium Antheridia Sporangia }	{ Perisporium Sporules. Spores. Spermatozoids Zoospores. }
		{ Perisporium Sporules. Spores. Spermatozoids Zoospores. }	

The student having arrived thus far, and being desirous of prosecuting his studies still farther, may consult *Henfrey's Elementary Course of Botany* (Van Voorst), *Bentley's Manual of Botany* (Churchill), *Balfour's Manual of Botany* (Griffin & Co.), or *Lindley's Elements of Botany* (Bradbury & Evans) : all are nearly the same in price and equally valuable.

N.B.—More minute and special information on *Ferns* will be found in Mrs. Lankester's little volume; and on *Fungi* in the author's two volumes on that subject.

INDEX.

ABORTIVE branches	<i>Page</i> 43	Botany	<i>Page</i> 2
Abortive stamens	66	Bracts	56
Achenium	81	Bractlets	56
Achlamydeous	62	Bromine	7
Acids, organic	11	Bulbs	40, 42
Aconitine	14	Bulbils	42
Acrogens	37	CAFFEINE	14
Adnate	69	Calceolate	65
Adventitious buds.. .. .	43	Calcium	7
Adventitious roots	32	Calyptra	96
Æstivation	57	Calyx.. .. .	61
Albumen	13, 85	Cambium	34
Albumum	34	Campanulate	64
Alga, reproduction of	89	Camphor	11
Alkaloids	14	Cane Sugar	10
Alternate leaves	52	Caoutchouc	12
Amentum	58	Capitulum	60
Ampulla	54	Capsule	83
Angiocarpous	91	Carbon	5
Angiospermous	84	Carbonaceous compounds	8
Animals and Plants	2	Cardamine pratensis	111
Annulated root	31	Carpellary leaves	72
*Annulus	94, 96	Carpels	72
Anther	68	Carpophore	73
Antheridia	90, 95, 97, 99	Caryophyllaceous	66
Anthocarpous	80	Caryopsis	81
Anthotaxis	58	Casine	13
Apocarpous.. .. .	72, 80	Catkin	58
Apophysis	97	Cell-development	23
Apothecia	91	Cells	18
Arabine	10	Cellulares	19
Archegonia	99	Cellulose	8, 18
Ariilus	85	Centrifugal inflorescence	60
Ascending axis	32	Centripetal ditto	60
Asci	95	Ceramidium	89
Ascidium	54	Cerasing	10
Assimilation:	101	Chalaza	76
Atropine	14	Chlorine	6
Axis	29	Chlorophyll.. .. .	14
BACCA	8.	Chlorenchyma	23
Bark	35	Circulation	100
Basidia	95	Circumscription	51
Bassorine	10	Citric acid	11
Berry	71	Coated roots	103
		Coccidium	80

Collum	Page 29	Epigynous	Page 60
Columella	79, 96	Epipetalous	68
Compound leaves	49	Epiphloeum	35
Concentric rings	34	Epiphragm	96
Conceptacles	90	Epiphytes	103
Conditions of germination	87	Episperm	84
Cone	59	Epithelium	26
Conical root	30	Essential organs	67
Connective	69	Etærio	81
Corns	40	Exciple	91
Corolla	63	Exogens	32
Corymb	59	Exorhizal	88
Cotyledons	85	Exosome	31
Cruciform	66	Extrorse	70
Cuckoo-flower	111	FALSE tracheæ	22
Cupule	56	Fasciculated root	31
Cyme	59	Favellæ	89
Cystidia	95	Feather-veined leaves	46
Cytoblast	23	Ferns, reproduction of	98
DECUSSATE	52	Fibrine	12
Dehiscence	79	Fibro-cellular tissue	20
Descending axis	29	Fibrous root	30
Description, botanical	109	Fibro-vascular tissue	21, 22
Descriptive models	111	Filament	68
Dextrine	9	Fissuration	24
Diadelphous	69	Fixed embryos	41
Diastase	13	Fixed oils	12
Dichlamydeous	61	Flagellum	39
Diclinous	67	Floral envelopes	57
Dicotyledons	85	Floral leaves	56
Didynamous	69	Flower-buds	55
Diceious	67	Follicle	82
Disk	66	Foramen	76
Dispersion of pollen	71	Fossil botany	3
Dissepiments	73	Fovilla	70
Distichous	52	Frons	38
Drupe	81	Fruit	78
Ducts	22	Fungi	94
Duramen	34	Funiculus	76, 85
ELATERS	96	Furcate-veined leaves	46
Elements of plants	4, 17	Fusiform root	30
Embryo	85	GALLIC acid	11
Embryo-buds	43	Galls and excrescences	104
Embryo-sac	99	Gemination	24
Emulsine	13	Germination	87
Endocarp	79	Gills	94
Endochrome	90	Glands	66
Endogens	36	Glandular hairs	28
Endophlœum	35	Glandular woody tissue	22
Endopleura	84	Glans	81
Endorhizal	88	Glomerulus	60
Endosome	31	Glucose	10
Epiblema	26	Gonidia	93
Epicarp	79	Grape sugar	10
Epidermis	25, 44	Gum	10

INDEX.

121

Gymnospermous	Page 84	Malic acid	Page 11
Gymnocarpous	91	Manganese	8
Gynandrous.. .. .	68	Mannite	10
Gynophore	73	Medulla	33
		Medullary rays	35
H		Medullary sheath	34
Hairs	27	Mesocarp	79
Heat	101	Mesophloeum	35
Hermaphrodite	67	Micropyle	76
Hesperidium	82	Midrib	45
Heterorhizal	88	Mineral compounds	14
Hilum	78, 85	Model for description	111
Histology	2	Monadelphous	69
Hydrocyanic acid	11	Moniliform root	30
Hydrogen	5	Monochlamydeous	63
	60	Monocotyledons	85
Hypogynous	97	Morphine	14
Hypothecium	92	Morphology	2
		Mosses, reproduction of	96
I		Mucilage	10
INDUSIUM	98	Mycelium	94
Inferior ovary	74		
Inflorescence	58	NAPIFORM ROOT	30
Infundibuliform	64	Nectarines	67
Internodes	33	Nemathecchia	90
Introrse	70	Nitrogen	5
Inuline	10	Nitrogenous compounds.. .. .	12
Invoucre	56	Nodes	33
Iodine	6	Nodulose root	31
Iron	7	Nucleus and nucleoli	24
Irritability	102	Nucleus of ovule	76
LABIATE	65	OCUREA	34
Lacunæ	19	Odours	102
Latex	23	Oils, fixed	12
Laticiferous vessels	23	Oils, volatile	11
Leaf-appendages	53	Opposite leaves	52
Leaf-buds	41	Organic acids	10
Leaf-form	47	Ovary	72
Leaf-margins	51	Ovules	75
Leaf-order	51	Oxalic acid	11
Leaf-structure	43	Oxygen	4
Legume	82		
Legumine	13	PANICLE	59
Liber	35	Papilionaceous	66
Lichenine	9	Papillæ	99
Lichens, reproduction of.. .. .	91	Pappus	62
Lignine	8	Parallel-veined leaves	46
Ligulate	65	Paraphyses	95, 97
Lime	7	Parasites	103
Loculicidal	80	Parenchyma	19, 44
Locustæ	59	Parietal	73
Luminosity	102	Pectine	10
		Pedicels	88
MAGNESIUM	7		

Peduncle	Page 58	SCALARIFORM vessels	Page 29
Pepo	82	Scheme, descriptive	111
Perianth	63	Sclerogen	25
Pericarp	79	Seed	87
Perigynous	68	Seed-buds	75
Peristome	96	Sepals	61
Perithecium	92	Septicidal	80
Personate	65	Septifragal	80
Petals	63	Septum	73
Petiole	43, 53	Seta	96
Phenomena, vegetative	100	Setæ	28
Phosphorus	5	Silicon	6
Phylloidium	54	Siliqua	83
Phytozoa	97	Simple leaves	47
Pileus	94	Soboles	39
Piperine	14	Sodium and Soda	7
Pistil	72	Solanine	14
Pitcher	54	Soredia	93
Pith	33	Sorosis	83
Pitted tissue	20	Spadix	56, 58
Placenta	73	Spathc	56
Pleurenchyma	21	Spawn	94
Plumule	85	Spermagones	91, 93
Podosperm	76	Spermatia	93, 95
Pollen	68, 70	Spermatozoids	90, 95
Polydelphous	69	Spermoderm	84
Polyspermous	74	Spicules	95
Pome	82	Spike	58
Potassium and potash	7	Spines	43
Prickles	28	Spiral vessels	22
Primordial utricle	25	Spongioles	29
Prothallus	99	Sporangium	96, 98
Protoplasm	25	Spores	97
Pseudo-bulbs	40	Sporophores	95
Pulvinus	53	Stamens	67
Pycnides	91	Staminiidia	97
Pyxidium	83	Starch	9
QUININE	11	Stem	32
RACEME	59	Sterigmata	93
Rachis	58	Stichidia	90
Radicle	85	Stigma	72, 74
Raphe	76	Stipe	94
Raphides	18	Stipels	55
Receptacle	75, 98	Stipes	38
Replum	74	Stipules	54
Resins	10	Stomata	26
Reticulum	55	Strobilus	58, 33
Rhizome	39	Struma	97
Ringent	65	Strychnine	14
ROOT	29	Style	72
Rosaceous	66	Stylospores	91
Rotae	65	Subterranean stems	38
Rotation	100	Sugar	10
Runner	30	Sulphur	6
		Summary	113
		Superior ovary	74
		Suture	70

INDEX.

123

Syconus	Page 83	Umbel	Page 60
Syncarpous	72, 80	Unguis	63
Syngenesious	70	Urceolate	65
TANNIC acid	11	Utricle	81
Tartaric acid	11	VAGINA	55
Tendrils	54	Valves	70
Testa	84	Vascular tissue	21, 22
Tetradynamous	69	Vegetative phenomena	100
Tetraspores	89	Venation	45
Thalamium	91	Veratrine	14
Thalamus	75	Vernation	41
Thallogens	33	Verrucæ	106
Thallus	33	Versatile anthers	70
Thecæ	22, 92, 95, 99	Vertical	52
Theine	14	Vital action	106
Thorns	43	Volatile oils	11
Thyrus	50	WARTS	106
Torus	75	Water	15
Trachenchyma	22	Wax	12
Tuber	40	Whorl	52
Tubercular root	30	Woody layers	34
Tubular	64	Woody tissue	21
ULTIMATE ELEMENT	3		

SCHLEIDEN'S
PRINCIPLES OF SCIENTIFIC BOTANY;
OR, BOTANY AS AN INDUCTIVE SCIENCE.

Translated by DR. LANKESTER.

Hundreds of Woodcuts, and Six Pages of Figures, beautifully
engraved on Steel. Demy 8vo., price 10s. 6d.

**Every Botanical Library should possess this work, as it contains the
principles upon which all structural botany is based.**

London: ROBERT HARDWICKE. 192, Piccadilly.

