

**PAGES MISSING
WITHIN THE
BOOK ONLY**

**TEXT PROBLEM
WITHIN THE
BOOK ONLY**

UNIVERSAL
LIBRARY

OU_156727

UNIVERSAL
LIBRARY

OSMANIA UNIVERSITY LIBRARY

Call No. *SM/S61E* Accession No. *962*

Author *Snowball.*

Title *Elements... trigonometry*

This book should be returned on or before the date
last marked below.

CAMBRIDGE
ELEMENTARY SCHOOL CLASS BOOKS.

The Volumes of this Series of ELEMENTARY SCHOOL CLASS BOOKS, are handsomely printed in a form that, it is hoped, will assist the young student as much as clearness of type and distinctness of arrangement can effect. They are published at a low price to ensure an extensive sale in the Schools of the United Kingdom and the Colonies.

1. *An Elementary Latin Grammar.*

By H. J. ROBY, M.A., Under Master of Dulwich College Upper School, late Fellow and Classical Lecturer of St John's College, Cambridge. 2s. 6d.

2. *Euclid for Colleges and Schools.*

By I. TODD HUNTER, M.A., F.R.S., Fellow and Principal Mathematical Lecturer of St John's College, Cambridge. 3s. 6d.

3. *An Elementary History of the Book of Common Prayer.*

By FRANCIS PROCTER, M.A., Vicar of Witton, Norfolk, late Fellow of St Catharine's College, Cambridge. 2s. 6d.

4. *Algebra for Beginners.*

By I. TODD HUNTER, M.A., F.R.S. Nearly ready.

5. *The School Class Book of Arithmetic.*

By BARNARD SMITH, M.A., late Fellow of St Peter's College, Cambridge. In the Press.

6. *The Bible Word-Book.*

A Glossary of Old English Bible Words with Illustrations.

By J. EASTWOOD, M.A., St John's College, Cambridge, and Incumbent of Hope in Hanley, Stafford, and W. ALDIS WRIGHT, M.A., Librarian of Trinity College, Cambridge.

Preparing.

•• *Other Volumes will be announced in due course.*

By the same Author.

FOURTH EDITION. Crown 8vo. Cloth, 5s.

THE CAMBRIDGE COURSE OF
ELEMENTARY NATURAL PHILOSOPHY.

BEING THE PROPOSITIONS IN MECHANICS AND
HYDROSTATICS

In which those Persons who are not Candidates for Honours are examined
for the DEGREE OF B.A.

ADAPTED FOR THE USE OF COLLEGES AND SCHOOLS.

To which are now added Examples and Problems from the University Exami-
nation Papers and Original.

WITH HINTS FOR THEIR SOLUTION.

MACMILLAN AND Co. CAMBRIDGE AND LONDON.

PREFACE TO THE EIGHTH EDITION.

IN preparing the present edition for the press, the text has been subjected to a careful revision; the proofs of some of the more important propositions have been rendered more strict and general; and a considerable addition, of more than two hundred examples, taken principally from the questions set of late years in the public examinations of the University and of individual Colleges, has been made to the collection of Examples and Problems for practice.

The original design of the work has still been kept in view,—namely, to present to the student who is reading for University Honours, in a plain, clear, and intelligible form, the main Principles of Trigonometry; without, on the one hand, giving too much space to the niceties of the practical application of Theory; or, on the other, introducing investigations that require considerable proficiency in Algebraical Analysis.

The Book now appears in a more commodious form, and its price has been reduced.

INDEX.

PLANE TRIGONOMETRY.

CHAPTER I.

	ARTICLES.
ON THE METHODS OF MEASURING LINES AND ANGLES.	1—12

CHAPTER II.

DEFINITIONS OF THE GONIOMETRICAL RATIOS, AND FORMULÆ CONNECTING THEM WITH EACH OTHER	13—33
Definitions	14
Changes of the Sine, &c. in magnitude and sign	20, 21
Formulæ involving one angle only	22—29
Sines, &c. of 45° , 30° , 60°	32
EXAMPLES	33

CHAPTER III.

FORMULÆ INVOLVING MORE ANGLES THAN ONE	34—67
Sines and Cosines of $A \pm B$	34—36
$\text{Sin } 2A$ and $\text{Cos } 2A$ in terms of $\text{Sin } A$ and $\text{Cos } A$, and conversely	36—41
Formulæ deduced from the Sines and Cosines of $A \pm B$	43—56
Sines and Cosines of 18° , 72° , 36° , 54°	58
Increments of the Sine, &c. of an angle	59—63
Inverse Goniometric Ratios	64, 65
EXAMPLES	67

CHAPTER IV.

THE SOLUTION OF TRIANGLES AND OF PLANE RECTILINEAL FIGURES	69—96
SUBSIDIARY ANGLES	97, 98

CHAPTER V.

ANALYTICAL TRIGONOMETRY	99—125
The Circumference of a Circle varies as the Radius ..	99
CIRCULAR MEASURE of an angle	100—103
$\theta > \sin \theta$, and $< \tan \theta$	104
$\text{Sin } \theta > \theta - \frac{1}{4} \theta^3$	105

	ARTICLES.
Number of <i>seconds</i> in $\theta = \frac{\theta}{\sin 1''}$	107
Area of Circle of Radius $r = \pi r^2$	108
DEMOIVRE'S THEOREM	109, 110
Sine and Cosine in terms of the Sines and Cosines of the multiples of the angle	111, 112
Tan $n\theta$ in terms of Tan θ	113
Sin α and Cos α in terms of α	114, 115
Series for finding the value of π	116—118
MISCELLANEOUS PROPOSITIONS.....	119—125

CHAPTER VI.

ON THE SOLUTION OF EQUATIONS BY MEANS OF THE TRIGONOMETRIC TABLES, AND THE RESOLUTION OF $x^n \pm 1$, Sin θ , AND Cos θ INTO FACTORS

126—133

APPENDIX I.

ON THE LOGARITHMS OF NUMBERS, AND THE CONSTRUCTION AND USE OF TABLES OF LOGARITHMS OF NUMBERS.

APPENDIX II.

ON THE CONSTRUCTION OF THE TABLES OF NATURAL GONIOMETRICAL RATIOS.

APPENDIX III.

ON THE CONSTRUCTION OF THE LOGARITHMIC TABLES OF THE GONIOMETRICAL RATIOS.

APPENDIX IV.

THE GENERAL PROOF OF THE FORMULE for SIN ($A \pm B$) and Cos ($A \pm B$).

A COLLECTION OF EXAMPLES FOR PRACTICE.

The Student is recommended to confine his attention at first to the Articles,

1—27; 30—32; 34; 38—43; 46; 50; 51; 58; 64.

Appendix I. 1—13; II. 1—8; III. 1, 4, 5, 12.

Arts. 69—76; 78; 81—83; 86; 87; 90; 99—109 omitting Cor.;
110; 113—115; 118.

SPHERICAL TRIGONOMETRY.

CHAPTER I.

	ARTICLES.
EXPLANATION of the objects of SPHERICAL TRIGONOMETRY	1—5
Certain preliminary general properties of Spherical Triangles established by means of Solid Geometry	6—26
Recapitulation of the results established in this Chapter.	27

CHAPTER II.

GENERAL FORMULÆ CONNECTING THE SIDES AND ANGLES OF TRIANGLES WITH ONE ANOTHER	28—35
Values of the Angles in terms of the Sides of a Triangle, and <i>vice versa</i> , adapted to Logarithmic Computation	31. 32
NAPIER'S ANALOGIES	33
GAUSS' THEOREM	35

CHAPTER III.

THE SOLUTION OF RIGHT-ANGLED, AND QUADRANTAL, TRIANGLES	36—43
---	-------

CHAPTER IV.

ON THE SOLUTION OF OBLIQUE-ANGLED TRIANGLES	44—55
The radii of the small circles found which can be described within and about a given Triangle	56—59

CHAPTER V.

ON THE AREAS OF SPHERICAL TRIANGLES; AND THE SOLUTION OF TRIANGLES WHOSE SIDES ARE SMALL COMPARED WITH THE RADIUS OF THE SPHERE	60—67
CAGNOLI'S and LHULLIER'S THEOREMS for finding the Spherical Excess	63, 64
Solution of Triangles whose sides are small compared with the radius of the Sphere	66—76
LEGENDRE'S THEOREM	73
THE CHORDAL TRIANGLE.....	74, 75

CHAPTER VI.

	ARTICLES.
ON GEODETIC MEASUREMENTS; THE INSTRUMENTS EMPLOYED, AND THE MANNER OF USING THEM	77—86

CHAPTER VII.

ON THE SMALL CORRESPONDING VARIATIONS OF THE PARTS OF A SPHERICAL TRIANGLE; AND THE CONNEXION EXIST- ING BETWEEN CERTAIN FORMULÆ OF SPHERICAL TRIGO- NOMETRY AND ANALOGOUS FORMULÆ OF PLANE TRIGO- NOMETRY	87—96
--	-------

CHAPTER VIII.

THE REGULAR POLYHEDRONS: certain properties of them inves- tigated	92—103
---	--------


EXAMPLES FOR PRACTICE.

The Student is recommended to confine his attention at first to
these Articles; 1—41; 44—55; 60—62.

the direction of $Y'AY$ is *positive*, because it lies on the upper side of $X'AX$.

Similarly, the ordinates of P_3 and P_4 , measured in the direction of $X'AX$ are *negative* and *positive* respectively; and the ordinates measured in the direction of $Y'AY$ are *negative* in both cases.

4. DEF. If a straight line revolve in one plane round its extremity A from a given position, as AB , into any other position, as AC , the inclination of AC to AB is called an *angle* (\angle); and the angle is signified by the letters BAC or CAB , the middle letter being that placed at the point in which the two lines meet.



By continuing this revolving motion, the angle may be supposed to become of any magnitude whatever.

5. DEF. If AD be equally inclined to the parts AB, AB' of the straight line BAB' , each of the angles $BAD, B'AD$ is called a *right angle*.

6. DEF. An *acute angle* is less, and an *obtuse angle* is greater, than a *right angle*.

7. If the angles formed by AC revolving in one direction, (as BCD) from the fixed line AB , be considered *positive*, then if AC revolve in the contrary direction from AB , it will trace out *negative angles*.

If to the angle BAC , Fig. Art. 4, it be required to add a given angle, CA must move in the direction BCD through an angle CAE equal to the given angle, and the whole BAE will be the angle required. And if it be required to take a given angle from BAC , CA must evidently move in a contrary direction till it come into a position $E'A$, such that $\angle CAE'$ is equal to the angle to be subtracted.

$$\text{Then } \angle CAE' + \angle E'AB = \angle BAC;$$


$$\therefore \angle E'AB = \angle BAC - \angle CAE'.$$

Now if $\angle CAE'$ be greater than $\angle CAB$, $E'A$ lies on the other side of AB ,

$$\text{and } \angle E'AB = \angle BAC - \angle CAE'$$

$$= -(\angle CAE' - \angle BAC),$$

a negative quantity, whose magnitude is the difference between the angles CAE'



and BAC ; which difference lies in this case on the lower side of AB .

Hence, when an angle is called *negative*, it is meant that it is formed by the revolving line moving from the fixed line in a direction contrary to that in which it revolved to trace out positive angles.

8. A right angle is divided by the English into 90 equal angles which are called degrees; a degree is subdivided into 60 minutes, and a minute into 60 seconds. And the magnitude of an angle is expressed by stating how many degrees and subdivisions of a degree are contained in the angle. If great accuracy be required, the parts of an angle which are less than a second are expressed in decimal parts of a second.

A degree and its subdivisions are thus indicated, $24^{\circ}, 50', 34''.7$, which denotes an angle containing 24 degrees, 50 minutes, 34 seconds, and seven tenths of a second.

9. By the French and other Continental Mathematicians, a right angle is divided into 100 equal angles called grades, a grade into 100 minutes, and a minute into 100 seconds; and the divisions are thus marked, $26^{\circ}, 24', 32''.47$,

$$\text{Now since } 1^{\circ} = \frac{1^{\circ}}{100} = .01^{\circ}, \quad \text{and } 1'' = \frac{1''}{100} = \frac{1^{\circ}}{1000} = .0001^{\circ};$$

The above angle might have been written thus, $26^{\circ}.243247$.

Whence it appears that if the French division be adopted, arithmetical operations can be performed on angles in the same manner as on any other decimal fractions; an advantage which does not attend the English division.

10. To find the relation between E and F , the number of Degrees and of Grades contained in the same angle BAC . (Art. 14. Fig. 1.)

$$\text{In the English division, } \frac{E^{\circ}}{90^{\circ}} = \frac{\text{given } \angle BAC}{\text{right } \angle};$$

$$\text{In the French division, } \frac{F^{\circ}}{100^{\circ}} = \frac{\text{given } \angle BAC}{\text{right } \angle};$$

$$\therefore \frac{E}{90} = \frac{F}{100}; \text{ and } \begin{cases} E = \frac{9}{10} F = F - \frac{F}{10} \dots(1), \\ F = \frac{10}{9} E = E + \frac{E}{9} \dots(2). \end{cases}$$

NOTE. In employing the latter of these formulæ it will be necessary to express the minutes and seconds of the angle in decimal parts of a degree, since E represents the number of *degrees* in the angle.

Ex. 1. To find how many degrees, minutes and seconds are contained in the angle $42^{\circ}, 34', 56''$.

$$\begin{array}{r}
 F = 42.3456 \\
 \frac{F}{10} = 4.23456 \\
 \hline
 \therefore E = F - \frac{F}{10} = 38.11104 \\
 \phantom{\therefore E = F - \frac{F}{10} = 38.} 60 \\
 \phantom{\therefore E = F - \frac{F}{10} = 38.} 6.6624 \\
 \phantom{\therefore E = F - \frac{F}{10} = 38.} 60 \\
 \hline
 \phantom{\therefore E = F - \frac{F}{10} = 38.} 39.744
 \end{array}$$

Or $38^{\circ}, 6', 39''.74$, retaining the tenths and hundredths and neglecting the *thousandth* parts of a second.

Ex. 2. Find how many grades, minutes and seconds are contained in the angle $24^{\circ}, 51', 45''$.

First reducing the minutes and seconds to the decimal parts of a degree,

$$\begin{array}{r}
 60 \mid 45'' \\
 \hline
 60 \mid 51.75 \\
 \hline
 .8625;
 \end{array}
 \qquad
 \begin{array}{r}
 \therefore E = 24.8625 \\
 \frac{E}{9} = 2.7625 \\
 \hline
 \therefore E + \frac{E}{9} = 27.6250
 \end{array}$$

and $\therefore F = 27^{\circ}, 62', 50''$.

11. DEF. *The Complement of an angle is its defect from a right angle.*

Thus, $90^{\circ} - 24^{\circ}, 32' = 65^{\circ}, 28'$, is the complement of $24^{\circ}, 32'$.

$90^{\circ} - 110^{\circ}, 15' = -(20^{\circ}, 15')$ is the complement of $110^{\circ}, 15'$.

12. DEF. *The Supplement of an angle is its defect from two right angles.*

Thus, $180^{\circ} - 56^{\circ}, 20' = 123^{\circ}, 40'$, is the supplement of $56^{\circ}, 20'$.

$180^{\circ} - 186^{\circ}, 12' = -(6^{\circ}, 12')$, is the supplement of $186^{\circ}, 12'$.


A Collection of Examples and Problems is placed after the fourth Appendix.

CHAPTER II.

THE GONIOMETRICAL RATIOS, AND SOME FORMULÆ CONNECTING
THEM WITH EACH OTHER.

13. DEF. *Plane Trigonometry*, in its original meaning, implies the measuring of plane triangles; in its extended signification it treats of the formulæ connecting the relations of angles with each other, and of the determination of the parts of plane rectilineal figures from sufficient data.

14. Let a straight line revolve from the fixed line AB round the point A , in the direction of the letters B, D, B', D' , and come into the position AC .



From any point C in AC draw CN at right angles to AB ,—produced either way if necessary; and through A draw DAD' at right angles to AB .

Now, for the reasons given in Arts. 2 and 3, in these figures the signs of NC are +, +, −, − respectively, and those of AN are +, −, −, + respectively.

15. DEFINITIONS. See the figures of the last Article.

1. $\frac{NC}{AC}$ is the *Sine* of the $\angle BAC$; or, $\text{Sin } \angle BAC = \frac{NC}{AC}$;

2. $\frac{AN}{AC}$ is the *Cosine* of the $\angle BAC$; or, $\text{Cos } \angle BAC = \frac{AN}{AC}$.

3. $\frac{NC}{AN}$ is the *Tangent* of the $\angle BAC$; or, $\text{Tan } \angle BAC = \frac{NC}{AN}$.

4. $\frac{AC}{AN}$ is the *Secant* of the $\angle BAC$; or, $\text{Sec } \angle BAC = \frac{AC}{AN}$.

5. $1 - \cos \angle BAC$ is the *Versed-sine* of the $\angle BAC$;
or, $\text{Versin } \angle BAC = 1 - \cos \angle BAC$.

6. The *Tangent* of the Complement of the $\angle BAC$ is called the *Cotangent* of the $\angle BAC$;

$$\text{or, } \text{Cot } \angle BAC = \tan (90^\circ - \angle BAC).$$

COR. If $90^\circ - \angle BAC$ be the original angle, its Complement is $\angle BAC$, Art. 11,

$$\therefore \text{Cotan } (90^\circ - \angle BAC) = \tan \angle BAC,$$

$$\text{or, } \text{Tan } \angle BAC = \cotan (90^\circ - \angle BAC).$$

7. The *Secant* of the Complement of the $\angle BAC$ is called the *Cosecant* of the $\angle BAC$;

$$\text{or, } \text{Cosec } \angle BAC = \sec (90^\circ - \angle BAC).$$

COR. If $90^\circ - \angle BAC$ be the original angle, its Complement is $\angle BAC$, Art. 11,

$$\therefore \text{Cosec } (90^\circ - \angle BAC) = \sec \angle BAC;$$

$$\text{or, } \text{Sec } \angle BAC = \text{cosec } (90^\circ - \angle BAC).$$

16. The cosine of $\angle BAC$ might have been defined to be the Sine of the Complement of $\angle BAC$.

$$\text{For } \text{Cos } \angle BAC = \frac{AN}{AC} = \sin \angle ACN = \sin (90^\circ - \angle BAC). \quad \text{Art. 14: Fig. 1.}$$

$$\text{Also, } \text{Sin } \angle BAC = \frac{NC}{AC} = \cos \angle ACN = \cos (90^\circ - \angle BAC),$$

or the Sine of an angle is equal to the Cosine of its Complement.


For the sake of convenience an angle will generally hereafter be indicated by a single letter, as $\text{Sin } A$, $\text{Cos } A$, $\text{Tan } B$, where A , B respectively represent the number of degrees contained in the angle.

17. So long as the magnitude of the angle is unaltered, its Sine, Cosine, Tangent, &c. remain the same, whatever be the magnitude of AC .

For let D be any other point in AC , and DM perpendicular to AB .

Then, by definition, $\text{Sin } A = \frac{NC}{AC}$; or $\text{Sin } A = \frac{MD}{AD}$.

But by similar triangles $\frac{NC}{AC} = \frac{MD}{AD}$, or $\text{Sin } A$ is the same wherever in the line AC the point C be situated. Similarly it may be shewn that $\text{Cos } A$, $\text{Tan } A$, $\text{Sec } A$... are invariable quantities so long as the magnitude of A remains unaltered.



Hence, if any of the quantities $\text{Sin } A$, $\text{Cos } A$, $\text{Tan } A$, $\text{Sec } A$... be given, the angle A may be determined.

18. DEF. The Ratios which are called the Sine, Cosine, Tangent, &c., of any angle are termed "The Goniometrical Ratios," because when any one of them is given the angle may be determined to which it belongs. These Ratios are also called "Trigonometrical Functions" of angles.

19. To express $\text{Versin } A$, $\text{Cot } A$, $\text{Cosec } A$, in terms of the sides of the triangle ANC . (Fig. 1. Art. 20.)

$$(1) \quad \text{Versin } A = 1 - \cos A = 1 - \frac{AN}{AC}.$$

$$(2) \quad \begin{aligned} \text{Cot } A &= \tan(90^\circ - A) = \tan ACN \\ &= \frac{NA}{CN}, \text{ by def. of the tangent.} \end{aligned}$$

$$(3) \quad \begin{aligned} \text{Cosec } A &= \sec(90^\circ - A) = \sec ACN \\ &= \frac{CA}{CN}, \text{ by def. of the secant.} \end{aligned}$$

20. To trace the variation in the algebraic signs of $\text{Sin } A$, $\text{Cos } A$, $\text{Tan } A$, $\text{Sec } A$, as A increases from 0° to 360° .


(1) $\text{Sin } A = \frac{NC}{AC}$, and therefore has the same sign in any case as NC has; for AC which lies in the direction of neither of the

lines AB and AD , cannot change its sign, and is always to be reckoned as positive.

Hence (14) $\text{Sin } A$ is positive if A be an angle between 0° and 180° (figs. 1, 2); and is negative if A be between 180° and 360° . (Figs. 3, 4.)

$$(2) \quad \text{Cos } A = \frac{AN}{AC}, \text{ and therefore has the same sign as } AN.$$

Hence (14) $\text{Cos } A$ is positive if A be between 0° and 90° , or between 270° and 360° (figs. 1, 4); and is negative if A be between 90° and 270° . (Figs. 2, 3.)



$$(3) \quad \text{Tan } A = \frac{NC}{AN}, \text{ and is therefore positive or negative according as } NC \text{ and } AN \text{ have the same or different signs.}$$

Hence (14) $\text{Tan } A$ is positive if A be between 0° and 90° , or between 180° and 270° (figs. 1, 3); and it is negative if A be between 90° and 180° , or between 270° and 360° . (Figs. 2, 4.)

$$(4) \quad \text{Sec } A = \frac{AC}{AN}, \text{ and therefore has the same sign as } AN.$$

Hence $\text{Sec } A$ is positive if A be between 0° and 90° , or between 270° and 360° ; and it is negative if A be between 90° and 270° .

21. *To trace the variations in the magnitudes of the Sine, Cosine, Tangent, and Secant, as the angle increases from 0° to 360° . (Figs. Art. 20.)*

Since (17) the values of the Sine, Cosine, Tangent, and Secant are not affected by the magnitude of AC , suppose this line to remain of the same magnitude while the angle A increases from 0° to 360° .

Now (fig. 1) as AC revolves from the position AB into the position AD , NC increases in magnitude from 0 to AC , and is positive; and AN decreases from AC to 0, and is positive.

As (fig. 2) AC revolves from the position AD into the position AB' , NC decreases in magnitude from AC to 0, and is positive; and AN increases from 0 to AC , and is negative.

As (fig. 3) AC revolves from AB' to AD' , NC increases in magnitude from 0 to AC , and is negative; and AN decreases from AC to 0, and is negative.

As (fig. 4) AC revolves from AD' to AB , NC decreases in magnitude from AC to 0, and is negative; and AN increases from 0 to AC , and is positive.

Hence it appears, that as

A changes from	0° to 90°	90° to 180°	180° to 270°	270° to 360°
$\text{Sin } A \left(\frac{NC}{AC} \right) \dots$	$\frac{0}{AC}$ to $\frac{+AC}{AC}$	$\frac{+AC}{AC}$ to $\frac{0}{AC}$	$\frac{0}{AC}$ to $\frac{-AC}{AC}$	$\frac{-AC}{AC}$ to $\frac{0}{AC}$
$\text{Cos } A \left(\frac{AN}{AC} \right) \dots$	$\frac{+AC}{AC}$ to $\frac{0}{AC}$	$\frac{0}{AC}$ to $\frac{-AC}{AC}$	$\frac{-AC}{AC}$ to $\frac{0}{AC}$	$\frac{0}{AC}$ to $\frac{+AC}{AC}$
$\text{Tan } A \left(\frac{NC}{AN} \right) \dots$	$\frac{0}{+AC}$ to $\frac{+AC}{0}$	$\frac{+AC}{0}$ to $\frac{0}{-AC}$	$\frac{0}{-AC}$ to $\frac{-AC}{0}$	$\frac{-AC}{0}$ to $\frac{0}{+AC}$
$\text{Sec } A \left(\frac{AC}{AN} \right) \dots$	$\frac{AC}{+AC}$ to $\frac{AC}{0}$	$\frac{AC}{0}$ to $\frac{AC}{-AC}$	$\frac{AC}{-AC}$ to $\frac{AC}{0}$	$\frac{AC}{0}$ to $\frac{AC}{+AC}$

These changes in sign and magnitude of the Sine, Cosine, Tangent, and Secant may be thus exhibited; the signs which belong to them in each right angle being written in a bracket. The symbol ∞ indicates an infinitely large quantity.

A being } between {	0° and 90°	90° and 180°	180° and 270°	270° and 360°
$\text{Sin } A \dots\dots$	0 and 1, (+)	1 and 0, (+)	0 and -1, (-)	-1 and 0, (-)
$\text{Cos } A \dots\dots$	1 ... 0, (+)	0 ... -1, (-)	-1 0, (-)	0 1, (+)
$\text{Tan } A \dots\dots$	0 ... ∞ , (+)	∞ ... 0, (-)	0 ∞ , (-)	∞ 0, (-)
$\text{Sec } A \dots\dots$	1 ... ∞ , (+)	∞ ... -1, (-)	-1 ∞ (-)	∞ 1, (+)


Since $\text{Cos } A$ is never greater than unity, $\text{Versin } A$ (or $1 - \cos A$) is always positive; and its greatest value is when A becomes 180° , when $\text{Cos } A$ becomes -1 , and $\text{Versin } A$ becomes 2 .

22. To shew that $\text{Sin } A = \sin (180^\circ - A)$, or $= -\sin (180^\circ + A)$, or $= -\sin (360^\circ - A)$, or $= -\sin (-A)$; where A is an angle less than a right angle*.

Let $BAC_1 = A = B'AC_2 = B'AC_3 = BAC_4$;
and $AC_1 = AC_2 = AC_3 = AC_4$.

Join C_1C_4 , and C_2C_3 .

It may easily be shewn that the angles at M and N are right angles; and that NC_1, MC_2, MC_3, NC_4 , are equal in magnitude,—as are also AN and AM .



$$\begin{aligned} \text{Now, Sin } A &= \frac{NC_1}{AC_1} = \frac{MC_2}{AC_2} = \sin BAC_2 \\ &= \sin (BAD + DAB' - B'AC_2) \\ &= \sin (180^\circ - A) \dots\dots\dots(1). \end{aligned}$$

$$\begin{aligned} \text{Again, Sin } A &= \frac{NC_1}{AC_1} = -\frac{MC_3}{AC_3}, \text{ since } NC_1 = -MC_3, \\ &= -\frac{MC_3}{AC_3} \\ &= -\sin (BAD + DAB' + B'AC_3) \\ &= -\sin (180^\circ + A) \dots\dots\dots(2). \end{aligned}$$

$$\text{Again, Sin } A = \frac{NC_1}{AC_1} = -\frac{NC_4}{AC_4} = -\frac{NC_4}{AC_4};$$

but $\frac{NC_4}{AC_4}$ is either the sine of the positive angle

$$\{BAD + DAB' + B'AD' + (D'AB - BAC_4)\},$$

or the sine of the negative angle BAC_4 , (7);

$$\therefore \text{Sin } A = -\sin (360^\circ - A) \dots\dots\dots(3),$$

$$\text{or } = -\sin (-A) \dots\dots\dots(4).$$

* In strictness these angles ought to be written thus, $A^\circ, (180 - A)^\circ, (180 + A)^\circ$, &c.

23. In like manner it might be shewn that

$$(1) \cos A = -\cos(180^\circ - A), = -\cos(180^\circ + A), = \cos(360^\circ - A), = \cos(-A).$$

$$(2) \tan A = -\tan(180^\circ - A), = \tan(180^\circ + A), = -\tan(360^\circ - A), = -\tan(-A).$$

$$(3) \sec A = -\sec(180^\circ - A), = -\sec(180^\circ + A), = \sec(360^\circ - A), = \sec(-A).$$

24. If any angle, as BAC , be increased by 360° , the line which bounds it will come into the same position again, and the sine of the angle will therefore remain unaltered. Wherefore $\sin A$ is in all cases the same with $\sin(360^\circ + A)$; and in like manner, $\sin(360^\circ + A) = \sin(2 \times 360^\circ + A)$, and so on. If, therefore, n be any positive integer,

$$\sin A = \sin(n \cdot 360^\circ + A) = \sin(2n \cdot 180^\circ + A) \dots \dots \dots (1).$$

Similarly, $\sin A = \sin(180^\circ - A)$, Art. 22, (1),

$$= \sin\{2n \cdot 180^\circ + (180^\circ - A)\}$$

$$= \sin\{(2n + 1) \cdot 180^\circ - A\} \dots \dots \dots (2).$$

In like manner it appears from (2) and (4) of Art. 22, that

$$\sin A = -\sin\{(2n + 1) \cdot 180^\circ + A\} \dots \dots \dots (3),$$

$$\sin A = -\sin(2n \cdot 180^\circ - A) \dots \dots \dots (4).$$

25. By the same process of reasoning it may be proved from the formulæ of (23) that

$$\cos A = \cos(2n \cdot 180^\circ + A), \quad \text{or} = -\cos\{(2n + 1) \cdot 180^\circ - A\},$$

$$\text{or} = -\cos\{(2n + 1) \cdot 180^\circ + A\}, \quad \text{or} = \cos(2n \cdot 180^\circ - A).$$

$$\text{And, } \tan A = \tan(2n \cdot 180^\circ + A), \quad \text{or} = -\tan\{(2n + 1) \cdot 180^\circ - A\},$$

$$\text{or} = \tan\{(2n + 1) \cdot 180^\circ + A\}, \quad \text{or} = -\tan(2n \cdot 180^\circ - A).$$

Similarly it might be proved that

$$\begin{aligned} \sec A &= \sec (2n \cdot 180^\circ + A), & \text{or} &= -\sec \{(2n + 1) \cdot 180^\circ - A\}, \\ & \text{or} &= -\sec \{(2n + 1) \cdot 180^\circ + A\}, & \text{or} &= \sec (2n \cdot 180^\circ - A)^*. \end{aligned}$$

* The relations, similar to those given in Arts. 22—25, between the trigonometrical functions of $m \cdot 180^\circ \pm A$ and those of A may be established directly, whatever be the magnitude of A , as follows. [See Figure to Art. 22.]

All angles are supposed (Art. 4) to be described by a line revolving from the initial position AB to some other position AC . Let AB be called the initial line, and AC the terminal line of the angle BAC .

Then it will be seen that

1. The *Sines* of all angles whose terminal lines lie on the same side of $B'AB$ will have the same algebraical sign;

2. The *Cosines* will have the same sign for all angles whose terminal lines lie on the same side of $D'AD$;

3. The *Tangents* will have the same sign for all angles whose terminal lines lie in the same quadrant, or in the alternate (or opposite) quadrant.

Now A and $2n \cdot 180^\circ + A$, (where n is any integer, positive or negative), have the same terminal line, and therefore all the trigonometrical functions of $2n \cdot 180^\circ + A$ are the same as those of A .

Again, the terminal line of $(2n + 1) \cdot 180^\circ + A$ will be the terminal line of A produced, and will therefore be in the alternate quadrant and on the sides of both $B'AB$ and $D'AD$ that are opposite to that in which the terminal line of A lies. Hence, the magnitudes of the trigonometrical ratios remaining the same,

$$\sin \{(2n + 1) \cdot 180^\circ + A\} = -\sin A, \quad \cos \{(2n + 1) \cdot 180^\circ + A\} = -\cos A, \quad \tan \{(2n + 1) \cdot 180^\circ + A\} = \tan A.$$

Again, A and $-A$ will have their terminal lines in the adjacent quadrants that are on the same side of $D'AD$ but on opposite sides of $B'AB$. Hence,

$$\sin(-A) = -\sin A, \quad \cos(-A) = \cos A, \quad \tan(-A) = -\tan A.$$

Also $(-A)$ and $2n \cdot 180^\circ - A$ will have the same terminal line, and therefore

$$\begin{aligned} \sin(2n \cdot 180^\circ - A) &= -\sin A, & \cos(2n \cdot 180^\circ - A) &= \cos A, \\ \tan(2n \cdot 180^\circ - A) &= -\tan A. \end{aligned}$$

Again, A and $(2n + 1) \cdot 180^\circ - A$ have their terminal lines in the adjacent quadrants which lie on the same side of $B'AB$ and on opposite sides of $D'AD$. Hence

$$\begin{aligned} \sin \{(2n + 1) \cdot 180^\circ - A\} &= \sin A, & \cos \{(2n + 1) \cdot 180^\circ - A\} &= -\cos A, \\ \tan \{(2n + 1) \cdot 180^\circ - A\} &= -\tan A. \end{aligned}$$

Collecting the above results,

$$\begin{aligned} \sin A &= \sin(2n \cdot 180^\circ + A) = \sin \{(2n + 1) \cdot 180^\circ - A\} \\ &= -\sin(2n \cdot 180^\circ - A) = -\sin \{(2n + 1) \cdot 180^\circ + A\}; \\ \cos A &= \cos(2n \cdot 180^\circ + A) = \cos(2n \cdot 180^\circ - A) \\ &= -\cos \{(2n + 1) \cdot 180^\circ + A\} = -\cos \{(2n + 1) \cdot 180^\circ - A\}; \\ \tan A &= \tan(2n \cdot 180^\circ + A) = -\tan \{(2n + 1) \cdot 180^\circ - A\} \\ &= -\tan(2n \cdot 180^\circ - A) = \tan \{(2n + 1) \cdot 180^\circ + A\}. \end{aligned}$$

26. From Arts. 16, 22, 23, it will appear that

$$\begin{aligned} \sin A &= \cos(90^\circ - A) & \cos A &= \sin(90^\circ - A) \\ \sin A &= \sin(180^\circ - A). & \cos A &= -\cos(180^\circ - A). \\ \tan A &= \cot(90^\circ - A) & \sec A &= \operatorname{cosec}(90^\circ - A) \\ \tan A &= -\tan(180^\circ - A). & \sec A &= -\sec(180^\circ - A). \end{aligned}$$

That is,

The Sine of an angle = cosine of its complement,
or, = sine of its supplement.

Cosine of an angle = sine of its complement,
or, = - cosine of its supplement.

Tangent of an angle = cotangent of its complement,
or, = - tangent of its supplement.

Secant of an angle = cosecant of its complement,
or, = - secant of its supplement.

NOTE. These relations between the sine, cosine, tangent, and secant of an angle and the sine, cosine, tangent, and secant of its complement and supplement, are perpetually occurring in practice, and will often be made use of in the following pages.


These two formulæ also are often useful ;

$$\sin A = \cos(90^\circ - A) = -\cos\{180^\circ - (90^\circ - A)\} = -\cos(90^\circ + A);$$

$$\cos A = \sin(90^\circ - A) = \sin\{180^\circ - (90^\circ - A)\} = \sin(90^\circ + A).$$

27. It will be found necessary to carry in memory the following expressions,

$$(1) \quad \tan A = \frac{NC}{AN} = \frac{\overline{NC}}{\overline{AN}} = \frac{\sin A}{\cos A}.$$



$$(2) \quad \text{Sec } A = \frac{AC}{AN} = \frac{1}{\frac{AN}{AC}} = \frac{1}{\cos A}; \quad \therefore \text{Cos } A = \frac{1}{\text{sec } A}.$$

$$(3) \quad \text{Cot } A = \frac{AN}{NC} = \frac{AC}{\frac{NC}{AC}} = \frac{\cos A}{\sin A}.$$

$$(4) \quad \text{Cot } A = \frac{AN}{NC} = \frac{1}{\frac{NC}{AN}} = \frac{1}{\tan A};$$

$$\therefore \text{Tan } A = \frac{1}{\cot A}.$$

$$(5) \quad \text{Cosec } A = \frac{AC}{NC} = \frac{1}{\frac{NC}{AC}} = \frac{1}{\sin A}; \quad \therefore \text{Sin } A = \frac{1}{\text{cosec } A}.$$

$$(6) \quad AC^2 = NC^2 + AN^2; \quad \therefore 1 = \left(\frac{NC}{AC}\right)^2 + \left(\frac{AN}{AC}\right)^2,$$

$$\text{or } 1 = (\sin A)^2 + (\cos A)^2;^*$$

$$\therefore \text{Sin } A = \sqrt{(1 - \cos^2 A)}, \text{ and } \text{Cos } A = \sqrt{(1 - \sin^2 A)}.$$

$$(7) \quad AC^2 = AN^2 + NC^2;$$

$$\therefore \left(\frac{AC}{AN}\right)^2 = 1 + \left(\frac{NC}{AN}\right)^2, \text{ or } \text{Sec}^2 A = 1 + \tan^2 A;$$

$$\therefore \text{Sec } A = \sqrt{(1 + \tan^2 A)}; \text{ and } \text{Tan } A = \sqrt{(\text{sec}^2 A - 1)}.$$

$$(8) \quad AC^2 = AN^2 + NC^2;$$

$$\therefore \left(\frac{AC}{NC}\right)^2 = \left(\frac{AN}{NC}\right)^2 + 1; \text{ or } \text{Cosec}^2 A = \cotan^2 A + 1;$$

$$\therefore \text{Cosec } A = \sqrt{(1 + \cot^2 A)}; \text{ and } \text{Cot } A = \sqrt{(\text{cosec}^2 A - 1)}.$$

* The powers of the *goniometrical ratios*, as $(\sin A)^2$, $(\cos A)^3$, $(\tan A)^n$, are generally written thus, $\sin^2 A$, $\cos^3 A$, $\tan^n A$.

28. By means of the expressions proved in the last Article the value of any one of the quantities defined in (15) may be found in terms of any other of them. For example:

$$(1) \quad \text{Tan } A = \frac{\sin A}{\sqrt{(1 - \sin^2 A)}}.$$

$$\text{For Tan } A = \frac{\sin A}{\cos A}, \text{ Art. 27, (1); } = \frac{\sin A}{\sqrt{(1 - \sin^2 A)}}; \text{ (27, 6).}$$

$$(2) \quad \text{Tan } A = \frac{\sin A}{\cos A} = \frac{\sqrt{(1 - \cos^2 A)}}{\cos A}.$$

$$(3) \quad \text{Sin } A = \frac{\sin A}{\cos A} \cdot \cos A = \tan A \cdot \frac{1}{\sec A} = \frac{\tan A}{\sqrt{(1 + \tan^2 A)}}.$$

29. The formulæ proved in the last Article will often be found useful to the analyst. The same method of proof is applicable to all other questions of the same kind. Thus, required to express the cosine of an angle in terms of the cosecant, and the cosecant in terms of the versed-sine:

$$(1) \quad \text{Cos } A = \sqrt{(1 - \sin^2 A)} = \sqrt{\left\{1 - \frac{1}{\text{cosec}^2 A}\right\}} = \frac{\sqrt{(\text{cosec}^2 A - 1)}}{\text{cosec } A}.$$

$$(2) \quad \text{Cosec } A = \frac{1}{\sin A} = \frac{1}{\sqrt{(1 - \cos^2 A)}} \\ = \frac{1}{\sqrt{1 - (1 - \text{versin } A)^2}} \\ = \frac{1}{\sqrt{(2 \text{versin } A - \text{versin}^2 A)}}.$$


30. It will be found useful to remember the following values of Sin A , Cos A , Tan A , Sec A .

Sin A	Cos A	Tan A	Sec A
$\sqrt{(1 - \cos^2 A)}$	$\sqrt{(1 - \sin^2 A)}$	$\frac{\sin A}{\sqrt{(1 - \sin^2 A)}}$	$\frac{1}{\sqrt{(1 - \sin^2 A)}}$
$\frac{\tan A}{\sqrt{(1 + \tan^2 A)}}$	$\frac{1}{\sqrt{(1 + \tan^2 A)}}$	$\frac{\sqrt{(1 - \cos^2 A)}}{\cos A}$	$\frac{1}{\cos A}$
$\frac{\sqrt{(\sec^2 A - 1)}}{\sec A}$	$\frac{1}{\sec A}$	$\sqrt{(\sec^2 A - 1)}$	$\sqrt{(1 + \tan^2 A)}$

31. If A be less than half a right angle, or 45° , $\text{Cos } A$ is greater than $\text{Sin } A$.

Let $\angle NAC$ be less than 45° .

Then, since $\angle NAC + \angle NCA = 90^\circ$, $\angle NCA$ is greater than 45° . And in every triangle the greater side is opposite to the greater angle (Eucl. I. 19):



$$\therefore AN > NC; \therefore \frac{AN}{AC} > \frac{NC}{AC}, \text{ or } \text{Cos } A > \text{sin } A.$$

Similarly, for angles between 45° and 90° , it may be shewn that the Cosine is less than the Sine.

32. To find the Sines, Cosines, and Tangents of 45° , 30° , and 60° .

(1) (Fig. Art. 27) Let $\angle NAC = 45^\circ$; $\therefore \angle NCA = 90^\circ - \angle NAC = 45^\circ$;

$$\therefore \frac{AN}{AC} = \frac{NC}{AC}; \text{ or } \text{Sin } 45^\circ = \text{Cos } 45^\circ.$$

$$\text{Also, } AC^2 = AN^2 + CN^2 = 2AN^2;$$

$$\therefore \text{Sin } 45^\circ = \frac{AN}{AC} = \frac{1}{\sqrt{2}}; \text{ Cos } 45^\circ = \frac{1}{\sqrt{2}}; \text{ Tan } 45^\circ = \frac{NC}{AN} = 1.$$


(2) Let ABC be an equilateral and equiangular triangle; each of its angles, therefore, being $\frac{1}{3}$ of two right angles, contains 60° .

Let AD be perpendicular to BC ;

$$\therefore BD = DC = \frac{1}{2}BC = \frac{1}{2}AB;$$

$$\text{and } \angle BAD = \angle DAC = 30^\circ;$$

$$\therefore \text{Sin } 30^\circ = \frac{DB}{AB} = \frac{\frac{1}{2}AB}{AB} = \frac{1}{2}.$$



$$\text{Cos } 30^\circ = \sqrt{\{1 - \text{sin}^2 30^\circ\}} = \sqrt{\{1 - \frac{1}{4}\}} = \frac{\sqrt{3}}{2},$$

$$\text{Tan } 30^\circ = \frac{\text{sin } 30^\circ}{\text{cos } 30^\circ} = \frac{1}{\sqrt{3}}.$$

$$(3) \quad \sin 60^\circ = \cos(90^\circ - 60^\circ), \text{ by (16), } = \cos 30^\circ = \frac{\sqrt{3}}{2},$$

$$\cos 60^\circ = \sin(90^\circ - 60^\circ) = \sin 30^\circ = \frac{1}{2},$$

$$\tan 60^\circ = \frac{\sin 60^\circ}{\cos 60^\circ} = \sqrt{3}.$$

33. Equations like the following may often be solved by means of the relations established in (27) between the different Goniometrical Ratios.

Ex. 1. From the equation, $\sin^2 A + 5 \cos^2 A = 3$, required the value of $\sin A$.

Since $\cos^2 A = 1 - \sin^2 A$, the equation becomes $\sin^2 A + 5\{1 - \sin^2 A\} = 3$; whence $4 \sin^2 A = 2$; and $\sin A = \frac{1}{\sqrt{2}}$.

Ex. 2. From the equations $\sin A = m \sin B$, and $\tan A = n \tan B$, required the values of $\sin A$ and $\cos B$.

For $\sin A$ put x , and for $\cos B$ put y ; $\therefore \sin B = \sqrt{1 - y^2}$.

$$\tan A = \frac{\sin A}{\sqrt{1 - \sin^2 A}} = \frac{x}{\sqrt{1 - x^2}}, \quad (30),$$

$$\tan B = \frac{\sqrt{1 - \cos^2 B}}{\cos B} = \frac{\sqrt{1 - y^2}}{y}, \quad (30).$$

Making these substitutions, the equations become

$$x = m \sqrt{1 - y^2}; \quad \text{and} \quad \frac{x}{\sqrt{1 - x^2}} = n \cdot \frac{\sqrt{1 - y^2}}{y};$$

$$\text{whence, } x = \sin A = \sqrt{\frac{m^2 - n^2}{1 - n^2}}; \quad \text{and} \quad y = \cos B = \frac{n}{m} \sqrt{\frac{1 - m^2}{1 - n^2}}.$$

Ex. 3. Given $\begin{cases} m = \operatorname{cosec} A - \sin A \\ n = \sec A - \cos A \end{cases}$, required to find an equation between m and n in which the angle A shall not appear.

The equations severally give, $m = \frac{\cos^2 A}{\sin A}$, and $n = \frac{\sin^2 A}{\cos A}$;

$$\therefore \frac{n}{m} = \frac{\sin^3 A}{\cos^3 A} = \tan^3 A; \quad \therefore \tan A = \frac{n^{\frac{1}{3}}}{m^{\frac{1}{3}}}.$$

$$\text{Also, } m^2 = \frac{\cos^4 A}{\sin^2 A} = \frac{\cos^2 A}{\tan^2 A} = \frac{1}{\tan^2 A \sec^2 A}.$$

$$\therefore m^2 \tan^2 A \cdot (1 + \tan^2 A) = 1;$$

whence, by substituting its value for $\tan A$,

$$m^{\frac{2}{3}} n^{\frac{2}{3}} (m^{\frac{2}{3}} + n^{\frac{2}{3}}) = 1.$$

CHAPTER III.

GONIOMETRICAL FORMULÆ INVOLVING MORE THAN ONE ANGLE.

34. *Given the Sines and Cosines of two angles, to find the Sines and Cosines of the angles equal to their Sum and to their Difference.*


Let BAC, CAD be two angles represented by A and B respectively.

From D , any point in AD , draw DB and DC perpendiculars on AB and AC ; and from C draw CF and CE perpendiculars on AB and DB .

Then FE is a rectangle; $FB = CE$, and $FC = BE$.

$$\angle CDF = 90^\circ - \angle DCF = \angle FCA$$

$$= A, \text{ since } FC \text{ is parallel to } AE.$$




$$\begin{aligned} \text{Now, } \sin(A + B) &= \frac{BD}{AD} = \frac{BF + FD}{AD} = \frac{EC}{AD} + \frac{FD}{AD} \\ &= \frac{EC}{AC} \cdot \frac{AC}{AD} + \frac{FD}{DC} \cdot \frac{DC}{AD} \\ &= \sin A \cos B + \cos A \sin B \dots \dots \dots (1). \end{aligned}$$

$$\begin{aligned} \text{Also, } \cos(A + B) &= \frac{AB}{AD} = \frac{AE - EB}{AD} = \frac{AE}{AD} - \frac{FC}{AD} \\ &= \frac{AE}{AC} \cdot \frac{AC}{AD} - \frac{FC}{CD} \cdot \frac{CD}{AD} \\ &= \cos A \cos B - \sin A \sin B \dots \dots \dots (2). \end{aligned}$$

Again; let $\angle BAC = A$, and $\angle CAD = B$.

From D , any point in AD , draw DB and DC perpendiculars on AB and AC , CE a perpendicular from C on AB , DF perpendicular to CE .



FB is a rectangle; $FE = DB$, and $FD = EB$;

$$\angle DCF = 90^\circ - \angle ACE = A.$$

$$\begin{aligned} \text{Then, } \sin(A - B) &= \frac{BD}{AD} = \frac{EC - CF}{AD} = \frac{EC}{AD} - \frac{CF}{AD} \\ &= \frac{EC}{AC} \cdot \frac{AC}{AD} - \frac{CF}{CD} \cdot \frac{CD}{AD} \\ &= \sin A \cos B - \cos A \sin B \dots \dots \dots (3). \end{aligned}$$

$$\begin{aligned} \text{Also, } \cos(A - B) &= \frac{AB}{AD} = \frac{AE + EB}{AD} = \frac{AE}{AD} + \frac{EB}{AD} \\ &= \frac{AE}{AC} \cdot \frac{AC}{AD} + \frac{EB}{CD} \cdot \frac{CD}{AD} \\ &= \cos A \cos B + \sin A \sin B \dots \dots \dots (4). \end{aligned}$$

NOTE. The four formulæ of this Article are proved generally for all values of A and B in Appendix iv.

Ex. Given the Signs and Cosines of 45° and 30° , required the Sine and Cosine of 75° , and of 15° .

$$\sin 45^\circ = \cos 45^\circ = \frac{1}{\sqrt{2}}; \quad \sin 30^\circ = \frac{1}{2}, \quad \cos 30^\circ = \frac{\sqrt{3}}{2}, \quad (32).$$

$$\begin{aligned} \sin 75^\circ &= \sin(45^\circ + 30^\circ) = \sin 45^\circ \cdot \cos 30^\circ + \cos 45^\circ \cdot \sin 30^\circ \\ &= \frac{1}{\sqrt{2}} \cdot \frac{\sqrt{3}}{2} + \frac{1}{\sqrt{2}} \cdot \frac{1}{2} = \frac{1}{2\sqrt{2}}(\sqrt{3} + 1). \end{aligned}$$

$$\text{Similarly, } \cos 75^\circ = \cos(45^\circ + 30^\circ) = \frac{1}{2\sqrt{2}}(\sqrt{3} - 1).$$

$$\sin 15^\circ = \sin(45^\circ - 30^\circ) = \frac{1}{2\sqrt{2}}(\sqrt{3} - 1).$$

$$\cos 15^\circ = \cos(45^\circ - 30^\circ) = \frac{1}{2\sqrt{2}}(\sqrt{3} + 1).$$

35. In the figures attached to the last Article, each of the simple angles A and B was represented as *less* than a right angle,—as was also their sum. But of whatever magnitude these simple angles are, if the same construction be made, and proper attention be paid to the signs of the sines and cosines of A and B , the same result will invariably be arrived at. For example, let it be required to prove the formula

$$\sin(A - B) = \sin A \cos B - \cos A \sin B$$

from the annexed figure, where $BAC' = A$, and $C'AD = B$; each angle being greater than a right angle.

From D , any point in AD , draw DC perpendicular to $C'A$ produced; let CEF be perpendicular to AB , DF parallel to AB , DB perpendicular to AB .

Then FB is a rectangle, and $EF = DB$.

$$\text{Now, } \sin(A - B) = \frac{BD}{AD} = \frac{EF}{AD}$$

$$= \frac{CF - CE}{AD}$$

$$= \frac{CF}{CD} \cdot \frac{CD}{AD} - \frac{CE}{AC} \cdot \frac{AC}{AD}$$

$$= \cos FCD \cdot \sin DAC - \sin CAE \cdot \cos DAC,$$

but $\cos FCD = \cos EAC = -\cos(180^\circ - CAB)$, by Art. 26,

$$= -\cos C'AB = -\cos A.$$

$$\sin DAC = \sin(180^\circ - DAC), \text{ Art. 26,} = \sin C'AD = \sin B,$$

$$\sin CAE = \sin C'AB' = \sin A,$$

$$\cos DAC = -\cos(180^\circ - DAC) = -\cos B;$$

$$\therefore \sin(A - B) = -\cos A \sin B + \sin A \cos B$$

$$= \sin A \cos B - \cos A \sin B.$$


36. *If any one of the formulæ of Art. 34, as $\sin(A + B) = \sin A \cdot \cos B + \cos A \cdot \sin B$, be given, the others may be deduced from it.*

For let B become $(-B)$, then

$$\sin(A - B) = \sin\{A + (-B)\} = \sin A \cdot \cos(-B) + \cos A \cdot \sin(-B).$$

But $\cos(-B) = \cos B$, (23), and $\sin(-B) = -\sin B$, (22);

$$\therefore \sin(A - B) = \sin A \cos B - \cos A \sin B.$$



$$\begin{aligned}\text{Again, Cos } (A + B) &= \sin \{90^\circ - (A + B)\}, (16) = \sin \{(90^\circ - A) + (-B)\} \\ &= \sin (90^\circ - A) \cdot \cos (-B) + \cos (90^\circ - A) \cdot \sin (-B) \\ &= \cos A \cos B - \sin A \sin B.\end{aligned}$$

Similarly, $\text{Cos } (A - B)$ may be proved $= \cos A \cos B + \sin A \sin B$.

37. From the formulæ of (34) the Sine or Cosine of the sum of three or more angles may easily be found in terms of the Sines and Cosines of the simple angles.

Given the Sines and Cosines of the angles A, B, C, required the Sine of (A + B + C).

$$\begin{aligned}\text{Sin}(A + B + C) &= \sin \{(A + B) + C\} = \sin (A + B) \cos C + \cos (A + B) \sin C \\ &= (\sin A \cos B + \cos A \sin B) \cos C + (\cos A \cos B - \sin A \sin B) \sin C \\ &= \sin A \cos B \cos C + \sin B \cos A \cos C + \sin C \cos A \cos B \\ &\quad - \sin A \sin B \sin C.\end{aligned}$$

In like manner $\text{Sin}(A + B + C)$, and $\text{Cos}(A + B \pm C)$, may be found in terms of the Sines and Cosines of A, B, C ; and the same method may be applied to the sum of any number of simple angles.

COR. If $A + B + C = (2n + 1) \cdot 180^\circ$, where n is an integer,
since $\sin (2n + 1) 180^\circ = 0$,

the above equation becomes

$$\begin{aligned}\text{Sin } A \sin B \sin C &= \sin A \cos B \cos C \\ &\quad + \sin B \cos A \cos C + \sin C \cos A \cos B.\end{aligned}$$

If $n = 0$, $A + B + C = 180^\circ$, and therefore this equation expresses a relation which exists between the sines and cosines of the three angles of any plane triangle.

38. *To shew that $\text{Sin } 2A = 2 \sin A \cdot \cos A$.*

$$\text{Sin } (A + B) = \sin A \cos B + \cos A \sin B,$$

which becomes, by writing A for B ,

$$\text{Sin } 2A = \sin A \cos A + \cos A \sin A = 2 \sin A \cos A.$$

39. *To shew that (1) $\text{Cos } 2A = \cos^2 A - \sin^2 A$;*

(2) $\text{Cos } 2A = 2 \cos^2 A - 1$; (3) $\text{Cos } 2A = 1 - 2 \sin^2 A$.

$$(1) \quad \text{Cos } (A + B) = \text{cos } A \text{ cos } B - \text{sin } A \text{ sin } B,$$

and writing A for B ,

$$\text{Cos } 2A = \text{cos } A \text{ cos } A - \text{sin } A \text{ sin } A = \text{cos}^2 A - \text{sin}^2 A.$$

$$\text{Again ;} \quad \text{Cos } 2A = \text{cos}^2 A - \text{sin}^2 A,$$

$$\text{and } 1 = \text{cos}^2 A + \text{sin}^2 A ;$$

$$\therefore 1 + \text{cos } 2A = 2 \text{cos}^2 A, \text{ and } 1 - \text{cos } 2A = 2 \text{sin}^2 A.$$

$$(2) \quad \text{Therefore, } \text{Cos } 2A = 2 \text{cos}^2 A - 1.$$

$$(3) \quad \text{And, } \text{Cos } 2A = 1 - 2 \text{sin}^2 A.$$

$$40. \quad \text{To shew that } \begin{cases} \text{Cos } A + \text{sin } A = \pm \sqrt{(1 + \text{sin } 2A)}, \\ \text{Cos } A - \text{sin } A = \pm \sqrt{(1 - \text{sin } 2A)}. \end{cases}$$

Since, $\text{Sin } 2A = 2 \text{sin } A \cdot \text{cos } A$, and $1 = \text{cos}^2 A + \text{sin}^2 A$;
 \therefore by addition and subtraction,

$$1 + \text{sin } 2A = \text{cos}^2 A + 2 \text{sin } A \text{ cos } A + \text{sin}^2 A,$$

$$1 - \text{sin } 2A = \text{cos}^2 A - 2 \text{sin } A \text{ cos } A + \text{sin}^2 A ;$$

$$\therefore \text{Cos } A + \text{sin } A = \pm \sqrt{(1 + \text{sin } 2A)},$$

$$\text{and } \text{Cos } A - \text{sin } A = \pm \sqrt{(1 - \text{sin } 2A)}.$$

41. *To shew that if A be less than 45° ,*

$$\text{then } \begin{cases} \text{Cos } A = \frac{1}{2} \{ \sqrt{(1 + \text{sin } 2A)} + \sqrt{(1 - \text{sin } 2A)} \}, \\ \text{Sin } A = \frac{1}{2} \{ \sqrt{(1 + \text{sin } 2A)} - \sqrt{(1 - \text{sin } 2A)} \}. \end{cases}$$

By (31) if $A < 45^\circ$, $\text{Cos } A$ is $>$ $\text{Sin } A$, and they are both positive; therefore $\text{Cos } A + \text{sin } A$ and $\text{Cos } A - \text{sin } A$ are both positive quantities when A is $<$ 45° ;

$$\therefore \text{Cos } A + \text{sin } A = + \sqrt{(1 + \text{sin } 2A)},$$

$$\text{and } \text{Cos } A - \text{sin } A = + \sqrt{(1 - \text{sin } 2A)} ;$$

whence, by addition and subtraction,

$$2 \cos A = \sqrt{(1 + \sin 2A)} + \sqrt{(1 - \sin 2A)}, \text{ ..}$$

$$2 \sin A = \sqrt{(1 + \sin 2A)} - \sqrt{(1 - \sin 2A)};$$

$$\therefore \begin{cases} \cos A = \frac{1}{2} \{ \sqrt{(1 + \sin 2A)} + \sqrt{(1 - \sin 2A)} \}, \\ \sin A = \frac{1}{2} \{ \sqrt{(1 + \sin 2A)} - \sqrt{(1 - \sin 2A)} \}. \end{cases}$$

42. *The equations of Art. 40 can be applied in any case to determine the Sine and Cosine of A from the Sine of 2A.*

For example, (1) If A be an angle $> 3 \times 45^\circ$ but $< 4 \times 45^\circ$, (*i.e.* if it be any angle comprehended under the form $180^\circ - B$, where B is $< 45^\circ$), $\cos A$ is negative, and greater in magnitude than $\sin A$, which is a positive quantity.

In this case, therefore,

$$\cos A + \sin A = -\sqrt{(1 + \sin 2A)},$$

$$\cos A - \sin A = -\sqrt{(1 - \sin 2A)};$$

$$\therefore \cos A = -\frac{1}{2} \{ \sqrt{(1 + \sin 2A)} + \sqrt{(1 - \sin 2A)} \},$$

$$\sin A = \frac{1}{2} \{ \sqrt{(1 - \sin 2A)} - \sqrt{(1 + \sin 2A)} \}.$$

If at first sight this value of $\sin A$ appear to be negative, it is to be remembered that $\sin 2A$ is a negative quantity, ($2A$ being between 270° and 360°) and therefore $1 - \sin 2A$ is greater than $1 + \sin 2A$; wherefore the value of $\sin A$ is here a positive quantity, as it ought to be.

So (2), If A be a negative angle which is between -45° and -90° , $\cos A$ is a positive quantity, and is less in magnitude than $\sin A$, which is a negative quantity; wherefore the equations to be taken of Art. 40 are

$$\cos A + \sin A = -\sqrt{(1 + \sin 2A)},$$

$$\cos A - \sin A = +\sqrt{(1 - \sin 2A)}.*$$

* PROBLEM. *To determine the limits between which the values of A must lie, which satisfy the equations,*

$$\sin A + \cos A = -\sqrt{(1 + \sin 2A)},$$

$$\cos A - \sin A = -\sqrt{(1 - \sin 2A)}.$$

For positive values of A .

The former equation is fulfilled if A be between $90^\circ + 45^\circ$ and 180° : for the value of $\cos A$, which is negative, is greater for such angles than the value of $\sin A$, which is positive. So A may lie between 135° and 270° ; and between 270° and $270^\circ + 45^\circ$.

Wherefore, if A be between 135° and 315° , the former equation is fulfilled. And it may be shewn, in like manner, that the latter equation is fulfilled if A lie between 45° and 225° .

Wherefore both equations will be fulfilled if A be between 135° and 225° . And as the Sine and Cosine of any angle remain the same if the angle itself be increased by 360° , it follows that all the positive values of A which satisfy both the equations, lie between $n \cdot 360^\circ + 135^\circ$ and $n \cdot 360^\circ + 225^\circ$, [*i.e.* between $(8n + 3) \cdot 45^\circ$ and $(8n + 5) \cdot 45^\circ$], where n is 0 or any positive integer.

And in the same manner

43. Given the Tangents of two angles, to find the Tangents of their Sum and their Difference.

$$\text{Tan}(A + B) = \frac{\sin(A + B)}{\cos(A + B)} = \frac{\sin A \cos B + \cos A \sin B}{\cos A \cos B - \sin A \sin B}$$

and dividing the numerator and the denominator by $\cos A \cos B$,

$$\text{Tan}(A + B) = \frac{\frac{\sin A}{\cos A} + \frac{\sin B}{\cos B}}{1 - \frac{\sin A}{\cos A} \cdot \frac{\sin B}{\cos B}} = \frac{\tan A + \tan B}{1 - \tan A \tan B}$$

Similarly, $\text{Tan}(A - B) = \frac{\tan A - \tan B}{1 + \tan A \tan B}$

COR. 1. If $B = A$, $\text{Tan } 2A = \frac{2 \tan A}{1 - \tan^2 A}$

COR. 2. If $B = 45^\circ$, since $\text{Tan } 45^\circ = 1$(32);

$$\therefore \text{Tan}(A + 45^\circ) = \frac{\tan A + 1}{1 - \tan A} \dots\dots\dots(1).$$

$$= \frac{\frac{\sin A}{\cos A} + 1}{1 - \frac{\sin A}{\cos A}} = \frac{\sin A + \cos A}{\cos A - \sin A} \dots\dots\dots(2).$$

COR. 3. Similarly, $\text{Tan}(A - 45^\circ) = \frac{\tan A - 1}{\tan A + 1} \dots\dots\dots(3);$

$$\text{or} = \frac{\sin A - \cos A}{\sin A + \cos A} \dots\dots\dots(4).$$

COR. 4. $\text{Tan}(A + 45^\circ) + \text{Tan}(A - 45^\circ) = \frac{\tan A + 1}{1 - \tan A} + \frac{\tan A - 1}{1 + \tan A}$
 $= \frac{4 \tan A}{1 - \tan^2 A}, = 2 \tan 2A, \text{ by Cor. 1} \dots\dots\dots(5).$

And in the same manner the negative values of A which satisfy both the equations may be shewn to lie between $-m \cdot 360^\circ - 135^\circ$ and $-m \cdot 360^\circ - 225^\circ$, m being 0 or any positive integer;

i. e. between $-(8m + 3) \cdot 45^\circ$ and $-(8m + 5) \cdot 45^\circ$,

which are of the same form as the limits obtained for the positive values of A .

If A be $< 45^\circ$;

Since $\tan(A + 45^\circ) = \tan - (45^\circ - A) = -\tan(45^\circ - A)$, (23),

the last expression becomes

$$\tan(45^\circ + A) - \tan(45^\circ - A) = 2 \tan 2A \dots\dots\dots (6).$$

44. To shew that $\frac{\tan A + \tan B}{\tan A - \tan B} = \frac{\sin(A + B)}{\sin(A - B)}$.

$$\begin{aligned} \frac{\tan A + \tan B}{\tan A - \tan B} &= \frac{\frac{\sin A}{\cos A} + \frac{\sin B}{\cos B}}{\frac{\sin A}{\cos A} - \frac{\sin B}{\cos B}} \\ &= \frac{\sin A \cos B + \cos A \sin B}{\sin A \cos B - \cos A \sin B} = \frac{\sin(A + B)}{\sin(A - B)}. \end{aligned}$$

45. Given $\tan A, \tan B, \tan C$, to find $\tan(A + B + C)$.

$$\begin{aligned} \tan(A + B + C) &= \tan\{(A + B) + C\} = \frac{\tan(A + B) + \tan C}{1 - \tan(A + B) \tan C}, \quad (43), \\ &= \frac{\frac{\tan A + \tan B}{1 - \tan A \tan B} + \tan C}{1 - \frac{\tan A + \tan B}{1 - \tan A \tan B} \cdot \tan C} \\ &= \frac{\tan A + \tan B + \tan C - \tan A \tan B \tan C}{1 - \tan A \tan B - \tan A \tan C - \tan B \tan C}. \end{aligned}$$

In the same manner the tangent of the sum of four or more angles might be found in terms of the tangents of the simple angles.

COR. If $A + B + C = (2n + 1) \cdot 180^\circ$, n being 0 or an integer, $\tan(A + B + C) = 0$;

$$\therefore \tan A + \tan B + \tan C - \tan A \tan B \tan C = 0,$$

$$\text{or, } \tan A + \tan B + \tan C = \tan A \tan B \tan C.$$

And since if $n = 0$, $A + B + C = 180^\circ$, this relation between the tangents of A, B, C , is one which exists between the tangents of the angles of any plane triangle.

46. To find the values of $\sin 2A$, and $\cos 2A$, in terms of $\tan A$.

$$\sin 2A = 2 \sin A \cdot \cos A, \quad (38), \quad = \frac{2 \sin A}{\cos A} \cdot \cos^2 A.$$

$$= \frac{2 \tan A}{\sec^2 A}, \quad (27. 2), \quad = \frac{2 \tan A}{1 + \tan^2 A}, \quad (27. 7).$$

Again, $\text{Cos } 2A = 2 \cos^2 A - 1$, (39. 2), $= \frac{2}{\sec^2 A} - 1$.

$$= \frac{2}{1 + \tan^2 A} - 1 = \frac{1 - \tan^2 A}{1 + \tan^2 A}.$$

47. The following values of $\text{Sin } 2A$, $\text{Cos } 2A$, $\text{Tan } 2A$ are of frequent occurrence, and necessary to be remembered; those which have not been proved already are easily found after the method of the last Article.

- | | | |
|--|--|--|
| <p>1. $\text{Sin } 2A = 2 \sin A \cos A$.</p> <p>2. $\dots\dots = \frac{2 \tan A}{1 + \tan^2 A}$.</p> <p>3. $\dots\dots = \frac{2 \sqrt{(\sec^2 A - 1)}}{\sec^2 A}$.</p> <p>4. $\text{Tan } 2A = \frac{2 \tan A}{1 - \tan^2 A}$.</p> | | <p>1. $\text{Cos } 2A = \cos^2 A - \sin^2 A$.</p> <p>2. $\dots\dots = 2 \cos^2 A - 1$.</p> <p>3. $\dots\dots = 1 - 2 \sin^2 A$.</p> <p>4. $\dots\dots = \frac{1 - \tan^2 A}{1 + \tan^2 A}$.</p> <p>5. $\dots\dots = \frac{2 - \sec^2 A}{\sec^2 A}$.</p> |
|--|--|--|

48. In the same way the following values of $\text{Sin } 2A$ and $\text{Cos } 2A$ can be found in terms of $\text{Cot } A$, $\text{Cosec } A$, and $\text{Versin } A$.

$$\text{Sin } 2A = \frac{2 \cot A}{1 + \cot^2 A}, \text{ or } = \frac{2 \sqrt{(\text{cosec}^2 A - 1)}}{\text{cosec}^2 A},$$

$$\text{or } = 2(1 - \text{versin } A) \cdot \sqrt{2 \text{vers } A - \text{vers}^2 A}.$$

$$\text{Cos } 2A = \frac{\cot^2 A - 1}{\cot^2 A + 1}, \text{ or } = \frac{\text{cosec}^2 A - 2}{\text{cosec}^2 A}, \text{ or } = 1 - 2(2 \text{vers } A - \text{vers}^2 A).$$

49. The easiest method of deducing such formulæ as these, is first to express $\text{Sin } 2A$ and $\text{Cos } 2A$ in terms of $\text{Sin } A$ and $\text{Cos } A$.

Thus let it be required to prove that $\text{Cos } 2A = \frac{\text{cosec}^2 A - 2}{\text{cosec}^2 A}$.

Since $\text{Cos } 2A = \cos^2 A - \sin^2 A = 1 - 2 \sin^2 A$; and $\text{Sin } A = \frac{1}{\text{cosec } A}$, (27);

$$\therefore \text{Cos } 2A = 1 - \frac{2}{\text{cosec}^2 A} = \frac{\text{cosec}^2 A - 2}{\text{cosec}^2 A}.$$

50. Since $\text{Sin } (A + B) = \sin A \cos B + \cos A \sin B$,
and $\text{Sin } (A - B) = \sin A \cos B - \cos A \sin B$;

∴ by adding and subtracting,

$$\sin(A+B) + \sin(A-B) = 2 \sin A \cos B \dots\dots(1).$$

$$\sin(A+B) - \sin(A-B) = 2 \cos A \sin B \dots\dots(2).$$

Similarly,

$$\cos(A+B) + \cos(A-B) = 2 \cos A \cos B \dots\dots(3).$$

$$\cos(A-B) - \cos(A+B) = 2 \sin A \sin B \dots\dots(4).$$

51. To find the values of $\sin A \pm \sin B$, and $\cos A \pm \cos B$, in terms of the Sines and Cosines of $\frac{1}{2}(A+B)$ and $\frac{1}{2}(A-B)$.

Since

$$A = \frac{1}{2}(A+B) + \frac{1}{2}(A-B), \quad \text{and} \quad B = \frac{1}{2}(A+B) - \frac{1}{2}(A-B);$$

$$\therefore \sin A = \sin \frac{1}{2}(A+B) \cos \frac{1}{2}(A-B) + \cos \frac{1}{2}(A+B) \sin \frac{1}{2}(A-B);$$

$$\sin B = \sin \frac{1}{2}(A+B) \cos \frac{1}{2}(A-B) - \cos \frac{1}{2}(A+B) \sin \frac{1}{2}(A-B),$$

$$\therefore \sin A + \sin B = 2 \sin \frac{1}{2}(A+B) \cos \frac{1}{2}(A-B) \dots\dots(1).$$

$$\sin A - \sin B = 2 \cos \frac{1}{2}(A+B) \sin \frac{1}{2}(A-B) \dots\dots(2).$$

Similarly,

$$\cos A + \cos B = 2 \cos \frac{1}{2}(A+B) \cos \frac{1}{2}(A-B) \dots\dots(3).$$

$$\cos B - \cos A = 2 \sin \frac{1}{2}(A+B) \sin \frac{1}{2}(A-B) \dots\dots(4).$$

These four formulæ (which are of the very greatest utility) might have been deduced from the formulæ of the last Article, by making $A+B=S$ and $A-B=D$; in which case $A = \frac{1}{2}(S+D)$ and $B = \frac{1}{2}(S-D)$.

52. Dividing (2) of Art. 51 by (1),

$$\frac{\sin A - \sin B}{\sin A + \sin B} = \frac{2 \cos \frac{1}{2}(A+B) \sin \frac{1}{2}(A-B)}{2 \sin \frac{1}{2}(A+B) \cos \frac{1}{2}(A-B)} = \frac{\tan \frac{1}{2}(A-B)}{\tan \frac{1}{2}(A+B)}.$$

Similarly, dividing (4) by (3),

$$\frac{\cos B - \cos A}{\cos A + \cos B} = \tan \frac{1}{2}(A+B) \tan \frac{1}{2}(A-B).$$

$$\text{So also, } \frac{\sin A + \sin B}{\cos A + \cos B} = \tan \frac{1}{2}(A \pm B); \quad \text{and} \quad \frac{\sin A \pm \sin B}{\cos B - \cos A} = \cotan \frac{1}{2}(A \mp B).$$

(The upper of the double signs going together, and the lower together).

$$53. \quad \begin{aligned} \tan A \pm \tan B &= \frac{\sin A}{\cos A} \pm \frac{\sin B}{\cos B} \\ &= \frac{\sin A \cos B \pm \cos A \sin B}{\cos A \cos B} = \frac{\sin(A \pm B)}{\cos A \cos B} \end{aligned}$$

Similarly, $\cot B \pm \cot A = \frac{\sin(A \pm B)}{\sin A \sin B}$

(The upper signs going together, and the lower together).

$$54. \quad \begin{aligned} \sin(A + B) \sin(A - B) &= \sin^2 A \cos^2 B - \cos^2 A \sin^2 B \\ &= \sin^2 A (1 - \sin^2 B) - (1 - \sin^2 A) \sin^2 B \\ &= \sin^2 A - \sin^2 B. \end{aligned}$$

Similarly, $\sin(A + B) \sin(A - B) = \cos^2 B - \cos^2 A$.

And in like manner it may be shewn that

$$\cos(A + B) \cos(A - B) = \cos^2 A - \sin^2 B; \text{ or } = \cos^2 B - \sin^2 A.$$

55. To prove that

$$\begin{cases} \sin nA + \sin(n - 2)A = 2 \sin(n - 1)A \cdot \cos A, \\ \cos nA + \cos(n - 2)A = 2 \cos(n - 1)A \cdot \cos A. \end{cases}$$

$$\sin nA = \sin\{(n - 1)A + A\} = \sin(n - 1)A \cos A + \cos(n - 1)A \sin A.$$

So, $\sin(n - 2)A = \sin(n - 1)A \cos A - \cos(n - 1)A \sin A;$

$$\therefore \sin nA + \sin(n - 2)A = 2 \sin(n - 1)A \cos A \dots\dots\dots (1).$$

$$\cos nA = \cos(n - 1)A \cos A - \sin(n - 1)A \sin A,$$

and $\cos(n - 2)A = \cos(n - 1)A \cos A + \sin(n - 1)A \sin A;$

$$\therefore \cos nA + \cos(n - 2)A = 2 \cos(n - 1)A \cos A \dots\dots\dots (2).$$

Cor. If $n = 2$, then, from (1), $\sin 2A = 2 \sin A \cos A,$

..... from (2), $\cos 2A + 1 = 2 \cos^2 A$, or, $\cos 2A = 2 \cos^2 A - 1.$

If $n = 3$; from (1), $\sin 3A = 2 \sin 2A \cos A - \sin A = 4 \sin A \cos^2 A - \sin A$
 $= 4 \sin A (1 - \sin^2 A) - \sin A = 3 \sin A - 4 \sin^3 A.$

..... from (2), $\cos 3A = 2 \cos 2A \cos A - \cos A$
 $= 2 \cos A (2 \cos^2 A - 1) - \cos A = 4 \cos^3 A - 3 \cos A;$

and, by successive substitutions, $\sin 4A, \sin 5A \dots \cos 4A, \cos 5A \dots$ might be found in terms of the Sines and Cosines of A respectively.

56. In like manner,

$$\sin nA - \sin(n - 2)A = 2 \cos(n - 1)A \cdot \sin A \dots\dots(1),$$

$$\cos(n - 2)A - \cos nA = 2 \sin(n - 1)A \cdot \sin A \dots\dots(2).$$

57. From the latter of these formulæ, by means of successive substitutions, the Cosine of nA may be found in terms of the Sines of A and its multiples.

Suppose n to be an even positive integer and $= 2m$.

$$\therefore \cos 2(m-1)A - \cos 2mA = 2 \sin(2m-1)A \sin A,$$

$$\text{so } \cos 2(m-2)A - \cos 2(m-1)A = 2 \sin(2m-3)A \sin A,$$

$$\cos 2(m-3)A - \cos 2(m-2)A = 2 \sin(2m-5)A \sin A,$$

$$\&c. \qquad \qquad \qquad = \qquad \qquad \&c.$$

$$\cos 2(m-m)A - \cos 2\{m - (m-1)\}A = 2 \sin A \sin A.$$

Whence, by addition, Since $\cos 2(m-m)A$, or $\cos 0$, $= 1$, ...

$$1 - \cos 2mA = 2 \sin A \cdot \{\sin(2m-1)A + \sin(2m-3)A + \dots + \sin 3A + \sin A\}.$$

$$\therefore \cos 2mA = 1 - 2 \sin A \cdot \{\sin(2m-1)A + \sin(2m-3)A + \dots + \sin 3A + \sin A\} \dots (1).$$

In like manner, if n were odd and $= 2m+1$, it would appear that

$$\cos(2m+1)A = \cos A - 2 \sin A \cdot \{\sin 2mA + \sin 2(m-1)A + \dots + \sin 4A + \sin 2A\}. \quad (2).$$

COR. If $m=1$, these formulæ give,

$$\cos 2A = 1 - 2 \sin A \sin A = 1 - 2 \sin^2 A.$$

$$\begin{aligned} \cos 3A &= \cos A - 2 \sin A \sin 2A = \cos A - 2 \sin A \cdot 2 \sin A \cos A \\ &= \cos A - 4 \cos A (1 - \cos^2 A) = 4 \cos^3 A - 3 \cos A. \end{aligned}$$

58. To find the Sines and Cosines of 18° , 72° , 36° , and 54° .

$$\sin 36^\circ = \cos(90^\circ - 36^\circ) = \cos 54^\circ,$$

$$\text{or, if } 18^\circ = A, \quad \sin 2A = \cos 3A;$$

$$\therefore 2 \sin A \cos A = 2 \cos 2A \cos A - \cos A, \quad (55);$$

$$\therefore 2 \sin A = 2 \cos 2A - 1 = 2(1 - 2 \sin^2 A) - 1;$$

$$\therefore 4 \sin^2 A + 2 \sin A = 1.$$

And, solving this equation, $\sin A = \frac{\pm \sqrt{5} - 1}{4}$; of which the posi-

tive sign is to be taken, because $\sin 18^\circ$ is a positive quantity.

$$\therefore \frac{1}{4}(\sqrt{5} - 1) = \sin 18^\circ = \cos(90^\circ - 18^\circ) = \cos 72^\circ \dots \dots (1).$$

$$\text{And } \cos^2 18^\circ = 1 - \sin^2 18^\circ = 1 - \frac{6 - 2\sqrt{5}}{16} = \frac{10 + 2\sqrt{5}}{16};$$

$$\therefore \cos 18^\circ = \frac{1}{4} \sqrt{(10 + 2\sqrt{5})} = \sin 72^\circ \dots \dots \dots (2).$$

$$\text{Again; } \sin 54^\circ = \cos 36^\circ = \cos 2 \times 18^\circ = \cos^2 18^\circ - \sin^2 18^\circ.$$

$$\begin{aligned} &= \frac{10 + 2\sqrt{5}}{16} - \frac{6 - 2\sqrt{5}}{16} \\ &= \frac{1}{4} (1 + \sqrt{5}) \dots \dots \dots (3). \end{aligned}$$

$$\cos^2 54^\circ = 1 - \sin^2 54^\circ = 1 - \frac{6 + 2\sqrt{5}}{16} = \frac{1}{16} (10 - 2\sqrt{5});$$

$$\therefore \cos 54^\circ = \frac{1}{4} \sqrt{(10 - 2\sqrt{5})} = \sin 36^\circ \dots \dots \dots (4).$$

59. *If an angle receive any increment, to find the corresponding increment of the Sine of the angle.*

Let the angle A receive an increment α , and let the corresponding increment of the Sine of A be represented by $\Delta \sin A$.

$$\begin{aligned} \text{Then, } \Delta \sin A &= \sin(A + \alpha) - \sin A \\ &= \sin A \cos \alpha + \cos A \sin \alpha - \sin A \\ &= \cos A \sin \alpha - \sin A (1 - \cos \alpha) \\ &= \cos A \sin \alpha \left(1 - \tan A \cdot \frac{2 \sin^2 \frac{1}{2} \alpha}{\sin \alpha} \right) \dots \dots (39). \\ &= \cos A \sin \alpha \left(1 - \tan A \cdot \frac{2 \sin^2 \frac{1}{2} \alpha}{2 \sin \frac{1}{2} \alpha \cos \frac{1}{2} \alpha} \right) \\ &= \cos A \sin \alpha (1 - \tan A \tan \frac{1}{2} \alpha). \end{aligned}$$

Con. If α be very small, $\tan \frac{1}{2} \alpha$ is very small. In this case, if $\tan A$ be not exceedingly large, (that is, if A be not nearly equal to $(2n + 1) 90^\circ$, n being 0 or an integer), $\tan A \tan \frac{1}{2} \alpha$ is a very small quantity, and may be neglected in comparison with unity. When, therefore, α is very small, and A is also not nearly equal to $(2n + 1) 90^\circ$,

$$\Delta \sin A = \cos A \sin \alpha, \text{ very nearly.}$$

Hence it appears that when A is an angle of a triangle, this result cannot be applied to determine the corresponding increment of $\sin A$ which results from A receiving a small given increment, if A be nearly equal to a right angle.

60. *If an angle receive an increment, to find the corresponding decrement of the Cosine of the angle.*

$$\begin{aligned}\Delta \cos A &= \cos(A + \alpha) - \cos A = \cos A \cos \alpha - \sin A \sin \alpha - \cos A \\ &= -\sin A \sin \alpha - \cos A(1 - \cos \alpha) = -\sin A \sin \alpha \left(1 + \cot A \cdot \frac{2 \sin^2 \frac{1}{2} \alpha}{\sin \alpha}\right) \\ &= -\sin A \sin \alpha \left(1 + \cot A \tan \frac{1}{2} \alpha\right).\end{aligned}$$

COR. 1. And as before, if α be very small, and also $\cot A$ be not exceedingly large (i. e. if A be not $= 2n \cdot 90^\circ$ nearly), $\cot A \tan \frac{1}{2} \alpha$ may be neglected with respect to unity, and

$$\Delta \cos A = -\sin A \sin \alpha, \text{ very nearly.}$$

NOTE. Hence it follows, that unless (1st) α be a very small angle, and also (2nd) A be an angle which is not nearly equal to 0° or 180° , this result cannot be applied to any particular case where A is an angle of a triangle.

COR. 2. If A be less than 90° , $\cos A$ is positive, and $\sin A$ being also positive, in this case $\Delta \cos A$ is necessarily negative. Wherefore, in angles less than a right angle, as the angle increases its cosine decreases.

If A be greater than one right angle but less than two, $\cos A$ is negative, and, $\sin A$ being positive, $\Delta \cos A$ is negative. Wherefore, when the angle is greater than one right angle but less than two, as the angle increases the cosine also increases in magnitude, but is negative.

61. *If an angle receive any increment, to find the corresponding increment of the Secant of the angle.*

$$\begin{aligned}\Delta \sec A &= \sec(A + \alpha) - \sec A = \frac{1}{\cos(A + \alpha)} - \frac{1}{\cos A} = \frac{\cos A - \cos(A + \alpha)}{\cos A \cos(A + \alpha)} \\ &= \frac{\sin A \sin \alpha \{1 + \cot A \tan \frac{1}{2} \alpha\}}{\cos A (\cos A \cos \alpha - \sin A \sin \alpha)} \text{ by (60)} \\ &= \frac{\sin A}{\cos A} \cdot \frac{1}{\cos A} \cdot \frac{\sin \alpha}{\cos \alpha} \cdot \frac{1 + \cot A \tan \frac{1}{2} \alpha}{1 - \tan A \tan \alpha} = \tan A \sec A \tan \alpha \cdot \frac{1 + \cot A \tan \frac{1}{2} \alpha}{1 - \tan A \tan \alpha}.\end{aligned}$$

COR. If α be very small, and neither $\tan A$ nor $\cot A$ be very large (that is, if A be not nearly equal to $n \cdot 90^\circ$, when n is 0 or any positive or negative integer), $\cot A \tan \frac{1}{2} \alpha$ and $\tan A \tan \alpha$ will both be so small that they may be neglected when compared with unity. In this case therefore

$$\Delta \sec A = \tan A \sec A \tan \alpha, \text{ very nearly.}$$

NOTE. It is to be remarked that before this result can be applied in any case where A is an angle of a triangle, α must be a very small angle, and A must also not be nearly equal to 0, or 90° , or 180° .

62. If an angle receive any increment, to find the corresponding increment of the Tangent of the angle.

$$\begin{aligned} \Delta \tan A &= \tan(A + \alpha) - \tan A = \frac{\sin(A + \alpha)}{\cos(A + \alpha)} - \frac{\sin A}{\cos A} \\ &= \frac{\sin(A + \alpha) \cos A - \cos(A + \alpha) \sin A}{\cos A (\cos A \cos \alpha - \sin A \sin \alpha)}. \end{aligned}$$

But $\sin(A + \alpha) \cos A - \cos(A + \alpha) \sin A = \sin \{(A + \alpha) - A\} = \sin \alpha$;

$$\therefore \Delta \tan A = \frac{\sin \alpha}{\cos^2 A \cos \alpha \cdot (1 - \tan A \tan \alpha)} = \sec^2 A \tan \alpha \cdot \frac{1}{1 - \tan A \tan \alpha}.$$

COR. If α be very small, and also $\tan A$ be not very large, (that is, if A be not $(2n + 1)90^\circ$ nearly, ... n being 0, or any integer, ...), then

$$\Delta \tan A = \sec^2 A \tan \alpha, \text{ very nearly.}$$

NOTE. If A be an angle of a triangle, this result will not hold when A is nearly equal to a right angle.

63. For a given small increment of A , the increment of the Sine of the angle is $>$, $=$, $<$ the decrement of the Cosine, according as $\cos A$ is $>$, $=$, $<$ $\sin A$; A not being very small, or nearly a multiple of 90° .

For $\Delta \sin A = \cos A \sin \alpha$, if A be not nearly $(2n + 1)90^\circ$; Art. 59. Cor.

$\Delta \cos A = -\sin A \sin \alpha$, if A be not nearly $2n \cdot 90^\circ$; Art. 60. Cor. 1.

Wherefore (n being 0, or any integer), if A be not very small, or nearly a multiple of 90° ,

$$\Delta \sin A \text{ is } >, =, < (-\Delta \cos A), \text{ as } \cos A \text{ is } >, =, < \sin A.$$

COR. In angles less than 90° , $\Delta \sin A$ is $>$ or $<$ $(-\Delta \cos A)$

as A is $<$ or $> 45^\circ \dots (31)$.

64. DEF. By $\text{Tan}^{-1}t$ the angle is indicated of which the tangent is t ; i. e. if $t = \tan A$, then $A = \text{tan}^{-1}t$.

So $\text{Sin}^{-1}s$, and $\text{Cos}^{-1}c$, &c. respectively indicate the angle of which the sine is s , and that of which the cosine is c , &c.

$$65. \text{ To show that } \text{Tan}^{-1}t_1 + \text{tan}^{-1}t_2 = \text{tan}^{-1} \frac{t_1 + t_2}{1 - t_1t_2}, \text{ and}$$

$$\text{Tan}^{-1}t_1 - \text{tan}^{-1}t_2 = \text{tan}^{-1} \frac{t_1 - t_2}{1 + t_1t_2}.$$

Let $\text{Tan } A = t_1$, and $\text{Tan } B = t_2$.

Then, by definition (64), $A = \text{tan}^{-1}t_1$, and $B = \text{tan}^{-1}t_2$.

$$\text{Now } \text{Tan}(A + B) = \frac{\tan A + \tan B}{1 - \tan A \tan B};$$

$$\therefore \text{ by def. } A + B = \text{tan}^{-1} \frac{\tan A + \tan B}{1 - \tan A \tan B};$$

$$\text{Or, } \text{Tan}^{-1}t_1 + \text{tan}^{-1}t_2 = \text{tan}^{-1} \frac{t_1 + t_2}{1 - t_1t_2} \dots \dots \dots (1).$$

$$\text{Similarly, } \text{Tan}^{-1}t_1 - \text{tan}^{-1}t_2 = \text{tan}^{-1} \frac{t_1 - t_2}{1 + t_1t_2} \dots \dots \dots (2).$$

66. If t_1, t_2, \dots, t_n be the tangents of any angles, then

$$\text{Tan}^{-1}t_1 - \text{tan}^{-1}t_n = \text{tan}^{-1} \frac{t_1 - t_2}{1 + t_1t_2} + \text{tan}^{-1} \frac{t_2 - t_3}{1 + t_2t_3} + \dots + \text{tan}^{-1} \frac{t_{n-1} - t_n}{1 + t_{n-1}t_n}.$$

$$\text{For, } \text{Tan}^{-1}t_1 - \text{tan}^{-1}t_2 = \text{tan}^{-1} \frac{t_1 - t_2}{1 + t_1t_2},$$

$$\text{tan}^{-1}t_2 - \text{tan}^{-1}t_3 = \text{tan}^{-1} \frac{t_2 - t_3}{1 + t_2t_3},$$

$$\dots \dots \dots = \dots \dots \dots$$

$$\text{tan}^{-1}t_{n-1} - \text{tan}^{-1}t_n = \text{tan}^{-1} \frac{t_{n-1} - t_n}{1 + t_{n-1}t_n};$$

\therefore by addition, $\text{Tan}^{-1}t_1 - \text{tan}^{-1}t_n$

$$= \text{tan}^{-1} \frac{t_1 - t_2}{1 + t_1t_2} + \text{tan}^{-1} \frac{t_2 - t_3}{1 + t_2t_3} + \dots + \text{tan}^{-1} \frac{t_{n-1} - t_n}{1 + t_{n-1}t_n}.$$

67. Examples of questions solved by the application of formulæ proved in this Chapter and the preceding.

$$(1) \text{ To prove that } \frac{\cos A + \sin^2 A}{\cos A - \sin A} = \sec 2A + \tan 2A.$$

[It is here required to bring the proposed fraction into one of which the denominator shall be $\cos 2A$, or $\cos^2 A - \sin^2 A$. Multiply, therefore, the numerator and denominator by the numerator, and]

$$\begin{aligned} \frac{\cos A + \sin A}{\cos A - \sin A} &= \frac{(\cos A + \sin A)^2}{\cos^2 A - \sin^2 A} = \frac{\cos^2 A + \sin^2 A + 2 \cos A \sin A}{\cos^2 A - \sin^2 A} \\ &= \frac{1 + \sin 2A}{\cos 2A}, \text{ by (27, 6), and (38), (39).} \\ &= \frac{1}{\cos 2A} + \frac{\sin 2A}{\cos 2A} = \sec 2A + \tan 2A. \end{aligned}$$

$$(2) \text{ To prove that } \cos 2A = \frac{1}{1 + \tan 2A \tan A}.$$

[The equation becomes, when inverted, $\frac{1}{\cos 2A} = 1 + \tan 2A \tan A$; where if the right-hand side were expressed in terms of the sines and cosines of A and $2A$, and then put into a fractional form, the denominator would be $\cos 2A \cos A$, and the numerator would involve the sines and cosines of the same angles. The first step therefore is to express $\frac{1}{\cos 2A}$ in a fraction of such a form.]

$$\begin{aligned} \text{Now, } \frac{1}{\cos 2A} &= \frac{\cos A}{\cos 2A \cos A} = \frac{\cos(2A - A)}{\cos 2A \cos A} \\ &= \frac{\cos 2A \cos A + \sin 2A \sin A}{\cos 2A \cos A} = 1 + \frac{\sin 2A \sin A}{\cos 2A \cos A} = 1 + \tan 2A \tan A; \\ \therefore \cos 2A &= \frac{1}{1 + \tan 2A \tan A}. \end{aligned}$$

NOTE. In the following Examples it is required to determine an angle from some given relation between its Goniometrical Ratios and those of either a multiple of the angle sought, or of some given angle; and conversely.

(3) Determine a value of A that will satisfy the equation $\sin 2A = \sin A$.

$$\sin A = \sin 2A = 2 \sin A \cos A \dots \dots \dots (33);$$

$$\therefore 2 \cos A = 1, \text{ and } \cos A = \frac{1}{2}. \text{ Wherefore } A = 60^\circ \dots \dots \dots (32).$$

(4) Determine values of B that will satisfy the equation,

$$\sin A + \sin(2B + A) - \sin(2B - A) = \sin(B + A) - \sin(B - A);$$

$$\therefore \sin A + 2 \cos 2B \sin A = 2 \cos B \sin A; \quad \text{Art. 50, (2).}$$

$$\therefore 1 + 2 \cos 2B = 2 \cos B;$$

$$\therefore 1 + 2(2 \cos^2 B - 1) = 2 \cos B \dots \dots \dots (39).$$

$$\text{Whence } \cos B = \frac{1}{4}(1 \pm \sqrt{5});$$

$$\text{and, Art. 58, (3), } \frac{1}{4}(1 + \sqrt{5}) = \cos 36^\circ.$$

$$\text{Art. 58, (1), } \frac{1}{4}(1 - \sqrt{5}) = -\frac{1}{4}(\sqrt{5} - 1) = -\cos 72^\circ = \cos(180^\circ - 72^\circ) = \cos 108^\circ.$$

$$\therefore B \text{ is } 36^\circ, \text{ or } 108^\circ.$$

(5) To prove that $2 \cos 11^\circ, 15' = \sqrt{2 + \sqrt{\{2 + \sqrt{2}\}}}$.

$$\cos 45^\circ = \frac{1}{\sqrt{2}}; \therefore 2 \cos 45^\circ = \sqrt{2},$$

$$\therefore 2 \cdot \{2 \cos^2 \frac{1}{2} \cdot 45^\circ - 1\} = \sqrt{2}, \therefore 2 \cos \frac{1}{2} \cdot 45^\circ = \sqrt{2 + \sqrt{2}};$$

$$\therefore 2 \cdot \{2 \cos^2 \frac{45^\circ}{2} - 1\} = \sqrt{2 + \sqrt{2}};$$

$$\therefore 2 \cos \frac{45^\circ}{2}, \text{ or } 2 \cos 11^\circ, 15' = \sqrt{2 + \sqrt{2 + \sqrt{2}}}.$$

Cor. By repeating the same process n times, it would appear in like manner that $2 \cos \frac{45^\circ}{2^n} = \sqrt{2 + \sqrt{\{2 + \sqrt{\{2 + \sqrt{\{2 + \dots\}}\}}\}}}$, where 2, with the sign of a square root over it, appears $n + 1$ times in the second member of the equation, the square root every time reaching to the end of the expression.

(6) If $x \cdot \tan A = (\sqrt{1+x} - 1) \cdot (\sqrt{1-x} + 1)$, required to prove that $x = \sin 4A$ satisfies the equation,

$$x \cdot \tan A = (\sqrt{1+x} - 1) \cdot (\sqrt{1-x} + 1) \cdot \frac{\sqrt{1+x} + 1}{\sqrt{1+x} + 1} = x \cdot \frac{\sqrt{1-x} + 1}{\sqrt{1+x} + 1};$$

$$\therefore \sqrt{1-x} + 1 = (\sqrt{1+x} + 1) \tan A;$$

$$\therefore (1 - \tan A)^2 = \{\sqrt{1+x} \tan A - \sqrt{1-x}\}^2;$$

Whence, $2 \tan A = x \{1 - \tan^2 A\} + 2\sqrt{1-x^2} \tan A$;

$$\therefore 1 = x \cdot \frac{1 - \tan^2 A}{2 \tan A} + \sqrt{1-x^2}, \quad \therefore \sqrt{1-x^2} = 1 - x \cot 2A \dots (47);$$

$$\therefore 1 - x^2 = 1 - 2x \cot 2A + x^2 \cot^2 2A;$$

$$\therefore x^2(1 + \cot^2 2A) - 2x \cot 2A = 0;$$

$$\therefore x = 0; \quad \text{or } x = \frac{2 \cot 2A}{1 + \cot^2 2A} = 2 \cdot \frac{\cot 2A}{\operatorname{cosec}^2 2A} = 2 \sin 2A \cos 2A = \sin 4A.$$

(7) To prove that $\tan^{-1} \frac{1}{3} + \tan^{-1} \frac{1}{5} + \tan^{-1} \frac{1}{7} + \tan^{-1} \frac{1}{8} = 45^\circ$.

$$\tan^{-1} \frac{1}{3} + \tan^{-1} \frac{1}{5} = \tan^{-1} \frac{\frac{1}{3} + \frac{1}{5}}{1 - \frac{1}{3} \cdot \frac{1}{5}}, \quad (65). \quad = \tan^{-1} \frac{4}{7},$$

$$\text{So } \tan^{-1} \frac{1}{7} + \tan^{-1} \frac{1}{8} = \tan^{-1} \frac{\frac{1}{7} + \frac{1}{8}}{1 - \frac{1}{7} \cdot \frac{1}{8}} = \tan^{-1} \frac{3}{11};$$

$$\begin{aligned} \therefore \tan^{-1} \frac{1}{3} + \tan^{-1} \frac{1}{5} + \tan^{-1} \frac{1}{7} + \tan^{-1} \frac{1}{8} \\ = \tan^{-1} \frac{4}{7} + \tan^{-1} \frac{3}{11} = \tan^{-1} \frac{\frac{4}{7} + \frac{3}{11}}{1 - \frac{4}{7} \cdot \frac{3}{11}} = \tan^{-1} \frac{65}{65} = \tan^{-1} 1 = 45^\circ. \end{aligned}$$

68. The Appendices I, II, III, on the Logarithmic Tables of Numbers and of Goniometrical Ratios, ought to be read before entering on the next Chapter.

CHAPTER IV.


ON THE SOLUTION OF TRIANGLES.

69. A TRIANGLE consists of six parts, namely three sides and three angles. When three of these parts are given, (except they be the three angles), it will be shewn that the other three can, generally, be determined.

The number of degrees in the angles of a triangle will be designated by the letters A, B, C , which are placed at the angular points of the triangle; and the length of the sides respectively opposite to the angles A, B, C , by the letters a, b, c .

70. *The Sines of the Angles of a Triangle are proportional to the Sides respectively opposite to them.*

Let ABC be the triangle. From C draw CD perpendicular to AB , or AB produced either way.



$$\text{Then } \sin CAB = \sin CAD = \frac{DC}{CA}.$$

(With reference to the third figure, see Art. 26.)

$$\text{Also, } \sin CBA = \sin CBD = \frac{DC}{CB};$$

$$\therefore \frac{\sin CAB}{\sin CBA} = \frac{CB}{CA};$$

$$\therefore \frac{\sin A}{\sin B} = \frac{a}{b}; \quad \text{or, } \frac{\sin A}{a} = \frac{\sin B}{b}.$$

In like manner, if a perpendicular were let fall from B upon the side opposite to it, or that side produced, it might be proved that

$$\frac{\sin A}{a} = \frac{\sin C}{c}.$$

Wherefore, $\frac{\sin A}{a} = \frac{\sin B}{b} = \frac{\sin C}{c}$;—the magnitudes of the lines a, b, c being represented by the *number* of units of length they respectively contain: for otherwise $\sin A$ and a would not be quantities of the same kind, and consequently no ratio could exist between them.

71. Since (Euclid, I. 32), the interior angles of a triangle are together equal to two right angles; $\therefore A + B + C = 180^\circ$.

$$\text{Also, } \frac{\sin A}{\sin B} = \frac{a}{b}; \quad \text{and } \frac{\sin B}{\sin C} = \frac{b}{c}.$$

And three of the parts of the triangle being given, the remaining three parts may be determined by these three equations.

One, at least, of the three given parts must be *sine*, or the *ratios* only are given which a, b, c bear to each other, and their *magnitudes* cannot be determined, because there are but *two* equations given, namely, $\frac{\sin A}{\sin B} = \frac{a}{b}$, and $\frac{\sin B}{\sin C} = \frac{b}{c}$, for determining the *three* unknown quantities a, b, c .

And this is also apparent from the consideration that an indefinite number of triangles having the same angles, of all possible degrees of magnitude, may be formed by drawing lines parallel to the sides of a given triangle.

72. There is however one case, commonly called "*the ambiguous case*," in which the equations of the last Article are not sufficient to determine the triangle when three of the parts are given.

If two Sides be given, and an Angle opposite to one of them (a, b, A), the triangle can be determined only when the side opposite to the given angle is the greater of the two given sides; i. e. when a is greater than b .

The equations of the last Article are

$$(i) B + C = 180^\circ - A, \quad (ii) \sin B = \frac{b}{a} \sin A, \quad (iii) c = b \cdot \frac{\sin C}{\sin B}.$$

If B can be determined from (ii), C and c are known from (i) and (iii), and the triangle is determined. But since the sine of an angle is equal to the sine of its supplement, there are two values of the angle B which satisfy (ii), the one greater and the other less than 90° .

(1) Let a be greater than b ; $\therefore A > B$. Eucl. I. 18.


Now B cannot be greater than 90° ; for if it could, then $A + B$ would be greater than 180° , which is impossible, (Eucl. I. 17): $\therefore B < 90^\circ$, and the lesser angle which satisfies (ii) is to be taken for the value of B .

(2) Let a be less than b ; $\therefore A < B$.

In this case, since the only limitations, $B + A < 180^\circ$, and $B > A$, may be satisfied whether B be greater or less than 90° , it is impossible to determine from (ii) whether B be $>$ or $< 90^\circ$.

Thus,—taking the annexed figure,—if $CB = CB'$; it is evident that both the triangles $ABC, AB'C$ have a, b, A of the same values; also, in this case,

$\angle A$ is less than the exterior angle CBB' , or the exterior angle $CB'D$, i. e. A is less than $CB'A$, and therefore is less than CBA ; and also $a < b$. Which agrees with what has been asserted above.



73. To find the Cosine of an angle of a Triangle in terms of the Sides. (Fig. Art. 70.)

Let ABC be a triangle, and from C draw CD perpendicular to AB , or AB produced either way.

Then, figs. 1, 2 of Art. 70, $CB^2 = AC^2 + AB^2 - 2AB \cdot AD$,
Eucl. II. 13.

fig. 3, $CB^2 = AC^2 + AB^2 + 2AB \cdot AD$, Eucl. II. 12.

And $\frac{AD}{AC} = \cos CAD$, = $\cos CAB$ in figs. 1, 2.
= $-\cos CAB$ in fig. 3... (26).

Therefore, in each of these cases,

$$CB^2 = AC^2 + AB^2 - 2AB \cdot AC \cdot \cos CAB;$$

$$\therefore \cos CAB = \frac{AC^2 + AB^2 - CB^2}{2AB \cdot AC}; \quad \text{Or, } \cos A = \frac{b^2 + c^2 - a^2}{2bc}.$$

74. To shew that $\cos \frac{1}{2} A = \sqrt{\frac{S \cdot (S - a)}{bc}}$; where
 $S = \frac{1}{2}(a + b + c)$.

$$\begin{aligned} \text{For } 1 + \cos A &= 1 + \frac{b^2 + c^2 - a^2}{2bc} = \frac{b^2 + 2bc + c^2 - a^2}{2bc} \\ &= \frac{(b + c)^2 - a^2}{2bc} = \frac{(b + c + a)(b + c - a)}{2bc}. \end{aligned}$$

Now, $1 + \cos A = 2 \cos^2 \frac{1}{2} A$, (39, 2).

$$\text{And } S - a = \frac{1}{2}(a + b + c) - a = \frac{1}{2}(b + c - a);$$

$$\therefore 2 \cos^2 \frac{1}{2} A = \frac{(b + c + a) \cdot (b + c - a)}{2bc} = \frac{2S \cdot 2(S - a)}{2bc};$$

$$\therefore \cos \frac{1}{2} A = \sqrt{\frac{S \cdot (S - a)}{bc}}.$$

75. To shew that $\sin \frac{1}{2} A = \sqrt{\frac{(S - b) \cdot (S - c)}{bc}}$.

By Art. 39, (3),

$$\begin{aligned} 2 \sin^2 \frac{1}{2} A &= 1 - \cos A = 1 - \frac{b^2 + c^2 - a^2}{2bc} = \frac{a^2 - (b - c)^2}{2bc} \\ &= \frac{(a + c - b)(a + b - c)}{2bc}. \end{aligned}$$

$$\text{Now } S - b = \frac{1}{2}(a + b + c) - b = \frac{1}{2}(a + c - b),$$

$$\text{and so } S - c = \frac{1}{2}(a + b - c);$$

$$\therefore 2 \sin^2 \frac{1}{2} A = \frac{2(S-b) \cdot 2(S-c)}{2bc}.$$

$$\therefore \sin \frac{1}{2} A = \sqrt{\frac{(S-b)(S-c)}{bc}}.$$

The positive sign of the root must be taken both here and in the last Article; because A , being an angle of a triangle, is less than 180° , and therefore $\text{Cos } \frac{1}{2}A$ and $\text{Sin } \frac{1}{2}A$ are necessarily positive quantities.

$$\begin{aligned} 76. \quad \text{Sin}^2 A &= 1 - \text{cos}^2 A = (1 + \text{cos } A) \cdot (1 - \text{cos } A) \\ &= \left\{ 1 + \frac{b^2 + c^2 - a^2}{2bc} \right\} \cdot \left\{ 1 - \frac{b^2 + c^2 - a^2}{2bc} \right\} \\ &= \frac{1}{4b^2c^2} (b + c - a)(b + c + a)(a + b - c)(a + c - b) \\ &= \frac{1}{4b^2c^2} \cdot 2S \cdot 2(S-a) \cdot 2(S-b) \cdot 2(S-c) \\ &= \frac{4}{b^2c^2} S(S-a)(S-b)(S-c); \\ \therefore \text{Sin } A &= \frac{2}{bc} \sqrt{S(S-a)(S-b)(S-c)}. \end{aligned}$$

$$\text{So, } \text{Tan } \frac{1}{2} A = \frac{\sin \frac{1}{2} A}{\cos \frac{1}{2} A} = \sqrt{\frac{(S-b)(S-c)}{S(S-a)}}.$$

77. To explain the meaning of the double sign of the second member of the equation

$$\text{Cos } \frac{1}{2} A = \pm \sqrt{\frac{S(S-a)}{bc}}.$$

$$\text{Since } \text{Cos}(2n \cdot 180^\circ \pm A) = \text{cos } A, (25); \therefore \text{cos}(2n \cdot 180^\circ \pm A) = \frac{b^2 + c^2 - a^2}{2bc}.$$

Whence it may be proved in the same manner as in (74), that

$$\text{Cos}(n \cdot 180^\circ \pm \frac{1}{2} A) = \pm \sqrt{\frac{S \cdot (S-a)}{bc}}.$$

Now the first member of this equation ought to furnish two sets of angles whose cosines are of the same magnitude but are of different signs.

First. Let n be even, and $= 2m$.

Then $\text{Cos}(n \cdot 180^\circ \pm \frac{1}{2}A) = \text{Cos}(2m \cdot 180^\circ \pm \frac{1}{2}A) = \text{Cos}(m \cdot 360^\circ \pm \frac{1}{2}A)$; which, by making m equal to 0, 1, 2, 3 . . . successively, gives the series of angles $\pm \frac{1}{2}A$, $360^\circ \pm \frac{1}{2}A$, $2 \times 360^\circ \pm \frac{1}{2}A$, $3 \times 360^\circ \pm \frac{1}{2}A$, . . . all the cosines of which are of the same magnitude as that of $\pm \frac{1}{2}A$, or of $+\frac{1}{2}A$.

Second. Let n be odd, and $= 2m+1$.

Then $\text{Cos}\{n \cdot 180^\circ \pm \frac{1}{2}A = \text{Cos}(2m \cdot 180^\circ + 180^\circ \pm \frac{1}{2}A)\}$; which, by making m equal to 0, 1, 2, 3 . . . successively, gives the series of angles $180^\circ \pm \frac{1}{2}A$, $360^\circ + (180^\circ \pm \frac{1}{2}A)$, $2 \times 360^\circ + (180^\circ \pm \frac{1}{2}A)$, . . . all the cosines of which are of the same magnitude as that of $180^\circ \pm \frac{1}{2}A$, which

$$= \text{Cos}(180^\circ \pm \frac{1}{2}A) = -\text{Cos} \frac{1}{2}A.$$

Wherefore the first member of the equation *does* furnish two sets of angles whose cosines are of the same magnitudes but are of different signs.

And in like manner sets of angles may be determined corresponding to each of the signs affecting the expressions which have been obtained as the values of $\text{Sin} \frac{1}{2}A$, $\text{Tan} \frac{1}{2}A$, and $\text{Sin} A$.

I. ON THE SOLUTION OF RIGHT-ANGLED TRIANGLES.

78. *The right angle, a side, and another part being given, to determine the remaining parts.*

Let ABC be a right-angled triangle, C being the right angle.


(1) Let c and A be the other given parts.

$$\text{Then } \frac{b}{c} = \text{Cos } A; \quad \text{and } \frac{a}{c} = \text{Sin } A;$$

$$\therefore \begin{cases} \frac{1}{10}b = \frac{1}{10}c + L \text{Cos } A - 10; \text{ which determines } b, \\ \frac{1}{10}a = \frac{1}{10}c + L \text{Sin } A - 10; \text{ which determines } a. \end{cases}$$

Also, $\angle B = 90^\circ - A$; which determines B .

In like manner, if B were the given angle, A might be determined.



(2) Let Λ and b be given.

Then $\frac{c}{b} = \sec A$; and $\frac{a}{b} = \tan A$;

$$\therefore \begin{cases} l_{10}c = l_{10}b + L \sec A - 10; \text{ which determines } c, \\ l_{10}a = l_{10}b + L \tan A - 10; \text{ which determines } a. \end{cases}$$

Also, $B = 90^\circ - A$; which determines B .

(3) Let Λ and a be given.

Then $\frac{a}{b} = \tan A$; $\therefore b = \frac{a}{\tan A}$; and $l_{10}b = l_{10}a - L \tan A + 10$.

Again, $\frac{a}{c} = \sin A$; $\therefore c = \frac{a}{\sin A}$, or $c = a \cdot \operatorname{cosec} A$;

And $l_{10}c = l_{10}a - L \sin A + 10$. Also, $B = 90^\circ - A$.

(4) Let a and b be given.

Then, $\tan A = \frac{a}{b}$; $\therefore L \tan A = l_{10}a - l_{10}b + 10$.

Again, $B = 90^\circ - A$.

Again, $\frac{c}{b} = \sec A$; $\therefore l_{10}c = l_{10}b + L \sec A - 10$.

The equation $c = \sqrt{(a^2 + b^2)}$ would give c ; but the operation of determining c is tedious, particularly if a and b be large numbers.

(5) Let c and a be given.

Then, $\sin A = \frac{a}{c}$; and $\therefore L \sin A = l_{10}a - l_{10}c + 10$.

Again, $\frac{b}{c} = \cos A$; $\therefore l_{10}b = l_{10}c + L \cos A - 10$.

$$\text{Or } b^2 = c^2 - a^2 = (c + a)(c - a);$$

$$\therefore l_{10}b = \frac{1}{2} \{ l_{10}(c + a) + l_{10}(c - a) \};$$

which determines b without previously finding A .

79. Different methods must be used in different cases to determine the unknown quantities; for what is said in Appendix III. 11, must always be carefully borne in mind, and such a formula must be selected in each case as is calculated to give the result with the greatest practical degree of accuracy.

Thus in the last case, if b be very small compared with a and c , the angle A is nearly a right angle, and the increment of $\sin A$ corresponding to a small given increment α of the angle, [i. e. $\cos A \cdot \sin \alpha \cdot \{1 - \tan A \cdot \tan \frac{1}{2}\alpha\}, \dots (59)$], is inconsiderable, and does not, besides, vary nearly as the increment of the angle. In this case therefore the value of $\sin A$ cannot be determined from the Tables with any great degree of accuracy. (App. III. 11.)

A better way of determining A in such a case is first to find the value of b , and then to determine A from its cosine.

$$\begin{aligned} \text{Thus } \cos A &= \frac{b}{c}; \quad \therefore L \cos A = 1_{10}b - 1_{10}c + 10, \\ &= 1_{10}\sqrt{(c^2 - a^2)} - 1_{10}c + 10, \\ &= \frac{1}{2} \{1_{10}(c+a) + 1_{10}(c-a)\} + 10 - 1_{10}c. \end{aligned}$$

80. EXAMPLE. Given $c = 365.1$, and $a = 348.3$, to find A .

(The Logarithms employed in this Example and others, will generally be found in the three pages of Logarithms subjoined to Appendices I. and II.)

Here, $c + a = 713.4$, and $c - a = 16.8$.

Performing the operations indicated in the last line of the last Article,

$$\begin{array}{r} 1_{10} 713.4 = 2.8533331 \\ 1_{10} 16.8 = 1.2253093 \\ \hline 2) 4.0786424 \\ \hline 2.0393212 \\ 10 \\ \hline 12.0393212 \\ 1_{10} 365.1 = 2.5624118 \\ \hline 9.4769094 \\ \text{And } L \cos 72^\circ, 33' = 9.4769380 \text{ See App. III. 9, Ex. 4.} \\ \hline \text{Difference} = \quad 286 \end{array}$$

Now Diff. for $1''$ is in this case 67.016 ;

$$\text{and } \frac{286}{67.016} = 4.267 = 4.27 \text{ nearly,}$$

$$\therefore A = 72^\circ, 33', 4'' 27, \text{ nearly.}$$

II. ON THE SOLUTION OF OBLIQUE-ANGLED TRIANGLES.

81. *Let two angles and the side between them be given.* (A, C, b)

Since $A + B + C = 180^\circ$, $\therefore B = 180^\circ - (A + C)$;
which determines B .

$$\text{Again, } a = b \cdot \frac{\sin A}{\sin B};$$

$$\therefore l_{10}a = l_{10}b + L \sin A - L \sin B; \text{ which determines } a.$$

$$\text{Again, } c = b \cdot \frac{\sin C}{\sin B};$$

$$\therefore l_{10}c = l_{10}b + L \sin C - L \sin B; \text{ which determines } c.$$



If $A + C$ be $< 90^\circ$, the value of B is not required in order to determine a and c ;

$$\text{For since } \sin B = \sin \{180^\circ - (A + C)\} = \sin (A + C);$$

$$\therefore l_{10}a = l_{10}b + L \sin A - L \sin (A + C),$$

$$\text{and } l_{10}c = l_{10}b + L \sin C - L \sin (A + C).$$

82. *Let two angles and a side opposite to one of them be given.* (A, C, a)

$$\text{Then, } B = 180^\circ - (A + C).$$

$$\text{Again, } b = a \cdot \frac{\sin B}{\sin A}; \therefore l_{10}b = l_{10}a + L \sin B - L \sin A,$$

$$\text{or } = l_{10}a + L \sin (A + C) - L \sin A.$$

$$\text{Again, } c = a \cdot \frac{\sin C}{\sin A}; \therefore l_{10}c = l_{10}a + L \sin C - L \sin A.$$

83. *Let two sides and the angle included between them be given.* (c, A, b)

First, to determine B and C .

$$B + C = 180^\circ - A; \quad \therefore \frac{1}{2}(B + C) = \frac{1}{2}(180^\circ - A).$$

Also, $\frac{b}{c} \frac{\sin B}{\sin C}$; $\therefore \frac{\frac{b}{c} - 1}{\frac{b}{c} + 1} = \frac{\frac{\sin B}{\sin C} - 1}{\frac{\sin B}{\sin C} + 1}$;

$\therefore \frac{b-c}{b+c} = \frac{\sin B - \sin C}{\sin B + \sin C} = \frac{\tan \frac{1}{2}(B-C)}{\tan \frac{1}{2}(B+C)}$, (52).

And $\text{Tan } \frac{1}{2}(B+C) = \text{tan } \frac{1}{2}(180^\circ - A) = \text{cot } \frac{1}{2}A$;

$\therefore \text{Tan } \frac{1}{2}(B-C) = \frac{b-c}{b+c} \cdot \text{cot } \frac{1}{2}A$; •

$\therefore L \tan \frac{1}{2}(B-C) = \frac{1}{10}(b-c) - \frac{1}{10}(b+c) + L \cot \frac{1}{2}A$. (2).

$\frac{1}{2}(B-C)$ being thus determined, and $\frac{1}{2}(B+C)$ being known,

$B = \frac{1}{2}(B+C) + \frac{1}{2}(B-C)$
and $C = \frac{1}{2}(B+C) - \frac{1}{2}(B-C)$ are known;

And $a = c \cdot \frac{\sin A}{\sin C}$ determines a .

84. *The side a can be determined without previously determining B and C. (c, A, b.)*

For $a^2 = b^2 + c^2 - 2bc \cdot \cos A$
 $= b^2 + c^2 - 2bc (1 - 2 \sin^2 \frac{1}{2}A) \dots \dots 39, (3).$
 $= (b-c)^2 + 4bc \sin^2 \frac{1}{2}A$
 $= (b-c)^2 \left\{ 1 + \frac{4bc}{(b-c)^2} \cdot \sin^2 \frac{1}{2}A \right\}$
 $= (b-c)^2 \left\{ 1 + \left(\frac{2\sqrt{bc}}{b-c} \cdot \sin \frac{1}{2}A \right)^2 \right\}.$

Now $\frac{2\sqrt{bc}}{b-c} \cdot \sin \frac{1}{2}A$ may be of any magnitude and sign, and therefore there is *some* angle of which the tangent is equal to this quantity. Let θ be the angle.

Then $\text{Tan } \theta = \frac{2\sqrt{bc}}{b-c} \cdot \sin \frac{1}{2}A \dots \dots \dots (1).$

$$\text{And } a^2 = (b - c)^2 (1 + \tan^2 \theta) = (b - c)^2 \sec^2 \theta;$$

$$\therefore a = (b - c) \sec \theta \dots \dots \dots (2).$$

$$\begin{aligned} \text{From (1), } L \tan \theta &= \text{I}_{10} 2 + \frac{1}{2} \text{I}_{10} b + \frac{1}{2} \text{I}_{10} c - \text{I}_{10} (b - c) + L \sin \frac{1}{2} A, \\ &= \frac{1}{2} (\text{I}_{10} b + \text{I}_{10} c) + \text{I}_{10} 2 + L \sin \frac{1}{2} A - \text{I}_{10} (b - c); \end{aligned}$$

which determines θ .

$$\text{From (2), } \text{I}_{10} a = \text{I}_{10} (b - c) + L \sec \theta - 10; \text{ which determines } a.$$

85. If $b = c$ nearly, $b - c$ is very small, and $\frac{2\sqrt{bc}}{b+c} \cdot \sin \frac{1}{2} A$, the value of $\tan \theta$, is very large, (unless $\frac{1}{2} A$, and therefore $\sin \frac{1}{2} A$, be very small.) And since the tangents of angles which are nearly right angles are very large, θ in this case is nearly a right angle.

Now if it be required to find from its tangent, an angle which does not consist of a certain number of degrees and minutes exactly, the additional seconds have to be determined on the principle that the increment of the tabular logarithm varies as the increment of the angle. But when the angle is equal to $(2n + 1) \cdot 90^\circ$ nearly, this principle does not obtain for the tangent, 62, Cor., App. III. 11; and therefore θ cannot in this case be determined near enough to find a with any great degree of exactness from the equation $a = (b - c) \sec \theta$.

When therefore c is nearly equal to b , and A is not a very small angle, the following method will give a with more exactness than the last Article does.

$$\begin{aligned} a^2 &= b^2 + c^2 - 2bc \cos A = b^2 + c^2 - 2bc \{2 \cos^2 \frac{1}{2} A - 1\} \dots \dots 39, (2) \\ &= (b^2 + c^2 + 2bc) - 4bc \cos^2 \frac{1}{2} A = (b + c)^2 \left\{ 1 - \left(\frac{2\sqrt{bc}}{b+c} \cdot \cos \frac{1}{2} A \right)^2 \right\}. \end{aligned}$$

Now $(\sqrt{b} - \sqrt{c})^2$, or $b - 2\sqrt{bc} + c$, being a square, is necessarily a positive quantity; therefore the positive part of it is greater than the negative part, or $b + c > 2\sqrt{bc}$; $\therefore \frac{2\sqrt{bc}}{b+c}$ is fractional, and *a fortiori* $\frac{2\sqrt{bc}}{b+c} \cdot \cos \frac{1}{2} A$ is fractional.

$$\text{Let therefore } \phi \text{ be the angle whose sine} = \frac{2\sqrt{bc}}{b+c} \cdot \cos \frac{1}{2} A \dots \dots \dots (1).$$

$$\text{Then, } a^2 = (b + c)^2 (1 - \sin^2 \phi), \text{ and } \therefore a = (b + c) \cos \phi \dots \dots \dots (2).$$

$$\begin{aligned} \text{From (1), } L \sin \phi &= \text{I}_{10} 2 + \frac{1}{2} \text{I}_{10} b + \frac{1}{2} \text{I}_{10} c - \text{I}_{10} (b + c) + L \cos \frac{1}{2} A, \\ &= \frac{1}{2} (\text{I}_{10} b + \text{I}_{10} c) + \text{I}_{10} 2 + L \cos \frac{1}{2} A - \text{I}_{10} (b + c). \end{aligned}$$

$$\dots (2), \quad \text{I}_{10} a = \text{I}_{10} (b + c) + L \cos \phi - 10;$$

which give ϕ and a respectively.

86. *Let two sides be given and an angle opposite to one of them.* (a, b, A.)

It has been shewn in (72) that with these data the solution is ambiguous unless a be greater than b . But if a be greater than b , then

$$\sin B = \frac{b}{a} \cdot \sin A, \text{ where } B \text{ is an angle less than } 90^\circ.$$

$$\text{Also, } C = 180^\circ - (A + B); \text{ and, } c = a \cdot \frac{\sin C}{\sin A}.$$

87. *Let all the sides be given.* (a, b, c.)

$$\text{Now, } \sin A = \frac{2}{bc} \sqrt{\{S(S-a)(S-b)(S-c)\}},$$

$$\sin \frac{1}{2} A = \sqrt{\frac{(S-b)(S-c)}{bc}},$$

$$\cos \frac{1}{2} A = \sqrt{\frac{S(S-a)}{bc}},$$

$$\tan \frac{1}{2} A = \sqrt{\frac{(S-b)(S-c)}{S(S-a)}}.*$$

* When the three sides of a triangle are given, the angle A is more easily found in practice by dividing the triangle into two right-angled triangles in the following manner, than from the formulæ in the text.

If a perpendicular CD be let fall from the vertex on the base or the base produced, it may be shewn from the results arrived at in Euclid II. 12, 13, that

Base

: Sum of the other two sides

:: Difference of the sides

: Difference, or Sum, of the segments of the base;

the fourth term of the proportion being the Difference of the segments of the base, or their Sum, according as the perpendicular cuts the base or the base produced.


And the fourth term of this proportion being found, $AD (= AB \pm BD)$ is known, and thence $\cos A \left(= \frac{AD}{AC} \right)$ may be determined.

88. **Obs.** If A be nearly 90° , the first formula of (87) will not give the value of A very exactly, because the increment of $\sin A$ does not in that case vary as the increment of A , and it is also very small; App. 111. 11. In this case any one of the last three forms may be used, and the second or the third form must be taken according as $\cos \frac{1}{2}A$ or $\sin \frac{1}{2}A$ is the greater, i. e. as $\frac{1}{2}A$ is less or greater than 45° , (63). The fourth form is applicable in all cases except where $\frac{1}{2}A$ is nearly 90° , (62).

89. **EXAMPLES.** 1. If BC be a perpendicular object standing on a horizontal plane, its height may be determined by measuring in that plane a line AC , which is called a *base*, and observing the angle BAC with a proper instrument.

For $BC = AC \cdot \tan BAC$;


$$\therefore l_{10} BC = l_{10} AC + L \tan BAC - 10.$$



2. If it be not practicable to come to the foot of the object, let a base DA be measured, such that the points D, A, C may be in the same straight line; and let the angles BDA, BAC be observed.

Here two angles and a side of the $\triangle BAD$ are known. By first determining the side BA the height BC can be found from the right-angled triangle BAC .

$$\text{Thus, } \frac{BA}{AD} = \frac{\sin BDA}{\sin DBA} = \frac{\sin BDA}{\sin (BAC - BDA)}.$$




$$\text{And } BC = BA \cdot \sin BAC = AD \cdot \frac{\sin BDA \sin BAC}{\sin (BAC - BDA)};$$

$$\therefore l_{10} BC = l_{10} AD + L \sin BDA + L \sin BAC - L \sin (BAC - BDA) - 10.$$

3. If D be not in the line AC , the height BC can still be determined.

At A let the angles BAC and BAD be observed, and at D the angle BDA .

Then in the $\triangle BDA$, the angles BDA, BAD and the side AD are given. If then BA be determined from these data, BC can be found from the right-angled triangle BAC .



$$\text{Thus, } \frac{BA}{AD} = \frac{\sin BDA}{\sin DBA} = \frac{\sin BDA}{\sin \{180^\circ - (BDA + BAD)\}};$$

$$\therefore BA = AD \cdot \frac{\sin BDA}{\sin(BDA + BAD)}$$

$$\text{And } BC = BA \cdot \sin BAC = AD \cdot \frac{\sin BDA \sin BAC}{\sin(BDA + BAD)}$$

$$]_{10} BC =]_{10} AD + L \sin BDA + L \sin BAC - L \sin(BDA + BAD) - 10.$$

[It is evident that this determination of BC is not affected by the circumstance of D lying out of the horizontal plane which passes through A and C . Hence it follows that if a straight base AD be measured in any direction from A , and the angles BAC , BAD , BDA be observed, these data will be sufficient for finding the height of B above the horizontal plane passing through A and C .]

4. Required to find the breadth of a river AD , from observations made from the top of a tower BC of which the height is known. (Figure to Ex. 2.)

At B let there be observed the angles of depression of the points D and A below a horizontal line passing through B and parallel to CD . The angles BDC and BAC are equal to these angles of depression; and

$$\begin{aligned} DA &= BA \cdot \frac{\sin DBA}{\sin BDA} = BA \cdot \frac{\sin(BAC - BDA)}{\sin BDA} \\ &= \frac{BC}{\sin BAC} \cdot \frac{\sin(BAC - BDA)}{\sin BDA}. \end{aligned}$$

5. Required the error in height arising from a small given error in an observation of the angle in Example 1.

Let $BC = h$, $AC' = a$, $\angle BAC = A$.

Let k be the error in height produced by an error α of the observed angle.

$$\text{Then } h = a \cdot \tan A, \quad h + k = a \cdot \tan(A + \alpha);$$

$$\therefore k = a \cdot \{\tan(A + \alpha) - \tan A\} = a \cdot \left\{ \frac{\sin(A + \alpha)}{\cos(A + \alpha)} - \frac{\sin A}{\cos A} \right\}$$

$$= a \cdot \frac{\sin(A + \alpha) \cos A - \cos(A + \alpha) \sin A}{\cos(A + \alpha) \cos A} = a \cdot \frac{\sin\{(A + \alpha) - A\}}{\cos(A + \alpha) \cos A}$$

$$= a \cdot \frac{\sin \alpha}{\cos^2 A}; \text{ since } \cos A = \cos(A + \alpha) \text{ nearly, when } \alpha \text{ is very small, (60).}$$

COR. Hence it can be determined when the error in height, arising from a small given error in the observed angle, will be the least.


$$\text{For } k = a \cdot \frac{\sin \alpha}{\cos^2 A} = \frac{h}{\tan A} \cdot \frac{\sin \alpha}{\cos^2 A} = \frac{h \sin \alpha}{\sin A \cos A} = \frac{2h \sin \alpha}{2 \sin A \cos A} = \frac{2h \sin \alpha}{\sin 2A}.$$

Now h being constant, and α being given, this expression, (which is the error in altitude,) will be the least when $\sin 2A$ is the greatest; that is, when $2A = 90^\circ$ or A is 45° .

The observer, therefore, ought to move along the base until $\angle BAC = 45^\circ$, and then by measuring AC he will determine CB (which is in this case equal to AC) with the least chance of error.

90. To find the Area of a triangle, the sides being given.

Let ABC be the triangle, and from C draw CD perpendicular to AB , or to AB produced either way.



Then Area of the triangle ABC , being half of the rectangle on the same base and between the same parallels,

$$= \frac{1}{2} AB \cdot CD = \frac{1}{2} AB \cdot AC \cdot \sin CAB$$

$$= \frac{1}{2} cb \cdot \frac{2}{bc} \sqrt{S(S-a)(S-b)(S-c)}, \text{ by (76).}$$

$$= \sqrt{S(S-a)(S-b)(S-c)}.$$

91. The Area of the triangle also $= \frac{1}{2} a^2 \cdot \frac{\sin B \sin C}{\sin(B+C)}$.

$$\text{For Area} = \frac{1}{2} AB \cdot CD = \frac{1}{2} AB \cdot BC \cdot \sin B$$

$$= \frac{1}{2} BC \cdot \frac{\sin C}{\sin A} \cdot BC \cdot \sin B = \frac{1}{2} a^2 \cdot \frac{\sin B \sin C}{(\sin B + C)};$$


$$\text{since } \sin A = \sin \{180^\circ - (B+C)\} = (\sin B + C).$$

92. To find the Radii of the circles described within and about a regular polygon of any number of sides.

Let AB be a side of a regular polygon of n sides.

Since the polygon is regular, it may have a circle described in and about it; and each of the sides will subtend the same angle at the common center C of these circles.

Draw CD perpendicular to AB . Then $AD = DB$, and CD is the radius of the inscribed circle. Let $r = CD$, and $R = CA$.



Now the sum of all the angles which the sides subtend at $C = n \times \angle ACB = 360^\circ$;


$$\therefore \angle ACB = \frac{360^\circ}{n}; \quad \therefore \angle ACD = \frac{1}{2} \angle ACB = \frac{180^\circ}{n}; \quad \text{and } \frac{AD}{CD} = \tan ACD;$$

$$\therefore r, = CD, = AD \cdot \cot ACD = \frac{1}{2} AB \cdot \cot \frac{180^\circ}{n} \dots\dots\dots (1).$$

$$\text{Again, } R, = AC, = \frac{AD}{\sin ACD} = \frac{\frac{1}{2} AB}{\sin \frac{180^\circ}{n}} = \frac{1}{2} AB \cdot \operatorname{cosec} \frac{180^\circ}{n} \dots\dots\dots (2).$$

93. To find the Area of a regular polygon of any number of sides which is described within or about a circle of given radius.

Let AEB be an arc of the circle whose center is C ; AB a side of the inscribed regular polygon of n sides; CE at right angles to AB , and therefore bisecting it; FG a tangent through E , meeting CA and CB in F and G ; then $F'G'$ is a side of the circumscribed regular polygon of n sides. Let $CA = r$.



$$AB = 2AD = 2r \sin ACD = 2r \sin \frac{360^\circ}{2n} = 2r \sin \frac{180^\circ}{n}.$$

$$FG = 2FE = 2CE \cdot \tan ECF = 2r \tan \frac{180^\circ}{n}.$$

A , = area of inscribed polygon, = $n \cdot \Delta CAB$

$$= n \cdot \frac{CA \cdot CB \cdot \sin ACB}{2} = \frac{n}{2} \cdot r^2 \cdot \sin \frac{360^\circ}{n}.$$

A' , = area of circumscribed polygon, = $n \cdot \Delta CFG$

$$= n \cdot CE \cdot FE = nr^2 \tan \frac{180^\circ}{n}.$$

CON. These areas may be thus compared ;

$$\begin{aligned} \text{Area of inscribed polygon} &= \frac{\Delta CAD}{\Delta CFE} = \frac{CD^2}{CE^2}, \text{ since the } \Delta s \text{ are similar;} \\ \text{Area of circumscribed polygon} &= \left(\frac{CD}{CA}\right)^2 = \cos^2 \frac{180^\circ}{n}. \end{aligned}$$

94. To find the Area of a regular polygon of n sides in terms of a side of the polygon. (Fig. Art. 92.)

AB being a side of the polygon,

$$\begin{aligned} \text{Area} &= n \cdot \Delta CAB = n \cdot \frac{1}{2} AB \cdot DC = \frac{1}{2} n \cdot AB \cdot AD \cdot \cot ACD \\ &= \frac{1}{2} n \cdot AB \cdot \frac{1}{2} AB \cdot \cot \frac{180^\circ}{n} = \frac{1}{4} n (AB)^2 \cot \frac{180^\circ}{n}. \end{aligned}$$


95. To find the Radii of the circles described in and about a triangle of which the sides are given.

Let the lines bisecting the angles A and B meet in O , and from O draw OD, OE, OF perpendiculars to the sides.

Then, Euclid iv. 4, O is the center of the inscribed circle; and its radius $r = OD = OE = OF$.

Now, Area of $\Delta ABC = \Delta AOB + \Delta BOC + \Delta COA$;

$$\begin{aligned} \therefore \sqrt{\{S(S-a)(S-b)(S-c)\}} &= r \cdot \frac{1}{2} c + r \cdot \frac{1}{2} a + r \cdot \frac{1}{2} b = rS; \\ \therefore r &= \frac{\sqrt{\{S(S-a)(S-b)(S-c)\}}}{S} \dots\dots\dots (1). \end{aligned}$$




Again; bisect the sides of the triangle in D, E, F , and draw perpendiculars from these points which will meet in a point O which is the center of the circumscribed circle; R , its radius, $= OA = OB = OC$, Euclid iv. 5.

And $\therefore A = \frac{1}{2} BOC$, Euclid 111. 20;

$$\therefore \sin A = \sin \frac{1}{2} BOC = \sin BOE = \frac{BE}{BO} = \frac{\frac{1}{2} a}{R};$$

$$\therefore, \text{ by (76), } \frac{2}{bc} \sqrt{\{S(S-a)(S-b)(S-c)\}} = \frac{1}{R};$$

$$\therefore R = \frac{abc}{4 \sqrt{\{S(S-a)(S-b)(S-c)\}}} \dots\dots\dots (2).$$



96. To find the Area of a quadrilateral figure of which the opposite angles are supplements to each other.

Let $ABCD$ be the quadrilateral.

Let $AB = a$, $BC = b$, $CD = c$, $DA = d$; Join AC .

Then Area $ABCD = \triangle ABC + \triangle ADC$

$$= \frac{1}{2} ab \sin B + \frac{1}{2} cd \sin D$$

$$= \frac{1}{2} (ab + cd) \sin B,$$

since $\sin D = \sin(180^\circ - D) = \sin B$.

Now from $\triangle ABC$, $a^2 + b^2 - AC^2 = 2ab \cos B$;

and from $\triangle ACD$, $c^2 + d^2 - AC^2 = 2cd \cos D = -2cd \cos B$;

\therefore , by subtracting, $a^2 + b^2 - c^2 - d^2 - 2(ab + cd) \cos B$;

$$\therefore \sin^2 B = 1 - \cos^2 B = 1 - \left\{ \frac{a^2 + b^2 - c^2 - d^2}{2(ab + cd)} \right\}^2$$

$$= \frac{4(ab + cd)^2 - (a^2 + b^2 - c^2 - d^2)^2}{4(ab + cd)^2}.$$

And (area $ABCD$)² = $\frac{1}{4} (ab + cd)^2 \sin^2 B$;

$$\therefore = \frac{1}{16} \{4(ab + cd)^2 - (a^2 + b^2 - c^2 - d^2)^2\}$$

$$= \frac{1}{16} \{2(ab + cd) + (a^2 + b^2 - c^2 - d^2)\} \cdot \{2(ab + cd) - (a^2 + b^2 - c^2 - d^2)\}$$

$$= \frac{1}{16} \{(a + b)^2 - (c - d)^2\} \{(c + d)^2 - (a - b)^2\}$$

$$= \frac{1}{16} (a + b + c - d)(a + b + d - c)(c + d + a - b)(c + d + b - a),$$

and if $S = \frac{1}{2}(a + b + c + d)$,


$$\text{Area } ABCD = \sqrt{\{(S - a)(S - b)(S - c)(S - d)\}}.$$

97. The determination of the unknown quantity in an equation may often be facilitated by breaking the equation up into two others by means of the Goniometrical Ratios. This artifice has been employed in Arts. 84, 85, and the following are additional examples of its use.

EXAMPLES. (1) Having given a , α , and δ , to determine l in a form adapted to logarithmic calculation from the equation,

$$\sin a = \cos l \cos \delta \cos \alpha + \sin l \sin \delta.$$

(Hymers' Astronomy Art. 154)



The equation may be thus written,

$$\sin a = \sin \delta \left(\sin l + \cos l \cdot \frac{\cos \delta \cos \alpha}{\sin \delta} \right).$$

Now since there are tangents of every magnitude and sign, there is *some* angle ϕ , such that

$$\tan \phi, \text{ or } \frac{\sin \phi}{\cos \phi}, = \frac{\cos \delta \cos \alpha}{\sin \delta} = \cot \delta \cos \alpha \dots \dots \dots (1);$$

$$\therefore \sin a = \sin \delta \left(\sin l + \cos l \cdot \frac{\sin \phi}{\cos \phi} \right)$$

$$= \frac{\sin \delta}{\cos \phi} (\sin l \cos \phi + \cos l \sin \phi) = \frac{\sin \delta}{\cos \phi} \cdot \sin (l + \phi),$$

$$\therefore \sin (l + \phi) = \frac{\sin a \cos \phi}{\sin \delta} \dots \dots \dots (2).$$

From, (1), $L \tan \phi = L \cot \delta + L \cos \alpha - 10$, which gives ϕ .

$$(2), \quad L \sin (l + \phi) = L \sin a + L \cos \phi - L \sin \delta,$$

which gives $l + \phi$, and thence l .

(2) To express $a \cdot \cos \theta + b \cdot \cos (\theta + \alpha)$ under the form

$$A \cdot \cos (B + \theta).$$

$$\text{Let } \rho = a \cos \theta + b \cos (\theta + \alpha)$$

$$= a \cos \theta + b \cos \alpha \cos \theta - b \sin \alpha \sin \theta$$

$$= a \left(1 + \frac{b}{a} \cdot \cos \alpha \right) \cos \theta - b \sin \alpha \sin \theta.$$

Let ϕ be the angle whose tangent is $\frac{b}{a} \cdot \cos \alpha$; or, $\tan \phi = \frac{b}{a} \cdot \cos \alpha \dots \dots \dots (1).$

$$\therefore 1 + \frac{b}{a} \cdot \cos \alpha = 1 + \tan \phi = \frac{\cos \phi + \sin \phi}{\cos \phi} = \sqrt{2} \cdot \frac{\frac{1}{\sqrt{2}} \cdot \cos \phi + \frac{1}{\sqrt{2}} \cdot \sin \phi}{\cos \phi}$$

$$= \sqrt{2} \cdot \frac{\sin 45^\circ \cos \phi + \cos 45^\circ \sin \phi}{\cos \phi} = \sqrt{2} \cdot \frac{\sin (45^\circ + \phi)}{\cos \phi};$$

$$\therefore \rho = a \sqrt{2} \cdot \frac{\sin (45^\circ + \phi)}{\cos \phi} \cdot \left\{ \cos \theta - \frac{b \sin \alpha \cos \phi}{a \sqrt{2} \sin (45^\circ + \phi)} \sin \theta \right\}.$$

$$\text{Let } \tan \phi' = \frac{b \sin \alpha \cos \phi}{a \sqrt{2} \sin (45^\circ + \phi)} \dots \dots \dots (2).$$

$$\text{Then } \rho = a \sqrt{2} \cdot \frac{\sin (45^\circ + \phi)}{\cos \phi \cos \phi'} \cdot \cos (\phi' + \theta) \dots \dots \dots (3)$$

a result of the required form.

(3) Adapt $\sqrt{\frac{a-b}{a+b}} + \sqrt{\frac{a+b}{a-b}}$ to logarithmic computation.

$$\sqrt{\frac{a-b}{a+b}} + \sqrt{\frac{a+b}{a-b}} = \frac{a-b+a+b}{\sqrt{(a^2-b^2)}} = \frac{2a}{\sqrt{(a^2-b^2)}} = \frac{2}{\sqrt{\left(1-\frac{b^2}{a^2}\right)}}$$

Now $\frac{b^2}{a^2}$ is necessarily a positive quantity, and it may be of any magnitude; let therefore θ be an angle such that $\text{Tan}^2 \theta = \frac{b^2}{a^2}$ (1).

$$\begin{aligned} \therefore \frac{2}{\sqrt{\left(1-\frac{b^2}{a^2}\right)}} &= \frac{2}{\sqrt{\left\{1-\left(\frac{\sin \theta}{\cos \theta}\right)^2\right\}}} = \frac{2 \cos \theta}{\sqrt{\cos^2 \theta - \sin^2 \theta}} \\ &= \frac{2 \cos \theta}{\cos^{\frac{1}{2}} 2\theta}; \text{ an expression of the required form.....(2).} \end{aligned}$$


98. DEF. The angles introduced to assist the solution of an equation, by breaking it up into other two equations or more, are called *Subsidiary Angles*.

CHAPTER V.

ANALYTICAL TRIGONOMETRY.

99. *The Circumference of a Circle varies as its Radius.*

Let P, p be the perimeters of two regular polygons of n sides each, which are inscribed in two circles whose radii are R, r and circumferences C, c respectively.



$$\text{Let } X = C - P, \quad x = c - p.$$

Now when n increases, P and p increase; therefore X and x are variable quantities dependent on the value of n .

Let AB, ab be sides of the polygons; O, o the centers of the circles.

Then $\angle AOB = \frac{360^\circ}{n} = \angle aob$; and AOB, aob are similar triangles:

$$\therefore \frac{R}{r} = \frac{AB}{ab} = \frac{n \times AB}{n \times ab} = \frac{P}{p} = \frac{C - X}{c - x};$$

$$\therefore R \times c - R \times x = r \times C - r \times X.$$

Now since $R \times c, r \times C$ are constant, and $R \times x, r \times X$ are variable quantities, (by *Francœur's Pure Mathematics*, Art. 167,)

$$R \times c = r \times C, \quad \text{and } R \times x = r \times X;$$

$$\therefore \frac{c}{C} = \frac{r}{R}, \quad \text{or } c \propto r.$$

COR. 1. Since $c \propto r$; $\therefore \frac{c}{r}$, (or $\frac{\text{the circumference of a circle}}{\text{radius}}$), is a constant quantity.


This constant quantity is always represented by 2π ; the approximate value of π , (3.14159...), will be determined hereafter.

COR. 2. Since $\frac{c}{r} = 2\pi$; $\therefore c = 2\pi r$; or $2\pi r$ represents the circumference of a circle whose radius is r .

100. **THE CIRCULAR MEASURE OF AN ANGLE.** *If an arc be traced out by a point in the line CB, by the revolution of which from the position CA an angle ACB is described, the angle ACB may be properly measured by the ratio*

$$\frac{\text{arc AB}}{\text{radius AC}}$$

Since in equal circles (and therefore in the same circle) angles at the center have the same ratio to each other as the arcs on which they stand, Euclid vi. 33,



$$\begin{aligned} \therefore \frac{\angle ACB}{4 \text{ right angles}} &= \frac{\text{arc AB}}{\text{whole circumference ABDA}} \\ &= \frac{AB}{2\pi CA}, \quad (99, \text{Cor. 2}); \\ \therefore \angle ACB &= \frac{4 \text{ right angles}}{2\pi} \cdot \frac{AB}{CA}, \end{aligned}$$

Now, since $\frac{4 \text{ right angles}}{2\pi}$ is a constant quantity, $\frac{AB}{CA}$ increases or decreases in the same ratio as the angle ACB increases or decreases, and therefore $\frac{\text{arc}}{\text{radius}}$ is a proper measure of the magnitude of an angle.

Wherefore an angle may be said to be equal to $\frac{\text{the subtending arc}}{\text{radius}}$.

COR. 1. If $\theta = \frac{\text{arc}}{\text{radius}}$, then $\text{Arc} = r\theta$.

COR. 2. Since the circular measure of an angle (arc \div radius) becomes unity when the arc is equal to the radius, the angle which is subtended by an arc that is equal to the radius is the *Unit of the Circular measurement of angles**.

* The theory of Angular Units may perhaps be rendered more intelligible in the following manner.

From Art. 100, it appears that $\text{Angle} \propto \frac{\text{arc}}{\text{radius}}$; wherefore $\text{Angle} = c \cdot \frac{\text{arc}}{\text{radius}}$, where c is a constant quantity, whose value may be determined if a particular angle be taken for the unit of measurement; and, conversely, if a particular value be assigned to the constant quantity c , the magnitude of that angle may be determined which, in consequence of such assumption, will become the unit of measurement.

For the purposes of analytical calculations it is convenient that the above equation should be of the simplest form possible, and this will be the case if c be taken = 1; the equation then becomes $\text{Angle} = \text{arc} \div \text{radius}$.

Now to determine the angular unit implied in the assumption $c = 1$, make in this last equation, the $\text{Angle} = 1$, the arc then which subtends this angular unit is equal to the radius. In the case therefore, that any angle is represented by the ratio $\text{arc} \div \text{radius}$, the unit of measure is that angle which is subtended by an arc which is equal to the radius with which it is described.

Again, if c have any particular value (as a) assigned to it, then the angle that is the unit of measurement, is such that its $\text{arc} = \frac{1}{a} \cdot \text{radius}$; for in that case the general equation becomes, $\text{Angle} = a \times \frac{1}{a} = 1$.

Example. The unit of measurement being the fourth of a right angle, find the relation between any angle, its arc, and the radius.

$$\text{Generally } \text{Angle} = c \cdot \frac{\text{arc}}{\text{radius}}.$$

In this case, when the angle taken is the unit of measurement,

$$\begin{aligned} \frac{\text{the subtending arc}}{\text{radius}} &= \frac{\text{subtending arc of } \frac{1}{4} \text{ of a right angle}}{\text{radius}} \\ &= \frac{\frac{1}{8} \text{ arc subtending two right angles}}{\text{radius}} = \frac{1}{8} \pi. \end{aligned}$$

$$\text{Wherefore } 1 = c \cdot \frac{1}{8} \pi, \text{ and } \therefore c = \frac{8}{\pi},$$

\therefore the general equation would become, on the supposition that $\frac{1}{4}$ of a right angle is the Angular Unit, $\text{Angle} = \frac{8}{\pi} \cdot \frac{\text{arc}}{\text{radius}}$.

101. In the preceding Chapters the magnitude of an angle has been measured by the number of times it contains a fixed and definite angle (which is the ninetieth part of a right angle, and is called a degree), and its subdivisions; for several *analytical* investigations, however, the circular measure $\frac{\text{arc}}{\text{radius}}$ is much more convenient. The circular measures of angles will, generally speaking, be denoted by the letters of the Greek Alphabet.

102. *Having given the circular measure $\left(\frac{\text{arc}}{\text{radius}}\right)$ of an angle, to determine how many degrees the angle contains; and conversely.*

Let θ be the circular measure of the angle which contains A degrees.

Since $\frac{\text{circumference}}{\text{radius}} = 2\pi$, and the circumference subtends four right angles, therefore 2π is the circular measure of four right angles.

$$\text{Now } \frac{A^\circ}{360^\circ} = \frac{\text{given angle}}{4 \text{ right angles}} = \frac{\theta}{2\pi};$$

$$\therefore \theta = 2\pi \cdot \frac{A}{360} = \frac{\pi}{180} \cdot A \dots \dots \dots (1)$$

$$= 0.017453 \dots \times A.$$


$$\text{And } A = 360 \cdot \frac{\theta}{2\pi} = \frac{180}{\pi} \cdot \theta \dots \dots \dots (2)$$

$$= 57.29577 \dots \times \theta.$$

Ex. 1. If $A = 60$,

$$\theta = \frac{60}{180} \cdot \pi = \frac{1}{3} \times 3.14159 \dots = 1.04719 \dots$$

Let θ be an angle less than a right angle, and $= \angle BAC, = \angle CAB'$. From any point C in AC draw CB, CB' perpendicular to AB and AB' ; then the triangles CAB, CAB' are similar and equal in every respect. With center A and radius AB describe a circle; this will pass through B' , and $BC, B'C$ will touch it at B and B' . Join BB' , cutting AC in N .



Then (*Legendre's Geometry*, iv. 9), arc BB' is $> BNB'$, and $< BC + CB'$.

$$\therefore \frac{\frac{1}{2} \text{arc } BB'}{AB} > \frac{\frac{1}{2} BNB'}{AB} \text{ and is } < \frac{\frac{1}{2} (BC + CB')}{AB},$$

$$\text{or } \theta > \frac{BN}{AB}, \text{ and is } < \frac{BC}{AB};$$

that is, $\theta > \sin \theta$, and is $< \tan \theta$.

AGAIN; since θ lies between $\sin \theta$ and $\tan \theta$,

$$\frac{\theta}{\sin \theta} \text{ lies between } 1 \text{ and } \frac{\tan \theta}{\sin \theta}, \left(\text{i. e. } \frac{1}{\cos \theta} \right).$$

But when θ is diminished, $\cos \theta$ is increased; and when θ is diminished indefinitely and vanishes, $\cos \theta$ becomes equal to unity. Wherefore $\frac{\theta}{\sin \theta}$ becomes ultimately equal to unity, and consequently $\frac{\theta}{\tan \theta}$, which $= \frac{\theta}{\sin \theta} \div \frac{1}{\cos \theta}$, likewise ultimately becomes equal to unity. [*Lefebure de Fourcy*.]

105. If θ be an angle less than a right angle,

$$\sin \theta < \theta - \frac{1}{4} \theta^3.$$

$$\sin \theta = 2 \sin \frac{1}{2} \theta \cdot \cos \frac{1}{2} \theta = \frac{2 \sin \frac{1}{2} \theta}{\cos \frac{1}{2} \theta} \cdot \cos^2 \frac{1}{2} \theta;$$

$$\text{but } \frac{\sin \frac{1}{2} \theta}{\cos \frac{1}{2} \theta}, (= \tan \frac{1}{2} \theta), \text{ is } > \frac{1}{2} \theta. \quad (104).$$

$$\therefore \sin \theta > 2 \left(\frac{1}{2} \theta\right) \cos^2 \frac{1}{2} \theta > \theta \{1 - \sin^2 \frac{1}{2} \theta\}$$

$$\text{a fortiori, } > \theta \left\{1 - \left(\frac{1}{2} \theta\right)^2\right\} > \theta - \frac{1}{4} \theta^3.$$

106. *The Sine of a very small angle is equal to the Circular Measure of the angle, very nearly.*

$$\text{Since, (105), } \theta - \frac{1}{4} \theta^3 < \sin \theta, \therefore \theta - \sin \theta < \frac{1}{4} \theta^3.$$

Now if the angle be very small, θ is a very small fraction and $\frac{1}{4} \theta^3$ a still less quantity, so that the circular measure may be written in the place of the sine in any numerical calculation into which such an angle enters without introducing errors of consequence by the substitution.

COR. If α and β be circular measures of two very small angles,

$$\frac{\sin \alpha}{\sin \beta} = \frac{\alpha}{\beta} = \frac{\text{the number of seconds there is in the } \angle \alpha}{\text{number of seconds in } \angle \beta}.$$

$$\text{Hence } \frac{\sin 2''}{\sin 1''} = \frac{2}{1}, \quad \frac{\sin 3''}{\sin 1''} = \frac{3}{1};$$

And, generally, if n be any small number, *but not otherwise*,

$$\frac{\sin n''}{\sin 1''} = \frac{n}{1};$$

$$\therefore \sin 2'' = 2 \sin 1'', \quad \sin 3'' = 3 \sin 1'', \quad \sin n'' = n \sin 1''.$$

107. *Required the number of seconds contained in the angle of which θ is the Circular Measure.*

Let a be the required number of seconds.

$$\text{Now, in the same circle, } \frac{\text{Arc subtending } a''}{\text{Arc subtending } 1''} = \frac{a}{1},$$

$$\therefore a = \frac{\frac{\text{arc subtending } a''}{\text{radius}}}{\frac{\text{arc subtending } 1''}{\text{radius}}}.$$

But, by hypothesis, $\frac{\text{arc subtending } a''}{\text{radius}} = \theta$; and since $1''$ is a very small angle,

$$\therefore \frac{\text{arc subtending } 1''}{\text{radius}} = \sin 1'', \text{ nearly; (105).}$$

$$\therefore a = \frac{\theta}{\sin 1''};$$

a result which is often of great practical use.

108. *The Area of a Circle of radius r is πr^2 .*

The Area of a Regular Polygon of n sides inscribed in the Circle is n times the area of a triangle two of whose sides are r, r and the included angle is $\frac{2\pi}{n}$;

$$\therefore \text{Area of the Polygon} = n \cdot \frac{r^2}{2} \sin \frac{2\pi}{n} = \pi r^2 \frac{\sin \frac{2\pi}{n}}{\left(\frac{2\pi}{n}\right)}.$$

Now as the number of the sides increases the Area of the Polygon approaches nearer and nearer to that of the Circle, and when n is infinite becomes identical with it; in which case $\frac{2\pi}{n}$ becomes an indefinitely small angle, and therefore $\sin \frac{2\pi}{n} \div \frac{2\pi}{n}$ becomes 1. (104.)

$$\therefore \text{Area of the Circle} = \pi r^2.$$

109. DEMOIVRE'S THEOREM. *To shew that for any value of m,*

$$(\cos \theta \pm \sqrt{-1} \sin \theta)^m = \cos m\theta \pm \sqrt{-1} \sin m\theta;$$

the upper signs being taken together, or the lower together.

$$\begin{aligned} (\cos \theta \pm \sqrt{-1} \sin \theta) (\cos \theta \pm \sqrt{-1} \sin \theta) \\ = \cos^2 \theta - \sin^2 \theta \pm \sqrt{-1} 2 \sin \theta \cos \theta. \end{aligned}$$

$$\text{Or, } (\cos \theta \pm \sqrt{-1} \sin \theta)^2 = \cos 2\theta \pm \sqrt{-1} \sin 2\theta.$$

$$\begin{aligned} \text{Again, } (\cos \theta \pm \sqrt{-1} \sin \theta)^2 (\cos \theta \pm \sqrt{-1} \sin \theta) \\ = (\cos 2\theta \pm \sqrt{-1} \sin 2\theta) (\cos \theta \pm \sqrt{-1} \sin \theta) \\ = (\cos 2\theta \cos \theta - \sin 2\theta \sin \theta) \pm \sqrt{-1} (\sin 2\theta \cos \theta + \cos 2\theta \sin \theta); \end{aligned}$$

$$\therefore (\cos \theta \pm \sqrt{-1} \sin \theta)^3 = \cos 3\theta \pm \sqrt{-1} \sin 3\theta.$$

Suppose this law to hold for m factors, so that

$$(\cos \theta \pm \sqrt{-1} \sin \theta)^m = \cos m\theta \pm \sqrt{-1} \sin m\theta;$$

then $(\cos \theta \pm \sqrt{-1} \sin \theta)^{m+1}$

$$\begin{aligned} &= (\cos m\theta \pm \sqrt{-1} \sin m\theta) (\cos \theta \pm \sqrt{-1} \sin \theta) \\ &= \cos m\theta \cos \theta - \sin m\theta \sin \theta \pm \sqrt{-1} (\sin m\theta \cos \theta + \cos m\theta \sin \theta) \\ &= \cos (m+1)\theta \pm \sqrt{-1} \sin (m+1)\theta. \end{aligned}$$

If therefore the law hold for m factors, it holds for $m+1$ factors; but it has been shewn to hold when $m=3$, it therefore holds when $m=4$, and so by successive inductions it may be proved to be true when the index is any positive integer.

$$\begin{aligned} \text{Again, } (\cos \theta \pm \sqrt{-1} \sin \theta)^{-m} &= \left\{ \frac{1}{\cos \theta \pm \sqrt{-1} \sin \theta} \right\}^m \\ &= \left\{ \frac{\cos^2 \theta + \sin^2 \theta}{\cos \theta \pm \sqrt{-1} \sin \theta} \right\}^m. \end{aligned}$$

$$= (\cos \theta \mp \sqrt{-1} \sin \theta)^m, \text{ by actual division;}$$

$$= \cos m\theta \mp \sqrt{-1} \sin m\theta = \cos (-m\theta) \pm \sqrt{-1} \sin (-m\theta);$$

which proves the theorem for negative integral indices.

$$\text{Again, } (\cos \theta \pm \sqrt{-1} \sin \theta)^m = \cos m\theta \pm \sqrt{-1} \sin m\theta.$$

$$\text{But } \left(\cos \frac{m}{n} \theta \pm \sqrt{-1} \sin \frac{m}{n} \theta \right)^n = \cos m\theta \pm \sqrt{-1} \sin m\theta;$$

$$\therefore (\cos \theta \pm \sqrt{-1} \sin \theta)^m = \left(\cos \frac{m}{n} \theta \pm \sqrt{-1} \sin \frac{m}{n} \theta \right)^n;$$

$$\therefore (\cos \theta \pm \sqrt{-1} \sin \theta)^{\frac{m}{n}} = \cos \frac{m}{n} \theta \pm \sqrt{-1} \sin \frac{m}{n} \theta;$$

which proves the theorem for fractional indices.

COR. By the theorem just proved,

$$(\cos \phi \pm \sqrt{-1} \sin \phi)^m = \cos m\phi \pm \sqrt{-1} \sin m\phi,$$

m being positive or negative, whole or fractional.

Let $\phi = 2p\pi + \theta$, where p is any integer;

$$\left. \begin{aligned} \therefore \cos \phi &= \cos (2p\pi + \theta) = \cos \theta, \\ \sin \phi &= \sin (2p\pi + \theta) = \sin \theta, \end{aligned} \right\} \dots (103).$$

First, let the index of $(\cos \theta \pm \sqrt{-1} \sin \theta)$ be integral, as m ;

$$\text{Then } \cos m\phi = \cos (2mp\pi + m\theta) = \cos m\theta,$$

$$\sin m\phi = \sin (2mp\pi + m\theta) = \sin m\theta.$$

Second, let the index be fractional, as $\frac{m}{n}$; Then p , being an integer, may be represented by $qn + r$, where q may be 0 or any integer, and r may be 0 or any integer less than n ;

$$\begin{aligned} \therefore \cos \frac{m}{n} \phi &= \cos \frac{m}{n} (2p\pi + \theta) = \cos \frac{m}{n} \{ 2(qn + r)\pi + \theta \} \\ &= \cos \{ mq \cdot 2\pi + \frac{m}{n} (2r\pi + \theta) \} = \cos \frac{m}{n} (2r\pi + \theta). \end{aligned}$$

$$\text{Similarly, } \sin \frac{m}{n} \phi = \sin \frac{m}{n} (2r\pi + \theta).$$

From these two cases, therefore, it appears that the theorem might have been thus enunciated;

If the index be an integer,

$$(\cos \theta \pm \sqrt{-1} \sin \theta)^m = \cos m\theta \pm \sqrt{-1} \sin m\theta;$$

If the index be fractional,

$$(\cos \theta \pm \sqrt{-1} \sin \theta)^{\frac{m}{n}} = \cos \frac{m}{n} (2r\pi + \theta) \pm \sqrt{-1} \sin \frac{m}{n} (2r\pi + \theta);$$

where r may be 0, or any integer less than n .

[NOTE. It is to be observed that by giving r all values from 0 to $n - 1$ there will be obtained from the second member of the second equation the n different values of

$$(\cos \theta \pm \sqrt{-1} \sin \theta)^{\frac{m}{n}}.$$

Also these values will recur if r have values given it which are greater than $n - 1$. For taking $r = pn + q$, where p is any integer, and q any integer less than n ,

$$\text{Since } \frac{m}{n} \{2(pn + q)\pi + \theta\} = 2m\pi + \frac{m}{n} (2q\pi + \theta),$$

therefore the cosine and the sine of $\frac{m}{n} \{2(pn + q)\pi + \theta\}$ are respectively equal to the cosine and the sine of $\frac{m}{n} (2q\pi + \theta)$, and the numbers q and $pn + q$ when substituted for r give the same value for $(\cos \theta \pm \sqrt{-1} \sin \theta)^{\frac{m}{n}}$.

110. If $2 \cos \theta$ be represented by $x + \frac{1}{x}$, then $2 \sqrt{-1} \sin \theta$ will be represented by $x - \frac{1}{x}$, $2 \cos m\theta$ by $x^m + \frac{1}{x^m}$, and $2 \sqrt{-1} \sin m\theta$ by $x^m - \frac{1}{x^m}$.

$$\text{For } -\sin^2 \theta = \cos^2 \theta - 1 = \frac{1}{4} \left(x + \frac{1}{x}\right)^2 - 1 = \frac{1}{4} \left(x^2 - 2 + \frac{1}{x^2}\right);$$

$$\therefore 2 \sqrt{-1} \sin \theta = x - \frac{1}{x};$$

And since $2 \cos \theta = x + \frac{1}{x}$, \therefore by addition and subtraction,

$$x = \cos \theta + \sqrt{-1} \sin \theta; \quad \text{and } \frac{1}{x} = \cos \theta - \sqrt{-1} \sin \theta;$$

$$\therefore \begin{cases} x^m = (\cos \theta + \sqrt{-1} \sin \theta)^m = \cos m\theta + \sqrt{-1} \sin m\theta \\ \frac{1}{x^m} = (\cos \theta - \sqrt{-1} \sin \theta)^m = \cos m\theta - \sqrt{-1} \sin m\theta, \end{cases}$$

$$\therefore x^m + \frac{1}{x^m} = 2 \cos m\theta; \quad \text{and } x^m - \frac{1}{x^m} = 2 \sqrt{-1} \sin m\theta.$$

Cor. By taking the equations of the corollary to (109), the following results are obtained;

(1) The index being an integer,

$$2 \cos m\theta = x^m + \frac{1}{x^m}; \quad 2 \sqrt{-1} \sin m\theta = x^m - \frac{1}{x^m};$$

(2) The index being a fraction $\left(\frac{m}{n}\right)$,

$$2 \cos \frac{m}{n} (2r\pi + \theta) = x^{\frac{m}{n}} + \frac{1}{x^{\frac{m}{n}}}; \quad 2 \sqrt{-1} \sin \frac{m}{n} (2r\pi + \theta) = x^{\frac{m}{n}} - \frac{1}{x^{\frac{m}{n}}};$$

where r may be 0 or any integer less than n .

111. To express any positive integral power of the Cosine of an angle in terms of the Cosines of the multiples of the angle.

Let $2 \cos \theta = x + \frac{1}{x}$; $\therefore 2 \cos n\theta = x^n + \frac{1}{x^n} \dots\dots(110)$.

Now

$$(2 \cos \theta)^n = \left(x + \frac{1}{x}\right)^n = x^n + nx^{n-2} + n \cdot \frac{n-1}{2} \cdot x^{n-4} + \dots + n \cdot \frac{n-1}{2} \cdot \frac{1}{x^{n-4}} + n \cdot \frac{1}{x^{n-2}} + \frac{1}{x^n} \quad (1)$$

$$= \left(x^n + \frac{1}{x^n}\right) + n \left(x^{n-2} + \frac{1}{x^{n-2}}\right) + \dots\dots\dots \quad (2)$$

If n be even; the last term of (2), or the $(\frac{1}{2}n + 1)^{th}$ term of (1), is

$$n \cdot \frac{n-1}{2} \cdot \frac{n-2}{3} \dots \frac{n - \frac{1}{2}n + 1}{\frac{1}{2}n}$$

If n be odd; the sum of the two middle terms of (1), viz.

$$\text{the } \left\{\frac{1}{2}(n-1) + 1\right\}^{th} \text{ and the } \left\{\frac{1}{2}(n-1) + 2\right\}^{th},$$

(which have the same coefficient, being terms equidistant from the extremities of the series (1)), is

$$n \cdot \frac{n-1}{2} \dots \frac{n - \frac{1}{2}(n-1) + 1}{\frac{1}{2}(n-1)} \cdot \left(x + \frac{1}{x}\right)$$

Hence (2) becomes

$$(2 \cos \theta)^n = 2 \cos n\theta + n \cdot 2 \cos (n-2)\theta + \dots$$

$$+ \begin{cases} n \cdot \frac{n-1}{2} \dots \frac{n - \frac{1}{2}n + 1}{\frac{1}{2}n}; & \text{when } n \text{ is even,} \\ n \cdot \frac{n-1}{2} \dots \frac{n - \frac{1}{2}(n-1) + 1}{\frac{1}{2}(n-1)} \cdot 2 \cos \theta; & \text{when } n \text{ is odd;} \end{cases}$$

$$\therefore 2^{n-1} \cos^n \theta = \cos n\theta + n \cos (n-2)\theta + n \cdot \frac{n-1}{2} \cdot \cos (n-4)\theta + \dots$$

the last term of the series being

$$+ \frac{1}{2}n \cdot \frac{n-1}{2} \dots \frac{n - \frac{1}{2}n + 1}{\frac{1}{2}n}; \quad n \text{ being even,}$$

$$+ n \cdot \frac{n-1}{2} \dots \frac{n - \frac{1}{2}(n-1) + 1}{\frac{1}{2}(n-1)} \cdot \cos \theta; \quad n \text{ being odd.}$$

- COR.** Thus, when $n = 2$, $2 \cos^2 \theta = \cos 2\theta + 1$,
 $n = 3$, $4 \cos^3 \theta = \cos 3\theta + 3 \cos \theta$,
 $n = 4$, $8 \cos^4 \theta = \cos 4\theta + 4 \cos 2\theta + 3$.

112. To express any positive integral power of the Sine of an angle in terms of the Sines and Cosines of the multiples of the angle.

$$\text{Let } 2\sqrt{-1} \sin \theta = x - \frac{1}{x}; \quad \therefore 2\sqrt{-1} \sin n\theta = x^n - \frac{1}{x^n}.$$

$$\text{And } (2\sqrt{-1} \sin \theta)^n = \left(x - \frac{1}{x}\right)^n = \left\{x + \left(-\frac{1}{x}\right)\right\}^n$$

$$= x^n - nx^{n-2} + n \cdot \frac{n-1}{2} \cdot x^{n-4} - \dots + n \cdot \frac{n-1}{2} \cdot \frac{(-1)^{n-2}}{x^{n-4}} + n \cdot \frac{(-1)^{n-1}}{x^{n-2}} + \frac{(-1)^n}{x^n} \dots (1)$$

$$= \left\{x^n + \left(-\frac{1}{x}\right)^n\right\} - n \cdot \left\{x^{n-2} + \left(-\frac{1}{x}\right)^{n-2}\right\} + n \cdot \frac{n-1}{2} \cdot \left\{x^{n-4} + \left(-\frac{1}{x}\right)^{n-4}\right\} - \dots (2).$$

1. Let n be even;

$$\text{then } (\sqrt{-1})^n = (\sqrt{-1})^{2 \cdot \frac{n}{2}} = \{(\sqrt{-1})^2\}^{\frac{n}{2}} = (-1)^{\frac{n}{2}};$$

and the middle, or $\left(\frac{n}{2} + 1\right)^{\text{th}}$, term of the expansion (1) is

$$(-1)^{\frac{n}{2}} n \cdot \frac{n-1}{2} \dots \dots \dots \frac{n - \frac{1}{2}n + 1}{\frac{1}{2}n}.$$

Wherefore when n is even the series (2) becomes

$$\begin{aligned} (-1)^{\frac{n}{2}} 2^n \sin^n \theta &= \left(x^n + \frac{1}{x^n}\right) - n \left(x^{n-2} + \frac{1}{x^{n-2}}\right) + n \cdot \frac{n-1}{2} \cdot \left(x^{n-4} + \frac{1}{x^{n-4}}\right) - \dots \\ &\quad + (-1)^{\frac{n}{2}} n \cdot \frac{n-1}{2} \dots \dots \frac{n - \frac{1}{2}n + 1}{\frac{1}{2}n} \\ &= 2 \cos n\theta - n 2 \cos (n-2)\theta + n \cdot \frac{n-1}{2} \cdot 2 \cos (n-4)\theta - \dots \\ &\quad \dots \dots + (-1)^{\frac{n}{2}} n \cdot \frac{n-1}{2} \dots \dots \frac{n - \frac{1}{2}n + 1}{\frac{1}{2}n}; \\ \therefore (-1)^{\frac{n}{2}} 2^{n-1} \sin^n \theta &= \cos n\theta - n \cos (n-2)\theta + n \cdot \frac{n-1}{2} \cdot \cos (n-4)\theta - \dots \\ &\quad \dots \dots + (-1)^{\frac{n}{2}} \frac{1}{2} n \cdot \frac{n-1}{2} \dots \dots \frac{n - \frac{1}{2}n + 1}{\frac{1}{2}n}. \end{aligned}$$

2. If n be odd, then $(\sqrt{-1})^n = \sqrt{-1} (\sqrt{-1})^{n-1} = (-1)^{\frac{1}{2}(n-1)} \sqrt{-1}$; and the last binomial in (2) is

$$(-1)^{\frac{1}{2}(n-1)} n \cdot \frac{n-1}{2} \dots \frac{n - \frac{1}{2}(n-1) + 1}{\frac{1}{2}(n-1)} \cdot \left(x - \frac{1}{x}\right).$$

Wherefore, when n is an odd number, the series (2) becomes

$$\begin{aligned} (-1)^{\frac{1}{2}(n-1)} 2^n \sqrt{-1} \sin^n \theta &= \left(x^n - \frac{1}{x^n}\right) - n \left(x^{n-2} - \frac{1}{x^{n-2}}\right) + n \cdot \frac{n-1}{2} \cdot \left(x^{n-4} - \frac{1}{x^{n-4}}\right) - \dots \\ &\dots \dots (-1)^{\frac{1}{2}(n-1)} n \cdot \frac{n-1}{2} \dots \frac{n - \frac{1}{2}(n-1) + 1}{\frac{1}{2}(n-1)} \cdot \left(x - \frac{1}{x}\right), \\ &= 2 \sqrt{-1} \sin n\theta - n 2 \sqrt{-1} \sin(n-2)\theta + \dots \\ &\dots \dots + (-1)^{\frac{1}{2}(n-1)} n \cdot \frac{n-1}{2} \dots \frac{n - \frac{1}{2}(n-1) + 1}{\frac{1}{2}(n-1)} \cdot 2 \sqrt{-1} \sin \theta; \\ \therefore (-1)^{\frac{1}{2}(n-1)} 2^{n-1} \sin^n \theta &= \sin n\theta - n \cdot \sin(n-2)\theta + n \cdot \frac{n-1}{2} \cdot \sin(n-4)\theta + \dots \\ &\dots \dots + (-1)^{\frac{1}{2}(n-1)} n \cdot \frac{n-1}{2} \dots \frac{n - \frac{1}{2}(n-1) + 1}{\frac{1}{2}(n-1)} \cdot \sin \theta.* \end{aligned}$$

113. *Having given Tan θ , to find Tan $n\theta$.*

$$\begin{aligned} \cos n\theta + \sqrt{-1} \sin n\theta &= (\cos \theta + \sqrt{-1} \sin \theta)^n = (\cos \theta)^n (1 + \sqrt{-1} \tan \theta)^n \\ &= (\cos \theta)^n \left\{ 1 + n \sqrt{-1} \tan \theta - n \cdot \frac{n-1}{2} \cdot \tan^2 \theta - n \cdot \frac{n-1}{2} \cdot \frac{n-2}{3} \cdot \sqrt{-1} \tan^3 \theta + \dots \right\} \quad (1). \end{aligned}$$

And equating the possible and impossible parts of this equation,

$$\cos n\theta = \cos^n \theta \left\{ 1 - n \cdot \frac{n-1}{2} \cdot \tan^2 \theta + n \cdot \frac{n-1}{2} \cdot \frac{n-2}{3} \cdot \frac{n-3}{4} \cdot \tan^4 \theta - \dots \right\} \dots (2).$$

$$\sin n\theta = \cos^n \theta \left\{ n \tan \theta - n \cdot \frac{n-1}{2} \cdot \frac{n-2}{3} \cdot \tan^3 \theta + \dots \right\} \dots \dots \dots (3).$$

* $\cos n\theta$ can be expanded in terms of $\cos \theta$ alone, when n is a positive integer, by an algebraical calculation, but the problem is solved more easily by means of the Differential Calculus. [See *Gregory's Examples*.]

$$\therefore \tan n\theta = \frac{\sin n\theta}{\cos n\theta} = \frac{n \tan \theta - n \cdot \frac{n-1}{2} \cdot \frac{n-2}{3} \cdot \tan^3 \theta + \dots}{1 - n \cdot \frac{n-1}{2} \cdot \tan^2 \theta + \dots}$$

Cor. If n be a positive integer, the series (1) will terminate, and the last term is $(\sqrt{-1} \tan \theta)^n$.

1. Let n be even; then

$$(\sqrt{-1} \tan \theta)^n = (\sqrt{-1})^{2 \cdot \frac{n}{2}} \tan^n \theta = (-1)^{\frac{n}{2}} \tan^n \theta;$$

therefore the last term is $(-1)^{\frac{n}{2}} \tan^n \theta$, and

$$\tan n\theta = \frac{n \tan \theta - n \cdot \frac{n-1}{2} \cdot \frac{n-2}{3} \cdot \tan^3 \theta + \dots + (-1)^{\frac{n}{2}-1} n \tan^{n-1} \theta}{1 - n \cdot \frac{n-1}{2} \cdot \tan^2 \theta + \dots + (-1)^{\frac{n}{2}} \tan^n \theta}.$$

2. Let n be odd; then

$$(\sqrt{-1} \tan \theta)^n = \sqrt{-1} (\sqrt{-1})^{n-1} \tan^n \theta = \sqrt{-1} (-1)^{\frac{1}{2}(n-1)} \tan^n \theta;$$

therefore the last term of (3) is $(-1)^{\frac{1}{2}(n-1)} \tan^n \theta$, and

$$\tan n\theta = \frac{n \tan \theta - n \cdot \frac{n-1}{2} \cdot \frac{n-2}{3} \cdot \tan^3 \theta + \dots + (-1)^{\frac{1}{2}(n-1)} \tan^n \theta}{1 - n \cdot \frac{n-1}{2} \cdot \tan^2 \theta + \dots + (-1)^{\frac{1}{2}(n-1)} n \tan^{n-1} \theta}.$$

114. To shew that $\cos \alpha = 1 - \frac{\alpha^2}{1 \cdot 2} + \frac{\alpha^4}{1 \cdot 2 \cdot 3 \cdot 4} - \dots$

and $\sin \alpha = \alpha - \frac{\alpha^3}{1 \cdot 2 \cdot 3} + \frac{\alpha^5}{1 \cdot 2 \cdot 3 \cdot 4 \cdot 5} - \dots$

By 113, (2)

$$\cos n\theta = \cos^n \theta \left\{ 1 - n \cdot \frac{n-1}{2} \cdot \tan^2 \theta + n \cdot \frac{n-1}{2} \cdot \frac{n-2}{3} \cdot \frac{n-3}{4} \cdot \tan^4 \theta - \dots \right\}.$$

* The sign of $n \tan^{n-1} \theta$ is the same as that of $\tan^n \theta$, because every *odd* term in the expansion (1) has the same sign as the term immediately following it; and when n is odd, $n \tan^{n-1} \theta$ is an odd term, and has therefore the same sign as $\tan^n \theta$.

Let $n\theta = \alpha$, and $\therefore n = \frac{\alpha}{\theta}$,

$$\begin{aligned} \therefore \cos \alpha &= \cos^n \theta \left\{ 1 - \frac{\alpha}{\theta} \cdot \frac{\alpha}{2} \cdot \tan^2 \theta + \frac{\alpha}{\theta} \cdot \frac{\alpha-1}{2} \cdot \frac{\alpha-2}{3} \cdot \frac{\alpha-3}{4} \cdot \tan^4 \theta - \dots \right\} \\ &= \cos^n \theta \left\{ 1 - \alpha \cdot \frac{\alpha-\theta}{2} \cdot \left(\frac{\tan \theta}{\theta}\right)^2 + \alpha \cdot \frac{\alpha-\theta}{2} \cdot \frac{\alpha-2\theta}{3} \cdot \frac{\alpha-3\theta}{4} \cdot \left(\frac{\tan \theta}{\theta}\right)^4 - \dots \right\}. \end{aligned}$$

Now this is true whatever be the value that is given to θ .

Let then 0 be written for θ ;

In this case $\frac{\tan \theta}{\theta} = 1$, and $\cos \theta = 1$, Art. 104,

and with regard to the value of $\cos^n \theta$ when $\theta = 0$ and n is infinite,

$$\begin{aligned} \left(\cos \frac{\alpha}{n}\right)^n &= \left(1 - \sin^2 \frac{\alpha}{n}\right)^{\frac{n}{2}} = 1 - \frac{n}{2} \cdot \sin^2 \frac{\alpha}{n} + n \cdot \frac{n-2}{2} \cdot \frac{1}{2^2} \cdot \sin^4 \frac{\alpha}{n} - \dots \\ &= 1 - \frac{\alpha^2}{2n} \cdot \left(\frac{\sin \frac{\alpha}{n}}{\frac{\alpha}{n}}\right)^2 + 1 \cdot \frac{1-\frac{2}{n}}{2} \cdot \frac{\alpha^2}{2^2 n^2} \cdot \left(\frac{\sin \frac{\alpha}{n}}{\frac{\alpha}{n}}\right)^4 - \dots \end{aligned}$$

but the limit of $\sin \frac{\alpha}{n} \div \frac{\alpha}{n}$, when α is constant and n is increased indefinitely, is 1;

Wherefore all the terms of the above series which is the value of $\left(\cos \frac{\alpha}{n}\right)^n$ vanish in the limit, except the first,

$\therefore 1 = \text{limit of } \left(\cos \frac{\alpha}{n}\right)^n$, when α is constant and n is increased indefinitely.

$$\therefore \cos \alpha = 1 - \frac{\alpha^2}{1 \cdot 2} + \frac{\alpha^4}{1 \cdot 2 \cdot 3 \cdot 4} - \dots \tag{1}$$

By a similar substitution in 113, (3),

$$\sin \alpha = \alpha - \frac{\alpha^3}{1 \cdot 2 \cdot 3} + \frac{\alpha^5}{1 \cdot 2 \cdot 3 \cdot 4 \cdot 5} - \dots \tag{2}$$

If α be less than $\frac{1}{2} \pi$ (or a right angle), the series (1) and (2) here obtained are immediately convergent.

NOTE. To arrive at these expressions $\frac{\tan \theta}{\theta}$ has been supposed to become 1 when 0 is written for θ , which is the case only when the angle is referred to the *circular* measure. Hence α , which is equal to $n\theta$, is also referred to that measure in the series (1) and (2).

COR. If α be an angle so small that α^2 and higher powers of α may be neglected when compared with unity, equation (1) becomes

$$\text{Cos } \alpha = 1.$$

If α^2 , α^3 be retained, but higher powers of α be neglected, (1) and (2) give,

$$\text{Sin } \alpha = \alpha - \frac{\alpha^3}{6}; \quad \text{Cos } \alpha = 1 - \frac{\alpha^2}{2};$$

approximations which are often practically very useful.

115. If $\epsilon^{\theta\sqrt{-1}}$ and $\epsilon^{-\theta\sqrt{-1}}$ be expanded in terms of $\theta\sqrt{-1}$ and $-\theta\sqrt{-1}$ in the same manner as ϵ^x is expanded in terms of x in the series,

$$\epsilon^x = 1 + x + \frac{x^2}{1.2} + \frac{x^3}{1.2.3} + \dots \quad (\text{App. I. 18. Cor.})$$

then,

$$(1) \text{ Cos } \theta = \frac{1}{2} \cdot (\epsilon^{\theta\sqrt{-1}} + \epsilon^{-\theta\sqrt{-1}}). \quad (2) \text{ Sin } \theta = \frac{1}{2\sqrt{-1}} \cdot (\epsilon^{\theta\sqrt{-1}} - \epsilon^{-\theta\sqrt{-1}}).$$

$$(3) \text{ Tan } \theta = \frac{1}{\sqrt{-1}} \cdot \frac{\epsilon^{2\theta\sqrt{-1}} - 1}{\epsilon^{2\theta\sqrt{-1}} + 1}.$$

For, $\epsilon^{\theta\sqrt{-1}} = 1 + \theta\sqrt{-1} - \frac{\theta^2}{1.2} + \frac{\theta^3}{1.2.3}\sqrt{-1} + \frac{\theta^4}{1.2.3.4} + \dots$

and $\epsilon^{-\theta\sqrt{-1}} = 1 - \theta\sqrt{-1} - \frac{\theta^2}{1.2} + \frac{\theta^3}{1.2.3}\sqrt{-1} + \frac{\theta^4}{1.2.3.4} - \dots$

$$\therefore \epsilon^{\theta\sqrt{-1}} + \epsilon^{-\theta\sqrt{-1}} = 2 \left(1 - \frac{\theta^2}{1.2} + \frac{\theta^4}{1.2.3.4} - \dots \right) = 2 \text{ cos } \theta; \quad (114),$$

$$\text{and } \epsilon^{\theta\sqrt{-1}} - \epsilon^{-\theta\sqrt{-1}} = 2\sqrt{-1} \left(\theta - \frac{\theta^3}{1.2.3} + \dots \right) = 2\sqrt{-1} \text{ sin } \theta;$$

$$\therefore \text{ Cos } \theta = \frac{1}{2} (\epsilon^{\theta\sqrt{-1}} + \epsilon^{-\theta\sqrt{-1}}) \dots\dots\dots (1).$$

$$\text{And, Sin } \theta = \frac{1}{2\sqrt{-1}} (\epsilon^{\theta\sqrt{-1}} - \epsilon^{-\theta\sqrt{-1}}) \dots\dots\dots (2).$$

$$\text{Also, Tan } \theta = \frac{\text{sin } \theta}{\text{cos } \theta} = \frac{1}{\sqrt{-1}} \cdot \frac{\epsilon^{\theta\sqrt{-1}} - \epsilon^{-\theta\sqrt{-1}}}{\epsilon^{\theta\sqrt{-1}} + \epsilon^{-\theta\sqrt{-1}}};$$

and multiplying the numerator and denominator by $\epsilon^{\theta\sqrt{-1}}$,

$$\text{Tan } \theta = \frac{1}{\sqrt{-1}} \cdot \frac{\epsilon^{2\theta\sqrt{-1}} - 1}{\epsilon^{2\theta\sqrt{-1}} + 1} \dots\dots\dots (3).$$

SERIES FOR DETERMINING THE VALUE OF π .

116. GREGORIE'S SERIES. To prove that

$$\theta = \tan \theta - \frac{1}{3} \tan^3 \theta + \frac{1}{5} \tan^5 \theta - \dots ;$$

where θ lies between $-\frac{1}{2} \pi$ and $+\frac{1}{2} \pi$.

Let $\theta = n\pi + \theta_0$; where θ_0 is any angle between $-\frac{1}{2} \pi$ and $+\frac{1}{2} \pi$, and n any positive or negative integer.

Then $\cos \theta = \cos (n\pi + \theta_0) = (-1)^n \cos \theta_0$.

Now $\epsilon^{\pi \sqrt{-1}} = \cos \pi + \sqrt{-1} \sin \pi = -1$; $\therefore (-1)^n = \epsilon^{n\pi \sqrt{-1}}$;

$\therefore \cos \theta = \epsilon^{n\pi \sqrt{-1}} \cos \theta_0$;

And from (1) and (2) of Art. 115,

$$\epsilon^{\theta \sqrt{-1}} = \cos \theta + \sqrt{-1} \sin \theta = \cos \theta (1 + \sqrt{-1} \tan \theta)$$

$$\frac{\epsilon^{\theta \sqrt{-1}}}{1 + \sqrt{-1} \tan \theta} = \epsilon^{n\pi \sqrt{-1}} \cos \theta_0,$$

$$\therefore 1_\epsilon \epsilon^{\theta \sqrt{-1}} = 1_\epsilon \epsilon^{n\pi \sqrt{-1}} + 1_\epsilon \cos \theta_0 + 1_\epsilon (1 + \sqrt{-1} \tan \theta)$$

$$\therefore \theta \sqrt{-1} = n\pi \sqrt{-1} + 1_\epsilon \cos \theta_0 + \sqrt{-1} \tan \theta + \frac{1}{2} \tan^2 \theta - \frac{1}{2} \sqrt{-1} \tan^3 \theta - \dots$$

and equating the impossible parts of this equation,

$$\theta = n\pi + \tan \theta - \frac{1}{3} \tan^3 \theta + \frac{1}{5} \tan^5 \theta - \dots \tag{1}$$

If θ lie between $-\frac{1}{2} \pi$ and $+\frac{1}{2} \pi$, $n = 0$ and the above becomes

$$\theta = \tan \theta - \frac{1}{3} \tan^3 \theta + \frac{1}{5} \tan^5 \theta - \dots ; \tag{2}$$

which is Gregorie's Series.

And generally n must be an integer so taken that $\theta - n\pi$ shall lie between $-\frac{1}{2} \pi$ and $+\frac{1}{2} \pi$.

The series (1) is convergent only when $\tan \theta$ lies between -1 and $+1$; that is, when $\theta - n\pi$ lies between $-\frac{1}{4} \pi$ and $+\frac{1}{4} \pi$.

COR. Tangent of half a right angle, (or $\tan \frac{1}{4} \pi$), = 1;

$$\therefore \frac{\pi}{4} = 1 - \frac{1}{3} + \frac{1}{5} - \frac{1}{7} + \frac{1}{9} - \frac{1}{11} + \dots = \frac{2}{1.3} + \frac{2}{5.7} + \frac{2}{9.11} + \dots ;$$

$$\therefore \pi = 8. \left(\frac{1}{1.3} + \frac{1}{5.7} + \frac{1}{9.11} + \dots \right),$$

a series not convergent enough to be of much use for calculating the numerical value of π .

117. EULER'S SERIES. *To prove that*

$$\frac{\pi}{4} = \left(\frac{1}{2} - \frac{1}{3 \cdot 2^3} + \frac{1}{5 \cdot 2^5} - \dots \right) + \left(\frac{1}{3} - \frac{1}{3 \cdot 3^3} + \frac{1}{5 \cdot 3^5} - \dots \right).$$

$$\text{By (65), } \tan^{-1} 1 - \tan^{-1} \frac{1}{2} = \tan^{-1} \frac{1 - \frac{1}{2}}{1 + \frac{1}{2}} = \tan^{-1} \frac{1}{3};$$

$$\therefore \tan^{-1} 1 = \tan^{-1} \frac{1}{2} + \tan^{-1} \frac{1}{3},$$

$$\text{and } \tan^{-1} \frac{1}{2} = \frac{1}{2} - \frac{1}{3 \cdot 2^3} + \frac{1}{5 \cdot 2^5} - \dots; \quad \tan^{-1} \frac{1}{3} = \frac{1}{3} - \frac{1}{3 \cdot 3^3} + \frac{1}{5 \cdot 3^5} - \dots; \dots (116).$$

$$\therefore \tan^{-1} 1, \text{ or } \frac{\pi}{4}, = \left(\frac{1}{2} - \frac{1}{3 \cdot 2^3} + \frac{1}{5 \cdot 2^5} - \dots \right) + \left(\frac{1}{3} - \frac{1}{3 \cdot 3^3} + \frac{1}{5 \cdot 3^5} - \dots \right),$$

a series which converges much more rapidly than that deduced in the Corollary to the last Article.

118. MACHIN'S SERIES. *To prove that*

$$\frac{\pi}{4} = 4 \left(\frac{1}{5} - \frac{1}{3 \cdot 5^3} + \frac{1}{5 \cdot 5^5} - \dots \right) - \left(\frac{1}{239} - \frac{1}{3 \cdot (239)^3} + \frac{1}{5 \cdot (239)^5} - \dots \right)$$

$$4 \tan^{-1} \frac{1}{5} = 2 \times 2 \tan^{-1} \frac{1}{5} = 2 \tan^{-1} \frac{\frac{1}{5} + \frac{1}{5}}{1 - \frac{1}{5} \cdot \frac{1}{5}} = 2 \tan^{-1} \frac{5}{12}$$

$$= \tan^{-1} \frac{\frac{10}{12}}{1 - \frac{25}{144}} = \tan^{-1} \frac{120}{119},$$

$$\text{and } \tan^{-1} \frac{120}{119} - \tan^{-1} 1 = \tan^{-1} \frac{\frac{120}{119} - 1}{1 + \frac{120}{119}} = \tan^{-1} \frac{1}{239};$$

$$\text{or } 4 \tan^{-1} \frac{1}{5} - \frac{\pi}{4} = \tan^{-1} \frac{1}{239};$$

$$\therefore \frac{\pi}{4} = 4 \tan^{-1} \frac{1}{5} - \tan^{-1} \frac{1}{239};$$

$$\therefore \frac{\pi}{4} = 4 \left(\frac{1}{5} - \frac{1}{3 \cdot 5^3} + \frac{1}{5 \cdot 5^5} - \dots \right) - \left(\frac{1}{239} - \frac{1}{3 \cdot (239)^3} + \frac{1}{5 \cdot (239)^5} - \dots \right).$$

In this way it is found that $\pi = 3 \cdot 141592653589793 \dots$

Cor. Since $\tan^{-1} \frac{1}{99} + \tan^{-1} \frac{1}{239} = \tan^{-1} \frac{\frac{1}{99} + \frac{1}{239}}{1 - \frac{1}{99} \cdot \frac{1}{239}}$

$$= \tan^{-1} \frac{338}{23660} = \tan^{-1} \frac{1}{70},$$

$$\therefore \frac{\pi}{4} = 4 \tan^{-1} \frac{1}{5} - \tan^{-1} \frac{1}{70} + \tan^{-1} \frac{1}{99};$$

a series by means of which Mr Rutherford (*Phil. Trans.* 1841) succeeded in determining the value of π to 200 places of figures.

MISCELLANEOUS PROPOSITIONS, SHEWING THE APPLICATION OF THE THEOREMS AND FORMULÆ OF THIS CHAPTER.

119. *Having given Tan α , Tan β ,...Tan λ , to find Tan $(\alpha+\beta+\dots+\lambda)$.*

$$\begin{aligned} & (\cos \alpha + \sqrt{-1} \sin \alpha) (\cos \beta + \sqrt{-1} \sin \beta) \\ &= (\cos \alpha \cos \beta - \sin \alpha \sin \beta) + \sqrt{-1} (\cos \alpha \sin \beta + \cos \beta \sin \alpha) \\ &= \cos (\alpha + \beta) + \sqrt{-1} \sin (\alpha + \beta) \dots\dots\dots (1). \end{aligned}$$

Suppose this law to hold for n factors, so that

$$\begin{aligned} & (\cos \alpha + \sqrt{-1} \sin \alpha) (\cos \beta + \sqrt{-1} \sin \beta) \dots (\cos \kappa + \sqrt{-1} \sin \kappa) \\ &= \cos (\alpha + \beta + \dots + \kappa) + \sqrt{-1} \sin (\alpha + \beta + \dots + \kappa); \end{aligned}$$

by introducing another factor,

$$\begin{aligned} & (\cos \alpha + \sqrt{-1} \sin \alpha) \dots (\cos \kappa + \sqrt{-1} \sin \kappa) (\cos \lambda + \sqrt{-1} \sin \lambda) \\ &= \{ \cos (\alpha + \beta + \dots + \kappa) + \sqrt{-1} \sin (\alpha + \beta + \dots + \kappa) \} (\cos \lambda + \sqrt{-1} \sin \lambda) \\ &= \cos (\alpha + \beta + \dots + \lambda) + \sqrt{-1} \sin (\alpha + \beta + \dots + \lambda), \text{ by (1);} \end{aligned}$$

and therefore the law holds for $n+1$ factors. .

But by actual multiplication the law has been shewn to hold for two factors, it must therefore hold for three factors, and thus by successive inductions it is concluded that it holds for any number of factors.

Let S_1 = the sum of the quantities $\text{Tan } \alpha, \text{Tan } \beta, \dots$; S_2 = the sum of the products of every two of them; S_3 = the sum of the products of every three of them; and so on.

$$\begin{aligned} \text{Then, } \quad & \text{Cos } (\alpha + \beta + \dots + \lambda) + \sqrt{-1} \sin (\alpha + \beta + \dots + \lambda) \\ & = (\cos \alpha + \sqrt{-1} \sin \alpha) (\cos \beta + \sqrt{-1} \sin \beta) \dots (\cos \lambda + \sqrt{-1} \sin \lambda) \\ & = \cos \alpha (1 + \sqrt{-1} \tan \alpha) \cos \beta (1 + \sqrt{-1} \tan \beta) \dots \cos \lambda (1 + \sqrt{-1} \tan \lambda) \\ & = \cos \alpha \cos \beta \dots \cos \lambda (1 + \sqrt{-1} S_1 - S_2 - \sqrt{-1} S_3 + S_4 + \dots). \end{aligned}$$

Wood's Algebra, Art. 271.

Hence, by equating the possible and impossible parts of this equation,

$$\begin{aligned} \text{Cos } (\alpha + \beta + \dots + \lambda) &= \cos \alpha \cos \beta \dots \cos \lambda (1 - S_2 + S_4 - S_6 + \dots) \\ \text{Sin } (\alpha + \beta + \dots + \lambda) &= \cos \alpha \cos \beta \dots \cos \lambda (S_1 - S_3 + S_5 - \dots) \\ \therefore \text{Tan } (\alpha + \beta + \dots + \lambda) &= \frac{\sin (\alpha + \beta + \dots + \lambda)}{\cos (\alpha + \beta + \dots + \lambda)} = \frac{S_1 - S_3 + S_5 - \dots}{1 - S_2 + S_4 - \dots}. \end{aligned}$$

If there be n of the angles $\alpha, \beta, \dots, \lambda$, it may be shewn as in (113. Cor.) that

$$\begin{aligned} \text{Tan } (\alpha + \beta + \dots + \lambda) &= \frac{S_1 - S_3 + S_5 - \dots + (-1)^{\frac{n}{2}-1} S_{n-1}}{1 - S_2 + S_4 - \dots + (-1)^{\frac{n}{2}} S_n}, \quad n \text{ even.} \\ &= \frac{S_1 - S_3 + S_5 - \dots + (-1)^{\frac{1}{2}(n-1)} S_n}{1 - S_2 + S_4 - \dots + (-1)^{\frac{1}{2}(n-1)} S_{n-1}}, \quad n \text{ odd.} \end{aligned}$$

120. *If* $\text{Sin } p = \sin P \cdot \sin (z + p)$, *required to expand* p *in terms of* $\text{Sin } P$ *and of the Sines of* z *and its multiples.*

Hymers' Astronomy, Art. 247.

Since $\text{Sin } p = \frac{1}{2\sqrt{-1}} (\epsilon^p \sqrt{-1} - \epsilon^{-p} \sqrt{-1})$ by (115), the equation becomes by substituting such values of $\text{Sin } p$ and $\text{Sin } (z + p)$,

$$\frac{1}{2\sqrt{-1}} (\epsilon^p \sqrt{-1} - \epsilon^{-p} \sqrt{-1}) = \sin P \frac{1}{2\sqrt{-1}} (\epsilon^{(z+p)} \sqrt{-1} - \epsilon^{-(z+p)} \sqrt{-1})$$

and, by multiplying both sides of the equation by $2\sqrt{-1} \cdot \epsilon^p \sqrt{-1}$,

$$\begin{aligned} \epsilon^{2p} \sqrt{-1} - 1 &= \sin P (\epsilon^{(z+2p)} \sqrt{-1} - \epsilon^{-z} \sqrt{-1}) \\ &= \sin P \epsilon^z \sqrt{-1} \cdot \epsilon^{2p} \sqrt{-1} - \sin P \epsilon^{-z} \sqrt{-1}; \end{aligned}$$

$$\therefore \epsilon^{2p} \sqrt{-1} (1 - \sin P \epsilon^{\epsilon \sqrt{-1}}) = 1 - \sin P \epsilon^{\epsilon \sqrt{-1}};$$

$$\therefore \epsilon^{2p} \sqrt{-1} = \frac{1 - \sin P \epsilon^{\epsilon \sqrt{-1}}}{1 - \sin P \epsilon^{\epsilon \sqrt{-1}}};$$

$$\begin{aligned} \therefore 2p \sqrt{-1} &= l_{\epsilon} (1 - \sin P \epsilon^{\epsilon \sqrt{-1}}) - l_{\epsilon} (1 - \sin P \epsilon^{\epsilon \sqrt{-1}}) \\ &= -\sin P \epsilon^{\epsilon \sqrt{-1}} - \frac{1}{2} \sin^2 P \epsilon^{2\epsilon \sqrt{-1}} - \frac{1}{3} \sin^3 P \epsilon^{3\epsilon \sqrt{-1}} - \dots \\ &\quad + \sin P \epsilon^{\epsilon \sqrt{-1}} + \frac{1}{2} \sin^2 P \epsilon^{2\epsilon \sqrt{-1}} + \frac{1}{3} \sin^3 P \epsilon^{3\epsilon \sqrt{-1}} + \dots \end{aligned}$$

Appendix 1. 14. iii.

$$\begin{aligned} \therefore p &= \sin P \frac{1}{2 \sqrt{-1}} (\epsilon^{\epsilon \sqrt{-1}} - \epsilon^{-\epsilon \sqrt{-1}}) + \frac{1}{2} \sin^2 P \frac{1}{2 \sqrt{-1}} (\epsilon^{2\epsilon \sqrt{-1}} - \epsilon^{-2\epsilon \sqrt{-1}}) + \dots \\ &= \sin P \sin \alpha + \frac{1}{2} \sin^2 P \sin 2\alpha + \frac{1}{3} \sin^3 P \sin 3\alpha + \dots \end{aligned}$$

COR. The number of seconds contained in the angle p , which is expressed by the circular measure, is (107)

$$\begin{aligned} \frac{p}{\sin 1''} &= \frac{\sin P}{\sin 1''} \sin \alpha + \frac{1}{2} \frac{\sin^2 P}{\sin 1''} \sin 2\alpha + \frac{1}{3} \frac{\sin^3 P}{\sin 1''} \sin 3\alpha + \dots \\ \text{or} &= \frac{\sin P}{\sin 1''} \sin \alpha + \frac{\sin^2 P}{\sin 2''} \sin 2\alpha + \frac{\sin^3 P}{\sin 3''} \sin 3\alpha + \dots \end{aligned}$$

121. In the same manner if $\tan l' = n \cdot \tan l$, and the tangents be expressed in terms of $l \sqrt{-1}$ and $l' \sqrt{-1}$ by the formula of (115), it may be proved that,

$$l' = l - m \sin 2l + \frac{1}{2} m^2 \sin 4l - \frac{1}{3} m^3 \sin 6l + \dots \text{ where } m = \frac{1 - n}{1 + n}.$$

COR. The number of seconds in l' , is (107)

$$\begin{aligned} \frac{l'}{\sin 1''} &= \frac{l}{\sin 1''} - \frac{m}{\sin 1''} \sin 2l + \frac{1}{2} \frac{m^2}{\sin 1''} \sin 4l - \dots \\ \text{or} &= \frac{l}{\sin 1''} - \frac{m \sin 2l}{\sin 1''} + \frac{m^2 \sin 4l}{\sin 2''} - \dots \end{aligned}$$

[The angle l , which is the observed latitude of a place, is read off in degrees, minutes, and seconds from the instrument by which the observation is made; therefore to find the degrees, minutes, and seconds in l' , the only computation necessary is to determine the number of seconds in the latter part of the series, viz.

$$- \frac{m}{\sin 1''} \sin 2l + \frac{m^2}{\sin 2''} \sin 4l - \dots]$$

122. To expand $\frac{1}{1 - e \cos \theta}$ in a series of the Cosines of θ and its multiples; e being less than 1.

By hypothesis, e is less than 1; and since $(1-b)^2$, or $1-2b+b^2$, being a square, is necessarily positive, $1+b^2$ is greater than $2b$.

$$\text{Let } \therefore c = \frac{2b}{1+b^2}.$$

$$\therefore 1 - e^2 = 1 - \frac{4b^2}{(1+b^2)^2} = \left(\frac{1-b^2}{1+b^2}\right)^2; \quad \therefore \frac{1-b^2}{1+b^2} = \sqrt{1-e^2};$$

$$\text{Whence, } b^2 = \frac{1 - \sqrt{1-e^2}}{1 + \sqrt{1-e^2}} = \frac{1 - \sqrt{1-e^2}}{1 + \sqrt{1-e^2}} \cdot \frac{1 + \sqrt{1-e^2}}{1 + \sqrt{1-e^2}};$$

$$\therefore b = \frac{e}{1 + \sqrt{1-e^2}}.$$

$$\text{Let } 2 \cos \theta = x + \frac{1}{x}.$$

$$\text{Then } \frac{1}{1 - e \cos \theta} = \frac{1}{1 - \frac{b}{1+b^2} \left(x + \frac{1}{x}\right)} = (1+b^2) \cdot \frac{x}{(1-bx)(x-b)}.$$

$$\text{Assume } \frac{x}{(1-bx)(x-b)} = \frac{A}{1-bx} + \frac{B}{x-b};$$

$$\therefore x = A(x-b) + B(1-bx).$$

Whence, by putting x successively equal to $\frac{1}{b}$ and b ,

$$A = \frac{1}{1-b^2}, \quad B = \frac{b}{1-b^2}.$$

Wherefore,

$$\begin{aligned} \frac{1}{1 - e \cos \theta} &= \frac{1+b^2}{1-b^2} \cdot \left(\frac{1}{1-bx} + \frac{b}{x-b} \right) \\ &= \frac{1+b^2}{1-b^2} \cdot \left(\frac{1}{1-bx} + \frac{b}{x} \cdot \frac{1}{1-\frac{b}{x}} \right) \\ &= \frac{1+b^2}{1-b^2} \cdot \left\{ 1 + bx + b^2x^2 + b^3x^3 + \dots \right. \\ &\quad \left. + \frac{b}{x} + \frac{b^2}{x^2} + \frac{b^3}{x^3} + \dots \right\} \end{aligned}$$

$$\begin{aligned} & \sqrt{\frac{1+b^2}{1-b^2}} \{1 + 2b \cos \theta + 2b^2 \cos 2\theta + \dots\} \\ &= \frac{1}{\sqrt{1-e^2}} \{1 + 2b \cos \theta + 2b^2 \cos 2\theta + \dots\}. \end{aligned}$$

Since e is less than 1, b , or $\frac{e}{1+\sqrt{1-e^2}}$, is small, and therefore this series converges rapidly.

COR. The equation to an ellipse is $r = a \frac{1-e^2}{1-e \cos \theta}$, the focus being the pole and θ being measured from that vertex which is the further from the focus;

$$\therefore r = a \sqrt{1-e^2} \{1 + 2b \cos \theta + 2b^2 \cos 2\theta + \dots\}; \quad \text{where } b = \frac{e}{1+\sqrt{1-e^2}}.$$

123. The approximations of 114, Cor. to the values of the sine and the cosine of a very small angle, may often be applied to determine the magnitudes of Astronomical Corrections.

Ex. 1. If $\sin(\omega - y) = \sin \omega \cdot \cos y$, where y is very small, required an approximate value of y .

$$\text{Here, } \sin \omega \cos y - \cos \omega \sin y = \sin \omega \cos u;$$

$$\therefore (1 - \frac{1}{2}y^2) \sin \omega - y \cos \omega = \sin \omega \cos u;$$

$$\begin{aligned} \therefore y \cos \omega + \frac{1}{2}y^2 \sin \omega &= \sin \omega - \sin \omega \cos u \\ &= \sin \omega (1 - \cos u) = 2 \sin \omega \sin^2 \frac{1}{2} u; \end{aligned}$$

$$\therefore y \{1 + \frac{1}{2}y \tan \omega\} = 2 \tan \omega \sin^2 \frac{1}{2} u;$$

$$\begin{aligned} \therefore y &= \frac{2 \tan \omega \sin^2 \frac{1}{2} u}{1 + \frac{1}{2}y \tan \omega} = 2 \tan \omega \sin^2 \frac{1}{2} u \{1 + \frac{1}{2}y \tan \omega\}^{-1} \\ &= 2 \tan \omega \sin^2 \frac{1}{2} u \{1 - \frac{1}{2}y \tan \omega + \dots\}. \end{aligned}$$

By neglecting the second and all the succeeding terms of the expansion, as being small when compared with 1, a first approximation (y_1) to the value of y is obtained,

$$y_1 = 2 \tan \omega \sin^2 \frac{1}{2} u;$$

And by putting for y in the second member of the equation this its first approximate value, a second approximation (y_2) is obtained;

$$\begin{aligned}
 y_2 &= 2 \tan \omega \sin^2 \frac{1}{2} u \{1 - \tan \omega \sin^2 \frac{1}{2} u \tan \omega\} \\
 &= 2 \tan \omega \sin^2 \frac{1}{2} u \{1 - \tan^2 \omega \sin^2 \frac{1}{2} u\}.
 \end{aligned}$$

The number of seconds contained in y_2 , is (107)

$$\frac{y_2}{\sin 1''} = 2 \left(\frac{\tan \omega}{\sin 1''} \sin^2 \frac{1}{2} u - \frac{\tan^3 \omega}{\sin 1''} \sin^4 \frac{1}{2} u \right);$$

and the two terms of the expression having been separately determined by means of logarithms, the number of seconds in y_2 is known.

Hymers' Astronomy, Art. 176.

Ex. 2. In the same manner it may be shewn from

$$\cos(z + y) = \sin n \cdot \sin z \cdot \cos m + \cos z \cdot \cos n,$$

where y and n are very small, that

The number of seconds in y is

$$\frac{y}{\sin 1''} = -\frac{n \cos m}{\sin 1''} + \frac{n^2}{\sin 2''} \cot z \sin^2 m,$$

nearly. The terms of this expression are, as in Ex. 1, determined separately by means of logarithms.

Hymers' Astronomy, Art. 161.

124. The expressions of Arts. 115, 110, have been employed in Arts. 120, 122, to expand certain quantities in the form of a series. The operation can be reversed, as in the following instances.

To find the sum of the series,

$$(1) \quad \sin a + \sin 2a + \sin 3a + \dots \quad + \sin na.$$

$$(2) \quad \cos a + \cos 2a + \cos 3a + \dots \quad + \cos na.$$

$$(1) \quad \text{Let } 2\sqrt{-1} \sin a = x - \frac{1}{x};$$

$$\therefore 2\sqrt{-1} \sin 2a = x^2 - \frac{1}{x^2}; \quad (110).$$

$$2\sqrt{-1} \sin 3a = x^3 - \frac{1}{x^3}$$

.....

$$2\sqrt{-1} \sin na = x^n - \frac{1}{x^n};$$

∴ representing the sum of the series (1) by S ,

$$\begin{aligned}
 2\sqrt{-1} S &= x + x^2 + x^3 + \dots + x^n - \frac{1}{x} - \frac{1}{x^2} - \frac{1}{x^3} - \dots - \frac{1}{x^n} \\
 &= x \cdot \frac{x^n - 1}{x - 1} - \frac{1}{x} \cdot \frac{\frac{1}{x^n} - 1}{\frac{1}{x} - 1} = \frac{x^{n+1} + \frac{1}{x^n} - x - 1}{x - 1} \\
 &= \frac{\left(x^{n+\frac{1}{2}} + \frac{1}{x^{n+\frac{1}{2}}}\right) - \left(x^{\frac{1}{2}} + \frac{1}{x^{\frac{1}{2}}}\right)}{x^{\frac{1}{2}} - \frac{1}{x^{\frac{1}{2}}}} = \frac{2 \cos \frac{1}{2}(2n+1)a - 2 \cos \frac{1}{2}a}{2\sqrt{-1} \sin \frac{1}{2}a};
 \end{aligned}$$

$$\therefore S = \frac{1}{2 \sin \frac{1}{2}a} \cdot \left\{ \cos \frac{1}{2}a - \cos \frac{1}{2}(2n+1)a \right\} = \frac{\sin \frac{1}{2}na}{\sin \frac{1}{2}a} \cdot \sin \frac{1}{2}(n+1)a.$$

(2) Similarly, $2(\cos a + \cos 2a + \cos 3a + \dots + \cos na)$

$$\begin{aligned}
 &= \frac{\left(x^{2n+1} - \frac{1}{x^{2n+1}}\right) - \left(x^2 - \frac{1}{x^2}\right)}{x^{\frac{1}{2}} - \frac{1}{x^{\frac{1}{2}}}} = \frac{\sin \frac{1}{2}(2n+1)a - \sin \frac{1}{2}a}{\sin \frac{1}{2}a};
 \end{aligned}$$

$$\therefore \text{the series (2)} = \frac{\sin \frac{1}{2}na}{\sin \frac{1}{2}a} \cdot \cos \frac{1}{2}(n+1)a.$$

125. The formulæ of Chapters II. and III. may sometimes be employed for the like purpose.

To sum the Series,

$$(1) \quad \sin a + \sin(a+b) + \sin(a+2b) + \dots + \sin\{a+(n-1)b\},$$

$$(2) \quad \cos a + \cos(a+b) + \cos(a+2b) + \dots + \cos\{a+(n-1)b\}.$$

$$(1) \quad \text{Cos} \left(a - \frac{1}{2}b \right) - \text{cos} \left(a + \frac{1}{2}b \right) = 2 \sin \frac{1}{2}b \sin a, \dots (51, 4).$$

$$\text{so } \text{cos} \left(a + \frac{1}{2}b \right) - \text{cos} \left(a + \frac{3}{2}b \right) = 2 \sin \frac{1}{2}b \sin \left(a + b \right)$$

$$\text{cos} \left(a + \frac{3}{2}b \right) - \text{cos} \left(a + \frac{5}{2}b \right) = 2 \sin \frac{1}{2}b \sin \left(a + 2b \right)$$

.....

$$\text{cos} \left\{ a + \frac{1}{2}(2n-3)b \right\} - \text{cos} \left\{ a + \frac{1}{2}(2n-1)b \right\} = 2 \sin \frac{1}{2}b \sin \left\{ a + (n-1)b \right\};$$

\therefore by addition,

$$2 \sin \frac{1}{2}b \left[\sin a + \sin \left(a + b \right) + \sin \left(a + 2b \right) + \dots \right.$$

$$\left. + \sin \left\{ a + (n-1)b \right\} \right] = \text{cos} \left(a - \frac{1}{2}b \right) - \text{cos} \left\{ a + \frac{1}{2}(2n-1)b \right\};$$

$$\therefore \text{ the series (1) } = \frac{\sin \frac{1}{2}nb}{\sin \frac{1}{2}b} \cdot \sin \left\{ a + \frac{1}{2}(n-1)b \right\}.$$

(2) In like manner, beginning with $\text{Sin} \left(a + \frac{1}{2}b \right) - \text{sin} \left(a - \frac{1}{2}b \right)$ = $2 \sin \frac{1}{2}b \cos a$ and proceeding as in the last case,

$$\text{Cos } a + \text{cos} \left(a + b \right) + \text{cos} \left(a + 2b \right) + \dots + \text{cos} \left\{ a + (n-1)b \right\}$$

$$= \frac{\sin \frac{1}{2}nb}{\sin \frac{1}{2}b} \cdot \text{cos} \left\{ a + \frac{1}{2}(n-1)b \right\}.$$

Cor. Writing a for b in these results, the results of (124) are immediately obtained.

CHAPTER VI.

ON THE SOLUTION OF EQUATIONS AND THE RESOLUTION OF CERTAIN EXPRESSIONS INTO FACTORS.

126. To solve a Quadratic Equation.

1. Let $x^2 + px - q = 0$ be the equation.

$$\text{Then } x = -\left\{ \frac{p}{2} \mp \sqrt{\left(\frac{p^2}{4} + q \right)} \right\} = -\sqrt{q} \left\{ \frac{p}{2\sqrt{q}} \mp \sqrt{\left(\frac{p^2}{4q} + 1 \right)} \right\}.$$

$$\text{Let } \frac{4q}{p^2} = \tan^2 \theta \dots \dots \dots (1);$$

$$\begin{aligned} \therefore x &= -\sqrt{q} \left\{ \frac{1}{\tan \theta} \mp \frac{\sqrt{(\tan^2 \theta + 1)}}{\tan \theta} \right\} \\ &= -\sqrt{q} \cdot \frac{1 \mp \sec \theta}{\tan \theta} = -\sqrt{q} \cdot \frac{\cos \theta \mp 1}{\sin \theta}. \end{aligned}$$

$$\text{Now } \frac{\cos \theta - 1}{\sin \theta} = \frac{-2 \sin^2 \frac{1}{2} \theta}{2 \sin \frac{1}{2} \theta \cos \frac{1}{2} \theta} = -\tan \frac{1}{2} \theta,$$

$$\text{and } \frac{\cos \theta + 1}{\sin \theta} = \frac{2 \cos^2 \frac{1}{2} \theta}{2 \sin \frac{1}{2} \theta \cos \frac{1}{2} \theta} = \frac{1}{\tan \frac{1}{2} \theta}.$$

If therefore x_1 and x_2 be the two roots of the equation,

$$x_1 = \sqrt{q} \tan \frac{1}{2} \theta, \dots \dots (2); \quad x_2 = \frac{-\sqrt{q}}{\tan \frac{1}{2} \theta}, \dots \dots (3).$$

From (1), $L \tan \theta = 1_{10} 2 + \frac{1}{2} \cdot 1_{10} q - 1_{10} p + 10$, which determines θ ,

$$(2), \quad 1_{10} x_1 = \frac{1}{2} \cdot 1_{10} q + L \tan \frac{1}{2} \theta - 10,$$

$$(3), \quad 1_{10} (-x_2) = \frac{1}{2} \cdot 1_{10} q - L \tan \frac{1}{2} \theta + 10.$$

11. If the equation be $x^2 - px - q = 0$, the roots are the same as those of the preceding equation with their signs changed; for the product of the roots with their signs changed ($-q$) remains the same, and the sum of the roots with their signs changed (the coefficient of the second term) changes its sign, but continues of the same magnitude. (*Wood's Algebra*, 8th Ed., Art. 281.)

III. Let the equation be $x^2 - px + q = 0$; then

$$x = \frac{p}{2} \left\{ 1 \pm \sqrt{\left(1 - \frac{4q}{p^2}\right)} \right\}.$$

If $\frac{4q}{p^2}$ be less than 1, assume $\frac{4q}{p^2} = \sin^2 \theta$ (1).

Then, x_1 and x_2 being the roots of the equation,

$$x_1 = \frac{1}{2} p (1 + \cos \theta) = p \cos^2 \frac{1}{2} \theta \text{(2).}$$

$$x_2 = \frac{1}{2} p (1 - \cos \theta) = p \sin^2 \frac{1}{2} \theta \text{(3).}$$

From (1), $L \sin \theta = \text{I}_{10} 2 + \frac{1}{2} \cdot \text{I}_{10} q - \text{I}_{10} p + 10$,

$$(2), \quad \text{I}_{10} x_1 = \text{I}_{10} p + 2L \cos \frac{1}{2} \theta - 20,$$

$$(3), \quad \text{I}_{10} x_2 = \text{I}_{10} p + 2L \sin \frac{1}{2} \theta - 20.$$

IV. If the equation be $x^2 + px + q = 0$, the roots are the same as those of the last equation with their signs changed. (*Wood's Algebra*, Art. 281.)

In the last two cases, if $\frac{4q}{p^2}$ be > 1 , let $\frac{4q}{p^2} = \sec^2 \theta$;

$$\text{then } x = \frac{1}{2} p \{1 \pm \sqrt{-(\sec^2 \theta - 1)}\} = \frac{1}{2} p \{1 \pm \tan \theta \sqrt{-1}\},$$

and both roots of the equation are impossible.

These solutions may be employed in preference to the common method of solution when p and q are very large numbers.

Ex. *Required the roots of* $x^2 + 365.42x - 3469.1 = 0$.

By case (I.) $L \tan \theta = \text{I}_{10} 2 + \frac{1}{2} \cdot \text{I}_{10} 3469.1 - \text{I}_{10} 365.42 + 10$.

By the tables,

$$\text{I}_{10} 2 = .3010300$$

$$\frac{1}{2} \cdot \text{I}_{10} 3469.1 = \frac{1}{2} \times 3.5402168 = 1.7701084$$

10

$$\text{12.0711384}$$

$$\text{Subtract } \text{I}_{10} 365.42 = 2.5627923$$

$$9.5083461 = L \tan 17^\circ, 52', \text{ nearly};$$

$$\therefore \text{I}_{10} x_1 = \frac{1}{2} \cdot \text{I}_{10} 3469.1 + L \tan 8^\circ, 56' - 10.$$

$$\frac{1}{2} \cdot \text{I}_{10} 3469.1 = 1.7701084$$

$$L \tan 8^\circ, 56' = 9.1964302$$

$$\text{10.9665386}$$

Subtract 10

$$.9665386 = \text{I}_{10} 9.2584, \text{ nearly.}$$

Again, $1_{10}(-x_2) = \frac{1}{2} \cdot 1_{10} 3469 \cdot 1 - L \tan 8^\circ, 56' + 10.$

$$10 + \frac{1}{2} \cdot 1_{10} 3469 \cdot 1 = 11 \cdot 770 \cdot 084$$

$$L \tan 8^\circ, 56' = \frac{9 \cdot 1964302}{2 \cdot 5736782} = 1_{10} 374 \cdot 69 \text{ nearly.}$$

Hence the approximate values of the roots of the proposed equation are

$$9 \cdot 2584 \quad \text{and} \quad -374 \cdot 69.$$

After x_1 was found, x_2 might have been more easily determined from the equation

$$-(x_1 + x_2) = 365 \cdot 42. \quad (\text{Wood's Algebra, Art. 271.})$$

127. To solve a Cubic Equation.

Let the equation when transformed, if necessary, to another which wants the second term (*Wood's Algebra*, Art. 284), be $x^3 - qx - r = 0$; if $x = \frac{y}{n}$, this equation becomes $y^3 - qn^2y - rn^3 = 0$.

$$\text{Now } \cos^3 \phi - \frac{3}{4} \cos \phi - \frac{1}{4} \cos 3\phi = 0. \quad (55, \text{Cor.})$$

And this equation is identical with the equation $y^3 - qn^2y - rn^3 = 0$, if

$$(1) \quad \cos \phi = y.$$

$$(2) \quad \frac{3}{4} = qn^2; \quad \text{and} \quad \therefore n = \frac{1}{2} \sqrt{\frac{3}{q}}.$$

$$(3) \quad \frac{1}{4} \cos 3\phi = rn^3; \quad \text{and} \quad \therefore \cos 3\phi = 4rn^3 = \frac{r}{2} \sqrt{\frac{27}{q^3}}.$$

Let α be the least value of 3ϕ which satisfies the equation (3); then one value of y is $\cos \phi$, or $\cos \frac{1}{3} \alpha$.

Also since, (103), $\cos \alpha = \cos(2m\pi \pm \alpha)$, the two other values of y are contained among the values of $\cos \frac{1}{3}(2m\pi \pm \alpha)$.

Now m , being an integer, must be of one of the forms $3p, 3p \pm 1$,

$$\text{and } \cos \frac{2 \cdot 3p \cdot \pi \pm \alpha}{3} = \cos(2p\pi \pm \frac{1}{3} \alpha) = \cos \frac{1}{3} \alpha. \quad (103).$$

$$\cos \frac{2(3p \pm 1) \pi \pm \alpha}{3} = \cos\left(2p\pi \pm \frac{2\pi \pm \alpha}{3}\right) = \cos \frac{2\pi \pm \alpha}{3}.$$

Wherefore the three values of y are

$$\cos \frac{1}{3} \alpha; \quad \cos \frac{1}{3} (2\pi + \alpha); \quad \cos \frac{1}{3} (2\pi - \alpha);$$

and the three values of x , or $\frac{y}{n}$, are

$$2 \sqrt{\frac{q}{3}} \cdot \cos \frac{1}{3} \alpha; \quad 2 \sqrt{\frac{q}{3}} \cdot \cos \frac{1}{3} (2\pi + \alpha); \quad 2 \sqrt{\frac{q}{3}} \cdot \cos \frac{1}{3} (2\pi - \alpha).$$

128. *The solution of a cubic equation given in the last Article applies to those cases only where all the roots are possible: i. e. to the irreducible case of Cardan's Rule. [Wood's Algebra, Art. 331.]*

Since $\cos \alpha$ is necessarily less than 1,

$$\therefore \frac{r}{2} \sqrt{\frac{27}{q^3}} < 1, \quad \frac{r}{2} < \sqrt{\frac{q^3}{27}}, \quad \frac{r^2}{4} < \frac{q^3}{27};$$

which is the irreducible case of Cardan's Rule.

129. *To resolve the equation $x^{2n} - 1 = 0$ into its quadratic factors, n being a positive integer.*

Here $x^{2n} = 1$.

$$\text{Now } (\cos \theta + \sqrt{-1} \sin \theta)^{2n} = \cos 2n\theta + \sqrt{-1} \sin 2n\theta,$$

and if $\theta = \frac{m\pi}{n}$, m and n being positive integers,

$$\begin{aligned} \left(\cos \frac{m\pi}{n} + \sqrt{-1} \sin \frac{m\pi}{n} \right)^{2n} &= \cos 2m\pi + \sqrt{-1} \sin 2m\pi \\ &= 1, \quad \text{since } \sin 2m\pi = 0; \\ &= x^{2n}; \end{aligned}$$

$$\therefore x = \cos \frac{m\pi}{n} + \sqrt{-1} \sin \frac{m\pi}{n}.$$

Now m , being an integer, must be of the form $p \cdot 2n + r$, where p is 0 or any positive integer, and r is 0 or any positive integer less than $2n$.

$$\begin{aligned} \text{Wherefore, } x &= \cos \frac{(p \cdot 2n + r)\pi}{n} + \sqrt{-1} \sin \frac{(p \cdot 2n + r)\pi}{n} \\ &= \cos \left(2p\pi + \frac{r\pi}{n} \right) + \sqrt{-1} \sin \left(2p\pi + \frac{r\pi}{n} \right) \\ &= \cos \frac{r\pi}{n} + \sqrt{-1} \sin \frac{r\pi}{n}, \quad \text{by (103).} \end{aligned}$$

And for r writing $0, 1, 2, \dots, 2n-1$, successively in

$$x = \cos \frac{r\pi}{n} + \sqrt{-1} \sin \frac{r\pi}{n};$$

(1) If $r = 0$, $x = 1$,

(2) $r = 1$, $x = \cos \frac{\pi}{n} + \sqrt{-1} \sin \frac{\pi}{n}$,

(3) $r = 2$, $x = \cos \frac{2\pi}{n} + \sqrt{-1} \sin \frac{2\pi}{n}$,

.....

($n+1$) $r = n$, $x = -1$,

.....

($2n-1$) $r = 2n-2$, $x = \cos \frac{2n-2}{n} \pi + \sqrt{-1} \sin \frac{2n-2}{n} \pi$
 $= \cos \frac{2\pi}{n} - \sqrt{-1} \sin \frac{2\pi}{n}$, (163).

($2n$) $r = 2n-1$, $x = \cos \frac{2n-1}{n} \pi + \sqrt{-1} \sin \frac{2n-1}{n} \pi$
 $= \cos \frac{\pi}{n} - \sqrt{-1} \sin \frac{\pi}{n}$.

Now the equations (1) and ($n+1$) give the quadratic factor $(x-1)(x+1) = x^2-1$; the equations (2) and ($2n$) give

$$\left(x - \cos \frac{\pi}{n} - \sqrt{-1} \sin \frac{\pi}{n}\right) \left(x - \cos \frac{\pi}{n} + \sqrt{-1} \sin \frac{\pi}{n}\right) = x^2 - 2x \cos \frac{\pi}{n} + 1;$$

and so on;

Wherefore, $x^{2n} - 1 = \{x^2 - 1\} \left\{x^2 - 2x \cos \frac{\pi}{n} + 1\right\} \dots \left\{x^2 - 2x \cos \frac{n-1}{n} \pi + 1\right\}$.

130. To resolve the equation $x^n + 1 = 0$ into its quadratic factors, n being a positive integer.

$$x^{2n} + 1 = 0; \quad \therefore x^{2n} = -1.$$

Now $(\cos \theta + \sqrt{-1} \sin \theta)^{2n} = \cos 2n\theta + \sqrt{-1} \sin 2n\theta$. And making $2n\theta = (2m+1)\pi$, and proceeding as in the last Article,

$$x = \cos \frac{2m+1}{2n} \pi + \sqrt{-1} \sin \frac{2m+1}{2n} \pi.$$

Also, assuming $m = p \cdot 2n + r$, where p is 0 or anⁿ positive integer, and r is 0 or any positive integer less than $2n$, as before,

$$x = \cos \frac{2r+1}{2n} \pi + \sqrt{-1} \sin \frac{2r+1}{2n} \pi;$$

by means of which it may be proved that

$$\begin{aligned} x^{2n} + 1 &= \{x^2 - 2x \cos \frac{\pi}{2n} + 1\} \{x^2 - 2x \cos \frac{3\pi}{2n} + 1\} \dots \\ &\dots \{x^2 - 2x \cos \frac{2n-1}{2n} \pi + 1\}. \end{aligned}$$

131. *To resolve the equation $x^{2n+1} - 1 = 0$ into its quadratic factors, n being a positive integer.*

As in the last two Articles,

$$\begin{aligned} (\cos \theta + \sqrt{-1} \sin \theta)^{2n+1} &= \cos (2n+1) \theta + \sqrt{-1} \sin (2n+1) \theta \\ &= 1, \text{ if } (2n+1) \theta = 2m\pi, \text{ or } \theta = \frac{2m\pi}{2n+1}, \\ &= x^{2n+1}; \end{aligned}$$

$$\therefore x = \cos \frac{2m}{2n+1} \pi + \sqrt{-1} \sin \frac{2m}{2n+1} \pi;$$

and making $m = p \cdot (2n+1) + r$, this becomes

$$x = \cos \frac{2r\pi}{2n+1} + \sqrt{-1} \sin \frac{2r\pi}{2n+1}.*$$

Whence

$$x^{2n+1} - 1 = \{x - 1\} \{x^2 - 2x \cos \frac{2\pi}{2n+1} + 1\} \dots \{x^2 - 2x \cos \frac{2n\pi}{2n+1} + 1\}.$$

132. *To resolve the equation $x^{2n+1} + 1 = 0$ into its factors, n being a positive integer.*

Here $x^{2n+1} = -1$,

$$(\cos \theta + \sqrt{-1} \sin \theta)^{2n+1} = \cos (2n+1) \theta + \sqrt{-1} \sin (2n+1) \theta,$$

$$\text{and let } (2n+1) \theta = (2m+1) \pi, \quad \text{or } \theta = \frac{2m+1}{2n+1} \pi;$$

* Here if $r = 0$, $x = 1$, and one factor is $x - 1$; the values of x corresponding to $r = 1$ and $r = 2n$ form one quadratic factor; those corresponding to $r = 2$ and $r = 2n - 1$ form another; and so on.

$$\therefore (\cos \theta + \sqrt{-1} \sin \theta)^{2n+1} = -1 = x^{2n+1};$$

$$\therefore x = \cos \frac{2m+1}{2n+1} \pi + \sqrt{-1} \sin \frac{2m+1}{2n+1} \pi;$$

and making $m = p \cdot (2n+1) + r$, this becomes

$$x = \cos \frac{2r+1}{2n+1} \pi + \sqrt{-1} \sin \frac{2r+1}{2n+1} \pi.$$

Whence, as before,

$$x^{2n+1} + 1 = \{x + 1\} \left\{x^2 - 2x \cos \frac{\pi}{2n+1} + 1\right\} \dots\dots$$

$$\dots\dots \left\{x^2 - 2x \cos \frac{2n-1}{2n+1} \pi + 1\right\}.*$$

133. To resolve $\sin \theta$ and $\cos \theta$ into factors.

The values of θ which satisfy the equation $\sin \theta = 0$ are $0, \pm \pi, \pm 2\pi, \pm 3\pi, \dots, \pm n\pi$; n being any integer whatever. Assuming then that the series which expresses the value of $\sin \theta$ is divisible by $\theta, \theta - \pi, \theta + \pi, \theta - 2\pi, \theta + 2\pi, \&c.$, let $\sin \theta = a \cdot \theta (\theta - \pi) (\theta + \pi) (\theta - 2\pi) (\theta + 2\pi) \dots$, where a is some constant quantity whose value is to be determined.

$$\therefore \sin \theta = \pm a \cdot \theta (\pi - \theta) (\pi + \theta) (2\pi - \theta) (2\pi + \theta) \dots\dots$$

$$= \pm a \cdot \theta \cdot \pi \left(1 - \frac{\theta}{\pi}\right) \cdot \pi \left(1 + \frac{\theta}{\pi}\right) \cdot 2\pi \left(1 - \frac{\theta}{2\pi}\right) \cdot 2\pi \left(1 + \frac{\theta}{2\pi}\right) \dots\dots$$

$$= \pm a \cdot \pi^2 \cdot 2^2 \pi^2 \cdot 3^2 \pi^2 \cdot \dots \times \theta \left(1 - \frac{\theta^2}{\pi^2}\right) \left(1 - \frac{\theta^2}{2^2 \pi^2}\right) \dots\dots$$

$$\therefore \frac{\sin \theta}{\theta} = \pm a \cdot \pi^2 \cdot 2^2 \pi^2 \cdot 3^2 \pi^2 \cdot \dots \times \left(1 - \frac{\theta^2}{\pi^2}\right) \left(1 - \frac{\theta^2}{2^2 \pi^2}\right) \dots\dots$$

now if θ become 0, $\frac{\sin \theta}{\theta}$ becomes 1, (104), and this equation becomes

$$1 = \pm a \cdot \pi^2 \cdot 2^2 \pi^2 \cdot 3^2 \pi^2 \cdot \dots$$

* Here the values of x corresponding to the values 0 and $2n$ of r form one quadratic factor; those corresponding to the values 1 and $2n-1$ form another; and so on. When r is n the value of x is -1 , and $x+1$ is therefore a factor.

$$\therefore \sin \theta = \theta \left(1 - \frac{\theta^2}{\pi^2}\right) \left(1 - \frac{\theta^2}{2^2 \pi^2}\right) \left(1 - \frac{\theta^2}{3^2 \pi^2}\right) \dots \quad (1)$$

Again; the values of θ which satisfy the equation $\cos \theta = 0$ are $\pm \frac{\pi}{2}$, $\pm \frac{3\pi}{2}$, $\pm \frac{5\pi}{2}$, ... $\pm \frac{2n+1}{2} \pi$,; n being any integer whatever.

Hence, making an assumption like that in the former case,

$$\cos \theta = \pm b \cdot \frac{\pi^2}{2^2} \cdot \frac{3^2 \pi^2}{2^2} \cdot \dots \left(1 - \frac{2^2 \theta^2}{\pi^2}\right) \left(1 - \frac{2^2 \theta^2}{3^2 \pi^2}\right) \dots$$

Now if $\theta = 0$, this equation becomes $1 = \pm b \cdot \frac{\pi^2}{2^2} \cdot \frac{3^2 \pi^2}{2^2} \dots$

$$\therefore \cos \theta = \left(1 - \frac{2^2 \theta^2}{\pi^2}\right) \left(1 - \frac{2^2 \theta^2}{3^2 \pi^2}\right) \left(1 - \frac{2^2 \theta^2}{5^2 \pi^2}\right) \dots \quad (2)$$

COR. If $\theta = \frac{1}{2} \pi$, the expression for $\sin \theta$ becomes,

$$\begin{aligned} 1 &= \frac{\pi}{2} \left(1 - \frac{1}{2^2}\right) \left(1 - \frac{1}{4^2}\right) \left(1 - \frac{1}{6^2}\right) \left(1 - \frac{1}{8^2}\right) \dots \\ &= \frac{\pi}{2} \cdot \frac{2^2 - 1}{2^2} \cdot \frac{4^2 - 1}{4^2} \cdot \frac{6^2 - 1}{6^2} \cdot \frac{8^2 - 1}{8^2} \dots \\ &= \frac{\pi}{2} \cdot \frac{(2-1)(2+1)}{2^2} \cdot \frac{(4-1)(4+1)}{4^2} \cdot \frac{(6-1)(6+1)}{6^2} \dots \\ \therefore \pi &= 2 \cdot \frac{2^2}{1 \cdot 3} \cdot \frac{4^2}{3 \cdot 5} \cdot \frac{6^2}{5 \cdot 7} \cdot \frac{8^2}{7 \cdot 9} \dots \end{aligned}$$

This is Wallis's Theorem for the determination of π ; in which the successive factors become more and more nearly equal to 1.

APPENDIX I.

ON THE LOGARITHMS OF NUMBERS, AND THE CONSTRUCTION OF THE LOGARITHMIC TABLES OF NUMBERS.

1. **DEF.** If $n = a^x$, x is called the *Logarithm* of the Number n to the Base a ; or the *Logarithm of a Number to a given Base* is that power to which the base must be raised to give the number.

If a logarithmic formula be generally true whatever may be the value of the base, the logarithms of the quantities involved will be written thus, $\log m$, $\log n$; but if the logarithms are calculated to some particular base, (as 10 for instance), they will be written thus, $\log_{10} m$, $\log_{10} n$; or thus, $l_{10} m$, $l_{10} n$.

If $n = a^x$, and, while a remains the same, successive values be given to n , and the corresponding values of x be registered, the tables so formed are called "*Tables of a System of Logarithms to the Base a.*"

It will hereafter (10) be shewn that a system of logarithms calculated to the base 10 is attended with peculiar advantages.

2. Since if $n = a^x$, $x = l_a n$;
therefore, in all cases, $n = a^x = a^{l_a n}$.

COR. 1. If $x = 1$, n becomes $= a$, and $\therefore l_a a = 1$.

If $x = 0$, a^x (or n) becomes 1; and $\therefore l_a 1 = 0$.

COR. 2. If a be the base of any system of logarithms,

since $m = a^{l_a m}$, and $n = a^{l_a n}$,

$$\therefore m^{\frac{1}{l_a m}} = a, \quad \text{and} \quad n^{\frac{1}{l_a n}} = a,$$

$$\therefore m^{\frac{1}{l_a m}} = n^{\frac{1}{l_a n}}; \quad \text{and} \quad m^{l_a n} = n^{l_a m};$$

a true result, whatever be the value of a .

Wherefore, $m^{\log n} = n^{\log m}$.

3. *Required from tables of logarithms calculated to a given base, as ϵ , to form tables of logarithms to any other base, as 10.*

$$\text{By (2), } n = \epsilon^{l_\epsilon n}; \quad n = 10^{l_{10} n}; \quad 10 = \epsilon^{l_\epsilon 10};$$

$$\therefore \epsilon^{l_\epsilon n} = 10^{l_{10} n} = (\epsilon^{l_\epsilon 10})^{l_{10} n} = \epsilon^{l_\epsilon 10 \cdot l_{10} n};$$

and equating the indices of ϵ ,

$$l_\epsilon n = l_\epsilon 10 \cdot l_{10} n; \quad \text{and } l_{10} n = \frac{1}{l_\epsilon 10} \cdot l_\epsilon n.$$

Hence the logarithms of any number n in two systems calculated to any bases, as 10 and ϵ , are connected by a constant multiplier, (viz. $\frac{1}{l_\epsilon 10}$) which is called the "modulus;" and therefore from tables of logarithms calculated to a base ϵ , tables may be formed to the base 10.

COR. By writing a for 10 in this proof, $l_a n = \frac{1}{l_\epsilon a} \cdot l_\epsilon n$; or, the modulus connecting the logarithms of a number in the two systems whose bases are a and ϵ , is $\frac{1}{l_\epsilon a}$.

4. *The logarithm of the product of any number of factors is equal to the sum of the logarithms of the several factors.*

$$\text{For } m \cdot n \cdot r \dots = a^{l_a m} \cdot a^{l_a n} \cdot a^{l_a r} \dots = a^{(l_a m + l_a n + l_a r + \dots)}$$

$$\text{But } m \cdot n \cdot r \dots = a^{l_a(m \cdot n \cdot r \dots)};$$

$$\therefore l_a(m \cdot n \cdot r \dots) = l_a m + l_a n + l_a r + \dots$$

$$\text{or, } \log(m \cdot n \cdot r \dots) = \log m + \log n + \log r + \dots$$

Hence if there be found in the tables the number whose logarithm is the sum of the logarithms of the several factors, the product of those factors is obtained.

5. *The logarithm of a quotient is equal to the logarithm of the dividend minus the logarithm of the divisor.*

$$\text{For } a^{l_a\left(\frac{m}{n}\right)} = \frac{m}{n} = \frac{a^{l_a m}}{a^{l_a n}} = a^{l_a m - l_a n};$$

$$\therefore l_a\left(\frac{m}{n}\right) = l_a m - l_a n; \quad \text{or, } \log\left(\frac{m}{n}\right) = \log m - \log n.$$

Hence if there be found in the tables the number whose logarithm is the logarithm of the dividend minus the logarithm of the divisor, the quotient is obtained.

6. *The logarithm of the cth power of any number is equal to c times the logarithm of the number;—c being either whole or fractional.*

For $a^{l_a(m^c)} = m^c = (a^{l_a m})^c = a^{c \cdot l_a m}$;

$\therefore l_a(m^c) = c \cdot l_a m$; or, $\log(m^c) = c \cdot \log m$.

Also, $a^{l_a(m^{\frac{1}{c}})} = m^{\frac{1}{c}} = (a^{l_a m})^{\frac{1}{c}} = a^{\frac{1}{c} \cdot l_a m}$;

$\therefore l_a(m^{\frac{1}{c}}) = \frac{1}{c} \cdot l_a m$; or, $\log(m^{\frac{1}{c}}) = \frac{1}{c} \cdot \log m$.

Hence if any number be given, the cth power of it is that number in the tables whose logarithm is c times the logarithm of the given number; and the cth root of it is that number in the tables whose logarithm is $\frac{1}{c}$ th of the logarithm of the given number.

7. From the last three Articles it appears that the operations of multiplication, division, involution, and evolution can be performed with greater facility by means of tables of logarithms than by the common arithmetical methods, particularly when the numbers are large. The easier arithmetical operations of addition and subtraction cannot be performed by logarithms.

8. *In the common (or Briggs') system of logarithms, the base of which is 10, the logarithms of $10^n \times N$ and $\frac{N}{10^n}$ may be determined from the logarithm of N.*

For $l_{10}(10^n \times N) = l_{10} 10^n + l_{10} N$, by (4)

$= n \cdot l_{10} 10 + l_{10} N, \dots$ (6)

$= n + l_{10} N \dots \dots \dots$ (2, Cor. 1).

And $l_{10}\left(\frac{N}{10^n}\right) = l_{10} N - l_{10} 10^n, \dots \dots \dots$ (5).

$= l_{10} N - n.$

Thus from the pages of logarithms printed at the end of this Appendix,

$$\log 6 = 0.7781513;$$

$$\therefore \log 60 = \log 10 + \log 6 = 1 + 0.7781513 = 1.7781513,$$

$$\log 600 = \log(10^2) + \log 6 = 2 + 0.7781513 = 2.7781513,$$

$$\log 0.6 = \log 6 - \log 10 = 0.7781513 - 1,$$

$$\log 0.06 = \log 6 - \log(10^2) = 0.7781513 - 2.$$

The last two logarithms are written thus, $\bar{1}.7781513$, $\bar{3}.7781513$.

DEF. The *integral* part of the logarithm is called the "*Characteristic*" of the logarithm of the number; the *decimal* part is called the "*Mantissa*"* of the *significant digits* of which the number is composed.

Thus, in $\log 600 = 2.7781513$, 2 is the *Characteristic* of the *Logarithm of the Number 600*, .7781513 is the *Mantissa* of the *significant digit 6*.

9. In the common system, to determine the *Characteristic of the Logarithm of any given Number*.

If a number be between

1 and 10,	its log. is between 0 and 1;	\therefore the characteristic is 0.
10 and 100, 1 and 2;	\therefore 1.
100 and 1000, 2 and 3;	\therefore 2.
10^{n-1} and 10^n $n-1$ and n ;	\therefore $n-1$.

Hence the characteristic of the *Logarithm of a Number of n integral places* (and which therefore lies between 10^{n-1} and 10^n), is $n-1$, or is less by one than the number of integral places of figures in the number.

Again, if the number be between

1 and $\frac{1}{10}$,	its log. is between 0 and -1 ;	\therefore the characteristic is 1,
$\frac{1}{10}$ and $\frac{1}{100}$, -1 and -2 ;	\therefore $\bar{2}$,
$\frac{1}{10^{n-1}}$ and $\frac{1}{10^n}$, $-(n-1)$ and $-n$;	\therefore \bar{n} .

* "MANTISSA;" a handful thrown in over and above the exact weight; an overplus.

Hence, the Characteristic of the logarithm of a decimal fraction having $n - 1$ cyphers after the decimal point, is \bar{n} .

Generally therefore, *the Characteristic of the Logarithm of any Number is the number of its digits minus one*; where if the number be a decimal fraction, the cyphers which follow the decimal point are alone counted, and are reckoned negatively.

Wherefore, conversely, if logarithms be given having characteristics 1, 2, 3... $\bar{1}$, 2, 3... there are in the integral parts of the numbers to which these logarithms belong 2, 3, 4.....0, - 1, - 2... digits respectively.

Thus the logarithms of 245 and 25400 have for Characteristics 2 and 4, and the Characteristics of the Logarithms of 2.54, 25.4, 0.254, .000254 are 0, 1, 1, 4.

The Mantissa given in the tables for $l_{10} 3652$ is .5625308.

$$\begin{aligned} \therefore l_{10} 3652 &= 3.5625308, & l_{10} 36.52 &= 1.5625308, \\ l_{10} 365200 &= 5.5625308, & l_{10} .3652 &= \bar{1}.5625308, \\ l_{10} .003652 &= 3.5625308. \end{aligned}$$

10. *The advantages of Briggs' system of logarithms.*

From the last Article it appears, that if the *Base* of the system be 10, it is requisite to register the *Mantissæ* only in the tables; because the *Characteristics* can be determined by counting the digits in the integral part of the numbers whose logarithms are required. This omission of the characteristics renders the common tables *less bulky* than those calculated to any other base.

Also, from (8) it appears that in this system the Mantissa of N is also the Mantissa of $10^n \times N$, and of $\frac{N}{10^n}$,—where n is any integer: this circumstance renders the common tables *more comprehensive* than if any other base were taken; for if any other base were used, the Mantissa of N would *not* be the Mantissa of $10^n \cdot N$, or of $\frac{N}{10^n}$.

In the same way it might be shewn that if the system of arithmetic in common use were duodecimal instead of being decimal, tables calculated to the base 12 would possess the same advantages which have been here shewn to belong to the tables in common use.

11. The tables of logarithms in common use register, some to *five*, and others to *seven*, places of decimals, the mantissæ for numbers from 1 to 100000. Two pages of logarithms are printed at the end of this Appendix, in which the mantissæ are calculated to *seven* places of decimals. The line at the top of the second of the pages begins with the number 3650 (or 36500), and its mantissa .5622929 is placed opposite to it. And because the mantissæ of all numbers from 36500 to 36599, (comprised in the first six lines of the page), have the same initial

three figures, viz. 562, these three figures are registered, once for all, opposite to the number 3650, and the last four figures of each succeeding mantissa are placed under the number to which they belong. Thus the first line of the page is

Num.	0	1	2	3	4	5	6	7	8	9
3650	5622929	3048	3167	3286	3405	3524	3642	3761	3880	3999

Whence is obtained, Mantissa of 36500 = 562 2929
 36501 = 562 3048
 36502 = 562 3167
 36503 = 562 3286,

and so on.

In the same manner, the next line gives the mantissæ of numbers from 38510 to 36519 inclusive.

12. Since a change in the value of the third figure may not take place at the *beginning* of one of the horizontal lines, whenever a mantissa is sought from the tables, care must be taken to get the *right* initial figures. Thus (see p. 111), the mantissa of 36643 is 5639910, and the last four figures of the mantissa of 36644 are put down as 0029. Now if the mantissæ of these two numbers had the *same* third figure, the mantissa of 36644 would be less than that of 36643, which cannot possibly be. A change in the value of the third figure does, in point of fact, take place here; and the mantissa of 36644 (as do those of the numbers immediately following 36644) begins with 564, and not with 563.

Similar changes of the third figure of the mantissa occur at the numbers 36729, 36813, 36898, 36983, and are indicated by printing in a smaller type the fourth figure of the mantissæ of those numbers.

The construction and use of the small tables in the last column of page 111 will be explained hereafter.

13. EXAMPLES. (1) *To multiply 23 by 16. Art. 4.*

By the tables, p. 110, Mantissa of 23 is 3617278,

Mantissa of 16 is 2041200;

$$\therefore l_{10} 23 = 1.3617278$$

$$l_{10} 16 = 1.2041200$$

$$2.5658478$$

And, p. 111, the significant digits corresponding to the mantissa 5658478 are 3680;

$\therefore l_{10} 3680$ or $l_{10} 368 = 2.559478$; and $\therefore 368$ is the product sought.

(2) To multiply $\cdot 0172$ by $\cdot 00214$.

$$\begin{aligned} 1_{10} \cdot 0172 &= \bar{2} \cdot 2355284 \\ 1_{10} \cdot 00214 &= \bar{3} \cdot 3304138, \\ &\quad \bar{5} \cdot 5659422, \\ &= 1_{10} \cdot 000036808; \quad \text{p. 111;} \end{aligned}$$

Wherefore $\cdot 000036808$ is the product sought.

(3) To divide 3672 by 51000 . Art. 5.

$$\begin{aligned} 1_{10} 3672 &= 3 \cdot 5649027 \quad \text{p. 111.} \\ 1_{10} 51000 &= 4 \cdot 7075702 \quad \text{p. 110.} \\ &\quad \underline{2 \cdot 8573325} \\ &= 1_{10} \cdot 072; \end{aligned}$$

Wherefore $\cdot 072$ is the quotient sought.

(4) To find the values of $(15 \cdot 4)^3$, and $(650)^{\frac{1}{3}}$. Art. 6.

$$\begin{aligned} 1_{10} 15 \cdot 4 &= 1 \cdot 1875207 \quad \text{p. 110.} \\ &\quad \underline{\quad \quad \quad 3} \\ &3 \cdot 5625621 = 1_{10} 3652 \cdot 3, \text{ nearly,} \quad \text{p. 111.} \end{aligned}$$

$\therefore 3652 \cdot 3$ is the approximate cube of $15 \cdot 4$.

$$\text{Again, } 1_{10} 650 = 2 \cdot 8129134, \quad \text{p. 110.}$$

$$\therefore \frac{1}{5} \cdot 1_{10} 650 = \cdot 5625826 = 1_{10} 3 \cdot 6524, \text{ nearly,} \quad \text{p. 111.}$$

$\therefore 3 \cdot 6524$ is the approximate fifth root of 650 .

(5) To find the values of $(\cdot 085)^4$, and of $(\cdot 000065)^{\frac{1}{4}}$.

$$\begin{aligned} 2 \cdot 9294189 &= 1_{10} \cdot 085 \\ &\quad \underline{\quad \quad \quad 4} \\ &\bar{5} \cdot 7176756 \end{aligned}$$

and the mantissa is that of 522006 , nearly. Therefore $\cdot 0000522006$ is the number sought, nearly.

$$\text{Again, } 1_{10} \cdot 000065 = 5 \cdot 8129134 = 6 + 1 \cdot 8129134;$$

$$\therefore 3) 5 \cdot 8129134$$

$$\underline{\quad \quad \quad 2 \cdot 6043044} = 1_{10} \cdot 040207, \text{ nearly.}$$

NOTE. In dividing it is to be remembered that $\bar{5} \cdot 8 \dots = \bar{6} + 1 \cdot 8 \dots$

14. To expand $\log_a(1+x)$ in a series ascending by powers of x .

Let p denote the logarithm of $(1+x)$ to the base a , so that $1+x = a^p$. Then $(1+x)^m = a^{mp} = \{1 + (a-1)\}^{mp}$. And expanding both sides of this equation by the binomial theorem,

$$1 + mx + m \cdot \frac{m-1}{2} \cdot x^2 + m \cdot \frac{m-1}{2} \cdot \frac{m-2}{3} \cdot x^3 + \dots \\ = 1 + mp(a-1) + mp \cdot \frac{mp-1}{2} \cdot (a-1)^2 + \dots;$$

and arranging according to powers of m ,

$$1 + (x - \frac{1}{2}x^2 + \frac{1}{3}x^3 - \dots)m + Pm^2 + Qm^3 + \dots \\ = 1 + \{(a-1) - \frac{1}{2}(a-1)^2 + \frac{1}{3}(a-1)^3 - \dots\} pm + P'm^2 + \dots$$

This relation being true for all values of m , the coefficients of like powers of m on each side of the equation will be equal; wherefore

$$\{(a-1) - \frac{1}{2}(a-1)^2 + \frac{1}{3}(a-1)^3 - \dots\} p = x - \frac{1}{2}x^2 + \frac{1}{3}x^3 - \dots;$$

$$\therefore \log_a(1+x) = p = \frac{x - \frac{1}{2}x^2 + \frac{1}{3}x^3 - \dots}{(a-1) - \frac{1}{2}(a-1)^2 + \frac{1}{3}(a-1)^3 - \dots} \dots \dots \dots (i.)$$

Let ϵ be the quantity which when substituted for a makes the denominator of this fraction equal to unity; then

$$\log_\epsilon(1+x) = x - \frac{1}{2}x^2 + \frac{1}{3}x^3 - \dots, \dots \dots \dots (ii.)$$

$$\text{and } \therefore \log_a(1+x) = (a-1) - \frac{1}{2}(a-1)^2 + \frac{1}{3}(a-1)^3 - \dots \dots \dots (iii.)$$

And hence

$$\log_a(1+x) = \frac{1}{\log_\epsilon a} (x - \frac{1}{2}x^2 + \frac{1}{3}x^3 - \dots) \dots \dots \dots (iv.)$$

Cor. The modulus $\frac{1}{\log_{\epsilon} 10}$ of the common system of logarithms may be calculated thus; by taking for granted the formulae (ix.) and (vi.) which are proved hereafter.

$$\begin{aligned}
 \text{By (ix.), } l_{\epsilon 5} &= l_{\epsilon 4} + 2 \left\{ \frac{1}{8+1} + \frac{1}{3} \cdot \frac{1}{(8+1)^2} + \frac{1}{5} \cdot \frac{1}{(8+1)^3} + \dots \right\} \\
 &= 2 \cdot l_{\epsilon 2} + 2 \left\{ \frac{1}{9} + \frac{1}{3} \cdot \frac{1}{9^2} + \frac{1}{5} \cdot \frac{1}{9^3} + \dots \right\}; \\
 \therefore l_{\epsilon 10} &= l_{\epsilon 2} + l_{\epsilon 5}, = 3 \cdot l_{\epsilon 2} + 2 \left\{ \frac{1}{9} + \frac{1}{3} \cdot \frac{1}{9^2} + \frac{1}{5} \cdot \frac{1}{9^3} + \dots \right\} \\
 &= \left\{ \begin{array}{l} 6 \left\{ \frac{1}{3} + \frac{1}{3} \cdot \frac{1}{3^2} + \frac{1}{5} \cdot \frac{1}{3^3} + \dots \right\} \text{ by (vi.),} \\ + 2 \left\{ \frac{1}{9} + \frac{1}{3} \cdot \frac{1}{9^2} + \frac{1}{5} \cdot \frac{1}{9^3} + \dots \right\} \end{array} \right\} \\
 &= 2 \cdot \left\{ \begin{array}{l} 1 + \frac{1}{3} \cdot \frac{1}{9} + \frac{1}{5} \cdot \frac{1}{9^2} + \dots \\ + \frac{1}{9} + \frac{1}{3} \cdot \frac{1}{9^2} + \frac{1}{5} \cdot \frac{1}{9^3} + \dots \end{array} \right\} \\
 &= 2 \cdot 30258509\dots
 \end{aligned}$$

Whence is obtained, $\frac{1}{l_{\epsilon 10}} = 0.434294819\dots$

15. Some rapidly-converging formulæ for the calculation of logarithms will next be investigated.

I. By (ii.), $l_{\epsilon} \frac{1}{x} = l_{\epsilon} (1 - \frac{x-1}{x}) = -\frac{x-1}{x} - \frac{1}{2} \left(\frac{x-1}{x} \right)^2 - \dots$

But $l_{\epsilon} \frac{1}{x} = l_{\epsilon} 1 - l_{\epsilon} x = -l_{\epsilon} x;$

$\therefore l_{\epsilon} x = \frac{x-1}{x} + \frac{1}{2} \left(\frac{x-1}{x} \right)^2 + \dots \dots \dots$ (v.)

II. Again; $l_{\epsilon}(1+x) = x - \frac{x^2}{2} + \frac{x^3}{3} - \dots;$ and $l_{\epsilon}(1-x) = -x - \frac{x^2}{2} - \frac{x^3}{3} - \dots;$

$\therefore l_{\epsilon}(1+x) - l_{\epsilon}(1-x), \text{ or } l_{\epsilon} \frac{1+x}{1-x}, = 2 \left\{ x + \frac{x^3}{3} + \frac{x^5}{5} + \dots \right\}.$

And if $\frac{1+x}{1-x} = x,$ and $\therefore x = \frac{x-1}{x+1},$ this becomes,

$l_{\epsilon} x = 2 \cdot \left\{ \frac{x-1}{x+1} + \frac{1}{3} \left(\frac{x-1}{x+1} \right)^3 + \frac{1}{5} \left(\frac{x-1}{x+1} \right)^5 + \dots \right\} \dots \dots \dots$ (vi.)

If x be a little greater than 1, this series converges very rapidly.

III. Again, $1_\epsilon x = 1_\epsilon (x^m)^{\frac{1}{m}} = m \cdot 1_\epsilon x^{\frac{1}{m}}$
 $= m \{ (x^m - 1) - \frac{1}{2} (x^m - 1)^2 + \dots \}$, by (ii.)

and, (x being greater than 1,) by assuming m of sufficient magnitude $x^{\frac{1}{m}}$ may be made to differ from 1 by any definite quantity, however small the quantity may be; in which case the succeeding terms of the series may be neglected as being of inconsiderable magnitude with respect to the first, and

$$1_\epsilon x = m (x^{\frac{1}{m}} - 1) \dots\dots\dots (vii.)$$

16. Having given $1_\epsilon x$, to find $1_\epsilon (x + z)$; z being small when compared with x .

$$1_\epsilon (x + z) = 1_\epsilon \left\{ x \cdot \left(1 + \frac{z}{x} \right) \right\} = 1_\epsilon x + 1_\epsilon \left(1 + \frac{z}{x} \right).$$

Expanding $1_\epsilon \left(1 + \frac{z}{x} \right)$ by (vi.), [since $\frac{\left(1 + \frac{z}{x} \right) - 1}{\left(1 + \frac{z}{x} \right) + 1} = \frac{z}{2x + z}$],

$$1_\epsilon (x + z) = 1_\epsilon x + 2 \left\{ \frac{z}{2x + z} + \frac{1}{3} \left(\frac{z}{2x + z} \right)^3 + \dots \right\} \dots\dots\dots (viii.)$$

• COR. If $z=1$; $1_\epsilon (1+x) = 1_\epsilon x + 2 \left\{ \frac{1}{2x+1} + \frac{1}{3} \cdot \frac{1}{(2x+1)^3} + \frac{1}{5} \cdot \frac{1}{(2x+1)^5} + \dots \right\} \dots\dots (ix.)$

which is useful in computing $1_\epsilon (1+x)$ from $1_\epsilon x$, particularly when x is large.

17. Having given the Napierian Logarithms of two successive numbers, $x - 1$ and x , to find that of the number next following.

$$1_\epsilon (x + 1) = 1_\epsilon \frac{x^2 - 1}{x - 1} = 1_\epsilon \frac{x^2 \left(1 - \frac{1}{x^2} \right)}{x - 1}$$

$$= 2 1_\epsilon x - 1_\epsilon (x - 1) + 1_\epsilon \left(1 - \frac{1}{x^2} \right).$$

Expanding $\left(1 - \frac{1}{x^2} \right)$ by (vi.), [since $\frac{\left(1 - \frac{1}{x^2} \right) - 1}{\left(1 - \frac{1}{x^2} \right) + 1} = -\frac{1}{2x^2 - 1}$],

$$1_\epsilon (x + 1) = 2 1_\epsilon x - 1_\epsilon (x - 1) - 2 \left\{ \frac{1}{2x^2 - 1} + \frac{1}{3} \cdot \frac{1}{(2x^2 - 1)^3} + \dots \right\} \dots\dots\dots (x.)$$

18. To expand a^x in a series ascending by powers of x ; i. e. to expand the number in a series ascending by powers of the Logarithm.

$$a^x = \{1 + (a - 1)\}^x$$

$$= 1 + x(a - 1) + x \cdot \frac{x-1}{2} \cdot (a - 1)^2 + x \cdot \frac{x-1}{2} \cdot \frac{x-2}{3} \cdot (a - 1)^3 + \dots$$

let the coefficient of x {which is $(a - 1) - \frac{1}{2}(a - 1)^2 + \frac{1}{6}(a - 1)^3 - \dots$ } be represented by p_1 , and let the coefficients of x^2, x^3, \dots be represented by p_2, p_3, \dots ; then

$$a^x = 1 + p_1x + p_2x^2 + p_3x^3 + \dots$$

$$\therefore a^z = 1 + p_1z + p_2z^2 + p_3z^3 + \dots$$

Now $a^{(x+z)} = a^x \cdot a^z$,

$$\text{or } 1 + p_1(x+z) + p_2(x+z)^2 + \dots + p_n(x+z)^n + \dots$$

$$= \{1 + p_1x + p_2x^2 + \dots + p_nx^n + \dots\} \times \{1 + p_1z + p_2z^2 + \dots + p_nz^n + \dots\},$$

and by equating the coefficients of the terms involving $xz, x^2z, x^3z, \dots, x^{n-1}z, \dots$

$$2p_2 = p_1 \cdot p_1, \quad \therefore p_2 = \frac{p_1^2}{2}$$

$$3p_3 = p_1 \cdot p_2, \quad \therefore p_3 = \frac{p_1^3}{1 \cdot 2 \cdot 3}$$

$$4p_4 = p_1 \cdot p_3, \quad \therefore p_4 = \frac{p_1^4}{1 \cdot 2 \cdot 3 \cdot 4}$$

.....

$$n \cdot p_n = p_1 \cdot p_{n-1}, \quad \therefore p_n = \frac{p_1^n}{1 \cdot 2 \cdot 3 \dots n}$$

Now $p_1 = (a - 1) - \frac{1}{2}(a - 1)^2 + \frac{1}{6}(a - 1)^3 - \dots = l_e a$; by (iii.)

$$\therefore a^x = 1 + \frac{l_e a}{1} \cdot x + \frac{(l_e a)^2}{1 \cdot 2} \cdot x^2 + \frac{(l_e a)^3}{1 \cdot 2 \cdot 3} \cdot x^3 + \dots + \frac{(l_e a)^n}{1 \cdot 2 \cdot 3 \dots n} \cdot x^n + \dots \text{(xi.)}$$

Cor. If the base be e , (xi) becomes

$$e^x = 1 + x + \frac{x^2}{1 \cdot 2} + \frac{x^3}{1 \cdot 2 \cdot 3} + \dots + \frac{x^n}{1 \cdot 2 \dots n} + \dots \text{(xii.)}$$

19. To find the value of ϵ , the base of the Napierian system of logarithms.

If $x = 1$, and the base be ϵ , the series for a^x in the last Article becomes

$$\epsilon = 1 + 1 + \frac{1}{1 \cdot 2} + \frac{1}{1 \cdot 2 \cdot 3} + \dots + \frac{1}{1 \cdot 2 \cdot 3 \dots n} + \dots$$

$$\begin{aligned} \text{Now, } 1 + 1 + \frac{1}{1 \cdot 2} &= 2.5 \\ \frac{1}{1 \cdot 2 \cdot 3} &= .1666666666 \\ \frac{1}{1 \cdot 2 \cdot 3 \cdot 4} &= .0416666666 \\ \frac{1}{1 \cdot 2 \cdot 3 \cdot 4 \cdot 5} &= .0083333333 \\ \frac{1}{1 \cdot 2 \cdot 3 \cdot 4 \cdot 5 \cdot 6} &= .0013888888 \\ \frac{1}{1 \cdot 2 \cdot 3 \cdot 4 \cdot 5 \cdot 6 \cdot 7} &= .0001984126 \\ \frac{1}{1 \cdot 2 \cdot 3 \cdot 4 \cdot 5 \cdot 6 \cdot 7 \cdot 8} &= .0000248015 \\ \frac{1}{1 \cdot 2 \cdot 3 \dots 8 \cdot 9} &= .0000027557 \\ \frac{1}{1 \cdot 2 \cdot 3 \cdot 4 \dots 9 \cdot 10} &= .0000002755 \\ & \underline{\hspace{1.5cm}} \\ & 27182818 \end{aligned}$$

This result gives the correct value of ϵ , the base of the Napierian system, so far as the figures are put down. By taking more terms and a greater number of figures in each, the value of ϵ might be determined to any degree of accuracy required.

20. On the construction of the common tables of logarithms.

From one of the series (v), (vi) (vii), the Napierian logarithms of low prime numbers may be found. The logarithm of a high number which is not a prime may be determined from the logarithms of its factors, by resolving the number into powers of its prime factors;

$$\text{Thus, } 1_{\epsilon} 288 = 1_{\epsilon} 2^5 \cdot 3^2 = 1_{\epsilon} 2^5 + 1_{\epsilon} 3^2 = 5 \cdot 1_{\epsilon} 2 + 2 \cdot 1_{\epsilon} 3.$$

And the expressions (viii), (ix), (x), will greatly facilitate the computation of the logarithms of high numbers.

The Napierian logarithms having been determined, the tables to base 10 are

deduced from them by multiplying each by $\frac{1}{1_e 10}$, which is equal to .434294819...

Arts. 3 and 14, Cor. 4.

Some of the artifices employed in computing the tables may be found in Sharpe's "Method of making Logarithms" prefixed to Sherwin's Tables.

21. In the common tables are registered the mantissæ of the logarithms of numbers of five places of figures, these mantissæ being computed to seven places of decimals. At the side of each page is placed a "Table of proportional parts," by which, as it will be shewn, the mantissa may be found of the logarithm of a number containing six or seven places of figures; and conversely, if a logarithm be given whose mantissa is not contained exactly in the tables, the number corresponding to it may be determined, by means of these additional tables, to six or seven places of figures.

22. *On the construction and use of the Tables of Proportional Parts.*

Let m_1 and m_2 be the mantissæ of two consecutive integral numbers, n and $n + 1$, which contain five digits each; and let m be the mantissa of the number $n + \frac{a}{10}$, which contains six digits, the last of which (a) is after the decimal point.

Now, since n and $n + \frac{a}{10}$ have the same number of integral places, their logarithms have the same characteristic;

$$\therefore m - m_1 = l_{10} \left(n + \frac{a}{10} \right) - l_{10} n = l_{10} \frac{n + \frac{a}{10}}{n}, \text{ Art. 5.} = l_{10} \left(1 + \frac{a}{10n} \right)$$

which, by expanding $l_{10} \left(1 + \frac{a}{10n} \right)$ by (iv.) and neglecting the succeeding terms as being small compared with the first term, becomes $\frac{1}{1_e 10} \cdot \frac{a}{10n}$, nearly.

$$\text{Similarly, } m_2 - m_1 = l_{10} (n + 1) - l_{10} n = \frac{1}{1_e 10} \cdot \frac{1}{n};$$

$$\therefore m - m_1 = (m_2 - m_1) \cdot \frac{a}{10}.$$

Whence m may be found, if m_1, m_2, a , be given; or a may be determined, when m_1, m_2, m , are known.

Should $n + 1$ be a number of the form 10^p , where p is an integer, the above proof will hold, if the characteristic and the mantissæ of $l_{10}(n + 1)$ be taken to be $p - 1$ and 1 respectively.

23. By the Tables, p. 111,

$$\text{Mantissa of } 36633 = m_2 = \cdot 5638725$$

$$\dots\dots\dots 36632 = m_1 = \cdot 5638606$$

$$\therefore m_2 - m_1 = \cdot 0000119$$

And in the expression $m - m_1 = (m_2 - m_1) \frac{a}{10}$, writing for a the numbers 1, 2, 3, 4, 5, 6, 7, 8, 9 successively, the former of these tables is obtained.

a	$m_2 - m_1$	
1	$\cdot 0000119 \times \frac{1}{10}$ or $\cdot 00000119$	$\cdot 0000012$ nearly.
2	$\cdot 00000238$	$\cdot 0000024$
3	$\cdot 00000357$	$\cdot 0000036$
4	$\cdot 00000476$	$\cdot 0000048$
5	$\cdot 00000595$	$\cdot 0000060$
6	$\cdot 00000714$	$\cdot 0000071$
7	$\cdot 00000833$	$\cdot 0000083$
8	$\cdot 00000952$	$\cdot 0000095$
9	$\cdot 00001071$	$\cdot 0000107$

119	
1	12
2	24
3	36
4	48
5	60
6	71
7	83
8	95
9	107

Now the "difference" put down in the Tables for numbers near 36600 is 119, and the "Table of Proportional Parts" is the latter of the Tables above.

It appears then, that the *significant digits only* of the whole difference, and of the differences corresponding to the several digits, are inserted in the table of proportional parts. Hence the following Rule for constructing Tables of Proportional parts is evident:

Of the *significant* part of the whole difference point off the last digit as a decimal; (this is the same thing as multiplying the whole difference by $\frac{1}{10}$, the whole difference being treated as an integer). Multiply the resulting number by 1, 2, 3, . . . 9 successively, and the *whole* numbers thus obtained (the last digit in the integral part being increased by unity where the decimal part is not less than $\cdot 5$) are the significant parts of the differences for the digits respectively.

Thus, let the significant part of the whole difference be 156.

$15.6 \times 1 = 15.6 = 16$ nearly.	$15.6 \times 6 = 93.6 = 94$ nearly.
$\dots \times 2 = 31.2 = 31$	$\dots \times 7 = 109.2 = 109$
$\dots \times 3 = 46.8 = 47$	$\dots \times 8 = 124.8 = 125$
$\dots \times 4 = 62.4 = 62$	$\dots \times 9 = 140.4 = 140$.
$\dots \times 5 = 78.0 = 78$	

If the digit be given, the difference is immediately known from such a table, or *vice versa*. To avoid the necessity of performing the operation of subtraction in any particular case in order to find the whole difference, there is a line in the tables marked at the top with "Diff.," in which the difference is placed opposite to that logarithm at which such difference begins. To know what the difference therefore is in any particular case, it is merely requisite to take the number in this line next above the logarithm in question.

Ex. 1. *To find the Number whose Logarithm is 3.5677766.*

By the Tables, p. 111, the mantissa next below the given mantissa is that of $1_{10} 36963$, and the whole difference put down is 117.

$$\text{Mantissa of the given Logarithm} = m = .5677766$$

$$\text{Mantissa of } 1_{10} 36963 = m_1 = .5677672$$

$$\therefore m - m_1 = \underline{\quad 94}$$

By the Table of Proportional Parts to "Diff." 117, the difference 94 corresponds to the digit 8; therefore the significant part of the Number sought is 369638. Also, since the given Logarithm has 3 for its characteristic, the Number required is 3696.38.

Ex. 2. *Required the Logarithm of 367.654.*

$$\text{By p. 111, } 1_{10} 367.650 = 2.5654346$$

$$\text{And "Diff." being 118, Part for } \quad 4 = \quad 47$$

$$\therefore 1_{10} 367.654 = 2.5654393$$

24. *To find the Mantissa of the Logarithm of a Number which has seven places of digits.*

Let m_1 and m_2 be the mantissæ of n and $n + 1$, two successive integers of five digits each: let M be that of $n + \frac{a}{10} + \frac{b}{100}$, which has the same number of

integral places as n and $n + 1$, and has also one digit (a) in the place of the tenths, and another (b) in that of the hundredths.

Since a and b are digits after the decimal point, the numbers n and $n + \frac{a}{10} + \frac{b}{100}$ have the same number of integral places, their logarithms have the same characteristic, and therefore the difference of their mantissæ is the same as the difference of their logarithms;

$$\begin{aligned} \therefore M - m_1 &= \log_{10} \left(n + \frac{a}{10} + \frac{b}{100} \right) - \log_{10} n = \log_{10} \frac{n + \frac{a}{10} + \frac{b}{100}}{n} \\ &= \log_{10} \left\{ 1 + \frac{1}{n} \left(\frac{a}{10} + \frac{b}{100} \right) \right\} = \frac{1}{10} \cdot \frac{1}{n} \left(\frac{a}{10} + \frac{b}{100} \right) \text{ by (ii),} \end{aligned}$$

neglecting the terms of the series after the first.

Similarly, since n and $n + 1$ are integers of the same number of digits,

$$m_2 - m_1 = \log_{10} (n + 1) - \log_{10} n = \log_{10} \left(1 + \frac{1}{n} \right) = \frac{1}{10} \cdot \frac{1}{n};$$

$$\therefore M - m_1 = (m_2 - m_1) \left(\frac{a}{10} + \frac{b}{100} \right) = (m_2 - m_1) \frac{a}{10} + \frac{1}{10} \left\{ (m_2 - m_1) \frac{b}{10} \right\}.$$

Now the first part of the expression, $(m_2 - m_1) \frac{a}{10}$, is the quantity to be added to m_1 for the *first* additional digit a ; Art. 22. And $(m_2 - m_1) \frac{b}{10}$ is what *would* have been added, had b been the *first* additional digit, instead of being the *second*; wherefore the quantity which is to be added for b when it is the *second* additional digit, namely, $\frac{1}{10} \cdot \left\{ (m_2 - m_1) \frac{b}{10} \right\}$, is the tenth part of what would have been added, had b been the *first*, instead of being the *second*, additional figure.

The following Examples will explain what has been said.

Ex. 1. To find $\log_{10} 3684 \cdot 286$.

By the tables, p. 111, $\log_{10} 3684 \cdot 2 = 3 \cdot 5663432$, and "Diff." being 118, the part for a *first* additional digit 8 is 94; and for a *first* additional digit 6 the part is 71, and therefore the part when 6 is a *second* additional digit is $\frac{71}{10}$, or 7.1. The operation is thus performed;

	$\log_{10} 3684 \cdot 2$	$= 3 \cdot 5663432$
Part for	8	= 94
Part for	6	= 7.1

$$\therefore \log_{10} 3684 \cdot 286 = 3 \cdot 5663533$$

Ex. 2. To find the Number whose Logarithm is 2.5656560.

2.5656560

Now $2.5656471 = 1_{10}367.83$, p. 111.

89

$\frac{83}{89} =$ Part for digit 7; "Diff." being 118.

$\frac{6}{89} =$ Part for a second digit 5.

$\therefore 2.5656560 = 1_{10}367.8375$,

and therefore 367.8375 is the number sought.

25. On the adaptation of formulæ to logarithmic computation.

After a table of logarithms has once been constructed, the labour of certain arithmetical operations can be materially diminished, while at the same time the chance of committing errors is lessened. But by referring to Articles 4, 5, 6 of this Appendix, it will appear that the sole arithmetical operations which can be performed by logarithms, are those of Multiplication, Division, Involution, and Evolution. Before, therefore, the value of an expression can be calculated by means of logarithms, the expression must be put into such a form that no other arithmetic operations than these have to be performed. Such an arrangement of an expression is called the *adaptation of it to logarithmic computation*.

Thus if a, b, c , the three sides of a triangle, be given, logarithms cannot be *directly* applied to determine the value of $\cos A$ from the equation.

$$\cos A = \frac{b^2 + c^2 - a^2}{2bc};$$

but if from this equation the formula,

$$\cos \frac{1}{2}A = \sqrt{\frac{S(S-a)}{bc}}, \text{ where } S = \frac{1}{2}(a+b+c),$$

be deduced, logarithms can be immediately applied to determine the value of $\cos \frac{1}{2}A$.

For a, b, c being given, S and $S-a$ are easily determined; and these being known, $\cos \frac{1}{2}A$ is determined from the equation

$$1_{10} \cos \frac{1}{2}A = \frac{1}{2} \{ 1_{10} [S(S-a)] - 1_{10} bc \} = \frac{1}{2} \{ 1_{10} S + 1_{10} (S-a) - 1_{10} b - 1_{10} c \}.$$

The two accompanying pages of logarithms are taken from Babbage's Tables, the most correct and the best arranged, perhaps, of any which have been published. The columns have been omitted which in those Tables are given to determine the number of seconds in an angle containing a given number of degrees, minutes, and seconds, and conversely.

1	0000000	51	7075702	101	0043214	151	1789769	201	3031961
2	3010300	52	7160033	102	0086002	152	1818436	202	3053514
3	4771213	53	7242759	103	0128372	153	1846914	203	3074960
4	6020600	54	7323938	104	0170333	154	1875207	204	3096302
5	6989700	55	7403627	105	0211893	155	1903317	205	3117539
6	7781513	56	7481880	106	0253059	156	1931246	206	3138672
7	8450980	57	7558749	107	0293838	157	1958997	207	3159703
8	9030900	58	7634280	108	0334238	158	1986571	208	3180633
9	9542425	59	7708520	109	0374265	159	2013971	209	3201463
10	0000000	60	7781513	110	0413927	160	2041200	210	3222193
11	0413927	61	7853298	111	0453230	161	2068259	211	3242825
12	0791812	62	7923917	112	0492180	162	2095150	212	3263359
13	1130434	63	7993405	113	0530784	163	2121876	213	3283796
14	1461280	64	8061800	114	0569049	164	2148438	214	3304138
15	1760913	65	8129134	115	0606978	165	2174839	215	3324385
16	2041200	66	8195439	116	0644580	166	2201081	216	3344538
17	2304489	67	8260748	117	0681859	167	2227165	217	3364597
18	2552725	68	8325089	118	0718820	168	2253093	218	3384565
19	2787536	69	8388491	119	0755470	169	2278867	219	3404441
20	3010300	70	8450980	120	0791812	170	2304489	220	3424227
21	3222193	71	8512583	121	0827854	171	2329961	221	3443923
22	3424227	72	8573325	122	0863598	172	2355284	222	3463530
23	3617278	73	8633229	123	0899051	173	2380461	223	3483049
24	3802112	74	8692317	124	0934217	174	2405492	224	3502480
25	3979400	75	8750613	125	0969100	175	2430380	225	3521825
26	4149733	76	8808136	126	1003705	176	2455127	226	3541084
27	4313638	77	8864907	127	1038037	177	2479733	227	3560259
28	4471580	78	8920946	128	1072100	178	2504200	228	3579348
29	4623980	79	8976271	129	1105897	179	2528530	229	3598355
30	4771213	80	9030900	130	1139434	180	2552725	230	3617278
31	4913617	81	9084850	131	1172713	181	2576786	231	3636120
32	5051500	82	9138139	132	1205739	182	2600714	232	3654880
33	5185139	83	9190781	133	1238516	183	2624511	233	3673559
34	5314789	84	9242793	134	1271048	184	2648178	234	3692159
35	5440680	85	9294189	135	1303338	185	2671717	235	3710679
36	5563025	86	9344985	136	1335389	186	2695129	236	3729120
37	5682017	87	9395193	137	1367206	187	2718416	237	3747483
38	5797836	88	9444827	138	1398791	188	2741578	238	3765770
39	5910646	89	9493900	139	1430148	189	2764618	239	3783979
40	6020600	90	9542425	140	1461280	190	2787536	240	3802112
41	6127839	91	9590414	141	1492191	191	2810334	241	3820170
42	6232493	92	9637878	142	1522883	192	2833012	242	3838154
43	6334685	93	9684829	143	1553360	193	2855573	243	3856063
44	6434527	94	9731279	144	1583625	194	2878017	244	3873898
45	6532125	95	9777236	145	1613680	195	2900346	245	3891661
46	6627578	96	9822712	146	1643529	196	2922561	246	3909351
47	6720979	97	9867717	147	1673173	197	2944662	247	3926970
48	6812412	98	9912261	148	1702617	198	2966652	248	3944517
49	6901961	99	9956352	149	1731863	199	2988531	249	3961993
50	6989700	100	0000000	150	1760913	200	3010300	250	3979400

Log. 562. N. 365.

Num.	0	1	2	3	4	5	6	7	8	9	Diff.
3650	5622929	3048	3167	3286	3405	3524	3642	3761	3880	3999	119 119 1 2 3 4 5 6 7 8 9
1	4118	4237	4356	4475	4594	4713	4832	4951	5070	5189	
2	5308	5427	5546	5664	5783	5902	6021	6140	6259	6378	
3	6497	6616	6734	6853	6972	7091	7210	7329	7448	7567	
4	7685	7804	7923	8042	8161	8280	8398	8517	8636	8755	
5	8874	8993	9111	9230	9349	9468	9587	9705	9824	9943	
6	5630062	0181	0299	0418	0537	0656	0775	0893	1012	1131	
7	1250	1368	1487	1606	1725	1843	1962	2081	2200	2318	
8	2437	2556	2674	2793	2912	3031	3149	3268	3387	3505	
9	3624	3743	3861	3980	4099	4218	4336	4455	4574	4692	
3660	4811	4930	5048	5167	5285	5404	5523	5641	5760	5879	118 118 1 2 3 4 5 6 7 8 9
1	5997	6116	6235	6353	6472	6590	6709	6828	6946	7065	
2	7183	7302	7421	7539	7658	7776	7895	8013	8132	8251	
3	8369	8488	8606	8725	8843	8962	9081	9199	9318	9436	
4	9555	9673	9792	9910	0029	0147	0266	0384	0503	0621	
5	5640740	0858	0977	1095	1214	1332	1451	1569	1688	1806	
6	1925	2043	2162	2280	2398	2517	2635	2754	2872	2991	
7	3109	3228	3346	3464	3583	3701	3820	3938	4056	4175	
8	4293	4412	4530	4648	4767	4885	5004	5122	5240	5359	
9	5477	5595	5714	5832	5951	6069	6187	6306	6424	6542	
3670	6661	6779	6897	7016	7134	7252	7371	7489	7607	7726	118 118 1 2 3 4 5 6 7 8 9
1	7844	7962	8080	8199	8317	8435	8554	8672	8790	8908	
2	9027	9145	9263	9382	9500	9618	9736	9855	9973	0091	
3	5650209	0328	0446	0564	0682	0800	0919	1037	1155	1273	
4	1392	1510	1628	1746	1864	1983	2101	2219	2337	2455	
5	2573	2692	2810	2928	3046	3164	3282	3401	3519	3637	
6	3755	3873	3991	4109	4228	4346	4464	4582	4700	4818	
7	4936	5054	5173	5291	5409	5527	5645	5763	5881	5999	
8	6117	6235	6353	6471	6590	6708	6826	6944	7062	7180	
9	7298	7416	7534	7652	7770	7888	8006	8124	8242	8360	
3680	8478	8596	8714	8832	8950	9068	9186	9304	9422	9540	117 117 1 2 3 4 5 6 7 8 9
1	9658	9776	9894	0012	0130	0248	0366	0484	0602	0720	
2	5660838	0956	1074	1192	1310	1428	1545	1663	1781	1899	
3	2017	2135	2253	2371	2489	2607	2725	2843	2960	3078	
4	3196	3314	3432	3550	3668	3786	3903	4021	4139	4257	
5	4375	4493	4611	4728	4846	4964	5082	5200	5318	5435	
6	5553	5671	5789	5907	6025	6142	6260	6378	6496	6614	
7	6731	6849	6967	7085	7203	7320	7438	7556	7674	7791	
8	7909	8027	8145	8262	8380	8498	8616	8733	8851	8969	
9	9087	9204	9322	9440	9557	9675	9793	9911	0028	0146	
3690	5670264	0381	0499	0617	0734	0852	0970	1087	1205	1323	117 117 1 2 3 4 5 6 7 8 9
1	1440	1558	1676	1793	1911	2029	2146	2264	2382	2499	
2	2617	2735	2852	2970	3087	3205	3323	3440	3558	3675	
3	3793	3911	4028	4146	4263	4381	4499	4616	4734	4851	
4	4969	5086	5204	5322	5439	5557	5674	5792	5909	6027	
5	6144	6262	6379	6497	6615	6732	6850	6967	7085	7202	
6	7320	7437	7555	7672	7790	7907	8025	8142	8260	8377	
7	8495	8612	8729	8847	8964	9082	9199	9317	9434	9552	
8	9669	9787	9904	0021	0139	0256	0374	0491	0608	0726	
9	5680843	0961	1078	1196	1313	1430	1548	1665	1782	1900	

APPENDIX II.

ON THE CONSTRUCTION AND USE OF TABLES OF GONIOMETRIC RATIOS.


1. IF it be required to find the value of a trigonometrical formula in which the sines, cosines, tangents, secants, &c., of given angles enter, much labour will be avoided if the values of these quantities be determined once for all, and registered in Tables.

In very small angles the sines and tangents are exceedingly small quantities, and if they be expressed as decimal fractions, two or three cyphers will follow the decimal point before the significant digits are arrived at. Now in order to avoid the inconvenience of printing these cyphers, the real values of all Goniometrical Ratios are multiplied by 10,000 (or the decimal point is moved four places to the right) before they are registered in the tables; and the tables so formed are called, "Tables of natural sines, cosines, &c.*"

* The tables of Goniometric Ratios are sometimes said to be "calculated to a radius of 10,000." To explain the meaning of this expression.

If with C as centre, and radius CA , an arc AB be described, and BN , AT be \perp to CA , the lines NB , CN , AT , CT , AN respectively, are sometimes defined to be the *sine*, *cosine*, *tangent*, *secant*, and *versed sine*, of the angle ACB to the radius CA ,—or the *sine*, *cosine*, *tangent*, *secant*, and *versed sine*, of the arc AB .

The expression "to the radius CA " is necessary to these definitions, because the lines NB , AT ...depend on the magnitude of CA as well as upon that of the angle ACB ; in fact, for a given value of the angle ACB , those lines vary directly as CA .



Now

2. To find the sine and cosine of 10".

Let θ be the Circular Measure of an angle of 10".

$$\text{Then } \frac{\theta}{\pi} = \frac{10}{180 \times 60 \times 60} = \frac{1}{64800};$$

$$\therefore \theta = \cdot 000048481368110.$$

$$\text{But } \sin \theta > \theta - \frac{1}{4} \theta^3, \text{ Art. 105.}$$

\therefore , a fortiori,

$$\begin{aligned} \sin 10'' &> \theta - \frac{1}{4} (\cdot 00005)^3 \\ &> \cdot 000048481368110 - \cdot 000000000000032 \\ &> \cdot 000048481368078. \end{aligned}$$

$$\text{Also, } \sin 10'' < \theta, \text{ or, } \cdot 000048481368110.$$

Wherefore a near approximation to the value of $\sin 10''$ is obtained by taking the first twelve places which these two quantities have in common, and therefore

$$\sin 10'' = \cdot 000048481368 \text{ very nearly.}$$

By substituting this value of $\sin 10''$ in the formula $\cos 10'' = \sqrt{1 - \sin^2 10''}$, there is obtained

$$\cos 10'' = \cdot 9999999988248.$$

Now by the definition of the sine which has been adopted in this treatise,

$$\sin ACB = \frac{NB}{CB};$$

$$\therefore 10,000 \times \sin ACB = NB \cdot \frac{10,000}{CA}.$$

But $10,000 \times \sin ACB$ is the tabular, or natural, sine of ACB ;

$$\therefore \text{tab. } \sin ACB = NB \cdot \frac{10,000}{CA} = NB, \text{ if } CA = 10,000.$$

Wherefore the tab. sine of ACB expresses the magnitude of the line NB , (the Sine of ACB to the radius CA), the magnitude of CA being represented by 10,000.

Similarly, the tab. cosine, tab. tangent, &c., of ACB , express the magnitudes of the lines CN , AT , &c., the magnitude of CA being 10,000.

3. *The Sine and Cosine of 10'' being known, the Sines of all angles between 10'' and 90° may be calculated.*

$$\sin(A + B) = 2 \sin A \cdot \cos B - \sin(A - B);$$

and writing $n \cdot 10''$ for $\angle A$, and $10''$ for $\angle B$,

$$\sin(n + 1)10'' = 2 \sin n10'' \cdot \cos 10'' - \sin(n - 1)10''.$$

$$\begin{aligned} \text{Now } 2 \cos 10'' &= 1.9999999976496 \\ &= 2 - .0000000023504 \\ &= 2 - k \text{ suppose;} \end{aligned}$$

$$\begin{aligned} \therefore \sin(n + 1)10'' &= 2 \sin n10'' - k \cdot \sin n10'' - \sin(n - 1)10'' \\ &= \{\sin n \cdot 10'' - \sin(n - 1)10''\} + \sin n10'' - k \cdot \sin n10''. \end{aligned}$$

And by writing successively for n the numbers 1, 2, 3...

$$\sin 20'' = (\sin 10'' - \sin 0'') + \sin 10'' - k \cdot \sin 10'',$$

$$\sin 30'' = (\sin 20'' - \sin 10'') + \sin 20'' - k \cdot \sin 20'',$$

$$\sin 40'' = (\sin 30'' - \sin 20'') + \sin 30'' - k \cdot \sin 30''.$$

.....

This method is not very laborious. In the last line, $\sin 30''$ and $(\sin 30'' - \sin 20'')$ are known from the two lines preceding, and the chief labour is in multiplying $\sin 30''$ by k .

The Sines of angles up to 60° having been successively calculated by the above method, those of angles between 60° and 90° may be thus determined.

$$\sin(60^\circ + A) - \sin(60^\circ - A) = 2 \cos 60^\circ \cdot \sin A = \sin A;$$

$$\therefore \sin(60^\circ + A) = \sin A + \sin(60^\circ - A).$$

So that if A be made to increase by $10''$ at a time from 0° up to 60° , this last formula, by addition merely, will give the sines of angles from 60° to 90° .

4. The Sines of angles up to 90° having been determined, their Cosines are also known.

$$\text{For } \cos A = \sin(90^\circ - A).$$

$$\text{Thus } \cos 25^\circ = \sin(90^\circ - 25^\circ) = \sin 65^\circ; \quad \cos 72^\circ = \sin(90^\circ - 72^\circ) = \sin 18^\circ; \quad \&c.$$

5. The Tangents, Cotangents, Secants, and Cosecants can be determined from the Sines and Cosines.

$$\text{For } \tan A = \frac{\sin A}{\cos A}, \quad \cot A = \frac{\cos A}{\sin A}, \quad \sec A = \frac{1}{\cos A}, \quad \operatorname{cosec} A = \frac{1}{\sin A}.$$

6. Since,

$\sin A = \cos(90^\circ - A)$, $\tan A = \cot(90^\circ - A)$, $\sec A = \operatorname{cosec}(90^\circ - A)$,
 the Sines, Tangents, and Secants of angles from 45° to 90° are
 respectively the same as the Cosines, Cotangents, and Cosecants
 of angles from 45° to 0° . Wherefore it is unnecessary to carry
 the tables further than to the angle 45° .

$$\begin{aligned} \text{Thus } \quad \cos 72^\circ, 20' &= \sin(90^\circ - 72^\circ, 20') = \sin 17^\circ, 40', \\ \sin 72^\circ, 20' &= \cos 17^\circ, 40', \quad \tan 72^\circ, 20' = \cot 17^\circ, 40', \\ \sec 72^\circ, 20' &= \operatorname{cosec} 17^\circ, 40'. \end{aligned}$$

At the bottom of the page containing the Sines, &c. of angles
 from 17° to 18° is placed the angle 72° , and the column which at
 the top of the page is marked to indicate the Sines of angles
 from 17° to 18° , is marked at the bottom to shew the Cosines
 of angles from 72° to 73° ; and so for the other Goniometrical
 Ratios. See page 117.

7. *Formula of Verification.* Since the Goniometrical Ratios
 are determined *successively* one from another, one error will affect
 every successive result. As checks against the possibility of errors,
 several formulæ (of *verification* as they are called) are used to
 examine the accuracy of the results; and the values registered
 in the tables are presumed to be correct if they satisfy these
 formulæ.

The following are the principal formulæ of verification.

$$\begin{aligned} (1) \quad \sin A &= \frac{1}{2} \{ \sqrt{1 + \sin 2A} - \sqrt{1 - \sin 2A} \} \dots\dots\dots \text{Art. 41.} \\ (2) \quad \cos A &= \frac{1}{2} \{ \sqrt{1 + \sin 2A} + \sqrt{1 - \sin 2A} \} \end{aligned}$$

A being an angle less than 45° .

Again, $\cos 36^\circ = \frac{\sqrt{5} + 1}{4}$; $\cos 72^\circ = \frac{\sqrt{5} - 1}{4}$; Art. 58, (3), (1).

$$\begin{aligned} \therefore \sin(36^\circ + A) - \sin(36^\circ - A) &= 2 \cos 36^\circ \cdot \sin A = \frac{\sqrt{5} + 1}{2} \cdot \sin A, \\ \text{and } \sin(72^\circ + A) - \sin(72^\circ - A) &= 2 \cos 72^\circ \cdot \sin A = \frac{\sqrt{5} - 1}{2} \cdot \sin A; \end{aligned}$$

\therefore by subtraction,

$$\sin(36^\circ + A) + \sin(72^\circ - A) - \sin(36^\circ - A) - \sin(72^\circ + A) = \sin A.$$

(3) $\therefore \sin A + \sin (36^\circ - A) + \sin (72^\circ + A) = \sin (36^\circ + A) + \sin (72^\circ - A)$,
which is *Euler's formula*.

$$\text{Again, } \sin 54^\circ = \frac{\sqrt{5+1}}{4}; \quad \sin 18^\circ = \frac{\sqrt{5-1}}{4}; \quad \text{Art. 58, (3), (1),}$$

$$\left. \begin{aligned} \text{and } 2 \sin 54^\circ \cdot \cos A &= \sin (54^\circ + A) + \sin (54^\circ - A) \\ 2 \sin 18^\circ \cdot \cos A &= \sin (18^\circ + A) + \sin (18^\circ - A) \end{aligned} \right\}.$$

$$(4) \therefore \cos A, \text{ or } \sin (90^\circ - A),$$

$$= \sin (54^\circ + A) + \sin (54^\circ - A) - \sin (18^\circ + A) - \sin (18^\circ - A),$$

which is *Legendre's formula*.

This formula might have been proved by writing $90^\circ - A$ for A in Euler's formula.

Ex. To exemplify the use of these formulæ. By making $A = 13^\circ$ in *Legendre's formula*;

$$\cos 13^\circ = \sin 67^\circ + \sin 41^\circ - \sin 31^\circ - \sin 5^\circ.$$

Now the tables give for the quantities in the second member of the equation 9295.049 + 6560.590 - 5150.381 - 871.557, which = 9743.701, the quantity given by the tables as the Cosine of 13° .

Since, therefore, these quantities satisfy the relation which ought to exist between the Sines of 67° , 41° , 31° , 5° , and the Cosine of 13° , it may be concluded, without much chance of error, that the values of these Goniometric Ratios are correctly given by the tables.

8. The values of the Goniometric Ratios having been thus calculated, multiplied by 10,000, verified, and registered in tables, are called "Tables of Natural Sines, Cosines, Tangents, Cotangents, Secants, and Cosecants." To find the *real* Goniometric Ratios from these tables, the tabular numbers have to be divided by 10,000; that is, the decimal point has to be removed four places to be left.

9. The *logarithmic* sines, cosines, tangents, &c. of angles will next be treated of, by which, rather than by the natural goniometric ratios, mathematical calculations are most frequently made.

A page from Sherwin's Logarithmic Tables, calculated for angles which differ from one another by $1'$, is here subjoined, the Natural cosines, tangents, secants, and cosecants being omitted. The column following that of the *N. sines*, which is marked "*N.D. 1''*," will be explained in the next Appendix.

M	N. Sine.	D. 1'	L. Sine.	Diff 1 sec.	Co-secants.	L. Tan.	Diff. 1 sec.	L. Sec.	Diff 1 sec.	Co-sines.	M
0	2923 717		9-4659353	68 833	10-5340647	9 4855380	75 283	10 0194037	6 433	9 9805963	60
1	2926 499	46 36	9 466 483	68 786	10 5336517	9 4857907	75 200	10 0194423	6 430	9 9805577	59
2	2929 280		9 4677690	68 683	10-53 12391	9 4862419	75 150	10 0194810	6 440	9 9805190	58
3	2932 061		9 467 170	68 633	10 5326270	9 4868928	75 083	10 0195197	6 466	9 9804804	57
4	2934 842		9 4675848	68-533	10-5324152	9 4871433	75 000	10 0195580	6 466	9 9804415	56
5	2937 623		9 4674960	68-483	10-5320040	9 4874933	74 950	10 0195973	6 466	9 9804027	55
6	2940 403	46 33	9 4674069	68 400	10 5315631	9 4880430	74 900	10 0196361	6 483	9 9803639	54
7	2943 183		9 4673173	68 333	10-5311827	9 4884924	74 816	10 0196750	6 500	9 9803250	53
8	2945 963		9 4672273	68 266	10 5307727	9 4889413	74 750	10 0197140	6 483	9 9802860	52
9	2948 743		9 4671369	68-200	10-5303631	9 4893908	74 700	10 0197529	6 500	9 9802471	51
10	2951 522		9 4700461	68-116	10 5299539	9 4898390	74 633	10 0197919	6 516	9 9802081	50
11	2954 302	46 31	9 4704 46	68 050	10-5295452	9 4902868	74 566	10 0198310	6 516	9 9801690	49
12	2957 081		9 4708931	67 983	10-5291369	9 4907342	74 500	10 0198701	6 531	9 9801299	48
13	2959 859		9 4712710	67 983	10 5287290	9 4911802	74 616	10 0199092	6 516	9 9800908	47
14	2962 638		9 4716785	67 916	10 5283215	9 4916269	74 366	10 0199484	6 533	9 9800516	46
15	2965 418		9 4720856	67 766	10-5279144	9 4920731	74 316	10 0199876	6 533	9 9800124	45
16	2968 197	46 29	9 472 922	67 716	10 527 078	9 4925190	74 260	10 0200468	6 550	9 9799732	44
17	2970 974		9 472 898	67 633	10-5271015	9 4929646	74 183	10 0200851	6 550	9 9799340	43
18	2973 749		9 473043	67 566	10-5266957	9 4934067	74 133	10 0201434	6 566	9 9798946	42
19	2976 526		9 4737097	67 483	10-5262903	9 4938545	74 050	10 0201858	6 566	9 9798552	41
20	2979 303	46 28	9 4741146	67 433	10-5258854	9 4942981	74 016	10 0202346	6 566	9 9798158	40
21	2982 079		9 4745192	67 366	10-5254800	9 49474 9	74 060	10 0202836	6 583	9 9797764	39
22	2984 856		9 4749234	67 303	10-5250756	9 4951965	73 983	10 0203322	6 600	9 9797369	38
23	2987 632		9 4753271	67 216	10-5246729	9 4956438	73 883	10 0203807	6 583	9 9796973	37
24	2990 408	46 26	9 4757304	67 166	10 5242696	9 4960927	73 816	10 0204292	6 600	9 9796578	36
25	2993 184		9 4761334	67 083	10-5238666	9 4965412	73 700	10 0204818	6 616	9 9796182	35
26	2995 959	46 25	9 4765359	67 016	10-5234641	9 4969897	73 616	10 0205412	6 616	9 9795785	34
27	2998 734		9 4769380	66 933	10-5230620	9 4974391	73 543	10 0206016	6 616	9 9795388	33
28	3001 509		9 4773396	66 833	10 5226604	9 4978866	73 500	10 0206609	6 633	9 9794991	32
29	3004 284	46 23	9 4777404	66 816	10-5222591	9 4983316	73 450	10 0207207	6 633	9 9794593	31
30	3007 058		9 4781418	66 750	10-5218582	9 4987723	73 383	10 0207805	6 650	9 9794195	30
31	3009 832	46 23	9 4785423	66 666	10 5214577	9 4992162	73 333	10 0208404	6 633	9 9793796	29
32	3012 606		9 4789423	66 616	10-5210577	9 4996606	73 266	10 0209002	6 633	9 9793398	28
33	3015 380		9 4793420	66 533	10-5206580	9 5001042	73 200	10 0209600	6 650	9 9792998	27
34	3018 153	46 21	9 4797412	66 483	10-5202588	9 5005484	73 150	10 0210201	6 633	9 9792599	26
35	3020 926		9 4801401	66 400	10-5198599	9 5009920	73 083	10 0210802	6 666	9 9792198	25
36	3023 699	46 22	9 4805385	66 350	10-5194615	9 5014358	73 016	10 0211402	6 683	9 9791798	24
37	3026 471		9 4809366	66 266	10 5190634	9 5018799	72 966	10 0212003	6 683	9 9791397	23
38	3029 244		9 4813342	66 216	10 5186658	9 5023237	72 900	10 0212604	6 700	9 9790996	22
39	3032 016	46 20	9 4817313	66 133	10-5182685	9 5027671	72 850	10 0213206	6 700	9 9790594	21
40	3034 788		9 4821283	66 083	10-5178717	9 5032109	72 800	10 0213808	6 716	9 9790192	20
41	3037 559	46 20	9 4825248	66 000	10 5174752	9 5036549	72 716	10 0214411	6 716	9 9789789	19
42	3040 331		9 4829208	65 950	10 5170792	9 5040982	72 666	10 0215014	6 716	9 9789386	18
43	3043 102		9 4833165	65 866	10-5166835	9 5045412	72 600	10 0215617	6 733	9 9788983	17
44	3045 872	46 19	9 4837117	65 816	10-5162883	9 5049838	72 550	10 0216212	6 733	9 9788579	16
45	3048 643		9 4841066	65 733	10 5158934	9 5054281	72 483	10 0216815	6 750	9 9788175	15
46	3051 413	46 18	9 4845010	65 683	10 5154980	9 5058720	72 433	10 0217418	6 750	9 9787770	14
47	3054 183		9 4848951	65 616	10 5151049	9 5063158	72 366	10 0218023	6 750	9 9787365	13
48	3056 953		9 4852888	65 533	10-5147112	9 5067592	72 316	10 0218628	6 766	9 9786960	12
49	3059 723	46 17	9 4856820	65 483	10 5143180	9 5072027	72 250	10 0219234	6 766	9 9786554	11
50	3062 492		9 4860749	65 416	10 5139251	9 5076460	72 183	10 0219839	6 783	9 9786148	10
51	3065 261	46 15	9 4864674	65 350	10 5135326	9 5080893	72 133	10 0220445	6 783	9 9785741	9
52	3068 030		9 4868595	65 267	10 5131405	9 5085326	72 083	10 0221051	6 783	9 9785334	8
53	3070 798	46 15	9 4872512	65 223	10 5127488	9 5089758	72 033	10 0221657	6 800	9 9784927	7
54	3073 566		9 4876426	65 150	10 5123574	9 5094190	71 950	10 0222264	6 800	9 9784519	6
55	3076 334		9 4880335	65 083	10 5119665	9 5098622	71 916	10 0222870	6 816	9 9784111	5
56	3079 102	46 11	9 4884240	65 000	10 5115760	9 5103054	71 833	10 0223477	6 816	9 9783702	4
57	3081 869		9 4888142	64 966	10 5111858	9 5107486	71 783	10 0224084	6 833	9 9783293	3
58	3084 636		9 4892041	64 900	10 5107960	9 5111915	71 733	10 0224691	6 816	9 9782883	2
59	3087 403	46 10	9 4895934	64 833	10 5104066	9 5116340	71 683	10 0225298	6 850	9 9782474	1
60	3090 170		9 4899824		10 5100176	9 5120766		10 0225905		9 9782063	0

APPENDIX III.

ON THE LOGARITHMIC TABLES OF GONIOMETRIC RATIOS.

1. WHEN the Sines, Cosines, &c., of angles have been determined, their logarithms may be found from the tables of the logarithms of numbers. There are, however, methods by which the logarithms of the Goniometric Ratios can be found independently.

2. To find $l_{10} \sin \theta$, $\sin \theta$ not being given.

$$\sin \theta = \theta \cdot \left(1 - \frac{\theta^2}{\pi^2}\right) \cdot \left(1 - \frac{\theta^2}{2^2\pi^2}\right) \cdot \left(1 - \frac{\theta^2}{3^2\pi^2}\right) \dots \text{Art. 133.}$$

By making $\theta = \frac{m}{n} \cdot \frac{\pi}{2}$, and taking the logarithms of both sides of this equation,

$$\therefore l_{10} \sin \left(\frac{m}{n} \cdot \frac{\pi}{2}\right) = l_{10} \left(\frac{m}{n} \cdot \frac{\pi}{2}\right) + l_{10} \left(1 - \frac{m^2}{2^2 n^2}\right) + l_{10} \left(1 - \frac{m^2}{4^2 n^2}\right) + l_{10} \left(1 - \frac{m^2}{6^2 n^2}\right) + \dots$$

{and $l_{10} \left(1 - \frac{m^2}{4^2 n^2}\right)$, $l_{10} \left(1 - \frac{m^2}{6^2 n^2}\right)$, &c. being expanded by (ii), p. 100}

$$\begin{aligned} &= l_{10} \left(\frac{m}{n} \cdot \frac{\pi}{2}\right) + l_{10} \left(\frac{2^2 n^2 - m^2}{2^2 n^2}\right) \\ &= \frac{1}{l_{10}} \cdot \left(\frac{m^2}{4^2 n^2} + \frac{1}{2} \cdot \frac{m^4}{4^4 n^4} + \frac{1}{3} \cdot \frac{m^6}{4^6 n^6} + \dots\right) \\ &\quad - \frac{1}{l_{10}} \cdot \left(\frac{m^2}{6^2 n^2} + \frac{1}{2} \cdot \frac{m^4}{6^4 n^4} + \frac{1}{3} \cdot \frac{m^6}{6^6 n^6} + \dots\right) \\ &\quad - \frac{1}{l_{10}} \cdot \left(\frac{m^2}{8^2 n^2} + \frac{1}{2} \cdot \frac{m^4}{8^4 n^4} + \frac{1}{3} \cdot \frac{m^6}{8^6 n^6} + \dots\right) \\ &\quad - \&c. \end{aligned}$$

$$\text{Now } l_{10} \left(\frac{m}{n} \cdot \frac{\pi}{2}\right) = l_{10} m + l_{10} \pi - l_{10} n - l_{10} 2,$$

$$\begin{aligned} \text{and } l_{10} \frac{2^2 n^2 - m^2}{2^2 n^2} &= l_{10} \{(2n + m) \cdot (2n - m)\} - l_{10} (2^2 n^2) \\ &= l_{10} (2n + m) + l_{10} (2n - m) - 2 \{l_{10} 2 + l_{10} n\}; \end{aligned}$$

$$\therefore l_{10} \sin \left(\frac{m}{n} \cdot \frac{\pi}{2} \right) = l_{10} m + l_{10} (2n + m) + l_{10} (2n - m) + l_{10} \pi - 3 \{ l_{10} n + l_{10} 2 \}$$

$$- \frac{1}{10} \times \left\{ \begin{aligned} & \left(\frac{1}{4^2} + \frac{1}{6^2} + \frac{1}{8^2} + \dots \right) \frac{m^2}{n^2} \\ & + \frac{1}{2} \left(\frac{1}{4^4} + \frac{1}{6^4} + \frac{1}{8^4} + \dots \right) \frac{m^4}{n^4} \\ & + \frac{1}{3} \left(\frac{1}{4^6} + \frac{1}{6^6} + \frac{1}{8^6} + \dots \right) \frac{m^6}{n^6} \\ & + \&c. \end{aligned} \right\}$$

by giving m and n different values, the logarithmic sines of all angles may be found by this formula.

3. In like manner from the series

$$\cos \theta = \left(1 - \frac{2^2 \theta^2}{\pi^2} \right) \cdot \left(1 - \frac{2^2 \theta^2}{3^2 \pi^2} \right) \cdot \left(1 - \frac{2^2 \theta^2}{5^2 \pi^2} \right) \dots \dots \text{Art. 133,}$$

the following formula is obtained;

$$l_{10} \cos \left(\frac{m}{n} \cdot \frac{\pi}{2} \right) = l_{10} (n + m) + l_{10} (n - m) - 2 l_{10} n$$

$$- \frac{1}{10} \times \left\{ \begin{aligned} & \left(\frac{1}{3^2} + \frac{1}{5^2} + \frac{1}{7^2} + \dots \right) \frac{m^2}{n^2} \\ & + \frac{1}{2} \left(\frac{1}{3^4} + \frac{1}{5^4} + \frac{1}{7^4} + \dots \right) \frac{m^4}{n^4} \\ & + \frac{1}{3} \left(\frac{1}{3^6} + \frac{1}{5^6} + \frac{1}{7^6} + \dots \right) \frac{m^6}{n^6} \\ & + \&c. \end{aligned} \right\}.$$

4. The logarithms of the Sines and Cosines having been thus determined, those of

$$\tan \theta = \frac{\sin \theta}{\cos \theta}, \quad \sec \theta = \frac{1}{\cos \theta}, \quad \cot \theta = \frac{\cos \theta}{\sin \theta}, \quad \operatorname{cosec} \theta = \frac{1}{\sin \theta},$$

may be severally found.

5. Since all Sines and Cosines are, generally, less than 1, their logarithms to base 10 are negative. In order to avoid the inconvenience of printing negative characteristics, the logarithms to base of 10 of *all* Goniometrical Ratios are increased by 10, and the resulting numbers being registered are called "Tables of Logarithmic Sines, Cosines," &c.

Hence if any one of these *tabular logarithmic* quantities be given, by subtracting 10 from it the *real* logarithm of the goniometric ratio may be obtained.

These tabular logarithmic quantities will be indicated by the letter L ; thus the tabular logarithmic sine of A , or $10 + \log_{10} \sin A$ will be written $L \sin A$.

6. The common Logarithmic Tables of Goniometric Ratios are calculated for angles which differ from one another by one minute. If, beside degrees and minutes, the angle contain some seconds, its tabular logarithmic function may, with certain exceptions, be found on the principle proved in the next Article.

7. *The increments of tabular logarithmic sines, &c., of angles vary, except in certain cases, as the increment of the angle.*

Let the angle A receive the increments α'' , and $60''$, successively.

$$\begin{aligned} \text{Then } \sin(A + \alpha'') &= \sin A \left\{ 1 + \frac{\sin(A + \alpha'') - \sin A}{\sin A} \right\} \\ &= \sin A \cdot \left\{ 1 + \frac{\cos A \sin \alpha''}{\sin A} \right\}, \text{ unless } A = 90^\circ \text{ nearly; Art. 59, Cor.} \end{aligned}$$

$$\therefore \log_{10} \sin(A + \alpha'') = \log_{10} \sin A + \log_{10} (1 + \cot A \sin \alpha'');$$

$$\therefore \{10 + \log_{10} \sin(A + \alpha'')\} - \{10 + \log_{10} \sin A\} = \log_{10} (1 + \cot A \sin \alpha'');$$

$$\therefore L \sin(A + \alpha'') - L \sin A$$

$$= \frac{1}{\log_e 10} \cdot \left\{ \cot A \sin \alpha'' - \frac{1}{2} \cot^2 A \sin^2 \alpha'' + \dots \right\} \quad \text{App. I. (14) (iv.)}$$

$$= \frac{1}{\log_e 10} \cdot \cot A \sin \alpha''; \text{ by neglecting the higher powers of } \cot A \cdot \sin \alpha'';$$

which may be done unless $A = 2n \cdot 90^\circ$ nearly.

And writing $60''$ for α'' in this equation, it becomes

$$L \sin(A + 60'') - L \sin A = \frac{1}{\log_e 10} \cdot \cot A \sin 60'';$$

$$\therefore \frac{L \sin(A + \alpha'') - L \sin A}{L \sin(A + 60'') - L \sin A} = \frac{\sin \alpha''}{\sin 60''} = \frac{\alpha}{60}. \quad \text{Art. 106, Cor.}$$

And in an exactly similar manner it may be shewn that for the Cosine, Tangent, &c., of an angle, the increment of the tabular logarithm varies as the increment of the angle, except in those cases mentioned in the Corollaries to Arts. 60, 61, 62.

8. To explain the meaning and use of the columns of differences for one second (Diff. 1''), which are placed after the columns of logarithmic sines, tangents, &c.

If α'' become 1'', the equation arrived at in the last Article becomes,

$$L \sin (A + 1'') - L \sin A = \{L (\sin A + 60'') - L \sin A\} \cdot \frac{1}{60},$$

which is the difference for $L \sin A$ corresponding to one second.

Now, if this quantity be computed and registered, $L (\sin A + \alpha'') - L \sin A$ may be determined, when α is given, by merely multiplying this registered difference by α ; and when $L \sin (A + \alpha'')$ is given, α may be found by dividing $L \sin (A + \alpha'') - L \sin A$ by this difference. For

$$\begin{aligned} L \sin (A + \alpha'') - L \sin A &= \{L \sin (A + 60'') - L \sin A\} \cdot \frac{\alpha}{60} \\ &= \frac{L \sin (A + 60'') - L \sin A}{60} \cdot \alpha, \end{aligned}$$

$$\text{and } \alpha = \frac{L \sin (A + \alpha'') - L \sin A}{\frac{1}{60} \cdot \{L \sin (A + 60'') - L \sin A\}}.$$

$$\text{Thus, } L \sin 17^\circ, 1' = 9.4663483$$

$$L \sin 17^\circ, \quad = 9.4659353$$

$$\therefore L \sin 17^\circ, 1' - L \sin 17^\circ = .0004130$$

Now $\frac{4130}{60} = 68.833$, which is the quantity put down in the Tables as the difference for one second to L sines of angles between 17° and $17^\circ, 1'$.

The significant part of the difference is considered as a whole number, or the *real* difference is multiplied by 10^7 , in order to avoid the necessity of printing the three or four cyphers which in nearly every case precede the significant part of the difference*.

9. Ex. 1. To find $L \sin 17^\circ, 14', 12''$.

$$L \sin 17^\circ, 14', \quad = 9.4716785$$

Now Diff. for 1'' = 67.850

$$\therefore \text{Diff. for } 12'' = 814.200 = \quad 814.2$$

$$\therefore L \sin 17^\circ, 14', 12'' = 9.4717599.$$

* The column of differences for the natural sines, &c. of angles are computed after a manner similar to this, and the differences themselves are all multiplied by 1000, to avoid the necessity of printing the cyphers immediately following the decimal point.

Ex. 2. If $L \sin A = 9.4685537$, required A .

$$L \sin A = 9.4685537$$

$$L \sin 17^{\circ}, 6' = 9.4684069$$

$$\text{Diff.} = \quad 1468$$

Now *Diff. for 1''* is in this case 68.400 , and $\frac{1468}{68.400} = 21.46$; $\therefore A = 17^{\circ}, 6', 21''.46$.

Ex. 3. If $L \cos A = 9.9784328$, required A .

In this case, because the increase of the angle is attended by the decrease of the L cosine, Art. 60, Cor. 2, the given L cosine must be *subtracted* from that in the tables which is next greater than it.

$$\text{Now } L \cos 17^{\circ}, 15' = 9.9784519$$

$$L \cos A = 9.9784328$$

$$\text{Diff.} = \quad 191$$

Now *Diff. for 1''* in this case is 6.800 , and $\frac{191}{6.800} = 28.088 = 28.09$ nearly; and the angle required is $17^{\circ}, 54', 28''.09$.

Ex. 4. Required the L cosine of $72^{\circ}, 5', 8''$.

By the Tables, p. 117,

$$L \cos 72^{\circ}, 5' = 9.4880335, \text{ and } \text{Diff. for } 1'' = 65.15;$$

$$\therefore \text{Diff. for } 8'' = \quad 521.2$$

$$\therefore L \cos 72^{\circ}, 5', 8'' = 9.4879814$$

the difference for the additional seconds being in this case *subtracted* from $L \cos 72^{\circ}, 5'$. Art. 60.

NOTE. It may here be observed, that the difference for additional seconds must be *added* for L sines, L tangents, and L secants, Arts. 59, 61, 62; and *subtracted* for L cosines, Art. 60, L cotangents, and L cosecants.

10. To shew that the same columns of "Differences for 1''" serve for $L \sin A$ and $L \text{ cosec } A$, for $L \cos A$ and $L \text{ sec } A$, and for $L \tan A$ and $L \text{ cot } A$.

$$\text{For } \sin A = \frac{1}{\text{cosec } A}; \quad \therefore 1_{10} \sin A = -1_{10} \text{ cosec } A,$$

$$\therefore L \sin A, = 10 + 1_{10} \sin A, = 20 - (10 + 1_{10} \text{ cosec } A) = 20 - L \text{ cosec } A.$$

$$\text{Similarly, } L \sin (A + 1'') = 20 - L \text{ cosec } (A + 1'');$$

$$\therefore L \sin (A + 1'') - L \sin A = -\{L \text{ cosec } (A + 1'') - L \text{ cosec } A\}.$$

Hence a column of "differences for 1''" is printed *between* the column of logarithmic sines and cosecants; serving to the former as a column of *increments* for 1'', and to the latter as a column of *decrements* for 1''.

In like manner it may be shewn that

$$L \cos (A + 1'') - L \cos A = - \{L \sec (A + 1'') - L \sec A\}$$

$$L \tan (A + 1'') - L \tan A = - \{L \cot (A + 1'') - L \cot A\}.$$

Wherefore the columns of cosines and secants have the same differences for 1'', as also have the tangents and cotangents: and it is to be observed that these differences serve respectively as increments to the secants (Art. 61), and to the tangents (Art. 62), and as decrements to the cosines (Art. 60), and to the cotangents.

11. Before the increment of the tabular logarithm of a Goniometrical Ratio can be determined from the *small* given increment of the angle, or conversely, these two conditions must be fulfilled;

I. The logarithmic increment must in that particular case vary as the increment of the angle;

II. The increment of the logarithm must not be an exceedingly small quantity.

Thus, if it were required to determine $L \sin 89^\circ, 40', 3''$, from Tables in which L sines were registered for all angles from 0° to 90° which differed from one another by 1', it would be found that,

$$L \sin 89^\circ, 41' = 9.9999934$$

$$L \sin 89^\circ, 40' = 9.9999927$$

$$\therefore \text{Difference for } 60'' = \quad 7$$

Wherefore, if even the first of these conditions held for the L sines of angles about $89^\circ, 40'$ in magnitude (which it does not, Art. 59, Cor.), yet a difference of 1 in the L sine would produce a difference of $\frac{60''}{7}$, or $9''$ nearly, in the angle; and therefore any increment of $89^\circ, 40'$ which was not greater than $8''$, would produce no change at all in the first seven figures following the decimal point of $L \sin 89^\circ, 40'$.

12. To determine the degree of accuracy to which additional seconds may be calculated in a given case by assuming that the increase of the angle is proportional to the increase of some logarithmic function of the angle.

Let n be the number of seconds by which a difference of unity is produced in a certain logarithmic function of a given angle, and let l be the difference for $60''$;—the differences being considered in both cases as whole numbers;—

$$\text{Then } \frac{n}{60} = \frac{1}{l}, \text{ and } n = \frac{60}{l}.$$

The quantity $\frac{60}{l}$ therefore gives the number of additional seconds corresponding to the least possible increase of the given logarithmic function of the angle, and is consequently the measure of the degree of accuracy to which small increments of the angle may be calculated on the principle that the increments of the angle vary as the increments of the logarithm of some particular Goniometric function of it.

13. It has been observed, App. III. 5, that the *real* logarithm of a Goniometrical quantity is obtained by subtracting 10 from the tabular logarithm. It is therefore necessary

To establish a general rule for supplying the *TENS* when the tabular logarithms of goniometrical quantities are used.

Let $\text{Cos}^n A = a \cdot \sin^m B \cdot \tan^p C$ be any trigonometrical formula adapted to logarithmic computation, App. I. 25.

$$\text{Then } n \cdot \text{I}_{10} \cos A = \text{I}_{10} a + m \cdot \text{I}_{10} \sin B + p \cdot \text{I}_{10} \tan C;$$

$$\therefore n \cdot (10 + \text{I}_{10} \cos A) - n \cdot 10 = \text{I}_{10} a + m \cdot (10 + \text{I}_{10} \sin B) - m \cdot 10 \\ + p \cdot (10 + \text{I}_{10} \tan C) - p \cdot 10;$$

$$\therefore n \cdot L \cos A = \text{I}_{10} a + m \cdot L \sin B + p \cdot L \tan C + \{n - (m + p)\} \cdot 10.$$

And here n, m, p may be whole or fractional.

Whence the *RULE*; *Add to the second member of the equation as many tens as the number of times the tabular logarithms of the goniometrical ratios have been taken in the former member of the equation exceeds the number of times they have been taken in the latter member.*

Ex. 1. $\tan^5 A = \csc B \cdot \cos^2 C$;

$$\therefore 5 \cdot L \tan A = L \cos B + 2 \cdot L \cos C + \{5 - (1 + 2)\} \cdot 10 = L \cos B + 2 \cdot L \cos C + 20.$$

Ex. 2. $\tan^2 A \cdot \sin^6 B = \frac{a}{b} \cdot \sec^4 C$,

$$\begin{aligned} 2 \cdot L \tan A + 6 \cdot L \sin B &= 1_{10} a - 1_{10} b + 4 \cdot L \sec C + \{(2 + 6) - 4\} \cdot 10 \\ &= 1_{10} a - 1_{10} b + 4 \cdot L \sec C + 40. \end{aligned}$$

Ex. 3. $\tan^3 A = \frac{2 \sin^5 B}{\cos^3 C}$;

$$\begin{aligned} \therefore \frac{3}{2} L \tan A &= 1_{10} 2 + \frac{5}{2} L \sin B - \frac{2}{3} L \cos C + \left\{ \frac{3}{2} - \left(\frac{5}{2} - \frac{2}{3} \right) \right\} 10 \\ &= 1_{10} 2 + \frac{5}{2} L \sin B - \frac{2}{3} L \cos C - \frac{1}{3} \cdot 10. \end{aligned}$$

[Had both sides of this equation been raised to the sixth power, the fractional indices would have disappeared, and the value of $\tan A$ would have been practically determined much more easily.]

14. Lastly, the methods will be explained by which small angles are determined from their L sines, and conversely.

When an angle receives a small increment, the Differential Calculus affords the means of determining with facility the consequent increase of the Goniometric Ratio.

Required the increase of $L \sin \theta$ arising from θ receiving a small increment $\delta \theta$.

By TAYLOR'S THEOREM,

$$L \sin (\theta + \delta \theta) = L \sin \theta + d_{\theta} L \sin \theta \cdot \delta \theta + d^2_{\theta} L \sin \theta \cdot \frac{(\delta \theta)^2}{1 \cdot 2} + \dots$$

$$\text{Now } L \sin \theta = 10 + 1_{10} \sin \theta;$$

$$\therefore d_{\theta} L \sin \theta = \frac{1}{1_{\epsilon} 10} \cdot \frac{\cos \theta}{\sin \theta} = \frac{1}{1_{\epsilon} 10} \cdot \cot \theta,$$

$$\text{whence, } d^2_{\theta} L \sin \theta = -\frac{1}{1_{\epsilon} 10} \cdot \operatorname{cosec}^2 \theta;$$

$$\therefore L \sin (\theta + \delta \theta) - L \sin \theta = \frac{1}{1_{\epsilon} 10} \cdot \cot \theta \cdot \delta \theta - \frac{1}{1_{\epsilon} 10} \cdot \operatorname{cosec}^2 \theta \cdot \frac{(\delta \theta)^2}{1 \cdot 2} + \dots$$

15. Now if θ be very small, $\operatorname{cosec} \theta$ is very large, and therefore (unless $\delta\theta$ be exceedingly small also) the second difference, $-\frac{1}{1_e 10} \cdot \operatorname{cosec}^2 \theta \cdot \frac{(\delta\theta)^2}{1 \cdot 2}$, is of such a magnitude that it cannot be neglected in comparison with the first difference, $\frac{1}{1_e 10} \cdot \cot \theta \cdot \delta\theta$.

In this case, therefore, since the increment of $L \sin \theta$ does not vary as $\delta\theta$, the simple power of the increment of θ , the quantity to be added to $L \sin \theta$ for a small increment of θ cannot be obtained by a simple proportion, but will have to be determined approximately by the tedious process of computing the first two terms of the series which gives the value of the increment of $L \sin \theta$.

There are three methods of escaping this inconvenience.

16. *The first method* is to construct tables for the first few degrees to intervals of a *second*, instead of to intervals of a *minute* as the tables in ordinary use are constructed.

Here $\delta\theta$ is less than one second, and $L \sin(\theta + \delta\theta)$ may be roughly computed to decimal parts of a second by neglecting the second term of the series obtained in the last Article but one, in which case,

$$L \sin(\theta + \delta\theta) - L \sin \theta = \frac{1}{1_e 10} \cdot \cot \theta \cdot \delta\theta;$$

Therefore, for any particular value of θ , The increment of $L \sin \theta \propto \delta\theta$, and

$$\frac{\text{Increment of } L \sin \theta \text{ for } \delta\theta''}{\text{Increment of } L \sin \theta \text{ for } 60''} = \frac{\delta\theta}{60}.$$

17. *Second method.* By the following formulæ, which are given by Maskelyne in his introduction to Taylor's Logarithms, a small angle may be determined very accurately to decimal parts of a second from its L sine and conversely, by the aid of tables of L sines and L cosines which are calculated to every second for a few degrees,

$$\sin \theta = \theta - \frac{\theta^3}{2 \cdot 3} + \dots = \theta - \frac{\theta^3}{2 \cdot 3} \text{ nearly.}$$

$$\cos \theta = 1 - \frac{\theta^2}{1 \cdot 2} + \dots = 1 - \frac{\theta^2}{2} \text{ nearly.}$$

$$\therefore \frac{\sin \theta}{\theta} = 1 - \frac{\theta^2}{2 \cdot 3} = \left(1 - \frac{\theta^2}{2}\right)^{\frac{1}{2}} \text{ nearly,} = \cos^{\frac{1}{2}} \theta.$$

Let θ be an angle containing n'' , then $n = \frac{\theta}{\sin 1''}$, or $\theta = n \cdot \sin 1''$;

$$\text{and } \therefore \sin \theta = n \cdot \sin 1'' \cdot \cos^{\frac{1}{2}} \theta.$$

Writing n'' for θ , and taking the logarithms,

$$L \sin n'' = l_{10} n + L \sin 1'' + \frac{1}{3} L \cos n'' - \frac{1}{3} \cdot 10;$$

$$\therefore L \sin n'' = l_{10} n + L \sin 1'' - \frac{1}{3} \cdot (10 - L \cos n'') \dots\dots\dots (i.),$$

$$\text{and } l_{10} n = L \sin n'' + \frac{1}{3} (10 - L \cos n'') - L \sin 1'' \dots\dots\dots (ii.)$$

DEF. The quantity $10 - L \cos n''$ is called "the arithmetic complement" of $L \cos n''$.

18. It is to be remarked, that in using these formulæ to determine $L \sin n''$ when n is given, or conversely, an *approximate* value of $L \cos n''$ may be taken from the tables and written in the second member of the equations without sensibly affecting the result, because the variation of $L \cos n''$ is exceedingly small when n is small, as may easily be shewn*.

The whole matter will be rendered more clear by an example.

Ex. If $L \sin n'' = 7.3217783$, required n .

Taylor's tables give,

$L \sin 7', 12''$	7.3210583		$L \cos 7', 12''$	9.9999990
$L \sin 7', 13''$	7.3220624		$L \cos 7', 13''$	9.9999990

Therefore the angle is $7', 12''$ nearly; and $7', 12''$ is the approximate value of the angle which must be taken in the second member of (ii.) of the last Article.

Now, by (ii.),

$$L \sin n'' = 7.3217783$$

$$\text{or } \left. \begin{array}{l} \frac{1}{3} (10 - L \cos 7', 12'') \\ \frac{1}{3} \times .0000010, \end{array} \right\} = \frac{.0000003}{.0000003}$$

$$\underline{7.3217786}$$

$$L \sin 1'' = 4.6855749$$

$$\therefore l_{10} n = 2.6362037 = l_{10} 432.717;$$

$$\therefore n = 432.717; \text{ or the angle required is } 7', 12''.717.$$

$$* \cos \theta = 1 - \frac{\theta^2}{1.2} + \frac{\theta^4}{1.2.3.4} \dots\dots = 1 - \frac{\theta^2}{2} \text{ nearly;}$$

$$\therefore l_{10} \cos \theta, \text{ or } L \cos \theta - 10, = -\frac{1}{1_e 10} \cdot \left\{ \frac{\theta^2}{2} + \frac{1}{2} \left(\frac{\theta^2}{2} \right)^2 + \dots \right\};$$

\therefore , by the Differential Calculus, $d_{\theta} (L \cos \theta - 10) = -\frac{1}{1_e 10} \left(\theta + \frac{\theta^3}{2} + \dots \right)$, a very small quantity, if θ be an angle of a few minutes only.

19. In like manner a formula may be established for finding $L \tan n''$ from n , and conversely.

$$\tan \theta = \frac{\sin \theta}{\cos \theta} = \frac{\theta - \frac{\theta^3}{2 \cdot 3}}{1 - \frac{\theta^2}{2}} \text{ nearly;}$$

$$\therefore \frac{\tan \theta}{\theta} = \frac{1 - \frac{\theta^2}{2 \cdot 3}}{1 - \frac{\theta^2}{2}} = \frac{\left(1 - \frac{\theta^2}{2}\right)^{\frac{1}{2}}}{1 - \frac{\theta^2}{2}} = \left(1 - \frac{\theta^2}{2}\right)^{-\frac{1}{2}} = (\cos \theta)^{-\frac{1}{2}};$$

$$\therefore \frac{\tan n''}{n \cdot \sin 1''} = (\cos n'')^{-\frac{1}{2}};$$

$$\therefore L \tan n'' = 1_{10} n + L \sin 1'' - \frac{2}{3} \cdot L \cos n'' + \frac{2}{3} \cdot 10;$$

$$\therefore L \tan n'' = 1_{10} n + L \sin 1'' + \frac{2}{3} (10 - L \cos n'') \dots\dots\dots(iii.)$$

$$\text{And } 1_{10} n = L \tan n'' - L \sin 1'' - \frac{2}{3} (10 - L \cos n'') \dots\dots\dots (iv.)$$

20. *Delambre's Tables.* The third method alluded to, (15, p. 126,) is to construct tables as far as an angle of one degree, which give $1_{10} \frac{\sin \theta}{\theta}$, (or tables which give $1_{10} \frac{\sin \theta}{\theta} + L \sin 1''$), for every second.

Such tables are printed in no collection, perhaps, except those of Callet; they may be easily constructed in the following manner:

Let θ be an angle of n seconds; $\therefore \theta = n \cdot \sin 1''$.

$$\begin{aligned} \text{Then } 1_{10} \frac{\sin \theta}{\theta} &= 1_{10} \frac{\sin n''}{n \cdot \sin 1''} = 1_{10} \sin n'' - 1_{10} n - 1_{10} \sin 1'' \\ &= L \sin n'' - 1_{10} n - L \sin 1''; \end{aligned}$$

$$\therefore 1_{10} \frac{\sin \theta}{\theta} + L \sin 1'' = L \sin n'' - 1_{10} n.$$

Similarly, if θ be an angle of n minutes,

$$1_{10} \frac{\sin \theta}{\theta} + L \sin 1' = L \sin n' - 1_{10} n.$$

21. To determine the Sine of a given small angle, or conversely, from Delambre's Tables.

Since $\sin \theta = \frac{\sin \theta}{\theta} \cdot \theta = \frac{\sin \theta}{\theta} \cdot n \cdot \sin 1''$;

$\therefore L \sin \theta = (l_{10} \frac{\sin \theta}{\theta} + L \sin 1'') + l_{10} n$;

$\therefore L \sin n'' = (l_{10} \frac{\sin \theta}{\theta} + L \sin 1'') + l_{10} n \dots\dots\dots (i).$

And $l_{10} n = L \sin n'' - (l_{10} \frac{\sin \theta}{\theta} + L \sin 1'') \dots\dots\dots (ii).$

The most convenient tables are evidently those which give $(l_{10} \frac{\sin \theta}{\theta} + L \sin 1'')$ for every second.

22. Ex. 1. To determine $L \sin n''$ by Delambre's Tables.

Since, as shewn below*, $l_{10} \frac{\sin \theta}{\theta} + L \sin 1''$ increases very slowly as θ increases, the value of $L \sin n''$ is obtained without sensible error by taking for $l_{10} \frac{\sin \theta}{\theta} + L \sin 1''$ the quantity which is given in the tables for the angle containing that number of seconds which is the nearest integer to the given number (n) and adding $l_{10}n$ to it.

Thus; If $n = 546 \cdot 25$, required $L \sin 546'' \cdot 25$.

By Taylor's Tables. $L \sin 546''$, or $L \sin 9', 6''$, is $7 \cdot 4227670$;
and $l_{10} 546 = 2 \cdot 7371926$.

Therefore, when θ is an angle of $546''$,

$$l_{10} \frac{\sin \theta}{\theta} + L \sin 1'' = L \sin n'' - l_{10} n. \quad \text{Art. 20: p. 128.}$$

$$= 7 \cdot 4227670 - 2 \cdot 7371926 = 4 \cdot 6855744,$$

* For $\frac{\sin \theta}{\theta} = 1 - \frac{\theta^2}{1 \cdot 2 \cdot 3} + \frac{\theta^4}{1 \cdot 2 \cdot 3 \cdot 4 \cdot 5} - \dots = 1 - \frac{\theta^2}{6}$ nearly;

$\therefore l_{10} \frac{\sin \theta}{\theta} = -\frac{1}{l_{10}} \cdot \left\{ \frac{\theta^2}{6} + \frac{1}{2} \left(\frac{\theta^2}{6} \right)^2 + \dots \right\}$;

\therefore , by the Differential Calculus, $d_{\theta} l_{10} \frac{\sin \theta}{\theta} = -\frac{1}{l_{10}} \cdot \left(\frac{\theta}{3} + \frac{2\theta^3}{6^2} + \dots \right)$,

a very small quantity, if θ be an angle of a few minutes only.

the quantity corresponding to the angle $546''$ which would be given in the tables of $1_{10} \frac{\sin \theta}{\theta} + L \sin 1''$.

∴ By (i) of the last Article,

$$\begin{aligned} L \sin 564'' \cdot 25 &= 4 \cdot 6855744 + 1_{10} 564 \cdot 25 \\ &= 4 \cdot 6855744 + 2 \cdot 7373914 \\ &= 7 \cdot 4229658. \end{aligned}$$

Ex. 2. *To determine n when L sin n'' is given.*

By referring to the tables of L sines in common use the integral number of seconds is found which is contained in the angle whose L sine is next below the proposed quantity: suppose this number to be m . Then substituting in (ii) of the last Article, the quantity given in the tables for that value of $1_{10} \frac{\sin \theta}{\theta} + L \sin 1''$ which corresponds to the angle m'' , a near approximation to the value of n is obtained.

Thus, If $L \sin n'' = 7 \cdot 4230612$, required n .

By Taylor's Tables,

$$L \sin 546'' = 7 \cdot 4227670, \text{ and } L \sin 547'' = 7 \cdot 4235617;$$

∴ the value of m in this case is 546.

Now, as in the last example, the value of $1_{10} \frac{\sin \theta}{\theta} + L \sin 1''$ for an angle $564''$ is $4 \cdot 6855744$.

Therefore by (ii) of the last Article,

$$\begin{aligned} 1_{10} n &= 7 \cdot 4230612 - 4 \cdot 6855744 = 2 \cdot 7374868 = 1_{10} 564 \cdot 37; \\ &\text{and } \therefore 546'' \cdot 37 \text{ is the angle sought.} \end{aligned}$$

APPENDIX IV.


THE GENERAL PROOF OF THE FORMULÆ

$$\sin(A \pm B) = \sin A \cdot \cos B \pm \cos A \cdot \sin B,$$

$$\cos(A \pm B) = \cos A \cdot \cos B \mp \sin A \cdot \sin B.$$

On the Theory of Projections.

1. Let $X'OX$ be an indefinite straight line, and PQ a finite straight line from which PM , QN are drawn perpendicular to $X'OX$; the length MN intercepted on $X'OX$ between these perpendiculars is termed the *Orthogonal Projection** (or simply the "projection") of PQ on $X'OX$, and $X'OX$ is termed the *Line of Projection*.




Let lengths on $X'OX$ be reckoned positive if measured in the direction OX , and negative in the opposite direction OX' ; and let the sign of the projection of any line on $X'OX$ be determined in accordance with this convention. Then in the above figure the projection of PQ (being MN , measured from M to N) will be positive; while the projection of QP (being measured from N to M) will be negative.

2. From this explanation of projections it will be seen that the projection of a line on a line parallel to it is equal to the line itself,—that the projection of a line on a line perpendicular to it is zero,—and that the projections of two equal and parallel lines, taken in the same direction, are equal in magnitude and of the same sign.

* An *orthogonal projection* is made by means of straight lines that are *perpendicular* to the line of projection. There are other modes of projecting a line on another line, as for instance, by means of lines drawn from a fixed point through the extremities of the first line.

3. If two points be joined by any series of straight lines, the algebraical sum of the projections of the lines taken in order from one point to the other is constant.


Thus, let AB, BC, CD, DE, EF be any series of straight lines commencing at A and terminating at F . Their projections are MN, NP, PQ, QR, RS , of which the algebraical sum
 $= MN + NP - QP + QR - SR$
 $= MS$, the projection of AF .



This is a proposition of very frequent application; and it may be remarked that it will be equally true if the straight lines do not all lie in the same plane. It is sometimes convenient to express this result by saying that *the sum of the projections on any straight line of the sides taken in order, of a closed polygon, is zero.*

4. If PR (Fig. to Art. 1) be drawn parallel to $X'OX$, it will be seen that $MN = PR = PQ \cdot \cos QPR$; and therefore the length of the projection of a line is obtained by multiplying the length of the line by the cosine of its inclination to the line of projection. The same will be true with regard to *sign* as well as *magnitude*, if the following conventions be made.


5. From a fixed point O in $X'OX$ draw a line parallel to PQ , the line to be projected, and in the same direction as PQ ; and let the angle which this line makes with the positive portion (OX) of the line $X'OX$ be considered as the inclination of PQ to the line of projection; then will the sign of the projection be the same as that of the cosine of the inclination.



Thus the inclination of PQ to $X'OX$ is XOT , an angle whose terminal line lies in the first quadrant, and whose cosine therefore is positive, as is also MN the projection of PQ . But the inclination of QP is XOT' , an angle whose terminal line lies in the

third quadrant, and whose cosine therefore is negative, which is also the case with NM , the projection of QP .

Again, in the annexed figure, where OT , OT' lie in the fourth and second quadrants respectively, the same will be seen to be true.




6. The line to be projected has not in the preceding Articles been considered as affected by any sign*; but if it lie on a line on which the positive and negative directions are assigned, the inclination of the line to be projected must be taken to be the inclination of the *positive side* of the line on which it lies to the *positive side* of the line of projection.

Thus, if PQ be considered as negative, (that is, if PQ lie on a line on which lengths measured in the same direction as from P towards Q are taken as negative), its inclination to the line of projection must be considered as XOT' in both the figures to Art. 5; and the cosine being in both cases negative, $PQ \times$ cosine of the inclination will be positive as before.

7. From the preceding considerations, a perfectly GENERAL PROOF may be derived of the formulæ which give the values of $\text{Sin}(A \pm B)$ and $\text{Cos}(A \pm B)$ in terms of the Sines and Cosines of the simple angles.

Suppose a pair of rectangular axes $x'Ox$, $y'Oy$ to be originally coincident with the axes XOX , $Y'OY$, having their positive sides Ox , Oy coincident with the positive sides OX , OY ; and then suppose them to revolve through any angle A , in the positive or negative direction of revolution



according as A is a positive or negative angle. Thus, in the figure, Ox and Oy have revolved from OX , OY through an angle

* A distinction has been made throughout between the lines PQ and QP , but we have not before supposed that distinction has not been supposed to be expressed by means of the algebraical signs $+$ and $-$.

in the positive direction greater than two right angles and less than three, or through an angle in the negative direction greater than one right angle and less than two.

Also, suppose a line OP , initially coincident with Ox , to revolve through any angle B , measured from Ox in a positive or negative direction according as B is positive or negative. Thus, in the figure, OP has revolved through a positive angle xOP that is greater than one right angle and less than two, or through a negative angle greater than two right angles and less than three.

Let ON , NP be the projections of OP on $x'Ox$, $y'Oy$;

$\therefore ON = OP \cdot \cos B$, and $NP = OP \cdot \cos (B - 90^\circ) = OP \cdot \sin B$;
as well with regard to sign as to magnitude (Art. 5).

Now projecting OP , and also the lines ON , NP on $X'OX$, the projection of the first will be equal to the algebraic sum of the projections of the other two (Art. 3).

But the angle between OP and OX (the positive part of $X'OX$, is $A + B$),

\therefore the projection of OP on $X'OX = OP \cdot \cos (A + B)$.

Also, by Art. 6,

the projection of ON on $X'OX$

$$= ON \cdot \cos A = OP \cdot \cos A \cdot \cos B;$$

and the projection of NP on $X'OX$

$$= NP \cdot \cos (A + 90^\circ) = -OP \cdot \sin A \cdot \sin B.$$

Therefore,

$$OP \cdot \cos (A + B) = OP \cdot \cos A \cdot \cos B - OP \cdot \sin A \cdot \sin B,$$

$$\text{or } \cos (A + B) = \cos A \cdot \cos B - \sin A \cdot \sin B;$$

for all values of A and B , positive or negative.

8. The formula for $\sin (A + B)$ may be found in like manner, by projecting on $Y'OY$, but it may be derived from the formula just proved by writing in it $90^\circ + A$ for A , when

$$\cos (90^\circ + A + B) = \cos (90^\circ + A) \cdot \cos B - \sin (90^\circ + A) \cdot \sin B,$$

$$\therefore -\sin (A + B) = -\sin A \cdot \cos B - \cos A \cdot \sin B,$$

$$\therefore \sin (A + B) = \sin A \cdot \cos B + \cos A \cdot \sin B.$$

9. The demonstration given above is due to Professor De Morgan: see Chap. III. of his "Trigonometry and Double Algebra." And it ought to be remarked in his words, that "it can be convincing only to those who enable themselves to understand in the most general sense the preliminary theorems. Any want of such mastery over the universal character of theorems in projection will follow the student through all his course, particularly in the higher Geometry and in Mechanics."

10. EXAMPLES. (1) Shew, by projecting the sides of a regular polygon on any line, that

$$\cos \theta + \cos \left(\theta + \frac{2\pi}{n} \right) + \dots + \cos \left\{ \theta + \frac{2(n-1)\pi}{n} \right\} = 0,$$

$$\sin \theta + \sin \left(\theta + \frac{2\pi}{n} \right) + \dots + \sin \left\{ \theta + \frac{2(n-1)\pi}{n} \right\} = 0,$$

whatever be the value of θ , n being an integer.

(2) If x , y , and x' , y' be adjacent sides of two parallelograms described on the same diagonal, shew that

$$y \cdot \sin \widehat{yx} = x' \cdot \sin \widehat{x'x} + y' \cdot \sin \widehat{y'x},$$

$$x \cdot \sin \widehat{xy} = x' \cdot \sin \widehat{x'y} + y' \cdot \sin \widehat{y'y};$$

where \widehat{yx} denotes the angle which y makes with x , and so for the other expressions $\widehat{x'x}$, &c.

[These are the general formulæ for the transformation of co-ordinates in Analytical Geometry.]

(3) If the inclinations of the sides a , b , c of a triangle, taken in order, to a given line be α , β , γ respectively, prove that

$$a \cdot \cos \alpha + b \cdot \cos \beta + c \cdot \cos \gamma = 0,$$

and that $a \cdot \cos (\beta + \gamma) + b \cdot \cos (\gamma + \alpha) + c \cdot \cos (\alpha + \beta) = 0$.

EXAMPLES.

I. PROVE that $45^{\circ}, 15', 20'' = 50^{\circ}, 28', 39'' \cdot 50$; $10^{\circ}, 15', 37'' = 11^{\circ}, 40', 3'' \cdot 09$; $18^{\circ}, 10', 48'' = 20^{\circ}, 20'$; $\frac{180^{\circ}}{\sqrt{3}} = 115^{\circ}, 47'$.

II. The Complements of $17^{\circ}, 36', 43''$; $29^{\circ}, 27', 6'' \cdot 32$; and $216^{\circ}, 45'$; are $72^{\circ}, 23', 17''$; $60^{\circ}, 32', 53'' \cdot 68$; and $-126^{\circ}, 45'$.

III. The Supplements of $37^{\circ}, 4', 3''$; $115^{\circ}, 13', 24'' \cdot 66$; and $226^{\circ}, 14', 17''$; are $142^{\circ}, 55', 57''$; $64^{\circ}, 46', 35'' \cdot 34$; and $-(46^{\circ}, 14', 17'')$.

IV. 1. If $\text{Cot } A = \frac{3}{2}$, find the values of $\text{Sin } A$, $\text{Cos } A$, $\text{Cosec } A$, $\text{Versin } A$, and $\text{Sec } A$.

2. What angles have the same sine as 320° ? *Ans.* The form $2m \cdot 180^{\circ} + A$...see Art. 24. (1),...gives $320^{\circ}, 680^{\circ}, 1040^{\circ}$...for the values 0, 1, 2...of m , and $-40^{\circ}, -400^{\circ}, -760^{\circ}$...for the values -1, -2, -3,...of m . The form $(2m + 1) 180^{\circ} - A$, see Art. 24. (4), gives $-140^{\circ}, 220^{\circ}, 580^{\circ}$...for the values 0, 1, 2,...of m ; and $-500^{\circ}, -860^{\circ}, -1220^{\circ}$...for the values -1, -2, -3,...of m .

3. What are the angles that have their tangents of the same magnitude as that of -110° , but affected with a different sign? *Ans.* $110^{\circ}, 290^{\circ}, 470^{\circ}, 650^{\circ}, \dots -70^{\circ}, -250^{\circ}, -430^{\circ}, \dots$ The angles are comprised under the form (see Art. 25) $m \cdot 180^{\circ} + 110^{\circ}$, where m is 0 or any positive or negative integer.

V. Prove the formulæ,

1. $\text{Sec}^2 A \text{ cosec}^2 A = \text{sec}^2 A + \text{cosec}^2 A$.

2. $\text{Cot}^2 A \cos^2 A = \text{cot}^2 A - \cos^2 A$.

3. $\text{Cos } A = \frac{\cot A}{\sqrt{(1 + \cot^2 A)}}$ 4. $\text{Versin } A = \frac{\sec A - 1}{\sec A}$.

5. $\text{Sin } A \cos A = \frac{1}{\tan A + \cot A}$.

VI. 1. If $\tan^2 A + 4 \sin^2 A = 6$; $A = 60^\circ$.

2. If $m = \tan A + \sin A$, and $n = \tan A - \sin A$; $\cos A = \frac{m-n}{m+n}$.

3. If $m \sin A = n \cos A$; $\sin A = \pm \frac{n}{\sqrt{m^2 + n^2}}$.

4. If $1 = \left(\frac{\sin A}{\sin B}\right)^2 + (\cos A \cos C)^2$; $\sin C = \frac{\tan A}{\tan B}$

5. If $\cos x = \frac{\cos A}{\sin C}$, and $\cos(90^\circ - x) = \frac{\cos B}{\sin C}$;

then $\cos^2 A + \cos^2 B + \cos^2 C = 1$.

VII. Prove the following formulae;

1. $\tan A + \cot A = 2 \operatorname{cosec} 2A$. 2. $\cot A - \tan A = 2 \cot 2A$.

3. $\tan A = \frac{\sin 2A}{1 + \cos 2A}$. 4. $\sqrt{\frac{1 + \sin A}{1 - \sin A}} = \frac{1 + \tan \frac{1}{2} A}{1 - \tan \frac{1}{2} A}$.

5. $\operatorname{Cosec} 2A + \cot 2A = \cot A$. 6. $2 \operatorname{cosec} 2A = \sec A \operatorname{cosec} A$.

7. $\frac{\sin A}{1 - \cos A} = \cot \frac{1}{2} A$. 8. $\frac{\operatorname{Versin} A}{\operatorname{versin}(180^\circ - A)} = \tan^2 \frac{1}{2} A$.

9. $8 \cot 2A \operatorname{cosec}^2 2A = \cot A \operatorname{cosec}^2 A - \tan A \sec^2 A$.

10. $\operatorname{Versin}(180^\circ - A) = 2 \operatorname{vers} \frac{1}{2}(180^\circ + A) \operatorname{vers} \frac{1}{2}(180^\circ - A)$.

11. $\sec 2A = \frac{\cot A + \tan A}{\cot A - \tan A}$. 12. $\tan^2 \frac{1}{2} A = \frac{2 \sin A - \sin 2A}{2 \sin A + \sin 2A}$.

13. $\operatorname{Cosec} 2A = 2 \cdot \frac{\cos 2A + \sin 2A}{(\cos A - \sin A) - (\cos 3A - \sin 3A)}$.

14. $\cos^2 A = (\cos \frac{1}{2} A - \sin \frac{3}{2} A)^2 + 2 \cos \frac{1}{2} A \sin \frac{3}{2} A (\cos \frac{1}{2} A - \sin \frac{1}{2} A)$.

15. $\sqrt{1 + \sin A} = 1 + 2 \sin \frac{1}{4} A \cdot \sqrt{1 - \sin \frac{1}{2} A}$. Shew also that the radicals have their proper signs in this equation if A be between -90° and 180° .

$$16. \quad \cot A + \cot 2A + \cot 4A = \frac{1}{\sin 4A} \cdot (2 + 2 \cos 2A + 3 \cos 4A).$$

$$17. \quad \text{Prove that } \frac{\sec(2^{2n+1}A) - 1}{\sec(2^{2n}A) - 1} = \frac{\tan(2^{2n+1}A)}{\tan(2^{2n}A)};$$

and thence shew that

$$\frac{(\sec 2A - 1)(\sec 2^2A - 1)(\sec 2^3A - 1) \dots \text{to } n \text{ factors}}{(\sec A - 1)(\sec 2^2A - 1)(\sec 2^4A - 1) \dots \text{to } n \text{ factors}} = \cot \frac{1}{2}A \tan(2^{2n-1}A).$$

$$18. \quad \operatorname{Cosec} 2A + \cos 4A = \cot A - \operatorname{cosec} 4A.$$

$$19. \quad \sin 3A \sin^3 A + \cos 3A \cos^3 A = \cos^3 2A.$$

$$20. \quad \frac{\sin nA}{\cos 2nA + \cos A} = \frac{\sec(n + \frac{1}{2})A - \sec(n - \frac{1}{2})A}{4 \sin \frac{1}{2}A};$$

and thence shew that

$$\frac{\sin A}{\cos 2A + \cos A} + \frac{\sin 2A}{\cos 4A + \cos A} + \frac{\sin 3A}{\cos 6A + \cos A} + \dots (\text{to } n \text{ terms}) = \frac{\sin \frac{1}{2}nA \cdot \sin \frac{1}{2}(n+1)A}{\sin A \cdot \cos \frac{1}{2}(2n+1)A}.$$

VIII. Prove that, whatever be the values of the angles,

$$1. \quad \cos 2A + \cos 2B = 2 \cos(A+B) \cos(A-B).$$

$$2. \quad 1 + \cos 2A \cos 2B = 2(\sin^2 A \sin^2 B + \cos^2 A \cos^2 B).$$

$$3. \quad \cos^2(A+B) - \sin^2 A = \cos B \cos(2A+B).$$

$$4. \quad \frac{\sin(A-B)}{\sin A \sin B} + \frac{\sin(B-C)}{\sin B \sin C} + \frac{\sin(C-A)}{\sin C \sin A} = 0.$$

$$5. \quad \cos(A+B) \sin(A-B) + \cos(B+C) \sin(B-C) + \cos(C+D) \sin(C-D) + \cos(D+A) \sin(D-A) = 0.$$

$$6. \quad \cos(A+B) \sin B - \cos(A+C) \sin C = \sin(A+B) \cos B - \sin(A+C) \cos C.$$

7. $\sin(A+B)\cos B - \sin(A+C)\cos C = \sin(B-C)\cos(A+B+C).$

8. $\sin(A+B-2C)\cos B - \sin(A+C-2B)\cos C$
 $= \sin(B-C)\{\cos(B+C-A) + \cos(A+C-B) + \cos(A+B-C)\}.$

9. $\sin A \sin(B-C) + \sin B \sin(C-A) + \sin C \sin(A-B) = 0;$
 and $\cos A \sin(B-C) + \cos B \sin(C-A) + \cos C \sin(A-B) = 0.$

10. $\cos 2A + \cos 2B + \cos 2C$
 $= \cos(B+C)\cos(B-C) + \cos(C+A)\cos(C-A) + \cos(A+B)\cos(A-B);$
 and $\sin 2A + \sin 2B + \sin 2C$
 $= \sin(B+C)\cos(B-C) + \sin(C+A)\cos(C-A) + \sin(A+B)\cos(A-B).$

11. If $2S = A + B + C,$
 $4 \cos A \cos B \cos C = \cos 2(S-A) + \cos 2(S-B) + \cos 2(S-C) + \cos 2S;$
 and $4 \sin A \sin B \sin C$
 $= \sin 2(S-A) + \sin 2(S-B) + \sin 2(S-C) - \sin 2S.$

12. $\sin B \sin(A-B) + \sin C \sin(A-C)$
 $= \cos(B-C)\cos 2(S-A) - \cos A.$

13. $(\sin^2 A + \sin^2 B + \sin^2 C)\{\sin^2(A-B) + \sin^2(B-C) + \sin^2(C-A)\}$
 $= \{\cos A - \cos(B-C)\cos 2(S-A)\}^2 + \{\cos B - \cos(C-A)\cos 2(S-B)\}^2$
 $+ \{\cos C - \cos(A-B)\cos 2(S-C)\}^2;$ where $2S = A + B + C.$

14. $\frac{\sin(A-B)}{\sin C} + \frac{\sin(B-C)}{\sin A} + \frac{\sin(C-A)}{\sin B}$
 $+ \frac{\sin(A-B)\sin(B-C)\sin(C-A)}{\sin A \sin B \sin C} = 0.$

15. $\left(\frac{\tan A}{\tan B} - \frac{\tan B}{\tan D}\right) + \left(\frac{\tan B}{\tan C} - \frac{\tan C}{\tan B}\right) + \left(\frac{\tan C}{\tan A} - \frac{\tan A}{\tan C}\right)$
 $= 2 \cdot \frac{\sin 2(B-A) + \sin 2(C-B) + \sin 2(A-C)}{\sin 2A \sin 2B \sin 2C}.$

16. The formula in the last Example is also equal to
 $\frac{8 \sin(A-B)\sin(B-C)\sin(C-A)}{\sin 2A \sin 2B \sin 2C}.$

IX. Prove that

$$1. \frac{1 + \sin A}{1 - \sin A} = \tan^2(45^\circ + \frac{1}{2}A).$$

$$2. \sec(45^\circ + A) \sec(45^\circ - A) = 2 \sec 2A.$$

$$3. 2 \sec A = \tan(45^\circ + \frac{1}{2}A) + \cot(45^\circ + \frac{1}{2}A).$$

$$4. \tan(30^\circ + A) \tan(30^\circ - A) = \frac{2 \cos 2A - 1}{2 \cos 2A + 1}.$$

$$5. \sin(60^\circ + A) - \sin(60^\circ - A) = \sin A.$$

$$6. \sqrt{(1 - \cos A)} = \frac{\sin A}{\sin(45^\circ - \frac{1}{2}A) + \cos(45^\circ + \frac{1}{2}A)}.$$

$$7. \cos A + \cos(120^\circ - A) + \cos(120^\circ + A) = 0.$$

$$8. \cos^2 A + \cos^2(60^\circ - A) + \cos^2(60^\circ + A) = \frac{3}{2}.$$

$$9. \cos^4 A + \cos^2(72^\circ + A) + \cos^4(2.72^\circ + A) + \cos^4(3.72^\circ + A) \\ + \cos^4(4.72^\circ + A) = \frac{15}{8}.$$

X. Find the numerical values of $\sin 15^\circ$, $\sin 9^\circ$, $\cos 12^\circ$, $\text{Versin } 15^\circ$, $\tan 22^\circ$, $12'$, $\sin 150^\circ$, $\cos 135^\circ$, $\sin 3^\circ$, $\sin 75^\circ$, $\sec 225^\circ$; and prove that

$$1. \tan 50^\circ + \cot 50^\circ = 2 \sec 10^\circ.$$

2. $2 \cos \frac{A}{2^n} = \sqrt{2 + \sqrt{2 + \dots \&c. + \sqrt{(2 + 2 \cos A)}}}$; where n is a positive integer, and the symbol $\sqrt{\quad}$ is repeated n times, each time affecting all the quantities which follow it.

XI. Determine A in the following equations:

$$1. \sin A = \sin 2A.$$

$$2. \tan 2A = 3 \tan A.$$

$$3. \tan A = \text{cosec } 2A.$$

$$4. \cos A = \tan A.$$

$$5. \tan^2 2A = 3 \tan 2A.$$

$$6. \tan A + 3 \cot A = 4.$$

PLANE TRIGONOMETRY.

(3) $\text{Log} \left(1 - \frac{1}{x^2} \right) = \log(x+1) + \log(x-1) - 2 \log x.$

11. If $\log x + \log 6 = \log 12 - \log 2 - \log x$, then $x^2 = 1$.

12. From $1_{10}2$, and $1_{10}3$, find $1_{10}72$.

13. From $1_{10}14$, and $1_{10}16$, find $1_{10}2$, $1_{10}5$, $1_{10}7$, $1_{10} \frac{1}{1900}$.

14. $1_{10} \left\{ (1+N)^{\frac{1+N}{2}} \cdot (1-N)^{\frac{1-N}{2}} \right\} = \frac{N^2}{1 \cdot 2} + \frac{N^4}{3 \cdot 4} + \frac{N^6}{5 \cdot 6} + \dots$

15. If $\log x = \log a - \log b$, shew that

$\log(a+b) = \log a + \log \left(1 + \frac{1}{x} \right)$, and $\log(a-b) = \log a - \log \frac{x}{x-1}$.

16. If $y = \epsilon^{1-\frac{1}{\epsilon^x}}$, and $z = \epsilon^{\frac{1}{1-\frac{1}{\epsilon^y}}}$; then $x = \epsilon^{1-\frac{1}{\epsilon^z}}$.

17. If $m^{ax-\beta} - n^{\gamma x+\delta} = 0$, then $x = \frac{\beta \log m + \delta \log n}{a \log m - \gamma \log n}$.

18. If $a^x + b^y = c$, and $a^x - b^y = d$, then

$x = \frac{\log \frac{1}{2}(c+d)}{\log a}$, and $y = \frac{\log \frac{1}{2}(c-d)}{\log b}$.

19. Prove that

$\log(\tan 2^n \theta) = 2 \log(2 \sin 2^n \theta) - \log(2 \sin 2^{n+1} \theta);$

and thence shew that

$\frac{1}{2} \log(\tan 2^n \theta) + \frac{1}{2^2} \log(\tan 2^2 \theta) + \dots$ (to n terms)

$= \log(2 \sin 2 \theta) - \frac{1}{2^n} \log(2 \sin 2^{n+1} \theta).$

SPHERICAL TRIGONOMETRY.

CHAPTER I.

ON CERTAIN PROPERTIES OF SPHERICAL TRIANGLES.

1. **DEF.** A *Sphere* is a solid bounded by a surface of which every point is equally distant from a fixed point which is called the *center* of the sphere.

DEF. The straight line joining the center with any point in the surface is called the *radius* of the sphere.


2. *Every section of a sphere made by a plane is a circle.*

Let $ABCD$ be a sphere of which the center is O ; $AFCG$ the curve in which a plane cutting the sphere intersects its surface; OE a perpendicular from O upon the cutting plane.

Join E with F any point in $AFCG$, and join FO . Then since OE is perpendicular to the cutting plane, it is perpendicular to EF , a line meeting it in that plane;

$\therefore EF = \sqrt{(OF^2 - OE^2)}$, a constant quantity.

Now E is a *fixed* point in the cutting plane, and F is *any* point in the curve AFC . Therefore AFC is a circle whose center is E and radius is EF . Euclid, I. Def. 15.



Cor. If the cutting plane pass through the center of the sphere, EO vanishes, and EF becomes equal to OF , the radius of the sphere.

SPHERICAL TRIGONOMETRY.

3. DEF. The circle in which a sphere is cut by a plane is called a *great*, or a *small* circle, according as the cutting plane passes, or does not pass, through the center of the sphere.

COR. Since the radius of every great circle is equal to the radius of the sphere (2, Cor.), all great circles of the sphere must be equal.

NOTE. Unless the contrary be expressly mentioned, when hereafter an arc of a sphere is spoken of, an arc of a *great* circle is meant.

4. DEF. SPHERICAL TRIGONOMETRY investigates the relations subsisting between the angles of the plane faces which form a solid angle, and the angles at which the plane faces themselves are inclined to one another.

For the sake of convenience the angular point is made the center of a sphere, which, being cut by the plane faces containing the solid angle, presents on its surface a figure of which the sides are arcs of great circles.


Let ABC be a triangle of this kind, whose sides AB, BC, CA are formed by the intersection of the planes AOB, BOC, COA , with the surface of a sphere of which O is the center. The angle of any face, as AOB , is $\frac{\text{arc } AB}{AO}$, Pl. Trig. Chap. vi. ;

and the angle contained between any two faces (as BOA and COA), is the angle contained between the lines AD and AE which are drawn in the planes BOA and COA at right angles to their intersection OA . Euclid, xi. Def. 6.

The lines AD, AE , being at right angles to the radius OA , and lying in the planes of the arcs AB and AC , are tangents to those arcs.

5. After certain properties of spherical triangles have been proved, it will not be requisite, in pursuing further investigation, to represent in the figures attached to the Propositions the center of the sphere on which the triangles are described.

It must not, however, be forgotten, that when the words "the angle BAC " or "the angle A " occur, the angle meant is that of the inclination of the two planes which pass through O the center of the sphere and the arcs AB and AC ; and that this is the angle contained between two lines drawn from any point in AO at right angles to it, and respectively lying in the planes AOB and AOC . Also, whenever the expression "the sine of AB " occurs, the sine is meant of the angle which the arc AB subtends at the center of the circle of which it is a portion. If this circle be a great circle, its center is also the center of the sphere.



SPHERICAL TRIGONOMETRY.

6. In the following pages the angle BAC will commonly be indicated by the letter A , and the angle subtended by BC , the side of the triangle opposite to $\angle BAC$, will be indicated by a . The other parts of the triangle ACB will be represented in like manner.

7. DEF. If OE (fig. Art. 2), which is perpendicular to the plane AFC , be produced both ways to meet the surface of the sphere in B and D , these points are respectively called *the nearer and the further poles* of the circle AFC ; and the straight line BOD is called *the axis of the circle AFC*.

8. *The pole of a circle is equally distant from every point in its circumference.* (Fig. Art. 2.)

Join B with F any point in $AFCG$. Then, BEF being a right angle, $BF^2 = BE^2 + EF^2 = BE^2 + (OF^2 - OE^2)$, a constant quantity. And F being any point in $AFCG$, B is equally distant from every point in the circumference of that circle.

Similarly, $DF^2 = DE^2 + EF^2 = DE^2 + (OF^2 - OE^2)$, a constant quantity. Hence D is equally distant from every point in the circumference of $AFCG$.

Again; let BNF be an arc of a great circle passing through B and F , any point in AFC . Then since the chord BF is the same for every point in the arc AFC , and that in equal circles equal circumferences are subtended by equal straight lines (Euclid, III. xxix.), the arc BNF is constant; and also, because the radii of all great circles are equal, the angle BOF subtended at the center of the sphere is constant.


Hence it appears that every point in the circumference of a circle of a sphere is equally distant from the pole of the circle, whether the distance be measured by the length of a straight line joining the point with the pole, or by the arc of a great circle connecting the same points, or by the angle which such an arc subtends at the center of the sphere.

9. Since BO is at right angles to the plane AFC , every plane through BO is at right angles to that plane. Hence, *the angle between any circle whatever and a great circle passing through its pole is a right angle.*

10. If FB be a quadrant drawn at right angles to the great circle AFC from any point F in it, B is the pole of AFC .

Join B and F with O the center of the sphere, and in the plane of AFC draw OA at right angles to FO .

Then, AO being perpendicular to OF , and (since FB is a quadrant) BO being perpendicular to the same line, therefore $\angle BOA = \angle$ between the planes AOF and BOF , whose intersection is OF ; an angle which is by hypothesis a right angle.



And BO , being at right angles to OA and OF , is at right angles to the plane AOF in which they lie; and therefore B is the pole of AFC .

COR. If B be the pole of AFC , AO and FO , any two lines in the plane $CFAG$, are at right angles to BO , and AOF is the inclination of AOB and FOB ;

$$\therefore \text{Art. 5, } \angle AF = \angle ABF.$$

DEF. Great circles which pass through the pole of a great circle are called *secondaries* to that circle.

11. If from a point on the surface of a sphere there can be drawn two arcs (not parts of the same circle) which are at right angles to a given circle, that point is the pole of the circle. (Fig. Art. 10.)

Let B be the point, BA and BF arcs (not parts of the same circle) through A and F , points in the circle AFC ; and let BA and BF be at right angles to AF .

Then since BA and BF are at right angles to AF , and are not parts of the same circle, the planes BOA and BOF must intersect, and their intersection BO is at right angles to AOF , Eucl. xi. 19. Therefore B is the pole of AFC .

[The proof holds whether AFC be a great or a small circle.]

COR. Hence also it appears that the intersection of two arcs drawn at right angles to a given circle through any two points in it, is the pole of that circle.

12. *The angle subtended at the center of the sphere by the arc of a great circle which joins the poles of two great circles, is the inclination of the planes of those circles.*

Let the given circles be FD and FE intersecting in F , A and B their respective poles, and $ABDE$ the circle through A and B .

Now AO is perpendicular to OF , which is a line in the plane DOF ,

And BO is perpendicular to OF , a line in the plane EOF ;

$\therefore OF$ is perpendicular to the plane AOB ; and therefore to OD and OE , which are lines in that plane;

$\therefore DOE$ is the angle of inclination of the planes FOD , FOE .

$$\begin{aligned} \text{And } AOB &= AOD - BOD = 90^\circ - BOD \\ &= BOE - BOD = DOE. \end{aligned}$$

COR. Hence also it appears that $\angle AB = \angle DE = \angle DFE$
Arts. 5, 4.


13. *Two great circles are bisected by their intersections.*
(Fig. Art. 10.)

Let BAD and BFD be arcs of two great circles intersecting each other in B and D . Join BD .

Then since BAD is a great circle, the center of the sphere is a point in its plane; similarly, it is a point in the plane of BFD ; therefore the center of the sphere is a point (O) which is in the intersection of these planes, that is, in the line joining B and D . Therefore BAD and BFD are semicircles, having BOD for their common diameter, and the two great circles are bisected by the points B and D .


14. *Any side of a spherical triangle is less than a semicircle, and any angle is less than two right angles.*

Since Euclid takes two right angles as the limit of a plane angle, this is also his limit for the angle of any plane face of a solid



angle. Hence in a spherical triangle no side can be equal to a semicircle.

Thus, if ACB intersect $AEBF$ in the points A and B , and CD be any other arc, the triangle connecting the points B, C, D is *not* the figure formed by $BC, CD, DAFB$ (of which the side $DAFB$ is greater than a semicircle), but the figure formed by BC, CD, DEB .



If a side, as $ADEB$, become a semicircle, then the arcs which join A and B with any other point C , are portions of the semicircle BCA , and the arcs joining the points A, B, C cease to form a triangle.

Wherefore the *side* of a triangle is less than a semicircle.

COR. 1. Hence it follows that an *angle* of a triangle must be less than two right angles.


For if possible let $BFADC$ be a triangle having the angle BCD greater than two right angles. Produce BC to A ; then BFA is a semicircle, and $BFAD$ is greater than a semicircle; which is impossible.

If the angle BCD become equal to two right angles, BC and CD become parts of the same circle, and the figure ceases to be a triangle.

Wherefore the *angle* of a triangle is less than two right angles.

COR. 2. Hence also it follows, that if the great circle be completed of which any side AB of a triangle ABC forms a part, the triangle lies within it.

For if possible, let C lie without the circle, and let CA cut it in D ; then since DA is a semicircle (13), a side CDA of a triangle is greater than a semicircle; which is impossible.



15. If P be the pole of a great circle BAC and of a small circle bac which are cut by the great circles PaA and PbB , then

$$\frac{\text{Arc } ab}{\text{Arc } AB} = \text{Sin } Pa.$$

Let O be the center of the sphere, and OP , which is at right angles to the planes of both circles, cut the plane of the small circle in D . Join AO, BO, aD, bD . Since these lines lie in planes which are perpendicular to OP , each of them is perpendicular to OP . Join aO .


Then aD , being perpendicular to PO , is parallel to AO , a line which lies in the same plane with itself.

Similarly, bD is parallel to BO ;

$$\therefore \angle aDb = \angle AOB. \text{ Eucl. XI. 10,}$$

$$\text{that is, } \frac{\text{arc } ab}{aD} = \frac{\text{arc } AB}{AO};$$

$$\therefore \frac{\text{Arc } ab}{\text{Arc } AB} = \frac{aD}{AO} = \frac{aD}{aO} = \sin POa, \text{ or } \sin Pa.$$



COR. 1. Since $POA = 90^\circ$, $\sin POa = \cos aOA$;

$$\therefore \frac{\text{Arc } ab}{\text{Arc } AB} = \cos aOA, \text{ or } \cos aA.$$

COR. 2. If AB be any circle of which P is the pole, it may be proved in like manner that

$$\frac{\text{Arc } ab}{\text{Arc } AB} = \frac{\sin Pa}{\sin PA}.$$

16. DEF. If the angular points A, B, C of a spherical triangle ABC , Fig. Art. 17, be the poles of the three great circles DE, EF, FD respectively, the triangle ABC is called with respect to DEF the *Primitive Triangle*, and DEF is called with respect to ABC the *Polar Triangle*.


17. *The angular points of the polar triangle are the poles of the sides of the primitive triangle.*

Join AE and BE , by arcs of two great circles.

Then, since A is the pole of ED , $AE = \frac{1}{2}\pi$,
Art. 10.

And, since B is the pole of EF , $BE = \frac{1}{2}\pi$.

Therefore the great circle of which E is the pole passes through A and B , or E is the pole of AB . Similarly D and F are the poles of AC and BC respectively.



18. The *Polar Triangle* is called also the *Supplemental Triangle*, from the following property it possesses.

The sides and angles of the polar triangle are the supplements of the angles and sides respectively of the primitive triangle. (Fig. Art. 17.)

$$\begin{aligned}\text{For } \angle A &= HG, \text{ Art. 10, Cor.,} = EG - EH \\ &= EG - (ED - DH) = EG + DH - ED;\end{aligned}$$

but EG and DH are quadrants, or each subtends an angle $\frac{1}{2}\pi$,

$$\therefore \angle A = \pi - ED.$$

Similarly, $\angle B = \pi - EF$, and $\angle C = \pi - FD$.

$$\begin{aligned}\text{Again, } AB &= AG - BG = AG - (IG - BI) \\ &= AG + BI - IG \\ &= \frac{1}{2}\pi + \frac{1}{2}\pi - E, \text{ since } E = IG. \text{ (10, Cor.)} \\ &= \pi - E.\end{aligned}$$

Similarly, $BC = \pi - F$, and $AC = \pi - D$.

19. *If a general equation be established between the sides and the angles of a spherical triangle, a true result is obtained if in the equation the supplements of the sides and of the angles respectively be written for the angles and sides which enter into the equation.*

For if the equation be proved for any triangle whatever, the sides a', b', c' , and the angles A', B', C' , of the *polar triangle* may respectively be substituted in it, in the place of the sides a, b, c , and the angles A, B, C , of the *primitive triangle*. And in the equation as it then stands, putting for the sides and angles of the polar triangle their equivalents drawn from the primitive triangle, viz.

$$\begin{aligned}\pi - A &= a', & \pi - B &= b', & \pi - C &= c', \\ \pi - a &= A', & \pi - b &= B', & \pi - c &= C',\end{aligned}$$

a true result is obtained, which differs from the original equation in having the supplements of the sides and of the angles respectively written for the angles and the sides of the triangle.

Ex. If A, B, C be the angles, and a, b, c be the sides, of any triangle, it will hereafter be proved (29) that

$$\text{Cos } A = \frac{\cos a - \cos b \cos c}{\sin b \sin c},$$

in which the cosine of an angle of a spherical triangle is expressed in terms of the sides.

$$\text{Therefore } \text{Cos } A' = \frac{\cos a' - \cos b' \cos c'}{\sin b' \sin c'}.$$

Now $A' = \pi - a$, $a' = \pi - A$, $b' = \pi - B$, $c' = \pi - C$;

$$\therefore \text{Cos}(\pi - a) = \frac{\cos(\pi - A) - \cos(\pi - B) \cos(\pi - C)}{\sin(\pi - B) \sin(\pi - C)};$$


$\therefore \text{Cos } a = \frac{\cos A + \cos B \cos C}{\sin B \sin C}$; in which the cosine of a side of a triangle is expressed in terms of the angles.

20. Any two sides of a spherical triangle are together greater than the third; and the sum of the three sides is less than the circumference of a great circle.

For, Eucl. xi. 20, any two of the angles AOB, BOC, COA ,

$$\left(\text{or } \frac{\text{arc } AB}{AO}, \frac{\text{arc } BC}{AO}, \frac{\text{arc } CA}{AO}\right),$$

which form the solid angle at O , are together greater than the third. Therefore any two of the arcs AB, BC, CA are together greater than the third.



Also, Eucl. xi. 21, the three angles forming the solid angle at O are less than four right angles;

$$\therefore \frac{\text{arc } AB}{AO} + \frac{\text{arc } BC}{AO} + \frac{\text{arc } CA}{AO} < 2\pi.$$

$\therefore \text{Arc } AB + \text{arc } BC + \text{arc } CA < 2\pi \cdot AO$, which is the circumference of a circle whose radius is AO .

COR. 1. Since the sum of the plane angles which contain *any* solid angle is less than four right angles, Eucl. xi. 21, it follows from the same mode of proof, that the sum of the sides of *any* polygonal figure which is described on a sphere, and whose sides are arcs of great circles, is less than the circumference of a great circle.

[The polygonal figure, however, must be such that the angle between any two adjacent sides is less than two right angles, and also that its area is contained on the surface of a hemisphere: for by Euclid, the angle between two planes is less than two right angles, and in the proof of the proposition referred to, it is supposed that *all* the plane faces which contain the solid angle may be cut by *one* plane; which cannot be the case unless all the edges of the solid angle lie within the same hemisphere.]

COR. 2. Also, let $ABCDEA$ be a five-sided figure described on a sphere, and let it be divided into triangles by the arcs AC , AD . (A figure is easily drawn.)

Then $AB + BC > AC$;

$\therefore AB + BC + CD > AC + CD > AD$, for $AC + CD > AD$;

$\therefore AB + BC + CD + DE > AD + DE > AE$.

And the same method of proof being applicable to a polygon of any number of sides, it follows that the sum of all the sides but one of a polygon described on a sphere is greater than the remaining side.

COR. 3. If a and b be two sides of a spherical triangle, since each is less than π ,

$$\therefore a + b < 2\pi, \quad \therefore \frac{1}{2}(a + b) < \pi.$$

$$\text{And } a - b < \pi, \quad \therefore \frac{1}{2}(a - b) < \frac{1}{2}\pi.$$

$$\text{So, } \frac{1}{2}(A + B) < \pi, \text{ and } \frac{1}{2}(A - B) < \frac{1}{2}\pi.$$

COR. 4. a' , b' , c' , being the sides of the polar triangle,

$$a' + b' > c'; \quad \therefore (\pi - A) + (\pi - B) > (\pi - C);$$

$$\therefore \pi > A + B - C; \quad \text{or, } A + B - C < \pi.$$

21. *The sum of the angles of a spherical triangle is greater than two right angles, and less than six.*

Let A, B, C be the angles, and a, b, c the sides, of a triangle; A', B', C' , and a', b', c' , the angles and sides of its polar triangle.

$$\text{Now } 2\pi > a' + b' + c'. \quad (20)$$

$$> (\pi - A) + (\pi - B) + (\pi - C) > 3\pi - (A + B + C);$$

$$\therefore A + B + C > \pi.$$


Also, since each of the angles A, B, C is less than π ,

$$\therefore A + B + C < 3\pi.$$

22. *The angles at the base of an isosceles triangle are equal to each other.*

Suppose ABC to be an isosceles triangle, having $AB = AC$, and therefore $\angle AOB = \angle AOC$.

From D , any point in OA , draw DG perpendicular to the plane BOC ,—and therefore at right angles to every line it meets in that plane; and from G draw GE and GF perpendicular to OB and OC ; join DE, DF, GO .



Then

$$OE^2 = OG^2 - GE^2 = (OD^2 - DG^2) - (ED^2 - DG^2) = OD^2 - ED^2;$$

$\therefore DE$ is perpendicular to OF , and $\therefore \angle DEG =$ inclination of the planes BOC and BOA , $= \angle B$.

Similarly, DF is perpendicular to OC , and $\angle DFG = \angle C$.

$$\text{Now } DE = OD \cdot \sin AOB = OD \cdot \sin AOC = DF.$$

$$\text{And } EG^2 = DE^2 - DG^2 = DF^2 - DG^2 = FG^2.$$

Hence, since GE , ED are equal to GF , FD , each to each, and GD is common,

$$\therefore \angle DEG = \angle DFG, \quad \text{or } \angle B = \angle C.$$

23. Conversely, if $\angle B = \angle C$, or $\angle DEG = \angle DFG$, it may be shewn from the same figure that $AB = AC$; that is, the angles at the base being equal, the sides opposite to them are equal.


COR. Hence also it follows that every equilateral triangle is also equiangular; and conversely, that every equiangular triangle is also equilateral.

24. *Of the two sides which are opposite to two unequal angles in a triangle, that is the greater which is opposite to the angle which is the greater.*

Let $\angle ABC$ be greater than $\angle BAC$.
Then $AC > BC$.

Make $\angle ABD = \angle BAD$; $\therefore DA = DB$.

And $AC = CD + DA = CD + DB$;



But $CD + DB$ is $> BC$ (20); $\therefore AC > BC$.

25. Conversely, it may easily be shewn that of two angles in a triangle which are opposite to unequal sides, that is the greater which is opposite to the side which is the greater.

COR. Hence $A - B$ and $a - b$ have the same sign.

26. Article 24 has been proved after the manner of Euclid, 1. 19. The following propositions may be enunciated for spherical triangles in the terms used for plane triangles, and may be proved in nearly the same words. Euclid, 1. Props. 4. 8. 24. 25. 26. &c.

27. To recapitulate the more important properties of Spherical Triangles which have been proved in this Chapter.

Art. 14.	1. A side must be less than a semicircle.
14.	2. An angle must be less than two right angles.
20.	3. Any two sides are together greater than the third.
24.	4. The greater side is opposite to the greater angle; and conversely.
25, Cor.	5. $A - B$ and $a - b$ are of the same sign.
20.	6. $a + b + c < 2\pi$.
21.	7. $A + B + C > \pi$, and $< 3\pi$.
20, Cor. 3.	8. $a + b < 2\pi$, $a - b < \pi$.
20, Cor. 3.	9. $A + B < 2\pi$, $A - B < \pi$.
20, Cor. 4.	10. $A + B - C < \pi$.

The following properties will also be proved hereafter:


34.	11. $\frac{1}{2}(A + B)$ and $\frac{1}{2}(a + b)$ are "of the same affection;" i. e. they are both $> \frac{1}{2}\pi$, or both $< \frac{1}{2}\pi$.
39.	12. If ABC be a right-angled triangle, C being the right angle, then A and a (as also B and b) are of the same affection.

CHAPTER II.

FORMULÆ CONNECTING THE SIDES AND ANGLES OF A SPHERICAL TRIANGLE.

28. *The Sines of the Sides of a Spherical Triangle are as the Sines of the Angles to which they are respectively opposite.*

Let ABC be the triangle, O the center of the sphere. From D , any point in OA , draw DG perpendicular to the plane BOC , and from G draw GE and GF perpendicular to OB and OC . Join DE , DF .



Then a line through G parallel to OB would be perpendicular to DG and GE , and \therefore to the plane DGE : wherefore, Euc. xi. 8, OE is perpendicular to the plane DEG , and \therefore to DE and EG ; and $\therefore \angle DEG = \angle B$.

Similarly, $\angle DFG = \angle C$.

$$\text{Now } DE \cdot \sin DEG = DG = DF \cdot \sin DFG;$$

$$\therefore OD \cdot \sin AOB \cdot \sin DEG = OD \cdot \sin AOC \cdot \sin DFG;$$

$$\therefore \sin AB \cdot \sin B = \sin AC \cdot \sin C;$$

$$\therefore \frac{\sin B}{\sin AC} = \frac{\sin C}{\sin AB}.$$

Similarly it may be proved that

$$\frac{\sin B}{\sin AC} = \frac{\sin A}{\sin BC}.$$


Wherefore

$$\frac{\sin A}{\sin BC} = \frac{\sin B}{\sin AC} = \frac{\sin C}{\sin AB} \dots\dots\dots(i.)$$

The Proof will be found to hold whether G fall within the angle BOC or without it. For if, for instance, G fall without OB , the angle DEG will be the supplement of the angle B , and $\sin DEG = \sin B$. Also if E be in BO produced, the proof will still be found to hold.

29. To express the cosine of an Angle of a Triangle in terms of the Cosines and Sines of the Sides.

From any point D in OA draw DE and DF , in the planes AOB and AOC , at right angles to AO . Therefore $\angle EDF$ is the inclination of those planes to each other, that is, the angle A . Join EF . Then, from the triangles FOE and FDE ,



$$OF^2 + OE^2 - 2OF \cdot OE \cdot \cos FOE = FE^2$$

$$= FD^2 + DE^2 - 2FD \cdot DE \cdot \cos FDE;$$

$$\therefore 2OF \cdot OE \cdot \cos FOE$$

$$= (OF^2 - FD^2) + (OE^2 - DE^2) + 2FD \cdot DE \cdot \cos FDE$$

$$= 2OD^2 + 2FD \cdot DE \cdot \cos FDE;$$

$$\therefore \cos FOE = \frac{OD}{OF} \cdot \frac{OD}{OE} + \frac{FD \cdot DE}{OF \cdot OE} \cdot \cos FDE;$$

$$\text{or, } \cos a = \cos b \cos c + \sin b \sin c \cos A \dots \dots \dots \text{(ii.)}$$

$$\text{Whence } \cos A = \frac{\cos a - \cos b \cos c}{\sin b \sin c} \cdot *$$

* In the proof DE and DF are supposed to cut OB and OC , which will be the case only if AB and AC be less than quadrants. The result arrived at is notwithstanding general, as may be thus shewn. [The figures may easily be drawn from the descriptions.]

1st. If AC be greater than 90° , and AB less than 90° , produce CA and CB until they meet in C' . Then $AC'B$ is a triangle in which AC' , AB are each less than 90° , and therefore by the proof in the text,

$$\cos C'B = \cos AC' \cdot \cos AB + \sin AC' \cdot \sin AB \cdot \cos BAC'$$

$$\text{or } -\cos BC = -\cos AC \cdot \cos AB + \sin AC \cdot \sin AB \cdot (-\cos BAC);$$

$$\therefore \cos BC = \cos AC \cdot \cos AB + \sin AC \cdot \sin AB \cdot \cos BAC.$$

2nd.


Cor. By writing $\pi - a$ for A , &c. (19), .

$$\cos A + \cos B \cos C = \sin B \sin C \cos a \dots\dots\dots(\text{iii.})$$

$$\text{Whence, } \cos a = \frac{\cos A + \cos B \cos C}{\sin B \sin C} .$$

30. To shew that $\cos a \sin b = \sin a \cos b \cos C + \sin c \cos A$.

Produce the side CA to P , making AP a quadrant. Then the formula (ii.) applied to the triangles PCB and PAB gives



$$\begin{aligned} \cos BP &= \cos(90^\circ + b) \cos a \\ &\quad + \sin(90^\circ + b) \sin a \cos C \\ &= -\sin b \cos a + \cos b \sin a \cos C, \end{aligned}$$

$$\text{and } \cos BP = -\sin c \cos A ;$$

Equating these values of $\cos BP$,

$$\cos a \sin b = \sin a \cos b \cos C + \sin c \cos A \dots\dots(\text{iv.})$$

2nd. If AC and AB be both greater than 90° , produce AC and AB until they meet in A' . Then $A'BC$ is a triangle in which AB and AC are each less than 90° .

$$\begin{aligned} \therefore \cos BC &= \cos A'B \cdot \cos A'C + \sin A'B \cdot \sin A'C \cdot \cos A', \\ &= \cos(180^\circ - AB) \cdot \cos(180^\circ - AC) + \sin(180^\circ - AB) \cdot \sin(180^\circ - AC) \cdot \cos A \\ &= \cos AB \cdot \cos AC + \sin AB \cdot \sin AC \cdot \cos A. \end{aligned}$$

3rd. Let one of the sides, as AB , be 90° .

From B draw BD perpendicular to AC or AC produced. Then the angles at D are right angles; and from the $\triangle BCD$,

$$\cos BC = \cos DB \cdot \cos DC + \sin DB \cdot \sin DC \cdot \cos D,$$

where $\cos DB = \cos A$; $\cos DC = \cos(90^\circ \sim AC) = \sin AC$; $\cos D = 0$, and the equation becomes

$$\cos a = \cos A \cdot \sin b,$$

which is the form (ii.) assumes when $c = 90^\circ$.

4th. If both the sides AB, AC be quadrants, $\cos A = \cos BC = \cos a$; and this is the form which (ii.) assumes when b and c each become 90° .

COR. 1. By (iv.), $\text{Cos } a \sin b = \sin a \cos b \cos C + \sin c \cos A$,

$$\begin{aligned} \therefore \frac{\cos a}{\sin a} \cdot \sin b &= \cos b \cos C + \frac{\sin c}{\sin a} \cdot \cos A \\ &= \cos b \cos C + \frac{\sin C}{\sin A} \cdot \cos A; \end{aligned}$$

$$\therefore \text{Cot } a \sin b = \cos b \cos C + \sin C \cot A \dots \dots \text{(v.)}$$

COR. 2. By writing $\pi - a$ for A , &c. (19), the formula (iv.) becomes,

$$\text{Sin } C \cos a = \cos A \sin B + \sin A \cos B \cos c \dots \dots \text{(vi.)}$$

31. To find the values of $\text{Cos } \frac{1}{2} A$, $\text{Sin } \frac{1}{2} A$, $\text{Tan } \frac{1}{2} A$, and $\text{Sin } A$ in terms of a , b , c .

By (ii.), $\text{Sin } b \sin c \cos A = \cos a - \cos b \cos c$;

$$\therefore \sin b \sin c (1 + \cos A) = \cos a - (\cos b \cos c - \sin b \sin c);$$

$$\begin{aligned} \therefore \sin b \sin c 2 \cos^2 \frac{1}{2} A &= \cos a - \cos (b + c) \\ &= 2 \sin \frac{1}{2} (a + b + c) \sin \frac{1}{2} (b + c - a). \end{aligned}$$

And if $S = \frac{1}{2} (a + b + c)$, $S - a = \frac{1}{2} (a + b + c) - a = \frac{1}{2} (b + c - a)$,

$$\therefore \cos^2 \frac{1}{2} A = \frac{\sin S \sin (S - a)}{\sin b \sin c};$$

$$\therefore \text{Cos } \frac{1}{2} A = \sqrt{\frac{\sin S \sin (S - a)}{\sin b \sin c}} \dots \dots \text{(vii.)}$$

Again, $2 \sin^2 \frac{1}{2} A = 1 - \cos A = 1 - \frac{\cos a - \cos b \cos c}{\sin b \sin c}$;

$$\text{Whence, Sin } \frac{1}{2} A = \sqrt{\frac{\sin (S - b) \sin (S - c)}{\sin b \sin c}} \dots \dots \text{(viii.)}$$

$$\text{Also, Tan } \frac{1}{2} A = \frac{\sin \frac{1}{2} A}{\cos \frac{1}{2} A} = \sqrt{\frac{\sin (S - b) \sin (S - c)}{\sin S \sin (S - a)}} \dots \dots \text{(ix.)}$$

And $\text{Sin } A = 2 \sinh \frac{1}{2} A \cos \frac{1}{2} A$
 $= \frac{2}{\sin b \sin c} \cdot \sqrt{\{\sin S \sin (S - a) \sin (S - b) \sin (S - c)\}} \dots\dots\dots(x.)$

The positive signs of the square roots are taken in these cases, because $\frac{1}{2} A$ is necessarily less than $\frac{1}{2} \pi$.

32. To find the values of $\text{Cos } \frac{1}{2} a$, $\text{Sin } \frac{1}{2} a$, $\text{Tan } \frac{1}{2} a$, and $\text{Sin } a$ in terms of A , B , C .

By (iii.), $\text{Cos } a = \frac{\cos A + \cos B \cos C}{\sin B \sin C}$;

$\therefore 2 \cos^2 \frac{1}{2} a = 1 + \cos a = \frac{\cos A + \cos B \cos C + \sin B \sin C}{\sin B \sin C}$
 $= \frac{\cos A + \cos (B - C)}{\sin B \sin C} = \frac{2 \cos \frac{1}{2} (A + C - B) \cos \frac{1}{2} (A + B - C)}{\sin B \sin C}$.

And if $S' = \frac{1}{2} (A + B + C)$, then $S' - B = \frac{1}{2} (A + C - B)$, &c. ;

$\therefore \text{Cos } \frac{1}{2} a = \sqrt{\frac{\cos (S' - B) \cos (S' - C)}{\sin B \sin C}} \dots\dots\dots(xi.)$

Again, $2 \sin^2 \frac{1}{2} a = 1 - \cos a = 1 - \frac{\cos A + \cos B \cos C}{\sin B \sin C}$
 $= \frac{\cos A + \cos (B + C)}{\sin B \sin C}$;

Whence, $\text{Sin } \frac{1}{2} a = \sqrt{\frac{-\cos S' \cos (S' - A)}{\sin B \sin C}} \dots\dots\dots(xii.)$

Also, $\text{Tan } \frac{1}{2} a = \frac{\sin \frac{1}{2} a}{\cos \frac{1}{2} a} = \sqrt{\frac{-\cos S' \cos (S' - A)}{\cos (S' - B) \cos (S' - C)}} \dots\dots\dots(xiii.)$

And $\text{Sin } a = 2 \sin \frac{1}{2} a \cos \frac{1}{2} a$
 $= \frac{2}{\sin B \sin C} \cdot \sqrt{\{-\cos S' \cos (S' - A) \cos (S' - B) \cos (S' - C)\}} \dots(xiv.)$

NOTE. Since the angles of a spherical triangle are together greater than two right angles and less than six, S' is greater than one right angle and less than three, and its cosine is therefore a negative quantity. Also since (27) $A + B - C < \pi$, therefore $\cos \frac{1}{2}(A + B - C)$, or $\cos(S' - C)$, is a positive quantity; and in like manner $\cos(S' - B)$ and $\cos(S' - A)$ are positive. Wherefore the last three formulæ are not *impossible* quantities, as at first sight they appear to be, but *real* quantities.

33. To prove Napier's Analogies.

$$\text{By (iii.), } \cos A + \cos B \cos C = \cos a \sin B \sin C;$$

$$\text{so } \cos B + \cos A \cos C = \cos b \sin A \sin C.$$

By addition,

$$(\cos A + \cos B)(1 + \cos C) = (\cos a \sin B + \cos b \sin A) \sin C;$$

$$\therefore (\cos A + \cos B) 2 \cos^2 \frac{1}{2} C$$

$$= (\cos a \sin A \cdot \frac{\sin b}{\sin a} + \cos b \sin A) 2 \sin \frac{1}{2} C \cos \frac{1}{2} C;$$

$$\therefore \cos A + \cos B = \frac{\sin A}{\sin a} \cdot (\cos a \sin b + \cos b \sin a) \tan \frac{1}{2} C$$

$$= \frac{\sin A}{\sin a} \cdot \sin(a + b) \tan \frac{1}{2} C.$$

$$\text{Again, } \sin A + \sin B = \sin A \left(1 + \frac{\sin B}{\sin A}\right) = \sin A \left(1 + \frac{\sin b}{\sin a}\right)$$

$$= \frac{\sin A}{\sin a} \cdot (\sin a + \sin b);$$

$$\therefore \frac{\sin A + \sin B}{\cos A + \cos B} = \frac{\sin a + \sin b}{\sin(a + b)} \cdot \cot \frac{1}{2} C,$$

$$\frac{2 \sin \frac{1}{2}(A + B) \cos \frac{1}{2}(A - B)}{2 \cos \frac{1}{2}(A + B) \cos \frac{1}{2}(A - B)} = \frac{2 \sin \frac{1}{2}(a + b) \cos \frac{1}{2}(a - b)}{2 \sin \frac{1}{2}(a + b) \cos \frac{1}{2}(a + b)} \cdot \cot \frac{1}{2} C;$$

$$\therefore \tan \frac{1}{2}(A + B) = \frac{\cos \frac{1}{2}(a - b)}{\cos \frac{1}{2}(a + b)} \cdot \cot \frac{1}{2} C \dots \dots \dots (\text{xv.})$$

In like manner,
$$\frac{\sin A - \sin B}{\cos A + \cos B} = \frac{\sin a - \sin b}{\sin(a+b)} \cdot \cot \frac{1}{2} C;$$

Whence,
$$\tan \frac{1}{2} (A - B) = \frac{\sin \frac{1}{2} (a - b)}{\sin \frac{1}{2} (a + b)} \cdot \cot \frac{1}{2} C \dots\dots (\text{xvi.})$$

By writing $\pi - A$ for a , &c. (19), the last two formulæ become

$$\tan \frac{1}{2} (a + b) = \frac{\cos \frac{1}{2} (A - B)}{\cos \frac{1}{2} (A + B)} \cdot \tan \frac{1}{2} c \dots\dots (\text{xvii.})$$

$$\tan \frac{1}{2} (a - b) = \frac{\sin \frac{1}{2} (A - B)}{\sin \frac{1}{2} (A + B)} \cdot \tan \frac{1}{2} c \dots\dots (\text{xviii.})$$

These four formulæ are called **NAPIER'S ANALOGIES**. The latter two may be easily proved independently by beginning with the formula

$$\cos a = \cos b \cos c + \sin b \sin c \cos A.$$

34. Since $\cos \frac{1}{2} (a - b)$ and $\cot \frac{1}{2} C$ are necessarily positive quantities, it follows from (xv.) that $\tan \frac{1}{2} (A + B)$ and $\cos \frac{1}{2} (a + b)$ have the same algebraic sign; i. e. $\frac{1}{2} (A + B)$ and $\frac{1}{2} (a + b)$ must both be greater, or both less, than a right angle.

DEF. When two angles are both greater or both less than a right angle, they are said to be *of the same*, or *of the like*, affection.

Wherefore in any Spherical Triangle,

$$\frac{1}{2} (A + B) \text{ and } \frac{1}{2} (a + b) \text{ are of the same affection.}$$

35. **GAUSS' THEOREM.** *To shew that*

$$\cos \frac{1}{2} c \sin \frac{1}{2} (A + B) = \cos \frac{1}{2} C \cos \frac{1}{2} (a - b) \dots\dots (\text{xix.})$$

$$\cos \frac{1}{2} c \cos \frac{1}{2} (A + B) = \sin \frac{1}{2} C \cos \frac{1}{2} (a + b) \dots\dots (\text{xx.})$$

$$\sin \frac{1}{2} c \sin \frac{1}{2} (A - B) = \cos \frac{1}{2} C \sin \frac{1}{2} (a - b) \dots\dots (\text{xxi.})$$

$$\sin \frac{1}{2} c \cos \frac{1}{2} (A - B) = \sin \frac{1}{2} C \sin \frac{1}{2} (a + b) \dots\dots (\text{xxii.})$$

Since $\text{Sin } \frac{1}{2}(A+B) = \sin \frac{1}{2}A \cos \frac{1}{2}B + \cos \frac{1}{2}A \sin \frac{1}{2}B$, taking the values of the cosines and sines of $\frac{1}{2}A$ and $\frac{1}{2}B$ from (vii.) and (viii.),

$$\begin{aligned} \text{Sin } \frac{1}{2}(A+B) &= \sqrt{\frac{\sin(S-b) \sin(S-c) \cdot \sin S \sin(S-b)}{\sin b \sin c \cdot \sin a \sin c}} \\ &\quad + \sqrt{\frac{\sin S \sin(S-a) \cdot \sin(S-a) \sin(S-c)}{\sin b \sin c \cdot \sin a \sin c}} \\ &= \left\{ \frac{\sin(S-b)}{\sin c} + \frac{\sin(S-a)}{\sin c} \right\} \cdot \sqrt{\frac{\sin S \sin(S-c)}{\sin a \sin b}} \\ &= \frac{2 \sin \left\{ S - \frac{1}{2}(a+b) \right\} \cos \frac{1}{2}(a-b)}{\sin c} \cdot \cos \frac{1}{2}C \\ &= \frac{\cos \frac{1}{2}(a-b)}{\cos \frac{1}{2}c} \cdot \cos \frac{1}{2}C, \quad \text{since } S - \frac{1}{2}(a+b) = \frac{1}{2}c; \end{aligned}$$

$$\therefore \cos \frac{1}{2}c \sin \frac{1}{2}(A+B) = \cos \frac{1}{2}C \cos \frac{1}{2}(a-b).$$

And the formulæ (xx.), (xxi.), (xxiii.) can be proved in like manner.

There will be no ambiguity respecting the algebraical signs in these formulæ, if it be borne in mind that if A be $> = < B$, then a is $> = < b$, and that $\frac{1}{2}(A+B)$ and $\frac{1}{2}(a+b)$ are of the like affection.

CHAPTER III.

ON THE SOLUTION OF RIGHT-ANGLED TRIANGLES.

86. RIGHT-ANGLED triangles may in all cases (with an exception which will be pointed out, Art. 40), be solved by means of the following formulæ, when, besides the right angle, two other quantities are given out of the three sides and the two remaining angles.

If A, B, C be the angles of any spherical triangle, and a, b, c the sides respectively opposite to them, it has been proved that

$$(ii.) \quad \text{Cos } c = \text{cos } a \text{ cos } b + \sin a \sin b \text{ cos } C \dots\dots\dots(a)$$

$$(iii.) \quad \text{Cos } c \sin A \sin B = \text{cos } C + \text{cos } A \text{ cos } B \dots\dots\dots(b)$$

$$\text{Cos } a \sin B \sin C = \text{cos } A + \text{cos } B \text{ cos } C \dots\dots\dots(c)$$

$$(i.) \quad \text{Sin } a \sin C = \sin c \sin A \dots\dots\dots(d)$$

$$(v.) \quad \text{Cot } a \sin b = \text{cos } b \text{ cos } C + \sin C \text{ cot } A \dots\dots\dots(e)$$

$$\text{Cot } c \sin a = \text{cos } a \text{ cos } B + \sin B \text{ cot } C \dots\dots\dots(f)$$

By making $C = 90^\circ$, there will be obtained,

From (a), $\text{Cos } c = \text{cos } a \text{ cos } b.$

(b), $\text{Cos } c = \text{cot } A \text{ cot } B.$

(c), $\text{Cos } A = \text{cos } a \sin B.$

$\text{Cos } B = \text{cos } b \sin A.$

From (d), $\text{Sin } a = \sin A \sin c.$

$\text{Sin } b = \sin B \sin c.$

(e), $\text{Sin } b = \tan a \text{ cot } A$

$\text{Sin } a = \tan b \text{ cot } B.$

From (f), $\text{Cos } B = \text{cot } c \tan a.$

$\text{Cos } A = \text{cot } c \tan b.$

87. These results are comprised under the following formulæ, which the Student will find it necessary to keep in his memory.

- (1). *Cos hyp* = product of cosines of *sides*.
- (2). *Cos hyp* = product of cotangents of *angles*.
- (3). *Sin side* = *sin opposite angle* × *sin hyp*.
- (4). *Tan side* = *tan hyp* × *cos included angle*.
- (5). *Tan side* = *tan opposite angle* × *sin the other side*.
- (6). *Cos angle* = *cos opposite side* × *sin the other angle*.

38. An artificial method of remembering these formulæ is by **NAPIER'S RULES**.

The formulæ of the last Article are comprised under two Rules, which take their name from Napier, who first gave them.

The right angle being left out of consideration, the *two Sides which include the right angle*, and the *Complements of the Hypothenuse and of the other Angles*, are called the *Circular parts* of the triangle. Any one of these being fixed upon as *the middle part (M)*, the two circular parts next to it and *immediately joining it* are called *the adjacent parts (A₁, A₂)*, and the other two parts are called *the opposite parts (O₁, O₂)*.

Thus in the triangle *ABC* whose right angle is *C*;

If <i>M</i> be	The adjacent Parts are	The opposite Parts are
<i>a</i> , (one of the sides including the right angle).	$\frac{1}{2}\pi - B$, <i>b</i> ;	$\frac{1}{2}\pi - A$, $\frac{1}{2}\pi - c$.
$\frac{1}{2}\pi - A$, (the complement of an angle),	<i>b</i> , $\frac{1}{2}\pi - c$;	$\frac{1}{2}\pi - B$, <i>a</i> .
$\frac{1}{2}\pi - c$, (the complement of the hypothenuse),	$\frac{1}{2}\pi - A$, $\frac{1}{2}\pi - B$;	<i>a</i> , <i>b</i> .

And **NAPIER'S RULES** are,

$\text{Sin } M = \text{product of the Tangents of the Adjacent parts} = \tan A_1 \cdot \tan A_2$,

$\text{Sin } M = \text{product of the Cosines of the Opposite parts} = \cos O_1 \cdot \cos O_2$;

with which the formulæ of the last Article will, on trial, be found to agree.

39. To shew that in a triangle ABC in which C is a right angle, A and a are of the like affection, as are also B and b .


$$\text{By 37, (5), } \sin b = \frac{\tan a}{\tan A}.$$

Now since b is less than π , $\sin b$ is positive; therefore $\tan a$ and $\tan A$ must be of the same sign. And because π is the limit both of a and of A , these angles must be both greater or both less than a right angle; that is, A and a must be of the like affection.

Similarly, from $\sin a = \frac{\tan b}{\tan B}$ it appears that B and b are of the like affection.

40. If, in a right-angled triangle, an Angle and the Side opposite to it be the only quantities given, the triangle cannot be determined.

For if the circles AB and AC intersect again in A' , and C be a right angle, it is evident that ACB and $A'CB$ have the angles A, A' equal, and CB , the side opposite to these angles, is the same in both triangles. It is therefore ambiguous whether ABC or $A'BC$ be the triangle sought.



This ambiguity will also be found to exist, if it be attempted to determine the triangle by 37, (5). For it cannot be determined from the equation $\sin AC = \tan CB \cot A$ whether the angle AC is to be taken, or its supplement $A'C$.

41. The solutions of the other cases of a right-angled triangle from two given parts are not ambiguous, if attention be paid to these two principles;

- (1) The greater side is opposite to the greater angle,
- (2) An angle and the side opposite to it are of the like affection.

[For example: Let c and A be given, to find a, B, b .

Now $\sin a = \sin A \sin c$; and since a and A are of the like affection (39), the greater or lesser angle which satisfies this equation is to be taken for a , according as A is greater or less than $\frac{1}{2}\pi$.

Again, $\cos c = \cot A \cot B$; $\therefore \tan B = \cot A \sec c$. And B is $< 90^\circ$, or $> 90^\circ$, according as the second member of the equation is positive or negative; that is, as A and c are of like or unlike affection.

Again, $\cos c = \cos a \cos b$; $\therefore \cos b = \cos c \sec a$. And b is $< 90^\circ$, or $> 90^\circ$, according as the second member of the equation is positive or negative; that is, as a and c are of like or unlike affection.]

42. In selecting a formula, attention must be paid to the principles laid down in Appendix II. to Pl. Trig. The following formulæ may be used with advantage, when the side or angle required is small, or nearly equal to one right angle, or to two.

$\cos c = \cot A \cot B$; whence c cannot be accurately determined, if it be either a very small angle or nearly equal to two right angles.

$$\text{Now } 2 \sin^2 \frac{1}{2} c = 1 - \cos c = 1 - \cot A \cot B = 1 - \frac{\cos A \cos B}{\sin A \sin B};$$

$$\therefore \sin \frac{1}{2} c = \sqrt{\frac{1 - \cos A \cos B}{2 \sin A \sin B}}.$$

$$\text{So, } \cos \frac{1}{2} c = \sqrt{\frac{1}{2}(1 + \cos c)} = \sqrt{\frac{\cos(A - B)}{2 \sin A \sin B}}.$$

In like manner there is obtained from $\cos a = \frac{\cos A \sin(\frac{1}{2}\pi - A)}{\sin B}$,

$$\left\{ \begin{array}{l} \cos \frac{1}{2} a = \sqrt{\frac{\sin\{\frac{1}{2}(B - A) + \frac{1}{4}\pi\} \cos\{\frac{1}{2}(B + A) - \frac{1}{4}\pi\}}{\sin B}}; \\ \sin \frac{1}{2} a = \sqrt{\frac{\sin\{\frac{1}{2}(B + A) - \frac{1}{4}\pi\} \cos\{\frac{1}{2}(B - A) + \frac{1}{4}\pi\}}{\sin B}}. \end{array} \right.$$

* Since (21), $A + B + C > \pi$, \therefore if $C = \frac{1}{2}\pi$, $A + B > \frac{1}{2}\pi$,
and since (20, Cor. 4), $A + B - C < \pi$, \therefore $A + B < \frac{3}{2}\pi$;

wherefore, $\cos(A + B)$ is necessarily a negative quantity here.

Also since (20, Cor. 4), $A + C - B < \pi$, $\therefore A - B < \frac{1}{2}\pi$, and $\cos(A - B)$ is positive.

So from $\text{Cos } A = \cot c \tan b$,

$$\begin{cases} \text{Sin } \frac{1}{2} A = \sqrt{\frac{1}{2} (1 - \cot c \tan b)} = \sqrt{\frac{\sin (c - b)}{2 \sin c \cos b}}, \\ \text{Cos } \frac{1}{2} A = \sqrt{\frac{1}{2} (1 + \cot c \tan b)} = \sqrt{\frac{\sin (c + b)}{2 \sin c \cos b}}. \end{cases}$$

When c and A are given, if a be nearly a right angle it cannot be accurately determined from its sine. In this case $\text{Cot } B = \cos c \tan A$ determines B , and a may be found from the formula for $\text{Sin } \frac{1}{2} a$, or from that for $\text{Cos } \frac{1}{2} a$, (32.)

43. DEF. A triangle is called *Quadrantal*, if any one of its sides be a quadrant.

A QUADRANTAL TRIANGLE may be solved by applying to its Polar Triangle the formulæ employed for solving a Right-angled Triangle.

A Collection of Examples for practice is added at the end of this Treatise.

CHAPTER IV.

ON THE SOLUTION OF OBLIQUE-ANGLED TRIANGLES.

44. *Let the three Sides be given.* (a, b, c.)

The *Angles* may be determined from one of the formulæ (vii.), (viii.), (ix.), (x.).

45. *Let the three Angles be given.* (A, B, C.)

The *Sides* may be determined from one of the formulæ (xi.), (xii.), (xiii.), (xiv.).

46. *Let two Sides and the included Angle be given.* (a, C, b.)

By Napier's first and second Analogies, (xv.) and (xvi.),

$$\left\{ \begin{aligned} \tan \frac{1}{2} (A + B) &= \frac{\cos \frac{1}{2} (a - b)}{\cos \frac{1}{2} (a + b)} \cdot \cot \frac{1}{2} C, \\ \tan \frac{1}{2} (A - B) &= \frac{\sin \frac{1}{2} (a - b)}{\sin \frac{1}{2} (a + b)} \cdot \cot \frac{1}{2} C, \end{aligned} \right.$$

$\frac{1}{2} (A + B)$, and $\frac{1}{2} (A - B)$, are determined;

$$\therefore \left\{ \begin{aligned} A &= \frac{1}{2} (A + B) + \frac{1}{2} (A - B), \\ B &= \frac{1}{2} (A + B) - \frac{1}{2} (A - B), \end{aligned} \right\} \text{ are known.}$$

And *A* and *B* being known, *c* is found from

$$\sin c = \sin a \cdot \frac{\sin C}{\sin A}.*$$

* The easiest practical method of solving this case is by letting fall from *A* a perpendicular (*AD*) on *BC* or *BC* produced either way, and then determining the right-angled triangles *ABD* and *ACD*. Supposing *D* lies between *B* and *C*, then

$$\tan CD = \cos C \tan b, \text{ which gives } CD.$$

$$\text{And } \therefore DB, = a - CD, \text{ is known.}$$

$$\text{Also } \cos c = \cos DB \cos AD = \cos DB \cdot \frac{\cos b}{\cos CD}.$$

47. To determine c independently of A and B , by forms adapted to logarithmic computation.

$$\begin{aligned} \cos c &= \cos a \cos b + \sin a \sin b \cos C \\ &= \cos b (\cos a + \sin a \tan b \cos C). \end{aligned}$$

Let θ be an angle such that $\tan \theta = \tan b \cos C \dots\dots\dots(1)$.

$$\begin{aligned} \text{Then } \cos c &= \cos b \left(\cos a + \sin a \frac{\sin \theta}{\cos \theta} \right) \\ &= \frac{\cos b}{\cos \theta} \cdot \cos (a - \theta) \dots\dots\dots (2). \end{aligned}$$

From (1), $L \tan \theta = L \tan b + L \cos C - 10$; which gives θ .

(2), $L \cos c = L \cos b + L \cos (a - \theta) - L \cos \theta$; which gives c .

[On comparing this solution with that given in the foot-note to Art. 46, it will be found to be identically the same, if CD be represented by θ .]

48. Let two Angles and the included Side be given. (A, c, B .)

From Napier's third and fourth Analogies, (xvii.), (xviii.),

$$\begin{cases} \tan \frac{1}{2} (a + b) = \frac{\cos \frac{1}{2} (A - B)}{\cos \frac{1}{2} (A + B)} \cdot \tan \frac{1}{2} c, \\ \tan \frac{1}{2} (a - b) = \frac{\sin \frac{1}{2} (A - B)}{\sin \frac{1}{2} (A + B)} \cdot \tan \frac{1}{2} c, \end{cases}$$

$\frac{1}{2} (a + b)$ and $\frac{1}{2} (a - b)$ are determined;

$$\therefore \left. \begin{cases} a = \frac{1}{2} (a + b) + \frac{1}{2} (a - b), \\ b = \frac{1}{2} (a + b) - \frac{1}{2} (a - b), \end{cases} \right\} \text{ are known.}$$

And a and b being known, C is found from

$$\sin C = \sin A \cdot \frac{\sin c}{\sin a}.$$

49. To find C , independently of a and b , by forms adapted to logarithmic computation.

$$\cos C - \cos c \sin A \sin B - \cos A \cos B;$$

$$\begin{aligned} \therefore 2 \cos^2 \frac{1}{2} C - 1 &= (1 - 2 \sin^2 \frac{1}{2} c) \sin A \sin B - \cos A \cos B \\ &= -\cos(A + B) - 2 \sin^2 \frac{1}{2} c \sin A \sin B; \end{aligned}$$

$$\therefore 2 \cos^2 \frac{1}{2} C = 1 - \cos(A + B) - 2 \sin^2 \frac{1}{2} c \sin A \sin B;$$

$$\therefore \cos^2 \frac{1}{2} C = \sin^2 \frac{1}{2} (A + B) - \sin^2 \frac{1}{2} c \sin A \sin B.$$

Now $\sin^2 \frac{1}{2} c \sin A \sin B$ is necessarily positive, and less than unity; wherefore there is an angle θ such that

$$\sin^2 \theta = \sin^2 \frac{1}{2} c \sin A \sin B \dots \dots \dots (1);$$

$$\therefore \cos^2 \frac{1}{2} C = \sin^2 \frac{1}{2} (A + B) - \sin^2 \theta$$

$$= \sin \left\{ \frac{1}{2} (A + B) + \theta \right\} \sin \left\{ \frac{1}{2} (A + B) - \theta \right\}. \text{ Pl. Trig. Art. 54.} \dots \dots (2).$$

From (1), $L \sin \theta = \frac{1}{2} (L \sin A + L \sin B) + L \sin \frac{1}{2} c - 10$; which gives θ .

From (2), $L \cos \frac{1}{2} C = \frac{1}{2} [L \sin \left\{ \frac{1}{2} (A + B) + \theta \right\} + L \sin \left\{ \frac{1}{2} (A + B) - \theta \right\}]$; which gives C .

50. Let two Angles and a Side opposite to one of them be given. (A, B, a .)

$$\sin b = \sin a \cdot \frac{\sin B}{\sin A};$$

$$\text{from (xv.), } \tan \frac{1}{2} C = \frac{\cos \frac{1}{2} (a - b)}{\cos \frac{1}{2} (a + b)} \cdot \cot \frac{1}{2} (A + B);$$

$$\text{from (xvii.), } \tan \frac{1}{2} c = \frac{\cos \frac{1}{2} (A + B)}{\cos \frac{1}{2} (A - B)} \cdot \tan \frac{1}{2} (a + b).$$

And b having been determined from the first of these equations, C and c may be found from the other two.

51. To determine C and c independently of b , by forms adapted to logarithmic computation. (A, B, a .)

$$\begin{aligned} \cos A &= \cos a \sin B \sin C - \cos B \cos C \\ &= \cos B (\cos a \tan B \sin C - \cos C). \end{aligned}$$

Let θ be an angle such that $\cot \theta = \cos a \tan B \dots\dots\dots(1)$;

$$\begin{aligned} \therefore \cos A &= \cos B \left(\frac{\cos \theta}{\sin \theta} \cdot \sin C - \cos C \right) \\ &= \frac{\cos B}{\sin \theta} \cdot \sin (C - \theta) \dots\dots\dots(2). \end{aligned}$$

From (1), $L \cot \theta = L \cos a + L \tan B - 10$; which gives θ .

(2), $L \sin (C - \theta) = L \cos A + L \sin \theta - L \cos B$; which gives $C - \theta$, and thence C .

Again, from (v.), $\sin B \cot A = \cot a \sin c - \cos c \cos B$
 $= \cot a (\sin c - \cos B \tan a \cos c),$

and, if ϕ be such that $\tan \phi = \cos B \tan a, \dots\dots\dots(1)$,

$$\therefore \sin B \cot A = \frac{\cot a}{\cos \phi} \cdot \sin (c - \phi);$$

$$\therefore \frac{\sin (c - \phi)}{\cos \phi} \cdot \frac{1}{\tan \phi} = \sin B \cot A \tan a \cdot \frac{1}{\cos B \tan a},$$

$$\text{or } \frac{\sin (c - \phi)}{\sin \phi} = \frac{\tan B}{\tan A} \dots\dots\dots(2).$$

From (1), $L \tan \phi = L \cos B + L \tan a - 10$; which gives ϕ .

(2), $L \sin (c - \phi) = L \sin \phi + L \tan B - L \tan A$; which gives $c - \phi$, and thence c .*

[The method used in Art. 49 might have been employed here.]

* If from C an arc CD be drawn cutting AB , or AB produced either way, at right angles in the point D , the solution in the text is the same as determining the right-angled triangles CBD and ACD , the angle BCD being represented by θ , and the arc BD by ϕ .

52. Let two Sides and an Angle opposite to one of them be given. (a, b, A.)

$$\sin B = \sin A \cdot \frac{\sin b}{\sin a},$$

$$\tan \frac{1}{2} C = \frac{\cos \frac{1}{2}(a-b)}{\cos \frac{1}{2}(a+b)} \cdot \cot \frac{1}{2}(A+B);$$

$$\tan \frac{1}{2} c = \frac{\cos \frac{1}{2}(A+B)}{\cos \frac{1}{2}(A-B)} \cdot \tan \frac{1}{2}(a+b).$$

And B having been determined from the first equation, C and c are found from the other two.

53. To determine C and c independently of B , by forms adapted to logarithmic computation.

$$\begin{aligned} \text{From (v.) } \cot a \sin b &= \cos b \cos C + \sin C \cot a \\ &= \cos b \left(\cos C + \frac{\cot a}{\cos b} \cdot \sin C \right); \end{aligned}$$

$$\text{Whence, if } \theta \text{ be such that } \cot \theta = \frac{\cot a}{\cos b} \dots\dots\dots(1),$$

there is got, $\sin(C + \theta) = \cot a \tan b \sin \theta \dots\dots\dots(2);$
from which two equations θ and C can be found.

$$\begin{aligned} \text{Again, } \cos a &= \cos b \cos c + \sin b \sin c \cos A \\ &= \cos b (\cos A \tan b \sin c + \cos c); \end{aligned}$$

$$\text{Whence, if } \phi \text{ be such that } \cot \phi = \cos A \tan b \dots\dots\dots(1),$$

there is got, $\sin(c + \phi) = \frac{\cos a \sin \phi}{\cos b} \dots\dots\dots(2),$

and from these two equations ϕ and c can be found*.

* If $\theta = \frac{1}{2}\pi - \theta'$, $\tan \theta' = \frac{\cot a}{\cos b}$, and θ' and $(C - \theta')$ are the segments of the angle C made by a perpendicular let fall from C on AB . Also if $\phi = \frac{1}{2}\pi - \phi'$, ϕ' and $(c - \phi')$ will be the segments of AB .

54. In (50) the triangle has apparently been solved when two angles and a side opposite to one of them are given; and in (52) the triangle has apparently been solved from two sides and the angle opposite to one of them.

It is, however, doubtful in some cases, in determining one of the quantities A, B, a, b from the equation $\frac{\sin A}{\sin B} = \frac{\sin a}{\sin b}$ when the other three are given, whether the angle to be taken should be less or greater than $\frac{1}{2}\pi$. The ambiguous cases will now be distinguished from those which are not so.

(1) Given A, B, b , to determine in what cases a may be found from the equation,

$$\frac{\sin a}{\sin b} = \frac{\sin A}{\sin B}.$$

I. Let $A + B$ be greater than two right angles, or π .

Then since, (34), $\frac{1}{2}(a + b)$ and $\frac{1}{2}(A + B)$ are of like affection,

$$\therefore a + b \text{ is } > \pi.$$

If therefore b be $< \frac{1}{2}\pi$, a must be $> \frac{1}{2}\pi$. In this case, therefore, a may be determined, and the triangle may be solved.

But if b be $> \frac{1}{2}\pi$, the condition $a + b > \pi$ affords no means of determining whether a be $>$ or $< \frac{1}{2}\pi$.

II. If $A + B$ be $< \pi$, then $a + b$ is $< \pi$; and a is therefore $< \frac{1}{2}\pi$ if b be $> \frac{1}{2}\pi$, but cannot be determined if b be $< \frac{1}{2}\pi$.

III. If $A + B = \pi$; then, (by xv.), $a + b = \pi$, and therefore $a = \pi - b$.

Whence it appears that, (A, B, b being given),

(1), when $A + B > \pi$, and $b < \frac{1}{2}\pi$; then a is greater than $\frac{1}{2}\pi$,

(2), $A + B < \pi$, and $b > \frac{1}{2}\pi$; then a is less than $\frac{1}{2}\pi$,

(3), $A + B = \pi$; then $a = \pi - b$;

and in no other case can a be found, and the triangle determined, from these data.

(2) Given two Sides, and an Angle opposite to one of them, to determine when the remaining parts of the triangle may be determined. (a, b, B.)

If $a + b > \pi$; then, as before, $A + B > \pi$, and $\therefore A > \pi - B$.
 If $a + b < \pi$; then, as before, $A + B < \pi$, and $\therefore A < \pi - B$.

Whence it is collected, as in the last proposition, that

(1), when $a + b > \pi$, and $B < \frac{1}{2}\pi$; then A is greater than $\frac{1}{2}\pi$,

(2), $a + b < \pi$, and $B > \frac{1}{2}\pi$; then A is less than $\frac{1}{2}\pi$,

(3), $a + b = \pi$; then $A = \pi - B$;


and in no other case can A be determined.

55. By letting fall a perpendicular from any angle upon the side opposite, an oblique-angled triangle may be divided into two right-angled triangles, which in most cases may be solved by Napier's Rules with not less facility than by the methods just given. The very same ambiguities, however, will arise (40) when this construction is used for the determination of the triangle as have been pointed out in (54).

56. PROB. To find the radius of the small circle described about a given triangle in terms of the Angles of the triangle.

Let ABC be the triangle; bisect CA and CB in D and E , and draw from those points at right angles to AC and to CB arcs intersecting in P . Join PA , PB , and PC .

Then, from the right-angled triangles PCD and PAD ,



$$\cos PC = \cos PD \cos DC = \cos PD \cos DA = \cos PA;$$

$$\therefore PA = PC. \text{ Similarly, } PB = PC.$$

Therefore P is the pole of the circumscribing circle.

$$\text{Now } \cos PBE = \cot PB \tan BE = \cos PB \tan \frac{1}{2} a,$$

$$\therefore \cot PA, = \cot PB, = \cos PBE \cot \frac{1}{2} a ;$$

and since PAC, PCB, PBA are isosceles triangles,

$$\therefore 2 \angle PAC + 2 \angle PAB + 2 \angle PBE = A + B + C = 2S' ;$$

$$\therefore \angle PBE = S' - (\angle PAC + \angle PAB) = S' - A.$$

$$\text{Also, by (xiii.), } \cot \frac{1}{2} a = \sqrt{\frac{\cos(S' - B) \cos(S' - C)}{-\cos S' \cos(S' - A)}} ;$$

$$\therefore \cot PA = \sqrt{\frac{\cos(S' - A) \cos(S' - B) \cos(S' - C)}{-\cos S'}}$$

57. PROB. *To determine the radius of the circumscribing circle in terms of the Sides of the triangle.*

$$\text{As in (56), } \cot PA = \cot \frac{1}{2} a \cos PBE ;$$

And as before,

$$\cos PBE = \cos(S' - A) = \cos \frac{1}{2} \{(B + C) - A\}$$

$$= \cos \frac{1}{2} (B + C) \cos \frac{1}{2} A + \sin \frac{1}{2} (B + C) \sin \frac{1}{2} A$$

$$= \frac{\sin \frac{1}{2} A \cos \frac{1}{2} A}{\cos \frac{1}{2} a} \cdot \{\cos \frac{1}{2} (b + c) + \cos \frac{1}{2} (b + c)\}, \text{ by (xx.) and (xix.),}$$


$$= \frac{\sin A}{\cos \frac{1}{2} a} \cdot \cos \frac{1}{2} b \cos \frac{1}{2} c ;$$

$$\therefore \tan PA = \frac{\tan \frac{1}{2} a}{\cos PBE} = \frac{\sin \frac{1}{2} b}{\cos \frac{1}{2} b \cos \frac{1}{2} c \sin A}$$

$$= \frac{2 \sin \frac{1}{2} a \sin \frac{1}{2} b \sin \frac{1}{2} c}{\sqrt{\{\sin S \sin(S - a) \sin(S - b) \sin(S - c)\}}}, \text{ by (x.)}$$

58. PROB. To find the radius of the circle inscribed in a given triangle in terms of the Sides of the triangle.

Let ABC be the triangle; bisect $\angle A$ and $\angle C$ by AP and CP , arcs of great circles meeting in P ; and from P draw the arcs PD , PE , PF perpendicular to the sides.



Then it⁴ may be proved that $PE = PD = PF$; and therefore P is the pole of the inscribed circle. Also it may be shewn that $CE = S - (AF + FB) = S - c$, and thence that

$$\begin{aligned} \text{Tan } PE &= \sin CE \tan PCE = \sin (S - c) \tan \frac{1}{2} C \\ &= \sqrt{\frac{\sin (S - a) \sin (S - b) \sin (S - c)}{\sin S}}. \end{aligned}$$

59. PROB. To determine the radius of the inscribed circle in terms of the Angles of the triangle.

As in (58), $\text{Tan } PE = \sin CF \cdot \tan \frac{1}{2} C$,

$$\begin{aligned} \text{and } \sin CE &= \sin (S - c) = \sin \frac{1}{2} \{(a + b) - C\} \\ &= \sin \frac{1}{2} (a + b) \cos \frac{1}{2} c - \cos \frac{1}{2} (a + b) \sin \frac{1}{2} c; \end{aligned}$$

Whence, by means of GAUSS' THEOREM, (xxii.) and (xx.), it may be proved that

$$\text{Cot } PE = \frac{2 \cos \frac{1}{2} A \cos \frac{1}{2} B \cos \frac{1}{2} C}{\sqrt{\{-\cos S' \cos (S' - A) \cos (S' - B) \cos (S' - C)\}}}.$$


CHAPTER V.

ON THE AREAS OF SPHERICAL TRIANGLES, AND THE SOLUTION
OF TRIANGLES WHOSE SIDES ARE SMALL COMPARED
WITH THE RADIUS OF THE SPHERE.

60. DEF. THE portion of the surface of a sphere which is contained within two great semicircles is called a *lune*.

61. *To find the Area of a Lune.*

If $ACBDA$, $ADBEA$, $AEBFA$ be lunes each having the same angle at A , any one may be placed on another so as to coincide, and therefore be equal with it. Thus if the angle CAD be repeated any number of times, the area $ACBDA$ will be repeated the same number of times. Wherefore the Area of a Lune varies as its angle.




$$\therefore \frac{\text{Area of the lune whose angle is } A^\circ}{\text{Area of the sphere (whose angle is } 360^\circ)} = \frac{A}{360};$$

And Area of a sphere = $4\pi r^2$, if r = radius of the sphere;
(See Hymers' Integral Calculus.)

$$\therefore \text{Area of the Lune whose angle is } A^\circ = \frac{A}{360} \cdot 4\pi r^2 = \frac{A}{180} \cdot 2\pi r^2.$$

62. To find the Area of a Spherical Triangle.

Let ABC be a triangle upon a hemisphere $ABDEGA$ (14, Cor. 2); and let AC , BC be produced until they meet again in F , which is a point on the side of the sphere turned from the spectator.



Then since CDF = a semicircle = ACD ,

$$\therefore RF = AC;$$

Similarly, $FE = CB$; and $\angle DFE = \angle ACB$;

$$\therefore \triangle DFE = \triangle ACB \text{ in every respect.}$$

Now Σ (= area of $\triangle ABC$)

$$= \text{surface of hemisphere} - BIIDC - AGE C - DCE$$

$$= 2\pi r^2 - (\text{lune } AIIDA - \Sigma) - (\text{lune } BGECB - \Sigma) - (\text{lune } CDFEC - \Sigma)$$

$$= 2\pi r^2 \left(1 - \frac{A}{180} - \frac{B}{180} - \frac{C}{180} \right) + 3\Sigma; \quad (61)$$

$$\therefore \Sigma = (A + B + C - 180) \cdot \frac{\pi r^2}{180}$$

$$= \frac{E}{180} \cdot \pi r^2, \quad \text{if } E^0 = A^0 + B^0 + C^0 - 180^0.$$

DEF. The quantity $A^0 + B^0 + C^0 - 180^0$, by which the sum of the degrees in the angles of the spherical triangle exceeds 180^0 , is called the *Spherical Excess* of the triangle.

The Spherical Excess is generally written thus, $E = A + B + C - 180^0$.

COR. 1. Hence for all triangles described on the same sphere, $\Sigma \propto A^0 + B^0 + C^0 - 180^0$, and on this account the Spherical Excess has been taken as the measure of the surface of a triangle.

COR. 2. To find the Area of a Spherical Polygon. Divide the Polygon into as many triangles as it has sides, by means of arcs of great circles drawn from each of the angular points to any point within the polygon. Let n be the number of the sides of the polygon.

Then Area = area of the n triangles = $\frac{\pi r^2}{180} \times$ (number of degrees in the angles of the triangles - $n \cdot 180$)

$$= \frac{\pi r^2}{180} \cdot \{\text{number of degrees in the angles of the polygon} - (n - 2) 180\}.$$

63. CAGNOLI'S THEOREM. *To shew, that if E be the Spherical Excess,*

$$\text{then } \sin \frac{1}{2} E = \frac{\sqrt{\{\sin S \sin (S-a) \sin (S-b) \sin (S-c)\}}}{2 \cos \frac{1}{2} a \cos \frac{1}{2} b \cos \frac{1}{2} c}.$$

$$\begin{aligned} \sin \frac{1}{2} E &= \sin \frac{1}{2} (A+B+C-180^\circ) = \sin \left\{ \frac{1}{2} (A+B) - \frac{1}{2} (180^\circ - C) \right\} \\ &= \sin \frac{1}{2} (A+B) \sin \frac{1}{2} C - \cos \frac{1}{2} (A+B) \cos \frac{1}{2} C. \end{aligned}$$

$$\text{And by (xix.), } \sin \frac{1}{2} (A+B) = \frac{\cos \frac{1}{2} C}{\cos \frac{1}{2} c} \cdot \cos \frac{1}{2} (a-b),$$

$$\text{(xx.), } \cos \frac{1}{2} (A+B) = \frac{\sin \frac{1}{2} C}{\cos \frac{1}{2} c} \cdot \cos \frac{1}{2} (a+b).$$

$$\begin{aligned} \therefore \sin \frac{1}{2} E &= \left\{ \cos \frac{1}{2} (a-b) - \cos \frac{1}{2} (a+b) \right\} \cdot \frac{\sin \frac{1}{2} C \cos \frac{1}{2} C}{\cos \frac{1}{2} c} = \frac{\sin \frac{1}{2} a \sin \frac{1}{2} b}{\cos \frac{1}{2} c} \cdot \sin C \\ &= \frac{\sin \frac{1}{2} a \sin \frac{1}{2} b}{\cos \frac{1}{2} c} \cdot \frac{2}{\sin a \sin b} \cdot \sqrt{\{\sin S \sin (S-a) \sin (S-b) \sin (S-c)\}} \\ &= \frac{\sqrt{\{\sin S \sin (S-a) \sin (S-b) \sin (S-c)\}}}{2 \cos \frac{1}{2} a \cos \frac{1}{2} b \cos \frac{1}{2} c}. \end{aligned}$$

64. LLHULLIER'S THEOREM. *To shew that*

$$\tan \frac{1}{4} E = \sqrt{\{\tan \frac{1}{2} S \tan \frac{1}{2} (S-a) \tan \frac{1}{2} (S-b) \tan \frac{1}{2} (S-c)\}}.$$

By Pl. Trig. (51),

$$\frac{\sin \frac{1}{2} (A+B) - \sin \frac{1}{2} (180^\circ - C)}{\cos \frac{1}{2} (A+B) + \cos \frac{1}{2} (180^\circ - C)} = \frac{\sin \frac{1}{4} (A+B+C-180^\circ)}{\cos \frac{1}{4} (A+B+C-180^\circ)} = \tan \frac{1}{4} E;$$

$$\begin{aligned} \therefore \tan \frac{1}{4} E &= \frac{\sin \frac{1}{2} (A+B) - \cos \frac{1}{2} C}{\cos \frac{1}{2} (A+B) + \sin \frac{1}{2} C} = \frac{\cos \frac{1}{2} (a-b) - \cos \frac{1}{2} c}{\cos \frac{1}{2} (a+b) + \cos \frac{1}{2} c} \cdot \frac{\cos \frac{1}{2} C}{\sin \frac{1}{2} C} \text{ by (xix.) and (xx.)} \\ &= \frac{\sin \frac{1}{4} (c+a-b) \sin \frac{1}{4} (c+b-a)}{\cos \frac{1}{4} (a+b+c) \cos \frac{1}{4} (a+b-c)} \cdot \sqrt{\frac{\sin S \sin (S-c)}{\sin (S-a) \sin (S-b)}}, \text{ by (ix.)} \\ &= \sqrt{\{\tan \frac{1}{2} S \tan \frac{1}{2} (S-a) \tan \frac{1}{2} (S-b) \tan \frac{1}{2} (S-c)\}}, \end{aligned}$$

by expressing the sines under the root in terms of the sines and cosines of half the angles.

65. [There are some other values of E , the Spherical Excess, which are symmetrical functions of the sides of the triangle; but not being adapted to logarithmic computation, the demonstrations of them will not be given. The expressions will be found among the Examples which are given at the end of the book, and may be proved without much trouble by Gauss' formulæ.]

66. In the following Articles x, y, z will generally be taken to represent the *lengths* of the arcs opposite to the angles A, B, C of the spherical triangle ABC whose sides are small compared with the radius of the sphere on which it is described. Also, the angles of the *plane* triangle whose sides are x, y, z , will be represented by A', B', C' ; and the angles which are, in practice, observed for the angles of the spherical triangle ABC , by A_1, B_1, C_1 ,—the errors α, β, γ being made at the respective observations; so that $A = A_1 + \alpha, \quad B = B_1 + \beta, \quad C = C_1 + \gamma$.

67. *The Area of a triangle whose sides are small compared with the radius of the sphere being approximately known, required the number of seconds in the Spherical Excess of the triangle.*

Let the triangle be described on the surface of the Earth, and let the Earth be supposed to be a sphere.

Let r = number of linear feet in the Earth's radius,

$$n = \text{square feet in the area of the triangle} = \frac{E}{180} \cdot \pi r^2, \quad (62);$$

$$\therefore n = E \cdot 60 \cdot 60 \cdot \frac{\pi r^2}{180 \cdot 60 \cdot 60}.$$

Now $180 \times$ length of an arc of 1° = semi-circumference of the circle = πr ;

$$\therefore \frac{\pi r}{180} = \text{length of } 1^\circ, \text{ in feet,}$$

$$= 60859 \cdot 1 \times 6 \text{ feet, by actual measurement;}$$

$$\therefore r = \frac{180}{\pi} \times 60859 \cdot 1 \times 6;$$

$$\begin{aligned} \therefore n &= E \cdot 60 \cdot 60 \cdot \frac{180}{\pi \cdot 60 \cdot 60} (60859 \cdot 1 \times 6)^2 \\ &= E \cdot 60 \cdot 60 \cdot \frac{180 \cdot (6085 \cdot 91)^2}{3 \cdot 141592}; \end{aligned}$$

$$\begin{aligned} \therefore I_{10}(E \cdot 60 \cdot 60) &= I_{10}n - (I_{10}180 + 2 \cdot I_{10}6085 \cdot 91 - I_{10}3 \cdot 141592) \\ &= I_{10}n - 9 \cdot 3267736 \dots \dots \dots (1), \end{aligned}$$

whence $E \cdot 60 \cdot 60$, the number of *seconds* in the Spherical Excess, may be found.

[NOTE. This reasoning being general, if the radius (r) of the sphere be known, the number of seconds in the Spherical Excess may always be determined from

$$n = (E \cdot 60 \cdot 60) \cdot \left[\frac{\pi r^2}{180 \cdot 60 \cdot 60} \right].$$

68. The number n may be determined approximately in the following manner.

n = area of the spherical triangle, in square feet.

= area (nearly) of the plane triangle whose sides are x, y, z , and whose angles are the observed angles A_1, B_1, C_1 .

$$= \frac{1}{2} y \cdot z \cdot \sin A_1, \quad \text{or} \quad = \frac{1}{2} x^2 \cdot \frac{\sin B_1 \sin C_1}{\sin (B_1 + C_1)} \cdot \text{Pl. Trig. (91).}$$

69. Ex. *If the observed angles of the spherical triangle ABC be $A_1 = 42^\circ, 2', 32''$, $B_1 = 67^\circ, 55', 39''$, $C_1 = 70^\circ, 1', 48''$, and the side opposite to the angle A be 27404.2 feet, required the number of seconds in the sum of the errors made in observing the angles.*

Here the *apparent* Spherical Excess is

$$A + B + C - 180^\circ = 179^\circ, 59' - 180^\circ = -1''.$$

* General Roy, in the Trigonometrical Survey of England, approximately determined the area of the triangle to a sufficient degree of accuracy, by laying down on paper the base and the observed angles at the base, and measuring the perpendicular from the vertex to the base by the compasses on a scale.

Now representing the given side by a ,

$$n = \frac{1}{2} a^2 \cdot \frac{\sin B_1 \sin C_1}{\sin (B_1 + C_1)},$$

$$\text{and } I_{10}(E. 60. 60) = I_{10} \left\{ \frac{1}{2} a^2 \cdot \frac{\sin B_1 \sin C_1}{\sin (B_1 + C_1)} \right\} - 9.3267736$$

$$= I_{10} \left\{ \frac{1}{2} (274.4.2)^2 \cdot \frac{\sin 67^\circ. 55'. 39'' \times \sin 70^\circ. 1'. 48''}{\sin 137^\circ. 57'. 27''} \right\} - 9.3267736.$$

$$\text{Add, } 2 \times I_{10} 274.4.2 = 2 \times 4.4378172 = 8.8756344$$

$$I \sin 67^\circ. 55'. 39'' = 9.9669434$$

$$L \sin 70^\circ. 1'. 48'' = 9.9730685$$

$$28.8156163$$

$$\text{Subtract, } I_{10} 2 = .3010300$$

$$L \sin 137^\circ. 57'. 27'',$$

$$\text{or } L \sin 42^\circ. 2'. 33'' = 9.8258684$$

$$10.$$

$$9.3267736$$

$$29.4536720$$

$$\bar{1} 3619743$$

$$= I_{10}(.23), \text{ nearly;}$$

and therefore the *computed* Spherical Excess (which, for all practical purposes, may be *supposed* to be the *real* Spherical Excess), is $.23''$.

Hence it appears that the whole error of observation, viz. *real* Spherical Excess - *apparent* Spherical Excess, is $.23'' - (-1'')$, or $1''.23$, which the observer must add to the three observed angles, A_1, B_1, C_1 , in such proportions as his judgment may direct. (See the next Article.)

70. To shew how the observed angles of a Spherical Triangle whose sides are small compared with the radius of the sphere may be best freed from the errors of observation.

Let A, B, C , be the *real* angles of the triangle,

A_1, B_1, C_1 , the *observed* angles,

α, β, γ , the errors made in observing A, B, C , respectively.

$$\begin{aligned} \text{So that } \alpha + \beta + \gamma &= (A - A_1) + (B - B_1) + (C - C_1); \\ &= (A + B + C - 180^\circ) - (A_1 + B_1 + C_1 - 180^\circ) \\ &= \text{real Spherical Excess} - \text{computed Spherical Excess.} \end{aligned}$$

Now if a value of the Spherical Excess which differs very slightly from the real value, be found by the method of the last Article, the above equation will give the *sum* of the errors of observation which have been made. The *distribution*, however, of this sum—(that is, the determining what part of the whole error is to be assigned to each angle individually)—must evidently be left to the judgment of the observer, who, from knowing the state of the atmosphere at the times of the observations, may be able to form an opinion how far his optical observations can be depended on, and may then assign to each of the observed angles such a portion of the whole error as he thinks will be the most likely to lead to a correct solution of the triangle.

[If $\alpha + \beta + \gamma$ (the sum of the errors of observation) be found by means of (67), and thence the several angles A, B, C be determined by the arbitrary assignment of the several parts of this whole error to each of the observed angles A_1, B_1, C_1 , the *sum* of the errors of observation may be supposed to be got rid of. Yet it is highly probable that the judgment of the observer has not been *absolutely* correct in this arbitrary assignment of the parts of the whole error. The following theorem will point out what relation the sides of the triangle ought to bear to each other, in order that the small quantities by which the *corrected* angles differ from the *real* angles of the triangle may have the least possible effect in producing errors when the other two sides of the triangle have to be determined from a measured side and the *corrected* angles.]

71. *Having given the Corrected Angles and one Side of a Spherical Triangle whose sides are small compared with the radius of the sphere, required the relation which the Sides of the Triangle ought to bear to each other in order that the other sides may be determined from these data with the least probable amount of error.*

Let A, B, C be the real angles of the triangle, and x, y, z the sides respectively opposite to them; α', β', γ' the errors of the corrected angles; therefore $A + \alpha', B + \beta', C + \gamma'$ are the corrected angles.

Then, since the Spherical Excess (and therefore the sum of the real angles of the triangle) is supposed to be known *exactly*, the sum of the corrected angles is known exactly, and therefore the sum of the separate errors α', β', γ' must vanish;

$$\therefore \alpha' + \beta' + \gamma' = 0, \quad \text{or } \alpha' = -(\beta' + \gamma').$$

[Throughout this investigation, powers of α', β', γ' of any order above the first will be neglected.]

Now, considering the spherical triangle to be very nearly a plane triangle,

$$\begin{aligned} z &= x \cdot \frac{\sin(C + \gamma')}{\sin(A + \alpha')} = x \cdot \frac{\sin(C + \gamma')}{\sin\{A - (\beta' + \gamma')\}} \\ &= x \cdot \frac{\sin C \cos \gamma' + \cos C \sin \gamma'}{\sin A \cos(\beta' + \gamma') - \cos A \sin(\beta' + \gamma')} \\ &= x \cdot \frac{\sin C}{\sin A} \cdot \frac{1 + \gamma' \cot C}{1 - (\beta' + \gamma') \cot A}, \text{ nearly,} \\ &= x \cdot \frac{\sin C}{\sin A} \cdot \{1 + \gamma' \cot C\} \{1 + (\beta' + \gamma') \cot A\} \\ &= x \cdot \frac{\sin C}{\sin A} \cdot \{1 + \gamma' (\cot A + \cot C) + \beta' \cot A\} \\ &= x \cdot \frac{\sin C}{\sin A} + x \left\{ \gamma' \cdot \frac{\sin C}{\sin A} \cdot \frac{\sin(A + C)}{\sin A \sin C} + \beta' \cdot \frac{\cos A \sin C}{\sin^2 A} \right\}. \end{aligned}$$

And $\sin B = \sin(\pi - B) = \sin(A + C)$ nearly;

Therefore the error in the value of z , (or $z - x \cdot \frac{\sin C}{\sin A}$), is

$$x \left\{ \gamma' \cdot \frac{\sin B}{\sin^2 A} + \beta' \cdot \frac{\cos A \sin C}{\sin^2 A} \right\} \dots \dots \dots (1).$$

Similarly the error in the value of y is

$$\beta' \left\{ \beta' \cdot \frac{\sin C}{\sin^2 A} + \gamma' \cdot \frac{\cos A \sin B}{\sin^2 A} \right\} \dots \dots \dots (2).$$

The question is now reduced to determine what are the values of A , B , and C which make *both* the quantities (1) and (2) the least possible.

Since $\alpha' + \beta' + \gamma' = 0$, two of the three quantities α' , β' , γ' must be of the same sign and one of the contrary sign. Wherefore the probability is that any particular two, as β' and γ' , are of different signs. If then β' and γ' be of different signs, the expressions (1) and (2) are diminished in magnitude by giving $\cos A$ a positive value*; that is, by supposing A to be less than a right angle.

And if it is further supposed,—that the errors β' and γ' , though different in sign, are yet nearly equal in magnitude,—it is clear that (1) and (2) satisfy this hypothesis if B be nearly equal to C .

This conclusion is therefore arrived at,—that there is the greatest probability of a small spherical triangle having been correctly solved from three observed angles and a measured side, if the angle opposite to the known side be less than a right angle and the other two sides be nearly equal. And these conditions will be best fulfilled for a series of triangles if each triangle be nearly equilateral.

72. [It may be as well to recount the gratuitous suppositions made in the last Article.

- I. The Spherical Excess has been accurately determined.
- II. The Errors β and γ' are of different signs.
- III. These errors are of the same magnitude.

* It is evidently more advantageous in determining a series of triangles from one another, that the errors should be inconsiderable and equally diffused through all, than that any one calculated side, by differing much from its real value, should affect with considerable errors all the triangles successively determined from it. If, β' and γ' being of different signs, $\cos A$ become negative, the errors (1) and (2) are increased, and considerable inaccuracies might so be introduced into the calculations.

Now for any *particular* triangle it is very probable that some of these suppositions may not be true. It may happen that none of them may be correct. In very few cases indeed will they *all* be fulfilled.

With respect to the Spherical Excess, it may generally be supposed to be known accurately.

With respect to the signs of β' and γ' , since two of the quantities α' , β' , γ' , are of the same, and the third is of the contrary sign, the probability of β' and γ' being of *different* signs is twice as great as the probability that they are of the *same* sign. And if they be of different signs, then (1) and (2) of (71) shew that the errors will not be so great if $\text{Cos } A$ be positive, as if it be negative. This consideration renders it advisable that each of the angles of the triangle should be less than a right angle; which will be the case if the triangle be nearly equilateral.

But if the third hypothesis be admitted, since α' vanishes in this case, the angle A is supposed (which is highly improbable) to have been determined with mathematical exactness. Besides, it is very unlikely that an observer after making an error $+\beta'$ in observing an angle, should make an error $-\beta'$ (for $\gamma' = -\beta'$) in observing, with the same instrument, an angle C which is nearly of the same magnitude as B ; and again, using still the same instrument, and observing a third angle A , (nearly equal to B or to C), that he should make no error at all.]

73. **LEGENDRE'S THEOREM.** *If each of the angles of a Spherical Triangle whose sides are small when compared with the radius of the sphere be diminished by one third of the Spherical Excess, the triangle may be solved as a Plane Triangle, whose sides are equal to the sides of the Spherical Triangle, and whose angles are these reduced angles.*

Let x, y, z be the *lengths* of the sides respectively opposite to the angles A, B, C of a small *spherical* triangle, and A', B', C' the angles of that *plane* triangle ($A'B'C'$) whose sides are x, y, z .

$$\begin{aligned} \text{Then } \sin A + \sin A' &= \frac{\sin a}{\sin b} \cdot \sin B + \frac{x}{y} \cdot \sin B' \\ &= \frac{x}{y} \cdot (\sin B + \sin B') \text{ nearly;} \end{aligned}$$

$$\therefore 2 \sin \frac{1}{2}(A + A') \cos \frac{1}{2}(A - A') = \frac{x}{y} \cdot 2 \sin \frac{1}{2}(B + B') \cos \frac{1}{2}(B - B');$$

$$\therefore \cos \frac{1}{2}(A - A') = \frac{x}{y} \cdot \frac{\sin B'}{\sin A'} \cdot \cos \frac{1}{2}(B - B'), \text{ nearly;}$$

$$\therefore \cos \frac{1}{2}(A - A') = \cos \frac{1}{2}(B - B'), \text{ or } A - A' = B - B'.$$

Similarly $A - A' = C - C'$.

Now $E = A + B + C - 180^\circ$,

and $0 = A' + B' + C' - 180^\circ$;

$\therefore E = (A - A') + (B - B') + (C - C') = 3(A - A')$;

$\therefore \frac{1}{3}E = A - A'$, or $= B - B'$, or $= C - C'$.

Hence if BC (x) be measured, and one third of the computed Spherical Excess be subtracted from each of the observed angles of the triangle A, B, C , the other two sides (y and z) of that triangle can be determined by solving the *plane* triangle $A'B'C'$ whose angles are $A - \frac{1}{3}E, B - \frac{1}{3}E, C - \frac{1}{3}E$, and whose sides are x, y, z .

NOTE. It may here be remarked that the determination of E is a matter of considerable importance when triangles have to be solved whose sides are small compared with the radius of the sphere on which they are described. The observed angles have to be corrected by means of it (70), and this Article shews that it is employed in Legendre's approximate method of solution. How it may be determined from three measured or observed parts of the triangle is given in (68).

74. *To find the angle contained between the chords of two spherical arcs which subtend given angles at the center of the sphere, the angle between the arcs themselves being also given.*

Let AB and AC be the arcs, O the center of the sphere. Let the straight lines AO, AB, AC meet the surface of a sphere which is described with center O and any radius AD , in the points D, E, F respectively. Then the angle EDF is the inclination of the planes BAO and CAO , i. e. the angle contained between the arcs AB and AC , or $\angle A$.

Let $\frac{\text{Arc } AB}{AO} = c, \quad \frac{\text{Arc } AC}{AO} = b.$


Now $\angle CAO = \frac{1}{2}(\pi - \angle AOC) = \frac{1}{2}(\pi - b).$

So $\angle BAO = \frac{1}{2}(\pi - c).$

And from the triangle EDF ,

$\cos EF = \cos DF \cos DE + \sin DF \sin DE \cos EDF$;

Or $\cos EAF = \sin \frac{1}{2}b \sin \frac{1}{2}c + \cos \frac{1}{2}b \cos \frac{1}{2}c \cos A \dots\dots\dots(1).$



75. Two formulæ will now be deduced from (74), which are convenient for determining the angle EAF practically.

Let $\angle EAF = A - \theta$;

Then

$$\cos(A - \theta) - \cos A = \sin \frac{1}{2}b \sin \frac{1}{2}c + (\cos \frac{1}{2}b \cos \frac{1}{2}c - 1) \cos A ;$$

And since, Pl. Trig. (54),

$$\begin{cases} \sin \frac{1}{2}b \sin \frac{1}{2}c = \sin^2 \frac{1}{4}(b+c) - \sin^2 \frac{1}{4}(b-c), \\ \cos \frac{1}{2}b \cos \frac{1}{2}c = \cos^2 \frac{1}{4}(b+c) - \sin^2 \frac{1}{4}(b-c) \\ \qquad \qquad \qquad = 1 - \sin^2 \frac{1}{4}(b+c) - \sin^2 \frac{1}{4}(b-c); \end{cases}$$

$$\begin{aligned} \therefore 2 \sin(A - \frac{1}{2}\theta) \sin \frac{1}{2}\theta &= \{\sin^2 \frac{1}{4}(b+c) - \sin^2 \frac{1}{4}(b-c)\} \\ &\quad - \{\sin^2 \frac{1}{4}(b+c) + \sin^2 \frac{1}{4}(b-c)\} \cos A \\ &= \{1 - \cos A\} \sin^2 \frac{1}{4}(b+c) - \{1 + \cos A\} \sin^2 \frac{1}{4}(b-c). \end{aligned}$$

For an approximation to the value of θ , take $\sin \frac{1}{2}\theta = \frac{1}{2}\theta$, and $\sin(A - \frac{1}{2}\theta) = \sin A$; then, since

$$\frac{1 - \cos A}{\sin A} = \tan \frac{1}{2}A, \text{ and } \frac{1 + \cos A}{\sin A} = \cot \frac{1}{2}A,$$

the above equation becomes

$$\theta = \tan \frac{1}{2}A \sin^2 \frac{1}{4}(b+c) - \cot \frac{1}{2}A \sin^2 \frac{1}{4}(b-c).$$

The number of seconds in this angle is $= \frac{\theta}{\sin 1''}$

$$= \frac{1}{\sin 1''} \cdot \tan \frac{1}{2}A \sin^2 \frac{1}{4}(b+c) - \frac{1}{\sin 1''} \cdot \cot \frac{1}{2}A \sin^2 \frac{1}{4}(b-c) \dots \dots (1);$$

and by determining the two terms of the second member of this equation separately by means of tables of logarithms, the number of seconds in θ is obtained, and the angle $A - \theta$, contained by the straight lines BA and CA , may then be found.

COR. If y, z be the measured lengths of the arcs AC, AB ,

$$\sin \frac{b \pm c}{4} = \sin \frac{y \pm z}{4r} = \frac{y \pm z}{4r}, \text{ nearly.}$$

∴ the number of seconds in θ

$$= \frac{1}{\sin I''} \left\{ \left(\frac{y+z}{4r} \right)^2 \tan \frac{1}{2} A - \left(\frac{y-z}{4r} \right)^2 \cot \frac{1}{2} A \right\}, \text{ nearly.....(2).}$$

If then the radius of the Earth and the sides of the spherical triangle be approximately known, the chordal triangle can be determined; since its sides can be found by the expression, $\text{Chord} = 2r \cdot \sin \left(\frac{1}{2} \frac{\text{arc}}{\text{rad.}} \right)$, and its angles can be determined by one of the last two Articles.

76. Another method is to solve the triangle by the rules laid down for the solution of spherical triangles whose sides are *not* small in comparison with the radius of the sphere. In this case the logarithms of the sines and tangents of the sides, which are very small, must be found by the methods pointed out in Appendix III. to Pl. Trig.

CHAPTER VI.

ON GEODETIC MEASUREMENTS.

77. THE object of Geodetic Measurements is to obtain a correct representation of a part of the Earth's surface which is too large in extent to be considered as lying in one plane.

A horizontal line of considerable length is first measured, which is called a *base*.

Next an object is fixed upon, so situated that it forms with the extremities of the base a triangle which is nearly equilateral (Arts. 71, 72). The angles of this triangle are then measured by a Theodolite, (an instrument described hereafter in Art. 84), and the remaining parts of the triangle computed by some one of these three methods:


1. By the common processes of Spherical Trigonometry, Chap. IV.
2. By the Chordal Triangle (74), (75.)
3. By diminishing each of the observed angles by one third of the Spherical Excess, and then treating the figure as a plane triangle*. (78.)

78. *The use of Geodetic Measurements.*

The form and position of the polygon being thus determined, it may be represented on paper according to any projection of the sphere (See *Hymers' Astronomy, Appendix 1.*), and a map of a

* The first of these methods was preferred by Delambre, the second was employed in the English Survey, and the third in the French Survey.

country obtained. If the object in view be to determine the figure and dimensions of the Earth with great accuracy, the length of an arc mn , which passes through two given points of the polygon, may be found by calculation. If these points, m and n , be situated on the same meridian, and the difference of the zenith distances of the same fixed star when on the meridian be noted by observers at m and n , the Earth's radius of curvature at the middle point of the calculated arc may be approximately found. From arcs thus measured in different latitudes the figure of the Earth has been determined with great exactness. (*Hymers' Astronomy*, 2nd Edit. Chap. II. Arts. 123—142.)*



Next, one of the computed sides is taken as the base of another triangle, whose angles are observed and sides computed; and thus, by a series of triangles, the figure and dimensions of a polygonal area on the Earth's surface are determined. The form of the Earth is, in fact, spheroidal, but it is so nearly spherical, that each triangle may individually be supposed, without appreciable error, to be described on the same sphere. If the triangulation be carried over a very extensive tract of country, it will become necessary to take into consideration the alteration which a change of latitude produces in the Earth's radius. (*Hymers' Astronomy*, Art. 125.)

79. Base of Verification.

A side of the last triangle of the series, after being computed in this manner, is carefully measured. The degree of exactness with which the *measured* coincides with the computed length tests the accuracy of the survey. Any considerable error is easily detected in the course of the calculations, in the following manner. (Fig. Art. 78.) If the value of b , as calculated from the original

* The student will find this part of the subject treated clearly and concisely in the Articles on "Trigonometry" and "The Figure of the Earth," written by the Astronomer Royal for the Encyclopædia Metropolitana. From Section 179 of the former of these Articles, 71 of this treatise has been taken.


For a more particular description of the details of Geodetic operations any of the published accounts of the English Survey may be consulted.

base a through the triangles A, B, C, D, E , be found to agree closely with its value as computed from the same base through a different set of triangles, as A, B', C', E , it may be presumed to have been accurately determined*.

83. *Corrections of Measurements.*—There are several causes productive of error in Geodetic Measurements. (1) The refraction of light is affected by the perpetual variations of the atmosphere in density and temperature, and thus the observed angles cannot always be relied on. (2) It is almost impossible to find a line perfectly straight and perfectly horizontal, to measure for a base. (3) Neither can a portion of the Earth be found which is altogether free from inequalities of surface. To get rid of the errors thus introduced several corrections are used. Some of these, which do not depend on experiment but are capable of mathematical investigation, will now be given.

(1) *If there be a slight rise in the line of the base, to determine the reduction to the horizon.*

Let CB be horizontal, CE the line of the base, ED the *small* rise,—which is obtained by levelling with a spirit-level (83). Describe a circle with center C and radius CE .



Then $BD = CB - CD = CE - CD$
= the "Reduction to the horizon."

$$\text{And } BD = \frac{BE^2}{AB} = \frac{BE^2}{2CE} = \frac{DE^2}{2CE} \dagger, \text{ nearly;}$$

the formula made use of in the English Survey of 1784.

* In the English Trigonometrical Survey of 1784 and succeeding years, the original base on Hounslow Heath was by admeasurement 27404.2 feet; and the base of verification on Salisbury Plain was, as measured, 36574.4 feet, and as computed through three different series of triangles, 36574.3, 36574.6, and 36574.9; any one of which is an approximation sufficiently near for all practical purposes.

† If $BD = x$, $CE = a$, $DE = b$, $x = \frac{BE^2}{2a} = \frac{b^2 + x^2}{2a}$; and neglecting powers of x above the first, $x = \frac{b^2}{2a}$, nearly.

(2) Let the measured base, instead of being one straight line, consist of two straight lines, a and b , enclosing an angle $\pi - \theta$, where θ is very small. Required the correction to find c .

$$c^2 = a^2 + b^2 + 2ab \cos \theta$$

$$= a^2 + b^2 + 2ab \left(1 - \frac{1}{2} \theta^2\right), \text{ nearly,}$$

$$= (a+b)^2 \left\{1 - \frac{ab}{(a+b)^2} \cdot \theta^2\right\};$$

$$\therefore c = (a+b) \left\{1 - \frac{ab}{2(a+b)} \cdot \theta^2\right\}, \text{ nearly;}$$

$$= (a+b) - \frac{ab}{2(a+b)} \cdot \theta^2.$$

$$\therefore \text{Correction} = (a+b) - c = \frac{ab}{2(a+b)} \cdot \theta^2.$$

Or if θ be an angle containing n'' , where n is very small,

$$\text{the Correction, in seconds,} = \frac{1}{\sin 1''} \cdot \frac{ab (n \sin 1'')^2}{2(a+b)} = \frac{ab n^2 \sin 1''}{2(a+b)}.$$

[In practice it is seldom necessary to apply this correction.]

(3) From observations made at a point D which is at a small known distance from a signal C , required to find the angle which A and B subtend at C .

Let the angles BDA and ADC be observed at D . The distances CB and CA are known approximately from the base AB and the observed angles BAC, ABC .

Then $\angle BDA + \angle DBE$

$$= \text{exterior angle } BEA = \angle BCA + \angle CAD,$$

$$\therefore \text{The reduction, viz. } \angle BCA - \angle BDA,$$



$$= \angle DBE - \angle CAD$$

$$= \sin \angle DBE - \sin \angle CAD, \text{ nearly,}$$

$$= \frac{CD}{CB} \cdot \sin BDC - \frac{CD}{CA} \cdot \sin ADC$$

$$= CD \left\{ \frac{\sin BDC}{CB} - \frac{\sin ADC}{CA} \right\};$$

$$\text{or} = \frac{CD}{\sin 1''} \left\{ \frac{\sin BDC}{CB} - \frac{\sin ADC}{CA} \right\}, \text{ when expressed in seconds.}$$




(4) *The angles of elevation or depression, being small quantities, of two objects having been observed, and also the angle the objects subtend at a certain point, to find the horizontal angle they subtend. (Encycl. Metrop.)*

Let Z be the Observer's zenith, $CDRO$ his horizon, A and B the two objects; $\angle BOA = \theta$; $\angle DOC$, its reduction to the horizon, $= \theta + x$; $CA = h$, $DB = h'$.

From the $\triangle AZB$,

$$\begin{aligned} \cos CZD &= \frac{\cos \theta - \cos ZA \cos ZB}{\sin ZA \sin ZB} \\ &= \frac{\cos \theta - \sin h \sin h'}{\cos h \cos h'}; \end{aligned}$$



Making $\sin h = h$, $\cos h = 1 - \frac{1}{2}h^2$, &c., and neglecting powers of the angles above the second, this becomes

$$\begin{aligned} \cos CZD &= \frac{\cos \theta - hh'}{1 - \frac{1}{2}(h^2 + h'^2)} \\ &= (\cos \theta - hh') \left\{ 1 + \frac{1}{2}(h^2 + h'^2) \right\}, \text{ nearly,} \\ &= \cos \theta - hh' + \frac{1}{2}(h^2 + h'^2) \cos \theta. \end{aligned}$$

$$\begin{aligned} \text{But } \cos CZD &= \cos(\theta + x) = \cos \theta \cos x - \sin \theta \sin x \\ &= \cos \theta - x \sin \theta, \text{ nearly.} \end{aligned}$$

And from these two equations,

$$x = \frac{hh'}{\sin \theta} - \frac{1}{2}(h^2 + h'^2) \frac{\cos \theta}{\sin \theta}.$$

Now if $p = h + h'$, and $q = h - h'$; then $hh' = \frac{1}{4}(p^2 - q^2)$, $h^2 + h'^2 = \frac{1}{2}(p^2 + q^2)$;

$$\begin{aligned} \therefore x &= \frac{1}{4} \left\{ \frac{p^2 - q^2}{\sin \theta} - (p^2 + q^2) \cdot \frac{\cos \theta}{\sin \theta} \right\} \\ &= \frac{1}{4} \left\{ p^2 \cdot \frac{1 - \cos \theta}{\sin \theta} - q^2 \cdot \frac{1 + \cos \theta}{\sin \theta} \right\} \\ &= \frac{1}{4} \left\{ p^2 \tan \frac{1}{2} \theta - q^2 \cot \frac{1}{2} \theta \right\}. \end{aligned}$$

In seconds, this correction

$$= \frac{1}{4} \frac{1}{\sin 1''} \cdot \{(p' \sin 1'')^2 \tan \frac{1}{2} \theta - (q' \sin 1'')^2 \cot \frac{1}{2} \theta\}, \text{ where}$$


p', q' are the seconds in the angles p and q ,

$$= \frac{1}{4} \sin 1'' \{p'^2 \tan \frac{1}{2} \theta - q'^2 \cot \frac{1}{2} \theta\}.$$

NOTE. This reduction is not required when a Theodolite is used.

INSTRUMENTS USED IN SURVEYING.

81. DEF. THE VERNIER is a contrivance for subdividing equal graduations that have been made on a straight line or a circle.



AB is a portion of a straight line, or a circle, which is divided by straight lines at right angles to it into any number of equal parts. CD , the Vernier, is another scale, which slides along AB when AB is a straight line, and revolves round the center of AB when it is a circular arc.

If it be required to subdivide each of the divisions of AB into n equal parts, take $QM = n - 1$ of these parts, and divide QM into n equal parts by straight lines at right angles to it; if a be the length of a division of AB , the magnitude of each of these parts will be $\frac{n-1}{n} \cdot a$.

Let R , the r^{th} division from Q , coincide with a division on AB ; then $PQ = PR - QR = ra - r \cdot \frac{n-1}{n} \cdot a = r \cdot \frac{a}{n}$, and the length of PQ is known.

Ex. If each of the original divisions were an inch, and nine inches were divided into 10 parts on the Vernier, then, supposing that the extremity of a line AQ which it was wanted to measure came to Q , and that the inches marked at P were p , if the third division of the Vernier coincided with a division on the scale, the length required would be $(p + \frac{3}{10})$ inches.

82. THE SPIRIT LEVEL is a glass tube $DABE$ of circular bore, which is ground into the form of a circular arc of very large radius—sometimes 800 feet. It is then nearly filled with some fluid, and the ends are closed.

If the instrument be placed in a vertical plane, and the extremities D and E rest on a horizontal surface, the bubble (AB) of air left in the tube will be at the highest part of it; and if E be gradually raised the bubble will continually keep moving towards E .*



If there be a plane of an instrument (such as a Theodolite) which it is necessary to bring into a horizontal position, it is provided with two levels, as nearly equal to one another in every respect as possible, which are placed at right angles to each other and permanently attached to the plane. The instrument-maker marks the positions of the bubbles when the plane is horizontal, and therefore if the bubbles occupy these positions on any occasion, the plane to which the levels are attached must then be horizontal.

If the plane be inclined at any angle to the vertical and the positions of the bubbles be noted, then if at a second observation they occupy the same positions, the plane will have the same inclination to the vertical which it had before.

83. *To level between two points.*

For the purpose of finding the altitude of one point above another point, a spirit-level is attached to a telescope, and so


* The grinding the bore of a Spirit-level is done with a plug of metal covered with emery. The elasticity of the glass, assisted probably by that of the metal of the plug, enables the workman, by means of pressure on the outside, to wear away any particular portion of the interior surface he chooses. If, after the grinding is finished, the bubble be found to move through equal lengths of the tube for equal increments of inclination, and to be always of the same length, the bore of the tube must be uniform, and the form of the tube a truly circular arc.

If l be the length through which the bubble moves in consequence of an increase of n'' in the inclination (n being a small quantity), the radius of the circular arc = $\frac{l}{n \sin 1''} = \frac{l}{n} \times 206265$.

Ether is the best fluid with which spirit-levels can be filled; because in ether the bubble is found to come into a state of rest in the shortest time after a sudden displacement.

adjusted that the optical axis of the telescope is horizontal when the bubble is at the middle of the level.


Let there be two staffs set up vertically, AB , CD ; and when the instrument is at B and its axis is horizontal, let the point C be seen on the cross wires; and when the instrument is placed at D with its axis horizontal let E be seen on the cross wires; then, by measuring AB and CD , EF , the altitude of E above the horizontal line AF passing through A , is found. The operation can be repeated as often as it may be necessary.



84. THE THEODOLITE. This is an instrument for measuring the angles of elevation of objects above a horizontal plane; and also the horizontal angle which two objects subtend at the observer's eye.

The accompanying figure is an elevation (or projection on a vertical plane by lines perpendicular to the plane) of a Theodolite.

The two circles I and K fit close to each other, the lower one having attached to it three horizontal bars, making angles of 120° with each other, on which the instrument rests. The feet of the instrument are three screws, M , M , M , working in these bars; and by moving the screws in one direction or the other, the planes of the circles I and K , which are parallel to each other, can be brought into a horizontal position.



The upper circle I is graduated, and revolves with the utmost possible nicety upon the lower circle by means of an axis attached to the upper circle, and working within the collar L which forms a part of the frame of the instrument. The circle K has two Verniers engraved upon it, and in the best instruments the divisions are read off by microscopes attached to the Verniers.


To the revolving circle I , which is called "the limb" of the Theodolite, two levels at right angles to each other are attached.

To this circle also two stands are fixed diametrically opposite to each other (one of which, DOE , is here represented) supporting an axis parallel to the revolving circle I . To this axis a circle ABC is permanently attached at right angles to it; and a telescope (GH) is fastened to the circle, with its line of collimation perpendicular to the axis of the circle. The circle and attached telescope revolve along with the axis in such a manner that the whole can turn completely round without touching "the limb" of the instrument. The rim BB of the circle is graduated; and there are two Verniers, A and C , unconnected with the axis, but carried by a support which is seen to enter the circle I (but not the circle K) at F .

85. *To explain how the Theodolite is used.*

The two circles I and K are first rendered horizontal by lengthening or shortening the screw-feet. The limb is then turned round upon the lower circle until the plane of the vertical circle passes through the object whose altitude it is required to find, and the axis carrying the vertical circle is made to revolve until the observer, on looking through the telescope, perceives the object on the cross wires with which the telescope is furnished. The graduations at the points of the rim of the vertical circle which are opposite to the beginning of the scales engraved on the Vernier plates are then read off.

Next, without touching the axis to which the telescope is attached, the limb is made to revolve through 180° . The plane of the vertical circle again passes through the object, but the direction of the telescope ($A'B'$) is now as much depressed below the horizon as it was before (AB) elevated above it. Let the vertical circle and its axis be turned in the direction $A'AB'$ until the object is again seen on the cross wires of the telescope; the telescope $A'B'$ is therefore now brought into the same direction (AB) that it had at the first observation. The graduations are again read off at the Verniers, and the number of degrees marked on the arcs AA' , BB' of the instrument, (which are the arcs that have passed between the two Verniers), is four times the zenith distance of the object.



[For let O be the real center of the graduated circle; then twice the zenith distance is BPB' , or APA' ; join BA' , AB' ;

$$\begin{aligned} \therefore 4 \times \text{zenith distance} &= BPB' + APA' = (PAB' + PB'A) + (PBA' + PA'B) \\ &= BAB' + A'B'A + ABA' + B'A'B \\ &= 2BAB' + 2ABA' = BOB' + AOA'; \text{ Euclid, III. 20} \end{aligned}$$

= sum of the differences of readings at the Verniers at the two observations.

It appears, therefore, that any errors arising from the axis of rotation of the circle not coinciding exactly with the center of its graduation are wholly avoided by using the instrument in this manner. (*Hymers' Astronomy*, Art. 117.)]

To observe the *horizontal* angle between two objects, the limb of the Theodolite is made horizontal, and the angle is noted (by four readings off, as in the last case) through which the limb revolves to bring the two objects successively on the cross wires of the telescope.

86. THE REPEATING CIRCLE was the instrument used in the French Surveys for observing the angles. The observations, however, which are made by it, are of questionable value for this reason, that although the errors of imperfect graduation (which are but slight, if the instrument be a good one) may *possibly* be destroyed, as they *probably* are, by repeating the observations and taking the mean of them, yet, for anything that can be known to the contrary, the errors arising from the inability of the observer to distinguish the position of a point with perfect exactness, may have been accumulating all the while.

Also, from the construction of the Repeating Circle, it is scarcely possible to avoid errors arising from the *instability* of the instrument.

CHAPTER VII.

ON (I.) THE SMALL CORRESPONDING VARIATIONS OF THE PARTS OF
A SPHERICAL TRIANGLE; AND (II.) THE CONNEXION EXISTING
BETWEEN SOME FORMULÆ IN SPHERICAL TRIGONOMETRY AND
ANALOGOUS FORMULÆ IN PLANE TRIGONOMETRY.

I. THE process of Differentiation can be applied to determine the errors introduced in determining the other parts of a Spherical Triangle from three given parts, when one of the given parts is affected by a small known error.

87. *If C and c remain constant, the corresponding small variations δa and δb of the sides a and b are connected by the equation,*

$$\delta b \cdot \cos A + \delta a \cdot \cos B = 0.$$

Considering C and c constant, and differentiating with respect to a, the formula $\text{Cos } C \sin a \sin b = \cos c - \cos a \cos b$,

$$\text{Cos } C \{ \cos a \sin b + d_a b \cdot \cos b \sin a \} = \sin a \cos b + d_a b \cdot \sin b \cos a ;$$

$$\therefore \text{Cos } C \{ \delta a \cdot \cos a \sin b + \delta b \cdot \cos b \sin a \}$$

= $\delta a \cdot \sin a \cos b + \delta b \cdot \sin b \cos a$, nearly ; δa and δb being small corresponding increments of a and b ;

$$\therefore 0 = \delta a \cdot \{ \sin a \cos b - \text{Cos } C \sin b \cos a \}$$

$$+ \delta b \cdot \{ \sin b \cos a - \text{Cos } C \sin a \cos b \}$$


$$= \delta a \cdot \cos B \sin c + \delta b \cdot \cos A \sin c ; \quad \text{by (iv.)}$$

$$\therefore 0 = \delta a \cdot \cos B + \delta b \cdot \cos A.$$

This method is always applicable. A small spherical triangle, however, is frequently treated as a plane triangle after the following manner ; particularly in establishing formulæ for calculating the corrections used in Astronomical investigations.

88. If c , the side opposite the right angle in a right-angled spherical triangle, receive a small known increment, to determine the corresponding increments of the sides enclosing the right angle.

Let $AB = c$, $BD = \delta c$, $\angle BAC = \omega$. Draw DE perpendicular to AE , and let BF be an arc of a small circle which has the same pole as ACE ; BF is therefore parallel to AE , and perpendicular to DE .



Now, considering BDF as a small plane triangle,

$$\frac{BF}{\delta c} = \cos DBF = \sin ABC,$$

for $\angle CBF = 90^\circ$, and $\therefore \angle ABC = 90^\circ - \angle DBF$;

$$\therefore BF = \delta c \cdot \frac{\cos \omega}{\cos BC}, \quad \text{by (37).}$$

$$\therefore \delta(AC), \quad \text{or } CE, = \frac{BF}{\cos BC}, \quad (15), = \delta c \cdot \frac{\cos \omega}{\cos^2 BC} \dots\dots (1).$$

$$\begin{aligned} \text{Again, } \delta(BC) &= FD = \delta c \cdot \sin DBF \\ &= \delta c \cdot \cos ABC = \delta c \cdot \sin \omega \cos AC \dots\dots\dots (2). \end{aligned}$$

[Cor. If ω be constant,

- { $\delta(AC)$ has its greatest value, when BC is the greatest.
- { $\delta(AC)$ least , when $BC = 0^\circ$.
- { $\delta(BC)$ has its greatest value, when $AC = 0^\circ$, or 180° .
- { $\delta(BC)$ least , when $AC = 90^\circ$.

This result shews, that if the Sun's daily motion (δc) in longitude be nearly constant (as it is in fact), his greatest daily change in Right Ascension, $\delta(AC)$, is at the Solstices, and his greatest daily change in Declination, $\delta(BC)$, is at the Equinoxes; and his least daily changes in Right Ascension and in Declination, are at the Equinoxes and Solstices respectively.

Also; since when BC and AC are small angles the variation in magnitude of their cosines is then the least, it appears from (1) and (2) that the daily change, both in Right Ascension and in Declination, is more nearly proportional to the change in longitude, (that is, to the time), when the Sun is near the Equinoxes, than when he is in any other part of his apparent orbit. (*Hymers' Astronomy*, Art. 171.)]

II. 89. From the formula, $\frac{\text{Sin } A}{\text{Sin } B} = \frac{\sin a}{\sin b}$, which is true for a Spherical Triangle, to deduce the analogous formula in the case of a Plane Triangle.

Let a' , b' be the lengths of the arcs subtending the angles A and B ; r the radius of the sphere.

$$\text{Then } \frac{\text{Sin } A}{\text{Sin } B} = \frac{\sin \frac{a'}{r}}{\sin \frac{b'}{r}} = \frac{\frac{a'}{r} - \frac{1}{2.3} \cdot \frac{a'^3}{r^3} + \dots}{\frac{b'}{r} - \frac{1}{2.3} \cdot \frac{b'^3}{r^3} + \dots} = \frac{a' \cdot \left(1 - \frac{1}{2.3} \cdot \frac{a'^2}{r^2} + \dots\right)}{b' \cdot \left(1 - \frac{1}{2.3} \cdot \frac{b'^2}{r^2} + \dots\right)}$$

And if a' and b' be indefinitely small compared with the radius of the sphere, the formula becomes

$$\frac{\text{Sin } A}{\text{Sin } B} = \frac{a'}{b'}; \text{ a property of Plane triangles.}$$

90. And, in like manner as in the last Article, from the formula

$$\text{Cos } A = \frac{\cos a - \cos b \cos c}{\sin b \sin c}, \text{ and } \text{Tan } \frac{1}{2}(A-B) = \frac{\sin \frac{1}{2}(a-b)}{\sin \frac{1}{2}(a+b)} \cdot \cot \frac{1}{2}C,$$

there may be deduced these analogous formulæ of Plane Trigonometry.


$$\text{Tan } \frac{1}{2}(A-B) = \frac{a' - b'}{a' + b'} \cdot \cot \frac{1}{2}C, \quad \text{and } \text{Cos } A = \frac{b'^2 + c'^2 - a'^2}{2b'c'}$$

91. PROB. If two arcs of great circles intersect each other in a small circle, the product of the tangents of the semi-segments of the one is equal to that of the tangents of the semi-segments of the other.

Let AB , CD be the arcs of great circles intersecting in a point E within the small circle whose pole is P ; $FEPG$ an arc of the great circle through E and P ; PQ perpendicular to AB .

Then AB is bisected in Q , and APQ , BPQ are triangles equal in every respect.

$$\text{Also } PA = PB = PF = PG.$$



By (37),

$$\cos PE = \cos EQ \cos PQ, \quad \text{and} \quad \cos PA = \cos AQ \cos PQ;$$

$$\therefore \frac{\cos EQ - \cos AQ}{\cos EQ + \cos AQ} = \frac{\cos PE - \cos PA}{\cos PE + \cos PA};$$

$$\therefore \tan \frac{1}{2}(AQ + EQ) \tan \frac{1}{2}(AQ - EQ) = \tan \frac{1}{2}(PA + PE) \tan \frac{1}{2}(PA - PE),$$

$$\begin{aligned} \text{or} \quad \tan \frac{1}{2}BF \tan \frac{1}{2}EA &= \tan \frac{1}{2}GE \tan \frac{1}{2}EF \\ &= \tan \frac{1}{2}CE \tan \frac{1}{2}ED, \text{ in like manner.} \end{aligned}$$

COR. If a', b' represent the lengths of the segments of the arc AB , and c', d' represent the lengths of the segments of the arc CD ; and these quantities be indefinitely small compared with the radius of the sphere, this formula becomes, in the case where the radius of the small circle vanishes with respect to the radius of the sphere, $a'b' = c'd'$; which agrees with Euclid, III. 35.

CHAPTER VIII.

ON THE REGULAR SOLIDS.

92. **DEFS.** (1) A *Polyhedron* is a solid bounded by plane rectilinear figures.

If the bounding surface be composed of any similar and equal regular rectilinear figures, the polyhedron is called a *Regular Polyhedron*.

(2) A *Tetrahedron* is bounded by four equal and equilateral triangles.

(3) A *Hexahedron*, or *Cube*, is bounded by six equal squares.

(4) An *Octahedron* is bounded by eight equal and equilateral triangles.

(5) A *Dodecahedron* is bounded by twelve equal and equilateral pentagons.

(6) An *Icosahedron* is bounded by twenty equal and equilateral triangles.

It will be proved hereafter that no more Regular Polyhedrons exist than these five.

93. In any regular *Polyhedron*, if

F = number of Faces, S = number of Solid Angles,

E = number of Edges, m = number of Sides in each Face;

then $2E = mF$, and $S + F = E + 2$.

(1) Since every edge is made by two sides, the whole number of sides in the polyhedron is $2E$, and this

$$= \text{Number of faces} \times \text{number of sides in a face,}$$

$$\therefore 2E = mF.$$

(2) Take any point within the polyhedron as the center of a sphere whose radius is r , and join it with each of the angular points of the polyhedron. Let the points in which these lines meet the surface of the sphere be joined by arcs of great circles; the surface of the sphere will then be divided into as many polygons as the polyhedron has faces, and the Area of one of these polygons

$$= \frac{\pi r^2}{180} \cdot \left\{ \begin{array}{l} \text{number of degrees in the angles of the polygon} \\ - (\text{number of sides of polygon} - 2) \cdot 180 \end{array} \right\}$$

(62, Cor. 2.)

\therefore Area of all these polygons

$$= \frac{\pi r^2}{180} \cdot \left\{ \begin{array}{l} \text{number of degrees in the angles of all the polygons} \\ \text{on the sphere} \\ - (\text{number of all the sides} - 2F) \cdot 180 \end{array} \right\}$$

$$= \frac{\pi r^2}{180} \cdot \{S \cdot 360 - (E - F) \cdot 360\} = 2\pi r^2 \cdot (S - E + F).$$

But Area of all the polygonal areas = area of the sphere = $4\pi r^2$;

$$\therefore S - E + F = 2,$$

$$\text{and } S + F = E + 2.$$

[These results are evidently true whether the Polyhedron be Regular or Irregular.]

94. *The sum of all the Plane Angles which form the Solid Angles of a Regular Polyhedron = $(S - 2) \cdot 360^\circ$.*

For the Sum of the Plane Angles = sum of all the Interior Angles of each face.

$$= F \cdot (m - 2) \cdot 180^\circ \quad \text{Eucl. I. 32, Cor. 1.}$$

$$= 2(E - F) \cdot 180^\circ$$

$$= (S - 2) \cdot 360^\circ \quad \text{by (93).}$$

CHAPTER VIII.

ON THE REGULAR SOLIDS.

92. DEFS. (1) A *Polyhedron* is a solid bounded by plane rectilinear figures.

If the bounding surface be composed of any similar and equal regular rectilinear figures, the polyhedron is called a *Regular Polyhedron*.

(2) A *Tetrahedron* is bounded by four equal and equilateral triangles.

(3) A *Hexahedron*, or *Cube*, is bounded by six equal squares.

(4) An *Octahedron* is bounded by eight equal and equilateral triangles.

(5) A *Dodecahedron* is bounded by twelve equal and equilateral pentagons.

(6) An *Icosahedron* is bounded by twenty equal and equilateral triangles.

It will be proved hereafter that no more Regular Polyhedrons exist than these five.

93. In any regular *Polyhedron*, if

F = number of Faces, S = number of Solid Angles,

E = number of Edges, m = number of Sides in each Face;

then $2E = mF$, and $S + F = E + 2$.

(1) Since every edge is made by two sides, the whole number of sides in the polyhedron is $2E$, and this

= number of faces \times number of sides in a face,

$$\therefore 2E = mF.$$

(2) Take any point within the polyhedron as the center of a sphere whose radius is r , and join it with each of the angular points of the polyhedron. Let the points in which these lines meet the surface of the sphere be joined by arcs of great circles; the surface of the sphere will then be divided into as many polygons as the polyhedron has faces, and the Area of one of these polygons

$$= \frac{\pi r^2}{180} \left\{ \begin{array}{l} \text{number of degrees in the angles of the polygon} \\ - (\text{number of sides of polygon} - 2) \cdot 180 \end{array} \right\}.$$

(62, Cor. 2.)

\therefore Area of all these polygons

$$= \frac{\pi r^2}{180} \left\{ \begin{array}{l} \text{number of degrees in the angles of all the polygons} \\ \text{on the sphere} \\ - (\text{number of all the sides} - 2F) \cdot 180 \end{array} \right\}$$

$$= \frac{\pi r^2}{180} \cdot \{S \cdot 360 - (E - F) \cdot 360\} = 2\pi r^2 \cdot (S - E + F).$$

But Area of all the polygonal areas = area of the sphere = $4\pi r^2$;

$$\therefore S - E + F = 2,$$

$$\text{and } S + F = E + 2.$$

[These results are evidently true whether the Polyhedron be Regular or Irregular.]

94. *The sum of all the Plane Angles which form the Solid Angles of a Regular Polyhedron = $(S - 2) \cdot 360^\circ$.*

For the Sum of the Plane Angles = sum of all the Interior Angles of each face.

$$\begin{aligned} &= F \cdot (m - 2) \cdot 180^\circ \quad \text{Eucl. I. 32, Cor. 1.} \\ &= 2(E - F) \cdot 180^\circ \\ &= (S - 2) \cdot 360^\circ \quad \left. \vphantom{\begin{aligned} &= F \cdot (m - 2) \cdot 180^\circ \\ &= 2(E - F) \cdot 180^\circ \end{aligned}} \right\} \text{ by (93).} \end{aligned}$$

95. *To prove that in a Regular Polyhedron*

$$S = \frac{4m}{2(m+n) - mn}, \quad E = \frac{2mn}{2(m+n) - mn}, \quad F = \frac{4n}{2(m+n) - mn},$$

where m and n are respectively the number of Sides in every Face and the number of Plane Angles in every Solid Angle.

Since every face has m plane angles,

\therefore number of the Plane Angles which form all the Solid Angles $= mF$, and $\therefore = Sn$.

Hence, (93,) $Sn = mF = 2E$; and since $S + F = E + 2$,

$$\therefore 2 = S + F - E = S \left(1 + \frac{n}{m} - \frac{n}{2} \right);$$

$$\therefore S = \frac{4m}{2(m+n) - mn}; \quad E = \frac{2mn}{2(m+n) - mn}; \quad F = \frac{4n}{2(m+n) - mn}.$$

96. *There can be but five Regular Polyhedrons.*

In any regular polyhedron, m , n , S , E , F must each be a positive integer.

In order that the values of S , E , F obtained in the last Article may be positive, $2(m+n)$ must be greater than mn ; and that each of them may be integral, $4m$, $2mn$, and $4n$ must be severally divisible by $2(m+n) - mn$.

Now if $2(m+n)$ be greater than mn ,

$$\frac{1}{m} + \frac{1}{n} > \frac{1}{2}, \quad \text{or} \quad \frac{1}{m} > \frac{1}{2} - \frac{1}{n};$$

but n cannot be less than 3,

$$\therefore \frac{1}{m} \text{ cannot be so small as } \frac{1}{2} - \frac{1}{3}, \quad \text{or } \frac{1}{6};$$

Therefore, since m must be an integer, and cannot be less than 3, it can only be 3, 4, or 5.

Similarly, since $\frac{1}{n} > \frac{1}{2} - \frac{1}{m}$, and m cannot be less than 3;

\therefore the values of n can only be 3, 4, and 5.

It will be found, on trial, that the only values of m and n which satisfy all the required conditions are the following. Each regular solid takes its name from the number of its plane faces.

m .	n .	S .	E .	F .	Name of the Regular Solid.
3	3	4	6	4	Tetrahedron. (Regular Pyramid.)
4	3	8	12	6	Hexahedron. (Cube.)
3	4	6	12	8	Octahedron.
5	3	20	30	12	Dodecahedron.
3	5	12	30	20	Icosahedron.

COR. If $2(m+n) = mn$, S , E , and F become infinite quantities, and the solid itself becomes a sphere.

97. If I be the inclination of two contiguous faces of a Regular Polyhedron,


$$\text{then, } \sin \frac{1}{2} I = \frac{\cos \frac{180^\circ}{n}}{\sin \frac{180^\circ}{m}}, \quad \text{or } = \frac{\cos \frac{S}{E} \cdot 90^\circ}{\sin \frac{F}{E} \cdot 90^\circ}.$$

Let C and E be the centers of the circles inscribed in two adjacent faces whose common edge is AB ; bisect AB in D , and join A , B , and D with the points C and E ; CD and ED are manifestly perpendicular to AB , and $\therefore \angle CDE = I$.

In the plane CDE draw CO and EO at right angles to CD and ED respectively; let these lines meet in O ; join OA , OB , OD .

About O as center describe a spherical surface which is cut by the planes AOD , DOC , COA in ad , dc , ca .

Now since AB is perpendicular to CD and to ED , it is perpendicular to the plane CDE , and therefore the plane AOB , in which AB lies, is perpendicular to the plane COE ; $\therefore \angle adc$ is a right angle;



And, $\angle cad$

$= \frac{1}{2} \cdot 360^\circ \div$ number of edges which meet in a solid angle

$= 180^\circ \div$ number of plane angles which meet in a solid angle $= \frac{180^\circ}{n}$;

Also $\angle acd$

$= \left\{ \begin{array}{l} \frac{1}{2} \angle \text{ which each side of a plane face subtends} \\ \text{at the center of the circle inscribed in the face} \end{array} \right\} = \frac{1}{2} \cdot \frac{360^\circ}{m} = \frac{180^\circ}{m}$.

Now by (37), $\text{Cos } \angle cad = \text{cos } dc \sin \angle acd$,

$$\therefore \text{Cos } \frac{180^\circ}{n} = \text{cos } dc \sin \frac{180^\circ}{m}$$

And $\text{Cos } dc = \text{csc } DOC = \text{cos } \frac{1}{2} COE = \text{cos } \frac{1}{2} (180^\circ - CDE) = \sin \frac{1}{2} I$;

$$\therefore \text{Sin } \frac{1}{2} I = \frac{\text{cos } \frac{180^\circ}{n}}{\sin \frac{180^\circ}{m}}$$

Again, Since $n = \frac{2E}{S} \dots (95)$, and $m = \frac{2E}{F'} \dots (93)$,

$$\therefore \text{Sin } \frac{1}{2} I = \frac{\text{cos} \left(\frac{S}{E} \cdot 90^\circ \right)}{\text{sin} \left(\frac{F'}{E} \cdot 90^\circ \right)}$$

98. *To find the Radius of the Sphere which may be inscribed in a Regular Polyhedron.* (Fig. Art. 97.)

$OC = OE$, ($= r$), is this radius. Let $AB = 2a$.

Then $CD = AD \cdot \cot ACD = a \cdot \cot ACD = a \cdot \cot \frac{180^\circ}{m}$;

And $r = CD \cdot \tan CDO = CD \cdot \tan \frac{1}{2} I = a \cdot \tan \frac{1}{2} I \cot \frac{180^\circ}{m}$.

99. *To find the Radius of the Sphere described about a Regular Polyhedron.* (Fig. Art. 97.)

$OA = OB$, ($= R$), is this radius.

$$\text{And } r = R \cdot \cos ac = R \cdot \cot acd \cot cad = R \cdot \csc \frac{180^\circ}{m} \cot \frac{180^\circ}{n}$$

$$\therefore R = r \cdot \tan \frac{180^\circ}{m} \tan \frac{180^\circ}{n} = a \cdot \tan \frac{1}{2} I \tan \frac{180^\circ}{n}, \text{ by (98).}$$

100. *If a Hexahedron and an Octahedron be described about a given sphere, the sphere described about those Polyhedrons will be the same; and conversely.*

Let R and r , R' and r' , be the radii of the inscribed and circumscribed spheres for a hexahedron and an octahedron respectively.

$$\text{Then } \frac{R}{r} = \tan \frac{180^\circ}{3} \cdot \tan \frac{180^\circ}{4} = \frac{R'}{r'}$$

Wherefore, if R' be equal to R , r' is equal to r ; or if r' be equal to r , R' is equal to R . That is, if a hexahedron and an octahedron be described about the same sphere, the spheres circumscribing them will also be the same; and conversely.

In like manner it may be proved that if a dodecahedron and an icosahedron be described in a given sphere, they will have the same circumscribing sphere; and conversely.

101. *To find the Volume of a regular Polyhedron.*

From O in fig. Art. 97, draw OA , OB , &c. to all the angles of the polyhedron. The solid will thus be divided into F pyramids, whose common altitude is r , and common base, being the area of a face, = $m \cdot \frac{AB}{2} \cdot CD = m \cdot a \cdot CD = m \cdot a \cdot r \cdot \cot \frac{1}{2} I$;(98).

$$\therefore \text{Whole Volume} = \frac{1}{3} \cdot F \cdot m \cdot a \cdot r^2 \cdot \cot \frac{1}{2} I$$

$$= \frac{1}{3} \cdot F \cdot m \cdot \tan \frac{1}{2} I \cdot \cot^2 \frac{180^\circ}{m} \cdot a^3, \text{ by (98).}$$

COR. Therefore in similar Polyhedrons, Volume $\propto a^3$.


102. In a Parallelopiped, given the three Edges which meet, and the Angles between them, to find the Altitude, Surface, and Volume of the solid.

Let AO, BO, CO be the three edges meeting in the point O ; and let

$$\angle BOC = A, \angle COA = B, \angle AOB = C;$$

$$OA = a, \quad OB = b, \quad OC = c;$$

$$S = \frac{1}{2} (A + B + C).$$



CD perpendicular to the plane AOB . Join OD, BD, DA .

About O describe a spherical surface, and let it be cut by the planes AOB, BOC, COA, DOC in ab, bc, ca, dc .

Then, the angles at d being right angles,

$$\text{Sin } dc = \text{sin } ac \text{ sin } cad, \quad \text{or} \quad \text{Sin } DOC = \text{sin } B \text{ sin } cad;$$

And, by (x.),

$$\text{Sin } cad = \frac{2}{\text{sin } B \text{ sin } C} \sqrt{\{\text{sin } S \text{ sin } (S - A) \text{ sin } (S - B) \text{ sin } (S - C)\}};$$

$$\therefore DC = c \cdot \text{sin } DOC = \frac{2c}{\text{sin } C} \sqrt{\{\text{sin } S \text{ sin } (S - A) \text{ sin } (S - B) \text{ sin } (S - C)\}}.$$

The Surface = $2 \{bc \text{ sin } A + ac \text{ sin } B + ab \text{ sin } C\}$.

The Volume = $\left\{ \begin{array}{l} \text{Volume of rectangular parallelopiped on the} \\ \text{same base, and of the same altitude.} \end{array} \right.$

= area of base \times altitude. Euclid, XI. 31.

$$= 2abc \sqrt{\{\text{sin } S \text{ sin } (S - A) \text{ sin } (S - B) \text{ sin } (S - C)\}}.$$

103. The same things being given, to determine the Diagonal which passes through the Solid angle O of the Parallelopiped.

Let D be the diagonal required; and now suppose OdD in the fig. Art. 102, to be in the direction of the diagonal of the face AOB . Let this diagonal = d .

Then $d^2 = a^2 + b^2 + 2ab \cos C,$

and $D^2 = c^2 + d^2 - 2cd \cos (180^\circ - DOC)$

$= c^2 + d^2 + 2cd \cos DOC,$

$\therefore = a^2 + b^2 + c^2 + 2cd \cos DOC + 2ab \cos C.$

Now $\cos DOC = \cos cd = \cos ac \cos ad + \sin ac \sin ad \cos bac$

$= \cos B \cos ad + \sin B \sin ad \cdot \frac{\cos A - \cos B \cos C}{\sin B \sin C}$

$= \frac{1}{\sin C} \cdot \{ \cos B \sin C \cos ad - \cos B \cos C \sin ad + \cos A \sin ad \}$

$= \frac{1}{\sin C} \cdot \{ \cos B \sin (C - ad) + \cos A \sin ad \}$

$= \frac{1}{\sin C} \cdot \{ \cos B \sin BOD + \cos A \sin AOD \}.$

But $\frac{a}{d} = \frac{\sin BOD}{\sin C},$ and $\frac{b}{d} = \frac{\sin AOD}{\sin C};$

$\therefore \cos DOC = \frac{a \cos B + b \cos A}{d}.$

And $D^2 = a^2 + b^2 + c^2 + 2bc \cos A + 2ac \cos B + 2ab \cos C.$

$\therefore D = \sqrt{a^2 + b^2 + c^2 + 2bc \cos A + 2ac \cos B + 2ab \cos C}.$

COR. 1. At the solid angle $C,$

The Cosine of the angle between c and $b = -\cos COB = -\cos A,$

..... c and $a = -\cos B,$

..... a and $b = \cos C;$

Therefore the square of the diagonal through C

$= a^2 + b^2 + c^2 - 2bc \cos A - 2ac \cos B + 2ab \cos C.$

Similarly, the squares of the diagonals through the other solid angles, B and A , are,

$$a^2 + b^2 + c^2 - 2bc \cos A + 2ac \cos B - 2ab \cos C.$$

$$a^2 + b^2 + c^2 + 2bc \cos A - 2ac \cos B - 2ab \cos C.$$

CoR. 2. Hence it appears that the sum of the squares of the four diagonals

$$= 4(a^2 + b^2 + c^2).$$

EXAMPLES FOR PRACTICE.

NOTE. In the following Examples, unless it be expressly stated otherwise, the word "Triangle" is to be understood to mean a Spherical Triangle, whose sides are arcs of great circles.

1. Two arcs of great circles intersect at right angles in a point in the circumference of a small circle. If one of them touch the small circle, the other bisects it.

2. On the surface of a sphere draw a great circle passing through a given point, and touching a given small circle.

IN A RIGHT-ANGLED SPHERICAL TRIANGLE, C BEING
THE RIGHT ANGLE.

3. $2 \cos c = \cos(a + b) + \cos(a - b)$.

4. $\tan \frac{1}{2}(c + a) \tan \frac{1}{2}(c - a) = \tan^2 \frac{1}{2}b$.

5. $\sin^2 \frac{1}{2}c = \sin^2 \frac{1}{2}a \cos^2 \frac{1}{2}b + \cos^2 \frac{1}{2}a \sin^2 \frac{1}{2}b$.

6. $\sin^2 b \cos^2 a = \sin(c + a) \sin(c - a)$.

7. $\sin a \tan \frac{1}{2}A - \sin b \tan \frac{1}{2}B = \sin(a - b)$.

8. If $\cos A = \cos^2 a$, shew that $b + c$ is equal to $\frac{1}{2}\pi$ or $\frac{3}{2}\pi$, according as b and c are both less or both greater than $\frac{1}{2}\pi$.

9. $OA A_1$ is a spherical triangle, A_1 being a right angle; $A_1 A_2$ is an arc perpendicular to OA cutting it in A_2 , $A_2 A_3$ an arc perpendicular to OA_1 , $A_3 A_4$ an arc perpendicular to OA_2 , and so on; prove that $A_n A_{n+1}$ ultimately vanishes when n becomes infinite, and shew that $\cos AA_1 \cdot \cos A_1 A_2 \cdot \cos A_2 A_3 \dots$ (*ad infinitum*) = $\cos OA$.

10. Deduce the property of a plane right-angled triangle which corresponds to the formula $\cos c = \cos a \cos b$.

Ans. $c^2 = a^2 + b^2$.

11. If there be two right-angled spherical triangles ABC , $AB'C'$, having the angle A common to both,

$$\tan \frac{1}{2}(a + a') \tan \frac{1}{2}(c - c') = \tan \frac{1}{2}(a - a') \tan \frac{1}{2}(c + c').$$

12. If α, β be the arcs drawn from C , respectively perpendicular to c and bisecting c ;

$$\cot \alpha = \sqrt{(\cot^2 a + \cot^2 b)}; \quad \cot \beta = \frac{\cos a + \cos b}{\sqrt{(\sin^2 a + \sin^2 b)'}}$$

$$\sin \frac{1}{2}c = \frac{\sin \beta}{\sqrt{(1 + \sin^2 \alpha)'}}$$

13. Through the vertical angle A of an isosceles triangle there is drawn an arc of a great circle meeting the base in D ; shew that

$$\tan \frac{1}{2}BD \cdot \tan \frac{1}{2}CD = \tan \frac{1}{2}(BA + AD) \cdot \tan \frac{1}{2}(BA - AD).$$

14. If $A = a$, and $B = b$, then $C = 180^\circ - c$.

IN SPHERICAL TRIANGLES NOT RIGHT-ANGLED.

15. If each of the three sides be quadrants, and α, β, γ be the distances of a point within the triangle from the angular points;

$$\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma = 1.$$

16. If d be the length of the arc which bisects C and is terminated by the opposite side,

$$\tan d \sin(a + b) = 2 \sin a \sin b \cos \frac{1}{2}C.$$

17. If a and b be nearly equal,

$$a = \frac{1}{2}(a + b) + \tan \frac{1}{2}(a + b) \tan \frac{1}{2}(A - B) \cot \frac{1}{2}(A + B), \text{ very nearly.}$$

18. If one angle of a triangle, plane or spherical, be equal to the sum of the other two angles, the greatest side is double of the distance of its middle point from the opposite angle.

19. If O be any point in which arcs of great circles drawn through the angular points of ABC intersect, then

$$\begin{aligned} \frac{\sin A}{\sin a} &= \frac{\sin B}{\sin b} = \frac{\sin C}{\sin c} \\ &= \frac{\sin AO \sin BO \sin CO}{\sin a \sin b \sin c} \cdot \{ \cot AO \sin BOC + \cot BO \sin COA \\ &\quad + \cot CO \sin AOB \}. \end{aligned}$$

20. Find the locus of the vertices of all right-angled spherical triangles which have the same hypotenuse; and from the equation prove that the locus is a circle when the radius of the sphere is infinite.

21. Divide, by drawing an arc from an angle to the side opposite, a given triangle into two others whose areas are in a given ratio.

22. The sides of a spherical triangle are each $111^\circ, 28'$; find its angles, and shew that its area = $\frac{1}{4}$ surface of the sphere.

$$\{111^\circ, 28' = 2 \times (55^\circ, 44'), \text{ and } \tan \frac{1}{2} (55^\circ, 44') = \sqrt{2}\}.$$

23. If C be a right angle, E (the Spherical Excess)

$$= 2 \tan^{-1} (\tan \frac{1}{2} a \tan \frac{1}{2} b).$$

Also $\sin \frac{1}{2} E = \frac{\sin \frac{1}{2} a \sin \frac{1}{2} b}{\cos \frac{1}{2} c}$, and $\cos \frac{1}{2} E = \frac{\cos \frac{1}{2} a \cos \frac{1}{2} b}{\cos \frac{1}{2} c}$.

24. If E be the Spherical Excess,

$$\cot \frac{1}{2} E = \frac{\cot \frac{1}{2} a \cot \frac{1}{2} b + \cos C}{\sin C},$$

$$\text{or} = \frac{1 + \cos a + \cos b + \cos c}{2\sqrt{\sin S \sin (S-a) \sin (S-b) \sin (S-c)}}.$$

$$\cos \frac{1}{2} E = \frac{1 + \cos a + \cos b + \cos c}{4 \cos \frac{1}{2} a \cos \frac{1}{2} b \cos \frac{1}{2} c}.$$

In seconds, $\frac{1}{2} E = \frac{\tan \frac{1}{2} b \tan \frac{1}{2} c \sin A}{\sin 1''} - \frac{\tan^2 \frac{1}{2} b \tan^2 \frac{1}{2} c \sin 2A}{\sin 2''}$
 $+ \frac{\tan^3 \frac{1}{2} b \tan^3 \frac{1}{2} c \sin 3A}{\sin 3''} - \dots$

25. If P be the Perimeter and E the Spherical Excess of the triangle ABC , then

$$2 \sin \frac{1}{2} P \sin \frac{1}{2} A \sin \frac{1}{2} B \sin \frac{1}{2} C$$

$$= \{\sin \frac{1}{2} E \sin (A - \frac{1}{2} E) \sin (B - \frac{1}{2} E) \sin (C - \frac{1}{2} E)\}^{\frac{1}{2}}.$$

26. Determine the area of a spherical triangle from the data a, b, C ; and shew that if a and b be constant, and also $a + b$ be less than π , the area admits of a maximum value.

27. The angles of a spherical triangle, of which the area is $\frac{7}{12}\pi r^2$, where r is the radius of the sphere, form an arithmetic progression of which the common difference is 45° . Find them.

28. If the sides AC , BC of a triangle be produced to D and E , points such that $\tan \frac{1}{2}AC \cdot \tan \frac{1}{2}BC = \tan \frac{1}{2}DC \cdot \tan \frac{1}{2}EC$, and DE be joined by an arc of a great circle, the triangles ABC , CDE are of equal area.

29. If S be the surface of a spherical triangle whose angles are each 120° , and S' that of its polar triangle,

$$\tan \frac{1}{3}S : \tan \frac{1}{3}S' = 6\sqrt{2} + \sqrt{3} : 2\sqrt{2} - \sqrt{3}.$$

30. If a be one of the n sides of a regular spherical polygon, its surface (S) may be found from the equation

$$\cos \frac{\pi - \frac{1}{2}S}{n} = \cos \frac{\pi}{n} \sec \frac{1}{2}a.$$

31. The Spherical Excess of a triangle is $1''5$. Find its area, the radius of the Earth being taken to be 7757 miles.

32. If the three sides of a spherical triangle measured on the Earth's surface be 12, 16, and 18 miles, find the Spherical Excess.

33. A plane triangle whose sides are a , b , c , is placed in a sphere of radius r . Prove that the angle between the arcs of the great circles of which a and b are the chords is a right angle, if $2r\sqrt{(a^2 + b^2 - c^2)} = ab$.

34. If P be the pole of the small circle circumscribing a triangle ABC , prove that $\angle APB$ is double of the angle between the chords of AC and BC .

35. The middle points of the sides AB and AC of a triangle are D and E respectively, and P is the pole of DE ; shew that $\angle BPC$ is double of $\angle DPE$.

36. If an arc of a great circle be bisected, its segments will subtend equal angles at any point on the great circle of which its middle point is the pole.

37. A lune is formed by two great circles which intersect at a right angle; prove that from any point in one of the circles two arcs of great circles can be drawn to the other cutting it at equal angles, and find the least value of these angles. [The points are equidistant from the extremities of the lune.]

38. ABC and $A'B'C'$ are equal triangles; prove that arcs drawn at the middle points of the arcs of the great circles AA' , BB' , CC' , and at right angles to those lines, meet in a point at which AA' , BB' , CC' subtend equal angles. What limitation is there to this proposition? [The triangles ABC , $A'B'C'$ must be such that if placed one upon the other they would coincide.]

39. There is a great circle ABC , and AA' , BB' , CC' are arcs at right angles to it, which are reckoned positive on one side and negative on the other: prove that the condition of A' , B' , C' lying in a great circle is

$$\tan AA' \sin BC + \tan BB' \sin CA + \tan CC' \sin AB = 0.$$

40. Two quadrants (OA , OB) of great circles include a right angle; a great circle meets them in C , D respectively, and through P , any point in it, arcs of great circles APY , BPX are drawn meeting OB , OA respectively in Y , X ; if $OX = \theta$, $OY = \phi$, $OC = \alpha$, $OD = \beta$, shew that

$$\frac{\tan \theta}{\tan \alpha} + \frac{\tan \phi}{\tan \beta} = 1.$$

41. Two great circles, inclined at an angle ω , intersect at O . NN' and MM' are equal arcs on the two circles respectively, and NM and $N'M'$ are arcs perpendicular to OMM' ; if $NM = \delta$, $N'M' = \delta'$, shew that $\cos \delta \cos \delta' = \cos \omega$; also that

$$\cos MM' = \frac{\sin \delta \sin \delta'}{1 - \cos \omega}, \quad \cos NN' = \sin ON \cdot \sin ON' \cdot (1 + \cos \omega).$$

42. P is the pole of a small circle; $S_1 S_2 S_3 \dots S_n$ a series of points in this circle equidistant from one another; if Z be any other point and ZS_1, ZS_2, \dots be joined, shew that the sum of the cosines of $ZS_1, ZS_2, \dots = n \cos PZ \cdot \cos \delta$, where δ is the radius of the small circle.

43. Three small circles are inscribed in a spherical triangle whose angles are each 120° , in such a manner that each circle touches each of the other circles and also two sides of the triangle; prove that the radius of each circle is 30° , and that the centers of the circles coincide with the angular points of the polar triangle.

44. Three small circles, whose radii are ρ_1, ρ_2, ρ_3 , touch one another in P, Q, R . If A, B, C be the degrees in the angles of a spherical triangle formed by joining their centers, prove that

$$\text{Area } PQR = (A \cos \rho_1 + B \cos \rho_2 + C \cos \rho_3 - 180) r^2,$$

r being the radius of the sphere.

45. If in a triangle, R , r be the radii of the small circumscribing and inscribed circles, and r_1 , r_2 , r_3 the radii of the circles touching one side of the triangle and the other two sides produced, prove that $\cot r_1 + \cot r_2 + \cot r_3 - \cot r = 2 \tan R$, and shew from this result that the corresponding property in a plane triangle is

$$\frac{1}{r_1} + \frac{1}{r_2} + \frac{1}{r_3} = \frac{1}{r}.$$

46. Determine the points in the sides of a triangle at which tangents being drawn, they will meet two and two and form a triangle. If A' , B' , C' be the angles of this triangle, prove that

$$\tan \frac{1}{2} A' = \cos(S-a) \tan \frac{1}{2} A.$$

47. Prove that the square of the area of the triangle formed by the tangents as in the last question is equal to

$r^4 \{ \tan(S-a) + \tan(S-b) + \tan(S-c) \} \tan(S-a) \tan(S-b) \tan(S-c)$,
 r being the radius of the sphere.

48. If the vertical angle of a triangle be equal to the sum of the angles at the base, the locus of the vertex, while the base remains fixed, will be the small circle described with the middle point of the base as pole and the base as diameter.

49. $ABCD$ is a quadrilateral whose sides are arcs of great circles, E and F the middle points of AC and BD ; prove that
 $\cos AB + \cos BC + \cos CD + \cos DA = 4 \cos \frac{1}{2} AC \cdot \cos \frac{1}{2} BD \cdot \cos EF$.

50. If $ABCD$ be a spherical quadrilateral, P the intersection of AB and DC , Q that of AD and BC , R that of AC and BD , shew that

$$\begin{aligned} & \sin AB \cdot \sin CD \cdot \cos P - \sin AD \cdot \sin BC \cdot \cos Q \\ &= \cos AD \cdot \cos BC - \cos AB \cdot \cos CD = \sin AC \cdot \sin BD \cdot \cos R. \end{aligned}$$

51. If α , β , γ be the angles which three diameters of a sphere of radius a make with one another, prove that the volume of the parallelepiped formed by tangent planes at their extremities, (2σ being $= \alpha + \beta + \gamma$), is

$$\frac{4a^3}{\sqrt{\{\sin \sigma \sin(\sigma - \alpha) \sin(\sigma - \beta) \sin(\sigma - \gamma)\}}}$$

52. If i be the inclination of a plane to the horizon, and α the inclination of a line in it to the intersection of the plane with the horizontal plane, the inclination θ of the line to the horizon will be found from the equation, $\sin \theta = \sin i \sin \alpha$.

53. The shadow of a cloud is observed to fall upon a spot at a known distance on the side of a hill. Given the altitudes and the azimuths of the cloud and shadow, and the azimuth of the sun, find the distance of the cloud.

Ans. If $\alpha_1, \alpha_2, \alpha_3$ be the azimuths of the Sun, cloud, and shadow respectively, and a_1, a_2 the altitudes of the cloud and shadow, d the known distance, the distance of the cloud is

$$d \cdot \frac{\cos a_2 \cdot \sin (\alpha_3 - \alpha_1)}{\cos a_1 \cdot \sin (\alpha_2 - \alpha_1)}.$$

54. If Z be the zenith, K the pole of the limb (which is *not exactly* horizontal) of a theodolite, and S be an object whose azimuth is observed, the error is known from the equation,

$$\angle SKZ - \angle SZK = KZ \cdot \cot SK \cdot \sin SZK.$$

55. If Z be the zenith, K the pole of the circle of a theodolite, which is *not exactly* vertical, and KZ be produced to Q till KQ is a quadrant, then if S be an object whose zenith distance is to be observed, the error of observation is known from the equation,

$$SQ - SZ = \cot SQ \cdot \sin \frac{1}{2} QZ.$$

56. Given two sides, a and b , of a triangle, spherical or plane, and the included angle C , to find the variation produced in A corresponding to a small given variation in C .

57. If a solid be bounded by plane figures, of which some have an odd and some have an even number of sides, shew that there must be an even number of those faces which have an odd number of sides.

58. In a triangle C and c remain constant, and a, b receive small increments $\delta a, \delta b$ respectively; shew that

$$\frac{\delta a}{\sqrt{(1 - n^2 \sin^2 a)}} + \frac{\delta b}{\sqrt{(1 - n^2 \sin^2 b)}} = 0; \text{ where } n = \frac{\sin C}{\sin c}$$

59. A solid is formed of an equal number of faces bounded by 3, 4, and 5 sides; find the least number of faces in such a solid, and the numbers of its edges and solid angles.

$$\text{Ans. } F = 6, \quad E = 12, \quad S = 8.$$

60. Every solid in which four or more edges meet in each solid angle must have at least eight triangular faces; and those in which five or more meet in each angle must have at least twenty triangular faces. Also no solid can be formed so that not less than six edges meet in each solid angle.

61. No solid is entirely composed of faces all of which have more than five sides; and if there be neither quadrilateral nor pentagonal faces, there must be more than four triangular faces, unless the faces be all triangles.

62. The base of a pyramid is an equilateral hexagon whose alternate angles are equal and adjacent angles are unequal. Given the angles between any face of the pyramid and two adjacent faces, find the angle between a normal to any face and the axis of the pyramid. *Ans.* The angle required is equal to that which any plane face makes with the base. If α, α' be the given angles, and θ be the angle required, then $\text{Sin } \theta = 2 \cos \frac{1}{4}(\alpha' + \alpha) \cdot \cos \frac{1}{4}(\alpha' - \alpha) \cdot \sec \phi$, where ϕ is known from the equation

$$\text{Tan } \phi = \frac{1}{\sqrt{3}} \cdot \tan \frac{1}{4}(\alpha' + \alpha) \cdot \tan \frac{1}{4}(\alpha' - \alpha).$$

63. If r, R be the radii of the spheres described within and about a regular tetrahedron, r', R' the radii of the spheres to which the edges, and one face and the planes of the three others produced, are respectively tangents, prove that

$$r' = \sqrt{rR}, \quad \text{and} \quad R' = \sqrt{2rR}.$$

64. Given the six edges of a triangular pyramid, to find its volume.

65. Of all triangular pyramids of given volume, the regular tetrahedron has the least surface.

66. A cube is turned round one of its diagonals through 180° ; shew that $2\sqrt{2}$ is the tangent of the angle at which any one of the faces is inclined to its original position.

67. The diagonal of a cube is produced until the length of the part produced is equal to the diagonal, and from the extreme point as pole the cube is projected on a plane perpendicular to the diagonal. Shew that the projection will be an equilateral hexagon, in which the alternate angles are equal and the adjacent angles are unequal, their sines being in the ratio of 8 to 5.

London and Cambridge.

MACMILLAN AND CO.'S

DESCRIPTIVE CATALOGUE OF CAMBRIDGE CLASS BOOKS.

The Works in this Series of CAMBRIDGE CLASS-BOOKS FOR THE USE OF SCHOOLS AND COLLEGES, which have been issued at intervals during the last ten years, are intended to embrace all Branches of Education, from the most Elementary to the most Advanced, and to keep pace with the latest discoveries in Science.

Of those hitherto published the large and ever increasing sale is a sufficient indication of the manner in which they have been appreciated by the public.

A SERIES of a more Elementary character is in preparation, a list of which will be found on page 2 of this Catalogue.

ELEMENTARY SCHOOL CLASS BOOKS.

The volumes of this Series of ELEMENTARY SCHOOL CLASS BOOKS are handsomely printed in a form that, it is hoped, will assist the young student as much as clearness of type and distinctness of arrangement can effect. They are published at a moderate price to ensure an extensive sale in the Schools of the United Kingdom and the Colonies.

1. *Euclid for Colleges and Schools.*

By I. TODHUNTER, M.A., F.R.S., Fellow and Principal Mathematical Lecturer of St. John's College, Cambridge. 18mo. 3s. 6d.

2. *An Elementary Latin Grammar.*

By H. J. ROBY, M.A., Under Master of Dulwich College Upper School, late Fellow and Classical Lecturer of St. John's College, Cambridge. 18mo. 2s. 6d.

3. *An Elementary History of the Book of Common Prayer.*

By FRANCIS PROCTER, M.A., Vicar of Witton, Norfolk, late Fellow of St. Catharine's College, Cambridge. 18mo. 2s. 6d.

4. *Mythology for Latin Versification.*

A Brief Sketch of the Fables of the Ancients, prepared to be rendered into Latin Verse, for Scholars. By F. C. HODGSON, B.D., late Provost of Eton College. *New Edition.* Revised by F. C. HODGSON, M.A., Fellow of King's College, Cambridge. 18mo. 3s.

5. *Algebra for Beginners.*

By I. TODHUNTER, M.A., F.R.S. [In the Press.

6. *The School Class Book of Arithmetic.*

By BARNARD SMITH, M.A., late Fellow of St. Peter's College, Cambridge. [In the Press.

7. *The Bible Word-Book.*

A Glossary of old English Bible Words with Illustrations. By J. EASTWOOD, M.A., St. John's College, Cambridge, and Incumbent of Hope in Hanley, Stafford, and W. ALDIS WRIGHT, M.A., Trinity College, Cambridge. [Preparing.

*** Other Volumes will be announced in due course.

CAMBRIDGE CLASS BOOKS

FOR

SCHOOLS AND COLLEGES.

WORKS by the Rev. BARNARD SMITH, M.A.

Fellow of St. Peter's College, Cambridge.

1.

Arithmetic & Algebra

In their Principles and Applications.

With numerous Examples, systematically arranged.

Eighth Edit. 696 pp. (1861). Cr. 8vo. strongly bound in cloth. 10s. 6d.

The first edition of this work was published in 1854. It was primarily intended for the use of students at the Universities, and for Schools which prepare for the Universities. It has however been found to meet the requirements of a much larger class, and is now extensively used in *Schools and Colleges both at home and in the Colonies*. It has also been found of great service for students preparing for the MIDDLE-CLASS and CIVIL AND MILITARY SERVICE EXAMINATIONS, from the care that has been taken to elucidate the *principles* of all the Rules. Testimony of its excellence has been borne by some of the highest practical and theoretical authorities, of which the following from the late DEAN PEACOCK may be taken as a specimen:

“Mr. Smith's Work is a most useful publication. The Rules are stated with great clearness. The Examples are well selected and worked out with just sufficient detail without being encumbered by too minute explanations; and there prevails throughout it that just proportion of theory and practice, which is the crowning excellence of an elementary work.”

2. **Arithmetie**

For the Use of Schools.

New Edition (1862) 348 pp.

Crown 8vo strongly bound in cloth, 4s. 6d. Answers to all the Questions.

3. **Key** to the above, containing Solutions to all the Questions in the latest Edition. Crown 8vo. cloth. 392 pp. Second Edit. 8s. 6d.

To meet a widely expressed wish, the ARITHMETIC was published separately from the larger work in 1854, with so much alteration as was necessary to make it quite independent of the ALGEBRA. It has now a very large sale in all classes of Schools at home and in the Colonies. A copious collection of Examples, under each rule, has been embodied in the work in a systematic order, and a Collection of Miscellaneous Papers in all branches of Arithmetic is appended to the book.

4. **Exercises in Arithmetic.**

104 pp. Cr. 8vo. (1860)

2s. (Or with ANSWERS, 2s. 6d.)

Also sold separately in 2 Parts
1s. each. **Answers, 6d.**

These EXERCISES have been published in order to give the pupil examples in every rule of Arithmetic. The greater number have been carefully compiled from the latest University and School Examination Papers.

WORKS by ISAAC TODHUNTER, M.A. F.R.S.

Fellow and Principal Mathematical Lecturer of St. John's College, Cambridge.

1. Algebra.

*For the Use of Colleges and Schools.*Third Edition. 512 pp. (1862).
Strongly bound in cloth. 7s. 6d.

This work contains all the propositions which are usually included in elementary treatises on Algebra, and a large number of *Examples for Exercise*. The author has sought to render the work easily intelligible to students without impairing the accuracy of the demonstrations, or contracting the limits of the subject. The Examples have been selected with a view to illustrate every part of the subject, and as the number of them is about *Sixteen hundred and fifty*, it is hoped they will supply ample exercise for the student. Each set of Examples has been carefully arranged, commencing with very simple exercises, and proceeding gradually to those which are less obvious.

2. Plane Trigonometry

*For Schools and Colleges.*2nd Edit. 279 pp. (1860). Crn. 8vo.
Strongly bound in cloth. 5s.

The design of this work has been to render the subject intelligible to beginners, and at the same time to afford the student the opportunity of obtaining all the information which he will require on this branch of Mathematics. Each chapter is followed by a set of Examples; those which are entitled *Miscellaneous Examples*, together with a few in some of the other sets, may be advantageously reserved by the student for exercise after he has made some progress in the subject. As the Text and Examples have been tested by considerable experience in teaching, the hope is entertained that they will be suitable for imparting a sound and comprehensive knowledge of Plane Trigonometry, together with readiness in the application of this knowledge to the solution of problems. In the Second Edition the hints for the solution of the Examples have been considerably increased.

3. Spherical Trigonometry.

*For the Use of Colleges and Schools.*112 pp. Crown 8vo. (1859).
Strongly bound in cloth. 4s. 6d.

This work is constructed on the same plan as the *Treatise on Plane Trigonometry*, to which it is intended as a sequel. Considerable labour has been expended on the text in order to render it comprehensive and accurate, and the Examples, which have been chiefly selected from University and College Papers, have all been carefully verified.

The Elements of Euclid

*For the Use of Schools and Colleges.*COMPRISING THE FIRST SIX BOOKS AND
PORTIONS OF THE ELEVENTH AND
TWELFTH BOOKS, WITH NOTES, AP-
PENDIX, AND EXERCISES

384 pp. 18mo. bound. (1862). 3s. 6d.

As the Elements of Euclid are usually placed in the hands of young students, it is important to exhibit the work in such a form as will assist them in overcoming the difficulties which they experience on their first introduction to processes of continuous argument. No method appears to be so useful as that of breaking up the demonstrations into their constituent parts, and this plan has been adopted in the present edition. Each distinct assertion in the argument begins a new line; and at the end of the lines are placed the necessary references to the preceding principles on which the assertions depend. The longer propositions are distributed into subordinate parts, which are distinguished by breaks at the beginning of the lines. The Notes are intended to indicate and explain the principal difficulties, and to supply the most important inferences which can be drawn from the propositions. The work finishes with a collection of *Six hundred and twenty-five Exercises*, which have been selected principally from Cambridge Examination papers and have been tested by long experience. As far as possible they are arranged in order of difficulty. The Figures will be found to be large and distinct, and have been repeated when necessary, so that they always occur in immediate connexion with the corresponding text.

FOR SCHOOLS AND COLLEGES.

WORKS by ISAAC TODHUNTER, M.A., F.R.S.—continued.

5.

The Integral Calculus

And its Applications.

With numerous Examples.

Second Edition. 342 pp. (1862).

Crown 8vo. cloth. 10s. 6d.

In writing the present *Treatise on the Integral Calculus*, the object has been to produce a work at once elementary and complete—adapted for the use of beginners, and sufficient for the wants of advanced students. In the selection of the propositions, and in the mode of establishing them, the author has endeavoured to exhibit fully and clearly the principles of the subject, and to illustrate all their most important results. In order that the student may find in the volume all that he requires, a large collection of Examples for exercise has been appended to the different chapters.

6. Analytical Statics.

With numerous Examples.

Second Edition. 330 pp. (1858).

Crown 8vo. cloth. 10s. 6d.

In this work will be found all the propositions which usually appear in treatises on Theoretical Statics. To the different chapters Examples are appended, which have been selected principally from the University and College Examination Papers, these will furnish ample exercise in the application of the principles of the subject.

7. EXAMPLES OF

Analytical Geometry of Three Dimensions.

76 pp. (1858). Crn. 8vo. cloth. 4s.

A collection of examples in illustration of Analytical Geometry of Three Dimensions has long been required both by students and teachers, and the present work is published with the view of supplying the want.

8. The

Differential Calculus.

With numerous Examples.

Third Edition, 398 pp. (1860).

Crown 8vo. cloth, 10s. 6d.

This work is intended to exhibit a comprehensive view of the Differential Calculus on the method of Limits. In the more elementary portions, explanations have been given in considerable detail, with the hope that a reader who is without the assistance of a tutor may be enabled to acquire a competent acquaintance with the subject. More than one investigation of a theorem has been frequently given, because it is believed that the student derives advantage from viewing the same proposition under different aspects, and that in order to succeed in the examinations which he may have to undergo, he should be prepared for a considerable variety in the order of arranging the several branches of the subject, and for a corresponding variety in the mode of demonstration.

9. Plane Co-Ordinate Geometry

AS APPLIED TO THE STRAIGHT LINE
AND THE CONIC SECTIONS.

With numerous Examples.

Third and Cheaper Edition.

Crn. 8vo. cl. 326 pp. (1862). 7s. 6d.

This *Treatise* exhibits the subject in a simple manner for the benefit of beginners, and at the same time includes in one volume all that students usually require. The Examples at the end of each chapter will, it is hoped, furnish sufficient exercise, as they have been carefully selected with the view of illustrating the most important points, and have been tested by repeated experience with pupils. In consequence of the demand for the work proving much greater than had been originally anticipated, a large number of copies of the *Third Edition* has been printed, and a considerable reduction effected in the price.

By ISAAC TODDINGTON, M.A.

AN ELEMENTARY TREATISE ON THE
Theory of Equations.

With a Collection of Examples.

Crown 8vo. cloth. 279 pp. (1861).
7s. 6d.

This treatise contains all the propositions which are usually included in elementary treatises on the Theory of Equations, together with a collection of Examples for exercise. This work may in fact be regarded as a sequel to that on Algebra by the same writer, and accordingly the student has occasionally been referred to the treatise on Algebra for preliminary information on some topics here discussed. The work includes three chapters on Determinants.

11. **History of the Progress
of the
Calculus of Variations**

During the Nineteenth Century.

8vo. cloth. 532 pp. (1861). 12s.

It is of importance that those who wish to cultivate any subject may be able to ascertain what results have already been obtained, and thus reserve their strength for difficulties which have not yet been conquered. The Author has endeavoured in this work to ascertain distinctly what has been effected in the Progress of the Calculus, and to form some estimate of the manner in which it has been effected.

A TREATISE ON

**Mechanics and Hydro-
statics.**

With Solutions of Questions

PROPOSED IN THE CAMBRIDGE SENATE HOUSE

By W. H. GIRDLESTONE, M.A.

Christ's College.

8vo. cloth. 100 pp. 1862.

By J. H. PRATT, M.A.

Archdeacon of Calcutta, late Fellow of
Gonville and Caius College, Cambridge.

**A Treatise on
Attractions,**

*La Place's Functions, and the Figure
of the Earth.*

Second Edition. Crown 8vo. 126 pp.
(1861). cloth. 6s. 6d.

In the present Treatise the author has endeavoured to supply the want of a work on a subject of great importance and high interest—La Place's Coefficients and Functions, and the calculation of the Figure of the Earth by means of his remarkable analysis. No student of the higher branches of Physical Astronomy should be ignorant of La Place's analysis and its result—"a calculus," says AIRY, "the most singular in its nature and the most powerful in its application that has ever appeared."

By G. B. AIRY, M.A.

Astronomer Royal.

1. **Mathematical Tracts**

*On the Lunar and Planetary Theories,
Figure of the Earth, the Undulatory
Theory of Optics, &c.*

Fourth Edition. 400 pp. (1858).
8vo. 15s.

2. **Theory of Errors of
Observations**

And the Combination of Observations.

103 pp. (1861). Crown 8vo. 6s. 6d.

In order to spare astronomers and observers in natural philosophy the confusion and loss of time which are produced by referring to the ordinary treatises embracing both branches of Probabilities, the author has thought it desirable to draw up this work, relating only to Errors of Observation, and to the rules derivable from the consideration of these Errors, for the Combination of the Results of Observations. The Author has thus also the advantage of entering somewhat more fully into several points of interest to the observer, than can possibly be done in a General Theory of Probabilities.

By *GEORGE BOOLE, D.C.L., F.R.S.*
Professor of Mathematics in the Queen's
University, Ireland.

Differential Equations

468 pp. (1859). Crn. 8vo. cloth. 14s.

The Author has endeavoured in this treatise to convey as complete an account of the present state of knowledge on the subject of Differential Equations as was consistent with the idea of a work intended, primarily, for elementary instruction. The object has been first of all to meet the wants of those who had no previous acquaintance with the subject, and also not quite to disappoint others who might seek for more advanced information. The earlier sections of each chapter contain that kind of matter which has usually been thought suitable for the beginner, while the latter ones are devoted either to an account of recent discovery, or to the discussion of such deeper questions of principle as are likely to present themselves to the reflective student in connection with the methods and processes of his previous course.

2. The Calculus of Finite Differences.

248 pp. (1860). Crown 8vo. cloth.
10s. 6d.

In this work particular attention has been paid to the connexion of the methods with those of the Differential Calculus—a connexion which in some instances involves far more than a merely formal analogy. The work is in some measure designed as a sequel to the Author's *Treatise on Differential Equations*, and it has been composed on the same plan.

Elementary Statics.

By the Rev. *GEORGE RAWLINSON*
Professor of Applied Sciences, Elphinstone Coll., Bombay.

Edited by the Rev. *E. STURGES, M.A.*
Rector of Kencott, Oxfordshire.

(150 pp.) 1860. Crn. 8vo. cl. 4s. 6d.

This work is published under the authority of H. M. Secretary of State for India for use in the Government Schools and Colleges in India.

By *P. G. TAIT, M.A., and*
W. J. STEELE, B.A.

Late Fellows of St Peter's Coll. Camb.

Dynamics of a Particle.

With numerous Examples.

304 pp. (1856). Cr. 8vo. cl. 10s. 6d.

In this Treatise will be found all the ordinary propositions connected with the Dynamics of Particles which can be conveniently reduced without the use of D'Alembert's Principles. Throughout the book will be found a number of illustrative Examples introduced in the text, and for the most part completely worked out; others, with occasional solutions or hints to assist the student are appended to each Chapter.

By the Rev. *G. F. CHILDE, M.A.*
Mathematical Professor in the South
African College.

Singular Properties of the Ellipsoid

And Associated Surfaces of the n th
Degree.

152 pp. (1861). 8vo. boards. 10s. 6d.

As the title of this volume indicates, its object is to develop peculiarities in the Ellipsoid; and further, to establish analogous properties in unlimited congeneric series of which this remarkable surface is a constituent.

By *J. B. PHEAR, M.A.*
Fellow and late Mathematical Lecturer of
Clare College.

Elementary Hydrostatics

With numerous Examples and
Solutions.

Second Edition. 156 pp. (1857).
Crown 8vo. cloth. 5s. 6d.

"An excellent Introductory Book. The definitions are very clear; the descriptions and explanations are sufficiently full and intelligible; the investigations are simple and scientific. The examples greatly enhance its value."—*ENGLISH JOURNAL OF EDUCATION.*

This Edition contains 147 Examples, and solutions to all these examples are given at the end of the book.

By Rev. S. PARKINSON, B.D.

Fellow and Prætor of St. John's Coll.
Cambridge.

By R. D. BEASLEY, M.A.

Head Master of Grantham School.

**1. Elementary Treatise
on Mechanics.***With a Collection of Examples.*Second Edition. 345 pp.ⁿ (1861).
Crown 8vo. cloth. 9s. 6d.

The Author has endeavoured to render the present volume suitable as a Manual for the junior classes in Universities and the higher classes in Schools. With this object there have been included in it those portions of theoretical Mechanics which can be conveniently investigated without the Differential Calculus, and with one or two short exceptions the student is not presumed to require a knowledge of any branches of Mathematics beyond the elements of Algebra, Geometry, and Trigonometry. A collection of Problems and Examples has been added, chiefly taken from the Senate-House and College Examination Papers—which will be found useful as an exercise for the student. In the Second Edition several additional propositions have been incorporated in the work for the purpose of rendering it more complete, and the Collection of Examples and Problems has been largely increased.

2. A Treatise on Optics

304 pp. (1859). Crown 8vo. 10s. 6d.

A collection of Examples and Problems has been appended to this work which are sufficiently numerous and varied in character to afford useful exercise for the student; for the greater part of them recourse has been had to the Examination Papers set in the University and the several Colleges during the last twenty years.

Subjoined to the copious Table of Contents, the author has ventured to indicate an elementary course of reading not unsuitable for the requirements of the First Three Days in the Cambridge Senate House Examinations.

**AN ELEMENTARY TREATISE ON
Plane Trigonometry.***With a numerous Collection of
Examples.*106 pp. (1858), strongly bound in
cloth. 3s. 6d.

This Treatise is specially intended for use in Schools. The choice of matter has been chiefly guided by the requirements of the three days' Examination at Cambridge, with the exception of proportional parts in logarithms, which have been omitted. About *Four hundred* Examples have been added, mainly collected from the Examination Papers of the last ten years, and great pains have been taken to exclude from the body of the work any which might dishearten a beginner by their difficulty.



By J. BROOK SMITH, M.A.

St. John's College, Cambridge.

**Arithmetic in Theory
and Practice.***For Advanced Pupils.*

PART I. Crown 8vo. cloth. 3s. 6d.

This work forms the first part of a Treatise on Arithmetic, in which the Author has endeavoured, from very simple principles, to explain in a full and satisfactory manner all the important processes in that subject.

The proofs have in all cases been given in a form entirely arithmetical: for the author does not think that recourse ought to be had to Algebra until the arithmetical proof has become hopelessly long and perplexing.

At the end of every chapter several examples have been worked out at length, in which the best practical methods of operation have been carefully pointed out.

By G. H. PUCKLE, M.A.

Principal of Windermere College.

Conic Sections and Algebraic Geometry.*With numerous Easy Examples Progressively arranged.*

Second Edition. 264 pp. (1856).

Crown 8vo. 7s. 6d.

This book has been written with special reference to those difficulties and misapprehensions which commonly beset the student when he commences. With this object in view, the earlier part of the subject has been dwelt on at length, and geometrical and numerical illustrations of the analysis have been introduced. The Examples appended to each section are mostly of an elementary description. The work will, it is hoped, be found to contain all that is required by the upper classes of schools and by the generality of students at the Universities.



By EDWARD JOHN ROUTH, M.A.

Fellow and Assistant Tutor of St. Peter's College, Cambridge.

Dynamics of a System of Rigid Bodies.*With numerous Examples.*

336 pp. (1860). Crown 8vo. cloth. 10s. 6d.

CONTENTS: Chap. I. Of Moments of Inertia.—II. D'Alembert's Principle.—III. Motion about a Fixed Axis.—IV. Motion in Two Dimensions.—V. Motion of a Rigid Body in Three Dimensions.—VI. Motion of a Flexible String.—VII. Motion of a System of Rigid Bodies.—VIII. Of Impulsive Forces.—IX. Miscellaneous Examples.

The numerous Examples which will be found at the end of each chapter have been chiefly selected from the Examination Papers set in the University and Colleges of Cambridge during the last few years.

• • The

Cambridge Year Book

AND UNIVERSITY ALMANACK

For 1863.

Crown 8vo. 228 pp. price 2s. 6d.

The specific features of this annual publication will be obvious at a glance, and its value to teachers engaged in preparing students for, and to parents who are sending their sons to, the University, and to the public generally, will be clear.

1. The whole mode of proceeding in entering a student at the University and at any particular College is stated.

2. The course of the studies as regulated by the University examinations, the manner of these examinations, and the specific subjects and times for the year 1863, are given.

3. A complete account of all Scholarships and Exhibitions at the several Colleges, their value, and the means by which they are gained.

4. A brief summary of all Graces of the Senate, Degrees conferred during the year 1861, and University news generally are given.

5. The Regulations for the Local Examination of those who are not members of the University, to be held this year, with the names of the books on which the Examination will be based, and the date on which the Examination will be held.



By N. M. FERRERS, M.A.

Fellow and Mathematical Lecturer of Gonville and Caius College, Cambridge.

AN ELEMENTARY TREATISE ON

Trilinear Co-Ordinates*The Method of Reciprocal Polars, and the Theory of Projections.*

154 pp. (1861). Cr. 8vo. cl. 6s. 6d.

The object of the Author in writing on this subject has mainly been to place it on a basis altogether independent of the ordinary Cartesian System, instead of regarding it as only a special form of abridged Notation. A short chapter on Determinants has been introduced.

By J. C. SNOWBALL, M.A.

Late Fellow of St. John's Coll. Cambridge.

Plane and Spherical Trigonometry.

*With the Construction and Use of
Tables of Logarithms.*

Ninth Edition. 240 pp. (1857).
Crown 8vo. 7s. 6d.

In preparing a new edition, the proofs of some of the more important propositions have been rendered more strict and general; and a considerable addition of more than *Two hundred Examples*, taken principally from the questions in the Examinations of Colleges and the University, has been made to the collection of Examples and Problems for practice.

By W. H. DREW, M.A.

Second Master of Blackheath School.

Geometrical Treatise on Conic Sections.

With a copious Collection of Examples.

Second Edition. Crown 8vo. cloth.
4s. 6d.

In this work the subject of Conic Sections has been placed before the student in such a form that, it is hoped, after mastering the elements of Euclid, he may find it an easy and interesting continuation of his geometrical studies. With a view also of rendering the work a complete Manual of what is required at the Universities, there have been either embodied into the text, or inserted among the examples, every book work question, problem, and rider, which has been proposed in the Cambridge examinations up to the present time.

Solutions to the Problems in Drew's Conic Sections.

Crown 8vo. cloth. 4s. 6d.

Senate-House Mathematical Problems.

With Solutions.

1848-51. By FERREUS and JACKSON. 8vo.
15s. 6d.

1848-51. (RIDEEN) By JAMISON. 8vo.
7s. 6d.

1854. By WALTON and MACKENZIE.
10s. 6d.

1857. By CAMPION and WALTON. 8vo.
8s. 6d.

1860. By ROUTH and WATSON. Crown
8vo. 7s. 6d.

The above books contain Problems and Examples which have been set in the Cambridge Senate-house Examinations at various periods during the last twelve years, together with Solutions of the same. The Solutions are in all cases given by the Examiners themselves or under their sanction.

By H. A. MORGAN, M.A.

Fellow of Jesus College, Cambridge.

A Collection of Mathematical Problems and Examples.

With Answers.

190 pp. (1858). Crown 8vo. 6s. 6d.

This book contains a number of problems, chiefly elementary, in the Mathematical subjects usually read at Cambridge. They have been selected from the papers set during late years at Jesus College. Very few of them are to be met with in other collections, and by far the larger number are due to some of the most distinguished Mathematicians in the University.

Cambridge University Examination Papers.

Crown 8vo. 184 pp. 2s. 6d.

A Collection of all the Papers set at the Examinations for the Degrees, the various Triposes, and the Theological Certificates in the University, with List of Candidates Examined and of those Approved, and an Index to the Subjects. 1860-61.

**A Treatise on
Solid Geometry.**

By *PERCIVAL FROST, M.A.*,
St. John's College, and
JOSEPH VOLSTENHOLME, M.A.,
Christ's Coll. Cambridge.

472 pp. 8vo. cloth. 18s. 1863.

The authors have endeavoured to present before students as comprehensive a view of the subject as possible. Intending as they have done to make the subject accessible, at least in the earlier portion, to all classes of students, they have endeavoured to explain fully all the processes which are most useful in dealing with ordinary theorems and problems, thus directing the student to the selection of methods which are best adapted to the exigencies of each problem. In the more difficult portions of the subject, they have considered themselves to be addressing a higher class of students, there they have tried to lay a good foundation on which to build, if any reader should wish to pursue the science beyond the limits to which the work extends.

AN ELEMENTARY TREATISE ON
The Planetary Theory.

WITH A COLLECTION OF PROBLEMS.

By *C. H. H. CHEYNE, B.A.*
Scholar of St. John's College, Cambridge.
148 pp. 1862. Crn. 8vo. cloth. 6s. 6d.

In this volume, an attempt has been made to produce a Treatise on the Planetary Theory, which being elementary in character, should be so far complete, as to contain all that is usually required by students in the University. A collection of Problems has been added, taken chiefly from Cambridge Examination papers of the last twenty years.

By *JOHN E. B. MAJOR, M.A.*
Fellow and Classical Lecturer of St John's
College, Cambridge.

1. Juvenal.

With English Notes.
464 pp. (1854). Crown 8vo. cloth.
10s. 6d.

"A School edition of Juvenal, which, for really nice scholarship, extensive acquaintance with Latin literature, and familiar knowledge of Continental criticism, ancient and modern, is unsurpassed, we do not say among English School-books, but among English editions generally"—
EDINBURGH REVIEW.

**2. Cicero's
Second Philippic.**

With English Notes.
168 pp. (1861). Fcp. 8vo. cloth. 5s.

The Text is that of Halm's 2nd edition, (Leipzig, Weidmann, 1858), with some corrections from Madvig's 4th Edition (Copenhagen, 1858). Halm's Introduction has been closely translated, with some additions. His notes have been curtailed, omitted, or enlarged, at discretion; passages to which he gives a bare reference, are for the most part printed at length; for the Greek extracts an English version has been substituted. A large body of notes, chiefly grammatical and historical, has been added from various sources. A list of books useful to the student of Cicero, a copious Argument, and an Index to the introduction and notes, complete the book.

By *P. FROST, Jun., M.A.*
Late Fellow of St. John's Coll. Cambridge.

Thucydides. Book VI.

With English Notes, Map and Index.
8vo. cloth. 7s. 6d.

It has been attempted in this work to facilitate the attainment of accuracy in translation. With this end in view the Text has been treated grammatically.

By B. DRAKE, M.A.

Late Fellow of King's Coll. Cambridge.

1. Demosthenes on the Crown.

With English Notes.

Second Edition. To which is prefixed ÆSCHINES AGAINST CTESIPHON. With English Notes. 287 pp. (1860). Fcap. 8vo. cl. 5s.

The first edition of the late Mr. Drake's edition of Demosthenes de Corona having met with considerable acceptance in various Schools, and a new edition being called for, the Oration of Æschines against Ctesiphon, in accordance with the wishes of many teachers, has been appended with useful notes by a competent scholar.

2. Æschyli Eumenides

With English Verse Translation, Copious Introduction, and Notes.

8vo. 144. pp. (1853). 7s. 6d.

"Mr. Drake's ability as a critical Scholar is known and admitted. In the edition of the Eumenides before us we meet with him also in the capacity of a Poet and Historical Essayist. The translation is flowing and melodious, elegant and scholarlike. The Greek Text is well printed: the notes are clear and useful"—GUARDIAN.

By C. MERIVALE, B.D.

Author of "History of Rome," &c.

Sallust.

With English Notes.

Second Edition. 172 pp. (1858). Fcap. 8vo. 4s. 6d.

"This School edition of Sallust is precisely what the School edition of a Latin author ought to be. No useless words are spent in it, and no words that could be of use are spared. The text has been carefully collated with the best editions. With the work is given a full current of extremely well-selected annotations."—THE EXAMINER.

The "CATILINA" and "JUGURTHA" may be had separately, price 2s. 6d. each, bound in cloth.

By J. WRIGHT, M.A.

Head Master of Sutton Coldfield School

1. Help to Latin Grammar.

With Easy Exercises, and Vocabulary

Crown 8vo. cloth. 4s. 6d.

Never was there a better aid offered alike to teacher and scholar in that arduous pass. The style is at once familiar and strikingly simple and lucid; and the explanations precisely hit the difficulties, and thoroughly explain them."—ENGLISH JOURNAL OF EDUCATION.

2. Hellenica.

A FIRST GREEK READING BOOK.

Second Edit. Fcap. 8vo. cl. 3s. 6d.

In the last twenty chapters of this volume, Thucydides sketches the rise and progress of the Athenian Empire in so clear a style and in such simple language, that the author doubts whether any easier or more instructive passages can be selected for the use of the pupil who is commencing Greek.

3. The Seven Kings of Rome.

A First Latin Reading Book.

Second Edit. Fcap. 8vo. cloth. 3s.

This work is intended to supply the pupil with an easy Construing-book, which may, at the same time, be made the vehicle for instructing him in the rules of grammar and principles of composition. Here Livy tells his own pleasant stories in his own pleasant words. Let Livy be the master to teach a boy Latin, not some English collector of sentences, and he will not be found a dull one.

4. Vocabulary and Exercises on "The Seven Kings of Rome."

Fcp. 8vo. cloth. 2s. 6d.

* * * The Vocabulary and Exercises may also be had bound up with "The Seven Kings of Rome." 5s. cloth.

By EDWARD THRING, M.A.
Head Master of Uppingham School.

**Elements of Grammar
Taught in English.**

With Questions.

Third Edition. 136 pp. (1860).
Demy 18mo. 2s.

**2. The Child's English
Grammar.**

New Edition. 86 pp. (1859). Demy
18mo. 1s.

The Author's effort in these two books has been to point out the broad, beaten, every-day path, carefully avoiding digressions into the byways and eccentricities of language. This Work took its rise from questionings in National Schools, and the whole of the first part is merely the writing out in order the answers to questions which have been used already with success. The study of Grammar in English has been much neglected, nay by some put on one side as an impossibility. There was perhaps much ground for this opinion, in the medley of arbitrary rules thrown before the student, which applied indeed to a certain number of instances, but would not work at all in many others, as must always be the case when principles are not put forward in a language full of ambiguities. The present work does not, therefore, pretend to be a compendium of idioms, or a philological treatise, but a Grammar. Or in other words, its intention is to teach the learner how to speak and write correctly, and to understand and explain the speech and writings of others. Its success, not only in National Schools, from practical work in which it took its rise, but also in classical schools, is full of encouragement.

3. School Songs.

A COLLECTION OF SONGS FOR
SCHOOLS.
WITH THE MUSIC ARRANGED FOR
FOUR VOICES.

*Edited by Rev. E. THRING and
H. RICCIUS.*

Music Size. 7s. 6d.

By EDWARD THRING, M.A.

**4. A First Latin Con-
struing Book.**

104 pp. (1855). Fcap. 8vo. 2s. 6d.

This Construing Book is drawn up on the same sort of graduated scale as the Author's *English Grammar*. Passages out of the best Latin poets are gradually built up into their perfect shape. The few words inserted, or inserted as the passages go on, are printed in Italics. It is hoped by this plan that the learner, whilst acquiring the rudiments of language, may store his mind with good poetry and a good vocabulary.

By C. J. VAUGHAN, D.D.
Head Master of Harrow School.

**St. Paul's Epistle to
the Romans.**

The Greek Text with English Notes.

Second Edition. Crown 8vo. cloth.
(1861). 5s.

By dedicating this work to his elder Pupils at Harrow, the Author hopes that he sufficiently indicates what is and what is not to be looked for in it. He desires to record his impression, derived from the experience of many years, that the Epistles of the New Testament, no less than the Gospels, are capable of furnishing useful and solid instruction to the highest classes of our Public Schools. If they are taught accurately, not controversially; positively, not negatively; authoritatively, yet not dogmatically; taught with close and constant reference to their literal meaning, to the connexion of their parts, to the sequence of their argument, as well as to their moral and spiritual instruction; they will interest, they will inform, they will elevate; they will inspire a reverence for Scripture never to be discarded, they will awaken a desire to drink more deeply of the Word of God, certain hereafter to be gratified and fulfilled.

By C. J. VAUGHAN, D.D.

**Notes for
Lectures on Confirmation.**
With Suitable Prayers.
4th Edition. 70 pp. (1862). Fcp.
8vo. 1s. 6d.

This work, originally prepared for the use of Harrow School, is published in the belief that it may assist the labours of those who are engaged in preparing candidates for Confirmation, and who find it difficult to lay their hand upon any one book of suitable instruction at once sufficiently full to furnish a synopsis of the subject, and sufficiently elastic to give free scope to the individual judgment in the use of it. It will also be found a handbook for those who are being prepared, as presenting in a compact form the very points which a lecturer would wish his hearers to remember.

2.
The Church Catechism Illustrated and Explained. By
ARTHUR RAMSAY, M.A.
18mo. cloth. 2s.

3.
Hand-Book to Butler's Analogy. By C. A. SWAINSON, M.A. 55 pp. (1856). Crown 8vo. 1s. 6d.

4.
History of the Christian Church during the First Three Centuries, and the Reformation in England. By W. SIMPSON, M.A. Fourth Edition. Fcp. 8vo. cloth. 3s. 6d.

5.
Analysis of Paley's Evidences of Christianity. By CHARLES H. CROSSE, M.A. 115 pp. (1855). 18mo. 3s. 6d.

FORTHCOMING BOOKS.

1.
An Elementary Treatise on Natural Philosophy.

By WILLIAM THOMSON, LL.D., F.R.S., late Fellow of St. Peter's Coll. Cambridge, Professor of Natural Philosophy in the University of Glasgow, and PETER GUTHRIE TAIT, M.A., late Fellow of St. Peter's College, Cambridge, Professor of Natural Philosophy in the University of Edinburgh. With numerous Illustrations
[In the Press]

2.
The Narrative of Odysseus.

Homer's *Odyssey*, Books IX—XII. The Greek Text with English Notes. For Schools and Colleges. By JOHN E. B. MAYOR, M.A., Fellow and Principal Classical Lecturer of St. John's College, Cambridge.
[Nearly Ready.]

3.
First Book of Algebra. For Schools. By J. C. W. ELLIS, M.A., and P. M. CLARKE, M.A., Sidney Sussex College, Cambridge.
[Preparing]

4.
Aristotelis de Rhetorica.

With Notes and Introduction. By E. M. COPE, M.A., Fellow and Assistant Tutor of Trinity College, Cambridge.

5.
The New Testament in the Original Greek. Text revised by B. F. WESTCOTT, M.A., and F. J. HORT, M.A., formerly Fellows of Trinity College.

CAMBRIDGE MANUALS

FOR THEOLOGICAL STUDENTS.

- 1. History of the Christian Church during the Middle Ages.** By ARCHDEACON HARDWICK. Second Edition. 482 pp. (1861). With Maps. Crown 8vo. cloth. 10s. 6d.

This Volume claims to be regarded as an integral and independent treatise on the Mediæval Church. The History commences with the time of Gregory the Great, to the year 1520,—the year when Luther, having been extruded from those Churches that adhered to the Communion of the Pope, established a provisional form of government and opened a fresh era in the history of Europe.

- 2. History of the Christian Church during the Reformation.** By ARCHDN. HARDWICK. 459 pp. (1856). Crown 8vo. cloth. 10s. 6d.

This Work forms a Sequel to the Author's Book on The Middle Ages. The Author's wish has been to give the reader a trustworthy version of those stirring incidents which mark the Reformation period.

- 3. History of the Book of Common Prayer.** With a Rationale of its Offices. By FRANCIS PROCTER, M.A. Fifth Edition. 464 pp. (1860). Crown 8vo. cloth. 10s. 6d.

In the course of the last twenty years the whole question of liturgical knowledge

has been reopened with great learning and accurate research, and it is mainly with the view of epitomizing their extensive publications, and correcting by their help the errors and misconceptions which had obtained currency, that the present volume has been put together.

- 4. History of the Canon of the New Testament during the First Four Centuries.** By BROOKE FOSS WESTCOTT, M.A. 594 pp. (1855). Crown 8vo. cloth. 12s. 6d.


The Author has endeavoured to connect the history of the New Testament Canon with the growth and consolidation of the Church, and to point out the relation existing between the amount of evidence for the authenticity of its component parts and the whole mass of Christian literature. Such a method of inquiry will convey both the truest notion of the connexion of the written Word with the living Body of Christ, and the surest conviction of its divine authority.

- 5. Introduction to the Study of the GOSPELS.** By BROOKE FOSS WESTCOTT, M.A. 458 pp. (1860). Crown 8vo. cloth. 10s. 6d.

This book is intended to be an Introduction to the *Study* of the Gospels. In a subject which involves so vast a literature much must have been overlooked; but the Author has made it a point at least to study the researches of the great writers, and consciously to neglect none.

This Series of THEOLOGICAL MANUALS has been published with the aim of supplying Books concise, comprehensive, and accurate; convenient for the Student, and yet interesting to the general reader.

Uniformly printed in 18mo.
with Vignette Titles by
T. Woolner, W. Holman
Hunt, &c.



Handsomely bound in ex-
tra cloth, 4s. 6d. Morocco
plain, 7s. 6d. Morocco ex-
tra, 10s. 6d. each Volume.

1. THE GOLDEN TREASURY
OF THE BEST SONGS AND LYRICAL POEMS IN THE ENGLISH
LANGUAGE.

Selected and arranged, with Notes, by F. T. PALGRAVE.

TWELFTH THOUSAND, with a Vignette by T. WOOLNER.

"There is no book in the English language which will make a more delightful
companion than this . . . which must not only be read, but possessed, in order to
be adequately valued."—SPECTATOR.

2. THE CHILDREN'S GARLAND.
FROM THE BEST POETS.

Selected and Arranged by COVENTRY PATMORE.

FOURTH THOUSAND, with Vignette by T. WOOLNER.

"Mr. Patmore deserves our gratitude for having searched through the wide field
of English poetry for these flowers which youth and age can equally enjoy, and
woven them into 'The Children's Garland.'"—LONDON REVIEW.

3. THE PILGRIM'S PROGRESS.
By JOHN BUNYAN.

With Vignette by W. HOLMAN HUNT.

Large paper copies, crown 8vo. cloth, 7s. 6d., half morocco, 10s. 6d.

"A prettier and better edition and one more exactly suited for use as an elegant
and inexpensive Gift Book is not to be found."—EXAMINER.

4. THE BOOK OF PRAISE.
FROM THE BEST ENGLISH HYMN WRITERS.

Selected and arranged by ROUNDELL PALMER.

EIGHTH THOUSAND, with Vignette by T. WOOLNER.

"Comprehending nearly all that is excellent in the hymnology of the language.
. . . . In the details of editorial labours the most exquisite finish is manifest."—THE
FREEMAN.

**5. BACON'S ESSAYS AND COLOURS OF GOOD
AND EVIL.**

With Notes and Glossarial Index, by W. ALDIS WRIGHT, M.A.,
Trinity College, Cambridge.

And a Vignette of Woolner's Statue of Lord Bacon.

Large Paper Copies, Crown 8vo. cloth, 7s. 6d., half-morocco, 10s. 6d.

"Edited in a manner worthy of their merit and fame, as an English classic ought
to be edited."—DAILY NEWS.

Jonathan Palmer, Printer, Cambridge.

