

Historic, archived document

Do not assume content reflects current scientific knowledge, policies, or practices.



GLUTAMIC ACID REVIEW AND BIBLIOGRAPHY

H. S. Olcott and G. H. Brother

Western Regional Research Laboratory*, Albany, California


REVIEW

Production and use as a condiment of the monosodium salt of glutamic acid originated, according to Han, in Japan about 1908. Previously, protein hydrolysates were widely used throughout the Orient as flavoring materials for predominantly vegetarian diets. The desirable meat-like flavor of these hydrolysates was found to be due largely to the glutamic acid, the monosodium salt of which is so effective that the taste threshold is 1:3000, as compared with 1:200 for cane sugar and 1:400 for table salt. The development of what has been a laboratory curiosity into an article of commerce was therefore quite justified.

Little interest was shown in this development by American industry until the early 1930's, although it had been mentioned in the patent literature as early as 1912. The industrial capacity to produce monosodium glutamate was not sufficient to meet the increased requirements for canned and dehydrated foods in World War II; consequently, interest in the product and in its production have been widespread. It is obtainable both from Steffen's waste, a by-product of beet sugar manufacture, and from wheat gluten, by-product of starch and alcohol production from wheat. Wheat, frequently grown in excess of food requirements in prewar years, has presented a persistent problem of economical utilization.

Chemical structure

Glutamic acid is alpha-amino glutaric acid:



The starred carbon atom is assymmetric; hence glutamic acid occurs in two forms, one levo-rotatory, the other dextro-rotatory. However, because the dextro-rotatory form has been shown to belong configurationally to a group of compounds which contain the naturally occurring amino acids, most of which are levo-rotatory, it is correctly designated as l(+) glutamic acid, the "l" indicating the family compounds to which it belongs and the (+) the direction of rotation. The optical isomer is called d(-) glutamic acid. In many patents and papers the dextro compound is written simply d-glutamic acid; l(+)glutamic acid is meant.

A mixture of equal parts of l(+) and d(-)glutamic acid is optically inactive and is called racemic or d \pm -glutamic acid.

Occurrence

Glutamic acid occurs in nature mostly as the l(+) isomer. A major constituent of most proteins, it occurs to some degree in practically all (See Table I). The monoamide, l-glutamine, is found in plant and animal tissues. Folic acid, a vitamin, contains glutamic acid. A major commercial source is beet extract from which the sucrose has been removed (Steffen's waste). A tri-peptide, glutathione, containing glutamic acid, is also a constituent of living tissue.

* Bureau of Agricultural and Industrial Chemistry, Agricultural Research Administration, U. S. Department of Agriculture.

The occurrence of d(-)glutamic acid is limited. The only proven natural source is a polypeptide of d(-)glutamic acid elaborated by a specific strain of B. subtilis. Other bacterial end-products are known to contain the unnatural forms of other amino acids and it may be expected that the d(-)glutamic acid will be isolated from some of these.

Kogl and his collaborators reported that cancer tissues contain appreciable amounts of unnatural amino acids, particularly d(-) glutamic acid. The results of considerable research investigation that was stimulated by the theory have not confirmed this claim.

Synthesis of glutamic acid

A number of investigators have succeeded in synthesizing glutamic acid from simpler organic substances. The more recent syntheses are based on malonic ester and acetoacetic ester methods. The products are optically inactive and hence contain only 50% of the desired l(+) isomer. At the present time it appears that l(+)glutamic acid can be obtained more economically from natural sources than by synthesis in the laboratory.

Resolution

The separation of optical isomers of amino acids is usually a lengthy and laborious task. It can be accomplished by careful fractionation of the salts of amino acids with optically active alkaloids. Sugasawa resolved dl-glutamic acid by converting it to dl-pyrrolidone carboxylic acid, preparing and then fractionating the quinine salts, and reconverting the isomers to glutamic acid. d(-) Glutamic acid has been prepared from dl-glutamic acid by enzymatic resolution.

Racemization

Racemization is the term applied to the change of an optically active substance to its inactive form. This transformation is to be avoided in the manufacture of l(+)glutamic acid inasmuch as d(-)glutamic acid does not possess the unique flavor of its isomer, and the net result is a loss of flavor. Racemization occurs much more readily in alkaline than in acid solutions. Boiling 20% hydrochloric acid will racemize approximately 2-4% of l(+)glutamic acid in 24 hours. With the same time and temperature of heating, 30% is racemized by 20% sodium hydroxide.

Preparation from wheat gluten

In one method of preparing glutamic acid from wheat gluten, the protein is completely hydrolyzed by boiling or autoclaving with strong hydrochloric acid. The humin is then removed by filtration. Decolorization may be effected by heating with carbon. The excess hydrochloric acid is removed by evaporation at reduced pressure. Glutamic acid hydrochloride crystallizes from the concentrated, strongly acid hydrolysate.

The crystalline hydrochloride is dissolved in water, treated with charcoal to remove residual color, filtered, and carefully adjusted to pH 3.3, the isoelectric point of glutamic acid. Glutamic acid crystallizes.

In order to obtain the monosodium salt, the glutamic acid is brought to pH 7.0 in aqueous solution with sodium hydroxide. Upon concentration, the monosodium salt crystallizes. This salt is the article of commerce. Its taste and purity depend upon the care which has been taken in the previous steps. There is only

meager information about yields. Commercial recoveries possibly amount to approximately 20% by weight of the gluten used, or from $\frac{1}{2}$ to $\frac{2}{3}$ of the glutamic acid known to be present.

Preparation from other proteins

Corn gluten and soy bean protein have been advocated as sources of glutamic acid. With these proteins, the percentage of recovery appears to be even lower than with wheat gluten.

Preparation from Steffen's waste

Steffen's waste is the spent liquor remaining after the extraction of sugar from sugar beet molasses by Steffen's process. Some Steffen's waste contains amounts of glutamic acid which make practical its isolation from this source. Other lots may contain little or no glutamic acid. The reasons for the differences are obscure although it is known that beets from certain geographical localities are consistently high, and from others low, in glutamic acid.

Careful alkaline hydrolysis is used with Steffen's waste to obtain the glutamic acid from its precursor, possibly pyrrolidone carboxylic acid. Several patented procedures are described for the isolation steps.

Condiment

The taste of most samples of monosodium glutamate is meat-like. When a few crystals are placed in the mouth, there is no immediate sensation. Then the taste of good beefsteak develops slowly, mostly in the rear of the mouth, and persists for some time. It is claimed that monosodium glutamate, like salt, enhances other flavors, in addition to possessing one of its own. A very highly purified sample of monosodium glutamate has a more delicate flavor than those usually marketed, suggesting that the more gross, meat-like flavor may be an impurity.

Nutrition

Glutamic acid is usually classed as one of the non-essential amino acids. However, it has been maintained that its inclusion in the diet of chickens is necessary for best growth.

Approximate glutamic acid content of natural materials.

<u>Protein</u>	<u>Glutamic Acid Content*</u>
	per cent
Gliadin	46
Gluten	36
Zein	26
Casein	22
Insulin	20
Beta-lactoglobulin	19
Edestin	19
Bovin serum albumin	17
Egg albumin	16
Fibrin	13
Gelatin	12

Protein

Steffen's waste
Silk fibroin

Glutamic Acid Content*

per cent

0-15**
2

* Dry basis

**The glutamic acid content of Steffen's waste may vary widely depending upon geographical location of the beet-growing area and other as yet unidentified factors.

BIBLIOGRAPHYGeneral

Abderhalden, E., Biochemisches Handlexicon, vol. XII, p. 486-510, J. Springer, Berlin, 1930. (lithoprinted by Edwards Brothers, Ind., Ann Arbor, 1944).

Beilsteins Handbuch der Organischen Chemie, 4, 488-494 (1922); H4, 902-911 (1942) Springer, Berlin ; reproduction by Edwards Bros., Ann Arbor, Mich., 1943.

Thorpe's Dictionary of Applied Chemistry, vol. VI, p. 31-28, Longmans, Green and Co., New York, 4th Ed., 1943.

Determination

Arhimo, A. A. & Laine, T., Determination of glutamic acid, Suomen Kemistilehti, 12B, 18 (1939); Chem. Abs. 33, 6198.

Baumgarten, W., Mather, A. N. & Stone, L., Glutamic acid content of feed materials, Cereal Chem., 22, 514-21, (1945); Chem. Abs. 40, 968.

Cannan, R. K., The estimation of the dicarboxylic amino acids in protein hydrolysates, J. Biol. Chem., 152, 401-10, (1944); Chem. Abs. 38, 2679.

Cohen, P. P., Microdetermination of glutamic acid, Biochem. J., 33, 551-8 (1939), Chem. Abs. 38, 7835.

Darling, S., Chromatographic method for the determination of glutamic and aspartic acid in mixtures of different amino acids, Acta. Physiol. Scand 10, 91-6 (1945); Chem. Abs. 40, 2182.

Dunn, M. S., Camien, M. N., Rockland, I. B., Shankman, S. & Goldberg, S. C., Amino acids, peptides and proteins. XVII. Determination of glutamic acid in protein hydrolysates by a microbiological method, J. Biol. Chem., 155, 591-603 (1944), Chem. Abs. 39, 950.

Foreman, F. W., Rapid Volumetric Methods for the estimation of amino acids, organic acids and organic bases, Biochem. J., 14, 451-73 (1920), Chem. Abs. 14, 3044.

Gale, E. F., Studies on amino acid decarboxylases. V. Use of specific decarboxylase preparation in the estimation of amino acids and in protein analysis, Biochem. J., 39, 46-52 (1945), Chem. Abs. 39, 5261.

Hac, L., Snell, E. E., and Williams, R. J., Microbiological determination of amino acids. II. Assay and utilization of glutamic acid and glutamine by *L. Arabinosus*. J. Biol. Chem. 159, 273-280 (1945)

Jones, D. B. & Moeller, O., Some recent determinations of aspartic and glutamic acids in various proteins, J. Biol. Chem., 79, 429-41 (1928), Chem. Abs. 23, 62.

Kihara, K., Decolorizing method for melanin-containing amino acid solution and its application, J. Agr. Chem. Soc. Japan 16, 206-8; Bull. Agr. Chem. Soc. Japan 16, 42-3 (in English) (1940), Chem. Abs. 34, 5868.

Kuiken, K. A., Norman, W. H. Lynman, C. M. & Hale, F., Microbiological Determination of Amino Acids, Science 98, 266 (1943), Chem. Abs. 37, 6296.

Lewis, J. C. & Olcott, H. S., A Lactobacillus assay method for 1(+)-Glutamic acid, J. Biol. Chem., 157, 265-85. 1945, Chem. Abs. 39, 1657.

Lynman, C. M., Kuiken, K. A., Blotter, L., and Hale, R., Microbiological determination of amino acids. II. Glutamic acid, J. Biol. Chem., 157, 395-405 (1945), Chem. Abs. 39, 1658.

Olcott, H. S., A method for the determination of glutamic acid in proteins, J. Biol. Chem. 153, 71-82 (1944), Chem. Abs. 38, 3685.

Opsahl, J. C. & Arnow, L. E., The optical configuration of glutamic acid isolated from casein hydrolysates by six procedures, J. Amer. Chem. Soc. 64, 2035-9 (1942), Chem. Abs. 36, 6555.

Prescott, B. A., and Wafsch, H., Macro determination of glutamic acid and the application to protein analysis, J. Biol. Chem., 164, 331-343 (1946).

Rittenberg, D. & Foster, G. L., A New procedure for quantitative analysis by istope dilution, with application to the determination of amino acids and fatty acids, J. Biol. Chem. 133, 737-44 (1940), Chem. Abs. 34, 4093.

Shankman, S., Dunn, M. S. & Rubin, L. B., The analysis of eight amino acids by a microbiological method., J. Biol. Chem., 150, 477-8 (1943), Chem. Abs. 38, 394.

Schales, O., Mino, U., and Schales, S. S., Glutamic acid decarboxylase of higher plants, Fed. Proc. 5, 152 (1946).

Shemin, D., Amino acid determinations on crystalline bovine and human serum albumin by the istope-dilution method, J. Biol. Chem., 159, 439-43 (1945), Chem. Abs. 39, 4905.

Shankman, S. Dunn, M. S. & Rubin, L. B., Microbiological analysis of seven amino acids with lactobacillus casein, J. Biol. Chem., 151, 511-14 (1943), Chem. Abs. 38, 1257.

Vickery, H. B., Continuation and extension of work on vegetable proteins, Carnegie Inst. Washington Yearbook 35, 308-14 (1936), Chem. Abs. 31, 3074.

Wieland, T. & Wirth, L., Quantitative separation of aspartic and glutamic acids with the acid aluminum oxide column, Ber. 76B, 823-5 (1943), Chem. Abs. 38, 2060.

Wieland, T., and Paul V., Determination of l-and d-glutamic acid in hydrolysates of Brown-Pearce tumors with N^{15} glutamic acid, Ber. 77B, 34-44 (1944), Chem. Abs. 39, 120.

Woodward, G. E. & Smith, L. D., Attempts to analyze for δ (-)-glutamic acid in tissue hydrolysates, J. Franklin Inst., 231, 89-91 (1941), Chem. Abs. 35, 1823.

Technic of isolation

Braun, G., Process of glutamic acid recovery from solutions, Brit. 508, 841, July 5, 1939, Chem. Abs. 34, 2865.

Braun, G., Improvements in the production of glutamic acid and its salts, Brit. 508, 842, July 5, 1939, Chem. Abs. 34, 2865.

Braun, G., Process of producing glutamic acid, U. S. 2,267,971, Dec. 30, 1941, Chem. Abs. 36, 2568.

Cross, R. J., Electroosmotic Process and Apparatus, U. S. 1,986, 920, Jan. 8, 1935, Chem. Abs. 29, 1333.

Ikeda, K. & Suzuki, S., Method of making a nutritive and flavouring substance, Brit. Pat. 9440, Apr. 21, 1910, Chem. Abs. 5, 836.

Ikeda, K. & Suzuki, S., Process of separating glutamic acid and other products of hydrolysis of albuminous substances from one another by electrolysis, U.S. Pat. 1,015,891, Jan. 30, 1912, Chem. Abs. 6, 717.

Oka, K., A process of purifying sodium glutamate, Brit. 290,657, Sept. 2, 1929 Chem. Abs. 23, 851.

Schildneck, P. R., Preparation of monosodium glutamate, U. S. 2,306,646, Dec. 29, 1942. Chem. Abs. 37, 3107.

Wei, Y. S. & King, C. K., Neutralization of d-glutamic acid hydrochloride and crystallization of monosodium d-glutamate during the manufacture of glutamate condiments, Science (China), 19, 354-66 (1935), Chem. Abs. 29, 4849.

Preparation of glutamic acid
from:

CASEIN

Kemmerer, K. S., Amino acids product and method of manufacture, U. S. 2,180,636, Nov. 21, 1939. Chem. Abs. 34, 1821.

Marshall, A. E., Process for manufacturing and recovering glutamic acid, U.S. 1,973,574, Sept. 11, 1934, Chem. Abs. 28, 6726.

Masuda, R., Royal C. L. & Marshall, A. E., Process for Manufacturing and recovering glutamic acid and its compounds, U. S. 1,947,563, Feb. 20, 1934, Chem. Abs. 28, 2728.

CORN PROTEIN

Bartow, E. & Albrook, R. L., Process of producing glutamic acid and compounds, U. S. 1,992,804, Feb. 26, 1935, Chem. Abs. 29, 2548.

Bishop, J. P., and Tucker, F. L., Process of making mono-sodium glutamate from gluten, U. S. 2,214,115, Sept. 10, 1940, Chem. Abs. 35, 755.

Corn Products Refining Company, Procede de fabrication de chlorhydrate de l' acide glutamique et de ses derives a partir de gluten de mais, Fr. 844,306, July 24, 1939, Chem. Abs. 34, 7300.

Corn Products Refining Company, Improvements in and relating to the production of glutamic acid and its salts, Brit. 522, 365, June 17, 1940, Chem. Abs. 36, 1048.

Corn Products Refining Company, Improvements in and relating to the production of amino acids, Brit. 529, 612, Nov. 25, 1940, Chem. Abs. 36, 1048.

Mark, A. M., Isolation and production of leucine and tyrosine from corn gluten, U. S. 2,178,210, Oct. 31, 1939, Chem. Abs. 34, 1336.

Shildneck, P. R., Preparation of glutamic acid, U. S. 2,347,220, Apr. 25, 1944, Chem. Abs. 39, 89.

SEaweeds

Schindelmeiser, J., Improvements in or relating to the production of glutamic acid and glutamates, Brit. 481,898, Mar. 21, 1938, Chem. Abs. 32, 6264.

STEFFEN'S WASTE

Amino Products Company, Verfahren zur Herstellung von d-glutaminsäure Hydrolysis of Steffen's waste or other protein material with alkali of 3 to 7% NaOH to produce glutamic acid, Ger. 693,145, July 3, 1940, Chem. Abs. 35, 4782.

Bartow, E., Coleman, G. H. & Redman, H. E., Variations in the composition of Steffen's Waste, II, Proc. Iowa Acad. Sci. 44, 111-16 (1937), Chem. Abs. 33, 425.

Bartow, E. U. Wetzstein, H. L., Preparation of tyrosine from Steffen's Waste, Proc. Iowa Acad. Sci., 47, 193-95 (1940), Chem. Abs. 35, 7750.

Ikeda, K. Process of utilizing waste waters from beet-sugar molasses, U. S. 1,721,820, July 23, 1929. Chem. Abs. 23, 4591.

Larrove Construction Company, Improvements in or relating to the manufacture or recovery of substances from residual liquors of the beet sugar industry, Brit. 265,831, Feb. 17, 1927, Chem. Abs. 22, 509.

Larrove-Suzuki Company, Perfectionnements aux procedes pour la fabrication et la recuperation de l'acide glutamique et de ses composes. Fr. 742,750, May 2, 1932, Chem. Abs. 26, 4828.

Larrove-Suzuki Company, Improvements in Processes of manufacturing and recovering glutamic acid and its compounds, Brit. 385,054, Dec. 22, 1932, Chem. Abs. 27, 4248.

Masuda, R., Manufacture of glutamic acid hydrochloride, betaine acid sulphate, etc., U. S. 1,928,840, Oct. 3, 1933., Chem. abs. 27, 5756.

Nees, A. R., and Bennett, A. N., Recovery of nitrogenous products from organic wastes. U. S. 2,375,165, May 1, 1945, Chem. Abs. 39, 3457.

O'Day, D. W. & Bartow, E., Glutamic acid content of Steffen's Waste, Ind. Eng. Chem. 35, 1152-55 (1943), Chem. Abs. 37, 6921.

Phikin, S. S., The production of acidol and glutamic acid from the waste products of the sugar industry, Trudy Beningrad Farm. Inst., 3, 85-98 (1940); Khim. Referat. Zhur., 4, 130 (1941), Chem. Abs. 37, 5884.

Royal, C. L., Manufacture of glutamic acid, U. S. 2,373,342, April 10, 1945, Chem. Abs. 29, 4510.

Tressler, D. K., Betaine hydro-chloride, Can. 268,728, Mar. 1, 1927, Chem. Abs. 21, 2395.

Tressler, D. K., Manufacture of betaine hydrochloride, etc., U.S. 1,634,221, June 28, 1927. Chem. Abs. 21, 2814.

Tressler, D. K., Process of Preparing Betaine Hydrochloride, U. S. 1,634,222, June 28, 1927. Chem. Abs. 21, 2814.

SOYBEANS

Bersworth, F. C., Method of hydrolyzing protein materials, U. S. 2,334,986, Nov. 23, 1943, Chem. Abs. 38, 2770.

Bersworth, F. C., Method of hydrolyzing proteins, U. S. 2,334,987, Nov. 23, 1943, Chem. Abs. 38, 2770.

Cheng, Y-C. & Adolph, W. H., Preparation of d-glutamic acid, J. Chinese Chem. Soc. 2,221-4 (1934), Chem. Abs. 29, 740.

Itikawa, S. & Huzi, S., Treatment of soybean cake, Jap. 129,862, Apr. 21, 1939, Chem. Abs. 35, 2024.

Mashino, M. & Shishido, T., Effect of carbohydrates and other impurities on the crystallization of glutamic acid hydrochloride from decomposition products of soybean protein, J. Soc. Chem. Ind. (Japan) 33, Suppl. binding 421-2 (1930), Chem. Abs. 25, 746.

Okano, K. & Beppu, I., Soybean oil foots. II. Isolation of free amino acids, J. Agr. Chem. Soc. Japan 14, 248-50 (1938), Chem. Abs. 32, 6086.

Rokusho, B., Tanaka, R. & Miyahara, C., Preparation of glutamic acid., J. Agr. Chem. Soc. Japan, 13, 916-53, (1937), Chem. Abs. 32, 1245.

Takayama, Y., Utilization of soy beans, III. J. Soc. Chem. Ind. Japan, 33, Suppl. binding 91-2 (1930), Chem. Abs. 24, 2810.

WHEAT PROTEIN

Barnett, H. M., Amino acids, U. S. 1,990,769, Feb. 12, 1935, Chem. Abs. 29, 2179,

Barnett, H. M., Manufacture of flavoring materials, U. S. 1,992,462, Feb. 26, 1935, Chem. Abs. 29, 2618.

Braun, G., Process of glutamic acid recovery from solutions, Brit. 508,927, July 5, 1939, Chem. Abs. 34, 2865.

Braun, G., Process of glutamic acid production, U. S. 2,176,785, Oct. 17, 1939, Chem. Abs. 34, 1037.

Braun, G., Process of glutamic acid recovery from solutions, U. S. 2,186,162, Jan. 9, 1940, Chem. Abs. 34, 2865.

Corti, A., Process to separate glutamic acid from other amino acids, U.S. 1,255,390, Feb. 5, 1918. Chem. Abs. 12, 909.

Fong, Y. S., Procede pour la fabrication de glutamate de sodium, Fr. 669,815, Nov. 21, 1929. Chem. Abs. 24, 1868.

Hardie, H. D. & Company, Ltd., An improved process for the manufacture of mono-sodium glutamate and like glutamic acid salts, Brit. 347,258, Apr. 23, 1931, Chem. Abs. 26, 2750.

Ikeda, K. & Suzuki, S., Nutritive and flavoring substance and process of making same, U. S. Pat. 1,035,591, Aug. 13, 1912. Chem. Abs. 6, 3299.

Ito, K., Glutamic acid hydrochloride, Jap. Pat. 98,725, Dec. 19, 1932, Chem. Abs. 27, 4817.

King, H., d-Glutamic acid, organic syntheses, Collective Vol. I, 286-289 (1941) John Wiley & Sons, New York.

Komatsu, S., Glutamic acid or its salts, Jap. 93,152, Oct. 8, 1931, (Addn. to 92, 068), Chem. Abs. 27, 1364.

Kumagai, G. S., Method of preparing sodium glutamate, U. S. 2,050,491, Aug. 11, 1936, Chem. Abs. 30, 6763.

Masuda, R., Process for recovering glutamic acid, U. S. 1,940,428, Dec. 19, 1933 Chem. Abs. 28, 1364.

Olcott, H. S., Outlook for wheat gluten as an industrial material, Food Ind. 16, 541, 576-78, (1944).

Olcott, H. S., & Elish, M. J., Wheat gluten as an industrial protein, Trans. Am. Assoc. Cereal Chemists 2, 20-25 (1944), Chem. Abs. 38, 6045.

Sen, F. Y., Manufacture of glutamic acid compounds, U. S. 1,890,590, Dec. 13, 1932, Chem. Abs. 27, 1641.

Standard Brands, Inc., Improvements in process of glutamic acid production, Brit. 501,542, Feb. 28, 1939, Chem. Abs. 33, 6348.

Standard Brands, Inc., Improvements in process of glutamic acid production, Brit. 501,635, Feb. 28, 1939, Chem. Abs. 33, 6348.

Town, B. W., The separation of amino acids by means of their copper salts. III, The hydrolysis of gliadin. An investigation of the dicarboxylate fraction, including the isolation of α -glutamic acid as a hydrolysis product, Biochem. J. 35, 417-32 (1941), Chem. Abs. 35, 6930.

Tseng, C. L. & Chu, E.J.H., Optimum conditions for preparing d-glutamic acid hydrochloride from wheat gluten. I. Effect on concentration and proportion of hydrochloride acid used in the hydrolysis, Trans. Sci. Soc. China, 7, 217-24 (1932), Chem. Abs. 26, 5543.

Tseng, C. L. & Chu, E.J.H., Optimum conditions for preparing d-glutamic acid hydrochloride from wheat gluten. II. Effect of purity of acid, time of reaction, climatic conditions and other experimental circumstances, Science Quart. Natl. Univ. Peking 3, 53-68 (1932), Chem. Abs. 27, 1867.

Tseng, C. L. & Chu, E.J.H., The decolorization of the solution in the preparation of d-glutamic acid hydrochloride by hydrolysis of wheat gluten, Science Quart. Natl. Univ. Peking 3, 69-74 (1932), Chem. Abs. 27, 1867.

Tseng, C. L. & Chu, E.J.H., Use of some American brands of decolorizing carbon in the preparation of d-glutamic acid hydrochloride by hydrolysis of wheat gluten, J. Chinese Chem. Soc. 1, 35-6, (1933), Chem. Abs. 27, 3197.

Tseng, C. L. & Hu, M.J., Gluten hydrolysis and preparation of α -glutamic acid hydrochloride, J. Chinese Chem. Soc. 3, 154-72 (1935), Chem. Abs. 29, 6212.

Yanagigawa, T. & Nishida, Y., Studies on proteins. IV. On the taste-producing amino acids, Repts. Imp. Ind. Research Inst. Osaka (Japan), 16, No. 10, 1-26 (1936), Chem. Abs. 30, 3004.

Sugiura, W., Glutamic acid, Jap. 100,805, Apr. 28, 1933, Chem. Abs. 28, 4747.

Waters, V., Preparation of glutamic acid hydrochloride, U. S. Pat. 2,380,890, July 31, 1945, Chem. Abs. 40, 91.

Synthesis

Hamlin, K. E., Jr. & Hartung, W. H., The Synthesis of alpha amino acids from substituted acetoacetic esters, J. Biol. Chem. 145, 349-57 (1942), Chem. Abs. 37, 1387.

Marvel, C. S. & Stoddard, M. P., Convenient synthesis of dl-glutamic acid, J. Org. Chem. 3, 198-203 (1938), Chem. Abs. 33, 965.

Redemann, G. E. & Dunn, M. S., A general method for the synthesis of alpha amino acids with ethyl benzamidomalonate, J. Biol. Chem. 130, 341-8 (1939), Chem. Abs. 33, 9285.

Snyder, H. R., Shekleton, J. F., and Lewis, C. D., Synthetic amino acids and their derivatives. VIII. Synthesis from acetamidomalonate ester, J. Amer. Chem. Soc. 67, 310-12 (1945), Chem. Abs. 39, 1871.

Resolution

Chargaff, E., The chemistry of bacteria. IX. The configuration of glutamic and aspartic acids from pathogenic bacteria (*phytomonas tumefaciens* and *corynebacterium diphtheriae*), J. Biol. Chem. 130, 29-33 (1939), Chem. Abs. 33, 8672.

Fruton, J. S., Irving, G. W., Jr. & Bergmann, M., Preparation of d(-)-glutamic acid from dl-glutamic acid by enzymatic resolution, J. Biol. Chem., 133, 703-5 (1940), Chem. Abs. 34, 4382.

Sugasawa, S., Synthesis of glutamic acid. III. Optical separation of dl-glutamic acid, J. Pharm. Soc. Japan, No. 537, 934-41 (1926), Chem. Abs. 21, 2664.

Condiment

Esselen, G. J., Food Product, U. S. 2,270,582, Jan. 20, 1942, Chem. Abs. 36, 3284.

Fujii, M., Improvements relating to the preparation of flavouring, nourishing, and similar substances from proteins, Brit. 279,985, Nov. 10, 1927, Chem. Abs. 22, 3002.

Han, J. E. S., Monosodium glutamate as a chemical condiment, Ind. Eng. Chem., 21, 984-7 (1929), Chem. Abs. 23, 5248.

Tseng, C. L. & Chu, E. J-H., Tastes of some derivatives of d-glutamic acid, Acad. Sinica Mem. Natl. Research Institute Chem., No. 5, 1-18 (1931), Chem. Abs. 26, 484.

Tseng, C. L. & Chu, E. J-H., Tastes of glutamic acid and related compounds. II. Tastes of some derivatives of dl-glutamic acid, J. Chinese Chem. Soc. 1, 188-98, (1933), Chem. Abs. 28, 1719.

Wang, S. M. & Tseng, C. L., Glutamic acid and sodium glutamate as condiments, Science (China) 17, 1018-48 (1933), Chem. Abs. 27, 5837.

Derivatives

Abderhalden, E. & Bohm, O., Formation of alpha aminobutyric acid on warming l-glutamic acid with sodium hydroxide, Z. Physiol. Chem. 266, 41-2 (1940), Chem. Abs. 35, 2475.

Anslow, W. K. & King, H., Neutral salt addition compounds of alkaline earth glutamates and aspartates, Biochem. J. 21, 1168-78 (1927), Chem. Abs. 22, 196.

Aubel, E. & Reich, W. S., Synthetic preparation of some phosphoryl derivatives of some amino acids, Compt. rend., 195, 183-5 (1932), Chem. Abs. 26, 5548.

Bovarnick, M. R., Isolation of the nicotinamide formed from Asparagine and glutamic acid, J. Biol. Chem. 153, 1-3 (1944), Chem. Abs. 38, 3673.

Braun, G., Process of obtainment of the lactam of glutamic acid, U. S. 2,112,329, Mar. 29, 1938, Chem. Abs. 33, 5870.

Braun, G., Improvements in process for production of the lactam of glutamic acid, Brit. 501,543, Feb. 28, 1939, Chem. Abs. 33, 5870.

Braun, G., Improvements in process for treatment of solutions containing the lactam of glutamic acid, Brit. 501-563, Feb. 28, 1939, Chem. Abs. 33, 5870.

Crebert, T., Dyeing cellulose fibers with acid wool dyes in the presence of synthetic resins, Zellwolle u. Deut. Kunstseiden-Ztg., 5, 146-58; 182-7; 227-31 (1939), Chem. Abs. 34, 1177.

Frieden, E. H., Dunn, M. S. & Coryell, C. D. Quantitative investigations of amino acids and peptides. XI. Equilibrium between amino acids and formaldehyde: Glutamic acid, J. Phys. Chem. 47, 20-4 (1943), Chem. Abs. 37, 2754-5.

Gavrilov, N. I., Elagyna, N. V., Dudykina, N. V. & Kornilov, I. V., The effect of heating certain amino acids in neutral and acid solutions in an autoclave, Bull. Soc. Chem., Series 5, 5, 442-54 (1938), Chem. Abs. 32, 4946.

Harrington, C.R. & Moggridge, R. C. G., Experiments in the glutamic acid series, J. Chem. Soc. (1940) 706-12, Chem. Abs. 34, 6615-6.

Harris, L. J., The combination of proteins, amino acids, etc. with acids and alkalis. Part II. Titration curves of amino-acids in presence of formaldehyde, Proc. Roy. Soc. London, B104, 412-39 (1929), Chem. Abs. 23, 4124.

Lang, K., Solubility of nickel salts of amino acids, Biochem. Z. 301, 368-70 (1939) Chem. Abs. 33, 9091.

Lichtenstein, N., Production of glutamine by amination of pyrrolidone-carboxylic acid, Enzymologia 7, 383 (1939), Chem. Abs. 34, 6229.

Lichtenstein, N., Preparation of dl-glutamine, Enzymologia 9, 185-6 (1941), Chem. Abs. 36, 756.

Lichtenstein, N. & Gertner, S., Preparation of gamma-alkylamides of glutamic acid, J. Amer. Chem. Soc. 64, 1021-22 (1942), Chem. Abs. 36, 4097.

Lifschitz, I. & Schauteden, F. L. M. Metallic complex compounds of optically active alpha amino acids, Rec. Trav. Chim. 58, 411-22 (1939), Chem. Abs. 33, 6248.

Sugasawa, S., Synthesis of glutamic acid. IV., New derivatives of glutamic acid, J. Pharm. Soc. Japan, No. 543; 359-69 (1927), Chem. Abs. 21, 3044.

Tadokoro, T. & Saito, T., Biochemical studies of reductinic acid, I. J. Chem. Soc. Japan, 62, 615-17, (1941), Chem. Abs. 36, 7043.

Toennies, G. & Homiller, R. P., The oxidation of amino acids by hydrogen peroxide in formic acid, J. Amer. Chem. Soc., 64, 3054-6 (1942), Chem. Abs. 37, 1136.

Voss, W. & Guttman, R., Preparation of alkali salts of amino acids, Ber. 63B 1726-31 (1930), Chem. Abs. 24, 5022.

Wilson, H. & Cannan, R. K., The glutamic acid-pyrrolidonecarboxylic acid system, J. Biol. Chem. 119, 309-31 (1937), Chem. Abs. 31, 6089.

Woodward, G. E. & Reinhart, F. E., The effect of pH on the formation of pyrrolidinecarboxylic acid and glutamic acid during enzymatic hydrolysis of glutathione by rat kidney extract, J. Biol. Chem., 145, 471-80 (1942), Chem. Abs. 37, 2025.

Physical properties

Arnow, L. E. & Opsahl, J. C., Partial racemization of glutamic acid in boiling hydrochloric acid solutions, J. Biol. Chem. 133, 765-6 (1940), Chem. Abs. 34, 4727.

Arnow, L. E. & Opsahl, J. C., Racemization of glutamic acid with heat, J. Biol. Chem. 134, 649-51 (1940) Chem. Abs. 34, 5826.

Arnow, L. E. & Opsahl, J. C. Racemization of glutamic acid with alkalies, Science 92, 214-15 (1941), Chem. Abs. 35, 2862.

Cahill, W. M. & Burton, I. F., Racemization of amino acids and dipeptides on acetylation with ketene, J. Biol. Chem., 132, 161-9 (1940), Chem. Abs. 34, 1617.

Dalton, J. B. & Schmidt, C. L. A., The solubilities of certain amino acids in water, the densities of their solutions at twenty-five degrees, and the calculated heats of solution and partial molal volumes, J. Biol. Chem. 103, 549-78 (1933), Chem. Abs. 28, 1913.

Duclaux, J. & Durand-Gasselin, A., The perchlorates of the lyotropic series. II, J. Chim. Phys. 35, 189-92 (1938), Chem. Abs. 33, 22-23.

Dunn, M. S. & Brophy, T. W., Decomposition points of the amino acids, J. Biol. Chem. 99, 221-29 (1932), Chem. Abs. 27, 1371.

Dunn, M. S. & Stoddard, M. P., Crystalline anhydrous and monohydrated dl-glutamic acid, J. Biol. Chem. 121, 521-29 (1937), Chem. Abs. 32, 918.

Dunn, M. S. & Ross, F. J., Quantitative investigations of amino acids and peptides. IV. The solubilities of the amino acids in water-ethyl alcohol mixtures, J. Biol. Chem., 125, 309-32 (1938), Chem. Abs. 32, 6882.

Froentjes, W., Anomalous rotation dispersion of pyroglutamic acid and its ethyl ester, Rec. Trav. Chim., 62, 97-115, (1943); Chem. Abs. 38, 2255.

Huffman, H. V., Ellis, E. L. & Barsook, H., Thermal data. XI. The heat capacities and entropies of guanidine carbonate, glutamic acid hydrochloride, ornithine dihydrochloride, d-lactic acid and l-lactic acid, J. Amer. Chem. Soc. 62, 297-99 (1940) Chem. Abs. 34, 1904.

Jeannerat, J., Racemization of l(+)-glutamic acid, Bull. soc. Chim., 9, 637-8 (1942), Chem. Abs. 38, 719.

Johnson, J. M., The racemization of glutamic acid, J. Biol. Chem., 134, 459 (1940) Chem. Abs. 34, 5827.

Okinaka, C. & Sexagint, Y., Proteins, V. Glutamic acid, Osaka Chem. Inst. Dept. Science Kyoto Imp. Univ. (1927) 27-59, Chem. Abs. 22, 946.

Patterson, J. W. & Brode, W. R., A rotary-dispersion study of alpha amino acids, Arch Biochem. 2, 24757 (1943), Chem. Abs. 37, 5302.

Pertzoff, V. A., The solubility of glutamic acid in water and certain organic solvents, J. Biol. Chem. 100, 97-104 (1933), Chem. Abs. 27, 2365.

Ratner, S., Rittenberg, D. & Schoenhimer, R., The stability of hydrogen-carbon linkages in glutamic acid, J. Biol. Chem. 135, 357-58 (1940), Chem. Abs. 34, 6575.

Ryabinovskaya, A. M., Influence of sodium fluoride, sodium maleinate, glutamic acid and glyceraldehyde on muscular activity, Compt. rend. acad. sci. USSR, 23, 958-61 (1939), Chem. Abs. 34, 1078.

Shih, T. K., A study on the solubility of d-glutamic acid hydrochloride, Ind. Research (China) 4, 542-7, (1935), Chem. Abs. 30, 2826.

Takayama, Y., Utilization of soy beans. III., J. Soc. Chem. Ind. Japan, 33, Suppl. binding 91-2 (1930), Chem. Abs. 24, 2810.

Tseng, C. L., & Chu, E. J.-H., Constitution of monosodium d-glutamate, Science Quart. (Peking) 3, 1-5 (1932), Chem. Abs. 27, 708.

Vellinger, E., Rotary power of some amino acids as a function of the acidity (pH), Compt. rend., 194, 718-20 (1932), Chem. Abs. 26, 2910.

Dehydrogenase

Adler, E., Hellström, V., Günther, G., & Euler, H., Components of dehydrogenase systems. XXII. Enzymic synthesis and decomposition of glutamic acid. 3. In *Bacterium Coli*, Z. Physiol. Chem., 255, 14-26 (1938), Chem. Abs. 32, 9107.

Adler, E. Günther, G. & Everett, J. E., Components of dehydrogenase systems. XXII. Enzymic synthesis and decomposition of glutamic acid. 4. In yeast, Z. Physiol. Chem. 255, 27-35 (1938), Chem. Abs. 32, 9108.

Agren, G., Transamination with peptide substrates in cattle diaphragm muscle, Acta Physiol. Scand. 1, 233-45 (1940), Chem. Abs. 35, 4047.

Boulanger, P., Identification of d(-)glutamic acid by means of d-amino acid dehydrogenase, Compt. rend. Soc. Biol. 137, 521-2 (1943); Chem. Abs. 39, 2774.

Chepinoga, O. P., Dehydrogenase of muscle tissue during training and fatigue. III. Capacity of minced trained and fatigued muscles to oxidize malic, glutamic and citric acids, Biochem. J. (Ukraine) 15 45-54 (1940) (In Russian), (55-56 in English), Chem. Abs. 35, 172.

Chepinoga, O. P., Oxidation mechanism of malic, citric and glutamic acids in minced rabbit muscle. Biochem. J. (Ukraine) 15, 57-64 (1940), Chem. Abs. 35, 172.

Cohen, P. P., Transamination in pigeon breast muscle, Biochem. J. 33, 1478-87 (1939), Chem. Abs. 34, 2001.

Damodaran, M. & Nair, K. R., Glutamic acid dehydrogenase from germinating seeds, Biochem. J. 32, 1064-74 (1938), Chem. Abs. 32, 8484.

Dewan, J. G., The l(+) glutamic dehydrogenase of animal tissues, Biochem. J. 32, 1378-85 (1938), Chem. Abs. 32, 9120.

Dewan, J. G., Coenzyme-linked reactions involving l(+)glutamic acid dehydrogenase, Biochem. J. 33, 549-50 (1939), Chem. Abs. 33, 7831.

Euler, H. V., & Adler, E., Gunther, G. & Elliot, L., Isocitric acid dehydrogenase and glutamic acid synthesis in higher plants and in yeast, Enzymologia 6, 337-41 (1939), Chem. Abs. 34, 458.

Euler, H. V., & Adler, E., Stearic specificity of glutamic acid dehydrogenase, Enzymologia, 8, 21-4 (1939), Chem. Abs. 34, 459.

Felix, K., and Naká, S. The action of liver and kidney slices on l(+)Aspartic Acid, Z. Physiol. Chem. 264, 123-34 (1940), Chem. Abs. 34, 5101.

Franke, W., The biologic decomposition and synthesis of amino acids, Angew. Chem. 52, 695-8; 703-8 (1939), Chem. Abs. 34, 1337.

Grodzenskii, D. E., The formation and breakdown of amino acids by intra-molecular transfer of amino groups. IX. Glycolysis and transamination, Bull. Biol. Med. Exptl. (USSR) 9, 116-18 (1940), Chem. Abs. 35, 2535-36.

Klein, J. R., Oxidation of l(-)-aspartic and l (+)-glutamic acids by *Hemophilus parainfluenzar*. Preparation of pyridine nucleotides from Baker's yeast by the method of Warburg and Christian, J. Biol. Chem. 134, 43-57 (1940), Chem. Abs. 34, 6320.

Konikova, A. A., Kritsman, M. G. & Teis, R. V., Study of the mechanism of transamination by means of deuterium. XVII. Formation and decomposition of amino acids by intermolecular transfer of the amino group, Biokhimiya, 7, 86-92 (English summary, 92), (1942), Chem. Abs. 37, 4411.

Krebs, H. A., & Cohen, P. P., Glutamic acid as a hydrogen carrier in animal tissues. Nature 144, 513-14 (1939), Chem. Abs. 33, 9415.

Kritsman, M. G., Formation and breakdown of amino acids by intermolecular transfer of amino groups. V. The enzyme transferring the amino group of glutamic acid, Biokhimiya 3, 603,15 (1938), Chem. Abs. 33, 2916.

Roloff, M., Ratner, S. & Schoenheimer, R., The biological conversion of Ornithine into proline and glutamic acid, J. Biol. Chem., 136, 561-2 (1940), Chem. Abs. 35, 167.

Sharzynski, P. & Euler, H. V., Deamination of d-and l-glutamic acid by slices and pulp of sarcoma and kidney, Arkiv Kemi, Mineral., Geol., 13F, No. 17, 8pp., (1940), Chem. Abs. 34, 5095.

Virtanen, A. I., Rintala, P. & Laine, T., Decarboxylation of aspartic and glutamic acids, Nature 142, 674 (1938), Chem. Abs. 33, 2165.

Zorn, K., Tyrosine catabolism and transamination in liver, kidney and muscle, Z. Physiol. Chem., 266, 239-48 (1940), Chem. Abs. 35, 2583.

Bacteria

Barker, H. A., The use of glutamic acid for the isolation and identification of Clostridium cochlearium and Clostridium tetanomorphum, Arch. Microbiol. 10, 376-84 (1939), Chem. Abs. 34, 3304.

Bloch, K. & Schoenheimer, R., The biological precursors of creatine, J. Biol. Chem., 138, 167-94 (1941), Chem. Abs. 35, 2584.

Povarnick, M., The formation of extracellular d(-)-glutamic acid polypeptide by *Bacillus subtilis*, J. Biol. Chem. 145, 415-24 (1942), Chem. Abs. 37, 901.

Bruckner, G., Jr., Ivanovics, G. & Oskolas, M. K., The occurrence of the poly-peptide of d(-)-glutamic acid in nature. Chemical investigation of the substance of the capsules of the *Bacillus* of anthrax, Magyar Chem. Folyoirat, 45, 131-46 (1939), Chem. Abs. 34, 3766.

Chargaff, E., The chemistry of bacteria. XI. The configuration of glutamic and aspartic acids from pathogenic bacteria, (*phytomonas tumefaciens* and *corynebacterium diphtheriae*), J. Biol. Chem. 130, 29-33 (1939), Chem. Abs. 33, 8672.

Clifton, C. E., The utilization of amino acids and related compounds by *Clostridium tetani*, J. Bact. 44, 179-83 (1942), Chem. Abs. 36, 6571.

Coghill, R. D., Configuration of glutamic acid from scarlet fever antitoxin, Science 89, 535-6 (1939), Chem. Abs. 33, 6435.

Feeney, R. E. & Strong, F. M., Growth Stimulating Substances for *Lactobacillus casei*, J. Amer. Chem. Soc. 64, 881-84 (1942), Chem. Abs. 36, 3823.

Hill, J. H., & Mann, E. F., Studies on the interference of certain substances of biological importance with the action of sulfanilamide, J. Urol. 47, 522-30 (1942), Chem. Abs. 36, 4538-9.

Nielsen, N. & Hartelius, V., Growth effect of amino acids. V. Studies on the cooperation of aneurin and beta alanine in combination with asparagine and glutamic acid as growth substances for yeast, Biochem. Z., 301, 125-34 (1939: Compt. rend. trav. lab. Carlsberg, Ser. physiol. 22, 375-86 (1939)), Chem. Abs. 33, 9355-6.

Nielsen, N., Growth effect of amino acids. IX. Supplementary studies on yeast, Compt. rend trav. lab. Carlsberg, Ser. Physiol. 23, 147-53 (1941), Chem. Abs. 35, 8011.

Nielsen, N., The action of amino acids as growth substances with respect to yeast, Biochem. Z. 307, 187-93 (1941), Chem. Abs. 36, 3819.

Pollack, M. A. & Lindner, M., Glutamine and glutamic acid as growth factors for lactic acid bacteria, J. Biol. Chem. 143, 655-61 (1942), Chem. Abs. 36, 4852.

Tumors

Abderhalden, E., Presence of d-glutamic acid in proteins from malignant tumors and normal organs, Z. Physiol. Chem. 275, 135-54 (1943), Chem. Abs. 37, 6324.

Abderhalden, E., The optical behavior of the glutamic acid from the proteins of carcinoma and its metastasis and tissue of origin, Ber. 75B, 1800-02 (1942), Chem. Abs. 38, 1559.

Arnow, L. E. & Opsahl, J. C., The configuration of the glutamic acid of adenocarcinoma protein science 90, 257-58 (1939), Chem. Abs. 33, 8783.

Arnow, L. E. Opsahl, J. C., & Watson, C. J., Configuration of glutamic acid isolated from subacute lymphatic leukemic tissue proteins, Proc. Soc. Exptl. Biol. Med. 43, 766-7 (1940), Chem. Abs. 34, 4791.

Arnow, L. E. & Opsahl, J. C., The attempted characterization of malignant tissue protein with d-amino acid oxidase, Science 91, 431-2 (1940), Chem. Abs. 34, 7401.

Bayerle, H. & Borter, G., Optical rotation of glutamic acid from necrotic tissue, Biochem. Z. 305, 418-21 (1940), Chem. Abs. 35, 510.

Bayerle, H. & Borger, G., Digestibility of tumor tissue by normal enzymes. III. The Enzymatic isolation of glutamic acid, Biochem. Z. 316, 87-95 (1943), Chem. Abs. 38, 4626.

Burger, M. & Plotner, K., The chemical composition of liver and cancer tissue, Klin. Wochschr. 20, 1209-12 (1941), Chem. Abs. 38, 3340.

Cedrangolo, F. & Carandante, G., The transamination of l(+)-and d(-)-glutamic acid in normal tissues and malignant tumors, Arch. Sci. Biol. (Italy) 27, 6-47 (1941), Chem. Abs. 38, 2104.

Chibnall, A. C., Rees, M. W., Williams, E. F. & Boyland, E., The glutamic acid of normal and malignant tissue proteins, Biochem. J. 34, 285-300 (1940), Chem. Abs. 34, 4451.

Chibnall, A. C., Rees, M. W., Williams, E. F., & Boyland, E., Glutamic acid of proteins, Nature, 145, 311-12 (1940), Chem. Abs. 34, 3356.

Clark, W. G., Arnow, L. E. & Opsahl, J. C., Configuration of glutamic acid isolated from proteins of pig and chick embryo tissues, Proc. Soc. Exptl. Biol. Med. 43, 767-69, (1940), Chem. Abs. 34, 474.

Dittmar, I. C., The presence of d(-)glutamic acid in tumor proteins, Z. Krebsforsch. 49, 397-98; 441-42 (1939), Chem. Abs. 34, 1731.

Euler, H. V., Hellström, H., Günther, G., Elliot, L. & Elliot, S., Enzymic experiments on sarcoma and chlorophyll-deficient barley mutants, Z. Physio. Chem. 259, 201-3, (1939), Chem. Abs. 33, 6362.

Euler, H. V., Skarzynski, B. & Höglberg, F. The Biochemistry of tumors; II. Pyroglutamic acid in the blood of rats after implantation of Jensen Sarcomas, Z. Krebsforsch. 50, 230-44 (1940), Chem. Zentr. (1941), I, 215. Chem. Abs. 36, 7121.

Euler, H. V., Günther, G. & Forsman, N.: The biochemistry of tumors. Enzyme systems in the Jensen Sarcoma, Z. Krebsforsch. 49, 49-56 (1939), Chem. Abs. 33, 9423.

Euler, H. V., Hellström, H. & Gunther, G., Enzymic sarcoma studies, Arkiv Kemi, Mineral, Geol. 13B, No. 8, 6pp. (1939), Chem. Abs. 33, 9424-25.

Euler, H. V. & Gunther, G., Transamination and glutamic acid dehydrogenation by Jensen sarcoma extract, Naturwissenschaften 27, 214-5 (1939), Chem. Abs. 33, 9424.

Graff, S., Glutamic acid in malignant tumors, J. Biol. Chem. 130, 13-17 (1939), Chem. Abs. 33, 8778.

Graff, S., Rittenberg, D. & Foster, G. L., The glutamic acid of malignant tumors, J. Biol. Chem. 133, 745-52 (1940), Chem. Abs. 34, 4140.

Kermack, W. O., Recent advances in science: Biochemistry, Science Progress 34, 320-28 (1939), Chem. Abs. 34, 118.

Klingmuller, V., The glutamic acid in tumors, Z. Physiol. Chem. 278, 97-119 (1943) Chem. Abs. 37, 5774.

Kogl, F. & Erxleben, H., Chemistry of tumors. Etiology of malignant tumors, Z. Physiol. Chem. 258, 57-95 (1939), Chem. Abs. 33, 5060.

PLATZIA
Kogl, F. & Erxleben, H., The glutamic acid of tumor proteins, Nature 144, 111 (1939), Chem. Abs. 33, 7376.

Kogl, F. & Erxleben, H., Isolation of glutamic acid from tumor proteins, Naturwissenschaften, 27, 486-7 (1939), Chem. Abs. 33, 9421.

Kogl, F., Erxleben, H. & Akkerman, A. M., Chemistry of tumors. II. Isolation of glutamic acid from tumor proteins, Z. Physiol. Chem. 261, 141-53 (1939), Chem. Abs. 34, 509.

Kogl, F. & Erxleben, H., Chemistry of tumors, III. Stereo chemistry of myoma and other tumors, Z. Physiol. Chem. 261, 154-71 (1939), Chem. Abs. 34, 510.

Kogl, F., Erxleben, H. & Herken, H., Chemistry of tumors. IV. Further stereochemical analyses of tumor proteins, Z. Physiol. Chem. 263, 107-24 (1940), Chem. Abs. 34, 3353.

Kogl, F. & Erxleben, H., Chemistry of tumors. V. The behavior of normal digestive enzymes on tumor proteins, Z. Physiol. Chem. 264, 108-22 (1940), Chem. Abs. 34, 5159.

Kogl, F. & Erxleben, H., Chemistry of tumors, VI. Methods of isolating glutamic acid antipodes, Z. Physiol. Chem. 264, 198-219 (1940), Chem. Abs. 35, 796.

Kogl, F., A chemical theory of cancer, Chem. Weekblad., 38, 450 (1941), Chem. Abs. 36, 5880.

Kogl, F., Erxleben, H. & Veersen, G. J. van, The chemistry of Tumors. IX. Determination of d-glutamic acid in tumor hydrolysates with D as indicator, Z. Physiol. Chem. 277, 251-83 (1943), Chem. Abs. 37, 5482.

- Kogl, F., Notes on recent development in chemistry of tumors. Naturwissenschaften, 30, 46-47 (1942), Chem. Abs. 37, 6738.
- Konikova, A., Glutamic acid of proteins, Nature, 145, 312 (1940), Chem. Abs. 34, 3356.
- Konikova, A., Role of d-glutamic acid in the development of malignant tumors, Biokhimiya, 5, 316-20 (1940), Chem. Abs. 35, 4837.
- Mathers, R.G., Glutamic acid in tumor hydrolysates, Cancer Research 1, 146-47 (1941), Chem. Abs. 35, 4470.
- McDonald, E., Progress of the biochemical research foundation of the Franklin Institute, J. Franklin Institute 230, 727-48 (1940), Chem. Abs. 35, 1831-14.
- Ottawa, H., The Optical activity of glutamic acid isolated from tumor protein, Z. Krebsforsch, 49, 677-8 (1940), Chem. Abs. 35, 4833.
- Town, B. W., Glutamic acid of proteins, Nature 145, 312-13 (1940), Chem. Abs. 34, 3356-7.
- Wieland, T., Glutamic acid from tumors, Ber. 75B, 1001-7 (1942), Chem. Abs. 37, 4793.
- Wieland, T., and Wolfgang, P., Determination of l-and d-glutamic acids in hydrolysates of Erown-Pearce tumors with N^{15} -glutamic acid, Ber. 77B, 34-44 (1944), Chem. Abs. 39, 120.
- White, J. & White, F. R., The nature of the glutamic acid of malignant tumor tissue, J. Biol. Chem. 130, 435-36 (1939), Chem. Abs. 33, 8778.
- Woodward, G. E., Reinhart, F. E. & Dohan, J. S., The glutamic acid of malignant tissue proteins, J. Biol. Chem. 138, 677-88 (1941), Chem. Abs. 35, 3705-6.
- Glutamine
- Archibald, R. M., Chemical characteristics and physiological roles of glutamine, Chem. Reviews, 37, 161-208, (1945), Chem. Abs. 40, 908.
- Ferdman, D. L., Frenkel, S. R. & Silakova, A. I., Glutamine in the tissues of animal organisms, Biokhimiya, 7, 43-58 (1942), Chem. Abs. 37, 4451.
- Fife, J. M. & Ferguson, W. C., The effect of high concentrations of carbon dioxide on the glutamine and asparagine metabolism of the sugar beet, Plant Physiol. 16, 677-90 (1941), Chem. Abs. 37, 1156.
- Fildes, P., Glutamine and the growth of bacteria. Rept. Proc. 3rd Intern. Cong. Microbiol. (1939) 495, Chem. Abs. 35, 1086-7.
- Lankford, C. E. & Snell, E. E., Glutamine as a growth factor for certain strains of *Neisseria gonorrhoeae*, J. Bact. 45, 410-11 (1943), Chem. Abs. 37, 5102.
- Leuthardt, F., Formation of hippuric acid from glutamine, Z. Physiol. Chem. 270, 113-24 (1941), Chem. Abs. 37, 6680.
- Leuthardt, F., Glutamine metabolism, Schweiz. Med. Wochschr. 71, 322-23 (1941), Chem. Abs. 36, 5869.

Leuthardt, F., Formation of urea from glutamine, *Z. Physiol. Chem.* 265, 1-8 (1940), *Chem. Abs.* 35, 1418.

Mothes, K., The synthesis of glutamine and asparagine in vivo, *Planta*, 30, 726-56 (1940), *Chem. Abs.* 35, 4800.

Neuberger, A. & Sanger, F., The nitrogen of the potato, *Biochem. J.* 36, 662-71, (1942). *Chem. Abs.* 37, 1203.

"Orstrom, A., Chemical reactions, especially the ammonia exchange with the stimulation of development of sea-urchin eggs, *Z. Physiol. Chem.* 271, 1-176 (1941), *Chem. Abs.* 37, 2081.

Pucher, G. W. & Vickery, H. B., A method for determining glutamine in plant tissues, *Ind. Eng. Chem., Anal. Ed.* 12, 27-29 (1940).

Schlenker, F. S., Determination of ammonia, glutamine and asparagine amide nitrogen in plant juice, *Plant Physiol.* 15, 701-09 (1940), *Chem. Abs.* 36, 6111.

Vickery, H. B. & Pucher, G. W., Amide metabolism in etiolated seedlings. I. Asparagine and glutamine formation in *Lupinus angustifolius*, *Vicia satropurpurea* and *Cucurbita pepo*, *J. Biol. Chem.* 150, 197-207 (1943), *Chem. Abs.* 38, 6707.

Virtanen, A. I., & Laine, T., Transamination in green plants, *Biochem. Z.* 308, 213-15, (1941), *Chem. Abs.* 37, 2044.

Nutrition

Almquist, H. M., and Grau, C. R., Amino acid requirements of the chick, *J. Nutrition*, 28; 325 (1944), *Chem. Abs.* 39, 1204.

