

Historic, archived document


Do not assume content reflects current scientific knowledge, policies, or practices.

S B
115
B66

LIBRARY OF CONGRESS.

Clas. LIBRARY OF CONGRESS
Copyright No. _____
DUPLICATE
974.366
EXCP

UNITED STATES OF AMERICA.



62 61

1878.

The Potato.

— HOW TO CULTIVATE. —

Chemistry of the Potato.

PREMIUM CROPS AND HOW TO GROW THEM.

ILLUSTRATIONS OF THE LEADING VARIETIES. BEST VARIETIES
FOR SEED.

Centennial Collection of 500 Varieties.



SNOW-FLAKE.—Showing habit of growth in the hill.

B. K. BLISS & SONS,

Seed and Horticultural Warehouse,

NO. 34 BARCLAY STREET,

P. O. Box 5,712.

NEW YORK.

PRICE, 10 CENTS.

FOUR GRAND PRIZE MEDALS, WITH DIPLOMAS,

Awarded to **B. K. BLISS & SONS,**

For their Exhibition of Roots, Vegetables, Potatoes and Garden Requisites, at the Centennial Exhibition, 1876.



WARRANTING SEEDS.

We are frequently asked if we warrant our seeds, and in reply we would say that we do not, any farther than to refund the price paid for the seeds, should they prove to be different from what they are represented. There are so many causes which operate unfavorably in the germination of seeds, over which we have no control, that to do more than this cannot be expected of us. Among the causes of failure to germinate may be mentioned unfavorable weather, which is one of the most important. The soil may be in proper condition when the seed is planted, but the weather which follows may be cold and wet, which will cause the seed to rot, or it may be hot and dry, which destroys the germ before it shows itself. The soil may also be unfavorable for the variety of seed planted. The seeds may be and are frequently destroyed by vermin of various kinds. And lastly, changes not unfrequently occur, especially among newly hybridized varieties, by which frequent sports are produced, which give the planter the idea that his seeds are badly mixed. Such occurrences are beyond the power of man to prevent. For these and other reasons, we wish to state explicitly that, while we exercise the greatest care to have all our seeds pure and reliable, it is hereby mutually agreed between the purchasers of our seeds and ourselves, that we do not warrant the same, and we are not in any respect liable or responsible for seeds sold by us, or for any loss or damage arising from any failure thereof in any respect.

THE FOLLOWING CATALOGUES, PUBLISHED DURING THE YEAR, WILL BE MAILED TO ALL APPLICANTS ON RECEIPT OF PRICES AFFIXED. REGULAR CUSTOMERS SUPPLIED GRATIS.

No. 1. BLISS'S SEED CATALOGUE AND GUIDE TO THE FLOWER, KITCHEN GARDEN AND THE FARM—Enlarged and improved, contains upwards of 200 pages, with many BEAUTIFUL ILLUSTRATIONS, including a splendid Colored Lithograph of a group of favorite flowers, and a descriptive list of upwards of TWO THOUSAND VARIETIES OF FLOWERS AND VEGETABLE SEEDS including many CHARMING NOVELTIES, now offered for the first time in this country, with explicit directions for their culture. Also, a list of upwards of ONE HUNDRED VARIETIES OF FRENCH HYBRID GLADIOLUS, and other SUMMER FLOWERING BULBS. To which is added a list of a few of the choicest varieties of GRAPES, STRAWBERRIES, RASPBERRIES and other SMALL FRUITS, AGRICULTURAL IMPLEMENTS, FERTILIZERS, etc., etc., with much useful information upon the subject of Gardening generally, which will be found useful to the experienced amateur as well as those about to commence the delightful occupation of gardening, published early in January, 35 cents; an edition elegantly bound in muslin, \$1.00.

No. 2. BLISS'S ABRIDGED CATALOGUE AND GARDENER'S ALMANAC.—Published annually; January 1st, profusely illustrated; 128 pages; 10 cents.

No. 3. BLISS'S ILLUSTRATED POTATO CATALOGUE.—Contains a list of 500 varieties Potatoes, embracing several new and very promising varieties, now offered for the first time, with explicit directions for culture, and much other valuable information respecting this valuable esculent. 10 cents.

No. 4. BLISS'S BULB CATALOGUE.—(Illustrated.) Published September 1st, containing a Choice Collection of DOUBLE and SINGLE HYACINTHUS, arranged in their several colors; TULIPS, in many varieties, both double and single; POLYANTHUS NARCISSUS, CROWN IMPERIALS, JONQUILS, SNOW-DROPS, LILIES, etc. 10 cents.

No. 5. BLISS'S ILLUSTRATED PLANT CATALOGUE.—Contains a list of all the most popular varieties of plants for the GREENHOUSE, CONSERVATORY and the FLOWER GARDEN, BEDDING and DECORATIVE PLANTS, ROSES, ORNAMENTAL and FLOWERING SHRUBS, HERBACEOUS PLANTS, FRUIT and ORNAMENTAL TREES, EVERGREENS, SMALL FRUITS, etc., etc., with many beautiful illustrations, mailed to all applicants enclosing three-cent postage stamp.

No. 6. BLISS'S ILLUSTRATED HAND-BOOK FOR THE FARM AND GARDEN.—Contains a catalogue of all the leading varieties of Seeds for the Farm, Kitchen, and Flower Garden, Agricultural Implements, Fertilizers, etc., with many valuable cultural directions; about 100 pages. Sent to applicants enclosing a three cent stamp for return postage.

SEP 15 1876

ILLUSTRATED AND DESCRIPTIVE
CATALOGUE
—OF—
POTATOES FOR SEED,

FOR SALE BY

B. K. BLISS & SONS,

At their Seed and Horticultural Warehouse,

No. 34 BARCLAY STREET,

P. O. Box 5,712.

New York.

•••••
THE POTATO.

This valuable and well-known esculent, now so widely cultivated, is a native of the mountainous parts of tropical America, and was taken to Spain and Italy by early adventurers in the sixteenth century, for we read of its cultivation in those countries in the year 1550. The usual size of the potato in its wild state is rarely more than an inch in diameter, and the flavor very insipid and almost unpalatable.

When first introduced into Europe it created great excitement, of a similar nature to that caused by the introduction of tobacco and coffee; for many years it was only to be found on the tables of the opulent, where it was used as a dessert either in the form of a sweetmeat or as a fruit. The first varieties grown in the United States were brought from Europe. The quality was very poor, and not a variety then in use would at the present time be deemed fit for the table. It is only within a comparatively recent period that it has found its way into both continents as a general article of food for man and beast, or has received attention from agriculturists. Many of our most practical and foremost gardeners are now directing their attention and energies to its improvement and propagation.

In the year 1844, the disease known as the rot appeared and nearly destroyed the whole crop. About this time a few persons, among others the late C. E. Goodrich, of Utica, imported a lot of the wild varieties directly from South America, and proceeded to raise seedlings by crossing with the various kinds then in use. Many thousand seedlings were then produced, but few of them were ever brought to any state of perfection.

One of the principal sorts saved was the Garnet Chili, which had a great reputation, and is the parent of many of our new sorts which are now attracting so much attention. Twenty-five years ago, a farmer who raised one hundred bushels of potatoes was looked upon as having an enormous stock; while to-day, many growers in the vicinity of our large cities raise from ten to twenty thousand bushels, without exciting any unusual attention. Within this period nearly all the numerous varieties with which we are now acquainted have been brought to notice, and every year adds its score of new seedlings to the already overgrown lists. Hundreds of millions of bushels are now raised annually in this country, and the demand is always greater than the supply—the crop of this State alone being twenty-five million bushels, raised on a little over two hundred and fifty thousand acres of land.

The uses of this tuber are numerous, aside from their principal use as an article of food. Thousands of bushels are annually manufactured into starch, and alcohol is distilled from its juices. Even sugar has been made; but with so much competition in this line, it has never proved a great success. Too much cannot be said in favor of this valuable staple, and we shall endeavor to instruct our readers in these few pages as to the methods used to increase the supply, without increasing the expense of its cultivation.

CULTIVATION.

The soil acknowledged to be the best for the potato is a rich loam, sandy, and neither too wet or too dry. A cool, moist soil will produce larger potatoes, but the danger of too much wet is so great that a warmer soil is preferable in this climate. Early potatoes reach their maturity sooner in a quick, light soil, and present a brighter and cleaner appearance, than when dug from that which is heavier. A calcareous formation generally yields a sure crop.

Old sod land, well turned under in the fall, and lightly plowed and harrowed in the Spring, will produce a sound crop, and often an astonishingly large one. Clover sod for this purpose is excellent, and furnishes a large amount of vegetable substance to the ground. When turned under in August or September it will rot by the following Spring, and only a top-dressing of some well-established fertilizer will be required to carry through the crop. Wet land produces a coarse, unpalatable article, and one of little value even as food for cattle. Barnyard manures are of little benefit to such land. Lands should never be plowed while wet and heavy; it injures the soil and does more harm than the manure can offset. Prepare the ground as carefully and thoroughly for potatoes as for any other crop. Attention in this particular well repays the farmer.

Land intended for potatoes requires but little manure, and that should be old and well rotted. By many, spreading the manure before plowing in the Spring is thought to be the best mode. At the time of planting, bone-dust, ashes, plaster, marl and like fertilizers can be used to great advantage with this crop, as they are of a dry or absorbent nature. On wet soils they are very beneficial, as they prevent disease as well as promote the growth of the tubers. On warm, dry, light land, muck compost may advantageously be used; decayed leaves are excellent. In seasons of disease among potatoes, fields where ashes have been used have suffered but little from the rot.

Potatoes are usually planted in drills or hills, the latter being the more common method in this country. Some varieties require more space than others. For drills, two and a half feet by three is ample; while, when planted in hills, three to three and a half feet is the usual distance. The latter method has some advantages, as the cultivator or horse-hoe can be used both ways of the field. Some of the earlier varieties may be planted closer without loss to the crop. Cover about four inches in light soil, and not so deep in the heavier. Cultivation should be commenced soon after the shoots appear above the ground, and weeds should be kept down with as much care as in a carrot-bed. The earth should be drawn a little towards the hill at each hoeing, that the rootlets may gain strength and nutriment from the surrounding earth. When the blossoms appear, hoeing should be discontinued, and in fact is rarely necessary, as the vines then cover the ground and discourage the growth of weeds. An excellent plan is to go over the field occasionally and remove carefully any weeds that appear in the hill, for they draw largely from the sustenance required for the developing tubers.

Many farmers still continue to plant three to five eyes to the hill. In our opinion this is a mistake, and the last few years' experience confirms our views. Potatoes cut *carefully* to a single eye, the hills, perhaps, a trifle closer, will yield a larger crop than the careless way of throwing in seed by wholesale. We all know the effects of too close a growth of carrots, turnips and such roots on the crop, and we think the same rule applies equally to potatoes. Two good eyes to the hill, with a proper proportion of the flesh to each eye, will produce a far larger crop of merchantable tubers than two whole ones; will yield from one-third to one-half more in weight—an item of no small consequence—to say nothing of the saving of nearly one-half the seed. Any intelligent farmer who is up with the times and reads the leading agricultural papers will corroborate this statement.

In order to secure an early crop, the seed must be planted as soon as the ground is in *proper* working order. This time differs in the various parts of our extended country, so that no set time can well be given. The earliest varieties can be marketed in sixty-five to seventy days from planting, thus competing successfully with Southern crops of the more common kinds. A second crop can thus be planted, as described elsewhere, allowed to thoroughly mature, and thus have good seed for another year and a sound article of food for Winter use. The yield of the first planting will give a handsome profit, for *early* potatoes always command a high price when near a city or town. The later varieties do better when planted some weeks after the early kinds; and the farmer has this advantage, he can get the former well started and cultivated before he turns his labor and attention to the latter.

A common method of forcing potatoes is to select whole, sound tubers of some early variety of medium size, placing them close together in a moderately heated bed, composed of either light loam or partially decayed leaves. This should be done several weeks—say three or four—before the time of planting. By that time the tubers will have started sufficiently to be set out. Cut the potatoes into pieces as has been directed, and use care not to injure the young shoots. Plant three inches deep, apply a little horse manure to the bottom of each hill, to afford warmth and to urge forward and nourish the young starting tubers.

The decay of the tops indicate the maturity of the roots. The later kinds often continue to grow till checked by autumn frosts. They should, however, be dug before the ground is touched by frost at night.


HOW TO CUT TO A SINGLE EYE.

Take any potato and hold before you, with the stem end (the place where it was joined to the vine) down. It will then be noticed that the eyes are arranged around the tuber in regular ascending rotation from the bottom to the top, similar to the thread of a corkscrew, each eye being a little above and further around the side than the one next below it. Now take the potato in the left hand, with the stem end down, keeping it in a perpendicular position throughout the entire cutting. Take a sharp, thin-bladed knife and remove the first eye, by placing the knife about equally

distant between it and the eye next in rotation above it, sloping it to the indentedure left by the stem (see dotted lines in center cut), removing the flesh with it. When the first eye is removed, turn the potato around in the hand until the next eye above appears. Remove this one in the same manner, and keep on turning the potato, removing each eye as it appears in exact rotation, always sloping the knife to the stem. After three or four eyes are thus removed, the bottom part of the tuber will have a somewhat pyramidal form (see center cut). It will be noticed that each eye removed has a similar form to that represented by the cut on the left, and has its proportionate share of the flesh attached.

After the first two eyes are removed no further trouble will be found until the seed end is reached, and only a little extra care will be required to remove these closely clustered eyes. The cut on the right represents what remains of the potato after all but the small eyes are removed, while the dotted lines show how to separate each of these. It will be noticed that the base retains the same form throughout, and by sloping the knife each time, and cutting down to the apex of this inverted pyramid (which is the center

of the tuber), each eye will be supported by an equal amount of the flesh which is to start it into a strong, healthy growth. With common varieties, where seed is cheap, many will think this manner of cutting too troublesome. But if every farmer should save one-half his seed, as he easily could do (for it will only require two barrels of sets for an acre, instead of four, as is usually planted, the increase being equal), thousands of bushels would annually be saved, worth many thousands of dollars. But to those who wish to obtain a large increase from a small quantity of seed, as is necessary to those who buy the new and valuable varieties, this manner of cutting so that every eye is saved will prove invaluable.



CUTTING.

This is one of the most important subjects to be considered in the propagation of potatoes, and there is such a diversity of opinion regarding the manner and method of cutting, that many pages could be filled in giving the different experiences of the professors in this art. While we do not attempt to decide this question to the satisfaction of every one, we shall give our own views, and claim that in our method an enormous quantity of the tubers now annually planted may be thrown into the market, causing a reduction in the prices charged for this common and necessary crop. Without discussing the respective merits of planting whole potatoes, or half a dozen pieces, each piece containing three and four eyes, in a hill, we shall state, what has been proven by so many cultivators, that two good eyes are ample for one hill, and the yield of large, marketable potatoes is larger than when more are planted. With the aid of the cut and remarks on page 3, we trust our readers may be enabled to understand our method.

HOW TO RAISE FROM CUTTINGS.

For this purpose any ordinary hot-bed may be used. About the 1st of March take the potatoes to be propagated, dividing them lengthwise, and laying these pieces with the cut side down upon the soil of the hot-bed. Keep them perfectly dry until the cut part has healed over and the sprouts have commenced to start. When the sprouts reach the height of three or four inches, cut them off about half an inch above the eye, and insert the end of the cuttings thus obtained into the soil of the hot-bed. Shade them from the sun, and water carefully until they are well rooted and the leaves begin to develop. The old pieces of the potato will continue to throw up shoots to an almost incredible number, and these are all to be removed as soon as strong enough in the same manner as the first ones. In order to increase the crop still more, as soon as these cuttings have reached the height of eight inches, their tops may also be removed and planted in the same manner as the slips from the potato. As soon as the ground becomes warm and can be worked, prepare it as is usual in planting the tubers, and set out these young plants. It is best to transplant them on some cloudy day or towards evening, as the hot sun withers them and destroys many if planted in the hot part of the day. These plants will be found to grow very rapidly, and can be propagated indefinitely from cuttings of the older plants. No cuttings should be taken after the 1st of August, as they will likely be destroyed by the cold weather before the crop is matured. The immense increase of stock by the use of this method may be illustrated as follows: A pound usually contains four medium-sized potatoes, and there are from twelve to twenty eyes on each tuber. When cut and sprouted they will give, at least, five hundred plants. From each of these plants *three* cuttings may be taken, which gives a total of two thousand plants to be set in the ground. With the ordinary yield, each hill, at the lowest estimate, would give one and one-half pounds, or three thousand pounds; in all making about eighteen barrels of good, sound potatoes, or a year's supply for a large family. We do not claim that this is either profitable or advisable with the common sorts of potatoes; but with the many new and high-priced varieties which are now being disseminated at the prices of one, two, or three dollars a pound, it is almost invaluable, as for a slight expenditure a large stock may be obtained, paying to the propagator a thousand-fold. This is no new experiment, but has been practiced by the initiated for the last few years, and has ever proved a success. It is not confined to hot-beds, but many of our most prominent nurserymen have devoted whole greenhouses to this use, and we would confidently recommend it to our readers.

TWO CROPS A YEAR.

Take good, sound, early potatoes, and cut them into single eyes, as is shown in the article on cutting. Allow these pieces to dry for a day or two, and then plant as early as the ground can be worked (a slight frost will not injure the potato after being well planted). With ordinarily favorable weather the new crop of tubers will mature in from eight to ten weeks. As soon as they are ripe, dig them, and after remaining a day or two in some dry and warm place, proceed to cut them into single eyes as before. Place the pieces thus obtained into pans or boxes containing dry plaster or gypsum. This absorbs the abundant moisture, which would otherwise greatly check the growth if it did not destroy the sets entirely. Allow them to remain in the plaster for ten or twelve days, or until the eyes commence to start, when they are to be taken out and planted as before. In the latitude of New York this is only applicable to early varieties, like the famous Early Rose, or Extra Early Vermont, which are of quick growth, and early maturity; but in many parts of the South, where the growing season is long, it may be practiced indiscriminately upon all varieties. A gentleman has raised *two* crops of Early Rose, a short time since, in this vicinity,

the two crops yielding an aggregate weight of twenty-five hundred pounds. He planted his pound, cut into single eyes, early in March, and dug his first crop about the middle of May. These were then treated as above described and planted the 10th of June, and the second crop dug the 1st of September. The yield from the one pound at the first digging was fifty pounds, and the second crop of this increase was twenty-five hundred pounds, or over forty bushels. This method is within the reach of all, and there is no extra expense incurred for hot-bed sashes or any other forcing requisites.

HOW TO RAISE SEEDLINGS.

Save any well-ripened seed-balls from a good variety, and plant in early Spring, in well-drained boxes of sandy loam. Sow the seed on the surface, and sift fine soil over them to the depth of one-quarter to one-half an inch; water sparingly, and when the seedlings are three inches high, remove them from the seed box without disturbing the earth around them more than is necessary, and plant in more roomy quarters. Many successful growers, however, prefer sowing the seed in open ground, when a partially shaded spot may be selected, and the seeds may be sown in drills about ten inches apart; cover with half an inch of soil. When the plants are strong enough, transplant in rows three feet apart, two feet in the rows, and keep down the weeds until the tubers ripen. Some few strong growing varieties, will form tubers weighing from six to eight ounces the first year. As a general rule they will be about the size of a walnut. The seed we offer this season, "**Pringle's hybrid**," is far in advance of any hitherto offered, as will be seen by referring to the description on page 21. Store the tubers carefully until the next season, keeping them as cool as may be without freezing, when they may be planted in the same manner as any mature potato.

It usually takes three years to ascertain the true value of a seedling, and if a person is favored by finding one really good variety among the many seedlings, he may feel well repaid for his time and trouble. Many new varieties are raised by hybridization, which is a more difficult method, although it generally secures a greater number of good varieties. The manner of procedure is as follows: Remove all flowers excepting those you wish to hybridize, then with a pair of sharp scissors remove all the anthers from the stamens in the flowers to be impregnated, just before they commence to discharge their pollen. When the flowers are dry, shake the flower containing the stamens of the variety which you wish to cross with it, being careful to do it when they are ready to discharge their pollen. Fit a piece of fine netting over the impregnated flower, to prevent the bee and other insects from leaving the pollen of other varieties upon the exposed pistil. The covering may be removed after two or three days. Do not disturb them again until the seed-ball has ripened, when the treatment as given in the first part of this article may be applied.

Instances have been known, though rare, where one potato would produce two distinct sorts from its different buds or eyes. The White Peachblow, for example, has been found growing on the same stalk with the Jersey Peachblow. As so much interest is now excited in the growth and propagation of new seedlings, and many of the new varieties command such high prices—it behoves our farmers and amateur gardeners to avail themselves of the latest and, by actual tests, the best method of producing new varieties.

HOW TO STORE AND KEEP.

It is a matter of no small importance to the farmer to be able to keep his crop of potatoes in good condition through our long Winters, and to present them for sale, free from blemish or mildew, in the Spring. A well-kept potato brings three or four times its value in market in early Spring than the same stock will if sold in the Fall, paying an extra profit over and above the cost of storing, handling and care required.

Of the three methods of storing in general use, each has its champions. They are: storing in barrels, bins, or heaps or pits. The advantage of placing in barrels is, they can be easily handled, do not suffer from abrasion, can be readily looked over, and if *disease presents itself* it can be checked or removed.

When thousands of bushels are raised on one farm, this method cannot be followed on account of the time and expense involved. Bins are largely used by our fore-handed farmers, especially those near large cities, as the roots can at any time be reached and got ready for market. A dry, cool, well-ventilated cellar, with the light excluded, is the best place to store potatoes. It has been found very advantageous in preventing decay to sprinkle lime in the barrels or bin at the rate, say, of one pound to each barrel. It acts as an absorbent and neutralizes the earthy odors, thus directly acting as a preventive of decay to the roots.

The importance of excluding light from potatoes and keeping them as cool as possible, cannot be over-estimated as means of preserving the crop.

The Chemistry of the Potato.

The Variations in its Composition, Average Analyses according to standard authorities.

Dr. Voelker, on Differences of Composition, especially in proportion of Starch.

Manurial requirements: Action of Potash in Potato Culture. Quantity of Nitrogen, Phosphoric Acid and Potash required to produce 100 bushels; also one bushel of Potatoes.

Commercial Sources of Fertilizing Materials required, etc., etc.

The importance of the Potato (*solanum tuberosum*) as one of the most valuable and profitable of the staple food-crops, entitle it to much more attention than it has heretofore received at the hands of agricultural chemists. In England and Germany, where the potato has been made the subject of considerable study and of numerous analyses, it has by no means had the benefit of the exhaustive investigations accorded to the grain and root crops. While much valuable information has been gained, as will be hereafter seen, as the result of these researches, yet there is much to be learned, especially among our own farmers, of the relative values of the new varieties, so popular among us, their capacity for improvement in quality for food, more economical culture, etc. It is to be hoped that the system of experiments so successfully started last season by Professor W. O. Atwater, among farmers, will include trials on the different varieties of potatoes.

The variations in the composition of the potato, as shown by hundreds of analyses made or endorsed by such standard authorities as Voelker, Johnston, Fresenius, Payen, Wolff, Knobbe and S. W. Johnson, are sufficiently great to even lead these chemists to differ somewhat as to what fairly represents an average composition of this important vegetable. As will be seen by examining the data hereafter given, there are large differences in the percentage of the most important ingredient contained in the tuber, namely, the starch, and this difference will be found in the same variety of potatoes grown upon different soils, as well as in different varieties grown upon the same soil.

The feeding value of the potato, to the extent of producing the animal heat of the body and of the formation of fat, is almost entirely due to the amount of starch it contains—and it is equally well known that it is owing to the presence of starch in greater or lesser quantity that some potatoes are, when cooked, *white, mealy* and *fine flavored*, and others, deficient in starch, are unattractive, waxy and soggy.

When the conditions of growth are favorable and the variety a good one, the percentage of starch in the potato tuber is twenty per cent. and upwards, while inferior specimens are found to yield less than eleven per cent. One hundred pounds of the former, when thoroughly dried, would contain over eighty pounds of starch, and the latter some forty-four pounds. In other words, the good potatoes quoted are nearly or quite twice as valuable as food for man or stock as the poor variety.

The variations in the albumen, casein and fibrin are by no means so great as with the starch. The total quantity of these ingredients (called the albuminoids) found in an average healthy potato, is only some two pounds in every one hundred pounds of tubers, in the air-dry or natural state; and it is due to this fact that potatoes, while they have but few superiors in power to maintain life in man or beast, possess so little power in building up the solid tissues of the body, muscle, flesh, etc.

The ability of the farmer to increase the feeding value of the potato, in this direction, by any process of manuring is probably very limited, as the formation of the albuminoids (albumen, casein and fibrin) is due to the presence of nitrogen in the plant, and as the potato seems to have the power of deriving, under ordinary circumstances, through its full leaf and vine development, much of the entire supply of nitrogen it requires or can use advantageously, any addition of this ingredient in considerable quantity, as in the form of nitrogenous or putrescent manures, invariably results in developing an undue growth of vines at the expense of both the quality and quantity of the tubers. In any but a dry season the entire plant is almost sure to

become in an abnormal condition, and prove a prey to attacks of fungus, resulting in disease or "rot." The attempt, on the other hand, to improve the quality of the potato as food, so far as depends upon the presence of starch in the tuber, will be found to be more encouraging when the action of potash in potato manure is considered. As will be seen further on, potash invariably increases the starch formation, and without its presence no starch can be found. When grown upon newly recovered forest land, or upon any land of suitable mechanical texture, containing available potash, or supplied with the same in the form of unleached wood ashes or potash salts (of proper kinds) potatoes are invariably of superior quality.

The variations in the ash or mineral constituents of the potato are remarkable, and probably exceed those of any other crop. With the exception of phosphoric acid and potash in the tubers, and lime, magnesia and sulphuric acid in the vines, the mineral ingredients, soda, iron, pure silica, chlorine, etc., vary to such an extent that they almost seem to be taken up by the roots for convenience rather than for any actual needs of the plants. Soda is found to exist in the ash of the tubers in quantity quite or next to nothing, up to twelve per cent. Oxide of iron from nothing to six per cent. Silica from nothing to over six per cent. Chlorine from nothing to nearly nine per cent.

In all these ash analyses, however, the proportions of potash and phosphoric acid are uniformly large; in the case of potash, from forty-two to seventy-three per cent., while the phosphoric acid is found to rarely fall below twelve per cent., and often rises to over twenty-seven per cent. (average say twenty per cent).

We are thus enabled to see, so far as the development of the tuber of the potato is concerned, that notwithstanding the large variations in the proportions of its ash constituents, the presence of potash and phosphoric acid, but particularly the former is essential. Practice fully corroborates this view, for the good effects resulting from the use of wood ashes and bone, (well known sources for these ingredients,) have caused their almost universal adoption as special manures for potato culture.

The increased yield per acre from the addition of potash alone has been, in several cases, from 150 to 200 bushels in excess of the quantity produced on another portion of the same land to which the potash was not supplied, but otherwise well manured.

The potato vines, as indicated by their analyses, seem to require full supplies of lime and magnesia as well as of sulphuric acid, and as these ingredients are generally found to exist in the vines in increased proportion as the season advances towards October, it would indicate that they are required more particularly for use by the vines themselves. The remaining ingredients found in the vines at the end of their growing season, show, on the contrary, a decreased percentage. This is particularly the case with the potash and phosphoric acid, showing that these ingredients, so important in the growth of the potato, have been transmitted from the vines for use by the tubers.

ANALYSIS OF POTATO VINES (WOLFF).

	AUGUST.	OCTOBER.
	Per 100 lbs. of ash.	
Potash, - - - - -	14.50	6.30
Soda, - - - - -	2.70	0.80
Magnesia, - - - - -	16.80	22.60
Lime, - - - - -	39.00	46.20
Phosphoric acid, - - - - -	6.15	5.50
Sulphuric acid, - - - - -	5.60	5.50
Silica, - - - - -	8.10	4.20
Chlorine, - - - - -	4.60	3.00
Percentage of ash, - - - - -	8.90	5.12

The benefit arising from the use of plaster (sulphuric acid and lime) so successfully practiced by potato growers, as well as from magnesia salts, is probably due to their effects upon the vines, in increasing their power to sustain the entire plant in sufficient strength and vigor to enable it to withstand, especially during seasons of sudden and extreme changes of temperature and moisture, all parasitical attacks or fungus growth. It has been stated that a strong healthy vine is not liable to become affected even when brought into actual contact or rubbed with a diseased plant. The predisposition towards disease, through impaired vitality, must exist before there is danger of "rot."

The success claimed for the good action of common salt (chlorine and soda) on potatoes can hardly be due to any direct manurial action on the plant, but rather to its indirect action in aiding in rendering soluble and available the other ingredients on the soil. As Professor S. W. Johnson states: "Soda is an extremely variable ingredient of the ash of plants, and though gen-

erally present in some proportion, has been observed to be absent in weighable quantities in the seeds of grain and in the tubers of potatoes."

The evidence of the importance of the presence of potash in the soil for potato culture, as given by the fact of its uniform presence in large quantities in the tubers, as well as by the good effects that always attend its use in a fertilizer for potatoes, is made even much stronger when it is remembered that *without the presence of potash in the leaves of a plant no starch can be formed*. Starch, as we have stated, forms from nearly one-half to over two-thirds the entire weight of air-dry tubers.

The following interesting summary of the experiments, made by Dr. Nobbe and others in Germany, is taken from a lecture recently delivered by Professor W. O. Atwater, before the Connecticut State Agricultural Society, and will be found to fully cover the importance of the presence of potash for starch formation.

"If we examine a green leaf with a microscope, we may find in it thousands of minute sacs or cells as they are called. Some of these contain the green substance called chlorophyll. Inside these grains of chlorophyll appear still smaller grains of starch. Starch is composed of the elements carbon, oxygen, and hydrogen. It is produced inside the leaves. The carbon is taken from the carbonic acid of the air by the agency of the chlorophyll, is united with hydrogen and oxygen, the elements of water, and thus starch is produced. After the starch is so formed, it is conveyed to the other parts of the plant, and either stored away, as in potatoes or grain, which consist largely of starch, or transformed into the other materials of which the plant is made up. And this formation of starch in the chlorophyll grains, and transportation, transformation and storage to build up the plant is going on continually as the plant grows. If then, no starch is formed by the chlorophyll in the leaves, normal growth is impossible."

Now, in every one of Dr. Nobbe's plants grown in artificial soil without potash, after the starch of the seeds was used up, no more appeared in the leaves. As Dr. Nobbe says at the conclusion of over a hundred pages of description of his experiments, "**without potash the plant cannot assimilate** (the materials needful for its growth) and shows no increase in weight, because, without the co-operation of potash in the chlorophyll grains, no starch is formed."

The extent of the variations of the leading organic ingredients in potatoes, and some of the reasons for the same as given by Dr. Voelcker of England, are here given as comprising some of the most reliable data offered at the present time.

"The variation in proportion of starch in different samples of potatoes are chiefly caused by the variety cultivated, by the soil upon which and the locality where the plants were grown, and by the time during which the tubers have been kept. Thus the following varieties grown in Scotland yielded respectively of starch:"

Connaght Cups,	- - - - -	21	per cent.
Irish Blacks,	- - - - -	16½	" "
White Dons,	- - - - -	13	" "
Red Dons,	- - - - -	10½	" "

The influence of soil and locality in affecting proportion of starch in the same variety of potato is seen from the following determinations, according to which

Bufs gave in Mid Lothian,	- - - - -	14.89	per cent.
Bufs gave in Forfarshire,	- - - - -	20.71	" "
Cups gave in Argyleshire,	- - - - -	15.14	" "
Cups gave in Mid Lothian,	- - - - -	23.82	" "

On keeping, the proportion of starch diminishes in potatoes. Thus Payen found the same variety of potato to yield in

October,	- - - - -	17.2
November,	- - - - -	16.8
December,	- - - - -	15.6
January,	- - - - -	15.5
February,	- - - - -	15.2
March,	- - - - -	15.
April,	- - - - -	14.2

"The amount of the albuminous or flesh forming constituents of potatoes, according to Boussingault, is greater in newly dug potatoes than in others which have been kept some time. Thus in newly dug potatoes he found them to amount to 2¼ per cent., and in long kept potatoes to only 1½ per cent. of their weight. The first, according to this determination, when dried, contained 9 and the latter only 6 per cent. of albuminous substances."

"In diseased potatoes the proportion of albuminous matters is smaller than in healthy, as the disease primarily attacks the albuminous matters, which, when once attacked are readily decomposed, and changed into ammonia and other compounds."

Dr. Voelcker also states that the analysis of the ash of a diseased potato failed to show any material difference from that of a healthy one. A small increase in the amount of silica in the diseased potato was the only noticeable difference.

PROXIMATE ANALYSIS OF THE POTATO.

Potato Tuber (air dry).

	Average Composition (100 lbs.), as given by				Extreme percentages. (Various authorities.)	
	Voelcker.	Johnston.	Fresenius.	S. W. Johnson.	Lowest.	Highest.
Water,	75.	75.52	71.	76.	68.94	77.
Starch,	15.	15.72	15.	20.	10.75	21.
Albumen, } Gluten, } Casein, }	2.	2.20	2.43	3.	1.50	2.85
Fat,	.24	.24	.20		.10	.42
Fibre,	5.	2.47	7.		3.	7.
Sugar, Gum, etc.,	1.76		3.40			
Ash,	1.		.97	1.	.651	2.

1 Maximum percentage of water in 27 varieties Young Potatoes was 82 per cent. (Voelcker.)

2 Calculated on basis 75 per cent. water in air dry potatoes; from 2.60 to 8 per cent given as ash in potatoes (dried).—(How Crops Grow.)

THE VARIATIONS IN THE ASH INGREDIENTS OF different varieties of Potatoes grown upon the same soil and precisely similar circumstances are shown in the following analyses by Herapath.

	White Apple.	Prince's Axbridge Beauty.	Kidney.	Maggie.	Forty-fold.
Potash,	69.7	65.2	70.6	70.	62.1
Chloride Sodium,	-	-	-	-	2.5
Lime,	3.	1.8	5.0	5.	3.3
Magnesia,	6.5	5.5	5.0	2.1	3.5
Phosphoric Acid,	17.2	20.8	14.9	14.4	20.7
Sulphuric Acid,	3.6	6.	4.3	7.5	7.9
Silica,	-	-	0.2	-	-

The extreme variations in the ash ingredients of Potatoes of different varieties and grown upon various soils may be seen in the annexed table, taken from Prof. Johnson's standard work, "How Crops Grow." (page 156).

Per cent. of ash,	Lowest Percentages, Highest Percentages.	
	38 Analyses.	39 Analyses.
Potash,	Contained in 100 lbs. of ash.	2.60
Soda,		8.05
Magnesia,		42.9
Lime,		0.0
Oxide Iron,		12.80
Phosphoric Acid,		2.5
Sulphuric Acid,		0.5
Silica,		6.20
Chlorine,		0.0
		6.
	11.2	
	0.1	
	27.10	
	0.0	
	18.00	
	0.0	
	6.50	
	0.0	
	8.70	

Average composition of 1000 lbs. of Potato Tubers, natural condition (air dry); also of 100 bushels and one bushel.

	1000 lbs.	100 bushels. 6000 lbs.	1 bush. 60 lbs.
Water,	750.00	4500.	45.
Starch,	150 to 200.00	1200.	12.
*Albumen, Casein, Fibrin,	20.00	120.	1.200
Fibre, fat, etc.,	70 to 20.00	120.	1.200
Ash, 9 to 10 lbs. containing—			
Potash,	5.60	33.60	.336
Soda,	.10	.60	.006
Magnesia,	.40	2.40	.024
Lime,	.20	1.20	.012
Phosphoric Acid,	1.80	10.80	.108
Sulphuric Acid,	.60	3.60	.036
Silica,	.20	1.20	.012
Chlorine,	.30	1.80	.018
Sulphur,	.20	1.20	.012
	.60	3.60	.036
	1000 lbs.	6000 lbs.	60 lbs.

*Nitrogen exists to the extent of sixteen per cent. (about one-eighth entire weight) in these albuminoids. Therefore 1000 lbs. potato tubers would contain 3.20 lbs. of nitrogen.

MANURING OF POTATOES.

The supply of lime and sulphuric to potatoes is cheaply given in the form of plaster, which as before stated, serves an excellent purpose in practice in protecting vines against extreme changes

in weather, liability to rot, etc. Soda and chlorine, if found to act well, will cost but little in the form of refuse salt or brine. The main fertilizing ingredients, however, with which the potato-grower, need concern himself are the following: **Nitrogen, Phosphoric Acid and Potash.** The following gives the quantity of each contained in 1,000 pounds, as well as in 100 bushels, and a single bushel of tubers:

Contained in - - - - -	1,000 lbs.	6,000 lbs. (100 bushels).	60 lbs. (Single bushel).
Nitrogen, - - - - -	3.20	19.20	0.192
Phosphoric acid, - - - - -	1.80	10.80	0.108
Potash, - - - - -	5.60	33.60	0.336
	10.60 lbs.	63.60 lbs.	0.636 lbs.

To supply *all* the phosphoric acid and potash required for the growth of each 100 bushels of potato tubers (the vines supposed to remain on the land) without exhausting the soil of any of its resources in these ingredients, would therefore require say 11 pounds of phosphoric acid, and say 31 pounds of potash, and for *continued potato culture* an application of an equal quantity for *each crop*. In the case of nitrogen, however, the quantity may safely be reduced to 10 pounds, instead of using nearly 20 pounds, as called for by the composition of 100 bushels of tubers. The reasons for this reduction in the quantity of nitrogen in the manure are as before explained, viz.: The natural power of the potato plant to derive nitrogen from the air, (a cheap source,) the tendency as shown in practice, of nitrogen to induce, except when used in very limited quantity, an **over-rank vine-growth** and a **poor quality and a limited quantity of tubers**, besides an almost certain tendency on the part of the potato crop towards "disease."

Taking the quantities of nitrogen, phosphoric acid and potash, as above advised, we have the cost of the same at the current rates, furnished in the best known forms, (Peruvian Guano, blood or flesh, dissolved-bone and potash salts), as follows:

For 100 bushels of potatoes:

	Pounds.	Price per lb.	
Nitrogen, - - - - -	9.60	21 $\frac{1}{4}$ c.	\$2.05
Phosphoric acid, - - - - -	10.86	9c.	.97
Potash, - - - - -	33.60	7 $\frac{1}{2}$ c.	2.52
Cost for 100 bushels, - - - - -			\$5.54

Equal to a cost for single bushel. 5 $\frac{1}{2}$ cents.

To grow 300 bushels of potatoes, irrespective of resources of the soil, would require of:

Nitrogen, - - - - -	29 lbs.
Phosphoric acid, - - - - -	32 lbs.
Potash, - - - - -	100 lbs.

The following table gives the leading commercial sources for the above named ingredients, and also the quantities of each required to supply the necessary quantity of potash, phosphoric acid and nitrogen required for 300 bushels of potatoes.

COMMERCIAL SOURCES OF POTASH.

Quantity of Each Required to Yield 100 lbs. of Potash.

		Potash.	Phosph acid.
		lbs.	lbs.
Wood ashes (unleached, 25 bushels), - - - - -	1,200 lbs. will yield	100	50
Wood ashes (leached, 117 bushels), - - - - -	6,670 lbs. " "	100	100
*Sulphate of Potash, - - - - - "Kainit"	800 lbs. " "	100	
" " " - - - - -	50 per cent. 361 lbs. " "	100	
" " " - - - - -	72 per cent. 250 lbs. " "	100	
" " " - - - - -	80 per cent. 228 lbs. " "	100	
†Muriate of Potash, - - - - -	80 per cent. 200 lbs. " "	100	

*In the use of the low grade potash salts known as "Kainit," as a source of potash for potatoes, it is necessary to take precautions against damage arising from the presence of chloride of magnesium which this grade of potash salt contains to the extent of some ten per cent. (200 lbs. to the ton). These "Kainit" salts average only some 13 per cent. of actual potash, it therefore requires some 700 lbs. per acre, provided this salt is used alone, to give 100 lbs. of actual potash, (the quantity required for 300 bushels of potatoes). The use of this quantity per acre would be attended with serious danger to the crop owing to the presence of some 70 lbs. (10 per cent.) of chloride magnesium. The only safe method of application is to sow broadcast and harrow in several weeks before planting.

Professor S. W. Johnson advises the mixing of fifteen pounds of air-slaeked lime with each one hundred pounds of "Kainit" before use. This is said to counteract any injurious effects that might otherwise arise from the presence of the chloride of magnesium. In Germany they seldom use this grade in quantities exceeding 500 to 600 lbs. per acre.

†The **muriate of potash** is open to the objection for use on potatoes, that it has a tendency to produce tubers of an inferior quality—watery and waxy. It is recommended to use as a source

of potash, when the formation of starch or sugar is desired, either wood ashes or some grade of sulphate of potash (as free as possible from any chloride salt.)

COMMERCIAL SOURCES OF PHOSPHORIC ACID.

Quantity of Each Required to Yield 32 lbs. of Phosphoric Acid.

		Nitrogen.	Phos. acid.	Potash.
		lbs.	lbs.	lbs.
Pure Fine Bone, (dissolved in sulphuric acid),	267 lbs. will yield	6	32	
Pure Bone Meal, - - - - -	160 lbs. " "	6	32	
Dissolved Bone Black, - - - - -	230 lbs. " "	" "	32	
Rectified Peruvian Guano, - - - - -	228 lbs. " "	16.	32	4½

COMMERCIAL SOURCES OF NITROGEN.

Quantity Required of Each to Yield 30 lbs. of Nitrogen.

		Nitrogen.	Phos. acid.	Potash.
		lbs.	lbs.	lbs.
Dried Flesh or Blood, - - - - -	300	30		
Rectified Peruvian, - - - - -	375	30	52½	6½
‡Nitrate of Soda, - - - - -	192	30		
§Sulphate of Ammonia, - - - - -	145	30		
Pure Bone Meal, - - - - -	800	30	160	

‡These articles, nitrate of soda and sulphate of ammonia, as they furnish nothing of value but nitrogen, should be used in potato culture with caution, and in small quantities. Their effect upon potatoes, except when used on soil abounding in the mineral ingredients (particularly phosphoric acid and potash), will be to produce an excessive vine-growth and a yield of tubers small in quantity and inferior in quality.

***A COMPLETE MANURE FOR POTATOES.**

The composition of a good potato manure for average soils, and suitable for continued culture of this crop, would be:

	Per 100 lbs.
Nitrogen, - - - - -	4 to 5 per cent.
Phosphoric acid (from dissolved bone or Peruvian Guano), - - - - -	5 to 6 " "
Potash (in the form of a Sulphate of Potash), - - - - -	13 to 14 " "

*This formula will be found to correspond in composition with the potato manure offered for sale by us. Price per ton, \$52.00; per bag (200 lbs.), \$5.40; cartage (50c) charged only on less than half ton. One to three bags sufficient per acre. Directions for use accompany each bag. See our Price List of Fertilizers.

Eureka Tree and Post-Hole Digger.

Indispensable to Every Farmer, Nurseryman, and Railroad and Telegraph Co.

The Grand Medal of the Centennial Exhibition has been awarded to us for our Digger for "distinguished merit" and great "excellence." We have also received the prize at six State, and twenty County fairs, and have always received the premiums wherever we have exhibited.

This tool has been thoroughly tested during the past three years, and as yet no place has been found where it has failed to do its work satisfactorily. The principle on which it works prevents adhesion in sticky soils, hence it always works free and easy; no clogging, no knee work, no backache. It is far superior to all Plungers, Augers and Boring Machines, as it works equally well in stony, sandy and clay soils. Quicksand under water is as certainly handled and removed as though no water existed. Hundreds have used them for setting curbs in wells, where there was quicksand. For setting out trees and transplanting it is unequalled; also for cleansing out wells and for tiling.

One man with this Digger can do five times as much work in the same time, as can be done in any other way. One great advantage in using the Eureka Digger is, that you need make the post-hole only a little larger than the post, thus causing it to set firmer than where a larger hole is dug. The size of the hole dug is bounded only by the requirements of its purpose, and can be made of any inclination or shape, and anything that can be reduced or loosened to five inches or less in diameter, can be easily removed. As constructed for ordinary use, it will dig readily four feet deep. Its durability will equal any tool made for any purpose, the material being cast steel. But should any part fail from use or accident, duplicate parts can be furnished; in this way it can be kept good for a generation. Price, \$5.00. Send for Circular.




NEW VARIETIES
OF
POTATOES FOR 1878.

WE take much pleasure in offering two new Potatoes, which we are confident are destined to occupy a prominent position in the list of varieties heretofore introduced by us.

We refer with pleasure to the great popularity of most of the varieties sent out by us within the past ten years, among which are the **Early Rose**, **King of the Earlies**, **Bresee's Prolific**, **Peerless**, **Climax**, **Late Rose**, **Extra Early Vermont**, **Compton's Surprise**, **Brownell's Beauty**, **Eureka**, **Snowflake**, **Alpha**, **Ruby**, **Centennial**, **Superior** and **Bliss's Improved Peach Blow**. There is no section of the country in which these varieties have been tested, where they have not proved superior to the older varieties in cultivation, so much so, that in our leading markets, by far the greater portion of those now offered for sale were first introduced by us. Many of our varieties are also exceedingly popular in foreign markets, where they have been successfully hybridized with their finest sorts, producing varieties far superior to those formerly in cultivation.

Notwithstanding the many fine sorts now in cultivation, we believe the Potato to be susceptible of still greater improvement, and our numerous growers in various parts of the country, are constantly exerting themselves, by careful hybridization, to introduce some new variety which shall exceed, in some important respect, those already in cultivation. We have many very promising varieties on trial, and shall continue to add to our list, from time to time, such as we think worthy of dissemination.




TROPHY.

This new and exceedingly fine variety is, like most of those heretofore sent out by us, a native of northern Vermont, and is one of several very promising varieties raised from a packet of Pringle's Hybridized Potato seed, by a most successful cultivator, and is sure to become a general favorite. It is a seedling of the Ruby impregnated with the Excelsior, and is a decided improvement upon both of these varieties.

Tubers of medium size, very regular in form, elongated oval, somewhat flattened, eyes very few, almost flat upon the surface, resembling, in many respects, the well-known Snowflake, skin reddish, slightly russeted, flesh fine grain, white, and of excellent quality, either baked or boiled. Vines are stout and vigorous, foliage dark green. It matures its crop about two weeks later than the Early Rose. The tubers are compactly clustered around the base of the stalk, and easily dug. It is a good keeper, very productive, and we are confident will become a general favorite.

This variety was exhibited in London at the great International Potato Show, in October last, and received a first-class certificate.

Price, \$1.00 per pound; three pounds to one address, \$2.50; by mail, post-paid.



BLISS'S TRIUMPH.

This new and beautiful variety is, without exception, the most attractive in appearance of any that we have yet offered, and will also compare favorably with the best of them in quality. It originated in the State of Connecticut, several years since, and has been faithfully tested by the originator, and found to maintain its good qualities from year to year. It was raised from a seed ball of the well-known Peerless crossed with a seedling of the Early Rose. It combines the wonderful productiveness of the Peerless, with all the good qualities of the Early Rose, is much more productive, and matures its crop at least ten days in advance of that favorite sort, before the second crop of beetles appear. In color and form it resembles the Garnet Chili, though greatly improved in form and quality. Tubers of medium size, round and uniform in shape, with but a very few small ones, eyes slightly depressed, color a beautiful light red, strongly resembling the early varieties from Bermuda; flesh fine grain and of excellent flavor. Vines about two feet in height, erect, with but a few lateral branches, covered with long, dark-green foliage.

The tubers grow compactly in the hill, are easily harvested. It is an excellent keeper; not inclined to sprout early. Its great beauty, productiveness, and fine quality, will make it one of the best market varieties in cultivation.


Price, \$1.00 per pound; three pounds to one address, \$2.50 by mail, post-paid. By express or freight, charges to be paid by the purchaser, one-half peck, \$3.00; one peck, \$5.00.

HARLEQUIN,

OR VARIEGATED-LEAVED EARLY ROSE.

This singular variety was discovered, several years since, in a field of Early Rose, and only differs from that variety in the foliage, which is beautifully variegated with different shades of white and yellow, resembling many of the ornamental varieties of the Coleus now so extensively grown in the flower garden. It has been cultivated for several years, and the variegations are quite constant. In some soils it is almost entirely white or straw color, with an occasional green leaf, and in others the green predominates. The variegation is much more marked in light soil but little manured. By planting this variety, you not only have a desirable crop for domestic use, but a decorative plant worthy of a place in every flower garden.

Price per pound, \$1.00; single tubers, 25 cents each.



MANHATTAN.

In this variety we have what may be called an "Improved Compton's Surprise." It combines all the good qualities of that favorite variety, but is much more regular in form, and consequently more desirable as a market variety. In shape it is nearly round, sometimes a little oblong, skin dark purple, occasionally blotched with white; origin unknown; flesh white, very solid, fine grain, cooks through evenly, either in baking or boiling; dry and mealy, and of excellent quality. Size, medium to large, eyes slightly depressed, vines vigorous, a little spreading, of a dark green color; one of the most productive in cultivation. It is an excellent keeper, and can be strongly recommended as a main crop variety.

Price, 75 cents per lb.; 3 lbs. \$2.00, by mail post-paid; by express, freight paid by the purchaser, \$3.00 per peck, \$10.00 per bushel.

EARLY OHIO.

A seedling of the Early Rose, and similar in color, habit of growth, and appearance, with the exception of being a round oblong, while the former is more of an oval oblong, so that side by side it is readily distinguishable. It is several days earlier and more productive, and of very fine quality. It has given general satisfaction wherever it has been tested.

By mail, 60 cents per pound; 2 pounds \$1.00.

By express or freight at expense of purchaser, 1 peck \$1.00; 1 bushel \$2.25; 1 bbl. \$5.00.


BURBANK'S SEEDLING.

A seedling of the Early Rose, tubers large, long and slim, eyes few and but little depressed; flesh white, fine grain, dry and floury; cooks through readily. It is a second early variety, very productive, and superior for the market, as its size and general appearance are very much in its favor.

By mail, 60 cents per pound; 2 pounds \$1.00.

By express or freight at purchaser's expense, 1 peck \$1.00; 1 bushel \$2.25; 1 bbl. \$5.00.

Varieties for 1877.



SUPERIOR. (Brownell.)

The parentage of this variety is the same as that of the "Centennial." It was produced in 1873, from a seed ball of Brownell's Beauty, fertilized with Peachblow. The yield of this new seedling potato is enormous; six hundred and seventy-three pounds were grown from one pound of seed, during a season of unprecedented drouth, when many older varieties did not yield enough to pay for the digging. Its tubers are medium to large, elongated oval or cylindrical, of a peculiar dark copper color, and very uniform and handsome in appearance. Skin very fine and smooth, eyes few and small. The vines are strong and healthy, and the growth of the roots and tubers, close around the stalks. It ripens second early or medium late; keeps well during Winter, and retains its mealiness and excellent table qualities through the entire season. A certificate of merit was awarded to this variety at the great International Potato Exhibition at London, last September.

Price, 60 cts. per pound, 2 lbs., \$1.00, by mail to one address, post-paid. By express or freight, charges to be paid by purchaser; 1 peck \$1.00, bush., \$3.00, bbl. \$7.00.

DUNMORE.

This new seedling—a white skinned and white fleshed variety, which originated in Vermont, tested side by side with over forty varieties, in every requisite of a first-class potato, ranks but second to the Burbank. It is superior in its yield, size of the tubers, handsome appearance and fine floury quality either boiled or baked, of the varieties that have recently become famous. As the same potato varies on different soils, possibly on some soils the Dunmore may give greater satisfaction than the Burbank Seedling.


The past season the crop has been remarkably large. Some of the potatoes weighing two pounds each, and were perfect in shape. Its general appearance somewhat resembles the Peerless, but it surpasses that well known variety in both yield and quality.

By mail 60 cts. per lb., 2 lbs. \$1.00. By express or freight, at purchaser's expense; 1 peck, \$1.00, 1 bush. \$2.25, 1 bbl. \$5.00.

WHITE ROSE.

Similar in many respects to the Late Rose, said by the raiser to be of very superior quality.

By mail, 60 cts. per lb., 2 lbs. \$1.00. By express or freight, 1 peck \$1.00, 1 bush. \$2.25, 1 bbl. \$5.00.



CENTENNIAL. (Brownell.)

This new seedling was raised by E. S. Brownell, the originator of that well-known and favorite variety, "Brownell's Beauty." It was produced in 1874, by fertilizing the blossoms of the Brownell's Beauty with pollen from the White Peachblow. The vines are upright, stout, vigorous and of medium height; foliage dark green, strongly resembling the leaves of a Raspberry; very healthy. The tubers are compactly clustered around the base of the stalks, consequently easy to dig; are of a good medium and uniform size; shape nearly round, somewhat flattened, very symmetrical, remarkably uniform and handsome, never rough or prongy, eyes few and quite small, and but slightly depressed near the seed end; stem set in a shallow, round basin; skin is of a deep red color, smooth and uniform in coloring; season second early or medium. Its flesh is of exceedingly fine grain, white, and when boiled or baked of a lightness and porosity seldom equaled; cook through evenly without any hard or watery core.

During the past two seasons it has proved to be one of the most productive varieties; its tubers were perfect in every respect, never hollow or false-hearted; and their excellent and delicate flavor places them in the first ranks of our best table potatoes.

Per lb., 60 cts.; 2 lbs. to one address, \$1.00 by mail, prepaid. By express or freight, charges paid by the purchaser, 1 peck, \$1.00; bush., \$3.00; bbl., \$7.00.

TWO NEW VARIETIES OF SWEET CORN.

Dolly Dutton.—The earliest sweet corn ever introduced. A very dwarf-growing sort, stalks from three to four feet high. Ears small, averaging from four to five inches in length, kernels of good size; tender, very sweet and delicious. It ripens from seven to ten days earlier than the Early Minnesota, and is of much better quality.

Package containing sufficient for fifty hills, 25 cents; five packages for \$1.00; selected ears, 15 cents each; two for 25 cents; \$1.25 per dozen.


Washington Market.—This is, without exception, the best of the large varieties of sweet corn, and when better known, will, we are confident, become a standard variety for general cultivation. See description, page 120, in our Guide to the Flower and Kitchen Garden.

Select ears, 30 cents each. Packages containing sufficient seed for one hundred hills, 25 cents; 50 cents per pint; 75 cents per quart. By express, freight paid by the purchaser, or when delivered at our counter, 50 cents per quart; \$3.00 per peck.

EARLY LIMA BEAN.

This variety has been in cultivation for several years by a skillful horticulturist, but is now offered for the first time. A careful trial the past season proved it to be ten days earlier than the ordinary variety, besides being very productive and of extra fine quality.

Packets, 25 cents each; per pint, \$1.00.



IMPROVED PEACHBLOW.


A cross between the "Jersey Peachblow" and "Excelsior," grown in 1873 by an experienced hybridizer of Vermont, who has devoted many years in experimenting with the Potato. It partakes some characteristics of each of its parents, the vines and leaves having the appearance of the "Excelsior," while its tubers resemble the "Peachblow." The form of the tubers, however, is more round and regular than that of the "Peachblow." While in quality it fully equals that old standard market sort, in productiveness it far excels the same, yielding nearly double as much per acre. Its season of ripening is somewhat earlier than that of the "Peachblow," and the growth of its tubers is more compact and closer in the hills. Taking all its points together, it may be considered a rejuvenated and reinvigorated "Peachblow," combining all the best qualities of that general favorite, in its best days, with the additional advantages of earlier ripening and compacter growth.

Price, \$0.60 per lb.; 2 lbs. for \$1.00, by mail, to one address, postpaid. By express or freight, charges to be paid by the purchaser, 1 peck, \$1.00; bush., \$3.00; bbl., \$7.00.

Bliss's Improved Long Orange Carrot.

This superior variety is the result of a careful selection, for successive years, of the best formed, largest and deepest colored roots of the Improved Long Orange Carrot, by which it has attained a perfection hitherto unknown in this useful vegetable, being larger, better flavored, and of a deeper orange color, and more sure to produce a crop. Butter makers will find this variety very useful in giving to their butter a rich, deep yellow color. We unhesitatingly pronounce it the best variety in the market, and one which will not fail to give satisfaction to the purchaser. One ounce, 15 cents; four ounces, 40 cents; one lb., \$1.25; by mail, postpaid.

Varieties of 1875 and 1876.



RUBY. (Pringle.)

1,982 lbs. (33 bushels) grown from one pound of Seed.

A new red kidney-shaped Potato of superior quality, raised in 1871 from the Early Rose hybridized with the White Peachblow. Its hybrid origin is quite apparent. The shape is oblong, slightly flattened, resembling that of the Early Rose. In its coloring the red of the latter variety is deepened by the earmine which shows in the blotches of the White Peachblow. The eyes, which are but slightly sunken, are carmine like those of this variety. The flesh possesses much the character of the Peachblows, being white, fine-grained, firm, and of excellent flavor. The tubers are of approved medium size; and are clustered close about the foot of the stalks. These are short and stout, with foliage broad, thick, and of a very dark green color. The time of maturity is the same as that of the Early Rose, and it is equally productive.


In short, we believe this variety combines in a rare manner the good points of its parents; and to those with whom the Peachblows are favorites we can recommend it with confidence as an early variety of the Peachblows without any of their faults. At the recent International Potato Exhibition held in the Alexandra Palace, London, in September, this seedling, in common with a few others sent by us, received a certificate of merit. To those who prefer a red-skin Potato, we can confidently recommend this excellent variety.

The following extract from the report of the committee who selected the premiums offered by us for this variety, gives their opinion of its merits:

"The Ruby, although not presenting as striking characteristics as the *Alpha*, has proved to be a valuable introduction. It is declared to be a 'No. 1 Potato in every particular,' and seems to be especially valued for its remarkable exemption from rot. Although there was so much rain in some of the Western States, that Potatoes generally failed to grow, and those that grew generally rotted, yet no rotten ones were found among these, and their quality was excellent. Its large yield and very handsome appearance, combined with excellent quality, make it a valuable market variety, where red-skinned Potatoes are in demand." In Europe it has likewise succeeded exceedingly well. Mr. P. Robertson of Scotland, says: "they were pronounced of excellent quality by every person who has tried them."

For further information see the reports of the successful growers in the following pages:

Per lb., 60 cts.; 2 lbs. to one address, \$1.00, by mail, prepaid. By express or freight, charges paid by the purchaser, 1 peck, \$1.00, $\frac{1}{2}$ bush., \$1.50, bush., \$2.25, bbl., \$5.00.



ALPHA. (Pringle.)

The Earliest Variety in Cultivation. 1,707 lbs. (28 1-4 bush.) grown from one pound of Seed.

Raised in 1870, from seed borne on Early Rose and impregnated by pollen of Sebee. A very early variety for farm and garden culture, also for forcing under glass; fit for the table ten or fifteen days before the Early Rose. Tubers of medium size, oblong, somewhat flattened, with eyes but slightly depressed; color a clear white, with the slightest tinge of red about the eyes; flesh very white, fine grained, dry and firm, and possessed of a decided and excellent flavor; stalks short and close jointed, seldom exceeding a foot in height; leaf broad, light green and shining above; tubers clustered about the base of the stalk; quality of the highest excellence. A first-class certificate was awarded by the Royal Horticultural Society, of London, in 1874. A silver medal was also awarded by the Massachusetts Horticultural Society. It has been thoroughly tested the past season in various sections of this country, and has given perfect satisfaction. We can recommend this with the greatest confidence as *the very best early variety in cultivation.*

(From the Report of the Royal Horticultural Society of London.)

ALPHA—(B. K. BLISS & SONS.)—Haulm compact, about 12 inches long; ripening off very early, stem pale green, leaflets broad, flat, very pale green; tuber medium size, half round, flat; eyes large, skin smooth, very clear, pale straw-colored; flesh firm, white, of excellent quality for early use. Moderate cropper; one of the very earliest of potatoes. *First-class certificate.*

The committee who awarded the premiums offered by us last Spring, after examining the various communications received from competitors for premiums, report as follows:

"The Alpha has by many growers been declared "much the earliest of any seedlings." It was found to be "fit for use, in sixty days from the day of planting," "of excellent quality when cooked in any way, and gaining steadily in quality and yield." In this latter respect the Alpha differs from most new seedlings. But few improve after the third year, while many deteriorate rapidly. The Alpha, when first brought to notice, was below medium size, and so delicate that it was thought only suitable for garden culture. But gradually we found it increasing in size and productiveness, while it retains its earliness and excellent quality. That it will henceforth rank as the earliest Potato for the field as well as the garden, and that it yields enormous crops, even under ordinary culture, has been sufficiently proved by Mr. Clute's 1,535 pounds, grown without manure whatever." For further information respecting this variety, see the reports of the successful competitors in the following pages.

Per lb., 60 cents; 2 lbs. to one address, \$1.00, by mail, prepaid. By express or freight, charges paid by the purchaser, 1 peck, \$1.25; ½ bushel, \$2.00; 1 bushel, \$3.00; 1 barrel, \$7.00.

THE SNOWFLAKE. (Pringle.)

1,417 lbs. (23 6-10 bushels,) grown from 1 pound Seed.

This new variety, first sent out by us in the Spring of 1873, has been thoroughly tested, both in this country and in Europe the past two seasons, and we have yet to learn of the first instance where it has failed to give entire satisfaction. The superior quality claimed by the originator, when first offered, has been confirmed in every case as far as heard from.

It is one of the earliest varieties, ripening about the same time as the Early Rose. The tubers are of a good medium and uniform size; shape elongated oval, compressed, exceedingly symmetrical and remarkably uniform; eyes few, entirely flat on the base and body of the tuber, and but slightly and sharply depressed near the seed end; skin white with a russety tinge, and somewhat roughish and tessellated. Its flesh is of exceedingly fine grain, snow-white when boiled, and of a lightness and porosity almost approaching a snowflake. In quality, we do not hesitate to say nothing can surpass this new variety; its mealiness, its pure, delicate flavor, and the evenness with which it cooks through, have never been eclipsed by any Potato. As a baking Potato, it is equally valuable, and as such is distinguished for its pure starchy texture, and delicate nutty flavor. The tubers have attained the full development of their quality as soon as they are fit to dig, and do not lose it during Winter; samples kept till the first of June, did not show the least deterioration. The vines are of medium height, stout and vigorous; leaves medium, and of dark green color. The tubers are compactly clustered around the base of the stalks,—an important consideration in digging the crop. The variety has been tested on widely varying soils—sand, gravel, loam, as well as heavy clay—and has, in every case, given the same favorable results, and often produced a yield of from 300 to 400 bushels per acre. In every case it has proved healthy and hardy, while other varieties alongside of it failed to give satisfactory results.

We could fill quite a volume with the many letters of commendation that have been received from various growers throughout the country in favor of this fine variety, but for want of room we publish those only which have been received from the successful competitors for the prizes offered by us last Spring. These will be found in the report of the Committee in the last pages of this Catalogue.

We take pleasure in submitting the following extract of that report to our friends, which cannot but satisfy the most incredulous of its superiority.

"The Snowflake has received more and higher praise than has probably ever been bestowed upon any Potato. There is no dissenting voice among the whole list of reports, nearly every one of which contains '*It is the best Potato I ever saw.*' Its quality and uniformity of size are especially commended. In many cases, 25 to 40 perfect Potatoes were found in every hill planted, and 'tubers of two and three pounds each cooked readily and completely through.' Mr. Perkins could select 1000 tubers weighing 1000 pounds from a gross product of 1304 pounds, and finds them preferable to any Potato out of over a hundred varieties he grew. Mr. Salter 'never saw so fine a Potato; beautiful in color and shape, firm in texture, flesh white; luscious cooked in any way; it stands unrivalled.' There is certainly within our knowledge no variety which combines all the essential points of a Potato in as high a degree as the Snowflake. Quality, shape, size, color, yield, are all that can be desired, and it is difficult to perceive in what direction further improvement can be obtained."

By mail postpaid, per pound, 60 cents; two pounds, \$1.00. By express or freight, charges paid by the purchaser, 1 peck, \$1.00; ½ bushel, \$1.50; 1 bushel, \$2.25; 1 barrel, \$5.00.

(From the Report of Messrs. Carter's Royal Metropolitan Root Show, in Bell's Weekly Messenger, Nov. 20 1876.)

A dish of Snowflake Potatoes, sent by Mr. Penny, head gardener to his Royal Highness the Prince of Wales, cannot be adequately described in words; but we may advise any professional or amateur grower to secure a supply of this stock for trial next year, that they may put it to a practical test under their own eye.

(From the London Gardeners' Chronicle of Jan. 16, 1875.)

"It is not many years ago, since our American cousins introduced us to the first of their wonderful 'Taters,' and just see what a revolution they have caused! On the exhibition tables, frequently no other than American varieties are to be found—they are unmistakably great croppers and of fine, taking appearance. We cannot say in points of quality they are always so superior, yet, we can say, from personal experience, that some of the best potatoes we tasted during the past season were of the American sorts. Of this year's introduction, we would first especially note Snowflake. The tubers of this sort are of a long ovate form, the eyes very full, skin rough, pale straw, the most handsomely formed of all, and of excellent quality."



Mr. Burnett's Crop of Snowflakes from One Pound of Seed.

SKANEATELES, N. Y., September 27, 1875.

Messrs. B. K. BLISS & SONS:—Accompanying this please find Stereoscopic view of the **Snowflake Potato** as grown in my garden the past Summer. It exhibits the product of one pound of seed purchased of you last Spring. The earth was carefully removed, leaving each potato attached to the root *exactly as it grew*. They were grown in good garden soil, without extra manure, and had the same cultivation as the other potatoes. No one could have been more surprised than I at the wonderful production. I had them on exhibition all one day, and a large number of farmers and amateur gardeners called to see them. For size and quantity all admitted they had never seen its equal. At the nearest end of the row, as shown in the picture, the yield was affected by a shade tree, but at the other end they were very large and in one solid mass—in fact, there was no room for a jack-knife between them. It makes quite a pretty picture, with rose bushes, geraniums and tuberose on the left and my hoy "Frank" and the grapevines for a background. I have created no little excitement by exhibiting the Snowflakes at our Town Fair.

Yours respectfully,

JOSEPH H. BURNETT.

A Trial of the Snowflake in June.

In order to show the wonderful keeping properties of the Snowflake Potato, we annex a letter from the well-known horticulturist, Chas. Downing, Esq., to whom a sample grown in 1874, was sent in June for trial.

NEWBURGH, June 28, '75.

Messrs. B. K. BLISS & SONS—*Dear Sirs*: The box of potatoes was received in due time. The quality is equal and I think superior to any potato I ever ate so late in the season, and being of good size and so smooth and regular in form, it will be an acquisition if the production is sufficiently good.

Please accept my best thanks for your kindness, and for the opportunity to test them.

Very respectfully,

CHAS. DOWNING.

From Nash & Crook, proprietors of the well-known and popular restaurant in the *Times* building, opposite the new post-office on Park Row:


NEW YORK, July 1, 1875.

Messrs. B. K. BLISS & SONS:—We have been using the Snowflake potatoes, supplied by you in our restaurant for the past month, and find them superior in every respect to the Bermuda or any other early variety in the market. They cook dry and mealy, and are of excellent quality; and on account of the eyes being so nearly even with the surface, there is but little or no waste in peeling. They give general satisfaction to our customers. We have no hesitation in pronouncing them the best potato we ever used in our business at this time of the year.


NASH & CROOK.



SNOWFLAKE.—Showing Habit of growth in Hill.



SNOWFLAKE. (Pringle.) Page 20.



BROWNELL'S BEAUTY. (Brownell.)


This is another of the varieties first sent out by us in the Spring of 1873, the beauty and superior keeping qualities of which, together with its fine quality as a table variety and productiveness, places it in the front rank of those recommended for general cultivation. We know of no varieties whose good qualities can be retained for the entire year, as this has done. Potatoes of the crops of 1873 and 1874 have been exhibited side by side at several State and County exhibitions, those of 1873 having been kept in an ordinary cellar without any especial care, being equally fair and sound as those of this year's growth. Samples were sent in 1873 to the gardens of the Royal Horticultural Society of London, where they received a *first-class certificate*, and have also received many premiums at various Agricultural Fairs in this country. Size, medium to large, growing very fair and smooth. Eyes few and small, nearly even with the surface; shape oval, somewhat flattened; skin reddish, or a deep flesh color; flesh white, fine-grained and very delicate. For the table, they cook either by baking or boiling equal to the very best, and with ordinary boiling they cook through to the center evenly, dry and mealy, and are never hard, hollow, watery, or discolored at the center; flavor unexceptionable. Vine of medium growth; foliage deep green, and very healthy in all respects. The tubers grow compactly in the hill and are easily dug, ripening in about three months from time of planting, though suitable for cooking about two weeks later than the Early Rose, with the same culture. They are very productive, with but few small tubers. Its beautiful appearance, fine quality, extraordinary productiveness and remarkable keeping qualities render it a most valuable variety for the market.

To show the wonderful productiveness of this favorite variety—are among the following from the successful competitors for the first prize offered by us in 1874.

[From H. C. Pearson, Pitcairn, St. Lawrence Co., N. Y.]

I hereby certify that I ordered of you, March 25, 1874, 1 lb. of Brownell's Beauty Potatoes; planted them May 16, and dug them September 28, and from the 1 lb. planted, I raised ten hundred eighteen pounds, (1018 lbs.) The soil was light loam, with some gravel, with a sand and gravel subsoil, not under-drained but having good natural drainage. The land is new, having produced only one crop before. Applied broadcast a two-horse load of barnyard manure three years old, plowing it under 7 inches deep; then went over the ground 3 times with a pulverizing harrow. Placed in each hill before planting, 2 quarts of a compost, composed of 10 bushels decayed manure, 2 bushels of ashes, 4 quarts salt, and 2 lbs. sulphur. Cut the tubers to single eyes, dividing some of the stronger into 3 pieces, making in all 112 pieces, and planted them in rows 3½ feet apart and 3 feet apart in the rows planting one set in each hill and covering them about 3 inches deep. They grew to an enormous size, 491 selected tubers weighing 500 lbs., and 37 fair sized tubers weighing 15 lbs., were dug from one hill. They are the most beautiful potato I ever saw. Have tested their quality by cooking them in various ways and find them dry and mealy, fine-grained, and of a flavor not surpassed by any potato I have ever tried. They matured about September 18.

(Signed,) H. C. PEARSON.
PRICE:—One pound, 60 cents; two pounds, \$1.00, by mail to one address, postpaid. By express or freight, charges to be paid by the purchaser, 1 peck, \$1.00; ¼ bushel, \$1.25; 1 bushel, \$2.25; 1 barrel, \$5.00.



EXTRA EARLY VERMONT.

The superior merits of this variety, first sent out by us in the Spring of 1872, may now be considered as fully established. It has been largely cultivated over every section of the country, the past two seasons, and fully sustains the high character given it by the committee for awarding the premium offered by us in 1873, in the following statement, after examining the reports of the various competitors:

"**The Early Vermont**, as proved by the numerous reports before us, more than sustained its previous reputation. *Nearly all the competitors declare it from one to two weeks earlier than the Early Rose, and many even more.* Its uniform and large size is recognized by every one. Mr. McLeod says: 'There are more than 100 in the amount I raised that would weigh from one to two pounds each;' and Mr. Salter raised one tuber that weighed **THREE POUNDS TWELVE OUNCES.** Its superior cooking and eating qualities are unanimously commended, as well as its compact growth in the hill and its freedom from disease, and with the thousands of cultivators who have grown it alongside the Early Rose, there seems to be no doubt left that in *quality, hardness, earliness and yield, it far surpasses that celebrated variety.*"

A first-class certificate was awarded this variety by the Royal Horticultural Society of London, 1873.

Caution.—In consequence of the great similarity between the **Extra Early Vermont** and the **Early Rose**, many of the latter will doubtless be offered by unprincipled persons as the **Early Vermont.** To avoid imposition, we caution purchasers to beware of itinerant peddlers, and purchase their stock of reliable parties only. We employ no peddlers or agents. Address all orders directly to our house.

PRICE:—By mail, postpaid, one pound, 60 cents; two pounds, \$1.00. By express or freight, charges paid by purchaser, 1 peck, \$0.75; $\frac{1}{2}$ bushel, \$1.25; 1 bushel, \$2.00; 1 barrel, \$4.00.

Tomato "Little Gem" (the earliest variety known).

This new variety was raised by Mr. Pringle, the originator of the Conqueror Tomato, heretofore considered the earliest in cultivation, and is a combined hybrid or cross of that popular variety with two other unnamed early varieties of fine quality. The vines are of medium length, compact growth, and excessively loaded with bright red fruit of medium size, round and of uniform shape, varying from four to six inches in circumference, of superior quality flavor, either raw or cooked. Its bright glossy red color, uniform size, shape, and otherwise beautiful appearance, make a dish of **LITTLE GEMS** almost as tempting as a plate of Strawberries. It may be relied upon to yield several pickings, a week or ten days earlier than any other variety. 15 cts. per packet; half oz. packets, 75 cts.



POTATOES—EXTRA EARLY VERMONT.—Showing Habit of growth in the Hill.



BROWNELL'S BEAUTY.—Showing Habit of growth in the Hill. Page 23.



Compton's Surprise Potatoes.

Showing the growth from a single eye in the grounds of F. Seiler, Verona, New Jersey, who raised 384 lbs. from one lb. of the tubers, with ordinary culture.

Remarkable for its size, quality and productiveness. Its shape is oval-oblong, eyes sunken, brow prominent, skin smooth, color reddish-purple, flesh white; grows to a large size, and is invariably sound to the center. It retains its quality *perfectly* throughout the year, appearing on the table like a ball of flour. Remaining plump and free from sprouts when kept until June and never having that wilted appearance common to early sorts. Thousands have testified that they never ate a better Potato. Its uniform mealiness of grain, combined with the purest flavor and its snowy whiteness of flesh, which is not in the least affected by its blue skin, cannot fail to make it highly valuable.

The following extract from the report of the Committee awarding the premiums offered in 1873, confirms all that was claimed for them by the originator when first introduced.

“Compton's Surprise has received the unanimous verdict for the most prolific Potato cultivated at present. Yields of from twelve to twenty pounds to the hill are reported by the hundreds, and in one instance 28½ pounds were dug from one hill. As a rule, the most prolific varieties are not of the best eating quality, but to this, Compton's Surprise is an exception. Thousands have testified that they never ate a better Potato. Its uniform mealiness of grain, combined with the purest flavor, and its snowy whiteness of flesh, which is not in the least affected by its blue skin, cannot fail to make it highly valuable.

One pound, 60 cents; 3 pounds, \$1.25 by mail, prepaid; by express or freight, charges paid by the purchaser, 1 peck, \$0.75; ½ bushel, \$1.25; 1 bushel, \$2.00; 1 barrel, \$4.00.

Early Paragon.

A new early variety introduced last season, and said to be a few days earlier than the Early Rose. The vines are of dwarf habit, shape oblong, eyes very shallow, skin smooth and general appearance very attractive: productive and of good quality.

By mail, prepaid, one pound, 60 cents; 3 pounds, \$1.25; by express or freight, charges paid by the purchaser, 1 peck, \$1.00; 1 bushel, \$2.25; 1 barrel, \$5.00.

General Collection.

Two pounds of either of the following varieties will be mailed, post-paid, to any address in the United States, upon receipt of \$1.00, or one pound for 60 cents.

Not less than one pound, or more than one variety in one package, will be mailed.

Early Rose.—This was the first of Mr. Bresee's Seedlings, offered by us in January, 1868, and has now become the standard variety for earliness, quality and productiveness. *Per peck, 75 cents; bush., \$2.00; bbl., \$4.00.*

Bresee's Prolific.—Vines of medium height, quite bushy, somewhat spreading. Tubers large, regular in shape, and very smooth, slightly oblong, somewhat flattened. Skin dull white, inclined to be russeted, eyes but little depressed and slightly pinkish, flesh white, cooks quickly, is very mealy and of excellent quality, yield very large, matures about three weeks later than the Early Rose; a valuable variety for field culture. *Per peck, \$1.00; bush., \$2.25; bbl., \$5.00.*

Brownell's Eureka.—This seedling resulted from a seed-ball grown on an Excelsior potato vine, blossoms fertilized with pollen from the White Peachblow. Vines of strong and vigorous growth; tubers of good medium and uniform size; shape elongated oval, somewhat flattened; eyes few, skin white and fair, season second early. It is one of the most productive in cultivation, besides being an excellent keeper; its flesh is exceedingly fine grained, white, and when boiled or baked, mealy and of excellent flavor, cooking through uniformly without fault at center. A silver medal was awarded this variety by the Massachusetts Horticultural Society in 1874.

Climax.—Early; uniformly large; long, cylindrical; skin white; eyes sharp, shallow; flesh white and solid. This has been the most prolific early variety during the past season, and has proved hardy and healthy. *Per peck, \$1.00; bush., \$2.50; bbl., \$6.00.*

Late Rose.—This variety, first offered by us in the Fall of 1871, has been largely cultivated in various parts of the country, and has given universal satisfaction. It ripens two or three weeks later than the Early Rose, and has proved to be much more productive, hardier, healthier, and a better keeper, retaining its good quality till new potatoes come in. *Per peck, 75 cents; bush., \$2.00; bbl., \$4.00.*

Peerless (Bresee's No. 6.)—Skin dull white, occasionally russeted; eyes shallow; oblong, flesh white, mealy, grows to a large size, often weighing from one and a half to two pounds, and enormously productive, frequently producing from 100 to 150 barrels to the acre. Its great beauty, superior quality, and enormous productiveness, place it among the best varieties for general culture. *Per peck, 75 cents; bush., \$2.00; bbl., \$4.00.*

Lapstone Kidney.—Of English origin. Medium early; very long, kidney-shaped; skin very smooth and white; eyes small and entirely flat; flesh white, finely grained, sound and solid, and is not excelled in its qualities for baking or salad. *Per peck, 75 cents; bush., \$2.00; bbl., \$5.00.*

Jackson White.—A northern variety. Medium late; large; irregular, round to longish; skin white and smooth; eyes deep; flesh white, finely grained, and of good table quality; is a good keeper, and very productive in some localities. *Per peck, 75 cents; bush., \$2.00; bbl., \$4.00.*

White Peachblow.—A seedling of the old Peachblow. Very late; medium to large; round; skin white, with bright pink eyes; flesh white, cooks very dry and mealy. This variety has for years been the principal market potato in New York, and has proved a remunerative crop to the producers. *Per peck, 75 cents; bush., \$2.00; bbl., \$4.00.*

Peachblow (Jersey).—A well-known variety, very productive, superior for the table, and one of the best for the market and shipping purposes. *Per peck, 75 cents; bush., \$2.00; bbl., \$4.00.*

Any other varieties not in our list will be furnished at lowest market prices.

A NEW EARLY TOMATO—ACME.

This new and beautiful variety far exceeds anything ever before introduced among the many varieties of this valuable esculent. It has been carefully tested the past season, in many localities, and all agree that it is not only one of the earliest, but the handsomest variety ever introduced. It seems to possess all the good qualities that can be desired in a Tomato, and will eventually supersede many others now in the field. The plants are of a strong and vigorous growth, very productive; fruit of medium size, large enough for any use, form perfect, round, slightly depressed at the ends, very smooth; color a glossy dark red with a sort of purplish tinge; ripens all over and through at the same time; bears continuously until frost; delicious in flavor, has no green core and but few seeds; unequaled for canning, preserving, or as salad; a splendid market variety, and well adapted for culture in the Southern States, for shipping to the North in the early part of the season. *Price, 15 cts. per packet of about 100 seeds; ½ oz. packets, 75 cts. each.*

THE GREAT CENTENNIAL EXHIBIT

OF

500 Named Varieties of Potatoes.

The Largest Collection of Potatoes in the World, for which was Awarded

THE GRAND CENTENNIAL PRIZE MEDAL AND DIPLOMA!

Acme.	Bulkeley's Prince of Wales.	Davis' Seedling.
Adirondack.	Bulkeley's Seedling.	Delmahoy.
Advance.	Busam's Early.	Dickinson's Saybrook.
Agawan.	Busam's Late Prime.	Dover.
Akron.	Byram.	Dover Seedling.
Alaska Blue.	Calico.	Dr. Bretonnean.
Albion.	California.	Duke of Cumberland.
Alexandra.	California (purple).	Dutch Pinkeye.
Alpha.	California Mercer.	Dykeman.
Amazon.	Callao.	Dyright.
American Breadfruit.	Campbell's Late Rose.	Early Blue.
Anderson.	Carmel.	" Cottage.
Audes.	Carpenter.	" Dexter.
Angola.	Carter.	" Dimnick.
Armonk.	Carter's Early Forcing.	" Don.
Armstrong.	Caseoe.	" Durham.
Ashleaf Finke.	Cayuga.	" Golden.
Ashleaf Kidney.	Central City.	" Goodrich.
Ashtop Fluke.	Champion of England.	" Handsworth.
Badger.	Chase.	" Henry.
Baldwin.	Checker.	" Indiana.
Baltimore.	Chenango.	" June.
Banks.	Chenery.	" Kidney.
Barron's Perfection.	Cherry Blow.	" Lilae.
Beaty.	Chester.	" Manly.
Bellaire.	Chili No. II.	" Minnesota.
Ben. Merritt.	Circassian Kidney.	" Mohawk.
Berkshire.	Climax.	" Ohio.
Black Diamond.	Cluster.	" Oneida.
Black Kidney.	Coldstream.	" Paragon.
Black Mercer.	Colebrook.	" Peachblow.
Black Prince.	Collin's Seedling.	" Pearson.
Blanchard.	Colmbus.	" Pinkeye.
Blanche.	Colorado.	" Prince.
Blue Kidney.	Colorado I.	" Purple.
Blue Pinkeye.	Colorado II.	" Queen.
Blue Western.	Compton.	" Racchorse.
Blush.	Compton's Surprise.	" Rose.
Bolton.	Con.	" Russet.
Bonutiful Kidney.	Concord.	" Samaritan.
Bradford.	Conn. Blue.	" Scotch Cottage.
Bradford Seedling.	Conover.	" Shaw.
Breakfast.	Coppermine.	" Snowball.
Bresec's Prolific.	Cornell's Kid.	" Sovereign.
British Queen.	Cornish.	" Stevens.
Brownell's Beauty.	Cowhorn.	" Vietor.
Brownell's Nonesuch.	Cullock.	" Violet.
Brownell's Success.	Cuzco.	" Wendell.
Brownell's Superior.	Dagger.	" White.
Buckeye.	Dana's Seedling.	" York.
Bucyrus.	Davenport Seedling.	Eastport.

East Branch.	Heins.	Massasoit.
Eclipse.	Hemlock.	Mereer.
Economist.	Hickory.	Mereer Seedling.
Egg.	Hinman.	Merino.
Elder.	Holly Hoek	Mexican.
Empire.	Holmes.	Michigan Red.
Empire State.	Hoosier.	Miller.
English Kidney.	Hugh.	Miller's Seedling.
Enreka.	Huntington Seedling.	Minnesota Seedling.
Excelsior.	H. Ward.	Mission White.
Extra Early Rose.	Ice Cream.	Model.
" " Rose Seedling.	Improved Ashleaf Kidney.	Monas Pride
" " Seedling.	Improved Gem.	Monitor.
" " Vermont.	Improved Shaw.	Monon.
" " White.	Iona Red.	Motley.
" Large Seedling.	Irish American.	Mountain Blue.
" Peerless.	" Blue.	Mount June Pinkeye.
Fancy Red.	" Cup.	Multiplier.
Farmer's Delight (red).	" Cup White	Multiply.
Farmer's Delight (white).	" Orange.	Myatt's Prolific Ashleaf.
Favorite.	J. Quinby.	Nansemond.
Fenn's Early Market.	Jackson Seedling.	Nanuet.
Fenn's Early White.	Jackson White.	Napoleon.
Fenn's Perfection.	Jenny Lind.	New Hampshire Seedling.
Ferris' Seedling.	Jersey White.	New Hartford.
Fersey White.	John Bright.	New Kidney.
Field Crop.	Jones' Seedling.	New Nutmeg.
Findlay.	Jug.	New Peachblow.
Fisher.	Kalamazoo.	Noblow.
Fluke.	Kansas.	Nonesuch.
Forest Rose.	Kearsarge.	North Castle.
Forfarshire Red.	Kemyon.	Nova Scotia.
Fortune.	Ketchum's Seedling.	Noyes.
Foster's Late Rose.	Keystone State.	Niggerhead.
Fox Seedling.	Kid.	Niggertoe.
Frankfort.	King of Jacksons.	Nutmeg.
Freeman.	King of Potatoes.	Ohio.
Fremont.	King of Sweden.	Ohio Beauty.
French.	King of the Earlies.	" Chenango.
Galva.	King's Seedling.	" Mercer.
Gardner.	Kinney.	" Red.
Garnet Chili.	Kruger.	" Russet.
Gen.	Lackawanna.	" Victor.
Gen. Grant.	Lady Finger.	Old Fleshcolored.
Genesee Co. King.	Laing's Seedling.	" Kidney.
German Russet.	Land's Prolific.	" Red.
Giant.	Lapstone Kidney.	" White.
Gilman.	Late Pinkeye.	Oneida.
Globe.	Late Rose.	Orange Peel.
Golden Gem.	Lenthereoat.	Orono.
Golden's Seedling.	Ledding's Seedling.	Oscar.
Goodrich Seedling.	Lincoln Red.	Pale-Blush Pinkeye.
Granite State.	London White.	Palmer.
Gray Nonesuch.	Long Island.	Palmynra.
Gray Russet.	Long Pond.	Para.
Great Britain.	Mahopae Seedling.	Paterson's Albert.
Great Western.	Maiden's Blush.	" Blue.
Greenfield.	Maine Mercer.	" Early White.
Guernsey.	Manatan.	" Golden Don.
Hall.	Marchioness of Lorne.	" Queen.
Hamburg.	Marjolin Letard.	" Regent.
Harison.	Massachusetts White.	Patoka.

- | | | |
|--------------------------|------------------------|-------------------------|
| Peachblossom. | Rose Bud. | True Lady Finger. |
| Peerless. | Roso of the West. | Triumph. |
| Penn. Searchwarrant. | Rose Seedling. | Tuberose. |
| Philadelphia. | Rough and Ready. | Tucker. |
| Philbrick's Early White. | Round White. | Union. |
| Pierpont. | Roxbury. | Utica Pinkeye. |
| Pigeon Eye. | Royal Ashleaf Kidney. | Vanderveer. |
| Pink Blow. | Ruby. | Van Tassel Seedling. |
| Pinkeye. | Russet Kidney. | Vermont Beauty. |
| Pinkeye Rustycoat. | Sandy Brown. | " Seedling. |
| Pinkeye Strawberry. | Scotch Blue. | Wampoo. |
| Pink Peachblow. | " Russet. | Ward I. |
| Plymouth Rock. | " White. | " II. |
| Porter's Excelsior. | Sebee. | Weeks' Seedling. |
| Prairie Flower. | Sedilla. | Weston Chief. |
| Prairie Seedling. | Seedling Cuzeo. | " Russet. |
| President. | " Early Goodrich. | West-Fairview Seedling. |
| Prince Albert. | " Mercer. | West Windsor. |
| Prince Arthur. | " Peachblow. | Wheeler's Milky White. |
| Prince of Wales. | " Prolific. | Whig. |
| Purple Kidney. | " Rock. | Whipple's Seedling. |
| " Mercer. | Sharon. | White. |
| " Peerless. | Shaker's Fancy. | " Apple. |
| Putnam. | " Russet. | " Clinton. |
| Quarantine Violette. | Shaw. | " Cownhorn. |
| Quadroon. | Sheridan. | " Eyed Peachblow. |
| Queen of the Earlies. | Sherwood. | " Lily. |
| Queen of the Roses. | Silverskin. | " Mountain. |
| Queen of the West. | Six Weeks. | " Mountain Early. |
| Quinby's Seedling. | Skerry Blue. | " Neshamock. |
| Rand's New Peachblow. | Smith's New Seedling. | " Peachblow. |
| " Red. | Snapdragon. | " Pinkeye. |
| " White. | Snowball. | " Rock. |
| Rasperry-Leaved. | Snowflake. | " Rose (Wainwright) |
| Rathbone. | Soisoto. | " Rose (Young's.) |
| Rattails. | South Bend. | " Russet. |
| Rector of Woodstock. | St. Helene. | " Sprouts. |
| Red Carter. | Starch. | " Uink. |
| " Chmax. | Stanton's Premier Kid. | " Utica. |
| " Emperor. | State of Maine. | Willard. |
| " Fluke. | Stevens. | Williams. |
| " Gem. | Stonehouse. | Wm. R. Prince. |
| " Jacket. | Strawberry. | Willow Tree. |
| " Kidney. | " Gem. | Wilson. |
| " Neshamock. | " Mercer. | Wisconsin Blue. |
| " Orange. | Supper. | " Seedling. |
| " Peachblow. | Sutton's Flourhall. | Wonderful Red. |
| " Streak. | " One Hundred Fold | Woodard. |
| " Uink. | Temple. | Worcester. |
| " Utica. | Tippee canoe. | York Seedling. |
| Ridgefield Seedling. | Titicaea. | Yorkshire Hero. |
| Rochester Seedling. | Toledo. | Young. |
| Rodes. | Topeka. | Young America. |
| Rogmore Rose. | Trip's Garnet. | |

One Tuber of each variety of the entire collection, correctly labeled, carefully packed, expressage or postage prepaid, for **\$100.**

Collections of 250 varieties, one tuber of each, for \$50.00 ; 100 varieties for \$25.00 ; 50 varieties, \$15.00 ; 25 varieties, \$8.00.

—•••—

Collection of Seedling Potatoes.

We have on hand 200 varieties **unnamed Seedlings** raised within the past two years from **Pringle's Hybridized Potato Seed**, which have been selected from upwards of a **thousand varieties** for further trial. Among them are many that are exceedingly promising and well worthy of the attention of Farmers. They will be offered in collections as follows, and mailed postpaid :

200 varieties, 1 tuber each,	-	-	-	-	-	-	-	-	-	\$25 00
100 " 1 " " "	-	-	-	-	-	-	-	-	-	15 00
50 " 1 " " "	-	-	-	-	-	-	-	-	-	8 75
25 " 1 " " "	-	-	-	-	-	-	-	-	-	5 00

N. B. Orders for Potatoes, received during Winter, will be forwarded in Spring as soon as the weather will permit, which is usually about the first of April. They can be forwarded earlier, if desired, at the risk of the purchaser.

POTATOES.

ENGLISH AND FRENCH VARIETIES.

The following varieties, the very best cultivated in Europe, were in nearly all of winning stands at the International Potato Exhibition at the Alexandra Palace, London, for the past two years, and are highly recommended both for their extra fine quality and productiveness. One pound of the following-named varieties will be mailed postpaid for 75 cents. They are also offered in collections of one tuber each, as follows:

Collections of 5 varieties, 1 tuber each, \$1.50. | Collections of 20 varieties, 1 tuber each, \$5.00.
 " " 10 " 1 tuber " 2.75. | " " 40 " 1 tuber " 8.00.

Ashtop Fluke.
 Barrows' Perfection.
 Blanchard.
 Bountiful.
 Coldstream.
 Dawes' Matchless.
 Early Dimnick.
 Early Union.
 Excelsior Kidney.
 Fenn's White Kidney.
 Fenn's Early White.
 Fenn's Early Market.
 Fenn's Perfection.
 Hayes' Superb Kidney.
 Headley's Nonpareil.

Jersey Blue.
 King of Potatoes.
 Mareean.
 Marchioness of Lorne.
 Marjolin Cetaud.
 McKinlay's Lady Webster.
 Model.
 Myatt's Prolific Ashleaf.
 Prince Arthur.
 Prince Teck.
 Prince of Wales Kidney.
 Patterson's Victoria.
 Patterson's "The Queen."
 Porter's Excelsior.
 President.

Quarantaine Violette.
 Rector of Woodstock.
 Red Emperor.
 Red Fluke.
 Rivers' Royal Ashleaf.
 Rognon Rose.
 Sedila.
 Scotch Regent.
 Stanton's Premier.
 St. Helena.
 Sutton's Red Skin Flourball.
 Sutton's Exhibition Kidney.
 Sutton's New Hundredfold Fluke.
 Wonderful Red Kidney.
 Yorkshire Hero.

Pringle's Hybridized Potato Seed.

The extraordinary success which has attended Mr. Pringle in his attempts to improve this valuable esculent—which has resulted in the production of the **Snowflake and Alpha**, and last, though not least, the **Ruby**—has encouraged him to still greater efforts in his favorite pursuit of hybridization, and we have now the pleasure of offering a very choice strain of seed saved by him, which is the product of numerous hybridizations between the above named and many of the best new and old varieties in cultivation, both English and American, and includes every strain, which Mr. Pringle will himself sow the coming Spring. Full directions for sowing accompany each packet. Numerous testimonials have been received during the past season in favor of the superior quality of this seed, as well as of its extraordinary productiveness, 5 to 22 pounds of good sized potatoes having been grown by several parties from a single seed.

Directions for Cultivation.

Sow in pans or shallow boxes, of a mixture of leaf mold and loam, and place under glass or a gentle hot-bed; as the plants appear, they should be partially shaded from the full sunlight; if they begin to crowd, transplant, giving them ample room; when danger from frost is over, set them in the open ground, giving them the full space usually allowed to potatoes. With this treatment they will, with good attention attain to full size and maturity the first year.

The following letters are selected from a number received from our customers, who have given this seed a trial, showing its wonderful productiveness. Samples of the tubers were also sent us, which were marvels of beauty, and will be sure to create a sensation when they become known.

CHARLOTTE, VT.

From Pringle's Hybridized Potato Seed, purchased of you last Spring, I raised from one seed, 45 potatoes, weight seven and one-quarter pounds; from another, 95 potatoes, twenty-two and one-half pounds; from another, 115 potatoes, twenty-seven and one-half pounds.

O. H. ALEXANDER.

PITCAIRN, N. Y., September 28, 1876.

B. K. BLISS & SONS:—The packet of potato seeds I received from you the first of April last, were planted the 9th day of April, in boxes, transplanted the 13th day of May in rich soil; used some bone dust in each hill; the plants were under glass most of the time until June; hoed them often, and they grew fast; watered several times in July and August with liquid manure. The packet contained 29 seeds; 24 grew, and one was destroyed by worms; the largest product from a single hill was 9 5/16 pounds of handsome, smooth, red potatoes.

Yours truly,

H. C. PEARSON.

Price 25 cents per packet; 5 packets \$1.00.

Report of Committee

For awarding PREMIUMS offered by B. K. BLISS & SONS to growers of the largest quantity of POTATOES, from One Pound of Seed.

• • •

After becoming fully satisfied of the superiority of the new varieties of Potatoes, introduced by us during the past five years, we have offered, from year to year, premiums to those who should grow the largest quantity from one pound of seed, of the different varieties with ordinary farm culture, and have paid out within that time upwards of **Three Thousand Dollars** to the successful competitors. The premiums offered for each variety were as follows:

\$250.00 in Premiums.

For the largest quantity of Potatoes grown from one pound of seed, - - - - -	\$100 00
For the second largest, - - - - -	50 00
For the third largest, - - - - -	40 00
For the fourth largest, - - - - -	30 00
For the fifth largest, - - - - -	20 00
For the sixth largest, - - - - -	10 00

Competitors for the prizes were required to give the date on which they gave their order for the potatoes, date of planting, date of digging, with a written statement of their mode of culture, characteristics of the soil—whether clay, alluvial, sandy or loam—nature of the subsoil, whether underdrained or not; also the kind and quantity of fertilizers used, how and when applied, the number of hills, and distance apart each way, with the weight of the crop when dug, and the number of square feet occupied by the crop, which must be witnessed and sworn to before a justice of the peace, notary, or any other one competent to administer the oath.

The awards were made by a committee composed of the following well-known gentlemen in the agricultural community: PROF. GEORGE THURBER, editor American Agriculturist, DR. F. M. HEXAMER, and P. T. QUINN, Horticulturists. They were published in the leading Agricultural Papers, and a copy mailed to each competitor. The prizes were awarded the first of January, of each year.

N. B.—To prevent misunderstanding, we wished it distinctly understood that no prizes would be awarded unless the above requirements were complied with in every particular. Competitors for premiums were placed under no restrictions, as to their mode of culture, excepting that *they must not be grown from slips or forced by artificial heat*, our object being to ascertain their respective merits with such culture as is usually given to crops in a well-managed vegetable garden or farm.

REPORT.

The number of competitors, who, tempted by these liberal prizes, tried their skill and industry in producing a premium crop, was naturally very large. Thousands of reports—representing every Potato-producing State and Territory of the United States, and even many Foreign Countries—nearly all from practical farmers, and many from men who have made Potato-growing their life's work, and may well be called the Champion Potato growers of the world. The practical information embodied in these Reports must naturally be of great value to every tiller of the soil, and at the request of the committee I have endeavored to condense the leading features of the modes and conditions under which these immense crops were grown.

YIELD FROM ONE POUND OF

Compton's Surprise in 1873.	Early Vermont in 1873.	Early Vermont in 1874.	Compton's Surprise in 1874.	Brownell's Beauty in 1874.	Snowflake in 1877.	Eureka in 1877.	Alpha in 1876.	Ruby in 1876.
511	607	708	900	1,048	1,417	1,666	1,707	1,982
450	437	698	874	811	1,304	1,403	1,665	1,694
390	393	690	832	782	1,125	1,149	1,535	1,576
386	380	674	811	719	1,090	1,145	1,511	1,571
		629	684	720	1,089	1,087	1,426	1,534
		615	588	696	1,069	1,066	1,280	1,353

The gradual increase in the yield, from year to year, is at once apparent, and when we consider that several of the largest yields in 1876, were grown by the same persons to whom the highest awards were given at the first trial, and when we consider that this increase in the yield is not due to the greater productiveness of the newer varieties, but solely to the increasing skill of the competitors, and the better management given their crops, we are forced to appreciate the great amount of good which these competitive trials have produced. I am aware that the largest yield from one pound is not always in proportion to the largest yield per acre; nor do these immense yields, produced by the lavish application of manures, regardless of expense, give a criterion of the profits of the crops, after deducting the premisses. Many growers who have produced less than one thousand pounds from one, may have obtained larger profits per acre, than the successful competitors yet no one who has grown a thousand, or even five hundred pounds from one, can have done so without learning something which will be of value to him in growing Potatoes as a field crop, nor can any one read an intelligent report of his methods of cultivation, without benefit.

Soil.—Many of the premium crops were grown on new lands, and on soil of almost unequalled quality. The largest yield was produced on "a mixture of sand and clay, very rich in vegetable matter to the depth of *eighteen feet*, and underlying this is a gravelly subsoil. For three years the ground was used as a stock-yard, the straw being left on the ground to rot and be burned." The second largest product, was grown "on a very rich, sandy loam, rich in decayed vegetable matter to the depth of between two and three feet, and lying upon a compact formation known as hard-pau, which has never been underdrained." Another competitor describes his soil as "black loam, four feet deep, on the bank of a creek, and it has been used as a cattle-yard for ten years." Another, "as vegetable mould and sandy loam, three feet deep, never cultivated before." Most crops however, were raised on deep, alluvial lands, underlaid with gravel; others on light loam with clay or gravelly subsoil, and in a few cases on heavy clay highly manured. In but very few instances was the land artificially underdrained, which seems to be a noteworthy fact, as most writers consider drained soil indispensable for the production of good crops.

Manures.—The large quantities of Fertilizers used by most competitors is something astonishing, and may well serve to disprove the general belief that heavy manuring is injurious to potatoes. Mr. Pearson added to an already very rich soil, about 60 two-horse loads of manure, nearly 200 bushels of wood-ashes, and 24 bushels of lime, per acre, together with bone-dust and other fertilizers in smaller quantities. Mr. Rose, after covering his land three inches thick with rotten barn-yard manure, and three bushels of wood-ashes per square rod, applied in addition, a large shovelful of rotted hen manure, and two handfules of ashes to each hill, besides several surface dressings with other fertilizers. But all this must appear but a small attempt at enriching the land, to our Scotch friend, Mr. Robertson, who would not entrust his seed to a black sandy loam, four feet deep, underdrained and trenched, to probably the entire depth, before spading under a coat of *five inches* of well rotten cow-dung, and applying afterwards to the hills three cart-loads of wood-ashes, two of sheep droppings, and several other fertilizers.

About the value of wood ashes and especially in mixture with hen manure and plaster, there seems to be no doubt left; they were used by a large majority and may be considered the Special Potato Fertilizer. Even when applied in very large quantities they have produced no injurious effect. In a few cases, as much as one pint of ashes has been applied to each hill as top-dressing, and in one instance the sets were actually planted in and covered with ashes. Plaster, lime and salt have likewise been extensively used and with great advantage. Strong nitrogenous manures have generally been considered as detrimental to the potato, but here we find that many successful growers have used large quantities of Blood, Fish and Peruvian Guano; hen manure as well as barn-yard manure, at the rate of fifty loads per acre, to the greatest advantage and without producing diseased potatoes.

Planting.—The time of planting in nearly all cases was between the 10th and 26th of May, and in the majority of these before the 15th. That single eyes and eyelets will, with good care, produce large crops, has been sufficiently proved. All the large yields were grown from very small sets. In some cases, single eyes were divided into ten pieces, and in one instance two hundred and ninety (290) sets were made from one pound, nearly all of which grew well. The sets, with few exceptions, were planted singly, yet we find a product of nine hundred and seventy (970) pounds raised from fifty-two (52) hills, two sets to each, nearly nineteen (19) pounds per hill, and six hundred and seventy-seven (677) bushels per acre. Whether this large yield is due only to the very favorable soil they grew in—a rich black loam, formerly used as a hog yard—and the immense quantities of ashes applied in the hills and as top-dressing—one peck to the hill—or to the two-set system, does not appear. It is to be regretted that a part of the plat was not planted with one set to the hill, and the products weighed separately. These practical tests, of the feasibility of raising large crops from small sets, become of much importance in seasons of scarcity of seed

potatoes. For it is shown here that, even without carrying the division of the eyes to extremes, nine-tenths of the seed may be saved.

A comparison of the distances between the hills with the average yield per acre gives a most interesting and valuable table, as follows:

The sets planted at a distance of

2 x 3 feet gave a yield of 378 bushels per acre.	3 x 4 feet gave a yield of 372 bushels per acre.
2 x 4 " " " 462 " " "	3½ x 4 " " " 342 " " "
3 x 3 " " " 651 " " "	4 x 4 " " " 332 " " "
3 x 3½ " " " 441 " " "	4 x 8 " " " 88 " " "

The large number of data of which the above figures form an average, give these statistics a special value. It will be seen that although the greatest yields from one pound grew from hills four feet apart, the largest crops per acre were raised at distances of three feet each way, and that as the distances between the hills are increased or decreased, the yield diminishes in regular proportion. In the first case, there remains wasted ground which is not reached by the roots of the plants, and in the latter, the roots are so crowded that they cannot obtain all the nourishment they are capable of consuming.

Cultivation.—It will be hardly necessary to state that in many cases the cultivation and care these growing potatoes received were unremitting and indefatigable. In no case were weeds to be seen in the patch; some of the plantations received semi-weekly hoeings, and all were kept scrupulously clean, and the soil loose and mellow. Yet many large crops were grown with but ordinary care. Mr. J. I. Salter swears "that the cultivation, manuring, etc., was the same he has given his general crop of potatoes for the last five years, and in no respect had there been extra care and labor bestowed." Mr. H. C. Pearson certifies "that these potatoes were grown with the most ordinary farm culture, being hoed only twice and receiving no extra treatment in any respect."

The mode of planting and cultivating with a larger number of the best cultivators consists in crossing their fields with furrows six and more inches deep. The sets are dropped at the crossings and immediately covered with about two inches of soil or compost. The vines as they grow are hilled up gradually and frequently to a final height of twelve to eighteen inches. Then large, broad hills are made, using all the soil between the rows.

Irrigation.—The fact that the largest yields were produced in the Eastern States, in regions which suffered from one of the severest droughts known, where the general crops failed almost entirely, induced us to investigate more closely the manner in which these premium potatoes were grown. As the result we find that all these competitors attribute their success to the judicious application of water during the dry season. One of the competitors had arranged an ingenious system of irrigation, by inserting six inches from each hill, two inch drain tiles, six inches deep, and filling these with water, twice a week, during the dry weather. In reply to a letter he writes: "I attribute my success the past season to the fertilizers applied, but more especially to the mode of applying water. I find that, even without manure, this plan gives great results. Give me drouth and this means of applying water, and I can grow a much larger crop than with the most favorable weather and no watering. This I have proved by two years' experience." Many may not be so situated that they can apply as complete a system of irrigation, but thousands of farmers have runing through their lands, brooks and streams which might, without much cost, be made to furnish nutriment to their parching crops. One acre of potatoes would in many cases pay for more than the entire expense of a permanent system of irrigation. The information about this important subject, derived from the accompanying reports, cannot but prove of the greatest value to all cultivators of the soil.

Although these short extracts, from material sufficient for a large volume, can do but little justice to the importance of the subject, yet they may serve as an outline of the important and valuable information which could in no other way be obtained.

Trusting that the increasing interest in potato culture, stimulated largely by these competitive trials, may be still more productive of information and progress in this important branch of Agriculture,

I am yours respectfully,

F. M. HEXAMER, *Secretary.*

Extracts from the Report of Mr. Nardy, delegate from the French Government to the Centennial Exhibition at Philadelphia in 1876.

"The collection exhibited by the house of Messrs. B. K. Bliss & Sons of New York, is the most remarkable of its kind in the Exposition. The varieties of Potatoes in their exhibit, and the evident care and attention bestowed upon their collection, manifest constant study on the part of the exhibitors. In our opinion theirs was the most complete display of this valuable esu-

lent in the exhibition. To this remark we desire to add a few observations. We sincerely hope that the influence of this horticultural exhibition at Philadelphia will long be beneficially felt among farmers and growers of plants and vegetables. That were in reality all the fruits and vegetables which we to-day enjoy, when Asia, Africa and America sent them to us in their wild state? unfit for eating! Now, after having been cultivated and improved by the gardeners, they have become succulent and nourishing, and pleasant to the taste. By patience and careful cultivation, horticulture has been enabled to transform the type of many species of vegetation into a great number of varieties. Let me take as an example of this, the collection of Potatoes exhibited by these gentlemen which consists of 500 varieties, and we can thus appreciate the patience and researches which must have been necessary to obtain so many useful varieties. The service rendered to the farmers of the United States by the house of B. K. Bliss & Sons is so evident, that we do not hesitate to call attention to them, for agriculture in general derives an immense benefit from these great improvements in useful vegetables, which are mainly due to their initiative. The cultivation of the Potato takes an important rank in the agricultural productions of this country. As a source of revenue, it comes immediately after grain and before tobacco. It is now many years since the senior partner, having a special aptitude for horticulture, devoted himself to that study. In 1815 he founded his first establishment, and year by year, by steady application, and especially by his energy in collecting the rarest seeds and best varieties of useful vegetables from all countries, the reputation of the house constantly increased. He was the first to introduce the Early Rose Potato, which he did in 1867, and his establishment then acquired the important position which it now enjoys. Animated by the spirit of the amateur he has not only sought out varieties in this product, but has endeavored in every way to render a benefit to agriculture by inducing the farmer to improve the cultivation of this esculent.

But if we were to dwell longer on this question, we should be carried too far from our object, which is to call the attention of our French Government to the importance of this house who were so highly distinguished by the jury at Philadelphia, and close this report by expressing our desire that they will send some of their rare specimens to our Paris Exhibition of 1878."



Reports of Three of the Successful Competitors

FOR THE PREMIUMS OFFERED BY US IN 1876.

How to Grow Twenty Bushels and Upwards of Potatoes from One Pound of the Seed.

The following reports have been properly witnessed and sworn to before a Justice of the Peace—in their respective residences—they have also been carefully examined by the committee and found to conform to the rules prescribed in our offer for the Premiums.

(From H. C. Pearson.)

PITCAIRN, N. Y. October 4, 1876.

MESSRS. B. K. BLISS & SONS:—I ordered of you April 1st, 1876, one pound each of **Ruby** and **Alpha** Potatoes, and planted them May 10th, 1876, and dug them September 28th. The **Ruby** produced 1,982 pounds and the **Alpha** produced 1,707 3-4 pounds. The soil was light loam with some gravel with a sand and gravel subsoil not underdrained. The soil was very rich, and its fertility was increased in the Fall of 1875, by spreading broadcast fifty-two horse loads of well rotted manure, three years old, and 150 bushels of ashes per acre, and plowed it under about eight inches deep; plowed and harrowed until perfectly pulverized eight inches deep last May. Placed in each hill before planting, two quarts of compost, composed of thirty bushels of decayed manure, five bushels of ashes, three bushels slacked lime eight quarts salt, and four pounds of sulphur; the tubers were cut, some of the eyes divided into as many as seven and eight parts, planted one set in each hill, and covered them about three inches deep with soil mixed with some bone dust, making in all 155 hills of the **Ruby**, 210 of the **Alpha**; they were planted three and one-half feet apart each way, the number of square feet occupied by **Ruby**, was 2,361 feet, the number occupied by **Alpha**, was 2,516 feet. I hoed them three times, making very high, broad hills, watered them several times during July and August, with liquid manure, dug from one hill of **Rubys**, forty-three potatoes, weight 17 1-2 pounds; used no slips or artificial heat. H. C. PEARSON.

(From J. I. Salter.)

ST. CLOUD, MINN., October 24, 1876.

MESSRS. B. K. BLISS & SONS:—On the 7th of April, 1876, I ordered of you, one pound each of the "**Ruby**" and "**Alpha**" Potatoes. Planted the **Ruby** on the 31st day of May. Prepared the ground as follows: I spread as evenly as I could on the sward, about equal parts of hen and barn-yard manure, at the rate of five heaping garden wheelbarrowfuls to the square rod,

before plowing, then plowed, turning a furrow ten inches deep and ten inches wide. I then planted my sets, after dividing every cluster of eyes into from three to twelve pieces, in rows four feet apart and as near as convenient, three and one-half feet apart in the rows, dropping but one piece in a place. I covered from two to three or four inches with loose earth. From the pound I made two hundred and ninety (290) sets, the space planted was sixty by seventy feet, (4,200 square feet); hoed but once, when the plants were about ten inches high, making a broad and continuous ridge, flat on the top, and a little depressed in the center; covered the vines all but a very little of the top. Before hoeing, I applied to each plant a large shovelful of fermenting hen manure; it was so hot that it would turn the leaves of plants black, and give them the appearance of being sealded, in less than a minute; I would manure three or four plants, and then cover as quickly as possible. The plants, in a day or two, began to grow and retained a dark green, almost black color until they were killed by the frost. I did nothing more in the way of cultivation.

On the 19th of August, I dug one hill of the "Ruby," and had ten and one-half (10½) pounds; on the 9th of October, I dug of the Ruby eight hundred and twelve (812) pounds; on the 10th of October, I finished digging the Ruby, digging eight hundred and seventy-one and one-half (871½) pounds, making a total of sixteen hundred and ninety-four (1,694) pounds. The "Alpha" was planted the same distance apart each way, had the same cultivation, manuring, etc., as the Ruby, except about fifty hills, that I covered the sets about four inches deep with coarse, unfermented horse manure, and applied no other manure afterward. I covered this manure with earth, the same as I covered the hen manure. I did not get as many large potatoes from these so treated, but about the same in weight per hill; made two hundred and fifty sets. The ground occupied by the Alpha was the same as the Ruby, sixty feet by seventy (4,270 square feet). I also planted two rows of this Potato in hills four feet apart each way. The land in both cases, a black sandy loam, rich in decomposed vegetable matter, usually about two feet deep, under which is hard pan. On the 28th of August, I dug two hills of the Alpha, and had 15 3-4 pounds; on the 12th of October, I dug 1,010 pounds of the Alpha, and on the 13th of October, 640 pounds, making altogether 1,665 3-4 pounds.

These amounts I dug, were grown alone from the one pound each of the above named varieties, purchased from you as I stated above, and without any sprouting or any other means being used than as stated. The land was not drained in any way. J. I. SALTER.

(From Peter Robertson.)

THE GARDENS—HARTRIGGE HOUSE, JEDBURGH,

ROXBOROUGH, SCOTLAND, October 4, 1876.

Messrs. B. K. BLISS & SONS:—Gentlemen. I beg to forward to you my report. I ordered of you, March 21, 1876, one pound each of Alpha and Ruby Potatoes. I planted them on April 21, 1876; they were dug October 2 and 3, 1876. The Ruby produced 1,534 pounds, and the Alpha produced 1,246 pounds. The soil is a mixture of black, sandy loam, four feet deep, with a red clay subsoil and underdrained; it has been lawns or short grass for many years; it was trenched on or about December, 1875, and remained in Winter fir until about April 10, 1876, it was enriched by about five inches of well-rotted cow dung and gas lime rubbish; the ground was then dug in the usual way, care being taken to mix and make it as fine as possible; the rows were six feet apart, and the sets were planted three feet apart, a mixture of about three cart-loads of wood-ashes, two cart-loads of sheep droppings, one lime, two hundred weight salt, all mixed. The tubers were cut and planted in the presence of Mr. Kerr and others. Some of the eyes divided into seven parts; they were very small. Planted one set three feet apart each way; under each, three spadefuls of the above mixture was pulverized three inches deep with the soil, and each set planted one inch deep; they all grew; at first they looked very weak, but gained strength something extraordinary; they were kept free of weeds, and the soil drawn to them as they grew; they were two inches high on May 29, 1876; they got nothing but kept free of weeds, and the soil drawn to them as they grew, until the tubers began to swell; they were then watered with a mixture of sheep droppings and hen manure a few times until they were lifted; there were 176 sets of Ruby, 164 Alpha, single eyes and very small; from one set single eye I had twenty-one pounds; many of the tubers were two and three pounds each. I may also say that I had some very fine, early Cauliflowers and Early Cabbage on the same ground; they were planted between the rows of the Potatoes, and cut before the Potato vines got up to cover the space between; the ground they occupied was forty-six feet by seventy feet, and would be about 3,220 square feet for each variety. Many of the vines remained green until they were lifted. The season was generally good for Potato culture; no forcing process was used whatever.

PETER ROBERTSON.

The Potato Crop.

A Paper Read at the American Institute Farmers' Club.

By Conrad Wilson.

THE annual yield of potatoes in the United States, according to the returns of the last census, was nearly one hundred and fifty million bushels. In view of the increasing attention lately given to this crop, it will doubtless show, at the close of the present decade, a large increase in the amount of the yield, as well as in the number and improvement of varieties. In fact, it may safely be assumed that, in spite of the fears of many in regard to the ravages of the beetle, this crop will still show a yield, at the next census, of over two hundred million bushels.

By its valuable qualities, and its large consumption, the potato ranks in nearly all countries as a leading food staple. It is therefore in every view important, and may always be discussed with interest and profit. There are, of course, some other crops that are grown on a larger scale, and some that foot up a larger yearly aggregate. Yet no product of husbandry is more variously useful, or more generally raised by farmers, and none, if we except wheat, more universally consumed by the people.

A few examples illustrating the rate of yield and the cost per bushel for this crop will tend to show what possibilities belong to it, and thus perhaps stimulate the average farmer to aim at higher results.

Some experiments in potato culture were reported in January of last year, of which the following are a part of the results:

H. C. Pearson, of Pitcairn, St. Lawrence Co., N. Y., produced from one pound of seed, 1,982 pounds of potatoes, of the Ruby variety. He also raised the same year, from one pound of seed, 1,707 pounds of the Alpha variety. J. I. Salter, of St. Cloud, Minn., got the same year, from one pound of seed, 1,694 pounds of the Ruby, and 1,665 pounds of the Alpha; while Alfred Rose, of Penn Yan, N. Y., got 1,576 pounds, and P. L. Wood, of Ill., 1,571 pounds, each from a pound of seed of the Ruby variety. The year previous J. L. Perkins, of Little Sioux, Iowa, obtained 1,666 pounds from one pound of the Eureka variety.

These amazing products were some of the results obtained by farmers competing for the premiums offered by B. K. Bliss & Sons of New York.

The inducements thus held out by this enterprising firm has led to valuable consequences, and the competition has developed the prolific quality of this vegetable to a degree that surpasses all previous conception.

But while duly crediting the liberality and public spirit of B. K. Bliss & Sons, and the grand success of the winning parties, we cannot help regretting that the most important question in potato culture was not included among the offers. It is easy to see that

The Cost of Production

is the underlying question which measures the importance of all the others. Yet this does not at all diminish the value of the test made by Mr. Bliss. The Rubies raised by Mr. Pearson from a pound of seed fell short of a ton by only 18 pounds. This fact has arrested the attention of potato growers as well as consumers in both hemispheres, and confers merited distinction on both Pearson and Bliss; and certainly the tendency of such facts is to help forward the solution of the still greater problem of cost. The man who gets the bottom figures in this problem, though he may not "double the blades of grass," will more than double the product of potatoes.

Since these prizes were awarded I have received statements of cost from several competitors, and hope still to hear from others.

Alfred Rose has reported his cost, in one competition, at less than 15 cents per bushel, and J. L. Perkins about the same. J. I. Salter obtained a yield in one case at about 20 cents per bushel. His latest results I have not learned. Henry V. Rose and M. M. Rose have each reported a cost slightly above the figures obtained by J. L. Perkins and the elder Mr. Rose. For the previous year, Alfred Rose reported his cost at \$55 per acre, and 9½ cents per bushel. Mr. Perkins has also reported a later crop, since the competition, in which he makes the cost, without manure, \$5.40 per acre, and 5½ cents per bushel.

Various other farmers, since the Bliss competition, have reported to me their yield and cost for potatoes. Lyman Alexander, of Minnesota, got 400 bushels per acre of the Peerless variety, at a cost of about 7 cents per bushel.

Azro Smith, of Rock Bluffs, Nebraska, raised a crop of Extra Early Vermont, of which the yield was 824 bushels per acre, and the cost, as he made it, $4\frac{1}{2}$ cents per bushel. But the cost in this instance, as also in the two previous cases, requires some correction. Every crop should be charged for interest on the land, and when no manure is used, a charge should also be made for exhaustion of soil. This would probably add 2 or 3 cents per bushel to the cost of the three last named crops.

C. C. Holton, of Rochester, N. Y., has reported potatoes at 400 bushels per acre, and at a cost of $12\frac{1}{2}$ cents per bushel. A crop of 600 bushels per acre has been reported by one of the editors of the *Agriculturist*, 640 bushels by C. L. Bragdon, and 700 bushels by P. F. Rice of Polo, Ill. In one of the competitions for the Bliss prizes, Mr. Perkins obtained 376 bushels per acre, and Alfred Rose got 600 bushels of Brownells and 746 bushels of Eurekas.

Let us now take another view of the potato crop that may perhaps shed some light on its utility and value as a food staple.

Feeding Value of Potatoes.

Like Indian corn, this vegetable is everywhere extensively utilized as an article of food, both for the human family and for nearly all domestic animals. Hence it is clear that the nutritive properties of the potato are a question of no little importance to the farmer, and one on which the light of further experience is still greatly needed.

How to convert potatoes, with the best economy, and with the largest profit, into other forms of food, into milk, butter and meat, is a problem not yet fully solved. Various estimates have been made by practical men as to the effective value of the potato in the production of beef, mutton and milk; and though opinions still differ, the proportion of other food staples that potatoes are capable of yielding is nearly indicated in the following statement:

A bushel of potatoes when judiciously fed to animals of a good breed will produce:

Of Beef, - - - - -	from 2 to 3 pounds.
" Mutton, - - - - -	" 4 to 5 "
" Pork, - - - - -	" 4 to 5 "
" Milk, - - - - -	" 35 to 40 "
" Butter, - - - - -	" 2 to $2\frac{1}{2}$ "

It is of course not supposed that potatoes are fed *exclusively* in producing these results, but in a suitable combination with other kinds of feed.

Now the practical value of the above table depends materially upon the amount of potatoes produced from an acre. The average yield of the crop for the whole country is probably not over 100 bushels per acre. And yet it will be seen, from the examples above reported, that over 800 bushels have been produced, and even that is not by any means the final limit of yield. Considering the recent progress of potato culture, and conceding the fact that 1,000 bushels per acre have been occasionally achieved by brilliant farmers, it seems hardly credible that the average yield for the United States is only 100 bushels per acre. What shall we say then? Shall we distrust the return of the last census? or shall we rather credit some progress to the eight intervening years, and claim that the average is now very much better. One thing at last seems clear. If the progress of the next two or three years shall correspond to the same period of the past, it may safely be assumed that the average yield of potatoes for the coming decade will not be less than 200 bushels per acre. I venture therefore to assume this figure as the coming average, and if any of our twenty million farmers are disposed to quarrel with me for claiming this yield, my best revenge will be to prove the possibility and then to tell them how it may be done.

If now we apply to this estimate the figures of the above table, we shall discover

What an Acre of Potatoes Means,

when expressed in the form of other food staples. We shall find that it is measured

In Beef, - - - - -	by 400 to 500 pounds.
" Pork, - - - - -	" 800 to 1,000 "
" Mutton, - - - - -	" 800 to 1,000 "
" Milk, - - - - -	" 7,000 pounds, or over.
" Bread, - - - - -	" 4,000 "
" Butter, - - - - -	400 "

Again, as I have already shown that the total potato product of this country is nearly certain to reach 200 million bushels as the average for the next decade, it will be seen that if the above figures are extended so as to meet this case of the total product, then it will be found that

The Annual Potato Crop of the United States


is equivalent

In Beef, - - - - -	to 400,000,000 pounds.
" Pork, - - - - -	" 800,000,000 "
" Mutton, - - - - -	" 800,000,000 "
" Milk, - - - - -	" 7,000,000,000 "
" Bread, - - - - -	" 4,000,000,000 "
" Butter, - - - - -	" 400,000,000 "

Let us now take another view of this subject. If Mr. Pearson had taken his crop of Ruby potatoes that were produced from a pound of seed, and by a right mode of feeding, had converted them into butter, according to the above estimate, the outcome of his experiment would have been about sixty pounds of butter in the Fall, as the legitimate result of one pound of potatoes planted in the Spring. Or if he had converted the potatoes into mutton, the result would have been 120 pounds.

Again, when Azro Smith harvested over 800 bushels of potatoes from one acre, if he had fed that crop to a good breed of cattle, it would have produced, according to the above estimate, at the rate of two pounds of beef for each bushel of potatoes, making a total of 1,600 pounds of beef, as the product of one acre, or if, instead of beef, he had converted the crop into milk, the result would have been over 20,000 pounds.

These results, of course, depend in part on the above estimate for the feeding value of potatoes. On this point there is room for some difference of opinion. But it would be easy to show that when potatoes are combined with other well-selected elements, and fed in the right proportion to animals of good breed, and good capacity, the estimate given is not far out of the way. But to place the matter beyond any question, if we reduce the estimate by 50 per cent., it would still be possible for Pearson to plant his pound of potatoes in the Spring, and harvest in the Fall either 30 pounds of butter, or 60 pounds of mutton; and Azro Smith would still be able to show from his acre of potatoes, that an acre of beef is equivalent to 800 pounds, and an acre of milk to 10,000 pounds.



EGYPTIAN, EAST INDIAN OR PEARL MILLET.—(*Pencillaria spicata*.)

This new Millet is quite distinct from all other species, and is without doubt, destined to take a place in the front rank of valuable forage plants. Sown in light, sandy soil, the plants at first appear feeble, resembling broom-corn; but when a few inches above ground they begin to tiller, and new shoots appear very rapidly from the original root, until they number a half dozen to a dozen or more. The stems at first are nearly prostrate, but when about two feet long they begin to assume an upright position, reaching a height of eight to ten feet, not differing in color or substance from our common Indian corn. In fact, it is one of the most "leafy" plants we have ever met in the great family to which it belongs. When the stems have reached nearly their full height, the seed or flower spikes appear at the summit. As soon as the first or principal flower spike appears, the stems throw out lateral branches from every joint, these in turn producing leaves and flower spikes. When cultivated for fodder, the seed should be dropped in drills, and given plenty of room on account of the peculiar habit of tillering; and, judging from our brief experience with it, we think a quart, or at most two quarts, of seed would be abundant for an acre. The fodder is in the best condition for cutting and curing when the stalks are five or six feet high; but, if used for sowing, it might be cut earlier or later, at the convenience of the cultivator; the stumps, sprouting and throwing up a new growth, continue to grow until killed by frosts. Cows, horses and other farm stock are exceedingly fond of this kind of millet, eating it with as much avidity and apparent relish as they do the green leaves of Indian corn; and we are inclined to think that it is fully as nutritious. The seed will not mature in the Northern States. *Price, 25 cts. per packet; 5 packets, \$1.00.*

THE CHUFA, OR EARTH ALMOND.—(*Cyperus esulentus*.)

This plant is extensively grown in the Southern States, and for sheep, hog and poultry feed, it is claimed to be one of the most profitable crops grown. Chufas may be planted in rows two and a half or three feet apart, and two tubers should be dropped twenty inches or two feet apart in the row, and covered two inches deep. In this way, about a half bushel is sufficient for an acre. In the South they are planted during February and March. North they should be planted when the ground is warm; about the same time as corn, or a little earlier.

Price per packet 15 cts.; per qt., 40 cts.; [by mail 60 cts.]; per peck, \$2.50; ½ bush., \$4.50; bush. \$8.00.

For a more detailed description of the above and mode of culture, see our circular of **Novelties for the Farm and Garden**—mailed free to all.

New Sweet Potato.

EARLY PEABODY.

This variety, first introduced by Mr. C. A. Peabody of Georgia, in the Spring of 1877, has proved a valuable acquisition, and occupies the same position towards other varieties of sweet potatoes that the Early Rose does among our common varieties. At the South they have been ready for the table six weeks from the time they appeared above ground. The following report from the editor of the *American Agriculturist* shows how well they have succeeded at the North:

"We started the potatoes in a hot-bed, and treated the slips, or plants, as we have heretofore treated others. The season was unusually wet with us, and the vines grew luxuriantly. With the ordinary sweet potatoes, we think it fortunate if we can steal a few from the rows two or three weeks before the vines are stopped by the frost, when the whole must be dug. We had potatoes abundantly large for the table by August 10th, but we did not have them on the table as they were not fit to eat. So abundant had been the rains, that while the vines gave us a plenty of potatoes, they forgot all about the *sweet*, and they were simply useless, as probably any other variety would have been under the same conditions. After this trial we let the crop alone until frost in early October, when it was dug. There were three rows across the garden, each 180 feet long, and they gave, without any unusual culture, 9 barrels of Early Peabody. And such potatoes for size! Three of them, selected, of course, weighed 12 pounds. We are now having them on the table, and while they are not quite so sweet as those brought from Virginia and Delaware, they are sweet enough to suit the writer's taste, and of a most excellent flavor and texture. Our land is very light and suffered so much from drouth in previous seasons, and especially the last, that a steam pump was put in to enable us to water the garden from the river, but so unlike to the average seasons was that just past, that we have not needed to apply a gallon of water, the rains giving us more than enough. From our experience, under these conditions, we have no doubt that in an ordinarily dry summer, the Early Peabody will not only prove early, but good and productive, and it is especially valuable because, on account of its early maturing, it will allow the cultivation of the sweet potato in localities much further North than has heretofore been possible. From our experience this season, adverse as it was in some respects, we feel that we speak within bounds when we say that the Early Peabody will prove to be at least a month earlier than any other variety—and we have cultivated all that are grown in the Northern States."

Different Methods of Propagating.

The usual method is to place the potatoes in a hot-bed, and cover them with a few inches of fine soil; the buds will soon start and form shoots; these, as they grow, make roots of their own, and when they are 8 inches to a foot long, and well rooted, the most forward ones are removed, and the others not yet large enough are allowed to grow on. It is customary to split the larger potatoes lengthwise, and lay them flatwise down in the bed. After the shoots appear, they should have the same care as other hot-bed plants, to prevent them growing weak and drawn up. These shoots, known in the Southern States as "draws" and "slips," are at the North called "sets" and "plants." Do not set the plants until settled warm weather; then set them well down to the first leaf, 15 inches apart in the ridge. There will be below the leaf some joints without leaves. Should a frost nip the top, then a new shoot will start from one of the barren joints. The sides of the ridges and the spaces between them must be kept clear of weeds until the vines cover the ground; move the vines once a week or so at the North to prevent them from taking root; this is quickly done by means of a rake handle or other stick. Do not break or cut the vines. At the South the season is long enough to let them take root at the joints, as they will make potatoes then, but at the North we wish to prevent their taking root, and to throw all the strength of the vines into the potatoes that form on the original plant. Be careful in working among the plants not to hoe too deeply, as the earliest potatoes lie immediately beneath the surface. Upon digging, the potatoes they should be carefully handled and put away in boxes or barrels with alternate layers of leaves or cut straw, in a warm dry place to keep.

The Land and How to Cultivate Them.

The sweet potato may be cultivated on any soil but a heavy one. We have seen fine crops on white sand that would blow into drifts with every heavy wind. Mr. Peabody says that he has never seen a sand-hill so poor but what, with judicious manuring, it would bring a good crop; and that any soil which will produce a fair crop of corn, will produce one of sweet potatoes, provided it is a sandy loam. He says that he has known a crop in his county of 800 bushels of sweet potatoes to the acre, upon land that would yield 50 bushels of corn, and was fertilized with 400 lbs. of Peruvian Guano to the acre. The manure should be placed on the surface and the ridges turned upon it from $3\frac{1}{2}$ to 4 feet from center to center. Plant the slips on the top of the ridges as above recommended.

In addition to the above we offer the Nansemond and Southern Queen varieties.

Prices of Sweet Potato Plants delivered in May or June.

Early Peabody.—25 cents per dozen; 75 cents per 100 by mail; by express 60 cents per hundred, \$5.00 per thousand.

Nansemond.—25 cents per dozen; 75 cents per hundred, by mail; 50 cents per hundred, \$4.00 per thousand, packed, by express.

Southern Queen.—25 cents per dozen; 75 cents per hundred by mail; 50 cents per hundred, \$5.00 per thousand, by express.

Dioscorea Batatas—Chinese Yam.



DIOSCOREA BATATAS—BULBLETS FROM THE AXILS OF THE LEAVES.

One of the most valuable esculents in cultivation, though but little known. Stem twelve to twenty feet in length, of rapid growth, of creeping or climbing habit, forming an excellent covering for a screen; flowers small, white in clusters; leaves, heart-shaped. The root is of a pale russet color, oblong, regularly rounded, club-shaped, largest at the lower end. Plant eight inches apart, in a deep, light soil, tolerably rich, and thoroughly stirred two feet deep. A well grown root, two years from the bulblets, will measure two feet in length, and two and one-half inches in its broadest diameter, and is quite hardy, remaining in the ground over Winter without protection. The flesh is remarkably white, and very mucilaginous in its crude state. They may be boiled or roasted, and when cooked possess a rich-like taste; are quite farinaceous, nutritive, and valuable for food. It is also a very desirable climbing plant, suitable for covering screens, arbors, and unsightly places.

Extract from a letter in the *Rural New Yorker* from a correspondent who has cultivated this plant for many years.

"The Chinese Yam possesses merits that should commend it to the agricultural classes of the United States, yet there are many who don't even know that it is grown in this country, although it is more than twenty years since its introduction here, from China. The difficulty all new beginners have to contend with in the culture of this esculent, is a proper understanding of its wants. I have grown them for twelve or fifteen years, and will give the readers of the *RURAL* the benefit of my experience:

In the first place, select a soil that is moderately dry, deep and rich, and prepare it by spading or plowing deeply; then plant the bulblets eight inches apart each way. When the large roots are used for planting they should be cut up in pieces about one inch in length. They will produce much stronger roots than those grown from the bulblets. They will need to be cultivated a few times the first year, to keep weeds down and soil loose. After the first year, keep the weeds heded off or mowed down, as you prefer. The Yams are perfectly hardy, standing in the ground all Winter and growing again when Spring comes, increasing in size for a number of years. They increase naturally from the small tubers that grow on the vines just above each leaf. These should be saved in the Fall, and kept during Winter where they will not freeze, as freezing injures their growth, and planted in the Spring as soon as the ground gets warm.

As to the product per acre, in the best of soil, with three years' growth and tubers planted eight inches apart each way, we could safely calculate on getting six and a quarter tons of Yams per acre. Some may think this a large yield, but I believe it is not as large as I should put it, and think it is not an overestimate. I have only cultivated them on a small scale, but have come to the above conclusion with regard to their productiveness. A few years ago, I was digging in an old bed of them where they had been left to grow for several years, and such a sight! The ground was literally full of roots, measuring one and a half to two

DIOSCOREA BATATAS. feet in length, and one and a half to two inches in diameter at the
 CHINESE YAM—ONE largest end, and this too where the soil was not more than eight or ten
 YEAR OLD ROOT. inches deep.

They grow in a perpendicular position with the large end downwards. The vines are ornamental, having heart shaped leaves edged with scarlet and are very pretty. A few tubers planted near a door or window and the vines trained over and about it, make an ornament worthy the admiration of all. The flowers are numerous and have a cinnamon fragrance, but the vines do not bloom until the roots are two years old. In garden culture I let the vines run on the ground, except those I wish to save tubers from for seeds. These I set stakes or poles to, as I think by this method the tubers are produced in greater abundance and of large size.



There is scarcely any difference perceptible to the taste between the Chinese Yam, when properly cooked, and the Irish potato, although the Yam is much whiter and somewhat finer grained.

Some object to growing them on account of the great depth to which the roots penetrate. But to those I find no difficulty. I commence digging at the end of a row, take out what I want, and leave the hole open. The next time I dig I commence where I left off, and throw the dirt where I dug before; by this way I have the soil stirred so deep that it is in fine condition for planting out more tubers.

They have no insect enemy, and drought affects them but very little, as they root so deeply. There is no necessity for their being dug at any time of the year, except when wanted for immediate use. They are suitable for cooking any time in the year. A person can plant enough at one planting to do his family for years, each year bringing him larger Yams. I consider them safer to depend on than the Irish potato, which has its enemies, and is so sensitive to a little freeze.


I believe their proper cultivation would be attended with success anywhere in the United States. If the masses of the people better understood their nature and were better acquainted with their many fine qualities, they would be more generally grown. In conclusion I will say that they are the most profitable crop that I cultivate.

Small bullets, which form roots about a foot in length, in one year, in packets of one dozen, 25 cents. Five dozen, \$1.00. One year old roots \$2 00 per dozen; \$12.50 per 100.

J. L. True's Improved Potato Planter.

This Machine will cut the Potato and Plant Six Acres in Ten Hours.

We invite the attention of the farming community to this machine, which is beyond doubt one of the greatest and most successful labor-saving machines of the age.



Having been submitted to thorough practical tests on all kinds of soil, in this country and in Europe, we offer it to farmers for the season of 1878, with entire confidence in its ability to do the work intended for it, in a thorough and proper manner.

No intelligent farmer can fail to see that an implement with which a man can, with the aid of a horse, cut the potatoes, and plant six acres in one day, doing the work in many respects better than it is

usually done by hand, is of inestimable value.

The Machine is very simple in construction, and its arrangement is such as to render it very easy to operate.

The Hopper holds about one bushel of potatoes.

The knife is placed in such a relative position to the seed-boxes that the potatoes are cut into pieces of such a size and shape as to render it almost impossible for any to drop without eyes on them.

The potatoes drop through the plow, the back of which being open, the operator can see each piece as it falls, and they are still exposed to his view for some distance as they lay in the furrow, before the coverers reach and cover them, thus affording him the satisfaction of knowing that the work is being well done.

Gauge rings are provided for diminishing the size of the seed-boxes, when planting different sizes of potatoes separately.

Any kind of dry, concentrated fertilizer may be dropped with the potato in such quantities as may be desired.

The Plow and Coverers can be adjusted to cover the potatoes in a most satisfactory manner, at any depth desired.

The drive wheels mark the rows, which can be made as far apart as the farmer chooses, while the seed is dropped 9, 18, or 36 inches apart in the rows.

Prices of Machines on cars in New York:

No. 1—Machine with Fertilizer Attachment, complete (drops 9, 18 or 36 inches apart),	\$40 00
No. 1—Without Fertilizer Attachment,	37 00
No. 2—Machine has no Fertilizer Attachment, and drops 18 or 36 inches apart,	30 00
Steel Covers, extra,	2 00

Antidote for Colorado Potato Beetle.

From a treatise on the "Pest and its Remedy," published by C. T. Reynolds & Co. New York.

The restless progress of that dreaded pest, the Colorado Potato Beetle, has for several years past been watched with anxiety and alarm by the farmers of the Middle and Eastern States. It was foreseen that it would eventually make its way north and east, and the appearance of this farmer's enemy in Pennsylvania, New York, and portions of New England, last Summer, leaves but little room to doubt that the bug will be out in full force this coming season, and rapidly extend its ravages. In fact, we consider its crossing the ocean itself and becoming naturalized in the old world, as merely a question of time. Already several of the European governments have taken the alarm, and fully alive to the magnitude of the danger with which they are threatened, have endeavored to forestall it by prohibiting the importation into their dominions of American potatoes from the district devastated by the bug. Whether this measure will prove effectual in warding off the evil from their shores remains to be seen, *we believe it will not.*

ITS GREAT PROLIFICACY.

The Colorado beetle propagates itself with astonishing rapidity; several broods (at least three) following each other in the course of the year. The first batch of larvæ makes its appearance towards the end of May, and sometimes even in April, if the weather be warm. Each female lays from 700 to 1200 eggs, in clusters of twelve or thirteen, on the under side of the leaves. In about six days the larvæ are hatched, and immediately begin the work of devastation, which continues for about nineteen days. They then descend to the ground, where they are transformed into pupæ, at the surface of the earth. The perfect beetle appears in ten to fourteen days after the pupæ is formed—begins to pair in about a week, and on the fourteenth day commences to deposit her eggs.

THE VORACITY OF THE BEETLE,

especially in its larval condition, is truly marvellous. When once a field of potatoes has been attacked, unless very prompt remedial measures are resorted to, all hopes of a harvest will speedily vanish, as in a very few days nothing will remain but a barren waste of dried-up stalks.

VARIOUS PLANS

for the destruction of the Colorado potato beetle have from time to time been devised, very few of them have proved satisfactory, however, while many of them were wholly impracticable. Picking the larvæ and bugs from off the plants by hand, for instance, has been thoroughly tried but leaving out of the question the great loss of time this plan involves, and the danger of poisoning from handling the larvæ, it was generally found that after the field had been thus gone over, the insects would be apparently as numerous as ever in twenty-four hours, so that the operation had to be constantly repeated!

PARIS GREEN THE TRUE REMEDY,

is the general verdict of the Western farmer. This is not only the case with respect to efficacy, but it has also the additional merit of being the cheapest in first cost, and the most convenient and economical in use of all the remedies that have been tried. When first proposed as an antidote, some seven or eight years ago, Paris Green met with considerable opposition, but its triumphant success has overcome nearly all prejudice, and while some still hesitate to resort to so virulent a poison, experience has demonstrated that with *due care* no harm can result from its use, as it will not poison the soil or injure the plant or the Potato.

Prof. CHARLES V. RILEY, State Entomologist of Missouri, in his fourth annual report referring to Paris Green, says: "Properly mixed I have used it without the slightest trace of evil on the leaves or tubers; and I know hundreds of others who have done likewise; so that with present experience I should not hesitate to recommend its judicious use." In this connection we take occasion to acknowledge our indebtedness to Prof. Riley's able reports for much valuable information. We also refer to the testimony of the same high authority to Paris Green as "THE remedy for the Colorado potato beetle."

MODES OF APPLICATION.

In its application the Western farmers generally use Paris green *dry*, and as it is a highly concentrated poison it is necessary to mix it with some other substance, such as flour, plaster, or ashes, in proportion varying with the strength of the green, and thus reducing its cost. We give the preference to flour as a vehicle, as combining with the dew on the plant, it forms a paste which adheres with greater tenacity than either plaster or ashes. Another method is by mixing with water, say a large tablespoonful of green to a pailful of water. This is in some respects a convenient way, and has the advantage of being free from dust; it can also be used at any time of day. It has some disadvantages, however: 1st, as the green is not soluble—though it quickly gives a green tint to the water when stirred, it soon settles to the bottom, and needs continual agitation to keep it in suspension. 2d. It *settles in spots on the leaves*, the natural tendency of water in finding its level being to carry and concentrate in wherever a drop finds rest and evaporates. 3d. Much of it is wasted on the ground in sprinkling.

THE UNPARALLELED DEMAND FOR PARIS GREEN,

caused by the rapidity with which the Colorado beetle has overrun the country, has greatly stimulated its production; and, as usual in such cases, the market is flooded with various brands, representing all grades, from the strictly pure down to the highly adulterated article of merely nominal strength and value. There are several qualities manufactured, in order to meet the views and conveniences of all who are compelled to resort to its use.

PROPORTIONS FOR MIXING.

To those who have suitable appliances for mixing, we recommend the "strictly pure," but where these are wanting, and it is an object to avoid much mixing, on account of the danger arising from the dust (and this is the greatest danger attending its use) the cheaper grades are more desirable. These, of course, contain more or less adulteration, but the price is made to correspond, and the buyer has but *comparatively* little mixing to do. The following are the proportions recommended by experienced Western farmers, and also by scientific men who have arrived at the same conclusions by careful experiment:

To 1 lb.	"Strictly Pure"	use 30 to 35 lbs. Flour.
1	"Genuine"	use 25 to 30 "
1	"King's County"	use 20 to 25 "
1	"Long Island"	use 15 to 20 "
1	"Montauk"	use 10 to 15 "

Relatively speaking, the "Genuine" is the cheapest to use, but the "Montauk" has the advantage of being less liable to poison from dusting.

We have made arrangements with the manufacturers of the above to furnish our patrons at the lowest market rates. In consequence of frequent fluctuations, we do not publish prices, but will give prices upon application.

We have on hand a small quantity ready for use for those who wish to use it in their gardens in *one pound packages, at 15 cents each, mixed ready for use.*

There is great danger in mixing this green for potato bug and cotton worm poison, owing to the fine dust which arises in the process, which is inhaled, and also rapidly absorbed by the pores of the skin, especially if the person using it should be in a state of perspiration. To guard against this, the hands and face, (particularly the nostrils) should be protected as much as possible, and should be carefully washed after working in it, or in any of the preparations of which it is an ingredient. As it penetrates and poisons *wood*—gets into the seams and crevices of articles made of metal—and even into earthenware that is at all porous. *All household utensils, or anything in barn or stable (which cattle or horses could have access to) in which the article may have been mixed, or from which it has been used, should be carefully set aside, and never again used for any other purpose.*



APPARATUS FOR APPLYING.

Several ingenious machines have been invented for applying the poison,—three of which are here illustrated. The cheapest and most simple is made in the form of a common dredging-box, holding about two pounds each, and is attached to the end of a pole, as seen in the accompanying engraving. It is used by gently shaking over the plants, taking care to walk to windward, so as to avoid any dust that may arise. It should be applied in the morning, when the dew is on the vines. When mixed with water, it is usually applied by means of an ordinary watering-pot, or sprinkled on the vines with a broom, taking care to keep it well stirred. *Price, 35 cents each.*

Randolph's Fertilizer Distributor.

A Dropper for all Hill Crops, such as Corn, Potatoes, Tobacco, Cotton, Etc.

THE LATEST AND BEST THING OUT. SOMETHING THAT NO FARMER CAN AFFORD TO BE WITHOUT.

For dropping Phosphates, Bone Dust, Fish and Peruvian Guano, Poudrette, Ashes, Plaster, and all kinds of Concentrated Fertilizers.



The manner of using the dropper will be readily understood from an inspection of the cut.

The sack G, having been filled with the fertilizer, is attached to the person by passing the adjustable strap E, over the shoulder as shown. The handle O of the connecting shaft S is then held by the left hand, and the handle H of the plunger J, by the right, unless the person is left-handed, in which case the arrangement is just the reverse, the dropper being adapted for use in either way.

When thus arranged, the operator walks over the field, and as he walks, places the ball step F of the dropper, successively, upon the spots which he has selected for planting and as the plunger J, being unsupported, is necessarily depressed at each rest of the implement, a charge of fertilizer will be deposited in a circle about each hill, its quantity being determined by the previous adjustment of the valves C, C.

It will thus be seen that the implement is used after the manner of a cane, it being operated by simply swinging it from hill to hill, no exercise of care or judgment being required other than to select the spots upon

which to rest its step F. For windy weather a detachable skirt (not shown in the cut, is provided which ensures the accurate placing of the fertilizer in any required circle even under such unfavorable circumstances.

SPECIAL ADVANTAGES.

It is not necessary to touch the fertilizer with the hands, a matter of some importance, when caustic substances are used.

The fertilizer will not stick or clog, even though it be damp and has a tendency to pack, the vibrating hopper being so effective in its operation, that nothing suitable for use, can withstand its action.

The quantity sown may be determined to a nicety, by properly adjusting the valves upon the plunger, so that every hill in the field may be supplied with a uniform amount.

The entire quantity between the valves is always discharged; the construction of the lower valve with its conical surface, and the action of the plunger, which is suddenly arrested at a limit of its movement by a stop, serving to ensure this result.

The amount delivered is accurately deposited in a circle about the planting spot or hill, no more being delivered upon one side than another, and the center of the hill covered by the ball step being, of course, without any deposit, these being the precise conditions desired in using concentrated fertilizers. These effective results, also, can be obtained in windy weather, by using the protecting skirt.

For many other recommendations, and a more complete description—send for circular. Price, \$5.00 each. A liberal discount to agents.

Allen's Potato Bug Destroyer.



Allen's Potato Bug Destroyer, sent out late last Summer, has been greatly improved, and will be found efficient, handy and inexpensive, and an important acquisition where the potato beetle is either established or expected. It is especially contrived for the purpose of quickly and economically throwing a mixture of Paris Green and Flour, or other destructive compound in a penetrating cloud, among Potato and other plants, and upon bushes, vines and trees, to destroy Bugs, Fleas, Worms, Slugs, Caterpillars, etc.

It consists of a peculiarly constructed double coned reservoir, readily filled, and by means of bellows, capable of throwing either a jet or cloud of fine powder, thoroughly dusting a row of potatoes or other plants at the speed of a walk. It is also especially effective for Pear and Rose Slugs, Cotton Worms, Plant Lice and other insects.

Paris Green, before use, should be carefully mixed with five or six times its weight of *dry flour or plaster*; the quantity applied being easily regulated by the quantity of flour added.


Though the Powder is a dangerous poison, this machine discharges at so great a distance from the hands and face, as with reasonable care to make its use *entirely safe*. It is desirable to apply in the *early morning*, the dew then forming a paste, and that time being *less windy*. Price, \$3.00.

The Planet, Jr. Potato Beetle Destroyer.

as with reasonable care to make its use *entirely safe*. It is desirable to apply in the *early morning*, the dew then forming a paste, and that time being *less windy*. Price, \$3.00.

HEXAMER'S PRONG HOE.

The Best Hand Potato Digger in Use.



"HEXAMER'S PRONG HOE"

(TRADE MARK)


The highest premium in the Horticultural Department of the Great American Institute Fair of 1867, was awarded to it. The *American Agriculturist*, in speaking of this implement, gives it the following favorable notice:

HEXAMER'S PRONG HOE.—Dr. HEXAMER, the well-known Horticulturist, has such a way of making his investigations and coming at his facts, that we accept his results with almost the same confidence we would have in our own. He makes use of expensive labor, and the monthly pay-roll is so large as to lead him to employ his men to the very best advantage. The problem was what tool to place in the hands of the laborers for hoeing out between rows of strawberries or other small fruits, root crops, nursery stock, etc. The common hoe is a slow, hard, old-fogy tool, and, of course, its use is out of the question. The potato hook, or pronged hoe with round prongs, good, but not sufficiently rapid for the outlay of strength, yet vastly superior to the hoe for the same purposes, except after weeds have grown large, which ought rarely or never to occur. After having made numerous experiments with tools made expressly for him, he decided upon this implement. The six teeth or prongs are eight inches in length, the outer ones being ten inches apart, which is the width of the actual cut. The prongs are square, of the best steel, and inserted in pairs into a malleable iron head, in which they are firmly wedged. They are delicate, but very strong and elastic.

It stirs the soil thoroughly, more than a foot wide, and from two to four inches deep, killing all small weeds, lifting out stones of small size, removing weeds and all obstructions as effectually as a rake. On light soil it is as easily worked as a hoe, and on heavy soils, if dry enough to work at all, very much easier. It is safe to say that a man, with one of these, can do several times as much work as with a hoe. We think it will prove more useful as a potato digger on account of its breadth, than the implement made for the purpose.

PRICE.—\$1.50 each; \$15.00 per dozen.

Allen's Potato Digging Plow.



ALLEN'S POTATO DIGGING PLOW.

This implement weighs one hundred pounds, and is of very light draft. A pair of small horses or oxen, with a boy to drive will easily dig potatoes as fast as twenty men can pick up. It turns them out so cleanly that scarcely one bushel in fifty, whether small or large, is left uncovered.

The standard is high, so as to allow of its working freely, with-

out clogging from weeds and potato vines, but in harvesting for an early market, when the vines are long and still green, the work will be much facilitated by cutting these and removing them from the rows.

Price of Potato Plow, with cast-iron Mould and Share, \$15.00. The same, with steel Mould and Share, \$25.00.

The Planet Horse Hoe and Cultivator.

This is a new and most perfect tool for horse cultivation. It is especially adapted to market garden and farm-hoeed crops, and to working *Tobacco, Broom-Corn, Cotton, Rice and Roots*. It greatly reduces the *labor of hoeing*, often thus saving cost in a day. Blades tempered and polished steel. Price, \$12.00. With extra cross-piece and pair of hoes, \$15.00.

~ Planet No. 3.—A Fertilizer sower. Price, \$16.75.

Complete directions for use accompany each machine. Send for a full descriptive circular, with cuts of each implement and testimonials.

IMPORTANT TO WHEAT GROWERS.**PRINGLE'S NEW HYBRID SPRING WHEATS.**

We take much pleasure in announcing two new and distinct varieties of Spring Wheat, produced in 1870 and 1871 by Mr. Pringle of Vermont, whose skill and judgment as a hybridizer of Cereals and Potatoes have already won for him a world wide reputation. After a careful trial of six years, we now offer these varieties to the public, having entire confidence in their superiority over all others, feeling assured that they will prove to be valuable acquisitions to the wheat growers of the country.

CHAMPLAIN

Was produced in 1870 by Mr. Pringle, in his endeavors to unite the remarkable hardness of the Black Sea with the fine and superior quality of the Golden Drop. Several varieties were the result of this hybridization, from which this one was chosen, as realizing the end in view, showing greatly increased vigor and productiveness over both its parents. A careful selection from this for the past seven years, has now fully established its character, and we have a wheat bearded like the Black Sea, with the white chaff of the Golden Drop, free from rust and smut, yielding a lighter colored grain than the former, which makes a flour of superior quality. Its strong and vigorous straw, growing 6 to 12 inches higher than its parent varieties, stands erect, frequently bearing, even in very ordinary culture, heads from 5 to 6 inches in length, containing from 50 to 75 kernels each.

We confidently recommend this new wheat as among the earliest, promising to give the growers of this most important crop better results than are produced by the old and "run out" varieties now sown. Price, \$1.00 per lb; 3 lbs., \$2.50, by mail post-paid; peck, \$9.00.

DEFIANCE.

Another variety of Spring Wheat of the highest promise, the result of a series of experiments by Mr. Pringle in 1871 to incorporate superior qualities upon the hardy stock of our common Clib Wheat, by hybridizing it with one of the finest, whitest, and most extensively grown sorts of the Pacific Coast.

This variety displays great productiveness, vigor and hardness. It is a beardless, white chaff wheat, with heads frequently five to six inches long, very closely set with large white kernels, frequently numbering 75 to 80 to the single head. Its white, stiff, erect straw, exempt from the attack of rust, its earliness combined with great vigor and superior qualities, should claim for it universal trial. Price, \$1.00 per lb; 3 lbs., \$2.50, by mail post-paid.

\$250.00 in Premiums.*

In order to induce Wheat Growers to give these new sorts a thorough and universal trial, we offer **\$250.00 in premiums** for the largest quantities grown from one pound of seed purchased from us. Parties will please state in their order that they desire to compete for these premiums.

For the largest quantity of Champlain Wheat grown from one pound of seed,	\$50 00
For the second largest quantity,	25 00
For the third largest quantity,	15 00
For the fourth largest quantity,	10 00
For the largest quantity of Defiance Wheat grown from one pound of seed,	50 00
For the second largest quantity,	25 00
For the third largest quantity,	15 00
For the fourth largest quantity,	10 00

We also offer additional premiums of \$50.00 for the best 20 heads of either of the above varieties as follows, \$25.00 for each variety.

For the best 20 heads of Champlain,	\$10 00
For the second best 20 heads,	7 00
For the third best 20 heads,	5 00
For the fourth best 20 heads,	3 00
For the best 20 heads of Defiance,	10 00
For the second best 20 heads,	7 00
For the third best 20 heads,	5 00
For the fourth best 20 heads,	3 00

Competitors for the prizes will be required to give the date on which they gave their order for the wheat, date of sowing, time of harvesting, with a written statement of their mode of culture, characteristics of the soil, whether clay, alluvial, or sandy loam,—nature of the subsoil, whether under-drained or not; also, the kind and quantity of fertilizers used, how and when applied, with the weight of the crop when harvested and cured, which must be witnessed and sworn to before a Justice of the Peace, Notary, or any one competent to administer the oath, and sent to our address before the first of November, 1878. The awards will be made by a competent committee of Agriculturists, and will be printed and a copy mailed to each competitor. The prizes will be paid in December, 1878.

The engravings are exact representations of some of the heads grown with ordinary farm culture.

Competitors for the premiums for the best heads of wheat must deliver them at our office, 34 Barclay street, New York, (all charges prepaid), previous to the 10th of September, 1878. The straw must be cut not less than eight inches in length exclusive of the heads, and carefully packed to prevent injury in the transportation. They can be sent by mail or express prepaid, at the option of the grower. If sent by mail, they should be boxed in a light wooden or a stiff paste-

*Circulars explaining the mode of awarding the Premiums for the largest quality grown from one pound mailed to applicants.

board box, the cover of which must be secured by strong twine, and not nailed or fastened in any way to prevent the examination of the parcels at the New York post-office. No writing of any kind must be enclosed in the packages, as in that case letter postage would be charged on the whole package, and would not be taken from the office. The name of the sender must be written on the *outside* of the package, with our address. Notify us by letter when the wheat is sent. All those who wish their samples returned, in the event of their not receiving a premium, must enclose in their letter at the time of sending the heads a sum sufficient to pay return charges; otherwise the wheat will not be returned. The premium samples will be retained by us.

A NEW WINTER WHEAT.

Arnold's Victor Wheat.—This wheat is the final result of a large number of intercrossings, and selections of the best varieties grown in America. After carefully experimenting for a number of years, it is claimed that a variety has been obtained with a hardness in resisting the effects of a Canadian winter, entirely superior to the Diehl when grown side by side, and subjected to the very same conditions. One bag, containing 150 lbs., \$9.00. One bushel, including bag, \$4.50. One peck, including bag, \$1.50. 3 lbs., by mail, \$1.00.

COMPTON'S EARLY FIELD CORN.

This corn, which has made such a sensation in agricultural circles, is the result of years of patient effort by one of our most intelligent experimenters. Seeing the great deterioration of the common sorts, and comprehending the magnitude of the losses frequently sustained in consequence of the partial or total destruction of the crop by early Autumn frosts, he instituted a series of experiments with the view of obtaining a new variety in which extreme earliness and great productiveness should be combined. A large plot was planted with seed from the small huskless ears sometimes found on the tops or blossom branches. The kinds so obtained proved to be different from each other, and each distinct from any known variety. One stalk on which were four fine ears, ripened by the middle of August. The crop of this stalk is the source of the variety offered. It is a seedling of the Dutton, and resembles its parent somewhat in appearance. Stalks grow eight to ten feet in height, ears from ten to seventeen inches in length, well filled to the end. Kernel medium, bright yellow and of the flinty order.

This variety was sent to all parts of the country the past season, and we have yet to hear one unsatisfactory report. All who have tried it strongly recommend it. It is an acquisition which farmers cannot fail to appreciate.

Price, one pint, by mail prepaid, 50 cents; one quart, by mail prepaid, 80 cents; by express, charges to be paid by purchaser, peck, \$2.50; bush., \$8.00. Select ears, by mail, 25 cents each; by express, \$1.50 per doz.; \$8.00 per 100.

WHITE RUSSIAN SPRING WHEAT.

The White Russian Wheat was first grown in this country in Wisconsin. It was increased from a small quantity received from Russia, and has been known as the White Russian, although it is not exactly white, but is of much lighter color than most varieties of Spring Wheat.

It is a bold white chaff wheat, and has proved itself to be one of the best Spring wheats ever grown in Wisconsin. Such long, straight, strong, healthy yellow straw, bearing large long white chaff heads, well filled with plump kernels, weighing oftentimes from 60 to 62 lbs. to the measured bushel, and the wheat producing 5 to 10 bushels more per acre than other once well thought of varieties that are now fast running out, as all wheat does after being sown a long series of years under the influence of the same soil and climate. As soon as it can be got in sufficient quantity, it is destined to crowd out to a great extent other varieties, and will become the *Staple Spring Wheat* of the United States, until other new varieties take its place.

The White Russian Wheat stands well after being ripe, and is not liable to lodge when green, or rust; it is decidedly a healthy and sure crop wheat, that has in many cases produced a full average crop, where other varieties along side of it have failed. 3 lbs., by mail, \$1.00; 1 peck, \$1.00. 1-2 bush., \$1.75; 1 bush., \$3.00.

GOLDEN MILLET.

Golden Millet is of extraordinary value, and coming into great favor as an annual hay and fodder crop. It is mistakenly called German Millet, which is a very indefinite name for seed imported from Europe under this name proves to be the *Panicum miliaceum* or common Millet of our catalogues and persons ordering German Millet of ordinary seedsmen, will probably get *Sytaria Germanica*, or Hungarian grass. Hence Golden Millet is the only definite name, for no other plant bears the name. The experience of those to whom we sold seed last year, warrants our speaking of it in the highest terms.

Prices: We can furnish the seed at the following prices, by express or freight, charges paid by purchaser; per bushel, \$2.00; per $\frac{1}{4}$ bushel, 60 cents; by mail, postage prepaid by us, 40 cents per quart.

KENNEY'S EARLY AMBER SUGAR CANE.

In this variety we have a plant that is destined to prove of immense value to the northern portions of our country. It has been tested in Minnesota for several years past. A large grower gives the following statement to the State Commissioner:

I planted a piece of this cane this year, and the 27th of May, on new land, and before any frosts it was entirely ripe. Its saccharine qualities are of the first order, fine and rich, and its earliness highly recommend it to the whole country. It has another desirable quality, it stands up well, does not easily blow down, usually grows eleven and twelve feet high with us. When planted between the first and tenth of May, the seed almost invariably ripens, thus giving one the advantage of a double crop in one year, from one planting, some 20 or 25 bushels of seed, and 150 or more gallons of fine syrup per acre, that is, if it is skillfully handled. All kinds of farm stock are fond of the seed; when ground and mixed with bran it makes a good heavy feed for horses; hens are very fond of the seed, and I believe it makes them produce eggs more abundantly than almost any other kind of grain. We have seen sugar produced from this variety equal to the best coffee sugars in the market.

By mail, 75 cents per lb. When sent by express at the expense of purchaser, 60 cents per lb.

PRICE LIST OF MANURES.

No. 1 PERUVIAN GUANO, PURE BONE, Mapes' Complete and Special Manures,

(Including the celebrated Formulas of Prof. George Ville.)

PREPARED BY

THE MAPES' FORMULA AND PERUVIAN GUANO CO.

CASH MUST ACCOMPANY ALL ORDERS FOR FERTILIZERS.

The following quotations include cartage, excepting when total quantity ordered is less than half a ton, then add fifty cents for cartage.

PERUVIAN GUANO, at Gov't Stores, Brooklyn, N. Y.

	Per Ton, 2000 lbs.	Single Package, Per Pound
No. 1 PERUVIAN GUANO —"STANDARD," or "GUANAPE." Ammonia, 10 per cent.; Phosphoric Acid, 12 to 15; Potash 2 to 3. Average weight of Bags 180 lbs.	\$56.50	3c.
No. 1 PERUVIAN GUANO —"LOBOS." Ammonia, 6 to 7 per cent.; Phosphoric Acid, 15 to 20; Potash, 3 to 5. Average weight of Bags 210 lbs.	47.50	2½c.
No. 1 PERUVIAN GUANO —GUARANTEED. Cargo B....Ammonia, 11.50; Phosphoric Acid, 17.10; Potash, 2.30.....Bags 200 lbs.	70.00	\$7.35 Per 200 lbs.
Cargo F....Ammonia, 6.30; Phosphoric Acid, 19.70; Potash, 3.20.....Bags 200 lbs.	58.00	5.90
No. 1 PERUVIAN GUANO —RECTIFIED. Cargo.....Ammonia, 9.70; Phosphoric Acid, 14.25; Potash, 2.....Bags 200 lbs.	69.00	7.25
Cargo.....Ammonia, 3.40; Phosphoric Acid, 16.90; Potash, 3.40.....Bags 200 lbs.	51.00	5.35

No. 2 PERUVIAN GUANO.

Ammonia, 3 per cent.; Phosphoric Acid, 15; Potash, 2.....Bags 200 lbs. 38.00 4.00

No charge for Delivery at Government Stores, at Brooklyn, to vessel.

Official descriptive Pamphlets on Peruvian Guano furnished to dealers, with their name on cover. Liberal concessions made to manufacturers and dealers, from even the official price list of the Agents of the Peruvian Government

CHEMICALS, FURNISHING AMMONIA, ETC.

NITRATE OF SODA.

STANDARD 95 per cent., equal to 19 per cent. Ammonia.....Bags.....lbs. \$75.00 4¼c.

SULPHATE OF AMMONIA.....

STANDARD 25 per cent. Ammonia..... { Bbls.....lbs. 90.00 5½c.

{ Casks.....lbs.

GROUND DRIED BLOOD—(Pure).

GRADE AA...Ammonia, 14 to 15.....Bags.....lbs. 50.00 3c.

GROUND DRIED FLESH—(Odorless).

GRADE AA...Ammonia, 14½ to 15½.....Bags.....lbs. 50.00 3c.

CASTOR PUMICE—Ammonia.....Bags.....lbs.

SULPHATE OF MAGNESIA.

GRADE.....per cent. Sulphate Magnesia.....Bags.....lbs. 22.50 1¼c.

PLASTER, Nova Scotia—Very finely ground.....Bags 200 lbs.

{ Bbls. 250 lbs. 8.00

All Chemicals, grades of Bone, Formulas of Prof. Ville, Complete and Special Crop Manures, Super-Phosphates, and Potash Salts, sold by us subject to test by Prof. Geo. H. Cook, New Jersey State Chemist; Prof. W. O. Atwater, Connecticut Experiment Station; Dr. Wm. M. Habirshaw, Chemist New York State Agricultural Society.

PRICE LIST OF MANURES.

PROF. VILLE'S FORMULAS.

	Per Ton, 2000 lbs.	Per 200 lbs.
COMPLETE MANURE —(Ville Formula)—Ready Mixed.		
A substitute for barn-yard manure, adapted for general use, especially on light or sandy soils, or those deficient in potash. Ammonia, 7.69 per cent.; Phosphoric Acid, 5; Potash, 7.59.....Bags 200 lbs.	\$52.28	\$5.50
FOR ONIONS and Vegetables —Use the above Complete Manure.		
TURNIP MANURE —(Ville Formula)—Ready Mixed. Ammonia, 2.53 per cent.; Phosphoric Acid, 7.50; Potash, 7.59.....Bags 200 lbs.	38.72	4.00
POTATO MANURE —(Ville Formula)—Ready Mixed. Ammonia, 4.50 per cent.; Phosphoric Acid, 5.97; Potash, 13.64.....Bags 200 lbs.	51.04	5.40
These formulas are now furnished by us at the above quoted prices, in a ready-mixed and finely ground, dry condition.		
Any of these formulas, when ordered in lots of two tons and upwards, may be obtained in the form of the materials of which they are composed, all shipped in their separate and original packages, at a reduction of five dollars per ton from the above prices.		
The prices charged for these formulas in the unmixed form, will be found to be based upon the actual per centages of the following ingredients at the annexed low rates.		
Ammonia, from Peruvian Guano, Ground Dried Blood or Flesh. 17½c. per lb.		
Phosphoric Acid, all in form of Peruvian Guano.....	9c.	“
Potash, as Phosphate, Sulphate and Muriate.....	7½c.	“

MAPES' COMPLETE AND SPECIAL MANURES.

In the preparation of these Manures, particular attention is paid to supplying all the required ingredients to meet the demand of each crop; not only in the proper proportion, but also in those forms and conditions which practical and varied experience has shown to develop the healthiest growth and best quality of product, as well as the largest yield.

This will be readily appreciated in the case of such crops as Tobacco, also Spring Wheat as compared with Fall Wheat, &c.

The materials from which these Manures are prepared are Peruvian Guano, Nitrate of Soda, Sulphate of Ammonia, Dried Blood or Flesh, Fine Bone, Sulphuric Acid, and Potash Salts. They contain no plaster. They are all supplied in a ready-mixed state, finely ground, and easy of distribution.

MAPES' COMPLETE MANURE.

Ammonia, 6 to 7; Phosphoric Acid, 10 to 12; Potash, 3 to 4.....Bags 200 lbs. 50.00 5.25

Adapted for general crops on heavy soils or average clay-loams, or those requiring an increased per centage of Phosphoric Acid as compared with Potash, such as fair Corn or Potato lands.

This Manure for clay lands, as well as the Ville Complete Manure for light or sandy soils, is recommended as our WHEAT MANURE, and for seeding down to permanent mowing or pasture lands, also for Oats, Barley, Rye, and Hops, as well as for general use, as a substitute for barn-yard or stable manure.

MAPES' CORN MANURE.

Ammonia, 4.50 to 5; Phosphoric Acid, 10 to 12; Potash, 6 to 7.....Bags 200 lbs. 49.50 5.20

MAPES' FODDER CORN MANURE.

Ammonia, 4; Phosphoric Acid, 6; Potash, 12 to 13.....Bags 200 lbs. 49.00 5.15

MAPES' BEET AND MANGOLD MANURE.

Ammonia, 6.50 to 7.50; Phosphoric Acid, 4 to 5; Potash, 7 to 8.....Bags 200 lbs. 52.00 5.45

The above Formulas and Manures should be reduced in strength by mixing with plaster or dry earth. This will secure better distribution and guard against injury to the young roots, especially if used in the hills or close to the seed.

PRICE LIST OF MANURES.

	Per Ton, 2000 lbs.	Per 200 lbs.
MAPES' TOBACCO MANURE. Ammonia, 5.00; Phosphoric Acid, 5.00; Potash, 11.90.....Bags 200 lbs.	\$49.34	\$5.20
The Potash in this manure is all in the form of a Sulphate.		
MAPES' FRUIT AND VINE MANURE. Ammonia, 1.50 to 2.00; Phosphoric Acid, 7.50 to 11.00; Potash, 6 to 7..Bags 200 lbs.	35.00	3.70
MAPES' SPRING WHEAT MANURE. Ammonia, 6 to 7; Phosphoric Acid, 7 to 8; Potash, 5 to 6..... .Bags 200 lbs. <i>May be drilled in with the seed.</i>	50.00	5.25
MAPES' CAULIFLOWER AND CABBAGE MANURE. Ammonia, 5.50 to 6.00; Phosphoric Acid, 4 to 5; Potash, 10 to 12..... .Bags 200 lbs.	49.00	5.15
MAPES' GRASS AND GRAIN SPRING TOP-DRESSING.Bags 200 lbs.	51.00	5.35
This Manure, owing to the solubility of its Phosphoric Acid, will be found very useful as a spring top-dressing to winter grain which has suffered injury from winter exposure; also, as a dressing to any crop around the hills or rows to promote rapid growth.		
MAPES' SUGAR CANE OR SORGHUM MANURE. Ammonia, 2.00 to 3.00; Phosphoric Acid, 10.00 to 12.00; Potash, 6.00 to 8.00.Bags 200 lbs.	45.00	4.75
We prepare this Manure to meet the special requirements of the sugar lands of South America, West Indies, and the Southern States. The effect of this Manure is particularly to increase the proportion of SACCHARINE MATTER.		
MAPES' LAWN TOP-DRESSING. <i>For Lawns, Croquet Grounds, Grass Plots, &c.....</i> Bags 200 lbs.	60.00	6.30
FOR COTTON —Use Mapes' Nitrogenized Super-Phosphate.		

MAPES' SUPER-PHOSPHATES, ETC.

MAPES' NITROGENIZED SUPER-PHOSPHATE —(with Potash). Fine and Dry. Ammonia, 2½ to 3½ per cent; Phosphoric Acid (soluble and available), { Bags 200 lbs. 10 to 14; Potash, 2 to 3..... } Bbls. 250 lbs.	45.00	4.75
<i>The Phosphoric Acid, ALL from Bone and Peruvian Guano.</i>		
MAPES' NITROGENIZED SUPER-PHOSPHATE (Old Brand) { Bags 200 lbs. Ammonia, 3 to 3½ per cent.; Phos. Acid (soluble and available), 10 to 14.. } Bbls. 250 lbs.	43.00	4.50
<i>The Phosphoric Acid, ALL from Bone and Peruvian Guano.</i>		
AMMONIATED DISSOLVED BONE —(Moosehead Brand)..... { Bags 200 lbs. Ammonia, 2½ to 3 per cent.; Phos. Acid (soluble and available), 10..... } Bbls. 250 lbs.	40.00	4.20
<i>The Phosphoric Acid, ALL from Bone.</i>		
DISSOLVED BONE-BLACK —(Mapes Brand)..... { Bags 200 lbs. Phosphoric Acid (soluble and available), 15 to 17..... } Bbls. 250 lbs.	35.00	3.68
ACID PHOSPHATE. Phosphoric Acid (soluble and available), 11 to 14.....:..... Bbls. 250 lbs. <i>The Phosphoric Acid from South Carolina Phosphate.</i>	30.00	3.15
PREPARED DRY FISH GUANO —(Mapes Brand)..... { Bags 180 lbs. Ammonia, 5 to 6½ per cent.; Phosphoric Acid, 6 to 7½..... } Bbls.....lbs.	40.00	4.20
The dry, fine condition of this preparation of Fish Guano, renders it convenient for Transportation, especially when packed in barrels. It is in considerable demand for shipment to the Truck Growers in Charleston, Florida, and Bermuda.		
MAPES' POTASH BONE PHOSPHATE. Ammonia, 2 to 3 per cent.; Phosphoric Acid, 11 to 13; Potash, 6 to 7.....Bags 200 lbs. <i>Prepared from Bone and Potash Salts.</i>	35.00	3.70

This Manure is intended to meet the demand for a **cheap and effective fertilizer** for restoring worn-out lands, such as old orchards, side-hill stony pastures; to take the place, in fact, of finely ground Pure Bone and Wood Ashes. May be used broadcast as a top-dressing, spring or fall.

PRICE LIST OF MANURES.

GROUND BONE—STRICTLY PURE.

	Per Ton. 2000 lbs.	Per Bag, 200 lbs.
PURE BONE MEAL —(Mapes Brand). Ammonia, 4 to 5 per cent.; Phosphoric Acid, 20 to 26.....Bags 200 lbs.	\$42.00	\$4.40
PURE GROUND BONE —(Mapes Brand)— Extra Fine } Ammonia, 3.50 to 4.50 per cent.; Phosphoric Acid, 20 to 28 } { Bags...200 lbs. { Bbls.....lbs.	40.00	4.20
This grade of bone will be found to be nearly as fine as ordinary Bone Meal .		
PURE GROUND BONE —(Mapes Brand)— Fine } Ammonia, 3.50 to 4.50 per cent.; Phosphoric Acid, 20 to 28 } { Bags...200 lbs. { Bbls.....lbs.	38.00	4.00
This grade of Bone designated Fine , consists of the three other grades, viz: Meal, Extra Fine, and Medium, in about equal proportions, and is therefore well adapted to meet the wants of those who desire permanent effects as well as immediate action.		
PURE GROUND BONE —(Mapes Brand)— Medium } Ammonia, 2 to 4 per cent.; Phosphoric Acid, 20 to 30..... } { Bags...200 lbs. { Bbls.....lbs.	36.00	3.80
PURE FINE BONE, Dissolved in Sulphuric Acid —(Mapes Brand). Ammonia, 2.50 to 3 per cent.; Phosphoric Acid, soluble and available, 12 to 16.	42.00	4.40
This brand is prepared from only bone and new Sulphuric Acid , and will be found to be more economical to the farmer than any home-made preparation.		
PURE BONE MEAL, for Feeding	60.00	6.30
PURE BONE, for Chickens	50.00	5.25
All of the above grades consist of absolutely pure Bone , reduced by improved machinery to fine condition, without admixture or treatment with any dryer, salt or nitre-cake, plaster, natural phosphate, "Kettle Bottoms" or adulterant.		
The Ammonia is therefore all intimately blended (in the form of the gelatine and organic matter of the Bone) with the phosphates of the Bone, and thus secures rapid decomposition and best action on plant growth.		
Any desired grade of Bone prepared to order. Deliveries in bulk made at our Works on Morris Canal, Newark, N. J.		

POTASH SALTS, Etc.—Our own Importation.

		Single Package, Per lb.
MURIATE OF POTASH —(High Grade). STANDARD 80 per cent., equal to 50 p. c. actual Potash. Bags about 200 lbs.	\$45.00	2½c.
MURIATE OF POTASH —(High Grade) "Douglasshall." (This Grade contains about 10 per cent. Sulphate Magnesia). STANDARD 80 per cent., equal to 50 p. c. actual Potash. Bags even weight 225 lbs	45.00	2½c.
SULPHATE OF POTASH —(High Grade). Test 70 to 85 per cent., equal to 33½ to 46¼ per cent. actual Potash..... { Bags.....lbs. STANDARD 80 per cent., equal to 44 p. c. actual Potash..... { Bbls...340 lbs.	65.00	4c.
SULPHATE OF POTASH —(German). Test 40 to 60 per cent., equal to 22 to 33 per cent. actual Potash.....Bags.....lbs.	35.00	2c.
GERMAN POTASH SALTS — <i>Kainit</i> —"Leopoldshall," "Stassfurt." Test 20 to 32 per cent. Sulphate of Potash equal to 11 to 18 per cent. actual Potash.....Bags 200 lbs.	15. to 18	1½c.
STANDARD 24 per cent., equal to 13 p. c. actual Potash.		
NITRATE OF POTASH . Test 90 to 95 per cent., equal to 42 to 46 per cent. actual Potash. Contains also some 12½ per cent. of Nitrogen, (Ammonia, 15).....Bags about 200 lbs.	170.00	0½c.
STANDARD 95 per cent., equal to 46 per cent. actual Potash. Ammonia, 15 per cent.		

SPECIAL RATES will be given on all of the above grades of Potash Salts, delivered in bulk or packages direct from vessel in which imported, (involving no expense to us of storage, etc.) Dealers can obtain their supplies in this way at very low figures.

Extract from Circular of the Mapes Formula and Peruvian Guano Company, New York.

CLASSIFICATION AND REQUIREMENTS OF CROPS.

Crop.	Yield.	Weight Per Bushel.	Lbs.		Phos. Acid Lbs.	Potash Lbs.
			Nitrogen Lbs.			
Wheat	20 bushels.....	60 lbs.	{ Grain.....	1,270... 25	9.50	6.40
			{ Straw.....	3,000 .. 14.40		
				39.40	16.10	25.30
"	30 bushels.....	60 lbs.	{ Grain ...	1,800 }	59.10	21.15
			{ Straw ...	4,500 }		
Barley	30 bushels.....	48 lbs.	{ Grain.....	1,440... 27.50	9.64	6.33
			{ Straw.....	2,000... 6.26		
				33.76	15.04	32.13
"	40 bushels.....	48 lbs.	{ Grain.....	1,920 }	45.01	20.05
			{ Straw.....	2,666 }		
Rye.....	15 bushels	56 lbs.	{ Grain.....	840... 15.00	7.03	4.69
			{ Straw.....	1,780... 4.21		
				19.21	10.31	18.28
"	30 bushels.....	56 lbs.	{ Grain.....	1,680 }	38.43	20.62
			{ Straw.....	3,560 }		
Oats.....	30 bushels	32 lbs.	{ Grain.....	96) 18.40	6.00	4.20
			{ Straw.....	2,000... 11.20		
				29.60	9.80	22.00
"	50 bushels	32 lbs.	{ Grain.....	1,600 }	49.34	16.34
			{ Straw.....	3,333 }		
Average Meadow Hay...	2 tons	(Dry)	Yield.....	4,000... 60.00	16.40	52.80
Timothy Hay	2 tons	(Dry.)	Yield.....	4,000... 62.00	28.80	81.60
Hops	600 bushels.....	50 lbs.	Yield.....	30,000... 63.60	46.50	50.40
Sugar Beets	10 tons		{ Roots.....	20,000... 32.00	16.00	78.00
			{ Tops.....	5,000... 15.00		
				47.00	22.50	109.50
Carrots.....	364 bushels.....	55 lbs.	{ Roots.....	20,000... 44.00	20.00	56.00
			{ Tops.....	8,000... 40.80		
				84.80	29.60	79.20
Corn (Shelled).....	60 bushels.....	56 lbs.	{ Grain.....	3,350 .. 53.60	18.40	11.20
			{ Stover.....	6,400... 30.40		
				84.00	42.80	117.60
"	75 bushels.....	56 lbs.	{ Grain.....	4,200 }	105.00	53.50
			{ Stover.....	8,000 }		
Sugar Cane.....						
Sorghum						
Cotton.....	500 lbs.....	(Lint)	(Seed Cotton) 1,500...	46.50	14.40	16.60
Turnips	10 tons.....		{ Roots.....	20,000... 36.00	18.00	66.00
			{ Tops.....	6,000... 18.00		
				54.00	23.40	82.80
Rutabagas.....						
Tobacco	1,800 lbs....	(Green Leaf)	{ Dry Leaf..	1,260 .. 49.00	7.50	71.00
			{ Stalks.....	1,100... 33.00		
				82.00	22.50	118.00
Potatoes.....	300 bushels.....	66 lbs.	Tubers.....	18,000 .. 58.00	32.00	101.00
Cabbages	20 tons.....		Yield.....	40,000... 104.00	80.00	240.00
Peas.....	25 bushels.....	60 lbs.	{ Seed.....	1,500 .. 63.70	12.90	14.70
			{ Straw.....	3,600... 37.44		
				91.14	25.50	51.06
Beans.....	25½ bushels.....	62 lbs.	{ Seed.....	1,581... 61.50	18.81	20.71
			{ Straw.....	2,800... 45.64		
				110.14	27.77	72.51
Buckwheat	30 bushels	48 lbs.	{ Grain.....	1,440... 20.73	8.20	3.88
			{ Straw.....	2,000... 26.00		
				46.73	20.40	52.28
Green Crop Fodder Corn	20 tons		Yield.....	40,000... 76.00	52.00	172.00
			Fodder Rye	5 tons	Yield.....	10,000... 53.00
Red Clover (in blossom)	10 tons.....		Yield.....	20,000... 102.00	28.00	88.00
Red Clover Hay.....	3 tons		Yield.....	6,000... 118.20	33.60	109.80

Group A.—The dominant element required in the Fertilizer is Nitrogen, next Phosphoric Acid and Potash.
Group B.—Dominant requirement is P. A. Am., Pot.
Group C.—Dominant requirement is Potash, next Ammonia and Phosphoric Acid.

Can Concentrated Manures be Used with Profit?

The following table gives the estimated cost [per bushel, etc.] to grow farm produce in excess of the natural yield of the soil. The estimates include all the Phosphoric Acid and Potash contained in the plant, together with the full proportion of Nitrogen, as recommended by Prof. GEORGE VILLE.

							Cents.
One Bushel Wheat	60 lbs.	Nitrogen	0.625	lbs.	at 21	c. per lb.	13.25
		Phosphoric Acid	0.475	"	at 9	"	4.26
		Potash	0.329	"	at 7	"	2.40
							20 cents.
The Natural Proportion of Straw	150 lbs.	Nitrogen	0.369	"	at 21	"	7.67
		Phosphoric Acid	0.330	"	at 9	"	2.97
		Potash	0.945	"	at 7	"	7.08
							18 "
One Bushel Barley	48 lbs.	Nitrogen	0.158	"	at 21	"	9.56
		Phosphoric Acid	0.321	"	at 9	"	2.88
		Potash	0.211	"	at 7	"	1.58
							14 "
The Natural Proportion of Straw	67 lbs.	Nitrogen	0.104	"	at 21	"	2.21
		Phosphoric Acid	0.180	"	at 9	"	1.62
		Potash	0.860	"	at 7	"	6.45
							10 "
One Bushel Rye	56 lbs.	Nitrogen	0.500	"	at 21	"	10.62
		Phosphoric Acid	0.168	"	at 9	"	4.21
		Potash	0.312	"	at 7	"	2.34
							17 "
The Natural Proportion of Straw	118 lbs.	Nitrogen	0.140	"	at 21	"	2.97
		Phosphoric Acid	0.218	"	at 9	"	1.96
		Potash	0.906	"	at 7	"	6.79
							12 "
One Bushel Oats	32 lbs.	Nitrogen	0.306	"	at 21	"	6.49
		Phosphoric Acid	0.200	"	at 9	"	1.80
		Potash	0.140	"	at 7	"	1.05
							9 "
The Natural Proportion of Straw,	67 lbs.	Nitrogen	0.186	"	at 21	"	3.95
		Phosphoric Acid	0.126	"	at 9	"	1.13
		Potash	0.593	"	at 7	"	4.44
							19 "
One Bushel Shelled Corn	56 lbs.	Nitrogen	0.223	"	at 21	"	4.73
		Phosphoric Acid	0.306	"	at 9	"	2.75
		Potash	0.186	"	at 7	"	1.39
							9 "
The Natural Proportion of Stover	107 lbs.	Nitrogen	0.126	"	at 21	"	2.67
		Phosphoric Acid	0.406	"	at 9	"	3.65
		Potash	1.773	"	at 7	"	13.29
							20 "
One Bushel Potatoes (Tubers)	60 lbs.	Nitrogen	0.097	"	at 21	"	2.06
		Phosphoric Acid	0.106	"	at 9	"	0.95
		Potash	0.336	"	at 7	"	2.52
							5 1/2 "
One Bushel Buckwheat	48 lbs.	Nitrogen	0.172	"	at 21	"	3.66
		Phosphoric Acid	0.273	"	at 9	"	2.45
		Potash	0.129	"	at 7	"	.96
							7 "
The Natural Proportion of Straw	67 lbs.	Nitrogen	0.216	"	at 21	"	4.59
		Phosphoric Acid	0.496	"	at 9	"	3.65
		Potash	1.613	"	at 7	"	12.99
							20 "
One Bushel Onions	50 lbs.	Nitrogen	0.053	"	at 21	"	1.12
		Phosphoric Acid	0.077	"	at 9	"	0.69
		Potash	0.084	"	at 7	"	0.63
							2 1/2 "
Dry Leaf Tobacco	100 lbs.	Nitrogen	1.94	"	at 21	"	41.22
		Phosphoric Acid	0.59	"	at 9	"	5.31
		Potash	5.63	"	at 7	"	42.22
							89 "
The Natural Proportion of Stalk	87 lbs.	Nitrogen	1.30	"	at 21	"	27.62
		Phosphoric Acid	1.18	"	at 9	"	10.62
		Potash	3.72	"	at 7	"	27.96
							66 "
Seed Cotton (3/4 lbs. Lint.)	1,000 lbs.	Nitrogen	2.50	"	at 21	"	.531
		Phosphoric Acid	9.60	"	at 9	"	.874
		Potash	11.06	"	at 7	"	.829
							\$2.22

The cost of raising the staple products with Mapes' Complete and Special Manures has proved in practice to be much less than shown by the figures of the above table. This has been uniformly the case even upon poor light soils.

Owing to the difficulty of bringing the fertilizer, especially during the first season of its application, in contact with the roots of the plants, there will always remain a large proportion unused and ready for future crops. As before stated, the marked effects of the fertilizer on subsequent crops, particularly on grass after grain crops, give evidence of the increased stock of available plant food left in the soil after cropping.

Again, the natural fertility of the soil is increased by the action of the fertilizer in liberating and rendering available, for plant use, the ingredients which it already contains, but which without the presence of the manure, would remain inert and useless.

For these reasons the cost of raising farm-products with Mapes' Manures, including fertilizer, seed, labor, interest on land, etc., on poor as well as good land, is found to be less than the estimated cost of the fertilizing ingredients alone required as per above table.

With 300 lbs. "Complete Manure" per acre on 60 acres poor land this season in Connecticut, the cost of raising corn per bushel was less than *twenty-five* cents. This included all expenses of fertilizer, cultivation, interest, etc. The yield of corn with 300 lbs. per acre of this same manure, has been from 80 to 96 bushels shelled corn per acre. In one instance, where the yield was at the rate of over 90 bushels shelled corn per acre, the writer stated that the "land was so poor that without the fertilizer it would have produced nothing."

On potatoes, the potato manure, 300 to 400 lbs. per acre, has produced from 350 to 375 bushels per acre under ordinary culture. The cases reported were principally "Late" and "Early Rose." In one instance, during the past season, where land heavily manured with barn-yard manure, produced only 150 bushels of indifferent quality, the addition of seven dollars worth of potash fertilizer (13 per cent. Potash) raised the yield to 350 bushels of potatoes of large size and very fine quality.

$\frac{27}{100}$ of a Cent per Day, 8 1-2 Cents a Month, or \$1.00 a Year,

PAYS FOR A TRUE FARMER'S PAPER,

The Scientific Farmer,

A Handsomely Illustrated Monthly Journal in the interests of Profitable Agriculture.

For all farmers gardeners, stock raisers, grain growers and dairymen. For all who wish to learn about fertilizers and manures, about the principles underlying successful practice in the field and stable. It is full of the most valuable and practical information. Its articles on PLANT FEEDING have intensely interested thousands of farmers; for it is an acknowledged authority on all subjects relating to manures and fertilizers; and its articles are so *useful*, so *practical*, and so *reliable*, that no farmer can afford to be without it.

In HORTICULTURE and BOTANY it publishes the latest and best. Its DAIRY and STOCK department is in the hands of *practical* and *successful* dairymen and breeders. It employs the best talent on the subjects of TILLAGE and GENERAL FIELD OPERATIONS, DISEASES OF FARM STOCK, INSECT PESTS, POULTRY, RURAL ART AND ARCHITECTURE, WINDOW GARDENING, etc. All its departments are of every-day interest, and are conducted in a manner far above that of the common newspaper.

The Scientific Farmer aims to be the most thoroughly instructive farmer's magazine ever published. It strives to be entirely accurate in statement; to make science *practical*; knowledge *useful*; and the farm *profitable*.

Ever dignified and seeking the highest, it is still bright and cheery, and always on the look out for the brighter and pleasanter phases of farm life. Brimful of *meat*, and without a trashy line in it; and all for about **one quarter of a cent a day.**

There are few better uses to which an intelligent farmer can put a dollar than to send for this "*Farmer's Magazine*." A specimen number will be sent for ten cents, or 3 postage stamps. Address,

Scientific Farmer Co., Boston, Mass.

Feed the Plant and the Plant will Feed You.

Stockbridge Manures,

Originated by Prof. Levi Stockbridge, Professor of Agriculture in the

MASSACHUSETTS AGRICULTURAL COLLEGE.

These manures are made according to the analysis of plants for different crops by formulas worked out by Prof. Stockbridge, and supply the plant food in an available form, which is not obtained from the soil or air in sufficient quantity. These manures also recognize the fact that plants, like animals, differ and require different food or in different proportions to insure the best results at the least cost. They have been extensively used for FIVE YEARS and were used in 1877 on over 10,000 acres with very satisfactory results, many farmers raising from 75 to 100 BUSHELS of CORN per acre, and with like results on other crops. As they supply the principal elements of plant food which crops remove. **THEY DO NOT EXHAUST THE LAND, but actually leave it richer**; for being composed largely of chemicals they have a strong chemical action on the soil, and liberate other plant food. In short, the **Stockbridge Manures**, based as they are on the analysis of the crop, does away with *guess work* in fertilization, for they supply just what the crop wants.

They contain no *weed seeds*, and often cost less than the *hauling* of stable manure and are much more quickly applied. Send for Pamphlet for 1878, Mailed Free, containing PRICES, the experience of farmers all over the country who have used these manures, and other valuable information.

W. H. BOWKER & CO., Sole Agents and Manufacturers.

Also proprietors of Bowker's Hill and Drill Phosphates, and dealers in High Grade Agricultural Chemicals.

BOSTON AND NEW YORK.

Orders will also be received by

B. K. BLISS & SONS, 34 Barclay Street, New York.

NEW AGRICULTURAL BOOKS.



A book that ought to be in the hands of every owner of a farm, large or small, or of a garden plot.

POTATO PESTS.

Being an Illustrated account of the
COLORADO POTATO BEETLE,

And the other
Insect Foes of the Potato

IN NORTH AMERICA,

With Suggestions for their Repression
and Methods for their Destruction.

By **CHARLES V. RILEY, M. A., Ph. D.,**
(State Entomologist of Missouri.)

Finely Illustrated.

Price, postpaid, paper covers, 50 cents;
cloth, 75 cents.

AN EGG FARM. THE MANAGEMENT OF POULTRY IN LARGE NUMBERS.

By **H. H. STODDARD.**

Being a Series of Articles written for the *American Agriculturist*, with
Other Articles. Illustrated.

CONTENTS.—Introduction.—Plan of Farm.—Manner of Feeding.—Location of Farm.—Klud of Soil.—Crops on the Farm.—Supplying Water and Food.—Collecting and Storing Dry Earth.—Houses for Layers.—Feeding House for Winter.—Houses for Sitters.—Arrangements for Breeding Stock.—Fowls for Layers.—Fowls for Sitters.—Management of Breeding Stock.—Coops for Chickens.—Feeding Chickens.—Setting the Eggs.—Management of Sitting Fowls.—Testing the Eggs.—Winter Management.—House for Early Hatched Pullets.—Shelters for Fowls and Chickens.—Kind of Food.—Building for Storing and Cooking Food.—Management of Young Chickens.—Feeding and Sheltering Chickens.—Additional Buildings.—General Conclusions.—Farm Poultry House.—Poultry Farming.—Poultry Keeping as a Business.

Price, postpaid, paper covers, 50 cents; cloth 75 cents.

THE POULTRY YARD AND MARKET;

Or, A Practical Treatise on Galliniculture.

BY **PROF. A. CORBETT,**

Inventor of the New Process for Hatching Eggs and Raising Poultry, by means of Horse Manure alone, for which Gold and Bronze Medals, and several Diplomas, have been awarded by State and County Fairs, and the American Institute. Leading newspapers endorsed the valuable discovery of this simple process, which is fully described in this book, giving the valuable information acquired by 20 years' experience in Poultry Breeding.

CONTENTS.—Origin of Artificial Incubation.—Its Prolificness.—Particulars of what has been done in Artificial Incubation up to the present time.—The discovery of how Eggs could be Hatched in Manure.—The Researches of Prof. Corbett and his Discovery.—Prof. Corbett's Success.—Amount of Profit to be Made Annually by Anybody with Twelve Hens.—How Poultry Breeders can Raise Chickens in Manure, with Slight Expense, All the Year Round.—The Raising Poultry with Large Profits.—The Kind of Manure to be Used.—The Best Breed of Profitable Fowls.—Advice to the Ladies.—Important Hints How to Keep Poultry and Make Money.—Diseases and their Cure.—How to Fatten and Prepare for Market.—Opinions of the Press and Rewards given for this Process. Price, postpaid, 50 cents.

THE SHEPHERD'S MANUAL.

A practical treatise on The Sheep. Designed especially for American Shepherds. By Henry Stewart. Illustrated. Price, postpaid, \$1.50.


PLAY AND PROFIT IN MY GARDEN.

By **E. P. Roe.** Author of "Barriers Burned Away," etc. Price, postpaid, \$1.50.

Valuable Books for sale by B. K. Bliss & Sons.

Sent by mail, post-paid, at following prices.

Allen's (R. L. & L. F.) New American Farm Book, - - - - -	\$2 50	Jennings on the Horse and his Diseases, -	\$1 75
Allen's (L. F.) American Cattle, - - - - -	2 50	Jennings on Sheep, Swine and Poultry, -	1 75
Allen's (R. L.) American Farm Book, - - - - -	1 50	Jersey, Aiberney, and Guernsey Cow, - - -	1 50
Allen's (L. F.) Rural Architecture, - - - - -	1 50	Johnson's How Crops Feed, - - - - -	2 00
Allen's (R. L.) Diseases of Domestic Animals, - - - - -	1 00	Johnson's How Crops Grow, - - - - -	2 00
American Bird Fancier, - - - - -	30	Johnson's Peat and its Uses, - - - - -	1 25
American Rose Culturist, - - - - -	30	Johnston's Agricultural Chemistry, - - - -	1 75
American Weeds and Useful Plants, - - - -	1 75	Johnston's Elements of Agricultural Chemistry, - - - -	1 50
Atwood's Country and Suburban Houses, - - -	1 50	King's Beekeeper's Text Book. Paper. 40	
Barry's Fruit Garden, - - - - -	2 50	cts.; cloth, - - - - -	75
Bement's Rabbit Fancier, - - - - -	30	Klippart's Wheat Plant, - - - - -	1 75
Boumber's Method of Making Manures, - - - -	25	Lakey's Village and Country Houses, - - - -	5 00
Breck's New Book of Flowers, - - - - -	1 75	Leuchar's How to Build Hot-Houses, - - - -	1 50
Brill's Farm-gardening and Seed-growing, - - -	1 00	Lyman's Cotton Culture, - - - - -	1 50
Broom-Corn and Brooms, paper, 50c., cloth,	75	McClure's American Gentleman's Stable	
Brown's Taxidermist's Manual, - - - - -	1 00	Guide, - - - - -	1 00
Bruckner's American Manures, - - - - -	1 50	McClure's Diseases of the American Horse,	
Buel's Cider-maker's Manual, - - - - -	1 50	Cattle and Sheep, - - - - -	2 00
Burr's Vegetables of America, - - - - -	3 00	Miles on the Horse's Foot, - - - - -	75
Butler's Family Aquarium, - - - - -	75	Mohr on the Grape Vine, - - - - -	1 00
Butler on the Dog, - - - - -	2 00	My Vineyard at Lakeview, - - - - -	1 25
Canary Birds. Paper, 50 cts.; cloth, - - - -	75	Nichol's Chemistry of the Farm and Sea, - - -	1 25
Chorlton's Grape-grower's Guide, - - - - -	75	Norris' American Fish Culture, - - - - -	1 75
Cobbett's American Gardener, - - - - -	75	Norton's Scientific Agriculture, - - - - -	75
Coburn's Swine Husbandry, - - - - -	1 75	Onions—How to Raise them Profitably, - - -	20
Cole's American Fruit Book, - - - - -	75	Our Farm of Four Acres, Paper, 30c., cloth,	60
Cole's American Veterinarian, - - - - -	75	Pardee on Strawberry Culture, - - - - -	75
Cooked Food for Domestic Animals, - - - - -	20	Parsons on the Rose, - - - - -	1 50
Cook's Manual of the Apiary, - - - - -	30	Pedder's Land Measurer, - - - - -	60
Corbett's Poultry Yard and Market, paper,		Percheron Horse, - - - - -	1 00
50 cts.; cloth, - - - - -	75	Potato Culture. (Prize Essay.) - - - - -	25
Dadd's Modern Horse Doctor, 12mo., - - - -	1 50	Quincy's Mysteries of Bee-keeping, - - - -	1 50
Dadd's American Cattle Doctor, 12mo., - - - -	1 50	Quincy (Hon. Josiah) on Soiling Cattle, - - - -	1 25
Dadd's American Reformed Horse Book, - - - -	2 50	Quinn's Money in the Garden, - - - - -	1 50
Dana's Muck Manual, - - - - -	1 25	Quinn's Pear Culture for Profit, - - - - -	1 00
De Voe's Market Assistant, - - - - -	2 51	Randall's Fine Wool Sheep Husbandry, - - - -	1 00
Downing's Landscape Gardening, - - - - -	6 50	Randall's Sheep Husbandry, - - - - -	1 50
Dwyer's Horse Book, - - - - -	2 00	Rarey and Knowlson's Horse Tamer, - - - -	50
Eastwood on Cranberry, - - - - -	75	Richardson on the Dog, Paper, 30c.; cloth,	60
Entfield's Indian Corn, - - - - -	1 00	Riley's Potato Pests, Paper, 50 cts.; cloth,	75
Every Horse Owner's Cyclopaedia, - - - - -	3 75	Rivers's Miniature Fruit Garden, - - - - -	1 00
Every Woman Her Own Flower Gardener,		Roe's Manual on the Culture of Small	
Paper, 50 cts.; cloth, - - - - -	1 00	Fruits, - - - - -	50
Field's Pear Culture, - - - - -	1 25	Roe's Play and Profit in my Garden, - - - -	1 50
Flax Culture, [Seven Prize Essays.] - - - -	50	Saunders's Domestic Poultry, Paper 40 cts.;	
Flint (Charles L.) on Grasses, - - - - -	2 50	cloth, - - - - -	75
French's Farm Drainage, - - - - -	1 50	Schenek's Gardener's Text-Book, - - - - -	75
Fuller's Forest Tree Culturist, - - - - -	1 50	Skillful Housewife, - - - - -	75
Fuller's Grape Culturist, - - - - -	1 50	Stewart's American Farmer's Horse Book,	3 00
Fuller's Illustrated Strawberry Culturist,		Stewart's Irrigation for the Farm, Garden,	
Fuller's Small Fruit Culturist, - - - - -	1 50	and Orchard, - - - - -	1 50
Fulton's Peach Culture, - - - - -	1 50	Stewart's Sorghum and its Products, - - - -	1 50
Gevelin's Poultry Breeding, - - - - -	1 25	Stewart's Stable Book, - - - - -	1 50
Grant's Beet Root Sugar, - - - - -	1 25	Stoddard's An Egg Farm. 50 cts.; cloth, - - -	75
Gregory on Cabbages, - - - - -	30	Stonehenge on the Horse in Stable and	
Gregory on Carrots, - - - - -	30	Field. American edition, 12mo., - - - - -	2 00
Gregory on Mangold Wurtzels, etc. - - - - -	30	Tegetmeier's Poultry Book, - - - - -	9 00
Gregory on Onion Raising, - - - - -	30	The Rifle: Its Theory and Practice, - - - -	50
Gregory on Squashes, - - - - -	30	Thomas's American Fruit Culturist. New	
Guenon on Milch Cows, - - - - -	75	edition, - - - - -	3 75
Harlan's Farming with Green Manures, - - - -	50	Thomas's Farm Implements and Machinery,	
Harris's Insects Injurious to Vegetation,		- - - - -	1 50
Plain, \$4; colored engravings, - - - - -	6 50	Thompson's Food of Animals, - - - - -	1 00
Harris on the Pig, - - - - -	1 50	Tobacco Culture. By fourteen experienced	
Hazard's Butter and Butter Making, - - - -	25	cultivators, - - - - -	25
Hedges on Sorgho, or the Northern Sugar		Turner's Cotton Planter's Manual, - - - - -	1 50
Plant, - - - - -	1 50	Ville's Chemical Manures, - - - - -	50
Hemslley's Hardy Trees, Shrubs and Plants		Warder's American Pomology, - - - - -	3 00
Hemlerson's Gardening for Pleasure, - - - -	1 50	Warder's Hedges and Evergreens, - - - - -	1 50
Henderson's Gardening for Profit, - - - - -	1 50	Waring's Draining for Profit and Health, - - -	1 50
Henderson's Practical Floriculture, - - - - -	1 50	Waring's Earth Closets and Earth Sewage,	50
Hooper's Dog and Gun. Paper 30c.; cloth,	60	Waring's Elements of Agriculture, - - - - -	1 00
Hoopes's Book of Evergreens, - - - - -	3 00	White's Cranberry Culture, - - - - -	1 25
Hop Culture. By nine experienced culti-		White's Gardening for the South, - - - - -	2 00
vators, - - - - -	30	Wright's Brahma Fowl, - - - - -	2 50
Howard's Grasses and Forage Plants at the		Wright's Practical Poultry-keeper, - - - -	2 00
South, - - - - -	30	Yonatt and Spooner on the Horse, - - - - -	1 50
How to Make Candy, - - - - -	50	Yonatt and Martin on Cattle, - - - - -	1 50
Hunter and Trapper, - - - - -	1 00	Yonatt and Martin on the Hog, - - - - -	1 00
Jennings on Cattle and their Diseases, - - - -	1 75	Yonatt on Sheep, - - - - -	1 00
Jennings' Horse Training Made Easy, - - - -	1 25	Yonatt on the Dog, - - - - -	2 50



SMALL FRUITS

FERNERIES

WINDOW-GARDENS

SEED SOWER

LAWN MOWERS

GARDEN IMPLEMENTS

AGRIC. BOOKS

BIRD SEED

PLANTS B. K. BLISS & SONS. BULBS.

GARDEN REQUISITES & RUSTIC WORK.

SEED POTATOES, GRASS SEEDS & FERTILIZERS.

34 B. K. BLISS & SONS 34 GARDEN, FIELD & FLOWER SEEDS.

PATTERSON ENG.

V. WOLBERGER

B. K. BLISS & SONS'
Seed Warehouse,
34 Barclay St., New York.

