

THE  
AMERICAN  
NATURAL  
HISTORY



W. T. HORNADAY


LABORATORY  
OF ORNITHOLOGY  
LIBRARY



*Dept of  
Patricia Sedde*

CORNELL UNIVERSITY LIBRARY





# Cornell University Library

The original of this book is in  
the Cornell University Library.

There are no known copyright restrictions in  
the United States on the use of the text.

<http://www.archive.org/details/cu31924022552511>





THE AMERICAN  
NATURAL HISTORY

“The concise and precise phraseology of science, admirable though it be for the use of those who have been trained to employ it, is to others not only misleading, but it may be repulsive.”—G. BROWN GOODE.

“The highest type of scientific writing is that which sets forth useful scientific facts, in language which is interesting, and easily understood by the millions who read.”—L. A. MANN.







Painted by CARL RUNGUIS.

THE AMERICAN MOOSE, IN NEW BRUNSWICK.





# THE AMERICAN NATURAL HISTORY

A FOUNDATION OF USEFUL KNOWLEDGE OF  
THE HIGHER ANIMALS OF NORTH AMERICA

BY

WILLIAM T. HORNADAY

DIRECTOR OF THE NEW YORK ZOOLOGICAL PARK; AUTHOR OF  
"TWO YEARS IN THE JUNGLE," ETC.

ILLUSTRATED BY 227 ORIGINAL DRAWINGS BY BEARD, RUNGIUS,  
SAWYER, AND OTHERS, 116 PHOTOGRAPHS, CHIEFLY BY SANBORN,  
KELLER, AND UNDERWOOD, AND NUMEROUS CHARTS AND MAPS

CHARLES SCRIBNER'S SONS  
NEW YORK :::::::::::::::::::: MCM VI

OPM TH  
OL  
151  
H81  
1906

COPYRIGHT, 1904, BY  
WILLIAM T. HORNADAY

Published, April, 1904

SPECIAL NOTICE

The publishers hereby give warning that the unauthorized use of illustrations, charts, or maps from this book is expressly forbidden.



## PREFACE

By natural inclination, every child is interested in animals. Whenever a grown person is not so interested, it is positive proof that the natural instincts of childhood either have been turned aside, or stifled by lack of opportunity to live and grow. The love for animals is, I believe, even more universal than the love for music.

Whenever I try to sum up the amount of living interest, and also genuine delight, that is yielded by even a very modest acquaintance with the higher forms of life, "I would that my tongue could utter the thoughts that arise in me!" It seems a pity that so many appreciative persons should lose so much of life through lack of acquaintance with about three hundred important and well-chosen species of animals.

In these days of struggle and stress for Place and Power, and in these nights of insomnia and nerves, there are few side issues more restful or more pleasantly diverting to a tired brain than an active interest in some branch of natural history. A hunt for the life history of a fine animal species is next in restfulness to a real hunt, over the fields and far away, with all cares and worries left behind.

The foregoing is for the eyes of adult readers. Argument is not necessary to convince young people that a mighty host of interesting things awaits every one who sets foot in the field of Nature. To-day, the all-absorbing question is—how can Nature be made available to the young?

This book is one of my two answers to that question; and it is particularly addressed to teachers and parents. It is intended to be a plain, practical, common-sense answer, presented in a systematic and scientific way. The author assumes that fifteen years of earnest thought, and conferences with scores of teachers on the subject of natural history teaching in American public schools, may fairly entitle him to a hearing.

Briefly stated, the situation to-day is as follows:

The scientific "zoology" is suitable only for students in the higher colleges and universities. Between it and the "nature study" books of the grammar schools there exists a chasm that is wide and deep.

The "nature studies" of some of our city schools are good for young pupils, from ten to fourteen years of age; but they are insufficient for those between fourteen years and university age.

Students in the highest grammar-school grades, the high schools, normal schools, academies and small colleges are so inadequately equipped for the study of natural history that *fully ninety-five per cent. of them, including also the great mass of students from the higher colleges and universities, enter active life ignorant of even the most important forms of the wild life of our own country!* If this statement can be disproved, the author will be delighted to withdraw it, and apologize.

While the "nature-study" teaching of the present day is acceptable and commendable for very young pupils, tending to arouse their interest and prepare their minds for more serious work, its sphere is strictly limited, and it is a mistake to carry it too far. Valuable

and permanent results in the study of animal life cannot be achieved by turning in the class-room a kaleidoscope filled with a chaotic mass of birds, butterflies, flowers, frogs and trees. Object-teaching is excellent, if rightly conducted. But the object can easily become a fetich; and all fetich-worship is dangerous to its devotees. Twenty-five years hence, some of the courses of study of the year 1903 will be regarded as educational curiosities. Even the finest lobster or grasshopper should not be held so close to the eye that it obscures all the remainder of the animal kingdom.

There is no royal road to a real acquaintance with living animals. Entertaining and truthful story-books about quadrupeds and birds are excellent in their way, but they do not, and cannot, go down to bed-rock, and lay foundations on which the pupil can build for aye. It has been decreed by Nature that he who will not work shall not know her. There is no process by which the secrets of Nature can be placed automatically in a giddy mind.

The author maintains in this volume, and also out of it, that *System* is the only master-key by which the doors of Animate Nature can be unlocked. Even with boys and girls fifteen years of age, the foundations of natural history classification must not be ignored! Let them but begin right, and the structure is bound to rise. But *beware of all chaotic jumbles of unrelated facts!*

This volume is intended as builder's "filling" in the chasm that now exists between the technical "zoology" of the college and the "nature-study" lessons of the common schools. To-day, I am certain that many nature-study teachers dislike their work solely because they lack suitable sources of information. Surely it is unnecessary to suggest to any intelligent and sincere teacher that it is possible to utilize only a portion of this book, by selecting the subjects best adapted to each particular class, and passing over the others.

Among the writers of manuals of zoology, it is now customary to begin with the lowest and least interesting forms of life, and work upward toward the highest. That will answer for the advanced student—if he chooses to have it so; but for middle-grade students and readers at home it is decidedly wrong. All elementary lessons in natural history should begin with Nature's most important facts, and first bring forward her most interesting animals. To begin with the grasshopper, and struggle through a hundred dreary pages of anatomy and low forms of life, before reaching a creature of personality and intelligence, is too much for the patience of any active school-boy who wishes "to know about animals."

Anatomy is necessary to the advanced student; but in a book for schools and the general reader, it is easily carried too far. As with human beings, the first thing to be learned about an animal is its place in Nature, and after that, its personality. It is only the scientific specialist who wishes to know first about its mandibular symphysis, the geography of its sutures, and the size of its auditory bullae.

As the reader will observe, I have striven to accomplish two ends: (1) to make clear each animal's place in the great system of Nature, and (2) to introduce the animal in such a manner as to enable the reader to become personally acquainted with it. The subjects chosen for introduction are not confined to any one section of our country, but represent all North America, and even lands beyond. For the purpose of avoiding wide gaps, several important foreign animals have been included.

At this point I wish to record a grateful acknowledgment to Mr. Andrew Carnegie, for



his interest in the author's plans for introducing the study of natural history in schools, and for encouragement at a time when it was most needed.

The manuscripts and proofs relating to mammals have been read, criticised and corrected by Dr. T. S. Palmer, Assistant Chief of the Biological Survey, Washington, D. C. Through Dr. Palmer's advice, the author's old-fashioned preferences on certain points of nomenclature were abandoned, and the names of orders, families, genera and species were brought down to date. It is due to him that in our nomenclature we are in reality a trifle in advance of the times rather than behind them.

Similar valuable service has been rendered the section on Birds by Mr. C. William Beebe, Curator of Birds, and those on the Reptiles and Amphibians were read and corrected by Mr. Raymond L. Ditmars, Curator of Reptiles, in the New York Zoological Park. The portion treating of Fishes received critical attention from Mr. Charles H. Townsend, Director of the New York Aquarium, but in fairness to him it must be stated that he is in no way responsible for the author's arrangement of the Orders of Fishes.

To each of the gentlemen named above I offer a most grateful acknowledgment for timely and valuable services, and desire to assure the reader that for any shortcomings that may appear in the finished book, they are not in the least responsible.

In the text of this work I have endeavored to give due credit for the noteworthy facts quoted from other authors. Practically the only instances wherein this has not always been possible are those involving the geographic ranges of species, wherein combinations of authorities are the rule rather than the exception. To cover all possible omissions, I desire to mention here the names of the authors from whom I have derived many facts, but chiefly regarding distribution, and I gratefully acknowledge indebtedness to Mr. D. G. Elliot's "Synopsis of the Mammals of North America and the Adjacent Seas"; to the many papers on our Mammalia by Dr. C. Hart Merriam, Dr. T. S. Palmer, and Mr. Vernon Bailey; to Mrs. Florence Merriam Bailey's "Birds of the Western United States." Mr. Frank M. Chapman's "Birds of Eastern North America," Dr. A. K. Fisher's "Hawks and Owls," and Mrs. Mabel Osgood Wright's "Birdcraft"; to Dr. Leonhard Stejneger's "Poisonous Snakes of North America"; Prof. E. D. Cope's "Crocodylians, Lizards and Snakes of North America," and Dr. H. Gadow's "Amphibia and Reptiles"; to Dr. David S. Jordan and Dr. Barton W. Evermann's "Fishes of North and Middle America," and "American Food and Game Fishes," and to Mr. Richard Lydekker's "Royal Natural History."

Naturally, I have drawn freely upon the zoological knowledge that has been accumulated in the New York Zoological Park during its existence.

A final word must be added regarding the illustrations. Probably no other author ever had a more tempting opportunity for completely filling a volume with photographs of animals. But, while I am an ardent admirer of the best results in animal photography, and a diligent user of them, I also recognize the limitations of the camera.

The demands of a zoological illustration are inexorable; and all too often the camera ignores some of them. A *perfect* zoological portrait of an animal must possess clear and distinct outlines, showing a side view, and perfect details. A picture *sans* feet, tail, ears, eyes or legs, is not a portrait; and a ball of fur, even though photographed, is not necessarily an animal. Very often, also, the most perfect photograph of a spiritless animal in captivity utterly fails to convey a just and adequate impression of the species as it is seen at its best, on its native heath.

Because of the limitations of the camera, several thousand dollars have been expended upon the beautiful drawings by Messrs. J. Carter Beard, Carl Rungius, Edmund J. Sawyer and a few other artists, which adorn as well as illustrate this work. In addition to these, about one hundred and sixteen particularly excellent photographs have been made, of specially selected subjects, by Messrs. Elwin R. Sanborn, Ernest F. Keller, W. Lyman Underwood, R. J. Beck, and a few other experts in animal photography. With but very few exceptions, the illustrations which appear in this book have been made expressly for it, and now appear for the first time. The author is indebted for the loan of several from the publications of the New York Zoological Society.

Now that the last page save the preface has been set up, locked fast and turned into a plate of cold metal, the hour for regret has struck. I know that my proof-reading has not been perfect, and that various errors may be found by those who watch for them. In view of the patient and even tireless efforts and the generous expenditures which Messrs. Charles Scribner's Sons have bestowed upon this volume, the author deeply regrets that his own share of the work is not as perfect as theirs. For the reader's sake, also, he wishes that he could have done better.

W. T. H.

BEDFORD PARK, New York City.

# CONTENTS

INTRODUCTION . . . . .	Page xix
------------------------	-------------

## BOOK I—MAMMALS

CHAPTER I.—ORDERS OF MAMMALS . . . . .	3
--	---

CHAPTER II.—ORDER OF APES AND MONKEYS . . . . .	<i>PRIMATES</i> 7
---	-------------------

	<i>Page</i>		
ANTHROPOID APES . . . . .	7	NEW WORLD MONKEYS . . . . .	14
OLD WORLD MONKEYS AND BAROONS . . . . .	13	FAMILY OF MARMOSETS . . . . .	16
		SUBORDER OF LEMURS . . . . .	17

CHAPTER III.—ORDER OF FLESH-EATING MAMMALS . . . . .	<i>FERAE</i> 18
--	-----------------

	<i>Page</i>		
CAT FAMILY . . . . .	18	BEAR FAMILY . . . . .	32
DOG FAMILY . . . . .	22	FULL LIST OF THE BEARS OF NORTH AMER-	
NORTH AMERICAN FOXES . . . . .	24	ICA . . . . .	35
SMALL FUR-BEARERS . . . . .	27	RACCOON FAMILY . . . . .	41

CHAPTER IV.—ORDER OF SEALS AND SEA-LIONS . . . . .	<i>PINNIPEDIA</i> 43
--	----------------------

	<i>Page</i>		
SEA-LION FAMILY . . . . .	44	SEAL FAMILY . . . . .	52
REVIEW OF FUR SEAL HISTORY . . . . .	48	WALRUS FAMILY . . . . .	53

CHAPTER V.—ORDER OF MOLES AND SHREWS . . . . .	<i>INSECTIVORA</i> 56
--	-----------------------

	<i>Page</i>		
MOLE FAMILY . . . . .	57	SHREW FAMILY . . . . .	58

CHAPTER VI.—ORDER OF BATS . . . . .	<i>CHIROPTERA</i> 59
-------------------------------------	----------------------

	<i>Page</i>		
FAMILY OF LEAF-NOSED BATS . . . . .	62	FAMILY OF FALSE VAMPIRES . . . . .	65
FAMILY OF FREE-TAILED BATS . . . . .	63	FAMILY OF HORSESHOE BATS . . . . .	66
FAMILY OF COMMON BATS . . . . .	64	FAMILY OF FRUIT-EATING BATS . . . . .	66

CHAPTER VII.—ORDER OF GNAWING ANIMALS . . . . .	<i>GLIRES</i> or <i>RODENTS</i> 68
---	------------------------------------

	<i>Page</i>		
SQUIRREL FAMILY . . . . .	68	JUMPING MOUSE FAMILY . . . . .	92
SEWELLEL FAMILY . . . . .	80	POCKET GOPHER FAMILY . . . . .	93
BEAVER FAMILY . . . . .	80	PORCUPINE FAMILY . . . . .	94
FAMILY OF MICE AND RATS . . . . .	83	PIKA FAMILY . . . . .	95
TYPICAL NORTH AMERICAN MICE AND RATS . . . . .	84	HARE AND RABBIT FAMILY . . . . .	95
CHEEK-POUCH MICE AND RATS . . . . .	91		

CHAPTER VIII.—ORDER OF HOOFED ANIMALS . . . . .	<i>UNGULATA</i> 99
---	--------------------

	<i>Page</i>		
CATTLE AND SHEEP FAMILY . . . . .	99	MEASUREMENTS OF LARGE CARIBOU ANTLERS . . . . .	138
MEASUREMENTS OF MOUNTAIN SHEEP HORNS . . . . .	112	PECCARY FAMILY . . . . .	143
PRONG-HORNED ANTELOPE FAMILY . . . . .	115	TAPIR FAMILY . . . . .	144
DEER FAMILY . . . . .	118		

CHAPTER IX.—ORDER OF WHALES AND PORPOISES . . . . .		<i>CETE</i> 1 <sup>a</sup>
<i>Page</i>		
FAMILY OF BALEEN WHALES . . . . .	147	SPERM WHALE FAMILY . . . . . 1 <sub>4</sub> DOLPHIN AND PORPOISE FAMILY . . . . . ' 1
CHAPTER X.—ORDER OF SEA-COWS . . . . .		<i>SIRENIA</i> 1
FAMILY OF MANATEES . . . . .	153	FAMILY OF THE RHYTINA . . . . . 154
FAMILY OF DUGONGS . . . . .	154	
CHAPTER XI.—ORDER OF TOOTHLESS MAMMALS . . . . .		<i>EDENTATA</i> 156
ARMADILLO FAMILY . . . . .	156	SLOTH FAMILY . . . . . 159
FAMILY OF ANT-EATERS . . . . .	158	
CHAPTER XII.—ORDER OF DIGGERS . . . . .		<i>EFFODIENTIA</i> 161
PANGOLIN FAMILY . . . . .	161	AARD-VARK FAMILY . . . . . 162
CHAPTER XIII.—ORDER OF POUCHED MAMMALS . . . . .		<i>MARSUPIALIA</i> 1 <sup>c</sup>
KANGAROO FAMILY . . . . .	164	OPOSSUM FAMILY . . . . .
CHAPTER XIV.—ORDER OF EGG-LAYING MAMMALS . . . . .		<i>MONOTREMATA</i> 167
DUCK-BILL FAMILY . . . . .	167	ECHIDNA FAMILY . . . . . 168

**BOOK II—BIRDS**

CHAPTER XV.—INTRODUCTION TO THE BIRD-WORLD . . . . .		171
DECREASE IN BIRD LIFE . . . . .	171	ORDERS OF LIVING BIRDS . . . . . 175
CHAPTER XVI.—ORDER OF PERCHERS AND SINGERS . . . . .		<i>PASSERES</i> 179
THRUSH FAMILY . . . . .	181	WAXWING FAMILY . . . . . 192
KINGLET FAMILY . . . . .	183	SWALLOW FAMILY . . . . . 193
NUTHATCH AND TITMOUSE FAMILY . . . . .	184	TANAGER FAMILY . . . . . 194
TREE-CREEPER FAMILY . . . . .	185	FINCH AND SPARROW FAMILY . . . . . 195
WRENS AND CAT-BIRDS . . . . .	186	BLACKBIRD FAMILY . . . . . 199
DIPPER FAMILY . . . . .	187	CROW FAMILY . . . . . 202
WARBLER FAMILY . . . . .	188	HORNED LARK FAMILY . . . . . 206
VIREO FAMILY . . . . .	191	FLYCATCHER FAMILY . . . . . 206
SHRIKE FAMILY . . . . .	191	
CHAPTER XVII.—ORDER OF ODD FAMILIES . . . . .		<i>MACROCHIRES</i> 207
GOATSUCKER FAMILY . . . . .	207	HUMMING-BIRD FAMILY . . . . . 208
SWIFT FAMILY . . . . .	208	
CHAPTER XVIII.—ORDER OF WOODPECKERS . . . . .		<i>PICI</i> 210
CHAPTER XIX.—ORDER OF CUCKOOS AND KINGFISHERS . . . . .		<i>COCCYGES</i> 214
CUCKOO FAMILY . . . . .	214	KINGFISHER FAMILY . . . . . 215
CHAPTER XX.—ORDER OF PARROTS AND MACAWS . . . . .		<i>PSITTACI</i> 216

	<i>Page</i>
CHAPTER XXI.—ORDER OF BIRDS OF PREY . . . . .	<i>RAPTORES</i> 218
	<i>Page</i>
HORN-OWL FAMILY . . . . .	218
BARN-OWL FAMILY . . . . .	220
	HAWKS AND EAGLES . . . . . 225
	VULTURE FAMILY . . . . . 232
CHAPTER XXII.—ORDER OF PIGEONS AND DOVES . . . . .	<i>COLUMBAE</i> 237
CHAPTER XXIII.—ORDER OF UPLAND GAME-BIRDS . . . . .	<i>GALLINAE</i> 241
GROUSE FAMILY . . . . .	242
	PHEASANT FAMILY . . . . . 250
CHAPTER XXIV.—ORDER OF SHORE-BIRDS . . . . .	<i>LIMICOLAE</i> 251
CHAPTER XXV.—ORDER OF CRANES, RAILS, AND COOTS . . . . .	<i>PALUDICOLAE</i> 255
CRANE FAMILY . . . . .	255
	FAMILY OF RAILS . . . . . 257
CHAPTER XXVI.—ORDER OF HERONS, STORKS, AND IBISES . . . . .	<i>HERODIONES</i> 259
HERON FAMILY . . . . .	259
STORK FAMILY . . . . .	263
	IBIS FAMILY . . . . . 263
	SPOONBILL FAMILY . . . . . 264
CHAPTER XXVII.—ORDER OF FLAMINGOES . . . . .	<i>ODONTOGLOSSAE</i> 266
CHAPTER XXVIII.—ORDER OF DUCKS, GEESE, AND SWANS . . . . .	<i>ANATIDAE</i> 267
ORDERS OF SWIMMING BIRDS . . . . .	267
	AN OBJECT LESSON IN BIRD PROTECTION . 276
CHAPTER XXIX.—ORDER OF FULLY WEB-FOOTED BIRDS . . . . .	<i>STEGANOPODES</i> 284
PELICAN FAMILY . . . . .	284
CORMORANT FAMILY . . . . .	287
DARTER FAMILY . . . . .	287
	GANNET FAMILY . . . . . 288
	MAN-O'-WAR-BIRD FAMILY . . . . . 290
CHAPTER XXX.—ORDER OF TUBE-NOSED SWIMMERS . . . . .	<i>TUBINARES</i> 292
ALBATROSS FAMILY . . . . .	292
	FULMAR FAMILY . . . . . 294
CHAPTER XXXI.—ORDER OF LONG-WINGED SWIMMERS . . . . .	<i>LONGIPENNES</i> 296
GULLS AND TERNS . . . . .	296
SKIMMER FAMILY . . . . .	298
	SKUA AND JAEGER FAMILY . . . . . 299
CHAPTER XXXII.—ORDER OF WEAK-WINGED DIVING BIRDS . . . . .	<i>PYGOPODES</i> 300
GREBE FAMILY . . . . .	300
LOON FAMILY . . . . .	301
	CLIFF-DWELLERS OF THE SEA . . . . . 302
CHAPTER XXXIII.—ORDER OF FLIGHTLESS DIVERS . . . . .	<i>IMPENNES</i> 307
CHAPTER XXXIV.—ORDER OF WINGLESS LAND BIRDS . . . . .	<i>RATITAE</i> 309

**BOOK III—REPTILES**

CHAPTER XXXV.—INTRODUCTION TO THE CLASS OF REPTILES . . . . .	313
ORDERS OF LIVING REPTILES . . . . .	314

		<i>Page</i>
CHAPTER XXXVI.—ORDER OF CROCODILES AND ALLIGATORS		<i>CROCODILIA</i> 317
SYNOPSIS OF THE CROCODILIANS . . . . . 317		
	<i>Page</i>	
	CROCODILE FAMILY . . . . .	319
	AMERICAN SPECIES OF CROCODILIANS . . . . .	320
CHAPTER XXXVII.—ORDER OF TORTOISES, TERRAPINS, AND TURTLES		<i>CHELONIA</i> 323
SYNOPSIS OF THE ORDER OF TORTOISES AND TURTLES . . . . . 324		
TORTOISE FAMILY . . . . .	SNAPPING TERRAPINS . . . . .	328
MUD-TERRAPIN FAMILY . . . . .	SOFT-SHELLED "TURTLES" . . . . .	329
SMOOTH-SHELLED TERRAPINS . . . . .	HARD-SHELLED SEA TURTLES . . . . .	330
	LEATHERY-SHELLED SEA TURTLES . . . . .	331
CHAPTER XXXVIII.—ORDER OF LIZARDS . . . . .		<i>LACERTILIA</i> 333
CHAPTER XXXIX.—ORDER OF SERPENTS . . . . .		<i>OPHIDIA</i> 337
GENERAL CHARACTERS OF SERPENTS . . . . .	HARMLESS SNAKES OF THE UNITED STATES . . . . .	343
FOOD OF SERPENTS . . . . .	POISONOUS SNAKES OF NORTH AMERICA . . . . .	347
POPULAR QUESTIONS AND MISAPPREHENSIONS . . . . .	SPECIES OF RATTLESNAKES . . . . .	349
LARGEST SPECIES OF SERPENTS . . . . .	SNAKE POISONS, AND THEIR TREATMENT . . . . .	353

**BOOK IV—AMPHIBIANS**

CHAPTER XL.—INTRODUCTION TO THE CLASS OF AMPHIBIANS . . . . .		359
GENERAL CHARACTERS OF AMPHIBIANS . . . . .	BIRD'S-EYE VIEW OF THE AMPHIBIANS . . . . .	360
CHAPTER XLI.—ORDER OF FROGS AND TOADS . . . . .		<i>ECAUDATA</i> 361
FAMILY OF WATER FROGS . . . . .	TOAD FAMILY . . . . .	364
TREE-FROG FAMILY . . . . .	TONGUELESS FROGS . . . . .	364
CHAPTER XLII.—ORDER OF TAILED AMPHIBIANS . . . . .		<i>URODELA</i> 366
FAMILY OF SALAMANDERS . . . . .	FREE-GILLED SALAMANDERS . . . . .	370
NEWTs, OR TRITONS . . . . .	TWO-LEGGED SALAMANDERS . . . . .	370
FAMILY OF AMPHIUMAS . . . . .	ORDER OF WORM-LIKE AMPHIBIANS . . . . .	371

**BOOK V—FISHES**

CHAPTER XLIII.—INTRODUCTION TO THE CLASS OF FISHES . . . . .		375
FISHERY INDUSTRIES AND FISH PROPAGATION . . . . .	THE ORDERS OF LIVING FISHES . . . . .	378
DISTRIBUTION OF EGGS AND LIVE FISH . . . . .		377
CHAPTER XLIV.—ORDER OF THE CONNECTING-LINK FISHES		<i>SIRENOIDEI</i> 380
CHAPTER XLV.—ORDER OF THE SPINY-FINNED FISHES . . . . .		<i>ACANTHOPTERI</i> 382
BASSES AND SUNFISHES . . . . .	MISCELLANEOUS SPINY-FINNED FISHES . . . . .	388
SEA-BASS FAMILY . . . . .	SNAPPER FAMILY . . . . .	391
PERCH AND PIKE-PERCH FAMILY . . . . .	ODD FISHES OF THE SPINY-FINNED ORDER . . . . .	392
CHAPTER XLVI.—ORDER OF PIKES . . . . .		<i>HAEPLIOMI</i> 394

CONTENTS

		<i>Page</i>
CHAPTER XLVII.—ORDER OF TROUT AND SALMON . . .	<i>ISOSPONDYLI</i>	396
	<i>Page</i>	
SALMON FAMILY . . . . .	396	NORTH AMERICAN TROUT . . . . . 397
SUBDIVISION OF NORTH AMERICAN TROUTS AND		THE SALMON GROUP . . . . . 400
CHARRS . . . . .	397	AMERICAN SALMON . . . . . 400
CHAPTER XLVIII.—ORDER OF FLYING-FISHES . . .	<i>SYNENTOGNATHI</i>	409
CHAPTER XLIX.—ORDER OF SOLID-JAW FISHES . . .	<i>PLECTOGNATHI</i>	410
CHAPTER L.—ORDER OF SUCKERS AND MINNOWS . .	<i>PLECTOSPONDYLI</i>	412
CHAPTER LI.—ORDER OF HALF-GILLED FISHES . . .	<i>HEMIBRANCHII</i>	415
CHAPTER LII.—ORDER OF CATFISHES . . . . .	<i>NEMATOGNATHI</i>	416
CHAPTER LIII.—ORDER OF FLAT-FISHES . . . . .	<i>HETEROSOMATA</i>	418
CHAPTER LIV.—ORDER OF FOOT-FISHES . . . . .	<i>PEDICULATI</i>	420
CHAPTER LV.—ORDER OF EELS . . . . .	<i>APODES</i>	421
CHAPTER LVI.—ORDER OF PIPE-FISHES AND SEA-HORSES .	<i>LOPHOBRANCHI</i>	423
CHAPTER LVII.—ORDER OF THE DOGFISH . . . . .	<i>HALECOMORPHI</i>	424
CHAPTER LVIII.—ORDER OF GAR-FISHES, OR GANOIDS . .	<i>GINGLYMODI</i>	425
CHAPTER LIX.—ORDER OF STURGEONS . . . . .	<i>GLANIOSTOMI</i>	427
CHAPTER LX.—ORDER OF THE PADDLE-FISH . . . . .	<i>SELACHOSTOMI</i>	429
CHAPTER LXI.—ORDER OF THE CHIMERAS . . . . .	<i>CHIMAEROIDEI</i>	431
CHAPTER LXII.—ORDER OF SHARKS . . . . .	<i>SQUALI</i>	432
CHAPTER LXIII.—ORDER OF RAYS AND SKATES . . . . .	<i>RAIAE</i>	434
CHAPTER LXIV.—LOWEST CLASSES OF VERTEBRATES . . . . .		437
LAMPREYS . . . . .	437	LANCELETS . . . . . 438





# LIST OF ILLUSTRATIONS

	PAGE		PAGE
Albatross, Black-Footed .....	293	Canvas-Back Duck .....	275
Alligator .....	316	Cardinal .....	198
Alligator, Skull of .....	318	Caribou, Woodland .....	133
Anaconda, Yellow .....	341	Carp, German Scaled .....	413
Angel Fish .....	387	Cassowary, Ceram .....	309
Angler .....	420	Cat-Bird .....	187
Ant-Eater, Great .....	158	Chickadee .....	184
Antelope, Prong-Horned .....	116	Chipmunk, Eastern .....	72
Antlers of Alaskan Moose .....	142	Chipmunk, Western .....	73
Antlers? Do Elk Shed Their—4 figures .....	119	Chimpanzee, A Dressed-Up .....	10
Antlers of Greenland Caribou—2 figures .....	135	Chimpanzee, Young Female .....	9
Antlers of Kenai Caribou .....	134	Chimera, Spotted .....	431
Armadillos, Three-Banded and Six-Banded .....	157	Coach-Whip Snake .....	345
Auklet, Rhinoceros .....	304	Condor .....	234
Axolotl, Two Lives of the .....	367	Congo "Snake" .....	369
		Coot .....	258
Baboon, Dead Gelada .....	14	Copperhead Snake .....	352
Badger .....	32	Cormorant .....	289
Bass, Black Sea .....	385	Coyote .....	23
Bass, Calico .....	383	Crane, Whooping .....	256
Bass, Striped .....	386	Creper, Brown .....	186
Bass, Small-Mouthed Black .....	383	Crocodile, Florida .....	321
Bat, Bonneted .....	63	Crocodile, Skull of Indian .....	318
Bat, Bornean Naked .....	59	Crocodile, Skull of Florida .....	318
Bat, California Leaf-Nosed .....	62	Crocodile, Skull of Orinoco .....	318
Bat, Flower-Nosed .....	62	Cross-bill, American .....	195
Bats, Fruit-Eating .....	67	Cuckoo, Yellow-Billed .....	214
Bat, Hammer-Headed .....	66		
Bat, Red .....	65	Deer, Mule, in the Bad-Lands .....	125
Bear, Alaskan Brown .....	33	Deer, Mule, with Antlers in Velvet .....	127
Bear, American Black .....	39	Deer, White-Tailed .....	128
Bear, Glacier .....	40	Deer, White-Tailed, "Freak" antlers of .....	131
Bear, Grizzly, at Home .....	38	Deer, Young White-Tailed .....	130
Bear, Polar .....	36	Devil-Fish .....	436
Beaver, Skull of .....	82	<i>Dicrostonyx hudsonius</i> , Skin of .....	85
Beavers, American, and their Work .....	81	<i>Dipodomys merriami</i> , Skin of .....	85
Bittern, American .....	262	Dogfish .....	424
Black Duck, Head of .....	269	Dolphin, Common .....	151
Blackbird, Red-Winged .....	200	Dove, Mourning .....	239
Bluefish .....	387		
Bluebird .....	183	Eagle, Bald .....	170
Blue-Jay .....	204	Eel, Electric .....	421
Boa Constrictor .....	340	Egret, Great White .....	262
Bobolink in Spring .....	178	Eider, American .....	277
Bob-White .....	242	Eider, King, Head of .....	279
Box-Fish .....	374	Eider, Spectacled, Head of .....	279
Buffalo, American Bison, or .....	101	Elk, American .....	121
Buffle-Head Duck .....	275	Elk, Winter Home of the .....	123
Bullhead, Common .....	416	<i>Evotomys gapperi</i> , Skin of .....	85

	PAGE		PAGE
Fer-de-Lance .....	353	Kinglet, Ruby-Crowned .....	184
Ferret, Black-Footed .....	29	Kite, Swallow-Tailed .....	232
Fins of a Typical Fish (Black Grunt).....	376	Lark, Meadow .....	200
Fisher.....	28	Lemming, Hudson Bay.....	86
Flamingo .....	266	Lemur, Ruffed .....	17
Flying-Fish, Common .....	409	Lizard, Blue-Tailed.....	334
Fox, Arctic.....	26	Loon.....	301
Fox, Black or Silver .....	25	Lung-Fish.....	380
Fox, Gray .....	27	Lynx, Bay.....	22
Fox, Red .....	25	Lynx, Canada .....	22
Frog, Leopard .....	362		
Gadwall, Head of .....	269	Mackerel, Spanish .....	388
Gar Pike .....	425	Maggie, American .....	203
Garter-Snake, Common.....	345	Mallard Duck.....	268
Gavial, Skull of .....	318	Manatee, Florida .....	155
Gila Monster .....	335	Man-o'-War Birds.....	290
Glass "Snake".....	336	Marten .....	28
Goat, Rocky Mountain .....	113	Martin, Purple.....	193
Golden-Eye Duck, Head of .....	269	Marmoset, Common.....	16
Goldfinch, American .....	196	Massasauga Snake .....	352
Goose, Canada .....	280	Master of the Trail, The.....	109
Gopher, Red Pocket .....	94	Menobranchnus, or Mud-Puppy .....	370
Gorilla .....	6	Menopoma, or Hellbender .....	368
Grackle, Purple .....	202	Merganser, American, Head of .....	279
Grouse, Canada .....	245	Merganser, Head of Hooded .....	279
Grouse, Eastern Ruffed .....	244	Merganser, Red-Breasted .....	278
Grouse, Pinnated .....	246	<i>Microdipodops megacephalus</i> , Skin of.....	85
Grouse, Sage .....	247	<i>Microtus pennsylvanicus</i> , Skin of.....	85
Grosbeak, Rose-Breasted .....	199	Mink.....	28
Gull, Herring .....	297	Moccasin, Water .....	352
Halibut, Common .....	418	Mocking-Bird .....	188
Hare, Polar .....	97	Mole, Digging Muscles of a.....	57
Hare, Prairie.....	97	Mole, Common.....	57
Hare, Varying .....	97	Mole, Fore Foot of Star-Nosed.....	58
Harlequin Duck, Head of.....	269	Mole, Nose of Star-Nosed.....	58
Haven of Refuge for Ducks .....	276	Mole, Star-Nosed.....	57
Hawk, Cooper's .....	231	Monkey, Black-Faced Spider .....	15
Hawk, Sharp-Shinned .....	230	Monkey, Diana .....	13
Hawk, Sparrow.....	227	Monkey, Japanese Red-Faced.....	13
Heron, Great Blue .....	260	Monkey, White-Throated Sapajou.....	14
Heron, Little Green .....	261	Moose in New Brunswick.....	<i>Frontispiece</i>
Hog-Nosed Snake .....	347	Mouse, Field .....	86
Horned Lizard; Horned "Toad" .....	336	Mouse, Jumping .....	93
Horns of Asiatic and American Mountain Sheep .....	114	Mouse, Le Conte's Harvest (lower figure).....	90
Humming-Bird, Ruby-Throated.....	209	Mouse, Mole .....	90
Ibis, White .....	263	Mouse, Rice-Field .....	89
Iguana, Common.....	333	Mouse, Red-Backed .....	87
Iguanas, Marine, on Narborough Island.....	332	Mouse, Typical Pocket .....	92
Jaguar .....	19	Mouse, White-Footed (upper figure) .....	90
Kingfisher, Belted .....	215	Mullet, Silver .....	390
King Snake .....	343	Murre, Common .....	301
		Muskallunge .....	394
		Musk-Ox, Wild Herd of .....	106
		Musk-Ox, Young Female .....	104

	PAGE		PAGE
Narwhal.....	152	Ptarmigan, Willow.....	249
<i>Neotoma floridana</i> , Skin of.....	85	Puffin, Common.....	304
Nighthawk.....	207	Puffin, Tufted.....	304
Nutcracker, Clarke's.....	205	Puffer Fish.....	374
Nuthatch, White-Breasted.....	185	Puma, or "Mountain Lion".....	20
Ocelot.....	21	Python, Reticulated.....	337
Old Squaw Duck, Head of.....	269	Rabbit, Cotton-Tail.....	97
<i>Onychomys leucogaster</i> , Skin of.....	85	"Rabbit," Jack.....	97
Opossum, Murine, and Young.....	166	Raccoon.....	41
Opossum, Virginia.....	165	Rail, Virginia.....	257
Orang-Utan, Female and Young.....	11	Rat, Cotton.....	89
Orang-Utans "Fight in the Tree-Tops".....	10	Rat, Florida Wood.....	88
Oriole and Nest.....	201	Rat, Kangaroo.....	91
<i>Oryzomys palustris</i> , Skin of.....	85	Rat, Kangaroo.....	92
Osprey, American.....	226	Rattlesnake, Banded, yellow phase.....	351
Otter.....	28	Rattlesnake, Banded, dark phase.....	351
Owl, Barn.....	220	Rattlesnake, Diamond.....	350
Owl, Barred.....	220	Rattlesnake, Prairie.....	351
Owl, Great Horned.....	223	Ray, Sting.....	436
Owl, Screech.....	221	Redhead Duck.....	274
Owl, Snowy.....	224	<i>Reithrodontomys leconti</i> , Skin of.....	85
Owl, Young Great-Horned.....	223	Ring-Necked Duck, Head of.....	269
Owl, Young Screech.....	222	Robin.....	181
Paddle-Fish.....	429	Ruddy Duck, Head of.....	279
Paddle-Fish, Under View of.....	429	Salmon, Quinmat.....	401
Pangolin, Rolled Up.....	161	Salmon, Sebago.....	404
Parrakeet, Carolina.....	216	Sand-Piper, Least.....	253
Partridge, California Mountain.....	242	Sawfish.....	435
Partridge, California Valley.....	243	Scaup Duck, Head of.....	269
Peccary, Collared.....	144	Scoter, Head of American.....	269
Pelican, California Brown.....	285	Scoter, Head of Surf.....	269
Pelicans, Florida Brown, on Pelican Island.....	284	Sea-Horse.....	423
Pelican, Great White.....	286	Seal, Harbor.....	44
Penguin, Emperor.....	306	Seal, Harp.....	51
Perch, Yellow.....	383	Seal, Head of Hooded.....	53
<i>Perognathus fasciatus</i> , Skin of.....	85	Seal, Ribbon.....	52
<i>Peromyscus leucopus</i> , Skin of.....	85	Seals, Fur, on "Hauling Grounds".....	49
<i>Perodipus richardsoni</i> , Skin of.....	85	Sea-Lions, California.....	44
Petrel, Stormy.....	294	Sea-Lion, Steller's.....	44
<i>Phenacomys orophilus</i> , Skin of.....	85	Sea-Lion, Steller's.....	46
Pickering, Chain.....	386	Shad, Common.....	407
Pigeon, Band-Tailed.....	238	Shark, Hammer-Head.....	432
Pike-Perch, Yellow.....	386	Shark, Mackerel.....	432
Pin-tail Duck.....	272	Shark-Ray.....	434
Pine Snake.....	344	Sheep, Black Mountain.....	111
Pipe-Fish, Great.....	423	Sheep, Head of White, front view.....	110
Platypus.....	167	Sheep, Head of White, side view.....	110
Plover, Kildeer.....	251	Sheep, White Mountain.....	111
Plumage of a Bird.....	180	Shoveller Duck.....	271
Porcupine, Canada.....	95	Shrew, Common.....	58
Porcupine Fish.....	374	Shrew, Short-Tailed.....	58
Prairie-"Dogs".....	76	Shrike, Loggerhead.....	191
Prairie-"Dog" Burrow.....	78	<i>Sigmodon hispidus</i> , Skin of.....	85

	PAGE		PAGE
Siren Salamander, or Mud-"Eel".....	371	Tortoise, Box.....	325
Skeleton of an American Bison.....	100	Tree-Duck, Head of Fulvus.....	269
Skeleton of a Bird of Prey.....	219	Tree-Frog, Northern.....	363
Skeleton of a Turtle.....	323	Trigger-Fish.....	374
Skeletons of Man and Gorilla.....	8	Trout, Brook.....	399
Skeleton of Pale Bat.....	60	Trout, Rainbow.....	398
Skunk, Common.....	31	Tuna.....	389
Skunk, Spotted.....	31	Turkey, Virginia Wild.....	250
Sloth, Two-Toed.....	160	"Turtle," Musk.....	326
Snake-Bird.....	287	"Turtle," Painted.....	327
Snapper, Red.....	391	"Turtle," Soft-Shelled.....	329
Snipe, Wilson's.....	253	"Turtle," Wood.....	328
Snow-Bunting.....	196		
Sparrow, White-Throated.....	197	Vireo, Red-Eyed.....	190
Spermophile, Say's.....	73	Vole, Northwestern.....	87
Spermophile, Thirteen-Lined.....	74	Vulture, California.....	233
Spermophile, Richardson's.....	75	Vulture, Young California.....	234
Spoonbill, Roseate.....	265		
Squirrel, Eastern Red.....	71	Wallabay, Rock.....	164
Squirrel, Flying.....	80	Walrus, Pacific.....	44
Squirrel, Gray.....	69	Walrus, Pacific.....	54
Squirrel, Southern Fox.....	70	Walrus, Young Atlantic.....	55
Steller's Duck, Head of.....	279	Warbler, Yellow.....	189
Stickleback, Two-Spined.....	415	Water-Snake, Red-Bellied.....	346
Sturgeon, Lake.....	427	Waxwing, Bohemian.....	193
Sucker, Common.....	412	Whales Attacked by Killers.....	150
Sunfish, Common.....	383	Whale, Bow-Head.....	147
Swallow, Barn.....	195	Widgeon, Head of American.....	269
Swallow, Cliff, and Nest.....	194	Wolf, Gray.....	22
Swan, Trumpeter.....	282	Wolverine.....	30
Swordfish.....	392	Woodcock, American.....	252
<i>Synaptomys cooperi</i> , Skin of.....	85	Woodcock on Nest.....	252
		Woodchuck.....	79
Tadpole to Frog, From.....	361	Wood-Duck.....	273
Tails of American Deer.....	129	Woodpecker, Downy.....	213
Tamandua Ant-Eater.....	158	Woodpecker, Golden-Winged.....	211
Tanager, Scarlet.....	195	Woodpecker, Red-Headed.....	212
Tarpon.....	406	Wood Thrush.....	182
Teal, Blue-Winged.....	270	Wren.....	187
Teal, Head of Green-Winged.....	269		
Tern, Common.....	297	<i>Zapus hudsonius</i> , Skin of.....	85
Terrapin, Alligator.....	328		

## MAPS AND CHARTS

	PAGE		PAGE
Map of North America.....	<i>Third page of cover</i>	Distribution of Mountain Sheep in North America.....	108
Landscape Chart of the Orders of Living Mammals.....	5	Distribution of the Prong-Horned Antelope.....	117
Map of Annual Migration of the Fur Seal Herd.....	48	Distribution of the Moose in North America.....	141
Chart of the Hare and Rabbit Family.....	97	Landscape Chart of the Orders of North American Birds.....	177
Range of the Musk Ox.....	105		

# INTRODUCTION

## THE GROUND-PLANS OF NATURE

**Science** is a collection of facts concerning natural objects or phenomena, arranged in good order, and made useful.

**Natural Science** is the study of Nature's works and forces, and embraces all things not made by man. Among its grand divisions may be mentioned natural history, chemistry, and physics.


**Natural History** is the study of Nature's common objects; but by most persons, this name is applied only to the study of animal life. Natural history treats of three great kingdoms—the animal, vegetable, and mineral.

The **Animal Kingdom** embraces not only all the living creatures which now inhabit the earth, but also those which have died, become extinct, and left only their buried remains, called fossils. Of the animal kingdom, three great groups of subjects may be recognized, as follows:

MAN, the study of whom is called . . . . . AN-THRO-POL'O-GY  
 THE LOWER ANIMALS, the study of which is called . . . . . ZO-OL'O-GY  
 EXTINCT, or FOSSIL ANIMALS, the study of which is called . . . . . PA-LE-ON-TOL'O-GY

In strict reality, Paleontology is only a branch of Zoology, for the two are inseparably dovetailed together. The living animals of to-day are the standards by which the paleontologist studies and determines those of the past.

This diagram illustrates the relations which the grand divisions of Natural History bear toward each other:



In its broadest sense, Natural History includes Chemistry and Physics; but as that term is now commonly used, it is intended to refer only to the life histories of living creatures.

**An Animal** is a living creature belonging to the animal kingdom; but this word is commonly, though incorrectly, used to designate mammals alone.

The animals of the world are so vast in number, and so varied in form, that these lessons will treat only of the higher forms of life, known as *Ver'te-brates*.

A **Vertebrate** is an animal having (usually) a bony skeleton, and a spinal column, or backbone, composed of a series of bones called *ver'te-brae*. This division of life is called a *Branch*.

The **Branch Ver-te-bra'ta** is divided into seven grand divisions, called *Classes*; which are known as *Mam'mals*, *Birds*, *Reptiles*, *Am-phib'ians*, *Fishes*, *My'zonts*, and *Lance'lets*.<sup>1</sup>

<sup>1</sup>Two other Classes, Enteropneusts and Tunicates, are, by some modern zoologists, regarded as Vertebrates. These low forms, however, lack a complete backbone, or notochord, and are therefore omitted.

A **Mam'mal** is a warm-blooded creature, that brings forth its young alive, and nourishes it with milk from its own body. All land mammals, save a few species, are covered with hair; and all sustain life by breathing air with the aid of lungs. Except man, the mammals which live upon land are also called *quad'ru-peds*.

A **Quad'ru-ped** is a mammal which possesses four feet, or, having two hands and two feet, like the apes, yet walks upon all-fours.

Man is a bi'ped, or two-footed animal. Land mammals generally are quad'rupeds, or four-footed, and monkeys are *quad-ru'ma-nous*, or four-handed.

The term **quadrumana** is often applied to apes and monkeys because the long great-toe on the hind foot makes the foot quite hand-like in its grasping power.

A **Bird** is a warm-blooded animal, which comes from an egg that usually is laid and hatched by the parent. It breathes air, is covered with feathers, usually is provided with wings, and all save a few species can fly.

A **Reptile** is a cold-blooded, egg-laying animal, usually covered with scales or a bony shell. All have lungs and breathe air, but some are able to live in water so comfortably they are called *am-phi'b'i-ous*.

An **Am-phi'b'i-an** is a member of the Class of animals which forms a connecting link between reptiles and fishes. Some breathe air, and live alternately on land and in water, like frogs. Others have gills, and live in water all their lives. A few are capable of developing *either gills or lungs*, according to the presence or absence of water, like the wonderful *Ax-o-lotl'* of Mexico.

A **Fish** is a cold-blooded animal, possessing gills, fins, and (usually) scales. All save a very few species live permanently in water. The exceptions are certain fishes in the East Indies which for short intervals hop about on land, or even climb rocks or trees!

**GRAND DIVISIONS OF THE ANIMAL WORLD**

		BRANCHES.			CLASSES.
<b>THE ANIMAL WORLD.</b>	}	<b>Ver'te-brates:</b> .....	}	Mammals	
				Birds	
				Reptiles	
				Amphibians	
				Fishes	
				Myzonts	
				Lancelets	
		<b>Insects:</b> —Body in segments, reproduce by a complete change in form.			
		<b>Crus-ta'ce-ans</b> (Crabs, Lobsters, etc.):—Skeleton external; gill-breathing, chiefly aquatic.			
		<b>Mol'lusks</b> ("Shell-Fish"):—Soft-bodied, usually covered by a hard, limy shell.			
<b>Worms:</b> —True worms, and other forms not fitting in elsewhere.					
<b>Star-Fishes:</b> —Salt-water animals, with star-like structure.					
<b>Corals:</b> —Minute, salt-water animals, which build up solid masses of their limy skeletons.					
<b>Jelly-Fishes:</b> —Disk-shaped, jelly-like sea animals, with no hard parts.					
<b>Sponges:</b> —Stationary aquatic animals, which look like plants; skeletons of tough, fibrous cells.					
<b>Pro-to-zo'ans:</b> —Lowest forms of life, beginning with the single cell; mostly microscopic.					

**HOW ANIMALS ARE CLASSIFIED**

In order to know and appreciate even a small proportion of the world's animals, their correct arrangement into groups is as necessary as a systematic arrangement of the books in a vast library. By their forms and characters, animals are divided into natural groups and subdivisions, and in

order that we may understand their proper relationships, and their places in Nature, we must learn and remember the general principles of animal **classification**. Without this foundation knowledge, a clear view of the splendid domain of animal life is impossible, and the life histories of our living creatures will be but a jumble of disconnected facts, of very slight practical use.

When properly simplified, the classification of the principal groups of our vertebrate animals is as easily learned and remembered as the leading facts of geography. Once learned, each animal observed thereafter can be located in the group to which it belongs, and its place in Nature understood. This helps toward exact knowledge of its anatomy and habits.

**No-men'cla-ture** is the naming of animals, and the groups to which they belong. The object of popular nomenclature, or naming, is to make the place and character of an animal clearly and correctly understood by the greatest possible number of people.

Scientific nomenclature relates to the use of technical names, in Latin or Greek, in which the general student is not often interested. Whenever through frequent or frivolous changes of scientific names, or by the giving of too great a number of them, our knowledge of animals becomes confused and uncertain, scientific classification defeats its own object, and becomes worse than useless. The observance by technical writers of the fatal rule of priority, by which the most obscure names often are exalted at the expense of more appropriate names in universal use, is rapidly debasing the legitimate value of Latin names generally, and creating wide-spread uncertainty and confusion.

Latin words are used for most scientific names, because Latin is the universal language of scientific men, the world over; and Latin names are used by all educated nations without change in form.

In the development of animal classification, the various classes of animals are subdivided into groups which gradually grow smaller, until at last each species is named and placed, thus:

Classes	are	divided	into	Orders:
Orders	"	"	"	Families:
Families	"	"	"	Genera (singular = genus):
Genera	"	"	"	Species (singular = species):
Species	"	"	"	Individuals.

As an example, take the Puma, or Mountain "Lion."

Its Order is FE'RAE, the wild beasts.  
 " Family is *Fel'li-dae*, the Cats.  
 " Genus is *Fel'lis*, the true Cats.  
 " Species is *concolor*, gray.  
 " Scientific name, therefore, is *Felis concolor*.

All these groups are divided into subdivisions, such as suborders, subfamilies, subgenera, and even subspecies; but in the writer's opinion there is very little excuse for their creation, or for their continued existence, and the student will do well to let them alone—until he feels the need for them.

A *tau'to-nym* is a scientific name in which the name of the genus is repeated as the name of the species. Thus, some authors write the Latin name of the American Bison as *Bison bison*; and the Anhinga is *Anhinga anhinga*. In America, the tautonym habit is merely another step toward the complete demoralization of zoological nomenclature.

A *tri-no'mial* is a name in three sections, applied to a subspecies; such as *Felis concolor oregonensis*.

By scientific authors, *species* are frequently divided into *subspecies*, or *races*, because in widely separated localities, animals of the same parent stock sometimes are so influenced by differences in climate, food, and surroundings that they assume different colors, or grow larger or smaller than the type. But, no matter how much individuals may differ in size and color, if it is possible to bring together a collection of specimens which will show *all stages of variation from the type* to

*the extremes*, then the specimens all belong to the same species. Thus, in passing from New York to Ohio, specimens of the Gray Squirrel show all shades of variation, from the typical gray to black; but all belong to the same species, called in Latin, *Sci-u'rus car-o-li-nen'sis*.

A **Species** is an assemblage of individual animals which in at least one respect are distinctly different from all others, and whose peculiarities are so well marked and so constant that they can be distinguished from all others without the aid of locality labels.

When a new kind of animal is found, adult specimens of which are distinctly different from those of all known species, an average specimen is taken as a *type*, and it is *described*, and christened by its describer. Every species should be distinguishable by external characters; and any animal which requires to be killed and dissected before it can be named, is of no practical value as an independent form.

To secure **recognition** among zoologists, it is important that the first description of a new species should appear in a regular publication of some scientific society, or in a scientific journal. In case the creature has not already been described, and the proposed species has just claims to stand alone, this name is entitled to stand, by right of **priority**, or first christening.

Many times it happens that through ignorance of what has been done by others, or by errors in judgment, a new name is bestowed upon an animal or plant that has already been named. Sometimes, also, it is found that the name bestowed has already been used for some other animal. A name applied to an animal or plant already named is called a *syn'o-nym*. In scientific books, synonyms sometimes are printed in a list under the correct name, followed by the names of their respective authors. A zoological synonym always stands for a published error, and scientific authors should be chary of describing as "new" any species which are likely to prove mere synonyms.

**The type** of any species is a carefully selected specimen which in size and color may fairly be considered the standard, or average, for that species. Among zoologists, this term is applied to the identical skin, or other specimen, described by its discoverer. Because of the many scientific names that are erroneously bestowed upon animals, the name of the author who is responsible for a name is usually printed, in abbreviated form, immediately after the name itself, thus:

*Popular name. Scientific name. Authority.*

**Coyote. Canis latrans. Say.**

A parenthesis enclosing a Latin name and the name of its author is a sign that the name has been changed somewhat from the form originally chosen and put forth by the author of the species.

Taken as a whole, this name means (1) that the "popular" name of the animal is Coy'ote; (2) that its scientific name (Latin) is *Canis* (=dog) *la'trans* (=barking); and (3) that it was first correctly described and named in print by a man named Say. If we consult our books, we will find that Thomas Say was a Philadelphia naturalist, and his description of this animal appeared in "Long's Expedition to the Rocky Mountains," published in 1823, Vol. I, page 168.

Whenever the name of an animal has been so long in use that it has become familiar to millions of people, any attempt to change it tends to create confusion. A slightly incorrect name in universal use is often better than the confusion and doubt inseparable from attempting a change. Thus, the American buffalo, considered in connection with the world's bovine animals generally, is really a bison; and the prairie-"dog" is really a prairie marmot; but since nearly all the inhabitants of America know these animals by their incorrect names, and any effort to force a universal change would be quite fruitless, it would be unwise to attempt it.

It is very important to the student that the names of the various Orders of vertebrate animals should be learned and remembered; for they are the keys with which to unlock and reveal all systematic knowledge of mammals, birds, reptiles, amphibians, and fishes.

### THE INTELLIGENCE OF ANIMALS: A WARNING

During the past two years, so many persons have requested my views regarding the mental capacity of animals, that I feel impelled to enter here a brief statement, coupled with a warning. Unfortunately, it cannot be written otherwise than in the first person.



While I have no desire to exploit my personal experiences among wild creatures, it is at least fair to state, for the benefit of the millions to whom the writer is unknown, that of wild creatures in their haunts, and also in captivity, he has seen as much as most men of his tastes.

The tendency of the present is to idealize the higher animals, to ascribe to them intelligence and reasoning powers which they do not possess, and in some instances to "observe" wonderful manifestations that take place chiefly in the imagination of the beholder. For example, to a ruffed grouse, having mingled blood and mud on a broken leg, is ascribed a deliberate and well-considered attempt at "surgery," and the intentional making of a clay jacket, re-enforced with pieces of grass. To my mind, all such "observations" as the above are too absurd for serious consideration; and when put forth for the information of the young, they are harmful.

There exists to-day a tendency to ascribe to wild animals a full measure of human intelligence. But wild creatures must not be taken too seriously. With all their "schools" in the woods, they are not yet as intelligent as human beings; and the strain that is being put upon them by some of their exponents is much too great. With the most honest intentions, a naturalist may so completely overestimate and misinterpret the actions of animals as to reach very ridiculous conclusions.

Judging from all that I have seen and heard of wild creatures of many kinds, from apes to centipedes, both in captivity and out, I believe that practically all their actions are based upon natural, inborn instinct—nearly all of it in the line of self-preservation, and the exceptions are due to the natural tendency to imitate leaders. Of hereditary knowledge—another name for instinct, some animals have an abundance. Of special knowledge, acquired by systematic reasoning from premise to conclusion, most animals have very little, and very few ever exhibit powers of ratiocination.

It is *not* true that young animals know things only as their parents teach them. The assertion that all young birds must be "taught" to fly, or run, or swim, or catch insects, is ridiculous, and not even worthy of discussion. It is just as natural for a one-week-old lion cub to spit, and claw at a human hand, as it is for it to breathe and suck. There are no deer in a captive herd so insanely wild and fearful of keepers as the fawns.

No; even the higher animals are not yet as wise as human beings. In matters involving intelligence, such as in the treatment of wounds, or disease, below the higher Primates there is not more than one out of every hundred which has sense enough to comprehend a relief measure, or which will not fight the surgeon to the utmost. Some apes do indeed learn to be doctored; but there are many which never grasp the idea, and fight until they die. Of mammals generally, not more than one out of every hundred will permit a bandage to remain on a broken leg when they have the power to tear it off. "Animal surgery," indeed!

In the matter of disposition, wild mammals and birds are no more angelic than human beings. In every family, in every herd, and in every cage, from tigers to doves, the strong bully and oppress the weak, and drive them to the wall. Of all quadrupeds, deer are the greatest fools, wolves are the meanest, apes the most cunning, bears the most consistent and open-minded, and elephants the most intellectual.

Of birds, the parrots and cockatoos are the most philosophic, the cranes are the most domineering, the darters are the most treacherous, the gallinaceous birds have the least common-sense, and the swimming birds are by far the quickest to recognize protection, and accept it.

The virtues of the higher animals have been extolled unduly, and their intelligence has been magnified about ten diameters. The meannesses and cruelties of wild animals toward each other form a long series of chapters which have not yet been written, and which no lover of animals cares to write.

I can see no possible objection to the writing of good fiction stories in which animals are the characters and the actors throughout. I love a good story, and I enjoy a wild-animal hero, even when the entire plot and all its characters are imaginary. To such there can be no objection, *so long as the reader knows that fiction is fiction!* But the realms of fact and fiction are very distinct, and the boundary should be maintained, openly and visibly. In books for children, espe-

cially, fantastic imaginings should not be offered as serious facts; but such stories as "Raggylogg," "Redruff," and "Krag," by Mr. Ernest T. Seton, deserve to live forever. "Mooswa" is a fiction story of animals that is one of the best of its kind.

The most marvellous doings of wild animals are to be found in books and newspapers. Only in books do porcupines roll down steep hills in order to gather dead leaves upon their quills, and thereby be able to do more wonderful things. Only in books do kingfishers catch fish, carry them a mile or less, and place them in a brook in order to give their nestlings object lessons in ichthyology, and in the gentle art of angling. You or I may spend years in the forests and fields, observing and collecting wild creatures, and see only a very few acts of the wild folk which we can call wonderful. But then, somehow, our animals rarely have been as large, or as well educated, as those of some other observers.

Try all questions of animal action and intelligence with the touchstone of common-sense. Be not startled by the "discovery" that apes and monkeys have "language"; for their vocabulary is not half so varied and extensive as that of barn-yard fowls, whose language many of us know very well. Take no stock in the systematic and prolonged "duels" of wild animals who meet and fight to the death, under Marquis of Queensberry rules. A fight between two wild animals is usually a very brief event,—so say reliable men who have seen them in the wilds,—and unless there is an accidental death-lock of antlers, the vanquished party usually shows his heels long before he is seriously wounded.

Animal psychology is a most interesting study, and its pursuit is now engaging the serious attention of scientific men. If the general public could know the plain and simple basis on which they are proceeding, this warning against the idealization of animals would hardly be necessary. Men of science who study the minds of animals do not idealize their subjects, or ascribe to them super-human intelligence; nor are they always on the alert to ascribe to every simple action some astoundingly intelligent and far-fetched motive. In the study of animal intelligence, the legitimate Truth is sufficiently wonderful to satisfy all save those who crave the sensational, regardless of facts.

### RULES FOR MEASURING MAMMALS, HORNS, ETC.

The increasing amount of attention that is being paid to the measurements and weights of animals renders necessary the adoption of a uniform system, in order that species and individuals may be compared on a fair basis. To promote this end the following rules are offered:

#### SMALL MAMMALS GENERALLY

1. Record all measurements in feet and inches, and leave the metric scale for those who prefer a foreign system.
2. Measurements of skins are of very slight value; therefore, always measure a specimen before skinning it.
3. Lay every mammal on its side, pull the head straight forward, and measure from the tip of the nose to the point where the tail joins the body. This is the "*Length of head and body.*"
4. From the last-mentioned point, measure to the *end of the tail vertebrae, not the hair*, for "*Length of tail.*" If the tail-tuft is important, measure it separately.
5. Weigh large examples of species that are larger than rats and mice; and in each case, *weigh the whole of the specimen.*

#### LARGE MAMMALS

1. The "*Height at the shoulder*" is the most important measurement. To obtain this, hold the uppermost foreleg *as nearly as possible in the position it occupied when supporting the animal.* Do *not* measure from the "point of the hoof"; for that means nothing. Hold the hoof with its bottom parallel with the body, as when the animal stood upon it; erect there a stick to mark the

bottom line, and another to mark the top of shoulders, at the skin. The distance between the two perpendiculars, in a straight line, will be the true height of the animal. Do not follow any curves.

2. The "*Length of head and body*" must be obtained in a *straight line* between root of tail and end of nose, *with the head drawn straight forward*, and not following any curves. The "*Length of tail*" is from its base to the end of the vertebrae.

3. The "*Girth*" is the tight circumference of the animal immediately behind the forelegs.

4. The "*Depth of the body*" is the distance in a straight line from the top of the shoulders to the brisket, or lower line of the breast, immediately behind the foreleg. To artists, sculptors, and taxidermists, this is a very important measurement.

5. The "*Circumference of the neck*" is taken half way between the ears and shoulders, close to the skin.

6. The "*Length from head of femur to head of humerus*" is also a highly valuable figure for artists, and it is easily taken by feeling through the skin for the high points of those joints.

7. Weigh an animal before it is "dressed"; but if the dressed weight of a deer is known, a close approximation to its live weight can be obtained by the aid of the rule given on page 124.

#### ANTLERS AND HORNS

1. The "*Length on outer curve*" is obtained by starting the tape line at the base of the horn, at its lowest point on the face, and following the curves or windings of the horn, quite to the tip. In horns that are deeply ringed, such as those of the large African antelopes, the tape must not be pressed into the hollows between the ridges.

2. The "*Greatest spread*" is taken from outside to outside of the antlers *where they spread widest!* This should *not* be taken inside the horns, for that does not represent the real width of the horns, any more than interior measurements would represent the spread of a tree.

3. The "*Distance between tips*" needs no explanation.

4. The "*Circumference at base*" should, for all bovines, sheep, goats, ibex, and deer, be taken in a circle around the largest diameter of the horns. The tape should *not* follow the meanderings of the end of a sheep's horn. With the antlers of all members of the Deer Family, the circumference should be measured immediately above the burr.

5. "*Width of palmation*" of moose and caribou should always be measured where the palmation is widest.

6. A "*Point*" on an antler is any pointed projection of sufficient length that a watch can hang upon it without falling off.

7. The "*Weight of horns*" must state whether it be with "entire skull," or "with skull-piece" only.

8. Shed antlers that have been set artificially on a manufactured skull, or frame, are not entitled to measurement for "spread"; but where a skull has been sawn in two lengthwise by a clean cut, and bolted together again *without alteration of the sawn surfaces*, it is entitled to measurement for "spread" and "distance between tips."



BOOK I  
MAMMALS



## CHAPTER I

### THE ORDERS OF MAMMALS

The living mammals of the world, as distinguished from those which are extinct, or fossil, may be divided into thirteen grand divisions, called **Orders**. The order is the foundation of mammalian arrangement. Without adequate knowledge of these divisions, a clear understanding of the relationships of mammals is quite impossible.

It is customary with technical writers to begin with the lowest forms of life, and toil upward toward the highest; but it is very discouraging to the young student to find the most interesting forms the farthest away. Frequently the most interesting animals are never reached! For many reasons, it is best that the general student should study first the forms that are most important, and also most interesting, and thus make sure of them. We therefore begin our studies of the animal kingdom with the highest forms, and adopt the latest names that have come into use amongst zoologists.

While the great majority of the examples cited will be North American, a few from other continents will be introduced to complete the chain of important facts.

#### THE ORDERS OF LIVING MAMMALS.

ORDER.	PRONUNCIATION.	MEANING.	EXAMPLES.
PRIMATES . . . . .	<i>Pri'matz</i> . . . . .	First order . . . . .	Man ; apes and monkeys.
FERAE, or Carnivora . . . . .	<i>Fe're</i> . . . . .	Flesh-eating Wild Beasts . . . . .	Cats, dogs, bears, weasels.
PINNIPEDIA . . . . .	<i>Pin-ni-pe'dia</i> . . . . .	Fin-footed . . . . .	Sea-lions, seals, walrus.
INSECTIVORA . . . . .	<i>In-sec-tiv'o-rah</i> . . . . .	Insect-eaters . . . . .	Moles and shrews.
CHIROPTERA . . . . .	<i>Ki-rop'ter-ah</i> . . . . .	Wing-handed . . . . .	Bats and flying-"foxes."
GLIRES, or Rodentia . . . . .	<i>Gli'rez</i> . . . . .	Gnawers . . . . .	Hares, gophers, rats, squirrels.
UNGULATA . . . . .	<i>Un-gu-la'tah</i> . . . . .	Hoofed . . . . .	Cattle, deer, sheep, swine, tapirs.
CETE . . . . .	<i>Se'te</i> . . . . .	Whales . . . . .	Whales, porpoises, dolphins.
SIRENIA . . . . .	<i>Si-re'ne-a</i> . . . . .	Sea-cows . . . . .	Manatee and dugong.
EDENTATA . . . . .	<i>E-den-ta'ta</i> . . . . .	Toothless . . . . .	Armadillos, sloths and ant-eaters.
EFFODIENTIA . . . . .	<i>Ef-fo-de-en'shia</i> . . . . .	Diggers . . . . .	Pangolin and aardvark.
MARSUPIALIA . . . . .	<i>Mar-su-pi-a'li-a</i> . . . . .	Pouched . . . . .	Opossum, kangaroo.
MONOTREMATA . . . . .	<i>Mon-o-trem'a-ta</i> . . . . .	Single duct . . . . .	Platypus and echidna.

## EXPLANATION OF THE CHART OF THE ORDERS OF MAMMALS.

To the beginner in Natural History studies, the Order is the master-key to classification.

This Chart is based on the well-known fact that in the pursuit of a difficult study, any scheme which properly and truthfully appeals to the eye is an aid both to the understanding and the memory. It shows the relative importance of the various Orders of Mammals, but *not* their relative sizes, based on the number of species in each, as has been done later on with the birds. If number of species were given precedence over economic importance, the Order Glires would dominate, and the Order Ungulata would appear small and insignificant.

It is impossible to construct a diagram which will show correctly the relations which the various Orders bear toward each other, anatomically. This is because some Orders are characterized by their teeth, some by their feet, or hands; others by their wings, and two by their mode of producing their young.

It will be noted that:

The Primates, of the tree-tops, have the highest position.

The Cete, which in some respects are the lowest of the Mammalia, occupy the lowest position.

The Bats are shown in mid-air, and the Insectivores appear under ground, where they live out their lives.

The Seals and Sea-Lions appear both on the shore and in the sea, and the Sirenians are located in an estuary.

The Ferae, Glires and Ungulata spread throughout the whole visible earth, covering forest and plain, sea, pond and stream, from the sea to the most distant mountains.

The Monotremates, or egg-laying mammals, are quite apart from all other land mammals, and appear low down, near the home of the ducks, as shown on the bird chart. The space allotted to this strange Order has been made egg-shaped, to suggest the leading characteristic of its members.



## CHAPTER II

# THE ORDER OF APES AND MONKEYS

### PRIMATES

This Order includes all creatures with hands, and hand-like feet. With the exception of the Japanese red-faced monkey, the tscheli monkey of China, and two or three other Chinese species, all its members inhabit the tropics, far below the frost line. It is on or near the Equator that the lower Primates reach their highest development, and the great apes approach nearest to man. Let it not be supposed, however, that the chain of evolution from the aye-aye to the gorilla is complete; for the gap between the gibbons and the monkeys is much greater than that between the gorilla and man.

All men, even savages, are specially interested in apes and monkeys, because they are the highest of the lower animals, and stand nearest to man. There is no human being of sound mind to whom their human-likeness does not appeal. For this reason, we will introduce here several species which are not found in the New World, for the reason that without them our Foundation for the Mammalia would be incomplete.

Although tropical America contains a very respectable number of species of monkeys, they

are, as a whole, both structurally and mentally, far lower than the monkeys and baboons of the Old World. Structurally they are weak, in spirit they are timid and cowardly, and intellectually they are dull to the point of stupidity. With the exception of the sapajous, they are in general so ill fitted to survive that if they are on exhibition it is a difficult matter to keep any of them alive in captivity much longer than one year. If not exhibited, they survive longer.

On the other hand, very many of the monkeys and baboons of the Old World have developed first-class fighting powers, and pugnacious tempers. They have dangerous canine teeth, wide-spreading jaws, strong muscles, and keen wits for either attack or defence. The Lemuroids, however, the lowest of the Primates, are as mild-mannered and harmless as rabbits.

With Ethnology, the study of the races of Mankind, we have here nothing to do. That subject is so interesting, and so vast in its extent, that nothing less than an entire volume can adequately set it forth. The grand divisions of the Primates in general are as shown below.

		FAMILIES.		EXAMPLES.	
ORDER PRIMATES.	SUB-ORDER ANTHROPOIDEA:	MAN,	<i>HOM-IN'I-DAE.</i>		
		ANTHROPOID APES,	} <i>SIM-PI-DAE.</i> . .	<b>Gorilla,</b> <i>Gorilla gorilla.</i> <b>Chimpanzee,</b> <i>Pan troglodytes.</i> <b>Orang-Utan,</b> <i>Simia satyrus.</i> <b>Gibbon,</b> <i>Hylobates leuciscus.</i>	
		OLD-WORLD MONKEYS AND BABOONS,	} <i>CER-CO-PI-THE'CI-DAE.</i> . . .	<b>Japanese Red-Faced Monkey,</b> <i>Macacus speciosus.</i> <b>Diana Monkey,</b> <i>Cercopithecus diana.</i> <b>Gelada Baboon,</b> <i>Theropithecus gelada.</i>	
		NEW-WORLD MONKEYS,	} <i>CEB'I-DAE.</i> . . .	<b>White-Throated Sapajou,</b> <i>Cebus hypoleucus.</i> <b>Black Spider-Monkey,</b> <i>Ateles ater.</i> <b>Howlers,</b> <i>Alouatta.</i>	
		MARMOSETS,	} <i>CAL-LI-THR'CI-DAE.</i>	<b>Common Marmoset,</b> <i>Callithrix jacchus.</i>	
		SUB-ORDER LEMUROIDEA:	LEMURS,	<i>LE-MURI-DAE.</i> . .	<b>Ruffed Lemur,</b> <i>Lemur varius.</i>
			TARSIER,	<i>TAR-SII-DAE.</i> . .	<b>Tarsier,</b> <i>Tarsius tarsius.</i>
			AYE-AYE,	} <i>DAU-BEN-TON-I-DAE.</i>	<b>Aye-Aye,</b> <i>Daubentonia.</i>

**The Apes.**—The three great man-like (or an'thro-poid) apes—gorilla, chimpanzee and orang-utan—are so much like human beings that, to most persons, they are the most won-

derful of all living creatures below man. Their points of resemblance to man are so many and so striking that they are a source of wonder even to savages.

As will be observed from a comparison of the skeletons of man and gorilla, below the skull their parallelism is remarkably close. Both in kind and in number the bones are the same, and they differ only in their proportions. The hands and feet of the gorilla are designed for a life that is half terrestrial and half arboreal, while those of man are for life on the ground. The long thumb and great toe of the gorilla are far superior to those members in the chimpanzee and orang-utan.

The widest differences between man and the gorilla are in their skulls. In the gorilla, the high forehead and intellectual faculties so characteristic in man are totally wanting, indicating a very low order of intelligence. The long and powerful canine teeth are alone sufficient to proclaim the savage wild beast.

To many persons it seems strange that notwithstanding the seemingly wide differences between the various races of men, all mankind be referable to a single species. In spite of the vast differences in intellect between the native Australian—not yet out of the stone age—and a Caucasian philosopher, both belong to *Homo sapiens*, and between them there is not even a sub-specific difference.

Even if the great apes could talk as well as the Veddahs of Ceylon, whose vocabulary consists of about two hundred words, their anatomical differences from the genus *Homo* would separate them quite as widely as they now are. To segregate a species requires a *structural difference that is constant*.

**The Gorilla**<sup>1</sup> is the largest, the ugliest, the most fierce in temper, and by reason of its shorter arms and longer legs, it is really the nearest to man. It is the only ape that walks erect without being taught, and that spends a considerable portion of its life upon the ground. In bulk it is larger than an average man, and its

<sup>1</sup> *Go-ri'la gorilla*.



By permission of J. F. G. UMLAUFF.

SKELETONS OF MAN AND GORILLA.

- |                        |                       |
|------------------------|-----------------------|
| 1, cervical vertebrae. | 12, carpals.          |
| 2, collar bone,        | 13, metacarpals,      |
| 3, humerus.            | 14, phalanges,        |
| 4, sternum,            | 15, cavity of pelvis, |
| 5, ribs,               | 16, sacrum,           |
| 6, rib cartilages,     | 17, femur,            |
| 7, dorsal vertebrae,   | 18, patella,          |
| 8, lumbar vertebrae,   | 19, fibula,           |
| 9, pelvis,             | 20, tibia,            |
| 10, radius,            | 21, tarsals,          |
| 11, ulna,              | 22, metatarsals,      |
|                        | 23, phalanges.        |

arms and chest are of enormous proportions.<sup>1</sup> The countenance of the Gorilla is very ugly and repulsive, and the shape of its skull is much farther from that of man than are those of the chimpanzee and orang-utan. Its skin is black, and the hair of full-grown specimens is grizzly gray.

The Gorilla inhabits only a very small area in West Africa, directly on the equator, between the Gaboon and Congo Rivers, and extending only two hundred miles back from the coast. It is very shy, and so difficult to approach in those dark and tangled forests that very few white men ever have seen one wild.

One of the most remarkable specimens ever secured was the huge old male killed and photographed by Mr. H. Paschen, a German trader, near Tsonu Town, German Cameroon country, two hundred and forty miles north of the equator, in 1901. This animal, photographed in the flesh, with three natives beside it for comparison, to show its immense size, was shot in a tree, without difficulty or danger. It measured 66 inches in height, its chest, arms and shoulders were of gigantic proportions, and its weight was estimated at 500 pounds. Twelve men were required to carry it from the jungle to the village, where it was photographed.

On account of the sullen, sulky disposition of the Gorilla in captivity, only one of the four or five young specimens that have been brought to Europe has lived longer than about eighteen months. They sulk, often refuse food, will not exercise, and die of indigestion. Up to this date (1903) only one live Gorilla, and that a tiny infant, has ever landed in the United States; and it lived only five days after arrival. Showmen sometimes label a baboon "Gorilla," or "Lion-Slayer," and it is well to remember that the Gorilla has no tail whatever.

**The Chimpanzee**<sup>2</sup> is about one-third smaller than the gorilla. Its brain, face, ears and hands are more man-like than those of any other ape, and its large brain and keen mind render it in thought and habit much more man-like than the

gorilla. It is an animal of bright and cheerful disposition, though subject to sudden fits of bad temper, and having a good memory, it is easily taught. Young Chimpanzees are affectionate and child-like, but when large and strong, the males are usually dangerous, and not to be trusted. Some individuals have displayed remarkable intelligence. "Sally," of the London Zoological Gardens, could count correctly up to five, whenever bidden, and hand out the correct number of straws.

After several years of observation of living Chimpanzees and orang-utans, in daily comparison, I am convinced that the only substantial psychological differences between the two species are (1) that the temperament of the Chimpanzee



N. Y. Zoological Park.

YOUNG FEMALE CHIMPANZEE.

is of the nervous type, and its mind is more alert and prompt in action than that of the orang, while on the other hand (2) the temperament of the orang is sanguine, its disposition is more serene, and while its mind may be somewhat less showy on exhibition, its capacity is quite equal to that of the Chimpanzee. The greater quickness of the Chimpanzee, both in thought and action, renders it on the whole the best show animal in public performances.

Many persons consider the Chimpanzee superior in intelligence to the orang-utan, but thus far the only real difference appears to be that the

<sup>1</sup> The average man of the Anglo-Saxon race is 5 feet 6 inches in height and weighs 160 pounds.

<sup>2</sup> *Pan troglodytes*. Described in most books under the untenable and more unwieldy name of *Anthropopithecus troglodytes*. This animal has been described under nine different generic names, but *Pan* is the oldest one available and the best.



By permission of EDWARDS BROS.

A DRESSED-UP CHIMPANZEE.

mind of the former is more alert, and acts more quickly than that of the orang.

In walking, the Chimpanzee does not place the palms of its hands flat upon the ground, but bends its fingers at the middle joint, and walks upon its knuckles.

It does not, as so often asserted on hearsay evidence, build a hut or a roof of branches under which to sleep. Its home is the heavy forest region of equatorial Africa, from the Atlantic ocean to Lake Tanganyika. Like the gorilla, its skin is black, and when young its hair also, but when fully grown its hair is dark iron-gray. This animal can at one glance be distinguished from the orang-utan by the greater size of its ears, and its black color.

**The Orang-Utan** (from two pure Malay words, "orang" = man, and "utan" = jungle) is also about two-thirds the size of the gorilla, and is easily recognized by its brick-red hair, brown skin and small ears. The largest specimen on record stood 4 feet 6 inches in height from heel to head, measured 42 inches around the chest, and between finger tips stretched 8 feet. The old males develop a strange, flat expansion of the cheek, called "cheek callosities,"

13 inches across; but in young animals this is seldom developed. The hand is  $11\frac{1}{2}$  inches long, the foot  $13\frac{1}{2}$  inches, but the width of each across the palm is only  $3\frac{3}{4}$  inches. The weight of a large, full-grown male Orang is about 250 pounds.

The black gorilla and chimpanzee both inhabit the land of black men; the brown Orang-Utan lives only in Borneo and Sumatra, the land of the brown-skinned Malay. The latter prefers the belt of level, swampy forest near the coast, lives wholly in the tree-tops, and rarely descends to the earth except for water. Orangs travel by swinging underneath the large branches with their long, muscular arms. Because of their great weight, they cannot leap from tree to tree, as monkeys do, but they swing with wonderful rapidity and precision. They eat all kinds of wild fruit, fleshy leaves, and the shoots of the screw pine.


In proper hands, young Orang-Utans are very susceptible to training. In 1901 the New York



Drawn by C. B. HUDSON.


A FIGHT IN THE TREE-TOPS.

Old male Orang-Utans, with cheek callosities.



Copyright, 1903, by W. T. HORNADAY.

LANDSCAPE CHART OF THE ORDERS OF LIVING MAMMALS.



GORILLA.

By permission of J. F. G. UMLAUFF.

Shot and photographed at Tsonu Town, West Africa, by H. PASCHEN, 1901



FEMALE ORANG-UTAN AND YOUNG.

Drawn from specimens living in the New York Zoological Park, by A. G. DORING.

Zoological Park contained four Orangs, all of which were easily taught to wear clothes, sit in chairs at table, eat with fork and spoon, drink from cups and bottles, and perform many other human-like actions without nervousness, in the presence of two thousand visitors. Each of the Orangs learned its part in about two weeks' training, and at the dinner-table acted with gravity and decorum. "Rajah," the senior member of the quartette, never once suffered from stage fright, or lost his nerve during a public performance.

In captivity, young Orang-Utans are as affectionate as human children, and very fond of their human friends. In the jungles of Borneo the full-grown males often fight savagely by biting each other's faces, and by biting off fingers and toes. At night the Orang makes a nest to sleep upon, by breaking off leafy branches, and laying them cross-wise in the forked top of a sapling. On this huge nest-like bed it lies flat upon its back, grasps a branch firmly in each hand and foot, and is rocked to sleep by the cradle-like swaying of the tree-top.

Unless attacked at close quarters, in their forest homes, none of the great apes is dangerous to man. All of them flee quickly from the dreaded presence of Man, the Destroyer. They never fight with clubs, but when attacked at close quarters they bite, just as do human roughs. When enraged, the gorilla *does* beat its breast with its fists, just as Du Chaillu said; and it does this even in captivity.

"**The Missing Link.**"—For thirty years at least, Science has been seeking in the earth for fossil remains of some creature literally standing between man and the great apes, but at present unknown. In 1879, Mr. A. H. Everett made for the Zoological Society of London a thorough examination of the deposits on the floors of some of the caverns of Borneo. To-day, some naturalists are straying toward the lemurs in search of the parent stem of man's ancestral tree. Vain quest! The gap between Man and Lemur is too great to be bridged in this world. A coincidence between skull bones is a long way from man-likeness.

Place upon the shoulders of a gorilla the head of a chimpanzee, and we would have—what? The Missing Link, no less,—a hairy, speechless man! The man-apes we have. Let those who

seek the undiscovered ape-man search the Tertiary deposits of the fertile uplands that lie between the gloomy equatorial forests of the black apes and the Bushmen of South Africa; for there, if anywhere, will the Missing Link be found.

**The Gibbons.**—From the three huge, coarsely-formed and unwieldy man-like apes described above, the line of descent drops abruptly and far. Their nearest relatives are the Gibbons—creatures of small size, marked delicacy of form, no weight or strength to speak of, but of marvellous agility in the tree-tops. Their heads are small and round, their teeth are weak, and their faces are like those of very tiny old men.

Their arms and hands are of great length in proportion to their body size, yet so very slender are their muscles that a live Gibbon seems like a hairy skin drawn over a skeleton. The largest specimen I measured in Borneo had the following remarkable dimensions: head and body, 19 inches; extent of outstretched arms and hands, 5 feet 1 inch; entire reach of arms and legs, 5 feet 1 inch; hand, 6½ inches long by 1 inch wide; weight, 10¼ pounds.

Of Gibbons there are about six species, and they inhabit Borneo, Sumatra, the Malay Peninsula, Burmah and Siam. With the **Gray Gibbon**,<sup>1</sup> of Borneo, I am well acquainted; and after the three great man-like apes, it is to me the most wonderful of anthropoids. They are very timid, the shyest of all Primates that I ever hunted, and wonderfully successful in eluding the hunter. Nevertheless, so strong is their affection for their young, I have seen a whole troop that had made good its escape, return at the call of an infant Gibbon in trouble, and all reckless of their own safety come down within twenty feet of their deadly enemy. Very few other mammals will do this.

The most wonderful habit of the Gibbon is its flight down hill when pursued. Of course it never dreams of descending to the earth, but in the half-open hill forests of Borneo I have seen these creatures go downward through the tree-tops, in a straight course, leaping incredible distances, catching with their hands, swinging under, catching with their feet, turning again, and so on by a series of revolutions, almost as fast as the flight of a bird.

<sup>1</sup> *Hyl-lo-bates leu-cis'cus*.



The **Slamang**,<sup>1</sup> of Sumatra, is the largest and rarest of the Gibbons. It is jet black, all over, face as well as fur, and it has a throat pouch which is distended to astounding proportions when it utters its peculiar, piercing cry. This species is as rare in captivity as the gorilla, and the only specimen seen alive in the New World up to 1903 was exhibited at the New York Zoological Park in that year.

**OLD-WORLD MONKEYS AND BABOONS.**

*Cercopithecidae.*

**Typical Old-World Monkeys.**—Asia, Africa and the islands of the Malay Archipelago contain a great number of species of monkeys. The most northern is the sturdy **Japanese Red-Faced Monkey**, with no tail to speak of. It is

catching cold. Their tempers are quite as warm as their blood.

From Japan, monkey-land extends southward through China, and southern Asia generally, the



DIANA MONKEY.

Malay islands almost to Australia, and throughout the whole of Africa except its great deserts, to the extreme south.

Of all these Old-World species, none have prehensile (grasping) tails, like many American species. Many of them are beautifully colored, however, and the markings of some are quite fantastic. The **Diana Monkey**, of West Africa, is elaborately marked with black, white, gray and brown, and it is one of the most beautiful of all monkeys. An old-world monkey can nearly always be recognized by the very narrow space between the nostrils.

**Short-Tailed Monkeys.**—It must not be supposed that because the tail of a monkey is so short as to be scarcely visible, the wearer is therefore a true ape. There are several baboon-like animals with tails exceedingly short and insignificant, but which are far removed from the true apes. Some of these are called apes, but they are all much lower in the scale. Of these, the most important are:

The Black "Ape" of Celebes;

The Barbary "Ape" of Gibraltar and North Africa;

The Pig-Tailed Ma-caque' (pronounced Ma-cak') of the East Indies, east of Ceylon, and

The Japanese Red-Faced Monkey.

**The Baboons.**—In nearly every portion of Africa abounding in rocky hills covered with scanty vegetation may be found Baboons,—fierce of aspect, domineering in temper, strong of limb, and sometimes very ugly in countenance.



SANBORN, Photo., N. Y. Zoological Park.

JAPANESE RED-FACED MONKEY.

Note the narrow space between the nostrils.

clothed with long, shaggy hair, and those in the New York Zoological Park live outdoors all winter, and gallop about in the snow without

<sup>1</sup> *Sym-pha-lan'gus syn-dac'tu-lus.*



N. Y. Zoological Park.

**DEAD GELADA BABOON.**

Note the lion-like aspect.

Their noses are long and dog-like. They live on the ground, travel in troops of ten to twenty individuals, and rob grain-fields with great boldness. It is asserted by African explorers that even hungry lions prefer to let them alone. The canine teeth of an adult Baboon are so long and sharp that they are dangerous weapons. Without exception, Baboons are the most fierce-tempered animals of all the Primates, not even excepting the great apes, which never fight when they can run away.

All told there are about sixteen species of Baboons, all of which are found in Africa outside of the dark forests of the equatorial regions. The great **Gelada Baboon**,<sup>1</sup> of Abyssinia, is one of the most remarkable of all animals. It is like a small lion, with a Baboon's feet and hands; but its wonderful grimaces are peculiar to itself.

A Baboon of average size stands 24 inches in height at the shoulders, and weighs about 45 pounds. The majority of the species are of a yellowish color, mixed with brown. The **Mandrill** is known everywhere by its brilliant blue and scarlet muzzle, and yellow chin beard.

<sup>1</sup> *The-ro-pith'e-cus ge-la'da.*

**NEW-WORLD MONKEYS.**

All the monkeys of the New World are marked by the wide space between the nostril openings, and nearly all the larger species possess prehensile, or grasping, tails that are as useful as a fifth arm and hand. Most of the species which do not have prehensile tails are quite small. Of the clinging-tailed monkeys there are three important groups, which are represented in North America. They are the Sapajous, the Spider Monkeys and Howlers.

The American monkey most frequently seen in captivity is the **White-Throated Sapajou**<sup>1</sup>

**WHITE-THROATED SAPAJOU.**

Note the wide space between the nostrils.

(sap'a-jew) or **Cap'u-chn**, called by animal dealers and showmen, the "Ring-Tail." This monkey is a kind-spirited and affectionate little creature, and rarely gives way to bad temper.

<sup>1</sup> *Ce'bus hy-po-leu'cus.*

It has a wrinkled and care-worn face, as if burdened with sorrows—which most captive monkeys certainly are! Its forehead, throat and shoulder-points are white, and the remainder of the body is either gray, brown or jet black. The Sapajous inhabit Central America and northern South America. About two hundred specimens are brought to New York every year,



BLACK-FACED SPIDER MONKEY.

*At'c-les a'ter.*

where they are sold by dealers at prices ranging from \$10 to \$15 each.

**The Spider Monkeys**<sup>1</sup> may easily be recognized by their very long, slender legs and tails, and small, round heads. In color they are usually either black or gray, and rarely reddish brown. As they swing on their way through life, always using their prehensile tails to cling or to swing by, they have a very uncanny look, and it is no wonder that they are called "Spider" monkeys. They can come as near tying themselves into knots as living mammals ever can.

<sup>1</sup> *At'e-les.*

When fully grown, they are much larger than the sapajous, but are weak, unable to fight, and therefore timid. In a cage containing several species of monkeys, they are always the greatest cowards, and often are heard shrieking from fright at imaginary terrors. They are dainty feeders, and very difficult to keep in health in captivity. Four species are found north of Panama. The **Mexican Spider Monkey** occurs up to Lat. 23°, and is the most northern monkey on this continent.

**The Owl Monkeys.**—Next to the spider monkeys is found a group often represented in captivity, the members of which are distinguished by their small size, their round heads, very large, owl-like eyes, and long, hairy tails, which are not prehensile. As their staring eyes suggest, these creatures are of nocturnal habits, and in daylight hours are as inactive and uninteresting as opossums. Because of this, they make rather uninteresting pets; but being good-tempered creatures, they are frequently kept. They are sometimes called **Do-rou-cou'lis**. They are found from Central America to southern Brazil.

**The Squirrel Monkeys** of northern South America and Central America are next in order, and in activity and general liveliness of habit they make up for all that the owl monkeys lack. They are the most active of all the small American monkeys, and so nervous and unmanageable they are unfit for captive life elsewhere than in cages. The **Common Squirrel Monkey**,<sup>1</sup> sometimes, though erroneously, called the **Teetee**, is a trim little yellow fellow, with a very long cranium, close-haired head, and a very long tail, which it gracefully curls up over its own shoulders whenever it sits down. This species comes from the Guianas and Venezuela, and is very common in captivity.

On board ship a Squirrel Monkey of my acquaintance once furnished constant entertainment and amusement. Its favorite food was big, fat cockroaches, contributed by the sailors from their collection in the fore-castle. Each morning a sailor would bring a jacket, and shake it over a clear space on the deck. As the cockroach shower struck the deck, the agile little monkey dashed after the insects like a terrier at rats, cramming them into his mouth as fast as

<sup>1</sup> *Sai-mi'ri sci-u're-a.*

possible, and meanwhile seizing and holding in his hands as many more of the struggling insects as his absurd little paws could grasp.

This creature is a skilful climber, and it is the only mammal I ever saw which could exert sufficient lateral pressure with its hands and feet to enable it to climb with ease a perfectly smooth, right-angled corner of wood to a height of six feet.

This particular animal was so fond of its owner that it loudly and vociferously refused to sleep elsewhere than in his bunk, cuddled against his feet. With its piercing cries it controlled the situation as effectually as any spoiled child.

**The Saki Monkeys**, of tropical South America east of the Andes, are of medium size, mostly black and shaggy-haired, and sometimes possessed of a *long, black chin beard*. They are always marked by their big, heavily-haired tails, which are long, but not prehensile. They are often mistaken for howling monkeys. They are difficult to keep alive, seldom live to reach the United States, and for this reason are likely to remain but little known. The most remarkable species is the **Black Saki**,<sup>1</sup> two specimens of which were placed on exhibition in the New York Zoological Park in 1903.

**The Uakari, or Yarkee, Monkeys**, of which there are three species, all found in Brazil, have the shortest tails to be found amongst American monkeys. The **Bald Yarkee**<sup>2</sup> of the Upper Amazon is an excellent imitation of the Japanese red-faced monkey, having not only the same stubby tail, and long, shaggy hair, but also a *red face!* Unfortunately this species is one of the rarest in all America.

**The Howlers** are rarely seen in captivity, because it seems almost an impossibility for man to find food which they will eat, and which agrees with them.

Between the two sides of the lower jaw, the Howler possesses a large sound-box of cartilage—a development of the hyoid bone—which gives to the creature's voice a deep resonance, of a very unusual character. These monkeys delight to indulge in vocal concerts, and the deep roar of their unearthly voices can be distinguished at a distance of a mile or more.

In all there are six species of Howling Monkeys. Occasionally young specimens of the **Golden Howler** are brought from Venezuela

<sup>1</sup> *Pithecia sa-tan'as*.

<sup>2</sup> *U-a-ka'ri-a cal'va*.

and Guiana to New York, but in confinement their digestive organs are easily disturbed, and they seldom, if ever, live to reach maturity.

## THE FAMILY OF MARMOSETS.

### *Callithricidae.*

Lowest in the scale of all the American monkeys, and in fact next to the lemurs, we find a collection of small and odd-looking creatures, some of which are so strangely formed that it often is necessary to state that they belong to the Order of Apes and Monkeys. This is the Family of **Marmosets**, the members of which are distributed variously from southern Mexico to southern Brazil. They are frequently found in the stores of animal dealers, and by ladies who have abundant time for their care are often



Photo. by JENNESS RICHARDSON.

COMMON MARMOSET.

prized as household pets. But they are very delicate, and do not long endure the strain of being on public exhibition. Their market price varies from \$3.50 to \$8.

Without exception these are all very small,

delicately-formed creatures, with hairless faces, eyes that are large and bright, and long tails. Their hair is long, abundant and silky, and in some species it stands up on the top of the head like a white ruff. As these frail little creatures perch motionless in their cages, and focus their brown eyes upon the visitor, they seem more like little toys than living animals of Man's own Order. They are really very odd, picturesque and interesting.

The **Pinche Marmoset**<sup>1</sup> is a good representative of this group. It comes from the United States of Colombia, is about as large as a small chipmunk, and can be recognized anywhere by the jaunty bonnet of white hair which stands stiffly erect on the top of its head.

Of marmosets there are altogether about twenty-one species. The best-known are the **Common Marmoset**,<sup>2</sup> with a fan of white hairs standing stiffly erect above each ear, and the **Silky Marmoset**,<sup>3</sup> which is half buried in a mop of long, silky, yellowish hair.

### THE SUBORDER OF LEMURS.

#### *Lemuroidea.*

On the great island of Madagascar there are no fewer than thirty species of lemurs, many of them very beautiful creatures, all very kind-spirited and inoffensive, and so numerous that some travellers have declared that "every bush has its lemur." And yet, in America, these creatures are about as little known as if they inhabited Mars instead of Madagascar. During the first six months following the opening of the Primates' House in the Zoological Park, at least twenty educated and intelligent young men asked how to spell the word "lemur."

The lemurs, tarsiers and aye-aye constitute the lowest grand division of the Ape-and-Monkey Order—Primates. Their low position is due chiefly to their long, fox-like muzzles, and their teeth, which are not monkey-like. Their hands and feet, however, define their position.

The **Ruffed, or Black-and-White Lemur**<sup>4</sup> is the handsomest and most conspicuous animal in this strange group. It is the size of a large house cat, its tail is very long, and the creature is abundantly clothed with long, soft, silky-fine fur, jet black and pure white.

<sup>1</sup> *Mi'das aed'i-pus.*

<sup>2</sup> *Mi'das ros-a'li-a.*

<sup>3</sup> *Cal'li-thrix jac'chus.*

<sup>4</sup> *Le'mur va'ri-us.*



SANBORN, Photo., N. Y. Zoological Park.

THE RUFFED LEMUR.

Although lemurs have large eyes, and are supposed to be night-prowlers, they are fairly active in the daytime, and are not at all disturbed by daylight. They are charming pets, very affectionate, easily kept, and even with twenty in one large cage they do not quarrel, as monkeys are so prone to do.

**Keeping Monkeys in Captivity.**—Large monkeys need large cages, with means to climb and swing. Fine hay should cover the floor. Cages should always stand three feet above the floor of a room, and while the ventilation should be good, there should be freedom from draughts. The temperature should be 75°, kept as even as possible. *Food*: boiled rice or tapioca, baked or boiled potatoes, ripe bananas or apples; a little raw meat, finely chopped; dried or parched sweet corn that is easily chewed; a little stale bread; occasionally, a small raw onion. Permit no teasing; feed regularly, water frequently, and keep cages clean. When monkeys become ill, carefully ascertain their trouble, then treat them the same as one would sick children.

## CHAPTER III

# THE ORDER OF FLESH-EATING MAMMALS

### FERAE, OR CARNIVORA

North America contains a fine array of animals belonging to the **Order Ferae**,<sup>1</sup> numbering about ninety species north of Mexico, not counting subspecies. They are divided into the following groups:

	FAMILIES.	APPROXIMATE NUMBER OF SPECIES NORTH OF MEXICO.
}	THE CATS . . . . .	8 Species
	THE DOGS . . . . .	22 "
	THE MARTENS . . . . .	46 "
	THE BEARS . . . . .	12 "
	THE RACCOONS . . . . .	3 "

### THE CAT FAMILY.

#### *Felidae.*

In the order of their size, the five largest cat-like animals of North America are the following: Jaguar, Puma, Canada Lynx, Red Lynx, and Ocelot.

Of the Cat Family, the **Jaguar**<sup>2</sup> (pronounced Jag' you-ar) is not only the largest, but also the handsomest species in America. Of yellow-and-black Cats it stands next in size to the tiger, but in form it is not so finely proportioned as the leopard. It is of massive build, throughout, and its head is very large for the height and length of the animal. Its tail, however, is disproportionately short.

This creature has a golden-yellow coat, marked on the back and sides by large, irregular hollow islands of black, called rosettes—quite different from the smaller and more solid black spots of the leopard. Between these rosettes run the narrow lines of yellow ground-color, like the streets of an oriental city on a map. The legs, head and under-parts are marked with solid black spots. An animal of this species can always be

*recognized by its large rosettes, large head, heavy build, and short tail.*

The Jaguar, which in Mexico and South America is called "el Tigre" (tee'gree), is found as far north as southern Texas, and from that region southward to the limit of tropical forests in South America. A female specimen which once lived in the New York Zoological Park, measured 48 inches in length of head and body, its tail was 20 inches long, it stood 24 inches high at the shoulders, and weighed 120 pounds. The big and burly male which murdered the female above mentioned is fully one-fourth taller, and larger in every way.

In killing pigs, cattle, horses, deer and other wild animals, the Jaguar is a fierce, powerful and dangerous beast; but, like all other wild creatures, it is afraid of man.


It is my belief that the strength of the jaws of the Jaguar is greater in proportion to its size than that of any other member of the Cat Family. Of this power we once witnessed in the Zoological Park a tragic illustration. A full-grown female Jaguar was purchased as a cage-mate for a large and powerful male, named "Lopez," from the interior of Paraguay. After two days' preliminary introduction through their cage-fronts, the two animals were placed together. No sooner had the female entered the cage of Lopez

<sup>1</sup> From Latin *fe-rus*, meaning a wild beast. This is a much older name than *Carnivora*, which heretofore has been generally applied to this group.

<sup>2</sup> *Felis on'ca*.

than he rushed upon her, seized her neck between his jaws, and by a square bite crushed two of the neck vertebrae, and killed her instantly—as quickly as if her head had been cut off with an axe.

adventures with Pumas have been written and printed, but in reality this animal is less to be dreaded than a savage dog. It appears to be true, however, that it occasionally follows belated hunters or travellers, out of curiosity. It



Drawn by J. CARTER BEARD.

#### JAGUAR.

**The Puma**, also called **Mountain "Lion"** and **Cougar**,<sup>1</sup> is the most widely-known cat animal of North America. It is found in all the great western mountain ranges of the United States, in many tracts of "bad-lands" in Wyoming and Montana, British Columbia, and in the Adirondacks and Florida. Southward it ranges over table-lands and through tropical forests, all the way to Patagonia. In the United States it is most abundant, and also most accessible, in Routt Co., Colorado, where it is easily found by dogs, chased into low trees, and shot without danger. In this manner Mr. John B. Goff has killed nearly three hundred Pumas, "only two of which fought courageously."

Hundreds of thrilling stories of (imaginary)

<sup>1</sup> *Felis concolor*, and other species and races recently described.

is now a well-established fact that prowling Pumas do sometimes scream, in a manner calculated to inspire terror, just as caterwauling cats frequently do. I have heard Pumas scream precisely like terrified women or boys, but they always flee from man when the way is open.


The Puma is a *thin-bodied, flat-sided animal, tall for its weight, and of a brownish drab color*. It has a beautiful face, and is a handsome creature. Of all the large cats of the world, it is by far the best climber. A large specimen is from 7 to 8 feet in total length, from nose to tail tip, and weighs about 225 pounds.

The Puma makes its den among rocks, in "wash-out" holes, or in very thick brush or forests, and preys upon every living creature that can be killed and eaten, except man. In settled regions they frequently destroy much young

stock. Throughout the Rocky Mountains, it is a dangerous enemy of the mountain sheep and mule deer. In the "bad-lands" of Montana I once saw a mule deer killed which had on its neck a twelve-inch scar, a torn ear, and the beam of

are not possessed by any other animal. But no two Ocelots are ever marked exactly alike.

This animal is the size of a cocker spaniel, and being a good climber, when in its native forests it spends much of its time on the lower



From a photograph.

By permission of *Outdoor Life Magazine*.

PUMA, OR MOUNTAIN "LION."

one antler broken off half-way up. Apparently these injuries were received in an encounter with a Puma, and a fall over a cut bank, which evidently released the deer from its savage assailant.

The young of the Puma vary in number from two to five, and are *spotted*. Living specimens vary in value from \$30 to \$75, according to age and size.

At first glance the **Ocelot**, or **Tiger-Cat**,<sup>1</sup> seems to be a small leopard with a pale-yellow body-color. Its legs are spotted, but instead of having spots on its body, its back and sides are marked with irregular stripes and bands of black which run *lengthwise*. It may be instantly recognized by its *horizontal stripes*, for the like


<sup>1</sup> *Felis pard-a'lis*. See page 42.

branches of trees, watching for prey. It feeds chiefly upon small quadrupeds and birds. The following are the dimensions of an average specimen: Height, 13 inches; head and body, 30 inches; tail, 15 inches; weight, 36 pounds. It is frequently taken in southern Texas—its northern limit—and its range is about the same as that of the jaguar. In the New York Zoological Park it has been kept out-doors all winter, and has bred and reared young very successfully. Like most small yellow cats, Ocelots are usually bad-tempered. The value of a living specimen is about \$30.

**The Lynxes** of North America form a very distinct group of short-tailed, heavily-furred, tree-climbing cats, the members of which are spread throughout nearly all portions of the con-



continent north of Mexico, which are yet sufficiently wild to shelter them from man. They inhabit with equal facility forests, mountains, canyons,



Drawn by J. CARTER BEARD.

CANADA LYNX.

sage-brush plains, and even deserts. They prey chiefly upon rabbits and hares, grouse, prairie-"dogs," ground squirrels, and any other living creatures, except porcupines, which they can catch and kill. They are not courageous, or disposed to fight except when cornered, and so far as voluntarily attacking human beings is concerned, Lynxes are no more dangerous than rabbits.

In North America the genus *Lynx* is represented by two well-marked types.

**The Canada Lynx**<sup>1</sup> is a heavily-furred, short-bodied, long-legged bob-tailed wild cat of a pepper-and-salt gray color, standing about 18 inches high at the shoulders. It is readily recognized by the *long pencil of stiff, black hair rising from the tip of each ear, and its huge, hairy paws*. Its big eyes and long side whiskers give it a really terrifying countenance, particularly when it snarls. To the lone hunter who camps in the dark and gloomy forests inhabited by this creature, it seems a very dangerous animal; but in reality it is not so. Those who have hunted it say it is not courageous, and at close quarters is easily killed with a stick. It is a good climber, swims well, but on land runs rather poorly, with a galloping gait. Although found in a few localities in the northern United States, its real home is in the provinces of Quebec, Ontario, and the

<sup>1</sup> *Lynx can-a-den'sis*.

Northwest, up to Latitude 60°. A good average-sized male specimen collected by Professor Dyche in British Columbia measured as follows: Height, 17½ inches; head and body length, 32 inches; tail, 5 inches; girth, 17½ inches.

The weight of a full-grown specimen is 22 pounds, and the young are two in number. This species is rarely seen in captivity, and is always desired by zoological parks and gardens. Living specimens are worth from \$10 to \$40 each.

**The Bay Lynx**<sup>2</sup> is also called the **Red Lynx**, **Wild Cat** or **Bob Cat**, according to the locality in which it is found. Owing to variations in its color, and in some other characters, several subspecies have been described, but these are too closely related to the type to be set forth separately here. This species is marked by the absence of the long ear-pencil of the Canada lynx (although sometimes a small pencil is present), by the small feet and the warm brown tone in the color of the fur.

Western specimens are sometimes so strongly marked with round black spots that we feel impelled to recognize the "Spotted Lynx" as a distinct species; but when we find others from the Atlantic coast also spotted, besides others of the standard reddish gray, we are compelled to refer all of them to the species of the Bay Lynx. In the Atlantic states, the standard color for this



E. R. SANBORN, Photo., N. Y. Zoological Park.

THE OCELOT.

animal is a mixture of rusty red, gray and blackish brown, with the red so prevalent as to have given a name to the creature. In the West, the

<sup>2</sup> *Lynx rufus*.

spotted coat is more common, and occasionally the spots are strongly marked all over the animal.

The face of the Bay Lynx is really very beauti-




Photo. and copyright by W. L. UNDERWOOD, 1902.

BAY LYNX.

ful, and when not too fat from overfeeding in captivity, the body is lithe and graceful. When kept in large cages in the open air and sunlight, sheltered from storms, and not overfed, this animal is easily kept in fine condition. In artificially heated buildings they do not thrive.

This species is found in nearly all the states east of the Mississippi which contain large areas of rough forests, but are most numerous in Maine, the Carolinas, Florida, Virginia and Tennessee. In the "bad-lands" and mountains of Montana, Wyoming, Colorado and Texas they are really numerous, and feed luxuriously on the cottontail rabbits that are now so abundant in that region. Varieties of this species extend westward to the Pacific coast states. East of the Mississippi River, an average of about twenty specimens are caught alive each year, and offered for sale. Their value when caught is \$10 each, and the supply exceeds the demand.

By measurement the Red Lynx is fully as large as the Canada lynx. The largest specimen that ever came into my hands (on Pryor Creek, Montana) measured in length of head and

body 31 inches, tail 7 inches, height at shoulder 18 inches, and weighed 18 pounds. The largest of nine specimens killed by Mr. Roosevelt's party in Routt Co., Colorado, in 1901, weighed 39 pounds. One killed near Asheville, North Carolina, in 1900, is reported to have weighed 51 pounds.

No lynxes are found in the lowlands of the tropics, or in South America.

### THE DOG FAMILY.

#### *Canidae.*

Of all the wild creatures of North America, none are more despicable than wolves. There is no depth of meanness, treachery or cruelty to which they do not cheerfully descend. They are the only animals on earth which make a regular practice of killing and devouring their wounded companions, and eating their own dead. I once knew a male wolf to kill and half devour his female cage-mate, with whom he had lived a year.

In captivity, no matter how well yarded, well fed or comfortable, a wolf will watch and coax for hours to induce a neighbor in the next cage to thrust through tail or paw, so that he may instantly seize and chew it off, without mercy. But in the face of foes capable of defence, even gray wolves are rank cowards, and unless cornered in a den, will not even stop to fight for their own cubs.



GRAY WOLF.

**The Gray Wolf, or Timber Wolf,<sup>1</sup>** is really a formidable animal, but in its dealings with men,

<sup>1</sup> *Canis occidentalis.*

it has learned to fear the deadly rifle, the poison pot, and the trap. Storms, cold and fatigue affect it but little, and its powerful teeth, strong jaws and wide gape enable it to bite with great cutting power. In fighting with dogs, every well-aimed snap means either a deep wound or a piece of flesh bitten out.

The type of this species is a strong, robust animal, cunning and merciless. Its winter coat is long, shaggy and coarse-haired. Its standard color is mixed black and white, but it varies greatly, and unaccountably. In Florida it is often black, in Texas reddish brown, and in the far North it varies from black to white. Although in some localities it is called the Timber Wolf, it is equally at home on the treeless prairies of the West, in the dark, evergreen forests of British Columbia, and on the desolate barren grounds of Arctic America.

Although once very abundant on the great plains, the coming of the cattle ranch and sheepherder provoked against the Gray Wolf and coyote a relentless war of extermination, which still is being waged. Several states in the cattle country of the great plains offer cash bounties on wolf scalps ranging from \$2 to \$10, and large sums of money have been paid out for them. In Montana the number of wolves has so greatly diminished that in the course of a month in the saddle in 1901, in wild country, no Gray Wolves were seen, and only four coyotes. Wolves have now become so scarce that the occupation of the professional "wolfer" is almost gone.

Nevertheless, even on the cattle plains, the Gray Wolf is very far from being extinct; and as long as the "bad-lands" remain, with their thousands of wash-out holes, and tens of thousands of rabbits, the gray marauder will remain. In the far North, above the Arctic Circle, and in the land of the musk-ox, in 1899, Mr. C. J. Jones and his companion were so beset by packs of huge and fierce White Wolves, seeking to devour their five living musk-ox calves, that for over forty-eight hours they fought them continuously at short range, killing a wolf at every shot.

The young of the Gray Wolf are usually five in number, and are born early in May. At first they are of a sooty brown color, and are distinguishable from coyote puppies by the large size of the head. One which was examined

when four days old measured  $9\frac{1}{2} + 3$  inches, and weighed 16 ounces. When twenty days old, it was  $15 + 4$  inches, and weighed  $4\frac{1}{2}$  pounds.

The cry of the Gray Wolf is a prolonged, deep-chested howl, corresponding with B-flat below middle C, not broken into a bark, like the cry of the coyote. When seen at home, the Gray Wolf can readily be distinguished from the coyote, even at a distance, by the way it carries its tail,—pointing above the horizon.

Gray Wolves hunt in packs, often in relays, and successfully pull down deer, antelope, and wounded animals of all sizes. In the cattle country their specialty is the destruction of calves and colts. Except in the far North, they know well what firearms are, and are very careful to keep out of rifle-shot.



N. Y. Zoological Park.

COYOTE.

To-day the range of the Gray Wolf embraces the Great Plains and the Rocky Mountain region from Mexico to the northern limit of land. Lockwood and Brainard found tracks of a Gray Wolf at Latitude  $83^{\circ} 24'$ . In Alaska, animals of this species grow larger than in the United States, and frequently are white instead of gray. A fairly large Gray Wolf is  $48\frac{1}{2} + 15\frac{1}{4}$  inches long, stands 26 inches high at the shoulders, and has a girth measurement of  $29\frac{1}{2}$  inches. (L. L. Dyche.)

**The Coyote, or Prairie Wolf,**<sup>1</sup> is about one-third smaller than the gray wolf, but in form and color the two species look very much alike. It carries its tail low—humbly—as befits a cow-

<sup>1</sup> *Canis latrans*, and related forms.

ardly animal. It is not dangerous to man, and never was, and is bold only in the persistence with which it hangs upon the outskirts of civilization, and prowls around ranches in quest of food.

The delicacy of the Coyote's judgment in keeping always beyond fair gun-shot is truly wonderful. If he is not a mind-reader, his actions belie him. Twice in Montana, each time for two weeks, have I tried my utmost to shoot a Coyote; but during those periods not one would offer more than a running shot at three hundred yards or more. Twice, however,—and immediately after the above,—when riding quite unarmed, have Coyotes sat down beside the trail, waited for me to approach within forty yards, then yawned in a bored manner, and slowly trotted off. It is my belief that those animals knew perfectly well my inability to shoot.

The food of Coyotes consists chiefly of prairie-"dogs," ground-squirrels, sage grouse, hares and rabbits. The largest animals ever killed by them are deer and prong-horned antelope. From the ranchman they steal poultry, pigs, lambs and sheep. They "den" in "wash-outs," or deep holes in the cut banks of ravines,

and rear from five to seven puppies every May.

The cry of the Coyote is a dog-like yelping, half howl and half bark; whereas, the call of the gray wolf is a prolonged and steady deep-bass howl. As far as they can be heard, these wolves can be distinguished by their cries, and to those who have camped on the plains, or in the wild and weird "bad-lands" of the great West, the high-pitched, staccato cry of the Coyote as he announces the coming dawn, is associated with memories of vast stretches of open country, magnificent distances, fragrant sage-brush and freedom. The specific name of this animal (*latrans*) means "barking," and was bestowed on account of its peculiar dog-like cry.

The Coyote ranges from the latitude of the City of Mexico northward through the Great Plains and Rocky Mountain region to Alberta. The size of my best Montana specimen was 37 $\frac{3}{4}$  + 16 inches in length, and 20 $\frac{3}{4}$  inches in height at shoulders.

Coyotes vary in color from the typical pepper-and-salt gray to yellowish gray, the latter being found in the Southwest. At rare intervals, black specimens occur.

	GENUS.	ENGLISH NAME.	LATIN NAME.	LOCALITY.
<b>NORTH AMERICAN FOXES,</b> north of Mexico.	<b>Red Fox Group:</b>  GENUS <i>Vulpes</i> .	<b>Red Fox,</b> . . . .	<i>Vulpes fulvus</i> (Desmarest).	Virginia to Alaska.
		<b>Cross Fox,</b> . . .	<i>Vulpes fulvus decussatus</i> (Desmarest).	New York to Manitoba.
		<b>Black Fox,</b> . . .	<i>Vulpes fulvus argentatus</i> (Shaw).	Northwest Territory, Alaska.
		<b>Plains Fox,</b> . . .	<i>Vulpes macrourus</i> (Baird).	Great Plains.
		<b>Kadiak Fox,</b> . . .	<i>Vulpes harrimani</i> (Merriam).	Kadiak I., Alaska.
		<b>Newfoundland Fox,</b>	<i>Vulpes deletrix</i> (Bangs).	Newfoundland.
		<b>Swift Fox,</b> . . . .	<i>Vulpes velox</i> (Say).	The Great Plains.
	<b>Gray Fox Group:</b>	<b>Large-Eared Fox,</b> .	<i>Vulpes macrotis</i> (Merriam).	Southern California.
		<b>Arctic, or Blue Fox,</b>	<i>Vulpes lagopus</i> (Linnaeus).	Polar regions of both hemispheres.
		<b>Hall Island Fox,</b> .	<i>Vulpes hallensis</i> (Merriam).	Hall Island, Bering Sea.
	<b>GENUS</b> <i>Urocyon</i> .	<b>Gray Fox,</b> . . . .	<i>Urocyon cinereoargenteus</i> (Schreber).	Southeastern States.
		<b>Florida Gray Fox,</b>	<i>Urocyon cinereoargenteus floridanus</i> (Rhoads).	Florida.
		<b>Scott's Gray Fox,</b> .	<i>Urocyon cinereoargenteus scottii</i> (Mearns).	New Mexico to Southern California.
		<b>Texas Gray Fox,</b> .	<i>Urocyon cinereoargenteus texensis</i> (Mearns).	Texas.
<b>Coast Gray Fox,</b> .		<i>Urocyon cinereoargenteus californicus</i> (Mearns).	Southern California.	
	<b>Townsend's Gray Fox,</b> . . . . .	<i>Urocyon cinereoargenteus townsendi</i> (Merriam).	Northern California.	

**The Red Fox.**<sup>1</sup>—Of the many handsome and valuable species of foxes inhabiting North America, our wise old friend, the Red Fox, is the one most widely distributed, and the best



SANBORN, Photo., N. Y. Zoological Park.

RED FOX.

known. Between the southern Alleghenies and Point Barrow it appears in coats of many different shades, but everywhere it is recognizable by the prevailing yellowish-red color from which it derives its name. It is palest in the desert regions, where shade is scarce, and brightest in the forest regions and Alaska, where the bleaching power of the sun is not so great. The largest and finest skins come from Alaska.

The range of the Red Fox is very wide. From North Carolina and Tennessee it extends through the whole northeastern United States, westward to Montana and northward to the limit of trees. It is the most common fox in Alaska, wherever there are trees. It is so cunning, and so well able to take care of itself, even in populous countries, that it refuses to be exterminated. The length of an average specimen is 24 + 13 inches; height, 13 inches.

There is little pleasure to be derived from foxes kept in captivity as pets. They are very nervous, easily frightened, and, as a rule, are totally lacking in all the sentiments which resemble affection. Nevertheless, we have seen, and also owned, Red Foxes that were tame, and trustworthy when handled.

**The Cross Fox** is really a color phase of the red fox, marked by black legs and under parts, a dark-colored cross on the shoulders, steel-gray body and head, and a big black tail with a snow-

<sup>1</sup> *Vulpes fulvus*.

white tip. There is a reddish patch behind the fore-leg, and another on the side of the neck. In my opinion a really typical Cross Fox is the handsomest fox in the world, far more beautiful than the much-sought "silver fox." Some day it will attract the appreciation it deserves, and be sought accordingly. It stands between the red and the black foxes, and grades into both. It is found in Manitoba, Alberta, British Columbia, and Alaska, and occasionally in Idaho and Utah.

**The Black Fox**, commonly called the "Silver Gray" Fox (although there is no silvery color about it, save its tail-tip), enjoys the distinction of having the highest price on his head that is offered for any fur-bearer. In March, 1900, a single skin of this animal sold at auction in London for \$2,784; and it is not at all uncommon for extra fine skins to sell in this country at from \$600 to \$1,200. They are worth so much as furs for the very wealthy that zoological gardens cannot afford to purchase live specimens for exhibition. Their exhibition value is far below their fur value.

Like the cross fox, this is only a color phase of the typical red fox, but commercially the two forms are so distinct, and so sharply defined in dollars and cents, that they demand separate notice.



Drawn by J. CARTER BEARD.

BLACK, OR "SILVER" FOX.  
A subspecies of the Red Fox.

With the exception of its snow-white tail-tip, and a few scattering white hairs on the top of the hind quarters, a typical Black Fox is jet black. This form inhabits the same localities as the cross fox, and is much given to mixing with it, which causes many variations from their

standard colors toward the typical red fox. Both these animals are somewhat larger than the typical red fox found in New England.

On account of the great value of the fur of the Black Fox, many persons have desired to establish farms for breeding it in confinement, and several attempts in that direction have already been made. Thus far, however, none of them have proved successful. In Alaska, on the blue-fox farms, the Black Foxes are such dainty feeders that they will not eat the corn bread and fish which so well meet the wants of the other species, but require live game for food. Neither will they enter box traps, or permit themselves to be caught in any way other than in steel traps, which of course seriously injure them.

**The Swift Fox, or Kit Fox,**<sup>1</sup> is the smallest and daintiest of all our foxes. Its color is a beautiful silver-gray, with a tinge of yellow. It is strictly an inhabitant of the Great Plains region from the Rio Grande to the Saskatchewan, but owing to the readiness with which it eats poisoned meat that has been put out for wolves, it has already become very scarce. In spite of its name, it does not run with remarkable swiftness.

**The Arctic Fox.**<sup>2</sup>—This creature of the polar world is a striking example of climatic influence on a species, and also of the danger that lies



ARCTIC FOX.

in describing a species from a single specimen. In the far North, the Arctic Fox is snow-white all the year round. Farther south it is white in winter, but in summer is bluish-brown. In the southern part of its range, the Aleutian

<sup>1</sup> *Vulpes velox*.

<sup>2</sup> *Vulpes lagopus*.

Archipelago for example, except for an occasional white individual, it is dark all the year round, and is known only as the **Blue Fox**. At first it may seem difficult to believe that these two widely-different extremes are only color-phases of the same species; but it is quite true. The dark-colored animal is not even accorded subspecific rank.

The Arctic-Blue Fox is a simple-minded creature, of sanguine temperament, easily trapped and handled, and ever ready to adopt the prepared food of civilization. In its white phase, the finest skins sell in London at \$12 each. In its blue-brown coat, it has a very comical countenance, characterized by much hair, close-cropped ears, and a total absence of beauty; but its fur, when taken in season, is worth in the London market from \$25 to \$50 per skin.

On various islands along the Alaska coast, especially in the Aleutian Archipelago, about forty commercial companies are engaged in breeding Blue Foxes for their fur, some of them with satisfactory success. The foxes are fed daily, on cooked corn meal and dried fish. They come up to be fed, and when the time comes to handle and sort them previous to killing the annual allotment, they greatly facilitate matters by the readiness with which they enter box traps.

In the New York Zoological Park, three pairs of Blue Foxes that were received in 1902 from Alaska have taken kindly to captivity. The great decrease in the annual supply of good fur has caused many persons to hope that fox-breeding may be developed into a remunerative industry. Except in Alaska, no successful experiments in that line have been made, and it is quite desirable that fox-breeding in the United States should be taken up under state or national auspices, and wrought out to a successful issue. There is good reason to hope and believe that it might be developed into an important industry.

**The Gray Fox**<sup>1</sup> is the fox of the South, but it ranges northward far into the home of the red fox. It is noticeably smaller than the latter, pepper-and-salt gray above, and rusty-brown underneath, with a red patch on the side of its neck. For a fox it is very agile, and when hard pressed by dogs it can climb small trees up to a height of twenty feet or more.

<sup>1</sup> *Urocyon cinereo-argenteus*.

The five subspecies of the gray fox extend throughout the southern United States from Florida to California.

Besides the foxes already mentioned, several other species and races are recognized.



SANBORN, Photo., N. Y. Zoological Park.

GRAY FOX.

**THE SMALL FUR-BEARERS.**

*Mustelidae.*

A majority of the valuable fur-bearing animals of North America are found in a group of flesh-eaters known as the Marten Family. It contains about fifty full species, and its conspicuous types are the following:

**THE MARTEN FAMILY:**  
*Mustelidae.*

- |  |   |   |
|--|---|---|
| <p>                 Otter;<br/>                 Mink;<br/>                 Weasel;<br/>                 Marten;             </p> | } | <p>These four types are marked by long, slender bodies, very short legs, flattened heads, and general activity on foot.</p> |
|  | } | <p>Wolverine; the greatest glutton and pest in this Family.</p>   |
|  | } | <p>Skunk; aggressive and destructive pests; valuable fur-bearers.</p>   |
|  | } | <p>Badger; a fat-bodied, inert and practically harmless burrower.</p>   |

The great demand for fur, both for ornament and use, has brought about the systematic destruction of all fur-bearing animals. Many species that once were numerous have now become very rare. Formerly the wearers of fur ac-

cepted nothing less desirable than beaver, otter, mink and marten. To-day, the fur of the skunk, raccoon, fox, lynx, black bear and even the despised rabbit are in active demand, for garments and for trimmings.

**The Otter**<sup>1</sup> is as fond of water as a seal, and quite as much at home in fresh water as on land. Its regular food consists of fish, in the capture of which it is very expert. It has webbed feet, a thick, pointed tail distinctly flattened for use in swimming, and it is clothed with a thick coat of very fine, dark brown fur. Strange to say, when fairly treated, the Otter is a good-tempered animal, tames easily when caught young, and makes an interesting pet. In a public park, one Otter is worth more to the public than twenty beavers.

In the days when they were numerous, and less persecuted than now, it was no uncommon thing for a party of Otters to select a steep and slippery river-bank, and slide down it repeatedly, as small boys slide down hill on sleds, except that each slide of the Otter always ended in a plunge into the water.

The Otter of North America still is found occasionally in Florida and the Carolinas, the Canadian provinces, in a few localities in the Rocky Mountain region, and from British Columbia to central Alaska. Outside of Alaska, its fur is taken so rarely that it has ceased to be regarded as an article of commerce. Its value alive for exhibition purposes is from \$10 to \$30. The length of a large northern Otter, head and body, is 27 inches and tail 16 inches.

The Otter builds no house, but lives in a bank burrow, usually under the spreading roots of some large tree growing near the water. The young are usually two in number.

**The Sea Otter**,<sup>2</sup> one of the most valuable of all fur-bearing animals, is literally a child of the ocean surges and the surf-beaten rocks of the rugged north Pacific coast. It is born at sea, on a bed of kelp, and literally "rocked in the cradle of the deep." It was formerly found from California to the Aleutian Islands, but is now very rare except in certain parts of Alaska. Here its pursuit is strictly limited by law to the natives, to whom it is vitally important, and a white man may not kill a Sea Otter except under penalty of a fine of \$500.

<sup>1</sup> *Lutra can-a-den'sis.*

<sup>2</sup> *La'tax lu'tris.*

The fur of this creature is extremely valuable. In March, 1900, the finest skin in the London market sold for \$1,344. A full-grown specimen

Otter usually is quite dangerous, but to the natives of the Alaskan Peninsula, this creature is far more important than the fur seal. For-



1. OTTER.  
2. FISHER.

3. MARTEN.  
4. MINK.

measures from 3½ to 4 feet in length (head and body) and has a tail 11 inches long. Its fur is very dense and fine, and in color is a shimmering, lustrous black. The pursuit of the Sea

merly between five thousand and six thousand skins, worth from \$100 to \$500 each, were taken annually, and formed practically the sole dependence of the natives along nearly 2,000 miles



of coast line. But with the introduction of fire-arms, and the sealing schooners, the Sea Otter has been almost exterminated. The few individuals that remain are widely scattered, and are the wildest and wariest of all wild creatures.

**The Mink**<sup>1</sup> is much smaller than the otter, yellowish brown or dark brown in color, and while it prefers to live along the banks of streams, it is not an aquatic animal like the otter. When possible, it feeds chiefly upon birds, because they are easily caught and killed, and when opportunity offers, it is a wanton murderer. It also preys upon small mammals and fish, whenever it can procure them. In the Beaver Pond of the New York Zoological Park a murderous Mink once killed six wild geese in one night, and another slaughtered ten herring gulls.

A full-grown Mink looks very much like a large weasel, having a long, slender body and very short legs.

The Mink is by no means as rare as the otter, and even to-day is found scattered throughout nearly the whole of North America, as far as the limit of trees. The round, hairy tail, chocolate-brown or yellowish-brown color and smaller size of this animal quickly distinguish it from all other animals of its Family. The body of a full-grown specimen is about as thick as the wrist of a medium-sized man. The length of the head and body is 19 inches, tail 7 inches.

**The Black-Footed Ferret,**<sup>2</sup> of Kansas, Colorado, Wyoming and Montana, is to many persons who live in its home country, an enigma. In 1849 this pretty creature was described and illustrated by Audubon and Bachman, after which it totally disappeared, and remained a mystery until it was re-discovered in 1886. In its home it is often called the **Prairie-“Dog” Hunter**, because its specialty is the killing of prairie-“dogs;” and it is nearly always found in the towns of that jolly little animal. It can be recognized at a glance by its black feet, brown legs and black tail-tip, and the cream-yellow color of its head and body. Next to the skin, the fur is white, and there is a broad black or dark-brown patch across the nose, including both eyes. Its length of head and body is 19 inches, tail 4 inches. Regarding its habits and life history, much re-

mains to be ascertained by the young naturalists who live in the country it inhabits.

**The Weasel**, of which many species and races have been described, is the smallest animal in the marten family.<sup>1</sup> Its legs are very short and far apart, and its body is no thicker than a



BLACK-FOOTED FERRET.

man's thumb, but it is of such great length that the animal is positively snake-like in its proportions. In life it is very odd to see the front legs walk to and fro quite independently of the hind quarters. Fifteen full species have been described, several of them being very much alike. **The Common Weasel**, or **Ermine**<sup>2</sup> is brown in summer, and white in winter.

The Weasel is one of the most courageous and aggressive of all animals. It kills rabbits, grouse, chickens and ducks of ten or twelve times its own size, and often kills ten times as many chickens as it can eat, purely to gratify its murderous disposition. It is as savage as a tiger, but on farms it often does good service in destroying rats and field-mice. Weasels are so small their fur has little value, but the time is coming when it will eagerly be sought and used.

<sup>1</sup> The Least Weasel (*Putorius rixosus*), which is found from the Saskatchewan to Alaska, is said to be the smallest carnivore in the world.

<sup>2</sup> *Pu-to'ri-us nig'ri-e-a*.

<sup>1</sup> *Lu-tre-o'la vi'son*, and related species.

<sup>2</sup> *Pu-to'ri-us nig'ri-e-a*.

The **Marten**<sup>1</sup> looks very much like a young red fox, and in size it is about as heavy as an ordinary domestic cat. Its head and body length is 17 inches, and its tail 7 inches. The body is brownish yellow, the legs are two or three shades darker, and it has three kinds of hair. It loves timber, and spends much of its time in trees. It is rarely found in open country, and is most abundant on rugged and rocky forest-covered mountains.

The Marten is not a poultry-killer, nor a wan-

in America. It is a bold, active tree-climber, an industrious hunter, an aggressive fighter, and as a stealer of baits it is almost as great a nuisance to trappers as the hated wolverine. With this animal, "all's fish that cometh to net," and with equal relish it devours dead fish, rabbits, squirrels, chipmunks, ground birds, snakes, toads and frogs. Occasionally it murders its own cousin, the pine marten, and even feeds upon the Canada porcupine.

The Fisher is at home in the swamps or the



THE WOLVERINE.

ton murderer of more game than he can eat, but he lives by honest hunting of wild game. His food consists of small rodents, birds, eggs, or even an occasional reptile. In the United States this animal is now rare, for its fur has always been highly prized. It is often called the **Pine Marten**.

The **Fisher**, or **Pennant's Marten**,<sup>2</sup> is one of the largest members of the Marten Family

<sup>1</sup> *Mus-le-la americana*.

<sup>2</sup> *Mus-le-la pen'nant-i*.

rocky mountain-sides of northern New York, and in the forest regions of North America generally from Maine and southern Labrador to the Pacific coast. Northward it ranges to Great Slave Lake and the Yukon River. In color it varies from glossy black to dark brown, with occasional gray, or grayish white, on head and neck, chin, chest and abdomen. Its average length is 23 + 14 inches. The young vary in numbers from two to three.

**The Wolverine, or Carcajou,**<sup>1</sup> is one of the most remarkable animals in North America. It is about the size of a full-grown bull-dog, has a ravenous appetite, great strength, a fierce temper, and the combined cunning of many generations of criminals. It is the greatest thief amongst animals, and is such a greedy feeder that it is known to many as the **Glutton**. It will follow a trapper's "line" of marten traps, for miles, destroy every animal it finds in them,

the *Skunk-Bear*, and in Washington the Indians call it the *Mountain Devil*. It inhabits the northern Cascades and the Rocky Mountain region of the United States as far south as Great Salt Lake, and the whole of arctic and subarctic America to the northern limit of trees. It is especially abundant on the Kuskowim River, Alaska. Its length is 32 x 10 inches.

**The Skunks** form a large group, widely distributed, but all the species, however much they



COMMON SKUNK.

LITTLE SPOTTED SKUNK.

devour baits, and sometimes steal the traps also.

It breaks open caches, raids cabins, and systematically destroys everything it encounters. It is the only animal living which maliciously and deliberately destroys property, and soils food which it can neither eat nor carry away. It steals articles which it cannot possibly use, and more than once has been known to strip a cabin of nearly its entire contents.

In form this animal resembles a cross between a badger and a bear. In Wyoming it is called

<sup>1</sup> *Gu'lo lus'cus*.

differ in size or color, are arranged in three genera.

**The Common Skunk,**<sup>1</sup> of which nine species are recognized, is very well known, chiefly because of its powerful odor, its wide distribution, and its very conspicuous jet-black color, divided on the back by one or two broad bands of white.

This type of skunk is practically confined to the United States and Mexico, and is most abundant in the North. The very offensive fluid which constitutes its defence against all

<sup>1</sup> *Meph'i-tis*.

enemies, is contained in two glands situated near the base of the tail, and can be thrown several feet. Its odor is so offensive and so stifling that neither man nor beast can long endure it.

The Skunk is a bold marauder, and destructive to poultry, but nevertheless of value as a destroyer of white grubs and other noxious insects. Owing to the disappearance of the otter, beaver, mink and marten, the fur of the Skunk has become valuable, and is now very extensively used, the white portions being first dyed black.

**The Little Spotted Skunks**<sup>1</sup> are found chiefly in our southern states, and can immediately be recognized by the alternating bands of black and white which extend lengthwise along the body. Of these there are about a dozen species, but some of them are very much alike. They range from the Gulf coast north to West Virginia and Kansas, but on the Pacific slope they are found in Washington, Oregon, California and Utah.

**The Badger Skunks**<sup>2</sup> resemble the common Skunks in size, but may be readily distinguished by the broad white stripe on the back, and the powerful claws on the fore feet. As indicated by their name, they are more badger-like than



THE BADGER.

the other skunks, and are expert diggers. They are the only skunks which occur in South America, and their range extends from the Straits of Magellan northward along the west coast, through Central America and Mexico into southern Texas and Arizona.

**The Badger** is an animal of strange form, its body being very broad and flat, and its legs very

<sup>1</sup> *Spillo-gale.*

<sup>2</sup> *Co-ne-pa'tus.*

short. In size it stands midway between the common skunk and the wolverine. It has a savage and sullen disposition, and as a pet is one of the worst imaginable. It lives in burrows, and feeds on ground squirrels, prairie-"dogs," and ground game of every description. Often Badgers will be found living in deserts where it would seem an impossibility for any carnivorous animal to find a supply of food. Its home is the Great Plains, the Rocky Mountains and westward thereof to the Pacific coast, from Mexico to Manitoba and Alaska.

## THE BEAR FAMILY.

### *Ursidae.*

That nearly all young people, the whole world over, are greatly interested in bears, is no cause for wonder. Under proper conditions, young bears are the most merry-hearted wild animals that come into captivity, not even excepting monkeys, and in some respects the most interesting. Of all wild animals kept in zoological parks, there are none that more fully repay the care bestowed upon them, and excepting apes and monkeys, none that furnish more amusement. With plenty of sun-lit space in which to romp and play, good bathing pools, and no stone walls to depress their spirits, *if not fed by visitors*, bears are more playful and mirth-provoking than most monkeys. If immured in gloomy "bear-pits," or confined in small cages, their spirits are correspondingly depressed. They are then like unhappy prisoners, rather than care-free wild creatures. If tantalized with bits of food, they quarrel and fight, and their tempers become savage and dangerous.

Contrary to general belief, a bear is naturally cheerful and good-tempered. Elk, deer, buffalo, elephants and large cats often attack their keepers, but bears that have been properly reared in captivity seldom do so.

The bear dens of the New York Zoological Park, contained (in 1903) thirty-four bears, of eleven different species, living in peace and harmony, in nine paved yards. Fully one-half of their waking hours are spent in romping, wrestling, boxing and swimming, and ill-temper is rarely shown. The keepers go amongst these bears with only brooms for defence, and the great brutes are hustled about and driven to and fro



Drawn from life, in the New York Zoological Park, by J. CARTER BEARD.

ALASKAN BROWN BEAR.

From near the head of Cook Inlet, Alaska, and believed to represent *Ursus middendorffi*.

as if they were so many sheep. At the same time, any visitor who is so unwise as to thrust a hand between the bars within reach of the jaws of any of the inmates is certain to be very severely bitten,—in playfulness rather than rage! In their rough play these bears continually bite each other, without inflicting injury; and they do not appreciate the difference between a tender human hand and a tough, hairy paw.

Never offer a finger to a carnivorous animal, unless you really wish to have it bitten off. And do not feed pea-nuts, candy, peaches, or tobacco to animals in captivity. If you wish to kill any of them, a gun is far more respectable, and also more merciful.

**Structure and Habits of Bears.**—Bears are plantigrade, or flat-footed, animals, with long claws that are not retractile. They live on the ground, and eat all kinds of food, from green grass to elk steaks. A few species only are able to climb trees. In their food habits they are om-niv'o-rous, and devour almost everything they can chew, except wood and foliage. The bears of the Alaskan coast eat great quantities of marsh grass, and berries, but salmon is their regular food. All bears eat succulent roots, insect larvae, honey, frogs and also reptiles, fish, and every other kind of flesh they can obtain. In captivity they thrive best on a variety of food consisting of stale bread, raw meat, cooked meat, rice, raw fish, boiled potatoes, raw carrots, and fruit.

In the temperate zone, where the snow falls to a depth of a foot or more, bears are unable to procure food in winter, and pass that season in a sort of sleep, or hibernation. With its stomach and intestines empty, or nearly so, a bear enters its den in December, curls up, and with some of the functions of Nature entirely suspended, sleeps until spring! In reality, the creature lives upon the fat that has been secreted under its skin and elsewhere during the summer days of good living. Ordinarily, bears in captivity that are supplied with daily food, do not hibernate in winter, but one cinnamon bear which I knew personally, at Mandan, North Dakota, dug a hole in the prairie, entered it on December 17, and did not reappear until March 14, of the following year. In the tropics, bears never hibernate.

Naturally, the dens of hibernating bears are of several kinds, according to conditions. In the Adirondacks, of New York, the black bear often chooses the base of a hollow tree, or digs a cavity under the roots of a tree. In the "bad-lands" of the West, bears easily find warm and comfortable dens in the wash-out holes of rugged ravines. In the mountains, amongst rocks, small caves are easily found. In Washington, "Grizzly" Adams caught "Lady Washington" and "Ben Franklin" in a deep den that had been dug by their mother in a steep hillside.

All the world over, two bear cubs usually constitute a litter. In America, they are usually born in January, and at birth are ridiculously small, almost hairless, and as helpless as newly-born mice. Although they grow rapidly during the first year, they are seven years in reaching full maturity. In captivity bears seldom breed and rear their young, chiefly because of the lack of satisfactory seclusion for the female. Mr. Arthur B. Baker, who has recently inquired into the habits of the American black bear in captivity, states that "at Cuyahoga Falls, Ohio, are two specimens which regularly hibernate, and also a pair, born in 1888, which, with the exception of three years, have had cubs each January (21st to 27th) up to 1903, all of which were raised, excepting a few which met death by accident."

Bears have bred in captivity in the zoological gardens and parks of Philadelphia, Cincinnati, Washington and New York, but few of the cubs have been reared.

The dimensions of a Russian brown bear cub—a species that is an excellent understudy of our silver-tip grizzly, and but slightly inferior in size—was when two days old as follows: Length, head and body,  $9\frac{1}{4}$  inches, tail,  $\frac{1}{2}$  inch; height, 5 inches, circumference of chest,  $6\frac{3}{4}$  inches; hind foot  $1\frac{1}{8}$  inches by  $\frac{7}{8}$  inch; weight 15 ounces. This cub was born on January 17.

All American bears, except the polar, show great changes in the color of their pelage at different seasons of the year. In the late summer the new pelage is darkest, but by the following spring, the old coat has grown so much lighter in color that the wearer seems like a different individual. The shedding period is from May 1 to August 1.

**North American Bears.**—Leaving out of count the subspecies, and the species of which we know little or nothing, the world contains fourteen well-marked types of bears. Of these, eight inhabit Asia and Europe, four are found in North America, one is found all around the north pole, and one in South America. From both the Old World and North America, quite a number of additional species and subspecies have been described; but it must be remembered that at present we are dealing only with conspicuous types.

Owing to puzzling variations in color, claws and skulls, and the great difficulty of bringing together several hundred adult skins with skulls, it is at present impossible to state precisely how many different kinds of bears inhabit this continent, or how they are related. In the near future, however, many existing questions will be settled; and until then the wisest course for the student and the general reader is to accept only well-known facts, and wait with patience for more.

The bears of North America constitute four distinct groups, as follows:

**Polar Bear**, of the far North. White. Very large.

**Big Brown Bears**, of Alaska. Light brown. Very large.

**Grizzly Bears**, Mexico to Alaska. Gray or brown. Medium to very large.

**Black Bears**, North America generally from Mexico to Alaska. Black or brown. Medium size, and large.

To most persons, the second group of this series is quite new, and for several reasons its members are of unusual interest.

**The Polar Bear.**

**The Polar Bear** stands alone in its genus. It is the king of the frozen North, and its robe is pure white, all the year round. It inhabits the coasts of the Arctic Ocean, all around the pole, and wanders over the arctic islands and the great ice-fields almost as far north as man has ever gone. Nansen saw its tracks at Latitude 84°,—its farthest north.

As a rule, the Polar Bear follows the edge of the great ice-pack, for the sake of the seals and walruses which move with it, north in summer, and south in winter. He seldom travels more than a day's journey inland on any shore. His food consists chiefly of seals, walruses, fish and dead whales; at times of vegetable matter.

**FULL LIST OF THE BEARS OF NORTH AMERICA.**

*Corrected to December 1, 1903.*

	<b>Polar Bear</b> , . . .	<i>Thalarctos maritimus</i> (Phipps), . . .	Arctic regions generally.
<b>The Big Brown Bears.</b>	<b>Kodiak Bear</b> , . . .	<i>Ursus middendorffi</i> (Merriam), . . .	Kodiak Island, Alaska.
	<b>Yakutat Bear</b> , . . .	<i>Ursus dalli</i> (Merriam), . . .	Yakutat Bay, Alaska.
	<b>Peninsula Bear</b> , . . .	<i>Ursus dalli gyas</i> (Merriam), . . .	Pavlof Bay, Alaska.
	<b>Merriam's Bear</b> , . . .	<i>Ursus merriami</i> (Allen), . . .	Portage Bay, Alaska.
	<b>Kidder's Bear</b> , . . .	<i>Ursus kidderi</i> (Merriam), . . .	Chinitna Bay, Alaska.
	<b>Sitka Bear</b> , . . .	<i>Ursus sitkensis</i> (Merriam), . . .	Alaska coast, near Sitka.
<b>The Grizzly Bears.</b>	<b>Silver-Tip Grizzly</b> ,	<i>Ursus horribilis</i> (Ord), . . .	Wyoming to Alaska.
	<b>Sonora Grizzly</b> , . . .	<i>Ursus horribilis horriaeus</i> (Baird),	S-W. New Mexico.
	<b>Alaskan Grizzly</b> , . . .	<i>Ursus horribilis alascensis</i> (Merr),	Norton Sound, Alaska.
	<b>Barren-Ground Grizzly</b> , . . . . . }	<i>Ursus richardsoni</i> (Swainson), . . . }	Great Slave Lake and Barren Grounds.
<b>The Black Bears.</b>	<b>Black Bear</b> , . . .	<i>Ursus americanus</i> (Pallas), . . .	North America.
	<b>Labrador Bear</b> , . . .	<i>Ursus americanus sornborgeri</i> (Bangs)	Labrador.
	<b>Louisiana Bear</b> , . . .	<i>Ursus luteolus</i> (Griffith), . . .	Louisiana and Texas.
	<b>Everglade Bear</b> , . . .	<i>Ursus floridanus</i> (Merriam), . . .	Florida.
	<b>Glacier Bear</b> , . . .	<i>Ursus emmonsii</i> (Dall), . . .	St. Elias Alps, Yakutat Bay, Alaska.
	<b>Queen Charlotte Bear</b> , . . . . . }	<i>Ursus carlottae</i> (Osgood), . . . }	Queen Charlotte Islands, British Columbia.

In 1874, when Mr. Henry W. Elliott and Lieutenant Maynard visited St. Matthew Island, a lonely bit of treeless land in the northern portion of Bering Sea, they found upon it between 250 and 300 Polar Bears! The animals were basking in the warm sunshine, shedding their winter coats, and growing fat on the roots of the plants and mosses that grew there. On one occasion *twenty bears* were in sight simultaneously. The bears literally overran the island, grazing and



E. R. SANBORN, Photo, N. Y. Zoological Park.

POLAR BEAR.

rooting about like hogs on a common. They showed no disposition to fight, but always ran when approached.

The Polar Bear is a tall animal, with long legs, flat sides, and paws that are very wide and flat. The largest specimen in the New York Zoological Park is 50½ inches in height, 7 feet 2 inches in length, and weighs about 800 pounds. When standing erect on his hind legs, the end of his nose is 8 feet 8 inches from the ground. If properly and comfortably caged, and provided with a swimming pool five feet deep, Polar Bears in the temperate zone do not suffer from the heat of summer, and can endure hot weather fully as well as our black bears. Of course they require shade in summer; but it is not necessary to put ice in their pool to cool the water.

The power of this active, warm-blooded animal to resist cold is one of the wonders of Nature. With the temperature many degrees below zero, the Polar Bear cheerfully leaps into the Arctic Ocean, amid the broken ice, and swims for hours.

Of all bears, it is the best swimmer, and it dives with great ease. Thanks to the limitations imposed by the Frost King on hunting in the arctic regions, it is not very probable that the Polar Bear ever will be exterminated by man.

**The Big Brown Bears.**

In 1896 the specimens collected by the United States Biological Survey, at Washington, revealed to Dr. C. Hart Merriam the presence in Alaska of two or three species and subspecies of huge brown bears, totally different in character from all the American bears previously known. These bears range from Sitka around to the extremity of the Alaskan Peninsula, Kodiak Island, and inland for unknown distances. They are marked by their light brown color, high shoulders, massive heads of great breadth, short, thick claws, and shaggy pelage. In their high shoulders, they resemble the grizzly bear, but otherwise differ from them in many ways. Of these bears, Dr. Merriam has published preliminary descriptions of four new species and one subspecies, but additional collections and information may possibly result in the consolidation of some of these.

It is sufficient for our purpose to set forth only the species which seems most sharply defined, and which may be considered representative of the whole group.

**The Kodiak Bear,**<sup>1</sup> of Kodiak Island, and probably also of the Alaskan Peninsula and the mainland for some distance eastward, is not only the largest of all living bears, but also the largest carnivorous animal in the world. Several skins of immense size, and skulls 19 inches in length, have been collected. The largest specimen ever killed and measured by a naturalist was a female killed at Chinitna Bay, by Mr. James H. Kidder, which had a shoulder height of 51 inches.

A very large flat skin measured at Kodiak by Mr. J. A. Loring, was 9½ feet long by 10½ feet wide across the fore legs.

Immediately after shedding, the new coat of the Kodiak Bear is dark-brown, like that of a grizzly, but it soon changes to a beautiful golden-brown tint. In March and April, the old coat is of a golden-yellow color, and really very beautiful. The full coat is long, thick and shaggy, and

<sup>1</sup> *Ursus mid'den-dorf-fi.*



except when shedding is in full progress, the animal makes a very imposing appearance. This species is recognized by its uniform brown or golden color, its high shoulders, broad and massive head, flat forehead, short, square nose, and a drop in the upper line of the head in front of the eyes. Mr. Kidder states that the bears on Kodiak Island are uniformly colored over the body and legs, but those on the mainland are darker on the legs than on the body.

The Kodiak Bear catches and devours great numbers of salmon, which are so abundant in many Alaskan streams that it can throw them out with its paws. It also eats quantities of the rank marsh grass which grows along many salmon streams where they flow through alluvial plains before discharging into the sea. It inhabits the most rugged mountains, and is seldom killed save when it leaves the shelter of the timber and comes into the open river valleys and bay heads to feast on freshly-caught salmon, with tender grass for dessert.

Just how far eastward this bear ranges on the mainland, remains to be determined; but I believe it will be found as far as the Copper River. The big animal found in the Yukon valley, and commonly called the "Red Bear," undoubtedly belongs to the group of big brown bears, and in all probability is the same as the Kodiak Bear. The illustration shown on page 33 is a portrait of a fine Alaskan brown bear living in the New York Zoological Park, which came from the country between Cook Inlet and the Copper River. Inasmuch as all the descriptions of the species composing the brown bear group have been based chiefly upon skulls, the exact identity of our specimens can not be determined while they are alive. In the month of September its entire pelage is of the uniform dark-brown color characteristic of the bears of Kodiak Island at the season when the majority of them are killed, but later on the pelage of the body becomes lighter than that on the legs.

### The Grizzly Bears.

**The Grizzly Bear.**<sup>1</sup>—Of all the bears of the world, this species is certainly one of the most celebrated. During the days of muzzle-loading rifles, its name and fame inspired terror through-

out the mountains and foot-hills of the wild western domain which constituted its home. For many years it held the old-fashioned Kentucky rifle of the pioneer in profound contempt, and frequently when it was used to annoy him, the user met a tragic fate. I believe that Grizzlies have killed and maimed a larger number of hunters than all other bears of the world combined.

Down to the advent of the breech-loader, the Grizzly was a bold, aggressive and highly dangerous animal. When attacked, he would charge his enemies with great ferocity, striking terrible blows with paws that were like sledgehammers armed with huge hooks of steel. The combined swiftness and strength with which any large bear can strike must be seen or felt to be fully appreciated.

I have made many observations on the temper of the Grizzly Bear, and am convinced that naturally the disposition of this reputedly savage creature is rather peaceful and good-natured. At the same time, however, no animal is more prompt to resent an affront or injury, or punish an offender. The Grizzly temper is defensive, not aggressive; and unless the animal is cornered, or *thinks he is cornered*, he always flees from man.

Either in captivity or freedom, the Grizzly responds to fair treatment as well as any well-armed wild animal ever does, and far better than any other species with which I am personally acquainted. In the Yellowstone Park, where for several years past all bears have been fully protected, both the Grizzly and black bears now live in close touch with man, without breaking faith with him. Although they frequently visit the hotels, and steal food from the wagons and camps of tourists, I believe *no bear has yet broken faith with the Government* by molesting either his human neighbors or domestic animals! This fact speaks volumes for the moral character of our bears.<sup>1</sup>

The Grizzly is an animal of commanding appearance, and amongst other wild beasts it

<sup>1</sup>Since the above was written, the truce of the Yellowstone Park has been broken. Two horses belonging to a party of tourists have been killed by bears, and the aggressiveness of the latter has become so serious that it will be necessary for the government to take measures which will teach them to keep their place.

<sup>1</sup> *Ursus hor-ri'bi-lis*.

acknowledges no superior. A small Grizzly cub which we once set free in a mixed company of five or six bears of other species, all of which were larger than he, boldly stalked into the centre of the group, with an air of conscious superiority and courage that was both characteristic and amusing. It was the other bears who were frightened, not he!

Specimens of this species are readily recog-

very gray. The huge brown Grizzly of southern California, now very rare, has been described as a species distinct from the Rocky Mountain Silver-Tip. I once measured the dry skin of one of these animals, which was 9 feet 4 inches in length, and 10 feet 3 inches wide across the shoulders, between the ends of the front claws.

So far as I am aware, the largest Grizzly Bear



A GRIZZLY BEAR AT HOME.

Copyright, 1902, by F. C. Wolcott.

Photographed in the mountains of western Wyoming, by F. C. Wolcott. The bear was enticed by a bait to within thirty feet of the camera, and taken by flashlight.

nized by their high shoulders, powerful proportions, grizzly-gray hair, and long curved claws. The standard color (in winter) is brown next to the skin, the extremities of the hair being tipped with silvery gray, from which has come the common name of "Silver-Tip."

From Mexico and southern California to the Yukon valley, especially along the main ranges of the Rocky Mountains, the Grizzly shows about six different shades of color, from brown to sil-

ver actually weighed was one that lived and died in the Lincoln Park menagerie, Chicago, and was weighed by Mr. G. O. Shields. Its weight was 1,153 pounds; yet when alive, western hunters who saw it frankly admitted that it was larger than bears killed by them which they "estimated" at 1,800 pounds! Thus far the Rocky Mountains have not produced a wild Grizzly actually weighing over 800 pounds, and the average weight of the adult Grizzlies killed

in the United States during the last fifteen years has been between 500 and 600 pounds.

In all parts of the United States save the Yellowstone Park and the Clearwater Mountains of Idaho, the Grizzly is now a rare animal, and so difficult to find that it is almost useless to seek it this side of British Columbia. Like other large mammals of this continent, the long-range, high-power rifles leave them absolutely no chance

but is quite unable to climb trees. Like all other bears, he eats nearly everything he can chew, and is very partial to berries, and fruit of all kinds.

### The Black Bears.

**The Black Bear**<sup>1</sup> is the best known bear in North America. It is found in nearly all the mountains and great tracts of forest between



Photo. by E. R. SANBORN, N. Y. Zoological Park.

AMERICAN BLACK BEAR

From northern Wisconsin.

for their lives, and in a short time none will exist in the United States outside of the Yellowstone Park and the zoological gardens. In the wilds of Alaska, they may survive for perhaps a quarter of a century longer. Unfortunately, the Grizzly loves to roam over treeless mountains, on which his huge bulk makes him conspicuous for miles, and invites the attacks of his enemies. He loves water, swims well, and is a great traveller,


Florida and Alaska, and from Nova Scotia to the Pacific coast. During the past twenty years it has been seen or killed in forty states of the United States, in Mexico, Alaska, and eleven of the British provinces. Its farthest south is the mountains of Costa Rica.

Its standard color is jet black, all over, except the nose, which is dirty white or light brown. A

<sup>1</sup> *Ursus a-mer-i-can'us.*

very confusing fact about the Black Bear is the frequency with which it runs into brown or cinnamon colors. Sometimes black and brown cubs have been found in the same litter. Very curiously, however, this color is found only in the Rocky Mountains, and farther west. In its brown phase, this animal is called the **Cinnamon**

round on the hind quarters, low at the shoulders, and also by the fact that in walking it usually carries its head low. It is a smaller animal, and its claws are short and well adapted to tree-climbing. It conceals itself from its enemies much more successfully than the grizzly, and therefore still survives in such places as the forests of the



GLACIER BEAR.

Drawn from a specimen in the United States National Museum.

**Bear**, and in the Rocky Mountain regions and Alaska, brown specimens are almost as numerous as black. Sometimes it is difficult to believe that both kinds belong to the same species, but this seems to be a fact.

Some grizzlies are very dark brown, but they are never inky black, like the true Black Bear. The latter differs in form from the grizzly in being highest in the middle of the back, very

Adirondacks, the Catskills, in West Virginia, and the swamps of the southern states.

When properly treated, small Black Bears are good-tempered and playful in captivity; and some are easily tamed, and taught to perform tricks. Cubs are very interesting when small, but by the time they are a year old, they become so strong and troublesome, as well as dangerous, that private owners nearly always are heartily

glad to get rid of them. Never buy a Black Bear cub in the belief that it can be kept for amusement and resold at a profit; but if thine enemy offend thee, present him with a Black Bear cub.

The Black Bear is a timid animal, and always runs when observed by man. It is a good climber, runs quite swiftly when pursued, but in a rough and tumble fight it bawls, roars, and coughs.

**The Glacier Bear,**<sup>1</sup> found on the glaciers around Yakutat Bay, near Mt. St. Elias, Alaska, is one of the recent discoveries in the Northwest, but it is so clearly distinct as to merit special notice. Thus far no living specimens have found their way into zoological parks or gardens, and the only mounted skin on exhibition is in the United States National Museum. It is exactly reproduced in the accompanying illustration.

The species is known to-day only by the single specimen referred to, and a few flat skins. As mounted it is only 24 inches in height at the shoulders, and is beyond question the smallest species of bear in America. Its color is a peculiar bluish gray, on all parts save the muzzle from the eyes forward, which is dark brown or black. The hair is long, very thick, woolly in texture, and stands out straight all over the body. The rarity of this animal in collections, and the long delay in its discovery, are due to the rough, inhospitable and dangerous character of the country in which it lives.

**THE RACCOON FAMILY.**

*Procyonidae.*

**The Raccoon,**<sup>2</sup> placed next to the bears, is also plantigrade in its manner of walking. It is a cheerfully persistent animal, and no amount of hunting discourages it, or drives it away from its favorite haunts. It is at home in the timbered regions of the southern and eastern United States, especially where there are swamps,—for the Raccoon loves to play in water. In the West it ranges from Arizona to British Columbia.

Its favorite dwelling-place is a hollow tree, and its yearly family consists of five or six young. In its appetite, it is as omnivorous as any bear, and eats everything that it can chew,—from live rabbits down to green corn,—fish, flesh, or fowl. The only point on which the Raccoon is particu-

<sup>1</sup> *Ursus em'mous-i.*

<sup>2</sup> *Procyon lotor* and related species.

lar, regarding its food, is in soaking it in water before eating it.

Excepting the cacomistle or "civet cat" of the Southwest, this is the only animal in the



Photo. and copyright, 1902, by W. L. UNDERWOOD.

THE RACCOON.

United States which has black and gray rings around its tail. A live "Coon" makes one of the most satisfactory carnivorous pets that a boy can keep in confinement.

**The Cacomistle, "Civet Cat," or Bassarisk**<sup>1</sup> is a strange little creature like a small pine marten with a long, bushy tail, and many common names. It is spread over so wide an area of our country that its personality should be better known. It inhabits Mexico and the southwestern United States from Texas to California and north to southern Oregon. These are the names by which it is called and miscalled: in Mexico, *Cacomiztli*, or in English *Ca-co-mis'tle*; in Texas, *Texas Civet Cat*, and *Cat Squirrel*; in California, *Mountain Cat* and *Ring-Tailed Cat*; in Arkansas, *Raccoon Fox*; by various scientific authors from Audubon to Allen, *Civet Cat*, *Ring-Tailed Bassaris*, and *Northern Civet Cat*.

Now, as to the facts regarding this pretty little creature, it is not a "cat" of any kind, and there is about it not a trace of "civet." Dr. Coues proposed *Bassarisk* as a name that

<sup>1</sup> *Bas-sa-ris'cus as-tu'tus* and related species.

was appropriate, and entitled to use. Let it be so called henceforth, and the misnomers relegated to obscurity, where they belong. Its original Mexican name is so ill adapted to our wants it never will be generally used.

The Bassarisk is, after the true raccoon, the only animal in the United States possessed of a long, bushy tail with alternating black and white rings around it. It climbs trees, and nests in hollow branches like a squirrel; it scratches

and bites, and catches rats, mice and small birds like a cat; and it has a many-sided appetite, like a raccoon. Its length of head and body is 16 inches, tail about the same, and its general color is a brownish gray. It is a night prowler, and often makes its home in outbuildings and deserted ranch houses. In California it is occasionally kept in captivity by the miners, and is said to make a very attractive and interesting pet.

## CHAPTER IV

# THE ORDER OF SEALS AND SEA-LIONS

### PINNIPEDIA

Some students may feel that it is useless for land dwellers to try to become acquainted, at long range, with sea animals. Toward many sea animals, this feeling is justified; but it should not be entertained toward the bold and hardy fin-footed children of the surf. The seals and sea-lions of our shores are well worth knowing.

From the warm and luxurious shore of southern California to Oregon's storm-beaten Tillamook Rock, and on up to the inhospitable, rock-bound edge of western Alaska, you will find at intervals, where Nature has done some of her grandest work in shore-building, colonies of bold and hardy sea-lions. On the Pribilof Islands lives the most valuable of all the fur-bearing animals of the world, the fur-"seal," which has contributed millions of dollars to our national treasury, and more than repaid the whole price of Alaska.

On the low shores and adjacent ice floes of the North Atlantic live the seal herds that annually yield an immense store of valuable oil, and furnish employment for thousands of Newfoundland sailors and sealers.

The reader may rest assured that even though his home be in the centre of the Great Plains, the North American seals and sea-lions are well worth knowing; for, sooner or later, travel surely will bring him into visual contact with many of them, either in museums, zoological gardens, or alive in their haunts. Let us, then, lay the foundation for a profitable acquaintance with them.

By some writers, these animals are classed with the Ferae, because they eat flesh; but to associate in the same Order such widely different

creatures as sea-lions and cats seems incongruous, if not incorrect.

The Order Pinnipedia<sup>1</sup> contains three groups of sea-faring animals, distributed widely through the ocean waters of the world, and in some instances, in fresh water, also. They are the Sea-Lions, Seals and Walruses.


A **Sea-Lion** has a long, supple neck, and long, triangular front flippers that have neither hair nor claws, but are simply living paddles. Their hind limbs are web-toed flippers. They have very small, sharp-pointed ears, carry their heads high, and all are lively, active animals, both in swimming and in climbing rocks. The males of some species grow to enormous size, and have faces so lion-like in appearance that this resemblance has given the group its popular name, —Sea-Lion.

A **Seal** is a short-necked, fat-bodied, low-lying, clumsy animal, not nearly so active on land nor so intelligent as a Sea-Lion. Its front flippers are short, square-ended, fully covered with hair, and provided with claws. They have no external ears of skin and cartilage. Their hair is short, close, and stiff, and of no value as fur save to the Eskimo, to whom every Seal is a Godsend, and utilized in a great variety of ways.

A **Walrus** is a sea mammal of great size, formed somewhat like a Sea-Lion, and it is the clumsiest living creature that ever comes upon land. It has two long ivory tusks that grow downward from the upper jaw, a very thick skin which lies in deep folds, no hair worth mentioning, and a very dull brain.

The following are the groups and species which every American should know :

<sup>1</sup> *Pin-ni-pe'di-a* means "fin-footed."



PACIFIC WALRUS,



STELLER'S SEA-LION,



HARBOR SEAL.

On the same scale.

FAMILIES.

EXAMPLES DESCRIBED.<sup>1</sup>

<b>ORDER PINNI- PEDIA.</b>	}	SEA-LIONS, <i>O-TA-RI'-DAE</i> , . . .	{	California Sea-Lion, . . .	<i>Zalophus californianus</i> .
				Steller's Sea-Lion, . . .	<i>Eumetopias stelleri</i> .
				Fur "Seal," . . .	<i>Callotaria ursina</i> .
		SEALS, . . . <i>PHO'CI-DAE</i> , . . .	{	Ringed Seal, . . .	<i>Phoca foetida</i> .
				Harbor Seal, . . .	<i>Phoca vitulina</i> .
				Harp Seal, . . .	<i>Phoca groenlandica</i> .
				Hooded Seal, . . .	<i>Cystophora cristata</i> .
				Ribbon Seal, . . .	<i>Histiophoca fasciata</i> .
		WALRUSES, <i>OD-O-BEN'I-DAE</i> , . . .	{	Pacific Walrus, . . .	<i>Odobenus obesus</i> .
				Atlantic Walrus, . . .	<i>Odobenus rosmarus</i> .

**THE SEA-LION FAMILY.**

The California Sea-Lion,<sup>2</sup> or Barking Sea-Lion, is the most familiar representative of the first group, for the reason that this species is easiest to catch alive, and keep in captivity. In zoological gardens and travelling shows, this is the animal which cries out so frequently, and with ear-piercing clearness and volume, "How-woo! Hook! Hooook! Hook!" It inhabits nearly the entire coast of California, the Farallone Islands, the famous Cliff House rocks, and the Lower California peninsula. Full-grown males are about 7 feet in length, weigh about 450 pounds, and all are of a uniform dark-brown color. An adult female which died in the Zoological Park weighed 112 pounds and measured length of head and body, 56½ inches, tail, 2¾ inches, total length from nose to end of hind flippers, 70¼ inches, girth, 31½ inches. These creatures are very active in the water, and can climb rocks,



KELLER, Photo., N. Y. Zoological Park.

CALIFORNIA SEA-LION.

<sup>1</sup> The most important of these species will be found well described and commented upon in Mr. Henry W. Elliott's interesting volume entitled "Our Arctic Province." (Charles Scribner's Sons.)

<sup>2</sup> *Zal'ophus cal-i-for-ni-an-us*.

and even high cliffs, with surprising agility. When frightened, Captain Scammon says they will leap from a height of sixty feet into the sea.

The hair of this animal is very short, coarse, and of no value. The California Sea-Lions rarely eat fish, but live chiefly upon squids, shell-fish



and crabs. For reasons known only to themselves, they swallow many round pebbles, from one to two inches in diameter. We once took 16 pounds (half a pailful) from the stomach of a medium-sized specimen.

In captivity all kinds of seals and Sea-Lions live contentedly in fresh water. The value of a living California Sea-Lion in New York City is about \$150. This species possesses great intelligence, and quite recently several specimens have been trained to go through a show performance which is really wonderful, including a most remarkable act in which a Sea-Lion successfully balances a large ball on the point of its nose.

An important incident in the life history of the California Sea-Lion furnishes a good illustration of the folly of condemning a wild species to destruction on insufficient evidence.

For several years the fishermen of San Francisco complained that the Sea-Lions of the California coast devoured such enormous quantities of salmon and other fish that they were seriously affecting the available supply; besides which, they caused great damage to nets and impounded fish. They demanded that the Sea-Lions be destroyed, and finally convinced the state authorities that their contentions were well founded.

It was decided that the animals should be destroyed, by systematic shooting, down to a comparatively small number; and the slaughter was duly ordered. Men were engaged to do the work, in a business-like way, and an official request for permission to kill on the light-house reservations of the government was granted.

But there were certain naturalists who doubted the entire accuracy of the charges made against the Sea-Lions, and asked for proof in detail. When no evidence of a specific and convincing nature was brought forward, they requested that the slaughter proposed on the Farallone Islands, and other light-house reservations, be deferred, pending a careful inquiry; and this was done.

However, where the state authorities had full power to act, the killing proceeded in a few localities. It happened that during the killing of California Sea-Lions on the shore of Monterey Bay, and vicinity, Professor L. L. Dyche, of the University of Kansas, arrived on the scene to pursue studies in marine life. He examined the

stomachs of twenty Sea-Lions which were washed ashore, and of five more which he killed for the purpose of mounting their skins.

*Every stomach examined contained the remains of squids and devil-fish (Octopus), one or both; and both of which are among the fisherman's enemies! Not one of the twenty-five stomachs which he carefully examined and reported upon contained any portion of a scaled fish.*

In 1901, the United States Fish Commission conducted a systematic investigation of the food habits of the Sea-Lions of the Pacific coast and the report of Messrs. Rutter, Snodgrass and Starks appears in the Report of the Fish Commissioner for 1902. At six points on the coast of California, the investigators killed a total of twenty-four specimens of the California Sea-Lion, and eighteen of the Steller Sea-Lions. The report says:

"Of thirteen California Sea-Lions whose stomachs contained food, five had eaten fish and eleven had eaten squid. The quantity of fish was inconsiderable, seventeen small fishes being the maximum, while the remains of one hundred to three hundred squid were found in each of five stomachs.

"All the thirteen Steller Sea-Lions whose stomachs contained food had eaten fish, and five had eaten squid, or octopus. The number of squid eaten was small, six being the maximum number in one Sea-Lion, while the quantity of fish was large, at least thirty-five pounds being taken from one stomach."

The detailed report of the kinds of fishes consumed as food by these animals reveals an assortment of very little value, and not one salmon or shad. Professor Dyche's discovery—that the California Sea-Lion feeds almost exclusively upon squid—was fully confirmed, for the twenty-four animals killed contained only three rock-fish, two hake, twenty-four "small fish" and one chimera,—but over eleven hundred squid! The stomachs of the Steller's Sea-Lions contained fourteen rock-fish, two perch, thirty clupeoid fish, seventeen "large fishes of 12 to 18 inches," and a few skates, sharks and squids.

"The testimony of the fishermen was so contradictory it is of no value. . . . One man claims that the Sea-Lions are becoming more numerous and destructive every year, while another claims that they are rapidly becoming

exterminated." There was "practically no complaint" of fish destruction "at the time of the investigation. Sea-Lions were scarcely ever seen in the vicinity of the salmon nets during 1901."

At the mouth of the Columbia River, "the fishermen were unanimous in their denunciation of the Sea-Lions." "The shallow water and the large number of salmon make that point a favorite feeding ground, and there is no doubt that the Sea-Lions are doing much damage there." "It

the strength of general opinions; for a supposed enemy may, on careful investigation, prove to be a friend.

**Steller's Sea-Lion,**<sup>1</sup> the largest Sea-Lion in the world, inhabits a few isolated spots on the Pacific coast, from Santa Cruz, California to Berling Strait. Large male specimens attain an average length of 10 to 11 feet, stand 6 feet high, and attain a weight estimated by competent observers at 1,400 pounds. The full-grown male



STELLER'S SEA-LION.

appears that the Sea-Lions are doing very little damage anywhere excepting at the mouth of the Columbia River." (Report, page 117.)

A summary of the results of the investigation establishes three facts:

1. The California Sea-Lion is not guilty of destroying fish to any great extent, and deserves protection, not death.

2. Steller's Sea-Lion eats miscellaneous fish; but on the coast of California does nothing to merit destruction. At the mouth of the Columbia it is destructive, and there deserves to be kept in check.

3. Wild animals never should be destroyed on

has a girth of 8 to 9 feet, a lion-like head, coarse neck hair 4 inches long, and canine teeth like a grizzly bear, which are much used in fighting.

The full-grown females are from 8 to 9 feet long, weigh from 400 to 500 pounds and are more finely formed. The hair is coarse, and the animal is now of practically no commercial value, save for its oil. This species is readily distinguished from the California sea-lion by its far greater size, its hoarse voice, the very large neck, and the long, coarse neck hair of the males.

In its habits, this great Sea Lion is very peculiar. Amongst themselves the old males

<sup>1</sup> *Eu-me-to'pi-as stel'ler-i.*

fight fiercely, and with their big canine teeth inflict upon each other many severe skin wounds. I have seen specimens whose necks bore scores of large scars. In the presence of man, however, they are timid, and easily frightened.

This giant among Sea-Lions is found on the coast of California, in small numbers only, at Point Ano Neuvo, near Santa Cruz, at Purissima, the Farallone Islands, Point Reyes, and Point Arena. On the coast of Oregon it is found about the mouth of the Columbia and Tillamook Head. The agents of the United States Fish Commission, reporting observations made in 1901, stated that "probably half of the Sea-Lions of California (of both species) are found at the Farallone Islands, and it seems doubtful whether the total number on the coast amounts to five thousand." A large colony of Steller's Sea-Lions inhabits Bogoslof Island, Alaska, living almost in the shadow of that celebrated volcano.

In October, 1903, the New York Zoological Society's agents succeeded, after many fruitless efforts, in capturing six young specimens in the sea off San Miguel Island, California, and safely transporting them to New York, where the experiment of keeping this species in captivity is now being tried in the Zoological Park.

**The Fur Seal,**<sup>1</sup> which yields the beautiful and costly fur so highly prized for ladies' garments, is not a true seal, but a sea-bear or sea-lion, quite similar in form, size and general habits to the California species already described. It is found on the Pribilof or Seal Islands, in Bering Sea, where during the Russian occupation it was twice nearly exterminated for its fur; on Copper and Robben Islands, off the coast of Siberia; and in the open sea from the Pribilof Islands southeastward to the thirty-fifth parallel, thence northward along the coast, back to the Seal Islands.

The size of the Fur Seal has been carefully observed by Mr. Henry W. Elliott, and recorded as follows:

MALES	{	At birth (June 20)	Length 12 to 14 in.	Girth 10 in.	Weight 6 to 7½ lbs.
AND		At six months,	" 24 "	" 25 "	" 39 "
FEMALES.	{	At one year,	" 38 "	" 25 "	" 39 "
		At two years,	" 45 "	" 30 "	" 58 "
MALES	{	At three years,	" 52 "	" 36 "	" 87 "
ONLY.		At six years,	" 72 "	" 64 "	" 280 "
	{	At 8 to 20 years,	" 75 to 80 "	" 70 to 75 in.	" 400 to 500 lbs.

<sup>1</sup> *Cal-lo-ta'ri-a ur-si'na.*

The Fur Seal has two kinds of hair. Its outer coat is long, stiff, coarse, and gray in color. In preparing skins for market, all this is plucked out and thrown away, leaving only the fine, soft, brown under fur, which before manufacture is dyed a rich, blackish-brown color. Fur Seal garments vary in price from \$200 to \$700.

The Fur Seal has strange and interesting habits. It spends about two-thirds of each year far at sea, making a circuit of 6,000 miles in the open ocean without touching land. For some strange reason, the herd in American waters has chosen the two Pribilof Islands, St. Paul and St. George, as the only spots in our waters whereon they are willing to land and rear their young. To these favorite breeding-places, on these islands known as "hauling-grounds," the Fur Seal millions were wont to repair in the early summer of each year, to rear their young. The returning herd begins to arrive between May 1 and 15, the breeding season is over by September 15, and by the end of November all the Seals are gone on their great winter cruise southward into warmer waters. By a long series of inquiries the winter cruise of the herd has been mapped out by Dr. D. S. Jordan and his associates, and is shown on the next page.

On the breeding grounds, each large and hard-fighting old male gathers round him a harem of from six to ten females, fights off all rivals, young or old, and elects himself the head of an imposing family. The three-year-old male Seals—called "bachelors"—were killed for their fur, to the number of about 100,000 each year. The females bear only one "pup" annually, immediately after landing in May.

The mother Seals leave their young, go to sea in search of food, remain several days perhaps, or even a fortnight, then return and go straight to their own respective offspring. It was the killing of the mothers at sea that produced an enormous falling-off in the number available each year. The persistent slaughter of mothers


will exterminate any species of animal, no matter how numerous.

The accompanying map graphically illustrates the remarkable sea-going habits of the Pribilof Fur Seal herd after the close of the breeding season, and during the intensely cold and fear-

Commercial Company, for twenty years, the exclusive right to kill each year on the Pribilof Islands, 100,000 young male Fur Seals, receiving therefor, annually, the sum of \$317,500.

**1872.**—The Alaska Commercial Company began to expend \$100,000 in cash, chiefly in London, in making the wearing of sealskin fashionable. This effort was entirely successful.

**1873.**—After a careful survey of the Pribilof Islands, and an elaborate computation of the number of Fur Seals then inhabiting them, Mr. Henry W. Elliott, a special agent of the Treasury Department, announced the total number of Seals to be 3,193,420. He says: "No language can express adequately your sensations when you first stroll over the outskirts of any one of those great breeding grounds of the Fur Seal on St. Paul's Island. . . . Indeed, while I pause to think of this subject, I am fairly rendered dumb by the vivid spectacle which rises promptly to my view. It is a vast camp of parading



ANNUAL WINTER MIGRATION OF THE FUR SEAL HERD.

fully windy winters that annually render life on the Seal islands a serious task.

The combined political and commercial importance of the Fur Seal demands a brief summary of the most important facts of its rise to favor, its decline, and finally its fall. The end, however, is not yet; but it looms very near.

**REVIEW OF FUR SEAL HISTORY.**

For the past seventeen years, the Fur Seal has been to the United States, England and Canada a source of well-nigh constant anxiety, contention, and at times irritation. Inasmuch as the fate of that animal is still pending, it seems desirable to set forth the most important facts in its case, in chronological order. The history of the Fur Seal since our acquisition of Alaska is divided into two periods, one of revenue, and one of contention.

**The Period of Revenue.**

**1867.**—When Alaska became a United States possession, by purchase from Russia at a cost of \$7,200,000, the fur of the Fur Seal was almost unknown to fashion, and outside of Russia was neither used nor particularly desired.

**1870.**—The United States leased to the Alaska

squadrons which file and deploy over slopes from the summit of a lofty hill a mile down to where it ends on the south shore. Upon that area before my eyes, this day and date of which I have spoken, were the forms of not less than three-fourths of a million seals, moving in one solid mass from sleep to frolicsome gambols, backward, forward, over, around . . . until the whole mind is so confused and charmed by the vastness of mighty hosts that it refuses to analyze any further." ("Our Arctic Province," p. 313.)

Some observers estimated the number of Seals at a figure higher than Mr. Elliott's. Others have recently contended that it must have been less.

**1880.**—"Pelagic sealing" means the killing of Fur Seals, male or female, in the open sea, by means of guns or spears. It is an exceedingly wasteful and destructive method, but it had been going on in a quiet way for many years. On land, only male Seals are killed. In the sea, about four females were killed to every male taken, and the pups on shore were left to starve. In 1880, the total number of Seals taken at sea in Bering Sea was only 8,418; but from that time on, the killing increased rapidly, and became fearfully destructive.



Drawn by J. Carrer Beard.

FUR SEALS ON THEIR "HAULING GROUNDS," PRIBILOF ISLANDS.

The large central figure is that of an old male, surrounded by the members of his "harem." In the foreground are two young "bachelor" seals, such as were regularly killed for their fur.

**1882.**—Up to this time, the great Seal herd of Bering Sea was in a state of equilibrium, and yielded on the islands its annual quota of 100,000 "bachelor" Seals without sensible variation. The number killed at sea in 1882 was 15,551.

#### The Period of Contention.

**1886.**—The catch of Seals at sea rose to 28,494. Of the large fleet of vessels then hunting Seals in Bering Sea, a number were seized by the United States government vessels which were guarding the Islands. These were chiefly Canadian schooners, but some were American.

**1887.**—The pelagic sealing fleet was increasing each year. The United States began negotiations with six foreign governments with a view to securing co-operation in saving the Seals from the extermination which threatened them at the hands of the "poachers."

**1890.**—The lease of the Alaska Commercial Company terminated, and the North American Commercial Company bid successfully for the new lease of the Seal-taking privilege on the Pribilof Islands. According to the calculations of Mr. Elliott, the Seals on the Islands now numbered 959,455. Except four years, from 1871 to 1889, over 100,000 male Seals had been taken annually, on the Islands, and paid for. The total revenue derived by our government during that twenty-year period was \$6,350,000. In 1890, the Seals killed and secured at sea numbered 40,814, while the number killed and lost was unknown.

**1891.**—An agreement called a *modus vivendi* (or way of living in peace) was made between England and the United States, for three years, designed to close Bering Sea to pelagic sealing pending the result of the Paris Tribunal. Practically, it amounted to nothing.

**1893.**—The case of the pelagic sealers was tried before the Paris Tribunal, and through the ineffective management of our case, we lost on practically all our contentions. The pelagic sealers emerged from the contest with full license to kill Seals at sea everywhere outside a sixty-mile radius of the Pribilof Islands. Because Japan, China and Russia were not parties to the Tribunal, the people of those nations were not bound by the award which keeps American, Canadian and English sealing vessels sixty miles away from the Seal islands!

**1894.**—In this year 61,838 Seals were killed at sea and secured, while an unknown number were killed and lost.

**1895.**—Mr. J. B. Crowley (Member of Congress in 1903), as a special agent of the Treasury Department, assisted in counting the dead bodies of about 30,000 Fur Seal "pups," on the Seal islands, which had starved that year by reason of the killing of their mothers while at sea in search of fish. (*Congressional Record.*) There were 56,291 Seals killed at sea, by the eighty-one vessels engaged in pelagic sealing. On land the number killed was, by order of the government, reduced to 14,846.

From 1890 to the end of 1895 (six years) the cost to the United States Government of its efforts to patrol the waters of Bering Sea, with war vessels and revenue cutters, and protect—as far as possible—the Seal herd from complete annihilation, was \$1,410,721. Besides this, the government expended \$227,163 on its Treasury Agents, and \$473,000 was paid by the decision of the Paris Tribunal, as "damages," to the men who stole our Seals, and were caught in the act!

**1897.**—The number of dead pups counted on the breeding grounds, by Mr. Frederic A. Lucas and others, was 21,750, and in October the number of seals remaining alive of our herd was estimated at 343,746. (D. S. Jordan. "Report Fur Seal Investigation," 1896-97, p. 100.)

**1898.**—By a law passed December 29, 1897, all citizens of the United States were absolutely prohibited from killing or capturing Fur Seals at sea anywhere in Bering Sea, the Sea of Okhotsk, or anywhere north of the 35th parallel of north latitude. The ownership of any Fur Seal skins taken in those waters was also prohibited, under severe penalties. All skins from female Seals, either raw or dressed, were also excluded from our markets.


From that date (December 29, 1897), pelagic sealing ceased to be an *American* industry. It is now for England and Japan to say whether or not it shall continue until all the mothers are slaughtered, and all the pups starved to death.

**1903.**—The situation of the Fur Seal has grown desperate, and its fate is wavering in the balance. The number now alive is about 200,000. While Americans cannot now engage in pelagic sealing under our flag, and no Canadians may inside the sixty-mile limit, dozens of well-equipped sealing

vessels are sent out from Yokohama, and other ports in Japan, under the Japanese flag, *which hunt seals within three miles of the Pribilof Islands!* Canadian Sealers still hunt outside the protected zone, and kill many seals, annually.

Up to this date, our government has done everything in its power to prevent the extermination of the Fur Seal, and afford it a just measure of protection. England fears that she can go no farther without giving grave offence to

of him who can take it. Patriotism, and the desire for the greatest good of the greatest number, does not enter into their calculations. The American or Canadian pelagic sealer claims that the open sea is his, and he cares only for the \$10 or \$15 that each raw skin is worth. England cannot reasonably be expected to quarrel with Canada because of our desire to perpetuate our Seal herd, and derive from it a revenue of a million dollars a year,—which is the sum which the



Drawn by J. CARTER BEARD.

THE HARP SEAL.

Young and old specimens, showing changes in pelage at different periods.

Canada. But in England, about \$2,000,000 of capital are invested in the business of dyeing and dressing Fur Seal skins, and this work employs—or *did* employ—between two thousand and three thousand operatives. It has always been impossible for Seal skins to be satisfactorily dyed and dressed in America.

The insurmountable obstacle to the protection of the Fur Seal is its fatal habit of going to sea, far from its hauling grounds, coupled with the belief of a large number of Canadians and Americans that a Seal at sea is the lawful prize

Fur Seals would yield to-day, but for the slaughter of 1,000,000 females at sea, and the murder or starvation of 1,000,000 pups, at sea and on shore.

Just what events will make up the next and possibly the final chapter in the life history of this interesting and valuable species, it is at present impossible to foretell. Judging by the past, and the indications of the present, the Alaskan Fur Seal is doomed to practical annihilation, but not total extinction. Let us hope, however, that even yet the statesmen of the United States, England, Canada and Japan will join in the

enactment and enforcement of a humane measure of protection which really will protect.

### THE SEAL FAMILY.

*Phocidae.*

The **Little Ringed Seal**<sup>1</sup> is the Seal of the Farthest North, and the friend of the northern Eskimo all round the pole. It is the smallest North American species, and looks very much

bear, with two small cubs, was closely following up the seals as they worked north through the ice pack.

The **Common Harbor Seal**,<sup>1</sup> of both our ocean coasts, is a good representative of the Seal Family, chiefly because it is the species most frequently seen. It ascends rivers far above tidal influence, and has been taken in Lake Champlain. In the Columbia River a closely related species



THE RIBBON SEAL.

like the common harbor seal. It goes as far north as it can find breathing-holes. Nansen found it on May 31, at 82° 21', or within 460 miles of the pole, living in the narrow lanes of water that were then forming in the great polar ice pack. It was a **Bearded Seal**,<sup>2</sup> however, which, on June 22, afforded the brave explorers a good supply of food when men and dogs were almost starved. And, true to its nature, an old polar

has been taken above The Dalles, 200 miles from the sea.

The **Harp Seal**<sup>2</sup> is not only one of the handsomest of all Seals, but it is also the species most valuable to man. It is found on both sides of North America, but always in cold waters. In the year 1900, five sealing steamers of Newfoundland took nearly 100,000 seals, mostly Harps, on the coast of Labrador and northward

<sup>1</sup> *Pho'ca fo'e'ti-da.*

<sup>2</sup> *Er-i-gnath'us bar-ba'tus.*

<sup>1</sup> *Pho'ca vit-u-li'na.*

<sup>2</sup> *Pho'ca green-land'ic-a.*



thereof, and the value of the catch was over a quarter of a million dollars.

This species passes through several strongly marked changes of pelage and color. The baby is covered from nose to flipper-tips with a thick coat of long, woolly hair of snowy whiteness. This, when shed at six months after birth, is replaced by a coat of bluish gray hair, with light trimmings. On reaching adult age, in its fifth year, this animal is very strikingly marked by black or dark-brown patches grouped together on the sides and back, on a white or yellowish ground-color apparently in the shape of a harp. This Seal is also called the **Saddle-Back**, and **Greenland Seal**.

**The Hooded Seal**<sup>1</sup> of the North Atlantic is a large species, often attaining 8 feet in length. The old males are distinguished by the possession of a flexible bag of skin on top of the nose, which is capable of being inflated with air until it forms a lofty and remarkable excrescence on the creature's face. This sac is sometimes 10 inches long and 6 inches high. The color of this Seal is dark bluish-gray, marked with irregular light spots. It once came as far south as New Jersey.

**The Ribbon Seal**, or **Harlequin Seal**,<sup>2</sup> in its color pattern is the most remarkable of all living Pinnipeds, and there are many persons who consider it the most beautiful member of its Order. On a smooth ground-color, either of blackish-brown or yellowish-gray, Nature has sportively arranged several yards of broad, yellowish-white ribbon. One strip goes around the neck, and ties under the throat. From a point low down on the breast, another starts upward, curves gracefully over the shoulder, drops down in front of the pelvis, where it comes together, then turns and crosses over the body. In many specimens the uniformity of the width of the ribbon is remarkably well maintained.

This Seal is from 4 to 6 feet in length. Its home is on the eastern shore of Bering Sea, and in the fresh waters of Lake Iliamna, in the upper end of the Alaskan Peninsula.

## THE WALRUS FAMILY.

### *Odobenidae.*

Of all living monsters that ever move upon land, the **Pacific Walrus**<sup>3</sup> is one of the most

<sup>1</sup> *Cystophora cristata*.

<sup>2</sup> *His-tri-o-pho'ca jas-ci-a'ta.* <sup>3</sup> *O-do-ben'us o-be'sus.*

wonderful. A full-grown male is a living mountain of heaving flesh, wrinkled, furrowed and seamed, ugly as a satyr, and as strange in habits as in appearance.

Its form is that of a sea-lion with a neck enormously thickened. Its upper jaw is provided with two long, strong tusks of ivory, and its skin is almost destitute of hair. A full-grown male measures from 10 to 12 feet in length from nose to tail, the top of its head is about 5 feet from



HEAD OF HOODED SEAL.

the ground, the girth of its neck is from 12 to 14 feet, and it weighs from 1,800 to 2,000 pounds. Its skin varies from half an inch to two inches in thickness; it is of a dirty yellow color, and lies on a mass of fat which often is six inches thick. The largest pair of tusks known to the author measure 24½ inches in exposed length, and are in the British Museum.

The Pacific Walrus eats more or less of aquatic-plant food, but its principal food is shell-fish and crustaceans. These it digs up from the muddy bottoms of the broad, shallow bays along the coast, crushes between its powerful jaws, and swallows in great quantities, shells and all! Crabs and shrimps form a pleasing variety, and for salad it devours the bulbous roots and tender stalks of marine plants which in summer grow in its home waters.

In former times, the Pacific Walrus existed in great herds on the coast of Alaska, from the north shore of the Alaskan Peninsula northward through Bering Strait, and thence eastward as far as Point Barrow. There the herds encountered the edge of the great permanent ice-pack, and could go no farther. In winter the Walrus

herds float about on the ice-fields, retreating southward as the edge of the ice advances. In the open sea, the sleeping posture of the Walrus is floating bolt upright in the water. He grunts and bellows, and many times vessels have been warned off dangerous, fog-hidden rocks by the Walrus lying upon them.

On land the Walrus is the most clumsy and

In 1900, steamers bearing gold-miners to Cape Nome passed through herds of Walrus in Bering Sea, and many of the animals were killed, wastefully and wantonly, by passengers firing from the decks, with no possibility of securing a single victim. As elsewhere, the instinct of man in the far north is to slay and slay, and preserve no living thing.



THE PACIFIC WALRUS.

An old male of the largest size. Drawn from a mounted specimen in the United States National Museum.

helpless of all land animals, and is easily approached and killed. In the water, it becomes a danger to be avoided, on account of its proneness to wreck small boats. A full-grown Walrus has never been seen in captivity. Two or three very young specimens have reached Europe, and in September, 1902, Commander Robert E. Peary brought one to New York for the Zoological Park, where it was exhibited until it died.

The Walrus has been hunted so diligently for its oil that to-day very few remain, and the natives who once depended solely upon this animal for food, fuel, lights, boats, dog harness, and leather for all purposes now are on the verge of starvation, and are really kept alive by public bounty. Previous to our purchase of Alaska, about 10,000 Walrus were killed annually by the Eskimo, and utilized. In the long, hard winter

of 1879-80, when the sea was frozen all around St. Lawrence Island, for many miles in every direction, the Walrus herds were forced to re-

length as the Pacific species, but it has a shorter and much smaller neck. Its tusks, also, are much smaller. It is still found in considerable numbers in Smith's Sound, and is quite abundant north of Franz Joseph Land, where Nansen photographed and killed many. Its most northerly latitude is 82°. A specimen killed by Commander Robert E. Peary was 9 feet in length, and weighed 1,569 pounds. The skin alone weighed 220 pounds.

Professor L. L. Dyche has kindly furnished the measurements of the largest male Walrus out of eighteen taken by him on the coast of northern Greenland:

Length (straight line), end of nose to end of body, 129 inches.

Tail, exposed, none.

Length of rear flippers, 26 inches.

Girth of animal when suspended by the neck, 129 inches.

Exposed length of tusk, 19 inches.

Circumference of tusk at base,  $8\frac{1}{4}$  inches.

The largest cow Walrus measured 116 inches in length, 113 in girth, exposed tusk,  $10\frac{3}{4}$  inches.



YOUNG ATLANTIC WALRUS.

Captured by Commander R. E. PEARY, and exhibited in the New York Zoological Park.

main so far away that all the inhabitants of the Island, save one small settlement, died of starvation.

The Atlantic Walrus<sup>1</sup> is of about the same

<sup>1</sup> *O-do-ben'us ros-ma'rus*.

## CHAPTER V

# THE ORDER OF MOLES AND SHREWS

### INSECTIVORA

In the dark and cold embrace of Mother Earth, away from the cheering sunlight, and the beautiful upper world that we enjoy, there dwells a group of mammals so strange, and yet so useful to man, that they excite our admiration for the wise purpose which developed and placed them there. Pass not unthinkingly the moles and shrews, for they have been most cunningly designed to serve a definite and important purpose in the economy of Nature.

In farming countries, the top soil of the earth is a vast incubator for the development of destructive insect larvae. In soil that is rich and productive, "grub-worms," "cut-worms," and "wire-worms" abound; and in regular rotation they greedily devour the seeds, roots and leaves of growing crops. But for the enemies which keep them in check, there would be a hungry grub for every sprouting seed.

And how can man wage war successfully against insect life in the soil? Impossible. To meet this difficult proposition, we need a vigorous living creature with a nose like a gimlet, sharp-pointed teeth, soft fur, feet specially designed for digging, and eyes so small that to them sunlight is an unnecessary luxury. Such animals are found in the moles and shrews, of the Order In-sec-tiv'o-ra, humble but faithful workers in man's interest. Neither the horse nor the ox is more diligent in our service than are these toilers of the soil. Yet what is their reward?

In his mole-like blindness, man frequently discovers things that are not true. Often a perfectly honest farmer concludes that a mole is eating his seed corn in the ground, or the vegetables in the garden; and straightway the mole is killed. His accuser has found a runway following up a row of newly-planted corn, and when the seed fails to sprout, the mole is accused of having eaten it!

In all such cases, the mole is a victim of circumstantial evidence, and suffers through the

lack of counsel to cross-examine the witnesses for the prosecution. Did anyone ever find much vegetable food in a mole's stomach? Not often. Did anyone ever see a mole eat vegetable food? Probably not. A mole placed in a box and supplied with vegetable food alone soon starves to death. Moles do not eat seed corn, or garden vegetables; but they do visit corn-hills to eat the grubs that come to devour the corn.

Every young naturalist must learn early what constitutes *direct evidence*. Far too long have the mole and shrew been convicted and slain on circumstantial evidence. Meadow mice sometimes attack seed corn by utilizing the runways that have been made by moles in reaching the corn-hills to secure the grubs that attack the seeds; and almost invariably the testimony is that the moles have done the damage. In France the value of the mole is recognized by law, and the killing of one is punishable by a fine of five francs.

The shrews and moles not only find their food underground, but live the entire cycle of their lives in subterranean darkness. Moles seek their food by digging tunnels in ground that is loose and dry, the roof being raised into a ridge which in smooth lawns is an annoying disfigurement. Gardeners are apt to forget that they always work where insect larvae are thickest, and the need for their help is most urgent. The tunnel-makers are driven from lawns by persistently trampling down their runways.

The Order Insectivora is represented in the United States by two Families, the members of which are easily recognized by the following well-marked characters:

**The Moles** have pointed heads; *extremely large spade-like front feet*, that always are held with the outer edge up; no neck; the front legs are exceedingly short; there is no external ear, and no external eye; the body is short, thick and clumsy, and the tail is hairless.

**The Shrews** have pointed heads, but *small*,



1. COMMON MOLE.

2. STAR-NOSED MOLE.

rat-like feet; there is a very small eye, an external ear, and a distinct neck. The body is rather slender, and as a whole, the animal looks much like a short-tailed mouse.

### THE MOLE FAMILY.

#### *Talpidae.*

This Family contains twelve full species, all quite interesting. Their skins and skulls have been studied closely, but our information regarding their habits is very meagre. As a rule, moles are larger than shrews. The largest of all is an Oregon species, which measures 7 inches in length of head and body, and tail  $1\frac{1}{2}$  inches,—an unusual size for a mole.

On all moles the fur is fine, thick, very soft and velvety, and faultlessly smooth and clean. All these creatures love sandy soil, which they can easily burrow.

**The Common Mole**<sup>1</sup> is known to the majority of country dwellers by its upheaved tunnels on the surface of the ground. In appearance the animal is a flattened, oblong ball of fine, soft, shimmering gray fur,  $6\frac{1}{4}$  inches long,


<sup>1</sup> *Sca'lops a-quat'i-cus.*

to which the naked, little pink-white tail—which looks like a small angle-worm—adds  $1\frac{3}{4}$  inches. Its nose projects half an inch beyond its mouth, and on the end it feels as hard as if it contained a bone. It terminates in a broad, flattened point, shaped quite like a rock-drill.

The fore foot is three-quarters of an inch wide, but less than an inch in length, including the claws, which measure half an inch. In your hand, a Mole is a wriggling, restless creature. Place it upon ground that is not packed hard, and in about one second it has found a suitable spot for an opening. Its nose sinks into the earth as if it were a brad-awl, with a combined pushing and boring motion, and in three seconds your Mole's head is no longer in sight.

Up comes the powerful right foot, sliding close along the side of his head, edgewise and palm outward, to the end of the nose. The living chisel cuts the earth vertically, and then, with a quick motion it pries the earth sidewise from its nose. Instantly the left foot does the same thing on the other side, while the brad-awl nose goes right on boring. In ten seconds, by the watch, the Mole's body has entirely disappeared, and in three minutes our Mole will tunnel a foot, unless interrupted.

When skinned for dissection, it is found that the eye is merely a small, dark speck under the skin, suitable only to distinguish light from darkness. The eye-ball is about the size of a pin-head. The arm and forearm is a big, hard bundle of tough muscles and powerful tendons, shaped like an Indian club, of enormous size in proportion to the creature's body.




DIGGING MUSCLES OF A MOLE.

The Mole is a wonderful example of energy and power. Desiring to observe their methods of working when undisturbed, I once placed one in a five-acre clover-field, at 11 o'clock A.M. During the first seven hours it had tunneled twenty-three feet, in a zig-zag line. During the next seventeen hours it dug thirty-five feet, and

during the next hour, ten feet more. The total work consisted of sixty-eight feet of main line, and thirty-six and a half feet of branches, making in all one hundred and four and a half feet.

An observing farmer-boy, named Lawrence



1. End of nose.



2. Left forefoot.

STAR-NOSED MOLE.

Miller, once gave me a clear and intelligent description of a Mole's burrow which he uncovered and observed closely. It was a dome-shaped hole, two feet below the surface of the ground, reached from above by a hole that ran down slanting into its top. The burrow was a foot wide at the bottom, where three small galleries ran off about six inches, in different directions. Near the top of the chamber was a sort of shelf, supporting a bed of soft material, and on this lay a Mole. The young are usually two in number.

Besides the Common Mole, of the Eastern United States generally, we have the *Prairie* or *Silver Mole* of the prairie regions of the Mississippi Valley; the *Hairy-Tailed Mole* of the Eastern United States, and the *Oregon Mole* of the Pacific slope. The *Star-Nosed Mole*, of the northeastern United States and Canada, is quickly recognized by the remarkable star-like appendage of eighteen ray-like points, with four more between them, on the end of its nose.

## THE SHREW FAMILY.

### *Soricidae.*

North of Mexico, this Family contains about thirty-five full species, distributed throughout nearly every portion of North America south of a line drawn from the mouth of the Mackenzie River to Labrador. With most cheerful indifference, they inhabit mountains, plains, swamp lands and sandy sea-coasts, hot countries and cold. Everywhere, however, their noses are long and sharp, their eyes and ears minute, and the colors of all species are very sober, ranging

from dull gray to brown, and ending in black. There are two species which are so widely distributed they may well be taken as types of the entire thirty-five.

**The Common Shrew**<sup>1</sup> is found on the Atlantic coast, from New England northwestward to Alaska, and southward through the Appalachian Mountains to Tennessee and North Carolina. Its color is brown above, and dull gray underneath; head and body,  $3\frac{1}{4}$  inches long, tail,  $1\frac{1}{2}$  inches. The ground plan of its skull is a perfect triangle spreading thirty-five degrees, and is very flat. Although very soft and fine, its fur is not so velvety as that of a mole. This creature is very small, and quite mouse-like in appearance.

Unlike the mole, Shrews occasionally emerge from their burrows, and wander about near their entrances. But they are exceedingly shy, and although they are frequently thrown out by the spade or plough, they are very rarely seen moving about. Above ground they are very helpless, and being unable to run rapidly, they try in a feeble way to hide. When taken in the hand, the musky odor they emit is rather disagreeable.

**The Short-Tailed Shrew**<sup>2</sup> is another type worthy of special mention. It is readily recognized by its very short tail, only 1 inch in length, while its head and body measure 4 inches. Its color is smoky brown above, and dull gray underneath, and in size it is the largest



1. COMMON SHREW. 2. SHORT-TAILED SHREW.

of the Shrews. It is found from the eastern edge of the Great Plains to the Atlantic coast, and is one of the largest members of the Shrew Family.

<sup>1</sup> *So'rex per-son-a'tus.*    <sup>2</sup> *Bla-ri'na bre-vi-cau'da.*

## CHAPTER VI

# THE ORDER OF BATS

### CHIROPTERA

The strange wing-handed, flying mammals composing this Order exhibit differences in form that are fairly bewildering. They range all the way from the beautiful to the fantastic and the hideous, and some of them are well worthy of study.

members of the Bat Order as a whole are almost as little known as the whales and porpoises of the deep sea. Our lack of acquaintance with bats is due chiefly to their nocturnal habits, and the consequent difficulty in observing them. To-day, bats are so little known that there are



Drawn by J. CARTER BEARD.

From a specimen in the Philadelphia Academy of Sciences.

#### BORNEAN NAKED BAT.

The young are carried in two dorsal pouches, from one of which, under the left elbow, a small head protrudes.

The great majority of bats are useful to man in destroying the insects which, without the aid of the birds and beasts, very soon would overwhelm him. The harmful species are those which destroy fruit, and a few which suck the blood of domestic animals.

Owing to certain natural conditions, the

perhaps a million persons who only know that they fly at night, and are "awful things to get into your hair."

I have seen thousands of bats, flying in many different places, but never yet saw one alight upon a woman's hair; and I believe they are no more given to doing so than are humming-birds.

From the bats of the United States, there is nothing to fear, for their claws and teeth are pitifully weak. One cross old "bumble-bee," angrily bumbling, is more dangerous to a peaceful community than all the bats of our country taken together. In some portions of South America, however, the vampire bats cause serious trouble.

Keen-eyed boys and girls all over the world should know that little is known concerning



• SKELETON OF PALE BAT.  
*Antrozous pallidus.*

the habits of bats, and much remains to be found out. These creatures are therefore excellent subjects for original investigation.

The Order of Bats as a whole contains about four hundred and fifty species, but it is safe to say that three-fourths of them are known only by their dry skins and skulls, and that their habits are quite unknown. The questions are,—why do bats live? Upon what do they feed? Are they useful to man, or injurious? What are their friends and their enemies? Do they migrate, and at what times? Where do they nest, or take shelter; and what are the facts about their young? What parasites and diseases have they?

Although the bat is a true mammal, it is almost as wide a departure from the ordinary, four-legged, land-going type as is a whale or manatee. Its hand reveals an extreme degree of what is called "specialization." For a mammal, the arms are of great length. The bones of the fingers are enormously extended, and connected with hairless skin as flexible as india rubber, to form a wing for flight. This wing membrane is extended on up the arm to the body and the legs, and is continued between

the legs and tail, where it forms a supporting parachute in flight.

The thumb of a bat is very short and free; and its nail is developed as a hooked claw, by the aid of which the creature can comfortably climb about or support itself. The favorite position of a bat at rest is hanging by its feet, head downward.

To be "as blind as a bat" is not to be blind at all, but rather to possess powers of vision that are uncommonly good in semi-darkness, or at night, and fairly good even in the broad light of day. When disturbed at midday, all the bats I have ever seen alive (perhaps twenty species in all) have flown away to places of security as briskly and successfully as so many swallows. The eyes of all night-flying bats are small, jet black, and look like tiny black beads, but those of the day-flying fruit-bats are very much larger in proportion.

The teeth of bats of different species show wide variation. In nearly all of the four hundred and fifty species, the canine teeth are as strongly developed as in the cat, and in some bats their proportions are really formidable. A careless examination of a bat's skull might easily lead one to believe that it belonged to a carnivorous animal. But the molar teeth will always tell the true story.

The insect-eating bats, which far outnumber all others, have cheek-teeth which terminate in sharp points, and are specially designed for cutting to pieces the hard parts of hard-shelled insects. The fruit-bats, however, have molars of a very different sort, with rather smooth crowns, for crushing instead of cutting. The blood-sucking vampire bats of South America have very large canine teeth with sharp, cutting edges, and even the molar teeth are formed with scissor edges, very much like the teeth of cats.

The teeth and skulls of bats exhibit many interesting and even extraordinary variations, but it is impossible to enumerate them here. The accompanying figures show the characters of two species found in the United States.

As previously remarked, very little is known regarding the habits of bats, chiefly because their nocturnal habits make it very difficult to find them, or to observe them. We know that in winter some of our species live in caves, in a semi-dormant condition. Dr. C. H. Eigenmann



says, of the thousands that inhabit Mammoth Cave, "they fly readily if disturbed in summer, but in winter they hang apparently dead. If disturbed, a few respiratory movements may be seen, and they may utter a few squeaks, when they again remain apparently lifeless. If knocked from the roof some of them fall to the bottom of the cave and flap about, others fly away. I have seen them leave a cave in mid-winter, after being disturbed, but fly no further than a hundred yards, then turn and enter the cave again."

In central Montana, where there are no trees, I once found a large colony of bats inhabiting a cave that a subterranean stream had washed under the prairie. In Arizona there is a cave which is said to contain "a million" bats. Once while hunting elephants in the Malay Peninsula, the attention of my companion and myself was arrested by a strange, pungent odor which filled the air. Upon investigating the cause of it, we discovered a large cave of a very interesting character, inhabited by thousands of bats, and floored with a layer of bat guano a foot or more in depth, representing the accumulation of a century.

In warm countries, bats inhabit hollow trees. But do they inhabit such homes, and actually hibernate in them in winter, in the temperate zone? On this point, direct evidence is desirable. Dr. C. Hart Merriam has proved that some bats of the North American temperate zone do mi-

grate, as birds do, going south in winter and returning in spring.

The conditions of wild life in the temperate zone are rather unfavorable to the development of large bats, and for this reason none of the bats of the United States are of large size or commanding importance. The large fruit-bats, or "flying foxes," can exist only where they can procure a good supply of fruit all the year round; and for this reason they are mainly confined to the tropics. During our northern winter, a true vampire bat could indeed prey upon the blood of domestic animals; but in zero weather, the naked wings of such a creature would freeze stiff in a very few moments. The large vampire bat of India, for some reason called the "false" vampire (*Meg-a-der'ma ly'ra*), which devours small frogs, fishes, small birds, and even bats smaller than itself, could live in our southern and southwestern states, but it would be impossible for it to go far north of the frost line. All bats inhabiting the colder regions of the temperate zone, within the snow limit, must either hibernate in winter, without food, or migrate.

Owing to the great number of species of bats, and of the many groups into which they have been divided, it is desirable to mention here only a few examples with which every intelligent person should be acquainted.

The bats have been divided by Nature into two Suborders, and six Families, as follows :


THE ORDER OF BATS

SUBORDERS.	FAMILIES.	EXAMPLES.	
<b>ORDER CHIROPTERA.</b> { Insect-Eating Bats: <i>Micro-chiroptera.</i> { Fruit-Eating Bats: <i>Meg-a-chiroptera.</i>	LEAF-NOSED BATS, . . .	<i>PHYL-LOS-TO-MAT'I-DAE.</i> . . .	{ Leaf-Nosed Bat. Blainville's Bat. Javelin Bat. Great Vampire.
	FREE-TAILED BATS,	<i>EM-BAL-LO-NU'R'I-DAE.</i> . . .	{ Bonneted Bat. Naked Bat.
	COMMON BATS, . . .	<i>VES-PER-TIL-I-ON'I-DAE.</i> . . .	{ Red Bat. Gray Bat. Big-Eared Bat.
	FALSE VAMPIRES, . . .	<i>MEG-A-DER-MAT'I-DAE.</i> . . .	False Vampire.
	HORSESHOE BATS, . . .	<i>RHI-NO-LOPH'I-DAE.</i> . . .	
	FLYING FOXES, . . .	<i>PTER-O-POD'I-DAE.</i> . . .	{ Flying Fox. Hammer-Headed Bat.

## THE FAMILY OF LEAF-NOSED BATS.

### *Phyllostomatidae.*

The members of this Family bear on their noses thin leaves of naked skin that stand erect behind, or partly around, the nostrils. These wonderful nose-leaves are pear-shaped, heart-shaped, wedge-like, and of many other forms. The ears are large, or very large; the wing mem-



CALIFORNIA LEAF-NOSED BAT.

(After Harrison Allen.)

brane reaches down to the foot; the tail is long, and sometimes extends a short distance beyond the interfemoral membrane. On the whole, the bats of this Family form an astonishing exhibit of facial oddities. All save a few species are confined to South America.

The California Leaf-Nosed Bat<sup>1</sup> may be taken as a very modest example, because it bears what is really a very simple form of nose-leaf. It is found in southern California and Mexico, and its pelage is very light-colored.

The most remarkable of all bat faces is that of a small, brown-colored West Indian species known as **Blainville's Bat**.<sup>2</sup> As a sport of Nature it stands fairly unrivalled, and shows what is possible in the fashioning of skin into ornamental forms. The ears are large and of most fantastic form, the chin is bedecked with a highly convoluted bib of skin, and the eyes and nostrils are almost lost amid the leaves and tubercles which cover the muzzle. As a whole, the appearance of the face of this bat suggests a highly complicated flower, like a double pansy. The skull is only five-eighths of an inch in length.

<sup>1</sup> *O-top'ter-us cal-i-for'ni-cus.*

<sup>2</sup> *Mor'moops blain'vill-ii.*

This species is quite uncommon, and practically nothing is known of its habits.

In fashioning the noses and ears of bats, Nature has done some very odd and curious work. The flowers of orchids are not more oddly fashioned than the heads and faces of some species.

Let it not be supposed, however, that these queer facial appendages and long ears of the leaf-nosed bats are purely ornamental. Dr. George E. Dobson, one of the greatest authorities on bats, has pointed out two very curious facts. (1) The bats with small ears and *no nose-leaves* fly most in the early twilight; and many, such as the fruit-bats, fly in the daytime. (2) The long-eared and leaf-nosed bats prefer darkness, and seek their food only at night.

Let us see if we can find a reason for this. A cruel investigator of the eighteenth century, named Spallanzani, once destroyed the eyesight of several bats, then suspended many silken threads from the ceiling of a room, and liberated the creatures. Although totally blind, the bats flew to and fro between the threads, without once striking them, and were equally successful in avoiding branches of trees that were introduced. It now seems certain that some bats possess a sixth sense, of which at present we know nothing, by which they are able to fly in total darkness, and avoid even the smallest obstructions.

It seems quite probable that the long ears and nose-leaves of the night-going bats aid their owners in guiding their flight; but the precise manner in which it is done remains to be discovered.

**The True Vampire Bats.**—By this name we seek to distinguish the bats which actually suck the blood of living creatures, from the so-called vampires which live on fruit. In South America there are five species of true vampires, three of which are known as the javelin



BLAINVILLE'S FLOWER-NOSED BAT.

(After Peters.)

bats, the others as the short-nosed vampires. The centre of abundance of these creatures appears to be the valleys of the Amazon and the Rio Negro, and the adjacent regions; but one of the species ranges all the way from Chile to Mexico.

Of the true vampires, the **Javelin Bat**<sup>1</sup> is the one which is most aggressive, and most dreaded. It bites horses and cattle, usually on the shoulders, neck or hindquarters, and makes a wound in the skin of sufficient depth to cause blood to flow freely, even after the bat has flown away. Naturally, an animal that is thus preyed upon soon grows thin in flesh, and becomes visibly weakened. On the island of Mucina, in the delta of the Amazon, the serious injuries inflicted by the Javelin Bats upon domestic animals have long been known.

But where true vampires are abundant, they do not confine their attacks to domestic animals. Human beings are occasionally called upon to pay blood tribute to the small wing-handed demons of the air. Men are bitten at night, when asleep, usually either upon the nose, or the feet. With its sharp-edged teeth, the creature makes a very small round hole in the skin, and by means of mouth suction which must be quite powerful, the blood is soon flowing freely. Fortunately, blood-poisoning is not an attendant evil of the Vampire's bite, and the wound seldom becomes painful.

The common Javelin Bat measures a little less than 4 inches in length of head and body, and in color is reddish brown. All the other true vampires are smaller, and all are practically tailless, the parachute membrane stretching between the legs, quite down to the feet, without the support of tail vertebrae. Naturally, these creatures are widely known; for any bat which lives upon warm blood, always drawn from a living fountain, is bound to acquire wide notoriety and a very evil reputation. The skull of a Javelin Bat, seen in profile, looks very much like the skull of a miniature wolf.

In order to illustrate once more how easily a harmless animal can acquire an evil reputa-

<sup>1</sup> *Phyl-los'to-ma, has-ta'tum.*

tion, and further emphasize the necessity of taking direct evidence before pronouncing a verdict, we introduce a 28-inch bat from South America, most unjustly called the **Great Vampire**,<sup>2</sup> but not really belonging to the genus of blood-suckers. Mr. H. W. Bates, the "Naturalist on the Amazon," lived for a time where this species was quite abundant, and of it he wrote in his book as follows:

"Nothing in animal physiognomy can be more hideous than the countenance of this creature when viewed from the front; the large, leathery ears standing out from the sides and top of the head; the erect, spear-shaped appendage [nose-leaf] on the tip of the nose, the grin, and the



BONNETED BAT.

*Promops californicus.* (After Harrison Allen.)

glistening black eye, all combining to make up a figure that reminds one of some mocking imp in a fable. [The very savage-looking canine teeth might well have been mentioned, also.] No wonder that imaginative people have inferred diabolical instincts on the part of so ugly an animal. The Vampire, however, is the most harmless of all bats." Mr. Bates opened the stomachs of a number of specimens, and found that "they had been feeding chiefly on fruits," and wild fruits, at that, obtained by honest hunting in the depths of the forest.

Moral: Never make an affidavit on the food habits of wild animals without first examining the stomachs of several specimens.

## THE FAMILY OF FREE-TAILED BATS.

### *Emballonuridae.*

The bats belonging to this Family have no nose-leaves, and the tail is partly free from the

<sup>2</sup> *Vam-py'rus spec'trum.*

membrane between the legs, either rising from its upper surface, or projecting beyond its end. The muzzle is rather blunt, and the nostrils open beyond the upper lip.

**The Bonneted Bat,**<sup>1</sup> of California and Mexico, is one of the largest of our species of free-tailed bats. Above the shoulders it looks like a rat wearing a poke bonnet. Its head-and-body length is  $2\frac{3}{4}$  inches, tail  $1\frac{1}{4}$ , total length of ear,  $1\frac{5}{8}$  inches. One-half the tail is free.

**The Naked Bat,**<sup>2</sup> of Borneo, Java and Sumatra, is one of the most remarkable species of the entire Order of Bats, and in some respects is the widest departure from the typical bat. In the interior mountains of Sarawak, Borneo, I once secured ten fine specimens, and to me they are as wonderful to-day as when handled for the first time.

As its name implies, this bat is practically destitute of hair, the only hair noticeable being a few stiff, black bristles on the neck, and a little microscopic fuzz on the breast and hind-quarters. The skin is thick and leathery, lying in numerous creases and folds, and on the living animal it is very elastic. There is no nose-leaf, and the lips are very thick and fleshy. The tail is free of parachute membrane for two-thirds of its length, and is quite like the tail of a mole. On the joint at the base of the thumb there is a large, callous tubercle, which indicates that this bat is much given to crawling about on "all fours," on rocks and tree-trunks.

Around the neck, the skin lies in two thick folds, and in these, directly under the chin, is situated a deep gland or sac which secretes a gummy substance with an odor both strong and disagreeable. Clearly, like the scent-gland of the skunk, it is for defence.

The most wonderful feature of the Naked Bat is yet to be noticed. On seeing this species for the first time, one's first thought is, how do the young bats cling to the parents during flight?

Nature, ever wise and provident, has answered this question by placing under each arm of this bat a *deep, wide pocket of rubber-like skin, in which the young are carried until they are able to fly!* The mouth of this pocket is on a line between the elbow and the knee, and it

extends upward and backward, over the entire shoulder, quite to the back-bone, where the two sacs are separated by a thin partition of skin. The pouch is  $1\frac{3}{4}$  inches deep, and in its lower portion, against the ribs, is located the mammary gland. On the whole, this is the most wonderful infant-pouch possessed by any living creature, not even excepting that of the marsupials, which is much more simple.

My largest specimen of this bat had a head-and-body length of  $5\frac{1}{2}$  inches, tail 2 inches long, and a wing expanse of 22 inches. In the skin were many curious folds. The face of the Naked Bat is coarse and ugly, and the body is quite devoid of grace and beauty; but ere one has time to scoff at such homeliness, the creature seems to say,—“Study me; for I am fearfully and wonderfully madé!”

This bat lives upon fruit and vegetation, and nests in hollow trees, rock crevices, or in holes in the earth. The illustration on page 59 was drawn from one of my Bornean specimens.

## THE FAMILY OF COMMON BATS.

### *Vespertilionidae.*

These are the bats that are most widely known, and also the most numerous. Dr. E. L. Trouessart recognizes more than 200 species. They range over all portions of the world that are habitable by small bats.

The distinguishing characters of the members of this Family are chiefly negative. There are no nose-leaves, the nostril openings are simple, and the tail is not produced to any extent beyond the interfemoral membrane.

All the bats of the United States are of small or medium size, and the majority of them belong to this Family. Along the Atlantic coast, they are so common that nearly every person living beyond the confines of the great cities is personally acquainted with at least one species. The commonest is the beautiful little **Red Bat**<sup>3</sup> which appears in the early twilight, gliding on swift yet noiseless wings up and down the shaded streets and roads, and occasionally making a friendly diversion into an open window, or through your veranda, partly for business purposes, and partly as an evidence of friendly regard.

<sup>1</sup> *Pro'mops cal-i-for'ni-cus.*

<sup>2</sup> *Cnc-i-ro-me'les tor-qua'tus.*

<sup>3</sup> *Las-i-u'rus bo-re-al'is.*

In midsummer, sharp eyes sometimes find this bat hanging close in amongst the leaves of a chestnut tree, its delicate fur as red as the brightest iron-rust. Touch it ever so gently and whisk! it is off as swiftly as a swallow, to seek another and a better hiding-place.

From sunset until it grows quite dark, it is very busy, and constantly on the wing. The



THE RED BAT.

Red Bat is a swift flyer, and much more of an aerial gymnast than any bird I know. In its flight it can turn abruptly with marvellous precision, and to me it is a constant source of wonder that it can fly so rapidly, turn and double so quickly, and dart in all possible directions without striking something. Almost any bird attempting to fly over the course of a Red Bat, and at the same speed, would probably come to grief in a very short time.

The only mistake that Red Bats are prone to make is in flying into houses through open windows, and instantly forgetting the location of the means of escape. Once in a room, the bat flies slowly, and frequently is so bewildered by the sudden change from semi-darkness to light that it strikes a wall, and falls to the floor. Although many persons are nervous about bats, I have noticed that whenever one flies in, some

kind-hearted and sensible person generally cries out, "Don't kill it!"

While crossing the Atlantic quite recently, a British **Long-Eared Bat** was found on board the steamer, thirty miles from the nearest land, clinging to the rail, wet and weary. At that time there was no breeze from the land.

When taken into the library, its wet fur soon dried, and it began to fly to and fro. In a short time the room was well filled with passengers, who watched the exhibition with great interest. When caught and held for close examination, it did not squeak shrilly and protest as the red bat usually does. After having served as a useful object lesson for a large number of young people, our strange visitor was brought safely to New York harbor, and liberated.

**The Gray Bat**<sup>1</sup> is one of the largest and handsomest species inhabiting the northeastern United States and Canada. It is also found throughout the middle West from Ohio to California, and from Manitoba to New Mexico. This is a species well worth looking for. It has small ears, a head-and-body length of 3 inches, tail 2 inches, and it is readily distinguished by its dark brown hair tipped with silvery white.

**The Big-Eared Bat**<sup>2</sup> of the south Atlantic states has ears of incredible height and width for a creature so small. In comparison with the size of the wearer, these ears are the largest worn by any American mammal. They are one-half as long as the entire head and body, being  $1\frac{1}{4}$  inches in height and nearly 1 inch wide, while the head and body measure only  $2\frac{1}{2}$  inches.

#### THE FAMILY OF FALSE VAMPIRES.

##### *Megadermatidae.*

This Family is absent from America, but is mentioned here to fill what otherwise would be a gap. The members of one genus, *Megaderina*, are noted for their carnivorous habits. The most noteworthy species is well worthy of mention.

**The "False" Vampire Bat**, of India and beyond, bears a name which is quite misleading; for in its habits, this creature is far from being a "false" Vampire. It devours frogs, small fishes, bats smaller than itself, and even

<sup>1</sup> *At-a-la'pha cin'e-re-a.*

<sup>2</sup> *Co-ry-norhi'nus ma-cro'tis.*

small birds. It has very large ears, an elaborate nose-leaf, a head-and-body length of 3 inches and a wing expanse of 16 inches.

### THE FAMILY OF HORSESHOE BATS.

#### *Rhinolophidae.*

This Family contains thirty species of small bats, all of which are restricted to the Old World.

### THE FAMILY OF FRUIT-EATING BATS.

#### *Pteropodidae.*

The members of this Family are bats of very large size, with fox-like heads, dense and abundant pelage, large eyes, and free tails when tails are present. They are quite diurnal in their habits, and feed almost exclusively upon fruit. They inhabit India, Ceylon, the Malay Archipelago and eastern Australia, and are almost the only bats that find their way into captivity for exhibition purposes. They are very sociable in their habits, and live in colonies of from five to fifty individuals.

**The Flying "Fox."**<sup>1</sup> The largest of the bats which we occasionally see darting through the gloaming with irregular, jerky flight, are about as large as purple martens,—tiny creatures, weak, and quite incapable of offence. In the East Indies, however, and also Australia, there are bats of enormous size. These are known as **Fruit Bats**, or **Flying "Foxes."** Some of those shot by the author in Ceylon had wings which spread forty inches.

On one occasion I found the top of a small tree, about fifty feet high, filled with these animals. They hung head downward from the upper branches, in places so thickly as to crowd each other,—quarrelling, squealing shrilly, and climbing about. To see nearly a hundred bats of such huge size hanging in one tree-top, quite at home in the broad glare of a tropical afternoon sun, was a strange and impressive sight. I had been asked to procure and preserve for American museums six dozen specimens of that species, and when after long observation I finally fired into the bunch, the black and brown cloud of giant bats that rose in the air, and slowly

<sup>1</sup> *Pteropus edwardsi.*

flapped away, was one of the most grewsome sights I ever saw in animal life. Of all creatures that fly, none are so thoroughly uncanny when outlined against the sky as the big, black-winged, half-naked Flying "Fox." They suggest demons and calamities.

The Flying "Fox" derives its name from the resemblance of its head to that of a very small fox. It feeds wholly upon fruit, and when it inhabits well-settled districts it is cordially disliked by every person who owns a fruit-tree. In some portions of Australia, these creatures have done great damage to fruit, and energetic measures, such as the explosion of dynamite among them, have been resorted to for their destruction.

Some of the fruit-growers of California are so apprehensive of this creature, and so fearful that it might be "introduced," they have secured the passage of a law, by which the importation of the Flying "Fox" is prohibited so rigidly that not one specimen can be imported, even for exhibition in a zoological garden. As a matter of fact, this fear of the presence of the Flying "Fox" in the United States is quite as groundless as the old fear of being quill-shot by Canada porcupines. It certainly would be very difficult to introduce that species, and keep it from being exterminated, except possibly in some of our insular possessions.

In the Flying "Fox" Family is found another remarkable variation in bat physiognomy, the **Hammer-Headed Bat**,<sup>2</sup> a species discovered in the land of the gorilla, by Du Chaillu. The head of the animal is of large proportions as compared with the body, and the muzzle is enormously enlarged. In general outline, the head in profile is much like the head of a moose. This is quite a large bat, its wing expanse being 28 inches.

<sup>2</sup> *Epomops*.



HAMMER-HEADED BAT.  
(After Joseph Wolf.)



FRUIT-EATING BATS, OR FLYING "FOXES."

## CHAPTER VII

# THE ORDER OF GNAWING ANIMALS

### GLIRES, OR RODENTS


The Order of Gnawing Animals contains a great many species, and to persons who have not studied it with some attention, it is a chaotic jumble of living creatures. This unsatisfactory condition is entirely unnecessary. A few hours' diligent study—under helpful conditions—will give any intelligent person a fair knowledge of the subdivisions of this Order, and an acquaintance with a sufficient number of examples so that each strange North American rodent met with can be referred to its proper Family.

The first step is to learn the names of the Families, which are as follows:

	FAMILIES.	APPROXIMATE NUMBER OF FULL SPECIES.
<b>ORDER GLIRES, OR RODENTIA.</b> (North of Mexico.)	SQUIRREL FAMILY, . . . . .	SCI-U'RI-DAE, . . . . . about 72 Species.
	SEWELLEL FAMILY, . . . . .	AP-LO-DONT'I-DAE, . . . . . 4 “
	BEAVER FAMILY, . . . . .	CAS-TOR'I-DAE, . . . . . 1 “
	MOUSE AND RAT FAMILY, . . . . .	MUR'I-DAE, . . . . . 171 “
	POUCHED MOUSE AND RAT FAMILY, . . . . .	DI-POD'I-DAE, . . . . . 42 “
	JUMPING MOUSE FAMILY, . . . . .	ZA-POD'I-DAE, . . . . . 10 “
	POCKET GOPHER FAMILY, . . . . .	GE-O-MY'I-DAE, . . . . . 33 “
	PORCUPINE FAMILY, . . . . .	E-RETH-I-ZONT-I-DAE, . . . . . 2 “
	PIKA, OR “CHIEF HARE” FAMILY, . . . . .	O-CHO-TON'I-DAE, . . . . . 6 “
HARE AND RABBIT FAMILY, . . . . .	LE-POR'I-DAE, . . . . . 30 “	
	371 <sup>1</sup>	

### THE SQUIRREL FAMILY.

In order to avoid recognizing a large number of Families for animals that are closely related, zoologists have agreed that the Squirrel Family shall contain the marmots, and a number of other animals that are closely related to squirrels. To make this point clear, observe this diagram:



All these creatures appeal strongly to persons who live in the country, or visit city parks. Go anywhere in the temperate zone, and you will find some of them, ready to greet you, and

make friends with you if you choose. You have but to use your eyes, and you will see them. In the East you have the gray squirrel and chipmunk; in the Mississippi Valley the fox squirrel; on the Great Plains, the ground-squirrels and prairie-“dogs”; in the West the Douglas squirrel, and a bewildering array of chipmunks and ground squirrels. He who fails to learn their names, and make friends with them, loses much pleasure.

The members of the Squirrel Family are so widely distributed, and have grown so accustomed to man and his ways, that there are few persons who have not seen at least two or three wild species in their haunts. Their lives are full of incident and interest, and to the young naturalist, animal artist or sculptor, they are usually the most available of all wild animal subjects.

A very attractive book might be written

<sup>1</sup>The subspecies recorded number about 260!



about the many beautiful and interesting species of squirrels that are found throughout North America, the number of which is surprisingly great. The total number of species and subspecies described is as follows.

In Mexico and Central America, species, about 25, subspecies, about 18, total 43; in the United States and Canada, species, about 60, subspecies, about 67. The total for North America is about 170 species and geographic races. Many of these, however, resemble each other so closely that their differences are too slight for our consideration; and there may be a number that are not entitled to stand as independent forms.

Nature has divided the many species of North

graveyard. There is no other animal of equal size that can add so much of life and cheerfulness to a hardwood forest or a meadow as a good healthy squirrel. *Why is it* that American men and boys kill them so eagerly? Surely the flesh of their little bodies is not needed as food. It has a taste so "gamey" and rank that to many persons it is decidedly unpalatable. Americans are the only white men on earth who eat squirrels. An Englishman would as readily eat a rat!

Possibly their flesh was necessary to the hardy but hungry pioneers of the early days; but today we have no excuse for shooting any squirrels, save the quarrelsome red squirrel. Surely no true sportsman or right-minded boy can



Photographed by E. R. SANBORN, N. Y. Zoological Park.

GRAY SQUIRREL.

American squirrels into three easily remembered groups, as follows:

**Tree Squirrels**, which live in the tree-tops. Example: Eastern Gray Squirrel.

**Rock Squirrels**, which live in rocks, fences and among the roots of large trees. Example: the Common Chipmunk.

**Ground Squirrels**, of prairie countries, which burrow deeply in the earth. Example: the Striped Spermophile.

In each of these three groups there are several important types which must be noticed.

### The Tree-Squirrel Group.

A patch of timber or a wood lot without squirrels always conveys an impression of lonesome solitude and something gone,—like a country

find any real "sport" in "potting" squirrels out of the tree-tops.

Take the common gray squirrel, for example. It is one of the most beautiful and graceful of our native mammals. It is perfectly harmless, and *as soon as it learns that it is protected, it becomes so tame as to be a delightful companion on the farm.* Thousands of American farmers would fight, were it necessary, to save their squirrels from slaughter. Except the red squirrel, all tree squirrels should be protected, both by public sentiment and by law.

Excepting the chickarees, the squirrels which live in the tree-tops are considerably larger than those of other groups, and their tails are much longer. Their characteristic colors are gray, rusty-brown, yellow and black; and as a rule

they are devoid of spots or stripes. They are very strong and active climbers, and keen of eye and ear.

**The Gray Squirrel**<sup>1</sup> is chosen as the leading type because it represents an average size, the most frequent color, and is widely distributed. This is the most prominent squirrel of southern Canada, New England, and the eastern and

The Gray Squirrel frequently consents to live in city parks, and becomes quite tame. It spends much of its time upon the ground, searching for nuts, roots, or anything which can be eaten. A very large specimen measures  $9\frac{1}{2} + 8\frac{1}{2}$  inches. Northern specimens are larger, and have longer and finer fur than those of the southern states.

**The California Gray Squirrel**<sup>2</sup> is the Pacific coast counterpart of the eastern gray squirrel, except that it is larger, and its colors are brighter. Its color above is bluish gray and black, and underneath it is pure white. It is the largest squirrel in the far West, its maximum length being  $12 + 10$  inches. Its home extends from the state of Washington to southern California, and it is in every way a worthy product of that fertile and healthful region.

**The Fox Squirrels.**—We have now reached two important species, to which the student must give close attention in order to avoid confusing them with each other, and with the gray squirrel. The southern species will be presented first, because it has two points by which it can be recognized at a glance.

**The Southern Fox Squirrel**<sup>3</sup> is the only Squirrel in America which has a *pure white nose* and *white ears*. No matter how much the remainder of the animal may vary in color from the standard, in adult specimens the white nose and ears are constant. Typical specimens of this species are colored as follows: top of head, black; upper surface, blackish brown; lower surface, lighter brown; tail, dark brown, margined with black.

Variations occur, of every shade from the above to jet black all over the body, head and tail; but the ears and nose still are white.

This animal measures  $13 + 12$  inches. Its home is east of the Alleghanies from Virginia to Florida, and westward along the Gulf Coast to Louisiana. On the map its range looks like an arm bent around the range of the next species.

**The Northern Fox Squirrel**,<sup>4</sup> or **Cat Squirrel**, is smaller than the southern species ( $12 + 11$  inches), but very much like it in color, save that its nose and ears *never* are *white*. The standard color is rusty brown, washed with black on the upper surface, and bright brown underneath.

<sup>1</sup> *Sci-u'rus gris'e-us.*

<sup>3</sup> *Sci-u'rus ni'ger.*

<sup>4</sup> *S. lu-do-vi-ci-an'us.*



SOUTHERN FOX SQUIRREL.

southern states, southward to Florida. It ranges westward to Minnesota, Kansas and Texas. Above its color is clean iron-gray, which in southern specimens is mixed with dull yellow. The lower surface is white, varying to yellowish brown. Usually it nests in hollow trees, but when crowded for room builds an open nest of green leaves, or strippings of cedar bark made into a round ball. The young are usually five in number.

<sup>1</sup> *Sci-u'rus car-o-li-nen'sis.*

**Variations.**—This squirrel is the most variable in color of all our species, and in fifty specimens it may be difficult, or even impossible, to find two exactly alike. Often it has a beautiful gray coat, and looks like a genuine gray squirrel with a brown back and head. Often it is dark gray above, and black on the legs and under surface,—a strange combination of colors,—and occasionally a pure white specimen is found.

This species inhabits the Mississippi Valley from the Alleghanies to Arkansas, western Iowa, and northward to Michigan and New York. In captivity it seems to be more hardy in winter than the gray squirrel. In the New York Zoological Park it blithely runs about in the snow when the latter takes pains to avoid it. Often the Northern Fox Squirrel will be out when none of the other occupants of the Rodents' cages are visible. It seems to me, however, that the Fox Squirrels are not as nimble on foot, or as active and daring in the tree-tops, as the gray squirrels.

**The Red Squirrel, or Chickaree,**<sup>1</sup> represents a large group of species containing the smaller of the tree squirrels. Its length is  $7\frac{3}{4}$  +  $5\frac{1}{2}$  inches, weight  $7\frac{1}{2}$  ounces. What it lacks in size it makes up in courage and activity. In New York and New England, it often drives all the gray squirrels out of any grove which they have undertaken to inhabit as tenants in common. Many observers believe the habits of the Red Squirrel to be so bad that the species deserves to be exterminated; but to this we are not prepared to agree. The complete destruction of any species of mammal or bird is a doubtful experiment, and never should be entered upon without most careful investigation.

In its normal colors, this little animal is readily recognized by its brown upper surface and outer surface of its legs, and its white under parts. It must be remembered, however, that it undergoes important seasonal changes in pelage,—from winter coat to summer coat, and the reverse,—and sometimes its standard colors are greatly changed.

Its legs are long and thin in proportion to the size of its body, and its form is not as graceful

<sup>1</sup> *Sci-u'rus hud-son'i-cus.*



EASTERN RED SQUIRREL.

as that of the gray or fox squirrels. It is readily recognized by its markings, and the fact that it is the smallest of our northern tree squirrels.

Three species and fifteen subspecies of Red Squirrels are recognized, and their combined ranges cover about two-thirds of North America, from Alaska and Labrador to North Carolina and southern Arizona.

In California and Oregon this group is represented by the sprightly and interesting **Douglas Squirrel,**<sup>2</sup> showing a mixture of colors,—dark gray, yellowish, and black. This is the most familiar squirrel of the great coast forests, in which it uses the sides of the giant spruces and redwoods as play-grounds. In Colorado and Utah occurs the third full species, known as **Fremont's Squirrel,**<sup>3</sup> which is colored gray, yellowish brown and white, much mixed.

Of the forty-three species and races of squirrels inhabiting Mexico and Central America, the most conspicuous is the **Red-Bellied Squirrel.**<sup>4</sup> Its upper surface is pale grizzled gray, and its under parts bright rusty red. It inhabits the forests of eastern Mexico, ascending the high mountains to an elevation of 8,000 feet.

The *largest squirrel* in the world is the great **Malabar Squirrel**<sup>5</sup> of southwestern India, which is yellowish brown above, reddish brown or black below, and measures, head and body, 17

<sup>2</sup> *Sci-u'rus doug'las-i.*    <sup>3</sup> *S. fre-mont'i.*

<sup>4</sup> *S. ery-thro-gas'ter.*    <sup>5</sup> *Sci-u'rus mal-a-bar'i-cus.*

inches, tail, 14½ inches, and it weighs 4½ pounds.

The *most beautiful* squirrel in the world is **Prevost's Squirrel**<sup>1</sup> of the Malay Peninsula, a species about the size of a small gray squirrel. Its colors form a beautiful pattern of gray, brown, black, white and buff.

### Rock Squirrels, or Chipmunks.

Next below the tree squirrels comes a large group of small squirrels which live on the ground,



EASTERN CHIPMUNK.

preferably amongst rocks, in which they find refuge from their enemies. In the absence of rocks, they live along fences, where any exist; but their favorite nesting-places are in hollow trees which can be entered directly from the ground.

These little creatures are about one-third the size of large tree squirrels, and inasmuch as their small size renders them secure from the deadly attentions of man, they have become the most tame and confiding of all the wild mammals of civilization. They are graceful in form, beautiful in color-markings, and exceedingly pert and

<sup>1</sup>*Sci-u'rus pre-vost'i.*

quick in their movements. When fully protected, as they are in some public parks, they become so tame and confiding that they dart about on the walks in search of food, and often allow persons to pass within three feet of them.

For convenience and clearness, we shall designate all the chipmunks as **Rock Squirrels**, because of their well-known preference for rocks, whenever any are available. It is a mistake to call these animals "ground squirrels." That name does not properly apply to them, but belongs to the next group.

**The Eastern Chipmunk**<sup>2</sup> is widely known, and will serve admirably as the key to the group. When you walk in the country, almost anywhere in the eastern states, this pretty little creature darts in front of you like a flash of brown light, and says, "Chip, chip, chip, chip!" most gleefully. If you stop to observe him, he pauses and looks at you very intently, wide-eyed and with ears erect, and save for the quick heaving of his tiny sides, remains as motionless as a stuffed squirrel.

To him, every fence is a fortress. Whether it be of stone or wood, the Chipmunk knows its best runs when danger threatens, and carries in his active little brain a complete check-list of burrows and hiding-places. When pursued by dog, boy or wild animal, he darts swiftly along the top or the lower rails of his stockade, until he reaches a satisfactory hiding-place, when a flash of brown fur shoots into it, and he is seen no more.

When hard pressed, Chipmunks frequently climb tree-trunks up to the lower branches, but such situations are very dangerous for them, because they are so seriously exposed to attack. Next to the birds of prey, the weasel, mink and fox are their worst enemies. The weasel is the worst of all, because it follows them into the remotest recesses of their burrows, and kills every inhabitant without mercy.

Although the Chipmunk burrows in the ground below the frost line, and has roomy cheek-pouches in which it carries astonishingly large quantities of grain and small nuts, it is more nearly related to the tree squirrels than to the true ground squirrels. In the autumn it stores in its burrow a quantity of grain or nuts, which ever is most abundant,—a habit which has sug-

<sup>2</sup>*Tam'i-as stri-a'tus.*

gested its generic name, *Tamias*, meaning a steward. It does not become dormant, but on the warm, sunny days of winter, when the rocks are free from snow, it hastens above ground to enjoy the light and warmth.

The length of an Eastern Chipmunk is  $6\frac{1}{2} + 4\frac{1}{2}$  inches. Its ground color is bright reddish brown above, light underneath, and along each side runs a conspicuous yellow-brown stripe between two black stripes. A black stripe runs from the head backward along the centre of the back, almost to the tail. The home of this animal extends from southern Canada and New York to Georgia and Louisiana, and westward to Iowa.

There are eighteen full species of Chipmunks, several of which are very much alike, distributed throughout nearly the whole of the United States. The greater number are marked by two or more black lines extending along the side, frequently alternating with lines of a yellowish-gray color.

It is impossible to mention even the majority of these species without risk of confusing the reader, but it is desirable to note a few important and strongly marked types inhabiting widely separated localities in the United States.

**The California Chipmunk**<sup>1</sup> is a merry-hearted little elf, particularly pert and beautiful. Its high, sharp-pointed ears and harlequin stripes of white give it a very roguish and saucy look. To judge by the lively actions of this little creature, it seems to regard life as a long play-spell. There are many in the Zoological Park, and in some respects they are the most satisfactory of all our burrowing rodents. Only the severest weather drives them into their burrows, and in the dead of winter, when a thick blanket of snow keeps all other animals of the Burrowing Rodents' Quarters snug under ground, the first hour of clear sunshine will see half a dozen of the California Chipmunks above ground, and sunning themselves on their logs. Having an abundance of room, they enjoy their life in the Park, and are much interested in visitors who notice them.

This species could easily and safely be introduced in any region suitable for it. Its home is in the San Bernardino and San Jacinto Mountains, California, but the limits of its range are yet to be defined. It is one of the smallest spe-

<sup>1</sup> *Eutamias spe-ci-osus*.

cies of its genus, its total length being 6 + 3 inches.

**The Antelope Squirrel**<sup>2</sup> is readily recognized by the broad and conspicuous band of white, which extends along the middle of the side, and its pale buff color. It has the pale colors of a



Photo. by E. D. WARREN.

SAY'S SPERMOPHILE.  
(*Callospermophilus lateralis*.)

WESTERN CHIPMUNK.  
(*Eutamias quadrivittatus*.)

desert animal. It is found in the desert regions of the southwest from western Texas to southern California, and northward to Nevada and Utah.

It is larger than the eastern species, and is strikingly different in appearance from all other chipmunks.

**Ground Squirrels.**

We have now reached a large group of burrowing squirrels which to the farmers west of the Mississippi are of very serious importance, on account of the grain they destroy. All these animals may be known under the name of Spermophiles. The word Spermophile means "seed-lover"; and as this very appropriate general term implies, the animals which bear it feed chiefly upon seeds or grain.

No ground squirrel, or spermophile, ever should be called a "gopher," as is frequently done in the Dakotas and Minnesota. The latter name should be reserved for the clumsy, burrowing pocket gophers, of the genera *Geomys* and *Thomomys*.

Ground squirrels live by preference on prairies,

<sup>2</sup> *Am-mo-sper-moph'i-lus leu-cu'rus*.

and burrow deeply in the ground. They seldom frequent rocks, and seldom climb trees. They are essentially dwellers in open country, where they can range freely, and behold a goodly portion of the world about them. Even fields of standing grain are distasteful to them, and they move to the open country around their borders.

Of spermophiles north of Mexico there are thirty-one full species and forty-two subspecies, or races. Going westward, they are first found in western Indiana and Michigan, from which they spread northwest and southwest throughout the whole western half of the United States, save the timbered areas. They also range into Mexico, Canada, and Alaska. They are at home on the rich, rolling prairies of the Dakotas, the level, floor-like plains of Nebraska, the alkali flats of Utah, the hot deserts of Arizona, and the dry valleys and mountain regions of California. They seem to be most numerous in California and the Dakotas, where they do much damage to crops.

All the ground squirrels have cheek-pouches, dig deep burrows (unless the earth is too rocky), store quantities of grain in the autumn for winter food, and in cold latitudes live all winter in their burrows. If forced to do so, they will live amongst rocks, and it is surprising to note how they can live in situations both high and low, dry and wet. Their favorite food is grain, seeds of every description, green grass, and hay, and their worst habit is digging up seed grain.

Some species eat quantities of destructive insects, such as grasshoppers, beetles, cut-worms, and crickets, and in this way partly compensate the farmer for the grain they devour. In fact, from all observations made thus far it seems that in the insect season, insects form a considerable proportion of the daily food supply of these industrious little animals. Not only do they eat all kinds of ground insects, but they also devour mice, and almost any other flesh that comes within their reach, particularly dry meat adhering to the bones of large animals which have died near their holes.

Ground squirrels are prolific, and bring forth from seven to ten young in each litter. Their enemies are coyotes, foxes, badgers, skunks, hawks and owls.

The spermophiles of North America are so wide-spread, so numerous and so important it

is necessary that two or three of the leading species should be specially noticed.

**The Thirteen-Lined, or Leopard Spermophile,**<sup>1</sup> is the most familiar and widely distributed species, and although one of the smallest, it is also the most strangely marked. Nature was in a sportive mood when she marked the back and sides of this little creature with seven broad stripes of dark brown, then laid between them six narrow stripes of pale yellow, and finally marked each of the seven brown stripes with a row of large, pale yellow spots. The yellow spots on the brown lines are the first feature of the color scheme to catch the eye, and they distinguish this animal almost as far as it can be seen. Its under parts are pale yellow, and its size is  $6\frac{1}{2} + 3\frac{1}{2}$  inches.

Do not call this animal the "Striped" Spermophile, because that name would apply to several other species, and be worthless; and do not call it the "Striped Gopher," because it is not a "gopher" of any kind.

The Thirteen-Lined Spermophile inhabits



THIRTEEN-LINED SPERMOPHILE.

about one-third of the United States, extending from Fort Wayne, Indiana, southwestward to Fort Worth, Texas, and northwestward to the plains of the Saskatchewan. Its western limit is the Rocky Mountains, but nowhere does it live in timbered regions, being strictly a prairie animal.

Its burrow is a hole about two inches in diameter, which descends quite steeply into the earth until it passes below the frost line (two to three feet), after which it runs off in a more or less horizontal course for ten or fifteen feet farther. If the burrow is an old one, and much used, it is a long and difficult task to dig to the end of it, and few boys undertake it more than once.

<sup>1</sup> *Citellus tri-decem-lineatus*.

As in the case of nearly all burrowing rodents of cold latitudes, nature has so adjusted the life of this animal that it survives the long and dreary winter in the strange, half-dead condition called *hibernation*. To make this possible, the young are born early in the year, and mature early, and during summer and autumn, take on a great quantity of fat. At the approach of winter, it curls up in its burrow for a sleep of from three to four months' duration.

By the investigations of Dr. P. R. Hoy, it has been discovered that in the case of the Thirteen-Lined Spermophile, the action of the heart is reduced from two hundred to only four feeble beats per minute, the temperature is reduced from 105° to 58°, and there is no visible breathing. The circulation of the blood was so feeble that when a limb was amputated, only a few drops of blood slowly oozed from the wound, while the nerves showed no sensitiveness. In fact, the animal was in a condition of suspended animation, as if under the influence of chloroform. In the northern portions of its range, this spermophile hibernates from about November 20 to April 1.

**Franklin's Spermophile**<sup>1</sup> looks very much like a slender-bodied, short-tailed tree-squirrel; and very often it is called the *Gray Ground Squirrel*. It should *not*, however, be called the "*Gray Gopher*," or "*Scrub Gopher*," for both these names are erroneous. It is best to call each animal by a name peculiarly its own, even though the beginning of correct naming involves a little trouble.

On an open prairie, especially in spring when the young grass is short, this spermophile is a conspicuous animal, and strongly resembles the gray squirrel of the East. Its upper surface is of a yellowish-gray color marked with fine, wavy, cross-wise lines of black or brown. Its under surface is distinctly gray, and its hair is coarse and stiff. In size it is about 9 + 5 inches. Its home is the central portion of the range of the Thirteen-Lined Spermophile. The western limit follows the eastern boundary of the arid plains northward from southeastern Kansas to the Saskatchewan, Alberta, and from thence south-eastward to southern Wisconsin, eastern Illinois and northern Missouri.

Whenever numerous in farming regions, this

<sup>1</sup> *Ci-tel'lus frank'lin-i*.

animal is very troublesome, not only in destroying grain in the ground and in the stack, but also in destroying young chickens. They are very venturesome in locating permanently near farm-houses and barns, and sometimes they are very destructive in gardens. As an offset to the valuable farm products destroyed by these creatures, Franklin's Spermophile destroys great numbers of noxious insects, such as grasshoppers, caterpillars, beetles, and also field mice. In the



RICHARDSON'S SPERMOPHILE.

United States Department of Agriculture, twenty-nine stomachs were examined with the following result: animal matter present, 30.3 per cent.; vegetable, 68.5 per cent., and undetermined, 1.2 per cent. Out of the whole twenty-nine stomachs examined, twenty-six contained the remains of insects! Thus the grain consumed by this animal is at least partially paid for by the destruction of insects that prey upon crops; but farmers everywhere are diligent in destroying it with poisoned wheat placed in its burrow.

**Richardson's Spermophile**,<sup>2</sup> of northern Montana, North Dakota and the region immediately northward as far as the Saskatchewan, has a short body, short legs, and a short tail, and looks very much like a thin prairie-"dog." In color it is like the preceding species, except that its tail is darker; but in size it is a trifle smaller (9 + 3 inches). Its habits are practically identical with those of Franklin's Spermophile, but if there is any difference, it is more destructive to grain than is the latter, and consumes less insect food. It is fortunate that this species inhabits so small an area of the wheat country of the Northwest.

<sup>2</sup> *Ci-tel'lus rich'ard-son-i*.

### Marmots.

The group of marmots consists of burrowing rodents which in structure are quite squirrel-like, but are distinguished by their large size and general heaviness of body. As befits their portliness of form, they are not active and lively, like squirrels, but live quietly and unobtrusively. By reason of the good sense they manifest in keeping out of mischief, some of them are tolerated in farming communities when more aggressive rodents would be exterminated.

The woodchuck is our most perfect type of Marmot, from which the prairie-"dog," or prairie marmot is slightly removed by the possession of a large and perfect fifth claw. It is desirable, however, that the latter should be included in the group of marmots.

### The Prairie-"Dogs."

The Prairie-"Dog"<sup>1</sup> is a plump and sociable little Rodent, not a Carnivore,—well known to every dweller in the plains region of the great



PRAIRIE-"DOGS."

West, and to every trans-continental traveller. His explosive, yapping cry is the most cheerful sound of the western plains. He hates solitude, and always lives in colonies of from 40 to 1,000 individuals. Unlike most other burrowing Rodents, the darkness and silence of a burrow easily pall upon his vivacious nature; therefore he spends the greater portion of his waking hours above ground, visiting his neighbors, and observing what goes on in his small world.

<sup>1</sup> *Cy-no'mys lu-do-vi-ci-an'us*.

When no enemies are in sight, he and his fellow-townsmen roam about for short distances from their homes, and feed upon grass blades and stems. At the approach of an enemy,—man, coyote, badger, fox, gray wolf, eagle or hawk,—the sentry cries out sharply, "*Skip! Skip! Skip!*"

Instantly every "Dog" halts, motionless and alert. If the sentry again cries "Skip!" each "Dog" scurries to his hole, and poises himself over its wide mouth, in readiness for a dive to subterranean safety. If the danger approaches quite near, the alarm cry resounds shrilly from all sides, stubby tails jerk nervously as if worked by wires, and down goes every Prairie-"Dog."

Just how far down the burrows go, it is difficult to say, for they probably vary greatly in depth. The mouth of a burrow is a miniature model of a volcano,—a conical mound of bare earth, a foot high and three or four feet in diameter, with a four-inch crater in the centre, going down at a slight angle. The crater prevents water from running into the burrow.

In making a crater the "Dogs" press the earth into shape on the inside with their noses. Once when an inmate of the Prairie-"Dog" Village in the New York Zoological Park incurred the hostility of four of his mates, they drove him into his burrow, filled up the mouth of it with moist earth, and with their noses tamped it down quite hard, the prisoner scolding vigorously meanwhile.

Prairie-"Dogs" are easily introduced into almost any open country where the ground is dry, but they are very difficult to exterminate. Under fair conditions they breed readily in captivity, and usually produce four young at a birth. In 1899, a free colony was established in the New York Zoological Park in the Antelope Range, where it existed for two years, and its saucy members attracted far more attention than those confined in the fenced village. Knowing that guns and dogs are not allowed in the Park, they often permitted visitors to pass within six feet of them. But it proved impossible to keep those industrious diggers from spreading far beyond the limits fixed for them, and seriously damaging walks and lawns, so they were finally caught by placing sand in boxes over their burrows, and transferred to the village whose walls of solid masonry go down to bed rock.



Some plainsmen claim that these interesting little creatures are able to locate their towns away from streams because they burrow down until they strike water, but Dr. Merriam points out the fact that in some regions they live where the nearest veins of artesian-well water are 1,000 feet below the surface. As a matter of fact *they can live without drinking.*

The Prairie-"Dog" is at home—where not exterminated by poisoned wheat put into his burrow—from Texas, New Mexico and Arizona northward to the Canadian boundary, and on the western slope of the Rocky Mountains in Utah and Colorado. It is most abundant in Montana, Wyoming and western Kansas. One of the largest Prairie-"Dog" towns yet reported begins in Trego County, Kansas, five miles west of the one-hundredth meridian, and extends along the divide north of the Smoky Hill River, practically without a break, to Colorado, a total distance of about one hundred miles. This town varies in width from half a mile to five miles, and on the top of the divide the nearest water is believed to be 350 feet below the surface. (Arthur B. Baker.)

It is now (1903) reported that because of the wholesale destruction of wolves and foxes, the enormous increase of Prairie-"Dogs" in Kansas, Oklahoma, Texas and Colorado has become a genuine scourge to farmers and cattlemen. The number of "Dogs" in that region is now estimated at several millions, and a general campaign against them has been begun. The method employed for their destruction is a spoonful of poisoned wheat placed in the mouth of each burrow. Beyond doubt, this will soon reduce their numbers to reasonable limits.

When he is not too numerous, I am the friend of the Prairie-"Dog." He is as bright and cheerful as the day is long, and he enlivens many a dreary landscape, but at the same time he often changes fine, grass-covered cattle ranges into dreary wastes, and causes great losses to cattle owners. I hope, however, that he will be tolerated at least to the extent that systematic destruction will stop short of extermination.

It is *not* true that the Prairie-"Dog" lives in peace and harmony in the same burrow with the rattlesnake and burrowing owl. The snakes would make short work of the young Prairie-"Dogs," and the latter would quickly kill the

owl! It is safe to surmise that when a deadly and quarrelsome rattler invades the home of a Prairie-"Dog" family, the latter speedily seeks a home elsewhere. The burrowing owl is in the habit of taking refuge in *abandoned* burrows, and nesting in them, to save the labor of digging a burrow for itself. In the Philadelphia Zoological Garden Mr. A. E. Brown once tried the experiment of associating burrowing owls and Prairie-"Dogs." The owls were immediately killed and torn to pieces by the "Dogs."

### A Prairie-"Dog" Burrow.

At last a Prairie-"Dog" burrow has been completely exposed by digging, and reported upon in full in one of the publications of the Biological Survey. In the "Yearbook of the Department of Agriculture" for 1901, Dr. C. Hart Merriam publishes a valuable paper on "The Prairie-Dog of the Great Plains," which contains the following illustrated description:

"The holes go down for some distance at a very steep angle and then turn at nearly a right angle and continue horizontally, rising somewhat toward the end. The nests are in side chambers connecting with the horizontal part of the burrow, and usually, if not always, at a somewhat higher level. (See H in figure.) Recently, at Alma, Nebraska, W. H. Osgood dug out a burrow, of which he made a careful diagram, accompanied by measurements.

"In this case the burrow went down nearly vertically to a depth of 14½ feet below the surface, when it turned abruptly and became horizontal as shown in the diagram. The horizontal part was 13½ feet in length. One-third of the horizontal part (the terminal 4 feet, F) and two old nests and passageways (E) were plugged with black earth brought in from the surface layer, which was very different from the light-colored clayey earth in which the greater part of the burrow lay.


"Four or five feet below the entrance was a diverticulum, or short side passage (G), probably used as a place in which to turn around when the animals come back to take a look at the intruder before finally disappearing in the bottoms of their burrows. It is also used, apparently, as a resting-place where they bark and scold after retreating from the mouths of the

burrows. As elsewhere noted, they are often heard barking after they have gone in.

"The burrow was opened the day after bisulphide of carbon had been used for destroying the animals, and the material carrying the bisulphide was found at the bottom of the vertical

serve to hold its numbers in check. The most inveterate of these appear to be the coyote, badger, black-footed ferret and rattlesnake."

**The Woodchuck, or Ground-"Hog,"** is tolerated on the farms of New England because he is wise enough to live on clover and other grass,



PRAIRIE-"DOG" BURROW.

From Dr. C. Hart Merriam, "Yearbook," Department of Agriculture, for 1901.

part, just where the horizontal part turns off. Two dead animals were found, one in the horizontal part, the other in the nest, as indicated by the letter K in the diagram.

"The Prairie-Dog has several natural enemies which, when not interfered with by man, usually

and let the vegetable gardens alone. In the East he is the only representative of the marmots. In form he is short and stout, and his flat head and beady, black eyes give him a surly look. He is not lively and cheerful in his habits, like a

<sup>1</sup> *Mar-mo'ta mo'nax.*

prairie-“dog,” and it is seldom that anyone speaks well of him. His favorite home is a burrow in a gravelly hillside in a “swamp lot,” or woods pasture, and while he likes to come out and bask in the warm sunshine, he never ventures far from his front door.

In the autumn, instead of storing up vegetables for winter, he takes on a quantity of fat, under his skin. Early in November he blithely goes to sleep in his burrow, and does not waken

York to Georgia, and westward to Kansas and South Dakota.

A much larger species called the **Gray Marmot**,<sup>1</sup> or **Whistler** (22 + 7 inches), is an important northwestern form, strongly marked by its light, grizzly-gray color, with certain dark markings. It is found from the Columbia River northward to about 63° North Latitude and eastward to Hudson Bay. It derives one of its names from the fact that its alarm cry consists of



WOODCHUCK.

until February 2,—“Ground-Hog day.” Then,—so runs the popular legend,—he emerges, and looks about him. If he sees his shadow, he again retires to his burrow, and sleeps six weeks longer,—which betokens a cold, wintry spring.

The eastern Woodchuck is a typical marmot, short-legged, heavy-bodied, flat-headed, and brownish gray in color. The length of its head and body is 14 inches, and of its tail 5 inches. It inhabits the eastern United States from New

a shrill whistle, which is repeated by the various members of the colony threatened with danger.

**The Yellow-Bellied Marmot**,<sup>2</sup> easily distinguished by the bright red hair on its under parts, is a southern species, found in California, Arizona, New Mexico and Texas. High up, on the Olympic Mountains of western Washington, is found still another species of marmot, as large as the

<sup>1</sup> *Mar-mo'ta pru-in-o'sus.*

<sup>2</sup> *Mar-mo'ta flav'i-ven-ter.*

Whistler, which is yellowish in summer, and bluish-gray in winter. This is called the **Olympic Marmot**.

### Flying Squirrels.

The **Flying Squirrel**<sup>1</sup> is a very beautiful little creature, but its strictly nocturnal habits, and strong dislike to daylight, almost rob us of its acquaintance. This is to be regretted, because it is the only native tree-dwelling quad-



THE FLYING SQUIRREL.

ruped which has been provided by Nature with a parachute, consisting of a thin fold of skin stretched between the fore and hind legs, to partly sustain the animal in a long downward flight. Neither the Flying Squirrel, nor the flying lemur of the East Indies, can actually fly; but they leap from a tree-top, go sailing gently downward and outward, and when near the ground curve upward and are carried by their momentum on an ascending plane to the side of an adjoining tree. Anything like horizontal flight is quite out of the question.

<sup>1</sup> *Sci-u-rop'te-rus vo'lans*.

The Flying Squirrel is one of the most exquisite little mammals in North America. Its legs are very delicately formed; its fur is as fine and soft as silk; and when at rest the edge of its flying membrane looks like the edge of a lace ruffle. The head and body (of the eastern species) is about 5 inches long, and the tail 4 inches. These little creatures are quite sociable, and nest in hollow trees, where from five to seven young are born. They come out to play about sunset, and are as sportive as schoolboys playing tag. In captivity they are quite worthless for exhibition, for in the daytime there is nothing to be seen save a small and wholly uninteresting ball of fur.

Three species (and nine subspecies) have been described, and their range covers the eastern United States from Canada to Florida, and westward to Louisiana. On the Pacific Coast, they are found from southern California to Alaska, even to the Mackenzie River basin, but they are not found in the desert regions.

### THE SEWELLEL FAMILY.

#### *Aplodontidae.*

The Sewellel,<sup>2</sup> Mountain "Beaver," or Show't'l of the Indians is a strange and little known animal of the Northwest, with which at least every person in that region should be acquainted. It is reddish-brown in upper color (sometimes grayish-brown), and looks like a tailless woodchuck. It feeds like a beaver, fights fiercely when cornered, is sociable in habit like the prairie-"dog," can climb bushes four feet high, and can burrow and live comfortably either in ground that is low and boggy, or high and dry. Usually it prefers wet ground! A large specimen weighs 4 pounds, measures about 13 inches in length of head and body, and tail a little more than one inch. Strange to say, this once rare animal has recently been discovered inhabiting the grounds of the University of Washington, at Seattle.

### THE BEAVER FAMILY.

#### *Castoridae.*

The Beaver<sup>3</sup> easily leads the mammals of the world in mechanical and engineering skill, and also in habits of industry. Being chiefly nocturnal in its habits, it sleeps by day, and after nightfall carries on its work unmolested.

<sup>2</sup> *Ap-lo-don'ti-a ru'ja*. <sup>3</sup> *Cas'tor can-a-den'sis*.



AMERICAN BEAVERS AND THEIR WORK.

The dam, and house of sticks in the middle of the pond, are exact reproductions of those works in the Beaver Pond of the New York Zoological Park, as they were at the time this drawing was made.

It is seldom that anyone sees a live Beaver in its haunts during the middle of the day, but it is possible to do so during the hour before sunset. In public zoological gardens and parks, the persistence and success of this animal in avoiding observation is very disappointing to visitors, and exasperating to directors and keepers.

This is the largest gnawing animal in North America. A huge specimen caught in Maine, in 1900, weighed a trifle over 50 pounds. A large one in the New York Zoological Park is 31 inches long, has a tail 12 inches long and weighs 44 pounds.

The American Beaver is still found in a few localities,—but in very small numbers,—from the Rio Grande in Texas throughout the Rocky Mountains, Sierra Nevada and Cascade Mountain regions northward to the limit of trees, and southeastward through Canada to northern New England. The number now remaining in Colorado has been estimated at one thousand.

The Beaver's efforts are directed toward its own preservation and comfort. It builds extensive dams of mud, grass and sticks, in order to create ponds in which it can hide from its enemies, maintain a safe refuge close by the wood on which it feeds, and have an under-water doorway to its house or burrow. More than this, the pond serves as a refrigerator, in the bottom of which the animal stores its supplies of food-wood for winter use, when the surface is frozen for a long period.

Sometimes when food-wood on a beaver pond becomes scarce, the animals dig canals into places where fresh supplies can be cut, and floated down to the pond. These canals are usually about two feet wide.

A Beaver is readily recognized by its very flat, hairless and scaly tail, which beyond the hair of the body is about 9 inches long by 4 inches wide. The tail is never used as a trowel in building dams, but only as a propeller in swimming.

Dam-building is done in two ways. With his front feet the animal digs up soft mud, holds the mass with his fore legs against his breast, and swims with it to the dam. There he deposits it where it is most needed, and pats it down with his front feet. To strengthen the structure, he brings sticks four or five feet long, and one or two inches in diameter, from which he has eaten the bark. These he usually lays upon the dam,

crosswise or nearly so, and fills between them with mud.

When Beavers have to build a dam exceeding fifty feet in length, to flood low ground, they usually lay it out with a curve up-stream. The dam built by the Beavers in the New York Zoological Park is about forty feet long, and three feet high, and quite sharply curved up-stream.

In most localities inhabited by Beavers, the banks of the streams are so low that the animals cannot burrow into them, and consequently they build houses for themselves. The ordinary Beaver house is a huge pile of neatly trimmed six-foot poles, with all spaces between the sticks plastered full of mud. The one in the Zoological Park is about fifteen feet in diameter, and five feet high, with a central chamber above high-water-mark, and its only entrance is well under water. If a beaver house is attacked, the occupants immediately seek refuge in deep water.



SKULL OF BEAVER, A TYPICAL RODENT.

The trees which furnish bark most prized by the Beaver as food are the poplar, cottonwood, willow, birch, elm, box-elder and aspen. The bark of the oak, hickory, or ash is not eaten.

The Beaver's front teeth (incisors) are very strong and sharp, and the muscles of the jaw are massive and powerful. It is no uncommon thing for a Beaver to fell a tree a foot in diameter in order to get at its branches. It is said by some observers that large trees are made to fall as the Beavers prefer to have them,—toward their pond. In felling a tree, they first remove the bark from a circle a foot in width, just above

the spur roots, standing on their hind legs while they work. Then, with their huge, chisel-like incisors they cut out chips, circling round the trunk all the while, until only the heart of the trunk remains, and the tree falls.

## THE FAMILY OF MICE AND RATS.

### *Muridae.*

When their groups and relationships are fairly understood, the wild mice and rats will be found quite interesting. They are so widely distributed it is very desirable that country-dwellers should know something about them, and appreciate their good points as well as their bad ones. A moderate effort, properly aided and encouraged, will give anyone a fair conception of the grand divisions of this great group; and there the general student can stop, if he so elects.

In approaching this assemblage of North American mammals, the first thought is that its members are difficult to deal with. In some respects they are, but they are by no means as difficult as might be supposed. Like many other new subjects, they yield to a little old-fashioned study. It is not necessary for the general student to enter into the study of a large number of species. Lay the foundation first by becoming acquainted with each genus, and one typical species. Observe the following injunctions:

1. Treat this bit of study with serious attention.
2. Learn first the names of the Families, and the approximate size of each Family.
3. Next learn *by rote*, in regular order, the *common names* of the typical examples given.
4. Learn some of the distinguishing characters of each example.
5. Study the comparative sizes of the various types.
6. Finally, in determining the name of a strange species, *do not feel that you must name it instantly*, or be disgraced! Take time to think over it, and to "look it up." Snap judgments on small creatures have a most annoying habit of proving to be wrong. It is a wise judge who knows when to hand down a decision.

In order to make the genera of North American rats and mice clear to the student, I have procured from Dr. C. Hart Merriam, the highest

living authority on these creatures, a fine, perfect, adult specimen of the best known (or most typical) species of each genus. Figures of these skins are here reproduced to show their relative sizes, and a life-like illustration of each of these types is also given. In the text, the most striking distinguishing characters are printed in italics.

With these aids to the text, it should be possible for a clear-headed, keen-eyed student to refer any adult North American rat or mouse to its proper genus. But beware of *young* specimens! Often they are so puzzling that Solomon himself could not place them with any degree of certainty. In determining the species of mice and rats, mammalogists depend largely upon the characters of the teeth; but that is a subject too intricate for the general student.

The table on page 84 shows the various Families of rats and mice, the North American genera, and the typical species of each. It is not necessary for young students to memorize the Latin names of the genera and species; but those who become specially interested in natural history will very soon desire to know them.

**The Muskrat,**<sup>1</sup> which received its name from its very pronounced musky odor, is the largest native representative of the Mouse and Rat Family. It is readily recognized by its *flat, hairless tail, carried on its edge*. It is of large size, measuring about 21 inches in length. It is of aggressive habit, an admirable diver and swimmer, an industrious and intelligent house-builder, and the only native rat whose fur is valuable. It is found from Labrador and Newfoundland to Alaska, and southward to Arizona and Louisiana.

It is very shrewd in preserving its own life, and even in the large forest parks of New York City, it refuses to be exterminated. When three bogs in the New York Zoological Park were dug out and converted into ponds, the wild Muskrats in the Bronx River found them as soon as they were completed, immediately took possession of them, and there they still remain. Being very destructive to lily bulbs, and most other aquatic plants, their presence in ornamental ponds is very objectionable.

Muskrats are rarely, if ever, found away from ponds or good-sized streams. They are quite as much at home in the water as beavers, and their

<sup>1</sup> *Fiber zibethicus*.

TYPICAL NORTH AMERICAN MICE AND RATS (north of Mexico).

	COMMON NAME OF		SCIENTIFIC NAME.		APPROXIMATE	
	GENUS.		GENUS.	TYPE SPECIES.	FULL SPECIES.	SUB-SPECIES.
<b>MOUSE AND RAT FAMILY.</b> ( <i>Mu'ri-dae.</i> )	Muskrat, . . . . .	<i>Fiber</i> , . . . . .	<i>zibethicus</i> , . . . . .		4	3
	Lemming, . . . . .	1. <i>Di-crosta'omyx</i> , . . . . .	<i>hud-so'ni-us</i> , . . . . .		3	5
	Lemming Mouse, . . . . .	2. <i>Syn-ap'to-mys</i> , . . . . .	<i>coop'er-i</i> , . . . . .		8	1
	Field Mouse, . . . . .	3. <i>Mi-crot'us</i> ( <i>Ar-vic-o-la</i> ), . . . . .	<i>penn-syl-van'i-cus</i> , . . . . .		48	18
	Red-Backed Mouse, . . . . .	4. <i>E-vot'o-mys</i> , . . . . .	<i>gap'per-i</i> , . . . . .		17	5
	Vole, . . . . .	5. <i>Phe-nac'o-mys</i> , . . . . .	<i>o-ro'phi-lus</i> , . . . . .		7	
	Wood Rat, . . . . .	6. <i>Ne-o-to'ma</i> , . . . . .	<i>flor-i-dan'a</i> , . . . . .		17	19
	Harvest Mouse, . . . . .	7. <i>Reith-ro-don'to-mys</i> , . . . . .	<i>le-cont'i</i> , . . . . .		10	6
	Rice-Field Mouse, . . . . .	8. <i>O-ryz'o-mys</i> , . . . . .	<i>pa-lus'tris</i> , . . . . .		2	3
	Cotton Rat, . . . . .	9. <i>Sig'mo-don</i> , . . . . .	<i>his'pi-dus</i> , . . . . .		3	5
	White-Footed Mouse, . . . . .	10. <i>Per-o-mys'cus</i> , . . . . .	<i>leu-co'pus</i> , . . . . .		42	27
Grasshopper Mouse, . . . . .	11. <i>O-ny-cho'mys</i> , . . . . .	<i>leu-co-gas-ter</i> , . . . . .		6	6	
Domestic Rat, . . . . .	<i>Mus</i> , . . . . .	<i>nor-veg'i-cus</i> , . . . . .		4		
<b>THE CHEEK-POUCHED FAMILY OF MICE AND RATS.</b> ( <i>Het-e-ro-my'i-dae.</i> )	Subfamily of the Pocket Mice, . . . . . (Species small.)	{ 12. <i>Per-og-na'thus</i> , . . . . .	<i>fas-ci-a'tus</i> , . . . . .		26	15
		{ 13. <i>Mi-cro-dip'o-dops</i> , . . . . .	<i>meg-a-ceph'a-lus</i> , . . . . .		1	
	Subfamily of the Kangaroo Rats, . . . . . (Species larger.)	{ 14. <i>Di-pod'o-mys</i> , . . . . .	<i>mer'ri-am-i</i> , . . . . .		5	8
	{ 15. <i>Per-o-di'pus</i> , . . . . .	<i>rich'ard-son-i</i> , . . . . .		9	1	
<b>JUMPING MOUSE FAMILY.</b> ( <i>Za-pod'i-dae.</i> )	Jumping Mouse, . . . . .	16. <i>Za'pus</i> , . . . . .	<i>hud-so'ni-us</i> , . . . . .		10	10
					256	145

habits are strictly aquatic. The tail furnishes the motive power for swimming. The feet are small, and but very slightly webbed, and the body is completely covered with soft, brown fur an inch or more in length, which is much sought by furriers. When taken at the best season, plucked, dressed and dyed a rich brown-black, it is known to the trade as "French seal."

Musk rats that inhabit streams with high banks do not trouble themselves to build houses, but merely burrow into the banks. In rivers and ponds with low margins, however, they gather coarse grass, reeds and mud, and build dome-shaped houses, about five feet in diameter, which rise from two to four feet above the water. All such houses are entered below the surface of the water, so far down that ice does not close their doors, and within there is a floor raised well above the water, on which the inmates eat their food, and sleep.


When too many captive Muskrats are kept in

the same enclosure, say twelve in a fenced pool thirty feet square, they fight viciously, and not only kill each other, but sometimes partly devour one of the victims. Although often disputed, it is nevertheless a fact that they eat flesh on very slight provocation. They are very unsatisfactory animals to keep in captivity, no matter what the conditions may be.

**The Hudson Bay Lemming**<sup>1</sup> is worthy of special notice, because it is the most widely-distributed and noteworthy rat-like animal of the far North. It is strictly a mammal of the cold northland, and like many other arctic animals, its winter coat is pure white, and its fur is dense and warm. Among the west Alaskan Eskimo, skins are very common, and the children delight in using them for doll clothes. (Charles H. Townsend.)

This animal is about the size of a large mole,  
<sup>1</sup> *Dicrostonyx hudsonius*.





1 LEMMING

2 LEMMING MOUSE

3 FIELD MOUSE

4 RED-BACKED MOUSE

5 VOLE

6 WOOD RAT

7 LITTLE HARVEST MOUSE

8 RICE-FIELD MOUSE

9 COTTON RAT

10 WHITE-FOOTED MOUSE

11 MOLE-MOUSE

*Pocket-Mouse Sub-Family.*

POCKET-MOUSE

*Kangaroo-Rat Sub-Family.*

CHEEK-POUCH FAMILY.

KANGAROO RAT

*Heteromyidae.*

MOUSE AND RAT FAMILY.

*Muridae.*

JUMPING-MOUSE FAM.

*Zapotidae.*


THE FAMILIES OF NATIVE MICE AND RATS, ILLUSTRATED BY SKINS OF TYPICAL SPECIES.

The scientific names of the above specimens will be found on the opposite page, against the corresponding numbers.

thick-bodied, short-legged, and sharp-nosed. The ears are extremely short, and quite hidden in the fur; the legs are short, the feet rat-like, and the tail is so very short that it also is half hidden by the fur. The fur is long, fluffy and fine; brown, brownish-gray, or mottled in summer, but snow-white in winter. The length of the head and body is 4 to 5 inches, and of the tail,  $\frac{1}{2}$  inch.

The Lemming is found from Latitude 56° northward to the whole arctic coast; in Labrador, Greenland, the arctic islands, and on as far north as man has ever gone on land. It prefers

*Its ears are very small, and do not rise above the fur on the head.* The type species, known as **Cooper's Lemming Mouse**,<sup>1</sup> is only two-fifths the size of the Hudson Bay Lemming. It inhabits the northeastern United States, from Massachusetts to Minnesota, and southward to North Carolina, Tennessee, Indiana and Iowa. Its color above is yellowish-brown washed with black, with bluish-gray or whitish underparts. Length,  $3\frac{1}{4}$  to 4 inches; tail,  $\frac{3}{4}$  inch. Other species of Lemming Mice inhabit Canada, Labrador, New Hampshire, Washington, Kansas and Alaska.



HUDSON BAY LEMMING.  
Winter and summer pelage.

FIELD MOUSE.

open, dry, moss-covered uplands, and is not found in timbered regions. Often a district of acceptable ground is covered with a wide-spreading network of runways, just below the surface. Mr. C. H. Townsend, who has kept them in captivity, says they are kind-spirited and sociable little creatures, fond of attention, and much given to standing up and hopping about on their hind legs. In summer they store up supplies of vegetable food in their runways for use in winter.

**The Lemming Mouse, or False Lemming,** is interesting chiefly because it is a connecting link between the true lemmings and the mice.

**The Field Mouse, or Meadow Mouse,**<sup>2</sup> stands as a murine monument to scientific endeavor. Since 1798, the genus of this group—long known as *Arvic'o-la*—has been described under twenty-four different names, and the type species has received nineteen names besides its own! But, through a century of misnaming in Latin, its original English name, Meadow Mouse, has stood unchanged!

The trouble with this genus seems to have been due to exaggerating the importance of trivial characters, molar teeth and claws. Externally

<sup>1</sup> *Synaptomys cooperi*.

<sup>2</sup> *Microtus pennsylvanicus*.

its species and varieties are so much alike that very few of them can be distinguished from the general mass.

The typical Field Mouse is a *short-eared, short-tailed, thick-set little animal*. It averages  $4\frac{1}{2}$  inches long, with a tail  $1\frac{1}{2}$  inches long. Its color above is reddish-brown, while beneath it is whitish-gray.

It is found from the Atlantic coast to the Dakotas, feeding on roots and grasses.

In severe winters, when the ground remains frozen for a long period, Field Mice are sometimes forced to feed on bark, and frequently kill

**The Red-Backed Mouse**<sup>1</sup> is, in form, very much like the meadow mouse, but in size it is smaller, and in habit it is quite different. It prefers to live in cool, damp woods and timbered regions, varying all the way from dark swamps and valleys to timbered mountain-tops; but it is seldom found in open country.

They are found from Ontario, New England and New Jersey westward to California, and northward through Canada and Alaska, sixteen species and five subspecies. They are all very much alike, rather slender, and more graceful in form than the field mice, and the majority



GAPPER'S RED-BACKED MOUSE.



NORTHWESTERN VOLE.

young fruit trees by barking them near the surface of the snow. When shocks of corn are available these mice live high, literally, feeding well, and being well housed at the same time. In husking shock corn in winter, many a nestful of Field Mice have we helped to turn out into the cold world; but the amount of grain they consumed was so insignificant we never grudged them their food.

Taken as a whole, the Field Mice of various species inhabit nearly the whole of North America north of Mexico and the Gulf, even to the remote islands of Bering Sea. I do not know of a state or province from which they have not been recorded.

are reddish-brown above and grayish underneath. The species most common in the eastern United States, often called **Gapper's Field Mouse**, is found westward to the Rocky Mountains. It is  $3\frac{3}{4}$  inches in length of head and body, tail,  $1\frac{3}{4}$  inches. In scientific lists of the mammals of North America, Red-Backed Mice are sometimes called Red-Backed "Voles."

**The Voles** of the genus *Phe-nae'o-mys*, are small brown mice, mostly of recent discovery, about the size of the red-backed mouse, in color

<sup>1</sup> Until recently this species has been considered identical with *Evotomys rutilus* of the Old World, and has been so called. Now, however, our species is considered quite distinct, and is called *E. gapperi*.

usually dark brown mixed with black. Seven species are known, extending in range from Labrador westward to Oregon, Washington and northern British Columbia, and also down to Colorado. None are found in the eastern half of the United States. There is no special mark by which it is easy to distinguish them from their nearest relatives, the red-backed mice.

The species most widely distributed, and best known, is the **Northwestern Vole**,<sup>1</sup> the largest member of this group,—a grayish-brown creature, with feet and all under parts white, or nearly so. It inhabits Alberta, British Columbia, Idaho, Wyoming, Washington and south central Oregon. Length of head and body, 4 inches, tail, 1 $\frac{3}{4}$  inches.

In mental capacity the **Wood Rat**, **Pack Rat**, **Trading Rat** or **Bushy-Tailed Rat**<sup>2</sup> is the most wonderful member of the whole Rat-and-Mouse Family, at least in North America. The true stories of its pranks are almost beyond belief. Seemingly its chief object in life is to play practical jokes on mankind; and any rat which manifests a spirit of toleration toward man surely is entitled to special consideration.

The typical Wood Rat is a large-sized, big-eyed, large-eared and rather handsome creature,



FLORIDA WOOD RAT.

without the mean, vicious look of a common rat, with fine yellowish-gray fur, white feet, and white under parts. In some species, the tail is covered with long hair, and by this fact alone it is possible to distinguish many members of the genus. The Wood Rats are distributed very

<sup>1</sup> *Phenacomys orophilus*.

<sup>2</sup> *Neotoma*.

generally throughout the southern and western part of the United States, and are also found in British Columbia and Mexico. Frequently their presence is indicated by the huge, mound-like nests, from two to three feet high, which they build of twigs, grass, leaves and bark.

These animals are nocturnal in their habits, and their nest-building and other work is done at night. The most remarkable thing about them is their habit of entering houses and playing practical jokes upon the inmates. A pair of Wood Rats that I knew by reputation at Oak Lodge, in Florida, first carried a lot of watermelon seeds from the ground floor upstairs, and hid them under a pillow. Then they took from the kitchen a tablespoonful of cucumber seeds, and placed them in the pocket of a vest which hung upstairs on a nail. In one night they removed from a box eighty-five pieces of bee-hive fixtures, and hid them in another box, and on the following night they deposited in the first box about two quarts of corn and oats.

Western frontiersmen, and others who live in the land of the Wood Rat, relate stories innumerable of the absurd but industrious doings of these strange creatures. In general they are rather harmless. One of the best known species is the **Florida Wood Rat**.<sup>3</sup> It belongs to the round-tailed group and does not have the hairy, squirrel-like tail of some of the western wood rats. Its upper color is tan mixed with brown, feet and under parts white. The length of the head and body is 8 $\frac{1}{2}$  inches, tail 6 $\frac{3}{4}$  inches. Distribution: the southern states from the Carolinas to Texas.

The **Little Harvest Mouse** looks so much like a small house mouse, 2 $\frac{1}{2}$  + 2 inches long, that only an expert can readily recognize it at first sight. The ten or more species are scattered throughout the southern, southwestern and Pacific states, but none of them are found in northeastern North America. The usual color is gray-brown above, and lighter underneath, and the best known example is **Le Conte's Harvest Mouse**<sup>4</sup> of the south Atlantic states, from Virginia to Florida.

The **Rice-Field Mouse**<sup>5</sup> should have been called a rat, for it is 5 inches long, with a five-inch tail. It is strictly a southern animal, in-

<sup>3</sup> *Neotoma floridana*. <sup>4</sup> *Reithrodontomys lecontei*.

<sup>5</sup> *Oryzomys palustris*.


habiting the wet rice-fields and swamps of the Gulf states from Texas up to southern New Jersey, its northern limit. It has a long head, a sharp nose, a shapely body, prominent ears, and a long tail. Its color above is bleached brown, but its under surface is grayish, or dull white.

This mouse is partial to the vicinity of water, especially the banks of rice-fields. It swims and dives well, and sometimes builds its nest and rears its young in interlaced marsh grass, over water, and far from dry ground.

**The Cotton Rat, or Marsh Rat,**<sup>1</sup> is a species

homa, New Mexico, Arizona, and Mexico; and wherever found their destructiveness causes them to be cordially disliked.

**The White-Footed Mouse, or Deer Mouse,**<sup>2</sup> is well worthy of acquaintance. It is distributed over nearly the whole of upper North America, except the arctic islands, and the Barren Grounds. On account of the changes it has undergone, chiefly in color shades, and length of tail, naturalists now recognize in the United States and Canada about *seventy species and subspecies!* But the student need not be discouraged by this fact.



RICE-FIELD MOUSE.

COTTON RAT.

which any country may well be without. It is small for a rat, but courageous, vicious in temper and voracious in appetite. It is fond of flesh, and when several are caged together, the stronger ones do not scruple to kill and eat weaker rats of their own kind. In length it is the size of a large chipmunk, 6 + 4 inches. The upper surface of the body and head, and outside of the legs, are dark mottled yellowish-brown, the under surface and inside of legs dull white, or brownish-gray. Cotton Rats are found from North Carolina to southern Florida, and also in Texas, Okla-

<sup>1</sup> *Sigmodon hispidus*.

Every White-Footed Mouse can be recognized by the clean white or light gray color on the under half of its body, head, tail and inner surfaces of the legs, its white feet, and its long tail. The color of the back is usually gray, or brown, or a mixture of the two.

Of all the small mice of North America, I consider this the most beautiful, and one of the most interesting. In the eastern states, where small quadrupeds and birds are numerous, it attracts little attention, but on the western plains, and in the desert regions, where animal life is very

<sup>2</sup> *Peromyscus leucopus*.

scarce (and rapidly becoming more so!) these pretty little creatures seem much more worthy of notice. I have many times found them nesting in cavernous and ill-smelling buffalo carcasses, and in the brain cavity or between the jaws of buffalo skulls from which the skin had not been removed by the hide-hunters.

In some places I have lain awake at night to hate mice, for cause, and wish them all dead, by all manner of violent deaths; but on a bleak and wind-shaven Montana plain where the bleaching skulls of thousands of slaughtered buffalo lie

elled over smoothly-shaven prairie divides miles away from all proper shelter. In the West, however, they are found most frequently in the brush and timber of stream valleys, where the rank weeds and grasses produce seed on which they feed. In the eastern United States they are found in nearly all agricultural regions. They are active climbers, possess a wide range of intelligence, and nest in all sorts of places, from ground burrows up to hollows in trees twenty feet from the ground. Of all mice, they are probably the most active climbers, and in fleeing



1. WHITE-FOOTED MOUSE.

2. LE CONTE'S HARVEST MOUSE.

MOLE MOUSE.

staring heavenward in mute protest against man's inhumanity, an agile White-Footed Mouse, scurrying out of its warm nest of buffalo-hair between the jaws of a buffalo skull, appeals not in vain for my sympathy and protection. Out on the Great Plains the world always seems large enough to contain us both. The great buffalo range of 1883 is now so barren of wild life that to-day even wild mice are objects of interest. From the buffalo to the White-Footed Mouse the time has been less than twenty years.

Many times in their wanderings from one buffalo carcass to another, these mice have trav-

eled over smoothly-shaven prairie divides miles away from all proper shelter. In the West, however, they are found most frequently in the brush and timber of stream valleys, where the rank weeds and grasses produce seed on which they feed. In the eastern United States they are found in nearly all agricultural regions. They are active climbers, possess a wide range of intelligence, and nest in all sorts of places, from ground burrows up to hollows in trees twenty feet from the ground. Of all mice, they are probably the most active climbers, and in fleeing

from a disturbed home the mother often carries her brood of young clinging to her body. Their food is seeds, small nuts and acorns, grain, and dried meat when available. Once in the wilds of Montana, we hauled some old logs to camp, for fire-wood. When one was cut up, we found in it a nest, made chiefly of feathers, containing five White-Footed Mice, snugly housed in the hollow. Packed close against the nest was a pint and a half of fine, clean seed, like radish seed, from some weed of the Pulse Family. While the food-store was being examined, and finally deposited in a pile upon the open ground, near the tent door, the five

mice escaped into the sage-brush. Near by stood an old-fashioned buggy.

Next morning, when the photographer lifted the cushion of his buggy-seat, and opened the top of the shallow box underneath, the five mice, with their heads together in a droll-looking group, looked out at him in surprise and curiosity, without attempting to run away. But very soon it became our turn to be surprised.

We found that those industrious little creatures had gathered up every particle of their nest, and every seed of their winter store, and carried all of it up into the seat of that buggy! The nest had been carefully re-made, and the seed placed closed by, as before. Considering the number of journeys that must have been necessary to carry all those materials over the ground, and climb up to the buggy-seat, the industry and agility of the mice were amazing.

By way of experiment, we again removed the nest, and while the mice once more took to the sage-brush, we collected all the seed, and poured it in a pile upon the ground, as before. During the following night, those indomitable little creatures *again* carried nest and seed back into the buggy-seat, just as before. Then we gathered up the entire family of mice with their nest and seed, and transported them to New York.

**The Grasshopper Mouse,**<sup>1</sup> originally described by Audubon and Bachman as the **Missouri Mouse**, and often called the **Mole Mouse**, is mentioned in order to caution western observers against confusing it with the preceding species. In some respects it strongly resembles the white-footed mouse, being all white underneath, including its legs. It can readily be distinguished by its large fore claws and its *short, stumpy tail*, which is only about one-third as long as the head and body. Its upper surface is brownish-gray. *Its fur is very fine and soft*, and hence it is sometimes called the Mole Mouse. Its length, head and body, is  $4\frac{1}{2}$  inches, tail,  $1\frac{3}{4}$  inches.

## CHEEK-POUCH MICE AND RATS.

### *Heteromyidae.*

This is strictly a Family of the West and Southwest, its members being found only west of Arkansas, Iowa and Minnesota. It does *not* contain the pocket gophers. Many of its twenty-six spe-

<sup>1</sup> *Onychomys leucogaster.*

cies are desert dwellers, even inhabiting Death Valley, California. All its members are distinguished from other North American animals (except the jumping mouse and pocket gopher) by the presence of a large and very serviceable hair-lined pouch in the skin of each cheek. Barring the two exceptions noted, this character alone is sufficient for the recognition of any American member of this Family.

As clearly shown in the full-page diagram, this family may be divided into two Subfamilies, an arrangement which is very convenient and helpful. The first we must call the **Pocket Mouse** Subfamily and its leading genus (*Perognathus*) contains twenty-six full species, and fifteen sub-species. All are distinguished by the following characters: head large; body slender and graceful; *hind legs long, and fitted for jumping*; *tail long*; *large external cheek pouches, hairy inside, and not connected with the interior of the mouth*; hair smooth and compact, sometimes intermingled with spines. These mice are quick and active in movement, and some species leap with considerable power.



KANGAROO RAT.  
Showing the very large cheek-pouches.

Since 1839 the **Typical Pocket Mouse**<sup>2</sup> has been described again and again, but none of its describers have taken the trouble to give it an English name! Hereafter, let us call it by the name given above, because it is the type of its genus. It inhabits Montana, Wyoming and the Dakotas. Its color above is sandy-yellowish, lined with black; underneath, white; and these two color areas are divided low down along the side by a lengthwise band of pale yellow. Length,  $3 + 2\frac{3}{8}$  inches.

**The Kangaroo Rat**<sup>3</sup> Subfamily, of fifteen full species, is fitly represented by an elf-like creature which is one of the most beautiful and at-

<sup>2</sup> *Perognathus fasciatus.*

<sup>3</sup> Typical species, *Perodipus richardsoni* of western Kansas, Oklahoma and Indian Territory.

tractive of all our native rats. In the dry and sterile regions of the great Southwest, from the Indian Territory to Arizona and California, where seemingly the deserts produce nothing but sand, cacti, yuccas and sage-brush, these pert little creatures hold forth. Apparently they are both fire-proof and water-proof, for no amount of heat affects them, and the absence of water does not seem to depress their spirits in the least. Like most mice and rats, they are nocturnal. Some of the species build for themselves large mounds of dirt and gravel, from one to three feet high and five to ten feet in diameter, which are honeycombed with burrows



1. AND 2, KANGAROO RAT.  
3. TYPICAL POCKET MOUSE.

and runways. These dwellings are often inhabited by rattlesnakes and lizards, and doubtless the Kangaroo Rat is an important item of food in the diet of the desert rattler.

The Kangaroo Rat is very unlike the members of the Mouse-and-Rat Family; and in temper no creature could be more unlike the domestic rat. Unlike most mice and rats, they do not bite when handled, but they are so delicate that they do not live long in captivity, unless tended with extreme care and intelligence. They stand high on their hind legs, like pigmy kangaroos, and hop about with their front paws tucked up close under the chin, almost hidden by their fur. The tail is very long, has a showy tuft of long hair on the end, and is used by the animal in

balancing itself when in motion. The fur is soft, silky, rather long, and of a tawny-brown color above. Length of head and body,  $4\frac{1}{2}$  inches, tail,  $5\frac{3}{4}$  inches. The cheek-pouches are large, and are of great use in carrying sand out of burrows.

### JUMPING MOUSE FAMILY.

#### *Zapodidae.*

The **Jumping Mouse**<sup>1</sup> is one of the most remarkable of all our small animals. In form it is a *slender-bodied mouse*, with an *exceedingly long tail*, *kangaroo-like hind legs*, and *cheek-pouches*. Its average length of head and body is about 3 inches, and tail 5 inches. In color it is dark reddish-brown above, white underneath, with smooth compact hair. Although no larger than a house mouse, it can jump from eight to ten feet.

When a farmer boy is hauling in sheaves of wheat, and a small animal suddenly makes a tremendous flying leap from the bottom of the shock, he may know that he has disturbed a Jumping Mouse, and the chances are that he cannot capture it by hand. In these long jumps—perhaps the longest on record for an animal of equal size—the tail is as necessary as a stick is to a sky-rocket, to enable the little creature to preserve its balance, and go straight ahead. If the tail is cut off, the Jumping Mouse turns over and over in the air, and perhaps lands upon its back.

The Jumping Mouse is quite nocturnal in its habits, and is seldom seen in the daytime. It feeds on seeds and grain, and while it devours great quantities of weed seeds, it inflicts upon the farmer no damage worthy of mention. In the autumn it stores in the ground quantities of food for winter use, but despite this fact, under certain conditions it becomes so thoroughly dormant in winter that it seems to be quite lifeless. It is found throughout the northern United States and Canada, in wooded regions, from New York to California, and as far north as Lake Nushagak, Alaska.

### Opinions Regarding Rats and Rat-like Animals.

The largest rat-like animal in America is the **Coy'pu Rat**,<sup>2</sup> of Central and South America, which stands 9 inches high at the shoulders, attains a length of 19 inches head and body, tail,

<sup>1</sup> *Zapus hudsonius.*    <sup>2</sup> *Myo-cas'-tor coy'pus.*



13 inches, and weighs 8 pounds. It is a water-loving animal, almost as much so as the muskrat, and its thick, brown fur is valuable. Under proper conditions it is easily kept in captivity.

*The smallest rodent in America is the Least Pocket Mouse,<sup>1</sup> of the Rocky Mountain region, which has a total length of head and body, 1½ inches; tail, 2¾ inches.*

*The best swimmer of all rat-like animals is the Muskrat.<sup>2</sup>*

*The best climber is the Tree Rat,<sup>3</sup> of southern India.*

*The handsomest rat or mouse in the New World is the Kangaroo Rat, of the southwestern United States, figured on the opposite page.*

*The most humorous of all rat-like animals is the Trading Rat, described on page 89, which delights in playing practical jokes upon its human neighbors.*

*The meanest of all rodents is the brown-coated Domestic Rat, the pest of civilization everywhere, which was sent to man as a perpetual punishment for his crimes against harmless wild creatures all over the world.*

### THE POCKET GOPHER FAMILY.

#### *Geomysidae.*

**The Red Pocket Gopher<sup>4</sup>** is the most important representative of a large Family of burrowing rodents which does great damage to the crops and lands of American farmers. Whenever you see a brown-coated burrowing animal, the length of a small rat, but twice as thick, with a big pouch in the skin of each cheek, a swinish appetite, a set of long claws like burglar's tools on each fore foot and a most villainous countenance and temper, you may know that it is a Pocket Gopher. The pockets in his cheeks are to enable him to carry extra large quantities of stolen potatoes and seeds. When once you have learned the true character and habits of this creature, you will, without being asked, carefully refrain from calling any ground-squirrel a "Gopher."

Most wild animals have some redeeming qualities, but this cannot make good a claim to one. Gophers are not only thieves and robbers, but they are so ill-tempered that they even hate each other, and the old ones usually are found living

alone. When two captives are placed together, they usually fight fiercely until one is killed. Their teeth and front claws are very powerful, and working together they do great damage, in many different ways.

As a Family, Pocket Gophers inhabit the whole United States west of Indiana and the lower Mississippi, and also a large part of Alabama,



JUMPING MOUSE.

Georgia and Florida. Three genera and about thirty-three species are recognized, and while some are smaller than others, and some are gray or black instead of brown, their appetites and habits are all equally objectionable. They spoil meadows by throwing up innumerable hillocks of loose earth; they devour great quantities of vegetable crops, and also corn and small grain; they eat the roots of young fruit-trees of nearly all kinds, and they destroy canals and irrigating ditches by honeycombing their banks. With incisor teeth that in sharpness and strength are like steel chisels, a Gopher can pare off all the roots from a young tree quite as neatly as a man pares potatoes.

Our type species, the Red Pocket Gopher "is," says Mr. Vernon Bailey, "of much greater economic importance than all the other species combined, for the reason that its home is in the fertile prairie region of the Mississippi valley,"

<sup>1</sup> *Perognathus flavus.*    <sup>2</sup> *Fiber zibethicus.*  
<sup>3</sup> *Mus rupestris.*        <sup>4</sup> *Geomys burbanki.*

embracing Iowa,—which is its centre of distribution,—Illinois, Missouri, Wisconsin, Minnesota, and the eastern parts of the Dakotas, Nebraska and Kansas. Its length is about  $7\frac{1}{2} + 3$  inches. The young are either two or three in number, and there is only one litter each year.

The enemies of the Gopher are the weasel and the gopher snake.<sup>1</sup> Because of the damage done by Gophers, farmers generally wage war upon them with traps, strychnine, and poisoned grain. In Iowa, Minnesota and other states, many thousands of dollars have been paid out by county treasurers in bounties on Gopher scalps and tails. No animal in the West is more universally disliked, nor more diligently destroyed.

My acquaintance with the Gopher Family began when I was a farmer boy, in Iowa, the storm centre of *Ge'o-mys bur-sa'ri-us*. Having



RED POCKET GOPHER.

trapped a few, I made the mistake of supposing that I knew more about the habits of those creatures than did my elders, who had not. In an evil moment, I announced that any strong boy could catch a Gopher by digging it out of its burrow, and my large brother offered me twenty-five cents if I could prove that claim within a week.

That evening, with mattock and spade, I repaired to my father's corn-field, into which strange Gophers were rapidly migrating and settling; and finding a fresh hole with the owner inside, I began to dig. My shepherd dog, Rover, assisted me all he could, chiefly by keeping me company, but also by digging when I rested.

We dug into the twilight, and later on we dug into the night; but the Gopher kept well ahead of us. Whenever we paused to listen, we could

<sup>1</sup> *Pituophis*.

hear him digging hard, and to our dismay we found that he knew a thing or two about getting on in the world. With the descent of black darkness, our hopes of overtaking that Gopher descended also; and then pride, not hope of reward, was all that spurred us on. *Would* we have to give up beaten, by an ugly, pig-eyed old Gopher?

When for about the thirtieth time I paused to wipe the accumulation of perspiration and prairie loam from my brow, Rover suddenly rushed off into the darkness. In the corn-rows thirty yards away, he seized something, shook it vigorously, and a moment later came trotting back to me, carrying in his mouth a large Gopher! The beast had been migrating into the corn-field, and Rover simply caught him on the fly.

Digging operations ceased abruptly at that point. Thanking Rover for his timely assistance, I accepted his contribution, and we marched home together. When I exhibited to my brother the Gopher that we had secured "by digging," he was profoundly surprised, but promptly paid the money. Rover looked on smilingly, and said not a word; but we both knew then that in catching Gophers, steel traps are better than spades.

## THE PORCUPINE FAMILY.

### *Erethizontidae.*

The Porcupine is at home either in tree-tops or on the ground, but it is always a slow-moving and dull-witted animal. It is easily captured or killed by man, but not so readily overcome by wild animals. In the woods, it loves to prowl around camps, and eat every scrap of leather or greasy board that it can find. It is fond of the bark of hemlock, beech and cottonwood, and often a Porcupine will remain in a good tree until he entirely strips it of its bark.

The **Canada Porcupine**,<sup>2</sup> which is black, with a gray-tipped storm-coat, is found in New England, New York, Pennsylvania, Ohio and thence northward and northward to Fort Churchill on Hudson Bay. The West and Northwest is the home of another species, known as the **Yellow-Haired Porcupine**.<sup>3</sup> Large specimens weigh from 25 to 30 pounds. The flesh is not palatable to white men, but is eaten by Indians.

The Canada Porcupine *never* should be called  
<sup>2</sup> *Er-c-thi'zon dor-sa'tus*.      <sup>3</sup> *E. cp-i-xan'thus*.

a "Hedgehog," because the latter is not a gnawing animal, but a small, weak, insect-eater, which does not inhabit America. A full-grown Porcupine is about twenty times as large as the common European hedgehog.

Porcupines can *not* shoot their quills, not even for one inch; and the idea that they can—or ever have—is entirely erroneous. When attacked, their defence consists in erecting their quills, and striking quickly a strong sidewise blow with the tail, which often drives many quills into its enemy. Strange to say, wild animals are about as lacking in original information, or "instinct," regarding this creature as dogs are. Several pumas and lynxes have been killed in a starving condition, with their mouths and throats so filled with porcupine quills that eating had become almost impossible.

**THE PIKA FAMILY.**

*Ochotonidae.*

The **Pika**, commonly called the **Little Chief "Hare,"** or **Crying "Hare,"**<sup>1</sup> looks very much like a small, gray-brown rabbit, 7 inches long; but it is neither a rabbit nor a hare, and represents an independent Family. It lives high up on the great mountain ranges of the West, from just below timber line up to the line of perpetual snow. It finds shelter in the crevices of rugged masses of rock, and its sharp little cries often seem to come from so many different points that the hunter is completely confused. In form this strange little creature is about half way between a gray rabbit and a guinea-pig; and it has neither speed nor activity.

**THE HARE AND RABBIT FAMILY.**

*Leporidae.*

This group is very clearly subdivided and there need be no confusion of ideas regarding its North American members. Nevertheless, early writers have made a confusing error in the improper adoption, for one important group, of the misleading name Jack "Rabbit." It should be Jack *Hare*.

All the American members of this Family are separated into two general groups, the *Hares* and the *Rabbits*. The accompanying diagram shows these subdivisions, and their relations to each other.

<sup>1</sup> *O-cho-to'na prin'ceps.*

A typical **Hare** is *big, long-eared, long-legged,* and a swift runner. Very often its color changes according to the season. It does not burrow, but rears its young in a nest or "form."

The **Rabbit** is *small, short-eared, short-legged,* a weak runner for a long distance, its color is fairly constant, and it lives in a burrow.

The **Varying Hare Group** is the key to the entire Family; or, in other words, it stands on middle ground between the Rabbits, the Polar Hare, and the Jack Hare, and is related to all three. Naturally this group should be studied



SANBORN, Photo., N. Y. Zoological Park.

CANADA PORCUPINE.

first. Its type species is the **Northern Varying Hare,**<sup>2</sup> of northern New York, New England, Canada and the Northwest Territories. Its name is due to the fact that its color varies according to the season, being *pale cinnamon brown in summer,* and *white in winter,* with only a narrow back line of brown.

It is nearly twice as large as the cotton-tail rabbit, but its ears and legs are about half way in proportionate length between those of the

<sup>2</sup> *Le'pus a-mer-i-can'us.*

common rabbit, and the jack hare of the Southwest. Large male specimens measure 18 inches in length of head and body, tail, 2 inches, and weigh 6 pounds.

Like the true fur-bearing animals, Varying Hares have two kinds of fur,—a dense, fine and soft under fur through which grows a storm-coat of thin, coarse, straight hair. It is the latter which gives an animal its color. In the summer these long hairs are black, but as winter approaches they turn white.

The habits of the Varying Hares and Rabbits are so nearly the same that it is unnecessary to describe them separately. They all require brushy ground, broken rocks, rugged ravines or tree-holes in which to hide from the foxes, dogs, men, mink, martens, lynxes, skunks and birds of prey which constantly hunt them as food. But for their keenness of sight, hearing and scent, their swiftness in running to cover, and their marvellous agility in doubling and turning when pursued, their numerous enemies would soon exterminate them.

**The Polar Hare**<sup>1</sup> is the most northern species of this Family. Colonel Brainard found its tracks at 83° 24', which for fifteen years remained man's "farthest North." In the southern portion of its home, this hare is gray and white in summer, but in the higher polar regions it is white all the year round, like the majority of true arctic animals,—the owl, fox, bear and wolf.

**The Prairie Hare**<sup>2</sup> of the western plains is generally supposed to be of the same species as the so-called jack "rabbit" of the Southwest; but it is not. In form, size and color, it may be considered a connecting link between the varying hare group and the jack hare group, and its separate identity should be remembered. Its home is the great sage-brush plains of the Northwest, from Kansas to the Saskatchewan, and westward to Oregon, and northern California. It is gray in summer, but changes to white in winter. It is a large species (23 inches long), with ears longer than its head, long, strong hind legs, and a white tail unmarked with black, a character by which it can be readily distinguished from other jack "rabbits."

On the treeless plains of the great West, where it is often seen without any other objects to furnish comparisons, it sometimes seems of immense

size, and a Prairie Hare 200 yards away has often been mistaken for an antelope supposed to be 600 yards distant.

**The Jack Hare**<sup>3</sup> (commonly called **Jack "Rabbit"**) is easily recognized by his extremely large ears,—five to six inches long,—slender body, long legs and athletic build, and the black mark on the upper surface of the tail. There are seven species, all very much alike, which inhabit the southwestern quarter of the United States, extend northward to Oregon, eastward to Nebraska and Kansas and southward to Tehuantepec, Mexico. In many localities wherein wolves and foxes have been exterminated, these hares have multiplied until they have become a great pest. In several localities in California, and also in eastern Colorado, great rabbit-drives are made, in which many thousand Jacks are slaughtered, and given away in large cities for food.

The Jack Hare is a very swift runner. In eastern Kansas, Professor L. L. Dyche once saw a good greyhound chase a Jack on fair ground for about two and a half miles, and in the whole distance the hound gained only about twenty-five yards. The hare finally escaped by running into a *hollow log* that had been left on the prairie by accident, and was the only shelter within five miles!

**The Gray Rabbit, or Cotton-Tail**,<sup>4</sup> is a typical representative of the Rabbit Family, which contains twelve species. Throughout the extensive region which forms its home,—from New England and Minnesota to Yucatan,—it refuses to be exterminated, and is perhaps more frequently seen and more widely known than any other quadruped.

All the true rabbits are small, and for long running their legs are short and weak; but what they lack in endurance they make up in cunning and quickness. To aid in their preservation, Nature has given them colors that blend so perfectly with their surroundings that a rabbit crouching low often is compelled to run to avoid being trodden upon. When hard pressed for a nesting place in a city, a Gray Rabbit has been known to dig a shallow hole in the smooth lawn of the Smithsonian grounds at Washington, line it with her own fur, and rear her young in it, within forty feet of the National Museum building, and a busy roadway, without discovery by

<sup>1</sup> *Lepus arcticus*.

<sup>2</sup> *Lepus campestris*.

<sup>3</sup> *Lepus texianus*.

<sup>4</sup> *Lepus sylvaticus*.



**POLAR HARE**  
(*LEPUS ARTICUS*)

**POLAR HARE GROUP.**

LARGE. EARS SHORTER THAN HEAD: PELAGE PURE WHITE, OR BUT SLIGHTLY COLORED IN SUMMER.

ARCTIC REGIONS.  
6 SPECIES.

**VARYING HARE, (*LEPUS AMERICANUS*).**



**VARYING HARE GROUP.**

NORTHERN UNITED STATES AND CANADA.  
4 SPECIES.

**RABBIT GROUP.**

SMALL: COLORS CONSTANT; NEVER WHITE: LEGS SHORT WEAK RUNNERS. EARS SHORT.

UNITED STATES GENERALLY, MEXICO, AND SOUTH TO COSTA RICA.  
12 SPECIES



**PRAIRIE HARE.**

**JACK HARE (OR JACK RABBIT) GROUP.**

LARGE: SLENDER; LEGS LONG AND STRONG: SWIFT. EARS VERY LONG AND LARGE. COLORS CONSTANT.

S:W. UNITED STATES, PACIFIC STATES & MEXICO.  
7 SPECIES.

**COTTON-TAIL RABBIT, (*LEPUS SYLVATICUS*)**



**PRAIRIE HARE,**  
(*LEPUS CAMPESTRIS*),

LARGE: EARS LONGER THAN HEAD: HIND LEGS LONG AND STRONG: COLOR VARIABLE.  
N:W. UNITED STATES.



**JACK RABBIT,**  
(*LEPUS CALLOTIS*).

**DIAGRAM OF THE HARE AND RABBIT FAMILY IN NORTH AMERICA,**  
SHOWING THE RELATIONSHIPS AND COMPARATIVE SIZES OF THE FIVE GROUPS.

Copyright 1903 by W. T. Hornaday.

dogs or men until the mowers found the nest almost under their feet. Every year one or two pair breed in the adjoining grounds of the Department of Agriculture.

When a rabbit can have his choice of hiding-places, he chooses a burrow directly beneath a large tree, the roots of which render it difficult, or it may be impossible, for man or beast to dig him out. Crevices in rock ledges are equally good, but he often finds that hollow logs, hollow trees and brush piles only lead to swift destruction.

He never sleeps in daylight, when enemies are afoot. If the Man-With-a-Gun approaches, he crouches low and lies as still as a stuffed rabbit, breathing seldom, winking never, but with legs all ready to spring. His keen eyes and ears measure every yard of his enemy's approach, until the dead line is crossed when—Zip! Out flashes a long, gray streak,—flying over logs, and darting through openings so swiftly that in two or three seconds a snow-white signal flag waves an adieu, and disappears.

In summer hares and rabbits feed on green twigs, soft bark, buds, grass, leaves and berries. In winter they are forced to subsist chiefly on the bark of bushes and the berries of the wild rose. Whenever they gnaw the bark from young fruit-trees, it is a sign that they are hard pressed for food.

Rabbits breed very rapidly, often raising three litters a year, and if not kept in check by birds of prey and carnivorous animals, would soon be-

come altogether too numerous. In Australia and New Zealand, the rabbits "introduced"<sup>1</sup> from Europe have multiplied until they have become a fearful scourge, and are now so numerous it is impossible even to keep them in check. Possibly the use of their flesh as food, and their skins as "fur," may lead to an abatement of the evil. The moral of the rabbit in Australia, the mongoose in the West Indies, and the English sparrow in America, is, before "introducing" a foreign species of bird or mammal into America, take expert advice,—and then don't do it! This refers to species able to live wholly by their own efforts when imported and set free.

### Bibliographical.

The following popular papers are of special interest and value:

*On Jack Rabbits.*—The Jack Rabbits of the United States. By Dr. T. S. Palmer; pamphlet, 88 pages. Bulletin No. 8, Biological Survey, Department of Agriculture. Washington, 1897.

*On Gophers.*—The Pocket Gophers of the United States. By Vernon Bailey; pamphlet, 47 pages. Bulletin No. 5, as above, 1895.

*On Prairie-Dogs.*—The Prairie-Dog of the Great Plains. By Dr. C. Hart Merriam; pamphlet, 14 pages. Yearbook of the Department of Agriculture, 1901.

*On Ground Squirrels.*—The Prairie Ground Squirrels of the Mississippi Valley. By Vernon Bailey; pamphlet, 69 pages. Bulletin No. 4, Biological Survey, Department of Agriculture, 1893.

<sup>1</sup> A species transplanted from one country to another is said to be "introduced."

## CHAPTER VIII

# THE ORDER OF HOOFED ANIMALS

### UNGULATA

The Order which includes the hoofed animals of the world is called Un-gu-la'ta, a Latin word which means "hoofed." In North America, it is represented by a great variety of forms, several of which are of special importance.

Before seeking to become acquainted with these animals, the student must pause long enough to gain a bird's-eye view of the groups into which they are divided, and thereby understand their relationships, clearly and correctly.

The following diagram of arrangement is very simple, and the animals it sets forth are in some respects the most important in America.

### THE GROUPS OF NORTH AMERICAN HOOFED ANIMALS.

	FAMILIES.	GROUPS.	EXAMPLES.	
<b>ORDER UNGULATA.</b> HOOFED ANIMALS (Of North America only).	CATTLE AND SHEEP FAMILY, or BOVIDAE:	Cattle: {	<b>American Bison :</b> <b>Buffalo,</b>	<i>Bos americanus.</i>
		Sheep-Cattle: {	<b>Musk-Ox,</b>	<i>Ovibos moschatus.</i>
		Sheep: {	<b>Big-Horn,</b> <b>White Sheep,</b> <b>Black Sheep,</b>	<i>Ovis canadensis.</i> <i>Ovis dalli.</i> <i>Ovis stonoi.</i>
		Goat: .	<b>Mountain Goat,</b>	<i>Oreamnos montanus.</i>
	ANTELOPE FAMILY, or ANTILOCAPRIDAE:	. . .	<b>Prong-Horned Antelope,</b>	<i>Antilocapra americana.</i>
		DEER FAMILY, or CERVIDAE:	Round-Horned Groups: {	<b>Elk, or Wapiti,</b> <b>White-Tailed Deer,</b> <b>Mule Deer,</b> <b>Black-Tailed Deer,</b>
	Flat-Horned Groups: {		<b>Barren-Ground Caribou,</b> <b>Woodland Caribou,</b> <b>Moose,</b>	<i>Rangifer arcticus.</i> <i>Rangifer caribou.</i> <i>Alces americanus.</i>
	. . .		<b>Collared Peccary,</b>	<i>Tayassu tajacu.</i>
	PECCARY FAMILY, or TAYASSUIDAE:	. . .	<b>Dow's Tapir,</b>	<i>Tapirus dowi.</i>
		TAPIR FAMILY, or TAPIRIDAE:		

### THE CATTLE AND SHEEP FAMILY.

*Bo'vi-dae.*

**General Characters.**—The Cattle Family of the world contains a grand array of large animals, such as the wild cattle, bison, buffalo,

musk-ox, mountain sheep, ibex, and wild goats. There are about fifty species in all, scattered over all continents save South America and Australia. All the members of this Family have divided hoofs, and simple horns (i.e., not branching) consisting of a hollow sheath growing over


a pointed core of very porous bone. The horns grow until the animal reaches old age, and are never shed. If knocked off by accident, the new horn material presently covers the horn core, but never succeeds in forming a perfect weapon like the original. Such a growth is called a "crumpled" horn. The members of this Family eat vegetable food, preferably grass and herbage, and have no upper front teeth.

**The American Buffalo.**

**The American Bison or Buffalo.**<sup>1</sup>—Because of its great size, imposing appearance, former

complete extinction, by appropriating \$15,000 for the purpose of purchasing and establishing under fence in the Yellowstone Park, a herd of captive Buffaloes. This undertaking has very wisely and appropriately been placed in charge of the Department of Agriculture.

At this date (1903) there are about 634 wild Buffaloes alive, of which about 600 inhabit a desolate and inhospitable region southwest of Great Slave Lake. In 1890, the Yellowstone Park herd contained about three hundred head; but through inadequate protection, and killing done by unprincipled poachers in quest of



SKELETON OF AN ADULT MALE AMERICAN BISON.

abundance and value to mankind, this is the most celebrated of all American hoofed animals. Its practical extermination in a wild state is now a source of universal regret. In 1902, Congress took the first step toward its preservation from

<sup>1</sup> A true "Buffalo" is an animal with *no hump* on its shoulders; and is found only in Africa and Asia. Our animal, having a high hump, is really a *bison*; but inasmuch as it is known to seventy-three millions of Americans as the "Buffalo," it would be quite useless to attempt to bring about a universal change in its popular name. There is but one living species.

heads to sell, to-day *less than thirty buffaloes remain!* The weakness of the efforts to protect that herd is a national disgrace. Through lack of sufficient laws and patrol service the poachers were permitted to rob the American people of a wild herd which no expenditure of money ever can replace.

There were in captivity, in March, 1903, 1,119 pure-bred Buffaloes, and the number is slowly increasing. Of these, the majority are in large private game preserves, and every zoological



park and garden contains as many head as it can properly accommodate. It is useless to give a list of these animals, because owners and figures are constantly changing.

The Buffalo breeds readily in captivity, and is easily cared for. The majority of captive animals are reasonably tractable, but occasionally an individual becomes savage and dangerous, and requires either solitary confinement or

contains one hundred and twenty-eight head of pure-blood animals, and the number is steadily increasing. The largest herd on public exhibition is that of the New York Zoological Park, which in 1903 contained thirty-four head of pure-breed animals representing all ages, presented by the Hon. William C. Whitney from his October Mountain preserve.

The value of a full-grown Buffalo cow in New



E. R. SANBORN, Photo.

#### AMERICAN BISON, OR BUFFALO.

An adult male, "Apache," of the Whitney herd. Photographed in the New York Zoological Park, near the end of the shedding season.

shooting. The best place in which to exhibit a savage Buffalo is a museum. Full-grown males must be watched closely for signs of permanent ill temper, and a savage Buffalo should be treated the same as a tiger. Frequently the first serious sign of danger in a Buffalo is the murder of a weaker member of the herd.

The largest herd in a fenced game preserve is that of Blue Mountain Park, in New Hampshire, established by the late Austin Corbin. It

York is from \$400 to \$500, and an adult bull is worth about \$100 less. Exceptionally fine mounted heads are worth from \$300 to \$500.

The Buffalo was first seen by white men in Anahuac, the Aztec capital of Mexico, in 1521, when Cortez and his men paid their first visit to the menagerie of King Montezuma. In its wild state it was first seen in southern Texas, in 1530, by a ship-wrecked Spanish sailor. The Buffalo once roamed over fully one-third of the

entire continent of North America, and its numbers far exceeded those of any other large mammal of recent times.

Not only did it inhabit the plains of the West, but also the hilly hard-wood forests of the Appalachian region, the northern plains of Mexico, the "Great American Desert," the Rocky Mountain parks on the continental divide to an elevation of 11,000 feet, and the bleak and barren plains of western Canada, up to the land of the musk-ox. From north to south it ranged 3,600 miles, and from east to west about 2,000 miles.

The centre of abundance of the Buffalo was the Great Plains lying between the Rocky Mountains and the Mississippi valley. When the herds assembled there, they covered the earth seemingly as with one vast, brown buffalo-robe.

It is safe to say that no man ever saw in one day a greater panorama of animal life than that unrolled before Colonel R. I. Dodge, in May, 1871, when he drove for twenty-five miles along the Arkansas River, through an unbroken herd of Buffaloes. By my calculation, he actually saw on that memorable day nearly half a million head. It was the great southern herd, on its annual spring migration northward, and it must have contained a total of about three and one-half million animals. At that date, the northern herd contained about one and one-half millions. In those days, mighty hosts of Buffaloes frequently stopped or derailed railway trains, and obstructed the progress of boats on the Missouri and Yellowstone rivers.

In 1869, the general herd was divided, by the completion of the Union Pacific Railway, into a "northern herd" and "southern herd." The latter was savagely attacked by hide hunters in the autumn of 1871, and by 1875, with the exception of three very small bunches, it had been annihilated.

In 1880, the completion of the Northern Pacific Railway led to a grand attack upon the northern herd. In October, 1883, the last thousand head were killed in southwestern Dakota, by Sitting Bull and about a thousand Indians from the Standing Rock agency, leaving only the Yellowstone Park bunch of two hundred head, a band of forty in Custer County, Montana, and the Great Slave Lake herd of about five hundred head.

The largest Buffalo ever measured by a nat-

uralist is the old bull which was shot (by the author) on December 6, 1886, in Montana, and which now stands as the most prominent figure in the mounted group in the United States National Museum. A very good picture of him adorns the ten-dollar bill of our national currency. His dimensions in the flesh were as follows:

	Ft.	In.
Height at shoulders . . . . .	5	8
Length of head and body, to root of tail . . . . .	10	2
Depth of chest . . . . .	3	10
Girth, behind forelegs . . . . .	8	4
Circumference of muzzle, behind nostrils . . . . .	2	2
Length of tail vertebrae . . . . .	1	3
Length of hair on shoulders . . . . .	6	½
"    "    "    "    forehead . . . . .	1	4
"    "    "    "    chin beard . . . . .	11	½
Estimated weight . . . . .	2,100 pounds.	

The shoulder height of wild Buffaloes of various ages, and both sexes, as taken by me on the Montana buffalo range, are as follows:

	Ft.	In.
Male calf, 4 months old . . . . .	2	8
"    one year old . . . . .	3	5
"    two years old . . . . .	4	2
"    five years old (average size) . . . . .	5	6
Female, three years old . . . . .	4	5
"    eight years old . . . . .	4	10

The Buffalo begins to shed its faded and weather-beaten winter coat of hair in March, and during April, May and June it presents a forlorn appearance. The old hair hangs to the body like fluttering rags, and at last, when it finally disappears, the body is almost bare. At this time the flies are very troublesome. By October, the new coat is of good length and color, and in November and December, it is at its finest. The animal is then warmly clad for the worst storms of winter, and the shaggy head is so well protected that the animal faces all storms instead of drifting before them. A bull Buffalo in perfect pelage is an animal of really majestic presence, and is far more imposing in appearance than many animals of larger bulk, but less hair.

The calves are born in May and June, and at first are of a brick-red color. This coat is shed in October, except in calves born late in the season.

The flesh of the Buffalo so closely resembles domestic beef of the same age and quality that it

is impossible for any one to distinguish a difference.

The future of the Buffalo depends solely upon the owners of the great private game preserves, such as that of the late Austin Corbin, and Mr. James J. Hill. If the perpetuation of the species depended solely upon the efforts possible in zoological gardens and parks, within twenty-five years the species would become extinct. Even in a range of twenty acres, the largest in any zoological institution, the Buffalo becomes a sluggish animal, and rapidly deteriorates from the vigorous standard of the wild or semi-wild stock. In the close confinement of a thirty-acre zoological garden, the loss in physique is still greater. Mr. Arthur E. Brown, Superintendent of the Philadelphia Zoological Garden, and a very close observer, has drawn the writer's attention to the striking difference in size and back outline between a Buffalo born on a great range, and another of the same age born of a line of closely confined ancestors.

Interesting as have been the experiments made by Mr. C. J. Jones and others in the cross-breeding of Buffaloes and domestic cattle, it is now quite time that all such experiments should cease. It has been proven conclusively that it is impossible to introduce and maintain a tangible strain of buffalo blood into the mass of western range cattle. This is admitted with great regret, but inasmuch as it is absolutely true, the existing herds of Buffalo should not be further vitiated and degraded by the presence in them of animals of impure blood.

In an adult animal, the presence of domestic blood is readily perceived in the lower hump, longer tail, shorter pelage on the head, neck, shoulders and fore legs, and the longer and more slender horns. In the calf under one year of age, it is not always possible for even the best judges to detect a strain of domestic blood. In the year 1900, a male calf was inspected and passed by four men who were with good reason considered qualified judges of the points of Buffaloes; but two years later that animal stood forth unmistakably as a cross-breed, one-quarter domestic.

In judging Buffaloes, the finest animals are those with the greatest height of hump, heaviest and longest pelage in front of the armpit, shortest tails, and horns curving with the shortest radius.

If the recent action of the national government

toward establishing a herd in the Yellowstone Park is liberally and intelligently sustained by future administrations, it will go far toward perpetuating the species for a century. But it should be conceded at the beginning that the effort can succeed only by giving the animals a great area to roam over at will. In addition to that herd, however, another should immediately be established in the Plains region, in a fenced reservation of not less than 100 square miles, with choice grazing, water and ravine shelters. It is only by such methods that the American people can in a small measure atone for the annihilation of the great herds between 1870 and 1885, and the subsequent brutal slaughter by poachers of the Yellowstone Park herd of three hundred head.

On March 1, 1903, Dr. Frank Baker completed a count of all the pure-blood captive Buffaloes alive at that date, with the following result:

*Captive Buffaloes:*

In the United States.....	969
In Canada.....	41
In Europe.....	109
	<hr/>
	1,119

*Wild Buffaloes (estimated):*

In the United States.....	34
In Canada.....	600
	<hr/>
	634
	<hr/>
	1,753

**The Musk-Ox.**

**The Musk-Ox**<sup>1</sup> is an inhabitant of the frozen North, the land of snow and ice, of howling storms and treeless desolation. In 1901, Commander Peary killed a specimen within half a mile of the most northerly point of land in the world,—the northeastern extremity of Greenland.

How this animal finds food of any kind during the dark and terrible arctic winter, is yet one of the secrets of Nature. After making all possible allowance for the grass, willow and saxifrage obtainable by pawing through the snow, and on ridge-crests that are swept bare by the blizzards, it is still impossible to explain how the Musk-Ox herds find sufficient food in winter, not only to sustain life, but actually to be well-fed.

I gaze upon each living Musk-Ox to be seen

<sup>1</sup> *O'vi-bos mos-cha'tus.*

in captivity with a feeling of wonder, as if it were a creature from another world. There are times, also, when I wonder whether many of the visitors who see them quietly munching their clover hay, appreciate the effort that has been put forth to capture them in the remote and desolate regions of the far North, keep them alive, and bring them to civilization for public exhibition.

The Musk-Ox is one of the strangest of all our large animals, and its appearance is so odd and striking that when once seen by an observant person it is not easily forgotten. In it one sees an oblong mass of very long and wavy brown hair, 4½ feet high by 6½ feet long, supported upon



YOUNG FEMALE MUSK-OX.

In the New York Zoological Park, 1902.

very short and post-like legs that are half hidden by the sweeping pelage of the body. The three-inch tail is so very small and short it is quite invisible. There is a blunt and hairy muzzle, round and shining eyes, but the ears are almost invisible.

The whole top of the head is covered by a pair of horns enormously flattened at the base, and meeting each other in the centre line of the forehead. From the meeting point they sweep downward over the edge of the cranium, close to the cheeks, but finally recurve upward before coming to a point, like the waxed mustache of a boulevardier.

The iris of the Musk-Ox is of a chocolate brown

color, the pupils are elongated, and bluish-purple. The lips and tip of the tongue are also bluish-purple.

The outer hair is a foot or more in length, and often touches the snow when the animal walks. In the middle of the back is a broad "saddle-mark," of shorter, dull-gray hair. Next to the body is a woolly coat of very fine, soft, light brown hair, very clean, and so dense that neither cold nor moisture can penetrate it. This is for warmth. The longer and coarser hair that grows through it is the storm-coat, to shed rain and snow. Our first Musk-Ox began to shed its woolly under-coat on April 10. On April 26, it was loose all over the body, and beginning to hang in rags; therefore, for both the comfort and the appearance of the animal, we threw her upon the ground, held her securely, and combed it all out. It was very fine, curly, free from oil, and the entire mass weighed six pounds.

Although known for more than a century, the Musk-Ox is one of the last of the large land mammals of the world to come into captivity for public exhibition, and it was not until 1900 that its soft anatomy was studied for the first time.

Anatomically, this animal presents a few sheep-like features. By some writers their importance has been so much exaggerated that the name "Musk-Sheep" has been proposed as a substitute for Musk-Ox. But the sheep-like characters are insignificant in comparison with those that are clearly ox-like.<sup>1</sup>

Two species have been described. That of the Barren Grounds of the mainland of North America has long been known as *Ovibos moschatus*. In 1901, the animal of Greenland and northern Grinnell Land was described as *Ovibos wardi*, the White-Fronted Musk-Ox, because of a band of gray or dirty-white hair, extending across the top of its head.

Although this animal is called a Musk-Ox, it has neither the odor nor taste of musk, and its flesh is excellent food. General Greeley, Commander Peary and many other explorers have feasted on its flesh. In their native desolation, these animals go in herds of from twenty to fifty head, are easily brought to bay by dogs, and under such circumstances they stupidly stand


<sup>1</sup> See E. Lonnberg on the *Anatomy of the Musk-Ox*, in the Proceedings of the Zoological Society of London, 1900.

facing their enemies until killed. This habit, so fatal in the presence of man, is all that saves the herds from being exterminated by the hordes of big white wolves which infest the Barren Grounds.

General A. W. Greely states that the average weight of ten Musk-Oxen, dressed, was 360 pounds, while the heaviest weighed 432 pounds. This would indicate an average live weight of 404 pounds, and a maximum live weight of 604 pounds.

The accompanying map shows the range of

Natural History Museum of Stockholm, made important and valuable contributions to the life history of *Ovibos wardi*. On the barren, rocky hillsides and level upland pastures surrounding Scoresby Sound and Liverpool Bay, from latitude 70° to about latitude 74°, the expedition found Musk-Oxen in herds of from three to sixty-seven individuals, until the total number observed amounted to between two hundred and thirty and two hundred and forty. For the first time, this remarkable species was photographed



RANGE OF THE MUSK-OX.

Heavy black spots signify actual occurrences. The dotted area indicates the probable range of the genus. The species north of Great Slave Lake is *Ovibos moschatus*, and that of Greenland and Grant Land is the White-Fronted Musk-Ox, *Ovibos wardi*.

the Musk-Ox, the southern limit of which is 64°. During the last fifteen years whole herds have been killed in the Barren Grounds north of Great Slave and Great Bear Lakes, at Lady Franklin Bay, and on the eastern and northeastern coasts of Greenland.

During the year 1899, a Swedish scientific expedition to the east coast of Greenland, under the leadership of Prof. A. G. Nathorst, of the

in its wild haunts, by Prof. Nathorst, Mr. Johannes Madsen and Mr. E. Nilson, and with very gratifying success. Of these pictures the most perfect is that which shows the leader of the expedition closely approaching a herd.

Prof. Nathorst states that to the leeward of a herd, the odor of the animals was noticeable at a distance of 100 metres, but that when a freshly slain animal is promptly and properly eviscerated,

the flesh is free from musky flavors, and very good.

One of the most important discoveries of the expedition was the fact that the region visited had once been inhabited by Eskimo, but their kitchen-middens contained no remains of Musk-Ox, from which, and from other evidence, Prof. Nathorst concludes that the presence of that animal on the eastern coast of Greenland is due to

ing Island, on the east coast of Greenland. Both were purchased by the Duke of Bedford.

In 1900, thirteen living specimens were captured on the eastern coast of Greenland, between Latitude 70° and 74° and taken alive to Europe.

One male in Woburn Park, England, owned by the Duke of Bedford, survived until 1903.

Of the specimens mentioned above, the following were alive in December, 1903:



Reproduced by permission of A. G. NATHORST.

WILD MUSK-OX HERD AT FRANZ JOSEPH FJORD, E. GREENLAND, 1899.

The figure in the foreground is that of Prof. Nathorst. Photographed by E. NILSON, Lat. 73° 30'.

a southward migration along the coast which has taken place since 1823.<sup>1</sup>

A complete count of all the living Musk-Ox specimens that thus far have reached Europe and the United States should be entered here.

In 1899, a Swedish expedition carried to Europe two male specimens captured on Claver-

<sup>1</sup> See *Le Loup polaire et le Boeuf Musque*, par A. G. Nathorst, Bulletin de la Société Géographique, Paris, 1901.

One male in the Copenhagen Zoological Garden;

One male in the Berlin Zoological Garden;

Three in Norrland, Sweden (one male and two females), practically at liberty on pasture closely resembling their Greenland home.

Of the other specimens, five died in Antwerp when very small, and three in Sweden, in wild pastures.

In March, 1902, the New York Zoological

Park received, as a gift from Mr. William C. Whitney, a female Musk-Ox twenty-one months old, captured on the Barren Grounds north of Great Bear Lake, about Latitude 69°. This specimen died of acute pneumonia on August 16, 1902.

In September, 1902, a very small female Musk-Ox calf, captured by Commander Robert E. Peary, at Fort Conger (Latitude 81°), was received in the New York Zoological Park, as a gift from the Peary Arctic Club. It died in October.

In 1903 (July) five Musk-Ox calves, one male and four females, arrived at Tromsø, Norway, from Greenland, and were offered for sale to zoological gardens generally.

The first specimen exhibited in the New York Zoological Park, in 1902, was captured in March, 1901, thirty miles from the Arctic Ocean, directly north of Great Bear Lake, by a party of Eskimo hunters and whalers sent by Captain H. H. Bodfish, from the steam whaler *Beluga*. Its price, delivered in New York in good health, was \$1,600. When two years old it stood 3 feet 2 inches high at the shoulders, and was 4 feet 10 inches in length. Its food was clover hay, raw carrots or potatoes, a little green grass when in season, and occasionally a few apples.

### The Mountain Sheep.

High on the mountain's frowning crest,  
Where lines of rugged cliff stand forth,  
Where Nature bravely bares her breast  
To snowy whirlwinds from the north;  
High in the clouds and mountain storms,  
Where first the autumn snows appear,  
Where last the breath of springtime warms,  
—There dwells my gallant mountaineer.

And truly he is a gallant mountaineer. Wherever found, the mountain sheep is a fine, sturdy animal, keen-eyed, bold, active and strong. It fears no storm, and defies all enemies save man and domestic sheep. From the former it receives bullets, from the latter, disease. Whether its home is the highest crags of the saw-tooth ranges, the boldest rim-rock of the mountain plateaus, or the most rugged "bad-lands," it is always found amid the scenery that is grandest and most inspiring.

In summer, its favorite pastures are the treeless slopes above timber-line, where, on our northern mountains, grasses and wild flowers

grow in astonishing profusion. When the raging storms and deep snows of winter drive the elk and deer down into the valleys for shelter and food, the mountain sheep makes no perceptible change in altitude.

All the year round, this animal is well fed, and its savory flesh invites constant pursuit by the mountain lion, and by hunters both white and red. The massive, curving horns and handsome head of the adult ram, taken amid grand mountain scenery, with much difficulty and no little danger, constitute, in my judgment, one of the finest trophies that a true sportsman can win. But it must be clean, and not haunted by the ghosts of slaughtered ewes and lambs! One of the greatest days of my life was that on which I pursued and killed, alone, amid the grandeur of the Shoshone Mountains, my first big mountain ram. It was then that I learned how much a mountain sheep needs to be seen in its native cloudland in order to be fully appreciated. It is an animal for which my admiration is as boundless as the glories of its mountain home.

The mountain sheep is a bold and even reckless climber. It is robust and strong on its legs, yet active withal, and capable of feats of endurance that really are astonishing. It cannot, and never did, "leap from a height, and alight upon its horns,"—save by some neck-breaking accident. When pursued it can, however, dash down an appalling declivity, touching here and there, and land in safety, when to the observer it seems certain to be dashed to death.


The young are born in May or June, above timber-line if possible, among the most dangerous and inaccessible crags and precipices that the mother can find. Her idea is to have her offspring begin its life in places so steep and dangerous that a very slight effort on its part will suffice to keep it beyond the reach of foes. The lamb's most dangerous enemy is the eagle, against which the mother successfully guards it.

Except the burriel and aoudad, any adult mountain sheep, from either the Old World or the New, can readily be recognized by its massive, round-curving horns, which, when seen in profile, describe from one-half to three-fourths of a circle, or more. *No wild animals other than wild sheep have circling horns.* The largest species of wild sheep are found in Asia, and are

known respectively as the argali, and Marco Polo's sheep. The horns of the argali are the greatest in size and weight, and those of Marco Polo's sheep have the widest spread.

Six species of mountain sheep are found in

has been known for exactly a century, and it is the species which is most widely known in America. Once quite abundant throughout the Rocky Mountains from Mexico to Latitude 60° in northern British Columbia, it has been so persistently



KNOWN DISTRIBUTION OF MOUNTAIN SHEEP IN NORTH AMERICA IN 1903.

The black dots represent actual observations.

1. Big-Horn Sheep, (*Ovis canadensis*).
2. Mexican Sheep, (*O. mexicanus*).
3. Nelson's Sheep, (*O. nelsoni*).
4. White Sheep, (*O. dalli*).
5. Black Sheep, (*O. stonei*).
6. Fannin's Sheep, (*O. fannini*).


North America, of which five have been described since 1883. They are scattered from the northern states of Mexico through the Rocky Mountains almost to the shore of the Arctic Ocean, and throughout one-half of Alaska, a range fully 3,600 miles long. The accompanying map shows actual occurrences of the various species during the past twelve years.

Of our six species, four are so interesting they deserve separate notice.

**The Big-Horn, or Rocky Mountain Sheep,<sup>1</sup>**  
<sup>1</sup> *O'vis can-a-den'sis*.

hunted and slain that now it exists only in small bands, in widely-separated localities. In most of our western states and territories, the killing of Mountain Sheep is now prohibited for a term of years, and it is hoped that these laws will be





From a painting by CARL RUNGES.

THE MASTER OF THE TRAIL.

A Mountain Lion lying in wait on a narrow sheep-trail, high up on a mountain side, is dismayed by the great size and threatening aspect of the leader of the band of Big-Horn.

enforced and respected. Wherever they are ignored, the wild sheep are doomed to extinction.

The general color of the Big-Horn is gray brown, with a large white or cream-yellow patch on the hind quarters, completely surrounding the tail. A large ram killed by the author in the Shoshone Mountains, Wyoming, on November 16, 1889, stood 40 inches high at the shoulders, was 58 inches in length from end of nose to root of tail; its tail was 3 inches long, and its weight was about 300 pounds. Although the snow on its wild pasture was knee deep, and the sheep were pawing through it to reach the tallest blades of dry grass, they were as well fed and fat as if they had been feeding at a manger.

The largest horns of this species ever taken are

male specimen from Lower California, fifteen months old, which in 1902 was exhibited in the New York Zoological Park was as follows:

Height, 29 inches.

Length, head and body, 39½ inches.

Tail, 3 inches.

Length of horn, 10 inches.

Spread at tips, 13¼ inches.

Weight of animal, 65 pounds.

Cause of death, pneumonia.

In the State of Chihuahua, Mexico, is found the **Mexican Mountain Sheep**,<sup>2</sup> in color much like the Californian species, but larger, and with large ears. The horns of a fine old ram, killed by Mr. Charles Sheldon, measured 16½ inches in basal circumference.



Copyright, 1902, by HARRY PIDGEON.

HEAD OF WHITE MOUNTAIN SHEEP.

Shot and photographed on the Kenai Peninsula, Alaska, by HARRY PIDGEON.

said to measure 18½ inches in circumference, and 52½ inches in length on the curve; but any horns which are 14 inches in basal circumference may be considered large. All female wild sheep have horns, but they are small, short, erect and much flattened. They vary in length from 5 to 8 inches.

Southward of the range of the Big-Horn are found two new species which appear to be offshoots of it. In southern California and the peninsula of Lower California is found the **California**, or **Nelson's Mountain Sheep**,<sup>1</sup> a smaller animal than the big horn, short haired, and of a pale salmon-gray color. The size of the

<sup>1</sup> *Ovis nelsoni*.

**The White Mountain Sheep**, or **Dall's Sheep**,<sup>3</sup> of Alaska, discovered and described by E. W. Nelson in 1884, is an animal of very striking appearance. When its hair has not been stained by mud or dirt, it is everywhere pure white, and its horns have a yellowish, amber-like appearance. From May to September, during the shedding period, the hair is so short and so often stained by reddish earth, that the skin is almost worthless as a trophy. From October to February, however, the pelage is very long and thick, and snow white. This species is noticeably smaller than the Big-Horn, and the horns are smaller and more slender in pro-

<sup>2</sup> *Ovis mex-icanus*.

<sup>3</sup> *Ovis dalli*.



Painted by CARL FUNGIUS.

WHITE MOUNTAIN SHEEP AND BLACK MOUNTAIN SHEEP.

portion. A large adult ram measures 39 inches high at the shoulder, and the ewe 33½ inches.

By reference to the map, it will be seen that this species is very widely distributed throughout Alaska and the Yukon Territory. Ten years ago it was abundant on the Kenai Peninsula, and the head of Cook Inlet, but many have been killed, and the number greatly reduced. Recently Congress has passed a law protecting not only the White Sheep, but all the large game animals of Alaska.

**The Black Mountain Sheep, or Stone's Sheep,**<sup>1</sup> of northern British Columbia, is distinguishable by the wide spread of its horns, the dark-brown color of its sides and upper parts generally, and white abdomen. It is of the same size as the white sheep, but the two species together form a striking contrast. The precise range of the Black Sheep is south of the head waters of the Stickeen River. Although this species and the white sheep have not yet been found inhabiting the same locality, it is probable that they will be, and we have ventured to show both in one plate.

**Fannin's Mountain Sheep**<sup>2</sup> is also a new species, found first on the Klondike River, Yukon Territory, in 1900. It is about the size of the white sheep, and has a snow-white head, neck, and tail-patch, and a bluish-gray body, like a white sheep covered with a gray blanket. It

<sup>1</sup> *Ovis stone'i.*                      <sup>2</sup> *O. fan'nin-i.*

also has a blue-gray tail, and a band of brown running down the front of each leg. The first specimen was sent from Dawson City to the Provincial Museum at Victoria, B. C., in 1900, and since then others have reached New York.

In the table below are given the measurements in inches of some of the largest and finest wild sheep horns with which I am personally acquainted.


**Origin of American Mountain Sheep.**—It seems highly probable that a number of species of North American mammals and birds were acquired by immigration from the Old World. Of this there is no stronger evidence than that furnished by the genus *Ovis*, which was cradled in the mountains of Central Asia. Western Mongolia and Thibet have produced the colossal Argali, the wonderful, wide-horned Polo sheep and the robust Siar sheep.

As the genus spread southward, it produced the small Urial and Burrehel, and stopped short at the northern edge of the super-heated plains of India. But northward, its fate was very different. From the place of its nativity,—let us say the Altai Mountains,—there stretches north-eastward through Siberia and Kamchatka, Alaska, and British Columbia to northern Mexico a practically unbroken range of mountain sheep, 7,500 miles long. From northern India to northern Mexico, the species stand in the following order: burrehel and urial; Argali and Polo's sheep; Siar sheep; Kamchatkan sheep; white

**MEASUREMENTS OF MOUNTAIN SHEEP HORNS.**

			BASAL CIRCUM-FERENCE.	LENGTH ON OUTER CURVE.	SPREAD.	OWNER.
<b>Siberian Argali,</b>	<i>Ovis ammon,</i>	Central Asia,	19½	59½	40	W. T. Hornaday.
<b>Marco Polo's Sheep,</b>	<i>Ovis poli,</i>	Central Asia,	15½	59½	39	Robert Gilfort.
<b>Siar Sheep,</b>	<i>Ovis siarensis,</i>	Central Asia,	15½	47¼	30½	W. T. H.
<b>Big-Horn,</b>	<i>Ovis canadensis,</i>	British Columbia,	16½ <sup>1</sup>	40½	17	G. O. Shields.
<b>Black Sheep,</b>	<i>Ovis stonci,</i>	British Columbia,	14¼	32½	28¾	Unknown.
<b>White Sheep,</b>	<i>Ovis dalli,</i>	Kenai Pen., Alaska,	12¾	38½	23½	W. T. H.
<b>Mexican Sheep,</b>	<i>Ovis mexicanus,</i>	Chihuahua, Mexico,	16½	35	18¼	Charles Sheldon.

<sup>1</sup> Circumference half way between base and tip, 16 inches ! Weight, skull and horns, 38 lbs.



Painted by CARL RUNGUIS.

ROCKY MOUNTAIN GOAT.



HORNS OF ASIATIC AND AMERICAN MOUNTAIN SHEEP.

1. Siberian Argali. No. 1 in list on page 112.
2. Marco Polo's Sheep. A specimen of medium length, only.
3. Big-Horn. No. 4 in list. A very large pair.
4. White Sheep. No. 5 in list; of unusual length.

sheep; black sheep; Big-Horn; Nelson's and Mexican sheep.

It requires no stretch of the imagination to behold Bering Strait choked with the great polar ice-pack, and hardy, strong-limbed bears, wolves, mountain sheep and reindeer crossing over the sixty miles that now separate Asia from Alaska, and spreading in all directions over North America. I fully believe that the parent stock of our mountain sheep, caribou, moose, wolves and bears came from Asia by this route.

**The Rocky Mountain Goat, or White Goat,<sup>1</sup>**

<sup>1</sup> *O-re-am'nos mon-tan'us.*

is the only American representative of the numerous species of wild goats, ibexes and other goat-like animals so numerous throughout the Old World from Japan to India, southern Europe and northern Africa. Thus far with but one exception all the rumors of "ibex" that have come from Wyoming, Colorado, Montana and British Columbia have proven entirely without foundation. In one case a Colorado hunter discovered a small band of once-tame goats running wild and reported it to *Recreation* magazine, with a photograph of a mounted specimen. While it is possible that a genuine *Capra* may yet be found inhabiting some unexplored region, like the Romanzoff Mountains, such an occurrence is very improbable.

The only use or value thus far found in the Mountain Goat is as "game" for sportsmen who like difficult and dangerous tasks. With but few exceptions, it inhabits the grassy belt of the high mountains just above timber-line, and it particularly loves the dangerous ice-covered slopes and "hog-backs" over which only the boldest hunters dare follow it. This,

however, specially applies to its haunts in the Rocky Mountains, and the Coast Range. On the coast of British Columbia, the White Goat sometimes descends so near to tide water that more than one specimen has been shot from a canoe.

For a large Ungulate, the Mountain Goat is said to be phenomenally stupid. It is quite true that any hunter who has the nerve and strength to climb to where it lives will there find no great difficulty in killing it. From all accounts, it is both erratic and stupid. Several times goats have approached the camp-fires of explorers, and

SEARCHED THE CAMP \_\_\_\_\_  
 SEARCHED THE CAMP \_\_\_\_\_  
 SEARCHED THE CAMP \_\_\_\_\_  
 SEARCHED THE CAMP \_\_\_\_\_

on one occasion an individual whose "partner" had been shot deliberately sat down, dog-like, thirty yards away and watched the hunter skin and cook a portion of his mate. In Idaho two miners killed a large Mountain Goat with an axe. While exploring in Alaska, unarmed, a member of the United States Geological Survey was once vigorously attacked by an old male goat, which attempted to drive him from a narrow mountain path.

The White Goat is quite as odd in appearance as in mind and habit. Judging merely from its appearance an observer would be justified in considering it a slow, clumsy creature, safe only upon level ground. Instead of being so, it is the most expert and daring rock-climber of all American hoofed animals. Its hoofs are small, angular and very compact, and consist of an ingenious combination of rubber-pad inside and knife-edge outside, to hold the owner equally well on snow, ice or bare rock.

Professor L. L. Dyche declares that Mountain Goats will cross walls of rock which neither man, dog nor mountain sheep would dare attempt to pass. He has seen them cross the face of a precipice of apparently smooth rock, to all appearances entirely devoid of ledges or shelves of any kind, and so nearly perpendicular that it seemed an impossibility for any creature with hoofs to maintain a footing upon it. And yet, the goats not only passed safely across, but they did it with perfect composure, frequently looking back, and turning around whenever they saw fit to do so.

In general outline this animal has the form of a pigmy American bison, and were its pelage dark brown instead of pure white, the external resemblance would indeed be striking. It has high shoulders, low hind-quarters, stocky legs, a thick-set body, and shaggy pelage. Its head is carried low, the crown seldom rising above the upper line of the shoulders and back, and the face is too long for beauty. The horns are so small, short and severely plain they are neither beautiful nor imposing.

The weight of this animal is about that of the Virginia deer. The shoulder height of a good average size male is 37 inches, length of head and body, 66 inches, tail, 4 inches, and girth 51 inches (L. L. Dyche). The females average about one-fourth smaller. Except in length and color of

pelage the Mountain Goat is clad after the style of the musk-ox. Next to the skin it wears a dense coat of fine wool, through and far beyond which grows a long, outside thatch of coarse hair. When free from dirt, both these coats are yellowish-white, and contain no patches of color. Behind each horn is a peculiar bare patch of black, oily skin, the size of a half-dollar. The horns are small, smooth, very sharp-pointed, jet black, and the longest on record measure  $11\frac{1}{2}$  inches. The cannon bone is proportionately the shortest to be found in any large ungulate.

Professor Dyche thinks this animal is not likely to be exterminated very soon, chiefly because of its inaccessibility, its lack of beauty as a trophy, and the expenditure of time, money and muscle that is necessary to win within gunshot of it. Its flesh is so musky and dry that it is not palatable to white men save when they are exceedingly hungry, and its skin has no commercial value. Nevertheless, in the United States, the White Goat has been so much sought by sportsmen and others who like difficult hunting that now it is found only in Washington, Idaho and northwestern Montana. Northward of our boundary, it is scattered very thinly, and at long intervals, throughout British Columbia and Alaska as far as the head of Cook Inlet.

In 1900 a new species was discovered on Copper River, Alaska, and named **Kennedy's Mountain Goat**. It is marked very plainly by horns that are no longer, but are more slender, more strongly ringed, and spread farther at the tips than those of the original species.

Up to the year 1903, only four white goats had ever been exhibited alive in the United States east of the Rocky Mountains. Of these, two were shown at Boston in 1899, and two are now alive in the Philadelphia Zoological Gardens. As might be expected, it is a difficult matter to keep such creatures alive and in good health on the Atlantic coast. In 1902 a very fine adult male specimen was on exhibition in the London Gardens.

#### PRONG-HORNED ANTELOPE FAMILY.

##### *Antilocapridae.*


This unique Family of one species and one subspecies, must not be confused nor in any way connected with the large and important group of African antelopes, which contains a grand array

of animals of all sizes, many of them odd, and many of them noted for their beauty. The student who has a special liking for the large hoofed animals surely will find pleasure in making the acquaintance of such superb creatures as the sable antelope, the koodoo, the water-buck, the eland, the oryx, the gnu, the pallah, and the hartebeest of Africa. We have reason to envy Africa her exclusive possession of all those fine creatures, not to mention her other hoofed animals, great and small.

**The Prong-Horned Antelope<sup>1</sup>** is found only

cent bullet flies true to the mark, it will destroy an animal more wonderful than the rarest orchid that ever bloomed.

Remember the ages which Nature has spent in fashioning this wonderful combination of keen eye, fleet foot and graceful limb, and preserving it from the extermination which overtook the great reptiles, rhinoceroses, and toothed birds of the vast inland sea now known as the Uintah Basin. Surely this animal is worth perpetual protection at our hands, rather than needless, cruel and inexcusable slaughter. *It cannot*



Painted by CARL RUNGJUS.

#### PRONG-HORNED ANTELOPE.

in North America, and it possesses so many anatomical peculiarities, found in no other animal, that zoologists have created for it a separate family, which it occupies in solitary state. It is like an island in a vast sea, unrelated. Let him who hereafter may be tempted, either lawfully or unlawfully, to raise a death-dealing rifle against one of these beautiful prairie rovers, remember two things before he pulls the trigger: In this land of plenty, no man really needs this creature's paltry pounds of flesh; and if his two-

<sup>1</sup> *An-ti-lo-cap'ra americana*.

*be perpetuated by breeding in captivity; and unless preserved in a wild state, it will become extinct.*

Behold the list of characters, in which this animal differs from all other antelopes: Although its horns grow over a bony core, they are *shed and renewed every year*; the horn bears a prong, and is placed directly over the eye; the feet have no "dew-claws"; the hair consists of a hollow tube filled with pith, coarse, harsh, straw-like and easily broken; and all the hair on the rump is fully erectile, like the bristles of swine.



When fighting, or alarmed, this white hair is instantly thrown up, and on a fleeing animal it forms a dangerously conspicuous and inviting mark. To my mind, the white rump-patch of the Prong-Horn is one of Nature's errors. It enables a pursuer to mark the animal long after it should really become invisible.


The Prong-Horned Antelope is next in size to the smaller species of our mountain sheep. It is smaller than the white-tailed deer of the north, but as large as the southern forms. The largest specimen in the Zoological Park herd measures 37½ inches high at the shoulders, has a head and body length of 47¾ inches, tail, 3½ inches, and chest circumference of 35 inches. Its horns are 12¾ inches long and 12½ inches wide between the tips. The longest horns on record are 17 inches in length, but any that measure 12 inches may fairly be considered large. The female has no horns.

The colors of this animal are usually two, consisting of a cloak of light yellowish-brown thrown over the back and neck of an otherwise white animal. On the throat the brown is laid on in a curious collar-like pattern, and the adult males usually have a wash of black on their cheeks. The ears are very shapely, and from the neck an erect mane rises from four to five inches in length. The legs are exceedingly trim and delicately formed, and the erect horns and high pose of the head give the animal a very jaunty appearance.

In running it has three very distinct gaits. When fleeing from danger, it carries its head low, like a running sheep, and gallops by long leaps; when showing off, it holds its head as high as possible, and trots forward with stiff legs, and long strides, like German soldiers doing the goose-step. Occasionally, it gallops with high head, by stiff-legged leaps, like the mule deer.

In captivity the Prong-Horn is always affectionate, trustful, and very fond of being noticed;

but the bucks soon become too playful with their sharp horns, and push their human friends about until the play becomes more dangerous than amusing. They readily come at call, and at times become very playful with each other. They cannot live on the rich, green grasses of the country east of the Great Plains, and are very difficult to keep in captivity. At the New York



DISTRIBUTION OF THE PRONG-HORNED ANTELOPE IN 1903.

Zoological Park it has been found that they survive and breed only when kept in a paved corral, and fed on rolled oats, clover hay, and a very limited amount of fresh grass. Those who have attempted to preserve and breed the Prong-Horn in captivity have met with many discouragements, and failure has been the result of many experiments that deserved success. At present, our herd seems well established, and on June 2, 1903, two fawns were born.

Owing to the extreme difficulty of maintaining this species in captivity, its total extinction at an early date seems absolutely certain, unless it is fully and permanently protected in a wild state, on its native ranges, for a long period. To-day it exists only in small, isolated bands, widely scattered, in a few localities in Montana, Wyoming, Colorado, Kansas, Texas, New Mexico, Nevada, Idaho, Utah, Oregon and Califor-

nia. In all these states save three its destruction has been absolutely prohibited for periods ranging from five to ten years, and it is hoped and believed that all will very shortly provide for its absolute protection. But has protection come to this species early enough to save it? It is very doubtful. Says Mr. A. G. Wallihan, in *Outdoor Life*, "Look at the Antelope! But I don't know whether you can find any to look at; for I don't think there are 50 in Routt County [Colorado], where ten years ago there were probably 50,000. They have almost completely disappeared here. No doubt a small herd of a thousand or so went north into Wyoming, but they will fare no better there."

The destruction of this beautiful and interesting creature is now absolutely inexcusable, and for the good name of Americans generally it is to be hoped that wherever a wild Prong-Horned Antelope is now to be found, public sentiment will protect it more powerfully and more permanently than can any statute law.

## THE DEER FAMILY.

### *Cervidae.*

**General Observations.**—The Deer Family is well represented on all continents, and on all large islands, save Africa, Australia and New Zealand. There are about forty-five well-defined species, and many subspecies. With but one or two exceptions, the species found in the tropics and subtropics are scantily antlered, dull in color, and covered with coarse, thin hair. There is but one tropical deer which is really beautiful, and that is the axis, or spotted deer, of India and Ceylon.

The following facts regarding the deer of the world are worth remembering:

The American Moose is the largest member of the Deer Family, living or extinct.

The American Elk, or Wapiti, is the largest and finest of all the round-horned deer.

The Axis Deer is the most beautiful in color of all deer.

The Moose has the heaviest and most massive antlers, with the widest spread.

Male deer of most species have solid antlers, of bone, branching into several tines.

Deer shed their antlers, and renew them completely, every year.

The young of nearly all round-horned deer are spotted at birth.

All adult male deer are dangerous in the mating season, when their antlers are new and perfect.

The female Caribou is the only female deer with antlers.

The best deer to keep in captivity in a park is the Fallow Deer, of Europe; and outside of its own home, the worst is the Columbian Black-Tail.

Except as already stated, nearly every country in the world is provided with representatives of the Deer Family, according to conditions. Nature has fitted the caribou to live in the awful lands of desolation in the far north, and the moose in the forests fringing the Arctic barrens. The elk is fashioned for the plains, the foothills and open-timbered mountains of western America and central Asia. The white-tailed deer skulks in safety through the thickest forests of temperate North America, and in India and the far East the axis deer, the sambar, and the tiny muntjac, with only one or two tines on each antler, have been formed to slip through the tangled jungles with ease and safety.

North America has the good fortune to be rich in Cervidae. It has six prominent types, and at this date (1903), a full count reveals twenty-four recognized species and subspecies, which form a group combining the grand, the beautiful and the picturesque, and of very decided value to man. In the exploration and settlement of the United States, and the exploration of Alaska and the far North, the wild herds have played an important part.

The unvarying distinctive mark by which any American representative of the Deer Family can be recognized is the presence on the male of *solid horns of bone, called antlers, which are shed once a year, close down to the skull, and are fully renewed by rapidly growing out in a soft state called "the velvet."* When fully grown, the antlers branch several times; but the first pair, which are grown during the second year, are only two straight and slender spikes, called "dag antlers." The grouping of animals with antlers brings together in the Deer Family not only the true deer, but also the moose and caribou.

**Shedding and Renewal of Antlers.**—At this point it is necessary to emphasize certain facts regarding the antlers of deer, elk, moose and caribou.

Many persons find it difficult to believe that the antlers of all these creatures drop off close to the skull, every year, and are completely renewed in about four months; but such is the fact. It is Nature's special plan to absorb the

seems incredible—unless watched from week to week—that the enormous antlers of full-grown moose or elk can be dropped and completely renewed again in as short a period as four months; but it is true.



“DO ELK SHED THEIR ANTLERS?”

An answer from the New York Zoological Park.

1. March 21.  
3. April 30.

2. April 8.  
4. May 15.

surplus strength of the males, and render them weak and inoffensive during the period in which the mothers are rearing their young, when both the does and their fawns would be defenceless against savage males with perfect antlers. It

The antlers of North American deer are usually dropped in March, but occasionally in February. Sometimes a day or two passes between the fall of the first antler and the loss of the second. The root, or *pedicle*, exposed is a rough

disc of bone belonging to the frontal bone of the skull. No blood flows. Dropped antlers are sometimes gnawed by rodents until destroyed; but many are picked up by those who look for them. At the end of the first week, the bony disc or seat of the antler is covered over by the dark brown skin of the head. At the end of two weeks, a rounded bunch, like a big brown tomato, has risen on the pedicle of each antler. It is soft, full of blood, and easily injured.

Gradually this elongates into the form of a thick, blunt-ended club, in color brown or pink, shiny, and thinly covered with minute hairs. When fairly started, the antlers of a healthy and vigorous elk or caribou grow at the rate of one-third of an inch per day, or even more. They are soft, spongy, warm, full of blood, are easily injured, and if cut will bleed freely. The material of which they are composed, internally, is the same as that which forms the hair. The drain upon the animal's vitality during this period is very severe, and it is not strange that the animal is then meek and spiritless.

A large pair of elk antlers, dropped in the Zoological Park on March 21st, had been renewed to their full length by June 21st, but the tips were flat and club-like. The first sign of the hardening process was the shrinkage of the blunt tips of the tines to sharp points. Gradually the diameter of the entire antler decreased in size, and at the same time the hair composing the velvet grew longer. The surface now assumed a gray appearance. On August 1st all the points were sharp, and the antlers were in perfect form, but the velvet was all on. (See "The Elk's Calendar," page 122.)

**Deer as Dangerous Animals.**—The rapid multiplication of deer parks, and small collections of captive animals, renders it necessary to offer a few words of warning regarding deer of all species. During the season immediately following the perfect development of the new antlers,—say September, October and November,—male deer, elk, caribou and moose sometimes become as savage as whelp-robbed tigers. The neck swells far beyond its natural size, the eye-pits distend, and the buck goes stalking about with ears laid back and nostrils expanded, fairly spoiling for a fight. I have seen stags that were mild and gentle during eight or nine months of the year suddenly transformed into murderous

demons, ready and anxious to stab to death any unarmed man who ventured near.

At first a buck walks slowly up to his victim, makes a wry face, and with his sharp, new antlers makes believe to play with him. Not wishing to be punctured, the intended victim lays hold of the antlers, and seeks to keep them out of his vitals. On finding himself opposed, the buck begins to drive forward like a battering ram, and then the struggle is on.

Heaven help the man thus attacked, if no other help be near! He shuts his teeth, grips the murderous bone spears with all his strength, leans well forward, and with the strength and nimbleness of desperation, struggles to maintain his grasp and keep his feet. Each passing instant the rage of the buck, and his joy of combat, increases. If the man goes down, and help fails to come quickly, his chances to escape the spears are few.

Once when unarmed and alone, I saved myself from an infuriated buck (fortunately a small one), by suddenly releasing one antler, seizing a fore-leg low down, and pulling it up so high that the animal was powerless to lunge forward as he had been doing. In this way I held him at bay, and at last worked him to a spot where I secured a stout cudgel, with which I belabored him so unmercifully that he was conquered for that day.

The strength and fury of a buck of insignificant size are often beyond belief. The loving "pet" of May readily becomes the dangerous, fury-filled murderer of October. With a large deer of any species, a man not fully armed has little chance. In the winter of 1902, at Helena, Montana, a man armed with a pitchfork entered an elk corral, to show a friend that the large male elk feared him. The elk attacked him furiously, and killed him before he could be rescued.

Men who have charge of deer herds must keep the bucks in a perpetual state of fear. Do not make a pet of any male member of the Deer Family after it is two years old. It is dangerous. In the autumn or winter, never enter an enclosure containing deer, elk or caribou unless armed with a pitchfork, or a long pole of tough wood, with an iron spike in the end. If a buck threatens to attack you, strike him *across the nose*; for that is his tender spot. When angry he can take any amount of punishment on the

forehead, neck and shoulders, without its diminishing his energy in the least.

Solitary bucks in small corrals are most dangerous. Where deer run in a large herd, the danger is much less; but if a herd-buck begins to approach people with the slow stride of a pugilist, lips and nose turned up, ears laid back, and snorting defiantly, shut him up at once, or saw off his antlers close to his head, before he does mischief.

locked, wild deer are much given to fighting during the rutting season. It is to be remembered, however, that male deer are in the habit of playfully sparring with their horns, and it is very likely that many a death-lock has been due to a pushing-match rather than to deadly combat. The antlers of our white-tailed and mule deer are peculiarly adapted to the fatal interlocking that has caused many a fine buck to perish miserably by slow starvation. In cap-



Photographed by E. R. SANBORN, New York Zoological Park, 1903

A MODEL AMERICAN ELK, IN OCTOBER.

**Fighting Among Deer.**—Even among themselves, deer are murderous brutes. It is quite a common thing for one buck to treacherously assassinate another; and some are such thorough degenerates they will murder their own does and fawns. The largest and handsomest bucks are not always the best fighters, for they often lack the activity and youthful vigor which gives supremacy to a younger animal.

Judging by the number of pairs of deer that have been found dead with their antlers tightly

tivity, pushing-matches amongst deer are quite common.

#### The Round-Horned Deer.

The American Elk, or Wapiti,<sup>1</sup> is as tall as a horse, handsomely formed, luxuriantly maned, carries its head proudly, and is crowned by a pair of very imposing antlers. Even the doe Elk is a handsome and stately creature; and

<sup>1</sup> *Cervus can-a-den'sis*. In Europe, this animal is called the Wapiti; and the European Moose is called the "Elk."

from the second week after its birth, the fawn steps about with the air of a game-cock. If you will observe a seven-year-old male Elk in October or November, when the modelling of his form is handsomest, his pelage long, bright and immaculate, his neck swollen with pride, and his fine new antlers ready for admiration or for battle, I think you will say, "This is the king of the *Cervidae*!"

Even the moose, giant though he be, is not a creature of regal presence, like the Elk. Although the latter is a large and heavy animal, it has the small and shapely legs and hoofs of a thoroughbred. It is strictly a creature for solid ground, and while very fond of bathing in ponds during hot weather, it avoids swamps and low situations.

It is both a grazing and browsing animal. Although up to twenty-five years ago it often ranged far out into the western edge of the Great Plains, and loves to frequent mountain parks, it is also a forest animal. Originally, its range coincided to a remarkable extent with that of the buffalo, covering fully three-fourths of the United States, from the Adirondacks and the eastern foothills of the Alleghenies to California and Vancouver Island. It was not found, however, on the Great Plains north of the Saskatchewan.

In summer it ascends the Rocky Mountains to the very crest of the Continental Divide, 11,000 feet above the sea. The species reaches its highest physical development on the backbone of the continent, between northwestern Wyoming and southern Colorado.

From nineteen-twentieths of its original range, this grand animal has been exterminated. Today it is abundant in one locality only, the Yellowstone National Park and the country immediately surrounding it, where about 20,000 Elk find a safe retreat.

Every winter the Elk herds of the Yellowstone Park migrate southward to feed in the sheltered valleys of Jackson Hole. During these migrations, which usually are made through deep snow, Mr. S. N. Leek and others have made many fine photographs of the herds. One of Mr. Leek's striking pictures is reproduced on the opposite page.

Elk are found in small numbers in the Olympic Mountains of Washington, in Oregon, sparingly in Colorado, western Montana and Idaho, in one

small area of Manitoba, and at one point in south central California. On Vancouver Island the species is now extinct.

It is probable that within a few years the Elk will disappear from all the localities mentioned save the Yellowstone Park, for in the other wild and thinly settled regions which it inhabits to-day, the measures for its protection from illegal slaughter are by no means adequate. Some Americans who go hunting—I will not call them sportsmen—are so greedy, so lawless and so wasteful of animal life, that we frequently hear accounts of Elk slaughter which are enough to disgust all decent men.

Fortunately, Elk are easily bred in confinement, and during the last twenty years many good herds have been established in the great private game preserves that are scattered from New Hampshire and Massachusetts to Minnesota. In addition to these, there are many smaller herds in small private parks. Nearly every city north of the Potomac has a herd of Elk in one of its parks, and other hardy native animals in an establishment known either as a "zoo," a zoological garden, or a zoological park. Thanks to this constantly increasing public demand for living collections of wild animals, the American Elk and buffalo are now familiar objects to the children of at least twenty American cities.

#### The Elk's Calendar in the New York Zoological Park.

- Jan. 1. Pelage has grown perceptibly paler.
- Feb. 1. Pelage has lost its lustre, and begins to look weathered.
- Mar. 21. Antlers of the largest male dropped, 9 hours apart.
- Apr. 8. Each budding antler looks like a big brown tomato.
- Apr. 18. New antlers about 5 inches long, thick and stumpy.
- Apr. 30. Each antler has developed three branches. Young elk born, well spotted. Closely hidden in the rocks. Height 26 inches; length 35 inches; weight 30½ pounds.
- May 10. Shedding in full progress; the Elk look their worst.
- June 1. Shedding about half finished.
- June 18. Antlers now full length, but club-like;



THE WINTER HOME OF THE ELK.  
Photographed in Jackson Hole, Wyoming, by S. N. LEEK.

well haired. Tips flat. Large male has finished shedding.

- July 20. Antlers are now sharp at the tips. Flies troublesome. Herd bathes in the pond frequently and long.
- Aug. 1. Entire herd now free from winter pelage. Animals look well in short, red summer coat, but smaller! Velvet still on antlers. Spots on young are all gone, and white rump-patch is fully developed.
- Aug. 15. Two big males began to rub velvet from antlers, against trees.
- Aug. 22. Antlers of one bull almost clean, but velvet still hangs in tatters, like carpet rags. Tips pure white, base looks bloody.
- Sept. 15. The summer coat has been completely shed.
- Oct. 1. The herd is at its best. All antlers clean and perfect. Pelage long, full, and rich in color. Mating season now on. Bulls aggressive and dangerous. Fawns active and playful. The "bugle" of the bull is a shrill shriek, sliding down the scale into a terrific bawl.

**Size of Elk.**—Professor L. L. Dyche, an exceedingly careful observer, has contributed a striking illustration of the difficulty of obtaining from a dead Elk an accurate measurement of the animal's standing height when alive. The largest and finest male Elk ever taken by him (for the State University of Kansas) fell in Colorado on October 21, 1891. I can testify that it is a grand representative of its species.

As is frequently done, the guide of the party measured its height in a line from the *point of the hoof* to the top of the shoulder, and recorded 65 inches. This being ruled out, the bottom of the hoof was held parallel with the axis of the body, and the elbow *even* with the lower line of the brisket. This gave 62 inches. Professor Dyche then pushed the elbow up to the position it occupies in a standing Elk—about five inches above the lower line of the body—and found the actual standing height at the shoulders to be 57 inches. The head and body length was 97 inches; girth, 73 inches; circumference around

abdomen, 81 inches; circumference of neck, 36 inches.

On October 3, 1903, a fine bull Elk in the New York Zoological Park was suffering so intensely from a horn wound in the hock joint that it was chloroformed. Being in fine condition, its measurements and weight were carefully noted, with the following result:

Height at the shoulders.....	56½ inches.
Length of head, body and tail... 86¾	"
Circumference of chest.....	78 "

*Weight.*

Trunk.....	300 lbs.
Skin, head and legs.....	255 "
Viscera.....	151 "

Total live weight..... 706 lbs.

*Antlers.*

Length, following curves.....	53 inches.
Widest spread.....	35 "
Circumference above bez tine... 7½	"
Points.....	7+7=14.
Age about 8 years.	


**Rule for Obtaining the Live Weight of Deer from Dressed Weight.**—So many records of the "dressed" weight of deer are published, it is desirable to offer a simple rule by which anyone can accurately calculate the weight of the animal when alive. Taking an antlered Elk (*Cervus canadensis*) as a basis, we find that the dressed weight represents .21388 of the live weight, or  $\frac{2}{9}\frac{5}{6}$  of the whole animal.

*The dressed weight being given, in pounds, add to it five ciphers, divide by 78612, and the result will be the live weight, in pounds.*

While this rule will often prove convenient, the author desires to state that none of the weights recorded in this volume were obtained by it; and any weight so obtained and published always should be marked "as calculated."

The longest and widest Elk antlers are not necessarily the handsomest. Usually, antlers that are of great length are slender, whereas the finest pairs are those of massive proportions, fairly symmetrical, and about 60 inches long. The longest pair of reliable record to this date was purchased in Colorado Springs, Colorado, in 1897 for the Emperor of Germany. Their length of beam was 67½ inches, and their points were twelve





Painted by CARL FUNGUS.

MULE DEER IN THE BAD-LANDS OF SNOW CREEK, MONTANA.

in number. They were obtained from an animal killed in White River County, Colorado. A very large pair from the Shoshone Mountains, in the author's collection of horns, has a beam length of 58 inches, a spread of 49½ inches, and burr circumference of 11 inches.

Elk hunting is not always as fine sport as the noble individuality of this animal would naturally lead the hunter to expect. Very often the Elk is unsuspecting, to the point of stupidity. There have been many times when attacking a herd was too much like attacking a herd of cattle. It is not an animal of "highly-wrought-nervous" temperament, like the deer, but when startled is too much given to hesitating, and seeking knowledge, before it dashes away to safety.

During the last three years important steps have been taken, by private individuals only, toward restoring the Elk to the Adirondack forests, which it once inhabited. In 1901, the Hon. William C. Whitney caused twenty-two head to be liberated there, and in 1902, forty more were set free. In August and September, 1903, five car-loads of Elk, sixty-eight head in all, were shipped from Mr. Whitney's game preserve on October Mountain, near Lenox, Mass., and liberated at Saranac Lake, Floodwood Station and near Paul Smith's Station. All these animals had become fully acclimatized on the Atlantic coast, were in fine physical condition, and if not killed by poachers will no doubt multiply at a reasonably rapid rate. That many of these fine animals will from time to time be killed and eaten by lawless and unprincipled persons seems absolutely certain, and the great danger is that they will be killed more rapidly than they breed.

**The Mule Deer, or Rocky Mountain "Black-Tail,"**<sup>1</sup> is a large and handsome animal, and the largest of the North American species that are universally known as "deer." It is easily recognized by its very large ears, the two Y's on each antler, a short, white tail with a small tip of black, and a white patch around the base of the tail. Its antlers are much larger than those of the white-tailed deer. Owing to their size and width, and their more erect poise on the head, the appearance of this animal is more stately than that of any other round-horned American deer, save the elk.

<sup>1</sup> *O-do-coi'le-us hem-i-o'nus*.

In the region it inhabits, this fine animal is known as the "Black-Tailed" Deer; but that name is not appropriate to a creature which has a snow-white tail with only a tiny tip of black. It rightfully belongs to the Pacific coast species, which has a black tail, and is known by no other name than Columbian Black-Tail. To avoid further confusion and misunderstandings, students are urged to speak of the Rocky Mountain species as the Mule Deer.

The winter color of the Mule Deer is a steel gray, to match the gray rocks and vegetation amongst which it lives. Its summer coat is gray-brown, and it is shed in September.

The Mule Deer chooses for its home the most picturesque "bad-lands" and foot-hills of the Rocky Mountain region, and the deep ravines along rivers, but it also ascends the mountain plateaus of its home to an elevation of 12,000 feet. It is a proud-spirited, high-headed animal, a bold traveller, and like the mountain sheep, is often found where the scenery is wild and picturesque. In this respect it differs from the white-tailed deer, which prefers low ground, and either brush or timber in which to hide.

A large Mule Deer buck, shot by the author on Snow Creek, Montana, measured 42 inches high at the shoulders, and 62 + 6 inches in length. A large pair of antlers (in the author's collection) have a beam length of 27¼ inches, spread 29 inches, and have 14 points.

In the United States, the present scarcity of really large antlers in the possession of taxidermists is a sure sign of the approaching end of this species.

In February, 1903, Mr. A. G. Wallihan, the famous photographer of wild animals in their haunts, made the following prediction regarding the impending extermination of the Mule Deer in Colorado, its centre of abundance in the United States:

"Unless we have a close season on deer, five years will see the finish of these animals. Five years would give them a good start again. I will cite you some figures: In 1897 I was on the big trail here for nine days, and I counted within a few of a thousand deer. In 1901 I was on the same trail for eighteen days, and counted two hundred and twenty-eight deer. In 1902 I was out fourteen days, and counted fifty-two deer. More deer passed in a single twenty-four hours in

1892-3-4-5 and 6 than passed during the whole month of October, 1902.

"There are a lot of deer, it is true, on the north slope of the divide at Pagoda and Sleepy Cat mountains, and eastward in the Williams Fork country; but they are practically the remnant. People here say, 'You can't enforce a close-season law.'" (*Outdoor Life Magazine*.)

The Mule Deer reaches its largest and finest antler development in the Rocky Mountains, from Colorado to southern British Columbia. The few widely-scattered survivors of this species are found to-day in central Chihuahua and Sonora, Mexico; western Colorado and Wyoming, south-eastern Idaho, central Montana, and eastern British Columbia. One fact which militates most strongly against the perpetuation of this species is that states and provinces sufficiently wild and unsettled to afford it a home are financially unable to maintain the large force of salaried game-wardens which alone could really protect it from final annihilation.



KELLER, Photo. Copyright, 1900, N. Y. Zoological Society.

MULE DEER WITH ANTLERS IN THE VELVET.

This species ranges as far east as western Dakota, and westward to the Blue Mountains of Oregon. Formerly it was most numerous in Routt County, Colorado, where about forty-five hundred were slaughtered as late as the winter of 1900. Unfortunately, on account of its preference for open country, its ultimate extinction in the United States is only a question of about ten years; for everywhere, save in the Yellow-

stone Park, it is being destroyed very much faster than it breeds.

The Mule Deer nearly always produces two fawns at a birth, and sometimes three. In feeding it is much given to browsing on twigs and foliage, but it also grazes freely when good grass is available. In the Snow Creek country of central Montana I found that its October bill of fare consisted almost solely of the long-leaved mugwort (*Artemisia tomentosa*), a species of very pungent and spicy sage, which was eaten greedily to the complete exclusion of the finest grasses I ever saw in the West.

In running, this deer often progresses by a series of stiff-legged leaps, in which it touches the ground lightly with its hoofs, bounds upward as if propelled by steel springs, and flies forward for an astonishing distance. In Manitoba and a few other localities this remarkable gait has caused this animal to be called the **Jumping Deer**. Owing to the fact that it lives in a dry climate and rarefied atmosphere, and subsists on very dry foods, it is difficult to acclimatize it anywhere outside of its own home. East of the Mississippi most Mule Deer die of gastro-enteritis, but in the Hon. William C. Whitney's great park on October Mountain, near Lenox, Mass., this species has actually become acclimatized.

**The Columbian Black-Tailed Deer**,<sup>1</sup> of the Pacific Coast, is smaller than the typical white-tailed deer, and very much smaller than the mule deer. The outer surface of its tail is black all over, and constitutes the best distinguishing character of the species. The antlers are very variable. Occasionally those of old bucks exhibit the double Y on each beam which is so characteristic of the mule deer; but in most cases the double bifurcation is wanting, and the antlers look very much like those of the white-tailed deer. In its body colors it resembles the latter species more closely than the mule deer.

This species inhabits the well-watered and densely-shaded coniferous forests of the Pacific coast from the north end of Vancouver Island to central California. It feeds freely upon evergreen foliage, and I have seen a captive animal, in its native forest in the great natural park at Vancouver, partake freely of the foliage of spruce, Douglass fir and juniper, in rapid succession.


Because of some diatetic peculiarity as yet un-

<sup>1</sup>*O-do-coil'e-us co-lum-bi-an'us*.

known, the Columbian Black-Tailed Deer cannot live on the Atlantic coast. After persistent efforts, with at least fifteen specimens drawn from Oregon, Washington and British Columbia, and the loss of all through gastro-enteritis, the New York Zoological Society has abandoned its attempt to transplant the species.

In Alaska, this species dwindles still lower, into the **Sitka Deer**,<sup>1</sup> in stature and antlers

it freely risks its life in the thin fringes of cotton-wood timber, quaking-asp and willow brush that border the banks of small rivers and large creeks. Unlike the elk and mule deer, the White-Tail is a great skulker. When hiding, it crouches and carries its head low, and by clinging persistently to the friendly cover of brush or timber, saves itself under circumstances that would be fatal to any high-headed, open-ground species.



Painted by CARL RUNGUIS.

WHITE-TAILED DEER.

even smaller than the Florida white-tail. It is very abundant on Admiralty Island, but as late as 1901 was being slaughtered in great numbers.

**The Virginia Deer, or White-Tailed Deer**,<sup>2</sup> was the first member of the Deer Family met by the early settlers of America when they went hunting along the Atlantic coast. It will also be the last of the large hoofed animals of North America to become extinct. It is a forest animal, but in many portions of the Great Plains region

<sup>1</sup> *O-do-coil'e-us sit-ken'sis*.

<sup>2</sup> *Odocoileus vir-gin-i-an'us*.

The White-Tailed Deer derives its name from its very long, bushy, wedge-shaped tail, which is snowy-white underneath, and also on the edges. When alarmed and running away, this white brush is held stiffly aloft, and with every stride of the bearer, it sways from side to side, in a startling and highly conspicuous manner. While the peculiar mixed gray color of the pelage makes it difficult to see this animal in brushy surroundings, the moment the creature starts to run, its white flag waves as if purposely inviting bullets, and in total defiance of all the laws of "protective

coloration" amongst animals. Indeed, so very flag-like is this creature's waving tail that in the West many hunters call it the **Flag-Tailed Deer**.

There are two points in which this deer differs from all others, and by which it can easily be recognized.

1. Its antlers rise a short distance from the forehead, then suddenly *drop forward, with the beam almost horizontal, and from the beam three long, sharp tines rise perpendicularly*. The antlers of nearly all other deer point backward as they rise.

2. *The tail is very long, pointed at the end, bushy near the body, and white underneath, as described above.*

The White-Tailed Deer is the best known of all our hoofed animals except the buffalo, because it is the one most widely distributed, and has been longest known. Generally speaking, it is a United States species, for it inhabits at least a portion of every state and territory save Delaware, Oregon, Nevada, California and Arizona. To-day it is most abundant in the Adirondacks, Maine, Vermont, northern Minnesota and Michigan. Closely related forms of White-Tailed Deer are fairly abundant in Florida, on the Lower Rio Grande, and in northeastern Washington.

As might naturally be expected, this wide distribution, throughout such a diversity of country and variety of available food, has produced such variations in size that several subspecies have been described. Of the latter, the most important is the dwarf **Arizona White-Tailed Deer**, extending from southern Arizona southeastward into Mexico to Latitude 25°. This animal, like the **Florida White-Tailed Deer**, seems to be nothing more than a diminutive race of the more robust northern type, with very small antlers, and the short, scanty pelage which is necessary to the comfort of deer in the tropics.

In such forests as those which cover the Adirondack Mountains of northern New York, where small lakes are numerous, there are three methods of hunting deer.

*Hounding deer* consists in beating through the forest surrounding a body of water, with a pack of hounds, and chasing the deer until they leap into the water, where they are shot at very short range by men in boats or posted on the shore. It is no credit to anyone, save an invalid

or a cripple, to kill a deer in this manner, any more than to kill a buck out of season, whose antlers are in the velvet. Any person, no matter how stupid, can be paddled up to a swimming deer and permitted to blow its head to pieces at short range. Pot-hunters have even been known to catch swimming deer, and cut their throats.

In forests like the Adirondacks, frequented by



TAILS OF AMERICAN DEER.

1. Columbian Black-Tail.
2. Mule Deer.
3. White-Tailed, or Virginia Deer. (Small specimen.)

a great many people, hounding deer never should be permitted; and in the wilderness mentioned it is now prohibited by law. In the West Virginia mountains, the hunters are posted on the runways of the deer, and are obliged to kill them on the run. This requires good judgment and excellent marksmanship, and is legitimate sport.

*Jacking or fire-lighting* is a very picturesque and romantic method of hunting deer, but inasmuch as it gives the game no chance, and calls for very little skill or exertion on the part of the hunter, it is by some considered unsportsmanlike. In the prosecution of this plan the hunter requires a canoe, a skilful paddler, and a good light. With a flaring jack-light held aloft in the bow, the paddler, or guide, sits in the stern of the boat, and noiselessly paddles it through the darkness, around the shores of lake or river. The hunter sits under the light, and waits for its beams to emblazon the eyeballs of deer standing

on the shore, or feeding in shallow water. Often the boat approaches so near a wonder-struck deer that to miss it is almost impossible.

*Still-hunting* is the true sportsman's method of outwitting deer which for keenness of eye, ear and nose, have, I believe, no superior in the




Photo. and copyright, 1902, by W. L. UNDERWOOD.

YOUNG WHITE-TAILED DEER.

Showing the conspicuous appearance of the tail when held erect.

whole Family. One fine old White-Tailed buck killed by fair and square trailing and stalking is equal to two mule deer or three elk. When first alarmed, the mule deer and elk are prone to halt from curiosity, and stare at the hunter for that fatal ten seconds which so often ends with a ringing "bang," and a fatal bullet.

But not so the White-Tail. Time after time the trailing still-hunter, stealing forward ever so cautiously, sees ahead of him and far beyond fair rifle shot a sudden flash of white, a pillar of cloud swaying from side to side between the tree-trunks, and the vanishing point of a scurrying White-Tail. This creature knows right well that as a discourager of cervine curiosity, nothing

in the world equals a breech-loading rifle. When he hears behind him a rustle of dry leaves, or the snap of a twig, nothing else is so dear to him as space, judiciously distributed between himself and his pursuer. I have sometimes made so bold as to consider myself a fairly good deer-stalker; but I have still-hunted White-Tailed Deer in November, on dry leaves and without snow, when for days and days together I found it utterly impossible to come within fair rifle shot of a buck worth having. At such times, a light snow means a fair chance, and properly evens up the game.

During the summer, while the antlers are in the velvet, the coat of this species is short, thin, and of a bright sandy color, often called "red." In Canada, the Virginia Deer is frequently called the "Red Deer"; but this is a mischievous misnomer, for its use always suggests the red deer of Europe. The red coat is worn about three months, say from May 1 to August 1, and then it rapidly gives place to the beautiful mottled brown-gray suit, so long and thick that the owner looks like quite a different creature, and is fitted to withstand the severest winter weather.

The White-Tailed Deer is one of the most persistent species of the entire Deer Family. Give it suitable ground and full protection, and there is no limit to its increase. On Long Island, where deer hunting is lawful on only four days of each year, the animals are increasing with surprising rapidity.

In the northern portions of its range from Minnesota to the Adirondacks, where it attains its most perfect development, it is next in size to the mule deer, or Rocky Mountain "black-tail," and is really a fine animal. A large buck stands 36 inches high at the shoulders, is 53 inches in length of head and body, its tail is 7 inches long to the end of the vertebrae, and 5 inches more to the end of the hair. A fairly large pair of antlers from central Montana are 23½ inches in length from burr to tip of beam, spread 18 inches, and have 13 points. A heavy Maine buck is reported to have weighed, before being dressed, 278 pounds.

Usually but one fawn is born each year, in May, which at birth is beautifully spotted, stands 15½ inches high, and weighs 4½ pounds.

Let it not be supposed, however, that in the

South the White-Tailed Deer of the North necessarily becomes a small or inferior animal. A collection of more than one hundred pairs of antlers from Texas, recently inspected by the writer, contained a surprisingly high percentage of large and heavy specimens, fully equal in length, spread and weight to the best examples from Montana, Minnesota and Maine.

**Wild Game as a Source of Revenue.**—All persons who pay state taxes in states or territories inhabited by "big game," and game fishes, will do well to bear in mind that under certain conditions wild animal life can be made an important and legitimate source of revenue. The United States Supreme Court has decided (*Ward vs. Race Horse*, 163 U. S. 507) that all wild game on unoccupied lands is the property of the state, and that even the national government may not, either by treaty with Indians, or in any other manner save actual sequestration, convey any rights or privileges affecting it adversely.

The states of New York and Maine long since discovered that their wild deer constituted valuable state property, and both entered seriously upon the task of preserving them from the annihilation that everywhere follows swiftly upon the heels of non-protection. New York elected to preserve the great Adirondack wilderness as a free hunting-ground for her citizens. Maine, with perfectly proper thrift, decided that her game should not only pay the cost of its preservation, but also be made for her citizens a legitimate source of annual income. All guides must be licensed by the state, no visitor may hunt without a guide, and every non-resident hunter must procure a license, at a cost of \$15. This permits the killing of one bull moose and two deer, but no caribou, nor female moose.

As a result of the game and fish laws of Maine that state becomes every autumn a vast hunting-ground, visited by perhaps ten thousand sportsmen who desire to fish, or to procure deer or moose in their haunts. The army of recreationists annually expends within that state a total sum which is usually estimated at one *million* dollars, or more. And yet, the supply of deer is maintained so successfully that to-day there are in Maine a greater number of deer than anywhere else in the United States, unless it be in the Adirondacks.

The records of the Maine railways show accurately the number of White-Tailed Deer transported by them annually for hunters leaving the state, and afford a very fair index of the abundance of the species. The following are the figures for the last nine years:

1894.....	1,001
1895.....	1,581
1896.....	2,245
1897.....	2,940
1898.....	3,377
1899.....	3,379
1900.....	3,756
1901.....	3,882
1902.....	4,495
Total.....	26,656

Of course these figures do not take into account the great number of deer that are killed and consumed in camp, or by residents of the state who live in or near the great hunting grounds. The whole number of deer in Maine is roughly estimated at 100,000, and the total number killed annually at between 15,000 and 20,000.

#### The Flat-Horned Deer.

**The Caribou.**—In general terms it may be stated that a caribou (pronounced car'y-boo) is a wild deer-like animal, which bears a general



REMARKABLE "FREAK" ANTLERS OF WHITE-TAILED DEER.

Total number of points, 78. Owned by Albert Friedrich, San Antonio, Texas.

resemblance to the domestic reindeer of Europe. Its antlers are long, branching, partly round and partly palmated. Considered as a whole, caribou occupy the upper half of the continent of North America, over which they are widely scattered above the 45th parallel of Latitude.

Next to the musk-ox, the caribou is the most northerly of all hoofed animals. It is not only at home on the vast arctic waste above Great Slave Lake, known as the Barren Grounds, but it ranges on northeastward through Ellesmere Land, crosses to the west coast of Greenland, swings around the northern rim of that island, along the edge of the great ice cap, and down the eastern coast, at least as far as Liverpool Bay, Latitude 70°. Doubtless it inhabits the whole coast of Greenland, wherever the naked ridges and valleys of the terminal moraines yield a supply of food: but there is no evidence that it wanders over the vast sheet of lifeless inland ice which covers the interior of Greenland.

At all times, a caribou is an odd-looking creature. Even a very brief inspection is sufficient to reveal the special provisions which Nature has made to enable it to brave the terrors of an arctic climate. The legs are thick and strong, and the hoof is expanded and flattened until it forms a very good snow-shoe. The caribou walks over snow-fields and quaking muskegs, when the moose sinks in and ploughs through them.

Its pelage consists of a thick, closely-matted coat of fine, wool-like hair, through which grows the coarse hair of the rain-coat. It is the warmest covering to be found on any hoofed animal except the musk-ox, or on any animal of the Deer Family. To the touch, the new coat of a caribou feels like a thick felt mat.

The natural food of the caribou is moss and lichens, and in captivity very few survive many months without the former. The supply of moss for the caribou and reindeer of the New York Zoological Park comes from Maine, and costs in that state seventy-five cents per hundred pounds. A full-grown woodland caribou consumes about seven pounds daily.

Although up to this date nine species of caribou have been described, there are but two well-defined groups, the woodland and Barren Ground. In each of these, several species have been described, but it must be admitted that so

effectually do they run together it is not always an easy matter to distinguish them.

In common with many members of the Deer Family, caribou are distinguished chiefly by their antlers. But even here, great difficulties are encountered. With their many tines and points, varying size and forms of palmation, their antlers are subject to thousands of variations. As a result, no two pairs ever are found exactly alike. Between the very long, few-pointed and scarcely palmated antlers of the Greenland caribou, and the short, many-pointed and widely palmated antlers of the mountain caribou, every conceivable form may be found.

If ten pairs of adult antlers of each so-called species were collected in its type locality, and the whole ninety mixed in one heap, the utmost that even an expert could hope to accomplish without a heavy percentage of error would be to separate the collection into two groups, one containing the four species of Barren Ground caribou, the other the five woodland species.

It is useless to enter here into details regarding each of these nine tentative species. Without a very large collection of specimens, and prolonged study of them, it is impossible to define the boundaries between the various species that have been proposed. Let it suffice to present a brief outline of the two great groups into which all our caribou seem to be rationally divisible.

#### The Woodland Caribou Group.

Roaming through the pine and spruce forests, and also the prairies of Newfoundland, Nova Scotia, New Brunswick, northern Maine, Quebec, Ontario and Manitoba, are the caribou longest and best known to us. A typical specimen<sup>1</sup> living in the Zoological Park is a strong lusty animal, 48 inches high at the shoulders, weighing 280 pounds, and endowed with sufficient energy to vanquish the strongest man in about one minute. Its shoulders are high and sharp, its head is held low and thrust straight forward, and as it walks on hard ground its dew-claws and hoofs click like castanets. Its head is long and cow-like, and its muzzle is too large for beauty; but the large, liquid, dark brown eyes appeal successfully against all adverse decisions on questions of beauty.


<sup>1</sup> *Ran'gi-fer cor'i-bou*, from Maine.



When a caribou walks, its long stride and swinging gait proclaim a born traveller and migrant. And truly, the strangest of all caribou habits is that which impels these creatures, particularly the Barren Ground species, to assemble in immense throngs, and for climatic reasons migrate *en masse*, for long distances. In the

*are short in the main beam, liberally palmated both on brow-lines and tips, and have more than thirty points. As a whole, the antlers have a tree-top appearance.*

2. Antlers of Barren Ground caribou, generally, *are long in the main beam, scantily palmated, especially on the tips, and have less than thirty*



E. F. KELLER, PHOTO.

Reproduced from the Seventh Annual Report of the N. Y. Zoological Society.

#### WOODLAND CARIBOU.

Adult male specimen in the Zoological Park. Height at shoulders, 48 inches, weight, 280 pounds. For a Caribou as large as this the antlers are small.

woodland species, however, this habit is not nearly so pronounced.

**Character of Antlers.**—A comparison of many antlers of Woodland caribou with those of Barren Ground animals reveals one or two points of difference which seem sufficiently distinct to be accepted as constant.

1. Antlers of Woodland caribou, generally,

*points. As a whole, the antlers have an arm-chair appearance.*

If these distinctions between the two great groups of caribou will not hold good, none will.

**The Woodland Caribou** of Maine, Ontario and Quebec (*Rangifer caribou*), is the original type of what recently has become a group of species. Its body color is bluish-brown and

gray, which color also suffuses the neck, head and hind-quarters. In October the new coat is of a dark color known as seal brown, quite different from the same pelage in spring.

Originally the **Newfoundland Caribou** were referred to the species named above, but in 1896 they were given rank as an independent species (*R. terracovae*) chiefly on account of their very light color. They are the whitest of all caribou.

In 1899, Mr. Ernest Thompson Seton described

Osborn, the distinguished zoologist of the American Museum of Natural History. This species attains a shoulder height of 55 inches, and is said to be the largest of all caribou. In September its coat is so brown the animal has been described as a brown caribou.

**The Kenai Caribou** of the Kenai Peninsula—but, in 1903 almost extinct in that locality—was described in 1901 as a distinct species, and christened *Rangifer stonci*. In September, 1903, the Secretary of Agriculture issued an order prohibiting for five years the killing of caribou on the Kenai and Alaska Peninsulas.

Regarding the distribution and habits of caribou in the Canadian Northwest, Mr. J. B. Tyrrell, who, while a member of the Canadian Geological Survey, travelled over a greater area of the range of that animal than any other observer known to me, has kindly furnished the very interesting facts quoted below. His letter is dated at Dawson City, September 10, 1903.

“Regarding the portions of the districts of Alberta, Athabasca and Saskatchewan spoken of by you, I am reasonably certain that the Woodland Caribou may be found in all the thickly wooded tracts. This deer is known to the Cree Indians of that country as the ‘Muskeg-Atik,’ or Swamp Deer, in recognition of the fact that it lives in the swamps and coniferous forests, and not on the plains, or on the country studded with groves of poplar. Now, much of Alberta, and a great part of Saskatchewan, is dry, open country, and into such country caribou rarely wander.

“This dry, ‘bluffy’ country extends northward through the western part of Athabasca, but throughout all the *thickly wooded* parts of Athabasca I have no hesitation in saying that Woodland Caribou are not uncommon. They certainly occur along the Churchill River, and I think that their tracks were common along the banks of the Athabasca River, though I cannot definitely remember this, and I have not my note-books here to help me.

“The Indians told me that the Woodland Caribou of the Churchill River and vicinity move northward, and the Barren-Ground Caribou southward in autumn, and that both winter in the same region, in a country where the trees are festooned by a long, black, hair-like lichen (*Alectoria jubata?*). However, I believe that the



ANTLERS OF KENAI CARIBOU.

From photograph of specimen taken on the Kenai Peninsula in 1900, by HARRY E. LEE.

the **Black-Faced Caribou** of southeastern British Columbia (Revelstoke) as *Rangifer montanus*, or **Mountain Caribou**. The new September coat is almost black. The antlers are short, but throw off a surprising array of long tines.

In 1902 the large, dark-colored caribou of the Cassiar Mountains, in northern British Columbia, was described by Dr. J. A. Allen as **Osborn's Caribou** (*Rangifer osborni*), the name bestowed being in honor of Professor Henry Fairfield

Woodland Caribou are not numerous anywhere in the Canadian Northwest Territories, for in all my travels for the Geological Survey of Canada, extending over the period from 1883 to 1898, I did not see a dozen of those animals, though on hundreds of different occasions I saw their great wide-spreading tracks. The only one I ever shot was feeding on a rocky hill, beside a stream that flows into the east side of Lake Winnipeg; and his head is now hanging in the Museum of the Geological Survey, in Ottawa.

"The smaller species of Caribou lives on the Barren Grounds during the summer. On the approach of winter most of the animals migrate

est and value. To many Indian tribes, such as the Dog-Ribs and Yellow Knives, and to many of the Eskimo tribes also, it has been an important source of subsistence, both in food and clothing. It is so peculiarly a creature of treeless and inhospitable regions, and is so independent of the conditions which are essential to the existence of all round-horned members of the Deer Family, that its desolate home has been inseparably connected with its popular name. Species may come, and species may go, but we hope that the brave and hardy Barren Ground Caribou will go on forever.

It is natural that in any animal species which



ANTLERS OF GREENLAND CARIBOU (*R. groenlandicus*).

Showing the form characteristic of the Barren Ground Caribou group. Specimen from the northwest coast of Greenland, in author's collection.

southward to the edge of the forest, though some remain throughout the winter on the open barrens.

"Twice, in 1893 and 1894, I met what is known as 'the herd,' on its way southward, once on a good feeding ground, where hundreds of thousands were collected together, and again on a rough, rocky tract where the individual bands rarely exceeded a few hundred in number, and all were on the run."

#### Barren Ground Caribou Group.

Throughout a vast and very hungry sweep of northlands, the **Barren Ground Caribou**<sup>1</sup> long has been, and still is, an animal of leading inter-

<sup>1</sup>*Ran'gi-fer arc'ti-cus.*

ranges from the east coast of Greenland to the west coast of Alaska (3,500 miles in an air line), and from Grant Land to the Churchill River (1,800 miles), some variations in form, color and horn architecture should occur. Indeed, in a range so immense, it could scarcely be otherwise. While it is probable that some of these variations justify the creation of specific divisions, we are at present less concerned with these details than with a consideration of the group as a whole. Moreover, it may be said with entire truth that naturalists have but recently begun to study the caribou of America; and until far more material has been gathered, it is impossible to set forth the true status and life history of this genus.

The characters which serve to distinguish

Barren Ground Caribou from the woodland groups have already been pointed out,—smaller size, antlers that are longer in the main beam, less palmated and with fewer points. The following forms have been described as independent species of this group; but whether all of them are entitled to specific rank remains to be seen.

#### BARREN GROUND CARIBOU SPECIES.

Greenland Caribou, *Rangifer groen-land'i-cus*, Greenland Coast.

Barren Ground Caribou, *Rangifer arct'i-cus*, Canadian Barren Grounds.

Grant's Caribou, *Rangifer granti*, Alaska Peninsula.

Peary's Caribou, *Rangifer pearyi*, Ellesmere Land.

In view of the tens of thousands of Barren Ground Caribou that have been seen by white men, and the thousands that have been killed by and for them, the scarcity of definite observations upon this group, and of preserved specimens is, as a whole, very unsatisfactory. At present, therefore, the many undetermined questions regarding the component parts of the group render it impossible to do much more than to define the assemblage as a whole.

In general terms it may be said that the average Barren Ground Caribou is a close under-study of the average reindeer of Siberia and Lapland, and also a smaller animal. That all our caribou have descended from the reindeer of Asia, and came to us by crossing Bering Strait on the ice, seems more than probable.

In surveyor's parlance, the head of Cook Inlet is the "point of departure" of the woodland caribou from the reindeer—Barren Ground type. It would be difficult to find on land a clearer or sharper line of cleavage between two groups of animals than that between *Rangifer granti* of the Alaska Peninsula, and *Rangifer stonci* of the Kenai Peninsula. One moment's examination of the types is sufficient to place those species in their respective groups. The antlers of the Kenai caribou are massive, with many long tines on the terminal half of the main beam. They have 36 points, and a tree-top effect when seen from the front. Grant's caribou, however, has a long and naked main beam running up to a terminal bunch of short tines, a wide-open, arm-

chair appearance, and only twenty-seven points, all strongly characteristic of the Barren Ground type. The superior size of the Kenai caribou is confirmatory of the testimony of the antlers of both.

**Geographic Range.**—The centre of abundance of the Barren Ground Caribou group is midway between the eastern end of Great Slave Lake and the southeastern extremity of Great Bear Lake. This, however, is not the geographic centre of its distribution. The great semi-annual migration is about on a line that might be drawn between Cape Bathurst and the eastern extremity of Great Slave Lake, and undoubtedly the great mass of caribou on the mainland east of the Mackenzie assemble along that route.

Another line of migration, also from northwest to southeast, passes eastward of Dawson City, and sufficiently near it that great numbers of caribou carcasses have been sledded in to the meat markets of that city. In 1901 a search of those markets revealed 5,225 pounds of moose and caribou meat on hand at one time. Along the arctic coast between Point Barrow and the mouth of the Mackenzie, tens of thousands of caribou have been killed by natives, and sold to whaling ships wintering along that coast. As a natural consequence, the herds have nearly disappeared from that locality.

Up to the time that Alaska was purchased by the United States, the natives had few firearms, or none at all, and caribou were abundant. Along the west coast, caribou once were so numerous that a cannon from the fort at St. Michael was fired at a herd that passed within half a mile of the settlement. As usual, we immediately supplied the natives with firearms and ammunition; and as a first result, the only caribou now remaining in western Alaska are the few stragglers that the hunters have not yet overtaken. A few herds of Grant's caribou still inhabit the treeless wastes of the Alaskan Peninsula, but on the Kenai Peninsula, the caribou is now believed to be almost extinct. In 1903 it was estimated that only thirty individuals remained alive.

The great herd seen by Mr. Tyrrell at Carey Lake, west of Hudson Bay, will be mentioned in detail later on. On the Labrador Peninsula, there are said to be three distinct herds, on Hudson Straits, Ungava Bay, and the Atlantic coast

down to Hamilton Inlet. From Ellesmere Land, five skins of a white animal with a gray back have been described as **Peary's Caribou**,<sup>1</sup> and from at least four points in Ellesmere Land, Caribou have been reported.

Along the northwest coast of Greenland, especially between Melville Bay and Kane Basin, Commander Peary found a fair abundance of caribou, and at Liverpool Bay, on the east coast, a number were killed by a Danish expedition, in 1900.

**Habits.**—One of the habits of the Barren Ground Caribou is particularly striking. At stated periods, in spring and autumn, they assemble in immense herds, and migrate *en masse* with the compactness and definiteness of purpose of an army of cavalry on a march. This is most noticeable on the Canadian Barren Grounds, which by reason of its summer pasturage and the absence of water barriers, encourages the display of natural instinct. The observations of several travellers north of the Great Slave Lake have resulted in the belief that "in spring the Barren Ground Caribou seek the coast of the Arctic Ocean, and remain near the salt water until about September." But this idea is much too circumscribed.

The explorations of Mr. J. B. Tyrrell, of the Canadian Geological Survey, have proved conclusively that the universal herd of the Great Slave Lake region does exactly as did the universal buffalo herd of 1871. It moves northward in spring for a given distance only, stops at will, spends the summer, and in the early winter moves southward. On July 30, 1893, Mr. Tyrrell saw a vast assemblage of Barren Ground Caribou at Carey Lake (Latitude 62° 10' and Longitude 102° 45'), nearly 500 miles from the Arctic coast. A herd of several thousand animals was composed of females with young fawns, young females and males of all ages, the lofty antlers of the latter being noticeably prominent. This herd was then only sixty miles north of the southern edge of the Barren Grounds.

The most impressive published description of a caribou migration is from the pen of Mr. Warburton Pike. It is a relation of what he saw on Lake Camsell, sixty miles north of the eastern end of Great Slave Lake, in 1889, and refers to the southward movement to the timbered regions,

<sup>1</sup> *Rangifer pearyi*.

where the lichens growing upon the trees afford subsistence in winter when the ground mosses are buried under snow and ice.

"From what I could gather from the Yellow-Knife Indians," says Mr. Pike in "The Barren Grounds of Northern Canada," "and from my own personal experience, it is late in October that the great bands of Caribou, commonly known as *La foule*, mass upon the edge of the woods, and start for the food and shelter afforded by the stronger growth of pine farther southward.

"Scattered bands of Caribou were almost always in sight from the top of the ridge behind the camps, and increased in numbers till the morning of October 20, when little Baptiste, who had gone for firewood, woke us before daylight with the cry, '*La foule! La foule!*' (The throng.) Even in the lodge we could hear the curious clatter made by a band of travelling Caribou. *La foule* had really come, and during its passage of six days, I was able to realize what an extraordinary number of these animals still roam the Barren Grounds.

"From the ridge we had a splendid view of the migration. All the south side of Mackay Lake was alive with the moving beasts, while the ice seemed to be dotted all over with black islands, and still away on the north shore, with the aid of the glasses, we could see them coming like regiments on the march. In every direction we could hear the grunting noise that the Caribou always makes when travelling.

"The snow was broken into broad roads, and I found it useless to try to estimate the number that passed within a few miles of our encampment. We were just in the western edge of their passage, and afterward we heard that a band of Dog-Ribs, hunting some forty miles to the west, were at this very time in the last straits of starvation, only saving their lives by a hasty retreat to the woods. This is a common danger in the autumn, as the Caribou, coming in from the Barren Grounds, join together in one vast herd, and do not scatter much till they reach the thick timber.

"The Caribou, as is usually the case when they are in large numbers, were very tame, and on several occasions I found myself right in the middle of a band, with a splendid chance to pick out any that seemed in good condition. . . .

Notwithstanding all the tall stories that are told of their numbers [the buffaloes], I cannot believe that the herds on the prairie ever surpassed in size *La foule* of the Caribou."

**Size and Antlers.**—At present the size of the Barren Ground Caribou appears to be a matter of opinion rather than of observation and record. In the hope that some one will come forward and disprove it, I venture to make the assertion that no one ever has weighed a whole, full-grown male specimen. We have a few figures of "dressed" weight, and various "abouts," but really useful facts are lacking. It is currently believed that the Barren Ground Caribou of northern Canada is about one-third lighter than the woodland species of Ontario and Quebec. If this be true, and we may judge by our own woodland bull, which unquestionably was a large one (48 inches high, weight, 261 pounds), then the male Barren Ground animal may be set down as weighing 174 pounds. For the Greenland caribou and Grant's caribou, this weight surely is too low; for the skulls and skins of both those species indicate a greater weight. On the Alaska Peninsula Mr. C. H. Townsend weighed a dressed specimen of *Rangifer granti* and estimated very carefully the weight of the viscera, with the conclusion that the live weight of the animal was 410 pounds.

For their body size, Barren Ground Caribou

have very large antlers. They sweep back so far, rise so high and spread so widely that they have the effect of magnifying the height and bulk of the wearer. As will be seen by the following measurements, the antlers of the Barren Ground species are longer than those of the woodland, but with fewer points, and in most cases less palmation. In the series of plates of all species published by Mr. Madison Grant in his valuable paper on "The Caribou" (Report of the New York Zoological Society, 1902), one of the most striking differences between the two groups is the tree-top appearance of all woodland antlers, and the open, arm-chair effect of the Barren Ground types.

**The Reindeer in Alaska.**—In 1887 Mr. Charles H. Townsend advised the government<sup>1</sup> that it would be a very beneficial and humane proceeding toward the Eskimo tribes of western Alaska to import a large number of domestic Reindeer from Siberia, and teach the natives how to care for and use them. Through the heroic efforts of Dr. Sheldon Jackson, General Agent of Education in Alaska, this advice was promptly followed under the auspices of the Bureau of Education; but the first fund of \$2,000 came from private sources, and was expended in 1892-3. The initial Congressional appropriation, of \$6,000, was expended in 1894, but since

<sup>1</sup> The Cruise of the *Corwin* in 1885, p. 88.

### MEASUREMENTS OF LARGE ANTLERS.

All measurements in inches.

			LENGTH OF MAIN BEAM.	WIDEST SPREAD.	POINTS.	OWNER.
GREENLAND CARIBOU . . .	<i>R. groenlandicus</i> ,	W. Greenland, . .	52	41½	21	Author's collection.
BARREN GROUND CARIBOU . . .	<i>R. arcticus</i> ,	. . N. Labrador, . .	60	34	18	National Museum. <sup>2</sup>
BARREN GROUND CARIBOU . . .	<i>R. arcticus</i> ,	. . Circle City, Alaska,	54½	36	16	G. R. Anchors.
GRANT'S CARIBOU . . .	<i>R. granti</i> ,	. . . Alaska Peninsula,	33⅞	35⅞	27	American Museum. <sup>2</sup>
NEWFOUNDLAND CARIBOU . . .	<i>R. terraenovae</i> ,	. Newfoundland, .	41	36	36	Madison Grant.
WOODLAND CARIBOU . . .	<i>R. caribou</i> ,	. . Northern Canada,	35½	21	30	Robert Gilfort.
MOUNTAIN CARIBOU . . .	<i>R. montanus</i> ,	. S. Brit. Columbia,	35	21	31	
OSBORN'S CARIBOU . . .	<i>R. osborni</i> ,	. . N. Brit. Columbia,	44½	38½	43	Daniel Carter Beard.
KENAI CARIBOU . . .	<i>R. stonei</i> ,	. . . Kenai Peninsula,	45½	34½	56	Harry E. Lee.

<sup>2</sup> From "The Caribou," by Madison Grant. Report of the New York Zoological Society, 1902.

1899, the amount granted annually has been \$25,000.

From 1892 to 1902, 1,580 Reindeer were imported from Siberia and 144 from Lapland, from which 6,116 fawns have been born in Alaska. Dr. Jackson states that "the animals born in Alaska are developing into larger and stronger animals than their parents." Of the whole number of Reindeer, 2,692 have been sold, butchered or lost by death. On May 1, 1903, the total number remaining alive in Alaska was 5,148. The number of fawns born in 1902 was 1,654.

The Reindeer experiment has been wisely conducted, on good business principles, and is an unqualified success. There are nine Reindeer stations, extending from Point Barrow, on the Arctic Ocean, to Eton Station, near St. Michael, on Norton Sound. The Laplanders who were taken to Alaska to educate the natives in the care and use of Reindeer, have done their work conscientiously, and the Eskimo have eagerly embraced the opportunity to acquire a domestic animal, good for use and for food, to take the place of the vanished walrus and Barren Ground caribou.

On the whole, the systematic introduction of Reindeer along the northwest coast of Alaska—now almost barren of wild life fit for human food—is one of the most humane and sensible measures ever undertaken for the children of the cold. If this industry is further fostered, and diligently pursued, its ultimate value in the promotion of the moral and material welfare of the Eskimo is beyond calculation. The multiplication of the herds in the hands of private owners means a great increase in the animal food supply, less dependence upon the foods of civilization, a greater measure of general prosperity and contentment, and in the end, far less expense to the government in the form of annual maintenance for starving natives.

**The Moose**<sup>1</sup> is the largest animal of the Deer Family, living or extinct. Even the Irish elk, with antlers which, in at least one specimen, spread 9 feet 3 inches, was a smaller animal. It is a satisfaction to know that the most colossal deer that ever trod the earth is alive to-day, and an inhabitant of our continent.

<sup>1</sup> *Alces americanus*. Called in Europe, the "Elk"; and our Elk is there called the "Wap'ti." See Frontispiece.

It is not, however, an easy matter to convey a truthful and adequate impression of this antlered giant of the north. The young specimens occasionally seen for a brief season in zoological parks and gardens are scarcely more than suggestions of the adult animal. The mounted groups in our large museums do indeed represent its full size; but to be fully appreciated, the Moose must be seen alive, adult, full of strength and purpose, striding like a four-legged colossus through the evergreen forests of Canada or Alaska, or swinging away at incredible speed from the dangers of the chase.

Imagine, if you can, an antlered animal standing between six and seven feet high at the shoulders, its legs quite four feet long, its neck and body covered with a heavy thatch of coarse, purplish-gray hair from three to six inches long, and its huge head crowned with massive antlers spreading from five to six feet in width. Its head is among the lower branches of the forest, and its long legs stride with indifferent ease over fallen tree-trunks which to the hunter are barriers to be climbed over, slowly and laboriously.

The Moose can instantly be recognized by its broad, square-ended, overhanging nose, large ears, high hump on the shoulders, and long, coarse, smoky-gray hair. The adult male is further distinguished by antlers that are enormously flattened and expanded, in a form popularly known as "palmation."

The Moose is not a grazing animal, like the elk, and most other members of the Deer Family. It lives by "browsing," or eating the bark, twigs and leaves of certain trees, and also moss and lichens. It is strictly a forest animal, and is never found on open, treeless plains. It is very fond of still water, and is much given to frequenting the small lakes and ponds which abound in some portions of its home. It is as fond of wading in shallow water as a boy, and is a ready and powerful swimmer. It loves to feed upon lily pads and stems, and moose hunters have assured me that it even seeks the bulbs growing in the muddy bottom.

Except in Alaska, the majority of Moose killed by hunters are shot from ambush beside ponds, or from canoes. Frequently, Moose that are surprised when wading and feeding in shallow water, make the mistake of rushing into deep water, to escape

by swimming, when they are easily overtaken, and either killed, captured, or photographed.

In the autumn months, the northeastern Moose hunter sometimes makes a horn of birch bark, at nightfall conceals himself beside a pond, and imitates the call of the cow Moose until a bull is actually attracted within shooting distance. The cry of this animal is a prolonged, resonant bawl, ending in three or four hoarse grunts.

The accompanying map shows that the Moose is yet found in northern Maine, New Brunswick, Canada, Manitoba, northern Minnesota, northwestern Wyoming, Idaho, British Columbia, Alberta, Athabasca, Yukon and Alaska. It shows only localities known to have been inhabited in 1902. In none of these, however, are Moose so abundant as in Alaska, around Cook Inlet. The southern limit of the Moose in North America is the head of Green River, Wyoming, Latitude 43°, Longitude 110° W., corresponding to the latitude of Albany, New York.

Below Alaska, the favorite hunting-grounds for Moose are Maine, New Brunswick, the upper Ottawa River country of Canada, and northwestern Manitoba. In view of the great number of hunters—estimated at ten thousand—who annually hunt and fish in Maine, of whom a large proportion hold licenses that permit the killing of one bull, the persistence of the Moose in Maine is really wonderful. During the past nine years the number of Moose transported by the railways of Maine have been as follows:

1894.....	45
1895.....	112
1896.....	133
1897.....	139
1898.....	202
1899.....	166
1900.....	210
1901.....	259
1902.....	244

Total ..... 1,510

In all probability, fifteen hundred more were killed and consumed within the state, and not accounted for in any permanent records.

The young of the Moose—always spoken of as a "calf," its mother being called a "cow"—is born in May, and at first is a very grotesque-looking creature. Its enormously long, loose-jointed legs are attached to an abnormally short and diminutive body. The neck is so short the

creature cannot put its nose to the ground without kneeling. Its hair is woolly and brick red, or "sandy," like that of a buffalo calf.

A Moose calf which I once owned, and measured when seven weeks old, had the following dimensions:

Height at shoulders.....	37 inches.
"    "    hips.....	31    "
Length of head and body.....	42    "
Depth of chest.....	11    "
Length of foreleg to elbow.....	26    "
Weight.....	79 pounds.

At one year of age, if not stunted in growth, a Moose stands from 4 feet 9 inches to 5 feet in height at the shoulders, where it has developed a lofty hump. On August 14, 1901, the largest of six Moose in the New York Zoological Park, each one about fifteen months old, measured as follows:

Height, 5 feet 3 inches; length, head and body, 5 feet 9 inches. Length of tail, 3½ inches; depth of chest, 2 feet 2 inches. Horns, 4 inches long; weight, 330 pounds.

Any Moose which stands 6 feet 6 inches in height at the shoulders may be considered a very large one, a prize, in fact. The largest Moose of which I have a reliable record, was killed in New Brunswick, in 1901, by Carl Rungius, the justly-celebrated animal painter, and carefully measured by him with the following result:

Height of shoulders, 7 feet, exactly.
Length of head and body, 9 feet 7 inches.
Girth, 8 feet.
Length of head alone, 2 feet 9 inches.
Antlers small for so large an animal.

The largest antlers recorded up to this date (1903) came from the Kenai Peninsula, are now in the Field Columbian Museum, and have the following dimensions:

Spread at widest point, 78½ inches.
Greatest width of palmation, 16 inches.
Circumference of burr, 15 inches.
Greatest thickness of palmation, 2½ inches.
Length of skull, 28¾ inches.
Total number of points, 34.
Weight of antlers and dry skull, 93¼ pounds.

From the above figures, one can imagine the strength necessary to enable an animal to carry such an unwieldy load upon its head, and run at great speed for long distances over the roughest kind of timbered country.




Regarding the weight of adult Moose, very few exact observations have been recorded, or otherwise made available. A large Maine Moose killed by W. L. Miller of Bangor, weighed 1,123 pounds. A dressed carcass weighed by S. L. Crosby showed a weight of 1,009 pounds. (*Recreation Magazine*, IV, p. 89.)

By the time a Moose calf is a year old, it has taken on the colors of adult life, which consist of a mixture of blackish-brown on the head, neck and body, and yellowish-gray on the legs and under parts. The hair and mane is long, coarse

sense than any other species of deer with which I am acquainted.

Owing to the peculiar nature of the digestive organs of this animal, it cannot live long upon ordinary grass or hay, even when supplemented with the best tree-branches that its own native forests can supply. It is my belief that vigorous daily exercise is vitally necessary to the proper digestion and assimilation of their food. In captivity, even when fed on fresh green browse of the choicest variety, which they eat with relish, they usually die of gastro-enteritis, or inflamma-



DISTRIBUTION OF THE MOOSE IN NORTH AMERICA, IN 1903.

and stiff, and lies more like a thatch of straw than genuine hair. On the neck and shoulders it is six inches long. Under the throat hangs a long, ornamental strip of hair-covered skin, four inches long, called a "bell." In the adult male animal this bell is sometimes a foot in length.

The female Moose has no antlers, but in bulk she almost equals the proportions of the male. Out of every thousand females, only one has a "bell."

In captivity the Moose is naturally a docile animal, not foolishly nervous like most deer, but steady, confiding and affectionate. Moose are easily handled, and trained to drive in harness, and in contact with man manifest more common-

tion of the stomach and intestines. Green grass is fatal to them, and when fed on grain, hay and vegetables they soon become emaciated and die. Thus far the best results achieved in the maintenance of captive Moose on public exhibition have been in the Cincinnati Zoological Garden, where Superintendent S. A. Stephan has succeeded in keeping a pair for about five years. In great forest preserves, such as Blue Mountain Park, in New Hampshire, Moose do live, thrive and increase.

In a wild state, Moose browse upon many kinds of trees, but particularly upon birch, hemlock, spruce, alder, aspen, willow and maple. They reach the tender tops of tree saplings by

walking astride of them, and "riding them down," and in the manipulation of small branches, the use of the overhanging and prehensile nose is strikingly apparent. With their strong lower front teeth, used chisel-fashion, they gouge the bark off large branches, and feed upon it. In grazing on grass, or feeding upon ground mosses, a Moose must kneel in order to reach them.

During the deep snows of winter, Moose herd together in sheltered spots in the forest; and



Copyright by DALL DEWEESE.

ANTLERS OF AN ALASKAN MOOSE.

Spread, 68 inches. From an animal killed on the Kenai Peninsula, by DALL DEWEESE.

through their moving about in a small area, the snow is trodden down until they form what is called a "Moose yard."

Naturally, because of its grand proportions, and its massive antlers, the Moose has been to every hunter of big game a grand prize. Although difficult to find and approach within easy rifle-shot, when approached it is killed easily and without danger. During the past five years, this species has been fairly protected throughout the eastern half of its range, and in 1902 this protection was by Act of Congress extended over

the whole of Alaska. Without real protection, ten years' time surely would see this magnificent animal, which Nature has been millions of years in bringing to perfection as we now see it, practically exterminated throughout North America. In 1900 the legislature of the state of New York appropriated \$5,000 to be expended in restoring wild Moose to the Adirondack wilderness, from which the species was exterminated by man, forty years ago. Up to September, 1903, fifteen head of young Moose had been purchased, chiefly in Canada, taken to the Adirondacks, and liberated. Although the responsible guides and guides' associations are using all their influence to secure the protection of the liberated Moose and elk, already have individuals of both those species been shot. It is greatly to be feared that the well-meant efforts of the state, and also of public-spirited private individuals, will accomplish little else than to furnish more meat for lawless persons who kill until they are caught, and then plead that they killed their Moose and elk "by mistake!" It is also to be feared that the Adirondack Moose will migrate northward into Canada, and remain there.

It remains to be seen how much the real men of the Adirondacks are going to accomplish against the Moose-killers and their supporters.

The Alaskan Moose has obtained a place in the annals of natural history to which its title is, at the least, very questionable. It has been described as a new species (*Alce gigas*), and a giant besides; and because of this, and its really immense antlers, it has dwarfed prevailing ideas regarding the more southern species (*A. americanus*).

For the exaggerated ideas of this animal that now quite generally prevail, its antlers are perhaps chiefly responsible. Occasionally they are of great size and weight, exhibiting enormous spread (from 70 to 78 inches), wide palmations and also great thickness (from 1½ to 2 inches). Their maximum dimensions considerably surpass those of antlers from more southern individuals. In addition to all this, they occasionally show freaky development in the shape and set of the brow antlers; and occasionally the main shovel throws out a palmated spur of striking form and size. Seen from the front, it often happens that the antlers of an Alaskan Moose present a chaotic

jumble of tines and palmations. Occasionally these odd forms are also found among the moose of Ottawa and New Brunswick.

But in Alaskan Moose antlers, freaky development is exceptional, and the real type is the same as that found on the moose of Nova Scotia, Manitoba and Minnesota. The largest antlers on record up to this date are perfectly regular. Apparently the Alaskan Moose find in summer an abundant supply of some food which is particularly rich in horn-producing properties, and their enormous and freaky antlers are the result.

Regarding the size of Alaskan and other moose, it is well to weigh the best available evidence.

So far as I am informed, the largest moose ever killed and measured by thoroughly experienced and reliable hands is the one already referred to which was shot in New Brunswick by Mr. Carl Rungius, the painter of American animals, whose knowledge of the external anatomy of that animal is, as many believe, second to that of no other man. The accuracy and fairness of Mr. Rungius' measurements of the animals he has so long studied in their wild haunts, is beyond question. According to Mr. Rungius, the moose referred to above stood precisely 84 inches high at shoulders, and had a girth of 96 inches; but "for so large an animal its antlers were rather small."

The following measurements of moose, in inches, are of interest in determining the real value of prevailing impressions regarding the Alaskan animal, and its right to specific rank by reason of its great size:

BY WHOM SHOT AND MEASURED.	LOCALITY.	SEX.	HEIGHT AT SHOULDERS.	GIRTH.	LENGTH OF HEAD AND BODY.
Carl Rungius, . . . .	New Brunswick, <sup>1</sup> . .	Male	84	96	115
Dall DeWeese, . . . .	Alaska, <sup>2</sup> . . . . .	Male	80 $\frac{3}{4}$	91 $\frac{3}{4}$	119 $\frac{3}{4}$
L. L. Dyche, . . . . .	Minnesota, <sup>1</sup> . . . .	Male	78 $\frac{1}{2}$		
A. J. Stone, . . . . .	Alaska, <sup>2</sup> . . . . .	Male	77 $\frac{1}{2}$		106

Until the enactment of the national law of 1902 for the preservation of wild animal life in Alaska, the huge antlers of the moose of Alaska threatened to cause the annihilation of the species in that territory. "Record heads" and "record antlers" began to be sought for by those who were able to buy them at high prices, and very promptly moose-killing for heads and horns

became an established industry. The unfortunate fact that in many portions of southwestern Alaska Moose were easily found and killed, bore heavily against them. The Kenai Peninsula partook of the character of a moose "preserve," in everything save preservation.

In 1902, through the combined efforts of naturalists and sportsmen, Congress enacted a law for the protection of the wild animals of Alaska, very wisely charged the Secretary of Agriculture with its enforcement, and vested him with wide discretionary power. It was a great day for big game, and for all persons interested in the preservation of our grandest wild animals, when the fauna of Alaska came under the protection of Drs. C. Hart Merriam and T. S. Palmer, of the United States Biological Survey, who are specially charged with the enforcement of the Alaska game law. The killing of moose for salable heads promptly decreased. Excepting by prospectors and natives in great need of food, no moose, white sheep, goat, caribou or big brown bear may be killed in close season without a special license signed by the Secretary of Agriculture; nor can any skins, heads or antlers of protected game be transported from Alaska without permits.

The only thing now necessary for the protection of the valuable animals of Alaska is an annual appropriation of \$25,000 for the pay of game wardens, and legal expenses, and the placing of the entire salmon industry under the control of the United States Fish Commission.

**THE PECCARY FAMILY.**

*Tayassuidae.*


The wild swine of the world form a group which contains several remarkable forms.

The wart hog, of Africa, is the ugliest of all land animals, and its head is of such a remarkable form that at first sight it seems like one of male specimens collected on the Kenai Peninsula.

<sup>1</sup> *Alces americanus*. <sup>2</sup> Largest of several very large

the sports of nature. The red river-hog, of West Africa, is the most beautiful of all swine, and its immaculate red coat, and long, slender ears produced to infinity in the form of a waving pencil of threadlike hairs, renders this animal acceptable in any zoological garden.

**The Collared Peccary** is our nearest and best-known representative of the wild swine. Its northern limit is the Red River, and the valley of the Rio Grande, in Texas, and southward it ranges to Patagonia. In northwestern Sonora, it has recently been obtained by Dr. D. T. MacDougall in regions so dry, hot and barren of vegetation that it was a surprise to find it there. Its preference is for brushy, upland jungles, but



COLLARED PECCARY.

at the same time it frequents all available cover, from the fruitful hard-wood forests of Arkansas and Texas to the moist and hot jungles of Central and South America.

In Texas this animal is called the "Javelina," and hunting it on horseback with dogs is a sport not to be despised. When hotly pursued, the Peccary of Texas gladly dives into any rocky crevice or hole that is large enough to receive it. Both jaws of this animal are provided with tusks, of sufficient length and sharpness to make them dangerous weapons.

The courage and pugnacity of the Peccary are well known, and when threatened with attack by a drove, the boldest hunter does not hesitate

<sup>1</sup> *Tay'as-su ta'ja-cu.*

to climb the best tree that happens to be available. An enraged Peccary, athirst for blood, is to any one not armed with a rifle or a first-rate spear a formidable antagonist. But for their tusks and dauntless courage when attacked, these animals could not have long survived in forests infested by savage jaguars, pumas, wolves and ocelots. Truly, it seems as if this species represents the survival of one of the fittest.

In our southwestern states the regular food of the Peccary consists of acorns, pecans, farmers' crops, seeds and edible roots of many kinds, and (it is said) also frogs, lizards, snakes, and all other ground animals it can catch. If the musk gland situated on the top of the hind-quarters is cut out as soon as a Peccary is killed, the flesh will be saved from the musky flavor and odor which without this precaution would soon render it unpalatable.

The Collared Peccary derives its name from a ribbon-like band of white which encircles the animal about where the neck joins the shoulders. Other than this, the hair is of a black color, sprinkled with gray.

**The White-Lipped Peccary**<sup>2</sup> is a larger species than the preceding, with white hair on its upper lips. It is found only as far north as southern Mexico, but ranges southward to Paraguay.

## THE TAPIR FAMILY.

### *Tapiridae.*

In all the world there are at least five species of tapirs, only one of which is found in the Old World. In southern Mexico and Central America, we know of the existence of a species called Baird's Tapir (*Tapirus bairdi*), and in Central America one known as Dow's Tapir (*Tapirus dowi*), but of their life histories very little is known, and at present it is impossible to describe them adequately.

**The South American Tapir**<sup>3</sup> is so frequently seen in captivity, and is already so well known, that it may well be chosen as the representative of the only Family of odd-toed Ungulates existing on this continent. It takes kindly to captivity, grows rapidly, and always manages to look well-fed, and as sleek as a seal. Its color is a rich mahogany brown, its head is long and triangular, and its long, prehensile nose, ever soliciting something to eat, is strongly suggestive

<sup>2</sup> *Tay'as-su al-bi-ros'-tre.*    <sup>3</sup> *Tap'i-rus ter-res'tris.*

of the end of an elephant's trunk. The shoulder height of a full-grown animal is about 37 inches.

The species best known to the world inhabits Venezuela, the Guianas, Brazil, Paraguay, Uruguay and the northern portion of the Argentine Republic. Although tapirs are usually found along small and well-shaded rivers in the hot lowlands of the tropics, they are also frequently found on forest-covered mountains. They are exceedingly shy and wary, and under all circumstances are difficult to find. Without dogs it is almost impossible to outwit them. When at-

tacked they always head for the nearest stream, and plunge into the water for concealment. Their food consists of soft and fleshy plants that grow in or within easy reach of streams, and in dense forests where the humidity is great. The flesh of all tapirs is said to be very palatable, and in South America it is much sought by hunters.

The South American Tapir thrives in captivity, either with a bath-tank or without, and breeds. In 1903 a pair bred in the National Zoological Park, at Washington, and the offspring survived.

## CHAPTER IX

# THE ORDER OF WHALES AND PORPOISES

### CETE

To some persons who are beyond the reach of large museums, or a complete work on natural history, the whales, dolphins and porpoises seem very far away. To those who live far from the sea, it might seem justifiable to omit them from our list; but, inasmuch as all Americans travel, and nearly every reader of this book is certain to observe some of the great sea-mammals disporting in the waves of their ocean home, it is necessary to give them a brief notice.

The salt waters of the world are inhabited by what is really a great array of species of fish-like mammals, some of which are the largest creatures that ever inhabited the earth. It is a satisfaction to know that even the largest of the great extinct lizards of North America did not equal the gigantic bulk of a ninety-foot sulphur-bottom whale of our Pacific coast.

Although the Cetaceans are very fish-like in form, and also in mode of life, they are warm-blooded mammals, which breathe air instead of water, drown if submerged too long, bring forth their young alive, and nourish them with milk from their own bodies. For the protection of their flesh and vital organs from the cold of Arctic waters, they are completely enveloped in a thick layer of fat, called "blubber," which lies under the skin, and is impervious to cold. It is as if a man had a layer of felt an inch thick under his skin.

All Cetaceans are destitute of hair, and in most cases the skin is as smooth as plate-glass. The great majority of them have teeth, but many are toothless. Except the whales of greatest commercial value, little is known of the habits of Cetaceans generally. It is very difficult to study creatures that make their home in the sea, and can be closely studied only when killed. Nevertheless, quite a number of interesting facts regarding these strange animals have been brought together, chiefly by observing whalers. Their four Families are as follows :

#### FAMILIES.

- |                   |              |   |  |   |  |
|-------------------|--------------|---|--|---|--|
| <b>CETACEANS:</b> | <i>Cete.</i> | } | <ol style="list-style-type: none"> <li>1. <b>Baleen Whales:</b><br/>(without teeth)<br/><i>Balaenidae.</i></li> <li>2. <b>Sperm Whales:</b><br/>(with teeth)<br/><i>Physeteridae.</i></li> <li>3. <b>Dolphins and Porpoises:</b><br/><i>Delphinidae.</i></li> <li>4. <b>Fresh-Water Dolphins:</b><br/><i>Platinistidae.</i></li> </ol> | } | <p>"Whalebone" Whales, of large size, without teeth. The mouth is provided with "baleen," commercially called "whalebone." This group includes the <i>Sulphur-Bottom</i>, largest of all whales, and about fifteen other species.</p> <p>Whales with a narrow, beak-like lower jaw, and <i>formidable</i> teeth. There are four species, varying in size from the <i>Pigmy Sperm Whale</i>, 12 feet long, to the great <i>Sperm Whale</i>, 80 feet long.</p> <p>This Family includes about thirty species of <i>Dolphins</i>, <i>Porpoises</i>, <i>Grampuses</i>, <i>Blackfish</i> and <i>Narwhals</i>. They vary in size from the five-foot common Porpoise to the thirty-foot Orca, or "Whale-Killer." All save a very few are harmless, but the Killer is the most savage and dangerous creature that swims the seas.</p> <p>The narrow-beaked dolphins of the Amazon and Ganges.</p> |
|-------------------|--------------|---|--|---|--|

## THE FAMILY OF BALEEN WHALES.

*Balenidae.*

If seen on land, any member of this Family would recall Falstaff's graphic reference to his own fleshy self,—“A mountain of mummy!”

In one respect, a large whale is like an iceberg. When seen in the water, only a small fraction of its bulk appears, and the remainder must be imagined. On the ocean, one sees nothing of a whale save a rather flat back, and a jet of dense vapor rising and curving back into the sea. Startling indeed would be the sight of a whale's bulk, if it could be seen in its entirety.

The largest and also the swiftest of all whales is the great **Sulphur-Bottom Whale**,<sup>1</sup> of the Pacific Ocean, found from northern California to Central America. So far as we know, this is the largest animal that ever lived upon this planet. Captain C. M. Scammon, one of the most observant and scholarly of all whalers, records the measurements of a specimen taken by him as follows: Total length, 95 feet; length of jawbone, 21 feet; girth, 39 feet; length of longest “whalebone,” 4 feet; weight of “whalebone,” 800 pounds; calculated weight of whole whale, 294,000 pounds; barrels of oil yielded, 110—not a large quantity.

The accompanying illustration shows the form of a baleen whale, and the peculiar outline of its enormous mouth. The whales of this Family live upon minute shrimp-like crustaceans, and swimming mollusks (shell-fish) belonging to the group known as pteropods (ter'o-pods) which float in myriads on or near the surface of the sea. To enable the sea-monster to feed upon these very small organisms, and secure them in a wholesale way, the roof of the mouth is provided with two great masses of thin, horny plates set edgewise on each side, and very close together. The lower edges of these plates (of “whalebone”) are frayed out into a mass of what looks like coarse, bristly hair, and these frayed edges unite into a web of filaments as long and as wide as the whole inside of the mouth.


In feeding, the whale swims through a mass of floating pteropods, with its mouth open; and the fringe of the baleen, hanging down upon the sides of the lower jaw, forms a perfect strainer for catching even the smallest creatures afloat.

<sup>1</sup> *Bal-ae-nop'ter-a sul-fu're-us*

The pteropods gather in a mass on the tongue, and presently are swallowed. When the mouth is shut, the plates of baleen fold in diagonally.

Captain David Gray has stated that sometimes the whale finds its food under water, at a depth of from sixty to ninety feet. In gathering it the animal dives, holds its breath like any air-breathing animal, and after an interval reappears at the surface to breathe, swallow the food collected, and rest before diving again. When whales are feeding in this manner, it is comparatively easy for whalers to approach them within striking distance, and harpoon them.

One of the most astonishing statements recorded of this animal is that sometimes when harpooned, and sometimes in sport, as well, it leaps out of the water, for practically its entire length! Captain Scammon states that a pair of Sulphur-Bottom Whales have been known to float side by side at the surface of the water,



BOW-HEAD WHALE.

*Balacna mysticetus.*

and caress each other by striking each other's bodies with their flippers, “the sound made by these gigantic love-pats being audible for miles.”

The young of a whale is called a “calf,” and usually the mother is very solicitous for the welfare of her offspring. She suckles it until it is able to seek other food than her milk.

**The Bow-Head Whale**, also called **Greenland**, and **Polar Whale**,<sup>2</sup> of the polar seas around the north pole, is known by the immense size of its head and the semicircular arch of its jaws. Its individual plates of baleen are sometimes 10 to 12 feet in length. This material is now scraped very fine, and mixed with the silk fibre of dress silks, to make the cloth rustle when worn, and also give it stiffness. It is now of such high value commercially that the baleen whales are being pursued as far north as vessels can go. When a vessel is having a run of luck, and striking Bow-Head Whales frequently, the oil is some-

<sup>2</sup> *Bal-ae'na mys-ti-ce'tus.*

times completely ignored, and the quest settles down to a hunt for whalebone alone.

Whale oil is no longer the valuable commodity it was forty years ago, but the hunt for baleen will ultimately exterminate all the whales of this Family. The Bow-Head Whale is of medium size, rarely attaining 65 feet, and usually runs under 50; yet it is uncommonly rich, both in baleen and oil. A large whale of this species is said to yield 275 barrels of oil, and 3,500 pounds of whalebone.

On the coast of Newfoundland there are now five whaling stations which during the summer season do a thriving business. Small whales of two or three species are killed in adjacent waters, towed to the stations, and hauled up on ways. In a single day a whale forty feet long is completely worked up, and practically every part of the animal yields a commercially-valuable product.

When a whale is struck by a harpoon, it dives deeply to escape its foes, and remains under water as long as possible. The comfortable period for a whale to remain under water is fifteen minutes, but in feeding below the surface, this is often extended to twenty-five minutes. Harpooned whales sometimes descend 300 feet and lie on the muddy bottom of a shallow sea for a period of from fifty minutes to an hour and twenty minutes.

But whalers know that their victim must sooner or later come to the surface, or drown. As a whale reaches the surface, it immediately discharges its breath from the blow-holes situated on top of its head. A whale does not spout water, but the breath which comes from its lungs is so heavily laden with moisture that at a little distance it looks like water, especially when it curves over and falls into the sea. It is this "spouting" which reveals the whale to its enemy in the "crow's-nest" of the whaling vessel, and causes him to shout joyously to those on deck, "There she blows!"

In addition to the above, the most important species of baleen whales are these:

**The Right Whale** (*Balaena glacialis*), of the cool waters around the north pole and the Atlantic Ocean, north and south, attains 70 feet, but usually runs under 50 feet.

**The Pacific Right Whale** (*Balaena sibboldii*) inhabits the North Pacific.

**The Humpback Whale** (*Megaptera nodosa*), of the Atlantic, off the United States coast, is the species most frequently seen from the decks of passenger steamers and stranded on our coast. Its usual length is from 45 to 60 feet.

**The Finback Whale** (*Balaenoptera physalus*), of the North Atlantic coast, attains 60 feet, but yields little oil, and is difficult to kill.

**The California Gray Whale** (*Rachianectes glaucus*), from the arctic seas to Lower California, attains 45 feet. It is fond of shallow water, and is savage and dangerous.

## THE SPERM WHALE FAMILY.

### *Physeteridæ.*

It is impossible to give in a few words a clear and adequate conception of the various localities inhabited by the great **Sperm Whale**.<sup>1</sup> It may be said, however, that it is a habitant of the warm seas of the globe, from the North Atlantic, around Cape Horn, to the North Pacific.

The Sperm Whale has an enormous, square-ended head, which constitutes one-third of its entire bulk. Under this great mass is the lower jaw of solid bone, shaped like a letter Y, the stem being fully armed with a double row of huge, conical teeth. In comparison with the great bulk of the head, the lower jaw seems absurdly small; but it is a formidable weapon, and whalers dread it.

In seizing a whale-boat, a man struggling in the water, or any other dangerous enemy, a Sperm Whale turns on its side or back, like a shark, in order to bring its lower jaw over its victim.

The largest Sperm Whales have measured from 80 to 84 feet. At birth they are from 11 to 14 feet long. Their food consists of fish of various kinds, and also squid. A young whale, only twenty feet long, which was taken on the coast of Cornwall, had in its stomach about 300 mackerel. The head of the Sperm Whale yields sperm oil, spermaceti, and teeth which are valuable for ivory. A substance called ambergris, of much value to druggists and perfumers, is obtained from the intestinal canal.

**The Sperm Whale Porpoise**, or "Pygmy Sperm Whale" (*Ko'gi-a*), is found on both the

<sup>1</sup> *Phys'e-ter mac-ro-ceph'a-lus.*



Atlantic and Pacific coasts of the United States. It is a true pygmy, adult specimens being but 15 feet long. They are so rare that their existence in the western Atlantic Ocean was not known until 1883, when a specimen was washed ashore at Spring Lake, New Jersey, and secured by the United States National Museum.

## THE DOLPHIN AND PORPOISE FAMILY.

### *Delphinidae.*

This Family contains a number of different groups of animals, some of which are sharply distinct, and are not called by either of those names. The porpoises are distinguished by their blunt noses, and the dolphins by their long, pointed noses and elongated, beak-like jaws. Unfortunately for our purpose, there are a few porpoises with long snouts, and a few dolphins with short, blunt noses; and consequently the two groups run together so confusingly that it is impossible to lay down any rules by which one may always be distinguished from the other. We shall therefore shorten our work by setting forth the species most worth knowing, and leaving the anatomical details of the different genera to be learned in the future.

**The White "Whale,"** or *Beluga*,<sup>1</sup> of the upper half of the northern hemisphere, is not really a whale, but a member of the Dolphin Family. It is creamy white all over, and 16 feet long; has several times been exhibited in aquaria and shows, and is known personally to millions of Americans. One of the fine specimens exhibited in the New York Aquarium in 1897 met its death from suffocation caused by a live eel becoming immovably fixed in its blow-hole, and shutting off its breath so suddenly that the mammal died before the fish could be removed. This species ascends the Yukon River, Alaska, for 700 miles, and is also an inhabitant of the St. Lawrence. Dr. Goode states that the food of the White "Whale" consists of such fish as flounders, halibut, cod, salmon, and eels, and also of squids and prawns. In the St. Lawrence River there is a fishery of considerable importance.

**The Blackfish**<sup>2</sup> is not a fish, but a jet black member of the Dolphin Family, 15 to 18 feet long, and is shaped very much like a small sperm

whale. The head has the same square-ended, sawn-off appearance, and a barely perceptible snout. It is one of the most abundant and important of the small cetaceans of the east coast of North America. Thousands of them have been stranded, or deliberately driven into shallow water, on Cape Cod, sometimes over a hundred in one school. The yield of oil from a single Blackfish varies from ten gallons to ten barrels. The jaw yields a fine quality of oil much used for sewing-machines, and known as porpoise-jaw oil. The value of a stranded Blackfish on Cape Cod varies from \$5 to \$40. (G. Brown Goode.)

Once on a voyage from South America to New York, we sighted a large school of Blackfish, travelling south, and playing by the way. Some chased each other, lazily, and half a dozen of them stood on their tails in the water, perfectly erect, with their heads six or seven feet high in the air, as if to look at the ship. Those so standing looked like big, black posts, all ready for wharf building.

**The Grampus, or Cow-"Fish,"**<sup>3</sup> of our Atlantic coast inhabits the same waters as the preceding species, but is not nearly so numerous, nor so stupid in getting stranded in shallow water.

Its color is slaty gray, variegated with irregular white markings, and its length is from 15 to 20 feet.

**The Killer "Whale,"** or *Orca*,<sup>4</sup> is the demon of the seas. This creature has the appetite of a hog, the cruelty of a wolf, the courage of a bulldog, and the most terrible jaws afloat. Its teeth are surpassed in size only by those of the sperm whale. It attacks whales of the largest size, and devours sea-lions, seals and small porpoises as a hungry longshoreman destroys saddle-rock oysters.

A full-grown Killer is from 16 to 20 feet in length, and can always be recognized by the great height of its back fin. The all-black High-Finned Killer, of the Pacific only, has a back fin six feet high. The colors of *Orcinus orca* are those of the pirate's flag of skull-and-cross-bones,—black and white, disposed as shown in the accompanying illustration. This species is found on both coasts of North America, and in the Arctic Ocean.

<sup>1</sup> *Delphinapterus leucas.*

<sup>2</sup> *Globicephala melas.*

<sup>3</sup> *Grampus griseus.*

<sup>4</sup> *Orcinus orca.*

The following quotation from Captain Scammon is the testimony of an eye-witness of the Orca in action:

“Three or four of these voracious animals do not hesitate to grapple with the largest baleen whale. The attack of these wolves of the ocean upon their gigantic prey may be likened in some respects to a pack of hounds holding a stricken deer at bay. They cluster about the animal’s head, some of their number breaching over it, while others seize it by the lips, and draw the bleeding monster under water; and when capt-

to the bottom where the water was five fathoms deep. During the struggle the mother became nearly exhausted, having received several deep wounds about the mouth and lips. As soon as their prize had settled to the bottom, the three Killers descended, bringing up large pieces of flesh in their mouths, which they devoured after coming to the surface. While gorging themselves in this wise, the old whale made her escape, leaving a track of gory water behind.”

The swiftness of the Killer is very great, and to all small Cetaceans this savage monster is a



CALIFORNIA GRAY WHALES ATTACKED BY KILLERS.

Drawn by J. CARTER BEARD, from Captain Scammon's narrative.

ured, should the mouth be open, they eat out its tongue.

“We once saw an attack made by three Killers upon a cow whale and her calf, in a lagoon on the coast of California, in the spring of 1858. The whale was of the California gray species, and her young was grown to three times the bulk of the largest Killers engaged in the contest, which lasted an hour or more. They made alternate assaults upon the old whale and her offspring, finally killing the latter, which sank

genuine terror. An eminent naturalist named D. F. Eschricht, who devoted much attention to the Cetaceans, states that he knew one of these animals to capture and swallow alive, and in quick succession, four small porpoises, while from the stomach of another Killer, but sixteen feet long, were taken fourteen seals! In Bering Sea the Killer destroys large numbers of fur seals, and when walrus were plentiful, even made war on them, also. On the Atlantic coast, it was, until recently, a common occurrence for

a band of Killers to chase large schools of blackfish and porpoises into shallow water. They also persecuted the horse-mackerel, or tunny. The Killer is widely distributed, and his deeds of destruction have made him widely known and feared.

**The Dolphin.**—Few persons cross the Atlantic, or make a voyage of half a dozen days in any direction, without seeing a school of dolphins. In fact, it might almost be said that every voyage has its dolphins. As a rule, they do not appear until the passengers have recovered from sea-sickness, and are on deck, eagerly scanning the surface of the sea for living things.


To most voyagers, the sudden appearance of a school of dolphins is a thrilling sight. Hour after hour the eye scans the watery expanse, eager for a sign of life, or gazes with awe and fear into the dark, watery abyss below. Suddenly, out of the steep side of a green-topped wave leap forth a dozen shining, sharp-pointed forms. They seem joyous and full of power, like acrobats entering an arena. In sublime ignorance of man's rapacious nature, they confidently swim within twenty feet of the ship's side. They curve up to the surface, frequently leaping clear of the water, arch their bodies, breathe quickly, and dive again. For a few yards, perhaps, they race along under water, but in plain view, then some leap out again. How easily they keep pace with the ship! Their mastery of old Ocean is so complete that it is a wonderful thing to see.

Sometimes the animals are so near the ship that the species can be determined to a certainty, especially those which are marked by light colors. However, it is no disgrace to any naturalist to declare his inability to say positively what species is alongside.

Dolphins are particularly fond of playing around the bow of a ship; but for some reason best known to themselves, they evince a decided preference for the out-thrusting bow of a sailing ship, and are not attracted so much by the high, perpendicular cutwater of a steamer, with no bowsprit or jib-boom.

A swift ocean steamship is not escorted very far, for such a promenade soon becomes tiresome; but I have seen a school of these interesting creatures circle about a sailing ship, and play around its cutwater for half an hour. It is a simple matter for an expert sailor to take a position on the martingale-guys of a ship, under the bowsprit, and harpoon a dolphin; but to me it has never seemed like a fair thing to do.

In North American waters there are about twelve species of dolphins, most of which are



Drawn by J. CARTER BEARD.

THE COMMON DOLPHIN.

from 6 to 7 feet in length, and but two or three species exceed 10 feet. The Short-Beaked Dolphin of the Pacific is the most beautiful species.

**The Common Dolphin,**<sup>1</sup> of the Atlantic Ocean, may well be taken as the type of the family of true Dolphins. It is the species that is most frequently seen, and the one longest known. It has a slender, cigar-shaped body, a small head, and its beak is long and narrow. Its length is from 6½ to 7½ feet, and in color it is dark gray above and dull white below. Dolphins generally feed upon small fish, and at times destroy great numbers of mackerel.

**The Common Porpoise,**<sup>2</sup> of the Atlantic Coast, is a jet black creature, blunt-headed, heavy in action, a veritable pig of the sea. It loves to roll about in the breakers, and loaf lazily in harbors and sheltered bays, and at river mouths. As before stated of porpoises gener-

<sup>1</sup> *Delphi'nus del'phis.*    <sup>2</sup> *Pho-cae'na com-mu'nis.*

ally, this animal does not leap from the water, in sheer enjoyment of a "life on the ocean wave," but heaves itself to the surface just high enough to bring its blow-hole out of the water, gives a loud puff or snuff, and then rolls heavily below.

This Porpoise is the species most frequently seen by summer visitors on the Atlantic coast, and in various localities it is variously named. It is known as the **Herring Hog**, **Snuffling**

ivory tusk, which is from 6 to 8 feet long, is twisted throughout its length, from left to right, and is developed only in the male.

The Narwhal's teeth, aside from a few that are merely rudimentary, are reduced to a single pair, lying horizontally in the upper jaw. In the female they remain permanently concealed. In the male the right tooth usually remains similarly concealed, but the left is enormously developed into the tusk just mentioned. Hav-



THE NARWHAL, ADULT AND YOUNG.

**Fig, Puffer and Snuffer.** Its length seldom exceeds 4 feet 6 inches. It feeds upon fish, particularly on species like the herring and menhaden, which run in schools, and is said to be very destructive. Its flesh is very dark, its blood is almost black, and on the dissecting table it reeks of oil.

One of the strangest of all Cetaceans is the **Nar'whal**,<sup>1</sup> a creature 16 feet long, mottled black and gray, with a blunt-ended head, no back fin, and with a very long, straight tusk of ivory projecting straight forward from its head. This

<sup>1</sup> *Mon'o-don mon-o'-ce-ros.*

ing no other teeth, the creature is obliged to feed upon squids, jelly-fish generally, and small fishes that can be swallowed whole. It is found in the polar waters of the North Atlantic, and the Arctic Ocean north of the Old World, but is now rare in accessible waters. When Nansen and Johansen were retreating southward over the ice, after their dash toward the pole, each man with three dogs dragging a sledge with a kyak upon it, the first living creature actually observed by them was the Narwhal, in the lanes of water then rapidly forming in the great ice-pack, in Latitude 83° 36'.

## CHAPTER X

# THE ORDER OF SEA-COWS

### SIRENIA

In certain warm and deep rivers of the tropics and sub-tropics, where water plants grow abundantly and all nature seems at peace, there live certain species of water mammals of strange form and habits. The manatees and dugongs differ so widely from even their nearest relatives in other Orders, that it is not an easy matter to introduce them.

The body of a Sirenian is like that of a long-bodied seal. The neck is very large, but extends straight forward, and terminates in a small, blunt-ended head with very small eyes and lips so extensible and mobile in the manipulation of food that the artist who tries to draw their moods and tenses soon finds himself quite bewildered. There are no incisor or canine teeth, and the serrated molars are intended only for the bruising and cutting of tender plants.

There are front flippers of good dimensions, but they are wellnigh useless, and about as shapely and graceful as a pair of old shoes. Apparently they are made for use in gesturing rather than in work, for when the animal rests upon the ground, the flippers break squarely at two joints, and are folded under the body, backs downward! There are times, however, when the flippers are of some use in feeding, in holding food and conveying it to the mouth. Instead of hind legs, there is a broad, flat tail, nearly as wide as the body of the animal at its widest point. The skin is almost as naked as that of an elephant, and about one inch in thickness. When twisted and dried strips of it make practicable canes. The flesh is well-flavored, and wherever taken is eaten with relish.

Usually the Sirenians live in the lower reaches of rivers that flow into the sea, sometimes in water that is bitterly salt, frequently in brackish water, but in most cases quite above tidal influence, where the water is fresh and sweet. Never, so far as we know, do they live in shallow water, and as a rule they prefer a depth of about fifteen feet. So far as we know, only one species of the Order has ever inhabited a land of ice and snow. The divisions of the Order are as follows:

### THE ORDER OF SIRENIANS.

	FAMILIES.	SPECIES.	
<b>ORDER SIRENIA :</b>	<b>MANATEES,</b> <i>Triche'chidae :</i>	{	<i>Trichechus Latiostris</i> . . . . . Florida, Central America, Mexico, Cuba.
			<i>Trichechus Americanus</i> . . . . . South America to the Amazon.
			<i>Trichechus Senegalensis</i> . . . . . West Africa.
	<b>DUGONGS,</b> <i>Dugong'idæ :</i>	{	<i>Dugong Dugon</i> . . . . . Africa, Ceylon, India.
			<i>Dugong Australis</i> . . . . . Australia.
	<b>RHYTINA,</b> <i>Hydrodam-</i> <i>al'i-dæ :</i>	{	<i>Hydrodamalis</i> (or <i>Rhytina</i> ) <i>Gigas</i> . . . . . Bering Island ( <i>now extinct</i> ).

**The Manatee, or Sea-Cow,**<sup>1</sup> will not often be seen outside of museums, but it must be introduced here in order that the readers of this book never need ask, as do thousands of other persons —“What is a Manatee?”

<sup>1</sup> *Tri-che'chus lat-i-ros'tris*.

This creature, the only American representative of its Order except the extinct Steller's sea-cow, is a large and heavy water mammal, from 9 to 13 feet in length, and in form very much like a seal. It has a blunt muzzle, small eyes, and rather feeble, clumsy front flippers. Its tail is a

rounded disc, which in swimming forms a powerful propeller. When dry its skin is of a clean, slaty-gray color, but in the water it seems almost black. The bones are solid and heavy, and the ribs are very thick. The largest specimen ever taken and preserved in the United States was 13 feet in length, and must have weighed about 1,200 pounds. In the summer of 1903, a fine specimen about eight feet long was captured under a state permit in the Banana River, Florida, and placed on exhibition in the New York Aquarium. From time to time, others have been exhibited at various watering-places along the Atlantic coast.

The Manatee never comes upon land. Usually its home is chosen in the upper waters of some deep, quiet tropical river, above the influence of the tide, where there is an abundance of manatee grass and other water plants acceptable to it for food. It is herbivorous, and because its molar teeth are weak, and there are no front teeth, it is compelled to live upon aquatic plants which are tender as well as nourishing. Its food is always eaten under water, and when at home, its presence is generally revealed by the bits of plant stems and grass blades which escape and float to the surface. In captivity, the Manatee feeds upon lettuce, cabbage, canna leaves, celery tops, water-cress, spinach, eel-grass and ocean sea-weed.

Even to-day the Manatee is found in Florida, in the Banana, Sebastian and St. Lucie Rivers, and its wanton destruction is prohibited by state laws, under penalty of \$500 fine. Occasionally, however, a specimen is netted alive, under a state permit, for exhibition purposes. In the Sebastian River two of the great cold waves of the past ten years unfortunately killed several individuals. Farther south it is found about the Isle of Pines, Cuba, and along the east coast of Mexico, and Central America, while another species occurs in South America as far down as southern Brazil. The flesh of this animal is light-colored, and both looks and tastes like lean fresh pork.

As the result of several years of inquiry, I am convinced that, strange as it may appear, in Florida the Manatee really is being perpetuated. The sentiment in favor of its preservation is almost universal, and there is ground for the belief that this is largely due to the wise liber-

ality of the state authorities in granting a reasonable number of permits to capture specimens alive when the animals are ordered at high prices for public exhibition. I believe that there are more Manatee alive in Florida to-day than there were twenty years ago, even though at one time the species seemed doomed to speedy extinction in the state.

**The Dugong** is the only living Old-World representative of the Order Sirenia, and between it and the manatee the chief difference is found in the whale-like tail of the former. The Australian Dugong, which attains a length of 14 feet, once was so abundant along the coast of Queensland, between Moreton Bay and Cape York, that a regular fishery was established at Moreton Bay.

**The Rhytina, or Arctic Sea-Cow,** is of special interest to Americans because of the important part it played about the middle of the eighteenth century in the discovery of Alaska. In 1741, the Russian navigator, Captain Vitus Bering, was shipwrecked on Bering Island, and compelled to winter there. The majority of the crew of the *St. Peter* died of hardship, and the remainder also would have perished but for the presence of the great Arctic Sea-Cow, then seen for the first time. To George William Steller, the official naturalist of the ill-fated expedition, the world owes all it ever will know of the life history of this animal. Despite the sufferings he endured, he faithfully and laboriously reduced to writing everything that he observed of the ponderous animal whose flesh sustained the lives of the castaways.

The Rhytina was an animal closely resembling the dugong and manatee, but greatly exceeding the maximum size of either. Steller declared that "the full-grown animal weighs about 8,000 pounds," and from the skeletons that were collected on Bering Island in 1883 by Dr. Leonard Stejneger, and now on exhibition in the United States National Museum, we know that full-grown animals attained a length of between 20 and 30 feet.

This species was exterminated by whalers who sought it for food, aided by the natives who used both its flesh and skin. It was practically exterminated about 1780, but the last animal was not killed until 1854. (Nordenskiöld's "Voyage of the Vega.")



THE MANATEE (*Trichechus latirostris*).

Drawn by J. CARTER BEARD from a living specimen in the New York Aquarium

## CHAPTER XI

# THE ORDER OF TOOTHLESS MAMMALS

### EDENTATA

Near the bottom of the scale of terrestrial warm-blooded quadrupeds, is found the Order Edentata, so called because several of its members are toothless, and others are nearly so. It contains perhaps a greater proportion of odd and remarkable forms than any other Order, and all are found on the American continent. Many of them are so wonderful in form and habit that they well repay the effort necessary to make their acquaintance. The species fall into three Families, as follows:

	FAMILIES.	EXAMPLES.
ORDER EDENTATA.	ARMADILLOS, . . . . . <i>DAS-Y-POD'I-DAE</i> , . . . . .	{ Nine-Banded Armadillo. Six-Banded Armadillo. Three-Banded Armadillo. Giant Armadillo.
	ANT-EATERS, . . . . . <i>MYR-ME-CO-PHAG'I-DAE</i> , . . . . .	} Great Ant-Eater. Tamandua.
	SLOTHS, . . . . . <i>BRAD-Y-POD'I-DAE</i> , . . . . .	{ Three-Toed Sloth. Two-Toed Sloth.

### THE ARMADILLO FAMILY.

#### *Dasypodidae.*

With a few exceptions, armadillos are found only in South America. The southern half of that continent was once the home of a wonderful array of gigantic animals belonging to this Order. In the La Plata Museum of Natural History is a procession from the Past. It is a long row of earth-colored, dome-like shells of great thickness, some of them as large as small hogsheads, and curiously ornamented by a scalloped lower edge. Some are provided with huge tails that are studded with many big, pointed knobs, called tubercles. These curious objects are the remains of gigantic armadillos, now extinct, called Glyp'to-dons, which once roamed over the pampas of South America.<sup>1</sup> In many American museums, casts of the remains of one of these weird creatures may be seen in what is known as the "Ward Casts of

Fossils." The shell of the Glyptodon copied in plaster by Professor Ward is a nearly perfect dome, 5½ feet long, 4 feet wide and 40 inches high.

With but one exception, the armadillo of to-day is a small creature, finding shelter in burrows which it digs for itself in the earth. Its movements are nervous and spasmodic, and for a short distance it scurries over the ground quite rapidly, running on the ends of its claws, and dodging quite skilfully. Its legs are so short, however, it cannot run far, and when about to be overtaken by a dangerous enemy, it halts, and burrows in the ground with wonderful rapidity. It is not equipped for fighting, for it has no front teeth. Its claws are fit only for digging, and since it cannot climb trees, it prefers to live in burrows, on open prairies.

But Nature has not left these creatures without protection from their numerous enemies. The body is incased in a hard shell, composed of small plates of bone very cunningly joined together, which covers every portion save the breast and abdomen.

The head is protected by a plate placed on its upper surface, and the tail is incased in a chain


<sup>1</sup> A large Glyptodont, 7 feet long, has recently been discovered in Texas, and described by Professor H. F. Osborn as *Glyptotherium texanum*.



of bony rings. When attacked by a savage animal, the armadillo tucks its legs under the edge of the shell alongside its body, rolls into a ball, and as nearly as possible leaves nothing exposed save its shell. The creature thus becomes a living nut that is not to be cracked and eaten by every enemy that comes along.

If the shell is strong enough, the armadillo is safe; but if it is not strong and hard, nor en-

ranged northward, until in southern Texas and Arizona we find the northern limit of the group, and the only species found in the United States. There are three species of armadillos that from time to time appear, alive, in zoological parks, the nine-banded, six-banded, and three-banded. The largest species now living is so rare it is very seldom seen in captivity. It is the giant armadillo, of northeastern South America.



THE THREE-BANDED ARMADILLO (1-3), AND SIX-BANDED ARMADILLO,<sup>1</sup> (4).

Figures 1-3 represent half-grown specimens.

tirely perfect as an envelope, a jaguar or puma may possibly kill the animal and devour it.

The armadillos with the weakest armor have found it wise to avoid the forest home of the jaguar and puma, and live on the open plains, where they are less liable to be killed. To enable them to do this, Nature has provided them with long and powerful front claws, with which to dig burrows in the hardest soil.

It was in Argentina that the great armadillos of the past reached their highest point in size and abundance. From thence, smaller species

**The Three-Banded Armadillo,**<sup>3</sup> of Argentina, represents the highest degree of perfection attained, either past or present, by any member of the Family.

Its shell is very strong, and so perfect is its mechanism that when the animal is in danger, it makes of itself a round ball, so completely incased in horn that no four-footed enemy can penetrate it. Even the top of the head is protected by a shield which acts as a shutter when the animal rolls up, and wishes to close the only

<sup>1</sup> *Tol-y-peu'tes tri-cinc'tus.*    <sup>2</sup> *Das'y-pus sex-cinc'tus.*

opening leading into the shell. It gives one a very queer sensation to handle one of these living nuts, and note the marvellous ingenuity in design and skill in mechanical execution which has been displayed in providing this special means of protection for an otherwise defenceless creature.

Having such excellent defensive armor, the

our taste. The Nine-Banded Armadillo has a total length, from nose to end of tail, of about 26 inches, and in bulk is about the size of our opossum. In captivity its food is milk, boiled eggs, and chopped meat, but in a wild state it feeds upon a mixed diet of worms, ants, snails, beetles, small lizards, grasshoppers, and other insects. The young in a litter vary from six to ten.



THE GREAT ANT-EATER (LOWER FIGURES) AND THE TAMANDUA (UPPER FIGURE).

Three-Banded Armadillo does not often burrow in the ground, and it ranges freely by daylight. In running it touches only the ends of its claws to the ground, and the shell is held high. The head-and-body length of the adult animal is about 14 inches, and the tail measures  $3\frac{1}{2}$  inches.

The **Nine-Banded Armadillo**<sup>1</sup> ranges all the way from southern Texas and Arizona to Paraguay, and along the Rio Grande is so common that living specimens are sold at \$2 each. In Venezuela I found it burrowing on the open savannas, going down about four feet, in a hole seven inches in diameter. The flesh of this creature is well-flavored, and is generally esteemed as palatable food. Being in a state of perpetual hunger, we found Armadillo stew very much to

<sup>1</sup> *Ta'tu no'vem-cinctum*.

## THE FAMILY OF ANT-EATERS.

### *Myrmecophagidae.*

The ant-eaters form another Family of Edentates, also confined to South and Central America, and all its members are absolutely toothless. The most celebrated member of the group is the **Great Ant-Eater**.<sup>2</sup> Although it is very unlike a bear, it is sometimes called the **Ant-“Bear”**; and when once seen it is never forgotten. The most peculiar thing about it is the extraordinary length of its head, which in front of the eyes is prolonged into a slender beak, with the mouth and nostrils situated at its tip end. The opening of the mouth is just large enough to admit the blunt end of a lead-pencil.

<sup>2</sup> *Myr-me-coph'a-ga ju-ba'ta*.

The feature which comes next in oddity is the big, fleshy tail, covered with an enormous brush of coarse, wavy hair. The popular belief in South America that the Ant-Eater sweeps up ants with its tail in order to devour them in a wholesale way, is quite erroneous, for the tail serves a very different purpose. Its use is to cover the owner when asleep. When the animal lies down to sleep, the tail is flung over the body, and the long, wavy hair forms a thatch so thick that no other portion of the creature is visible. It looks like a pile of brown hay. A medium-sized specimen that lived for about a year in the New York Zoological Park measured 12 inches in length of head, the neck and body, 31 inches, and tail vertebrae, 26 inches.

In its wild state, the Ant-Eater feeds upon ants, which it devours in great quantities. In fact, Nature has provided this Family of animals to restrict the number of plague-like ants which, even with Ant-Eaters in the forests, are entirely too numerous. Its long and powerful front claws are very useful in tearing open ant-hills, and dissecting decayed logs, but as a means of defence they are quite inadequate. Neither are they well-formed to walk upon. The tongue is very long and slender, and can be thrust out 9 inches; but, contrary to innumerable misstatements, it is as clean and smooth as the tongue of a dog, and is *not* coated with sticky saliva, or anything like it.

This animal is very clumsy on its feet, and being defenceless, unable to climb, and too large to live in a burrow, it is a wonder that all the Great Ant-Eaters were not killed and devoured long ago, by jaguars and pumas. Although quite rare, even in South America, a goodly number of specimens find their way into captivity. Until settled down sensibly to a diet of chopped meat, milk, and eggs, they are difficult to keep alive. Our specimen persistently refused to eat ants.

The **Tamandua**<sup>1</sup> is a smaller Ant-Eater than the preceding species, of tree-climbing habits, with a proportionately shorter head, no long hair on its tail, and extremely large front claws. It is found in Venezuela, the Guianas, Brazil, and in fact the greater portion of the region of tropical forests on this continent south of Mexico. Its tail is prehensile, or grasping, and in climbing

<sup>1</sup> *Tam-an'du-a tet-ra-dac'ty-la.*

is used almost constantly. One of these creatures which I once kept in South America as a camp pet, became very friendly, and even affectionate, and when permitted would climb all over me, as if I were a new and very soft species of tree. In the accompanying picture, the Tamandua is represented by the small central figure. Its head-and-body length is about 24 inches, tail, 18 inches.

## THE SLOTH FAMILY.

### *Bradypodidae.*

The sloths inhabit the New World only; and the so-called "sloth" of Ceylon is not a sloth, but a slow lemur. All the real sloths belong to the Order of Edentates, and inhabit the tropical forests of Central and South America, from Costa Rica southward. The sloths are not really toothless, for they have five pairs of teeth in the upper jaw, and four in the lower.

One cannot look at a live sloth without thinking that Nature has but poorly equipped this animal to live in this murderous world. Its countenance is a picture of complete and far-reaching stupidity, its bodily form the acme of four-footed helplessness. It can neither fight, hide, nor run away. It has no defensive armor, nor even spines. It is too large to live in a hole in a tree, and too weak to dig a burrow in the earth. It is too tired to walk on its feet, as the monkeys do, so throughout its queer life it hangs underneath the branches of the trees in which it finds its food. Its feet are merely four hooks by which to hang. Since it feeds wholly upon leaves and buds, it lives in the tropical forests, where green leaves are plentiful and cheap.

The sloth dwells only in the tree-tops, among the monkeys and macaws. On the ground, it would be more helpless than a tortoise, and easily killed by any carnivore, or wild pig. In the tree-tops, it escapes the climbing ocelot by living far out on the ends of the branches; and it is fortunate for him that hawks, owls, and eagles are scarce in the forests wherein he dwells.

At this point, however, it is a pleasure to point out that Nature has done one special thing for the preservation of these odd creatures. The hair of a sloth is long, wavy and coarse, rather more like grass than hair, and in color and general appearance it is the best imitation of tree-

bark that has been given to any quadruped. This resemblance to bark is heightened by the fact that the back hair of many a sloth in its native forest has a greenish tint, like moss on a tree-trunk, due to the presence on the hair of living vegetable algae. This aids the sloth in escaping observation.

On the mighty Essequibo River, in British Guiana, I once made a special hunt for sloths. Having found it useless to hunt them by stalking through the dense and lofty forests, I took a leaky old canoe, an Indian to help furnish power, and paddled fifteen miles and back. We followed the shores, going and coming, and secured eight specimens of the **Three-Toed Sloth**,<sup>1</sup> the one with a brown saddle-mark of short hair in the middle of its back.

We found them in the tops of low trees at the water's edge, spread-eagled on the outer branches, or hanging upside down, but always eating leaves. They did not know what it was to "take alarm," and try to escape. Judging by the awful deliberation of those that we saw in motion, I estimated that a really swift sloth could travel half a mile in twenty-four hours, if not sidetracked.


We shot some of our specimens, and others we took alive by cutting down their trees. One tree fell with its top in the river, and the sloth was carried four feet under water. But even the prospect of drowning did not make him hurry to the surface. To my amazement, he climbed up through the branches, slowly and deliberately, until at last, with dignity entirely unruffled, he appeared above the surface, and looked at me with a most disgusted expression on his wooden countenance.

Sloths eat so slowly that before one meal is over it is time for the next, so that their meals overlap one another.

The Three-Toed Sloth is not found above the Isthmus of Panama, but two other species inhabit Central America as far north as Nicaragua. It is considerably smaller than the next species, having a head-and-body length of 21 inches, while the spread of its outstretched arms, exclusive of the claws, is 32½ inches. The tail is so very short that it seems to be wanting entirely, but in reality its length is 1½ inches.

<sup>1</sup> *Brad'y-pus tri-dac'ty-lus*.

The **Two-Toed Sloth**,<sup>2</sup> also called **Hoffman's Sloth**, ranges northward as far as Costa Rica. It is the largest living member of the Sloth Family, and its appearance is well shown in the accompanying picture of a specimen kept in the Zoological Park. It inhabits the same regions as the preceding species, but is less common. It is occasionally seen alive in large zoological gardens, and when once properly acclimated, lives in captivity very well. Usually, however, it is difficult to keep alive. In captivity its food is chopped carrots, cabbage, lettuce, and boiled rice. A sloth usually sleeps sus-



SANBORN, Photo. N. Y. Zoological Park.

TWO-TOED SLOTH.

pended from a branch, but at the same time it always seeks a position in which it can rest its body on a branch below parallel with the one to which it clings.

In prehistoric times, a Family of gigantic ground sloths, called **Meg-a-the'ri-ums**, creatures as large as the largest rhinoceros, lived on the pampas of southern South America and also in the southern United States. Plaster casts of the entire skeleton of the most celebrated species (*Megatherium euvicri*), from South America, 17 feet 9 inches long, are now to be found in many American museums.

<sup>2</sup> *Cho-loe'pus hoff'man-i*.

## CHAPTER XII

# THE ORDER OF DIGGERS

### EFFODIENTIA

This Order contains only a very small number of genera and species, all of which are confined to the Old World. They are the pangolins of Africa and the Far East, and the aard-varks of Africa. Until very recently, these animals have been classed with the ant-eaters, sloths and armadillos, in the Order Edentata, or toothless mammals. But both in internal and external anatomy they differ widely from their very distant American relatives.

The latest and most exact classification assigns them to a new and wholly independent Order, called Ef-fo-di-en'-tia, which means "Diggers." Its divisions are as below:

	FAMILIES.	EXAMPLES.
<b>ORDER EFFODIENTIA.</b>	{ PANGOLINS, . . . <i>MANI-DAE</i> , . . . . .	{ Manis, or Pangolin, of India. <i>Manis pentadactyla</i> .
	{ AARD-VARKS, . . . . . <i>O-RYC-TE-RO-PODI-DAE</i> ,	{ Aard-Vark, of South Africa. <i>Orycteropus a'fer</i> .

### THE PANGOLIN FAMILY.

#### *Manidae.*


One good look at a pangolin, or manis, is enough to arouse curiosity, and provoke inquiry. Like the armadillo, it is one of the wonders of the living world,—absolutely toothless, dwelling upon the earth, surrounded by savage and merciless enemies, but safe in the protection of a complete suit of plate armor, and powerful claws for digging. There are about seven species in this Family, scattered all the way from China and Borneo to South Africa, excepting the break in the chain caused by the deserts of North Africa and Arabia. Of the three African species, two are distinguished by the extreme length of their tails, and one by its great size, six feet in length, which entitles it to the name **Giant Pangolin**.

The **Indian Pangolin**, or **Manis**,<sup>1</sup> of Ceylon and India, generally in the lowland forests, may be chosen as the representative of this Family. My first feeling toward it is that of friendship and gratitude, for in the jungles of Ceylon a living specimen once furnished me entertainment, anxiety and sustenance.

My first Manis was brought by a native, who carried it in a bag over thirty-five hot and dusty

<sup>1</sup> *Man'is pent-a-dac'ty-la.*

miles. While in transit on man-back, the animal kept himself comfortably coiled, but when set free upon the ground he promptly uncoiled



AN INDIAN PANGOLIN, ROLLED UP.

and stood up for inspection. He was 36 inches long, including the tail, which measured 17 inches, and his weight was 18 pounds.

From the tip of his nose to the end of his tail, he was covered with broad, flattened shield-shaped plates, or scales, of clear, gray horn.<sup>2</sup>

<sup>2</sup> Museum specimens are usually of a yellowish-brown color.

Those plates, which were concave underneath and convex above, lay close down upon the skin and upon each other, and were arranged in rows or courses, perfectly imbricated (*i.e.*, joint-breaking) like the scales of a big fish, or a hawk's-bill turtle. We presently discovered that they were fully controlled by the voluntary muscles of the skin. The tail was very broad, measuring  $5\frac{1}{2}$  inches across where it joined the body, slightly hollowed underneath, and rounded on the top. It was a most useful appendage, and its special function was to protect the head.

In walking, the *Manis* carried his back very highly arched in the middle. The long and powerful front claws were bent under the feet, until they pointed directly backward, and were literally walked upon. The heavy tail barely cleared the ground, and the nose was always carried low, as if slyly searching for something. Often the creature stood erect on its hind legs, like a kangaroo, especially when looking about for insect food, and as it walked its armor clanked like that of an ancient mail-clad knight.

Whenever he found a colony of ants, he would begin to dig most industriously. After digging a short distance into an ant-hill, and exposing the interior, he would thrust his long and slender tongue into the passage-ways, and draw it out thickly covered with ants.

To me, the most wonderful thing about the animal was its means of protection from its enemies, for it cannot truthfully be called defence. Without some very special provision of Nature, a slow-moving, toothless and hornless terrestrial animal would fare badly in jungles inhabited by leopards, tigers, wolves, jackals and wild swine.

When I first endeavored to become acquainted with my *Manis*, he immediately tucked his head down between his four legs, brought his tail under his body and up over his head, held it there closely, and thus formed of himself a flattened ball completely covered with scale armor. When I undertook to uncoil him, I could not manage it alone, and called a servant to help me; but the tail clung to the body as tightly as if it had been riveted there. Then I called another man, and while I held the body, the other two pulled on the tail with all their strength, to uncoil it. But in vain. We wrestled with that small animal until we were fairly exhausted, and so great was the power of the tail that we gave up beaten.

From the very first, I had no end of trouble with my scaly pet. I could not tie him, for on no part of his body or limbs would a rope hold ten minutes without hurting him. During the day, he was reasonably quiet, but at night he was very restless, and anxious to go out ant-hunting. For the first night, I shut him up in the main room of the Rest House; and in the morning I found him fully ready to break through a hole he had dug with his big front claws in the ten-inch wall of solid masonry. Well may naturalists assign the Pangolins to the independent Order of Diggers!

The next night, I placed the Pangolin in a large tin box, well covered with boards. At three o'clock in the morning the village dogs raised such a row at the edge of the jungle that my servant went to them to investigate; and it was that animal. It had torn a hole in one corner of its tin prison, and escaped; and but for the very dogs that had so often annoyed me by trying to steal my specimens, it would have been lost to me forever.

#### THE AARD-VARK FAMILY.

This Family contains but one genus and two species, the **Cape Aard-Vark**,<sup>1</sup> of South Africa, and the **Ethiopian Aard-Vark**, of East Central Africa.

With their usual facility in misnaming wild animals, the Boer pioneers in Cape Colony bestowed upon the species found there the name Earth-"Pig," and it has become a fixture.

The Cape Aard-Vark is as much like a pig as it is like a jack-"rabbit," but no more. Cut off its extremely long and rabbit-like ears, cover it with imbricated scales to fit its body, and externally we will have a rather tall pangolin, about 5 feet long. Unlike the pangolins, the jaws are provided with teeth. The tail is long, thick and heavy, and its special use is not quite apparent.

In the usually wise economy of Nature, these insect-eating animals were developed in Africa for the special purpose of checking the ants of that region. Their powerful front claws enable them to dig with great success into the tall and also numerous ant-hills of Africa, and before the days of universal game destruction, the Aard-Vark was oftenest found where ant-hills were most numerous.

<sup>1</sup> *Orycteropus afer*.

CHAPTER XIII

THE ORDER OF POUCHED MAMMALS

MARSUPIALIA

An animal is said to be "low" in zoological rank according to the distance of its position below the highest types of animal life. Thus, a hairless, fish-like mammal, with very simple teeth, like a porpoise, is far lower than the monkeys and carnivores.

As we approach the Orders of mammals which we have been taught to place at the end of the list, we encounter some very strange forms, which are of greater interest to the special student than some higher forms which are duplicated many times over. Fortunately for our purpose, all the Orders of living mammals, save two, are represented in North America.

Although the Order Marsupialia is too extensive, and the majority of its members too far away, to justify its full exposition here, it is desirable to mention all its Families:

	FAMILIES.	HABITAT.	
<b>ORDER MARSUPIALIA.</b>	KANGAROOS, . . . .	<i>MA-CRO-PODI-DAE.</i> . . . .	{ Australia, New Guinea, Aru Is.
	PHALANGERS, . . . .	<i>PHAL-AN-GERI-DAE.</i> . . . .	Australia.
	WOMBATS, . . . .	<i>PHAS-CO-LO-MY' DAE.</i> . . . .	S. Australia.
	CAENOLESTES, . . . .	<i>E-PAN-OR'THI-DAE.</i> . . . .	South America.
	BANDICOOTS, . . . .	<i>PER-A-MELI-DAE.</i> . . . .	{ Australia and New Guinea.
	DASYURES AND TAS- MANIAN WOLVES, . . . .	<i>DAS-Y-U'RI-DAE.</i> . . . .	Australia.
	OPOSSUMS, . . . .	<i>DI-DEL-PHY'I-DAE.</i> . . . .	} North, Central and South America.
	MARSUPIAL MOLES, . . . .	<i>NO-TO-RYC'TI-DAE.</i> . . . .	Australia.

Of these eight Families, only two, the first and seventh, will be specially noticed.

Marsupials are distinguished from all other mammals by the fact that the female possesses in the skin of her abdomen a large, flexible pocket, or pouch, in which the nursing glands are situated, and in which the young are carried for a time after birth, until more fully developed. They differ from ordinary mammals in being without what is called a *pla-cent'a*, which is an arrangement of veins by which the blood of the mother circulates through the veins of the unborn young. In other words, in a marsupial, the blood of the mother does *not* circulate through the veins of the unborn young. As a result, at the time of its birth, the young marsupial

is a tiny creature, hairless, blind, and utterly helpless. Even the young of a large kangaroo looks more like a little lump of jelly than a highly organized living creature. One which I saw in the London Zoological Gardens was less than an inch in length, and no thicker than a lead-pencil.

The newly born young is taken by the mother, in her front paws, and placed in her pouch; and the half-formed creature, with a mouth specially formed for suction, attaches itself to the nursing gland, and so remains for many days, or even weeks. Slowly it grows, until it develops hair, and its eyes open. At length it becomes large enough so that it ventures to stick its little head out, and view the world. By and by it climbs

out, to take exercise, but jumps back again at the first alarm. In an animal which travels as far each day as the kangaroo, a pouch for the conveyance of the young is a great convenience.

### THE KANGAROO FAMILY.

#### *Macropodidae.*

In Australia, the land of queer things, nearly all the land mammals are marsupials. The Order includes the kangaroos, large and small,



E. F. KELLER, Photo.. National Zoological Park.

BRUSH-TAILED ROCK WALLABY (*Petrogale penicillata*).

Length, head and body, 28 inches ; tail, 24.

wombat, Tasmanian wolf, Tasmanian devil, koala, and many others. All kangaroos come either from Australia, Tasmania, or New Guinea, but one group of small wallabies extends its range to New Britain and the Aru Islands. The great majority of these creatures dwell on the ground in the open plains, or in the "bush"

of Australia. In northern Queensland and New Guinea are four species of Tree Kangaroos, which actually climb trees, and inhabit them.

The largest species is the great **Gray Kangaroo**,<sup>1</sup> also called "Old man" and "Boomer," which stands over 4 feet high, weighs nearly 200 pounds, and when frightened can leap twenty feet or more. The smallest species are the **Rat Kangaroos**, some of which are but 14 inches high. Despite their nocturnal habits specimens are frequently seen in captivity. One of the handsomest of all the species is the **Red Kangaroo**,<sup>2</sup> a creature about 4 feet high, frequently seen in captivity, and quickly recognized by its brick-red color, and fine, silky hair. Several small species of Kangaroos are called **Wallabies**, and the species figured herewith is a good representative of this whole Family.

The Kangaroo is a strange variation in form from the ordinary terrestrial mammal. Its extremely long, strong hind legs, and massive tail, also of great length, form a wonderful jumping machine. The tail not only assists the animal in leaping, but it also serves as a balancing pole, and keeps its owner from losing his proper position when in mid-air. It is reasonably certain that a Kangaroo without a tail would frequently overbalance when leaping, and turn somersaults. Kangaroos were once very abundant in Australia, but the general settlement of that country, and the systematic killing of the animals for their skins, which are used as leather for shoes, has so greatly reduced the number that now

one must go far inland in order to find them wild.

Most pouched mammals are strictly herbivorous, but some, like the opossum and Tasmanian wolf, are true flesh-eaters.

<sup>1</sup> *Mac-ro'pus gi-gan'te-us.*

<sup>2</sup> *Macropus ru'fus.*



## THE OPOSSUM FAMILY.

*Didelphyidae.*

The New World contains more than twenty species of omnivorous animals, varying in size from a large cat to a small rat, mostly provided with long, hairless tails that are fully prehensile, and always well clad with fine and abundant hair. In all species save a few, the female possesses the abdominal pouch to which every marsupial female is entitled, but in some species it is either rudimentary or wholly lacking. These animals are the **Opossums**, and while the majority of the species are confined to South America, our North American representative is about as widely known as all the tropical species combined.

The **Virginia Opossum**<sup>1</sup> is a typical marsupial, but differs widely from all the Australian members of that Order. Seemingly it is a dull-witted, slow-moving creature, and so ill-fitted by Nature either to fight or to run away, that it might be considered almost defenceless. But let us see what use this odd little animal makes of the physical and mental equipment which Nature has given it.

It eats almost everything that can be chewed,—wild fruit, berries, green corn, insect larvae, eggs, young birds and quadrupeds, soft-shelled nuts, and certain roots. It is a good climber, and has a very useful prehensile tail. It forages on the ground quite as successfully as a raccoon. Usually it burrows under the roots of a large tree, where it is impossible for a hunter to dig it out, but sometimes it makes the mistake of entering a hollow log. Like the bear and woodchuck, it stores up under its skin a plentiful supply of fat for winter use, when food is scarce and dear. Above all, the female has a nice, warm pouch in which to carry and protect her helpless young, instead of leaving them in the nest to catch their death of cold, or be eaten by some enemy.

The young of the Opossum vary in number from seven to eleven. Not until they are about five weeks old do they begin to venture away from the mother; but for a season they are very careful not to get beyond grabbing distance of her shaggy coat.

The Opossum is a very hairy animal. Its under fur is woolly and white, and the outer coat

<sup>1</sup> *Di-del'phis vir-gin-i-an'a.*

is straight, coarse, and tipped with black. The nose, lips, and half the ear are pinkish white, and the eyes are like a pair of shoe-buttons. The tail is naked, white, and strongly prehensile.

A large specimen has a head-and-body length of 15 inches, tail 12 inches, and the weight of a large specimen is 12 pounds. In the South, the flesh of this animal is much prized as food, and I can testify that when properly roasted, and served with nicely browned sweet potatoes and yellow corn bread, it is an excellent dish.

One habit of this animal is so remarkable and so widely known it has passed into a proverb,—



VIRGINIA OPOSSUMS.

“playing 'Possum.” When found by hunters, the Opossum deliberately feigns death, hoping to escape by being “left for dead.” Give it a tap on the head or back, and it stretches out, limp, motionless, and seemingly quite dead. Its breath is so short and feeble the thick fur almost conceals the chest movement.

When but a lad I killed my first Opossum in an Indiana forest, and had carried it by the tail for half a mile when we came to a rail fence. In climbing through, I noticed that the front claws of my Opossum caught on a rail, and held fast in a manner highly unbecoming in an animal that was honestly and sincerely dead. A close examination revealed the fact that my victim was only nominally dead. In other words, it was fully alive, and sharply watching for a chance to escape. This discovery led me to keep the animal alive in confinement, until finally it did escape.

The Virginia Opossum is the species found in the United States, from New York to Florida, and westward through the southern States to

Texas. In Mexico and tropical America several other species are found. Notwithstanding the persistent destruction of the Opossum, both for moonlight sport and for food, it still manages to survive throughout its entire original range, and bids fair to outlive the native American.

means *Mouse-Like Opossum*—is a South American species which is remarkable because of its diminutive size. The full-grown female specimen shown in the accompanying illustration, with a brood of seven hairless young clinging to the fur of her body, is about the size of an eastern chip-



E. R. SANBORN, Photo., New York Zoological Park.

MURINE OPOSSUM AND YOUNG.

About one-half life size.

As a pet, or cage animal, the Opossum shows off very poorly, and is rather uninteresting. In the daytime, its sole desire is to curl up into a furry ball, and sleep. If disturbed, it opens its pink mouth very widely, in silent protest, and as soon as the trouble is over, again tucks its head under its body, out of sight, and sleeps on.

**The Murine Opossum**<sup>1</sup>—a name which

<sup>1</sup> *Mar-mo'sa murina*.

munk. The abdominal pouch is wholly wanting in this species, and from birth the naked and almost helpless young must either cling to the fur of the mother or die. As they grow larger, they travel on the back of the mother, with their tendril-like tails clinging to her tail.

The specimen shown reached New York just as a score of others have before it,—hidden in the interior of a bunch of bananas!

CHAPTER XIV

THE ORDER OF EGG-LAYING MAMMALS

MONOTREMATA

*"There are more things in heaven and earth, Horatio,  
Than are dreamt of in your philosophy."*

There are two Families of mammals the members of which lay eggs, from which their young are hatched as are those of birds. They form the lowest order of mammals, and in one respect this group forms a good connecting link between mammals and birds:

FAMILIES.


<b>ORDER MONOTREMATA:</b> <i>Egg-Layers.</i>	}	DUCK-BILL, . . . <i>OR-NI-THO-RHYN'CHI-DAE.</i> . . .	}	The Ornithorhynchus or Duck-Bill.
		ECHIDNAS, . . . <i>TACH-Y-GLOS'SI-DAE.</i> . . . . .		}

**The Platypus, or Duck-Bill,**<sup>1</sup> is found only in Australia,—a land of queer things. Not only is it bird-like in laying eggs, but it also possesses webbed feet, and a flat, duck-like bill, from which it derives one of its popular names. The beak is of black horn, and the food is crushed between the cross-ridged plates of the lower jaw and the roof of the mouth.

This animal is about as large as a prairie-"dog," and its body is similarly shaped; but there the resemblance ends abruptly. Its front feet are webbed quite beyond the ends of the toes, and in digging, the outer edge of the web is rolled back underneath the foot, to expose the claws. The hind feet are webbed only to the base of the claws, and each is provided with a strong, sharp spur an inch long, which is said to be connected with a poison gland.

The tail is broad and flattened, well haired on the upper side, and almost naked below. The hair of the Platypus is dark brown in color. The outer coat is stiff and harsh outside, but the

<sup>1</sup> *Or-ni-tho-rhyn'chus an-a-ti'nus.*



THE PLATYPUS, OR DUCK-BILL.

inner is fine and soft. The length of head and body is 13 inches, tail, 5 inches.

The habits of the Duck-Bill are very similar to those of our old friend the muskrat. It inhabits quiet but deep pools of fresh water, burrows deeply into the banks, and is seldom seen save at nightfall. In its burrow it builds a nest for its young, and deposits two eggs, which are enclosed in a strong, flexible shell three-fourths of an inch in length by two-thirds of an inch in greatest diameter. When first hatched the young are blind and hairless, and the beak

is very short. The food of this creature consists of aquatic insects, crustaceans and worms.

The other Family of egg-laying mammals belonging to this Order contains the **Echidnas** (pronounced *E-kid'nas*) of Australia and New Guinea. These animals are arranged in two genera, the Five-Toed Echidna (*Tachyglossus*),

consisting of a single species which occurs in Australia, Tasmania and New Guinea, and the Three-Toed Echidnas (*Zaglossus*), comprising two species, are confined to New Guinea. The Five-Toed Echidna is covered with strong spines set very thickly all over its outer surface, and its nose is a slender and narrow beak.

**BOOK II**

**BIRDS**



THE BALD EAGLE.

## CHAPTER XV

### AN INTRODUCTION TO THE BIRD WORLD


**Bird Destruction.**—There are many things to be learned about birds besides their names, and their length in millimetres. To-day the first thing to be taught is the fact that from this time henceforth *all birds must be protected, or they will all be exterminated.*

To-day, it is a safe estimate that there is a loaded cartridge for each living bird. Each succeeding year produces a new crop of gun-demons, eager to slay, ambitious to make records as sportsmen or collectors. If a bird is so unfortunate as to possess plumes, or flesh which can be sold for ten cents, the mob of pot-hunters seeks it out, even unto the ends of the earth. Quite recently two "plume-hunters" went at the risk of their lives to Tiburon Island, Gulf of Lower California, to kill egrets for their plumes; and both were killed by the savage Indians there.

In 1897-98 the writer made for the New York Zoological Society a careful inquiry into the volume of bird life in the United States, with special reference to its increase or decrease during the fifteen years prior to that date. From one hundred and eighty competent and conscientious observers, representing thirty-four states and territories, reports were received in answer to a series of questions, all of which were carefully tabulated.<sup>1</sup> Throughout my calculations, wherever a doubt existed, the living birds were given the full benefit of it.

Four states, Kansas, Wyoming, Utah and Washington, show an increase in bird-life. Thirty states show decreases varying from ten per cent. to ninety per cent., but with a general average decrease from 1883 to 1898 of *forty-six per cent.* In the adjoining detailed statement, the shaded portions show the percentages of decrease throughout the states named during the period reported upon:

#### DECREASE IN BIRD LIFE IN 30 STATES, IN 15 YEARS.



<sup>1</sup> "The Destruction of Our Birds and Mammals." By William T. Hornaday. Second Annual Report (1898) of the New York Zoological Society. Until the present edition is exhausted, copies of this paper will be mailed to teachers, on application.

Since the above inquiry was made, the volume of bird-life appears to have changed so slightly that in 1903 conditions are practically as they were in 1898.

**Causes of Decrease in Bird Life.**

The temptation to offer a full statement of the causes and means of prevention of bird-slaughter is very great; but those subjects must be left to other pages. There is, however, much food for thought in the following summary of causes of destruction, as reported by the one hundred and forty-four observers who entered into this branch of the subject. They are listed very nearly in the order of their importance according to the reports:

No.	Reports.
1. Sportsmen, and "so-called sportsmen"...	54
2. Boys who shoot.....	42
3. Market-hunters and "pot-hunters".....	26
4. Plume-hunters, and milliners' hunters....	32
5. "Shooters, generally" .....	21
6. Egg-collecting, chiefly by small boys....	20
7. English sparrow.....	18
8. Clearing off timber, development of towns and cities.....	31
9. Italians, and others, who devour song- birds.....	12
10. Cheap firearms .....	5
11. Drainage of marshes.....	5
12. Non-enforcement of laws.....	5
13. Gun-clubs and hunting contests.....	5
14. Collectors (ornithologists and taxider- mists).....	5
15. Colored population .....	4
16. Indians (for decrease of game quadrupeds)	4

**The Slaughter of Birds for Food.**—The craze for the destruction of bird-life is almost beyond belief. No matter how much the bird-protectors may say about the destruction of our birds, and their impending extermination, far more than the half will remain untold. As our game-birds become fewer and fewer, the market-shooters begin to slaughter birds of song and beauty, which twenty years ago were safe because they were not considered "game." Even ten years ago, no self-respecting American would have lowered himself to the level of the hawk and buzzard by killing and eating the poor little sand-piper and snow-bunting. But mark what is going on to-day:

There is now pending (1903) in the courts the case of the People of the State of New York against two men of New York City, to enforce the payment of fines amounting to \$1,168,315 for

having in their possession contrary to law, in a cold storage warehouse, certain dead birds out of season, game and not game. When the state game wardens searched the premises of the defendants, it is stated that they found the following appalling mass of birds:

8,058 Snow-Buntings!	7,560 Grouse,
7,607 Sand-Pipers!	4,385 Quail,
5,218 Plover!	1,756 Ducks,
7,003 Snipe,	288 Bobolinks,
788 Yellow Legs,	96 Woodcock.

And all this in one cold storage warehouse, for poor, starving New York!

To the public it was a profound surprise to find that snow-buntings and sand-pipers were being slaughtered by thousands for food. At least half a dozen species of song-birds are served on bills of fare under the name of reed-bird. This fact is equivalent to a notice that hereafter no bird is safe from the deadly "market-shooter," and only the strictest watch and the severest measures will save any considerable portion of our birds.

**Protect the Birds.**—Young reader, learn to-day that the birds are the natural protectors of man and his crops from the hordes of insects which without them ravage leaf, flower and fruit. But for the hawks and owls, the wild mice and rats soon would multiply into an intolerable pest. But for the insectivorous birds, destroying grubs and perfect insects by the million, the life of the farmer, fruit-grower and forester would be one long battle against the pests of the insect-world.

Learn that it is wise to encourage birds, as well as to protect them from slaughter. A little food intelligently bestowed is always accepted as a token of friendship and hospitality. Any country dweller can draw birds around him, if he will. Why grudge a few simple shelter-boxes, a few handfuls of grain, and a few pounds of fat pork when in exchange for them you may have, even in winter's dreariness, the woodpeckers, chickadees, crows, and many other winter "residents" and "visitants"? Surely, no right-hearted man or boy can prefer solitude to the company of cheerful and beautiful feathered friends.

**Don't make Bird or Egg "collections."**—Learn to take broad views—bird's-eye views,



if you please—of the bird-world. Consider how you can promote its enjoyment, its betterment, and its perpetuation. Think not that in order to take an interest in birds it is necessary to buy a gun and a bushel of cartridges. Don't think that a badly made bird-skin in a smelly drawer is as pleasing an object in the sight of God or man as the living bird would be. Do not, I beg of you, make a "collection of bird-skins;" for the "bird-skin habit," when given free rein, becomes a scourge to the bird-world.

Do not think that ornithology is the science of dead birds, named in a dead language; or that an attic room is the best field for the study of birds. Study *bird-life*, not merely the mummied remains of dead birds. And, finally, *don't* collect eggs! They teach no useful lesson. The majority of them have no beauty, and are as meaningless as marbles. The pursuit of them is interesting, I grant, but the possession nearly always palls. The collector of eggs destroys life, fearfully, and has for all his labors and his pains only such as this:—**○ ○ ○ ○**.

If you think enough of birds *to mount, or have mounted*, every fine specimen that you kill—aside from legitimate game—then you will be justified in forming a collection. There is some excuse for collections of well-mounted birds, especially those that *are presented to schools*, where thousands of young people may study them; but wild life is now becoming so scarce that the making of large private collections, for the benefit of one man, is a sin against Nature.

**Don't be narrow.**—In studying birds, do not be narrow! Use the field-glass, the camera and pencil, rather than the shot-gun and the microscope. Any fool with a gun can kill a bird; but it takes intelligence and skill to photograph one.

The time was when the analysis and classification of our American birds were important work, because the bird fauna was only partially discovered and written up. In their days, Audubon, Wilson, Baird and Coues did grand work, because so many birds were strange, and needed introducing. The time was when analyzing, naming, and working up geographical distribution were desirable and necessary. But in North America that period has gone by. There is no longer any real need for new technical books on the birds of this continent north of

Mexico. The describing, and re-describing, the naming, re-naming and tre-naming of microscopic varieties, has been done enough, and in places overdone.

**What to do.**—Henceforth, these are the things to be done with and for our American birds:

1. Join actively in protecting the few birds that remain, and help to save them from complete extermination.

2. Aid in teaching the millions how to know and enjoy the beautiful and useful birds without destroying them.

It is not at all necessary that people generally should be able to name correctly every bird that the forest and field may disclose. Many species of warblers, and sparrows, and larger birds also, are so much alike that it is very difficult for any one save a trained ornithologist to analyze them correctly. The general public is not interested in differences that are nearly microscopic. When birds and mammals cannot be recognized without killing them, and removing their skulls, it is quite time for some of us to draw the line.

It is entirely possible for any intelligent person to become well acquainted with at least one hundred and twenty-five of our birds without killing one; and any person who can at sight recognize and claim acquaintance with that number of bird-species may justly claim to be well informed on our birds. Because birds are more common than quadrupeds, bird-books are also more common, and now the most of them are beautifully illustrated. The road to ornithology is now strewn with flowers, and the rough places have been made smooth.

**The Vastness of the Bird-World.**—Go where you will upon this earth—save in the great deserts—some members of the bird-world will either bear you company, or greet you as you advance. Some will sing to cheer you, others will interest and amuse you by the oddities of their forms and ways. On the mountain back-bone of the continent, you will meet the spruce-grouse, the raven, and the mountain-jay. In the foothills and on the great sage-brush plains, the stately sage-grouse and the garrulous magpie still break the monotony.

In the fertile regions of abundant rain, bird-life is—or rather *was* once—bewildering in its variety. In the tropics, the gorgeous colors and harsh voices of the birds remind you that

you are fairly within another world. In mid-ocean, the stormy petrel causes you to wonder how it survives the storms. On the bald mountains of Alaska, or the barren shores of the Arctic Ocean, the snow-white ptarmigan may be the means of saving you from death by starvation; and when you discover new lands in the mysterious and forbidding waters of the Antarctic, the huge and helpless emperor penguin will be there to greet you.

The greatest wonders of bird-life are the immense variety of its forms, and the manner in which the members of the various groups have been equipped to perform so many functions in the economy of life. It seems as if Nature has undertaken to furnish birds for every portion of the globe, and provide food and shelter for each in its own place. This is why different birds fly, wade, swim, dive, scratch, run and climb.

**How we Study Birds.**—To-day, in the primary schools, little children learn something of the wild-birds by which they are surrounded. These studies of Nature are but contributions of bricks and mortar toward what must be the complete building. It is now our purpose to lay the foundation for a structure of bird-knowledge which may be built upon all through life, as elaborately as the builder may choose. But, even those who wish to build only one story in height need just as correct a foundation as those who build the highest.

Our purpose now is to offer the student a general introduction to the bird-world of North America, and illustrate its groups by about one hundred prominent types, all so typical and so representative that every one should know them all. Herein, the student is urged to pay special attention to the systematic groups set forth. Once these are permanently fixed in the mind, the detailed study of the different species of birds becomes a genuine joy.

Learn well the various *Orders* of our birds, the prominent *Families*, and the prominent types representing them. Details regarding anatomy, seasonal changes, migration, breeding-habits, distribution and exact food-habits can be sought later on, and found in great abundance in the wealth of beautiful bird-books now available at small cost. In presenting herein the individual birds which have been chosen to represent the

different groups, we shall strive to give in a few words an accurate and clearly defined general impression of each, but no more.

**Remembering the Orders of Birds.**—The birds of North America are divided into seventeen Orders, besides which two additional Orders exist elsewhere. Under different circumstances, the student might find some difficulty in remembering these Orders, and the relations they bear toward each other. In this, however, we find ourselves aided by Nature in a remarkable way.

By a very simple and natural arrangement, with fair regard to the forms and habits of birds, and their haunts upon the earth, it is possible to show upon a chart, the following facts:

1. The various Orders of North American birds;
2. The relative size of each Order, in number of species;
3. The haunts of each Order, on land or water, and
4. Approximately, the rank of each order, from lowest to highest.

On the accompanying chart of bird-life, an ideal panorama of land and water is divided between the various Orders of North American birds, just as we find them in Nature. By a fortunate coincidence, the Orders that are lowest in the scale of natural classification are those containing the sea-birds, of deep water, which therefore belong at the bottom of the chart. On the other hand, the birds that are highest in the zoological scale—the perching birds—are also the birds of the tree-tops, and must be placed at the top of the chart.

The birds of the shore, the river-bank, and the uplands have their respective areas in the middle portion of the scale, and we are thus enabled to see almost at a glance the geography of the bird-world, at least as we find it in North America.

Beginning with the highest, we shall endeavor to point out the leading characters of the various Orders, and the examples which best represent them. Just at present, however, it is not wise for the student to go too far into the subdivisions of the Orders, and only the most important Families will be mentioned by name.

Any student who is unwilling to devote a few hours to learning the names and places of the

various Orders of birds may as well refrain from attempting to know our feathered friends; for that knowledge is quite as necessary as foundation-stones are to a tall building. *The names of the Orders must be learned, and remembered!*

For the purpose of making the contents of each Order familiar to the reader, representatives of the most important Families it contains will be mentioned, and illustrated by the presentation of at least one species.

### THE ORDERS OF LIVING BIRDS.

ORDER.	PRONUNCIATION.	CHARACTER OF BIRDS INCLUDED.	EXAMPLES.
PASSERES	<i>Pas'se-rez</i>	Perching-Birds	Robin, Warbler and Jay.
MACROCHIRES	<i>Mac-ro-chi'rez</i>	{ Goatsuckers, Swifts and Humming-Birds	{ Nighthawk, Swift, Ruby-Throat.
PSITTACI	<i>Sit'ta-si</i>	Parrots and Macaws	Carolina Parakeet, Macaw.
PICI	<i>Pi'si</i>	Woodpeckers	Golden-Winged Woodpecker.
COCCYGES	<i>Coe'si-gez</i>	{ Cuckoos and Kingfishers	{ Belted Kingfisher, Cuckoo.
RAPTORES	<i>Rap-to'rez</i>	Birds of Prey	Eagle, Owl and Vulture.
COLUMBAE	<i>Co-lum'be</i>	Pigeons and Doves	Band-Tailed Pigeon, Mourning-Dove.
GALLINAE	<i>Gal-li'ne</i>	Scratching-Birds	Quail, Grouse, Wild-Turkey.
LIMICOLAE	<i>Li-mic'o-le</i>	Shore-Birds	Plover, Woodcock, Snipe.
PALUDICOLAE	<i>Pal-u-dik'o-le</i>	Cranes and Rails	Whooping Crane, Virginia Rail.
HERODIONES	<i>Her-o-de-o'nez</i>	Herons and Egrets	Great Blue Heron, Snowy Egret.
ODONTOGLOSSAE	<i>O-don-to-glos'se</i>	Flamingoes	American Flamingo.
ANSERES	<i>An'se-rez</i>	{ Swimmers with Comb-Edge Bills	{ Mallard, Canada Goose, Swan.
STEGANOPODES	<i>Steg-a-nop'o-dez</i>	{ Fully-Webbed Swimmers	{ Pelican, Darter, Cormorant.
TUBINARES	<i>Tu-bi-na'rez</i>	{ Tube-Nosed Swimmers	{ Albatross.
LONGIPENNES	<i>Lon-gi-pen'nez</i>	{ Long-Winged Swimmers	{ Gull and Tern.
PYGOPODES	<i>Py-gop'o-dez</i>	Diving-Birds	Loon, Grebe, Auk, Murre.
IMPENNES	<i>Im-pen'ez</i>	Flightless Divers	Penguin.
RATITAE	<i>Ra-ti'te</i>	Flightless Runners	Ostrich, Cassowary.

## EXPLANATION OF THE CHART OF THE ORDERS OF NORTH AMERICAN BIRDS.

The Orders of North American birds lend themselves with gratifying readiness to the purposes of a landscape chart. In this way more than any other known to the author can the greatest number of facts regarding the Orders and their relationships be set forth in a manner easily understood, and calculated to appeal to the eye.

As with the mammals, the highest Orders are found in the tree-tops and the air; and as nearly as possible the relative sizes of the various Orders are shown. The birds of the highest and most perfect organization appear at the top of the chart, and the lowest forms are those of deep water, farthest from the land.

The great size of the Order Passeres is strikingly apparent; and it is situated in the tree-tops where its members live.

The curious shape of the Order Macrochires is due to the fact that the Goatsuckers, Swifts and Humming-Birds have so little in common that they are wellnigh separated; but the larger body—the Hummers—are closely related to the Perching-Birds.

The Order Coccyges is composed of two groups equally ill matched, the Cuckoos and Kingfishers. The former touch the Perching-Birds, the latter the sharp-beaked fishers; but the association of the two in one Order is not satisfactory, and not likely to stand.

The Orders Columbæ, Gallinæ and Paludicolæ are found on the uplands, immediately above the Limicolæ, or Shore-Birds.


The Herodiones (Herons, Egrets and Bitterns) range along the shore from the sea, up the river, to the interior lake, while the Anseres—Ducks and Geese—cover lake, river and sea.

The Flamingo's Order—Odontoglossæ—is of the shallow water of an estuary, connecting the Herons and Ducks.

The Steganopodes (Cormorants, Pelicans, etc.) prefer the shallow waters of the sea, while the Gulls and Terns (Longipennes) range from shallow to deep waters.

The Tubinares (Albatrosses, Fulmars, etc.) and Pygopodes (Auks, Murres, and other weak-winged divers) are birds of deep water.

For obvious reasons, it has not been considered a practicable matter to include on a landscape chart the birds of the world, or even those of South America.



Copyright, 1903, by W. T. HORNADAY.

LANDSCAPE CHART OF THE ORDERS OF NORTH AMERICAN BIRDS.



S. SAWYER

THE BOBOLINK IN SPRING.

## CHAPTER XVI

# THE ORDER OF PERCHERS AND SINGERS

### PASSERES


This Order is the highest in the scale of birds, and it is by far the largest of the whole nineteen Orders. In the zone of agriculture it contains the birds which are of the greatest importance to mankind, the insect-eaters. It also contains all the real song-birds of the world, and its North American Families are as shown below:

	FAMILIES.	SCIENTIFIC NAMES.	EXAMPLES.
ORDER PASSERES.	THRUSHES, . . .	<i>TUR'DI-DAE.</i> . . .	Robin, Thrush, Bluebird.
	KINGLETS, . . .	<i>SYL-VI-DAE.</i> . . .	Kinglet and Gnatcatcher.
	NUTHATCHES, . .	<i>PA'RI-DAE.</i> . . .	Nuthatch, Chickadee, Titmouse.
	TREE-CREEPERS,	<i>CER-THY-DAE.</i> . .	Brown Creeper.
	DIPPERS, . . .	<i>CIN'CLI-DAE.</i> . . .	Water-Ouzel.
	WRENS, . . .	<i>TRO-GLO-DY'TI-DAE.</i>	Wren, Cat-Bird, Mocking-Bird, Thrasher.
	WAGTAILS, . . .	<i>MOT-A-CIL'LI-DAE.</i> .	Wagtail and Pipit.
	WARBLERS, . . .	<i>MNI-O-TIL'TI-DAE.</i> .	Warbler, Water-Thrush, Redstart, Chat.
	VIREOS, . . .	<i>VI-RE-ON'I-DAE.</i> . .	Red-Eyed Vireo.
	SHRIKES, . . .	<i>LAN'I-DAE.</i> . . .	Butcher-Bird and Loggerhead Shrike.
	WAXWINGS, . . .	<i>AM-PEL'I-DAE.</i> . . .	Bohemian Waxwing, Cedar-Bird.
	SWALLOWS, . . .	<i>H'RUN-DIN'I-DAE.</i> . .	Swallow and Martin.
	TANAGERS, . . .	<i>TAN-A-GR'I-DAE.</i> . .	Scarlet Tanager.
	FINCHES, . . .	<i>FRIN-GIL'LI-DAE.</i> . }	Sparrow, Finch, Grosbeak, Cardinal, Snow-Bunting, Redpoll.
	BLACKBIRDS, . .	<i>IC-TER'I-DAE.</i> . . }	Blackbird, Oriole, Meadow-Lark, Bobo- link.
CROWS, . . .	<i>CORVI-DAE.</i> . . .	Crow, Raven, Jay, Nut-Cracker.	
HORNED LARKS,	<i>A-LAU'DI-DAE.</i> . .	Horned Lark.	
FLYCATCHERS, .	<i>TY-RAN'NI-DAE.</i> . .	Flycatcher, Pewee, Phœbe, Kingbird.	

The majority of perchers are birds of plain feather, quite as if Nature had intended that these, the best friends of the farmer and fruit-grower, should be the last to be destroyed by the merciless Man-With-a-Gun.

It will be a sad day for the American farmer when the last insect-eating bird of our country is brought fluttering and lifeless to the ground. When the armies of destroying insects begin to multiply unchecked, and send forth their millions and tens of millions, then will the husbandman realize the value of the allies he has lost, and vainly wish to exchange any number of grapes and cherries for the once-despised robin, thrush and blackbird.

Quite apart from their cash value to the agriculturist, it is the song-birds that appeal most strongly to the ear and heart of man. Even the exquisite plumage of the resplendent trogon, most beautiful of all American birds, does not thrill the soul as does the song of the robin, the brown thrasher and the mocking-bird. Next to sunshine and green verdure, the most cheering thing in Nature is the song of a bird. At this moment (early spring) a robin, in the big maple in front of my windows, is pouring forth a song that is at once restful and inspiring. It reminds me that we who live in the temperate zone are *greatly favored* by the presence in our bird-life of the sweetest singers in the world. Shall



THE PARTS AND PLUMAGE OF A PERCHING-BIRD.

we, then, be so utterly barbarous and mean as to engage in, or permit, the killing of our song-birds in order that they be used either as food for biped pigs, or to adorn(?) the cheap millinery of servant-girls? Never!

Let it not be thought, however, that the Order Passeres has not a good share of birds of beautiful plumage. In our own fields and forests, behold the waxwing, the oriole, the cardinal, the tanager, the grosbeak, the magpie, the jay and the bobolink. The tropics contain the wonderful birds of paradise, and a bewildering array of humming-birds, cotingas, finches, ground-thrushes and many others.

If the temperate zone lacks anything in perching-birds of brilliant plumage, that lack is more than made up by the singing-birds. With all its wealth of bird-life, brilliant and plain, the tropics are generally silent, and a joyous or musical bird-song is rarely heard. Of the bird-cries that one occasionally hears, the majority are harsh and unpleasant squawks. The tropical day has neither robin nor mocking-bird, the night no whippoorwill. True, there is the awful "brain-

fever" bird of the Indian night, but it is neither musical nor joyous. One may spend months in the tropics, both of America and of the Far East, and in all that time hear less of real bird-song than can be heard on many an American farm in one day.

As might be expected in a large Order of birds, the food habits of the perchers cover a wide variety of foods. The great majority prefer to live upon insects, and the young of all species are absolutely dependent upon soft-bodied insects, larvae and earth-worms. Many birds are really limited to insect-food, and can subsist on no other kind. Next in importance, and for the longest period, perhaps, come seeds and grain, especially the seeds of weeds that are a pest to the farmer. As a rule, fruit is taken in its brief season more as a dessert than as a staff of life.

A very few species, like the crow, magpie and jay, eat meat whenever opportunity offers it, and welcome the discovery of raw meat or eggs.

The great value of the perching-birds lies in the enormous quantities of insects which they con-



sume as food.<sup>1</sup> These birds have been specially developed by Nature to combat and destroy the hordes of insects destructive to fruit, grain and tree life, which otherwise would in a short time increase to such enormous numbers that no vegetation could withstand their attacks.

To young pupils, the Order of Perching and Singing Birds may at first seem difficult to grasp; but in reality it is not. A knowledge of forty birds will give one a very good idea of its various Families; and any one can learn about forty birds. After this Order has been mastered, all others will be found quite easy. The examples introduced have been selected with great care, and concerning those illustrated, the pictures will tell of their forms and markings far better than wordy descriptions could do.

### THE THRUSH FAMILY.

#### *Turdidae.*

**The Robin.**<sup>2</sup>—All lovers of birds should agree in placing this dear old friend at the head of the list of the birds of this continent. This is because it is the highest avian type. It has typical plumage, it flies well, it perches, it sings beautifully, it migrates, and its anatomy is thoroughly representative. Moreover, it quickly discerns a friend and protector, and it is not driven away by the English sparrow.

Of all our birds, the Robin comes the nearest to being "folks." It is always one of the first birds to arrive in the spring, it remains all summer, and it is one of the last to depart at the approach of winter. Often the late spring snows catch it on its early migration, and its staying powers are put to the test. It is a good plan to scatter food for these early birds. Nothing save the sun itself is more gladdening on a raw March day than the joyous note by which the Robin announces the arrival of himself and spring.

Who is there who can know the Robin and not love him? Few indeed; and those persons

<sup>1</sup> Up to 1903 the Biological Survey of the United States Department of Agriculture had published twenty important bulletins and shorter papers on the food habits of our birds, with especial reference to the species either most beneficial or most harmful to the farmer and fruit-grower. A list of those now available, and the terms on which they are procurable, will be furnished by the Department upon application.

<sup>2</sup> *Me-ru'la mi-gra-to'ri-a.* Length, from end of beak to end of tail, 9 to 10 inches.

around New York and in some parts of the South who shoot Robins for food, are wholly unfit to inhabit the Robin's country, unless they reform.<sup>3</sup>

The Robin is one of the sweetest and most joyous songsters I know. As well try to describe the glories of a sunset as to set forth in words the liquid melody, clear and sweet, which pours from his throat when he feels particularly joyous.

Everywhere, the Robin is a very sociable bird, and exceeding quick to distinguish a friend from a foe. Give it absolute protection, and security from cats, and it will cheerfully nest on your window-sill. This is what one actually did in Buffalo, under our roof,—built her nest on the sill of an upper window, close against the glass, and reared her brood there. We went many times to see



ROBIN.

how she was getting on, and she, knowing well that glass is a barrier, permitted us to put our faces within two inches of her head.

In the Zoological Park, the Robins were the first wild creatures to learn, in 1900, that the reign of the poacher was over; and they quickly told it to the crows, and thrushes, and other birds. In an eight-foot pine-tree, that was

<sup>3</sup> "In central Tennessee are large tracts of cedars, the berries of which serve to attract myriads of Robins in the winter. One small hamlet in this district sends to market annually enough Robins to return \$500, at five cents per dozen, equal to 120,000 birds." They are killed at night by torchlight, with sticks. An officer of the Louisiana Audubon Society states that a conservative estimate of the number annually killed in Louisiana for food purposes is a quarter of a million when they are usually plentiful.—William Dutcher, in *Educational Leaflet No. 4, of the National Committee of Audubon Societies.*

planted six feet from the edge of the main walk, and directly in front of our head-quarters, a Robin built her nest, only five feet from the ground; and there she reared her young. To many visitors who loved birds, her nest was shown, but to the Robin-killers and the nest-robbers no one said a word. On Gardiner's Island, where cats live not, the Robins nest on fence-rails only two feet from the ground, in full view of the bird-loving inhabitants of that small world.



WOOD-THRUSH.

Often we have been greatly interested by the keenness of sight of the Robins which visit our lawn. After every shower, certain Robins of our acquaintance take possession of the lawn, and stride over the grass with an air of great importance and earnestness of purpose. After several wise and sidewise cocks of the head, a Robin will suddenly drive his bill far down into the grass, and brace himself for a hard struggle. By dint of many hard tugs, out comes the earthworm, to be borne away in triumph to a certain nest. Often I have tried to see worms down among the roots of the grass, as the Robins do, but never once have I succeeded. Evidently my objectives never were focused just right for worms in green grass.

In all save a very few localities in North America, the Robins are treated as friends. In the "grape belt" of western New York, they are a great annoyance to some grape-growers because of the bunches they disfigure. Elsewhere they are of great benefit to farmers, and the few cherries they take in cherry time are very modest

compensation for the noxious grubs they pick out of the freshly ploughed fields.

The investigations of the Biological Survey of the Department of Agriculture have demonstrated the great economic value of the Robin as a destroyer of harmful insects. The contents of three hundred and thirty stomachs of birds taken in all seasons revealed the fact that in the course of an entire year, insects make up 40 per cent of the food of Robins, wild fruit 43 per cent, cultivated fruit 8 per cent, and miscellaneous vegetable food 5 per cent.

Regarding the killing of Robins, and other song-birds, and also doves, as food for man in a land of plenty, there cannot be two opinions. It is not necessary; it is not "sport"; it is very injurious to our farmers and fruit-growers, and entirely reprehensible. No self-respecting boy or man can be guilty of such wrong-doing; no civilized community should tolerate it for one moment, and *no farmer can afford to permit it!* I would rather that any friend of mine should be caught stealing a sheep than killing Robins, either for food or "sport."

Let us protect the great American Robin, and all other perching-birds, even at the point of the bayonet if it be necessary.

**The Wood-Thrush**<sup>1</sup> is one-fifth smaller than a robin, and is easily recognized anywhere by its beautifully spotted breast. It has about fifty dark-brown spots, often arranged in rows up and down its breast, belly and throat, on a creamy-white ground color. Other thrushes have dark spots on the breast, but not down to the legs. The head and shoulders of this bird are of a bright cinnamon color.

This graceful creature often works overtime to make the woods melodious, and it is one of our sweetest singers. It is not so bold and confident as the robin, and is much given to following the robin's lead. Its favorite haunt is the sweet seclusion of shady woods and thickets, where the half-bare earth affords good hunting-grounds, and a fair degree of safety from observation. Its nesting habits are very much like those of the robin, and its range includes the whole eastern half of the United States, to the Great Plains beyond the Mississippi.

**The Common Bluebird.**<sup>2</sup>—The United

<sup>1</sup> *Hy-lo-ci'chla mus-tel-i'na*. Length, 8 inches.

<sup>2</sup> *Si-a'li-a si-a'lis*. Length, 6.75 inches.

States is a country of such vast extent it is a physical mosaic of different elevations, soils, and climates. Roughly speaking, these are its physical divisions:

1. The *eastern half*, of ideal rainfall, bountiful harvests, and abundant shade.

2. The *Great Plains*, fine for grazing, but mostly too dry for agriculture.

3. The *Rocky Mountain region*, embracing a perfect medley of physical conditions, mostly high, rugged, and rather lacking in insect-life.

4. The *arid regions*, of the country between the Rockies and Sierra Nevadas, extending from southern Washington to the City of Mexico, and including southern California.

5. The *region of great rainfall*, on the north-western Pacific coast (northern California, Oregon and Washington).

It is not strange, therefore, that we find typical species of eastern animals developing westward into different colors, and also different pelage, and designated scientifically by different names. Take these examples by way of illustration:

In the East we have the *Common Bluebird*.

In the Rockies we have the *Chestnut-Backed Bluebird*, and also the *Mountain Bluebird*.

In Arizona we have the *Azure Bluebird*.

In the Pacific states we have the *Western Bluebird*,

And in Lower California, the *San Pedro Bluebird*.

Is it at all necessary that the general student should know about all these different species in order to not be accounted ignorant? Let us see.

Any sensible civilized person knows a cow at sight, also something of its place in Nature, and its habits. No one, however, save the special student of domestic cattle, is expected to be able to say, without "looking it up," whether a particular cow is an Alderney, a Jersey, a Short-Horn, a Hereford, or a Durham.

The case of the Bluebird is quite similar. He who knows one Bluebird well, may justly claim a bowing acquaintance with all the others, and feel at home when in their company.

Here in the East, the Bluebird is a thing of beauty, and a joy until the abominable English sparrows drive it away. It comes with the robin, to help chase winter away; and though we have heard it a hundred times, it is always welcome

news, late in February or early in March, to hear some one say triumphantly, "I saw a Bluebird to-day!" It is as needless to describe this feathered beauty, with the brown breast, and back of heaven's bluest sky-tint, as it would be to describe a rainbow.

Unfortunately, the Bluebirds are not good fighters, and the English sparrows harry them shamefully. They are timid, and easily driven away. Worse than this, they are easily killed by cold weather. The cold wave which visited the South in 1895 killed so many thousands of Bluebirds, especially in North Carolina and Arkansas, that for some time afterward the number visible in the North was alarmingly small. If not molested by the English sparrow, the Blue-



BLUEBIRD.

bird takes readily to boxes erected on poles near farm-houses, similar to those frequently erected by the farmer boys to attract the purple marten. A good way to encourage robins and Bluebirds is to kill the English sparrows.

## THE KINGLET FAMILY.

### *Sylviidae.*

**The Ruby-Crowned Kinglet**<sup>1</sup> is one of our smallest birds, and it is easily recognized by the tiny tuft of ruby-red feathers on the crown of its

<sup>1</sup> *Reg'ulus cal-en-du'la*. Length, 4.25 inches.



RUBY-CROWNED KINGLET.

head. In life it is a dainty little feathered gem, but it is so modest and retiring that it is seen only by sharp eyes. "Kinglet" means "Little King." Mrs. Mabel Osgood Wright testifies strongly to its value as an insect-destroyer, especially in the late autumn, when other insectivorous birds have gone, when it works industriously upon the trunks of evergreens. Dr. Coues considered the Kinglet an exquisite singer, but I must confess that its vocal powers have quite escaped me.

#### THE NUTHATCH AND TITMOUSE FAMILY.

##### *Paridae.*

The birds of this Family deserve to rank as prime favorites. They remain with us through "the long and dreary winter," when all save a corporal's squad of the grand army of birds have fled southward, and left us to our fate. They are exceedingly industrious, and their efforts are directed against insects of very destructive habits, the tree-borer and the bark-louse. In their work they are not continually "playing to

the gallery," and telling people how busy they are.

**The Chickadee, or Black-Capped Titmouse,**<sup>1</sup> is one of the dearest little fellows that flies. It always reminds me of a forest-elf, in a black cap and a feather cloak. Instead of making a great show of fright, like a girl in the presence of a ferocious mouse, little Black-Cap perches on a tiny twig growing low down on the trunk of a big tree, and cocks his head at you, while he looks you over with a fraternizing air. His attitude and manner say as plainly as English, "You are a good fellow, and I'm another. We understand each other perfectly, don't we?"

And then his greeting. If you have never before had the pleasure of meeting him, he pipes out cheerily, "CHICK-a-dee-dee-dé!" Watch for him the next time you go into the woods in winter,—a jet-black cap with a white waistcoat below it; a black necktie, bluish-gray overcoat, and a very pert and saucy air. You can hardly fail to recognize him, but in case you hesitate, and think his "face is familiar," he will up and tell you his name, as plainly as print.



CHICKADEE.

<sup>1</sup> *Parus at-ri-cap'il-lus.* Length, 5.25 inches.

Six well-marked types and several races of Chickadees inhabit North America from Alaska to Mexico, but the one most widely known is that just named.

**The White-Breasted Nuthatch**<sup>1</sup> deserves the most perfect protection and encouragement that the people of this country know how to offer. One good look at this bird on the trunk of a valuable tree, searching as if with a magnifying glass for the trees' deadly enemies,—the borers,—ought to convert any person to the cause of bird-protection. Like the chickadee, the Nuthatch remains in the north all winter, because he feels that he has not a moment to lose in his war on the borers.

The tree-trunks are his favorite hunting-ground, and he goes over them, literally inch by inch. He becomes so absorbed in his work that he forgets all about himself, and works half the time head downward, or oblique, or horizontal, as it may happen to be. Rarely does he stop to talk, and even then he only clucks in his throat, "not necessarily for publication, but as a guarantee of good faith."

Often in the silent and snowy woods, when your feet go *rip! rip! rip!* through the frozen crust, you hear close overhead a scratching, digging sound, as of some one gouging into rough bark with a pocket-knife. Look up, and it will be a Nuthatch, working away as if his job depended upon the doing of a daily stint. He thinks that in his case it is the late bird that catches the worm! His beak is like that of a small woodpecker, and although his friend the chickadee has more style than he, he himself is much better fitted for digging in bark. The top of his head is black, his sides, throat and breast are pure white, while his back is dull blue, or gray-blue. As a climber, this bird surpasses the woodpecker, because in clinging to a tree-trunk it makes no use of its tail.

Nuthatches are easily encouraged to make your trees their head-quarters. In December, nail to a tree-trunk here and there, about twelve feet from the ground, some lumps of suet, or fat pork on the rind, or beef bones with a little raw meat upon them, and see how quickly the birds find them out. The "winter residents" will feast upon them until the last morsel has disappeared, and they will appreciate your thought-

<sup>1</sup> *Sitta carolinensis*. Length, about 6 inches.

fulness thus displayed precisely when tree-borers burrow deepest, and are most difficult to get at.

### THE TREE-CREEPER FAMILY.

#### *Certhiidae.*

**The Brown Creeper**<sup>2</sup> represents a small Family of small birds of tree-climbing habits, but with bills that are rather too slender for work in bark. They are not fitted by nature for digging a modest and retiring borer from the bottom of his tunnel, and therefore they make a specialty of bark-lice and other surface wood-workers which can be picked off without hard digging.

As an example of protective coloration, this



WHITE-BREASTED NUTHATCH.

little creature is worthy of special note. Its back is brown, marked by about twelve broad stripes of dull gray, and between the two colors the striations of bark are surprisingly well imitated. On the side of an oak, or elm, or chestnut, this little bird is almost invisible until it moves. It does not work head downward, like the nuthatch, but creeps about with its head up, braced by its tail, like a woodpecker. Like

<sup>2</sup> *Certhia familiaris americanus*. Length, 5½ inches.

both the preceding species, it is a winter resident, and in fact is not much in evidence at any other season. The four species of this group cover the United States, and extend from Alaska to Guatemala.

### THE WRENS AND CAT-BIRDS.

#### *Troglodytidae.*

In some respects, the wrens are but a short step from the tree-creepers, but in others they are widely apart. For its size the **House-Wren**<sup>1</sup>



BROWN CREEPER.

is the most pert and saucy bird in North America. Forty years ago, a pair of these merry little sprites took up their abode in the wild fastnesses of the grape-arbor that sheltered our well; and I can hear their shrill chatter yet. It was like the piping of a piccolo. For eight years, they and their children and grandchildren possessed the outskirts of our dwelling, and it was a great day when we discovered a beautiful, feather-lined nest, nearly six inches deep, that the Wrens had built in an old-fashioned lantern that hung in the wood-house. I wish it were possible to have

<sup>1</sup> *Tro-glo-dy'tes ae'don*. Length, 4.75 to 5.25 inches.

Wrens around a city dwelling, or in a Zoological Park.

A Wren is known by the way it carries its tail, so very straight up in the air that sometimes it tilts forward. The House-Wren is the most sociable of all our wild birds, and also the one most confident of its place in the hearts of its countrymen. I never knew of a Wren being killed by any one save a collector of bird-skins. As for myself, I would go Wrenless forever rather than take the life of a creature so winsome and trustful. Even the cats of our household used to respect the family Wrens. In the country, where there are no English sparrows, it is easy to attract these interesting birds by putting up nesting-boxes for them. Five species of Wrens occupy the United States, from ocean to ocean, the Pacific species, west of the Rockies, being the *Tule Wren*.

**The Brown Thrasher.**<sup>2</sup> — Vocally, this bird is practically the northern understudy of the mocking-bird. When, after a warm spring shower and a sudden burst of sunshine, an able-bodied Brown Thrasher perches on the tip-top of a red-haw bush, and for fifteen minutes pours forth a steady stream of delicious melody, in bewildering variations, one is tempted to declare that no mocking-bird can surpass it. It is simply indescribable. Often when sadly toiling in the Iowa fields, I have been stopped and held by this feathered spellbinder for what seemed to my brothers like very long intervals.

In form this bird is very much like the mocking-bird, but its back is colored a rich iron-rust brown, and its under surface is dull white, strongly spotted with large, triangular brown spots. Its home is the whole of the United States east of the Rocky Mountains, and it is the sweetest singer of the North. Unfortunately, its song-period is rather short, and terminates about the end of June.

**The Cat-Bird**<sup>3</sup> of the North bears a strong resemblance to the mocking-bird, in form, color and movement. It is also a good singer, though hardly in the same class as its southern relative. It is very sociable in its habits, and loves the orchards, gardens, fruit-trees, and berry-bushes of the country dweller. Its name is derived from its favorite exclamatory cry, which sounds

<sup>2</sup> *Har-po-rhyn'chus ru'fus*. Length, 11.25 inches.

<sup>3</sup> *Gal-e-os-cop'tes carolinensis*. Length, 8.75 inches.



HOUSE-WREN.

like the plaintive mew of a half-grown kitten. Its prevailing color is dark, slaty gray.

**The Mocking-Bird,**<sup>1</sup> of the states south of the Ohio River, is a singing wonder. It is a little bundle of nerves, covered with modest drab feathers, and its throat is tuned up to concert pitch. When it is silent, it can be recognized by its slender body, long legs and long tail; but when it is singing, only a deaf man needs an introduction. This bird can also be recognized by its nervous and irregular movements, hopping and darting about, up, down and sidewise. If the Mocker feels well, he sings as he darts about, as jerkily and impulsively as he moves.

The Mocking-Bird loves to sing almost as well as some persons love to hear him. His typical song is a bewildering medley of warbling, chirping and twittering, many passages being very clever imitations of other birds, but the majority of it is improvised for the occasion. Next to

<sup>1</sup> *Mimus pol-y-glot'tos*. Length, about 10 inches.

the marvellous variety of his vocal exercises is the clearness and sweetness of his notes; for this singer never sharps nor flats. The amount and variety of the melody that comes from that insignificant little gray midget in feathers are truly marvellous. Every person who has heard the free, wild bird performing in its home thicket knows that the singing of caged specimens is but a spiritless imitation of the wild song.

Strange to say, this bird not only sings in the daytime, but there are periods, especially during the breeding season, when the male sings at night.

As usual, man's destructiveness reaches out for this the greatest of all American singers. Thousands of nestlings are caged, the majority of them in Louisiana. Those that do not die in the process of rearing, live for brief periods in wretched little 12 by 14-inch cages, and die without having known one happy, joyous hour. It is reported that in most portions of the South, the Mocking-Birds are rapidly decreasing in number, especially in Arkansas. The killing of a bird of this species, on any pretext, should be made a penal offence.

#### THE DIPPER FAMILY.

##### *Cinclidae.*

**The Water-Ouzel, or Dipper,**<sup>2</sup> is one of the most remarkable little birds on this continent.



CAT-BIRD.

It is a genuine water-elf, and the things it can do are almost beyond belief. I first saw it in late

<sup>2</sup> *Cinclus mex-i-can'us*. Length, about 8 inches.

November, on the strip of ice which fringed the edge of the roaring, swirling, icy-cold water which plunges into the Shoshone Canyon at the forks of the Shoshone River. Man or beast stepping into that foaming torrent would have been crushed against the rocks, and drowned at the



MOCKING-BIRD.

same moment,—two deaths in one. In that grim and terrible solitude, fast in the embrace of early winter, we saw on the snow-white brink of the ice-bank a tiny dark object, which closer inspection revealed to be a bird. It looked like a large gray wren.

As we paused to regard it, it blithely flew down into mid-stream, and dived head foremost into a chilly wave that ran ten miles an hour. An instant later it reappeared, all unruffled and unwet, blithely flew back to the edge of the ice, and alighted once more. Then we knew well what it was; for it could be nothing else than the Water-Ouzel. Afterward, we saw others along the line of the Denver & Rio Grande Railway where it winds its way through the Rocky Mountains. Where the walls of the Royal Gorge almost crowd the train into the Arkansas River is a good place to watch for them.

This bird is a *diving thrush*! Nature has

fitted it to dive boldly into the coldest and most turbulent water, or through a water-fall, and even to walk on the bottom of a still pool, without being at all disturbed. Both in form and size this little creature is like a large wren, but it is so peculiar it occupies a genus quite alone. Of course it is not web-footed; and in appearance it exhibits not one feature suggestive of a semi-aquatic life. Its home is along the foaming torrents of the Rocky Mountains, and Sierra Nevadas, from Alaska to Guatemala. It nests close beside swift-running streams, sometimes beside or even behind a cascade. It is known that this strange bird gives forth a song both clear and sweet, but I have never seen one elsewhere than near a roaring torrent, where no ordinary bird-song could be heard.

### THE WARBLER FAMILY.

#### *Mniotiltidae.*

From the middle of April to the middle of September, the woods and thickets of the northern states are inhabited by a very considerable number of tiny bird-forms. They are trim-built little creatures, quiet and business-like, and they take themselves very seriously. A few of them are clad in refined shades of yellow, but—most fortunately—the great majority wear dull olive, gray or brown colors, and thereby escape the hostile attention that bright plumage always attracts.

These are the warblers, grand in the destruction of insects, but the most elusive and difficult little creatures with which bird-students have to deal.

The difficulty lies in studying them effectively without killing them. As for myself, I have not yet seen the day wherein I could find myself willing to slaughter from five hundred to a thousand of these exquisite little creatures for the sake of becoming sufficiently acquainted with them to name them *when they are dead*! I blush not in admitting that I have gone half way through life knowing less than a score of warblers to the point of naming them, accurately, as they fly before me. My exhortation to all young people is—*do not slaughter birds, of any kind, merely to become acquainted with their names.* Some of the wild flowers can endure that method without extermination, but the wild birds and mammals cannot.



It is *not at all essential* that such tiny, inconspicuous creatures as warblers should be recognized and correctly named at sight. Already a million warblers have died to make holidays for collectors. Not long since I received from an egg-dealer a circular advertising the following eggs for sale:

Worm-Eating Warbler.....	84 sets,	416 eggs.
Yellow Warbler.....	94 "	388 "
Oven-Bird.....	105 "	458 "
Yellow-Breasted Chat.....	139 "	521 "
Kentucky Warbler.....	210 "	917 "

Total for 51 species. . 1,274 sets, 5,433 eggs.

It is such wanton destruction as this which makes me "down" on egg-collecting. It is safe to say that the taking of those 5,433 warbler eggs, robbed the farms and forests of New York state of that number of useful birds, not counting possible progeny, and did not one dollar's worth of good to the "cause of science," or any other public interest. Already, poor "Science" has an awful load of crimes against Nature to answer for. Do not add to it without very strong justification.

The members of the Warbler Family, commonly called wood warblers, are distributed all over North America, wherever insects abound, from the southern edge of the arctic Barren Grounds to southern Mexico. In her very scholarly and useful book entitled "Birds of the Western United States," Mrs. Florence Merriam Bailey enumerates forty species; and Mr. Frank M. Chapman, in his "Birds of Eastern North America," gives fifty-two. Of these, however, twenty-one are duplicated, and therefore the whole number of warblers described in the two handbooks is seventy-one. When we consider the fact that about sixty of those species are very small birds, of uniform size, and many of them quite unmarked by striking special colors, the difficulty of becoming acquainted with the different species will begin to appear. For present purposes, the whole Family can be very fairly represented by three species. Two of them are of universal distribution, and the third (the chat) is nearly so.

**The Yellow Warbler, or Summer Yellow-bird,**<sup>1</sup> is chosen as the type of about sixty species

<sup>1</sup> *Den-dro'i-ca aesti-va*. Length, 5 inches.

of small wood warblers each of which is called "Warbler" with a descriptive name prefixed, such as palm warbler, prairie warbler, Calaveras warbler, etc. It is of a bright, greenish-yellow color, and is easily recognized on the wing. On the Western prairie farms, the boys call it a "Wild Canary," because it strongly resembles the orange yellow phases of that popular cage-bird. As if courting acquaintance with man, it loves to frequent the roadside thickets, the edges of woods, and even the orchard and garden.

The beauty of this bird far surpasses its minstrelsy, for it is but an indifferent singer. The fact is, however, that it has so much work to do in catching insects it has little time for music; for it will be noticed throughout the bird-world that the most diligent insect-catchers are not in the habit of singing over their work. This is due to the same reason that a good deer-hunter does not talk and tell stories while following a trail.

The Yellow Warbler ranges from the Atlantic



YELLOW WARBLER.

to the Pacific, and over practically the whole of North America save the arctic barrens, Alaska, and our arid southwestern states. Mrs. Mabel Osgood Wright says "it is one of the particular victims which the cow blackbird selects to foster its random eggs, but the Warbler puts its intelligence effectively to work, and some-

times builds a floor over the unwelcome egg." (*Birdcraft*, p. 95.)

**The Yellow-Breasted Chat**<sup>1</sup> is much larger than the typical wood warblers, being  $7\frac{1}{4}$  inches long to their 5 or  $5\frac{1}{2}$  inches. It has an olive-green back and a sulphur-yellow breast and throat, with a white line extending from its beak above and around its eye. By these colors, and its



RED-EYED VIREO.

erect tail, it may easily be recognized. It is a very pert and saucy bird, and much given to frequenting the haunts of country dwellers.

The Chat is not a great singer. He has no regular song, and the notes he utters are jerky, erratic and elusive. Its voice has some peculiar quality which renders this bird very difficult to locate by sound alone. Many times I have been completely misled by its call notes coming from a thicket, and finally found the bird yards away from the spot whence its go-as-you-please voice seemed to come.

"A Chat courtship," says Mr. A. C. Webb, in "Some Birds, and their Ways," "is a sight never to be forgotten. In the spring, when birds begin housekeeping, the male Chat charms himself and his mate by some remarkable performances in the air. Launching himself from the top of some tall tree, he flutters from side to side, flirts his tail, stops, stands on his head, dangles his

<sup>1</sup> *Icteria virens*. Length, 7.25 inches.

legs as if they were broken, turns somersaults, and makes a monkey of himself generally, as he descends to the thicket below, where his mate is perched among the briars. Sometimes he starts from the low bushes and rises almost straight up into the air until he is above the tree-tops. He chatters and screams as he goes, telling her to watch him now as he comes down, and see if in all her life she ever saw a bird that could do such wonderful feats. No doubt to her eyes he is the picture of grace and elegance as he performs on his flying trapeze, but to us his clown-like antics seem ridiculous."

The Chat of the East is represented in the far West by a long-tailed variety, and between the two their range covers nearly the whole of the United States, British Columbia and Mexico.

**The American Redstart**<sup>2</sup> looks like a small, pinkish-yellow understudy of the Baltimore oriole,  $5\frac{1}{2}$  inches long. Its colors and color-pattern are very similar to those of our old friend of the elm-trees, velvety black on the back and head, reddish-orange on the sides and breast, and white on the belly. The tail is orange and black, and the colors are very prettily disposed.

On the whole, this bird has (in my estimation) the most beautiful color-pattern to be found in all our long procession of warblers and ground-thrushes. The female is so different in color it is at first difficult to believe her of the same species. Her body-colors are brownish-olive above with sides of pale yellow, and the head is gray instead of black.

This beautiful bird is to be looked for all over North America from Labrador and Fort Simpson to northern South America. In the North it arrives in May, and abides until September.

**The Water-Thrushes.**—Beginners in bird-study are warned to note the fact that in the Warbler Family are several birds called "*Water-Thrushes*," which do *not* belong to the Thrush Family. It is a pity that they have not been distinguished by some other name. There are two species, the **Common Water-Thrush**,<sup>3</sup> and the **Louisiana Water-Thrush**,<sup>4</sup> the first a northern, the latter a southern bird. Both live in the dark recesses of virgin forests, where clear brooks gurgle over mossy stones, between fern-

<sup>2</sup> *Se-toph'a-ga ru-ti-cil'la*. Length, 5.50 inches.

<sup>3</sup> *Se-i-u'rus no-re-bo-ra-ccn'sis*. Length, 6 inches.

<sup>4</sup> *S. mol-a-cil'la*.

covered banks. They are watchful and suspicious, but when flushed they do not immediately fly beyond gunshot, as nowadays every bird should do. The Louisiana Water-Thrush strongly resembles the wood-thrush, but is one-fourth smaller.

### THE VIREO FAMILY.

#### *Vireonidae.*

It is quite difficult to point out peculiarities by which the vireos can be distinguished from the warblers. They are placed next to the shrikes because of a supposed resemblance to those birds in the shape of the upper mandible—hooked and notched. The vireos look so much like warblers that only an expert can distinguish them.

The **Red-Eyed Vireo**<sup>1</sup> is distinguishable at close range by its red eye with a white line over it, and the **White-Eyed Vireo**<sup>2</sup> also is marked by the white color of its eyes. Both are fairly good songsters, and the former is about as "domestic," in its habit of frequenting the haunts of man, as the yellow warbler. The former ranges from New York northwestward across the continent, the latter only as far as the Rocky Mountains.

### THE SHRIKE FAMILY.

#### *Laniidae.*

The **Great Northern Shrike**, or **Butcher-Bird**,<sup>3</sup> is a bird of very striking personality. In appearance he is a high-headed, well-dressed dandy. In disposition, he is to-day a fierce little bird of prey, feeding solely upon flesh food; but to-morrow he will change into a modest insect-eater. It seems very odd to find a bird of prey among the Perching-Birds.

The Butcher-Bird is a bird of the North, breeding from Labrador to Alaska and visiting the United States only in winter, when it is almost impossible to obtain food at home. The species which we find in the United States in summer is the **Loggerhead Shrike**,<sup>4</sup> which closely resembles its northern relative, both in form and habits.

In the fields, you can easily recognize a Shrike

<sup>1</sup> *Vireo olivaceus*. Length, 6 inches.

<sup>2</sup> *Vireo noveboracensis*. Length, 5 inches.

<sup>3</sup> *Lanius borealis*. Length, about 10 inches.

<sup>4</sup> *L. ludovicianus*.

by his bluish-gray back, and large head. His strong, hooked beak has a notch, or tooth, near the end of the upper mandible. He is deliberate and dignified in his movements, and like the true sportsman that he is, he is happiest when hunting. He catches and feeds upon small frogs, mice, small snakes and even birds (so it is said), and has the odd trick of hanging up, impaled upon a thorn, dead game which he cannot eat as soon as caught. Once I saw a Butcher-Bird seize a large field-mouse out of a freshly upturned furrow, and fly away with it, struggling vigorously. The mammal was so large and heavy it was surprising to see the bird bear it away. Many times I have seen dried frogs hanging upon thorns, where they had been placed when fresh, by Shrikes.

Every Shrike is a feathered Jekyll and Hyde. In summer and autumn, the harvest of insects is everything that could be expected. In Dr. Judd's Bulletin No. 9, Biological Survey, Department of Agriculture, the list of groups of insects destroyed by the Loggerhead Shrike fills a



LOGGERHEAD SHRIKE.

page, and includes such pests as caterpillars, cut-worms, canker-worms, grasshoppers, crickets and weevils.

But mark the winter and early spring record. Thirteen species of small birds are numbered among the Loggerhead's victims, of which five are sparrows, and others are the ground-dove, chimney-swift, Bell's vireo and snow-bunting. The Butcher-Bird is known to kill twenty-eight species of birds, some of them valuable insect-

destroyers, and none of them to be spared without loss except the English sparrow. On the other hand, this bird is a great destroyer of wild mice, which in cold weather formed one-fourth of its entire food. The Loggerhead also feeds freely upon lizards, snakes, frogs and fish, when they are obtainable. The Butcher-Bird is a deadly enemy of the English sparrow, and kills and eats them so industriously that in Boston certain city officials once felt called upon to order the Shrikes to be shot.

The accompanying table is a very full exposition of the food habits of the two members of the Shrike Family referred to.

**THE WAXWING FAMILY.**

*Amplidae.*

**The Bohemian Waxwing.**<sup>1</sup>—Once, on a certain cold and bleak Thanksgiving spent on the banks of the Musselshell River in Montana, when the mercury stood at 8° below zero and the face of nature was a “gray and melancholy waste,” a flock of birds settled in the top of a dead cedar that stood near our camp. They were like so many exquisite gems, found ready cut and polished in a desert of rocks; and the whole camp quickly turned out to admire the exquisite creatures at short range.

TABLE SHOWING PERCENTAGES OF PRINCIPAL ELEMENTS OF FOOD OF THE BUTCHER-BIRD AND LOGGERHEAD SHRIKE, CALCULATED BY VOLUME.

By SYLVESTER D. JUDD, United States Biological Survey, Bulletin No. 9.

NAME.	VERTEBRATES.	INVERTEBRATES.	MAMMALS.	BIRDS.	OTHER VERTEBRATES.	GRASSHOPPERS AND CRICKETS.	SEIDERS.	WASPS.	CATERPILLARS AND MOTHS.	GROUND BEETLES.	OTHER BEETLES.	MONTH.	NO. OF STOMACHS.
Butcher-Bird . . . .	77	23	22	55	..	8	3	1	6	4	1	} Dec., Jan. and Feb. . . . .	33
Loggerhead . . . . .	76	24	55	14	7	13	1	..	2	4	6		14
Butcher-Bird . . . .	77	23	46	31	..	9	1	3	5	3	2	} March and April. . . . .	17
Loggerhead . . . . .	20	80	9	9	2	24	9	4	11	14	18		16
Do. . . . .	11	89	3	..	8	71	3	..	4	7	4	May and June . . . . .	17
Do. . . . .	2	98	..	..	2	67	2	4	1	5	19	July, Aug. and Sept. . . . .	22
Butcher-Bird . . . .	27	73	11	16	..	57	4	..	6	4	2	} October and Nov. . . . .	17
Loggerhead . . . . .	10	90	..	10	..	50	5	7	3	13	12		19
Butcher-Bird . . . .	60	40	26	34	..	24	3	2	6	4	1	} Average Oct. to April, inclusive . . . . .	67
Loggerhead . . . . .	28	72	16	8	4	39	4	3	4	9	13		} Average for the whole year . . . . .

The Great Northern Shrike is able to sing, but seldom does so; and many of his friends think he sings not at all. In summer it ranges all the way to Cook Inlet, Alaska, and in winter it migrates as far south as Virginia. In the southern states it meets the Loggerhead Shrike, and the two species so strongly resemble each other they are like two feathered Dromios.

I think that the Bohemian Waxwing, when alive and in perfect plumage, is one of the most exquisite perching-birds I know. It is not gorgeous or resplendent; but in dainty prettiness of form, immaculateness of plumage, and delicate refinement in color-scheme combined, it has few equals. The red wax-like tip on the

<sup>1</sup> *Am-pel'is gar-rul'us*. Length, 8 inches.



BOHEMIAN WAXWING.

end of each secondary feather gleams like a ruby. No picture of this bird ever can fairly portray its beauties. The **Cedar Waxwing** or **Cedar Bird**<sup>1</sup> of the eastern United States is but a fair understudy of its more robust and also more beautiful brother of the Northwest and the far North. Any one can instantly identify one of these birds by its jaunty top-knot, and the little drops of vermilion wax on the tips of its secondaries, eight on each side.

### THE SWALLOW FAMILY.

#### *Hirundinidae.*

The members of the Swallow Family are among the most sociable of our feathered friends, and also the most conspicuous.

The **Purple Martin**<sup>2</sup> loves the little house atop of a tall pole, which the country boy who loves birds takes pleasure in erecting for it. Forty years ago, thousands of the prairie farms

<sup>1</sup> *A. ce-dro'rum*. Length, about 7 inches.

<sup>2</sup> *Prog'ne su'bis*. Average length, 8 inches.

of the Middle West bore these tall monuments to the love of wild birds which *is born in every right-minded boy!* And how gracefully the glossy-black Martins used to circle, and swoop, and gyrate about them. Sometimes the blue-birds took possession of the martin-boxes, and then George or John was troubled; for having designed and erected on high a dwelling especially for the Martins, it seemed morally wrong that they should be forestalled, or crowded out.

And then came Ahab, the English sparrow, a homely, quarrelsome, low-minded and utterly uninteresting little wretch, a gutter-rat among birds. Unless coerced with a shot-gun, he steals the nesting-boxes of all other small birds, driving before him the Martins, bluebirds, and many others who used to love our company. In the North the Purple Martin does not seem to thrive away from the haunts of man, and I believe their great decrease in number has been due almost wholly to the English sparrow. It is really a bird of the South, but there was a time when it was common in at least some of the northern states.

The **Eave**, or **Cliff Swallow**<sup>1</sup> is still more sociable than the purple martin, and also more enterprising. With complete confidence in man's



PURPLE MARTIN.

good-will toward the bird-world, it chooses a barn that is big and high, and prosperous-looking, and calls it home. From the edge of

<sup>3</sup> *Pet-ro-chel'i-don lu'ni-frons*. Length, 5½ inches.

the nearest pond, it brings pellets of mud, and sticks a lot of them in a solid circle, against the outside wall of the barn, and close up under the eaves. Upon this, working most industriously to finish before previous layers have had time to dry, the cup-shaped nest is built out, pellet by pellet. At the last, the cup is narrowed down to a tube barely large enough to admit the bird, and the opening thrusts out into the air, usually tilted slightly upward.


All the members of a flock of Swallows build close together, nest joined to nest very frequently,

are the **Cliff, Bank** and **Tree Swallows**. The **Barn-Swallow** can be distinguished from these three by its very long and deeply forked tail, the tails of all the others being rather short.

### THE TANAGER FAMILY.

#### *Tanagridae.*

The male **Scarlet Tanager**<sup>1</sup> is one of the most showy small birds of our American Passeres. Excepting its wings, which are jet black, its entire plumage is of a clear scarlet hue, as bright



CLIFF-SWALLOW AND NESTS.

Nests under eaves of log house, photographed by E. R. WARREN.

and thus depends a most interesting Swallow town, usually called a "colony." Surely, any one who is not pleased and cheered by their sweet chattering and chirping under the eaves is "fit for treason, stratagems and spoils." Their flight is poetry expressed in motion. In catching the insects which constitute their food, they love to skim close to the surfaces of ponds and streams.

There are three swallows which so much resemble each other it requires a reference to a good handbook of birds to identify them. These

as the brightest ribbon. There is no precious stone which compares with it, for beside it the ruby is dull. The cardinal grosbeak is not nearly so bright as the male Tanager.

Wherever seen, the male Scarlet Tanager fixes the attention of the observer, and challenges admiration. It is an early spring arrival from the South, and in Washington, D. C., I have seen it in the parks while the trees were yet leafless. Some of those which came last year to the Zoological Park, New York, felt so secure

<sup>1</sup> *Pi-ran'ga e-ryth-ro-me'las*. Length, 6.50 inches.



BARN-SWALLOW.

*Hi-run'do e-ryth'ro-gas-tra.*

in our protection that they permitted their admirers to approach within ten feet of them.

The female of this species is widely different in color from the male, being dull olive-green above and greenish-yellow below.

**THE FINCH AND SPARROW FAMILY.**

*Fringillidae.*

This Family is a large one, and it embraces the perching-birds with strong beaks, such as the finches, sparrows, snow-birds and their near relatives, and one group of grosbeaks. By their beaks you shall know them,—short, and wide at the base, like the jaws of a pair of pliers. They are made for cracking all seeds which the owner does not wish to swallow entire.



**The American Cross-bill<sup>1</sup>** is a dull-red bird with brown wings and tail, and its bill is so emphatically crossed it seems like a deformity which must necessarily

be fatal to a seed-eater. But Nature has her own odd ways; and it seems that the scissor arrangement of this bird's beak is to promote the husking of pine cones, and the cracking of the seeds.

This is a bird of the North, and in the East comes no farther south than a line drawn from <sup>1</sup>*Lox'ia cur-vi-ros'tra minor.* Length, 6.50 inches.

Colorado to Washington, D. C. In the West it descends to Arizona, but everywhere in the United States it is only a winter visitor. With an opera-glass it is always easily recognized by its crossed bill.

**The American Goldfinch<sup>2</sup>** is a conspicuously yellow bird, though quite small. It is a plump-bodied, fluffy little bird, all sulphur yellow except a circular black cap atop of its head, and black trimmings on its wings and tail. It is exquisitely pretty, and, like a feathered coquette, loves to pose on the steep side of a tall mullein stalk, with no leaves about to cut off the admirers' view. It is sociable, also, and loves the garden, orchards and meadows of the self-elected "lord of creation," man.

As a weed-destroyer, this bird has few equals. It makes a specialty of the seeds of members of the Order Compositae, and is especially fond of thistles, ragweed, wild lettuce and wild sunflower. (Sylvester D. Judd.)

**The Snow-Bunting<sup>3</sup>** comes down from the far North, in the dead of winter, when the snow



SCARLET Tanager.  
Male and female.

falls fluffy and deep, and the song-birds of summer are basking in the sunshine of the South. They do not appear every winter, however.

<sup>2</sup> *As-trag-a-li'nus tris'tis.* Length, 5 inches.  
<sup>3</sup> *Pas-ser-i'na ni-val'is.* Length, 6½ inches.



AMERICAN GOLDFINCH.

They come in flocks of from ten to twenty birds, and settle in the snow as if they love it. But for a few dark streaks on back and wings, they are the color of snow, and generally have the plump outlines which betoken good feeding and contentment.

When you see this bird, remember that it belongs to the *polar world*, quite as much as the arctic fox and musk-ox, and in summer it goes to the "farthest north" on our continent. Rarely indeed does it breed in even the most northerly portions of the United States, and seldom enters a southern state.

In winter the food of this pretty bird consists chiefly of the seeds of weeds that send tall fruit-stalks above the level of the snow. In our park grounds, we scatter wheat for it, on the

tops of granite ledges from which the wind has blown the snow.

The **Slate-Colored Junco**,<sup>1</sup> often called the **Snow-Bird**, is also a bird of the snow-fields; but it is a home product rather than a visitor from the desolate Barren Grounds. When seen on snow, its slaty-blue back makes it appear like a dark-colored bird, but underneath it is dull white. Like the snow-bunting, it goes in small flocks, and in winter feeds chiefly upon weed-seeds and grain. It breeds in our northern states, and in winter migrates southward almost to the Gulf of Mexico. Altogether, thirteen species and varieties of Juncos are recognized in North America, and they are at home all the way from Alaska to Mexico and the Gulf.

**The Sparrows.**—There was a time when in America it was not only respectable but even honorable to be a Sparrow; but during the past ten years, the doings of one alien species, most unwisely introduced here have tended to bring the name into disrepute. How our native species must hate the interloper! But we protest that our native Sparrows are as sweet-voiced and interesting as ever they were; and as wholesale destroyers of noxious weeds, they are unsurpassed. After a careful investigation of the quantity of weed-seeds consumed in Iowa by the **Tree-Sparrow**,<sup>2</sup> Professor F. E. L. Beal calculated the total amount for one year to be 1,750,000 pounds, or about 875 tons! Practi-



SNOW-BUNTING.

cally without exception, all our Sparrows are diligent consumers of the seeds of noxious weeds.

If you doubt the vocal powers of Sparrows go

<sup>1</sup> *Jun'co hy-e-mal'is*. Length, 6 inches.

<sup>2</sup> *Spi-zel'la mon-ti-co'la*. Length, 6 inches.



with me to the country roads, and listen for three minutes to the delicious melody that pours from the quivering throat of a **Song-Sparrow**.<sup>1</sup> When he feels well, he will perch on the top of a hedge, secure a good grip on a comfortable twig, point his beak skyward at an angle of sixty degrees, and sing as if trying to burst his little throat. Mrs. Mabel Osgood Wright justly calls him "the darling among the song-birds," and "the most constant singer among our northern birds." In some localities, at least, they sing all summer long. In Iowa I have heard them a thousand times, bravely piping and trilling in the sweltering heat of July and August, when other birds were silent, and have been moved to wonder at the amount of energy stored up in their little bodies.

I think the best way to identify this bird is by its singing. Pick out the sparrow in gray and brown which sings to surpass all others, and it will be a Song-Sparrow. Its home is the eastern half of North America, from northern Manitoba to Mexico. West of the Rocky Mountains it becomes the *Mountain Song-Sparrow*. In the southwestern deserts it grows pale,—to match its environment,—and becomes the *Desert Song-Sparrow*. There are thirteen species of the Song-Sparrow genus,—or at least that number have been described, and Alaska is yet to be heard from.

**The White-Throated Sparrow**<sup>2</sup> is the species which comes next in general attractiveness. It is a very pert and pretty bird—for a sparrow, and its oddly marked head is easily identified. It wears a white goatee and a black cap, and on the latter is laid a broad arrow, in white. A white line comes down along the centre line of the head, and another comes forward over each eye, until the three come together at the base of the upper mandible. The song of this bird is pleasing, and nearly every self-respecting ornithologist translates it into English to suit his or her fancy; but, to tell the truth, the White-Throat never will win a prize as a great singer.

**The English Sparrow**.<sup>3</sup>—Let me dip my pen in blue vitriol; for my temperature rises at the thought of writing the name. Daily we see the unclean little wretches grubbing in the filth

and microbes of the street, where a native American bird will humble itself to feed. After twenty years of acquaintance, I am obliged to say that I never saw one catch a worm, a caterpillar, or an insect of any kind. When the elm-trees are loaded with tent caterpillars, an English Sparrow will let them crawl all over him, and not kill one. Instead of ranging out into the open fields and hunting for clean weed-seeds, this bird revels in the foulest dirt of the street. It does, however, manage to eat the seeds of the



WHITE-THROATED SPARROW.

dandelion, when the heads are filling, in April and May.

The English Sparrow is not beautiful, either in form or plumage, and it cannot sing a note. Its tastes are low and vulgar. It is quarrelsome, and crowds out many other species of small perching-birds. In Cheyenne, Wyoming, when Mr. Frank Bond killed all the English Sparrows, and kept them killed, other perching-birds flocked into the city in great numbers, and many species bred there. The more persistently these

<sup>1</sup> *Mel-o-spi'za fas-ci-a'ta*. Length, 6½ inches.

<sup>2</sup> *Zo-no-tri'chi-a al-bi-col'l'is*. Length, 6½ inches.

<sup>3</sup> *Pas'ser do-mes'ti-cus*. Length, about 6 inches.

interlopers are killed off, the better for all other birds. They can be made to serve well as subjects for dissection in the school-room, and for amateur taxidermists; and they make excellent food for captive hawks, owls, small carnivores, and live snakes of several species.

The introduction of this bird may well serve as a solemn warning against any further meddlings with Nature on that line. In the first place, there never existed the slightest reason



CARDINAL.

or need for this importation. Without serious consideration, or consultation with the persons most competent to advise, this bird was imported and planted in twelve widely separated localities in the United States. To-day it is a feathered nuisance that spreads over one-half the United States, and excepting locally cannot be abated. Nevertheless, it is within the power of western towns and cities wherein it has not yet gained a foothold to follow the example of Mr. Bond in Cheyenne, and destroy every colony that enters before it has time to breed.

**The Cardinal**, or **Cardinal Grosbeak**,<sup>1</sup> also called the **Cardinal Redbird**, is the pride of the South. From New Orleans to New York it is persistently trapped and "limed,"—not to "keep" as a cage-bird, but to sell as such. Poor, unhappy Cardinal! How much better its fate

<sup>1</sup> *Car-di-nal'is car-di-nal'is*. Length, 8.50 inches.

had it been created black instead of bright cardinal red, with no jaunty top-knot, and no fatal gift of song!

In a cage 6 by 9 feet, or even 4 by 4, a bird like this flies to and fro, and in company with a dozen other small birds finds life far from dull. But if you put a wild song-bird in a cage barely large enough for a canary, the bird is wretched, it dies soon, and the keeping of it is a sin against Nature. Excepting canaries and a very few other species, if you cannot keep birds (and mammals, also) in big cages, *do not keep them at all!* The way thousands of song-birds are caught in some portions of the South, to sell as cage-birds, is a sin and a shame. At this date, New Orleans in particular has before her an imperative duty in breaking up this business. Children everywhere should be taught that it is *almost impossible for any one save an expert bird-man to take young song-birds, and rear them successfully*. Young insectivorous birds require specially compounded bird-food, and it must be given to them *every hour*, with small forceps—a very tedious operation.

In the kindness of their little hearts, children often take young song-birds from the nest, cage them, and try to feed them on what some little folks like best—cake and cream! They might as well give them poison! For any one ignorant of the precise methods necessary in rearing insectivorous birds, to take such birds from their parents is *cruelty and destruction!*

The sight of a wild Cardinal always compels attention. The bird is not only beautiful in color, but it is aristocratic in form and manner. It comes up from the South into New York state, and the Ohio River region, and extends westward to the edge of the plains region.

**The Rose-Breasted Grosbeak**<sup>2</sup> is, in all respects save one, a very beautiful bird. It has a big, clumsy-looking, conspicuously white beak, which almost spoils the whole bird. But the pink-sunset flush on the clear sky of its breast, its glossy-black head and tail, and black-and-white wings, are so beautiful a combination they lead one to forgive the homely beak. The delicate pink-rose tint on the breast renders the identification of this bird very easy, even at first sight.

I must confess that I remember nothing of

<sup>2</sup> *Zam-e-lo'di-a lu-do-vi-ci-an'a*. Length, 8 inches.

the Grosbeak's song, and that it made no impression on me, even when these birds were around me. Certainly it is no great singer, not more than third-rate, at the best, or its song would be more in evidence. It is celebrated as an enemy of the potato-bug, and it feeds omnivorously upon other insects, buds, blossoms, seeds and fruit.



ROSE-BREADED GROSBEAK.

The range of this species is bounded by the great Rocky Mountain barrier. Westward thereof is found the *Black-Headed Grosbeak*, and the arid lands of Texas, Arizona and southern California are inhabited by the *Western Blue Grosbeak*.

The bluest bird that flies in North America is the **Indigo Bunting**,<sup>1</sup> a trim little craft, built and rigged like a warbler, and of warbler size. Like the ocean, it is

“Deeply, darkly, beautifully blue,”

—not the sky-blue of the jay, but like indigo. In the East you cannot possibly mistake it. The deep-blue bird of the Far West is the *Lazuli Bunting*, our bird's nearest relative.

<sup>1</sup> *Cy-a'-no-spi-za cy-a'-ne-a*. Length, 5.50 inches.

## THE BLACKBIRD FAMILY.

*Icteridae.*

This Family includes several showy species of birds which are very much in evidence, and quite generally known to country dwellers. Five representative and very interesting species will be noticed.

The **Bobolink**<sup>2</sup> is a bird with two very distinct characters. It has a name and a suit of feathers for the North, another suit and another name in the South; and it has three reputations.

When in springtime a certain jolly and vigorous little song-bird comes up from the South, he puts on a dress-suit of marvellous design, in black, white, brown and gray, as shown on page 178. He is then a regular swell, and his name is Bobolink. His mate, however, is a plain little bird, clad in yellowish-brown, with slight trimmings of yellow and white. They frequent the marshes and low meadows, nest on the ground, and rear from five to seven young.

That accomplished, the male bird doffs his pretty spring suit, acquires plumage like that of the female, and then they go South. There they become **Rice-Birds**, and they raid the rice-fields of the southern states until they grow quite fat. Next, enters the Man-With-a-Gun; and the birds fall easy victims. The birds are shot for two reasons: The rice-planters find them a nuisance in their fields, and many people think Rice-Birds are good eating.

Consider the “**Reed-Bird** on toast,” or, worse still, “on a skewer.” It is a trifle too large for one mouthful, but by no means large enough for two. A healthy, able-bodied American at work upon this two-ounce bird with a ten-inch knife is a sad but impressive spectacle. It is to be hoped that it will be long ere the people of this country really have cause to turn to this tiny song-bird—or any other song-bird—as a source of food with which to satisfy hunger. How can any self-respecting man deliberately order so pusillanimous a dish as “Reed-Birds on a skewer?” There is a land so populous and poor that its people eat sparrows because they need them for food; but it is far from America.

The Bobolink is really a very acceptable singer,

<sup>2</sup> *Dol-i-cho'nyx o-ry-ziv'o-rus*. Length, 6.75 inches.



RED-WINGED BLACKBIRD.

and has furnished a theme for several poets, of whom Bryant was the most celebrated.

**The Red-winged Blackbird**<sup>1</sup> is a bird that delights my soul. No marsh or cat-tail swamp is complete without him. No spring ever can be perfect without his sweet, liquid gurgle—"O-ka-leé." You hear that flute-like call when the sky is clear, the warm sunshine is flooding field and stream, and you are glad that you are alive.

The jet-black plumage of this bird, with epaulettes of scarlet and white, make a brave show among the dull green blades of the cat-tails. As a rule, bird-songs translated into English do not appeal to me very strongly; but the Red-Wing does say "O-ka-leé" to perfection!

The typical Red-Wing is an eastern bird, but its half-dozen subspecies are so well dispersed throughout the United States that almost every region possesses one.

**The Yellow-headed Blackbird**<sup>2</sup> is a very conspicuous species throughout the West, from Indiana almost to the Pacific. Its entire head, neck and breast are of a dull-yellow color, but elsewhere it is wholly black, save a white patch on the wing. In Montana it is very common in summer.

**The Meadow-Lark**<sup>3</sup> is one of the most trustful and sociable of our birds. As its name implies, it is partial to open grass-lands, and its fa-

<sup>1</sup> *Agelaius phoeniceus*. Length, 9 inches.

<sup>2</sup> With an apology to the reader, it is stated that the Latin name of this bird is *Xanthocephalus xanthocephalus*: and its length is about 9 inches.

<sup>3</sup> *Sturnella magna*. Length, about 10 inches.

vorite hunting-grounds are the bits of waste land in sloughs (called "slews" in the West), that are full of low weeds. This bird does not like tall grass or weeds, for it is very curious to know all that is going on in the world about it. It is an indifferent flyer,—slow and short,—and manifests a decided preference for the haunts of man.

The Lark contributes much to the pleasures of life on a farm. Its bright-yellow breast and throat, with a jet-black neck-scarf, are as cheerful as an April sunburst. The long, conical beak, rather long legs and erect carriage of this bird give him an air of cheerful confidence which says to you, "I'm a good fellow, and you're another!" His song is nothing to boast of, but he always pipes up cheerfully, and does the best he can. I always liked this bird, and count him as one of the dear friends of my boyhood. To me, his plumage is beautiful, especially when seen on a fresh, dewy morning, when the sun is newly risen, and the song-birds are greeting the new day.

According to the investigations of the Biological Survey, the Meadow-Lark is one of the most valuable of all birds that frequent farming re-



MEADOW-LARK.

gions. Throughout the year, insects make up 73 per cent of its food, grain, 5 per cent, and weed-seeds, 12 per cent. During the insect season, insects constitute over 90 per cent of this bird's food supply. As a destroyer of insects and weeds, this bird is entitled to the most perfect protection that laws and public sentiment combined can afford.

In Montana, the **Western Meadow-Lark**<sup>1</sup> quite wearied me by the tiresome iteration, day after day, of its one short, seven-word song. This was it:



As our "outfit" pulled over the smoothly shaven Missouri-Yellowstone divide, in the month of May, I think we heard that song repeated a thousand times, or less; and when the wind blew hard for five long days without intermission, even that cheerful welcome at last became irritating.

The eastern Meadow-Lark inhabits the eastern half of the United States, and the western species begins at the western edge of Iowa and Missouri; *but neither of them belongs to the Lark Family!*

**The Baltimore Oriole**,<sup>2</sup> or **Hang-Nest**, has beautiful plumage of orange and black, a very pleasing song, good habits, and therefore is one of our feathered favorites. Either when perching or on the wing, it is a very graceful bird. It is the most skilful builder in North America, and constructs a strong and durable hanging nest which is a marvel of intelligent and skilful effort. The Oriole does not believe in having boys make collections of Oriole eggs. The outermost branches of a very tall and very drooping elm are particularly suited to its views of an ideal building site.

The nest of this Oriole is bound to create in the mind of any one who examines it attentively a high degree of admiration for the mental capacity of its builder. Its superstructure is composed very largely of long, spring-like horse-hairs, so tightly woven together that even when the end of a hair waves freely in the air, it is im-

<sup>1</sup> *Sturnella neglecta*. Average length, about 9.50 inches.

<sup>2</sup> *Icterus galbula*. Length, 8 inches.

possible to pull it out. Here is genuine *weaving*, done with hair and fibrous fragments of soft, weathered bark. Let it be remembered at this point that not even the higher apes know how to *weave* a nest or a roof.

The mouth of the Oriole's bag-like nest is thin but strong, and terminates in an edge as thin and firm as hair-cloth. A nest now before me is



BALTIMORE ORIOLE AND NEST.

five inches long, four inches in outside diameter at a point half-way between bottom and top, and its opening is two inches in diameter. For a space of two inches, the horse-hairs of the upper margin are wrapped around an elm-twig the size of a slate-pencil. *At no point* are the walls more than a quarter of an inch in thickness, and the inside is as symmetrical and shapely as if the nest had been woven around a form.

The usefulness of the Baltimore Oriole is fully equal to its beauty. As a destroyer of cater-

pillars it has few equals among birds. In May, insects constitute 92 per cent of its food, and in April and July 70 per cent. For the entire year, animal food, chiefly caterpillars and beetles, constitute 83.4 per cent of its food, and vegetable matter the remainder of 16.6 per cent.

**The Purple Grackle, or Crow Blackbird,**<sup>1</sup> has prompted scores of persons to ask, "What is the name of that very shiny, jet-black bird with a long tail?" No wonder it attracts attention, especially in contrast with the lustreless rusty blackbird. Its color is deep purple-black, and it is as shiny as if it had been varnished all over. It loves to follow the plough, and pick up the big, fat grubs that are exposed to view, before they have had time to burrow out of sight. Often in their eagerness not to miss a chance, these birds will approach within ten feet of the plough-handles. It is then that one notices that their eyes are light yellow, and very odd-looking. This bird has no song, and its sign of contentment with life is like a great asthmatic wheeze. The tail of this bird is creased lengthwise along the middle, or "keeled."



PURPLE GRACKLE.

Prior to the systematic investigations of the Department of Agriculture the value or harmfulness of the Crow Blackbird was in dispute. The examination of 2,346 stomachs revealed that during an entire year the food supply of this bird is made up in the following percentages: insect food, 26.9; other animal food, 3.4; corn, 37.2; oats, 2.9; wheat, 4.8; other grain, 1.6; domestic fruit, 2.9; wild fruit, 2.1; weed-seed,

<sup>1</sup> *Quis'ca-lus quis'cu-la*. Length, about 12 inches.

4.2; mast, 14; total, 100. "The charge that the blackbird is a habitual robber of birds' nests is disproved by the stomach examinations." (F. E. L. Beal.)

### THE CROW FAMILY.

#### *Corvidae.*

Take them all in all, there is no Family in the whole Order of Perching-Birds whose members have more striking individual traits, or more commanding personality than the Family which contains the ravens, crows, jays and magpies. All these birds are bold and conspicuous, and fond of entering into the affairs of man. The crow feels it to be his duty to assist in planting-operations. The blue-jay robs you, and scolds while he does it. The magpie will hold a fifteen-minute conversation with you, and tell you of all his troubles. Go where you will in the United States, some of the twenty species of birds of this Family will cheerfully bear you company.

**The American Magpie,**<sup>2</sup> of the somewhat "wild West," is a beautiful and showy bird, and in winter especially it bravely strives to adorn the bare and bleak valleys, foothills, divides and mountain-sides of the Rocky Mountain region. In the whole of the West, I know of no bird more beautiful in flight than this. Its plumage is half glossy-purple black, and half snow-white; and this, with its extremely long tail streaming after it in its flight, makes it a very striking object. In winter the absence of other birds renders the Magpie trebly conspicuous and welcome. Its flight is slow, dignified, and as straight as an arrow.


The Magpie is fatally fond of fresh meat, and many a fine bird meets its death by devouring poisoned meat laid out for wolves. If hospitably received, this bird will come close to the haunts and camps of man, investigating everything, and looking for scraps of food. If not fired at, it soon becomes very friendly, and a small cabin easily becomes the haunt of a score of birds. Some of those in the Flying Cage of the New York Zoological Park are at times as amusing as monkeys. They come close up to the wires, and when the visitor bends down, to listen or converse, they actually *talk*—in their language. In low, confidential tones they tell

<sup>2</sup> *Pi'ca pi'ca hud-son'i-ca*. Length, about 18 inches.

of their fear of the big condor, the painful pecks they get from the herons, and the greediness of the ducks in devouring all of their kind of food.

In the days of elk and buffalo slaughter, the Magpies feasted continually upon fresh meat. Now they make friends with the ranchmen, and eat all kinds of food. This interesting bird ranges from Alaska, and the edge of the arctic barrens, southward through the great plains and mountains to the arid regions of the South-

Besides his harsh "Jay," a crow is a sweet songster. He will take your cherries right before your eyes, and then scold you roundly for not looking pleasant about it! He robs the nests of other birds, eating eggs or young, whichever may be there; and to that extent he is a pest. During the closed season on eggs and young nestlings, he lives on insects—until berries and small fruits ripen. If Jays were as numerous as English sparrows, it would be necessary to



AMERICAN MAGPIE.

west. It is easily kept in confinement, if provided with a large cage and a suitable house, *out-of-doors*.

The **Blue-Jay**<sup>1</sup> needs no description—only toleration; for his reputation would be all the better for washing. He is a bird of unbounded assurance, and being well known as a marauder, it is only his audacity which saves him from extermination. Externally, he is really a beautiful bird, but his voice is strident and unmusical.

<sup>1</sup> *Cy-an-o-cit'ta cris-ta'ta*. Length, 11.50 inches.

reduce their number; but they are not so numerous or so destructive that we need to attack them.

**Steller's Jay**<sup>2</sup> is one of the handsomest birds of the moist and dark forest region of the Pacific coast, which extends from Mount St. Elias to San Francisco Bay. It is also the type of three subspecies, or varieties, found farther east and south. It is the Pacific coast counterpart of our blue-jay,—high-crested, barred with black

<sup>2</sup> *Cy-an-o-cit'ta stel'ler-i*. Length, 12.50 inches.

on wings and tail, and with blue as its prevailing color.

**The Piñon Jay**<sup>1</sup> (pronounced pin'yone) is a bird well worth knowing. On the Sierra Nevada mountains and adjacent plateaus, where the piñon pine, juniper and cedar bravely struggle against the scarcity of water, and only half clothe the rugged nakedness of Nature, this Jay is a welcome habitant. I think it safe to say that you will find it wherever you find the piñon



BLUE-JAY.

pine, whose big, husky cones furnish a generous quantity of seeds, called "nuts," which are good for man, and grand food for all the wild creatures that can crack their delicate shell.

I have never seen the Piñon Jay so numerous that it could be called a "common" bird throughout an extensive region. At the same time, it is a bird of social habit, and given to flocking, quite like our eastern crow. It is really a connecting link between the crows and jays. It has a short, square tail, no crest or "top-knot;" its predominating color is grayish-blue, and its cry is a crow-like "caw."

**Clarke's Nut-Cracker**<sup>2</sup> is a bird of the western mountain-tops and canyons, and a companion of the mountain-sheep. Wild creatures that love to dwell on high mountains, amid grand scenery, appeal to my affection more strongly than some others. To me, this bird recalls

<sup>1</sup> *Cyanocitta stelleri* *cy-an-o-ceph'a-lus cy-an-o-ceph'a-lus*. Length, 11 inches.

<sup>2</sup> *Nucifraga columbiana*. Length, 12 inches.

pictures of mountain-parks, "rim-rock," "slide-rock," pines and cedars bravely climbing up steep acclivities, gloomy canyons, and rushing streams of icy-cold water below all.

I first made acquaintance with this bird while hunting elk and mountain-sheep, on a fearfully steep mountain-side, with a magnificent panorama spread out below. It greeted me in friendly fashion with the rasping "Kurr, Kurr!" which, when heard amid such surroundings, is not soon forgotten. It has been my misfortune, however, never to see the remarkable habit thus graphically described by Mrs. Florence Merriam Bailey in her delightful "Handbook":

"Living mainly on the crests of the ranges, the birds fly to the high peaks to get the first rays of the sun, and when warmed go for food and water to the lower slopes. Their method of getting down is startling at first sight. Launching out from a peak, with bill pointed downward and wings closed, they drop like a bullet for a thousand feet, to the brook where they wish to drink. Sometimes they make the descent at one long swoop, at other times in a series of pitches, each time checking their fall by opening their wings, and letting themselves curve upward before the next straight drop. They fall with such a high rate of speed that when they open their wings there is an explosive burst which echoes from the canyon walls."

The head, neck and body of this bird are uniform ashy gray, and the wings and tail are black, with a white patch half-way down the former. The Nut-Cracker is really a small crow, twelve inches long, and much resembles the common gray and black crow of Europe. It is found in all the mountains of the West, from Alaska to Mexico, and straggles eastward to the eastern edge of the Great Plains. It is often called **Clarke's Crow**.

**The Canada Jay, Whiskey-Jack, or Moose-Bird**,<sup>3</sup> is by reason of its personal oddities and assertiveness perhaps the most conspicuous and widely known of all the perching-birds of the great coniferous forests of Canada. Every man who has trailed moose or caribou, or for any reason has camped in the Laurentian wilderness, knows well this audacious camp-follower, and remembers him with interest, if not even friendship. He has no real song, and his cries are

<sup>3</sup> *Perisoreus canadensis*. Length, 12 inches.



rather harsh and strident ; but in his native solitudes, where bird-sounds are so seldom heard, the voyageur is always glad to hear his call. And surely, every perching-bird that chooses to brave the rigors of the northern winter instead of migrating is entitled both to respect and admiration.

The plumage of the Canada Jay has a peculiar fluffy appearance, suggestive of fur. Its prevailing color is ashy-gray. The nape and back of the head are black, but the forehead is marked by a large white spot. The wings and tail are of a darker gray than the body. The home of this interesting bird—the companion of the moose, as well as of forest-haunting man—extends from Nova Scotia, and northern New England, throughout Canada to Manitoba, and northward to the limit of the great forests.

**The Common Crow**<sup>1</sup> needs no description. When finer birds were abundant, we cared little for him; but now that bird-life generally has so greatly diminished, we feel like welcoming him as a friend. His cheerful "Caw" is a welcome sound, and his services to the farmer overbalance the bad things he perpetrates. The Department of Agriculture, through Professor F. E. L. Beal, has officially investigated him, published the court records of his case, and pronounced him a bird worthy of protection. It is declared, after an examination of the stomachs of specimens, that the noxious insects destroyed by the Crows—cut-worms, caterpillars, grasshoppers, and also mice—represent a saving of more grain than the bird consumes.

It must be admitted, however, that the Crow does many things he should not. He is too fond of eggs, and also of young birds. He will pull up, by the roots, altogether too much newly planted corn ; which is very unfair toward the farmer. While the damage is seldom serious, it is always annoying; but when the Crow passes the limit of human endurance, powder and lead are his portion. For example: when a Crow nesting in Beaver Valley elected to make visits to our duck-pond where young wild-ducks were hatching, and take three mallard ducklings in one morning, Curator Beebe was compelled to choose quickly between ducks and Crows, and provide for the survival of the fittest.

<sup>1</sup> *Corvus a-mer-i-can-us*. Length, 18 to 20 inches.

**The American Raven**<sup>2</sup> is a bird of the "wild West," quite rare, and seldom seen beyond the mountains. Even when you see it for the first time, you will readily recognize it by its all-black plumage, large size, slow and heavy flight, and its hoarse and seldom "Quock!" The crow is at all times a cheerful citizen, but the Raven always has a sore throat, and is always going to a funeral.

He lives with Clarke's nut-cracker and the other dwellers on the mountain-tops north of the arid regions of Arizona and New Mexico, and nests in the crevices of high, rugged cliffs or canyon walls that are as completely inaccessible as can be found. He is suspicious of all attentions, wants no companions save of his own kind, and



Photographed by E. R. WARREN.

CLARKE'S NUT-CRACKER.

mighty few of those. The "Quock" of a Raven in a rock-ribbed and gloomy canyon is anything but a cheerful sound.

Like the vulture, this bird feeds upon dead animals, dead fish, and sometimes also upon the poisoned meat that wolfers distribute so generously.

<sup>2</sup> *Corvus co-rax sin-u-a-tus*. Length, 22 to 24 inches.

**THE HORNED LARK FAMILY.***Alaudidae.*

There is a Lark Family which we regret to say does not include the *meadow-lark*; for this separation of birds bearing the same general name tends to create confusion. In Europe the Lark Family is a very large one, and contains about a hundred species, the most celebrated of which is the unfortunate skylark. It is unfortunate because of the wholesale and heartless manner in which it is caught and kept in pitiful captivity as a "cage-bird." In London these wretched little creatures are sold by the thousand, sometimes at sixpence each, or even less.

Strange to say, in America the Lark Family is represented by only twelve species and subspecies, of which the **Horned Lark** or **Shore Lark**<sup>1</sup> is the best type. It is called "Horned" Lark because of a small, pointed tuft or brush of feathers which lies along the side of the head above the eye, pointing backward and thrusting its tip through the regular outline of the back of the head. The resemblance of these points to horns is quite far-fetched, but it seems to have been brought in to stay.

This bird looks very much like a small plover. Our eastern species is by habit a shore-bird, whence its second name. It comes to us in winter, in flocks of from six to twenty individuals, and at that season its plumage is not so bright and pleasing as in spring.

The West and Southwest are inhabited by nine subspecies of Horned Larks, ranging all the way from Mexico to British Columbia, some of them necessarily living in hot countries, and far from large bodies of water.

**THE FLYCATCHER FAMILY.***Tyrannidae.*

There are many little birds, in size next above the sparrows, which look as if they ought to sing;

<sup>1</sup> *O-toe'o-ris al-pes'tris*. Length, 7.50 inches.

but in reality they do not. They are very expert at catching insects, however, and nothing that flies can escape them in mid-air. These birds make up the Family of Flycatchers, and to the farmers of this country every flycatcher is worth double its weight in pure silver. Altogether there are about thirty species.

**The Kingbird**,<sup>2</sup> also called the **Bee "Martin"** and **Bee-Bird**, may well stand as the representative of this Family. Whenever you see a small bird swiftly and actively chasing a large crow in mid-air, darting down upon the back of the black fellow every hundred feet or so, with a peck that sends a thrill of life along his keel, you may know that the gallant little warrior is a Kingbird, and it is driving the crow away from the vicinity of its nest. The performance is like that of a man and a mad hornet. The crow thinks not of battle, but only of getting on in the world, and giving the nestlings of his tormentor a good square mile of crowless space in which to grow.

Look long enough, and you will see the Kingbird return from the chase, perch on his favorite dead limb at the edge of the field, smooth his feathers and renew his watch for flying insects. Presently you will see him dart from his perch, swoop to a certain point in space, and then return to his place. He has caught some flying insect, and like Oliver, "wants some more."

Never shoot a Kingbird. It is easier to "identify the species" on the wing than lying dead, all shot to pieces. Without killing this most courageous of all birds—which can whip almost anything that wears feathers, but attacks only crows and hawks—you can see that its colors are bluish-gray, trimmed with black and white.

The *Crested Flycatcher*, the dear little *Phabe-Bird*, and the *Wood-Pewee* belong to the Flycatcher Family.

<sup>2</sup> *Ty-ran'nus ty-ran'nus*. Length, 8 inches.

## CHAPTER XVII

# THE ORDER OF ODD FAMILIES

### MACROCHIRES

With certain exceptions, the different Orders of American birds are founded on reasonable grounds and built up of homogeneous materials. As a rule, a few moments' examination of a bird enables one to name the Order to which it belongs. There is no difficulty about the birds of prey, swimmers, fishers, waders or woodpeckers.

Unfortunately, however, Nature has turned out of her workshop so many odd forms that it has been found necessary to have a certain number of Orders for them. In mammals we have seen that the Order *Ungulata* is of this character. In birds, there are two such Orders. One is that which contains the cuckoos, road-runners and kingfishers, and the other is that which forms the subject of this chapter.

The Order Macrochires means literally "odd ones," and its members do not belie the name. On the strength of certain resemblances in anatomical structure, observable only after the birds are dead and dissected, our humming-birds, swifts and goatsuckers (*i.e.*, birds like the whippoorwill and night-hawk) are grouped together in this Order, in three Families, as follows:

	FAMILIES.	EXAMPLES.
ORDER MACROCHIRES.	GOATSUCKERS, . . . . . <i>CAP-RI-MUL'GI-DAE.</i>	. . . Nighthawk, Whippoorwill.
	SWIFTS, . . . . . <i>MI-CRO-POD'I-DAE.</i>	. . . Chimney-Swift.
	HUMMING-BIRDS, <i>TRO-CHILI-DAE.</i>	. . . Ruby-Throated Humming-Bird.

### THE GOATSUCKER FAMILY.

#### *Caprimulgidae.*

The Nighthawk<sup>1</sup> is far from being a true hawk. It belongs to a Family of birds which have soft, owl-like plumage, and enormous mouths, fringed above with a row of stiff bristles, for use in capturing insects on the wing. Many years ago, when people believed many things that were not true, some believed that these big-mouthed birds sucked goats; hence the absurd name applied to the Family.

Whenever, during the hour just before sunset, you see a good-sized bird with dark plumage, long, sharp-pointed wings, and a big white spot on the under surface of each wing,—wheeling, soaring, dropping and circling through the air,—you may know that it is a Nighthawk, catching insects. Its flight is graceful and free, and when

<sup>1</sup> *Chor-dei'les virginianus.* Length, about 9.50 inches.



THE NIGHTHAWK.

on the aerial war-path it is a very industrious bird. Some people compare this bird on the wing with bats; but I see no resemblance save the bare fact of semi-nocturnal flight.

When this bird alights upon a tree to rest, it chooses a large and nearly horizontal limb, on which it usually sits lengthwise. As it sits motionless on a large limb, the bird strongly resembles a knot. This is a trans-continent bird, being found from the Atlantic to the Pacific, in wooded regions, and northward to the Mackenzie River.

**The Whippoorwill<sup>1</sup>** needs no introduction. It is more than a bird. It is a national favorite.

When the mantle of night has fallen, and the busy world is still, we who are in the country in summer often hear a loud, clear, melodious whistle from somewhere near the barn. As plainly as print, it exclaims, "*Whip-Poor-Will!*" and repeats it, again and again. Before each regular call, there is a faint "*chuck,*" or catching of the breath, strong emphasis on the "*whip,*" and at the end a piercing whistle which is positively thrilling.

Sometimes the bird will come and perch within thirty feet of your tent-door, and whistle at the rate of forty whippoorwills to the minute. Its call awakens sentimental reflections, and upon most persons exercises a peculiar, soothing influence. It has been celebrated in several beautiful poems and songs.

The range of this interesting bird is the same as that of the nighthawk. In the South, both are replaced by another goatsucker called, from its whistle, the **Chuck-Will's-Widow**. Until actually hearing it, one can scarcely believe that any bird of this Order can say things as plainly as this bird says "*Chuck Will's Wid-ow!*" The Pacific states, from British Columbia to Mexico, and eastward to Nebraska, have the **Poor-Will**.

### THE SWIFT FAMILY.

#### *Micropodidae.*

**The Chimney-Swift, or Chimney-"Swallow,"**<sup>2</sup> has been for a century or more classified with the swallows and martins, but recent studies of its anatomy have caused its removal from their group. This is the bird whose nest and young sometimes tumble down into your fireplace in spring or summer, and cause commotion.

To me, the nesting habits of this bird seem

<sup>1</sup> *An-tros'to-mus vo-cif'er-us*. Length, about 9.50 inches.

<sup>2</sup> *Chac-tu'ra pe-lag'i-ca*. Length, 5 inches.

like faulty instinct. A chimney is a poor place of residence for a bird, and the habitants frequently come to grief. If the aperture is small, the householder objects to having the chimney stopped by nests; and if it is large, so many Swifts may nest there that their noise is an annoyance. These birds get up and out before daylight, to hunt insects that fly at night, and doubtless many a "ghost" in a "haunted-house" is nothing more frightful than a colony of these birds in the chimney.

This bird has the ability to fly straight up, or straight down, else it could not enter or leave a chimney. It is quite an aerial gymnast, and feeds only when on the wing. Its flight is very graceful, and both in manner of flight and personal appearance it so closely resembles a short-tailed swallow that there are few persons who can distinguish the difference in the flying birds.

One strongly marked peculiarity of this bird is that the tip of each tail-feather ends in a sharp, wire-like point, caused by the shaft of the feather being projected considerably beyond the vane. The eastern Chimney-Swift ranges westward to the Great Plains. On the Pacific slope is found another species, a close parallel to the preceding, called the *Vaux Swift*. The *White-Throated Swift* of the Pacific States is distinguished by its white throat and breast, and a few white patches elsewhere.

### THE HUMMING-BIRD FAMILY.

#### *Trochilidae.*

**The Ruby-Throated Humming-Bird<sup>3</sup>** represents the Family which contains the smallest of all birds. When the trumpet-vine on your veranda is in flower, you will see this delicate creature dart into view, like a large-winged insect, and poise itself easily and gracefully in mid-air at the mouth of the most conspicuous flower. Its tiny wings beat the air with such extreme rapidity and machine-like regularity that you see only a gray, fan-shaped blur on each side of the living bird. It holds itself in position with the greatest exactitude, thrusts its long and delicate beak into the heart of the flower, and, with the skill of a surgeon probing a wound, extracts the tiny insects or the honey so dear to its palate.

<sup>3</sup> *Troch'i-lus col'u-bris*. Length, 3.25 inches.

As the bird poises in mid-air, the sunlight catches the patch of brilliant ruby-red feathers on its throat, and sets it aflame. To make up for their diminutive size, and give them a fair share of beauty, Nature has clothed the throats and

mental tail-feathers, ruffs, and other showy decorations in feathers.

The Humming-Birds are so very diminutive one never ceases to wonder how such frail and delicate creatures, feeding only upon the smallest insects and the nectar of flowers, can make long journeys over this rough and dangerous earth, withstand storms, build their wonderful little nests, rear their young, and migrate southward again without being destroyed. Of course their diminutive size enables them to escape the attention of most of the living enemies which gladly would destroy them.

The nest of a Humming-Bird is about as large in diameter as a lady's watch, and the eggs, of which there are two, are the size of adult peas. The food of these birds generally consists of minute insects, many of which they find in large flowers. When at rest, perching, the average Hummer is not beautiful in form. Its head seems too large, its neck and body much too short, and its wings too long. It seems top-heavy, and as if destitute of legs. It is on the wing that these creatures look their best.

What Humming-Birds lack in size, they try to make up in number. There are nearly five hundred species, and they are found only in the New World. They are thoroughly tropical, but in warm weather, and the season of flowers, they migrate as far north as Alaska, and as far south as Patagonia. Our country makes an acceptable summer home for about sixteen species.

The Ruby-Throat is the only one inhabiting the eastern half of the United States, all the others being found west of Arkansas, and the Rocky Mountains.



RUBY-THROATED HUMMING-BIRD.

breasts of many Humming-Birds with feather-patches of the most brilliantly iridescent colors,—ruby-red, scarlet, green, blue and gold,—which flash like jewels. Others again have long, orna-

## CHAPTER XVIII

# THE ORDER OF WOODPECKERS

### PICI

The Woodpeckers are the natural protectors of the forests of the temperate zone. But for them, tree-borers would multiply without limit, and the number of trees that would fall before the insect pests is quite beyond computation. While the robin, the thrush and the warblers take care of the caterpillars and the leaf-insects generally, the woodpecker sticks to the business of his own guild, and looks after the pests that attack the bark and the wood. The tree-creepers assist by picking off insects from the outside, but when it comes to the heavy work of digging borers out of the bark by main strength, the woodpecker is the only bird equal to it.

There are about twenty-five species of woodpeckers in the United States.

Usually, the long, barbed tongue of this bird is sufficient to spear a borer, and drag it forth to meet the death it deserves. When this will not do the work, the woodpecker's claws take a good grip on the bark, and serious work begins.

Do not think, however, that because a rolling tattoo beaten on a hard dead limb can be heard a quarter of a mile, that the bird making the noise is working unusually hard. Quite the contrary. The loud tattoo is a signal, like the "certain whistle" of a small boy. In our Beaver Pond, the golden-winged woodpeckers sometimes beat on the galvanized-iron drums which protect the bases of the trees from the teeth of the beavers.

When a woodpecker is working hardest, you hear only a faint "chuck! chuck! chuck!" as he drives his sharp, wedge-like beak into the bark, or soft wood. Often the falling chips are your first notice that a winged forester is at work aloft, digging out and devouring the larvae that, if left alone, bring decay and death to trees.

You may be sure that whenever you find one of these valuable birds at work, there is need for him. To-day, a great many persons know their

value, and protect them. Occasionally, however, men who are so thoughtless or so mean as to engage in the brutal pastime known as a "side hunt," do lower themselves, and injure the land-owners about them, by killing every woodpecker that can be found,—for "points." If all farmers only knew what a loss every "side hunt" means to them, such wicked pastimes would not be tolerated.

Although the woodpeckers are not counted as birds of song, to me the loud, joyous cry of the flicker, the downy and the red-head, ringing through the leafy forest aisles, is genuine music. One species cries "*Cheer-up! Cheer-up!*" and it cheers-up and thrills me to hear it. Even in summer, when other birds are plentiful, it is a welcome sound. In bleak winter, when the great bulk of bird-life has vanished southward, and you toilsomely tread the silent forest, ankle-deep in snow, the world seems lifeless and drear—until you hear the clarion greeting of the golden-winged woodpecker. It is enough to stir the soul of a Digger Indian with a pleasing sense of companionship in life.

It is only the children of the cities who need to be told that woodpeckers have two toes in front and two behind, to enable them to cling to tree-bark; that the natural perch of such a bird is the perpendicular trunk of a tree; that sometimes they store acorns in holes which they dig in the sides of decayed trees, not in order that worms in those acorns may develop, but in order to eat the acorns themselves. They nest high up in hollow tree-trunks, which they enter through round holes of their own making.<sup>1</sup>

<sup>1</sup> Those who are specially interested in the habits of woodpeckers may profitably consult a report on "The Food of Woodpeckers" by Prof. F. E. L. Beal, published by the Department of Agriculture in 1895. The exact proportions of the various kinds of food consumed by seven species have been determined by examination of the stomachs of several hundred birds, and the figures quoted later on are from that report.

It is a good thing to feed wild birds of all species that are either useful or beautiful. The woodpeckers are the largest insectivorous birds that remain in the North over winter, and they appreciate friendly offerings of suet or fat pork, nailed high up on conspicuous tree-trunks. In the Zoological Park we put up every winter at least twenty-five two-pound strips of fat pork, for the woodpeckers and chickadees which live with us all the year round.

**The Golden-Winged Woodpecker<sup>1</sup>** is my favorite of the members of this Order. It is a bird of good size, dignified in bearing, decidedly handsome, and a great worker. He loves to hunt insects on the ground, occasionally, but is very alert and watchful, meanwhile. If you approach too near, he leaps into the air, and with a succession of wave-like sweeps upward and downward, his golden wings flash back one of his names as he flies to safety on some distant post or tree. Unlike most birds of this Order, this species frequently perches crosswise on a limb, like a true perching-bird.

This is the woodpecker of many names, some of which are *Flicker*, *High-Hole* and *Yellow-Hammer*. His regular call sounds like "Cheer up!" but in spring he gives forth a call which comes very near to being a song. When written out, it is like "Cook-cook-cook-cook!" At that season, also, you hear this bird beat the "long roll," on a drum which Nature provides for him in the shape of a hollow tree with a thin, hard shell. The rapidity and force with which the bird strikes the blows producing this sound are almost beyond belief.

An examination of the stomach contents of many specimens of this species showed 56 per cent of insect food, 39 vegetable, and 5 mineral. Of the insect food, ants made up 43 per cent and beetles 10 per cent. The vegetable food represented two kinds of grain (corn and buckwheat), eighteen kinds of wild berries, and fifteen kinds of seeds, mostly of weeds. Out of ninety-eight stomachs examined in September and October only four contained corn. Practically, this bird does no damage to man's crops, but destroys great quantities of harmful insects.

The range of the Golden-Wing embraces the eastern half of the United States to the Rocky

Mountains, where it is met by the *Red-Shafted Flicker* of the Pacific slope.

**The Red-Headed Woodpecker<sup>2</sup>** need not be described, because, in "Hiawatha," Longfellow has immortalized it. This bird, "with the crimson tuft of feathers," was the identical Mama which gave Hiawatha the timely tip which enabled him to put the finishing touch to old Megissogwon, and so end in triumph "the greatest battle that the sun had ever looked on."



GOLDEN-WINGED WOODPECKER.

As a return for this kindness, Hiawatha did the one mean act of his life. He took Mama's little red scalp, and "deckerd" his pipe-stem with it,—as coolly as if he had been a modern servant-girl decorating a forty-nine-cent hat.

This is a very showy bird, and recognizable almost as far as it can be seen,—brilliant crimson head and neck; white breast, sides and rump, and jet-black back and tail. In the Mississippi Valley, thirty years ago, this was one of the most common birds. Now, thanks to man's insatiable desire to "kill something" that is un-

<sup>1</sup> *Co-lap'tes au-ra'tus lu'te-us*. Length, about 12 inches.

<sup>2</sup> *Mel-an-er'pes e-ryth-ro-ceph'a-lus*. Length, 9½ inches.

protected, it has been so greatly reduced in number that it is seldom seen. It is an omnivorous feeder, eating insects, fruit, beech-nuts, corn and other grain, according to necessity. Its cry is loud and far-reaching, and sounds like "Choor! Choor!" As to migrating, it seems unable to make up its mind whether to become a "regular migrant" or a "winter resident."



RED-HEADED WOODPECKER.

Sometimes it migrates southward during the early winter, and sometimes it winters in the North.

An examination of the stomachs of one hundred and one Red-Headed Woodpeckers revealed 50 per cent of animal food and 45 per cent vegetable. Of the former, ants made up 11 per cent, and beetles 31 per cent. The fruit and vegetable food represented five kinds of

cultivated fruit (strawberries, blackberries, cherries, apples and pears), and fifteen kinds of wild fruit and seeds. The insect food consisted of ants, wasps, beetles, bugs, grasshoppers, crickets, moths, caterpillars, spiders and thousand-legged worms. In the fruit season, the Red-Head undoubtedly does considerable damage to fruit crops, more by mutilating fruit, perhaps, than by actual loss through fruit wholly consumed; and if these birds were as numerous as sparrows, it would be necessary for fruit-growers to take precautions against them during the fruit season. The damage done to corn appears to be quite insignificant. (Professor F. E. L. Beal's report.)

The great fondness of the Red-Head for beech-nuts, and its habit of storing them up for winter use, in holes and crevices, are well known.

**The Ant-Eating Woodpecker**<sup>1</sup> of the Pacific slope is the most conspicuous and interesting bird of this Order in that region, either around the suburban home, on the ranch, or in the mountain forests. This is the species which is now celebrated in word and picture for its habit of digging hundreds of holes in soft bark or dead tree-trunks, and "storing" an acorn in each hole, for future food.

**The Downy Woodpecker**<sup>2</sup> is a small gray-and-black species, modest and quiet in demeanor, but quite as common about the haunts of man as the golden-wing. It is the smallest species found in the United States and is the one which is most in evidence in winter.

This bird ranks high as a destroyer of insects, and in the percentage of insect food consumed leads all other woodpeckers that have been studied by the Biological Survey of the Department of Agriculture. An examination of one hundred and forty stomachs revealed 74 per cent of insect food and 25 of vegetable. The vegetable food consisted chiefly of seeds of the poison ivy, poison sumac, mullein, poke berries, dogwood and woodbine. The fruits consisted of service berries, strawberries and apples.

Apparently this bird is almost worth its weight in gold to the farmer who has valuable trees and fruit; and in winter, the farmer who is wise will put up suet, fat pork, and bones bearing some raw meat, on the trees in his orchard and woods.

<sup>1</sup> *Melanerpes formicivorus*.

<sup>2</sup> *Picus pubescens meridionalis*. Length, 7 inches.



**The Hairy Woodpecker**<sup>1</sup> is so close a counterpart of the downy, in appearance and habits, that it is unnecessary to describe both. The former is larger, but its rank as an insect exterminator is a little lower. Its proportion of insect food is 68 per cent, and vegetable, 31 per cent. Of the former, ants make up 17 per cent, beetles 24 per cent, and caterpillars 21 per cent. The only cultivated fruits found in eighty-two stomachs were blackberries; but wild fruits were well represented.

This bird inhabits practically the same region as the downy woodpecker, and belongs in the ranks of the farmer's best friends.

**The Yellow-Bellied Sapsucker**<sup>2</sup> is practically the only woodpecker which inflicts serious damage upon man's property; and possibly it may in some localities become so numerous as to require thinning out. Any bird which deliberately girdles a tree and kills it is a bird entitled to serious consideration, and punishment according to the harm it does.

This bird eats great quantities of insects, but as dessert it is fond of the sap of certain trees, among which are the maple, birch, white ash, apple, mountain-ash and spruce. Into the soft, green bark of these trees, this Sapsucker drills small, squarish holes, that look like gimlet holes. Usually they are placed in a horizontal line, and sometimes in mathematical groups. Occasionally several lines of these holes will quite girdle a tree. The bird not only drinks the sap that exudes, but he lies in wait to catch the winged insects and ants that are attracted to the sweet fluid, and devours great numbers of them.

Dr. C. Hart Merriam, who has closely observed the work of the Sapsucker, states that frequently mountain-ash trees are girdled to death by this bird, but that trees of greater endurance, like the apple and thorn-apple, are more able to survive its attacks. Another observer, Mr. Frank Bolles, declares that in well-wooded regions the damage it does is too insignificant to justify its destruction. Mrs. Mabel Osgood Wright states that in Connecticut "where these birds are plentiful, many orchard-owners cover the tree-trunks with fine wire netting."

"This species," says Professor Beal, "is probably the most migratory of all our woodpeckers,

<sup>1</sup> *Dry-o-ba'tes vil-lo'sus*. Length, 10.50 inches.

<sup>2</sup> *Sphy-ra-pi'cus va'ri-us*. Length, 8.25 inches.

breeding only in the most northerly parts of the United States, and in some of the mountains farther south. In the fall it ranges southward, spending the winter in most of the eastern states. It is less generally distributed than some of the other woodpeckers, being quite unknown in some sections, and very abundant in others."

In its general color-scheme, this is a bird of many and much-mixed colors—black, white and yellowish indescribably varied—both above and below. The top of the head and the throat



DOWNY WOODPECKER.

are bright red; and the sides of the head have two broad streaks of white, and two of black. The name of the bird is derived from the predominating greenish-yellow color of its breast and abdomen.

The Pacific coast has the *Red-Naped Sapsucker*, a subspecies of the above, of similar tree-girdling habits; the *Red-Breasted Sapsucker*, one of the commonest woodpeckers found from Oregon to Lower California, and two others,—the *Northern Red-Breasted* and *Williamson's*.

## CHAPTER XIX

# THE ORDER OF CUCKOOS AND KINGFISHERS

### *COCYGES*

This Order (pronounced Coc'-si-jēz) represents an effort to find a place for three familiar Families of birds whose members have something in common, yet in their most noticeable features are widely different. Both in their structure, habits and mode of life, the kingfisher and cuckoo are widely different from each other; and if there is one really good reason why these birds should be placed in the same Order, the writer would be pleased to have it pointed out. Their feet are totally different, and so are their beaks, their tails and their plumage. Any future revision of the classification of birds should strike this Order, early and hard.

#### THE CUCKOO FAMILY.

##### *Cuculidae.*

The **Yellow-Billed Cuckoo**,<sup>1</sup> or **Rain-Crow**, will fitly represent the Cuckoo Family. It looks like an insect-eating perching-bird, and



YELLOW-BILLED CUCKOO.

in reality it is one! You can easily recognize it by its extreme length and slenderness, the fan-like shape of its tail when spread, its upper surface of glossy drab—or gray-brown—and its white

under surface from throat to tail. To carry out this color-scheme to its logical sequence, the upper mandible is dusky brown, and the lower one is yellow.

This bird derives one of its common names—Rain-“Crow”—from the fact that its peculiar cry is heard oftenest on still and cloudy summer days,—two conditions which to the weather-wise farmer always portend rain. Its cry is a weird, gurgling note which sounds like “Cow-cowk-cowk-cowk!” and usually it comes from the heart of a thick bush or tree which effectually screens the bird. It seems to be fully aware of the dangers which beset all birds which attempt to live in the open with civilized man, for it lives amid the forest shadows.

This bird, and also its twin species, the **Black-Billed Cuckoo**, lives almost wholly upon insects. Of one hundred and fifty-five Cuckoo stomachs examined by the Department of Agriculture, only one contained any vegetable food—two small berries. Nearly half the Cuckoo's food proved to be caterpillars, 2,771 of which were found in 129 stomachs. It was not uncommon for one bird to contain more than 100 of them. “During May and June, when tent-caterpillars are defoliating the fruit-trees, these insects constitute half of the Cuckoo's food.”

The stomachs examined contained remains of sixty-five species of insects, in the following percentages: beetles, 6; bugs, 6½; grasshoppers, 30; caterpillars, 48½; other insects, such as web-worms, tussock-moths, army-worms, and moth larvae, 9.

From the results of this investigation it is clear that our two species of Cuckoo are to be numbered with the farmers' best friends among birds. As an estimate, I should say that each of these birds that enters a section devoted to farming and fruit-growing is worth to that section about \$10 per season. The charge that Cuckoos devour

<sup>1</sup> *Coccyzus americanus*. Length, about 12 inches.

the eggs or egg-shells of other birds was proven by the finding of shells "in several stomachs, but only in very small quantities—no more than was found in the stomachs of nearly every species that has been examined." Thus the offence charged proves to be too trivial to consider.

The Yellow-Billed Cuckoo inhabits the eastern half of the United States to the Great Plains, and the Black-Billed ranges westward to the Rocky Mountains, from Canada to the tropics. From the Rockies to the Pacific, and up to British Columbia, is found the *California Cuckoo*, a close counterpart of the Yellow-Billed species.

**The Road-Runner, or Chaparral Cock,**<sup>1</sup> is a very strange bird; and many strange "yarns" have been told of it. It is remarkably odd in form, and also in its habits. It is about the size of a small crow, with a tail as long as its entire body and head, and legs that are so long and strong they seem like those of a grouse, save that the toes are longer. The body is slender, but the neck and head are large, and the head has a conspicuous crest. The beak is large. Although this bird has wings, it seldom uses them, and they must be constantly growing smaller through disuse.

This strange bird is a habitant of the Southwest, from Texas to southern California and southward, and lives on the ground, in the low, dry brush which is called chaparral (shap-a-ra!'). It feeds upon every living thing inhabiting that region which it can catch and swallow,—mice, lizards, small snakes, centipedes and insects. It is one of the most nervous birds imaginable,—suspicious of everything that moves, and ready to make off without stopping to reason why.

It exhibits a decided preference for the smooth trails and paths through its beloved chaparral, and when alarmed it does not rise and fly, but makes off running, in the trail. It runs with great swiftness and seeming ease, but Dr. D. T. MacDougal has been informed that Mexican boys sometimes run them down, on foot, and either kill them with sticks or catch them alive.

This bird is also great at leaping, as we have seen in keeping it in captivity. Instead of flying to the top of a cedar-tree perch six feet high, and down again, it always leaps, with closed wings; but in leaping up it prefers to take a short


<sup>1</sup> *Ge-o-coc'cyx cal-i-for-ni-an'us*. Length, 21 to 23 inches.

run to acquire momentum. If this bird goes on ten thousand years in its present habits, by the end of that period its descendants probably will be without the power of flight, but provided with legs and feet so strong and full of spring that they can leap twenty feet.

## THE KINGFISHER FAMILY.

### *Alcedinidae.*

This family is widely and beautifully represented in the Malay Archipelago, but only three species are found in the United States. The **Belted Kingfisher**<sup>2</sup> is of almost universal distribution throughout North America, from the arctic Barren Grounds to Panama and the West Indies. Go where you will, in its season, where small fish abide, there will you find it. It is dignified, handsome, alert, and a true sportsman. Its favorite perch is a dead limb over



THE BELTED KINGFISHER.

still water, from which it can command a wide view, and swoop to the surface of the water in five seconds of time. You will know it by its bright blue upper surface; high and saucy crest; long, dagger-like beak; white under surface and broad belt of blue around the upper breast. Its cry is a metallic rattle, like "Churr-r-r-r-r-r!" and its food is small fish. It nests in a hole dug several feet horizontally into a perpendicular bank of earth, near water, or in a hollow tree.

<sup>2</sup> *Cer'y-le al-'cy-on*. Length, about 12 inches.

## CHAPTER XX

# THE ORDER OF PARROTS AND MACAWS

### PSITTACI

The parrots, parrakeets, macaws and cockatoos form a large group, containing in all more than 500 species. Of these, about 150 inhabit the New World, but only one species is found in the United States. South America contains the greatest number of species; Africa and Asia are but poorly supplied, and Europe has none. The widest departures from the standard types are found in New Zealand and Australia.



Drawn by EDMUND J. SAWYER.

CAROLINA PARRAKEET.

Although these birds are by nature thoroughly tropical, some of them range far into the temperate zones. This Order contains a larger proportion of beautifully colored birds than any other. Among the parrots, parrakeets, macaws and lorries, there is a lavish display of brilliant scarlet, crimson, blue, green, yellow and purple, while all save a few of the cockatoos are snowy white.

The members of this Order are specially distinguished by their bills and feet. Of the former, the lower mandible is a short but powerful gouge, while the upper mandible is a big hook, with a thick and heavy base, and a long, sharp point.

The foot of a bird of this Order is evenly divided, with the second and third toes pointing forward, and the first and fourth pointing back. The tails of most parrots are rather short, and square at the end, and the legs are very short. With but one or two exceptions, all the 500 species of this Order feed upon fruit, seeds and flowers.

**The Parrots** are celebrated by reason of the natural inclination of some species to mimicry, and their ability to learn to talk. They are naturally sedate and observant, possess excellent memories, and are fond of the companionship of man. The broad, fleshy tongue of a parrot renders possible the articulation of many vocal sounds, and when a certain phrase is endlessly repeated to a parrot that is secluded from other sounds, the bird is sometimes moved to remember and repeat them. The *African Gray Parrot* is the most celebrated talker, and its value is from \$15 upward. Next in rank comes the Mexican *Double Yellow-Head*, although the *Carthage Parrot*, being a good talker and a more hardy bird, is rapidly becoming more popular. Of both these species, the price in the New York bird-stores is from \$10 to \$12.

The parrot of the most remarkable habits is the **Kea**, of New Zealand, a bird with very large and strong feet, which not only loves fresh mutton, but sometimes kills sheep on its own account, for food purposes.

**The Parrakeets** are really small, trim-built parrots, with long, sharp-pointed tails. Excepting the Thick-Billed Parrot, which has been seen in southern Arizona, this Family contains the only member of the Order Psittaci which

inhabits the United States. The **Carolina Parakeet**<sup>1</sup> once ranged northward in summer to Maryland, Lake Erie and Iowa, and as far west as Colorado; but now all that is only so much history. The hand of the destroyer has been heavy upon this pretty bird. To-day it is found only in a few localities in Florida, and the prospects are that in a very few years it will be totally extinct. To illustrate: In 1893, a colony of about thirty birds which nested on the Sebastian River was completely destroyed in one night by a local hunter, who captured the entire flock, and sent the birds to a New York dealer, in whose hands all those which reached him alive died in a short time.

In color this bird has a bright green body, and yellow head and neck. It feeds upon fruit and seeds, and nests in hollow trees.

**The Macaws** are large, showy birds with very long, pointed tails, and the most awful voices for screeching ever made for feathered folk. They are found only in the New World from Mexico to Paraguay, and in the Andes up, to 10,000 feet. Either in flight, or at rest in the green tree-tops, they are exceedingly showy and attractive birds, and to find a flock in the depths of a tropical forest is an event to be remembered. In hunting macaws in the delta of the Orinoco, about every fourth bird that was

<sup>1</sup> *Co-nu'rus carolinensis*. Length, about 12 inches.

mortally wounded would hook its beak over a small branch, die, and hang there until I would be reluctantly compelled to make my fellow-collector, who was a good climber, climb up to the bird and throw it down, with much anger and unnecessary violence.

It is a pity that such beautiful birds should have such ear-splitting, nerve-racking voices. Although they seldom can be taught to talk, never cease to scream until dead, and are very apt to bite most unexpectedly, they are often kept as household pets.

**The Blue-and-Yellow Macaw**,<sup>2</sup> orange yellow below and cobalt blue above, is one of the species most frequently seen in captivity. In the bird-stores of New York, they sell at from \$10 to \$15 each. The *Red-and-Blue Macaw* is another common species. The beautiful plum-colored bird occasionally seen is the *Hyacinthine Macaw*, from Brazil.

**The Cockatoos** are mostly—but not all—snow-white birds, with lofty and beautiful triangular crests which can be erected at will, with striking effect. They inhabit Australia, Celebes, the Philippines and the southern islands of the Malay Archipelago. They are easily tamed, talk readily, take kindly to training, and become very affectionate and satisfactory companions.

<sup>2</sup> *Ar'a ar-a-rau'na*. Length, about 30 inches, of which the tail constitutes about 18 inches.

## CHAPTER XXI

# THE ORDER OF BIRDS OF PREY

### RAPTORES

To every farmer and poultry-raiser, the birds of this Order are divided into two groups, friends and enemies. Inasmuch as feathered friends are to be encouraged, and all enemies slain, the standing of each species becomes a life-or-death matter. America is a wide and populous country, and despite the labors of the Biological Survey of the Department of Agriculture, there are yet millions of persons who desire precise information regarding our hawks and owls. Because of the economic importance of the subject, we will devote a liberal amount of space and effort to the important members of this group. The Families of the Order are as follows:

	FAMILIES.	EXAMPLES.
ORDER RAPTORES.	BARN-OWLS, . . . <i>STRIGIDAE</i> , . . .	Barn, or Monkey-Faced Owls.
	HORNED OWLS, <i>BUBONIDAE</i> , . . .	Horned, Burrowing, Snowy and Screech-Owls.
	HAWKS, . . . . <i>FALCONIDAE</i> , . . .	Hawks, Kites, Buzzards and Eagles.
	VULTURES, . . . . <i>CATHARTIDAE</i> , . . .	California, Turkey and Black Vultures.

### THE BARN-OWL FAMILY.

#### *Strigidae.*

It is now a well-established fact that "owls are among the most beneficial of all birds," inflicting little damage upon the producers of poultry, and conferring vast benefits upon the farmer by the destruction of mammal and insect pests. Inasmuch as their regular working hours are from sunset to sunrise, they wage successful war on the nocturnal mammals which remain quiet during the daytime in order to escape hawks and other daylight enemies.

Owls are exceedingly interesting birds, and in them there is also much to admire. They take life seriously; they have but few nerves, and seldom use them. Rarely do they become really tame or affectionate, but easily become very indignant at real or fancied affronts. Like many people of few words and solemn manner, they are not nearly so wise as they look. They are easily caught in steel traps, or shot; and they are much given to nesting in situations that are wide open to attack.

Omitting the subspecies,—which are only geographic races,—there are eighteen species of owls in North America, north of Mexico. They vary in size from the tiny elf-owl, of Arizona, only 6 inches in total length, to the great gray owl, of the arctic regions, 30 inches long.

With the exception of the great horned owl, the owls of our country are by no means so destructive to poultry and wild bird life as is generally supposed. The great majority of the species feed upon wild mice, rats, squirrels, shrews, fish, crustaceans and insects; and some of them render great service to man. Nearly all owls are night-flyers, and by reason of their soft, fluffy plumage, which renders their flight quite noiseless, they are specially fitted to keep in check the grand army of destructive rodents that roam abroad under cover of darkness.

Owls do very well in captivity, provided they are properly housed and fed, and have comfortable perches to sit upon. Naturally, they are most active at night, and quiet in the daytime. Be it known, however, that they cannot live long on a steady diet of beefsteak. Every

owl must have a liberal allowance of small birds, like English sparrows, and, if possible, an occasional small mammal, in each case with the feathers or hair upon it. Nature has constructed the owl to devour its prey entire,—feathers, hair, bones and all, on the spot where it is captured.

By a curious rotary action of the stomach, all the desirable elements are extracted and assimilated, and the indigestible refuse—hair, feathers, bones, claws, etc.—is rolled into a ball called a “pellet,” which is cast up, and expelled through the mouth. These pellets are sometimes collected at roosting-places, and when carefully examined by expert zoologists, it is possible to identify most of the animal remains, and tell what the bird has fed upon.

**The Barn-Owl, or Monkey-Faced Owl,**<sup>1</sup> is the most oddly shaped of all the owls; it has the smoothest and most compact plumage, and proportionately the longest legs. Its general color is that of scorched linen—light brownish-yellow. Each small black eye is the centre of a sunburst of radiating feathers, and the whole face is surrounded by a heart-shaped ring of brown.


The Barn-Owl is to rats and mice as the cuckoo is to the caterpillar. As a destroyer of the meanest vermin on earth (rats) this bird has no equal. Whether North or South, in the tropics or the temperate zone, it loves to live under the roofs of civilized man, especially in church bell-fries, where it is not molested. In the town of Barrancas, at the head of the Orinoco delta, some Venezuelan boys piloted me into the best church in the place, showed me two Barn-Owls nesting over the altar, and urged me to shoot them then and there. My refusal because the birds were very thoroughly in sanctuary was with difficulty comprehended.

Many observations on the food habits of this bird have been made by examining the pellets that have been gathered from its roosting place. In June, 1890, Dr. A. K. Fisher collected 200 pellets that had accumulated from two birds that roosted and nested in one of the towers of the Smithsonian building. These contained 454 skulls, of which 225 were of meadow-mice, 2 of pine-mice, 179 of house-mice, 20 were of

<sup>1</sup>*Strix pra-tin'co-la*. Length, from 15 to 17 inches.

rats, 6 of jumping mice, 20 shrews, 1 star-nosed mole and 1 vesper-sparrow.


The Barn-Owl rarely molests birds—probably never does so except when forced by hunger—and all over the world, wherever it is found, its favorite food is rats and mice. The number an industrious pair will destroy in a year is really very great, and this species deserves the most



SKELETON OF A BIRD OF PREY. (BALD EAGLE.)

- |                        |                      |
|------------------------|----------------------|
| 1, Upper mandible.     | 11, Radius,          |
| 2, Lower mandible,     | 12, Carpals,         |
| 3, Hyoid,              | 13, Metacarpals,     |
| 4, External nostril,   | 14, Digits,          |
| 5, Orbit,              | 15, Sternum,         |
| 6, Occiput,            | 16, Keel of sternum, |
| 7, Cervical vertebrae, | 17, Pelvis,          |
| 8, Clavicles,          | 18, Fibula,          |
| 9, Coracoid,           | 19, Tibia,           |
| 10, Ulna,              | 20, Tarsus,          |
|                        | 21, Digits of foot.  |

careful protection that man can give it. Fortunately, it and its subspecies are very widely distributed,—more cosmopolitan, in fact, than any other owl, save the short-eared.



N. Y. Zoological Park.  
BARN-OWL.

### THE HORNED-OWL FAMILY.

#### *Bubonidae.*

The Long-Eared Owl<sup>1</sup> looks like a small and imperfect imitation of the great horned owl. It can always be distinguished by its small size, and the fact that its horns appear to have been set too close together on the top of its head, and do not fit very well. Its total length is about 15 inches, and its general color is a fine mottling of gray, tawny and black, which produces a brownish-gray bird. It is found all over the United States.

The food of this very useful bird consists mainly of mice. In April, 1888, at Munson Hill, Virginia, Dr. Fisher collected about 50 pellets from under a tree in which a Long-Eared Owl had roosted, and found that they contained the following remains: 95 meadow-mice, 19 pine-mice, 15 house-mice, 5 white-footed mice, 3 Cooper's mice, 26 short-tailed shrews and 13 birds. Of the birds, there were 11 sparrows, 1 blue-bird and 1 warbler. Of this species Dr.

<sup>1</sup>*A'si-o wil-son-i-an'us.*

Fisher says: "It is both cruel and pernicious to molest a bird so valuable and innocent as the one under consideration."

The Short-Eared Owl<sup>2</sup> is of about the same size as the preceding species, but its ears are so short that they look like two small feathers that have been thrust carelessly into the plumage directly above the eyes. Above it is a brownish-yellow bird, and buffy white underneath. It is found from the arctic regions of North America to Patagonia, and throughout nearly the whole of the Old World except Australasia. Its food habits are very similar to those of the long-eared owl, and it is equally deserving of a perpetual close season.

The Barred Owl<sup>3</sup> has not quite so good a reputation as the three noticed above, but its record is by no means bad. Out of 109 stomachs examined by the Biological Survey, three contained domestic fowls, one a ruffed grouse and one a pigeon. Six contained screech-owls, one a saw-whet owl, three held sparrows, one a woodpecker, and two small birds were not identified. Against this debit was a credit of 46 mice, 18 other small mammals, 4 frogs, 1 lizard, 2 fishes, 2 spiders, 9 crawfish and 20 empties. The eighteen small mammals consisted of 5 red



Photo. and copyright, 1902, by W. L. UNDERWOOD.

#### BARRED OWLS.

squirrels, 1 flying squirrel, 1 chipmunk, 4 rabbits, 2 shrews, 2 moles, 1 weasel and 2 rats.

From this very exact evidence, the reader

<sup>2</sup>*A'si-o ac-cip-i-tri'nus.* Length, from 14 to 16 inches.

<sup>3</sup>*Syr'ni-um va'ri-um.*



can judge of the value or lack of value of this bird to the country at large. It does not seem as if the forty-six mice are a fair equivalent for the useful birds and small mammals destroyed.

Dr. Fisher's conclusion is as follows: "If a fair balance be struck, it must be considered that this Owl is on the whole beneficial, and hence should occupy a place in the list of birds to be protected."

The Barred Owl is next in size to the great horned owl. It is from 20 to 22 inches long, heavy-bodied, round-headed, and quite without "horns," or "ears." Its head, neck and breast are marked by many black horizontal bars on a gray or creamy-white ground, and the breast and abdomen have a few thick, perpendicular bars. Many times a big Barred Owl of my acquaintance has exclaimed to me through the darkness, in a fearfully hollow and sepulchral voice,—“Whò? Whó-whó-whó-whó-whó-whó? Ah!” It is like the war-cry of an angry ghost.

This bird ranges throughout the eastern half of the United States, and westward almost to the Rocky Mountains; and it frequently finds its way into captivity.

The **Great Gray Owl**<sup>1</sup> is the largest member of this Family found in the New World. It is an arctic bird, one-fourth larger than the great horned owl, and even in winter has never wandered farther south than the Ohio River. In Alaska and British Columbia it inhabits the timbered regions, and does not wander far into the treeless Barren Grounds. Anyone who captures a very large owl of a dusky brown or dusky gray color, larger than a great horned owl, but with *no ear-tufts*, may know that he has secured a specimen of the rare and handsome Great Gray Owl.

The **Saw-Whet Owl**<sup>2</sup> is a very small Owl, and so shy that few people ever see it; but it feeds almost *exclusively upon mice*, and any bird which wages perpetual war on those pests deserves honorable mention in these pages. In appearance it looks very much like a small gray-phase screech-owl without ears. It may be looked for—but it will seldom be found—almost anywhere in the United States from the international boundary to the Gulf States and California.

<sup>1</sup> *Scio-ti-ap'tex neb-u-lo'sa*. Length, 25 to 30 inches.

<sup>2</sup> *Nyc'ta-la a-ca'di-ca*. Length, 8 inches.

The **Screech-Owl**<sup>3</sup>—with an awful shiver in its voice, but *no screech* whatever—is so widely distributed, and so easily affected by climatic variations, that the original species has been split up into eight varieties, or subspecies. Thus we now have the Texas, California, Rocky Mountain, Mexican, and Florida Screech-Owls, and others too numerous to mention. The differences between all these are not very great. Let each American know his own Screech-Owl, and study its habits, and he will then know the others, quite well enough for all practical purposes.

To me, the cry of this little Owl is one of the most doleful sounds in animated nature, not even excepting the howl of a wolf. It is like the



N. Y. Zoological Park.

SCREECH-OWL.

quivering, shivering, heart-broken wail of a lost spirit, and suggests chattering teeth and freezing vocal chords. Written out it is “*Woe-woe-woe-woe-woe-woe-woe ah!*” But no phonetic spelling can even suggest the high-pitched mental and physical anguish expressed in the cry that one hears.

The Screech-Owl is a round-bodied little fellow, sometimes almost as broad as it is high; and its head is surmounted at the corners by very respectable ears. In its gray phase, this bird looks very much like a dwarf great horned

<sup>3</sup> *Meg'as-cops a'si-o*. Length, 7 to 9 inches.

owl; but of course the black markings are not the same.

This Owl exhibits a peculiarity in color which must be specially noted. It has two distinct and widely different colors, red and gray. In the same locality will be found owls that are of a cold, black-and-white gray color, and others that are pale, rusty red, with white mottlings on the abdomen. For this very odd development, we are quite unable to account; and such



SANBORN, Photo., N. Y. Zoological Park.

#### YOUNG SCREECH-OWLS.

lawless color-variations are called "phases," possibly because they phase the naturalists who try to study out their whys and wherefores.

In its food habits, the Screech-Owl prefers, if it can procure them, mice, grasshoppers, locusts, cut worms, beetles, caterpillars, crickets, spiders, lizards, frogs and crawfish. If these are lacking, it attacks the English sparrow and almost any other small bird that comes handy, usually other sparrows. To show that when very hungry all birds look alike to him, he occasionally kills and eats a bird of his own species! Dr. A. K. Fisher's ever useful and scholarly report on the "Hawks and Owls of the United States" sets forth in full detail the results of the examination of 255 stomachs

of Screech-Owls, of which the following is a summary of contents: 100, insects; 91, mice; 12, English sparrows; 26, other birds; 11, miscellaneous mammals; 9, crawfish; 7, miscellaneous food; 5, spiders; 5, frogs; 2, lizards; 2, scorpions; 2, earth-worms; 1, poultry; 1, fish, and 43 were empty. The following is a full list of the birds found: 12 English sparrows, 9 other sparrows, 3 juncos, 2 Screech-Owls, 1 shore-lark, 1 water thrush and 15 unrecognized.

Leaving out the two Screech-Owls, of the birds that were identified, the English sparrows formed practically one-half. On this basis we will allow that of the unrecognized birds, seven were song-birds. Add these to the fifteen recognized-song birds and we have a total of twenty-one song-birds out of two hundred and fifty-five stomachs examined.

The question is, what shall be the fate of the Screech-Owl,—encouragement, toleration, or limitation? To me it seems that the number of Screech-Owls should be *limited*, for the benefit of the song-birds; but I do *not* believe in their extermination.

**The Great Horned Owl**<sup>1</sup> is, by necessity, an aerial pirate and highway robber—the tiger of the air. Its temper is fierce and intractable, and if you attempt to make friends with one in captivity, it will hiss like a snake, snap its beak like an angry peccary, and dare you to come on. Of all the birds I know, there is no other so persistently savage in captivity as this bloody-minded game-killer. Of course, the Owl is not to blame for the raw-meat appetite which Nature gave him, and for which he feels bound to provide; but there is no reason why he should have a temper like a black leopard toward those who feed him.

"Of all the birds of prey, with the exception possibly of the goshawk and Cooper's hawk," says Dr. A. K. Fisher, "the Great Horned Owl is the most destructive to poultry. All kinds of poultry seem to be taken, though when Guinea-fowls and turkeys are obtainable, it shows a preference for these. In sections of the country where it is common, the inhabitants complain bitterly of its ravages." In the museum of the Philadelphia Academy is an Owl which carried off from one farm twenty-seven individuals of various kinds of poultry before it was shot.

<sup>1</sup> *Bu'bo virginianus*. Length, from 20 to 24 inches.



GREAT HORNED OWL.

With "horns" laid back in anger.

But let us give even the Horned Owl its just due. Mr. O. E. Niles, of Ohio, once found in a nest of this bird "several full-grown Norway rats with their skulls opened and brains removed," and on the ground under the tree which contained the nest he found "the bodies of *one hundred and thirteen rats*, most of them full grown!" Now, in the course of a year, would not one hundred and thirteen Norway rats consume and destroy enough grain to feed one hundred and ten head of poultry?

This is the summary of the contents of 127 stomachs of Great Horned Owls examined by the Biological Survey: 31 contained poultry or game-birds; 8 contained other birds; 13 contained mice; 65 contained other mammals; 1 contained a fish; 1 contained a scorpion; 10 contained insects, and 17 contained nothing.

The bird-food represented the following: 21 domestic birds, 11 song-birds, 3 ruffed grouse,

2 quail, 1 pinnated grouse, 1 pigeon, 1 rail, 1 wild duck, 1 Cooper's hawk, and 2 unknown.

The mammals found were as follows: 46 mice and rats, 32 rabbits and hares, 7 shrews, 5 squirrels, 3 chipmunks, 4 pocket-gophers, 2 skunks, 1 weasel and 1 bat.

Beyond question, the debit balance against this bird is heavy, and justifies its destruction, wherever found; but at the same time, it goes against the grain to kill a bird which destroys so many rats.

The Great Horned Owl, or Hoot-Owl, as it is frequently called, is a bird of dignified and imposing appearance. Its big, round-topped horns of feathers are singularly like cats' ears in shape, and when with these are seen the fiercely-glaring eyes of yellow and black, the half-yellow face and fluffy white feathers on the throat, the whole head of this bird is singularly like that of a Bengal tiger. The body plumage is a complex mottling and barring of black and brown, dull yellow and white, impossible to describe successfully.

But this bird can always be recognized by its large size, cat's-ear "horns," and the fine, black horizontal bars across its breast-feathers. From wing to wing, across its upper breast there is an assemblage of heavy splashes of black.

The eastern Great Horned Owl is the type species on which are based the *Western*, *Arctic*, *Dusky* and *Pacific* Horned Owls, which in com-



Photographed by E. R. WARREN.

YOUNG GREAT HORNED OWLS.

ination cover practically the whole of North America down to Costa Rica. By reason of the live food available in winter, these birds are not migratory.

**The Snowy Owl**<sup>1</sup> is a bird of the Arctic wastes, and reaches the northern United States only as a winter visitor. Its occurrence with us varies from a total scarcity during some years to an abundance during others. During December, 1886,—the beginning of the awful winter which killed over 90 per cent of the range cattle in Montana,—we saw in the country in which we



Photo. by C. WILLIAM BEEBE, N. Y. Zoological Park.

SNOWY OWL.

were hunting buffalo, in central Montana, at least twenty-five Snowy Owls. They were living on hares, rabbits, and sage-grouse, out in the open, twenty miles from the nearest timber. It was their habit to alight upon the tops of the low buttes, in reality upon the ground, from which they could survey a wide circle of sagebrush plains. Whenever there is an annual "flight" of Snowy Owls, they are always particularly numerous in Minnesota.

But for its perfectly round and rather comical-looking head, this bird would be the most beautiful of all American owls. Its plumage varies

<sup>1</sup> *Nyc'tea nyc'tea*. Average length, about 23 inches, the female being larger than the male.

from almost spotless snow-white, in some individuals, to white barred all over with narrow horizontal bands of black—which is really the standard color-plan. The number and width of the black bands vary exceedingly in different individuals, some birds being rendered much darker than others.

The food of this species consists of every kind of wild bird or small mammal it can catch; but there is no evidence that it ever destroys poultry. In summer, when its far-northern home is full of migratory birds, nesting and rearing their young, its bill of fare is quite varied, but in winter it is confined to such winter residents as the ptarmigan, hare, rabbit, sage-grouse, and such small rodents as dare to venture forth from their burrows.

With the **Burrowing Owl**<sup>2</sup> of the western plains, the Owl Family may justly be regarded as "run to earth." This odd little owl indeed takes shelter in the mouths of prairie-"dog" holes, but as far as I am aware there is no proof that it ever descends to the bottom of a deep burrow, or that it is chummy with the rattlesnake. It is reasonably certain that no owl in its right mind ever would fraternize with a rattlesnake, and neither would a prairie-"dog."

The Burrowing Owl lives in the plains of the West and Southwest, from North Dakota to southern California. A closely related species is found in Florida, where it easily digs burrows in the sandy soil.

Many persons have the idea that this Owl is unable to dig, and is therefore dependent upon prairie-"dogs" and badgers for a home. This is entirely erroneous. In soil that is reasonably loose, the Burrowing Owl is a most industrious and successful digger, and with his feet flings out the loose dirt and gravel in a shower. A pair of western birds which we kept in the Bird-House of the New York Zoological Park for two years burrowed so deeply into the big pile of solid gravel in their enclosure that its interior became a perfect cavern.

In the land of plains and prairie-"dogs," the Burrowing Owl is a frequent corollary to a "dog" town, sitting on the highest point of a burrow mound, or, if alarmed, taking short flights to the suburbs. Between bird and rodent there ap-

<sup>2</sup> *Spe-ot'i-to cu-nic-u-la'ri-a hy-po'ga-a*. Average length, about 10 inches.

pears to exist a *modus vivendi*, which is good so long as the bird does not come within reach of the legitimate owner of the soil. As already mentioned (page 77), when the two are intimately mixed, the prairie-"dog" quickly kills the Burrowing Owl. It seems practically certain that the bird inhabits only the mouth of the prairie-"dog's" burrow, or burrows that have been abandoned.

This owl is far too small to kill even a half-grown "dog;" besides which, its favorite diet is grasshoppers, locusts, other insects, lizards and scorpions. It is to be noticed that in thirty-two stomachs examined in Washington, one really did contain a portion of a prairie-"dog," and two contained one mouse each, but thirty-three contained insects only, some of them showing from forty-nine to sixty each of locusts and grasshoppers.

The color of a Burrowing Owl is a grayish mixture, darkest on the back, and lighter below, and the legs are long and naked, like those of a sharp-shinned hawk. In captivity our specimens dug extensive burrows for themselves, in doing which they threw out gravel and earth with astonishing force. They are savage little wretches, and murder each other at a shocking rate. The males fight savagely, and the western species will not live peaceably with that of Florida.

## THE HAWKS AND EAGLES.

### *Falconidæ.*

This section of the Order Raptores contains a remarkable assemblage of forms, and the wide differences between some of the groups add zest to the study of them. Some are expert in fishing, some are of dignified and imposing bearing, some have beauty of plumage, and one is the most beautiful flyer in all the bird-world. Until only ten years ago, most people regarded all hawks as so many robbers, deserving death.

In 1893, the investigations of the Department of Agriculture revealed the surprising fact that of all the forty-one species of day-flying birds of prey in North America, there were only four species whose destructiveness so far outweigh their useful services that they deserve to be destroyed. The others are either harmless to man's interests, or else so positively beneficial that they deserve careful protection. Beyond

doubt, the careful and thorough investigations made by the Biological Survey, under the direction of Dr. C. Hart Merriam, and the publication of the results, have resulted in the correction of popular errors which if persisted in would have caused enormous losses to the farmers of the United States.

As an object lesson, take the case of Pennsylvania.

In 1885, the legislature of that State enacted a law aimed at the wholesale destruction of hawks and owls, and authorizing the various counties to pay cash bounties for the "scalps" of those birds, at the rate of fifty cents each. Immediately the work of slaughter began. Many thousand scalps of hawks and owls were brought in, and over \$90,000 were paid out for them. It has been estimated that the "saving" to the agricultural interests of the state amounted to \$1 for every \$1,205 paid out as bounties! In this manner the balance of Nature was quickly and completely destroyed.

The awakening came even more swiftly than anyone expected. By the end of two years from the passage of the very injudicious "hawk law," the farmers found their field-crops and orchards so completely overrun by destructive mice, rats and insects, they appealed to the legislature for the quick repeal of the law. This was brought about with all possible haste. It was estimated by competent judges that the "hawk law" cost the farmers and fruit-growers of Pennsylvania not less than \$2,000,000 in actual losses on valuable crops.

The moral of this episode is,—*it is always dangerous, and often calamitous, to disturb violently the balance of Nature, either by the destruction of existing species of birds or mammals, or by the introduction of new ones.*

**The American Osprey, or Fish-Hawk,<sup>1</sup>** is, by common consent, regarded as a sort of connecting link between the Owl and Falcon Families. It is a good bird to lead a large Family, and it is to be regretted that those who dwell far from the sea-coast and large rivers lack opportunities for becoming well acquainted with it. Surely this bold fisher, who thinks nothing of dropping a hundred feet into ice-cold water, seizing a fish of nearly half his own weight, and

<sup>1</sup> *Pan'di-on hal-i-ae-e'tus carolinensis*. Average length, about 24 inches; weight, 3 pounds.

flying five miles with it, must appeal to every man and boy who loves the grasp of a good rod, and the musical click of a reel.

The boat trip up the Shrewsbury River, from New York to Long Branch, is worth taking in midsummer solely for the sight of the Ospreys, winging slowly over the still lagoon, stalking their finny prey, and anon plunging with a loud



AMERICAN OSPREY.

splash into the water. Sometimes the bold fishers go quite out of sight. The most surprising thing about such performances is the size of the fish that an Osprey can lift and carry away.

In carrying a fish, an Osprey always grasps it on the back, with one talon well ahead of the other, and the head of the fish pointing straight forward. This is to secure a minimum of resistance from the air, and render it an easy matter to steer the prize to the home-nest, or to a tall tree on which it may be devoured at leisure. It is no wonder that a three-pound Osprey carrying a one-pound fish is moved to jettison his cargo when he sees a hostile bald eagle bearing down upon him with empty claws and his decks cleared for action.

The story of the Ospreys of Gardiner's Island is a most interesting chapter in bird-life. The owner of that island is a relentless enemy to cats and gunners, and a fierce protector of all the wild life on the island, which is wholly his. His weapons are loaded for hunters only, and for several years the Ospreys have bred regularly around Mr. Gardiner's house, and all over the island. A pair of birds occupies the same nest year after year, adding to the mass each year, until the nest contains a wagon-load of sticks of many sizes, and measures six feet in diameter. To-day, strange to relate, some of the Ospreys are nesting practically upon the ground, serenely confident of their security from all harm.

The Osprey is built like a light-weight athlete, all bone, tendon, hard muscle and wing-power, and no fat. Its long, half-naked legs and powerful claws remind one of patent grappling-hooks. The wings are long and acutely pointed, going well beyond the end of the tail. The whole neck and lower surface of the bird is white, but the back, wings, and upper surface of the tail is dark colored, as also is the upper half of the head. The plumage is compact, smooth and oily, as befits a diving-bird.

In summer this bird is at home on the sea-coast from Alaska and Hudson Bay to the Gulf of Mexico, and along a few rivers, but in winter it migrates to southern Florida, the West Indies and northern South America.

The jaunty little **Sparrow-Hawk**<sup>1</sup> is the smallest American hawk, and also the most beautiful. Its form is elegant, and its colors are varied and pleasing. As if desirous of admiration, it tolerates man at shorter range than any other hawk I know. Its cap is dull blue, its throat white with black side-patches, and its upper neck and back are bright rusty brown. Its breast is salmon color, sparingly spotted, its knickerbockers are white, and its tarsi and feet are bright yellow. It inhabits the whole United States, and on northward to Great Slave Lake, but I think it is most plentiful on the prairie farms of the middle West.

As a destroyer of grasshoppers, beetles, crickets, caterpillars and other insect enemies, this little Hawk deserves to rank with the birds most beneficial to man. For so small a bird, the

<sup>1</sup> *Falco spar-ve'ri-us*. Length, 9 to 10 inches.

number of grasshoppers it consumes in a year is enormous. It never molests poultry, and when insects are obtainable never kills a song-bird, but it does destroy great numbers of mice. Dr. Fisher reports that of 320 stomachs examined, 215 contained insects; 29, spiders; 89, mice; 12, other mammals; 53, small birds; 1 game-bird, and 29 were empty. Many stomachs contained from 10 to 35 grasshoppers each, and of other insects, from 25 to 40 in one bird was of common occurrence.

It must be noted at this point that when the Sparrow-Hawk is rearing its young, it does sometimes catch young chickens; but the extreme infrequency of this may be judged from the fact that in the entire series of 320 specimens examined at Washington, taken at all seasons from January to December, and throughout a wide range of localities, not one stomach contained any remains of a domestic bird. In the early spring, before grasshoppers come, Sparrow-Hawks often follow a plough very closely, to capture the mice that are ploughed up. Sometimes this bird is half domestic in its habits, and nests in buildings erected by man. Wherever it is found, it should be a welcome visitor.

The Pigeon-Hawk<sup>1</sup> is a slightly larger bird than the preceding, very destructive to song-birds, of little use to man, and deserves to be shot wherever found. It kills sparrows, thrushes, goldfinches, vireos, bobolinks, swifts, and a host of other species. Out of 56 specimens examined by Dr. Fisher, 41 contained small song-birds, and 2 poultry; 2 only had mice, and 16 insects. This is a bird of plain colors, being bluish-gray or brownish above, and lighter below.

Apparently the Duck Hawk,<sup>2</sup> a geographic race of the Peregrine Falcon, never devours a mouse or an insect save by mistake. Out of 20 specimens, 7 contained game-birds or poultry, 9 had eaten song-birds, only 2 contained insects, and 1 a mouse. You may know this bird by the great size and strength of his "pickers and stealers." It can best be studied with a rope, a basket, and a chokebore shot-gun loaded with No. 6 shot.

First shoot both male and female birds, then

<sup>1</sup> *Fal'co col-um-ba'ri-us*. Length of male, about 10 inches; female, 2 to 3 inches more.

<sup>2</sup> *Fal'co per-e-gri'nus an-a'tum*. Length of male, 17 inches; female, 19 inches.

collect the nest, and the eggs or young, whichever may be present. In doing this, however, be careful not to shoot the *Red-Tailed* or *Red-Shouldered Hawk*,—both good friends of ours, who are entitled to protection. A Duck-Hawk has *no red nor decided brown* upon it, anywhere. In general effect it is a dull black bird with a white breast and throat, and white abdomen cross-barred with black. It inhabits all of America north of Chili.

The time was when the **Bald Eagle**,<sup>3</sup> or **White-Headed Eagle**, was known to every human being within the limits of the United States. Today there are probably two million men in this country, speaking foreign languages only, but voting regularly and persistently, who do



SPARROW-HAWK.

not know an Eagle from a parrot, nor the number of stripes there are in Old Glory. It is related by a reliable eye-witness that when an escaped parrot recently perched in one of the trees of City Hall Square, New York City, a dispute

<sup>3</sup> *Hal-i-ae-e'tus leu-co-ceph'a-lus*. Average length of male, about 34 inches; female, 38 inches; spread of wings, from 7 to 8 feet. See plate on page 170.

as to its identity was ended satisfactorily by some who oracularly pronounced it an "eagle bird."

But, no matter how many persons there are in this country who do not know our national bird, I will not humiliate "Old Baldy" by formally introducing him. To every intelligent American, the perfect bird, with its snow-white head, neck and tail, is recognizable at a distance of a mile or more. To see one perching on the topmost branch of a dead tree, overlooking a water prospect, with its snowy head shining in the sunlight like frosted silver, is enough to thrill any beholder. Even when in flight, an eagle can be distinguished from all other birds by its slow and powerful wing-strokes, and the great breadth of its wings, especially near their extremities.

It is unfortunate that this Eagle does not acquire its white head and tail until its fourth year. The head is fully feathered, and the name "Bald" refers solely to its white appearance. Up to three years of age it is of the same general color as the golden eagle, and to distinguish the two species it is necessary to look at the lowest joint (tarsus) of the leg. If it is *naked*, the bird is a Bald Eagle; but if it is covered with feathers quite down to the toes, it is a golden eagle.

As a rule—to which there are numerous exceptions—the White-Headed Eagle is found along rivers, and the shores of lakes and ponds containing fish. Fish are its favorite food, and lambs are purely supplementary. As a regular thing, it catches fish out of the water, with neatness and despatch; but when it sees an osprey flying by with a large fish in its talons, the Eagle does not hesitate to levy tribute on the subject bird. Taken thus at a great disadvantage, the fish-hawk has no option but to drop its fish, and go away to catch another, while the Eagle catches the prize before it touches the water and bears it away.

This act of the Eagle, and the extra trouble it puts upon the fish-hawk in catching duplicate fish, is by a few writers taken seriously to heart. So is the additional fact that Eagles—like many human beings—often eat dead fish that are found floating upon the water, or are cast up on the shore. For these, and other reasons equally weighty (!), it has become almost a fashion among writers to denounce the Bald

Eagle, and declare it a shame that such a bird ever was chosen as our national standard-bearer. Some have asserted that the brave and high-minded wild turkey would have been more appropriate!

Against all of this, I have nothing to say. The American Eagle needs no defence from me. Whether

"He clasps the crag with hooked hands,  
Close to the sun in lonely lands,"

or perches defiantly on the United States coat-of-arms, with a brow to threaten or command, he is beloved by at least seventy-two million people who will rise as one whenever he is really in need of defenders. Abroad, it once was well-nigh an international fashion to flout this bird, and the standard he bears; but since May 1, 1900, that fashion has gone out. Abroad, those who do not respect this bird fear him, wholesomely. At home, it is quite time for all strangers to secure an introduction to him, and for some of those who should be his friends but are not, to write him down no longer.

In its distribution, this Eagle ranges over the whole of North America from Mexico to Kamchatka. Considering the size of this bird, it holds its own remarkably well, even in New England. In Florida it is very abundant all along Indian River, and in one locality in the State of Washington it is so numerous that its depredations on the flocks of sheep-raisers are cause for serious complaint and reprisals.

In the East so many Eagles are caught alive and offered for sale that it is a difficult matter to find sale for one at \$10. This bird so seldom destroys domestic animals, or game-birds, there is no excuse for its destruction save possibly in a few far-western localities where it happens to be very numerous, and evinces a particular fondness for lambs.

About every six months there appears in some newspaper an account of a child having been attacked by a fierce Eagle, and rescued by a heroic mother, or else actually carried off to the top of a tall tree or rocky cliff, from which the child was finally rescued unhurt, etc., etc. It is quite time that this absurd yarn, which is nearly as old as the Swiss Alps in which it originated, were consigned to the oblivion it deserves. Eagles know what guns are, and nothing is



farther from their thoughts than attacking the children of Man, the destroyer of life.

**The Golden Eagle**<sup>1</sup> is in no sense whatever a golden-colored bird. Its plumage is dark brown, with a very slight outside wash of lighter brown. It would be much more appropriate to call it the "brown eagle." In appearance it looks very much like a white-headed eagle in its second year, except that its tarsi are *feathered quite down to the toes*. By this point it can always be distinguished from its nearest relative.

This bird has a very bad record as a destroyer of lambs, poultry, game-birds, young deer, antelope, rabbits, and other small mammals. It cares very little for fish, and prefers to frequent interior regions, where either domestic animals or wild species of good size are abundant. By preference it is a bird of the mountains, and although found all the way from the Atlantic to the Pacific, and from Mexico to the Arctic Ocean, it is most abundant in the great mountain-ranges of the West. In the cattle country east of the Rockies, many a Golden Eagle dies ignominiously from eating poisoned meat that is intended for wolves.

**The Hawks** of North America above Mexico form a group of about thirty-four species, not counting subspecies, and the conspicuous types are well worth serious attention.<sup>2</sup> Some of them are useful to man, and some are so destructive and generally useless that they deserve death. It is highly important that hawk enemies should be distinguishable from hawk friends.

**The Red-Tailed Hawk**<sup>3</sup> is the greatest of all destroyers of noxious four-footed animals. It might well be called the *Mammal-Eater*, instead of being universally miscalled the **Hen-Hawk**, or **Chicken-Hawk**.

The species of the above name inhabits the entire eastern half of the United States, and ranges westward to the Rocky Mountains, where it meets the subspecies known as the *Western*

<sup>1</sup> *A-quil'a chrysa-a-ē'tos*. Size, about the same as the white-headed eagle.

<sup>2</sup> To avoid the possibility of confusion, attention is called to the fact that the sparrow-hawk, pigeon-hawk and duck-hawk, already described, belong to *Falco*, the genus of the falcons, a group quite distinct from those of the hawks now to be introduced.

<sup>3</sup> *Bu'teo bo-re-al'is*. Average length of male, about 21 inches; female, 24 inches.

*Red-Tail*. By reason of the abundance of this bird, and its undoubted influence for good or evil upon agricultural communities, the Department of Agriculture has made a study of it which was particularly thorough. From Arizona to Connecticut, and in all seasons of the year, collections were made, until finally 562 stomachs had been collected and examined.

The result was a complete vindication of the moral character of the previously despised and persecuted "Hen-Hawk." *Two hundred and seventy-eight specimens contained mice; 131, other mammals; 54, poultry or game-birds; 51, other birds; 47, insects; 37, amphibians and reptiles; 13, offal; 8, crawfish, and 89 were empty.* It was found that poultry and game did not constitute 10 per cent of the food of this Hawk, and that all other beneficial creatures preyed upon, including snakes, did not increase this proportion to 15 per cent. Against this small debit stands a credit of 85 per cent, made up chiefly of destructive rodents.

"It is not to be denied," says Dr. Fisher, "that a good deal of poultry is destroyed by this Hawk; but the damage done is usually among the less vigorous fowls, in the late fall, and in view of the great number of injurious rodents as well as other noxious animals which this Hawk destroys, *it should seem equivalent to a misdemeanor to kill one, except in the act of carrying off poultry.* The fact that there are robbers among Hawks is no sound argument for exterminating any and every one."

This bird is very omnivorous in its habits. In the examination noted above, the remains of 35 species of small mammals were found, of which 30 were rodents, 5 were insectivores and 1 (a common skunk!) was a carnivore. Of birds there were only 20 species.

The important markings of the Red-Tailed Hawk are its rusty-brown tail, back and head of blackish-brown, white throat, and light-colored breast streaked with dusky or brown. *The immature bird has a gray tail, crossed by from 6 to 10 dark bands, and the rusty-red tone of the adult bird is everywhere absent.* The head is large, and rather square in outline at the back.

There are varieties of this bird scattered all over the United States, and under most circumstances it is rather difficult to tell them apart.

The **Red-Shouldered Hawk**<sup>1</sup> has not only "red" shoulders, but also a red head, neck, back and breast. But there are many shades of red, and the so-called red on this bird is as widely different from the red of a cardinal as blue is from green. The so-called "red" on this Hawk is really a rusty brown; and by the great amount of it, the small, round head of the bird, and its *black* tail crossed by about six bands of white, this species may easily be distinguished from the preceding.



SHARP-SHINNED HAWK.

This Hawk is to be counted with the farmer's best friends. Mr. J. Alden Loring knew a pair which for two years nested within fifty rods of a poultry-farm on which were about 800 young chickens and 400 ducks, but never attempted to catch one. Mice constitute two-thirds of its food, but it is very fond of frogs and toads. In the 220 specimens which he examined, Dr. Fisher found the remains of creatures representing eleven classes of life. The food exhibit was made up as follows: 3 stomachs contained domestic fowls; 12, other birds; 102, mice; 40, other small mammals (16 species in all); 20, reptiles; 3, fish; 39, amphibians (frogs and toads);

<sup>1</sup> *Buteo lineatus*. Average length of male, 18 inches; female, 20 inches.

92, insects; 16, spiders; 7, crawfish, and 1, earth-worms.

The service rendered by the Red-Shouldered Hawk consists chiefly in the destruction of mice and grasshoppers; and birds of all kinds are touched very lightly. This species inhabits eastern North America from Nova Scotia and Canada to the Gulf, and westward to the Plains. The Pacific coast contains a variety known as the *Red-Bellied Hawk*, which is quite as honest about poultry as the eastern species.

The **Sharp-Shinned Hawk**<sup>2</sup> is a swift flyer, a keen hunter, and a great murderer of small birds. Like all the hawks, its upper surface is dark, and its lower surface light. Its tail is long, and has three or four narrow, dark-colored bands across it, far apart, with the widest band nearest to the end. The wings, back, upper neck surface and upper tail are all bluish-gray. The throat and under parts of the body are white, plentifully cross-banded with rusty brown.

This is a small hawk,—next in size to the pigeon-hawk. Its beak seems rather small and weak, but its legs are long and its feet large, and these, backed up by swift flight and great courage and impudence, render this bird a winged terror. It hunts along fences like a dog hunting rabbits, and pursues song-birds into their thickets and out again. Its principal food is song-birds, and only at long intervals does it capture a mouse. This bird is rather too small to handle poultry with complete success.

The complete list of the bird-remains found in 159 stomachs of Sharp-Shinned Hawks constitutes a tale of slaughtered innocents that is appalling. Six stomachs contained poultry, and 99 contained *song-birds*, woodpeckers and a few others. Only six contained mice, and 5, insects; and 52 were empty. Of the wild birds, 56 species were identified. There can be no question regarding the necessity for the destruction of this bird, wherever it is found. It breeds throughout the entire United States, northward to the arctic circle, and southward to Guatemala. (Fisher.) In some localities it is quite abundant.

**Cooper's Hawk**<sup>3</sup> is a companion in crime to the preceding species, and equally deserving an

<sup>2</sup> *Accipiter velox*. Average length of male, 10.50 inches; female, 13 inches.

<sup>3</sup> *Accipiter cooperii*. Average length of male, 15.50 inches; female, 19 inches.

early and violent death. By a strange coincidence, it bears a strong resemblance to the sharp-shinned hawk, both in form and color, but it is a much larger bird. Leaving size out of consideration, it is difficult to describe in words the slight differences that exist between the two.

Being a bird of strong and rapid flight, much strength and activity, and great boldness, it is well equipped for raiding poultry-yards, and carrying off almost anything except geese and turkeys. Of 133 stomachs examined by Dr. Fisher, 34 contained poultry or game-birds; 52, other birds; 11, mammals; 1, a frog; 3, lizards; 2, insects, and 39 were empty. The game-birds found were 1 ruffed grouse, 8 quails and 5 pigeons. Altogether, 21 species of useful birds had been eaten, but only 4 mice, 1 rat and 1 grasshopper.

No record could be much blacker than this, and Cooper's Hawk is a pest whose career deserves to be ended by three drachms of powder and an ounce and a half of No. 6 shot, whenever opportunity offers. If gunners could *only discriminate*, the killing off of this species would make great sport for them; but the trouble is, many innocent birds would be killed by mistake.

This bird inhabits the whole United States, but stops at the Canadian boundary, and goes south to southern Mexico.

**The American Goshawk<sup>1</sup>** is to Canada and Alaska what Cooper's hawk is to the United States,—a wholesale destroyer of game-birds, serving no useful purpose whatever. To the unprotected flocks of ptarmigan it is a genuine scourge, and it merits destruction. Fortunately this hawk visits the United States only in winter, and even then is by no means numerous. Those who have had opportunities to observe it in action consider it the boldest and most audacious hawk in America. It has been known to seize a freshly killed chicken from the side of the farmer who had slain it for dinner, and also to follow a hen into a house, and seize it in the presence of its owner. (Fisher.)

The length of the Goshawk is from 21 to 25 inches. The top of its head is black, and its upper surface is bluish-slate color. Its whole under surface is white, with many gray cross-bars, in addition to which it is lined up and down with short, black lines, rather far apart. The lower tail surface is crossed by four gray bands.

<sup>1</sup>*Accipiter atricapillus*.

**The Marsh-Hawk<sup>2</sup>** is essentially a prairie-hawk; and in the open and fertile uplands of the Mississippi valley, it is one of the most conspicuous species. It loves farming regions wherein members of the Mouse Family are plentiful and cheap. In hunting it flies low, in a very business-like way, just above the grain or tall grass, and its intentions are so apparent that the American farmer gave it credit for its good work, years before the true value of the once-despised "Hen-Hawk" became known.

This hawk is not beautiful, either in form, color or movement. To me it always seems to have too much sail area for the size of its hull. Its adult color is drab, or bluish-gray, but the



COOPER'S HAWK.

females and immature males are rusty brown; much like the red-shouldered hawk. However, this hawk can always be distinguished by the *large white patch on the rump*, just above the tail.

One of the first facts about the nesting of hawks that comes to a Western farmer boy by personal

<sup>2</sup>*Circus hudsonius*. Average length, about 22 inches.

observation is that the Marsh-Hawk nests on the ground, preferably in tall grass, in a nest that is anything but a workmanlike affair. When I found my first nest of this bird,—a patch of trampled grass in the head of a slough, with four big, downy nestlings wallowing around upon it,—the Marsh-Hawk fell several points in my estimation.

This species ranges all the way from Alaska, Hudson's Bay and Ontario to Panama and Cuba. Regarding its value, Dr. Fisher has this to say:

"The Marsh-Hawk is unquestionably one of the most beneficial as it is one of our most abundant hawks, and its presence and increase should be encouraged in every way possible, not only

try or game-birds; 7, reptiles; 2, frogs; 1, unknown and 8 were empty.

**The Swallow-Tailed Kite,**<sup>1</sup> or, as the boys of the prairies call it, the **Forked-Tailed "Hawk,"** is in flight the most graceful bird I ever saw on the wing. No matter whether the sky be blue or gray, the snow-white head, neck and body, and glossy black tail and wings are sharply outlined in the heavens, drawing attention as a magnet draws nails. The bird is instantly identified by its long and deeply V-shaped tail, and its striking colors, which divide evenly between themselves the under surface of the wing.

In the golden days of boyhood, I saw scores of these birds in Iowa, but never saw one alight and perch, even for a moment. Several times we saw them with snakes in their talons, devouring them as they sailed through the air, and we also saw two or three seizures of prey. But it is the flight of this bird that makes the most lasting impression. In hunting and prospecting it never flies in a straight line, but always in graceful curves, and reverse curves, circles, parabolas, and spirals, like an expert skater "showing off." Its flight is indeed the poetry of motion in mid-air.

Unfortunately, this beautiful bird is not of wide distribution in the North, for its real home is in the tropics. In the United States it migrates northward in April into Iowa, Minnesota, Illinois, southern Michigan, and at rare intervals farther east and west to the Carolinas and the plains. So far as known, its food consists exclusively of small reptiles and large insects.

This bird fitly represents the whole group of Kites, of which the *White-Tailed Kite* is the Pacific coast species. The *Mississippi Kite* inhabits the Gulf states, and the *Everglade Kite* reaches our country only in Florida.

## THE VULTURE FAMILY.

### *Cathartidæ.*

This Family ranks at the bottom of the list of the birds of prey, because its members are less intelligent, less active and resourceful in obtaining their food, and less able to take care of themselves than the hawks and owls. Although

<sup>1</sup> *El-a-noi'des for-fi-ca'tus*. Average length, about 23 inches.



SWALLOW-TAILED KITE.

by protecting it by law, but by disseminating a knowledge of the benefits it confers. It is probably the most active and determined foe of meadow-mice and ground-squirrels, destroying greater numbers of these pests than any other species, and this fact alone should entitle it to protection, even if it destroyed no other injurious animals."

One hundred and twenty-four specimens of this species were examined, and the stomachs revealed the following contents. 57, mice; 27, other mammals; 34, birds; 14, insects; 7, poul-

not so highly developed as the hawks, the vultures serve a most useful purpose in the economy of Nature, and exhibit some traits that are really wonderful. The broad-minded student will not turn from these birds with aversion merely because their heads are bare, and they feed on dead

seen the **Common Turkey Vulture**<sup>1</sup> sailing and circling on wide-spread but motionless pinions, so high in the heavens that its distance from the earth seemed to be two miles or more.

Clearly, these aerial promenades, often continued until the observer is weary of watching



Photographed by E. F. KELLER, National Zoological Park.

CALIFORNIA VULTURE.

food. Their heads are naked for professional reasons.

Two things about vultures are particularly striking. One is the enormous heights to which they soar, the other is their marvellous quickness in discovering the body of a dead animal. Many times, in clear summer weather, I have

them, are taken for pleasure. One great circle succeeds another in a series that seems unending, but all the while the wings are as motionless as if wired in position. On such occasions, even a homely and unlovely Buzzard can become

<sup>1</sup> *Ca-thar'tes au'ra*. Average length, about 29 inches.

an object of admiration, and a reminder of William Tell's Alpine eagle, which—for sentimental reasons, only—he "could not shoot."

"His broad, expanded wings  
Lay calm and motionless upon the air,  
As if he floated there without their aid,  
By the sole act of his unlorded will,  
That buoyed him proudly up."

The flight of the Vulture, by which it gains enormous heights without any serious exertion after getting well clear of the earth, is an interesting illustration of what a perfect arçodrome might accomplish if it could flap its wings for a lofty rise, sail with abundant wing-power, and be intelligently guided. Beyond doubt, the bird keeps aloft by properly utilizing the lifting power of air-currents.

By a strange coincidence, the bird which flies highest and longest, and soars most majestically, is also the bird of lowest tastes on the earth. Although it has strong talons and a strong beak, it kills nothing, and feeds upon dead animals. In every country on earth, vultures are treated as highly useful creatures. In the tropics, where their services really are of great value, they are fully protected by law.

The species found farthest north, with a bright-red head and neck, is the Turkey Vulture, and it ranges across the continent from the plains of the Saskatchewan to Patagonia.

**The Black Vulture,**<sup>1</sup> marked by a head and plumage which is perfectly black, is seldom seen in the northern portions of the United States, but is abundant in the Gulf states, and southward far down into South America. In appearance this bird is most funereal. It is a smaller bird than the turkey vulture, but does not fly so well, and flaps its wings oftener. Around the cities of the South it is a great domestic economist and labor-saver.

In Bombay, India, the Parsees expose their dead in two great, shallow, open-topped towers, called the Towers of Silence, and the vultures regularly devour them,—all except the bones, which fall down into a central pit.

**The California Vulture, or California "Condor,"**<sup>2</sup> is, among naturalists, the most

<sup>1</sup> *Cath-ar-is'ta ur'u-bu*. Average length, about 25 inches.

<sup>2</sup> *Gym'no-gyps californianus*.

celebrated bird of this Family, partly because it is our largest bird of prey, and also because of its great rarity. The "collectors" are certain to exterminate it in a very few years. Its appearance depends upon its attitude. With its wings spread, it is a grand bird; but with them closed its personality is far less impressive. On the wing, in the wild, rocky fastnesses of its native mountains, those who have seen it there say it is a grand and imposing object, and it is not to be wondered at that its pursuit is quite as exciting as the chase of the big-horn.



E. F. KELLER, Photo., National Zoological Park.

YOUNG CALIFORNIA VULTURE.

Mrs. Florence Merriam Bailey<sup>3</sup> gives the following as the dimensions of this bird: "Length, 44 to 55 inches; wing-spread, 8½ to nearly 11 feet; weight, 20 to 25 pounds. Distribution: coast ranges of southern California from Monterey Bay south to Lower California, and east to Arizona."

This great Vulture breeds in the most inaccessible crags it can find, but of course collectors find it. In 1894, Mr. Stephens actually encountered a flock of twenty-six of these magnificent

<sup>3</sup> "Handbook of Birds of the Western United States," p. 144.



Photo. by E. R. SANBORN, N. Y. Zoological Park.

THE CONDOR.

birds. For three years, a very fine specimen has lived in the National Zoological Park, at Washington, shut up at night in an elevated sleeping-box. In the morning when liberated in its enclosure, it perches aloft, spreads its wings and holds them out to catch the sun's rays, in true vulture fashion.

Largest of all the Birds of Prey is the **Condor**<sup>1</sup> of the Andes, a bird of lofty home but lowly habits. In the Andes of Chili and Peru, its range is from 9,000 to 16,000 feet above the sea, and it not only feeds upon dead guanacos and vicunias, horses and other domestic animals, but it also ventures to attack living calves and old horses

<sup>1</sup>*Sar-co-rham'pus gry'phus*. Length of male, 48 inches; spread of wings, 8½ to 9½ feet.

that are almost incapable of defence. Condors are so easily captured alive that the zoological gardens of the world are always well stocked with them.

By nature the Condor is a peace-loving bird, and for two years visitors to the New York Zoological Park have witnessed the strange spectacle of the world's largest bird of prey—the fine adult male shown in the accompanying plate—living in the great Flying Cage in peace and harmony with about eighty flamingoes, herons, egrets, ibises, ducks, other water-birds and various land-birds. Encouraged by the success of the Condor experiment, a large griffon vulture has been added to the “happy family,” with very satisfactory results.



## CHAPTER XXII

# THE ORDER OF PIGEONS AND DOVES

### COLUMBAE

**The Passenger Pigeon**<sup>1</sup> was until very recently only a bird of history; and, until 1899, it was regarded as a species practically extinct. The men who lived in the Mississippi Valley forty years ago remember the flocks that flew swiftly over the farms, sometimes fifty and sometimes two hundred or more birds together. It was a wonderful sight to see the perfect mechanical precision with which they kept together, wheeling and circling in as perfect formation as the slats of a Venetian blind.

This very rare bird is much larger than a dove. Its color is bluish above, and reddish-brown underneath, and the feathers of its neck have a rich metallic lustre. Its tail is *long and pointed*, and its feet and legs are red. It never was found in the far West, and never will be. The pigeon of the Pacific coast is a totally different species.

In the early days, Ohio seemed to be the centre of abundance of this bird, and the accounts that have been written of that period relate how the Pigeons sat so thickly upon the trees that branches were broken by their weight; how they covered the earth when they alighted in the fields to feed, and darkened the sky when they flew.

As usual, that great abundance of wild life

provoked great slaughter. Migrating Pigeons were killed by wholesale methods. While breeding they were attacked in their nesting-places, and in an incredibly short time the great flocks vanished. As in the case of the blotting out of the great northern buffalo-herd, in 1884, many persons have wondered, and do still, whether the great flocks of Pigeons have not migrated, and found a permanent home elsewhere. There is not a single fact on which to base either belief or supposition that the Passenger Pigeon exists abundantly in Mexico, Central America or elsewhere.

Among naturalists, the blotting out of this interesting species has been a source of sincere regret. As usual, no one thought of protecting it until it was entirely too late. But it seems as if we are to be given another opportunity to count this bird in our avifauna. Beginning about 1891, a few small flocks began to appear in the United States, first four or five birds together, and then larger flocks. Mr. George O. Shields, Editor of *Recreation Magazine*, has carefully sought out and published the details of every Pigeon occurrence that came to his knowledge. Up to January, 1901, the following observations of the occurrence of Passenger Pigeons were reported in the magazine mentioned above:

Wisconsin, Milton.....	1891 to '99.....	Several annually.....	<i>Recreation</i> , Sept., 1899.
Canada, Ft. QuAppelle.....	July, 1898.....	A flock.....	“ Dec., 1898.
Illinois, Edinburg.....	“.....	3 birds.....	“ May, 1899.
Kentucky, Caldwell Co.....	Oct., “.....	30 “.....	“ Feb., “
Michigan, Ann Arbor.....	Oct., “.....	200 “.....	“ Apr., “
Wisconsin, Lime Ridge.....	April, 1899.....	100 “.....	“ July, “
Indiana, Sullivan.....	May, “.....	25 “.....	“ “ “
Ohio, Litchfield.....	April, “.....	150 “.....	“ June, “
Wisconsin, Amherst.....	“ “.....	100 “.....	“ Aug., “
Illinois, Chadwick.....	Oct., “.....	50 “.....	“ Sept., “
Wisconsin, Milwaukee.....	“ “.....	17 “.....	“ “ “
“ Norway.....	May, “.....	200 “.....	“ “ “
Manitoba, Southern.....	“.....	A few.....	“ Jan., 1900.
New York, Willsville.....	Sept., “.....	10 “.....	“ “ “
Canada, Three Rivers.....	Dec., 1899.....	1 bird.....	“ “ “
New York, Willowemoc.....	Nov., “.....	200 birds.....	“ Feb., “
Minnesota, Dumont.....	July, “.....	45 “.....	“ “ “
Michigan, Lowell.....	1900.....	40 “.....	“ Dec., “

<sup>1</sup> *Ec-to-pis'tes mi-gra-to'ri-us*. Average length, about 16 inches.

The latest and most gratifying information on this subject is contained in a letter dated Nov. 9, 1903, transmitted by Mr. Shields. In Pennsylvania, in a locality that shall be nameless here, three flocks of Passenger Pigeons, containing in all about 300 birds, have been feeding for three weeks on the farm of a sportsman and nature-lover who is protecting them.



THE BAND-TAILED PIGEON.

So it seems that our old friend is striving to ignore the black record of the past, and come back to us, to live and breed. Wherever it elects to be seen, or to breed, it should be accorded the most thorough protection, both by public sentiment, and by law.

The **Band-Tailed Pigeon**,<sup>1</sup> of the Pacific states from British Columbia to Guatemala, and eastward to the Rocky Mountains, yet exists in fair abundance, and it is earnestly hoped that it never will be annihilated without reason or mercy, as was the sad fate of its eastern rela-

<sup>1</sup> *Co-lum'ba jas-ci-a'ta*. Average length, 15 inches.

tive. Wherever found it should be accorded legal protection, without delay.

This fine bird is conspicuously marked by a white collar around its neck, and a *square-ended tail which terminates with a dull-white band from one to two inches wide*. The head and under parts are purplish-pink, fading downward to a lighter color. The back is brownish-gray, fading out toward the tail into a dull-blue tone.


This Pigeon subsists upon acorns, seeds and berries, and attracts attention to itself by its noisy flight. Its strange vocal utterances are graphically described by Mrs. Florence Merriam Bailey:

"If you follow the pigeons to their breeding-grounds in some remote canyon you will be struck by the owl-like hooting that fills the place, and you will locate the sound here and there along the sides of the canyon at dead tree-tops, in each of which a solitary male is sunning himself, at intervals puffing out his breast and hooting. The hooting varies considerably. Sometimes it is a calm *whoo'-hoo-hoo, whoo'-hoo-hoo*, at others a spirited *hoop-ah-who'*, and again a two-syllabled *whoo'-ugh*, made up of a short, hard hoot and a long coo, as if the breath was sharply expelled for the first note and drawn in for the second." ("Handbook," p. 139.)

To me the **Mourning-Dove**<sup>2</sup> has always seemed like a sacred bird; and although I could have killed thousands, I never took the life of one. When a very small boy at my mother's knee, she related to me the story of the winged messenger which Noah sent out of the ark, over the waste of waters, to look for real estate. She told me that Doves were innocent and harmless little birds, and that I must never harm one in the least. Had my good mother issued an injunction covering the whole animal kingdom, I think I would have grown up as harmless to animals as any Hindoo; for her solemn charge regarding Doves has always seemed as binding as one of the ten commandments.

I mention this in order to point out to mothers the far-reaching extent of their power in behalf of our wild creatures, and the vast influence which they can easily wield in behalf of birds and mammals in sore need of protection. Is it not a good thing to teach all boys that it is mor-

<sup>2</sup> *Ze-na-i-du'ra ma-crou'ra*. Average length, 12 inches.



MOURNING-DOVE.

ally wrong (*which it is!*) to kill wild creatures without reason, mercy and common-sense?

The Mourning-Dove received its "given" name from the mournful sound of its call-notes. Its sad-voiced "*Coo, coo, coo,*" suggests moaning, and, next to the awful, storm-beaten wail of the screech-owl, it is, under certain conditions, the most doleful sound uttered by an American bird. I knew one sensitive woman who was so affected by the daily "mourning" of a neighboring Dove that she begged a sportsman to frighten it away.

Another peculiar fact about this bird is the strange musical note that is sounded by the vibration of its wings. As the bird springs from the ground in flight, or wings its way overhead, the pulsations of its wings give forth a ringing, metallic sound, like the twanging of a tight wire.

This Dove loves country roads, more than any other bird, and to those who love beautiful things, its exquisitely moulded form and immaculate plumage is always a pleasing touch of Nature. One might as well try to describe in words the colors in a fire opal as those of this bird. There is pink iridescence, and brownish, and grayish, and blackish, and other shades too numerous to mention, but the combination baffles description.

This Dove breeds throughout the United States from the international boundary to the Gulf, and migrates as far south as Panama. In California it is now counted as a "game-bird," and killed by sportsmen, and in the South also it is killed by the negroes for food. A great "game-bird" this, truly! A genuine sportsman must be very hard pressed for gun victims when he can seriously call this tamest of all birds "game." And can any farmer in his senses afford the expense of having Doves shot on his farm, or in his neighborhood? Let us see.

When the Biological Survey of the Depart-

ment of Agriculture took up the case of the Mourning-Dove, and examined the stomachs of 237 specimens, the summary of results proved that as a weed-destroyer this bird is one of the most valuable in North America. Weed-seeds constitute 64 per cent of its food, all the year round, with little monthly variations. In order to arrive at an exact determination, the seeds in three stomachs were carefully identified and counted. One contained the following:

Orange hawkweed ( <i>Hieracium aurantiacum</i> ).....	4,820 seeds.
Slender paspalum ( <i>Paspalum setaceum</i> ).....	2,600 "
Hoary vervain ( <i>Verbena stricta</i> )...	950 "
Panicum.....	620 "
Carolina cranesbill ( <i>Geranium carolinianum</i> ).....	120 "
Yellow wood-sorrel ( <i>Oxalis stricta</i> )..	50 "
Miscellaneous weeds.....	40 "
	9,200

The second specimen of the three contained 6,400 seeds of the farmers' ancient and persistent enemy, fox-tail (*Chactocloa*), while the third turned out 7,500 seeds of the yellow wood-sorrel. The grand total of weed-seeds for those three Doves was 23,100! And this for only one day's supply. Assuming that those three Doves had been killed as "game" by some "sportsman (!)," previous to their meal, and those seeds had produced 23,100 weeds, how much would it have cost in labor at \$1.50 per day to destroy them?

Besides the 64 per cent of weed-seeds in the 237 stomachs, there was found 32 per cent of grain, but of this three-fourths was waste grain, gleaned in the fields after harvest.

Whoever does aught for the protection of Doves, does well; and a word to the wise is sufficient.

CHAPTER XXIII

THE ORDER OF UPLAND GAME-BIRDS

GALLINAE

It is natural that a country possessing the wide diversity of uplands that exists in the United States should possess a great variety of ground-dwelling birds. In response to the inviting fields and forests, plains and mountains,—cold and warm, wet and dry,—the birds of the Order Gallinae have greatly multiplied, both in number and in species.

It is no wonder that men and boys like to hunt upland game-birds; and when the conditions are properly observed, it is right that they should do so. The natural death of a game-bird or quadruped is by shot or bullet, from the gun of a true hunter, who hunts only at the proper time, in a fair manner, and kills sparingly. Wherever game-birds are most plentiful, each hunter is in honor bound to kill only a small number, and give others a chance.

If you are a boy, or man, don't be a "game-hog!" Shoot like a gentleman, or don't shoot

at all. If any species becomes so rare that it is threatened with extinction, *stop killing it*, and take measures for its complete protection until it has had time to recover. Above all, *never engage in a "side-hunt,"* which is a wholesale slaughter of wild creatures "for points," and never tolerate one in your neighborhood. Side-hunting should be prevented, at the muzzle of breech-loaders, if necessary.


Some of the most interesting hunting experiences ever recorded have been in hunting game-birds with the camera. If space were available, it would be a pleasure to record here the names of some of those who have made beautiful pictures of ruffed grouse, pinnated grouse, woodcock, ptarmigan and many other species. A fine bird photograph is a joy forever, but a bagful of dead birds disappears in an hour.

The table below affords a bird's-eye view of this Order as it exists north of Mexico:

	FAMILIES.	GROUPS.	EXAMPLES.
ORDER GALLINAE, North of Mexico.	GROUSE FAMILY: . . . <i>TET-RA-ON'I-DAE.</i>	Quail <sup>1</sup> and Partridges: . . .	{ Virginia "Quail," or Bob-White. California Mountain "Quail." California Valley "Quail." Mearns' Partridge. Scaled Partridge.
		Grouse: . . . . .	{ Ruffed Grouse. Canada Grouse. Pinnated Grouse. Sharp-Tailed Grouse. Sage-Grouse.
	PHEASANT FAMILY: . . . <i>PHAS-I-AN'I-DAE.</i>	Ptarmigan: . . . . .	Willow Ptarmigan.
		Turkeys: . . . . .	Wild-Turkey.
	Pheasants: . . . . .	{ Ring-Necked Pheasant. ant. Golden Pheasant. Silver Pheasant. }	Introduced.
	All of the Old World only.		

<sup>1</sup> By technical writers the name "quail" is now considered as applying only to the members of a group of Old-World birds, much smaller than our quails; and our quails are called "partridges," because they are related to the Old-World birds of that name. But this is only another "buffalo"

case. The good old name "quail" is so universally known that no power on earth could supplant it, and in a work of this kind it would be folly to ignore it in favor of "partridge," even though the latter be more correct.



BOB-WHITE.

As the preceding diagram shows, there are no true pheasants in America save those that have been introduced from China and Japan. All the birds to which that name correctly applies inhabit the Old World.

### THE GROUSE FAMILY.

#### *Tetraonidae.*

Our dear old friend the **Common "Quail"** is now called **Bob-White**<sup>1</sup> in all the modern bird-books, but to about fifty million Americans it is yet, and ever will be, *the* Quail. It is our longest-known and most widely known American game-bird, and it is almost wholly a United States bird. It is at home from Maine and Florida to Texas, the western border of Oklahoma and South Dakota. In very many eastern localities, however, it has been almost exterminated by excessive shooting, and during the past ten years, Mr. Charles Payne, of Wichita, Kansas, has caught and shipped east fully *two million* live quail for use in restocking quailless game-


<sup>1</sup> *Co-livus virginianus*. Average length, 10 inches.

preserves. The extermination of desirable species always costs money.

The call of this bird is one of the most cheerful sounds in nature, and for carrying qualities it is far-reaching. From the heart of a hazel thicket one hears his loud, shrill whistle, saying "CLERK-it! CLERK-it! CLERK-it!" until everything rings again. On the hurricane deck of a high stump, or the top rail of a fence, he poises himself, points his bill at the sky, swells out his white throat and whistles long and loud, "Bob! BOB! WHI-EET!" But the feathered rascal knows very well when the close season is on; and when the "law is off" he sings very small.

That many men enjoy Quail shooting is no cause for wonder or reproach. The birds lie close in the edge of the brush until the dogs are ready to tread upon them, when "*Burr-r-r-r!*" the covey explodes in the air like a bomb, the gray and brown fragments fly in half a dozen directions, and the young sportsman is so "rattled" he is almost sure to miss. A well-scared Quail is no easy mark.

Quail are rapid breeders, and in protected localities they increase rapidly. A good bird-law in Kansas has resulted in bringing back the vanished flocks, to a surprising extent. Unfortunately, however, it is not possible to breed Quail in large numbers in confinement, even with a quarter section of land for the experiment.



CALIFORNIA MOUNTAIN PARTRIDGE.

The flesh of this bird is a great table delicacy, — provided it has not been kept in cold storage. A cold-storage Quail is as good to the taste as a chunk of pressed sawdust, but no better; and

as human food an eminent New York physician, Dr. Robert T. Morris, pronounces it unwholesome and dangerous. In flavor, cold-storage Quail is far inferior to fresh chicken or turkey. In a court of law, a cooked Quail can easily be identified from squab, reed-bird, "rail-bird" and many others by the fact that the meat on its breast is *white*, while all the others wear dark meat.


**The California Mountain "Quail," or Mountain Partridge,**<sup>1</sup> is a bird of most pleasing appearance, which inhabits California, Oregon and Washington. Wherever protected it is spreading rapidly in the settled portions of the Northwest. It loves moist regions, wherein the rainfall is abundant. This is the bird with a black throat, a white crescent running down from the eye, two rows of white markings on each side, and a long, *drooping* plume on its head running back on the same curve as the forehead. This bird goes in small flocks, of ten to twenty, hides well, and is not easily flushed without a dog.

**The Valley "Quail," or Valley Partridge,**<sup>2</sup> is the bird of the Pacific coast which has the very jaunty, *erect* black plume, rising from the top of its head and gracefully curving forward. Its color markings are rich and beautiful, but not gaudy, and in form as well as color, it is very handsome. In fact, it is the most beautiful of all our small upland game-birds. It inhabits Oregon, Nevada, the whole of California and the Lower California peninsula, and in some places ascends the mountains to 9,000 feet. It has been acclimatized in Utah, and there are many other localities in which it might well be introduced.

This partridge is the most widely distributed and frequently seen game-bird in California, not only in the mountains, but also in the cultivated valleys, everywhere, and even in Golden Gate Park, San Francisco. It breeds readily in confinement in the New York Zoological Park, and when safe from rats is not difficult to keep.

**The Mearns' Partridge,**<sup>3</sup> of Mexico, western Texas and southern New Mexico and Arizona, must be mentioned because it is too odd and

striking in appearance to be ignored. It may be known by the numerous large white spots on the sides of its body just below the wings, and its harlequin head of black and white bars and collars. It is of great interest to Americans residing in Mexico, and many attempts have been made to acclimatize it in captivity in the United States. I once had in my possession two of these birds whose white spots had been artificially changed by some enterprising Mexican to a beautiful golden-yellow color. Until the trick was discovered, the birds were quite a puzzle, for the



CALIFORNIA VALLEY PARTRIDGE.

fact that they had been dyed was not proven until they moulted.

**The Ruffed Grouse**<sup>4</sup> is the dandy of American game-birds. In various places it is called by various names, some of which are mischievously confusing. By many persons it is called a "**Pheasant**," and by others a "**Partridge**"; but both these names are entirely incorrect, and when applied to this bird create confusion. Often it is impossible to converse understandingly about this bird without first defining boundaries, and coming to an agreement regarding the names "Pheasant" and "Partridge." Now that a real pheasant (the ring-necked) has been introduced from China into many portions of the United States, it is all the more imperative that

<sup>4</sup> *Bo-na'sa um-bel'lus*. Average length, 16 inches.

<sup>1</sup> *Or-e-or'tyx pic'tus*. Average length, 11 inches.

<sup>2</sup> *Lo-phor'tyx californicus*. Average length, 9 inches.

<sup>3</sup> *Cyr-to'nyx mon-te-zu'mae mearnsi*. Average length, 8.50 inches.

the Ruffed Grouse should be called by that name, *and no other!* It is called "Ruffed" because of the ruff of feathers that it wears just in front of its shoulders, and under the name "Redruff" this bird has been most charmingly introduced by Mr. Ernest Thompson Seton to many thousand readers who never had known it previously.

This Grouse is in every respect a forest-bird. Its ideal home is mixed forest of hardwood and coniferous trees, with the white-tailed deer and



EASTERN RUFFED GROUSE.

gray squirrel for company. Its home extends from Massachusetts and northern New York to northern Georgia, and westward very sparingly beyond the Mississippi to the Dakotas. Besides being beautiful, it is a bird of interesting habits, and its flesh is entirely too fine for its own good. In size it is smaller than the pinnated grouse, or prairie-chicken, but in intelligence it is second to no other grouse living.

The prevailing color of the Ruffed Grouse is rusty brown, but the mottlings of black, gray and white defy intelligent description. Open or shut, the tail is a dream—cross-barred, banded and mottled most exquisitely. It is no wonder that the male bird is fond of strutting, with spread tail; but besides this it has a still more effective means of attracting the female. It perches on a log, secures a good grip with its feet, then *beats the air* with its wings until you hear at the end of the performance a long, quivering resonance disturbing the solitude, like beating upon a Hindoo tom-tom.

The beats start slowly, but quickly increase in rapidity to the end, thus: "*Dum!-dum!-dum!-dum-dum-dumdumdum.*" The bird does not

beat the log, and it does not beat its own sides. Thoreau declares that its wings strike together behind its back! This "drumming" of the Ruffed Grouse is heard oftenest in spring, and is a signal to the female; but it is also heard occasionally in summer and autumn.

This Grouse is a strong flier, and gets up before the hunter with such a tremendous "*burr-r-r-r*" of wings, and goes off so explosively, that it takes a quick eye and hand to bring it down. It can dash off through timber like a feathered rocket, dodging trees and branches, and zig-zaging in all directions leading away from danger, with a degree of speed and certainty that is really marvellous. No wonder the young hunter who kills one, fairly and squarely, feels proud of his skill, and hastens away to have the trophy mounted for his den.

Unfortunately, in most eastern states, where the Ruffed Grouse should hold its own for a hundred years, this bird is doomed to complete extinction—*unless its sale for the table is immediately and effectually stopped!* So long as it is lawful to sell it, pot-hunters will shoot it, and *snare it*, in season and out of season, as "food" for the already over-fed patrons of fashionable hotels and restaurants of the large cities. As food for the hungry, this beautiful bird is not needed in the least. As a means of inducing thousands of brain-weary men to take healthful exercise in the woods, it will serve a highly useful and important purpose—if not meanly and foolishly exterminated.

The following subspecies, closely related to the typical Ruffed Grouse, are found in North America:

*The Oregon, or Sabine's Grouse*, is found on the mountains of the Pacific coast, west of the Coast Range, from northern British Columbia to California. This species possesses rich red plumage, and is quite beautiful.

*The Canadian Ruffed Grouse* belongs to Canada and Maine, but in the Northwest it ranges south of the international boundary. *The Gray Ruffed Grouse* inhabits the Rocky Mountains from the Yukon to Colorado.

**The Dusky Grouse**<sup>1</sup> is a conspicuous type which inhabits the Rocky Mountains from Idaho and Montana to Arizona. Its other

<sup>1</sup> *Den-drag'a-pus ob-scu'rus*. Average length of male, about 21 inches; female, 18 inches.



names are **Blue, Pine,** and **Gray Grouse,** and also **Pine-Hen.** I first saw it alive in the Shoshone Mountains, while skirting a very steep mountain-side in search of mountain-sheep. The stunted pines that struggle with the slide-rock for existence, were not more than thirty feet high, but in them perched, dangerously near the ground, this handsome slaty-blue Grouse. Its nearest neighbors were the mountain-sheep, elk, magpie, Clarke's nut-cracker, and golden eagle.

This fine bird ranges up to timber-line, but loves rough mountain-sides that are partially covered with pines, cedars and firs. It usually lives alone, but sometimes forms very small flocks. The crop of a specimen which I shot was stuffed full of fresh, green pine needles, some of them two inches long. At that time, however, the snow was a foot deep.

This bird is recognizable by the broad, white band across the end of its tail, and its slaty-blue color. From Alaska to California is found a subspecies, very much like the preceding, called the *Sooty Grouse*. From western Montana to the Coast Range in Oregon and Washington, and northward to Alaska, is found the *Franklin Grouse*, known very generally as the "*Fool Hen*," because it trusts too much to man's humanity, and often finds itself a victim of misplaced confidence. This is one of the last American birds to learn that man is a very dangerous animal, and often devoid both of mercy and of appreciation of the beautiful in bird-life.

**The Canada Grouse,**<sup>1</sup> also called the **Spruce-Grouse** and **Black "Partridge,"** is, as its most acceptable name implies, *the* grouse of Canada and the Northwest. It has the widest range of any American member of the Grouse Family,—from the Alaskan Peninsula southeastward to northern Minnesota, Michigan, New York and New England. It inhabits the evergreen forests of that vast region, usually in very small flocks. It does not really migrate, but by reason of seasonal changes which affect its food supply, it often shifts from one locality to another. (D. G. Elliot.)

In many localities it is known as the "*Fool Hen*,"—a name which is applied in various

<sup>1</sup> *Ca-nach'i-tes canadensis can-a'-ce*. Length, about 14 inches.

places to several other species. Man is so conscious of his own insensate destructiveness, and so accustomed to seeing all wild creatures fly in terror before his baneful presence, he naturally feels that any bird which trusts its life to his tender mercies, and does not live in constant fear of him, must indeed be a feathered fool! For some strange reason, several members of the Grouse Family are surprisingly slow to comprehend man's true nature, and acquire the flight instinct, which most other species learn by experience in a few generations of contact with the Universal Killer.

The male Canada Grouse is readily recognized by its black breast and throat, and black tail, which handsomely set off the barred gray back and sides.

**The Pinnated Grouse, or Prairie-Chicken,**<sup>2</sup> lives chiefly in the memories of those who from 1860 to 1875 were "Western men," or boys. At that time, Illinois, Wisconsin and Iowa, and the states adjoining, were the "West." Railroads were few, all guns were muzzle-loaders, and the game-dealers of Chicago were not stretching out their deadly tentacles, like so many long-armed octopi, to suck the last drop



CANADA GROUSE.

of wild-game blood from prairie and forest. The "market-shooter" was a species of game-butcher then unknown, and the beautiful, fertile prairies, and prairie-farms of Illinois, Wisconsin, Minnesota, Iowa, Missouri, Kansas and


<sup>2</sup> *Tym-pa-nu'chus americanus*. Average length of male, 18 inches.

Nebraska were well stocked with Prairie-Chickens.

In spring they courted openly, and even proudly. The cocks strutted, and inflated the bare, salmon-yellow air-sacs on the sides of their necks, bowed low, and "*Boo-hoo-hooed!*" until the sound rolled over the bare earth in great waves. Then they scattered, to nest and rear their young. In summer, they hid themselves closely; and no self-respecting farmer

early spring, and the long, flaming days of July and August. If the farmers only had been far-sighted, and diligent in protecting for their all-too-scanty recreation, and for their own tables, *the game that was theirs*, they might have had Prairie-Chickens to hunt for a century.

But the game-devouring octopi began to reach out, from Water Street, Chicago, and from New York and Boston. An army of men began to "shoot for the market," and the Pin-



PINNATED GROUSE, OR PRAIRIE-CHICKEN.

dreamed of such a low act as killing one, or meddling with a nest.

In the fall, after the harvesting, and just before the corn-cutting and corn-husking, the young broods were ready to fly, and the flocks began to gather. They first ranged through the wheat and oat stubble, gleaning; and the sport they furnished there,—dear me! Those were the golden days of life on a prairie-farm. The flocks of Pinnated Grouse and quail were the rightful heritage of the boys and men who toiled in the fields through the raw cold of

nated Grouse and quail began to "go east," by the barrel. Some markets were so glutted, time after time, that unnumbered barrels of dead birds spoiled. That was before the days of cold storage.


The efforts that were made to stop that miserable business were feeble to the point of imbecility; and absolutely nothing permanent was accomplished. Had farmers generally stopped all shooting on their farms, as every farmer should, the war on those birds would have stopped also; but the barn was not locked

until after the horse had been stolen. A species destroyed is rarely regained.

To-day, the Prairie-Chicken is to be numbered with the buffalo and passenger-pigeon. It is so nearly extinct that only a few flocks remain, the most of which are in Kansas and Nebraska. If hunting them with dogs continues, five years hence the species will probably be quite extinct.

Even as late as 1874, many birds were killed every winter by flying against the telegraph wires along the railways.

**The Prairie Sharp-Tailed Grouse<sup>2</sup>** inhabits the Great Plains, from the states bordering the Mississippi to the Rocky Mountains. It is the plains counterpart of the pinnated grouse, and like it, is rapidly disappearing before the set-



SAGE-GROUSE.

It is useless to describe this bird. The chances are that no reader of this book ever will see one outside of a museum, or a large zoological garden.<sup>1</sup> The great flocks of from one to three hundred that from 1860 to 1875 were seen in winter in the Iowa cornfields, are gone forever.

<sup>1</sup> During the first four years of its existence, the N. Y. Zoological Park was able to secure only four living specimens.

lements that are fast filling up its home. The neck of the male lacks the side tuft of long, pointed feathers and the naked air-sac so conspicuous on the male pinnated grouse.

To-day, this bird is seldom seen in the open sage-brush plains and bad lands of Montana and Wyoming, but is occasionally found in or

<sup>2</sup> *Ped-i-æ-cet'es phas-i-an-el'lus cam-pes'tris*. Average length, about 17 inches.

near the foot-hills of the Rocky and Big Horn Mountains. When flushed, it makes the mistake of its life in alighting in the low, isolated cottonwood-trees that straggle along the creeks, for when thus perched it takes a strong man to resist the temptation to cut off its head with a rifle ball,—or try to do so. This bird will fly out of the most impregnable cover, and perch aloft to be shot at in a manner indicating a total absence of the most ordinary instinct of self-preservation.

**The Sage-Grouse,** or "**Cock-of-the-Plains,**"<sup>1</sup> is a superb bird—big, handsome and showy. It is one of the very few creatures which can eat with pleasure and benefit the leaves of the common sage-brush, and subsist upon that food indefinitely. Naturally, however, this diet often imparts to the flesh of the bird an excess of sage flavor which renders it quite unpalatable. On this fact alone, the Sage-Grouse can base a hope of a better fate than that of its more edible relatives in the Grouse Family.

Of the really conspicuous members of the Plains fauna,—buffalo, antelope, elk, coyote, gray wolf, swift fox, jack "rabbit," prairie-"dog," and Sage-Grouse,—all have vanished from frequent sight, save the last two, and some have wholly disappeared. In riding in October, 1901, from Miles City to the Missouri River and back, about 250 miles all told, we saw only three coyotes, one gray wolf, and four prairie-hares. Cotton-tail rabbits abounded in the bad-lands, and we saw about six flocks of Sage-Grouse—a very small number for so much territory.

One of those flocks, however, was a sight to be remembered. In the valley of the Little Dry, it spread out, in open order, on a level flat that was covered with short, gray buffalo-grass, and dotted here and there with low clumps of sage-brush. Halting the outfit wagon, I slowly rode forward until within thirty feet of the vanguard of the flock. There were in all forty-six birds, and all were on dress parade. They stood proudly erect, headed across the trail, marched forward in a slow and stately manner, and every weather eye was kept on me. The majority were big, long-tailed cocks. At last the parade terminated in the flight of

the birds nearest me, gradually followed by all the others.

In size, the Sage-Grouse is the largest member of the Grouse Family in America,—next in fact to the magnificent black cock of Europe. When a whole flock suddenly rises out of the sage-brush and takes wing, it is an event to remember. The rush and beat of wings makes a startling noise, and the size of the bird is also highly impressive. This grouse is so large that, as it flies away, you see its body rock violently from side to side, and note the effort of the wings to carry the bird, and maintain a true balance.

The male has an air-sac on each side of its neck, which it inflates in the courting season, when it struts to attract the attention of the females. Recently Mr. Frank Bond has observed, and reported in *The Auk*, that the male also rubs its breast along the ground, as a part of its strutting performance, which accounts for the mysteriously worn condition of the breast-feathers.

It is no more necessary to describe a Sage-Grouse than an elephant. Its size, and its extremely long and pointed tail proclaim its identity anywhere. According to Mrs. Bailey, it ranges "from Assiniboia and British Columbia to Utah, Nevada and California, from the Sierra Nevadas and Cascades east to the Black Hills, Nebraska and Colorado." I will only add the earnest wish that every one who reads these notes may some day have the pleasure of seeing at close range this glorious bird in its ideal home,—on a sage-brush flat in the land of buttes, where the world is big and free, and full of sunshine.

**The Ptarmigans** (pronounced tar'mi-gans) form a sharply distinguished group of the Grouse Family, with which, in view of the different species we possess in Alaska, and also nearer home, every American should become acquainted. The most striking and peculiar character about these birds is that at the approach of winter they turn snow-white. They prefer to nest on the tops of rugged mountains, above timber-line, and in Alaska are at home either on the lofty snow-fields of the mountains, or the desolate barrens.


There are four well-defined species, and six varieties. The only species which is at home in the United States is the **White-Tailed**

<sup>1</sup> *Cen-tro-cer'cus u-ro-phas-i-an'us*. Length of male, 27 inches; female, 22 inches.

**Ptarmigan**<sup>1</sup>—in Colorado sometimes called the "White Quail,"—which lives in the Rocky Mountains from the Liard River, British Columbia, to New Mexico. It is said that another species (the *Willow*) does occasionally wander down into northern New England. The majority of the species are found in Alaska, but the *Rock Ptarmigan* covers nearly the whole of

Lena River, their last food was one of these birds, shot with a rifle by Alexy, the Eskimo. In northern Greenland and Grinnell Land Peary and Greely ate it, and in the Kenai Peninsula, flocks of it were photographed by Dall DeWeese and others.

This bird is almost constantly busy in changing its clothes. In the spring it goes by slow



Summer plumage.

WILLOW PTARMIGAN.

Winter plumage.

Drawn from photographs made in Alaska by DALL DEWEESE.

Arctic America from Alaska to Labrador and Greenland. Two of its subspecies inhabit Newfoundland.

The **Willow Ptarmigan**<sup>2</sup> may well be chosen as the typical representative of the whole group, for its distribution covers the Arctic lands entirely around the pole. When De Long and his party fought starvation at the mouth of the

degrees from winter white to chestnut brown, barred with black. By July the dark plumage of midsummer is fully developed; but not for long. By the first of September, the trouble begins once more, and feather by feather the plumage gradually changes to snowy-white. In winter the legs and feet of Ptarmigans generally are heavily clothed with feathers, and often only the ends of the toes are visible.

<sup>1</sup> *La-gopus leu-cu'rus*. Length, about 12 inches.

<sup>2</sup> *La-gopus lagopus*. Length, about 14 inches.

As might be expected, this bird and its rela-

tives often constitute an important source of food supply for the Indians and Eskimos of the arctic regions.

### THE PHEASANT FAMILY.

#### *Phasianidae.*

The Pheasant Family was originally represented on this continent only by the wild-turkeys; but during recent years certain foreign species have been successfully introduced, and



WILD-TURKEY, FROM VIRGINIA.

are now becoming so numerous as to require notice.

The Ring-Necked Pheasant<sup>1</sup> has been introduced from China, and acclimatized in Washington, Oregon, California, British Columbia, and elsewhere with pronounced success. In many localities it has become so abundant that now it is shot by sportsmen as upland game-birds once were killed in New York state. From Portland, Oregon, to Vancouver the taxidermists are annually called upon to mount scores of these birds, because they are so beautiful that many of the sportsmen who shoot them cannot consent to see their skins destroyed.

Following the examples of the Pacific states, Ohio, New York, Massachusetts, Pennsylvania, and several other states both east and west have entered seriously upon the business of breeding, rearing and introducing this valuable bird at state expense.

The Silver Pheasant, and the very beautiful

<sup>1</sup> *Phas-i-an'us tor-quat'us.*

Golden Pheasant, both natives of China, have also been acclimatized in Washington and Oregon. In view of the strong and hardy natures of both these birds, there should be little difficulty in introducing them in any well-wooded farming region east of the Mississippi, and south of the fortieth parallel.

The Wild-Turkey<sup>2</sup> once inhabited nearly one-half of the United States; and, considering the great size of the bird, the earnestness of our efforts to exterminate it, and the very little that has been done toward its protection, its survival to-day is cause for wonder. It is yet found in a few heavily timbered regions in the East and South,—such as Florida, the Virginias, Pennsylvania, and a few of the southern states. It is doubtful if even one flock exists in the north anywhere west of Pennsylvania. In Oklahoma and Texas it still lives, but the gunners of the cattle-ranches are fast killing off the few flocks that remain.

The Wild-Turkey is the king of upland game-birds. It has been given to but a few hunters to seek this bird in its native forests, witness its splendid flight, and afterward shoulder a giant gobbler weighing from twenty-five to thirty pounds for a ten-mile carry. He who has done this, however, will thereafter rank this bird as second to none on earth. In the United States only one species exists, but three geographic races have been described. The wild bird so closely resembles the domestic turkey that almost the only difference observable is the white upper tail coverts of the tame bird.

The Ocellated Turkey,<sup>3</sup> of Yucatan, British Honduras and Guatemala is a bird of more splendid plumage but smaller size than our northern species. Its name refers to the beautiful eye-spots of blue, green and purple which adorn the tail-feathers. The prevailing color of the body plumage is a rich metallic green, exhibiting the brilliant iridescence and burnished-bronze effects so strongly displayed in most turkeys in full plumage. On account of its great beauty, several attempts have been made to establish this species in zoological gardens, but thus far quite without success. It is to be hoped that future efforts will succeed.

<sup>2</sup> *Me-le-a'gris gal-lo-pa'ro.* Length of large male, about 46 inches; weight, 28 pounds.

<sup>3</sup> *Me-le-a'gris oc-cl-la'ta.*

## CHAPTER XXIV

# THE ORDER OF SHORE-BIRDS

### LIMICOLAE.

There are many genera and species of birds in this Order, but for certain reasons it is difficult to form an acquaintance with more than a very few of them. The majority of them reach us only as birds of passage, on the way to or from their breeding grounds farther north, and during the year are with us only a few weeks. Others are so few in number, and live in such remote localities that they, also, are beyond our acquaintance. As usual, therefore, we will introduce only those species that are sufficiently abundant, long-tarrying and generally interesting to make them worth knowing.


As the name of the Order indicates, these birds live on the ocean and lake beaches, and the banks of rivers, ponds and pools, where they find many kinds of queer things to feed upon. On the boundary line betwixt sea and land they find many insects, shell-fish, crustaceans and worms. The Turnstones make a business of turning over pebbles and small stones, in order to capture the worms and insects that take shelter under them.

**The Kill-Deer Plover**<sup>1</sup> makes an excellent representative of a large section of this Order. It is of average size, and handsome appearance, and is such a loud and frequent caller its presence is always well advertised. It is so widely distributed that millions of people may know it if they will. It is a bird of the inland ponds and pools, not of the sea-shore, and it is found throughout the whole temperate portion of North America, from the Atlantic to the Pacific. It is not a bird of heavily timbered regions, however, and is most abundant in the lake regions of the Mississippi valley. On the prairies of the Middle West, wherever there are small, shallow ponds, or even pools in wet meadows, all through the season of mild weather you will hear its clear and rather strident cry of "*Kill-d-e-e-r!*"

<sup>1</sup>*Ox-y-e'chus vo-ci'fer-a.* Length, 10½ inches.

*Kill-d-e-e-r!*" And it is always a pleasing sight to see this immaculate bird in snow-white, brown and black plumage standing at the edge of a bit of water—a stroke of living high-light in the landscape. I always liked the Kill Deer, and, although I have seen hundreds, and heard its cry a thousand times, I never wearied of its companionship. In my opinion it is our most beautiful shore-bird.

**The American Golden Plover**,<sup>2</sup> also called **Green and Field Plover**, is (or, at least *was* until recently) the Plover most frequently seen in the Atlantic states, and in the markets. It frequents the banks of marshes and tide pools along the sea-shore, but it is equally fond of the pools and ponds of the uplands, particularly in



KILL-DEER PLOVER.

old meadows. They are seldom seen during the spring migration; they do not remain with us during the summer, and it is only during the months of their fall migration, from August 15,

<sup>2</sup>*Char-a-dri'us do-min'i-cus.* Average length, 10 inches.



AMERICAN WOODCOCK.

to November 1, that they are really in evidence. During the open season they are much sought by gunners,—which is the reason why there is now only one bird where formerly there were ten.

The American Woodcock<sup>1</sup> is the oddest-looking land-bird in North America. Its legs are too short for so large a body, its tail is only half as long as it should be, its neck is too short and too thick, and its head is entirely out of drawing. The eyes are placed too far back, and the bill is too long and too straight. In appearance, the Woodcock looks like an avian caricature.

But, odd or not, this bird is very dear to the heart of the great American sportsman, and its plump brown body is a genuine delicacy. It has a long array of local names, some of which are so uncouth that the less said concerning them the better.

The long, sensitive beak of this bird is really a probe and a pair of forceps combined, for probing in soft earth or mud after earthworms, and dragging them out when found. In order to feed, the Woodcock has no option but to frequent the moist banks of wooded streams, or wet grounds in the shelter of bushes or timber, where it can work unobserved. During the day, it lies low to escape observation, and does the most of its feeding at night. It is seldom found in open ground, and Woodcock shooting is much like shooting quail among brush—quick and difficult.

<sup>1</sup> *Phi-lo-he'la mi'nor*. Average length, about 10.50 inches.

This bird ranges throughout the United States from the Atlantic coast to the edge of the Great Plains. In the course of much hunting in central Iowa I never but once shot a specimen of this species.

As a highly esteemed game-bird, **Wilson's Snipe**, or the **Jack Snipe**,<sup>2</sup> is a close second to the woodcock. Like the latter, it has a long, straight bill with a sensitive tip, with which to probe down in mud or soft earth of pond margins or spring holes, to the home of the angle-worm. Unlike the woodcock, however, this Snipe is a very well-formed bird, and it feeds more in the open, which renders its pursuit more fruitful of results. On the wing, it is awkward and angular looking. It flies in a very angular course, but so rapidly it is a difficult mark to hit. When it rises, it utters a shrill cry, half scream and half squawk, and in windy weather it often flies quite high.

This Snipe has a very wide range—from Alaska and Hudson's Bay through all the United States, except the arid regions, to northern South America. Its most conspicuous color is brown, striped on the back with black, which in brushy ground protects the bird so well it is difficult to distinguish it.

Whenever at the sea-shore in warm weather



WOODCOCK ON NEST.

Photographed at a distance of 6 feet, by LE ROY M. TUFTS, and copyright, 1903.

you wander "far from the madding crowd," you may make the acquaintance of the **Semi-palmated Sand-Piper**,<sup>3</sup> or possibly it will be

<sup>2</sup> *Gal-li-na'go del-i-co'ta*. Length, about 11 inches.

<sup>3</sup> *Er-e-un-e'tes pu-sil'lus*. Length, 6 inches.



the **Least Sand-Piper**,<sup>1</sup>—a trifle more minute, and with no web at the base of its toes. At a distance of ten feet the two species look precisely alike, and there is no need to worry about an exact identification. They are also called "Peeps," and "Ox-eyes," and the toes of the Semipalmated Sand-Piper are partly webbed.

As the green-topped surf dashes to pieces on the pebbles, and goes sliding in a silvery sheet up the yellow sand, you will notice just above its frothy edge a flock of little gray sprites, their tiny legs twinkling as they patter swiftly over the smooth floor. Sometimes the sliding sheet of water overtakes them. If it is nearly spent, they mind it not; but if the rush is too strong, up springs the flock, all members at the same instant, and with quick flashes of light gray wings, it skims the surf-sheets or the sand, to a point farther on. The unison of action in the rising, flight, and landing of the flock is as perfect as if each little pair of wings were worked by the same wires. How does each bird know the impulses of all the others? Watch them, and see if you can guess the secret.

At the sea-shore I never weary of watching these busy little creatures, and never fail to be amused by the twinkling of their tiny legs as

it has brought to them from the sea, or uncovered on the sand.

Fortunately these birds are so small the gunners are not slaughtering them—as yet.



LEAST SAND-PIPER.

But their day of doom is not far distant. There is in every country on earth a lawless, miscreant element which is devoid of all love for Nature and wild life, and which sticks at nothing in the line of destruction. It pollutes streams, dynamites fish, poisons dogs, steals ash-cans and swill-pails, and kills song-birds for "food."

Some day, alas! the evil eye of this bad breed will fall upon the flocks of Sand-Pipers by the sea, and on the shores of inland lake, pond and stream. Then the little gray clouds will quickly vanish forever. To-day, however, both the species mentioned above are found sprinkled throughout the whole eastern United States, and they breed northward quite up to the arctic Barren Grounds. Wherever they are, they are interesting birds, and worthy of your friendship.

**The Long-Billed Curlew**<sup>2</sup> is a bird which has caused much wonderment and many guesses in the Middle West, where on the virgin prairies it once was frequently seen. This bird's trick of holding its wings high above its back for two or three seconds after it alights upon the ground always attracts special attention. Its cry, also, oft repeated in spring, is very weird and peculiar, and well calculated to make the bird remembered.

This bird once was common on the rolling  
<sup>2</sup> *Nu-men'i-us lon-gi-ros'tris*. Average length, about 23 inches; bill of adult bird, about 8 inches.



WILSON'S SNIPE.

they run before the water. As the sheet of surf recedes, down they run after it, to pick up whatever of insect or other edible animal life

<sup>1</sup> *Ac-to-dro'mas min-u-till'a*. Average length, 5.50 inches.

prairies of Iowa, regardless of ponds or streams, where it sought every sort of animal life small enough to be swallowed. It is easily recognized, even in flight, by its long, curved bill. In its form, its beak and its legs, it is almost a perfect counterpart of a typical ibis, but it has the mechanically mottled plumage of a typical shore-bird. Although by some ornithologists this bird is credited to the whole length and breadth of the United States, there certainly are some very wide regions from which it is totally absent. In various localities it has various names, some of which are Sickle-Bill, Saber Bill, Smoker, Spanish Curlew and Mowyer.

This bird is very sympathetic toward its wounded mates, and in response to the cries of a bird that has been shot, a flock sometimes will return, and with loud cries circle near the gunner, at close range, until several more have been brought down. (D. G. Elliot.)

Besides the shore-birds mentioned above, there are several groups which are of interest chiefly to the special student, and which there is no space to introduce here, save by name. There are the oyster-catchers, turnstones, godwits, stilts, and phalaropes. In the *Order Limicolae* as a whole there are in North America, north of Mexico, about seventy-five species and subspecies.

## CHAPTER XXV

# THE ORDER OF CRANES, RAILS, AND COOTS

### PALUDICOLAE

The name of this Order, *Pal-u-dic'o-lae*, means "marsh-dweller," and the presence in it of the cranes is enough to make it notable. It must be admitted, however, that from the stately and commanding crane down to the humble coot, the scared gallinule, and the diminutive rail, is a long step downward. But it is inevitable that the efforts of science to classify the birds of the world in as few Orders as possible, should bring together many widely divergent forms. To have a greater number of Orders would be still more confusing to the general student than the present number.

In the order of Marsh-Dwellers there are only two Families which are entitled to notice here. These are the Cranes, and the Rails, Gallinules and Coots.

### THE CRANE FAMILY.

#### *Gruidae.*

The Cranes of the world form a group of about eighteen species, which, in stateliness, beauty and oddity of habit, are second only to the ostriches and their allies. Every zoological garden which possesses a good collection of cranes has good reason to be proud of it. The *Crowned Cranes* of Africa are the most beautiful species of all, the *Paradise Crane* is the oddest in appearance, the little *Demoiselle Crane*, of the Nile region, has the most amiable disposition. The big, red-headed *Saras Crane* of India is the most quarrelsome, and the stately *Whooping Crane* of North America is the species which comes nearest to being pure white.

Through some mischievous and unfortunate circumstance, the great majority of the people who live in the eastern United States have become almost fixed in the habit of calling the great blue heron the "blue crane." The former is common enough along watercourses and tidal

rivers, but it is probable that not more than one person out of every ten thousand has ever seen in America a living wild crane. As applied to wild-birds, the word "crane" should be used most sparingly. Along the Atlantic coast, the only locality in which it might correctly be used afield is on the interior savannas of Florida.

**The Whooping Crane**<sup>1</sup> is now one of the rarest of all living North American birds. Three years of diligent quest for living specimens have produced but one bird. There were in captivity on January 1, 1903, exactly six specimens, four of which were in the United States. Inasmuch as this bird is of no value save to zoological gardens, it must be believed that it has been wantonly shot, down to the verge of extinction. Since it is a practical impossibility to induce it to breed in captivity, the species seems almost certain to disappear from our fauna at an early date.

As seen with its wings closed, the visible plumage of this grand bird is all snowy white. When the wings are spread, however, it is found that the largest feathers, called the primaries, are jet black. The upper tail coverts form a plume that arches upward over the tail, and gives the bird a very jaunty air. The top of the head is bare of feathers, and the rough skin has a dull-red glow. The eye is big and keen, and the bill is long, strong and rather blunt on the end, for digging angle-worms out of the ground, not for spearing fish.

The strength of the beak and neck of the Whooping Crane in the New York Zoological Park is truly remarkable. The bird roams at will in a grassy meadow of about two acres in extent. Soon after it attained full growth, it was noticed that after every rain, it would vigorously attack the grass. With mandibles

<sup>1</sup> *Grus americana.*

two inches apart at the tips, it would drive its beak into the earth to a depth of from two to three inches, grasp a tuft of grass between them, and by main strength deliberately pull it up by the roots. A few vigorous shakes sidewise dislodged any angle-worms which might have been brought up, after which the roots of the tuft would be carefully looked over before being cast aside. Next in order, the wounded earth would be carefully probed and picked



New York Zoological Park.

WHOOPING CRANE.

over. In a few hours, this bird sometimes pulled up the grass on a space fifteen feet square, and finally disfigured the ground so seriously that after every rain the Crane had to be shut up.

A living full-grown Whooping Crane stands 4 feet, 3 inches high. Its name is due to its wonderfully clear, powerful, and trumpet-like call, which is uttered with the beak pointing straight upward. When properly delivered, the crane's call consists of two notes, an octave apart, one following the other so closely that there is no interval, thus: "Quah-KEE-E-

E-oo!" I believe that a Crane's trumpet-call will carry as far as the roar of a lion.

All our Cranes are strictly open-country birds, and formerly inhabited the fertile, froggy prairies and cornfields of the Mississippi valley; but the species named above never was really numerous anywhere. In travelling, cranes always fly in single file, with their long necks and legs in a straight line, and in that position the length of the bird seems very great.

**The Sandhill Crane**<sup>1</sup> is a smaller bird than the preceding, always has been more numerous, and therefore is much more widely known. In color it is a dull bluish-slate, and it has a half-bald, dull-red head, like a whooping crane. The pioneers who were on the western prairies from 1850 to 1870 occasionally saw long lines of enormously long birds sailing high in the heavens, trumpeting their identity to those unable to see them, or alighting on stilt-like legs in the cornfields. In springtime, when the birds alighted in the bare fields, and stalked about with majestic stride, they seemed fairly gigantic. They went far north in spring to breed, and on their return trips sought their winter home in Texas, Florida, and elsewhere along the Gulf coast.

Cranes in captivity, and wild ones, also, often indulge in strange antics. Suddenly, and for no apparent reason, one will half-open its wings, leap into the air, and begin to dance. It bobs and bows, salaams, and courtesies almost to the ground, and in sheer delight repeatedly leaps into the air. Often the lead of one bird is followed by several others, and occasionally (as I have myself seen), a whole wild flock of fifteen or twenty birds will join in the fandango.

Whenever the days are cool and clear,

The sandhill crane goes walking  
Across the field by the flashing weir,  
Slowly, solemnly stalking.

The little frogs in the tules hear,  
And jump for their lives if he comes near;  
The fishes scuttle away in fear  
When the sandhill crane goes walking.

The field folk know if he comes that way,  
Slowly, solemnly stalking,  
There is danger and death in the least delay,  
When the sandhill crane goes walking.

<sup>1</sup> *Grus mexicana*. Height, about 3 feet, 10 inches.

The chipmunks stop in the midst of play;  
 The gophers hide in their holes away;  
 And "Hush, oh, hush!" the field-mice say,  
 When the sandhill crane goes walking.

MRS. MARY AUSTIN, in *St. Nicholas*.<sup>1</sup>

## THE FAMILY OF RAILS.

### *Rallidae.*

From the stately crane to the timid, self-effacing **Virginia Rail**<sup>2</sup> is going at one step from the sublime to the ridiculous. To the latter, which is a bird about half the size of a bob-white, a crane must seem like a giant whose head is in the clouds. The crane can either fight, run or fly away; but the rail is safe only when threading the mazes of a reedy marsh, where no enemy can follow it far. When boating on a marsh filled with cat-tails, or reeds, or tall grass, you may hear a score of rails clucking and calling in the heart of the green tangle about you without seeing one. There are times when it seems as if this bird is a deliberate and intentional ventriloquist, for its voice seems to come from all directions save that which points toward its owner. A marsh is as necessary to rails as water is to fishes.

When a rail flies up out of a marsh or a meadow, you can recognize it by its feeble, fluttering flight, and its hanging legs. Often in alighting it seems to fall helplessly into the tall cover.

In the mosquito-ridden marshes along the New Jersey shore, dwells a species known as the **Sora Rail**<sup>3</sup> in numbers sufficiently numerous to attract gunners. The moment the "law is off," the flat-bottomed boats are brought out, and the fusillade begins. With no larger game available, even a small Rail can form an excuse for a day's outing on the marshes, bringing the grip of the gun-stock, the dull "boom" that is music to the desk-weary man, and the welcome snell of gunpowder. Therefore, rail not at all those who shoot rails; for there be some who do not shoot "for revenue only."

As may be inferred, rails are good to eat, though not very good; for they are several sizes too small for real comfort. There are only

<sup>1</sup> By courtesy of THE CENTURY CO., and of the Author.

<sup>2</sup> *Rallus virginianus*. Average length, 9 inches.

<sup>3</sup> *Porzana carolina*. Length, about 9 inches.

about twelve species in North America, of which the *King Rail*, 15 inches long (of eastern North America), is the largest, and the *Virginia Rail* is the most widely distributed. The latter has a long bill (1½ inches), and is found from Long Island to British Columbia, breeding everywhere that marshy lands occur. It is an olive-brown bird, streaked and barred with black, and in places with white, also.

While the most typical rails have long bills, some species are short-billed.

A **Gallinule** is a bird which lives, acts and looks like a rail, and is easily mistaken for either a rail or a coot; but it stands midway between



VIRGINIA RAIL.

the two. It is distinguished from the rails by the bare, horny shield upon the forehead, and from the coots by the long, slender, unwebbed toes. The **Florida Gallinule**<sup>4</sup> is also called the *Blue "Rail,"* and *Red-Billed "Mud-Hen,"* and its general color-effect is bluish-gray. It is found in localities adapted to its habits throughout temperate North America, north to Canada, and as far south as Brazil.

The **Purple Gallinule**<sup>5</sup> of the southern half of the eastern United States, is a bird of beautiful plumage. Its colors are a rich, dark purple on the head, neck and shoulders, lightening to

<sup>4</sup> *Gal-li-nu'la gal-e-a'ta*. Length, about 13 inches.

<sup>5</sup> *I-o-nor'nis mar-tin'i-ca*. Length, 12 inches.

peacock blue on the back and lower breast. Even as it rises beside your railway train you can easily recognize it before it is lost to view. It still breeds on the head waters of the St. Johns, opposite Melbourne.

**The Coot, or Mud-Hen,**<sup>1</sup> is a bird of the small creeks, and the shores of shallow lakes and ponds where cat-tails, lizard-tails, iris and



THE COOT.

rushes grow abundantly. It is natural for any one who writes about a bird to think of it as he saw it most impressively. My memory goes back to my first days of alligator and crocodile hunting, in the little creeks that flow from the Florida Everglades into the head and western side of Biscayne Bay. Then and there, Mud-

<sup>1</sup>*Fu-li'ca americana*. Average length, 14.50 inches.

Hens were so numerous and so tame they became positively monotonous. As we rowed silently along Snake Creek, or Arch Creek, the man in the bow ready for the next "big, old 'gator" found sunning himself at the edge of the saw-grass, up would go three or four slaty-blue birds of the size of bantam hens. With feeble flight, and feet pattering on the water to help along, they would fly ahead of the boat in a most offensively ostentatious manner. Of course any old alligator knows that a scared Coot usually means a boat; and since every boat is known to be loaded, the natural sequence of a frightened Coot is the bottom of the creek.

The foot of the Coot is very curiously formed. It looks as if originally it had been fully webbed, but some one in sportive mood took a pair of scissors, cut out the centre of the web, and cut deep scallops in the web along each side of each toe. The foot, therefore, is half webbed,—an excellent arrangement for running on water when the wings lend their assistance. This bird never rises on the wing without a preliminary run on the water of from fifty to one hundred and fifty feet. It swims and dives quite well, but as a rule it prefers to live as do the rails and gallinules, in the edges of heavy marsh vegetation, where it can pick up its living of buds, blossoms, seeds, aquatic insects and snails, and also hide from its enemies.

As yet the Coot is not considered a "game-bird," and is not slaughtered for food; but, once let the evil eye of the Epicure fall with favor upon this bird—or any other—and its doom will be sealed.

The distribution of this species is given as "from Greenland and Alaska southward to the West Indies."

## CHAPTER XXVI

# THE ORDER OF HERONS, STORKS, AND IBISES

### HERODIONES.

All the members of this Order are either sturdy fisherfolk, or loughshorem. They wait not for bud or blossoms, or ripening grain, but when hunger calls they go a-fishing. Then woe betide the small fish or frog of any size which is tempted to stray into the warm shallows, and linger there.

The neck of the heron is specially formed by Nature for quick lunging. At rest, it folds upon itself, in angular kinks, until the neck totally disappears, and the bird's head seems to rest down upon its shoulders. But alarm this neckless bird, and presto! it is another creature. Up goes the head into the air, borne on a long, flat-sided neck, which curves like a capital S.

When a heron is fishing, it stalks slowly and silently along the shore, preferably in water about six inches deep, its head carried well forward but about on a level with the top of its shoulders, while its big eyes keenly scrutinize every object in the water. It takes long steps, and plants each foot softly, in true still-hunter fashion, to avoid alarming its game. When a fish is found within range, the kinks of the neck fly straight, and the fish is seized between the mandibles. The fish is not stabbed through and through, as is generally supposed. In swallowing a fish, it is, of course, taken head first.

Hérons, egrets and ibises are gregarious, or sociable, in their nesting habits. In other words, they are fond of nesting together; and a place of many nesting birds is called either a "heronry," or a "rookery." The nesting sites are chosen with due regard to seclusion and food supplies. Usually the heronry is located in low trees that stand on a small island, or else grow up out of a swamp or bayou, so that without a boat they are almost inaccessible.

Thirty years ago, the greatest and most numerous heronries in the United States were in

Florida, on the head waters of the St. Johns, on the edge of the Everglades, and the small rivers and creeks that run down to the sea. To-day it is difficult to find in Florida a heronry worthy of the name, or one which belongs to a large assemblage of birds. Herons, egrets and ibises have been so persistently destroyed for their "plumes" that not one-fiftieth of the original number remain.

As will be seen by the following table, the Order Herodiones contains quite a number of important water-birds which are not herons:

	FAMILIES.	EXAMPLES.
ORDER HERODIONES.	HERON, <i>AR-DE'I-DAE</i> , . . .	{ Herons, Egrets and Bitterns.
	STORK, . . . <i>CIC-O-N'I-DAE</i> ,	Wood Ibis.
	IBIS, . . . <i>I-BID'I-DAE</i> , . . .	{ White Ibis and Scarlet Ibis.
	SPOONBILL, <i>PLAT-A-LE'I-DAE</i> ,	{ Roseate Spoonbill.

### THE HERON FAMILY.

#### *Ardeidae.*

The Great Blue Heron<sup>1</sup> is the largest, handsomest and most conspicuous Heron in North America—if not the world. This is the bird so persistently called the "*Blue Crane*"; and one of the first things for the beginner to learn about birds is to call this bird a *Heron*, instead of a "crane!"

Whether fishing in the shallows along the shore, or perching on a dead tree, or winging his way slowly and majestically through the air, this is a fine, handsome bird, and a welcome sight to see. Its height when standing fairly erect is 3 feet, 3 inches. It has plumes on its head, breast and back, which American cranes

<sup>1</sup> *Ar-de'a her-o'di-as*. Length, from 40 to 48 inches.

do not have. It is never seen away from water-courses, and, it may be added, in warm weather no river-scene is truly complete and perfect without one!

When seen with closed wings, its upper neck and body are of a bluish-slate color, and its under surfaces are white, streaked up and down with black. In the North, this bird is shy, and afraid of being shot at; but in the tropics, where they are not persecuted, I have some-



New York Zoological Park.

GREAT BLUE HERON.

times approached within thirty feet of full-grown birds without alarming them.

The range of this bird is from the arctic regions southward wherever the conditions of water, timber and food are suitable, to the West Indies and South America; but there are many arid and treeless regions from which it is totally absent.

**The Little Green Heron**, or "Fly-up-the-Creek,"<sup>1</sup> is found throughout the well-watered regions of the United States, wherever timber

<sup>1</sup> *Bu-tor'i-des vi-res'cens*. Average length, about 18 inches.

is plentiful. In many localities of the Middle West and the Mississippi valley from which the great blue heron is now absent, this is the only heron to be found; and away from the Atlantic coast it is the most familiar member of its Order.

Its body is about as large as that of a sparrowhawk, and when in a crouching attitude it is a very proper-looking bird. With its neck stretched, however, and its head held high, the

body seems much too small, and the neck makes the bird seem top-heavy. Start it off in flight, however, and it is one of the most ill-fitting herons that ever took wing. It is so angular and loose-jointed it seems ready to fall to pieces, and its flight is slow and feeble. The prevailing color of its plumage is a beautiful metallic green, but the flat shape of its neck, and the peculiar set of the feathers thereon have caused many young taxidermists some very sad hours.

The food of the Green Heron consists of minnows, small frogs, tadpoles and insects.

**The Little Blue Heron**<sup>2</sup> is yet fairly abundant in Florida, because it bears no fatal "plumes." In summer, this species sometimes wanders northward as far as Illinois and Maine. One striking peculiarity of its plumage is worthy of special mention. Until one year old, the young birds are snow-white, and

look precisely like young snowy egrets which are of corresponding size and form. Sometimes it is a matter of difficulty to convince a person that a *snow-white bird* is a Little Blue Heron, in its first year. But the moulting finally tells the story. First the plumage is flecked with blue, then it is half blue, and at last the solid blue color prevails. It seems to me that in clothing young and inexperienced birds in snow-white robes, which attract all eyes to them, Nature forgot all about "protective coloration!"

<sup>2</sup> *Ar-de'a cae-ru'le-a*. Average length, 24.50 inches.



**The Black-Crowned Night-Heron<sup>1</sup>** breeds all around New York City, and every summer two or three come and try to break into the great Flying Cage of the New York Zoological Park. As its name implies, this bird has a crown of glossy black feathers, with two or three long white occipital plumes. This is a southern bird, but it breeds as far north as Massachusetts and Illinois. Like its twin, the *Yellow-Crowned Night-Heron*, it is half nocturnal in its habits. When at night in Florida you hear a bird say "*Quawk!*" and repeat it to you from the depths of the mangroves as your boat glides by, you know it is a Night-Heron. Both these species have beautiful plumage, and are handsome birds. Their distinguishing marks are, thick bodies, and short, thick necks; short legs (for herons), and two or three round, wisp-like plumes from five to seven inches long growing out of the top of the head, and drooping backward.

**The Snowy Heron, or Snowy Egret,<sup>2</sup>** when fully adult, is the most beautiful white bird in all the avian world. Its form is the embodiment of symmetry and grace, its plumage is immaculate, and the filmy "plumes" on its head and back are like spun glass. Its black legs and bill merely serve to intensify the whiteness of its feathers.

But the vanity of women has been the curse of the snowy egret. Its plumes are finest during the breeding season, and it was then that the hunters sought them, slaughtering the parent birds in the rookeries by thousands (when they were abundant), and leaving the nestlings to die of starvation. If all women could know the price in blood and suffering which is paid for the accursed "*aigrettes*" of fashion, surely but few could find any pleasure in wearing them.<sup>3</sup> It is strange that civilized women—the tender-hearted, the philanthropic and compassionate—should prove to be the evil genius of the world's most beautiful birds.

<sup>1</sup> *Nyc-ti-co'rax nycticorax nae'vi-us*. Length, 24.50 inches.

<sup>2</sup> *E-gret'ta can-di-dis'si-ma*. Length, about 23 inches.

<sup>3</sup> Thanks to the efforts of the Audubon Societies, the American Ornithologists' Union and the United States Biological Survey, the laws of the United States now prohibit the sale of aigrettes throughout the United States, irrespective of the countries from which they come.

In Florida, this bird once lived and bred, in thousands, on the head waters of the St. Johns, around the Everglades, and the heads of the streams that run down to the sea. At the first shot fired in a rookery, a white cloud would arise, and old residents tell how "the savannas were sometimes white" with these beautiful creatures. To-day, not half enough



LITTLE GREEN HERON.

remain to stock our zoological gardens. The slaughter has been exasperatingly complete, and protection has come too late.

In the United States the Snowy Egret exists now only by accident, and the "plume-hunters" are pursuing this and the next species in Central and South America, to their most remote haunts, sometimes even at the risk of their lives. Fashion, cruel and remorseless, has decreed that the egrets must go!

**The American Egret, or Great White Egret,<sup>4</sup>** is, when adult, our second largest bird

<sup>4</sup> *Her-o'di-as e-gret'ta*. Length, about 40 inches.



New York Zoological Park.

## GREAT WHITE EGRET.

of the Order of Herons with pure white plumage, the great white heron being the first. Much to the misfortune of this species, it possesses about fifty "aigrette" plumes which droop in graceful curves from the middle of its back, far beyond the tail and wing tips. For these beautiful feathers this bird also has been pursued by plumehunters, to the point of total extermination. A very few individuals are yet living in Florida, but they will all be blotted out within a short period.

The American Bittern<sup>1</sup> is a fairly large bird, of a yellowish-brown color, elaborately mottled and streaked with various shades of light and dark. When standing in concealment, it draws in its neck until it wholly disappears in its plumage. The result is an egg-shaped bird, with a beak at the small end, pointing heavenward, and short, thick legs below. I have seen a Bittern stand motionless in that idiotic atti-

<sup>1</sup> *Bo-tan'rus len-tig-i-no'sus*. Length, 26 inches.

tude, for nearly an hour at a time. Even in the whirling gayety of a big Flying Cage, it takes life sadly, and never makes merry, as do all other birds, even the funeral vultures. Standing erect, however, the Bittern is a bird with a fair length of neck; but its neck seems much too large and heavy for its body.

Because of the peculiar sound it utters, the Bittern is called the "*Stake-Driver*," and "*Thunder-Pumper*." I have never heard thunder pumped, but with stake-driving am quite familiar, and must say that I never heard a Bittern give forth a cry that sounded like it. I think also that the "booming" of the Bittern should be taken subject to inspection and approval; for to at least one tympanum there is a wide difference between a real "boom" and the alleged "boom" of the Bittern.

This bird inhabits sloughs and marshes of tall, rank grass, in which it hides most successfully by standing very erect, and pointing its



AMERICAN BITTERN.

beak toward the zenith. It feeds chiefly upon frogs, small snakes, lizards, and crawfish.

**The Least Bittern**<sup>1</sup> is the smallest member of the Heron Order,—a queer little brownish-yellow and black creature, duly mottled, of course, with a sparrow-like body, and a wide, flat neck several sizes too large for the body of the bird. On the whole, it is a pretty little creature, associated by habit with the long-billed marsh-wren, the rail, and the red-winged blackbird.

### THE STORK FAMILY.

#### *Ciconiidae.*

The real Storks are found only in the Old World; but the **Wood Ibis**<sup>2</sup> is a member of the Stork Family, and he looks it. He is a big, burly, bald-headed, good-natured bird, standing 31 inches high. No matter what goes on around him, he is as solemn as an owl. Although large enough to do much damage to birds smaller than himself, he associates with herons, ducks, geese, and ibises, of all sizes, without the slightest desire to harm any of them, or even to rule them. In a large bird, capable of much mischief, such perpetual good temper is worthy of note.

When this bird is adult and clean, its plumage is pure white, and it is a noteworthy member of any feathered community. Specimens are nearly always procurable in Florida at a reasonable price (\$8), and there are always several in the New York Zoological Park. This species "breeds in Florida and the Gulf states, after which it wanders north as far as Kansas, Indiana and New York." (F. M. Chapman.)

### THE IBIS FAMILY

#### *Ibididae.*

In North America this Family contains three species of birds that are heron-like in general form, but are quite differently provided as to their bills. The bill of a true ibis is long, slender and curved, much like that of a long-billed curlew, and it is fitted for probing in soft earth, or shallow water. The neck is round, and the head also, instead of being flat-sided like that of a heron.

**The White Ibis**<sup>3</sup> is yet found in Florida, and excepting the four outer wing-feathers (prima-

<sup>1</sup> *Ar-del'ta ex-i'lis*. Length, 13 inches.

<sup>2</sup> *Tan'ta-lus loc-u-la'tor*. Average length, 38 inches.

<sup>3</sup> *Guar'a al'ba*. Average length, 24 inches.

eris), which are black, it is a pure-white bird. Specimens in the first year are grayish-brown and white, and in color do not even suggest the pure-white plumage of the second year, and thereafter. This species rarely comes into any of the northern states.

The beautiful and brilliant **Scarlet Ibis**,<sup>4</sup> once a habitant of southern Florida and Louisiana, is no longer found in the United States. In color it is one of the most brilliant birds in all America, though by no means so beautiful as the resplendent trogon. I saw it in great numbers on the mud-flats at the mouth of the



WHITE IBIS.

Orinoco, and shot it on the coast of British Guiana. On Marajo Island, in the delta of the Amazon, it breeds in hundreds—a sight worth a long journey to see. Unfortunately, it is impossible to keep specimens of this species in confinement and have them retain their color. In a few months they fade until they are pale pink.

**The Glossy Ibis**<sup>5</sup> is a dark-colored bird, its prevailing color being rich brownish-purple with metallic-green reflections, and abundant iridescence. It seems smaller than the two light-colored species mentioned above, but in reality it is not. In 1899 two specimens were captured on the St. Johns River, opposite Melbourne, Florida, and one of them lived two years in the Zoological Park. This species is rare, even in Florida, but in Texas and the Southwest the *White-Faced Glossy Ibis* is of more frequent occurrence.

<sup>4</sup> *Guar'a ru'bra*. Length, 23 inches.

<sup>5</sup> *Pleg'a-dis au-tum-nal'is*. Length, 23 inches.

## THE SPOONBILL FAMILY.

*Plataleidae.*

**The Roseate Spoonbill,<sup>1</sup> or Pink "Curlew,"** is the only member of the Spoonbill Family in America, and it is also the farthest from the type of the Order Herodiones. It is really an ibis with a wide bill which terminates in two rounded, flat plates, nearly two inches wide. When standing erect, it is about 16 inches high. Its body plumage is either rosy gray or white, and its wing coverts and secondaries are tinted a delicate and very beautiful rose-madder pink, the color being most intense on the lesser coverts.

Once quite abundant throughout the lagoons, streams and swampy districts of Florida, this beautiful bird is now so nearly extinct there that no live specimens have been obtainable nearer than the Gulf coast of Mexico. Indeed, until very recently there were good reasons for the belief that not one Roseate Spoonbill remained alive anywhere in Florida. Now, however, it is a pleasure to record the fact that this species has not wholly disappeared from our avifauna.

In *The Auk* for January, 1904, Mr. A. C. Bent describes the finding of a few small flocks of these birds near Cape Sable, which he found nesting in two localities. "The principal breeding-ground of the Roseate Spoonbills was a great morass on the borders of Alligator Lake, a few miles back from the coast near Cape Sable, where the mangrove islands in which the birds were nesting were well protected by impenetrable jungles of saw-grass, treacherous mud-holes, and apparently bottomless creeks. . . . The Spoonbills were here in abundance, and had eggs and young in their nests, in all stages, as well as fully grown young climbing about in the trees. The old birds were tamer than at Cuthbert Lake, and allowed themselves to be photographed at a reasonable distance.

<sup>1</sup> A-ja'i-a a-ja'i-a.

"The Spoonbills," continues Mr. Bent, "will probably be the next to disappear from the list of Florida water-birds. They are already much reduced in numbers and restricted in habitat. They are naturally shy and their rookeries are easily broken up. Their plumage makes them attractive marks for the tourist's gun, and they are killed by the natives for food. But fortunately their breeding-places are remote, and almost inaccessible; and through the earnest efforts of the A. O. U. wardens they are now protected. It is to be hoped that adequate protection in the future will result in the preservation of this unique and interesting species."

The nests found by Mr. Bent on Cuthbert Lake, almost on the edge of the Everglades, were built in red mangrove-trees on the edge of the water, all on nearly horizontal branches, from 12 to 15 feet from the ground. "They were well made, of large sticks, deeply hollowed, and lined with strips of bark and water-moss. One nest contained only a single, heavily incubated egg, one a handsome set of three eggs, and the other held two downy young, not quite half grown."

In my opinion, there is no "cause," either existent or creatable, not even the "cause of science," which could justify the killing or capture of any of the birds composing those last small flocks of Spoonbills. Not even the necessities of a zoological garden should for one moment be accepted as an excuse for meddling with that avian remnant; and let no hunter think of offering a bargain in live Spoonbills from Cape Sable, or of now writing to ask "What will you give?"

It is to be hoped that the people of Florida will see to it that the Spoonbill is absolutely protected for the next twenty years, with the same intelligent interest and humane reason that has saved the manatee down to 1903.



ROSEATE SPOONBILL.

## CHAPTER XXVII

# THE ORDER OF FLAMINGOES—A CONNECTING LINK

### ODONTOGLOSSAE

The long-legged, long-necked **Flamingo** is a very perfect connecting link between the wading-birds and the swimmers, and a most curiously formed bird. It has enormously long, stilt-like legs, like a heron; but its feet are fully webbed, like the feet of a duck. Its standing height is from 48 to 54 inches. It has a long, slender, crane-like neck; but its thick, broken-

In 1902, Mr. Frank M. Chapman, of the American Museum of Natural History, visited a great breeding-place of the **American Flamingo**,<sup>1</sup> on Andros Island, Bahamas, where he saw about 700 birds in one flock, and about 2,000 mud-nests in one group, situated on a level mud-flat close beside a shallow lagoon.

The nest of this queer bird proved to be a low, flat pillar of mud from 6 to 12 inches in height, 13 inches in diameter at the bottom, and 10 inches across the top—which is flat, and slightly depressed.<sup>2</sup> The eggs are two in number.

Up to 1890, the Flamingo flocks still visited southern Florida, near Cape Sable, and it is possible that at rare intervals they still do so. Besides those in the Bahamas, flamingoes are found in Cuba. Every year from twenty to fifty live birds are brought to New York by the dealers in live animals, and sold at prices ranging from \$12 to \$20 each. When they arrive they are all over bright red, but in captivity all gradually fade out until they are pale pink.

In all the world there are eight species of flamingoes. While our species is bright scarlet, all over, those of Europe and Africa are almost white, with pink wing coverts. The food of this bird in captivity is dried shrimps, boiled rice, and cubes of stale bread, fed in water. In a room which is warmed to 60° F., it can live all winter, wading half the time in water that is almost icy cold, without catching cold. The voice of this bird is fearfully and wonderfully made. It is a resonant, deep bass, utterly unmusical "*honk*," like a rasping blast on a large tin horn, blown by an amateur.

<sup>1</sup> *Phoenicopterus ruber*. Length, 45 inches; spread of wings, 62 inches; tarsus, 12.50 inches.

<sup>2</sup> *Bird Lore Magazine*. IV., p. 180.



New York Zoological Park.

THE FLAMINGO.

backed bill is provided with lamellae along the edges, like the bill of a shoveller-duck. The anatomy of the bill and tongue of this bird is particularly interesting.

This bird is by habit a true wader, and lives and breeds near shallow lagoons, where it can walk in the water, and feed on the bottom.

CHAPTER XXVIII

THE ORDER OF DUCKS, GEESE, AND SWANS

ANATIDAE

We have now reached the first Order of a great group of birds which might well stand as a Sub-class—the Web-Footed Swimmers. It embraces six different Orders, and before touching any one of them, it is highly necessary that the student should take a bird's-eye view of the whole subdivision. A clear conception of these six Orders, and the characters on which they are based, will be of great and perpetual service to every person who desires a comprehensive view of the avian world:

THE ORDERS OF SWIMMING-BIRDS.

<p><b>THE WEB-FOOTED BIRDS.</b></p>	}	<p><b>FLYING SWIMMERS:</b> with good wings.</p>	}	<p><b>Ducks and Geese</b> (three toes webbed). <i>AN-A-TI'DAE.</i></p>	<p><b>Fully Palmated Birds</b> (four toes webbed). Cormorants, Pelicans, Snake- Birds, etc. <i>STEG-AN-OP'O-DES.</i></p>
		<p><b>Tube-Nosed Swimmers.</b> Albatrosses and Petrels. <i>TU-BI-NA'RES.</i></p>	<p><b>Long-Winged Swimmers.</b> Gulls, Terns, etc. <i>LON-GI-PEN'NES.</i></p>		
		<p><b>DIVING SWIMMERS:</b> with small wings, or none for flight.</p>	}	<p><b>Weak-Winged Divers.</b> Loons, Grebes, Auks, Puf- fins. <i>PY-GO'P'O-DES.</i></p>	<p><b>Flightless Divers.</b> Penguins. <i>IM-PEN'NES.</i></p>

This group is not only extensive, but its members show a wide diversity in form and habits, and they are fitted for life in all climates, on waters great and small. Having before us such a host of swimming-birds that six Orders are necessary to classify them, it is difficult to select only a few examples, and resolutely exclude all others. However, the student who becomes permanently acquainted with about thirty-five web-footed birds specially chosen to represent these Orders, will have a very good foundation on which to build higher, with the aid of special books and specimens.

As heretofore, we will take up the selected examples in the order in which it is easiest for


the student to receive them,—the highest types first,—rather than in the very curious sequence adopted by the A. O. U., and most technical writers on birds.

Once a year, the grand army of birds of the Order *Anatidae* take wing, and sweep northward from the tropics and sub-tropics. Many halt in the temperate zone, where food is abundant, but many more press on to the arctic circle, and far beyond it. Wherever they pause for the summer, they nest and rear their young; and many pages might be filled with descriptions of the different kinds of nesting-sites and nests.

One would naturally suppose that in any civilized country, birds in flight to their breeding-

grounds, or in occupancy of them, would be immune from the attacks of gunners. In some states (of which New York now is one!) the laws prevent "spring shooting," but in others it does not. In view of the changes for the better that are being made year by year, it is best not to particularize; but it is surprising that in some states a prolonged fight should be necessary to secure laws prohibiting spring shooting! The need for absolute protection for birds while

Atlantic coast and the Mississippi valley, literally teem with roaring guns and flying shot, and to-day the wonder is not that the wild-fowl have become "so scarce," but rather that so many have escaped slaughter! In view of the enormous annual output of new gunners, guns and ammunition, nothing but the strongest kind of public sentiment for bird-protection, backed by stringent laws, rigidly enforced, can save the ducks, geese and swans of North Amer-



Female.

Male.

THE MALLARD DUCK.

they are breeding, or about to breed, is so imperative that it is difficult to see how any sensible and honest person can oppose the enactment of laws to provide it. The killing of wild-fowl in spring, or at any time during their breeding-season, should everywhere be made a penal offence.


During the autumn migration southward, the flocks run a gantlet of guns a thousand miles long. Whenever and wherever a duck or goose alights to rest and feed, the guns begin to roar. The more important migration routes, like the

ica from becoming as extinct as the great auk and the dodo.


To-day, we are advised that *automatic repeating shot-guns* are about to be put upon the market,—to hasten the total extinction of all our game-birds. Their manufacture, sale and use should be rigidly prevented by law.

North America is—or *was*—particularly rich in species of birds belonging to the Order *Anatidae*, and once was richly stocked with individuals. Even yet, a very interesting remnant remains. Of the whole assemblage of species,







FULVOUS TREE-DUCK.  
*Dendrocygna fulva.*



BLACK DUCK.  
*Anas obscura.*




GADWALL: GRAY DUCK.  
*Chauliasmus strepera.*




AMERICAN WIDGEON.  
*Mareca americana.*




GREEN-WINGED TEAL.  
*Nettion carolinensis.*




SCAUP DUCK.  
*Aythya marila.*



RING-NECKED DUCK.  
*Aythya collaris.*



BARROW'S GOLDEN-EYE.  
*Clangula islandica.*




OLD SQUAW.  
*Harelda hyemalis.*



HARLEQUIN DUCK.  
*Histrionicus histrionicus.*



SURF SCOTER.  
*Oidemia perspicillata.*



AMERICAN SCOTER.  
*Oidemia americana.*

great, medium and small, I think the **Mallard Duck**<sup>1</sup> is the highest type, and the best average. It is one of the largest ducks; it is one of the handsomest; it is strong on the wing, and highly intelligent. It is a joy unto the sportsman who finds it in its haunts, and a delight to the epicure who finds it upon the bill of fare. Sluggish indeed must be the pulse which does not beat faster at the sight of a flock of wild Mallards, free in its haunts, and ready to leap into the air and speed away at the slightest alarm.

The Mallard is recognizable by its large size, and the brilliant metallic-green head and neck, and pearl-gray body, of the male. The female is a very different-looking bird, of a modest brown color, streaked with black. There is only one thing at all annoying about this bird, and that



BLUE-WINGED TEAL.

is its close resemblance to our domestic duck; but for this there is a very good reason. It is the wild ancestor of all our domestic ducks, save one or two varieties.

The Mallard is found throughout the temperate zone in both the Old World and the New, and therefore it is known by many names. In

<sup>1</sup> *An'as boschas*. Average length, 22 inches.

England it is called the *Stock Duck*, because it was the original stock from which the domestic duck has descended. In North America its range covers practically the whole continent down to Panama, and in Asia it reaches to India. It breeds persistently throughout the greater portion of its immense range—in the long grass of pond margins; in the woods, between the spur roots of trees; and on the prairies, beside streams of the smallest size.

Once while collecting in Montana, late in May, I found a tiny water-hole, barely ten feet in diameter, hiding in the sunken head of a very dry coulee. For miles in every direction stretched a billowy sea of sage-brush, already shimmering in the heat of early summer. As I dismounted to scramble over the edge of the bank for a drink, up rose a Mallard Duck from her nest in a thick patch of sage-brush, within a yard of my feet.

The nest was the old, familiar type,—a basin of grass lined with a thick layer of down from the breast of the prospective mother, and a bunch of eggs that almost overflowed the boundaries of their resting-place. As I gazed in astonishment at this nest and its contents beside an insignificant bit of water in a landscape that certainly was not made for ducks, I understood how it is that this bird has been able to spread itself all around the northern two-thirds of the globe.

In captivity the Mallard is the best of all ducks, and the most persistent and prolific breeder. Put a flock on any pond having long grass or timber about it, keep away the rats, raccoons, mink, thieves, and other vermin, and each female will do her utmost to surround herself with a downy flock of about fifteen small Mallards, regularly every summer. In the Zoological Park, several nests have been built within twenty-five feet of walks that are in daily use by crowds of visitors, the immunity of their builders being due in each case to their wonderful color resemblance to the dead oak-leaves which surrounded them, and with which they almost covered themselves.

**The Blue-winged Teal**<sup>2</sup> represents with us a group of three species which contains the smallest ducks found in North America.

<sup>2</sup> *Quer-quad'u-la dis'cors*. Average length, 15 inches.

Throughout its home, which embraces the whole United States east of the Rockies, and also far north and far south, it is so common—and also so small—it is not highly prized by sportsmen, and its worst enemy is the sordid market-hunter. Like the other teal, it prefers quiet, inland waters to the wide expanses that back up from the sea.

All the teal are quick risers, and also speedy on the wing; but they are rather dull of sense, and easy to approach. The Blue-Wing is known by the conspicuous white crescent in front of and half encircling the eye, and the bright blue patch, called the "speculum," on its wing.

**The Cinnamon Teal<sup>1</sup>** is a cinnamon-brown bird of the western half of the United States, once common, but rapidly diminishing in numbers. This species is very difficult to keep long in captivity, being very sensitive to all adverse influences.

**The Green-Winged Teal<sup>2</sup>** has a very noticeable crest, and a beautiful emerald-green speculum on each wing. It is found scattered over practically the whole of North America, from the Atlantic to the Pacific, and from the Barren Grounds to Cuba and Honduras.

**The Shoveller,<sup>3</sup>** also called the **Spoonbill**, is a handsome duck, recognizable by its extremely broad and spoon-shaped bill—the broadest of any American duck. The head and neck of the male are either black, or dark metallic-green; and the body-colors are black, white, blue, and green, handsomely disposed.

The bill of this bird shows the limit of development in width, and the comb-like lamellae along the outer edges, which are designed for use in straining minute particles of food out of water, are very pronounced. These minute plates are set cross-wise at the edges of the mandibles, and perform the same function as the plates of hairy baleen, or "whalebone," in the mouth of a baleen whale. All the members of the Order *Anatidae* are provided with lamellated bills, as also are the flamingoes.

This fine duck is a bird of inland waters, and

<sup>1</sup> *Quer-quad'u-la cy-an-op'ter-a*. Average length, 16 inches.

<sup>2</sup> *Nel'ti-on carolinensis*. Average length, 13.50 inches.

<sup>3</sup> *Spat'u-la cly-pe-a'ta*. Average length, 19 inches.

appears to dislike salt water. It is found sparingly "pretty much everywhere throughout the northern hemisphere . . . but is not common in the eastern states, and breeds from Alaska to Texas." Its flight is much like that of a teal, but less swift, and in cruising about for good feeding-grounds it is irregular and hesitating. "The body of the Shoveller is not large, and its apparent size in the air is made up chiefly of wings and head. . . . As a bird for the table, I have held it in very high esteem." (D. G. Elliot.)



Male.

Female.

THE SHOVELLER-DUCK.

In captivity it is a difficult bird to acclimatize and keep alive, which for several reasons is to be regretted. The females and immature birds are colored very differently from the adult and perfect males. The following local names of this bird have been recorded by Mr. Elliot in his admirable book on "The Wild Fowl of North America": Blue-Winged Shoveller, Red-Breasted Shoveller, Spoonbilled "Teal," Spoonbilled "Widgeon," Broad-Bill, Broady, Swaddle-Bill and Mud-Shoveller.

I regard the **Pintail**, or **Sprigtail**,<sup>4</sup> as the most beautiful duck in America, not even excepting the wood-duck. On land its outlines are trim, graceful and finely drawn, and on the water it makes one think of a finely modelled yacht. In beauty of form it far surpasses all

<sup>4</sup> *Daf'i-la a-cu'ta*. Average length of male, 27 inches; female, 22 inches.


other American ducks; and nowhere among wild-fowl is there to be found a more charming color-scheme than in the plumage of the drake. It is a harmony of delicate drabs, grays and white, used to set off several pleasing shades of brown, black, and iridescent green. None of the colors are gaudy or cheap-looking, and as a whole the combination of form and colors produces a bird that is in every way an exquisite creature.

It is in recognition of its beauty that this

southward before the advance of snow and ice begins in September. On our Atlantic coast, many of the flocks winter in the labyrinth of sounds, bays and channels that fringe the coast of Virginia and the Carolinas.

During recent years, quite a number of these birds have been caught alive near Water Lily, North Carolina, which is a locality famous for its wild ducks, geese and swans.

Fortunately the Pintail is easily acclimatized, and although not a good breeder, like the



Female.

THE PINTAIL DUCK.

Male.

duck is sometimes called the *Water-Pheasant*. Its correct name, however, has been bestowed in honor of its 7-inch long, finely pointed tail.

This bird ranges over nearly the whole of North America, but its favorite breeding-grounds are in the subarctic regions, particularly in the Yukon valley, and in the lake regions of the Canadian Barren Grounds. It is equally at home on the fresh-water lakes and rivers of the interior, and the salt-water inlets and channels of the Atlantic coast. The annual migration

mallard, it does well in captivity, and is truly a thing of beauty, and a joy as long as it lives.

The beauty of the **Wood-Duck**, or **Summer-Duck**,<sup>1</sup> depends almost wholly upon its brilliantly colored plumage; for its form is quite commonplace. It may be wrong to make a cold-blooded analysis of its points, but for beauty of form, the neck of this bird is too small and too short, its head is too large, and its body is very ordinary. Its plumage, however, presents

<sup>1</sup>*Aix sponsa*. Average length of male, 19 inches.

a color-scheme of brilliant reds, greens, blacks, browns, yellows and whites which is quite bewildering. Even its weak little bill is colored scarlet and white, and its iris is bright red.

In my opinion the claims of the two duck species which are rivals for the prize for web-footed beauty may fairly be expressed by the following proportion:

The Pintail is to the Wood-Duck as a well-gowned American Woman is to a Chinese Mandarin.

The Wood-Duck needs no description. Among ducks it is equalled in gorgeous colors only by its nearest relative, the mandarin duck of China—a painted harlequin. Our species is a tree-duck, and not only perches on trees, but also makes its nest in them, and rears its young at an elevation of from ten to thirty or forty feet. The nesting-site is always above water, in order that as the ducklings finally scramble out of the nest and fall, they will alight in the water without injury, and quickly learn to swim.

In captivity the best nesting arrangement for this bird consists of a long, narrow box set on end on a stout post, well out in a pond, roofed over to keep out the rain. There must be a hole in one side, near the top, and a slanting board with cross slats reaching up to it from the water, for use as a ladder. The Wood-Duck will sometimes nest on the ground, either in captivity or out. This species is being bred in captivity in England in large numbers, and also with some success in this country. Duck fanciers find no difficulty in purchasing live specimens of this interesting bird at \$15 per pair.

During the summer of 1902, a pair of wild Wood-Ducks made daily visits to the Ducks' Aviary in the New York Zoological Park, and in the autumn of that year a small flock settled with the Wood-Ducks, mallards and pintails on the Aquatic Mammals' Pond, and remained there permanently. In the spring of 1903, a fine drake manifested a fixed determination to break into the great Flying Cage, and become a member of the happy family within. After he had flown around the cage two or three times, Keeper Gannon opened wide the wire gates at the north end, drove him in, and he is there now, serene and happy.

The Wood-Duck is a bird of great discernment.

Although this bird is called the Summer-Duck, and migrates far in advance of winter, it winters very comfortably in the northern states if it is fed and continuously provided with open water to keep its feet from freezing. The natural range of this species is from Hudson Bay to the Gulf of Mexico, chiefly on fresh water; but often it is found on brackish sounds and channels along the Atlantic coast where food is plentiful.

Like all other wild ducks that are impera-



WOOD-DUCK.

Male and female.

tively needed to keep the American people from starving, there remains to-day about one Wood-Duck where formerly there were from thirty to fifty. Apparently, the only winged creatures that are too beautiful or too good to be shot and eaten are angels; but I doubt if even a white-winged seraph with webbed feet would be safe for half an hour anywhere between Cape Cod and Charleston.


**The Redhead Duck**<sup>1</sup> is one of our largest and best species, and one of the most satisfactory to keep in captivity. It belongs to the same genus as the canvas-back, and in size, habits, table value and beauty it is in no sense

<sup>1</sup>*Aythya americana*. Average length, 19 inches

whatever inferior to its more famous relative. When shot in the same locality, I think there is no one who could distinguish the two species by a difference in the flavor of their flesh.

In the color of their plumage, the Redhead and canvas-back look so much alike that the casual observer might easily mistake one species for the other. Both have heads and necks of solid rusty brown, but the head-color of the Redhead is the more intense and conspicuous.

The head of the Redhead has a *high and well-rounded forehead and crown*, while that of the canvas-back is *wedge-shaped*, the forehead forming a *straight line with the top of the bill*. The Redhead has a short bill, with a blue band across it; the other species has a long bill, with no band.



THE REDHEAD DUCK.

The Redhead (like the canvas-back) feeds chiefly upon aquatic plants, its favorite food being the *vallisneria*, a kind of trailing water-weed which grows in many of the inlets along the Atlantic coast.

Through countless generations of diving after food-plants, the Redhead has become a deep diver. It is accustomed to seeking its food in mid-stream of deep rivers, and in the open water of lakes and sounds, where many other ducks would be quite unable to reach the bottom. Reliable lake fishermen at Lakeside, Orleans County, New York, have informed me that they have taken drowned Redhead Ducks from nets that had been set on the bottom of Lake Ontario, at a depth of *ninety feet*, where the

ducks could not possibly have become entangled save in going to the bottom for food. It also appeared that those Ducks sought their food and became entangled only at night. It takes a bold and energetic bird to feed successfully *at night* in ninety feet of water!


Naturally this fine bird has ever been a prime favorite with sportsmen and "market-shooters," and during the past fifteen years its numbers have diminished to about one-fiftieth of what they were prior to 1885. It is as easily deceived by decoys as green hunters are; and in preparing to alight the Redhead flock has a fatal habit of coming together in a manner called "bunching," which is as deadly to the birds as "close formation" is to soldiers in a modern battle.

Much more might be noted regarding this interesting bird, which must be left to the special works on birds. For many reasons it is very desirable that the Redhead should be semi-domesticated, and by protection and breeding in captivity saved from the final blotting out which otherwise will be its fate. While it does not breed in captivity as bravely as the mallard, it *can be taught* to do so, and the prices at which living birds can be procured (\$5 each) is so very moderate that experiments with it are not costly.

The distribution of this bird is given as "North America, breeding from California, southern Michigan, and Maine northward;" but in North America there are to-day more lands and waters without this duck than with it. In addition to its best and most appropriate name it is also called *Raft-Duck*, and *American Pochard*.

**The Canvas-Back Duck**<sup>1</sup> had the misfortune, early in its history, to attract the evil eye of the deadly epicure, whose look of approval is a blighting curse to every living creature upon which it is bestowed. Because of this, the unfortunate Canvas-Back is now little more than a bird of history. It is of no present interest, outside of museums and the zoological parks and gardens which have been so fortunate as to secure a very few specimens. Unfortunately, it has been impossible for even the most energetic duck-fanciers to secure a sufficient number

<sup>1</sup>*Aythya valis-neria*. Average length, 22 inches.



THE CANVAS-BACK DUCK.

of unwounded specimens to carry out the experiments necessary to determine the precise conditions under which this species will breed in captivity. No one ever sees more than two or three living Canvas-Backs together in an aviary, and thus far I believe none have bred.

It is unnecessary to describe this species, for it is probable that no one of the readers hereof ever will see one wild and unlabelled. Its range was once the same as that of the redhead, and its habits also were quite similar.

**The Buffle-Head Duck, or Butter-Ball,**<sup>1</sup> is a small, tree-nesting duck, so pretty and so very odd-looking that when seen every one wishes to know its name; and when named, it is not soon forgotten. When you see a short-bodied, plump-looking little duck, black above and white below, with a head that is a great round mass of soft feathers, half snow-white, and half a rich metallic mixture of purple, violet and green,—that is a Butter-Ball, and nothing else. Wherever seen, it commands instant attention.

Unfortunately, this picturesque little creature does not like our country as a summer residence, for it breeds from Maine, Iowa and British Columbia, northward, and returns to us only when snapping cold weather heralds the approach of winter. On the water it is the most nervous and watchful duck that I know, and its habit of constantly turning from side

<sup>1</sup> *Char-i-ton-el'ta al-be-o'la*. Average length, 14.50 inches.

to side is certainly in the interest of self-preservation. But after all, what is the alertness of any duck against the deadly, cold calculation of the greedy "market-shooter" with a choke-bore gun?

The Buffle-Head is one of the ducks that is rarely seen in captivity. A specimen that is so seriously wounded that it can be caught, usually dies a few days later. So far as I know, it has not yet been induced to breed in captivity; but that is no reason for believing that it never will. We hold that if conditions are made satisfactory, any wild species will breed in captivity. Usually it is a question of sufficient seclusion, and immunity from disturbance. The range of this bird is said to include all North America, from the Arctic Ocean to Cuba. And so it does, all save those localities wherein it does *not* occur.



THE BUFFLE-HEAD, OR BUTTER-BALL.

**The Harlequin Duck**<sup>2</sup> is most fantastically marked. The prevailing colors of the male are dark blue, blue-black and violet, with various white collars, stripes and patches that seem to have been laid on with a paint-brush. This bird is to be looked for along the Pacific coast above Oregon to Japan, and on the Atlantic coast from Newfoundland northward. It is nowhere common, rather solitary, but frequents costal rivers as well as the sea. As a rarity to be prized, one Harlequin is equal to twenty ducks of almost any other species in America.

<sup>2</sup> *His-tri-on'i-cus his-tri-on'i-cus*. Length, 16 inches.



Reproduced from *Recreation Magazine*.

A HAVEN OF REFUGE.

**An Object Lesson in Bird-Protection.**—As a fitting conclusion to our studies of the ducks of our interior rivers, lakes and ponds, we present a remarkable instance of what bird-protection can accomplish. The picture of the pond described might well be entitled—"An Oasis in the Great American Desert of Game Destruction!" By the courtesy of Mr. G. O. Shields, Editor of *Recreation Magazine*, we reproduce from that periodical for June, 1903, the above illustration, and the following description by Mr. Charles C. Townsend, which appeared under the caption, "A Haven of Refuge."

"One mile north of the little village of Mosca, Colorado, in San Luis valley, lives the family of J. C. Gray. On the Gray ranch there is an artesian well which empties into a small pond about 100 feet square. This pond is never entirely frozen over and the water emptying therein is warm even during the coldest winter.

"Some five years ago Mr. Gray secured a few wild-duck eggs, and hatched them under a hen. The little ducks were reared and fed on the little pond. The following spring they left the place, to return in the fall, bringing with them broods of young; also bringing other ducks to the home where protection was afforded them, and plenty of good feed was provided. Each year since, the ducks have scattered in the spring to mate and rear their families, returning again with greatly increased numbers in the fall, and again bringing strangers to the haven of refuge.

"I drove out to the ranch November 24, 1902, and found the little pond almost black with the birds, and was fortunate enough to secure a picture of a part of the pond while the ducks were thickly gathered thereon. Ice had formed around the edges, and this ice was covered with ducks. The water was also alive with others,

which paid not the least attention to the party of strangers on the shore.

"From Mr. Gray I learned that there were some 600 ducks of various kinds on the pond at that time, though it was then early for them to seek winter quarters. Later in the year, he assured me, there would be between 2,000 and 3,000 teal, mallards, canvas-backs, redheads and other varieties, all perfectly at home and fearless of danger. The family have habitually approached the pond from the house, which stands on the south side, and should any person appear on the north side of the pond the ducks immediately take fright and flight. Wheat was strewn on the ground and in the water, and the ducks waddled around us within a few inches of our feet to feed, paying not the least attention to us, or to the old house-dog which walked near.

"Six miles east of the ranch is San Luis lake, to which these ducks travel almost daily while the lake is open. When they are at the lake it is impossible to approach within gunshot of the then timid birds. Some unsympathetic boys and men have learned the habit of the birds, and place themselves in hiding along the course of flight to and from the lake. Many ducks are shot



in this way, but woe to the person caught firing a gun on or near the home-pond. When away from home, the birds are as wild as other wild-ducks and fail to recognize any members of the Gray family. While at home they follow the boys around the barn-yard, squawking for feed like so many tame ducks.

"This is the greatest sight I have ever witnessed, and one that I could not believe existed until I had seen it. Certainly it is worth traveling many miles to see, and no one, after seeing it, would care to shoot birds that, when kindly treated, make such charming pets."



AMERICAN EIDER.

**The Group of Eider-Ducks.**—The arctic and subarctic regions contain a group of about seven species of large sea-ducks, called eiders (i'ders). The representative species are distinguished by their flat foreheads and wedge-shaped heads; by a long, wedge-shaped point of the cheek-feathers which extends forward and *divides the base of the upper mandible*; and by the possession of more or less bright green color on the head.

On land, the eiders are heavy and clumsy birds, but on the sea they are at home, and dive with great ability. The females line their nests very liberally with down from their own

breasts, and this when gathered and utilized becomes the well-known "eider-down" of commerce. Unfortunately, the natives of arctic America are unable to make use of eider-down, save on the skin, and this leads to the slaughter of great numbers of the birds.

Eiders nest on the tops of rocky islets, using sea-weed or grass for a foundation, and covering this with down plucked from their own breasts. So abundantly is the nest lined that by the time the eggs are all deposited they are fairly embedded and covered in the softest of beds. In Iceland, the eider-ducks are half domesticated. The inhabitants collect the down from the nests for sale, and therefore they are much interested in preserving the birds. Nesting-places are made for the birds by building thick stone walls with spacious crevices along each side, at the base, or by scooping out shallow cavities in the hard earth. The Eiders permit their human friends to go among them, and even to handle their eggs.

On the Atlantic coast, from Labrador to Delaware in winter, we have the **American Eider**,<sup>1</sup> which appears to be the best type for the eider group. Fortunately for our chances of close acquaintance with it, this species occasionally penetrates westward along the great lakes to Illinois and Wisconsin—a very unusual proceeding for a sea-duck. Any bird which will go so far out of its natural range in order to become acquainted with interocean Americans surely is worth knowing. Moreover, the eider of the Old World so closely resembles this bird in all essential details that to know one species is to know the other also.

The colors of this bird are black and white, as shown in the illustration, except that the nape and the rear portion of the region around the ear are sea-green, and the tail and the primaries are pale brown. The bill and feet are olive-green.

**The Spectacled Eider**,<sup>2</sup> of northwestern Alaska, is a bird easily remembered by its name, and the large, white spot around each eye which at once suggests a pair of spectacles. This bird is limited to our arctic territory, and is said, by Mr. E. W. Nelson, to be threatened with extinction by man at no very distant day.

<sup>1</sup> *So-ma-le'ri-a dres'ser-i*. Length, about 23 inches.

<sup>2</sup> *Arc-ton-et'ta fisch'er-i*. Length, about 21 inches.

Our occupation of Alaska, after the Russians, has led to the arming of the natives with modern rifles and shot-guns, before which wild life generally is rapidly being swept out of existence.

**The White-Winged Scoter**<sup>1</sup> (sko'ter) quite acceptably represents a group of sea-ducks and deep divers, called Scoters, and of which there are three species resident in North America. These are the blackest of all our ducks. The species known as the **American Scoter**

southern California, northern Missouri, Illinois and Maryland. Like most of our ducks, it breeds in the far north, and returns to us only for the winter. It is a deep and persistent diver, and it is said that when wounded and pursued it will sometimes dive to the bottom, even fifty feet if necessary, seize a bunch of grass or weeds with its bill, and hold on until it has quite drowned. Its food consists of fish, crustaceans and mollusks.



Female.

Male.

RED-BREASTED MERGANSER.

is glossy black throughout, without a single patch of color save the bright orange-yellow which colors the basal half of the bill and its knob.

The White-Winged species has a white patch on each wing, technically known as a "speculum," and a white patch of variable shape under or in rear of the eye. Above and in rear of the nostrils the bill and skull together are raised into a conspicuous hump, half covered by feathers.

Like all the scoters, this bird is a fish-eating duck, and its flesh is so fishy in flavor it is not considered fit for the table. It is widely distributed throughout North America down to

<sup>1</sup> *Oi-de-mi-a deg-land'i*. Average length, 21 inches.

**The Red-Breasted Merganser**<sup>2</sup> bravely and handsomely represents what is structurally the lowest group of ducks, known as the Mergansers, embracing three species. The bill of this bird is long, narrow, and set along the edges with lamellae that look quite like sharp teeth—a most admirable arrangement for seizing fish under water. The bill of a Merganser always reminds me of two things: the jaws of the gavial, or Gangetic crocodile, and Professor Marsh's toothed bird, the *Hesperornis*, from the great extinct inland sea of the Middle West. One of the common names of this bird is the *Saw-Bill*; and it is peculiarly appropriate.

<sup>2</sup> *Mer-gan'ser ser-ra'tor*. Average length, 22 inches.

Among other ducks this fine bird has the bold, confident air of a born free-booter. The back of its head is ornamented with several long feathers which form a crest, like the war-bonnet of a Sioux Indian. The whole head and upper neck are black, with green and purple reflections. Around the middle of the neck is a conspicuous white collar, and under that is the pale rusty-red breast, streaked with black, which gives the bird its name.

nervous, and difficult to keep alive in captivity. A fine specimen which we cherished for a time in the Flying Cage of the New York Zoological Park, along with many other water-birds of good size, at first seemed inclined to accept the situation, and become acclimatized; but it lived only two months. With several Mergansers together, the result might be more satisfactory.


The Hooded Merganser<sup>1</sup> is distinctly



STELLER'S DUCK.  
*Eniconetta stelleri.*



SPECTACLED EIDER.  
*Arctonetta fischeri.*



KING EIDER.  
*Somateria spectabilis.*



RUDDY DUCK.  
*Erismatura jamaicensis.*



AMERICAN MERGANSER.  
*Merganser americanus.*



HOODED MERGANSER.  
*Lophodytes cucullatus.*

This sea-going bird-craft is at home—under many names—in both the Old World and the New. On our continent it breeds from our northern states as far as the Aleutian Islands and western Alaska, where the Aleuts prize it for food above all other ducks. In winter it migrates along our two ocean coasts to southern California and Florida. It feeds entirely on fish, and the flavor of its flesh is rank and disagreeable.

Nearly all sportsmen admire this duck, and it is much to be regretted that it is so shy and

marked by a striking, black-and-white semi-circular crest of great height, standing stiffly erect, and jaunty beyond compare amongst water-fowl. By that crest and the slender Merganser bill any one may know this bird out of ten thousand species, whether seen in New York or New Zealand. It ranges all over North America, wherever there is water enough to float it, down to Mexico and Cuba, and as a result has been burdened with an appalling

<sup>1</sup> *Lo-phod'y-tes cu-cul-la'tus*. Average length, 17 inches.

collection of names. It nests in hollow trees, near good fishing-grounds, and whenever it makes its summer camp near a trout stream, the fry fare badly.

**The Geese.**—Those who have not looked into the subject usually are surprised to find what a fine collection of geese is found in North America. The continent is so large it requires an effort to come in touch with representatives of all the species of wild-geese which inhabit it. While they are somewhat lacking in the fine coloring that characterizes a few foreign species, such as the spur-winged goose of Africa,



CANADA GOOSE.

they form, as a whole, a highly interesting group, well worth the acquaintance of all Americans save the market-hunters, and others who shoot not wisely but too well.

Fortunately for those who live where wild-geese dare not show themselves for fear of being killed, all these species take kindly to captivity, and are easily kept in parks and zoological gardens. In 1903, five species were living quite contentedly in the New York Zoological Park.

In writing of geese, we would not think of mentioning any species ahead of our old favorite and most faithful friend, the **Canada**

**Goose.**<sup>1</sup> Where is the country dweller who has not heard, far aloft, the well-known trumpet "Honk," and the prompt answers all down the two lines as the V-shaped flock winged swiftly forward? In the raw, windy days at winter's end, from the Gulf to Hudson Bay, the old gander's cry is accepted as a guarantee of spring, and hailed with joy. Dull indeed is the mind that is not moved to wonder and admiration by the remarkable V-formation in which the wild-geese flock cleaves the air.

Although wild-geese in transit through the Mississippi valley frequently alighted in cornfields to rest and feed, as a rule they were so wary and wide-awake it was next to impossible to bag one. In Minnesota and the Dakotas, however, they often flocked on the ground in such numbers that goose-shooting was as regular a sport as chicken-shooting, and during a brief period of slaughter yielded weighty results. Thousands of geese alighting in cornfields to feed have been shot from the interior of innocent-looking corn-shocks.

The Canada Goose is not only the largest of the wild-geese of North America, but also the most important and valuable member of the group. There are times, also, when it seems to be the most savory bird that finds its way to the platter. One of those times was when a flock alighted near our camp, on the ice of the Musselshell, in Montana, the day before a certain whizzing cold Thanksgiving, and a fat young gander was shot, and beautifully roasted over the camp-fire in a large Dutch oven.

In captivity the Canada Goose is an all-around philosopher; and even when wild, he often knows a good thing when he sees it. In October, 1901, a flock of nine geese flying southward over the New York Zoological Park suddenly espied our flock of the same species on the Aquatic Mammals' Pond. Without a moment's hesitation, the wild-birds sailed down and alighted on the shore beside their relatives, and invited themselves to the banquet of cracked corn.

On the following day, Mr. H. R. Mitchell coaxed seven of the visitors into a huge wire cage that was set up on the shore, where they

<sup>1</sup> *Bran'ta canadensis*. Average length, about 35 inches; but individuals vary greatly in size.

were caught and wing-clipped to prevent further wandering into danger. The seven are still there; but the two unclipped birds, after remaining all winter, flew away north the following spring, and it is quite likely that their bad judgment has ere now cost them their lives.

Apparently, all the North American geese are almost as easy to keep in captivity as domestic geese. Their favorite food is cracked corn and whole wheat, but they will eat almost any kind of grain. In winter they require low shelter coops, open toward the south; and a small portion of their pond must be kept open all winter, by frequently removing the ice, to keep their feet from freezing. Not all these birds, however, care to seek shelter in a humble coop.

The Canada Goose is known by its large size, and its jet-black head and neck, with a conspicuous white crescent encircling the throat. The black on the neck ends abruptly where the neck joins the body, and the general tone of the latter is gray-brown. Its neck is longer, and also more slender as a rule, than those of other birds.

This fine bird winters in Texas, along the Gulf of Mexico, and in the sounds and bays of Virginia and the Carolinas, and goes north early in spring. Its nesting-grounds begin in our northern tier of states, and extend northward to Labrador, the Barren Grounds and Alaska. Throughout much of that vast area, the shot-guns and rifles are ever ready, and the number of geese that still survive are eloquent testimony to the wariness, the keenness of vision and the good judgment of this much-prized bird. A bird of equal desirability, but with a dull brain and poor vision, would have been exterminated long ago.

One of the most interesting things about the Canada Goose is the energy and courage of the male in defending the female on her nest. Last spring two of our geese paired off, and built a nest on the south bank of the Mammals' Pond, in a very exposed situation. From that time until the young were hatched, the gander never once wavered from his post. It was his rule never to go more than sixty feet from the nest, and whenever any one approached it, he immediately hastened to intercept the intruder, hissing and threatening with his wings

in a most truculent manner. Had any one persisted in disturbing the female, he would willingly, even cheerfully, have shed his blood in her defence. His unswerving devotion to his duty attracted the admiring attention of thousands of visitors, and the proudest day of his life was when the first live gosling was led to the water, and launched with appropriate ceremonies.

There are three subspecies of the Canada Goose, all smaller, but otherwise very similar. The **White-Cheeked Goose** inhabits the Pacific coast, north to Sitka; and the **Cackling Goose** is found in the same region, and on up to the Yukon. **Hutchin's Goose** is merely a small edition of the Canada.

**The Black Brant**<sup>1</sup> is a very distinct bird, noticeably smaller than the Canada goose, and readily recognized by its blackness and its small size. Its head, neck, and breast are entirely black, save for a white collar going two-thirds of the way around the upper neck. The black of the neck does not end abruptly at the shoulders, but spreads back over the back and under parts until the final effect is that of a bird which is two-thirds black.

Although this bird is generally accounted rare on the Atlantic coast, the New York Zoological Society has secured a number of fine living specimens from Carrituck Sound, on the coast of North Carolina. Beyond doubt, however, it is rare everywhere in the eastern United States. It is remarkable for the fact that it migrates northward not only to the desolate shores of the Arctic Ocean, but far beyond, and must nest and rear its young far out on the great polar ice-pack.

**The Brant Goose**<sup>2</sup> is quite a different species from the preceding. The black of its neck ends abruptly at the shoulders, and the white collar is a mere broken patch, without decided character. The body is everywhere much lighter than the color of the black brant, with which this species is often confounded, because the two are often found together, though not on the Pacific coast. Once the Brant Goose was plentiful along the Atlantic side, but it is now rare, and fast disappearing.

<sup>1</sup> *Bran'ta ni'gri-cans*. Average length, about 24 inches.

<sup>2</sup> *Bran'ta ber'ni-cla*.

**The American White-Fronted Goose<sup>1</sup>** is, in my opinion, the most handsomely colored goose we possess. Contrary to expectations that are often based upon its name, it has not a white breast, nor white shoulders. Its white "front" is limited to an inch-wide frill of white immediately surrounding *the base of its bill*.

arrangement of the plumage, and as a whole the bird is decidedly beautiful.

This fine bird is even yet abundant on the Pacific coast, from southern California to Alaska, where it crosses over to the Asiatic side. It appears that Alaska is its favorite nesting-ground. On the Atlantic coast it is no longer



New York Zoological Park.

TRUMPETER SWANS.

Other than this the head and the neck are dark brown, and the back, sides, breast and abdomen are covered with a scale-like arrangement of feathers that are various shades of brown or black, strongly edged with white or gray. The effect of the white edges of the feathers is to bring out in strong relief the immaculate ar-

<sup>1</sup> *An'ser al'bi-frons gam'bel-i*. Average length, 28 inches.

seen. The specimens living in the Zoological Park were taken in southern Texas, on the Rio Grande, where the species is yet a winter visitant.

**The Snow-Goose<sup>2</sup>** is, excepting its large wing-feathers (the primaries), an all-white bird. Based on the tape-line, two species have been

<sup>2</sup> *Chen hy-per-bo're-a*. Average length, about 30 inches.

described and recognized by ornithologists, the "Greater" Snow-Goose, and the "Lesser." If the specimen under the tape is a large one, it is the former species; but if it is smaller than the average, it is booked as the "Lesser." Obviously, the wisest course is to discard both adjectives of size, and recognize the Snow-Goose only, be it more or less.

This easily recognized bird, like the majority of our other wild geese and ducks, wanders over almost the whole of the well-watered portion of North America down to Cuba and Mexico; but where the guns of civilization are most numerous it is now a rare and lonesome bird. To-day it is more abundant—or it were better to say, *less scarce*—in the Mississippi valley, Texas, and the Pacific states than elsewhere. Where they were permitted to do so, these birds often assembled in large flocks, and often made themselves conspicuous around the prairie-ponds of the Dakotas and Minnesota. When you are travelling over the Northern Pacific Railway, or the Great Northern, and see on the smooth prairie a flock of rather large white birds, it is safe to declare that they are Snow-Geese.

**The Swans.**—Last of the Order of Ducks, and farthest from the type of the Order, are the Swans. Although two species are recognized, the difference between them is not always visible to the naked eye.

**The Trumpeter Swan**<sup>1</sup> is one of our largest birds, and considering its great size it is strange that it has not been exterminated ere this. Its existence speaks highly for its wariness. Living specimens are purchasable at from \$20 to \$30 each, and the majority of them come from Texas and the plains region. To my mind, this is the least attractive of all the large swimming-birds, and it certainly is one of the most pugnacious and quarrelsome. In captivity, Trumpeter Swans always wish to do the wrong thing. Even when policy demands that they at least appear friendly, they are always truculently hissing at and threatening their human neighbors, friends as well as enemies. This Swan's voice is like a short blast

<sup>1</sup> *Olor buc-cin-a'tor*. Length, 4 feet 8 inches; height, when standing erect, 3 feet 9 inches; expanse of wings, 7 feet 10 inches; weight, 22 pounds.

on a French horn, but when a large flock rises from a pond in a wilderness, and gets fairly under way, the chorus given forth on such occasions I know to be decidedly musical, and also heart-breaking when out of range.

With birds smaller than themselves, Swans often are so quarrelsome and murderous they require to be separated, and yarded by themselves.

On level ground, the Swan is the most ungainly of all the American members of the Order of Ducks; and even afloat, its bows lie much too deep in the water.

The central line of migration and distribution of this species is the western boundary of the states forming the western bank of the Mississippi. It breeds from Iowa northward to the Barren Grounds, and in the United States straggles eastward and westward to both shores of the continent. I have seen specimens taken in 1885 in the Potomac River, and it has often been observed near Los Angeles, southern California.

Thus far, only one naturalist (so far as we know) ever has heard the "Song of the Dying Swan." Mr. D. G. Elliot, in "Wild Fowl of North America," records the following interesting observation:

"Once, when shooting in Carrituck Sound, . . . a number of Swan passed over us at a considerable height. We fired at them, and one splendid bird was mortally hurt. On receiving his wound the wings became fixed, and he commenced at once his song, which was continued until the water was reached, nearly half a mile away. I am perfectly familiar with every note a Swan is accustomed to utter, but never before nor since have I heard any like those sung by this stricken bird. Most plaintive in character, and musical in tone, it sounded at times like the soft running of the notes in an octave."

**The Whistling Swan**<sup>2</sup> is accorded rank as a species chiefly on the strength of a small yellow patch on the base of the bill—which is not always present! Young Swans of both species are of a dirty-gray color—not white; but the plumage of the adult bird is perfectly white. The bill and feet are jet black.

<sup>2</sup> *Olor co-lum-bi-an'us*.

## CHAPTER XXIX

# THE ORDER OF FULLY WEB-FOOTED BIRDS

### STEGANOPODES

To recognize a member of this Order, look at its foot, and see that the web of the three large toes *is also united to the fourth, or rear toe*. This may seem like a small peg on which to hang an Order; but it is a very useful one, nevertheless. As usual, the best and most conspicuous example will be mentioned first. The Families are as follows:

	FAMILIES.	EXAMPLES.
<b>ORDER STEGANOPODES.</b>	PELICANS, . . . . . <i>PEL-E-CAN'I-DAE.</i> , . . . . .	} Brown Pelican; White Pelican.
	CORMORANTS, . . . . . <i>PHAL-A-CRO-CO-RAC'I-DAE.</i> , . . . . .	} Common Cormorant.
	DARTERS, . . . . . <i>AN-HING'I-DAE.</i> , . . . . .	} Darter, or Snake-Bird.
	GANNETS, . . . . . <i>SU'LI-DAE.</i> , . . . . .	} Common Gannet.
	MAN-O'-WAR BIRDS, <i>FRE-GAT'I-DAE.</i> , . . . . .	} Frigate-Bird.



Photographed by C. WILLIAM BEEBE

From the *Zoological Society Bulletin*.

FLORIDA BROWN PELICANS, ON PELICAN ISLAND.

### THE PELICAN FAMILY.

*Pelecanidae.*

The Brown Pelican<sup>1</sup> is known to every tourist who knows Florida thoroughly, or <sup>1</sup>*Pel-e-ca'nus fus'cus*. Length, 49 inches; spread of wings, 6 feet 9¼ inches.

southern California. Somehow this bird appeals to every one,—possibly by reason of its cheerful confidence in man,—and for a wonder it has not been exterminated. It takes to captivity not only willingly, but gladly, and its motto is, "All's fish that cometh to net."



It is an amiable bird, sociable to an unlimited degree, harms no one, and makes no enemies.

Pelican Island, in Indian River, Brevard County, Florida, is the most interesting sight in the land of flowers. On an area of about three acres, raised only two or three feet above high-water mark, destitute of trees because the Pelicans have nested them to death, live about 2,000 Brown Pelicans, and in 1902 they made 976 nests. During every breeding-season they

babies, as large as their parents, but covered all over with down as white as cotton.

It is no uncommon thing for a young Pelican to have from six to nine mullet in its neck and crop at one time, as we have discovered by catching some of them with a search-warrant, and searching their premises.

To feed these hungry and appallingly capacious pouches, the old birds fly about fifteen miles up the coast to fishing-grounds where silver



Photographed by R. J. BECK.

CALIFORNIA BROWN PELICAN.

Galapagos Islands.

inhabit that islet, nesting in small nests of grass plucked on the spot, and arranged on the ground. The few dead mangroves that still stand are loaded with stick-made nests, to the point of breaking down.

Egg-laying begins about the first of February, and straggles along until the end of May. By March 15, the breeding-grounds contain in close proximity, unfinished nests, and nests with fresh eggs (usually three); young just out of the shell; half-grown young, and, finally, full-grown young. The latter are great hulking

mullet are plentiful and cheap; and there each old bird fills its neck and crop with from six to nine fish, each from seven to ten inches in length. At evening, just before sunset, in groups of from three to seven they slowly wing their way back along the beach, flying low over the saw palmettos that fringe the shore. They give about six wing-beats, then sail as far as possible, each little company winging in unison. Several times I have lain low in the palmettos, to watch their flight at a distance of only a few feet as they approached and passed over me.

Truly they are fine birds,—rich in coloring, remarkably odd in form, and very well set up. Unfortunately they do not acquire their full colors until in their third year. The neck of the adult bird is in two colors, rich blackish-brown and white, and the back is a beautiful silvery gray-brown effect, composed of many tints. The top of the head of the adult bird is yellow. The bill is a foot long, the pouch is of a bluish-purple

of flying spray tells the story of the plunge with open pouch for the luckless fish.

For several years the fate of the great Pelican colony in Indian River has been in doubt, and its preservation has been due more to public sentiment in Brevard County than to the arm of the law. In 1903, however, Pelican Island was formally declared to be a government reservation, and placed under the absolute control of the Biological Survey, thus insuring the permanent protection of its occupants.

**The California Brown Pelican<sup>1</sup>** so closely resembles the Florida species that the differences between the two are not easily recognized. The accompanying illustration is from a photograph taken on the Galapagos Islands, directly under the equator; and from that locality this species ranges northward along the Pacific coast to British Columbia.

**The Great White Pelican<sup>2</sup>** is a grand bird,—big, clean, immaculate, and with the dignity of a newly appointed judge. About him there are two bad things. In captivity his appetite for fresh fish makes him a costly luxury, and his Latin name always frightens timid people.

The curious horn seen in winter and spring atop of the bill of this bird is purely a sexual ornament, found only on the male in the breeding-season, after which it drops off. It begins to grow about February 15, is perfect by May 1, and drops off not later than July 1.

To-day, as a matter of course, the Great White Pelican is a rare bird. On the west coast of Florida, where once it was abundant, I believe it is no longer found. It is yet found inland in certain western localities, where there are lakes large enough to shelter it, and supply it with fish, and it is to be hoped that it will be many years ere this grand bird is exterminated. Fortunately, a colony has become established on an island in Yellowstone Lake, in the Yellowstone Park, where it breeds regularly every summer, to the great delight

<sup>1</sup> *Pel-e-ca'nus californicus*.

<sup>2</sup> *Pel-e-ca'nus er-yth-ro-rhyn'chos*. Length, 61 inches; spread of wings, 8 feet 10 inches; weight, 16½ pounds.



New York Zoological Park.

GREAT WHITE PELICAN.

color, and calls for about four pounds of fish daily.

It is very interesting to watch Pelicans fishing. On calm days when the surface of Indian River is like a mirror, the eruption of silvery spray that rises high when the big bird plunges into the water, attracts attention at a distance of two or three miles. It is finest, however, to see them fishing in the breakers on the ocean side of the Indian River Peninsula, about 200 feet from shore. They sail along so near the water it seems a wonder it does not strike them; but they rise over the incoming waves, and lower again into the trough with the utmost precision, always keenly alert. All of a sudden, the wings are thrown out of gear, and a fountain

of all tourists who care for the sight of what is called a "pelicanery." In winter, southern Texas is the haven for this bird, as well as for so many other swimming-birds.

### THE CORMORANT FAMILY.

#### *Phalacrocoracidae.*

**The Cormorant**<sup>1</sup> is to me a most uninteresting bird. Month in and month out I have seen them perching, and perching,—on spar buoys in harbors, on mud-bank stakes, and on dead trees along shore and up stream. For days together have Cormorants fled up stream before my boat, yet never once have I seen a wild Cormorant do an interesting thing. Instead of getting out and hustling for fish, like the pelican, or taking delight in architecture, like the osprey, the Cormorant tiresomely perches, and waits, Micawber-like, for something to turn up.

In captivity it does better. In our Flying-Cage pool, the Cormorants play with sticks, and dive for amusement, more than any other bird, except the brown pelican. In fact, it seems like a different creature from the wild bird.

The Cormorant is, in general terms, a dull black bird, wholly devoid of colored plumage. Its range is given in the check list of the American Ornithologists' Union as "coasts of the North Atlantic, south in winter on the coast of the United States, casually, to the Carolinas." It lives upon fish, and wanders inland much farther than might be supposed.

**The Double-Crested Cormorant**<sup>2</sup> is the bird of the interior of the United States, from Texas northward into Manitoba, but also ranging to the Atlantic coast. Its color is glossy black. On the Pacific coast, from Washington to Alaska, is found the **Pelagic Cormorant**,<sup>3</sup> with an erect crest rising from its forehead, and by which this bird is easily recognized.

**Pallas' Cormorant**, which once inhabited the northern shore of Bering Sea, was the largest and handsomest bird of this Family. Its prevailing color was dark metallic-green, set off with blue and purple reflections. It was discovered by Bering in 1741, but is now quite extinct.

<sup>1</sup> *Phal-a-cro-co'rax car'bo*. Average length, 34 inches.

<sup>2</sup> *P. di-lo'phus*. <sup>3</sup> *P. pe-lag'i-cus*.

### THE DARTER FAMILY.

#### *Anhingidae.*

**The Snake-Bird, Darter, or Water-"Turkey,"**<sup>4</sup> is a web-footed bird, with many peculiarities. Its most popular name—Snake-Bird—has been bestowed in recognition of the fact that in this bird the neck and head are so long and slender they suggest the body and head of a snake. When not in action, the head and upper neck are only an inch in diameter, yet so rubber-like is the skin I have seen a Darter swallow a mullet 8 inches long, and 1½ inch in diameter—a truly snake-like stretch. Frequently when the head of a fish is in this bird's crop, the tail fin will protrude from a corner of the mouth.

The beak is like a Spanish dagger, and at all times is decidedly a dangerous weapon. One well-aimed stroke is enough to stab any ordinary bird to death, or destroy an eye. In a cagel of Darters the presence of a quarrel-



Drawn by EDMUND J. SAWYER.

SNAKE-BIRD.

<sup>4</sup> *An-hin'ga an-hin'ga*. Average length, 33 inches.

some bird is usually made known by the dead body of a cagemate that has been foully murdered.

In its home, the habits of the Snake-Bird interested me greatly. Almost invariably it perches on a dead tree, or a branch which overhangs water, preferably a small running stream. Its neighbors are the two white egrets, the Louisiana and little blue herons, and an occasional black vulture. Seldom indeed is one of these birds found swimming in the water, but Mr. C. E. Jackson once very dexterously speared one from his boat, as it was diving under him.

When your boat approaches a Snake-Bird and crosses his danger-line, the bird slides off its perch, falls straight down, and sinks out of sight. It goes down head erect, and "all standing," as if weighted with a bag of shot. This is the queerest of all bird ways in diving. If you halt, and watch sharply for the bird to reappear at the surface, for three or four minutes you will see nothing.

At the end of a long wait you will notice a sharp-pointed stick, half as long as an adult lead-pencil, sticking up out of the water. It looks so queer you watch it sharply. Presently you see the point of it turn a few degrees; and then you discover a beady black eye watching you. It is one of the neatest hiding-tricks practised by any water-bird I know.

The Snake-Bird has the power to submerge its body at any depth it chooses, and remain for any reasonable length of time. It is a very expert diver, and the manner in which it can pursue and capture live fish under water is enough to strike terror to the hearts of finny folk. The bird swims with a sharp kink in its neck, driving forward by powerful strokes of its cup-shaped feet. On overtaking a fish, the kink in its neck flies straight, and like the stab of a swift dagger the finny victim is transfixed. Then the bird rises to the surface,—for it is unable to swallow its food under water,—tosses the fish into the air, catches it head first, and in an instant it is gone.

In the United States this bird is most at home in the rivers and creeks of southern and central Florida, but it is also found farther west, along the Gulf. It is abundant in the delta of the Orinoco, in the Guianas, and far-

ther south. It lives well in captivity, and when provided with a large glass tank is quite willing to give daily exhibitions in diving after live fish. In color the adult male is a glossy black bird, and so is the female, except that her entire neck is light brown.

## THE GANNET FAMILY.

### *Sulidae*

**The Common Gannet**<sup>1</sup> is, in many respects, a bird of very striking appearance. It is a goose-like bird, as large as a medium-sized goose, and its prevailing colors are white and a very beautiful ecru. Its plumage is as smooth and immaculate as the surface of a wooden decoy; it has a slow and solemn manner, and has the least suspicion of man of any swimming-bird I know. Its head, neck and bill are massive, the latter especially being long and very thick at the base. The total length of this bird when adult is only a trifle under three feet.

Although the Common Gannet is strictly a bird of the ocean coasts, and apparently never is seen inland, it is a bird of such striking personality it well deserves to be introduced in these pages. Any large bird which once existed in countless thousands on our coast, and has not yet been exterminated, may well be known to every intelligent American.

Although the Gannet wanders as far south as Long Island, its real home is where it breeds. "While there are many points along the coast from Maine to Labrador where the Gannets might breed, they are found, so far as I have been able to ascertain, only at three places, an island in the Bay of Fundy, the Bird Rocks near the geographical centre of the Gulf of St. Lawrence, and Bonaventure Island, at Percé, Canada, the colony at Mingan being too small and too nearly exterminated to be taken into consideration." (Frederic A. Lucas.)

In 1860, Dr. Bryant estimated the total number of Gannets on the Bird Rocks at 150,000.

In 1872, Mr. William Brewster estimated the number then living there at 50,000.

In 1887, Mr. Lucas found not a single Gannet nesting on Little Bird Rock, and not over 10,000 on Great Rock.

Although the Gannets, and other sea-birds, make their homes on the most inaccessible

<sup>1</sup> *Su'la bas-sa'na*.



THE CORMORANT.

spots they can find, there is no bird which man cannot reach with a gun, no nest to which he cannot climb, or be lowered at the end of a rope.

Sea-birds everywhere are persecuted by man, either for their eggs or for themselves. In their breeding-season the Gannets are continually visited by Indians and whites, who take their eggs. "Scarce a day passes," says Mr. Lucas, "without a visit from fishermen in search of eggs, or murre. Many barrels of eggs are gathered during the season, and altogether the birds lead a rather precarious existence. There is a law regulating the taking of eggs, and if this were observed, or could be strictly enforced, a large number of eggs could be gathered annually, while at the same time the number of birds would steadily increase."

As will be inferred, the Gannet lives wholly upon fish, and is an expert deep-water diver. In his report on his "Explorations in Newfoundland and Labrador," Mr. Lucas gives the following interesting account:

"While lying at Grindstone Island we first made the acquaintance of the Gannets, whose head-quarters are at Bird Rocks, and had a good opportunity to watch them fishing. The birds are usually associated in small, straggling flocks, and with outstretched necks, and eyes ever on the lookout for fish, they fly at a height of from 75 to 100 feet above the water, or occasionally somewhat more. The height at which the Gannet flies above the water is proportioned to the depth at which the fish are swimming beneath, and Captain Collins tells me that when fish are swimming near the surface, the Gannet flies very low, and darts obliquely instead of vertically upon its prey.

"Should any finny game be seen within range, down goes the Gannet headlong, the nearly closed wings being used to guide the living arrow in its downward flight. Just above the surface, the wings are firmly closed, and a small splash of spray shows where the winged fisher cleaves the water to transfix his prey. Disappearing for a few seconds, the bird reappears, rests for a moment on the water, long enough to swallow his catch, then rises in pursuit of other game. The appetite of the Gannet is limited only by the capacity of its stomach, and a successful fisher may frequently be seen

resting on the water, too heavily laden to rise without disgorging a part of its cargo, which it sometimes must do to escape from the pathway of an approaching vessel."

Any person who is accustomed to diving, even from a very moderate height, knows well the serious disturbance to vision caused by the shock of impact with the water. That a Gannet—or any other bird—can fall from even a height of twenty-five feet, saying nothing of a hundred, take the water plunge, and *retain its gaze* upon its prey sufficiently to follow and capture it, surely betokens a special optical provision which as yet we know nothing about,



Photo. by R. J. BECK.

Galapagos Islands.

#### MAN-O'-WAR BIRDS.

and which remains to be discovered and described.

Besides the species described above, there are five other species of gannets, called **Boobys**, with various prefixes, which touch the coasts of the continent of North America.

### THE MAN-O'-WAR BIRD FAMILY.

#### *Fregatidae.*

Whenever at sea in the tropics your attention is arrested by the flight far aloft of a big, dark-colored bird with long, sharp-pointed wings, and a *long tail* that is *deeply forked*, know that it is a **Frigate-Bird**,<sup>1</sup> or, as the sailors call it,

<sup>1</sup> *Fre-ga'ta a'quil-a.* Length, about 40 inches.

**Man-o'-War "Hawk."** It is a long-distance flyer, and goes out far from land. Its beak is long, hooked at the end, and really very strong, but its legs are so short and stumpy they seem to be deformed. Under the throat there is a patch of skin quite devoid of feathers, which really is a sort of air-sac.

I once found the roosting-place of a colony of about forty of these birds, on the top of a perpendicular cliff seventy-five feet high on the seaward side of an island at the northwestern point of Trinidad. The birds came there regularly every night, to roost in some small dead trees that almost overhung the precipices. They were not nesting at that time, however, and were so very wakeful that even though I

went to their roost before daylight, I did not succeed in killing even one bird.

This bird inhabits the warm oceans of the Old World, as well as the New, and Mr. H. O. Forbes states that in the Cocos-Keeling Islands they are regular pirates, and gain their livelihood by remaining inactive, and forcing honest fisherfolk, like the gannets, and noddy terns, to disgorge for their lazy benefit the fish they bring home from distant fishing-grounds.

Mr. R. J. Beck found Frigate-Birds nesting in the Guadaloupe Archipelago, which were so tame and unsuspecting that he was able to approach quite near, and make the photograph which is reproduced on the opposite page.

## CHAPTER XXX

# THE ORDER OF TUBE-NOSED SWIMMERS,—MID-OCEAN BIRDS

### TUBINARES

These are indeed strange birds. To a landsman, it requires an effort to imagine a series of birds, some of them small and seemingly weak, which prefer to live in the watery solitudes of mid-ocean, indifferent to calms, and defying both tempests and cold. To my mind, there is no section of the bird-world so strange and so awe-inspiring as this. Just how the albatrosses and the petrels ride out the long, fierce gales, and keep from being beaten down to the raging surface of the sea, and drowned, I believe no one can say. It is no wonder that sailors hold the albatross in superstitious reverence, or that Coleridge has immortalized it in the "Rhyme of the Ancient Mariner." Well may a sailor feel that any large bird which lives only at sea, and follows his ship day after day, is the bird "that makes the breezes blow."

The members of this small group of mid-ocean birds are distinguished by the curious fact that the nostrils, instead of opening through the side of the upper mandible, near its base, are *carried well forward through two round tubes* that either lie along the top of the bill or along its sides. By this arrangement, the nostril opening is about half way between the base and tip of the bill. The bill terminates in a strong, serviceable hook, like the beak of a bird of prey.

This Order consists of the albatrosses, fulmars, shearwaters and petrels,—all of them deep-water birds, strong of wing, and brave spirited beyond all other birds. Of the thirty-five species and subspecies recognized by the American Ornithologists' Union, only two or three ever wander to inland lakes, even for three hundred miles from salt water. The variation in size from the largest albatross to the smallest petrel is very great; but at least half the species of the Order are to be classed

as large birds. Three species will suffice to represent the group.

### THE ALBATROSS FAMILY.

#### *Diomedeidae.*

**The Wandering Albatross**<sup>1</sup> is a bird of the southern oceans of the New World; it is the largest and handsomest species in the Order *Tubinares*. It has the longest wings, but the narrowest for their length, and the greatest number of secondary feathers (over thirty in number) of any living bird. The weight of an adult bird is from 15 to 18 pounds, and when the wings are fully extended, they have a spread of from 10 to 12 feet. Either when on the wing at sea, or mounted with spread wings as a museum exhibit, the wings of an Albatross are so exceedingly long and narrow that they have a very odd and unfinished appearance. They seem to be out of proper proportion, like wings lacking a proper outfit of secondary feathers. But they have their purpose. The Albatross can sail for hours, to and fro, without resting, and with wings so motionless they might as well be mechanically fixed.

Mr. Charles H. Townsend, who, as Naturalist of the United States Fish Commission Steamer *Albatross*, has had exceptional opportunities for studying Albatrosses at sea in all kinds of weather, has kindly furnished the following account of the most conspicuous species that inhabits the North Pacific:

**"The Black-Footed Albatross**<sup>2</sup> is a common bird almost anywhere in the Pacific Ocean, from the latitude of California northward. This dark species is frequently seen the first day

<sup>1</sup> *Di-o-me'de-a ex'u-lans.*

<sup>2</sup> *Di-o-me'de-a ni'gri-pes.*



out, and can usually be depended upon to follow vessels in increasing numbers. On many voyages between San Francisco and the Aleutian Islands, the average attendance of Albatrosses, or '*Gonies*,' as they are usually called, was from fifteen to twenty. Whether the same individuals stayed with the vessel during the whole

flock of birds would alight upon the water, often coming close enough to be caught on cod-hooks baited with pork. When on the wing, sometimes all the birds would assemble at once to feed on the waste thrown overboard from the galley, alighting in a confused manner, with much squawking and fluttering of wings.



BLACK-FOOTED ALBATROSS.

run, or were replaced from time to time by other birds encountered along the way, we could not determine.

"The birds were with us from daylight to dark, and in all sorts of weather. The *S. S. Albatross*, being engaged in deep-sea investigations, made frequent stops for the purpose of sounding and dredging. At such times the

"We often hooked specimens while the ship was under way, by paying out the line rapidly enough to leave the bait lying motionless, and buoyed on the surface with a cork. The birds were not able to pick up a bait while on the wing, or while it was moving. When hooked they would set their wings rigidly at an angle, and a rapid hauling-in of the long line would

send a bird skyward like a kite, which position it would retain until hauled down on the deck.

"Fishing for 'Gonies' was a common amusement on the *Albatross*, and specimens were often photographed alive on the decks, or marked in some way to determine if possible whether the same individuals followed the vessel throughout the voyage. Marked birds, however, never were seen again. The handling which they received probably disinclined them to follow the vessel.

"The arrival of an Albatross on deck was usually followed by the disgorging of more or less food. They could not rise from the deck, and frequently were kept on board for several days. They walk with great difficulty, and bite savagely.

"Albatrosses rise easily from the sea, and when the wind is blowing it is done very quickly. In calm weather, several strokes of the wings, and a rapid movement of the feet are necessary for the bird to clear the water. No bird can exceed the Albatross in the gracefulness of its flight. Usually following in the wake, it has, however, no difficulty in passing ahead of the vessel, always on rigid, motionless wings, rising, descending, or turning without a wing movement that is visible to the eye.

"On voyages southwestward from California, the Black-Footed Albatross did not usually follow the vessel more than two-thirds of the way to the Hawaiian Islands. A species known as *Diomedea chinensis* breeds in great numbers on the chain of islands extending northwestward from Hawaii. So far as I am aware, the breeding-place of *Diomedea nigripes* is not known. It probably breeds during the winter months on islands in the southern hemisphere. It is sometimes found in Bering Sea, particularly in the Bristol Bay region, and is met with all summer long in the Pacific south of the Aleutian Islands. During many visits to the Aleutian and other American islands, it was never found on land, and the natives were not acquainted with it as a nesting bird.

"In Bering Sea we sometimes met with the **Short-Tailed Albatross** (*Diomedea albatrus*). This species is nearly white, and in calm weather was usually observed resting on the sea, near the great flocks of fulmars. While the steamship *Albatross* was dredging off the south-

ern coast of Chili, the great wandering albatross was frequently to be seen resting upon the water about the vessel, and we had no difficulty in taking specimens with hook and line."

Perhaps the most wonderful sight in Albatross life is to be found on Laysan Island, in the Pacific Ocean, where thousands of these birds nest close together on an open plain. There are acres and acres of living Albatrosses, stretching away as far as the camera can include them, until the plain is white with them. They manifest little fear of man, even when iron rails are laid down, and small iron box-cars are



Drawn by J. CARTER BEARD.

#### STORMY PETREL.

pushed over them, to load with eggs from the nests.

### THE FULMAR FAMILY.

#### *Procellariidae.*

**The Fulmars** are like so many understudies of the Albatrosses; and the **Shearwaters** bring the Tube-Nosed group still nearer to the gulls and terns. The habits of all these are very much alike. All are strong-flying, mid-ocean birds, following ships for miles in order to pick up whatever edible food is thrown overboard. In one respect they are marine vultures, for some of the species make haste to feed upon any dead animal found floating on the sea, or stranded on the shore.

No one with eyes ever need cross the Atlantic

without seeing the dear little **Stormy Petrel**,<sup>1</sup> or "**Mother Carey's Chicken**," as it is called by sailors. After the last gull has been left far behind, and there are about two miles of water under the ship, in the trough between two waves there suddenly glides into view a pair of small black wings, fluttering rapidly, while two little webbed feet work violently to pat the concave surface of the deep blue water. Those who do not know the creature exclaim in surprise, "What in the world is *that*?"

"That" is one of the wonders of the ocean world. The cause for surprise is that so small and weak a creature—the smallest of all the web-footed birds, no larger, and seemingly no stronger than a cat-bird—should live on the watery wastes of a landless ocean, eating, sleeping and enjoying literally "a life on the ocean wave, and a home on the rolling deep."

<sup>1</sup> *Pro-cel-la'ri-a pe-lag'i-ca*. Length, 5.50 inches.

Even when seas are calm, and skies are clear, one cannot easily imagine how this creature can live, and find its food. But when a prolonged storm sets in, and for ten days, or two weeks at a stretch the surface of the sea is a seething, boiling caldron, with every wave a ragged "white-cap" and every square foot of the sea fretted like a fish-net by the force of the wind, how does the frail little Stormy Petrel survive?

You nearly always see this bird in the trough of the sea, skimming so low that its feet can paddle upon the surface of the water, and assist the wings. It is a black bird, with a large white patch on the rump, just above the tail. It rests upon the water fully half its time, I should say, and aside from the table and galley refuse thrown overboard from vessels, the bulk of its food must consist of the tiny crustaceans that inhabit the floating bunches of sargasso weed.

## CHAPTER XXXI

# THE ORDER OF LONG-WINGED SWIMMERS

### LONGIPENNES

The members of the Order of Gulls and Terns appeal to a greater number of admirers than any other group of web-footed birds. The reasons are, their wide distribution, both on salt water and fresh water lakes; their conspicuous and graceful flight; their partial immunity from wholesale slaughter, and their friendliness toward the arch-destroyer, man. Every harbor and every steamer track is a safe feeding-ground for these birds, and along thousands of miles of shore line, they are the most beautiful wild creatures that greet the eye.

The three North American Families of this Order are as follows:

	FAMILIES.	EXAMPLES.
ORDER LONGIPENNES.	GULLS AND TERNS, . . . . .	<i>LA'RI-DAE.</i> , . . . . . Herring-Gull; Common Tern.
	SKIMMERS, . . . . .	<i>RYN-CHOP'I-DAE.</i> , . . . . . Black Skimmer.
	SKUAS AND JAEGERS, . . . . .	<i>STER-CO-RAR-I-DAE.</i> , . . . . . Parasitic Jaeger.

### THE GULLS AND TERNS.

#### *Laridae.*

**The Herring-Gull,**<sup>1</sup> an old and familiar friend which ranges far inland, and also far outward on the sea, is the best and most interesting type of this Family. It is an ideal Gull,—long-winged, large, white and pearl-gray in color, strong, yet graceful on the wing, a good fighter, and sufficiently plentiful in number to be known to millions of people. It inhabits the whole sea-coast, and all the salt-water bays and inlets of North America, the great lakes, the lakes and ponds of Michigan, Minnesota, Iowa, and several of our larger rivers, such as the Potomac, Mississippi, Missouri and Columbia. From all their regular routes of travel and places of residence, they stray inland for an indefinite number of miles.

The Herring-Gull nests from southern Maine and the great lakes northward to the Arctic Ocean, and makes its winter home in the United States. All trans-Atlantic voyagers have seen it far out at sea, almost half way between Sandy Hook and Queenstown.

In Georgian Bay the sight of Gull life on the

<sup>1</sup> *La'rus ar-gen-ta'tus*. Average length, 24 inches.

crystal-clear waters, and clean, bare islets of pink granite near Owen Sound was one of the most enchanting I ever beheld. Going down Puget Sound on a cold and windy day in November, a large flock of the same old friends followed the steamer for twenty miles, sailing along beside us, sometimes within ten feet of the rail of the hurricane-deck,—a sight which well repaid one for half-freezing in order to see it to the most perfect advantage.


But why wander so far from home to see Gulls? Half a mile from the Zoological Park is the Williamsbridge Reservoir of the New York City water-works. Not long since, curiosity to see if any winter birds were being attracted by that very small but high basin of water, led me to climb up and see. To my great astonishment, I found a distinguished company of sixty-seven Herring-Gulls, standing and sitting in serene contentment on the sheet of ice that covered one-half the surface of the water. It was a nice, quiet, genteel place, well below the sweep of the wind; there was plenty of water for the birds to soak their feet in when the ice made them too cold, and what more could a Gull ask, except a daily delivery of fresh fish?

The voice of this Gull is not melodious; and some persons call it harsh and strident. But opinions differ, even on as small a matter as the voice of a Gull. I never yet heard the cry of a wild gull, either on the booming sea-shore, or over the silvery mirror of an inland lake, which was not music to my ears.

In captivity the Gull is badly handicapped. With the primaries of one wing clipped to prevent escape, and without the power of flight,

two enterprising Gulls decided to nest and rear a family. Accordingly they built a nest under a bush which stood on a point of the island, in a position that strategically was well chosen for purposes of defence. The two birds made a very wise division of the labor. The female built the nest, laid the eggs and hatched them, and the male did the screaming and fighting that was necessary to protect the family from molestation.

3



THE HERRING-GULL (1, 2) AND COMMON TERN (3, 4).

it is not seen at its best; for no Gull is perfect save in flight. Our flock is continually shrieking protests against unlawful detention, and with perfect wings every one would quickly fly away, as did those bred in the park and reared to adolescence with perfect wings. We tried to colonize them, but once away they never came back.

In an enclosure which embraced a pond and an island inhabited by about twenty Gulls, twelve Canada geese, and a few other birds,

Never was there a more bonnie fighter than that male bird. During that whole nesting-period, lasting from April 1 to May 15, he either bluffed or fought to a stand-still everything that came within ten feet of that nest. Before his defiant and terrifying screams, and his threatening beak and wings, no other Gull could stand for a moment. When a Canada goose crossed his dead-line, the Gull would rush at him, seize him by the nearest wing, wing-beat him, and hang on like a bull-dog, regard-

less of being dragged about by the stronger bird, until the goose was glad to purchase peace by retreating. During all these battles, the female sat firmly on her eggs, but pointed her bill at the sky, and screamed encouragement with all the power of her vocal machinery. Eventually the three eggs were hatched, and the young were reared successfully.

On certain islands along the coast of Maine, where Gulls nest in considerable numbers, the Bird Protection Committee of the American Ornithologists' Union, under the leadership of Mr. William Dutcher, has done important and effective work in securing the protection of the birds by the owners of the islands. As if to reward Mr. Dutcher for his labors in their behalf, the Gulls permit him to photograph them on their nests, at very short range. In England, the Zoological Society of London has awarded its medal to several persons for noteworthy services in protecting Gulls from destruction.

**The Common Tern,**<sup>1</sup> but for the timely interference of the Lacey Law, would ere now have become the very Uncommon Tern. The persons who for years slaughtered birds wholesale and without check for "millinery purposes" would have exterminated this species, at least all along the Atlantic coast.

In an evil hour, some person without compassion, and with no more taste for the eternal fitness of things than a Texas steer, conceived the idea of placing stuffed Terns on women's hats, as "ornaments." Now, unfortunately, woman's one universal weakness lies in the belief that whatever the Fashion Fetish commands that she shall wear, that is necessarily a beautiful thing for her to deck herself withal. As a result, we have seen thousands of angular, dagger-beaked, sharp-winged, dirty-plumaged, rough-looking and distorted Terns, each one a feathered Horror, clamped to the fronts and sides of the hats of women, and worn as head ornaments!

Those objects spoke very poorly for their wearers; for since the daughters of Eve first began to wear things on their heads, the Rumped Tern is the ugliest thing ever devised for head-gear. Thus has been developed a new bird species, which we will christen as above, with *Sterna horrida* as its Latin name. Thanks to the

<sup>1</sup> *Sterna hi-run'do*. Average length, 14.50 inches.

Lacey Law, however, the wearing of stuffed birds has, with fashionable people, quite gone out of fashion, and the only exceptions now seen are on the heads of servants, who, for motives of economy, are wearing the cast-off millinery of their mistresses.

The Tern is much smaller than the herring-gull; it has a very short neck, very long and angular wings, and when on the ground is not a bird of beautiful form. On the wing, however, and especially over the breakers, its appearance is graceful and pleasing. It is a white and gray bird, excepting the black bonnet which covers the upper half of its head and neck; and its bill, feet, and legs are coral red.

Along our Atlantic coast, and especially from Nantucket to Hatteras, it was once a very familiar bird, and its escape from annihilation has been of the narrowest. The Lacey Law, and the anti-bird-millinery laws passed by New York and other states, effectually stopped the sale of wild-birds and their plumage for "millinery purposes," and the Terns are no longer slaughtered as heretofore. In several places where they breed they are now protected, and henceforth should slowly increase in number.

There are now but few localities on our Atlantic coast between New Jersey and Nova Scotia where the Common Tern, or "Sea Swallow," breeds. Two of these are Muskeget Island, northwest of Nantucket, and Gardiner's Island. The once numerous colony that formerly inhabited Gull Island, near the eastern end of Long Island, was broken up and driven off by a "military necessity," no less important than the building of a modern fort to protect the City of New York. By a strange coincidence, it was the 12-inch guns of our coast-defence artillery that drove these much-persecuted birds from one of their favorite nesting-grounds.

## THE SKIMMER FAMILY.

### *Rynchopidae.*

**The Black Skimmer**<sup>2</sup> is a tern in form, but without the spear-like bill of the latter for spearing fish. Its lower mandible is formed for use as a cut-water,—long, thin, rather broad, and flattened *vertically*. The upper mandible is similarly shaped, but is shorter.

<sup>2</sup> *Rynchops nigra*. Length, about 16 inches.

When seeking food, the Skimmer looks for calm water, and then with most dexterous and well-balanced flight, it slowly wings its way close down to the surface, so low that the lower mandible is actually held *in the water* while the bird is in full flight. Any small edible object that happens to lie on the surface is shot into the mouth, through what is really a very narrow opening.

This is a bird of the tropics, and is much more at home on the coast of British Guiana, among the scarlet ibises, than it is on the coast of the United States anywhere north of Florida. I have never seen it elsewhere than in South America, and on our shores it is a visitor of great rarity.

#### THE SKUA AND JAEGER FAMILY.

##### *Stercorariidae.*

The members of this family are habitants of the cold northern seas and high latitudes. They

are strong-winged, bold and hardy, and so frequently rob other sea-birds of their prey that they are sometimes called the hawks of the sea. Living examples are rarely seen save by persons who are voyaging northward above the 49th parallel. Of the four species inhabiting North America, the following is the one most frequently seen in the United States:

**The Parasitic Jaeger<sup>1</sup>** is quoted geographically in the Check-List of the American Ornithologists' Union as follows: "Northern part of northern hemisphere, southward in winter to South Africa and South America. Breeds in high northern districts, and winters from New York and California southward to Brazil." A description of the colors of this bird would be a formidable affair, for both adults and young birds have each two color-phases. The beak of the adult is strongly hooked at the end, like that of a cormorant, but still more pronounced.

<sup>1</sup> *Ster-co-ra'ri-us par-a-sit'i-cus*. Length, about 17 inches.

CHAPTER XXXII

THE ORDER OF WEAK-WINGED DIVING-BIRDS

PYGOPODES

With this group, the Class of Birds enters upon a very marked and swift decline from the high types. Another step beyond this Order, and we land among birds so nearly wingless that they are without the power of flight. The birds of the present Order have wings that are small and weak; and while they are able to fly, and also to migrate, they fly feebly in comparison with the cloud-cleaving goose, duck, gull and albatross. Their legs are set far back on their bodies, and on land they have no choice but to stand erect—a posture which is strikingly characteristic of the wingless sea-birds, generally.

This Order, as represented in North America, contains but three Families:

	FAMILIES.	EXAMPLES.
<b>ORDER PYGOPODES.</b>	{ GREBES, . . . . .	<i>POD-I-CIP'I-DAE</i> , Pied-Billed Grebe.
	{ LOONS, . . . . .	<i>GAV-I-I-DAE</i> , . . Great Northern Diver.
	{ AUKS AND PUFFINS, <i>AL'CI-DAE</i> , . .	Razor-Billed Auk; Tufted Puffin; Murre.

Of these, the first and second are comparatively well known. The third is composed of birds that are strangers to the great majority of us; but inasmuch as Alaska is constantly being brought nearer to us, it is quite necessary that we should become acquainted with its most prominent forms of bird-life.

**The Pied-Billed Grebe,** or "**Hell-Diver,**" also called the **Carolina Grebe,**<sup>1</sup> is well qualified to stand as the representative of the Grebe Family, which in North America contains about six species. It is usually seen in the geographical centre of a quiet pond, sharply watching in every direction for enemies. It is a sad and uncomfortable-looking little creature, destitute of bright and pleasing colors, and also devoid of beauty. At a distance, the hunter is thrilled by the sight of what he gladly thinks is a duck; but on approaching nearer he sighs regretfully, and admits that it is "only a Grebe." If he fires at it, in revenge for the disappointment, the bird is gone before the charge of shot is half way to it, and only an innocent ripple marks its disappearance.

All the Grebes are expert long-distance

<sup>1</sup> *Pod-i-lym'bus pod'i-ceps*. Average length, 12 inches.

divers. They can either sink straight down, or dart down head first in a fraction of a second, and remain under water so long a time, and wim so far while submerged, that it is very difficult to follow their movements. Sometimes a Grebe will insinuate only its bill above the surface, in order to breathe without exposing even its head and neck. It is a waste of time, ammunition and self-respect to shoot and actually kill one of these birds; for they are very commonplace and useless.

The only redeeming feature about this bird is its breast, which is covered with a thick mass of very persistent feathers, set so tightly in a very tough skin that the evil-eyed milliners once used Grebes' breasts for hat trimmings.

The nesting habits of the Grebe are remarkable and interesting. Instead of choosing a dry situation, where incubation might proceed under the best possible conditions, it frequently chooses a clump of rushes in deep water and builds a floating nest, attached to the rushes. Sometimes, however, it selects a spot where the water is very shallow, and builds from the bottom up, using rushes when possible to procure them. In either case, the sodden mass rises only two or three inches above high-water mark, and



how the eggs ever receive warmth sufficient to hatch them is a mystery.


Occasionally a clump of rushes with a floating nest breaks loose from its moorings, and floats away. Some friends of mine once discovered a derelict nest, with the Grebe sitting serenely upon it, floating about in Lake Ontario, whither it had evidently been borne on the current of Johnson's Creek. Doubtless it is a real grief to Grebes that they cannot hatch their eggs under water!

Its prevailing color is brownish-gray, with black throat and chin. Its bill is dull white, with a broad, perpendicular band of black crossing it at the middle, like a rubber band to hold the mandibles together. In size this bird is about as small as a green-winged teal.

### THE LOON FAMILY.

*Gaviidae.*

**The Loon, or Great Northern Diver,<sup>1</sup>** is a large, showy, black-and-white bird, of such



COMMON MURRE.

THE LOON.

The Pied-Billed Grebe, also called *Dabchick*, and *Diedipper*, is a Pan-American bird, being found throughout North and South America from Cape Horn to the Mackenzie River, and from the Atlantic to the Pacific. Its phenomenally wide range includes Cuba, several others of the larger islands of the West Indies, and the Bermudas.

striking personality that when once well seen it is not easily forgotten. In bulk it is as large as an ordinary goose, and when standing erect, on land, its height is about 25 inches. Its neck and head are large and jet black, and the upper portion of the former is encircled by a white collar which is formed of upright lines of white

<sup>1</sup> *Gav'i-a im'ber.*

dots. The breast is pure white, and the jet-black back is marked by rows of rectangular white dots, or broken bars. The legs join the body far down, near the tail, and when the bird takes to the land, it rests on its feet, the lowest joint of the legs (tarsi), and the tail, which lies flat upon the ground.

Either on land or water, this Loon is a very showy bird, and also a bird possessing many of the mental traits which when combined form what we call "character." Usually it is very wide-awake, suspicious, and difficult to approach; but there are times when it will approach danger as if bent on suicide. Its cry is loud and far-reaching. Sometimes it is like a distressful howl, and again it resembles wild, uncultivated laughter. It is an expert diver and fisher, and in summer is at home all over the upper two-thirds of North America, breeding from our northern states to the Arctic Circle, quite across the continent. In winter it migrates southward to the Gulf and the Mexican boundary.

Its eggs are two in number, of a dull green color. The newly hatched birds are covered with black down, and in travelling the mother-bird often swims with them upon her back. The Loon rises from the water with considerable effort, and flies heavily, but in migrating its powers of flight are sufficient to carry it wherever it wishes to go.

In the Potomac River, and along the Virginia coast, this bird is called the "War Loon."

### THE CLIFF-DWELLERS OF THE SEA.

There is a Family of weak-winged birds whose members are all fisher-folk, and live high up on the ledges of the bold and precipitous cliffs which hem in the northern oceans. They are sociable birds, and where not destroyed by man, live in great companies varying from hundreds to thousands. They form, as a whole, a great and diverse company, divided into twenty-two well-defined species. Collectively, they are known as the Auk Family, and include 4 puffins, 6 auklets, or little auks, 5 murrelets, 3 guillemots, 2 murrees, 2 auks, and 1 dovekie.

Whenever you visit Alaska, or the arctic regions, almost anywhere on salt water, you will be surprised by the abundance of the birds belonging to this Family. Wherever rocky cliffs

rise out of blue water, you will find them tented by these interesting creatures. Doubtless, also, you will find that when such great gatherings of bird-life are to be studied and recorded, one good camera is better than ten guns.

Like the Aztecs who, like eagles, built high up in the crevices of the rock-cliffs of the gloomy Canyon de Chelly, to be inaccessible to the hostile enemies who gave no quarter, for similar reasons the feathered cliff-dwellers of the sea build in similar situations. Dearest of all spots to the nesting sea-bird is a precipitous islet of rock rising out of the sea, wholly inaccessible to the prowling wolf, fox, and wolverine, and if not actually inaccessible to man, at least so very difficult that he looks for easier conquests.

But let it not be understood that the birds of the Auk Family confine themselves to high cliffs and precipices. On the contrary, they congregate in thousands on rocky ridges, or on the tops of sandy hills—called dunes—at the sea-shore, where their nests are easily accessible to all their enemies. Just why their enormous colonies do not attract foxes and wolves by hundreds, we cannot imagine, unless it be for the reason that the general abundance of animal life dulls the edge of appetite and enterprise.

To any one interested in sea-birds, of which there is really a great variety, a trip to Alaska is replete with interest. Within a few hours after leaving Seattle, or, let us say at Port Townsend, the bird-life around the ship fairly compels attention. A flock of gulls fly so close to the rail of the hurricane-deck that some of them might be caught with a dip-net. Pigeon guillemots, and ducks of several species afloat on the cold waters of the Sound ostentatiously swim out of the steamer's track. On the ocean, it will be strange if an albatross does not sail out of space, and with far-stretching wings swoop and soar, and sail after you, hour after hour, without once flapping its wings!

In Bering Sea, no matter where you land, the chances are that thousands of murrees and puffins are there to greet you with noisy cackle, and spread a cloud of wings overhead when you disturb them. Really, the rookeries of Alaska—of seals as well as birds—are alone sufficient to repay a trip to that arctic wonderland, aside from the wonderful scenery, flora, and big

game. There are dozens of birds there which we would gladly introduce to the reader, but owing to uncontrollable limitations, only the most interesting examples can be accorded space.

Of all arctic and northern sea-birds, the **California Murre**<sup>1</sup> (pronounced *mur*) deserves to be mentioned first, for the reason that it is and ever has been most in the public eye. This is really a subspecies of the **Common Murre**<sup>2</sup> of the North Atlantic, which nests on Bird Rocks in the Gulf of St. Lawrence, and sometimes comes as far south as Massachusetts. There is another North Atlantic species, called **Brunnich's Murre**,<sup>3</sup> also nesting on Bird Rocks, which occasionally strays down to Long Island. Both the Atlantic species are black above, and white underneath.

The California Murre is the bird which once nested on the Farallone Islands, about thirty-five miles west of San Francisco, in countless thousands, and furnished between 1880 and 1890, according to Mr. W. E. Bryant, from 180,000 to 228,000 eggs per annum to the San Francisco market. Like true Americans, the eggers always endeavored to make "a clean sweep," regardless of the future of the rookery, and under their ministrations the Murres rapidly declined in number.

Finally an appeal was made to the United States Light-House Board. The admirable record of that body in the preservation of wild life was sustained by an order which at once put a stop to all egg-gathering on the Farallones. It has already been noted in the chapter on seals and sea-lions that the only localities on the California coast where sea-lions are now safe from annihilation are the light-house reservations, the most important of which are the Farallones.

The following vivid pen-picture of the California Murre at home, on Hall Island, Bering Sea, Alaska, is from the pen of Mr. John Burroughs (Harriman Alaska Expedition, p. 109):

"The first thing that attracted our attention was the Murres—'urries' the Aleuts call them—about their rookeries on the cliffs. Their numbers darkened the air. As we approached, the faces of the rocks seemed paved with them,

with a sprinkling of gulls, puffins, black cormorants and auklets.

"On landing at a break in the cliffs where a little creek came down to the sea, our first impulse was to walk along the brink and look down upon the Murres, and see them swarm out beneath our feet. On the discharge of a gun, the air would be black with them, while the cliffs apparently remained as populous as ever. They sat on little shelves, or niches, with their black backs to the sea, each bird covering one egg with its tail-feathers. In places one could have reached down and seized them by the neck, they were so tame and so near the top of the rocks. I believe one of our party did actually thus procure a specimen. It was a strange spectacle, and we lingered long looking upon it. To behold sea-fowls like flies, in uncounted millions, was a new experience.

"Everywhere in Bering Sea the Murres swarm like vermin. It seems as if there was a Murre to every square yard of surface. They were flying about over the ship, or flapping over the water away from her front at all times. I noticed that they could not get up from the water except against the wind; the wind lifted them as it does a kite. With the wind, or in a calm, they skimmed along on the surface, their heads bent forward, their wings beating the water impatiently. Unable to rise, they would glance behind them in a frightened manner, then plunge beneath the waves until they thought the danger had passed. Their tails are so short that in flying their two red feet stretched behind them to do the duty of a tail."

Mrs. Florence Merriam Bailey says that "When incubating, one bird stays on the nest during the day, and the other during the night, and when the exchange is made a great commotion ensues, the air being filled with quarrelling, screaming masses of bird-life." ("Handbook," p. 17.)

In its breeding plumage, the California Murre has a jet-black head and neck, the back is dull black, or slate color, and the under parts are white. In winter the sides of the head and throat are white. The range of the species is from California to Hall Island, Bering Sea.

**The Puffins** are the clowns of the bird-world. Without exception, they are the drollest-looking

<sup>1</sup> *U'ri-a tro'i-le californica.*

<sup>2</sup> *U. troile.*

<sup>3</sup> *U. lom'vi-a.*

things in feathers. The countenance of a Puffin always reminds one of a face in a comical mask, while in manner they are so solemn, and take life so seriously, their clown-likeness is all the more pronounced.

The most remarkable feature of a Puffin is its huge, triangular beak, which is flattened out into two high, thin plates, set edgewise against the head, and gorgeously colored. After the breeding-season, certain plates at

mies of several kinds. In the defence of their homes they are quite courageous, and often an angry or well-frightened Puffin will seize an offending nose, or human hand, bite it severely, and hang on like a bull-dog. In places where these birds nest in burrows, sentinels are always posted outside, to give the alarm of any approaching enemy.


It is to be observed, however, that Puffins do not always nest in burrows, but frequently they find rock-ledges so rugged and broken that they can find good nesting-sites in deep and narrow crevices, wherein they are reasonably safe from molestation. A Puffin lays but one egg, which is large and white, and placed at the end of its burrow. Of course all these birds dive and swim well.

**The Tufted Puffin**<sup>1</sup> is the most widely distributed member of this genus, being found from southern California all the way up the Pacific coast to Alaska, Bering Strait, Siberia and on down to Japan. It is (or at least *was*) abundantly represented on the Farallone Islands<sup>3</sup> from April to July, when they breed there.

This species is instantly distinguishable by its black plumage, its big, triangular bill colored bright red and olive green, white eye and white triangular cheek-patch. In the breeding-season, a beautiful flowing tuft of soft, yellow feathers, thick as a lead-pencil, comes forth just behind the eye, and flows backward and downward in a graceful curve.

On the Atlantic side, from Maine to Greenland, and also from Great Britain to North Cape, lives the **Common Puffin**,<sup>2</sup> or "**Sea Parrot.**" Of this bird, the whole side of the head, and the breast and abdomen are white, the remainder of the plumage being deep black. Wherever found, it is one of the most interesting birds to be met with near the sea, and its comical appearance, queer movements and fierce temper when disturbed never fail to amuse the observer.

**The Auks and Auklets** are really birds of the cold northern waters; but on the Pacific side there are four species which touch the coast of the United States, and two of them even push their way down to Lower California. These birds are much like puffins with rational



1. COMMON PUFFIN.                      2. TUFTED PUFFIN.  
3. RHINOCEROS AUKLET.

the base of the beak are shed. The bird is about the size of a summer-duck. Its wings are short, and very scantily feathered, and its tail is so short as to be practically invisible. In flight its wings look very much like the wings of a penguin as it swims with them under water.

In many respects Puffins are wise birds, and if there is aught in the survival of the fittest, they should live long and prosper. They have the remarkable habit of nesting in burrows, which they dig deeply, usually about three feet, in the steep sides of sandy hills. In these retreats they can defend themselves against ene-

<sup>1</sup> *Lun'da cir-ra'ta*. Length, 15 inches.

<sup>2</sup> *Fra-ter'cu-la arc'ti-ca*. Length, 13 inches.

beaks, and I believe all existing species are black above and white below. The beaks show but little tendency to the sportive flattening so characteristic of the puffins.

These birds are very strong divers, and get a great portion of their food from the bottom of the sea. The two species found all along our Pacific coast, on the Farallone Islands and Santa Catalina, are the *Rhinoceros Auklet*<sup>1</sup> (14 inches long), and the *Cassin Auklet*, the former so called because of an erect horny shield at the base of its beak. The *Least Auklet*<sup>2</sup> is only 6½ inches long—about the bulk of a small, thinly feathered screech-owl.

**The Razor-Billed Auk**,<sup>3</sup> of the North Atlantic Ocean, sometimes wanders in summer to the coast of Maine, and in winter even migrates as far south as New Jersey. (Robert Ridgway.) It is 17 inches long, and is the largest living member of the group of auks. As might be expected, it is a distinguished resident of the Bird Rocks.

**The Great Auk** is now a bird of history and museums only. It met its fate on Funk Island, a treeless dot in the sea, about thirty miles northwest of Newfoundland, which was the first land met with as the Auks swam south-

ward on their annual migrations. The wings of this bird were so little developed that it was wholly unable to fly, and while on land it was any one's prey.

The thousands of Great Auks that visited Funk Island naturally attracted men who wished to turn them to account. Whalers were landed, and left there to kill Auks and secure their feathers. The birds were either driven into pens and slaughtered there, or else the pens were used to contain their dead bodies. Apparently great numbers of the bodies were burned for fuel. About 1844, the species became entirely extinct.

When Funk Island was visited by Mr. F. A. Lucas in 1887, in quest of Auk remains, he found deposits of bones several feet in thickness, evidently where the bodies of slaughtered birds had been heaped up, and left to decay. Out of these deposits, several barrels of mixed bones and peaty earth were taken which yielded several complete skeletons of that species.

Had the Great Auk possessed wings for flight, the chances are that it would not have fallen such easy prey to its exterminators. The moral lesson of its fate is—in these days of fire-arms and limitless ammunition, no bird should be hatched without steel-plate armor, strong wings for flight, and swift legs for running away.

<sup>1</sup> *Cer-o-rhin'ca mo-no-cer-a'ta.*

<sup>2</sup> *Sim-o-rhyn'chus pu-sil'lus.*

<sup>3</sup> *Al'ca tor'da.*



W. Carter Beard

EMPEROR PENGUIN.

## CHAPTER XXXIII

# THE ORDER OF FLIGHTLESS DIVERS

### IMPENNES.

No matter where man may go, on land or sea, or polar ice-pack, Nature holds birds in readiness to welcome him.

When Peary reached the point of land that is nearest the north pole, at the northeastern extremity of Greenland, on July 4, 1892, he found there the snow-bunting, sand-piper, raven, Greenland falcon, and ptarmigan. On the great arctic ice-floe, at Lat.  $82^{\circ} 40'$ , Nansen saw the fulmar (*Procellaria glacialis*), and the black guillemot, and a little later the ivory gull, little auk, and Ross's gull. When the steamer *Belgica* penetrated the awful solitudes of the antarctic archipelago, in 1898, and spent there the "First Antarctic Night"<sup>1</sup> ever endured by man in that region, Dr. Frederick A. Cook and his companions found, in close proximity to their ice-bound ship, flocks of large and very strange birds. They had an opportunity to study the wonderful **Emperor Penguin**<sup>2</sup> in its haunts, such as never before had been secured by naturalists.

This species is the largest of the wingless and flightless swimming-birds. In bulk it is about the size of our great white pelican. Its height is  $3\frac{1}{2}$  feet, and it stands as erect as any soldier on parade. In its erect posture its wings seem like arms, and its queer manner of talking, scolding, and prying into man's affairs, makes this bird seem more like a feathered caricature of a big, fat human being than an ordinary diving-bird. Its head is black, its abdomen is white, and its legs and feet are feathered quite down to the claws. The wings are covered with feathers that are more like fish-scales than feathers, and the feathers of the back also are very close and scale-like.

To a naturalist or bird-lover, the sight of

<sup>1</sup> Dr. Cook's valuable narrative of the exploration bears this title.

<sup>2</sup> *Ap-te-no-dy'tes fos'ter-i*.

great flocks of Emperor Penguins, and of the smaller *Pack Penguins*, on the antarctic ice-floes, must be sufficient to repay the explorer for many of the long, dark hours of the voyage that is required to reach their haunts. Says Dr. Cook:

"A number of royal and small penguins, and some seals, were led by curiosity to visit us. They called, and cried, and talked, and grunted as they walked over the ice about the ship."

I have seen and heard the **Black-Footed Penguin**,<sup>3</sup> of South Africa, scold and complain in a most human-like manner. On land, or on an ice-floe, this bird is so awkward and helpless that any blood-thirsty observer can walk up and kill it with a stick. Place it in water, however, and what a transformation! Immediately it will give an exhibition of diving which is astonishing.

In an instant, a waddling, slow-moving, almost helpless bird is transformed into a feathered seal. With its feet floating straight behind, and of no use save in steering, it points its beak and head straight forward, and swims wholly with its wings. Those flipper-like members reach forward simultaneously, work in perfect unison, and strike the water like living paddles—which they are. The quickness and dexterity of this bird in chasing and capturing live fishes, swallowing them under water, and instantly pursuing others, is one of the most wonderful sights in bird-life. The bird always dives with its lungs full of air, and during the middle of its period under water, it exhales. When it does so, bubbles of air issue from each corner of the mouth and float upward like two strings of pearls.

It is strange that the feet perform very little service while the Penguin is diving; but such is

<sup>3</sup> *Sphe-nis'cus de-mer'sus*.

the fact. Of all birds that love water, I think the Penguin loves it most. It will lie on its side at the surface, and in sheer playfulness and excess of joy, beat the water with its uppermost wing, wriggle about, then turn over and splash with the other.

In the sea, a flock of Penguins is readily mistaken for a school of dolphins, because they dive so persistently, in order to swim with their wings, and thus get on in the world very much

faster than if they sat up and paddled with their feet.

There are about twenty species of Penguins, of which the Emperor is the largest, and the King Penguin second. All are found in the southern hemisphere. The largest Emperor Penguin ever weighed and recorded, weighed 78 pounds! Needless to say, these birds live almost wholly upon fish, in the capture of which they are the most expert of all birds.



## CHAPTER XXXIV

# THE ORDER OF WINGLESS LAND-BIRDS

### RATITAE

Lowest of the Orders of living birds is that which contains the birds which are so nearly wingless that they are wholly unable to fly, but are provided with long and powerful legs, which enable them to run swiftly. Of these, there are a larger number of species than might be supposed, but our purpose requires here only the briefest introduction of a few important forms. The majority of the birds of this group are birds of great size, and their legs are so long and powerful they are able to kick or strike quite dangerously. These are the ostriches, rheas, cassowaries, and emeus.

**The African Ostrich<sup>1</sup>** is the largest living bird, and in every respect it is a worthy descendant of the still more gigantic but now extinct moa of New Zealand. Our full-grown male Ostrich stands, when fully erect, exactly 8 feet in height to the top of its head, and weighs about 275 pounds. The manager of the Florida Ostrich Farm at Jacksonville states that the average weight of adult African Ostriches is about 300 pounds.

Once abundant in nearly all the dry and open country of Africa, except the Sahara and Libyan deserts, this noble bird has shared the fate of the elephant, rhinoceros, buffalo and giraffe. To-day it is to be found but sparingly, and only in those regions of southern and eastern Africa wherein it has been impossible for man to exterminate it. The value in America of a full-grown African Ostrich is \$250.

<sup>1</sup> *Struthio cam'el'us.*

Fortunately the Ostrich farms of South Africa and southern California have proven completely successful, and bid fair to perpetuate this grandest of all feathered creatures long



New York Zoological Park.

CERAM CASSOWARY.

after the last wild flock has been destroyed. If many Ostriches still remain in the Egyptian Soudan, the stringent game-laws recently enacted to protect the wild life of that region will go far toward perpetuating them.

**The Rhea, or South American Ostrich,<sup>2</sup>**

<sup>2</sup> *Rhe'a americana.*

is a bird which is so constantly overshadowed by the larger and more showy African ostrich that it is not appreciated at its true zoological value. In height it stands about 5 feet, its bulk is only about one-half as great as that of the African ostrich, and its plumage has much less value. Nevertheless, the adult bird, in full plumage, is a fine creature, of a beautiful bluish-gray or drab color, and when it opens its wings they seem surprisingly long. A fine, male Rhea "showing off" its plumage is an object which always commands admiration.

This bird inhabits Patagonia, the Argentine Republic, and the more remote plains of Uruguay and Paraguay. Frequently, half-grown birds find their way into the wild-animal markets so easily that they sell at from \$40 to \$50 each.

**The Emeu**<sup>1</sup> stands half way, literally, between the ostrich and cassowary, being considerably larger than the latter. Its neck and head are ostrich-like, but in the shape of its body it is more like the cassowary. Like the latter, its feathers seem like long, coarse hair, of a gray-brown color. The lower outline of an Emeu's body is almost a straight line, with the legs in the centre, and the highest point of the back curve comes directly above the insertion of the legs. Thus the Emeu appears to be, and is, a very well-balanced bird. Its home is the upland plains of Australia, so far back in the interior that it is now found only with great difficulty.

Like the cassowary, the Emeu is easily kept in captivity, and is not expensive to buy. In Woburn Park, England, owned by the Duke of Bedford, troops of these birds stalk freely over the vast green lawn; and surely no birds could be more striking, or picturesque in such situations. Strange to say, a fully grown Emeu can be bought in New York for \$125.

<sup>1</sup> *Dro'mae-us no-vae-hol'land-ae.*

**The Ceram Cassowary**<sup>2</sup> is a big, purplish-black bird, with highly-colored patches of naked skin on its upper neck, and an elevated helmet or casque on the base of its upper mandible. Its feathers look like coarse and stiff hair from three to six inches in length, and its legs and feet are very thick and heavy for its stature. The height of a Cassowary is about 5 feet.

Cassowaries are forest-loving birds. They inhabit Australia, Ceram, and other islands of the Malay Archipelago. Because they take kindly to captivity, they are frequently seen in zoological parks and gardens, and travelling shows.

**The Apteryx, or Kiwi**,<sup>3</sup> of New Zealand is the lowest species in the scale of living birds. It is absolutely without wings, and it lives upon the ground in dark forests, where it can hide. Unfortunately, it has no means of defence, and is too small to escape from a dangerous enemy by running away. It is about the size of a Cochin-China hen, covered with long, stringy, hair-like feathers of a dark-brown color, and it has a long, curved beak like that of an ibis, for probing in the earth. Undoubtedly, the civilized development of New Zealand will cause the total extinction of this very shy but interesting species at no distant day.

In captivity in a zoological garden it is as shy and retiring as a beaver. In order to keep it from fretting itself to death, it is necessary to place in a corner of its cage a sheaf of straw, or a bundle of leafy branches, behind which it can retreat from observation, and lie concealed.

Outside of its New Zealand home, this bird is rarely seen in captivity, which is to be regretted, because it is one of the most interesting forms of the whole avian world.

<sup>2</sup> *Cas-u-a'ri-us gal-e-a'ta.*

<sup>3</sup> *Ap'te-ryx aus-tral'is.*

BOOK III

REPTILES



## CHAPTER XXXV

### INTRODUCTION TO THE CLASS OF REPTILES

**The Point of View.**—In studying or not studying the world of reptiles, everything depends upon the point of view. With persons in middle life, who hold up their hands and shudder at the mention of the word “reptile,” there is nothing to be done. They are victims of an unreasoning prejudice that often is deliberately taught to young people, both by precept and example, until at last it becomes bone of their bone and flesh of their flesh. Human children are *not* born with the inherited fear of reptiles which is so characteristic of the apes and monkeys of the jungles; and it is not fair to terrorize their innocent souls with awful “snake stories,” any more than with the “ghost stories” which most careful parents forbid.

With young people whose minds have not been artificially warped by older persons who abhor all reptilian life, much may be done.

Now, come! Let us reason together.

Despite electricity and steam, this world is yet a fairly large place. That it has existed through countless ages, and that its animal life has gone through many marvellous transformations, no one can deny, without being put to shame by the silent and immutable testimony of the rocks. This world, the animals now living upon it, and those lying within it, entombed by Nature’s hand, have been *millions of years* in forming. If you doubt it, go into an Arizona canyon, half a mile in depth, and at the bottom of a mountain-wall of rock, dig out the remains of a fossil, then ask yourself this question: “How long has it taken Nature to pile half a mile of solid rock upon the grave of this creature, *and then cut down to it again?*”

In the evolution of the birds of to-day, the reptiles of the past have played an important part; and the study of the Class *Reptilia* is very much worth while, if for no other reason than to learn the nearness of the relationships between its members and the birds.

Remember, first of all, that the reptiles of to-day are actually insignificant in comparison with those which existed ages ago, the bones of which are now fast coming to light. A twenty-four-foot python or anaconda of to-day, lying beside a sixty-foot dinosaur, with a hind leg ten feet high, would be like a garter-snake beside a kangaroo.

In this day of liberal thought and broad reasoning, any person whose knowledge of the world of reptiles is limited to the false notion that all these creatures are either “slimy” or dangerous, is to be pitied. A persistence in that all-too-common estimate is a distinct loss to all those who entertain it. It means the shutting out, with the black curtain of Ignorance, of a whole world of interesting forms and useful facts, and also a lifetime of cringing fear, largely without cause.

Young Americans, I exhort you to take a broad and sensible view of the reptilian world,—as of every other great subject. Many of these creatures are worth knowing, some because they are wonderfully interesting, some because they are useful, and others because they are dangerous. *None of them, however, are “slimy”!* A snake may be cold to the touch, but its skin is as clean and free from slime as a watch-chain. What is more, there is no living creature, not even a dolphin, dripping from the sea, which possesses a skin displaying the beautiful pattern of colors and the rainbow iridescence of the reticulated python, of the East Indies. In reality there are a great number of reptiles that are undeniably beautiful.

I would it were possible to touch upon all the Orders of Reptiles, extinct as well as living, and introduce some of the gigantic and wonderful lizards that were like kangaroos, rhinoceroses, and sea-lions, and also like nothing else under the sun; but in this volume it is impossible. There is space available only for the four Orders

of living Reptiles; the seven that are extinct can be studied elsewhere by those who become specially interested in this subject.

**The Grand Divisions of Living Reptiles.**—There are, all told, eleven Orders of the Class *Reptilia*; but seven of them are extinct, and for the present these will be left out of consideration. The four Orders of living reptiles are made up as shown in the following synopsis:

wide range of variation, beginning with the clumsy-flipped harp-turtle, passing the gila monster, the swift-footed monitor, the kangaroo-like collared lizard (of Arizona), the gliding serpents, and ending with the flying dragon.

In their food habits, the range of the world's reptiles is infinitely great, embracing fruit, vegetables, herbage, and all forms of flesh, living and dead. Oddly enough, however, no modern reptile has been provided with molar teeth for the

THE ORDERS OF LIVING REPTILES.

ORDER.	PRONUNCIATION.	GROUPS INCLUDED.	EXAMPLES.
CROCODILIA.....	<i>Croc-o-dil'i-a</i> .....	Gavials, Crocodiles, Alligators...	Florida Crocodile, Alligator.
CHELONIA.....	<i>Ke-lo'ni-a</i> .....	{ Tortoises, Terrapins and Sea- { Turtles.	{ Box Tortoise, Painted Ter- { rapin, Hawksbill Turtle.
LACERTILIA.....	<i>La-ser-til'i-a</i> .....	Iguanas, Slow-Worms, Skinks.	{ Marine Iguana, Glass "Snake," { Blue-Tailed Lizard.
OPHIDIA.....	<i>O-fid'i-a</i> .....	{ Colubrine Snakes, Rattlesnakes, { Harlequin Snakes.	{ Anaconda, Timber Rattle- { snake, Coral Snake.

**General Characters of Reptiles.**—Chiefly through certain extinct species, the reptiles lead so directly into the birds that the two Classes overlap each other.

In the Berlin Museum are the well-preserved fossil remains of a bird called the *Archaeo-pteryx*, which had a long, lizard-like tail fully covered with feathers, and lizard-like teeth in its beak. In 1873, Professor Marsh discovered in the chalk-beds of western Kansas, a low-formed, penguin-like bird, called the *Hesperornis*, also provided with teeth.

All reptiles are cold-blooded animals, and breathe air by means of lungs. Because of the low temperature of their blood, and their slow heart-action, many of them are able to remain under water for quite lengthy periods—of minutes, not hours. Some turtles and terrapins become so thoroughly dormant at the approach of winter that the vital organs actually suspend their functions, for a period of from one to three months. It is then that these creatures bury themselves in the mud at the bottom of ponds, and so pass the winter months.

The majority of reptiles are covered with scales, or armor of solid bone, and are provided either with teeth for conflict and offence, or with armor for defence. Their means of locomotion show a

mastication of food. The saurians, lizards and serpents have teeth for seizing and holding their living prey. The turtles, however, are quite toothless, and in place of teeth their horny jaws have sharp, cutting edges for clipping up their food into pieces small enough to be swallowed without mastication.

The teeth of serpents and crocodylians generally are perpetually renewed, as fast as old teeth are worn out, and disappear. By reason of this, the lives of these reptiles are indefinitely prolonged, and it is believed that some of them continue to grow almost as long as they live.

The great majority of reptiles reproduce by laying eggs, which are hatched either by the heat of the sun, or by the fermentation of muck-heaps. Many species of serpents hatch their eggs in their own bodies, and bring forth their young alive. Such species are called viviparous. Those which lay eggs are called oviparous.

Some reptiles, notably the crocodiles and tortoises, continue to grow almost as long as they live. Doubtless this is also true of some large species of serpents, such as the great constrictors of India and South America.

**Distribution.**—Reptiles reach their maximum development in the tropics, and the subtropics, between the isothermals of 32° F. North and

south of that zone, reptilian life still is abundantly represented, but chiefly by small species. The largest land-serpents are found in the low-lying, moist and hot forests of the equatorial regions; but crocodilians of the largest size are found several hundred miles from the equator, both north and south. The largest tortoises live directly on the equator.

**Poisonous Species.**—Among our reptiles only one lizard and a few species of serpents are venomous,—an exceedingly small proportion of the whole number. Indeed, so few in number are the dangerous species of North America, it is an easy matter for any intelligent person to learn to recognize all of them at sight. In a few hours of diligent and conscientious study, aided by a text-book that has been properly designed, any clear-headed person over fourteen years of age can learn to determine almost at a glance whether any fully grown serpent of North America is poisonous or harmless. This is possible from the fact that more than half of the venomous species possess rattles, and those which have not are few in number.

**Useful Species.**—Many reptiles are of decided value to mankind, by reason of the rats, mice and other destructive vermin which they destroy. Others diligently devour insects. Quite a number furnish useful food, and some yield skins and other commercial products of much value.

**Lack of General Knowledge Regarding Reptiles.**—While birds have been well taken care of in books, museums, zoological gardens and lectures, and mammals are now coming in for a small proportion of the attention they deserve, the reptiles have been greatly neglected. Very few zoological institutions contain collections of reptiles worthy of the name, and the books on this Class are mostly to be written. As a result of this well-nigh universal lack of opportunity for study, the great majority of persons possess very little precise and clear information regarding these creatures. The following chapters are offered merely as a foundation on which to build an acquaintance with a world of living creatures concerning which we are assured that a large number of persons sincerely desire information.



E. F. KELLER, Photo.

MISSISSIPPI ALLIGATOR, "OLD MOSE."

N. Y. Zoological Park.

Captured in Indian River, Florida. Length, 12 feet 5 inches.



CHAPTER XXXVI

THE ORDER OF CROCODILES AND ALLIGATORS

CROCODILIA

The warm regions of the world contain nineteen species of big, burly, bony-armored reptiles, with long tails, powerful jaws, and tempers as ugly as their own rough backs. These creatures are known collectively as Croc-o-dil'i-ans, and two Families embrace all the gavials, crocodiles, alligators and caimans of both the Old World and the New.

So pointed is the need for a clear bird's-eye view of this important group of large reptiles, it is necessary to set forth a synopsis of the entire Order. The species will be arranged in a regular series according to the width of their heads, beginning with the narrowest.

A SYNOPSIS OF THE CROCODILIANS.

The measurements given are believed to represent the maximum size attained by each species.

FAMILY.	GENUS.	SPECIES.	COMMON NAME.	LOCALITY.	
GAVIAL FAMILY:	<i>Gav-i-al'is</i> .....	<i>gan-get'i-cus</i> .....	Indian Gavial, 17 feet.....	Northern India.	
		<i>To-mis'to-ma</i> .....	Bornean Gavial, skull 3 feet 3 inches....	Borneo and Sumatra.	
CROCODILE FAMILY:	<i>Croc-o-di'lus</i> .....	<i>cat-a-phrac'tus</i> .....	Sharp-Nosed African Crocodile.....	W. Africa.	
		<i>in-ter-me'di-us</i> .....	Orinoco Crocodile, 12 feet.....	Venezuela.	
		<i>johns'ton-i</i> .....	Australian Crocodile.....	Australia.	
		<i>rhom'bi-fer</i> .....	Cuban Crocodile, 7 feet.....	Cuba only.	
		<i>a-cu'tus</i> .....	American Crocodile, 14 feet.....	Central and South America.	
		<i>a. flor-i-dan'us</i> .....	Florida Crocodile, 14 feet 6 inches.....	Florida.	
		<i>nil-ot'i-cus</i> .....	Nile Crocodile.....	Africa generally.	
		<i>po-ro'sus</i> .....	Salt - Water Crocodile, 16 feet.....	Malayana.	
		<i>pa-lus'tris</i> .....	Mugger, 12 feet.....	India.	
		<i>Os-te-o-lae'mus</i> .....	<i>te-tras'pis</i> .....	Broad-Nosed African Crocodile, 6 feet..	Equatorial W. Africa.
		<i>Cai'man</i> .....	<i>tri-go-na'tus</i> .....	Rough-Backed Caiman, 6 feet.....	Upper Amazon.
			<i>pal-pe-bro'sus</i> .....	Banded Caiman.....	South America.
			<i>scl'e'rops</i> .....	Spectacled Caiman .	Central and South America.
			<i>ni'ger</i> .....	Black Caiman, 20 feet (Bates).....	Guianas: Brazil.
		<i>All'i-ga-tor</i> .....	<i>lat-i-ros'tris</i> .....	Broad - Nosed Caiman.....	Amazon to Rio de la Plata.
<i>si-nen'sis</i> .....	Chinese Alligator, 6 feet.....		China.		
	<i>miss-is-sip-pi-en'sis</i> .....	Common Alligator, 16 feet.....	United States.		


**General Characters of Crocodylians.**—A crocodylian is a lizard-like reptile, of very large size, with short, thick legs, a long tail, and the most highly developed vascular system to be found among reptiles. Its back and neck are protected by powerful armor consisting of rough, lozenge-shaped plates of solid bone set in a very thick and tough skin, and arranged in rows, both lengthwise and crosswise.

Both the tail and the abdomen and throat are covered by a regular arrangement of tough scales. The whole animal is covered by a thin, transparent epidermis which is impervious to water. The tail is long, flattened vertically, and fringed

The eyelids are movable, and the ear opening closes tightly by a flap of skin controlled by voluntary muscles.

Most saurians are voiceless or nearly so; but the alligator emits a very deep bellow, or roar, which in animals over ten feet in length is much lower on the scale than any fog-horn.

"The difference between a crocodile and an alligator" (a question that has been asked a countless number of times) consists chiefly in the shape of the head, and the manner in which the teeth are placed in the lower jaw. The typical crocodile has a narrow, triangular head, terminating in a rounded point. The head of an



1. GAVIAL.

2. ORINOCO  
CROCODYLE.3. FLORIDA  
CROCODYLE.4. INDIAN  
CROCODYLE.5. MISSISSIPPI  
ALLIGATOR.

along the top with a row of lofty, saw-toothed scales of great use in swimming.

The head is a mass of well-nigh solid bone, overlaid by the same thin layer of scaly epidermis which covers the body, of the thinness of writing-paper. The nostrils are placed far forward, near the end of the snout. The jaws possess great strength, and are armed with rows of sharp-pointed, conical teeth which are shed when worn out, and renewed.

The tongue is not free, but is firmly attached to the bottom of the mouth. Its color *never* is red, but usually is yellowish-white, and sometimes pinkish. The iris of the eye is dark green, and the pupil is very narrow, and vertical.

alligator is broad, with almost parallel sides, and at the end it is broadly rounded off. The canine tooth in the lower jaw of a crocodile fits on the outside of the upper jaw, in a notch close behind the nostrils; whereas in the alligator, the same tooth fits into a pit in the upper jaw, just inside the line of the upper teeth.

The heads of living crocodylians show wide but progressive variations in breadth, as the annexed series of figures reveal. The gavial, of the Ganges and Jumna, in northern India, has a snout like the handle of a saucepan, set with four rows of long and very sharp teeth. After the gavial of Borneo, its nearest relative is the Orinoco crocodile. At intervals come in the

Florida crocodile, the mugger of India, followed by the broad-headed West African crocodile, and ending with the alligator, widest of all.

## THE CROCODILE FAMILY.

### *Crocodilidae.*

**Erroneous Impressions Corrected.**—Regarding these reptiles, a number of the erroneous impressions which are now prevailing should be corrected. Some of them are as follows:

The true crocodiles are *not* confined to the Old World, four species being found in America.

Alligators are not wholly confined to America; for a small species exists in China.

The "movement" of a crocodile's jaws differs in no manner whatever from that of an alligator.

Only a very few species of crocodilians are dangerous to man.

So far as the author is aware, there is no authentic record of the loss of a human life by our common alligator.

All crocodilians swim with their tails, not their feet.

The skin of a large crocodilian is by no means impervious to rifle bullets. A bullet sometimes strikes a bony plate and glances off; but a proper bullet, properly placed, will penetrate the skin or armor of the largest alligator or crocodile, at any point.

The author believes that no crocodile or alligator of to-day exceeds 20 feet in length, by actual measurement; and one of that length is one out of ten thousand.

**Food.**—Crocodilians are not epicures, and some species devour all kinds of vertebrate animals that they can capture, from man to mud-hens. But the supply of obtainable mammals and birds is very limited, and fish constitutes by far the greater portion of their daily food. If all the scaly monsters of this Order were limited in food to the mammals and aquatic birds which can be seized when drinking at the water's edge, or swimming in mid-stream, they would indeed go hungry.

It is a comparatively easy matter for a large crocodilian to seize a quadruped of medium size, draw it into deep water while struggling, and drown it.

In the Reptile House of the Zoological Park, during a fight between two large alligators in the

pool, it was discovered how an alligator dismembers a bulky victim in order to devour it. An alligator seized a fighting enemy by one leg, and using his tail as a propeller, whirled himself round and round like a revolving shaft, until in about five seconds the leg was twisted off, close up to the body! That deadly rotary movement would have torn a leg from a small elephant.

On another occasion, a twelve-foot alligator named "Moses" became angry at an eight-foot companion, seized it by the body, lifted it clear of the water, and shook it until the tough skin of the back was torn in two at the joint immediately in front of the hind legs.

In the course of work among the crocodiles of Ceylon, I found that some crocodiles will eat the flesh of their own kind, and do so with genuine relish. Crocodiles which I skinned and left beside a pool were promptly eaten by their relatives, who in their turn were also killed, dissected and eaten.

**Man-Eating Crocodiles.**—Out of the nineteen species of crocodiles and alligators (eight of which I have observed in their haunts), so far as I can learn only three are dangerous to man. The most dangerous man-eater is the salt-water crocodile of the Malay Peninsula, Borneo and surrounding regions. This reptile attains a size of sixteen feet, and in the Territory of Sarawak, Borneo, it devours so many people the government has for years paid a cash reward for its destruction. Its method is to take advantage of the murky waters of the rivers, swim up to a village bathing-place, seize any man or woman found bathing in the shallow water, or filling a water-jar, and back off into deep water.

The West African crocodiles,<sup>1</sup> of Angola and other portions of West Africa, are the boldest of all crocodilians, sometimes attacking people who are in canoes, and dragging a victim from a boat. (William Harvey Brown.)

The gavial and mugger of India are harmless to man, and so are the American crocodiles, and the alligator. I have gone swimming in the home waters of both the gavial and alligator,—the two extremes in jaw development,—and therefore feel sure that both are harmless.

**Nesting-Habits.**—All the crocodilians repro-

<sup>1</sup>This is the Nile crocodile, which is widely distributed throughout Africa.

duce by laying from thirty to sixty oblong, perfectly white eggs, in layers, in a low mound of muck, or vegetable mould, or sand. The female lies in wait to defend her eggs while they hatch through the heat of the sun, or by regular fermentation. From the nest of the salt-water crocodile I have taken fifty-five eggs, from the gavial, forty-one and forty-four, from the Florida crocodile, twenty-six, and from the alligator, thirty-eight. The nest of the alligator is about two feet high and four feet in diameter.

At birth, young alligators are about eight inches long. As soon as they are out of the shell, they are wide-eyed and alert, and ready to take to the water. At this period, the muzzle is short, abnormally broad, and the arch of the forehead very high.

**Growth and Size.**—In the Reptile House of the New York Zoological Park, we have recorded the following facts regarding the rate of growth of our alligators:

	Inches.	Weight.
Length when hatched,	8	1 3/4 oz.
“ “ one year old,	18	9 1/4 “
“ “ 22 months old,	23	3 lbs.
“ “ 29 “ “	45	14 “

An alligator when received measured 6 ft. 11 in.  
 During the first year it grew 1 ft.

3 in. and measured. . . . . 8 “ 2 “  
 During the second year it grew 1 ft.

1 1/2 in., and measured. . . . . 9 “ 3 “  
 During the third year it grew 1 ft.

7 in. and measured. . . . . 10 “ 11 “  
 Length of “Old Mose,” July, 1899, 12 feet.

Length of “Old Mose,” July, 1903, 12 feet 5 in.

Judging by the rate of growth of specimens of all sizes under constant observation in the Zoological Park, where they probably are growing as rapidly as they could in a wild state, I have reached the conclusion that, under ordinary circumstances, a wild crocodile or alligator is about ten years in attaining a length of twelve feet. The average rate of growth up to twelve feet appears to be about 1.4 inches per month. After twelve feet has been attained the rate is much slower, being (in the case of our largest specimen) about two inches per year.

The secret in securing rapid growth in captive crocodilians lies in giving them a pool four feet

deep, of water warmed to a temperature of between 80 and 90 degrees F. If kept in cold water, and but little of it, they are uncomfortable, they feed sparingly, and grow either very slowly, or not at all.

**AMERICAN SPECIES OF CROCODILIANS.**

The Florida Crocodile<sup>1</sup> is the type which represents the midway average between the two extremes of the crocodilian series,—narrow-beaked gavial and broad-snouted alligator. It is a subspecies of the so-called “American” crocodile (*Crocodilus acutus*), of Central and South America, and is not found elsewhere than in southern Florida. It is the only crocodile which inhabits a country that is visited by killing frosts.

The presence of a true crocodile in Florida was not discovered until 1875, when a pair of specimens of large size were collected in Arch Creek, at the head of Biscayne Bay, by Mr. C. E. Jackson and the writer. The male measured 14 feet 2 inches (with 4 inches of his tail missing), and the female 10 feet 8 inches. Since that date, at least seventy specimens have been taken between Lake Worth and Cape Sable. Lake Worth is the northern limit of the species, but it is most abundant in the watery labyrinth of low land and shallow water where the mainland of Florida reluctantly sinks into the Gulf.

The alleged “big ‘gator” of Arch Creek was very wary, and permitted no boat to approach within rifle-shot. Even a boat completely masked by green branches, and innocently floating with the current, was enough to send the old fellow quickly sliding from his basking-place on the bank into deep water. At last, however, we shot him from an ambush in the mangroves opposite his mid-day lair, and secured him. His mounted skin is now to be seen in the United States National Museum.

The adult male Florida Crocodile is very rough, externally, and usually its natural colors have been so far obliterated by age and exposure that on its upper surfaces its color is a dull, weather-beaten gray. The females, and males under eleven feet, are of a clean, grayish-olive color,—or dull yellowish-green,—very different indeed

<sup>1</sup> *Cro-co-di'lus a-cu'tus flor-i-dan'us.*

from the funereal black of the alligator. This difference in color between our crocodiles and alligators is so marked it is quite noticeable at a distance of 200 feet, or more.

The Florida Crocodile digs burrows in the sandy banks of the Miami River, and other deep streams where the ground is suitable. These lairs are used as hiding-places, resting-places, and doubtless also as warm retreats in which to escape the cold waves from the north, which about once every five years produce killing frosts as far south as Miami.

that he has become very expert in making captures. For \$50 he will at any time take out a party of "tourists," go to a Crocodile's burrow, and with a noose, capture the reptile alive and unhurt. In each case he guarantees that the Crocodile shall exceed nine feet in length. He locates the burrows in advance, by probing for them in the sand, with a sharp-pointed iron rod. With this iron rod, the reptile is driven out of its lair, and rarely does Joe fail to make a capture "as advertised."

Many other persons in Florida have captured



E. R. SANBORN, Photo, New York Zoological Park.

#### FLORIDA CROCODILE.

The entrances to these burrows are either under water, or half submerged, and they extend into the bank from ten to thirty feet. At their extremity, they are widened out sufficiently to permit the owner to turn around. Usually, the banks are so low that the top of a burrow is only about two feet below the surface.

This burrowing habit of the Florida Crocodile has led to a very droll and uncommon industry. A young man named "Alligator Joe," of Palm Beach and Miami, knows his game so thoroughly

crocodiles and alligators in their burrows, by means of a long pole of tough wood with a strong and very sharp iron hook lashed on one end. When this pole is thrust into a burrow the reptile bites it viciously, and holds on stubbornly. But even if inclined to let go, the sharp hook engages the tongue or other portions of the mouth, and thus the creature is dragged by sheer force into the hands of his captors, and bound with ropes.

**The Cuban Crocodile**<sup>1</sup> is a small species,

<sup>1</sup> *Croc-o-dil'us rhom'bi-fer.*

with a narrower head than the preceding, and with two more rows (six in all) of bony plates along its back. It is the smallest species of crocodile now living, and so far as I have observed, also the most savage in disposition. It is olive green in color, slender in form, quick as lightning in some of its movements, and much given to roaming overland, or following up tiny watercourses, in search of new hunting-grounds. I once shot a full-grown specimen in a very small brook, near the geographical centre of the Isle of Pines, Cuba, and saw others in a salt-water lagoon on the north shore of that island. So far as known, it is not found elsewhere than in Cuba.

**The American Crocodile** inhabits the northern coast of South America, and the Gulf coast of Central America, up to Mexico. In the lagoons along the coast of Colombia, a short distance eastward from the mouth of the Magdalena River, there are small bays so thickly infested with reptiles of this species, and of such great size, that very courageous men of my acquaintance have not dared to enter in a small boat.

**The Orinoco Crocodile**<sup>1</sup> is marked by a very narrow snout, by which character it is but two places removed from the slender-beaked gavials of India and Borneo. In 1876 I found this species abundant in the Orinoco River, seven miles below Ciudad Bolivar, and killed a twelve-foot male specimen which was undoubtedly very old.

Of the **Caimans**, there are five species, all of which strongly resemble our alligator, and inhabit Central and South America, and portions of the West Indies. The *Eyebrowed Caiman* has the widest distribution, and is found from southern Mexico to the Argentine Republic. The *Black Caiman*, of the Guianas and Brazil, is the largest, and is said to attain a length of twenty feet. (Bates.) The *Rough-Backed Caiman*, of the Upper Amazon, is said to be quite small—only six feet in length. (H. Gadow.)

**The Alligator**<sup>2</sup> is so well known it needs no particular description. In individuals over eight years of age, and ten feet in length, the eight yellow bands around the tail practically though not wholly disappear, and from that time on the animal is of a uniform dull black color above, and dirty yellow or white below. I never

saw a living specimen larger than "Old Mose" (12 ft. 5 in.), and only one mounted skin which exceeded fourteen feet. That one measured 16 feet 3 inches, and is believed to be in a museum in Louisiana.

The Alligator finds its northern limit in southeastern North Carolina. From thence its range extends southward along the Atlantic and Gulf coasts to Cape Sable, the southern point of Florida, and westward through the Gulf states to the Rio Grande in southern Texas. Twenty-five years ago, this reptile existed in certain portions of its range, especially Florida, in great abundance; but about that time Alligator leather became fashionable, and the demand thus created has reduced the visible supply of Alligators by about 98 per cent. To-day you may travel from Jacksonville to Miami without once seeing the black line upon the water which betokens the existence of an Alligator; and an experienced Florida hunter has declared his belief that there is not now living in that state a specimen as large as "Old Mose" of the Zoological Park.

The habits of this reptile are quite similar to those of crocodilians generally. In Florida it burrows in sand-banks precisely like the Florida crocodile, and builds a mound of earth, moss, and grass about two feet high, in which it lays from twenty to forty eggs.

The Alligator is the only Crocodilian I ever heard utter a vocal sound of any kind. The bellow of this animal, however, is well known. Every day, regularly when the whistles blow, the five Alligators in our Reptile House lift their heads out of the water at an angle of 45°, and bellow, or roar, in concert, four or five times, making a truly unearthly noise. "Old Mose" is an excellent living understudy of "Pfafner," the bellowing dragon of Wagner's "Siegfried."

**The Chinese Alligator** was discovered in 1870 by Swinhoe, and described by Fauvel in 1879. It is quite strange that the nearest living relative of our Alligator should live in the Yangtse-Kiang River, in China; but it appears to be true. It is a small species, only about six feet in length, of a greenish-black color, dotted with yellow. A specimen in the author's possession so closely resembles our American species that specific differences are difficult to point out.

<sup>1</sup> *Croc-o-di'lus in-ter-me'di-us.*

<sup>2</sup> *Al-li-ga'tor miss-is-sip-pi-en'sis.*

CHAPTER XXXVII

THE ORDER OF TORTOISES, TERRAPINS, AND TURTLES

CHELONIA

Surely there are few intelligent persons to whom a live turtle does not appeal. From the impregnable box tortoise to the grim alligator terrapin, the giant tortoise of Galapagos, and the marine monsters of the Gulf Stream, the diversity in form and habit is very great. Fortunately, however, a fixed knowledge of twelve species will give a very good foundation on which to build up this Order.


is called the *plastron*. The carapace of a male tortoise is hollowed out underneath, but that of the female is flat. The shell has an opening at the front end to receive the head, neck and fore-legs, and the rear is open to afford space for the hind legs and tail. The shell of a turtle is a city of refuge, into which its owner withdraws its head and feet whenever threatened by enemies. In some species the shell is a remarkably perfect means of defence.

These reptiles have no teeth, but the edges of their strong, horny jaws are sufficiently sharp and chisel-like to enable them to cut up vegetable food. The head and neck move freely, in and out. The skin is very tough and leathery. Like other reptiles, the members of this Order reproduce by laying eggs and burying them, to be hatched by the sun. Some of the large tortoises live to the greatest age attained by any living creatures now on the earth.

In the original classification of the land-going tortoises, and the water-loving turtles and terrapins, it was an unfortunate mistake that the name "tortoise" was not limited to the dry-land species, "terrapin" to the hard-shelled species inhabiting fresh water, and "turtle" to the species with flippers which inhabit the sea. To-day the names "tortoise" and "turtle" are applied so indiscriminately through all three of the groups mentioned, they are useless as distinctive titles and the mixture is mischievously confusing. In the interest of common sense I therefore propose the following revised system of these common names:

1. All Chelonians of the land only shall be called . . . . . *Tortoises*.
2. All Chelonians of fresh-water shall be called . . . . . *Terrapins*.
3. All Chelonians of the sea to be called . . . . . *Turtles*.

To this at least one person will henceforth try to adhere.



SKELETON OF A FALSE GEOGRAPHIC TURTLE  
(*Graptemys pseudogeographicus*.)

P, plastron,	Hu, humerus,
C, carapace,	Tib, tibia,
Ra, radius,	Fib, fibula,
Ul, ulna,	Fe, femur,
	Sc, scapula.

**General Characters.**—A member of the Order of Turtles is a reptile which has its skeleton on the outside of its body, and its vital organs completely encased in a box of bone, called a *shell*. The top half of the shell is called the *carapace*, and it is formed by the widening of the ribs until they grow together and firmly unite wherever their edges meet. The lower portion of the shell

The following is a common-sense grouping of the members of the Order Chelonia, as found in North America and the seas adjacent:

3 inches, with a height of shell of 20 inches, was estimated to be 400 years old! Its age was estimated by comparison with other Giant Tor-

**SYNOPSIS OF THE ORDER OF TORTOISES AND TURTLES.**

ORDER.	SUBORDERS.	FAMILIES.	EXAMPLES.
AMERICAN CHELONIA.	LAND TORTOISES:	COMMON TORTOISES, BOX TORTOISES, <sup>1</sup>	<i>TES-TU-DINI-DAE.</i> . . . { Giant Tortoise. Gopher Tortoise. Common Box Tortoise.
			<i>KIN-O-STER'NI-DAE.</i> . . . Musk-Terrapin.
	FRESH-WATER TERRAPINS:	SMOOTH-SHELLED TERRAPINS,	<i>E-MYD'I-DAE.</i> . . . . { Painted Terrapin. Wood-Terrapin. Diamond-Back.
			SNAPPING TERRAPINS,
		SOFT-SHELLED TERRAPINS,	<i>TRI-O-NYCH'I-DAE.</i> . . . Soft-Shelled Terrapin.
	SEA-TURTLES:	HARD-SHELLED,	<i>CHE-LON'I-DAE.</i> . . . . { Hawksbill. Green Turtle.
		LEATHERY-SHELLED,	<i>DER-MO-CHE-LYD'I-DAE.</i> Harp-Turtle.

**THE TORTOISE FAMILY.**

*Testudinidae.*

The group of tortoises contains many species that are either beautiful, or curious, or remarkable for their size and age. Quite a number of species are handsomely colored, but the majority are perfectly plain.

Two distinct types have been developed. The ordinary, thick-shelled, uncolored tortoises, some of them of great size, constitute the majority of the species. The smaller section is made up of small tortoises, some of which have a practical hinge in the centre of the lower shell. These are strictly land-going animals, and some of them even burrow in the earth, in sandy situations where digging is easy.

The Giant Tortoise<sup>2</sup> is a good species to lead this entire Order. If there be aught in the theory of "the survival of the fittest," then this creature is clearly entitled to the leading position. A specimen at the New York Zoological Park, which weighed 310 pounds, and whose shell measured on its curves 4 feet 7½ inches by 4 feet

toises which, according to authentic history, have been in captivity over 100 years.

This wonderful creature lived all save the last two years of its life on the Galapagos Islands, a group of burnt-out volcanoes, and mountains of rock covered with brush, cacti and lava, directly on the equator, 500 miles west of Ecuador. Six species of Giant Tortoises inhabit that archipelago, living chiefly upon cacti and coarse grass, but all of them are now being exterminated at a very rapid rate either for the paltry amount of oil they contain, or a few pounds of meat from each. An ignorant cattle-herder thinks nothing of killing a Tortoise 300 years old for three pounds of meat, nothing more! In the interests of science, and her own reputation, Ecuador should prohibit henceforth the wanton and wasteful killing of those remarkable creatures.

With the exception of the crocodilians, the Giant Tortoises inhabiting the Galapagos Islands, and two islands in the Indian Ocean, are the only survivors of the famous reptilian age, when a warm atmosphere heavily charged with moisture called forth luxuriant vegetation, which nour-

<sup>1</sup> By some authorities on the classification of reptiles, the Box Tortoises are placed in the Family *Kinosternidae*, one of the divisions of the Fresh-Water Terrapins. If this arrangement should be followed, it would take the Box Tortoises out of the group of Land Tortoises, where they really belong. With this explanation the author elects to preserve the very useful arrangement into land, fresh-water and marine groups, as set forth above, and leave the Box Tortoises in the Family *Testudinidae*.

<sup>2</sup> *Tes-tu'do vi-ci'na.*



ished a marvellous series of gigantic reptilian forms. Beside some of these extinct creatures our largest reptiles are mere pygmies, and to-day they are equalled in bulk only by the rhinoceros, hippopotamus, elephant, and whale. The great Brontosaurus, whose fossil remains were found in the bad-lands of Wyoming, was sixty feet long, and some of the great Dinosaurs, or kangaroo-like lizards, stood over thirty feet in height!

Beside the Giant Tortoises, our **Gopher Tortoise**,<sup>1</sup> the largest allied species of tortoise we possess, seems insignificantly small. The largest specimens weigh only fifteen pounds. This

Excepting these and similar forms, the small Chelonians find refuge from danger in the watery depths of the ponds and streams they inhabit. The Box Tortoise, however, formed for life on land, is so small it has required a special invention for its protection.

Its shell is high, and contains sufficient room to permit the head, legs and tail to be fully withdrawn within it. Across the centre of the lower shell, or plastron, a practical double hinge has been provided. Thus, in time of danger, the animal completely withdraws its head, legs and tail, at both ends it draws the lower shell tightly against the upper, and all the soft parts are en-



BOX TORTOISES.

species is found from South Carolina to Florida, and westward to Texas. It has a very thick and strong shell, and burrows in the earth of the sandy pine-forests in which it lives. Its shell is smooth, and unmarked by bright colors, and its flesh is palatable food.

**The Box Tortoise**<sup>2</sup> is, to my mind, one of the small wonders of Nature, the special purpose of which is to point out how far "specialization" can go in fitting an animal to survive. After all, the most interesting things about animals are the lessons they teach bearing upon the development of the world and its inhabitants.

<sup>1</sup> *Tes-tu'do pol-y-phe'mus*.

<sup>2</sup> *Cis-tu'do carolina*.

tirely out of reach, behind strong walls of bone. The box of bone is as tightly closed at all points as a strongly made cigar-box with the cover nailed down.

The Box Tortoise is an illustration of the fact that several species of tortoises are quite handsomely colored, in geometric patterns of black or red lines, on lighter ground-colors. A representative specimen of this species is covered with an open fret-work of black bands laid in a mechanical pattern on a lemon-yellow ground-color. North of the range of the gopher tortoise, the Box Tortoise is our only genuine tortoise,—living only upon land, and never inhabiting water. It is common all around New York City, and is

found even in the large northern parks, where it inhabits the well-shaded forests in situations as remote as possible from the paths of men. The moist valleys of the Zoological Park have yielded many fine specimens to the Reptile House collections, where they live contentedly. The Carolina Box Tortoise is found throughout the eastern United States from the Atlantic coast to the Mississippi River, and in the South is called the Pine-Barren "Terrapin."


### THE MUD-TERRAPIN FAMILY.

#### *Kinosternidae.*

The Family *Kinosternidae* was invented for the special accommodation of the box tortoises, with plastrons hinged across the middle; but in an unguarded moment the Mud-"Turtle," Musk-

is aquatic insects, minnows, fish-eggs, worms, and in fact any fleshy creature slow enough to be caught and small enough to be eaten.

The Musk-"Turtle," or Terrapin, is possessed of a very noticeable musky odor, which serves better as a distinguishing character in the living specimen than its very dull color and general commonplace-ness of external appearance. Sometimes it shows a few spots; and the neck bears two stripes, one starting above the eye, the other below it. The plastron shows a slight tendency toward a practicable hinge, but it is only a suggestion, for the shell is practically rigid, and incapable of closing. This species, like all the terrapins of the North, burrows into the mud of pond-bottoms at the approach of winter weather, and lies dormant, with the functions of Nature suspended, until spring. It is found abundantly in the eastern United States, and ranges westward into Illinois.



MUSK-"TURTLE."

"Turtle" and similar terrapins with fixed plastrons were included. To-day, oddly enough, there is a decided inclination to leave the Box Tortoise in the Tortoise Family—where they belong, and leave the Musk-Turtle and his nearest relatives in possession of the abandoned order. But to the general student, all this is of but momentary interest.

The Musk-"Turtle,"<sup>1</sup> or Stink-Pot, has been loaded down with names in two languages which proclaim a smelly character. It is a commonplace little terrapin about six inches long, inhabiting quiet ponds or sluggish streams, basking in the sun when it is safe to linger above high-water mark. Occasionally it so far forgets itself as to swallow a worm-baited hook, and bring on trouble of two or three kinds. Its regular food

### SMOOTH-SHELLED TERRAPINS.

#### *Emydidae.*

Numerous indeed is the company composing the group of pond and river Chelonians, which live half in and half out of the water. They vary in size from the little musk-terrapin, no larger than the palm of your hand, to the big alligator-terrapin, of Louisiana, with a shell 23 inches long, and a gross weight of 115 pounds, or more. There are many species that are valuable as food, and one which is now accepted as the symbol of epicurean luxury. As usual, only the types of greatest importance and widest distribution will be mentioned here.

If it were necessary to choose a single species to represent the many species of North American Terrapins, that choice might well fall upon the **Red-Bellied Terrapin**,<sup>2</sup> or "Slider." This is a species above the average size. The largest specimen in our collection weighs 10 pounds, and its shell is 13 inches long by 9 inches wide, axial measurement. It is handsomely and plainly marked by its back of umber brown, and reddish-white under-surface. It is alert and active, its distribution is wide, and its flesh is excellent. When you go to a restaurant and order diamond-backed terrapiu, at a dollar a plate, you may know to a certainty what you are eating and paying for.

<sup>1</sup> *Ar-o-mo-chel'ys o-dor-o'tus.*

<sup>2</sup> *Pseu'de-mys ru-bri-ven'tris.*

Nine times out of ten it is Slider, no more, no less; and a very good dish it makes, too.

Of the genus to which this animal belongs, there are in North America at least six other species, all of them habitants of the southern half of our country. The Slider ranges northward only as far as Delaware, and the Susquehanna River in Pennsylvania, but is frequently seen in the New York markets. Of the terrapins that are in the habit of sunning themselves on logs within diving distance of rivers, creeks or ponds, this species is, I think, the largest we are accustomed to see. Even at quite a distance it can be recognized by the height and narrowness of its shell, as compared with species of other genera.

**The Painted Terrapin,**<sup>1</sup> hitherto called at random the **Painted "Turtle"** and **Pond-"Tortoise,"** is perhaps the most widely distributed species, and the one available to the greatest number of school-rooms, in the United States. It inhabits the whole region east of the Mississippi River except the extreme southeastern states, or about one-half of the entire country. Its shell is from 6 to 8 inches in length, and its contour is rather flat. The plates of the carapace are greenish-black, edged with yellow, and those around the margin are marked with bright red. The under shell (plastron) is yellow with brown markings; and the legs and tail are dark brown, marked with bright red lines. The upper jaw is notched in front.

This small boy's favorite is a very common species, and nine times out of ten when a nice, well-behaved little Terrapin is seen sunning itself on the hurricane-deck of a derelict log, ready to drop into the water with a gentle *plash* when Small Boy approaches dangerously near, that is it. It is called the Pond-Terrapin because it dislikes the nerve-wrecking hilarity of a river which rushes past at two or three miles per hour, but prefers a nice, quiet little 4 × 5 pond, where it can vegetate quite unmolested. In captivity its food consists of chopped fish and meat and angle-worms.

**The Ellachick,**<sup>2</sup> of the Pacific slope, from the Sierra Nevadas to the coast, and from southern California to Vancouver, is the most important species in that region. It is good for food, and

<sup>1</sup> *Chrysemys picta*.

<sup>2</sup> *Chelonia mydas*.

is frequently seen in the markets of the large cities on or near the coast. It is about the size of the painted terrapin.<sup>3</sup>

**The Diamond-Backed Terrapin**<sup>3</sup> of the salt marshes is, most unfortunately, famous for the flavor of its flesh, and its association with champagne. From the unlucky day when the epicures of Maryland pronounced terrapin stew a particularly delicious dish, the doom of this species has been sealed. Its price has risen from the original 25 cents each for large ones to \$70 per dozen for small ones, and the supply is rapidly dwindling to nothing. It is now a difficult matter for a zoologist to procure for exhibition a specimen that is more than half grown.

In appearance the Diamond-Back is neither beautiful nor striking, and in flavor I think it has been greatly overpraised. At the same time



PAINTED "TURTLE."

A good example of the Smooth-Shell Terrapins.

as reptiles go (for human food), its flesh is really very good; but, with all the good things that go into a terrapin stew, and champagne for sauce at three-fifty a bottle, almost any animal would taste good.

The Diamond-Back Terrapin is a habitant of salt water, and at one time was found in the shallow bays and salt marshes along our Atlantic and Gulf coast from Massachusetts to Texas. Chesapeake Bay has always been a sort of centre of abundance of this species, and when it flourished the markets were supplied chiefly from the region lying between New York and Pamlico Sound.

This Terrapin is small, rather flat, rounded

<sup>3</sup> *Malaclemmys palustris*.

in outline, and its scales are marked by independent black patterns composed of many geometric figures, placed one within another. A



WOOD-“TURTLE” (*Chelopus insculptus*).

Back rugose. An exception to the rule of Smooth-Shelled Terrapins.

specimen with a plastron seven inches long; and weighing a pound is a large one. Formerly the great majority measured between 4 and 5 inches; but now, it is difficult to find one large enough to make a “count” by the old standard. A “count” Terrapin must measure 5 inches (in some markets it is 6 inches) in the length of the lower shell.

Beyond reasonable doubt, the continual destruction of the largest specimens will erelong render the species unproductive, and it will cease to exist. The persistent destruction of fathers and mothers will soon wipe out the stroughest species in existence. It is reported, however, that in the South there are several terrapin “farms” on which this species is being bred and reared for the markets, in large numbers.

**THE SNAPPING TERRAPINS.**

*Chelydridae.*

**The Alligator-Terrapin,**<sup>1</sup> of Louisiana, and other states bordering on the Gulf between Flor-

<sup>1</sup> *Mac-ro-chel'ys tem-minck'i.*


ida and Texas, is, when adult, a huge, rough-backed, big-headed creature, weighing from 100 to 125 pounds, and even attaining on rare occasions to 150 pounds. This is the largest terrapin in North America, and also the ugliest. The broad and rather flat table of its upper surface rises in a series of brown hillocks, earthy-looking, and often actually covered with moss.

The head is of huge proportions, and the strength of the jaws is very great. The tail is very long and fleshy,—which is rather unusual in Chelonians. Notwithstanding the rough exterior of this creature, its flesh is eaten by many persons who share its habitat.

This remarkable reptile is found only in the semi-tropical fresh-water bayous and streams of the South. A specimen now living in the Reptile House at New York measures as follows:

Length of head and neck . . . . .	12 in.
“ “ shell . . . . .	23 “
“ “ tail . . . . .	19 “
Total length . . . . .	54 “
Width of shell . . . . .	18 “
Weight . . . . .	113½ lbs.

It is a shy animal, and if not permitted to live under the crocodile’s raft which floats in the pool, it will not eat its usual daily ration of raw meat or fish. It never attempts to leave the water, and can remain submerged, without breathing, for periods which are so long we can only describe



ALLIGATOR-TERRAPIN.

From Louisiana. Weight, 113½ lbs.

them as “indefinite.” In its home, this burly reptile feeds upon fish, frogs, and other water-animals.

**The Snapping Terrapin, or Snapping "Turtle,"**<sup>1</sup> which is found in the northern states as well as in the South, is a very cross-tempered and savage understudy of the preceding species, and it is ugly in more senses than one. It has a humpy, moss-covered back, a mean eye, a dangerously sharp and hooked beak like a horned owl, and a tail that reminds one of the terminal half of a bloated water-moccasin.

This reptile seldom leaves the waters of the ponds in which it lives. It believes most thoroughly in the survival of the fittest, and to it the Fittest is "Number One." It is a chronic fighter, and inasmuch as its jaws are very strong,—and, like some men, never know when to let go,—it is a reptile to be either mastered or avoided. It is wholly carnivorous in its habits, and is very destructive to fish and young water-fowl.

Strange to say, the Snapping Turtle is regularly consumed as food, and is often sold in the Centre Market at Washington.

## THE SOFT-SHELLED "TURTLES."


### *Trionychidae.*

This Family is of ancient lineage, and wide distribution, its members being found in the rivers of Asia, Africa and North America. Wherever found they may be recognized by very flat and nearly circular shells that are imperfectly ossified, both above and below, and which terminate at the edges in thin plates of leathery skin. The nose is prolonged into a decided proboscis, and the neck is long and flexible. In some species (found in Australia) the neck is so very long it cannot be withdrawn into the shell, but in times of danger it is laid away snugly under the upper edge of the shell, passing over one fore-leg.

The members of this Family present many anatomical exceptions to the regular order of form among tortoises and terrapins, and by some authors they are placed at the foot of the Order Chelonia. The shell is really very im-

perfect, the bones being literally few and far between, and the upper and lower shells are quite unconnected by bony structure. The feet are large and strongly webbed, but only the three inner toes are provided with claws. In habit these creatures are persistently aquatic, rarely going upon dry land, and they are both voracious and carnivorous. They live upon fish, fish-eggs, frogs, angle-worms, and small mollusks generally.

**The Soft-Shelled "Turtle"**<sup>2</sup> is perhaps the most common representative of this Family in



SOFT-SHELLED "TURTLE."

*Aspiderochelys ferox*, from Florida.

the United States. It is found from South Carolina westward through the Gulf states to Texas; up the Mississippi to Indiana, Illinois and the Great Lakes, north and westward up the Missouri to the Rocky Mountains.

I never shall forget those I encountered in central Indiana, when fishing with hook and line. The provoking Soft-Shells would persist in swallowing hooks that were not baited for them, and the difficulties we had in cutting off their leathery heads and dissecting out our hooks tried our patience very sorely. It was not until many years later that we squared accounts with this species. At Miami, Florida, fine large specimens were fried in batter, and eaten with great relish. When properly cooked, the shell of this reptile is tender, tasty and desirable.

<sup>1</sup> *Che-ly'dra ser-pen-ti'na.*

<sup>2</sup> *As-pi-do-nee'tes fe'rox.*

A large specimen has a shell 16 inches long by 14 inches wide, and weighs from 20 to 30 pounds. The upper surface is olive brown mottled with black, and underneath is clear white. On account of its widely palmated feet, these "turtles" are the most active swimmers of all the fresh-water terrapins and turtles. In North America this Family is represented by five species.

### THE SEA-TURTLES.

The sea is so vast, it is but natural that we should look to it for the largest species of Chelonians. There is one character by which any one can recognize a sea-turtle, anywhere. The front limbs are developed as *long, flat, triangular flippers, without separate toes and claws, like the flippers of a sea-lion.*

Nearly all the sea-going Turtles are large, and one species is the largest of all living Chelonians. Without exception, all are habitants of tropical waters; but occasionally an individual is lulled into fancied security, and borne northward in the warm waters of the Gulf Stream until it wanders out of the track, and suddenly finds itself in the chilly arctic current. Then, benumbed with cold, it falls an easy prey to the first predatory fisherman who sails near it, and promptly lands in Fulton Market.

### HARD-SHELLED SEA-TURTLES.

#### *Chelonidae.*

**The Green Turtle**<sup>1</sup> is the most important and valuable of the sea-turtles, and in the Atlantic it is the species that is most widely distributed. It is of large size, its flesh is excellent food, and wherever found it is regarded as a prize. It is said that sometimes it attains a weight of about 600 pounds; but those which now find their way to market in our large cities are steadily diminishing in size, and rarely exceed fifty pounds.

This turtle is found from Long Island down the Atlantic to Cuba, throughout the Gulf of Mexico, the Caribbean Sea, the West Indies, and on southward to Brazil. It is also found in the Indian Ocean, and is common on the coast of Ceylon. I should say that on our coast Key West is its centre of greatest abundance and maximum size. The favorite haunts of this creature are in the shallow channels that lie be-

tween the keys, where they find quiet waters and plenty of food, but no security from the sharp eyes of the turtle-catchers. It feeds upon aquatic plants that grow on the bottom of shallow seas.

A large proportion of the Green Turtles captured on the Florida coast are sent north, by steamer and rail, to supply the ever-greedy and high-priced city markets from Baltimore northward.

And really, it is not surprising that the flesh of this animal is considered most excellent food, and much sought after, both for soups and steaks. It is tender, fine-grained, dark colored, not too fat and very agreeable in flavor. Moreover, this is a clean-looking animal, its shell is smooth, its head is small and neatly formed, and the front flippers are scaled, quite down to their extremities. The shell is of no commercial value.

**The Hawksbill Turtle, or Tortoise-Shell Turtle,**<sup>2</sup> furnishes the valuable tortoise-shell of commerce, and it is the most beautiful of all the Chelonians. Its name is derived from the strongly hooked beak which terminates its upper jaw. Its back is covered with a roof of very beautiful curved plates of tortoise-shell, overlapping like shingles, each scale terminating in a saw-tooth point. The scales are clear yellowish horn, beautifully mottled with black and brown.

This species is yet found occasionally around the Bahama Islands, where the sea is very clear, and the white-sand bottom is liberally garnished with sea-fans, corals, and other beautiful invertebrate forms. Its range as a whole is from the coast of southern Florida, the Bahamas and the Gulf of Mexico, southward through the West Indies to the Amazon. It also inhabits the tropical waters of the Old World.

Formerly it often grew to a weight of between twenty and thirty pounds, but it has been so persistently sought after, on account of the commercial value of its shell, that all those now seen in the markets are very small. The largest shell on record is 34 inches long. Another species is found on the Pacific coast, and it bears so strong a resemblance to its eastern relative that for a considerable period the two species were believed to be identical.

**The Loggerhead Turtle**<sup>3</sup> looks like a coarse

<sup>1</sup> *Che-lo'ne im-bri-ca'ta.*

<sup>3</sup> *Thal-tas-so-chel'ys car-el'ta.*

<sup>1</sup> *Che-lo'ne my'das.*

and large-headed understudy of the green turtle. It is readily distinguished, however, by its massive head, and thick, heavy shell. It is a turtle of coarser quality every way than the green turtle, and sells at a lower price. Like its handsomer relative, it is widely distributed, but does not inhabit the Indian Ocean.

The flesh of this animal bears so close a resemblance to beefsteak that even a butcher cannot always detect the difference. One Christmas morning, at Key West, I dissected a large Loggerhead. The flesh was fresh, and very tempting, and when a choice lot of steaks were offered to the landlady of a certain small hotel, they were gratefully accepted.

It happened that the butcher who supplied the hotel with beef and mutton was a boarder thereat; and, as became his calling, he sat at the head of the long table, and served the meat. Although he was an able butcher, he had one weakness; and it lay in the fact he could *not* eat turtle-meat. It was "too oily," too "musky," and too far removed from beefsteak.

With no unnecessary announcements, the turtle-steaks were fried, *à la* beefsteak, and set before the butcher. He served them as beefsteak, ate his own portion with evident relish, and all the other guests ate theirs. The butcher had nearly finished his second instalment, without having discovered the substitution, when he was asked how he liked turtle-steaks, for a change.

The sandy beach on the east coast of Florida, along the Indian River Peninsula, is a favorite spot for both Loggerhead and green turtles to lay their eggs. Mrs. C. F. Latham, of Oak Lodge, ninety miles above Palm Beach, has made careful observations on the habits of these

turtles. In the months of May and June, when the summer heat is becoming severe, on moonlight nights the turtles crawl up out of the water, dig holes in the sand high above tide-mark, from 15 to 18 inches deep, and in them lay their eggs, to the number of from 80 to 220. The period required for incubation is about sixty days. When first hatched the young are only 2½ inches long, but the moment they emerge from the nest they start for the ocean.

### LEATHERY-SHELLED SEA-TURTLES.

#### *Dermochelydidae.*

The **Harp-Turtle**, or **Lyre-Turtle**,<sup>1</sup> is the giant of the Chelonians of the present day. Sometimes it is called the **Leather-Backed Turtle**. I once dissected and preserved a specimen which weighed 740 pounds, and the oil and the toil of it are yet vividly remembered.

This remarkable creature has a very feeble bony shell, which is buried under a one-inch layer of fatty material which looks quite like the blubber of a whale. It is easily cut with a knife, and contains about a pint of oil for every square foot. The back of this strange creature is marked by five sharp ridges that run lengthwise, and are separated by concave, wave-like depressions. The front flippers are very long, and it seems quite certain that even in its native element this great animal is slow and clumsy. Its flesh is quite unfit for food.

This turtle is found very sparingly along the Atlantic coast from Long Island southward, but is abundant nowhere. One may travel all around Florida, and all through the West Indies without seeing even one specimen.

<sup>1</sup> *Sphar'gis co-ri-a'ce-a.*



Photographed and copyright, 1903, by R. J. Beck.

MARINE IGUANAS ON NARBOROUGH ISLAND, GALAPAGOS ARCHIPELAGO.



CHAPTER XXXVIII  
THE ORDER OF LIZARDS

LACERTILIA

Of all the world's reptiles, the lizards are the most elusive, and the most difficult to know personally. With the exception of the large iguanas, monitors, and a very few others, the vast majority of the species are tiny creatures, lightning-quick in movement, and very much opposed to being caught.

And the little sprites are difficult to keep in captivity, beyond all other reptiles. Being children of the sun and sands, they demand quarters that are of desert dryness, roasting heat,

partly or wholly covered with spines. The majority of lizards live upon the ground, or near the earth, but quite a number of species live in trees. Those called flying dragons possess parachute wings, and can fly as a flying squirrel does. Some of the legless lizards live in the earth.

Most lizards have teeth, but usually of a very simple character, setting in each jaw in a long and rather even row, like the teeth of a saw.

There are eighteen Families of lizards, provided, with eighteen formidable names, and it is not



COMMON IGUANA.

and flooded with sunshine. Without these conditions they refuse to eat, and quickly die. If every student of lizards had a private desert which he could keep heated up to 100 degrees, a sun all his own to shine upon it sixteen hours a day, and meal-worms without limit, it would be quite possible to keep small lizards long enough to become well acquainted with them. Without such an equipment, the path of the student is beset with difficulties.

Because of these conditions, we will introduce here only a very few of what we may call the practicable lizards,—those which it is possible to know, and worth while to note.

**General Character.**—Most of the lizards are four-footed creatures, many have long, whip-like tails, and nearly all are covered with scales, mostly very fine. Sometimes the scales are large and horny. Quite a number of species are either

possible to consider each one. For the present we will omit all references to the Families, and merely present a few examples which will illustrate the Order as a whole.

**The Iguanas** are among the largest and most interesting of the Lizards, being surpassed in size only by the *Monitors*. In their food habits they are omnivorous. Although feeding chiefly upon vegetable food, many species devour birds and eggs with great avidity. In their habits they are partly tree-climbing and partly terrestrial. By reason of their saw-toothed backs they are so odd and showy they always attract attention. Were it necessary to select but one species to represent all the species of Lizards, that one should be the **Common Iguana**<sup>1</sup> of the West Indies, Central and South America. It is from 4 to 5 feet in length, in color it is an irregular

<sup>1</sup> *I-guan'a tu-ber-cu-la'ta*.

mixture of green, black and yellow, and it may be recognized at a glance by the row of long, slender, fringe-like scales which rise along the centre of its back. One good look at its extremely long and slender toes is enough to suggest the idea that it is a climbing animal. It makes its home in thick tree-tops, and feeds chiefly upon fruit and soft vegetation. I can testify that its flesh is palatable food, for in the hungry Orinoco country we ate it more than once.

Iguanas generally possess one good trait which is sufficient to forever endear them to zoological garden people. They are good-tempered animals, and never fight, no matter how many are placed in one cage, nor how many species of Iguanas are represented. Owing to the ease with which these creatures are captured, their price in New York is about \$2 each.

**The Marine Iguana,**<sup>1</sup> or **Sea-Lizard**, of the Galapagos Islands, is a creature of gre-ga'ri-ous habits, which means the habit of flocking or assembling together in companies of noteworthy size. So far as we know, this is the only lizard which elects to assemble in companies of several hundred individuals. When Mr. R. J. Beck visited Narborough Island of the Galapagos group, in 1902, in quest of giant tortoises, he found on the clean lava-bed which formed the shore, a truly wonderful assemblage of Marine Iguanas. An area of at least three acres, destitute both of soil and vegetation, was literally covered by these reptiles, all wide-awake and fully interested in life, but serenely waiting for something to turn up.


Owing to their lymphatic temperament, and previous immunity from persecution by man, these strange creatures were quite tame, and willingly permitted Mr. Beck to make the photograph that is reproduced herewith. It represents one of the most wonderful views of reptilian life to be found anywhere on the earth to-day.

The Marine Iguana is a stockily built, dull-colored animal, about four and one-half feet in length, frugivorous in its habits, and very much at home in the water. It subsists almost wholly upon sea-weed.

**The Rhinoceros Iguana,**<sup>2</sup> of the same form as the preceding, but much lighter in color, and

marked by half a dozen horny tubercles on the upper surface of its head and snout, is found in Hayti and San Domingo.

Leaving the large lizards, of which be it remembered there are many interesting species in the Old World,—called Monitors,—impossible to mention here, we reach the small lizards, of which there are a legion of species. The warm and dry countries of the world literally swarm with these tiny creatures, which dart over rocks and fences like streaks of green or brown light. If you try to catch one by its long tail, and close upon it, the lizard leaves its tail between your thumb and finger, as a souvenir, and gayly streaks away *to grow another*, without loss of time! The power possessed by lizards to reproduce a miss-



BLUE-TAILED LIZARD.

ing tail is one of the strangest things in animal growth; but it is to be observed that the second edition of a lizard's tail is far from being the shapely and perfect member that is seen in the first.

Many lizards, like much study, are a weariness to the flesh; and we will limit our exhibit to a very few prominent and interesting types which are well fitted to represent the entire group.

**The Blue-Tailed Lizard**<sup>3</sup> is not only a common species throughout a wide area of the United States, but it is also representative of a large number of species which resemble it. It is found throughout the eastern half of the United States, from Nova Scotia and Canada to Florida and the Gulf, westward in the South to Arizona, and in the North to Wyoming. It is often called the **Skink**, and "**Blue-Tail**," and **Blue-Tailed Skink**, and in summer it is available for study

<sup>1</sup> *Am-bly-rhyn'chus cris-ta'tus*.

<sup>2</sup> *Met-o-poc'e-ros cor-nu'tus*.

<sup>3</sup> *Eu-me'ces quin-que-lin-e-at'us*.

purposes to a larger number of school pupils than any other lizard known to the author.

The colors of this creature vary with age to an extent that is apt to be very confusing. Observe the programme:

During the first year the body is black, with bright yellow stripes, and the tail is brilliant blue. In the second, the tail is slaty gray, and the black of the body is less intense. In the third, the body becomes brownish, and the stripes are indistinct. In the fourth, and thereafter, the body is brown, the head vermilion, and the stripes have disappeared. The length of a large specimen is about eight inches.

All the small lizards and skinks are insect-eaters, and in captivity thrive best upon meal-worms and insects generally. Their quickness of movement is almost beyond belief, and even with a long-handled net it is very difficult to capture one alive and unhurt.

**The Ring-Necked Lizard,**<sup>1</sup> which should be called the **Kangaroo-Lizard**, represents a group quite different from the skinks, and also nearer to the iguanas. It is a creature of the canyons, deserts and dry mountains of the Southwest, from Texas to southeastern California, and northward into Utah and Nevada. It is often found on mountains up to 5,000 and even 6,000 feet. (Merriam.)

This is a plump-bodied creature, and its colors vary to an extent that is apt to create confusion. It is either dark green or bluish above, and the sides, back and thighs are covered with light spots. The under surface is yellowish-white, sometimes tinged with pale green. This lizard derives its name from two bands of black which stretch across the shoulders between the fore-legs.

The most interesting feature about it appears never to have been observed and recorded until Mr. Barnum Brown sent several specimens to the Zoological Park. When one was liberated in a large sanded cage, it rose on its hind-legs, in the position of an erect kangaroo, and in that strange posture ran rapidly. It held its head well erect, carried its fore-legs *à la* kangaroo, and ran, not by hopping, but by taking long steps. In experimenting with the different individuals received from Mr. Brown, it was found that under similar provocation, all of them ran in the remarkable

<sup>1</sup> *Cro-ta-phy'tus col-lar'is*.

attitude described,—highly suggestive of a pygmy dinosaur.

**The Gila Monster**<sup>2</sup> is perhaps the most famous lizard of North America, and its first name is pronounced *He'la*.

It is big, odd-looking and very showy, and therefore is dear to the heart of nearly every collector of reptiles. A large specimen has a total length of 20 inches, girth around the middle, 9½ inches, and weighs, 43 ounces. When in robust health, the body and tail seem stuffed to the point of discomfort. Externally the whole of the creature appears to be covered with round glass beads, jet black and orange yellow in color, and laid on in a Navajo pattern.

This remarkable lizard inhabits the desert regions of Arizona and the adjoining state of



From the Zoological Society Bulletin.

GILA MONSTER.

Sonora, Mexico. It is more sluggish in its movements than a box tortoise, and the very slow and clumsy manner in which it partakes of its daily meal of raw eggs and chopped meat leads the observer to pity its helplessness. How it manages to secure a sufficient quantity of acceptable food on the deserts where it lives is a puzzle.

Whether the bite of this creature is poisonous or not is yet a debated question among naturalists. Several authorities cite the deaths of various small animals bitten by it, but others point to other victims which were bitten, but did not die. At the United States National Museum, Mr. A. Z. Schindler was bitten by a Gila Monster, but aside from a very natural degree of irritation

<sup>2</sup> *Hel-o-der'ma sus-pec'tum*.

and soreness of the wound during two or three days, he experienced no permanent ill effects from it. It is quite certain that the bite of this creature is seldom fatal to man, even if it ever is; but it can cause the death of small and weak creatures, like frogs and guinea-pigs.

This reptile lives well in captivity, and half a dozen of them in a desert cage make a very showy exhibit.

**The Horned "Toad,"**<sup>1</sup> so dear to the heart of every eastern traveller on his first visit to the great Southwest, where deserts are plentiful and cheap, is not a "toad" at all! Observe its long tail, such as real toads never have, then call it forever after by its real name—**Horned Lizard**. There is much excuse, however, for the universal name; for, saving the presence of the tail, the little living cactus is quite toad-like in its form.

Professor Cope recognized eleven species of Horned Lizards, any one of which, wherever found, will serve as well as another to represent this genus. They are all habitants of the deserts and arid regions, where cacti, cat's-claw, and other thorny things possess the land. They are frequently seen in the roads and trails of the Southwest, and are easily captured. If meal-worms are abundant, they are easily kept in



HORNED LIZARD.

captivity, on dry sand, in warm sunshine. The length of a large specimen is only  $5\frac{1}{2}$  to 6 inches; and, strange as it may seem, these odd creatures are related to the iguanas.

<sup>1</sup> *Phryno-so'ma cor-nu'tum*.

No! **The Glass "Snake"**<sup>2</sup> does *not* join itself together again after it has once been broken in two. And it is not by any means a snake! It is a smooth-bodied, legless lizard, but so scaly,



GLASS "SNAKE."

and so snakelike in general appearance that any stranger is quite excusable for calling it a snake. As a matter of fact, the tail of this creature is so feebly attached to the body that a very moderate blow with a stick breaks the connection, and the reptile lies in two pieces. If left until doomsday, the severed parts will not reunite, but the body does its utmost to repair the injury by growing another tail. As a matter of fact, the new growth of tail is but a short and very imperfect substitute.

This creature inhabits the southern states from the Carolinas westward to Texas, and northward up the Mississippi valley to Kansas and Wisconsin. It feeds chiefly upon insects, and being quite without legs, it forms an excellent connecting link between the lizards and serpents. There are quite a number of species of legless lizards.

<sup>2</sup> *O-phi-o-sau'rus ven-tral'is*.



New York Zoological Park.

TWENTY-TWO-FOOT RETICULATED PYTHON (DEAD).

## CHAPTER XXXIX

### THE ORDER OF SERPENTS

**General Characters.**—A serpent, commonly called a “snake,” is a very slender, long-bodied, legless reptile, cold-blooded, covered with scales, and breathing air. It moves by a sinuous motion, in which the scales under the body grip the earth, while the extension of the body muscles push the body forward. To afford a good hold upon the earth, the abdominal scales are very broad, set crosswise with the body, and the rear edge of each scale is free and sharp, like a blade.

**The backbone** contains a great number of vertebrae, sometimes nearly 300, and there is one for each crosswise scale under the body. There are also a great number of ribs, but the tail vertebrae are of course without them. The ribs are

quite loosely attached to the vertebrae, in order that they may have the very free play that is absolutely necessary to the life of a serpent.

**The head** is usually flat and broad, and entirely covered with scales. The jaws are long, and well armed with long, sharp-pointed teeth, which point backward, in the direction of the throat. There are no molars for masticating food, and therefore all food is swallowed whole. Excepting in the injection of poison, the only function of the teeth is to seize and hold fast the serpent's prey while it is being swallowed. Poisonous serpents have special teeth, called fangs, for making deep wounds and filling them with poison. These are set in the roof of the

mouth, well forward, and while not in use they lie up against the roof of the mouth. The tongue of a serpent is very extensible, and capable of being thrust out fully half the length of the head. Its greatest use is in examining food, or possible food. From the fact that when travelling the tongue is so frequently thrust out, even when there is no excitement, it seems highly probable that it is used to detect vibrations in the air. (R. L. Ditmars.) The tongue is forked, and being entirely harmless, its sole use in defence is to threaten and intimidate its enemies.

**The lower jaws** are loosely attached to the skull, and to each other at their front end, by ligaments so elastic that when prey is being swallowed, the gape expands to enormous proportions. Mammals, birds and fishes to be swallowed are always seized head first, in order that the limbs, and also the feathers or scales, if there be any, will lie snugly against the body. Frogs and toads are usually taken hind feet first. The lower jaw is forced forward and over the animal, always one side at a time, as far as it will go; and when the teeth are inserted, that side is drawn back. The upper part of the head slides forward as far as possible, one side at a time, to match the lower jaw. Sometimes it seems as if the lower jaw will be torn loose from the head. Often after an animal has vanished, the jaws are a bad misfit, and do not come back into shape for half an hour.

The skin stretches like India-rubber, and over a heavy meal the scales are widely separated. The manner in which serpents feed in a wild state is certainly one of the most cruel processes of Nature.

**The eyes** of a serpent have no lids, and the eyes never close; but they are protected by a thin and perfectly transparent section of the outer skin, or epidermis, which is shed and renewed periodically.

**The epidermis**, or outer skin, is completely renewed about three times per year. To free itself from the old skin, the serpent usually crawls through a small aperture, the edges of which catch the old skin at the head and hold it fast while the owner crawls out of it. The first intimation of an impending change of epidermis is found in the dull appearance of the eye, over which a glassy film seems to be forming. Strange to say, even the eye sheds its outer

surface, and emerges clear and brilliant. Most snakes shed their skins about three times a year.

A serpent is always most beautiful immediately after it has shed its epidermis, for then its colors are brightest and most iridescent. In captivity it often happens that the atmosphere in which a snake lives is not sufficiently moist to enable the old skin to loosen and be cast off. In such cases, if the serpents are non-venomous species, the owner must moisten the old skin, and peel it off by hand, or with forceps.

**Reproduction.**—Some snakes lay eggs, with soft, tough shells, that are hatched by the sun. A serpent which develops in an egg of this sort is provided with a special, temporary tooth, set on the tip end of its jaw, with which it easily punctures the shell sufficiently to escape. Others do not develop eggs with shells, but instead retain their eggs in their own bodies until the young are fully developed. Finally they are brought forth, each fully enclosed in a thin, membranous sac, which the little serpent quickly bursts. Snakes that lay eggs are called *o-vip'-a-rous*, and those that bring forth their young alive are called *vi-vip'-a-rous*.

Although serpents are cold-blooded animals, they reach their highest development in warm latitudes, and in regions of arctic cold they do not survive. In the temperate zone and the tropics, Nature has fitted them for life upon the ground, in the water, and in the tree-tops; and they inhabit swamps, uplands and deserts. They live under stones and logs, in hollow trees and stumps, and in holes in the earth; and they seldom attack man wilfully, and without provocation.

**Food of Serpents.**—In a wild state, snakes feed chiefly upon frogs and toads, fish, other snakes, small birds and mammals. Large serpents feed upon mammals of all sizes, up to small deer and goats. Water-snakes feed chiefly upon fish and frogs. Land species find frogs, toads and small lizards their cheapest prey, but the extent to which snakes feed upon each other is quite surprising. For example, the king cobra,<sup>1</sup> a large, athletic, and very deadly land-serpent of the Malay Peninsula, feeds exclusively upon other snakes and lizards, and while a greedy feeder upon what it prefers, it persistently refuses all other food. During the three years that

<sup>1</sup> *Na'ja bun-gar'us*.

one of these serpents has been kept in the Zoological Park, it has persistently refused to eat any of the moccasins or rattlesnakes which have been offered to it.

This fine specimen, which is nearly 11 feet long, became, toward the end of its first winter, so difficult to provide for, when the special supply of food-snakes had become well-nigh exhausted, that Curator Ditmars and Keeper Snyder tried a novel experiment. They killed a six-foot snake, stuffed it with frogs to the number of half a dozen, then offered it to the cobra. It was immediately accepted, and devoured in good faith; and since that time the experiment has often been repeated.

A large collection of captive reptiles requires many different kinds of food, and plenty of it. It is not necessary that food should be given alive. Very naturally, a serpent cannot swallow a bird or a mammal which is stiff in death, and unyielding. Swallowing is not possible unless the legs or wings are folded very closely against the body. All that a serpent requires is that the animal be offered while yet warm, and before *rigor mortis* has set in. The practice is to kill the food in the Reptile House, and offer it immediately afterward, while it is yet warm.

During the year 1902, the Reptile House contained 33 Crocodilians, 112 Lizards, 134 Chelonians, 381 Serpents, and 112 Amphibians, and the animal food they consumed during the year was as follows:

3,550 Rats and Mice,	512 Guinea-Pigs,
1,456 English Sparrows,	About 18,000 Meal-
624 Small Chickens,	Worms,
208 Large Chickens,	About 25,500 Live Fish,
210 Pigeons,	About 2,000 Toads,
1,300 Eggs,	About 2,000 Frogs,
272 Rabbits,	About 2,500 lbs. Vegeta-
	bles and Fruit.

**Classification of Serpents.**—Unfortunately, it is impossible to offer the general student a diagram of the Families of living serpents, based on the highest scientific authorities, which would be either simple or understandable. The species are many, and their teeth, scales, bones and other features are diversified. Thus far no scientific authority has succeeded in dividing the world's serpents into logical groups without basing the divisions upon anatomical features, and describ-

ing them in technical terms which only the special student of reptiles can understand.

By way of example, take Professor Gadow's simple statement of the distinguishing characters of the Family *Colubridae*: "ectopterygoids are present: the squamosals are loosely attached to the skull, and carry the quadrates, which are not reached by the pterygoids: the prefrontals are not in contact with the nasals; the maxillaries are horizontal, and form the greater portion of the upper jaws: the mandibles lack the coronoid process or element: both jaws are toothed."

Under the circumstances, our wisest course will be to select and set forth a series of small groups of serpents which will introduce the species most worth knowing, and at the same time convey a fair amount of general information regarding serpents as a whole.

**Popular Questions and Misapprehensions.**—Regarding the habits of serpents there are many unsettled questions, and many disputes. The perennial "Hoop Snake" delusion, for example, will not down, and probably it never will lack exponents and defenders.

The question "Do snakes swallow their young?" is also a perpetual storm-centre; and there is plenty of reliable evidence on all sides of it.

Snake disputes between truthful persons are due either to deceptions of the eye (*an organ easily deceived!*), a misinterpretation of things seen, or imperfect observations.

For example, men of the highest truthfulness have been deceived into the fixed belief that they have "seen horse-hairs turn into worms."

Without attempting to settle out of hand any of the snake disputes that are "rock-ribbed, and ancient as the sun," I will at least state what experienced men, who have observed and studied reptiles all their lives, and gathered facts regarding them, *believe to be true*.

The "Hoop-Snake," which is said to travel by taking the end of its tail in its mouth, and rolling along like a hoop, is believed to be an absolute myth.

It is believed that snake mothers do *not* swallow their young in order to protect them, and emit them all as good as new, when the danger is over.

Many snakes do hiss, some of them as loudly as a red-hot poker thrust into cold water.

The tongue of a snake is not capable of in-

flicting a wound, nor of conveying poison into the blood of another creature.

Snakes never are "slimy."

Removing the fangs of a poisonous serpent does not necessarily render it harmless; for new fangs promptly grow out to take the place of those removed.

The rattle of the rattlesnake contains more than one joint for each year of life,—usually two or three.

### THE LARGEST SPECIES OF SERPENTS.

#### *Family Boidae.*

The Family *Boi-dae*, containing the boas, anacondas and pythons, embraces between sixty and seventy species.

It is as natural for human interest in ani-

before swallowing it, are *constrictors*, because of their method; but all big serpents are not necessarily *Boa constrictors*. That title applies to but a single species, found in South America; and, curiously enough, its Latin name is also its popular name.

In seizing its prey, this serpent instantly reveals its name by its method. The jaws open widely, fly forward with electric quickness, close on the animal, and hold fast. Instantly thereafter, a coil of the body near the head is flung completely around the victim and drawn tight, to suppress struggling, and prevent possible escape from the jaws. From the oldest and largest to the youngest and smallest *Boa Constrictors*, all seize their prey with precisely the same action, and the flinging of the first coil fol-



BOA CONSTRICTOR.

New York Zoological Park.

mals to be greatest toward those that are the largest of their kind, as it is for sparks to fly upward. It is well to see what Nature can do when she puts forth her best efforts. No one need apologize for a keen interest in pythons, boas and anacondas, provided that interest is kept down to bed-rock truth, and all exaggerations and overestimates are rigidly eliminated. Unfortunately, however, the makers of sensations about wild animals regard all large serpents as their lawful prey, and often stretch them unmercifully.

**The Boa Constrictor.**—The serpents which seize their prey, and crush it into compact shape

lows so quickly after the strike of the jaws that the two acts seem almost simultaneous.

The *Boa Constrictor* is much smaller than its neighbor, the anaconda, and not more than one-half the size of the gigantic reticulated python of the East Indies. Its maximum length is about 12 feet. It inhabits South America, from the Caribbean Sea to Paraguay, but only in forested regions, where animal food is plentiful, and cover for concealment is abundant. This species is readily recognized by its bright, reddish-brown tail, which is much more highly colored than the head and body. It is also marked by the prevalence of reddish, iron-rust brown in its color-



scheme, and the very large oval patches of light color, divided by black bands, that are laid along its back with regularity and precision. The sides are beautifully marked by light-colored diamonds and bars.

When at home, this serpent feeds upon pacas, agoutis, capybaras, tamanduas, young peccaries and tapirs, and any bird that is large enough to justify attention. Considering the excellent climbing powers of the Boa Constrictor, and the dulness of certain South American monkeys, it is highly probable that monkeys furnish many a meal for this serpent. The sloth is protected in two ways. It prefers the small and weak outer branches of a tree, and it moves so slowly and unostentatiously a Boa would be long in finding one.

If a twelve-foot Boa once wrapped itself around an unarmed man, it undoubtedly could suffocate him, or crush him to death, but it would be impossible for it to swallow him. There is at hand no authentic record of a Boa Constrictor ever having killed a man or a horse. In South America I was assured by native hunters that Boas and anacondas swallow antlered deer, but when direct proof of this was called for, it never came.

**The Anaconda**<sup>1</sup> is the great water-constrictor of South America, and it so loves the aqueous element that some captive specimens never leave their bathing-tanks unless forced to do so. This serpent is strongly marked for identification by the very large black spots, round or nearly so, which cover its back from head to tail, laid on a dark olive ground. Sometimes these are arranged in pairs, and suggest dumb-bells.


This species attains very great size, and being fully equal to the reticulated python of the East Indies, it is one of the largest of living serpents. Of course it can hardly happen that specimens of the largest size would find their way into zoological gardens. The largest thus far exhibited in the Zoological Park measured 18 feet 6 inches, and came from the Berbice River, British Guiana. In the British Museum there is a stuffed specimen which is 29 feet long.

In British Guiana I was assured by local hunters that the "Camudie," as this serpent is commonly called, often attains a length of 35 feet. There is, however, no proof that it exceeds 30 feet; and any traveller or observer who has the good

fortune to meet with a specimen exceeding that length will do well to back up his tape measure with either the preserved skin or skeleton. One snake-skin is more convincing than a hundred snake-stories.

I believe the delta of the Orinoco is the northern limit of the Anaconda, where it is called the "Culebra de Agua," and regarded with profound respect. It inhabits the Guianas and Brazil, and probably extends to the head-waters of the Amazon, in eastern Peru. Of its regular food, the capybara (a water-loving rodent, as large as a good-sized hog) undoubtedly stands first, followed by the tapir, otter, deer and large water-birds generally.

**The Reticulated Python**,<sup>2</sup> of the Malay Peninsula, Sumatra and Borneo, is the largest



YELLOW ANACONDA.  
*Eunectes notaeus.*

serpent of the Old World, and the only rival of the anaconda for first place. A surprisingly large number of specimens of this species are captured alive each year, and sold to dealers in wild animals. As a result, the largest serpent with which the animal-loving public becomes familiar in the zoological gardens and parks is this handsome Python. Specimens exceeding 20 feet in length, and running up to 25 feet, are really common in the possession of the animal dealers of Singapore, but about three-fourths of them die from lack of proper care before they are finally disposed of in Europe or America, and placed on exhibition.

The largest specimen which thus far has died in the Zoological Park measured 22 feet 10 inches, and weighed 170 pounds; but a larger unmeasured specimen is now living there.

<sup>1</sup> *Eunectes murinus.*

<sup>2</sup> *Python reticulatus.*

This splendid Python is at home in the hot and moist jungle which from Burmah to Java covers the land with a dense mantle of trees, thorny palms, rattans and tangled underbrush.

The temperature is practically stationary all the year round, and varies little save between 82° and 98° F. The frequent rains, and the moist, hot-house air of that region, with abundant animal food and ample cover, constitute ideal conditions for the rapid growth of reptiles, and the triennial shedding of their epidermis. It is no wonder that Pythons' and king cobras grow large there, or that they are so numerous that many of the former are caught alive by the Malays.

But the term "numerous" is capable of several interpretations, and in this case we enjoin a strict limitation. Although between forty and fifty Pythons of two large species<sup>1</sup> leave Singapore every year, let it not for one moment be supposed that anywhere in the East Indies are these serpents so numerous that they constitute a danger to human life, or that it is even possible to find them by hunting for them. Quite the contrary.

I spent several months in the Far East, roaming through jungles of all kinds, some of them so dense and so full of deadly bogs and miasma that now I recall them with a shudder. I never once found a wild Python, great or small; nor a cobra, even in cobra-ridden Hindustan; nor did any of my own native followers ever find a specimen of either for me. The only wild Python I ever saw or handled in its home jungle was one that was brought to me in the Malay Peninsula. It was hiding in a hollow tree, and when it looked out at a Malay who was passing, he whipped out his parang, cut off its head at one blow, and came to me calmly dragging behind him twelve feet of dead snake.

So far as I could learn, even the largest Pythons are harmless to man. They sometimes visit native villages, crawl through the frail fences which very feebly protect the domestic animals, and swallow—chickens and ducks! It is in these humble raids that some Pythons come to grief by being caught alive. But jungle people have no fear that a Python would make such

<sup>1</sup> The Black-Tailed Python (*Python mo-lu'rus*), although smaller than the Reticulated, attains a length of 20 feet.

a blunder as to attempt to make a conquest of a man. To be sure, in the Far East, people do not often go poking around in the jungles at night, in thick darkness. It is not considered the proper thing to do so.

The food of the Pythons of the East Indies must consist chiefly of the muntjac, hog-deer and other deer of small size; young wild pigs, pheasants and jungle-fowl. Our captive Pythons prefer large chickens—full-feathered—and rabbits. A Python should voluntarily eat a full meal every two weeks.

Until quite recently it was generally believed that if a large serpent would not feed voluntarily there was nothing to be done for it save to watch it commit suicide by starvation. Two years ago, Mr. Raymond L. Ditmars, Curator of Reptiles in the Zoological Park determined upon a very bold experiment. He decided that a starving twenty-foot Python should be fed artificially. Accordingly, a smooth bamboo pole was procured, and a string of four rabbits was tied up so that the pole would thrust the first one far into the serpent's interior, and drag the others after it. The next question was, how could the snake be controlled?

Summoning Keepers Snyder and Dahl, and five other men, the cage-door was opened. As the reptile raised its head to strike the intruders, a stream of cold water from a hose struck it full in the face. When it recoiled in confusion, the plucky keepers seized it by the neck, and quickly dragged it from its cage. As its form emerged, the waiting men seized it at proper intervals, and held it nearly straight.

The Curator presented the pole-strung rabbits, the first of which was angrily seized in the Python's jaws. With this auspicious beginning, it was the work of only a few moments to gradually push the string of wet rabbits down the serpent's throat, to a distance of seven feet, and withdraw the pole. Finally the tail and body of the snake was thrust into the cage, and with a careful toss from the hands of Mr. Snyder, the head landed on the coils, sufficiently distant that the door could be closed without accident.

Since that time, all large serpents that fast too long are fed in this manner, and the food thus mechanically placed in the stomach is properly assimilated.

## HARMLESS SNAKES OF THE UNITED STATES.

Of the grand army of harmless snakes inhabiting North America, the **King-Snake**<sup>1</sup> is unquestionably the king. It is also called the **Chain-Snake** and **Thunder-Snake**. It is the most courageous of all snakes, and in proportion to its size it is also the strongest. Toward man it is by no means especially vicious; but on the contrary, its manner is quite tolerant.

Toward all other serpents, however, it manifests as great aversion as any snake-hating woman, and it is pugnacious and aggressive to an astonishing degree. The King-Snake is, for its size, the most powerful of all the constrictors, and does not hesitate to attack a snake of another species several times larger than itself. It is cannibalistic in its tastes, and not only attacks and kills other snakes, but devours them.

In our Reptile House, a snake of this species once attacked a Cuban boa, fully three times its own size, and tried to swallow it! Had not the boa been rescued, it would undoubtedly have been quickly suffocated by the coils which its antagonist had wrapped tightly around its body. On another occasion a King-Snake that was placed for a very short time in the cage of the water moccasins, attacked one of the latter, wrapped around it, and killed it. Several times the moccasin bit its assailant, but the King-Snake is immune to the venom of serpents, and paid no attention to the counter-attack.

In some portions of the South, the King-Snake is believed to be a special enemy of rattlesnakes and moccasins, and on this account it is preserved from general slaughter. It is well attested that it does sometimes kill and devour snakes of both those species.


This bold serpent is found from Maryland to southern Florida, thence westward through the Gulf states to the Indian Territory, Texas and Matamoras, Mexico. Its average length is about 3½ feet, and it rarely exceeds 4 feet. From Maryland to Georgia it is a black snake with thirty white bands or rings around it, and is called the Chain-Snake. Farther south its body-color is greenish, with white rings, and is called the Thunder-Snake. Its favorite food is rats,

<sup>1</sup> *O-phi-bo'lus ge-tu'lus*.

mice, lizards, birds, and other snakes; but no frogs are eaten.

It reproduces by laying eggs. In Texas, New Mexico and Sonora, Mexico, the *Splendid King-Snake* is found. In Arizona, California and Nevada occurs *Boyle's King-Snake*, a conspicuous black serpent, marked by thirty broad, cream-colored bands. The latter sometimes predominate so effectively as to give the snake a general cream-colored appearance, with black rings. An entirely black variety, without rings, is found in Indiana and Illinois.

**The Corn-Snake**,<sup>2</sup> sometimes called the **Red Racer**, is one of the handsomest serpents in North America. Its general color-tone is mottled yellowish-red, or reddish-yellow. In detail its color-pattern consists of about forty squarish blocks along the back, each of which is dull



KING-SNAKE.

brick-red, with a deep margin of black, outside of which is a lighter ground-color. Its length is a little over three feet, and its form is slender and graceful.

Like the king-snake, this serpent is a powerful constrictor, a good climber, and seldom is seen on the ground. In the fields and forests, it is usually found in or upon low bushes. It frequents the habitations of man, and the roofs of old out-buildings are its favorite hunting-grounds for rats and mice. It is fond of rats, and because of this is considered a useful ally of the southern farmer, by whom it is often called the **Rat-Snake**. (Raymond L. Ditmars.)

<sup>2</sup> *Co-lu'ter gut-ta'tus*.


This serpent is courageous, but not particularly aggressive. Its food consists of rats, birds, eggs, small rodents, and warm-blooded creatures generally. In South Carolina, Mr. Ditmars captured a specimen which but a few minutes previously had finished swallowing a bob-white.

The home of this interesting and beautiful serpent is practically the same as that of the king-snake,—along the Atlantic coast from Maryland to Florida, and westward through the Gulf states to Arkansas. This snake is an egg-layer.

The **Gopher-Snake**<sup>1</sup> is our representative of the rat-snakes of South America and India, that make a business of catching rats in and around dwellings and out-buildings. In the South, it is often called the "**Black-Snake**,"—because it is black; but when it is particularly well polished, it takes on a gun-barrel blue appearance, when it is also called the **Indigo-Snake**.

This is a large and showy serpent, often attaining 8 feet in length, very docile and good-natured, and easily tamed.

At Oak Lodge, Florida, we once saw a very large wild Gopher-Snake emerge from the saw-palmetto jungle, and crawl directly toward the house. When Mrs. Latham was informed, she cried out reassuringly, "Oh, that is my pet snake! It keeps the place clear of rats." Forthwith she



PINE-SNAKE.

laid hold of it and picked it up, which the serpent did not resent in the least, even when it was passed from hand to hand for close examination. When finally released, it leisurely crawled under the house, quite as if nothing had happened.

<sup>1</sup> *Spi-lo'tes co'ra-is cou'per-ii*.

This is one of the best of all serpents to keep in captivity. In four years we have not lost a specimen by death, and Mr. Ditmars has one which he has kept in good health for eleven years. It is next in hardness to the water-moccasin. It is an omnivorous feeder, and, named in the order of choice, its food consists of rats, mice, birds, snakes, eggs, frogs, fish, lizards and even raw meat! (R. L. Ditmars.)

The Gopher-Snake is not a constrictor, it does not climb frequently, and does not care for water except to drink. It is strictly a warm-country species, and inhabits our Gulf states, from Florida to Matamoros, Mexico.

The typical **Pine-Snake**<sup>2</sup> inhabits the sandy pine-woods along the Atlantic coast from New Jersey to Florida; but other species of this genus are found throughout nearly every other portion of the United States except New England.

This species is quite harmless, even to other snakes, but for all that, it is a powerful constrictor. It lays eggs, and feeds upon birds, small rodents, and eggs. In devouring eggs it has a very odd but intelligent trick. It swallows an egg whole, and after it has passed a few inches down the throat, where it forms a large swelling, the serpent lifts its head, elevates its back, and exerts downward pressure directly upon the egg until the shell breaks!

A striking peculiarity of the Pine-Snake is found in the structure of its epiglottis, first observed and described by Dr. C. A. White, by means of which the hiss of this creature is so loud and so well sustained that it is like the hiss of red-hot iron in water. The maximum length of this snake is about 7½ feet. Its ground-color is whitish, the head is spotted with black, and along the back there is a series of about twenty-four very large brown patches, margined with black. Sometimes these blotches of color take shape as bands. The abdomen is dull yellow, with blackish-brown patches.

The **Black-Snake** of the East is a serpent of narrow form, but wide distribution. Westward it changes color, and is known at first as the **Blue Racer**, and then as the **Green Racer**. Although its Latin name is *Za-me'nis con-stric'tor*, it is not a constrictor, it is badly misnamed, it is perfectly harmless to man, and its bite is never more than a mere scratch. It is very cowardly, and

<sup>2</sup> *Pit-y-o'phis me-lan-o-leu'cus*.

will leap wildly from the edge of a rock or a steep bank in order to escape. If cornered, it makes a fierce but often absurd fight, sometimes be-



WESTERN COACH-WHIP SNAKE, OR RED RACER.

coming so frantic that it bites its own body. (R. L. Ditmars.)

This snake is a good climber, swims well, and is active and quick in movement, but it has no real power to speak of. It is not an enemy of the rattlesnake, as many persons suppose, but it devours snakes that are smaller and weaker than itself. Its favorite food consists of small rodents, young birds, eggs and frogs, but it does not eat fish. It is a great destroyer of mice and moles, and deserves well of the farmer on that account.

The young differ in color from adult specimens, being slaty gray, with chestnut-brown saddles on the back. In the third year, these colors fade, and the snake assumes its adult color. Speaking generally, the black form of this species occurs nearly everywhere throughout the United States east of the Mississippi into New England. What is called the *intermediate color* is too widely scattered to be defined, while the *green-and-yellow* form is found from Nebraska and Louisiana westward to the Pacific coast, and from Puget Sound to San Diego.

The length of this snake, when adult, varies from 40 to 58 inches.

**The Coach-Whip Snake**<sup>1</sup> is closely related

<sup>1</sup> *Za-me'nis fla-gel'lum*.

to the preceding species (both being members of the same genus), and has similar habits. It is even more slender than the black-snake. Its standard color is, toward the head, black or light yellowish-brown, fading out rapidly backward, until the tail becomes nearly white. But these colors vary exceedingly in widely separated localities.

This is a southern snake, and extends from Florida quite across the continent to California. In the far Southwest, its colors are so much suffused with pinkish it becomes the **Red Racer** (*Zamenis flagellum fre-na'tum*).

**The Garter-Snake**,<sup>2</sup> our oldest and most familiar friend among the snakes, is as harmless as a house-fly, and any one who exerts himself to crush one simply makes a pitiful exhibition of ignorance and folly. This is the most prolific and generally abundant snake in North America, and no amount of persecution seems to diminish its numbers to any noticeable degree. During the month of March, 1903, about 450 specimens were collected in and around the Zoological Park.

This serpent is viviparous, and sometimes forty-five are born in one brood. Out of a brood of thirty-eight born in our Reptile House, there was one double-headed specimen and three albinos. The standard length of this snake is from 24 to 30 inches, and one 36 inches long is a large specimen. Of the genus to which the Garter-Snake belongs, twenty-four species have been described, covering the whole of the United



COMMON GARTER-SNAKE.

States, and much contiguous territory. From the species named above, twelve tiresome sub-

<sup>2</sup> *Eu-tae'ni-a sir-tal'is*.

species have been evolved, which are of no interest whatever to the general student.

**The Red-Bellied Water-Snake**<sup>1</sup> is a highly colored variety of the common Water-Snake that merits special attention. It is the most showy and handsome representative of an interesting group of water-snakes, comprising about ten species, all of which are harmless, but very much in evidence in small streams and other bodies of water. They bring forth their young alive. They love to lie upon low bushes that overhang water, and bask in the sun. They are very suspicious, however, and when disturbed drop head first into the water, like a stream of oil running

Ditmars took three sunfish, one catfish, about a dozen tiny suckers and a crawfish. This interesting fish collection had filled the serpent so full it could hold no more. The species referred to is prominently marked by its shiny red belly, and rusty-brown upper surface. It is from 3½ to 4 feet long, and like all Water-Snakes, emits a disagreeable odor when handled. It inhabits the southern states generally, and extends northward into Illinois and Michigan.

**The Common Water-Snake**<sup>2</sup> inhabits all of the Gulf states and the Mississippi valley up to Iowa. In the New England states as far up as Connecticut, and also in the southeastern states



RED-BELLIED WATER-SNAKE.

New York Zoological Park.

down. The way to catch them is with a wire noose on the end of a light pole about ten feet long.

The species named above is widely known amongst the negroes of the Carolinas and other portions of the South as the **Copper-Belled "Moccasin,"** and it is feared accordingly. To the negroes of South Carolina, all water-snakes are "Moccasins." The Red-Bellied is held to be very deadly, and its bite is said to be "fatal" unless counteracted with large doses of good whiskey! (R. L. Ditmars.)

Water-snakes feed chiefly upon small fishes and frogs. From the stomach of one Red-Bellied Water-Snake collected in South Carolina, Mr.

<sup>1</sup> *Na'trix fas-ci-a'ta er-yth'ro-gas-ter.*

and the Mississippi valley is found a subspecies called *Natrix fasciata sipedon*.

**The Hog-Nosed Snake**<sup>3</sup> is a serpent of many names and remarkable habits. It is often called the **Blowing "Viper," Spreading "Adder,"** and other combinations of "Viper" and "Adder," all erroneous. This is the snake that is such a bold bluffer, and often saves its life by pretending to be very fierce and dangerous. Instead of fleeing from an intruder, this creature comes straight forward, with savage determination, hissing and darting out its tongue, and pretending to be a serious proposition. It looks as ugly and deadly as any real viper. It inflates the skin of

<sup>2</sup> *Na'trix fas-ci-a'ta.*

<sup>3</sup> *He-ter'o-don plat-y-rhi'nus.*

its neck with air, and hisses until it can be heard twenty-five feet.

In spite of all this bluffing, however, the Hog-Nosed Snake is really a harmless creature. It strikes viciously, but always with its mouth closed! Mr. Ditmars says it is almost impossible to induce one of these snakes to bite. When greatly annoyed, or tickled on the back, it will



HOG-NOSED SNAKE.

turn over on its back, open its mouth, allow its tongue to hang out, and permit the experimenter to hang it over a stick, as if dead. If thrown upon the ground on its back, it will slowly turn back again, take in its tongue, and crawl away.

When a small boy I once had a thrilling encounter on a bare prairie with one of these snakes, which sought to take refuge in its hole while I fought it off with my hat. At last the snake fled, and I blocked up the mouth of the hole. While I was ploughing the next round, the snake returned, and with its nose dug a new opening running diagonally down into the old one, and entered.

This snake is flat-headed and thick-bodied, and varies in length from 30 to 37 inches. Its colors are a mixture of brown, yellow and black, with no definite pattern, and are almost impossible to describe successfully. This species lays eggs, which are about one and one-half inches in length, covered with a thick, tough, flexible shell. When hatched the young are from 7 to 8 inches long, and they hiss very soon after they emerge. The embryo serpent possesses an "egg-tooth," for cutting the shell of the egg, but it loosens and drops out within a day or two after the serpent is hatched.

## THE POISONOUS SNAKES OF NORTH AMERICA.

Fortunately for us, all save one of our species of poisonous serpents are so peculiarly marked it is possible for any intelligent person to know them all, and recognize their dangerous character in a moment. This knowledge once acquired, all the other snakes of North America cease to be objects of dread or terror, and become merely so many interesting specimens of natural history.

A bird's-eye view of our venomous serpents reveals the following assemblage:

### Venomous Serpents of North America.

Rattlesnakes, 11 species,  
Massasaugas, 3 species,  
Water-Moccasin,  
Copperhead,  
Harlequin Snake,  
Sonoran Coral Snake.

Out of the 75,000,000 people in the United States, probably not more than two die each year as the result of snake-bites. The number of timid people who are frightened by harmless snakes, each year, must be about 1,000,000. Now, if all the latter could be so fully informed as to be free for all time from *groundless* fear, what a relief to suffering nerves it would be.

And why should any one remain in ignorance? In reality, there are only five types to learn, all the rattlesnakes and massasaugas being referable to one group by reason of the rattles and "buttons" on their tails.

Come, then! Let us address ourselves to the very simple task of learning from a book how to recognize the venomous serpents of North America, as readily as one recognizes the dogs and horses of our next-door neighbor. Excepting the water-moccasin, they are all so plainly marked that all persons except those who are blind may know them; and there is no excuse for forgetting them. Instead of going into their anatomy at length, our efforts for this occasion will be concentrated upon their external characters, habits and homes.

Fortunately, we have not in North America any house-haunting serpents of great cunning and unflinching deadliness like the **Hooded Cobra**, or **Cobra-de-Capello**,<sup>1</sup> of India. The bite of this species is very deadly, and whether wholly guilty or not, in India it is debited annually

<sup>1</sup> *Na'ja tri-pu'di-ans.*

with the deaths of between 18,000 and 22,000 persons. It is said, however, that many persons are murdered on the sly, and their deaths are charged up to the account of the Cobra-de-Capello.

The reasons why so many persons are bitten by Cobras are, (1) that in the rainy season, the serpents take refuge in and about the huts; (2) that practically all the natives go bare-footed and bare-legged; (3) that many of them are compelled to go about at night, without lights of any kind, and (4) the warning of the Cobra—spreading the hood, and hissing—is more frequently given after the bite than before it! Moreover, the Cobra is naturally much more irritable and vicious than the rattlesnake, or any other American serpent.

Of all the serpents that have entered the Reptile House, the Hooded Cobras are the most vicious, and eager to do mischief. At the slightest excuse, they spring to an erect posture, spread their hoods, and try their utmost to bite. One of them struck the glass of its cage front so frequently that it brought on a disease of the jawbone, which finally rendered it necessary to remove one entire side of the lower jaw. To keep the three Cobras from seriously injuring their heads by striking against the glass, it is necessary to keep the lower portion of the plate painted white.

The Hooded Cobra is a slender-bodied, nervous and active serpent, with a maximum length of about 48 inches. When the rainy season is on in India, it seeks refuge in and about human dwellings, especially under floors, and is also partial to thatched roofs. For its bite there is no sure antidote.

**The King-Cobra, or Snake-Eating Cobra,**<sup>1</sup> of the Malay Peninsula is the largest of all venomous serpents, easily attaining a length of ten feet. It is a very athletic serpent, slender-bodied and strong-muscled, able to erect its head three feet, perpendicularly, and strike nearly a yard. It is a very expert and vigorous climber, swims nearly as well as a water-snake, and is a thorough believer in the survival of the fittest. It feeds only upon other serpents and lizards, but it would be better if harmless serpents fed upon it.

No matter where you find him, the **Rattlesnake** is a fair fighter, and entitled to far more

<sup>1</sup> *Na'ja bun-gar'us*.

respect than he is likely to receive in this snake-terrified world. He strikes only in self-defence, when he thinks he is about to be trodden upon. Instead of lying in ambush, and striking in deadly silence, like the cobra and the moccasin, he rattles loudly when man or beast approaches, and gives fair warning to "keep off!" He rattles to save himself from injury, and his persistent *whirr* has saved thousands of persons, and tens of thousands of domestic animals, from being bitten. A western cow-pony, a government mule, or a range steer will spring sidewise from a warning whirr in the sage-brush quite as quickly as man himself, and almost as far.

If Rattlesnakes generally (of which there are fifteen species) were disposed to be mean, and treat man as many human beings treat all serpents, the annual death-list from Rattlesnake bites would be a long one. Despite the few exceptional cases, however, it is a ruling fact that Rattlesnakes do not go pestering around camps, or frequently crawl under the blankets of men sleeping upon the ground. Every year thousands of cow-boys sleep on the ground, literally among these reptiles, without a single Rattlesnake accident.

Thanks to a long-standing acquaintance with this serpent, I have myself on numberless occasions "bedded down in the open" in Montana, Wyoming, Florida, and elsewhere, with not a moment's fear of snakes. Depend upon it, a Rattlesnake does not go about looking for trouble. His best efforts are devoted to the promotion of peace and longevity.

Beyond question, the Rattler is a serpent of timid and retiring disposition. It has not one-half the courage of the hog-nosed snake, nor a quarter of the cobra's vicious aggressiveness. If you encounter one at a fair distance, say ten feet, it will either crawl away, slowly and defensively, or coil and warn you to keep off. In its feeding habits, in captivity, it is one of the most timid and nervous of all reptiles, and seldom eats save when safe from observation and interruption. When darkness falls, and the Reptile House is entirely quiet, the Rattler bashfully swallows his freshly killed rat or guinea-pig.

My first experiment with a captive Rattlesnake, a huge Diamond Rattler from Florida, was to catch and place in its cage a live rat. The



rat ran over the snake several times, and greatly annoyed it. The snake endeavored to get away from its disreputable associate, but in vain.

At last the rat flew at the Rattler, and bit him severely on the lips! This was too much to be endured. In a great rage the snake drew back, seized the body of the rat in its wide jaws, and held on while it drove its fangs through the tough skin of the rodent, and far into its body. After one could have counted ten, the rat was released; and thirteen minutes later it was dead.

**Species of Rattlesnakes.**

Fourteen valid species of Rattlesnakes are found in North America, one in South America, and there are none elsewhere. Our most prominent species are as follows:

their young alive, the normal number being between nine and fourteen. As soon as an infant Rattler bursts the thin transparent sac in which it is born, it is ready to coil and strike. Even at birth it is fully equipped with poison and fangs. Wild or captive, the favorite food of a full-grown Rattler is small mammals; but what they feed upon in a wild state when very young, remains to be ascertained. From our six species of captives, we have learned that Rattlers climb bushes with almost as much ease as professional tree-climbers, but in a wild state it seems fairly certain that they rarely do so.

The tail of the Rattlesnake is ornamented at the end with a rattle consisting of a number of joints of horny material developed out of the skin, one section dovetailed into another. The exact age of a Rattler is not indicated by the number

ENGLISH NAME.	LOCALITY.	LATIN NAME.
<b>Dog-Faced Rattlesnake</b> .....	New Mexico .....	<i>Crotalus molossus.</i>
<b>Timber Rattlesnake</b> .....	Eastern half of United States .....	<i>Crotalus horridus.</i>
<b>Diamond Rattlesnake</b> .....	Florida and Gulf States .....	<i>Crotalus adamanteus.</i>
<b>Texas Rattlesnake</b> .....	The Southwest .....	<i>Crotalus atrox.</i>
<b>Prairie Rattlesnake</b> .....	The Plains Region .....	<i>Crotalus confluentus.</i>
<b>Pacific Rattlesnake</b> .....	The Pacific States .....	<i>Crotalus lucifer.</i>
<b>Tiger Rattlesnake</b> .....	Extreme Southwest .....	<i>Crotalus tigris.</i>
<b>Horned Rattlesnake</b> .....	Extreme Southwest .....	<i>Crotalus cerastes.</i>
<b>Green Rattlesnake</b> .....	Mexican Boundary .....	<i>Crotalus lepidus.</i>
<b>White Rattlesnake</b> .....	Southern and Lower California .....	<i>Crotalus mitchelli.</i>
<b>Massasauga</b> .....	Nebraska to New York .....	<i>Sistrurus catenatus.</i>
<b>Edwards' Massasauga</b> .....	The Southwest .....	<i>Sistrurus edwardsi.</i>
<b>Ground Rattlesnake</b> .....	Atlantic States South .....	<i>Sistrurus miliaris.</i>

Among the Rattlesnake species are several striking examples of color-development to suit their surroundings, or what is known in well-worn phrase as "protective coloration." The Banded or Timber Rattlesnake is a good imitation of the color of dead leaves and damp earth. The color-pattern of the Diamond Rattler is made up of rich though quiet tones of brown and yellow, dark and light, like the shadows of saw-palmetto leaves falling upon yellow sand. The Texas Rattler and the Horned Rattlesnake of the Southwest are so pale and bleached one instantly associates them with naked deserts shimmering in fierce sunshine.

In their habits, so far as known, the various species are very much alike. They bring forth

of joints in the rattle at the rate of one for each year. On the contrary, under favorable circumstances about three joints will be developed each year, until the snake reaches maturity. We have now, in the Reptile House, Rattlesnakes three years old which already have in their rattles from seven to nine joints.

The rattles are not shed when an old skin is cast off, nor are they ever shed; but they are frequently broken off, usually about three joints each year after more than nine or ten joints have been acquired. It is very seldom that more than ten joints are found on a living snake.

It is possible to lengthen a snake's rattles, after they have been cut off, by joining on other joints of the same size, up to the number desired.

The slow vibration of a large set of rattles gives a sort of clicking sound, but when the wearer is thoroughly alarmed and angry, the spiteful "whirr" sounds like meat frying. The motion then is so rapid the eye cannot follow it.

Rattlers are not fond of bathing, but when swimming is necessary they swim well. The species which live in the North, pass the cold months in burrows below the frost line, either in the earth, or among rocks. If the situation chosen proves to be a cold one, the serpent becomes so torpid that it seems lifeless.

I once found a Prairie Rattlesnake abroad in northern Montana on October 10, two weeks

is the skin of the largest individual known to me. The wearer measured, before it was skinned, 8 feet 5 inches, and its girth at the thickest part of its body was 1 foot 3 inches.

This brown-and-gold species is most at home in Florida, on clean sand, among the cabbage-palmettos, saw-palmettos, and long-leafed pines. Although it rarely takes to water, it is sometimes called the **Water-Rattler**. It ranges northward into the Carolinas, westward through the Gulf states to the Mississippi River, and probably beyond. In Texas begins the home of the big **Texas Rattlesnake**,<sup>2</sup> of the same size and appearance as the Diamond, color-pattern



DIAMOND RATTLESNAKE.

New York Zoological Park.

after the first fall of snow. When brought to a realizing sense of its weakness and unworthiness, it crawled into a hole like a shallow post hole, and lay on the bottom completely exposed. This species is very wise in sheltering in the burrows of the prairie-"dog," but where none of those are to be found, the wash-out holes in cut banks can always be relied upon to furnish warm shelter for Rattler, bob-cat or wolf.

The **Diamond Rattlesnake**<sup>1</sup> is a royal serpent, the largest of the rattlers, and the handsomest snake in North America. A specimen 6 feet long, in good condition, will be accepted anywhere as a large one, but the largest specimens far exceed that size. At Oak Lodge, Florida, in the possession of Mrs. C. F. Latham,

<sup>1</sup> *Cro'ta-lus ad-a-man'te-us*.

and all, but of a very light color, as becomes a serpent of the arid regions.


In captivity the Diamond Rattler is, like all members of its genus, a timid and erratic feeder. Unless all conditions are entirely to its liking—perfect quietness, choice food, and no one looking, it will not swallow a morsel. When its views on the subject of food and service have been fully met, it will partake of a young rabbit, a rat or a guinea-pig.

The **Timber, or Banded, Rattlesnake**<sup>3</sup> of the eastern United States shows a wide range in color, varying from a handsome sulphur yellow to brown, and finally to almost black. Young specimens are always lighter in color than old ones. One of the popular names of this creature

<sup>2</sup> *Cro'ta-lus a'trox*.

<sup>3</sup> *Cro'ta-lus hor'ri-dus*.

is derived from the broad bands of brown color which encircle the light-colored specimens. Often the hinder half of an adult or old specimen



PRAIRIE RATTLESNAKE.

has a black-velvet appearance. The length of a large specimen is  $4\frac{1}{2}$  feet.

This Rattlesnake has suffered more from civilization than any other species. Throughout many vast areas of rich and closely cultivated agricultural regions, it is now totally extinct. Although it is believed to exist within fifty miles of New York City, a living specimen would be about as difficult to find as a mastodon.

Originally the home of this species embraced the entire territory from the Atlantic coast to western Iowa, Kansas, and into Texas. In many portions of this region it still exists in small numbers, and is said to be "fairly common in the Allegheny Mountains," from Pennsylvania southward.

**The Horned Rattlesnake, or Side-Winder,<sup>1</sup>** of the far Southwest is a creature of the deserts, and the oddest member of this group. It has a small horn over each eye, and in crawling it moves sidewise, in very deep curves, totally different from the straightforward course of most rattlesnakes when on the war-path. This is the smallest of our rattlers. Its general color is yellowish-gray, marked by small round spots, and its home is in southern Arizona, California, Nevada, and probably Sonora, Mexico.

**The Massasauga<sup>2</sup>** is the type of a genus of rattlesnakes containing only three species, distinguished by various anatomical characters, but from neck to tail well marked, for the general student, by a succession of very dark brown saddle-bag patches of color laid upon lighter brown.

<sup>1</sup> *Cro'ta-lus ce-ras'tes.*

<sup>2</sup> *Sis-tru'rus cat-e-na'tus.*


The joints of the rattles never exceed ten in number. This species is found at long intervals from the swamps of western New York to Nebraska, but it is so rare that living specimens are difficult to obtain.

**The Copperhead<sup>3</sup>** is a rather short and small serpent, seldom exceeding three feet in length. Its colors look like two shades of copper—broad bands of old copper laid on a background of new copper. When the skin is new and fresh, or when a specimen has been reared in the shadows of captivity, this serpent is beautiful. Strangely enough, it is in some respects the direct opposite of its nearest relative, the water-moccasin.

The Copperhead is a serpent of the woods and rocks, and is not found in open grass lands. It is found from Indiana eastward (but not northward) to the Atlantic coast, and well up into New England. It ranges southwestward to



BANDED RATTLESNAKE. (YELLOW PHASE.)




BANDED RATTLESNAKE. (DARK PHASE.)

Texas, and in different portions of its home it is known as the *Pilot-Snake*, *Upland "Moccasin"*

<sup>3</sup> *An-cis'tro-don con-tor'trix.*

and *Deaf "Adder."* It is decidedly poisonous, and its venom is second in virulence only to that of the rattlesnake.




MASSASAUGA.

In captivity, the food of this species consists of small mammals, young birds and frogs. It brings forth its young alive, and the usual number is between seven and nine.

The **Water-Moccasin**, or **Cotton-Mouth**,<sup>1</sup> is the ugliest snake in North America. Its body is about as lithe and graceful as a Bologna sausage, and its skin resembles the surface of sun-cracked mud. It is so ugly that stuffing it with tow does not make it look any worse. It has a piggish appetite for fish, but if no fish or frogs are handy, it eats other snakes. It is quite as ready to bite a friend as an enemy, and when Mr. Percy Selous was bitten by his "pet" Moccasin, he died in fifty hours, despite medical treatment.

The Moccasin is a southern snake, and it is a pity the species is not confined to Tierra del Fuego. It lives along the grassy margins of bay-



COPPERHEAD.

ous and swamps, and is most frequently found lying at the shore line, with its head and a small

<sup>1</sup> *An-cis'tro-an pis-ci-vo'ras.*

portion of its body out of the water. It is also much in the habit of lying upon logs, on bushes overhanging water, or in the vicinity of dried-up pools. When disturbed, it starts up, opens its mouth very wide, holds it open, moves its tail in slow vibrations, and stares wickedly at the intruder. It is the whiteness of the interior of the mouth that has given rise to the name of "Cotton-Mouth Moccasin."

This serpent does not coil itself in a round, tight coil, like a rattlesnake. As a rule, it holds its ground tenaciously, and does not retreat unless deep water is near. The fangs are shorter in proportion than in the rattlesnake, and the action of the poison is not so quick and violent as that of the rattler. But the bite must be taken seriously, and treated with the utmost vigor, if a fatal result is to be avoided.

This serpent attains an extreme length of about 5 feet, and a diameter of 3 inches. Usually,



WATER-MOCCASIN.

however, specimens are about  $3\frac{1}{2}$  feet by 2 inches. When adult, it is a snake *absolutely devoid of bright colors*, its scales being the color of dried mud, and very rough. The head is flat, the body thick and puffed out, and the tail is very blunt.


The young of the Moccasins are born alive, each one being enclosed in a thin, transparent sac, which bursts immediately upon reaching the outer air. The young are usually from 7 to 8 in number, but the last family born in the Reptile House contained 14. The young are strongly marked by light and dark bands, on account of which they are easily mistaken for young copperheads. They also resemble young hog-nosed snakes.

About the only redeeming feature in this serpent is the fact that in captivity it is very hardy. In four years, not one has died in our

Reptile House. It is a serpent of the Gulf states, coming as far north as North Carolina and southern Illinois, and extending westward to Texas.

**The Harlequin Snake**<sup>1</sup> is a small, shiny, delicately formed serpent, of rather quiet habits and retiring disposition. It belongs to the same Family (*Elapidae*) as the deadly king-cobra of India! As far as it can be seen, it is instantly recognizable by the alternation of brilliant coral-red, yellow and jet-black rings which encircle its body from head to tail-tip. Unlike the broad-headed pit vipers,<sup>2</sup> the head of this serpent is no wider than its neck, and as a special feature, its head is quite insignificant in size, but is always crossed by a broad yellow band. It is well to remember from this species that *not all venomous serpents have lance-shaped heads*.

The range of this beautiful but rather stupid little serpent begins in South Carolina, and in-



FER-DE-LANCE.

cludes all the Gulf states southward and westward to the Pecos River in Texas. It ascends the Mississippi states to southern Indiana. It is a very persistent ground-dweller, and in captivity it spends three-fourths of its time buried in the sand of its cage, quite out of sight. It eats garter-snakes and black-snakes, voraciously. Although its bite is undoubtedly poisonous, I have never known of any one having been bitten. In fact, it is difficult to see how any one can be bitten by this serpent without having it done by special appointment.

**The Sonoran Coral Snake**,<sup>3</sup> of southern Arizona and northern Mexico, is in appearance

<sup>1</sup> *E'laps ful'vi-us*.

<sup>2</sup> So called because of the existence of a round and deep pit on the side of the head, about half way between the eye and the end of the nose. In the rattlesnakes this character is very noticeable.

<sup>3</sup> *E'laps eu-ryx-an'thus*.

much like the harlequin snake, and it is mentioned only because it is so little known, and to remark that it is a good subject for observation.

**The Fer-de-Lance, or Lance-Head Snake**,<sup>4</sup> is the serpent terror of the West Indies. It is a small snake, only about 6 feet in length when fully grown, and 2 inches in diameter. Its head is very wide, and it has very long fangs in proportion to its size. Its color-pattern strongly suggests the light phase of the timber rattlesnake—brown, with black markings. On two occasions that we know of, travellers returning from the West Indies have brought with them in pasteboard boxes, as indifferently as if they were frogs, living and healthy specimens of this venomous creature! One specimen was brought to us by a lady and her child, for identification; and the keepers of reptiles shudder even yet when they think what might easily have occurred.

Fortunately, this serpent is not particularly aggressive, or hostile toward those about it. When it seizes its prey, however, it buries its fangs, and holds on determinedly. A female specimen in our collection gave birth to twenty-four young, but they one and all refused to eat, and failed to survive.

## SNAKE-POISONS AND THEIR TREATMENT.

The Rattlesnake's defensive equipment of fangs and poison has been perfected by Nature with as much care as the horns of hoofed animals, or the defensive armor of an armadillo. The ordinary jaw teeth have nothing to do with the poisoning process, and wounds from them would prove fatal only under exceptional conditions.

The venom of a serpent is a rather thick fluid, secreted in two glands that are situated on the side of the upper jaw, under the skin, behind the eye. In the stomach of an animal it is supposed to be harmless, and we know that in many cases it is so. To produce death, it must be injected into the blood, by a method that is practically instantaneous, and very effective. First there must be a puncture, then the injection of the poison.

To pierce the skin and flesh, the rattlesnake has two special teeth, called *fangs*, which are very long, slender, slightly curved, and exceedingly

<sup>4</sup> *Bo'throps lan-ce-o-la'tus*.

sharp at the point. A slender tube traverses the axis of the fang, from the root almost to the point, for the passage of the venom. Around each fang is a flexible sheath of tough, white skin, evidently for its protection.

The fang of a diamond-backed rattlesnake—the largest species—is about an inch in length. The small bone in which it is set at the root (maxillary) is so hinged by tough ligaments attaching to the roof of the mouth that it has some freedom of motion. When the jaws are closed, the fangs lie against the roof of the mouth. When the serpent strikes an enemy with the intention of poisoning it, the mouth is opened widely, the pterygoid bone pushes hard against the maxillary, and the sheathed fangs are thrown forward until they look like great hooks of white skin.

A serpent cannot be rendered permanently harmless by the removal of its fangs, because the fangs are *constantly renewed*. Each operating fang is backed up by a series of smaller ones, of different sizes, growing and awaiting their turn to do duty, and drop away. An adult fang is shed every six or eight weeks. The old tooth does not drop out until the new one is close beside it, duly connected with the poison gland, and ready for duty. Then the old fang either drops out, or is left sticking in the next animal bitten.

Even if fangs were pulled out, the poison sac would remain, and a scratch from the jaw teeth, duly poisoned, would endanger the life of the patient.

In striking to do mischief, the function of the lower jaw is to get under the part to be bitten, and press it up firmly against the attack of the fangs. The mechanism by which the fangs are thrown forward consists of a series of levers, and the special student will be greatly interested in the published drawings which illustrate its details. It is admirably shown in "Amphibia and Reptilia," by Dr. H. Gadow.

**Effect of the Poison.**—It is obviously impossible in a work of this nature to enter into this subject at length. In lieu of this, we will offer a very brief digest of what we believe to be absolute facts. These have been gleaned with care from several sources, but I make special acknowledgment to Dr. Leonhard Stejneger's presentation of the subject in his admirable

monograph on "The Poisonous Snakes of North America."<sup>1</sup>

There are two ways for the introduction of snake-poison into the system of a warm-blooded animal: (1) *through the blood*, by direct connection with a vein or artery, and (2) *through the skin and muscles*, one or both.

Although some of the great investigators differ somewhat on this point, it now seems reasonably certain that the manner in which snake-poison acts is by *paralyzing* the circulation of the blood, the breathing organs, the nerves, and even the digestive organs. The effect on the blood is a decrease in the strength and rapidity of the flow. In the nerves (after the first period of excitement), drowsiness ensues, which in fatal cases often lasts until death. The breathing is gradually diminished in strength and volume. The brain is usually the last organ to succumb. Dr. Stejneger's conclusion is that "the death which follows the introduction of the venom into the circulation must be attributed to gastro-intestinal apoplexy, and the stupefying action exercised directly upon the nervous system."

Venom introduced directly into the blood acts with great rapidity. When introduced hypodermically, through the skin and muscles, its action is much slower, and if the case is treated with great vigor from the very start, the patient has a fair chance to recover. Except from cobra bites, very many do recover.

The most dangerous snake bites are those inflicted upon the neck or face. The least dangerous are those upon the feet, the legs below the knees, and the hands and forearms.

**Treatment.**—There is small need to apologize for recording here the fundamental principles that should be carried out in case of accident. In the first place, any one who expects to campaign in a country infested with poisonous snakes should expend \$5.00 in the purchase of a small pocket-case containing a hypodermic syringe, a bottle of chromic acid 1 to 100, and another of liquid strychnine. Only the boldest and most enterprising travellers ever get beyond the sphere of influence of whiskey and brandy.

During the last ten years, medical men have been conducting investigations and making experiments to produce a universal antidote for

<sup>1</sup> Government Publication. For sale by the Bureau of Public Documents, Washington, D. C.

snake-poisons. These efforts have produced the now celebrated anti-venomous serum, discovered by Dr. Calmette, of the Pasteur Institute of Lille, France. It is obtained by very gradually injecting cobra-venom into the flesh of a living domestic animal, and giving Nature time to counteract the poison by her own methods. Eventually the subject becomes immune to these injections, and produces within itself a product which when injected into other animals renders them immune.

This material, now popularly known as anti-venine, is prepared in large quantities, and sent all over the civilized world for use against animal poisons generally.

Aside from the use of the antitoxin referred to, the key-notes of the treatment of a snake-bitten patient are, bleeding the wound, isolation of the bitten part if it be possible, the application of an antidote, and stimulation. In case of an accident, the regular medical treatment appears to be about as follows:

1. Cut cross the wound, or stab it, and compel it to bleed freely.

2. Tie a ligature, of cloth, rope or string, around the bitten member, above the wound, to keep back, as long as possible, the poisoned blood from the veins of the body.

3. If anti-venomous serum is at hand, inject it according to the directions which accompany it.

4. Give any alcoholic stimulant that may be available, in *small doses*, at frequent intervals; but remember that a *quantity* of any strong stimulant will do more harm than good, and may actually hasten complete paralysis, and death. Ammonia is of very little use, if any; and its use depends so much upon conditions that it should be employed only by a physician.

5. If the serum is not available, inject directly into the wound, as quickly as possible after the accident, a solution of chromic acid, or permanganate of potash, 1 to 100, and see to it that the hypodermic needle penetrates to the bottom of each wound. In the absence of a syringe, bathe the wound with the solution.

6. Having done all possible at the wound itself, then give hypodermic injections, on leg or arm, of "15 to 20 minims of *liquid strychnine*, every 20 minutes, until slight tetanic spasms appear." (Stejneger.)

7. The ligature must be loosened from time

to time, to permit a limited circulation of fresh blood, or mortification will ensue.

8. If medical aid is within reach, it should be procured as speedily as possible, but in most cases, the life of the patient depends upon what is done for him during the first hour following the accident.

**Mr. Gruber's Treatment.**—A practical method by which to escape death from the bite of a rattlesnake can be learned of Mr. Peter Gruber, of Rochester New York, who has been bitten about twenty times. His method of treating himself was described, to the writer as follows:

"I no longer suck the venom from a wound. Unless a man's mouth is in very perfect condition, it is dangerous. My first act is to take my knife, and cut a slit an inch and a half long straight from my body into the wound, and continue it the same distance beyond; and I make these two cuts bleed freely. This is to make the poisoned blood flow out of my veins, instead of farther into them, to poison my whole system. After the wound has bled as much as I think it should, I inject the permanganate above and around the wound. The proper proportion is one five-grain tablet of permanganate of potash dissolved in two ounces of water, and I inject about thirty minims—the capacity of a hypodermic syringe—about three times around and above the wound. I always have it ready, and I bathe the wound with this solution, using absorbent cotton to cover the wound so that it is not exposed to the air.

"During this time I take two or three small doses of whiskey,—but not much. After the permanganate has had a chance to take effect, I bathe the wound freely with a solution of two ounces of laudanum and two ounces of Goulard's extract in two quarts of water, and keep it moist with this until all unnatural colors leave it. And I drink quantities of milk—all I can swallow. After a time my stomach ejects it, and at first it comes up the color of snake venom. But I continue to take milk, again and again, until I am sure my stomach has been washed free from the poison. If the action of my heart grows weak, I inject strychnine into my arms with a hypodermic syringe."

Mr. Gruber bears on his forearms and hands a number of scars, as ocular proof of the success of his method in the treatment of rattlesnake bites.





BOOK IV

AMPHIBIANS



## CHAPTER XL

### INTRODUCTION TO THE CLASS OF AMPHIBIANS

Among the many wonders of Nature, few are more interesting to the thoughtful mind than those forms which connect the great groups of vertebrate animals by bridging over what otherwise would seem like impassable chasms.

For example, between the classes of Mammals and Birds we have the Platypus, or Duck-Bill, an Australian mammal the size of a small muskrat, which has webbed feet, and a duck-like bill, and which reproduces by laying eggs. Between the classes of Birds and Reptiles, there is a fossil bird called the *Archaeopteryx*, with a long, vertebrated, lizard-like tail, covered with feathers. The *Hesperornis* was a water-bird with teeth, but no wings, which inhabited the shores of a great western lake which now is a vast stretch of arid bad-lands.

Between the Reptiles and the Fishes stretches a wonderful chain of living links by means of which those two Classes are united. So numerous are these forms, they make an independent Class, containing about 1,040 species. Originally this group was called *Batrachia*, but recently the fact has been recognized that that term is too limited in its application, and by the latest authorities the term *Amphibia* has been adopted instead.

In the transition from the water-habiting Fishes, with gills and fins, to the land-going Reptiles, with lungs and legs, Nature has made some strange combinations. In some instances, fins, legs, lungs and gills have become so mixed that several notable misfits have resulted. In some cases we see legs going with gills, and in others fins and lungs are associated. Many of the Amphibians will serve teachers as very striking object lessons in the evolution of animal forms.

The **Class Amphibia** contains the cold-blooded vertebrates known as frogs, toads, salamanders, newts, proteans, and sirens.

In the insect-world, we are familiar with the three stages of existence by which the larva

passes into the chrysalis stage, and later on emerges as a perfect insect. Here, among the vertebrates, we find creatures which also pass through two very distinct and sharply defined stages.

An **Amphibian**, if literally translated from the Greek, is a creature of "two lives." A typical amphibian begins life as a legless, fish-like creature, possessed of perfect gills, an eel-like tail, and living wholly in water. This is the *larval stage* of the animal. Later on, four legs make their appearance, the tail disappears by absorption into the body, the digestive organs change from simple to complex form, and lungs take the place of gills. In this *adult stage*, the creature (usually) is fitted for life on land if it so elects.

Owing to the bewildering variations of form and anatomy that are exhibited by various species, it is almost impossible to formulate a general statement regarding amphibians which will not be open to exceptions. If the reader will bear this in mind, we may venture to state the leading characters of the members of this Class.

**General Characters.**—All save a very few amphibians are hatched from soft, translucent, jelly-like eggs that are laid in shallow water, usually in stringy masses. Sometimes the larval stage of a species is passed in the egg, but usually this period forms an important part of the active life of the animal, and may be observed at length before the change to the adult stage takes place. Amphibians are (usually) covered with smooth skins, quite destitute of scales, and have minute teeth, or none at all. During the larval stage they feed chiefly upon vegetable food, but when adult the majority require animal food. Their skeletons are much more simple in structure than those of reptiles. The majority are aquatic. Some species permanently retain their gills, and live wholly in water; others, like the frogs and toads, lose their gills, acquire practical lungs and

legs, and live upon land at will. Of the 1,040 species of amphibians, only forty are without legs.

An *am-phis'i-ous* animal is not necessarily an amphibian. The hippopotamus, the seal, sea-lion, otter and crocodile are indeed very much at home in water, but they are far above the Class *Amphibia*. They are by no means creatures of two lives, and they do not pass through a larval stage before attaining perfect form.

Like the reptiles, the amphibians are confined to the torrid and temperate zones, but a surprising number of species permanently inhabit some very cold and inhospitable portions of the temperate zone. With but very few exceptions, the

amphibians are quite useless to man. The legs of certain large species of frogs are prized by epicures, but with this exception, civilized man regards amphibians generally as inedible. Scientifically, the Class is highly interesting, chiefly by reason of the striking changes which so many of its members undergo. As a subject for the classroom and laboratory, frogs and toads are of well-nigh universal availability. Unfortunately, however, too many courses in elementary zoology do not forge beyond the frog.

As usual in seeking an acquaintance with Nature, a very simple diagram places this Class of animals on a clear and comprehensible basis.

**A BIRD'S-EYE VIEW OF THE CLASS AMPHIBIA.**

ORDERS.	FAMILIES.	TYPICAL SPECIES.
<b>ORDER ECAUDATA:</b> The Tailless Amphibians; Frogs and Toads.	WATER-FROGS, <i>RA'NI-DAE.</i> . . .	{ <b>Common Frog,</b> <i>Rana clamata.</i> <b>Bull-Frog,</b> <i>Rana catesbiana.</i> <b>Wood-Frog,</b> <i>Rana sylvatica.</i>
	TREE-FROGS, <i>HY'LI-DAE.</i> . . .	<b>Tree-Frog,</b> <i>Hyla versicolor.</i>
	TOADS, <i>BU-FON'I-DAE.</i> . . .	{ <b>American Toad,</b> <i>Bufo lentiginosus.</i>
	BURROWING TOADS, { <i>PEL-O-BATI-DAE.</i> . . . . .	{ <b>Spade-Foot Toad,</b> <i>Scaphiopus holbrooki.</i>
<b>ORDER URODELA:</b> Tailed Amphibians.	SALAMANDERS, { <i>SAL-A-MAN'DRI-DAE.</i> . . . . .	<b>Axolotl,</b> <i>Amblystoma mavortium.</i> <b>Spotted Salamander,</b> <i>Amblystoma punctatum.</i> <b>Newt,</b> <i>Triton viridescens.</i>
	AMPHIUMAS, { <i>AM-PHI-U'MI-DAE.</i> . . . . .	<b>Hellbender,</b> <i>Cryptobranchus (Menopoma) allgahaniensis.</i> <b>Congo "Snake,"</b> <i>Amphiuma means.</i>
	MUD-PUPPIES, <i>PRO-TE'I-DAE.</i> . . .	{ <b>Menobranchus or Mud-Puppy,</b> } <i>Necturus maculatus.</i>
	SIRENS, <i>SI-REN'I-DAE.</i> . . . . .	<b>Mud-"Eel,"</b> <i>Siren lacertina.</i>
<b>ORDER APODA:</b> Legless and worm-like Amphibians.	CAECILIANS, <i>COE-CI-LI'I-DAE.</i> . . .	{ <i>Typtonectes compressicauda.</i>

## CHAPTER XLI

# THE ORDER OF FROGS AND TOADS

### ECAUDATA

The members of this Order are the most numerous, most widely dispersed and the best known of the amphibians. In all there are about 900 species; and it may be added that the habits of some of them are very strange and interesting.


In their modes of life, the frogs and toads exhibit great diversity of inclination. The tree-frogs live in trees, the toads seldom leave dry land, the burrowing toads burrow in the earth, and the water-frogs live in water at least half the time.

Some of these creatures begin active life in water, as ugly, little fish-like tadpoles, and their transformation into the perfect frogs may easily be watched from beginning to end. In some of the toads, however, the tadpole stage is passed *in the egg*, and at hatching-time a fully developed but very minute toad emerges, and begins to hop about. Others again develop from the tadpole stage, much the same as frogs.

The larva of a species fairly typical of this Order as a whole may be found in the tadpole of any aquatic frog. It possesses a big, purse-like head,—like that of a goose-fish,—and a long, eel-like tail, surrounded by a continuous fin. At first there is no sign of legs. The intestinal canal is very long and simple, as befits the vegetable diet of the creature. In the transformation process, the tail is absorbed into the body, and long before it has disappeared, two pairs of legs have grown out. The front legs are weak, but the hind legs are long and powerful, and being attached at the extreme end of the body they have great freedom of movement. They are adapted both for leaping and swimming.

Of the adult creature, the body is short and

broad, covered with a smooth skin, destitute of scales, and there is no tail whatever. The mouth



FROM TADPOLE TO FROG.

A series of specimens showing the development of the Common Frog. Prepared by RAYMOND L. DITMARS.

is wide and capacious. The tongue is not free, being attached at the sides to the lower jaw.

The eyes are placed high up, quite above the upper surface of the head, so that the creature can float with only its eyes and nostrils above water.

The frog skeleton possesses several marked peculiarities, some of which must be noted, even though briefly. There are *no ribs*. The vertebrae are very few in number, but very wide in comparison with those of other vertebrates. The pelvis is of great size, and so long that it forms nearly one-half of the axis of the body. Instead of being attached at its sides, midway from top to bottom, the thigh bones (femora) are attached



LEOPARD-FROG.

*Ra'na vi-res'cens.*

at the extreme lower end,—the portion called the is'chi-um. In comparison with other vertebrates, the hind limbs and feet are of enormous proportions; and when these members are flexed, and then suddenly straightened out, the frog flies forward through the air as if thrown by a powerful steel spring. Some frogs can leap eight feet.

Although there are no ribs, there is a well-developed breast-bone, or sternum, for the attachment of the fore-legs; and it is said that in the frog the sternum appears for the first time in the development of the vertebrates from the lower forms.

The members of some groups of the frogs and

toads have teeth in the upper jaw, on a bone called the *vomer*; others have teeth in both jaws, but the majority are toothless.

The hibernation habits of these creatures sometimes produce unexpected and remarkable results. Occasionally the public is startled by the publication of a story of a living frog or toad being dug out of solid rock, many feet below the surface of the earth. I have never had an opportunity to investigate any of these alleged occurrences, but a personal experience has at least furnished food for thought.

In a hot and dry jungle in the interior of Ceylon, I once made a search for elephant bones in the dry bed of what in wet weather was a shallow brook. The larger bones were found upon the surface, but so many of the smaller ones had become embedded in the sand that it was necessary to dig for them. The sand had become so hard and solid it was half-way toward sandstone, and our spades and mattocks loosened it with difficulty.

About eighteen inches below the surface, we came upon several small frogs, three species in all, closely and solidly entombed. Even the ignorant and stolid coolies were amazed and excited by the discovery. The sides of the animals were greatly distended by water, but from the first moment they were in full possession of their faculties.

As we released these creatures from their tombs and placed them upon the grass, each one disgorged a quantity of water, and hopped away. Evidently they had filled themselves with water and burrowed into the sand during the previous monsoon, then six months past, in order to live until the next rainy season; and had the annual water-supply of that little stream been permanently diverted, no one can say how many years these frogs would have continued to live in their solid tomb of sand. The natives said that excepting in their wells, there was no water anywhere for many miles around.

## THE FAMILY OF WATER-FROGS.

### *Ranidae.*

**The Common Frog**<sup>1</sup> is the most popular and well-known species in North America. It is the first to be heard in spring, it gathers in the

<sup>1</sup> *Ra'na cla-ma'ta.*

most numerous companies, and is one of the most cheerful and industrious croakers we know. Sometimes its cry becomes almost a warble; and when about fifty voices are raised in tuneful chorus from the surface of one small pond, each one trilling and piping at the rate of sixty to the minute, without missing a note, the effort is certain to attract attention, in case there is any to be bestowed.

This species is one of the handsomest of our water-frogs, and is colored to match its marshy home. Its upper ground-color is a brilliant-green, broken up by irregular black blotches that are bordered with dull white, with dark bars across the legs. The head-and-body length is about 2½ inches.

The **Bull-Frog**<sup>1</sup> is known by its deep-bass voice, and its great size when adult. Beside the preceding species, this creature is a giant, and it is small wonder that the eyes of epicures rest covetously upon its massive thighs. Its upper color varies from bright green to dark olive-brown, marked with small and rather inconspicuous dark spots. Its length varies from 5 to 8 inches, and it is so well known that further description is unnecessary.

As an indication of the extent to which frogs' legs are consumed as food in the United States, the latest statistics of the United States Fish Commission are interesting. In 1899, the total quantity of frog meat recorded in the markets was 472,415 pounds, valued at \$74,690. The following were the chief sources of the supply:

	Pounds.	Worth.
Missouri. . . . .	237,608	\$29,313
Arkansas . . . . .	79,760	10,162
California. . . . .	20,687	20,638

In 1895, New York handled 69,774 pounds, valued at \$6,572.

The **Wood-Frog**<sup>2</sup> is not often found without specially seeking it. In the spring, when you are searching for early flowers, and are startled by seeing a small dead leaf suddenly take life and leap about four feet, you may know that it is one of these small creatures. It is only 1½ inches long, being next in smallness to the tree-frog. Although for a frog so small it can leap a very long distance, its strength is soon exhausted, and its final capture is easily made.

<sup>1</sup> *Ra'na cates-bi-an'a.*

<sup>2</sup> *Ra'na syl-val'i-ca.*

**THE TREE-FROG FAMILY.**

*Hylidæ.*

If tree-frogs were of great rarity, and inhabited only one remote island of a far-distant archipelago, their arboreal habits would be accounted as much of a wonder as the flying-frog of Borneo. But being fairly abundant in the eastern United States, the tree-frogs are regarded with but a mild degree of interest.

These creatures, which vary in length from one inch to five inches, have been provided with an opposable thumb, and a very effective sucking disc on the end of each toe, by which they are



NORTHERN TREE-FROG.

Natural size. Photographed at the instant of croaking, and copyright, 1903, by W. LYMAN UNDERWOOD.

able to climb trees, and live very comfortably upon their branches. Of all vertebrates that live in trees, these tiny frogs are the most difficult to see. Even when one is chirping boldly and cheerfully within six feet of your eyes, it is necessary to look keenly in order to locate it. There are few kinds of rough bark with which the colors of a tree-frog do not combine with startling accuracy. The opposable thumb, which appears in frogs and tree-toads for the first time in Nature's ascending scale, is of great use, and in all probability it is the principal factor in the arboreal life of these animals.

In South America there are several species of tree-frogs whose females carry their eggs, during incubation, in pouches or cells upon their backs. It is believed that the eggs are placed in

position and embedded there by the male frogs. Other species attach their eggs to leaves that are afterward rolled together at the edges. Others deposit their eggs at the bases of large leaves where water collects, and some are credited with placing them where they will fall into pools, to be hatched there. A Brazilian species called the "Smith"<sup>1</sup> constructs, at the edge of a pool, a really wonderful circular wall or fortress, of mud, in which its eggs are deposited.

**The Northern Tree-Frog<sup>2</sup>** is our best and most common representative of this large Family. It is two inches in length, and in cloudy weather, especially when storms are gathering, its cheerful, bird-like call is universally regarded as a harbinger of rain. It is not a high climber, seldom ascending more than twenty feet from the ground. Its colors match tree-bark so closely it requires very sharp eyes to find it, and when seen it usually is believed to be a knot.

In croaking, its vocal sac swells to enormous proportions. Mr. W. Lyman Underwood has been successful in photographing this animal at the instant of utterance, and his very interesting picture is reproduced herewith.

## THE TOAD FAMILY.

### *Bufo*nidae.

North American toads are distinguished from frogs by their rough, wart-covered backs, their dull colors, large and puffy bodies, smaller hind feet, shorter hind legs, lack of agility, and their land-going habits. The hopping amphibians which every summer shower brings out on side-walks and country paths, usually are toads. Altogether, there are about eighty-five species, mostly tropical. The majority live upon land, a few burrow into the earth, and a few live in the water. There are many species so frog-like that it is difficult to note the characters (chiefly of internal anatomy) which distinguish them.

**The Common Toad<sup>3</sup>** may stand as the representative of the Toad Family of North America. The long-legged, lightly built frog leaps gracefully and far; but the plethoric Toad is content to wriggle or hop briefly through life. Its existence depends largely upon the fact that

as yet man finds no value in it, and does not regard it as worth killing. When Toads become salable at five cents each, their extermination will follow soon.

The Toad deposits its eggs in water, in long strings, and after the transformation they grow so slowly that even in August the toadlets are so minute that about three could sit upon a copper cent. They seem more like insects than amphibians with bony skeletons. In winter, these creatures hide away in the deepest crevices they can find, or the cavities of hollow trees, or holes in the earth, and lie dormant until spring recalls them to life.

## THE BURROWING TOADS.

### *Pelobatidae*.

**The Spade-Foot Toad<sup>4</sup>** of the eastern and southern United States represents this large Family, of which two species only are found in the United States, and eighteen elsewhere. In the North, it is rarely seen, and little known. Personally I know nothing of it in life. Holbrook states that it is the commonest toad in the South; that it digs for its burrow a small hole about six inches deep, in which it lies in wait for every insect that may be tempted to enter. It seldom leaves its hole except in the evening, or after long-continued rains. This animal is two inches long, and its color is brown above, with dark patches.

## THE TONGUELESS FROGS.

Excepting the members of two small Families, all frogs and toads have tongues. Of the Tongueless Frogs, one species—which is universally called a "Toad"—is of special interest in illustrating a very curious feature of frog life.

**The Surinam Toad,<sup>5</sup>** of Dutch Guiana, is celebrated among naturalists all over the world because of the remarkable manner in which its eggs are cared for and hatched. Just previous to the egg-laying period, the skin of the back of the female is specially prepared by Nature for a remarkable proceeding. It becomes very thick, spongy and soft. The eggs are taken by the male Toad, and one by one are *embedded in the skin on the back of the female*, so effectually that the skin closes over them, and each egg becomes partially

<sup>1</sup> *Hy'la fo'ber*.

<sup>2</sup> *Hy'la ver'si-co-lor*.

<sup>3</sup> *Bu'fo len-tig-i-no'sus*.

<sup>4</sup> *Sca-phi-o'pus hol'brook-i*.

<sup>5</sup> *Pi'pa americana*.



encysted, and retained in a cell of its own<sup>1</sup>. There they remain until they are fully incubated, the tadpole stage is passed, and a tiny, but perfect Toad emerges *from the skin of its mother's back!*

The number of young usually produced at one hatching is from sixty to seventy, and the period of incubation is from seventy-five to eighty-five days. At the close of this process, the thickened layer of skin on the back of the female loses its vitality, and is shed very much as a snake sheds

a dead epidermis. Although the front feet of the Surinam Toad are small and webless, the hind feet are of great size, fully webbed, and so much drawn in at the ends of the toes that in swimming the foot is saucer-shaped.

There are other frogs which display remarkable intelligence in the production of their young, their methods going far beyond what one would expect in creatures as low in the vertebrate scale as the amphibians. As a whole, the members of this Order offer a wide field for the specialist.

<sup>1</sup>For a full description of the process, see the Proceedings of the Zoological Society of London, 1896, p. 595.

## CHAPTER XLII

# THE ORDER OF TAILED AMPHIBIANS

### URODELA

The members of this Order are readily distinguished from the preceding group by the possession of tails, which they retain throughout their lives; by their gills, which most of them retain permanently; by the absence of scales, and by the fact that with very few exceptions they are strictly aquatic. It is safe to say that any four-legged aquatic creature having a tail but no scales, is either a salamander, newt, mud-puppy or siren.

**The Order U-ro-de'-la** is the dividing line between the finny, gill-breathing fishes, and the four-legged, lung-breathing, land-going lizards. Strange to say, its members are most abundant in the temperate regions of the earth, and except in two or three small areas, are absent from the tropics. In Australasia there are none, and in South America and Africa there are none save in their extreme northern portions.

Of all countries, the region embracing the United States and the southern provinces of Canada is by far the richest in species belonging to this Order, the total number present being fifty-two. Mexico and Central America contribute fourteen more, all salamanders. In this total of sixty-six species, eighteen genera are represented, fourteen of which are found only in the New World. The total number of species in the Old World is only thirty-six. In North America, the northern boundary of the Order Urodela is a line extending due east and west across the continent about on the 52d parallel of latitude. ("Amphibia and Reptiles," Dr. Hans Gadow, pp. 95-6.)

### THE FAMILY OF SALAMANDERS.

#### *Salamandridae.*

The members of this Family seem to be engaged in a continuous struggle at the dividing line between lungs and gills, and exhibit all pos-


sible variations between perfection in both those organs. One species (the axolotl) changes from water to land with neatness and despatch. Another (the striped-backed salamander) lingers for two or three years in its larval state, in the water, while the blue-spotted salamander lives upon land, in moist forests. But one or two illustrations must suffice for all.

**The Axolotl**,<sup>1</sup> of Mexico, is in some respects the most striking—even theatrical—example of salamandrine life and character. Its transformation is so rapid and complete that it is highly impressive. As an Axolotl, it is either a dark gray or a perfectly white and almost translucent animal, about 7 inches long, with external gills divided into three long, ragged branches; a long tail with a continuous fin above and below, and four very practical legs. This is the larva.

If the pond in which this creature lives threatens to dry up, the gills and the fins on the tail and back begin to shrink, and disappear, and the animal begins to breathe air at the surface of the water. Finally, when the transformation is complete, a lizard-like animal with very serviceable lungs, no gills whatever, and not a vestige of fins on tail or back, emerges upon the land, and thereafter leads a terrestrial life. It is then known as a **Spotted Salamander**; and it is no wonder that for many years these two forms were considered creatures of different species. It was in the Jardin des Plantes, in Paris, that the process of birth, growth and transformation was finally discovered.

It is not difficult to bring about the transformation of the Axolotl, by gradually diminishing the water-supply, and thus observing from day to day the progress of the change. More than this, the transformation can be arrested by gradually diminishing the allowance of air, thus forcing the

<sup>1</sup> *Am-bly'sto-ma ma-vor'ti-um.*



THE TWO LIVES OF THE AXOLOTL.

The lower figure shows the wholly aquatic larval form, with gills and tail fins, called the Axolotl. The upper figure shows the same creature fitted for life on land, and known as the Spotted Salamander.

imperfect Spotted Salamander back into aquatic life. At first there is a struggle against life under water, but finally the animal becomes adjusted to it. (R. L. Ditmars.)

By keeping the larval Axolotl in an aquarium, with an abundance of water but with no encouragement nor facilities for breathing air, it not only remains in that stage indefinitely, but it breeds successfully.

This species is most abundant in the shallow lakes around the City of Mexico, but it inhabits nearly the whole of Mexico and also a considerable area in the southwestern United States. Unquestionably, the wonderful mobility—as it may truly be called—of this creature is for the purpose of enabling it to survive in a region wherein droughts are common, and where the life of an aquatic animal depends upon its ability to change from water to land. Of all members of the Order Urodela, this is to me the most wonderful.

**Salamanders, Generally.**—In shallow brooks, in still pools of all kinds, from the shaded woods of the East to the wind-swept, sun-bathed prairies and bad-lands of the West, and both on and in the damp earth of forests high and low, we occasionally find little smooth-skinned, lizard-like animals. They are slow in movement, weak and incapable of either defence or flight, and are at the mercy of almost any species larger than themselves. These are Salamanders, and in view of the fact that some are wholly aquatic and others wholly terrestrial, it is difficult to choose from our sixteen species one which may stand for the majority. The diversity of habit of these animals is greater than their differences in form. The various members of the group inhabit all sorts of quiet situations, from the rocks and dry ground of the *Blotched Salamander*<sup>1</sup> to the mountains of the *Blue-Spotted Salamander*,<sup>2</sup>

<sup>1</sup> *Am-bly'sto-ma o-pa'cum.*

<sup>2</sup> *Pleth'o-don glu-ti-no'sus.*

and the swift-running streams of the *Dusky Salamander*.<sup>1</sup>

Very frequently, salamanders are found underneath fallen trees, or stones, or under the bark of decaying logs; and on the western prairie farms the plough-share turns into the broad light of day many a burrowing amphibian.

On the whole, the **Spotted Salamander**<sup>2</sup> appears to be the best species with which to introduce the North American group. It is distinctly marked, and of wide distribution. Its length is 6½ inches, its body is broad and full, and its tail is



MENOPIOMA, OR HELLBENDER.

shorter than its body. Above, its ground color is dark brown or black, on which is laid about thirty irregular yellow spots. The Spotted Salamander of Europe is a different species, its light markings being in the form of elongated patches or bands. Except for its external gills, the larva of this species looks much like an ordinary tadpole; but with transformation the gills disappear. Occasionally this species is found in spring-houses and cellars.

### THE NEWTS, OR TRITONS.

#### *Pleurodelidae.*

Although quite abundant in the Old World, (sixteen species), the newts are represented in America by only two species. All these tiny creatures inhabit water during the breeding sea-

<sup>1</sup> *Des-mog-na'thus fus'ca.*

<sup>2</sup> *Am-blys'to-ma punc-ta'tum.*

son, but at its close, some species leave it, and live for a period upon land, where their habits are much like those of terrestrial salamanders.

Most species of newts look very much like small, weak, scaleless lizards, except that in some species the males, and in others both sexes, have broad fins on the tail, above and below. In some cases the upper fin is prolonged forward along the back, quite up to the head.

Of our two species of Newts, the **Crimson-Spotted Newt**<sup>3</sup> endeavors to make up by its abundance for the scarcity of species of the Genus *Triton* in America. It is quoted by herpetologists as "very common in ponds everywhere" in the State of New York, and its known range embraces the northern and eastern portions of the United States. It is about 3½ inches long. Its color above is brown, or greenish-brown, with two rows of bright vermilion spots, in all from 6 to 12. Its under surface is orange, marked with small black dots. Half-grown specimens are brownish-red, with the characteristic spots of bright red.

This puny little animal inhabits deeper water than most salamanders, and swims freely, often in an upright position, in

which the hind legs hang motionless while the tail does all the work. It feeds upon the larva of aquatic insects, worms, and very small mollusks. For schoolroom aquaria, Newts are more easily obtained than any other of the tailed amphibians, and they are easily kept.

Our Newt has long been of much interest to American naturalists, and its complicated series of changes from the egg to adolescence have been carefully studied and reported upon.<sup>4</sup>

The Newt of western North America (*Triton torosus*) is one of the largest of the genus, and attains a length of 6 inches. The tail is longer than the body, much flattened vertically, and is provided with a dorsal and ventral fin. The under parts are colored yellow.

<sup>3</sup> *Di-e-myc'ty-lus vir-i-des'cens.*

<sup>4</sup> See L. J. Gage in the *American Naturalist*, 1891, p. 1084.

## THE FAMILY OF AMPHIUMAS.

*Amphiumidae.*

Unfortunately there is no English name which properly applies to the members of this Family. Some are like salamanders, and some are like eels; but none are "fish-like" salamanders, as they are sometimes called. In the perfect state they are without gills, the gill-clefts being in a vanishing stage, either reduced to a pair of small holes, or totally absent. Both jaws are provided with teeth, but the eyes are without lids.

This Family consists of two genera and three species, two of which are found in the United States, the other in Japan.

**The Hellbender, or Menopoma,**<sup>1</sup> is one of the ugliest looking creatures on this continent. When fully adult it is from 18 to 20 inches in length, its head and body are much flattened, while its tail is flattened vertically and completely finned. Its legs and feet are short and thick, and all along the middle of each side is a wide, convoluted fold of skin.

Its color is a uniform dull brown, accentuated by a few dark blotches of very irregular shape. On the left side there is a gill opening, but on the right there is none; and there are four pairs of gill-arches. The skin is smooth, but the head bears many wart-like tubercles.

This unpleasing animal is found in many of the streams that flow into the Ohio River, and the Mississippi also, but it is most abundant in Pennsylvania, especially in streams whose sources are in the Alleghanies. In its food habits it is very voracious, feeding upon worms, minnows and crayfish, and often taking the hooks of fishermen in quest of that most repulsive of all American fishes, the catfish. Fishermen hate

<sup>1</sup> *Cryp-to-bran'chus* (or *Men-o-po'ma*) *al-le-gha-ni-en'sis*.

the Hellbender; but between catfish and Hellbender there would seem to be small choice.

The Hellbender is very tenacious of life, and it is said that it can live on land for twenty-four hours without perishing. On this point, Mr. William Frear offers the following testimony:

"One specimen, about eighteen inches in length, which had lain on the ground exposed to a summer sun for forty-eight hours, was brought to the museum, and left lying a day longer before it was placed in alcohol. The day following, desiring to note a few points of struct-



Drawn by J. CARTER BEARD.

THE CONGO "SNAKE," OR EEL-LIKE SALAMANDER.

ure, I removed it from the alcohol in which it had been completely submerged for at least twenty hours, and had no sooner placed it on the table than it began to open its big mouth, vigorously sway its tail to and fro, and give other undoubted signs of vitality."

**The Giant Salamander,**<sup>2</sup> of Japan, is a brother species to the Hellbender, but is very much larger. It is the largest of all the amphibians, and sometimes attains a length of three feet. Specimens may always be seen in the Reptile House of the Zoological Park.

<sup>2</sup> *Cryp-to-bran'chus max'i-mus*.

**The Congo "Snake," or Eel-like Salamander,**<sup>1</sup> is in many ways related to the foregoing species, but in external appearance it seems widely different. In appearance it looks like a blunt-nosed, wide-mouthed eel, with a tiny pair of legs close behind its head, and another pair about four-fifths of the way back to the end of its tail. Even in a small aquarium tank, in a well-lighted reptile house, these tiny legs are so thread-like and so short they are seldom noticed save by those who know they are present, and look specially for them. The absurd little feet on these ridiculous legs have but three toes, and the wonder is that such useless or "aberrant" appendages have not long since disappeared altogether.

The color of the Eel-like Salamander is a uniform gray-brown, and its length when adult is usually about two feet. There is a gill opening on each side of the neck, and there are four internal gill-arches. There are no external gills.



THE MENOBRANCHUS, OR MUD-PUPPY.

These creatures inhabit the muddy streams and stagnant waters of our southeastern states, and are in the habit of burying themselves in the mud, sometimes to a surprising depth. They feed upon every form of aquatic animal life which is small enough to be seized and swallowed—insects, worms, crustaceans, shell-fish and even small fishes. In the South they are occasionally found in the ditches which irrigate the rice-fields.

<sup>1</sup> *Am-phi-u'ma means.*

## FREE-GILLED SALAMANDERS.

### *Protidae.*

This very small Family contains only three genera, with but one species in each. One of these, the **Olm**<sup>2</sup> of Europe, is recognizable at sight by its general eel-like appearance, its tiny legs far apart (like the Congo "Snake"), and the big bunch of external gill-branches waving on each side of the neck, close to the head. This animal is totally blind, and is found only in the caverns of the Alps.

**The Mud-Puppy, or Menobranchus**<sup>3</sup>, bears a strong external resemblance to the Hellbender, but is readily distinguished from the latter by the conspicuous mass of external gill-branches with which the animal fans the water at every breath. It inhabits many of the rivers of Ohio, Pennsylvania, Indiana, the Great Lakes and northern New York, and is often taken in fishermen's nets.

### **The Subterranean Protean of Texas.**

—Very recently a rather startling discovery was made in Texas, near San Marcos. From the bottom of an artesian well 188 feet deep, there came up with the water several blind Proteans, colorless and white, which up to that time had lived only in the subterranean streams and pools of the earth's crust, and were quite unknown. Along with them came four new species of Crustacea (crab-like creatures); and doubtless it was upon those that the Proteans lived. Unfortunately, thus far those who received the new amphibians did not succeed in inducing them to eat, and none have survived.

The species has been christened *Typh-lo-mol'ge rath'bun-i*.

## THE TWO-LEGGED SALAMANDERS.

### *Sirenidae.*

Near the foot of the Class *Amphibia*, we find the **Two-Legged Salamanders**, of which there are only two species, both American. Both look

<sup>2</sup> *Pro'te-us an-guin'e-us.*

<sup>3</sup> *Nec-tu'rus mac-u-la'tus.*

very much like the Congo "Snake"; but the hind legs are totally absent, and external gills are conspicuously present. The front legs, which are close behind the gills, are larger than in any other of the eel-like salamanders, and are of some slight use.

The **Siren Salamander**, or **Mud-"Eel,"**<sup>1</sup> of our southeastern states, has four toes on its feet, three pairs of gill openings, a smooth skin of a dull black color, and when fully adult a length of about 24 inches. The habitat and habits of this creature correspond closely to those of the Congo "Snake" of the southern rice-fields and swamps.

### THE ORDER OF WORM-LIKE AMPHIBIANS.


#### *Apoda.*

Last and lowest in the Class *Amphibia*, we find a group of creatures that externally seem more like worms than vertebrates. It is interesting to know that there are true vertebrates so very worm-like that they have neither legs, feet, nor fins. Some, however, have overlapping scales, like fishes.

Collectively, these animals are called **Caecilians** (pronounced se-sil'i-ans). There are forty

<sup>1</sup> *Siren la-cer-ti'na*.

species, inhabiting the lower half of Mexico, Central and South America, equatorial Africa, India, Burma and northern Australasia, but not the United States. They are of burrowing habits, and their skulls exhibit a degree of solidity and strength quite in keeping with the necessities of creatures which can burrow only with their



SIREN SALAMANDER, OR MUD-"EEL."


heads. Many of them are totally blind—by the concealment of their eyes under the skin, or the maxillary bones. (Gadow.) The exact relationships of the Apoda are yet to be determined conclusively.





**BOOK V**

**FISHES**



Drawn by J. CARTER BEARD.

REMARKABLE MEMBERS OF THE ORDER OF SOLID-JAW FISHES. (Page 410.)

1. TRIGGER-FISH (*Balistes capriscus*).
2. BOX-FISH (*Ostracion tricornis*).
3. PORCUPINE FISH (*Chilomycterus geometricus*), inflated.
4. PUFFER (*Tetodon turgidus*), with air-sac inflated.

## CHAPTER XLIII

# INTRODUCTION TO THE CLASS OF FISHES

The study of fishes is called *ich-thy-ol'o-gy*.

So great is the number of species that the mass is, at first thought, fairly bewildering. During the last twenty years the researches of the men who devote their lives to the study of fishes (called ich-thy-ol'o-gists) have brought to light hundreds of new forms.

The inhabitants of the waters of North America, alone, form a great multitude. Of the fishes found north of Panama, marine and inland, the "Descriptive Catalogue" of Drs. Jordan and Evermann, Part IV, completed in 1900, enumerates the following groups, species and subspecies as recognized by those authors :

Orders of Fishes. . . . .	30
Families. . . . .	225
Genera . . . . .	1113
Species. . . . .	3263

The four volumes comprising the work mentioned above make a pile nine-and-a-half inches high, and contain 3,313 fine-print pages of text, and 392 plates. The "Systematic Arrangement," or table of contents, is wholly in Latin, and fills 95 closely-printed pages. The work has been carefully devised to be of no use whatever to anyone save an ichthyologist.

When this array confronts the general student, the prospect is rather appalling. From the first page to the last, every technical work on fishes abounds in descriptive terms that to most persons are about as attractive as the spines on a porcupine fish. If the general reader attempts to master them, he soon finds himself involved and discouraged, and the desired general view of our finny tribes is obscured in fog.

But the whole subject of fish study is merely a matter of method. With fishes, as with the other vertebrates, the *Orders* are the master keys by which a proper exhibit can be unlocked and displayed. At the same time, the Subclass divisions are of great importance, and must constantly be kept in mind. Leaving out the deep-sea fishes, which we can well spare for the present, there are twenty well-defined Orders, the types of which are almost as easily known and

remembered as a score of pictures in an art gallery. *The Orders must not be lost sight of*, for when they are firmly grasped by the understanding and the memory, the fog begins to rise.

**General Characters.**—A typical fish is a cold-blooded animal, with a bony skeleton, an elongated body which is covered with overlapping scales, and an outfit of fins for balancing, steering and propulsion. It has gills instead of lungs, fixed eyes, and a swimming-bladder, and is specially fitted for a wholly aquatic life. It is provided with teeth, it hears sounds by the transmitting power of the bony plates of the skull, and usually it lays eggs for the production of its young. The body of a typical fish is wedge-shaped, narrowest at the tail, thin from side to side, and the head tapers to a blunt point. This form is specially designed for rapid and easy progress through water.

The Black Bass may fairly be regarded as a perfectly typical fish.

The variations from the perfect type are almost innumerable. For example:

The Lung-Fish has foot-like fins, and practical lungs.

The Catfish has no scales.

Some Sharks and a few other fishes bring forth their young alive.

The Rays and Skates are the flattest of all vertebrates.


The Climbing Perch can climb.

The Flying-Fish can rise from the sea, and fly.

The Lantern Fish, of the deep sea, carries a phosphorescent light upon its head.

The anatomy of fishes is a special branch of knowledge in which the general reader can scarcely be concerned, but for the young ichthyologist there are many special works. Books for the identification of all the known species of fishes in North America are now available for those who desire them. At present, however, we are concerned only with the twenty great groups, or Orders, and the fifty or sixty important types which represent them. Of these there must be some serious study.

Up to this date, nearly every systematic writer



THE NAMES OF THE FINS OF A TYPICAL FISH.

The species shown is the Black Grunt (*Haemulon plumieri*), and it represents the large and commercially important Family of Grunts (*Haemulidae*), represented in our warm waters by about 55 species.

upon the Fishes as a Class has chosen either to alter or ignore previous classifications, and adopt the arrangement which to him has seemed most logical and reasonable. In order to conform to this time-honored custom, I have elected to do likewise!

With the subdivisions of the Orders, we are not at present seriously concerned, our main object being to block out the larger groups, only. The arrangement of Orders set forth on pages 378-9 is called a "practical arrangement" because it can be understood, and is available for practical, every-day use.

### THE FISHERY INDUSTRIES, AND FISH PROPAGATION.

Says Mr. Charles H. Townsend, late Chief of the Division of Statistics, United States Bureau of Fisheries, "The commercial fisheries of the United States employ about 200,000 persons, the amount of capital invested is \$60,000,000, and the annual value of the products to the fishermen is approximately \$50,000,000."

As a source of supply of cheap and wholesome flesh food, the fishes of our waters are almost as vitally important as coal. The best fish rarely costs more than one-half the price of the best beef and mutton, and often only one-third as much.

In 1871, Professor Spencer F. Baird, Secretary of the Smithsonian Institution, induced Congress to create and perpetuate the United States Bureau of Fisheries, for the propagation of food fishes, and the preservation of the fisheries. The appropriation for 1902 was \$543,120.

To-day the United States Bureau of Fisheries, as the propagator and preserver of food fishes, is engaged in a hand-to-hand struggle with the 200,000 destroyers. There are comparatively few laws which are intended to limit the catch of commercial fishes; but the sportsmen have provided many statutes for the preservation of the high-class "game" fishes. Nearly every state maintains a state fish commission, for the special benefit of its own citizens, and some of these are doing very important work.

The United States Bureau of Fisheries has not entered into the business of procuring legislative enactments for the regulation of fisheries, but has left that work to the various states concerned. Its greatest efforts have been put forth in stocking new waters with desirable food fishes, and in restocking waters that have been depleted of their natural supply of fishes.

The importance of the fish-propagating measures of the national government can hardly be overestimated. The map of the United States is dotted over, from corner to corner, with the fish-hatching stations of the Bureau. In number they are thirty-nine, and they have been located with a view to the propagation and distribution of practically all the most desirable species which by their habits of life are available for such operations. It is of general interest to state the locations of the fish-hatching stations now (1903) actively at work, taking eggs, hatching them, and distributing both eggs and young fish. They are situated as follows:

Maine, 3:	{ Green Lake. Craig Brook. Grand Lake Stream.	North Carolina: Edenton. Virginia: Wytheville. Tennessee: Erwin. Ohio: Put-in-Bay.
Vermont: St. Johnsbury. New Hampshire: Nashua. Mass. 2: { Gloucester. Wood's Holl.		Michigan, 4: { Northville. Detroit. Alpena. Sault Ste. Marie.
New York: Cape Vincent.		Minnesota: Duluth.
Maryland, 2: { Battery Station. Bryan Point.		Illinois: Quincy. Iowa: Manchester. Missouri: Neosho. Texas: San Marcos. Colorado: Leadville.
District of Columbia: { Central Station. Fish Lakes.		

Of still greater interest to all catchers and consumers of fish is the answer to the question, "What are the fishes that are being propagated and planted by the United States Bureau of Fisheries?" A full answer will constitute an excellent showing of the Bureau's estimate of the comparative values of our best food fishes; but at the same time due allowance must be made for the things which are and are not possible in fish hatching.

**Distribution of Eggs and Live Fish by the United States Bureau of Fisheries, during the Year which ended July 1, 1902.**

Shad. . . . .	106,986,000
Quinnat Salmon. . . . .	48,683,718
Atlantic Salmon. . . . .	638,765

Landlocked Salmon. . . . .	822,220
Silver Salmon . . . . .	424,530
Blueback Salmon. . . . .	3,371,000
Steelhead Trout. . . . .	534,882
Lock Leven Trout. . . . .	96,760
Rainbow Trout. . . . .	1,675,121
Black-Spotted Trout . . . . .	1,868,500
Brook Trout. . . . .	6,579,762
Lake Trout. . . . .	27,260,490
Scotch Sea Trout. . . . .	24,531
Golden Trout. . . . .	69,950
Grayling. . . . .	1,803,258
Whitefish. . . . .	594,490,000
Pike-Perch. . . . .	237,099,575
Pickereel . . . . .	805
Catfish . . . . .	95,970
Yellow Perch. . . . .	1,700
Buffalo Fish. . . . .	200,000
Black Bass. . . . .	262,157
Crappie. . . . .	735,120
Strawberry Bass. . . . .	3,551
Rock Bass. . . . .	37,170
Warmouth Bass . . . . .	100
Sunfish and Bream. . . . .	623,739
Cod. . . . .	212,001,000
Flat-Fish. . . . .	168,133,000

---

Total. . . . .	1,414,523,374
Lobster. . . . .	81,020,000

---

1,495,543,374

Of the above, 99 per cent were in the interest of the commercial fisheries, and 1 per cent, or 14,900,000, were game fishes. The number of applications for fish to stock interior waters was 73,814, and the distributing cars of the Bureau of Fisheries travelled 95,259 miles, and sixty-eight railways furnished free transportation for 29,616 fish cars and 68,940 trips of messengers.

In the matter of fish propagation and distribution for the stocking of new or depleted waters, the national government stands pre-eminent. The only defect in its policy lies in its failure to protect existing fisheries from over-fishing, and from such reckless waste as is now destroying the salmon fisheries of Alaska.

# A PRACTICAL ARRANGEMENT OF BASED CHIEFLY UPON VISIBLE CHARACTERS; DEEP-SEA

SUBCLASSES	ORDERS AND CHARACTERS	TYPES AND EXAMPLES	PAGE
<b>LUNG FISHES:</b> Nearest to the <i>Amphibia</i> .	<b>LUNG-FISHES,</b> . . . . . <i>SI-REN-OI'DE-I</i> . . . Fishes with partial lungs, rudimentary legs, and molar teeth.	<b>Australian Lung- Fish</b> . . . . . 380 <b>South American Mudfish</b> . . . . . 381	
	<b>SPINY-FINNED FISHES,</b> . . . . . <i>AC-AN-THOP'TE-RI</i> Typical fishes, with perfect gills, fins, scales, and bony fin-rays.	<b>Bass</b> . . . . . 382 <b>Sunfish</b> . . . . . 384 <b>Perch</b> . . . . . 386 <b>Bluefish</b> . . . . . 387 <b>Mackerel</b> . . . . . 388 <b>Tuna</b> . . . . . 489 <b>Mullet</b> . . . . . 390 <b>Red Snapper</b> . . . . . 391 <b>Dolphin</b> . . . . . 392 <b>Swordfish</b> . . . . . 392 <b>Remora</b> . . . . . 393	
	<b>PIKES,</b> . . . . . <i>HA-PLO'MI</i> . . . . . Head flattened and scaly. Only one dorsal fin, far back.	<b>Pike</b> . . . . . 394 <b>Muskallunge</b> . . . . . 394 <b>Pickrel</b> . . . . . 395	
	<b>TROUT and SALMON,</b> <i>I-SO-SPON'DY-LI</i> . . . Differential characters relate wholly to bony anatomy.	<b>Trout</b> . . . . . 396 <b>Salmon</b> . . . . . 398 <b>Tarpon</b> . . . . . 406 <b>Shad</b> . . . . . 407 <b>Whitefish</b> . . . . . 408 <b>Herring</b> . . . . . <b>Menhaden</b> . . . . .	
<b>BONY FISHES:</b> Typical Fishes, high and low forms.	<b>FLYING FISHES,</b> . . . . . <i>SYN-EN-TOG'NA-THI</i> Pectoral fins greatly enlarged ; some species able to fly.	<b>Flying Fish</b> . . . . . 409	
	<b>SOLID-JAW FISHES,</b> <i>PLEC-TOG'NA-THI</i> . . . With solidified teeth and strong jaws. Mostly with rough, file-like skins.	<b>Trigger-Fish</b> . . . . . 410 <b>Box-Fish</b> . . . . . 410 <b>Puffer</b> . . . . . 410 <b>Poreupine Fish</b> . . . . . 411	
	<b>SUCKERS and MINNOWS,</b> . . . . . <i>PLEC-TO-SPON'DY-LI</i> Differential characters based wholly upon bony anat- omy.	<b>Common Sucker</b> 412 <b>Buffalo Fish</b> . . . . . 413 <b>Carp</b> . . . . . 413 <b>Minnows</b> . . . . . 414	
	<b>HALF-GILLED FISHES,</b> . . . . . <i>HEM-I-BRAN'CHII</i> . . . With imperfect or incomplete gills.	<b>Stickleback</b> . . . . . 415	
	<b>CATFISHES,</b> . . . . . <i>NEM-A-TOG'NA-THI</i> Scaleless ; head broad and flattened ; barbels around mouth ; defensive spines in dorsal and pectoral fins.	<b>Mississippi Cat- fish</b> . . . . . 416 <b>Bullhead</b> . . . . . 417	

**THE ORDERS OF LIVING FISHES**  
**ORDERS OMITTED. TYPES CHIEFLY NORTH AMERICAN**

SUBCLASSES	ORDERS AND CHARACTERS	TYPES AND EXAMPLES	PAGE
<b>BONY FISHES:</b> <i>Continued.</i>	<b>FLAT-FISHES, . . . . .</b> <i>IET-E-RO-SO'MA-TA</i> Without bilateral symmetry. Both eyes on one side. Flat, oval. Swim in horizontal plane.	Common Hal- but . . . . .	418
	<b>FOOT-FISHES, . . . . .</b> <i>PE-DIC-U-LA'TI</i> Mouth enormous; body broad, flattened, bag-like. Pectoral fins long at base.	Angler . . . . .	420
	<b>EELS, . . . . .</b> <i>AP'O-DES</i> Body long, slender, snake-like. No ventral fins, no scales.	Electric Eel . . . . .	421
	<b>PIPE-FISHES and SEA-HORSES, . . . . .</b> <i>LO-PHO-BRAN'CHI</i> Gills tufted; mouth tubular; body covered with scale armor. Very unlike true fishes.	Pipe-Fish and Sea-Horse . . . . .	423
<b>GANOIDS:</b> Armored Fishes, and their allies.	<b>The DOGFISH, . . . . .</b> <i>HAL-E-CO-MOR'PHI</i> Air bladder cellular, acting as rudimentary lung. Helmet-headed.	Dogfish or Bow- fin . . . . .	424
	<b>GAR-FISHES, . . . . .</b> <i>GING-LY-MO'DI</i> Ancient forms, covered with formidable bony armor.	Gar Pike . . . . . Alligator Gar . . . . .	425 425
	<b>STURGEONS, . . . . .</b> <i>GLAN-I-OS'TO-MI</i> Body with rows of large, bony plates. Mouth with barbels.	Lake Sturgeon . . . . .	427
	<b>PADDLE-FISH, . . . . .</b> <i>SEL-A-CHOS'TO-MI</i> Scaleless, shark-like. Broad, bony paddle projecting from nose.	Paddle-Fish . . . . .	429
<b>CHIMERAS:</b>	<b>CHIMERAS, . . . . .</b> <i>CHI-MAE-ROI'DE-I</i> Odd, shark-like forms.	Chimera collel . . . . .	431
<b>CARTILAGINOUS FISHES:</b> With soft skeletons. Lowest Fishes.	<b>SHARKS, . . . . .</b> <i>SQUA'LI</i> Scales minute; skeleton cartilaginous. Many species bring forth their young alive.	Mackerel Shark . . . . . Hammerhead . . . . .	432 433
	<b>RAYS and SKATES, . . . . .</b> <i>RA'I-AE</i> Excessively flattened, but otherwise shark-like. Many species with long, whip-like tails.	Shark-Ray . . . . . Sting Ray . . . . . Devil-Fish . . . . .	434 435 435

## CHAPTER XLIV

# ORDER OF THE CONNECTING-LINK FISHES, WITH LUNGS AND LEGS

### SIRENOIDEI

As in the preceding sections of this work, we will begin our studies of the Class of Fishes with the highest forms, and run down in regular course to the lowest. Of the 144 Families composing this class, as it occurs in North America, it is impossible to mention separately more than a

congeners lie embedded in Jurassic rocks 500,000 years old; and how this poor orphan of the Past escaped with its life down to the Present, many have wondered, but nobody knows. As you stand before the glass tank in the end of the Reptile House of the London Zoo, and behold a



THE AUSTRALIAN LUNG-FISH.

very few of those which are of greatest importance.

The **Lung-Fishes** are introduced because they are the highest of all the fishes, and form the connecting link between that class and the amphibians. Of the three genera that are known, one is found in Australia, two in Africa, one in South America, and in North America, none.

To some ichthyologists, the great **Australian Lung-Fish**<sup>1</sup> is the most interesting of all fishes. It is not only an intermediate form between the amphibians and fishes of to-day, but it is a creature that has far outlived its natural fate. Its

<sup>1</sup> *Ceratodus forsteri*.

magnificent living *Ceratodus* four feet long, with an ancestry running back half a million years without a break, it makes one's brain whirl to reel in the idea. This creature's ancestors lived in the days when many fishes were struggling to develop legs and lungs, with which to go on land, and become salamanders first, then lizards. It is said that this fish sometimes leaves the water and goes about on adjacent mud-flats, like the jumping fish of the Malay Peninsula; but the statement needs confirmation.

The Australian Lung-Fish is from 4 to 5 feet long, and it is said that its maximum weight is about 20 pounds. It breathes air over its palate



like a reptile, and its swimming-bladder is so developed that it does duty as low-class lungs. Its gills are very small and imperfect, and of little use. The top of its skull is quite unlike those of other fishes, and its scales are very large. Its pectoral and ventral fins are very long and leg-like, and are covered with scales everywhere save on the edges, where the fin-rays are situated.

One of the most extraordinary features of this strange fish is the possession of large and very remarkable molar teeth, those above being set in the palate (vomer), and evidently designed for the cutting up of vegetable food. Leaving bony anatomy out of consideration, it is quite clear that the living fish which stands nearest to

*Ceratodus* is the jumping fish or mud-skipper, of the Malay Peninsula, which hops about on land with surprising independence and agility. Its long pectoral fins are really foot-like in usefulness.

Both in the Burnett and Mary Rivers of Queensland, where it lives, and also in captivity, this Lung-Fish frequently rises to the surface of the water to take breath, like a porpoise.

The allied *Mud-Fish* (*Lepidosiren*) of the Amazon, and the *African Mud-Fish* (*Protopterus*) of the River Gambia, have legs that are mere wisps of skin and flesh, and strongly resemble our *Amphiuma*, of the Class *Amphibia*. They are rarely seen alive in captivity.

## CHAPTER XLV

# ORDER OF THE SPINY-FINNED FISHES

### ACANTHOPTERI

Even of forms classed as North American, this gigantic and rather unwieldy Order contains 45 Families and 483 species. Fortunately the groups which are of general interest are sufficiently limited in number that it is possible to place representatives of them before the reader.

#### THE BASSES AND SUNFISHES.

##### *Cen-trar'chi-dae.*

The Bass and Sunfish Family enjoys, on the whole, the widest popularity of all the finny Families of North America. With due respect to the justly distinguished Trout Family, I believe its members are known personally to a much smaller number of people than those of the Bass Family. The reason is that the latter are abundant in the most densely populated portions of the United States, while the human neighbors of the trout are comparatively few.

This Family (of thirty species) leads from the narrow-bodied and athletic black bass, by regular gradations in breadth through the rock bass, calico bass and their allies down to the little gem-like sunfish, with the extreme width of body and the limit of smallness and timidity. The black bass fights like a wild-cat, the sunfish can be taken on a bent pin, at the end of a cotton string; but observe this proportion:

The Sunfish is to the Small Boy as the Black Bass is to the Man.

It is good to find in Nature a Family whose members run from top to bottom in a stair-like series; for if so studied, the natural sequence is a great aid to the memory. We therefore begin with the narrowest fish, and descend to the broadest.

Surely, the **Black Bass**, be his mouth large or small, is a fish fit to head a Family. You can catch an eight-pound yellow pike-perch, and think you have hooked a bunch of weeds; but if

you hook a two-pound Black Bass you know at once that you have engaged a Fish.

For its size, this is the bravest and the gamiest fish that swims in our waters. In size and in silver the tarpon is truly the silver king of game fishes; but if he had Black-Bass energy and courage in proportion to his size, no hook-and-line angler in a small boat would bring him alive up to the end of a twelve-ounce rod.

The Black Bass has the narrowest body and the darkest color found in the Bass Family. It is built for speed and strength, and colored for concealment. There are two species, so very much alike that there is practically but one point of difference—the size of the mouth; and naturally their habits are quite identical. It is important to remember, however, that in color and markings, individuals vary most strangely and unaccountably. Some are uniform dark and light; others are mottled, much and little.

The **Small-Mouthed Black Bass**<sup>1</sup> is the fish of the East and North, from western New Hampshire to Manitoba, and southward to South Carolina and the northern Gulf states to Arkansas.

It is a pity that so fine a fish should not be handsomely colored, but it is really very plain and unattractive. Its back is usually a uniform dull olive-green, the sides being somewhat lighter. A Bass of three pounds weight may fairly be counted a large one, but this species has been known to attain a length of  $18\frac{3}{4}$  inches, and a weight of 5 pounds.

This is strictly a clear-water fish, and for this reason its capture is a source of pleasure beyond anything that can be drawn from muddy waters. It takes live minnows, or worms, or a neat trolling spoon, but resists the hook and the dip-net to the last extremity. Its flesh is excellent, and its propagation a matter of both state and national


<sup>1</sup> *Mi-crop'ter-us dol'o-mieu.*

importance. It has been planted successfully in so many bodies of water outside its original range that the limits of the latter are likely to be lost to view.

The Small-Mouthed Black Bass has the corner of its mouth directly under the front angle of the eye, while the mouth of the **Large-Mouthed**<sup>1</sup> species terminates under the rear corner of the eye. The range of the latter is from Manitoba, southward to the Gulf states, and spreads

bass is so close as to be at first sight a little confusing. But spread the dorsal fin to its full extent, and it will tell the story. In the Rock Bass it is long, rather low, and its front half contains eleven stout spines, of nearly equal length. The calico bass has a short and high dorsal fin, with only seven large spines; and the body of the fish is of greater depth.

The Rock Bass is a fish of the Great Lakes region and Mississippi valley—a clear-water fish,



Drawn by J. CARTER BEARD.

1. SMALL-MOUTHED BLACK BASS.  
2. CALICO BASS.

3. COMMON SUNFISH.  
4. YELLOW PERCH.

through the latter to Texas and Florida. In twelve months of 1897-9 the catch of Black Bass for market amounted to 1,785,418 pounds, valued at \$100,095.

**The Rock Bass, or Red-Eye,**<sup>2</sup> stands next in width of body to the black bass, and intermediate between it and the next species. Sometimes the resemblance between the Rock and calico

of habits quite similar to the black bass. Every way considered, it is a very perfect connecting link between the black bass and the next species. Its weight seldom exceeds 1½ pounds.

**The Warmouth Bass**<sup>3</sup> is a fish of the South, and in form is an intermediate between the rock bass and calico bass.

**The Calico Bass, or Strawberry Bass,**<sup>4</sup> is a

<sup>1</sup> *M. sal-moi'des*.      <sup>2</sup> *Am-blop'li-tes ru-pes'tris*.

<sup>3</sup> *Chae-no-bryl'tus gu-lo'sus*.      <sup>4</sup> *Po-mox'is spa-rox'des*.

handsome and substantial fish. Its bright, silvery coat is beautifully mottled with olive-green blotches, so regularly splashed on as to suggest the pattern of a piece of calico.

Take, if you please, a beautiful bay on the southern shore of Lake Ontario, a sunny day in May, no hotels or cottages in sight, with red-winged blackbirds singing "*O-ka-lee'*" in the cat-tails, and the Calico Bass becomes one of the prettiest fish you can pull out of the water. Each time, it gives a firm and vigorous bite, and leaves the water with a *swish* that once heard under proper conditions lives long in the memory.

I like the Calico Bass because it is so handsome, so well set-up, so substantial on the string, and so delicious on the table. A large specimen measures only about ten inches in length, but by reason of its great depth of body, and its thickness, too, it is a fish well worth having. Its weight never exceeds two pounds, and usually is about one pound. Besides the names given above, it is called the Grass Bass, Bar-Fish and "Crappie"; but the latter name belongs to another species.

The Calico Bass is at home throughout the whole region of the Great Lakes, the valley of the Mississippi to Louisiana and Texas, and along the Atlantic side down to the Carolinas and Georgia. In the beautiful lakes and ponds of Michigan, Wisconsin and Minnesota it is abundant, and highly valued. It can be taken still-fishing with worms, minnows, and grasshoppers, and also with a small trolling spoon.

It dislikes warm and muddy waters, it is a clean feeder, not quarrelsome or destructive to weaker species, and is said to increase rapidly. Strange to say, the propagation of this fine fish has received scanty attention from American fish-culturists, and in 1900 only 7,544 were distributed by the United States Bureau of Fisheries. It seems to me that for stocking northern lakes and ponds this is one of the most desirable of all the smaller fishes; and I wish long life and prosperity to the Calico Bass!

**The Crappie**<sup>1</sup> is a muddy-water understudy of the preceding species. In some portions of the North, the two species overlap each other, but in the main the Crappie is a southern fish.

<sup>1</sup> *Po-mox'is an-nu-lar'is*.

In 1900, the number distributed by the United States Bureau of Fisheries was 151,653.

**The Sunfishes** are divided into fifteen species, and as a group their range covers the whole of the United States eastward of the Great Plains. Poor indeed in fish life is the pond or stream between Maine and Texas, Dakota and Florida which contains no sunfish, bream or blue-gills, pumpkin-seed or dollaree. In about nine cases out of ten, the first fish that dangles from the first hook-and-line of the very small American angler is a sunfish. Small though it be, and feeble, it is yet a *Fish*; and it is large enough to open to Childhood the door to a great wonder-world of fish and fishing. Where is the veteran fresh-water angler who does not recall the electric thrills of his first "bite," and his first living, wriggling, scintillating Sunfish! Blessings be upon their rainbow-tinted sides for the joys they have been, are, and yet will be to Childhood!

Out of so many species it is difficult to select representatives, but it seems that first choice should fall upon the following:

**The Common Sunfish, or Pumpkin-Seed.**<sup>2</sup>

—This is the brilliant olive-green, blue and orange-yellow fish which when taken dripping from the water has all the colors of a green opal, and several more. It is distinguishable by the touch of bright scarlet on the lower portion of its gill-covers. It is found in clear ponds, large brooks and other streams from Florida, northward and eastward of the Appalachian chain to Maine, thence westward through the Great Lakes region to Iowa and Manitoba. It is subject to considerable variations in color markings.

In the Great Lakes, this fish attains a weight of 1½ pounds, but elsewhere a specimen 6 inches in length and weighing 8 ounces is considered a large one.

**The Blue-Gill or Black-Gill**<sup>3</sup> is the largest of the sunfishes. Its opercle, or gill-cover, terminates on the side in an ear-like flap which is of a deep black color; and this conspicuous character at once proclaims the species. This fish is found throughout the Great Lakes region and Mississippi valley. It sometimes attains a length of 12 inches, and a weight of 1½ to 2 pounds, and in some localities it is an important market fish.

<sup>2</sup> *Eu-po-to'mis gib-bo'sus*.

<sup>3</sup> *Le-po'mis pal'li-dus*.

## THE SEA-BASS FAMILY.

*Ser-ran'i-dae.*

In the ocean and its dependencies there exists a Family which, in general form, and anatomy also, so closely resembles the fresh-water Bass Family that it is almost impossible to base distinctive characters upon skeletal differences. The Sea-Bass Family, of North American waters, contains 104 species, some of which are of colossal size. Whenever you go a-fishing in tropical or subtropical waters, and catch a large, thick-bodied, big-scaled fish that you cannot name, it is generally a safe hazard to call it a Sea-Bass.

The great **Jewfish**, or **Black Sea-Bass**,<sup>1</sup> of Santa Catalina anglers, is one of the largest of the spiny-finned fishes. During the last ten years it has become celebrated because it permits itself to be outwitted so easily by the amateur angler. A very large fish can be caught with rod and line that seem absurdly light for such work.

What must we think of the courage of a 300-pound fish which will permit itself to be caught and gaffed on a line which will break under a strain of 50 pounds dead weight?


With heavy tackle, the catching of a large Jewfish would be no more of an event than would the pulling in of a Greenland halibut; but to go with one companion miles out from shore in a boat weighing from 125 to 150 pounds, catch a 300-pound fish on a sixteen-ounce rod, and kill it, without even getting upset, is a feat worth while. If a large Sea-Bass possessed the courage and fighting qualities, pound for pound, of the fresh-water black bass, it would take a hawser and a donkey engine to handle the line, a tug-boat to withstand the shock, and a bomb-lance to kill the fish when alongside.

In the Tuna Club of Santa Catalina (southern California), the holders of cups and records for the capture of Sea-Bass during the past five seasons were as follows: H. T. Kendall, Pasadena, 1902, 419 pounds; A. C. Thompson, Pomona, 1901, 384 pounds; F. S. Schenck, Brooklyn, 1900, 384 pounds; T. S. Manning, Avalon, 1899,

<sup>1</sup> *Ster-e-o-le'pis gi'gas*. It should be remembered that in another genus of this Family, called *Centro-pris'tes*, there is another species, found along our Atlantic coast, that is also called the Black Sea-Bass.

370 pounds; and F. V. Rider, Avalon, 1898, 327 pounds.

On our Atlantic coast, from Charleston to Brazil, occurs another huge fish to which the popular name "**Jewfish**" is applied. It is really the **Black Grouper**.<sup>2</sup> Its normal weight is 500 pounds, and "only one specimen weighing less than 100 pounds" has been recorded. (Jordan and Evermann.) I have reason to



Photographed by IRONMONGER.

## BLACK SEA-BASS.

Caught at Santa Catalina, with rod and reel, by MRS. A. W. BARRETT, of Los Angeles. Weight, 416 pounds; length, 7 feet 10 inches; girth, 5 feet 11 inches. Time, 2 hours and 15 minutes.

know the appearance of this fish quite well. It is a great, hulking, coarse-grained creature, unattractive to the eye, except that of the successful light-tackle angler, and very inferior on the table. In no point has it the look of a high-class animal, for every line is coarse and plebeian; but it has the avoirdupois of a Wonder.

**The Striped Bass**,<sup>3</sup> **Rock-Fish**, or **Rock**, is the finest representative of the whole great Family of Sea-Basses. It is a fish of handsome

<sup>2</sup> *Gar-ru'pa ni-gr'i'ta*.

<sup>3</sup> *Roc'cus lin-e-a'tus*.

form and colors, its table qualities are excellent, and it is a persistent breeder. Its ground color is silver-white, on which is laid, along the upper two-thirds of the body, a series of seven



STRIPED BASS, OR ROCK-FISH.

straight, equidistant stripes of black. It is a fish of large size, often attaining a weight of 85 to 90 pounds, and its flesh is most excellent. In the markets it stands next in desirability to the shad and bluefish. The greatest weight recorded for this species (by Dr. G. Brown Goode) is 112 pounds.

The centre of abundance of this fine fish is now from Fire Island, New York, to Albemarle Sound, on the coast of North Carolina. Many great catches have been reported, the most notable of which were the following: At Bridgehampton, N. Y., in 1874, 8,000 in less than a week; by Charles Ludlow, 1,672 bass at one set of a seine; at Norfolk, Va., 1,500 at one haul; in eight days of June, 1879, off Fire Island, one fisherman caught 10,164 pounds.

The full range of the Striped Bass is from the St. Lawrence River to New Orleans, both along the coast and in all the great rivers which flow into that region. At Cuttyhunk Island, and in scores of other places also, angling for this fish by heaving and hauling through the surf is pursued as one of the most fascinating kinds of sport.

One of the greatest hits ever made by the United States Bureau of Fisheries in the planting of fish in new localities was the introduction of the Striped Bass into the coast waters of California. In 1879, 135 live fish were deposited

in Karquines Strait, at Martinez, and in 1882, 300 more were planted in Suisun Bay, near the first locality chosen.

Twelve years after the first planting in San Francisco Bay, the markets of San Francisco handled 149,997 pounds of Striped Bass. At that time the average weight for a whole year was eleven pounds and the average price was ten cents per pound. Fish weighing as high as forty-nine pounds have been taken, and there are reasons for the belief that eventually the fish of California will attain as great weight as those of the Atlantic and the Gulf.


The San Francisco markets now sell, annually, about *one and one-half million pounds* of Striped Bass. This fish has taken its place among anglers as one of the game fishes of the California coast, and affords fine sport. Strange to say, however, it has not as yet spread beyond the shores of California.

Regarding this species, the latest records of the United States Bureau of Fisheries are of interest. In 1897, the California markets handled 2,949,642 pounds, worth \$225,527. In 1897, which is the last year fully reported upon, the catch for the whole United States amounted to 5,996,882 pounds, worth \$440,222.

### THE PERCH AND PIKE-PERCH FAMILY.

*Per'ci-dae.*

The festive little **Yellow Perch**<sup>1</sup> is, to the small angler, the next step upward from the



1. YELLOW PIKE-PERCH.  
2. CHAIN PICKEREL.

sunfish—a sort of half-way fish on the road to the black bass and tarpon. And there is many a

<sup>1</sup>*Per'ca fla-ves'cens.* See illustration on page 383.

worse thing in the fishing line than a good string of golden-yellow and umber-brown Perch. When crisply and daintily fried in a small modicum

$4\frac{1}{2}$ ; and in length it measures from 7 to 12 inches.

**The Yellow Pike-Perch<sup>1</sup>** is frequently called the **Yellow "Pike"** and **Wall-Eyed "Pike"**; but it is not a real pike at all. The real pike is a blood brother to the muskallunge. The Pike-Perches have *two prominent dorsal fins*, the real pikes only *one*.

Twice in trolling with hand-lines I have caught my spoon full of eel-grass. On hauling in to clear the tackle, each time the eel-grass turned out to be an eight-pound Yellow Pike-Perch. The first one came into the boat like a bunch of wet weeds. The second finally roused to a realizing sense of its position, and made quite a demonstration, but chiefly in the boat, endeavoring to climb out.

In the eastern United States, this is a northern fish that goes southward almost to the Gulf States. It is abundant in Lakes Ontario, Erie and Huron, and in many of the bays and




Photo. by E. F. KELLER.


#### BLUE AND YELLOW ANGEL-FISH.

*Hol-a-can'thus cil-i-ar'is*, a tropical species, about 15 inches in length, which is one of the most beautiful fishes in the world. It represents the Family of Sealy-Finned Fishes, *Chac-to-don'ti-dac*.

of meal, and laid on hissing from the spider, they are "pan-fish" worth while; and they make up in delicacy and richness of flavor all that they lack in size. Except in famine times, an ounce of Yellow Perch is worth a pound of pike, carp or catfish.

Like egg-rolling rights on the White House lawn every Mayday, this neat little fish belongs to the small citizen; but in the great lakes and a few other places it is so numerous and so large, that it takes rank as a desirable market fish. It is at home in the north-eastern quarter of the United States, north of the Ohio and Missouri valleys from Maine to Iowa and Minnesota. In most of the lakes, ponds and fresh-water bays of New England generally it is fairly abundant. Its rule of life is to bite at everything that is offered at the end of a line—angle-worm, minnow, grasshopper, frog-leg, trolling spoon, and fly, either natural or hand-made. The size of this fish varies from half a pound to three pounds, with a possible

larger streams attached to them, in which the water is clear, and the bottom of sand and gravel. By very many persons this fish is well liked as a food fish, and immense numbers are propagated every year. In 1900 the United




THE BLUEFISH.

States Bureau of Fisheries distributed, of this species, 89,700,000 eggs and live fish.

<sup>1</sup> *Sti-zos-tes'd'i-on vil're-um*.

### MISCELLANEOUS SPINY-FINNED FISHES.

**The Bluefish<sup>1</sup>** is a fish for men. To take it in orthodox fashion, go to treeless but delightful Block Island, pay your dollar-fifty, take deck passage on a low-browed, broad-beamed cat-boat, don a full suit of oil-skins, and set sail for blue water. If the wind is so light that the sailing is uninteresting, you get no fish. But if there is a stiff breeze, and you go up and down the eastern side of the island at racing speed, the Bluefish will come chasing after you to bite at your dummy fish, and give you a hundred thrills to the minute while you are hauling them in. If it happens that the bite of a bear has



THE SPANISH MACKEREL.

put two of the fingers of your right hand out of commission, that hand will have all it can possibly do to grasp the line adequately, and haul in.

Fishing for Bluefish in a good breeze—with not too much sea on—is like hunting mountain-sheep amid grand scenery. Half the sport is in the fine surroundings.

Drs. Jordan and Evermann say that this fish is found all the way from our coast to the Cape of Good Hope, the Mediterranean, the Indian Ocean and the Malay Archipelago, “a wandering fish . . . sometimes disappearing from certain regions for many years at a time.” Professor Baird always considered it, of all our coast fishes, one of the most destructive to marine life, a genuine wolverine of the sea.

<sup>1</sup> *Po-mal'o-mus sal-la'trix.*

The Bluefish swim in schools, ready to pounce upon anything edible that comes along. Once a cat-boat from which four of us were fishing sailed swiftly through a school. Within about five seconds, four fish struck in a rush that was practically simultaneous, and amid flying spray and general excitement, four vigorous victims of misplaced confidence were hauled aboard. A fish which is so greedy that it kills more fish-prey than it can use surely is a good fish to pursue for sport.

On our coast this fine fish is fairly common from Florida to northern Maine, ranging in size from 5 to 20 pounds. As a food fish it ranks on the bill of fare next to shad. Owing to its known voracity, it is debited with the annual destruction of an enormous quantity of other fishes. On the hook it is savagely courageous, and fights to the last.

Of all North American fishes, this species stands fifth in commercial value, being surpassed only by the Pacific salmon, cod, shad, and mullet. In 1897—the last year fully reported—the catch amounted to twenty million pounds, worth \$643,705.

**The Spanish Mackerel<sup>2</sup>** may stand as a typical representative of the Mackerel Family (*Scombridae*), in which we find the *Common Mackerel* of the North, the *Kingfish* of our tropical waters, and the *Tuna*. It is a large and showy fish, colored silvery white and dark metallic blue, and no cruise in Floridian or Cuban waters is complete without it. It is a favorite in all markets reached by it, and in flavor it is a fair rival of the bluefish.

To every sportsman, the finest thing about this fish is the catching of it, on a one-hundred-foot line and a hook baited with that least appetizing of all baits this side of angle-worms—a white rag! Like the bluefish, the Spanish Mackerel and kingfish both bite best when the sails are well filled, and the boat is making about twelve miles per hour. In 1902 the total catch for the United States amounted to 1,703,224 pounds, valued at \$112,342.

It would require many pages to contain a really adequate life-sketch of this interesting fish, which ranges most erratically, in great schools—or in none at all—from the Gulf of Mexico to Block Island. It comes north only in the spring and

<sup>2</sup> *Scom-be-rom'o-rus mac-u-la'tus.*



summer, and does not go far into waters that are colder than 65°. (G. Brown Goode.)

Apparently, specimens taken in northern waters average much smaller than those taken around the two coasts of Florida. Dr. Goode says this fish "sometimes attains a weight of 8 or 9 pounds, though it rarely exceeds 3 or 4 pounds." A specimen of 3 pounds, 5 ounces, measured 26½ inches in length. Drs. Jordan and Evermann give its weight as "6 to 10 pounds," with a maximum of all "seen" of 25 pounds weight, and 41 inches in length. ("American Food and Game Fishes".)

The great leaping **Tuna**<sup>1</sup> of the enchanted waters of Santa Catalina, "the tiger of the California seas," is, on our Atlantic Coast, the big but commonplace **Horse Mackerel, Tunny** or **Great Albacore**,—no more, no less. It is the largest and now the most interesting member of the Mackerel Family.

At Santa Catalina, bold men, and women, too, go out with rod, reel and line, to angle for this monster, and vanquish Strength and Weight by Tackle and Skill.

This is hook-and-line fishing with a vengeance. The beginner hopes to catch a Tuna heavier than 100 pounds, in order to gain membership in the Tuna Club. The club member always hopes either to improve his own record, or break all others; but, record or no record, the button of the Tuna Club is a good thing to wear by right of conquest.

Beyond question, when treated as a game fish, and fairly challenged with rod and line in the watery arena of Santa Catalina, in more senses than one the Tuna is great! Mr. C. F. Holder—for two years literally the holder of the Tuna championship with a 183-pound fish which fought four hours, and towed his captor ten miles—says that the Tuna, "when played with a rod that is not a billiard cue or a club will give the average man the contest of his life. My idea of a rod is a 7 or 8½ foot greenheart or split bamboo, with a good cork grip above the reel, the latter of Edward vom Hofe's make, with a leather pad, brake and click."


The sport in catching a Tuna *à la* Santa Catalina consists in bringing the monster within gaffing distance by the aid of the rod and reel alone. The hooked fish leaps into the air, or

<sup>1</sup> *Thun'nus thyn'nus*.

rushes seaward, or to the bottom, or plays on the surface like an escaped fire-hose,—in all directions at once.

The game consists in tiring out the fish without a break, and sometimes ten miles and ten hours of strenuous struggle are reeled off between the start and the finish.

The beautiful waters of Avalon Bay, the bare and frowning mountain-sides rising like the walls of a rock-built coliseum, and the houses of the little town clustering at its foot like a gathering of living and interested spectators,



Photographed by IRONMONGER.

#### THE TUNA.

Caught at Santa Catalina, with rod and reel, by MRS. E. N. DICKERSON, of New York. Weight, 216 pounds. Time, 1 hour and 55 minutes.

make up a stage setting for the Tuna fisherman sufficiently romantic to quicken the sporting instinct of the most blasé tourist who ever swung a rod.

Concerning the kind of tackle in use by the members of the Tuna Club, and by himself, Col. C. P. Morehouse, of Pasadena, holder of the

Tuna championship record, has kindly furnished the following statement:

"The most of the Tuna fishermen use a green-heart rod, as per the rules of the Club, viz., 6 feet 9 inches long, and a 16-ounce tip. As for myself, I prefer a split bamboo of the very best quality made. I caught the large Tuna (251 pounds) with the longest and lightest rod ever used for Tuna, viz., split bamboo, 7 feet 4 inches long, tip 12 ounces, with a 21-strand Cuttyhunk line on a reel made to order, to carry 300 yards. The time was 3 hours and 20 minutes. I do not think a lighter rod than the above would stand the strain necessary to capture a Tuna of 251 pounds, or even 150 pounds.


"The Tuna are hooked by trolling from light

sustain a dead weight not exceeding forty-eight pounds.

On our Atlantic coast, the Horse Mackerel is not sought by anglers as a game fish. Its average length is put down as "about 8 feet." It feeds chiefly upon menhaden, and inasmuch as its appetite is in proportion to its size, it is considered very voracious. In its turn, this great fish is to the killer (*Orea gladiator*) an ideal food fish, and from the latter it receives special attentions which the Tunny would gladly forego.

One of the largest specimens on record, as vouched for by Dr. Storer, was taken in 1838, off Cape Ann, and measured 15 feet in length. Its weight of "one thousand pounds" was undoubtedly an estimate, only.

**The Pompanos.**—Following closely after the members of the Mackerel Family comes a large Family of deep-bodied fishes, with very small and narrow scales, deeply forked tails, and with the dorsal and anal fins prolonged to nearly, if not quite, one-third the entire length of the fish. They are really warm-water fishes, but often stray out of their regular haunts into colder waters. This Family includes the amber-jack, the cavallas, the moon-fishes, and several others. Of this Family, the



THE SILVER MULLET.

naphtha launches, and flying-fish are used for bait."

At this date (1903), the five heaviest catches of Tuna by members of the Tuna Club stand as follows:

	Pounds.
Col. C. P. Morehouse, Pasadena, 1900..	251
John E. Stearns, Los Angeles, 1902....	197
C. F. Holder, Pasadena, 1899.....	183
F. S. Schenck, Brooklyn, 1901.....	158
F. V. Rider, Avalon, 1901.....	158

The rules of the annual tournaments in which such records are made are very severe and strict. The angler must make his catch unaided, the fish must be reeled in, and a broken rod constitutes a disqualification. The rod must measure not less than six feet nine inches, the tip must not exceed sixteen ounces, the line must not contain more than twenty-four threads, and

following species is the best type:

**The Common Pompano<sup>1</sup>** is a delicious fish for the table, but unfortunately its mouth is so small it is next to impossible to take it with a hook. Once when penned up by bad weather in the mouth of New River, Florida, where this fish was abundant, we fished for Pompano until we almost starved. The "Silver King" tantalized us daily by showing himself at the surface, but his vagrant pounds of flesh were almost as far beyond our reach as the stars.

The Pompano is essentially a fish of the two coasts of Florida, and the northern half of the Gulf of Mexico. It is the most highly prized fish in the markets of its home waters, and as a rule the supply seldom is equal to the demand. The *Jacks* are more common. Several of the species found in this Family are characterized by the enormous thickness of their ribs,—a

<sup>1</sup> *Trach-i-no'tus car-o-li'nus.*

very peculiar character, which makes them look like ribs afflicted with elephantiasis.

Mr. John T. Granger, of Washington, regards the *Permit*, or *Great Pompano*<sup>1</sup> as a game fish well worthy of the attention of salt-water anglers, and believes that it will become a general favorite. A struggle with a 27-pound fish, taken with rod and reel at Miami, Florida, revealed to Mr. Granger the game qualities of the Permit.

**The Mullet.**—Throughout the sounds, and bays, and half-salt rivers of the Carolinas, Florida, and the Gulf states, the mullets are omnipresent and highly prized. When better fishes fail you, they can be depended upon to fill the dish; and you may go far without finding a more toothsome morsel than a **Silver Mullet**,<sup>2</sup> or **White Mullet**, freshly snatched from its native element with a fling of the cast-net that experience alone can give. If you wish to beguile the Silver King, you first catch a Mullet, or buy one, for bait.

The name of this fish brings vividly to mind the balmy air and placid waters of Indian River, Florida, in February; a little, mangrove-clad archipelago along its eastern shore; herons quawking hoarsely in the green tangle, and small fishes of glistening silver jumping a yard high in front of a lotus-eater's boat. The Mullet leaps high in the air, gleaming and dripping, from pure joy in being alive amid such beautiful surroundings; and, having attained his zenith, he relaxes and falls back broadside upon the water, with a startling "slap." In one quiet evening hour afloat, you may see thirty or forty Mullet leap out of water, and to some persons the sight is even more welcome than the flight of a bird.


The Silvery Mullet is a very trim little fish,—big-scaled, round-bodied and swift. In external appearance, it is very much like a pygmy tarpon, and quite as silvery. It is really a small fish, averaging about 9 inches in length, and as food for other fishes, and fish-eating birds, it is ideal. The brown pelicans of Pelican Island delight in this fish. When Mrs. Latham playfully squeezed the neck of a big, clumsy young pelican in the down, it promptly disgorged nine good-sized Mullet. I have seen a darter, with a neck one inch in diameter, swallow a nine-inch Mullet with relish and despatch.

<sup>1</sup> *T. goodii*

<sup>2</sup> *Mu'gil bra-sil-i-en'sis*.

The Mullet genus (*Mugil*) contains about seventy species, widely distributed throughout the warm waters of the world. Besides the species mentioned and figured above, the *Striped Mullet* is also abundant in the waters of our southeastern coast. Both are important food fishes, and are caught in great numbers for the southern markets. They are taken in gill-nets and cast-nets, and the largest specimens rarely attain a weight of 6 pounds.

Of all North American fishes, the Mulletts are fourth in commercial value. In 1897, the total catch amounted to 21,402,624 pounds, which sold for \$332,090. Of this, the yield to Florida alone amounted to 16,700,094 pounds.



THE RED SNAPPER.

### THE SNAPPER FAMILY.

*Lu-ti-an'i-dae.*

**The Red Snapper**<sup>3</sup> brilliantly represents a large and important Family of valuable food fishes, which in our waters contains about 35 species. Many of these fishes are handsomely and showily colored, red being the commonest and most conspicuous tint, with yellow tints of frequent occurrence. A typical Red Snapper is recognizable a hundred feet distant by the clear and beautiful crimson color which completely suffuses it.

The average market specimen is about 16 inches long, but it is stated that this species

<sup>3</sup> *Lu-ti-an'us ay'a.*

attains sometimes a length of 3 feet, and a maximum weight of 40 pounds. In the Gulf of Mexico, says Mr. Silas Stearns, they very seldom exceed 30 pounds weight, and the average is 8 or 9 pounds. It happens, however, that one can spend months on the coast of Florida, and around Key West, without even once seeing a Red Snapper reaching 25 pounds in weight.


This fish prefers to live on a rocky bottom, in holes and gullies where all kinds of marine animals and fish are abundant. These gullies occur at a depth of from twelve to forty-five fathoms, and are most numerous in the northern border of the great level plain of sand which

487 pounds of Red Snappers, worth to them \$171,234.

### ODD FISHES OF THE SPINY-FINNED ORDER.

The "Dolphin" of this Order is a fish, not a cetacean of the Class of Mammals; and its unfortunate popular name and sea-going habits cause between it and the true dolphins much confusion.

This is the mid-ocean fish with a long, paddle-like body, a dorsal fin which reaches in one unbroken sweep from head to tail, and which pos-



THE SWORDFISH.

stretches out as the Gulf bottom from Cedar Keys toward the delta of the Mississippi.

Within easy reach of Jacksonville, Florida, are fishing-banks so well populated by the Red Snapper, and other fishes, also, that excursions are made to them with great success. Dr. C. J. Kenworthy described for Dr. Goode ("Game Fishes of North America") a day's sport twelve miles off shore from Mayport, which for eighteen fishermen yielded 208 Red Snappers averaging 25 pounds each. The bait used was bluefish, young shark or skip-jack. The only serious drawback to this fish is the fact that "it should always be boiled, or cooked in a chowder."

In 1897, the fishermen of Florida caught 5,314,-

sesses when alive the wonderful iridescent colors which have tested the descriptive powers of so many voyagers.

This is the terror and destroyer of the flying fish. The "Dolphin" pursues it with tremendous speed and perseverance, often taking long leaps out of the water, until the victim is exhausted, overtaken and devoured.

The colors of the "Dolphin" are a mixture of all the colors of the solar spectrum, revealed with the metallic lustre and iridescence of the opal and the reticulated python. The fully grown fish is from 5 to 6 feet in length, and in contrast with the ordinary sailing-vessel diet

*Cor-y-phac'na hip-pu'rus.*

of salt meat, its flesh is a delicacy. To the writer it was a red-letter day when with a new artificial flying-fish, fresh from the horny hand of an old sailor named "Porpoise George," he caught his first "Dolphin," in mid-ocean, from the deck of the *Golden Fleccc*.

**The Swordfish**<sup>1</sup> needs neither preface nor introduction, for his sword serves all such purposes.

In the government museum at Singapore is a three-inch-thick section of copper-sheathed oak plank, cut from the side of a ship, which has sticking through it the sword of a Swordfish. Now, the material of such a sword is by no means so hard that it could by ordinary means be forced through three inches of the hardest kind of oak planking, sheathed with copper. The fact of clean penetration implies a speed of not less than sixty miles per hour, and perhaps more. With such locomotive powers, and such a weapon for slaughter, it is fortunate that its owner has not been fitted out with the teeth and appetite of a killer, else the cetaceans would soon be exterminated.

The Swordfish well understands the offensive and defensive value of his sword, and there are on record many well-authenticated instances wherein this pugnacious creature has driven its formidable weapon through the sides or bottoms of small boats, to the peril of the occupants. The majority of such incidents have occurred to boats regularly engaged in swordfishing, which is a noteworthy industry on our Atlantic coast.

Broken swords have been found in the sides and bottoms of quite a number of ships. In 1871, the fishing yacht *Redhot*, of New Bedford, was pierced and sunk by a Swordfish which had been hauled alongside to be killed. In 1875, a Swordfish drove its sword through the bottom of a fishing schooner off Fire Island, and stuck fast. Before the fish had time to free itself by breaking off its sword, the fishermen cast ropes about it, and secured it. Its length was over 11 feet, its weight 390 pounds, and the length of its sword, 3 feet 7 inches.

The Swordfish is a food fish of very good standing in New England, where it is sliced and salted, and widely esteemed. In 1898, the total catch was 1,617,331 pounds, valued at \$90,130.

The food of this fish consists of menhaden, mackerel, bonitoes, bluefish, herring, whiting and squids.

**The Sucking-Fish, or Remora**,<sup>2</sup> is a high-class parasite, who does much of his travelling at the expense of sharks who would eat him if they could. In one of her odd freaks of merry-making, Nature fashioned on this creature's head a large, flat disk, set crosswise with rows of delicate spines, all pointing backward. It is really a peculiar development of the first dorsal fin. When the Sucking-Fish desires to travel and see the sea-world, it hunts up the nearest shark, swims alongside from the rear, claps its head to the shark's side, and sticks fast. The faster the shark glides through the water, the more tightly clings the automatic tramp. Like a passenger in a Pullman sleeping-car, the Remora can bid the world good-night, and go to sleep serenely confident that he will get on in the world, even while he sleeps. It is as if a human tramp were provided by Nature with means enabling him to cling automatically and comfortably to the side of a moving freight-car, instead of walking in dust and sorrow upon the ties.

The Remora is not a large fish, its usual length being under two feet. Not only is it a parasite of sharks and other large fishes, but it attaches itself to the sides of ships. It is said that sometimes sharks actually become emaciated through prolonged labor in furnishing free transportation for lusty Remoras. The parasite is himself a good swimmer, and the best reason assignable for its strange habit in clinging to sharks is its desire to gather in fragments of the feast when the latter makes an important killing. The Remora is an inhabitant of our Atlantic coast, the Gulf of Mexico, and the West Indies generally, but it is not considered a food fish.

<sup>1</sup> *Xiphias gladius*. The pronunciation of the generic name is Zif'e-as.

<sup>2</sup> *Remora brachyptera*. See figure on page 432, of an individual attached to a mackerel shark.

## CHAPTER XLVI

# THE ORDER OF PIKES

HAPLOMI

After the Order of Spiny-Finned Fishes, with its great array of genera and species, it is a relief to reach an Order which contains but one Family, and only five species. The so-called Yellow "Pike" is not a member of this aristocratic and exclusive Family; for, as already stated, it is only a pike-perch.

Look at any member of the Pike Family, and tell me whether it does not make you think of a pirate. Observe that yawning sepulchre of a

amphibians generally, ducklings, other small aquatic birds, and also small aquatic mammals. Worse than this, they even devour their own kind. That they are found living with the bass, perch and other fishes is generally due to the fact that it is impossible for them to devour all their neighbors.

**The Pike**<sup>1</sup> is a fish of very wide—almost world-wide—distribution. In America it is found from Kodiak Island, Alaska, southward through British Columbia, Canada, the upper Mississippi valley and the Great Lakes region, to Europe and Asia. Dr. Jordan says ("Food and Game Fishes") that it reaches a length of 4 feet, a weight of 40 pounds, and that the Kankakee River, in Illinois, is one of the best streams for great Pike fishing of which he knows. As a food fish the Pike ranks low.

**The Muskallunge**<sup>2</sup> is a game fish of high rank, and its Indian name is spelled in eight different ways. Its standing may be expressed in the following proportion: *The Muskallunge is to the fresh-water angler as the tuna*



THE MUSKALLUNGE.

mouth, that evil eye, and low, flat forehead—all indicating a character replete with cunning and ferocity. Note the total absence of a dignified and respectable *front dorsal fin*, which nearly every fish of proper moral character possesses and displays with pride.

Like scaly assassins, the pikes and pickerels lie in wait for their prey; and whenever one rushes like a green streak from under the lily-pads, and bolts a trolling- spoon in one great, ill-mannered gulp, the angler feels a savage delight in thinking that it serves him right. These fishes are the most voracious creatures that inhabit our inland waters. Their ambition is to devour every living creature that comes in sight, and they prey upon all other fishes, frogs and

*is to the salt-water angler.*

Its great size makes it a great prize, and the taking of a large fish with sportsman-like tackle, and a very good chance to be upset in deep water during the struggle, makes the Muskallunge the king of fresh-water game fishes. The northern species—of the Great Lakes, the St. Lawrence and southern Canada—reaches a length of 7½ feet, or more, and attains a maximum weight of about 90 pounds. Its centre of abundance seems to be in the Thousand Islands of the St. Lawrence, where it affords grand sport. Usually it is caught by trolling,—a most delightful scheme by which the twin pleasures of boating and fishing are combined.

<sup>1</sup> *E'sox lu'ci-us.*

<sup>2</sup> *E'sox mas-quin-on'gy.*

**The Chautauqua Muskallunge**<sup>1</sup> is a species quite distinct from its more northern relative. It is confined to Chautauqua Lake and a few localities in the Ohio valley—a comparatively small area. In that landlocked region, far from the shad and the bluefish, it is by many persons considered a fine fish for the table.

**The Chain Pickerel**<sup>2</sup> is so common throughout the region bounded by Maine, Florida, Arkansas and Minnesota, that it is difficult to say where it is not found. It is so well and so widely known that it requires neither introduction nor description. On the lovely lakes of central Michigan, and New York, to stand up in a boat that is properly handled, and throw a trolling-spoon along the borders of the lily-pad archipelagoes,

where the Pickerel hide, is good sport. In the crystal-clear water the whirling, glittering spoon is in sight every moment, and you can see the rush of the Pickerel when he flies straight as an arrow at the lure. This fish is so voracious that several kinds of bait are effective for it; but I see no reason for calling its flesh a delicacy. Its maximum size is about 28 inches, which is considerably larger than the little *Brook Pickerel* of the northern Mississippi valley.

In a period of twelve months ending in 1899, the total market catch of Pike, Muskallunge and Pickerel, as reported to the United States Fish Commissioner, amounted to 1,041,293 pounds, worth \$47,773. The Fish Commission makes no serious efforts to propagate these species.

<sup>1</sup> *E'sox o-hi-en'sis*.

<sup>2</sup> *E'sox re-tic-u-la'tus*. See illustration on page 387.

## CHAPTER XLVII

# THE ORDER OF TROUT AND SALMON

### ISOSPONDYLI

This grand Order is represented in North American waters by 135 full species of fishes, all decidedly edible, and the majority of them are classed as "game" fishes. It includes not only some of the most choice of all our finny tribes, but also others whose commercial value is of the highest rank. In it are found the trout, salmon, whitefish, shad, herring, menhaden and tarpon. Despite the great number of species in the Spiny-Finned Order (446), it seems highly probable that their combined value in the markets falls far below the aggregate for the Order now under consideration. On the Pacific coast, the value of the annual salmon catch alone is, at this date, about \$13,000,000, whereas the annual value of the cod, the most valuable food fish of the Atlantic, is only \$2,000,000.

Reminding the reader once more that we are endeavoring to present groups in the order of their natural rank and importance, we present first in this Order of fishes the Family of highest interest and value.

### THE SALMON FAMILY.

#### *Sal-mon'i-dae.*

The Salmon Family contains all the trout, salmon and whitefishes, to the number of thirty-two full species and twenty-nine subspecies. Of these three groups, the first is celebrated for the beauty of form, picturesque surroundings, and gamy qualities of its members. The salmon and whitefish are noted chiefly for their great value as food.

Few persons, it is safe to say, know either the size or the subdivisions of the group of American trout and charrs. The species are numerous, beautiful, and widely distributed north of a line drawn from New York City to San Diego, California. For a clear and correct understanding of these fishes, a diagram is absolutely necessary. The world is indebted to Dr. D. S. Jordan, Pres-

ident of Leland Stanford University, for the researches which have made him the leading authority on this large and extremely interesting group of fishes, and by means of which it has been made comprehensible.

Of North American trout, generally, the centre of abundance is certainly west of the Rocky Mountains, and the group as a whole is decidedly of the Far West. The trout of the eastern United States are but the advance guard of the main body which fills the swirling mountain streams and lakes of the Rocky Mountain region and the Pacific coast. Our famous and well-beloved speckled trout of the East is but a trifling incident in comparison with the many fine species found in the true home of the Trout Family.

Dr. Jordan believes that our original stock of trout came to us from Asia, and "extended its range southward to the upper Columbia, thence over the continental divide via Two-Ocean Pass to the Yellowstone and the Missouri Rivers, the Platte, Arkansas, Rio Grande and Colorado." He actually caught Yellowstone trout in Two-Ocean Pass, on the top of the great continental divide, "in the very act of going from Pacific into Atlantic drainage."

**The Mountain Trout, or Black-Spotted Trout.**<sup>1</sup>—Like many others, this is a fish of many names,—Spotted, Black, Silver, Salmon, Steelhead and Cut-Throat,—all ending with Trout. The last mentioned,—“Cut-Throat Trout,”—Dr. G. Brown Goode characterized as “a horrible name, which it is hoped will never be sanctioned in literature.” And why “Cut-Throat,” any more than Ripper Trout, or Wife-Beater Trout?

Surely this fine fish, which Dr. Jordan considers probably the parent from which all others of this group have been derived, is worthy

<sup>1</sup> *Sal'mo clark'ii.*




of a dignified and respectable name. It is a fish of large size, attaining a length of 3 feet, and a weight of 30 pounds. It is *the* fish of the Rocky Mountain region, and occurs in nearly every lake and important stream of Montana, Wyoming, Utah, Colorado, New Mexico, Idaho, Oregon, Washington and northern California. It reaches the sea from Mount Shasta northward to Puget Sound, and beyond. "Those that live in the depths of shady lakes are almost black, while others are pale. Those in the sea are sil-

is regarded as the greatest of all game fishes." It "reaches a weight of half a pound to 5 or 6 pounds, though in most of the streams in which it is found it rarely exceeds 2 or 3 pounds." It bites readily, but when hooked makes a gallant fight to escape, rushing, leaping, and shaking its head vigorously to expel the barb.

In appearance, this typical Rainbow Trout is like an elegant little salmon from 15 to 18 inches long, with spots along its upper body like those of the eastern brook trout, and sides like a section

**SUBDIVISIONS OF THE NORTH AMERICAN TROUT AND CHARRS.**

(Species in italics are introduced in the text.)



very, or only faintly spotted." (G. Brown Goode.)

In the most representative specimens of this species, the upper half of the body is abundantly spotted with small, round, and rather regular black spots.


**The Rainbow Trout<sup>1</sup>** is a fish of real beauty, comfortable size, fine flavor, and easy to propagate artificially. On this side of the Rocky Mountains, however, it is not politic to assert that it is more beautiful than the brook trout; but Dr. Jordan says that "by many anglers it

of a rainbow. It is found only in the small brooks of the coast ranges of California, from Klamath River to San Diego. It takes a fly with a degree of readiness which "will please the most impatient of inexperienced amateurs."

The group of Rainbow Trout contains six species all told, the others being the *Western Oregon Brook Trout*; the *McCloud River Rainbow Trout*, which is the species propagated by the United States Bureau of Fisheries; the *Kern River Trout*; *Golden Trout* of Mt. Whitney and Kern River, which Dr. Jordan considers the most beautiful of all, and *Stone's Trout*. All

<sup>1</sup> *Sal'mo ir-id'e-us.*

these species are found only along the Pacific coast, between Washington and southern California.



RAINBOW TROUT.

**The Steelhead Trout,<sup>1</sup>** and its group.—The fish which represents this group is of commanding size, and of high value as a food fish. Its other names are **Salmon Trout** and **Hard-head**. It reaches a maximum weight of 14 pounds, but usually its weight is between 5 and 8 pounds. It “ranks very high as a game fish, and trolling for Steelheads in the bays, sounds and river mouths along our Pacific coast affords excitement and pleasure exceeded among the *Salmonidae* only by trolling for Chinook Salmon.” (Jordan and Evermann.)

This fish is regularly propagated by the United States Bureau of Fisheries, by which it has been successfully planted in Lake Superior. Great numbers are caught every year in the Columbia River, and canned for the eastern markets. It is found in the streams flowing into the sea along the coast of California, from southern California to Alaska. Its scales are small, its form is salmon-like, and its color is silvery, with a wash of rose-pink down the sides.

**The Great Lake Trout, or Mackinaw Trout,<sup>2</sup>** and its group.—This fish is the largest of all trout. Its usual weight is from 15 to 20 pounds, but it reaches a maximum of 125 pounds. Its color is dark gray, varying most erratically from pale gray to almost black. Its irregular

and very numerous spots of gray mark this fish very distinctly, for they cover not only the body but all the fins save those under the body.

As its name implies, this is essentially a fish of the Great Lakes, and for many years has been the principal source of fresh-fish supply for a large area in that region. In its own field its only rival in commercial importance is the white-fish. Usually the flesh of the latter is supposed to be a greater delicacy than the other.

The Lake Trout has passed through two or three very interesting periods. From 1880 to 1886, commercial fishing for Lake Trout was carried on so persistently that the supply showed alarming signs of exhaustion. Here the United States Bureau of Fisheries stepped in, and along with state hatcheries began to propagate and distribute this species. This work was continued until many millions of fish eggs had been planted in the lakes. After that, the supply of Lake Trout increased so rapidly that presently the markets were overstocked, and the price dropped accordingly.


More recently, however, the pendulum has swung the other way. All around the Great Lakes the demand for food fishes is now so great and so permanent, that the natural supply has proven unable to meet it. Nature cannot produce food fishes in the lakes in the enormous quantities required, even though in 1899 the yield of Lake Trout was ten million pounds (10,611,588). To-day the United States Bureau of Fisheries is doing its utmost to help maintain the supply, and in 1900 distributed 337,838,000 eggs and young of the Lake Trout.

“Lake Trout spawn on the reefs, and at other times live in deep water. In Lake Superior the spawning season begins in late September. In Lakes Huron and Michigan, the height of the season is early November, and spawning continues until December. The spawning grounds are on the reefs of ‘honey-comb’ rock, 10 to 15 miles off shore, and in water from 6 to 120 feet deep. The number of eggs produced is not large. A 24-pound fish produced 14,943 eggs, but the usual number does not exceed 5,000 or 6,000.” (Jordan and Evermann.)

The range of this fish is from New Brunswick and Maine westward throughout the Great Lakes to Vancouver Island, B. C., and northward to Labrador, Hudson Bay and northern Alaska.

<sup>1</sup> *Sal'mo gaird'ner-i*.

<sup>2</sup> *Cris-ti-vo'mer nam'ay-cush*.



DRAWN BY W. L. STEWARD.

THE BROOK TROUT.

**Deep-Water Fishing for Lake Trout.**—“The Siscowet of Lake Superior is taken by the commercial fishermen in very deep water, the nets being lifted by steam power. The nets are set well out toward the centre of the lake, at depths frequently as great as 500 feet. About forty nets, each over 600 feet long, are set in one ‘gang,’ constituting practically a single gill net considerably over four miles in length. Each end of each gang is buoyed.

“The average steam fishing-boat attends to five gangs of nets, lifting one each day. Each gang, therefore, remains in the water five days before it is lifted. As the net comes up around the steam windlass forward, it is passed aft and immediately reset by being paid out over the stern by two members of the crew. The nets are about eight feet wide, and the mesh is  $4\frac{1}{2}$  inches.

“The largest Lake Trout I observed on the *Currie* was 2 feet 10 inches long, and its weight was 21 pounds. The average length of the fishes taken during my inspection was less than 2 feet.” (Charles H. Townsend.)

**The Brook Trout, or Speckled Trout, and**

its group.—Concerning this beautiful and high-spirited creature, so much has been written it would now seem that there is nothing untold. But this is a very wide country; and I ween that in the real West there may be a million of good citizens who are strangers yet to *Sal-ve-ti-nus jon-ti-nal’is*.

After all has been said, I think it must be conceded that this is the most beautiful of all our game fishes. Its back and dorsal fins are elegantly marbled, its sides have about fifteen or twenty crimson and black spots, and its pectoral, ventral and anal fins are bright crimson, edged in front with white. Its general ground color down to the latitude of the pectoral fin is dark olive, below that comes sunset pink, and underneath all is the silver white of the belly.

Along with its beauty, agility, and general gaminess, this fish makes its home in the most picturesque and beautiful streams its range affords. Its ideal haunt is a deep, clear pool at the foot of a picturesque rush of water over mossy boulders. Usually this forest jewel is delightfully set in the foliage of overhanging

birches, beeches and maples, and well backed by the forest shadows that painters love. Usually the music of rushing water pervades the haunt of the Brook Trout; and the only cloud upon it all is that, ever and anon, Man, the supposedly high-minded, savagely bends every energy to kill an unduly-great number of these beautiful creatures, and fills a sordid creel entirely too full.

Most unluckily for the Trout, it is its habit to be ever on the alert for insects on the surface of its pool, and "rise to a fly." To the high-class sportsman who scorns the humble angle-worm, the accurate throwing of a small fly for a very long distance, solely by the exercise of great skill and judgment, is the crowning attraction in seeking the Brook Trout in its haunts. The skill required in fly-fishing is enough to tempt any man who has ever felt the electricity that every good fly-rod is charged with; and it is no wonder that men love to fish for this very beautiful fish, in the most charming of all sylvan situations.

The Brook Trout was once a habitant of the northeastern United States, northward of a line drawn from New Jersey to Minnesota, into Labrador, Canada and Manitoba; but to-day, where is it? Ask the "fish-hog" who spares no Trout that is big enough to lift from a platter. Ask the market fishers, who fish and fish to supply hotels and restaurants, in season and out of season.

In its wild state, this fish is doomed to disappear at an early date. We have now in this country a large and rapidly increasing element the members of which have come to us to slay and eat. To them, the preservation of wild life *to look at* seems like childish folly. These, and others like-minded, are raking our trout-streams with fine-toothed combs, and mean that nothing larger than a trout egg shall escape. And the end will be that in a very few years the wild Brook Trout will be as nearly extinct as the wild buffalo.

**THE SALMON GROUP.**

The salmon were made for the millions. The Siwash Indian eats them fresh in summer, dries them, and later on freezes them, for himself and his dogs in winter. The epicure pays for

having the fresh fish shipped in ice to his table, wherever that table may happen to be. In mid-ocean, the great American canned salmon is often the best and only fish afloat. In the jungles of the Far East, in the frontier bazaar of the enterprising Chinese trader, it "bobs up serenely" to greet and cheer the lonesome white man who is far from home and meat markets. Even in the wilds of Borneo its name is known and respected; and he who goes beyond the last empty salmon-tin, truly goes beyond the pale of civilization. The diffusion of knowledge among men is not much greater than the diffusion of canned salmon; and the farther Americans travel from home, the more they rejoice that it follows the flag.

The common salmon of Europe, and also of Labrador and New England, was accounted a wonderful fish, both for sport and for the table, until the discovery of the salmon millions of the Pacific coast effectually cheapened the name. To hold their place in the hearts of sportsmen, game fishes positively must not inhabit streams so thickly that they are crowded for room, and can be caught with pitchforks. Yet this once was true of the salmon in several streams of the Pacific coast. The bears of Alaska grow big and fat on the salmon which they catch with the hooks that Nature gave them.

The salmon species of North America are as follows:

AMERICAN SALMON.	ATLANTIC Species:	Atlantic Salmon. (Of Europe and N. America.)
		Ouananiehe. (The leaping fighter.)
		Sebago.
	PACIFIC Species:	Quinnat, or Chinook. (Most valuable species.)
		Blueback, or Sockeye. (Second in value.)
		Silver, or Coho. (Third; flesh white.)
		Humpback, or Gorbusha. (Of little value.)
		Dog, or Kayko. (Of least value.)

The five species of Pacific coast salmon form a remarkable group. They lead all fishes in annual commercial value (\$13,000,000); they are the most abundant of all fishes that inhabit fresh water; they traverse very great distances

to reach their spawning beds, and they all die immediately after spawning!

The sea is the home of all the Pacific salmon, and except when the young are floating toward it from their birth-place, it contains their food. Of their life in the sea, little is known. They are nowhere numerous, and trolling for them in salt water is interesting sport.

Throughout the months of spring and summer, the salmon leave the sea, enter the large rivers,—and many small ones, also,—and proceed upward for hundreds of miles, to deposit their eggs as far as possible from salt water. In the Columbia and Yukon Rivers, and their tributaries, it “is the habit of salmon to ascend for a *thousand miles* or more before spawning!”


The “run” begins with the advent of spring, when the salmon travel up the rivers until they can ascend no farther. It is on these runs that the fish congregate in such incredible numbers that sometimes they actually crowd each other, and can be photographed *en masse*. They rush up rapids, and if cascades or low water-falls are encountered, they leap atop of them with a display of energy and activity that is, when first heard of, almost beyond belief.

“When the Pacific salmon reach maturity,” says Mr. Cloudsley Rutter, in “Country Life,” “they seek fresh water to spawn. As soon as they leave their accustomed salt-water food, they stop eating. It is not uncommon for fishes of the Salmon Family to fast during the breeding season, but the Pacific salmons never taste food after leaving salt water, and their fast ends only with death. This is true of all species of Pacific salmons, and is without a parallel among the higher fishes.

“As the salmon advances into fresh water, the digestive organs shrivel to one-tenth their natural size, all the fat disappears from the tissues, the flesh turns white, and the skin thickens and embeds the scales till they cannot be seen. By the time spawning begins the fish has lost about twenty per cent of its weight, and sometimes much more. In fresh water, the jaws of the males become much prolonged and hooked,

and large canine teeth appear. The body becomes deep and slab-sided; and the skin turns reddish in most species. No individual of either sex of any species of Pacific Salmon ever returns to the ocean after spawning.”

Concerning the Chinook salmon, Drs. Jordan and Evermann say that the run begins in the Columbia River as early as February or March. The fish move up without feeding, travel leisurely at first, but as they advance farther they move more rapidly. Many of them pause not until they have found satisfactory spawning beds in the Snake and Salmon Rivers, among the Sawtooth Mountains of Idaho, more than 1,000



THE QUINNAT SALMON.

miles from the sea. “Those which go to the head waters of the Snake River spawn in August and early September; those going to the Big Sandy in Oregon, in July and early August; those going up the Snake to Salmon Falls, in October; while those entering the lower tributaries to the Columbia, or small coastal streams, spawn even as late as December.”

“The spawning extends over several days, the eggs being deposited upon beds of fine gravel, in clear, cold mountain streams.” The temperature of the water must be about 54°, and if on arrival it is much above that, the fish wait until it lowers. (“American Food and Game Fishes.”)

A very remarkable feature about the spawn-

ing of the salmon is that after it is completed, both males and females *die!* "This," says Jordan and Evermann, "is true of all, whether spawning remote from salt water, or only a few miles, or yards, from the sea," and whatever the cause may be, it "is general in its application to all the Pacific coast salmon."

Inasmuch as the bodies of many dead salmon show injuries of many kinds, the belief has become prevalent that the fish injure themselves by striking against rocks on the run upstream, and ultimately die from wounds so received. But the investigations of Drs. Jordan and Evermann have completely disproved this. It was found that of the many salmon examined immediately after arrival on their spawning grounds in central Idaho, not one showed any bruises or mutilations, and all were in excellent condition. The mutilations which subsequently were observed were obtained either by fighting, or by pushing the gravel about on the spawning beds.

Salmon eggs hatch in about fifty days. During the first six weeks, the egg-sac supports the life of the alevin, which lives quietly on the spawning bed. By the end of six weeks the yolk-sac is absorbed, and the young fry begins to float down stream toward the sea. When the journey is very long, the trip occupies several months, or even a year; and when the young salmon at last reaches salt water, it is four or five inches long, and is known as a "parr." Of course the young salmon feed all the way down, on a fresh-water menu.

Naturally the salmon millions of the Pacific streams early attracted the attention and aroused the avarice of men who exploit the products of Nature for gain. As usual, the bountiful supply begat prodigality and wastefulness. The streams nearest to San Francisco were the first to be depleted by reckless over-fishing, and now some of the fishermen of California solemnly aver that the sea-lions are largely to blame for the depletion of the Sacramento salmon fishery! It is the rapacious and deadly net and salmon-wheel, not the squid-eating sea-lion, that is to blame. Regarding the conditions that in 1901 prevailed in Alaska, the following notes by Mr. George Bird Grinnell in the "Harriman Alaska Expedition" are of interest:

"The salmon of Alaska, numerous as they

have been and in some places still are, are being destroyed at so wholesale a rate that before long the canning industry must cease to be profitable, and the capital put into the canneries must cease to yield any return.

"The destruction of salmon comes about through the competition between the various canneries. Their greed is so great that each strives to catch all the fish there are, and all at one time, in order that its rivals may secure as few as possible. . . . Not only are salmon taken by the steamer load, but in addition millions of other food fish are captured, killed and thrown away. At times, also, it happens that far greater numbers of salmon are caught than can be used before they spoil. A friend of mine told me of the throwing away of 60,000 salmon at one time, near a cannery in Prince William Sound, in 1900; and again the similar throwing away of 10,000 fish. So something like 700,000 pounds of valuable fish were wasted."

In the Kodiak and Chignik districts, the catch of salmon decreased from 360,000 cases in 1896 to 90,000 in 1898, and in 1899 it was almost a failure. In many of the small Alaskan streams, the canning companies built dams or barricades to *prevent the fish from ascending to their spawning beds*, and to catch all of them! In some of the small lakes, the fishermen actually haul their seines on the spawning grounds.

The laws passed by Congress to prevent the destruction of the Alaskan salmon fisheries are "ineffective, and there is scarcely a pretence of enforcing them." To-day, the question is,—will lawless Americans completely destroy an industry which if properly regulated will yield annually \$13,000,000 worth of good food? Will the salmon millions of the Pacific share the fate of the buffalo millions of the Great Plains? At present *it seems absolutely certain to come to pass!* In the preservation of fish and game, ours is one of the weakest of civilized nations. Very shortly we may expect to see the salmon-hogs knocking at the doors of Congress to report that the salmon of Alaska are "all gone," and hear them plaintively petition for government appropriations with which to restock those waters, by artificial propagation.

The time for strong, effective and far-reaching action for the protection of that most valuable source of cheap food for the millions, is *now!*

**The Quinnot Salmon,**<sup>1</sup> also called *Chinook*, *California*, *King*, *Columbia River* and *Sacramento Salmon*, is the largest, the most widely distributed and the most valuable of the Pacific Salmon. Frequently it attains a weight of 50 pounds, and specimens have been taken in Alaskan waters weighing about 100 pounds. Its average weight is between 20 and 30 pounds. It is found from Monterey Bay, California, up the whole Pacific coast to Bering Strait, and down the Asiatic coast to China.

**The Blueback Salmon,**<sup>2</sup> also called *Sockeye*, *Nerka*, *Redfish* or *Red Salmon* is the most abundant species, and in flavor and general importance it stands next to the preceding. In Alaska it is of greater value than all other species combined. Its flesh is of a rich red color, full of oil, and of fine flavor. In size it is small for a salmon, its average weight being only about 5 pounds. Its geographic range is from the Columbia River to Japan, and it is the species most abundant in the canneries along the Fraser River and the shores of Puget Sound. In 1901 the number of Red Salmon canned in Alaska and elsewhere on the Pacific coast was 19,615,310.

The *Little Red Fish* of various lakes in Idaho, Oregon and Washington, wherein they reside continuously, are now regarded by Dr. Jordan as small and immature specimens of the Blueback Salmon. Like all others of their genus, they die immediately after spawning, sometimes bearing body bruises, and again quite free from them.

In Alaska the abundance of the Blueback is almost beyond belief. A catch of 10,000 fish at one haul is of common occurrence; 25,000 at a haul is not uncommon, and 50,000 are taken at least once every year. The record haul was made in 1896 when about 100,000 were taken, of which 60,000 were used and the remainder liberated. (Cloudsley Rutter.)

**The Silver Salmon**<sup>3</sup> stands third in the list. Its other names are *Kisutch*, *Hoopid*, *Skowitz*, *Coho Salmon* and "*White Fish*."

The range of this fish is from California to

Japan. It is next in size to the quinnot, but inferior in flavor, and its flesh is pale. It is a good fish to ship fresh, and despite the fact that when canned it is not highly esteemed, great quantities are canned in Oregon and Washington, and marketed as "medium-red Salmon." In 1901, the number canned in Alaska was 523,713.

**The Humpback Salmon,**<sup>4</sup> also called the *Gorbusha*, *Holia*, *Honc* and *Haddoh Salmon*, ranges from the Sacramento to Kamchatka. It derives its best name from the fact that "upon the approach of the breeding season, the back of the male grows higher than it usually is, and forms an abrupt hump back of the head," at which time the flesh is valueless.

Formerly this fish was not highly regarded by the canning establishments, and was but little used. As a matter of fact, its flavor when fresh, in the spring, is by some experts, of whom Mr. C. H. Townsend is one, considered fully equal in flavor to that of any other salmon. To-day it is receiving its full share of attention from the canning establishments, and is now quite on the market. Its place now and in the future is clearly indicated by the fact that in 1898, 109,000 cases, in 1899, 150,000, and in 1900, 232,022 cases were prepared for the market. In 1901 the number of fish of this species and of the dog salmon canned in Alaska was 11,301,886.

**The Dog Salmon**<sup>5</sup> is the least valuable of the Pacific Salmon. Its flesh is of such poor flavor that formerly it was ignored by the canners. Now, however, it is being put up under various names that are not its own, such as "Chum Salmon." This fish is also called *Hayho*, *Lekai*, *Qualoh*, and *Calico Salmon*. Its range is from Sacramento to Kamchatka; and in the rivers of Japan it is the most common species. Its weight is from 10 to 12 pounds.

As previously observed, the Salmon of the Pacific coast far surpass in commercial value all other American fishes. Their accessibility renders their capture little more than a mechanical operation, and eventually it will result in the practical destruction of the salmon industry. Americans seem utterly unable to conserve for perpetual benefit any particularly valuable form of wild life.

The records of the salmon industry for the

<sup>1</sup> The scientific name of this fish, *On-co-rhyn'chus tscha-wyts'cha*, is presented with an apology to the reader. The specific name is useful only as an example of the disgusting barbarism to which an ill-balanced mind can sometimes descend in choosing names.

<sup>2</sup> *On-co-rhyn'chus ner'ka*.

<sup>3</sup> *O. ki'sutch*.

<sup>4</sup> *On-co-rhyn'chus gor-bus'cha*.


<sup>5</sup> *O. ke'ta*.

year 1899 are of universal interest. They are as follows:

Of Pacific Salmon, Alaska	produced	118,622,230 pounds,	worth	\$6,773,876
Of Pacific Salmon, { California, } { Oregon and } { Washington }	produced	130,004,835 pounds,	worth	3,504,622
		248,627,065		\$10,278,498

To-day the question is, shall we permit this industry to go by default, and be ruined in a few years? Or shall we conserve it sensibly and properly, both for ourselves and future generations?

**The Atlantic Coast Salmon.**—It is now necessary to call this fish the **Atlantic Salmon**<sup>1</sup> in order to distinguish it from the Pacific species; but for two centuries it held its place in literature as *the* Salmon. It once inhabited many



Drawn by W. L. STEWARD.

THE SEBAGO SALMON.

portions of northwestern Europe, and in some it still survives.

In North America, its natural habitat was originally from the mouth of the Hudson River northward throughout the costal rivers of New England, Canada, New Brunswick, Nova Scotia, Newfoundland and Labrador to Greenland. Once very abundant in the Connecticut River, it was driven out of that stream in 1798 by the erection of a sixteen-foot dam in Miller's River, 100 miles from the sea, which cut off the fish from their spawning beds. In 1872 there were twenty-eight rivers in the United States which once contained Salmon, but from twenty of them that fish had totally disappeared. To-day the nearest Atlantic Salmon are found in Maine and northern New Hampshire, New Brunswick and Nova Scotia.

As a game fish, *Salmo salar* is fit to rank with the kings of the animal world. He who catches

<sup>1</sup> *Sal'mo sa'lar*.

one with a fifteen-foot rod weighing twenty ounces, with a Silver Doctor at the end of five-

foot leader, and brings it to the gaff, may well call himself an angler. So far as I know, this is the largest fish that rises to a fly.

The greatest weight on record for the Atlantic Salmon is 83 pounds. The maximum weight of those now taken in Maine is about 25 pounds, and the average is about 10 pounds. In 1900, the catch of the Bangor anglers in Penobscot Pool was 67 fish, weighing 970 pounds. The largest weighed 23½ pounds, and the average was nearly 14½ pounds.

The most wonderful thing about the Atlantic Salmon is its leaping power, in surmounting waterfalls that lie in its course to its spawning grounds. To a fish of this species, a rock-studded cascade three hundred feet long and thirty feet high, down which the water plunges and tears with murderous speed and violence, is a fine highway, up which it gayly promenades without pause or accident.

But a waterfall, with a perpendicular drop of ten or twelve feet, is a more serious proposition, and requires a special effort. To clear such a barrier, the Salmon makes a rush in the pool below it, leaps out of the water, and if possible lands on the edge of the fall. If he falls short by no more than one or two feet, but strikes the descending torrent squarely head on, so that he is not at once swept down, it is said that by a strong flirt of the tail and a wriggle of the body, the gallant fish actually can force itself on up to the edge of the fall, and over it into the coveted waters of the upper level.

The following graphic description of the leap of the Salmon is from the pen of Dr. Robert T. Morris, whose opportunities for observing and photographing the scenes he describes have been of the best:<sup>2</sup>

"It is a most impressive and inspiring sight to watch the untamed Salmon on a wild river mak-

<sup>2</sup> *Country Life Magazine*, 1903, p. 356.



ing his display of strength and agility in surmounting a crashing torrent that threatens with instant death anything that dares to approach its mad tumult of waters. A Salmon can make his way upward through a sheer fall of water so long as the water is in a solid mass, but the moment that it becomes admixed with air the white water no longer gives a sufficiently firm hold for the broad caudal fin, and the Salmon must leap entirely over the fall. There are pretty well authenticated instances of Salmon clearing a fall of twenty feet. I have measured leaps to nearly this length on falls where almost every Salmon that flew through the air over the fall fairly took one's breath away, and they were going up at the rate of three or four to the minute at that. I know of nothing short of watching a house on fire that is of more engaging interest than watching the Salmon throwing themselves over wicked waters. The Salmon must have some advantages, to be sure, for accomplishing their best feats. If the water beneath a fall is much broken with rocks and rapids a fish cannot gain sufficient momentum and velocity for hurling himself far into the air; but given a deep and fairly quiet pool to start from, and the Salmon look more like great birds than like fish as they sail upward. One can sometimes find a place to stand at the edge of a fall, and if he remain quiet for a few moments the Salmon will begin to go through the air over his head in quick succession."

Dr. Morris states that from the Penobscot River, in Maine, to Hudson Bay, Salmon enter almost every river on the coast, but south of the Straits of Belle Isle the sawdust and dams in the streams of the lumber region constitute most serious obstacles to their progress and existence. But "the time is coming when twenty rivers on the Maine coast will have their mills so managed in the interest of the Salmon that they will rival the historical streams of Europe. In Washington County alone there are six rivers that Salmon now ascend every year."

**The Ouananiche,**<sup>1</sup> whose name is of Indian origin, and is pronounced *win-nan-ish'*, is a fresh-water Salmon, dear to the heart of every angler who has ever brought one to gaff.

It is fondly spoken of as the "*Leaping Ouananiche*," and frequently as the *Landlocked Salmon*. It is neither more nor less than a

<sup>1</sup> *Sal'mo ouan-an-iche'*.

fierce-fighting, fresh-water understudy of the Atlantic salmon, which if not self-restricted to fresh water would hardly be described as an independent species. When first taken from the water, it has "a beautiful peacock-blue" color, which disappears at death, changing to the light-gray back and sides and silvery belly of the Salmon. Although called "landlocked," this fish can, and sometimes does, live in salt water,—in the mouth of the Saguenay River, for example.

The Ouananiche is a fish which loves rapids and rushing water as a mountain sheep loves crags and precipices. Because of the strenuous life it leads, it is beyond doubt the most vigorous and athletic fish that inhabits our waters.

Says Mr. Eugene McCarthy: "None of the fresh-water fish can equal its fighting powers, and, pound for pound, it will outfight even the salmon. Ouananiche are great smashers of rods and tackle, unless one understands how to play them, especially when they make their numerous high jumps from the water. It is not an exaggeration to state that these jumps will average at least five to six, and frequently will number ten to twelve feet. And such leaps! Two or three feet out of the water, often toward the fisherman, then a rush deep down, a pause, a succession of jerks that would seem to tear the hook loose, a wild rush of varying distance, and a run back, almost to the angler's feet. A fish weighing 3½ or 4 pounds will make a fight lasting ten or fifteen minutes, often longer; and that means hard work for every moment for the fisherman." ("Familiar Fish," p. 126.)

This fish is best taken with a fly, on a rod of from six to eight ounces, with No. 4 or 5 hooks. Its home is in Lake St. John, Province of Quebec, and its tributaries; its outlet, the Saguenay, and no one knows how many of the rivers of southern Quebec that flow into the Gulf of St. Lawrence; and also the rivers of Labrador.

**The Sebago Salmon,**<sup>2</sup> of Maine, is a strictly fresh-water, or "landlocked," species, which takes its name from Sebago Lake, its type locality. It is essentially a 15-pound fish, with an average in Sebago Lake of from 8 to 10 pounds. Owing to the quiet waters it inhabits, and the powerful tackle used in fishing for it, this fish does not manifest the vigor and fighting quali-

<sup>2</sup> *Sal'mo se-ba'go*.


ties that have made the ouananiche famous. This species bears to Atlantic salmon and the ouananiche so close a resemblance that it is yet an open question whether the three species should not be merged into one (*Salmo salar*). At all events, a picture of the Sebago Salmon might easily, under other names, be made to do duty in representing the other two!

**The Tarpon<sup>1</sup>** is one of the very few large fishes to which it is proper to apply the word *magnificent*. Either in the water or out, or hanging upon the wall of a dining-room, it is as its pet name implies, a Silver King, entitled to royal honors. Its enormous scales, its back of

Its flesh is excellent, and will always hold its place in the markets of the South.

In cruising around the coast of Florida, you first see the Tarpon breaking water, back in air, like an undulating porpoise. You may see fifty of them, and sail and fish for days before you catch one; but one big Silver King pays for a long journey, and ten days of cruising.

Twenty-five years ago, no one attempted to catch a 100-pound Tarpon with rod, reel and line of light weight. To-day, angling for this grand creature has become an established recreation, and on the Florida coast is regularly pursued as such at Fort Myers, Punta Rassa, Boca Grande



Drawn by J. CARTER BEARD.

#### THE TARPON.

royal blue and sides of burnished silver proclaim it at a glance, and in the presence of such external splendor we cease to care whether its flesh is savory or not. How the Romans would have doted upon this fish, had they but lived within its realm!

To-day it is beloved of every American sportsman who can get in touch with it, first because of its imposing personality, and next because of the difficulty in catching it with hook and line. It is taken by rod-and-reel fishing in lagoons, and also by trolling in "the passes" between islands.

<sup>1</sup> *Tar'pon at-lan'ti-cus.*

Pass, Marco, and Bahia Honda, on the adjacent coast of Cuba. Besides the above, Corpus Christi, Texas, and Tampico, Mexico, have become famous as resorts for Tarpon fishermen.

The size of this fish is entirely satisfactory. Usually its weight is between 100 and 200 pounds, but it is credited with a maximum record of 383. Specimens six feet long are by no means rare.

So far as known on January 1, 1904, the championship of Tarpon angling was then held by Mr. Edward vom Hofe of New York, with a fish of 210 pounds weight, a length of 6 feet 11 inches, and a girth measurement of 45 inches. Its larg-

est scales measured  $3\frac{3}{4} \times 4$  inches. The tackle used in the capture of this fish consisted of a short-butt snakewood rod seven feet long, of which the tip weighed thirteen ounces, a vom Hofe Universal reel, 600 feet of No. 24 vom Hofe line, and a No. 1 Van Vleck hook.

The Tarpon is not to be caught in deep water with hook and line. As a rule, the waters of the east coast of Florida are unsuitable for successful adventures with the Silver King; but at several points on the west coast, where the level beach of clear sand shelves far out into the Gulf before it drops into deep water, this grand fish loves to bask in the sunshine, and linger in the warm, placid waters along the shore.

The Tarpon fisherman goes out early, and casts his bait—a small mullet—upon the shallow waters. For hours he floats upon a sea of molten silver, bathed in a flood of dazzling sunshine, and at times grilling in the heat which comes with it. The clean leap out of water of a big Tarpon firmly hooked is a sight that no sportsman ever can forget.

In a few localities, Tarpon are really plentiful, and easily caught. Off Useppa Island, Florida, between March 5 and May 31, 1903, the total catch of Tarpon was 336.

**The Common Shad**<sup>1</sup> is, to many persons, the most savory of all American fishes. It possesses the maximum number of bones to the cubic inch, but its flesh is fine-grained, juicy, and of exquisite flavor. The freshest Shad is "the finest Shad," but when treated with even a show of culinary fairness, every fresh Shad is good.

Like the salmon, the Shad spends half its life in the sea, and enters the rivers of its choice only to spawn. Owing to the practical impossibility of taking Shad in the ocean, the shad-fishing season is limited to its spawning-season. This is one of the most prolific of our fishes, a single fish sometimes yielding 150,000 eggs. It is easily propagated by artificial means, and a decrease in the annual supply can in a measure be made good by the hatcheries of the United States Bureau of Fisheries. During the spring of 1900, the Agents of that Bureau planted 291,056,000 young Shad and eggs in the rivers of the Atlantic coast that are accepted by the species as breeding-grounds.

This fish is found all along our Atlantic coast

<sup>1</sup> *Al-o'sa sap-i-dis'si-ma.*

from Florida to Newfoundland, but it is most abundant from the Hudson River to the Potomac. Of all our fishes, it stands third in commercial value, being surpassed only by the quinnat salmon and the cod.

Including both the Atlantic and Pacific coasts, the value of the Shad catch for 12 months ending in 1899 was 49,780,530 pounds, worth \$1,519,946.

Originally, the Shad was not a habitant of Pacific waters; but in 1871, Mr. Seth Green, of Rochester, made for the California State Fish Commission the initial experiment of transporting 10,000 young Shad across the continent, and planting them in the Sacramento River. From



THE COMMON SHAD.

that year up to 1880, about 60,000 more fry were deposited in that stream by the United States Bureau of Fisheries. In 1885 and 1886, 910,000 Shad fry were planted in the Columbia and Willamette Rivers.

To-day, on the Pacific coast the Shad ranges from southern California to southern Alaska, and is one of the most valuable food fishes of that region. In 1899, the fish dealers of California alone handled 1,137,801 pounds, worth \$14,303.

The average length of the Shad is from 24 to 30 inches, and its weight is from 3 to 4 pounds. The color of the fish is a soft, silvery white, all over, but the scales are easily detached, and an immaculate specimen is rarely seen in a fish market.

To landlocked Americans of the upper Mississippi valley and the shores of the Great Lakes,

the **Common Whitefish**<sup>1</sup> is an undisguised blessing. To them it is all that the shad is to the East, or the salmon to the Pacific coast. Whenever the traveller between Cleveland and Omaha discovers before him a large fish of excellent flavor, he may be sure that it is either a Whitefish or a lake trout, from one of the great lakes, and worthy of profound respect.

But for the fact that this fish is so well and so widely known, many pages might be written of it without exhausting the subject. Dr. Jordan considers the Whitefishes the most important group of fresh-water fishes of North America, and probably of the world.

<sup>1</sup>*Co-re-go'nus clu-pe-i-for'mis.*

The home of this group extends from Niagara to Chicago and Duluth. The average weight of a typical fish is about 4 pounds, but specimens weighing 20 pounds have been taken. In 1899, the catch of Whitefish (all species) amounted to 6,862,094 pounds, worth \$345,640. In 1898 the catch in Canadian waters, say Jordan and Evermann, amounted to about 18 million pounds, worth \$877,000.

In winter, the Whitefish retires to the deepest portions of the great lakes, and is beyond the reach of fishermen. In the spring, it frequents the shallower waters, near shore, where it spawns, and lingers to fall a prey to the gill-net fishermen, even until late in the autumn.

## CHAPTER XLVIII


# THE ORDER OF FLYING-FISHES

### SYNENTOGNATHI

**The Common Flying-Fish**<sup>1</sup> is as necessary to a perfect ocean voyage as a whale and a school of "dolphins." Suddenly and unexpectedly it breaks out of the side of a wave, and with a tremulous flutter of wing-like pectoral fins,—that from the ship's fore-castle seem to be ultramarine blue,—it feebly guides its course away from the disturbing mountain of throbbing steel. The flight of a Flying-Fish is usually from four to six

in schools so near to the island of Barbadoes that the fishermen capture it in great quantities, for the markets. It is not unusual to see 2,000 in the market at one time. I have heard much of the pursuit of the Flying-Fish by the "dolphin" (*Coryphæna hippurus*), but have seen nothing of it.

**The Flying Gurnard** or **Sea Robin** (*Dactylopterus volitans*), is a wonderful pink fish, 8



Drawn by J. CARTER BEARD.

THE COMMON FLYING-FISH.

feet above the water, and is sustained for from 50 to 100 feet. The greatly enlarged pectoral fins act as wings, and furnish the motive power.

Someone has raised the question, "Does a Flying-Fish move its wings in flight?" Of course it does. On all up grades it gives a stiff wing-stroke about every three feet, rises to over-top each advancing wave, and drops as the wave rolls on, like a stormy petrel.

This is distinctly a mid-ocean fish, but it swims

<sup>1</sup> *Ex-o-cæ'tus vol'i-tans.*

inches long, that is found from Cape Cod to Brazil. It is often picked up on the sea-shore near pound-nets, because fishermen throw it away as unmarketable; but as fish become more scarce, *it will be eaten.* Its pectoral fins are of enormous size, but useless for flight. This fish is *not* closely related to the flying-fish, but belongs in the Order of Spiny-Finned Fishes. It is the only representative of its Family in the New World, and only one other species exists elsewhere.

## CHAPTER XLIX

# THE ORDER OF SOLID-JAW FISHES

### PLECTOGNATHI

The characters on which the members of this Order have been brought together are, for the general reader, rather obscure. They are strictly anatomical, and relate to the manner in which the teeth and bones of the jaw are grown together, and solidified. On the whole, it will be about as easy to become acquainted with the various groups of fishes composing the Order as to learn fully and correctly the precise anatomical characters which are common to all.

This Order contains some very odd and picturesque forms; and, fortunately for the student, good examples of them are fairly common along the Atlantic coast.

**The Trigger-Fish,<sup>1</sup> or File-Fish,** is a very good species to represent this entire group. It derives one of its names from the large, movable spine of solid bone (a fin-ray of the front dorsal fin), which stands upon the foremost point of its back, with a smaller trigger behind it, like that upon an old-fashioned hair-trigger rifle. The large spine can be set quite rigidly by a neat interlocking device supplied by the second spine.

This fish is a thin-bodied creature, and its skin has the toughness, the rigidity and even the external appearance of stamped leather, with the roughness of fine sand-paper. It is a fine fish for the first efforts of the amateur taxidermist, for it has ingrowing scales that cannot possibly come off, and its colors are equally fast.

All the Trigger-Fishes are habitants of tropical and subtropical waters, and feed chiefly upon small shell-fishes (mollusks) which their strong jaws and teeth enable them to masticate successfully. Some of them, like the *Orange File-Fish*, are brilliantly colored. In the tropics they are considered edible, but the few that exist along our Atlantic coast are not ranked as food fishes. The species shown in the illustration is the one

<sup>1</sup> *Ba-lis'tes ca-pris'eus*. See figure on page 374.

most widely known along our Gulf coast, and also the Atlantic coast up to the mouth of the Potomac. In the Bahamas and the Bermudas, the skins of Trigger-Fishes are extensively used by carpenters in place of sand-paper for smoothing the surface of wood previous to polishing.

**The Box-Fish, or Trunk-Fish,<sup>2</sup>** is one of the curiosities of the tropic seas, and of curio-shops generally. Its skin is a rigid, triangular box, shaped in cross-section like an isosceles triangle, and consists of large hexagonal plates of thin bone, joined firmly together by the regular integument.

Of these fishes we have four species on our Atlantic and Gulf coasts, and one off California. According to Dr. G. Brown Goode, all the species of Box-Fishes were so thoroughly and correctly studied by the fathers of natural history two hundred years ago, that their classification of the group has stood the test of time, and come down even into these troublous times unchanged and unimproved.

**The Bellows-Fish,<sup>3</sup> or Rabbit-Fish,** is possessed of many local names, such as *Globe*, *Bottle*, *Blower*, and even *Egg Fish*. When taken from the water, and scratched smartly on the abdomen against the grain of the small spines which cover that region, it begins to pump air into its interior, the skin expands like india-rubber, and in a moment it assumes balloon-like proportions. If the fish is then thrown into the water, it floats belly upward for a moment, then suddenly the air is expelled, the fish collapses, instantly turns right side up, and disappears.

This species ranges from Cape Cod to the Gulf of Mexico, and may be looked for with confidence in the pound-nets at nearly all our sea-side resorts.

<sup>2</sup> *Os-tra'd'i-on quad-ri-cor'nis*. See illustration on page 374.

<sup>3</sup> *Lag-o-ceph'a-lus 'lae-vi-ga'tus*.

**The Porcupine Fish,**<sup>1</sup> also known as *Puffer*, *Ball-Fish*, *Swell-Fish* and *Toad-Fish*, is another sea-side "curio," although usually it is stuffed not wisely, but too syell. A tow-filled balloon of fish-skin, with spines upon it, is not necessarily a Porcupine Fish; and the sea-side taxidermist should sometimes put a curb upon his zeal for expansion.

Like the bellows-fish, this species can expand itself with air to about twice its normal size. Its back is covered with strong, bony spines, which in some species are an inch in length. It is a fish of tropical waters, and in Cuba is considered a food fish. The species figured is one of four which in summer visit our Atlantic coast, while two others are found on the coast of California.

<sup>1</sup> *Chi-lo-myc'te-rus ge-o-met'ri-cus*. See illustration on page 374.

## CHAPTER L


# THE ORDER OF SUCKERS, CARP, AND MINNOWS

### PLECTOSPONDYLI

#### THE SUCKER FAMILY.

*Cat-os-tom'i-dae.*

This huge Order contains 60 genera and 311 species, divided into 4 Families. Of these Families, the Sucker Family is the most important. It contains about 70 species, all of which save two are inhabitants of North America. Besides the Suckers themselves the Family includes the buffalo fish, the red-horse and fresh-water "mullet." These fishes have the mouth placed underneath the head, and fitted with very fleshy, tubular lips, well adapted for sucking food from the bottom. They have been specially formed to live upon mud bottoms, and in murky water,—



THE COMMON SUCKER.

precisely the conditions that high-class fishes abhor.

There are times when a sucker (or a carp) seems like a good fish for the table; and that is when one is very fish-hungry, and there is no other kind of fish to be had. To my mind, the flavor of the flesh is either barely tolerable, or verging closely upon disagreeable. The very numerous and wholly unnecessary bones seem like a positive affront. Although these fishes are seldom eaten by choice, by the landlocked dwellers in the interior of our great continent, to whom clear streams and good fishes are only long-distance memories, the sucker, carp and bull-head are eaten with real relish, and a feeling of thankfulness that they are no worse. And after all, men who can eat musky squirrels, and

call them "game," ought to be pleased with suckers and carp.

The Common Sucker,<sup>1</sup> Brook, or White Sucker, is qualified to represent a large section of this Family. In the home of this fish, acquaintance with it nearly always begins in the month of June, when, if ever, come perfect days, and the annual spring "run" of Suckers, up river and creek to their spawning beds, brings them prominently into notice.

I remember one wildly hilarious day of boyhood, when a great run of Suckers came up Eagle Creek, Indiana, from the Ohio, via White River.

Now Eagle Creek is a very beautiful stream, flowing over a fine bed of clear gravel and sand.

Its waters are as clear as the sea, and the big sycamores that reach their long white arms across them are truly grand. All the young men and small boys turned out *en masse*, and rushed to a shallow, rock-bound channel above a big "drift." Each able-bodied "angler" was armed with a snare of soft brass wire loaded with enough lead to kill an elephant, and a pole

that would have driven a real angler to a mad-house.

The Suckers moved about restlessly in the swift current, and occasionally paused, head up-stream. That was the snarer's only opportunity; for the fish refused all baits. The heavily loaded snare was set as a hoop five inches in diameter, gently lowered ahead of the fish, and with a very steady hand and correct eye steered downward over the victim until it passed his pectoral fins. Then, at precisely the proper instant, steam was turned on, the erstwhile quiet fisherman became a raging demon of activity, and if the snare held just "so," a 16-inch Sucker weighing 4 pounds would be yanked high in air by a human derrick, amid the shouts of a de-

<sup>1</sup> *Ca-tos-to-mus com-mer-son-i.*



lighted and strong-lunged populace. The string of fish caught on that halcyon day by my tall brother reached from my shoulders to the ground, and for three days the cooks of that countryside had Suckers "to burn."

This Sucker is perhaps the most widely distributed and the most common fish species inhabiting the United States. It ranges "from Quebec and Massachusetts westward to Montana and Colorado, and southward to Missouri and Georgia." (Jordan.) It is one of the best of its Family for the table, it is universally eaten, and is much superior to any carp the writer has ever encountered. In one year (1899-1900) the catch of Suckers in twenty-three states yielded 655,637 pounds, worth \$115,512.

**The Red-Horse,**<sup>1</sup> or so-called "Mullet," which makes Ohio the centre of its distribution, is an abundant and well-known fish in the region west of the Alleghenies. It is rather handsome in colors, and, although its flesh is coarse and insipid, it is really an important food fish in its region.


**The Buffalo Fishes**<sup>2</sup> comprise three species, all big and burly, ranging in maximum weight from 35 to 50 pounds, and from 2 to 3½ feet in length. They inhabit the Mississippi and its tributaries, and in the spawning season push their way even into the larger lakes and flooded marshes of Wisconsin, Iowa and Minnesota. I have seen specimens weighing between 30 and 40 pounds caught in the Mississippi, at Burlington, Iowa, by hand-line fishing between lumber rafts, with about as much interest and enthusiasm on the part of the fisherman as usually attends the capture of a good strawberry bass. One fat and fearless "angler" sat on a chair, and baited his hook with cheese.

But let no one underrate the economic importance of the Buffalo Fish. The catch of 1899, chiefly in Illinois, Arkansas, Mississippi and Missouri, in the order named, amounted to 14,221,988 pounds, worth \$350,026.

**The German Carp**<sup>3</sup> was introduced into the United States by Mr. R. Poppe in 1872, and in 1877 by the United States Bureau of Fisheries, because of the fact that in Germany it is con-

sidered a good food fish, and can live and thrive in muddy ponds and streams. By thousands of prairie dwellers, it was received gladly, especially throughout the great plains, where any fish with scales is welcomed. The free distribution of young Carp led a great many persons to apply for them, and plant them in ponds, from which they afterward found their way into streams that contained fishes infinitely their superior.

Between the years 1877 and 1885, the streams of very nearly the whole Pacific coast of the United States were stocked with Carp. At first



GERMAN SCALED CARP.

they were placed in ponds, but through "moving accident, by flood and field," they reached the rivers, and impregnated them and all their tributaries. At first they were highly esteemed, and sometimes greatly overpraised. It was claimed that they were hardy, prolific, harmless to other fishes, rapid in growth, persistent under adverse conditions, and acceptable on the table. Beyond question, *under certain conditions* nearly all these claims are justified by the facts!

But when the novelty wore off the Carp, the cold-blooded critic began to say things. By him it was pointed out that Carp stir up the mud in all mud-bottomed ponds inhabited by them, and keep the water murky. This is quite true; and to keep the mud-loving Carp from perpetually soiling and disfiguring the once clear and beautiful waters of the Merced Lakes, in California, first sea-lions, and then muskallunge, were introduced to exterminate the Carp.

In California, the Carp is now ranked with the introduced catfish, as an unwelcome guest. It is claimed that Carp consume to a serious extent the wild celery and grasses on which wild ducks feed, and the duck supply is diminished

<sup>1</sup> *Mox-os-to'ma au-re-o'lum.*

<sup>2</sup> The Common Buffalo Fish is *Ic-ti'o-bus cyp-ri-nell'a.*

<sup>3</sup> *Cy-pri'nus car'pi-o.*

thereby; but this charge remains to be proven. The chances are as ninety-nine to one that the choke-bore shot-gun is the real and the only cause of the decrease in wild ducks.

It is also claimed that Carp eat the eggs of other fishes; which is extremely probable, for very many fishes do that.

Whatever may be said for or against the desirability of the Carp in America, one important fact remains unassailed. That fish is now thoroughly established in our waters, and is here to stay, just as much as the English sparrow. It is rapidly coming into demand as a market fish. "Over *seven million pounds* are consumed yearly in New York City. From the Illinois River over six million pounds are taken annually; and over seventeen million pounds are now marketed annually in the United States. At Port Clinton, near the western end of Lake Erie, great quantities are taken, and placed in large ponds until the market is ripe for them, when they are taken out and sold. Hundreds of tons are skinned, sent to the markets of Cincinnati, Louisville and St. Louis, and sold as buffalo fish." (C. H. Townsend.)

**Minnows.**—No common fishes of our country, it is safe to say, are so little understood, or so generally misunderstood, as those classed under the above name. To most persons a "Minnow" is a tiny young fish, from one to three inches in length, useful only as bait for bass, and other fishes.

The Minnow Family contains (says "American Food and Game Fishes") 200 genera, and more than 1,000 species, of which about 225 are found in our waters.

Many a Minnow only two inches in length is a fully-grown fish; but some species of Minnows attain a length of from one to two feet. One of the Pacific coast species (the Squaw-Fish) sometimes reaches a length of 4 feet.

For obvious reasons, it is impracticable to attempt to set forth even the leading species of this extensive Family, but it is proper to mention that to it belong the *Hornyhead*, of the Ohio and Mississippi valleys, the *Fallfish* of the northern Atlantic states, the *Common Chub* of the northeastern states, the *Columbia Chub* of the far northwestern states, and the *Utah Lake Chub* of Utah and northwestern Wyoming.

## CHAPTER LI

# THE ORDER OF HALF-GILLED FISHES


### HEMIBRANCHII

Because of the fact that a few very small fishes have less than their rightful number of gill-arches, and shoulder-girdles with one bone only instead of two, the Order of Half-Gilled Fishes has been created.

**The Sticklebacks** are very small fishes, only a few inches in length, and derive their name from the formidable dorsal spines that stand upon the back in front of the dorsal fin. We have Two-Spined, Four-Spined and Ten-Spined

observers say that the fish first brings a few stems and bits of vegetation, and by means of his gelatinous secretion practically ties them fast to the upright stalks, to use as a foundation.

The fish then proceeds to exude its secretion and dispose it in commingling rings, vertically, around a space sufficiently large for the female Stickleback to pass through. In a manner nothing short of marvellous, a hood-like nest is spun, of the fish's own secretion, which well retains its



Drawn by J. CARTER BEARD.

TWO-SPINED STICKLEBACK.

Sticklebacks, all three being found in brackish water along the Atlantic coast from Cape Ann to New Jersey.

All the Sticklebacks are celebrated for their nest-building habits. The abdomen of the male fish has been provided with a large gland which is "filled with a clear secretion which coagulates into threads" when it comes in contact with water. At first the fluid is colorless, but after contact with water it becomes whitish, and its many fibres hang together like strings of spaghetti. (Ryder.)

The entire work of nest-building is performed by the male Stickleback. It begins by selecting a bottom situation, in a gentle current, wherein the nest can be attached to two or more stems of growing vegetation, and anchored fast. Some

shape for some weeks. In this the female deposits her eggs, all the time jealously watched by the male, to prevent her from eating them! The male guards the eggs until they are hatched, and it is said that if the current flowing through the nest does not meet his views as to strength, the fish increases the volume of it by moving its pectoral fins to and fro. Sticklebacks are sometimes kept in aquaria in order that they may show their wonderful intelligence in nest-building.

**The Two-Spined Stickleback**<sup>1</sup> will serve as a type for the whole Order. It is only about 7 inches long, and has no commercial value. It is said to occur in quiet brackish waters along our coast, but is seldom brought into notice outside of aquaria.

<sup>1</sup> *Gas-ter-os'te-us a-cu-le-a'tus.*

## CHAPTER LII

# THE ORDER OF CATFISHES

### NEMATOGNATHI

Acquaintance with this numerous Family usually begins with the bullhead, which is merely a pygmy catfish.

Even when a lad in prairie-land, thirsting for open water and aquatics, and looking upon every mile of running water as an enchanted realm, the bullhead did not appeal to me as a genuine fish. Even when most eager to "quit, and go a-fishing, and call it half a day," we drew the line at that ill-shaped, skinny body, ugly head and

and hooks, represents the lowest depths of depravity in fishing with hook and line. It is even lower than fishing with four poles.

With a tenacity of purpose worthy of a better species, the bullhead ramifies throughout the muddiest rivers and creeks of the United States, and in the heat of midsummer holds on whence all but him have fled. He was built for mud bottoms and murky waters, and so long as the mud is thin enough to swim in, and deep enough to float him, he remains. When removed from his native element, the tenacity of life of this creature is astonishing. A bullhead will lie on the bank in midsummer sunshine, and breathe hot air for an hour without giving up.


The species of catfishes found in the United States number about thirty, but it is recorded that elsewhere there are about 970 more, representing in all about 100 genera. Of our series, all save four are confined to the eastern half of the United States.

**The Mississippi Catfish,<sup>1</sup> or Blue Cat,** of the Mississippi River and Gulf States is the giant of its genus. Even when alive and in good health, it is a very ugly fish,—heavy-paunched and

mud-colored. It looks like a fish modelled out of river-mud. I saw a specimen taken at Burlington, Iowa, which weighed 93 pounds, and have heard of others exceeding 100 pounds. Jordan and Evermann say the "record specimen weighed 150 pounds," and was caught at St. Louis; but the mischievous evenness of the figure casts doubt upon the reliability of the record.

Very naturally, the tons of edible flesh annually contributed by this fish to our national food supply are not wasted. Thousands of persons

<sup>1</sup> *Ictalurus furcatus*.



Drawn by J. CARTER BEARD.

COMMON BULLHEAD.

wide-gaping mouth with barbels that suggest dripping saliva. To me it was, and still is, a repulsive creature, and its only feature worthy of respect is the outfit of sharp and dangerous spines with which its dorsal and pectoral fins are furnished.

Excepting the big Mississippi catfish, it is the most unattractive fish inhabiting our fresh waters, and as an angler's proposition, it is worse than an eel. It is easily taken on a trot-line; and the "trot-line," set for all night across a stream, and hung with about twenty short lines

like the flesh of Catfish and bullheads, and in twelve months of 1899-1901, twenty-six states and six great lakes yielded twelve and a half million pounds, worth \$503,562. Illinois headed the list with 1,569,615 pounds, worth \$68,535.

**The Channel Catfish**<sup>1</sup> is the large Catfish of the North, and also the Mississippi valley, which so closely resembles the preceding species that it is at best very difficult—and sometimes impossible—to distinguish them. It is, however, much smaller than the blue cat, and instead of frequenting sluggish waters, it displays a decided preference for river channels, and clear water when it can be found. Naturally enough, its flesh is said to be of better flavor than the more sluggish, mud-inhabiting blue cat.

**The Common Bullhead**,<sup>2</sup> or **Horned Pout**, is merely a small, cheap catfish, whose room is better than his company. It ranges from the Atlantic well into the eastern edge of the great plains, and from the great lakes to the Gulf.

<sup>1</sup> *Ic-tu-lu'rus punc-ta'tus.*

<sup>2</sup> *Amei'u--rus neb-u-lo'sus.*

Much to the displeasure of many persons in California, three species of catfish have been introduced into many streams on the Pacific coast. Concerning them, the *San Francisco Evening Bulletin* has thus recorded the facts, and its views thereon:

“Then the fish commissioners made another unfortunate experiment, against the strongest protests that could be put forth. They introduced the hated and almost worthless Catfish to the waters of California. These fish, like the carp, have multiplied rapidly. It was reported, in answer to protests made at the time, that only a superior kind of Catfish would be introduced, against which there could be no objection. But they turned out to be the same old toughs that have occupied western rivers and bayous to the exclusion of better fish. These Catfish are voracious feeders on young trout and salmon. Their value is so low that very few seek them. The Chinese sell them occasionally, as they do carp; if they can find a customer. But most consumers turn away from these fish in disgust.”

## CHAPTER LIII

# THE ORDER OF FLAT-FISHES

### HETEROSOMATA

The flounders, halibuts, soles, plaice, and turbot make up the very desirable and important Order of Flat-Fishes. When in doubt about an English or continental breakfast, order a fried sole and you are safe; for so trustworthy is this fish that only the most bungling cook can spoil it. In England, the sole is almost a national institution, but on our side, its counterpart, the small flounder, is not so plentiful that it attains equal importance on the daily bill of fare.


The Order of Flat-Fishes, all the world over, is very large, "containing about fifty-five genera

in many instances it is so skilfully colored and mottled in imitation of the sandy bottom on which it lives, that when at rest on the floor of the ocean or aquarium the fish is almost invisible. On the other hand, the under side of the fish is white, or cream color, in order that to enemies below it, looking upward, it will match the light of the upper world.

As food fishes, the majority of the Flat-Fishes are very desirable. Their flesh is excellent, and their bones are few and far between. The flesh of the halibut is very white and firm, and whether fresh or smoked, it is highly palatable.

The common flounders are so well known they require no special notice. The species most common on our coast is the **Winter Flounder**,<sup>1</sup> which is caught in great numbers, and of all our Flat-Fishes is next in value to the great halibut. It is a small species, with an average weight of about 3 pounds, and a maximum of 5 pounds, or thereabouts. It has been extensively propagated by the United States Bureau of Fisheries.

**The Common Halibut**<sup>2</sup> is a cold-water fish of commanding importance. It is widely dispersed throughout both the North Atlantic, North Pacific and circumpolar waters, not only in shallow waters and the off-shore banks, but also on the sides of the sea-bottom slopes down to 1,500 feet. In the Atlantic, fishermen say the species stops at the latitude of the Delaware River. The fisheries along the west coast of Greenland are so important that regularly every year a number of schooners from Connecticut and Massachusetts go north, sometimes beyond the



THE COMMON HALIBUT.

and nearly five hundred species." Among its members, some of the halibuts attain great size. Almost any member of this Order is recognizable at one glance, by its broad, oval form, almost completely encircled by the fringe-like dorsal and anal fins, and the presence of both eyes on the upper side of the body. The body is so thin that "flat as a flounder" is a standard comparison wherever the English language is spoken.

The Flat-Fishes are good examples of protective coloring. All these fishes swim and rest with their bodies in a horizontal position. The upper surface, or back, is always darkest, and

<sup>1</sup> *Pseudo-pleuro-nectes americanus*.

<sup>2</sup> *Hippoglossus hippoglossus*.

Arctic Circle, and return loaded with Halibut to within three feet of their deck-beams.

On the Pacific coast, according to Dr. T. H. Bean, the Common Halibut ranges from the Farallone Islands, opposite San Francisco, to Bering Strait, its centre of abundance being found in the Gulf of Alaska, near Kodiak.

In point of size this fish is surpassed in our waters by no other good food fish, the 500-pound jewfishes being out of that class. A large Halibut is one which weighs 250 pounds or more. The largest of reliable record (at least from our waters) was observed by Captain Atwood, at Provincetown, Massachusetts. It weighed 401 pounds gross (we are thankful for that odd one

pound!) and 322 pounds dressed. Dr. G. Brown Goode states that a Halibut weighing 350 pounds is from 7 to 8 feet long, by nearly 4 feet wide.

The roe of a fish weighing nearly 200 pounds, which was caught at a depth of 200 fathoms, in water only 4° above freezing point, weighed 17 pounds, 2 ounces. A careful calculation made at the laboratory of the United States Bureau of Fisheries showed that the number of eggs in the mass was about 2,182,773.

The Halibut catch in twelve months of 1898-9 amounted to a total of nineteen million pounds, having a market value of \$797,222, and all credited to Maine, Massachusetts, Washington and Oregon.

CHAPTER LIV  
THE ORDER OF FOOT-FISHES


*PEDICULATI*

The strange creatures which form the group of so-called foot-fishes are introduced here, not in the expectation of close acquaintance with many of them, but rather that they may not remain absolute strangers to us. They live on the bottom of the sea, are not edible, and, being devoid of all value to mankind, they are safe from extermination. The most of them are also safe from close observation. Structurally, they stand next to the foot of the Subclass of Bony Fishes.

**The Angler, or Goose-Fish,**<sup>1</sup> is the typical representative of this Order. Among fishermen, it is sufficiently known that it has received twenty-

By taste and habit the Angler is in the same class as the human fish-hog who fishes with three poles at once. He lies on the bottom of the sea, where the muddy mottlings of his skin give him the appearance of mud and sand, opens his head widely, and props it open, for the free admission of any fish, crustacean, reptile or aquatic bird that chooses to enter.

Dr. Goode observes that the Goose-Fish derived that name from the swallowing of live geese, and that there is an authentic record of the capture of one which contained seven wild ducks.



THE ANGLER.

one English names, and in the languages of continental Europe about fifty more. (G. Brown Goode.)

It is the glutton of the sea, and its body is merely a purse-like attachment to a mouth that is fearful and wonderful to behold. It has a mouth and an appetite like an old-fashioned carpet-bag, and to it no living thing comes amiss. At present the body of this creature is painfully small for a mouth so ambitious and all-absorbing, but evolution is doing its perfect work, and eventually the maw of the Angler will be developed on the same scale as its mouth.

A fully-grown Angler is about four feet long, and its mouth is a little more than a foot wide. From snout to tail its lower jaw and the median line of the body are fringed with tiny barbels most cunningly calculated to lure unsuspecting fishes within seizing distance.

The weight of a large specimen is from 35 to 40 pounds. In our longitudes it is used only for bait, but Dr. Goode says that "in Italy it is much esteemed as an article of food." No doubt of it. In Naples, they eat stewed octopus; which I can testify is as tender and palatable as rubber hose stewed in brine, but not any more so.

<sup>1</sup> *Loph'i-us pis-ca-to'ri-us.*



## CHAPTER LV

# THE ORDER OF EELS

### APODES

Whenever a fish-like creature looks so much like a snake that it becomes necessary to inform people "it is not a snake, but a fish," then it is time to place it and all such creatures at the foot of the class of Bony Fishes. But for the good, hard bones in its skeleton, its descent to a position below the Order of Rays would be swift and sure.

As a real fish, an eel is little more than a caricature, and he who eats it must first skin it, just as the Dyaks of Borneo do their water-snakes before they roast them. It is the vulture of the waters, and prefers to feed upon things dead.

But, again are we reminded that there is no accounting for differences in taste. Both in Europe and America, they have been eaten ever since the days of the Cave-Dweller and Mound-Builder. And even to-day they are devoured, not with toleration, but with a degree of avidity worthy of better meat.

A German writer who catalogued the good points of the eel set forth prominently the fact that it is an excellent scavenger, and devours dead fish, crabs, and any fleshy prey, living or dead, that it can secure. Those who wish to pursue the subject of the food-habits of the eel to its logical conclusion can find it in a notable epic by Canon Ingoldsby, entitled "The Knight and the Lady."

Nevertheless, in times past, the eel has contributed a great store of edible flesh to the people of New England,—where some of the finest of fishes have always been abundant! There, eels are eaten—stewed, fried, pickled and salted. The flavor of an eel is not half bad, but its choice of food is decidedly objectionable. If eels are to be eaten by civilized people, then why draw the line at sharks, whose flesh is far superior to that of eels?

The United States Bureau of Fisheries has taken the eel quite seriously, and been at

considerable pains to introduce it in the upper Mississippi valley, the great lakes above Niagara Falls, and on the Pacific coast. And yet, Professor Baird recorded this very pertinent statement:

"It [the eel] is, however, a very undesirable inmate of rivers in which fish are taken by means of gill-nets, the destruction of shad and herring in the waters of the Susquehanna and others farther south being enormous. *It is not unfrequent that, when a gill-net is hauled up, the greater part of the catch consists simply of heads and back-bones, the remainder being devoured by myriads of eels in the short time the net is left out.*"



THE ELECTRIC EEL.

Is such a rapacious scavenger as this a species worthy of introduction in any new waters save those of an avowed enemy?

The maximum length of the **Common Eel**<sup>1</sup> is about four feet, but the average length is less than three feet. The female lays an enormous number of eggs,—estimated at ten millions,—preferably in salt water; but the young enter fresh water to develop, and ascend as far as they can go.

**The Electric Eel**<sup>2</sup> of South America is an Eel worth knowing. Having had with it some thrilling experiences, I can speak of it feelingly.

Once while canoeing for zoological specimens

<sup>1</sup> *An-gu-il'la vul-gar'is.*

<sup>2</sup> *Gym-no'tus e-lec'tri-cus.*

in the delta of the Orinoco, we entered a large creek flowing into the main stream from the south, and ascended it to the head of canoe navigation. It was a clear and beautiful stream, full of zoological wonders, and its Venezuelan name was *Canyó del Toro*, or Bull Creek. On the way up, our bow boatman checked the speed of the canoe, pointed to a straight, round stick of wood floating in the water about a foot below the surface, and said in an awe-struck tone, "*Tremblador! Grande!*"

The stick of wood was smooth, barkless, and of a bluish-gray color; and in reality it was a large specimen of the renowned and dreaded Electric Eel.

Acting on the collector's principle that the first specimen seen must be the first one taken, my companion poised his capybara spear, and drove it into the creature's body. The detachable head promptly came off, and the spearman held fast to the handle.

Instantly the big Eel became a storm centre of the first magnitude; and it writhed and struggled, and thrashed about until it struck against the handle of the spear. Mr. Jackson received such a shock that he cried out from the pain of it, and dropped the spear-handle, which floated on the water.

But not for long. My friend recovered his spear-handle, and drew the fiercely struggling Eel within striking distance of the canoe. When-

ever it struck the side of the boat, either with head or tail, we were thrilled by a shock. At last, two or three severe blows on the head, with the club used for killing capybaras, seemed to settle matters, and against the protests of Antonio, the creature was dragged aboard.

To all appearances, the Eel was dead; but a few moments later when Antonio chanced to touch it with his bare foot, at once he broke out in a torrent of anathemas upon all "*trembladors.*" As an experiment, I touched its head with the tip of my finger, and instantly received a shock so severe that my nerves tingled for an hour. A more vigorous application of the capybara club finally killed the creature, and its electric power died with it.

This specimen measured 6 feet 4 inches in length, and I believe that when delivered to advantage its electric power was sufficient to administer a severe shock to the largest elephant. Woe to the crocodile or shark which attempts to dine or sup at the expense of *Gym-notus electricus!* While on the *Canyo del Toro* we saw about ten specimens, always of the same floating-stick appearance, and captured four.

**The Lamper "Eel,"** as the Lamprey is very frequently called, is not a true eel of any sort, and it will be found in its proper place, immediately following the fishes. It is so low in the zoological scale that for it and kindred forms a separate Class has been provided.

## CHAPTER LVI

# THE ORDER OF PIPE-FISHES AND SEA-HORSES

### LOPHOBANCHI

At the foot of the Subclass of Bony Fishes, stand certain small creatures, each of which is so fantastic in form that it requires to be introduced with the solemn assurance, "This is a fish!" At first glance, any one wholly unacquainted with them might from their hard external shells be inclined to regard them as particularly odd crustaceans; but the presence of tiny fins without, and skeletons more or less bony within, place them fairly within the confines of the Bony Fishes.

**The Great Pipe-Fish<sup>1</sup>** is a long, slender stalk of jointed bone, with queer little fins very far apart, and a head that terminates in a long, hollow tube. But for this very tough and persistent bony armor, other small fishes would devour the Pipe-Fishes, bit by bit, as children bite off sticks of candy. Its armor is so stiff, however, that the wearer moves slowly and with difficulty, and the prey usually sought by this fish is found very small and weak, hiding in the branches of sea-weed, coral clusters, sponges, and the sea-grasses generally. It was for insertion into such hunting-grounds as these that the long, tubular snout of this fish has been developed.

The Pipe-Fishes swim in a half vertical position, as if literally leading up to the introduction of the next species, which swims bolt upright in the water, and fairly caps the climax in fishes. All the Pipe-Fishes are small creatures. Our largest species is found on the Pacific coast, and "reaches a length of 18 inches." (Jordan and Evermann.) There exists in North American waters about thirty species.

**The Sea-Horse<sup>2</sup>** bears not the faintest resem-

<sup>1</sup> *Syng-na'thus a'cus.*

<sup>2</sup> *Hip-po-cam-pus hep-tag'o-nus.*

blance to a typical fish, and is the strangest-looking creature of the whole fish world. It looks like a Chinese dragon, reduced about a thousand diameters. Its minute pectoral fins are so inconspicuous they are at first quite unnoticed, and the fan-shaped dorsal fin seems when in action like a stationary fan with which the outlandish creature frequently tries to fan itself.

At all times the Sea-Horse swims in a perpendicular attitude, and with its prehensile tail it



1. GREAT PIPE-FISH.  
2. THE SEA-HORSE.


holds itself stationary by grasping any inanimate object that either grows upon the bottom or floats in the water. Like the pipe-fish, it is completely encased in a strong suit of bony plate-armor. The average aquarium Sea-Horse is seldom more than 4 inches in length, but the Gigantic species (*H. ingens*) of the Pacific coast "reaches a length of nearly a foot." (J. & E.) The smallest species, found abundantly about Pensacola, is only 2 inches long.

## CHAPTER LVII

# THE ORDER OF THE DOGFISH

### HALECOMORPHI

To naturalists, the **Dogfish**<sup>1</sup> is a creature of much interest. Like the prong-horned antelope, it is so unique and peculiar that it has been necessary to create for it a grand division of classification which it occupies all alone. The antelope is only a Family, but this fish is a whole Order. Its other English names are *Mudfish*, *Boufin*, *Grindle* and *Lawyer*; and since Linnaeus christened it *Amia calva*, in 1766, eleven other naturalists have given it eleven other names in Latin.



THE DOGFISH.

The Dogfish has an air-bladder that is divided into cells, and is a half-developed lung. At intervals it ascends to the surface of the water, gulps down a mouthful of air, just as a turtle does, and descends again. If hindered from rising when the time comes to take in a supply of fresh air, the fish struggles violently, like a mammal about to be drowned; but it can expel air while below the surface. This character indicates that lungs were first developed in fishes, from modifications of their air-bladders. Other characters establish a distinct relationship with the gar fishes, and place it in the Subclass *Ganoidea*. The dorsal fin is low, of uniform height throughout, and is about one-half as long as the entire fish.

By its general anatomy, this fish appears to stand midway between the true lung-fishes and

<sup>1</sup> *Am-i'a cal'va*.

the gar pikes. It is of scientific interest, only, for, save to the negroes of the South, its flesh is quite unpalatable, and valueless as food. It is an inhabitant of sluggish fresh waters, attains a length of 2 feet, and 12 pounds weight. It is found in the great lakes, the Mississippi valley generally, and in a few fresh-water streams on the southern Atlantic coast.

The individuality of the Dogfish is very positive and interesting. Among the small fry of other fishes its voracious appetite renders it very destructive to species of more value than itself. Mr. Charles Hallock, who knows it well, has thus set forth the salient points of its moral character:

“They take frogs, minnows, and sometimes the spoon. Their habitat is deep water, where they drive everything before them. They are

very voracious and savage. Their teeth are so sharp and their jaws so strong they have been known to bite a two-pound fish clean in two the very first snap. They are as tenacious of life as an eel. The young, when about six inches long, make a famous bait for pickerel and pike. To use it, run the hook into the mouth right up through the centre of the head, through the brains, cast a hundred times, catch several fish, and at the end of three to six hours he will kick like a mule.”

“Put a hundred in a rain-barrel, and you can keep them all summer without change of water. For the aquarium, the young have no equal, and on account of the spot in the tail they are quite attractive; but nothing else than snails can live in the tank. He will kill a lizard or any other living thing the instant it touches the water.”

## CHAPTER LVIII

# THE ORDER OF GAR FISHES, OR GANOIDS

### GINGLYMODI

To the scientific student, the **Gar Pike** of the middle eastern states, and the big **Alligator Gar** of the Gulf states are two of the most interesting fishes of our whole finny fauna. They are the living representatives of a wonderful lot of dead-and-gone species which many thousand years ago laid the foundations of the fish world. By means of the impregnable bony armor with which Nature wisely provided them, they have been able to withstand the attacks of the enemies that otherwise would have exterminated them.

The simplest, and therefore the earliest, forms of fishes are some of the Gan'oids,—as the armored fishes are called,—whose remains now exist only in the rocks of the Devonian age, far down toward the strata which were formed before life was. The first of these fishes—and they were well-nigh the first of all fishes—had their heads completely encased in solid bone, their eyes were placed in the tops of their heads, and they must have lived upon the bottom of the sea. And who shall say how many years have passed since the days when their dead bodies sank in the mud along the shores they frequented? To-day they are found high up in the rocky cliffs of Devonshire, England.

It must be remembered, however, that the armored fishes were not the only ones which existed in those early days. The same rocks have yielded to science the remains of lung-fishes, sharks, and sturgeons; but the so-called "bony fishes" of to-day were undoubtedly of later development than the foregoing.


Our two Gar Fishes are therefore to be regarded as living relics of the Devonian age, or "Age of Fishes." There are others; but for an introduction to them, as well as the fossil forms, the reader is referred to Le Conte's "Geology."

**The Long-Nosed Gar Pike**<sup>1</sup> is the species

<sup>1</sup> *Lep-i-dos'te-us os'se-us*.

which is nearest at hand, and most accessible to teachers and students. It is found in the great lakes, and in large streams generally from New Jersey to Mexico, and northward in the Mississippi valley to Minnesota. It is frequently called the *Bill-Fish* and the *Gar*. It is said to be destructive to the young of other fishes, but Dr. Goode declares that fish remains are "rarely found in its stomach." Its flesh is unfit for food, and, except to educators, the fish is valueless. It is said to attain a maximum length of from 5 to 6 feet, but specimens exceeding 3 feet are very rare, and the majority are certainly under that length.

The armor of this fish is more perfect than any



LONG-NOSED GAR PIKE.

plate armor that man could make for it. It consists of diagonal whorls of solid and highly polished plates of bone, each divided into scale-like sections, and so hinged together that while fully protected the fish has abundant freedom of movement. The dried skin of a Gar Pike is as hard and unyielding as a cylinder of sheet iron.

In about the same waters as the preceding species, and very much like it, lives the **Short-Nosed Gar Pike** (*Lepidos'teus platys'tomus*).

**The Alligator Gar**<sup>2</sup> is a giant in comparison with both the above species, sometimes attaining 6 feet in length. It is essentially a fish of the South, and inhabits the large streams—and also many small ones—of all the Gulf states, Mexico and Cuba. It is readily recognized by its short and broad snout, which is strongly suggestive of the head of an alligator.

As an instance of the manner in which fishes

<sup>2</sup> *Lep-i-dos'te-us spat'u-la*.

sometimes perish through natural causes, and become fossil, Mr. Frederic S. Webster tells the story of a death pool near the Rio Grande. While collecting birds near Brownsville, Texas, he discovered a large pool which had been filled by the overflow of the river, but afterward entirely cut off by the receding of the flood waters. A muddy pool seventy-five feet long by twenty-five feet wide was crowded full of Alligator Gars, living, dying and dead, varying in size from two feet to six. Mr. Webster estimated that that tiny area of water and mud, no larger than a fair-sized ball-room, contained between 700 and 800 fishes, all doomed to speedy annihilation by the evapora-

tion of the remaining water. When he discharged his shot-gun into the mass, pandemonium ensued. The pool became a seething mass of frantic life, and the wild rushing to and fro of the large fishes actually threw smaller ones into the air.

A million years from now, the few men of science who have not yet perished from cold may discover on the summit of a lofty, rock-ribbed mesa at the edge of a great desert, a marvellous deposit of fossil Alligator Gars, and wonder how so many fishes chose to die in the same spot. But only the rocks will then be able to tell the story of Mr. Webster's Pool, and the world will be too cold to care for it.

## CHAPTER LIX

# THE ORDER OF STURGEONS

### GLANIOSTOMI


A sturgeon is a big, shark-like, wedge-headed fish, which looks as if Nature had once decided to cover it with a bullet-proof suit of bony armor, but, after setting three or four rows of plates on each side, had grown weary of the task, and abandoned it. Had the plan been wrought out to a finish, it would now be necessary to skin every sturgeon with an axe.

The mouth of a sturgeon is situated underneath the head, and is provided with long, sucker-

sturgeons are distributed at intervals throughout the northern portion of the north temperate zone, across America, Europe and Asia. The American species are but four in number.

**The Lake Sturgeon<sup>1</sup>** is from 5 to 6 feet in average length, weighs from 30 to 40 pounds, and inhabits the great lakes and adjacent connecting waters of good depth.

**The Short-Nosed Sturgeon<sup>2</sup>** is a salt-water species, found along our Atlantic and Gulf coasts,



LAKE STURGEON.

like lips, for taking food off the bottom. The principal food of sturgeons is small, thin-shelled mollusks, and other fishes are not eaten save on occasions so rare they are not worthy of note.

From the coast of California to the Caspian Sea, wherever they are found, sturgeons are fishes of desirability, and of commercial value in direct proportion to their size. Their smoked flesh is by many considered equal in flavor to halibut, and "caviar" is only the society name of air-tight sturgeon eggs. The twenty living species of

from Cape Cod to Texas. This is a small species, only about 2 feet in length, and is of no importance.

**The Common Sturgeon<sup>3</sup>** of our Atlantic coast is the largest and most valuable member of this Order in American waters. It attains a length of 10 feet, and 500 pounds in weight, and to-day at Wilmington, Delaware, its centre of abundance, a large specimen represents about \$75 worth of commercial value. The most valuable part is the

<sup>1</sup> *Ac-i-pen'ser ru-bi-cun'dus*.

<sup>2</sup> *A. bre-vi-ros'tris*.

<sup>3</sup> *A. stu'ri-o*.

roe, a cask of which, weighing 130 pounds, is worth \$110.

**The White Sturgeon**<sup>1</sup> inhabits the waters of the Pacific from southern California to Alaska, and the records show it to be a giant among food fishes. Jordan and Evermann quote it up to 13 feet in length, and weighing 1,000 pounds; but the weight of any animal, dead or alive, which ends with two ciphers is certain to be a weight of Estimate, and not of Fairbanks. Strangely enough numerous specimens of this Sturgeon

have been taken in Idaho, in the Snake River, weighing from 100 to 650 pounds. "An example 11 feet 2 inches long measured 2 feet across the head." (Jordan and Evermann.)

The latest reports on the Sturgeon industry generally are for 12 months during 1897 and 1898. During that period, 17 states participated in a catch which amounted to 5,726,830 pounds, which sold for \$321,036. The catch in Oregon was nearly two million pounds, that of New Jersey 868,326, and Virginia next.


<sup>1</sup> *Ac-i-pen'ser Atrans-mon-tan'us.*



## CHAPTER LX

# THE ORDER OF THE PADDLE-FISH

### SELACHOSTOMI



THE PADDLE-FISH.

To some persons, the big **Paddle-Fish**,<sup>1</sup> or **Shovel-Nosed "Sturgeon,"** as it is more commonly called, is one of the wonders of fresh water. Here we find a case of what naturalists call "specialization" which has gone to an astonishing extreme. This is a scaleless fish, with a body very much like a shark, and a half-cartilaginous, shark-like skeleton. It has a low-browed, armor-plated head that runs forward into a broad, thin paddle of bone, one-third the length of the entire fish.

Beyond doubt, this remarkable implement is used in turning up the mud and gravel of the bottoms of the streams in which the owner lives, in searching for food. It is unfortunate that we never can see it in action, and still more so that this fish has not yet been kept successfully in aquaria. Mr. Charles H. Townsend says that in captivity they always injure their paddles against the sides of their tanks, and do not live longer than two or three weeks.


<sup>1</sup> *Po-ly'o-don spath'u-la.*

In "American Food and Game Fishes," Drs. Jordan and Evermann give a number of size records of this fish which will be a surprise to many persons who, like the writer, have seen and handled only medium-sized specimens. The figures given show length in inches, and weight in pounds.

	In. Lbs.
Lake Manitou, Ind., heaviest on record. <sup>2</sup>	163
Lake Tippecanoe, Ind. (W. C. Harris), length .....	74 150
Chautauqua Lake, N. Y., length .....	74 25½
St. Louis (Dr. Engelman), length .....	70 79
White River, S. Dakota (J. and E.), length .....	53 18

The latest record is interesting as showing the light weight of what was a long, but very slender specimen. Judging from all available evidence,

<sup>2</sup> This fish and the one next noted was 4 feet in girth.



UNDER VIEW OF THE PADDLE-FISH.

and personal observations, I should place the average length of the Paddle-Fish at 45 inches, and weight 25 to 30 pounds.

The U. S. Bureau of Fisheries' records show that this fish is now coming into use as food, and is finding a ready sale in the markets of the region it inhabits. In some places its flesh is smoked and sold as sturgeon. Its eggs, which are very numerous, and greenish-black in color, make excellent caviar, and are being so utilized at Louisville, Kentucky, and along the Mississippi, in Mississippi and Tennessee.

In 1899, sixteen states participated in the catching of Paddle-Fish, Mississippi leading with 981,080 pounds, and followed by Arkansas, Tennessee, Illinois and Missouri, in the order named. The total catch was 2,543,950 pounds, valued at \$82,343.

In a limited sense the Paddle-Fish inhabits the Mississippi valley, from Louisiana to Minnesota, the Ohio, and the Missouri to South Dakota, which is a wide range for a fish so peculiarly formed.


CHAPTER LXI  
THE ORDER OF THE CHIMERAS

CHIMAEROIDEI

The Chimeras are introduced for the purpose of making our series of fish Orders reasonably complete, and not because of anticipated personal acquaintance with them. For fifteen or twenty years one may live on the Atlantic coast, frequent its fish markets, and fish occasionally at first hand, without once seeing either a live Chimera, or one freshly caught. They inhabit blue water

**The Spotted Chimera,**<sup>2</sup> figured herewith, is said to be extremely abundant just off the borders of the submerged plateau that extends all along the northwest coast of the United States. It was frequently taken in the dredge hauls made by the steamship *Albatross*, the majority of the specimens being under 2 feet in length.

Like all the members of this Order,—the total



SPOTTED CHIMERA.

only, have no commercial value save as scientific specimens, and in our Atlantic waters are rarely caught elsewhere than on the off-shore fishing banks of New England.

As a natural result of these conditions, the shark-like chimaeroids are the least known of all the fishes that inhabit our shore waters. Indeed, there are several species of deep-sea fishes that are much more common in fish collections than they appear to be elsewhere. One species, however, of the Pacific coast, has been studied by Dr. Bashford Dean, and it will be set forth on the strength of his description.<sup>1</sup>

<sup>1</sup>“Fishes, Living and Fossil.” Columbia University Biological Series, page 100.

number of which is very small,—this species resembles a big-eyed shark with a cutlass-fish tail.

The head is blunt and very thick, and from it the body gradually tapers down to the whip-like tail. The skin is smooth, and the paired fins are shark-like.

The front dorsal fin is provided with anterior spine-folds, like a fan, and may be depressed into a sheath in the body-wall.

The sense organs are similar to those of sharks, and the visceral parts also are shark-like. The skeleton is cartilaginous, and the vertebral axis is notochordal. Of the embryology and life history of the Chimeras generally, practically nothing is known.

<sup>2</sup>*Chi-me'ra col'le-i.*

## CHAPTER LXII

# THE ORDER OF SHARKS

### SQUALI

We have now reached the subclass of Cartilaginous Fishes.

And what is a "car-ti-lag'i-nous fish?"

Cartilage is a bloodless tissue, commonly called *gristle*, flexible but not elastic, quite colorless, of the consistency of cheese-rind, and of use in the

is a tree-like development of cartilage designed to bridge together the outer ends of the principal ribs, protect with some firmness the vital organs within, and yet permit the rise and fall of the chest in breathing.


The Cartilaginous Fishes, embracing the sharks, rays, skates and intermediate forms, are those whose skeletons are largely composed of plates and stems of cartilage, or gristle and but little bone. Instead of bony rays, the fins of these creatures are supported by cartilaginous rays so closely joined together that they form plate-like structures.

#### General Characters of Sharks.—

With few exceptions, sharks have externally the same general form as the typical fishes. Instead of broad, flat scales that overlap each other like shingles, their scales are very minute, horny, sharp-pointed and closely packed together. When the skin of a shark is stroked from head to tail, it feels like a hair-cloth sofa, but when stroked the other way, it is like the sharpest sand-paper. For centuries shark-skin has been used for smoothing and polishing wood and other substances; and when prepared for that use it is called "shagreen."

Instead of one very large gill-opening, as in typical fishes, a shark has usually five small slits in the skin behind the gills, which are capable of being tightly closed. In nearly all species the mouth is situated underneath the head, and often it is of

enormous proportions. The jaws are composed of cartilage, the teeth are usually triangular, and set along the edge of the jaw, in rows, crosswise with the edge of the mouth. Behind each active



1. MACKEREL SHARK, WITH REMORA ATTACHED.  
2. HAMMER-HEAD SHARK.

anatomy of animals for sustaining or connecting softer parts. The external ear of man consists chiefly of a convoluted wing of cartilage covered with skin. The so-called "breast-bone" of man

and visible tooth there is a line of reserves, from three to five in number, always growing outward and crowding to the front, so that as soon as a tooth in the line of battle becomes much worn, or in any way weakened or broken, it is crowded off the jaw, and a new tooth is thrust forward into its place.

Many sharks bring forth their young alive; but others (the majority, perhaps) lay eggs. Some of the egg cases are of remarkable form. Some of them are rectangular, flattened, and provided at each corner with a long, threadlike tendril with which to attach to any fixed object.

Sharks very rarely exhibit color patterns, or bright colors of any kind. As befits pirates and freebooters, they are mostly ashy gray, or drab,—the most inconspicuous colors at sea, both for sharks and men-of-war. The small Tiger Shark, of Ceylon (*Stegastoma tigrinum*), is one of the few sharks of variegated colors, and its handsome pattern of yellow and black is a welcome variation.

Only a few of the whole 150 species of sharks can rightly be classed as “man-eaters.” A typical “man-eating shark” is one which is very large, exceedingly voracious, practically devoid of fear of mankind, and so aggressive that it will attack a swimmer at the surface of the water, and devour him regardless of his resistance. The standard prey for sharks consists of small fishes, squid, jelly-fishes, crabs, lobsters and other non-combatants.

Occasionally, however, the big Tiger Shark<sup>1</sup> of the Atlantic chooses a victim in his own class as a fighter. Dr. Goode notes the capture, by Captain Atwood at Provincetown, Mass., of a specimen which contained “nearly a whole full-grown sword-fish;” and “ten or twelve wounds in the skin of the shark gave evidence of the contest that must have occurred.”

The “man-eater shark” is not a myth, for that name is applied to the great white shark, a species which ranges from our Atlantic coast to Australia, and on to California. In the tropics it attains a length of 30 feet. With us this creature is rated

<sup>1</sup> *Gal-e-oc'er-do ti-gru'nus*.

as “exceedingly rare,” and judging from Dr. Goode's notes, not more than a dozen specimens are caught and identified in a century. The only loss of life from it on our coast, so far as recorded, occurred in 1830.

It is indeed fortunate, and merciful to mankind, that sharks generally are harmless to man. Were they otherwise, the terrors of the sea would be greatly increased.

**The Mackerel Shark**<sup>2</sup> is a fair type of the sharks of the world. It is common along both coasts of the United States, and the length of fully-grown specimens is between 9 and 10 feet.

**The Hammer-Head Shark**<sup>3</sup> is a genuine curiosity. With no intermediate forms leading up to this strange departure, the head of this creature suddenly thrusts out on each side a great shelf of cartilage and skin, in the outermost edge of which the eye is situated! It is like a flat-headed shark with a seven-inch board twenty inches long placed squarely across its forehead. This species is found in the seas of the tropics and subtropics, practically around the world. Once when the writer was approaching the coast of Barbadoes, on a sailing vessel, a large Hammer-Head swam for fifteen minutes close to the bow of the ship, and quite near the surface. In the Havana market I once obtained a specimen nearly ten feet long. This species brings forth its young alive, and occasionally specimens are taken as far north as New Jersey.

**Sizes of Sharks.**—The majority of the species of sharks are under 8 feet in length, and a few are as small as 2 feet, when adult. The largest species are the following:

The <b>Basking Shark</b> ( <i>Rhinodon typicus</i> ) . . . . .	45 ft.
The <b>Bone Shark</b> ( <i>Cetorhinus maximus</i> ) 36 “	
The <b>Man-Eater or White Shark</b> ( <i>Carcharodon carcharias</i> ) . . . . .	30 “
The <b>Great Tiger Shark</b> ( <i>Galeocerdo tigrinus</i> ) . . . . .	30 “
The <b>Hammer-Head</b> ( <i>Sphyrna zygaena</i> ) . . . . .	15 “
The <b>Blue Shark</b> ( <i>Carcharias caeruleus</i> ) . . . . .	15 “
The <b>Thresher Shark</b> ( <i>Alopias vulpes</i> ) . . . . .	15 “
The <b>Mackerel Shark</b> ( <i>Lamna cornubica</i> ) 10 “	

<sup>2</sup> *Lam'na cor-nu'bi-ca*.

<sup>3</sup> *Sphyr'na zy-gae'na*.

CHAPTER LXIII  
THE ORDER OF RAYS AND SKATES


RAIAE

The rays and skates are merely flat-bottomed, side-wheel sharks, built to navigate very shallow waters. From the typical shark down to the broadest and flattest ray, the change of form is shown by a beautifully complete series of living links, several of which it has been my privilege to handle and dissect fresh from their home waters.

Of these connecting links, the most interesting is the rare and wonderful **Shark-Ray**,<sup>1</sup> of the

and thinness. But the long, fleshy body and triangular head still proclaim very unmistakably the line of relationship with the sharks. Several species representing this intermediate type are found in our waters, but they are not common, and the real home of the genus is in the tropics and subtropics.

**The Sawfish**,<sup>3</sup> of the Florida coast, and many portions of the tropics farther south, is celebrated



SHARK-RAY.

Indian Ocean, a fine specimen of which was caught for me in the shallow waters between Ceylon and India. It is as nearly as possible half shark and half Ray, and is shown in the accompanying illustration.

Between this and the typical ray stands the **Beaked Ray**,<sup>2</sup> much flatter than the preceding, and with the pectoral fins taking on ray-like spread

among fishes because of the very long, flat beak of bone which projects forward from its snout, armed on both sides with formidable teeth. The length of this saw is more than one-third the length of the head, body and tail. It is, we may safely assert, strictly a weapon of defence, not offence; for unless it is used as a shovel in searching for mollusks and other food on the bottom of the sea, it is useless in the search for food.

When the Sawfish is threatened with attack, however, it defends itself by quickly curving side-

<sup>1</sup> *Rhamphobatis ancylostomus*.

<sup>2</sup> Of the genus *Rhinobatis*. The species sometimes seen on the coast of Florida is *R. lentiginosus*.

<sup>3</sup> *Pristis pectinatus*.


wise, thereby giving a sweeping sidewise stroke with its saw, and swiftly repeating it in the opposite direction. On a Sawfish fourteen feet in length, the saw is about four and a half feet long, and the teeth project about one and a half inches from the bone.

This creature is an intermediate form between the sharks and the typical rays, and in reality it is a shark-ray. Its eyes are a-top of its head, its mouth is underneath, its body in front of the dorsal fin is quite well flattened, and its pectoral fins have "ray" written all over them. The maximum

tends outward to the very tips of the wing-like fins. Upon this is laid a thin layer of flesh, and over all is spread the rough and tough skin. The tail is like a long, stiff whip, with a many-barbed bone stiletto midway,—a very dangerous weapon to be so carelessly exposed.

To a taxidermist, the mounting of a large ray is about the most calamitous task he can possibly encounter. The trouble lies in the perpetual shrinking after mounting.

**The Sting Ray,**<sup>1</sup> or, by corruption, "Stingaree," is one of the greatest pests of the eastern



THE SAWFISH.

length attained by it is said to be 15 feet. Because of the long, flat beak of this creature, it has become associated in many minds with the swordfish, but structurally the two are as far apart as a deer and a bear.

Notwithstanding the fact that there exists a group called the Order of Flat-Fishes (halibuts, flounders and soles), the rays are by far the flattest of all fishes. For example, the Spotted Ray of Ceylon is about 5 feet across, 5 inches thick at the centre of the body, and at the edges its great wings flatten out into thin air. From the body, which really is quite small, and centrally located, a thin sheet of cartilage, consisting of a great number of very long, jointed rays firmly joined together, ex-

coast of the American continent. From Cape Cod to the Orinoco, and I know not how much farther beyond, this vindictive and cruel fish lies, assassin-like, half buried in the sand along shore, ready and anxious to drive its spine into any naked foot that comes within striking distance. The upper surface of the animal closely resembles the loose sand in which it hides, and the spine makes a ragged and ugly wound. The spine is long, dagger-like, and barbed like an arrow all along both edges, so that the withdrawal of it from a wound is very painful. On the lower Orinoco I saw a strong man who was then in the seventh week of disability from the stroke of a

<sup>1</sup> *Trygon sa-bi'na*.

Sting Ray in his foot; and in the Malay Peninsula I treated a Malay fisherman whose hand had been completely transfixed by the spine of a huge ray.



STING RAY.

Fortunately, this abominable creature is averse to cold, or even moderately cool waters, and is rarely encountered even as far north as Florida. On our coast, one may bathe for a lifetime without seeing even one; and in all waters they carefully avoid crowds of bathers.

The gigantic creature known as the **Devil-Fish**<sup>1</sup>


<sup>1</sup> *Man'ta bi-ros'tris*.

is the largest of all rays, and to many persons, even the most truthful accounts of some of its doings will seem beyond belief. To begin with, its maximum size of *twenty feet* across its "wings" is almost incredible. The towing of a good-sized fishing smack by a harpooned Devil-Fish, going for miles at race-horse speed, is another wonder of the deep.

Many years ago, the planters on the coast of South Carolina found royal sport in harpooning this monster, and conquering it. In a volume entitled "Carolina Sports," the Hon. William Elliott has drawn this picture: "Imagine a monster many feet across the back, having powerful flaps or wings with which he drives himself furiously through the water, or vaults high in the air, his horns projecting several feet beyond his mouth!"

If a Devil-Fish could leap out of water,—which there is good reason to suppose that it could do,—it would look as Mr. Beard has represented it in his illustration.

So far as can be learned, large examples of this creature are now rarely observed, and still more rarely captured. Its centre of abundance now appears to be off the Gulf coast of Florida; but it is also found on the coast of southern California.



THE DEVIL-FISH.



## CHAPTER LXIV

### THE LOWEST CLASSES OF VERTEBRATES

There are a few creatures which, by reason of their internal skeletons and jointed back-bones, are justly entitled to stand with the vertebrates, but yet are lower in the scale than the lowest fishes. For these it has been necessary to create two grand divisions of the first rank; and they stand as two small and very low Classes. It is because of their very low position in the zoological scale of vertebrates that it becomes important to know them.

#### THE LAMPREYS.

##### *Class Marsipobranchii.*

A **Lamprey** is an aquatic creature which bears so strong a resemblance to an eel that for a long period all lampreys were regarded as true eels. Even to-day, the most important of our species is, by unscientific persons, almost universally called the "**Lamper Eel.**" In view of the general external resemblance of these creatures to eels of similar size, it is not strange that their true character remained for a long period quite unknown. As a matter of fact, these creatures forcibly illustrate the unwisdom in animal classification of attaching too much importance to external characters.

*The lampreys are the lowest and last creatures that have the spinal cord expanded at its upper end into a brain, and encased in a skull.* But the skull is imperfectly developed, and without jaws; there is no shoulder girdle, no pelvis, no limbs, no ribs, and no paired fins. There is a single median nostril, the gills are purse-shaped, the skin is naked like that of an eel, and the skeleton is cartilaginous. The gills are in the form of a fixed sac, the gill openings consist of a row of tiny round holes along the side of the body, and the mouth is specially formed for suction.

It is evident from the foregoing characters that the lampreys are creatures of very simple form, lacking almost all the evidences of special development which characterize the higher fishes. Ex-

ternally, their very modest median fins are the only visible signs that they are not marine worms.

The **Sea Lamprey**<sup>1</sup> is the best and most available example of the Class *Mar-si-po-branch'ii*. "The mouth is completely circular, and forms a great and powerful sucker, surrounded by fleshy lips that are supported on a framework of cartilage and studded with tentacles. This mouth is covered over its entire interior surface with strong teeth arranged in concentric circles. A large double tooth, situated above the aperture of the mouth, indicates the situation of the upper jaw, and seven or eight great teeth represent the lower jaw. Even the tongue carries three large teeth, deeply serrated on their edge."<sup>2</sup>

With a mouth specially formed and savagely equipped for suction, it is no surprise to find that this creature is a blood-sucking parasite, preying upon other forms of marine life. It is often found attached to shad, sturgeon, sharks, cod, halibut and mackerel. It fastens to its victim beneath the pectoral fins, tears at its flesh with its rasping circles of teeth, and sucks its blood "until the flesh becomes as white as paper." Beyond doubt, these creatures destroy a very considerable number of valuable food fishes. Fishermen charge to the account of the Lamprey the raw spots and sores frequently found upon the bodies of sturgeons.

Formerly the Lamprey was greatly esteemed by the people of Massachusetts as a food "fish." In the Merrimac River it was captured in great numbers, and salted down for winter use. While this industry, and its object, have both greatly decreased, in some portions of Connecticut the Lamprey is yet taken, as often as it can be found, and thankfully consumed. The species specially mentioned varies in length from two to three feet, but the *Brook Lamprey*, and all the

<sup>1</sup> *Pet-ro-my'zon ma-ri'nus.*

<sup>2</sup> "Fishery Industries of the United States," Part I, p. 677.

fresh-water species are much smaller. Fortunately, none of the fresh-water species are so injurious to fishes as the Sea Lamprey.

According to Jordan and Evermann's "Fishes of North and Middle America," there are in American waters ten species of lampreys, and two of their very near, but still lower, relatives, the *Hag-Fishes*. They are scattered at intervals from Alaska to New England, in brooks, rivers, lakes, estuaries and various other bodies of shallow water. They are most accessible in fresh water, on a stony or gravelly bottom; and whenever in such a situation you find an eel-like creature holding fast to a stone by the suction of a big flat mouth on the end of its head, know of a surety that it is a Lamprey.

### THE LANCELETS.

#### *Class Leptocardii.*

The long and interesting chain of Vertebrates ends in a very weak and insignificant link. The great work entitled "Fishery Industries of the United States" dismisses this creature with only

And thus ends our bird's-eye view of the Vertebrates, setting forth the prominent types and examples which every intelligent American should know. *It is here, and here only, that "specialization" may properly begin!* Behind lie the Mammals, Birds, Reptiles and Fishes; beyond lie the mighty hosts of the Invertebrates,—Crustaceans, Insects, Mollusks and others. In any one of these grand divisions of life, the special student may wander for a lifetime in a wonderland of his own, and to the last find each day filled with new light and new joys in the unending revelations of Nature.

two and a half lines, and leaves three-fourths of the page blank.

And truly, the **Lancelet**, or **Amphioxus**,<sup>1</sup> is not a creature calculated to arouse enthusiasm. Its skeleton is composed of membranes and cartilages. It has no brain, nor even a skull in which to develop one. It is neither eel-like nor worm-like, but as its name implies, it is shaped like the head of a lance. The middle line of the body is provided with weak and indifferent fins. There is no proboscis, and the mouth is slit-like, and fringed with hair-like filaments. All the above characters, and many others of a purely technical nature, are set forth in "The Fishes of North and Middle America," where eight species are recognized.

These small, naked, colorless and translucent creatures are found "embedded in the sand in the shallow waters of warm coasts throughout the world." They are of special interest only because they are the lowest of the Vertebrates, and on the whole they constitute a very ignominious ending for the highest grand division of Nature.

<sup>1</sup>The West Indian Lancelet (*Brach-i-os'to-ma-car-i-bae'um*), is found from Beaufort, N. C., to the mouth of the La Plata.

THE END.

# INDEX

- Aard-Varks, 162  
 Acanthopteri, Order, 382  
 Accipiter atricapillus, 231  
     " cooperii, 230  
     " velox, 230  
 Acipenser brevirostris, 427  
     " rubicundus, 427  
     " sturio, 427  
     " transmontanus, 428  
 Aetodromas minutilla, 253  
 "Adder," Blowing or Spreading, 346  
 Adams, "Grizzly," 34  
 Agelaius phoeniceus, 200  
 Agriculture, Department of (*see* Biological Survey), 225, 240  
 Aix sponsa, 272  
 Ajaia ajaia, 264  
 Alaska Commercial Company, 48  
 Alaudidae, Family, 179, 206  
 Albacore, Great, 389  
 Albatross, Black-Footed, 291, 294  
 Albatross, Nesting on Laysan Island, 294  
 Albatross, Short-Tailed, 294  
     " Wandering, 291  
*Albatross*, S. S., 293, 431  
 Alca torda, 305  
 Alcedinidae, Family, 215  
 Alces americanus, 99, 139, 143  
 Alcidae, Family, 300, 302  
 Allen, Dr. J. A., 134  
 Alligator, Chinese, 317, 322  
     " Mississippi, 317, 322  
 "Alligator Joe," 312  
 Alopias vulpes, 433  
 Alosa sapidissima, 407  
 Amazon, "Naturalist on the, 63  
 Ambergris, 148  
 Ambloplites rupestris, 384  
 Amblyrhynchus cristatus, 334  
 Amblystoma mavortium, 360, 366  
 Amblystoma opacum, 367  
     " punctatum, 360, 368  
 Ameiurus nebulosus, 417  
 Amia calva, 424  
 Ammospermophilus leucurus, 73  
 Ampelis cedrorum, 193  
     " garrulus, 192  
 Amphiuma means, 360, 370  
 "Amphibia and Reptiles," Gadow's book on, 354, 366  
 Amphibians, Bird's-Eye view of, 360  
 Amphibians, Order of Tailed, 366  
 Amphibians, Order of Worm-Like, 371  
 Amphioxus, 438  
 Anas boschas, 270  
     " obscura, 269  
 Ancistrodon contortrix, 351  
     " piscivorus, 352  
 Angel Fish, 387  
 Anguilla vulgaris, 421  
 Anhinga anhinga, 287  
 Anser albifrons gambeli, 282  
 Anseres, Order, 175  
 Ant-"Bear," 158  
 Ant-Eater, Great, 156, 158  
     " Tamandua, 159  
 Antelope, Prong-Horned, 99, 116  
 Antelope Squirrel, 73  
 Antilocapra americana, 99, 116  
 Antlers, Caribou, 133, 138  
     " Columbian Black-Tailed Deer, 127  
 Antlers, Moose, 142  
     " Mule-Deer, 126  
     " Record Elk, 124  
     " Sitka Deer, 128  
     " White-Tailed Deer, 130  
 Apes and Monkeys, Order of, 7  
     " Anthropoid, 7, 8  
 Aplodontia rufa, 80  
 Aplodontidae, Family, 68, 80  
 Apoda, Order, 360, 371  
 Apodes, 421  
 Aptenodytes forsteri, 307  
 Apteryx australis, 310  
 Aquila chrysaetus, 229  
 Ara ararauna, 217  
 Archaeopteryx, 359  
 Arctic Province, "Our, 44, 48  
 Arctonetta fischeri, 277, 279  
 Ardea caerulea, 260  
     " herodias, 259  
 Armadillo, Giant, 156  
     " Nine-Banded, 156, 158  
 Armadillo, Six-Banded, 156  
     " Three-Banded, 156, 157  
 Ardetta exilis, 263  
 Aromochelys odoratus, 326  
 Arvicola, 84, 86  
 Asio accipitrinus, 220  
     " wilsonianus, 220  
 Aspidonectes ferox, 329  
 Astragalinus tristris, 195  
 Atalapha cinerea, 65  
 Ateles ater, 7, 15  
 Atwood, Capt., 419, 433  
 Audubon Society, 181  
 Auk, "The (magazine), 264  
 Auk, 302  
     " Great, 305  
     " Razor-Billed, 300, 305  
 Auklet, Cassin's, 305  
     " Least, 305  
     " Rhinoceros, 304  
 Axolotl, 360, 366  
 Aye-Aye, 7  
 Aythya americana, 273  
     " collaris, 269  
     " marila, 269  
     " vallisneria, 274  
 Austin, Mrs. Mary, Poem by, 256  
 Baboons, 9, 13  
     " Gelada, 14  
 Badger, 32  
 Bailey, Mrs. Florence Merriam, 204, 234, 238, 303  
 Bailey, Vernon, 98  
 Baird, Prof. Spencer F., 376, 413, 421  
 Baker, Arthur B., 34, 77  
     " Dr. Frank, 103  
 Balaelna glacialis, 148  
     " mysticetus, 147  
     " sieboldii, 148  
 Balaeloptera physalus, 148  
     " sulfureus, 147  
 Balistes capricus, 410  
 Bandicoots, 163  
 Barren Grounds of Northern Canada, "The, 137  
 Barrett, Mrs. A. W., 385  
 Bass and Sun-Fish Family, 382  
     " Calico or Strawberry, 383  
     " Large-Mouthed Black, 383  
     " Rock, 383  
     " Small-Mouthed Black, 382  
     " Warmouth, 383  
 Bassariscus astutus, 41  
 Bassarisk, 41  
 Bates, H. W., 63  
 Batrachia, 359  
 Bat, Big-Eared, 65  
     " Blainville's Flower-Nosed, 62  
 Bat, Bonneted, 64  
     " California Leaf-Nosed, 61, 62  
 Bat, Common, 61  
     " False Vampire, 61, 65  
     " Free-Tailed, 61  
     " Fruit, 66  
     " Gray, 65  
     " Great Vampire, 63  
     " Hammer-Headed, 66  
     " Horseshoe, 61, 66

- Bat, Javelin, 63  
 " Long-Eared, 65  
 " Naked, 64  
 " Red, 64  
 " True Vampire, 62  
 Beal, F. E. L., 202, 205, 210, 212  
 Bean, Dr. T. H., 419  
 Beard, J. Carter, 436  
 Bear, Black, 35-39  
 " " Cinnamon, 40  
 " " Everglade, 35  
 " " Glacier, 35, 41  
 " " Labrador, 35  
 " " Louisiana, 35  
 " " Queen Charlotte, 35  
 " Big Brown, 35, 36  
 " " Kidder's, 35  
 " " Kodiak, 35, 36  
 " " Merriam's, 35  
 " " Peninsula, 35  
 " " Sitka, 35  
 " " Yakutat, 35  
 " Grizzly, 35, 37  
 " " Alaskan, 35  
 " " Barren-Ground, 35  
 " " Silver-Tip, 35  
 " " Sonora, 35  
 Beaver, American, 82  
 " Mountain, 80  
 Beck, R. J., 291, 334  
 Bedford, Duke of, 106  
 Bee-Bird, 206  
*Belgica*, S. S., 307  
 Bent, A. C., 264  
 Bering, Capt. Vitus, 154  
 Big-Horn or Rocky Mt. Sheep, 108  
 Blackbird, Crow, 202  
 " Red-Winged, 200  
 " Yellow-Headed, 200  
 Black-Gill Sunfish, 384  
*Blarina brevicauda*, 58  
 Bluebird, 182  
 Bluefish, 388  
 Biological Survey, 36, 181, 192, 212, 220, 240, 286  
 Bird collecting condemned, 172, 189  
 Bird Destruction, 171  
 " Skeleton of, 219  
 " Snake, 267, 287  
 " Frigate, 290  
 "Bird-Lore" (magazine), 266  
 Birds, Chart of, 176  
 " Fully-Palmated, 267  
 " Man-o'-War, 284, 290  
 " Swimming, 267  
 " Weak-Winged Diving, 267  
 " Web-Footed, 267, 284  
 Bird Life, Decrease in, 171  
 Birds and Mammals, The Destruction of, 171  
 Bird-World, An Introduction to the, 171  
 Bison, American, 99  
 Bittern, American, 262  
 " Least, 233  
 Boa Constrictor, 340  
 Bob-Cat, 21  
 Bobolink, 199  
 Bob-White, 242  
 Bolles, Frank, 213  
*Bonasa umbellus*, 243  
*Bos americanus*, 99  
*Botaurus lentiginosus*, 262  
*Bothrops lanceolatus*, 353  
 Bovidae, 99  
*Brachistoma caribaeum*, 438  
 Bradyrodidae, 156  
*Bradypus tridactylus*, 160  
 Brant, Black, 28  
*Branta canadensis*, 280  
 " *nigricans*, 281  
 Brewster, William, 288  
 Brown, A. E., 77, 103  
 " Barnum, 335  
 " William Harvey, 319  
 Bryant, W. E., 303  
*Bubo virginianus*, 222  
 Buffalo, American, 99, 100  
*Bufo lentiginosus*, 360, 364  
 Bullhead, Common, 417  
 Bunting, Indigo, 199  
 " Snow, 195  
 Burroughs, John, 303  
*Butorides virescens*, 260  
 Butcher-Bird, 191  
*Buteo borealis*, 229  
 " *lineatus*, 230  
 Caecilians, Family of, 360, 371  
*Cacomistole*, 41  
*Caenolestes*, 163  
 Caiman, Banded, 317  
 " Black, 317, 322  
 " Broad-Nosed, 317  
 " Eye-Browed, 322  
 " *latirostris*, 317  
 " niger, 317  
 " *palpebrosus*, 317  
 " Rough-Backed, 317, 322  
 Caiman *sclerops*, 317  
 " *trigonatus*, 317  
 Callithricidae, 7, 16  
*Callithrix jacchus*, 7, 17  
*Callotaria ursina*, 44, 47  
 Calmette, Dr., treatment of snake wounds by, 355  
 "Camudie" (boa constrictor), 341  
 Canidae, 18, 22  
*Canis latrans*, 23  
 " *occidentalis*, 22  
 Caprimulgidae, Family, 207  
 Capuchin, 14  
 Capybara, 341  
 Carcajou, 31  
*Carcharias caeruleus*, 433  
*Carcharodon carcharias*, 433  
 Cardinal, 198  
*Cardinalis cardinalis*, 198  
 Caribou, 99, 131  
 " Black-Faced, 134  
 " Barren-Ground, 135, 136  
 Caribou, Grant's, 136  
 " Greenland, 136  
 " Kenai, 134  
 " Mountain, 134  
 Caribou, Newfoundland, 134  
 " Osborn's, 134  
 " Peary's, 136, 137  
 " Woodland, 132, 133  
 Carnivora, 3, 18  
 Carp, 413  
 " Minnows and Suckers, Order of, 412  
 Cassowary, Ceram, 310  
*Casuarius galeata*, 310  
*Castor canadensis*, 80  
 Cat-Bird, 186  
 Catfish, Channel, 417  
 " Mississippi, 416  
 Catfishes, Order of, 416  
*Catharista urubu*, 234  
*Cathartes aura*, 233  
*Catostomus commersoni*, 412  
 Cattle and Sheep Family, 99  
 Cebidae, 7  
*Cebus hypoleucus*, 7, 14  
 Cedar Bird, 193  
*Centrocercus urophasianus*, 248  
 Century Company, The, 257  
*Ceratodus forsteri*, 380  
*Cercopithecus diana*, 7, 13  
*Cerorhinca monocerata*, 305  
*Certhia familiaris americanus*, 185  
*Cervus canadensis*, 99, 121, 124  
*Ceryle alcyon*, 215  
 Cetaceans, 146  
*Cetorhinus maximus*, 433  
*Chaenobryttus gulosus*, 383  
*Chaetura pelagica*, 208  
 Chaparral Cock, 215  
 Chapman, F. M., 263, 266  
*Charadrius dominicus*, 251  
*Charitonetta albeola*, 275  
 Chart of Birds, 176  
 " Mammals, 4  
 Chat, Yellow-Breasted, 190  
*Chauleasmus strepera*, 269  
*Cheiromeles torquatus*, 64  
*Chelone mydas*, 330  
 " *imbriata*, 330  
 Chelonia, Order, 314, 323  
 Chelonians, 325, 326, 330  
*Chelopus marmoratus*, 327  
 " *insculptus*, 328  
*Chelydra serpentina*, 329  
 Chelydridae, Family, 324, 328  
*Chen hyperborea*, 282  
 Chickadee, 184  
*Chilomycterus geometricus*, 411  
*Chimera collei*, 431  
 Chimeras, Order of, 431  
 Chipmunk, 79  
 Chipmunk, California, 73  
 " Eastern, 72  
 Chiroptera, 3, 59, 60  
*Choloepus hoffmani*, 160  
*Chordeiles virginianus*, 207  
*Chrysemys picta*, 327  
 Chuck-Will's-Widow, 208  
 Ciconiidae, 263  
*Cinclus mexicanus*, 187  
*Circus hudsonius*, 231  
*Cistudo carolina*, 325  
*Citellus tridecemlineatus*, 74

- Citellus franklini*, 75  
     " *richardsoni*, 75  
 Civet Cat, 41  
*Clangula islandica*, 269  
*Cobra-de-Capello*, 347  
     " Hooded, 347  
     " King, or Snake-Eating, 348  
*Coccyges*, Order, 175, 214  
*Coccyzus americanus*, 214  
*Cockatoo*, 217  
*Colaptes auratus luteus*, 211  
*Colinus virginianus*, 242  
*Collins*, Capt. J. W., 290  
*Coluber guttatus*, 343  
*Columba fasciata*, 238  
*Columbae*, Order, 175, 237  
*Condor*, 236  
*Conepatus*, 32  
 Congo "Snake," 360, 370  
 "Congressional Record," 50  
*Conurus carolinensis*, 217  
*Cook*, Dr. Frederick A., 307  
*Coot*, 258  
*Cope*, Prof. E. D., 336  
*Copperhead*, 351  
*Corbin*, Austin, 101, 103  
*Coregonus clupeiiformis*, 408  
*Cormorant*, 267, 284, 289  
     " Double-Crested, 287  
     " Pallas', 287  
*Corvus americanus*, 204  
     " *corax sinuatus*, 205  
*Corynorhinus macrotis*, 65  
*Coryphaena hippurus*, 392  
*Cotton-Mouth Moccasin*, 352  
*Cougar*, 19  
*Coyote*, 23  
*Crane*, Whooping, 255  
     " Sandhill, 256  
*Crappie*, 384  
*Creeper*, Brown, 185  
*Cristivomer namaycush*, 398  
*Crocodile*, American, 317, 322  
     " Australian, 317  
     " Broad-Nosed African,  
     317  
*Crocodile*, Cuban, 317, 321  
     " Florida, 317, 320  
     " Nile, 317  
     " Orinoco, 317, 322  
     " Salt-Water, 317  
     " Sharp-Nosed African,  
     317  
*Crocodiles and Alligators*, Man-  
     Eating, 319  
*Crocodiles and Alligators*, Nest-  
     ing Habits of, 319  
*Crocodiles and Alligators*, Order  
     of, 317  
*Crocodylia*, Order, 314, 317  
*Crocodylus acutus*, 317  
     " *cataphractus*, 317  
     " *floridanus*, 317, 320  
     " *intermedius*, 317, 322  
     " *johnstoni*, 317  
     " *niloticus*, 317  
     " *palustris*, 317  
     " *porosus*, 317  
     " *rhombifer*, 317, 321  
*Crossbill*, American, 195  
*Crotalus adamanteus*, 349  
     " *atrox*, 349, 350  
     " *cerastes*, 349, 351  
     " *confluentus*, 349  
     " *horridus*, 349, 350  
     " *lepidus*, 349  
     " *lucifer*, 349  
     " *mittelli*, 349  
     " *molossus*, 349  
     " *tigris*, 349  
*Crotaphytus collaris*, 335  
*Crow*, Clarke's, 204  
     " Common, 205  
*Crowley*, J. B., 50  
*Crustacea*, 370  
*Cryptobranchus alleghaniensis*,  
     360  
*Cryptobranchus maximus*, 369  
*Cuckoo*, Black-Billed, 214  
     " Yellow-Billed, 214  
 "Culebra de Agua" (anaconda),  
     341  
*Curllew*, Long-Billed, 253  
*Cyanocephalus cyanocephalus*,  
     204  
*Cyanocitta cristata*, 203  
     " *stelleri*, 203  
*Cyanospiza cyanea*, 199  
*Cynomys ludovicianus*, 76  
*Cyprinus carpio*, 413  
*Cyrtonyx montezumae mearnsi*,  
     243  
*Cystophora cristata*, 44, 53  
 "Dabchick" (grebe), 301  
*Dactylopterus volitans*, 409  
*Dafila acuta*, 271  
*Dahl*, Keeper Frederick, 342  
*Darter*, 284, 287  
*Dasyus sexcinctus*, 157  
*Dasyures*, 163  
*Deer*, Arizona White-Tailed, 129  
     " Axis, 118  
     " Columbian Black-Tailed,  
     99, 127  
*Deer*, Florida White-Tailed, 129  
     " "Jumping," 127  
     " Mule or "Black-Tailed,"  
     126  
*Deer*, Sitka, 128  
     " White-Tailed or Virginia,  
     99, 128  
*Deer Family*, 99, 118  
*Delphinapterus leucas*, 149  
*Delphinus delphis*, 151  
*Dendragapus obscurus*, 244  
*Dendrocygna fulva*, 269  
*Dendroica aestiva*, 189  
*Desmognathus fusca*, 368  
*Devil-Fish*, 436  
*De Weese*, Dall, 142, 143, 249  
*Diana Monkey*, 13  
*Dickerson*, Mrs. E. N., 389  
*Dicrostonyx hudsonius*, 84  
*Didelphis virginiana*, 165  
*Diedipper* (grebe), 301  
*Diemyctylus viridescens*, 368  
*Diggers*, Order of the, 161  
*Dinosaur*, 325  
*Diomedea albatrus*, 294  
     " *chinensis*, 294  
     " *exulans*, 292  
     " *nigripes*, 292  
*Dipodomys merriami*, 84  
*Dippers*, 179, 187  
*Ditmars*, Raymond L., 338, 339,  
     342, 343, 346, 361, 367  
*Diver*, Great Northern, 300  
*Divers*, Order of Weak-Winged,  
     267, 300  
*Divers*, Order of Flightless, 267,  
     307  
*Dobson*, Dr. G. E., 62  
*Dodge*, Col. R. I., 102  
*Dog Family*, 22  
*Dogfish*, Order of the, 424  
*Dolichonyx oryzivorus*, 199  
*Dolphin*, Common, 151  
*Dolphin and Porpoise Family*,  
     149  
 "Dolphin," 392  
*Doroucoulis*, 15  
*Dove*, Mourning, 238  
*Dovekie*, 302  
*Dromaeus novaehollandae*, 310  
*Dryobates villosus*, 213  
*Du Chaillu*, Paul B., 66  
*Duck-Bill*, 167, 359  
*Duck*, American Scoter, 269, 278  
     " American Widgeon, 269  
     " Barrow's Golden-Eye, 269  
     " Black, 269  
     " Blue-Winged Teal, 270  
     " Buffle-Head or Butter-  
     Ball, 275  
*Duck*, Canvasback, 274  
     " Cinnamon Teal, 271  
     " Eider, 277  
     " Fulvus Tree, 269  
     " Gray, 269  
     " Green-Winged Teal, 269  
     " Harlequin, 269, 275  
     " Mallard, 268, 270  
     " Old Squaw, 269  
     " Pintail or Sprigtail, 271  
     " Red-Breasted Merganser,  
     278  
*Duck*, Red-Head, 273  
     " Ring-Necked, 269  
     " Scaup, 269  
     " Shoveller or Spoonbill,  
     271  
*Duck*, Spectacled Eider, 277  
     " Surf Scoter, 269  
     " White-Winged Scoter,  
     278  
*Duck*, Wood, 272  
*Ducks*, Geese, and Swans, Order  
     of, 267  
*Dugong*, 153, 154  
*Dutcher*, William, 181, 298  
*Dyche*, Prof. L. L., 45, 55, 96,  
     115, 123, 143  
*Eagle*, Bald, 227  
     " Golden, 229  
*Ecaudata*, Order, 360  
*Echidnas*, 167, 168  
*Ectopistes migratorius*, 237  
*Edentata*, Order, 3, 156

- Eel, Common, 421  
 " Electric, 421  
 " Lamper, 422  
 Eels, Order of, 421  
 Effodientia, Order, 3, 161  
 Egret, American, 261  
 " Snowy, 261  
 Egretta candidissima, 261  
 Eider, American, 277  
 " Spectacled, 277  
 Eigenmann, Dr. C. H., 60  
 Elanoides forficatus, 232  
 Elaps euryxanthus, 353  
 " fulvius, 353  
 Elk or Wapiti, 99, 118, 121  
 Ellachiek, 327  
 Elliot, D. G., 254, 271, 283  
 Elliott, Henry W., 44, 47, 50  
 " William, 436  
 Emeu, 310  
 Engelman, Dr., 429  
 Eniconetta stelleri, 279  
 Epomophorus, 66  
 Erethizon dorsatus, 94  
 " epixanthus, 94  
 Ereunetes pusillus, 252  
 Erignathus barbatus, 52  
 Erismatura jamaicensis, 279  
 Esehricht, D. F., 150  
 Esox lucius, 394  
 " masquinongy, 394  
 " ohioensis, 395  
 " reticulatus, 395  
 Eumeces quinquelineatus, 334  
 Eumetopias stelleri, 44, 46  
 Euneetes murinus, 341  
 " notaeus, 341  
 Eupotomis gibbosus, 384  
 Eutamias speciosus, 73  
 "Evening Bulletin," San Francisco, 417  
 Evermann, Dr. Barton W., 375, 385, 388, 398, 401, 428, 429, 438  
 Evotomys gapperi, 84, 87  
 " rutilus, 87  
 Exocactus volitans, 409  
 Falco columbarius, 227  
 " peregrinus anatum, 227  
 " sparverius, 226  
 Felis concolor, 19  
 " onca, 18  
 " pardalis, 20  
 Ferae, Order, 3, 18  
 Fer-de-Lance or Lance-Head Snake, 353  
 Ferret, Black-Footed, 29  
 Fiber zibethicus, 83, 84, 93  
 Finches, 179, 195  
 Fish, Angel, 387  
 " Angler or Goose, 420  
 " Bellows or Rabbit, 410  
 " Blue Cat-, 416  
 " Box or Trunk, 410  
 " Buffalo, 413  
 " Channel Cat-, 417  
 " Devil, 436  
 " Flying, 409  
 " Gar or Bill, 425  
 Fish, Hag, 438  
 " Mud-, 381  
 " Orange File-, 410  
 " Paddle, 429  
 " Porcupine, 411  
 " Sucking, 393  
 " Trigger or File, 410  
 Fish Commission, U. S. (See Fisheries Bureau.)  
 Fish-Hawk, 225  
 Fisher, Dr. A. K., 219, 220, 222, 227, 229, 231  
 Fisher, 30  
 Fishes, Class of, 375  
 " Order of Flat, 418  
 " Order of Foot, 420  
 " Order of Gar or Ganoid, 425  
 Fishes, Order of Half-Gilled, 415  
 " Order of Solid-Jaw, 410  
 " Order of Spiny-Finned, 382  
 Fishes, Pipe, and Sea-Horses, 423  
 Fishes," "Descriptive Catalogue of, 375  
 "Fishes, Living and Fossil," 431  
 "Fishes of North and Middle America," 375, 438  
 Fisheries, U. S. Bureau of, 376, 386, 398, 430  
 "Fishery Industries of the United States," 437  
 Flamingo, American, 266  
 " Order of, 266  
 Flounder, Winter, 418  
 Flycatchers, 179, 206  
 Flying-Fish, 409  
 " Gurnard, 409  
 Food and Game Fishes," "American, 429  
 Forbes, H. O., 291  
 Fox, Arctic, 24, 26  
 " Blue, 24, 26  
 " Coast Gray, 24  
 " Cross, 24, 25  
 " Florida Gray, 24  
 " Gray, 24, 26  
 " Hall Island, 24  
 " Kit, 24, 26  
 " Kodiak, 24  
 " Large-Eared, 24  
 " Newfoundland, 24  
 " Red, 24, 25  
 " Scott's Gray, 24  
 " Swift, 24, 26  
 " Texas Gray, 24  
 " Townsend's Gray, 24  
 Fratercula arctica, 304  
 Frear, William, 369  
 Fregata aquila, 290  
 Frigate-Bird, 284, 290, 291  
 Fringillidae, 179, 195  
 Frog, Bull, 360, 363  
 " Common, 360, 362  
 " Leopard, 362, 363  
 " Northern Tree, 364  
 " Tree, 360, 363  
 " Tongueless, 364  
 " Wood, 360, 363  
 Frogs and Toads, Order of, 361  
 Fulica americana, 258  
 Fulmar Family, 294  
 Fur-Bearers, The Small, 27  
 Fur Seal, 45, 48, 50  
 Gadow, Dr. H., 354, 366  
 Gadwall, 280  
 Gage, L. J., 368  
 Galeocерdo tigrinus, 433  
 Galeoscoptes carolinensis, 186  
 Gallinae, Order, 175, 241  
 Gallinago delicata, 252  
 Gallinula florida, 257  
 " galeata, 257  
 Game-Birds, Order of Upland, 241  
 Gannet, 284, 291  
 " Common, 288  
 Ganoids, Order of, 425  
 Gar, Alligator, 425  
 Gardiner's Island, Ospreys on, 226  
 Gar Pike, Long-Nosed, 425  
 " Short-Nosed, 425  
 Garrupa nigrita, 385  
 Gasterosteus aculeatus, 415  
 Gavia imber, 301  
 Gavial, Bornean, 317  
 " Indian, 317  
 Gavialis gangeticus, 317  
 Geese, Ducks, and Swans, 267  
 Gelada Baboon, 14  
 Geomys bursarius, 93, 94  
 Gibbons, 12  
 Gila Monster, 335  
 Ginglymodi, Order, 425  
 Glaniostomi, Order, 427  
 Glires, 3, 68  
 Globicephala melas, 149  
 Glutton, 31  
 Glyptodont, 156  
 Gnawing Animals, Order of, 68  
 Goat, Kennedy's, 115  
 " Rocky Mountain or White, 114, 399  
 Goatsuckers, 207  
 Goldfinch, American, 195, 196  
 "Gonies" (Albatross), 293  
 Goode, G. Brown, 149, 386, 396, 410, 419, 420, 425, 433  
 Goose, American White-Fronted, 282  
 Goose, Black Brant, 281  
 " Brant, 281  
 " Cackling, 281  
 " Canada, 280  
 " Hutehins's, 281  
 " Snow, 282  
 " White-Cheeked, 281  
 Gopher Family, Pocket, 68, 93  
 " Red Pocket, 93  
 Gorilla, 8  
 Goshawk, American, 231  
 Grackle, Purple, 202  
 Grampus griseus, 149  
 Granger, John T., 391  
 Grant, Madison, 138  
 Gray, Capt. David, 147  
 Gray Duck, 269

- Gray, J. C., 276  
 Grebe, Pied-Billed or Carolina, 300  
 Greely, Gen. A. W., 101, 105  
 Grinnell, George B., 402  
 Grosbeak, Cardinal, 198  
 " Rose-Breasted, 198  
 Grouper, Black, 375  
 Grouse, Blue, 245  
 " Canada Spruce, 245  
 " Canadian Ruffed, 241  
 " Dusky, 245  
 " Franklin, 245  
 " Gray Ruffed, 244  
 " Oregon or Sabine's, 244  
 " Pine, 245  
 " Pinnated, 245  
 " Prairie Sharp-Tailed, 247  
 " Ruffed, 243  
 " Sage, 248  
 " Sooty, 245  
 Gruber, Peter, 355  
 Grunt, Black, 376  
 Grus americana, 255  
 " mexicana, 256  
 Guara alba, 263  
 " rubra, 263  
 Guillemots, 302  
 Gull, Herring, 296  
 Gulo luscus, 31  
 Gurnard, Flying, 409  
 Gymnogyphus californianus, 234  
 Gymnotus electricus, 421  
 Haemulon plumieri, 376  
 Haliaeetus leucocephalus, 227  
 Halibut, Common, 418  
 Hallock, Charles, 424  
 " Handbook of Birds, Western United States," 234  
 Hang-Nest, 201  
 Hare and Rabbit Family, 68, 95, 97  
 Hare, Jack, 96  
 " Little Chief or Crying, 95  
 " Northern Varying, 95  
 " Polar, 96  
 " Prairie, 96  
 Harelda hyemalis, 269  
 Harporhynchus rufus 186  
 Harris, W. C., 429  
 " Haven of Refuge," for Ducks, 276  
 Hawk, Chicken, 229  
 " Cooper's, 230  
 " Duck, 227  
 " Fish, 225  
 " Forked-Tailed, 232  
 " Hen, 229, 231  
 " Marsh, 230  
 " Pigeon, 227  
 " Red-Shouldered, 230  
 " Red-Tailed, 229  
 " Sharp-Shinned, 230  
 " Sparrow, 226  
 Hawk and Eagle Family, 218, 225  
 Hellbender, 360, 368, 369  
 Hell-Diver (grebe), 300  
 Heloderma suspectum, 335  
 Hemibranchii, Order, 415  
 Herodiones, Order, 175, 259, 264  
 Heron Family, 259  
 " Great Blue, 259  
 " Little Blue, 260  
 " Little Green, 260  
 " Snowy, 261  
 Herring Gull, 296  
 Hesperornis, 278, 359  
 Heterodon platyrhinus, 346  
 Heterosomata, 418  
 Hill, James J., 103  
 Hippocampus heptagonus, 423  
 Hippoglossus hippoglossus, 418  
 Hirundo erythrogaster, 195  
 Histrionicus histrionicus, 269, 275  
 Histriophoca fasciata, 44, 53  
 Hog, Red River-, 144  
 " Wart, 143  
 Holacanthus ciliaris, 387  
 Holder, C. F., 389  
 Homo sapiens, 8  
 Hoofed Animals, Order of, 99  
 Hornaday, W. T., 112  
 Horns, Mountain Sheep, 112  
 " Prong-Horned Antelope, 116, 117  
 Hoy, Dr. P. R., 75  
 Humming-Birds, 207  
 " Ruby-Throated, 208  
 Hydrodamalis, 153  
 Hyla faber, 364  
 " versicolor, 360, 364  
 Hylobates leuciscus, 7, 12  
 Hylcichla mustelina, 182  
 Ibis Family, 263  
 Ibis, Glossy, 263  
 " Scarlet, 263  
 " White, 263  
 " Wood, 263  
 Icterus furcatus, 416  
 " punctatus, 417  
 Icteria virens, 190  
 Icterus galbula, 201  
 Ictiobus cyprinella, 413  
 Iguana, Common, 333  
 " Marine, 332, 334  
 " Rhinoceros, 334  
 Inupennes, Order, 175, 267, 307  
 Insectivora, Order, 3, 56  
 Ionornis martinica, 257  
 Isospondyli, Order, 396  
 Jackson, Chester E., 288  
 " Dr. Sheldon, 138  
 Jaguar, 18  
 Jaeger, Parasitic, 296, 299  
 Jaegers and Skuas, 296  
 Japanese Red-Faced Monkey, 13  
 Jay, Blue, 203  
 " Canada, 204  
 " Pinon, 204  
 " Steller's, 203  
 Jewfish, 385  
 Jones, C. J., 23, 103  
 Jordan, Dr. David S., 375, 385, 388, 398, 401, 428, 429, 438  
 Judd, Sylvester D., 192, 195  
 Jumping Mouse Family, 68  
 Junco hyemalis, 196  
 Junco, Slate-Colored, 196  
 Kangaroo, Boomer, 164  
 " Gray, 164  
 " Old Man, 164  
 " Rat, 164  
 " Red, 164  
 " Tree, 164  
 Kea, 216  
 Kenworthy, Dr. C. J., 392  
 Kidder, James H., 36  
 Kingbird, 206  
 Kingfisher, Belted, 215  
 Kinglets, 179, 183  
 Kite, Swallow-Tailed, 232  
 Kiwi, 310  
 Kogia, 148  
 Lacertilia, Order, 314, 333  
 " Lacey Law," 298  
 Lagocephalus laevis, 410  
 Lagopus lagopus, 249  
 " leucurus, 249  
 Lamna cornubica, 433  
 Lamper "Eel," 422, 437  
 Lamprey, Brook, 437  
 " Sea, 437  
 Lancelets, The, 438  
 Laridae, 296  
 Lark, Horned, 179, 206  
 " Meadow, 200  
 " Shore, 206  
 Larus argentatus, 296  
 Lasiurus borealis, 64  
 Latax lutris, 27  
 Latham, Mrs. C. F., 331, 344, 350  
 Laysan Island, 294  
 Leek, S. N., 122  
 Lemming, 84  
 " False, 86  
 " Hudson Bay, 84  
 Lemming Mouse, or False Lemming, 86  
 Lemming Mouse, Cooper's, 86  
 Lemur varius, 7, 17  
 Lemurs, 7  
 " Ruffed, or Black and White, 17  
 Lemurs, Suborder of, 17  
 Lepidosiren, 381  
 Lepidosteus osseus, 425  
 " platystomus, 425  
 " spatula, 425  
 Lepomis pallidus, 384  
 Leporidae, 68, 95  
 Leptocardii, 438  
 Lepus americanus, 95  
 " arcticus, 96  
 " campestris, 96  
 " sylvaticus, 96  
 " texianus, 96  
 Light-House Board, 303  
 Limicolae, Order, 175, 251  
 Lizard, Blue-Tailed, or Skink, 334  
 Lizard, Horned, 336  
 " Ring-Necked or Kangaroo, 335

- Lizards, Order of, 333  
 Longipennes, Order, 175, 267, 296  
 Lonnberg, E., 104  
 Loon, or Great Northern Diver, 267, 300  
 Lophius piscatorius, 420  
 Lophodytes cuculatus, 279  
 Lophortyx californicus, 243  
 Loring, J. Alden, 36, 230  
 Loxia curvirostra minor, 195  
 Lucas, Frederic A., 50, 288, 290, 305  
 Lunda cirrata, 304  
 Lung-Fish, Australian, 380  
 Lutianus aya, 391  
 Lutra canadensis, 27  
 Lutreola vison, 29  
 Lynx, Bay or Red, 21  
 " Canada, 21  
 Lynxes, 20  
 Macacus speciosus, 7  
 Macaw, Blue, 217  
 " Yellow, 217  
 MacDougal, Dr. D. T., 144, 215  
 Mackereel, Horse, 389  
 " Spanish, 388  
 Macrochelys temmincki, 328  
 Macrochires, Order, 175, 207  
 Macropus giganteus, 164  
 " rufus, 164  
 Madsen, Johannes, 105  
 Magpie, American, 202  
 Malacoelmys palustris, 327  
 Mammals, Chart of, 4  
 " Orders of, 3  
 " Order of Egg-Laying, 167  
 Mammals, Order of Flesh-Eating, 18  
 Mammals, Order of Pouched, 163  
 " Order of Toothless, 156  
 Man, 7  
 Manatee or Sea-Cow, 153  
 Mandrill, 14  
 Manis pentadactyla, 161  
 Man-o'-War Birds, 284, 290  
 " "Hawk," 291  
 Manta birostris, 436  
 Mareca americana, 269  
 Marmoset, Common, 17  
 " Pinche, 17  
 " Silky, 17  
 Marmot, 68  
 " Gray, or Whistler, 79  
 " Yellow-Bellied, 79  
 Marmota, 68, 76  
 " flaviventer, 79  
 " monax, 78  
 " prinosus, 79  
 Marsh, Prof. O. C., 278  
 Marsipobranchii, 437  
 Marsupialia, Order, 3, 163  
 Marten Family, 27, 29, 30  
 Marten, Pennant's, 30  
 " Pine, 30  
 Martin, Bee, 206  
 " Purple, 193  
 Massasauga, 349, 351  
 " Edward's, 349  
 Maynard, Lieut., 36  
 McCarthy, Eugene, 405  
 Meadow-Lark, 200  
 " Western, 201  
 Megachiroptera, 61  
 Megaderina, 65  
 Megaderma lyra, 61  
 Megaptera nodosa, 148  
 Megascops asio, 221  
 Melanerpes erythrocephalus, 211  
 " formicivorus, 212  
 Meleagris gallopavo, 250  
 " ocellata, 250  
 Melospiza fasciata, 197  
 Menobranhus, 360, 370  
 Menopoma alleghaniensis, 360, 369  
 Mephitis, 31  
 Merganser americanus, 279  
 " serrator, 278  
 Merganser, Hooded, 279  
 " Red-Breasted, 278  
 Merriam, Dr. C. Hart, 36, 61, 77, 83, 98, 143, 213, 225, 335  
 Merula migratoria, 181  
 Metopoceros cornutus, 334  
 Mice and Rats, Cheek-Pouched, 84, 91  
 Mice and Rats, Family of, 83  
 " " Typical North American, 84  
 Microchiroptera, 61  
 Microdipodops megacephalus, 84, 86  
 Micropterus dolomieu, 382  
 " salmoides, 383  
 Microtus (Arvicola) pennsylvanicus, 84  
 Midas aedipus, 17  
 " rosalia, 17  
 Mimus polyglottos, 187  
 Mink, 27, 29  
 Minnows, Carp, and Suckers, Order of, 412  
 Mitchell, H. R., 280  
 Moccasin, Copper-Bellied, 346  
 " Water or Cotton-Mouth, 343, 352  
 Mocking-Bird, 187  
 Mole, Common, 57  
 " Hairy-Tailed, 58  
 " Marsupial, 163  
 " Prairie or Silver, 58  
 " Star-Nosed, 58  
 Monotremata, Order, 3, 167  
 Monitors, 333  
 Monkey, Black-Faced Spider, 15  
 " Black Saki, 16  
 " Capuchin, 14  
 " Common Squirrel, 15  
 " Diana, 13  
 " Golden Howler, 16  
 " Japanese Red-Faced, 13  
 Monkey, Marmoset, 16  
 " Mexican Spider, 15  
 " Owl, 15  
 " Saki, 16  
 Monkey, Sapajou, 14  
 " Spider, 15  
 " Squirrel, 15  
 " Teetee, 15  
 " Uakari, 16  
 " Yarkee, 16  
 Monkeys, New World, 7, 14  
 " Old World, 7, 13  
 " Short-Tailed, 13  
 Moose, 99, 118, 122, 139  
 " Alaskan, 142  
 Monodon monoceros, 152  
 Moose-Bird, 204  
 Morehouse, Col. C. P., 389  
 Mormoops blainvillii, 62  
 Morris, Dr. Robt. T., 404  
 "Mother Carey's Chickens," 295  
 Mountain "Beaver," 80  
 " Lion, 19  
 " Sheep, 99, 107  
 " " Alaskan, 142  
 " " Big-Horn, 99, 108, 112  
 Mountain Sheep, Black, 98, 108, 112  
 Mountain Sheep, California or Nelson's, 110  
 Mountain Sheep, Fannin's, 108, 112  
 Mountain Sheep, Marco Polo's, 112  
 Mountain Sheep, Mexican, 108, 110, 112  
 Mountain Sheep, Siar, 112  
 " " White, or Dall's, 99, 108, 110, 112  
 Mouse, Cooper's Lemming, 86  
 " Field, 84, 86  
 " Gapper's Field, 87  
 " Grasshopper, 84, 91  
 " Harvest, 84  
 " Jumping, 68, 84, 92  
 " Le Conte's Harvest, 88  
 " Little Harvest, 88  
 " Lemming, 84  
 " Meadow, 86  
 " Missouri or Mole, 91  
 " Pocket, 84, 91, 93  
 " Red-Backed, 84, 87  
 " Rice-Field, 84, 88  
 " White-Footed, 84, 89  
 Mouse and Rat Family, 68, 84  
 " Cheek-Pouched, 84  
 Moxostoma aureolum, 413  
 Mud "Eel," 360, 371  
 " Hen, 258  
 " Puppy, 360, 370  
 Muger Crocodile, 317  
 Mugil brasiliensis, 391  
 Mullet, White or Silver, 391  
 Murre, 300, 302  
 " Brunnich's, 303  
 " California, 303  
 " Common, 303  
 Murrelets, 302  
 Museum, British, 341  
 Muskallunge, 394  
 Musk-Ox, 99, 103  
 Muskrat, 84, 93



- Mus rufescens*, 93  
*Mustela americana*, 30  
     " *pennanti*, 30  
 Mustelidae, 27  
*Myocastor coypus*, 92  
*Myrmecophaga jubata*, 158  
  
*Naja bungarus*, 338, 348  
     " *tripudians*, 347  
 Nansen, F., 55, 152  
 Narwhal, 152  
 Nathorst, Prof. A. G., 105, 106  
*Natrix fasciata*, 346  
*Necturus maculatus*, 360, 370  
 Nelson, E. W., 105, 110, 277  
*Nematognathi*, 416  
*Neotoma floridana*, 84, 88  
*Nettion carolinensis*, 269, 271  
 Newtons, 360, 368  
     " *Crimson-Spotted*, 368  
 Nighthawk, 207  
 Night-Heron, Black-Crowned,  
     261  
 Niles, E. O., 223  
*Nucifraga columbiana*, 204  
*Numenius longirostris*, 253  
 Nut-Cracker, Clarke's, 204  
 Nuthatches, 179, 184  
*Nyctala acadica*, 221  
*Nyctea nyctea*, 224  
*Nycticorax nycticorax naevius*,  
     261  
  
 Ocelot, 20  
*Ochotona princeps*, 95  
*Odobenus obesus*, 44, 53  
     " *rosmarus*, 44, 55  
*Odocoileus columbianus*, 99, 127  
     " *hemionus*, 99, 126  
     " *sitkensis*, 128  
     " *virginianus*, 99, 128  
*Odontoglossae*, Order, 175, 266  
*Oidemia americana*, 269  
     " *deglandi*, 278  
     " *perspicillata*, 269  
*Olor buccinator*, 283  
     " *columbianus*, 283  
*Oncorhynchus gorbusha*, 403  
     " *keta*, 403  
     " *kisutch*, 403  
     " *nerka*, 403  
     " *tschawyttscha*,  
         403  
*Onychomys leucogaster*, 91  
*Ophidia*, Order, 314  
*Ophiosurus ventralis*, 336  
*Opossum*, Murine, 166  
     " *Virginia*, 165  
 Orang-Utan, 10  
*Orcinus orca*, 149  
 Orders of Amphibians, 360  
     " *Birds*, 175  
     " *Fishes*, 378  
     " *Mammals*, 3  
     " *Reptiles*, 314  
*Oreamnos montanus*, 99, 114  
*Oreortyx pictus*, 243  
*Oreole*, Baltimore, 201  
 Ornithological Union, American,  
     264, 267, 291, 298  
  
*Ornithorhynchus anatinus*, 167  
*Orycteropus afer*, 161, 162  
*Oryzomys palustris*, 84, 88  
 Osborn, Prof. Henry F., 134, 156  
 Osgood, W. H., 77  
 Osprey, American, 225  
*Osteolaemus tetraspis*, 317  
*Ostracion quadricornis*, 410  
     " *tricornis*, 374  
 Ostrich, African, 309  
     " *South American*, 309  
*Otocoris alpestris*, 206  
*Otopterus californicus*, 62  
 Otter, 27, 341  
     " *Sea*, 27  
 "Outdoor Life" (magazine), 127  
 Ouzel, Water, 187  
*Ovibos moschatus*, 99, 103  
     " *wardi*, 105  
*Ovis ammon*, 112  
     " *canadensis*, 99, 112  
     " *dalli*, 99, 110, 112  
     " *fannini*, 108  
     " *mexicanus*, 108, 110, 112  
     " *nelsoni*, 108, 110  
     " *poli*, 112  
     " *siarensis*, 112  
     " *stonei*, 99  
 Owl, Barn, 219  
     " *Barred*, 220  
     " *Burrowing*, 224  
     " *Great Gray*, 221  
     " *Great Horned*, 222  
     " *Horned Family*, 218, 220  
     " *Long-Eared*, 220  
     " *Monkey-Faced*, 219  
     " *Saw-Whet*, 221  
     " *Screech*, 221  
     " *Short-Eared*, 220  
     " *Snowy*, 224  
*Oxyechus vocifera*, 251  
  
*Paddle-Fish*, 429  
 Palmer, Dr. T. S., 98, 143  
*Paludicola*, Order, 175, 255  
*Pandion haliaeetus carolinensis*,  
     225  
 Pangolin, Giant, 161  
     " *Indian*, 161  
 Pan troglodytes, 7, 9  
 Parrakeet, Carolina, 216, 217  
 Parrots and Macaws, Order of,  
     216  
 Partridge, Black, 245  
     " *Mearns'*, 243  
     " *Mountain*, 243  
     " *Valley*, 243  
*Parus atricapillus*, 184  
 Paschen, H., 9  
*Passer domesticus*, 197  
*Passeres*, Order, 175, 179  
*Passerina nivalis*, 195  
 Pasteur Institute, of Lille, 355  
 Peary, Commander Robt. E., 54,  
     55, 104, 107, 137, 307  
 Peccary, Collared, 99, 144  
     " *White-Lipped*, 144  
 Peccary Family, 99, 143  
*Pediocætes phasianellus cam-*  
     *pestris*, 247  
  
*Pelecanus californicus*, 286  
     " *erythrorhynchos*, 286  
     " *fuscus*, 284  
 Pelican, Brown, 284  
     " *California Brown*, 285,  
         286  
 Pelican, Great White, 286  
 Pelican Island, 285  
 Penguin, 267, 306, 308  
     " *Black-Footed*, 307  
     " *Emperor*, 307  
     " *Pack*, 307  
*Perca flavescens*, 386  
 Perch, Pike-, 387  
     " *Yellow*, 386  
*Perisoreus canadensis*, 204  
*Perodipus richardsoni*, 84, 91  
*Perognathus fasciatus*, 84, 91  
     " *flavus*, 93  
*Peromyscus leucopus*, 89  
 Petrel, 267  
     " *Stormy*, 294, 295  
*Petrochelidon lunifrons*, 193  
*Petromyzon marinus*, 437  
*Phalacrocorax carbo*, 287  
     " *dilophus*, 287  
     " *pelagicus*, 287  
 Phalangers, 163  
*Phasianus torquatus*, 250  
 Pheasant Family, 241, 250  
 Pheasant, Golden, 250  
     " *Ring-Necked*, 250  
     " *Silver*, 250  
*Phenacomys*, 87  
     " *orophilus*, 84, 88  
*Philohela minor*, 252  
*Phocaena communis*, 151  
*Phoca foetida*, 44, 52  
     " *groenlandica*, 44, 52  
     " *vitulina*, 44, 52  
*Phoenicopterus ruber*, 266  
*Phrynosoma cornutum*, 336  
*Phyllostoma hastatum*, 63  
*Physeter macrocephalus*, 148  
*Pica pica hudsonica*, 202  
 Pici, Order, 175, 210  
 Pickerel, Chain, 395  
*Picus pubescens medianus*, 212  
 Pigeons and Doves, Order of, 237  
 Pigeon, Banded-Tailed, 238  
     " *Passenger*, 237  
 Pika Family, 68, 95  
 Pike, 394  
     " *Wall-Eyed*, 387  
     " *Mr. Warburton*, 137  
 Pike-Perch, Yellow, 387  
 Pine-Hen, 245  
 Pinnated Grouse, 245  
 Pinnipedia, Order, 3, 43, 44  
*Pipa americana*, 364  
 Pipe-Fishes, Order of, 422  
*Piranga erythromelas*, 194  
*Pithecia satanas*, 16  
*Pityophis melanoleucus*, 344  
*Platypus* or Duck-Bill, 359  
*Plectognathi*, 410  
*Plectospondyli*, 412  
*Plegadis autumnalis*, 263  
*Plethodon glutinosus*, 367  
 Plover, American Golden, 251

- Plover, Field, 251  
     " Kill-Deer, 251  
 Pocket-Gopher Family, 68  
 Podilymbus podiceps, 300  
 Polyodon spathula, 429  
 Pomatomus saltatrix, 388  
 Pomoxis annularis, 384  
     " sparoides, 383  
 Pompano, Common, 390  
 Porcupine, Canada, 94  
     " Yellow-Haired, 94  
 Porcupine Family, 68, 94  
 Porpoise, Common, 151  
 Porpoises and Whales, Order of, 146  
 Porzana carolina, 257  
 Pouched Mouse and Rat Family, 68  
 Prairie-Chicken, 245  
 Prairie-"Dog," 68, 76  
     " " pamphlet on the, 98  
 Prairie-"Dog" Burrow, 77  
 Prairie-"Dog" and Burrowing Owl, 224  
 Prairie-"Dog" Hunter, 29  
 Prairie Wolf, 23  
 Primates, 3, 7  
 Pristis pectinatus, 434  
 Procellaria glacialis, 307  
     " pelagica, 295  
 Procyon lotor, 41  
 Progne subis, 193  
 Promops californicus, 64  
 Protean, subterranean, 370  
 Proteus anguineus, 370  
 Protopterus, 351  
 Pseudemys rubriventris, 326  
 Pseudopleuronectes americanus, 418  
 Psittaci, Order, 175, 216  
 Ptarmigan, 241, 307  
     " White-Tailed, 249  
     " Willow, 249  
 Pteropus edwardsii, 66  
 Puffin, Common, or "Sea Parrot," 304  
 Puffin, Tufted, 300, 304  
 Puffins, 267, 300, 302, 303  
 Puma, 19  
 Putorius erminea, 29  
     " nigripes, 29  
     " rixosus, 29  
 Pygopodes, Order, 175, 267, 300  
 Python molurus, 340, 342  
     " reticulatus, 341  
 Quail, 241  
     " California Mountain, 243  
     " Common, 242  
     " Valley, 243  
 Querquedula cyanoptera, 271  
     " discors, 270  
 Quisculus quiscula, 202  
 Rabbit and Hare Family, 95  
 Rabbit, Jack, 98  
 Raccoon Family, 41  
 Rachianectes glaucus, 148  
 Raia, 434  
 Rail Family, 257  
 Rail, Sora, 257  
     " Virginia, 257  
 Rallus virginianus, 257  
 Rana catesbiana, 360, 363  
     " clamata, 360, 362  
     " sylvatica, 360, 363  
     " virescens, 362  
 Rangifer arcticus, 99, 135, 136  
     " caribou, 99, 132, 133  
     " granti, 136  
     " groenlandicus, 136  
     " osborni, 134  
     " pearyi, 136, 137  
     " stonei, 134  
 Raptores, Order, 175, 218  
 Rat and Mouse Family, 68, 83, 84, 91  
 Rat, Cotton or Marsh, 84, 89  
     " Coypu, 92  
     " Domestic, 84, 93  
     " Kangaroo, 84, 91, 93  
     " Pack or Trading, 88, 93  
     " Tree, 93  
     " Wood, 84, 88  
 Rats and Rat-like Animals, 92  
 Ratitae, Order, 175, 309  
 Rattlesnake, 348  
     " Diamond, or Water, 349, 350  
 Rattlesnake, Dog-Faced, 349  
     " Edwards', 349  
     " Green, 349  
     " Ground, 349  
     " Horned, 349, 351  
     " Massasauga, 349, 351  
 Rattlesnake, Pacific, 349  
     " Prairie, 349, 350, 351  
 Rattlesnake, Texas, 349, 350  
     " Timber or Banded, 349, 350  
 Rattlesnake, White, 349  
 Raven, American, 205, 307  
 Ray, Beaked, 434  
     " Devil-Fish, 436  
     " Shark-, 434  
     " Sting, 435  
 Rays and Skates, Order of, 434  
 "Recreation" (magazine), 237, 276  
 Red Horse, 413  
 Redstart, American, 190  
 Reed-Bird, 199  
 Regulus calendula, 183  
 Reindeer, in Alaska, 138  
 Reithrodontomys leonti, 84, 88  
 Remora, 393  
 Reptile House, 339  
 Reptiles, Food Consumed by, 339  
 Reptiles, Introduction to the Class of, 313  
 Reptiles, Orders of, 314  
     " Poisonous Species of, 315  
 Rice-Bird, 199  
 Ring-Tail Monkey, 14  
 Rhamphobatis ancylostomus, 434  
 Rhea, 309  
 Rhinobatis lentiginosus, 434  
 Rhinodon typicus, 433  
 Rhytina gigas, 153  
 Road-Runner, 215  
 Robin, 181  
 Roccus lineatus, 385  
 Rodentia, 3, 68  
 Rungius, Carl, 143  
 Rutter, Cloudsley, 401  
 Rynchops nigra, 298  
 Sage-Grouse, 247  
 Saimiri sciurea, 15  
 Salamanders, Eel-Like, 370  
     " Family of, 360, 366  
 Salamander, Free-Gilled, 370  
     " Giant, 369  
     " Siren, or Mud-"Eel," 371  
 Salamanders, Spotted, 360, 366, 368  
 Salamanders, Two-Legged, 370  
 Salamandridae, 360, 364  
 Salmo clarkii, 396  
     " gairdneri, 398  
     " irideus, 397  
     " ouananiche, 405  
     " salar, 404  
     " sebago, 405  
 Salmon and Trout, Order of, 396  
     " Atlantic, 404  
     " Blueback or Sockeye, 403  
 Salmon, Dog, 403  
     " Family of the, 396  
     " Groups of American, 400  
     " Humpback, 403  
     " Ouananiche, 405  
     " Quinnet, 401, 403  
     " Sebago, 405  
 Salvelinus fontinalis, 399  
 Sand-Piper, Least, 253  
     " Semi-Palmated, 252  
 San Francisco "Evening Bulletin," 417  
 Sapajou, White-Throated, 14  
 Sapsucker, Yellow-Bellied, 213  
 Sarcorhampus gryphus, 236  
 Sawfish, 434  
 Scalops aquaticus, 57  
 Scammon, Capt. C. M., 147, 150  
 Scaphiopus holbrooki, 360, 364  
 Scuriidae, 68  
 Sciuropertus, 68  
     " volans, 80  
 Sciurus carolinensis, 70  
     " douglasi, 71  
     " erythrogaster, 71  
     " fremonti, 71  
     " griseus, 70  
     " hudsonicus, 71  
     " ludovicianus, 70  
     " malabaricus, 71  
     " niger, 70  
     " prevosti, 72  
 Scomberomorus maculatus, 388  
 Scoter, American, 269, 278

- Scoter, Surf, 269  
     " White-Winged, 278  
 Scotiaptex nebulosa, 221  
 Sea-Bass, Black, 385  
     " Family of the, 385  
 Sea-Cow, Rhytina, or Arctic, 154  
 Sea-Cows, Order of, 153  
 Sea-Horses, Order of, 423  
 Sea-Lions, California, 44, 46  
     " Steller's, 44, 46  
 "Sea-Parrot" (puffin), 304  
 "Sea-Swallow" (tern), 298  
 Seal, Bearded, 52  
     " Fur, 48  
     " Greenland, 53  
     " Harbor, 44, 52  
     " Harp, 44, 52  
     " Hooded, 44, 53  
     " Ribbon or Harlequin, 44, 53  
 Seal, Ringed, 44  
     " Saddle-Back, 53  
 Seal Family, 52  
 Seals and Sea-Lions, Order of, 43  
 Seiurus motacilla, 190  
     " noveboracensis, 190  
 Selachostomi, Order, 429  
 Selous, Percy, killed by moccasins, 252  
 Serpents, Order of, 337  
 Serum, Anti-venomous, 355  
 Seton, Ernest T., 134  
 Setophaga ruticilla, 190  
 Sewellel, 80  
 Shad, Common, 407  
 "Shagreen," 432  
 Shark, Basking, 433  
     " Blue, 433  
     " Bone, 433  
     " Great Tiger, 433  
     " Hammer-Head, 432, 433  
     " Mackerel, 433  
     " Man-Eater or White, 433  
     " Thresher, 433  
     " Tiger, 433  
 Shark-Ray, 434  
 Sharks, Order of, 432  
 Sharp-Tailed Grouse, 247  
 Shearwaters, 294  
 Sheep, Big-Horn, 99, 108  
     " Black, 99, 108, 112  
     " Fannin's, 108, 112  
     " Mexican, 108, 110  
     " Nelson's, or California, 108, 110  
 Sheep, White, 99, 108, 110  
 Sheep and Cattle Family, 99  
 Sheldon, Chas., 110, 112  
 Shields, G. O., 38, 112, 237, 276  
 Shore-Birds, Order of, 251  
 Showt'l, 80  
 Shrew Family, 58  
 Shrew, Common, 58  
     " Short-Tailed, 58  
 Shrikes, 179, 191  
 Sialia sialis, 182  
 Siamang, 13  
 Sigmodon hispidus, 84, 89  
 Simia satyrus, 7  
 Simorhynchus pusillus, 305  
 Sirenia, Order, 153  
 Siren larcetina, 360, 371  
 Sirens, Family of, 360  
 Siscowet, 399  
 Sistrurus catenatus, 349, 351  
     " edwardsi, 349  
     " miliarius, 349  
 Sitta carolinensis, 185  
 Skaus and Jaegers, 296, 299  
 Skeleton of American Bison, 100  
     " " Bald Eagle, 219  
     " " Gorilla, 8  
 Skimmer, Black, 296, 298  
 Skink, 334  
 Skunk, Badger, 31  
     " Common, 32  
     " Little Spotted, 32  
 "Slider" Terrapin, 326  
 Sloth, Three-Toed, 156  
     " Two-Toed, 156  
 Snake, Black, 344  
     " Blue, or Green Racer, 344  
 Snake, Boyle's, 343  
     " Coach, Whip, or Red Racer, 345  
 Snake, Copperhead, 347  
     " Corn, Red Racer or Rat, 343  
 Snake, Fer-de-Lance or Lance-Head, 353  
 Snake, Garter, 345  
     " Gopher, Black or Indigo, 344  
 Snake, Harlequin, 347, 353  
     " Hog-Nosed, 346  
     " "Hoop," 339  
     " King, Chain or Thunder, 343  
 Snake, Massasauga, 347  
     " Pine, 344  
     " Poisons, 353, 354, 355  
     " Rattle, 347, 348  
     " Red-Bellied Water, 346  
     " Splendid, 343  
     " Sonoran Coral, 347, 353  
     " Water, 346  
     " Water-Moccasin, 347  
 Snake-Bird, 267, 287  
 Snake-bites, Treatment of, 354, 355  
 "Snake," Glass, 339  
 Snakes, captive, Food of, 339  
     " Harmless, of the United States, 343  
 Snakes, oviparous, 338  
     " Poisonous, of North America, 347  
 Snakes, Viviparous, 338  
 Snapper, Red, 391  
 Snow-Bird, 196  
 Snow-Bunting, 195, 307  
 Snipe, Wilson's or Jack, 252  
 Snyder, Keeper Chas. E., 339, 342  
 Somateria dresseri, 277  
     " spectabilis, 279  
 Sorex personatus, 58  
 Spallanzani, Experiments of, 62  
 Sparrow, English, 197  
     " Song, 197  
     " Tree, 196  
 Sparrow, White-Throated, 197  
 Spatula clypeata, 271  
 Speotito cucularia hypogea, 224  
 Spermophile or Ground Squirrel, 73, 98  
 Spermophile, Franklin's, 75  
     " Thirteen-Lined or Leopard, 74  
 Spermophile, Richardson's, 75  
 Sphargis coriacea, 331  
 Spheniscus demersus, 307  
 Sphyrapicus varius, 213  
 Sphyrna zygaena, 433  
 Spilogale, 32  
 Spilotes corais couperii, 344  
 Spizella monticola, 196  
 Spoonbill Family, 264  
 Spoonbill, Roseate, 264  
 Squali, 330  
 Squirrel, Antelope, 73  
     " California Gray, 70  
     " Douglas', 71  
     " Fremont's, 71  
     " Gray, 70  
     " Malabar, 71  
     " Northern Fox or Cat, 70  
 Squirrel, Prevost's, 72  
     " Red, or Chickaree, 71  
     " Southern Fox, 70  
 Squirrel Family, 68  
 Squirrels, Flying, 68, 80  
     " Fox, 70  
     " Ground, 68, 73, 98  
     " Rock, or Chipmunks, 68, 72  
 Squirrels, Tree, 68  
 Stake-Driver (bittern), 262  
 Stearns, Silas, 392  
 Steganopodes, Order, 175, 267, 284  
 Stegastoma tigrinum, 433  
 Stejneger, Dr. Leonhard, 154, 354, 355  
 Steller, George W., 154  
 Stephan, Supt. S. A., 141  
 Stercorarius parasiticus, 299  
 Stereolepis gigas, 385  
 Sterna hirundo, 298  
 Stickleback, Two-Spined, 415  
 Sting Ray, or "Stingaree," 435, 436  
 Stizostedion vitreum, 387  
 Stone, A. J., 143  
 Stork Family, 259, 263  
 Strigidae, Family, 218  
 Strix pratincola, 219  
 Struthio camelus, 309  
 Sturgeon, Common, 427  
     " Lake, 427  
     " Short-Nosed, 427  
     " Shovel-Nosed, 429  
     " White, 428  
 Sturgeons, Order of, 427  
     " Size records of, 429  
 Sturnella magna, 200  
     " neglecta, 201  
 Sucker, Common Brook or White, 412  
 Sucker, Red-Horse, 413

- Suckers, Carp, and Minnows,  
   Order of, 412  
 Sucking-Fish, 393  
 Sula bassana, 288  
 Sun-Fish, Black-Gill or Blue-  
   Gill, 384  
 Sun-Fish, Common, 384  
 Swallow, Bank, 194  
   " Barn, 194  
   " Chimney, 208  
   " Cliff, 194  
   " Eave, 193  
   " Tree, 194  
   " White-Throated, 208  
 Swallow Family, 179  
 Swan, Trumpeter, 283  
   " Whistling, 283  
 Swans, Ducks, and Geese, Order  
   of, 267  
 Swifts, 207  
   " Chimney, 208  
 Swimmers, Diving, 267  
   " Flying, 267  
   " Long-Winged, 267,  
   296  
 Swimmers, Order of Tube-Nosed,  
   267, 291  
 Swimming-Birds, Orders of, 267  
 Swordfish, 393  
 Sylviidae, Family, 179, 183  
 Symphalangus syndactylus, 13  
 Synaptomys cooperi, 84, 86  
 Syngnathus acus, 423  
 Synnium varium, 220  
  
 Tamandua tetradactyla, 159  
 Tamias, 68  
   " striatus, 72  
 Tanager, Scarlet, 194  
 Tanagers, 179, 194  
 Tantalus loculator, 263  
 Tapir Family, 99, 144  
 Tapiurus dowi, 99  
   " terrestris, 144  
 Tarpon, 406  
   " atlanticus, 406  
 Tarsier, 7  
 Tatu novemcinctum, 158  
 Tayassu alboirostre, 144  
   " tajaacu, 99, 144  
 Teal, Blue-Winged, 270  
   " Cinnamon, 271  
   " Green-Winged, 269, 271  
 Teetee, 15  
 Tern, 267, 296  
   " Common, 297, 298  
 Terrapin, Alligator, 328  
   " Diamond-Back, 327  
   " Ellachick, 327  
   " Painted, 327  
   " Pine-Barren, 326  
   " Pond, 327  
   " Red-Bellied or "Slid-  
   er," 326  
 Terrapin, Snapping, 328, 329  
 Terrapin, Fresh-Water, 324  
   " Smooth-Shelled, 326, 328  
 Testudo polyphemus, 325  
   " vicina, 324  
 Tetraonidae, 241, 242  
  
 Tetrodon turgidus, 410  
 Thalaretos maritimus, 35  
 Thalamocheilus caretta, 330  
 Theropithecus gelada, 7, 14  
 Thrasher, Brown, 186  
 Thrushes, 179, 181  
   " Water, 190  
 Thunder-Pumper (bittern), 262  
 Thunnus thynnus, 389  
 Tiger-Cat, 20  
 Titmouse, Black-Capped, 184  
 Toads and Frogs, Order of, 361  
 Toad, American, 360  
   " Burrowing, 360, 364  
   " Common, 364  
   " Horned, 336  
   " Spade-Foot, 360, 364  
   " Surinam, 364  
 Tolyteutes tricinctus, 157  
 Tomistoma schlegeli, 317  
 Tortoise, Box, 325  
   " Giant, 324  
   " Gopher, 325  
   " Pond, 327  
 Townsend, Chas. H., 84, 86, 138,  
   276, 292, 376, 399, 414, 429  
 Trachinotus carolinus, 390  
   " goodei, 391  
 Tree-Creepers, 179, 185  
 Tree-Frogs, Family of, 360, 363  
 Trichechus americanus, 153  
   " latirostris, 153  
   " senegalensis, 153  
 Triton torosus, 368  
   " viridescens, 360, 368  
 Trochilus colubris, 208  
 Troglodytes aedon, 186  
 Trouessart, Dr. E. L., 64  
 Trout, Brook or Speckled, 399  
   " Lake or Mackinaw, 398  
   " Mountain or Black Spot-  
   ted, 396  
 Trout, Rainbow, 397  
   " Steelhead or Salmon, 398  
 Trout and Salmon, Order of, 396  
 Trygon sabinia, 435  
 Tubinares, Order, 175, 267, 292  
 Tufts, Le Roy M., 252  
 Tuna, or Tunny, 389  
   " Club, 389, 390  
 Turkey, Ocellated, 250  
   " Wild, 241  
 Turtle, Green, 330  
   " Hard-Shelled Sea-, 330  
   " Harp or Lyre, 331  
   " Hawksbill or Tortoise-  
   Shell, 330  
 Turtle, Leather-Backed, 331  
   " Leathery-Shelled Sea-,  
   331  
 Turtle, Loggerhead, 330  
   " Mud, 326  
   " Musk or Stink-Pot, 326  
   " Snapping, 329  
   " Soft-Shelled, 329  
   " Wood, 328  
 Tympanuchus americanus, 245  
 Typhlomolge rathbuni, 370  
 Typtilonectes compressicauda,  
   360  
  
 Tyrannus tyrannus, 206  
 Tyrrell, J. B., 134, 136  
  
 Uakaria calva, 16  
 Underwood, William Lyman,  
   364  
 Ungulata, Order, 3, 99  
 Uria californica, 303  
   " lomvia, 303  
   " troile, 303  
 Urocyon californicus, 24  
   " cinereoargenteus, 24,  
   26  
 Urocyon floridanus, 24  
   " scotti, 24  
   " texensis, 24  
   " townsendi, 24  
 Urodela, Order, 360, 366  
 Ursidae, 32  
 Ursus americanus, 35, 39  
   " sornborgeri,  
   35  
 Ursus carlottae, 35  
   " dalli, 35  
   " " gyas, 35  
   " emmonsii, 35  
   " floridanus, 35  
   " horribilis, 35, 37  
   " " alascensis, 35  
   " " horriaeus, 35  
   " kidderi, 35  
   " luteolus, 35  
   " merriami, 35  
   " middendorffi, 35, 36  
   " richardsoni, 35  
   " sitkensis, 35  
  
 Vampyrus spectrum, 63  
 Vega, " Voyage of the (Norden-  
   skiold's), 154  
 Vertebrates, Lowest Classes of,  
   437  
 Vespertilionidae, 61, 64  
 Vireos, 179, 191  
 Vole, 84, 87, 88  
 vom Hofe, Edward, 406  
 Vulture, Black, 234  
   " California, 234  
   " Common Turkey, 233  
 Vultures, Family of, 218, 232  
 Vulpes deletrix, 24  
   " fulvus, 24, 25  
   " " argentatus, 24  
   " " decussatus, 24  
   " hallensis, 24  
   " harrimani, 24  
   " lagopus, 24, 26  
   " macrotis, 24  
   " velox, 24, 26  
  
 Wagtails, 179  
 Wallabies, 164  
 Wallihan, A. G., 118, 126  
 Walrus, 43  
   " Atlantic, 44, 55  
   " Pacific, 44, 53  
 Wapiti, 99, 118, 121  
 Warblers, 179, 188  
 Ward, Henry A., 156

- Warren, E. R., 194, 205  
 Water-Frogs, Family of, 360  
 Waxwings, 179, 192  
 Weasel, Common, 27, 29  
 " Least, 29  
 Webster, Frederic S., 426  
 Weeds Destroyed by Doves, 240  
 Whale, Bowhead, Greenland or Polar, 147  
 Whale, California Gray, 148  
 " Sulphur-Bottom, 147  
 Whales, various species, 148  
 Whales and Porpoises, Order of, 146  
 Whales, Family of Baleen, 146, 147  
 Whales, Family of Sperm, 148  
 Whaling Stations, 148  
 Whippoorwill, 208  
 Whiskey-Jack, 204  
 White, Dr. Charles A., 344  
 Whitefish, Common, 408  
 Whitney, William C., 101, 107, 126  
 Widgeon, 269  
 Wingless Land Birds, Order of, 309  
 Wolcott, F. C., 38
- Wolf, Gray or Timber, 22  
 " Prairie, 23  
 " Tasmanian, 163  
 Wolverine, 27, 31  
 Wombat, 163  
 Woodchuck, 68, 76, 78  
 Woodcock, American, 252  
 Woodpecker, Ant-Eating, 212  
 " Downy, 212  
 " Golden-Winged, 211  
 Woodpecker, Hairy, 213  
 " Red-Headed, 211  
 Woodpeckers, Order of, 210  
 Wood Thrush, 182  
 Wren, House, 186  
 Wrens, 179  
 Wright, Mrs. Mabel Osgood, 184, 189, 213
- Xanthocephalus xanthocephalus, 200  
 Xiphias gladius, 393
- Yellowbird, Summer, 189
- Zalophus californianus, 44  
 Zamelodia ludoviciana, 198
- Zamenis constrictor, 344  
 " flagellum, 345  
 " " frenatum, 345  
 Zapodidae, 68, 84, 92  
 Zapus hudsonius, 84, 92  
 Zenaidura macroura, 238  
 Zonotrichia albicollis, 197  
 Zoological Garden, Antwerp, 106  
 Zoological Garden, Berlin, 106  
 " " Cincinnati, 141  
 Zoological Garden, Copenhagen, 106  
 Zoological Garden, London, 9, 163  
 Zoological Garden, Philadelphia, 77, 115  
 Zoological Park, National, 226, 234  
 Zoological Park, New York, 37, 226, 263, 270, 273, 279, 320, 324, 335, 339, 341, 342, 345  
 Zoological Society, London, 12, 298, 365  
 Zoological Society, New York, 171, 281













