



CREW
ON A T.
OF
PLANTS

1682










25715/10/1




E. L. H. Lloyd










Compt. #

1684:

—
—
—
—

1684



I most humbly Present this Booke to the Right Hon^{le}. My Lord the
Earle of Winchilsea, for his Library at Eastwell, Whom
God preserve; & grant him health & long life; w^{ch} is & shall be
Constantly the prayers of

My Lord.

Your Lordps
Most Faithfull Humble, Oblidged


&
Most Obedient Servant

Samuel

J. Finch

August y^e. 6th

1684.



*At a Meeting of the Council of the ROYAL
SOCIETY, Feb. 22. 1681*

DR. Grew having read several *Lectures* of the *Anatomy of Plants*, some whereof have been already printed at divers times, and some are not printed; with several other *Lectures* of their *Colours, Odours, Tasts, and Salts*; as also of the *Solution of Salts in Water*; and of *Mixture*; all of them to the satisfaction of the said *Society*: It is therefore Ordered, That He be desired, to cause them to printed together in one Volume.

CHR. WREN P.R.S.

Stewart

THE
ANATOMY
OF
PLANTS.
WITH AN
IDEA
OF A

Philosophical History of Plants.

And several other

LECTURE'S,
Read before the
ROYAL SOCIETY.

By *NEHEMIAH GREW* M.D. Fellow
of the *ROYAL SOCIETY*, and of the
COLLEGE of PHYSICIANS.

Printed by *W. Rawlins*, for the Author, 1682.

23237



Hindlesham Hall
1521 6/09

TO HIS MOST
Sacred Majesty
CHARLES II.
King of Great Britain, & c.

May it please Your Majesty,



THE Dedication of one Part of the following Anatomy having been very graciously received by Your Majesty: I am now emboldened most humbly to present the Whole into Your Royal Hands.

By which Your Majesty will find, That there are *Terræ Incognitæ* in Philosophy, as well as Geography. And for so much, as lies here, it comes to pass, I know not how, even in this Inquisitive Age, That I am the first, who have given a Map of the Country.

Your

The Epistle Dedicatory.

Your Majesty will here see, That there are those things within a Plant, little less admirable, than within an Animal. That a Plant, as well as an Animal, is composed of several Organical Parts; some whereof may be called its Bowels. That every Plant hath Bowels of divers kinds, containing divers kinds of Liquors. That even a Plant lives partly upon Aer; for the reception whereof, it hath those Parts which are answerable to Lungs. So that a Plant is, as it were, an Animal in Quires; as an Animal is a Plant, or rather several Plants bound up into one Volume.

Again, that all the said Organs, Bowels, or other Parts, are as artificially made; and for their Place and Number, as punctually set together; as all the Mathematick Lines of a Flower or Face. That the Staple of the Stuff is so exquisitely fine, that no Silk-worm is able to draw any thing near so small a Thred. So that one who walks about with the meanest Stick, holds a Piece of Natures Handicraft, which far surpasses the most elaborate Woof or Needle-Work in the World.

That by all these Means, the Ascent of the Sap, the Distribution of the Aer, the Confection of several sorts of Liquors, as Lympha's, Milks, Oyls, Balsames; with other parts of Vegetation, are all contrived and brought about in a Mechanical way.

In

The Epistle Dedicatory.

In sum, Your Majesty will find, that we are come ashore into a new World, whereof we see no end.

It may be, that some will say, into another Utopia. Yet not I, but Nature speaketh these things: the only true Pallas, wherewith it is treasonable for the most curiously banded Arachne to compare. In whose Name, I, the meanest of her Pupils, do in all humility crave Your Majesties Gracious Patronage. Whereof I cannot doubt, since Your Majesty hath been pleased to be the Founder, and to style Your Self the Patron of that Society, of which I have the honour to be a Member. Your Majesty deeming it to be a more Noble Design, To enlarge the Territories of Knowledge, than those of Dominion: and the Highest Pitch of Human Glory, not to rule, in any sort, over many; but to be a Good Prince over Wise Men. I am

Your Majesties


most humble

and

most obedient

Subject

NEHEMIAH GREW.



T H E

P R E F A C E.



It is a *Politick* or *Civil Virtue* in every prudent mans Eye, To set himself. an example, in what he doth, unto others. And in so doing, he looks upon himself as accountable, in some sort, to all Men. To those therefore, who may either expressly, or tacitly, expect the Reasons, upon which I first undertook the *Anatomy of Plants*, and also made the after-progress therein; I shall summe them up as follows.

The first occasion of directing my Thoughts this way, was in the *Year 1664*, upon reading some, of the many and curious Inventions of Learned Men, in the *Bodies of Animals*. For considering, that both of them came at first out of the same *Hand*; and were therefore the *Contrivances* of the same *Wisdom*: I thence fully assured my self, that it could not be a vain Design; to seek it in both. And being then newly furnished with a good stock of *Seeds*, in order to raise a *Nursery of Plants*; I resolved, besides what I first aimed at, to make the utmost use of them for that purpose: that so I might put somewhat upon that side the *Leaf* which the best *Botanicks* had left bare and empty. And in which, notwithstanding some other Learned Men had inserted somewhat of this nature; as Dr. *Highmore* in his *Book of Generation*, Dr. *Sharrock* of the *Propagation of Plants*, and Mr. *Hook* in his *Micrography*: yet but collaterally, and without shewing any purpose of managing this *Part of Natural History*. And although it seemed at first an Ob-
a jection

The Preface.

jection in my way, That the first projectors seldom bring their business to any good end: yet I also knew, That if Men should stay for an Example in every thing; nothing extraordinary would ever be done.

But notwithstanding the reasonableness of the Design; yet I did not forget, that, in respect of the Undertaker, there might be *Impar congressus*. And therefore, before I had ventured very far, in the Year 1668, I imparted it to my Brother-in-Law, the Learned Dr. Henry Sampson, now Fellow of the Colledge of Physicians in London. Who not only very well liked the same; but also excited me to a vigorous and accurate prosecution of it. Which he did, partly, by mentioning a very pertinent passage of Dr. Glisson, in the Preface to his *Book de Hepate*, (a) which I had not then read.

Plantæ quoque in hunc censum (sc. Anatomicum) veniunt; variâ enim Partium texturâ, & differentiis constant: & proculdubio, ex acurata earundem dissectione, utiles valde observationes nobis exurgerent: præstaretque in illis (inferioris licet ordinis) rebus examinandis operam impendere, quam in transcribendis ut sæpe fit, aliorum laboribus, inutiliter ætatem transigere. Quippe hoc pacto, ignavarum apum more, aliena duntaxat alvearia expilamus, nihilque bono publico adjicimus.

After I had finished the *First Book*, that I might know the sense also of other Learned Men, whether the steps I had already taken, would warrant me to proceed any further: I put some part of it into the same Hand; who, in the Year 1670, communicated the same to Mr. Oldenburge, then Secretary to the Royal Society: and after he had read it over, it was, upon his motion, delivered to that excellent Person Dr. John Wilkins then Bishop of Chester; who produced it at a Meeting of the Royal Society, and desired, they might see the rest. Which, or the greatest part, being also presented to them, the Right Honourable the Lord Vicount Brouncker, then President of the Royal Society, was pleased to peruse the same. Presently, after which, at a Meeting of the Council
of

The Preface.

of the said *Society*, the following *Order* was made, and entred in their Council-Book with this Date, and in these words:

May 11th 1671.

Then was Licensed Dr. Nehemiah Grew's Book, Entituled, The Anatomy of Vegetables begun; together with an account of Vegetation grounded thereupon. And Ordered to be Printed by the Printer to the Royal Society.

Hereupon, I was obliged to send the Book to the Press. And upon the 9th of *November* following in the same Year 1671, when it was near being printed, my Lord *Brouncker* signed the forementioned *Order*: the Printer, whose Name was to be inserted therein, not having received his *Diploma* till that time.

The Book being quickly after printed off, I ordered it to be Presented to the *Royal Society*; which was accordingly done at one of their Meetings *December 7, 1671*. And also to be sent to the Bishop of *Chester*; who was pleased to signify his acceptance thereof by a Letter dated at *Chester, December 26th 1671*. now filed amongst others in the Custody of the *Royal Society*: part whereof, in regard it relates to matter of Fact, I shall here recite.

Sir,

I did yesterday receive your Book; and am very sensible of the Honour you have done me in the Dedication of it. You was very happy in the choice
a 2 of

The Preface.

of this Subject to write upon; one of the most Noble and the most Copious parts of Philosophy; and such an one, as hath hitherto lain uncultivated. And you have been very successful in your first Attempt about it, in so many remarkable Observations and Discoveries, as you have made already. I could heartily wish, that you would still apply your self to this kind of Enquiries. You will find that Additionals will come in more copiously and easily. And it is not fit, that any one should, by his Superfluities, carry away the praise from him, who was the first Inventor, and who laid the Foundations, wherein the greatest difficulty doth consist, &c.

Having thus submitted my self to the Judgment of many Learned Men; I saw that my Journey must not here end. So that, like one who is got into a Wood, I thought I might as fairly find my way out, by going on, as by making a retreat. Whereupon, I began to draw up a Scheme of the whole Design.

While I was doing this, I received news from London, that the same day, December 7. 1671, in which my Book, then printed, was presented to the Royal Society: there was also presented a Manuscript (without Figures) from Seignior Malpighi, upon the same Subject; dated at Bononia, November, 1st 1671. the same, which Mr. Oldenburge, when it came to be printed, calleth his *Idea*. And of this, entry was made in their Journal Book. So that the Royal Society having now a Prospect of the good service of an Ancient Member, and one, who had highly merited by his Works then extant; from thence forward, I looked upon my self to be excused.

But soon after, receiving another Letter from the Bishop of Chester, dated at London, Febr. 18. 1672. I found

The Preface.

found the matter otherwise; and that the *Society* were pleased to engage me to proceed. Whereof entry was made by the *Secretary* in their *Journal Book*, at one of their Meetings, *April*, 18. 1672, in these words:

The Society was made acquainted with one particular lately passed in the Council; sc. That the Bishop of Chester had there proposed Dr. Grew to be a Curator to the Royal Society for the Anatomy of Plants: and that the Council had approved of that Proposal. Upon which, it was Ordered, That the Thanks of the Society be returned to the Lord Bishop of Chester, for this Proposal, and to the Council for their Approbation of the same.

This they might be induced to do; upon considering, that it would be no disadvantage to the credit of those matters, which were so new and strange, to be offered to the World from a double Authority. For one, although he may have no mind to deceive; yet is it more likely for one, than for two, to be deceived. Likewise, that the same Subject, being prosecuted by two Hands, would be the more illustrated by the different Examples produced by both. And that, as in other matters, so here, the defects of both, would mutually be supplied.

Whether for these, or other Reasons also, they were pleased to pass the forementioned *Order*; that being done, it had been very ill manners in me, not to have answered their expectation therein. And therefore reassuming the Design I had laid by, and having reduced it to some intelligible *Idea*, it was submitted to the Censure of the *Royal Society*: and it was thereupon ordered it should be printed.

Not

The Preface.

Not long after, I received a Curious and Learned Book from *Monf. Dodart*, *Archiater* to the Prince of *Conde*, and *Fellow* of the *Royal Academy* at *Paris*; in pursuance of whose Order, it was by him composed and published. Which being a Design of a like Import, I was glad to see it so far justify'd by that *Illustrious Society*, as well as by our own.

In this *Idea*, one principal Thing I insist upon, for a *Philosophical History* of *Plants*, is *Anatomy*. And, agreeing to the *Method* therein proposed, all the Observations contained in the *First Book*, except one or two, were made with the *Naked Eye*. To the end, I might first give a proof, How far it was possible for us to go, without the help of *Glasses*: which many Ingenious Men want; and more, the patience to manage them. For the Truth of these Observations, *Seignior Malpighi*, having procured my Book to be translated into Latin for his private use, speaks his own sense, in some of his Letters to *Mr. Oldenburge*, printed at the end of his *Anatomy* of *Plants*. And some of them, have since been confirmed, both by our Learned Country-men *Dr. Wallis*, and *Mr. Lister*; and by the Ingenious *Mr. Lewenhoeck*, abroad.

Having thus begun with the bare Eye; I next proceeded to the use of the *Microscope*. And the Observations thereby made, first on *Roots*, and afterwards on *Trunks* and *Branches*, together with the *Figures*, were all exhibited to the *Royal Society* at several times from *May 15. 1672.* to *April 2. 1674*; being the *Materials* for the *Second* and *Third Parts*: and hereof *Memorials* were inserted in their *Journal Books*.

After this, the *Royal Society* received from *Seignior Malpighi* his *Second Part* of the *Anatomy* of *Plants*, together with the *Figures* therein described, and his *Letters* to their *Secretary*, dated at *Bononia Aug. 20th* of the same year *1674.* when, and not before, he gave leave that the two said *Parts* should be printed.

So

The Preface.

So soon as I had finished the *Second* and *Third Parts*, I proceeded to the *Last*, *sc.* of *Leaves, Flowers, Fruits* and *Seeds*: and those Things I met with, more remarkable, were presented to the said *Society* in the Years 1676 & 1677. And the publishing of the former *Parts* successively, as well as of all together, hath been done in pursuance of their several *Orders* for the same.

Having concluded the *History of Perfect Plants*; I intended to have subjoyned the *Description* of those which are *Imperfect*. As also of *Parasitical, Marine, and Sensitive Plants*. And lastly, a view of the chief Particulars, wherein the *Mechanisme* of a *Plant*, is different from that of an *Animal*. But these things I leave to some other Hand.

The *First Book*, a little after it came forth; was translated into the French *Tongue*, by *Monf. Le Vasseur* an Ingenious Gentleman in *Paris*; elegantly, and in the Judgment of those who are well skilled in that *Language*, with much exactness, as to the sense. He having taken special care, to have all the difficulties of our own, by Me, cleared to him. And in a late Book Entituled, *Philosophia vetus & nova* printed at *Noriberg* 1682. the Learned Author seems to have made use of this Translation, for all that he hath taken notice of in that my *First Book*.

By the Ingenious Collectors of the *German Ephemerides*, both my *First, Second, and Third Books*, are all published in *Latine*. But their unskilful Interpreter doth often fail of the *Grammatical Sense*. Whose Errors, many of them very gross, I desire may be imputed neither to them, nor to my self.

Besides these, the *Second Lecture of Mixture* is also translated into *French*, by *Monf. Mesmin* a Learned Physician in *Paris*: whose *Version* is very well approved by those who are competent Judges hereof.

This,

The Preface.

This, and the rest which follow, are placed, not in the order of *Time*; but more according to their Nature or Relation one to another. All of them intended as a Commentary upon some particulars mentioned, either in the *First Lecture*, or in the *Idea*.

In the *Plates*, for the clearer conception of the *Part* described, I have represented it, generally, as entire, as its being magnified to some good degree, would bear. So, for instance, not the *Barque*, *Wood*, or *Pith* of a *Root* or *Tree*, by it self; but at least, some portion of all three together: Whereby, both their *Texture*, and also their Relation one to another, and the *Fabrick* of the whole, may be observed at one *View*. Yet have I not every where magnify'd the *Part* to the same degree; but more or less, as was necessary to represent what is spoken of it. And very highly, only in some few Examples, as in *Tab. 40.* which may suffice to illustrate the rest. Some of the *Plates*, especially those which I did not draw to the *Engravers* hand, are a little hard and stiff: but they are all well enough done, to represent what they intend.

THE

A N
I D E A
O F A
Philosophical History
O F
P L A N T S.

Read before the

ROYAL SOCIETY,

January 8. and January 15. 1672.

By *NEHEMIAH GREW* M.D. Fellow of the
Royal Society, and of the College of Physicians.

The Second Edition.

L O N D O N;

Printed by *W. Rawlins*, 1682.

THE
JOURNAL

OF

PHYSIOLOGICAL ANATOMY

OR

PLANT ANATOMY

BY
JAMES H. HENNING

Author of "The Principles of Plant Anatomy"

BY
JAMES H. HENNING

NEW YORK

1881

Published by the Author

TO THE
Most Illustrious
THE
ROYAL SOCIETY,

The following

I D E A

Is most HUMBLY

P R E S E N T E D,

A N D,

In their NAMES also

P R O P O S E D

TO THE

C O N S I D E R A T I O N

Of other

Learned Men.

By the AUTHOR

NEHEMIAH GREW.

Mrs. Johnson

ROYAL SOCIETY

1 D B A

PRESS LTD

1871

1871

1871

1871

T H E

C O N T E N T S.

UNTO what Degree the knowledge of Plants is arrived, §. 1. Wherein defective, §. 2. Why concluded to be so, §. 3. Yet capable of Improvement, §. 4. And worthy of it, §. 5.

Divers Instances given, wherein; first of the Organical Parts, as to their external Accidents and Oeconomical Uses, 6. Then of their Contents, Qualities, and Powers, 7. And an Improvement of this Part, will further that of divers other parts of knowledge; whereof Instances are given, 8.

In order whereto, Five General Means are propounded, 9. The First, a particular and comparative Survey of whatever is of more External consideration about Plants, 10. Instanced as to their Figures, 11, 12. Proportions, 13. Seasons, 14. Places, 15. Motions, 16.

The Second, A like Survey of the Organical Parts by Anatomy, as that which is very necessary, 17. In what manner to be prosecuted, both without, and with the Microscope, 18. What thereupon to be observed, 19. And what, from observation made, probably attainable, 20.

The Third, A like Survey of the Contents of Plants; their several Kinds, 21. Of all which, their Receptacles, 22. Motions, 23. Qualities, 24. Consistence, 25. Colours, Smells, and Tastes, 26. Where also the same Qualities are to be inquired into, as generally belonging to Plants, 26. As their Colours, 27. Odours, 28. Tastes, 29. Also their Faculties, 30. All these to be further examined, 31. By Contusion, 32. Agitation, 33. Frigification, 34. Infusion, 35. Subsession, 36. Digestion, 37, 38. Decoction, 39. Destillation, 40. Arefaction, 41. Assation;

The Contents.

Affation, 42. Ustion, 43. Calcination, 44. By Composition with other Bodies, 45. And by Compounding the Experiment it self, 46. What hence attainable, 47.

The Fourth, A like Survey of the Principles, as well as of the Contents, of the Organical Parts, 48. The Difficulty hereof, in some respects cleared, 49. Further, by two Instances, 50, 51. Some Remarques hereupon, of the Principles of Plants, 52. From hence will be attainable a further knowledge of the Modes of Vegetation, 53. Of the Qualities of Vegetables, 54. And of their Powers, 55, 56.

The Fifth, A like Survey of those Bodies, either from which these Principles are derived, or wherewith they have any communion, 57. Which are Four in general, scil. Earth, and all solid Receptacles, 58. Water, and all liquid Receptacles, 59. Aer, 60. And Sun, 61.

A Sixth General Inquiry, only hinted, 62.

The Conclusion, 63.



AN
 IDEA
 OF A
 Philosophical History
 OF
 PLANTS.

IF WE take an account of the *Degrees* whereunto the Knowledge of *Vegetables* is Advanced, it appeareth, That besides the great Varieties, which the Successful *Arts* of *Florists*, or Transplantations from one *Climate* to another, have produced; we have very many *Species* brought to light, especially Natives of the *Indies*, which the *Ancients*, for any thing that appears in their *Writings* now extant, were ignorant of. In which particular *Clusius*, *Columna*, *Baubinus*, *Boccone*, and others, have performed much. Withall, That their *Descriptions* (of all Parts above ground) their *Places* and *Seasons*, are with good diligence and preciseness set before us. Likewise their Order and Kindred: for the adjusting whereof our Learned Countryman Mr. *Ray*, and Dr. *Morrison*, have both taken very laudable pains. As also the ordering of them with respect to their *Alimental* and *Mechanick* Uses; for which, amongst others, Mr. *Evelyn* and Dr. *Beal* have deserved many thanks, and great praise. We are also informed, of the *Natures* and infallible *Faculties* of many of them. Whereunto so many as have assisted, have much obliged their *Posterity*.

2. §. By

2. §. By due Reflection upon what hath been *Performed*; it also appears, what is left *Imperfect*, and what *Undone*. For the *Virtues* of most *Plants*, are with much *uncertainty*, and too *promiscuously* ascribed to them. So that if you turn over an *Herbal*, you shall find almost every *Herb*, to be good for every *Disease*. And of the *Virtues* of many, they are altogether silent. And although, for the finding out, and just appropriation of them, they have left us some *Rules*, yet not all. The *Descriptions* likewise of many, are yet to be perfected; especially as to their *Roots*. Those who are very curious about the other *Parts*, being yet here too remiss. And as for their *Figures*, it were much to be wished, That they were all drawn by one *Scale*; or, at most, by Two; one, for *Trees* and *Shrubs*; and another for *Herbs*. Many likewise of their *Ranks* and *Affinities*, are yet undetermined. And a great number of *Names*, both *English* and *Latine*, not well given. So what we call *Goat's-Rue*, is not at all of kin to that Plant, whose *Generical Name* it bears. The like may be said of *Wild-Tansy*, *Stock-July-Flowers*, *Horse-Radish*, and many more. So also when we say *Bellis Major*, & *Minor*, as we commonly do, these *Names* would intimate, That the *Plants* to which they are given, differ (as the great double *Marigold*, doth from the less) only in Bulk: whereas, in truth, they are two *Species* of *Plants*. So we commonly say, *Centaurium Majus* & *Minus*, *Chelidonium Majus* & *Minus*, and of others in like manner, which yet are distinct *Species*, and of very different *Tribes*. But for the *Reason* of *Vegetation*, and the *Causes* of all those infinite *Varieties* therein observable (I mean so far as *Matter*, and the various *Affections* hereof, are instrumental thereunto) almost all Men have seemed to be unconcerned.

3. §. That Nothing hereof remaineth further to be known, is a Thought not well Calculated. For if we consider how long and gradual a *Journey* the *Knowledge* of *Nature* is; and how short a Time we have to proceed therein; as on the one hand, we shall conclude it our ease and profit, To see how far Others have gone before us: so shall we beware on the other, That we conceive not unduly of *Nature*, whilst we have a just value for Those, who were but her *Disciples*, and instructed by Her. Their Time and Abilities both, being short to her; which, as She was first Designed by *Divine Wisdom*; so may Her vast Dimensions best be adjudged of, in being compared Therewith. It will therefore be our Prudence, not to insist upon the Invidious Question, Which of Her *Scholars* have taken the fairest measure of Her; but to be well satisfied, that as yet She hath not been Circumscribed by Any.

4. §. Nor doth it more behove us to consider, how much of the Nature of *Vegetation* may lie before us yet *unknown*; Than, to believe, a great part thereof to be *knowable*. Not concluding from the acknowledged, much less supposed *Insuccesfulness*, of any Mens *Undertakings*: but from what may be accounted *Possible*, as to the Nature of things themselves; and from *Divine Providence*, by Infinite Ways conducting to the knowledge of them. Neither can we determine how great a part This may be: Because, *It is impossible to Measure, what we See not*. And since we are most likely to under-measure, we shall hereby but intrench our Endeavours, which we are not wont to carry beyond the *Idea*, which we have of our *Work*.

5. §. And

5. §. And how far soever this kind of Knowledge may be attainable, its being so far also worthy our attainment will be granted. For beholding the Many and Elegant Varieties, wherewith a Field or Garden is adorned; Who would not say, That it were exceeding pleasant to know what we See: and not more delightful, to one who has *Eyes*, to discern that all is very fine; than to another who hath *Reason*, to understand *how*. This surely were for a Man to take a True Inventory of his Goods, and his best way to put a price upon them. Yea it seems, that this were not only to be *Partaker of Divine Bounty*; but also, in some degree, *To be Copartner in the Secrets of Divine Art*. That which were very desirable, unless we should think it impertinent for us to design the *Knowing of That*, which God hath once thought fit to *Do*.

6. §. If for these, and other *Reasons*, an inquiry into the Nature of *Vegetation* may be of good Import; It will be requisite to see, first of all, What may offer it self to be enquired of; or to understand, what our *Scope* is: That so doing, we may take our aim the better in making, and having made, in applying our Observations thereunto. Amongst other Inquiries therefore, such as these deserve to be proposed. First, by what means it is that a *Plant*, or any *Part* of it, comes to *Grow*, a *Seed* to put forth a *Root* and *Trunk*; and this, all the other *Parts*, to the *seed* again; and all these being *formed*, by continual Nutrition still to be *increased*. How the Aliment by which a *Plant* is fed, is duly prepared in its several *Parts*; which way it is *conveyed* unto them; and in what manner it is *assimilated* to their respective Natures in them all. Whence this Growth and Augmentation, is not made of one, but many differing Degrees, unto both extremes of *small* and *great*; whether the comparison be made betwixt several *Plants*, or the several *Parts* of one. How not only their *Sizes*, but also their *Shapes* are so exceeding various; as of *Roots*, in being *Thick* or *Slender*, *Short* or *Long*, *Entire* or *Parted*, *Stringed* or *Ramified*, and the like: of *Trunks*, some being more *Entire*, others *Branched*, others *Shrub'd*: of *Leaves*, which are *Long* or *Round*, *Even-edg'd* or *Escallop'd*, and many other ways different, yet always *Flat*: and so for the other *Parts*. Then to inquire, What should be the reason of their various *Motions*; that the *Root* should *descend*; that its descent should sometimes be *perpendicular*, sometimes more *level*: That the *Trunk* doth *ascend*; and that the ascent thereof, as to the space of *Time* wherein it is made, is of different *measures*: and of divers other *Motions*, as they are observable in the *Roots*, *Trunks*, and other *Parts* of *Plants*. Whence again, these *Motions* have their Different, and Stated *Terms*; that *Plants* have their set and peculiar *Seasons* for their *Spring* or *Birth*, for their *Full Growth*, and for their *Teeming*; and the like. Further, what may be the *Causes* as of the *Seasons* of their *Growth*; so of the *Periods* of their *Lives*; some being *Annual*, others *Biennial*, others *Perennial*; some *Perennial* both as to their *Roots* and *Trunks*; and some as to their *Roots* only. Then, as they pass through these several *Seasons* of their *Lives*, in what manner their convenient *feeding*, *housing*, *cloathing* or *protection* otherwise, is contrived; wherein, in this kind and harmonious *Oeconomy*, one *Part*, may be officious to another, for the preservation of the health and life of the *whole*. And lastly, what care is taken, not only for themselves, but for their *Posterity*; in

what manner the *Seed* is prepared, formed and fitted for *Propagation*: and this being of so great concernment, how sometimes the other *Parts* also, as *Roots*, in putting forth *Trunks*; *Trunks* in putting forth *Roots*; yea in turning oftentimes into *Roots* themselves; whereof, in the *second Book of the Anatomy of Plants*, I shall give some instances. With other *Heads* of Inquiry of this kind.

7. §. Nor are the *Natures*, *Faculties*, and *Contents* of *Vegetables* less various, or a particular Inspection hereinto, of less concernment. For since All, or Most, seem to grow in the same manner, with one *Sun*, one *Rain*, indifferently well upon one *Soil*, and, to outward appearance, to have the same *Common Parts*; it may be asked, *How* it comes to pass, that their *Liquors*, or other *Contained Parts*, are of such different *Kinds*; one being Watry, another Winy, a third *Oily*, a fourth *Milky*, and the like. *How* also there is such a variety in their *Sensible Qualities*, as their *Colours*, *Tastes*, and *Smells*; what those *Materials* are, which are necessary to the *Being* of these *Qualities*; and those *Formalities*, wherein their *Essence* doth consist; as what it is that makes a *Plant*, or *Flower*, to be *white* or *red*; *fragrant* or *fetid*; *bitter* or *sweet*; or to be of any other *Colour*, *Smell*, or *Taste*. In like manner, their *Faculties* and *Powers*, what that is, or those things are, by which they are constituted; as whence one becomes *Purgative*, another *Vomitory*, a third *Diaphoretick*, &c. These, I say, with many other particular Inquiries depending hereupon; as they cannot but much oblige the Reason of Man to be obsequious to them, so by bringing in, at least, some satisfaction, will no less reward it. Especially, if it be withal considered, that besides our satisfaction as to the *Nature* of *Vegetation*; some further Light, to divers other parts of Knowledge, may likewise hence arise.

8. §. For since the present Design will engage us, to an accurate and multifarious Observation of *Plants*; we may hereby be enabled to *range* and *sort* them with more certainty, according to the Degrees of their *Affinity*. And all *Exoticks*, *Plants* or *Parts* of *Plants*, may probably be reduced to some such *Domesticks*, unto which they may bear the best Resemblance. Again, it may frequently conduct our minds to the consideration of the *State* of *Animals*; as whether there are not divers material Agreements betwixt them both; and what they are. *Wherein* also they may considerably differ, and what those things are which are more essential to their distinguishment. And besides, not only to compare what is already known of both; but also, by what may be observed in the *one*, to suggest and facilitate the finding out of what may yet be unobserved in the *other*. So also the consideration of the *Colours*, *Smells* and *Tastes* of *Vegetables*, may conduce to the Knowledge of the same *Qualities* in General; or of what it is, that constitutes them such, in any other Body: not as they are *actually* received by *Sense*; but so far, as such *Materials* or external *Circumstances*, are requisite to their becoming the *Adequate Objects* thereof. It may lead us also to inquire into further Ways of *Cultivation*, with respect to the whole *Plant*, or to the *Flower*, *Fruit*, or other *Part*: To amend them as to their *Sizes*, *Colours*, *Tastes*, *Fruitfulness*, or otherwise: To think of other Ways of *Propagation*; or to apply those already known to other *Plants* than hath been used. Likewise the Knowledge of their *Mechanical* Uses may hereby be enlarged; both as to the

Rea-

Reason of their use; in such particular *Trades* and *Manufactures*, already known; and the discovery of other uses yet unknown. As also their *Alimental*, with respect both to *Meats* and *Drinks*; the preparation of some, and the finding out of others. But especially their *Medicinal*; some *Plants* which have hitherto been neglected, may be applied to use; the *Perverted* uses of some, and the *Confused* uses of others, may be rectified. What may best correct their *Malignancies*, or inforce their *Virtues*; When needful to add the preparations of *Art* to That of *Nature*; How to Enlarge those of *Art*, and Rectifie those which are indeed *Inartificial*, may hereby be better conjectured. The knowledge of all which, that we may know how far it is accessible, and what probable Approaches may be made towards it; those several Means I have thought of, and suppose necessary thereunto, are next to be proposed.

9. §. Reflecting then upon the present Design, and seeing this to lie wide; we shall, in the first place, conclude the *Means* attending thereon, should do so likewise. Wherefore, although some may present themselves unto us as more promising; yet let us suppose what several Persons, were they hereunto engaged, each according to his Sense and Genius, would possibly make choice of. Believing, that although Considering Men may vary, in the approval of their own Sense and Notion; yet not always nearly, because it is their own; but because each, may probably see somewhat more in his own, than others do. Wherefore it will be our surest Logick to conclude, Not because no *Mean* may be approved by all Men, that all *Means* should be rejected; but rather, because each may be approved by some, that therefore, all be made choice of. And these, I think, may be comprehended under Five General *Heads* of Enquiry. *First*, Of those Things, which are of more *External* Consideration about *Plants*, as their *Figures*, &c. *Secondly*, Of their Compounding *Parts*, as *Vessels*, &c. *Thirdly*, Of their *Liquors*, and other *Contents*. *Fourthly*, Of their *Principles*, as *Salts*, &c. *Fifthly*, Of their *Aliment*, as *Water*, and other Means of Growth.

10. §. AND FIRST of all, whatever is of more *External* Con- The First
sideration, as the *Figures*, *Proportions*, *Motions*, *Seasons*, *Situations* of General
Vegetables, and of their several *Parts*, should be observed. In doing Mean.
which, a particular survey of all their Varieties should be taken. And then a Comparison made betwixt these, and the several *Plants*, or *Parts* of *Plants*, whereof they are the *Properties*. To the end, We may, if possible, be thereby conducted to find out, what other, either sensible, or more reclude *Property*, any of them may agree together in. For it is not more certain, that the three Angles of every *Rectilinear Triangle*, because all ways equal to two *Right Angles*, are therefore, if put together, always the same: than that *one Property*, agreeing to divers *Vegetables*, should have one *Cause*: For although the *Scope* and *End* may vary; yet the *Cause*, as it is the *Cause* of that *Property*, must be *one*: and consequently, must also import some *Identity* in the *Nature* of all those *Vegetables* wherein it Acts. Wherefore by thus comparing of them, we shall be able more exactly to state the *Orders* and *Degrees* of their *Affinities*; Better to understand both the *Causes* and *Ends* of their *Varieties*: And more probably to conjecture of their *Natures* and *Vertues*.

11. §. First then the various *Figures* of their several *Parts* should be observed; and that with respect both to the *Forms*, and the *Positions*, by which their *Roots*, *Trunks*, *Branches*, *Leaves*, *Flowers*, *Fruits*, and *Seeds* may vary, or agree; and those several *Lines*, by which both the said *Varieties* are determin'd. In which of these *Parts*, the agreement chiefly lies; this being both more observable, and more material in some of them; less in the *Root*, more in the *Flower*, or *Seed*. And in how many of these *Parts* together; whether one, more, or all. By both which, the *Orders* and *Degrees* of *Affinity*, which are many, may be accounted; either as to what we strictly call *Kin-dred*, or else *Analogy*. For there are found, not only *Herbs* account-ed of several *Tribes*, which are ally'd; and some of the Smallest, which are of kin to the Greatest: But there are also, probably, some *Herbs*, which have a particular Relation, to many Kinds of *Shrubs*; and some *Shrubs*, to many Kinds of *Trees*. Thus the several sorts of *Lettuce*, are of Kin, together in the *First Degree*; with *Endive*, in the *Second*. The several *Clarys*, amongst themselves in the *First*; with *Horehound*, in the *Second*; with *Lanium*, in the *Third*. All *Strawberries* agree together, in the *First Degree*; with *Cinquefoyl*, in the *Second*; with *Tormentil* in the *Third*; and with *Avens*, &c. in other *Degrees* more remote. So *Agrimony*, hath alike *Analogy* unto *Strawberry*; as *Goats-Rue*, hath to *Claver*: And *Strawberry*, the like unto the *Rasp*; as *Goos-berry* to the *Vine*; or *Burnet*, to the *Rose*. Amongst the several *Sorts* of *Grass*, there are some which match all those of *Corn*; which is but a greater kind of *Grass*. So again all *Pulse*, are not only of kin, in their several *Degrees*, to one another; but likewise, to almost all kinds of *Trefoyls*, as *Melilot*, *Fenugreek*, and the common *Clavers* themselves; as by comparing not only their *Leaves*, but *Flowers*, *Seeds*, and *Cods* together, may be evident. For the several *parts* of the *Flower* of a *Trefoyl*, are so many more *Flowers*, containing so many *Cods* of small *Seeds*, all, in shape, agreeable to the *Flowers*, *Cods*, and *Seeds* of *Pulse*. The same Relation, which *Trefoyls* have to the *Peas* or other *Pulse*; *Colts-foot*, hath to *Buttyr-Bur*; *Chickweed* to *Leucanthemum*; *Ground-sell*, to *Jacobæa*, or *Scorodonia*, to *Foxglove*: Or, to go higher, as the *Leguminous* Kinds of *Herbs*, have to *Sena*, or some other of the *Lobed Shrubs* and *Trees*. And, as among *Animals*, there are some which connect several Kinds; as the *Batt* doth *Beasts* and *Birds*: So, among *Plants*, there are some also, which seem to stand between two *Tribes*; as *Lappa*, between *Knapweeds* and *Thistles*; *Lampfana*, between the *Intybaceous* Kind, and the *Mouse-ears*.

12. §. From hence likewise, the *Natures* of *Plants* may be conjectured. For in looking upon divers *Plants*, though of different *Names* and *Kinds*; yet if some affinity may be found betwixt them, then the *Nature* of any one of them being well known, we have thence ground of conjecture, as to the *Nature* of all the rest. So that as every *Plant* may have somewhat of *Nature* individual to it self; so, as far as it obtaineth any *Visible Communities* with other *Plants*, so far, may it partake of *Common Nature* with those also. Thus the *Wild*, and *Garden Cucumers*, have this difference; that the one purgeth strongly, the other, not at all: yet in being *Diuretick*, they both agree. The *Natures* of *Umbelliferous* *Plants*, we know, are various; yet 'tis most probable, that they all agree in this one, scil. in being *Carminative*.

The

The several sorts, both of *Corn* and *Grass*, are all akin; there is no doubt therefore, but that the *Seeds* of *Grass* themselves (of *Rye* and *Oats* it is tryed) if it were worth the while to order them, as *Barley*, would yield an inflammable Spirit. So likewise the several Kinds of *Pulse*, have some one community in their Form, as is said: for which reason, I question not, but that in some Cases, wherein *Cicers* are esteemed a good *Medicine*; a *Decoction* of the better sort of *Pease*, especially that we call the *Sugar-Pease*, may go beyond them. As doth also the *Flower* or *Meal* of *Beans*, that of the *seeds* of *Fænugreek*; even there, where they are accounted excellent. So *Tulips*, *Lillies*, *Crocuses*, *Jacynths*, and *Onions* themselves, with many others, in their several Degrees, are all allied. If therefore *Crocuses*, *Onions*, *Lillies*, agree in one or more *Faculties*, then why may not all the rest? as in being *Anodyne*; or in some other *Common Nature*; whereby, in their *Vegetation*, their *Parts* are Governed and Over-ruled, to one *Common* or *Analogous Form*.

13. §. The *Proportions* likewise, amongst the several *Parts* of *Vegetables*, for the same Reasons, deserve to be observed; the comparison being made, both betwixt the *Parts* of several *Plants*, and the several *Parts* of one. And here again, either betwixt any *Two* of the *Parts*, or any *One* of them, and the *Whole* besides, or all the rest put together. So some larger *seeds*, produce a small *Root*; as those of *Cucumber*: and others smaller, produce one very great; as those of *Bryony*. Some *Plants*, as the *Melon*, though themselves but very slender, yet have a vast and bulky *Fruit*; others again, as *Thistles*, and many yet more substantial, have no other *Fruit*, besides their *Seed*. So the *Seeds* of all *Pulse*, and especially, the *Garden Bean*, though large, yet produce but a small *Plant*: but those of *Foxglove*, *Mullen*, *Burdock*, *Sun-flower*, &c. being themselves much less, do yet produce a far greater. And especially, those *Seeds*, which are inclosed in the Thicker sort of *Cover*, (analogous to that I have elsewhere called the *Secondine*) as that of *Peony*; whose *seed*, so called, is only the *Nest* wherein the true and real *Seed* is lodged, no bigger than a little *Pins head*: which is also observable of the *Seeds* of divers other *Plants*. These, and the like *Proportions*, as they lie betwixt the several *Parts*, should be noted: and to what *Plants* or *Parts* especially, any of them may agree: comparing also in what other kind of *Properties* an agreement betwixt the said *Parts* may be found: that so doing, we may, if possible, amongst all their *Individual Natures*, be instructed to single out those *Common Ones*, which are concomitant to such Agreeing *Properties*.

Anat. Plant.
Book I. Chap.
ult.

14. §. The several *seasons* also of *Plants*, and of their *Parts*, should be considered. Observing at what particular *Times* of the *Year*, any of them chiefly *Spring*, *Early* or *Late*. The *Times* wherein they *Germinate*; whether for some *Space* only, or all the *Year* long. Wherein they *Spring*, after *Sowing*; or *Flower*, after *Springing*, sooner, or later. Which *Flower*, the *first* *Year*, or not till the *second*. Which after the *Leaves* are put forth, or before them; for so, some do, as the *Crocus Vernus*, *Bears-foot*, *Hepatica aurea*, and others; all the *Leaves*, at the time of their flowering, being old, or of the foregoing *Year's* growth. So likewise the *Maturation* of the *Fruit* or *Seed*; how long after the *Flower*, and the like. All or some of which *Varieties*, being laid

laid together, we may probably conjecture the *Causes* thereof; and the *Natures* of the *Plants* in which they are seen: *scil.* as such a Degree of Heat may be necessary for the Fermentation, or the better Distribution of the *sap* of such a *Plant*; or for the Impregnation of the *Aer*, to be mixed therewith; or the due Disposing of the *soil*, to render the most convenient Aliment thereunto. So the *Principles* of such *Plants*, which flower all the Year, may be more equally proportion'd. Those which flower before the *Leaves* put forth, as the *Crocus Vernus*, and those which flower in *Spring*, may be accounted *Rank*, and full of *Volatile salt*. But *Autumn Plants* especially, to abound with a *Fixed*: and the like.

15. §. The proper *Places* also of *Plants*, or such wherein they have, from their *Seeds*, or other way of *Propagation*, a Spontaneous growth, should be considered. And that as to the *Climate*; whether in one Colder, Temperate, or more Hot. The *Region*; Continent, or Island. The *seat*; as Sea, or Land, Watry, Boggý, or Dry; Hills, Plains, or Vallies; Open, in Woods, or under Hedges; Against *Walls*, rooted in them, or on their *Tops*: and the like. And perhaps the *Seeds* of some *Plants*, as of *Mosses*, (which, through their smallness, will ascend like Moths in the Sun) may fly or swim for some time, in the *Aer*, *viz.* till they begin to shoot, and so become heavy enough, to fall down upon the *Ground*. From whence, in like manner, as from their *Seasons*, their particular *Natures* may be directed unto. In that, so far as we may conjecture the nature of such an *Aer*, *Soil*, or *Seat*, we may also of such a *Plant*, to which they are *congenial*.

16. §. So likewise, those many Varieties observable in the *Motions* of *Plants*, and of their *Parts*, both *Kinds* and *Degrees*; *Ascending*, *Descending*, and *Horizontal*; *Rectilinear*, and *Spiral Motions*, should be noted; to what *Plants* they agree, and wherein any of these *Motions* may be analogous to those of *Animals*. And in a word, any other *Forensick Properties* of *Plants*. And then, to Compare them all together; both being necessary. For *Thoughts* cannot work upon nothing, no more than *Hands*. He that will build an House, must provide *Materials*. And on the contrary, the *Materials* will never become an House, unless, by certain *Rules*, we joyn them all together. So, it is not, *simply*, the Knowledge of *many things*, but a multifarious Copulation of them in the *Mind*, that becomes prolifick of further Knowledge. And thus much for the first *General Mean*.

The Second
General
Mean.

17. §. THE NEXT which I propose, and that a most necessary one, is *Anatomy*. For when upon the Dissection of *Vegetables*, we see so great a difference in them, that not only their *Outward Figures*, but also their *Inward Structure*, is so *Elegant*; and in all, so *Variouſ*; it must needs lead us thus to Think, That these *Inward Varieties*, were either to no *End*; or if they were, we must assign to what. To imagine the first, were exceeding vain; as if *Nature*, the Handmaid of *Divine Wisdom*, should with Her fine *Needle* and *Thred*, stitch up so many several *Pieces*, of so difficult, and yet so groundless a *Work*. But if for some *End*, then either only to be looked upon, or some other besides. If for this only, then this must be such as in respect whereof, Her *Work* is at no time, nor in any degree frustrate; the contrary whereunto, is most manifest. For although Men do every where, with frequent pleasure, behold the *Outward Elegancies* of *Plants*;

Plants; yet the Inward Ones, which, generally, are as Precise and Various as the Outward; we see, how usual it is, for the beholding of These, to be omitted by them. And besides, when we have observed Nature's Work, as well as we can; it may be no impediment to our best Endeavours, to believe, That some Parts of it, will still remain behind, *Unseen*. So that if to be *Seen*, were the only End of it, it must needs be wholly frustrate, as to the greater number of Men; and, in some part, as to all. Wherefore, we must suppose some other *Ends* of the said Varieties, which should have their *Effect*, and so These, not be in vain, whether Men beheld them or not; which, are, therefore, such as have respect to *Vegetation*: That the *Corn* might grow, *so*; and the *Flower*, *so*, whether or no Men had a mind, leisure, or ability, to understand *how*.

18. §. If then the *Anatomy of Vegetables* be so useful a *Mean*, we ought not to streighten it; but to force this, as well as the rest, to its utmost Extent. And therefore, first of all, To go through all the *Parts*, with equal care; examining the *Root, Trunk, Branch, Leaf, Flower, Fruit, and Seed*. Then to Repeat or Retrograde the Dissection, from *Part to Part*: in that, although the best Method of Delivery, for clear Discourse, can be but one, according to that of *Nature*, from the *seed* forward, to the *Seed*: yet can it not but be useful, for That of Dissection, to proceed *to* and *fro*; somewhat or other being more Visible in each several *Part*, from whence still an Hint may be taken, for the ushering in the observation of it in the others. To examine, again, not only all the *Parts*, but *Kinds of Vegetables*, and comparatively, to observe divers of the same *size, shape, motion, age, sap, quality, power*, or any other way the *same*, which may also agree, in some one or more particulars, as to their *Interiour Structure*: and to make this comparison, throughout all their *Parts and Properties*. To observe them likewise, in several *Seasons* of the Year, and in several *Ages* of the *Plants*, and of their *Parts*; in both which, divers of them may be noted to change, not only their *Dimensions*, but their *Natures* also; as *Vessels*, do into *Ligaments*; and *Cartilages*, into *Bones*, sometimes, in *Animals*. And to do all this by several *Ways of section*, Oblique, Perpendicular, and Transverse; all three being requisite, if not to Observe, yet the better to Comprehend, some Things. And it will be convenient sometimes to Break, Tear, or otherwise Divide, without a *Section*. Together with the *Knife* it will be necessary to joyn the *Microscope*; and to examine all the *Parts*, and every Way, in the use of That. As also; that both Immediate, and Microscopical Inspections, be Compared: since it is certain, That some things, may be demonstrated by Reason and the Eye conjunct, without a *Glass*, which cannot be discovered by it; or else the discovery is so dark, as which, alone, may not be safely depended on.

19. §. By these several Ways of Inspection, it will be requisite, To observe their *Compounding Parts*; as *Simply* considered, and as variously *Proportioned*, and *Disposed*. As *Simply* considered, to note their *Number*; what, and whether the same, in all: their *Kinds*, wherein different in the same, or divers *Vegetables*: their *Original*, in part, or in whole: *Structure*, as to their *Contexture* and their *Cavities*; Their *Contexture*, within themselves severally, and as joyned together: their *Cavities*, as to their *Size, Shape, and Number*; in which a great variety

riety will be found. Next their *Positions* one amongst another, which are also various; as Anterior, Posterior, Collateral, Surrounding, Mediate, Immediate, Near, Remote; both as they respect the several *Parts*, and the several portions of one: And all these, as few, or more; these or others of them, may be diversly Compounded together. And then the *Proportions* they bear one to another; whether as to Minority, Equality, or Excess; each *Part* compared with each, and that as to the several Degrees appearing in the said *Proportions*; the Varieties whereof may be exceeding numerous. For if we should suppose but *Four* considerable *Parts* generally constitutive of a *Vegetable*: These *Four*, produce a Variety *Four* ways. First, when One is Unequal; and then it produceth only *Four* Varieties: and those two ways, *scil.* when one is Greater, and the other three, Equal and Less; or when one is Less; and the other three, Equal and Greater. Secondly, when Two be Unequal; and then they produce *Six* Varieties. Thirdly, when Three be Unequal, which produceth *Twelve* Varieties. Or lastly, when all Four be Unequal; which produceth *Twenty four*: which general Varieties, may be further multiplied by their several Degrees.

20. §. From all, which, we may come to know, what the *Communities* of *Vegetables* are, as belonging to all; what their *Distinctions*, to such a Kind; their *Properties*, to such a Species; and their *Peculiarities*, to such Particular ones. And as in *Metaphysical*, or other Contemplative Matters, when we have a distinct knowledge of the *Communities* and *Differences* of Things, we may then be able to give their true *Definitions*: so may we possibly, here attain, to do likewise: not only to know, That every *Plant* Inwardly differs from another, but also wherein; so as not more surely to Define by the Outward *Figure*, than by the Inward *Structure*, What that is, or those things are, whereby any *Plant*, or Sort of *Plants*, may be distinguished from all others. And having obtained a knowledge of the *Communities* and *Differences* amongst the *Parts* of *Vegetables*; it may conduct us through a *Series* of more facile and probable *Conclusions*, of the ways of their *Causality*, as to the *Communities* and *Differences* of *Vegetation*. And thus much for the Second General Mean.

The Third
General
Mean.

21. §. HAVING THUS far examined the *Organical* and *Containing Parts* of *Vegetables*; it will be requisite, more designedly, to observe those also which are *Fluid*, or any others Contained in them: and that, for our better understanding both of the *Nature* of *Vegetation*, and of the said *Contained Parts*. And to make inquiry, *First* of their *Kinds*; as *Spirits*; both such as agree, in general, in being *Vinous*; and those that are Special, to particular *Plants*. *Aers* and *Vapours*; for the existence whereof, in all *Vegetables*, there are Arguments certainly concluding. And for the difference of their *Natures*, in being more dry, or moist, more simple or compounded, as they are existent in several *Parts*, there are probable ones. *Lympha's* or clear and watry *Saps*; which most *Plants*, in one *Part* or other, at some time of the Year, do Bleed *Mucilages*; as in *Mallow* and *Violet Leaves*; in many *Seeds*, as of *Quinces*, *Clary*; *Fruits*, as in *Cucumers*; distinct from the watry *Sap*, as by permitting it to stand and gelly upon the *Vessels* from whence it issues, is plain: And in the young *Berrys* of *White Bryony*, when about the bigness of a *Peppercorn*;

Corn; the juyce whereof is so Viscous, that the twentjeth part of a *Grain*, will draw out above a *Yard* in length. *Oyles*; not only in *seeds*, and some *Fruits*, but other *Parts*; as in certain little cavities in the *Leaves* of *Savine*, visibly collected while they are growing. *Gumms* or *Resines*; as in *Pine*, *Fir*, and others of this Kind. *Milks*; as in a vast number of *Plants*, and amongst them, many not suspected to yield any. For, of *Herbs*, not only most of the *Umbelliferous Kind*, are *Milky*; but all or most of the *Intybous*; *Poppys*; *Tracheliums*; *Perwinkles*; divers *Thistles*; and even *Onions*, if cut at the bottome; with a great many more. Of *Trees*, not only the *Little Maple*, but the young *shoots* of *Lawrel*, especially being crushed; as also those of *Elder*, and some others. To which may be added, such *Mucilages*, which though not so properly contained *within* the *Parts*, yet are found lying *over* them; as over the first *Spring-leaves* of all kinds of *Docks*; betwixt the *Leaves* and the *Veil* wherein they are involved. That fine white *Flower* or *Powder*, which lies over the *Leaves* of some *Plants*, as of *Bears-Ear*: And in *Princes-Feather*, about certain *Aper-tures* only on the edges of the *Leaves*.

22. §. Of all these should be observed, first their *Receptacles*; some of them, being proper to one; others, common to two or more of them: since it is certain, that some of them do *Transmigrate* from one, into another *Receptacle*, or that the same *Receptacle* is filled with *Fluid Bodies*, of a quite different *Nature*, at the different *Seasons* of the *Year*, and *Ages* of the *Vegetable*. And it is also very probable, That two of some of them, may, sometimes, be contained in one *Receptacle*, at the same time; as in *Animals*, the *Lympha* in the *D. Thoracicus*, and that, and the *Chyle*, in the *Sanguineous Vessels*.

23. §. Then their *Motions*; both *Natural*, and such as may be effected by *Art*: and those either by *Descent* or *Ascent*; And in ascending, through what different *Channels* or *Parts* of the *Trunk*; since it is certain, That there is a variety, both in respect of the *Season*, and of *Vegetables*. Where it will fall in, To observe the *Tapping* of *Trees*. As also their *Bleeding*: to what *Trees* it is proper to *bleed*: in those to which it is, with what difference of *Celerity*: and when their peculiar *Season*: for none will bleed at all times; neither will all *bleed* at the same. And then their *Collateral Motion*, together with the *Mode* of their *Transition* from one *Organical Part* to another.

24. §. Next their *Quantities*, either of *one*; as the *Comparison* is made betwixt several *Plants*, or betwixt the *Parts* of the same. So the true *Seed* of all *Plants*, containeth more *Oyl*, in proportion, than any of the other *Parts*. Or else of *divers*, as coexistent and bearing such a proportion one to another in the same *Part*: of most of which, it may be known by their respective *Receptacles*. Yet the *Computati-on* must not be made from the number of the said *Receptacles*, *simply*; but as that is in conjunction with their *Capacity*; and as their *Capacity* is proportioned to their surrounding *Sides*; the *Sides* of those of the least *Capacity*, being usually as thick, as those of the greatest: so that suppose *Ten* lesser, to lye within the compass of *One* greater; the *Con-tent* of these altogether, would scarce be equal to half the *Content* of that *One*.

25. §. Also their *Consistence*; *scil*: of so many of them as are discriminable by Touch; in being Soft or Hard; Thin or Thick; Mucilaginous, Gummous, Glutinous, Friable, &c. And these in their several Degrees; in which there is a Variety, as in the *Milks* of some *Plants*, which are more *Dilute*, than that of others: *Mucilages*; which in some, are very thick and *Viscous*, in others, more diluted and coming nearer to a *watry Sap*. And by This, to be compared in the same manner, as by their *Quantity*.

26. §. Likewise their *Colours*, *Smells*, and *Tastes*: The general and particular Kinds of all which should be noted. And to what *Contained Parts*, and in what Variety, they appertain. So most *Resinous Gums* are Tinctur'd, some, not; as that which drops from the *Dome-stick Pine*, is as clear as *Rock-water*. The *Milks* of some *Plants* are *Paler*, as in *Burdock*; of others *Whiter*, as in *Dandelyon*, *Scorzoner*; *Citrine*, as in the Root of *Trachelium*, *Angelica*; *Yellow*, as in *Lovage*. In some *Plants*, *Odorous*, as in *Umbelliferous*; in others not, as in *Cichoraceous*. That of *Little Maple*, *Tasteless*; of *Garden Chervil*, *Sweet*; of *Fenil*, *Hot*; of *Scorzoner*, *Astringent*; of *Dandelion*, *Bitter*; and generally, in other *Plants*; but with many Degrees of Strength, and in conjunction with other *Tastes*. But most *Mucilages*, have little either *Colour*, *Taste*, or *Smell*; and the like. Here also the same *Qualities* are to be inquired into, as, in general speaking, they are said to belong to a *Vegetable*. Since it is more than probable, that all *Colours* (excepting *White*, which is sometimes common both to *Containing* and *Contained Parts*) all *Odours*, and *Tastes*, which are more immediately, and without a resolution of their *Essential Principles*, perceptible in a *Plant*; are not ascribable either to the *Organical*, or *Containing Parts*; but only to *Those*, *Contained* in them; as from divers reasons hereafter may appear.

27. §. And first, their *Colours*; where, with respect to several *Plants* and *Parts*, they are more *Changeable*; as *Red*, in *Flowers*; or *Constant*, as *Green*, in *Leaves*. Which, with respect to several *Ages* of one *part*, are more *fading*, as *Green* in *Fruits*; or *durable*, as *Yellow* in *Flowers*. In what *Parts* more *Single*, as always in the *Seed*; or more *Compounded*, as in the *Flower*; and in what *Plants* more especially, as in *Pancy*. Which proper to *Plants* that have such a *Taste* or *Smell*, as both, in *White Flowers*, are usually less strong. To *Plants* that flower in such a *Season*, as a *Yellow Flower*, I think, chiefly, to *Spring Plants*. And to *Plants* that are natural to such a *Soil* or *Seat*, as to *Water-plants*, more usually, a *white Flower*. What, amongst all *Colours*, more *Common* to *Plants*, as *Green*; or more *Rare*, as *Black*. And what all these *Varieties* of *Colours* are upon *Cultivation*, but chiefly, in their natural *soil*. To observe also with their superficial *Colours*, those within: so the *Roots* of *Docks*, are *Yellow*; of *Bistort*, *Red*; of *Avens*, *Purple*; but of most, *White*. Where the *Inward*, and *Superficial Colours* agree; as in the *Leaves*; or vary, as in the other *Parts* frequently. And in what manner they are *Situated*; some universally spreading, others running only along with the *Vessels*, as in the *Leaves* of *Red Dock*, and the *Flowers* of *Wood-Sorrel*.

28. §. Next their *Odours*; what may be their principal *Seat*; whether one or divers *seats* in the same *Plant*. What the chief *Matter* out of which they are continually bred. What similitude betwixt the
the

the *Smells* of divers *Vegetables*; as betwixt *Baume*, and a *Limon*; the *Green Leaves* of *Meadow-sweet*, and the *green Rinds* of *Walnuts*. Or betwixt those of *Plants* and *Animals*; as the *Smell* of green and well-grown *Carduus*, is like to that rank *scent*, *ab aliquorum axillis spiranti*. Which have a more sensible *Smell*; as most have; and which have less, as *Corn*. Where the *green Leaf* is the most *Fragrant Part*, as in *Musk-Crabsbill*; where the *Flower*, as in *Roses*; the *Root*, as in sweet *Calamus*. Where all the *Parts* have some *Odour*, where some, or one, only; as in *Scurvy-grass*, only the *Flowers*, unless the *Leaves* are bruised; and in *Arum*, the *Pestil* only; for neither the *Leaf*, nor *Root* hath any *Smell*, unless cut; but this is strong enough, not much unlike to *Humane Excrements*.

29. §. But especially their *Tastes*, which it much importeth us more precisely to distinguish; *First*, by their general *Kinds*; for the number, even of these, may be computed greater than usually it is. I remember not, that *Heat* and *Acritude*, with respect to *Taste*, are distinguished; yet *Arum-Root* is very *Pungent*, without any proper *Heat*; and *Cloves*, are very *Hot*, without any proper *Pungency*. So the *White Roots* of *Yarrow*, have a *Taste*, hardly any other way perceptible, than by causing a gentle *glowing* and *continued Warmth* upon the *Tongue*. Also their *Respondencies* one to another; as that of *Zedoary*, and of the lesser *Cardamoms*, is somewhat like to *Camphire*. Likewise their *Degrees*; in which there is a great latitude, and may be extended from *One* to *Ten*, or with easie distinction, from *One* to *Five*: So the *Root* of *Sorrel*, is *Bitter* in the *first*; of *Dock*, in the *second*; of *Dog-Rose*, in the *third*; of *Dandelyon*, in the *fourth*; of *Gentian*, in the *fifth*: observing them, not only as they vary in several *Kinds* of *Plants*, but the several *species* of one, as in *Cichory*, *Hawk-weed*, *Dandelyon*. And then their *Compositions*; for *Tastes* are as truly *conjunct* in one *Part*, as *Colours*: by which, the latitude is still greater; In that all *Kinds* of *Tastes*, in all their *Degrees*, and in differing *Numbers*, may be variously *Compounded* together: For the most part, *Two*, as in the *Leaves* of *sharp-pointed Dock*, *Astringent*, and *Sowre*; in *Sorrel-Roots*, *Astringent* and *Bitter*; and in *Aloes*, *Bitter* and *Sweet*; the one in the *fifth*, the other, in the *first Degree*; as upon an unprejudiced tryal may be perceived: and yet more evidently in the *Gall* of any *Land-Animal*. Sometimes *three*, as in *Agrimony*, *Bitter*, *Rough*, and *Sowrish*; and in *Agarick*, *Bitter*, *Rough*, and *Sweet*. And sometimes, perhaps more. The *Sensible* distinctions of all which, may lye almost as wide, as of *Plants* themselves. Wherefore, although it may be thought rashness, to take away the distinctions of *Hot*, *Cold*, *Moist*, *Dry*, *Thin*, *Gross*, and other *Qualities*, in their several *Degree*, which the *Ancients* have affixed to particular *Plants*: yet since they have done it, to many of them, with much uncertainty; and that, withal, they are, more properly, the *Effects* and *Operations* of *Plants*, than their *Qualities*; *Practical Observation*, may therefore approve it useful, to add these *Sensible* Ones of various *Tastes*, precisely distinguishing their *Conjugations* and *Degrees*. Lastly, their several *Varieties* and *Mutations*, with respect to the *Subject* wherein they reside, should also be noted, As, of all *Tastes* found in *Plants*, *Bitter* and *Sowr*, are most common; *Sweet* and *Salt*, most rare. Which latter, is not only perceptible in some *Sea-Plants*; but upon some others, as upon the fresh

Leaves of *Tamarisk*; which being licked while they grow, or when immediately gathered, are plainly *saltish*. How they vary with the *Age* of the *Plant*, or *Part*; as the *Roots* of *Radishes*, growing up to *Seed*, lose the strength of their *Tast*; so most *Fruits* are first *Sowre*, then *Sweet*. What proper to the several *Parts* of any one *Plant*; so the *Leaves* of *Wormwood* are extraordinary *Bitter*; the *Root* scarcely so at all; of an *Hot*, but quite different *Taste*. What more *Common*, or *Rare*, to any *Part*; so no *Root* that I ever tasted, is *Sowre*. And how they *Alternate* in several *Plants*; as the *Root* of *Stock-July-flower* is biting, not the *Leaves*; on the contrary, the *Leaves* of the *Water-Arsmart*, are *Biting*; but not the *Root*; and the like. To which we may add the difference of *Time* wherein the *Tastes* of *Plants* are perceived; as those of *Arum*, and *Rape-Crowfoot*, are both *Biting*; but that of the first, as it is slowly perceived, so it continues long; that of the other, quickly comes, and quickly goes.

30. §. Amongst the other *Adjuncts* of the *Contained Parts*, though not of these only, the *Faculties* of *Vegetables* are to be reputed. For so the *Resin* of *Jalap*, which is *Purgative*, is as truly contained in the *Organical Parts* of that *Root*, as *Blood* is in *Veins*: It will be requisite therefore to make particular observation of these also. And first, what *Faculties* chiefly may reside in *Plants*, above others: so there is none of known use in *Salivation*, except by holding in the mouth: Although we may ask, Why some amongst them, may not (being Taken inwardly) have a power to evacuate by This, as well as other *Violent* ways? Where the *Faculty* is more universally spread over all the *Parts* of a *Vegetable*, as in *Asarum*. Where belonging chiefly or wholly to any particular *Parts* or *Part*; as chiefly to the *Root* of *Rhubarb*; and only to the true and proper *Seed* of *Barbado Nuts*. Whether some *Faculties*, may be proper to some *Parts* especially. What conjunction they may have with any sensible *Qualities*. So, many *Purgers*, are not only *Resinous* and *Gummous*; But also *Mucilaginous*; as *Bryony*, wild *Cucumber*, *Lapathum Sativum*; and therefore probably *Rhubarb*, when growing; *Mallows*, *Violets*, &c. Such as are *Purging* and *Vomitory*, though some of them have a strong *Taste*, yet the greater part, and of those, many of the stronger sort, have no *Taste*, or not *Great*; as *senna*, *Jalap*, *Scammony*, *Hellebore*, *Asarum*, and others. Amongst which, although *Hellebore* hath a very *Durable Taste*, yet is it not very *High* or *Great*. So also, those that are most sensibly tasted, are, I think, for the most part, more or less *Bitter*; either simply, as *Colocynthis*; or *Bitter* and *Astringent*, as *Rhubarb*; or *Bitter* and *Sweet*, as *Aloe*; or *Bitter*, *Astringent*, and *Sweet*, as *Agarick*. Few are *Hot*, as *Iris*. Or simply *Sweet*. And though some may be *Subacid*, that are *Mollifying* or *Lenitive*, yet no proper *Purge* or *Vomit* is *Sowre*. Such *Plants* as are of a soft and sweetish *Taste*, without *Viscosity*, may be accounted good *Antiscorbuticks*, especially against the *Sea*, or other *Salt-Scurvey*; as are good sweet *Pease*: And sometimes the *Water* or *Spirit* of the *Shells*; which may easily be drawn from them, being first duly fermented, and hath a true *Vinous Taste*; but very mild, and not unpleasant. Those *Plants*, whose *Parts* are not only *Hot* but *Volatile*, as *Onions*, are generally good for *Burns*. Such as have a *Balsamick Taste* or *smell*, with a little *Astringency*, as *Hypericum*, *Golden-Rod*, *Lanium Luteum*, &c. the best *Wound-Herbs*. And such as are gently *Bitter*,

Bitter, and *Penetrant* upon the *Tongue*, or in the *Throat*, as *Daisy*, *Anagallis*, good *Cleansers*. That such *Bodys*, principally, are *Anodyne*, which are *Yellow*, I think, is more than a conceit; *Yelks of Eggs*, *Fœnugreek Seeds*, *Lint-feed Oyl*, *May-Butyr*, *Marrow*, *Pinguedo Humana*, *Hyoscyamus luteus*, *Safron*, *Sulphur*, *Opium*, all *Anodyne* and *Yellow*. How likewise their *Faculties* and *Qualities* may vary their *Degrees*, either differently or together: so *Aloe* and *Colocynthis*, are both *Bitter* in the highest *Degree*; yet *Aloe*, which is also *sweet*, Purgeth more moderately; *Colocynthis*, which is *Bitter*, but not *Sweet*, molt Violently. How far the *Faculties* of *Vegetables*, as well as their *Qualities*, may be *Compounded*; where, and which chiefly; as *Astringive* and *Purgative* in *Rhabarb*. Where this *Question* may be put, Whether divers other, and yet more extreme *Faculties*, as well as these of *Astringive* and *Purgative*, may not somewhere or other be also found, or made, to meet: whereby the same *Plant*, or some *Preparation* of it, may be most *Potent*, and yet most *Innocent*; the *Malignity* thereof exerting its *Power*, and the *Virtue* its *Sovereignty* at the same time. And lastly, what *Affinity* there may be betwixt them; as most *Plants*, that are strong *Purgatives*, and especially *Vomitories*, I think, are also *Sternutatory*; as white *Hellebore*, *Jalap*, *Tobacco*: and on the contrary, such as are *Sternutatory*, are some of the most proper and most potent *Medicines* for the *Head*, *Brain*, and *Genus Nervosum*, Taken inwardly, as *Lilium convalle*, &c. and the like.

31. §. Thus far a particular observation of the *Qualities* and *Faculties* of the *Contents* of *Vegetables* may proceed, as they are existent in their *Natural Estate*. From which, although some probable *Conjectures* may be made, of their *Material* and *Formal Essences*, and of the *Causes* of their determinate *Varieties*, or the *Modes* of *Vegetation* necessary thereunto: yet will our *Conceptions* hereof be more facile, clear, and comprehensive, if by all other *Ways* of *Observation*, they be likewise examined, according as *Experiment* may be applicable to any of them.

32. §. As by *Contusion*; so some *Plants* give their *Smell*, not without *Rubbing*, or not so well; as the green *Leaves* of *Stramonium*, *Scurvygrass*, and many more: others lose it by *Rubbing*, as the *flowers* of *Violets*, *Carnations*, *Borage*, &c. others yield it both ways, as *Rosemary*, &c. So some *Apples* mend their *Taste*, by *Scoaping*, and *Pears* by *Rowling*, especially that called the *Rowling Pear*.

33. §. By *Agitation*; which doth that, sometimes, by *Force*, which *Digestion*, doth by *Heat*: so any cold *Oyl* and a *Syrup* being, in a due manner, agitated together, of two *Fluid bodies* will become one *Consistent*, as is known.

34. §. By *Frigifaction*; how far the *Juyces* of *Plants*, either without or within them, may be any of them, or some more than others, subject to *Cold*: and thereby to be deprived of their *Motion* or natural *Consistence*, or may suffer alteration in their *Colour*, *Taste*, or *Smell*.

35. §. By *Infusion*; where I mean *Infusion* only in *Common Water*; So both *Cassia Lignea*, and *Cinnamon* are a little *Mucilaginous*; but the former most. Some of the *Contents* of *Plants*, may be wholly dissolved in *Common Water*; some but in part, others not at all; or very little; which is proper to some *Milks*, as well as *Gums*. The *Colours*, *Smells* or *Tastes* they hereupon yield, are found various; and in some

some very unexpected: So the green *Leaves* of *Bawm*, being duly infused in common Water, without any other Body added, tincture it with a clear and deep Red, near that of *Claret Wine*, as I have often tryed.

36. §. By *Subsiding*; So the *Juyce* of *Sorrel*, being ordered as that of *Grapes*, will, in time, let fall a kind of *Tartar* or *Essential Salt*. And so perhaps will that of many other *Plants*, without any previous *Decoction*; although that be commonly thought to be necessary.

37. §. By *Digestion* with *Fermentation*; either of the entire *Vegetables*, or of the *Juyces*, or other *Contents*; and these by themselves, or with common Water. And hereby to note, what difference may be in the *Strength*, *Celerity*, or *Continuance* of the *Fermentation*. Likewise, how their *Qualities* may thereby be altered; as the *Smell* of *Violet-flowers*, from a most excellent *Fragrancy*, may, by *Digestion*, be reduced to an odious and abominable *Stink*, like that of the black *Mud* of *Gutters*.

38. §. By *Digestion* with *Calefaction*; so the *Colour* of the *Juyce* of *Lemons*; from *Transparency* (if that be a *Colour*) may be turned to a perfect *Red*. Whence it is that many are deceived in the *Preparation* called the *Tincture* of *Corals*; supposing the *Corals* to give the *Menstruum* its *Colour*. Whereas the *Menstruum* will obtain it, only by *Digestion*, without any *Corals*, mixed with it.

39. §. By *Decoction*; either of *Vegetables* themselves, or of their *Liquors*; and to observe what alterations follow. So *Turpentine* boiled becometh friable; *Sugar*, *Bitter*, and of a *Brown Red*. *Turneps* lose their *Biting Taste*; *Onions*, their *Picquancy*; yet neither of them convey those self same *Qualities* to the *Water*. The same may be observed in the *Decoction* of *sweet-Fennel-seeds*, *Aniseeds*, and others, losing much of their *Tastes* themselves, and yet conveying very little of them to the *Liquors* wherein they are boiled; the greater portion of their *Volatile parts*, and so their *Virtue* and *Taste* therewith, flying away. Whereof therefore it is much better to make an *Emulsion*, than to decoct them; or to make an *Emulsion* from them, with their own *Decoction*, especially if the *Medicine* be intended to be *Carminative*, as I have frequently observed. The *Decoction* should also be carried on throughout all degrees to that of an *Extract*; by which the *Qualities* thereof, sometimes, are much altered; as the *Colour* of all or most green *Leaves*, from a kind of *Yellow*, deepens at last into a dark one, as *Black* as *Pitch*.

40. §. By *Distillations*; both with the cold *Still*, *Alembick*, *Chappel-* and open *Furnace*: and to note what *Vegetables* thus give their *Smell* or *Taste*, and in what *Degrees* of strength, either under, or over their natural ones; as *Mint*, *Pennyroyal*, and the like, which are *Aromatick* and *Hot*, give their *Tastes* perfect: but *Wormwood*, which is *Aromatick* and *Bitter*; gives it but by halves, pretty fully as *Aromatick*, little as *Bitter*. And *Carduus*, though also so exceeding *Bitter*, yet not being *Aromatick*, yieldeth a much weaker *Taste*. Also what *Vegetables* yield *Oyl* most plentifully; and what difference may be in those *Oyls*, as to their *Colour*, *Weight*, or otherwise; as that of *Cloves* is sometimes *Red*; of *Cinnamon*, limpid; both *Ponderous*. So to distil *Juices*, *Gums*, or other *Contents*, with an *hot fire*; and to see, what *Bodies* they yield, and of what *Qualities*; as *Turpentine* is known to yield, besides its

its Oyl, a subacid Water; Vinegar, an Eager Spirit; as that part may be called, which Chymists are wont to call the Phlegm.

41. §. By *Arefaction*; so *Milks* which are Liquid, and White in their Natural Estate, in Standing, grow Gummous, Yellow, and otherwise different, so doth that of *Scorzouera*; and that of *Fenil* becomes a Balsamical, but Limpid Oyl. The *Roots* of *Angelica*, being dry'd, and cut by the length, exhibit their small *Veins* fill'd with an Aromatick *Rosin*. In the whiter parts of *Rhubarb*, is gathered a kind of *Saline Concret*; by which, this *Root*, in chewing, seems as if it were a little gritty. *Cabbage-stalks*, sliced, and laid in the Shade to dry, gather on them a kind of *Nitrous Hoar*. *Raisins* and *Corins* contain, not only a sweet Juyce, but also a true *Sugar*, which lies *curdled* in the *Pulp*, as the more *Saline* parts do in *Green Soap*. And the like is gather'd on the out-side of a *Fig*; saving, that it is more *Nitrous*, as lying next the *Aer*. The *Roots* of *Arum*, upon drying, lose much of the strength of their *Taste*; but the contrary may be noted of many other *Roots*, which, upon drying, increase it. Some, being cut and laid by, change their *Natural Colours*, into Red, Purple, Yellow, Green, or White; as *Liquorish*, into White, in some places; and *Peony*, into Red: and sometimes into two; as *Patience*, into Yellow and Red.

42. §. By *Affation*; thus *Apples*, by roasting, eat more Sowre. The *Root* of *Horse-Radish*, toasted, tasteth like a *Turnep*. *Potatoes*, *Onions*, and many other *Roots*, and *Parts*, have their *Tastes*, either Altered or Refracted; which chiefly, and in what manner, should be observed. There is one alteration, as remarkable, as commonly known; and is that which followeth upon roasting or baking in one kind of the *Waldensian Pears*, which, for a *Walden*, we corruptly call a *Warden*.

43. §. By *Ustion*; wherein some *Plants*, or *Parts* of them, burn very quietly; others, not without violent motions; so *Fenil-Seeds*, held in the flame of a *Candle*, will spit and spurtle, like the *serum* of *Blood*. Some *Vegetables* lose their *Smell*, as *Roses*; others, keep it, as *Rosemary*; and others, mend it, as *Lignum Aloes*. To note, not only the alteration of their *Qualities*, but what they yield; as *Turpentine*, which, in *Distillation*, yieldeth *Oyl* and *Water*, both limpid; upon *Ustion*, sheweth nothing but a black *soot*. So *Benzoine*, by *Distillation*, *Oyl*; by *Ustion*, white *Flowers*, as is known.

44. §. By *Calcination*; and here to observe, wherein the *Caput Mortuum* of one, may differ from, or agree in Nature with that of another; and also to compare these with those of *Animal Bodies*. As also in their *Quantities*. And to compare them with what they yield by *Distillation* and *Ustion* as to both. Thus far they have been tryed singly, or by themselves. They should also be examined,

45. §. By *Composition*; not only with *Water*, as in simple *Infusions*, &c. but with any other *Bodies*, which may have a power of acting upon them, or upon which, these may have a power to act. And so to make *Infusions*, *Destillations*, *Decoctions*, *Digestions*, in divers kinds of *Liquors*, as *Vinegar*, *Urine*, *Spirit of H. H. Wine*, *Blood*, *Milk*, or others. So in *Infusions*, some *Red Colours* are heightned by *Acids*; *Blews*, turned *Purple*. So fetid *Spirits* (as of *H. H.*) may be rendred much more grateful, by being *Rectified*, once or twice, with fresh *Aromaticks*. To observe also what follows, upon mixing the *Liquors*, or other *Parts*
of

of *Plants* together; as *Oyl* of *Turpentine*, by *Digestion* with a *Lixivial Salt*, extracteth thence a *Red Tincture*. Or with *Salts*, *Earths*, *Metals*, or any other *Bodies*; as the *Juyce* of the green *Leaves* of *Rasberry*, *Primrose*, and divers other *Plants* (I think principally such as are *Astringent*) expressed upon *Steel*, as it drieth, becometh of a *Purple Colour*.

46. §. Lastly, by *Compounding* the *Experiment* it self, or joyning two or more of them, upon the same matter: as *Fermentation* and *Destillation*, as is used for some *Waters*. *Infusion* and *Fermentation*, as in making of *Beer*. *Fermentation* and *Codion*, or rather *Affation*, as in making of *Bread*. *Arefaction* and *Destillation*, as may be tryed upon some *Herbs*; and with what difference from what may be noted, upon their being distilled, moist.

47. §. Having proceeded thus far, by all the above particular *Ways* of *Observation*; a *Comparative Prospect* must be taken of them: by which, at last, the *Communities* and *Differences* of the *Contents* of *Vegetables*, may be discerned; the manner of their *Causation* and *Original*, partly, be judged of; and wherein it is, that the *Essence* of their several *Natures* and *Qualities* doth consist, in some measure comprehended. And consequently, both from the knowledge of their particular *Natures*, and the *Analogy* found betwixt them; we may be able, better to conjecture, and try, what any of them are, or may be good for. For certainly, we shall then know, more readily, to apply things unto, and more fitly to prepare them for, their *Proper Uses*, when we first know, *what they are*. Notwithstanding, since the *Faculties* of *Plants*, do often lie more recluse; it is best, therefore, not wholly to acquiesce in such *Conjectures*, as their *Tastes*, or other *Sensible Properties* may suggest; but to subjoyn *Experiment*. In making of which, and in passing a *Judgment* thereupon, many *Cautions*, both in respect of the *Plant* whereof, and the *Subject* whereupon it is made, are requisite to be attended. Which yet, in regard they result not so directly from the *Matter* at present in hand; I shall not, therefore, here insist upon them, And thus much for the *Third General Mean*.

The Fourth
General
Mean.

48. §. THE *Contents* of the *Organical Parts* of *Vegetables*, having been thus duly *Examined*: it will be requisite to make the like *Inquiry*, into their *Principles*; or the *Bodys*, immediately concurrent and essential to their *Being*. And of these, we are to observe, First, their *Number*; whether well reducible to *five*, *six*, *seven*, or *more*, or *fewer*: and the *Special Differences* observable under any one *General*; since there are many *Bodies*, of very different *Natures*, confounded under one *Name*. Next their *Conjugation*; which they are, that either under or over those observable in *animal*, or other *Bodies*, are here joyned together in a *Plant*; How far common to the *Organical Parts* of divers *Plants*; or to the several *Organical Parts* of one; or how far different in them. So the predominant *Principle* of the *Parenchymous Parts* of a *Plant*, that it is an *Acid*, seems evident, From the general *Nature* of *Fruits*; and of *Corn*; and most *Parenchymous Roots*, which are either *Spirituos*, or *Sower*, or by *Digestion*, do easily become such. Likewise their *Proportions*; which stand in the greatest, which in the least, or in the meaner *Quantities*, and in what *Degrees*; both in divers *Vegetables*, and in the several *Organical Parts* of one. And then the *Concentration* and *Union* of them altogether; as to the degrees

degrees of their Closeness or Laxity; or the manner of their Implication and Coherency; or as to their Location, one being more Central, another more Exposed and Rampant over the rest; or otherwise different. To examine these Principles, by their Colour, Taste, Smell, Consistence, Fixedness, Volatility, Weight, Figures, or other Accidents. And to these purposes, to go through the formentioned Ways of Experiment; as Ustion, Calcination, Destillation, &c. as any of them may appear applicable hereunto. So the Essential Salt of Wormwood, which may be obtained from the Lixivial; is Bitter, transparent, and commonly, of a Cylindrick figure: whereas that which is obtained by Coction, or from the Extract, is tasteless, greyish, and almost Cubick: and that in the Extract of the Green Leaves of Violets, appears in fine transparent Shoots, like so many little Needles. And it is probable, That the Salts of most Kinds of Plants, whether Lixivial or Essential; and of these, whether obtained by Decoction, or otherwise, have either their Figure, or other Qualities, proper to themselves, whereby they are all distinguished one from another. And lastly, to make Experiment upon these Principles, mixing them with one another, or with other Bodies, or otherwise.

49. §. I know it will be difficult to make observations of this kind upon the Organical Parts of Plants, severally. Yet I have thought of some Ways, whereby true and undeceivable ones may be made. And the better to illustrate what I mean, I shall give one or two Instances of Tryal to this purpose. For the making of which, and some others of the like nature, I considered, That upon the Anatomical Analysis of all the Parts of a Plant, I had certainly found, (and shall hereafter shew) That in all Plants, there are Two, and only Two Organical Parts Essentially distinct, viz. The Pithy Part, and the Lignous Part, 'or such others as are analogous to either of These. So that, if we can think of any Plants, which will afford us either of these two, though not perfectly, yet in some good measure, simple and unmixed: We may then see, by putting them to a Chymical, Test, what Principles and Proportion of Principles, concur to specify their Substantial Forms.

50. §. To the Pithy Part, Starch, or pure Manchet is analogous, as having very little of the Lignous mixed with them. I therefore ordered lb ij of Starch to be put into a Retort, and with a Receiver affixed, to be set in a Sand Furnace; and that all it would yield, should, by degrees, be forced over; which, besides what was evaporated at the Neck of the Receiver, was about lb j. of an acid and eager Liquor, of a heavy and blackish Oyl ʒ ss, and of a light Oyl ʒ j. The Caput Mortuum could not be reduced to Ashes, by the strongest heat which a naked fire in that Furnace would produce.

51. To the Lignous Part, Hemp or Flax is analogous, having very little of the Pithy mixed with them. I caused therefore lb ij of Flax to be put into a Retort, and manag'd as the Starch: whereupon, it yielded a Liquor, as I remember, somewhat like the former, and about the same quantity; no Oyl which remained liquid, when cold; but instead of that a Butyr, almost of the Consistence and Colour of the Oyl of Mace; and of this above ʒ iij, or near six times the quantity of the Oyl which was yielded by the starch. The Caput Mortuum being burned to a white Ash, yielded some portion of a Lixivial Salt.

52. §. From whence, I shall, at present, only make these two Remarques; *First*, That although the chief portion, as to quantity, in both these *Bodys*, (as in most *Plants*) is an Acid Liquor; yet the latter, yields also some of an Alkaly, which the other doth not. So that they are the *Lignous Parts* of a *Plant*, generally, which yield the *Alkalick Salt*, or at least in the greatest Proportion. *Secondly*, That the *sulphurous or Oleous Principle*, is also much more predominant in the *Lignous Part*, than in the *Pithy*. To these, the like Tryals upon other *Plants*, should be added; and other ways. So, in regard the *Soot* of most *Woods*, yields a *Volatile Alkaly*; it were fit to examine, Whether the *Soot* which is made of the *Pithy Parts* and that, of the *Lignous*, afford the said *Alkaly*, in equal quantity; or whether, as is most likely, that of the *Lignous* doth afford it in a far greater: and the like.

53. §. The prosecution of what is here proposed, will be requisite, To a fuller and clearer view, of the *Modes of Vegetation*, of the *Sensible Natures of Vegetables*, and of their more Recluse *Faculties and Powers*. *First*, of the *Modes of Vegetation*. For suppose we were speaking of a *Root*; from a due consideration of the *Properties* of any *Organical Part* or *Parts* thereof; 'tis true, that the real and genuine *Causes* may be rendred, of divers other dependent *Properties*, as spoken generally of the whole *Root*. But it will be asked again, What may be the *Causes* of those *first* and Independent ones? Which, if we will seek, we must do it by inquiring also, What are the *Principles* of those *Organical Parts*? For it is necessary, that the *Principles* whereof a *Body* doth consist, should be, if not all of them the *active*, yet the *capacitating Causes*, or such as are called *Causæ sine quibus non*, of its becoming and being, in all respects, both as to *Substance* and *Accidents*, what it is: otherwise, their Existence, in that *Body*, were altogether superfluous; since it might have been without them: which if so, it might then have been made of any other; there being no necessity of putting any difference, if neither those, whereof it is made, are thought necessary to its Being. Wherefore if we will allow a *Body*, and so the *Organical Parts* of a *Vegetable* to have *Principles*, we must allow these *Principles* their necessary Use; and that the Shapes or other *Properties* of the said *Parts*, are as much dependant upon the *Nature* of These; as is the Roundness of a Drop of Ink, upon the Fluidity of Water, ingredient to it.

54. §. Again, the *Principles* of the *Organical Parts* being known, we may from thence obtain a further knowledge of the *Natures*, and *Causation* or *Original* of their *Contents*; since these *Contents* are not only included in the said *Organical Parts*, but also Created by them: and must needs be so, whether we will suppose the *Principles* of these *Contents* to be præ-existent to their reception thereinto, or not. For, if not præ-existent, what can be clearer, than that the said *Parts* give them their Existence? And if præ-existent, yet in regard they are distinguished, and such only of them admitted in such sort into an *Organical Part*, from amongst others, as are apt to combine and mix together in such a *Form*, and so to constitute such a *Liquor*; it is as clear, that the Existence, if not of those *Principles*, yet of that *Liquor*, is dependent on the said *Part*.

55. §. And if by means of the said *Organical Parts*, it is, that their *Contents* become *such* and *such* peculiar *Mixtures*; it is hence also manifest, That, by the same means, they are of *such* distinct *Faculties* and *Powers*: Because the *Faculty* or *Power* of a Body, lieth not in any of its *Principles* apart; but is a Resultance from them all; or from their being, in such peculiar sort and manner, United and Combined together. So the *Principles* of the Purgative Parts of a *Root*, as of *Rhubarb*, although we should suppose them to be existent in the surrounding *Earth*, yet we cannot say, That *that Earth*, or the *Principles* therein contained, are Purgative; but only that they are such, as by being combined together, in such a peculiar way, may become *so*. So the several parts of a *Clock*, although they are and must be all præ-existent to it, and it is their *Form*, by which they are, what they are; yet is it the *setting together* of such *Parts*, and in such a way only, that makes them a *Clock*. And since we see that the *Mixture* of two Bodies of two different *Qualities*, as of *Two Colours*, will produce a *Third Colour*, differing from them both; as Blue and Red, do a Murrey: Why should not *Two or More Bodies* of different *Natures*, be so combined together, as to produce a *Third Nature*? Or wherefore may not that be allowed to be performed by *Nature*, which by Artificial Compounding of *Medicines*, or other Bodies, is designed, and oftentimes effected? I'll give but one Instance; *Water*, *Grease*, and an *Alcalizate Salt*, may be easily so ordered as to be invested with new *Qualities*, *Nature*, and *Powers*; the *Salt*, to lose its extreme fiery Pungent *Taste*; the *Tallow*, its *Smell*; and being before unfociable with the *Water*, to mingle therewith: neither *Tallow*, *Salt*, nor *Water* alone, will fetch out a spot of *Grease*; but all united easily do it: the same Three Bodys united, are, in some Cases, as in the *Jaundies*, no ill *Medicine*; any of which, given alone, may rather prove prejudicial, than a cure: and all this done, only by duly boiling them together into one Body, which we call *Sope*.

56. §. Whence again, if it be such an *Union*, and *Proportion*, of such a *Sort* of *Principles*, which produceth such a *Faculty*; and that we may, by any means, come to know what these are; we may, possibly, also attain to the knowledge of such *Rules*, whereby any kind of *Faculty* may be made; as to Compound such Bodies, which are neither Purgative nor Vomitory, so together, as to be Invested with those *Faculties*. And if to Make them, then consequently, to Mend, Exalt, Strengthen, and Enoble them, with greater ease and certainty. And thus much for the *Fourth General Mean*.

57. §. HITHERTO, We have considered the *Materials* of a *Ve-*
getable, only as Ingredient to it: there yet remains a *Fifth Story* to be
ascended; which is, to consider these *Materials* as they are derived
from *abroad*: or as, after they are received and naturalized, they may,
with others yet abroad, have any kind of correspondence. And these
are *Four* in general, *scil. Earth, Water, Aer, and Sun*; all which, in
that they contribute so universally to *Vegetation*, and to whatsoever is
contained in a *Vegetable*, it is therefore requisite, that of These likewise,
Particular Observation should be made.

The Fifth
General
Mean.

58. §. And First, of the *Earth*, and of all Solid Receptacles of
Plants. Where we are to consider their several *Kinds*; as Mellow,
Sandy, Clayie, Chalky, and others. Their *Ingredients*; as Rank and

Mellow *Earth*, with Sand, or with Clay; or Sand with Clay; or altogether; and in what Proportions. The *Principles* whereinto any one of these *Ingredients*, separated from the rest, and put to the Test of *Distillation*, *Ustion*, *Calcination*, or other, either alone, or by mixture with other Bodies, may be Resolved. And by their *Qualities*, as *Colour*, *Smell*, *Taste*, &c. both *Ingredients* and *Principles* to be examined. To make tryal of the growth of *Plants*, in all kinds of *simple Soils*; either *Earthy* or *Mineral*, as Clay, Marl, Oker, Fullers Earth, Bole Armeniac, Vitriol, Allum, &c. or *Vegetable*, as Rotten Wood, Brans, Starch, or Flower, &c. or *Animal*, as Dungs, pounded Fleth, dried and powdered Blood, and the like; that it may appear, how far any of these may contribute to the growth of a *Plant*; or to one, above another.

59. §. Next of the *Water*, and of all Liquid Receptacles. Where the several kinds of *Water*, from Wells, Springs, Rain, and Rivers are, by their *Qualities* and *Faculties*, to be examined; as these, and by these, their *Principles*, either in their Natural State, or upon Digestion, or otherwise, may be observable: since Common *Water* it self, is undoubtedly compounded of several *Principles*; the simplicity thereof, not being argued, from its Clearness and Transparency; for a Solution of *Alum*, though it containeth a considerable quantity of *Earth*, is yet very Clear: nor from its seeming to have neither *Smell* nor *Taste*; for *Water-drinkers* will tell you of the varieties of both in different *Waters*. Besides, if these *Qualities* should be accounted rather Phantise, than Sense; the difference of *Waters* is yet more manifest, from their different Effects, observed by *Cooks*, *Laundresses*, *Brewers*, and others, that have occasion to use them: for not to mix with *Sope*, without curdling; not to boil Meat tender, or without colouring it red; and the like, are the vices of some *Waters*, not of others, which yet would seem, in Colour, Taste, and Smell, to be the same. Tryal should also be made of the growth of *Plants* in all kinds of Liquid Receptacles, as *Common Water*, *Snow Water*, *Sea Water*, *Urine*, *Milk*, *Whey*, *Wine*, *Oyl*, *Ink*, &c. Or any of these, with a solution of *Salt*, *Nitre*; *Sal prunellæ*, *Sope*, or other body. And hereby to observe what follows, either in the *Liquor*, or in the *Plant* it self: as if any *fixed* Body, being weighed before its dissolution in *Water*; and if the *Plant*, set herein, groweth; the *Water*, being then evaporated; whether the quantity of that dissolved body, continue the same, or is lessened. So, whether any *Vegetable* will become *Opiate*, by growing a considerable time in a plain Solution or *Water-tincture* of *Opium*; and the like. Which *Experiments*, what event soever they have, yet at least, for our further instruction in the *Nature* of *Vegetation*, may be of use.

60. §. Next of *Aer*, where it will be requisite to inquire, what sort of Bodies may be herein contained: It being probable, from the variety of *Meteors* formed herein; and of *Vapours* and *Exhalations* continually advanced hereinto; that some or other of them, may bear an Analogy, to all Volatile Bodys, whether *Animal*, *Vegetable*, or *Mineral*. The flourishing also of *Frozen Dew*; and the Green Colour, which the *Aer* gives the Ground or Water, when, for some time exposed to it; and other effects; seem to argue, that it is Impregnated with *Vegetable Principles*. To consider also the peculiar Nature of that Body, which is strictly called, *Aer*, And of that true *Aerial Salt*, which to me, seemeth probable,

bable, that it is dissolved in the *Æther*, as other *Salts* are in *Water*, or in the Vaporous parts of the *Aer*. As also to try, what different Effects, a diversity of *Aer* may have upon a *Vegetable*; as by setting a *Plant*, or *Seed*, either exceeding Low, as at the bottom of a deep Well; or exceeding High, as on the top of a Steeple. Or else by exposing some *Soil* to the *Aer*, which is assuredly free from any *Seed*, and so, as no *Seed* can light upon it; and to observe, whether the *Aer* hath a power of producing a *Vegetable* therein, or not: and the like.

61. §. Lastly of the *Sun*; as to which, it may be considered, What Influence it may have upon the *Plant*, it self; upon the *Soil*; Or upon the *Aer*. Whether that *Influence* is any thing else besides Heat: or may differ from that of a *Fire*, otherwise, than by being Temperate, and more Equal. That it doth, seems evident from an *Experiment* sometime since given us, in one of the *Parisian Journals des Scavans*, and which I therefore think very applicable to our present purpose. If you hold a *Concave* at a due distance, against a *Fire*, it will collect and cast the Heat into a burning *Focus*: but if you put a peice of plain *Glass* between them, the *Glass* will scatter the Heat, and destroy the *Focus*. Whereas the *Sun-Beams*, being gathered in like manner, will pass through the interposed *Glass*, and maintain their *Focus*. As for That, of the Collection of the *Sun-beams*, by the help of *Glasses*, in the form of a *Magistry*, or of *Flowers*, and such like, I desire to suspend my thoughts of them, till I see them. I will only say thus much further at present; That I do not understand why the *Sun* should not have some Influence upon Bodies, besides by Heat, if it may be granted, That the *Moon* hath; for which, it should seem, there are some good Arguments.

62. §. WE HAVE thus far examined the *Principles* necessary to *Vegetation*. The *Question* may be put once more, In what manner are these *Principles* so adapted, as to become capable of being assembled together, in such a *Number*, *Conjugation*, *Proportion*, and *Union*, as to make a *Vegetable Body*? For the comprehension whereof, we must also know, What are the *Principles* of these *Principles*. Which, although they lie in so great anabyss of obscurity; yet, I think, I have some reason to believe, that they are not altogether undiscoverable. How far they may be so, I am so far from Determining, that I shall not now Conjecture.

A Sixth General Means Only hinted

63. §. THIS is the *Design*, and these the *Means* I propose in order thereunto. To which, I suppose, they may all appear to be necessary. For what we obtain of *Nature*, we must not do it by commanding, but by courting of Her. Those that woo Her, may possibly have her for their *Wife*; but She is not so common, as to prostitute her self to the best behaved *Wit*, which only practiseth upon it self, and is not applied to her. I mean, that where ever Men will go beyond *Phansie* and *Imagination*, depending upon the Conduct of *Divine Wisdom*, they must Labour, Hope and Persevere. And as the *Means* propounded, are all necessary, so they may, in some measure, prove effectual. How far, I promise not; the Way is long and dark: and as Travellers sometimes amongst Mountains, by gaining the top of one, are so far from their Journeys end; that they only come to see another lies before them: so the Way of *Nature*, is so impervious, and, as I may say, down Hill and up Hill, that how far soever we go, yet the surmounting of one difficulty, is wont still to give us the prospect of another. We may there-

The Conclusion.

therefore believe, our attainments will be imperfect, after we have done all : but because we cannot attain to all, that therefore we should endeavour after nothing ; is an Inference, which looks so much awry from the Practical Sense of Men, that it ought not to be answered. Nor with better Reason, may we go about determining, what may be done. The greatest Designs that any Men undertake, are of the greatest uncertainty, as to their Success: which, if they appear to be of good Import, though we know not how far they are attainable, we are to propound the *Means*, in the utmost use whereof only, we can be able to judge : A *War* is not to be quitted, for the hazards which attend it ; nor the *Councils of Princes* broken up, because those that sit at them, have not the Spirit of Prophecy, as well as of Wisdom. To conclude, If but little should be effected, yet to design more, can do us no harm : For although a Man shall never be able to hit *Stars* by shooting at them ; yet he shall come much nearer to them, than another that throws at *Apples*.

F I N I S.

P 3

THE
ANATOMY
OF
PLANTS,
BEGUN.

WITH A
General Account
OF
VEGETATION,
Grounded thereupon.

The FIRST BOOK.

Presented in Manuscript to the ROYAL SOCIETY,
Sometime before the 11th. of *May*, 1671.

And afterwards in Print, *December 7.* of the same Year 1671.

By NEHEMIAH GREW M.D. Fellow of the
Royal Society, and of the *College of Physicians*.

The Second Edition.

LONDON,
Printed by *W. Rawlins*, 1682.

THE
ANATOMY

OF
PLANTS

General Account

OF
VEGETATION

By W. VAUGHAN

Author of the

1784

TO THE
RIGHT HONOURABLE
WILLIAM
Lord Vi-Count Brouncker,
THE
PRESIDENT,

And to the
Council and Fellows
OF THE
ROYAL SOCIETY,

The following
ANATOMY

Is most HUMBLY
PRESENTED

By the AUTHOR

NEHEMIAH GREW.

WILLIAM

PRESIDENT

ROYAL SOCIETY

ANNALS

1788

1789

1790



TO THE
Right Reverend
JOHN
Lord Bishop of
CHESTER.

MY LORD,

I Hope your pardon; if while you are holding
That best of Books in one Hand, I here
present some Pages of that of *Nature* in-
to your other: Especially since *Your Lord-*
ship knoweth very well, how excellent a
Commentary This is on the *Former*; by
which, in part, *GOD* reads the World his own Defini-
tion, and their Duty to him.

But if this Address, *my Lord*, may be thought con-
gruous, 'tis yet more just; and that I should let *Your*
Lordship, and others know, how much, and how deser-
vedly, I resent *Your* extraordinary Favours. Particularly,
that you were pleased, so far to animate my Endeavours,
towards the Publishing the following *Observations*. Ma-

Epistle Dedicatory.

ny whereof, and most belonging to the *First Chapter*, having now lain dormant, near seven years; and might still, perhaps, have so continued, had not *Your Lordships* Eye, at length, created Light upon them. In doing which, *You* have given one, amongst those many Tokens, of as well *Your* readiness to promote Learning and Knowledge by the hands of others; as *Your* high Abilities to do it by *Your* Own: Both which, are so manifest in *Your Lordship*, that, like the first Principles of *Mathematical Science*, they are not so much to be asserted, because known and granted by all.

The Consideration whereof, *my Lord*, may make me not only *Just*, in owning of your Favours; but also most *Ambitious* of your *Patronage*: Which yet, to bespeak, I must confess, I cannot well. Not that I think, what is Good and Valuable, is always its own best Advocate: for I know, that the Censures of Men, are humorous, and variable; and that one *Age*, must have leave to frown on those *Books*, which another, will do nothing less than kiss and embrace. But, chiefly, for this Reason, Left I should so much as seem desirous, of *Your Lordships* Solliciting my Cause, as to all I have said. For as it is your Glory, that you like not so to shine, as to put out the least Star; so were it to *Your* Dishonour, to borrow *Your* Name, to illustrate the Spots, though of the most conspicuous. I am,

My Lord,

Your Lordships

Most Obliged,

And

Most Humble Servant

NEHEMIAH GREW.

Coventry,
June 10. 1671.

THE

THE
CONTENTS.

CHAP. I.

Of the Seed in its State of Vegetation.

THE Method propounded, §. 1. The Garden-Bean, dissected, 2. The two Coats Described, 3, 4. The Foramen in the outer Coat, 5, 6. What generally observable of the Covers of the Seed, 7. The Organical Parts of the Seed, 8. The Main Body, 9, 10. The Radicle in the Bean, 11. In other Seeds, 12. The Plume, 13, 14. The Similary Parts, 15. The Cuticle, 16, 17. The Parenchyma, 18, 19, 20. The Inner Body, 21, to 29. No solid Account yet given, of Vegetation, 30. The Coats how in common subservient to the Vegetation of the Seed, 31. The Foramen, of what use herein, 32. The use of the Inner Coat, 33. Of the Cuticle, 34. Of the Parenchyma, 35. Of the Seminal Root, 36. How the Radicle first becomes a Root, 37. By what means, the Plume all this while preserved, 38. How after the Root the Plume vegetates, 39. How the Lobes, 40. But not in all Seeds, 41. That they do in most, demonstrated, 42, 43, 44. What hence resolvable, 45. The use of the Dissimilar Leaves, 46, to the end.

CHAP. II.

Of the Root.

THis also to be Dissected, §. 1. The Skin hereof, its Original, 2. The Cortical Body, its Original, 3. Texture, 4. Pores, 5. Proportions, 6. The Lignous Body, its Original, 7. Texture, 8. Proportions, 9. The Insetment, its Original, 10. Pores, 11. Number and Size, 12. A fuller description hereof, with that of the Osculations of the Lignous Body, 13, 14, 15. The Pith, its original sometimes from the Seed, 16. Sometimes from the Barque, 17. Its Pores, 18, 19. Proportions, 20. Fibres of the Lignous Body therein, 21. The Pith of those Fibres, 22. How the Root grows, and the use of the Skin, Cortical

The Contents.

tical and Lignous Body thereto, 23. How it groweth in length, 24. By what means it descends, 25. How it grows in breadth, 26. And the Pith, how thus framed, 27. The use of the Pith, 27. Of the Insertment, 28. The joynt service of all the Parts, 29, 30, 31.

CHAP. III.

Of the Trunk.

THE Coarcture, §. 1. The Skin, its original, 2. The original of the Cortical Body, 3. Of the Lignous, 4. Of the Insertment and Pith, 5. The Latitudinal Shooting of the Lignous Body, wherein observable, 6, 7. The Pores of the Lignous Body, where and how most remarkable, 8. A lesser sort of Pores, 9. A third sort only visible through a Microscope. Observed in Wood or Charcoal, 10. Observed in the Fibres of the Trunks of Herbs, 11. The Insertions, where more visible, 12, 13. Their Westage with the Lignous Body, 14. The smaller Insertions, only visible through a Microscope, 15. No Valves in a Plant, 16. The Ranks of the Pores of the Insertions, 17. The Pores of the Pith, 18, 19, 20. How the Trunk ascends, 21. The disposition of its Parts consequent to that Ascent, 22. Consequent to the different Nature of the Sap, 23. The effects of the said Differences, 24, to 28. Which way, and how the Sap ascends, 29, to the end.

The Appendix.

Of Trunk-Roots and Claspers.

Trunk-Roots of two kinds, §. 1, 2. Claspers of one kind, 3. The Uses of both, 4, to the end.

CHAP.

The Contents.

CHAP. IV.

Of the Bud, Branch, and Leaf.

THE Parts of the Germen and Branch the same with those of the Trunk, §. 1, 2. The manner of their growth, 3. How nourished, 4. And the use of Knots, 5. How secur'd 6. The Parts of a Leaf, 7. The Positions of the Fibres of the Stalks of Leaves, 8. For what Uses, 9, 10. The visible cause of the different circumference of Leaves, 11. And of their being flat, 12. And filamentous, 13. The Foulds of Leaves, their Kinds and Use, 14, 15, 16. The Protections of Leaves, 17. The use of the Leaf, 18, to the end.

The Appendix.

Of Thorns, Hairs and Globulets.

Thorns of two Kinds; the Lignous, §. 1. The Cortical, 2. An argument of the Magnetick Descent of the Cortical Body, 3. Hairs of divers Kinds, 4, 5. Their Use, 6. Globulets of two Kinds, 7, 8.

CHAP. V.

Of the Flower.

ITs three Parts, §. 1. The Impalement, of divers kinds, 2. Their use, 3, 4. The Foliation, its nature, 5. Foulds, 6. Protections, 7. Downs, 8, 9. Globulets, 10. Its Use, 11, 12. The Attire of two kinds. The Description of the first, 13, 14, 15, 16. Of the other, 17, 18, 19, 20, 21. Their use, 22, to the end.

CHAP. VI.

Of the Fruit.

THE Vital Parts of all, the same, §. 1. The Number, Description, and Original of the Parts of an Apple, 2. Of a Pear, 3, 4. Of a Plum, 5, 6, 7. Of a Nut, 8. Of a Berry, 9. The use of the Fruit, 10, to the end.

CHAP.

The Contents.

CHAP. VII.

Of the Seed in its State of Generation.

WHat here further observed, not in the First Chapter, §. 1. The Case, its Figures, 2. The outer Coat, its Figures, 3. Various Surface, 4. And Mucilages, 5. The nature of the outer Coat, 6. Its Apertures, 7. Next to which the Radicle usually placed, 8. The Original of the Outer Coat, 9. The Original of the Inner, 10. Its Nature, 11, 12. The Essential Parts of a Plant, 13, 14. The Secondine, 15. The Colliquamentum herein, 16. The Navel Fibres, 17. In the Generation of the Seed, the Sap first prepared in the Seed-Branch, 18, 19. Next in the inner Coat, 20. With the help of the Outer, 21, 22. The use of the Secondine, 23. Of the Ramulets of the Seed-Branch, 24. Of their Inosculation, 25. How the Colliquamentum becometh a Parenchyma, 26, to the end.

THE



THE
ANATOMY
OF
PLANTS,
BEGUN.

With a General Account of *Vegetation*,
Founded thereupon.

CHAP. I.

Of the Seed in its State of Vegetation.

BEING to speak of *Plants*; and, as far as Inspection, and consequent Reason, may conduct, to enquire into the visible *Constitutions*, and *Uses* of their several *Parts*: I choose that Method, which, to the best advantage, may suit with what we have to say hereon. And that is the Method of Nature her self, in her continued Series of *Vegetations*; proceeding from the *seed sown*, to the formation of the *Root, Trunk, Branch, Leaf, Flower, Fruit*, and last of all, of the *seed* also to be *sown again*; all which, we shall, in the same order, particularly speak of.

2. §. The *Essential Constitutions* of the said *Parts* are in all *Plants* the same: But for *Observation*, some are more convenient; in which I shall chiefly instance. And first of all, for the *Seed*, we choose the great *Garden-Bean*.

3. §. If then we take a *Bean* and dissect it, we shall find it cloath-
ed with a doubled *Vest* or *Coat*. These *Coats*, while the *Bean* is yet
green, are separable, and easily distinguished. Or in an old one, after
it hath lay'n two or three days in a mellow Soil; or been soaked as
long a time in Water: as in *Tab. I.* When 'tis dry, they cleave so closely
together, that the Eye not before instructed, will judge them but one;
the inner *Coat* (which is of the most rare contexture) so far shrinking up,
as to seem only the roughness of the outer, somewhat resembling *Wafers*
under *Maquaroons*.

Tab. I. f. 2.

4. §. The Inner *Coat*, in its Natural State, is every where twice,
and in some places, thrice as thick, as the Outer. Next to the *Radicle*,
which I shall presently describe, it is six or seven times thicker;
and encompasses the *Radicle* round about, as in the same *Figure* ap-
pears.

Tab. I. f. 2.

5. §. At the thicker end of the *Bean*, in the outer *Coat*, a very
small *Foramen* presents it self, even to the bare Eye. In Dissection
'tis found to terminate against the point of that *Part* which I call
the *Radicle*. It is of that capacity, as to admit a small *Virginal* Wyer;
and is most of all conspicuous in a green *Bean*. Especially, if a little
magnified with a good *Spectacle-Glass*. This *Foramen* is not a hole ca-
sually made, or by the breaking off of the Stalk; but designedly form-
ed, for the uses hereafter mentioned. It may be observed not only in
the great *Garden-Bean*, but likewise in the other *kinds*; in the *French-
Bean* very plainly; in *Pease*, *Lupines*, *Vetches*, *Lentiles*, and other
Pulse 'tis also found; and in many *Seeds* not reckoned of this kind-
red, as in that of *Fænugreek*, *Medica Tornata*, *Goats-Rue*, and others:
In many of which, 'tis so very small, as scarcely, without the help of
Glasses to be discovered; and in some, not without cutting off part
of the *Seed*, which otherwise would intercept the sight hereof.

Tab. I. f. 1--2

6. §. That this *Foramen* is truly permeable, even in old *Setting-
Beans*, and the other *Seeds* above named, appears upon their being
soaked for some time in Water. For then, taking them out, and crush-
ing them a little, many small bubbles will alternately arise and break up-
on it.

7. §. Of all *Seeds* which have thick or hard *Covers*, it is also ob-
servable, That they have the same likewise *Perforated*, as above said,
or in some other manner. And accordingly, although the *Coats* of such
Seeds as are lodg'd in *Shells* or *Stones*, being thin, are not visibly *per-
forated*; yet the *Stones* and *Shells* themselves always are; as in *Chap. 7.*
shall be seen how. To which *Chapter*, what is farther observable, ei-
ther as to the nature and number of the *Covers* of the *Seed*, I also
refer.

8. §. The *Coats* of the *Bean* being stripp'd off, the proper *seed*
shews it self. The parts whereof it is composed, are three; *sc.* the
Main Body, and two more, appendant to it; which we may call, the
Three Organical Parts of the *Bean*.

9. §. The *Main Body* is not one entire piece, but always divided,
lengthwise, into two halves or *Lobes*, which are both joyn'd together
at the *Basis* of the *Bean*. These *Lobes* in dry *Beans*, are but difficultly
separated or observ'd; but in young ones, especially boil'd, they easily
slip asunder.

Tab. I. f. 2, 3.

10. §. Some very few *Seeds* are divided, not into two *Lobes*, but into more; as that of *Cresses* into Six. And some are not at all divided, but entire; as the Grains of *Corn*. Excepting which few, all other *Seeds*, even the smallest, are divided, like as the *Bean*, into just two *Lobes*. Whereof, though in most *Seeds*, because of their minuteness, we cannot by dissection be inform'd; yet otherwise, we easily may, as in this *Chapter* shall be seen. Tab. 1. f. 4.
f. 5.

11. §. At the *Basis* of the *Bean*, the two other *Organical Parts* stand appendent; by mediation whereof, the two *Lobes* meet and join together. The greater of these two *Parts* stands without the two *Lobes*, and upon divesting the *Bean* of its *Coats*, is immediately visible. 'Tis of a white colour, and more glossie than the *Main Body*, especially when the *Bean* is young. In the *Bean*, and many other *Seeds*, 'tis situated somewhat above the thicker end, as you hold the *Bean* in its most proper posture for growth. In *Oak-Kernels*, which we call *Acorns*, *Apple-Kernels*, *Almonds*, and many other *Seeds*, it stands prominent just from the end; the *Basis* and the *End* being in these the same, but in the *Bean* divers. Tab. 1. f. 2.
e 3---a
Tab. 1. f. 6---c

12. §. This *Part* is found not only in the *Bean*, and the *Seeds* above mentioned; but in all others: being that, which upon the Vegetation of the *Seed*, becomes the *Root* of the *Plant*; which therefore may be called the *Radicle*: by which, I mean the Materials, abating the Formality, of a *Root*. In *Corn*, it is that *Part*, which *Malsters*, upon its shooting forth, call the *Come*. 'Tis not easie to be observed, saving in some few *Seeds*, amongst which, that of the *Bean* is the most fair and ample of all I have seen. But that of some other *Seeds*, is, in proportion, greater; as of *Fænugreek*, which is full as big as one of its *Lobes*. Tab. 1. f. 7---e

13. §. The lesser of the two said Appendents lies occult between the two *Lobes* of the *Bean*, by separation whereof only it is to be seen. 'Tis enclosed in two small *Cavities*, form'd in the *Lobes* for its reception. Its colour comes near to that of the *Radicle*; and it is founded upon the *Basis* thereof, having a quite contrary production, *sc.* towards the *Cone* of the *Bean*; as being that very *Part*, which, in process, becomes the *Body* or *Trunk* of the *Plant*. In *Corn*, it is that *Part*, which after the *Radicle* is sprouted forth, or *come*, shoots towards the smaller end of the *Grain*; and by many *Malsters*, is called the *Acrospire*. Tab. 1. f. 3---b

14. §. This *Part* is not, like the *Radicle*, an entire Body, but divided, at its loose end, into divers pieces, all very closely couched together, as Feathers in a Bunch; for which reason it may be called the *Plume*. They are so close, that only two or three of the outmost are at first seen: but upon a nice and curious separation of these, the more interiour still may be discovered. In the *Bean*, this may be done: but in very few other *Seeds*; because of the extreme smallness of the *Plume*. Now as the *Plume* is that *Part* which becomes the *Trunk* of the *Plant*, so these pieces are so many true, and already formed, though not display'd *Leaves*, intended for the said *Trunk*, and foulded up in the same *plicature*, wherein upon the sprouting of the *Bean*, they afterwards appear. In a *French Bean*, and especially in the larger white Kind, or in the great *Indian Phaseolus*, the two outmost are very fair and elegant. In the great *Garden-Bean* two extraordinary small *Plumes* often, if not always, stand one on either side the great one now describ'd: From Tab. 1. f. 8---b

which, in that they differ in nothing save in their size, I therefore only here just take notice of them. And these three Parts, *sc.* the *Main Body*, the *Radicle*, and the *Plume*, are concurrent to the making up of a *Seed*; and no more than these

15. §. Having thus taken a view of the *Organical Parts* of the *Bean*, and other *Seeds*; let us next examine the *Similary*, *sc.* those whereof the *Organical* are compos'd: a distinct observation of which, for a clear understanding of the *Vegetation* of the *Seed*, and of the whole *Plant* arising thence is requisite: To obtain which, we must proceed in our *Anatomy*.

16. §. Dissecting a *Bean* then, the first *Part* occurring is its *Cuticle*. The Eye and first Thoughts, suggest it to be only a more dense and glossy Superficies; but better enquiry discovers it a real *Cuticle*. 'Tis so exquisitely thin, and for the most part, so firmly continuous with the *Body* of the *Bean*, that it cannot, except in some small Rag, be distinctly seen, which, by carrying your Knife aslant into the *Bean*, and then very gently bearing upward what you have cut, will separate, and shew it self transparent. This *Cuticle* is not only spread upon the *Convex* of the *Lobes*, but also on their *Flats*, where they are contiguous, extending it self likewise upon both the *Radicle* and *Plume*, and so over the whole *Bean*.

17. §. This *Part*, though it be so far common with the *Coats* of the *Bean*, as to be like those, an *Integument*; yet are we in a quite different Notion to conceive of it: For whereas the *Coats*, upon setting the *Bean*, do only administer the *Sap*, and, as being superseded from their Office, then die; as shall be seen: this, on the contrary, with the *Organical Parts* of the *Bean*, is nourished, augmented, and by a real *Vegetation* co-extended.

18. §: Next to the *Cuticle*, we come to the *Parenchyma* it self; the *Part* throughout which the *Inner Body*, whereof we shall speak anon, is disseminated; for which reason I call it the *Parenchyma*. Not that we are so meanly to conceive of it, as if (according to the stricter sense of that word,) it were a meer concreted Juyce. For it is a *Body* very curiously *organiz'd*, consisting of an infinite number of extreme small *Bladders*; as in *Tab. 1.* is apparent. The *Surface* hereof is somewhat dense, but inwardly, 'tis of a laxer Contexture. If you view it in a *Microscope*, or with a very good *Spectacle-Glass*, it hath some similitude to the *Pith*, while *sappy* in the *Roots* and *Trunks* of *Plants*; and that for good reason, as in *Ch. 2.* shall be seen. This is best seen in green *Beans*.

19. §. This *Part* would seem by its colour to be peculiar to the *Lobes* of the *Bean*; but as is the *Cuticle*, so is this also, common both to the *Radicle* and *Plume*; that is, the *Parenchyma* or *Pulp* of the *Bean*, as to its essential substance, is the same in all three. The reason why the colour of the *Plume*, and especially of the *Radicle*, which are white, is so different from that of the *Lobes*, which are green, may chiefly depend upon their being more compact and dense, and thence their different Tinctures. And therefore the *Lobes* themselves, which are green while the *Bean* is *young*; yet when it is *old* and *dry*, become whitish too. And in many other *Seeds*, as *Acorns*, *Almonds*, the *Kernels* of *Apples*, *Plums*, *Nuts*, &c. the *Lobes*, even *fresh* and *young*, are pure white as the *Radicle* it self.

20. §:

20. §. But although the *Parenchyma* be common, as is said, to all the *Organical Parts*; yet in very differing proportions. In the *Plume*, where it is proportionably least, it maketh about three *Fifths* of the whole *Plume*; in the *Radicle*, it maketh above five *Sevenths* of the whole *Radicle*; and in each *Lobe*, is so far over-proportionate, as to make at least nine *Tenths* of the whole *Lobe*.

21. §. By what hath been said, that the *Parenchyma* or *Pulp* is not the only constituting *Part*, besides the *Cuticle*, is imply'd: there being another *Body*, of an essentially different substance, embosom'd herein: which may be found not only in the *Radicle* and *Plume*, but also in the *Lobes* themselves, and so in the whole *Bean*.

22. §. This *Inner Body* appears very plain and conspicuous in cutting the *Radicle* athwart, and so proceeding by degrees towards *Tab. I. f. 10*, the *Plume*, through both which it runneth in a large and strait *Trunk*, *11, & 12*. In the *Lobes*, being it is there in so very small proportion, 'tis difficultly seen, especially towards their *Verges*. Yet if with a sharp Knife you smoothly cut the *Lobes* of the *Bean* athwart, divers small *Specks*, of a different colour from that of the *Parenchyma*, standing therein all along in a Line, may be observ'd; which *Specks* are the Terminations of the *Branches* of this *Inner Body*, *Tab. I. f. 13*.

23. §. For this *Inner body*, as it is existent in every *Organical Part* of the *Bean*; so is it, with respect to each *Part*, most regularly distributed. In a good part of the *Radicle* 'tis one entire *Trunk*; towards the *Basis* thereof, 'tis divided into three main *Branches*; the middlemost runneth directly into the *Plume*; the other two on either side it, after a little space, pass into the *Lobes*; where the said *Branches* dividing themselves into other smaller; and those into more, and smaller again, are terminated towards the *Verges* of each *Lobe*; in which manner the said *Inner Body* being distributed it becomes in each *Lobe* a true and perfect *Root*, *Tab. I. f. 14*.

24. §. Of this *Seminal Root*, as now we'll call it, from the Description here given, it is further observable; That the two main *Branches* hereof; in which the several *Ramifications* in each *Lobe* are all united, are not committed into the *Seminal Trunk* of the *Plume*, nor yet stand at right angles with *That* and the *Radicle*, and so with equal respect towards them both: but being produced through part of the *Parenchyma* of the *Radicle*, are at last united therein to the main *Trunk*, and make acute Angles therewith: as may be seen in the same. *f. 14. Tab. I. f. 14*.

25. §. This *Seminal Root* being so tender, cannot be perfectly excarnated, (as may the *Vessels* in the *Parts* of an *Animal*) by the most accurate Hand. Yet by dissection begun and continu'd, as is above declared, its whole frame and distribution may be easily observ'd. Again, if you take the *Lobe* of a *Bean*, and lengthwise pare off its *Parenchyma* by degrees, and in extreme thin slices, many *Branches* of the *Seminal Root*, (which by the other way of Dissection were only noted by so many *Specks*) both as they are fewer about the *Basis* of the *Bean*, and more numerous towards its *Verges*, in some good distinction and entireness will appear. For this you must have new *Beans*: or else soaked in Water, or buried for some time.

26. §. As the *Inner Body* is branched out in the *Lobes*, so is it in the *Plume*: For if you cut the *Plume* athwart, and from the *Basis* proceed along the Body thereof, you'll therein find, first, one large *Trunk*

OR

or *Branch*, and after four or five very small *Specks* round about it, which are the terminations of so many lesser *Branches* therewith distributed to the several parts of the *Plume*. The distribution of the *Inner Body*, as it is continuous throughout all the *Organical Parts* of the *Bean*, is represented, *Tab. I. f. 14.*

27. §. This *Inner Body* is, by dissection, best observable in the *Bean* and great *Lupine*. In other larger *Pulse* it shews likewise some obscure Marks of it self. But in no other *Seeds*, which I have observed; though of the greatest size: as of *Apples, Plums, Nuts, &c.* is there any clear appearance hereof, upon dissection, saving in the *Radicle* and *Plume*; the reason of which is partly from its being, in most *Seeds*, so extraordinary little; partly from its Colour, which in most *Seeds*, is the same with that of the *Parenchyma* it self, and so not distinguishable from it.

28. §. Yet in a *Gourd-Seed*, the whole *Seminal Root*, not only its *Main Branches*, but also the *Sub-divisions* and *Inosculation*s of the lesser ones, are without any dissection, upon the separation of the *Lobes*, on their contiguous Flats immediately apparent.

And as to the existence of this *Seminal Root*, what Dissection cannot attain, yet an ocular inspection in hundreds of other *Seeds*, even the smallest, will demonstrate; as in this *Chapter* shall be seen how.

29. §. In the mean time, let us only take notice; That when we say, every *Plant* hath its *Root*, we reckon short. For every *Plant* hath really two, though not contemporary, yet successive *Roots*; its *Original* or *Seminal-Root* within the *Lobes* or *Main Body* of its *Seed*; and its *Plant-Root*, which the *Radicle* becometh in its growth: the *Parenchyma* of the *Seed*, being in some resemblance, that to the *Seminal Root* at first, which the Mould is to the *Plant-Root* afterwards; and the *Seminal Root* being that to the *Plant-Root*, which the *Plant-Root* is to the *Trunk*. For our better understanding whereof, having taken a view of the several *Parts* of a *Bean*, as far as Dissection conducts; we will next briefly enquire into the Use of the said *Parts*, and in what manner they are the Fountain of *Vegetation*, and concurrent to the being of the future *Plant*.

Account of the Vegetation of the Bean. 30. §. THE GENERAL Cause of the growth of a *Bean*, or other *Seed*, is *Fermentation*. That is, the *Bean* lying in the Mould, and a moderate access of some moisture, partly dissimilar, and partly congenerous, being made, a gentle *Fermentation* thence ariseth. By which, the *Bean* swelling, and the *Sap* still encreasing, and the *Bean* continuing still to swell, the work thus proceeds: as is the usual way of explicating. But that there is simply a *Fermentation*, and so a sufficient supply of *Sap* is not enough: but that this *Fermentation*, and the *Sap* wherein 'tis made, should be under a various Government, by divers *Parts* thereto subservient, is also requisite; and as the various preparation of the *Aliment* in an *Animal*, equally necessary: the particular process of the Work according whereto, we find none undertaking to declare.

31. §. Let us look upon a *Bean* then, as a piece of Work so fram'd and set together, as to declare a Design for the production of a *Plant*; which, upon its lying in some convenient Soil, is thus effected. First of all, the *Bean* being enfoulded round in its *Coats*, the *Sap* wherewith it is fed, must of necessity pass through these: By which means, it is not

not only in a proportionate quantity, and by degrees; but also in a purer body; and possibly not without some *Vegetable Tincture*, transmitted to the *Bean*. Whereas, were the *Bean* naked, the *Sap* must needs be, as over-copious, so but crude and immature, as not being *filtr'd* through so fine a *Cotton* as the *Coats* be. And as they have the use of a *Filtre* to the transient *Sap*; so of a *Vessel* to that which is still deposited within them; being alike accommodated to the securer *Fermentation* hereof, as *Bottles* or *Barrels* are to *Beer*, or any other *Fermentative Liquor*.

32. §. And as the *Fermentation* is promoted by some *Aperture* in the *Vessel*; so have we the *Foramen* in the upper *Coat* also contrived. That if there should be need of some more *Aiery* Particles to excite the *Fermentation*; through *this*, they may obtain their Entry. Or, on the contrary, should there be any such *Particles* or *Steams*, as might damp the genuine proceeding thereof, through this again, they may have easie issue. Or if, by being over copious, they should become too high a *Ferment*; and so precipitate those soft and slow degrees, as are necessary to a due *Vegetation*. The said *Aperture* being that, as a common *Passport*, here to the *Sap*, which what we call the *Bung-hole* of the *Barrel*, is to the new tunn'd *Liquor*.

33. §. And the *Radicle* being designed to shoot forth first, as presently shall be shew'd how; therefore is it distinctly surrounded with the *Inner* and more succulent *Coat*. That being thereby supplied on every side, its eruption may be the better promoted.

34. §. The *Sap* being passed through the *Coats*, it next enters the *Body* of the *Bean*; yet not indiscriminately neither; but, being filtr'd through the *Outer Coat*, and fermented in the *Body* of the *Inner*, is by mediation of the *Cuticle*, again more finely filtr'd, and so entereth the *Parenchyma* it self under a fourth Government.

35. §. Through which *Part* the *Sap* passing towards the *Seminal Root*, as through that which is of a more spacious content; besides the benefit it hath of a farther *percolation*, it will also find room enough for a more free and active fermenting and *maturation* herein. And being moreover, part of the true *Body* of the *Bean*, and so with its proper *Seminalities* or *Tinctures* copiously repleat; the *Sap* will not only find room, but also matter enough, by whose Energy its *Fermentation* will still be more advanced,

36. §. And the *Sap* being duly prepared here, it next passeth into all the *Branches* of the *Seminal Root*, and so under a fifth Government. Wherein how delicate 'tis now become, we may conceive by the proportion betwixt the *Parenchyma* and this *Seminal Root*; so much only of the best digested *Sap* being discharged from the whole *Stock* in that, as this will receive. And this, moreover, as the *Parenchyma*, with its proper *Seminalities* being endowed; the *Sap* for the supply of the *Radicle*, and of the young *Root* from thence, is duly prepared therein, and with its highest *Tincture* and *Impregnation* at last enriched.

37. §. The *Sap* being thus prepared in the *Lobes* of the *Bean*, 'tis thence discharg'd; and either into the *Plume*, or the *Radicle*, must forthwith issue. And since the *Plume* is a dependent on the *Radicle*; the *Sap* therefore ought first to be dispenced to this: which accordingly, is ever found to shoot forth before the *Plume*: and sometimes an inch or two in length. Now because the primitive course of the *Sap* into
the

the *Radicle*, is thus requisite ; therefore, by the frame of the *Parts* of the *Bean* is it also made necessary. The two main *Branches* of the *Seminal Root*, being produced, as is before observed, not into the *Plume*, but the *Radicle*. Now the *Sap* being brought as far as the *Seminal Root*, in either *Lobe* ; and according to the conduct thereof continuing still to move : it must needs immediately issue into the same *Part*, whereinto the main *Branches* themselves do ; that is, into the *Radicle*. By which *Sap*, thus bringing the several *Tinctures* of the *Parts* aforesaid with it, being now fed ; it is no longer a meer *Radicle*, but is made also *Seminal*, and so becomes a perfect *Root*.

Tab. I. f. 14.

38. §. The *Plume*, all this while, lyes close and still. For the sake of which, chiefly it is, that the *Bean* and other *seeds* are divided into *Lobes*, viz. That it might be warmly and safely lodged up between them, and so secur'd from the Injuries so tender a *Part* would sustain from the Mould ; whereto, had the *Main Body* been entire, it must, upon the cleaving of the *Coats*, have lay'n contiguous.

39. §. But the *Radicle* being thus impregnated and shot into a *Root* ; 'tis now time for the *Plume* to rouse out of its Cloysters, and germinate too : In order whereto, 'tis now fed from the *Root*, with laudable and sufficient *Aliment*. For as the Supplies and Motion of *Sap* were first made from the *Lobes*, towards the *Root* : so the *Root* being well shot into the Mould, and now receiving a new and more copious *Sap* from thence ; the motion hereof must needs be stronger, and by degrees proceed in a contrary course, sc. from the *Root* toward the *Plume* : and, by the continuation of the *Seminal Root*, is directly conducted thereinto ; by which being fed, it gradually enlarges and displays it self.

40. §. The course of the *Sap* thus turned, it issues, I say, in a direct Line from the *Root* into the *Plume* : but collaterally, into the *Lobes* also ; sc. by those two aforesaid *Branches* which are obliquely transmitted from the *Radicle* into either *Lobe*. By which *Branches* the said *Sap* being disburfed back into all the *Seminal Root*, and from thence, likewise into the *Parenchyma* of the *Lobes*, they are both thus fed, and for some time augmenting themselves, really grow : as in *Lupines* is evident.

Tab. I. f. 14.

41. §. Yet is not this common to all *seeds*. Some rot underground ; as *Corn* ; being of a laxer and less Oleous Substance, differing herein from most other *seeds* ; and being not divided into *Lobes*, but one entire thick Body. And some, although they continue firm, and are divided into *Lobes*, yet rise not ; as the great *Garden Bean*. In which, therefore, it is observable, That the two Main *Branches* of the *Lobes*, in comparison with that which runs into the *Plume*, are but mean ; and so insufficient to the feeding and vegetation of the *Lobes* ; the *Plume*, on the contrary, growing so lusty, as to mount up without them.

42. §. Excepting a few of these Two Kinds, all other *seeds* whatsoever, (which I have observed) besides that they continue firm ; upon the *Vegetation* of the *Plume*, do mount also upwards, and advance above the Ground together with it ; as all *seeds* which spring up with one or more *Dissimilar Leaves* : These *Dissimilar Leaves*, for the most part *Two*, which first spring up ; and are of a different shape from those that follow, being the very *Lobes* of the *Seed*, divided, expanded, and thus advanced.

43. §.

43. §. The Impediments of our apprehension hereof are the Colour, Size and Shape of the *Dissimilar Leaves*. Notwithstanding, that they are nothing else but the *Main Body* of the *Seed*, how I came first to conceive, and afterwards to know it, was thus. First, I observed in general, that the *Dissimilar Leaves*, were never jagg'd, but even edg'd: And seeing the even verges of the *Lobes* of the *Seed* here-to respondent, I was apt to think, that those which were so like, might prove the same. Next descending to particular *Seeds*, I observed, first, of the *Lupine*; that, as to its Colour, advancing above the Ground, (as it useth to do) it was always changed into a perfect Green. And why might not the same by parity of Reason be inferred of other *seeds*? That, as to its size, it grew but little bigger than when first set. Whence, as I discern'd (the Augmentation being but little) we here had only the two *Lobes*: So, (as some augmentation there was) I inferr'd the like might be, and that, in farther degrees, in other *Seeds*. Tab. 2. f. 1.

44. §. Next of the *Cucumber-Seed*, That, as to its Colour, often appearing above ground, in its primitive white, from white it turns to yellow, and from yellow to green; the proper colour of a *Leaf*. That, as to its size, though at its first arise, the *Lobes* were little bigger than upon setting; yet afterwards, as they chang'd their Colour, so their dimensions also, growing to a three-four-five-fold amplitude above their primitive size. But whereas the *Lobes* of the *Seed*, are in proportion, narrow, short and thick: how then come the *Dissimilar Leaves*, to be so exceeding broad, or long, and thin? The Question answers it self: For the *Dissimilar Leaves*, for that very reason are so thin, because so very broad or long; as we see many things, how much they are extended in length or breadth, so much they lose in depth, or grow more thin; which is that which here befalls the now effoliated *Lobes*. For being once dis-imprisoned from their *Coats*, and the course of the *Sap* into them, now more and more encreased; they must needs very considerably amplify themselves: and from the manner wherein the *Seminal Root* is branched in them, that amplification cannot be in thickness, but in length or breadth. In both which, in some *Dissimilar Leaves*, 'tis very remarkable; especially in length, as in those of *Lettice*, *Thorn-Apple*, and others; whose *Seeds*, although very small, yet the *Lobes* of those *seeds* growing up into *Dissimilar Leaves*, are extended an Inch, and sometimes more, in length. Though he that shall attempt to get a clear sight of the *Lobes* of *Thorn-Apple*, and some others, by *Disssection*, will find it no easie Task; yet is that which may be obtained; and in the Last *Book* shall be shew'd. From all which, and the observation of other *seeds*, I at last found, that the *Dissimilar Leaves* of a young *Plant*, are nothing else but the *Lobes* or *Main Body* of its *Seed*. So that, as the *Lobes* did at first feed and impregnate the *Radicule* into a perfect *Root*; so the *Root*, being perfected, doth again feed, and by degrees amplify each *Lobe* into a perfect *Leaf*. Tab. 2. f. 2.

45. §. The Original of the *Dissimilar Leaves* thus known, we understand; why some *Plants* have none; because the *Seed* either riseth not, as *Garden-Beans*, *Corn*, &c. Or upon rising, the *Lobes* are little alter'd, as *Lupines*, *Pease*, &c. Why, though the proper *Leaves* are often indented round; the *Dissimilar* like the *Lobes* are even-edg'd. Why, though the proper *Leaves* are often hairy, yet these are ever smooth. Why some have more *Dissimilar Leaves* than two, as *Cresses*, Tab. 2. f. 3.

Hist. of the
Prop. of Vege. which have six, as the Ingenious Mr. *Sbarrock* also observes. The reason whereof is, because the *Main Body* is not divided into Two, but Six, distinct *Lobes*, as I have often counted. Why *Radishes* seem at first to have four, which yet after appear plainly two: because the *Lobes* of the *Seed*, have both a little Indenture, and are both plaited, one over the other. To which, other Instances might be added.

46. The use of the *Dissimilar Leaves* is, first, for the protection of the *Plume*; which being but young, and so but soft and tender, is provided with these, as a double Guard, one on either side of it. For this reason it is, that the *Plume*, in *Corn*, is trussed up within a membranous *Sheath*: and that of a *Bean*, cooped up betwixt a pair of *Surfoyls*: But where the *Lobes* rise, there the *Plume* hath neither of them, being both needless.

Tab. 3.

47. §. Again, since the *Plume*, being yet tender, may be injur'd not only by the *Aer*, but also for want of *Sap*, the supplies from the *Root* being yet but slow and sparing; that the said *Plume* therefore, by the *Dissimilar Leaves*, may have the advantage likewise of some refreshment from Dew or Rain. For these having their *Basis* a little beneath that of the *Plume*, and expanding themselves on all sides of it, they often stand after Rain, like a Vessel of Water, continually soaking and suppling it, lest its new access into the *Ayr*, should shrivel it.

48. §. Moreover, that since the *Dissimilar Leaves* by their *Basis* intercept the *Root* and *Plume*, the greater and grosser part of the *Sap*, may be, by the way, deposited into those; and so the purest proceed into the yet but young and delicate *Plume*, as its fittest *Aliment*.

49. §. Lastly, we have here a demonstration of the being of the *Seminal Root*: which, since through the colour or smalness of the *Seed*, it could not by Dissection be observ'd, except in some few; Nature hath here provided us a way of viewing it in the now effoliated *Lobes*, not of one or two *Seeds*, but of hundreds; the *Seminal Root* visibly branching it self towards the Cone or Verges of the said *Lobes*, or now *Dissimilar Leaves*.

CHAP. II.

Of the R O O T.



HAVING Examined and pursued the Degrees of *Vegetation* in the *Seed*, we find its two *Lobes* have here their utmost period: and, that having conveyed their *Seminalities* into the *Radicle* and *Plume*; these therefore, as the *Root* and *Trunk* of the *Plant*, still survive. Of these, in their order, we next proceed to speak; and first, of the *Root*: whereof, as well as of the *Seed*, we must by *Dissection* inform our selves.

2. §. In *Dissection* of a *Root* then, we shall find it with the *Radicle*, as the *Parts* of an *Old Man* with those of a *Fœtus*, substantially, one. The first *Part* occurring is its *Skin*, the Original whereof is from the *Seed*: For that extreme thin *Cuticle* which is spread over the *Lobes* of the *Seed*, and from thence over the *Radicle*, upon the shooting of the *Radicle* into a *Root*, is co-extended, and becomes its *skin*.

3. §. The next *Part* is the *Cortical Body*. Which, when it is thin, is commonly called the *Barque*. The Original hereof, likewise is from the *Seed*; or the *Parenchyma*, which is there common both to the *Lobes* and *Radicle*, being by *Vegetation* augmented and prolonged into the *Root*, the same becomes the *Parenchyma* of the *Barque*. Tab. 2. f. 4.

4. §. The *Contexture* of this *Parenchyma* may be well illustrated by that of a *Sponge*, being a *Body* Porous, Dilative and Pliable. Its *Pores*, as they are innumerable, so, extream small. These *Pores* are not only susceptible of so much *Moisture* as to fill, but also to enlarge themselves, and so to dilate the *Cortical Body* wherein they are: which by the shriv'ling in thereof, upon its being expos'd to the *Air*, is also seen. In which dilatation, many of its *Parts* becoming more lax and distant, and none of them suffering a solution of their continuity; 'tis a *Body* also sufficiently pliable; that is to say, a most exquisitely fine-wrought *Sponge*.

5. §. The *Extention* of these *Pores* is much alike by the length and breadth of the *Root*; which from the shrinking up of the *Cortical Body*, in a piece of a cut *Root*, by the same dimensions, is argu'd.

6. §. The proportions of this *Cortical Body* are various: If thin, 'tis, as is said, called a *Barque*; and thought to serve to no other end, than what is vulgarly ascrib'd to a *Barque*; which is a narrow conceit. If a Bulky *Body*, in comparison with That within it, as in the young *Roots* of *Cichory*, *Asparagus*, &c. 'tis here, because the fairest, therefore taken for the prime *Part*; which, though, as to *Medicinal* use, it is; yet, as to the private use of the *Plant*, not so. The *Colour* hereof, though it be originally white, yet in the continued growth of the *Root*, divers *Tinctures*, as yellow in *Dock*, red in *Bistort*, are thereinto introduced.

7. §. Next within this Part stands the *Lignous Body*: This *Lignous Body*, lyeth with all its parts, so far as they are visible, in a Circle or Ring. Yet are there divers extreme small *Fibres* thereto parallel, usually mixed with the *Cortical Body*; and by the somewhat different colour of the said *Cortical Body* where they stand, may be noted. These *Fibres* the *Cortical Body*, and *Skin*, altogether, properly make the *Barque*. The Original of this *Lignous Body*, as of the two former, is from the *seed*; or, the *Seminal Roots* of both the *Lobes*, being united in the *Radicle*, and with its *Parenchyma* co-extended, is here in the *Root* of the *Plant*, the *Lignous Body*.

Tab. 2. f. 4. b.

8. §. The Contexture hereof, in many of its parts, is much more close than that of the *Cortical*; and their *Pores* very different. For whereas those of the *Cortical* are infinitely numerous, these of the *Lignous* are in comparison nothing so. But these, although fewer, yet are they, many of them, more open, fair and visible: as in a very thin Slice cut athwart the young *Root* of a *Tree*, and held up against the light, is apparent. Yet not in all equally; in *Coran-Tree*, *Goosberry-Tree*, &c. less, in *Oak*, *Plums*, and especially *Damascens*, more; in *Elder*, *Vines*, &c. most conspicuous. And as they are different in number and size, so also (whereon the numerousness of the *Pores* of the *Cortical Body* principally depends) in their shape. For whereas those of the *Cortical Body* are extended much alike both by the length and breadth of the *Root*; these of the *Lignous*, are only by the length; which especially in *Vines*, and some other *Roots* is evident. Of these *Pores*, 'tis also observable, that although in all places of the *Root* they are visible, yet most fair and open about the *filamentous Extremities* of some *Roots*, where about, the *Roots* have no *Pith*; as in *Fenil*. And in many *Roots*, higher.

Tab. 2. f. 5.

Tab. 2. f. 7 a. b.

9. §. The proportion betwixt this *Lignous Body* and the *Cortical*, is various, as was said; yet in this, constant, sc. that in the *filamentous* and smaller Parts of the *Root*, the *Lignous Body* is very much the less; running like a slender *Wyer* or *Nerve* through the other surrounding it. Whereas in the upper part, it is often times of far greater quantity than the *Cortical*, although it be encompass'd by it. They stand both together pyramidally, which is most common to *Infant Roots*, but also to a great many others.

10. The next Part observable in the *Root*, is the *Insertment*. The existence hereof, so far as we can yet observe, is sometimes in the *Radicle* of the *Seed* it self; I cannot say always. As to its substantial nature, we are more certain; that it is the same with that of the *Parenchyma* of the *Radicle*; being always at least augmented, and so, in part, originated from the *Cortical Body*, and so, at second hand, from the said *Parenchyma*. For in dissecting a *Root*, I find, that the *Cortical Body* doth not only environ the *Lignous*, but is also wedg'd, and in many Pieces inserted into it; and that the said inserted Pieces make not a meer Indenture, but transmit and shoot themselves quite through as far as the *Pith*: which in a thin Slice cut athwart the *Root*, as so many lines drawn from the Circumference towards the Center, shew themselves.

Tab. 2. f. 5.

11. §. The *Pores* of the *Insertment* are sometimes, at least, extended somewhat more by the breadth of the *Root*, as about the top of the *Root* of *Borage* may be seen; and are thus different from those of the

the

the *Cortical Body*, which are extended by the length and breadth much alike; and from those of the *Lignous*, being only by its length.

12. §. The number and size of these *Insertions* are various. In *Hawthorn*, and some others, and especially *Willows*, they are most extream small; in *Cherries* and *Plums* they are Bigger; and in the *Vine* and some other *Trees*, very fairly apparent. In the *Roots* of most *Herbs* they are generally more easily discoverable; which may lead to the observation of them in all. Tab. 2. f. 5.
& 6.

13. §. These *Insertions*, although they are continuous through both the length and breadth of the *Root*; yet not so in all Parts, but by the several shootings of the *Lignous Body* they are frequently intercepted. For of the *Lignous Body* it is (here best) observable; That its several *Shootings*, betwixt which the *Cortical* is inserted, are not, throughout the *Root*, wholly distinct, strait and parallel: but that all along being enarch'd, the *Lignous Body*, both in length and breadth, is thus disposed into *Braces* or *Osculations*. Betwixt these several *Shootings* of the *Lignous Body* thus osculated, the *Cortical* shooting, and being also osculated answerably *Brace* for *Brace*, that which I call the *Insertment* is framed thereof. Tab. 2. f. 8.

14. §. These *Osculations* are so made, that the *Pores* or *Fibres* of the *Lignous Body*, I think, notwithstanding, seldom or never run one into another; being, though contiguous, yet still distinct. In the same manner as some of the *Nerves*, though they meet, and for some space are associated together, yet 'tis most probable, that none of their *Fibres* are truly inosculated, saving perhaps, in the *Plexures*.

15. §. These *Osculations* of the *Lignous Body*, and so the interception of the *Insertions* of the *Cortical*, are not to be observ'd by the traverse cut of the *Root*, but by taking off the *Barque*. In the *Roots* of *Trees*, they are generally obscure; but in *Herbs* often more distinctly apparent; and especially in a *Turnep*: the appearance whereof, the *Barque* being stripp'd off, is as a piece of close-wrought *Network*, fill'd up with the *Insertions* from thence.

16. §. The next and last distinct *Part* of the *Root* is the *Pith*. The substantial nature thereof, is, as was said of the *Insertment*, the same likewise with that of the *Parenchyma* of the *Seed*. And according to the best observation I have yet made, 'tis sometimes existent in its *Radicle*; in which, the two main *Branches* of the *Lobes* both meeting, and being osculated together, are thus dispos'd into one round and tubular *Trunk*, and so environing part of the *Parenchyma*, make thereof a *Pith*; as in either the *Radicle*, or the young *Root* of the great *Bean* or *Lupine*, may, I think, be well seen.

17. §. But many times the *Original* hereof is immediately from the *Barque*. For in dissection of divers *Roots*, both of *Trees* and *Herbs*, as of *Barberry* or *Mallows*, it is observable, That the *Cortical Body* and *Pith*, are both of them participant of the same Colour; in the *Barberry*, both of them tinged yellow, and in *Mallows*, green. In cutting the smaller *Parts* of the *Roots* of many *Plants*, as of *Borage*, *Mallows*, *Parsley*, *Columbine*, &c. 'tis also evident, That the *Lignous Body* is not there, in the least *Concave*, but standeth *Solid*, or without any *Pith*, in the *Center*; and that the *Insertions* being gradually multiplied afterwards, the *Pith*, at length, towards the thicker parts of the *Root*, shews and enlarges it self. Whence it appears, that in all such *Roots*, Tab. 2. f. 9. -- 2
the

the *Pith* is not only of the same substantial nature, and by the *In-
sertions* doth communicate with the *Barque*; and that it is also aug-
mented by it; which is true of the *Pith* of all *Roots*; but is moreo-
ver, by mediation of the said *Insertions*, wholly originated from it;
that is to say, from the *Parenchymous* Part thereof. The various appea-
rances of the *Insertions* and *Pith* from the filamentous *Parts* to the
top of the *Root*, see in *Tab. 2.* The *Pores* of the *Lignous Body*, as
Tab. 2. f. 9. it stands entire in the said filamentous *Parts*, are best seen when they
have lain by a night to dry, after cutting.

18. §. A farther evidence hereof are the Proportions betwixt the
Cortical Body and *Pith*. For as about the inferiour *Parts* of the *Root*,
where the *Pith* is small, the *Cortical Body* is proportionably great; so
about the top, where the *Pith* is enlarged, the *Cortical Body* (now more
properly becoming a *Barque*) groweth proportionably less, *sc.* because
the *Insertions* do still more and more enlarge the *Pith*. Likewise the
peculiar frame of some *Roots*, wherein besides the *Pith*, the *Lignous
Body* being divided into two or more *Rings*, there are also one or more
thick *Rings*, of a white and soft substance, which stand betwixt them;
and are nothing else but the *Insertions* of the *Cortical Body* collected
into the said *Rings*; but, towards the top of the *Root*, being inserted
again, thus make a large and ample *Pith*; as in older *Fennel-Roots*, those
of *Beet*, *Turnep*, and some other *Herbs*, is seen.
Tab. 2. f. 8.

19. §. The *Pores* of the *Pith*, as those of the *Cortical Body*, are
extended both by the breadth and length of the *Root*, much alike; yet
are they more or less of a greater size than those of the *Cortical
Body*.

20. §. The Proportions of the *Pith*, are various; in *Trees*, but
small; in *Herbs*, generally, very fair; in some making by far the great-
est part of the *Root*; as in a *Turnep*: By reason of the wide circum-
ference whereof, and so the finer Concoction and Assimilation of its
Sap; that *Part* which in most old *Trunks* is a dry and harsh *Pith*, here
proves a tender, pleasant meat.

21. §. In the *Roots* of very many *Plants*, as *Turneps*, *Carrots*, &c.
the *Lignous Body*, besides its main utmost *Ring*, hath divers of its of-
culated *Fibres* dispersed throughout the *Body* of the *Pith*; sometimes
all alike, and sometimes more especially in, or near, its Center; which
Fibres, as they run towards the top of the *Root*, still declining the
Center, at last collaterally strike into its Circumference; either all of
them, or some few, keeping the Center still. Of these principally, the
Succulent part of the *Lignous Body* of the *Trunk* is often originated.

22. §. Some of these *Pith-Fibres*, although they are so exceeding
slender, yet in some *Roots*, as in that of *Flower de liz*, they are visibly
concave, each of them, in their several Cavities also embosoming a ve-
ry small *Pith*; the sight whereof, the *Root* being cut traverse, and laid
in a Window for a day or two to dry, may without *Glasses* be obtain'd.
And this is the general account of the *Root*; the declaration of the man-
ner of its growth, with the use and service of its several *Parts*, we shall
next endeavour.

23. §. I SAY THEN, That the *Radicle* being impregnate, and
shot into the Moulds, the contiguous moisture, by the *Cortical Body*, be-
ing a *Body* laxe and Spongy, is easily admitted: Yet not all indiscrimi-
nately, but that which is more adapted to pass through the surrounding
Cuticle.

An Account
of the
Growth of
the *Root*.

Cuticle. Which transient *Sap*, though it thus becomes fine, yet is not simple; but a mixture of *Particles*, both in respect of those originally in the *Root*, and amongst themselves, somewhat heterogeneous. And being lodg'd in the *Cortical Body* moderately laxe, and of a Circular form; the effect will be an easie Fermentation. The *Sap* fermenting, a separation of *Parts* will follow; some whereof will be impacted to the Circumference of the *Cortical Body*, whence the *Cuticle* becomes a *Skin*; as we see in the growing of the Coats of Cheeses, of the Skin over divers Liquors, and the like. Whereupon the *Sap* passing into the *Cortical Body*, through this, as through a *Manica Hippocratis*, is still more finely filtered. With which *Sap*, the *Cortical Body* being dilated as far as its *Tone*, without a solution of Continuity, will bear; and the supply of the *Sap* still renew'd: the purest part, as most apt and ready, recedes, with its due *Tinctures*, from the said *Cortical Body*, to all the parts of the *Lignous*; both those mixed with the *Barque*, and those lying within it. Which *Lignous Body* likewise super-inducing its own proper *Tinctures* into the said *Sap*; 'tis now to its highest preparaton wrought up, and becomes (as they speak of that of an Animal) the Vegetative *Ros* or *Cambium*: the noblest part whereof is at last coagulated in, and assimilated to the like substance with the said *Lignous Body*. The remainder, though not united to it, yet tinctur'd therein, thus retreats, that is, by the continual appulse of the *Sap*, is in part carried off into the *Cortical Body* back again, the *Sap* whereof it now tinctures into good *Aliment*. So that whereas before, the *Cortical Body* was only relaxed in its Parts, and so dilated; 'tis now increas'd in real quantity or number of parts, and so is truly nourish'd. And the *Cortical Body* being saturate with so much of this Vital *Sap* as serves it self; and the second Remainders discharged thence to the *Skin*; this also is nourish'd and augmented therewith. So that as in an *Animal Body* there is no instauration or growth of Parts made by the *Bloud* only, but the *Nervous Spirit* is also thereunto assistant; so is it here: the *Sap* prepared in the *Cortical Body*, is as the *Bloud*, and that part thereof prepared by the *Lignous*, is as the *Nervous Spirit*; which partly becoming Nutriment to it self, and partly being discharged back into the *Cortical Body*, and diffusing its Tincture through the *Sap* there, that to the said *Cortical Body* and *Skin*, becomes also true Nutriment, and so they all now grow.

24. §. In which growth, a proportion in length and breadth is requisite: which being rated by the benefit of the *Plant*, both for firm standing and sufficient *Sap*, must therefore principally be in length. And because it is thus requisite, therefore by the constitution of one of its *Parts*, *sc.* the *Lignous Body*, it is also made necessary. For the *Pores* hereof, in that they are all extended by its length, the *Sap* also according to the frame and site of the said *Pores* will principally move; and that way as its *Sap* moves, the same way will the generation of its Parts also proceed; *sc.* by its length. And the *Lignous Body* first (that is by a *priority causal*) moving in length it self; the *Cortical* also moves therewith. For that which is nourish'd, is extended: but whatever is extended, is mov'd: that therefore which is nourish'd, is mov'd: The *Lignous Body* then being first nourish'd, 'tis likewise first mov'd, and so becomes and carries in it the Principle of all Vegetative motion in the *Cortical*; and so they both move in length.

25. §. Yet as the *Lignous Body* is the *Principle* of Motion in the *Cortical*; so the *Cortical* is the *Moderator* of that in the *Lignous*: As in Animal Motions, the *Principle* is from the *Nerves*; yet being once given to the *Muscle* or *Limb*, and that moving proportionably to its structure, the *Nerves* also are carried in the same motion with it. We suppose therefore, that as the principal motion of the *Lignous Body* is in length, so is its *proper tendency* also to *Ascend*. But being much exceeded both in *Compass* and *Quantity* by the *Cortical*, as in the smaller parts of the *Root* it is; it must needs therefore be over-born and governed by it; and so, though not lose its motion, yet make it that way wherein the *Cortical Body* may be more obedient to it; which will be by descent. Yet both of them being sufficiently pliable, they are thus capable, where the *Soyl* may oppose a direct descent, there to divert any way, where it is more penetrable, and so to descend obliquely. For the same reason it may also be, that though you set a *Bean* with the *Radicle* upward; yet the *Radicle*, as it shoots, declining also gradually, is thus arch'd in form of an *Hook*, and so at last descends. For every declination from a perpendicular *Line*, is a mixed motion betwixt *Ascent* and *Descent*, as that of the *Radicle* also is, and so seeming to be dependent upon the two *Contrary Tendencies* of the *Lignous* and *Cortical Bodies*. What may be the cause of those *Tendencies* (being most probably external, and a kind of *Magnetisme*) I shall not make my *Task* here to enquire.

26. §. Now although the *Lignous Body*, by the position and shape of its *Pores*, principally groweth in length; yet will it in some degree likewise in breadth: For it cannot be supposed that the purest *Sap* is all received into the said *Pores*; but that part thereof likewise, staying about its *Superficial Parts*, is there tinctur'd and agglutinated to them. And because these *Pores* are prolonged by its length; therefore it is much more laxe and easily divisible that way; as in slitting a *Stick*, or cleaving of *Timber*, and in cutting and hewing them athwart is also seen. Whence it comes to pass, that in shooting from the *Center* towards the *Circumference*, and there finding more room, its said original *Laxity* doth easily in divers places now become greater, and at length in open *Partments* plainly visible. Betwixt which *Partments*, the *Cortical Body*, being bound in on the one hand, by the surrounding *Skin* and *Moulds*, and pressed upon by the *Lignous* on the other, must needs insert it self, and so move contrary to it, from the *Circumference* towards the *Center*. Where the said contrary motions continued as begun, they at last meet, unite, and either make or augment the *Pith*. And thus the *Root* is fram'd, and the *Skin*, the *Cortical* and *Lignous Bodies*, so as is said, thereunto concurrent. We shall next shew the use of the two other *Parts*, sc. the *Insertment* and *Pith*; and first of the *Pith*.

27. §. ONE true use of the *Pith* is for the better Advancement of the *Sap*, whereof I shall speak in the next Chapter. The use I here observe, is for the quicker and higher Fermentation of the *Sap*: For although the Fermentation made in the *Cortical Body* was well subservient to the first *Vegetations*, yet those more perfect ones in the *Trunk* which after follow, require a *Body* more adapted to it, and that is the *Pith*; which is so necessary, as not to be only common to, but considerably large in the *Roots* of most *Plants*; if not in their inferior parts,

parts, yet at their tops. Where though either deriv'd or amplify'd from the *Cortical Body*, yet being by its *Insertions* only, we may therefore suppose, as those, so this, to be more finely constituted. And being also from its coarctation, while inserted, now free; all its *Pores*, upon the supply of the *Sap*, will more or less be amplified: Upon which accounts, the *Sap* therinto received, will be more pure, and its fermentation therein more active. And as the *Pith* is superiour to the *Cortical Body* by its *Constitution*, so by its *Place*. For as it thus stands central, it hath the *Lignous Body* surrounding it. Now as the *Skin* is the Fence of the *Cortical Body*, and that of the *Lignous*; so is the *Lignous* again a far more preheminent one unto the *Pith*; the *Sap* being here a brisk Liquor, *tunn'd up* as in a wooden *Cask*.

28. §. And as the *Pith* subserves the higher Fermentation of the *Sap*; so do the *Insertions* its purer Distribution; that separation which the parts of the *Sap*, by being fermented in the *Pith*, were dispos'd for; being, upon its entrance into the *Insertions*, now made: So that as the *Skin* is a *Filtre* to the *Cortical Body*, so are the *Insertions* a more preheminent one to the *Lignous*. And as they subserve the purer, so the freer and sufficient distribution of the *Sap*: For the *Root* enlarging, and so the *Lignous Body* growing thicker, although the *Cortical* and the *Pith* might supply *Sap* sufficient to the nutrition of its *Parts* next adjacent to them; yet those more inward, must needs be scanted of their *Aliment*; and so, if not quite starv'd, yet be uncapable of equal growth: Whereas the *Lignous Body* being through its whole breadth frequently disparted, and the *Cortical Body* inserted through it; the *Sap* by those *Insertions*, as the *Bloud* by the disseminations of the *Arteries*, is freely and sufficiently convey'd to its intimate *Parts*, even those, which from either the *Barque* or from the *Pith*, are most remote. Lastly, as the consequent hereof, they are thus assistant to the *Latitudinal growth* of the *Root*; as the *Lignous Body* to its growth in Length; so these *Insertions* of the *Cortical*, to its better growth in Breadth.

29. §. Having thus seen the solitary uses of the Several *Parts* of the *Root*, I shall lastly propound my Conjectures of that Design where-to they are altogether concurrent, and that is the *Circulation* of the *Sap*.

30. §. That the *Sap* hath a Double, and so a *Circular Motion*, in the *Root*; is probable, from the proper Motion of the *Root*, and from its Office. From its Motion, which is Descent: for which, the *Sap* must likewise, some where, have such a Motion proper to it. From its Office, which is, To feed the *Trunk*: for which, the *Sap* must also, in some *Part* or other, have a more especial Motion of Ascent.

31. §. We may therefore suppose, That the *Sap* moving in the *Barque*, towards the *Pith*, through the *Insertions*, therinto obtains a pass, Which passage, the upper *Insertions* will not favour; because the *Pith* standing in the same height with them, is there large, the fermenting and course of the *Sap* quick, and so its opposition strong. But through the lower it will much more easily enter; because there, from the smalness of the *Pith*, the opposition is little, and from the shortness of the *Insertions*, the way more open. So that the *Sap* here meeting with the least opposition, here it will bestow it self (feeding the *Lignous Body* in its passage) into the *Pith*. Into which, fresh *Sap* still entring, this being yet but crude, will subside: that

first receiv'd, and so become a Liquor higher wrought, will more easily mount upwards. And moving in the *Pith*, especially in the *Sap-Fibers* there dispers'd, as in the *Arterys*, in equal altitude with the upper-*Insertions*; the most volatile parts of all will still continue their direct ascent towards the *Trunk*. But those of a middle nature, and, as not apt to ascend, so being lighter than those beneath them, not to descend neither; they will tend from the *Pith* towards the *Insertions* in a Motion betwixt both. Through which *Insertions* (feeding the *Lignous Body* in its passage) it is, by the next subsequent *Sap*, discharged off into the *Cortical Body*, and so into the *Sap-Fibres* themselves, as into the *Veins*, back again. Wherein, being still pursu'd by fresh *Sap* from the Center, and more occurring from the Circumference, towards the lower *Insertions*, it thus descends. Through which, together with part of the *sap* afresh imbib'd from the Earth, it re-enters the *Pith*. From whence, into the *Cortical Body*, and from thence into the *Pith*, the cruder part thereof, is reciprocally disbur's'd; while the most *Volatile*, not needing the help of a *Circulation*, more directly ascendeth towards the *Trunk*.

CHAP.

CHAP. III.

Of the TRUNK.



HAVING thus declar'd the degrees of *Vegetation* in the *Root*; the continuance hereof in the *Trunk* shall next be shew'd: in order to which, the *Parts* whereof this likewise is compounded, we shall first observe.

1. §. That which without dissection shews it self, is the *Coarcture*: I cannot say of the *Root*, nor of the *Trunk*; but what I choose here to mention, as standing betwixt them, and so being common to them both; all their *Parts* being here bound in closer together, as in the tops of the grown *Roots* of very many *Plants*, is apparent.

2. §. Of the *Parts* of the *Trunk*, the first occurring is its *Skin*: The Formation whereof, is not from the Air, but in the *Seed*, from whence it is originated; being the production of the *Cuticle*, there investing the two *Lobes* and *Plume*.

3. §. The next *Part* is the *Cortical Body*; which here in the *Trunk* is no new substantial Formation; but, as is that of the *Root*, originated from the *Parenchyma* of the *Plume* in the *Seed*; and is only the increase and augmentation thereof. The *Skin*, this *Cortical Body* Tab. 3. f. 1; or *Parenchyma*, and (for the most part) some *Fibers* of the *Lignous* & 4. mixed herewith, altogether make the *Barque*.

4. §. Next, the *Lignous Body*, which, whether it be visibly divided into many softer *Fibers* or small *Threads*, as in the *Bean*, *Fennel*, and most *Herbs*; or that its *Parts* stand more compact and close, shewing one hard, firm and solid piece, as in *Trees*; it is, in all, one and the same *Body*; and that not formed originally in the *Trunk*, but in the *Seed*; being nothing else but the prolongation of the *Seminal Root* distributed in the *Lobes* and *Plume* thereof. Tab. 3. f. 1.

5. §. Lastly, The *Insertions* and *Pith* are here originated likewise from the *Plume*, as the same in the *Root*, from the *Radicle*: So that as to their *Substantial Parts*, the *Lobes* of the *Seed*, the *Radicle* and *Plume*, the *Root* and *Trunk* are all one.

6. §. Yet some things are more fairly observable in the *Trunk*. First, the *Latitudinal* shootings of the *Lignous Body*, which in *Trunks* of several years growth, are apparent in so many *Rings*, as is commonly known. For several young *Fibers* of the *Lignous Body*, as in the *Root*, so here, shooting in the *Cortical* one year, and the spaces betwixt them being after fill'd up with more (I think not till) the next, at length they become altogether a firm compact *Ring*; the *Perfection* of one *Ring*, and the *Ground-work* of another, being thus made concomitantly. Tab. 3. f. 5, & 8.

7. §. From these Annual younger *Fibers* it is, that although the *Cortical Body* and *Pith* are both of the same substantial nature, and their *Pores* little different; yet whereas the *Pith*, which the first year is green, and of all the *Parts* the fullest of *Sap*, becomes afterwards white and dry: The *Cortical Body*, on the contrary, so long as the *Tree* grows, ever keepeth green and moist, *sc.* because the said *Sap-Fibers*, annually grow therein, and so communicate with it.

8. §. The *Pores* likewise of the *Lignous Body*, many of them, in well-grown Timber, as in Oaken boards, are very conspicuous, in cutting both lengthwise and traverse. They very seldom, if ever, run one into another, but keep, like so many several *Vessels*, all along distinct; as by cutting, and so following any one of them as far as you please, for a Foot or half a Yard, or more together, may be observ'd. And so, the like, in any *Cane*.

Tab. 3. f. 2.
C. 3.

9. §. Besides these, there are a lesser sort; which, by the help only of a good *Spectacle Glass* may be observ'd.

10. §. And these are all the *Pores* visible without a *Microscope*. The use of which, excepting in some few particulars, I have purposely omitted in this first *Book*. Mr. *Hook* sheweth us, besides these, a third, and yet smaller Sort; and (as a confirmation of what, in the Second *Chapter*, I have said of the *Pores* of the *Lignous Body* in general) that they are all continuous and prolonged by the length of the *Trunk*, as are the greater ones: whereof he maketh Experiment, by filling up, in a piece of *Char-coal*, all the said *Pores* with *Mercury*: which appears to pass quite through them, in that by a very good *Glass* it is visible in their Orifices at both ends; and without a *Glass*, by the weight of the *Coal* alone, is also manifest. All these I have seen, with the help of a good *Microscope*, in several sorts of *Woods*. As they all appear in a piece of *Oak*, cut transversely, See *Tab. 3.*

Micrography.

C. 2. §. 8.

Tab. 3. f. 7.

11. §. Upon further Enquiry, I likewise find, That the *Pores* of the *Lignous Body* in the *Trunks* of *Herbs*, which at first I only supposed, by the help of good *Glasses*, are very fairly visible: each *Fibre* being sometimes perforated by 30, 50, 100, or hundreds of *Pores*. Or what I think is the truest notion of them, That each *Fibre*, though it seem to the bare eye to be but *one*, yet is, indeed, a great number of *Fibres* together; and every *Pore*, being not merely a space betwixt the several parts of the *Wood*, but the *Concave* of a *Fiber*. So that if it be asked, what all that Part of a *Plant*, either *Herb* or *Tree*, which is properly called the *Woody-Part*; what all that is, I suppose, That it is nothing else but a *Cluster* of innumerable and most extraordinary small *Vessels* or *Concave Fibers*: as in a Slice of the *Trunk* of *Burdock* is apparent.

Tab. 3. f. 6.

12. §. Next the *Insertions* of the *Cortical Body*, which in the *Trunk* of a *Tree* saw'd athawrt, are plainly discerned as they run from the *Circumference* toward the *Center*; the whole *Body* of the *Tree* being visibly compounded of two distinct Substances, that of the several *Rings*, and that of the *Insertions*, running cross; shewing that in some resemblance in a *Plain*, which the *Lines* of *Latitude* and of the *Meridian* do in a *Globe*. The entrance of the *Insertions* into the *Wood*, is also, upon stripping off the *Barque*, very apparent; as in the same *Fig. 8.*

Tab. 3. f. 5.
C. 8.

13. §.

13. §. These *Insertions* are likewise very conspicuous in Sawing of *Trees* length-ways into Boards, and those plain'd, and wrought into *Leaves* for *Tables*, *Wainscot*, *Trenchers*, and the like. In all which, as in course *Trenchers* made of *Beech*, and *Tables* of *Oak*, there are many parts which have a greater smoothness than the rest; and are so many *inserted Pieces* of the *Cortical Body*; which being by those of the *Lignous*, frequently intercepted, seem to be discontinuous, although in the *Trunk* they are really extended, in continued Plates, throughout its Breadth.

Tab. 3. f. 2.
& Tab. 4. f. 1.

14. §. These *Insertions*, although as is said, of a quite distinct substance from the *Lignous Body*, and so no where truly incorporated with it, yet being they are in all parts, the one as the *Warp*, the other as the *Woof*, mutually *braced* and *interwoven* together, they thus constitute one strong and firmly coherent Body; as the Timber of any *Tree*.

Tab. 4. f. 1.

15. §. As the Pores or Vessels are greater or less, so are the *Insertions* also: To the bare eye usually the greater only are discernable: But through an indifferent *Microscope* there are others also, much more both numerous and small, distinctly apparent, as in a transverse piece of *Oak*.

Tab. 3. f. 7.

16. §. In none of all the Pores can we observe any thing which may have the true nature and use of *Valves*, which is, Easily to admit that, to which they will by no means allow a regress. And their non-existence is enough evident, from what in the first *Chapter* we have said of the *Lobes* of the *seed*: in whose *Seminal Root*, were there any *Valves*, it could not be, that by a contrary *Course* of the *Sap*, they should ever grow; which yet, where-ever they turn into *Dissimilar Leaves*, they do. Or if we consider the growth of the *Root*, which oftentimes is *upward* and *downward* both at once. And being cut transversely, will bleed, both the same ways, with equal freedom.

C. 1. §. 42.

17. §. The *Insertions* here in the *Trunk* give us likewise a sight of the position of their Pores. For in a plained piece of *Oak*, as in *Wainscot*, *Tables*, &c. besides the larger Pores of the *Lignous Body*, which run by the length of the *Trunk*; the *Tract* likewise of those of the *Insertions* may be observed to be made by the breadth, and so directly cross. Nor are they continuous as those of the *Lignous Body*, but very short, as those both of the *Cortical Body* and *Pith*, with which the *Insertions*, as to their substance, are congenerous. Yet they all stand so together, as to be plainly ranked in even *Lines* or *Rows* throughout the breadth of the *Trunk*: As the *Tract* of those Pores appears to the naked Eye, see in *Tab. 3. Fig. 9.* The Pores themselves may be seen in the *Root* of a *Vine* described and figured in the Second *Book*, as it appears through a good *Microscope*.

Tab. 3. f. 2.

Tab. 17.

18. §. The Pores of the *Pith* likewise being larger here in the *Trunk*, are better observable than in the *Root*: the width whereof, in comparison with their *sides* so exquisitely thin, may by an *Hony-Comb* be grossly exemplified; and is that also which the vast disproportion betwixt the Bulk and Weight of a dry *Pith* doth enough declare. In the *Trunks* of some *Plants*, they are so ample and transparent, that in cutting both by the length and breadth of the *Pith*, some of them through the transparency of the *Skins* by which they are bounded, or of which they consist, would seem to be considerably

bly extended by the length of the *Pith*; but are really discontinuous and short, and as 'tis said, somewhat answerable to the *Cells* of an *Hony-Comb*. This is the nearest we can come to them, by the bare Eye without the assistance of a good *Microscope*. Mr. *Hook* sheweth in his *Micrography*, That the *Pores* of the *Pith*, particularly of *Elder-Pith*, so far as they are visible, are all alike discontinuous; and that the *Pith* is nothing else but (as he calls them) an heap of *Bubbles*. Although, in regard they are not fluid, but fixed Parts, I shall choose rather to call them, *Bladders*. As they appear through a good *Glass*, in a piece of *Burdock*, See in *Tab. 3.* But a more particular Description of the *Sizes*, *Figures*, and admirable *Textures* hereof, I have given in several places in the following *Books*.

19. §. Besides what this Observation informs us of here, it farther confirms what in the *Second Chapter* we have said of the Original of the *Pith* and *Cortical Body*, and of the sameness of both their natures with the *Parenchyma* of the *Seed*: which is nothing else but a *Mass* of *Bladders*; as in the *First Chapter* hath been said.

Tab. 3. f. 6.
C. 2. §. 3, 16, 17.
C. 1. §. 18:
& Tab. 1. f. 9.

20. §. In the *Piths* of many *Plants*, the greater *Pores* or *Bladders* have some of them lesser ones within them, and some of them are divided with cross *Membranes*: And betwixt their several sides, have, I think, other smaller *Bladders* visibly interjected. However, that they are all permeable, is most certain. They stand together not confusedly, but in even *Ranks* or *Trains*; as those of the *Insertions* by the breadth, so these by the length of the *Trunk*. And thus far there is a general corresponding betwixt the parts of the *Root* and *Trunk*. Yet are there some considerable *Disparities* betwixt them; wherein, and how they come to pass, and to what especial *Use* and *End*, shall next be said.

An Account
of the
Growth of
the *Trunk*.

21. §. WE SAY then, that the *Sap* being in the *Root* by *Filtrations*, *Fermentations* (and in what *Roots* needful, perhaps by *Circulation* also) duly prepar'd; the prime part thereof passing through the intermediate *Coarcture*, in due moderation and purity is entertain'd at last into the *Trunk*. And the *Sap* of the *Trunk* being purer and more volatile, and so it self apt to ascend; the motion of the *Trunk* likewise will be more noble, receiving a disposition and tendency to ascend therewith. And what by the *Sap* the *Trunk* is in part dispos'd to, by the respective position and quantity of its *Parts* it is effectually enabled. For whereas in the *Root* the *Lignous Body* being in proportion with the *Cortical*, but little, and all lying close within its *Center*; it must therefore needs be under its controul: on the contrary, being here comparatively of greater quantity, and also more dilated, and having divers of its *Branches* standing more abroad towards the *Circumference*, as both in the *Leaves* and *Body* of the young *Trunk* and *Plume*, is seen; it will in its own *magnetical* tendency to ascend, reduce the *Cortical Body* to a compliance with it.

22. §. And the *Trunk* thus standing from under the restraint of the *Ground* in the open *Air*, the disposition of its *Parts*, originally different from that of the *Parts* in the *Root*, will not only be continued, but improved. For by the force and pressure of the *Sap* in its collateral *Motion*, the *Lignous Body* will now more freely and farther be dilated.

lated. And this being dilated, the *Cortical Body* also, must needs be *inserted*; and is therefore in proportion always, more or less, smaller here in the *Trunk*, than in the *Root*. And as the *Cortical Body* lessens, so the *Pith* will be enlarged, and by the same proportion is here greater. And the *Pith* being enlarged it self, its *Pores* (the *Lignous Body*, upon its dilatation, as it were tentering and stretching out all their sides) must needs likewise be enlarged with it; and accordingly, are ever greater in the *Pith* of the *Trunk*, than of the *Root*. And the dilatation of the *Lignous Body* still continued, it follows, that whereas the *Pith* descendent in the *Root*, is not only in proportion less and less, but also in the smaller extremities thereof, and sometimes higher, altogether absent: Contrariwise, in the *Trunk*, it is not only continued to its top and smallest *Twigs*, but also there, in proportion, equally ample with what it is in any other inferiour part.

23. §. But although the openness of the Aer permitting, be always alike; yet the Energy of the *Sap* effecting, being different; as therefore that doth, the dilatation of the *Trunk*, will also vary. If that be less, so is this; as in the *Trunks* of most *Trees*: If that be greater, so this; as in *Herbs* is common; the *Lignous Body* being usually so far dilated, that the utmost *Shootings* thereof may easily be seen to jut out, and adjoyn to the *Skin*. And if the *sap* be still of greater energy, it so far dilates the *Lignous Body*, as not only to amplify the *Pith* and all its *Pores*; but also so far to stretch them out, as to make them tear. Whereupon either running again into the *Cortical Body*, or shrinking up towards it, the *Trunk* thus sometimes becomes an *hallow Stalk*, the *Pith* being wholly, or in part voided. But generally it keeps entire; and where it doth, the same proportion and respect to the *Lignous* and *Cortical Bodies*, as is said. The Consequences of all which will be, the *Strength* of the *Trunk*, the *Security* and *Plenty* of the *Sap*, its *Fermentation* will be quicker, its *Distribution* more effectual, and its *Advancement* more sufficient.

24. §. First, the Erect Growth and Strength of the *Trunk*; this being, by the position of its several Parts, effected: for besides the slendering of the *Trunk* still towards the top, the *Circumferential* position of the *Lignous Body*, likewise is, and that eminently, hereunto subservient. So that as the *Lignous Body*, in the smaller parts, of the *Root* standing Central, we may thence conceive and see their pliableness to any oblique motion; so here, on the contrary, the *Lignous Body* standing wide, it thus becomes the Strength of the *Trunk*, and most advantageous to its Perpendicular Growth. We see the same Design in *Bones* and *Feathers*: The strongest *Bones*, as those in the *Legs*, are hollow. Now should we suppose the same *Bone*, to be contracted into a Solid; although now it would be no heavier, and in that respect, as apt for motion; yet would it have far less strength, than as its Parts are dilated to a *Circumferential* posture. And so for *Quills*, which, for the same Reasons, in subserviency to flying, as they are exceeding light; So, in comparison with the thinness of their *Sides*, they are very strong, and much less apt to bend, than if contracted into a Solid *Cylinder*. We see it not only in *Nature*, but *Art*. For hence it is, that *Joyners* and *Carpenters* unite and set together their Timber-pieces and several Works oftentimes with double *Joynts*; which, although

though they are no thicker, than a single one might be made; yet standing at a distance, have a greater strength than That could have. And the same Architecture, will have the same use, in the *Trunks* of *Plants*; in most whereof 'tis very apparent; as for instance, in *Corn*. For *Nature* designing its *Sap* a great Ascent; for its higher maturity, hath given it a tall *Trunk*: But to prevent its ravenous despoiling either of the *Ear*, or *Soyl*; although it be tall, yet are its sides but thin: And because again, it should grow not only tall and thriftily, but for avoiding propping up, strongly too; therefore, the same proportion as its height bears, to the thinness of its sides, doth the greatness of its Circumference also; being so far dilated as to parallel a *Quil* it self.

25. §. Besides the position of the *Lignous Body* within the compass of a *Ring*, there are some *shootings* thereof, often standing beyond the Circumference of the said *Ring*, making sometimes a triangular, oftner a quadrangular *Body* of the *Trunk*. To the end, that the *Ring*, being but thin, and not self-sufficient, these, like *Splinters* to *Bones*, might add strength and stability to it.

26. §. Next, the security and plenty of the *Sap*. For should the *Lignous Body*, as it doth in the smaller *Parts* of the *Root*, stand Central here also, and so the *Cortical* wholly surround it: the greater part of the *Sap* would thus be more immediately expos'd to the *Sun* and *Aer*; and being lodged in a laxe *Body*, by them continually be prey'd upon, and as fast as supplied to the *Trunk*, be exhausted. Whereas, the *Pith* standing in the Center, the *Sap* therein being not only most remote from the *Aer* and *Sun*, but by the *Barque*, and especially the *Wood*, being also surrounded and doubly immur'd, will very securely and copiously be convey'd to all the *Collateral Parts*, and (as shall be said how) the top of the *Trunk*.

27. §. And the *Sap* by the amplitude, and great porosity of the *Pith*, being herein more copious, its Fermentation also will be quicker; which we see in all *Liquors*, by standing in a greater quantity together, proceeds more kindly: And being tun'd up within the *Wood*, is at the same time not only secur'd from loss, but all extream mutations; the *Day* being thus, not too hot; nor the *Night*, too cold for it.

28. §. And the Fermentation hereof being quicker, its motion also will be stronger, and its *Distribution* more effectual, not only to the dilatation of the *Trunk*, but likewise the shooting out of the *Branches*. Whence it is, that in the *Bodies* of *Trees*, the *Barque* of it self, though it be Sappy, and many *Fibres* of the *Lignous Body* mixed with it, yet seldom sendeth forth any; and that in *Herbs*, those with the least *Pith* (other advantages not supplying this defect) have the fewest or smallest *Branches*, or other *collateral Growths*: and that *Corn*, which hath no *Pith*, hath neither any *Branches*.

29. §. Lastly, the Advancement of the *Sap* will hence also be more ready and sufficient. For the understanding where, and how, we suppose, That in all *Trunks* whatsoever there are two *Parts* joyntly hereunto subservient. In some, the *Lignous Body* and the *Cortical*, as in older *Trunks*; the *Pith* being either excluded, or dried: But in most, principally, the *Lignous Body* and *Pith*; as in most *Annual Growths* of *Trees*; but especially *Herbs*, where the *Cortical Body* is usually much and often wholly *Inserted*.

30. §. Of the *Lignous Body* it is so apparent by its *Pores*, or rather by its *Vessels*, that we need no farther Evidence. For to what end are *Vessels*, but for the conveyance of *Liquor*? And is that also, which upon cutting the young Branch of a *Sappy Tree* or *Herb*, by an accurate and steady view may be observed. But when I say the *Vessels* of the *Lignous Body*, I mean principally them of the *younger shootings*, both those which make the *new Ring*, and those which are mixed with the *Cortical Body* in the *Barque*: that which ascendeth by the *Pores* or *Vessels* of the *Wood*, being probably, because in less quantity, more in form of a *Vapour*, than a *Liquor*. Yet that which drenching into the sides of its *Pores*, is with all thereunto sufficient Aliment; as we see *Orpine*, *Onions*, &c. only standing in a moyster Aer will often grow. And being likewise in part supplied by the *Insertions* from the younger *Shoots*: But especially because as it is but little, so (considered as Aliment) it serveth only for the growth of the *Wood*, and no more; whereas, the more copious Aliment ascendent by the younger *Shoots*, subserves not only their own growth, but the generation of others; and is besides with that in the *Cortical Body* the Fountain of *Perspirations*, which we know even in *Animals* are much more abundant than the *Nutritive Parts*; and doubtless in a *Vegetable* are still much more.

31. §. But these *Pores*, although they are a free and open way to the ascending *Sap*; yet that meer *Pores* or *Vessels* should be able of themselves to advance the *Sap* with that speed, strength and plenty, and to that height, as is necessary, cannot probably be supposed. It follows then, that herein we must grant the *Pith* a joynt service. And why else is the *Pith* in all *Primitive Growths* the most *Sappy* part, why hath it so great a stock of *Sap*, if not, after due maturation within it self, still to be disbursed into the *Fibres* of the *Lignous Body*? Why are the *Annual Growths* of all both *Herbs* and *Trees*, with great *Piths*, the quickest and the longest? But how are the *Pores* or *Bladders* of the *Pith* permeable? That they are so, both from their being capable of a repletion with *Sap*, and of being again wholly emptied of it, and again, instead thereof fill'd with *Aer*, is as certain as that they are *Pores*. That they are permeable, by the breadth, appears from the dilatation of the *Lignous Body*, and from the production of *Branches*, as hath been, and shall hereafter be said. And how else is there a *Communion* betwixt *This* and the *Cortical Body*? That they are so also, by the length, is probable, because by the best *Microscope* we cannot yet observe, that they are visibly more open by the breadth, than by the length. And withall are ranked by the length, as those of the *Insertions* by the breadth of the *Trunk*. But if you set a piece of dry *Elder-Pith* in some tinged *Liquor*, why then doth it not penetrate the *Pores*, so as to ascend through the *Body* of the *Pith*? The plain reason is, because they are all fill'd with *Aer*. Whereas the *Pith* in a *Vegetating Plant*, as its *Parts* or *Bladders* are still generated, they are at the same time also fill'd with *Sap*; which, as 'tis gradually spent, is still repaired by more succeeding, and so the *Aer* still kept out; as in all *Primitive Growths*, and the *Pith* of *Elder* it self: Yet the same *Pith*, by reason of the following *Winter*, wanting a more copious and quick supply of *Sap*, thus once become, ever after keeps dry. And since in the aforesaid *Trial* the *Liquor* only ascends by the sides of the *Pith*, that

is of its broken *Bladder*, we should thence by the same reason conclude that they are not penetrable by the breadth neither, and so no way; and then it need not be ask'd what would follow. But certainly the *Sap* in the *Bladders* of the *Pith* is discharged and repaired every moment, as by its shriv'ling up, upon cutting the *Plant*, is evident.

32. §. We suppose then, that as the *Sap* ascendeth into the *Trunk* by the *Lignous Body*, so partly also by the *Pith*. For a piece of *Cotton* with one end immers'd in some tinged *Liquor*, and with the other erect above, though it will not imbibe the *Liquor* so far as to overrun at the top, yet so as to advance towards it, it will. So here, the *Pith*, being a porous and spongy *Body*, and in its *Vegetating* state, its *Pores* or *Bladders* being also permeable, as a curious *Filtre* of *Natures* own contrivance, it thus advanceth, or as people use to say, sucks up the *Sap*. Yet as it is seen of the *Liquor* in the *Cotton*; so likewise are we to suppose it of the *Sap* in the *Pith*; that though it riseth up for some way, yet is their some term, beyond which it riseth not, and towards which the motion of the ascending *Sap* is more and more broken, weak and slow, and so the quantity thereof less and less. But because the *Sap* moveth not only by the length, but breadth of the *Pith*; at the same time therefore as it partly ascendeth by the *Pith*, it is likewise in part pressed into the *Lignous Body* or into its *Pores*. And since the motion of the *Sap* by the breadth of the *Pith* not being far continued, and but collateral, is more prone and easie, than the perpendicular, or by its length; it therefore follows, that the collateral motion of the *Sap*, at such a height or part of the *Pith*, will be equally strong with the perpendicular at another part, though somewhat beneath it; and that where the perpendicular is more broken and weak, the collateral will be less; and consequently where the perpendicular tendency of the *Sap* hath its term, the collateral tendency thereof, and so its pressure into the *Pores* or *Vessels* of the *Lignous Body*, will still continue. Through which, in that they are small, and so their sides almost contiguous, the *Sap* as fast as pressed into them willeasily run up; as in very small *Glasse Pipes*, or betwixt the two halves of a *Stick* first slit, and then tyed somewhat loosely together, may also any *Liquor* be observed to do. By which Advantage the facility and strength of that ascent will be continued higher in the said *Vessels*, than in the *Pith*. Yet since this also, as well as that in the *Pith* will have its term; the *Sap*, although got thus far, would at last be stagnant, or at least its ascent be very sparing, slow and feeble, if not some way or other re-inforced. Wherefore, as the *Sap* moving by the breadth of the *Pith*, presseth thence into the *Vessels* of the *Lignous Body*; so having well fill'd these, is in part by the same *Collateral* motion disburfed back, into a yet higher *Region* of the *Pith*. By which partly, and partly, by that portion of the *Sap*, which in its perpendicular ascent was before lodged therein; 'tis thus here, as in any inferiour place equally replenished. Whereupon the force and vigour of the perpendicular motion of the *Sap* herein, will likewise be renew'd; and so its *Collateral* motion also, and so its pressure into the *Vessels* of the *Lignous Body*, and consequently its ascent therein: and so by a pressure, from these into the *Pith*, and from the *Pith* into these, reciprocally carried on; a most ready and copious ascent of the *Sap* will be continued, from the bottom to the top, though of the highest *Tree*. An

An Appendix.

Of Trunk-Roots and Claspers.

THE distinct *Parts* whereof these are composed, are the same with those of the *Trunk*, and but the continuation of them.

1. §. *Trunk-Roots* are of two kinds: Of the one, are those that vegetate by a direct descent: The place of their Eruption is sometimes all along the *Trunk*; as in *Mint*, &c. Sometimes only at its utmost point, as in the *Bramble*.

2. 2. The other sort are such as neither ascend nor descend, but shoot forth at right Angles with the *Trunk*; which therefore, though as to their *Office*, they are true *Roots*, yet as to their *Nature*, they are a *Middle Thing* betwixt a *Root* and a *Trunk*.

3. §. *Claspers*, though they are but of one kind, yet their Nature is double; not a mean betwixt that of the *Root* and that of the *Trunk*, but a compound of both; as in their *Circumvolutions*, wherein they often mutually ascend and descend, is seen.

4. §. The use of these *Parts* may be observed as the *Trunk* Mounts; or as it Trails. In the mounting of the *Trunk*, they are for Support and Supply. For Support, we see the *Claspers* of *Vines*: the *Branches* whereof being very long, fragile and slender; unless by their *Claspers*, they were mutually contain'd together, they must needs by their own weight, and that of their Fruit, undecently fall; and be also liable to frequent breaking. So that the whole care is divided betwixt the Gardener and Nature; the Gardener, with his Ligaments of Leather, secures the main *Branches*; and Nature, with these of her own finding, secures the *Leaves*. Their Conveniency to which end, is seen in their *Circumvolutions*, a motion, not proper to any other *Part*: As also in their toughness, though much more slender than the *Branches* whereon they are appendent.

5. §. The *Claspers* of *Bryony* have a retrograde motion about every *Third Circle*, to the form a *Doublet-Clasp*. Probably for the more certain hold; which, if it miss one way, it may be sure to take another.

6. §. For Supply, we see the *Trunk-Roots* of *Ivy*. For mounting very high, and being of a closer or more compact Substance than that of a *Vine*; the *Sap* could not be sufficiently supplied to the upper *Sprouts*, unless these, to the *Mother-Root*, were jointly assistant. Yet serve they for support likewise; whence they shoot out, not as in *Cresses*, *Brook-lime*, &c. reciprocally on each side, but commonly, all on one; that so they may be fastned at the nearest hand.

7. §. In the Trailing of the *Trunk*, they serve for stabiliment, propagation and shade. For stabiliment, the *Claspers* of *Cucumbers* are of good use. For the *Trunk* and *Branches* being long and fragile, the *Brushes* of the Winds would injuriously hoise them to and fro, to the damage both of themselves and their tender Fruits; were

they not by these Ligaments brought to good Association and Settlement.

8. §. As for this end, so for Propagation, the *Trunk-Roots* of *Chamemile* do well serve. Whence we have the reason of the common observation, that it grows better by being trod upon: the Mould, where too laxe, being thus made to lie more conveniently about the said *Trunk-Roots* newly bedded therein; and is that which is sometimes also effected in Rowling of *Corn*.

9. §. For both these ends, Serve the *Trunk-Roots* of *Strawberries*; as also for shade; for in that all *Strawberries* delight; and by the trailing of the *Plant* is well obtain'd. So that as we are wont to tangle the *Twigs* of *Trees* together to make an *Arbour Artificial*; the same is here done to make a *Natural one*: as likewise by the *Classpers* of *Cucumbers*. For the *Branches* of the one by the Linking of their *Classpers*, and of the other by the Tethering of their *Trunk-Roots*, being couched together; their tender Fruits thus lie under the Umbrage of a *Bower* made of their own Leaves.

CHAP. IV.

Of the GERMEN, BRANCH, and LEAF.



THE Parts of the *Germen* and *Branch*, are the same with those of the *Trunk*; the same *Skin*, *Cortical* and *Lignous Bodies*, *Insertment* and *Pith*, hereinto propagated, and distinctly observable herein.

2. §. For upon Enquiry into the Original of a *Branch* or *Germen*, it appears, That it is not from the *superficies* of the *Trunk*; but so deep, as to take, with the *Cortical*, the *Lignous Body* into it self: and that, not only from its Circumference, but from in *Inner* or *Central Parts*; So as to take the *Pith* in also. Divers of which *Parts* may commonly be seen to shoot out into the *Pith*; from which *Shoots*, the surrounding and more superior *Germens* are originated; in like manner as the Succulent Part of the *Lignous Body* of the *Trunk* is sometimes principally from those *Fibrous shoots* which run along the *Pith* in the *Root*.

3. §. The manner wherein usually the *Germen* and *Branch* are fram'd, is briefly thus: The *sap* (as is said, *Chap. 3.*) mounting in the *Trunk*, will not only by its length, but by its breadth also, through the *Insertions* partly move. Yet, its *Particles* being not all alike qualified, in different degrees. Some are more gross and sluggish; of which we have the formation of a *Circle of Wood* only, or of an *Annual Ring*. Others are more brisk; and by these, we have the *Germen* propagated. For by the vigour of their own motion from the Center, they impress an equal tendency on some of the inner *Portions* of the *Lignous Body* next adjacent

cent to the *Pith*, to move with them. And since the *Lignous Body* is not entire, but frequently disparted; through these *Dispartments*, the said interior *Portions*, upon their Nutrition, actually shoot; not only towards the Circumference, so as to make part of a *Ring*; but even beyond it, in order to the production of a *Germen*. And the *Lignous Body* thus moving, and carrying the *Cortical* along with it; they both make a force upon the *Skin*. Yet their motion being most even and gradual, that force is such likewise; not to cause the least breach of its parts, but gently to carry it on with themselves; and so partly, by the extension of its already existent parts, as of those of *Gold* in drawing of *Gilded Wyer*; and partly, by the accretion of new ones, as in the enlarging of a *Bubble* above the Surface of the Water; it is extended with them to their utmost growth. In which growth, the *Germen* being prolonged, and so displaying its several parts, as when a *Prospective* or *Telescope* is drawn out, thus becomes a *Branch*.

4. §. The same way as the propagation of the *Parts* of a *Germen* is contriv'd, is its due nutrition also. For being originated from the inner part of the *Lignous Body*, 'tis nourished with the best fermented *Sap* in the *Trunk*, *sc.* that next adjacent to it in the *Pith*. Besides, since all its *Parts*, upon their shooting forth, divaricate from their perpendicular, to a cross Line, as these and the other grow and thrive together, they bind and throng each other into a *Knot*: through which *Knot* the *Sap* being strain'd, 'tis thus, in due moderation and purity delivered up into the *Branch*.

5. §. And for *Knots*, they are so necessary, as to be seen not only where collateral *Branches* put forth; but in such *Plants* also, as shoot up in one single *Trunk*; as in *Corn*. Wherein, as they make for the strength of the *Trunk*; so by so many percolations, as they are *Knots*, for the transmission of the *Sap* more and more refined towards the *Ear*. So that the two general uses of *Knots* are, For *finer standing*, and *finer growth*.

6. §. Lastly, as the due Formation and Nutrition of the *Germen* are provided for, so is its security also; which both in its position upon the *Trunk*, and that of its *Parts* among themselves, may be observed: The position of its *Parts* shall be considered in speaking of the *Leaf*. As to its standing in the *Trunk*, 'tis alwayes betwixt the *trunk* or *older Branch*, and the *Basis* of the *Stalk* of a *Leaf*; whereby it is not only guarded from the Injuries of any contingent Violence; but also from the more piercing assaults of the Cold; so long, till in time 'tis grown larger, and more hardy. The maner and uses of the position of every *Germen*, considered as after it becomes a *Branch*; hath already been, by the Ingenious Mr. *Sbarrock* Hist. of the Prop. of Veget. very well observed; to whom I refer.

7. §. UPON THE prolongation of the *Germen* into a *Branch*, its *Leaves* are thus display'd. The *Parts* whereof are substantially the same with those of a *Branch*. For the *Skin* of the *Leaf*, is only the ampliation of that of the *Branch*; being partly by the accretion of new, and partly the extension of its already existent parts, dilated (as in making of *Leaf-Gold*) into its present breadth. The *Fibres* or *Nerves*, dispersed through the *Leaf*, are only the Ramifications of the *Branch's Wood*, or *Lignous Body*. The *Parenchyma* of the *Leaf*; which

which lies betwixt the *Nerves*, and as in Gentlewomens Needle-works, fills all up, is nothing else, but the continuation of the *Cortical Body*, or *Parenchymous* part of the *Barque* from the *Branch* into it self, as in most *Plants* with a thick *Leaf*, may easily be seen.

8. §. The *Fibers* of the *Leaf* neither shoot out of the *Branch*, or the *Trunk*, nor stand in the *Stalk*, in an *even Line*; but alwayes in either an *Angular* or *Circular* posture; and usually making either a *Triangle*, or a *Semi-Circle*, or *Chord* of a *Circle*; as in *Cichory*, *Endive*, *Cabbage*, &c. may be observed. And if the *Leaf* have but one main *Fiber*, that also is postur'd in a bowed or *Lunar Figure*; as in *Mint* and others. The usual number of these *Vascular Threds* or *Fibres* is 3, 5, or 7.

Tab. 4. f. 2.
to f. 11.

9. §. The reason of the said Positions of the *Fibers* in the *Stalk* of the *Leaf*, is for its more *Erect* growth, and greater *Strength*: which, were the position of the said *Fibers* in an *even Line*, and so the *Stalk* it self, as well as the *Leaf*, flat; must needs have been defective; as from what we have said of the *Circumferential* posture of the *Lignous Body* in the *Trunk*, we may better conceive.

C. 3. §. 24.

10. §. As likewise for the security of its *Sap*: For by this means it is, that the several *Fibers*, and especially the main or middle *Fiber* of the *Leaf*, together with a considerable part of the *Parenchyma*, are so disposed of, as to jut out, not from its upper, but its back, or neither *Side*. Whence the whole *Leaf*, reclining backward, becomes a *Canopy* to them, defending them from those *Injuries* which from colder *Blasts*, or an hotter *Sun*, they might otherwise sustain. So that by a mutual benefit, as These give *suck* to all the *Leaf*, so that again *protection* to These.

11. §. These *Fibers* are likewise the immediate *Visible Cause* of the *Shape* of the *Leaf*. For if the nethermost *Fiber* or *Fibres* in the *Stalk* (which thence runs chiefly through the length of the *Leaf*) be in proportion greater, the *Leaf* is long; as in *Endive*, *Cichory*, and others: If all of a more equal size, it spreads rounder, as in *Ivy*, *Doves-foot*, *Colts foot*, &c. And although a *Dock-Leaf* be very long, whose *Fibers* notwithstanding, as they stand higher in the *Stalk*, are disposed into a *Circle* all of an equal size; yet herein one or more peculiar *Fibres*, standing, in or near the *Center*, betwixt the rest, and running through the length of the *Leaf*, may be observed.

Tab. 4.

12. §. In correspondence also to the size and shape of these *Fibres*, is the *Leaf* flat. In that either they are very small, or if larger, yet they never make an entire *Circle* or *Ring*; but either half of one, as in *Borage*, or at most three parts of one, as in *Mullen*, may be seen. For if either they were so big, as to contain, or so entire, as perfectly to include a *Pith*, the *Energy* of the *Sap* in that *Pith*, would cause the said *Lignous Ring* to shoot forth on every side, as it doth in the *Root* or *Trunk*: But the said *Fibers* being not figur'd into an entire *Ring*, but so as to be open; on that hand therefore where open they cannot shoot any thing directly from themselves, because there they have nothing to shoot; and the *Sap* having also a free vent through the said opening, against that part therefore which is thereunto opposite, it can have no force; and so neither will they shoot forth on that hand; and so will they consequently, that way only, which the force of the *Sap* directs, which is only on the right and left.

Tab. 4.

13. §.

13. §. The several *Fibers* in the *stalk*, are all Inosculated in the *Leaf*, with very many Sub-divisions. According as these *Fibers* are Inosculated near, or at, or shoot directly to the edge of the *Leaf*, is it Even, or Scallop'd. Where these *Inosculations* are not made, there we have no *Leaves*, but only a company of *Filaments*; as in *Fennel*.

14. §. To the *Formations* of *Leaves*, the *Fouldings* immediately follow. And sometimes they have one *Date*, or are the contemporary works of *Nature*; each *Leaf* obtaining its distinct shape, and proper posture together; both being perfect, not only in the outer, but Central and minutest *Leaves*, which are five hundred times smaller than the outer: both which in the Cautious opening of a *Germen* may be seen.

15. §. Nor is there greater Art in the *Forms*, than in the *Foulds* or *Postures* of *Leaves*; both answerably varying, as this or that way they may be most agreeable. Of the *Quincuncial* posture, so amply instanc'd in by the Learned Sir *Thomas Brown*, I shall omit to speak. Others there are, which though not all so universal, yet equally necessary where they are, giving two general advantages to the *Leaves*, *Elegancy* and *Security*, *sc.* in taking up, so as their *Forms* will bear, the least room; and in being so conveniently couch'd, as to be capable of receiving protection from other *Parts*, or of giving it one to another; as for instance,

Treat. of the
Quincunx.

16. §. First, There is the *Bow-Lap*, where the *Leaves* are all laid somewhat convexly one over another, but not plaited; being to the length, breadth and number of *Leaves* most agreeable; as in the Buds of *Pear-tree*, *Plum-tree*, &c. But where the *Leaves* are not so thick set, as to stand in the *Bow-Lap*, there we have the *Plicature*, or the *Flat-Lap*; as in *Rose-Tree*, *Strawberry*, *Cinquefoyl*, *Burnet*, &c. For the *Leaves* being here plaited, and so lying in half their breadth, and divers of them thus also collaterally set together; the thickness of them all, and half their breadth, are much alike dimensions; by which they stand more secure within themselves, and in better consort with other *Germen-Growths* in the same *Truss*. If the *Leaves* be much indented or jagg'd, now we have the *Duplicature*; wherein there are divers *Plaits* in one *Leaf*, or *Labels* of a *Leaf*, but in distinct *Sets*, a lesser under a greater; as in *Souchus*, *Tansy*, &c. When the *Leaves* stand not collaterally, but single; and are moreover very broad; then we have the *Multiplicature*; as in *Gooseberries*, *Mallows*, &c. the *Plaits* being not only divers in the same *Leaf*, but of the same *Set* continuant, and so each *Leaf* gather'd up in five, seven or more *Foulds*, in the same manner as our *Gentlewomen's Fans*. Where either the thickness of the *Leaf* will not permit a *Flat-Lap*, or the fewness of their number, or the smallness of their *Fibers*, will allow the *Rowl*, there This may be observed. Which is sometimes single, as in *Bears-Ears*, *Arum*, *Flammula*, *Jerusalem Cowslip*, &c. Sometimes double, the two *Rowls* beginning at each edge of the *Leaf*, and meeting in the middle. Which again, is either the *Fore-Rowl*, or the *Back-Rowl*. If the *Leaf* be design'd to grow long, now we have the *Back-Rowl*, as in *Docks*, *Sorrels*, and the rest of this Kindred: as also in *Primrose*, and other like *Plants*. For the main *Fibers*, and therewith a considerable part of the *Cortical Body* standing prominent from the *Back-side* of the *Leaf*, they thus stand securely couch'd up be-

twixt

twixt the two *Rowls*; on whose security the growth of the *Leaf* in length depends. But those of *Bears-Ears*, *Violets*, *Doves Foot*, *Warden*, and many more, upon contrary respects, are rowled up inwards. Lastly, there is the *Tre-Rowl*, as in *Fern*; the *Labels* whereof, though all rowled up to the *main Stem*, yet could not stand so firm and secure from the Injuries either of the *Ground* or *Weather*, unless to the *Rowls* in breadth, that by the length were super-induc'd; the *Stalk* or *main Stem* giving the same Protection here, which in other *Plants* by the *Leaves*, or some particular *Mantling*, is contriv'd. These, and other *Foulds*, See in the *Figures* belonging to the *First Part* of the *Fourth Book*.

17. §. According to the *Form* and *Foulding* of every *Leaf* or *Germen*, is its Protection order'd; about six ways whereof may be observ'd; *sc.* by *Leaves*, *Surfoyls*, *Interfoyls*, *Stalks*, *Hoods* and *Mantlings*. To add to what we have above given, one or two Instances. Every *Bud*, besides its proper *Leaves*, is covered with divers *Leafy Pannicles* or *Surfoyls*; which, what the *Leaves* are to one another, are that to them all: For not opening except gradually, they admit not the *Weather*, *Wet*, *Sun* or *Aer*, to approach the *Leaves*, except by degrees respondent, and as they are gradually inur'd to bear them. Sometimes, besides *Surfoyls*, there are also many *Interfoyls* set betwixt the *Leaves*, from the *Circumference* to the *Center* of the *Bud*; as in the *Hasel*. For the *Fibres* of these *Leaves* standing out so far from a plain surface; they would, if not thus shelter'd, lie too much expos'd and naked to the *Severities* of the *Weather*. Where none of all the Protections above-named, are convenient, there the *Membranes* of the *Leaves* by continuation in their first forming (together with some *Fibres* of the *Lignous Body*) are drawn out into so many *Mantles* or *Veils*; as in *Docks*, *Snakeweed*, &c. For the *Leaves* here being but few, yet each *Leaf* and its *Stalk* being both exceeding long; at the bottom whereof the next following *Leaf* still springs up; the form and posture of all is such, as supercedes all the other kinds of Protection, and so each *Leaf* apart is provided with a *Veil* to it self. These, and other *Protections*, See in the *Figures* belonging to the *First Part* of the *Fourth Book*.

18. §. The *Uses* of the *Leaves*, I mean in respect of their service to the *Plant* it self, are these: First, for Protection; which, besides what they give one to another, they afford also to the *Flower* and *Fruit*. To the *Flower* in their *Foulds*; that being, for the most part, born and usher'd into the open *Aer* by the *Leaves*. To the *Fruit*, when afterwards they are display'd, as in *Strawberries*, *Grapes*, *Rasps*, *Mulberries*, &c. On which, and the like, should the *Sun-Beams* immediately strike, especially while they are young, they would quite shrivel them up; but being by the *Leaves* screened off, they impress the circumjacent *Aer* so far only as gently to warm the said *Fruits*, and so to promote their *Fermentation* and *Growth*. And accordingly we see, that the *Leaves* above-named are exceeding large in propotion to the *Fruits*: whereas in *Pear-trees*, *Apple-trees*, &c. the *Fruit* being of a solider *Parenchyma*, and so not needing the like protection, are usually equal with, and often wider in Diameter than the *Leaves*.

19. §. Another use is for Augmentation ; or, the capacity for the due spreading and ampliation of a *Tree* or other *Plant*, are its *Leaves*. For herein the *Lignous Body* being divided into small *Fibres*, and these running all along their lax and spongie *Parenchyma*; they are thus a *Body* fit for the imbibition of *Sap*, and easie *Growth*. Now the *Sap* having a free reception into the *Leaves*, it still gives way to the next succeeding in the *Branches* and *Trunk*, and the voyding of the *Sap* in these, for the mounting of that in the *Root*, and ingress of that in the *Ground*. But were there no *Leaves* to make a free reception of *Sap*, it must be needs be stagnant in all the *Parts* to the *Root*, and so the *Root* being clogg'd, its fermenting and other Offices will be voyded, and so the due *Growth* of the whole. As in the motion of a *Watch*, although the original term thereof be the *Spring*, yet the capacity for its continuance in a due measure throughout all the *Wheels*, is the free and easie motion of the *Ballance*.

20. §. Lastly, As the *Leaves* subsERVE the more copious advancement, so the higher purity of the *Sap*. For this being well fermented both in the *Root*, and in its Ascent through the *Trunk*, and so its *Parts* prepar'd to a farther separation; the grosser ones are still deposited into the *Leaves*; the more elaborate and essential only thus supplied to the *Flower*, *Fruit* and *Seed*, as their convenient *Aliment*. Whence it is, that where the *Flowers* are many and large, into which the more odorous *Particles* are copiously receiv'd, the green *Leaves* have little or no smell; as those of *Rose-tree*, *Carnations*, *French-Marigold*, *Wood-bind*, *Tulips*, &c. But on the contrary, where the *Flowers* are none, or small, the green *Leaves* themselves are likewise of a strong favour; as those of *Wormwood*, *Tansie*, *Baum*, *Mint*, *Ruc*, *Geranium Moschatum*, *Angelica*, and others.

An Appendix.

Of Thorns, Hairs and Globulets:

THorns are of two kinds, *Lignous* and *Cortical*. Of the first are such as those of the *Hawthorn*, and are constituted of all the same substantial *Parts* whereof the *Germen* or *Bud* it self, and in a like proportion: which also in their Infancy are set with the resemblances of divers minute *Leaves*. Of affinity with these are the *Spinets* or *Thorny Prickles* upon the Edges and Tops of divers *Leaves*, as of *Barbery*, *Holly*, *Thistle*, *Furze*, and others; all which I think are the filamentous extremities of the *Lignous Body* sheathed in the *Skin*. But this principal difference betwixt a *Bud* and these *Lignous Thorns*, is observable; That the *Bud* hath its Original from the Inner part of the *Lignous Body*, next the *Pith*: But these *Thorns*, from the outer, and less fecund Part; and so produceth no *Leaves*, but is, as it were, the *Mola* of a *Bud*.

2. §. *Cortical Thorns* are such as those of the *Raspberry Bush*, being not, unless in a most extraordinary small and invisible proportion propagated from the *Lignous Body*, but as, it seems, wholly from the *Cortical* and *Skin*, or from the exterior part of the *Barque*.

C. 2. §. 25. 3. §. The *Growth* of this *Thorn* may farther argue what in the *second Chapter* we supposed; *sc.* That as the proper *Tendency* of the *Lignous Body*, is to *Ascend*; so of the *Cortical* to *Descend*. For as the *Lignous Thorn*, like other *Parts* of the *Trunk*, in its *Growth* ascends; This, being almost wholly *Cortical*, pointeth downward. The use of *Thorns* the Ingenious Mr. *Sharrock* hath observed.

Hist. of the Prop. of Veget. 4. §. Upon the *Leaves* of divers *Plants* two *Productions* shew themselves, *sc.* *Hairs* and *Globulets*. Of *Hairs*, only one kind is taken notice of; although they are various. Ordinarily they are of a *Simple Figure*; which when fine and thick set, as on most *Hairy Buds*; or fine and long, as on those of the *Vine*, we call them *Down*.

5. §. But sometimes they are *Branched out*, from the bottom to the top, reciprocally on every side, in some resemblance to a *Stags Horn*; as in *Mullen*. And sometimes they are *Astral*, as upon *Lavender*, and some other *Leaves*, and especially those of *Wild Olive*; wherein every *Hair* rising in one round entire *Basis* a little way above the surface of the *Leaf*, is then disparted, *Star-like*, into several, four, five or six *Points*, all standing at right Angles with the said perpendicular *Basis*.


6. §. The *Uses* of *Hairs* are for *Distinction* and *Protection*. That of *Distinction* is but secondary, the *Leaves* being grown to a considerable size. That of *Protection* is the prime, for which they were originally form'd together with the *Leaves* themselves, and whose service they enjoy in their *Infant-estate*: For the *Hairs* being then in form of a *Down*, always very thick set, thus, give that *Protection* to the *Leaves*, which their exceeding tenderness then requires; so that they seem to be vested with a *Coat of Frize*, or to be kept warm, like young and dainty *Chickens*, in *Wool*.

7. §. *Globulets* are seen upon *Orach*, both *Garden* and *Wild*; and yet more plainly on *Mercury* or *Bonus Henricus*. In these, growing almost upon the whole *Plant*, and being very large, they are by all taken notice of.

8. §. But strict *Observation* discovers, that these *Globulets* are the natural and constant *Off-spring* of very many other *Plants*. Both these *Globulets*, and likewise the diversity of *Hairs*, I find that Mr. *Hook* hath also observed. I take notice, that they are of two kinds; *Transparent*, as upon the *Leaves* of *Hysop*, *Mint*, *Baume*, and many more *White*, as upon those of *Germander*, *Sage*, and others. All which, though the naked *Eye* will discover, yet by the help of *Glasses* we may observe them most distinctly. The use of these we suppose the same, in part, with those of the *Flower*, whereof we shall speak.

CHAP. V.

Of the FLOWER.



E next proceed to the *Flower*. The general *Parts* whereof are most commonly three; *sc.* the *Empalement*, the *Foliation*, and the *Attire*.

2. §. The *Empalement*, whether of one or more pieces, I call that which is the utmost *Part* of the *Flower*, encompassing the other two. 'Tis compounded of the three general *Parts*, the *Skin*, the *Cortical* and *Lignous Bodies*, each *Empaler* (where there are divers) being as another little *Leaf*; as in those of a *Quince-Flower*, as oft as they happen to be overgrown, is well seen. As likewise in the *Primrose*, with the *green Flower*; commonly so call'd, though by a mistake: For that which seems to be the *Flower*, is only the more flourishing *Empalement*, the *Flower* it self being *White*. But the continuation of all the three aforesaid *Parts* into each *Empaler*, is discoverable, I think, no where better than in an *Artichoke*, which is a true *Flower*, and whose *Empalers* are of that amplitude, as fairly to shew them all: As also, that the Original of the *skin* of each *Empaler* or *Leaf* is not distinct from that of the rest; but to be all one piece, laid in so many *Plaits* or *Duplicatures*, as there are *Leaves*, from the outermost to the inner and most Central ones.

3. §. The Design of the *Empalement*, is to be *Security* and *Bands* to the other two *Parts* of the *Flower*: To be their *Security* before its opening, by intercepting all extremities of *Weather*: Afterwards to be their *Bands*, and firmly to contain all their *Parts* in their due and most decorous posture: so that a *Flower* without its *Empalement*, would hang as uncouth and taudry, as a *Lady* without her *Bodies*.

4. §. Hence we have the reason why it is various, and sometimes wanting. Some *Flowers* have none, as *Tulips*; for having a fat and firm *Leaf*, and each *Leaf* likewise standing on a broad and strong *Basis*, they are thus sufficient to themselves. *Carnations*, on the contrary, have not only an *Empalement*, but that (for more firmitude) of one piece: For otherwise, the Foot of each *Leaf* being very long and slender, most of them would be apt to break out of compass: yet is the top of the *Empalement* indented also; that the *Indentments*, by being lapp'd over the *Leaves* before their expansion, may then protect them; and by being spread under them afterwards, may better shoulder and prop them up. And if the Feet of the *Leaves* be both long and very tender too, here the *Empalement* is numerous, though consisting of several pieces; yet those in divers *Rounds*, and all with a counterchangeable respect to each other (which also the Learned Sir *Thomas Brown* observes) as in all *Knapweeds*, and other *Flowers*; whereby, how commodious they are for both the aforesaid ends, may easily be conceiv'd; and well enough exemplified by the Scales of *Fishes*, whereunto, as to their position, they have not an unapt resemblance.

Treat. of the
Quincunx.

5. §. THE FOLIATION also, is of the same substantial Nature with the green Leaf; the Membrane, Pulp, and Fibres whereof, being, as there, so here, but the continuation of the Skin, the Cortical and Lignous Bodies.

6. §. The Foulds of the Flower or Foliation are various, as those of the green Leaf; but some of them different. The most general are, First, The Close-Couch, as in Roses, and many other double Flowers. Then the Concave-Couch, as in *Blattaria flore albo*. Next the Plait, as in some of the Leaves of Pease-Blooms, in the Flowers of Coriander, &c. which is either single, as in those nam'd; or double, as in *Blew-Bottle*, *Jacea*, and more of that rank. Next, the Couch, and Plait together in the same Flower, as in *Marigolds*, *Daisies*, and all others of an agreeing form: where the first apparent Fould or Composture of the Leaves is in Couch; but the Leaves being erect, each likewise may be seen to lie in a double Plait within it self. Then the Rowl, as in the Flowers of *Ladies-Bower*, the broad top of each Leaf being by a double Rowl foulded up inwardly. Next, the Spire, which is the beginning of a Rowl; and may be seen in the Flowers of *Mallows*, and others. Lastly, the Plait and Spire together, where the Part analogous to the Foliation, is of one piece, the Plaits being here laid, and so carried on by spiral Lines to the top of the Flower, as is in divers, and I think, in *Convolvulus Doronici folio*, more elegantly seen. These and other Foulds, See in the Figures belonging to the Second Part of the Fourth Book. The reason of all which varieties, a comparative consideration of the several Parts of the Flower may suggest. I'll only mention, That no Flower, that I find, hath a Back-Rowl, as hath the green Leaf. For two Reasons; because its Leaves have not their Fibres standing out much on their backside, as the green Leaves have; and because of its Attire, which it ever embosomes, and cannot so well do it by a Back-Rowl.

7. §. The usual Protections of Flowers by the Precedents are express'd, sc. Green Leaves and Empalements. Some have another more peculiar, that is a double Veil; as the *Spring-Crocus*. For having no Empalement, and starting up early out of the Mould, even before its Green Leaves, and that upon the first opening of the Spring; lest it should thus be quite starved, 'tis born swath'd up in a double Blanket, or with a pair of Sheets upon its Back.

8. §. The Leaves of divers Flowers at their Basiss have an hairy Tuft; by which Tufts the Concave of the Empalement is filled up; That, being very choice and tender, they may thus be kept in a gentle and constant Warmth, as most convenient for them.

9. §. The Leaves of the Flower, though they are not hairy all over, yet in some particular parts they are often set with a fine Downy Velvet; that, being by their shape and posture in those parts contiguous to their delicate and tender Attire, they may thus give it a more soft and warmer touch. Thus in the Flower of *Ladies Bower*, those parts of its Leaves which rowl inward, and lie contiguous to the Attire, are Downy; whereas the other Parts are smooth or bald: So the Flowers of *Pease*, *spanish Broom*, *Toad-Flax*, and many others, where contiguous to their Attires, are deck'd with the like Hairy Velvet.

10. §. As upon the Green *Leaves*, so upon the *Flowers* are *Globulets* sometimes seen; as upon the backside of that of *Enula*. On none more plainly than that kind of *Blattaria* with the white *Flower*; where they are all transparent, and growing both on the *Stalk* and *Leaves* of the *Flower*, each shewing likewise its *Peduncle* whereon it is erected.

11. §. The use of the *Flower*, or the *Foliation* whereof we now speak, (that is, as to its private service) is for the protection of the *Attire*; *This*, as its under, and the *Empalement* as its upper Garments. As likewise of the *Fruit*: The necessity of which Service, in some Cases, by the different situation of the *Flower* and *Fruit*, with respect to each other, is evident; *Apples*, *Pears*, and several other *Fruits*, standing behind or under the *Flower*; but *Cherries*, *Aprecots*, and divers others, within it. For these being of a very tender and pulpy Body, and withal putting forth with the colder part of the Spring; could not weather it out against the Variations and Extremities of the Air, (as those of a more solid *Parenchyma* can) except lodged up within their *Flowers*.

12. §. And as the *Flower* is serviceable to the safety of the *Fruit*, so is it to its growth; *sc.* in its Infancy, or *Embryo*-estate; for which purpose, as there is a *Flower*, so that *Flower* is greater or less, according as the nature of the *Fruit* to which it belongs, and the plenty of the *Sap* by which the *Fruit* is fed, doth require. Thus, where the young *Fruit* is of a solid Substance and the ascent of the *Sap* less copious, were there here no *Flower* to promote the said ascent thereof into the *Fruit* (in the manner as is effected by the Green *Leaves*) it must needs pine and die, or prove less kindly. On the contrary, should the *Flower* be over-large, it would not only promote the ascent of the *Sap* up to the *Fruit*, but being as yet over-proportionate to it, would likewise it self exhaust the same *Sap*, as fast as ascendent; like a greedy Nurse, that prepares the Meat for her Child, and then eats it up her self. Thus we see *Apples* and *Pears*, with a *Flower* of a moderate Size; like their *Body*, of a middle Constitution, and their *Sap*, of a middle quantity: But *Quinces*, being more solid, besides that they have as great a *Flower*, the *Impalers* of their *Flower* also thrive so far as to become handsome *Leaves*; continuing also after the *Flower* is fallen, firm and verdent a great while; so long, till the *Fruit* be able to provide for it self. On the other hand, *Plums* being more tender and Sappy than *Apples* and *Pears*, besides that their *Empalers* are much alike, their *Flower* is less. and *Gooseberries* and *Currans*, which are still more Pulpy, and the course of the *Sap* towards them more free, have yet a *Flower* far less. And *Grapes*, whose *Sap* is still of quicker Ascent, have scarce any *Flower* at all; only some small resemblance thereof, serving just upon the setting of the *Fruit*, and no longer.

13. §. THE ATTIRE, I find to be of two kinds, *Seminiforme*, and *Florid*. That which I call *Seminiforme*, is made up of two general Parts, *Chives* and *Semets*, one upon each *Chive*. These *Semets* (as I take leave to call them) have the appearance, especially in many *Flowers*, of so many little *Seeds*: but are quite another kind of *Body*. For, upon enquiry, we find, that these *Semets*, though they seem to
be

be solid, and for some time after their first formation, are entire; yet are they really hollow; and their side, or sides, which were at first entire, at length crack asunder: And that moreover the *Concave* of each *Semet* is not a meer vacuity, but fill'd up with a number of minute Particles, in form of a *Powder*. Which, though common to all *Semets*, yet in some, and particularly those of a *Tulip* or a *Lilly*, being larger, is more distinctly observable.

Tab. 4.f.12.

14. §. These *Semets* are sometimes fastned so, as to stand erect above their *Chive*, as those of *Larks-beel*. Sometimes, and I think usually, so as to hang a little down by the middle, in the manner and figure of a *Kidney*; as in *Mallows*. Their *Cleft* or *Crack* is sometimes single, but for the most part double: At these *Clefts* it is that they disburse their *Powders*; which as they start out, and stand betwixt the two Lips of each *Cleft*, have some resemblance to the common Sculpture of a *Pomegranate* with its *Seeds* looking out at the *Cleft* of its *Rind*. This must be observ'd when the *Clefts* are recently made, which usually is before the expansion of the *Flower*.

f. 12. -a.

15. §. The Particles of these *Powders*, though like those of Meal or other Dust, they appear not easily to have any regular shape; yet upon strict observation, especially with the assistance of an indifferent *Glass*, it doth appear, That they are a *Congeries*, usually, of so many perfect *Globes* or *Globulets*; Sometimes of other *Figures*, but always regular. That which obscures their *Figure* is their being so small: In *Dogs-Mercury*, *Borage*, and very many more *Plants*, they are extreamly so. In *Mallows*, and some others, more fairly visible.

16. §. Some of these *Powders*, are *yellow*, as in *Dogs-Mercury*, *Goats-Rue*, &c. and some of other Colours: But most of them I think are *white*; and those of *yellow Henbane* very elegant; the disburse'd *Powers* whereof, to the naked eye, are *white* as *Snow*; but each *Globulet*, through a *Glass*, transparent as *Crystal*; which is not a fallacy from the *Glass*, but what we see in all transparent *Bodies* whatsoever, lying in a *Powder* or small Particles together. The Parts of this *Attire*, see in *Tab. 4.* But especially, in the *Figures* belonging to the *Second Part* of the *Fourth Book*.

17. §. The *Florid Attire*, is commonly known by the blind and rude Name of *Thrums*; as in the *Flowers* of *Marigold*, *Tansie*, &c. How in adequate its imposition is, observation will determine. For the several *Thrums* or rather *Suits*, whereof the *Attire* is made up, however else they may differ in various *Flowers*, in this agree, that they are ever consistant of more than one, sometimes of Two, and for the most part of Three *Pieces* (for which I call them *Suits*) and each *Piece* of a different, but agreeable and comely form.

Tab.4.f.13.a.

18. §. The *outer Part* of every *Suit*, is its *Floret*: whose *Body* or *Tube* is divided at the top (like that of the *Cowslip*) into five distinct *Leaves*. So that a *Floret*, is the *Epitome* of a *Flower*: and is all the *Flower* that many *Plants*, as *Mugwort*, *Tansie*, and others, have. What the Learned Sir *Thomas Brown* observeth of the number *Five*, as to the *Leaves* of the *Flower*, is still more universally holding in these of the *Floret*.

f. 13. b.

Treat. of the
Quincunx.

19. §. Upon the Expansion of the *Floret*; the next *Part* of the *Suit* is from within its *Tube* brought to light; which we may (with respect to that within it) call the *Sheath*. For this also, like the *Floret*,

f. 13. c.

is

is a *Concave Body*; in its shape very well resembling the *Fistulous Pouches* of *Wake-Robin*, or of *Dragon*.

20. §. The *Sheath*, after some time, dividing at the top, from within its *Concave* the *Third* and innermost part of the *Suit*, *sc.* the *Blade* advanceth and displayes it self. This Part is not hollow, as the other two, but solid; yet at its *Point*, is commonly, divided into two halves. Tab. 4. f. 13. d

21. §. About the said *Point* especially, there appears, *Globulets*, which are of the same nature with those of a *Semet*, though not so copious. So that all *Flowers* have their *Powders* or *Globulets*. The whole *Attire* may in *Aster Per*, *Blew-bottle*, &c. where the *Suits* are large, be plainly observed without a *Glass*. The *Parts* of this *Attire*, See in *Tab. 4.* But especially in the *Figures* belonging to the *second Part* of the *Fourth Book*.

22. §. The use of the *Attire*, how contemptibly soever we may look upon it, is certainly great. And though for our own use we value the *Leaves* of the *Flower*, or the *Foliation*, most; yet of all the three *Parts*, this in some respects is the choicest, as for whose sake and service the other two are made. The use hereof, as to *Ornament* and *Distinction*, is unquestionable; but is not all. As for *Distinction*, though, by the help of *Glasses*, we may make it to extend far; yet in a *passant view*, which is all we usually make, we cannot so well. As for *Ornament*, and particularly in reference to the *Semets*, we may ask, If for that meerly these were meant, then why should they be so made as to break open, or to contain any thing within them? Since their *Beauty* would be as good if they were not hollow; and is better before they crack and burst open, than afterwards.

23. §. Other uses hereof therefore we must acknowledge, and may observe. One is, for food; for *Ornament* and *Distinction* to us, and for *Food* to other *Animals*. I will not say, but that it may serve even to these for *Distinction* too, that they may be able to know one *Plant* from another, and in their flight or progress settle where they like best: and that therefore the varieties of these small parts are many, and well observed by them, which we take no notice of. Yet the finding out of *Food* is but in order to enjoy it: Which, that it is provided for a vast number of little *Animals* in the *Attires* of all *Flowers*, observation perswades us to believe. For why else are they evermore here found? Go from one *Flower* to another, great and small, you shall meet with none untaken up with these *Guests*. In some, and particularly the *Sun-Flower*, where the parts of the *Attire*, and the *Animals* for which they provide, are larger, the matter is more visible. We must not think, that *God Almighty* hath left any of the whole *Family* of his *Creatures* unprovided for; but as the *Great Master*, some where or other carveth out to all; and that for a great number of these little *Folk*, He hath stored up their peculiar provisions in the *Attires* of *Flowers*; each *Flower* thus becoming their *Lodging* and their *Dining-Room*, both in one.

24. §. Wherein the particular parts of the *Attire* may be more distinctly serviceable, this to one *Animal*, and that to another, I cannot say: Or to the same *Animal*, as a *Bee*, whether this for the *Honey*, another for their *Bread*, a third for the *Wax*: Or whether all only suck from

from hence some *Juice*; or some may not also carry some of the *Parts*, as of the *Globulets*, wholly away.

25. §. Or lastly, what may be the Primary and Private Use of the *Attire* (for even this above said, though great, yet is but Secondary) I now determine not.

CHAP. VI.

Of the FRUIT.



THE general composition of all *Fruits* is one, that is, their *Essential* and truly *Vital Parts*, are in all the same, and but the continuation of those which in the other *Parts* of a *Plant*, we have already observed. Yet because by the different *Constitutions* and *Tinctures* of these *Parts*, divers considerably different *Fruits* result; I shall therefore take a particular view of the more known and principal of them, *sc.* *Apples, Pears, Plums, Nuts* and *Berries*.

2. §. AN APPLE, if cut traverse, appears constituted of four distinct *Parts*, the *Pilling*, the *Parenchyma*, *Branchery*, and *Coare*. The *Pilling* is only the spreading and dilatation of the *Skin*, or utmost part of the *Barque* in the *Branch*. The *Parenchyma*, when full ripe, is a tender delicate *Meat*. Yet as the *Pilling* is but the Continuation of the utmost part of the *Barque*; so is this, but the continuance and ampliation, or (as I may call it) the swelth and superbience of the *Inner Part* thereof; which upon observation of a young and *Infant-Apple* especially, is evident. Thus we see the *Pith*, which is often tough; in many *Roots*, as *Parsneps, Turneps, &c.* is tender and edible. So here, the *Parenchyma*, though originally no more than the *Barque*, yet the copiousness and purity of its *Sap* being likewise effectual to the largeness and fineness of its growth, it thus becomes a soft and tender meat. The *Branchery* is nothing else but the *Ramifications* of the *Lignous Body* throughout all the parts of the *Parenchyma*; the greater *Branches* being likewise by the *Inosculation*s of the less (as in the *Leaf*) united together. The main *Branches* are usually Twenty: Ten are spread and distributed through the *Parenchyma*, most of them enarching themselves towards the *Cork* or *Stool* of the *Flower*: The other Ten, running from the *stalk* in a directer Line, at last meet the former at the said *Cork*, and are there osculated with them. Of these latter, five are originated from one; which running along the Center of the *stalk*, and part of the *Parenchyma* of the *Fruit*, is therein at last divided. To these the *Coats* of the *Kernels* are fastned. So that whereas most of these *Branches* were originally extended even beyond the *Fruit*, and inserted into the *Flower* for the due growth there-

thereof; the *Fruit* afterwards growing to some head, and so intercepting and preying upon the *Aliment* of the *Flower*, starves that and therefrom supercedes the service of the said *Branches* to it self, fifteen for its *Parenchyma*, and five for its *Seed*. The *Coar* is originated from the *Pith*; for the *Sap* finding room enough in the *Parenchyma*, through which to dispence it self all abroad, quits the *Pith*, which thereby hardens into a *Coar*. Thus we see the *Insertions*, although originate from the *Cortical Body*, yet their *Parts* being, by the *Inosculation*s of the *Lignous*, so much compress'd and made to co-incide together, they become a *Body* very compact and dense. And in the *Barque* the same thing is effected by *Arefaction* only, or a meer *voydance* of the *Sap*; the *Inner Part* whereof, though soft and sappy, yet its superficial *Rind* is often so hard and smooth, that it may be fairly writ upon. The *Parts* of an *Apple*, See in the *Figures* belonging to the *Third Part* of the *Fourth Book*.

3. §. IN A PEAR there are five distinct *Parts*, the *Pilling*, the *Parenchyma*, *Branchery*, *Calculary*, and *Acetary*. The three former are here and in an *Apple* much alike; saving that here the *Inner* or *Seed-Branches* ordinarily stand double. The *Calculary* (most observable in rough-tasted, or *Choak-Pears*) is a *Congeries* of little stony *Knots*. They are many of them dispersed throughout the whole *Parenchyma*: But lying more continuous and compact together towards the Center of the *Pear*, surround the *Acetary* there, in a somewhat Globular Form: About the *Stalk* they stand more distant; but towards the *Cork* or *Stool* of the *Flower*, they still grow closer, and there at last gather (almost) into the firmitude of a *Plum-stone* it self. Within this lies the *Acetary*; 'tis allways sour, and by the bounding of the *Calculary* of a Globular Figure. 'Tis a simple *Body*, having neither any of the *Lignous* branched in it, nor any *Calculous Knots*. It is of the same substantial nature with the outer *Parenchyma*; but whether it be absolutely one with it, or be derived immediately from the *Pith*, my Enquiries yet made, determine not. Tab. 4. f. 14.

4. §. The Original of the *Calculary* I seem to have neglected. But hereof we may here best say, that whereas all the other *Parts* are *Essential* and truly *Vital*; the *Calculary* is not: but that the several *Knots* whereof it consists, are only so many meer *Concretions* or *Precipitations* out of the *Sap*; as in *Urines*, *Wines*, and other *Liquors*, we often see. And that the *Precipitation* is made by the mixture and re-action of the *Tinctures* of the *Lignous* and *Cortical Bodies* upon each other: Even as all *Vegetable Nutrition* or *Fixation* of *Parts* is also made by the joynt efficiency of the two same *Tinctures*, as hath been said. Hence we find, that as the *Acetary* hath no *Branches* of the *Lignous Body*, so neither hath it any *Knots*. Hence likewise it is, that we have so different and contrary a tast in the *Parenchyma* beyond the *Calculary*, from that in the *Acetary*: For whereas this is sour, that, wherein the said *Precipitations* are made, is sweet; being much alike effect to what we find in mixing of *Corals*, &c. with *Vinegar* or other *acid Liquors*. The *Parts* of a *Pear*, See in *Tab. 4.* But especially in the *Figures* belonging to the *Third Part* of the *Fourth Book*.

5. §. IN A PLUM (to which the *Cherry*, *Apricot*, *Peach*, *Walnut*, &c. ought to be refer'd) there are four distinct *Parts*, the *Pilling*, the *Parenchyma*, *Branchery* and *Stone*. The *Pilling* and *Parenchyma* are, as to their *Original*, with those of an *Apple* or *Pear*, both alike. As likewise the *Branchery*; but differently ramified. In *Plums* (I suppose all) there are five main *Out-Branche*s, which run along the *Surface* of the *Stone* from the *Basis* to the point thereof, four of them by one *Ridge*, and one by the other opposite to it. In an *Apricot* there is the same number, but the single *Branch* runs not upon the *Surface*, but through the *Body* of the *Stone*. There are likewise two or three smaller *Branche*s, which run in like manner under the other *Ridge* for some space, and then advancing into the *Parenchyma*, therein disperse themselves: These latter sort in *Peaches* are numerous throughout.

Tab. 4. f. 15.

6. §. But notwithstanding the different disposition of the *Branche*s of the *Fruits* aforesaid; yet is there one *Branch* dispos'd in one and the same manner in them all. The entrance hereof into the *Stone* is at its *Basis*; from whence running through its *Body*, and still inclining or arching it self towards its *Concave*, is at last, about its *Cone*, there-into emergent, where the *Coats* of the *Seed* are appendent to it. Of the *Seed-Branch* 'tis therefore observable that after its entrance into the *Fruit*, 'tis always prolonged therein to a considerable length; as is seen not only in *Apples*, &c. where the *Seed* stands a good distance from the *Stalk*; but in *Plums* likewise, where it stands very near it; in that here the *Seed-Branch*, as is said, never strikes through the *Stone* into the *Coats* of the *Seed* directly, but runs through a *Chanel* cut in the *Stone*, till it issues, near the *Cone*, into the *Concave* thereof.

Tab. 4. f. 15.

7. §. The *Stone* though it seem a simple *Body*, yet it is compounded of different ones. The *Inner Part* thereof, as it is by far the thinnest, so is it the most *dense*, *white*, *smooth* and *simple*. The *Original* is from the *Pith*; difficult, but curious to observe: For the *Seed-Branch*, not striking directly and immediately quite through the *Basis* of the *Stone*, but in the manner as is above described, carries a considerable *Part* of the *Pith*, now gather'd round about it, as its *Parenchyma*, along with it self; which upon its entrance into the *concave* of the *Stone* about its farther end, is there in part spread all over it, as the *Lining* thereof. The outer and very much thicker *Part*, consisteth partly of the like *Precipitations* or concrete *Particles*, as in a *Pear*; being gathered here much more closely, not only to a *Contiguity*, but a *Coalition* into one entire *Stone*; as we see in *Pears* themselves, especially towards the *Cork*, they gather into the like *Stoniness*; or as a *Stone*, *Mineral*, or *Animal*, is oftentimes the product of accumulated *Gravel*. But as the *Parenchyma* is mixed with the *Concretions* in the *Calculary*, so is it also, though not visibly, with these in the *Stone*, the *ground* of the *Stone* being indeed a perfect *Parenchyma*; but by the said *Concretions* so far alter'd, as to become dry, hard and undistinguishable from them. All which *Particulars*, are observable only in the several degrees of *Growth* in the young *Fruit*. And are represented in *Tab. 4.* But especially by the several *Figures* belonging to the *Third* and *Fourth Parts* of the *Fourth Book*.

8. §. IN A NUT (to which an *Akern* is analogous) there are three general *Parts*, the *Cap*, *Shell*, and *Pith*. The *Cap* is constituted of a *Pilling* and *Parenchyma*, derived from the *Barque*; and *Ramulets* from the *Lignous Body* of the *Branch*. The *Shell* likewise is not one simple *Body*, but compounded. The *Superficial Part* thereof is originated from the *Pilling* or *Skin* of the *Cap*, from the inside whereof it is, in a *Duplicature*, produc'd and spread over the *Shell*. Which, if you look at the *Basis* of the *Shell*, is farther evident: for that being continuous with the *Parenchyma* of the *Cap*, without the interposure of the *Skin*, the said superficial *Part* is there wanting. The thicker and inner *Part* of the *Shell* consisteth of the same *Parenchyma* as that of the *Cap*, with a *Congeries* of *Precipitations* filled up, as in a *Stone*. And as the *Lignous Body* is branched in a *Stone*, so, with some difference, in a *shell*. The *outer Branches* or *Ramulets* are numerous, each issuing out of the *Parenchyma* of the *Cap*, and entering the *shell* at the *Circumference* of its *Basis*, and so running betwixt its superficial and inner *Parts* towards the *Cone*, round about. The *Inner* or *Seed-Branch* is single, entering in, as do the other, at the *Basis* of the *shell*, but at the *Center* thereof: from whence it runs, not through the *Shell*, as in *Plums* through the *Stone*; but through the *Pith*, as far as the *Cone*; where the *Coats* of the *Seed* hang appendent to it. The *Pith* whether derived from the same part both in name and nature in the *Branch* and *Stalk*; or from the *Cortical Body*, I yet determine not. The *Parts* of a *Nut*, See in the *Figures* belonging to the *Third Part* of the *Fourth Book*.

9. §. A BERRY, as a *Gooseberry* (to which *Corinths*, *Grapes*, *Hips*, &c. are to be refer'd) consisteth, besides the *Seed*, of the three general *Parts*, *Pilling*, *Parenchyma* and *Branchery*. The *Pilling* is originated as in the foregoing *Fruits*. The *Parenchyma* is double, as likewise in some other *Berries*. The *outer* is commonly, together with the *Pilling*, call'd the *skin*, and is that part we spit out, being of a sour taste. Now as the *Pilling* is originated from the *outer*, so this from the *inner Part* of the *Barque*; and accordingly the *Pores* thereof may be observed plainly of a like shape with those both of the *Cortical Body* and *Pith*. The *Inner* or *Pulp* is of a sweet taste, and is the *Part* we eat: It is of a Substance so laxe and tender, as it would seem to be only a thicker or jellied *Juice*; although this likewise be a true *Parenchyma*, something like that of an *Orange* or *Limon*, with its *Pores* all fill'd up with *Liquor*. The *Branchery* is likewise double: The *Exterior* runs betwixt the *Pilling* and *Outer Parenchyma* in arched *Lines*, from the *Stalk* to the *Stool* of the *Flower*. These *outer Branches*, though of various number at the *Stalk*, yet at the *Cork* are usually ten principal ones; five for the five *Leaves* of the *Flower*, and five for the *Attire*. The *Inner main Branches* are two, diametrically opposite to each other, and at the *Cork* with the other inosculated. From these two are branched other smaller, every one having a *Seed* appendent to it, whose *Coats* it entreteth by a double *Filament*, one at the *Basis*, the other at the *Cone*. They are all very white and turgent, and by a slant cut, may be observ'd concave; thus representing themselves analogous to so many true *spermatick Vessels*. The

Parts of a Gooseberry, See in the *Figures* belonging to the *Third Part* of the *Fourth Book*.

10. §. The Uses of *Fruits* are for *Man*, (sometimes also other *Animals*, as are *Akerns* and *Haws*) and for the *Seed*. For *Man*, they are so variously desirable, that till our *Orchards* and *Store-Chambers*, *Confectioners-Stoves* and *Apothecaries-Shops*, our *Ladies Closets*, their *Tables* or *Hands* are empty of them, I shall not need to enquire for what. If it be asked, how the *Fruit* becomes, generally above all the other *Parts*, so pleasant a Meat? It is partly from the *Sap*, the grosser portion thereof being deposited in the *Leaves*, and so the purer hereunto reserved. Partly from the *Globular Figure* of the *Fruit*. For the *Sap* being thus in a greater quantity herein, and in all *Parts* equally diffus'd, the *Concoction* hereof, as in a *Vessel*, is with greatest advantage favoured and promoted. Wherefore all *Fruits*, which we eat raw, how small soever, are of a *Globular Form*, or thereunto approaching; and the nearer, the delicater; amongst *Apples*, the *Pipin*; amongst *Pears*, the *Burgundian*; and amongst all *Fruits*, the *Grape*; and amongst *Grapes*, the roundest, are of all, the most dainty.

11. §. The visible cause of this *Globular Figure*, is the *Flower*; or the Inosculation of all the main *Branches* at the *Stool* of the *Flower*; and upon the fall of the *Flower*, the obtuseness, and with *Wind* and *Sun*, as it were the *sealing* of their several ends: For thus the *Sap* entering the *Fruit*, being not able to effect, either a *Disunion*, or a *shooting* forth of the said *Branches*, and so to carry on their *Growth* in length; they must of necessity be enarch'd, and with the *Parenchyma* more and more expand themselves. Whereas were they disposed and qualified otherwise, than as is said; instead of forming a *Fruit* within bounds, they would run out into all extravagance, and even into another little *Tree* or *Leafy Growth*.

12. §. To the *Seed*, the *Fruit* is serviceable; First, in order to its being supply'd with a due and most convenient *Sap*, the greater part thereof, and that which is less elaborated, being, in its passage towards the *Seed*, thereinto received; the *Fruit* doing the same office to the *Seed*, which the *Leaves* do to the *Fruit*; the *Sap* in the *Fruit* being, in a laxer comparison, as the *Wine*; and that for the *Seed*, a small part of the highest Spirit rectified from it.

13. §. So likewise for its Protection, in order to the prosperous carrying on and perfecting of its generation, and security being perfected. Which protection it gives not only to the Seminal *Sap* and *Seed* it self, but ever also to its *Seed-Branch*. Thus we see an *Apple*, besides that it is it self of ample compass, for the sake of its *seed*, hath likewise its *Cour*; as if it were not sufficient, that the Walls of their Room are so very thick, unless also *wainscoted*. In a *Pear* again, where the *Parenchyma* is of less compass than that of an *Apple*, to what protection this affords, that of the *Calculary* is super-added. But in a *Plum*, where the *Parenchyma* is exceeding tender, and in a *Peach*, which hangs late, and till Autumn Frosts approach, we have not only the Rubbish of a *Calculary*, but stout Stone-Walls. Within which also, not only the *Seed* it self, but the *Seed-Branch* is evermore immur'd. Lastly, in a *Nut*, where the *Shell* being not surrounded with a *Parenchyma*, that protection is wanting without, 'tis answer'd by an ample
Pith

Pith within it; and the *Seed-Branch* likewise included, not meerly in the *Body* of the *Shell*, as in a *Plum*, but within the *Pith* it self. So necessary is this design, that what the *Hen* by *Incubation* or *Hovering*, is to the *Egg* or *Chick*; that the whole *Fruit*, by comprehension, is to the *Seed*.

CHAP. VII.

Of the SEED, in its State of Generation.



THE Original, so the Ultimate end and Perfection of *Vegetation* is the *Seed*. How it is the former, and in its state apt for *Vegetation*, hath already been seen. How the latter, and in its state of *Generation*, we shall now lastly enquire. In doing which, what in the other state, was either not distinctly existent, or not so apparent, or not so intelligible, will occur.

2. §. The two general *Parts* of the *Seed* are its *Covers* and *Body*. The *Covers* in this estate are usually *Four*. The outmost, we may call the *Case*. 'Tis of a very various form; sometimes a *Pouch*, as in *Nasturtium*, *Cochlearia*; a *Cod*, as in all *Pulse*, *Galega*; sometimes not entire, but parted, or otherwise open, as in *Sorrel*, *Knotgrass*; with many other forms: I think alwaies more heterogeneous to that of the *seed*, by which it differs from the proper *Coats*. To this the *Caps* of *Nuts*, and the *Parenchyma's* of other *Fruits* are analogous.

3. §. The two next are properly the *Coats*. In a *Bean* especially, and the like; from whence, to avoyd *Confusion*, the denomination may run common to the responding *Covers* of other *Seeds*. The *Colour* of the outer, is of all degrees, from *White* to the *Blackness* of *Jett*. It's *Figure* sometimes *Kidney'd*, as in *Alcea*, *Behen*, *Poppy*; *Triangular*, as in *Polygonatum*, *Sorrel*; *Spherically triangular*, in *Mentha*, *Melissa*; *Circular*, in *Leucoium*, *Amaranthus*; *Globular*, in *Napus*, *Asperula*; *Oval*, in *Speculum Veneris*, *Tithymalus*; *half Globe*, in *Coriander*; that which we take for *one* single round *Seed*, being a *Conjugation* of *two*; *half Oval*, in *Anise*, *Fennel*; *Haftal*, in *Lactuca*; *Cylindrical*, as, if I mistake not, in *Jacobæa*; *Pyramidal*, in *Geranium Althææ fol.* with many other differences. But the *Perfection* of one or two of the said *Figures* lieth in the *Case*. So that, as all *Lines* and *Proportions* are in the *Leaf* and *Flower*; so all *Regular solids* in the *Seed*; or rather in its *Covers*.

4. §. 'Tis sometimes *glistening*, as in *Speculum Veneris*; *Rough-cast*, in *Catanance*; *Studded*, in *Behen*, *Baltaria*; *Favous*, in *Papaver*, *Antirrhinum*, *Lepidum annuum*, *Alcea Vesicaria*, *Hyoscyamus*, and many more, before the *Seeds* have lain long by; *Pounced*, in *Phalangium Cretæ*, *Lithospermum*; *Ramified*, in *Pentaphyllum fragiferum Erectum majus*, resembling

resembling the *Fibers* of the *Ears* of the *Heart*; some just *Quinquenerval*, as in *Anisum*, and many more, the *Lignous Body* being in five main *Fibers* branched therein. The *Figures*, and *Surface*, of These, and other *Seeds*, See in the *Tables* belonging to the *Fourth Part* of the *Fourth Book*.

5. §. The *Covers* of not only *Quince-Seeds*, and those of *Psyllium* (more usually taken notice of) but those also of *Horminum*, *Nasturtium*, *Eruca*, *Camelina*, *Ocimum*, and divers others, have a *Mucilage*. Which, though it be not visible when the *Seeds* are throughly dry; yet lying a while in some warm *Liquor*, or only on the *Tongue*, it swells more or less, and upon them all fairly shews it self. On that of *Ocimum* it appears grayish; on the other, transparent; and on that of *Nasturtium Hortense* very large; even emulous of the inner *Pulp* surrounding a *Gooseberry-Seed*. The putting of *Clary-seed* into the *Eye*, may have been brought into use from this *Mucilage*, by which alone it may become Medicinal. And thus far of the *Superficies*,

6. §. The *nature* of the outer *Coat* is also various, *Membranous*, *Cartilaginous* and *Stony*; the like *Precipitations* being sometimes made herein, as in a *Stone* or *Shell*; as in that of the *Seeds* of *Carthamum*, *Lithospermum* and others. The *Designment* hereof, being either with respect to the *Seed* in its state of *Generation*; as where the *Case* is either wanting, or at least insufficient of it self, there for its due protection and warmth. Or, in its state of *Vegetation*, for the better *Fermenting* of its *Tinctures* and *Sap*; the *Fermentations* of some *Seeds* not well proceeding, unless they lie in their *Stony Casks* in the *Ground*, like *Bottled Liquors* in *Sand*.

7. §. All *seeds* have their outer *Covers* open; either by a particular *Foramen*, as in *Beans*, and other *Pulse*, as is said; or by the breaking off of the *Seed* from its *Peduncle* or *Stool*, as in those in *Cucumber*, *Cichory*; or by the entering and passage of a *Branch* or *Branches*, not only into the *Concave* thereof near the *Cone*, but also through the *Cone* it self; as in *Shells* and *stones*.

8. §. For the sake of this *aperture* it is, that *Akerns*, *Nuts*, *Beans*, *Cucumbers*, and most other *seeds*, are in their formation so placed, that the *Radicle* still standeth next to it; That So, upon *Vegetation*, it may have a free and ready passage into the *Mould*.

9. §. The *Original* of the outer *Coat*, though from *Parts* of the same substantial nature, yet is differently made. In a *Plum*, the *Seed-Branch* which runs, as is described, through the *Stone*, is not naked, but, as is said, invested with a thin *Parenchyma*, which it carries from the *Stalk* along with it; and which, by the *Ramification* of the said *Branch* within the *Stone*, is, in part, dilated into a *Coat*. That of a *Bean* is from the *Parenchyma* of the *Cod*; the superficial part of which *Parenchyma*, upon the large *peduncle* of the *Bean* becoming a thin *Cuticle*, and upon the *Bean* it self a *Cartilaginous Coat*.

10. §. The *Original* of the inner *Coat* of the *Bean* is likewise from the inner part of the said *Parenchyma*; which first is spread into a long *Cake*, or that which with the *Seed-Branch* maketh the *Peduncle* of the *Bean*; under which *Cake*, there is usually a black part or spot; by the length of which, the inner part of the *Cake* is next inserted into the outer *Coat*, and spread all over the *Concave* thereof, and so becomes the inner.

11 §. Of this Inner Coat it is very observable, That although when the *Seed* is grown old and dry, 'tis shrunk up, and in most *Seeds*, so far, as scarcely to be discern'd; yet in its first and juvenile Constitution, it is a very Spongy and Sappy body; and is then likewise (as the *Womb* in a Pregnant *Animal*) in proportion, very thick and bulky. In a *Bean*, even as one of the *Lobes* it self: And in a *Plum* or *Apricot*, I think I may safely say, half an hundred times thicker than afterwards, when it is dried and shrunk up, and can scarcely be distinguished from the upper Coat. Upon which Accounts it is, in this estate a true and fair *Parenchyma*. The Delineation hereof, See in the *Figures* belonging to the *Fourth Part* of the *Fourth Book*.

12. §. In this Inner Coat in a *Bean*, the *Lignous Body* or *Seed-Branch* is distributed: Sometimes, as in *French-Beans*, throughout the whole Coat, as it is in a *Leaf*. In the Great *Garden-Bean*, upon its first entrance, it is bipartite, and so in small *Branches* runs along the *Circumference* of the Coat, all meeting and making a kind of *Reticulation* against the Belly of the *Bean*. In the same manner the main *Branches* in the outer Coat of a *Kernel*, circling themselves on both hands from the place of their first entrance, at last meet, and mutually inosculate; as the *Veins* in the *Kidneys* of a Man or any *Quadrupede*; Or the *Carotick Arteries* in the *Braine*.

13. §. So that all the *Parts* of a *Vegetable*, the *Root*, *Trunk*, *Branch*, *Leaf*, *Flower*, *Fruit* and *Seed*, are still made up of *Two* Substantially different Bodies.

14. §. And as every *Part* hath *Two*, so the whole *Vegetable* taken together, is a composition of *Two* only, and no more: All properly *Woody Parts*, *Strings* and *Fibers*, are *One Body*: All simple *Barques*, *Piths*, *Parenchyma's* and *Pulps*, and as to their substantial Nature, *Pills* and *Skins* likewise, all but *One Body*: the several *Parts* of a *Vegetable* all differing from each other, only by the various *Proportions* and *Mixtures*, and variated *Pores* and *structure* of these *Two Bodies*. What from these two general Observations might reasonably be inferr'd, I shall not now mention.

15. §. The *Fourth* or *Innermost Cover* we may call the *Secondine*. The sight of which, by cutting off the *Coats* of an *Infant-Bean*, at the *Cone* thereof, in very thin *Slices*, and with great *Caution*, may be obtain'd. While unbroken, 'tis transparent; being torn and taken off, it gathers up into the likeness of a *Jelly*, or that we call the *Tredle* of an *Egg*, when rear-boyl'd: This *Membrane* in larger or elder *Beans*, is not to be found distinct. But (as far as our *Enquiries* yet discover) it may in most other *Seeds*, even full grown, be distinctly seen; as in those of *Cucumber*, *Colocynthis*, *Burdock*, *Carthamum*, *Gromwel*, *Endive*, *Mallows*, &c. 'Tis usually so very thin, as in the above-nam'd, as *Tab. 4. f. 16.* very difficultly to be discover'd. But in some *Kernels*, as of *Apricots*, 'tis very thick; and most remarkably such, in some other *Seeds*. That all these have the Analogy of one and the same *Cover*, which I call the *Secondine*, is most probably argu'd from their alike *Natures*; being all of them plain simple *Membranes*, with not the least *Fibre* of the *Lignous Body* or *Seed Branch*, visibly distributed in them: As also from their *Texture*, which is in all of them more close. See this *Part* in *Tab. 4.* As also amongst the *Figures* belonging to the *Fourth Part* of the *Fourth Book*.

16. §. The *Concave* of this *Membrane* is filled with a most transparent *Liquor*, out of which the *Seed* is formed; as in cutting a *petite* and *Infant-Bean*, may be seen; and yet better in a young *Walnut*. In *Beans* I have observed it to turn, upon boiling, into a tender white *Coagulum*.

17. §. Through this *Membrane*, the *Lignous Body* or *Seed-Bran-ches* distributed in the inner *Coat*, at last shoot downright two slender *Fibres*, like two *Navel-Strings*, one into each *Lobe* of the *Bean*.
 Tab. 4. f. 18. The places where the said *Fibres* shoot into the *Lobes*, are near the *Basis* of the *Radicle*; and by their *Blackishness* well enough remark'd: but the *Fibres* themselves are so very small, as scarcely to be discern'd. Yet in a *Lupine*, of the larger kind, both the places where the *Navel-Fibres* shoot into the *Lobes* (which here from the *Basis* of the *Radicle* is more remote) and the *Fibres* themselves, are fairly visible. For the *Seed-Branch*, upon its entrance into the *Coat* of the *Lupine*, is presently divided into two *main Branches*, and those two into other less; whereof some underly, others aloft, run along the *Coat*, and towards its other end meet and are inosculated: where about, two opposite, shallow, round, and most minute *Cavities*, answerable to two *Specks* of a *Cartilaginous* gloss, one in either *Lobe*, may be observed; which
 Tab. 4. f. 17. *Specks* are the ends of the said *Navel-Fibres*, upon the ripening of the *seed* there broken off. These *Fibres* from the *Superficies* of each *Lobe*, descend a little way directly down: presently, each is divided into two *Branches*, one distributed into the *Lobes*, the other into the
 Tab. 4. f. 18. *Radicle* and *Plume*, in the manner as in the *First Chapter* is described. And thus far the *History*. I shall now only with a brief account of the *Generation* of the *seed*, as hereupon dependent, conclude this *Dis- course*.

An Account of the Generation of the Seed. 18. §. LET US say then, that the *Sap* having in the *Root, Trunk* and *Leaves*, passed divers *Concoctions* and *Separations*, in the manner as they are said to be perform'd therein; 'tis now at last, in some good maturity, advanced towards the *Seed*.

19. §. The more copious and cruder part hereof is again separated by a free reception into the *Fruit*, or other *Part* analogous to it: being either sufficiently ample to contain it, or at least laxe enough for its transpiration, and so its due discharge. The more Essential part is into the *Seed-Branch* or *Branches* entertain'd. Which, because they are evermore of a very considerable length, and of a Constitution very fine, the said *Sap* thus becomes in its Current therein as in the *Spermatick Vessels*, still more mature.

20. §. In this mature estate, from the *Seed-Branch* into the *Coats* of the *seed*, as into the *Womb*, 'tis next delivered up. The meaner part hereof again, to the *Outer*, as *Aliment* good enough, is supplied. The finer part is transmitted to the *Inner*; which being, as is said, a *Parenechymous* and more spacious *Body*, the *sap* therefore is not herein, as in the *Outer*, a meer *Aliment*; but in order to its being, by *Fermentation*, farther prepared.

21. §. Yet the *Outer Coat*, being on the contrary hard and dense; for that reason, as it admitteth not the *Fermentation* of the *Sap* so well within it self; so doth it the more promote and favour it in the *Inner*; being *Bounds* both to it and its *sap*; and also quickneth the *process* of the whole *Work* in the formation of the *Seed*.

22. §. Nor doth the Outer Coat, for the same reason, more promote, than declare the purity of the Sap now contained in the Inner: For being more hard and dense, and so not perspirable, must needs suppose the Parts of the Sap encompassed by it, since thus incapable of any evacuation, to be therefore all so choice, as not to need it.

23. §. The Sap being thus prepared in the Inner Coat, as a Liquor now apt to be the *Substratum* of the future *Seed-Embryo*; by fresh supplies, is thence discharg'd. Yet that it may not be over-copious; which, because of the laxity of the Inner Coat, from whence it issues, it might easily be: therefore, as the said Inner Coat is bounded without, by the upper Coat; so by the *Secundine*, is it bounded within. Through which *Secundine* the Sap being filtr'd, or, as it were, transpiring; the depositure hereof, answerable to the *Colliquamentum* in an Egg, or to the *Semen Mulibre*, into its Concave at last is made.

24. §. The other part of the purest Sap embosom'd in the *Ramulets* of the *Seed-Branch*, runs a Circle, or some progress therein; and so becomes, as the *Semen Masculinum*, yet more elaborate.


25. §. Wherein also, lest its Current should be too copious or precipitant, by their *co-arture* and *divarication* where they are inosculated, it is retarded; the noblest portion only obtaining a pass.

26. §. With this purest Sap, the said *Ramulets* being supplied, from thence at last, the *Navel-Fibres* shoot (as the primitive *Artery* into the *Colliquamentum*) through the *Secundine* into the aforesaid *Liquor* deposited therein.

27. §. Into which *Liquor*, being now shot, and its own proper Sap or *Tinctures* mixed therewith, it strikes it thus into a *Coagulum*; or of a *Liquor*, it becomes a *Body consistent* and truly *Parenchymous*. And the supply of the said *Liquor* still continu'd, and the shooting of the *Navel-Fibres*, as is above described, still carried on, the said *Coagulation* or *Fixation* is therewith likewise.

28. §. And in the Interim of the *Coagulation*, a gentle *Fermentation* being also made, the said *Parenchyma* or *Coagulum* becometh such, not of any Texture indifferently, but is thus rais'd (as we see Bread in Baking) into a *Congeries* of *Bladders*: For such is the *Parenchyma* of the whole *Seed*.

F I N I S.



THE
ANATOMY
OF
ROOTS;

Presented to the ROYAL SOCIETY at several times,
in the Years, 1672 & 1673.

With an Account of the

VEGETATION OF ROOTS,

Grounded chiefly hereupon.

The SECOND BOOK.

By NEHEMIAH GREW M.D. Fellow of the
Royal Society, and of the College of Physicians.

The Second Edition.

LONDON,

Printed by *W. Rawlins*, 1682.

N 2

ANATOMY

OF THE

VEGETATION OF ROOTS

BY HENRY SHARPE

1857

Printed by M. G. Lewis

TO THE
Right Honourable
WILLIAM
Lord Vi-Count *BRONCKER*
THE
PRESIDENT
AND TO THE
Council and Fellows
OF THE
ROYAL SOCIETY.

MY LORD,

IF the Dedication of *Books* were not in use; yet here, I think, I might have been a Precedent. The promotion of *Phytological Science* is one Part of *Your* Work; and 'tis *You* have called me to the management of this Part; for sometime, have intrusted me herein; and by *Your* most favourable and candid acceptance of what I have performed thus far, have encouraged me hereunto: I therefore present but *Your* Own, into *Your* Hands.

The great Honour and Advantage of *Your* Fellowship, I first obtained, by Mediation of Dr. *Wilkins*, the late most Reverend *Bishop* of *Chester*. Whom I cannot name, without saying thus much of him, That He was a Per-
son

The Epistle Dedicatory.

son of that eminent and happy Worth, which, as it was too good, to fear envy; so is it too great, to need an Elogie.

With Him, it was, *You* were pleased to commit to Me, the further prosecution of this *Work*; the Beginnings whereof, were by *Your Order* formerly made publique. Had I consulted my own Abilities altogether, I should scarcely have ventured upon it; seeing very little, for which I could think well of my self, saving, That I had learned, upon good grounds, to think of *You* with greatest Honour. But I also considered, That to insist hereon too much, might be a reflection upon *Your* Judgments, who had thought fit to make choice of Me. And, That *You* were not more the Patrons of Wit, than of Industry; and of All, who shall endeavour to find out, or to confirm the Truth of Things. Withal, I looked upon *Nature*, as a Treasure so infinitely full; that as all Men together, cannot exhaust it; so no Man, but may find out somewhat therein, if he be resolved to Try.

In compliance therefore with *Your* Commands, I have hereunto devoted a very considerable part of my Time. These, adding force to my own Desires, of being somewhat instrumental to the Improvement of Medicinal, and other wholesom Knowledge: if peradventure, as we increase herein, we may become better, and more happy. As to which Improvement, though I could not hope; yet, I would not despair. I have already prepared the Soil, and made some Plantation: what remaineth behind, and the Vintage of the whole, will depend much upon the continued Influence of *Your* Beams: for how unpromising soever the Stock may be; yet the Fruit cannot but be somewhat matured, upon which *You* are pleased to shine. I am also confident, that the same Nobility and Goodness, which accept the endeavours, will likewise pardon the faults, of,

My Lord,

Your Lordships most humbly

and most sincerly

devoted Servant

September 1.
1673.

NEHEMIAH GREW.

THE

THE
CONTENTS.

The FIRST PART.

CHAP. I.

OF the Original of Roots, §. 1, 2, 3. Of their Figures, 4, to 8. Of their Motions, 9, to 15. And of their Ages, 16, to the end.

CHAP. II.

OF the Skin. Its external Accidents, and Original, §. 1, 2. Compounding Parts. Whereof the one Parenchymous, 3. The other Lignous, 4, to the end.

CHAP. III.

OF the Barque. Its Original and external Accidents, §. 1. Size, 2. Compounding Parts: Whereof the one Parenchymous, 3. The Bladders of the Parenchyma, 4, 5, 6. The Diametral Portions, 7, to 11. The other Part, Lignous, consisting of long Pipes or Vessels, 12, to 17. Of several Kinds, 18, to 23. In different Proportion, 24, 25. And in different and elegant Position. 26, to the end.

CHAP. IV.

OF that Part of the Root next within the Bark; in Trees and Shrubby Plants, called the Wood. Hereof the Parenchyma, §. 1, 2, 3, & 7. The Lignous Portion: of which, the Sap-Vessels, 4. The Aer-Vessels, 5, 6. The Position of the Former, 8, 9. Of the Latter, 10, 11, 12. Their Proportion, 13, 14, 15. The Latter, sometimes a little tapering. 16. Their Texture, 17, to 22. Content, 23.

CHAP.

The Contents.

CHAP. V.

OF the Pith. Found in the upper part of most Roots, §. 1. Its size and shape, 2. Sap-Vessels, 3. Original, 4, 5. Bladders, 6, Fibres and Texture, 7, to 11. That of the Insertions and Barque the same, 12. Hence, the Original of the Aer-Vessels conjectured, 13. What the whole Body of a Root, concluded, 14, 15. The Contents of the Pith, 16.

The SECOND PART.

Theology, the Beginning and End of Philosophy, §. 1, to 6.
The Divine Wisdom seen in the Growth of Plants, 7. If we observe,
How the Ground is Prepared, 8, to 14.
How the Sap is Imbibed, and Distributed to the several Parts of the Root, 15, to 28.
How the several Parts are Nourished and Formed, 29, to 35.
How the several Parts receive their respective Situation, 36, to 40.
How Roots receive their different Size and Shape, 41, to 47.
How Roots receive their different Motions, 48, to 53.
How Roots are differently Aged, 54, 55, 56.
How the Liquors and other Contents of the several Parts are made 57 to 63.
How the Odors of Roots are made, 64.
How their Colours, 65, to 67.
How their Tasts, 68, to the end.

THE
ANATOMY
OF
ROOTS;

PROSECUTED
With the bare EYE,
AND WITH THE
MICROSCOPE.

PART I.

CHAP. I.

Of the ORIGINAL, FIGURES, MOTIONS,
and AGES of ROOTS.



BEING TO speak of *Roots*; it is requisite, for our better understanding of what follows, that some things, as to their *Original*, *Figures*, *Motions* and *Ages*, be premised.

1. §. *Roots*, taken altogether, have a *Threefold Original*. Either from the *Radicle*; as all *Roots* which come of the *seed*: or from the *Trunk* or *Caulis*, above ground; as in *Strawberry*, *Chamemile*, and many other *Creepers*: or from the *Trunk* or *Caulis*, after it is sunk under ground; as in *Primrose*, *Bistort*, and many others; and presently shall be shewed how.

2. §. In the Growth of a *Bud*, and of a *Trunk-Root*, there is this observable difference; That the former, carries along with it, some portion of every *Part* in the *Trunk* or *Stalk*; whereof it is a *Compendium*. The latter, always shoots forth, by making a *Rupture* in the *Barque*, which it leaves behind, and proceeds only from the inner part of the *Stalk*.

3. §. As also, That in a *Bud*, the *Lignous Part* is spread abroad, so as to encompass a *Pith*. Whereas in a *Trunk-Root*, it makes a solid Thred standing in the Center. Which is the Cause of its descending into the Ground: as is already, in the *First Book*, and shall in This be further shewed.

4. §. **ROOTS** are generally distinguished, as to their *Figures*, in being more *Entire*, as is that of *Liquirish*; or *Parted*, as of *St. Johnswort*. *Parted* or *Forked*, either at the *Bottom*, as most *Roots*; or at the *Top*, as *Dandelyon*, and some others. A thing very odd, and unintelligible, without the knowledge of the *Motions* of *Roots*; whereof presently.

5. §. *Parted*, again, are either *Ramified*, as that of *Cumfry*; or *Manifold*, as of *Crowfoot*: both are *Parted*; but the former, by the subdivision of greater *Branches*, into lesser; these, when divers *Strings*, have all their distinct original from one *Head*. Some are *Straight*, as a *Radish*; others *Crooked*, as *Bistort*. *Smooth*, as *Bugloss*; or *Stringy* all round about, as *Columbine*. And to *Carnations*, this seems to be peculiar, That sometimes many of the *Strings* run parallell with the *Wood* of the great *Root*, through the *Barque*, or betwixt the *Wood* and the *Barque*.

6. §. Again, some are *Thick*, as *Rhubarb*; *Slender*, as the *Vine*. *Long*, as *Fenil*; *Short*, as a *Turnep*: which are distinct from *Great* and *Little*; in that these, are so called with respect to several *Roots*; those, with respect to the several *Dimensions* of one. *Short*, are *Stubbed*, as *Iris tuberosa*; or *Round*, as *Dracontium*. *Round* are *Tuberous*, or *Simply Knobbed*, as *Rape-Crowfoot*; *Bulbous*, that is *Scaled*, as some *Lilys*; or *Shell'd*, as an *Onion*. Where note, That all *Bulbous Roots*, are, as it were, *Hermaphrodites*, or *Root* and *Trunk* both together: for the *Strings* only, are absolute *Roots*; the *Bulb*, actually containing those *Parts*, which springing up, make the *Leaves* or *Body*; and is, as it were, a *Great Bud* under ground.

7. §. *Roots*, again, are *Even* or *Uneven*; *Even*, are *Cylindrical*, as *Eryngo*; or *Pyramidal*, as *Borage*. Growing smaller *Downwards*, as do most; or *Upwards*, as *Skirrets*. *Uneven*, are *Pitted*, as *Potato's*, where the *Eyes* or *Buds* of the future *Trunks* lie inward; or *Knotted*, as *Jerusalem-Artichoke*; where they stand out. These Differences, are also *Compounded*: so some *Roots* are both *Entire* and *Smooth*, as *Peony*; others *Entire*, but *Stringy*, as *Clary*: that is, neither *Ramifid*, nor yet *Brushy*, or divided at the *Top* into severall small *Strings*; but a *Single Root* surrounded with many *Hairy Threds*. Some both *Plain* in some parts, and *Knobbed* in others, as *Filipendula*, *Lilium non bulbosum*, and others.

8. §. Some also have two or more *Roots*; and those of one *Kind*: of which, some are distinctly fastend to the bottome of the *Stalk*, as in *Dogstones*; some stand one under another, so as only the uppermost is fasten'd to the *Stalk*, as in *Dragon*, *Crocus*, and others. And there are some, which have not only two *Roots*, at the same time; but those also of two distinct *Kinds*, as in *Bistort*; one of them, a slender strait *Cylindrick* and *horizontall Root*; the other large and crooked, and bred of the *Descending Trunk*; as in speaking next of the *Motions* of *Roots*, will be understood, how. All which, with other Differences by

by Those that undertake the *Descriptions* of *Plants*, are accurately to be Noted. But the Differences, above mentioned, will serve for our present Purpose.

9. §. THE MOTIONS of *Roots* are also divers. Sometimes Level, as are those of *Hops*, *Ammi*, *Cinquefoyle*; and all such as properly *Creep*. Sometimes Perpendicular, as that of *Parsnep*: Which is different from Straightness; for some Straight *Roots*, are Level. Both of them are either Shallow or Deep: some run Level, and near the *Turf*, as *Woodbind*, *Wild Anemony*; others lower, as *Dogs-Grass*. Some strike down, but a little way, as *Stramonium*; others grow deep, as *Horse-Radish*: Which is different from being Long; for many long *Roots*, are Level, as *Hops*.

11. §. Some again Descend, as *Tulips*, and other Bulbous *Roots*, which differs from growing only Downwards; in that here, the *Head* of the *Root* is Immoveable; but in Descending, the whole *Root* obtaineth different Places, running deeper, time after time, into the *Earth*. Some also Ascend, sometimes, and in some part, appearing above ground, as *Turneps*.

11. §. These *Motions* are also Compounded; both in respect of the several Parts of the *Root*, and of several Times. So the main *Root* of *Primrose*, is Level; the Strings are Perpendicular. The *Roots* of most *seedlings* grow Downward and Upward, or shoot out in length at both Ends, at the same time. Those of *Bistort*, *Iris*, and some others, grow, in part, both Downward and Upward at several times: Whence it is, that *Bistort* is Crooked, with some resemblance to an S, according to its *Name*; And that some Parts of *Iris-Root* appear oftentimes above the ground.

12. §. There is also another *Motion*, in some *Roots*, not heeded; and that is *Contortion*: whereby, without being moved out of their Place, they are Writhed or Twisted; as a piece of Cloath is, when the Water is wrung out of it; as in *Carduus*, *Sonchus*, and others: whether always I cannot say. This *Motion* cannot be noted, without stripping off the *Barque*; whereby the *Vessels* may be seen, sometimes, to make two or three Circumvolutions. This *Motion* seems to be governed by the winding of the *Stalk*; and therefore to begin at the *Head*, and terminate at the Poynt or lower end of the *Root*, which is immoveable.

13. §. BUT ABOVE all the *Motions* of *Roots*, not observed, the most remarkable is that of DESCENT. Which, although it hath been noted, by some *Botanicks*, of *Bulbous Roots*; yet of these only: Whereas it is the Property, of a great many more; and those, of very different Kinds; probably, of the far greater number of *Perennial Roots* of *Herbs*; as of *Arum*, *Rape-Crowfoot*, *Valerian*, *Brownwort*, *Bear-foot*, *Tansy*, *Lychnis*, *Sampier*, *Primrose*, *Ammi*, *Avens*, *Wood-sorrel*, *Iris*, and others. Of all which *Plants*, it is very observable, That their *Root*, is annually renewed, or repaired, out of the *Trunk* or *Stalk* it self. That is to say, The *Basis* of the *Stalk* continually, and by insensible Degrees, descending below the surface of the *Earth*, and hiding it self therein; is thus, both in Nature, Place, and Office changed into a true *Root*. Which *Root*, by the continuance of the said

Motion of the *Stalk*, also *Descends*; and so, according to the durability of its Substance, becomes a shorter or longer *Root*; the Elder or Lower Portion thereof, Rotting off, by the same Degrees with the Generation of the Upper, out of the *Stalk*. So in *Brownwort*, the *Basis* of the *Stalk* sinking down by degrees, till it lies under Ground, becomes the upper part of the *Root*; and continuing still to sink, the next year, becomes the lower Part; and the next after that, rots away; a new Addition being still yearly made out of the *Stalk*, as the elder Parts yearly rot away. So in *Dragon*, *Crocus*, and the like, where the *Root* is double; the *Basis* of the *Stalk*, this year; the next, becomes the Upper-*Root*; after that, the Lower-*Root*; and at the length dies and is consum'd.

Tab. 5. f. 6,
& 7.

14. §. The Demonstration hereof, is taken, more evidently, from some *Roots*, than from others; as from the Level and Knobed *Roots* of *Wood-sorrel*, *Primrose*, &c. For the *Leaves* of those *Plants* rotting off successively, and the *Bases* of those *Leaves* gradually descending into the Ground; each *Basis* is thus nourished with a more copious *Sap*, and so swelled into so many thick *Knots*. It may likewise be gather'd in some, from the like Position of the *Vessels* or Woody Parts, in the *Root*, as in the *Trunk*; as in *Bares-foot*. As also, from the *Root* of the *Iris Tuberosa*: where, although the *Leaves* fall off close to the Surface of the *Stalk*; yet after that is sunk down, and swell'd into a *Root*, the *Seats* of the perished *Leaves*, and the Ends of the *Vessels* belonging to them, are not obscurely visible; whereby the *Root* is wrought, as it were, with several *seames* and *Prickt-Lines*; the *seams* shewing the setting on of the *Leaves*; and the *Pricks*, the Terminations or broken Ends of the *Vessels*: which ends, are still more apparent, upon the stripping off the *Barque*. I considered likewise, That as among *Animals*, there are many, which are not Bred of *Eggs*, immediately; but are Transformed, one *Animal* into another: So, it is more than probable, That among *Plants*, there are not a few Instances of the like *Transformations*; whereof, this is one.

Tab. 5. f. 1,
& 2.

Tab. 5. f. 4.

15. §. The Cause of this *Descent*, so far as it is dependent on the Inward Conformation of the *Root*, I shall shew in the following Part. But the Immediate Visible one, are the *String-Roots*, which this kind of *Trunks* frequently put forth: which, descending themselves directly into the Ground, like so many *Ropes*, lug the *Trunk* after them. Hence the *Tuberous-Roots* of *Iris* upon the rotting or fading away of the *String-Roots* hanging at them, sometimes a little Re-ascend. Hence also the *Shape* of some *Roots* is Inverted: For whereas most are parted downwards, into several *Legs*; some are parted upwards into divers *Necks*, as *Dandelyon*, and others. For these *Roots* sending forth at the top several *Trunk-Buds*, the said *Buds* successively put forth new, and cast their old *Leaves*; and continually also making their *Descent*, are at length formed into so many *Necks*, of three, four, five, or more Inches long, under Ground.

Tab. 5. f. 5.

16. §. HENCE ALSO we understand, in what particular way, some *Roots* become *Perennial*. Some are wholly so, as those of *Trees*, *Shrubs*, and divers other woody *Plants*. Others, in part, or by a new *Progenies* of *Roots*, from the old Head or Body, in the room of those that die yearly, or after a certain Time; as of *Lilium non bulbosum*,

sum, Jerusalem Artichoke, Potato, Dog-stones, Monks-hood, little Celandine, and others. In which Plants, one or more of their Roots are firm, the other spongy and superannuated; and partly, by the ravine of the Trunk, and other younger Roots, reduced to a Consumption and Death.

17. §. With these, *Tulips*, and other *Bulbous-Roots* consort: For the several *Rindes & Shells*, whereof chiefly, the *Bulb* consists, successively perish and shrink up into so many thin and dry *Skins*: betwixt which, and in their Centre, other *Leaves* and *Shells*, being successively formed, the *Bulb* is thus perpetuated. In the same manner the *String-Roots* also succeed one another annually. So that at the end of divers Years, although it be still looked upon as the same *Individual Root*, yet it is, in truth, Another, as to every particle thereof.

18. §. Lastly, many other *Roots* are perpetuated by the aforesaid Descent of the *Trunk*; out of which, it is still annually Repaired, as by the gradual perishing of its lower parts, it is Diminished; as hath been said. Whence also we see the reason of the Rugged and Blunt extremities of these, and some other *Roots*, as of that *Plant* superstitiously called *Devils-bit*: because the end of it seems to be bitten off. Yet doth it not appear so originally; but the Lower part thereof rotting off, as the Upper descends; the living remainder, becometh stumped, or seemeth Bitten. Thus far of the *Original, Shapes, Motions, and Ages of Roots.*

Tab. 5, f. 3.

CHAP. II.

Of the SKIN.



NEXT proceed to the several *Parts* whereof a *Root* is Compounded. The outer *Part* of all is the *Skin*; which is common to all *Roots*. 'Tis diversly Coloured: Whiter in *Skirrets*; Yellow, in *Dock*; Red, in *Potato*; Brown, in *Lovage*; Black, in *Bugloss*. Its Surface, sometimes Smooth, as in *Hors-radish*; Rough, as in *Scorzonera*. And the *Skins* of the several *Shells* of a *Tulip-Root*, taken up fresh, look as if they were perforated with a great many small holes. 'Tis of various Size; very Thin, in *Parsnep*; somewhat Thick, in *Bugloss*; very Thick in *Iris*. Sometimes it is Opaque, as in *Thistle*; and sometimes Transparent, as in *Madder*.

2. §. Every *Root* hath successively two kinds of *Skins*: the one, Coëtaneous with the other *Parts*; and hath its original from that which involveth the *Parts* of the *Seed* it self. The other, Postnate, succeeding in the room of the former, as the *Root* ageth; and is ori-nated from the *Bark*. So in *Dandelyon*, the old *skin*, looked upon about the beginning of *May*, seems to have been one of those several

Rings

Rings, which the precedent year composed the *Cortical Body* of the *Root*: but by the Generation of a new *Ring*, next the *Wood*, is now thrust off and shrunk up into a *skin*. So also in the *Roots* of *Bugloss* and *Horse-Radish*, as far as the *Bladders* in the former, and the *Vessels* in the latter are Radiated; the *Cortical Body* seems either annually or oftener, to shrink up into another new *skin*, as, the old ones fall off. And sometimes, perhaps, as in *Asparagus*, the whole body of the *Perpendicular Roots*, except the woody *Fibre* in the Centre, becomes the second *skin*. So that the wearing away of the old *Skin*, succeeds the derivation of the new one; as in *Descending Roots*, the Consumption of the Lower *Parts*, doth the Generation of the Upper. Because the *Barque* swells, and grows sometimes faster than the *Skin* can fall off, or give way to it: therefore are the *Roots* of many *Herbs*, *Barquebound*, as well as the *Trunks* of *Trees*.

Tab. 14, 15.

Tab. 10.

3. §. This *Skin* is usually, if not always, compounded of two Kinds of Bodies: which also is probable of the *Coëtaneous*. The one, *Parenchymous*, and frequently constructed of exceeding little *Cells* or *Bladders*; which in some *Roots*, as of *Asparagus*, cut traverse, and viewed through a *Microscope*, are plainly visible. These *Bladders* are of different Sizes; in *Bugloss*, larger; in *Asparagus* less; and sometimes they coincide and disappear. But in these, and all other *Roots*, even where these *Bladders* appear not, the *Parenchyma* of the *skin*, is of the same Substantial *Nature*, with that other more vivid and bulky one of the *Bark*: As is manifest, from its being thence Originated; and alike Conformed, as shall be seen; and not only adjacent to it, as a *Glove* is to the *Hand*; but continuous therewith, as the parts of a piece of *flesh*, are one with another.

Tab. 10.

Tab. 14.

4. §. OF THIS *Parenchymous Body*, the *skin* consisteth chiefly, but not wholly; there being many *Lignous Vessels* which are *Tubulary*, mixed therewith: which, though hardly by the *Microscope*, yet otherwise, is demonstrable. For in tearing the *Skin*, you shall do it more easily by the length, than breadth; because, by the first way, the continuity only of the *Parenchyma*, is dissolved; but by the latter, both of this, and of the *Vessels*, these being posited by the length of the *Root*: So that, as by the smallness of the *Bladders* of the *Parenchyma*, the *Skin* is Dense; so by these *Vessels*, is it Tough.

5. §. Again, if you cut a *Root* traverse, and let it lie by for some time, all the parts, where there are no *Vessels*, shrink below the surface of the cut-end; but where-ever These are posited, there is no shrinking; which oftentimes, evidently appears also in the *Skin*: because the said *Vessels*, though, as the *Bladders*, they may coincide; yet they cannot visibly shorten or shrink up in length; no more than a *Straw*, whose sides may yet be easily crushed together.

6. §. Further, the *Root* being cut traverse, if, near the cut-end, you very gently press the side of the *Root* with the edge of your *Nail*, the *Sap* will thereupon arise sometimes from the *Skin*; in the same manner, as from any other part of the *Root*, where the like *Vessels* are posited. And although the *Sap* may likewise be expressed from the *Pith*, and other *Parts* where sometimes, there are none of these *Vessels*; yet not without a solution of their continuity; which here doth not follow; as appears, from the disappearing of the *Sap*, together with the intermission

termiffion of the preffure; the faid *Veffels* then dilating themfelves by a Motion of *Reftitution*, and fo fucking up the *Sap* again.

7. §. Hereunto may be added the Testimony of fight; the very *Veffels* themfelves, in many *Roots*, coming under an apparent view, and ftanding in the utmoft furface of the *Root* all round about, as in that of *Liquirish*, *Columbine*, *Scorzoner*a, and others. Which *Experiments*, I have here, once for all, more particularly fet down; becaufe I fhall have occafion, hereafter, to refer to them.

C H A P. III.

Of the B A R Q U E.



NEXT WITHIN the *Skin* lieth the *Barque*. 'Tis fometimes Yellow, as in *Dock*; Red, in *Biftort*; but ufually, and in *Seed-Roots*, I think, always White. It is derived from the *Seed* it felf; being but the extension or prolongation of the *Parenchyma* of the *Radicle*; One of the three *Organical Parts* of the *Seed*, described in the *First Chapter* of the *First Book*.

2. §. It is variously Sized; fometimes very Thin, as in *Jerufalem Artichoke*, *Goats-beard*, and in moft *Trees*; where it alfo retains the Name of a *Barque* or *Rind*. Sometimes 'tis more Thick, and maketh up the far greateft protion of the *Root*, as in the *String-Roots* of *Asparagus*, in *Dandelion*, and others. The thinnest and the thickest are all analogous, and obtain the fame general Ufes. The degrees of its Size, amongst all *Roots*, may be well reckoned about Twenty, and feen in the following examples, *fc.* *Beet*, *Dropwort*, *Jerufalem Artichoke*, *Tab. 7, 8, 9.* *Orpine*, *Valerian*, *Goats-beard*, *Nettle*, *Brownwort*, *Columbine*, *Celandine*, *Asparagus*, *Horse-Radish*, *Peony*, *Bryony*, *Eryngo*, *Borage*, *Lovage*, *Dandelion*, *Parsnep*, *Carrot*, &c. In the *Root* of *Beet*, fcarce exceeding a good thick *skin*: but in a *Carrot*, half the Semidiameter of the *Root*, or above half an Inch over in fome places: and that of *Dandelion*, fometimes, in proportion with the *woody Part*, twice as thick: the reft of Several intermediate Degrees: And to moft *Roots*, this is common, To have their *Barque* proportionably thicker, at the bottome than at the top.

3. §. IT IS Compounded of two *Bodies*. The one *Parenchymous*; Continuous throughout; yet fomewhat Pliable without a folution of its Continuity. Exceeding Porous; as appeareth from its fo much fhinking up, in drying. The *Pores* hereof are extended much alike both by the length and bredth of the *Root*; therefore it shrinketh up, by both thofe Dimensions, more equally. And they are very Dilative; as is alfo manifelt from its reftorablenefs to its former bulk again, upon
its

its infusion in Water: that is to say, *It is a most curious and exquisitely fine wrought Sponge.* Thus much the Eye and Reason may discover.

Tab. 10,
& sequent.

4. §. The *Microscope* confirms the truth hereof, and more precisely shews, That these *Pores* are all, in a manner, Spherical, in most *Plants*; and this *Part*, an Infinite Mass of little *Cells* or *Bladders*. The sides of none of them, are Visibly pervious from one into another; but each is bounded within it self. So that the *Parenchyma* of the *Barque*, is much the same thing, as to its Conformation, which the Froth of *Beer* or *Eggs* is, as a fluid, or a piece of fine *Manchet*, as a fixed Body. The Sides also of these *Bladders* are as transparent, as those of Water; or the Bodies of some *Insects*.

Tab. 13, 14.

6. §. But their Size is usually much smaller; and their Posture more Regular than those in *Bread* or *Water*. In all *Roots* they are so small, as scarcely, without the *Microscope*, to be discerned: yet are they of different Size, both in the same, and in divers *Roots*; the varieties whereof, amongst all *Roots*, may be reduced to about Ten or Twelve according to the *Standard*, in *Tab. 11*. Some of those in *Dandelion*, being of the Smallest; and in *Bugloss*, of the Greatest. They are posited, for the most part, at an Equal Height; and piled evenly one over another: So that, oftentimes, they visibly run in Ranks or trains, both by the length and breadth of the *Roots*, as in the *Root* of *Bugloss*, or of *Dandelion*, split through the middle, may be seen. Although they are usually Spherical, yet sometimes, and in some places, they are more oblonge, as in the outward part of the *Barque* of *Bugloss*. These *Bladders*, are sometimes best seen, after the *Root*, being cut traverse, hath layn by a while, to dry.

Tab. 14.

6. §. They are the Receptacles of *Liquor*; which is ever Lucid; and I think, always more Thin or Watery. They are, in all *Seed-Roots*, filled herewith; and usually, in those also which are well grown, as of *Borage*, *Radish*, &c.

Tab. 8, 9.

7. §. THIS *Parenchymous Part*, in many *Roots*, is of one Uniform *Contexture*; as in *Asparagus*, *Horf-Radish*, *Peony*, *Potato*, and others. In many others, it is, as it were, of a Diversified *Woof*; the *Bladders* being, though every where Regular, yet either in Shape, Size, or Situation, different in some Parts hereof, from what they are, in other intermediate ones. For these Parts, are like so many *White Rays*, streaming, by the Diameter of the *Root*, from the inward Edge toward the Circumference of the *Barque*; as in *Lovage*, *Melilot*, *Parshp*, &c. cut transversly, is apparent. They are, though not in direct Lines, continued also by the length of the *Root*; so that they are, as it were, so many *Membrances*, by which the other Parts of the *Barque*, are determined.

Tab. 9.

Tab. 8.

8. §. The Continuation of these *Diametral Rays*, or *Portions*, is divers: sometimes, but half through the *Barque*, or somewhat more, or less, as in *Melilot*. And it is probable, that to the *Roots* of all or most *Trefoyls*, and also of the *Leguminous* Kind, this is proper, To have their *Diametral Rays* come short of the Circumference. Sometimes, they run quite through to the very *Skin*, as in *Lovage*. And I think, in the *Roots* of all *Umbelliferous Plants*: In which therefore, the *Skin* seems to have a closer *Communion* with the *Diametral Rays*, and to be originated especially therefrom. They usually stand at an Equal Distance in the same *Root*: But with respect to divers *Roots*, their Distance

stance varies; so less, in *Parsnep*, greater in *Bugloss*. They are commonly *Tab. 7, 8.*
 Rectilinear, as in *Lovage*; but sometimes winding to and fro, as in
 a *Carrot*. *Tab. 8.*

9. §. They are not always of one Size: in a *Carrot* near the
 Inner Edge of the *Barque*, exceeding Slender, and scarcely discerna-
 ble; in others, Thicker, as in the Three greater ones of *Melilot*, and
 in common *Chervil*. Both by their Distance, and Size, they are also *Tab. 8, 9.*
 less or more Numerous; some, only as they are nearer; some, as smal-
 ler; others, as both. And 'tis proper, I think, to the *Intybous* kind,
 either to have none, or but a few. Sometimes they are of the same
 Thickness quite through the *Barque* from edge to edge, as in *Marsh-Tab. 7.*
Mallow. And sometimes are considerably spread or dilated as they
 approach the *Skin*, wherewith they are joyned, and whereinto they
 more visibly run, as in *Parsley*, or the smaller part of the *Root* of
Lovage. And in some *Roots*, as of *Scorzonera*, at some times of the *Tab. 8.*
 year, when less succulent, almost the whole *Parenchyma* seems to be
 of the Nature of the *Diametral Rays*, in other *Roots*. The *Bladders*
 of these *Diametral Portions*, are sometimes, greater than those of the
 other *Parenchymous Parts*, as in *Parsley*; and I think sometimes less.
 Yet as there, so here, variously sized; to about six or eight De-
 grees; and those of *Parsley* about the third, fourth, and fifth. Their
 Figure is Sometimes more oblong; and their direction or respect more
 towards the Center of the *Root*.

11. §. As the other *Parenchymous Parts* of the *Barque*, are the
 Receptacles of *Liquor*; so these, (where they are) of *Aer*. This is ar-
 gued, From their being more White, and not Transparent, as such
Roots and *Parts* use to be, which are more copiously and equally fil-
 led up with *Liquor*: as the *Pith* of *Elder*, which, in the old *Stalks*, is
 White; was once, and by being well soaked, will become, again
 Transparent. And from their being more dry and voyd of *Liquor*;
 whereupon their *Bladders*, which cannot be *Vacuities*, must be filled
 with more or less *Aer*, mixed with the *Sap* or the *Vaporous* parts there-
 of. This is more observable in those *Diametral Portions*, which ter-
 minate upon, and run into the *Skin*.

12. §. THE BARQUE is not only of a divers *Woof*, but as is
 said, of a Compounded Substance; there being a certain number of
Lignous Vessels, fewer or more, in some place or other, mixed with
 the *Parenchymous Part* above described; and some way or other, are
 demonstrable in all *Roots*. As by the Toughness of the *Barque*, when
 pulled by the length. By the visible Continuation of the said *Vessels*
 through the length of the *Barque*, in the resemblance of small *Threds*. *Tab. 6.*
 And by the rising up of the *Sap* in the traverse cut of the *Root*, in
 such places of the *Barque*, where these *Threds* terminate: as the ex-
 istence of the same *Vessels* in the *Skin*, was proved in the *Precedent*
 Chapter.

13. These Tubulary *Threds*, run not through the *Barque* in di-
 rect lines; but are frequently Braced together in the form of *Net-*
Work; The *Parenchymous Parts* every where filling up the spaces be- *Tab. 6.*
 twixt the Braced *Threds*; as in *Burnet*, *Scorzonera*, &c. the *Barque*
 being paired or striped off, is apparent.

14. §. They seem, at first, where they are Braced, to be Inosculated; so as to be pervious one into another. But a more accurate view, especially assisted by a *Miscroscope*, discovers the contrary. Neither are they woun'd any way one about another, as Threds are in a Rope: nor Implicated, as in ravled Yarn, or the Knots of a Net: but only contiguous or simply Tangent, as the several Chords in the Braces of a Drum: being thus joyned together by the *Parenchymous Parts*, as in speaking of the *Pith*, will be understood how. Yet do not always the same *Threds* belong and keep entire to one *Brace*; but are frequently parted into lesser *Threds*; which are transposed from *Brace* to *Brace*. Nor do they always, in whole or in part, presently after their contingence, mutually fall off again; but, oftentimes, run along collateral-ly joyned together for some space.

Tab. 6.

15. §. These *Braces* are of various number in divers *Roots*; more frequent in *Jerusalem Artichoke*, less in *Scorzoner*a, more rare in *Cumfry*. The *Threds* likewise are variously Divaricated; sometimes more, where the *Braces* are frequent, as in *Jerusalem Artichoke*; and sometimes less, where the *Braces* are rare, as in *Scorzoner*a, *Dandelion*: And in all *Roots*, more frequent towards the Inner Verge of the *Bark*.

16. §. By what is said, it is partly implied, That these *Threds*, are not Single *Vessels*; but a *Cluster* of them, Twenty, Thirty, or more or fewer of them together. Yet as the *Threds* are not Inosculated in the *Braces*; so neither are the *Vessels*, in the *Threds*. Nor yet Twisted; but only stand collateral together; as the several Single *Threds* of the *Silkworm*, do in Sleeve-Silk. Neither are these *Vessels* pyramidal, so far as the *Glass* will discover; or, from probable Reason, may be conjectured. Nor Ramified, so as to be successively propagated one from another, after the manner of the *Veins* in *Animals*: but Cylindrical, and Distinctly continued, throughout the length of the *Root*; as the several *Fibres* in a *Tendon* or *Nerve*.

17. §. THESE VESSELS are either themselves of divers kinds, or serve, at least, to constitute divers Kinds, in divers *Roots*: of the different Natures whereof, although there may be other ways whereby to judge; yet so far as by Inspection, we may do it, chiefly, by the Diversity of those *Liquors*, which they severally contain. Sometimes they yield a *Lympha*; and that Thin, as they do in a *Parsnep*; especially those that make a *Ring*, at the inward extremity of the *Bark*. See the *Root* it self. That this Clear *Sap* ascendeth only from these *Vessels*, is certain. Because no *Liquor* will do the like, from any *Parenchymous Part*, as *Chap. 2.* hath been said. And because it is of a different nature from the *Sap* contained in the *Bladders* of the *Parenchyma*; although of the same *Colour*, yet sensibly more *Sweet*.

18. §. Sometimes they yield a Thick and Mucilaginous *Lympha*, as in *Cumfry*, as appeareth by its tenacity. From the Mucilaginous Content of these *Vessels* it is, I suppose, that the *Sap* contained in the *Bladders* is rendred of the like nature, so far as it approaches hereto, which sometimes is more, as in *Marsh-mallow*; and sometimes but little as in *Borage*: For in pressing out the *Liquor* of this *Plant*, and then heating it over an indifferent fire; the far greater part hereof remaineth thin; only some certain strings and little bits of a gellied substance are mixed herewith; which as it seems, were originally the proper *Liquor* of these *Muciducts*.

19. §.

19. §. Oftentimes these Succiferous *Vessels* yield a *Milky* or *White Sap*; and sometimes *Yellow*, and of other colours as in *Sonchus*, and most *Cichoraceous Plants*; in *Angelica*, and most *Umbelliferous*; in *Burdock*; and divers *Thistles*, to which that is 'akin: in *Scorzonera*, *Common Bells*, and many other *Plants*, not commonly taken notice of to be *milky*. The *Milky Saps* of all which, although they differ in *Colour*, *Thickness*, and other *Qualities*; yet agree, in being more *Oily* than any of the *Lymphous Saps*. It being the mixture of the *Oily* parts with some other *Limpid Liquor*, but of a different *Nature*, which causeth them to be of a *Milky*, or other *Opacous Colour*, in the same manner as common *Oyl*, and a strong *Liquamen* of *Tartar*, shaked in a *Bottle* together, presently mix into a *White Liquor*. And although they will, for the greatest part, separate again; yet some of their parts, without any *Boiling*, or so much as the least *Digestion* with *Heat*, by *Agitation* only, or standing together for some time, incorporate in the form of a *Thin Milky-Sope*, which will also dissolve in *Water*. I suppose, therefore, That it is the *Volatile Salt*, chiefly, of these *Plants*, which being mixed with their *Oyl*, renders this *Liquor* of a *White* or other *Opacous Colour*.

20. §. Sometimes the *Oyl* will separate and discover it self: for if you cut a *Fenil-Root* traverse, after it hath layn some days out of the *Ground*; the same *Vessels*, which, in a fresh *Root*, yields *Milk*; will now, yield *Oyl*: the watery parts of the *Milk*, which in the drying of the *Root* are more evaporable, being spent.

21. §. All *Gums* and *Balsams* are likewise to be reputed the proper *Contents* of these *Vessels*: for These and *Milks*, are very near akin. So the *Milk* of *Fenil*, upon standing, turns to a *Clear Balsam*; of *Scorzonera*, *Dandelion*, and others, to a *Gum*. In the dried *Root* of *Angelica*, &c. being split, the *Milk*, according to the Continuation of these *Vessels*, appeareth, as *Blood* clodders in the *Veins*, condensed to an hard and shining *Rosin*. And the *Root* of *Helenium* cut *Tab. 9.* transversely, presently yields a curious *Balsame* of a *Citrine Colour*, and sometimes of the Colour of *Balsame* of *Sulphur*. I call it a *Balsame*; because it will not dissolve in *Water*. Yet not a *Terebinth*; because, nothing near so viscid or tenaceous as that is. But the *Root* of *Common Wormwood*, bleeds, from large *Vessels*, a true *Terebinth*, or a *Tab. 10. E.* *Balsame* with all the defining properties of a *Terebinth*; although that word be commonly used only for the *Liquors* of some *Trees*.

22. §. There is yet another kind of *Sap-Vessels*, which may be called *Vapour-Vessels*; as in *Docks*, at least some of them. For by the *Sap-Vessels* it is, that the *Barques* of *Roots* do *Bleed*. Of which, some *Bleed* quick and plentifully, as the *Umbelliferous* and the *Cichoraceous* Kinds. Some, very slowly and scarce visibly, as all or most *Trefoyls*, and of the *Leguminous* Kind. And some seem not to *Bleed*, as the *Dock*. Yet that this *Root*, hath also *Vessels* distinct from those that carry *Aer*; doth partly appear, from the different *Colour* they produce where they stand; as will better be understood anon, in speaking of the *Causes* of the *Colours* of *Roots*. As also from the *Toughness* of the *Barque*, in pulling it by the length; neither the *Parenchyma*; nor the *Aer-Vessels*, being of themselves *Tough*. But because the *Succus* or *Sap* they carry, seems to be a kind of *Dewy Vapour*, therefore, they may not improperly be called *Roriferous* or *Vapour-Vessels*.

23. §. THE *Sap-Vessels*, are not only of divers Kinds, in divers *Roots*, but in the same. Whether in all, I doubt: but in some it is certain they are: For if you cut a *Fenil-Root* traverse, both *Milk* and *Limpid Sap*, will presently ascend, and, upon accurate inspection, appear thereupon distinctly. So the *Roots*, both of *Trachelium* and *Enula*, Bleed both a *Lympha*, and a *Citrine Balsame*: and *Wormwood*, both a *Lympha*, and a *Terebinth*, at the same time. So also the *Root* of *Dandelion* being cut in *November*, seems to bleed both a *Milk* and a *Lympha*; the latter being drowned by the former at another time when it is more copious. Whether all *Roots* have *Lymphæducts*, is doubtful; but 'tis most probable, that they have, more or fewer; standing, for the most part, in a *Ring*, at the Inner Verge of the *Barque*: the *Sap* whereof, I suppose, is so far of common Nature in all *Roots*, as to be Clear, and less Oily.

24. §. THE Quantity of these *Vessels* is very different: In *Borage*, *Peony*, *Bistort*, but few; in *Asparagus*, fewer: in *Parsnep*, *Celandine*, many; in *Fenil*, *Marsh-mallow*, many more: and betwixt these extremes, there are many Degrees, as by comparing the *Roots* of *Horse-Radish*, *Turnep*, *Briony*, *Skirrets*, *Parsley*, *Goats-Beard*, and as many more as you please, may be seen. Amongst the several Sorts of *Docks*, they seem in *Patience*, to be the fewest; in *Red-Dock*, the most numerous. There are two ways of judging of their Number; Either as their Extremities are visible upon the traverse cut of the *Barque*; or as the *Barque* is diversly Brittle or Tough; being so, from the various Number of these *Vessels* therein, as in the *Second Chapter* hath been said.

25. §. The Quantity of the ascending *Sap*, is a doubtful argument, whether of the Number, or Size of these *Vessels*. For it is common to most *Milky-Roots*, for the *Milk* to ascend more copiously: yet in some of them, the *Vessels* seem, in proportion with the *Parenchymous Part*, not to be so numerous, as in some other *Roots*, where the ascending *Sap* is less; as by comparing the *Lacteals* of *Dandelion*, and the *Lymphæducts* of *Fenil* together, may appear: so that it should seem, that the bore of the *Lacteal Vessels*, is greater than that of the *Lymphæducts*.

26. §. THE Situation of these *Vessels*, as they appear, even to the naked Eye, in the transverse Section, is Various and Elegant. Sometimes they are posited only at the Inner Edg of the *Barque*, where they make a *Ring*, as in *Asparagus*. In which place and position, they stand in most, if not in all, *Roots*, how varioussly soever they are posited also otherwise. The Common *Crow-Foot* with numerous *Roots*, hath a *Ring* of *Sap-Vessels* next the *Skin*. So the *Barque* of *Monks-Hood*, is encompassed with a transparent *Ring* of *Sap-Vessels*. The *Ring* is either more Entire, as in *Eryngo*, *Brown-Wort*, *Valerian*, *Hop*, *Madder*, &c. Or it is a Prick'd *Ring*, as in *Buttyr-Bur*. Sometimes they are chiefly postur'd in a Prick-Ring, towards the outward part of the *Barque*, as in *Peony*: and some *Roots* are pricked all over the *Barque*, as of *Melilot*. In others, they stand not so much in Pricks, as Portions or *Columns*, as in *Cumfry*-

27. §. In others, again, they all stand in more continued Lines, either Rays or Diametral, as in *Borage*; or Peripheral, as in *Celandine*. The *Vascular* Rays are not equally extended in all *Roots*: in *Parsnep*, towards the Circumference of the *Barque*; in *Bugloss*, about half way. In all *Docks*, and *Sorrels*, the *Rays* are extended through about $\frac{3}{4}$ of the thickness of the *Barque*, towards the Circumference, whereabout, divers of them are always *arched* in, two and two together. In all or many *Trefoyls*, and of the *Leguminous* Kind, they are extended through no more than $\frac{1}{3}$ of the *Barque*. In the *Umbelliferous*, they are Ralled in betwixt the Diametral *Portions* of the *Parenchymia*. In *Borage*, the *Rays* are more Continuous; in a *Carrot*, more Pricked. Here also the *Pricks* stand in Even Lines; in *Lovage*, they are Divaricated. Of which, and those of some other *Roots*, it is also Observable, That they are not all meier *Pricks*, but most of them small, yet real Circles; which, after the *Milk* hath been frequently licked off, and ceaseth to ascend, are visible, even without a *Glass*. And note, that in observing all *Milk-Vessels*, the *Milk* is to be taken off, not with the Finger but the Tongue; so often, till it riseth no more, or but little. And some *Roots* may also be soaked in Water; whereby the Position of the *Milk-Vessels*, will be visible by the darker Colour of the *Barque*, where they stand.

Tab. 7, 8, 9.

Tab. 8.

28. §. The *Rays* sometimes, run more Parallel, and keep several, as in *Monkshood*; and sometimes, towards the Circumference of the *Barque*, they are occurrent; as not only in *Docks*, but other *Plants*: In *Eryngo*, in a termination more Circular; and in *Bryony*, angular, or in the form of a *Glory*, as also in *Horradish*, through a *Microscope*. The Peripheral Lines are in some, more entire Circles, as in *Dandelion*; in others, made up of shorter Chords, as in *Potato*, *Cumfry*, and the smaller part of the *Root* of *Monkshood*. In some, the *Pricks* are so exceeding small, and stand so close, that, to the bare eye, they seem to be continous *Rings*, which yet, through the *Microscope*, appear distinct, as in *Marsh-mallow* and *Liquirish*.

Tab. 7, 8, 9.

Tab. 15.

Tab. 12.

29. §. Sometimes Columns and Chords are compounded, as in *Burnet*; *Pricks* and Chords, in *Potato*; *Rays* and *Rings*, in *Monkshood*; where the *Ring* is Single. In *Fenil*, there is a double or treple order both of *Rays* and *Rings*, the *Lymphæducts* standing in *Rays* and the *Lacteals* in *Rings*. And in *Marsh-mallow*, the *Vessels* are so posited as to make both those kinds of Lines at once.

30. §. In *Celandine*, they seem all, to the bare eye, to stand in numerous *Rings* lying even one within another. As also in *Dandelion*; in which yet, being viewed through a *Microscope*, there is an appearance of very many small *Rays*; which streaming from the Inner Verge of the *Barque*, cross three or four of the smaller *Rings*, and are there terminated. Whence it should seem that *Lymphatick* *Rays* and *Milky* *Rings*, are in that *Root*, so far mixed together. Only the *Lympha*, being confounded with the *Milk*, cannot be discerned. And where the *Milky-Vessels* are evacuated, or at such Seasons, wherein they are less full, divers *Milky* *Roots* will yield a clear *Liquor* at the Inner Verge of the *Barque*, where, at other times, they seem to yield only *Milk*. And this is the Description of the *Barque*.

Tab. 13.

C H A P. IV.

Of the WOOD.



THAT Portion of the *Root* which standeth next within the *Barque*, and in *Trees*, and *Shrubby Plants*, is the *Wood*; is also compounded of Two Substantially different Bodies, *Parenchymous* and *Lignous*. The *Parenchymous*, is of the same Substantial Nature with that of the *Barque*. And is originated from it; being not only adjacent to it, but all round about continuous therewith; even as that, is with the *Skin*; the *Parenchyma* of the *Barque*, being distributed, from time to time, partly outward into the *skin*, and partly inward, into the *Wood*.

Tab. 9.

Tab. 8, 9.

Tab. 17.

Tab. 17.

2. §. The Position of the several parts hereof, is different. For the most part it hath a Diametral Continuation, in several Portions, running betwixt as many more of the *Lignous*, from the Circumference towards the Center of the *Root*: all together, constituting that, which in the *second Chapter* of the *First Book*, I call the *Insertment*. In the *Roots* of many *Herbs*, these Diametral or Inserted Portions are more observable, as in *Cumfry*; which leadeth to the notice of them in all others, both of *Herbs* and *Trees*. Sometimes part of this *Parenchymous* Body is disposed into *Rings*, as in *Fenil*. The Number and Size of which *Rings* differ: In *Fenil*, when the *Root* is grown large, they are in some places broader, but fewer; in *Beet* they are narrower, but more. The Diametral Portions are here, in like manner, much varied; in *Cumfry*, *Celandine*, larger; in *Beet*, *Bugloss*, meaner; in *Borage*, *Parsnep*, more, and smaller; and in most *Woody-Roots*, streaming betwixt the *Pith* and the *Barque*, as so many small *Rays*. Their Continuation is also different; in some *Roots*, to the Centre, as in *Columbine*; in others not, as in *Parsnep*. And sometimes different in the same *Root*, as in the *Vine*.

3. §. The Contexture of these *Parenchymous* Portions is sometimes Uniform, as in *Bugloss*, *Peony*; and sometimes also, as it is in the *Barque*, different; in part, more sappy, and transparent; in part, more white, dry, and aery, as in *Carrot*, *Lovage*, *Scorzonera*, and others; which yet cannot be observed without a wary view. But their general Texture is the same being all made up of many small *Bladders*. Which are here of different Sizes, like those of the *Barque*, but for the most part smaller. Their Shape likewise, is usually Round; but sometimes Oblong and Oval, as in *Borage*; or Oblong and Square, as in the *Vine*.

4. §. The *Lignous Part*, if not always, yet usually, is also Compounded of Two Kinds of Bodies, *scil.* *Succiferous* or *Lignous* and *Aer-Vessels*. The *Lignous* as far as discernable, are of the same Conformation and Nature with those of the *Barque*, and in the transverse cut
of

of the *Root*, do oftentimes, as those, emit a *Liquour*. They are also Braced; and many of them run in distinct *Threads* or *Portions*, collaterally together.

5. §. The *Aer Vessels* I so call, because they contain no *Liquor*, but an *Aery Vapour*. They are, more or less, visible in all *Roots*. They may be distinguished, to the bare Eye, from the *Parenchymous Parts*, by their Whiter Surface; and their standing more prominent, whereas those shrink below the transverse level of the *Root*, upon drying. They are frequently Conjugated divers of them together, sometimes fewer, and *Tab. 10.* for the most part single, as in *Asparagus*; sometimes many, as in *Horf* & 15. *Radish*. And their Conjugations are also Braced, as the *Threads* of the *Succiferous Vessels*. But they are no where Inosculated: nor Twisted one about another; but only Tangent or Collateral. Neither are they Ramified, the greater into less; but are all distinctly continued, as the *Nerves* in *Animals*, from one end of the *Root* to the other.

6. §. Their Braces, as those of the *Succiferous Vessels*, are also of various number: in *Jerusalem Artichoke*, *Cumfry*, *Scorzoner*, more rare; in *Borage*, *Burnet*, more frequent; as by stripping off the *Barque* of such *Roots*, where it is easily separable, may be seen. And they often *Tab. 6.* vary in the same *Root*; so in *Borage*, *Scorzoner*, &c. they are more frequent in the Centre, and next the *Barque*, than in the Intermediate space, as by splitting those *Roots* down the middle doth appear. They also vary from those of the *Succiferous Vessels*; those being usually more *Tab. 6.* frequent, as in *Jerusalem Artichoke*, than these of the *Aerial*.

7. §. Betwixt these Braced *Aer-Vessels*, and the rest, which make the true *Wood*, run the *Parenchymous Parts* above described; as they *Tab. 6.* do betwixt the *Succiferous* in the *Barque*: and so make up two Pieces of *Net Work*, wherof one is the filling up of the other.

8. §. The Position of both these Kinds of *Vessels*, is Various. The *Succiferous* or *Lignous*, are sometimes posited in diametral lines or portions; as in the *Vine*, and most *Trees*. Sometimes, oppositely to the *Aerial*, as in *Beet*; each Ring herein being double, and made both of *Tab. 8.* *Sap* and *Aer-Vessels*.

9. §. In *Nettle* the Position is very peculiar, from what it is in the *Tab. 8.* *Roots* of other *Herbs*; being curiously mixed; the *Succiferous* running cross the *Aerial*, in several, *viz.* *Five*, *Six*, *Seven*, or more *Rings*. In *Bryony* the several Conjugations of the *Aerial*, are distinctly surrounded with the *Succiferous*. In *Patience*, the *Succiferous* are disposed, *Tab. 7.* besides *Rays*, into many small *Rings*, of different *Sizes*, sprinkled up and down, and not, as in other *Roots* having one common Centre; within divers whereof, the *Aer-Vessels* are included: especially within those which are drawn, not into *Rings*, but, as it were, into little stragling *Hedges*.

10. §. That also of the *Aer-Vessels*, is Various and Elegant: especially in the upper part of the *Root*. In *Ammi*, *Lilium-non-bulbosom*, they make a Ring. In these, a Prick'd-Ring; in *Peony*, a Ring of *Rays*; in *Valerian*, a Ring of Pricks and *Rays*. In others, they make *Tab. 7, 8, 9.* not *Rings*, but longer *Rays*, extended either towards the Centre, as in *Scorzener*; or meeting in it, as in *Columbine*. In the Common *Dock*, they stand more in single *Rays*: in the other *Species* of *Decks*, both in *Rays*, and collateral Conjugations between.

11. §. In *Beet*, they stand in several Rings; and every Ring, made of Rays. In *Cumfry*, the Rays and Rings are separate; those stand without, these next the Centre. In *Dandelion*, they stand altogether, and make a little Rope, in the Center it self. In *Geranium*, and others of that Kindred, they make a little Thred, in the same place. And in *Skirret*, they stand in two Threds, near the Centre.

12. §. In *Celandine*, they stand in almost parallel Lines. In *Monks-hood*, of a wedged Figure; divided in the smaller part of the *Root*, into Three little Wedges, with their poynts meeting exactly in the Centre. In *Cinquefoyle*, and *Strawberry*, they are also postur'd in three Conjugations, triangularly. In the young *Roots* of *Oak*, they stand neither in Radiated, nor otherwise strait, but Winding Lines. And in *Borage* the position, of many of them, is Spiral. As likewise, sometimes, in *Mercury*, or *Lapathum unctuosum*. In *Horseradish*, they stand more confused neither in Rings nor in Rays; yet their several Conjugations, are radiated: with very many other differences.

13. §. The Quantity of these *Vessels*, as to the space they take up in the *Root*, is to be computed Two ways, By their Number, and Size. Their Number may, in some *Roots*, and in some measure, be judged of, by the bare Eye; having, frequently, a whiter surface than the other *Parts*. As also their Size; the Bore of these *Vessels* being greater than that of the *Lignous* in all *Roots*; especially in some. For if you take the *Roots* of *Vine*, *Fenil*, *Dandelion*, *Plum-tree*, *Elder*, *Willow*, &c. and lay them by, for some time, to dry; and then, having cut off a very thin Slice of each, transversely; if you hold up those Slices before your Eye, so as the Light may be trajected through the said *Vessels*, they hereby become visible, as notably different, both in Number and Size.

14. §. But undeceitful and accurate Observation of both their Number, and Size, must be made by the *Microscope*; and so they will appear to be much more various. In *Bistort*, *Skirret*, they are very few; in *Beet*, very many: betwixt which extreams there are all Degrees; as in *Orpine*, *Venus Looking-Glass*, *Scorzonera*, *Great Celandine*, *Peony*, *Borage*, *Fenil*, &c. may be seen. So their Size, in some is extrem small, as in *Strawberry*, *Bistort*, *Valerian*; in others very great, as in *Asparagus*, *Bugloss*, *Vine*. They are also of several Sizes in one and the same Numerical *Root*; but in some, are less varied, as in *Lilium non bulbosum*, *Asparagus*, *Bugloss*; in others, more, as in *Bryony*, *Lovage*. Amongst all *Roots*, they vary by about Twenty Degrees; as by comparing the *Roots* of *Vine*, *Thorn-Apple*, *Bryony*, *Lovage*, *Fenil*, *Wild Carrot*, *Saxifrage*, *Parsley*, *Peony*, *Hore-bound*, *Cinquefoyl*, *Strawberry*, &c. together, may be seen. Some of those in the *Vine*, being of the greatest Size; appearing through a good *Glass*, at least one Third of an Inch in Diametre: those in *Strawberry*, and that Kind, of the smallest; most of them appearing, in the same *Glass*, no bigger, than to admit the poynt of a small Pin, according to the Standard, in *Tab. 12*. See also the Figures of so many of them as are drawn.

15. §. In some *Roots*, they are Small, and Few; as in *Jerusalem Artichoke*; in others Small, but Many, as in *Horse-Radish*: in *Bugloss*, *Tab. 11, 14*, they are Great, but Few; in the *Vine*, Great and Many. So that the proportion, which those of a *Vine*, their Number and Size being taken together, bear to those of *Jerusalem Artichoke*, may be, at least, as Fifty, to One. Of the smallest Kinds, as those of *Cinquefoyl*, *Jerusalem Artichoke*, and the like; It is to be noted, That they are scarce ever visible in the fresh Slices of these *Roots*; but after they have layn by a while, at last, by a good *Glass*, Clear Light, and stedy View, are discernable.

16. §. In some *Roots*, the greater of these *Vessels* stand in or next the Centre, as in *Taraxacum*, or *Dandelion*; in others next the circumference, as in *Horse-Radish*. Sometimes each of them is from one end of the *Root* to the other, of a more equal Size, or more Cylindrical, as in *Marsh-mallow*; but usually, they widen, more or less, from the Top, to the Bottom of the *Root*, as in *Thorn-Apple*: about the Top of which, they are, for the most part, but of the Sixth, Seventh, and Eighth, Magnitude; some of the Fifth, but none of the Third; but about the Bottom, they are most of the Third, and Fifth: whence it is manifest, That some of them are, in the manner of *Veins*, somewhat Pyramidal. Yet is it observable, That their ampliation proceedeth not towards, but from their Original, as in *Nerves*. *Tab. 13, 15.*

17. §. Of these *Vessels* Seignior *Malpighi* hath observed; *Componuntur (saith he) exposita fistula Zona tenui & pellucida, velut argentei coloris lamina, parum lata; quæ, spiraliter locata, & extremis lateribus unita, Tubum, interius & exterius aliquantulum asperum, efficit.*

18. §. To whose Observation I further add, That the *Spiral Zone*, or *Lamina*, as he calls it, is not ever one Single *Piece*; but consisteth of Two or More round and true *Fibres*, although standing collaterally together, yet perfectly distinct. Neither are these Single *Fibres* themselves flat, like a *Zone*; but of a round forme, like a most fine *Thred*. According as fewer or more of these *Fibres* happen to break off, from their *Spiral* location, together; the *Zone* is narrower, or broader: usually, Narrower in the *Trunk*, and Broader in the *Root*.

19. §. Of these *Fibres* I also Observe, That they are not *Inosculated* side to side, but are *Knit* together by other smaller *Fibres*; those being, as it were, the *Warp*, and these the *Woof* of the *Aer-Vessels*. Yet I think the several *Fibres* are not interwoven just as in a *Web*; but by a kind of *Stitch*, as the several *Plates* or *Bredths* of a *Floor-Mat*. A clear and elegant sight of these *Fibres*, and of their *Interweftage*, by splitting a *Vine-Root*, or a piece of *Oak*, may, with a good *Glass* in the sides of their Greater *Aer-Vessels*, be obtained; having much of the resemblance of *Close Needle-work*.

20. §. The *Spiration* of the *Fibres* of these *Vessels*, may more easily be observed in the *Trunk*, than in the *Root*. And better in younger *Plants*, than other. And not so well by *Cutting* as by *Splitting*, or by *Tearing off* some small *Piece*, through which they run: their *Conformation* being, by this means, not spoiled. Yet this way, the *Vessels* are seen, chiefly, *Unresolved*.

21. §. But in the *Leaves* and *Tender Stalks* of all such *Plants*, as shew, upon breaking, a kind of *Downe* or *Wool*; they may be seen *Resolved* and *Drawn out*, and that sometimes even to the naked Eye,

an Inch or two Inches in length. This *Wool* being nothing else, but a certain number of *Fibres* Resolved from their Spiral position in these *Vessels*, and Drawn out in Length; and so clustred together, as so many *Threds* or little *Ropes*: appearing thus more or less, in the *Leaves* and some other *Parts* of most *Plants*; but more remarkably in some, as in the *Vine*, *Scabious*, and others. As also in the Scales of a *Squill*. In which last, for example, they are so easily separable, as further to shew, what before was observed; viz. That the *Plate* or *Zone*, into which the *Aer-Vessels* are usually Resolved, is not one Single Piece, or meer *Plate*; but made up of several Round *Fibres*, all standing and running parallel, and so knit together by other smaller ones, transversely, in the form of a *Zone*. For if you break or cut a *Leaf* or *Shell* of a fresh *Squill*, till you come to the *Aer-Vessels*, and having softly drawn them out, for about an Inch or more (to the naked Eye) in length, you then single out one or two of them from the rest, and rowl them, as they hang at the *Shell*, eight or nine times round, each *Vessel* will appear, through a *Glass*, to consist of 8, 10, or 12 small *Fibres*; which, in the Unresolved *Vessel*, run parallel; but by this means, are all separated one from another. See the *Figures* belonging to the *Third* and *Fourth Books*.

22. §. The Process of their Spiration, is not, so far as I have observed, accidental, but constantly the same; *scil.* In the *Root*, by *South*, from *West* to *East*: But in the *Trunk*, contrarily, by *South*, from *East* to *West*.

23. §. The *Content* of these *Vessels*, is, as hath already been intimated, more *Aery*. The Arguments for which, are, That upon a transverse Cut of the *Root*, the *Sap* ascendeth not there, where These stand. Being also viewed through a *Microscope*, they are never observed to be filled with *Liquor*. Besides a *Root* cut and immersed in Water, till the Water is in some part got into these *Vessels*, and then the *Root* taken out and crushed; the other *Parts* will yield *Liquor*, but These, only *Bubbles*: which *Bubbles* are made, by some small quantity of *Liquor* mixed with the *Aer*, before contained in the said *Vessels*. To which, other Arguments will arise out of those Things that follow in the *second Part*. As also for this *Content*, its not being a pure or simple, but *Vaporous Aer*. Whether these *Vessels* may not, in some *Vegetables*, and at some times, contain *Liquor*, is doubtful. (a) Thus far of the *Lignous Part*.

(a) See
Book 3.

C H A P. V.

Of the P I T H.



WITHIN the *Lignous Part* lyeth the *Pith*. This *Part* is not common to all *Roots*, for some have none, as *Nicotian*, *Srtamonium*, and others. Yet many which have none, or but little, throughout all their lower parts, have one fair enough about their tops, as *Mallow*, *Bourage*, *Dandelion*, and the like. See the *Roots*. And in many others there are *Parenchymous Parts*, of the same substantial nature with the *Pith*, distributed betwixt the several Rings of *Vessels*, and every where visible, from the top to the bottom, as in *Beet*, *Fenil*, &c. Tab. 6.

2. §. The Size of the *Pith* is varied by many Degrees, easily reckoned an Hundred; in *Fenil*, *Dandelion*, *Asparagus*, but small; in *Horfe-Radish*, *Valerian*, *Bistort*, great. The Shape hereof, in the lower parts of most *Roots*, is *Pyramidal*; but at the tops, *Various*, according to the different Distribution of the *Vessels*, as in *Carrot*, *Hyperbolick*, in *Parsley*, *Oval*; as appeareth, in cutting the *Roots* lengthways. Tab. 8.

3. §. The *Pith*, for the most part, especially in *Trees*, is a *simple Body*; but sometimes, it is, as the *Barque*, compounded; some certain number of *Succiferous Vessels* being mixed herewith; as in *Jerusalem Artichoke*, *Horfe-Radish*, &c. upon a traverse cut, by a strict view, may be discerned. Their Position is sometimes *Confused*, as in a *Carrot*; and sometimes *Regular*, as in *Parsley*; appearing, by the traverse cut, in Rings, and in cutting by the length, in *Arches*. And sometimes the *Pith* is hollow; as in the *Level-Roots* of *Bishops-Weed*: these *Roots* being made out of the *Stalk*, as in the *First Chapter* hath been shewed, how. Tab. 6, & 8.

4. §. As all the other Parts of the *Root*, are originated from the *seed*; so, sometimes, is the *Pith* it self. But sometimes, it hath its more immediate Derivation from the *Barque*. Hence it is, that many *Roots*, which have no *Pith* in their lower parts, have one at their top, as *Columbine*, *Lovage*, &c. For the *Parenchymous Parts* of the *Barque* being, by degrees, distributed into *Diametral Portions*, running betwixt those of the *Lignous Body*, and at length, meeting and uniting in the Centre, they thus constitute the *Pith*. In the same manner, at the top of some *Roots*, the *Pith* is either made or augmented, out of the *Parenchymous Rings* above described; these being gradually distributed to, and embodied in the Centre; as in *Fenil*, and some other *Roots*, their lower and upper parts compared together, may be seen. Even as in *Animals*, one Part, as the *Dura Mater*, is the original of divers others. Tab. 6.

5. §. From hence, it also appears, That the *Pith* is of the same Substantial Nature with the *Parenchyma* of the *Barque*, and with the *Diametral Portions*; and that therefore they are all one body, differing in no Essential Property, but only in their Shape and Place. The same is also evident from the Continuity of the *Pith* with the *Diametral Portions*, as of These, with the said *Parenchyma*. And from their Contexture, which, by a *Microscope*, appeareth to be of one and the same general kind, in all *Plants*, both in the *Parenchyma* of the *Barque*, in the *Insertment* or *Diametral Portions*, and in the *Pith*, all being made up of *Bladders*.

Tab. 9, & 15. 6. §. The *Bladders* of the *Pith*, are of very different Sizes; seldom less, than in the *Barque*, as in *Asparagus*; usually much bigger, as in *Horse-Radish*. They may be well reckoned to about fifteen or twenty degrees; those in *Jerusalem Artichoke*, of the largest; in *Valerian*, *Horse-Radish*, of the meaner; in *Bistort*, *Peony*, of the smallest. Their Position is rarely varied, as it is oftentimes, in the *Barque*; but more uniform, and in the transverse Cut, equally respective to all parts of the *Root*: yet being piled evenly, one over another, in the long cut, they seem to run, in Direct Trains, by the length of the *Root*. Their Shape also is, usually more orbicular; but sometimes, somewhat angular, in the larger kinds, as in *Jerusalem Artichoke*.

Tab. II.

7. §. THUS FAR the Contexture of the *Pith* is well discoverable in the *Root*. In the *Trunk*, farther, and more easily. Whereof therefore, in the next *Book*, I shall give a more particular *Description* and *Draught*. Yet since I am speaking of it, I shall not wholly omit here to observe, That the *sides*, by which the aforesaid *Bladders* of the *Pith* are circumscribed, are not meer *Paper-Skins*, or rude *Membranes*; but so many several Ranks or Piles of exceeding small *Fibrous Threads*; lying, for the most part, evenly one over another, from the bottom to the top of every *Bladder*; and running cross, as the *Threads* in the *Weavers Warp*, from one *Bladder* to another. Which is to say, That the *Pith* is nothing else but a *Rete mirabile*, or an Infinite Number of *Fibres* exquisitely small, and admirably Complicated together: as by cutting the *Pith* with a *Razor*, and so viewing it with a good *Glass*, may be seen. See the *Figures* belonging to the *Third Book*.

8. §. All *Plants* exhibit this Spectable, not alike distinctly; those best, with the largest *Bladders*. Nor the same *Pith*, in any condition; but best, when dry: Because then, the *Sap* being voided, the spaces betwixt the *Fibrous Threads*, and so the *Threads* themselves, are more distinctly discernable. Yet is it not to be dried, after Cutting; Because its several parts, will thereupon coincide and become deformed. But to be chosen, while the *Plant* is yet growing; at which time, it may be often found dry, yet undeformed; as in the *Trunks* of *Common Thistle*, *Jerusalem Artichoke*, &c.

9. §. Neither are these *Threads*, so far as I can observe, Single *Fibres*; but usually, consist of several together. Nor are they simply Collateral, but by the westage of other *Fibres*, in their natural Estate, knit together; much after the same manner as the *Spiral Fibres* of the
Aer-

Aer-Vessels. This Connexion I have no where so well seen, as in the White *Bottoms* of the *Bladders* of a *Bulrush*, being cut traverse; wherein they have the appearance, of very Fine and close *Needle-work*.

10. §. The *Fibres* by which the said *Threds* are knit together, I think are all Single: and are seldom and scarcely visible, except by obliquely Tearing the *Pith*; by which means, they will appear through the *Glass*, broken off, sometimes, a quarter or half an Inch, or an Inch in Length; and as small as one Single *Thred* of a *Spiders Webb*. In a *Bulrush*, they are sometimes discernable in cutting by the Length. These *Fibres*, and the *Threds*, they knit together, for the most part, are so pellucid, and closely situate, that they frequently seem to make One entire Body, as a picce of *Ice* or a film of *Water* it self: or even as *Animal Skins* sometimes shew, which yet are known to be *Fibrous*.

11. §. The Situation of these *Threds*, is contrary to that of the *Vessels*, as those by the Length, to these, chiefly, by the Breadth of the *Root*, or horizontally, from one edge of the *Pith* to the other. They are continued circularly; whereby, as oft as they keep within the compass of the several *Bladders*, the said *Bladders* are Round: But where they winde out of one *Bladder*, into another, they mutually Intersect a *Chord* of their several *Circles*; by which means, the *Bladders* become Angular.

12. §. The Contexture, likewise, both of the *Parenchymous Part* of the *Barque*, and of the *Diametral Portions* inserted betwixt the *Lignous*; is the same with this of the *Pith*, now described; that is, *Fibrous*. Whence we understand, How the several *Braces* and *Threds* of the *Vessels* are made: For the *Vessels* running by the length of the *Root*, as the *Warp*; by the *Parenchymous Fibres* running cross or horizontally, as the *Woof*: they are thus knit and as it were stitched up together. Yet their *westage* seemeth not to be simple, as in *Cloath*; but that many of the *Parenchymous Fibres* are wrapped round about each *Vessel*; and, in the same manner, are continued from one *Vessel* to another; thereby knitting them altogether, more closely, into one *Tubulary Thred*; and those *Threds*, again, into one *Brace*: much after the manner of the *Needle work* called *Back-Stitch* or that used in *Quilting* of *Balls*. Some obscure sight hereof, may be taken in a *Thred* of *Cambrick*, through a *Microscope*. But it is most visible, in the *Leaves* and *Flowers* of some *Plants*. The *Delineation* of these Things I shall therefore omit, till we come hereafter to speak of the other *Parts*.

13. §. From what hath been said, it may be conjectured; That the *Aer Vessels* successively appearing in the *Barque*, are formed, not out of any *Fluid Matter*, as are the original ones: But of the *Parenchymous Fibres*; sc. by changing them from a *Spherical* to a *Tubulary Forme*.

14. §. From the precedents, it is also manifest, That all the *Parenchymous Parts* of a *Root*, are *Fibrous*.

15. §. And lastly, That the whole Body of a *Root*, consisteth of *Vessels* and *Fibres*. And, That these *Fibres* themselves, are *Tubulous*,
or,

or so many more *Vessels*, is most probable: There only wanteth a greater perfection of *Microscopes* to determine.

16. §. The *Contents* of the *Pith* are, sometimes *Liquor*, and sometimes a *Vaporous-Aer*. The *Liquor* is always *Diaphanous*, as that of the *Parenchymous Part* of the *Barque*; and in nature, not much differing from it. The *Aer* is sometimes less, and sometimes more *Vaporous*, than that of the *Barque*. By this *Aer* I mean, that which is contained in the *Bladders*. Within the *Concaves* of the *Fibres* which compose the *Bladders*, I suppose, there is another different Sort of *Aer*. So that as in the *Bladders* is contained a more *Aqueous*; and in the *Vessels*, a more *Essential Liquor*: So sometimes, in the same *Bladders*, is contained a more *Vaporous*; and in the *Fibres*, a more *Simple* and *Essential Aer*.

An Account of the

VEGETATION

OF

ROOTS

Grounded chiefly upon the foregoing

ANATOMY.

PART II.



TO *Philosophize*; is, To render the *Causes* and *Ends* of Things. No man, therefore; that denieth *God* can do this, Truly. For the taking away of the *first Cause*, maketh all things *Contingent*. Now, of that which is *Contingent*, although there may be an *Event*; yet there can be no *Reason* or *End*: so that Men should then study, That, which is *not*. So the *Causes* of Things; if they are *Contingent*, they cannot be *Constant*. For that which is the *Cause* of This, now; if it be so *Contingently*, it may not be the *Cause* hereafter: and no *Physical* Proposition, grounded upon the *Constancy* and *Certainty* of Things, could have any foundation. He, therefore, that philosophiseth, and denieth *God*, playeth a childish Game.

2. §. Wherefore *Nature*, and the *Causes* and *Reasons* of Things, duly contemplated, naturally lead us unto *God*; and is one way of securing our Veneration of Him: giving us, not only a general Demonstration of his *Being*; but a particular one, of most of the several *Qualifications* thereof. For all *Goodness*, *Righteousness*, *Proportion*, *Order*, *Truth*, or whatever else is Excellent and Amiable in the *Creatures*; it is the Demonstration of the like in *God*. For it is impossible, that *God* should

Theology the
Beginning
and End of
Philosophy.

should ever make any thing, not like Himself, in some degree or other. These Things, and the very Notions which we have of them, are *Conceptions* issuing from the *Womb* of the *Divine Nature*.

3. §. By the same means, we have a greater assurance of the Excellency of his *Sacred Word*. That He, who hath *Done* all things so transcendently well; must needs *Speak* as well, as he hath *Done*. That He, who in so admirable a manner, hath *made* Man; cannot but know best, What his true *Principles* and *Faculties* are; and what *Actions* are most agreeable thereunto: and, that having adorned him with such *Beauteous* and *Lovely* ones; it is impossible, He should ever put him upon the Exercise of those *Faculties*, in any way *Deformed* and *Unlovely*. That He should do all things, so well *Himself*; and yet require his *Creatures*, to do otherwise, is unconceivable.

4. §. And as we may come, hereby, to rectifie our Apprehension of His *Laws*; so also, of His *Misteries*. For there are many Things, of the *Manner* of whose Existence, we have no certain Knowledge. Yet, of their *Existence*, we are as sure, as our *Senses* can make us. But, we may as well deny, what *God* hath Made, *To be*; as, what he hath Spoken, *To be true*, because we understand not *how*. And the knowledge of *Things* being gradually attained, we have occasion to reflect, That some *Things*, we can now well conceive, which we once thought unintelligible. I know, therefore, what I *understand* not; but, I know not, what is *unintelligible*: what I know not now, I may hereafter; or if not I, another; or if *no Man*, or other *Creature*, it is sufficient, *That God fully understandeth Himself*. It is not, therefore, the *Knowledge* of *Nature*, but they are the *wanton phantasies* of Mens minds, that dispose them, either to Forget *God*, or to Think unduly of Him.

5. §. Nor have we reason to fear going *too far*, in the Study of *Nature*; more, than the *entring* into it: Because, the higher we rise in the true Knowledg and due Contemplation of *This*; the nearer we come to the *Divine Author* hereof. Or to think, that there is any Contradiction, when *Philosophy* teaches that to be done by *Nature*; which *Religion*, and the *Sacred Scriptures*, teach us to be done by *God*: no more, than to say, That the *Ballance* of a *Watch* is moved by the next *Wheel*; is to deny that *Wheel*, and the rest, to be moved by the *Spring*; and that both the *Spring*, and all the other *Parts*, are caused to move together by the *Maker* of them. So *God* may be truly the *Cause* of *This Effect*, although a Thousand other *Causes* should be supposed to intervene: For all *Nature* is as one Great *Engine*, made by, and held in His Hand. And as it is the *Watch-makers Art*, that the *Hand* moves regularly, from hour to hour, although he put not his Finger still to it: So is it the Demonstration of *Divine Wisdome*, that the *Parts* of *Nature* are so harmoniously contrived and set together; as to conspire to all kind of Natural Motions and Effects, without the Extraordinary and Immediate Influence of the *Author* of it.

6. §. Therefore, as the *Original Being* of all Things, is the most proper Demonstration of *Gods Power*: So the *successive Generations*, and *Operations* of Things are the most proper Demonstration of his *Wisdom*. For if we should suppose, that *God* did now make, or do any Thing, by any Thing; then, no *Effect* would be produced by a *Natural Cause*: and consequently, He would still be upon the Work of *Creation*: which yet *Sacred Scripture* assureth us, He *resteth from*. And we might expect
the

the Formation of a *Child*, in an *Egg*, as well as in a *Womb*; or of a *Chicken*, out of a *Stone*, as an *Egg*: And all Sorts of *Animals*, as well as *Plants*, might propagate their *Species*, without Coition: and the like. For *Infinite Power*, needeth not make any difference in the Things it undertakes to manage. But in that, these Things are not only *made*, but *so made*, that is, according to such certain Natural Laws, as to produce their *Natural Effects*; here is the Sensible and Illustrious Evidence of his *Wisdom*. Wherefore as the Wisdom of Government, is not seen, by the King his interposing Himself in every Case; but in the contrivance of the *Laws*, and Constitution of *Ministers* in such sort, that it shall be as effectually determin'd, as if he did so indeed: So the more complicated and vastly Numerous, we allow the *Natural Causes* of Things to be; the more duely we conceive of that *Wisdom*, which thus disposeth of them all, to their several *Effects*: All Things being thus, as *Ministers* in the Hands of *God*, conspiring together a Thousand *Ways*, towards a Thousand *Effects* and *Ends*, at one time; and that with the same certainty, as if he did prepose to each, the same Omnipotent *Fiat*, which he used at the Creation of the World.

7. §. THIS *Universal Monarchy*, as it is eminently Visible in all other Particular *Oeconomies*; so is it, no less, in that of *Vegetables*. Infinite Occurrences, and secret Intrigues, 'tis made up of; of which we cannot skill, but by the help of manifold *Means*; and those, in the foregoing *Idea*, have been lately propos'd. Wherein, although some *Experiments* have been briefly touch'd: yet that which I have hitherto chiefly prosecuted, hath been the *Anatomical Part*; and that not thoroughly neither. Notwithstanding, so far as Observations already made will conduct us, I shall endeavour to go. And if, for the better clearing of the way, I have intermixed some Conjectures; I think they are not merely such, but for which I have layd down some Grounds, and of which, the *Series* also of the following *Discourse*, may be some further proof.

The Divine
Wisdom
seen in the
Growth of
Plants; if
we observe,

8. §. LET US say then, that the *Root* of a *Plant* being lodged in some Soil, for its more convenient growth; 'tis necessary the Soil should be duly prepared for it. The *Rain*, therefore, falling and soaking into the Soyl, somewhat diluteth the Dissoluble *Principles* therein contained; and renders them more easily communicable to the *Root*: Being as a *Menstruum*, which extracteth those *Principles*, from the other greater and useles part of the Soil.

First,
How the
Ground is
prepared:

9. §. And the warm *Sun*, joyned with the diluting *Rain*, by both, as it were a *Digestion* of the Soil, or a gentle *Fermentation* amongst its several Parts, will follow: whereby the Dissoluble Parts therein, will rot and mellow: that is, those *Principles* which as yet remained more fixed, will now be further resolved and unlocked, and more copiously and equally spread themselves through the Body of the Soil.

10. §. These *Principles*, being with the growth of *Plants* continually exhausted, and needing a repair; the successions, therefore, of Wet, Wind, and other Weather, beat down and rot the *Leaves* and other *Parts* of *Plants*. Whereby these (as *Weeds* which are wont to be buried under ground) become a *natural Manure*, and Re-impreg-

nate the Soil : Being thus, in part, out of their own Resolved Principles, annually Compounded again.

11. §. Many of these Principles, upon their Resolution, being by the Sun more attenuated and volatilized; continually ascend into the Aer, and are mixed therewith. Where, although they lose not their Vegetable Nature, yet being amongst other purer Principles; themselves also, depositing their Earthy feculencies, become more subtile, simple and Essential Bodies.

12. §. And the Aer being of an Elastic or Springy Nature, pressing, more or less, upon all Bodies; it thereby forceth and insinuateth it self into the Soil, through all its permeable Pores. Upon its own entrance, it carries also many of the said Vegetable and Essential Principles along with it; which, together with the rest, are spread all over the Body of the Soil. By which means, though a less Vehement, yet more Subtil Fermentation, and with the least advantage of warmth, continuable, will be effected.

13. §. The Principles being thus farther resolved and subtilized, would presently exhale away, if the Rain, again, did not prevent. Which, therefore, falling upon and soaking through the Ground, is as a fresh Menstruum, saturate or impregnate with many of them. And as it still sinketh lower, it carries them along with it self, from the Superficial, to the Deeper parts of the Ground: thus, not only maturing those parts also, which, otherwise, would be more lean and cold; but therein likewise, laying up and securing a Store, more gradually and thriftily to be bestowed upon the Upper parts again, as they need.

14. §. And Autumn having laid up the Store, Winter following thereupon, doth, as it were, lock the doors upon it. In which time, some warmer Intervals, serve further and gradually to mature the stored Principles, without hazard of their being Exhaled. And the Spring returning, sets the doors open again, with warmer and more constant Sun, with gentle and frequent Rain, fully resolves the said Principles; and so furnisheth a plentiful Diet, for all kinds of Vegetables: being a Composition of Water chiefly, wherein are resolved, some portions of Earth, Salt, Acid, Oyl, spirit, and Aer; or other Bodies of Affinity herewith.

Then, How the Sap is imbib'd, and distributed to the several Parts. (a) P. I. c. 3. §. 3. (b) §. 11, 12

15. §. THE ROOT standing in the Ground thus prepared, and being always surrounded with a Barque, which consisteth chiefly of a Parenchymous and Spongy Body; (a) it will thus, as Sponges do, naturally suck up the watry parts of the Soil impregnate with the said Principles. Which Principles notwithstanding, being in proportion with the watry parts, but few, and also more Essential; (b) therefore in this Parenchymous Part, are they never much discovered, either by Colour, Taste, or Smell. As it is probable, that some distilled Waters, which discover nothing, to Sense, of the Plants from which they are distilled, may yet, in part, retain their Faculties. And it is known, that many Bodies; as Crocus Metallorum, convey many of their parts into the Menstruum, without any sensible alteration thereof. So Frost and Snow have neither Taste nor Smell; yet from their Figures, 'tis evident, that there are divers kinds of Saline Principles incorporated with them; or at least, such Principles as are common to them and divers kinds of Salts.

16. §.

16. §. The entrance of this Impregnate Water or Sap is not without difference, but by the Regulation of the intervening *Skin*; being thereby *strained* and rendred more pure: the *Skin*, according to the thickness (*a*) or closeness thereof, becoming sometimes only as a *brown paper*, sometimes as a *Cotton*, and sometimes as a *Bag of Leather* to the transient *Sap*, as the nature of it doth require. By which it is also *moderated*, lest the *Barque*, being spongy, should suck it up too fast, and so the *Root* should be, as it were, surcharged by a *Plethora*. And divers of the *Succiferous Vessels* being mixed herewith (*b*) and lying next the *Soil*, usually more or less *mortified*, and so their *Principles* somewhat *resolved*; the *Sap* is hereby better *specified*, and further *tinctured*; such parts of the *Sap* best entring, as are most agreeable to those *Principles*; which the *Sap* also carries off, in some part, as it passeth into the *Barque*.

17. §. The *Sap* thus *strained*, though it be *pure*, and consisteth of *Essential* parts; yet being *compounded* of *heterogeneous* ones; and received into the *Parenchyma* of the *Barque* a laxe and spongy Body, they will now easily and mildly *ferment*. Whereby they will be yet further prepared, and so more easily insinuate themselves into all the *Bladders* of the said *Parenchyma*; swelling and dilating it as far as the *Continuity* of its parts will bear. Whereupon, partly from the continued entrance of fresh *sap*, and partly by a Motion or Pressure of *Restitution* in the swollen and Tensed *Bladders* of the *Parenchyma*, the *Sap* is forced thence into the other parts of the *Root*.

18. §. And because the *Parenchyma* is in no place openly and Visibly *Pervious*, but is every where composed of an Infinite Number of small *Bladders* (*c*); the *Sap*, therefore, is not only *fermented* therein; and fitted for Separation; but, as it passeth through it, is every part of it, *strained* an Hundred times over, from *Bladder* to *Bladder*.

19. §. The *Sap* thus *fermented*, and *strained*, is distributed to the other *Organical Parts*, according as the several *Principles* of This, are agreeable to those whereof the said *Organical Parts* consist. As the *Sap* therefore passeth from *Bladder* to *Bladder*, such *Principles* as are agreeable to those of the *Fibres* of the said *Bladders*, will adhere to, and insinuate themselves into the Body of the *Fibres*; *sc.* *Watry* chiefly, next *Acid*, then *spirituous*, *Earthy*, *Aery*, and *Oleous*. (*d*)

20. §. And the *sap* by its continual appulse and *percolation*, as it leaveth some parts upon the said *Fibres*; so as it is *squeezed* betwixt them from *Bladder* to *Bladder*, it licks and carries off some others from them, in some *union* together with it; and so is *Impregnate* herewith: as *Water*, by passing through a *Mineral Vein*, becomes *tinctured* with that *Mineral*.

21. §. The *Sap* thus *Impregnate* with some *united Principles* of the *Parenchymous Fibres*, passeth on to the *Lignous Vessels*, whereinto their correspondent *Principles* also enter; *sc.* *Watry*, *Saline*, *Oleous* and *Earthy* chiefly. (*e*) And because the *Parenchymous Principles* mixed with them, are in some degree *united*, and so more ready to *fix*; some of these therefore will likewise enter into the said *Vessels*. Whereupon, the *Alkali oleosum* of the one, and the *Acidum spirituosum* of the other, meeting together; These, with the other *Principles*, all *concentre*, and of divers *fluids*, become one *fixed Body*, and are gradually *agglutinated* to the *Vessels*; that is, The *Vessels* are now *nourished*.

22. §. The supply of the *Sap* still continued, the *Principles* thereof will not only enter into the *Body* of these *Parts*, but also their *Concaves*.
 (a) P. I. c. 5. §. 12. And the *Parenchymous Fibres* being wrapped about the *Vessels*, (a) as often as the said *Fibres* are more *turgid* with their own contained *Fluid*, they will thereby be somewhat *shortned*, or contract in length; and so must needs *bind* upon the *Vessels*, and thereby, as it were, *squeeze* some part of the *Fluid*, contained both within themselves and the *Vessels*, back again into the *Bladders*.

23. §. And the *Sap* herein, being thus *tinctured* with some of the *united Principles* of the *Vessels*, divers of them will now also insinuate themselves into the *Parenchymous Fibres*, and be incorporated with them: Whereby, the said *Fibres*, which before were only *relaxed* and *dilated*, are now also *nourished*, and not till now. Some portion of the *united Principles* both of the *Parenchymous* and *Lignous Parts*, being necessary to the true *nutrition* of Each: As the *Confusion* and *joynt assistance* of both the *Arterious* and *Nervous Fluids*, is to the *nourishment* or *coagulation* of the *Parts* in *Animals*.

24. §. Some portion of the *Sap* thus doubly *tinctured*, is at the same time transmitted to, and enters the *Body* of the *Aer-Vessels*; consisting chiefly of *Water*, *Aer*, and *Acid*; and, in like manner, as in the other *Parts* is herein *agglutinated*. And the *appulse* and *pressure* of the *Sap* still continued, some portion hereof is also trajected into the *Concaves* of the said *Vessels*; existing therein as a most *Compounded Fluid*; partaking, more or less, both of the *Principles* and *Tinctures* of the other *Organical Parts*, and of the *Aer-Vessels* themselves; being as it were, a *Mixed Resolution* from them all.

25. §. And the *Parenchymous Fibres* being wrapped about These, as about the other *Vessels*, (b) and, in like manner, *binding* upon them; they thus frequently *squeeze* part of the said contained *Fluid* out again: As necessary, though not to the immediate *Nourishment* of the *Parts*, yet the due *Qualification* of the *Sap*; being a *Constant Aerial Ferment*, successively stored up within the *Aer-Vessels*, and thence transfused to the *Sap*, in the other *Organical Parts*.
 (b) P. I. c. 5. §. 12.

26. §. And that there may be a better *Transition* of the *Sap* thus *tinctured*, to the several *Organical Parts*; therefore, none of them are close set and compact within themselves, severally: For so, they would be inaccessible to the *Sap*, and their inward *Portions*, wanting a due supply of *Aliment*, would be starved. But the *Vessels*, both of *Aer* and *Sap*, being every where divided into *Braced Portions*, and other *Parenchymous Portions*, filling up the spaces every where betwixt them (c); there is therefore a free and copious communication of the *Sap*, (and so of all the *Tinctures* successively transfused into it) from *Part* to *Part*, and to every *Portion* of every *Part*: The *Parenchymous Portions*, running betwixt the *Braces*, as the smaller *Vessels* do throughout the *Viscera*, in *Animals*. Whereby, none of them want that *Matter*, which is necessary either for their *Nutrition*, or for the good *Estate* of their *Contents*, or for the due period of their *Growth*.
 (c) P. I. c. 3. §. 13. & c. 4. §. 4, 5, 7.

27. §. For the better *Tempering* of the several parts of the *Sap*, serve the *Diametral Portions* of the *Parenchymous Body* which run sometimes directly through the *Barque*, as in *Lovage*, *Parsley*, &c. is described and figur'd (d) Which being, all or most of them, continued be twixt both the *Succiferous* and the *Aer-Vessels*, from the *Circumference*
 (d) P. I. c. 3. §. 7, 8.

to the Centre; they hereby carry off a more Copious and *Aerial Ferment* from the One, and communicate it unto the Other. For as the *Sap* enters the *Barque*, the more *liquid* part, still passeth into the *succulent* Portions thereof; the more *Aery*, is separated into those *White* and *Dryer Diametral* ones; and in its passage betwixt the *Portions* of the *Aer-Vessels*, is all along communicated to them. Yet is it not a pure or *simple Aer*, but such as carries a *Tincture* with it, from the *Succiferous Vessels*. And therefore it is observable, That when the *Diametral* Portions are more distant, the *Sap-Vessels* run not in a *Straight Line* betwixt them, but are *Reciprocally* so inclined, as to touch upon them; as in *Lovage* is visible: Thereby communicating their *Tincture* to the *Aer*, as it passeth by them, through the said *Diametral* Portions.

28. §. By the continual appulse of fresh *Sap*, some, both of the *aery*, and of all the other parts thereof are transmitted into the *Pith*; where, finding more room, it will yet more kindly be *digested*. Especially having the advantage herein of some degree of *Warmth*; being herein remoter from the *soil*, and, as it were, *Tunn'd* up within the *Wood*, or the *Mass* of surrounding *Vessels*. So that the *Pith*, is a *Repository* of better *Aliment* gradually supplied to those *Succiferous Vessels*, which are frequently scattered up and down therein, and which ascend into the *Trunk*. (a) But where no *succiferous Vessels* are mixed, herewith, it usually becomes *Dryer*, and is replenished with a more *Aerial* and *Warmer Sap*; whereby the growth of the *Caulis* is promoted, as by an *Hot Bed* set just under it. And in many *Plants* with divers knobbed *Roots*, the younger are more *succulent*, serving chiefly to feed the *Stalk*: the *Elder* are spongy and fill'd with *Aer*, for the fermenting of the *Sap*, and more early growth of the *Stalk*: as in little *Celandine*, *Dogstones* and all of that Kindred. And thus all the *Parts* have a fit *Aliment* provided for their *Nourishment*

(a) P. I. c. §.
§. 3.

29. §. IN THIS *Nourishment*, the *Principles* of the *Sap* are, as is said, *concentred* and *locked* up one within another: (b) Whence it is, that the *Organical Parts*, being cleansed of their *Contents*, have none of them any *Taste* or *Smell*, as in the *Piths* of *Plants*, *Paper* and *Linen Cloth* is evident. (c) Because till by *Digestion*, violent *Destillation*, or some other way, they are resolved, they cannot act upon the *Organs* of those *Senses*. For the same reason, they are never *tinctured*, excepting by their *Contents*: and although, to the bare *Eye*, they frequently shew *White*, yet viewed through a *Microscope*, they all appear *transparent*. In like manner, as the *serum* of *Blood*, *Whites* of *Eggs*, *Tendons*, *Hairs* and *Horns* themselves are *transparent*, and without much *Smell* or *Taste*, their *Principles* being, in all of them, more or less *concentred*: But when ever these *Principles*, are forcibly *resolved*, they are ever variously invested with all those *Qualities*.

How the several Parts are Nourish'd and Form'd.
(b) §. 21.
(c) *Idea*, §. 49, 51.

30. §. And as from the *Concentration* of the *Principles*, in every *Organical Part*, the said *Parts* do thus far, all agree: So, from the *Predominion* of the *Principles* of each *Part*, the rest are controuled; not only to a *Concentration*, but an *Assimilation* also; whereby, the *specifick Differences*, of the several *Organical Parts*, are preserved. Hence the *succiferous Vessels* are always *Tough* and very *Pliable*; for so are all *Barques*, wherein these *Vessels* abound; so is a *Handful* of *Flax*, which is nothing else but a heap of the *succiferous Vessels* in the *Barque* of
that

- that *Plant*. For besides *Water*, and *Earth*, an *Alkaline Salt* and *Oyl* are, as is said, the predominant *Principles* of these *Vessels*. (a) It is then the *Oyl*, chiefly, by which these *Vessels* are *Tough*: for being of a tenacious Nature, by taking hold of other *Principles*, it marries them together; and the *Alkaline Salt* and *Earth*, concentrated with it, addeth to it more *Strength*. Hence the *Caput Mortuum* of most Bodies, especially those that abound with *Oyl* and a *sal Alkali*, is brittle and friable; those *Principles*, which were the *Ligaments* of the rest, being forced away from them. From the same Cause, the *Parenchymous Parts* of a *Root*, even in their Natural State, are brittle and friable; *sc.* Because their *Earthy*, and especially *Oleous* and *Saline Principles* are, as is said, (b) so very few. Therefore all *Piths* and more simple *Parenchyma's*, break short, so *Corn*, and the *Roots* of *Potato's*, and divers other *Plants*, being dried, will easily be rub'd to *Meal*; and many *Apples*, after *Frosts*, eat mealy; the *Parenchymous Parts* of all which, are not only by *Analogy*, but in Substance or Essence, the self same Body. (c)
- (a) §. 21. 31. §. And as the *Consistence* of the several *Organical Parts*, is dependent on their *Principles*; so are their *Figures*. And first, the *Succiferous Vessels*, from their *Alkaline Salt*, (d) grow in Length. For by that Dimension, chiefly, This salt always shoots: And being a less moveable *Principle* than the rest, and so apt more speedily to fix or shoot: It thus overrules them to its own *Figure*. And even as the Shape of a *Button* dependeth on the *Mould*, the *Silk* and other Materials wrought upon it, being always conformable thereunto: so here; the salt is, as it were, the *Mould*; about which, the other more passive *Principles* gathering themselves, they all consort and fashion to it. Hence also the same *Sap-Vessels* are not *pyramidal*, as the *Veins* of *Animals*; but of an equal bore, from end to end; the *shootings* of the said Salt, being also figured more agreeably to that *Dimension*. And as by the *Saline Principle*, these *Vessels* are *Long*; so by the *Oleous*, (e) they are every where *Round*, or properly *Cylindrical*; without some joynt Efficacy of which *Principle*, the said *Vessels* would be *Flat*, or some way *Edged* and *Angular*, as all *saline shoots*, of themselves, are; as those of *Alum*, *Vitriol*, *sal Ammoniac*, *Sea Salt*, *Nitre*, &c. And because the *Spirituos* and more *Fluid* part of the *Principles*, is least of all apt to fix; while therefore, the other parts fix round about, This will remain moveable in the *Centre*; from whence every *Vessel* is formed, not into a *solid*, but *hollow Cylinder*; that is, becomes a *Tube*.
- (b) §. 19. (c) *Lib.* 1. c. 7. §. 14. (d) *P.* 2. §. 21. (e) *Ib.*
32. §. The *Lactiferous Vessels* are *tubulary*, as the *Lymphæducts*, but of a somewhat wider *Concave* or *Bore*. For being their *Principles* are less *Earthy* and *Oleous*, and also more loosely *Concentred*; as from their easie corruption or *Resolution* by the *Aer*, it appears they are: they are therefore more tender, and so more easily dilative, and yielding to the said *Spirituos* part in the *Centre*. And by this means, obtaining a wider *Bore*, they are more adapted to the free motion of the *Milky Content*: which being an *Oleous* and Thicker *Liquor*, than that in the *Lymphæducts*; and having no advantage of *pulsation*, as the *Blood* hath in *Animals*; might sometimes be apt to stagnate, if the *Vessels*, through which it moves, were not somewhat wider.

33. §. As the *saline Principle* is the *Mould* of the *Succiferous*, so is the *Aerial* of the *Aer-Vessels*. (a) Now the Particles of *Aer* strictly so called, at least of that part of it concerned in the Generation of the *Aer-Vessels*, I suppose, are crooked: and that by composition of many of those crooked ones together, some of them become *Spiral*, or of some other winding *Figure*: and that thereupon dependeth the *Elastic* Property of the *Aer*, or its being capable of *Rarefaction* and *Condensation* by force. Wherefore, the said *crooked* Particles of the *Aer*, first *shooting* and *setting* together, as the *Mould*, the other *Principle* cling and *fix* conformably round about them. So that, as by force of the *Saline Principles*, the rest of them are made to *shoot* out in *Long continued Fibres*; so by force of the *Aerial*, those *Fibres* are still disposed into *Spiral Lines*, thus making up the *Aer-Vessels*. And according as there are fewer of these *Aerial* Particles, in proportion to the *Saline*, the *Concave* of the *Aer-Vessels* is variously wider, or the *Fibres* continue their *shooting* by wider *Rings*; as those that come nearer to a *right Line*, and so are more compliant to the *Figure* and shooting of the *Saline* parts. And whereas the *Lympheducts*, shooting out only in length, are never sensibly *amplified* beyond their original size: These, on the contrary, always, more or less, enlarge their *Diameter*; because their *Fibres*, being disposed into *Spiral Lines*, must needs therefore, as they continue their growth, be still dilated into greater and greater *Rings*. And being at the bottom of the *Root* more remote from the *Aer*, and so having somewhat fewer Particles purely *Aerial*, there ingredient to them, then at the top; they fall more under the government of the *Saline*, and so come nearer to a *right Line*, that is into greater *Circles*; and so the *Aer-Vessels*, made up of those *Circles*, are there generally wider. (b)

34. §. By mediation of their *Principles*, the *Parenchymous Parts* likewise of a *Root* have their proper *Contexture*. For from their *Acid Salt* they are *Fibrous*; from their *Oyl*, the *Fibres* are *Round*, and in all parts even within themselves; and from their *spirit*, it is most probable, that they are also *hollow*. But because the *spirit* is, here, more copious than the *Aer*; and the *saline Principle* an *Acid*, (c) and so, more under the government of the *Spirit*, than is an *Alkali*; therefore are not the said *Fibres* continued in *straight Lines*, as the *Sap-Vessels*; or by one *uniform* motion, into *spiral* lines, as the *Fibres* in the *Aerial*; but *winding*, in a circular manner, to and fro a *thousand ways*, agreeable to the like *motions* of the *Spirit*, that most *active*, and here most *predominant Principle*. And the *spirituous Parts* being, as is said, here more copious and redundant, they will not only suffice to fill up the *Concaves* of the *Fibres*, but will also gather together into innumerable little spaces, without them: whence the *Fibres* cannot wind close together, as *Thred*, in a *Bottom* of *Yarn*; but are forced to keep at some distance, one parcel from another, and so are disposed, as *Bread* is in baking, into *Bladders*. (d)

35. §. And the *under Fibres* being *set* first, as the *Warp*, the *spirituous* parts next adjacent, will incline also to *fix*, and so govern an *over work* of *Fibres*, wrapping, as the *Woof*, in still smaller *Circles* round the other: whereby they are all knit together. (e) For the same reason, the *Lympheducts*, being first formed, the *Parenchymous Fibres* *set* and *wrap* about These also. (f) And the *Aer-Vessels* being formed

(a) P. 1. c. 4.
§. 23. & P. 2.
§. 24.

(b) P. 1. c. 4.

§. 16.

(c) §. 19.

(d) P. 1. c. 3.

§. 4.

(e) P. 1. c. 5.

§. 9.

(f) P. 1. c. 5.

§. 12.

formed in the Center, the *succiferous* run along those likewise (as *volatile salts* shoot along the sides of a Glass, or *Frost* upon a Window) and so are, as it were, *Incrustate* about them in a *Ring*.

How the several Parts come to be Situate or Dispos'd.

36. §. SOME OF THE more Ætherial and Subtile parts of the *Aer*, as they stream through the *Root*, it should seem, by a certain *Magnitisme*, do gradually dispose the *Aer-Vessels*, where there are any store of them, into *Rays*. This Attraction (as I take leave to call it) or *Magnetick* power betwixt the *Aer* and these *Vessels*, may be argued, From the nature of the *Principles* common to them both: From the *Electral* nature of divers other Bodies; the *Load-stone* being not the only one which is attractive: And from other Effects, both before (a) and hereafter mentioned. Wherefore in the inferiour parts of the *Root*, they are less Regular; (b) because more remote from the *Aer*. And in the upper parts of many *Roots*, as *Cumfery*, *Borage*, *Parfnep*, where those that are next the Centre are confused, or differently disposed; those next the *Barque*, and so nearer the *Aer*, are postured more Regularly, and usually into *Rays*. For the same reason it may be; that even the *sap-Vessels* in the *Barque*, as often as the *Aer Vessels* are more numerous, are usually disposed into *Rays*, as following the direction of the *Aer-Vessels*. And that the *Parenchyma* of the *Barque*, is disposed into *Diametral Portions*: and that where the *Aer Vessels* are fewer or smaller, these *Portions* are likewise smaller or none; as in *Chervil*, *Asparagus*, *Dandelion*, *Orpine*, *Bistort*, *Horse-Radish*,

Tab. 7, 8, 9, *Potato's*, &c.

37. §. The said Ætherial parts of the *Aer*, have a Power over the *Aer-Vessels* not only thus to Dispose them; but also to Sollicite and spread them abroad from the Center towards the Circumference of the *Root*. By which means, those *Roots* which have no *Pith* in their lower parts, obtain one in their upper. (c) And the same *Pith*, which in the lower part, is ratably, small, in the upper, is more or less enlarged. (d)

(c) P. I. c. 5. §. 1.

(d) Ib. §. 4.

38. §. The Spreading of these *Vessels* is varied, not only according to the Force the *Aer* hath upon them, but also their own greater or less *Aptitude* to yield thereto. As often therefore, as they are Slenderer, they will also be more Pliable and recessive from the Centre, towards the Circumference. Hence, in such *Roots* where they are small, they stand more distant; as in *Turnep*, *Jerusalem Artichoke*, *Potato's*, and others; and so their *Braces* are fewer: and in the same *Root*, where they are smaller, their distance is greater. Besides, in these smaller *Aer-Vessels*, the *Rings* being less, and the *Spiral Fibres* whereof they are made, continuing to shoot; the said *Rings* therefore, must needs be so many more, as they are smaller; and so take up more space by the length of the *Root*; and so, not being capable of being crowded in a *right line*, every *Vessel* will be forced to recede to a *crooked* or *bowed* one.

Tab. 2, & 6.

39. §. The *Sap Vessels*, being by the *Parenchymous Fibres* knit to these, will likewise comply with Their motion, and spread abroad with them. Yet being still smaller (e) and more pliable than the *Aer-Vessels*, and so more yielding to the intercurrent *Fibres* of the *Parenchyma*, their *braced Threds* will, sometimes, be much more divaricated, than those of the *Aer-Vessels*; as in *Jerusalem Artichoke*. And because the *Succiferous Vessels*,

(e) P. I. c. 3. §. 16.

Tab. 6.

Vessels, although they are joyned to the *Aerial* by the *Parenchymous Fibres*, (a) yet are not continuous with them; neither fall under (a) P. 1. c. 5. the like *Attractive Power* of the *Aer*, as the *Aerial* do; the *Aerial* §. 12. therefore, upon their spreading, do not always carry all the *Succiferous* along with them; but often, if not always, leave many of them behind them sprinkled up and down the *Pith*; as in *Parsley*, *Carrot*, *Tab. 6.* *Jerusalem Artichoke*, *Turnep*, &c. may be seen.

40. §. The spreading of the *Aer-Vessels* still continued, several of them, at length, break forth beyond the circumference of the *Root*; and so are distributed, either in the lower parts, into *Branches* and *Strings*; or at the top, into *Leaves*. And lest they should all spread themselves into *Leaves*, and none be left for the *Caulis*; as where they are very small, or the *Sap-Vessels* to bound them, are but few, they might; therefore divers of them are, oftentimes, more frequently braced in the *Centre*; for which reason, they cannot so easily separate and spread themselves from thence, but run more inwardly up into the *Caulis*, as in *Borage*. *Tab. 6.*

41. §. FROM THE various *sizes*, *Proportions*, and *Dispositions* of the *Parts*, *Roots* are variously sized, shaped, moved and aged. Those which, by their Annual Growth, are large; have fewer, both *Aerial*, and *Sap-Vessels*, and a more copious *Parenchyma*. So that the *Aer-Vessels*, or rather, the *Aery Ferment* contained in them, volatilizing only a smaller portion of the *Sap*; the said *Sap* is less capable of advancement into the *Trunk*; and so must needs remain and fix more copiously in the *Root*, which is thereby more augmented. And where the *Sap-Vessels* alone, are but few, the *Root* is yet, ratably, somewhat large: but where they are numerous, it is never so, as to its Annual Growth, in any proportion to their Number: Because their *Tincture*, which is *Alkaline*, will go farther in setting the *Parenchymous Parts*: than the *Tincture* of *These*, which is *Acidulate*, will go, in setting Them. (b) How the whole *Root* is differently sized and shaped.

42. §. When the *Aer-Vessels* are more pliable and sequent to the Attraction of the *Aer*, and so spread themselves, and the *Succiferous* together with them, more abroad; in the manner as hath been said; the *Root* also will grow more in *Breadth*; the nutrition of the *Parenchymous Parts*, to which the *Vessels* are adjacent, being thus, by the same dimension, more augmented; as in *Turnep*, *Jerusalem Artichoke*, &c. But where these are not spread abroad, the *Root* is but slender; as in *Asparagus*, *Dandelion*, &c. *Tab. 2, & 7.* *Tab. 7, & 8.* (b) P. 2. §. 31.

43. §. If the *Aer-Vessels* be contracted into, or near the *Centre*, and are somewhat Large or Numerous; and the *Succiferous*, also more copiously mixed with, or surrounding them; the *Root* grows very Long; as do those of *Fenil*, *Vine*, *Liquirish*, &c. For the *Aer-Vessels* containing a more copious *Ferment*, it will well digest and mature the *Sap*: Yet the *Succiferous* being over proportioned to them; the *Sap* will not therefore, be so far volatilized, as to ascend chiefly into the *Trunk*; but only to subserve a fuller Growth of their *Vessels*: which being more numerous, and so more sturdy, and less sequent to the expansive motion of the *Aerial*; this their own Growth, and consequently, that of all the other *Parts*, cannot be so much in *Breadth*, as Length. *Tab. 2, & 17.*

44. §. Where the same *Aerial Vessels* are Fewer, or more Contracted, or sheathed in a Thicker and Closer *Barque*; the *Root* is smooth; and less *Ramified*, as in *Asparagus*, *Peony*, *Dandelion*. But where more Numerous, sheathed in a Thinner *Barque*, Smaller, or more Dilated; the *Root* is more *Ramified*, or more *stringy*, as in *Columbine*, *Clary*, *Beet*, *Nicotian*. For being, as is said, by these means, more sequent to the Attraction of the *Aer*; approaching still nearer the circumference of the *Barque*, they at last strike through it, into the *Earth*. And the *Parenchymous Fibres* being wrapped about them, and the *Succiferous Vessels* knit to them by those *Fibres*; (a) therefore they never break forth naked, but always invested with some quantity of these *Parts* as their *Barque*: where by, whatever *Constitutive Part* is in the *main Body* of the *Root*, the same is also in every *Branch* or *String*.

Tab. 7, 8.

Tab. 2, & 7.

(a) P. I. c. 5. §. 12.

Tab. 6.

(b) P. I. c. 1. §. 2, 3. Tab. 11.

(c) P. I. c. 4. §. 15.

Tab. 6.

45. §. From the same Expansion and Pliability of the *Aer-Vessels*, the *Root* oftentimes putteth forth *Root-Buds*; which gradually shoot up and become so many *Trunks*. In the Formation of which *Buds*, they are pliable and recessive all kinds of ways; being not only invited Outward, toward the Circumference of the *Root*, as in *Root-strings*, but also spread more Abroad every way, so as to make a *Root-Bud*: Where as in the said *Root-strings*; they are always more Contracted. Which, in respect of the Disposition of the *Parts*, is the principal difference betwixt the *Root* and the *Trunk*, as hath been said. (b) Hence, those *Roots*, chiefly, have *Root-Buds*, which have the smallest *Aer-Vessels*; (c) these, as is said, being the most pliable and Expansive.

46. §. But because the expansiveness of the *Vessels*, dependeth also, in part, upon the Fewness of their *Braces*; therefore the said *Buds* shoot forth differently, in divers *Roots*. Where the *Braces* are fewer, the *Buds* shoot forth beyond the Circumference of the *Root*, as in *Jerusalem Artichoke*; where more close, as in *Potato's*, the *Buds* lie a little absconded beneath it; the *Aer-Vessels* being here, by their *Braces*, somewhat checked and curbed in, while the *Barque* continueth to swell into a fuller Growth.

47. §. If the *Aer-Vessels* are all along more equally sized, the *Root* is so also, or *Cylindrical*; as are those of *Eryngo*, *Horse-Radish*, *Marshmallow*, *Liquirish*, &c. But if unequal, growing still wider towards the *bottom* of the *Root*; then the *Root* is unequal also: But groweth, as is observable, quite contrarily to the *Aer-Vessels*; not Greater, as They do; but still smaller, or *pyramidally*; as in *Fenil*, *Borage*, *Nettle*, *Patience*, *Thorn-Apple*, &c. is apparent. For the *Aer-Vessels* being considerably wider about the *bottom* of these *Roots*; they there contain a more Copious *Ferment*: Whereby the *sap* is there also more volatilized, and plentifully advanced to the Upper Parts. Withal, thus receiving into themselves, and so transmitting to the upper Parts, a more plentiful *Vapour*, they hereby rob the *Parenchymous Parts* of their Aliment, and so stint them in their Growth.

How Roots are differently Mov'd. Tab. 8. 48. §. FROM THE different Proportions and Situation of the *Parts*, the *Motions* of *Roots* are also various. For where the *Aer-Vessels* are spread abroad and invested with a thinner *Barque*; the *Root* runs or lies *Level*, as in the *level-Roots* of *Primrose*, *Bishops-weed*, *Anemone*, &c. may

may be seen. So that these *Roots*, as by the *Perpendicular Strings*, which shoot from them into the *Earth*, and wherein the *Aer-Vessels* are contracted into their Center, they are Plucked down (a): So by the *Aer-Vessels*, which stand nearer the *Aer*, and more under its Attractive Power (b) they are invited upwards; whereby they have neither ascent nor descent, but keep level, betwixt both.

49. §. But if these *Vessels* are Contracted, standing either in, or near the Centre, and are invested with a *Barque* proportionably Thick; the *Root* striketh down *perpendicularly*, as doth that of *Dandelion*, *Bugloss*, *Parsnep*, &c. And therefore the said *Vessels*, although they are spread abroad in the *level Roots*, yet in the *perpendicular* ones of the same *Plant*, they are always contracted; as by comparing the *Level* and *Down-right Roots* of *Ammi*, *Primrose*, *Jerusalem Artichoke*, *Cowslip*, and others, is manifest.

50. §. If the *Aer-Vessels* are Contracted, and Environed with a greater number of *Succiferous*, the *Root* grows deep; that is, *perpendicular* and long. (c) *Perpendicular*, from the Contraction of the *Aer-Vessels*; (d) and long, from the Predominion of the *Succiferous*, which in their growth, are extended only by that Dimension, as in *Liquirish*, *Eryngo*, &c.

51. §. If the *Succiferous* are over proportioned to the *Parenchymous Parts*, but under to the *Aer-Vessels*; the *Root* is *perpendicular* still, but groweth shallow: The *Succiferous* being sturdy enough to keep it *perpendicular*; But the *Aer-Vessels* having a predominion to keep it from growing deep; as in *Stramonium*, *Nicotian*, *Beet*, &c.

52. §. If, on the contrary, the *Parenchymous Parts* are predominant to the *Aer Vessels*; and that, both in the *Root* and *Trunk*; then the whole *Root* changeth place, or descends. (e) For the said *Aer-Vessels*, having neither in the *Trunck*, nor in the *Root*, a sufficient Power to Draw it upwards; it therefore gradually yields to the Motion of its *String-Roots*; which, as they strike into the Soil, Pluck it down after them. And because the old *Strings* annually rot off, and new ones successively shoot down into the *Ground*, it therefore annually still descendeth lower; as in *Tulip*, *Lily*, &c. may be observed.

53. §. Where the *Aer-Vessels* are much spread abroad, and also numerous, the *Root* oftentimes, as to its several parts, descends and ascends both at once. So *Radishes* and *Turneps*, at the same time, in which their nether parts descend; their upper, (where the said *Vessels* are more loosely braced, and spread more abroad than in the lower parts) do ascend, or make their Growth upward. Hence also, the upper part of most young *Roots* from seed, ascends: Because the first *Leaves*, being proportionably large, and standing in a free *Aer*, the *Aer-Vessels* therein, have a dominion over the young *Root*; and so themselves yielding to the sollicitation of the *Aer*, upwards; they draw the *Root*, in part, after them.

54. §. BY THE *Situation* and *Proportions* of the *Parts*, the *Age* of the *Root* is also varied. For if the *Sap-Vessels* have the greatest Proportion, the *Root*, is *Perennial*, and that to the farthest extent, as in *Trees* and *Shrubs*. Because these *Vessels* containing a more copious *Oyl*; (f) and their several *Principles* being more closely Concentred, they are less subject to a *Resolution*, that is, a *Corruption* or *Mortification* by the *Aer*.

55. §. If the *Parenchymous Parts* have much the greatest, the *Root* seldom liveth beyond Two Years; but afterwards perisheth either in whole, or in part; as do divers *bulbous, tuberous,* and other *Roots*; whether they are more Porous and Succulent, or more Close and Dry. If Porous, all the *Liquid Principles* standing herein more abundant, either by a stronger Fermentation, or otherwise, Resolve the *fixed ones* of the *Organical Parts*; whence the whole *Root*, rots; as in *Potato's*. So also *Parsneps,* and some other *Roots*, which, in a hard and barren *soil,* will live several years, in another more rank, will quickly rot. If the *Parenchyma* be Close, then the *Aer,* chiefly, entring in and filling it up, thus mortifies the *Root*; not by Rotting the *Parts,* but over Drying them; as in *Satyrion, Rape-Crowfoot, Monks-*

(a) P. I. c. I.
§. 13, 16. &
P. 2. §. 28.

Tab. 9.

(b) P. 2. §.
25, 26.

56. §. But if the *Aer-Vessels* have the greatest Proportion, and especially if they are more large, and withall, are spread more abroad: the *Root* is *Annual,* as in *Thorn-Apple, Nicotian, Carduus Ben. &c.* And of the same Kindred, if any, those are *Annual,* which have the most *Aer-Vessels.* So *Endive* and *Sonchus,* which have store of *Aer-Vessels,* are both *Annual:* whereas *Cichory,* in which they are fewer, is a *Perennial Root.* For hereby a more copious *Aer* being Transfused into all the other *Parts*; (b) they are thus, by degrees, hardened, and become sticky; and so impervious to the *Sap,* which ought to have a free and universal Transition from *Part* to *Part.* As *Bones,* by *Precipitations* from the *Blood,* at length, cease to grow. Or the same more abundant *Aer,* so far volatilizeth the *Liquors* in the *Root,* that they are wholly advanced into the *Trunk,* and so the *Root* is starved. Whence also the *Aer-Vessels* of the *Trunk;* where they are numerous, and over proportioned to the *Bulk* of the *Root,* as in *Corn;* they so far promote the advance of the *Sap,* as to exhaust the *Root,* sucking it into a Consumption and Death.

How the
Contents of
the several
Parts are
made.

(c) Idea, §.
54.

57. §. FROM THE *Principles* of the *Parts,* their *Contents* and the several *Qualities* hereof are also various; (c) the *Fluid* of each *Organical Part,* being made, chiefly, by *Filtration* through the sides thereof; such of the *Principles* in the *Sap,* being admitted into, and transmitted through them, as are aptest thereunto. In the like manner, as when *Oyl* and *Water,* being poured upon a *Paper,* the *Water* passeth through, the *Oyl* sticks: or as the *Chyle* is strained through the *Coats* of the *Guts,* into the *Lacteal Vessels:* or as *Water* in *Purgations,* is strained through the *Glands* of the same *Guts,* from the *Mesenterical.*

(d) P. 2.
§. 19.

58. §. The *Principles* therefore of the *Parenchymous Fibres* being *spirituous, acid,* and *aerial,* they will also admit the like into them; excluding those chiefly which are *Alkaline* and *Oleous.* (d) And as by the *Conjugation* of such *Principles* in the *Fibres,* the like are capable of admittance into their *Body:* so the *Proportion* and *Union* of the same *Principles,* regulates the *transmission* hereof into their *Concave.* Wherefore, the predominant *Principles* of the *Fibres* being chiefly *acid,* next *spirituous,* and *aery,* the more *aery* ones will be transmitted. For if more of *them* should fix they must do so by similitude and adhesion: But where there are fewer similiary parts to adhere to, fewer must adhere. The *Fibres* therefore contain so many parts of *Aer,* as to admit many

many more into their Body; but not to *fix* them; which therefore must needs, upon admission, pass through into their *Concave*; where, together with some other more *spirituous* parts, they make an *Ætherial Fluid*. And because some *aqueous* or *vaporous* parts will also strain through with them; hence it is, that as more and more of these enter, they by degrees still thrust out the *aery* ones; which quitting the more *succulent Fibres* of the *Parenchyma*, are forced to betake themselves to the *dryer* ones, *scil.* all those, whereof the *Diametral Portions* do consist. For the same reason the *Aery* parts being gradually excluded the *succulent Fibres* of the *Barque*; they are forced to recede and transmigrate into those of the *Pith*. And the *Fibres* of the *Pith* themselves being filled, and the *Aery* parts still forced into them; they at length also strain through the *Fibres* into the *Bladders*: whence it comes to pass, that while the *Barque* is *succulent*, the *Pith* is often times filled with *Aer*.

59. §. The *Lympheducts* being more *earthy*, *Salinous*; *oleous*, and *(a) P. 2.*
aqueous, will both *admit* and copiously *fix* the like *Principles*, as their *§. 21.*
 proper *Aliment*. The *Water* being more *perfluent* than the rest, will therefore strain, with a lighter *Tincture* of them, into their *Concave*. Especially the *Oleous* parts of these being *rampant*, and less apt to *fix* and *seize* the *aqueous*, upon their entrance, than the *saline*.

60. §. The *Lactiferous*, appearing to be made, chiefly, by the *Constipation* of the *Parenchymous Parts* all round about their *Sides*; the *Liquor* contained in those *Parts*, although it may easily enough be transfused into the *Hollow* of these *Vessels*; yet seems it not, with equal facility, to be refunded thence: So that the *thinner* and more *aqueous* *Portion* only, passing off; the remainder, is, as it were, an *Oleous Elixyr*, or extract, in the form of a *Milk*.

61. §. The *Fluid Ferment* contained in the *Aer-Vessels*, is also in part, dependent on the *Principles* of those *Vessels*, being in their percolation *tinctured* therewith. But because the percolation is not made *through* the *Body* of the *Fibres* whereof the *Vessels* are composed, but only *betwixt* them; therefore the transient *Principles* more promiscuously, yet with an over proportion of *dryer Particles*, pass into the *Concaves* of these *Vessels*, and so are herein all immersed in a *Body* of *Aer*. *(b) P. 2.*
Parenchymia, admitting and containing a more *Aery* and *Ætherial* *§. 24.*
Fluid.

62. §. The *Contents* are varied, not only by the *Nature*, but also the *Proportion* and *Situation* of the *Parts*, whereby the said *Contents* are with different *Facility* and *Quantity*, communicated one to another. Hence it is, partly, that a *Vine*, or that *Corn*, hath so little *Oyl*: *sc.* Because their *Aer-Vessels*, in proportion with the other *Parts*, are so *Great* and *Numerous*: in *Corn*, the *Stalk* being also very hollow, and so becoming as it were, one *Great Aer-Vessel*. For the *Oily* parts of the *Sap*, are so exceedingly *attenuated* *(c) P. 2. §.*
 these *Vessels*; that they are, for the most part, so far *immersed* in the *25, & 56.*
Spirit, or mixed therewith, as not, by being collected in any considerable *Body*, to be distinguishable from it. And the affinity that is betwixt *Spirits* and *Oils*, especially *Essential*, is manifest: Both are very inflammable; Both will burn all away; The *Odors*, which we call the *Spirits* of *Plants*, are lodged in their *essential Oyl*; Both, being
 duly

duly Rectified, will mix as easily together, as *Water* and *Wine*. So that, although *Oyl*, by the separation of its *earthy* and *Saline* parts, which give it its sensibly *oleous* Body, may not be so far attenuated, as to produce a *Spirit*; yet that it may so far be attenuated, and so be mixed therewith, as *not to be discerned from it*, as in the forementioned *Plants*, will be granted.

63. §. Hence it is, that the *Lactiferous* standing more remote from the *Aer-Vessels*, and the *succiferous* interposing; (a) the *Liquor*, therefore, contained in them, is not so much under the government of the *Aerial Ferment*, and is thence, partly, more *Oily*. For the same reason, all *Roots* which are *Milky*, so far as I have observed, have an under-proportion of *Aer-Vessels*; these being either Fewer or Smaller.

How the Odours of Plants are made.

64. §. FROM what hath been said, we may receive some information, likewise, of the *Odours*, *Colours*, and *Tastes* of *Plants*. And for *Odours*, I suppose, That the chief Matter of them, is the *Aerial Ferment* contained in the *Aer-Vessels*. Not but that the *other Parts* do also yield their smell; but that *these* yield the *strongest* and the *best*, and *immediately* perceptible in *fresh*, *undried* and *unbruised Plants*. For the *Aer* entring into, and passing through the *Root*, and carrying a *Tincture*, from the several *Organical* and *Contained* Parts, along with it, and at last entring also the *Concaves* of the *Aer-Vessels*; it there exists the most *Compound*ed and *Volatile Fluid*, of all others in the *Plant*, and so the fittest matter of *Odour*: and such an *Odour*, as answers to that of all the *Odorous* parts of the *Plant*. (b) Wherefore the *Organical Parts*, being well censed of their *Contents*, smell not at all; Because the *Principles* hereof are, as hath been said, so far *fixed* and *concentred* together. Hence also the *Contained Parts* themselves, or any other *Bodies*, as their *Principles* are any way more *fixed*, they are less *Odorous*: So is *Rosin*, less than *Turpentine*, and *Pitch*, than *Tar*; and many the self same *Bodies*, when they are *coagulated*, less than when they are *melted*. So also *Musk*, which is not so liquid as *Civet*, is not so strong; nor *Ambergreece*, as *Musk*: For although it hath a more excellent smell, than *Musk* hath, yet yieldeth it not so easily; since it is a more *fixed* Body, and requireth some *Art* to be opened. Hence also the *Leaves* of many *Plants* lose their *Odour* upon *rubbing*: Because the *Aer-Vessels* being thereby broken, all their *contained odorous Fluid* vanisheth at once: which before, was only strained gradually through the *Skin*. Yet the *fixed* Parts themselves, upon drying, are so far altered by the *Sun* and *Aer*, as to become *resoluble*, and *volatile*, and thence *odorous*.

How their Colours.

65. §. SO ALSO of their *Colours*. As whence the *Colours* of the *Skins* are varied, For divers of the *Sap-Vessels*, together with the *Parenchymous Parts* successively falling off from the *Barque* into the *Skin* (c) by their proximity to the *Earth* and *Aer*, their *Sulphureous* or *Oleous Principle* is more or less *resolved*, and so produceth divers *Colours*. So those *Roots* which turn *purple* any where within, have usually a *black*er *Skin*; the one of those two *Colours* being, by a resolution and corruption of parts, easily convertible into the other, as in *Cumfry*, *Thistle*, &c. So the *Milk* of *Scorzonera*, contained in the *Vessels* of the *Barque*, upon drying, turneth into a *brown Colour*:

lour: Wherefore the *skin*, in which there are divers of those *Vessels*, is of the *same*. So both the *Milk* and *Skin* of *Lovage* is of a brownish *yellow*. But *Parsnep* hath a clearer *Sap* in all its *Vessels*, and a whiter *Skin*. So *Potato's*, being cut traverse, after some time out of ground, have divers *red specks* up and down where the *Vessels* stand; and their *skin* is accordingly *red*.

66. §. The reason, I say of these *Colours*, is the *resolution* or *re-feration* of the *Principles* of the several *Parts*, chiefly, by the *Aer*, and a *lighter mixture* of them consequent thereupon: whereby the *Sulphureous* or *Oily Parts*, which were before *concentred*, are now more or less *rampant*, discovering themselves in divers *Colours*, according as they are diversly mixed with the other *Principles*. Hence these *Colours* are observable, according to the nature of the *Parts* wherein they are, or whereunto they are *adjacent*: So where the *Lympheducts* doe run, there is a *Red*, or some other *Sulphureous Colour*; the *Oleous Principles* being, as is said, (a) more copious in these *Vessels*; as (a) P. 2. §. 21. in the *Bark* of *Peony*, the inward parts of *Potato's*, &c. may be seen. But the *Parenchymous Parts*, where more remote from the said *Vessels*, they are usually *White*, or but *Yellow*: the *Sulphureous Principle* of these *Parts*, being, as hath been said, but sparing. (b) The (b) P. 2. §. 20. same is seen in those *Roots* which shew both *Red* and *Yellow*: those *Parts*, principally, where the *Succiferous Vessels* run, being *Red*; but those *Parts*, where only the *Aer-Vessels* are mixed with the *Parenchymous*, being *Yellow*; as in *Patience*. So likewise the *pithy* part of a *Carrot*, where the *Aer-Vessels* have very few *Succiferous* mixed with them, is *Yellow*; but the *Barque*, where the *Succiferous* are very numerous, is *Red*. For the same reason, many *Roots*, which are *Whiter* in their upper parts, are *Purple* or *Reddish* in their inferior, as *Avens*, *Strawberry*, &c. Because those lower parts, having lain longer (c) under ground (these being *descending Roots*) their (c) P. 1. §. 13. *Principles* are, thereby; somewhat more *resolved*, and so the *Oleous*, ramp and spread all over the rest in that *Colour*.

67. §. And that the *Resolution* of the *Sulphureous* and other *Principles* is partly effected by the *Aer*, appears, In that, where the *Aer* hath a free access to the *Succiferous Vessels*, the *Colours* are there, chiefly produced, or are more conspicuous. So in *Potato's*, where the *Succiferous Vessels* are either next to the external *Aer*, as in the *Skin*; or contiguous with the *Aer-Vessels*, as in the *Ring* within the *Barque*; there, they produce a *Red*: but where more remote from both, as in the middle of the *Barque*, and *Centre* of the *Root*; there they produce none. Hence also it is, that the *Leaves* and *Flowers* of some *Plants*, as *Bloodwort*, *Wood-sorrel*, *Radish*, *Jacea*, &c. although *Green* or *White* in the greatest portion of their *Parenchymous Part*; yet where the *Succiferous* and *Aer-Vessels* run together, they are of *Red*, *Blue*, and other *Colours*; the *Oleous parts* of the one, being *unlocked* and *opened*, by the *aery* of the other.

68. §. AND LASTLY, of their *Tasts*. Most *Roots* which are *How their Tastes.* *acres* or *bitting*, have a very copious *Parenchyma* in proportion with the *Succiferous Vessels*, as of *Arum*, *Dragon*, and others: Because the *Saline* and other *Principles* are not so much *hot*, by any sufficient quantity of *Sulphureous*, from those *Vessels*, in which the *Sulphur*, as is said, is more abundant; (a) but rendered rather *pungent*, from (a) P. 2. §. 21. some

some Spirit and Aer. But divers *Umbelliferous Roots*, especially which abound with *Lactiferous Vessels*, are hot; as *Fenil*, *Lovage*, *Angelica*, &c. Yet is it not their Oyl alone that makes them hot, but the combination thereof with the *Saline Parts*: as is manifest, from the nature of the *Seed* of these *Plants*; wherein, as the Oyl is most copious; So being held to a Candle till they burn, constantly spit; which cometh to pass, by the eruption of the *saline Parts*: and is the very same effect, with that which followeth upon burning of *Serum* or *Blood*. And therefore, as these *seeds* are more hot, they also spit the more; So those of *Cumine*, which, though fulsom, yet are not so hot, spit less; *Fenil* and *Dill*, which are hotter, more; there being a greater quantity of *volatile Salt* contained herein. Hence all *Essential Oyls* are hot, the *Spirit* and *volatile salt*, being incorporated herewith. And some of them will shoot, and crystallize as *salts* do, as that of *Anise*; which argues a mixture of a considerable quantity of *volatile salt*. As also doth the Nature of these Oyls, in being amicable to the *Stomach*, *Carminative*, and sometimes *Anodyne*; scil. as they kill some *fetid*, or *corrosive Acid*: for *volatile Salts* themselves will have the like operation in some cases as these Oyls.

Tab. 13.

69. §. Many *Lactiferous Roots*, as *Taraxacum* and others of that kind, are not so much hot, as bitter. For although by the *Lactiferous Vessels* they are very Oily; yet those *Vessels* being posited in *Rings*, and not in *Rays*, and having no *Diametral Portions* running through their *Barque* to the *Aer-Vessels*; the *Acido-Aerial Parts* do hereby, although not mortifie, yet so far refract the *saline*, lightly binding up the *Oleous* therewith, as to produce a bitter Taste. So, many sweet Bodies, upon burning, become bitter; the *Acid Parts*, now becoming rampant, and more copiously mixed with the *Oleous*.

70. The *Roots*, or other *Parts*, of many *Umbelliferous Plants*, have a sweetish Taste, as both the *sweet*, and *Common Chervil*; both the *Garden*, and *wild Carrot*; *Parsnep*, *Fenil*, &c. the *Saline Principles* being concentrated in the Oily, and both of a moderate quantity with respect to the rest. For by the Oily, the *saline* is rendered more smooth and amicable; and both being moderate, they are not therefore hot, as in some other *Umbelliferous Roots*; but by the predominion of the other *Principles*, made mild. Hence it is, that *Sugar* it self is sweet, scil. because it is an *Oleous salt*; as is manifest, from its being highly inflammable; its easie dissolution by a moderate, *Fire*, without the addition of *Water*; and in that, being melted with *Turpentine*, and other *Oily Bodies*, it will mix together with them. So also the *Acid Parts* of *Vinegar*, being concentrated in the *Salino-sulphureous* of *Lead*, produce a *sugar*. Hence *Barley*, which upon *Distillation* or *Decoction* yeildeth only an *acid*; being turned into *Mault*, becomes sweet. Because, being steeped, couched, and so fermented, the *oleous* parts are thereby unlocked, and becoming rampant, over the other *Principles*, altogether produce that Taste. And the *Bile* it self, which, next to *Water* and *Earth*, consisteth most of *oily* parts, and of many both *saline* and *acid* is a bitter-sweet. Wherein, as some of the *saline* and *Acid* parts, smoothed by the *Oleous*, produce a *Sweet*: So, some of the *Oleous*, impregnated with the *saline*, and the *Acid*, doe hereby produce a *Bitter*.

THE
ANATOMY
OF
TRUNKS,

With an Account of their
VEGETATION.

Grounded thereupon.

The Figures hereunto belonging, Presented to the ROYAL
SOCIETY in the Years, 1673 & 1674.

The THIRD BOOK.

By NEHEMIAH GREW M.D. Fellow of the
Royal Society, and of the College of Physicians.

The Second Edition.

LONDON,

Printed by *W. Rawlins*, 1682.

T

THE UNIVERSITY OF CHICAGO

PHYSICS DEPARTMENT

RESEARCH REPORT

NO. 1234

BY

J. D. JONES

1955

PHYSICS DEPARTMENT

CHICAGO, ILL.

The Epistle Dedicatory.

ment: which must needs be Candid and Benign,
because it is Great. I have only this to say,

—— Ἐς Τροίαν περιώμενοι ἦνθον Ἀχαιοί;

Your Lordship will not disapprove the Enterprize,
although it falls short of perfection. It being the
result of *Your Lordships* manifold *Virtues* and *Abi-*
lities, That *You* know how far to Encourage the
meanest Attempts; as well as rightly to Value
and Assist the greatest Performances.

I am,

My Lord,

Your Lordships

most humble

and

most obsequious

Servant

London,
August 20.
1675.

NEHEMIAH GREW.

THE

THE
CONTENTS.

The FIRST PART.

CHAP. I.

A Description of several Stalks or Trunks, as they appear to the Naked Eye.

OF the stalk of Mize, §. 1, 2. Of Dandelion, 3, to 6. Of Borage, 7, to 10. Of Colewort, 11, to 16. Of Holyoak, 17, to 20. Of Wild Cucumer, 21, to 23. Of Scorzonera, 24, to 26. Of Burdock, 27, to 29. Of Endive, 30, 31. Of Vine, 32, to 35. Of Sumach, 36, to 38. Cautions to be had in observing the Parts, 39. Some Particulars better observed in cutting by the length, 40, 41.

CHAP. II.

Of the Barque, as it appears through a good Microscope.

First, a General Description of the several Parts of the Barque, 2, to 9. Next, a Particular Description of the Barques of 8 several Trunks; sc. Of Holly, Hazel, Barbary, Apple, Pear, Plum, Elm, Ash; The Vessels of all whose Barques are Lymphæducts: and those of two kinds, 10, to 13. Of 3 more, sc. Walnut, Fig, and Pine: the Vessels of the Barques of the Two first, being Lymphæducts and Lactiferous, Of the next, Lymphæducts and Resiniferous, 14, to 20. Of 3 more, sc. Oak, Common Sumach, and Common Wormwood; the Vessels of whose Barques are of 3 Kinds, 21, to 29. Some further Observations and Conjectures of the Sap-Vessels, 30, to 37.

CHAP. III.

Of the Wood.

WHat in all Trunks, §. 1. A Description of its Parts, in the several Trunks aforesaid. Of the Parenchymous Part, or Insertions, 2, to 9. Of the true Wood, 10, to 15. Of the Aer-Vessels, 16, to 26. Some further Observations and Conjectures of their Form, 27. Texture, 28, to 32. Nature, 33, 34. And Original, 35.

CHAP.

The Contents.

CHAP. IV.

Of the Pith.

A Description of the Pith, in General, §. 1. In the several Trunks or Brances aforesaid. As of the Size, 2, 3. Vessels, 4. Parenchyma and Bladders, 5, to 9. Apertures or Rupturss, 10. Some further Observations of the Pith. And of all the Pithy and Parenchymous Parts. And thence of the True Texture of a Plant, 11, to 15.

The SECOND PART.

CHAP. I.

Of the Motion and Course of the Sap.

CHAP. II.

Of the Motion and Course of the Aer.

CHAP. III.

Of the structure of the Parts.

CHAP. IV.

Of the Generation of Liquors.

CHAP. V.

Of the Figuration of Trunks.

CHAP. VI.

Of the Motions of Trunks.

CHAP. VII.

Of the Nature of Trunks, as variously fitted for Mechanical Use.

THE
ANATOMY
OF
TRUNKS;

PROSECUTED
With the bare EYE,
And with the
MICROSCOPE.

PART I.

CHAP. I.

The Descriptions of several Trunks, as they appear to the bare Eye.



At the end we may clearly understand, what the *Trunk, Stalk, or Branch* of a *Plant*, is; I shall by these *Figures* here before us, Describe the several *Parts*, whereof it is compounded.

1. §. And for examples sake, I shall in the first place, Describe the *Trunks* of some *Plants*, as being cut tranversly, and accurately observ'd, they appear to the naked Eye. And some others, as by the length. Which having done, I shall next proceed to a more particular Description of divers other *Trunks* and *Branches* as they appear through a good *Microscope*. In both shewing, not only what their several *Parts* are, as generally belonging to a *Branch*; but also, by a Comparative Prospect, in what respects they are *specifically* distinguished one from another, in the several *Sorts* of *Branches*.

2. §. I SHALL begin where the Work of Nature appears less Diversify'd: as in the *Stalk* of *Maze* or *Indian Wheat*. In which, although there are the same *Parenchymous* and *Lignous Parts*, as in all other *Plants*; yet is there neither *Barque*, nor *Pith*; the *Vessels* being dispersed and mixed with the *Parenchyma*, from the *Circumference* to the *Centre* of the *stalk*: Saving, that in and next the *Skin*, there seems to be no *Aer-Vessels*. Every where else, they run up, like fine *Threds*, through the length of the *stalk*: Each *Thred* being also surrounded with *sap Vessels*; which in a *Slice* cut transversly, appear in very small and dark colour'd *Rings*. The like structure may also be seen in the *Sugar-Cane*, and some other *Plants*.
- Tab. 18. 3. §. LET the next *Trunk* be that of *Taraxacum*, or *Dandelion*. In a *slice* whereof, being cut transversly, is seen next the *skin*, first, a simple, white, and close *Parenchyma* or *Barque*; made up of *Vesicles*; but such as are exceeding small; and hardly visible without a *Glass*.
- Tab. 18. 4. §. Within This, stand *Milk-Vessels* in seven or eight distinct *Columns*, of different size: each *Column* being also made up of seven or eight *Arched Lines*. Betwixt these *Columns*, run as many *Diametral Portions*, derived from the *Barque*, into or towards the *Pith*.
5. §. Next within These, stand the *Aer-Vessels*. Which are likewise divided, by the said *Diametral Portions*, into divers *Arched Lines*. The size of these *Vessels*, as well as their number, is small.
6. §. Within These, stands the *Pith*, consisting of very small *Vesicles* or *Bladders*, as the *Barque*. 'Tis very small, the *Diameter* hereof, being scarce one fifth, of that of the *Pith* of *Borage*. But the *Barque* of *Borage* is not half so thick as this of *Dandelion*.
7. §. FOR a Third *stalk*; we may take that of *Borage*; wherein there is some further Variety. For in a *slice* hereof, cut transversly, there appears, first a Tough, yet Thin and Transparent *Skin*. Within this *Skin*, and *Continuous* therewith, there is also a Thin *Ring* of *sap-Vessels*: which, without being crushed in the least, do yeild a *Lympha*.
- Tab. 18. 8. §. Next standeth the *Parenchyma* of the *Barque*. Which is made up of a great number of very small *Vesicles* or *Bladders*. Upon the inner *Verge* of this *Parenchyma*, standeth another *Ring* of *Sap-Vessels*: which also yeild a *Lympha*; and that different, as is probable, from the *Lympha* in the utmost *Ring*. Hitherto goes the *Barque*.
9. §. Adjacent to the *Ring* of *Sap-Vessels*, on the inner *Verge* of *Barque*, stand the *Aer-Vessels* on the outer *Verge* of the *Pith*. Not in a *Ring*; but in several *Parcels*; some *Parcels* or *Conjugations*, in the figure of little *Specks*; others, in little *Arched Lines*, almost like an *V Consonant*. And being viewed in a good *Glass*, there appears to be within the compass of every larger *speck* or *Parcel*, about 20 or 30 *Aer-Vessels* and within the smallest, about 8 or 10.
10. §. The *Pith*, in a well grown *stalk* of this *Plant*; is always hollow. But originally, it is entire. It is likewise wholly made up of a great number of *Vesicles*: of which, through a *Glass*, some appear *Pentangular*, others *Sexangular*, and *Septangular*. Most of them are larger than those of the *Barque*; so as to be plainly visible to a naked Eye.
11. §. A FOURTH *Trunk*, shall be that of *Colewort*, which seems likewise, to have at least, two Sorts of *Lymphaducts*. For being cut transversly, as the former, we may observe, next the *Skin*, a very

very close *Parenchyma*, of a darkish Green. Wherewith are mixed some few *Sap Vessels*, which give it that *Colour*.

12. §. Within This, stands a *scalloped Parenchymous Ring*, or a *Ring* of many short and slender white *Arches*. Which all round about *Tab. 18.* the *Barque*, meeting together, run in so many white *Diametral Portions*, or extream small *Rays*, into the *Pith*.

13. §. Betwixt these white *Rays*, and next of all to the said white *Arches*, stand as many small *Parcels* of *Sap-Vessels*, like so many little *Half-Ovals*. Within each of which, is included a white *Parenchyma*.

14. §. On the inner *Verge* of the *Barque*, stands another Sort of *Sap-Vessels*, in one slender and entire *Ring*. And so far goes the *Barque*.

15. §. Next within this *Ring* stand the *Aer-Vessels*, in several *Parcels*, diametrically opposite to the said white *Parenchymous Parcels* next without the *Sap-Ring*.

16. §. Last of all, and more within the *Pith*, stand the same kind of *Sap-Vessels*, as those of the *Half-Ovals*. Both these, by small lines, run one into another; thus, on both sides, hemming in the *Aer-Vessels*, and so making altogether, so many little *Pyramids*.

17. §. LET a *Fifth* be that of *Holyoake*. In which, the *Curiosity* of *Nature*, is still more copious: presenting us, as it is seems, with Three sorts of *Lymphæducts*; Of which, two yield a *Thin*; the Third, a *Thick Lympha*. For being cut, as before, next to the *Skin*, stands the *Barque*; somewhat close, and, in proportion, *Thick*.

18. §. Towards the inner *Verge* hereof, stand one sort of *Sap-Vessels*, postur'd in short *Rays*. These *Vessels* yield a *Mucilage*. And on the inner *Verge* of the *Barque*, stands a *Thin Ring* of other *Sap-Vessels*, which yield a *thinner Liquor*. *Tab. 18.*

19. §. Next within the *Barque* stand the *Aer-Vessels*, postur'd likewise in short *Rays*, diametrically opposite to those in the *Barque*. In every *Ray*, there are about twelve or sixteen *Vessels*.

20. §. Lastly, and more within the *Pith*, there stand other *Sap-Vessels*, all in very *Thin* or *Slender Arched-Lines*; thus hemming in the *Parcels* of *Aer-Vessels*.

21. §. FOR a *Sixth*, I will take that of *Wild Cucumber*: Wherein is also found a *Mucilaginous Lympha*. For first of all, next to the *Skin*, there is a *Ring* of *Sap-Vessels*. Which *Ring* is also radiated, the *Rays*, all poynting towards, and most of them terminating on, the *Skin*.

22. §. Next of all, there is a thick, and simple *Parenchymous Ring*. *Tab. 18.* On the inner *Verge* whereof, there are other *Sap-Vessels* standing in *Parcels*, also in a *Ring*. So far goes the *Barque*.

23. §. Next within, stand the *Aer-Vessels*, in as many *Parcels*, contiguous to those of the *Sap-Vessels* aforesaid. To which likewise are adjoyned as many more *Parcels* of *Sap-Vessels* within the *Pith*, opposite to the said *Sap-Vessels* within the *Barque*.

24. §. FOR a *Seventh*, we may, choose that of *Scorzonera*. In which, the *Vessels* are both *Lymphæducts*, and *Lactiferous*. All of them, with the *Aer-Vessels*, in a radiated posture. For first next the outer *Edg* of the *Barque*, stand the *Lactiferous*, in little *Specks*. Next to *Tab. 18.* these, on the inner *Edg* of the *Barque*, stand the *Lymphæducts*, in the same form.

25. §. Hereunto adjacent, on the outer Edg of the *Pith*, stand the *Aer-Vessels*, some in *Specks*, and some in extream short *Lines*; hardly distinguished, without a very nice Inspection.

26. §. Within These, are placed other *Lymphæducts*, opposite to those in the *Barque*. And within these *Lymphæducts*, still in the same radiated Line, run more of the *Milk-Vessels*.

27. §. AN EIGHTH, may be that of *Furdock*; Wherein first, there are a Sort of *Lymphæducts*, which stand in *Arched Parcels*, round the *Trunk*, adjacent to the *Skin*.

28. §. Within these, about the middle of the *Barque*, run the *Milk-Vessels*, in the form of small round *Specks*.

29. §. Next to these on the inner Edg of the *Barque* are placed other *Lymphæducts*. Which, together with more of the same in the *Pith*, and the *Aer-Vessels* betwixt them, stand all in Radiated Lines, of several Lengths, and all sharpening towards the Centre.

30. §. LET the *Ninth*, be that of *Endive*: In which there is also much curious Work. Next to the *Skin*, there is, first, a thick and simple *Parenchyma*. Then there is a kind of *Undulated Ring* of *Milk-Vessels*. Within which stand a Sort of *Lymphæducts*, in several *Parcels*; some, in *Arched Half-Ovals*; others, in short slender *Rays*. Betwixt these *Parcels*, many of the *Milk-Vessels* likewise stand.

31. §. Next there is an *undulated Ring* of other *Lymphæducts*, parting as in most *Trunks*, betwixt the *Barque* and the *Pith*. Within which, are the *Aer-Vessels*. And within These, more *Sap-Vessels*. Both of them, in small *Specks*, answerable, or opposite to the *Rays* in the *Barque*.

32. §. I SHALL give also one or two Examples of *Trees*, or *Arborescent Plants*; the *Vine* and *Common Sumach*. In a Slice of the former cut transversly, next the *Skin*, there is a *Thin Barque*. In the inner part whereof, adjacent to the *Wood*, stand the *Lymphæducts* in several *Half-Oval Parcels*, opposite to so many Radiated Pieces of the *Wood*.

33. §. The *Wood* is divided into the said Pieces, by as many *Parenchymous Rays*, inserted from the *Barque*, and so continuous therewith.

34. §. Within these Radiated Pieces of *Wood*, stand the *Aer-Vessels*; the largest of which, especially if held up against the light, are plainly visible to the bare Eye.

35. §. Within the hollow of the *Wood*, stands the *Pith*; in the young Growths always large. In the utmost *Verge* whereof, adjacent to the *Wood*, stand a few more *Sap-Vessels* of the same Sort with those in the *Barque*.

36. §. IN A like Slice of *Common Sumach*, contiguous to the hairy *Skin*, there is a *Ring* of *Lymphæducts*. Next to this a Simple *Parenchyma*. Then several *Arched Parcels* of *Lymphæducts*. Within these, a *Ring* of *Milk-Vessels*. And then a *Ring* of other *Lymphæducts*. Thus far the *Barque*.

37. §. Within the *Barque*, stands the *Wood*, divided into several *Portions*, by the *Diametral Insertions* divided from the *Barque*. In the Body of the *Wood*, stand the *Aer-Vessels*, very much smaller than in the *Vine*.

38. §. The hollow of the *Wood* is filled up with the *Pith*. In the Circumference of which, stands a *Ring of Lymphædētus*; of the same sort with those next to the *Wood* without.

39. §. All the *Parts* of these *Trunks*, may, as I have now described them, be observed without a *Microscope*: excepting the *Bladders* and number of *Aer-Vessels*. Yet Three things are hereunto necessary; viz. a good *Eye*, a clear *Light*, and a *Razor*, or very keen *Knife*, wherewith to cut them with a smooth surface, and so, as not to Dislocate the *Parts*.

40. §. UPON *Inspection* also by the length, there are some particulars, common, more or less, to most *Plants*, yet better observable in some, than in others. As first, the *Reticulation* of the *Vessels*, (formerly described) not only in the *Wood*, but in the *Barque*: which is evident in a young *Branch* of *Corin*, upon the very Surface thereof, when some of the *Vessels* begin to be cast off into the *Skin*. And so, by stripping off the *Skin*, upon the Surface of the *Wood*. Tab. 19.

41. §. In cutting by the length, as well as transversly, the young *Fibres*, which grow within the *Wood* in the Edg of the *Pith*, are also seen. As likewise the manner of the Derivation of the *Parts* of the *Bud* from the *Branch* or *Stalk*; as in *Sonchus*. There are also many Varieties in the *Pith*, such as those hereafter mentioned (a) which fall under observation only in cutting by the length. (a) Chap. 4.

C H A P. II.

Of the Barque, as it appears through a good Microscope.



NOW proceed to a more particular *Description* of several *Trunks* and *Branches*, as they appear through good *Glasses*.

1. §. Now the *Trunk*, or *Branch* of every *Tree*, hath Three General *Parts* to be described; sc. the *Barque*, the *Wood*, and the *Pith*. That likewise of every *Herbaceous Plant*, hath either the same Three *Parts*; or else Three *Parts* Analogous; sc. the *Cortical*, the *Lignous*; and the *Pithy Parts*.

2. §. The *Barque* consisteth of two *Parts*, sc. the outmost or *Skin*, and the *Main Body*. The *skin* is generally composed, in part, of very small *Vesicles* or *Bladders*, cluster'd together. That is, originally it is so; but as the *Plant* grows, the *Skin* dries, and the said *Bladders*, do very much shrink up and disappear.

3. §. Amongst these *Bladders* of the *Skin*, there are usually intermixed a sort of *Lignous Fibres*, or *Vessels*, which run through the length of the *skin*; as in *Mallow*, *Nettle*, *Borage*, *Thistle*, and most *Herbs*. Which is argued not only from the Toughness of the *skin* by means of the said *Vessels*; but in some *Plants*, may be plainly seen, as in *Teasle*. In which, Tab. 20.

the several *Fibres*, which run by the length of the *Stalk*, are also conjoined by other smaller ones, which stand transversely.

4. §. Whether they are *Aer-Vessels*, or *Sap-Vessels*, is dubious. For, on the one hand, because they emit no *Sap*, or *bleed* not, and also stand adjacent to the *Aer*; 'tis probable that they are *Aer-Vessels*. On the other hand, they may be *Sap-Vessels*; notwithstanding that they *bleed* not: Because the non-emission of *Sap* is not an infallible and concluding argument of an *Aer-Vessel*. For there are some *Plants* which *bleed* not. Which yet are furnished with *Sap-Vessels*, as certainly as any others

(a) B. 2. which bleed. (a)

P. I. c. 3. §. 22. 5. §. The *Skin* of the *Trunk* is sometimes visibly porous. But no where more, than in the better sort of walking *Canes*; where the *Pores* are so big, as to be visible even to the naked *Eye*: like to those, which are observable in several parts of the *Ball* of the *Hand*, and upon the ends of the *Fingers* and *Toes*.

Tab. 20.

6. §. THE *Main Body* of the *Barque* consisteth likewise of two *Parts*, sc. *Parenchyma*, and *Vessels*. The *Parenchyma* is made up of an innumerable company of small *Bladders* cluster'd together. Differing in nothing from those aforesaid in the *Skin*; saving, that they are much larger; and generally rounder.

Tab. 22. & Sequent.

7. §. This *Parenchyma* of the *Barque* is the same, as to its *Substance*, both in the *Root* and *Trunk*. Yet as to the *Texture* of its *Parts*, in the one, and in the other, there is This observable difference, viz. That in the *Barque* of the *Root*, cut transversely, the said *Parenchyma* (as hath been shew'd) is usually, more or less, disposed into *Diametral Rays*; running through the *Barque*, after the same manner, as do the *Hour-Lines* through the *Margin* of the *Dial-plate* of a *Clock* or *Watch*: as in *Marsh-Mallow*, *Lovage*, *Melilot*, and others. Whereas here in the *Barque* of the *Trunk*, the said *Parenchyma* is rarely thus disposed into *Diametral Rays*: Nor when it is, are those *Rays* continued to the *Circumference* of the *Barque*; as in the *Barque* of the *Root* they frequently are. So in *Rhus* or *Sumach*, although part of the *Parenchyma* be dispos'd into *Diametral Rays*: yet are those *Rays* extended not half way through the *Barque*. So also in *Fig-tree*, *Worm-wood*, *Thistle*, and others. What is further observable in the *Texture* of the *Parenchyma*, I shall shew in the description of the *Pith*.

Tab. 7, 8, 9.

Tab. 22. & Seq.

Tab. 31, 34, &c.

8. §. THE *Vessels* of the *Barque*, are, as I shall also shew, diversified many ways. But there are some Things, wherein, in all Sorts of *Plants*, they agree. *First*, in standing, most numerously, in or near, the inner *Margin* of the *Barque*. *Secondly*, in being always, and only *Sap-Vessels*. I have viewed so many, that at least, I can securely affirm thus much, That if there be any *Heteroclital Plants*, wherein they are found otherwise, there is not *One*, in *Five Hundred*. *Thirdly*, in being always *Conjugated* or *Braced* together in the form of *Net-work*. Although the *Number* and *Distances* of the *Braces*, are very different: as I have already shewed in the *Anatomy* of *Roots*.

Chap. 3. Tab. 6.

9. §. THE *Properties*, whereby the said *Vessels* of the *Barque* are specified and distinguished one from another, both in the same *Plant*, and in the several *Species* of *Plants*, are very many. Which *Properties*, are not *Accidental*, but such as shew the *Constant* and *Universal Design* of *Nature*. All which shall be demonstrated by the *Description* of several *Quarters* of the *Slices*, of so many *Kinds* of *Branches*

Branches, cut Transversly : and by the several *Figures* which represent them.

10. §. FIRST then, for the *Eleven* first *Quarters*, the *Vessels* of the *Barque* are only of *Two Kinds*. And these, in the first *Eight*, seem to be *Roriferous* (described also in the *Anatomy of Roots*) (*a*) and those which are common to most, if not to all *Plants*, sc. the *Lymphæducts*. Yet in all the *Eight*, they are, in respect both of their *Proportion*, and *Position*, very different. So in *Hazel* and *Ash* they are but few. In *Holly* and *Barberry* more. In *Apple*, *Pear*, *Plum*, *Elm*, still more numerous. And of those three *Fruits*, in an *Apple*, or *Plum*, more than in a *Pear*.

11. §. Again, as their *Proportion*, so likewise their *Position* is divers. For in *Holly*, the *Lymphæducts* or inner *Vessels* next to the *Wood*, stand in *Rays*. Yet so numerous and close together, as to make one *Entire Ring*. In *Hazel*, they stand more in *Oblong Parcels*. In *Barberry*, they stand likewise in *Parcels*, but they are so many *Half-Ovals*. The utmost *Vessels* or *Roriferous* of all Three, make a *Ring*.

12. §. Again, in *Apple*, *Pear*, and *Plum*, the *Lymphæducts* are *Radiated*. The *Roriferous* are neither *Radiated*, nor make an *entire Ring*; but stand in *Peripheral Parcels*. Much after the same manner, they also stand in *Elm*. In *Ash*, the *Vessels* make *Two Rings*; but neither of them *Radiated*: the inmost *Ring* or *Lymphæducts*, consisting of *Arched Parcels*, and the utmost or *Roriferous Vessels*, of *Round ones*. And whereas in all the foregoing, the *Lymphæducts* are still contiguous to the *Wood*; and the *Roriferous* more or less, distant from the *Skin*: here, on the contrary, the former are distant from the *Wood*, and the latter contiguous to the *Skin*.

13. §. And that these *Vessels* in each *Barque* of the said *Eight Branches*, are of *Two distinct Kinds*, seems evident, as from some other reasons, so from hence; In that their *Positions* are altogether *Heterogeneous*: Yet in both *Constant*, *Regular* and *Uniform*. I say, there seems to be no Reason, why the self *same Kind* or *Species* of *Vessels*, should have a different, yea a contrary *Position* in one and the same *Plant*; and that *Contrariety*, not *Accidental*, but *Regular* and *Constant*.

14. §. FOR the *Three* next *Quarters* sc. the *Ninth*, *Tenth* and *Eleventh*, the *Vessels* of the *Barque* are also different in *Number*, *Position*, *Size* and *Kind*. In *Pine*, which is the *Eleventh*, they are fewer. In *Walnut* the *Ninth*, more. In *Fig*, the *Tenth*, most numerous.

15. §. So for their *Position*. In *Pine*, the inmost make a *Radiated Ring*. The utmost stand *stragling* up and down, without any certain order. In *Wallnut* the inmost make also a *Radiated Ring*; The utmost make a *Double Ring*; not *Radiated*, but of *Round Parcels*. In *Fig*, the inmost make also a *Radiated Ring*. But the utmost make a *Double* and sometimes *Treble Ring*, not of *Radiated*, nor *Round*, but *Arched Parcels*.

16. §. Thirdly, they are also different in *Kind*. Those, I think, of the two former, *Wallnut* and *Fig*, are thus different: those certainly, of the *Fig*, are so; being *Lymphæducts* and *Lacteals*. The *Lymphæducts* make the inmost *Radiated Ring*. The outmost which make the other *Rings* in *Arched Parcels*, are the *Lactifers*.

17. §. That they are distinct *Kinds* of *Vessels*, is evident for two Reasons. First, from their *Position* in the *Barque*; which is altogether

ther different, as hath been said. *Secondly*, from the most apparent Diversity of the *Liquors* or *Saps*, which they contain, and which, upon cutting the *Branch* transversely, do distinctly *Bleed* from them. Which is one way, whereby we do distinguish the *Vessels* of *Animals* themselves. As in the *Liver*, it were hard to say, which is a *Blood-Vessel*, and which is a *Bile-Vessel*, where they are very small, if it were not for the *Contents* of them both.

18. §. Those in the *Barque* of *Pine*, are likewise of *Two Kinds*. The inmost are *Lymphæducts*, as in the two former. The utmost are not *Milk-Vessels*, but *Gum-Vessels*, or *Resiniferous*; which stand straggling, and singly, about the middle of the *Barque*. Out of these *Vessels* all the clear *Turpentine*, that drops from the *Tree*, doth issue.

19. §. Few, but very great. So that besides the difference of their *Number* and *Position*, and of the *Liquors* which they contain, and *Bleed*; there is yet a *Fourth*, and that is, their *Size*. Most of these *Turpentine Vessels*, being of so wide a *bore*, as to be apparent to the naked *Eye*: and, through a good *Glass*, above $\frac{1}{3}$ ^d of an *Inch* in *Diameter*. Whereas that of the *Lymphæducts*, can hardly be discovered by the best *Microscope*.

20. §. The same *Turpentine-Vessels* of *Pine*, are likewise remarkably bigger, not only than the *Lymphæducts*, but many times, than the *Milk-Vessels* themselves: as those of the *Fig*, which, in comparison, are exceeding small; every *Arch*, not being a *single Vessel*, but a *Parcel* or *Cluster* of *Vessels*; Whereas one *single Gum-Vessel* in *Pine*, is sometimes as big as two whole *Arched Clusters*, that is, as some *Scores* of the *Milk Vessels* in a *Fig-tree*. And the said *Gum-Vessels* of *Pine*, being compared with the *Lymphæducts* of the same *Tree*, one *Gum-Vessel*, by a moderate estimate, may be reckoned *three* or *four hundred* times *wider* than a *Lymphæduct*. The like prodigious difference may be observed in the *Size* of the several *Kinds* of *Vessels* of many other *Plants*.

21. §. THE *Three* next *Quarters* of *Branches*, are of *Oak*, *Common Sumach*, and *Common Wormwood*. In the *Barque-Vessels* whereof, there is observable some farther *Variety*. For in all or in most of the above named, there are only *Two Kinds* of *Vessels* in the *Barque*. But in *Each* of these, there are, at least, *Three Kinds*.

22. §. And first, in that of *Oak* there are *Lymphæducts*, *Roriferous*, and a Sort of *Resiniferous*. The inmost or *Lymphæducts*, make a *Radiated Ring*, contiguous to the *Wood*. The utmost or the *Roriferous* make also a *Ring*, but not *Radiated*. Those which are a sort of *Rosin-Vessels*, stand in *Round Parcels*; the greater *Parcels* betwixt the *Two Rings* of *Roriferous* and *Lymphæducts*; and the lesser, betwixt the *Roriferous* and the *Skin*.

23. §. That these last are different *Vessels* from both the other, seems evident, from the difference of their *Position*, as aforesaid. And that they are a sort of *Resiniferous*, is argued from hence; In that, not only *Galls* are very full of *Rosin*, but that the *Barque* of *Oak* it self is also somewhat *Resinous*. For the conveyance of whose *Resinous* parts, it is most unlikely that any other *Vessels* should subserve, but a peculiar *Kind*; which may therefore be properly called *Resiniferous*.

24. §. The next is a *Branch* of *Common Sumach*. In the *Barque* whereof, there are likewise *Three Kinds* of *Vessels*. First of all, there is a thick *Radiated Ring* of *Lymphæducts*; standing on the inner *Margin*

gin of the *Barque*, contiguous with the *Wood*. These *Vessels* exhibit their *Lympha* very apparently. A second kind of *Vessels*, sc. *Roriferous*, are situate towards the outer *Margin* of the *Barque*, and are composed into distinct *Arched Parcels*, all standing in a *Ring*.

25. §. Betwixt these Two *Kinds* stand the *Milk-Vessels*. Every single *Milk-Vessel* being *empaled* or hemmed in with an *Arch* of *Roriferous*. The *Milk-Vessels* are extraordinary large, almost as the *Gum-Vessels* of *Pine*; so as distinctly to be observed without a *Microscope*; after they are evacuated of their *Milk*; and without difficulty will admit a *Virginal Wye*; being two or three hundred times as big as a *Lymphæduct*. Besides these Three sorts of *Vessels*, there is also a *Ring*, adjacent to the *Skin*; which seems to be another sort of *Roriferous*.

26. §. The Last, is a *Branch* of *Common Wormwood*. In the *Barque* whereof, there are likewise Three *Kinds* of *Vessels*. First of all, there is a thin *Radiated Ring* of *Lymphæducts*, contiguous with the *Wood* or on the inner *Margin* of the *Barque*. Yet the *Ring* is not entire, but made up of several *Parcels*; which are intercepted by as many *Parenchymous* inserted into the *Pith*. Tab. 35.

27. §. A Second Sort of *Vessels*, which seem to be *Roriferous*, are situate about the middle of the *Barque*: and are composed into *Arched Parcels*, which likewise stand all even in a *Ring*.

28. §. Beyond these *Arches*, and towards the outer *Margin* of the *Barque*, stand a Third Sort of *Vessels*. Different from the *Milk-Vessels* in *Sumach*, both as to their *Situation*, *Size* and *Content*. For in *Sumach*, the *Milk-Vessels* stand within the *Arched Lymphæducts*: whereas these in *Wormwood*, stand without them. Likewise, being the *Vessels* of an *Herb*, they are far less; sc. about the compass or width of a small *Wheat-straw*. Their *Content*, is not a *Milk*, but a *liquid*, most *Oleous* and *viscid Gum*. Or which, for its pleasant *Flavour* may be called an *Aromatick Balsom*. For it perfectly giveth whatever is in the *Smell* and *Taste* of *Wormwood*: being the *Essence* of the whole *Plant*, which nature treasureth up in these *Vessels*. So that they are, in all respects, analogous to the *Turpentine Vessels* in *Pine*. There are divers other *Herbs* and *Trees*, which in the like *Vessels*, contain a *Turpentine*, or rather *Aromatick Balsom*; as *Angelica*; *Helenium* and others; the *Vessels* being so very large, that they may be easily traced with a knife, in cutting by the length of a *Branch* or *Saltk*.

29. §. Whether in some *Plants*, there are not more *Sorts* of *Vessels*, in the *Barque*, than have been now mentioned, I cannot say: Though we have not much reason to doubt of it. Because we see, there is so great variety in the *Viscera* of *Animals*. For what the *Viscera* are in *Animals*; the *Vessels* themselves are in *Plants*.

30. §. CONCERNING the *Form* and *Texture* of the *Lymphæducts*, there are some things, which though they are best observed in the *Wood*, yet in regard I am now describing the said *Vessels*, I shall here therefore add. I have already said, and shewed, in the former *Books*, That the *Lignous* and *Towy Parts* of all *Plants*, are *Tubulary*. And that the *Lympha* is conveyed, by the length of a *Plant*, through an innumerable company of small *Tubes* or *Pipes*.

31. §. The *Question* may be yet further put: If the *Towy Parts* of the *Barque* are made of *Tubes*, What are these *Tubes* themselves made up of? I answer, That these *Tubes* or *Lymphæducts*, are not only themselves

Tab. 40.

themselves *Organical*; but their very *Sides* also, seem to be composed of other *Parts*, which are *Organical*, *sc.* of *Lignous* or *Tomy* *Fibres*. Which *Fibres*; standing close or contiguous in a round *Figure*, they make one *Tubulary Body*, which I call the *Lymphæduct* of a *Plant*. And it is probable, That these *Fibres* themselves, are also *Tubulary*. That is, that a *Lymphæduct*, is a small *Tube*, made up or composed of other, yet much smaller *Tubes*, set round together in a *Cylindrick Figure*. As if we should imagine a company of *Straws*, which are so many small *Pipes*, to be joyned and set round together, so as to make another greater *Pipe*, answerable to a *hollow Cane*. The *Cane*, I say; is as the *Lymphæduct*; and the *straws* are as the *Fibres* whereof it is composed. By which also appears, the admirable smallness of these *Fibres*. For there are some *Lymphæducts*, which may be reckoned fifty times smaller than a *Horse-Hair*. Allowing therefore but Twenty of the aforesaid *Fibres* to make a *Thred* so big as one *Lymphæduct*; then one of the said *Fibres*, must be a Thousand times smaller than a *Horse-Hair*. That these *Fibres*, whereof the *Lymphæducts* are made, are themselves made up of other *Fibres*, is not altogether improbable.

32. §. These *Fibres*, although parallel; yet are they not *coalescent*, but only contiguous; being contained together in a *Tubulary Figure*, by the *Westage* of the *Cortical Fibres*, as in *Chapter the Fourth* will better be understood.

33. §. The first notice I took of the *Composition* and *Texture* of these *Vessels*, so far as the best *Glasses* yet known, will admit; was in a very *white* and *clear* piece of *Ash-wood* torn, with some care, by the length of the *Tree*, and objected to a proper *Light*. They seem also sometimes discernable in some other *clear Woods*, as in very *white Fir*, &c. And having formerly demonstrated, that the *Lignous Part* of a *Plant*, is annually made or augmented out of the inner part of the *Barque*, wherein the *Lymphæducts* always stand: we may reasonably suppose the same *Lymphæducts* to have the like *Conformation* in the *Barque*, as in the *Wood*.

34. §. And I am the rather induced to believe, that I am not mistaken in this Description, upon these two Considerations. First, that herein the Analogy betwixt the *Vessels* of an *Animal* and a *Plant*, is the more clear and proper. For as the *Sanguineous Vessels* in an *Animal* are composed of a number of *Fibres*, set round, in a *Tubulary Figure*, together: so are these *Lymphæducts* of a *Plant*. Secondly, in that herein, there is a more genuine response betwixt these, and the other *Vessels* of a *Plant* it self; *sc.* the *Aer-Vessels*; which are made up of a certain number of *Round Fibres*, standing collaterally, or side to side as I have already observed in the *Anatomy* of *Roots*. So that it is the less strange, that the *Lymphæducts* should be made up of *Fibres*, since the *Aer-Vessels* are evidently so made. Only with this difference, that whereas in the *Aer-Vessels*, the *Fibres* are postured or continued *Spirally*: here, in the *Lymphæducts*, they stand and are continued only in *straight Lines*.

Tab. 20.

35. §. THE STRUCTURE of the *Lactiferous* and *Gum-Vessels*, which have a very ample *Bore*, is more apparent. And, by the best *Glasses* I have yet used, they seem to be made, chiefly, by the *Constipation* of the *Bladders* of the *Barque*. That is to say, That they are so many *Channels*,

Channels, not made or bounded by any walls or sides proper to themselves, as a *Quil* thrust into a *Cork*, and as the *Aer-Vessels* are in the *Wood*: but only by the *Bladders* of the *Parenchyma*; which are so postured and crouded up together, as to leave certain *Cilyndrick Spaces*, which are continued by the length of the *Barque*.

36. §. One difference betwixt the *Vessels* or *Channels* now describ'd, and the *Tubulary Hollows* and other *Apertures* in the *Pith*, is this; That these never exist originally with the *Pith*; but are so many *Ruptures* supervening to it in its Growth. Caused, partly, by the Stretch or Tenter it suffers from the Dilatation of the *Wood*: (a) and partly, the drying, and so the Shrinking up of its *Bladders*, and of the *Fibres* whereof they are composed. Whereas the said *Vessels* in the *Barque*, are many of them originally formed therewith. And those which are *post-nate*, not made by any *Rupture*, but only such a Disposition of the *Parenchymous Fibres*, and Constipation of the *Bladders*, as is thereunto convenient.

(a) B. I. c.
3. §. 22, &
23.

37. §. In paring the *Barque* of a *Branch* of *Pine*, *Sumach*, &c. they appear, neither *parallel*, nor any where *Inosculated*: but run, with some little obliquities, distinct one from another, through the length of the *Branch*: and so, we may believe, through the length of the *Tree*.

C H A P. III.

Of the W O O D.



THE next general *Part* of a *Branch*, is the *Wood*; which lyeth betwixt the *Barque* and the *Pith*. And this likewise evermore consisteth of Two *General Parts*, *sc.* of a *Parenchymous Part*, and that more properly called *Lignous*. The *Parenchymous Part* of the *Wood*, though much diversified, yet in the *Trunks* of all *Trees* whatsoever, hath this property, To be disposed into many *Rays*, or *Diametral Insertions*, running be-

twixt so many *Lignous Portions*, from the *Barque* to the *Pith*: as in any of the *Quarters* here before us may appear.

2. §. But these *Insertions* are much diversified, according to the several *Sorts* of *Plants*. So in *Barberry*, *Ash*, *Pine*, *Worm-wood*, they are less numerous. In *Elm*, *Wallnut*, *Fig*, *Sumach*, they are more. And in *Holly*, *Pear*, *Plum*, *Apple*, *Oak*, *Hazel*, are most numerous.

3. §. The same *Insertions*, in *Barberry*, *Wormwood*, and some in *Oak*, are very *Thick*. In *Pine*, *Fig*, *Ash*, of a middle *Size*. In *Pear*, *Holly*, and most of them in *Oak*, are exceeding *Small*. Again, in *Barberry*, *Elm*, *Ash*, *sumach*, *Fig*, they are of an *Equal size*. In *Holly*, *Hazel*, *Pear*, *Plum*, *Oak*, they are very *Unequal*: some of those in *Holly*, being *Four* or *Five* times thicker than the rest; in *Plum*, *Six* or *Seven* times; and in *Oak*, *Ten* times at least.

4. §. In some *Plants*, they are Equidistant; in others, not: in some, the Great ones are Equidistant; in others, the Lesser; in others, both; in some, neither. Which *Varieties* are not accidental; but constant to the *species* in which they are severally found.

Tab. 34. 5. §. They are not always visibly continued from the Circumference to the Centre of the *Wood*: but in some *Branches*, as of *Sumach*; and in most *Trunks* of many years growth, declining, in some places, under or over, from a Level, are thereby, upon a Transverse Section, in part cut away.

Tab. 28,
34, 35.
Tab. 32. 6. §. They have yet one more Diversity, which is, That in divers of the aforesaid *Branches*, they run not only through the *Wood*; but also shoot out beyond it, into some Part of the *Barque*, as in *Elm*, *Sumach*, *Wormwood*, &c. Whereas in *Pine*, and some of the rest they either keep not distinct from the other parts of the *Parenchyma* of the *Barque*; or are so small, as not to be distinguished there form.

Tab. 35. 7. §. The *Texture* likewise of these *Insertions* is somewhat various. For in *Wormwood*, and most *Herbs*, they are manifestly composed of small *Bladders*: differing in nothing from those of the *Barque* or *Pith*, saving, in their being much less. Yet in *Herbs*, they are much larger than they are in *Trees*. And in many *Trees*, as *Apple*, *Pear*, *Plum*, *Pine*, &c. they are either quite lost, or so squeezed and pressed together by the hard *Wood* standing on both sides, as to be almost undiscernable.

Tab. 36,
37. 8. §. So that although the *Parenchyma* of the *Barque* or *Pith*, and the *Insertions* in the *Wood*, are of the same *specifick* Nature or Substance: yet there is this difference betwixt them; That the *Fibres* of the former, are so Netted together, as to leave several round *Vacuities*; or to make a great many little *Bladders*, whereas, in the latter, they are usually so far crowded up, as to run (as when a Net is stretched out) like a *Skein* of *Parallel Threds*.

Tab. 40. 9. §. Of these *Insertions* in the *Wood*, it is futher observable, That they do not only run betwixt the *Lignous Portions*; but that many of their *Fibres* are likewise all along distributed to the several *Fibres*, of which the *Lignous Portions* consist, and are interwoven with them; both together thus making a piece of *Linsy-Woolsy Work*, or like many other *Manufactures* in which the *Warp* and the *Woof* are of different Sorts of *Stuff*: as in the end of the *Fourth Chapter* is further explained.

10. §. THE WOOD is likewise compounded of Two Sorts of *Bodies*; That which is strictly *Woody*; and the *Aer-Vessels* mixed herewith. The true *Wood* is nothing else but a mass of antiquated *Lymphæducts*, viz. those which were originally placed on the inner Margin of the *Barque*. For in that place, there grows, every year, a new *Ring* of *Lymphæducts*. Which losing its original softness by degrees, at the latter end of the year, is turned into a dry and hard *Ring* of perfect *Wood*.

11. §. So that every year, the *Barque* of a *Tree* is divided into Two Parts, and distributed two *contrary* ways. The outer Part falleth off towards the *Skin*; and at length becomes the *Skin* it self. In like manner, as hath been observed of the *Skin* of the *Root*. Or as the *Cuticula* in *Animals*, is but the efflorescence of the *Cutis*. I say, that the elder *Skin* of a *Tree*, is not originally made a *Skin*; but was once, some of the *midle* part of the *Barque* it self, which is annually cast off, and

and dryed into a *Skin*: even as the very *Skin* of an *Adder*, upon the gradual generation of a new one underneath, in time, becomes a *Slough*. The inmost portion of the *Barque*, is annually distributed and added to the *Wood*: the *Parenchymous Part* thereof making a new addition to the *Insertions* within the *Wood*; and the *Lymphæducts* a new addition to the *Lignous pieces* betwixt which the *Insertions* stand. So that a *Ring* of *Lymphæducts* in the *Barque* this year, will be a *Ring* of *Wood* the next; and so another *Ring* of *Lymphæducts*, and of *Wood*, successively, from year to year. So the *Table*, for an *Apple-Branch*, sheweth a quarter of a *Slice* of a *Branch* cut transversely, of Three years growth: That of *Barbery*, of Two; That of *Sumach*, of One only; That of *Elm*, of Five.

12. §. Hereby two things maybe the better noted. First, the difference betwixt the degrees of the *annual* growths of several *Trees*: three years growth in an *Oak*, being as thick as five in an *Elm*. Secondly, the difference betwixt the *Annual* growths of the *same Tree*; being not of a constant proportion, but varying in thickness, as it should seem, according to the *season* of the year: whereby it may appear, what *season*, or kind of year, doth most of all favour, the latitudinal growth, or the *thickening* of any *Tree*.

13. §. The *Lymphæducts* thus antiquated or turned into *Wood*, do rarely, if ever, *Bleed*: but only transmit a kind of *Dewy* or *Vaporous Sap*. And some of them, as in the *Heart* of some *Trees*, it is probable, That they transmit not any *Sap*, either in the form of a *Liquor*, or a *Vapour*: and so being gradually deprived of their *Watery Parts*, become the *Heart*.

14. §. There is this further variety in the *Wood*; represented in *Walnut*, *Fig* and *Oak*. That some certain parcels hereof, make either several small and white *Rings*, as in *Oak*; or else divers white and crooked *Parcels*, transverse to the *Insertions*, as in *Walnut* and *Fig*. For it seemeth, that, at least, in many *Trees*, some portion of all the *Kinds* of *Vessels* in the *Barque*, are not only annually distributed to the *Wood*, but do likewise therein retain the same, or somewhat like *Position*, which they originally had in the *Barque*. So that as all those bigger and darker *Portions* of the *Wood*, were originally, the *Radiated Lymphæducts* of the *Barque*: so the little white *Circles*, or *Parcels* of *Circles*, in the same *Wood*, were originally another sort of *Sap-Vessels* in the *Barque*, *sc.* those which have a circular *Position* therein.

15. §. In the *Branches* of *Fir*, *Pine*, and others of the same *Kindred*; there are some few *Turpentine-Vessels* scattered up and down the *Wood*; and represented by the larger *Black Spots*. Which *Vessels* are *eadem numero*, the self same, which did once appertain to the *Barque*; and do even here also in the *Wood*, contain and yield a liquid *Turpentine*. Only, being pinched up by the *Wood*, they are become much smaller *Pipes*.

16. §. THE *Aer-Vessels*, with the *Insertions*, and true *Wood*, altogether make up That, which is commonly called, The *Wood* of a *Tree*. The *Aer-Vessels* I so call, not in that they never contain any *Liquor*; but, because all the principal time of the growth of a *Plant*, when the *Vessels* of the *Barque* are filled with *Liquor*, these are filled only with a *Vegetable Aer*.

17. §. In almost all *Plants*, not one in some hundreds excepted, this is proper to the *Aer-Vessels*; To have a much more ample *Bore* or *Cavity*,

vity, than any other in the *Wood*. In the *Wood*, I say; for in the *Barque*, there are many *Sap-Vessels* bigger than the biggest *Aer-Vessels* that be.

18. §. The Varities hereof are very many; in respect both of their *Number*, *Size*, and *Position*; being, as to these, the same, in no two *Sorts of Plants* whatsoever. First in respect of their *Number*. So in *Hazel*, *Apple*, *Pear*, they are very numerous; but in different degrees: and are represented in the *Figures* already referred to, by all the black spots in the *Wood*. In *Holly*, *Plum*, *Barberry* somewhat numerous. In *Oak*, *Ash*, *Walnut* fewer. In *Pine*, and others of that *Kindred*, very few; *sc.* fewer than in any other kind of *Plant*.

Tab. 23,
25, 26.
22, 24, 27.
29, 30, 33.
32.

19. §. Secondly, in respect of their *Size*; which from the first or greatest, to the least, may be computed easily to about *Twenty Degrees*. Thus, many of those in *Elm*, *Ash*, *Wallnut*, *Fig*, *Oak*, are very large. In *Barberry*, *Plum*, not so large. In *Hazel*, *Sumach*, smaller. In *Holly*, *Pear*, of a still smaller *Size*. So that many of those in *Elm*, or *Oak*, are *Twenty times* bigger, than those in *Holly* or *Pear*.

Tab. 28,
33.
22, 26.

20. §. In an ordinary joynted *Cane*, they are so wide, that if you take one a yard, or a yard and $\frac{1}{2}$ long, and putting one end into a *Basin* of *Water*, you blow strongly at the other; your *Breath* will immediately pass, through the *Aer-Vessels*, the length of the *Cane*, so as to raise up the *Water* into a great many *Bubbles*.

21. §. And as they have a different *Size* in divers *Kinds of Plants*; so likewise, according to the place where they stand, in the self same. So in *Holly*, *Hazel*, *Apple*, their *Size* is more equal throughout the breadth of the *Tree*. But in *Barberry*, *Elm*, *Oak*, *Ash*, very different: Not fortuitously, but always much after the same manner. For in all the last named *Branches*, the *Aer-Vessels* that stand in the inner *margin* of each annual *Ring*, are all vastly bigger, than any of those that stand in the outer part of the *Ring*.

Tab. 22,
23, 25.
24, 28, 29.
33.

22. §. Thirdly, these *Aer-Vessels* are also different in their *Situation*. So in *Apple*, *Wallnut*, *Fig*, they are spread all abroad in every annual *Ring*; not being posited in any one certain *Line*. In others, they keep more within the compass of some *Line* or *Lines*; either *Diametral*, or *Peripheral*. So in *Holly* they are *Radiated*, or run in even *Diametral Lines* betwixt the *Pith* and the *Barque*. So also are some of of them in *Hazel*; and some few in *Wallnut*.

Tab. 25,
30, 31.

Tab. 22.
23, 32.

23. §. Whether they stand *Irregularly*, or are *Radiated*, it is to be noted, That *Nature*, for the most part, so disposeth of them, that many of them may still stand very near the *Insertions*. So in *Apple*, she will rather decline making an even *Line*; or in *Holly*, will rather break that *Line* into *Parcels*, than that the *Aer-Vessels* shall stand remote from the *Insertions*. To what end this is done, shall be said hereafter.

24. §. Again, in *Ash*, the *Aer-Vessels* are none of them *Radiated*, but most of them stand in *Circles* on the inner *Margin* of every annual *Ring*. Which *Circle* is sometimes very thick, as in *Ash* and *Barberry*. In others but thin, the *Vessels* standing, for the most part, single throughout the *Circles*; as in *Elm*. Sometimes again, they both make a *Circle*, and are also spread abroad; as in *Pear* and *Plum*.

Tab. 24.
29.
28.
26, 27.

25. §. Those likewise which are spread abroad, are sometimes *Regularly* posited. So in *Barberry*, besides those larger, that make the *Circle*, there are other smaller ones, that stand, in oblique *Lines*, athwart

Tab. 24.

athwart one another; almost like a *Bend*, or sometimes, an entire or broken *Saltyr* in an *Escutcheon*. In *Oak*, they make rather certain *Columns*, in the posture of the *Pale*. And in *Elm*, they make, as it were, many *cross Parcels*, in the posture of the *Fess*. Tab. 33.
38.

26. §. This great difference in the *size* and *Position* of the *Aer-Vessels*, in the same individual *Plant*, is one ground, for which, I think it probable, That there are *divers Kinds* of *Aer-Vessels*; as well as of *Sap-Vessels*. Even as in *Animals*, there are *divers Kinds* of *Organs* for *Spiration*, and the separation of *Aer*: *Fishes* having their *Branchiæ*; *Land-Animals* their *Lungs*; and those in *Frogs*, &c. being of a somewhat peculiar *Kind*.

27. §. THE *Form* and *Texture* of these *Vessels*, and the various ways whereby they may be best observed, I have already described and shewed in my *Anatomy of Roots*. As to their *Form*, one thing P. i. c. 4. remarked was this; That they are never *Ramified*, but distinctly continued from one end of a *Plant*, small or great, to the other: as the *Nerves* are in *Animals*. A further and easie proof whereof, may be made, only by holding up a piece of an ordinary *Cane*, about $\frac{1}{2}$ a foot long, cut very smooth at both ends, against a full light: whereupon, if you keep it in a straight *Line* betwixt the *Light*, and the cast of your *Eye*, and then look steadily, you may see quite through it, that is, through the *Aer-Vessels*, which run straight along the *Cane* from end to end.

28. §. As to their *Texture*; whereas, oftentimes, the *Aer-Vessels* appear to be *unroaved* in the form of a very small *Plate*, it is to be noted, That it is not only of different breadth, in *divers Plants*, and usually much broader in the *Root*, than in the *Trunk*: but also, that in the *Trunk*, many times, the said *Vessels* are unroaved or resolved, not in the form of a *Plate*, but of a *Round-Thred*. The Causes of which Diversity, are principally Three; *viz.* The *Westage* of the *Fibres* of which the *Aer-Vessels* consist; The deference betwixt the said *Fibres*, or betwixt the *Warp* and the *Woof*; And the different *Kinds* of *Woof*.

29. §. By the *Westage* of the *Fibres*, it is, That the *Vessels*, oftentimes, *unroave* in the form of a *Plate*. As if we should imagine a piece of fine narrow *Ribband*, to be woun'd spirally, and Edg to Edg, round about a *Stick*; and so, the *Stick* being drawn out, the *Ribband* to be left in the *Figure* of a *Tube*, answerable to an *Aer-Vessel*. For that which, upon the *unroaving* of the *Vessel*, seems to be a *Plate*, Tab. 39. or one single *Piece*, is, as it were, a *Natural Ribband*, consisting of several *Pieces*, that is, a certain number of *Threds* or *Round-Fibres*, standing parallel, as the *Threds* do in an *Artificial Ribband*. And as in a *Ribband*, so here, the *Fibres* which make the *Warp*, and which are spirally continu'd; although they run parallel, yet are not coalescent; but contained together, by other *Transverse Fibres* in the place of a *Woof*.

30. §. And as the said *Fibres* are transversely continued, thereby making a *Warp* and *Woof*: So are they (as in *divers woven Manufactures*) of very different *Bulk*; those of the Former, being much bigger, and therefore much stronger, than those of the Latter. By which means, as *Cloth* or *Silk* will often Tear one way, and not another; so here, while the *Warp* or those *Fibres* which are spirally continued;

tinued, are usually *unroav'd* without breaking; those smaller ones, by which they are *stitched* or *woven* together, easily tear in sunder all the way.

31. And because the *Fibres* of the *Woof*, are themselves also of different *Bulk*; therefore it is, That where they are more sturdy, as usually in the *Root*, they require a greater quantity of *Warp*, that is, a broader *Plate*, to overmatch them. Whereas, where they are more extream small, as in the *Trunk* and *Leaves*, one *Thred* of the *Warp*, that is, one *Spiral Fibre*, will be strong enough of it self, and so, sometimes, be singly *unroav'd*.

32. §. From the extream Tenuity of these *Fibres*, it is, That they are very rarely discern'd, and not without the greatest difficulty. As also; from their great *Tenderness*; whereby not enduring to be drawn out, they all break off close to the Sides of the *Spiral ones*. In the *Pith*, the like *Transverse Fibres* are a little more visible: which first conducted Me to the notice of them here also.

33. §. All the *Fibres* of the *Aer-Vessels*, both the *Warp* and the *Woof*, are of the same Substantial Nature with the *Pith* and the other *Parenchymous Parts* of a *Plant*. From whence it is, That whereas the *Towy Parts* of a *Plant*, whereof all *Linen Manufactures* are made, are very *Strong* and *Tough*; these, as is abovesaid, are extream *Tender* and *Brittle*, like those of the *Pith* and all the *Pithy Parts*. To which therefore, the *Aer-Vessels* are to be referr'd. And the *Content* of both, is oftentimes the same.

34. §. From whence, we have a further proof of what I have formerly asserted, which is, That in all *Plants*, there are *Two* Substantially different *Parts*, and no more than *Two*, *viz.* the *Pithy*, and the *Towy* or *Lignous Parts*.

B. I. c. 7.
§. 13, 14.

35. §. From hence also we have some ground to conjecture, That so many of the *Aer-Vessels*, at least, which are not formed with the seed, but *post-nate*, are originated from the *Parenchymous Parts*; which seem by some alteration in the *Quality*, *Position* and *Texture* of the *Fibres*, to be Transformed into *Aer-Vessels*, as *Caterpillars* are into *Flies*. And as the *Pith* it self, by the Rupture and Shrinking up of several *Rows* of *Bladders*, doth oftentimes become *Tubulary*: So is it also probable, that in the other *Parenchymous Parts*, one single *Row* or *File* of *Bladders* evenly and perpendicularly piled; may sometimes, by the shrinking up of their *Horizontal Fibres*, all regularly break one into another and so make one *continued Cavity*; or a *Tube*, whose *Diametre* is the same with that of the *Bladders*, whereof it is composed. All which, will appear more probable, and what hath been said, be yet better understood, when we come, in the next *Chapter*, to the Description of the *Pith*.

C H A P. IV.

Of the P I T H.



THE Third General Part of a Branch is the *Pith*. Which though it have a different name from the *Parenchyma* in the *Barque*, and the *Insertions* in the *Wood*; yet, as to its *Substance*, it is the very same with them both. Whereof there is a double evidence, *sc.* their *Continuity*, and the sameness of their *Texture*. Their *Texture* shall be shewed presently.

As to their *Continuity*, it is to be noted, That as the *Skin* is continuous with the *Parenchyma* of the *Barque*; and this *Parenchyma* likewise, with the *Insertions* in the *Wood*; so these *Insertions* again, running through the *Wood*, are also continuous with the *Pith*. So that the *Skin*, *Parenchyma*, *Insertions*, and *Pith*, are all One entire piece of Work; being only filled up, in divers manners, with the *Vessels*.

2. §. The *Size* of the *Pith* is various, being not the same in any two *Branches* here represented. In *Wormwood*, *Sumach*, *Fig*, *Barbery*, 'tis very large; *sc.* betwixt 5, and 7 *Inches* *Diametre*, as it appears through the *Microscope*. In *Pine*, *Asb*, *Holly*, *Walnut*, not so large; from 3 *Inches* *Diameter* to 4. In *Oak*, *Apple*, *Pear*, *Hazel*, lesser, scarce from 2, to 3. In *Damascene*, not above an *Inch* and half. And in *Elm*, scarce an *Inch* *Diameter*. Note also, that of all *Plants*, both *Herbs*, and *Shrubs*, have generally the largest *Piths*, in proportion with the other *Parts* of the same *Branch*, as in *Sumach*, *Fig*, *Barbery*; is manifest.

Tab. 24,
31, 34, 35.
22, 29,
30, 32.
23, 25,
26, 35.
27.
28.

3. §. It is also worth the noting, That whereas, in most *Plants*, the *Barque* and *Wood* do both grow thicker every year: the *Pith*, on the contrary, groweth more slender; So that in a *Branch* of one years growth, it is apparently more ample, than in one of two; and in a *Branch* of two, than in one of three; and so on.

4. §. The *Pith*, for the most part, if not always, in the *Branch*, as well as the *Root*, is furnished with a certain number of *Sap-Vessels*. They are here usually so postur'd, as to make a *Ring* on the *Margin* of the *Pith*. Where they are more numerous, or large, they are more evident; as in *Walnut*, *Fig*, *Pine*, and others. They are also of divers *Kinds*, answerable to those in the *Barque*; as in *Wallnut*, *Lymphæducts*; in *Fig*, *Lactæals*; in *Pine*, *Resiniferous*.

Tab. 30,
31, 32.

5. §. The *Parenchyma* of the *Pith* is composed of *Bladders*. Which are the very same with those in the *Barque*, and oftentimes in the *Insertions* within the *Wood*. Only these in the *Pith*, are of the largest *Size*; those in the *Barque*, of a lesser; and those of the *Insertions* least of all: for which reason they are less obvious than in the *Pith*.

6. §. The *Bladders* of the *Pith*, though always comparatively Great; yet are of very different *Sizes*. Being easily distinguished, even as to their *Horizontal Area*, to *Twenty* Degrees. Those of *Fig*, *Barbery*, and some others, are somewhat large. And of many *Herbs*, as

Tab. 24,
31.
of

Tab. 39. of *Thistle*, *Borage*, and others, three times as big again; appearing in the *Microscope*, like to the largest *Cells* of an *Hony-comb*. Those of *Plum*, *Worm-wood*, *Sumach*, less. Of *Elm*, *Apple*, *Pear*, lesser. Of *Holly* and *Oak*, still less. So that the *Bladders* of the *Pith* in *Borage* or *Common Thistle*, are of that *Size*, as to contain, within the compass only of their *Horizontal Area*, about twenty *Bladders* of the *Pith* of *Oak*. Wherefore one whole *Bladder* in *Thistle*, is, at least an hundred times bigger, than another in *Oak*.

7. §. Of the *size* of these *Bladders* of the *Pith*, 'tis also to be noted, That it doth not at all follow the *Size* of the *Pith* it self; but is still varied, according as Nature designeth the *Pith* for various use. Thus, whereas the *Pith* of *Sumach*, is Larger than that of *Barberry*; it might be thought, that the *Bladders*, whereof it is composed, should be likewise Larger: Yet are they Three times as Small again in *Sumach*, as they are in *Barberry*. So the *Pith* of *Plum*, is far Less, than that of *Pear*; yet the *Bladders* of the former are Four or Five times as big, as those of the latter. So the *Pith* of *Hazel* is almost Three times as Little again, as that of *Holly*; yet the *Bladders* in *Hazel*, are Ten times bigger, than in *Holly*.

Tab. 24,
34.
26, 27.
22, 23.

8. §. The *Shape* of the *Bladders* hath also some Variety. For although, for the most part, they are more round; yet oftentimes they are angular: as in *Reed-grass*, a *Water-plant*; where they are *Cubical*; and in *Borage*, *Thistle*, and many others, where they are *pentangular*, *sexangular* and *septangular*.

9. §. Of the *Texture* of the *Bladders*, 'tis also to be noted, that many times, the *Sides* of the greater *Bladders* are composed of lesser; as is often seen in those of *Borage*, *Bulrush*, and some other *Plants*. In the same manner, as the *Sap-Vessels*, are but greater *Fibres* made up of lesser.

10. §. The *Pith*, though always originally composed of *Bladders*, and so *One Entire Piece*; yet in process, as the *Plant* grows up, it hath divers openings or *Ruptures* made in it: oftentimes very regularly, and always for good use, and with constancy observed in the same *Species* of *Plants*. In *Sharp-pointed Dock*, many of the *Pores* are considerably prolonged by the length, like small *Pipes*. In *Walnut* it shrinketh up into transverse *Filmes* or *Membranes*; as likewise sometimes in *Spanish-Broom*. Sometimes the *Pith* is hollow or *Tubulary*: either throughot the *Trunk*, as in *Thistle*, *Endive*, *Scorzoner*, *Marsh-Mallow*: or so, as to remain entire at every joynt; as in *Sonchus*, *Nettle*, *Teasle*; in which it is divided as it were into several *Stories*: and divers other ways.

Tab. 19.

11. §. I SHALL conclude this discourse with a further illustration of the *Texture* of the *Pith*, and of the whole *Plant*, as consequent thereupon. I say therefore, (and have given some account hereof in the *Anatomy* of *Roots*) That as the *Vessels* of a *Plant*, sc. the *Aer-Vessels* and the *Lympheducts* are made up of *Fibres*; according to what I have in this Discourse above said; so the *Pith* of a *Plant*, or the *Bladders* whereof the *Pith* consists are likewise made up of *Fibres*. Which is true also of the *Parenchyma* of the *Barque*. And also of the *Insertions* in the *Wood*. Yea, and of the *Fruit*, and all other *Parenchymous Parts* of a *Plant*. I say, that the very *Pulp* of an *Apple*, *Pear*, *Cucumber*, *Plum*, or any other *Fruit*, is nothing else but a *Ball* of most extream small transparent *Threds* or *Fibres*, all wrapped and stick'd

stitch'd up (though in divers manners) *together*. And even all those *Parts* of a *Plant*, which are neither formed into visible *Tubes*, nor into *Bladders*, are yet made up of *Fibres*. Which, though it be difficult to observe, in any of those *Parts* which are closer wrought and principally in the *Insertions* of some *Trees*: yet in the *Pith*, especially of some *Plants*, which consisteth of more open *work*, they are more visible. Which introduceth the observation of them in all other *Parenchymous Parts*. So in the *Pith* of a *Bulrush* of the *Common Thistle*, and some other *Plants*; not only the *Threds* of which the *Bladders*; but also the single *Fibres*, of which the *Threds* are composed; may sometimes with the help of a good *Glass*, be distinctly seen. Yet one of these *Fibres*, may reasonably be computed to be a *Thousand* times smaller than an *Horse-Hair*. Tab. 38.

12. §. The *Fibrosity* of the *Parenchyma* is also visible in some *Woods*, in which, it is apparently mixed with the *Lignous Parts*, not only by *Insertions*, but *per minimas Partes organicas*. That is to say, The *Parenchymous Fibres*, like smaller *Threds*, are either *wrapped round* about both the *Lignous* and the *Aer-Vessels*, or at least *interwoven* with them, and with every *Fiber* of every *Vessel*: as in very white *Ash* or *Fir-Wood*, with an advantageous posture and light, may be observed. Tab. 39.

13. §. WHENCE it follows, that the whole *Substance*, or all the *Parts* of a *Plant*, so far as *Organical*, they also consist of *Fibres*. Of all which *Fibres* those of the *Lymphæducts*, run only by the *Length* of the *Plant*: those of the *Pith*, *Insertions*, and *Parenchyma* of the *Barque*, run by the *breadth* or horizontally: those of the *Aer-Vessels*, fetch their *Circuit* by the *Breadth*, and continue it by the *Length*.

14. §. By which means, the said *Parenchymous Fibres*, in fetching their *horizontal Circles*, do thus *weave*, and make up the *Bladders* of the *Pith*, in *Open-Work*. And the same *Fibres* being thence continued; they also *weave* and make up the *Insertions*, but in *close-Work*. Betwixt which *Insertions*, the *Vessels* being likewise transversely interjected, some of the same *Fibres* *wrap* themselves also about these; thus *tying* many of them together, and so making those several *Conjugations* and *Braces* of the *Vessels*, which I have formerly described. And as some of these *Horizontal Fibres* are *wrapped* about the *Vessels*; so also about the *Fibres*, whereof the *Vessels* are composed. By which means it is, that all the *Fibres* of the *Vessels* are *Tacked* or *stitched* up close together into *One Coherent Piece*. Much after the same manner, as the *Perpendicular Splinters* or *Twigs* of a *Basket*, are, by those that run in and out *Horizontally*. And the same *Horizontal Fibres*, being still further produced into the *Barque*; they there compose the same *work* over again (only not so *open*) as in the *Pith*. Tab. 40.

15. §. SO THAT the most unfeigned and proper resemblance we can at present, make of the whole *Body* of a *Plant*, is, To a piece of *fine Bone-Lace*, when the *Women* are working it upon the *Cushion*. For the *Pith*, *Insertions*, and *Parenchyma* of the *Barque*, are all extream *Fine* and *Perfect Lace-Work*: the *Fibres* of the *Pith* running *Horizontally*, as do the *Threds* in a Piece of *Lace*; and bounding the several *Bladders* of the *Pith* and *Barque*, as the *Threds* do the several *Holes* of the *Lace*; and making up the *Insertions* without *Bladders*, or with very small ones, as the same *Threds* likewise do the *close* *Parts* of

the *Lace*, which they call the *Cloth-Work*. And lastly, both the *Lignous* and *Aer-Vessels*, stand all *Perpendicular*, and so cross to the *Horizontal Fibres* of all the said *Parenchymous Parts*; even as in a Piece of *Lace* upon the *Cushion*, the *Pins* do to the *Threds*. The *Pins* being also conceived to be *Tubular*, and prolonged to any length; and the same *Lace-Work* to be wrought many Thousands of times over and over again, to any thickness or height, according to the height of any *Plant*. And this is the true *Texture* of a *Plant*: and the *general composition*, not only of a *Branch*, but of all other *Parts* from the *Seed* to the *Seed*.

An Account of the
VEGETATION
 OF
TRUNKS
 Grounded upon the foregoing
ANATOMY.

PART II.



HAVING before given the *Anatomy of Trunks*; I shall next proceed to see, what *Use* may be made thereof; and principally, to explicate the manner of their *Vegetation*. In doing which, that former *Method*, which I used in shewing the manner of the *Growth of Roots*, I shall not exactly follow. For so, in regard the *Organical Parts* of the *Root* and *Trunk* are the same, and consequently their *Nutrition* and *Conformation* are effected in the same way; I should hereby be obliged to a nauseous and unprofitable repetition of many things already said. The Explication therefore of all those Particulars, which more especially belong to the *Trunk*, or are more Apparent therein, and not spoken of, or not so fully, in the former *Books*, will be my present Task. The chief *Heads* whereof, shall be these Seven following, *viz.*

- FIRST, the *Motion* and *Course* of the *Sap*.
- SECONDLY, The *Motion* and *Course* of the *Aer*.
- THIRDLY, The *Structure* of the *Parts*.
- FOURTHLY, The *Generation* of *Liquors*.
- FIFTHLY, The *Figuration* of *Trunks*.
- SIXTHLY, The *Motion* of *Trunks*.
- SEVENTHLY, And lastly the *Nature* of *Trunks* as variously fitted for *Mechanical Use*.

C H A P. I.

Of the Motion and Course of the Sap.



I R S T, as to the *Course* of the *Sap*, there are Three *Parts* in which it *moveth*; *sc.* the *Pith*, the *Wood*, and the *Barque* First the *Pith*; in which the *Sap* *moveth* the *First* year, and *only* the *First* year. Or, it is *Proprium quarto modo*, to the *Pith* of every *Annual Growth*, and to the *Pith* of such a *Growth* only, *To be succulent*. That is, whether of a *Sprout* from a *Seed*, or of a *Sucker* from a *Root*, or of a *Cyon* from a *Branch*; The *Pith* is always found the *First* year full of *Sap*. But the *Second* year, the same individual *Pith*, always becomes *dry*, and so it continues ever after.

2. §. One cause whereof is, that the *Lymphæducts* in the *Barque*, being the first year adjacent to the *Pith*; they do all that time, transfuse part of their *Sap* into it, and so keep it always *Succulent*. But the same *Lymphæducts*, the year following, are turned into *Wood*; and the *Vessels* which are then generated, and carry the *Sap*, stand beyond them, in the *Barque*. So that the *Sap* being now more remote from the *Pith*, and intercepted by the new *Wood*, it cannot be transfused, with that sufficient force and plenty as before, into the *Pith*; which therefore, from the first year, always continues dry.

3. §. THE SECOND *Part* in which the *Sap* moves, *sub forma liquoris*, is the *Wood*. Which yet, it doth not in all *Plants*, but only in some; and visibly, in very few; as in the *Vine*: In a *Vine*, I say, the *Sap* doth *visibly ascend* by the *Wood*. And this it doth, not only the first year, but every year, so long as the *Vine* continues to grow. But although this *ascent*, in or through the *Wood*, be every year; yet it is only in the *Spring*, for about the space of a Month; *sc.* in *March* and *April*.

4. §. There are many other *Trees*, besides the *Vine*, wherein, about the same time of the year, the *Sap* *ascendeth*, though not so copiously, yet chiefly, in the *Wood*. For if we take a *Branch* of two or three years growth, suppose of *sallow*, and having first cut the same transversely; if the *Barque* be then also transversely, and with some force, pressed with the back of the knife, near the newly cut end; the *Sap* will very plainly rise up out of the utmost *Ring* of *Wood*. And if it be pressed in the same manner, or a little more strongly, about an Inch lower, the *sap* will ascend out of every *Ring* of *Wood* to the Center. Yet at the same time, which is to be noted, there *ariseth* no *Sap* at all out of the *Barque*.

5. §. Whence appears the Error of that so Common Opinion, That the *Sap* always *riseth* betwixt the *Wood* and the *Barque*. The contrary whereunto is most true, That it never doth. For the greater part of the year, it riseth in the *Barque*, *sc.* in the inner *Margin* adjacent

cent to the *Wood*, and in *Spring*, in or through the *Wood* it self, and there only.

6. §. THE THIRD Part in which the *Sap* ascends, is the *Barque*, as was above hinted, and may be observ'd in almost any *Branch*, if cut cross, in the *late Spring* and in *Summer*; either as the *Sap* issueth spontaneously, or upon pressing, as aforesaid. So that when the *Sap* ceaseth to ascend, *sub forma liquoris*, by the *Wood*, then it begins to ascend by the *Barque*.

7. §. Besides the difference of *Time*, the *Organical Parts* likewise, in which these two *Saps* ascend, are divers. For in the *Barque*, it ascendeth *visibly*, only in the *Succiferous*, whereas in the *Wood*, it ascendeth only by the *Aer-Vessels*.

8. §. FROM what hath been said, we may understand, what is meant by the *Bleeding of Plants*. If we take it generally, it properly enough expresses, *The eruption of the Sap out of any Vessels*. And so, almost all *Plants*, in *Summer* time, do *Bleed*, that is, from *Sap-Vessels*, either in the *Barque*, or in the *Margin* of the *Pith*: the *Saps* they *Bleed*, having either a *Sower*, *Sweet*, *Hot*, *Bitter*, or other *Tast*. At which time, the *Vessels* also, in the *Barque* of a *Vine-Branch*, do *Bleed* a *Sower Sap*.

9. §. But that which is *vulgarly* called *Bleeding*, as in a *Vine*, is quite another thing; both as to the *Liquor* which issueth, and the *Place* where it issues: that is to say, it is neither a *Sweet*, nor *Sower*, but *Tasteless Sap*; issuing, not from any *Vessels* in the *Barque*, but from the *Aer-Vessels* in the *Wood*. So that there is as much difference betwixt *Bleeding* in a *Vine*, or the *Rising* of the *Sap* in any other *Tree*, in *March*, and in *July*; as there is betwixt *Salivation* and an *Hemorrhage*; or betwixt the *Course* of the *Chyle* in the *Lactiferous Vessels*, and the *Circulation* of the *Blood* in the *Arteries* and *Veins*.

10. §. NOW the Cause from whence it comes to pass, that the *early Spring-Sap* of a *Vine*, and other *Trees*, ascendeth by the *Wood*, is, In that the *Generation* of the young *Sap-Vessels* in the *Barque*, by which the *Sap* ascendeth all the *Summer*; is, in the beginning of *Spring*; but newly attempted. So that the *Sap* having not yet these *Vessels* to receive it, it therefore (*pro hac vice*) runs up the *Aer-Vessels* in the *Wood*. But so soon as the said *Vessels* in the *Barque* begin to be considerably encreased, the *Sap*, declining the *Aer-Vessels*, betakes it self to *These*, as its most proper *Receptacles*.

11. §. THE CAUSE also, why the *Vessels* of almost all *Plants*, upon cutting, do yield *Sap*, or *Bleed*; is the *Pressure* which the *Parenchyma* makes upon them. For the *Pith* and other *Parenchymous Parts* of a *Plant*, upon the reception of *Liquor*, have always a *Conatus* to *dilate* themselves. As is manifest from *Sponges*, which are a Substance of the same Nature, and have a somewhat like structure. As also from *Cork*, which is but the *Parenchyma* or *Barque* of a *Tree*. I say therefore, that the *Parenchyma* being fill'd and swell'd with *Sap*, hath thereby a continual *Conatus* to *dilate* it self; and in the same degree, to press together or contract the *Vessels* which it surroundeth. And the said *Vessels* being cut, their actual *Contraction* and the *Eruption* of the *Sap*, do both immediately follow.

12. §. IT may be also noted, That the *Trunk* or *Branch* of any *Plant* being cut, it always *bleeds* at both ends, or upwards and downwards, alike

alike freely. Which, as well as divers other *Experiments* plainly shews, That in the *Sap-Vessels* of a *Plant*, there are no *Valves*.

13. §. FROM what we have now above, and elsewhere formerly said, we may also understand the *manner* of the *Ascent* of the *Sap*. As to which, I say, *First*, That considering to what height and plenty, the *Sap* sometimes ascends; it is not intelligible, how it should thus ascend, by virtue of any one *Part* of a *Plant*, alone; that is neither by virtue of the *Parenchyma*, nor by virtue of the *Vessels*, alone. Not by the *Parenchyma* alone. For this, as it hath the Nature of a *sponge* or *Filtre*, to suck up the *Sap*; so likewise, to suck it up but to a certain height, as perhaps, about an Inch, or two, and no more.

14. §. Nor by the *Vessels* alone, for the same reason. For although we see, that small *Glass-Pipes* immersed in *Water*, will give it an ascent for some Inches; yet there is a certain *period*, according to the *bore* of the *Pipe*, beyond which it will not rise. We must therefore joyn the *Vessels* and the *Parenchyma* both together in this Service; which we may conceive performed by them in the manner following.

15. §. Let A B be the *Vessel* of a *Plant*. Let C E D F be the *Bladders* of the *Parenchyma*, wherewith, as with so many little *Cisterns*, it is surrounded. I say then, that the *Sap*, in the *Pipe* B A, would, of it self, rise but a few Inches; as suppose, from D to L. But the *Bladders* D P, which surround it, being swelled up and turgid with *Sap*, do hereby press upon it; and so not only a little contract its bore, but also transfuse or strain some *Portion* of their *sap* thereinto: by both which means, the *Sap* will be forced to rise higher therein. And the said *Pipe* or *Vessel* being all along surrounded by the like *Bladders*; the *sap* therein, is still forced higher and higher: the *Bladders* of the *Parenchyma* being, as is said, so many *Cisterns* of *Liquor*, which transfuse their repeated Supplies throughout the length of the *Pipe*. So that by the supply and pressure of the *Cisterns* or *Bladders* F D, the *sap* riseth to L; by the *Bladders* Q L, it rises to M; by the *Bladders* N M, it rises to I; by the *Bladders* O I, it rises to K; by the *Bladders* P K, it rises to E; and so to the top of the *Tree*. And thus far of the *Motion* of the *Sap*.

C H A P. II.

Of the Motion and Course of the Aer.

THE NEXT enquiry to be made, is, into the *Motion and Course of the Aer*. Where this question will first of all be asked; *sc.* Which way the *Aer* first enters the *Plant*; whether at the *Trunk*, *Leaves*, and other *Parts* above ground; or at the *Root*? I answer, *That it enters in part, at them all*. For the *Reception*, as well as *Extramission* whereof, the *Pores* are so very large, in the *Trunks* of some *Plants*, as in the better sort of thick walking *Canes*, that they are visible, to a good Eye, without a *Glass*; but with a *Glass*, the *Cane* seems, as if it were stuck top full of holes with great *Pins*: being so large, as very well to resemble the *Pores* of the *Skin* in the end of the *Fingers* and *Ball* of the *Hand*. Tab. 19.

2. §. In the *Leaves* of *Pine*, they are likewise through a *Glass*, a very *Elegant Show*; standing all most exactly, in *rank* and *file*, throughout the length of the *Leaves*. The *Figure* whereof shall be given hereafter, when we come to the *Anatomy* of the *Leaf*.

3. §. But although the *Aer* enters, in part, at the *Trunk* and other *Parts*, especially in some *Plants*; yet its chief entrance, is at the *Root*. Even as some *Parts* of *Aer*, may continually pass into the *Body* and *Blood*, by the *Habit*, or *Pores* of the *Skin*; but the chief entrance hereof, is at the *Mouth*. And what the *Mouth* is, to an *Animal*; that the *Root* is to a *Plant*.

4. §. Again, if the chief entrance of the *Aer*, were at the *Trunk*; then, before it could be mixed with the *sap* in the *Root*, it must descend; and so move not only contrary to its own Nature, but likewise in a contrary Course to the *Sap*, throughout the *Plant*. Whereas, by its *Reception* at the *Root*, and so its *Transition* from thence; it hath a more natural and easie motion of *Ascent*. For while the *sap* ascends, that the *Aer*, in the same *Plant*, should continually descend, cannot reasonably be supposed.

5. §. The same is further argued, From the fewness and smallness of the *Diametral Portions* in the *Trunk* in comparison with those in the *Root*. In which Nature hath plainly designed the same, for the *Separation* of the *Aer* from the *sap*, after they are both together received thereinto. So that the *Reception* and *Course* of the *Aer*, is made on this manner following.

6. §. THE *Aer* being a *springy Body*, it insinuates into all the *Holes* and *Cranies* of the *Earth*; and so is plentifully mixed therewith. Whereupon, as the *Sap* enters the *Root*, more or less *Aer* still intrudes it self together with it. The *Liquid* *Portion* of the *sap*, swells and fills up the *Succulent Parts* of the *Barque*. The *Aery Part*, is, as was said, separated from the *Liquid*, into the *Diametral Portions*. Which running

running from the *Barque* towards the Centre of the *Root*, and so passing along betwixt the *Aer-Vessels*; do hereby convey the *Aery Part* of the *Sap* from the *Barque*, into the same.

7. §. Being thus received into the *Aer-Vessels*, and the *Reception* thereof, by the same means continued; it is by them advanced into the *Trunk*. In which advance, it is again, more or less, disbursed into all the *Parts* of the *Trunk*, as it goes. *Partly*, inwards to the *Pith*. From whence, the *Pith* is always, at length, filled with *Aer*. *Partly*, into the *Insertions*; by which it is conveyed outward into the *Barque*. Wherein, it is in some part, transfused through the *Sap*: and so the rest, with part of the *Sap*, remitted, in *perspirations*, back again into the *Aer*.

8. §. So that, whereas the *Diametral Portions* in the *Root*, do serve to convey the *Aer* from the *Sap* in the *Barque*, into the *Aer-Vessels*, in the *Wood*: on the contrary, the *Insertions* here in the *Trunk*, serve to convey the *Aer* from the *Aer-Vessels* in the *Wood*, into the *Sap*, in the *Barque*. Wherefore, as the *Aer-Vessels* advance the *Aer*, or the *Aery Part* of the *Sap*, and so convey it by the *length* of the *Trunk*; so the *Insertions* filter it, and convey it by the *breadth*.

9. §. AND that the *Insertions* have this *Office* or *Subservience* unto both *Kinds* of *Vessels*; doth yet further appear, if we consider, That the *Aer-Vessels* are always so postured, as to touch upon the said *Insertions*, or at least to stand very near them. For either they are large, and so do frequently touch upon them on both sides; as in
 Tab. 28. *Elm, Ash, Walnut, &c.* Or if they are small; then they either run
 29. along in even lines collateral and oftentimes contiguous with the said
 22. *Insertions*, as in *Holly*: or at least, are reciprocally, some on one side,
 25. and some on another, inclined to them; as in *Apple*. By all which means, the *Aer* is more readily conveyed from the *Vessels* into the *Insertions*.

10. §. A further evidence hereof is this, That generally, the bigger and the more numerous the *Aer-Vessels* be; the bigger, or at least, the more numerous also are the *Insertions*: Especially, if the comparison be made (as in all other cases it ought to be, as well as here) betwixt the several *species* of the same *Kind*. So *Corin*, which hath small *Aer-Vessels*, hath also very small *Insertions*. But the *Vine*, hath both very large: and so for others.
 Tab. 17.

11. Wherefore, the *Insertions* minister betwixt the *Aer-Vessels*, and the *Succiferous*; in the same manner, as the *Vesiculae* of the *Lungs*, do betwixt the *Bronchia* and the *Arteries*. That is to say, as in an *Animal*, the *Bronchia* deposite the *Aer* into the *Vesiculae* of the *Lungs*; which administer it to the *Arteries*: so in a *Plant*, the *Aer-Vessels* deposite the *Aer* into the *Insertions*, that is into the *Vesiculae* of the *Insertions*; by which it is gradually *filtered* off into the *Barque* and the *Sap-Vessels* therein.

C H A P. III.

Of the Structure of the Parts.



THIRD enquiry, is into the *Generation* and *Structure* of *Parts*. The manner whereof I have already endeavoured to *explicate* (a) from the *A-* (a) *Lib. 2.*
natomy of the *Root*, throughout all particulars. *P. 2.*

Some whereof I shall yet further clear.

1. §. As *First*, the *Union* of the *Barque* to the *Body* of the *Tree*, *Contrary* to the common *Opinion*, *That they are not continuous*; but that the *Barque* only surrounds the *Body*, as a *Scabbard* does a *Sword*, or a *Glove* the *Hand*. As also seemeth to be proved, by the easy *slipping* of the *Barque* of *Willow*, and most other *Trees*, when full of *Sap*, from the *Wood*.

2. §. But, notwithstanding this, they are as truly continuous, as the *skin* of the *Body* is with the *Flesh*: *sc.* by means of the *Parenchyma*; which is one entire *Body*, running from the *Barque* into the *Wood*, and so uniting both together; as in a *Branch* of *Vine* or *Corin-Tree*, when the *Barque* is stripped off, is apparent; the *Spaces* between the several *Parts* of the *Wood*, being filled up with the *Parenchymous*, inserted from the *Barque*. *Tab. 19.*

3. §. Now the reason why the *Barque* nevertheless slips so easily from the *Wood*, is plain, *viz.* Because most of the young *Vessels* and *Parenchymous Parts*, are there every year successively formed; that is, betwixt the *Wood* and *Barque*: where the said *Parts* newly formed, are as tender, as the tenderest *Vessels* in *Animals*. And we may imagine, how easie it were at once to tear or break a thousand *Vessels* or *Fibres* of an *Embryo*, of a *Womb* or *Egg*.

4. §. THE same *Vessels* of the *Barque* being always braced, and gradually falling off, together with the *Parenchyma*, into the utmost *Rind*: Hence it is, that the *Barques* of many *Trees*, are as it were, laticed with several *Cracks* of divers *Sizes*, and sometimes in the *Figure* of *Rombs*: the said *Fissures* representing the *Position* and *Tract* of the *Vessels* in their *Braces*: Hence also it is, that the *Barque* of some *Trees*, as of *Corin*, *Cherry*, &c. falleth off in *Rings*, *sc.* because the *Sap-Vessels* are posited in the same manner in the *Barque*. *Tab. 19.*

5. §. The *Sap-Vessels*, as they are generated at the inner *Verge* of the *Barque*: so likewise, in a small quantity, at the utmost *Verge* of the *Pith*. These being not only fed with a more vigorous *Sap*, but with great caution, secured within the *Wood*, for the propagation of the succeeding *Buds*.

6. §. Hence also it is, that is, by the annual accretion of these *Vessels*, that the *Pith* is sometimes less in the *Trunk*, than in the *Branches*; *Tab. 18.* and less in the elder *Branches*, than in the younger; and sometimes 'tis almost wholly filled up. By which means, as the *Branches* carry every year a greater burthen; so they become still more sturdy the better to support it.

7. §. SOMETIMES also the *Pith* breaks and shrinks up, thus making the *Trunk* a *Pipe*. The cause whereof, is either the *Largeness* of its *Pores*, or the *Thinness* of the *Sides* of the said *Pores*; upon both

Z

which

which accounts, the *Pith* doth more easily tear, and upon tearing shrink up, and so become hollow: as in *Cichory*, *Lampfana*, *Sonchus*, *Teasel*, *Brownwort*, and others; wherein the *Pores* of the *Pith* are *Large*, and the *Sides* of the *Pores*, *Thin*. Whereas, upon contrary accounts, the *Piths* of most *Trees*, remain perpetually entire.

8. §. THE Reason why *Plants* are made thus to become *hollow*, is partly, for the ripening of the *Fruit* or *Seed*; which is the better effected by a more plentiful supply of *Aer* continually received into their hollow *Trunks*. For by means of that *Aer*, part of the *Sap*, is dried up, and the remaining part of it made warmer, and so sooner matured.

9. §. Partly, for the better determining the due *Age* of the *Plant*. Hence it is, that the greater part of *Annual Trunks*, are hollow: the *Aer* contained in that hollow, drying up the *Sap*, and shrinking up the *Sap-Vessels* so far, as to hinder the free motion of the *Sap* therein; from whence the *Plant* must needs perish. So that as the *Content* of the *Aer-Vessels*, is a kind of *Vegetable Aer*, whose Office is to *Attenuate*, and *Ferment* the *Juyces* of *Plants*: so the *Content* of these *Cavities*, cometh nearer to a more common *Aer*, designed chiefly, so soon as it is convenient, to *dry* them up.

10. §. A GAIN, as to the *Aer-Vessels*, divers questions may be asked. As how it comes to pass, that they are generally less in the *Trunk* of the same *Plant*, than in the *Root*? The Cause whereof is, that here in the *Trunk* they are more under the power of the *Aer*; both that which entrench in at the *Trunk*, and that which of its own *Nature* ascendeth up into it from the *Root*. For the *Aer*, as we have elsewhere said, is the *Mould* of the *Aer-Vessels*; to whose crooked or at least, *Acid Parts*, the *saline*, and other *Principles* concurring to their generation, do conform. To which they do best, the smaller they are: the *Fibres* of the larger *Aer-Vessels* making greater *Circles*, and so coming nearer to a *right Line*, answerable to the *Figure* of the *Particles*, not of the *Aerial*, but of the *Saline Principle*.

11. §. Wherefore as the *Aer-Vessels* may be observed still to be dilated or widened towards the lower parts of the *Root*; the *Aerial Principle* being there less predominant, and the *Saline* more: So towards the upper part of the *Trunk*, to be contracted or grow smaller; the *Aerial Principle* being here more predominant, and the *Saline* less.

12. §. FOR the same cause it may be observed, That the *Aer-Vessels* of the *Second* years *Growth*, and the several years succeeding, are usually nearer of one *Size*, than those of the *Second* and *First*; all being under a less power of the *Aer*, than the *First*. For the first year the *Pith* being full of *Liquor*, the *Aer-Vessels* themselves, are the only *Repositories* of the *Aer*. Whereas, after the first year, the *Pith* becoming dry, or another great *Repository* for the *Aer*; the *Aer-Vessels* are henceforth filled with a moister or more *Vaporous* and *Saline Aer*, and so made to grow wider.

13. §. Hence the very *Size* of the *Pith*, hath much influence upon the *Aer-Vessels*, and the manner of *Nutrition*, and the *Generation* of *Liquors* in *Plants*.

14. §. BUT for the most part, the *Aer-Vessels* are somewhat, more or less, amplified in every new *Annual Ring*; or at least to a certain number of years. Probably, because in the elder *Branches*, the *Spiral Fibres*, of which the *Vessels* consist, are more bulky; and so make a
Vessel

Vessel of a wider, as a more agreeable bore. Nature obtaining hereby, that the Quantity of *Aer*, shall always be answerable to the Growth of the *Plant*, or at least, be sufficient to maintain its *Vegetable Life* and *Vigour*.

15. §. And therefore, as is above hinted, it seems likely, That after a certain number of years, the *Aer-Vessels* are no longer amplified, but stand at a stay, and perhaps may grow smaller, according as the *Tree* is less or more *Longeve*; and that after this period, it is some way or other in its Declining State.

16. §. LASTLY, from the *Content* and *Governing Principle* of the *Aer-Vessels*, the *Time*, when they begin every year to be formed, or to appear, is always *later*; at least with respect to the *season* of the *Tree*. So that whereas the *Sap-Vessels* begin to be formed in *Spring*: these, not till the latter end of *Summer*, or there about; at least not till about that time to appear. That is, when the *Sap* begins to decrease, and to grow more *Aery*; and so more fit matter for the *Generation* of the said *Aer-Vessels*.

C H A P. IV.

Of the Generation of Liquors.



PON the *Structure* and *Formation* of the *Parts*, dependeth the *Generation* of *Liquors*, as was lately intimated. The *manner* whereof I have formerly shewed, in discoursing of the *Root*. Yet some things I shall here further *explicate*. And *First*, what we have formerly asserted, *sc.* That the concurrence of two *specifically* distinct *Fluids*, is as necessary to *Nutrition* in *Plants*, as in *Animals*. Which appears, as from divers other considerations, so from the very *Structure* of a *Plant*: where in all the *Organical Parts*, or the *Parenchyma* and the *Vessels*, are every where mixed together *per minima*, that is, *per minimas partes organicas*, or *Fiber* with *Fiber* of several *Kinds*. Every small part of a *Tree*, or of the *Barque* of a *Tree*, being as I may say, a sort of *Linsy-Woolsey*. So that there is not the least part of the *Sap*, which is not impregnate with divers *Essential Tinctures*, as it is continually *filtered* from the *Fibres* of *one Kind*, to those of another; standing every where *woun'd* and *stitch'd* up together for the same purpose.

2. §. FROM the special *Nature* and *Structure* of the *Parts*, the *Liquors* of *Plants* are likewise *specified*. The *Vessels* being the chief *Viscera* of a *Plant*. For all *Liquors* in a *Plant*, are certainly made by that *Plant*. And since the *Plant* hath no *Viscera* (so called) I would then know, what its several *Liquors* are made by? If in the *Parenchyma*, surely by that *Parenchyma*. If in the *Vessels*, by the *Vessels*. And if of divers *Kinds* by divers *Kinds* of *Vessels*. So that what the *Viscera* are in *Animals*, the *Vessels* themselves are in *Plants*. That is to say, as the *Viscera* of an *Animal*, are but *Vessels conglomerated*: so the *Vessels* of a *Plant*, are *Viscera drawn out at length*.

3. §. AGAIN, as the *specifying* of the *Sap* dependeth chiefly on the special *Nature* of the *Parts*: so partly, upon the *Structure* of the *Whole*. Whereby every *Part* is still better accomodated with its own *Juyce*. Thus the *Aer-Vessels* are necessary, not only and barely for a supply of *Aer*; but also by their *Number*, *Size*, and *Position* to adjust the quantity of that *Aer*, to the government of *Nutrition*, and the *Generation* of the *Specifick Liquors* of every *Plant*. Which is evident from hence, in that they do not follow the *Size* of the *Plant*; but are great and many, in some small *Plants*; and small and few, in some others that are large. So *Vines*, and *Corn*, as we have formerly observed, have proportionably a great number of *Aer-Vessels*, and those very large. By which means the *Sap* is *attenuated* and less *Oyly*, and more copiously impregnated with a *Subtle*, *Volatile* and *Winy Spirit*.

4. §. For the same reason, the *Stalk* of *Maze* or of *Indian Wheat*, which when it is *Green* yieldeth a very sweet *Juyce*; and the *Canes*, whereof *Sugar* (which aboundeth with a *volatile* and *inflammable spirit*) is made; these, I say, obtain the like over proportion of *Aer-Vessels*, to what we see in most other *Plants*. Hence also it is, that none of the said *Plants* have any considerable *Barque*; that so the attenuating and subtilizing *Aer*, may have a more easie and plentiful admission at the *Trunk* also. For which reason likewise the *Pores* of the *Skin* of some *Canes* are, as hath been said, remarkably wide.

5. §. Hence also it is observable, that of the same *Species* or *Kindred*, those *Plants* which have the most, and especially the largest *Aer-Vessels*; have also the greatest abundance either of a *sweet*, or of a *winy Liquor*. So in *Apple*; they are larger than in *Crab*; In *Warden*, larger than in *Quince*; and in *Pear-Tree*, larger than in *Warden*. So also in *Corin*, larger than in *Gooseberrey*; and in *Vine*, larger than in *Corin*: and so in others.

6. §. AND as the *Aer-Vessels*, by their *Multitude* and *Largeness*, are accomodated to the better making of a *Winy Sap*: so by their *fewness* and *smallness*, of an *Oylie*. As is remarkably seen in *Fir*, and other *Resiniferous Trees*: these having, if not the *smallest*, yet the *fewest Aer-Vessels* of all other *Trees*.

7. §. IF it be asked, how a *Plant* comes to have any *Oyl* at all in any *Part*? Since we see, that the *Sap* by which the *Root* is fed, seemeth to be nothing else but *Water*: and that many *Plants* which yield a great deal of *stillatitious Oyl*, as *Mint*, *Rue*, and others, will yet grow in *Water*: I say, if it be enquired how this *Water*, is made *Wine* or *Oyl*? I answer, that there is no such matter. But that the *Oyl*, and all other *Vegetable Principles* are actually existent in, and mixed *per minima*, though in an extraordinary small proportion, with the *Water*. Even as we see the distilled *Waters* of *Anise Seeds*, *Penyroyal*, and the like to be impregnated with their own *Oyls*, which give the *Taste* and *Smell* to such *Waters*.

8. §. Wherefore, as a certain quantity of any *Salt* may be dissolved in *Water*; beyond which, it will not mix therewith, but remains under its own *Form*: So is there a certain proportion of *Oyl*, though far less, which may also be perfectly mixed with *Water*; and is certainly so, more or less, with all the *Water* in the world. But if that proportion, or degree of impregnation be once exceeded; the particles of *Oyl* do then, and not till then, gather into a body, and appear under their own *Form*.

9. §.

9. §. I say therefore, that all kinds of *Vegetable Principles*, are either in or together with the *Water*, with less difference first received into a *Plant*. But when they are once therein; they are then *separated*, that is to say, *filtered*, some from others, in very different *Proportions* and *Conjunctions* by the several *Parts*; the *Watery* by one *Part*, the *Aery* by another, the *Oily* by another, and so the rest: and so every *Part* is the *Receptacle* of a *Liquor*, become peculiar, not by any *Transformation*, but only the *Percolation* of *Parts* out of the *common Mass* or *Stock* of *Sap*. And so all those parts of the *Sap*, which are *superfluous* to any kind of *Plant*, are at the same time, discharged back by *Perspirations*, into the *Aer*.

10. §. AND, that *Nature*, in the various *Percolations* and *Separations* of the *Sap*, may still the better answer her end; hence, it is, that she carefully seeth, not only to the special *Nature* and *Proportion* of the *Organs*, by which she doth her work; but likewise to their very *Position*. Thus it is observable, That whereas the *Lymphæducts*, which carry a more *Watery Liquor*; are still placed on the inner *Verge* of the *Barque*, next to the *Aer-Vessels*: the *Lactiferous* and *Resiniferous Vessels* of *Plants*, to whose *Oylie Liquor* a mixture of much *Aer* is incongruous; do usually stand, neither on the inner, nor the outer *verge* of the *Barque*; but in the *middle*. By which means, they are at the greatest distance, and so most secure, from the *Aer*; either that which enters the *Barque* at the *Circumference*, or from the *Wood* and *Pith*.

11. §. AND because the *Resinous Liquors* of *Plants* are more *Oily*, than their *Milky*; their security therefore, from the approach of the *Aer*, is yet further contrived. In that in *Pine*, and other *Resinous Trees*, the *Diametral Insertions* are never found; or at least, not *visible*: which yet in other *Trees*, are *conspicuous*; being those *Parts*, whose office it is, to introduce the *Aer* from the *Aer-Vessels* into the *Barque*.

12. §. AGAIN, the *Milky Liquors* of *Plants* being thinner than the *Resinous*, and having a considerable quantity of *Water* mixed with their *Oyl*; hence it is, that in *Milky Plants*, as in *Rhus*, there are a greater number of *Lymphæducts*; and those standing nearer to the *Milky Vessels*, than they do in *Pine* and the like, to the *Resinous*. By which means they are better fitted to affuse their *Aqueous Parts* more plentifully to the said *Milky Liquor*.

13. §. FROM the *Mixture* of *Watery Parts* with the *Oylie*, it comes to pass, that whereas all *Lymphæ's*, *Mucilages*, and *Resins* are transparent; the *Aquæ-oleous Liquors* of *Plants* are *Milky* or white, or otherwise *Opacous*. For the same thing is the cause of the whiteness of *Vegetable*, as of *Animal-Milk*: that is to say, a more copious mixture of *Watery* and *Oily Parts per minima*, or into one *Body*. For even the *Serous* and *Oylie Parts* of *Animal Milk*, when thoroughly separated one from the other, they become very transparent. So the *Stillatitious Oyl* of *Anise Seeds*, is most transparent and *limpid*, even as *Water* it self: yet there is a known sort of *White Anise-Seed Water*, as it is commonly called: that it is to say, wherein the *Oyl*, in distillation, ariseth and is mixed more plentifully with the *Water*. And the *Water*, wherein the stillatitious *Oyl* of any *Vegetable* is dissolved, becomes a perfect white *Milk*; as in this *Honourable and Learned Presence*, I have formerly had occasion to shew the *Experiment*. (a)

(a) See the
Discourse
of Mixture

14. §.

14. §. AND that the *Milky Liquors* of all *Vegetables* whatsoever, are more *Oylie* than their *Lympha's*, is most certain. For all those *Gums*, which dissolve either in *Oyl* or in *Water*, as *Galbanum*, and the like, are originally the *Milky Juices* of *Plants*. And if you take the *Milk* of any *Plant*, as for instance, the *Milk* of common *Sumach*, or of any *Taste*, *Bitter*, *Astringent*, *Hot*, *Cold*, or any other whatsoever; and having well *dried* it, and then fired it at a candle; it will thereupon burn with a very bright and durable flame, even like that of *Tar* or *Turpentine* it self.

15. §. FROM what hath been said, we may likewise gather the most genuine import of the word *Gum*, and the distinction thereof both from a *Rosin* and a *Mucilage*. First, a *Rosin*, is originally a *Turpentine*, or *Acidoleous Liquor*, having an exceeding small quantity of *Watery Parts* mixed therewith; and which, for that reason, will not be dissolved in *Water*, but only in *Oyl*. Of this kind are *Mastick*, *Benzoine*, *Taccamahacca*, and divers others, commonly, in our *Bills* to *Apothecaries*, called *Gums*. Yet in strict speaking they are all so many *Rosins*.

16. §. Secondly, a *Gum*, and every *Oylie Gum*, is originally a *Milky Liquor*, having a greater quantity of *Water* mixed with its *Oily Parts*; and which for that reason, will be made to dissolve either in *Water* or *Oyl*. Of this kind are *Sagapen*, *Opopanax*, *Ammoniac*, and others.

17. §. The third sort of *Gum*, is that which is *Unoylie*, and which therefore dissolveth only in *Water*, as *Gum-arabick*, the *Gum* of *Cherry-Tree*, and others such like. This *Gum*, though commonly so called, yet is properly but a dried *Mucilage*: being originally nothing else but the *Mucilaginous Lympha* issuing from the *Vessels* of the *Tree*. In like manner, as it doth from *Cumfry*, *Mallow*, and divers other *Plants*: and even from the *Cucumber*. The *Vessels* whereof, upon cutting cross, yield a *Lympha*, which is plainly *Mucilaginous*, and which being well dried, at length becomes a kind of *Gum*, or rather a hardened *Mucilage*. In like manner, the *Gums* of *Plum-tree*, *Cherry-tree* and the like, are nothing else but *dried Mucilages*. Or, if we will take the word in its widest sense, then all *Gums* are originally, either a *Terebinth*, or a *Milk*, or a *Mucilage*.

18. §. I have likewise made divers Observations of the *Tastes*, *Smells*, and *Colours* of *Plants*, and of their *Contents*, since those I last published: and that both for the finding out the true *Causes* of their *Generation*, and also the applying of them unto *Medical* and other *Uses*. Of which hereafter.

C H A P. V.

Of the Figuration of Trunks.



THE Fifth Head, shall be, of the Figuration of Trunks. Which also, as well as the making of Liquors, dependeth upon the Structure of the Parts. As First, almost all Shrubs (*ceteris paribus*) have a greater number of Aer-Vessels; and those of a smaller Size; and consequently much spread abroad, as most easily yielding to the magnetick Power of the Aer, according as we have more fully demonstrated, in speaking of the Vegetation of Roots: as in Elder, Hazel, Fig, Sumach, and the like. By which spreading, the said Aer-Vessels do sooner, and more easily strike into the Barque, and so produce collateral Buds and Branches, and that upon the first rising of the Body from the Root: that is, the Plant becomes a Shrub.

2. §. BUT if the said Aer-Vessels are very large, they will not yield so easily to shoot out collaterally; and so the Trunk grows up taller and more entire: as in Oak, Wallnut, Elm, &c. wherein they are exceeding large, is seen. Hence also the Vine, if supported, will grow to a prodigious length. And Hops and Bryony, are some of the tallest, amongst all Annual Growths: the Aer-Vessels of all which, are very large. Whereas Borage, and many other like Plants, although the Pores of their Parenchyma, are vastly wide, and filled with Sap; yet because their Aer-Vessels are small, they are therefore but Dwarf-Plants. Wherefore the tallness or advancement of a Plant or Tree, dependeth not upon the Plenty of Sap, how great soever, but on the Largeness of the Aer-Vessels.

3. §. AGAIN, as a Plant or Tree grows either Shrubby, or Tall and Entire, according to the Size of the said Vessels: so from their Position, doth it grow slender or Thick. So, where they keep more within the compass of a Ring, as in Elm, and Ash, the Tree, in proportion, usually grows taller, and less thick. But where the said Vessels are spread more abroad, and especially are postured in Rays, as they are in Oak, the Tree grows very thick. Because the said Vessels thus standing all along nearer to the Insertions, there is a more ready and copious passage of the Aer out of the one into the other; and so the Diametral growth of the Wood is more promoted.

4. §. LASTLY, from the same general cause it is, That the Trunks of Vegetables are either Round or Angular. Those of all Trees are Round. Because the Barque, being here thicker, and the Aer-Vessels bound up with a greater quantity of Wood; the Aer hath not sufficient power to move them, and the Barque with them, into those various Positions or Figurations, as the Trunks of Herbs do yield to.

5. §. Yet the cause of the various shapes of the Trunk, is not the Aer alone; but partly, the Principles of the Plants themselves, in conjunction therewith; according to the predomination whereof, and chiefly of some certain kind of Salt or Salts, as I shall hereafter (a) more particularly explicate) the Trunk is Square, Triangular, Pentangular, or otherwise Figured. And thus much in general of the Figuration of Trunks.

(a) B. 4. P. 1. Ch. 6.

C H A P. VI.

Of the Motions of Trunks.



THE *Motions* also of *Trunks* are various. Principally *Four*; *sc.* *Ascending, Descending, Horizontal,* and *Spiral.* The cause of the *Ascent* of a *Plant*, is a certain *Magnetick Correspondence* betwixt the *Aer* and the *Aer-Vessels* of a *Plant*; the *Motion* and *Tendency* whereof, the whole *Plant* follows. This I have asserted, and I think, clearly demonstrated in my *First* and *Second Books* of the *Anatomy of Plants.* I will here add this plain *Experiment.*

2. §. Take a *Box of Moulds*, with a hole bored in the bottom, wide enough to admit the *stalk* of a *Plant*, and set it upon stilts half a yard or more above ground. Then lodg in the *Mould* some *Plant*, for Example a *Bean*, in such sort, that the *Root* of the *Bean* standing in the *Moulds* may poynt upwards, the *stalk* towards the ground. As the *Plant* grows, it will follow, that at length the *stalk* will rise upward, and the *Root*, on the contrary, arch it self downward. Which evidently shews, That it is not sufficient, that the *Root* hath *Earth* to shoot into, or that its *Motion* is only an *Appetite* of being therein lodged, which way soever that be: but that its nature is, though within the *Earth* already, yet to change its *Position*, and to *move Downwards.* And so likewise of the *Trunk*, that it rises, when a *Seed* sprouts, out of the *Ground*, not meerly because it hath an *Appetite* of being in the open *Aer*; for in this *Experiment* it is so already; yet now makes a new *Motion* upwards.

3. §. BUT although the *Natural Motion* of the *Trunk* be to *Ascend*; yet is it forced oftentimes to *Descend.* For the *Trunk-Roots* growing out of some *Plants* near the ground, and shrinking thereinto, like so many *Ropes*, do pluck the *Trunk* annually lower and lower into the ground together with them; as may be seen in *Scrophularia, Jacobaea,* and many other *Plants.*

4. §. IF these *Trunk-Roots* break out only about the *bottom* of the *Trunk*, as in the aforesaid *Plants*, then the *Trunk* gradually *Descends* into the *Earth*, and is turned into a *Root.* But if it be very *slender*, and the *Trunk-Roots* break forth *all along* it, then it *Creeps* horizontally; the said *Roots* tethering it, as it trails along, to the ground; as in *Strawberry, Cinquefoyl, Mint, Scordium,* &c.

5. §. AS to their *Spiral Motion*, it is to be noted; That the *Wood* of all *Convolvula's* or *Winders*, stands more close and round together in or near the *Center*, thereby making a round, and slender *Trunk.* To the end, it may be more tractable, to the power of the external *Motor*, what ever that be: and also more secure from breaking by its winding *Motion.*

6. §. Wherefore, *Convolvula's* do not wind by any peculiar Nature or *Genius*, which other *Trunks* have not; but because their *Parts* are disposed so, as to render them more sequaceous to the external *Motor*. Even as the *Claspers* of a *Vine*, having the like *Structure*, have also a *Motion* of *Convolution*: whereas the *Branches* themselves upon a contrary account, move in a *straight Line*.

7. §. The *Convolution* of *Plants*, hath been observed only in those that *Climb*. But it seems probable, that many others do also *wind*; in which, the main *Stalk*, is as the *Axis* to the *Branches* round about. Of which number, I conceive, are all those whose *Roots* are twisted; a *Motion* we observed in speaking of the *Root*. Whether it be so, or not the Experiment may easily be made by tying a *Thred* upon any of the *Branches*; setting down the respect it then hath to any *Quarter* in the *Heavens*: for, if it shall appear in two or three Months, to have changed its *Situation* towards some other *Quarter*; it is a certain proof hereof. And that hereby the *Roots* of many *Plants* become twisted; the *Motion* beginning in the *Stalk*, and ending at the bottom of the *Root*, which stands always fixed in the same place.

8. §. The *Convolution* of *Trunks*, is made not *one*, but *divers* ways; some moving by *South* from *East* to *West*; and others from *West* to *East*. Wherefore it seemeth, that as the *Efficient Cause* of *Convolution*, is not within the *Plant*, but external: so also, that it is not *One*, but that there are *Two Great Efficients* of this *Motion*; *sc.* the *Sun* and the *Moon*. Some winding together with the *Sun*, in its *Diurnal Motion*, (or, if the *Earth* moves, then, *Inclining* to the *sun*) by *South* from *East* to *West*. And others winding with the *Moon*, in its *Monthly Motion*, from *West* to *East*.

9. §. This possibly, may also be one *sensible* way of distinguishing betwixt *Solar*, and *Lunar Plants*. Thus far, in general, of the *Motions* of *Trunks*.

CHAP. VII.

Of the Nature of Timber or Trunks, as they serve for Mechanick Use.

THE last thing I purposed to speak of, is, Those several *Qualities* of *Timber* or of *Trunks*, by which they are fitted for *Mechanical Use*. As *Hardness*, *Softness*, *Fastness*, *Cleavesomeness*, *Toughness*, *Brittleness*, *Durableness*, or any of the same *Qualities* compounded. The *Visible Causes* whereof are observable, *Partly*, in the *Structure* of the several *Parts*; *sc.* the *Insertions*, *Sap-Vessels* and *Aer-Vessels*; as to the *Number*, *Size*, or *Position* of any of them. And *partly*, in the *Nature* of the *Parts*; I mean such as is *manifest to sense*. According to our clear and distinct observing of all which *Causes*, we may understand, Wherefore any *Wood* is made use of for any certain purpose. And also, wherein fitly to apply it to further *Use*. In order to which, a

compleat History of the *Mechanical Uses* of *Vegetables* would very much conduce. I shall for the present give some *Instances*.

2. §. AS *First*, some *Woods* are *soft*, as *Deal*, and *sallow*. Yet from different *Causes*. *Deal*, from the great *Porosity* of the *Wood* it self, or the large *Pores* amongst the *Sap-Vessels*. But *Sallow*, from the great number of *Aer-Vessels* spread all over it. And therefore, though they are both *soft*, yet will not serve for the same purposes; *Sallow* being well wrought upon, which way soever you cut it: but *Deal*, especially the white *Deal*, if it be cut cross, it tears, and will never polish or work smooth.

3. §. Again, in *sallow*, by the equal spreading of the *Aer-Vessels*, the *Softness* is equal or alike in all *Parts*. For which cause it maketh an excellent *Coal* for *Painters Scribets*. Because it doth not only make a *light Stroak*, but every where *certain*; and so doth not disturb the even *Motion* of the *Hand*. For the same cause, *Shoemakers* also make use of it for their *Carving-boards*. Because being every where equally soft, it turns not the edge of their *Knives*, Which *Deal* would presently do; because though very soft in some places, yet in others 'tis hard; that is to say, on the inner *Verge* of every annual *Ring* of *Wood*, where the old *Sap-Vessels* grow much more compact and close together.

4. §. A G A I N, some *Woods* are *soft*, but not *fast*; others are both, as *Linn*: its *Softness*, depending on the numerousness and equal spreading of the *Aer-Vessels*: its *Fastness*, on the closeness of the true *Wood*, and the shortness, and smallness of the *Insertions*. For which cause, it is of excellent use for many purposes; and particularly, for *small Sculpture*: such as may sometimes be seen for the *Frames* of *Looking-Glasses*, or of smaller *Pictures* in *Water-Colours*.

5. §. S O M E *Woods*, again, are *fast*, and *hard*, as *Elm*. Its *hardness* depending upon the closeness of the *Wood*. Its *fastness*, Partly, upon the same cause; and partly, on the smallness of the *Insertions*; as also on the fewness of the *Aer-Vessels* in proportion with the *Wood*; and on the *thwart* and *cross* *Position* of many of them. Hence it is, that *Elm*, of all others, is the most *Cross-grain'd* *Timber*; that is, cleaveth so unevenly, to and fro, according to the *cross* *Position* of the said *Vessels*.

6. §. Hence also it cleaveth the most *Difficultly*. Even then, when it is without any *Knots*. For which reason it is always used, as best for the *Hub* of a *great Wheel*. As also for *Water-Pipes*, and for *Pumps*. Not because it is the most *durable* *Wood*; but because it will not *split* or *crack*, either in the *working*, or *afterwards*. For the very same reason, it is used for *Coffins*; that is, because, it will not *split* in *working*: not because it will endure longest under ground; for *Pales* are always made of *Oak*. So also the *Ladles* and *soles* of a *Mill-wheel* are always made of *Elm*; as also the *Keel* of a *Boat*, *sc.* lest they should *split*: but the other *Parts* are made of *Oak*.

7. §. It may here also be noted, That the *Planks* commonly called *Groaning-Boards*, lately exposed, as a kind of *Prodigy*, to the view and hearing of many *People*, were of *Elm*. The *Aer-Vessels* of this *Wood*, being, though not more numerous, yet more ample, than in any other *Timber*. So that upon the application of the *Red-hot-Iron*, as was usual, and thereby the *Rarification* of the *Aer* and *Watery* *Parts* in the *Timber*; every *Vessel* became, as it were a little *Wind-Pipe* for their

their *Expiration*. And as a great many Drops falling together in a shower of Rain; so a great many of these *Pipes* playing together, might make a kind of big or groaning noyse.

8. §. AS *Elm*, of all *Woods*, is one of the *fastest*; So, on the contrary, of all *hard Woods*, *Oak* is the most *Cleavesome*, or *splitteth* the most *easily*. The cause whereof is, *partly*, the *Largeness* of the *Insertions*; and *partly*, the *Diametral* or *Radiated Position* of most of the *Aer-Vessels*: upon both which accounts, wherever a *crack* is once begun, 'tis easily continued throughout the *Diameter* of the *Trunk*.

9. §. AGAIN, some *Woods* are *hard*, *fast*, and *tough*. So is *Ash*, and especially *Beech*. *Hard* and *fast*, from some of the same Causes, as *Elm*. *Tough* not from the *Structure*, but from the *Nature* of the *Parts*; whose *Principles* are united in a more exact proportion. Wherefore *London-Cars* have the *Rings* of their *Wheels* of *Beech*; because it *tears* more difficulty than even *Ash* it self. Whence also for *large Screws*, there is no *Wood* like it. But for *Small Screws*, of about an *Inch Diameter*, *Birch* is the best; as being, though not so *hard*, yet more *tough*.

10. §. THE more *Brittle* a *Wood* is, 'tis likewise usually the more *durable*. So *Oak*, which, with respect to its *hardness*, is not a *tough*, but very *brittle Wood*, is almost as *durable* as any. Whereas *Beech*, *Birch*, and the like, although very *tough*; yet for *Duration*, are of no service; for there are no *Woods* will rot sooner: and therefore, though strong enough, yet unfit to make any *Standing Parts* of *Building*, or of *Furniture*; especially in wet and moist places. Because, these *Woods*, having a less proportion of *Oyl*, than there is in *Oak*; they are apter to imbibe the moisture even of a *dank Aer*; by which moisture, they either *Rot*, or breed *Worms*, which destroy them.

11. §. HENCE it is, that what we call the *Heart* of *Timber*, as it is more *brittle*, so also more *durable*; *sc.* Because more *Oylie*. So that which is called the *sap* of *Oak*, is much more *tough* than the *Heart*, although the *Heart* be more *durable*. That is to say, the older the *Wood* is, the *Watery Parts* are the more *evapourated*, whilst the *Oylie* still remaine, as a kind of *Tincture* or *Extract* in the *Wood*. Even as we see, that the older *seeds* of any one *Kind*, are more *Oylie* than those that are green and young. So that the *Oylie* or *Rosinous Parts* of the *Sap*, are a kind of *Embalming* to the *Heart*, or older *Part* of a *Tree*, securing it from the destructive impressions of the *Aer*. For which Cause it is, that *Oak*, *Yew*, *Cocus*, *Guajacum*, &c. which are *Oylie Woods*, have always much *Heart*, whereas *Birch*, *Alder*, *Beech*, *Maple*, which are very *Unoylie*, have never any *Heart*.

12. §. FROM hence likewise we may understand the Cause of the *Toughness* of *Flax*: what we call *Flax*, being only the *Sap-Vessels*, or *Lignous Fibres* of the *Barque*. And generally, the *Barque* of any *Tree*, as of *Willow* (whereof are usually made a sort of *Ropes*) is very *tough*. The *Vessels* being here younger, and less *Oylie* than in the *Wood*. So likewise *Hemp*, is nothing else but the *Sap-Vessels* of the *Barque* of the *Plant* so called. And *Scotch-Cloath*, is only the *Housewifery* of the same *Parts* of the *Barque* of *Nettle*.

13. §. WHENCE it is very probable, that there are many other *Plants*, as well as the above named, whereof might be made good *Tow*. And of some, especially in some respects, better than of *Flax* it self. Because that even *Hemp*, although it will not make so

fine a *Staple*, as *Flax* (for all our fine *Hollands* are made of *Flax*) yet *Flax*, which is but of the same fineness as *Hemp*, will never, by all the Art yet known, be made so white as *Hemp* is made. The *Qualities* therefore of the best *Tow*, that can be in Nature, are that the *Staple* be long, small, tough, and white. So that if in the *Barque* of any *Plant*, we can find these *Qualities*, or any of them, to excell; we may be sure, it will be of better use, in some respects, for the making of *Cloath*, or other purpose, than *Flax* it self.

14. §. I WILL conclude with one *Instance* more, and that is as to *Grafting*. The good and happy success whereof, doth certainly depend upon the suitableness or response betwixt the several *Parts* of the *Stock* and *Cyon*; as the *Barque*, *Wood*, and *Pith*; and that both as to the *Number*, *Size*, and *Position* of the said *Parts*, and of their several *Pores* or *Vessels*: according to the degrees whereof, the *Conjunction* (*ceteris paribus*) will be more or less prosperous. So that of all such *Conjunctions* as are found to be apt and taking, and which some have learned not without long *Practice* and *Experience*; another, only by comparing the *Branches* of *Trees* together, may with little trouble, and in much less time, inform himself. By the same means, some *Conjunctions* which seem to be strange, as *Quince* and *Pear*, *White Thorn* and *Medlar*, &c. do yet, by the response of their *Parts*, as well as by *Experience*, appear to be good. And there is no doubt, but that many *Conjunctions* not yet tryed, or not known to have been so, may upon the same ground, be tryed with good success.

15. §. The chief Use of *Grafting* and *Inoculation*, is, That they *Accelerate* the growth of *Good Fruit*. The *Cause* whereof, is the *Knot*, which is always made in the *Conjunction*. By means of which, all the *Sap* is strained, and so ascendeth up into the *Graff* or *Bud*, both *Purer* and in less *Quantity*; and is therefore better and sooner concocted. Hence, the smaller the *Fruit* of any *Tree*, though it be not the best, yet the *Sap* being there, in less *Quantity*, is the sooner ripe. On the contrary, where the *Sap* ascendeth too freely, it doth not only retard the growth of the *Fruit*, but produceth *Barrenness*; as is seen in those *luxuriant Branches*, where it runs all up to *Leaves*. Hence also *Vines*, by *Bleeding*, become more *Fruitful*: that is, by the Effusion of *Part* of the *Sap*, there is a more easier *melioration* of that which remains. Even as *Phlebotomy* doth oftentimes produce a more healthful and better *Habit* of our own *Bodies*. To conclude, the lessening the *Quantity*, and thereby the *melioration* of the ascending *sap*, by *Knots*, is *Natures* own contrivance; as is seen in *Sugar-Cane*, *Corn*, and other *Plants*.

THE
ANATOMY
OF
LEAVES, FLOWERS,
FRUITS and SEEDS.

In Four Parts.

The FOURTH BOOK.

By *NEHEMIAH GREW* M.D. Fellow
of the *ROYAL SOCIETY*, and of the
COLLEGE of *PHYSICIANS*.

LONDON,

Printed by *W. Rawlins*, 1682.

THE
CONTENTS.
OF THE
First Part.

CHAP. I.

Of the Protections and Folds of Leaves.

CHAP. II.

Of those Things which appear upon the Surface of the Leaf.

CHAP. III.

Of the Figures of Leaves; and the Apparent Position of the Fibres.

CHAP. IV.

Of the Parts and Texture of the Leaf.

CHAP. V.

Of the Duration of Leaves, and the Time of their Generation.

CHAP. VI.

Of the Manner of the Generation of the Leaf. Where also, that of the Two General Parts of a Plant, sc. the Lignous and Parenchymous, is further explain'd.

To

To the Honourable

Robert Boyle Esq;

S I R,



AFTER I had finished the foregoing Books, In which, I conceive, as far as Glasses will yet lead us, I have clearly Describ'd and Delineated the Structure of a Plant; and have endeavour'd, in some part, to Unfold the Reason and Scope of Nature therein: I was willing to sit down, and leave what remained, to the Improvements of the Present and Succeeding Ages.

But in Discourse upon this Subject, You have been pleased frequently to insist, That I should by no means omit, to give likewise, some Examples of the Mechanisme of Nature in all the other Parts. The Performance whereof therefore, next to the Obedience I owe to the Royal Society, is to be looked upon, as a Due to the Authority which Your Judgment hath over me.

*This I have said, that, if what is herein done, shall prove acceptable unto Learned Men; they may know, To whom they are oncemore to give their Thanks: After they have so often done it, upon (a better score) the Publishing of Your own Excellent Works. In which, there seems to be a Question, Whether Your Continual Endeavours, to enlarge the Bounds of Natural Knowledge, or Your Successes therein, have been the Greater: So that, whereas Nobility in some, doth
only*

The Epistle Dedicatory.

only serve to lift them, like Jupiter's Satellites, out of sight: You, by giving a greater Light, have drawn all Mens Eyes upon You. And whilest there are many, in all Ages, fond of Preheminency in the Conduct of Popular Affairs; who yet rarely hit the Mark they aim at; or aim at That they pretend: You have thought fit, rather to separate Your Self, to that more Innocent, and more Noble Sort of Wisdom, which lieth, not in the Arts of Concealing, but in Discovering, the Truth of Things.

That we may have many to imitate You herein, cannot but be heartily wish'd by all, who regard the Honour of their own Country; as it is, with much Zeal, by

Sir,

Your most obedient

Servant

NEHEMIAH GREW.

THE

THE
ANATOMY
OF
LEAVES,

PROSECUTED

With the bare EYE,

And with the

MICROSCOPE.

Read before the *Royal Society*, Octob. 26. 1676.

PART I.

CHAP. I.

Of the Protections and Folds of Leaves.



IN THE *General Anatomy of Plants*, I have assigned one whole *Chapter* (a) to the *Germen* and (a) *Lib. i.*
Leaf. Since then, I have occasionally made divers *Ch. 4.*
Remarques of the same; both with the *Naked*
Eye, as there, and also with the *Microscope*. The
Principal whereof, I shall here set down; without
repeating any from thence; or obliging my self
strictly to the *Order* there used.

2. §. That which in a *Germen*, first occurs to the *Eye*, is the *Pro-*
tection of the *Leaves*, or the various *Methods* which *Nature* takes to
preserve them from the *Injuries* both of the *Ground*, and of the *Wea-*
ther. To the *Instances* formerly given, I shall add these that follow.

3. §. AND First, it is observable of the young *Buds* of *Ammi*, that lest they should be bruised, or starved, upon their first *Eruption* from under the Ground; they are couched, as *Firn* is rowl'd, inward; each *Bud*, against the *Base* of the *Stalk* of the foregoing *Leaves*, and most exactly laid up within the *Membranes* thence produced: Just as the Child in the Womb, lies with his *Head* against his *Knees*; or as it is afterwards embraced with the *Armes* of the Nurse. And it is a general Rule of Nature, where the *stalks* of the *Leaves* are so long, that they cannot lap one over another, and where no other special *Protection* is provided; for the bottoms of the *stalks* to be produced into broad *Membranes*, as Blankets to the succeeding *Buds*; as in *Crowfoot*, *Dovesfoot*, *Claver*, *Cransbill*, *Strawberry*, *Tarrow*, and others. And sometimes instead of two *Skins* lapped one over another, there is one entire *Skin*, produced from the *Stalk*, in which as within a *Secundine*, the *Bud* is safely shrowded; and which, in its Growth, it gradually breaks open.

Tab. 41.

4. §. THE same is also observable in *Dock*, *Sorrel*, *Bistort*, and all other *Plants* of this Kindred; with this difference, That every *Veil* or *Secundine* is not here produced from the *Stalk* of the *Leaf*; but hath its *Original Distinct* from it. And whereas in the former, every *Bud* hath only one to it self: in these *Plants*, every lesser *Leaf*, together with its own proper *Veil*, is always inclosed, with the next greater *Leaf*, in another *Veil* common to them both; and both these with the next, in another; and so on to the greatest. These *Veils* are extream thin, and have very few *Vessels*; being so many meer transparent *Skins*. For which reason, there is always found a *Mucilage* or clear *Gelly*, between every *Leaf*, and its *Veil*, and between *Veil* and *Veil*. The one, thus preserving the other, (as do the *Humors* and *Membranes* of the Eye) from drying and shrinking up, and thereby from becoming useles for the *Protection* of the *Plant*.

5. §. THE *Orchis*, and other *Plants* of this kindred, because they *Spring* and *Flower* early, when the mornings are cold, have a double *Sheath*, or *Blanket* over all. The *Buds* of some *Herbs* (as of *Plantain*) having no *Hairs* growing on them, are covered with *Hairy Thrums*. And the *Nettle* hath *Bastard-Leaves*, or *Interfoyls* between *Leaf* and *Leaf*, for the preservation of its *Stings*.

Tab. 41.

6. §. ANOTHER Sort of *Protection* is seen in *Wild Clary*, *White Archangel*, and other *Plants* of a like Shape. In which, the greater *Leaves* do still cover and inclose the lesser, not by being lapped over them, as where the *Leaves* are more numerous, is usual; but by a *Double Fore-Curl* at the bottom of every two greater *Leaves*; by which the little *Under-Bud* is embraced, and so kept safe and warm.

Tab. 42.

7. §. THE *Leaves* of *Onions* are all *Pipes* one within another. These *Pipes* are every where entire, saving about the middle, where they have a small *Aperture*; common to all of them, even the most minute in the Centre: not being a forced *Crack*, but a *Door* originally formed, for the issuing of every lesser *Pipe*, out of a greater.

Tab. 41.

8. §. THE LAST I shall give, is that which is remarkable in *Common Sumach*. The *Buds* whereof, being exceeding tender, Nature appears sollicitous in a peculiar manner, for their preservation. For whereas in other *Plants*, they are well enough secured only by standing behind the *stalks* of the elder *Leaves*: here they are lodged within the
very

very *Body* of the *Stalk*; as entirely, as a *Kernel* is within an *Apple*, or a *Fœtus* in the *Womb*. From whence it comes to pass that the *Basis* of every *Stalk* is extremely swelled, as going Great with a *Bud*.

9. §. UPON THE removal of those *Parts*, which are contrived for the *Protection*; the *Foulds* and *Composture* of the *Leaves* do next appear: all which are most aptly suited both to the *Number* and *Shape* of the *Leaves*, and also their *Position* upon the *Branch*. In the *First Book* (a) I have given Examples of these *Eight Sorts*, sc. the (a) *Ch. 4.*
Plain Lap, the *Plicature*, the *Duplicature*, the *Multiplicature*, the *Single Roll*, the *Double Back-Roll*, the *Double Fore-Role*, and the *Treble-Tab. 42.*
Roll. To which I shall add *Four* or *Five* more.

10. §. And *First*, in some *Plants*, as *Ground-Ivy*, *St. Johns Wort*, and divers others, where the *Leaves* are small, pretty numerous, and grow by pairs, they have no *Fould*, but stand Flat and Tangent, like a pair of *Battledores* clapt together.

11. §. They have the like *Posture* in *Baum*; saving, that here the *Edges* of the *Leaves* are a little curled backward. Not *Rolled*, a *Curl* being but the beginning of a *Roll*. So the several *Labels* of a *Grounsel-Leaf* are all laid in a *Back-Curl*.

12. §. The *Leaves* of some *Plants*, as *Horehound*, *White Lamium*, *Nettle*, and others; are likewise only Tangent, but are set with a *Fore-Curle*. And the several *Labels* or *Scallops* of the *Leaf* of *Common Crowfoot*, are all *Curled Inward*. But those of *Hepatica aurea*, are composed into *Double Fore-Rolls*.

13. §. THE *Leaves* of *Sage*, *Scabious*, *Red Lamium*, *Lychinis Sylvestris*, and others, are neither couched one over another, as in the *Bow-Lap*; nor plated, as in the *Flat Lap*; but being loosely foulded, of every pair of *Leaves*, the half of one is reciprocally received between the two halves of another, and may therefore be called the *Cleep. Tab. 42.*
A *Position* very well suited to the *Smallness* of their *Number*, and the *Equality* of their *Size*, not so well agreeing with the *Bow-Lap*; and the somewhat inward *Posture* of the *Fibres*, not allowing the *Flat Lap*. Sometimes, as in *Syringa*, where the *Leaves* are broader, the *Cleep* is joyned with a *Fore-Curle*.

14. §. THE last I shall mention, is the *Plait-Roll*, as in the *Lapathum Alpinum*, which some call *English Rhubarb*. The *Leaves* whereof are so very large, and the *Fibres* so prominent; that besides and under the two *Back-Rolls*, they are also laid in several *Plaits*, and under those *Plaits*, again with lesser ones, all most exquisitely Tucked up between the said *Fibres*: So, as neither to bruise the same, nor yet to leave any *Vacuity*: whereby every *Leaf*, and the whole *Bud*, lie close and round within their *Veils*.

C H A P. II.

Of those things which appear upon the Surface
of the Leaf.



THESE are Globular Excrecences, Spots, Hairs, Thorns and Prickles: of all which, except Spots, I have spoken in the Appendix to the Chapter of Leaves in the First Book.

Tab. 43.

2. §. Of the Globulets, it may here be further noted, That those which are white, and lie sometimes like a fine Powder upon the Leaf, were once transparent, as in Bears-Eear; their clear Liquor being now evaporated to an Extract or White Flowers. This, if licked off, will give you the Taste of the more Essential Content of the Plant; different from that perceived in chewing the Leaf.

3. §. For the observing of them, it may also be noted, That although they often grow on both sides the Leaf alike; yet sometimes, as in Ground-Ivy, only or chiefly on the Back-Side. And that in many Plants, where the elder Leaves have none; on the young Buds they are very numerous; as in Corin Tree, Sorrel, and others.

Tab. 43.

4. §. AS for spots, the smaller ones are observable not only in St. Johns-worts, (in which Plant only they are commonly taken notice of) but also in Rue, Ground-Ivy, Pimpernel or Anagallis, and divers other Plants, when held up against the Light. The original whereof seems to be, at least in some, from the Globulets above mentioned; that is, when they break and dry away. So the Spots of Rue-Leaves, which in the Reflection of Light look black, but upon the Trajection thereof are transparent, are so many little Holes, pounced half way through the thickness of the Leaf, and seem as made, by the breaking and drying away of as many Globulets. Whence also, as the Globulets are best seen in the younger Leaves, so these Spots in the elder.

Tab. 43.

5. §. BESIDES these, and some others (as those in Ladies-Thistle) which are Natural to the Leaf; there are also some Spots, or rather Streaks, which are Adventitious; as those in the Leaves of Sonchus. The Cause whereof, is a small flat Insect, of a grey Colour, and about $\frac{1}{8}$ th of an Inch long. Which neither ranging in breadth, nor striking deep into the Leaf; eats so much only as lies just before it, and so runs scudding along betwixt the Skin and the Pulp of the Leaf; leaving a whitish Streak behind it, where the Skin is now loose, as the measure of its Voyage.

6. §. THE Original and several kinds of Thorns, I have describ'd in the above said Appendix. I only add, that the very Leaves of some Plants, if they stand till the second year, are changed into so many Thorns, as in the Furz.

7. §.

7. §. They are of Use, not only for the *Protection* of the *Bud*; but likewise, for the support of the *Plant*; as is observable in those *Climbers*, which are neither strong enough to stand of themselves; nor yet, from their fragility, are capable of winding about another, without being torn all to pieces. For which end also, these *Thorns* grow not like *Buds*, erected; but poynt all downwards, like so many *Tenters* or *Hanging-hooks*: as in the *Bramble*, chiefly on the *stalks*; and in *Clivers*, also on the *Leaves* themselves; whereby they catch at any Thing that stands next them; and so, although such slim and feeble *Plants*, yet easily climb to a very great height. Tab. 43.

8. §. OF THE several *Figures* of *Hairs*, and their Use, I have *B. 1. Ch. 4.* also spoken. As to one Use, *sc.* the *Protection* they give to the *Leaf*, Tab. 43. I shall here further note, That the design of *Nature*, is the more evident if we consider, That all *Leaves* are not alike *Hairy*, nor at all times, nor in every part: but differently, according to their *Age*, *Substance*, *Texture*, and *Foulding* up. Their *Age*; for there are many young *Buds* covered with a thick warm *Hair*, which afterwards dries up and disappears, as useless; as those of the *Vine*, *Golden Liverwort*, &c. Their *Substance*; so those *Buds* which are tenderest, and would sooner feel the cold, if naked, have the fullest *Hair*; as of *Thistle*, *Mullen*, *Burdock*, and others. Their *Structure*; therefore those *Leaves*, whose *Fibres* stand more prominent or above their *Surface*, lest the cold should nip them, are covered with greater Store of *Hair*; as in *Moth-Mullen*, *Garden-Clary*, and the like. And their *Fould*; it being observable, That those *Leaves* which are folded up inward, have little or no *Hair* on their inner, but only on their *Back-Sides*, which are open to the *Aer*; as is visible in *Corin*, *Warden*, *Golden Liverwort*, and others.

9. §. Add hereto, That where there is Store of *Hair*, *Nature* is the less solicitous for other *Covers*; and where there is not, she is more. So the *Leaves* of *Beans* and *Peasen*, of *Nettle*, *Plantain*, &c. not being *Hairy*, have each a *Surfoyl*, or else certain *Hairy Thrums*, to protect them. And those *Plants* which have neither, are such as have a *Hotter Juice*, and so less subject to the impressions of *Cold*, as *Speerwort*, *Scurvygrass*, *Watercress*, *Fenil*, and most of the *Umbelliferous* Kind.

10. §. *Hair* is of use to preserve young *Buds*, not only, from the cold *Aer*, but also from too much *Wet*; which, if it were contiguous, especially in *Winter*, would often rot and destroy them. But being made to stand off in drops at the ends of the *Hair*, doth not hurt, but refresh them. Thus doth *Nature* make the meanest Things sometimes subserve to the best Ends.

CHAP. III.

Of the Figure of the Leaf; and the Apparent Position of the Fibres.



H A T which in the *Leaf* offers it self next to be observed, is its *Figure*. This is infinitely varied with the several *Kinds of Plants*: and there are some, which have *Leaves* (besides the two first *Dissimilar* ones) of Two *Kinds* or Two distinct *Figures*; as the *Bitter-Sweet*, the common *Little Bell*, *Valerian*, *Lady-smocks*, and others. For the *Under Leaves* of *Bitter-Sweet*, are Entire; the Upper, with two *Lobes*: the *Under Leaves* of the *Little Bell*, like those of *Pancy*; the Upper, like those of *Carnation*, or of *Sweet-William*. And in some *Plants*, Nature affecteth a *Kind of Irregularity*; the *Leaves* whereof are of no one certain *Figure*; as in *Dragon*, *Peony*, *Bishops-Weed*, &c.

2. §. B U T the *Leaves* of most *Plants*, have a *Regular Figure*; and this *Regularity*, both in *Length* and *Circuit*, always defineable. In *Length*; by the *Proportion* between the several *Leaves* upon one *Stalk*, or between the several *Lobes* upon one *Leaf*. So the *Leaves* of *Clematis sylv. major*, which stand by Ternaries, shorten by equal *Proportions*, that is to say, if, the chief *Fiber* of each, be divided into equal *Parts*; their several *Lengths* are not as Ten, Eight, and Four; but as Ten, Eight, and Six. So the *Lobes* and *Fibers* of *Clematis Virginiana* *Hederæ folio*, of *Artemisa*, &c. shorten in like manner by equal *Proportions*. The same is observable in measuring, upon a *Gooseberry-Leaf*, from the *Poynt* of the first *Lobe*, to the first *Angle*; from thence, to the second *Poynt*; from thence, to the second *Angle*; and from thence to the third *Poynt*.

Tab. 46.

3. §. B U T in many, the *Proportion* is different. So in the *Leaves* of the *Lesser-Maple*; the shortning of the *smaller Lobes*, with respect to the middlemost; is not Equal, but Double to that of the middlemost, with respect to the Greater. For if their chief *Fibres* be divided into Equal *Parts*, they are as Eleven, Nine, and Five. On the contrary, in the *Leaves* of *Althæa fruticosa Pentaphylloidea*, the middlemost *Lobes* shorten by a greater *Proportion* than the Least; all three being as Ten, Fourteen, and Twenty.

4. §. W I T H respect to the *Circumference*, the *Figure* of most *Leaves* is very Complex. Yet Two things are evident. First, that all *Regular Leaves*, are defined or measured out by *Circles*; that is, by the *Arches* or *Segments* of several *Circles*, having either the same, or divers *Centers* and *Diameters*. Secondly, That the *Length* of the *Leaf*, or of the chief *Fiber* thereof, is the *Standard Measure* for the *Diameters* of these *Circles*: these being either its full *Length*, or certain equal parts subtracted, or multiplied; as half its *Length*, or its *Length* and half, &c.

5. §.

5. §. TO make this appear, I shall give several Instances: of some, where both the Edges are of one Measure; and of others, where they are different. And of both kinds, where they are measured by fewer and where by more *Circles*.

6. §. The *Leaf* of *Lagopus major fol. pennat.* is measured by One *Circle*, the same on both Edges, whose *Diameter* is Thrice the Length of the *Leaf*.

7. §. That of *Syderitis Salviae fol.* by Two *Circles*: the *Diameter* of the Lower, being Twice the Length of the *Leaf*; of the upper, the Length and half. In both these the *Circles* are drawn Outward; that is, with their Centers some where upon the middlemost or chief *Fiber* of the *Leaf*. Tab. 44.

8. §. That of *Orange-Tree*, is also measured by Two *Circles*: but one of them repeated with *Opposite Centers*. That next the *Cone* of the *Leaf*, is drawn Inward; that is, with the *Center* no where upon the *Leaf*, but without it. The *Diameter* hereof is just the Length of the *Leaf*. The middle part of the Edge is measured by the same *Circle*, only drawn Outward. The lower *Circle* next the *Stalk*, is drawn Inward, as the upper; and its *Diameter* Three times the Length of the *Leaf*. Tab. 44.

9. §. The *Leaf* of the *Venetian Vetch*, is measured by Three *Circles*. That next the *Cone*, drawn Inward; the *Diameter* whereof, is Twice the Length of the *Leaf*; the next is drawn Outward; whereof the *Diameter*, is just the Length. The third or lowermost, is drawn also Outward; and its *Diameter*, half the Length. So that they all lessen by an Equal Proportion. Tab. 44.

10. §. The *Leaf* of *Great Laserwort*, is also measured by Three *Circles*; all drawn Outward, and one of them Repeated. The *Diameter* of that next the *Cone*, is Half the Length of the *Leaf*; of the next, Thrice the Length; of the Third, just the Length; the lowermost, is the same with the First. Tab. 45.

11. §. That of *Broad Leav'd Laserwort*, is also measured with Three *Circles*; and one of them repeated with *Opposite Centers*. The *Diameter* of the First, is Half the Length of the *Leaf*; of the Second, Twice the Length; of the Third, just the Length: all of them drawn Outward. That next the *Stalk*, is the same with the First; only drawn Inward. Tab. 44.

12. §. The *Figure* of the *Leaf* of the *Cornelian Cherry*, is exactly that of the foregoing, Inverted: the same measure there beginning at the *Base*, and ending at the *Cone*; which here begins at the *Cone*, and ends at the *Base*: as by comparing their Draughts together may be observ'd. Tab. 44.

13. §. IN ALL, the foregoing Examples, both the Edges of the *Leaves* have the same Measure. But they have oftentimes, different ones; as in these that follow.

14. §. The *Leaf* of *Althæa fruticosa*, is measured by Three *Circles*. The left Edge (as the *Leaf* lies with the backside upward) by One *Circle*, but Twice repeated. For the *Diameter* of the First, is the Length of the *Leaf*; the Second is the same, but drawn upon another Center; the Third also the same, but drawn Inward. The right Edg, is measur'd by Two *Circles*: the *Diameter* of the First, being the Length of the *Leaf*; of the Second, Half the Length. Tab. 45.

15. §.

15. §. That of *Black Poplar*, by Three; and each Edge by Three repeated. On the left, the *Diameter* of the First, is the Length of the *Leaf*; of the Second, Half the length; of the Third, the Length and Half. The *Measure* of the right Edge, is that of the left, Inverted: the same *Measure* there beginning at the *Base*, and ending at the *Cone*; which here begins at the *Cone*, and ends at the *Base*.

16. §. That of *Doronicum*, is measured by Three *Circles*, whereof, one is repeated Once; and another Thrice. The right Edge by Two, and One repeated. For the *Diameter* of the First or that next the *Cone*; is the Length of the *Leaf*; the next is the same, but drawn Outward; the *Diameter* of the Third, is Half the Length. The left Edge, by Three *Circles*; whereof One is repeated on the same Edge, and Two, the same, as on the other. For the *Diameter* of the first, is the Length of the *Leaf*; of the Second, Four times the Length; the Third, the same as the First; and of the Fourth, Half the Length.

17. §. Lastly, that of *Mountain Calamint* is measured by Four *Circles*. The left Edge, by Three *Circles*, of which, the lowermost is once repeated: the right Edge also by Two; whereof the nether is likewise once repeated.

18. §. It may seem, even from these Instances, no very unobvious Conclusion; That all *Crooked Lines*, *Spiral*, *Helick*, *Elliptick*, *Hyperbolic*, *Regular*, or *Irregular*; are made up of the *Arches* of *Circles*, having either the same, or divers *Centers* and *Diameters*. And, as otherwise, so from the *Contemplation* of *Plants*, men might first be invited to *Mathematical Enquirys*.

19. §. TOGETHER with the *Figure* of the *Leaf*, the *Position* of the *Fibers*, as it is apparent before *Dissection*, is observable; especially on the back of the *Leaf*. Whereof I shall add, to what I have said in the *First Book*, the following *Remarques*.

20. §. First, that there are some *Leaves*, in which the first *Collateral Fibres* make *Right Angles* with the Great one in the middle: as the *Great-Maple*, the *Great Celandine*, *Chondrilla*, and the rest, or many, of the *Intybous* Kind; with some few others. But that generally all the chief *Fibers* of a *Leaf*, make *Acute Angles* together: both where they stand collateral with the middle *Fiber*, as in *Strawberry*; and where they all part at the *Stalk*, as in *Mallow*.

21. §. Again, that of these, there are some few, any two of whose *Defining Fibres* making two *Rays* of equal Length, take in One Eighth Part of a *Circle*, as in *Mallow*; and in some one Tenth: but in most they take in either one Twelfth part, as in *Holy-Oak*; or one Sixth, as in *Sirynga*. So that where the *Fibres* stand Collateral with one in the middle, if you suppose them to be drawn out at *Opposite Angles*; or where the chief *Fibers* part at the *Stalk*, you only take in the *Stalk*; you will thereby divide a *Circle* into Eight, Twelve, or Six equal Parts; as in *Sirynga*, the *Vine* and others. And so likewise, where there are several *Sprigs* upon one *Stem*, as in *Fenil*, *Hemlock*, and the like: as will best be understood by the *Figures*.

C H A P. IV.

Of the Parts and Texture of the Leaf.

COME next to observe the several *Parts*, whereof the *Leaf* is composed: and first the *Skin*. This being stript off the *Leaf*, although to the bare Eye it looks no otherwise than a *skin* of *Isinglass*: yet being viewed through a good *Glass*, with a clear and true *Light*, and in an advantageous *Position*; it appears to consist not only of *Organical Parts*, as do the *Skins* of *Animals*; but these also Regularly mixed together; that is, of *Parenchymous* and *Lignous Fibres*, all very curiously interwoven as it were, into a piece of admirably fine white *Sarcenet*: as in *Flag*, *Tab. 48.* *Tulip*, and the like.

2. §. From hence, it is easy to conceive how the *Skins* of all *Plants*, as well as those of *Animals*, are perspirable; *sc.* between the several *Fibers* of which they consist. But as the *Skins* of *Animals*, especially in some *Parts*, are made with certain open *Pores* or *Orifices*, either for the Reception, or the Elimination of something for the benefit of the *Body*: so likewise the *Skins*, of at least many *Plants*, are formed with several *Orifices* or *Pass-ports*, either for the better *Evolution* of *Superfluous Sap*, or the *Admission* of *Aer*.

3. §. THESE *Orifices* are not in all *Leaves* alike; but varied in *Bigness*, *Number*, *Shape*, and *Position*: Serving to the different *Nature* of the *Plant*, or *Leaf*; and giving the *Leaf*, as it were, a different *Grain*. *Princes Feather*, *i. e.* a Sort of *Sanicle*, they stand only on the *Edges* of the *Leaf*; but are very ample. In the *White Lily*, they are *Oval*, very white, and each surrounded with a slender white *Border*. They stand about a 6th or 8th part of an *Inch* distant, as they appear through *Tab. 48.* a good *Glass*, all over the *Leaf*, but not in any regular *Order*. These *Orifices* are the cause of the *Greyish Gloss* on the upper side the *Leaf*: for the *Back-side*, in which there are none of them, is of a dark *Sea-Green*.

4. §. In the *Leaf* of *Pine*, they are also *Oval*, and about the same *Bigness* and *Number*, as in that of a *Lily*; yet without a *Border*. But their *Position* is very *Elegant*, standing all, most exactly, in *Rank* and *File* from one end of the *Leaf* to the other. *Tab. 48.*

5. §. NEXT TO the *Skin*, lies the *Pulpy* part of the *Leaf*; which by the same latitude, as *Use* hath taught us in many other *Words*, I call the *Parenchyma*. This *Parenchyma* or *Pulp* of the *Leaf*, like the *Pith*, and all other *Parenchymous Parts* of a *Plant* is made up of incomparably small *Cylindrick Fibres*: and these *Fibres*, in most *Leaves*, wov'n and woun'd up into little *Bladders*.

6. §. The *Bladders* are here of several *Sizes*, as in the *Pith*: but generally more visible in the *Stalk*, than in the *Body* of the *Leaf*. *Varied*, as in the *Pith*, so here, not according to the *Size*, but the *Nature* of the *Leaf*: So in *Common Dock*, and *Moth Mullein*, both Great *Tab. 49.*

Tab. 50. *Leaves*, they are Small ; in *Wild Clary*, a Lesser *Leaf*, they are very Large. In the *Body* of the *Leaf*, sometimes the *Sides* of the greater *Bladders*, are made up of lesser ones ; as in *Borage*.

7. §. In some *Leaves*, these *Parenchymous Fibres* are all drawn close up together. In the Former, they are as the *Threds* in the *Open-work* of *Bone-Lace* ; in These, as the same *Threds*, in the *Cloth-work*.

8. §. The *Pithy Part*, in the *Stalk*, and almost up to the *Top* of the chief *Fiber*, in many *Leaves*, is *Tubular* ; even whilst they are yet *Young* and *Sappy* : as in *Sweet Chervil*, *Hemlock*, *Endive*, *Cichory*, *Lampfana*, *Dandelion*, *Burdock*, *Daisye*, *Scorzoner*, and others. And sometimes the said *Pithy Part* is opened into several little *Pipes*, like so many *Aer-Vessels*, above $\frac{1}{2}$ a *Foot* long ; as in the *Common Dock* and the *Little spurge*, by some called *Wart-Wort*.

9. §. THE *Strings* of the *Leaf*, or those *Fibres* which are visible to the bare *Eye*, are composed of *Vessels* of the Two General Kinds, *sc*, for *Sap*, and for *Aer*. They are jointly distributed throughout the *Leaf* : Yet not so, as to run meerly parallel ; as in *Animals*, every *Artery* hath its *Vein* : but the *Aer-Vessels* are every where Inclosed, or as it were *sheathed* in the *Sap-Vessels*.

Tab. 49. 10. §. THEIR *Position* is various and regular, not only in the *Body* of the *Leaf*, as is above shewed ; but likewise in the *stalk* : of which also I have given several Instances in the *First Book*. I shall here note, and more particularly describe, One or Two more. In the *Stalk* of a *Mallow-Leaf*, they stand in Six Oblong *Parcels* of equal Size, and in a *Ring* near the *Circuit*. Whereby the *stalk* is stronger, the *Growth* hereof, before and behind, more equal, and so the posture of the *Leaf* more erect.

Tab. 49. 11. §. In *Dandelyon*, they stand in *Five Parcels* : of which the Greater stands a little *behind* the *Centre* of the *Stalk* ; figured into a very small *Half-Moon* or *Semi-Tube*, whose *Diameter*, through a *Glass*, is not above $\frac{1}{4}$ th of an *Inch*. The other Four, are extream small *Cylinders*. Altogether make an *Angle*, twice as big as that of a *V* Consonant. Whereby, although the *stalk* be strong enough to support the younger *Leaves* ; yet those which are grown longer, and so not only by their *Bulk*, but their farther *Extension* from the *Center* of *Gravity*, are become more weighty ; commonly lie flat on the *Ground*.

Tab. 49. 12. §. In *Wild Clary*, they stand also in *Five Parcels*, the Greater stands not *behind*, but *before* the *Center* ; making an *Arch*, whose *Chord* in a *Glass*, is above $\frac{1}{2}$ an *Inch* long ; and belongeth to a *Circle*, whose *Diameter* is an *Inch* and half. The other Four, are small *Cylinders*, also different from those in *Dandelion* ; the two bigger, there standing hindmost ; but here, the two Less, and the two Bigger, within the two round *Ridges* of the *Stalk*.

13. §. From hence it is, that the *Leaves* of this *Plant* have not only a *Prone* or *Horizontal Posture*, but also make that *Forceable Pressure* on the *Ground*, which can by no means be imputed to their *Weight*. For the *Great Arched-Fibre* standing before the *Centre* of the *Stalk*, and the two Longer Round ones being uppermost, in the *Ridges* of the *Stalk* ; they put on the upper parts thereof to a more full and forward *Growth*, and so to bow the *Leaf* back-ward. And the *Fibrous Arch* being, though broad, yet almost flat, doth hereby the more easily yield to that *Motion*.

14. §.

14. §. In *Borage*, and *Moth-Mullen*, they stand also in *Five Parcels*. In the former, the largest maketh still a more bulky *Arch*; than that of *Clary*; being thicker, as broad, and of a lesser *Circle* or more bowed. But in *Mullein*, it maketh almost an entire *Oval*. Tab. 49.

15. §. By means of this *Figuration*, a sufficient number of *Vessels* for such large *Leaves*, are not only more conveniently *Distributed* into them; but also stand more safely in the *Stalk*. For were the *Arch* contracted into a *solid Cylinder*, it could not so presently be resolved into small *Fibers*. And were it laid into a flat *Plate*, or straight out, either the *Figure* of the *Stalk*, and so of the *Leaf*, must be altered; or else, the two ends of the *Plate*, would come too near the *Circumference* of the *Stalk*, and so be more liable to the *Impressions* of the *Weather*: as may be observed in cutting the *Stalk* transversly, and by the *Figures*.

16. §. IN the *Body* of the *Leaf*, besides the *Positions* of the *Fibrous Strings* or *Threds*; above expessed; there is one *Thred*, bigger or less, which in all *Plants*, runs round the *Edge* of the *Leaf*, and hems in all the rest; but can hardly be well observed in any, without stripping off the *skin* of the *Leaf*. When the *Fibres* of the *Leaf*, are bigger, or less tender, as in *Holly*, the *skin* and the *Pulp* are sometimes found either rotted off, or eaten away with *Insects*; whereby, both the said surrounding *Fiber*, and the rest, are all very fairly visible. Tab. 50.

17. §. THE *Vessels* seem to be continu'd, in the *Leaf*, by being *Ramified* out of *Greater* into *Less*, as *Veins* or *Arteries* are in *Animals*. But if the *Skin* and *Pulp* of the *Leaf*, as suppose a *Borage-Leaf* be taken off, and the *Vessels* laid bare; by the help of a good *Glass*, it will appear; That they are all of the same *Size*, every where in the *Leaf*; and also continued throughout the same, all several and distinct *Pipes* one from another, as the *Threds* in a *Skein* of *Silk*. And that therefore the *Distribution* of the *Threds* which the *Vessels* compose, is not the *Ramifying* of *Greater Pipes* into *Less*; but the dividing a greater *Cluster* of *Pipes*, into several lesser *Clusters*, till at last they come to be single; as in the *Distribution* of the *Nerves*. Tab. 50.

18. §. The *Vessels* seem also to be *Inosculed*, not only side to side, but the ends of some into the *Sides* of others. But neither is this ever really done: the lesser *Threds*, being only so far diducted, as sometimes to stand at *Right-Angles* with the greater. So that they are *Inosculed* only *End to End* or *Mouth to Mouth*, after they are come at last to their final *distribution*. Tab. 50.

19. §. The *Aer-Vessels*, are not only, as is said, *Existent* in the *Leaves* of all *Plants*; but are herein also discoverable without the help of *Glasses*: For upon breaking the *Stalk* or chief *Fibers* of a *Leaf*; the likeness of a fine *Woolly* Substance, or rather of curious small *Cobwebs*, may be seen to hang at both the broken *Ends*. This is taken notice of; only in some few *Plants*, as in *Scabious*, where it is more visible. But may also be seen more or less, in most other *Plants*, if the *Leaves* be very tenderly broken: as I have noted near twenty years since; and thence conjectur'd them a Sort of *Vessel* common to *Plants*. Now this fine *Wool*, is really a *Skein* of *Aer-Vessels*, or rather of the *Fibers* of the *Aer-Vessels*, unroaved from their *Spiral Position*, and so drawn out in *Length*. As they appear thus unroaved and drawn out at *Length*, both to the bare *Eye*, and through a good *Microscope*, I have represented in two *Exemples*, the one a *Scabious Leaf*, the other that of a *Vine*. Tab. 51, & 52.

20. §. THE *Westage* of the *Strings* and *Parenchymous Fibers* together, is here made in the same manner, as hath been described in the *Anatomy* of the *Root*, and *Trunk*: the former being in some Sort as the *Warp*, the latter as the *Woof* of the *Leaf*.

21. §. And one Example we have (it may be more than one) wherein Nature shews, though not a greater, yet a different *Art*; and that is the *Palm-Net*. For whereas in other *Plants*, the *Webb* is made betwixt the *Lignous-Strings* and the *Fibers* of the *Parenchyma*, only visible through a *Microscope*: here the said *Strings* themselves are Interwoven, and the *Westage* apparent to the bare Eye. Of these *Palm-Nets* or *Sacks*, there are several Sorts. One of them is composed in this manner. It hath a *Fivefold Series* of *Lignous Strings* or *Fibers*. The greatest whereof swell out above the rest; and like so many *Ribs*, are obliquely produced on both hands, so as to encompass the *Sack*. Along each of these *Ribs*, on the inside the *Sack*, runs a small *Whitish Line*; being a *Thread* of *Aer-Vessels* growing thereto. Betwixt these *Ribs* or larger *Strings*, there are others much less, Two or Three betwixt *Rib* and *Rib*, *Parallelly* interjected. On the inside, there is a *Third Series*, which is also obliquely produced; but transversely to the former. The *Fourth* and *Fifth*, consist of the smallest *Strings*; not only *Transversely* produced, but also *Alternately*, from the outside to the inside of the *Sack*, & *vice versa*. By these two last, all the rest are most elaborately woven into one entire and strong piece of Work.

CHAP. V.

Of the Duration of Leaves, and the Time of their Generation.



N *Evergreen*, is one degree above a *Plant* which is simply *Perennial*: of This, only the *Trunk* and *Buds* live all the *Winter*; of That, also the *Expanded Leaves*. And an *Evergrow*, is a degree above an *Evergreen*: here, the *Buds* and young *Sprigs*, do only live; there, they grow and are put forth.

2. §. An *Evergreen*, is made such, either by the *Toughness* of the *Skin*, and *Closeness* or *Density* of the *Parenchyma*, whereby the *Leaf* is better able to endure *Cold*; as in *Holly*: or by the extream *Smalness* or *Fewness* of the *Aer-Vessels*, whereby the *Sap* is less dried up, and so sufficient, even in *Winter*, for the *Nourishment* of the *Leaf*; as in *Box*, and *Tew*, as also *Fir*, and all *Resiniferous Plants*.

3. §. The perpetual *Growth* of a *Plant*, seemeth to depend chiefly on the *Nature* of the *Sap*. For all *Juyces* will not ferment alike, nor with the same degree of *Heat*. So that whereas many *Plants* require a greater *Heat*, as that of *Summer*, for the fermenting and distribution of their *Juyces*, and so their growth; the *Warmth* of *Spring* is sufficient for many others; and for some few, that of *Winter* it self.

4. §. AS TO the *Time* wherein the *Leaves* are formed; First, it is very probable, That in those *Plants* which have *Leaves* (besides the *Dissimilar*) of Two distinct *Figures*, as hath the *Little Common Bell*,

Bell, and some others; the *Under-Leaves*, which differ in Shape from the rest, are all at first formed in the *Plume*, before it begins to sprout; and the rest afterwards; That is to say, that the former *Leaves*, are all formed (out of *sap* from the *Trunk*) with the *Seed* it self; and so compose one Principal Part thereof, *sc.* the *Plume*: the latter, not till after the *Seed* is sow'n, and so the *Plume* supply'd with *Sap* immediately from the *Root*. Which *sap*, it seems, is so far different from the former, as sometimes to produce a different Sort of *Leaves*.

5. §. SECONDLY, of the *Buds* of all *Trees*, and of *Perennial Stalks*, it appears, That they consist of a great number of *Leaves*, all perfectly formed to the Centre; where, notwithstanding, they are sometimes, not half so big as a *Cheese-Mite*. So that all the *Leaves* which stand upon a *Branch* or *Cien* of one whole Years Growth, were actually existent in the *Bud*. It is also very observable, That although these *Buds* begin to be expanded not till *Spring*, yet are they entirely Formed, as to all their *Integral Parts*, in the *Autumn* foregoing. So that the whole Stock of *Leaves* which grow upon a *Tree*, or any *Perennial Stalk*, this year; were made, or actually in being, the last year. A greater *Heat*, more subtilized *Aer*, and better concocted *Juyce*, being requisite for their *Generation*, than for their bare *Expansion* and *Growth*.

6. §. LASTLY, of all *Annual Plants*, in which there are several Successive *Generations* of *Buds*, one under another in one year; although I have not made the *Remarque*, yet am apt to believe, That as the *Leaves* in every *Bud* are all formed together, as in other *Plants*: so likewise, that the Successive *Generations* of the *Under-Buds*, begin at certain stated *Terms*: as in some *Plants*, at every *New Moon*; in others; at the *Full Moon*; and in some perhaps; with both, or every *Fourt'night*.

CHAP. VI.

Of the Manner of the Generation of the Leaf. Where also, that of the Two General Parts of a Plant, sc. the Lignous and Parenchymous, is further explain'd.



THE *Visible Causes* of the *Figures* of *Leaves*, have been formerly mentioned. It may here be further noted, That the greater *Fibers* of the *Leaf*, being never Braced in the *Stalk*; it is a good preparative for their better spreading in the *Leaf*. As also, that the same is much favour'd; by the extream smalness of the *Aer-Vessels* herein: whereby they are more easily divaricated, in the lesser *Fibers*, and so the *Leaf* dilated.

B.I. Ch. 4.

2. §. BUT these and the like are to be reckoned a secondary Order of *Causes*; which serve rather to carry on and improve, that which *Nature* hath once begun. And therefore, we must not only consider the visible *Mechanism* of the *Parts*; but also the *Principles* of which they are composed; wherewith, *Nature* seems to draw her first *Strokes*.

Idea; §. 53.

3. §.

3. §. Now of these, I have formerly, and as I conceive upon good ground, supposed, the chief Governing Principle, to be the Saline, whether Alkaline, Acid, or of any other Kind: being in some sort as the Mold of a Button, to which the other Principles, as its Attire, do all conform. Or the Salts are, as it were, the Bones; the other Principles, as the Flesh which covers them.

Lib. 2. P. 2.
§. 31, &c.

4. §. A further Argument hereof may be deduced from the Cuticular and other Concretions, commonly called Mothers, in Distill'd Waters, Vinegar, and other Liquors. For in these Concretions, there is always a tendency to Vegetation; and many of them are true Vegetables in their Kind; as shall hereafter be seen. Now the Liquors, in which these are generated, do always, wholly or in part, lose their Taste and Smell, and so become Vapid. The more sensible Principles therein having made their Transit from the Fluid, into the Concrete Parts. So, I have known, sometimes, Vinegar it self, to become by these Concretions, almost as Tasteless as Common Water. Whereby it seems evident, That of Vegetable Principles, there are some, more Masterly than others: and that of these, the Saline is the chief. The same is likewise argued, from the frequent Experiment of many good Husband-men; that most Bodies which abound with Salt, are the greatest Nourishers of Plants.

5. §. This saline Principle, as is above hinted, is to be understood, a Generick Name, under which divers Species are comprehended; and of some whereof, it is always compounded, as in other Bodies, so in Plants. As shall be made to appear, by divers Experiments, when we come, hereafter, to speak of Vegetable Salts. Whereby we are conducted, yet further to enquire, What are the Principles of this Principle?

6. §. NOW these seem to be Four; a Nitrous, an Acid, an Alkaline, and a Marine. The Admixture of the First, is argu'd from the Place, which Nature hath assigned for the Generation and Growth of most Plants, sc. neither in Caverns under Ground, as for Minerals; nor above it, as for Animals; but the Surface of the Earth, where this Sort of Salt is copiously bred. And doth therefore prove, not only a Mixture, but a good Proportion hereof with the other Principles of a Plant. Hence it is, that Dew or Water on Windows or Plain and Smooth Tables, by virtue of a Nitro-Aerial Salt, is often frozen into the resemblance of little Shrubs. And the like Figure I have often seen in a well filtered Solution of the Salt of any of our Purging Waters, as of Epsom, &c. being set to shoot. Produced, as I conceive, by the Nitre, which with the Rain or other Waters, is washed down from the Surface of the Earth, and so mixed with the Mineral Salts.

7. §. The other Three Salts are exhibited, by the several ways of Resolving the Principles of a Plant. Many Plants, even in their Natural Estate, do yield an Acid Juice. And the Juices of many more, by Fermentation, will become Acid. And most, by Distillation in a Sand-Furnace, yield an Acid Liquor.

8. §. By Calcination, all Sorts of Plants, yield more or less, both of a Fixed and a Volatile Alkaly: the former, in the Ashes; the latter, in the Soot. And, at least the generality, by Fermentation also, yield a Volatile one; or such a kind of Salt, which, whether we call an Urinous, or otherwise, hath the like Odour and Taste with that of Urine, Harts-Horn, Spot, and the like.

9. §.

9. §. The *Marine*, is obtained no other way, that I know of, but from a Solution of the *Alkaline*, upon its being exposed to the *Aer*. The process wherof, I shall particularly set down in a following *Discourse*. Of these *Salts*, mixed in a certain proportion, together, and also Impregnated with some of the other *Active Principles* of a *Plant*, and not without an *Admixture* of some *Parts* from the *Aer*; I suppose, that which I call the *Essential*, is produced: of which, I shall also give an account in the same *Discourse*.

10. §. ALL THE Four *Salts* above mentioned, seem in their *Order*, to have a share in the *Formation* of a *Leaf*, or other *Part* of a *Plant*: And first of all, the *Marine*. For all *Generations* are made in some *Fluid*: But in every *Fluid* there is a perpetual *Intestive Motion* of *Parts*. So that the first Intention of *Nature* is, That some of those *Parts* be disposed to *Rest*. Now of all the *Principles* of a *Plant*, there are none hereunto more disposed, than their *Salts*; whose *Particles*, being fig'rd with plain *Sides*, as often as they touch Side to Side, like two *Marbles* exquisitely polished, they will adhere together. And the *Particles* of *Marine Salt*, being *Cubick*; and so, with respect to their *Figure*, of greater *Bulk* than those of any other *Salt*; they will hereby, be most and first of all disposed to *Rest*; and so become, as it were, the *Foundation* of the following *Superstructure*.

11. THE Second Intention of *Nature* is, That the *Particles* be brought to *Rest*, in a certain *Position*, agreeable to the *Figure* of the *Parts* which are to be formed. And therefore in the next place, all those *Parts* of a *Plant* which are truly *Lignous*, by the *Marine Salt*, with the assistance of the *Alkaline*, but especially of the *Nitrous*, are made to shoot out in *Length*, or into an innumerable company of small *Cylindrick Fibres*: these *Salts* being, altogether, sturdy enough to resist those *Impulses* which might incline them to conform to any other *Figure*. Tab. 53.

12. §. THE next Intention is, That these *Fibres*, at the same time in which they are formed, may likewise receive such a *Posture*, as will best answer the indented *Shape* of the *Leaf*. Which *Posture*, although in the *Growth* of the *Leaf* it is much Govern'd by the *Aer-Vessels*; yet in the *Generation* hereof, seems to be first determined by the forementioned *Salts*, according to their several *Angles*, whereby they are differently applicable one to another.

13. §. Now all the *Sides* of the *Marine Salt*, and the *Sides* and the *Ends* of the *Nitrous*, properly so called, stand at *Right Angles*. And it is very probable, from the *Figure* of the *Crystals* in *Spirit of Blood*, and some other *Bodies*, that the *Particles* of the *Alkaline* are Square at one *End*, and Poynted at the other. And those of the *Acid*, at both; And that, withal, they are Shorter and more Slender. Tab. 53.

14. §. It should therefore seem, That where the *Alkaline Salt* is any way predominant, and that the *Particles* thereof are placed *End to End*; there the *Lignous Fibres* (as the larger ones in many *Leaves*) declining their parallel *Growth*, begin to shoot out obliquely, or at *Angles* one with another, and those *Acute*. Tab. 53.

15. §. If the same *Salt* be predominant, and some of its *Particles* placed, with the Pointed End of one, to the Side of another, or the Square End of one, to the Poynted End of another; there the said *Fibres* begin to shoot at *Angles* less *Acute*.

Tab. 53.

16. §. But if either the *Marine* or *Nitrous Salt* is predominant; or some *Particles* of the *Alkaline*, are placed with the Square End of one, to the Side of another; there the *Fibres* begin to make, not *Acute*, but *Right Angles*; as do the greater *Fibres*, in some *Leaves*; and the smaller, in all.

Tab. 53.

17. §. IN the same manner, the *Fibre* in the Circumference of the *Leaf* is also governed; the *Particles* of the said *Salt*, being reduceable, not only to any *Angle*, but also to any *Circle*, or other *Crooked Line*, as they are variously applyed. For if the *major part* be applied End to End, and only every Third or Fourth applied End to Side, they produce a great *Circle*. But if the Poynted End of each, be set to the Side of another, they make a less. And if the Application be the same, but to the contrary Side, they thence begin a new *Circle* with the same *Diameter*, but with another *Center*, answerable to the intended *shape* of the *Leaf*.

Tab. 53.

18. §. AFTER the same manner, the *Aer-Vessels* may be formed by the *Particles* of the *Acid Salt*. Which, without being supposed to be crooked (as those of the *Aer*, at least the compounded ones, probably be) only by applying the lesser Side of one, to the greater Side of another, will also be reduced to any either *Circular* or *Spiral Line*. And so, likewise, for the production of the winding *Fibres*, which compose the *Bladders* of the *Pith* and other *Parenchymous Parts* of a *Plant*.

Tab. 53.

19. §. Thus doth *Nature* every where γεωμετρεῖν. For what She appears in Her *Works*, She must needs be also in their *Causes*.

THE
ANATOMY
OF
FLOWERS,

PROSECUTED

With the bare EYE,

And with the

MICROSCOPE.

Read before the *Royal Society*, Novemb. 9. 1676.

The SECOND PART.

By *NEHEMIAH GREW* M.D. Fellow
of the *ROYAL SOCIETY*, and of the
COLLEGE of *PHYSICIANS*.

LONDON,

Printed by *W. Rawlins*, 1682.

Dd

THE
CONTENTS
OF THE
Second Part.

CHAP. I.

Of the EMPALEMENT.

CHAP. II.

Of the FOLIATURE.

CHAP. III.

Of the ATTIRE SEMINIFORM.

CHAP. IV.

Of the FLORID ATTIRE.

CHAP. V.

Of the USE of the ATTIRE.

CHAP. VI.

Of the TIME of the Generation of the Flower.

The Appendix.

*Being a METHOD proposed, for the ready finding, by the
Leaf and Flower, to what Sort any Plant belongeth.*

THE
ANATOMY
OF
FLOWERS.

PART II.

CHAP. I.

Of the EMPALEMENT.



NEXT proceed to the *Flower*. Where I intend not to repeat those things, which have been by Me already noted in the *First Book*. And the foregoing Discourse of *Leaves*, will excuse me from divers particulars, common to *These* and the *Flower*. I shall here therefore remarque some things not before mentioned, or but *in transitu*, and such as are more particular to the *Flower*.

Ch. 5.

2. §. And First, it may be noted; That where the *Leaves* of the *Flower* are few; those of the *Empalement* or *Green Border*, are either of the same Number, or just half as many, whether even, or odd. So in *Leucanthenum* and *Chickweed*, there are *Five Leaves*; in the former *Five Empalers*; in the latter, *Ten*. In *Great Celandine*, there are *Four Leaves*, and but *Two Empalers*; and so in *Poppy*. The *Arithmetick* of *Nature* being every where suitable to Her *Geometry*.

3. §. Of this *Part* of the *Flower* it is likewise observable, That it is rarely, if ever, entire or one piece, but parted into divers little *Leafy Pales*, especially in all *Flowers* with the *Florid Attire*, as of *Mari-gold*, *Daisy* and the like; being so numerous, as to make a *Double*, and often a *Treble*, *Quadruple* or *Quintuple Border*. Whereby they are aptly designed, not only to protect the *Leaves* of the *Flower* in the *Bud*; and after their Expansion, to keep them tite: but also, by receding, *Bredthways*, one from another, and so making a greater *Circle*, gradually to give way for the full Growth and safe spreading of the *Attire*. Which, in regard it consists of *Parts* so exquisitely tender, were

it pinched up too close, would be killed or spoyled before it came to the Birth. As *Teeming Women*, gradually slaken their *Laces*; or as *Taylors* use to split their *Stomachers* into several *Lappets*, to spread, as their *Belly* rises.

4. §. Nor is the *Posture* of the *Parts* in the *Empalement* less suitable: not being filed one just over another, but alternately. Whereby the *Pales* or *Pannicles* of every *Under-Order*, serve to stop up the gaps made by the *Recess* of the *Upper*. And so, notwithstanding they all make more roome, yet all conspire to keep the *Aer* out.

5. §. It is also worth the notice, That, for the same purpose, the *Edges* at least, of the several *Pales*, are neither *Fibrous*, nor *Pulpy*; but so many extream fine transparent *skins*, as in *Chamemile*. Whereby they close so exactly one over another, that it is impossible for any *Aer* to creep in, or any *steams* useful to the *Attire* or *Seed*, over hastily to perspire. As we use, when we have put a *Cork* into a *Bottle*, to tie a *Bladder* over it.

CHAP. II.

Of the FOLIATURE.

Tab. 54.

Ch. 5.



THE *Leaves* of the *Flower* are folded up in such Sort, as is most agreeable to their own *Shape*, and that of their inclosed *Attire*: whereof I have given Instances in the *Fist Book*. I shall here add some further *Remarques*.

Tab. 54.

Equilateral Pentangle.

2. §. The *Leaves* of the *Flower* of *Blattaria*, although of different *Size* and *Shape*; are so lapped one over another, as to make an

Tab. 54.

3. §. The *Spiral Fold*, which is proper to the *Flower*, and never seen in the *Green Leaves*; as it is it self immediately visible on the *Surface*, so by cutting off the top of the *Flower* before it is expanded, seems also to make a *Helix*; as in *Perwinckle*, the larger *Convolvulus*, &c.

4. §. In some *Flowers*, where the *Attire* is lofty or spreading, as in *Holioak*, together with the *Spiral Fold*, the *Leaves* are all at the top tacked down a little; thereby making a blunter *Cone*, and so a more ample *Pyramid* for the inclosed *Attire*.

Tab. 54.

5. §. In *Poppy*, although the *Leaves* are extraordinary broad, yet being but few, and inclosing a small *Attire*; they could not be well reduced to any regular *Fold*, without leaving such a *Vacuity*, as by being filled with *Aer*, might be prejudicial to the *Seed*. For which reason, they are cramb'd up within the *Empalement* by hundreds of little *Wrinckles* or *Puckers*; as if Three or Four fine *Cambrick Handcherchifs* were thrust into ones *Pocket*.

6. §.

6. §. In *Ladies-Bower*, the *Leaves* are neither laped one over another, as is most usual, nor set Edge to Edge, as sometimes, but Side to Side, answerable to their *Shape*, and the *Distribution* of their *Fibres*. Their broad *Tops* being also rowled up so as to make a *Cone*. In *Ladys-Looking-Glass*, they stand also Side to Side, but in a different manner: in the Former with the Sides standing inward, but here, bearing outward.

7. §. In the *Marvel of Peru*, the *Fold* is likewise very peculiar. For, besides the several *Plates*, about Six, whereby the *Flower* is gathered in the Middle; the *Top* of it is also gathered up by as many distinct *Plates*, underneath the former; and these rowled or wreathed up together so exactly, that the like could hardly be imitated by a very dextrous *Hand*.

8. §. OF the *Hairs* upon *Flowers* and their *Use* to the *Attire*, I have also spoken in the *First Book*. I shall here add, That they are likewise of *Use* to the *Leaves* themselves, that is, for their closer and faster *Conjunction*. For of some *Flowers* it is observable, That they are all over smooth, saving on their Edges, which are border'd with *Fringes* of *Hair*; as of *spanish Broome*, *Dulcamara*, and others: In which, the *Hairs* on the Edge of one *Leaf*, are so complicated, or at least indented, with those of another, that all the *Leaves* seem to be but one piece. *Nature* seeing it fit, by this means to tie them together, lest they should be expanded before it be due time.

9. §. Many *Flowers* instead of *Hairs*, are beset round about, with a great Number of small *Parts*, not ending in a *Poynt*, but having a *Head*. Sometimes oval, as in *Snap-Dragon*, like the *Horns* of a *Butterfly*, or a *Plummers Sodering-Iron*. But usually *Globular*, as in *Deadly Nightshade*, like so many little *Mushrooms* sprouting out of the *Flower*.

10. §. Out of these *Heads*, doth sometimes issue a *Gummy* or *Balsamick Juyce*. From whence proceeds that *Clamminess* of some *Flowers*, whereby, being handled they stick to our *Fingers*, as do those of *Blattaria*, and of *Marigold*; and those of *Colus Jovis*, where the said *Heads* are so soft and succulent, that they resemble so many little *Drops* of *Balsame*. The *Clamminess* which is felt upon fresh *Carduus*, may perhaps proceed from the like *Cause*.

11. §. THE *Number* of the *Leaves* of the *Flower* hath been noted by the Learned Sir *Thomas Brown*, to be usually *Five*. And this *Treat.* of *Nature* so far affecteth, that many times where the *Leaves* of the same the *Flower* are of a different *Size*, yet they keep to this *Number*, as in *Blattaria*.

12. §. I also add, That even those *Flowers*, which are not properly parted into *Leaves*, have yet their *Tops* usually divided into *Five* great *Scallops*; as those of *Toad-Flax*, *Snap-Dragon*, *Coded-Arsmart*, *Clary*, *Broom*, and others. And when the *Flower* hath more than *Five*, even many times *Five Leaves*; yet the *Top* of each *Leaf* is indented into *Five Parts*; as in *Scorzonera*, *Cichory*, and all the *Intybous Kind*, with many others.

13. §. From whence and other like *Instances*, it may seem, That there is some certain *Species* of *Salt* in *Nature*, and that in most *Plants*, of whose *Agency* there are still some *Footsteps* or other in the *Flower*.

14. §.

14. §. The Number of the *Leaves*, as hath been said, is commonly *Five*. Yet some *Flowers* have fewer, and some more, and that with *Constancy*, in divers *Numbers*, from *One* to *One* and *Twenty*; perhaps in all, so far. The *Flower* of *Acanthus Syriacus*, is in a manner one single *Leaf*, that of *Monks-Rubarb*, *Three-Leav'd*; of *Poppy*, *Crosswort*, *Radish*, and many others, *Four-Leav'd*; the greater Number of *Flowers*, *Five-Leav'd*; of *White Hellebore*, *Tulip*, *Onion*, and most *Plants* with *Bulbous Roots*, *Six-Leav'd*; of *Wild-Crowfoot*, *Seven-Leav'd*; of *French Marigold*, commonly *Eight-Leav'd*; of *Flower-de-luce*, *Nine-Leav'd*; of *Chickweed*, *Ladies-Mantle*, *Ten-Leav'd*; of *St. James's Wort*, *Thirteen-Leav'd*; and I think of *Febrifuga*, *Cotula*, *Ageratum*, *Corn-Marigold*, with others; and of *Chamemile*, *Euphthalmum*, and some few more, the *Leaves* are commonly *One* and *Twenty*. In that of *St. James's Wort*, the Number is so constant that there is scarce *One Flower* in *Forty*, wherein the *Leaves* are more or fewer than *Thirteen*. Divers of which *Numbers*, seem also to have some relation to the Number 5. For 9, is *Twice*; 13, *Thrice*; and 25, *Five times* 5 running into it self.

Tab. 55.

15. §. THE *Constituent Parts* of the *Flower* are the same as those of the *Leaf*, sc. the *Parenchyma* or *Pulp*, and the *Vessels*. But in the *Basis* or bottom of the *Flower*, the *Parenchyma* is commonly much more *spongy* and *dry*, than in the *Leaves*; containing, after the *Flower* is open'd, little or no *Sap*, but only a *dry* and *warm Aer*. Which standing continually under the *Seed*, hastens the *Maturation* or due *Exiccation* thereof: as we use to dry *Maulted Barly* over a warm *Killn*.

16. §. The *Vessels* of the *Flower*, are both for *Sap* and for *Aer*, as well as in other *Parts*. And both of them sometimes, even in the *Skin* of the *Flower*; as may be argued from its being stained with divers *Colours*; produced as hath formerly been shewed, by the mixed *Tinctures* of the said *Vessels*. These *Colours*, in many *Flowers*, as *Tulips*, as they are in the *Skin* it self, so therein only; the *Pulp* of the *Leaf* being white.

B. 2. P. 2.
§. 65, 66,
67.

17. §. The *Lignous* or *Sap-Vessels* are fewer, and the *Aer-Vessels* smaller in the *Flower*, than in the *Leaf*. And therefore it is very difficult to observe the latter by *Glasses*; especially the *Proportion* which they hold to the other *Parts*. But if you break the *Leaves* of some *Flowers*, with very great gentleness; they may hereby be *Unroaved* or *drawn out*, as in the *Green Leaves*, to some visible length; and their different *Number* in divers *Flowers* may be discerned.

B. 1. Ch. 5.

18. §. THE *Use* of the *Flower* or of the *Foliature* whereof we are speaking, is various; as hath formerly been shewed. I now only add, That one *Use* hereof seemeth to be, for the *Separation* of the more *Volatile* and stronger *sulphur* of the *Plant*. That so the *Seed*, which lyeth within or next it, may be so much the milder, and the *Principles* thereof more fixed and centred. And this, both for its better *Duration* till the time of *Sowing*; and also, that its *Fermentation*, when it is sow'n, may not be too hot and precipitate; but suitable to so slow and equal a motion, as is the *Vegetation* of a *Seed*.

19. §. And that this *Sulphur* is separated and discharged by the *Flower*, seems evident, not only from the *Strength* of its *Odour*, above that of the other *Parts*; but likewise, in that many times where there is no *Flower*, or that very small, the *Seed*, that is its *Cover*, as in the *Umbelliferous*

belliferous Kind, is the more odorous. And therefore also, the *Vine* hath no *Flower*, partly, that the most *Volatile Spirit* and *Sulphur* might all run into the *Fruit*.

20. §. THE *Figure* of the *Flower*, although it is often much more complex, than that of the *Leaf*: yet there is no doubt, but that the *Measure* hereof may be defined in some way, answerable to that exemplified in the foregoing *Part*. The difference is only this, That whereas the *Green Leaves*, and the *Plain Leaves* also of the *Flower*, are all measured by the *parts* of several *Circles*: those *Flowers* which are *Bellyed*, and those *Leaves* of the *Flower* which are not *Plain*, but *Convex*, are all measured by the *parts* of several *Spheres*. And as the *Diametres* of those *Circles*, bear a certain proportion to the middle *Stemm* of the *Leaf*; so the *Axes* of these *Spheres*, to an imaginary one in the *Centre* of the *Flower*.

21. §. NOW the reason why the *Figure* of the *Flower* is more *multiplex*, than that of the *Leaf*; may be, partly, because it is under the *Command* and *Government* of those *salts*, which are here more *refined* and *depurate*, than in the *Leaf*; and so more free to lay the *Foundation* of any kind of *Figure*, for which, of their own *Nature*, they are adapted. Partly, for that as the *Nitrous* and *Alkaline Salts* are chiefly regnant in the *Leaf*; so in the *Flower*, in which the *Parenchymous Part* hath a greater (a) proportion than in the *Leaf*; it is most reasonable, to assign the *Predominion* to the *Acid* (b): the *Particles* whereof, both as they are less, and also poynted at both ends, (c) seem to be more easily applicable one to another for the making of any Sort of *Line* or *Figure*.

(a) §. 17.

(b) *Idea*,

§. 52.

(c) *P. 1.**Ch. 6. §. 13.*

CHAP. III.

Of the Attire, and first of that sort which may be called
Seminiform.



WITHIN the *Foliature* stands the *Attire*; which is of *Two* general *Kinds*, every where *Various* and *Elegant*; according to the *Description* I have given of them in the *First Book*. I shall here add some *Ch. 5.* further *Remarques*.

2. §. And first, of that Sort of *Attire*, which may be called *Seminiform*; being usually, as it were, a little *Sheaf* of *seed-like Particles*; standing on so many *Pedicills*, as the *Ear* doth upon the *End* of the *Straw*.

3. §. Of their *Colour* it is observable, That for the most part, they are *White* or *Yellow*; sometimes *Blew*; but never *Red*, let the *Flower* or *Foliature* be of what *Colour* it will. Neither doth their *Colour* always follow that of the *Foliature*, although that be not *Red*. Whereby it appears, how very *Curious* and *Critical Nature* is, in the *Separation* of the *Juyces* in *Plants*: that such small *Parts* as these of the *Attire*, and so near the *Leaves* of the *Flower*, should yet receive a different *Tincture*.

4. §.

4. §. These *Parts* differ also in their *Position*; standing sometimes double upon each *Pedicil*, as in *Toad-flax*, *Snapdragon*, and some others; but usually single, as in *Blattaria*, *Clematis Austriaca*, &c. Sometimes fastned to their *Pedicils* at their middle, stooping down after the manner of *Poppy* and other hanging *Flowers*; as in *Spanish-Broom*, *Hysop*, *Scabeous*, *Behen*, &c. Sometimes they stand erected, as in *Clematis Austriaca*, *Ladyes-Looking-Glass*, *Rape-Crowfoot*, &c. Those of *Coded Arsmart* have no *Pedicils*, but stand upon a large *Base*.

Tab. 55.

5. §. Of the *Pedicils* themselves, it is to be noted, That they are rarely fastned to the *Top* of the *Repository* or *Case* of the *Seed*, but round about the *Bottom*. Partly, That hereby they may the better intercept and separate the *Incongruous Parts* of the *Sap* from the *Seed*. Yet in the *Coded Arsmart* they stand at the *Top*. Which is not the only thing peculiar in that *Plant*; it being the property thereof, to ejaculate its *Seed*, upon the least touch. Which property seemeth to depend, partly, upon the *Position* of the said *Pedicils*, as shall be shewed in speaking of the *Seed*.

6. §. These *seed-like Parts* are also of different *Number*. In *Great Celandine*, *Rose*, *Rape-Crowfoot*, numerous; in *Great Plantaine*, and some other *Herbs*, much more conspicuous than the *Foliature* it self. In *Germander-Chickweed*, they are always *Two*, and no more. Sometimes they follow the number of the *Leaves*, especially in the number 5; as in *Blattaria*, *Black Henbean*, &c. In *Stichwort* and *Lychnis Sylvestris*, they are 10, just double to the number of the *Leaves*.

7. §. They differ also in their *Bigness*, being in some smaller *Flowers*, large; as in *Borage*, *Ladyes-Looking-Glass*, and others: and in some larger *Flowers*, less; as in the *Rose*.

8. §. But especially in their *Shape*, which is always very *Elegant*, and with much *Variety*. In *Borage*, like the point of a *Spear*. In *Blattaria*, like a *Horse-shooe*. In *Clematis Austriaca*, like the *Spatula*, wherewith *Apothecaries* make their *Mixtures*. In *Mallow*, like a *Head-Roll*. In *Hysop*, they have one *Cleft* before; in *Blattaria*, one round about; in *Water Bettony*, one at the *Top*; in *Scabious*, they have a double *Cleft*, one on each side; and so in *St. Johns Wort*, *Hyoscyamus*, and others; before they open, in the *Shape* of a double *Purse*.

Tab. 56.

9. §. These *Parts*, are all hollow; each being the *Theca* or *Case* of a great many extream small *Particles*, either *Globular*, or otherwise *Convex*; but always regularly figur'd. They are all crowded together, and fastned in close *Ranks*, without any *Pedicils*, to the *Insides* of the *Theca*, like other lesser *Seeds* within a greater; or after the same manner as in *Hyoscyamus* and some other *Plants*, the true *Seeds* themselves grow all round about close to the *Bed* of the *Case*; as in *Clary*, and the *Figures* now referred to, may be seen. And when they are ripe, the *Case* also opens and admits them to the *Aer*, as the *seed-Case* doth the *seed*. The whole *Attire*, together with the *Foliature* and *Seed-Case*,

Tab. 55, 56.

Tab. 57.

See in one Example, amongst the *Figures*.
10. §. The *Colour* of these small *Particles* contained in the *Theca*, is also different. But as That is usually *White* or *Yellow*, so are These: sometimes *Blewish*; but never *Red*. And sometimes not of the same *Colour* with that of the *Theca*. Which further shews how scrupulous *Nature* is, in differencing the *Tinctures* of the several *Parts*.

11. §. They are also of different *Bigness* and *Figure*. Those in *Snap-dragon*, are of the smallest *size* I have seen; being no bigger through a good *Microscope*, than the least *Cheese-Mite* to the naked Eye. In *Plantain*, also through a *Glass*, like a *Scurvy-grass-seed*. In *Bears-foot*, like a *Mustard-seed*. In *Carnation*, like a *Turnep-seed*. In *Bindweed*, like a *Peper-Corn*. In all these of a *Globular Figure*. Tab. 58.

12. In *Devils-bit*, they are also *Round*, but depressed, like the *Seed* of *Goof-grass*, or a *Holland Cheese*. In the *Bean* and all sorts of *Puls*, and *Trefoyls*, as also in *Blew-bottle*, &c. they are *Cylindrick*. In *Orange Lilly*, *Oval*, one 5th of an *Inch* long, like an *Ants-Egg*. In *Deadly-Night-shade*, also *Oval*, but smaller at both *Ends*. And those of *Pancy*, *Cubick*. In all these and the former, they are *smooth*. Tab. 58.

13. §. But in *Mallow*, *Holyoak*, and all of that kind, they are beset round about with little *Thornes*; whereby each looks like the *seed-Ball* of *Roman Nettle*, or like the *Fruit* of *Thorn-Apple*, or the *Fish* called *Piscis orbis minor*, or the *Murices*, used antiently in *Wars*. They are also very great, shewing, through a *Glass*, of the bigness of a large *White Pease*; being 200 or 300 times bigger than those in *Snapdragon*; of which there are about a *Thousand* in each *Theca*, that is, in the space of about 1000th *Cubical Part* of an *Inch*. Tab. 58.


15. §. In some *Plants*, as in *Deadly Night-shade*, where these *Partiales* are *White*, they seem, by a very good *Glass* and advantagious *Position*, to be composed of *Parenchymous* and *Lignous Fibres*, stitched up together, as in the other *Parts*.

15. §. In *Colocynthis*, (and with some *Analogy* in *Wild Cucumer*, and I suppose all of that kind) the *Attire* is very peculiar, not consisting of several little *Theca*, upon so many *Pedicils*, as is described; but is all one entire *Part*, like a thick *Columna* in the midst of the *Flower*; having several little *Ridges*, and *Furrows* winding from the *Top* to the *Bottom* round about. In the middle of each *Ridge* runs a *Line*, where the *Skin*, after sometime, openeth into two *Lips*, presenting the *Globular Particles* contained in the hollow of every *Ridge*.

16. §. Where the *Attire* consists of several *Seed-like Parts*, as is described; there, another *Part* distinct, like a little *Columna* or *Pinnacle*, stands on the *Top* of the *Uterus* or true *Seed-Case*. Which is also regularly and variously *Figured*. In *Bindweed*, it hath a round *Head*, like that of a great *Pin*. In the *Common Bell*, *St. Johns wort*, it is divided into *Three Parts*. In *Gerarium*, into *Five*; In *Asarum*, into *Six*. Sometimes, the *Head* is *Smooth*, and sometimes beset with little *Thorns*, as in *Hyoscyamus*. Of the *Use* of these *Parts*, anon. Tab. 56, 57.

C H A P. IV.

Of the FLORID ATTIRE.



IN THIS Attire there is also much Elegant Variety, according to the Description we have given of it in the First Book. It always consists of several Suits; Ten, Twenty, Forty, a Hundred, or more, according to the Bigness of the Flower. And every Suit most commonly, of three distinct Parts, all of a Regular, but Different Figure. The utmost Part, is always like a little

Flower with Five Leaves and a Tubular Base, like that of Cowslip. So that every Flower with the Florid Attire, Embosomes, or is, a Posy of perfect Flowers.

2. §. In some Flowers, every one of these Florets, is encompassed with an Hedg of Hairs; and every Hair branched on both sides almost like a Sprig of Fir; as in Aster Atticus, Golden-Rod, and others.

3. §. The Base of the Floret is usually Cylindrick, but sometimes Square, as in French Marigold. And the Leaves hereof which, for the most part, are Smooth on the Inside, in the same Flower are all over Hairy. And the Edges of these little Flowers, are frequently Ridged, or as it were, He m'd, like the Edge of a Band.

4. §. The midlemost of the Three Parts, which I call the Sheath, is usually fastened towards the Top, or else at the Bottom of the Floret. This is rather indented, than parted into Leaves. The Surface seldom Plain or Even, but wrought with Five Ridges, and as many Gutters running almost Parallel from the Top to the Bottom.

5. §. The Inmost Part, which I call the Blade, runs through the hollow of the Two Former, and so is fastned, with the Floret, to the convex of the Seed-Case. The Head and Sides of this Part, is always beset round about with Globulets, commonly through a Glass, as big as a Turnep-seed, or a great Pins-Head. In some Plants growing close to the Blade, as in the common Marigold; in the French, and others, upon Pedicils or little slender Stalks. These, as the Blade springeth up from within the Sheath, are still rubed off, and so stand like a Powder on them both. And sometimes, as in Cichory, they seem to grow on the Inside the Sheath, if it be split with a small Pin: as also in Knapweed, in which they are numerous. Yet in the Seed-like Attire, always more numerous, than in the Florid.

6. §. The Head of the Blade is always divided into Two, and sometimes into Three Parts, as in Cichory; which, by degrees, curl outward, after the manner of Scorpion-Grass.

7. §. The Description now given, agrees principally to the Corymbiferous Kind, as Tansy, Chamemile, and the like. But in Scorzonera, as also Cichory, Hawk-Weed, Mouseear and all the Intybous Kind, with many more

more, the *Attire* is not separate from the *Foliature*, so as to stand within that in one entire *Posy*; but every *Leaf* of the *Flower* hath its own *Attire* apart. For the sake of which, the *Basis* of every *Leaf* is form'd into a little *Tube* or *Pipe*, whereby it embosoms its own *Attire* within it self. Consisting commonly of *Two Parts*, a *Sheath* and a *Blade*: the *Leaf* it self answering to the *Floret* in other *Flowers*. Tab. 62.


8. §. In some *Plants*, besides the *Attire* or *Posy* in the middle of the *Flower*; the *Leaves* also have each their own to themselves, as in *Marigold*: yet this, as I take it, consisting only of one single *Part*, which answers to the *Blade*; the *Leaf* it self being as the *Sheath*. Tab. 61.

9. §. In many *Plants*, this *Florid Attire* is very large; so that not only the *Suits*, but also the several *Parts* whereof every *Suit* consists, being throughly ripe and well blown open, are all visible to the bare *Eye*, as in *Knapweed*, and all the *Thistle Kind*. This *Attire* is all the *Flower*, that this sort of *Plants* have; being, though *Empal'd*, yet without any *Foliature*. Tab. 61.

10. §. And sometimes, there is little or no *Flower* besides this *Attire*, although extream small, as in *Golden Rod*, *Wormwood* and others. Where it may be noted, That the *Medicine* called *Wormseed* or *Semen Santonici*, is no Sort of *Seed*, but the *Buds* of small *Flowers*, or of the *Florid Attire* of that *Plant*.

CHAP. V.

Of the Use of the Attire.

F the *Secondary Use* hereof, I have spoken in the *First Book*; and particularly, of the *Globulets* or small *Particles* within the *Theca* of the *Seed-like Attire*, and upon the *Blades* of the *Florid*, I have conjectur'd, That they are that *Body* which *Bees* gather and carry upon their *Thighs*, and is commonly called their *Bread*. For the *Wax* they carry in little *Flakes* in their *Chaps*: but the *Bread* is a Kind of *Powder*; yet somewhat moist, as are the said little *Particles* of the *Attire*. Ch. 5.

2. §. But the *Primary* and chief *Use* of the *Attire* is such, as hath respect to the *Plant* it self; and so appears to be very great and necessary. Because, even those *Plants* which have no *Flower* or *Foliature*, are yet some way or other *Attir'd*; either with the *Seminiform*, or the *Florid Attire*. So that it seems to perform its service to the *Seed*, as the *Foliature*, to the *Fruit*.

3. §. In discourse hereof with our Learned *Savilian* Professor Sir *Thomas Millington*, he told me, he conceived, That the *Attire* doth serve, as the *Male*, for the *Generation* of the *Seed*.

4. §. I immediately reply'd, That I was of the same *Opinion*; and gave him some reasons for it, and answered some *Objections*, which might oppose them. But withall, in regard every *Plant* is ἀρρενώδης or *Male* and *Female*, that I was also of *Opinion*, That it serveth for

the *separation* of some *Parts*, as well as the *Affusion* of others. The sum therefore of my Thoughts concerning this *Matter*, is as follows.

5. §. And First, it seems, That the *Attire* serves to discharge some redundant *Part* of the *Sap*, as a *Work* preparatory to the *Generation* of the *Seed*. In particular, that as the *Foliature* serveth to carry off the *Volatile Saline Sulphur*: So the *Attire*, to minorate and adjust the *Aereal*; to the end, the *Seed* may become the more *Oily*, and its *Principles*, the better fixed. And therefore the *Foliature* generally hath a much stronger *Odour*, than the *Attire*: because the *Saline Sulphur* is stronger, than an *Aerial*, which is too subtile to affect the Sense. Hence also it is, that the *Colour* of the *Parts* of the *Attire*, is usually *White*, or *Yellow*, never *Red*: the former, depending upon a greater participation of *Aer*; the latter, of *Sulphur*. I add further, That the most *Volatile* and *Aerial Sulphur*; being by means of these *Parts* much discharged; it may hereby come to pass, not only that the *Seed* is more *Oylie*, and its *Principles* more fixed; but also, that the *Body* or *Parenchyma* thereof, is so compact and close: For although it consists of *Bladders*, yet such, as are Twenty times smaller than in any other *Part* of a *Plant* of the like bigness. Whereas, were the *Aer* copiously mixed with the *Sap* here, as in the *Pith*, *Fruit*, and other *Parenchymous Parts*; it would give so quick a *Ferment* to the *Sap*, as to dilate and amplify the *Bladders* of the *Seed*, beyond its present compact and durable *Texture*; and so expose it, either to a precipitant *Growth*, or sudden *Rot*. Wherefore, as the *Seed-Case* is the *Womb*; so the *Attire* (which always stands upon or round about it) and those *Parts* of the *Sap* herinto discharged; are, as it were, the *Menses* or *Flowers*, by which the *Sap* in the *Womb*, is duly qualified, for the approaching *Generation* of the *Seed*.

6. §. And as the young and early *Attire* before it opens, answers to the *Menses* in the *Femal*: so is it probable, that afterward when it opens or cracks, it performs the *Office* of the *Male*. This is hinted from the *Shape* of the *Parts*. For in the *Florid Attire*, the *Blade* doth not unaptly resemble a small *Penis*, with the *Sheath* upon it, as its *Præputium*. And in the *Seed-like Attire*, the several *Thecæ*, are like so many little *Testicles*. And the *Globulets* and other small *Particles* upon the *Blade* or *Penis*, and in the *Thecæ*, are as the *Vegetable Sperme*. Which, so soon as the *Penis* is exerted, or the *Testicles* come to break, falls down upon the *Seed-Case* or *Womb*, and so Touches it with a *Prolifick* Virtue.

7. §. *Consentaneous* hereto it is also observable, That those *Herbs* generally have the *Seed-like Attire*, which either produce a greater Quantity of *Seed*, or a *Perennial Root*: and that there is no *Tree*, with the *Florid Attire*. As if the other, because it contains a far greater Proportion of the abovesaid *Particles*, that is, of *Sperm*; 'tis able to beget a more *Numerous*, *Vivaceous*, or *Gigantick Birth*.

8. §. That the same *Plant* is both *Male* and *Female*, may the rather be believed, in that *Snails*, and some other *Animals*, are such. And the *Parts* which imitate the *Menses*, and the *Sperm*, are not precisely the same: the former, being the *External Parts* of the *Attire*, and the *Sap*, which feeds them; the latter, the small *Particles* or moyst *Powder* which the *External* inclose.

9. §. And that these *Particles*, only by falling on the *Uterus*, should communicate to it or to the *Sap* therein, a *Prolifick Virtue*; it may seem the more credible, from the manner wherein *Coition* is made by some *Animals*; as by many *Birds*, where there is no *Intromission*, but only an *Adosculation* of *Parts*: And so in many *Fishes*. Neither in others, doth the *Penis* ever enter any further than the *Neck* of the *Womb*. Nor doth perhaps the *Semen* it self: or if it doth, it can by no means be thought, bodily or as to its gross *Substance*, to enter the *Membranes*, in which every *Conception*, or the *Liquor* intended for it, before any *Coition*, is involved; but only some subtle and *vivifick Effluvia*, to which the visible *Body* of the *Semen*, is but a *Vehicle*. And the like *Effluvia* may be very easily transfused from the above said *Particles* into the *Seed-Case* or *Womb* of a *Plant*.

10. §. If any one shall require the Similitude to hold in every Thing; he would not have a *Plant* to resemble, but to be, an *Animal*.

CHAP. VI.

Of the Time of the Generation of the Flower.



THE *Time* in which the *Flower* is Generated or Formed is a Providence in Nature, whereof, I do a little wonder, that no one, amongst so many observers of *Plants*, hath ever yet taken any notice. It is therefore to be remarked, That all *Flowers* are formed or perfectly finished, in all their *Parts*, long before they appear in sight; usually Three or Four Months, and sometimes half a year, or more. And that in all *Perennial Plants*, those *Flowers* which appear and are called the *Flowers* of any one year; are not formed in that year; but were actually in *Being*, and entirely formed in all *Parts*, the year before; as in many *Herbs*, and in all *Shrubs* and *Trees*.

2. §. This will best be seen by some Instances. So the *Flower* of *Mezereon*, which opens in *January*, is entirely formed about the middle of *August* in the year foregoing. At which time, the *Green Leaves* of the *Bud* being cautiously removed, the *Leaves* of the *Flower*, and the *Theca Seminiformes* or seed-like *Attire*, encompassing the *Seed-Case*, through an indifferent *Glass*, are all distinctly visible. Tab. 63.

3. §. The like may be seen in *Sirynga*, and other *Shrubs*, and in *Trees*. In as many of which, as are *Frugiferous*, the *Fruit* also, which answers to the *Seed-Case* in other *Plants*, is about the same time entirely formed.

4. §. And so in *Herbs*; as the *Flower* of *Asarum*, which appeareth in *April* or *May*, is entirely formed in *August* or *July* of the foregoing year. For there are here, as well as in *Trees*, Two Sorts of *Buds*; some Tab. 64. which are composed only of *Green Leaves*; and some which also contain a *Flower* and the *Seed-Case*. So in *Bears-foot*, by some called the *January Rose*, the *Flower-Buds*, which open in *January* are all formed in or before the Month of *August* in the year preceding.

5. §.

Tab. 63.

5. §. The same may also be seen about the end of *August* or the beginning of *September* in a *Tulip-Root*. In which, the Two Inmost *Shells* dryer than the rest, stand hollow, with the little young *Flower* (which appears in *March* or *April* following) inclosed now in their *Centre*. Being thus kept *warm* and *dry*, lest it should either perish, or be precipitated upon the *Winter*, by sprouting too soon.

6. §. From hence it is plain, That although the *Flower* appears before the *Seed*; yet if the comparison be made betwixt the *Flower* and *Seed* of the same year; the *Seed* is first formed, and afterward the *Flower*. That is, the *Seed*, for which Nature chooses the First-born *Sap*, is formed in the fore part of the year: which work being finished, out of the less *fecund* part of the *Sap*, the *Flowers* intended for the *Sire* and *Matrix* of the next years *Seed*; is afterwards produced.

7. §. THE true *Time* of the *Generation* of the *Flower* being know'n, it may also be an Inducement to make *Tryal*, for the bringing of many *Flowers* to grow fairly in *Winter*, which are used to grow, that is, to appear, only in the *Spring* and *Summer*: *sc.* by keeping the *Plants* warm, and thereby enticing the young lurking *Flowers* to come abroad.

The Appendix.

Being a Method proposed, for the ready finding, by the Leaf and Flower, to what Sort any Plant belongeth.



ALTHOUGH many have bestowed extraordinary Care and Industry upon the searching out, and Description of *Plants*; and for the reducing of them to their several *Tribes*: yet I will take leave, here to propose a short *Method* whereby Learners, seeing a *Plant* they know not, may be informed to what Sort it belongs, and so be directed where to find it described and discoursed of. For, except they have a Master to conduct them, which few have; they must needs, by seeking at random, lose a great deal of time, which by a regular Enquiry might be saved. Besides, that what is learned by their own Observation, will abide much longer on their mind, than what they are only Poynted to, by another.

2. §. Now the most *Philosophick* way of distinguishing or sorting of *Plants*, were by the *Characteristick Properties* in all *Parts*, both *Compounded*, *Constituents*, and *Contents*. But of the *Compounded*, the *seeds*, and some other *Parts*, are oftentimes very *minute*: and the *Roots* always lie hid. As also the *Constituent Parts*, every where, without cutting and the use of *Glasses*. Nor can the *Contents* be accurately observed otherwise. So that for the *Use* here intended, those *Properties* are the fittest to be insisted upon, which are the most *Conspicuous*, and in those *Parts*, where the Learner may the most readily and without any difficulty take notice of them; as in the *Flower* and *Leaf*. The *Flower* hath Varieties enough of it self. But in regard it is often wanting, when the *Green Leaf* is not; it is therefore convenient, that he be assisted by

by both, and that the Varieties of both be distinctly reduced unto *Tables*. Which may be done, after the following, or some other like manner.

3. §. And First for the *Leaves*. The most obvious *Varieties* of which, are in their *Position*, *Size* or *Shape*.

4. §. *Leaves* are fastned with, or without a *Stalk*. Without, only close to the *Branch*, as in *Southistle*; or surrounding it, as in *Thorow-Wax*.

5. §. Both these ways, they stand either singly, that is, but one at the same height; or more together.

6. §. More together, in Even or Odd Numbers. In Even Numbers, commonly Two and Two, as in *Sage*, *Polium*; Sometimes Four, as in *Cross-wort*, *Madder*, *Herb True-Love*, *Pomum Majæ*; or more, as, I think, in *Woodrose*, &c. In Odd Numbers, Three, as in all *Trefblys*, *Strawberries*; Five, in *Pentaphil*, *Castanea Equina*; Seven, in *Tormentil*.

7. §. The *Sizes* of *Leaves* are innumerable. It is therefore necessary to reduce them to a Standard. And so, they may be reckoned, Three; *Small*, *Mean* and *Great*: with respect to the *Length* of the *Leaf*, the *Breadth*, or both. From one Inch and under, all *Leaves* may be accounted *Small*; from one Inch and over, to five Inches, *Mean*: from five and over, *Great*.

8. §. The *Shapes* of *Leaves* are also numberless. But the most obvious distinctions which they admit of, are such as these;

9. §. *Leaves* are *Membraneous*, as the greater part; *Squameous*, as *Abies*, or *Filamentous*. Which are *solid*, as in *Fenil*, *Meum*, *Buphtalmum*, *Chamemile*, *Groundpine*; or *hollow*, as in *Onion*.

10. §. *Membraneous*, have all their main *Fibres* produced either from the *Stalk*, as in *Holyoak*; or from the middle *stem* of the *Leaf*, as in most. From the middle *Stem*, reciprocally, as in *Scabious*, or oppositely, that is, one over against another, as in *Rose*: and both ways, at *Acute Angles*, as in most; or *Right*, as in *Dandelion*.

11. §. Again, they are different with respect to the *Top*, the *Bottom*, and the *Sides*. The *Top* is *Thorny*, as in *Furz*; or *Unarmed*. *Unarmed*, either *Produced*, that is, *Poynted*, or at least, *Roundish*, as in *Lanium*, *Ironwort*; or else *Reduced*, as in *Woodsorrel*. And so the *Bottom*, is either *Reduced* towards the *Top*, as in *Ground-Ivy*; or *Produced* upon the *Stalk*, as in *Poplar*, *Bay*, &c.

12. §. The *Sides* or *Edges* of the *Leaf*, are either of one and the same *Measure*, as commonly; or of divers, as in *Doronicum*. Both ways they are Even, as in *Syringa*, *Mous-ear*; or Uneven. The Uneven, are *Prickly*, as *Holly*, *Eryngium*, *Thistle*; or Unarmed. Unarmed, are *Infected*, or *Resected*. *Infected* deeply, that is, *Lobed*, as *Golden Liverwort*, *Clematis Peregrina*; or with shallow *Infections*, as in most. And so, *Indented*, or *Scalloped*: the former, when the *Angle* is made with *Straight Lines*, as in *Dandelion*; the latter, with *Crooked*, as in *Thalictrum*. *Resected*, that is, both *Lobed*, and *Infected*, or when upon the greater *Infections*, there are other lesser ones, as in *Wild-Clary*, *Lovage*, *Masterwort*.

13. §. THE most *Conspicuous Varieties* of *Flowers*, are in their *Position*, *Size*, *Shape*, and *Colour*.

14. §. Most are fastned with *stalks*; but many without. Sometimes, they are placed round about the *Branch*, that is, *Coronated*, as
in

in *Pulegium*; and sometimes, all on one side; either in *Ranks* only, as in *Bawm*; or in *Rank* and *File*, as in *Foxglove*. In *Saxifraga Aurea*, they grow on the *Leaf*.

15. §. Again, they either stand *Singly*, as in *Corn Marigold*; or *Clustur'd*. And so, either all upon one *Branch*, or on several little *Ramified Sprigs*. On one *Branch*, prolonged like a *Tail*, as in *Blattaria*; or *Contracted*. And so, either without *Stalks*, that is, *Capitated*, as in *Scabious*; or with *Stalks*, that is, *Umbellated*, as *Fenil*, &c. On several *Sprigs*, as in *Tanacetum*, *Yarrow*.

16. §. The *Sizes* of *Flowers*, as of the *Leaves*, may be reduced to *Three*. From $\frac{1}{2}$ an *Inch* and under, in *Diameter* or *Length*, may be accounted *small*. From $\frac{2}{3}$ an *Inch* and over to an *Inch* and $\frac{1}{2}$, may go for *Mean*. And from an *Inch* and $\frac{1}{2}$ and over, *Great*.

17. In respect of the *Shape*, *Flowers* are *Open* or *Belly'd*. *Open* have both *Leaves* and *Attire*, as most; or else are all *Attire*, as of *Burdock*, *Beta Cretica*.

18. §. The *Open*, consist of a *Certain Number* of *Leaves*, *One*, *Two*, *Three*, *Four*, *Five*, *Six*, *Seven*, *Eight*, *Nine*, *Ten*, *Thirteen*, or *One* and *Twenty*. Uncertain, commonly called *Double*. Those of a *Certain Number*, either *Uniform*, that is, all of a certain *size* and *Shape*, as usually; or *Biform*, or *Triform*, as in *Iris*, *Blattaria*. And these again, *Even Edged* or *Notched*; with *Three Poynts*, as in *Marigold*; or *Five*, in *Cichory*.

19. §. The *Belly'd*, are either so in whole; or in *Part*, that is, with the *Top* divided into *Leaves*, and the *Bottom*, *Hollow*: The former, are also *Even Edged*, as in *Convolvulus*; or *Notched*, as in *Trachelium*. The latter have their *Leaves* distinguished as before. Their *Bottom* or *Base*, either fastned to the *Seed-Case*, as in *Snap-dragon*; or standing below it. And so, either *Straight*, as I think in *Toad-flax*; or *Crooked*, as in *Violet*, *Lark-beel*.

20. §. In all these, the *Attire* is either *Seminiform*, or *Florid*. And both, *Clustur'd*, or *Divided*; as in *Mallow*, *St. Johns wort*; *Starmort*, *Hawkweed*.

21. §. The *Colours* of the *Flower*, are *White*, as in *Water-Crowfoot*; *Red*, as *Lychnis*; *Blew*, as *Borage*; *Purple*, as *Stock-July-Flower*; *Black*, as in some *Anemones*; *Yellow*, in *Wall-Flower*; *Tawny*, in *Colus Jovis*; *Green*, in *Laureola*. Which are either *Single*, or *Mixed*: *Two* together, as in *Butyr-Bur*, *White* and *Red*; in *White Hellibore*, *White* and *Green*; in *Monks Rubarb*, *Red* and *Green*; &c. Or *Three* together, as in *Pancy*, *Yellow*, *Blew*, and *Black*, i. e. *atro-purpureus*.

22. §. How far these, and some other like *Distinctions*, being reduced to *Tables*, would serve for the finding out of any Sort of *Plant*, may be conceived, if we consider, how great a *Variety*, a few *Bells*, in the ringing of *Changes*, will produce. And the search will be easy, and successfull, if in every foregoing *Table*, reference be made to those that follow; and in the *Tables* containing the last *Divisions*, the *Names* of the *Plants* therein poynted out, be expressed.

THE
ANATOMY
OF
FRUITS,

PROSECUTED

With the bare EYE,

And with the

MICROSCOPE.

Read before the *Royal Society*, in the Year 1677.

The THIRD PART.

By *NEHEMIAH GREW* M.D. Fellow
of the *ROYAL SOCIETY*, and of the
COLLEGE of *PHYSICIANS*.

LONDON,

Printed by *W. Rawlins*, 1682.

FF

THE
CONTENTS
OF THE
Third Part.

CHAP. I.

Of the APPLE; and of the LIMON, and CUCUMER, the Fruits of Plants vulgarly called POMIFEROUS.

CHAP. II.

Of the PEAR and QUINCE.

CHAP. III.

Of the PLUM, and some other Fruits of the same Kindred

CHAP. IV.

Of the GRAPE, and HAZEL-NUT; with some other Fruits analogous to each of them.

CHAP. V.

Of the SEED-CASE or MEMBRANEOUS UTERUS.

CHAP. VI.

Of the USE of the Parts to the Fruit.

CHAP. VII.

Of the USE of the Parts to the Seed. And the TIME, in which, the Uterus or Fruit and Seed-Case are formed.

THE

THE
ANATOMY
OF
FRUITS.

PART III.

CHAP. I.

Of the APPLE; and of the LIMON, and CUCUMER, the Fruits of Plants vulgarly called POMIFEROUS.



THE Description and Use of Leaves and Flowers, together with the Figures thereto belonging, were presented to this Honourable Society, the last year. I shall conclude this Subject with *Fruits* and *seeds*; beginning with *Fruits*, which will take up the present Discourse.

2. §. And First, I shall describe the Compounding *Parts* of some, more generally known. Which having done, I shall next observe the *Uses* of the same; either for the *Fruit* it self, or for the *Seed*. Some of the *Descriptions*, the Reader may be pleased to compare with those in the *First Book. Ch. 6.* I begin with the *Apple*; to which I shall subjoyn the *Limon*, and *Cucumber*, commonly reduced to the *Pome Kind*.

3. §. AN APPLE, besides the *Skin*, consisteth of a *Parenchyma*, *Vessels*, and *Coar*. The *Parenchyma* or *Pulp*, is the same with that of the *Barque* of the *Tree*. As is apparent, not only from the visible continuation thereof from the one, through the *Stalk*, into the other: but also from the Structure common to them both; being both composed of *Bladders*. In which, notwithstanding, there is this difference,

F f 2

That

That whereas in the *Barque*, they are *spherical*, and very small, most of them, through a good *Glass*, not exceeding $\frac{1}{10}$ th of an Inch in *Diameter*, and some of them, less: here, they are oblong and very large, most of them about $\frac{1}{3}$ ^d of an Inch in Length, or more, according to the largeness and tenderness of the *Fruit*; being all uniformly tenter'd or stretched out, by the *arching* of the *Vessels*, from the *Coar* towards the Circumference of the *Apple*.

Tab. 65.

4. §. The *Vessels*, as in the other *Parts* of a *Plant*, are *Succiferous*, and for *Aer*. Both the *Branches* of the former, and the single *Vessels* of the latter, are extream small. They run every where together, not collateral, as *Veins* and *Arteries* do in *Animals*; but the latter, sheathed in the former.

Tab. 65.

5. §. They are distributed into Twenty principal *Branches*. The Ten outmost, a little within the *Apple*, are diverted from a straight *Line*, into so many great *Arches*; from which a few small *Fibres* are without any order dispersed through the *Apple*. The Five middlemost, and the Five inmost, run in a straight *Line* as far as the *Coar*, and are there diverted into as many lesser *Arches*; the former, at the outer, and the latter at the inner *Angles* of the *Coar*. Upon these Five inmost hang all the *Seeds*.

Tab. 65.

6. §. These Ten, and the other Ten abovesaid, do all meet together at the top of the *Apple*, where originally, they all ran into the *Flower*. But betwixt them, there are scarce any intercurrent *Fibres*; so that they appear every where disjunct from the bottom to the top of the *Apple*.

Tab. 65.

7. §. A LIMON hath a Threefold *Parenchyma*; which seem to be derived one from another: the *Texture*, upon every derivation, being somewhat altered, and so made more close and elaborate. The utmost, called the *Rind*, hath the most open, and the coarsest *Texture*; being composed of the largest *Threds*, and those *Threds* woven up into larger *Bladders*. Those little *Cells*, which contain the *Essential Oyl* of the *Fruit*, and stand near the Surface of the *Rind*, are some of the said *Bladders* much more dilated.

Tab. 66.

8. §. From this utmost *Parenchyma*. Nine or Ten *Insertions* or *Lamells* are produced, betwixt as many *Portions* of the *Pulpy Part*, towards the Centre, where they all unite into one *Body*, answerable to the *Pith* in the *Trunk* or *Root* of a *Tree*; and is a conspicuous demonstration, of the communion betwixt the *Barque* and the *Pith*; which there, is much more obscure and difficult to observe. At the bottom, but especially the top of the *Fruit*, the *Pith* is so far expanded, as without the mediation of any *Lamels*, to be joyned to the *Rind*.

Tab. 66.

9. §. Throughout this *Parenchyma*, the *Vessels* are dispersed. But the chief *Branches* stand on the inner Edge of the *Rind*, and the outer Edge of the *Pith*, just at the two extremities of every *Lamel*. From those *Branches* on the Edge of the *Pith*, other little and very short ones shoot into the *Pulp* of the *Fruit*, upon which the *Seeds* are appendant. In the Centre of the *Pith*, are Eight or Nine, in a *Ring*, which run through the *Fruit* up to the *Flower*.

Tab. 66.

10. §. Between the *Rind* and the *Pith* and those several *Lamels*, which joyn them together, stands the second Sort of *Parenchyma*, different from the former, in being somewhat closer, and finer wrought Divided, by the *Lamels*, into several distinct *Bodies*; every one of them a great and entire *Bag*.

11. §.

11. §. Within every great *Bag*, is contained a *Third Parenchyma*, which is also a *Cluster* of other little *Bags*, about the bigness of an *Oate*, all disjoyned one from another, and having their distinct *Stalks*, *Tab. 66.* of several *Lengths*, by which they are all fastned to the utmost Side of the great *Bag*, wherein they are contained. Within each of these lesser *Bags* are contained many hundreds of *Bladders*, consisting of most extreme fine *Threds* woven up together into that *Figure*. Within these *Bladders* lies the *Acid Juice* of the *Limon*.

12. §. A *CUCUMER*, hath also a *Threefold Parenchyma*. The Utmost, is derived, from the *Barque*. In this, being exposed for some time to dry, and then cut transversly with a *Rasor*; not only the *Bladders*, but also the *Threds* whereof the *Bladders* consist, through a good *Microscope*, are apparent.

13. §. Throughout this *Parenchyma* the *Sap-Vessels* are dispersed; near the *Circumference*, in Ten or Twelve very large *Branches*. Each *Tab. 66.* of these larger *Branches*, embosoms another of *Aer-Vessels* in its Centre. Adjacent to the *Midle Parenchyma*, they stand in *Clustres* of much smaller *Branches*, but more numerous.

14. §. Out of all these *Sap-Vessels*, issues a transparent and viscous *Mucilage*; which being dried, becomes as hard and tough as *Gum Tragacanth*. Analogous to which, I suppose, is the truly purgative part of *Elaterium*.

15. §. The *Midle Parenchyma* is derived from the *Pith*; and divided into Three *Columns*, standing triangularly, and having each of them a *Triangular Figure*. Within these *Columns* stand a distinct Sort of *Sap-Vessels*: from whence, several small and short *Fibres* shoot into the *Inmost Parenchyma*, whereupon the *seeds* do hang. So that these *Columns* are as it were the *Beds* on which the *Seeds* grow. With each of the *Seed-Branchs* or *Fibres*, goes some part of the said *Parenchyma* or *Column*, out of which, the *Covers* of the *Seed* are formed. *Tab. 66.*

16. §. The *Inmost Parenchyma* wherein the *Seeds* lie, and which answers to the *Pulp* of a *Limon*, seems likewise to be derived from the *Columns*, that is, to be originally thence produced upon the *Seed-Fibres*, and afterwards spread and augmented into a *Pulp*. By Three *Insertions* from the *Columns*, and as many from the *Utmost Parenchyma*, and these re-inserted; it is divided into Six *Triangular Bodies*; and every *Triangle*, into Three *Ovals*. *Tab. 66.*

17. §. A near resemblance betwixt the *Garden* and *Wild Cucumer*, with respect to the *Inward Structure*, as well as the *Outward Figure*, *Tab. 66.* may be observed: Both of them having a *Threefold Parenchyma*. Yet with this difference, That the Three *White Triangular Bodies* or *Columns* in the one, is answered by a *White Ring* or *Tube* in the other.

C H A P. II.

Of the PEAR and QUINCE.



Tab. 67.

P E A R, besides the *Skin*, consisteth of a Twofold *Parenchyma*, of *Vessels*, *Tartareous Knots* or *Grains*, and a *Coar*. The *Skin* is lined with a great number of the said *Tartareous Grains*, through a *Glass*, about the bigness of small *Shot*: whereby it looks withinside, like the *Skin* of the *Scate* and some other *Fishes*. Besides those which grow to the *Skin*, there are also many more standing near adjacent to it all round about the *Fruit*: altogether about $\frac{1}{3}$ ^d of an Inch in thickness, through a *Microscope*; as in a *Slice* of a *Pear* cut transversly is apparent. Somewhat more or less, as I take it, according to the *Delicacy* or *Harshness* of the *Fruit*; as more in a *Burgamy*, or other soft and sweet *Pear*, than in those which are called *Strangulatoria*. As all *Vinous Liquors*, and those especially which are the most *Tartareous*, become more soft and sweet, according as they cast off their *Tartar*, in a greater quantity, upon the *Sides* of the *Vessel*.

Tab. 67.

2. §. The *Outer Parenchyma*, is of the same *Original*, and general *Structure*, as in an *Apple*. But the *Bladders*, answerable to the *Shape* of this *Fruit*, not altogether so long, with respect to their *Breadth*. Throughout this *Parenchyma*, are also dispersed many small *Tartareous Grains*; most of them somewhat round, as those next the *Skin*, and of a like *Size*; but nothing near so numerous.

Tab. 67.

3. §. The *Bladders* here, have also a different *Position* from that they have in an *Apple*: there, they are all so stretched out, as to have respect to one common *Centre*, which is that of the *Apple* it self. But here, they every where bear a respect to the said *Tartareous Grains*, every *Grain* being the *Centre* of a certain Number of *Bladders*; like a *Star*, in the middle of its *Vortex*. Whereby, so many of the *Tartareous parts* of the *Sap*, as cannot well be thrown off upon the *Skin*, are more commodiously discharged, upon every little *Knot* or *Grain*, nearer hand.

Tab. 67.

4. §. Throughout this *Parenchyma*, the *Vessels* likewise are dispersed. Of the Two general *Kinds*, for *Sap*, and for *Aer*. The *Aer-Vessels*, are here extream small, as well as in an *Apple*; yet one degree, larger. They are both together distributed into *Fifteen* principal *Branches*. The *Five* Utmost make as many *Arches*, but commonly not near so deep as in an *Apple*. From these, some small *Fibres*, yet a little more numerously than in an *Apple*, are dispersed throughout the *Parenchyma*. The *Ten* Inmost run along to the *Seed*, and from thence, with the other *Five*, to the *Flower*.

5. §. Next the *Coar*, stands the *Inner Parenchyma*, in divers respects different from the *Outer*. The *Bladders* of the latter, as hath been said, large and long; of the former, small and round, answerable to those of the *Pith*, of which it seemsto be derived. Throughout that, the

the *Vessels* and *Tartareous Grains* are dispersed; in this, there are neither. The *Effect* whereof is, that is sweet, this sower; for which reason, I have taken leave to name it, the *Acetary*. Tab. 67.

6. §. Betwixt this and the outer *Parenchyma*, the said *Tartareous Grains* begin, first to stand nearer together, to grow bigger, and of a more unequal Surface; and by degrees, to unite into a *Body*, in some *Pears*, and especially towards the *Cork*, almost as hard as a *Plum-Stone*; which I have thereupon, named the *Calculary*. So that a *Pear*, is Nature's *Preface* or *Introduction* to a *Plum*. Tab. 67.
B. I. Ch. 6.

7. §. This *Tartareous Body*, and those small *Grains* above said, I have formerly supposed, to be precipitated out of the *Sap*, by virtue of the *Vessels*. Which is not only argued from their growing, where the *Vessels*, only in the outer *Parenchyma*: but in that the very *Bounds* or *Figure* of the *Calculary*, is determined by the *Situation* of the chief of those *Vessels*; as in cutting a *Pear* smoothly through the *Centre* and by the *Length*, is apparent. B. I. Ch. 6.
Tab. 67.

8. §. The *Coar* as well as the *Acetary*, seems to be derived from the *Pith*. And is therefore lesser here, than in an *Apple*, where the whole *Pith* of the *stalk*, goes to the making of the *Coar* only.

9. §. In most *Pears*, at the bottom of the *Coar*, and a little below the *Centre* of the *Fruit*, there is a kind of small *Umbelical Knot*; from whence is extended a straight *Chanel* or *Ductus*, which opens at the middle of the *Cork* or *stool* of the *Flower*, scarce wide enough to admit the smallest *Pin*. Made for the *Use* hereafter mentioned. Tab. 67.

10. §. A *QUINCE*, is nearly allyed to a *Pear*. The differences betwixt them are these; In the *Quince*, the outer *Parenchyma* is more close, that is, the *Bladders* are smaller. The *Vessels* more numerous, and more deeply enarched; the *Calculary* greater, and more spread; according to the *Shape* of the *Fruit*: but the *Acetary*, less: The *Coar* stands higher or nearer to the *Cork*; divided, not into Five, but Four *Cells*. And the *Ductus* from the bottom of the *Coar* to the top of the *Fruit*, much more open and observable. Tab. 67.

CHAP. III.

Of the *PLUM*, and some other Fruits of the same Kindred.



PLUM consisteth of a *Parenchyma*, the Two general Kinds of *Vessels*, and a *Stone*. All which I have already described in the *First Book*. I shall here add, and further clear some things. And First, it is to be noted, That, in Proportion to the *Bulk* of the *Fruit*, there are more *Vessels* in a *Plum*, than in an *Apple*, *Pear*, or *Quince*. As also, That in *Plums*, all the *Vessels* are braced together into one Uniform Piece of *Net-Work*, every where terminating at an equal distance from the *Circumference*, sc. $\frac{1}{4}$ th of an Inch or thereabout. And as for the *Bore* of the *Aer-Vessels*, although the *Glass* I used, when I examined this *Fruit*, would not reach it; yet Ch. 6.
Tab. 68.

is it to be presumed, that they bear a just Proportion to those in the *Trunk* of the same *Tree*; and that therefore they are here larger, than in an *Apple* or *Pear*. The *Skin* likewise of a *Plum*, is more *fibrous*, *thick*, and *tough*, than in those *Fruits*. The *Ends* of these *Diversities*, we shall presently speak of.

2. §. Of the *Stone*, amongst other particulars wherein the contrivance of Nature is very admirable, I have formerly shewed, That it is composed of Two or rather Three distinct *Bodies*. One of them, the *Lining*; which answers to the *Coar* in a *Pear*. And is originated from the *Parenchyma*, which the *Seed-Branch* brings along with it, through the *Chanel* in the *Side*, and at last into the *Hollow*, of the *Stone*; and is there spread all over it: as when a small *Glass-Pipe*, is blown and expanded into a *Bubble*. Or as if a *Bladder*, being stretch-out, and put through the *Neck* of a *Bottle*; were then blown up, so as to be every where contiguous to the *Sides*, and become, as it were, the *Lining* of the *Bottle*.

3. §. The *Foundation* or *Ground* of the Outer and more Bulky *Part* of the *Stone*, is the *Iner Part* of the *Parenchyma*; and answers to the *Acetary* in a *Pear*. As the *Fruit* grows, the *Tartareous Parts* of the *Sap*, being continually precipitated upon this *Parenchyma*, it is hereby petrify'd. As will best be seen, by comparing the several *Agges* of the same *Fruit* together. And in some *Stones*; on the *Surface* whereof, some of the said *Tartareous Parts* appear in distinct *Grains*. So that whereas in a *Pear*, the *Calculary* and the *Acetary* are distinct, here in a *Plum*, they are thrown one into the other. Or, as some *Mineral Waters* only make a *Crust* about a *Stick* or other *Bodies* immersed in them; but others, by sinking into these *Bodies*, do hereby petrify them: So in a *Pear*, the *Tartareous Parts* of the *Sap*, only make a *Crust* about the *Acetary*; but in a *Plum*, they sink into the *Body* thereof, or that *Part* of the *Parenchyma*, which stands in the place of it, whereby it is converted into a *Stone*. The *Figures* of *Stones* shall hereafter be spoken of, when I come in the next *Part*, to the *Covers* of the *Seed*.

4. §. AN APRECOCK is of the *Plum-Kind*. But some things are herein better observed. As first, the *Position* of the *Bladders* of the *Parenchyma*. For the *Tartareous Parts* of the *Sap* not being here dispersed, in little *Grains*, throughout the *Fruit*, as in a *Pear*; but all thrown off into the *Stone*: the *Bladders* therefore are so disposed, as not to have respect to several *Centres*, as in a *Pear*; but only the *Stone*, to which they all do most exactly radiate; thereto conveying the *feculent Sap*, in so many little *Streams*. This is best seen, when the *Fruit* is full ripe.

5. §. In this *Fruit*, while it is young, the gradual transmutation of the *Inner Part* of the *Parenchyma* into a *Stone*, is also more apparent. And so are the *Three Coats*, which serve for the *Generation* of the *Seed*; being now all very distinct; and remarkable, not only for their *Bulk*; but also, the *Analogy* which they bear to the *Three Membranes* in many *Viviparous Animals*. Whereof I shall give a more particular *Description*, when I come, in the following *Part*, to the *Covers* of the *Seed*.

6. §. A PEACH hath a much bigger *Stone*, than either a *Plum*, or an *Aprcock*: and hath therefore, when full ripe, and especially in hot

hot Countries, a more defecated or better fined *Juyce*. For the reason why the *Stone* is so great, is because the *Vessels* run so very numerously through the Body of it; and so cause a more copious precipitation of the *Lees* of the *Sap* thereinto.

7. §. A *CHEERY* is likewise near related to a *Plum*. But the *Bracement* or *Reticulation* of the *Vessels*, is here carried out further, so as to be all round about contiguous to the *skin*. And as the *Aer-Vessels* in the *Branch* of a *Cherry-Tree*, are larger than those of an *Apple-Branch*, but less than those of a *Plum-Branch*; so may they be presumed, to bear the same Proportion here in the *Fruit*, Tab. 69.

8. §. A *WALNUT*, is a *Nuciprune*; or betwixt a *Plum* and a *Nut*, as a *Bat* is betwixt a *Beast* and a *Bird*. For the *Rind*, answers to the *Pulp*; and the *Shell*, as the *stone*, is also lined. But the *Seed-Vessels*, which in a *Plum* run through a *Chanel* made on purpose in the *Stone*; do here enter, as in a *Nut*, at the *Centre* of the *Shell*. By which means, they are invested with a more fair *Parenchyma*; which Nature hath provided, as her *Cloth*, for the making of the *Coats* wide enough for so vast a *Kernel*.

C H A P. IV.

Of the GRAPE, and HAZEL-NUT; with some other Fruits, analogous to each of them.



GRAPE, is a *Plum* with two *Stones*; for their thickness, as hard as any other. The *Distribution* of the *Vessels* is also somewhat different. For the principal *Fibres* running up directly betwixt the *Stones*; and the smaller, making only one single *Net*, near the *Circumference*; they all meet together at the *Top* of the *Grape*. It is also to be noted, That many *Lignous Fibres* are visibly mixed with the *Skinit* self: whereby it becomes very thick and tough. And as the *Aer-Vessels* in the *Trunk* of a *Vine*, are greater than in that of an *Apple*, *Pear*, or *Plum*: So is it to be presumed, that in a *Grape*, they are greater than in the *Fruits* of those *Trees*. Tab. 69.

2. §. The *Parenchyma* or *Pulp* of a *Grape*, seems to be derived; not from the *Barque*, as in an *Apple*; nor partly from the *Barque*, and partly from the *Pith*, as in a *Goosberry*: but wholly from the *Pith*; at least, as far as the *Reticulation* of the *Fibres*; and the *Skin* only from the *Barque*; whereby the *Pulp* becomes so tender and delicate a *Meat*.

3. §. A *GOOSBERRY*, hath a *Threefold Parenchyma*. The *Utmost* is derived from the *Barque*; of a *Greener Colour*, and very *sappy*. The *midlemost*, from the *Pith*; somewhat *white*, and more *dry*, as the *Diametral Insertions* in some *Roots*. In both of them, the *Bladders* are very conspicuous, above what they are in any *Fruit*, I at present think of; so as to be visible to a good *Eye* without a *Glass*. Tab. 69.

Tab. 69.

4. §. Betwixt these Two *Parenchymas*, do run most of the principal *Fibres*, or *Vascular Threads*. From which several smaller ones are branched into the Inmost *Parenchyma*; upon which, the *Seeds* do hang.

5. §. Each of these smaller *Branches* is invested with some part of the middle or white *Parenchyma*. Serving partly to make the *Covers* of the *Seed*; and partly, the *Pulp*, that is, the Inmost and finest *Parenchyma* of the *Berry*, in which the *seed* lies.

A *White CORIN*, without taking off the *skin*, sheweth not unpleasantly how the *Seeds* are fastned. For as the *Trunk* of the *Tree* continues not to any considerable Length, entire, as in a *Plum*, but is presently divided into several *Boughs*; nor are the *Edges* of the *Leaf* entire, as also in a *Plum*, but slit into several *Lobes*; and the *Fruit*, into a great many *Corins* in a *Bunch*: So again, the *seeds* do hang upon the *Fibres*, like Two other *Bunches*, in every *Corin*. As by *Refraction*, Objects of all *Sizes* are represented on the *Walls* of the *Eye*. The *Operations* of *Nature* being every where *Uniform*: and sometimes the same in small, transcribed from a greater *Copy*.

Tab. 69.

7. §. A *NUT*, is a *Plum* inverted, or turned inside outward. For the *Shell*, standing naked, includes the *Parenchyma*: the bearded *Cap*, not precisely answering to that, but to the *Empalement* of the *Flower*; which likewise in many other *Plants*, out-lives the *Foliature* and Embosomes the *Uterus* of the *Seed*. And whereas the *Stone* of a *Plum* is not *Faced*, but *Lined* with a *Parenchyma* derived at second hand from the *Pith*: The *shell* of a *Nut* is not *Lined*, but *Faced* with the inner *Skin* of the *Cap*.

8. §. AN *AKERN*, is the *Nut* of an *Oak*. Yet with this difference; That besides the *Cup*, it stands in, it hath only a *Leathern* or *Parchment Cover* instead of a *Shell*. From whence it come to pass, that whereas the *Kernel* of a *Nut* is sweet; that of an *Akern*, is of a very rough *Tast*: the *Austere Parts* of the *Sap*, which in a *Nut* are drained off into the *Shell*, being here imbibed by the *Kernel* it self.

C H A P. V.

Of the SEED-CASE or MEMBRANEOUS
UTERUS.

O the forementioned *Fruits*, I shall subjoyne, in some Examples, the *Description* of the *Seed-Case*, which is analogous to the *Fruit*. For the *Fruit*, strictly so called, is, A *Fleshy Uterus*, which grows more moist and *Pulpy*, as the *Seed* ripens. But the *Seed-Case*, whether it be called a *Cod*, *Pod*, or by any other name, is, A *Membraneous Uterus*, which as the *Seed* ripens, still grows the more dry and hard: as in most *Plants*.

2. §. THE *SEED-CASE*, is either originally open; Or only when the *Seed* is ripe; Or never opens at all, till the *Seed* be sown. Of the first Sort, is that of *Luteola*; as also of *Clary*, *Sage*, *Hysop*, and the

the like: wherein one and the same *Part*, is both the *Empalement* of the *Flower*, and when that is gone, survives as the *Case* of the *seed*.

3. §. Of the Last, is that of *Myagrum Monspermon*, *Lithosperme*, all the *Stones* of *Fruits*, with divers others. And some *Cases*, which are soft, as, I think, that of *Garden Radish*. The former, by cleaving in some *part* or other; these only by rotting under Ground.

4. §. THAT of *Garden Radish*, is a Light and Spongy or Pithy *Body*; originally, every where entire. But, as it ripens, breaks within, into several White and Dry *Membranes* round about the *Seed*. By *Tab. 70.* the Length and about $\frac{1}{4}$ th of an Inch distant from the *Sides* of the *Case*, do run a pair of little *Vascular Ropes*. Some smaller *Fibres* are from these transmitted to the *Sides* of the *Case*; by which they are kept tite and steady. Upon divers others produced towards the *Centre*, hang the *Seeds*, like Two *Ropes* of *Onions*.

5. §. Of those which open so soon as the *Seed* is ripe; some are made to open at the *Top*, as *Popy Heads*; Some on the *Side*, as most *Cods*; and some at the *Bottom*, as that of *Coded Arsmart*.

6. §. THE *Popy-Head*, is a little *Dove Coat*; divided by Eight or Ten *Partitions*, into so many *Stalls*. On both *Sides* the *Partitions*, *Tab. 70.* hangs a most numerous *Brood* of *Seeds*. The *Partitions* and *Sides* of the *Head*, are made of the *Barque*, and Lined with the *Pith*. While young, they are very *thick* and *spongy*; and together with the *Seeds*, do then fill all up. The *Head* is then also every where entire; but as it dries, it gradually opens at the *Top*, into several *Windows*, one for every *Stall*: which are all covered with a very fair *Canopy*. A *Fabrick* designed for several purposes, as shall hereafter be said.

7. §. Of those which open on the *side*; some are made to open, only on One *Side*; some, on both *Sides*; some, with Three *Sides*; some, with more; and some horizontally or round about.

8. §. THE *COD* of *Garden Bean* (and so the rest of the *Leguminous* kind) opens on one *side*. It hath a Twofold *Parenchyma*. The Utmost derived from the *Barque*: in which stand all the *Vessels*, in several *Parcels*; one whereof, at the *Back* of the *Cod*, is much larger than the rest, shaped like a *Copula* used in *Schemes*; from whence shoot those lesser *Fibres* upon which the *Beans* do grow. *Tab. 70.*

9. §. The Inner *Parenchyma* is derived from the *Pith*. Upon its Nativity, and for some time afterwards, entire and wholly composed of *Bladders*, as the Outer. From the *Base* of the *Cod* they are gradually enlarged, so as to compose this *Parenchyma* into a very soft and delicate *Sponge*. In which (the *Cod* being well grown) the very *Threds* where- *Tab. 70.* of the *Bladders* were woven, are many of them so loose and ample, as easily to be drawn out (as in the uroaving of *Knit-work*) to a considerable Length, fairly visible through an ordinary *Glass*.

10. §. This may further confirm all that I have formerly said of the *B. 2. P. 1.* *Fibrous Texture* of the *Pith*, and of all the other *Parenchymous Parts* of *Ch. 5. & B.* *Plants.* *3. P. 1. Ch. 4.*

11. §. THE *Seed-Case* of *Medica*, is a *Cod* wound up: in the *Echinata*, *Spirally*; in the *Tornata*, by an *Helix*. Not finished all together; but, upon the fall of the *Flower*, beginning to wind, continues its *Circles*, till it be come to its full *Growth*.

12. §. THE *Seed-Case* of *Yellow Henbean* opens on both *Sides*. On the *Top*, is erected a *Colum*, about $\frac{1}{2}$ an Inch long; which, as the *Case* swells, grows less, and at last falls off. On the *Sides* of the *Uterus* or *Case*, Two *Vascular Fibres* run oppositely from the bottom to the top, and so into the *Colum*. Along the *Tract* of these *Fibres*, the *Case*, as it ages, gradually cleaves on both *Sides* asunder.

13. §. The *Case* is lined with a dry and thin *Parchment*, as smooth as *Glass*. In the *Centre* of the *Case*, stands a great *Parenchymous Body*, which is, as it were, the *Bed* or *Placentula* of the *seeds*; which lie all over it, as in a *strawberry*. And so in many other *Plants*. Throughout this *Bed*, the *Vessels* for the *Generation* and *Nourishment* of the *seeds*, are distributed; one very small *Fibre*, shooting, from the direct ones, obliquely into each *seed*.

14. §. THE *Seed-Case* of *Tulip*, opens with *Three Sides*; being, when young, a *Prism* or long *Triangle*. From the middle of each *Side*, a *Partition* or *Boord* is produced; all three meeting in the *Centre* of the *Case*; and so parting it into *Six Stalls* for the *seed*. The insides hereof, are, lined with a thin smooth and glossy *Parchment*, like that in *Hen-bean*; derived from the *Pith*; as the outside, from the *Barque*: and so in many other *Seed-Cases*.

15. §. The *Vessels*, after they rise above the *Stalk*, are disposed with great *artifice*. For first, they are divided into *Three* principal *Branches*, which run a long the *Three Angles* of the *Case*; where the *Three Sides*, as it ages, gradually cleave asunder. From these chief *Branches*, at the *Three Angles*, divers lesser ones run horizontally, and meet at the middle of each *Side*. From whence again, many yet smaller ones are produced through the breadth of each *Partition* to their *Edges* in the *Centre* of the *Case*. Where, once more, they are distributed into very fine and short *Threads*, whereupon hang the *seeds*.

16. §. THE *Seed-Case* of *Stramonium* or *Thorn Apple*, is divided into *Four Closets*: Not open one into another, as in *Poppy*, *Tulip*, &c. but so many distinct *Inclosures*. In the midst of each *Closet* stands a *Colum*, joyned to the *Side* of the *Closet* by a *Wall* or *Lamine*. Through the Length of the *Colums* run several greater and lesser *Branches* of *Vessels*, from whence others are obliquely produced, upon which the *seeds* grow.

17. §. THE *Seed-Case* of *Anagallis* or *Pimpernel*, is a little *Globe*; which opens not by its *Meridian* or *Vertically*, as do the former; but by its *Horizon*. For divers very small *Fibres*, being produced from the *Stalk* to the middle of the *Case*; do there fetch a *Circle*, and so divide it exactly into *Two Hemispheres*: the *Uppermost* of which, when the *seeds* are ripe, falleth off; and so the wind sows them.

18. §. THE *Seed-Case* of *Coded Arsmart*, neither opens at the *Top*, nor on the *sides*, as do all the former; but at the *Bottom*. It is composed of *Four Sides*: the *Outer Part* of which, is softer and more succulent; the *Inner* a tite and strong *Membrane*. In the *Centre* of the *Case*, is erected a *Pole* or *Colum* upon which the *seeds* do all hang very loosely.

19. §. From this *Mechanism*, the manner of that violent and surprising *Ejaculation* of the *seeds*, is intelligible. Which is not a motion originally in the *seeds* themselves; but contrived by the *Structure* of the *Case*. For the *seeds* hanging very loose, and not on the *Sides* of the

the *Case*, as sometimes, but on the *Pole*, in the *Centre*, with their thicker end downward, they stand ready for a discharge: and the *Sides* of the *Case* being lined with a strong and Tensed *Membrane*, they hereby perform the office of so many little *Bows*: which, remaining fast at the *Top*, and (contrary to what we see in other *Plants*) opening or being lett off at the *Bottom*, forceably curl upward, and so drive all the *Seeds* before them.

CHAP. VI.

Of the Use of the Parts to the Fruit.



IN the forgoing *Descriptions*, I have already mention'd the *Use* of the *Parts* in some particulars. I shall now a little further explain the manner of their service, both to the *Fruit*, and to the *Seed*.

2. §. And first, the *Vessels* serve for the *Figuration* of the *Fruit*. So in an *Apple*, the Ten great and utmost *Branches* serve not only to nourish and feed it; but also, by the *Arched Lines* they draw, to direct and govern the *Growth* thereof into an orbicular *Figure*. The *Dilatation* of these *Vessels*, not being hindred by any *Braces* or *Conjunction* with the *Interior* ones. By the *Slenderness* of the *Aer-Vessels*, as in the *Root*, so here in the *Fruit*, much promoted. And by their *Saline Principle*, first begun.

3. §. The Five midlemost and the Five Inmost serve together, to figure the *Coar*; the former bounding the *Outer*, the Latter, the *Iner Angles*. For were they only Five, or were all Ten in the same *Circle*, they would only make a round *Cavity* like that of a hollow *Pith*. Hence it is that *Apples*, in which some small *Threads* of the *Vessels* strike out into the *Circumference*, are very Uneven with divers *Knobs* and *Ridges*. But *Plums*, *Cherries*, &c. where the *Vessels* all terminate at an Equal distance from the *Skin*, are Even all round about.

4. §. The *Bulk* of the *Fruit* dependeth also on the *Braces* of the *Vessels*. For in *Plums* and *Cherries*, they are more numerous; but in *Apples* and *Pears* they are very loose one from another, and so have liberty left them to spread abroad.

5. §. As also on their *Size*; that is, on the *Size* of the *Aer-Vessels*. Which, the less they are themselves, they serve to make a bigger *Fruit*. As the less they are in any *Root*, they serve to make it the more ample. For the less they are, the more pliable to the *Attraction* of the *Aer*: and in their *Growth* must make so many more spiral *Rings*: by both which means, they make the greater *Arches*. And therefore a *Pear* is commonly a smaller *Fruit* than an *Apple*; a *Plum* than a *Pear*; and a *Grape*, than a *Plum*; in all which the *Aer-Vessels* are still greater and greater.

6. §.

6. §. From the same Cause, it is also most agreeable, That the *Fruit* should not come before the *Leaves* or *Flower*, but last of all. For the *Aer-Vessels*, as hath been often noted, are not exactly *Cylindrick*, but tapered; that is, not only the *Fibres* consisting of divers of these *Vessels*, but the *Vessels* themselves, as they ascend into the *Trunck*, *Branches*, *Leaves*, *Flower*, and *Fruit*, grow still more and more slender. So that the smallest coming last, and being the most pliable; they are also best accommodated for the *Expansion* of the *Parenchyma* into that we call the *Fruit*.

7. §. It is likewise a proper Question to be asked, How it comes to pass, That some *Plants* bear a *Fruit*, and not all? I answer, That as the *Size* of the *Aer-Vessels* conduceth to the *Bulk* of the *Fruit*, and the *Order* of its *Growth*: So the *Number* of them, to their being, or not being, any *Fruit* at all. For the *Fruit*, as we have already defined it, is an *Uterus*, which grows moyster and softer, as the *Seed* ripens. The reason therefore, why the *Uterus* in some *Plants*, continues moist and soft after the *Seed* is ripe; and in some, dries up; is, Because in the former, there is a smaller, in the latter, a greater Quantity of the *Aer-Vessels* in proportion to the other *Parts* of the *Uterus*, and so a greater quantity of *Aer*. Which as in the *Pith* of most *Plants*, so here, by degrees excludes the *Sap*, or rendring it more evaporable, comes in the room of it; and so the *Uterus* is dried up: that is, there is no *Fruit* produced, but only a *Seed-Case*.

8. §. From the *Size*, *Number*, and *Position* of all the *Vessels* in *Fruits* a reason also may be given, for the diversity of their *Tasts*. Some Instances have before been given; to which I shall add one or two more. So the *Rind* of an *Orange*, is bitter; the *Pulp*, sower. Because the former is furnished with many *Lignous Vessels*, the *Sulphureous* or *Oily Tincture* whereof, being copiously mixed with the *Acid* of the *Parenchyma*, produce that *Tast*. Whereas the *Pulp*, which is very sower, is void of all manner of *Vessels*. But if the *Sap-Vessels* are either less numerous or less *Sulphureous*; they give so mild a *Tincture* to the *Parenchyma*, as not to produce a bitter, but a sweet or soft *Tast*; as in *Apples*, *Grapes*, *Goosberries*, &c. And of a *Goosberry*, it is particularly to be noted, that whereas, in a *Limon*, the *Pulp* only is sower, as being void of *Vessels*: here, on the contrary, the *Pulp* only is sweet, whereinto all the *Vessels* strike, and the *Rind* sower.

9. §. The diversities of the *Skin* it self, have their *Use*. And therefore, the more tender and delicate the *Fruit* is; the *Skin*, on the contrary, is thicker and more tough. So *Apples* have a thicker *Skin*, than *Pears*; *Plums*, than *Apples*; and *Grapes* than *Plums*; those having as it were, only a *Coat* of *Kid*, but this of good thick *Buff*. And therefore some *Fruits*, although tender, yet either not having so rich a *Juyce*, or coming early, and so not being exposed to excessive heats, have a very thin *Skin*, as *Mulberries*, *Strawberries*, &c.

CH A P. VII.

Of the *U S E* of the Parts to the Seed. And the *T I M E*, in which the Uterus or Fruit and Seed-Case are formed.



AND first, for example, in an *Apple*, the Five Inmost *Branches*, do best serve for the *Generation* of the *seed*; these running into the *Attire* of the *Flower*, and so carrying off the most *Aerial Spirit* from the *seed*; by which means, it becomes a more compact and denser *Body*, than the *Fruit*, and so more accommodate to the process of *Vegetation*; as *P. 2. Ch. 5. §. 5.* hath formerly been shewed.

2. §. The *Elongation* likewise of the *Seed-Vessels*, in the *Fruit* and *Case*, sometimes directly, as in *Plums* and *Nuts*, and sometimes by several *Ambages* before they shoot into the *Seeds*, as in *Tulip*; shewes a design for the highest refining and maturation of the *seminal Sap*.

3. §. Chiefly by means of the Inmost *Vessels*, is made that *Chanel* in some *Pears*, and especially in *Quinces*. For these perishing with the *Flower*, the circumjacent *Parenchyma* shrinks up, leaving the said *Chanel* in the midst. Designed for an inlet to the *Aer*, for the better drying of the *seeds*; which here stand the more in need of it, because encompassed with a *Mucilage*.

4. §. For the better drying of the *seed*, and the disburfing or sowing of it in due time, the opening of the *Case* is, in the same manner, also contrived: either at the *Top*, as in *Poppy*; or on the *Sides*, as in *Tulip*, *Pimpernel*; or at the *Bottom*, as in *Codded Arsmart*. All which openings are effected by the running of the *Aer-Vessels* along those places: for by drying the *Parenchyma* next adjacent, they cause it to chop and cleave asunder.

5. §. Of the *seed-Case* of *Poppy*, it is particularly to be noted, That as the several *Windows*, serve to let in *Aer*, for the drying of the *Seeds*, after their full Growth: So the *Canopy* over them, serves to keep out *Rain*. For here, the *Case* not cleaving down the *Side*, as it usually doth; should the *Rain* get in, it would stand in it, as in a *Pot*, and so rot the *Seeds*. And as the *Canopy* serves to preserve the *Seeds*; so the several *Partitions* or *Walls*, for their better *Stowage*. For by an easie survey of this little piece of Ground, it is plain, that as they stand on both *Sides* every *Wall*, there is as much more Ground for them to stand upon, as if there were no parting *Walls*, but the *Seeds* stuck all round about upon the *Ambit* or *Sides* of the *Case*; or upon a great *Bed* or *Placenta* within it, as in *Hyoscyamus*, *Anagallis*, &c. where there is a less numerous *Brood*.

6. §.

6. §. The *Coar* likewise, by standing betwixt the moyst *Parenchyma* and the *Seed*, and being hollow and so filled with *Aer*; doth much conduce to the ripening and drying of the *Seed*, and its greater fitness both for keeping, and sowing. So the *Parchment Lining* of the *Seed-Case*, as in *Hyoscyamus*, &c. is answerable to a *Coar*.

7. §. The *Parenchyma* serveth, amongst other purposes, for the *Generation* of the *Covers* of the *Seed*; as in some instances hath been shewed. For which intent, sometimes the *Exterior Parenchyma*, as in a *Limon*; sometimes the midlemost, as in a *Goosberry* or *Cucumber*, is subservient; both of them, in those *Fruits*, being more white and dry, than the rest, and so fiter to make the *Covers* of the *Seed*.

8. §. The *Parenchyma* is also of use for the warmth of the *Seed*; as in the *Seed-Case* of *Garden Radish*. Wherein, as it ripens the *Parenchyma* gradually dries, breaks, and shrinks up into several soft *Membranes*, in which the *Seeds*, in the Centre of the *Case*, lie swaddled, as in so many fine *Calico Clouts*.

9. I SHALL conclude with observing the *Time* of the *Generation* of the *Fruit* and *Seed-Case*. This hath hitherto been thought to be initiated upon the opening, I say not, the forming, but the opening of the *Flower*, or not long before. Notwithstanding which, what I have formerly said of the *Flower*; I now do the like, of the *Uterus* it self; *sc.* That in very many *Plants*, 'tis formed, with the *Flower*, the year before it appears and comes to its full *Growth*. As for instance, in *Azarum*, not only all the *Parts* of the *Flower*, but the *Uterus* it self, and there in also the outer *Cover* of the *Seed* of any one year, are perfectly formed in *August* or *September* of the year foregoing. The like may be seen in *Tulip*, *Mezereon*, *Corin*, and many other *Perennial Plants*.

Tab. 71.

THE
ANATOMY
OF
SEEDS,

PROSECUTED

With the bare EYE;

And with the

MICROSCOPE.

The Figures presented to the *Royal Society*, in
the Year 1677.

The FOURTH PART.

By *NEHEMIAH GREW* M.D. Fellow
of the *ROYAL SOCIETY*, and of the
COLLEGE of PHYSICIANS.

LONDON,

Printed by *W. Rawlins*, 1682.

Hh

THE
CONTENTS
OF THE
Fourth Part.

CHAP. I.

Of the FIGURES of Seeds.

CHAP. II.

Of the NUMBER and MOTIONS of Seeds.

CHAP. III.

Of the several COVERS of Seeds, and of the VITELLUM.

CHAP. IV.

Of the FOETUS or true SEED: and first of the RADICLE and LOBES.

CHAP. V.

Of the BUDS of Seeds. And of the PARTS of which these, the Radicle and Lobes are compounded.

CHAP. VI.

Of the GENERATION of the Seed.

THE
ANATOMY
OF
SEEDS.

PART IV.

CHAP. I.

Of the FIGURES of Seeds.



THE *Figures of Seeds*, or rather of their outward *Covers*, are made suitable, Partly to their *Collocation* in the *Uterus*, as the *End*. So those of *Mallow*, standing like a *Coronet* round the *Stalk*, are of a *wedged Figure*; whereby their sharp *Edges* do all meet together in one *Centre*. Partly; to the various distribution of the *Vessels* or *Fibers*, as one *Cause*: by which the *Measures* and *Surface* of *Seeds*, as well as of the *Leaves* of *Plants*, are diversified. And partly, to the *Nature* of the *Saline* and other *Principles* regent in a *Plant*; as another principal *Cause*. And therefore the more *stony*, *brittle*, or full of *salt* the *Covers* of any *Seeds* are, they are generally more *angular*, and their *Figure*, whether *angular* or not, more constantly observed. So the *Tartareous Stone* of a *Plum*, is not only more *angular*, but also more regular than the *Husk* of the *Kernel* of a *Pear* or *Apple*.

2. §. For all *Stones* are measured by several *Circles*, whose *Diametres* hold a certain proportion to the *Length* of the *Stone*; in the same manner as hath been shewed in the description of the *Leaf*. So *P. I. Ch. 3.* the *Stone* of the *Pease-Cod-Plum*, is measured by two *Circles*. That of the *Turkey-Plum* with Four. That of the *Aprekok-Plum*, with Two *Tab. 72.* repeated oppositely; being perfectly *Rhomboid*. To which, those also of the *Wheat-Plum*, *Damasceen*, and some others, allude. And some are measured by four *Circles*, and one repeated.

3. §. The *Figures*, not only of the larger sort of *Seeds*, but even of the smallest, have much and elegant variety. We will take the pleasure of comparing these which follow.

Tab. 73. 4. §. And first of all, some are perfectly *Spherick*, and with an even *Surface*; as that of little *Century*. That of *Spergula* is also *Spherick*; but hath a knobbed *Surface*, and is encompassed with a *Membraneous Rimm*, like the *Horizon* of a *Globe*. That of little *Celandine* is *Circular*, but compressed like a *Cheese*.

Tab. 73. 5. §. Others are *Nephroideous*, or as it were *Hemispherick*. Of which *Figure*, and hereunto approaching, there are a greater number than of any other; as that which agrees with the more frequent *Shape* and *Fold* of the *Lobes* and *Radicle* of the *Seed*, as shall be seen. Yet with some difference, as to their *Shape* and *Surface*. So, that of *Lychnis Sylvestris* is figur'd just like the kidney of a *Cat*; and hath a knobbed *Surface*. That of *Poppey* comes near it in *Shape*; but hath a *Surface* exactly like that part of the *Paunch* of a *Sheep*, called the *Hony-Coom*. That of great *Celandine*, is a little more oblong; and so, like the *Kidney*, not of a *Cat*, but of a *Sheep*: chequered with parallel *Rings* and other short *Lines* placed alternately betwixt them.

6. §. Where, by the way, we may see, as well by the *Seed*, as by the other *Parts*, of how different kinds, the *Great* and *Little Celandine*, notwithstanding their *Names*, are to be esteemed.

Tab. 73. 7. §. The *Seed* also of *Ben* or *spatling Poppey* is somewhat like a *Kidney*: but hath its *Circumference* raised up into a double *Ridg*: to which several small *Ridges* do in some sort also radiate from one *Centre* sc. the *Base* of the *Seed*.

Tab. 73. 8. §. The *Seed* of *Chickweed*, is partly like a *Kidney*, and partly like a little *Retort*. As also that of *Pentaphil. fragiferum*. But the former is rough cast with small pieces having as it were feet on each side, like little *Insects*. With which, the *Seed* of *Leuchanthemum* (which may be called, the *Giant-Chickweed*) doth much agree. The latter, hath several *Fibrous Ridges*, resembling the *Fibres* in the *Auricles* of the *Heart*; or runing from the nose to the *Circumference*, somewhat like the *Azymuth Lines* on a *Quadrant*.

Tab. 73. 9. §. Some are *Oval*; as that of the little *Bell*, and rough cast with *Fibres* almost parallel and produced by the *Length* of the *Seed*. In which latter respect, the *Seeds* also of *Trachelium* and some other like *Plants*, are agreeable. That of *Brooklime*, is also *Oval*, but encompassed with a thick *Rimm*, narrowing all the way to the *Base* of the *Seed*.

Tab. 73. 10. §. The *Seed* of *Dovesfoot* hath an *oval Cone*, and a flat *Base*. Its *Surface* favours, like that of *Poppy*, *Toad-flax*, and some other *Seeds*. That of *Sedum minus aestivum luteum*, is in a manner the *Figure* of the former inverted, being flat, not at the *Base*, but on the *Top*. And whereas that rises with a blunt *Angle*, this hath only a *Ridg*, raised above the *Surface* of the *Seed*.

Tab. 73. 11. §. The *Seeds* of divers sorts of *Grass*, are more *Conick*, as particularly of that, which for the likeness its *Seed* hath to a *Barly Corn*, may be called *Barley-Grass*. And I little doubt, but that among the several sorts of *Grass*, there are some which answer to all the kinds of *Esculent Grains*, as *Oat-Grass*, *Rice-Grass*, *Wheat-Grass*, *Rye-Grass*. And accordingly, that they may be more profitably sown in one *Ground*, than in another; and used with distinction, for the higher, or more wholesome

wholsome feeding of Cattle. A *Rush*, though it seems an imperfect *Plant*, yet besides its *Flower*, hath also a plentiful brood of *Seeds* of a *Conick Figure*.

12. §. Some *Seeds* are *Cylindrick*, as that of *St. Johns-wort*, as also of *Tutsan*, and some other like *Plants*, with some little diversity in the *Shape* or *Surface* of the *Seed*. That of *Vervain*, is in a manner, half a *Cylinder*: the true *Seed* lying in the *Covers*, like a *Child*, in a *Cradle* without an head. Tab. 73.

13. §. Others are rather *Conico-Cylindrick*, as that of *Jacobæa*; having a *Coronet* on the top, and several furrows by the *Length* round about. Answerable to which, is that of *Erygerum*; in *Shape* not unlike to a *Rowling-pin*. Tab. 73.

14. §. Some are *Plani-Conick*, as that of *Nettle*, which is shaped somewhat like the end of a *Speer*. That of *Eye-bright* is more *Eliptick*; with several *Ridges* running by the *Length*; and joyned together with short pieces transversly, as in the looping of *Lace*. That of *Worm-mood* not very unlike a little flat *Essence-Glass*: in which, the *Fibrés* are produced by the *Length*, as the *Ridges* are in *Eye-bright*. And so in *Yarrow*, which is also encompassed with a *Membraneous Rimm*. That of *Dandelyon*, is *Plani-Conick* towards the *Base*. And so those of *Lettice*, *Sonchus*, and some others. To which, those also of *Hieraceum*, *Tragopogon*, *Scorzonera*, &c. with respect to their *Surface*, do all allude. Tab. 74.

15. §. And some are *Conico-Triangular*. Of which, that of *Sorrel* is *Conick* at both ends; the sides equal; and upon every *Angle*, hath a narrow and sharp *Rimm*. As also that of *Anagallis*; but the *Sides* are *Spheri-conick*, and so the ends are blunt. They are also pounced with many little round *Cavities*. But have no *Rimm* upon the *Angles*. Tab. 74.

16. §. The *Seed* of *Nigella* is *Triangular*, and *Conick* only at the *Top*. On every *Angle*, hath a narrow *Rimm*; the three *Sides* equal, and *Spheri-conick*; surrounded with seven or eight *Ridges* by the girth, joyned together in some places with others transversly. That of *Ar-smart*, is also *Triangular* and *Conick* at the *Top*. But one of the *Sides* is almost equal to the other two; which stand low. That of *Knot-Grass* hath three *Sides*, one less than another; being as 5, 3, and 2, or thereabout. Tab. 74.

17. §. The next (which I take to be the *Seed* of a sort of *Bugloss*) is very oddly figured. The *Base*, *oval*; the *Top*, *conick*; the *Back*, swelling and round as an *Egg*; the *Belly* also swelling, but rising up into an obtuse *Angle* highest in the middle, somewhat like a *Breast-piece* of *Armour*: and is encompassed with a *Rimm* sloaped upward. Tab. 74.

18. §. That of *Moldavian Bann*, is *Triangular*, and *Conick* only at the *Base*. The place where it is fastned, shaped like the *Beard* of a *Dart*. Two of the *Sides* are *Plani-conick*, the *Third Sphericonick*, and near as big as both the other two. The *Head* flat, with a *Rimm* erected upon each *Side*, so as to make a *Spherical Triangle*. Approahing to this, are those of *sage*, *Horehound*, *Clary*, &c. Tab. 74.

19. §. That also of *Bellis Tanaceti folio*, hath two *Sides Planiconick*, and a third *Sphericonick*. The two first have several *Ridges* running to the *Base*. Which is not perfectly *conick*, but a little dilated into two obtuse *Angles*. The *Head Triangular*, with one *Side* convex, the Tab. 74.

the other Two straight, a little hollowd, and having a small pinnacle in the Center.

20. §. That of *Stachas Arabica*, as the former, saving, that the Head is oval, and the Base sloaped into a little Triangle. That of *Wartwort* or *Sun-Spurge*, hath a very complex Figure. The Belly consisteth of two *Planiconick* Sides, as the former; the Back, *Sphericonick*. The whole Seed, in a manner, *Conick-oval*. Yet the Base and Head both flat. In the middle of the former, a Peg by which the Seed is fastned; and of the latter, a poynted *Knob*. The middle of the *Belly-Sides*, hollowed, so as to make a flat *Rimm* of equal Breadth; and the hollows filled up with *Bladders* like those in all the *Parenchymus Parts* of a Plant.

21. §. Lastly, there are some *Seeds* which are square. Whereof some are straight, as that of *Fox-glove*; which hath also an even Surface: And that of *Blattaria*, in which there are several little hollows in even Rows. And so in *Brounwort*.

22. §. And some *Convex*, as that of *Chrysanthemum Americ*. 'Tis *Quadrati-conick*, or square and sharp at the Base, and big at the Head. The sides all plain; and a thin *Rimm* erected upon every Angle. As also on the four Sides of the Head, which is flat, with a little *Pinnacle* in the middle.

23. §. The Seed also of *Tansy*, is a *Conick* and bended square not with the Angle forward, as the former, but the Side. And in the place of every *Rimm*, hath a round *Ridge*. Somewhat like to this, are those of *Febrifuga*, *Mayweed*, and some others. Thus far of the Figures of Seeds.

CHAP. II.

Of the NUMBER and MOTIONS of Seeds.



NATURE hath secured the Propagation of Plants several ways, but chiefly by the Seed: for the Production of which, the Root, Leaves, Flower, and Fruit, do all officiate, as hath been shewed. And according as the Plant, or the Seed it bears, is more liable to be destroyed, Provision is made for Propagation, either by a greater number of Seeds, or other ways. So the Seeds of *Strawberry*, being gathered, or eaten by *Vermin*, with the Fruit; the Plant is therefore easily propagated by *Trunk-Roots*. So *Poppy*, being an annual Plant, is highly prolifick: for instance, the *White Poppy*; which commonly bears about four mature Heads, in each of which, there are at least ten Partitions, on both sides whereof, the Seeds grow; and upon $\frac{1}{4}$ th part of one side, about 100 Seeds; that is, 800 on one Partition: which being multiplied by 10 (the number of Partitions) makes 8000; and 8000 again by 4 (the number of Heads) makes 32000 Seeds, the yearly product of that Plant.

2. §. So in *Typha major*, the *Seeds* being blow'n off and low'n (as the Eggs of many *Fishes* spawn'd) with great hazard, they are strangely numerous. For as they stand altogether upon the *Spike*, they make a *Cylinder* at least six Inches long, and near $\frac{3}{8}$ ths of an Inch in Diameter, or an Inch and $\frac{1}{4}$ about. Now 9 of these *Seeds*, set side to side, as they stand on the *Spike*, make but $\frac{1}{8}$ th of an Inch; so that 72 make a line of an Inch in Length. But because upon the *Spike*, the *Hairs* belonging to the *Seeds* come between them; we will abate 10, and count but 62. To which $\frac{3}{4}$ ths of 62, that is (without the Fraction) 46. being added, makes 108 for the Circuit of the *Cylinder*. And the *Cylinder* being six Inches long, there are six times 62, that is, 372, for a Line the length of the *Cylinder*. Which number being multiplied by 108, produceth 40176 the number of *Seeds* which stand upon one *Stalk*; and so, upon three *Stalks*, which one *Plant* commonly bears, there are in one year, above a hundred and twenty Thousand *Seeds*.

3. §. SO SOON as the *seed* is ripe, Nature taketh several *Methods* for its being duly sow'n: not only in the opening of the *Uterus*, as in some Instances (a) hath already been seen; but also in the make (a) P. 3. of the *Seed* it self. For *First*, the *Seeds* of many *Plants*, which affect Ch. 5. Tab. a peculiar *Soil* or *seat*, as of *Arum*, *Poppy*, &c. are heavy and small 70, & 71. enough, without further care, to fall directly down into the *Ground*: and so to grow in the same place where themselves had their *Birth*.

4. §. But if they are so large and light, as to be exposed to the wind, they are often furnished with one or more *Hooks*; To stay them from straying over far from their proper place, till by the fall of *Leaves* or otherwise, they are safely lodged. So the *Seeds* of *Avens* have one single *Hook*, those of *Agrimony* and *Goose-grass*, many; both the former, loving a *Bank* for warmth, the latter, a *Hedge* for its support. Tab. 72.

5. §. On the contrary, many *Seeds* are furnished with *Wings* or *Feathers*. Partly, with the help of the *Wind* to carry them, when they are ripe, from off the *Plant*, as those of *Ash*, *Maple*, *Orach*, &c. least staying thereon too long, they should either be corrupted, or miss their season. And partly, to enable them to make their flight, more or less, abroad: that so they may not, by falling together, come up too thick; and that if one should miss a good *Soyl* or *Bed*, another may hit. So the *Kernels* of *Pine* have wings not unlike to those of some *Insects*; yet very short, in respect of the weight of the *Seed*; Tab. 72: whereby they flye not in the *Aer*, but like domestick *Fowls*, only flutter upon the *Ground*. But those of *Typha*, *Dandelion* and most of the *Pappous* kind, with many more, have very long and numerous *Feathers*, by which they are waisted every way, and to any distance necessary for the aforefaid purposes.

6. §. Again, there are some *Seeds*, which are scattered not by flying abroad, but by being either *Spurtd*, or *Slung* away. The first are those of *Woodforrel*; which having a running *Root*, Nature sees it fit to sow the *Seeds* at some distance. The doing of which is effected by a white thick and sturdy *Cover* of a *Tendinous* or *Springy* Nature, in which the *Seed* lies within the *Case*. This *Cover*, so soon as it begins to drye, bursts open on one side, in an instant, and is violently turned inside outward, as you would turn the *Gizzard* of a *Fowl*; and so smartly throws off the *Seed*. Tab. 72.

7. §.

Tab. 72.

7. §. The *Seeds* of *Harts-tongue*, and of all that *Tribe*, are *Slung* or *Shot* away. The doing of which is performed by the curious contrivance of the *Seed-Case*; as in *Codded Arsmart*, and some other like *Plants*. Only there, the *Spring* moves and curls up inward; but here it moves outward. I shall describe it, as well as the *Weather* (which when I observed it was cloudy) would permit. Every *Seed-Case*, as it appears through a good *Glass*, stands upon a *Pedicle* from $\frac{1}{2}$ an Inch to an Inch or more in Length; at the bottom about as thick again as a *Horse hair*, and a little thicker at the *Top*, on which stands the *Case*, of a *Silver Colour*; about the bigness of a *Cherry-stone*, of a *Spherick Figure*, and girded about with a sturdy *Tendon* or *Spring*, of the *Colour* of *Gold*: the whole *Machine* looking not much unlike a little *Padlock*. The *Surface* of the *Spring* resembles a fine *Screw*, or some of the *Aer-Vessels* in the *Wood* of a *Plant*. So soon as by the *Innate Aer* of the *Plant*, or otherwise, this *Spring* is become stark enough, it suddenly breaks the *Case* into two halves, like two little *Cups*, and so flings the *Seed*.

Tab. 72.

8. §. These *Cases* grow in oblique *Furrows* or *Trenches* on the back side the *Leaf*, from $\frac{1}{4}$ of an Inch to an Inch in Length, and about $\frac{1}{2}$ th of an Inch broad. In one of these *Trenches* an Inch long are more than 300 of the *Cases* above described; and allowing but 10 *Seeds* to every *Case*, above 3000 *Seeds*. Which being multiplied by the number of *Furrows* in one *Leaf*, with allowance for the lesser *Furrows*; and that sum by the number of *Leaves* commonly growing upon one *Root*, comes to above Ten Hundred Thousand *Seeds*, the annual product of this *Plant*. The *Seed* is of a *Tawny Colour*, through a good *Glass* about $\frac{1}{2}$ th of an Inch long, flat, and somewhat oval. Of these, ten Thousand are not so big as a white *Pepper Corn*.

C H A P. III.

Of the several COVERS of Seeds, and of the VITELLUM.



Tab. 72.

Tab. 72.

THE next step of *Natures Managery*, relates chiefly to the Growth of the *Seed* when it is sow'n. For which purpose, the outer *Covers* are somewhere furnished with *Apertures* sufficient for the reception of *Alimental Moyster* from the *Ground*; and *Divisions*, for the shooting forth of the young *Root* into it. As in the *Seed* of a *Gourd*, at the *Bottom*; in a *Bean*, on the *Side*; and in a *Chestnut*, at the *Top*: in which places the *Radicle* or young *Root* always lies and puts forth, in the said several *Seeds*. And the *Seed* of *Palma Christi*; which falls to the *Ground* not only in the usual *Covers*, but also in the *Seed-Case*, for the more plentiful admission of *Aliment*, hath a double *Aperture*. Not much unlike to this, is that found sometimes

in

in larger parcels of *Euphorbium*; for which Cause, I suspect it to be the Gumm of a Plant of the *Tithymal* kind.

2. §. If the Cover of the Seed be stony and very hard, it is also distinguished into several Pieces; whereby they easily cleave asunder without much resistance to the eruption of the Root. So the Shell of a Hazel-nut easily cleaves on the edge; and the cleft begins best at the poynt, where the Root stands and shoots forth. The Shell of some Walnuts cleaves into three Parts; and the Stone of the Bellerick Myrobalan into five: that so, being very thick and hard, if one piece should not yield, another may not fail to do it. And the Covers or Husks of some sorts of Grain, as of Millet, are only folded or laped one over another, the better to give way to their tender Sprouts. Tab. 75.

3. §. Besides the Kernels of Plums and some other Fruits, there are very many Seeds, even of the smaller sort, which have also stony Covers; as of Carthamum, Myagrum monospermon, Beet, Borage, Lithosperme, Amaranthus, Violet, &c. Sometimes, for the reception of the harsher and less matured Principles from the Seed, in its Generation, as in Borage. Commonly, to keep it warmer before and after its sow'n. For which purpose, the outer Covers of some Seeds, are as it were lined with Fur: in that of Great Maple, Short; of Gossipium, Long. And if the Seed requires a longer stay under ground, the hardness of the Cover serves to stint the Aliment; lest too much, should either rot it, or cause it to germinate, before its proper season, or full time for a more Masculine Growth.

4. §. On the contrary, many Seeds, as those of Clary, Garden-Cress, and others of that Tribe, have their upper Covers faced with a Mucilage: which being easily receptive of any Moisture in the Ground, gradually swells, till it lies like a Gelly round about the Seed. Either for a more plentiful supply of Aliment; or at least, to soften the Covers, the better to accelerate the Growth of the Seed.

5. §. The process of Nature in the several steps of the Vegetation of the Seed, hath formerly been explained. (a)

6. §. THE COVERS of all, or at least the far greater number of Seeds, are Three; some way or other derived from the Pith: as shall hereafter be seen. And sometimes, Four: even those of ston'd Fruits, have Three, besides the Stone. In that of Gossipium, there are Two Coats under that lined with the Cotton. The Seeds of Cucumier, Goats-beard, Broom, Scabious, Lettice, &c. although so small, have plainly Three Coats. But in some of these, and many more, there are only Two distinctly visible, except in the State of Generation. (a) B. 1.
Ch. 1.
Tab. 75.

7. §. In the Upper Coat, the Seed-Vessels are disseminated. The Second, is first a meer Pulp; but afterwards shrinks up and sticks close to the upper. The Third or Inmost is more dense; and if it be thin, for the most part, transparent; whereby the Seed seems sometimes to be naked while it lies therein; as in Almonds, Cucumers, and the like. For this sticks not to the middle Coat, as that doth to the outer; but commonly, remains entire, after those are stripp'd off, being as it were, the Smock of the Seed. Tab. 76.

8. §. In Melissa and some other small Seeds, it comes finely off upon soaking in warm Water or on the Tongue. In Fenugreek, 'tis soft, and of an Amber-Colour; and being moistened, looks almost like fine Glem. But commonly, 'tis a pretty tough Membrane, and often with

some thickness, as in *Plums*, *Borage*, *Scabious*. Yet always extream thin at the *Tip* of the *Radicle*; the more easily to break and yield to it, as the *Secundine* to the *Fœtus*, when it first shoots into the *Ground*. And sometimes, as in the *Seeds* of an *Orange*, it hath at one end, the resemblance of a *Placenta*. But of this, and the two upper *Coats*, I shall give a further Description in the last *Chapter*.

Tab. 75.

9. §. AS ALL *Seeds* are *ex Ovo*; so there are many with thin *Covers*, as of *Orach*, *Spinage*, *Beet*, and the rest of that *Tribe*, &c. which besides the *Albumen* or clear *Liquor* out of which they are bred; have also, a *Vitellum*, or a *Body* thereunto *Analogus*: being neither part of the *Seed*, nor part of the *Covers*, but distinct from them both. With respect to the *Bulk* of the *Seed*, very large, as white as *Starch*, and pretty friable, like good *Rice* or *Barley*: of a roundish *Figure*, and grooved on the *Girth*, so as to have a double *Edge*; Whereby the *Seed*, which is long and slender, lies round it, as a *Sack* of *Corn* upon a *Pack-Saddle* or a *Rope* upon a *Pully-wheel*. Upon my first notice hereof, it seemed to answer to a *Placenta*. But upon further consideration, the *Analogy* doth not hold betwixt them. For the *Placenta* lies without the *Membranes* in which the *Fœtus* is contained: whereas this body lies within the *Covers* contiguous to the *Seed*, and so becomes its first and finest *Aliment*, as the *Yelk* doth to the *Chick*. For which purpose, as in the *Generation* of the *Seed*, it is a pure *Milky Chyle*; So in its *Vegetation*, it is converted into the like again.

Tab. 75.

10. §. The same *Body* for *Sustance*, is observable in the *Seeds* of *Rhapontick*, *Dock*, *Sorrel*, and the rest of that kindred, with this difference; That whereas in *Orach*, &c. the *Seed* only lies upon it; here, the main *Body* or *Lobes* of the *Seed* are immerfed therein, the *Radicle* standing naked or above it. So that the said *Lobes*, and therein the *Seminal-Root* are beded herein, as in a *Tub* of *Meal* or a little pot of pure refin'd *Mould*, necessary for the first *Vegetation* of the *Radicle*.

Tab. 75.

11. §. BY THESE middle *Steps*, *Nature* proceeds from the *Thiner Covers* of *Seeds*; or those, which after the *Generation* of the *Seed* is finished, shrink up; to the *Bulky Kind*, or those which keep their *Bulk* after they are dry. Wherein, not only the *Lobes*, as in *Dock*, but the whole *Seed* is immediately lodged. Different in *Substance*, *Shape* and *Bulk*; but always many times bigger than the true *Seed* within it: for which it is commonly mistaken; but is no more the *Seed*, than is the *Stone* of a *Plum*, the *Kernel*.

12. §. In the *Barbado Nut*, 'tis *White*, *Soft*, *Conick-oval*, and taking all its *Dimentions*, 8 or 10 times bigger than the *Seed* within it. In *Ashen Keys*, 'tis of a sad *Colour*, *hard*, yet somewhat *Oily*, *Oval* and *flat*, and of the same *Bigness* as in the *Barbado Nut*, with respect to the *Seed*. In the *Fruit* commonly called *Nux Vomica Officinarum*, 'tis of the *Colour* and *Hardness* of a *Cows-Horne*; and makes almost the whole *Body* of the *Fruit*, being about 14 or 15 times the *Bulk* of the *Seed*. In *Goasgrass* or *Cliver* 'tis of the like *Horny Substance*, but shaped somewhat like a *Bonet* with the *Rimm* tuckt in. And so in a *Coffee-Berry*; but rowled or foulded up into a kind of *Oval Figure*, with a *Notch* or *Rima* through the *Length*, where the two *Ends* meet. With other diversities which will best be understood, when I come presently to the *Description* of the *Seed* herein contained.


Tab. 76.

Tab. 77.

13. §. With respect to the use of this *Cover*, it is observable, that where there is a *Stone* or *Shell* over it, as in the *Barbado Nut*, it is *soft*; but where there is none, as in *Nux Vomica*, *Asb*, &c. 'tis *hard*; and so it self instead of a *Stone*. As also, That it becomes *hard*, only by the proper Nature of its *Parenchyma*, and the exquisite *smallness* of the *Bladders* of which it consists. Whereas a *Stone*, is also hardened by the *Lees* or *Tartar* of the *Sap* which sinks into it, and thereby *petrifies* it (a) as hath been said. So that whereas a *Stone*, as it lies in the *Ground*, (a) P. 3: only cleaves in certain *Places*, but continues *hard*: This *Cover*, like some Ch. 3: *Horns*, upon the due accession of *Moisture*, doth gradually become *soft*. Whereby, as while it is *hard*, it performs the Office of a *Stone*, in guarding the *Seed* til the proper *season* for its *Growth*: So afterwards when it is *soft*, it answers, as in *Orach* or *Dock*, (b) to a *Vitellum*, (b) Ch. 3: from whence the *Seed* receiveth its first and purest *Aliment*. §. 8, 9:

CHAP. IV.

Of the FOETUS or true SEED: and first of the RADICLE and LOBES.



HAVING discoursed of the *Covers*, I come next to the *Seed* or *Fœtus* it self. Of the *Shape* and *Posture* whereof, I shall give some *Examples*; first, among those with the thinner sort of *Covers*; and then, of those with the *Bulky* one: where I shall speak only of the *Lobes*, or *Main Body*, and the *Radicle*. Next, I shall describe the several sorts of *Nodes* or *Buds* of *Seeds*. And lastly, the several *Parts*, of which the *Lobes*, *Radicle*, and *Buds* are compounded.

2. §. Among *Seeds* with the *Thinner Covers*, are those of all sorts of *Corn* and *Grass*. Of a different make, from that of most other *Seeds*: The *Main Body* being not divided into *Lobes*, but one entire *Piece*, doubled in the form of a *Pair of Lipps*. And whereas commonly, the whole *Seed* is very *soft* and *Oily*; here, only those two minute *Parts*, which become the *Root* and *Stalk*, are so: The *Main Body* being of a different *Substance*; when the *Corn* is ripe, *hard* and *friable*; but when it is sown, easily *colliquable* into a kind of *Milk* or *Chyle*; so that, in some respects, it hath a near *Analogy* to a *Vitellum*. For as that is gradually melted into a sort of *Chyle*, and by the *Branches* of the *Ductus Intestinalis* carryed into the *Bowels* of the *Chick*: So is this, into a like *Substance*, and by the *Branches* of the *Seminal Root* (formerly describ'd) convey'd to those *Parts*, which become the future *Plant*. B. I. Ch. I.

3. §. Of Relation to this Kind, the *Seeds* of *Dates*, and of some other like *Plants*, may be esteem'd. For that which is commonly call'd the *Stone*, seems indeed to be the *Main Body* of the *seed*, doubled or folded up in the same manner as in *Corn*. To which that *Part* which becomes the *Plant*, is annexed; But whereas in *Corn*, 'tis placed at Tab. 75.

the Bottom of the *Main Body*; here it lies in a small round *Cavity* in the middle of the *Back*. The *Stone*, or *Main Body*, where this Part grows to it, is not so hard, as more remote from it: and is therefore probably in some part dissolved, by lying in the *Ground*, as in *Corn*.

4. §. But for the most part, the *Main Body* is divided, as hath been said, into two *Lobes*; and those in Substance *Homogeneous* to the other Part or Parts, plainly distinguished in most *Kernels* and other large seeds; and not difficultly in many lesser ones, as in that of *Viola Lunaris*, *Scabious*, *Doves-Foot*, &c. if slipped out of their *Covers* before they are full ripe.

5. §. In *Hounds-Tongue*, they are of a circular figure, and very large in Proportion to the *Radicle*. In *Cucumber*, oblong, with some visible *Branches* of the *Seminal Root*; and the *Radicle* somewhat bigger. But in *Scorzonera*, very long, like the Leggs of a Pair of *Compasses*: and the two first, or *dissimilar Leaves* of the *Plant* into which they are converted, are of the same *Shape*. Of these and many more, the *Radicle* is short and pointed; and lies in one straight Line with the *Lobes*.

6. §. In *Viola Lunaria*, they are very large; and the *Branches* of the *Seminal Root*, fairly apparent, so as to resemble a Pair of *Leaves*. The *Radicle* pretty long, equally thick from end to end, and couched down upon the two *Lobes*, each of them having a little *Shoulder* for it to lie upon. In *Woad*, where it hath the like *Posture* and *Shape*, as also in *Chamelina*, *Eruca*, and many others, it is very *Bulky*, being bigger than both the *Lobes* put together.

7. §. Of this Part, I think it may be observed, That commonly those seeds, wherein it is very small with respect to the *Lobes*, produce a *Perennial Plant*: And so, *vice versa*, where it is very large, an *Annual* one. In the latter, the *Seminal Virtue* being more vigorous, and so tending more hastily to the *Business* of *Generation*, followed with the *Death* of the *Plant*.

8. §. IN THE former Seeds, the *Lobes* lie flat one against another. But in *Garden-Radish*, they are folded up, so as to receive the *Radicle* into their *Bosome*: as when a *Chicken* tucks his Head under his *Wing*.

9. §. In *Holyoak*, the *Lobes* are plated upwards, and re-plated down again. Being most agreeably composed to the *Shape* of the *Covers*, as those are to their *Posture* on the *Plant*. In *Maple*, they are plated one over another, and so rouled up.

10. §. In the *Cotton-Seed*, which consisteth almost wholly of two very broad and thin *Lobes* or *Leaves*, the *Folds* are yet more numerous; all curiously reduced to an exact and solid *Oval*.

11. §. It happens now and then, that instead of two, there are three *Lobes*, as in the *Kernels* of *Plums*, *Apples*, and other *Fruits*, and the smaller sorts of *Seeds*, will spring up sometimes with more than two *dissimilar Leaves*, originally the *Lobes* of the *seed*. These are observed by some, more frequently to produce a double *Flower*, which may be, because the *seminal Virtue* in such *Seeds*, is increased by a third Part.

12. §. IN many *Seeds*, the *Radicle* is of one and the same *Colour* from end to end. But in others, as in the *Lupine*, it is observable; That the upper and greater half, is *White*; the Lower to the *Point*, hath a kind of *Horny Gloss*, and seems to be of a somewhat different make. *Tab. 75.* Whereby it comes to pass, that after the *Radicle* is shot forth a little way, only this lower half descends and becomes the *Root*: The upper half is produced or raised above ground, as a *Pillar* upon which the *Lobes*, or *dissimilar Leaves* are erected.

13. §. This *Seed*, on the out side of each *Lobe*, and near the *Radicle*, hath a very small and round *Node*, like a *Navel*; whereof, in the first Book: the whole *Seed* looking not much unlike a *Pidgeons Head*; *Ch. 7.* the *Radicle* resembling the *Bill*, and the *Navel* the *Eye*.

14. §. IN the *Seed* of *Garden-Orack*, both the *Radicle* and *Lobes* *Tab. 75.* are very long and slender, and lie almost in a compleat Circle round about the *Vitellum* before describ'd. The *Lobes* of *Rhapontick* are shaped like the *Bitt* of a *Spade*; and the *Radicle* stands erected above *Ch. 3.* them like the *Handle*.

15. §. OF SEEDS also with the *Bulky Cover*, there are many not divided into *Lobes*; being in a manner, all one *Piece*; as all of the *Bulbous-Kind*. In some of which, though the inmost *Cover* be thin; yet compared either with the other *Covers*, or with the *Seed* it self, it may very well be accounted of the *Bulky-Kind*.

16. §. In *Flag*, it is above twenty times bigger than the *Seed* within it. Consisting of *Bladders* all *Radiated* towards the *Seat* of the *Seed*. The *Seed* it self is shaped somewhat like a *Penknife*. The lower Part *Tab. 76.* which becoms the *Bulb*, as the *Haft*, is thick, and cometh near to a *Cylindrick* Figure, and the end, round. The upper Part which becomes the first years *Leaf*, as the *Blade*, is rather flat, double edged, and pointed, and the *Point* a little bent. The *Fibers* and *Bladders* of which it consists, are all disposed into *Parallel Lines* running by the length. In *Lily*, where this *Cover* is thinner and more *Transparent*, without being cut, but only held up against the *Light*, the *Seed* may be seen within it.

17. §. BUT THE greater number of *Seeds* also with the *Bulky Cover*, are divided into two *Lobes*; which, for the most part, resemble a pair of little *Leavs*. In the *Purging Nut* of *Angola*, the *Shell* being taken off, the upper *Covers* (dry'd and shrunk up) seem to be but one. *Tab. 78.* In these, the *Spermatick Vessels* are Branched. Under these, lies the *Thick and Inmost Cover*; which being cut down the middle, exhibits the true *Seed*: Consisting of a couple of fair *Leavs*, Veined, and as white as *Milk*, joyned together with the *Radicle* at their *Base*; and let into a *Hollow*, made in the *Cover*, of an answerable shape. The like is observable in the *Barbado-Nut*, *Ricinus Americanus*, and some other *Indian Fruits*; with some little difference in the *Shape* of the *Root* and *Leavs*.

18. §. IN the foregoing *Fruits*, the *Bulky Cover* is very soft. But in the *Nux Vomica Officinarum*, 'tis near as hard as a *Date stone*. *Tab. 76.* In this, besides the hollow made for the reception of the *Seed*, or the two *Leavs* and *Root*; the *Sides* are separated or distinct almost to the *Edge* of the *Cover* round about, especially towards the *Root*: So that it may not unaptly be compared to a little *Pouch* with the *Sides* clapt together.

19. §. IN this and the *Nuts* above mentioned, the *Seeds* are all very large. But in some other *Plants*, they are extream small, so as to be hardly visible without a *Glass*; as in *Staphisagria*, *Peony*, &c. In *Staphisagria*, the *Thick* or *Inmost Cover*, is commonly a *Spherical Triangle*, conick towards the *Base*. At the poynt of which, there is a little *Cavity*, in which the *Seed*, about as big as a small pins head, is lodged. The *Root* whereof is a little poynted, and the two *Lobes* rounded at the *Top*.

20. §. In *Peony*, the same *Cover* is *Soft*, *White*, and of an *Oval Figure*; the part used in *Medicine*. Usually thought to be the *Seed* it self. But is near two hundred times bigger than the true *Seed*, which is almost invisible. It lies in a little *Cavity* near the bottom of the *Cover*; with a thick and blunt *Root*, and two poynted *Lobes* or *Leavs*.

23. §. IN the *Coffee-Berry*, the *Seed* lies in the *Inner* or *Cartilaginous Cover* (formerly described) where one would not expect to find it, *sc.* near the *Top* or *Surface* of the *Back*. The *Lobes* of the *Seed* are veined like two very minute *Leavs*, and joyned to a long *Root* like a *Stalk*. The end of which comes just to the bottom of the *Cover*, ready for its *exit* into the *Ground*.

22. §. In *Goosgrass*, where the *Inner Cover* is also *Cartilaginous* or *Horney*, the *Seed* is postured in much a like manner, and looks just like a couple of poynted *Leavs* with a very long *Stalk*.

23. §. THE *Seed* of *Stramonium*, is also inclosed in a *Bulky Cover*. Which being soaked in warm water, and very warily cut about the edges, with a *Rasor*, the *Seed* may be taken out of it entire. Shaped like that of *Orach*, but much longer. For the *Reception* whereof, the *Cover* is formed with a hollow, which runs round about it near the *Edge*; where in the *Seed* lies like a little winding *Snake*.

CHAP. V.

Of the BUDS of Seeds. And of the PARTS, of which these, the Radicle, and Lobes are compounded.



FROM between the two *Lobes*, rises up the *Stalk* of the *Plant*. The original whereof, either to the naked *Eye*, or by a good *Glass*, is always visible in the *Seed*.

2. §. In many *Plants*, *Nature* sees fit only to lay the foundation hereof in a small round *Node*; where upon the *Leavs*, in the *Vegetation* of the *Seed*, are superstructed: as in *Viola Lunaria*, and others.

3. §. But in the greater number of *Seeds*, is formed a true *Bud*, consisting of perfect *Leavs*; different from those, which grow upon the *Stalk*, only in *Bigness*; and so far in *Shape*, as the same *Parts* of an *Animal Fetus*, in its several ages in the *Womb*. In many *Seeds*, as well
small

small as great, and as well of *Herbs* as *Trees*, it is very apparent. But oftentimes lyeth so deep between the *Lobes* as to be almost undiscernable, as in *Maple*.

4. §. The *Leaves* of the *Bud*, in different *Plants*, are of a different *Number*; in some, Two; in others, Four, Six, and sometimes more. In the *Bay-Berry*, they are only two; very small, but thick or fat, and finely veined. In the *Seed* of *Carduus Benedictus*, they are also *Tab. 78.* Two; almost invisible; broad at the *Bottom*, poynted at the *Top*, thick or fat, yet plated inward, and postured a little distant one from the other; for the two next to rise up between them. The like may be seen in *Carthamum*; and so, I suppose, in all the *Carduus Kind*.

5. §. In some *Herbs*, although the *Bud* consisteth but of two perfect *Leaves*, yet they are very conspicuous. Not only in larger *Seeds*, *Tab. 78.* as in the *Phaseolus* or *French Bean*; but in those which are small, as in the *Seed* of *Hemp*. In this, the two *Leaves* are both plated, and so set *Edge to Edge*, with mutual *Undulations*. Of that *Length*, as to be extended beyond a third part of the *Lobes*.

6. §. In the *Seed* of *Sena*, the *Bud* consisteth of Four *Leaves*; of *Tab. 78.* which, the greater pair is the outer, and guards the less. Shaped not much unlike those in the *Seed* of *Carduus*; but are a little more visible.

7. §. In the *Bud* of an *Almond*, we may easily count six, or eight *Leaves*, and sometimes more; the Innermost being laid bare by a dexterous *Tab. 78.* Separation of the Outer. These are by much the greatest, doubled Inward, and so laped one over another; whereby they embosome all the rest, as a *Hen* spreads her *Wings* over her *Chickens*. The like is observable in many other large *Kernels*, as also in the *Garden Bean*, and some other *Plants*. With respect to which, I have taken leave (a) to call this *Part* the *Plume*. (a) B. t. Ch. i.

8. §. THE LOBES of the *Seed*, and so likewise the *Radicle* and *Bud* consist of a *Skin*, *Parenchyma*, and *Branched Vessels*: all which I have formerly described. (b) I shall now add the following *Re-* (b) B. t. *marques*. Ch. i.

9. §. And first of the *Skin*, which in some *Seeds*, as the *French-Bean* may easily be separated from the *Parenchyma*: especially if the *Bean* be soaked in water for some days; for then it will slip off, like the *Skin* in any part of ones *Body* where it is blistered. 'Tis woven into *Bladders*, as the *Parenchyma*; but into smaller ones, and upon the *Lobes* of a *Garden Bean*, all radiated towards the *Center*. With *Tab. 79.* these *Bladders*, there are also mixed a sort of *Lignous Fibres*, incomparably small, which give a *Toughness* to the *Skin*, and by which the *Bladders* are directed into *Rays*.

10. §. The *Bladders* of the *Parenchyma*, as is said, are much larger than those of the *Skin*, especially in the *Lobes*. In those of a *Garden Bean*, somewhat oval, about $\frac{1}{4}$ of an *Inch* *Diameter* by their *Breadth*, *Tab. 79.* and directed towards the *Branches* of the *Seminal Root*. In the *Radicle*, they are twenty times smaller, than in the *Lobes*: and so in the *Plume*.

11. §. Throughout the *Parenchyma* run the *Branched Vessels*, which *Tab. 79.* in the *Lobes* make the *Seminal Root*; in the *Radicle* and *Plume*, the *Wood* of the *Root* and *stalk*: In all of them, distributed as hath been (c) formerly shewed. (c) B. t. Ch. i.

12. §. I shall here further note, That the utmost divisions are no where extended to the Circumference of the *Lobes*, but are all inosculated together at a considerable distance from it, as in the *Leaves* of some *Plants*.

Tab. 79.

13. §. In the *Lobes* they all meet in one solid *Nerve*. But in the *Radicle*, are dilated into a hollow *Trunk*, filled up with a *Pith*; composed of *Bladders* somewhat bigger than those which make, as it were, the *Barque* of the *Radicle*. In the *Radicle* of a *French Bean*, the *Pith* is very conspicuous.

Tab. 79.

14. §. The *Vessels* are of two kinds, as in the other *Parts* of a *Plant*; for *Sap*, and for *Aer*. Not running collateral, as *Arteries* and *Veins*; but the latter every where sheathed in the former. From the *Aer-Vessels* it is, that if a *Bean* be steeped in water, and then the *Radicle* cut transversly and pressed, it will yield *Bubbles* as well as *Liquor*. These *Vessels* are admirably small, yet through a very good *Glass* become visible.

Tab. 79.

15. §. The *Liquor* contained in the *Seed*, when full ripe is chiefly *Oyl*; generally, found in a greater proportion here, than in any other part of a *Plant*. Being as the *Pickle*, in which the *Seminal Virtues*, *i. e.* the more *volatile* and *active Principles* of the *Seed*, are immersed for their *Preservation*: and to curb them from too great a *Luxuriance* in the *Vegetation* of the *Seed*.

C H A P. VI.

Of the GENERATION of the SEED.



S I made choice of a *Garden-Bean*, to shew the manner of the *Vegetation* of the *Seed*: so I shall take an *Aprecock*, as very apt and convenient, to observe and represent the *Method* which *Nature* taketh in its *Generation*.

2. §. In order to this, the first thing that is to be done, is to make a fit *Uterus*. Both to keep the *Membranes* of the *Fœtus* warm, and succulent, till it be formed: and to preserve and secure the *Fœtus* it self afterwards, till it comes to be born into the *Ground*.

3. §. For this purpose, the *Pulp* and *Stone* of the *Fruit* are both necessary; but primarily the *Stone*: the *Meat* or *Pulp* being no otherwise necessary, but because the *Stone* cannot be made without it; the petrifying of that *Parenchyma* which is the *Ground* of the *Stone*, being effected, by the sinking of the *Tartar* from the *Pulp* thereinto.

4. §. And that, at the first, the *Ground* of the *Stone*, is a distinct, but soft *Parenchyma*; is evident in the cutting of a young *Aprecock*. Of which, also a slice cut off, with a *Rasor*, and viewed through a good *Glass*, sheweth it to be composed of *Bladders*, as the *Pulp* it self. Only, whereas many of those of the *Pulp* are large, now about as big as a white *Pepper-Corn*: these are no bigger than a *Mustard-Seed*. But as the *Parenchyma* hardens into a *Stone*, these *Bladders* are all gradually filled up, and disappear. Tab. 82.

5. §. This *Parenchyma* is derived immediately from the *Pith*, as the *Pulp* is from the *Barque*; and makes the far greater part of the *Stone*. 'Tis covered all over within, with a very thin *Lining*; derived, not from the *Pith* but the *Parenchyma* which covers the *Seed-Branch*, upon its first entrance within the hollow of the *Stone*. This *Lining* is of a close substance; yet composed of *Bladders*, exquisitely small and hardly visible. By which means, it soon becomes a very hard and dry *Body*; and is hereby fitted, both to promote the induration of the rest of the *Stone*; and the seasonable drying, and so, the shrinking up, of the *Covers* of the *Seed*, to make room for its *Growth*. Tab. 83.

6. §. The *Stone* being made hard and dry; it could never be so sufficiently softened by lying under ground, but that, it would keep the *Seed* a perpetual prisoner, unless it were also made pretty easily to cleave in two. For which purpose, the *Skin* of the *Fruit* doth observably conduce. For in a *Slice* of a young *Aprecock* cut transversly with a very sharp knife, it may be seen, especially with the help of a *Glass*, to be doubled inward from the two *Lips* of the *Fruit*, and so to be continued,

tinued, not only through the *Pulp*, but also through the *Stone* it self, into the hollow of the same, where it meets, and is united with the *Lining* thereof. Whereby, as it further helps to the drying and hardning of the *Stone*; so also renders it cleavable in that part, where it runs through it. And therefore, whereas towards the *Stalk*, it goes no farther than to the *Seed-Branch*, and so but half way through the *Stone*: towards the *Top* of the *Fruit*, where the *Radicle* stands, and where the *Stone* begins to cleave, it runs quite through it.

7. §. *Nature* having thus provided a convenient *Uterus*, She next taketh care about the *Membranes* of the *Fœtus*. These are *Three* apparently distinct, and in many respects different one from another.

Tab. 80. 8. §. The outer *Membrane* is derived from the *Parenchyma* which surrounds the *Seed-Branch*; which, upon its entry into the hollow of the *Stone*, is expanded, as it were, into two *Bladders*, one within another; whereof, one becomes the *Lining* of the *Stone*; the other, this outer *Membrane*: as is best seen by cutting a young *Aprcock*, when it is about half an Inch long, down through the middle, or from the *Seat* of the *Flower* to the *Stalk*, between the two *Lips*.

Tab. 80. 9. §. This outer *Membrane*, at this age, hath a good full and firm *Body*, about $\frac{1}{2}$ th of an Inch thick, or through an ordinary *Glass*, half an Inch, where it is thickest, as at the *Sides* and the greater end: the *Poynt* being thinner, for the more easy eruption of the *Radicle* into the *Earth*. Composed of *Bladders*, through an ordinary *Glass*, about as big, as a *Colewort-Seed*.

Tab. 80. 10. §. Throughout this *Membrane*, the *Vessels* contained in the *Seed-Branch* are distributed. Beginning a little below the smaller end of the *Coat* or *Membrane*, they thence fetch their circuit both ways round about, just beneath the *Surface* of the *Membrane*, and at last, meet in the middle of the greater end, where they are all inosculated, so as to make a kind of *umbilical Node*. From whence they strike deeper into it, and at last, into the middle *Membrane*, in which they presently become invisible. By these *Vessels*, the *Sap* is brought and spewed into the middle *Membrane*. So that the outer *Membrane* seemeth, in some respects, to be answerable to the *Placenta* in *Animals*.

Tab. 80. 11. §. The middle *Membrane*, is derived from the bottome of the Outer. From whence especially, but also round about, the *Bladders* hereof (all angular) are more and more amplified towards the *Centre*; most of them being at least two hundred times bigger, than those of the Outer *Membrane*: whereby it looks, through a *Glass*, not unlike a *Cooe* full of *Hony*; or in regard of their great transparency, like a company of little *Crystal Pans* full of a pure *Lympha*.

12. §. This Middle *Membrane*, is properly so called, from the state and condition it hath, upon the *Augmentation* of the *Seed*, at which time, it obtains the nature of an *Involucrum*. But originally, it is every where entire, without any *Hollow*, filling up the *Cavity* of the Outer *Membrane*, like a soft and delicate *Pulp*. After a short time, there

there appears in it a small *Ductus* or *Chanel*; which runs from the bottom to the top, like an *Axis*, through the middle of it. At first, *Tab. 81.* no wider than to receive the *Hair* of a *Mans Head*; not visible, except in a slice hereof cut transversly, and viewed in a *Glass*. Being grown a little wider, it may be seen, if the *Membrane* be dexterously cut by the length. At which time, it is also dilated into two *Oval Cavities*, one at each end: which are as two little *Cisterns*, whereinto a most pure *Lympha* continually owzeth, and is therein reserved for the nourishment of the *seed*; and through the *Chanel* which runs between the *Cisterns* is emptied out of one *Cistern* into another, according as the *seed* or the *Inmost Membrane* hath need of it; *i. e.* as the *Weather* and other *Circumstances* do more or less accelerate their *Growth*, and so render the *Lympha* useful to them.

13. §. A few days after this, the *Innermost Membrane* begins to appear; growing, like a soft *Node* or *Bud*, out of the upper *Cistern*; to the lower end of which it is joyned by a short and tender *Stalk*, from *Tab. 81.* whence it is produced into a *Conick-oval Figure*, answerable to that of the *Cistern*.

14. §. This *Membrane*, though soft and full of *Sap*, yet being compared with the *midlemost*, is a close and compact *Body*, composed of *Bladders* above 300 times smaller than they are in that. Whereby, as the *Seed* is so well guarded, as not to be supplied with any part of the *Lympha*, but the purest: so neither with any more of this, than will suffice, without the danger of making an *Inundation* out of so great a *Lake*.

15. §. This *Membrane*, if it be pulled with a most steady hand, and very gently, upwards, it will draw a small transparent *String* after it to the bottom of the *Midle Membrane*: The said *String* though for the greater part, *Parenchymous*, yet being strengthened with the admixture of some *Lignous Fibres*; no otherwise visible in either of these two *Membranes*. So that they seem, to be a small portion of those which are inosculated at the bottom of the *Outer Membrane*, and thence produced through the *midlemost*, underneath the *Chanel*, till at last they break forth into the upper *Cistern*, where they form this *Inner Membrane*: a piece of close-wrought *Work*, suitable to the incomparable fineness of all the *stuff* out of which it is made.

16. §. The same *Membrane* is originally entire, as the *Midlemost*: but being grown to about the bigness of a *Carvi-Seed*, becomes a little hollow near the *Cone*. And the *Lignous Fibers* above said, fetching their compass from the *Base*, shoot forth into the *Cone*; and so make a very small *Node* therein, for the first *Essay* towards the *Generation* of the *Seed*. The said *Fibers* being thus spun out, to the utmost degree of fineness for this purpose. *Tab. 81.*

17. §. This *Node*, being grown about $\frac{1}{3}$ th part as big as a *Cheese-Mite*; it begins next to be divided by a little indenture at the *Top*. *Tab. 81.* Which growing by degrees still deeper, the *Node* is hereby at length distinguished into two *Lobes* or thick *Leavs*.

18. §.

18. §. So soon as these are finished, their *Basis* begins afterwards to be contracted, and so to be formed into a *Radicle* or that part of the *Seed* which becomes the *Root*. As the *Stalks* of *Fruits* do grow lesser, while the *Fruits* themselves are expanded. So that in this estate, the *Radicle* is, as it were, the *Stalk* of the *Seed*.

19. §. At this time, the *Seed* being extream small, the *Lobes* are not so manageable as to be separated one from the other. But it is most reasonable to suppose that so soon as the *Radicle* is finished, the next step, is the pushing forth of another *Node*, between the *Lobes*, in order to the making of a *Bud*, and so the perfection of the *Seed*.

20. §. This being done or in doing, the *Radicle* or *Stalk* of the *Seed*, contracting still more and more at the bottome, hangs at the *Inner Membrane*, only by an extream small and short *Ligament* or *Navel-String*. Which at last, also breaks; and so the *Seed*, as *Fruits* when they are ripe, falls off and lies loose in the *Iner Membrane*; this gradually shrinking up and so becoming more hollow, to make room for the further *Growth* of the *Seed*.

Several
LECTURES
Read before the
ROYAL SOCIETY.

By *NEHEMIAH GREW* M.D. Fellow
of the *ROYAL SOCIETY*, and of the
COLLEGE of PHYSICIANS.

LONDON,

Printed by *W. Rawlins*, 1682.

L 1

THE
TITLES
Of the following
LECTURES.

- I. **O** *F the Nature, Causes, and Power of MIXTURE.* The second Edition.
- II. *Of the LUCTATION arising upon the Mixture of several Menstruum's with all sorts of Bodies.* The second Edition.
- III. *An Essay, Of the various Proportions, wherein LIXIVIAL SALTS are found in Plants.*
- IV. *Of the ESSENTIAL and MARINE SALTS of Plants.*
- V. *Of the COLOURS of Plants.*
- VI. *Of the Diversities and Causes of TASTS ; chiefly in Plants. With an Appendix, Of the ODOURS of Plants.*
- VII. *Experiments in Consort, upon the SOLUTION of SALTS in Water.*
-

TO THE
Right Honourable
WILLIAM
Lord Vi-Count *BROUNCKER*,
PRESIDENT
OF THE
Royal Society.

MY LORD,



ONE Reason why I Dedicate the following Discourses to Your Lordship, is, For that by Your great und undeserved Respects, You have obliged me to do no less.

Another, my Lord, is, Because I could not but Publickly return Your Lordship Thanks, for minding the Royal Society of so good a Way, they are lately resolved upon, for the Management of a great part of their Business. Wherein, my Lord, I do more than presume, that I also speak the Sense of the whole Society; I think, not any one excepted.

I may with the same Confidence intimate, my Lord, how happy they account themselves, in having a Person so fit to preside their Affairs, as Your Lordship. The Largeness of your Knowledge, the Exactness of Your Judgment, the Evenness of Your Comport; being some of those necessary Qualifications, which His Majesty had in His Eye (as right well understanding what He did) when He fixed His Choice upon Your Lordship.

I know, my Lord, that there are some men, who have

just so much Understanding, as only to teach them how to be Ambitious: The Flattering of whom, is somewhat like the Tickling of Children, till they fall a Dancing. But I also know, that Your Lordship unconcerneth Your self as much, in what I even now spake; as Cæsar did himself, when his Souldiers began to style him King. For as he said, Non Rex, sed Cæsar: So let Your Lordship be but once nam'd, and all that follows, is but a Tautology to what You are already known to be. Your being President of the Royal Society, Your being the First that was Chosen, and Chosen by so Knowing a Prince; becomēs so real a Panegyrick to Your Lordship, as leaveth Verbal ones without any sound.

Whence, my Lord, I have a third Reason most naturally emergent, which is, That I dare to submit myself, as to what I have hereafter said, to Your Lordships Censure. You being so able and just an Arbitrator betwixt the same and all those Persons therein concern'd; that You can neither be deceived, nor corrupted, to make a Judgment in any Point, to the Injury of either.

And truly, my Lord, were it only from a Principle of self-Interest, yet I could not desire it should be otherwise. For the World, if it lives, will certainly grow as much more knowing than it is; as it is now more, than it was heretofore. So that we have as little Reason to despise Antiquity; as we can have willingness, that we our selves should be despised by Posterity.

Yet some difference there is to be made; viz. betwixt those of all Ages, who have been modestly ignorant; and those who have thought, or pretended, that they were Omniscient. Or if knowing and acknowledging that they were Ignorant; have yet not been contented to be so; unless, with as good manners, as sense, they did conjure all Mankind not to offer at the knowing any more than themselves.

Upon the whole, my Lord, I desire not You should be a Patron, any further than You are a Judge. For if this small Essay hath deserved the least acceptance, I am sure, that in being one, You will be both. I am,

My Lord,
Your Lordships most Faithful
and Obedient Servant,

NEHEMIAH GREW.

A
 DISCOURSE
 Read before the
 ROYAL SOCIETY
 Decemb. 10. 1674.
 Concerning the
 NATURE, CAUSES, and POWER
 OF
 MIXTURE.



HAVING the honour to perform the Task of this day; I shall endeavour to conform to the *Philosophy*, which this *Society* doth profess; which is, *Reasoning grounded upon Experiment, and the Common Notions of Sense*. The former being, without the latter, too subtle and intangible; the latter without the former, too gross and unmanageable: but both together, bearing a true analogy to our selves; who are neither

Angels, nor meer Animals, but Men.

The Subject I have chosen to speak of, is *Mixture*. Whereof, that our *Discourse* may be the more *consistent*, and the better *intelligible*; all I have to say, shall be ranged into this Method; *viz.*

1. First, I shall give a brief account of the received *Doctrine* of *Mixture*.

2. Next, lay down some *Propositions* of the *Principles* whereof all *Mixed Bodies* consist.

3. Then, open the true *Nature* of *Mixture*; or say, *What* it is.

4. And then enumerate the *Causes* of *Mixture*; or say, *How* it is made.

5. Lastly, I shall shew the *Power* of *Mixture*; or, *What* it can do.

CHAP.

C H A P. I.

Of the received Doctrine of Mixture.



I R S T, As to the received *Doctrine of Mixture*; not to trouble you with tedious quotations of what *Aristotle*, *Galen*, *Fernelius*, *Scaliger*, *Sennertus*, *Riverius*, and other Learned men say hereof; we may suppose the whole summed up in that *Definition* which *Aristotle* himself hath given of it, and which the greater number of his Followers, have almost religiously adhered to; viz. that 'tis, τῶν μικτῶν ἀλλοιωθέντων ἕνωσις. that 'tis, *Miscibilium alteratorum unio*. Which *Definition*, as it is usually explained, is both *Unintelligible*, and *Unuseful*.

Lib. 1. de *Miscibilium alteratorum unio*. Which *Definition*, as it is usually explained, is both *Unintelligible*, and *Unuseful*.
 Generat.
 & Corrupt.
 Cap. ult.

2. §. Two things are *unintelligible*; what they mean by *Alteration*; and what by *Union*. In this *Alteration*, they say, That the very *Forms* of the *Elements* are altered. And therefore lay it down for an *Axiom*, *Quod in Mixto, Formæ Elementares tantum sint in potentia*, But let us see the consequence. For if in a *mixed* body, the *Forms* of the *Elements* are but in *potentia*; then the *Elements* themselves are but in *potentia*: for we all say, *Forma dat esse*. And if the *Compounding* *Elements*, are only in *potentia*; then the *Compounded* *Body* it self can be only in *potentia*: yet to say it is no more, is most absurd.

3. §. As for the *Union* of *Elements* in a *mixed* *Body*; they make it such, as brings them at last to assert, the *Penetration* of *Bodies*, and that the *Union* of *mixed* *Bodies* is nothing else. For they say it is made in such sort, that every particle of the *mixed* *Body*, partaketh of the *Nature* of the whole. Which *Nature*, ariseth from the contempered *Qualities* of the four *Elements*. Whence they conclude, That every particle of the *mixed* *Body*, containeth in it self all the four *Elements*. Which is plainly to assert a *penetration* of *Bodies*. For every *Element* is, at least, one particle; if therefore every particle of the *mixed* *Body*, containeth four *Elements*; then four particles are but one. I conclude then, That the received *Doctrine of Mixture* is *Unintelligible*.

4. §. Whence it follows, That it is also *Barren* and *Unuseful*. For who can make any use of that which he understandeth not? And the experience of so many years, wherein it hath been ventilated by the disputes of men, proveth as much: Scarce any of them, except the Learned *Sennertus*, daring to venture upon Experiment, for fear they should come to understand themselves.

5. §. It is confessed, that many gallant things have been found out by artificial *Mixture*. But no thanks to this *Definition* of it. For as an *Ignorant* Man may make bad *Work*, and a good *Rule* be never the worse; so one that is *Ingenious* may make good *Work*, and a bad *Rule* be never the better. The question is not, what have men done? but what have they done upon this foundation, *Quod Mixitio sit miscibilium alteratorum unio*. Had this ever taught them to do any thing, even so much as to make the *Inke* wherewith they have wrote, all their *Disputes*; I confess, they would have had something to shew for it. But the truth is; their *notions* of *Mixture*, have been so far from doing us any good, that they have done us much harm: being, through their seeming subtlety, but real absurdity, as so many phantastick *spectrums*, serving only to affright men from coming near them, or the Subject whereof they treat.

6. §.

§. 6. I shall therefore endeavour to open the true Nature of Mixture. And I shall build my Doctrine upon the Common Notions of sense: which none can deny; and every one may conceive of. In order to which, I shall take leave to lay down some Propositions, of the Principles of all mixed Bodies.

C H A P. II.

Of the Principles of Bodies.



AND first, by Principles, I mean Atomes, or certain Sorts of Atomes, or of the simplest of Bodies. For otherwise they would not be Principles; for a compounded Principle, in strict speaking, is a Contradiction. Even as Fives, Threes, or Two's are not the Principles of Number, but Unites.

2. §. Whence, secondly; it follows, that they are also Indivisible. Not Mathematically; for the Atomes of every Principle have their Dimensions. But Physically; and so; what is but one; cannot be made two. If it be asked, Whether a Stick cut with a Knife, be not of one, made two? I say, that a Stick, is not one Body, but many millions of Bodies; that is, of Atomes; not any one whereof is divided within it self, but only they are separated one from another, where the Knife forceth its way. As in the drawing of a mans Finger through a Heap of Corn; there is no Division made in any one Grain, but only a separation of them one from another, all remaining still in themselves entire. I say, therefore, that what is Physically one, is also most firm, and Indivisible, that is, Impenetrable: for Penetration is but the Separation, not the Division of Atomes.

3. §. Hence, thirdly, they are also Immutable. For that which cannot be divided, cannot be chang'd. So that of the whole World of Atomes, not any one hath ever suffer'd, or can suffer the least mutation. Hereupon is grounded the Constancy of Causes and Effects. So that, in all Generations, it is not less certain, that the self same Principle is still propagated from the same; than, that Man is from Man. Wherefore, compounded Bodies are generated; but Principles are not; but only propagated; that is, in every Generation, they pass, in themselves unaltered, from one Body, into another.

4. §. If Principles, or Atomes are all Immutable; it again follows; That they are of Divers Kinds. For one and the same Principle, or Kind of Atomes, will still make the Same Thing, and have the same Effect: so that all Generations would then be the Same. Wherefore, since they are Immutable, they must be Divers.

5. §. This Diversity, for the same reason, is not small, but very Numerous. For as the World, taken together, is Natures Shop; so the Principles of Things are her Tools, and her Materials. Wherefore, as it speaks the goodness of a Shop; so the Perfection of the Universe, That it is furnished with many Tools wherewith, and many Materials whereupon to work. And consequently, that Philosophy beareth best its own name; which doth not strain all to two or three Principles, like two or three
Bells

Bells in a Steeple, making a pitiful *Chime*: but tryeth to rise up to Nature's own *Number*, and so to ring all the *Changes* in the World.

6. §. Yet doth not this vast *Diversity* take away the *Regiment* and *Subordination* of *Principles*. There being a certain lesser number of them, which either by their greater *quantity*, or other ways, have *Rule* and *Dominion*, in their several *Orders*, over all the rest. For where-ever the *Subject* is *Multitude*, *Order* is part of its *Perfection*. For *Order* is *Proportion*. And how can *Nature* be imagin'd to hold *Proportion* in all things else, and not here? Wherefore, as certainly, as *Order* and *Government* are in all the Parts of the *Rational*; so certainly, of the *Material World*. Whence it is, That although the *species* of *Principles* be very *numerous*; yet the *Principles* called *Galenical*, *Chymical*, or any others, which do any way fall under the notice of *Sense*, are notwithstanding *reduceable* to a *smaller number*: viz. according to the number of *Predominant Principles* in *Nature*; or, rather in this part of the *Universe* which is *near and round about us*. To the *Power* and *Empire* whereof, all other *Principles* do submit. Which *Submission*, is not the *quitting* of their own *Nature*; but only their appearance under the external *Face* or *Habit* of the said *Predominant Principles*.

7. §. As there can be no *Order* of *Principles*, without *Diversity*; so no *Diversity*, but what is *originally* made by these two ways; sc. by *Size* and *Figure*. By these they may be exceeding different: and all other *Properties* besides, whereby they differ, must be *dependent* upon these *Two*.

8. §. Nor therefore, can they be of any other *Figures*, than what are *Regular*. For *Regularity*, is a *Similitude continu'd*. Since therefore all kinds of *Atomes* are *divers* only by their *Size* and *Figure*; if the selfe same *Size* and *Figure* were not *common* to a certain number of *Atomes*, they could not be said to be of any *one kind*: and consequently, if there were no *Similitude* of *Atomes*, there could be no *Distinction* of *Principles*.

9. §. Hence also, these two *Modes* of *Atomes*, viz. their *Size* and *Figure*, are the true, and only *original Qualities* of *Atomes*. That is, an *Atome* is *such* or *such*, because it is of such a certain *Size* and *Figure*.

10. §. Lastly, As these two *Modes*, taken severally, are the *Qualities* of an *Atome*: so consider'd together, they are its *Form*. A *Substantial Form* of a *Body*, being an unintelligible thing. I say of a *Body*; for although the *Rational Soul* be a *substantial Form*, yet is it the *Form* of a *Man*, and not of a *Body*. For the *Form* of a *Body*, we can conceive of no otherwise, than as of the *Modification* of a *Body*, or a *Complexion* of all the *Modes* of a *Body*. Which also agrees with that *Definition* of a *Form*, which amongst the *Peripatetick Philosophers* is well enough accepted, viz. *Quod sit, Ratio ejus Essentia, quæ cuique Rei competit*. Which *Ratio*, if it be referred to a *Body*, what is it, but the *Modification* of that *Body*? Having thus proposed a Summary of my *Thoughts* about *Principles*; I shall next proceed to shew what their *Mixture* is.

C H A P. III.

Of the NATURE of Mixture.



AND first of all, from the *Premisses*, we arrive at this *Conclusion*; *sc.* That the *Formation* and *Transformation* of all *Bodies*, can be nothing else, but the *Mixture* of *Bodies*. For all *Principles* are *immutable*; as we have above proved: and therefore not *generable*, *formable*, or *transformable*. And the *Forms* of *Principles*, being but their *Modes*, are also *immutable*. So that the whole *Business* of the *Material World*, is nothing else, but *Mixture*.

2. §. Again, as *Nature* worketh every where only by *Mixture*; so is this *Mixture* every where but *one thing*, and can be but *one*. For whether it be the *Mixture* of *great Bodies*, or of *small*; of *Compounds*, or of *Atomes*; it is every where *Mixture*, and the *Mixture* of *Bodies*. Wherefore, *Mixture* is either an *intelligible Affection* of all *Bodies*, or of *none*; which later, no man will say. As many ways therefore, as we can see, or conceive the *Mixture* of any *gross Bodies*, which we hold in our hand; so many ways, we may, of the *subtilest Mixtures* which *Nature* maketh, or of *Atomes* themselves; and no other ways.

3. §. Now all the ways we can distinguish *Mixture* by, are, in general, these *Two*; either in respect of the *Bodies Mixed*, or else of the *Modes* of the *Mixture* it self.

4. §. In respect of the *Bodies Mixed*, *Mixture* is distinguished also *two* ways; *viz.* by *Conjugation*, and by *Proportion*.

5. §. By *Conjugation*, I mean, a *Mixture* of *some certain Principles*, and not of others. Which is *threefold*. *First*, As to *Number*: as when one *Body* may be compounded of *two Principles*, another of *three*, a third of *four*, a fourth of *five*, and so on. *Secondly*, As to *Kind*: where, though there be a conjunction of the same *Number*, yet not of the same *Kind*. *Thirdly*, When they differ from one another both in *Number* and *Kind*. So many ways the *Principles* of *Bodies* may be conceived to be *Conjugated*; and therefore are: for here, that which may be, is. The *Consequence* is clear. For *first*, *Nature* hath various *Materials* wherewith to make these *Mixtures*; as we have shewed. *Secondly*, By these *Mixtures* she may, and without the concurrence of any imaginary *Forms*, must produce all the varieties in the *material World*; as likewise hath been said. Wherefore, since all imaginable *Mixtures* may be made, and that to *some purpose*; if they should not be so, *Nature* would be *Imperfect*: because we our selves can think, how she might put her *Materials* to further use, then so she would do. To think therefore, that all *Kinds* of *Principles*, or all *Elements* go to make up every *Compounded Body*, as by the *Peripatetick Philosophy* we are taught; is a conceit, no more to be credited, than one that should tell us, all *Kind* of *Wheels* and other

parts of a *Watch*, were put into a *Clock*; or that there were no other *Materials* wherewith to build an *House*, then for a *Tent* or a *Ship*. For why should *Nature*, the great *Artificer* by which all perfect *Works* are made, be feigned to cram and ram all things into one, which we our selves look upon as absurd?

6. §. Secondly, The *Mixture* of *Principles* is diversifi'd, as by *Conjugation*, so also by *Proportion*. That is, by the divers *Quantities*, of the several *Principles* or *Parts* mixed together. As if the *Quantity* of one, were as five to ten; of a second, as five to fifteen; of a third, as five to twenty, &c. Or if that of one, be as five to six; of a second, as six to seven; of a third, as seven to eight. By which, and by other *Proportions*, *Mixture* may be varied innumerable ways.

7. §. Again, As *Mixture* is varied with respect to the *Bodies Mixed*; so likewise in respect of the *Mixture* it self, which I call the *Location* of *Principles*, or the *Modes* of their *Conjunction*. Which may be various, as well as their *Conjugation* and *Proportion*. Yet are they all reduceable unto two general *Modes*: all *Bodies*, and therefore all *Principles*, being mixed either by *Mediation*, or by *Contact*.

8. §. Now all *Contact*, whether of *Compounds*, or of *Atomes*, can be no other way, than such as is answerable to their *Figures*. Whereof, therefore, we can conceive but three general ways, viz.

First, By *Contact* in a *Point*, or some smaller part: as when two *Atomes* meet, which are globular or otherwise gibbose. Secondly, By *Contact* in a *Plain*: as in the conjunction of the sides of *Triangular* or *Quadrangular Atomes*, or otherwise flat. Thirdly, By *Contact* in a *Concave*: as when one *Atome* is admitted into the *Concave* or *hole* of another; as a *Spigot* is into a *Fosset*. The first may be called, *Apposition*; the second, *Application*; the third, *Reception* or *Intrusion*.

9. §. In the two last ways, *Atomes* may be joyned by *Mediation*; but best of all the last. As when the two extrems of one *Atome* are received into the *Concaves* or the *holes* of two others.

10. §. And these are all the general ways, whereby we can conceive *Bodies* to be *Mixed* together; sc. by their various *Conjugation*, *Proportion* and *Location*. So that the *Composition* of *Atomes*, in *Bodies*; is like that of *Letters*, in *Words*. What a *Thunderclap* would such a *Word* be, wherein all the four and twenty *Letters* were pack'd up? One therefore is compounded of more, another of fewer: this of some, and that of others: and both the *Conjugation*, *Proportion*, and *Location* of *Letters* is varied in every *Word*: whereby, we have many thousands of differing *Words*, without any alteration at all, in the *Letters* themselves; and might have ten times as many more. In like manner, therefore, or in the self same analogous way, as the *Letters* of the *Alphabet*, are the *Principles* of *Words*; so *Principles*, are the *Alphabet* of *Things*.

11. §. What we have said of *Principles*; and of *Mixture* as consequent thereupon; may be a *foundation* for an intelligible account, of the *Nature* and *Cause* of most of the *Intrinsic Properties*, and *Qualities* of *Bodies*: as of *Gravity*, *Levity*, *Fixity*, *Fluidity*, *Angularity*, *Roundness*, *Heat*, *Cold*, *Blackness*, *Whiteness*, *Soverness*, *Sweetness*, *Fragrancy*, *Fetidness*, and very many more. I say an intelligible account; sc. such as is grounded upon the *Notions* of *Sense*, and made out *Mechanically*. But the exemplification hercof, being too large a field for

for this, or any one *Lecture*, I shall, before I come to the *Causes* of *Mixture*, only deduce from the *Premises*, these following *Corollaries*.

12. §. *First*, That there is no *alteration* of *Principles* or of *Elements*, in the most *perfect Mixture* of *Bodies*. It *cannot* be; for *Principles* are *Immutable*, as we have said. And if it could be, yet it *needeth not* to be: for they are also *many*, and *compoundable infinite* ways; as hath been shewed. So that we have no need to perplex our selves with any of those difficulties, that arise from the *Doctrine* of the *Alteration* of *Elements*. The ground of which conceit, is that, of three being but four *Elements*, and all in every particle of the *mixed Body*. And so men being puzled, how from thence to make out the infinite *variety* of *Bodies*, they feigned them to be alterable, and *altered*, upon every *perfect Mixture*. Not considering, that if their four *Elements* be *alterable*; as few as they are, no fewer than *three* of them may be spared: for *one Element*, if *alterable*, may be made *any*.

13. §. Hence, *Secondly*, may be solved that great *Dispute*, Whether such as we call *Lixivial Salts*, are made by the *fire*? For *first*, No *Principle* is made by the *fire*: all *Principles* being *unalterable*; and therefore *unmakable*. *Secondly*, We must therefore distinguish betwixt the *Principle*, and its various *Mixture* with other *Principles*; from whence it may receive different *Shapes* and *Names*. Wherefore, a *Lixivial Salt*, *qua Lixivial*, is certainly made by the *fire*. But *quatenus Salt*, it is not: that *Principle* being *extractable* out of most *Bodies*; and by *divers* other ways, then by the *fire*. For whether you *Calcine* a body, or else *Ferment* it, (after the manner shewed by the *curious Improver* of *Chimical* Knowledge, *Dr. Daniel Cox*) or *putrifie* it under ground, or *drown* it in the *Sea*; it still yieldeth *some kind* of *Salt*. All which *Salts* are made, not by *making* the *saline Principle*; but only by its being differently *Mixed*, by those several ways of the *Solution* of *Bodies*) with other *Principles*: from which its different *Mixture*, it receives the *various Denominations*, of *Marine*, *Nitrous*, *Volatile*, or *Lixivial*.

14. §. Hence, *Thirdly*, the most *perfect Mixture* of *Bodies*, can go no higher than *Contact*. For all *Principles* are *unalterable*; and all *Matter* is *impenetrable*; as hath been said. In the most *visible* and *laxe Mixture*, there is *Contact*; and in the most *subtile* and *perfect*, as in *Generation* it self, there is *nothing more*.

15. §. Hence, *Fourthly*, we easily understand, how *divers* of the same *Principles*, belonging both to *Vegetables* and many other *Bodies*, are also *actually* existent in the *Body* of *Man*. Because even in *Generation* or *Transmutation*, the *Principles* which are translated from one *Body* to another, as from a *Vegetable* to an *Animal*, are not in the least *alter'd* in themselves; but only their *Mixture*, that is, their *Conjugation*, *Proportion* and *Location*, is *varied*.

16. §. Hence also the difference of *Mixture*, arising from the difference of *Contact*, is intelligible; *sc.* as to those *three degrees*, *Congregation*, *Union*, and *Concentration*.

Congregation, and *Inconsistent Mixture*, is when the several *Atomes* touch but in a *Point*, or *smaller part*. In which manner, I have *divers* arguments, inducing me to believe the *Atomes* of all *Fluid Bodies*, *qua Fluid*, do touch; and in *no other*.

Ch. 3. §. 8. *Union*, is when they touch in a *Plain*. As in the *Crystals* and *Shootings* of all *Salts*, and *other* like *Bodies*. For if we pursue their divided and subdivided parts, with our eye, as far as we can; they still terminate, on every side, in *Plains*. Wherefore, 'tis intelligible, That their very *Atomes* do also terminate, and therefore touch, in *Plain*.

Ch. 3. §. 8. *Concentration*, is when two, or more *Atomes* touch by *Reception* and *Intrusion* of one into another: which is the *closest*, and *firmest* *Mixture* of all; as in any *fixed unodorable*, or *untastable* *Body*: the *Atomes* of such *Bodies*, being not able to make any *Smell* or *Taste*, unless they were first *dissolved*; that is to say, *unpin'd* one from another.

17. §. Hence, *Sixthly*, we understand, how in some cases, there seemeth to be a *Penetration* of *Bodies*; and in what *sense* it may be admitted: *viz.* if we will mean no more by *Penetration*, but *Intrusion*. For the *Intrusion* of one *Atome* into the *Concave* or *hole* of another, is a *kind* of *Penetration*; whereby they take up less room in the *mixed* *Body*, then they would do by any other way of *Contact*. As a naked knife and its sheath, take up almost double room, to what they do, when the knife is sheathed. Whence we may assign the *reason*, Why many *Liquors* being *mixed*; take up less room or space, then they did *apart*; as the *Ingenious* Mr. *Hook* hath made it to appear by *Experiment*, that they do. I say the plain *reason* hereof, or at least one reason, is the *Intrusion* of many of their *Atomes* into one another. Which yet is not a *Penetration* of *Bodies* strictly so called.

Ch. 3. §. 1. 18. §. *Seventhly*, If all that *Nature* maketh, be but *Mixture*; and Ch. 3. §. 14. all this *Mixture* be but *Contact* 'tis then evident, That *Natural* and *Artificial* *Mixture*, are the *same*. And all those *seeming subtilties* whereby *Philosophers* have gone about to *distinguish* them; have been but so many *Scarcrows* to affright *Men* from the *Imitation* of *Nature*.

Ch. 3. §. 1. 19. §. *Eighthly*, Hence it follows, That *Art* it self may go far in doing what *Nature* doth. And who can say, how far? For we have nothing to *Make*; but only to *mix* those *Materials*, which are already *made* to our hands. Even *Nature* her self, as hath been said, Maketh nothing *new*; but only *mixeth* all things. So far, therefore, Ch. 3. §. 1. as we can govern *Mixture*, we may do what *Nature* doth.

Ch. 3. §. 18. 20. §. Which that we may still the better understand; let us before, and in the next place, see the *Causes* of *Mixture*. For since *Natural* and *Artificial* *Mixture* are the *same*; the *immediate Causes* of both, are and must be the *same*.

C H A P. IV.

Of the CAUSES of Mixture.



OW all the Causes of Mixture we can conceive of, must, I think, be reduced to these six in general; viz. Congruity, Weight, Compression, Solution, Digestion, and Agitation.

1. §. Congruity, or aptitude and response betwixt the Sizes and Figures of Parts to be mixed: whereby Bodies may be truly called the Instrumental Causes of their own Mixture. As when a Plain answers to a Plain, a Square to a Square, a Convex to a Concave, or a Less to a Greater or an Equal, &c. according to which Respondencies in the parts of Bodies, they are more or less easily mingable.

2. §. Weight, by means whereof all Fluid Bodies, upon supposition of the Congruity of their parts, must unavoidably mingle.

3. §. Compression; which either by the Air, or any other Body, added to Weight, must, in some degree, further Mixture. Because, that Weight it self, is but Pression. For further Proof of all the said Causes, I made this Experiment; Let Oyle of Aniseeds, and Oyl of Vitriol be put apart into the Receiver of an Air-Pump. And, having exhausted it of the Air, let the two said Oyls be then affused one upon the other. Whereupon, First, It is visible, that they here mix and coagulate together; that is, their parts are wedged and intruded one into another, without the usual compression of the Air; for that is exhausted, and therefore only by the Congruity of their receiving and intruding parts; and by their Weight; by which alone they are so compressed, as to make that Intrusion. Secondly, It is also evident, That although they do Coagulate; yet not altogether so much, as when poured together in the same manner, and quantity, in the open Air. Wherefore, Compression, whether made by the Air, or any thing else, as it doth further the Dissolution of some Bodies, so the Mixture of others, and the greater the Compression, the more.


4. §. Solution; For all Bodies mix best, in Forma fluida. And that for two reasons. First, Because the parts of a Body are not then in a state of Union, but of Separation; and therefore, in a more capable state, for their Mixture and Union with the parts of another Body. Secondly, because then they are also in a state of Motion, more or less; and therefore, in a continual tendency towards Mixture; all Mixture being made by Motion. Wherefore all Generations, and most perfect Mixtures in Nature, are made by Fluids; whether Animal, Vegetable, or Mineral. Which is also agreeable to the Doctrine of the Honourable Mr. Boyle, in his Excellent Treatise of the Nature and Vertues of Gems. And it is well known, That Bodies are ordinarily petrified, or Stones made, out of Water. That is, out of petrifying parts dissolved per minima in Water, as both their Menstruum and their Vehicle. Wherefore, if we will talk of making Gold; it must not be by the Philosophers Stone, but by the Philosophers Liquor.

5. §. *Digestion*. For which there is the same reason, as for *Mixture*, by *Solution*. For, *First*, All heat doth *attenuate*, that is, still further *separate* the parts of a Body; and so render them more *mingleable* with the parts of another. And therefore, *Secondly*, Doth also add more *Motion* to them, in order to their *Mixture*.

6. §. *Agitation*. Which I am induced to believe a great and effectual means of *Mixture*, upon divers Considerations. As, *First*, That the *making* of *Blood* in the *Bodies* of *Animals*, and the *mixing* of the *Chyle* therewith, is very much promoted by the same means; *sc.* by the *Agitation* of the parts of the *Blood* and *Chyle*, in their continual *Circulation*. Again, from the *making* of *Butter* out of *Milk*, by the same means: whereby alone is made a *separation* of the oleous parts from the *Whey*, and *Conjunction* of the *Oleous* together. Moreover, From the great *Effects* of *Digestion*; well known to all that are conversant in *Chymical Preparations*. Which *Digestion* it self, is but a kind of *insensible agitation* of the parts of *digested* Bodies. 'Tis also a known *Experiment*, That the readiest way to dissolve *Sugar* in *Wine* or other *Liquor*; is to give the *Vessel* a *hasty turn*, together with a *smart knock*, against any *hard* and *steady* Body: whereby all the parts of the *Sugar* and *Liquor*, are put into a vehement *Agitation*, and so the *Sugar* immediately dissolved, and *mixed* with the *Liquor*. And I remember, that having (with intent, to make Mr. *Matthews's Pill*) put some *Oyl* of *Turpentine* and *Salt* of *Tartar* together in a *Bottle*, and sent it up hither out of the *Country*; I found, that the continual *Agitation* upon the *Road*, for three or four days, had done more towards their *Mixture*; than a far greater time of *Digestion* alone had done before. And it is certain, That a vehement *Agitation*, especially, if continu'd, or joyned with *Digestion*; will accelerate the *Mixture* of some Bodies, ten times more, than any bare *Digestion* alone; as may be proved by many *Experiments*. I will instance in this one. Let some *Oyl* of *Turpentine* and good *Spirit* of *Nitre* be stop'd up together in a *Bottle*, and the *Bottle* held to the *Fire*, till the *Liquors* be a little heated, and begin to bubble. Then having removed it, and the *Bubbles* by degrees increasing more and more; the two *Liquors* will of themselves, at last fall into so impetuous an *Ebullition*, as to make a kind of *Explosion*; sending forth a *smoak* for the space of almost *two yards* high. Whereupon, the parts of both the *Liquors*, being violently *agitated*, they are, in a great portion, *incorporated* into a *thick Balsam* in a moment: and that without any *intense* heat, as may be felt by the *Bottle*. And thus much for the *Causes* of *Mixture*.

CHAP. V.

Of the P O W E R and U S E of Mixture.



HAVING enumerated the *general Causes*, we shall, lastly, enquire into the *Power and Use of Mixture*; or, into what it can *Do and Teach*. And I shall Instance in *six* particulars. *First*, to Render all Bodies *Sociable*, whatsoever they be. *Secondly*, To Make *Artificial Bodies* in Imitation of those of *Natures own* production. *Thirdly*, to make or imitate the *sensible Qualities* of Bodies; as *Smells*, and *Tasts*. *Fourthly*, To make, or imitate their *Faculties*. *Fifthly*, It is a *Key*, to discover the *Nature of Bodies*. *Sixthly*, To discover their *Use*, and the *Manner* of their *Medicinal Operation*.

I N S T A N C E I.

FIRST, To render all Bodies *Sociable* or *Mingleable*: as *Water* with *Oyl*, *Salt* with *Spirit*, and the like. For *Natural* and *Artificial Mixture*, are the same; as we have before proved. If therefore *Nature* can do it, as we see in the *Generation* of Bodies she doth, 'tis likewise in the *Power of Art* to do it. Ch. 3. §. 18.

2. §. And for the doing of it, two *general Rules* result from the *Premisses*, sc. The *Application* of *Causes*, and the *Choice* of *Materials*. As for the *Causes*, they are such as I have now instanc'd in. And for *Ch. 4.* the *Application* of them, I shall give these *two Rules*.

3. §. *First*, That we tread in *Natures* steps as near as we can; not only in the *Application* of such a *Cause*, as may be most proper for such a *Mixture*; but also in allowing it *sufficient time* for its effect. For so we see *Nature* her self, for her more *perfect Mixtures*, usually doth. She maketh not a *Flower*, or an *Apple*, a *Horse*, or a *Man*, in a *moment*; but all things *by degrees*; and for her more *perfect and elaborate Mixtures*, for the most part, she requireth *more time*. Because all such *Mixtures* are made and carri'd on *per minima*; and therefore require a greater time for the compleating of them.

4. §. A *second Rule* is, Not only to make a due *Application* of the *Causes*; but sometimes to *Accumulate* them. By which means, we may not only, *imitate Nature*, but in some cases go beyond her. For as by adding a *Graft* or *Bud* to the *Stock*, we may produce *Fruit sooner*, and sometimes *better*, than *Nature* by the *Stock* alone would do: So here, by *accumulating* the *Causes* of *Mixture*, that is, by joyning *two, three, or more* together; or by *applying more* in some Cases, where *Nature* applyeth *fewer*; we may be able to make, if not a more *perfect*, yet a far more *speedy Mixture*, than *Nature* doth. As by joyning *Compression*, *Heat*, and violent *Agitation*, and so continuing them all together, by some means contrived for the purpose, for the space of a *Week*,

Week, or Month, or longer, without cessation. Which may probably produce, not only *strange*, but *useful Effects*, in the *Solution* of some, and the *Mixture* of other Bodies. And may serve to *mix* such Bodies, as through the *small number* of their *congruous* parts, are hardly *mingleable* any other way. *Agitation* being, as carrying the *Key* to and fro, till it hit the *Lock*; or within the *Lock*, till it hit the *Wards*.

Ch. 3. §. 9. 5. §. Secondly, For the *Choice* of *materials*, if they are not *immediately*, that is, of themselves, *mingleable*; we are then to turn one *Species* of *Mixture* into a *Rule*; which is, To *mix* them by *mediation* of some *third*, whether more *simple* or *compounded* Body, which may be *congruous in part* to them *both*: as *Sulphurous Salts* are to *Water* and *Oyl*; and are for that reason *mingleable* with *either* of them. Or, By any *two* *congruous* Bodies, which are also, *in part*, *congruous* to *two others*: and other like ways. Whereby the *parts* of Bodies, though never so *heterogeneous*, may yet be all *bound* and *lock'd* up together. Even as *twenty Keys* may be *united*, only by *uniting* the *two Rings* whereon they hang.

6. §. The Consideration of these things, have put me upon making several *Experiments*, for the *mingling* of *heterogeneous* Bodies. I shall give two *Examples* of *Trial*; the one upon *Fluid*, the other upon *consistent* Bodies.

7. §. For the *first*, I took *Oyl* of *Aniseeds*, and pouring it upon another Body; I so order'd it, that it was thereby turned into a perfect *milk-white Balsam*, or *Butyr*. By which means the said *Oyl* became *mingleable* with any *Winy*, or *Watery Liquor*; *easily*, and *instantaneously dissolving* therein, in the form of a *Milk*. And note, That this is done, without the *least alteration* of the *Smell*, *Tast*, *Nature*, or *Operation* of the said *Oyl*. By somewhat the like means, not only *Oyl* of *Aniseeds*, but any other *stillatitious Oyl*, may be transformed into a *milk-white Butyr*; and in like manner be *mingled* with *Water* or any other *Liquor*. Which is of *various use* in *Medicine*; and what I find oftentimes very convenient and advantageous to be done.

8. §. Again, not only *Fluid* but *consistent* Bodies, which of themselves will *mix* only with *Oyl*; by due *mixture* with other Bodies, may be render'd *easily* dissoluble in *Water*; as may *Rosin*, and all *resinous* and *friable Gums*. As also *Wax*: and this without changing much of their *Color*, *Tast*, or *Smell*. Whereof likewise, whatsoever others may do, the *Physician* may make a manifold *Use*.

INSTANCE II.

BY *Mixture* also, we may be taught to *Imitate* the *Productions* of *Nature*. As to which, from what we have before said of *Mixture*, we may conclude; That there is no *Generation* of Bodies *unorganical*, but what is in the *Power* of *Mixture* to *imitate*. As of *Animals*, to *Imitate* *Blood*, *Fat*, *Chyle*, *Spittle*, *Flegm*, *Bile*, &c. Of *Vegetables*, to *Imitate* a *Milk*, *Mucilage*, *Rosin*, *Gum*, or *Salt*. Of *Minerals*, to *Imitate* *Vitriol*, *Allom*, and other *Salts*; as also *Metals*, and the like.

2. §. I do not say, I can do all this: yet if, upon good *Premises*, we can conclude this possible to be done; it is one step to the doing of it. But I will also give an *Instance* of somewhat that may be *done* in every *kind*. And,

3. §. *First*, For the *Imitation* of an *Animal Body*, I will instance in *Fat*. Which may be made thus; Take *Oyl Olive*, and pour it upon high *Spirit of Nitre*. Then *digest* them for some days. By degrees, the *Oyl* becomes of the colour of *Marrow*; and at last, is *congealed*, or hardned into a *white Fat* or *Butter*, which *dissolveth* only by the *fire*, as that of *Animals*. In converting *Oyl* thus into *Fat*, it is to be noted, That it *hardens* most upon the *exhalation* of some of the more *Sulphureous* parts of the *Spirit of Nitre*. Which I effected, well enough for my purpose, by unstopping the glass after some time of *digestion*; and so suffering the *Oyl* to *dissolve* and *thicken* divers times by successive *heat* and *cold*. Hence, The true *Congealing Principle*, is a *Spirit of Nitre separated from its Sulphur*. For the better doing whereof, the *Aer* is a most commodious *Menstruum* to the said *Spirit of Nitre*. Whence also, if we could procure such a *Spirit of Nitre*, we might *congeal Water* in the midst of *Summer*. We might also *refrigerate Rooms* herewith *Artificially*. And might *Imitate* all *frosty Meteors*. For the *making of Fat*, is but the *Durable Congelation* of *Oyl*: which may be done without *frost*, as I have shewed how.

Hence also it appears, That *Animal Fat* it self, is but the *Curdling* of the *Oily* parts of the *Blood*; either by some of its own *Saline* parts; or by the *Nitrous* parts of the *Aer* mingled therewith.

Hence likewise it is, That some *Animals*, as *Conies*, and *Fieldfares*, grow *fatter* in *frosty* weather: the *oily* parts of the *blood*, being then more than ordinarily *coagulated* with a greater abundance of *nitrous* parts received from the *Aer* into their *bodies*.

For the same reason it is, That the *Fat* of *Land-Animals* is *hard*; whereas that of *Fishes* is *very soft*, and runs all to *Oyl*, *sc.* Because the *Water*, wherein they live, and which they have instead of *breath*, hath but very few *nitrous* parts in it, in comparison of what the *Aer* hath.

4. §. *Secondly*, For the *Imitation* of a *Vegetable Body*, I will give three *Instances*; In *Rosin*, *Gum*, and a *Lixivial Salt*. The *first* may be made thus; Take good *Oyl of Vitriol*, and drop it upon *Oyl of Anise-seeds*; and they will forthwith *incorporate* together; and by degrees, will *harden* into a *perfect Rosin*; with the *general* and *defining Properties* of a *truly Natural Resinous Gum*. Being not at all *dissoluble* in *Water*; or at least, not any more, then any natural *Rosin* or *Gum*: yet very *easily* by *fire*: as also *highly inflamable*: and exceeding *friable*. Although this *Artificial Rosin*, be the result of *two Liquors*, both which very strongly affect the *Sense*: yet being well *washed* from the *unincorporated parts*, (which is to be done with some care) it hath scarce any *Tast* or *Smell*.

The *Concentration* of these *two Liquors*, is likewise so *universal*; that the *Rosin* is not made by *Precipitation*, but almost a *total Combination* of the said *Liquors*; and that with scarce so much, as any *visible fumes*.

5. §. *Again*, Having taken a certain *Powder* and a *Saline Liquor*, and mixed them together in a bottle, and so *digested* them for some time; the *Powder* was at last transmuted to a *perfect Oily Gum*; which will also *dissolve* either in *Oyl* or in *Water*; in the self same manner, as *Galbanum*, *Ammoniac*, and the like will do.

6. §. And *Lastly*, A *Lixivial Salt* may be *imitated* thus; Take *Nitre*, *Oyl of Vitriol*, and high *spirit of Wine*, of each a like quantity. Of these *three Bodies*, not any *two* being *put together*, that is to say

N n

neither,

neither the *Nitre* with the *Oyl*, nor the *Oyl* with the *Spirit*, nor the *Nitre* with the *Spirit*, will make the least *Ebullition*: yet all three mingled together, make a very conspicuous one. The *Spirit* of *Wine* being as the *Sulphur*; and so that, and the *Nitre* together, standing, as it were, in the stead of an *Alkalizate*, that is, a *Sulphurous Salt*, against the *Oyl* of *Vitriol*. Divers other *Experiments* may be shew'n of the like Nature.

7. §. In the last place, for the *Imitation* of a *Mineral Body*, I will instance in two, *sc.* *Nitre* and *Marine Salt*; if I may have leave to reckon them amongst *Mineral Bodies*. As for *Nitre*, by mixing of four *Liquors* together, and then setting them to shoot; I have obtained *Chrystals* of true and perfect *Salt*; which have had much of a *nitrous taste*; and would be melted with a gentle *Heat*, as *Nitre* is; and even as easily as *Butyr* it self: I mean not, by the addition of any sort of *Liquor*, or any other *Body*, to dissolve it; but only by the *fire*.

8. §. And as for a *Sea-salt*, that I might *Imitate Nature* for the making thereof, I consider'd, That the said *Salt* is nothing else but that of *Animals* and *Vegetables*, freed from its true *Spirit* and *Sulphur*, and some *Saline particles*, specifically *Animal* or *Vegetable*, together with them. For both *Animal* and *Vegetable Bodies* being continually carried by all *Rivers* into the *Sea*; and many likewise by *Shipwrack*, and divers other ways immersed therein: they are at last corrupted, that is, their *Compounding* parts are opened and resolved. Yet the *Resolution* being in the *Water*, is not made precipitately, as it is in the *Air*; but by degrees, and very gently; whence the *Sulphurous* and other *Volatile* parts, in their *Avolation*, make not so much haste, as to carry the more fixed *Saline* parts along with them; but leaveth them behind in the *Water*, which imbibeth them as their proper *Menstruum*.

And the *Imitation* of *Nature* herein, may be performed thus; Put as much of a *Lixivial Salt* as you please, into a wide-mouth'd *Bottle*, and with fair *Water* make a strong *Solution* of it; so as some part thereof may remain unresolved at the bottom of the *Bottle*. Let the *Bottle* stand thus for the space of about half or three quarters of a year, all the time unstopped. In which time, many of the *Sulphurous* and other *Volatile* parts gradually flying away; the top of the unresolved *Salt* will be incrustate, or as it were frosted over, with many small and hard *Concretions*, which, in their nature, are become a true *Sea-Salt*. Whereof there is a double Proof; First, In that most of the said *Concretions* are of a *Cubical*, or very like *Figure*. Especially on their upper parts; because having a fixed *Body* for their *Basis*, their under parts, therefore, contiguous thereto, are less regular. Whereas the parts of the *salt* in the *Sea*, being environed on all sides with a *Fluid*; their *Figure* is on all sides regular. Secondly, In that a strong *Acid Spirit* or *Oyl* being poured upon a full body'd *Solution* hereof; yet it maketh herewith no *Ebullition*, which is also the property of *Sea-Salt*. And thus much for the more *General Imitation* of *Bodies*.

I N S T A N C E. III, & IV.

FROM the aforesaid *Premisses*, and by the aforesaid *Means*, there is no doubt to be made, but that also the other *sensible Qualities* of Bodies may be *Imitated*, as their *Odors*, and *Tasts*. And that not only the *general* ones, as *Fragrant*, or *Astringent*: but also those which are *specific* and *proper* to such a *species* of Bodies.

2. §. Thus for *Example*, by *mixing Spirit of Nitre* or *Vitriol* with *rectified Oyl of Turpentine*, and some other *Vegetable Oyls*, severally, and in a due *Proportion* and *Time*, I have *Imitated* the *Smells* of *divers Vegetables*; as of *Tansy*, of *Lignum Rhodium*, and *others*. And I conclude it *feasible*, To *Imitate* the *Tast* or *Smell* of *Musk*, or *Ambergreece*, or *any otherbody* in the world.

3. §. Hence also we may be *Taught*, How to *Imitate* the *Faculties*, as well as other *Qualities* of Bodies. The reason is, because even *these* have no dependance upon any *substantial Form*: but are the meer result of *Mixture*; effected by the same *Causes*, whether in *Nature* or *Art*; as I think I have made to appear in the foregoing *Idea*. And as in the *Premisses* of this *Discourse* hath been shew'd.

Id. §. 55.

Ch.2. §.10.

Ch.3. §.10.

I N S T A N C E V.

FROM whence, *again*, it is likewise a *Key* to *Discover* the *Nature* of Bodies. For how far soever we can attain to *Mingle*, or to *Make* them, we may also know *what they are*.

2. §. For Bodies are *mingleable*, either of *themselves*, or by some *Third*. As to those which *mingle of themselves*, we may certainly conclude, That there is a *congruity* betwixt them, in some respect or other. So upon various *Tryals* I find, That *Essential Oyls* do more easily *imbibe* an *Acid*, than an *Alkaly*. Whence it is evident, That there is some *Congruity* and *Similitude* betwixt *Essentian Oyls*; and an *Acid*, which there is not betwixt the said *Oyls* and an *Alkaly*.

3. §. As to those that *mingle* only by some *third*; we may also certainly conclude, That though the *two extrems* are *unlike*; yet that they have both of them some *congruity* with that *third*, by which they are *united*.

4. §. Moreover, We may make a *Judgment* from the *manner* or *Degree* of *Mixture*. Thus the *Acid Spirit* of *Nitre*, as is said, will *coagulate Oyl-Olive*, and render it *consistent*. Whence it might be thought, That any other strong *Acid* will do the like; and that therefore, there is no great difference in the *Nature* of the said *Acid Liquors*. But the contrary hereunto, is proved by *Experiment*. For having *digested* the same *Oyl* in the same *manner*, and for a much *longer* time, with strong *Oyl of Sulphur*; although it thence acquired some change of *Colour*; yet not any *Consistence*.

5. §. *Again*, Because the said *Spirit* of *Nitre* *coagulates Oyl-Olive*; it might be expected, it should have the same effect upon *Oyl of Aniseeds*; or, at least, that if other *Acids* will *Coagulate Oyl* of *Aniseeds*, that this should do it *best*. But *Experiment* proveth the contrary. For of all I have tryed, *Oyl of Vitriol* is the only *Acid* that doth it *instantaneously*. *Oyl of Sulphur*, if very strong, will do it; but not so soon;

nor so much *Aqua fortis*, and *Spirit of salt*, for the present, do not at all touch it. And *Spirit of Nitre* it self will not coagulate it, under eight or ten hours at least.

I N S T A N C E VI.

LASTLY, and consequently, It is a Key To Discover the Medicinal Use and Operation of Bodies. Thus, for Example, by the Imitation of *Resins* and *Resinous Gums*, we certainly know what all of them are, and when, and wherefore to be used. For what are *Mastick*, *Frankincense*, *Olibanum*, *Benzoin*, and other like *Resins*, or *Resinous Gums*, for their principle and predominant parts, that is, *quà Resins*; but Bodies resulting from *Natural*, in like manner, as I have shewed, they may be made to result, from *Artificial Mixture*? That is to say, the *Oleous*, and *Acid* parts of *Vegetables*, being both affused and mingled together, *per minima*, in some one sort of *Vessels* in a *Plant*, they thus incorporate into one consistent and friable Body, which we call *Resin*.

2. §. Now from hence it is, That the said *Resins*, and *Resinous Gums*; as also *Amber* and *sulphur* for the same Reasons; are of so great and effectual Use against most *thin* and *salt Rheums*; *sc.* as they are *Acidoleous* Bodies. For by their *Acid* parts, which in all these Bodies are exceeding copious, they mortifie and refract those *Salt* ones, which feed the *Rheum*. And by their *oleous* parts, the same *salt* ones are also Imbibed. Whence, they are all, in some degree, incorporated together; that is, The *Rheum* is thickned: which is the desired effect.

3. §. Whereas, on the contrary, if the *Cough* proceed not from a *thin*, and specially a *Salt Rheum*, but from a *Viscous Flegm*; the use of many other Bodies which are also more *oleous*, and abound not so much with an *Acid* as these do, especially some of them, is more proper: such as these, in this Case, proving sometimes not only ineffectual, but prejudicial. Since the very Cause of the said *Viscousness* of *Phlegm*, is chiefly some great *Acidity* in the *Blood*, or in some other part, as may be proved by divers Arguments.

4. §. Many more Instances might be hereunto subjoynd: and may hereafter be offered to the acceptance of such, who are inquisitive into matters of this Nature. If I shall not herein anticipate, or reiterate the Thoughts and Observations, of those two Accurate and Learned Persons *Dr. Willis*, and *Dr. Walter Needham*, as to what the one hath already published, and both have put us in Expectation of. But the Instances already given, are sufficient to evidence what I have said. And, I hope, this present Discourse to prove, in some measure, thus much; That Experiment, and the Common Notions of sense are prolifick; and that nothing is Barren, but Phansie and Imagination.

An Appendix to the precedent discourse of Mixture.

A V I N G, in the first Edition of the foregoing Dis- *Ch. 5. Inst.*
course, made mention of the preparation of *Essen-* *1. §. 8.*
tial Oyls, so as to become easily mingleable with
any *unoyly Liquor*. I shall here acquaint the Rea-
der, That this may be done, by digesting any of
the said *Oyls* with about an equal quantity of the
Yelk of an *Egg*, with a very soft heat, like that of
the *Meridian Sun* in *Summer*, continued for the

space of three Weeks or a Month; and in the mean time, to be now
and then stirred a little together. The *Yelk* will by degrees, imbibe the
Oyl, and at length be incorporated with it, and become a *Balsam*, as
white as Milk, easily dissoluble in any *watery* or *winy Liquor*.

2. §. I confess, that it will be very difficult to prepare any good
quantity for use, this way. But this being a sufficient proof of the
possibility of such a *Mixture*; I considered, whether the application of
some other forementioned *Cause of Mixture*, might not supply the de-
fect of this: and hereupon, have made several successful tryals; not
only for the mixing of the said *Oyls*, but likewise of all sorts of *Rosins*
and *Gums* with any *winy* or *watery Liquor*, in great quantities, in a
short time, and without much trouble. But for the mixing of some of
them, the *Yelk* of an *Egg* alone will not serve, without the interve-
ning of some other sociable Body, according to one of the *Rules* given *Ch. 5. Inst.*
in the foregoing *Discourse*. *1. §. 6.*

3. §. In the same *Discourse*, upon certain premises, I have laid
down this following conclusion.

— By accumulating the *Causes of Mixture*, that is, by joyning *Ch. 5. §. 5.*
“two or three or more together; or by applying more in some cases,
“where *Nature* applyeth fewer; we may be able to make, if not a more
“perfect, yet a far more speedy *Mixture*, than *Nature* doth. As by joyning
“ing *COMPRESSION*, *Heat*, and *violent Agitation*, and so con-
“tinuing them altogether, by some means contrived for the purpose, for
“the space of a Week or Month, or longer without *Cessation*. Which
“may probably produce, not only strange, but useful effects, in the
“*SOLUTION* of some, and the *Mixture* of other *Bodies*.

4. §. For the proof whereof, and that I had thoroughly weighed
what I have said, Mr. *Pappin* hath since given us an ingenious Instance;
in his new *Digester*. Which is, a *Balneum Mariæ clausum*: all *Infusions*
and *Digestions* made with *Double Vessels*, having hitherto been made
with the outer *Vessel*, open. So that whereas by the old way of *Di-*
gestion, their is no other *Power* made use of but that of *Heat*: in this
way, that also of *Compression* is joyned therewith:

EXPERIMENTS
IN
CONSORT
OF THE
LUCTATION
Arising from the Affusion of several
MENSTRUUMS
Upon all sorts of
BODIES,

Exhibited to the *Royal Society*, April 13. and
June 1. 1676.

Ch. 5. Inst. 5



THE intent of the following Experiments is twofold. The one, to be as a Demonstration of the Truth of one, amongst other Propositions, laid down in the precedent Discourse of Mixture, *sc.* That it would be a Key to let us easily into the knowledge of the Nature of Bodies.

The other, and that consequently, To be as a Specimen of a Natural History of the *Materia Medica*: that is to say, a multifarious Scrutiny into the intrinsic Properties of all those Materials, which have been, or may be used in Medicine: for the performance whereof, the following Method is exhibited as one, amongst others, necessary to be insisted upon. For what Dominion a Prince hath over the Moral, that a Physician hath, as one of God Almighty's Vice-Roys, over the Corporeal World. Whom therefore nothing can more import, than a particular knowledge of the Genius of all his Subjects, those several Tribes of Matter, supposed to be under his Command.

There

There are some known Observations of this nature : but there is no Author, I think, who hath given us a Systeme of Experiments upon the Subject : The performance whereof is here intended.

The Experiments may seem too numerous to be of one make. But no less a number would have answered the design of an Universal Survey ; which, though less pleasing, proves the more instructive in the end : not being like angling with a single Hook ; but like casting a Net against a shoale : with assurance of drawing up something. Besides the advantage of comparing many together ; which being thus joyned, do oftentimes, like Figures, signifie ten times more, then standing alone, they would have done.

How far the Corollaries all along subjoyned have made this good, is left to the Reader to judge. And also, to add to them, so many more, as he pleases : for I make my own Thoughts no mans Measure.

C H A P. I.

What is generally to be observed upon the Affusion of the Menstruum ; and what, particularly of Vegetable Bodies.



THE Bodies whereupon I made tryal, were of all kinds, *Animal, Vegetable, and Mineral*. Amongst *Vegetables*, such as these, *scil. Date-stones, Ginger, Colocynthis, Pyrethrum, Hawthorn-stones, staphisagria, Euphorbium, the Arenule in Pears, Semen Milii Solis, Tartar, Spirit of Scurvygrass, Spirit of Wine, &c.*

2. §. Amongst *Minerals*, several sorts of *Earths, Stones, Ores, Metals, Sulphurs, and salts*.

3. §. Amongst *Animals* ; such as these, *scil. Hairs, Hoofs, Horns, shells, and shelly Insects, Bones, Flesh, and the several Viscera, Silk, Blood, Whites and Yelks of Eggs, Sperma Ceti, Civet, Musk, Castor, Gall, Urine, Dungs, animal Salts and Stones*.

4. §. The *Liquors* which I poured hereupon severally, were these, *sc. Spirit of Salt Armoniac, Spirit of Harts-Horn, Spirit of Nitre, Aqua fortis, Oyl of Salt, Oyl of Sulphur, and Oyl of Vitriol* ; commonly so called.

5. §. In the *Mixture* of these Bodies, two things, in general, are all along to be observed, *viz. First*, which they are, that make *any, or no Luctation*. For, as some which seem to promise it, make none : So, many, contrary to expectation, make a considerable one.

6. §. Next, the *manner* wherein the *Luctation* is made ; being with much variety in these *five sensible Effects*. 1. *Bullition* ; when the Bodies mixed produce only a certain quantity of froth or bubbles. 2. *Elevation* ; when, like Paste in baking, or Barm in the working of Beer, they swell and buff up. 3. *Crepitation* ; when, they make a kind

kind of hissing and sometimes a crackling noise. 4. *Effervescence*; then only and properly so called, when they produce some degree of heat.

5. *Exhalation*; when not only fumes, but visible steams are produced.

7. §. Of all these, sometime one only happens, sometimes two or more are concomitant. Sometimes the *Luſtation* begins presently upon mixture, and sometimes not till after some intermission. In some bodies, it continues a great while; in others, is almost instantaneous: Examples of all which I shall now produce; beginning with *Vegetables*, as affording the least variety.

8. §. And first, if we take Spirit or Oyl of Salt, Oyl of Vitriol, Spirit of Nitre, or Aqua fortis, and pour them severally upon the several parts of Vegetables, as Roots, Woods, Stones, &c. we shall find, that they are, generally far less apt to make a *Luſtation*, than either Animal, or subterranean Bodies. Whence, as from one argument, it seemeth evident, That in most Vegetables, and in most of their parts, the predominant Salt is an Acid. But that, on the contrary, the predominant Salt in most Minerals, and parts of Animals, is an Alkaly: in the former, usually a fixed; in the latter, a volatile Alkaly.

9. §. Again, although the *Luſtation* which most Vegetables, and most of their parts make with Acids, be but small, yet some they make; especially with some Acids, as with Spirit of Nitre and Aqua fortis. Whence it seemeth plain, That there is an Alkaline Salt existent in many Vegetables, even in their natural estate; and that it is not made Alkaline, but only Lixivial, by the fire. Or, there is some quantity of a Salt, call it what we will, in the said Bodies, which is so far different from an Acid, as to make a *Luſtation* therewith. But to give particular instances of the several proportions, or manner of Mixture, wherein it appears to be in several Plants.

10. §. And first, of all vegetable Bodies, Date-stones are amongst the least apt to make a *Luſtation* with Acids, if they may be said to make any at all. Hence they are not so potent Nephriticks, as many other Stones, which make a more sensible *Luſtation*.

11. §. Ginger makes a small Bullition with Aqua fortis, only observable by a Glass. Hence the pungency of Ginger lyeth in a sulphureous and volatile Salt, which yet is very little Alkalizate.

12. §. Scurvygrass-seeds make a very small Bullition with Aqua fortis, like that of Ginger. So doth also the Seed of Purslane. Hence, although there is much more of a certain kind of volatile Salt in Ginger or Scurvygrass, than in Purslane; yet there is little more of an Alkaly in any one, than in an other.

13. §. The Pulp of Colocynthis, Fruit-Stones, the stony Covers of the Seeds of Elder, of white Bryony, of Violets, and others, with Aqua fortis make a Bullition just perceivable without a Glass. Hence it appears, That the great Cathartick power of Colocynthis lieth not so much in an Alkaly, as an Acid; as making a much less Bullition, than some other vegetable Bodies, which are less Cathartick. For which reason likewise it is, That the best Correctors, or Refractors of the force of Colocynthis, are some kinds of Alkalies, as particularly that of Urine, as Riverius hath somewhere observed.

14. §. The Root of Pyrethrum, with Aqua fortis, makes a Bullition and buff, in a short time. Hence, the Cause of a durable Heat, upon the Tongue, is an Alkalizate Sulphur. For the Heat of Ginger, though greater;

greater; yet abideth nothing near so long as that of *Pyrethrum*; which, as is said, maketh also a more sensible *Bullition* with *Acids*.

15. §. *Kermes-berries*, commonly, but ignorantly, so called, with the said *Liquor*, buff up to an equal height, but in a somewhat longer time. Hence they are gently astringent; scil. as their *Alkaly* binds in with some preternatural *Acid* in the stomach.

16. §. *Hawthorn-stones*, with *Aqua fortis*, buff up equally with the former *Body*; but the *Bullition* is not so visible. The like is also observable of *Medlar-stones*. Hence, as they contain a middle quantity of an *Alkaly*, they are not insignificantly used against the *Stone*.

17. §. *Seeds of Staphisagria*, with *Aqua fortis*, make a *Bullition* still more visible. But it quickly ends. This confirms what was said before, sc. That the cause of a durable *Heat* is an *Alkaline Sulphur*; these *Seeds* producing a durable *Heat*, as doth the *Root of Pyrethrum*.

18. §. The *Seeds* also of red *Roses*, *Borage*, and *Comfrey* do all with *Aqua fortis* make a considerable *Bullition* and buff; and that very quickly. So that amongst all *Shells* and *Stones*, those generally make the greatest *Bullition*, which are the hardest and the brittlest, and so the fullest of *Salt*.

19. §. *Euphorbium* makes a *Bullition* yet more considerable, with much froth, and very quickly. From which Experiment, compared with two of the former, it appears, That *Euphorbium* is not an *Acid*, but an *Alkaline Gum*. As also, that the cause of its so very durable *Heat*, is an *alkaline Sulphur*, as of *Pyrethrum* and *Staphisagria* hath been said. It seems also hence evident, that the power of all great *Sternutatories* lyeth not in their *Acid*, but their *Alkalies*.

20. §. The *Arenula* or little stones in *Pears*, cluster'd round about the *Coar*, with *Aqua fortis*, presently buff up, and make a great *Bullition* and *Efferescence*, much greater than do any of the *Bodies* above-named. Whence, although, so far as I know, they have never yet been used in *Medicine*; yet it is probable, that they are a more potent and effectual *Nephritick*, than any of the *Bodies* aforesaid, some of which are usually prescribed. It is hence also manifest, That, according to what I have elsewhere said, for the sweetning of the *Fruit* and *Seed*, the *Tartareous* and *Alkaline* parts of the *Sap*, are precipitated into their *Stones*, stony parts, and *Shells*.

Anat. of
Plants, B.
I. Ch. 6.

21. §. The last Instance shall be in the shells of the *Seeds of Miliun Solis*; which not only with *Aqua fortis*, but some other *Acids*, make a greater and quicker *Bullition* and *Efferescence*, than any other vegetable *Body*, upon which I have yet made tryal, in its natural estate. Hence, as well as from divers of the last fore-going Instances, we have a clear confirmation of what I have, towards the beginning of this Discourse, asserted; sc. That there is some kind of *Alkaline Salt* in *Plants*, even in their natural estate. As also, that they are as significantly used against the *Stone*, quatenus alkalizate, as *Millipedes*, *Egg-shells*, or any other testaceous *Bodies* of the same strength. To these I shall subjoyn one or two Examples of *Vegetable Bodies* which are more or less altered from their natural estate.

22. §. Neither *Crystals of Tartar*, nor *Tartar* it self (although they have some store of *alkaline* mixed with their acid parts) make any *Efferescence* with *Acids*, but only with *Alkalites*, as *Spirit of Harts-Horn*, &c. Hence the *calculous sediment* or *Arenula* in *Urine*, may not so properly be called

the *Tartareous* part of the *Urine*; the events following the mixture hereof with the aforesaid *Salts*, being quite contrary; as will be seen in the *Last Chapter*.

23. §. *Spirit of Scurvy-grass* maketh no *Luētation* with any *Acid*. Hence (as from a former Experiment was above-noted) it seems, That there may be a kind of *volatile Salt*, which is neither *acid*, nor *alkaline*; such as this of *Scurvygrass* and other like *Plants* seems to be: yet contrary to an *acid*; as experience shews in their efficacy against the *acid Scurvy*.

24. §. *Rectified Spirit of Wine*, both with *Spirit of Nitre*, and with *Oil of Vitriol*, severally, maketh a little *Luētation*. Which argues, that there is contained, even in this *Spirit*, some portion of a *volatile Alkaly*.

25. §. *Spirit of Wine*, and double *Aqua fortis*, as the strongest is called, make an *effervescence* so vehement, as plainly to boil.

26. §. Besides the vehemency hereof, there is another surprizing circumstance. For whereas all other *Liquors* which make an *Effervescence* together, will do it in any proportion assigned, although but one drop to a thousand: these two, sc. *rectified Spirit of Wine* and *Aqua fortis*, require a certain proportion the one to the other. For if, suppose, into six drops of *Spirit of Wine* you put but two or three of *Aqua fortis*, they stir no more than if you put in so much *Water*: but drop in about seven or eight drops of *Aqua fortis*, and they presently boil up with very great vehemency. Hence we may conceive the reason of the sudden access of an acute *Disease*, and of its *Crisis*. These not beginning gradually with the *Cause*; but then, when the *Cause* is arrived unto such an *ακμή*, or such a certain *Proportion*, as is necessary to bring *Nature* to the contest. And these may serve for Examples upon *Vegetables*.

CHAP. II.

What may be observed of MINERALS.



HAVING given several Instances of tryal upon *Vegetables*; I next proceed to *Minerals*, which, for some orders sake, I shall distribute into five or six sorts, sc. *Earths*, *Stones* *Ores* and *Metals*, *Sulphurs*, and *Salts*.

2. §. First for *Earths*. *Oyl of Vitriol* upon *Fullers Earth*, doth not stir it, or cause the least *Bullition*. Nor upon *yellow Oker*. Nor upon the *Oker* which falls from *green Vitriol*. The same *Oyl of Vitriol* and *Spirit of Harts-Horn* poured severally upon *Bolus Armena* of two kinds, and upon one kind of *Terra sigillata*, stir none of them. Hence *Bolus's* are the *Beds*, or as it were, the *Materia prima*, both of opacous *Stones*, and *Metals*; into which the said *Bolus's* are transmuted, by being concentrated with divers kinds of *Salts* and *Sulphurs*, which successively flow in upon them.

3. §. *Aqua fortis*, and Oyl of *Vitriol* being poured severally upon another sealed Earth, which was vended by the name of *Terra Lemnia*; they both made a very considerable *Effervescence* herewith. Whence it appears, That there is no small difference in the nature, and therefore the operation of *Bolus Armena* and *Terra Lemnia*. As also, betwixt the sealed Earths themselves, one making a great *Effervescence*, another none at all. Whereto those that use them, are to have regard.

4. §. Next for *Stones*. And first, *Irish Slat*, with *Spirit of Hartshorn*, maketh a small, yet visible *Bullition*: and it presently ceaseth. So that it seems to be nothing else but a *Vitriolick Bole*. As is also argued from its taste, which is plainly acid, and somewhat rough. Whence also it is with good reason given upon any inward Bruises. Because by coagulating the Blood, it prohibits its too copious afflux into the affected part. Yet being but gently astringent, and so the Coagulations it makes, not great; they are likewise well enough carried off from the same part in the Circulation; by both which means an Inflammation may be either prevented, or the better over-ruled.

5. §. *Lapis Hematites* maketh no *Effervescence* at all either with *Alkalies* or *Acids*.

6. §. Powder of the green part of a Magnet with Oyl of *Vitriol* maketh some few bubbles, yet not visible without a Glass. But the powder of the black part of a Magnet, which is the said stone fully perfect, stirreth not with any acid. Neither doth the calcined Magnet. Hence there is some considerable difference betwixt *Iron* and the Magnet.

7. §. *Lapis Lazuli*, with Oil of *Vitriol*, and especially with *Spirit of Nitre*, maketh a conspicuous *Bullition*. Hence its *Cathartick* virtuelyeth in an *Alkaly*. For which reason it is also appropriate, in like manner as *Steel*, to the cure of *Hypochondriacal Affections*; originated from some kind of fermenting Acid.

8. *Osteocolla*, with *Spirit of Nitre* maketh yet a greater *Effervescence*. How it comes to be so great a knitter of broken Bones, as it is reputed, is obscure. It seemeth, that upon its solution by a *Nitrous Acid* in the body; it is precipitated upon the broken part, and so becomes a kind of *Cement* thereto.

9. §. *Lapis Tutia*, with *Spirit of Nitre*, maketh an *Effervescence* much alike. And with Oyl of *Vitriol* very considerably. But *Lapis Calaminaris* with Oyl of *Vitriol* grows stark; as the powder of *Alabaster* doth with water. With *Spirit of Nitre* it maketh a little *Bullition*, and quickly. But with *Aqua fortis*, a great one; beyond any of the *Stones* above named. Hence both *Tutty* and *Calamy* are *Ophthalmicks* from their *Alkaly*. Which is also confirmed, from the efficacy of some *Alkalies* of the like use. Hence also *Calamy* seemeth to partake somewhat of the nature of *Silver*: as by tryal made upon that also, will hereafter better appear.

10. §. *Chalk* and Oil of *Sulphur* or *Vitriol* make as strong an *Effervescence* as any of the rest. Whence it is sometimes well used against a *Cardialgia*.

11. §. *Whiting* makes as great an *Effervescence* as *Chalk*. So that it seems the saline parts are not washed away with the water, wherein the *Chalk*, for the making of *Whiting*, is dissolved.

12. §. Talk will not stir in the least either with Spirit of Nitre, or Oyl of Vitriol. But the Lead-Spar maketh a considerable Effervescence with both of them severally. Hence, however this be also called English Talk, yet there is no small difference betwixt this, and true Talk.

13. §. To these Stones may be added petrified bodies. As petrified wood; which (that upon which I made tryal) no acid stirreth in the least. Petrified shells; upon four or five several sorts whereof, Oyl of Vitriol being poured, produceth a great Effervescence. The Root or rougher part of the Stone called Glossopetra, with Spirit of Nitre, makes a conspicuous Bullition. Astoria, the Stone so called, and found in some places in England, with Oyl of Vitriol, maketh an Effervescence at the same degree. So doth the Belemnites, or Thunder-Stone, both the larger and the lesser kinds. So that none of these are acid, or vitriolick, but alkalizate Stones.

14. §. Coraline, with Oyl of Vitriol, makes a conspicuous Bullition, yet mild and gentle; that is, with very little, if any heat, and without any visible Fumes. And red and white Coral do the like. Hence they are all of a very gentle operation, and fit for Children, as the case requires.

15. §. Magistery of Coral (prepared the ordinary way) stirreth not in the least, either with Alkalies or Acids. Whence it is evident, That its active Principles are in its preparation destroyed and washed away: that is to say, It is an elaborate Medicine good for nothing. And thus far of Stones.

16. §. I next come to Metals and Ores. And first for Lead; upon which Spirit of Salt Spirit of Nitre, or Aqua fortis being dropped, it stirreth not in the least with any of them: but with Oyl of Sulphur, and especially with Oyl of Vitriol it maketh a slow Bullition and froth. Hence it seemeth to be the most alkalizate Metal. Which is also confirmed by a foregoing Experiment upon the Lead-Spar, which maketh a considerable Effervescence with any sort of acid. And which likewise, being calcined, yieldeth a good quantity of Lixivial Salt.

17. §. Lead-Ore stirreth not at all with Aqua fortis or Oil of Vitriol. But Spirit of Salt makes it bubble, and Spirit of Nitre makes it boil. Hence there is a considerable difference betwixt the perfect Metal and the Ore.

18. §. Burnt Lead and red Lead, make a very small Bullition with Oyl of Vitriol, with Spirit of Nitre a far greater.

19. §. Mercury, with Oyl of Vitriol, will not stir, nor with Oyl of Sulphur. But with Spirit of Nitre presently boyls up. Hence Mercury is a subacid Metal; Spirit of Nitre being a subalkaline Acid.

20. §. The filings of Iron or Steel, with Oyl of Vitriol, make a fair Bullition, like that of Minium. But Spirit of Nitre makes them boil with much celerity. Hence Iron is likewise a subacid Metal.

21. §. Steel prepared with Sulphur maketh a far less Effervescence with the same Spirit of Nitre, than do the filings. Hence there is a great difference in their strength. So that ten grains of the filings unprepared, will go as far as fifteen grains or more of those which are prepared, as above-said. Yet in some cases the weaker and milder may be the better.

22. §. There is one Circumstance in the mixture of Steel and Aqua fortis, which is surprizing; and that is this, That strong Aqua fortis, dropped upon Steel, will not, of it self, make the least Bullition: but if hereto

hereto you only add a drop or two of Water, they presently boil up with very great vehemency. The Cause is obscure; yet it is well known, that Water it self will dissolve Iron: so that it appeas, as well by this, as by some other Experiments, that even in common Water, as mild as it is, there is some kind of corrosive Principle.

23. §. *Antimony with Spirit of Nitre, and Aqua fortis severally, maketh an Effervescence; somewhat lower than Iron. With Oil of Vitriol the Bullition is so small, as difficulty to be perceived with a Glass.* Hence it seemeth to be of a very compounded nature; if I may so call it, a subacid-alkaline Metal.

24. §. *Antimonium Diaphoreticum, with Spirit of Nitre and Oil of Vitriol severally, makes a considerable Effervescence.* Wherefore it is not an useles Preparation; as from the Calcination and Ablution used therein, some have thought.

25. §. *Bezoardicum Minerale, (that upon which I made tryal) stirreth not at all either with Alkalies or Acids.* To which, let those who make use of it, have regard.

26. §. *Tin, with Spirit of Nitre, makes so hot and vehement an Effervescence, that it turns presently, as it were, into a Coal. It makes also a fair Bullition with Oyl of Vitriol. And a gentle one with Spirit of Salt.* Wherefore, it hath something of the nature both of Iron, Lead, and Copper.

27. §. *The like remarkable circumstance is seen in the mixture of Aqua fortis with Tin, as with Iron. For Tin and strong Aqua fortis of themselves will not stir; but add a few drops of water to them, and they boyl up with the greatest vehemency.*

28. §. *Copper, with Spirit of Salt, and Oyl of Vitriol severally, stirs not at all. Spirit of Nitre, and Aqua fortis, both boil it up vehemently. Neither Spirit of Harts-horn, nor Spirit of Salt Armoniac maketh any Bullition therewith. But both of them, by a gentle solution, that is, gently separating its Sulphur from its Salts, turn it blue.* Hence Copper hath a greater proportion of acid than any of the forementioned Metals.

29. §. *Silver, neither with Spirit of Salt, nor Oyl of Vitriol makes any Bullition. With Spirit of Nitre it makes one, but tis soon over: and then continues to dissolve slowly into white Coagulations. It also maketh with Spirit of Harts-horn, or of Salt Armoniac, a full and deep blue.* Hence there is a greater proportion of acid in Silver, than in Lead, Mercury, Iron, Antimony, Tin, or Copper.

30. §. *Litharge of Silver maketh the greatest Effervescence with Oyl of Vitriol. Yet some with spirit of Nitre. And with spirit of Salt Armoniac maketh some little huff or elevation. And being mixed with spirit of Nitre and Spirit of Salt Armoniac both together, produceth a faint blue.* Hence, although the far greater part of this Litharge be but Lead; yet, it seems, it hath some small mixture of Silver. But that of Gold seemeth, for contrary reasons, not to have any Gold.

31. §. *Gold maketh no Effervescence with any single salt I know of. But it is commonly dissolved with Aqua Regis, which is known to be an alkaline Liquor.* Whence it seemeth, That as Lead is the most alkalisate, so Gold the most acid of Metals.

32. §. These things considered, and other observations added hereunto, may possibly give some directions, not only for the ordering and using, but even for the making, imitating and transmuting of *Metals*. Thus far of *Metals*.

33. §. I will next give one or two Instances of tryal upon *Sulphurs*. And first *Sulphur vive*, with *Aqua fortis*, maketh an apparent *Bullition*, but it is some time, before it begins. But the factitious or common *Brimstone*, maketh scarce any, if any at all. So that there is no small difference betwixt them.

34. §. *White and yellow Arsenick* make no *Bullition* either with *Alkalies* or *Acids*. Wherefore the strength of its operation on the Body, lies more in a *sulphur* than a *Salt*; or in a *Salt* drowned in its *Sulphure*.

35. §. The ashes either of *Pit-Coal*, or *Sea-Coal*, make no *Efferescence* with *Alkalies* or *Acids*. Whence the saline Principle is altogether volatile, and sublimed away by the fire.

36. Lastly for *Salts*. And first of all, *Borax* maketh no *Efferescence* nor any *Fumes* with *Oyl* of *Vitriol* or *Spirit* of *Nitre*.

37. §. *Oyl* of *Vitriol* and *Nitre* make *fumes* or *steams*, though no *Efferescence*.

38. §. *Green Vitriol*, with *Spirit* of *Harts-Horn*, is scarcely moved. *White Vitriol*, with the same *Spirit*, maketh a conspicuous *huff*. And *Roman Vitriol* a vehement *Efferescence*. Whence the former is the least acid, and the latter the most of all. Which also confirms what I said before of the like natures of the several *Metals* to which they belong.

39. §. *Salt* of *Vitriol*, though a fixed *Salt*, and made by *Calcination*, yet maketh no *Efferescence* with the strongest acid; but only with *Alkalies*; as may be seen upon their mixture, but much better heard by holding the mixture to ones ear. Hence, there are fixed *Acids*. Which further confirms what I have above asserted concerning the nature of *Gold*, sc. That the predominant *Salt* thereof is a fixed *Acid*.

40. §. *Sal Martis*, with *Spirit* of *Harts-horn*, maketh a considerable *huff*. Hence it is much more acid than *green Vitriol*; and is therefore a cooler body.

41. §. *Alum* and *Spirit* of *Harts-horn* make a plain *Efferescence*.

42. §. *Saccharum Saturni*, with *Oyl* of *Vitriol*, stirs not at all. With *Spirit* of *Salt*, huffs a little. With *Spirit* of *Nitre* much more. Hence the acid of the *Vinegar*, and not the *Alkaly* of the *Lead*, is the predominant Principle.

43. §. *Common Salt* stirs neither with *Spirit* of *Salt*, nor with *Spirit* of *Nitre*; nor with *Aqua fortis*. But with *Oyl* of *Vitriol* it maketh a great *Efferescence* with noise and *steams*. Hence, even common *Salt*, though it be not reckoned amongst *alkaline Salts*, yet is far nearer in nature to that, than to an acid. Hence also the *Spirit* of *Salt* is a *subalkaline Acid*, and of a very different nature from *Oyl* of *Sulphur* or *Vitriol*.

44. §. *Salt Armoniac*, with *Spirit* of *Nitre*, stirreth not. But with *Oyl* of *Vitriol* it maketh a great *Efferescence*. Hence *Spirit* of *Nitre* is a *subalkalitate Spirit*.

45. §. *Oyl* of *Vitriol* and *Spirit* of *Nitre*, though both acids, yet make a great *smoak*; greater than that which the *Spirit* maketh of it self. Which confirms the last precedent *Corollary*.

46. §. Oyl of Vitriol and Spirit of Salt, though both acids, yet make a strong Effervescence, with noise and fumes. Which further confirms, what was noted before, sc. that Spirit of Salt is a subalkaline Acid.

47. §. Spirit of Salt Armoniac, with Oyl of Vitriol, makes an Effervescence so extraordinary quick, and as it were instantaneous, that nothing seemeth quicker. Whence it is probable, That if Gun-powder were made of Salt Armoniac, instead of Nitre, or with both mixed together; it would be far stronger, than any kind now in use. And thus far for Minerals.

48. §. I have only one Corollary to add, from the whole; which is, That whoever doth undertake the Natural History of a Country, (such as that the Learned Dr. Plot hath exceedingly well done of Oxfordshire) the foregoing Method, seemeth so easie, cheap, and indeceitful, for the finding out and well distinguishing the natures of all kinds of Metals, Ores, Salts, Earths, Stones, or other subterranean Bodies; as cannot, I think, be supply'd, but by others of greater difficulty and expence.

C H A P. III.

What may be observed of the PARTS of Animals.



NOW proceed to the several Parts of Animals; as Hairs, Hoofs, Horns, Shells and shelly Insects, Bones, Flesh and the several Viscera, Silk, Blood, Eggs, Musk, Castor, Gall, Urine, Dungs, Salts and Stones.

2. §. And first of all, the Hair of a mans head, with Oyl of Vitriol, maketh no Bullition at all. Nor yet with Spirit of Nitre. So that although it contains a good deal of volatile Salt; yet it seemeth either not to be alkaline, or else is centred in so great a quantity of Oyl, that the acid menstruum cannot reach it.

3. §. Hares Fur, with spirit of Nitre, maketh, although a short, yet very plain Bullition and huff. Hence the Hair, and therefore the Blood, of some Animals, is fuller of Salt, at least of an Alkaline Salt, than that of some others. And perhaps the Hair of some men, as of Black's, may be so full of salt, as to make a Bullition like Hares Fur.

4. §. The shavings of Nails stir not at all, either with Oyl of Vitriol, or Spirit of Nitre: only with the latter they turn yellow. But Elks Claws, with spirit of Nitre, make a small and slow Bullition.

5. §. Horses Hoof, with Oyl of Vitriol, stirs not of many hours. But with spirit of Nitre, allowing it some time, makes a very plain Bullition, and huffs up very high.

6. §. Cows Horn, neither with Oyl of Vitriol, nor with Spirit of Nitre, maketh any Bullition, only turneth to a yellow colour.

7. §. Rams Horn stirs not with Oyl of Vitriol; but with Spirit of Nitre, makes a small and slow Bullition.

8. §.

8. §. Harts-Horn makes a considerable Bullition and huff, even with Oyl of Vitriol, which the rest of the Bodies abovesaid, will not do. But with spirit of Nitre, it makes yet a greater. From the foregoing Experiments, and almost all that follow, what is before asserted of the salts of Vegetables and Minerals, is here also evident concerning that of Animals, scil. That it is not made, but only separated by the fire. It likewise hence appears, That the proportion of Salt in the forementioned parts is very different; and that therefore some of them are never, and none of them but with good discretion, to be substituted one for another in Medicine. As also, that there is a different proportion of salt in the several Animals themselves, to which the said Parts belong.

9. §. Next for shells; as those of Lobsters, Eggs, Snails and Oysters: all which make an Effervescence, both with Oyl of Vitriol, and Spirit of tre. But with Spirit of Nitre the greatest. Lobster-shells make a considerable Bullition and huff, but no noise nor steams. Egg-shells make a Bullition and huff, with some noise, but no steams. Snail-shells make an Effervescence with noise and steams. Oyster-shells make one with the greatest noise and thickest steams. Hence we may judge, in what case to administer one more appositely than another. As also in what proportion, according to their different strength. Some may be better for Children, as being milder. Or for a Body whose very sharp Blood or other Humors, are more easily kindled into Ferments. Or else may be safest, to avoid a sudden precipitation of the Humors; or for some other cause.

10. §. Oyster-shells, and the rest above-said, make a quicker Effervescence, not only with Spirit of Nitre, but even with spirit of Salt, than they do with Oyl of Sulphur, or Oyl of Vitriol. So that these bodies, as well as Metals, have their proper Menstruums whereby they are dissolved.

11. §. Egg-shells calcined, make with Oyl of Sulphur, or Oyl of Vitriol, or Spirit of Nitre, a greater Effervescence, than when uncalcined. As also with steams, which uncalcined, they produce not. The like is seen in calcined Oyster-shells. And the longer the Calcination is continued, the quicker and stronger will be the Effervescence. This I tryed at several terms, from a quarter of an hour, to five hours. So that after so long a Calcination, they make an Effervescence almost instantaneous. The reason hereof is, Because the several Principles whereof the Shells consist, being relaxed, and the Sulphur for the greatest part, driven away by the fire; the remaining Salt lies now more open and naked to the attaque of the Menstruum, so soon as ever they are mixed together. From hence it is plain, That Egg-shells, and the others above-said, being burnt, are far stronger Medicines, than when unburnt. It is hereby likewise evident, That a great portion of their Salt, is not a volatile, but a fixed Alkaly. To these may be subjoyned all kinds of shelly Insects. I will instance in three or four.

12. §. And first Bees, with Oyl of Vitriol, stir not in the least. With Spirit of Nitre they make an exceeding small Bullition, without any elevation.

13. §. Cockinele (the Nest of an Insect) makes some Bullition with Oyl of Vitriol, but very small: for the bubbles are not to be seen without a Glass. But with Spirit of Nitre the Bullition is more visible, and joyned with some elevation.

14. §. *Cantharides* make no visible Bullition with Oyl of Vitriol. But with Spirit of Nitre they do, and huff up rather more than *Cochinele*. Yet is this done very slowly, and comparatively with many other bodies, is not much. Hence it is not the quantity, but the quality of their volatile Salt, which makes them so strong an *Epispastick*. For most of those Bodies above, and hereafter named, make a greater Bullition, and yet are neither *Caustick* nor *Epispastick* in the least. It is hence also evident, as hath been before suggested, That there are divers kinds of *Volatile Salts*, eminently different; some being highly *alkaline*, others very little, and some scarce any thing so: such as those of *Scurvy-grass*, *Anemone*, *Crowfoot*, and many the like *Plants*; to whose salts, this of *Cantharides* seemeth to be very near of kin.

15. §. *Millepedes* make a Bullition and huff, much greater and quicker, than any of the *Insects* above-named: and that both with Spirit of Nitre, and Oyl of Vitriol it self. Yet is this *Insect* of a very temperate nature. Whereby is further demonstrated, That the being simply *alkaline*, is not enough to make a body to be *Caustick*.

16. §. Again, although *Millepedes* make a Bullition, greater than any of the *Insects* above named: yet is it much less, than that of *Oyster*, *Snail*, or even *Egg-shells*; and of divers other bodies above, and hereafter mentioned. Hence, being given to the same intent, as any of those bodies; it is the mildest and gentlest in its operation of them all.

17. §. *Millepedes* likewise calcined, makes a stronger *Efferescence*, than when uncalcined, as do the *Oyster-shells*, &c. So that it appears, That all *Testaceous salts*, are at least in part, *fixed Salts*.

18. §. I next proceed to *Bones*. And first *Whale-bone* maketh no Bullition at all with any acid. A *Cartilage*, with Spirit of Nitre, makes some very small bubbles, not to be seen without a *Glass*.

19. §. The Bone in the Throat of a *Carp*, makes a little and slow Bullition with Spirit of Nitre. The Spina of a *Fish* (that which I used was of a *Cod-fish*) maketh a Bullition one degree higher.

20. §. All sorts of *Teeth*, as *Dogs*, *Boars*, the *Sea-horse*, *Elephant*, make the like. As also the Bone of an *Oxes heart*. So that all these are very gentle in their operation, and fit for *Children*.

21. §. *Sheeps* and *Calves Bones* both of them make a Bullition yet a little higher, especially with Spirit of Nitre. *Cocks Bones* somewhat higher than the former. *Cranium humanum* a little higher than all the rest.

22. §. *Bones* likewise, being calcined, make a Bullition with *Acids*. And so doth also calcined *Harts-Horn*. But in neither of them, is the Bullition advanced by *Calcination*, any thing comparable to what it is in shells. Whence it appears, That the Salt of *Horns* and *Bones*, is much more volatile, than that of shells.

23. §. Next for *Flesh* and the several *Viscera*. And first, dried and powdered *Mutton*, with Oyl of Vitriol, stirs not at all. But with Spirit of Nitre makes a small Bullition and huff. *Sheeps Heart* doth the like somewhat more apparently. *Vipers flesh* produceth a froth, but huffs not. Powdered *Earthworms* make a great froth, and huff a little. Powdered *Tripe* makes only a little Bullition. *Lamb-stones* do the like. *Kidney*, *Spleen*, and *Liver*, with some elevation. *Lungs*, with bubbles very large; because extraordinary slowly. Dried *Brain* makes also a little

and slow Bullition. Hence there is a greater proportion of Sulphur or Oyl, and less of an Alkaly in all these parts, than there is in Bones, Shells, and divers other parts hereafter mentioned. And in some of them, as in the Brain, that Alkaline Salt which there is, may rather be lodged in some sanguineous parts mixed with them, than in their own proper substance.

24. §. I proceed to instance in all sorts of Animal Contents. And first, raw Silk, with Spirit of Nitre, makes a very small Bullition, but the elevation is considerable.

25. §. The grumous part of the Blood dried, with Oyl of Vitriol, stirs but little. But with Spirit of Nitre it huffs up considerably.

26. §. Serum of Blood dried, with the same Spirit makes a plain elevation, with a little Bullition. Herewith may be reckoned the White of an Egg, which is nothing but a pure Crystalline Serum separated from the common stock. This being dried, with Spirit of Nitre, huffs up rather more than even the grumous part of the Blood, the Bubbles are much larger, break oftner, and the elevation sooner made. Whence it seemeth, that there is a greater quantity of a volatile Alkaly in proportion to the sulphur, requisite to the Generation, than to the Nutrition of an Animal.

27. §. The Yelk of an Egg is scarce moved with spirit of Nitre, producing only a very few Bubbles. The Salt being either little alkalizate, or else immersed in so great a quantity of Oyl, that the Menstruum cannot reach it. For the same reason Sperma Ceti stirs not with any Acid. Neither doth Civet.

28. §. Russian Castor, with Oyl of Vitriol, stirs not. But with Spirit of Nitre makes a considerable huff and froth. Yet it requires time. Wherefore it seemeth, That Castor by virtue of its alkaline sulphur, becomes so good a Corrector of the acid-alkaline sulphur of Opium: so I take leave to call it, having some reasons to believe it such.

29. §. Musk, with Oyl of Vitriol, stirs not. But with Spirit of Nitre it makes a considerable and quick Bullition, with large bubbles, which often break and rise again. Whence there is a very eminent difference betwixt Musk and Civet. Hence also, Musk is Cordial, not only from its sulphur, but its Alkaly; by both directly opposite to preternatural Acidities.

30. §. Dried Gall with Spirit of Nitre, for some time, is still: but at length it makes a considerable Bullition and froth. The reason why it is so long before it begins, is because the salt, (as was observed of some other Parts) is locked up in so great a quantity of Oyl. The abundance whereof is manifest, not only from Destillation, but also from hence, In that the dried Powder, in lying by, incorporateth all together into one body, as Mirrh, and some other softer and oily Gums are used to do.

31. §. Extract of Urine, with spirit of Nitre, makes a Bullition with some Effervescence, which continues for a considerable time; and at last it huffs up with great bubbles. The Bullition begins presently: the salt being copious, and the Oyl but little.

32. §. The same Extract of Urine makes a considerable Bullition and froth, not only with Spirit of Nitre, but even with Oyl of Vitriol. Hence the salt of Urine is more alkaline than that in most of the afore-said Contents.

Contents. From this and some of the following *Experiments*, it also appears, That the salt which concurs to the generation of Gravel or of a Stone in the Kidneys or Bladder, is of a very different nature from the Salt of Urine.

33. §. Next for Dungs. And first, dried Goats-dung makes with Spirit of Nitre, a small Bullition, but no elevation. That of Mice the like. And that of Cows. So that of all I have tryed, these three stir the least.

34. §. Goose-dung, with Spirit of Nitre, makes a very small Bullition and some elevation. But it requires time. Oyl of Vitriol stirs it not.

35. §. Album Græcum, with Spirit of Nitre, besides innumerable small bubbles, rises up with some great ones, exactly resembling the huffing up of Yest or Barm. Also with Oyl of Vitriol it maketh some little froth, but slowly. So that it should seem, that the Bones are a little opened by some acid Menstruum in the Dogs stomach (as the body of Steel is in its preparation with Sulphur) whereby it becomes a good mild Topick in Quinzies.

36. §. Hens dung, with Spirit of Nitre, makes a very great bullition and huff: greater and quicker, than any of the rest above-named.

37. §. But of all I have tryed, Pigeons dung, with the same Spirit, maketh the greatest and the quickest Effervescence and huff; and that not without steams. Yet neither the same Dung, nor that of Hens, is moved in the least with Oyl of Vitriol. The Cause of so great an Effervescence in these, more than in the rest, is that white part which is here mixed in a great quantity with the Dung. Which white part, descendeth not from the Stomach, but is an Excrement separated from the Blood (as the Urine in other Animals) by a peculiar Organ, which evacuates it into the *Intestinum rectum*; whence, together with the Stercus it is excluded. Hence it is evident, That in the said white part of Hens, and especially Pigeons dung, is contained a great quantity of a volatile Alkaly.

38. §. I proceed to Salts. And first Salt of Blood and Urine both make a more durable Effervescence with Acids, than doth Salt of Wormwood, or Salt of Fern. Hence the former are more alkaline, than the latter.

39. §. Again, though divers other Animal Salts will not stir with Spirit of Salt, or with Oyl of Sulphur or Vitriol; yet the Salt of Blood will make an Effervescence with all kinds of Acids. Whence it is further argued to be highly alkaline, and very proper for the correction, of all sorts of preternatural Acids in the body. There is little doubt, but that Spirit of Harts-horn will do the like.

40. §. The Gravel which is precipitated out of Urine; with Oyl of Vitriol makes no bullition in the least. Nor with strong Spirit of Salt. But with Spirit of Nitre, it makes a very great one, with Effervescence and steams. From hence it appears, That there is much difference to be made in the use of acid Diureticks, Nephriticks, &c.

41. §. And that I may not altogether omit to mention, what may be so much for the good of mankind, I do here declare, That for preventing (I say not, the breaking, but preventing) the generation of the Stone, either in the Kidneys, or in the Bladder, there are not bet-

ter Medicines in the world, than some certain Preparations of Nitre, duly administred. Whoever shall think that any kind of acid, as Oyl of Sulphur, Oyl of Vitriol, Spirit of Salt, or the like, will have the same effects, will find themselves much deceived in their practice.

42. §. I conclude with Stones. And first, Spirit of Nitre droped upon a Stone of the Kidneys or Bladder, produceth the very same effect, as upon the Gravel in Urine. That is to say, it makes it boil and buff up, until at length it is perfectly dissolved into a soft Pulp; which neither Oyl of Sulphur, nor Oyl of Vitriol, nor Spirit of Salt will do; nor give the least touch towards its dissolution. This confirms what I said before of the use of Nitre and Nitrous Spirits, if duly prepared and administred, above any other Acids, against the breeding of the Stone.

43. §. Pearls, with any Acid, make the like Effervescence, as do Oyster-shells. But Magistery of Pearls, as usually prepared, stirs not at all, with any Alkaly or Acid. So that as to the effect frequently intended by it, it is very insignificant; as of that of Corals hath been said.

44. §. Crabs Eyes, with any Acid, make an Effervescence, almost as quick as that of Oyster-shells.

45. Crabs Eyes likewise calcined, make a stronger Effervescence, than when uncalcined. So that these, as well as Shells, contain a fixed Alkaly.

46. §. The Stones in Whittings heads make a strong Effervescence like that of Oyster-shells.

47. §. Stone of humane Gall, stirs not with Oyl of Vitriol. But with Spirit of Nitre maketh a little bullition just upon mixing, and after a considerable time, a little froth. Much less than what was observed before of the Gall it self. So that it seemeth to be generated of the Gall coagulated by some Acid, which hath already refracted the Alkaly wherewith the Gall abounds. This confirms the use of those Medicines in the Jaundies, or any other bordering Disease, which destroys those Acidities by which the Gall is curdled or coagulated, and so rendred more difficulty separable into the Guts.

48. §. Since the first publishing of these Observations, Mr. William Matthews an Apothecary in Ledbury, sent me part, as I take it, of a Stomach-stone, as big as a Walnut of the largest Size, voided by a woman about 82 years of age, sometime after an Autumn Fever. It consisteth of the same Striæ, as the Bezoar Stone; and maketh some Bullition with Spirit of Nitre.

49. §. Bezoar, neither the Western nor the Eastern, doth stir at all with Oyl of Vitriol.

50. §. Western Bezoar, with Spirit of Nitre, makes a very little thin froth, and that's all; and that it doth very slowly. But Oriental Bezoar, with Spirit of Nitre, after some time, maketh a very great Effervescence, froth, elevation, noise, and steams (as if you poured Oyl of Vitriol upon Salt of Tartar) till it be wholly dissolved by the affused Spirit, and turned into almost a blood-red. Hence it may seem to be no mean Remedy against such fretting and venenate acids, as oftentimes in Fevers, and other Distempers, lye about the Stomach, and are thence frequently translated to the Heart, Brain, Nerves, and other parts. The difference likewise betwixt the Western and the Eastern Bezoar, is so great, that in any case of danger, and where the Bezoar is relyed upon, it is an unpardonable

able fault, for the *Apothecary*, or any *Person*, to substitute the one for the other: unless he will take *ten times* as much, or *ten times* as little of the one, as he would have done of the other: if that will serve turn.

51. §. The *Stones* already mentioned, (except the great *Stomach-stone*) are ordinarily generated in the bodies of *Animals*. I have one Instance more of some other *Stones* which are extraordinary. In the City of *Hereford* lives a *Maid*, who often voids these *Stones*; and in the space of some years last past, hath voided several pounds, of several *Colours* and *Sizes*, not only *per vias urinarias*, but also by vomit, and by stool: The first mention made to me of them, was by *Mr. Diggs*, a worthy *Gentleman* of that City, as a thing that was there much wondred at. And some of them, upon my desire, were sent me by *Mr. Wellington*, an *Apothecary* in the same place. I have tryed what several acid *Menstruums* will work upon them; and find, That with *Oyl of Vitriol*, and especially with *Spirit of Nitre* the great ones make a very quick and conspicuous *Effervescence*. But the small ones, neither the white, nor the grey, make any *Bullition* in the least: for in truth, they are no other but little *Pebbles* and *Grit-stones*.

52. §. This being considerd, and the various *colours* and *mixture* of any one of the great *Stones*, being well observed; it seemeth plain, That although she be somewhat old (above thirty years) yet may she have a kind of *μαλακία*, or diseased *Appetite* to *Stones*, *Bones*, *Wood-ashes*, *Tobacco-Pipes*, *Chalk*, and such like things; which sometimes swallowing in little *lumps*, sometimes grossly, or finely ground betwixt her teeth; they are in her *Stomach* and *Bowels*, more or fewer of them, cemented together, either with a *pituitous*, *bilious*, or some other more or less *glutinous substance*. And that by virtue also of the said *Cement*, or any of the said, or other like *alkalizate Bodies*, the greater *Stones*, which consist of those partly, do make an *Effervescence* with acid *Liquors*. Thus far of *Instances* upon the parts of *Animals*. I shall close with some *Corollaries* deduced from the whole.

53. §. And first, since we find, that amongst all the *Menstruums* we have made use of, *Spirit of Nitre*, or any very *Nitrous spirit*, is the most *universal dissolver* of all kinds of *Animal Bodies*; the best *dissolver* of many others both *Vegetable* and *Mineral*, and the only *dissolver* of some: Hence it is probable, That the great *stomachick Menstruum*, which either *dissolves*, or *opens* almost all *Bodies* which come into the *Stomach*, is a kind of *Nitrous Spirit*.

54. §. Again, *Spirit of Nitre* being a *subalkaline Acid*, and working more evidently upon *Animal bodies*, than other *simpler Acids* do, which yet are as *strong*; It hence follows, That most of the *Salts* of *Animals* are *subacid Alkalies*. How far this *conclusion* may further instruct us, I shall have occasion to shew in another *Discourse*.

55. §. Lastly, there being so many, say twenty or thirty *degrees*, from the *slowest* to the most *vehement*, in the *Bullition* of *mixed Bodies*; it seemeth, That *Fermentation* it self, as to the *formal notion* of it, is nothing

nothing else: or that from the common *Luētation* of *mixed Bodies* whereof we have now been speaking, it differs not *in specie*, but only in the manner of its *causation*, and in *degree*: the *Aer*, or some certain *Menstruum* lodged therein, being of no greater *strength*, than to produce a *Bullition* or *Luētation* of that *low* and *soft degree*, which we call *Fermentation*.

56. §. I have thus endeavoured to prove, by various *Instances*, how instructive this most easie, plain and simple *Method* in the *Mixture* of *Bodies*, may become to us: and that meerly by observing the *Luētations* which thence arise betwixt them. How much more then, if a diligent remarque be made of all those various *Colours*, *Smells*, *Tastes*, *Consistencies*, and other *Mutations* thereupon emergent?

A. N

E S S A Y

O F T H E

Various Proportions

Wherein the

LIXIVIAL SALT

Is found in

P L A N T S.

Read before the *Royal Society*, March, 1676.

C H A P. I.

*Of the QUANTITIES afforded by several Plants
calcined in gross.*



T is the part of a *Physician*, knowingly and artificially to use and govern *Nature*. And therefore by every likely *Method*, to inspect the *State* and *Properties* of all sorts of *Bodies*. One *Method*, is that I have taken in the foregoing *Experiments*; *sc.* by mixing them with several *Mestruums* or *Liquors*: whereby we may be assisted to judge, both of the *Kinds* and the *Proportions* of *Principles* in any *Body*; and of the manner of their *Mixture* in the same.

Another is by *Calcining* them; or, as it were, by *mixing* them with the *Fire*, a potent and almost universal *Menstruum*. I shall here only set down some *Tryals* for an *Essay*, upon *Plants*; chiefly noting, The different *Proportions* of their *Lixivial Salts*. Of these *Tryals*, some were

were made upon the whole *Plant*, or some *Portion* of it wherein several *Parts* are mixed together: And others, upon some one *Part* of a *Plant* distinct from the rest. All of them answering to such *Queries*, as may seem proper to be proposed.

Query 1. As first, *Whether Trees or Herbs and Bushes, quantity for quantity & cæteris paribus, yeild the most Lixivial Salt?*

For this I took *Ash-Barque* and *Rosemary* of each ℥j . The latter yielded 5 *Scruples*; the former but 32 *Grains*; which is three times less. I took also the same quantity of the *Barque* of *Black-Thorn*, and of *Agrimony*. The latter yielded 5 *Scruples* and 6 *Grains*; the former, not above 1 *Scruple* and 5 *Grains*; which is four times less.

Although the *Barque* of a *Tree* be compounded of *Pithy* and *Lignous Parts*; yet to answer the *Query* exactly, the *Wood* of these *Trees* should be taken with the *Barque*, that there may be some portion of every *Part* of the *Tree*, as well as of the *Herb*.

But thus far the *Experiment* is *conclusive*, That the same quantity of *Lixivial Salt*, doth not always follow the same *Generical Taste*. For the *Barque* of *Ash* and *Rosemary*, are both equally *Bitter*; and the *Barque* of *Black-Thorn* and *Agrimony* are both *Astringent* and *Bitter*.

Quer. 2. *Whether any Plant growing in a Garden or the Field, doth not yield a lesser quantity of Lixivial Salt, than another of the same kindred growing on the Sea-Coast; and with what difference?*

For this, I took *Garden* and *Sea-Scurvygrafs*, of each ℥j . The former yields 2 *Drachms* and 1 *Scruple*; the latter, being well washed, 9 *Drachms*, which is more than 4 times as much. The like may be tryed upon others.

Quer. 3. *Whether the same Specifick Plant affords more Lixivial Salt, being only dried, and then calcin'd, or after it hath first been distilled, it is then dried and calcin'd?*

For this, was taken ℥j of *Mint* only dried and then calcin'd; and another first distilled. The former yielded $\frac{1}{2}$ an *Ounce* and $\frac{1}{2}$ a *Drachm* of *Salt*; the latter, 5 *Drachms* and a *Scruple*; which is almost $\frac{1}{3}$ th more. This also should be tryed on other *Plants*.

Quer. 4. *How far the proportion follows the different Tasts of Plants?* The first *Experiment*, relates to the same *Tast* in several *Plants*; this, to several *Tasts*. And so,

Of *Majorane*, which is *Aromatick*, ℥j affords but one *Scruple* of *Lixivial Salt*; which is but the 384th part of the whole pound.

Of *Oak-Barque* which is *Astringent*, ℥j yields $\frac{1}{2}$ a *Drachm* of *Salt*; or the 256th part of the whole.

Of *Liquirish*, which is sweet, ℥j yields about the same quantity. But *Anise Seeds* ℥j yields 2 *Scruples* or a 192^d part.

Of *Sorrel*, which is sower, ℥j yields one *Drachm*, or the 128th part.

Of *Garden Scurvygrafs*, which is *Hot*, ℥j yields 2 *Drachms* and $\frac{1}{2}$ a *Scruple*; or the 59th part.

Of *Mint*, which is *Hot* and *Bitter*, ℥j yields 5 *Drachms* and a *Scruple*, or the 24th part.

Of *Sea Scurvygrafs*, which is *Salt*, ℥j yields 9 *Drachms* and a *Scruple* or 28 *Scruples*; which is near $\frac{1}{7}$ th part of the whole. A greater proportion of *Salt*, than in any other *Plant* upon which I have hitherto made *Tryal*: Or even in *Tartar* it self. Yet is it not a *Marine*, but true *Lixivial Salt*: as is evident, both from its *Taste*; and in that it maketh

maketh an *Efferescence* with *Spirit of Salt*; which *Sea-Salt* will not do.

For the Experiment to be fully adequate to the *Query*; the *Tryals* should be made, either all on *Trees*, or all on *Herbs*; all on *Roots*, or all on *Stalks*, &c. Yet thus much is evident, That *Sorrel* yields Thrice as much as *Majorane*; *Sea-Scurvygrass*, Eight and Twenty times as much: *Mint*, Five times as much as *Sorrel*; and Sixteentimes as much *Majorane*, &c.

Quer. 5. *How far the Proportion follows the Faculties of Plants?* And so, it appears, that

Majorane, a *Cephalick*, hath a greater *Proportion* of *Volatile Parts*, than any of the *Plants* above mentioned, and so far, is more agreeable to the *Animal Spirits*, and *Genus Nervosum*.

Agrimony, (a) an *Aperient*, yields above Five times as much *Lixivial* (a) *Quer. 1.* *Salt*, as *Majorane*. Yet much less than many other opening *Plants* which are stronger.

Mugwort (fbj) yields two *Drachms* and two *Scruples*; or above half as much more as *Agrimony*. So that this *Plant*, though it hath no considerable *Taste*, and in that respect promiseth but little; yet yielding a good quantity of *Lixivial Salt*, seems no contemptible *Medicine* to subdue those *Acidities* which either by causing *Obstructions*, or immoderate *Fermentations*, frequently disorder the *Female Sex*.

Mint, yieldeth still a greater quantity; and is therefore, partly for the same cause so excellent a *Stomachick*: And *Rosemary*, (b) which is appropriated both to the *Head* and *Stomach*, yieldeth a middle quantity of *Salt*; more than the chief *Cephalicks*, and less than the chief *Stomachicks*. (b) *Quer. 1.*

Common Mallow (fbj) yields 5 *Drachms* and 2 *Scruples*. *i. e.* the 23^d part of the whole. So that this *Plant*, though of a very mild *Taste*, yet yields more *Salt* than *Mint* it self a *Bitter Plant*. Whereby it no longer seems strange, that a *Plant* of so soft a *Taste*, should be very *Diuretick*, and so evidently affect the *Reins*.

Rhubarb (2 *Ounces*) yieldeth scarce any fixed *Salt*, so far as can be judged by the *Taste* of the *Ashes*, not more than a *Grain* or two. So that its *Salt* is, in a manner, wholly *volatile*; and thereby apter to operate upon the *Bilious* parts of the *Blood*; which contain a far greater proportion of *Volatile Salt*, than do the *Serous*.

Of the *Caput Mortuum* or meer *Earth*, it is observable, that it was near $\frac{1}{2}$ an *Ounce* or $\frac{1}{4}$ th part of the whole; Which is almost Six times as much as the *Caput Mortuum* of *Common Dock*: and much more than that of any other *Root* I have yet calcin'd. Whereby it seemeth probable that *Rhubarb* looseth much of its *Volatile Part*, and therefore of its *Virtue*, before it comes to our *Shops*.


Sena (fbj) yields 4 *Scruples* and $\frac{1}{2}$ of *Salt*; or the 85th part.

Falap (fbj) yields but one *Drachm* and 15 *Grains*, or 102^d part.

Colocynthis (fbj) of the *Pulp* yields an *Ounce* and half of *Caput Mortuum*, which is almost all *Salt*. Yet allow half an *Ounce* of the *Salt*, and *Earth* to be wasted in filtering &c. the remaining *Ounce* is no less than $\frac{1}{6}$ th part of the whole. Which is more than in any of the above named *Plants*, except the *Sea-Scurvygrass*.

C H A P. II.

Of the QUANTITIES afforded by the Parts of several Plants distinctly calcin'd.



SHALL next set down some Tryals, upon one Part of a Plant, as well Organick, as Content, separated from the rest; in answer to these supposed Queries.

Quer. 1. What Proportion doth the Lixivial Salt of the Pith or Pithy Part of a Plant, bear to that of the Fibrous, or of the Woody Part? Or whether is there a Fixed Salt always found in either of them? A

sufficient Answer to which, must be built upon many Tryals. At present I shall mention only Two; one upon Starch, answerable to the Pithy Parts; the other upon Flax, consisting almost wholly of the Nervous or Tomy Fibres: of the Volatile parts whereof, chiefly, I have

§. 50, 51,
52.

given some account in the foregoing Idea. Of Starch, ℥ij yieldeth about ℥ $\frac{1}{4}$ not of Ashes, but of Black Coal. For though it be exposed in a Calcining Furnace to a vehement fire, for 5 or 6 hours, which is longer then will serve to calcine most Bodies: yet would it not in the least part, be reduced to Ashes; but to the last continued (though the fierceness of the Fire consumed part of it) as black, as when it was first burnt. So strangely was the remaining part of the Sulphur fixed to the Earth; that in flying away, it did volatilize and carry that away with it. In this Coal or Cinder, there is not the least of a Lixivial or other Taste. And although, upon Tryal I find, That the Pith of many Plants, as of a Cabbage Stalk, will yield some quantity of Lixivial Salt; yet it is probably, that generally, it yields less than the Wood.

Of Flax, ℥ij yields not above 50 Grains of Caput Mortuum or white Ashes, which are Salt. According to vulgar conceit, it would seem to be a very dry Body: yet of 153 parts, 152 are volatile, and being distilled would have been collected into Liquor. Hence also appears the great and unexpected Variety in the Proportion of the Earthy Parts, as well as the other Principles of Bodies. Or else, that there are divers kinds of Earths, even in Plants, of which, as well as of Salts &c. some are volatile. For of ℥ij of this Plant, there remaineth fixed but 50 Grains: whereas of ℥ij of Rhubarb, there will remain near 1920 Grains, i. e. 88 times as much as the former.

Quer. 2. In what proportion is the Lixivial Salt found in the Gums of Plants? and whether is it yielded, more or less, by all? For answer to which, I caused the Eleven following, of each two Ounces, to be calcin'd, and so observed,

That Common Rosin, yields but one Grain and $\frac{1}{2}$ of Caput Mortuum. So that ℥ij will yield but 12 Grains. In this Caput Mort. there is not the least particle of Salt, it being altogether insipid.

Mastick yields gr. 12 of Cap. Mort. But not the least part of salt. Of this Rosin, it is observable; That being set, in a Crucible, within the

the fire, before it comes to have thick fumes, it boyls up with a very great foame or froath; and is the only *Gum* or *Rosin* (of the Eleven) that hath this property. So that I suspect, there is a great quantity of some kind of *volatile Spirit*, which then flies away; and so, in breaking through the *Oily parts*, huffs them up to so great a froath.

Olibanum yields half a Drachm of *Caput Mortuum*. But it is to be noted, That the weight is encreased by certain little *Spar-Stones*, which in the burning of several parcels, I always found mixed with this *Gumm*. These being picked clean out, the *Cap. Mort.* weigheth not much more than that of *Mastick*. And is in like manner insipid, when the said *Stones* are picked out.

From hence it appears, how proper these *Gums* are for the *Concoction* of *Salt Rheums*; according to what I have formerly suggested from another Experiment.

It may also be noted, that *Rosin* and *Mastick*, seem to be more purely *Acidoleous Gums*; not only from their consistence which is uniform; and their *Smell*, which is less strong and more pleasant: but also from the *Acid Liquor* they yield by *Distillation*; and in that the young *Leaves* of *Fir*, and especially of *Pine*, are sower; and tis probable that those of *Mastick* are so likewise. Whereby these, and other like *Gums* are more especially fitted for the abovesaid purpose. But *Olibanum* seems, besides its *Acidity*, to contain some *Volatile Alkaly*, and so to be an *Acid-Alkaline Gum*. For as it hath a stronger *Smell* than the former, so a hotter *Taste*; both the ordinary effects of an *Alkaline Sulphur*. And being infused in several *Menstruums*, appears to consist of two Bodies, one of them more *Resinous* than the other. Of which, it is probable, that the one is made by the *Acid parts* as the other by the *Alkaline*. Whereby it is very well adapted in some *Cases*, as in a *Pleuresie*, for removing the *Coagulations* of the *Blood*, or its disposition thereunto.

Asa fetida yields no less than half its weight or an Ounce of *Caput Mort.* that is 8 times as much as that of the other *Gumms*, and 48 times as much as that of some of them. Yet doth it not contain one grain of *Salt*, so far as can be judged by its *Tast*. Yet the *Strength* and *Loathsomeness* of the *Smell* and *Tast* of the *Gumm* do argue it to be highly impregnated with some kind of *Volatile Alkaly* proper to arrest those offensive *Vapours* (to use the vulgar word) which flying, either by the *Blood* or *Nerves*, from part to part, do often prove so troublesome.

Gum Arabick yields one Scruple of *Cap. Mort.* whereof, by the *Taste*, about $\frac{1}{3}$ part is *Salt*.

Euphorbium yields one *Drachm* of *Caput Mort.* of which, by the strength of the *Taste*, two Scruples seem to be *Salt*. Which confirms a former conjecture (a) of its being an *Alkaline Gum*.

Myrrh also yields a *Drachm* of *Cap. Mort.* and at least two Scruples of *Salt*. Of the Eleven, these two *Gums* have the greatest quantity of a fixed *Alkaly*.

Opium yields half a *Drachm* of *Cap. Mort.* whereof the one half is *Salt*.

Aloe yields a *Drachm* of *Cap. Mort.* containing about one Scruple of *Salt*.

Scammony yields Two Scruples of *Cap. Mort.* of which, about half a Scruple is *Salt*.

Gutta Gamba yields but half a Scruple of *Cap. Mort.* of which four or five Grains are *salt*.

So that considering the *Dose* of any *Cathartick Gumm*, the quantity of the *Fixed Alkaly*, is extream small with respect to the *Volatile parts*: In which, therefore, its *Crihartick Power* doth chiefly reside.

Yet none of the *Cathartick Gumms* are without some portion, more or less, of a *Fixed Alkaly*; though some of the rest are. Which seemeth to prove, That the *Fixed* it self, hath some Interest in the business of *Purgation*: as by being a *Clog* to the *Volatile*, and so preventing its being deleterious; or some other way. But the manner of their *Operation* will better be understood, when the *Volatile Parts* have likewise been examined.

It may also be of good import, to know, what different quantities of *Salt*, are afforded by the *Tartars* of all sorts of *Wines* Whereby, partly, as well as by the quantity of the *Tartar*, we may be enabled the better to judge of the *Nature* of *Wines*.

A

DISCOURSE

Concerning the

ESSENTIAL and MARINE

Salts of Plants.

Read before the *Royal Society*, December 21st
1676.

CHAP. I.

In which is shewed the way of making both an ESSENTIAL and a MARINE Salt, out of the LIXIVIAL Salt of a Plant.



SOMETIME since, I took the boldness to present my thoughts to this Honourable and Learned Body in a *Discourse* concerning *Mixture*. Wherein I have endeavoured to lay such a *Foundation*, as might hereafter reduce the *Doctrine* hereof to *Experience* and *Practise*; and to demonstrate, the *Power* and *Use* of *Artificial Mixture*. And in further proof of what is therein asserted, I have since made a continuation of *Experiments* upon the same Subject, in *Two Methods*. One in the *Mixture* of several *Menstruums*; both *Acid* and *Alkaline*, with all Sorts of *Bodies*. The Other, by calcining them, or, as it were, *mixing* them with the *Fire*.

2. §. I shall now proceed to a Third, which is, the *mixing* them with the *Aer* or exposing them to it; another of Nature's grand *Menstruums*; which goes sometimes further than the *Fire* it self, in the *dissolution*

Discourse of Nature, in producing a *Marine* or *Muriatick Salt* out of the *Lixivial* of *Mixture Salt* of a *Plant*. But some Learned Persons then present, seeming to doubt of the Experiment; I thought it requisite to prosecute the same a little further; that so, if possible, it might become clear and unquestionable.

2.

And because that *Method* was imperfect, and required half a year, or a longer time: I bethought my self of an other way; which proved far better, and more expedite. And which, withall, afforded me, not only a true *Marine salt*, out of the *Lixivial salt* of a *Plant*; but also another kind of *salt*, different from them both: which may not be improperly called, an *Essential salt* or *Nitre of Plants*. The History or manner of the production of them both, is as follows.

3. §. December 15. 1675, I took about half a pound of a strong *Solution* of the *Lixivial Salt* of *Firne*: and pouring it into an *Earthen Pan*, well glazed, broad and shallow, exposed it therein to the open *Aer*, in a Chamber Window, to evaporate of it self.

Tab. 83.

4. §. This *Solution* or *Lee*, although it was very clear before, and having stood corked up in a bottle many days, had no sedement: yet standing now in the open *Aer*, within the space of 4 or 5 days, it began to let fall a very *white Sedement*, like fine *Chalk*; which encreased daily for 8 or 10 days; amounting at last to about half a Drachm of white, light and meer *Earth*, altogether insipid, and when it was well washed, stirring not upon the *Affusion* of *Acids*.

5. §. Within the space of a day or two after this *white Sedement* began to fall to the bottom; there was also gatherd on the top, a kind of soft *Scum* or *Cremor*, wherewith the *Solution* was covered all over.

6. §. Within 8 or 9 days after the first exposing of the *Liquor*, or 2 or 3 days after the gathering of the *Cremor*; that *Salt*, which I take leave to call, an *Essential salt* of *Plants*, began to appear; shooting into several little *Crystals*. These *Crystals*, as they grew bigger, began to sink, and at last fell down to the bottom of the *Pan*.

Tab. 83.

7. §. Upon their first generation or shooting, the said *Cremor* presently breaks, leaving a bare space round about each *Crystal*; and upon the bounds of every space is indented; the space growing bigger and bigger together with the *Crystal* in the Centre. And so, by that time the *Crystals* are grown to a considerable number and bigness, the *Cremor* vanishes away, the several *Circles* or bare places breaking at last one into another all over the *Surface* of the *Lee*. After which, it never comes again.

8. §. From whence it seemeth, That the several *Circles* or bare *Spaces* about the *Crystals*, are made for the more free admision of the *Aer*, requisite to their *Generation*. For as there is no *Crystal* begins to be formed before there is a breach made in the *Cremor*: so that breach is enlarged together with the *Crystal*. So that as the falling of the *Sedement* and the gathering of the *Cremor*, sheweth that the *Aer*, as a *Menstruum* separates some part from the *Lee*: so the breaking of the *Cremor* afterwards, that as a *Vehicle*, it brings something to it: both in order to the *Generation* of the *Crystals*. Nature taking a *Method* for the *Generation* of simpler Bodies, as well as of those which are *Compounded* and *Organical*.

9. §. The *Figure* of these *Crystals* is angular and oblong, most of them about the fifth, sixth or seventh of an Inch; but none of them very regular. Yet we are not hence to conclude, but that with the help of some Circumstances which might be wanting in the shooting of these; some portion of regular ones may be obtained from this, as well as other *Lixivial Salts* hereafter mentioned.

10. §. They are somewhat transparent, and of a dark *Ambar Color*, or like that of brown *Sugar-Candy*. Of a quite different *Taste* from that of the *Solution* or *Lee* out of which they are bred; being not at all *Lixivial*, but very weak and mild; not *Salt*, but *Bitter* in a good degree.

11. §. It is also observable, that *Alkaline* and *Acid Salts* being both poured severally upon these *Crystals*, they stir not, nor are any way affected with either of them. So that these *Crystals* are no sort of *Tartar*, or *Tartareous Salt*. As is plain, from the manner of their *Generation*; *Tartar* being still bred in close *Vessels*; these never, but by exposing the *Liquor* to the *Aer*. As also from their *Taste*, being not sower, in the least, but bitter. And in that *Tartar* will make a *Bullition* with *Alkaline Salts*, which these will not do. Upon which accounts it appears, that they are a *Salt* different in Nature from all other *Salts* hitherto known, or a new *Species* added to the *Inventory* of *Nature*.

12. §. These *Crystals* within the space of about a fortnight after their first *Generation*, did also cease to shoot any more, but only increased a little in their *Bulk*. After which time, I dayly expected to see the production also of a true *Marine Salt*. And about two months after the said *Essential Crystals* had done shooting, and not before, this also began to shoot, in many small *Crystals*, and at the top of the *Solution*, as the other did, still falling to the bottom as they grew bigger.

13. §. The *Size* of most of them was near that of the *Flakes* or *Grains* of *Bay-Salt*. The *Colour* of some of them *white*, of others *transparent*; and of others *white* in the *Centre*, with transparent *Edges*; as is also usual in the *Crystals* of *Common Salt*.

14. §. The *Figure* of most is a perfect *square*, and of very many coming near to a *Cube*; which is also the *Figure* of *Common Salt*, and seldome an exact *Cube*. An exact *Cube*, being the constant property of no *Marine Salt*, that I know of, except that of the *Dead Sea*. Divers *Tab. 83* of them were also raised as it were by several steps from a deep *Centre* to the *Top*: as is often seen in the common shooting of *Common salt*; and not in any other. Their *Taste* is neither *Lixivial*, as that of the *Solution* out of which they shoot; nor bitterish, as that of the *Essential Crystals*; nor sowerish, as that of *Tartar*; but the perfect *Taste* of *Common Salt*.

15. §. It is also to be noted, That if *Oyl* of *Vitriol*, and some other strong *Acids*, be poured upon this artificial *sea-salt*, they make an *Effervescence* together: but if *Spirit* of *Salt* or *Spirit* of *Nitre* either be poured on it, though it be never so strong it stirreth it not. In both which, and all the formentioned respects, it answers to the *Properties* of a *Marine* or *Common Salt*, which no other *Salt* doth. I conclude it therefore to be a true *Marine Salt* produced by *Art* in the imitation of *Nature*.

C H A P. II.

Wherein is shewed, That the said *ESSENTIAL* and *MARINE* Salts of Plants are both of different Sorts.



HAVING made the Experiment, that both an *Essential* and *Marine Salt* may be produced out of the *Lixivial Salt* of a *Plant*. I thought it probable, that neither the one nor the other, was always the same, but that as they had their general properties which made them to be of two general kinds; so they might have some special property, for the distinguishing of each kind into several Sorts. And withall, that in a warmer season, than before taken, the *Tryal* hereof might be finished in a shorter time.

2. §. For the making of which, I conceived it requisite to remove an Opinion which seemed to lye in my way; *sc.* That there is little or no difference between the several *Lixivial Salts* of *Plants*, as some Learned men have thought. But either there is a difference, or not: if not, it should be proved: and if there be, it should then be justly stated, what that difference is. For the doing of which, I chose this *Method*. I took an equal quantity of the whitest and purest *Salts* of divers *Plants*, all made by an equal degree of *Calcination*; and dissolved them all severally in an equal quantity of water. And pouring likewise an equal quantity, as about 10 or 12 drops of each into a spoon, I tasted them severally. Whereby it was very evident, that they were not all of one *Tast*, but of very different ones, both as to strength and kind: and therefore different in *Nature* also. The *Salts* I made tryal of were those of *Sorrel*, *Anise*, *Wormwood*, *Mallow*, *Ash*, *Tartar* and others: and upon half a Drachm of each I poured ʒijss of water. The *Solutions* are here present to be tasted. By which the differences will easily be observed, and particularly that the *Salt* of *Wormwood* or *Scurvygrafs*, is almost as strong again as the *Salt* of *Anise*, or *Sorrel*: and that the *Salt* of *Ash* is above twice as strong, and that of *Tartar* above thrice as strong, as that of *Sorrel*, and almost thrice as strong as that of *Wormwood* or *Scurvygrafs*. So that he who shall give half a Scruple, suppose of *Salt* of *Tartar*; instead of half a Scruple of *Salt* of *Wormwood*, or other like *Salt*; he may as well give a Scruple of *Rosin* of *Jalap*, for a Scruple of the powder, or almost three Drachms of *Rhubarb*, or other like *Purge*, instead of one. And the like is to be said of other *Lixivial Salts* in their degrees.

3. §. Having observed thus much, I proceeded to repeat the former Experiment, with some of the aforesaid, and some other *Vegetable Salts*, the best calcin'd, and the purest, that could be made for this purpose, being these Six *Salts*, *sc.* of *Rosemary*, *Garden Scurvygrafs*,
Black

Black Thorn, Common Wormwood, Ash, and Tartar. All which dissolved severally in fair water, I exposed in a Chamber window, and not in Winter, as before, but in the heat of Summer, *sc.* on the 19 of July, to evaporate of themselves.

4. §. The Effect was, That the third day after their being exposed, the *Essential Crystals* began to shoot in three of the *Solutions*, *sc.* in that of *Rosemary*, of *Garden Scurvygrass*, and of *Black Thorn*. On the fourth day, in that of *Wormwood*. On the fifth day, in that of *Ash*. In that of *Tartar*, not at all.

5. §. These *Essential Crystals* began, in all, to shoot at the top, and then to fall to the bottome; as in the Experiment before. But as there was very little of the *white Sedement* before mentioned, that preceded; So no *Scum* or *Cremor* at all. Which although a more perfect *Calcination*, it seems, did here almost prevent; yet did not in the least destroy the aforesaid *Essential Salt*, but rather make way for its more speedy and copious *Production*: exhibiting likewise a distinct *Species* in several of the *Solutions*.

6. §. For first, the *Crystals* of *Rosemary* (the largest of them) were about the bigness of a *Rice-Corn*. In *Figure* almost like a *Tip-Cat*, which Boys play with, split down the midle. Each *Tip* being cut into 5 sides all ending in a poynt: the midle part divided into 7, all drawn by parallel *Lines*; the topmost with the lowermost but one, on each side, being three exact *Squares*. Tab. 83.

7. §. The *Crystals* of *Black Thorn* are most of them poynted with just six sides of *Equal Measure*: very like to the shooting of true *Crystal* it self. From the topmost of which six Sides, a *Line* being drawn out, runs parallel to a broad *Base*, whereon each *Crystal* stands. So that they are in some sort of a *Rhomboid Figure*. Tab. 83.

8. §. The *Crystals* of *Scurvygrass* have also a very elegant and regular *Figure*, which is in a manner compounded of the two former now described. But they are nothing near so bigg, the largest of them, being no bigger than a *Grain* of that which we call *Pearl Barley*.

9. §. The *Crystals* of *Wormwood* have also very many of them a regular *Figure*; but quite different from that of the *Crystals* before mentioned; each *Crystal* being a little *Cylinder*, saving that it is constantly somewhat smaller at one end, than the other: as it were one half of a *Rowling-pin*. And not evenly *Circular*, but cut out by Six Sides of equal *Measure*: almost as in the *Crystal* of *Nitre*. So that contrary to what is seen in the forementioned *Crystals*, the ends of these of *Wormwood* are not poynted, but *flat*; and cut at *Right Angles* with the *Sides*. Tab. 83.

10. §. The *Crystals* of *Ash*, though by their properties they appear likewise to be *Essential*; yet are nothing near so regularly figur'd, as all the forementioned.

11. §. The *Colour* also of the said *Crystals* is somewhat different: Those of *Ash* being of a *brown* transparency, almost like those of *Firne*. Those of *Wormwood* being also *brownish*, but *paler*. Those of *Rosemary* and *Scurvygrass* having some little *Tincture*, yet very *clear*. But those of *Black Thorn* without the least *Tincture*, and as *clear* as *Crystal* it self.

12. §. None of these *Essential Crystals* have any hot fiery *Taste*, but are very *mild*, and sensibly *Bitter*; especially, about the *Root* of the *Tongue*: as is also observable of some *Plants* hereafter mentioned, in speaking of the different *Tastes* of *Plants*.

13. §. *Oyl* of *Vitriol* droped upon these *Crystals* doth not affect them in the least: yet droped into the several *Solutions* out of which the *Crystals* are produced, immediately causeth a great *Effervescence*.

14. §. Of the *Solutions* above named, that, of *Salt* of *Tartar* was the 6th. Whereof it is remarquable, That having waited several Months together, I could not observe the least *Essential Salt* to be therein produced in all that time. Whether there be any other *Vegetable Salts*, besides this of *Tartar*, which will not yield the *Essential* above described, I have not yet experimented.

15. §. In the mean time, from the *Premises* it is very probable, that most of them afford more or fewer of the said *Crystals*. In regard they are *Plants* of a very different kind, which I made tryal upon: as *Garden Scurvygrass*, very *Hot*; *Rosemary*, very *Aromatick*. *Wormwood* very *Bitter*; *Black Thorne*, *Astringent* and *Sower*. And it is also plain, That the said *Essential Salts* contained in the *Lixivial*, are not altogether one and the same, but of divers *Sorts*.

16. §. ABOUT 7 or 8 days after the *Essential Crystals* were produced; the *Marine Salt* did also begin to shoot; first in *Rosemary*; quickly after, in *Scurvygrass*; Next, in *Black Thorn* and *Wormwood*, &c. after the space of a week or 10 days more. And in all of them with some difference of *Size* and *Figure*.

17. §. The plainest of all, was that produced out of the *Salt* of *Black Thorn*, consisting for the most part of very small *Crystals*, not above the 15th of an *Inch* square, as also *thin*, shaped like a *Duch Tile* used for *Chimnies*. Many others were very *thick*, and near to a *Cube*. Most of which were a little hollowed in the middle, like a grinding *Marble* or *Salt-Celler*; and the hollow bounded by 4 plain and equal *Sides*, all descending a little towards the *Centre*; and measured by two cross *Lines*, which staid upon the four *Angles* of the *Square*, and so cut one the other at *Right Angles*. Both which *Properties* are likewise usually seen in the *Crystals* of *Common Salt*.

18. §. In *Wormwood*, many of these *Crystals*, besides the plain ones, were figur'd crossways like a *Dagger-Hilt*. Which was sometimes naked, and sometimes inclosed in a square and almost *Cubical Box*. Many others were figur'd into *Sprigs* made up of four chief *Branches* standing crosswise, and those *subbranched*; and all the *Branches* made up of little square *Crystals*, clustered together in that *Figure*. The *Sprigy Figure* of these *Crystals* is not accidental, but hath constantly come after they had been three times dissolved, and the *Solution* exposed to evaporate.

19. §. The *Marine Salt* of *Rosemary* hath also some variety. For besides the plain ones above described, there are some thick *Squares*, which have also a square hollow descending by five, six, or seven narrow steps, towards the *Centre*; being in *Figure*, saving these *Steps*, somewhat like the *Hoper* in a *Mill*.

20. §. Upon a second *Solution* of the same *Salt*, there shoots another sort of square; which is not plain on the edges, as the above-named, but scalloped or florid all round about, not unlike the *Leaves* of some *Plants*.

21. §. The *Crystals* of *Marine Salt* of *Scurvygrafs* are somewhat like to those of *Rosemary* now described.

22. §. As for the *Lixivial Salts* of *Ash* and *Tartar*, though in a Month or Five Weeks Space, they yield some *Crystals* of very clear *Salt*: yet of *Marine Salt* neither of them yieldeth the least particle. So that of these Six *Lixivial Salts*, *sc.* of *Rosemary*, *Scurvygrafs*; *Black Thorn*, *Wormwood*, *Ash* and *Tartar*, all, but that of *Tartar*, yielded an *Essential salt*. And all, but those of *Ash* and *Tartar*, yielded a *Marine*, such as is above described. All which *salts* both *Essential* and *Marine*, together with their *Models*, made of *white Alabaſtre*, I have here ready to be ſeen.

23. §. Of those that yield these *salts*, or either of them, it is further to be noted, That there is a considerable difference in the *Proportion* or *Quantities* which they yield. The *Rosemary* yields store both of *Essential* and *Marine*, but more *Essential*. *Wormwood* and *Scurvygrafs* more *Marine*. *Black Thorn* leſs of Either. The *Ash* no *Marine*, and the *Tartar* neither the *Essential* nor *Marine*, as hath been ſaid.

24. §. From what hath been ſaid, I deduce only at preſent theſe Three *Corollaie*. *Fiſt*, That a *Lixivial Salt*, is not only a compounded Body *sc.* of *Salt*, *Sulphur*, *Aer* and *Earth*; but even a *Compounded Salt*, containing both a *Vegetable Nitre*, and a true *Sea Salt*.

25. §. *Secondly*, That the *Expoſing* of *Bodies*, in the manner above ſhewed, may juſtly be accounted one *Part* of *Chymiſtry* hitherto *Deficient*, and much farther to be improved for the *Discovery* of the *Nature* of *Bodies*. For as *Nature* chiefly compoundeth *Bodies* by *Digeſting* them, and ſo either ſhutting out or keeping in the *Aer*: So ſhe *Diſſolveth* them by *Expoſing*, and ſo neither ſhutting in the *Aer*, nor keeping it out, but leaving it free to come and go; and thereby to bring, and carry off whatſoever is neceſſary for the *Separation* or *Solution* of *Bodies*. For the *Sea* it ſelf (to confine the ſimilitude to our preſent caſe) is but as a *Great Pan*, wherein all kinds of bodies being long expoſed, are throughly reſolved, ultimately yielding from the reſt of their *viſible Principles*, that which we call *Sea Salt*.

26. §. *Laſtly*, if by *Expoſing* and *Diſſolving* we can make one *Salt*; then by *Compounding* and *Digeſting* we may make another, yea any other *Salt*; either a *Fixed* of a *Volatile*, or a *Volatile* of a *Fixed*. That is to ſay, a *Volatile Salt* may be ſo ſeparated from other *Bodies*, as to become *Fixed*; or a *Fixed Salt* may be ſo *mixed* with other *Bodies* as to become *Volatile*. For that any *Salt* ſhould of it ſelf become *Fixed* or *Volatile*, is a *Fixion* not grounded upon *Experiment*.

27. §. As for the *Virtue* of the *Essential Salts* above deſcribed, I believe they will be found upon tryal, not contemptible in ſome *Cafes*. For which amongſt other reaſons, I have been the more punctual

in relating the manner of their *Generation*; that others also may have the opportunity of making proof hereof.

28. §. When I made the *Experiments* for this and the foregoing *Discourse*, not having so good conveniency at home for making the *Salts* I used: I procured them all (except that of *Firne*, which I made my self) to be purposely prepared by Mr. *John Blackstone* a *London Apothecary*, who assured me of his great care herein; and particularly, that he added no *Nitre* to whiten any of the *Salts* with, as is commonly done for that of *Tartar*.

I do declare, That all the *Lixivial Salts* mentioned in this and the foregoing *Discourse* except that of *Firne*, were faithfully prepared by me

John Blackstone.

A
 DISCOURSE
 OF THE
 COLOURS
 OF
 PLANTS.

Read before the *Royal Society*, *May 3. 1677.*

CHAP. I.

Of the COLOURS of Plants in their Natural Estate.



HAVING formerly made some *Observations* of the *Colours* of *Plants*; I shall now crave leave to add some more to them of the like *Nature*. None of which, nor any of the *Conclusions* thence deduced, will, if duly considered, appear contrary to the *Hypothesis* and *Experiments* of *Mr. Boyle*, *Mr. Des Cartes*, *Mr. Hook*, *Mr. Newton*, or any other, concerning *Colours*. As not having respect to the *Colours* of all *Bodies* in general. Nor to the *Body* of *Colour*, which is *Light*; Nor to the formal notion of *Colours* (*ad extra*) as the *Rays* of *Light* are moved or mixed: But to those *Materials*, which are principally necessary to their *Production* in *Plants*. Concerning which, the present *Discourse* shall be reduced to these Three general *Heads*,
scil.

Idea, §. 27.
 and *Anat.*
 of *R. P.* 2.
 §. 65, &c.

2. §. First, Of those several Colours, which appear in *Plants* in their *Natural Estate*.

3. §. Secondly, As they appear upon the *Infusion* of *Plants* into several Sorts of *Liquors*.

4. §. Thirdly, As upon the *Mixture* of those *Infusions*, or of any one of them with some other *Liquor*, or other *Body*.

5. §. As they appear in the *Plants* themselves, it may be observed in the first place, That there is a far less variety in the Colours of *Roots*, than of the other *Parts*: the *Parenchyma* being, within the *Skin*, usually *White*, sometimes *Yellow*, rarely *Red*. The Cause hereof being, for that they are kept, by the *Earth*, from a free and open *Aer*; which concurrerth with the *Juyces* of the several *Parts*, to the *Production* of their several Colours. And therefore the upper parts of *Roots*, when they happen to stand naked above the *Ground*, are often deyed with several Colours: so the tops of *sorrel Roots* will turn *Red*, those of *Mullen*, *Turneps* and *Radishes*, will turn *purple*, and many others *green*. Whereas those parts of the same *Roots* which lie more under *Ground*, are commonly *White*.

6. §. As *Roots* are most commonly *White*; so the *Leaves*, *Green*. Which Colour is so proper to them, that many *Leaves*, as those of *Sage*, the young *sprouts* of *St. Johns-wort*, and others, which are *Redish* when in the *Bud*; upon their full *Growth*, acquire a perfect *Green*.

7. §. The Cause of this Colour, is the *action* of the *Aer*, both from within, and from without the *Plant*, upon the *Juyces* thereof, whereby it strikes them into that Colour.

8. §. By the *Aer* from without, I mean that which surrounds the *Body* of the *Plant*: which is the Cause of its *Greeness*, not meerly as it is contiguous to it, but as it penetrates through the *Pores* of the *Skin*, thereinto; and so mixing with the *Juyces* thereof, plainly deys or strikes them into a *Green*.

9. §. By the *Aer* from within, I mean, that which entring, together with the *Aliment*, at the *Root*, thence ascends by the *Aer-Vessels*, into the *Trunk* and *Leaves*, and is there transfused into all the several *Juyces*, thereby likewise concurring to their *Verdure*. Whence it is, that the *Parts* of *Plants* which lie under *Water*, are *Green*, as well as those which stand above it; because, though the ambient *Aer*, contained in the *Water* be but little, yet the want of it is compensated, by that which ascends from the *Root*.

10. §. And therefore it is observable, that the *Stalks* of *Marsh-Mallow*, and some other *Plants*, being cut transversly, though the *Parenchyma* in the *Barque* be *white*, yet the *Sap-Vessels* which lie within that *Parenchyma*, are as *Green* as the *Skin* it self; scil. because they stand close to the *Aer-Vessels*. The *Parenchyma*, I say, which is intercepted from the *Aer*, without, by the *Skin*; and from the *Aer* within, by the *Sap-Vessels*, is *white*: but the *Skin*, which is exposed to the *Aer* without, and the *Sap-Vessels* which are next neighbours to that within, are both equally *Green*. So likewise if a *Carrot* be plucked up, and suffered to lie sometime in the open *Aer*; that part which standeth in and near the *Centre*, amongst the *Aer-Vessels*, will become *Green* as well as the *Skin*, all the other *Parts* continuing of a *Redish Yellow*, as before. The *Aer* therefore, both from without, and from within the *Plant*, together with the *Juyces* of the *Plant*, are all the concurrent Causes of its *Verdure*.

11. §. BUT how doth the *Aer* concur to the *Greeness* of *Plants*? I answer; Not as it is meerly either *cold* or *dry*, or *moist*, nor yet *quatenus Aer*; but as it is a mixed, and particularly, a *Saline Body*: that is, as there is a considerable quantity of *Saline Parts* mixed with those which are properly *Aereal*. It being plain from manifold Experience; That the several kinds of *Salts*, are the grand *Agents* in the *Variation* of *Colours*. So that, to speak strictly, although *Sulphur* be indeed the *Female*, or *Materia substrata*, of all *Colours*; yet *Salt* is the *Male* or *Prime Agent*, by which the *Sulphur* is determined to the *Production* of one *Colour*, and not of another.

12. §. If then it be the *Aer* mixed with the *Juyces* of a *Plant*, and the *Salt* of the *Aer*, that makes it *Green*; It may further be asked, what kind of *Salt*? But this is more hard to judge of. Yet it seemeth, that it is not an *Acid*, but a *Subalkaline Salt*; or at least some *Salt* which is different from a simple *Acid*, and hath an *Affinity* with *Alkalies*.

13. §. One reason why I so judge, is, Because that although all *Plants* yield an *Alkaly*, or other *Salt* different from an *Acid*, and some in good quantity; yet in most *Plants*, the *Prædominant Principle* is an *Acid*. So that the *Supply* of an *Acid Principle* from the *Aer*, for the *Production* of a *Green Colour*, as it would be superfluous; So also ineffectual: a different *Principle* being requisite to the striking of this, together with the *Sulphur*, into a *Green Colour*.

14. §. I suppose therefore, That not only *Green*, but all the *Colours* of *Plants*, are a kind of *Precipitate*, resulting from the concurrence of the *Saline Parts* of the *Aer*, with the *Saline* and *Sulphurions Parts* of the *Plant*; and that the *Subalkaline*, or other like *Saline Part* of the *Aer*, is concurrent with the *Acid* and *Sulphurions Parts* of *Plants*, for the *Production* of their *Verdure*; that is, as they strike altogether into a *Green Precipitate*. Which also seemeth to be confirmed by divers *Experiments* hereafter mentioned.

15. §. THE *Colours* of *Flowers* are various; differing therein not only from the *Leaf*, but one from another. Yet all seem to depend upon the general *Causes* aforesaid. And therefore the *Colours* of *Flowers*, as well as of *Leaves*, to result not solely from the *Contents* of the *Plant*, but from the concurrence likewise of the *ambient Aer*. Hence it is, that as they gradually open, and are exposed to the *Aer*, they still either acquire, or change their *Colour*: no *Flower* having its proper *Colour* in the *Bud*, (though it be then perfectly formed) but only when it is expanded. So the *Purple Flower* of *Stock-July Flowers*, while they are in the *Bud*, are *white*, or *pale*. So *Butchelors Buttons*, *Blew Bottle*, *Poppy*, *Red Daisies*, and many others, though of divers *Colours* when blown, yet are all *white* in the *Bud*. And many *Flowers* do thus change their *Colours* thrice successively; as the youngest *Buds* of *Ladys-Lookinglass*, *Engloss* and the like, are all *white*, the larger *Buds* are *purple* or *murrey*, and the open *Flowers*, *blew*: according as they come still neerer, and are longer exposed, to the *Aer*.

16. §. But if the *Colour* of the *Flower* dependeth on the *ambient Aer*; it may be asked: How it comes to pass then, that this *Colour* is various, and not one, and that one, a *Green*? that is to say that all *Flowers* are not *Green*, as well as the *Leaves*? In answer to this Three things are to be premised.

17. §. *First*, What was said before, is to be remembred, that here the *Aer* is not a solitary, but concurrent *Cause*. So that besides the *Efficacy* of this, we are to consider that of the several parts of the *Plant*, by which the *Contents* both *Aereal* and *Liquid* are supplied to the *Flower*.

18. §. *Secondly*, That in the *Lymphaducts* of a *Plant*, *Sulphur* is the predominant *Principle*, and much more abounding than in any other part of a *Plant*, as also hath been formerly shewed.

19. §. *Thirdly*, That it appears, according to what we have observed in the *Anatomy* of the *Flower*, That the quantity of *Lymphaducts* with respect to the *Aer-Vessels* is greater in the *Flower* than in the *Leaf*.

20. §. It seemeth therefore, that the *Aer-Vessels*, and therefore the *Aer*, being predominant in the *Leaf*; *Green*, is therein also the predominant *Colour*. I say predominant, because there are other *Colours* lye veiled under the *Green*, even in the *Leaf*, as will hereafter appear more manifest.

21. §. On the contrary, the *Lymphaducts*, and therefore the *Sulphur*, being more, and the *Aer-Vessels* and therefore the *Aer*, less, in the *Flower* than in the *Leaf*; the ambient *Aer* alone is not able to control the *Sulphur* so far, but that it generally carrys the greatest part in the *Production* of the *Colour*. Yet in different degrees; For if the proportion betwixt the *Lymphaducts* and the *Aer-Vessels* be more equal, the *Flower* is either *White* or else *Yellow*, which latter *Colour* is the next of kin to a *Green*. If the *Sulphur* be somewhat predominant, the *Flower* will shew it self *Red* at first; but the ambient *Aer* hath so much power upon it, as gradually to turn the *Red* into a *Blew*. But if the *Sulphur* be much predominant, then the *Acid* of the ambient *Aer* will heighten it to a fixed *Red*.

22. §. Hence it is, that *Yellows* and *Greens* are less alterable, upon the drying of *Plants* than other *Colours*; *sc.* Because the *Aer* being predominant in their *Production*, they are the less lyable to suffer from it afterwards. Whereas *Reds* and *Purples*, in the *Production* whereof *Sulphur* is predominant, are very changeable. So the *Red Flowers* of *Lysimachia*, upon drying, turn *Purple*, and the young purple *Flowers* of *gloss* turn *Blew*. So likewise the *Purple* of *Bilberries*, and the *Crimson* of baked *Damascens*, both turn *Blew*. For being gathered, and so wanting a continued supply of fresh *Sulphur*, to bear up the *Colour* against the force of the *Aer*; it strikes it down at last from *Red* to *Purple* or *Blew*. I conclude therefore, that one *Principal Cause* of the *Variety* of *Colours* in the *Flower*, is the over proportion of the *Lymphaducts* to the *Aer-Vessels*, and therefore the dominion of the *Sulphur* over the *Aer*, therein.

23. §. If it be objected, that the *Aer* doth not deepen, but highten the *Colour* of the *Blood*: I answer, *First*, That I am not now speaking of *Animal*, but of *Vegetable Bodies*; the same *Aer* which hightens the *Colour* of *Blood* one way, may deepen that of a *Flower*, another: nay and may highten that of some *Flowers* too, some other way.

24. §. And therefore, *Secondly*, it is to be considered, That as there is not one only, but divers *Saline Principles* in the *Aer*; so are there also in the several *Parts* of one *Plant*; as in the *Root*, of one sort; in the *Leavs*, of another; in the *Flower*, of another; and so in the other *Parts*. For since the *Figuration* of the *Parts* of a *Plant* dependeth chiefly

chiefly upon the *Saline Principles*: and that the *Flower* hath a different *Figure* from that of the *Leaf*: it follows, that there is some *Salinè Principle* in the one, which is not in the other, especially, all in such *Flowers*, whose *Figures* are cut out by a greater *Variety* and *Complication* of *Lines*. The *Leavs* therefore, though variously shaped, yet agreeing so far in one common *Figure*, as usually to be *flat*; it therefore seemeth plain, that there is a *Saline Principle* in them all, so far *one*, as to be the chief *Cause* of that common *Figure*: and in concurrence with the *ambient Aer*, to be likewise the chief *Cause* of one common *Colour*, *sc.* a *Green*.

25. §. Whereas the *Figure* of the *Flowers*, and therefore their *Saline Principle*, being more various, and commonly distinct from that of the *Leaf*; it will easily concur with as a great *Variety* of *Salts* in the *Aer*, whether *Acid*, *Alkaline*, *Nitrous*, *Urinous*, *Armoniacal*, or any other therein existent, to the *Precipitation* of the *Sulphur* into the like *Variety* of *Colours*. Thus far of the *Colours* of *Plants* as they appear in their *Natural Estate*.

C H A P. II.

Of the COLOURS of Plants by Infusion.



THE next general *Inquiry*, proposed to be made, was this, After what manner the *Colours* of *Plants* shew themselves, upon their *infusion* into *Liquors*. The *Liquors* I made use of for this purpose, were three, *sc.* *Oyl* of *Olives*, *Water*, and *Spirit* of *Wine*. The *Water* I used was from the *Thames*, because I could not procure any clear *Rain Water*, and had not leasure at present to distill any. But next to this, that yields as little *Salt*, as any.

2. §. As for *Oyl*, it is known, that most *Plants* either by *Cocktion* or long *Infusion*, will give it their *Green Colour*. I have likewise tryed some *Yellows*, and find they will do indifferently well; as *Saffron*, which, by *Infusion* in *Oyl*, gives it a light golden *Tincture*.

3. §. Divers *Aromatick Plants*, as *Mint*, *Majorane*, &c. being dryed and infused in *Oyl* give it a double *Tincture*, both *green* and *yellow*; one drop of the *Oyl* shewing *green*; but a good quantity of it held up against a candle looketh *redish* or of a deep *yellow*.

4. §. But there is no *Vegetable* yet known which gives a true *Red* to *Oyl*, except *Alkanet Root*: with which, some colouring either common or other *Oyl*, vend it under the name of the *Red Oyl* of *Scorpions*.

5. §. These things confirm what we have said concerning the *Causes* of *Colours* in the *Leavs* and *Flowers* of *Plants*, upon this twofold Consideration. *First*, that *Oyl* is the most proper *Menstruum* of *Sulphur*. *Secondly*, that *Oyls* have a greater congruity with *Acids* than with *Alkalies*; as I have formerly shewed.

Discourse
of Mixture
Ch. 5. Inst.
2. §. 3. &
Inst. 5.

6. §. I say therefore, that in *Blews*, *Purples* and especially *Reds*, the predominant Principles being *Sulphur* and *Acid*, the *Oyl* either abstracts the *Sulphur* of it self, or at least, unlocks it from the *Acid Parts*; whereby both of them are bestowed seperately to their like parts in the *Oyl*; upon which their disunion the *Colour* vanishes: that depending, not upon either of them alone, which of themselves are *Colourless*, but upon both united together.

7. §. On the contrary, a *Green Colour* not depending on a predominant *Acid*, but an *Alkaly*, or some *Saline Principle* different from an *Acid*; this will not so easily be imbibed seperately, into the *Pores* of the *Oyl*, but only by mediation of their *Sulphur*. So that being both imbibed without any disunion, they still retain the same *green Colour* they had before in the *Plant*.

8. §. Hence also it is, that *red Roses* being dried and infused some time in *Oyl* of *Anise Seeds*, a more potent *Menstruum* than *Common Oyl*; they wholly lose their own *Colour*, and turn *white*; the *Oyl* remaining *Limpid*, as at the first. That is the *Sulphur* or that part of it on which chiefly the *Red* depended, is absorbed seperately by the *Oyl*, and so the *Colour* vanishes.

9. §. A SECOND *Menstruum* I made use of, was *Water*. And *First*, *Alkanet Root*, which immediately tinctures *Oyl* with a deeper *Red*, will not colour *Water* in the least.

10. §. Next it is observable, That *Water* will take all the *Colours* of *Plants* in *Infusion* except a *Green*. So that as no *Plant* will by *Infusion* give a perfect *Blew* to *Oyl*; so there is none, that I know of, which, by *Infusion* will give a perfect *Green* to *Water*.

11. §. But although the *Green Leavs* will not give their visible *Colour*, by *Infusion* in *Water*; yet they will give most other *Colours*, as well as the *Flowers* themselves. So the *Green Leavs* of *Cinquefoyl*, give a *Tincture* no higher than to resemble *Rhenish Wine*; those of *Hyssop*, *Canary*; of *Strawberrey*, *Malaga*; of *Mint*, *Muscadine*; of *Wood-Sorrel*, *Water* and some drops of *Claret*; of *Blood-wort*, *Water* and a dash of *Claret*; and those of *Bawm* make a *Tincture* near as red as ordinary *Claret* alone. All *Aromatick* hot *Plants*, give a *yellow-red Tincture*, or *colorem ex luteo rubrum*. All *Plants* with a *yellow Flower* give either a pale *citrine* or *yellowish Tincture*; and the like. Yet all give not their *Tincture* in the same space of time; some requiring a fortnight, others a week, others five, three or two days, and some but one, or half a day. From hence it appears, that the *Colours* of most *Flowers* are begun in the *Leavs*; only *Green* being therein the predominant *Colour*, as a *veil* spread over them, conceals all the rest. But passing on into the *Flower*, where the *Aer-Vessels*, as is aforesaid, are under the dominion of the *Lymphaducts* they shew themselves distinctly.

12. §. A THIRD and the last *Menstruum* I made use of, was *Spirit of Wine*. And here it is to be remarked; That as *Oyl* rarely takes a *Red*, there being but one known *Instance* of it; nor *Water*, a *Green*: So neither *Spirit of Wine*, a *Blew*. I have tryed with several *blew Flowers*, as of *Lark-heel*, *Violet*, *Mallows*, *Burrage*, and others, whereof it will not take the least *Tincture*.

13. §. Again though no *Blew Flowers*, that I know of, will give a *Blew Tincture* to *Spirit of Wine*: yet having been for some days infused in

in the said *spirit*, and the *spirit* still remaining in a manner *Limpid*, and void of the least *Ray* of *Blew*; if you drop into it a little *Spirit* of *Sulphur*, it is somewhat surprizing to see, that it immediately strikes it into a full *Red*, as if it had been *Blew* before: and so, if you drop *Spirit* of *Sal Armoniac* or other *Alkaly* upon it, it presently strikes it *Green*. Which further confirms what have been before said of the *Causes* of *Vegetable Colours*.

14. §. It is also observable, That the *Green Leaves* of *Bawm*, which give a *Muscadine Red*, with some *Rays* of *Claret*, to *Water*, gives a pure and perfect *Green* to *Spirit* of *Wine*: and is the only *Plant* of all that I have yet tryed, which doth the like.

15. §. It is likewise to be noted, That both *Yellow* and *Red Flowers* give a stronger and fuller *Tincture* to *Water*, than to *Spirit* of *Wine*; as in the *Tinctures* of *Cowslip*, *Poppys*, *Clove-July-Flowers* and *Roses*, made both in *Water* and *Spirit* of *Wine*, and compared together, is easily seen. So that for *Tinctures* made with *Flowers*, whether for *Medicines*, or other purposes, *Water*, with respect to the *Colour*, is the better *Menstruum*. I say for *Tinctures* made with *Flowers*; for there are some other *Parts*, especially *Gumms*, as *Gamboja*, *Myrrh* and *Aloes*, which give their *Tinctures* full and clear, only to *Spirit* of *Wine*. Some of which are used by *Leather-Gilders*, and others, for the washing over of *Silver*, so as to give it the *Colour* of *Gold*. Thus far of the *Colours* of *Plants* as they appear upon *Infusion*.

CHAP. III.

Of the *COLOURS* of *Plants* produced by their *Mixture* with other *Bodies*.



THE last general Enquiry proposed to be made, was this, After what manner they would exhibit themselves upon the *Mixture* of those *Infusions*, or of any one of them with some other *Liquor*.

2. §. A strong *Infusion*, or the *Juyce* of the *Leavs* of *Rose-Tree*, *Raspis*, *strawberry*, *Cynquefoyle*, *Goosberry*, *Primrose*, *Jerusalem Cowslip*, *Bearseare*, *Bearsfoot*, *Peony*, *Bistort*, *Lawrel*, *Goats-beard*, dropped upon *Steel*, make a *Purple Tincture*. But that of *Vine Leaves* scarce maketh any *Tincture* at all. So that there is something else besides *Soverness* concurring to the *Purple* upon *Steel*.

3. §. *Saccharum saturni* dropped on a *Tincture* of *Red Roses*, turneth it to a *faint pale Green*.

4. §. *Salt* of *Tartar* dropped upon the same *Tincture*, turneth it to a *deeper Green*.

5. §. Spirit of Harts Horn droped upon a Tincture of the Flower of Lark-heel and Borage turn them to a verdegreefe Green.

6. §. Spirit of Harts Horn droped on most green Leavs doth not change them at all. The like Effects have Aq. Calcis, and spirit of S. Armoniac.

7. §. These Experiments seem to confirm, That it is some Alkaline or other like Salt in the Aer, which is predominant in the production of Green in the Leavs of Plants.

8. §. Salt of Tartar droped on the white Flowers of Daisy, changeth them into a light Green. Which as it further confirms the aforesaid Position; so likewise argues, That Whiteness in Flowers, is not always from the defect of Tincture: but that there may be White, as well as Yellow, Green, Red or Blew Tinctures.

9. §. Spirit of Sulphur droped on the green Leavs of Adonis Flower, Everlasting Pease, and Holy Oak, turns them all Yellow.

10. §. Spirit of Sulphur on a Tincture of Saffron changeth it not.

11. §. Spirit of Sulphur on the Yellow Flower of Crowfoot alters them not. Neither are they changed by the Affusion of Alkalies.

12. §. So that it seemeth, that in all Yellows, the Sulphureous Acid and Alkaline Parts are all more equal.

13. §. Spirit of Sulphur on a Tincture of Violets turns it from Blew to a true Lacke, or midle Crimson.

14. §. Spirit of Sulphur upon a Tincture of Clove-July-Flowers makes a bright blood Red. Into the like Colour, it hightens a Tincture of Red Roses.

15. §. So that as Alkalys, or other Analogous Salts, are predominant in Greens, so Acids in Reds, especially in the brighter Reds, in the Leavs and Flowers of Plants. Hence it is, that Spirit of Nitre droped upon the Blew Flower of Ladies Looking-Glass, Larkspur, Borage, turns them all Red, sc. into the Red of Common Lychnis. But (which is particularly to be noted) being droped on the said Red Flowers of Lychnis, alters them little or nothing: because, that very Colour is therein produced by a copious admixture of the like Principle.

16. §. The Summ therefore of what hath now been said, of the Causes of Vegetable Colours, is this: That while their Sulphur and Saline Principles, only swim together, and are not as yet united into one Precipitate, no Colour results from them, but the Contents are rather Limpid; as usually in the Root, and many other Parenchymous Parts.

17. §. When they are united, and the Alkaline are predominant, they produce a Green.

18. §. When the Sulphur and the Alkaline are more equal, they produce a Tanny.

19. §. When the Sulphur, Acid and Alkaline, there a Yellow.

20. §. When the Sulphur predominant, and the Acid and Alkaline equal, there a Blew.

21. §. When the Sulphur and Acid are predominant to the Alkaline, then a Purple.

22. §. When the Sulphur predominant to the Alkaline and the Acid to them both, a Scarlet.

23. §.

23. §. Lastly, When the *Acid* predominant to the *Alkaline*, and the *Sulphur* to them both, a *Blood-Red*: which is the highest and most *Sulphurous Colour* in Nature.

24. §. From the *Premises*, divers *Rules* do also result for the making of *Tinctures*, either for *Medicines*, or for any other purposes.

25. §. I shall only add one or two *Notes*. As first, that of all *Colours*, *Yellows* are the most fixed and unfading. As for instance, if you drop either a *Solution* of *Tartar*, or of *Spirit* of *Sulphur* upon a *Tincture* of the *Yellow Flowers* of *Crowfoot*, of *Adonis*, or of *Saffron*, neither of them will alter their *Colour*. Which shewes the strength of most *Yellows*, to resist all manner of impressions from the *Aer*.

26. §. Again, that the use of *Salts*, is not only to highten or deepen *Colours*, but also to fix and make them permanent. As for Instance, The *Tincture* of *Clove-July-Flowers*, made either with *Water* or *Spirit* of *Wine* being exposed to the *Aer*, will often turn into a *Blackish Purple*. But the addition of a few drops of *Spirit* of *Sulphur*, doth not only highten the *Colour*, but renders it stable and permanent.

27. §. Likewise, of *Salts* themselves there is choice to be made. For there are some, which although they fix the *Colour*, yet, will a little give, as we say, and not hold throughly dry; as most *Lixivial Salts*, and *Stillatious Acids*. But there are some *Salts*, which will not give in the least, as *Alum*, that in *Lime-Water* and some others; which latter, is so far from being moystened, that it is rather petrified by the *Aer*. For which reason I take it to be one of the best *Liquors* for a stable and permanent *Green*, and some other *Colours*.

28. §. Amongst all *Water-Colours*, the rarest, and most difficult to make clear bright and permanent, is a *Blew*. There are many *Flowers* of an excellent *Blew*, as those of *Bugloss*, *Lark-heel* and others; but they easily fade. And there are very few *Flowers* that will strike into a *Blew* by any *Liquor*; being almost all changeable into *Green*, *Purple* or *Red*. Yet some few there are, in which this *Colour* may be produced. As for instance, the *Flower* of *Lathyrus* or *Parseverlasting*; which upon the affusion of *Spirit* of *Harts-Horn* is changed from a *Peach*, to as pure a *Blew*, as the best *Ultramarine*: that which hitherto is, I think, wanting in *Water Colours*. *Spirit* of *Harts Horn* was the *Liquor* I used; but I question not, but that other *Alkalies*, and particularly *Lime-Water*, will have the like Effect, and so render it the more stable.

29. §. From what hath been said, we may likewise be confirmed in the use of the already known *Rules*, and directed unto others yet unknown, in order to the variation of the *Colours* of *Flowers* in their Growth. The effecting of this, by putting the *Colour* desired in the *Flower*, into the *Body* or *Root* of the *Plant*, is vainly talked of by some: being such a piece of cunning, as for the obtaining a painted face, to eat good store of *white* and *Red Lead*.

30. §. The best known *Rules* are these Two; First, that the *Seed* be used above any other part, if the variation of the *Colour* be intended. One reason whereof is, because that part being but very small, the *Tinctures* of the *Soyl* will have the greater over proportion to those of the *Seed*. Besides, the tender and *Virgin Seed*, being committed to the *Soyl*, will more easily take any peculiar *Tincture* from it, than an
other

other *Part*, which is not so susceptible, and hath been tinctur'd already. All the strange varieties in *Carnations*, *Tulips*, and other *Flowers* are made this way.

31. §. The other *Rule* is, To change the *Soyl*, or frequently to transplant from one *Bed* to another. By which means, the *Plant*, is as it were, *superimpregnated* with several *Tinctures*, which are prolifick of several *Colours*; which way is taken for *Roots* and *Slips*.

32. §. The consideration whereof, and of the foregoing *Experiments*, may direct us not only in changing the *Bed*, but also in compounding the *Soyl*, as by mixing such and such *Salts*, or *Bodies* impregnated with such *Salts*, I say by mixing these *Bodies* in such a proportion, with the *Soyl*, as although they have no *Colour* in themselves, yet may be effectual to produce a great variety of *Colours* in the *Plants* they nourish; supplying the *Plants* with such *Tinctures*, as shall concur with the *Aer*, to strike or precipitate their *Sulphur* into so many several *Colours*, after the manner above explicated: and so to bring even Natures Art of *Painting*, in a great part, into our own power.

A
 DISCOURSE
 OF THE
 DIVERSITIES and CAUSES
 OF
 TASTS
 CHIEFLY IN
 PLANTS.

Read before the *Royal Society*, March 25. 1675.

CHAP. I.

Of the several Sorts of SIMPLE and COMPOUNDED Tasts; and the DEGREES of both.



HAVE formerly published some Notes, concerning *Tasts*. Since then, I have made other *Observations* upon the same Subject: and these have produced further *Thoughts*. I will summ up all in giving an account, First, of the *Diversities*; and then, of the *Causes* of *Tasts*, chiefly in *Plants*.

2. §. The *Diversities* of *Tasts* are so many, and so considerable; that it seemeth strange, to see the matter treated of both by *Philosophers* and *Physicians*, with so much scantness and defect. For the *subject* is not barren, but yieldeth much and pleasant *Variety*. And doth also appear to be of great import unto *Medicine*. Besides, it is preposterous to discourse of the *Causes* of *Tasts*, before we have taken an account of their *Diversities*; Whereof therefore I shall in the first place, exhibit the following *Scheme*.

3. §.

Idea, §. 29.
 & *Anat.*
 of *R. P.* 2.
 §. 68, &c.

3. §. TASTS may be distinguished by these Three general ways. First, with respect to the *Sensation* it self. Secondly, with respect to its *Duration* and *Terms*. Thirdly, with respect to its *Subject*.

4. §. The *Sensation* it self is differenced two ways, by its *Species*, and by its *Degrees*. With respect to the *Species*, *Tasts* are *Simple*, or *Compounded*. By *Simple Tasts*, I mean not such, as are never found in conjunction with other *Tasts*: but the *Simple* or *Single Modes* of *Tast*, although they are mixed with divers others in the same *Body*. As for example, the *Taste* of a *Peppin*, is *Acidulcis*; of *Rhubarb*, *Amarastringens*; and therefore *Compounded* in both. Yet in the *Peppin*, the *Acid* is one *Simple Taste*, and the *Sweet* another; and so in *Rhubarb*, the *Bitter* is one *Simple Taste*, and the *Astringent* is another.

5. §. Two faults have here been committed; the defective *Enumeration* of *Simple Tasts*; and reckoning them indistinctly among some others which are *Compounded*.

6. §. SIMPLE *Tasts*, (of which, properly so called, there are commonly reckoned but Six or Seven Sorts,) are, at least Sixteen. First, *Bitter*, as in *Wormwood*: to which, the contrary is *Sweet*, as in *Sugar*. Thirdly, *Sewer*, as in *Vinegar*: to which, the contrary is *Salt*. Fifthly, *Hot*, as in *Cloves*: whereto, the contrary is *Cold*. For we may as properly say, a *Cold Taste*, as a *Hot Taste*: there being some *Bodies*, which do manifestly impress the *Sense* of *Cold* upon the *Tongue*, though not by *Touch*. So doth *Sal Prunelle*, although the *Liquor* wherein it is dissolved, be first warmed.

7. §. Seventhly, *Aromatick*. For it doth not more properly agree to an *Odour*, than a *Taste*, to be *Aromatick*. And that an *Aromatick Taste*, is distinct from an *Hot*, is clear; In that, there are many *Bodies* of a *Hot Taste*, some meanly and others vehemently *Hot*; which yet are not in the least *Aromatick*: as amongst others, is apparent in *Euphorbium*. So that although an *Aromatick Taste* be often conjoynd with *Heat*; yet it is not that *Heat* it self, but another distinct *Sense*.

8. §. Eighthly, *Nauseous* or *Malignant*, contrary to the former. Such as is perceived, together with the *Astringent* and *Bitter*, in *Rhubarb*; or with the *Bitter*, and *Sweet*, in *Aloes*. It may be called *Malignant*, because distastful although mixed in a low degree with other *Tasts*: whereas other *Tasts* will render one another grateful.

9. §. Again, *Tasts* may properly be said, to be *Soft* or *Hard*. A *Soft Taste*, is either *Vapid*, as in *Watery Bodies*, *Whites* of *Eggs*, *Starch*, *Fine Boles*, &c. Or *Unctuous*, as in *Oyls*, *Fat*, &c.

10. §. A *Hard Taste* is Fourfold, sc. *Penetrant*, *Stupifacient*, *Astringent*, *Purgent*. Contrary to a *Vapid*, are *Penetrant* and *Stupifacient*.

11. §. *Penetrant*, is a kind of *Taste*, which worketh it self into the *Tongue* (as some *Insects* into the *Skin*) without any *Pungency*; as in the *Root* and *Leavs* of *Wild Cucumber*.

12. §. *Stupifacient*, as in the *Root* of *Black Hellebore*. Which being chew'd, and for sometime retained upon the *Tongue*; after a few minutes, it seemeth to be benum'd and affected with a kind of *Paralytick Stupor*; or as when it hath been a little burnt with eating or supping of any thing too hot.

13. §. Contrary to an *Unctuous Taste*, are *Astringent*, and *Purgent*; as in *Galls*, and *Spirit* of *Sal Aromanick*.

14. §.

14. §. Again, *Tasts* are either *Continual*, as most commonly: or *Intermittent*; as that of *Dracontium*, especially in the *Root*. For after it seems to be lost and extinguished; it will then again (chiefly upon the *Collision* of the *Tongue* and *Goomes*) be plainly heightened and reviv'd.

15. §. Lastly, *Tasts* are either *Still*, as usually; or may be called *Tremulous*, as the *Heat* produced by *Pyrethrum*. *Distinct* from that of *Cloves*, *Ginger*, and many other *Hot Bodies*, in that there the *Heat* is *still*; but here in *Pyrethrum*, 'tis joyned with a kind of *Vibration*: as when a *Flame* is brandished with a *Lamp-Furnace*. Thus far of the *Sorts of Simple Tasts*.

16. §. COMPOUNDED *Tasts* are very numerous; being made by the various *Conjunction* of *Simple Tasts*, as *Words* are of *Letters*. Sometimes of two, as in *Saccharum Saturni*, of *Astringent* and *Sweet*. Sometimes three, as in *Aloes*, *Malignant*, *Bitter* and *Sweet*; in *Rhubarb*, *Malignant*, *Astringent* and *Bitter*. Sometimes four, as in *Agarick*, *Malignant*, *Astringent*, *Bitter* and *Sweet*. And in some *Bodies*, five or six *Species* may be joyned together.

17. §. For the more accurate *Observation* whereof, there are these easie *Rules*. That not too many be tasted at one time: least the *Tongue* being furcharged, become less critical. That the *Mouth* be washed with warm water betwixt every tasting. And that those things be first tasted which produce a less durable *Taste*; that so one may be thoroughly extinguished, before another be try'd.

18. §. Of the numerous *Conjunctions* of *Tasts*, which may thus be observed, there are only Six to which the penury of *Language* hath allowed (if I may call them) *Proper Names*, sc. *Acerbus*, *Austus*, *Acris*, *Muriaticus*, *Lixivus* & *Nitrosus*. Most of which are commonly taken in to make up the number of *Simple Tasts*. But very improperly; being all of them *Compounded* and *Decomposed Tasts*: to which *Class* they ought therefore to be refer'd. For

19. §. *Austere*, is *Astringent* and *Bitter*; as in the green and soft *Stones of Grapes*.

20. §. *Acerb*, properly so called, is *Astringent* and *Acid*; as in the *Juyce* of unripe *Grapes*.

21. §. *Acris*, is also *Compounded*. For first, simply *Hot*, it is not: because there are many *Hot Bodies*, which are not *Acria*; as the *Roots* of *Zedoary*, *Yarrow*, *Contrayerva*. Nor Secondly, is it simply *Pungent*, because there are also *Bodies*, which are *Non-acria pungentia*; of which kind is the *Root* of *Arum*. Wherefore *Acritude*, is *Pungency* joyned with *Heat*.

22. §. *Muriatick*, is *Saltness* joyned with some *Pungency*, as in common *Salt*.

23. §. *Lixivial*, is *Saltness* joyned with *Pungency* and *Heat*.

24. §. *Nitrous*, is *Saltness* joyned with *Pungency* and *Cold*.

25. §. Besides these Six, or perhaps one or two more, there are, as is said, a great number of *Conjunctions*, for which we have no *Proper Names*. For admit that there were but *Ten species* of *Simple Tasts*, sc. these *Ten*; *Amarus*, *Dulcis*, *Acidus*, *Salsus*, *Calidus*, *Frigidus*, *Aromaticus*, *Malignus*, *Astringens*, *Pungens*. And of these *Ten*, but *Two*, or at most, but *Three* to be compounded together in any one *Body*. If only *Two*, they produce 45 *Compounded Tasts*. For the *First*, may

be compounded with all the 9 following; the *Second*, with all the 8 following; and so, the rest: which together make 45. But if the same *Ten* be compounded by *Threes* together; they produce no less than 120 *Variations*: as by the *Table* made of them all doth plainly appear.


26. §. Some few of the *Conjunctions* therein set down, may not be found actually existent in *Nature*. The abatement of which, will be much more than compensated two ways. *First*, by the other *Six species* of *simple Tasts*, which are also sometimes compounded. And by other more complex *Conjunctions*, as of many *Quadruples*, and perhaps some *Quintuple* or *Sextuple* ones. Thus far of the *Simple Species*, and *Conjunctions* of *Tasts*.

27. §. THE DEGREES of *Tasts* are also numerous; and each *species*, in every *Conjunction*, capable of *Variation* herein. For the more accurate observing whereof, it will be best, To take those *Bodies*, whose *Tasts* are, as near as may be, the same in *specie*: and that those be first tasted, which are less strong; whereby the true *Degree* will be more precisely taken.

28. §. The *Tasts* of *Bodies* will thus appear to be varied, in most *Species* unto *Five Degrees*; and in some of them, unto *Ten*. So the *Root* of *Turmeric*, is bitter in the *First Degree*; of *Gentian*, in the *Tenth*. The *Root* of *Carduus Benedictus*, is *Hot* in the *First Degree*; the *Green Pods* or *Seed-Cases* of *Clematis peregrina*, in the *Tenth*. So that, allowing some to vary under *Five*; yet by a moderate estimate, we may reckon every *Species*, one with another, to be varied by at least *Five Degrees*. Which being added to the several *Species* of *Tasts*, in all the *Treble Conjunctions* of the aforesaid *Table*, come to 1800 sensible and defineable *Variations* of *Taste*. And these are the *Diversities* of *Taste*, with respect to the *sensation* it self.

CHAP. II.

Of the DURATION and several TERMES of Tasts.



THE next general way of distinguishing *Tasts*, is by their *Duration*, and their *Terms*, or their *Motion* of *Intension* and *Remission* from one *Degree* to another. For there are many *Tasts*, which have their *Motions* analogous to those of *Diseases*; and by those may be distinguished in the same manner. For as of *Diseases*, so of *Tasts*, there are *Four Times*, as *Physicians* call them, or *Terms* of *Motion*; sc. *Principium*, *Augmentum*, *Status*, & *Declinatio*.

2. §. For the distinct observing of which, those *Bodies* which are hard, and so their tastable parts less easily extractable by the *Tongue*, should be reduced to a fine *Powder*: otherwise, the true measure of the
Princi-

Principium will be lost. And for the precise measuring of all the *Four Termes*, it should be done by a *Minute-Watch* or a *Minute-Glass*. For so it will appear, that the *Variations* of each, are divers and remarkable.

3. §. To instance first in those of the *Principium*. Which I call, That space of time, betwixt the first *Contact* of the *Body* to be tasted, and the first manifest *Perception* of the *Taste*. For Example, those *Bodies* which are *Acid*, or *Bitter*, as *Vinegar* or *Wormwood*, are presently perceiv'd, *quatenus Acid* or *Bitter*, upon the first *Contact*; or have *Principium brevissimum*. Those *Bodies* which are *Acria*, have their *Principium* somewhat longer. So the *Seed-Cases* of *Clematis peregrina*, although they have a vehement *Acritude*, even in the Tenth Degree; yet is not that *Acritude* so soon tasted, as the *Bitterness* of *Roses*, which is but in the second. But the *Principium* of *Hot Tasts*, is generally longer than that of any other. So the *Bitterness* of the *Root* of *Black-Helebore*, which exceedeth not the second Degree, is yet presently tasted: but the *Heat* proceeding from the same *Root*, and which ascendeth to the third Degree, is not perceived at all, till after two full *Minutes*. And so the *Bitterness* of *Enula*, which exceedeth not the 4th Degree, yet is sooner tasted than its *Heat*, which ascendeth to the 8th.

4. §. Next, in those of the *Augment*. Which I call, That space, betwixt the first *Perception* of the *Taste*, till it be come to the heighth. So the *Heat* of *Galangale*, is not only presently perceived, but ariseth to the heighth within half a *Minute*. But the *Heat* of the *Root* of *Enula*, comes not to the heighth till after a whole *Minute*. And the *Heat* of *Black-Hellebore*, not till after four full *Minutes* from the first *Contact*.

5. §. The *Status*, or space wherein the *Taste* continues in its heighth, is also divers. So the *Heat* of the *Seed-Case* of *Helleboraster*, comes to its heighth, and begins to decline within half a *Minute*; that of the *Root* of *Garden-Scurvygrass*, not till after a *Minute*; and that of the *Root* of *Asarum*, not till after two full *Minutes*.

6. §. And Lastly, the *Declination*, or the space betwixt the first *Remission* of the *Taste*, and its total *Extinction*. For instance, The *Leavis* of *Millefolium*, are *Bitter* in the 4th Degree, and *Hot* only in the 1st. yet the *Heat* continues for sometime, and the *Bitter* presently vanishes. *Calamus Aromaticus*, is *Bitter* in the 4th Degree, *Hot* in the 1st, and *Aromatick* in the 3^d: yet the *Bitter* quickly vanishes, the *Heat* continues two *Minutes*, and the *Aromatick* seven or eight. The *Heat* of the *Root* of *Conrayerra*, is extended, almost to two *Minutes*; the *Pungency* of *Jalap*, almost to six; the *Heat* of *Garden Scurvygrass*, to seven or eight. And even the *Bitterness* of *Wild Cucumer*, to near a quarter of an hour. But the *Heat* of *Euphorbium* dureth much longer, as also that of *Black Hellebore*. *sc.* above half an hour.

7. §. So that the *Augmentum*, is seldom extended beyond Four or Six *Minutes*, from the first *Contact*: but the *Declination*, sometimes to Thirty, Forty, or more. Thus far of the *Terms* of *Tast*, or the manner of their *Intension* and *Remission*.

C H A P. III.

Of the SUBJECT or SEAT of Tasts.



THE Third and Last way of distinguishing *Tasts*, is by their *Subject*, or the *Part* or *Parts* where they are either wholly or chiefly perceived. And so, *Tasts* are either *Fixed*, or *Movable*.

2. §. A *Fixed Taste*, is that which keepeth within the compass of some one *Part*, all the time of its *Duration*; as upon the *Tip*, or the *Root* of the *Tongue*, or other *Part*.

3. §. A *Movable Taste*, is either *Diffusive* or *Transitive*.

4. §. A *Diffusive Taste*, I call that, which by degrees spreads abroad into divers *Parts*, and yet in the mean time, adheres to that *Part* in which it is first perceived. So the *Bitterness* in the dried *Roots* of *Black Hellebore*, is first felt on the *Tip* of the *Tongue*; from whence it spreads it self to the middle of the same. And the *Bitterness* of the *Leavs* of *Wild Cucumer*, spreads from the *Tip*, to the *Root* of the *Tongue*.

5. §. A *Transitive Taste*, is that, which after sometime, wholly quitting the *Part* wherein it is first perceived, is thence transfered into some other *Part*: as the *Bitterness* of *Gentian*, immediatly from the *Tip*, to the middle of the *Tongue*. And most of the *Diffusive*, are also *Transitive*.

6. §. The several *Parts* which these ways become, and with some latitude may be called, the *Seats* of *Tasts*, are, the *Lips*, *Tongue*, *Palate*, *Throat* and *Gullet*.

7. §. Upon the *Lips*, the *Root* of white *Hellebore*, as also of *Pyrethrum*, being chewed, make a sensible *Impression*; which continues (like the flame of a *Coal* betwixt in and out) for 9 or 10 *Minutes*. But the *Heat* in other *Parts* much longer.

8. §. Upon the *Tongue*, *Tasts* are perceived in Three places, as hath been intimated. On the *Tip* or *Cone* of the *Tongue*; as most commonly. On or near the *Basis* of the *Tongue*; where the *Taste* of the *Leavs* of *Wild Cucumer* chiefly fixeth it self. Or on the *Vertex* or middle of the *Tongue*; in which place it is observable, that the *Taste* of *Gentian*, *Colocynthis*, and divers other *Bodies*, is then considerably strong, when not at all perceived at the *Tip* of the *Tongue* or in any other *Part*.

9. §. Upon the *Palate* or *Roofe* of the *Mouth*, the *Root*, as I take it, of *Deadly Nightshade* maketh its chief *Impression*; and there continues about four *Minutes* in some degree.

10. §. The *Throat*, or the *Uvula*, *Larinx* and other adjacent *Parts* are oftentimes the *Seat* of *Taste*. For there are many *Bodies*, which although they have scarce any *Taste* upon the *Tongue*, or any other of the aforesaid *Parts*, yet make a strong *Impression* on the *Throat*: as the *Leavs* of little *Daisy*, little *Celandine*, and of *Pimpinell*; as also the *Roots* of *Jalap*, *Mercury*, *Asparagus* and others. Which being chewed makelittle or no *Impression* on the *Tongue*, but their *Juyce* being swallowed,

lowed, causeth a kind of pricking in the *Throat*; as when one is provoked by a sharp *Rheum*.

11. §. And that this *Taste* or *Sense*, is truly distinct from either the *Heat*, *Pungency*, or *Acritude* upon the *Tongue*, it is hence further manifest; In that *Pyrethrum*, which is very *Hot*, and *Cortex Winterinus* which is very *Pungent* upon the *Tongue*; yet their *Juyce* being swallowed, causeth no *Heat*, *Pungency* or *Exasperation* in the *Throat*.

12. §. *Lastly*, if we will take the word (*Tast*) in a larger sense, the *Oesophagus* it self may be said to be sometimes the *Subject* thereof; as of the *Heat* produced by the *Root* of *Common Wormwood*. For of this *Heat* it is remarquable, that being first perceived on the *Tip* of the *Tongue*, it thence maketh its *transit* to the *Root* of the *Tongue*, and so into the *Throat*, and by degrees descends into the very *Gullet*; where it seemeth to warm the *Stomach*; and so continues, in some degree, almost $\frac{1}{4}$ of an hour. And the *Transition* and *Descent* of this *Heat* is made, although none of the *Juyce* be swallowed. And in this maner *Tasts* are distinguished with respect to their *Subject*.

13. §. So that the general *Diversities* of *Tasts* are these. With respect to their *Species*, they are *Simplices vel Compositi*; To their *Degree*, *Remissi vel Intensi*; To their *Duration*, *Breves vel Diurni*; To the *Terms* of their *Motion*, *Celeres vel Tardi*; and lastly, To their *Subject*, *Fixi, Diffusivi & Transivi*.

14. §. I HAVE thus endeavoured to draw up a *Scheme* or *Inventory* of the several sorts of *Tasts*. In which, some may think, that I have over done: and that as *Galen* hath been censured for being too curious in the *Distinctions* of *Pulses*; so have I been, in these of *Tasts*. Not to enquire now, how far the *Differences* of the *Pulse* may be extended, or be fit to be taken notice of; I shall only say, That we have not so much reason to censure him, if he hath given us some few which are coincident; as we have to thank him, for observing so many which are really distinct.

15. §. By the *Scheme* of *Tasts* here represented, we may be able, so to enumerate the *Modes* of any *Tast*, as to make a *Scientifick Definition* of it. Which is pleasant *Instruction* to any inquisitive mind; these things being all matter of sense and demonstration; wherein lyeth, though not always the most plausible, yet the most satisfying *Philosophy*; and where men, after they are grown weary with turning round, are oftentimes contented to rest.

16. §. But the usefulness of this *Schem* will further appear, in two respects; *sc.* In conducting us to a clearer and more particular *Explication* of the *Causes* of *Tasts*: and the *Investigation* of the *Virtues* of those *Bodies* in which they reside. Whereof in the following *Chapters*.

C H A P. IV.

Of the CAUSES of Tasts.



O speak of the *Causes* of *Tasts*, before we have well enumerated and distinguished them; is to provide *Furniture* for a *House*, before the *Roomes* have been counted and measured out. But the *Varieties* of *Tasts* having been first laid down; it will induce us to believe, and investigate as great a variety in their *Causes*.

2. §. Now the *Causes* of *Tasts*, particularly of the *Tasts* of *Plants*, whereof we chiefly speak, are, in general, these Four or Five, *sc.* The *Bed* out of which they grow; The *Aer* in which they stand; The *Parts* of which they consist; The several *Fermentations* under which their *Juyces* pass; And the *Organs* by which their *Tastable Parts* are perceiv'd: as will appear upon *Instance*.

3. §. But the immediate *Causes*, besides the *Organs* of *Taste*, are the *Principles* of *Plants*. As many of which, as come under the notice of *Sense*, we have already supposed to be these Seven, *Alkaline*, *Acid*, *Aer*, *Water*, *Oyl*, *Spirit* and *Earth*. The *Particles* both of *Alkaline* and *Acid Salts*, are all *angular* and *pointed*. Those of *Aer*, properly and strictly so called, are *Elastick* or *Springy*; and therefore also *Crooked*; as I have likewise formerly conjectured. And I find the Learned *Borelli*, in a Book of his since then published, to be of the same Opinion. Those of all *Fluid Bodies*, *quà Fluid*, and therefore of *Water*, *Oyl* and *Spirit*, I conceive to be *Globular*, but *hollow*, and with holes in their Sides. Those of *Water*, to be larger *Globes*, with more *holes*; those of *Oyl*, to be lesser, with fewer *holes*; and those of *Spirit* the least. Lastly, that the particles of *Earth* are also *Round*; yet *angular*; and nearer to a *solid*.

4. §. These *Principles* affect the *Organs* of *Sense*, according to the variety of their *Figures*, and of their *Mixture*. So those which are sharp or pointed; and those which are *springy*; are fitted to produce any stronger *Taste*: and those which are round, are apt, of their own *Nature*, to produce a *weaker* or *softer* one. And so by the diversities of their *Mixture*; not only with respect to their *Proportion*, but also the very *Mode* of their *Conjunction*. Hence it is, that many *Bodies* which abound with *salt*, as *Ambar* with an *Acid*, and the *Bones* of *Land-Animals* with an *Alkaline*, have notwithstanding but a weak *Tast*; the *saline Parts* being in the former drowned in the *Oyl*, and in the latter also buried in the *Earth*.

5. §. The same is further confirmed by an *Experiment* mentioned in a former *Discourse*; *sc.* the *Transmutation* of *Oyl* of *Anise-seeds*, with the help of *Oyl* of *Vitriol*, into a *Rosin*. For both those *Liquors*, though so strongly tasted, apart; yet the *Rosin* made of them, being well washed, hath a very mild *Taste*, and without any smatch of that
in

Of Mixt.
Ch. 5.

in either of the *Liquors*. Whence it follows, that the very *Mode* of *Mixture* is sufficient, not only for the variation of the *Degrees* in any one *species* of *Tast*; but also for the destroying of one *species*, and the introducing of another.

6. §. THESE things being premised, I conceive, That as an *Unduous Tast* dependeth upon *Oyl*; so a *Vapid* either on *Water*, or *Earth*: or upon such an intimate *Mixture* of other *Principles*, as renders them indissoluble by the *saliva*, and so, in a manner, untastable.

7. §. That a *Pungent*, is made either by an *Alkaly* or an *Acid* sharpened or whetted; that is, cleared from the soyl of other *Principles*; as in the *Spirit* of *Sal Aromoniac* or of *Sulphur*. And so in those *Plants* which have a *Pungent Tast*; whose *Juyces* or *Tinctures*, although they consist of divers *Principles*, yet all so loosely mixed, that being dissolved by the *Saliva*, the *saline* are hereupon left naked. Wherefore *biting Plants*, quæ *biting*, are *Nitrous Plants*. So that the *Juyce* of such *Plants*, is a kind of *Spirit* of *Nitre*, made by the several *Parts* of the *Plant*. Hence *Arum* grows best under an *Hedg*; where the *Ground*, not being exposed to the *Sun*, but the *Aer* only, like those *Rooms* in *Houses*, which are covered, is impregnated with a greater quantity of *Nitrous salt*. And those *Roots* which are *Biting*, have but few or but small *Aer-Vessels*; whereby fewer parts of the *nitroaereal Sap* are carryed off into the *Trunk*. For the same *Cause*, it is no wonder, that many *Aquaticks* are *Biting*; *Water* being, though it self cold, yet the *Menstruum* by which all *Salts* are imbibed most easily, and in laxer state of *Commixture* with other *Principles*.

8. §. *Penetrant* (something flower than *Pungent*) is made by any *salt* that is also soiled or guarded with *Earth*. *Sower*, by an *Acid* only soyled with *Earth*. *Salt*, by an *Acid* guarded by an *Alkaly*, and soyled with *Earth*. *Cold*, by an *Acid* drowned in *Water*, and soyled with *Earth*.

9. §. In all these, the *salts* are *predominant*; In *Heat* the *Oyl* or *Sulphur*. The particles whereof being *Spherick* and bored with *holes*; those of *salt* stick in them, as the *Spokes* do in the *Hub* of a *Wheel*, or as the *Quills* in the *Skin* of a *Porcupine*. Whereby, as in *Common Fire* the *Sparks* of *sulphur* being agitated and whirled about by the *Aer*; with the help of the *Salts*, which stick in them, tear in pieces all kinds of *Bodies*: so here, being agitated by the *Circulation* of the *Blood*, they make a kind of hurry or combustion; and so, according to the degree and strength of their *Motion*, tear in pieces fewer or more of the *Fibers* of the *Tongue*; and in a greater quantity, would raise a *Blister* upon it; the common *Effect* of *Fire*, or any strong *Epispastick*. So that a *Hot Tast*, is produced by *Sulphur* toothed or armed with *Salts*. Wherefore all *Stillatitious Oyls* are *Hot*; being strongly impregnated or armed with the *Essential Salts* of the *Plants* from whence they are distilled. And as those *Plants* which are very *Parenchymous*, from the predominancy of their *Volatile Acid*, are *biting*: So those which are *Lignous*, that is, have a good quantity of *Lympheducts*, from the domiou of their *Sulphur* are commonly *Hot*. For the same reason it is, that many both *Biting* and *Hot Plants*, as the *Roots* of *Dragon*, *Garden-Radish*, *Onion*, *Iris*, *Rape-Crowfoot*, &c. being corked up in a bottle with *Water*, and set in a *Cellar* or other cool place; they do all of them turn *Sower* in a few days: The same *Fermentation*, at once
fullying

fullying the *Salts* of the one, and disarming the *Sulphurs* of the other. But some, wherein the *Sulphureous* parts are more copious, will hardly ever become *Sower*. Hence also, some *Plants*, whose *Roots* are neither *Hot*, nor of any strong *Taste*, as those of *Wild Anemone*; yet their *Leaves* and *Flowers* are plainly *Caustick*: So that it seems, that as their *Juyces* rise up into the *Trunk* or *Stalk*, and are therein further fermented, the *Sulphureous Parts* thereof, are at the same time relaxed from the other *Principles*, and acuated with an *Aereal Salt*.

10. §. A *Stupifacient Taste* (as the *Impression* which some *Hot Plants* make upon the *Tongue* may be called) is in some sort, analogous to the mortifying of any part of the *Body* by the application of a *Caustick*. For as there the mortification succeeds the burning pain, so here, the *Stupifaction*, neither comes before, nor with the *Heat*, but follows it.

11. §. *Sweetness* is produced, sometimes by an *Alkaly*; smoothed either by a *Sulphur*, as in *Lime-Water*; or by both a *Spirit* and a *Sulphur*, as in the *Stillatitious Oyls* of *Animals*. But most commonly, by a smoothed *Acid*; as in *Malt*, *Sugar*, *Hony*. Hence a *Sweet Taste*, is generally founded in a *Sower*; So *Sower Apples*, by mellowing, and harsh *Pears*, by baking become sweet; the *Spirit* and *Sulphur* being hereby at once separated from the other *Principles* and brought to a nearer union with the *Acid*. So the *Sower Leaves* of *Wood-Sorrel*, being dry'd, become sweet: and those of a sower *Codlin*, while they hang on the *Tree*, and even of a *Crab-Tree*, are neither *Astringent*, nor sower, but sensibly sweet. And so commonly, wherever the said *Principles* are a little exalted by a soft *Fermentation*; as in the *Juyce* of the *Stalk* of *Maze* or *Indian Wheat*, which is as sweet as *Sugar*; and in the *green Stalks* of all sorts of *Corn* and *Grass*, in several degrees. So likewise *Tulips* and some other *Roots*, being taken up, in open weather, sometime before they *sprout*; if tasted, are as sweet as *Liquirish* or *Sugar*; and at no other time: not only *Fruits*, but many *Roots*, *Seeds*, and other *Parts*, upon their first or early *Germination*, acquiring a curious *Mellowness*, wherein, all their *Principles* are resolved, and their most *Spirituons Parts* exalted and spread over the *Acid*. Wherefore also most *Roots*, which are not meerly long, but grow deep in the ground, have at least some of their *Juyces* of a sweet *Taste*; as *Liquirish*, *Eryngo*, *Hounds-Tongue*, *Garden-Parsnep*, *Black Henbeane*, *Deadly Nightshade*, &c. Even the *Juyce* of *Horse Radish*, which bleeds at the *Lympheducts*, is of a sweet *Taste*. And of the same kindred those which grow the deepest, are the sweetest; as a *Parsnep* is sweeter than a *Car-root*, especially if you tast the bleeding *Sap*; and the *Root* of *Common Tall Tresfoyl* tasteth somewhat like *Liquirish*, but is not near so sweet. For all deep *Roots*, are fed with a less *Nitrous Aliment*: and being remoter from the *Aer*, their *Juyces* pass under much more soft and moderate *Fermentations*.

12. §. *Bitterness* is produced by a *Sulphur* well impregnated, either with an *Alkaline*, or an *Acid Salt*, but also shackled with *Earth*. And therefore the *Bitterest Plants*, commonly yield the greatest quantity of *Lixivial Salt*. So also many *Stillatitious Oyls* digested with any strong *Acid*, will acquire a *Bitter Taste*. Wherefore this *Taste* is often founded either in a *Hot Taste*, or a *Sweet*. Hence it is, that the *Leaves* of all sweet *Roots* are *Bitter*. And that the *Fig-Tree*, which bears a sweet *Fruit*, bleeds a *Bitter Milk*. So likewise those *Plants*, which bear a
Bitter

Bitter Stalk, have not *Bitter*, but *Hot Roots*, as in *Tarrow*, *Primrose*, *Wormwood*, *Rue*, *Carduus benedictus* &c. is manifest. So the *Coats* of the *Seeds* of *Viola Lunaria* are of a *hot* and *biting Taste*; but the *Seeds* themselves, in which the *Salts*, though copious, yet are also immersed in a greater quantity of *Oyl*, are *Bitter*. And that the *Earthy Parts* do also contribute something more to this, than to most of the forementioned *Tasts*, is argued from its being more *Fixed*; that is, the *Body* in which it resides, is either more *Fixed*, or else flyeth not away in that same state of conjunction, by which it maketh a *Bitter Taste*. For whereas *Hot*, *Biting*, and divers other *Plants* lose the strength of their *Taste*, by drying; most of those which are *Bitter*, do hereby increase it. And although the *Extract* of *Dandelion* and some other *Roots*, which are very *Bitter*, hath scarce any *Tast*; yet generally, they are *Bitter Plants*, which are best for the making of *Extracts*. And the distilled waters of *Plants* which are *Hot* and *Bitter*, notwithstanding that they always tast high of the *Heat*, yet rarely and very faintly of the *Bitter*.

13. §. *Astringency*, is made, partly, by the further increase and more intimate union of the *Earth*. And therefore this is seated still in a more *Fixed Composition*, than a *Bitter*. And partly, by the diminution of the *Sulphur*. And therefore the *Acid Parts* ingredient to it, either by *Fermentation* or otherwise, are easily exposed. *Astringency* being the *Womb* or *Bud* of a *Sower*. For all or most *Astringent Roots* bear a *sower Leaf*, or a *sower Fruit*; as those of all *Docks* and *Sorrels*, *Black-Thorn*, *Dog-Rose*, and others. Wherefore also, *Astringency* is often found in conjunction with *Bitter*, *Sweet*, or *Sower*; but scarce ever with *Pungent*, or *Hot*.

14. §. An *Aromatick Taste*, seems to be produced, chiefly, by a *spirituous*, *acid*, and *volatile Sulphur*; as in *Ambar-griese*, *Cardamon- Seeds*, many *Stillatitious Oyls* &c. A *Nauseous*, by a *Sulphur* less *Spirituos* and *Volatile*, and more *Alkaline*; as in the *Root* of *Dog-stones*, *Sheep-scabious*, the young and green *Leaves* of *Coriander*, or the *Seeds* of *Cumine*. The *Spirit*, as it enters the *Nerves*, carrying the *Alkaline Sulphur* along with it; as when a *City* is betrayed by one of its *Inhabitants* to an *Enemy*.

25. §. An *Intermittent Taste*, as in *Arum*, seems to have its dependance upon a simple and very pure *Nitre*, which by its subtilty enters into the very *Concaves* of the *Nervous Fibers* of the *Tongue*: and so being lodged there, is little affected or stirred, by the *Motion* of the *Blood*; but only when the *Tongue* it self is moved, at which time it causeth a kind of pricking *Taste*.

16. §. A *Tremulous Taste*, as in *Pyrethrum*, dependeth probably, upon an *Aereal Sulphur*; which being agitated by the *Blood* in its *Circulation*, the *springy Motion* or *Vibration* of the *Aereal Parts* produce that *Taste*.

17. §. A *Tast* is *Lingual*, *Guttural*, &c. according to the grossness or fineness or other difference of the *Membranes* into which the *tastable parts* are admitted. For *Tasts* are made not meerly by the outward *Contact*, but the *Ingress* of the *tastable parts*. Now the outer *Skin* of the *Tongue*, which is commonly observed to pill off in boyling, like the *Cuticula* in other *Parts*, hath either no sense, or much less than that which lies under it; and is therefore, but a *Seive* or *Strainer* to the *tastable parts*. So that being of different fineness in the several parts of the


Tongue; it hereby comes to pass, that according as the *tastable parts* of any *Plant* are more or less penetrant, subtle, or dissoluble, they are admitted into one part of the *Tongue*, and not another. And in the *Throat*, the outer *Skin* it self, seems to be the immediate *Sensory*; and so, to be evidently affected with the *Juyces* of some *Plants*, from which the *Tongue* receiveth little or no sensible *Impression*.

18. §. When the *Tast* is *Permanent* and *Fixed* in some one *Part*; it is a sign, either that the *Gustable Parts* are less dissoluble; or more subtle, so as to enter the *Concaves* of the *Fibers*; and that there is an admixture of an *Aereal Salt*, or a like *Sulphur*; some of the parts whereof, being crooked, hang like *Hooks* on the *Fibers* of the *Tongue*. For the reception of such a *Tast*, is not to be looked upon as a wound made with a *Lancet*, and so the *Lancet* taken away; but with the *Lancet* sticking in the wound; until in time, 'tis carryed off by the *Circulation* of the *Blood*; which like the *Stream* of a *River* in a *Flood*, carries all before it, but those things last, which stick in the *Mud*.

19. §. But when the *Tast*, though *Permanent*, yet is *Diffusive* or *Transitive*; it seems probable, that as there is a less admixture of *Aer*; so a greater subtlety of the *Tastable Parts*, whereby they are conveyed, through the *Nervous Fibers*, from one *Part* to another.

CHAP. V.

Of the Judgment which may be made of the *VIRTUES* of *Plants*, from their *Tasts*.



By duly observing the *Tasts* of *Plants*, we may be directed to understand their *Causes*. So also the *Use* and *Virtues* of those *Plants* or *Parts* of *Plants* in which they reside. For the proof whereof, an *Instance* might be fetched from every particular difference of *Tast* before set down. But it may be enough, to give these which follow.

2. §. And first, we may make no ill guess *ex Analogia*, or where we find the same *Tast*, that there the same *Virtue* in some kind, and in some degree, may reside. So *Jalap*, *Mercury*, and *Daisy*, have all of them that exasperating *Tast* in the *Throat* before described; and they are all three more or less *Cathartick*. Wherefore, we may believe, that other *Plants* which make the like *Impression* on the *Throat*, and there are many others which do, that they are in some degree alike *Cathartick*. Those *Plants* which are reckoned amongst the chiefest *Cephalicks*, cause rather a durable, than a vehement *Heat* upon the *Tongue*, as *Pyrethrum*, *Euphorbium*, *Black-Hellebore*, &c. It seemeth therefore reasonable to rank with these, any other *Plant*, though not used, which produceth the like durable *Heat*. The young *Roots* of *Yarrow*, or *Millefolium*, have

have the same *Taste*, as the *Root* of *Contrayerva*: and may therefore be used for the same purpose, with a probability of the like success; if not a better, because they may be gotten fresher. But by drying the *Root*, the *Tast* and *Virtue*, which lie in its exhalable parts, are much lost. The *Seeds* of the lesser *Cardamom*, and of *Zedoary Root*, if sound, have both a smack of the *Tast* of *Camphire*. They may therefore all, so far, reach the same Case.

3. §. Again, as we may make no ill conjecture from the sameness of *Taste* in *Plants* of several *Tribes*; so from the diversity of *Taste*, in those of the same. So the *Flowers* of all the *Docks* are evidently *Astringent*, and not *Sower*; except those of the *Rha-pontick*, which are extream sower, even in the 5th degree. Which is no mean *Signature* of some more than ordinary *Virtue* in it, besides what it hath in common with the rest of the *Tribe*. The *Flowers* of *Pancy* have a kind of fulsome *Tast*, plainly different from that of *Violets*: and in some *Hypochondriacal Cases* may be more useful.

4. §. It likewise importeth much, to observe the difference of *Taste* in the several *Parts* of the same *Plant*. So the *Barque* of *Sassafras* is three times as strong, as the *Wood*: and the like may be observed in any other commonly known *Tree*. If therefore we could obtain the *Barques* of *Santalum*, *Lignum Rhodium*, *Lignum Aloes*, &c. they would doubtless, most of them, be of much greater use. And as the *Taste* is sometimes stronger; so, much more grateful, in one *Part* than in another: as in the *Flowers* or *Yellow Attire* in the *Heads* of *Carduus Benediclus*; which being infused in *Spirit of Wine*, or other convenient *Liquor*, make a pleasant *Cordial*. Nature having laped up the *Virtue* in the *Leavs*, as in a brown *Paper*; but in the *Flowers*, as in *Leaf-Gold*.

5. §. As also, how far the *Taste* of any *Plant* may alter, either in preserving, or preparing it. So the *Root* of *Arum*, when taken fresh out of the ground, is notably *Pungent*: but being thoroughly dried, and especially kept for some time, hath no more *Taste*, and therefore in all likelihood, no more *Virtue*, than a *Lump* of *Starch*. The like we are to judge of all other *Plants*, whose *Virtue* lieth in their exhalable *Parts*. The *Stillatitious Oyls* of many *Plants*, are stronger than the *Leavs* or other *Parts* from whence they are drawn: but some there are, which are weaker; as is that of *Euphorbium*, in which the *Heat* is neither pertinaceous, as in the *Gum* it self, nor so great.


6. §. We may make, moreover, a judgment from the *Nature* of the *Tast*. So those *Roots* which are *Bitter*, and not *Hot*, as of *Cichory*, and the rest of the *Intybous* kind, may be accounted *Nitro-Sulphureous*; and so, to be *Abstersive* without any *Heating Quality*. The *Marum Austriacum*, which is extream *Pungent*, as well as *Aromatick*, may be looked upon as the best *Cephalick* of that *Tribe*. Because we find, that *Jalap* hath a special property of imitating the *Glandulous Parts* of the *Mouth*, and *Throat*; we may gather, That it is a better *Purge* to all the other *Glandulous Parts*, than most other *Catharticks*. Which is also one reason of its operation, for the most part, with at least a tendency to vomit; the *Stomach* it self being *Glandulous* as well as the *Throat*, and thereby answerably affected with it. A strong *Infusion* of *white Sarzaparilla* in *Water*, botled up, and kept in a *Cellar* for the space of two months, becomes extream sower; far beyond any thing observed in the *Tasts* of the *Juyces* and *Infusions* of divers other

Plants kept as long and in the same manner. Which shews, how well Nature hath adapted a *Plant* of so mild a *Taste*, either by similitude of parts, for the carrying off of any *preternatural Acid*; or by contrariety, for the curbing of an exorbitant *Salt*. The *Barque* of the *Root* of common *Wormwood*, which impresseth a pertinacious and diffusive *Taste*, which descendeth from the *Tongue* into the *Gullet*, as is before described; may be justly ranked with the most excellent *Stomachicks*; and upon tryal, I find it one of the best: besides, that it is neither unpleasant, nor affecteth the *Head*, as the *Leavs*. Yet the *Gardener*, and every *Body* throws it away, as good for nothing.

7. §. I shall conclude with one note, which is this; That the *Specifick Virtue* of *Medicines*, which some *Physicians* positively deny, and most dispute; from some of the forementioned *Differences* of *Taste*, as well as for other reasons, may seem, at least, to be probable. For why should not a *Medicine* make an *Impression* upon one *Part*, and not upon another, within the *Body*, as well as we find it doth within the *Mouth*? especially, since the *Parts* of the *Mouth*, are of a less different *Nature*, than some of the *Viscera*.

An Appendix.

Of the ODOURS of Plants.



THE *Senses* of *Tasting* and *Smelling* being so nearly ally'd; many things already explained concerning the *Diversities* and *Causes* of *Tastes* in *Plants*, may easily be transferr'd to those of their *Odours*. I shall now therefore only remarque some particulars, not commonly taken notice of hitherto, and leave them as a *Specimen* to be Improved by other Hands.

§. 2. The *Root* of *Rape-Crowfoot* being cut, and held to the *Nose*, when it is newly taken out of the *Ground*, smelleth almost like *Spirit* of *Sal Armoniac*, or fresh *Scurvygrafs Juice*. And hath the property of making the *Eyes* to water, as *Onions* do. *Horse-Radish Root* is not so *Pungent* to the *Nose*, but gets pretty much into the *Eyes*. But that of *Dragon*, doth neither affect the *Eyes*, nor the *Nose*.

3. §. The *Succulent Roots* of *Dogstones*, and most of that *Tribe*, have a rank *Smell*. And that of *Crown Imperial*, being rub'd a little, smells as like a *Fox*, as one *Fox* smelleth like another.

4. §. The *Root* of *Patience* digested with *Water*, in a warm *Room*, for the space of three weeks, smels like *Spirit* of *Harts Horn*, or other *Urinous Spirit*. Of *Red Dock*, almost like *Aqua fortis* or *Spirit* of *Nitre*. That of *Dragon* bottled up with *Water*, and set in a *Cellar*, about a *Month*, stinks like the *pus* of the most *Fetid Ulcer*. At the end of five *Months*, more abominably, than either to be endured or expressed.

5. §. The *Leavs* of *Mountain Calamint*, smell like *Peny Royal*. Those of *Ulmaria*, like *Walnut Pills*. Of *Yellow Lamium*, like a *Balsame*. Of *Sena*, a good quantity being held to the *Nose*, of a rank *Smell*

Smell betwixt that of *Sweat* and *Urine*. Of *Coriander*, when green and young, stink so basely, that they can hardly be endur'd. Sometimes the *Leavs* have a stronger *Smell*, than the *Flower*, as in *Borage*, and sometimes the *Stalk*, a stronger than the *Leavs*, as in *Ulmaria*.

6. §. *Rue Leavs* corked up in a bottle and set in a *Cellar* for about ten weeks, smell like *Spirit of Harts Horn*, or of *Urine*. The green *Leaves* of *Roses* infused in water, have a mild, but pleasant *Smell*. Neither is that of *Savine* unpleasant, upon the like *Infusion*.

7. §. *Scurvygrafs Juice* kept about $\frac{1}{2}$ of a year in bottles, with the green *Sedement*, in a warm *Room*, stinks like *Humane Excrements*. And *Scurvygrafs Wine*, made only of the *Juyce*, smells like some *Issues*.

8. §. The *Flowers* of *Tarrow*, smell not much unlike to those of *Southernwood*. And the *Flowers* of *Crowfoot* almost like those of *Scurvygrafs*. Some *Flowers* are of a weaker *Smell* in the *Bud*, as those of *Mallow*. But many have a stronger, than when they are blown open; as those of *Lavender*, *Rosemary*, &c.

9. §. The *Buds* of *Vervaine Mallow*, while they are young, and the *Flowers* unseen, have a very pleasant *Smell*, like that of *Geranium Moschatum*: but when afterwards they are opened they have an unpleasant *Smell*. Common *Mallow Flowers* dried and bottled up for some time, acquire, though not a strong, yet very noysom *Smell*.

10. §. The *Purple Pouch* of *Dragon* which covers the *Seed*, being broken, smells just like a *Lobster*. But permitted to lie in a warm *Room* for some days, smells exactly like *Carrion*; and scents the *Room* with the same *Smell*.


11. §. Some *Seeds* as those of *Cumine*, *Daucus*, being powdered and laped up only in *Papers*, do notwithstanding retain their *Smell*. But many others, as of *Sweet Fenil*, in a short time, lose it. Some *Seeds*, when they first begin to sprout, become *Odorous*, which were not so before; as the *Garden Bean*.

*Tabula, quâ perspicuè videre est, quot Triplicati Sapores,
ex solummodo decem Simplicibus numerantur.*

	Am.du.ac. am.du.fa. am.ac.fa. am.du.ca. am.ac.ca. am.fa.ca. am.du.fr. am.ac.fr. am.fa.fr. am.ca.fr. am.du.ar. am.ac.ar. am.fa.ar. am.ca.ar. am.fr.ar. am.du.ma. am.ac.ma. am.fa.ma. am.ca.ma. am.fr.ma. am.ar.ma. am.du.af. am.ac.af. am.fa.af. am.ca.af. am.fr.af. am.ar.af. am.ma.af. am.du.pu. am.ac.pu. am.fa.pu. am.ca.pu. am.fr.pu. am.ar.pu. am.ma.pu. (am.af.pu.)
AMARUS	
	Du.ac.fa. du.ac.ca. du.fa.ca. du.ac.fr. du.fa.fr. du.ca.fr. du.ac.ar. du.fa.ar. du.ca.ar. du.fr.ar. du.ac.ma. du.fa.ma. du.ca.ma. du.fr.ma. du.ar.ma. du.ac.af. du.fa.af. du.ca.af. du.fr.af. du.ar.af. du.ma.fa. du.ac.pu. du.fa.pu. du.ca.pu. du.fr.pu. du.ar.pu. du.ma.pu. du.af.pu.
DULCIS	
	Aci.fal.cal. aci.fal.fri. aci.cal.fri. aci.fal.aro. aci.cal. ar. aci.fri.ar. aci.fal.mal. aci.cal.ma. aci.fri.mal. aci.ar.mal. aci.fal.ast. aci.cal.ast. aci.fri.ast. aci.ar.ast. ac.ma.ast. ac.fal.pu. aci.ca pun. aci.fr.pun. aci.ar.pun. ac.ma.pu. ac.ast.pu.
ACIDUS	
	Sal.cal.fri. fal.cal.aro. fal. fri. aro. fal.cal.mal. fal.fri.mal. fal.aro.mal. fal.cal.ast. fal. fri. ast. fal.aro.ast. fal.ma.ast. fal.cal.pun. fal.fri.pun. fal.aro.pun. fal.ma.pu. fal.ast.pun.
SALSUS	
	Cal.fri.aro. cal.fri.mal. cal.aro.mal. cal.fri.ast. cal.aro.ast. cal.mal.ast. cal.fri.pun. cal.aro.pun. cal.mal.pun. cal.ast.pun.
CALIDUS	
	Fri.aro.mal. frig.aro.ast. fri.mal. ast. fri.aro.pun. fri.mal.pun. fri.ast.pun.
FRIGIDUS	
	Aro.mal.ast. aro.mal.pun. aro. ast. pun.
AROMATICUS	
MALIGNUS	
ASTRINGENS	
PUNGENS	

Tabula,

Tabula, quæ Genericas omnes Saporum differentias comprehendit.



EXPERIMENTS
IN
CONSORT
UPON THE
Solution of Salts
IN
WATER.

Read before the *Royal Society*, January, 18. 167⁶/₇.

CHAP. I.

In which is shewed, the Compleat or Utmost Impregnation of WATER with several kinds of Salt, both together, and apart.



IN discourse upon a *Lecture* formerly read, concerning the *Lixivial Salts* of *Plants*; It was mentioned, as a thing asserted by some *Phylosophers*, That *Water* having been fully impregnated with one kind of *Salt*, so as to bear no more of that kind; it would yet bear, or dissolve some portion of another; and so of a third. And it was referred to Me by this Honourable Chair, to examine and produce the *Experiment*. The doing whereof brought into my mind divers other *Experiments* hereunto relating.

2. §. As next, With what difference of quantity this *Superimpregnation* would be made, upon the *Solution* of different *Salts*?

3. §. *Thirdly*, Whether the *Solution* of a smaller quantity of several *Salts*, doth consist with the *non-increase* of the bulk of the *Water*? Because this also is affirmed by some.

4. §.

4. §. *Fourthly*, What quantity of the several kinds of *Salt*, may be dissolved severally, in the same quantity of *Water*?

5. §. *Fifthly*, Whether by dissolving a *salt* in *Water*, there be any *Space* gained, or not? That is, whether the *Bulk* of the *Water* be greater, before the *Salt* lying in it be fully dissolved, than it is afterwards? Or if a *Cubick Inch* of *Salt* be dissolved in nine *Cubick Inches* of *Water*; Whether the *Water* will then fill a *Vessel* of ten *Cubick Inches* content?

6. §. *Sixthly*, Whether the *Space* be equally gained, by an equal encrease of the same *Salt*?

7. §. *Seventhly*, Whether upon the *solution* of several kinds of *Salts*, be gained so many several quantities of *Space*? That is, if the *solution* of common *Salt* gains, suppose, an *Inch*, whether the *solution* of *Salt Armoniack* gains as much, or more, or less? and so for other *Salts*.

8. §. *Eighthly*, What that just space may be, which any *salt* gaineth with respect to its own *Bulk*, or that of the *Water*?

9. §. And first, for the *superimpregnation* of *Water*; I put into a bottle ζij of fair *Water*; adding thereto, first half an Ounce of *Nitre*; and afterwards more, as the *Water* would dissolve it; and (that I might be sure the *Impregnation* was full) some portion above what the *Water* would bear. Then having separated this remaining portion; I put to this *solution* of *Nitre*, two Drachms of *Sal Armoniac*; which wholly and easily dissolved in the said *solution*; though it would not bear a grain more of *Nitre*. I then added a third Drachm of *Sal Armoniac*, after that a fourth, and a fifth; all which, within the space of half an hour, were perfectly dissolved in the said *solution*, without any precipitation of the *Nitre*.

10. §. In the making of this Experiment, two things, to render it infallacious, are to be noted. That the said *Salts* were not dissolved by the help of *Fire*, but only by a strong and continued *Agitation*. And that this was done upon a warm day: which I mention, because that even the changes of the weather will somewhat alter the *solubility* of the *Salts*.

11. §. Having made the Experiment upon two *Salts*, I proceeded to repeat it upon three. And first I dissolved as much common *Salt* in ζij of *Water*, as that quantity would bear. Then having separated the subsiding portion; I put to the *solution*, no less than five Drachms of *Nitre*, which by a continued *Agitation*, was wholly dissolved therein, neither the *Nitre* nor the common *Salt* being in the least precipitated. Then adding a Scruple more, it would not dissolve, but subsided. This second subsiding portion, I again separated; and then put to this *Superimpregnation*, near ζj of *Sal Armoniac*, which was also dissolved as the former. And if as many more *Salts* had been added, tis probable that the same *Water* would have born some quantity of them all.

12. §. From this Experiment, it is a Conclusion demonstrated, That not only the visible *Crystals*, but the very *Atomes* of every *Salt*, at least those *Particles* which are ultimately dissolved in *Water*, have a different *Figure* one from another. Because that if they were all of one *Figure*; there would be no *Superimpregnation*, but the *Pores* of the same *Water*, would imbibe as much of one *salt*, as answers to the total of two more *Salts* imbibed: that is to say, it would as well imbibe two Ounces of common *Salt*, as one Ounce of common *Salt* and

another of *Nitre*: which yet is contrary to the Experiment. And it is the same thing, whether we suppose the *Pores* of *Water* to be also different, or not. Because, that if the *Figure* of all the said *Atomes* be the same; then their respect to the *Pores* of the *Water* must be the same, how different so ever those *Pores* be: which is also contrary to the Experiment. Besides it is a great presumption, to say, that the *Pores*, and therefore the *Atomes* of *Water* have different *Figures*; and yet not those of *Salts*.

13. §. From the same Experiment we may go upon good ground in *Compounded Infusions*; whether of *Purgative*, or other *Materials*. As not doubting, but that the same *Menstruum* may be highly impregnated with several *Ingredients* at once, whose operative parts may be therein copiously dissolved, without hindring either an *Extraction*, or causing a *Precipitation* one of an other.

14. §. The *Second Enquiry* is, With what difference this *Superimpregnation* of *Water* is made? which I find considerable. For a *Solution* of above five Drachms of *Nitre* may be *superimpregnated* with no less quantity of *Sal Armoniac*. And a *Solution* of five Drachms of common *Salt*, may be *superimpregnated* with as much *Nitre*. Yet neither a strong *Solution* (as of five Drachms) of common *Salt*, will bear above two Scruples of *Sal Armoniac*: nor will a strong *Solution* (as of five Drachms) of *Sal Armoniac*, bear above a Drachm of common *Salt*: for if above the said quantities of either of them be mixed together: they are both copiously and forthwith precipitated to the bottome of the *Glass*.

15. §. Whence, notwithstanding the former Experiment, yet are we admonished, not to infuse all manner of *Ingredients* in any proportion. Because though some do not, yet others will precipitate one another.

16. §. The *Third Enquiry* was this, Whether the *Solution* of a smaller quantity of several *Salts*, doth consist with the *non-increase* of the *Bulk* of the *Water*? For this I took a *Bolt-head* with a slender *Neck*, containing about a pint and a quarter of *Water*; and dissolved therein about $\frac{3}{4}$ of *Nitre*. And marking the place to which the *Water* ascended in the *Neck* of the *Bolt-head*: I then dissolved in the same *Water* about a Drachm of *Sal Gemma*: which little quantity raised the *Water* above half an Inch higher then it was before. The like I observed in the addition of *Nitre* to a *Solution* of *Sal Armoniac*. So that to suppose the variation of the *salt* doth prevent the increase of the *Bulk* of the *Water*, is a manifest Error.

17. §. From the same Experiment it also appears, That the ascent of the *Water* upon a *superimpregnation*, is the same, by whatsoever *Salt* the first *Impregnation* be made. For instance, Let a *Solution* of *Nitre* ascend in the *Neck* of the *Bolt-head*, suppose, to 10 Inches, then add $\frac{1}{2}$ an Ounce more of *Nitre*, so as to raise the *Water*, suppose, 12 Inches or more, or less, according to the *Bore* of the *Neck*. In like manner, let a *Solution* of *Sal Armoniac* reach to ten Inches: then add again half an Ounce of *Nitre*; and it will reach just 12 Inches, or more or less, as before.

18. §. The *Fourth Enquiry* is, What quantity of the several kinds of *salt*, may be dissolved severally in the same quantity of *Water*: that is to say, by agitation alone, without the help of fire, as I noted before.

before. And upon tryal it appears, First, that two Ounces of *Water* will dissolve three Ounces of *Loaf-Sugar* and no more, except the *Water* be heated.

19. §. The same quantity of *Water* that is, two Ounces will dissolve above two Ounces of *Salt of Tartar*. I say above, for how much more, want of a greater quantity of *Salt* which I could confide in, made me that I could not finish the Experiment.

20. §. The same quantity, *sc.* two Ounces of *Water*, dissolveth an Ounce and a Drachm of *Green Vitriol*.

21. §. The like quantity dissolveth six Drachms and a Scruple or above $\frac{1}{4}$ of an Ounce of common *Salt*.

22. §. Of *Nitre*, Five Drachms two Scruples and an half.

23. §. Of *Sal Armoniac*, five Drachms and two Scruples.

24. §. Of *Alum*, not above two Drachms and a Scruple.

25. §. And of *Borax*, not above a Drachm and half a Scruple.

26. §. Of these note, That although Common *Salt* be very dissoluble, and will presently catch the moisture of the *Aer*: yet a much greater quantity not only of *Salt of Tartar*, but even of *Loaf Sugar*, and of *Green Vitriol* it self, may be dissolved in *Water* than of Common *Salt*.

27. §. Again, as the great *Solubility* of some, so the less *Solubility* of other *Salts* is also observable, as of *Alum*, and *Borax*. For the same quantity of *Water* will dissolve near four times as much of *Green Vitriol*, as it will of *Alum*. And of *Sugar* more than ten times as much. Of *Green Vitriol* near eight times as much as of *Borax*; and of *Sugar*, twenty times as much.

28. §. From this Experiment we are likewise cautioned, not only in the *Infusion* of several *Ingredients* together, but of any one singly; that such a proportion thereof to the *Menstruum*, be not exceeded. For all that is over and above what the *Menstruum* will bear, is either not extracted, or will be precipitated. As is evident not only in the *Dissolution* of the *Salts* above named, but in the *Infusion* of *Plants* themselves: as, for instance; of *senna*; two Drachms whereof will impregnate four Ounces of *Water* as strongly, as if twice the quantity were infused; because the *Water* will bear no more of the *Purgative Parts* of that *Body*.

29. §. There is only one *Salt* more remaineth to be spoken of under this Experiment; and that is, the *Crystals* of *Tartar*. Whereof, it is somewhat strange to observe, that it will scarce at all dissolve in *Water*: not more, than even divers *Resinous Gums*, as *Mastick*, *Tolu*, *Tacchamahacca*, and some others will do. For if two Drachms, suppose of these *Crystals*, of *Tartar* (commonly sold for *Cremor Tartari*) be put to one Ounce of *Water*, scarce five Grains thereof will, by *Agitation*, be therein dissolved.

C H A P. II.

In which is shewed, that by the Solution of Salts in Water, some certain space, more or less, is gained. That the space is different according to the Nature of the Salt. And what the just space is, which is gained.



THE Fifth Enquiry is, Whether by dissolving of a Salt in Water, there be any space gained, or not. That is, whether the Bulk of the Water be greater before the Salt lying in it be fully dissolved, than afterwards. For tryal whereof, I took a Bolt-head with a slender Neck, holding somewhat more than a pint; and filling it up to a certain place in the Neck; I then put in an Ounce or two of Salt. And observing the height of the Water, both before it was dissolved, and afterwards; It plainly appeared, that there was some, and that a considerable space, gained by the Dissolution; the Water thereby sinking several Inches below the place, where it stood after the Salt was first put into it.

2. §. From this Experiment it is plain, that there are Vacuities in Water. That is to say, that all the parts of Water are not contiguous, but that either betwixt, or in the Atomes of the Water themselves, there are certain Pores, either absolutely void, or at least filled up with another more subtile body which is easily excluded by the particles of Salt: by possessing the room of which the above said space is gained.

3. §. The Sixth Enquiry is, Whether the space be equally gained, by an equal encrease of the same Salt.

4. §. For this I made two tryals; the first was this. Two half Ounces of Salt Armoniac, being successively dissolved in the same Water; both of them raised up the Water in the Neck of the Bolt-head, equally; the first 3 Inches $\frac{7}{8}$, and so the second.

5. §. The other was this. Four half Ounces of Nitre, being successively dissolved in the same Water, they all of them raised up the Water in the Neck of the Bolt-head, equally; the first a little above two Inches, and the 2^d, 3^d, and 4th, just as much.

6. §. The Seventh Enquiry is, Whether upon the Dissolution of several kinds of Salts, be gained so many several quantities of space. For this I made tryal upon Eleven several Salts, sc. salt of Tartar, Common Salt, Sal Gemmeus, Roman Vitriol, Nitre, White Vitriol, Green Vitriol, Alum, Borax, Loaf-Sugar, and Sal Armoniac; of all which, I dissolved an equal quantity sc. two Ounces, in an equal quantity of Water, severally; that is, taking fresh Water for every Solution. The success was, That the Sal Armoniac raised the Water 15 Inches. The Loaf-Sugar, 13 Inches and $\frac{3}{4}$ ths. The Borax, a Foot. The Alum 11 Inches, and $\frac{5}{8}$ ths. Green Vitriol, 9 Inches and $\frac{5}{8}$ ths. White Vitriol, 9 Inches and $\frac{1}{4}$ th. Nitre, 8 Inches, and $\frac{3}{4}$ ths. Roman Vitriol, 7 Inches and

and $\frac{5}{8}$ ths. *Sal Gemma*, 6 Inches, and $\frac{5}{8}$ ths. *Common salt*, 6 Inches and $\frac{3}{4}$ ths. *Salt of Tartar*, not above 4 Inches and $\frac{1}{2}$ th. All which differences are plain, and most of them very remarkable: Two Ounces of *Sal Armoniac* raising the *Water* near four times as high, as the same quantity of *Salt of Tartar*.

7. §. From this and the fourth Experiment, compared, it also appears, That the several spaces gained by the several *salts*, though sometimes they do, yet do not always answer to the *Solubility* of the said *Salts*. As to give some Instances; *Loaf-Sugar* is the most dissoluble of any other *Salt*; yet it gaineth less space than all the rest, save only *Sal Armoniac*. So *Green Vitriol* is more dissoluble than either *Nitre* or *Common Salt*; yet gaineth less space than either, especially than the latter. And *Sal Armoniac*, which is more dissoluble than *Alum* or *Borax*, yet gaineth less space than either of them. The *Cause* whereof is not easily assigned.

8. §. Note also, that by the same Experiment, as well as by the *Taste* and other Circumstances, it is plain, That *sal Gemma* is nothing else but *Common Salt*, coagulated or *Crystalliz'd* under *Ground*.

9. §. Again, as the Fifth Experiment sheweth, That there are *Vacuties* in *Water*: so doth this Last, that those *Vacuties*, are of differing kinds. Because, otherwise, it should seem, That the *Bulk* of the *Water* would increase, more or less, according to the *Solubilitie* of every *Salt*, and not be alternately differenced as it is; Some *Salts*, more dissoluble, increasing the *Bulk* of the *Water* less, and others less dissoluble, increasing it more. I say, that this difference dependeth not only upon the different *Figures* of the *Atomes* of *Salt*; because then every *Salt* which is more dissoluble, would (quantity for quantity) take up less room in the *Water*: which is contrary to the Experiment.

10. §. From the same Experiment, howsoever *paradoxical* it may seem, yet is it also manifest, That although *Water* be a *Fluid*, yet the *Particles* thereof are *hard* and *consistent*, and unalterable in their *Figure*. Otherwise it is plain, That all manner of *Salts* would be dissolved in the same manner, and take up the same room in the *Water*. For let the *Figures* of the *Salts* be never so various, yet if the *Particles* of *Water* were themselves *Fluid* or *Inconsistent* and *Alterable*, they would always so conforme to those *Figures*, as to fill up all *Vacuties*; and so upon the *Solution* of several *Salts*, if of equal quantity, the *Water* would still retain an equal *Bulk*. As suppose an Ounce of *Iron* were drawn into *Wyer*, another beaten into *Plates*, a third made into *Hooks*, a fourth into *Needles*, a fifth into *Nails*; every one of these five Ounces, being put severally into *Water* will encrease its *Bulk* equally. I conclude therefore, That the *Atomes* of *Water* are hard and unalterable.

11. §. The *Eighth Enquiry* was this, What that just space might be, which any *Salt* gaineth upon *Dissolution*, with respect to its own *Bulk*, or the *Bulk* of the *Water*? For the making of this Experiment, *Water* will not serve, nor yet *Spirit of Wine*; because they both of them dissolve more or less of those *Salts* which are put into them; whereby the observation of the true *Bulk* of the *Salt*, and consequently of the just space it gaineth by *Dissolution* is lost. I took therefore *Oyl of Turpentine*, and pouring it into a *Bolt-head*, marked the place of its ascent in the *Neck*. Then pouring likewise into it two Ounces of
Common

Common Salt, I marked the second ascent of the *Oyl*; and found it to be 10 Inches and 6 eighths. Repeating the Experiment in like manner with two Ounces of *Nitre*, I found the ascent of the *Oyl* to be 11 Inches and $\frac{1}{8}$ th. Repeating it again with two Ounces of *Alum*, the ascent of the *Oyl* was 13 Inches and $\frac{2}{8}$ ths. And making it once more with *Sal Armoniac*, the *Oyl* ascended to 15 Inches: the said several ascents of the *Oyl* being the true spaces which the Four abovesaid *Salts* take. From which, the space which the same *Salts* take up upon *Dissolution*, being deducted; the remainder is the space gained by that *Dissolution*. And so it appears, first, that *Sal Armoniac* gaineth nothing; being the only *Salt* of all I have tryed, which causeth the equal ascent both of the *Water* and the *Oyl* *sc.* just 15 Inches in both. *Alum* causeth the ascent of the *Oyl* to 13 $\frac{2}{8}$ ths, of the *Water*, to 11 Inches and $\frac{5}{8}$ ths: So that it gains about 1 Inch and $\frac{1}{2}$ out of 13. *Nitre* causeth the ascent of the *Oyl*, to 11 Inches and $\frac{1}{8}$ th; of the *Water*, to 8 Inches and $\frac{3}{8}$ ths. So that *Nitre* by *Dissolution* gets almost the space of 3 Inches in 11. *Common Salt* causeth the ascent of the *Oyl*, to 10 Inches and $\frac{6}{8}$ ths; of the *Water*, 6 Inches and $\frac{3}{8}$ ths. So that *Common Salt* gains by *Dissolution* 4 Inches in 10, which is very considerable.

12. §. By this way the *Specifick Gravity* of all kinds of *Salts* may be easily taken, and the difference betwixt them is somewhat surprising. For it appears by the Ascent of the *Oyl*, that *Nitre*, quantity for quantity, is about a 22th part lighter than *Common Salt*. *Alum* about a 6th part lighter. And *Salt Armoniac*, almost a 4th part lighter than *Common salt*. The like estimate may be made of the *Gravity* of all other *Salts*.

13. §. By the same Experiment it also appears, That according to the *Specifick Gravity* of *Salts* they are many times at least more or less *Volatile*; as in the four last *Salts* is plain. For *Common Salt* which of all the four is the most fixed, is also the heaviest. *Nitre* which is somewhat less fixed is somewhat lighter. But *Alum* which is still less fixed is much lighter. And *Sal Armoniac* which is wholly *Volatile*, is the lightest of all the *Salts* above mentioned.

C H A P. III.

Wherein, from the Experiments in the foregoing Chapter, is shewed, the Cause of the Motion of the Mercury in the BAROMETRE.



FOR the doing of this, it will first be acknowledg'd, That not only several sorts of *Sulphur*, but also of *Volatile Salts*, are continually sublimed from most *Bodies* into the *Aer*. So *Lightning*, from the celerity of the accension, appears to be made of a *Meteor*, which is *Nitro-Sulphureous*. *Snow* dependeth upon a *Mixture* of *Nitrous*, and other *Salts*; as is evident, from the regularly and differently *Figur'd Parts*, which compose the whole *Body* of a *Snowy Cloud*, before it clusters into *Flakes*. And one reason, why *Rain* is the best *Water* for any *Soyl*, is because it is impregnated with divers *Volatile* and *Fruitful Salts*. And so from other *Meteors*.

2. §. And next, that these *Salts*, are not always in the same *Quantity*, *Proportion*, and *State*, in the *Aer*: but that sometimes they are more copious; at others, less: sometimes, one more copious, than another: sometimes, more plentifully dissolved; at others, more sparingly: and that, either as they are more or less pure and dissoluble; or according to the quantity of the *Vaporous Parts* in the *Aer*, in which they are incorporated or dissolved.

3. §. Thus much being granted, from the *Experiments* in the foregoing *Chapter* compared together, we may resolve our selves about some *Phenomena* in the *Barometre*. Which seems to vary, not so much with the meer *Weight* of the *Aer*, which hitherto hath been supposed: as by the different pressure it makes, in being *crowded* more at one time, than at another. That is, according as certain *Nitrous*, or other *Saline Bodies*, take up less *space* in the *Aer*, when dissolved in the *Watery Parts* therein, than while they are undissolved.

4 §. And therefore it is especially to be observed, That as the *Mercury* commonly riseth in the *Cylinder* for some days, but always for some time, before the change of the *Weather*, whether for *snow* or *Rain*: So, that then it presently falleth again, even before the *Snow* or *Rain* falls. Whereas, if the *Weight* of the *Aer*, were the only, or the chief *Cause* of the ascent of the *Mercury*; than as it riseth all the while the *Weather* is gathering, so it would keep its standing or heighth, until the *Weather* breaks and falleth down: which yet it never doth, but always falls before it; sometimes no less than a whole day. The *Cause* whereof is, in that all the while the *Mercury* riseth in the *Cylinder*, the *Aer* is *crowded* with more and more *Saline Parts*, which by
the

the *Winds*, or otherwise, are carryed into it; and so causeth it to press upon the *Mercury* in the *Box*: but after that in some time the *Salts* are dissolved or incorporated in the *Aqueous Parts* of the *Aer*, as in *Rain* or *Snow*; so soon as that is done, there is some *space* gaind; and so, before any *Weather* falleth, the *Aer* is less crowded, and presseth less upon the *Mercury* in the *Box*, which gives way to its descent in the *Cylinder*.

5. §. From hence also it is, that the *Mercury* riseth higher with *Cold Winds*, than it doth with those which are *Warme*. Both because that in *cold Winds* there is the greatest quantity of *Nitre*: and that the *coldest Winds*, are usually the dryest. So that the *Nitre* wanting *Moysture* fully to dissolve it; it takes up so much the greater space, and so causeth a greater pressure in the *Aer*, as hath been said.

6. §. *Lastly*, For the same reason it comes to pass, that the *Mercury* first riseth higher, and then falleth lower before *Snow*, than it doth before *Rain*. Because that for the production of *Snow*, the *Aer* is crowded with a greater quantity of *Nitre*, or some other like *Salts*; which before they are dissolved, take up so much the more space; and afterwards so much the less, even before the *snow* falls: as hath been proved.

F I N I S.

A N

I N D E X

O F T H E

Chief Matters,

In which, Id. signifies Idea. An. Anotomy. The Figures before §. the Page. The Figures following §. the Section in that Page.

A.

Acid, commonly the predominant Principle in Plants, 240. §. 8. That is of the Parenchyma. Id. §. 48.

Aer, how to be examined, as relating to Vegetation, Id. §. 60.

Aer, in Plants, How made, An. 93. §. 61. Where it enters the Plant, 127. §. 1. Its Motion and Course in Plants, *ibid.*

Aereal Salt, Id. §. 60.

Aer-Vessels, their Structure, 115. §. 16, &c. See Root and other Parts.

Affinities of Plants, Id. 6. §. 11.

Age of Roots, see Roots.

Agitation, a Cause of Mixture, 230. §. 6.

Akers, 186.

Albumen, see Seed.

Alkaline Salt, in many Plants in their natural estate. 240. §. 9. This the predominant Principle of the true Wood of a Plant, Id. §. 52.

Anagallis, of what Taste, 284. §. 10.

Angellica Roots, when dry, full of Rosin, Id. §. 41.

Anatomy of Plants, why fit to be made, Id. §. 17. In what manner, §. 18. What to be observed thereby, §. 19. Of what use, §. 20.

Animals, their Parts mixed with several Menstruums, 247. to 253. Cantharides, of what nature, 249.

Antimony, of what nature, 245. §. 23.

Apertures of Seeds, An. 2. §. 5. & 200. §. 1.

Apple described, An. 40. §. 2. & 179.

Aperecock, 148.

Aisnart, coded, how its Seed ejaculated,

Y y

The Index.

lated, 188. §. 18.

Arenulæ in Pears, An. 41 §. 4. & 241. §. 20.

Arum-Root, of what Taste, 281. §. 21
The Pestil of what Scent, Id. §. 28.

Aqua-fortis double, mixed with Spirit of Wine, what remarkable thereupon, 242. §. 26.

With Steel, 244 §. 22.

With Tin. 245. §. 27.

Afa scetida, of what nature, 258.

Query, 2.

Ascent of the Trunk, how made, An. 22. §. 21.

A Magnetick Motion, 136.

Ascent of the Sap, how made, An. 24. §. 29. & 126. §. 13.

Asparagus, of what Taste, 284. §. 10.

Attire of Plants see Flower.

B.

B Barbado Nut, Id §. 30.

Barque of the Root, see Root.

Of the Trunk, see Trunk.

Bawme, its Tincture in Water, 274. §. 11.

In Spirit of Wine, 275. §. 14.

Beams of the Sun, different from the Heat of Common Fire, Id. §. 61.

Bean dissected, An. §. 1.

Beech-Wood, An. 20.

Berry, see Fruits.

Bezoar, its nature, 252. §. 49.

Bezoardicum minerale, 245. §. 25.

Bleeding of Plants, Id. §. 23. An. 124. §. 3.

Bolus, what, 242. §. 2.

Bonus Henricus, of what Taste. 284. §. 10.

Bones, their different nature, 249. §. 18.

Branch, how made, An. 28. §. 3.

Its Claspers, An. 27. see Trunks.

Bud of a Branch, how originated, nourished, and kept, An. 28.

§. 1. How kept, 145. §. 2.

Bud of the Seed, see Seed.

Butyr of Flax, Id. §. 51.

C.

C Alamus Aromaticus, of what Taste, 283. §. 6.

Cantharides, their nature, 249. §. 14.

Case of the Seed, of several manners, An. 45. §. 2. & 186.

Carduus green Leavs, their scent, Id. §. 28.

Castor, 250. §. 28.

Celandine, little, where tasted, 284. §. 10.

Cherry, 185

Circulation of the Sap, An. 17. §. 30.

Claspers, An. 27.

Clematis peregrina, the Seed-Case of what Taste, 283. §. 3.

Coats of the seed, see Seed.

Colocynthis. Its nature, 240. §. 13. & 257. Query 5.

Where tasted, 284. §. 8.

Colours of Plants, To what Parts of Plants they belong, Id. 26.

How to be observed, Id. §. 27.

Colours of Roots, An. 94. §. 65. & 270. §. 5.

Of Leavs, 270. §. 6.

Of Flowers, 271. §. 15.

By Infusion, in Oyl, 273. In Water, 274. In Spirit of Wine, ibid.

By their Mixture with other Bodies, 375.

By Cultivation. 277.

Their Causes summed up, 276.

Compression a Cause of Mixture; and of Dissolution, 229. §. 3. 232.

§. 4. & 237. §. 3, 4.

Contents of Plants, in what Method to be examined, Id. §. 21 to 26. & 31 to 47. Of what kind, §. 21.

Their Receptacles, §. 22. Motions, §. 23. Quantities, §. 24.

Consistence, §. 25.

How made in the several Parts of a Plant, An. 92. §. 57.

What in the Seed, 208 §. 15.

Contrayerva, of what Taste, 283. §. 6.

Convolution of the Trunk a Magnetick

netick

The Index.

netick Motion, 137.
 Copper, *its nature*, 245. §. 28.
 Copperas, 246. §. 38.
 Coral, *the Magistery*, 244. §. 15.
Tincture, Id. §. 28.
 Corin *white*, 285.
 Cortical *Body*, *see* Barque.
 Covers of the *Seed*, *see* Seed.
 Cucumer, An. 181.
Leavs of the wild, of what Taste,
 280. §. 11. & 283. §. 6. *Where*
tasted, 284. §. 4, 8.

D.

Daisy *Leaves*, *where tasted*,
 284. §. 10.
Descent of the Root, how made, An.
 34. §. 3.
Diametral Rays, *see* Roots.
Digester, *the nature of that inven-*
ted by Mons. Pappin, 237.
Dissolution of Bodies promoted by
Compression, 237.
Dissimilar Leavs, *see* Leavs.
Dragon Root, 59. §. 13.
Dung of Pigeons, 251. §. 37.

E.

Earth, *how to be examin'd, as*
relating to Vegetation, Id.
 §. 57.
How nature prepares it for the
growth of Plants, 11. §. 8.
Empalement, *see* Flower.
Emulsions, sometimes for Glysters,
 Id. 39.
Enula, of what Taste, 283. §.
 3, 34.
Essential Salts of Plants, see Salts.
Evergreen, 156. §. 2.
Euphorbium, of what nature, 200.
 §. 1. 241. §. 19. & 258. Query
 2. *of what Taste*, 283. §. 6.

F.

FAt, *how made by Art or Na-*
ture, 233. §. 3.
Fermentation, 253. §. 55.
Fibers of the Leaf, see Leavs.
Of the Seed, see Seed.
Figures, of Plants, Id. §. 11.
Of Roots, An. 58. §. 4.
Of Trunks, 135. *Of Leavs*, 150.
 §. 1.
of Seeds, 195.
Figs, their sugar, Id. §. 41.
Flax, its nature, 258. Query 1.
Flower, its Empalement, An. 35.
 §. 2. & 163.
Foliature, its Foulds, An. 36. §.
 5. & 164. §. 1. *Protections*, An.
 36. §. 7. *Hairs*, An. 36. §. 8.
 & 168. §. 8. *Globulets*, An. 37.
 §. 10. & 165. §. 9. *Number of*
Leavs, 165. §. 11. *Parts of the*
Leavs, 166. §. 15. *Use*, An. 37.
 §. 11. & 166. §. 18. *Shape or*
Figure, 167. §. 20.
Attire, Seminiform, An. 37. §.
 13. & 167. *Florid*, An. 38. §.
 17. & 170. *Globulets or Sperme*
of both, An. 38. §. 15. 39. §. 21.
 168 §. 9. & 170. §. 5.
Use of the Atire, An. 39. §. 22.
 & 171.
Flower, when formed, 173.
Colours of the Flower, 271.
How by the Flower to find out to
what sort a Plant belongeth, 175.
 §. 13.
Fœtus, see Seed.
Foulds of Leavs, see Leavs.
Formation of the Root, see Root.
Fruits; Apple, An. 40. §. 2. & 179.
Limon, 180. *Cucumer*, 181.
Pear, An. 41. §. 3. & 182.
Quince, 183. *Plum*, An. 42.
 §. 5. & 183. *Aprecoik*, 184.
Peach, 184. *Cherry*, 185. *Wal-*
nut, ibid. *Grape, ibid.* *Goose-*
berry, An. 43. §. 9. & 185. *White*
Corin, 185. *Filbert*, An. 43. §.
 8. & 186. *Akern*, 186.
The Use of the Fruit, An. 44.
 §. 10.

The Index.

§.10. *Of its Parts to its self,*
189. *To the seed,* 191, & 209.
When the Fruit formed, 192. §.9.
Furr of a Hare, 247. §.3.

G.

G All-Stones, 252. §.47.
Generation of the Seed, and
other Parts, *see the Seed, and o-*
ther Parts.
Gentian Root, *where tasted,* 284.
§.5.
Germen, *see Bud.*
Glysters, *sometimes best made of E-*
mulsions, Id. §.39.
Globulets, *see Leavs and Flowers.*
Gold, *its nature,* 245. §.31.
Gooseberry, 185.
Grape, 185.
Gravel, *its nature,* 251. §.40.
Gums, *of three kinds,* 134. §.15.

H.

H Airs, *see Leavs and Flowers.*
Hares Furr, 247. §.3.
Harts-Horn, 248. §.8.
Hazel Nut, *see Fruits.*
Hellebore black, *of what Taste,*
280. §.12. & 283. §.3. &c. *where*
tasted, 284. §.4.
Hoglice, 249. §.15.
Horses Hoofe, 247. §.5.

I.

J Alap, *of what Taste,* 283. §.6.
284. §.10.
Insertions, *in the Root, and other*
Parts, *see Root and other Parts.*
Iris Root described, 60. §.14.
Irish slate, *its nature,* 243. §.4.
Iron, *see Steel.*

L.

L Apis Calaminaris, *its nature,*
243. §.9.
Lapis Lazuli, 243. §.7.

Lapis Tuthia, 243. §.9.
Lead, *its nature,* 244. §.16.
Lead Spar, 244. §.12.
Leavs, *the two first which come of*
every Seed, what, An. 8. §.42,
&c. *Their Use,* An. 10. §.46.
Leavs; *their Protections,* An.
32. §.17. & 145. §.2. *Foulds,* An.
31. §.14. & 147. §.9. *Shapes and*
Measures, An. 30. §.17. & 150.
§.1. *Globulets,* An. 34. §.7. &
148. §.1. *Hairs,* An. 34. §.4.
149. §.8. *Spots,* 148. §.4.
Thorns, 148. §.6.
Their Compounding Parts, An.
29. §.7. *Skin,* 153. §.1. *Pa-*
renchyma or Pulp, 153. §.5.
Fibers or Vessels, Their Position In
the Body of the Leaf, 152. §.19.
In the Stalk, 154. §.9. *The Lig-*
nous Vessels, 155. §.16. *The*
Aer-Vessels, 155. §.19. *Texture*
of a Palm Leaf or Bag, 156. §.20.
Duration of the Leaf, 156. §.2.
Time and manner of its Genera-
tion, 156. §.4. & 174.
Colour of the Leaf, 270.
How by this to find out to what
sort a Plant belongeth, 174. §.1.
Lignous Body, *see Trunk and other*
Parts
Lilium convalle, *its nature,* Id. §.
30.
Limon, described, 180.
Lithosperm the Seed, *its nature,*
241. §.21.
Lixivial Salts, *see Salts.*
Lobes of the Seed, *see Seed.*
Lympha out of which the Seed is
first nourished, *see Seed.*
Lympheducts *their Structure,* III.
§.30.

M.

M Agistery of Corals, 244. §.15
Of Pearls, 252. §.43.
Mallow, *its nature,* 257. Query 5.
Marine Salt of Plants, *see Salts.*
Mastick, *its nature,* 258. Query. 2.
Measures of Leavs, 150. §.1.
Mechanick uses of Timber, 137.
Membranes

The Index.

Membranes of the Seed, *see* Seed.
 Menstruum of the Stomach, 253.
 § 53.
 Metals; Lead, 244. § 16. Mercury, 244. § 16. Steel, 244. § 20. Antimony, 245. § 23. Tin, 245. § 26. Copper, 245. § 28. Silver, 245. § 29. Gold, 245. § 31.
 Milks of Plants, Id. § 21, & 26.
 How made, An. 67. § 19. & 93. §. 60. & 133. § 12.
 Milk-Vessels, their Structure, 112. § 35.
 Millipedes, 249. § 15.
 Minerals of all sorts, how easily tryed, 247. § 48.
 Mixture; the received Doctrine hereof, 222.
 Its nature explained, 225. Causes, 229. Power and Use, 231.
 Mixture of the Parts of Plants with several Menstruums, 239, &c. Of Minerals. 247, &c. Of Animals, 247, &c.
 Motions, Of Plants, Id. § 16. Of Roots, and other Parts, *see* Roots, and other Parts.
 Of the Sap, *see* Sap.
 Of the Aer, *see* Aer.
 Muciducts, An. 66. § 18.
 Mucilages, Id. § 21. & An. 210. § 4.
 Musk, its nature, 250. §. 29.

N.

Nature of Bodies, how discoverable, 235
 Navel-Fibers, *see* Seed.
 Nightshade deadly, of what Taste, 284. § 9.
 Nitre, of what Taste, 280. § 6.
 Noli me tangere, how the Seed ejaculated, 188. § 18.
 Number of Leavs in Flowers. 165. § 11.
 Number of Seeds, 198.
 Nut Barbado, Id. § 30. & 205. § 17. Hazel Nut, 43. § 8. & 186.

O

OAK-Wood, described, An. 20, & 21.
 Odors of Plants, how to be observed, Id. § 28.
 Some Instances how made, An. 44 § 46. Imitated, 235.
 Olibanum, its nature, 258. Query, 2.
 Oyls stillatitious, how mingled with Water &c. 232. § 7, & 237.
 Oylly Sap, how made, 132. § 6.

P

Parenchyma or Cortical, Pithy, and Pulpy parts of a Plant, their predominant Principle, Id. § 48. Described in the Root and other parts, *see* Root, &c.
 How formed, *see* Roots and Leavs.
 Peach, *see* Fruits.
 Pear, *see* Fruits
 Pearls, their magistry, 252. § 43.
 Philosophy, begins and ends with Theology. 79. § 1.
 Pimpinell, where tasted, 284. §. 10.
 Pith, its structure, 76, §. 7. & 120 § 11, &c.
 Plants, their Natural History how far cultivated, Id. §. 3. Wherein defective, § 2. Fit to be further improved, § 3. & § 63. What to be enquired of, § 6. The usefulness hereof §. 8.
 Plants, their Nature and Virtue how judged of, *see* Virtues.
 Plants, their places of Growth, Id. 15.
 Propotions, § 13.
 Plants, their Parts only Two Essentially distinct, 47. § 14.
 Plants; the general structure of their Parts, 120. § 11, &c.
 Plants, their Principles how to be observed, Id. §. 48.
 For what purpose, § 53. What

L z

predo-

The Index.

pr dominant therein, 240. § 8.
Plants, how to find out to what kind any one belongs, 174
Plum, see Fruits.
Principles of Bodys, 223. *which predominant in the true wood of a Plant.* Id. § 52.
Principles of Principles. Id. § 62.
Protections of the Leaf and Flower. See Leaf and Flower.
Pyrethrum the Root, of what Taste, 281. § --- & 284. § 7.

R.

R Adicle, *see Seed.*
 Raisins, *their Sugar*, Id. § 41.
Rings annual in the Trunk, An. 19. § 6.
Roots; their Original, 57. § 1 *Shapes or Figures, & Sizes* 58. § 4, & 89. § 41, *Motions*, An. 15 § 24. &c. 34. § 3. 59. § 9. &c. 90. § 48 &c. *Ages*, 60. § 16. & 91. § 54.
Parts, the Barque, its Skin, An. 11, § 2. & An. 61.
Parenchyma described, An. 11. § 3. & 63. § 2. *How form'd*, 87. § 34. *Its Diametral rays*, 64. § 7. *Vessls*, 65, 66, 67.
The Wood, Herein the Insertions, An. 12. § 10 & 17. § 28. & 70. § 2. *Lignous Fibers or Vessels*, 70. § 4, 8, 9. *Aer Vessels*, An. 12. § 7. & 71. § 5, 6, 10, &c.
The Pith, An. 13. § 16. & An. 16. § 27. & An. 75, 76. § 7.
Root, how it grows, An. 14. § 23.
The Sap, how imbibed and distributed to its several Parts, 82. § 15 &c. *How circulated*, An. 17. § 27.
How all the parts are form'd, 85. § 26. &c. *And differently disposed*, 88. § 36, &c. *The Colours of Roots*, 170. § 5. *How made*, 94. § 65.
Root of Wormwood, where tasted, 285. § 12.

Rosin, how made by Art, 233. § 4.
Rosin in dryed Roots of Angelica, Id. § 41.
Common Rosin, its nature, 258. Query 2.

S.

S Alt *aereal*, Id. § 60.
 Salt *Alkaline, in many Plants in their natural estate*, 240. § 9.
Salt ammoniac, 246, § 44.
Salt, essential of Plants, How made, 262. § 3, &c. & 265. § 3. &c. *Of several sorts*, Id. § 48. *Instanced also in those of Rosemary, Black Thorn, Scurvey-Grass, Wormwood, Ash*, 865, § 6 &c. *Taste le in good Roubarb*, Id. § 41.
Salt fixed, of what use in Purgation, 260.
Salt Lixivial of Plants, how imitated, 233. § 6. *Of different nature*, 264. § 2.
Salt of Ash, of what nature, 167. § 22. *of Tartar, lb. Yielded in different quantities by the Barque of Ash, Rosemary, Black-Thorn, Agrimony*, 256. Query 1. *Garden and Sea Scurvy-grass*, 256. Query 2. *Mint distill'd, and not*, 256, Query 3. *Majorane, Oak-Barque, Liquorish, Aniseeds, Sorell, Garden Scurveygrass, Mint, Sea Scurveygrass*, 256. Query 4. *Majorane, Agrimony, Mugwort, Mint, Mallow, Roubarb, Sena, Jalap, Colocynthis*, 257, Query 5. *Flax*, 258. Qu. 1. *Gum Arabick, Euphorbiurn, Myrrh, Opium, Aloe, Scammony, Gutta gamba*, 258. Query 2.
Salt Marine, its nature, 246. § 43.
Salt Marine of Plants, how made by Nature or Art, 234. § 8, 263 § 12, &c. 266. § 16. *Of several sorts, instanced in those of Rosemary, Scurvey grass, Black Thorn, Wormwood*, 266. § 17 &c.
Salt of the dead Sea, 263. § 14.

Saps

The Index.

- Saps of Plants, *how to be observed*, Id. § 21. to 26. and 31 to 47. Their several kinds, § 21. Receptacles, § 22. Motions, § 23. Quantities, § 24. Consistence, § 25.
- Sap, *how imbibed, and distributed to the several parts of the Root*. An. 82. § 15 &c. Its Circulation therein, An. 17. § 29. where, and how it ascends in the Trunk, An. 24. § 29. & An. 124.
- Sap and other contents of the several Parts *how made*, An. 92. § 57 &c. & 131. *How a Milky Sap*, An. 67. § 19. & 93. § 60. & 133. § 12 &c. *How a Winny*, 93 § 62. & 132. § 3 &c. *How one very Oily*, 132. § 6 &c.
- Scurveygrafs Garden, *of what Taste*, 283. § 6.
- Scurvey-grafs Sea, *its Nature*, 256. Query 4.
- Seasons of Plants, Id. § 14.
- Secundine *see Seed*.
- Seeds; *their Case or Uterus*, An. 45. § 2. *of several manners*, 186. Figures, An. 45. § 3. &c. 195. Numb r, 198. Motions, 188. § 18 & 199 § 3, &c. Stones, 201. § 2. & 209. Mucilages, 201, § 4. Coats or Membranes, An. 2. § 3. 45. § 3, &c. 46 § 10. 47. § 15 & 201. § 6. 210 &c. Apertures, An. 2. § 5. & 200 § 1.
- Vitellum, 20. 2. § 9. *The Vætus, or true Seed, its Radicle and Lobes*, An. 2 & 3. § 9 to 12. & 203. *Plum.e or Bud*, An. 3. § 13 & 206. *Skin*, An. 4. § 16. & 207. § 9. *Parenchyma*, An. 4. § 18. & 207. § 10. *Seminal Root or Vessels*, An. 5. § 21, &c. & 207. § 11, &c. *Navle Filer*, An. 48. § 17. & 212. Content. 208. § 15. *The manner of its Vegetation*, An. 6. § 30. &c. *Of its Generation*, An. 48. § 18 &c. & 209 &c.
- Shape of Roots, and other Parts. *see Roots and other parts*.
- Shells, *their Nature*, 248. § 9, 10, 11.
- Skin, *see Seed and other Parts*.
- Silver, *its Nature*, 245 § 29.
- Smell of green Carduus, Id. § 28. *Of the Pestil of Arum*. Ib.
- Soyl. *see Earth*.
- Sperme of Plants. *see Flower*.
- Spirit of Salt, 247. § 46.
- Spirit of Salt Armoniac, 247. § 47.
- Spirit of Peas-Cods, Id. § 30.
- Spirits urinous, *how made less offensive*, Id. § 45.
- Spirit of Wine mixed with Aqua fortis, *what thereupon remarkable*, 242. § 26.
- Stalks. *see Trunks and Branches*.
- Steel, *its nature*, 244. § 20. *Mixed with double Aqua fortis, what thereupon remarkable*, 244. § 22.
- Stillatitious Oyls, *how mixed with water*, 232. § 7. & 237.
- Stomachick menstruum, 253. § 53.
- Stones; *a strange one bred in the Stomach*, 252. § 48. *Others probably bred there*, 253. § 51. *Of the Kidneys or Bladder, of what nature*, 251. § 32. *How prevented*, 251. § 41. & 252. § 42.
- Gall Stone, *its nature*, 252. § 47.
- Bezoar, *its nature*, 252. § 49.
- Lead-Spar, 244. § 12.
- Lapis Calaminaris, 243. § 9.
- Tuthiæ, *ibid*.
- Lazuli, 244. § 12.
- Structure of a Plant. 120. § 11, &c.
- Sugar of Raisins and Figs. Id. § 41.
- Sulphur predominant in the true wood of a Plant, *id*. § 52.
- Sun, *its Influence on Plants how to be examined*. Id. § 61.

Tastes.

The Index.

T.

TAsts of Plants; how to be observed, Id. § 29. *Simple*, 280. § 6. *Compounded*, 281. *Their Degree*, 282. *Motions or Terms*, 283. *Seat*, 284. *Causes*, An. 95. § 68 &c. & 286 &c.

Tast of Arum Root, 281. §
Asparagus Root, 284. § 10. *Bonus Henricus*, 284. § 10. *Calamus Aromaticus*, 283. § 6. *Celandine little*, 284. § 10. *Clematis peregrina the Seed-Case*, 283. § 3. *Colocynthis*, 284. § 8
Contrayerva, 283. § 6. *Cucumber wild, the Leavs*, 284. § 4, 8.
Daisy Leavs, 284. § 10. *Enula*, 283. § 3, 4. *Euphorium*, 283. § 6. *Gentian Root*, 284. § 5, 8.
Hellebore black, the Root, 280. § 12. 283. § 3. 284. § 4.
Jalap, 283. § 6. 284. § 10.
Nightshade deadly, 284. § 9. *Nitre* 280. § 6. *Pimpinell*, 284. § 10. *Pyrethrum the Root*, 281. § & 284. § 7. *Tamarisk-Leavs*, Id. § 29. *Wormwood-Root*, 285. § 12. *Yarrow*, 283. § 6.

Texture of a Plant, 120. § 11, &c.

Thorns, their kinds, An. 33. § 1.

Timber, see Trunk.

Tin, its nature, 245. § 26.

Tin mixed with strong Aqua fortis, what thereupon observable, 245. § 27.

Tincture of Bawme in Water, 274. § 11. *In Spirit of Wine*, 275. § 14.

Tincture of Corals, a cheat, Id. § 28.

Trunks, and Branches several described as they appear to the naked Eye, sc. of Indian Wheat, Dandelion, Borage, Colewort, Holy-oak, wild Cucumer, Scorzonera,

Burdock, Endive, Vine, Sumach 103, &c.

Trunk, skin, An. 19. § 2. & 107. § 2 to 5.

The Barque, An. 19. § 3.

Its Parenchyma, 108. § 7.

Vessels, 108. § 8. to p. 113.

The Lympheducts their Structure, III. § 30, &c.

Milk-Vessels, their Structure, 112. § 35, &c.

Different Surface of the Barque how made, 129. § 4.

How united to the Wood, 129, § 2, 3. *How this always keeps moist, not the Pith*, An. 20. §

7. & 93. § 58. & 124. § 2.

The Wood, An. 19. § 4. to 11. & An. 20 & 21.

Its Annual Rings, An. 19. § 6.

Insertions, An. 19. § 5. 12. to 15. 17. & 128. § 8. &c.

True wood, 114. § 10, &c.

How dilated, An. 22. § 22, 23.

And why, § 24, &c.

Aer-Vessels, An. 20. § 8, 9. & 115. § 16, &c.

How less in the Trunk, than in the Root, and whence formed late in the year &c. 130. § 10. &

131. § 16.

The Pith, An. 19. § 5, 18, 19, 20. & 119. to 122 & 129. § 5. &c.

Trunks, their different structure whence, 129.

Shapes, whence, 135.

Motions, whence, An. 22. § 21. & 136.

Trunks, how fitted for Mechanick use, 137.

Trunks, of their bleeding, Id. § 23. & 124. § 3, &c.

Trunk-Roots, An. 27, 28.

Turnep, described, An. 13.

Valves,

The Index.

V.

VAlves, *no where in Plants,*
An. p. 21. §.16.
Vegetables, *see* Plants.
Vegetation of the Seed, *see* Seed.
The manner of Vegetation, how
judged of, Id. §. 53.
Vessels of the Root and other Parts,
see Root and other Parts.
Virtues of Plants, *how to be obser-*
ved and judged of, Id. §.12,30,
47. & p. 236. 290.
Imitable, 235.
The reason of them, how knowa-
ble, Id. §.55.
Vitriols, *their nature, 246. §.38.*
Uterus of Plants, *fleshy or mem-*
braneous, 186.

W.

WAllnut, 185.
Water, *how to be examined*
as relating to Vegetation.
Water, *how mingled with stilla-*
titious or other Oyls, 237.
Wood of the Root and other Parts,
see the Parts.
Wood of Beech, An. 20. & 21.
Of Oak, *Ibid.*
Its predominant Principle, Id.
§. 52.

Y.

YArrow Leavs, *their Tast, 283*
§. 6.

THE
EXPLICATION
OF THE
TABLES,

Reduced to a narrow compass; as serving to clear those Particulars, chiefly, which the *Descriptions* before given, have not reached.

The TABLES to the First BOOK, are Four.

- TAB. I. **F**igure 1, a, The Foramen.
 F. 2, a, the Radicle lodged in the Body of the Iner Coat.
 F. 3, a, the Radicle, b, the Plume or Bud.
 F. 4, the Seed covered; c, the Seed open; e, the same magnified.
 F. 5. a, the Corn covered; c, naked and a little magnified.
 F. 6. a, b, the two Lobes; e, the Radicle; c, the Radicle and Bud; d, the Hollow in which the Bud lies.
 F. 7. a, the Seed covered; c, naked; e, open.
 F. 8. a, one Lobe; --b, the Bud; b, magnified.
 F. 9. the Slice a little magnified.
 F. 10. The Radicle d, cut transversly c.
 F. 11. The Plume or Bud a, cut transversly c.

F. 12. Cut by the Length.

F. 13. A Lobe cut transversly.

F. 14. Both the Lobes pared by the Length, to shew the Seminal Root.

F. 15. a, the convex side of one Lobe, shewing the Seminal Root without cutting; c, the flat side.

TAB. II. F. 1, 2, & 3. shew the gradual conversion of the Lobes of the Seed, into Leaves.

F. 4. a, the Radicle cut by the length; b, transversly.

F. 5. The white Wedges, are the Insertions; the black, are the Wood; the pricks are the Aer-Vessels; and the black half ovals, the Lympheducts in the Barque.

F. 6. The three black Rings, are the terms of three years growth.

F. 7. a, the upper part; b, the lower.

F. 8. A Turnep cut transversly, and part of the Rind cut off.

F. 9. sheweth the gradual growth of the Pith.

TAB. III. F. 1. The Bud cut transversly, and part of the Radicle by

The Explication of the Tables.

by the Length, in a Bean newly sprung up.

F. 2. sheweth the Wood as it appears to the naked Eye.

F. 3. the Cane split down.

F. 4. the Corn newly sprouted.

F. 5. A Branch of five years growth. From the Circumference, to the utmost black Ring, goes the Barque.

F. 6. a, a piece of the Stalk; b, magnified.

F. 7. a, a piece of Oak wood cut transversly; b, the same magnified. The white Lines are the lesser and greater Insertions. The Pricks, are the Wood. The little and great Holes two sorts of Aer-Vessels.

F. 8. Part of a Branch ten years old, with the Barque stripped off, and cut both transversly and down the length, to shew how the Barque is inserted into the Wood.

TAB. IV. F. 1. shewing how the Insertions appear, in a piece of Beech-Tree split down, to be braced or woven in together with the Wood.

F. 2. to 11. shew the different position and Figure of the Lignous Fibers.

F. 12. a, one of the Theca Seminiformes in a Lily, with the spermatick Powder therein, as apparent to the naked Eye.

F. 13. a, one of the suits in the Florid Attire, as it appears to the naked Eye; b, the Floret; c, the Sheath; d, the Blade.

F. 14. Wherein the white Pentangular Acetary is bounded by the Calculary.

F. 15. The Branches which run through the Stone to the Flower and Seed.

F. 16. The Innermost Cover of the Seed, as shaped when it is ripe.

F. 17. The Coats cut open.

F. 18. The Seminal Root.

The TABLES to the Second BOOK are Thirteen.

TAB. V. sheweth the generation of Roots out of the Descending Trunk. So F. 6. is a treble Root of three years descent; the lowermost, half-roted off.

TAB. VI. F. 1. sheweth the Surface of the Barque.

F. 2. the middle part.

F. 3. the Barque striped.

F. 4. the Root cut down the length.

F. 5. the Barque striped off.

F. 6. the Network both of the Lympheducts, and of the Aer-Vessels.

F. 7. the Generation of a Bud.

F. 8, 9, 10, 11. The Root split down, to shew the Position of the Vessels, and the Figure of the Pith at the top of the Root.

TAB. VII. The Roots all cut transversly, and their Varieties described, in the second Book, as they appear to the naked Eye.

TAB. VIII. Other Roots cut transversly, and the varieties of their Parts also described in the second Book.

TAB. IX. More Roots cut transversly.

TAB. X. F. 1. A Slice of the Root cut transversly; but a little too big for the life.

F. 2. AA, One half of a like Slice.

bb, The Skin.

AA DD, The Barque or all that part of the Root analogous to it.

GD, The Lympheducts on the inner edge of the Barque.

GG, The Wood.

GT, The Aer-Vessels therein.

TT, The Pith.

TAB. XI. F. 1. The Neck of the

The Explication of the Tables.

the Root cut transversely.

F. 2. One half of the same split down.

F. 3. Magnified.

AB, The Skin.

AE, The Barque.

EE, The Lympheducts.

The black Columns under them, are the Wood.

The Holes in the Columns are the Aer-Vessels.

The white Columns EL, are Insertions betwixt the Barque and the Pith.

Le, The Pith.

ee, The angular Bladders of the Pith.

T A B. XII. A, one half of F. 1. magnified.

Ab, The Skin.

AG, The Barque, or all that part of the Root which answers to it.

In which the round black spots, are the Muciducts.

DG, The common Lympheducts.

DT, The Pithy Part of the Root.

TT, More Lympheducts.

In both which, the black Holes are the Aer-Vessels.

T A B. XIII. A, One half of F. 1. magnified.

AC, The Skin.

AG, The Barque, or that part of the Root which answers to it.

DD, The Milk-Vessels placed in Rings.

EE, The Parenchymous Rings betwixt them.

GT, The Bladders streaming in Rays, by the mixture of the Lympheducts with the Lacteals.

GG, To the Centre, the Wood.

In which the Holes are the Aer-Vessels.

T A B. XIII. A b., The Skin, which should have been thicker.

AF, The Barque.

Gb, The Bladders in the outer part of the Barque, oblong and

postured circularly.

SS The Bladders in the inner part, standing in Arches.

FF, A Ring of Sap-Vessels.

dd, Parenchymous Insertions.

dLd, The Wood.

In which; the Holes edged with white Rings are the Aer-Vessels.

T A B. XV. AA, The Skin.

AB, The Barque.

BL, The Sap-Vessels in the form of a Glory.

BE, The Wood.

In which, the Holes are the Aer-Vessels.

GE, A Ring of more Sap-Vessels.

EE, The Pith.

T A B. XVI. Ab, The Skin.

AC, The Barque.

In which the round Holes B, are Balsame-Vessels.

B. C. Parcels of Lympheducts.

In which there are more Balsame-Vessels.

CD, Parenchymous Insertions.

DE, Parcels of Wood,

In which the Holes are the Aer-Vessels.

T A B. XVII. A, the Skin.

AB, The Barque.

LS, A parcel of Sap-Vessels.

Ll, A Parcel of Wood.

In which the Holes great and small are Aer-Vessels.

BB, Parenchymous Insertions betwixt the parcels of Wood.

DD, Others within them.

The TABLES to the Third BOOK are 23.

T A B. XVIII. Hereof see the Description in the Third Book, Chap. 1.

T A B. XIX. F. 1. A Branch of Corin Tree.

A, sheweth the surface of the Barque. B.

The Explication of the Tables.

B, Of the wood.
 F. 2. Stalk of *Sonchus* split down.
 F. 3. Branch of *Vine* split down.
 In both, the several Storys or Chambers of the Pith.
 F. 4. Branch of *Walnut*. A. an older. B, a younger: in both, the Pith parted into transvers Membrans.
 T A B. XX. F. 1. Sheweth the Surface of a *Walking Cane*.
 And the Clusters of *Aer-Vessels*, surrounded with Rings of *Succiferous*.
 F. 2. The surface of the skin of *Borage Stalk*.
 F. 3. The *Turpentine Vessels* running through the length of the *Barque*; one of them cut down the middle, the other entire.
 F. 4. The *Milk-Vessels* shewed in the same manner.
 T A B. XXI. Sheweth the *Woody and Aer-Vessels* by the length of the Branch, part of the *Barque*, and wood, being taken away.
 T A B. XXII. A B, The *Skin*.
 A C, the *Barque*.
 Q, the *Parenchymous part*.
 H I, *Parcels of Muciducts* in a Ring.
 D C, *Common Lympheducts*.
 C D E F, the *Wood of 3 years growth*.
 K L M N, *The second years growth*.
 O z, the *great Insertions*.
 P z, the *smaller*.
 X X, *Lignous parcels*.
 Within which the *Holes* are the *Aer-Vessels*.
 E F G, the *Pith*.
 T A B. XXIII. A B, the *Skin*.
 A C, the *Barque*.
 Q, the *simple Parenchyma*.
 H I, a *Ring of special Vessels*.
 P, *common Sap-Vessels*.
 C D E F, the *Wood of 3 years growth*.
 K L M N, *one years growth*.
 X, *great Insertions*.
 P O, *lesser between them*.

The black parcels are the wood.
 In which the *Holes* are the *Aer-Vessels*.
 E F G, the *Pith*.
 T A B. XXIV. A B, the *Skin*.
 A B C D, the *Bark*.
 N N, the *Parenchyma*.
 H I, a *Ring of special Sap-Vessels*.
 D M C, *Parcels of Lympheducts*.
 C D E F, the *Wood*.
 E F L K, *one years growth*.
 K P Q L, the *larger Aer-vessels* in the several parcels of *Wood*.
 z z, the *lesser Aer-vessels*.
 M T, the *insertions*.
 E F G, the *Pith*.
 T A B. XXV. A B, the *Skin*.
 A B C D, the *Barque*.
 H I, *special Sap Vessels* in arched parcels.
 O O, the *common Sap-vessels* which begin to turn into *Wood*.
 C D E F, the *Wood*.
 K L M N, *one years growth*.
 The *Holes* are the *Aer-vessels* in the wood.
 z z, the *true wood*.
 O z, O y, the *Insertions*.
 E F, *other Sap-vessels*.
 E F G, the *Pith*.
 T A B. XXVI. A B, the *skin*.
 A B C D, the *Barque*.
 Q Q, the *Parenchyma*.
 H I, *special Sap-vessels* in arched parcels.
 D C, a *Ring of common Lympheducts*.
 D C F E, the *Wood*.
 K L M N, *one years growth*.
 The *Holes* are the *Aer-Vessels*.
 O O, the *greater Insertions*.
 P O, the *smaller*.
 E F, *other Sap-vessels*.
 E F G, the *Pith*.
 T A B. XXVII. A B, the *Skin*.
 A B C D, the *Barque*.
 W V, the *Parenchyma*.
 H I, *round parcels of Sap-Vessels*.
 D C, the *common Sap-Vessels*.
 D C E F, the *Wood of 5 years growth*.
 Bbb Q R F E,

The Explication of the Tables.

- Q R F E, one years growth.
 X X, the true wood.
 The Holes both great and small
 are the Aer-Vessels.
 S S, The great Insertions.
 T S, the smaller.
 E F G, the Pith.
 T A B. XXVIII. A B, the Skin.
 A B C D, the Barque.
 H T I, special Sap-Vessels in
 round Parcels.
 D S C, common Sap Vessels.
 D C E F, the Wood of five years
 growth.
 e e, the true wood.
 K L &c. the great Aer-Vessels.
 D C, the smaller.
 S S, the Insertions.
 E F S, the Pith,
 T A B. XXIX. A B C D, the
 Barque.
 A B, a Ring of Sap-Vessels in
 round parcels next the Skin.
 H I, the Parenchyma.
 Another Ring of round parcels.
 D O C, Common Lympheducts.
 D C E F, the wood.
 M N E F, one years growth.
 S S, the true wood.
 K L, the great Aer Vessels.
 P Q, the lesser.
 O O, the Insertions.
 E F G, The Pith.
 e, the Bladders of the Pith.
 T A B. XXX. A B, the Skin.
 A B C D, the Barque.
 R R, the Parenchyma.
 H R I, two Rings of special Sap-
 Vessels.
 D C, Common Lympheducts.
 D C E F, the wood of four years
 growth.
 d d, the true wood.
 Q d, part of it whiter, by the
 mixture of special Sap-Vessels repre-
 sented by the transvers Lines.
 M N, the great Aer-Vessels.
 c e, parcels of lesser ones.
 E F, a Ring of other Sap-Vessels.
 E F G, the Pith.
 T A B. XXXI. A B C D, the
 Barque.
- m m, the Parenchyma.
 H m I, Milk Vessels in arched par-
 cels.
 D K C, Lympheducts.
 D C E F, the wood of one years
 growth.
 S T, probably milk Vessels hereto-
 fore.
 The Holes in the Aer-Vessels.
 K K, the Insertions.
 E v F, other Milk-Vessels.
 E F G, the Pith.
 T A B. XXXII. A B C D, the
 Barque.
 M N, The Parenchyma.
 D L C, the Lympheducts.
 H I, The Vessels which carry the
 Turpentine.
 D C F E, the Wood.
 L L, the Insertions.
 E F G, the Pith.
 The greater Holes both in the Wood
 and Pith, are more Turpentine Ves-
 sels.
 T A B. XXXIII. A B C D, the
 Barque.
 X Y, The Parenchyma.
 K X Y L, Special Vessels in round
 parcels.
 H I, others in a Ring.
 D C, Common Lympheducts.
 D C E F, the wood.
 S Z T, probably one sort of Sap-
 Vessels heretofore in the Barque.
 Q M Q N, small Aer-Vessels.
 M N, great Aer-Vessels.
 R Q, the small Insertions.
 Q Q, the great ones.
 E F G, the Pith.
 T A B. XXXIV. A B a a, the
 hairy Skin.
 A B C D, the Barque.
 H w I, the Parenchyma.
 D M C, the common Lymphe-
 ducts.
 K L, the Milk Vessels. v, one Ves-
 sel.
 H I, Another sort of Lympheducts,
 arched over the Milk Vessels.
 X X, seems to be a third sort of
 Lympheducts.
 D C F E, the Wood.

The Explication of the Tables.

MM, the Insertions.
 XX, the true Wood.
 The Holes therein are the Aer Vessels.
 EF, a Ring of Lympheducts.
 EFG, the Pith.
 TAB. XXXV. ABCD, the Barque.
 AMB, the Parenchyma.
 HMI, Balsam Vessels.
 KL, another sort of Sap-Vessels in parcels.
 KLDC, Lympheducts.
 DCEF, The Wood.
 In which the Holes are the Aer-Vessels.
 MM, the Insertions.
 EF, more Balsame-Vessels.
 EFG, the Pith.
 TAB. XXXVI. aa, part of a Vine-Branch cut transversly, and also split half way down the middle.
 BB, The same magnified. Shewing the Position of the Bladders in the Barque and Pith in perpendicular Rows; in the Insertions, in Horizontal Rows.
 And the Vessels or Parcels of Wood not rased as in many other Trees.
 TAB. XXXVII. Sheweth the bracing of the Vessels. And how the several Parcels of Vessels or Wood are interwoven with the Insertions.
 TAB. XXXVIII. ABCD, the Barque.
 HI, The Parenchyma.
 ee, A sort of Sap-Vessels.
 aa, Another sort.
 cc, Milk Vessels.
 DCEF, the Wood.
 VV, the Aer-Vessels.
 tt, More Lympheducts.
 ff, More Milk-Vessels.
 at, The Insertions.
 EFG, The Pith, composed of angular Bladders, the Bladders of Threds, and the Threds of single Fibers.
 g, One of the single Fibers.
 TAB. XXXIX. Sheweth the structure of the Lympheducts or of the Lignous Fibers both in the

Barque, and the Wood.
 F. 1. a, & F. 2, A single Vessel in the Barque of Flax, composed of a great number of other Lignous Fibers; with which also the Parenchymous are intermixd. Not visible, except very highly magnified.
 F. 3. A parcel of the same Vessels in Wood.
 F. 4, & F. 5. shew the manner of the Ascent of the Sap, both in the Lympheducts, and in the Lactiferous and other larger Vessels.
 TAB. XL. The Fibers which hang down from the Barque are the Lympheducts; one of which is composed of a great many other smaller Fibers.
 The large Tubes are the Milk-Vessels composed of Bladders.
 The Fibers which hang down from the wood, are some of them the old Lympheducts turn'd to wood.
 And some, Aer-Vessels unroav'd.
 The thin Plate between the two wedges of wood, is one of the Insertions, composed of Bladders, and those Bladders of Threds.
 The remainder, is part of the Pith, composed of Thredy or Fibrous Bladders.

The TABLES to the
 Fourth BOOK are 42.

TAB. XLI. b, a Dock-Leaf covered with the Veil.
 d, the Leaf naked.
 ac, the Veil spread open.
 In Clary, the Bud is embraced by the Curled Leaves.
 In Sumach, the Bud lies within the Stalk, as an Egg or Kernel within a Shell.
 TAB. XLII. F. 1. sheweth how the Pipes are inclosed one within another.
 F. 2. a, the Leaf foulded up. b, opened.
 F. 3. a b, the Bud. b, a little magnified

The Explication of the Tables.

magnified.

F. 4. a b, the Leaf rowled up, inward. c, a little magnified and cut transversly, to shew the Rowl.

F. 5. a, the Leaf rowled up. b, magnified and cut transversly.

F. 6. a, the Leaf rowled backward. b, magnified and cut transversly.

T A B. XLIII. F. 1. sheweth the Tenter-Hooks, by which the Leaf climbs.

F. 2. sheweth the Globulets, turned to a white powder.

The Leaf of Jerusalem Cowslip. sheweth the Way of the Insect under the skin.

T A B. XLIV. & XLV. sheweth the Measures of Leavs by the Circumference.

T A B. XLVI & XLVII. sheweth the proportion between the chief Fibers; and also the Angles they usually make together.

T A B. XLVIII. F. 1, 2, & 3. shew the Apertures in several Leavs.

F. 4. sheweth the same. And likewise, the peculiar composure of the Bladders and Fibers of the Leaf.

T A B. XLIX. sheweth the difference in the Bladders, and in the Position of the Lignous Fibers in the Stalks of Leavs.

T A B. L. sheweth the Pulp of a Borage-Leaf and many others composed of Bladders; the sides of which Bladders, are made of other smaller ones.

And the distribution of the Lignous Fibers (and of the Aer-Vessels sheathed within them) not like that of Veins in Animals, but of the Nerus, &c. See the description of the Leaf.

T A B. LI. F. 1. The appearance of the Aer-Vessels like Cobwebs to the naked Eye, upon breaking the Leaf.

F. 2. A small peice cut off of the Leaf.

F. 3. The same magnified in which the same Vessels look like spiral wyers

stretched out.

F. 4. The same as they stand entire within the Wood.

T A B. LII. Representeth the Aer-Vessels of Scabious, as in Tab. LI.

T A B. LIII. Sheweth the manner of the Generation of the Leaf; chiefly, by the help of several Salts, where-with the Sap is impregnated.

F. 5. (1) The Foundation of the work.

F. 6. (1 & 2) strengthened.

F. 7. (1 & 3); in which (3) is set with the square end to end: and with the poynt-side of one, to that of another.

F. 8. The same, directing the Position of the Lignous Fibers at very Acute Angles.

F. 9. At less Acute Angles.

F. 10. The greater Fibers at Acute, and the smaller at Right Angles.

F. 11. The greater at Right Angles with the help of (1) (2) or (3).

F. 12. (3) directing the Fiber in the Edge of the Leaf into a greater Circle.

F. 13. Into a less, and with divers Diameters.

F. 14. (4) directing the Parenchymous Fibers in making the Bladders.

F. 15. In winding from one Bladder to another.

F. 17. Or about the Lignous Fibers.

F. 16. In making the Aer-Vessels. T A B. LIV. sheweth how Nature manages the Folds of Flowers according to their Shape.

T A B. LV. F. 1. sheweth the Edges of the Leaf fastned by their Indented Hairs.

F. 2. The Balsamick Knobs in the place of Hairs.

F. 3. The number 5 running 3 times into its self in 13.

F. 4. And five times in 21.

F. 5. &c. The Seminiform Attire in Clematis Austriac a. With one of the

The Explication of the Tables.

the *Theca* magnified; of which, there are about 30 or 40 in one Flower.

F. 8, &c. The same in *Blattaria*, with one of the *Theca* magnified; of which are there about 5 in one Flower.

T A B. LVI. The same in yellow *Henbane*.

With one of the *Theca* magnified; of which there are about 5 in one Flower.

And the Column on the top of the Seed-Case.

T A B. LVII. The same in *St. Johns wort*, entire, together with the Seed-Case or *Uterus*.

T A B. LVIII. The Varieties of the Spermatick Particles in the Seminiform Attire.

T A B. LIX. The Florid Attire of Golden Rod;

In which, the several suits consist but of two pieces. And of which Attire, the Flower doth almost wholly consist.

T A B. LX. F. 1, &c. The same Attire in French Marigold or *Flos Africanus*, with one suit magnified. Of which, there are about 12, in one Flower; and every suit consisting of 3 Pieces.

F. 5. One of another Flower, consisting also of 3 Pieces.

T A B. LXI. One suit of the same Attire in Marigold, and *Knapweed*, each of them consisting of three Pieces.

F. 5. a, The Attire of one Piece, proper to each Leaf in a Marigold Flower, besides that in the bosome of the Flower.

F. 8. a b, the Seed-Case or *Uterus* at the bottom of every suit.

T A B. LXII. The Attire (of 3 Pieces) proper to each Leaf in the Flower of *Cichory*.

T A B. LXIII. Sheweth the Flower of *Mezereon* perfectly formed in all its Parts, in the year before it appears. But differs in Shape, as a *Fætus* doth when newly formed.

T A B. LXIV. Sheweth the same

in the Flower of *Alarum*.

T A B. LXV. Sheweth the position of the 20 chief Branches in an Apple.

Their Production from the Stalk to the Seeds and Flower.

And a part of the *Parenchyma* magnified, sc. that which is pricked out from the Coar to the Skin; shewing the oblong Figure of the Bladders, and the Divisions in every Bladder.

T A B. LXVI. F. 1, &c. Sheweth the Bladders in the Rind of a Limon containing the Oyl.

The Bags and Bladders of the Pulp, containing the sower Juyce.

And the Position of the Vessels belongeth to the Fruit, Seed, and Flower.

F. 5. Shews the same Vessels, and treble *Parenchyma* in a Cucumer.

T A B. LXVII. Representeth the Parts of a Pear.

The position and production of the Vessels.

The Chanel from the top of the Pear to the botome of the Coar.

The Tartareous Knots.

And the Bladders radiated to them.

T A B. LXVIII. See the Descriptions of Fruit; and the last Chapter of the Generation of the Seed.

T A B. LXIX. F. 5. Sheweth the Parts of a Goosberry.

The darker part is the sower Rind. Consisting of two sorts of Bladders, of which some very small, and others very great.

The white pieces on the circumference of the Berry, are the Lignous Fibers.

The two opposite white and radiated Bodys are the Midle *Parenchyma*.

And the oblong Bags round about the several Seeds or Seed-Cases, are the sweet Pulp.

T A B. LXX. Sheweth the Seed-Case of Radish opened, and the Seeds hanging on two Ropes.

That of Poppey both entire, and
C c c split

The Explication of the Tables.

split down the middle.

A slice of the Cod of Garden-Bean, while very young; and therein the Bladders and Threds of the Spongy Parenchyma.

And the gradual ripening and opening of that of yellow Henbane.

T A B. LXXI. sheweth the Seed-Case of Tulip entire, cut transversely, and split downe.

A slice of Thorn-Apple, or of the Seed-Case of Stramonium, while young.

That of Pimpinell naturally divided into two Hemisphers; with the Button, on which the Seeds grow, erected in the middle.

The manner of the ejaculation of the Seed, in Coded Arsmart.

And the Coats of the Seed of Azarum formed the year before it ripens.

T A B. LXXII. sheweth the measures of Plum-stones.

The Apertures, and Divisions, of the covers of the Seed.

The Seed and Seed-Case of Harts Tongue, opened with a Spring.

And other contrivances both for the Motion, and Arrest of other Seeds.

T A B. LXXIII & LXXIV. See the Descriptions.

In Tab. 74. the corners and edges of that of Fox-glove should have been rounder.

The Figures are all done pretty near a Scale.

T A B. LXXV. The Belly and Back of a Datestone, and the small sprouting Node taken out of the Hole in the back cut open.

The Shapes and Foulds of divers Seeds.

The Vitellum of Orach, and Rhapsontick.

In great blew-Lupine, d, the Navle; b, the descending part of the Radicle.

T A B. LXXVI. Flag. 1, the Seed. 2, split open. 3, the true Seed which lies in the hollow made in the Cover (2) 4, one half of

(2) magnifyd. 5, the Seed (3) magnifyd.

Purging Angola Nut. 1, with the shell on.

2, taken off. 3, the soft Cover split down. 4, the Seed which lies in it; the Lobes hereof answerable to two Leaves, and Radicle to the Stalk.

And so in the rest.

T A B. LXXVII. Coffee Berry stone. 1, The belly of the Stone. 2, the Black. 3, pared a little. 4, the Kernel taken out of it. 5, the same magnifyd.

Goosgrass. 1, the entire Seed. 2, the back of the hard Cover. 3, the belly. 4, cut in two. 5, the same magnifyd. 6, the true Seed taken out of it.

Staphisagria. 1, the entire Seed. 2, the hard Cover. 3, Split in two. 4, the true Seed taken out of it. 5, The same magnifyd.

Peony, 1, the Seed commonly so call'd. 2, one half of it split down. 3, the other half. 4, the true seed taken out of it. 5, the same magnifyd.

Stramonium. 1, the Seed entire. 2, the inner thick cover. 3, the same split in two. 4, the true seed taken out of it. 5, half the thick Cover (3) magnifyd. 6, the Seed (4) magnifyd.

T A B. LXXVIII. Some examples of the Buds of Seeds before they are sown.

Sena. 1, the naked Seed. 2, the Lobes divided to shew the Bud. 3, one Lobe with the Bud magnifyd.

Carduus Benedictus, 1, the entire Seed. 2, with the outer Covers off. 3, naked. 4, divided. 5, that half with the Bud, magnifyd.

Hemp. 1, the naked Seed divided. 2, 3, the same magnifyd.

Almond. 1, one half of the Kernel. 2, the Radicle and Bud at the bottom of it. 3, the same broken off. 4, magnifyd. 5, opened.


T A B. LXXIX. F. 1. ab, Part of the outer Coat.

The Explication of the Tables.


cd, Part of the Inner Coat.
cde, one Lobe cover'd with the Skin.
fg, the other, with the skin and part of the Parenchyma pared off.
ff, the Skin.
hh, the Parenchyma.
ii, the Seminal Root.
kk, the Radicle.
kl, where it is cut off from the Lobes.
M, the Plume or Bud.
N, The Cavities in which it is lodg'd.
 F. 2. Sheweth the Barque, Vessels and Pith of the Radicle.
 T A B. LXXX. F. 1. A Slice of a young Apricock, cut transversly, near the lower end; shewing the duplicature of the Skin half way through the Stone.
 F. 2. A Slice, cut near the upper end; shewing the duplicature of the Skin quite thro' the Stone.
 F. 3. A well-grown Apricock cut by the length.
 F. 4, 5. The Membranes of a Filbert full ripe.
 F. 6. The Membranes of a young Apricock, with part of the Seed-branch.
 F. 7. the two Membranes cut by the length.
 T A B. LXXXI. F. 1. The outer and middle Coats or Membranes; with the Chanel, oval at both ends, now formed in the latter.
 f. 2, Part of the same, with the upper Oval grown larger, and the inmost Cover now also formed therein.
 f. 3. the same with the inmost cover grown larger.
 f. 4. the Inmost Cover more magnified, and the hollow in the smaller end, laid open, to shew the Seed it

self, newly begun in a round Node.
 f. 5. the same; in which the Node begins to be divided into two Lobes.
 f. 6, 7, 8, the gradual forming of the Lobes.
 f. 9, next the forming of the Radicle.
 f. 10, 11, Its gradual contraction at the point, into a short and slender Navle string.
 Which in the further growth of the Seed, breaks and disappears.
 T A B. LXXXII. *aa*, the Pulp, or open Parenchyma.
bb, the close Parenchyma or ground of the Stone.
cc, the Flower-Branch running through the body of the Stone.
dd, the Seed-Branch striking into the hollow of the Stone, and so running round the outer Membrane *ee*.
ff, the middle Membrane.
gg, the Chanel.
h, the inner Membrane, in which lies the Seed.
 T A B. LXXXIII. f. 1. the manner of the generation of the Essential Salts of Plants.
 f. 2, a Crystal of the Essential Salt of Rosemary, a little magnified.
 f. 3, *a*, *b*, two of Wormwood, *a*, upon the second Solution; *b* upon the first.
 f. 4, one of G. Scurvy-grass; *a*, one side; *b* the other.
 f. 5. a Crystal of the Marine Salt of Rosemary.
 f. 6, of Garden Scurvy-grass; *a* the upper side; *b* the nether.
 f. 7, of Wormwood.
 f. 8. of Black Thorne.
 f. 9. another of the same.
 f. 10, of Firne.
 f. 11, another of Wormwood.

F I N I S.



Cress-Seed
f. 4.



Gar




f. 2.




Beane

fig. 1.




Almond
f. 6.





Barley

f. 5.



Slice of a Beane

f. 9.



Foenugreek Seed

f. 7.




French Bean

f. 8.



Gourd

f. 15.



Seminal Root

f. 14.




Plume

f. 11.



Plume

f. 12.




Radicle

f. 10.



Lobe

f. 13.








fig. 1.
Lupine-Seed.



Tab. 2.
f. 2. Cucumber-S.




f. 3. Lettice-S.



f. 4.

Beane




f. 5.

Vine-Root





f. 6.

Plume-Root




f. 7.

Fenil-Root




f. 9.

Columbine-Root




f. 8. Turnep.







Tab. 3. fig. 1. a Beane



f. 2. Oak-Wood.




f. 3. Peice of Cane




f. 4. a Barley=Corne

f. 5. Plum-Tree Branch


f. 6. Bur-dock




f. 7. Piece of Oak magnifyd




f. 8. Piece of Beech







Tab. iv. Piece of Beech Wood
fig. 1.



Stalks of y^e Leaves of



Lily
f. 12




Peruvian Starwort


f. 13



f. 14 *Pear*



f. 15 *Aprcock*



f. 16
y^e *Secundine*




Lupine

f. 18



f. 17 *Coats of lupine*







fig. 1.
Primrose Root



Tab: 5.
f. 2.
Wood-sorrel R.



f. 3.
Devils-bitt R.



f. 4.

Tuberous Iris R.




f. 5.

Dandelion R.




f. 6.
Dragon R.



f. 7.


Spring-Crocus R.






Tab: VI.

The Root of Scorzonera.



Ierusalem Artichoake.



Burnet.

Dandelion.

Borage.

Carrot.

Parsley.



1843

1	2	3	4	5
6	7	8	9	10
11	12	13	14	15
16	17	18	19	20
21	22	23	24	25
26	27	28	29	30
31	32	33	34	35
36	37	38	39	40
41	42	43	44	45
46	47	48	49	50
51	52	53	54	55
56	57	58	59	60
61	62	63	64	65
66	67	68	69	70
71	72	73	74	75
76	77	78	79	80
81	82	83	84	85
86	87	88	89	90
91	92	93	94	95
96	97	98	99	100

1843

1	2	3	4	5
6	7	8	9	10
11	12	13	14	15
16	17	18	19	20
21	22	23	24	25
26	27	28	29	30
31	32	33	34	35
36	37	38	39	40
41	42	43	44	45
46	47	48	49	50
51	52	53	54	55
56	57	58	59	60
61	62	63	64	65
66	67	68	69	70
71	72	73	74	75
76	77	78	79	80
81	82	83	84	85
86	87	88	89	90
91	92	93	94	95
96	97	98	99	100


1843

1	2	3	4	5
6	7	8	9	10
11	12	13	14	15
16	17	18	19	20
21	22	23	24	25
26	27	28	29	30
31	32	33	34	35
36	37	38	39	40
41	42	43	44	45
46	47	48	49	50
51	52	53	54	55
56	57	58	59	60
61	62	63	64	65
66	67	68	69	70
71	72	73	74	75
76	77	78	79	80
81	82	83	84	85
86	87	88	89	90
91	92	93	94	95
96	97	98	99	100


1843

Tab: VII. Roots w:^{ch} Bleed little or nothing.


Marsh Mallow.




Patience.




Iris.



Peony.




Bistord.




Roots which Bleed a Lympha.


Bugloss.




Bryony.




Borage.




Black-Henbane.




Horse-Radish.



Deadly-Nightshade.



Brownwort.



Non-Bulbous Lily.



Asparagus.



Columbine.



1800

The first part of the book is devoted to a description of the various species of plants which are found in the country. The author has been very particular in his observations, and has given a full and accurate account of each. He has also been very diligent in his collection, and has secured a large number of specimens, which he has preserved in the most perfect manner.

The second part of the book is a description of the various species of animals which are found in the country. The author has been very particular in his observations, and has given a full and accurate account of each. He has also been very diligent in his collection, and has secured a large number of specimens, which he has preserved in the most perfect manner.

The third part of the book is a description of the various species of minerals which are found in the country. The author has been very particular in his observations, and has given a full and accurate account of each. He has also been very diligent in his collection, and has secured a large number of specimens, which he has preserved in the most perfect manner.

The fourth part of the book is a description of the various species of fossils which are found in the country. The author has been very particular in his observations, and has given a full and accurate account of each. He has also been very diligent in his collection, and has secured a large number of specimens, which he has preserved in the most perfect manner.

The fifth part of the book is a description of the various species of rocks which are found in the country. The author has been very particular in his observations, and has given a full and accurate account of each. He has also been very diligent in his collection, and has secured a large number of specimens, which he has preserved in the most perfect manner.

The sixth part of the book is a description of the various species of metals which are found in the country. The author has been very particular in his observations, and has given a full and accurate account of each. He has also been very diligent in his collection, and has secured a large number of specimens, which he has preserved in the most perfect manner.

The seventh part of the book is a description of the various species of stones which are found in the country. The author has been very particular in his observations, and has given a full and accurate account of each. He has also been very diligent in his collection, and has secured a large number of specimens, which he has preserved in the most perfect manner.


The eighth part of the book is a description of the various species of shells which are found in the country. The author has been very particular in his observations, and has given a full and accurate account of each. He has also been very diligent in his collection, and has secured a large number of specimens, which he has preserved in the most perfect manner.

The ninth part of the book is a description of the various species of bones which are found in the country. The author has been very particular in his observations, and has given a full and accurate account of each. He has also been very diligent in his collection, and has secured a large number of specimens, which he has preserved in the most perfect manner.


The tenth part of the book is a description of the various species of teeth which are found in the country. The author has been very particular in his observations, and has given a full and accurate account of each. He has also been very diligent in his collection, and has secured a large number of specimens, which he has preserved in the most perfect manner.

Roots which Bleed a Lympha.


f. 1. Parsnep



f. 2. Carrot




f. 3. Beet



f. 5. Nettle



f. 4. Ierusalem Artichoak



f. 7. Dropwort



f. 6. Valerian




f. 8. Lychnis




Roots which Bleed a Milk or Oily Sap

f. 10. Butyr Bur



Lovage f. 9.




f. 11. Dandelion



f. 12. Great Celandine



B



f. 13. Cychory




f. 15. Bishopsweed



f. 14. Goatsbeard








Tab: 9.

Roots with Milky or Balsamick Vessels, and Lymphæducts, both apparent.

F: 1. Fenil. F: 2. Emula camp:



F: 3. Tracheliu.




F: 4. Wormwood.




Roots with two sorts of Lymphaticks; in some, Aquæducts and Muciducts.


F: 5. Card: Scurvygrafs.




F: 6. Wild Cucumer.



F: 6. B.



F: 7. Melilote.



F: 8. Potato.




F: 9. Carduus beued.




F: 10. Eryngo.




F: 11. Cumbrey.




F: 12. Monks Hood.




F: 13. Cinquefoyle.



F: 13. Valerian.



F: 15. Burnet.



Tab. x.

Fig: 2. The same magnifyd






Fig: 1
Small Root of
Asparagus





Root of Ierusalem
 Artichoak cut trans-
 -versly fig. 1.



Tab. XI.


Half a Slice Magnifyd

Half a Slice of
 y^e same f. 2.

Fig. 3.




A




- 10
- 20
- 30
- 40
- 50
- 60
- 70
- 80
- 90
- 100
- 110
- 120
- 130
- 140
- 150
- 160
- 170
- 180
- 190
- 200

Standard
 of Bladders
 in Roots

71




The same
Magnified
Fig. 2.





Slice of a small
M. Mallow Root
Fig. 1.







Slice of Dandelion
Root f. i.








magnify'd












Fig 1
The small end of
Hors-Radish Root



The same magnifyd



49





Tab. XVI.

f. 2. The same Magnified.

Wormwood Root cut
transversely.

f. 1.





1797

1797

1797

1797

1797

1797

1797

1797

1797

1797

1797

1797

1797

1797

1797

1797

1797



1797


The same Magnified.


Fig: 2

Wine - Root
cut transversty

Fig. 1.









Tab 18 Stalks & Branches cut transuersly

Indian Wheat


f.1 Dandelyon



f.2





Borage f.3




Colewort f.5

f.4 Holyoak




f.6 Wild Cucumer




Burdock f.7



Scorzonera f.8




Endiue f.9




Vine f.10.A







B



f.11. Sumach



From the



Tab 19.


F.3. vine.





F.2. Sonchus.




F.1. Corin





F.4. Walnut ..




with y Barque off.






A. Magnifyd.




A


the Surface of Walking-Cane. Fig. 1.



the Skin of Borago-Stalk. F. 2.




B. Magnifyd.



B


Turpentine-Vessels in the Barque of Pine. F. 3.



b


a

Milk-Vessels in the Barque of Sumach. F. 4.



b


a




Tab. 21.

AB Piece Cut out of (ab) and Magnifyd to Shew y^e Lymphæducts & Aer-vessels.

a.b. Part of a Vine Branch Cut transversely







Tab. 22

Holly Branch
cut transversely

f. 2.
The same

f. 1.








Tab. 23

Hazel. Branch
cut transversly

f-2.
The same

f-v








Tab. 24

Barberry Branch
cut transversly

f 2
The Same

f. 1







Tab. 25


Apple Branch
cut transversly

f. 2.
The same

f. 1.









Tab. 27

Plum Branch
cut transversly

f. 2.
The same








Tab. 28

Elm Branch
cut transversly

f. 2.
The same

f. 1.






Tab- 29

Ash Branch
cut transversly

f. 2.
The same

f. 1.







Tab. 30

Walnut Branch
cut transversly

f. 2.
The same

f. 1.








Tab. 31

Figg Branch
cut transversly

f. 2.
The same







Tab. 32

Pine Branch cut transversly

f. 2.
The same

f. 1.







Tab. 33

Oak Branch cut
transversly

f. 2.
The same

f. 1.







Tab. 34

Sumach Branch
cut transversly

f. 2.
The same

f. 1.








Tab. 35

Worm wood stalk
cut transversly


f. 2.
The same


f. 1.







*Part of a Vine Branch cut transversely, and
split half way down y^e middle.*





Tab. 37. Part of a Corin Branch Cut as in Tab. 36.







Tab. 38

*Thistle stalk
cut transversly*


*f. 2.
The same*

f. 1.






The small particle of Fir-wood (a)
magnifyd.
f. 3.




One of y^e Lignus Fibers or
Lympheducts (a) in y^e Thred (bc)
f. 2.




a linen Thred
f. 1.



f. 5.




f. 4.




0

Tab: XL


Part of Tab. 34. represented somewhat Larger &
with several Breaks to shew y^e. Contexture both of y^e.
Perpendicular & Horizontal Fibers





TAB. XXXXI


Leaf of Dock.




Wild-Clary.



Branch of Sumach.















fig: 1. Magnifyd.



Leaf of Goosgrass.




Of Bears Ear.




Hairs on y Leaf of Tree Sage.



fig: 2. Magnifyd.



Of Herusalem Cowslip.




Hairs on y Leaf of Mullen. fig: 1.




fig: 2.






Tab XLIII.


Leaf of Venetian Vetch




Orange




Sage-Leav'd Ironwort




Cornelian Cherry




Broad Leav'd Laser-wort






Tab. XLV.

Leopards Bane




Great Laserwort




Mountain Calamint




Black Poplar



Shrubby Marsh-Mallow






Tab. 46.

Clematis Syl. m.




Leaves of

Mallows.




Vine.





TAB XLVII.


Leaf of *Syringa*.



Hemlock.




Holyoak.




Strawberry.







Leaf of Firr.




Rhamnus Salicis Folio.




The top of Pine Leaf.




The top of Lilly leaf.






Mallow.




Dock.




Dandelion.



Wild-Clary.




Borage.




Mullen.






tab. 50.



A young Borage Leaf



Tab: 51.

The Aer-Vessels unroaved in a Vine Leaf.

Fig: 3.




Fig: 1.






Fig: 2.





The Aer-Vessels unravell'd in a Scabions Leaf.

Tab. LII







Fig. 3.


4





Tab. LIII

Marine Salt










Nitrous




Alkaline





Acid





Blattana



Flowers of Starwort



TAB. LIV



Convolvulus Mallow





Poppey



Ladys Looking Glass



Ladys Bower



Marvel of Peru




Cichory



Tab. 55.


f. 1.

Flower of *Dulcamara* magnified.



f. 2.

Flower of *Colus Jovis* magnified.



f. 3.

Flower of *St James's wort*.



1 2 3 4 5 1 2 3 4 5 1 2 3 4 5

f. 7.

Theca (a) magnified.



f. 4.


Fl. of Chamemile.



1 2 3 4 5 1 2 3 4 5 1 2 3 4 5

f. 10.

The Theca (c) magnified.



f. 5.

Fl. of *Clematis Austriaca*.



f. 8.

Fl. of *Blattaria*.



f. 9.
One of y^e
Thecae.




f. 6.

One of y^e Thecae
in y^e flower.


a

Tab. 56.


f. 3.
The Backside of y^e Theca. (a)



f. 7.
The Column (c) Magnifid.




f. 4.
The Belly of the Theca. (a)



f. 1.
Flower of
Hyoscyamus.





f. 2.
One of y^e Spermatick
Theca.




f. 6.
The Column in y^e
Middle of y^e flower.



f. 5.
The Edges of the
Theca (a) open.






f. 1.
Flower of
St. Johns wort.




f. 2.
The same a little magnified.


Tab. 57.



f. 3.




f. 4.






Tab. 58.
Snapdragon
f. 1.




The Sperme of
Plantaine



Bearsfoot
f. 3.



Carnation
f. 4.




Derils - bit
f. 6.




Mallow
f. 14.
The Spermatick Glo-
bulets in f. 13.



Bindweed
f. 5.



f. 13.
One of y^e Thecæ (t)
in f. 12.




f. 12.
The Attire (e)
in f. 11.




f. 11.
The Flower
of Mallow



Beane
f. 7.




Lily
f. 8.



Pancy
f. 10.



Deadly Night Shade




Tab. 59.

f. 1.
A Sprig of Golden-Rod flowers.




f. 3.
The flower (a) magnified.




f. 2.
One flower.



f. 4. & 5.
One suit of y flower (a)



f. 6.
The Bladth. in y^c
sheath.(c).



Tab. 60.

f. 1.
A French Marigold.




f. 2.
One suit of which
about 12 are within
if circle. (a, b.)




f. 5.
One suit of Chrysanthemum Cret.
of iv. about 20. are in every flower.




f. 3.
The suit (c) magnified.



f. 4
The Blade (c, d) taken
out of y^e Sheath. (c).






Tab. 61.




f. 8.
The suit (d) Magnifid.

f. 2.
One of y^e suits, of
w^hich are about 40. in y^e circle.
(a b)



f. 7.
One suit of Knapweed.



f. 5.
Is (b) Magnifid.



f. 3.
The suit (c)
Magnifid.




f. 6.
The flower of Knapweed.




f. 4.
A Marigold leaf.





f. 1. Cichory flower.




Tab. 62.

f. 2. One Leaf.



f. 3. The Leaf (a) with its Attire, Magnified.




f. 5. The Blade. (c.d)




f. 4. The Sheath. (b)






Tab. 6.3.

f. 2.
One of y^e Flowering
Bud (a) magnified.



f. 4.
Sheweth a young Tulip
(a) as it is formed in y^e bottome
of y^e Root in September.




f. 3.
The same flower cut open,
shewning y^e spermatick Theca,
& y^e Uterus.





Tab. 04.


f. 6.
Is (c)ory Column Naked.



f. 2.
The Same: Opend.




f. 5.
One of y Theca (a. b.)
Surrounding. (c.)




f. 1.
A flower Bud of
A Sarum.



f. 3.
Is (a) Magnifid.




f. 4.
Is (a) with y Leaves Strip'd of.




Tab 65


f.2
Cut transversely



f.3
The piece (ab)
magnified




f.1
an Apple cut by the
length




Tab. 66.


f. 3 great
One of the Baggs, a.c.e.




f. 4
One of the little Baggs
c.e. cut transversly




f. 1.
a Limon cut downe




f. 2.
cut transversly



f. 5
Garden Cucumer




f. 6
Wile Cucumer



Pear-cut transversly:

fig. 1




Cut by y length


fig. 2



f. 4.
The peice (a,b) magnify'd.




f. 5
Quince



f. 3.
A Peice cut out of
f. 1.




f. 1.
a plum cut
transversly



Tab. 68.
(See Tab. 80. &c)


f. 3.
a piece taken
out of f. 2.




f. 4.
the piece a,
Magnifyd




f. 2.
an Aprcock cut
transversly




f. 5.
A young one wth eⁿow
bulky Coats of y^e Seed



f. 6.
the same by
y^e Length




f. 7.
One with the
kernell full grown



Tab. 69.

f. 3.
a Goosberry cut downe.




f. 2.
a Grape.



f. 4.
a Goosberry cut transversely.




f. 5
Is (f. 4.) magnified.




f. 1.
a Cherry with kind




f. 6.
a Filbert



Garden Radish.




Seed-Case of Red Poppy.





Tab. 70.



of Garden Bean.




Yellow Henbane.







Tulip.




Seed Case of.



Tab. LXXI.



Stramonium




Anagallis



Codeo Arsmart




Azarum.




Faint, illegible text, possibly bleed-through from the reverse side of the page.

Tab LXXII


Plum-Stones



French-Bean



Chesnut



Stone of Bellir: Myrobal



P. Christi S




Orange S.



The Small piece (a c) taken out of the Leaf, & magnified


Leaf of Harts tongue



The Seed

a c

The Seed Case open




Seed of Typha Maior




Seed of Pine



Wood Sorrel Seed




S. of Goos Grass




Seeds of.


Poppy.



Little Century.




Spergula.




Little Celandine.




Lychnis.




Great Celandine.



Chickweed.



Ben.



Pentaphyll. fragif.




Doves-foot.



Brooklime.



Rush.



Little Bell.




Sed. maj.




Ragwort.



Barley Grass.




Vervaine.




S^t. Iohns wort.






Nettle



Seeds of

Eyebright




Wormwood




Tab LXXIII


yarrow




Sorrel.



Anagallis




Nigella.




a sort of Bugloss



Melissa Moldavica.




Bellis Tanaceti folio annua.




Stachas Arabica.




Wart Wort.




Chrysanth. Americ.



Blattaria.



Foxe Glove.




Tansey.




Date.

Seeds of


Tab: 75.




Hounds tongue. Cucumer




Scorzonera.





Viola lunaris.




Woad.




Great blew Lupine.




Rhapontick




Garden Radish



Holyoak




Cotton Plant




Tab: 76

Seeds of




Purging Angola-Nut




Purging Barbado-Nut




Ricinus Americanus



Nux Vomica officin:





Tab. 77. Coffee.




Seeds of




Goosgrass






Staphusaria



Peony.




Stramonium.



Tab. 78.


Seeds of

Carduus bened



Sena


3



5



Hemp




3




2



Almond




5



4




2




Tab. 80 Sheweth y^e Structure of y^e Stone & y^e
two uper Membranes of y^e Seed

f. 1.




f. 2.



f. 4.




f. 5.




f. 3.




f. 6.






f. 7.





Tab. 81.
Sheweth y^e fur-
ther process
in the Gene-
ration of
y^e Seed





Tab 82.

The young Fruit, Three Membranes,
& Seed now loose,

