

Historic, archived document

Do not assume content reflects current scientific knowledge, policies, or practices.

On January 24, 1978, four USDA agencies—Agricultural Research Service (ARS), Cooperative State Research Service (CSRS), Extension Service (ES), and the National Agricultural Library (NAL)—merged to become a new organization, the Science and Education Administration (SEA), U.S. Department of Agriculture.

This publication was prepared by the Science and Education Administration's Federal Research staff, which was formerly the Agricultural Research Service.



Tachinid (*Winthemia?*) ovipositing on larva of whitelined sphinx moth, *Hyles lineata* (Fabricius). Courtesy of Robert W. Mitchell, Texas Tech University.

A Host-Parasite Catalog of North American Tachinidae (Diptera)

By Paul H. Arnaud, Jr.

Miscellaneous Publication No. 1319

Science and Education Administration
UNITED STATES DEPARTMENT OF AGRICULTURE

Washington, D.C.

Issued June 1978

ACKNOWLEDGMENTS

Thanks are extended to Dr. George E. Lindsay, Director of the California Academy of Sciences, for his administrative support of this project. Dr. Curtis W. Sabrosky, Systematic Entomology Laboratory, Science and Education Administration, USDA, was technical representative and I would like to acknowledge his continuous aid and advice on editorial matters and for taxonomic advice on the Tachinidae. Thanks are due to the entomologists of the Systematic Entomology Laboratory and the Smithsonian Institution at the National Museum of Natural History for their checking several thousand host names.

This catalog is dedicated to those persons who have published on the biology of the Tachinidae and to those who have contributed their time and skill to bring it to completion. The preparation was made possible by a United States Department of Agriculture contract with the California Academy of Sciences (RMA Contract No. 12-14-100-8315 (33)).

CONTENTS

	Page
Synopsis of parasite-host relationships	3
Parasites	47
Synopsis of host-parasite relationships	559
Hosts	589
Bibliography	687

USDA policy does not permit discrimination because of race, color, national origin, sex, or religion. Any person who believes he or she has been discriminated against in any USDA-related activity should write immediately to the Secretary of Agriculture, Washington, D.C. 20250.

A Host-Parasite Catalog of North American Tachinidae (Diptera)

By Paul H. Arnaud, Jr., *curator and chairman, Department of Entomology, California Academy of Sciences, and associate, California Agricultural Experiment Station, Science and Education Administration*

The Tachinidae are the most important of the dipterous families with entomophagous habits. Though often little noticed, these flies have an important role in controlling phytophagous Insecta.

In 1897 Coquillett published a list of tachinid-host relationships, in which he enumerated 84 species of Tachinidae from 183 host species.

The purpose of this catalog is to bring together the published host associations of the North American Tachinidae. Unpublished rearings are not included. Over 2,200 papers were reviewed. This catalog lists 519 tachinid species (with an additional 226 of incomplete or uncertain determination) in 261 genera from 1,049 different host species (with an additional 355 of incomplete or uncertain determination). Except for secondary references that furnish new information, or that might be more readily available than the original, only primary references have been included.

Thaddeus William Harris, M.D., in 1841 published the first North American host record for the family Tachinidae. The literature reported here is from 1841 through 1969.

The region covered is North America from Panama (Canal Zone) north, the West Indies (Greater and Lesser Antilles, excluding Trinidad and Tobago), the Hawaiian Islands, and Greenland. The Hawaiian Islands are included because their tachinid fauna consists primarily of introduced North American species.

This catalog is divided into five parts. Bibliographic information is contained only in the section on parasites. The other sections must be used in conjunction with it. Although no general index is provided, parasite and host names are indexed in their respective sections.

Many publications were used in compiling this catalog. Over 370 serial publications, including the following entomological journals, were examined:

Acadian Entomological Society Proceedings
American Entomological Society Transactions
The American Entomologist

Brooklyn Entomological Society Bulletin
The Canadian Entomologist
The Canadian Insect Pest Review
Cooperative Economic Insect Report
Entomologica Americana
Entomological News
Entomological Society of America Annals
Entomological Society of British Columbia Journal
Entomological Society of Canada Memoirs
Entomological Society of Nova Scotia Proceedings
Entomological Society of Ontario Proceedings
Entomological Society of Philadelphia Proceedings
Entomological Society of Washington Proceedings
The Florida Entomologist (The Florida Buggist)
Georgia Entomological Society Journal
Hawaiian Entomological Society Proceedings
Insect Life
The Insect Pest Survey Bulletin
Insecutor Inscitiae Menstruus
Journal of Economic Entomology
Kansas Entomological Society Journal
Lepidopterists' Society Journal
New York Entomological Society Journal
The Pan-Pacific Entomologist
The Practical Entomologist
Psyche, A Journal of Entomology
U.S. Department of Agriculture, Division of Entomology,
Bulletin
U.S. Department of Agriculture, Division of Entomology, Technical Series
University of California Publications in Entomology

The Entomological Society of America's "Index to the Literature of American Economic Entomology," volumes 1 through 18 (published 1917-59) provided hundreds of references.

Major articles cataloged include those published by Schaffner and Griswold (1934) on parasites of Macrolepidoptera from the Northeastern United States, Raizenne (1952) on forest Lepidoptera of southern Ontario, and Schaffner (1959) on parasites of Microlepidoptera from the Northeastern United States.

Important catalogs or lists not included in this publication and to which the reader is referred are a world catalog of parasites and predators of insect pests by Thompson (1943-58), a world list of Tachinidae and their hosts by Townsend (1942, pp. 211-246), and natural enemies of Canadian agricultural pests by Graham (1965).

SYNOPSIS OF PARASITE-HOST RELATIONSHIPS

This section is based entirely on information contained in the next section on parasites. The Tachinidae follow the classification proposed in "A Catalog of the Diptera of America North of Mexico" (see Sabrosky and Arnaud, 1965). Author names and specific names of hosts are omitted.

Subfamily RHINOPHORINAE

Tribe RHINOPHORINI

Melanophora

roralis: Oniscidae (*Oniscus*, *Porcellio*).

Subfamily PHASIINAE

Tribe GYMNOSOMATINI

Gymnoclytia

dubia: Pentatomidae (*Cosmopepla*).

immaculata: Pentatomidae (*Euschistus*); Noctuidae (*Pseudaletia*).

occidua: Pentatomidae (*Euschistus*); Pieridae (*Pieris*).

Gymnosoma

filiola: Pentatomidae (*Chlorochroa*).

fuliginosum: Pentatomidae (*Chlorochroa*, *Eurygaster*, *Euschistus*,
Wheat bug(-s)).

Tribe TRICHOPODINI

Acaulona

erythropyga: Pyrrhocoridae (*Dysdercus*).

peruviana: Pyrrhocoridae (*Dysdercus*).

Trichopoda

lanipes: Coreidae (*Acanthocephala*, *Archimerus*); Pentatomidae (*Nezara*, *Podisus*).

pennipes: Coreidae (*Acanthocephala*, *Anasa*, *Archimerus*, *Chelinidea*, *Euthochtha*, *Leptoglossus*, *Narnia*); Largidae (*Largus*); Pentatomidae (*Acrosternum*, *Alcaeorrhynchus*, *Euschistus*, *Murgantia*, *Nezara*, *Plautia*, *Thyanta*); Scutelleridae (*Coleotichus*).

pennipes probably: Coreidae (*Anasa*); Mantidae (*Tenodera*).

- pennipes* complex: Coreidae (*Archimerus*, *Leptoglossus*).
pilipes: Pentatomidae (*Nezara*); Scutelleridae (*Coleotichus*).
plumipes: Coreidae (*Acanthocephala*, *Leptoglossus*); Acrididae (*Dissosteira*, Unidentified Acrididae).
plumipes probably: Acrididae (*Dissosteira*).
subdivisa: Acrididae (*Dissosteira*).
spp. (*pennipes* and/or *pilipes*); Pentatomidae (*Nezara*).
spp.: Coreidae (*Acanthocephala*, *Anasa*, *Leptoglossus*); Pentatomidae (*Chlorochroa*, *Nezara*, Plant bug(-s)); Acrididae (*Dissosteira*).

Tribe PHASIINI

Alophorella

- aeneoventris*: Miridae (*Lygus*); Pentatomidae (*Cosmopepla*); Cercopidae (*Philaenus*).
fumosa: Miridae (*Lygocoris*, *Lygus*).
opaca: Miridae (*Lygus*).
pulverea: Miridae (*Lygocoris*, *Lygus*); Pentatomidae (*Holcostethus*).
spp.: Miridae (*Lygus*).

Androeuryops

- ecitonis*: Formicidae (*Eciton*).

Hyalomya

- aldrichii*: Lygaeidae (*Emblethis*, *Geocoris*, *Nysius*); Unidentified Hemiptera (Heteroptera) (Miscellaneous).
purpurascens: Cydnidae (*Sehirus*).

Paraphorantha

- pollinosa*: Pentatomidae (*Chlorochroa*).

Phasia

- nigra*: Pentatomidae (*Eurygaster*).

"Phorantha"

- "*occidentis*": Lygaeidae (*Nysius*); Miridae (*Leptopterna*).

Tribe CATHAROSIINI

Catharosia

- lustrans*: Lygaeidae (*Eremocoris*, *Thylochromus*).

Tribe CYLINDROMYIINI

Beskia

- aelops*: Pentatomidae (Pentatomid (including *Euschistus*, *Mormidea*, *Nezara*, and/or *Oebalus*)); Aphididae (*Macrosiphum*); Noctuidae (*Alabama*).

aelops probably: Tettigoniidae (*Neoconocephalus*).

Cylindromyia

armata: Pentatomidae (*Chlorochroa*).

binotata: Pentatomidae (*Euschistus*); Saturniidae (*Actias*).

“*carolinae*”: Noctuidae (*Pseudaletia*); Acrididae (Grasshopper(-s)).

euchenor: Pentatomidae (*Chlorochroa*, Pentatomid (including *Euschistus*, *Mormidea*, *Nezara*, and/or *Oebalus*)); Noctuidae (*Pseudaletia*); Acrididae (Unidentified Acrididae).

fumipennis: Pentatomidae (*Euschistus*).

Hemyda

aurata: Pentatomidae (*Podisus*).

Tribe LEUCOSTOMATINI

Leucostoma

acirostre: Coreidae (*Harmostes*).

aterrimum: Rhopalidae (*Liorhyssus*).

gravipes: Lygaeidae (*Oncopeltus*).

simplex: Nabidae (*Nabis*); Rhopalidae (*Liorhyssus*); Acrididae (*Melanoplus*, Grasshopper(-s)).

spp.: Lygaeidae (*Lygaeus*).

Tribe STRONGYGASTRINI

Hyalomyodes

triangulifer: Alleculidae (*Chromatia*); Chrysomelidae (*Altica*, *Chrysomela*, *Oulema*); Coccinellidae (*Coccinella*, *Coleomegilla*, *Epilachna*); Curculionidae (*Hypera*, *Listroderes*, *Sitona*); Pterophoridae (*Platyptilia*).

Tribe EUTHERINI

Euthera

tentatrix: Pentatomidae (*Euschistus*, *Oebalus*, Pentatomid (including *Euschistus*, *Mormidea*, *Nezara*, and/or *Oebalus*)).

Tribe PALPOSTOMATINI

Hamaxia

incongrua: Scarabaeidae (*Popillia*, Unidentified Scarabaeidae).

Subfamily PROSENINAE

Tribe ORMIINI

Euphasiopteryx

brevicornis nuttingi: Tettigoniidae (*Neoconocephalus*); Unidentified Orthoptera (Miscellaneous).

ochracea: Gryllidae (*Gryllus*).

Ormia

punctata: Tabanidae (*Tabanus*).

spp. probably: Pyralidae (*Crambus*).

Tribe TRIXINI

Phasiops

flavus: Tabanidae (*Tabanus*); Pyralidae (*Crambus*).

flavus probably: Tabanidae (*Tabanus*).

Tribe AULACEPHALINI

Eutrixa

exilis: Scarabaeidae (*Euetheola*, *Ligyris*, *Phyllophaga*, June beetle(-s), May beetle(-s)).

Tribe DEXILLINI

Dexilla

ventralis: Scarabaeidae (*Maladera*, *Phyllophaga*, *Popillia*).

Eutrixoides

jonesii: Scarabaeidae (*Chemarachis*, *Phyllophaga*, Scarabaeid (including *Phyllophaga* or *Phytalis*)).

Megapariopsis

opaca: Scarabaeidae (*Euetheola*).

Microphthalma

disjuncta: Scarabaeidae (*Euetheola*, *Phyllophaga*, June beetle(-s)), Scarabaeid (including *Cyclocephala* or *Phyllophaga*), Scarabaeid (including *Cyclocephala* and/or *Phyllophaga*), Scarabaeid (allied genera of *Phyllophaga*), White grub(-s)).

michiganensis: Scarabaeidae (*Phyllophaga*, White grub(-s)).

obsoleta: Scarabaeidae (*Phyllophaga*).

spp.: Scarabaeidae (White grub(-s)).

Tribe MYIOPHASIINI

Myiophasia

"aenea": Curculionidae (*Ampelogypter*, *Anthonomus*, *Chalcodermus*, *Conotrachelus*, *Curculio*, *Sphenophorus*, *Trichobaris*, Curculionid (including *Conotrachelus* or *Curculio*)); Unidentified Coleoptera (Miscellaneous); Noctuidae (*Pseudaetic*).

"aenea" probably: Curculionidae (*Conotrachelus*).

"atra": Curculionidae (*Curculio*); Arctiidae (*Halisidota*); Olethreutidae (*Laspeyresia*); Pyralidae (*Acrobasis*).

clistoides: Curculionidae (*Chalcodermus*).

globosa: Curculionidae (*Anthonomus*, *Chalcodermus*, *Conotrachelus*, *Trichobaris*, Unidentified Curculionidae).

metallica: Curculionidae (*Conotrachelus*, *Curculio*, *Sphenophorus*, Caneborer(-s)); Noctuidae (*Pseudaetia*).

nigrifrons: Curculionidae (*Conotrachelus*, *Curculio*).

robusta: Curculionidae (*Sphenophorus*).

Tribe PROSENIINI

Ateloglossa

cinerea: Elateridae (*Melanotus*).

Dinera

grisescens probably: Scarabaeidae (*Phyllophaga*).

Mochlosoma

lacertosum: Scarabaeidae (Scarabaeid (including *Phyllophaga* or other Scarabaeidae)).

mexicanum: Unidentified Lepidoptera (Miscellaneous).

"Phorostoma"

spp.: Tabanidae (*Tabanus*); Pyralidae (*Diatraea*, *Zeadiatraea*).

Prosenia

siberita: Scarabaeidae (*Popillia*).

spp.: Scarabaeidae (*Adoretus*, *Anomala*).

Ptilodexia

"abdominalis": Scarabaeidae (*Phyllophaga*).

canescens: Cerambycidae (*Saperda*, *Stenocorus*).

cerata: Scarabaeidae (*Phyllophaga*).

harpasa: Scarabaeidae (*Chemarachis*, *Phyllophaga*, May beetle(-s)).

"tibialis": Scarabaeidae (*Phyllophaga*).

sp.: Scarabaeidae (*Popillia*).

Tribe THERESIINI

Carinosillus

novaeangliae: Tabanidae (*Hybomitra*).

tabanivorus: Tabanidae (*Chrysops*, *Hybomitra*, *Tabanus*, Horse fly (flies)).

Eutheresia

monohammi: Cerambycidae (*Monochamus*); Lucanidae (*Ceruchus*).

monohammi probably: Cerambycidae (Unidentified Cerambycidae); Lucanidae (*Ceruchus*).

nipigonensis: Cerambycidae (*Stenocorus*).

satisfacta probably: Cerambycidae (*Stenocorus*).

trivittata: Cerambycidae (*Monochamus*, *Stenocorus*).

Opsotheresia

bigelowi: Tabanidae (Unidentified Tabanidae).

Paratheresia

claripalpis: Pyralidae (*Chilo*, *Diatraea*, *Ostrinia*, *Zeadiatraea*).

spp.: Pyralidae (*Diatraea*, *Zeadiatraea*).

Theresia

rutilans: Cerambycidae (*Enaphalodes*).

Gen. of Theresiini

spp.: Gryllidae (*Anurogryllus*).

Vibrissotheresia

pechumani: Tabanidae (*Tabanus*).

Subfamily TACHININAE

Tribe NEMORAEINI

"Nemoraea"

sp.: Arctiidae (*Turuptiana*).

Tribe MACROMYINI

Bombyliopsis

abrupta: Arctiidae (*Diacrisia*, *Euchaetias*, *Halisidota*, *Hyphantria*, Unidentified Arctiidae).

Tribe TACHININI

Nowickia

algens: Noctuidae (*Apamea*, *Euxoa*, Cutworm(-s)).

hispida: Noctuidae (*Polia*).

latifrons: Noctuidae (*Crymodes*).

Peleteria

aenea: Lymantriidae (*Byrdia*).

obsoleta: Noctuidae (Unidentified Noctuidae).

"*tessellata*": Noctuidae (*Euxoa*).

"*tessellata*" probably: Noctuidae (*Euxoa*).

"*texensis*": Lymantriidae (*Hemerocampa*); Noctuidae (*Agrotis*, *Euxoa*, *Heliophila*, *Leucania*, *Peridroma*, *Pseudaletia*, Armyworm(-s)).

thomsoni: Lepidoptera probably (Miscellaneous).

spp.: Noctuidae (*Agrotis*, *Pseudaletia*).

Tribe DEJEANIINI

Archytas

apicifer: Lasiocampidae (*Malacosoma*); Noctuidae (*Peridroma*, *Pseudaletia*, *Spodoptera*).

aterrimus: Arctiidae (*Cycnia*, *Hyphantria*); Lasiocampidae (*Malacosoma*); Megalopygidae (*Megalopyge*); Noctuidae (*Acronicta*, *Baileya*, *Catocala*, *Peridroma*, *Plathypena*, Unidentified Noctuidae); Notodontidae (*Cerura*); Psychidae (*Thyridopteryx*).

californiae: Arctiidae (*Euchaetias*); Noctuidae (*Spodoptera*, *Trichoplusia*).

cirphis: Noctuidae (*Agrotis*, *Cirphis*, *Leucania*, *Peridroma*, *Pseudaletia*, *Spodoptera*, Armyworm(-s) (including *Pseudaletia* and/or *Spodoptera*), Armyworm(-s), Cutworm(-s)).

cirphis probably: Noctuidae (*Peridroma*).

incertus: Noctuidae (*Spodoptera*).

lateralis: Lasiocampidae (*Malacosoma*).

marmoratus: Noctuidae (*Agrotis*, *Heliophila*, *Heliopsis*, *Laphygma*, *Leucania*, *Mocis*, *Pseudaletia*, *Spodoptera*, Armyworm(-s), Cutworm(-s)); Unidentified Lepidoptera (Corn stalk borer(-s), Miscellaneous).

metallicus: Arctiidae (*Ecpantheria*); Noctuidae (*Spodoptera*); Notodontidae (*Datana*).

sp. near *metallicus*: Noctuidae (Unidentified Catocalinae).

smaragdinus: Megalopygidae (*Megalopyge*).

spp. (*apicifer* and/or *californiae*): Lasiocampidae (*Malacosoma*, Tent caterpillar(-s)); Noctuidae (*Elaphria*, *Heliophila*, *Leucania*, *Peridroma*, *Pseudaletia*, *Spodoptera*, Armyworm(-s), Cutworm(-s)); Unidentified Lepidoptera (Miscellaneous).

spp. (*apicifer* or *californiae*) probably: Noctuidae (*Pseudaletia*).

spp.: Noctuidae (*Peridroma*, *Spodoptera*, Armyworm(-s)); Noto-
dontidae (*Datana*).

Juriniopsis

adusta: Arctiidae (*Ecpantheria*); Hesperidae (*Epargyreus*).

aurifrons: Arctiidae (*Halisidota*).

floridensis probably: Arctiidae (*Ecpantheria*).

Paradejeania

rutilioides nigrescens: Unidentified Lepidoptera (Miscellaneous).

Tribe JURINIINI

Epalpus

signifer: Noctuidae (*Lithophane*).

Xanthoepalpus

bicolor: Geometridae (Unidentified Geometridae).

Tribe CUPHOCERINI

Copecrypta

ruficauda: Noctuidae (*Ogdoconta*, *Plathypena*).

sp.: Aegeriidae (*Sylvora*).

Deopalpus

miscelli: Noctuidae (*Catabena*).

parksii: Unidentified Lepidoptera (Miscellaneous).

Tribe LINNAEMYINI

Bonnetia

comta: Scarabaeidae (*Phyllophaga*); Noctuidae (*Agrotis*, *Cop-
ablepharon*, *Euxoa*, *Feltia*, *Peridroma*, *Polia*, *Spodoptera*,
Armyworm(-s), Cutworm(-s), Unidentified Noctuidae).

sp.: Aegeriidae (*Sylvora*).

Chrysotachina

alcedo: Hesperidae (*Urbanus*); Lacosomidae (*Cicinnus*).

Gymnochaetopsis

fulvicauda: Noctuidae (*Heliothis*, *Mocis*).

Gymnocheta

ruficornis: Noctuidae (*Papaipema*).

Linnaemya

"*picta*": Noctuidae (*Agrotis*).

tessellata: Noctuidae (*Amathes*).

Thompsonomyia

anthracina: Arctiidae (*Parasemia*).

Tribe ERNESTIINI

Eucheirophaga

lugubris: Pieridae (*Eucheira*).

Mericia

ampelus: Arctiidae (*Diacrisia*, *Hyphantria*); Lasiocampidae (*Malacosoma*); Notodontidae (*Datana*); Unidentified Lepidoptera (Miscellaneous).

arcuata: Arctiidae (*Hyphantria*).

arcuata probably: Noctuidae (*Eupsilia*).

bicarina probably: Noctuidae (*Barathra*).

johnsoni: Arctiidae (*Hyphantria*).

spp.: Arctiidae (*Diacrisia*, *Estigmene*, *Euchaetias*, *Hyphantria*); Geometridae (*Ennomos*); Noctuidae (*Agrotis*, *Amphipyra*, *Cosmia*, *Euxoa*, *Lithophane*, Unidentified Noctuidae); Notodontidae (*Datana*); Olethreutidae (*Spilonota*).

spp. probably: Arctiidae (*Hyphantria*).

Xanthophyto

labis: Aegeriidae (*Vespa mima*); Lasiocampidae (*Malacosoma*); Olethreutidae (*Sciaphila*).

spp.: Geometridae (*Sabulodes*); Oecophoridae (*Depressaria*); Pyralidae (*Dioryctria*); Tortricidae (*Choristoneura*).

sp. probably: Geometridae (*Ennomos*).

Tribe APHRIINI

Aphria

ocypterata: Noctuidae (*Euxoa*); Pyralidae (*Nephopteryx*, *Tlascalala*).

Parafischeria

eucerata: Aegeriidae (*Synanthedon*).

Pyraustomyia

penitalis: Pyralidae (*Ostrinia*, *Pyrausta*).

Tribe LYPHINI

Lypha

intermedia: Gelechiidae (*Anacamptis*); Olethreutidae (*Sciaphila*).

setifacies: Tortricidae (*Choristoneura*, *Croesia*).

sp.: Tortricidae (*Sparganothis*).

Tribe GERMARIINI

Bigonicheta

spinipennis: Forficulidae (*Forficula*).

Coloradomyia

eucosmaphaga: Olethreutidae (*Eucosma*).

Exoristoides

harrisi: Geophilidae (*Geophilus* probably).

johnsoni: Gryllidae (*Gryllus*).

sp.: Gryllidae (*Anurogryllus*).

Helioplagia

slossonae: Coccinellidae (*Epilachna*); Hesperiididae (*Thymelicus*).

Petinarctia

stylata: Lymantriidae (*Byrdia*).

Tribe CAMPYLOCHETINI

Chaetophlepsis

eudryae: Aegeriidae (*Melittia*); Noctuidae (*Alypia*, *Autographa*, *Euthisanotia*, *Plathypena*, *Pseudoplusia*, Unidentified Noctuidae); Notodontidae (*Pheosia*).

orbitalis: Geometridae (*Cingilia*, *Protitame*); Noctuidae (*Cryphia*).

rindgei: Geometridae (*Drepanulatrix*).

semiothisae: Geometridae (*Itame*, *Lambdina*, *Nepytia*, *Pero*, *Semiothisa*); Tortricidae (*Acleris*).

spp.: Aegeriidae (*Melittia*); Noctuidae (*Catocala*, *Plathypena*); Tortricidae (*Choristoneura*).

Tribe VORIINI

Athrycia

cinerea: Unidentified Hymenoptera (Sawfly(-flies)); Noctuidae (*Pseudaletia*).

Cyrtophleba

coquilletti: Noctuidae (*Epiglaea*, *Eupsilia*, Unidentified Noctuidae).

Metaplagia

occidentalis: Noctuidae (*Heliothis*); Sphingidae (*Agrius*, *Sphinx*).

Metavoria

orientalis: Sphingidae (*Manduca*).

Plagiomima

cognata: Noctuidae (*Heliothis*).

spinosula probably: Noctuidae (*Heliothis*).

Voria

aurifrons: Noctuidae (*Autographa*, *Heliothis*).

ruralis: Noctuidae (*Anticarsia*, *Autographa*, *Autoplusia*, *Peridroma*, *Plusia*, *Spodoptera*, *Trichoplusia*, Phalaenid moth(-s), Unidentified Noctuidae); Noctuidae and/or Pyralidae (Noctuid and/or pyralid (including *Scotogramma* and/or *Loxostege*)); Unidentified Lepidoptera (Green looper(-s)).

ruralis probably: Noctuidae (*Autographa*, *Trichoplusia*).

Wagneria

major: Noctuidae (*Choephora*).

vernata: Noctuidae (*Protorthodes*, *Xylena*).

vernata probably: Noctuidae (*Orthosia*).

sp.: Noctuidae (*Choephora*).

Subfamily DEXIINAE**Tribe THELAIRINI****Spathidexia**

dunningii: HesperIIDae (*Ochlodes*).

Thelaira

americana: ArctIIDae (*Diacrisia*, *Estigmene*).

spp.: ArctIIDae (*Isia*, *Turuptiana*, Unidentified ArctIIDae); Pyralidae (*Diaphania*).

Tribe DEXIINI**Dexia**

limata: Tipulidae (Unidentified Tipulidae).

"Hypostena"

"*floridensis*": Argidae (*Sphacophilus*, *Sterictiphora*); Blastobasidae (*Valentinia*); Pyralidae (*Diatraea*, *Ostrinia*).

sp.: Chrysomelidae (*Erynephila*).

Tribe MACQUARTIINI**Athanatus**

californicus: Noctuidae (*Heliothis*).

Eubrachymera

debilis: Pamphiliidae (*Neurotoma*).

Eulasiona

comstocki: Oecophoridae (*Psilocorsis*); Olethreutidae (*Epinotia*, *Pseudexentera*).

genalis: Aegeriidae (Unidentified Aegeriidae).

nigra: Tortricidae (*Archips*).

Grisdalemyia

spp.: Chrysomelidae (*Chrysomela*).

Ictericophyto

spinosa: Geometridae (*Alsophila*, *Brephos*, *Erannis*); Noctuidae (*Pyreferra*); Tortricidae (*Pandemis*).

tibialis: Geometridae (*Melanolophia*).

"Macquartia"

spp.: Unidentified Coleoptera (Miscellaneous); Arctiidae (*Halisidota*).

Pelatachina

pellucida: Nymphalidae (*Nymphalis*, *Vanessa*).

Tribe HYALURGINA

Hyalurgus

sp.: Tenthredinidae (*Tomostethus*).

Tribe ICELIINI

Ervia

triquetra: Chrysomelidae (*Monocesta*).

Tribe ZELIINI

Metadexia

tricolor: Curculionidae (*Rhabdoscelus*).

Zelia

vertebrata: Cerambycidae (Unidentified Cerambycidae); Passalidae (*Popilius*); Scarabaeidae (*Osmoderma*); Tenebrionidae (*Meracantha*).

Tribe URAMYINI

Anaporis

limacodis: Limacodidae (*Limacodes*, *Lithacodes*, *Sisyrosea*, *Tortricidia*).

pristis: Arctiidae (*Halisidota*); Limacodidae (*Euclea*, *Phobetron*, *Sibine*, *Sisyrosea*); Megalopygidae (*Megalopyge*).

sp.: Limacodidae (*Sisyrosea*).

Uramya

halisidotae: Arctiidae (*Halisidota*); Lasiocampidae (*Malacosoma*).

Tribe CALODEXIINI

Calodexia

sp. near *agilis* or *interrupta*: Gryllidae (*Eneoptera*).

dives: Blattidae (Cockroach(-es)).

fumosa: Tettigoniidae (Unidentified Tettigoniidae).

interrupta: Gryllidae (*Ponca*).

venteris: Blattidae (*Eublaberus*).

spp.: Blattidae (*Epilampra*, Cockroach(-es)); Gryllidae (Crick-et(-s)).

Tribe URODEXIINI

Chaetona

sp.: Arctiidae (*Turuptiana*).

Chaetonodexodes

vanderwulpi: Chrysomelidae (*Monoxia*).

Cholomyia

inaequipes: Curculionidae (*Conotrachelus*).

sp.: Curculionidae (*Conotrachelus*).

Phyllophilopsis

evanida: Endomychidae (*Endomychus*).

Tribe ERIOTHRIXINI

Paradidyma

singularis: Unidentified Lepidoptera (Burrowing webworm(-s)).

Periscepsia

cinerosa: Noctuidae (*Euxoa*).

helymus: Noctuidae (*Actebia*, *Agrotis*, *Amathes*, *Autographa*, *Cirphis*, *Elaphria*, *Epiglaea*, *Eupsilia*, *Euxoa*, *Faronta*, *Leucania*, *Lithophane*, *Peridroma*, *Polia*, *Pseudaletia*, *Trichoplusia*, Armyworm(-s), Cutworm(-s), Unidentified Noctuidae).

laevigata: Noctuidae (*Elaphria*, *Euxoa*, *Lascoria*, *Lithophane*, *Peridroma*, *Pseudaletia*).

rohweri: Noctuidae (*Agrotis*, *Euxoa*, *Spaelotis*).

spp.: Noctuidae (*Agrotis* probably, *Glaea*); Unidentified Lepidoptera (Leaf roller(-s)).

Tribe LESKIINI

Dejeaniopalpus

texensis: Pyralidae (*Desmia*, *Pyrausta* probably); Tortricidae (Leaf roller(-s)).

Genea

aurea: Pyralidae (*Tetralopha*).

“Leskia”

“*analis*”: Pyralidae (*Diaphania*).

Leskiella

brevirostris: Olethreutidae (*Laspeyresia*, Unidentified Olethreutidae).

brevirostris probably: Olethreutidae (*Grapholitha*).

Leskiomima

tenera: Olethreutidae (*Eucosma*, *Grapholitha*, *Laspeyresia*); Pyralidae (*Acrobasis*, *Desmia*, *Dioryctria*, *Homoeosoma*, *Macrobotys*, *Tetralopha*, Unidentified Pyralidae).

sp.: Pyralidae (*Dioryctria*).

Myobiopsis

depilis: Pyralidae (*Macrobotys*).

flavipennis: Pyralidae (“*Blepharomastix*”); Unidentified Lepidoptera (Leaf-folder caterpillar(-s)).

Sipholeskia

occidentalis: Aegeriidae (*Synanthedon*).

Stomatodexia

cothurnata: Pyralidae (*Diaphania*).

Subfamily GONIINAE

Tribe ACEMYINI

Acemya

tibialis: Acrididae (*Melanoplus*, Grasshopper(-s)).

spp.: Acrididae (Grasshopper(-s)).

Ceracia

dentata: Acrididae (*Camnula*, *Chimarocephala*, *Chortophaga*, *Melanoplus*, Grasshopper(-s)).

dentata probably: Acrididae (*Chortophaga*, *Psoloessa*).

spp.: Acrididae (*Melanoplus*, Grasshopper(-s)).

Euhalidaya

genalis: Phasmatidae (*Diapheromera*).

Hemithrixion

oestriforme: Acrididae (*Boopedon*, *Camnula*, *Melanoplus*, Grasshopper(-s)).

plankii: Acrididae (Grasshopper(-s)).

sp.: Acrididae (Grasshopper(-s)).

sp. probably: Acrididae (*Dissosteira*).

Tribe BLONDELIINI

Adoryphorophaga

aberrans: Chrysomelidae (*Blepharida*, *Leptinotarsa*, Chrysomelid spp. (including *Lema* and/or *Leptinotarsa*)).

spp. and/or *Doryphorophaga* spp.: Chrysomelidae (*Leptinotarsa*).

"Anisia"

"nigrocincta" probably: Pyralidae (*Ostrinia*).

Aplomyiopsis

epilachnae: Coccinellidae (*Epilachna*).

epilachnae probably: Coccinellidae (*Epilachna*).

galerucellae: Chrysomelidae (*Pyrrhalta*).

Aporotachina

chaetoneura: Argidae (*Arge*); Tenthredinidae (*Nematus*).

dexina: Argidae (*Arge*).

Blondelia

eufitchiae: Arctiidae (*Halisidota*, *Hyphantria*); Geometridae (*Alsophila*, *Caripeta*, *Ennomos*, *Eupithecia*, *Itame*, *Lambdina*, *Nepytia*, *Semiothisa*, *Thera*); Noctuidae (*Hypena*); Olethreutidae (*Spilonota*); Tortricidae (*Acleris*, *Sparganothis*, Unidentified Tortricidae).

hyphantriae: Arctiidae (*Diacrisia*, *Halisidota*, *Hyphantria*); Noctuidae (*Bomolocha*, *Ceramica*, *Lithophane*); Notodontidae (*Heterocampa*).

polita: Pyralidae (*Loxostege*).

sp. near *polita*: Tortricidae (*Archips*).

spp.: Geometridae (*Anacamptodes*, *Coryphista*); Tortricidae (*Archips*).

spp. probably: Diprionidae (*Diprion*); Geometridae (*Cingilia*).

Celatoria

diabroticae: Chrysomelidae (*Acalymma*, *Cerotoma*, *Diabrotica*, *Trirhabda*).

setosa: Chrysomelidae (*Acalymma*, *Diabrotica*, Cucumber beetle(-s)).

Compsilura

concinata: Curculionidae (*Pissodes*); Cimbicidae (*Cimbex*); Diprionidae (*Diprion*); Pamphiliidae (*Neurotoma*); Tenthredinidae (*Croesus*, *Nematus*, *Pristiphora*, *Trichiocampus*); Unidentified Hymenoptera (Sawfly(-flies), Tenthredinoidea); Arctiidae (*Diacrisia*, *Estigmene*, *Euchaetias*, *Halisidota*, *Hyphantria*, Unidentified Arctiidae); Bombycidae (*Bombyx*); Citheroniidae (*Anisota*); Ctenuchidae (*Ctenucha*); Danaidae (*Danaus*); Geometridae (*Biston*, *Cingilia*, *Ennomos*, *Eugon-*

obapta, *Hydria*, *Itame*, *Lambdina*, *Melanolophia*, *Nemato-campa*, *Phigalia*, Unidentified Geometridae); HesperIIDae (*Epargyreus*, *Erynnis*, *Thymelicus*, Unidentified HesperIIDae); Lasiocampidae (*Epicnaptera*, *Malacosoma*, American tent caterpillar(-s)); Limacodidae (*Cnidocampa*); Lymantriidae (*Dasychira*, *Euproctis*, *Hemerocampa*, *Nygmia*, *Orgyia*, *Porthetria*, *Stilpnotia*, Tussock moth(-s), Unidentified Lymantriidae); Megalopygidae (*Megalopyge*); Noctuidae (*Acronicta*, *Alypia*, *Amphipyra*, *Calpe*, *Catocala*, *Ceramica*, *Euthisanotia*, *Lithophane*, *Phosphila*, *Plathypena*, *Plusiodonta*, *Polia*, *Pseudaletia*, *Rhodophora*, *Scoliopteryx*, *Simyra*, *Trichoplusia*, *Xylena*, Unidentified Catocalinae, Unidentified Noctuidae); Notodontidae (*Cerura*, *Datana*, *Fentonia*, *Heterocampa*, *Ichthyura*, *Nadata*, *Pheosia*, *Schizura*, *Symmerista*, Unidentified Notodontidae); Nymphalidae (*Agraulis*, *Chlosyne*, *Euphydryas*, *Limenitis*, *Nymphalis*, *Polygonia*, *Vanessa*, Unidentified Nymphalidae); Oecophoridae (*Dasystema*); Papilionidae (*Battus*, *Papilio*, Unidentified Papilionidae); Pieridae (*Pieris*); Pterophoridae (*Oidaematophorus*); Pyralidae (*Evergestis*, *Ostrinia*); Saturniidae (*Antheraea*, *Automeris*, *Callosamia*, *Hemileuca*, *Platysamia*, Unidentified Saturniidae); Sphingidae (*Ampeloeca*, *Amphion*, *Atreides*, *Ceratomia*, *Deidamia*, *Hemaris*, *Hyles*, *Manduca*, *Paonias*, *Pholus*, *Smerinthus*, *Sphecodina*, *Sphinx*, Unidentified Sphingidae); Tortricidae (*Archips*, *Choristoneura*); Unidentified Lepidoptera (Cabbage worm(-s), Miscellaneous).

sp. probably: Lasiocampidae (*Malacosoma*).

Cryptomeigenia

aurifacies: Scarabaeidae (*Chemarachis*, *Lachnosterna*, *Phyllophaga*, Scarabaeid (including *Phyllophaga* or *Phytalis*)).

prisca: Scarabaeidae (May beetle(-s)).

simplex: Scarabaeidae (*Phyllophaga*).

theutis: Scarabaeidae (*Phyllophaga*).

triangularis: Scarabaeidae (*Phyllophaga*).

spp.: Scarabaeidae (*Phyllophaga*, May beetle(-s)); Diprionidae (Diprionid (including *Diprion* and/or *Neodiprion*)).

Didyma

exigua: Pyralidae (*Acrobasis*, *Ostrinia*).

Doryphorophaga

australis: Chrysomelidae (*Leptinotarsa*, Chrysomelid (including *Lema* and/or *Leptinotarsa*)).

doryphorae: Chrysomelidae (*Leptinotarsa*, Chrysomelid (including *Lema* and/or *Leptinotarsa*)); Coccinellidae (*Epilachna*); Nymphalidae (*Nymphalis*).

macella: Chrysomelidae (*Leptinotarsa*).

spp.: Chrysomelidae (*Leptinotarsa*).

Elephantocera

greeni: Olethreutidae (*Grapholitha*).

Eribella

exilis: Chrysomelidae (*Chelymorpha*, *Plagiometriona*).

sp. near *exilis*: Diprionidae (*Neodiprion*); Olethreutidae (*Barbara*).

spp.: Pterophoridae (Unidentified Pterophoridae).

Erynniopsis

rondanii: Chrysomelidae (*Pyrrhalta*).

Eucelatoria

armigera: Diprionidae (*Neodiprion*); Arctiidae (*Estigmene*); Geometridae (Unidentified Geometridae); Noctuidae (*Achaea*, *Agrotis*, *Amyna*, *Autographa*, *Chrysodeixis*, *Heliothis*, *Leucania*, *Mocis*, *Plusia*, *Polydesma*, *Pseudaletia*, *Spodoptera*, *Trichoplusia*, Armyworm(-s)); Pieridae (*Pieris*); Pyralidae (*Hedylepta*, *Herpetogramma*, *Hymenia*).

armigera probably: Noctuidae (*Peridroma*).

australis: HesperIIDae (*Calpodes*); Noctuidae (*Leucania*); Pyralidae (*Agathodes*).

comosa: Noctuidae (Cutworm(-s)).

rubentis: Noctuidae (*Pseudaletia*, *Spodoptera*, Armyworm(-s)).

rubentis probably: Noctuidae (*Pseudaletia*).

spp.: Noctuidae (*Heliothis*); Pyralidae (*Hedylepta*, *Hymenia*).

Eucelatoriopsis

dimmocki: Chrysomelidae (*Agrioconota*, *Chelymorpha*, *Coptocycla*, *Metriona*, *Plagiometriona*).

dimmocki probably: Chrysomelidae (*Plagiometriona*).

"Gymnostylia"

sp.: Tenthredinidae (Unidentified Tenthredinidae).

Lixophaga

diatraeae: Arctiidae (*Utetheisa*); HesperIIDae (*Calpodes*); Noctuidae (*Spodoptera*); Pyralidae (*Diatraea*, *Zeadiatraea*); Unidentified Lepidoptera (Miscellaneous, Moth stalk-borer(-s)).

sp. near *fasciata*: Olethreutidae (*Petrova*).

grisea: Pyralidae (*Diatraea*).

jenei: Olethreutidae (*Laspeyresia*).

mediocris: Gelechiidae (*Frumenta*); Limacodidae (*Cnidocampa*); Olethreutidae (*Grapholitha*, *Laspeyresia*, *Rhyacionia*, *Sulleima*).

orbitalis: Olethreutidae (*Laspeyresia*).

parva: Curculionidae (*Lixus*); Olethreutidae (*Grapholitha*).

plumbea: Olethreutidae (*Grapholitha*, *Rhyacionia*); Phaloniidae (*Phalonia*); Pyralidae (*Pyrausta*); Tortricidae (*Xenotemna*).

sphenophori: Curculionidae (*Rhabdoscelus*); Pyralidae (*Diatraea*).

stenomae: Stenomidae (*Stenoma*).

variabilis: Bostrichidae (*Amphicerus*); Cerambycidae (*Oberea*); Curculionidae (*Lixus*, *Sphenophorus*); Tenthredinidae (Unidentified Tenthredinidae); Unidentified Hymenoptera (Sawfly(-flies)); Noctuidae (*Anomis*, *Papaipema*, *Stibadium*, *Trichoplusia*); Olethreutidae (*Ancylis*, *Epiblema*, *Eumarozia*, *Grapholitha*, *Laspeyresia*, *Rhyacionia*, *Suleima*); Pyralidae (*Desmia*, *Homoeosoma*, *Loxostege*, *Ostrinia*, Unidentified Pyralidae); Unidentified Lepidoptera (Miscellaneous).

sp. near *variabilis*: Curculionidae (*Hypera*).

spp.: Unidentified Coleoptera (Rhynchophora); Olethreutidae (*Laspeyresia*); Pyralidae (*Diatraea*).

Gen. near *Lixophaga*

spp.: Unidentified Hymenoptera (Sawfly(-flies)).

Lydellohoughia

tenella: Unidentified Lepidoptera (Tuber moth(-s)).

sp.: Pyralidae (Cone moth(-s) probably).

Gen. near *Lydinolydella*

spp.: Tortricidae (*Archips*).

Medina

barbata: Chrysomelidae (*Altica*, *Disonycha*, "*Luperodes*," *Plagiometriona*).

spp.: Chrysomelidae (Unidentified Chrysomelidae).

Meigenia

mutabilis: Chrysomelidae (*Crioceris*).

Myiopharus

canadensis: Chrysomelidae (*Calligrapha*).

dorsalis: Chrysomelidae (*Calligrapha*).

Oswaldia

albifacies: Noctuidae (*Parallelia*); Pyralidae (*Loxostege*).

anorbitalis: Noctuidae (*Parallelia*).

assimilis: Noctuidae (*Parallelia*).

aurifrons: Tenthredinidae (*Nematus*); Citheroniidae (*Anisota*).

sp. probably: Noctuidae (*Arzama*).

Oxynops

anthracinus: Pterophoridae (*Oidaematophorus*).

Gen. near Oxynops

spp.: Olethreutidae (*Petrova*); Pyralidae (*Dioryctria*).

Paralispe

infernalis: Chrysomelidae (*Crioceris*, *Lema*); Tortricidae (*Archips*).

Phrynofrontina

discalis: Lymantriidae (*Euproctis*).

spp.: Olethreutidae (*Petrova*); Pyralidae (*Dioryctria*).

"Prospheysa"

sp.: Sphecidae (*Bembix*).

Pseudomyothyria

ancilla: Chrysomelidae (*Diabrotica*); Scarabaeidae (*Serica*); Gelechiidae (*Filatima*).

spp.: Tenthredinidae (Unidentified Tenthredinidae); Unidentified Lepidoptera (Miscellaneous).

spp. probably: Unidentified Lepidoptera (Miscellaneous).

Schizocerophaga

leiby: Argidae (*Sphacophilus*); Tortricidae (*Archips*).

Sitophaga

calosomae: Carabidae (*Calosoma*, *Carabus*).

eleodivora: Tenebrionidae (*Eleodes*).

georgiae: Carabidae (*Calosoma*); Scarabaeidae (*Phyllophaga*).

georgiae probably: Carabidae (*Calosoma*).

leechi: Carabidae (*Carabus*).

neomexicana: Scarabaeidae (*Serica*).

nocturnal: Scarabaeidae (*Phyllophaga*).

sp.: Carabidae (*Calosoma*).

Spathimeigenia

aurifrons: Diprionidae (*Diprion*, *Monoctenus*, *Neodiprion*); Tenthredinidae (*Pikonema*, *Pristiphora*); Unidentified Hymenoptera (Sawfly(-flies) probably).

spp. (*aurifrons-erecta* complex): Diprionidae (*Neodiprion*).

erecta: Diprionidae (*Neodiprion*).

erecta probably: Diprionidae (*Neodiprion*).

hylotomae: Argidae (*Arge*); Diprionidae (*Neodiprion*); Unidentified Hymenoptera (Sawfly(-flies)).

hylotomae probably: Notodontidae (*Heterocampa*).

mexicana: Diprionidae (*Zadiprion*).

spinigera: Argidae (*Arge*); Diprionidae (*Diprion*, *Monoctenus*, *Neodiprion*, Diprionid (including *Diprion* and/or *Neodiprion*)); Tenthredinidae (*Pikonema*, *Pristiphora*); Unidentified Hymenoptera (Pine sawfly(-flies), Sawfly(-flies)); Unidentified Lepidoptera (Miscellaneous).

spinigera complex: Diprionidae (*Neodiprion*).

spp.: Diprionidae (*Diprion*, *Monoctenus*, *Neodiprion*).

Tachinophyto

floridensis: Pterophoridae (*Oidaematophorus*).

spp.: Gelechiidae (*Isophrictis*); Olethreutidae (*Epiblema*, *Laspheyresia*); Pyralidae (*Desmia*, *Diatraea*); Tineidae (Unidentified Tineidae).

Urophyllopsis

retiniae: Olethreutidae (*Rhyacionia*).

Tribe NEOMINTHOINI**Eupelecotheca**

celer: Limacodidae (*Packardia*).

Tribe EXORISTINI**Bessa**

harveyi: Argidae (*Arge*); Diprionidae (*Diprion*, *Neodiprion*); Tenthredinidae (*Amauronematus*, *Ametastegia*, *Anoplonyx*, *Cladius*, *Croesus*, *Hemichroa*, *Macremphytus*, *Nematus*, *Pikonema*, *Platycampus*, *Priophorus*, *Pristiphora*, *Strongylogaster*, *Trichiocampus*, Unidentified Tenthredinidae); Geometridae (*Eupithecia*, *Nepytia*, *Semiothisa*, Unidentified Geometridae); Nymphalidae (*Nymphalis*); Tortricidae (*Archips*, Unidentified Tortricidae).

selecta: Diprionidae (*Diprion*, *Neodiprion*); Tenthredinidae (*Hemichroa*, *Pikonema*, *Priophorus*, *Pristiphora*); Lymantriidae (*Hemerocampa*).

selecta probably: Tenthredinidae (*Pristiphora*).

spp.: Tenthredinidae (*Pristiphora*).

Chaetexorista

javana: Limacodidae (*Cnidocampa*).

Diplostichus

lophyri: Diprionidae (*Diprion*, *Neodiprion*, Diprionid (including *Diprion* and/or *Neodiprion*)); Tenthredinidae (*Pikonema*, *Pristiphora*); Lasiocampidae (*Malacosoma*).

sp. near *lophyri*: Diprionidae (*Neodiprion*).

sellersi: Diprionidae (*Neodiprion*).

Epiplagiops

littoralis: Curculionidae (*Listroderes*).

Euphorocera

claripennis: Chrysomelidae (*Calligrapha*); Coccinellidae (*Epilachna*); Unidentified Coleoptera (Miscellaneous); Diprionidae (*Diprion*, *Neodiprion*, Diprionid (including *Diprion* and/or *Neodiprion*)); Unidentified Hymenoptera (Sawfly(-flies));

Arctiidae (*Apantesis*, *Eubaphe*, *Euchaetias*, *Halisidota*, *Hyphantria*, *Utetheisa*); Citheroniidae (*Anisota*); Ctenuchidae (*Lymire*); Gelechiidae (*Anarsia*); Geometridae (*Cingilia*, *Itame*, *Philtraea*, *Semiothisa*); Hesperidae (*Epargyreus*); Lasiocampidae (*Malacosoma*, Apple tree tent caterpillar(-s)); Limacodidae (*Sibine*); Lymantriidae (*Euproctis*, *Hemerocampa*, *Nygma*, *Porthetria*, *Stilpnotia*, Tussock moth(-s)); Megalopygidae (*Lagoa*, *Megalopyge*); Noctuidae (*Acronicta*, *Actebia*, *Agrotis*, *Alabama*, *Amathes*, *Eubolina* probably, *Euxoa*, *Faronta*, *Feltia*, *Heliothis*, *Hydroecia*, *Lithophane*, *Mocis*, *Peridroma*, *Plathypena*, *Pseudaletia*, *Scotogramma*, *Spodoptera*, Armyworm(-s), Cutworm(-s)); Noctuidae and/or Pyralidae (Noctuid and/or pyralid (including *Scotogramma* and/or *Loxostege*)); Notodontidae (*Datana*, *Ichthyura*, *Schizura*); Nymphalidae (*Agraulis*, *Asterocampa*, *Chlosyne*, *Nymphalis*, *Polygonia*); Pieridae (*Colias*, *Pieris*); Psychidae (*Thyridopteryx*); Pyralidae (*Crambus*); Saturniidae (*Automeris*, *Callosamia*, *Hemileuca*, *Platysamia*); Sphingidae (*Ceratomia*, *Sphinx*); Zygaenidae (*Harrisina*); Unidentified Lepidoptera (Miscellaneous); Acrididae (*Melanoplus*, Grasshopper(-s)).

claripennis probably: Megalopygidae (*Megalopyge*).

sp. near *claripennis*: Diprionidae (*Diprion*).

edwardsii: Diprionidae (*Diprion*, *Neodiprion*); Lasiocampidae (*Malacosoma*); Lymantriidae (*Hemerocampa*); Noctuidae (*Euxoa*); Notodontidae (*Datana*, *Symmerista*); Nymphalidae (*Anaea*, *Asterocampa*, *Nymphalis*).

floridensis: Cerambycidae (*Saperda*); Coccinellidae (*Epilachna*); Arctiidae (*Apantesis*, *Hyphantria*); Citheroniidae (*Anisota*); Ctenuchidae (*Lymire*, *Syntomeida*); Geometridae (*Ennomos*, *Melanchroia*); Hesperidae (*Epargyreus*); Noctuidae (*Alabama*, *Anticarsia*, *Heliothis*, *Plathypena*, *Pseudaletia*, *Pseudoplusia*, *Spodoptera*); Notodontidae (*Datana*, *Ichthyura*); Sphingidae (*Ceratomia*); Zygaenidae (*Harrisina*).

sp. near *floridensis*: Pieridae (*Pieris*); Saturniidae (*Saturnia*).

omissa: Noctuidae (*Heliothis*, *Peridroma*, *Spodoptera*); Pieridae (*Colias*); Pyralidae (*Loxostege*).

subnitens probably: Zygaenidae (Gen. and sp. near *Triplocris*).

tachinomoides: Hesperidae (*Lerodea*); Lasiocampidae (*Malacosoma*, Apple tree tent caterpillar(-s)); Noctuidae (*Heliothis*, *Melipotis*, *Spodoptera*); Pieridae (*Pieris*); Psychidae (*Thyridopteryx*); Unidentified Lepidoptera (Green looper(-s)).

sp. near *tachinomoides*: Zygaenidae (*Harrisina*).

sp.: Diprionidae (*Neodiprion*).

Exorista

fasciata fasciata: Lymantriidae (*Byrdia*).

lobeliae: Lymantriidae (*Hemerocampa*); Noctuidae (*Acronicta*, *Alabama*).

sp. near *lobeliae*: Noctuidae (*Acronicta*).

mella: Tenthredinidae (*Macremphytus*); Arctiidae (*Apantesis*, *Cycnia*, *Diacrisia*, *Estigmene*, *Euchaetias*, *Halisidota*, *Hypphantria*, *Isia*); Bombycidae (Unidentified Bombycidae); Danaidae (*Danaus*); Lasiocampidae (*Epicnaptera*, *Malacosoma*, *Tolype*, Apple tree tent caterpillar(-s), Tent caterpillar(-s)); Lymantriidae (*Dasychira*, *Euproctis*, *Hemerocampa*, *Nygmia*, *Orgyia*, *Porthetria*, *Stilpnotia*, Tussock moth(-s)); Noctuidae (*Acronicta*, *Pseudaletia*, *Pyreferra*, *Simyra*); Notodontidae (*Datana*, *Ichthyura*, *Schizura*); Nymphalidae (*Nymphalis*, *Polygonia*); Saturniidae (*Callosamia*, *Hemileuca*).

mella probably: Arctiidae (*Estigmene*).

spp.: Arctiidae (*Diacrisia*, *Estigmene*, *Euchaetias*); Danaidae (*Danaus*); Geometridae (*Biston*); Lasiocampidae (*Malacosoma*); Lymantriidae (*Dasychira*, *Euproctis*, *Hemerocampa*, *Nygmia*, *Porthetria*, Tussock moth(-s)); Noctuidae (*Acronicta*, *Papaipema*); Notodontidae (*Cerura*); Olethreutidae and/or Pyralidae (Olethreutid and/or pyralid (including *Laspeyresia* and/or *Dioryctria* probably)); Pyralidae (*Acrobasis*, *Desmia*); Tortricidae (*Archips*).

spp. probably: Notodontidae (*Datana*); Unidentified Lepidoptera (Miscellaneous).

Guerinia

dydas: Diprionidae (*Diprion*); Tenthredinidae (*Allantus*, *Macremphytus*, *Pristiphora*); Unidentified Hymenoptera (Sawfly (-flies)).

rufostomata: Unidentified Hymenoptera (Sawfly(-flies)).

Palpexorista

coccyx: Limacodidae (*Phobetron*, *Sibine*); Notodontidae (*Datana*).

Parasetigena

agilis: Lasiocampidae (*Malacosoma*); Lymantriidae (*Porthetria*, *Stilpnotia*).

Pseudotachinomyia

slossonae: Geometridae (*Alsophila*, *Erannis*).

Spoggosia

gelida: Lymantriidae (*Dasychira* probably).

Stomatomyia

floridensis: Pyralidae (*Crambus*, *Elasmopalpus*, *Pyrausta*).

parvipalpis: HesperIIDae (*Hylephila*); Pyralidae (*Hulstia*, *Loxostege*); Gryllacrididae (*Stenopelmatus*); Unidentified Orthoptera (Pictured sand cricket(-s)).

sp.: Pyralidae (*Hellula*).

Tachinomyia

acosta: Noctuidae (*Lithophane*).

nigricans: Diprionidae (*Monoctenus*); Geometridae (*Alsophila*, *Ennomos*, *Erannis*, *Itame*, *Phigalia*); Lasiocampidae (*Malacosoma*); Lymantriidae (*Porthetria*); Noctuidae (*Acronicta*, *Baileya*, *Catocala*, *Lithophane*); Tortricidae (*Choristoneura*).

panaetius: Lasiocampidae (*Malacosoma*); Lymantriidae (*Hemerocampa*); Noctuidae (*Agrotis*, *Epiglaea*, Cutworm(-s)); Unidentified Lepidoptera (Moth(-s)).

sp. near *panaetius*: Noctuidae (*Acronicta*, *Catocala*, *Lithophane*, *Orthosia*).

similis: Arctiidae (*Halisidota*); Lasiocampidae (*Malacosoma*); Lymantriidae (*Hemerocampa*, *Stilpnotia*).

variata: Lasiocampidae (*Malacosoma*); Lymantriidae (*Porthetria*); Noctuidae (*Nephelodes*).

spp.: Geometridae (*Alsophila*, *Phigalia*); Lasiocampidae (*Malacosoma*); Lymantriidae (*Stilpnotia*); Noctuidae (*Acronicta*, *Catocala*, *Lithophane*, *Orthosia*, *Raphia*, Unidentified Catocalinae, Unidentified Noctuidae).

Tribe SIPHONINI

Subtribe CORONIMYIINA

Clausicella

floridensis: Olethreutidae (*Endothenia*, *Epiblema*, *Suleima*); Pyralidae (*Alberada*, *Homoeosoma*, *Plodia*).

neomexicana: Curculionidae (*Curculio*); Apidae (*Bombus*); Pyralidae (*Homoeosoma*, *Plodia*).

opaca: Noctuidae (*Peridroma*); Pyralidae (*Homoeosoma*).

turmalis: Geometridae (*Eupithecia*).

Coronimyia

geniculata: Pyralidae (*Moodna*, Unidentified Phycitinae); Pyralidae or Tortricidae (Unidentified Pyralidae or Tortricidae).

melitarae: Pyralidae (*Melitara*).

Drepanoglossa

amydriae: Tineidae (*Amydria*).

Ginglymia

acrirostris: Pyralidae (*Parargyractis*).

Philocalia

tenuirostris: Tineidae (*Amydria*).

Subtribe SIPHONINA

Actia

americana: Notodontidae (*Schizura*).

diffidens: Olethreutidae (*Spilonota*); Tortricidae (*Acleris*, *Choristoneura*).

flavipes: Dioptidae (*Phryganidia*); Geometridae (*Brephos*, *Cosymbia*).

interrupta: Geometridae (*Melanolopia*); Notodontidae (*Schizura*); Olethreutidae (*Epinotia*, *Grapholitha*); Tortricidae (*Acleris*, *Amorbia*, *Archippus*, *Archips*, *Choristoneura*, *Croesia*, *Phalonia*, *Xenotemna*, Unidentified Tortricidae).

sp. near *interrupta*: Tortricidae (*Archips*).

ontario: Geometridae (*Ennomos*).

palloris: Drepanidae (*Drepana*).

sp. near *palloris*: Geometridae (*Ennomos*, *Eugonobapta*).

spp.: Dioptidae (*Phryganidia*); Gelechiidae (*Gnorimoschema*); Noctuidae (*Catocala*, Unidentified Catocalinae); Notodontidae (*Heterocampa*); Olethreutidae (*Epinotia*, *Episimus*); Tortricidae (*Choristoneura*).

Eucoronimyia

hastata: Pyralidae (*Titanio*).

Pseudosiphona

brevirostris: Pterophoridae (*Oidaematophorus*).

Siphona

ceres: Noctuidae (Unidentified Noctuidae).

conata: Dioptidae (*Phryganidia*).

geniculata: Tipulidae (*Tipula*).

plusiae: Dioptidae (*Phryganidia*); Noctuidae (*Autographa*, *Trichoplusia*, Unidentified Noctuidae).

spp.: Tipulidae (Crane fly (flies)); Noctuidae (*Autographa*, *Trichoplusia*).

Strobliomyia

tibialis: Olethreutidae (*Grapholitha*).

Subtribe NEAERINA

Dichaetoneura

leucoptera: Nymphalidae (*Nymphalis*); Oecophoridae (*Depressaria*); Olethreutidae (*Epiblema*, *Grapholitha*); Tortricidae (*Archips*).

Elfia

aenea: Tortricidae (*Phalonia*).

sp. near *aenea*: Pyralidae (*Pyrausta*).

amplicornis: Glyphipterygidae (*Lotisma*).

curriei: Pyralidae (*Dioryctria*).

melissopodis: Curculionidae (*Curculio*); Olethreutidae (*Melissopus*).

palpiger: Pyralidae (*Acrobasis*).

sp. near *palpiger*: Tortricidae (*Argyrotaenia*).

setigera: Pyralidae (*Dioryctria*).

tarsalis: Pyralidae (*Pyralis*).

sp. near *usitata*: Oecophoridae (*Psilocorsis*).

spp.: Gelechiidae (*Coleotechnites*); Oecophoridae (*Psilocorsis*);
Olethreutidae (*Epinotia*, *Melissopus*); Pterophoridae (*Platyptilia*).

Phytomyptera

walleyi: Olethreutidae (*Zeiraphera*).

Schizactia

vitinervis: Gelechiidae (*Aristotelia*, *Coleotechnites*, *Exoteleia*);
Olethreutidae (*Epinotia*); Tineidae (Unidentified Tineidae).

Schizotachina

convecta: Pyralidae (*Acrobasis*); Tortricidae (*Croesia*).

sp.: Gelechiidae (*Chionodes*).

Tribe GRAPHOGASTRINI

Psalidopteryx

macdunnoughi: Scythridae (*Scythris*).

nuda: Olethreutidae (*Griselda*).

pollinosa: Gelechiidae (*Anarsia*).

pseudonuda: Tortricidae (*Acleris*).

psilocorsiphaga: Oecophoridae (*Psilocorsis*); Olethreutidae (*Epinotia*, *Eucosma*).

slossonae: Gelechiidae ("Telphusa" probably); Unidentified
Lepidoptera (Leaf tier(-s)).

sp.: Olethreutidae (*Eucosma*).

Tribe ADMONTIINI

Admontia

degeerioides: Tipulidae (*Tipula*); Noctuidae (*Spodoptera*); Olethreutidae (*Grapholitha*).

pergandei: Tipulidae (*Tipula*).

spp.: Tipulidae (*Ctenophora*).

Tribe HYPERECTEININI

Cartocometes

io: Lasiocampidae (*Malacosoma*).

Gilvella

gilvipes: Gryllidae (Cricket(-s)).

Hyperecteina

aldrichi: Scarabaeidae (*Maladera*, *Popillia*).

Miamimya

cincta: Blattidae (*Periplaneta*).

Oedematocera

dampfi: Acrididae (*Schistocerca*).

flaveola: Gryllacrididae (*Ceuthophilus*).

flaveola probably: Gryllacrididae (*Ceuthophilus*).

Phasmophaga

antennalis: Phasmatidae (*Diapheromera*).

antennalis probably: Phasmatidae (*Anisomorpha*).

meridionalis: Phasmatidae (*Anisomorpha*).

Roeseliopsis

americana: Phasmatidae (*Anisomorpha*).

Tribe ERYTHROCERINI

Elodia

flavipalpis: Olethreutidae (*Grapholitha*).

subfasciata: Olethreutidae (*Grapholitha*).

tragica: Olethreutidae (*Grapholitha*).

Erynnia

tortricis: Coleophoridae (*Coleophora*); Gelechiidae (*Anarsia*, *Pectinophora*); Noctuidae (*Arzama*); Olethreutidae (*Ancylis*, *Evora*, *Grapholitha*, *Laspeyresia*, *Melissopus*, *Proteoteras*, *Rhyacionia*); Psychidae (*Thyridopteryx*); Pyralidae (*Desmia*, *Etiella*, *Homoeosoma*, *Moodna*); Tortricidae (*Acleris*, *Archips*, *Choristoneura*, *Platynota*, *Sparganothis*, Unidentified Tortricidae, Unidentified Tortricidae probably); Unidentified Lepidoptera (Miscellaneous).

spp.: Tortricidae (*Acleris*).

Ocytata

pallipes: Forficulidae (*Forficula*).

Tribe FRONTININI

Eleodiphaga

caffreyi: Tenebrionidae (*Eleodes*).

pollinosa: Tenebrionidae (*Eleodes*).

Tribe GONIINI

Subtribe GONIINA

Acroglossa

hesperidarum: HesperIIDae (*Epargyreus*); Unidentified Lepidoptera (Butterfly(-flies)).

Eucnephalia

sp. probably: Noctuidae (*Protorthodes*).

Gonia

aldrichi: Noctuidae (*Agrotis*, *Crymodes*, *Euxoa*).

aldrichi probably: Noctuidae (*Euxoa*).

sp. near *aldrichi*: Noctuidae (*Agrotis*).

breviforceps: Noctuidae (*Euxoa*).

crassicornis: Scarabaeidae (*Phyllophaga*); Noctuidae (*Feltia*, *Prodenia*, *Spodoptera*).

frontosa: Meloidae (*Tetraonyx*); Noctuidae (*Crymodes*, *Feltia*).

fuscicollis: Noctuidae (*Euxoa*, *Feltia*).

longiforceps: Noctuidae (*Agrotis*, *Euxoa*).

longipulvilli: Noctuidae (*Agrotis*, *Feltia*, *Peridroma*).

porca: Noctuidae (*Peridroma*, *Pseudaletia*, Cutworm(-s)).

sagax: Geometridae (*Alsophila*).

"*sagax*": Arctiidae (Unidentified Arctiidae).

sequax: Noctuidae (*Agrotis*, *Peridroma*, *Spodoptera*).

texensis: Noctuidae (*Mocis*, *Spodoptera*).

spp.: Lymantriidae (*Nygmia*); Noctuidae (*Agrotis*, *Anticarsia*, *Crymodes*, *Euxoa*, *Heliothis*, *Lithophane*, *Peridroma*, *Scotogramma*, *Spodoptera*, Cabbage cutworm(-s)).

Spallanzania

hebes: Noctuidae (*Feltia*).

Subtribe CHAETOGAEDIINA

Chaetogaedia

analis: Arctiidae (Unidentified Arctiidae); Geometridae (*Ennomos*); Noctuidae (*Lacinipolia*).

sp. near *analis*: Noctuidae (*Heliothis*).

crebra: Noctuidae (*Agrotis*, *Protorthodes*, Cutworm(-s)); Pieridae (*Pieris*).

monticola: Danaidae (*Danaus*); Geometridae (*Eupithecia*, *Scotorythra*); Noctuidae (*Achaea*, *Actebia*, *Agrotis*, *Anomis*, *Autographa*, *Autoplusia*, *Carneades*, *Chrysodeixis*, *Elydna*, *Heliothis*, *Peridroma*, *Polydesma*, *Pseudaletia*, *Spodoptera*, Armyworm(-s) (including *Pseudaletia* and/or *Spodoptera*),

Armyworm(-s), Cutworm(-s) (including *Agrotis* and/or *Leucania*), Cutworm(-s), Unidentified Noctuidae); Nymphalidae (*Vanessa*); Pieridae (*Pieris*); Pyralidae (*Hedylepta*, *Hymenia*, *Oeobia*); Sphingidae (*Agrius*); Unidentified Lepidoptera (Miscellaneous).

monticola probably: Noctuidae (*Peridroma*).

sp. near *monticola*: Lasiocampidae (*Malacosoma*).

Gen. near *Chaetogaedia*

spp.: Noctuidae (*Litoprosopus*).

Tribe BELVOSIINI

Atacta

brasiliensis: Noctuidae (*Mocis*); Sphingidae (*Manduca*).

Belvosia

argentifrons: Unidentified Lepidoptera (Miscellaneous).

bicincta: Noctuidae (*Mocis*); Sphingidae (Hawk moth(-s)).

bifasciata: Citheroniidae (*Anisota*, *Citheronia*, *Eacles*); Notodontidae (*Datana*); Saturniidae (*Antheraea*, *Hemileuca*); Sphingidae (*Agrius*, *Ceratomia*); Unidentified Lepidoptera (Miscellaneous).

borealis: Sphingidae (*Ceratomia*).

ciliata: Hesperidae (Unidentified Hesperidae); Saturniidae (*Automeris*, *Copaxa*).

ferruginosa: Unidentified Lepidoptera (Miscellaneous).

formosa: Saturniidae (*Rothschildia*).

nigrifrons: Saturniidae (*Hyalophora*, *Rothschildia*).

recticornis: Saturniidae (*Hylesia*).

townsendi: Citheroniidae (*Citheronia*, *Eacles*); Sphingidae (*Ceratomia*); Unidentified Lepidoptera (Miscellaneous).

sp.: Citheroniidae (*Anisota*).

Choeteprosopa

hedemanni: Noctuidae (*Spodoptera*).

Triachora

luteola: Noctuidae (*Leucania*).

ochriventris: Noctuidae (Armyworm(-s)).

omissa: Noctuidae (*Feltia*).

unifasciata: Noctuidae (*Alabama*, *Pseudaletia*, Armyworm(-s)).

Tribe HARRISIINI

Leschenaultia

adusta: Arctiidae (*Arachnis*, *Epantheria*, *Estigmene*, *Halisidota*, *Hemihyalea*, Unidentified Arctiidae); Lasiocampidae (*Malacosoma*); Saturniidae (*Coloradia*).

- americana*: Arctiidae (*Halisidota*); Lasiocampidae (*Malacosoma*);
Noctuidae (*Pyrophila*).
- exul*: Lasiocampidae (*Malacosoma*, Tent caterpillar(-s)).
- fulvipes*: Lasiocampidae (*Malacosoma*); Saturniidae (*Hemileuca*).
- grossa*: Arctiidae (Unidentified Arctiidae).
- halisidotae*: Arctiidae (*Halisidota*).
- leucophrys*: Arctiidae (*Ecpantheria*, *Estigmene*, *Euchaetias*);
Lasiocampidae (*Malacosoma*); Lymantriidae (*Euproctis*);
Sphingidae (*Paonias*, *Sphinx*, Unidentified Sphingidae);
Unidentified Rodentia (Mouse (Mice) probably).
- spp.: Arctiidae (*Halisidota*, Woolly bear(-s)); Lasiocampidae
(*Malacosoma*); Lymantriidae (*Hemerocampa*); Unidentified
Lepidoptera (Miscellaneous).

Parachaeta

- fusca*: Arctiidae (*Isia*).
- spp.: Arctiidae (*Isia*); Unidentified Rodentia (Mouse (mice)).

Tribe STURMIINI

Blepharipa

- schineri*: Lymantriidae (*Porthetria*).
- scutellata*: Lasiocampidae (*Malacosoma*); Lymantriidae
(*Euproctis*, *Porthetria*, *Stilpnolia*); Noctuidae (*Catocala*,
Lithophane, Unidentified Catocalinae, Unidentified
Noctuidae); Notodontidae (*Datana*, *Symmerista*).
- sp.: Lymantriidae (*Porthetria*).

Drino

- austrina*: Pyralidae (Unidentified Pyralidae).
- sp. near *bakeri*: Sphingidae (*Amphion*).
- balloui*: Sphingidae (Sphinx moth(-s), Unidentified Sphingidae).
- bohemia*: Diprionidae (*Diprion*, *Neodiprion*); Tenthredinidae
(*Hemichroa*, *Pikonema*, *Pristiphora*); Unidentified Hymen-
optera (Sawfly(-flies)).
- cubaecola*: Pyralidae (*Hedylepta*).
- fraudulenta*: Hesperidae (*Erynnis*).
- inca*: Sphingidae (*Agrius*, Unidentified Sphingidae).
- incompta*: Citheroniidae (*Eacles*); Saturniidae (*Hemileuca*);
Sphingidae (*Agrius*, *Ceratonia*, *Hyles*, *Manduca*, *Pholus*,
Sphinx, Unidentified Sphingidae).
- inconspicua*: Diprionidae (*Diprion*, *Neodiprion*); Geometridae
(*Hydria*); Lasiocampidae (*Malacosoma*); Lymantriidae
(*Hemerocampa*, *Porthetria*); Noctuidae (*Alypia*).
- inconspicua* probably: Diprionidae (*Diprion*).

rhoeo: Psychidae (*Thyridopteryx*); Sphingidae (*Agrius*, *Manduca*, *Pholus*, Sphinx moth(-s), Unidentified Sphingidae).

rhoeo probably: Sphingidae (*Manduca*).

sp.: Sphingidae (Sphinx moth(-s)).

Mystacella

chrysoprocta: Scarabaeidae (*Xyloryctes*).

spp.: Arctiidae (*Isia*); Noctuidae (*Acronicta*).

Pales

spp.: Tenebrionidae (*Meracantha*); Notodontidae (*Cerura*).

Palexorista

laxa: Noctuidae (*Heliothis*, *Spodoptera*).

Siphosturmia

confusa: Nymphalidae (*Euphydryas*).

phyciodis: Nymphalidae (*Phyciodes*).

Sturmia

harrisinae: Zygaenidae (*Harrisina*).

harrisinae probably: Zygaenidae (*Acoloithus*, *Harrisina*).

sp. near *harrisinae*: Zygaenidae (*Harrisina*).

"*tipulensis*": Tipulidae (Unidentified Tipulini).

spp.: Argidae (*Arge*); Citheroniidae (*Anisota*); Sphingidae (*Manduca*); Tortricidae (*Choristoneura*).

sp. probably: Saturniidae (*Hemileuca*).

Gen. of Sturmiini

spp.: Noctuidae (*Anticarsia*).

Gen. near Thysanomyia

spp.: Sphingidae (*Hemaris*).

Townsendiellomyia

nidicola: Lymantriidae (*Euproctis*, *Nygmia*, *Porthetria*).

Tribe WINTHEMIINI

Hemisturmia

tortricis: Glyphipterygidae (*Anthophila*); Nymphalidae (*Nymphalis*); Olethreutidae (*Ancylis*); Pterophoridae (*Pterophorus*); Pyralidae (*Acrobasis*); Tortricidae (*Acleris*, *Archips*, *Argyrotaenia*, *Choristoneura*, Unidentified Tortricidae).

Nemorilla

pyste: Coccinellidae (*Epilachna*); Gelechiidae (*Filatima*); Glyphipterygidae (*Anthophila*); HesperIIDae (*Urbanus*); Lycaenidae (*Strymon*); Lymantriidae (*Porthetria*); Noctuidae (*Achatodes*, *Heliothis*, *Prodenia*, *Spodoptera*, *Trichoplusia*); Oecophoridae (*Depressaria*); Olethreutidae (*Ancylis*, *Episimus*, *Evora*, *Exartema*, *Grapholitha*, *Rhopobota*, *Rhyacionia*, *Spilonota*); Pyralidae (*Acrobasis*, *Asciodes*, *Desmia*, *Dia-*

phania, *Diatraea*, *Hedylepta*, *Hellula*, *Herpetogramma*, *Hulstia*, *Hymenia*, *Loxostege*, *Maruca*, *Nephopteryx*, *Ostrinia*, *Pachyzancla*, *Phlyctaenia*, *Pilocrocis*, *Pococera*, *Psara*, *Pyrausta*, *Tetralopha*, Unidentified Pyralidae); Tortricidae (*Acleris*, *Adoxophyes*, *Amorbia*, *Aphelia*, *Archippus*, *Archips*, *Argyrotaenia*, *Choristoneura*, *Pandemis*, *Sparganothis*, Unidentified Tortricidae); Yponomeutidae (*Yponomeuta*); Unidentified Lepidoptera (Miscellaneous).

pyste probably: Pyralidae (*Acrobasis*).

sp. near *pyste*: Pyralidae (*Acrobasis*, *Asciodes*).

"*trivittata*" probably: Noctuidae (*Acronicta*).

spp.: Tortricidae (*Choristoneura*).

Omotoma

fumiferanae: Geometridae (*Alsophila*, *Phigalia*); Noctuidae (*Amphipyra*, *Lithophane*, *Pyreferra*, *Spodoptera*); Tortricidae (*Choristoneura*).

Winthemia

abdominalis: Noctuidae (*Spodoptera*).

borealis: Noctuidae (*Scoliopteryx*).

cecropia: Saturniidae (*Actias*, *Antheraea*, *Platysamia*).

citheroniae: Citheroniidae (*Citheronia*, *Eacles*).

datanae: Arctiidae (*Diacrisia*, *Estigmene*, *Euchaetias*, Unidentified Arctiidae); Citheroniidae (*Anisota*); Lasiocampidae (*Malacosoma*); Lymantriidae (*Hemerocampa*); Noctuidae (*Pseudaletia*, Unidentified Catocalinae); Notodontidae (*Datana*, *Dicentria*, *Hyperaeschra*, *Pheosia*, *Schizura*); Saturniidae (*Platysamia*); Sphingidae (*Ceratomia*, *Cressonia*, *Paonias*, *Sphinx*).

datanae probably: Arctiidae (*Isia*); Noctuidae (*Acronicta*); Saturniidae (*Samia*).

sp. near *datanae*: Citheroniidae (*Anisota*).

deilephilae: Sphingidae (*Hyles*).

deilephilae probably: Sphingidae (*Sphinx*).

"*diversoides*": Geometridae (*Anacamptodes*); Noctuidae (*Achaea*).

infesta: Noctuidae (*Spodoptera*, Armyworm(-s)).

intermedia: Noctuidae (*Pseudoplusia*).

leucanae: Noctuidae (*Euthisanotia*, *Peridroma*, *Pseudaletia*, *Spodoptera*, Armyworm(-s), Cutworm(-s)); Pyralidae (*Pyralis*); Saturniidae (*Platysamia*); Sphingidae (*Manduca*).

leucanae lacking red at tip: Lasiocampidae (*Malacosoma*); Noctuidae (*Peridroma*, Agrotidian(-s)); Notodontidae (*Datana*); Saturniidae (*Platysamia*).

leucanae probably: Noctuidae (*Euthisanotia*, *Pseudaletia*).

- militaris*: Hesperidae (*Thymelicus*); Noctuidae (*Pseudaletia*, Armyworm(-s)); Saturniidae (*Platysamia*); Acrididae (Grasshopper(-s)).
- militaris* probably: Noctuidae (*Pseudaletia*).
- sp. near *militaris*: Lasiocampidae (*Malacosoma*); Noctuidae (*Pseudaletia*).
- occidentis*: Geometridae (*Lambdina*).
- ostensackenii*: Noctuidae (*Pseudaletia*).
- quadripustulata*: Curculionidae (*Curculio*, Unidentified Curculionidae); Diprionidae (*Diprion*); Arctiidae (*Halisidota*, White woolly bear(-s)); Bombycidae (Unidentified Bombycidae); Citheroniidae (*Anisota*); Geometridae (*Cingilia*, *Ematurga*); Lymantriidae (*Hemerocampa*, *Stilpnotia*, Tussock moth(-s)); Noctuidae (*Alypia*, *Amathes*, *Caenurgina*, *Cucullia*, *Euxoa*, *Faronta*, *Feltia*, *Heliothis*, *Leucania*, *Lithophane*, *Papaipema*, *Peridroma*, *Plathypena*, *Platysenta*, *Pseudaletia*, *Schinia*, *Spodoptera*, *Trichoplusia*, *Xylena*, Armyworm(-s), Cutworm(s), Cutworm(-s) and related Noctuidae, Unidentified Noctuidae); Notodontidae (*Datana*, *Heterocampa*, *Schizura*); Olethreutidae (*Spilonota*); Pyralidae (*Loxostege*); Saturniidae (*Antheraea*, *Callosamia*, *Platysamia*); Sphingidae (*Hemaris*, *Hyles*, *Manduca*, *Pachysphinx*, *Pholus*, *Sphinx*, Unidentified Sphingidae); Unidentified Lepidoptera (Miscellaneous).
- quadripustulata* probably: Noctuidae (*Ceramica*, *Epiglaea*, *Pseudaletia*); Nymphalidae (*Nymphalis*).
- rufopicta*: Geometridae (*Alsophila*, Cankerworm(-s)); Lasiocampidae (Tent caterpillar(-s)); Noctuidae (*Alypia*, *Anticarsia*, *Ceramica*, *Cucullia*, *Heliothis*, *Lithophane*, *Luperina*, *Oligia*, *Papaipema*, *Peridroma*, *Platysenta*, *Pseudaletia*, *Spaelotis*, *Spodoptera*, *Trichoplusia*, *Xylena*, Armyworm(-s)), Noctuid (including *Mamestra* and/or *Peridroma*), Noctuid (including *Heliothis*, *Prodenia* and/or *Spodoptera*); Sphingidae (*Hemaris*, Unidentified Sphingidae).
- rufopicta* probably: Noctuidae (*Amathes*, *Amphipyra*, *Epiglaea*); Notodontidae (*Heterocampa*).
- sinuata*: Geometridae (*Cingilia*, *Erannis*); Hesperidae (*Thymelicus*); Lasiocampidae (*Tolyte*); Noctuidae (*Luperina*, *Plathypena*, *Spodoptera*); Notodontidae (*Schizura*).
- sinuata* probably: Geometridae (*Cingilia*, Unidentified Geometridae); Noctuidae (*Calpe*); Nymphalidae (*Nymphalis*).
- tricolor*: Saturniidae (*Samia*).
- venusta* probably: Geometridae (*Lambdina*).
- spp. (*leucanae* probably or *militaris*): Noctuidae (*Pseudaletia*).

spp. (*quadripustulata* or allied sp.: *datanae*, *sinuata* or *vesiculata*): Chrysomelidae (*Leptinotarsa*).

spp.: Arctiidae (*Hyphantria*); Geometridae (*Lambdina*); Megalopygidae (*Megalopyge*); Noctuidae (*Catocala*, *Heliothis*, *Pseudaletia*, *Spodoptera*, Cutworm(-s)); Notodontidae (*Ichthyura*, *Schizura*, *Symmerista*); Pieridae (*Eucheira*); Saturniidae (*Callosamia*, *Hemileuca*, *Platysamia*); Sphingidae (*Manduca*); Tortricidae (*Choristoneura*).

spp. probably: Noctuidae (*Heliothis*).

Tribe CARCELIINI

Carcelia

amplexa: Arctiidae (*Ecpantheria*); Lymantriidae (*Euproctis*, *Hemerocampa*, *Orgyia*, *Porthetria*, Tussock moth(-s)); Megalopygidae (*Lagoa*, *Megalopyge*); Noctuidae (*Plathypena*, *Spodoptera*); Sphingidae (*Manduca*).

diacrisiae: Arctiidae (*Diacrisia*, *Estigmene*).

formosa: Noctuidae (*Acronicta*, *Agrotis*, *Lithophane*); Saturniidae (*Automeris*).

illota: Noctuidae (*Heliothis*).

lagoae: Arctiidae (*Ecpantheria*, *Halisidota*); Megalopygidae (*Megalopyge*); Pyralidae (*Eulepte*, *Omphalocera*).

laxifrons: Arctiidae (*Apantesis*, Unidentified Arctiidae); Lasiocampidae (*Malacosoma*); Lymantriidae (*Euproctis*, *Nygmia*, *Porthetria*); Notodontidae (*Schizura*).

malacosomae: Tenthredinidae (Unidentified Tenthredinidae); Arctiidae (*Hyphantria*, *Turuptiana*); Lasiocampidae (*Malacosoma*); Notodontidae (*Datana*); Saturniidae (*Hemileuca*).

malacosomae probably: Arctiidae (*Apantesis*, *Arachnis*, *Turuptiana*); Lasiocampidae (*Malacosoma*, American tent caterpillar(-s)); Tortricidae (*Archips*).

"ochracea": Pyralidae (*Ostrinia*).

olenensis: Lymantriidae (*Dasychira*).

perplexa: Lymantriidae (*Dasychira*, *Hemerocampa*).

protuberans: Arctiidae (*Hyphantria*); Lasiocampidae (*Malacosoma*).

reclinata: Arctiidae (*Diacrisia*, *Ecpantheria*, *Estigmene*, *Phragmatobia*, Unidentified Arctiidae); Unidentified Lepidoptera (Miscellaneous).

reclinata probably: Arctiidae (*Estigmene*, *Phragmatobia*).

yalensis: Arctiidae (*Halisidota*); Lymantriidae (*Hemerocampa*, *Orgyia*); Unidentified Lepidoptera (Miscellaneous).

spp.: Arctiidae (*Ecpantheria*, Unidentified Arctiidae); Limacodidae (*Adoneta*, *Phobetron*); Lymantriidae (*Dasychira*); Noctuidae (*Alypia*, *Peridroma*, *Pseudaletia*); Notodontidae (*Datana*); Olethreutidae (*Melissopus*); Pyralidae (*Pantographa*); Sphingidae (*Manduca*).

Eusisyropa

autographae: Noctuidae (*Trichoplusia*).

blanda: Arctiidae (*Hyphantria*); Citheroniidae (*Anisota*); Gelechiidae (Gelechiid webber(-s)); Geometridae (*Cingilia*, *Ennomos*, *Hydria*, *Isturgia*, *Lambdina*, *Nepytia*); HesperIIDae (*Calpodes*, *Epargyreus*, *Erynnis*); Limacodidae (*Euclea*); Lymantriidae (*Euproctis*, *Nygmia*, *Orgyia*, *Stilpnotia*); Noctuidae (*Alabama*, *Catocala*, *Hypena*, *Mocis*, *Phosphila*, *Plathypena*, *Spodoptera*, Unidentified Catocalinae); Nymphalidae (*Nymphalis*, *Polygonia*, *Vanessa*); Oecophoridae (*Depressaria*, *Psilocorsis*); Olethreutidae (*Gretchena*); Psychidae (*Thyridopteryx*); Pyralidae (*Acrobasis*, *Alatuncusia*, *Dichogama*, *Loxostege*, *Nephopteryx*, *Tetralopha*); Sphingidae (*Smerinthus*, *Sphexodina*); Tortricidae (*Archips*, *Argyrotaenia*, Unidentified Tortricidae); Unidentified Lepidoptera (Miscellaneous, Oakworm(-s)).

blanda probably: Gelechiidae (*Filatima*); Noctuidae (*Anomis*).

boarmiae: Arctiidae (*Hyphantria*); Geometridae (*Isturgia*); Noctuidae (*Alabama*, *Hypena*, *Plathypena*); Pyralidae (*Acrobasis*, *Dichogama*, *Loxostege*); Tortricidae (*Archips*); Yponomeutidae (*Urodus*).

boarmiae probably: Noctuidae (*Plathypena*); Pyralidae (*Acrobasis*).

virilis: Arctiidae (*Hyphantria*); Danaidae (*Danaus*); Diopsideae (*Phryganidia*); Geometridae (*Anacamptodes*, *Cingilia*, *Ennomos*, *Melanolophia*, *Phigalia*); HesperIIDae (*Epargyreus*); Limacodidae (*Euclea*); Lymantriidae (*Porthetria*); Noctuidae (*Acronicta*, *Papaipema*, *Phosphila*, *Polia*, *Simyra*); Notodontidae (*Gluphisia*, *Ichthyura*); Nymphalidae (*Polygonia*, *Vanessa*); Pieridae (*Eucheira*); Psychidae (*Thyridopteryx*); Saturniidae (*Callosamia*, *Platysamia*); Sphingidae (*Ceratomia*).

Gymnocarcelia

ricinorum: Arctiidae (*Apantesis*, *Diacrisia*, *Ecpantheria*, *Estigmene*, *Utethesia*, Unidentified Arctiidae); Noctuidae (*Pseudaletia*, Unidentified Noctuidae).

ricinorum probably: Arctiidae (*Ecpantheria*, *Estigmene*).

sp.: Arctiidae (*Estigmene*).

Hubneria

estigmenensis: Arctiidae (*Diacrisia*, *Estigmene*, *Isia*, *Phragmatobia*, Unidentified Arctiidae).

Hyphantrophaga

desmiae: Pyralidae (*Desmia*).

euchaetiae: Arctiidae (*Cycnia*, *Euchaetias*); Pyralidae (*Pyrausta*).

hyphantriae: Arctiidae (*Cycnia*, *Euchaetias*, *Hyphantria*); Citheroniidae (*Anisota*); Geometridae (*Eucaterva*, *Lygris*); Lasiocampidae (*Gloveria*, *Malacosoma*); Megalopygidae (*Megalopyge*); Noctuidae (*Heliothis*, *Spodoptera*); Nymphalidae (*Nymphalis*); Pyralidae (*Acrobasis*, *Desmia*, *Loxostege*, *Omphalocera*, *Pempelia*, Unidentified Pyralidae); Sphingidae (*Ceratonia*).

sp.: Nymphalidae (*Nymphalis*).

Sisyropa

alypiae: Noctuidae (*Alypia*, *Psychomorpha*).

alypiae probably: Noctuidae (*Alypia*).

eudryae: Arctiidae (*Estigmene*, Unidentified Arctiidae); Noctuidae (*Acronicta*, *Agrotis*, *Alypia*, *Euthisanotia*); Nymphalidae (*Vanessa*); Saturniidae (*Automeris*); Unidentified Lepidoptera (Moth(-s)).

eudryae probably: Noctuidae (*Acronicta*); Notodontidae (*Fen-tonia*); Nymphalidae (*Vanessa*).

spp.: Arctiidae (*Arachnis*); Lymantriidae (*Hemerocampa*); Megalopygidae (*Megalopyge*); Noctuidae (*Acronicta*, *Agrotis*); Notodontidae (*Heterocampa*).

Tribe EUMASICERINI**Argyrophylax**

albincisa: Pyralidae (*Eulepte*, *Hedylepta*, *Hymenia*, *Maruca*, *Pilemia*, *Pilocrocis*, *Psara*).

spp.: Pyralidae (*Etiella*); Saturniidae (*Hemileuca*); Sphingidae (*Agrius*, *Ceratonia*, *Pholus*).

sp. probably: Arctiidae (*Estigmene*).

Eumasicera

sternalis: Citheroniidae (*Anisota*).

sp. near *sternalis*: Noctuidae (*Acronicta*).

Metopiops

pyralidis: Nymphalidae (*Vanessa*); Pyralidae (Unidentified Pyralidae).

sp. near *pyralidis*: Pyralidae (*Tetralopha*).

Pacidianus

persimilis: Citheroniidae (*Anisota*).

sp.: Citheroniidae (*Anisota*).

Pseudochaeta

- argentifrons*: Arctiidae (*Cisthene*, *Tigrioides*); Bombycidae (Unidentified Bombycidae); Pyralidae (*Loxostege*).
siminina: Pyralidae (*Macrobotys*, *Palpita*).

Tribe ERYCIINI**Aplomya**

- caesar*: Arctiidae (*Diacrisia*); Gracillariidae (*Gracillaria*); Lymantriidae (*Nygmia*); Noctuidae (*Papaipema*); Olethreutidae (*Laspeyresia*); Pyralidae (*Crambus*, *Dioryctria*, *Loxostege*, *Ostrinia*, *Phlyctaenia*, *Pyrausta*); Tortricidae (*Amorbia*, *Archips*, *Argyrotaenia*, *Choristoneura*, *Pandemis*, *Phaecasiophora*, Tortricid spp. (including *Amorbia*, *Archips*, *Argyrotaenia*, *Choristoneura* and/or *Pandemis*)).
caesar probably: Tortricidae (*Archips*).
cerurae: Notodontidae (*Cerura*).
confusionis: Pyralidae (*Crambus*).
crassiseta: Tortricidae (*Choristoneura*).
epicydes: Cimbicidae (*Cimbex*).
fronto: Pamphiliidae (*Acantholyda*, *Neurotoma*).
helvina: Arctiidae (*Ecpantheria*); Geometridae (*Biston*, *Phaeoura*, Unidentified Geometridae).
mitis: Pyralidae (*Ostrinia*).
neurotomae: Pamphiliidae (*Neurotoma*).
pheosiae: Noctuidae (*Alypia*); Notodontidae (*Pheosia*).
theclarum: Lasiocampidae (*Gloveria*); Lycaenidae (*Brephidium*, *Celastrina*, *Lycaeides*, *Lycaena*, *Strymon*, Unidentified Lycaenidae); Unidentified Lepidoptera (Miscellaneous).
trichiosomae: Cimbicidae (*Trichiosoma*).
trisetosa: Noctuidae (*Autographa*, *Nephelodes*); Pyralidae (*Loxostege*).

Buquetia

- musca*: Papilionidae (*Papilio*).
obscura: Danaidae (*Danaus*); Papilionidae (*Papilio*).

Catagoniopsis

- meracanthae*: Tenebrionidae (*Meracantha*).
 sp. near *specularis*: Diprionidae (*Diprion*).

Ceromasia

- auricaudata*: Arctiidae (*Hyphantria*); Olethreutidae (*Sciaphila*); Pieridae (*Pieris*); Pyralidae (*Galleria*, *Ostrinia*); Tortricidae (*Archips*, *Choristoneura*).
aurifrons: Geometridae (*Cingilia*); Noctuidae (*Hypena*); Tortricidae (*Choristoneura*); Unidentified Lepidoptera (Miscellaneous).

spp.: Geometridae (*Lycia*, *Phigalia*); Noctuidae (*Archanara*, *Papaipema*); Tortricidae (*Choristoneura*).

Chrysoexorista

dawsoni: Geometridae (*Lambdina*); Olethreutidae (*Laspeyresia*).
marginata: Geometridae (*Hydria*); Olethreutidae (*Laspeyresia*).
ochracea: Olethreutidae (*Laspeyresia*).

Cyzenis

albicans: Geometridae (*Alsophila*, *Operophtera*); Lasiocampidae (*Malacosoma*); Olethreutidae (*Pseudexentera*).

“*festinans*” probably: Olethreutidae (*Pseudexentera*).

spp.: Olethreutidae (*Pseudexentera*).

Euceromasia

floridensis: Yponomeutidae (*Urodus*).

Euexorista

futilis: Arctiidae (*Apantesis*, *Euchaetias*, *Isia*); Geometridae (*Ennomos*); HesperIIDae (*Thymelicus*); Lasiocampidae (*Malacosoma*); Noctuidae (*Apamea*, *Autographa*, *Hyppa*, *Nephe- lodes*, *Nycteola*, *Sideridis*); Nymphalidae (*Nymphalis*, *Vanessa*); Pyralidae (*Ostrinia*); Saturniidae (*Antheraea*); Unidentified Lepidoptera (Miscellaneous).

Eufrontina

ethniae: Ethmiidae (*Ethmia*).

spectabilis: Pyralidae (*Canarsia*, *Tetralopha*).

Frontiniella

parancilla: Gelechiidae (*Gnorimoschema*); Pyralidae (*Tetra- lopha*).

Lespesia

aletiae: Coccinellidae (*Epilachna*); Arctiidae (*Diacrisia*, *Estig- mene*, *Halisidota*, *Hyphantria*, *Seirarctia*, *Utetheisa*, Unidenti- fied Arctiidae); Brassolidae (*Opsiphanes*); Ctenuchidae (*Ceramidia*, *Syntomeida*); Geometridae (*Ennomos*); Hesperi- idae (*Autochton*, *Epargyreus*, *Urbanus*); Lasiocampidae (*Malacosoma*); Limacodidae (*Sibine*); Lymantriidae (*Hemero- campa*, *Orgyia*, *Porthetria*, *Stilpnotia*, Tussock moth(-s)); Megalopygidae (*Megalopyge*); Noctuidae (*Alabama*, *Heliothis*, *Mocis*, *Plathypena*, *Pseudaletia*, *Pseudoplusia*, *Spodoptera*, *Trichoplusia*, *Xanthopastis*); Notodontidae (*Cerura*, *Dasy- lophia*, *Datana*, *Heterocampa*); Nymphalidae (*Asterocampa*, *Nymphalis*, *Polygonia*, *Vanessa*); Pieridae (*Pieris*); Pyralidae (*Evergestis*, *Olycella*); Saturniidae (*Hemileuca*); Sphingidae (*Ceratomia*, *Pholus*); Unidentified Lepidoptera (Miscellane- ous).

aletiae probably: Ctenuchidae (*Lymire*).

archippivora: Tenthredinidae (*Pristiphora*); Arctiidae (*Estigmene*, *Euchaetias*, *Halisidota*, *Tyria*); Citheroniidae (*Citheronia*); Danaidae (*Danaus*); Geometridae (*Anacamptodes*, *Scotorythra*); Lasiocampidae (*Malacosoma*, Tent caterpillar(-s)); Lycaenidae (*Lampides*); Noctuidae (*Agrotis*, *Alypia*, *Anomis*, *Autographa*, *Elydna*, *Euxoa*, *Feltia*, *Heliothis*, *Peridroma*, *Polydesma*, *Pseudaletia*, *Scotogramma*, *Spodoptera*, *Trichoplusia*, Armyworms (including *Pseudaletia* and/or *Spodoptera*), Armyworm(-s), Cutworms (including *Agrotis* and/or *Leucania*), Cutworm(-s), Noctuid (including *Mamestra* and/or *Peridroma*); Noctuidae and/or Pyralidae (Noctuid and/or pyralid (including *Scotogramma* and/or *Loxostege*)); Notodontidae (*Schizura*); Nymphalidae (*Anaea*, *Nymphalis*, *Vanessa*); Papilionidae (*Papilio*); Pieridae (*Colias*, *Pieris*); Pyralidae (*Hedylepta*, *Loxostege*); Yponomeutidae (*Mapsidius*); Unidentified Lepidoptera (Cabbage worm(-s), Miscellaneous).

archippivora probably: Noctuidae (*Spodoptera*); Nymphalidae (*Vanessa*); Sphingidae (*Sphinx*).

callosamiae: Saturniidae (*Callosamia*).

ciliata: Bombycidae (*Bombyx*); Lasiocampidae (*Malacosoma*); Pyralidae (*Loxostege*); Saturniidae (*Hyalophora*, *Platysamia*).

cuculliae: Noctuidae (*Cucullia*); Notodontidae (*Dasylophia*, *Datana*).

cuculliae probably: Notodontidae (*Dasylophia*).

sp. near *cuculliae*: Arctiidae (*Estigmene*).

datanarum: Citheroniidae (*Anisota*); Notodontidae (*Datana*); Saturniidae (*Antheraea*).

dimmocki: Noctuidae (*Simyra*); Saturniidae (*Automeris*); Unidentified Lepidoptera (Miscellaneous).

dubia: Nymphalidae (*Nymphalis*).

euchaetiae: Arctiidae (*Euchaetias*, *Halisidota*).

flavifrons: Sphingidae (*Paonias*, *Smerinthus*).

frenchii: Apidae (*Bombus*); Arctiidae (*Diacrisia*, *Euchaetias*, *Halisidota*, *Hyphantria*); Bombycidae (Unidentified Bombycidae); Citheroniidae (*Anisota*, *Citheronia*, *Eacles*); Geometridae (Unidentified Geometridae); Lasiocampidae (*Malacosoma*, American tent caterpillar(-s), Tent caterpillar(-s)); Lymantriidae (*Hemerocampa*, *Nygmia*, *Porthetria*, *Stilpnotia*); Noctuidae (*Faronta*, *Heliothis*, *Hydroecia*, *Hypso-ropha*, Unidentified Noctuidae); Notodontidae (*Datana*, *Ichthyura*); Nymphalidae (*Nymphalis*, *Vanessa*); Papilionidae (*Papilio*); Pieridae (*Pieris*); Pyralidae (Unidentified Pyralidae); Saturniidae (*Actias*, *Antheraea*, *Attacus*, *Automeris*, *Callo-*

samia, *Hyalophora*, *Platysamia*, *Samia*, *Syssphinx*); Sphingidae (*Ceratonia*, *Manduca*, *Pachysphinx*, *Paonias*, *Smerinthus*, *Sphinx*, Unidentified Sphingidae); Unidentified Lepidoptera (Miscellaneous); Acrididae (*Dissosteira*).

frenchii complex: Notodontidae (*Datana*).

laniiferae: Pyralidae (*Laniifera*).

sp. near *laniiferae*: Arctiidae (*Eupseudosoma*).

melalophae: Geometridae (*Metanema*); Lasiocampidae (*Malacosoma*); Noctuidae (*Nycteola*); Notodontidae (*Ichthyura*).

pholi: Sphingidae (*Pholus*).

rileyi: Papilionidae (*Papilio*).

sabroskyi: Saturniidae (*Antheraea*, *Automeris*, *Platysamia*).

schizurae: Arctiidae (*Euchaetias*); Danaidae (*Danaus*); Lasiocampidae (*Malacosoma*); Notodontidae (*Heterocampa*, *Schizura*); Sphingidae (*Pholus*).

testacea: Noctuidae (*Xerociris*); Sphingidae (*Pholus*).

spp.: Arctiidae (*Ecpantheria*); Citheroniidae (*Anisota*, *Citheronia*); Ctenuchidae (*Syntomeida*); Danaidae (*Danaus*); Lasiocampidae (*Malacosoma*); Lycaenidae (*Strymon*); Lymantriidae (*Hemerocampa*); Megathymidae (*Megathymus*); Noctuidae (*Acronicta*, *Autographa*, *Celama*, *Faronta*, *Heliothis*, *Hypso-ropha*, *Simyra*, *Spodoptera*, *Trichoplusia*, Unidentified Noctuidae); Notodontidae (*Datana*, *Heterocampa*, *Ichthyura*); Nymphalidae (*Vanessa*); Olethreutidae (*Eucosma*, *Grapholitha*); Pieridae (*Colias*, *Pieris*); Psychidae (*Oiketeticus*); Pyralidae (*Diatraea*, *Laniifera*, *Melitara*, *Pyrausta*, Unidentified Pyralidae); Saturniidae (*Antheraea*, *Hemileuca*, *Rothschildia*); Sphingidae (*Ceratonia*, *Manduca*); Acrididae (*Dissosteira*, *Melanoplus*, Locust(-s)).

Lydella

deckeri: Noctuidae (*Papaipema*).

radicis: Noctuidae (*Achatodes*, *Archanara*, *Arzama*, *Hydroecia*, *Luperina*, *Macronoctua*, *Papaipema*); Olethreutidae (*Epiblema*); Pyralidae (*Ostrinia*); Unidentified Lepidoptera (Miscellaneous).

radicis probably: Noctuidae (*Achatodes*, *Papaipema*).

thompsoni: Noctuidae (*Papaipema*); Pyralidae (*Galleria*, *Ostrinia*).

sp. probably: Argidae (*Arge*).

Madremyia

saundersii: Curculionidae (*Hypera*); Danaidae (*Danaus*); Geometridae (*Cingilia*, *Coryphista*, *Deuteronomos*, *Ennomos*, *Lambdina*); Lasiocampidae (*Malacosoma*); Lymantriidae (*Euproctis*, *Nygma*); Noctuidae (*Agrotis*, *Autographa*, *Epiglaea*,

Peridroma, *Pseudaletia*, *Trichoplusia*, *Xylena*); Nymphalidae (*Nymphalis*, *Speyeria*, *Vanessa*); Olethreutidae (*Sciaphila*); Pieridae (*Pieris*); Pyralidae (*Ostrinia*); Tortricidae (*Acleris*, *Choristoneura*); Unidentified Lepidoptera (Looper(-s), Miscellaneous).

“Masicera”

“*tantilla*”: Pyralidae (*Acigona*).

spp.: Citheroniidae (*Anisota*); Noctuidae (*Simyra*); Pieridae (*Colias*); Sphingidae (*Manduca*).

Metagonistylum

minense: Pyralidae (*Diatraea*).

Nilea

carpocapsae: Olethreutidae (*Laspeyresia*).

noctuiformis: Noctuidae (Unidentified Noctuidae); Olethreutidae (*Laspeyresia*).

sp. near *noctuiformis*: Noctuidae (*Acronicta*).

sternalis: Noctuidae (*Acronicta*); Tortricidae (*Archips*).

Obolocera

rapae: Pieridae (*Pieris*).

Palpozenillia

palpalis: Castniidae (*Castnia*); Pyralidae (*Diatraea*).

Patelloa

fuscimacula: Lymantriidae (*Hemerocampa*); Saturniidae (*Antheraea*).

leucaniae: Lymantriidae (*Euproctis*, *Hemerocampa*); Noctuidae (*Alabama*, *Pseudaletia*, Armyworm(-s)); Notodontidae (*Datana*, *Heterocampa*); Psychidae (*Thyridopteryx*); Pyralidae (*Loxostege*); Unidentified Lepidoptera (Miscellaneous).

leucaniae probably: Lymantriidae (*Euproctis*).

sp. near *leucaniae*: Chrysomelidae (*Lema*).

pachypyga: Lasiocampidae (*Malacosoma*); Lymantriidae (*Euproctis*, *Nygma*).

pachypyga probably: Nymphalidae (*Nymphalis*).

spp.: Geometridae (*Drepanulatrix*); Noctuidae (*Crocigraha*).

“Phorocera”

comstocki: Diprionidae (Diprionid (including *Diprion* and/or *Neodiprion*)); Cossidae (*Cossula*); Megathymidae (*Megathymus*); Pyralidae (*Melitara*, *Olycella*, *Ostrinia*).

halisidotae: Arctiidae (*Halisidota*).

incrassata: Tortricidae (*Choristoneura*).

“*parva*”: Pyralidae (*Loxostege*, *Udea*); Tortricidae (*Argyrotaenia*).

signata: Cossidae (*Cossula*); Noctuidae (*Papaipema*).

texana: Megathymidae (*Agathymus*); Pyralidae (*Melitara*, *Olycella*).

“*victoria*”: Arctiidae (Unidentified Arctiidae probably).

spp.: Diprionidae (*Neodiprion*); Tenthredinidae (*Pristiphora*); Unidentified Hymenoptera (Pitch pine sawfly(-flies)); Arctiidae (*Estigmene*); Lymantriidae (*Hemerocampa*); Noctuidae (*Trichoplusia*, Armyworm(-s)); Nymphalidae (*Chlosyne*, *Nymphalis*); Pieridae (*Colias*); Pyralidae (*Diatraea*, *Laniifera*, *Loxostege*); Zygaenidae (*Harrisina*); Unidentified Lepidoptera (Hog moth(-s)).

Phryxe

pecosensis: Tenthredinidae (*Pikonema*); Danaidae (*Danaus*); Geometridae (*Alsophila*, *Cingilia*, *Ennomis*, *Erannis*, *Lambdina*, *Operophtera*); HesperIIDae (*Thymelicus*); Noctuidae (*Heptagrotis*, *Lithophane*, *Pseudaletia*, Noctuid (including *Mamestra* and/or *Peridroma*), Unidentified Noctuidae); Pieridae (*Pieris*); Pyralidae (*Dioryctria*, *Nephopteryx*); Saturniidae (*Hemileuca*); Tortricidae (*Acleris*, *Aphelia*, *Archips*, *Choristoneura*); Unidentified Lepidoptera (Miscellaneous).

pecosensis probably: Geometridae (*Synchlora*); Noctuidae (*Nephelodes*); Pyralidae (*Ostrinia*); Tortricidae (*Choristoneura*).

vulgaris: Danaidae (*Danaus*); Geometridae (*Alsophila*, *Cingilia*, *Ennomos*, *Erannis*); Lymantriidae (*Porthetria*); Noctuidae (*Lithophane*, *Pseudaletia*, *Trichoplusia*, Unidentified Noctuidae); Pieridae (*Pieris*); Pyralidae (*Evergestis*, *Loxostege*); Saturniidae (*Hemileuca*); Tortricidae (*Archips*, *Choristoneura*).

sp.: Pyralidae (*Loxostege*).

Pilatea

arator: Tipulidae (Unidentified Tipulidae); Lasiocampidae (*Malacosoma*).

delecta: Noctuidae (*Autographa*).

Prophryno

marginalis: Geometridae (*Semiothisa*).

parviteres: Geometridae (*Melanchroia*); Noctuidae (*Plathypena*); Pieridae (*Ascia*).

“Prosopaea”

spp.: Noctuidae (*Aletia*); Notodontidae (*Cerura*); Sphingidae (*Smerinthus*).

Pseudoperichaeta

erecta: Meloidae (*Epicauta*); Noctuidae (*Nycteola*); Notodontidae (*Schizura*); Olethreutidae (*Episimus*, *Evora*, *Grapholitha*, *Melissopus*, *Spilonota*); Pyralidae (*Desmia*, *Loxostege*,

Macrobotys, *Ostrinia*, *Pachyzancla*, *Pantographa*, *Phlyctaenia*, *Pyrausta*); Tortricidae (*Amorbia*, *Archips*, *Argyrotaenia*, *Choristoneura*, Tortricid (including *Amorbia*, *Archips*, *Argyrotaenia*, *Choristoneura*, and/or *Pandemis*)); Unidentified Lepidoptera (Leaf roller(-s)).

erecta probably: Tortricidae (*Amorbia*).

sp. near *erecta*: Olethreutidae (*Melissopus*).

insidiosa: Olethreutidae (*Grapholitha*); Pyralidae (*Ostrinia*).

Thelymyia

mathesoni: Lymantriidae (Tussock moth(-s) probably).

Trepophrys

cinerea: Tineidae (*Amydria*).

Tsugaea

nox: Diprionidae (*Neodiprion*).

Zenillia

blandita: Noctuidae (*Nycteola*).

libatrix: Arctiidae (*Euchaetias*); Bombycidae (*Bombyx*); Lymantriidae (*Euproctis*); Notodontidae (*Ichthyura*).

spp.: Diprionidae (*Neodiprion*); Arctiidae (*Halisidota*, *Haploa*); Megalopygidae (*Megalopyge*); Noctuidae (*Acronicta*, *Psychomorpha*); Pyralidae (*Acrobasis*, *Ostrinia*); Sphingidae (Unidentified Sphingidae).

Gen. near Zenillia

spp.: Noctuidae (Unidentified Noctuidae).

Unplaced as to Subfamily

Unplaced as to Tribe

“Tachina”

“*desmiae*”: Pyralidae (*Desmia*).

spp.: Chrysomelidae (*Coptocycla*); Cicadidae (*Magiccada*); Apidae (*Bombus*); Unidentified Hymenoptera (Bee(-s)); Aegeriidae (*Vespa mima*); Arctiidae (*Hyphantria*); Geometridae (Cutworm(-s) (including *Alsophila* and/or *Paleacrita*)); Lasiocampidae (*Malacosoma*); Noctuidae (*Alabama*, *Papaipema*, *Pseudaletia*, *Trichoplusia*, Cutworm(-s)); Psychidae (*Thyridopteryx*); Pyralidae (*Acrobasis*, *Crambus*); Saturniidae (*Antheraea*, *Callosamia*, *Platysamia*); Sphingidae (*Hyles*); Unidentified Lepidoptera (Miscellaneous, Moth(-s)); Phasmatidae (*Bacillus*, *Diapheromera*).

Tachinidae

COLEOPTERA: Buprestidae (*Buprestis*); Carabidae (*Carabus*, *Pterostichus*); Cerambycidae (*Enaphalodes*, *Monochamus*,

Saperda); Chrysomelidae (*Acalymma*, *Altica*, *Chrysomela*, *Diabrotica*, *Leptinotarsa*, *Physonota*); Cleridae (*Thanasimus*); Coccinellidae (*Epilachna*); Curculionidae (*Conotrachelus*, *Rhodoabaenus*, *Sphenophorus*); Scarabaeidae (*Phyllophaga*, May beetle(-s)); Tenebrionidae (*Eleodes*).

DERMAPTERA: Carcinophoridae (*Anisolabis*).

DIPTERA: Cecidomyiidae (*Rhabdophaga*); Tabanidae (*Hybomitra*, Unidentified Tabanidae); Tipulidae (*Tipula*).

HEMIPTERA (HETEROPTERA): Coreida (*Anasa*); Pentatomidae (*Nezara*, Plant bug(-s)); Rhopalidae (*Liorhyssus*).

HEMIPTERA (HOMOPTERA): Cicadidae (*Magicicada*).

HYMENOPTERA: Argidae (*Sphacophilus*); Cimbicidae (*Cimbex*); Diprionidae (*Diprion*); Tenthredinidae (*Nematus*, *Tethida*).

LEPIDOPTERA: Arctiidae (*Cycnia*, *Diacrisia*, *Estigmene*, *Euchaetias*, *Halisidota*, *Hyphantria*, *Isia*, *Phragmatobia*, *Utetheisa*); Brassolidae (*Brassolis*); Citheroniidae (*Anisota*, *Eacles*); Ctenuchidae (*Ctenucha*); Danaidae (*Danaus*); Diop-
tidae (*Phryganidia*); Drepanidae (*Drepana*); Geometridae (*Alsophila*, *Cingilia*, *Coniodes*, *Deilinia*, *Ennomos*, *Erannis*, *Eucaterva*, *Eulype*, *Itame*, *Lambdina*, *Nematocampa*, *Nepytia*, *Paleacrita*, *Phaeoura*, *Phigalia*, *Sabulodes*); Hepialidae (*Hepialus*); Hesperiididae (*Cymaenes*, *Epargyreus*, *Erynnis*, *Thymelicus*, Hesperiid (including *Cymaenes*, *Nyctelius* and/or *Panoquina*)); Lasiocampidae (*Epicnaptera*, *Malacosoma*, American tent caterpillar(-s)); Limacodidae (*Euclea*, *Limacodes* probably, *Phobetron*, *Prolimacodes*, *Sibine*); Lycaenidae (*Brephidium*, *Plebejus*); Lymantriidae (*Hemerocampa*, *Orgyia*, *Porthetria*, *Stilpnotia*, Tussock moth(-s)); Megalopygidae (*Megalopyge*); Noctuidae (*Acronicta*, *Actebia*, *Agrotis*, *Alabama*, *Aletia*, *Alypia*, *Amphipyra*, *Callopietria*, *Catocala*, *Ceramica*, *Chrysodeixis*, *Cirphis*, *Cosmia*, *Crymodes*, *Cucullia*, *Epiglaea*, *Euparthenos*, *Eupsilia*, *Euthisanotia*, *Euxoa*, *Faronta*, *Heliothis*, *Lithophane*, *Mamestra*, *Nephelodes*, *Orthodes*, *Papaipema*, *Parallelia*, *Peridroma*, *Pseudaletia*, *Pyreferra*, *Spodoptera*, *Trichoplusia*, Armyworm(-s), Cutworm(-s), Unidentified Noctuidae); Notodontidae (*Cerura*, *Dasylophia*, *Datana*, *Heterocampa*, *Hyperaeschra*, *Ichthyura*, *Nadata*, *Schizura*, *Symmerista*); Nymphalidae (*Chlosyne*, *Limenitis*, *Nymphalis*, *Polygonia*); Oecophoridae (*Depressaria*, *Psilocorsis*); Olethreutidae (*Grapholitha*, *Rhyacionia*); Papilionidae (*Papilio*); Pieridae (*Pieris*, *Zerene*); Plutellidae (*Harpigteryx*); Psychidae (*Thyridopteryx*); Pterophoridae (*Oidaematophorus*, *Trichoptilus*); Pyralidae (*Acrobasis*,

Crambus, *Diaphania*, *Diatraea*, *Dioryctria*, *Hedylepta*, *Loxostege*, *Macrobotys*, *Nephopteryx*, *Phlyctaenia*, *Pyrausta*, *Tetralopha*); Saturniidae (*Actias*, *Antheraea*, *Automeris*, *Callosamia*, *Hemileuca*, *Platysamia*); Sphingidae (*Agrius*, *Ceratomia*, *Cressonia*, *Hemaris*, *Hyles*, *Pachysphinx*, *Paonias*, *Pholus*, *Smerinthus*, *Sphinx*); Stenomidae (*Antaeotricha*); Thyatiridae (*Pseudothyatira*); Tortricidae (*Archips*, *Choristoneura*, *Sparganothis*); Yponomeutidae (*Urodus*); Unidentified Lepidoptera (Miscellaneous).

ORTHOPTERA: Acrididae (*Ageneotettix*, *Camnula*, *Chimarocephala*, *Dissosteira*, *Melanoplus*, *Orphulella*, *Spharagemon*, Grasshopper(-s), Locust(-s), Unidentified Acrididae); Mantidae (*Stagmomantis*); Phasmatidae (*Bacillus*, *Diapheromera*).

PARASITES

This is the only section containing bibliographic information. The others must be used in conjunction with it in order to locate published records. Both the tachinid genera and species within each genus are arranged alphabetically. The subfamily and tribal assignment is given for each genus. The specific names are followed by broader categories, such as "spp." Under each tachinid species the host families are arranged alphabetically.

The Diptera Catalog (see Sabrosky and Arnaud, 1965) is the source of the tachinid names used in this section. Names not included in that publication were reviewed by Curtis W. Sabrosky for their current assignments. Quotation marks used with a parasite heading name indicate a questioned or unknown status. For example, "*Anisia nigrocincta* Wulp" is unrecognized both as to genus and species, whereas "*Phorocera*" *texana* Aldrich and Webber is a recognized species, but the present generic assignment is unknown.

Synonyms, obsolete names, and incorrect spellings of parasite names are cross indexed to currently accepted names, with misspellings enclosed by quotation marks. Such general references as tachina fly, Tachinidae, and Larvaevoridae are placed together at the end of this section under the heading Tachinidae.

Following the parasite name the statements "Most records as . . ." or "All records as . . ." are used if most or all the references are of a different combination from the accepted parasite name.

Under the heading "HOST," the orders (and within each order, the families, genera, and species) are arranged alphabetically. Nonscientific names assignable to a family follow the scientific names. The bibliographic citations are listed under the host species names.

Following the host name the statements "Most records ex . . ." or "All records ex . . ." are used if most or all the references are of a different combination from the accepted host name.

Each bibliographic citation lists the author(s), year, and page(s). Illustration(s), locality(ies), and comments follow if pertinent.

The most recent reference is listed first. If two or more references occur in one year, the authors' names are alphabetized.

Vide preceding an author's name means that he refers an earlier rearing to a different parasite species, as well as reporting a new rearing. If an asterisk precedes the vide, he refers an earlier rearing to a different parasite species, but he does not report a new rearing.

Abbreviations for the States of the United States and the Provinces of Canada are used without the country name and are alphabetized without distinction to country. The States of Mexico are also abbreviated, enclosed by parentheses, and listed after the United States-Canadian distribution. With the exception of Puerto Rico, the names of other countries are not abbreviated. The locality abbreviations are as follows:

Ala.—Alabama	Nay.—Nayarit
Alta.—Alberta	N.B.—New Brunswick
Amer.—America	N.C.—North Carolina
Ariz.—Arizona	N. Dak.—North Dakota
Ark.—Arkansas	Nebr.—Nebraska
B.C.—British Columbia	Nev.—Nevada
Calif.—California	Nfld.—Newfoundland
Camp.—Campeche	N.H.—New Hampshire
Cent.—central	N.J.—New Jersey
Chih.—Chihuahua	N. Mex.—New Mexico
Colo.—Colorado	N.S.—Nova Scotia
Conn.—Connecticut	N.Y.—New York
D.C.—District of Columbia	Okla.—Oklahoma
Del.—Delaware	Ont.—Ontario
D.F.—Distrito Federal	Oreg.—Oregon
Dgo.—Durango	Pa.—Pennsylvania
E.—east; eastern	P.E.I.—Prince Edward Island
Fla.—Florida	P.R.—Puerto Rico
Ga.—Georgia	prov.—province(-s)
Ill.—Illinois	Pue.—Puebla
Ind.—Indiana	Que.—Quebec
Kans.—Kansas	R.I.—Rhode Island
Ky.—Kentucky	S.—south; southern
La.—Louisiana	Sask.—Saskatchewan
Man.—Manitoba	S.C.—South Carolina
Mass.—Massachusetts	S. Dak.—South Dakota
Md.—Maryland	Sin.—Sinaloa
Mex.—Mexico	S.L.P.—San Luis Potosi
Mich.—Michigan	Son.—Sonora
Micho.—Michoacan	Tenn.—Tennessee
Minn.—Minnesota	Tex.—Texas
Miss.—Mississippi	U.S.—United States
Mo.—Missouri	Va.—Virginia
Mont.—Montana	V.C.—Vera Cruz
Mor.—Morelos	Vt.—Vermont
N.—north; northern	W.—west; western

Wash.—Washington
 Wis.—Wisconsin

W. Va.—West Virginia
 Wyo.—Wyoming

The following special terms are used:

“as” = parasite name as published. Only that part that differs from the heading is cited.

“ex” = host name as published. Only that part that differs from the heading is cited.

“incomplete rearing” = host parasitized without emergence of adult parasites.

“lab.” = laboratory rearing of specimens usually for release in the field.

“probable host” = rearing not fully documented.

“probable parasite” = rearing not fully documented.

References containing information on the “STAGE PARASITIZED,” “IMMATURES,” or “BIOLOGY” are listed under those headings at the end of each tachinid species section. Headings are omitted if no relevant data are found in the literature that was reviewed.

The section “STAGE PARASITIZED” reports only published data. No interpolation is made. For example, under Lepidoptera if the pupa is mentioned as being parasitized, only the pupal stage is cited, even though the larval stage would have been parasitized also.

The section “IMMATURES” cites descriptions and illustrations of the parasite immature stages in the following sequence: Egg, larva, first instar larva, second instar larva, third instar larva, and pupa.

A description is cited by “descr.” and an illustration by plate, figure numbers, or both.

Under “BIOLOGY,” biological information not included in the previous sections is cited. The pages in parentheses refer to the rearing.

Acaulona Wulp (Phasiinae: Trichopodini)

Acaulona erythropyga Sabrosky

HOST: HEMIPTERA (HETEROPTERA)

Pyrrhocoridae

Dysdercus andreae (Linnaeus)

Plank, 1951: 19, P.R.; Sabrosky, 1950: 371, P.R.

Dysdercus sanguinarius Stål

Sabrosky, 1950: 371, P.R.

Dysdercus spp.

*Vide Plank, 1951: 19 P.R. (Plank, 1950); Plank, 1950: 15, P.R. (as *Acaulona* n. sp.).

Acaulona peruviana Townsend

HOST: HEMIPTERA (HETEROPTERA)

Pyrrhocoridae

Dysdercus andreae (Linnaeus)

Bartlett, 1943: 15-16, P.R. (lab., preferred host).

Dysdercus sanguinarius Stål

Bartlett, 1943: 15-16, P.R. (lab., not preferred host).

Acaulona n. sp., see *Acaulona erythropygæ**Acemya* Robineau-Desvoidy (Goniinae: Acemyini)*Acemya dentata*, see *Ceracia dentata**Acemya tibialis* Coquillett

HOST: ORTHOPTERA

Acrididae

Melanoplus bivittatus (Say)

Smith, R. W., 1944: 29, Canada.

Melanoplus bruneri Scudder

Newton, 1954: 936, Mont. (as *Acemyia*).

Melanoplus femurrubrum (De Geer)

Smith, R.W., 1944: 29, Canada (ex *M. "f. rubrum"*).

Melanoplus sanguinipes (Fabricius)

Smith, R.W., 1965: 187, 189, Ont. (as *Euacemyia*); Smith, R. W., 1944: 29, Canada (ex *M. mexicanus* Saussure).

Grasshopper(-s)

Anon., 1956: 196, W. Canada (as *Euacemyia*); Smith, R. W., and Finlayson, 1950: 98-100, figs. D, J, 25, 49, Canada (as *Euacemyia*); Smith, R. W., 1939: 60-61, Ont., Sask.

STAGE PARASITIZED: Nymph, adult.

IMMATURES: Second-instar larva — Smith, R. W., and Finlayson, 1950: (descr., figs. D, J, 25, 49).

BIOLOGY: Smith, R. W., and Finlayson, 1950: (98-100).

Acemya spp.

HOST: ORTHOPTERA

Acrididae

Grasshopper(-s)

Anon., 1953a: 128, Man. (as *Euacemyia*).

- Acemyia dentata*, see *Ceracia dentata*
Acemyia tibialis, see *Acemya tibialis*
 "Achaetoneura" *archippivora*, see *Lespesia archippivora*
Achaetoneura, see *Lespesia* species of same name except as noted
Achaetoneura "achippivora", see *Lespesia archippivora*
Achaetoneura "aletia", see *Lespesia aletiae*
Achaetoneura anisotae, see *Lespesia datanarum*
Achaetoneura piperi, see *Lespesia schizurae*
Achaetoneura promiscua, see *Lespesia archippivora*
Achaetoneura samiae, see *Lespesia ciliata*
Achaetoneura similis, see *Tachinomyia similis*
Achaetoneura spectabilis, see *Eufrontina spectabilis*
Achaetoneura tenthredinidarum, see *Bessa harveyi*
Achaetoneura websteri, see *Lespesia archippivora*
 near *Achaetoneura* (n. sp.) (1) or (2), see *Lespesia* spp.
 "Achaetonura" *archippivora*, see *Lespesia archippivora*
 "Achaetopsis" *tortricis*, see *Erynnia tortricis*
Acroglossa Williston (Goniinae: Goniini)
Acroglossa hesperidarum Williston
 Most records as *Spallanzania hesperidarum*.
 HOST: LEPIDOPTERA

Hesperiidae

Epargyreus clarus (Cramer)

- Johnson, 1925a: 88, N.H. (ex *Hesperia tityrus* (Fabricius));
 Johnson, 1910: 783, N.J. (ex *Eudamus tityrus*); *Brauer
 and Bergenstamm, 1895: 542, 591 (as *Acroglossa* ex
Epargyreus tityrus, Williston, 1889); Williston, 1889:
 1914, 1917, N.H. (as *Acroglossa* ex *Epargyreus tityrus*).

Unidentified Lepidoptera

Butterfly(-flies)

Aldrich, 1915: 83.

STAGE PARASITIZED: Pupa.

Actia Robineau-Desvoidy (Goniinae: Siphonini)

Actia americana (Townsend)

HOST: LEPIDOPTERA

Notodontidae

Schizura concinna (J. E. Smith)

- Schaffner and Griswold, 1934: 81, 107, Conn., Maine, Mass.,
 N.H., N.Y., Pa., R.I., Vt. (as *Gymnophthalma*); Webber

and Schaffner, 1926: 19, New England (as *Gymnophthalma*); Johnson, 1925: 186, Conn., Maine, R.I. (as *Thryptocera* ex *S. "concina"*); Walton, 1914a: 91, Conn. (as *Polychaetoneura elyii* Walton).

BIOLOGY: Schaffner and Griswold, 1934: (81, 107).

Actia diffidens Curran

HOST: LEPIDOPTERA

Olethreutidae

Spilonota ocellana (Denis and Schiffermüller)
Stultz, 1955: 80, N.S.

Tortricidae

Acleris variana (Fernald)

Miller, 1966: 603, N.B.; Prebble, 1935: 216-227, figs. 4-19, 24-27, 30-31, N.S. (ex *Peronea*).

Choristoneura conflictana (Walker)

Prentice, 1955: 471, Man., Sask.

STAGE PARASITIZED: Larva.

IMMATURES: First-instar larva — Prebble, 1935: (descr., figs. 4-5, 30).

Second-instar larva — Prebble, 1935: (descr., figs. 6-7, 19, 31).

Third-instar larva — Prebble, 1935: (descr., figs. 8-18).

Pupa — Prebble, 1935: (descr., figs. 24-27).

BIOLOGY: Miller, 1966: (603); Prebble, 1935: (216-227).

Actia flavipes (Coquillett)

All records as *Thryptocera flavipes*.

HOST: LEPIDOPTERA

Diopsideae

Phryganidia californica Packard

Burke and Herbert, 1920: 9-10, Calif.; Burke, 1919: 125, Calif.

Geometridae

Brephos infans Möschler

Schaffner and Griswold, 1934: 90, 107.

Cosymbia pendulinaria (Guenée)

Schaffner and Griswold, 1934: 91, 107 (ex *C. lumenaria* Hübner).

STAGE PARASITIZED: Larva.

BIOLOGY: Schaffner and Griswold, 1934: (107).

Actia interrupta Curran

HOST: LEPIDOPTERA

Geometridae

Melanolophia imitata (Walker)

Evans, 1962: 597, B.C.

Notodontidae

Schizura concinna (J. E. Smith)

Felt, 1912: 100-101, N.Y. (as *A. pilipennis* (Fallén));

Johnson, 1907: 10, R.I. (as *A. pilipennis*).

Olethreutidae

Epinotia crenana (Hübner) probably

Powell, 1962: 134, Calif.

Epinotia emarginana (Walsingham)

Powell, 1962: 134, Calif. (as *A. "interruta"*).

Grapholitha molesta (Busck)

All records ex *Laspeyresia molesta*.

Steenburgh and Boyce, 1938: 73, Ont.; Steenburgh, 1935:

71, Ont.; Haeussler, 1930: 366, N.J. (as *A. pilipennis*).

Tortricidae

Acleris variana (Fernald)

Miller, 1966: 603, N.B.; Miller, 1963: 239, N.B.; Reeks,

Forbes and Cuming, 1948: 12, [maritime prov., Nfld.];

Reeks, Forbes and Cuming, 1947: 12, [maritime prov.]

(as *Gymnophthalma*).

Acleris variegana (Denis and Schiffermüller)

Powell, 1962: 134-135, Calif.

Amorbia cuneana (Walsingham)

Powell, 1962: 134, Calif. (as *A. "interruta"*).

Archippus packardianus (Fernald)

Daviault, 1950a: 150, Que. (ex *Tortrix*); Daviault, 1946:

117, Que. (ex *Tortrix "parckardiana"*).

Archips argyrospilus (Walker)

Prokopy, 1968: 350, Conn.; Powell, 1962: 134, 137, Calif.

Archips rosanus (Linnaeus)

Schaffner, 1959: 31, 48, Mass.

Archips spp. (including *A. argyrospilus* (Walker) and/or *A. griseus* (Robinson))

Prokopy, 1968: 350, Conn.

Choristoneura conflictana (Walker)

Prentice, 1955: 471, Man., Sask.

Choristoneura fumiferana (Clemens)

Blais, 1965: 131-134, Que.; MacDonald and Webb, 1963: 299, N.B.; Miller, 1963: 229, 237, 239, N.B.; Blais, 1960: 388, 390-392, Que. (ex spruce budworm); Coppel, 1960: 94, 97, B.C. (as *Gymnophthalma*); Carolin and Coulter, 1959: 553, Oreg.; McGugan and Blais, 1959: 765, Ont.; Anon., 1953c: 137, B.C.; Ross, 1952: 108, fig. 7, Canada; Daviault, 1950: 43-45, Que.; Dowden and Carolin, 1950: 778-779, N.Y. (as *A. "interruptor"* ex *Archips*); Wilkes, Coppel and Mathers, 1949: 140, 146, 149, 151, B.C. (as *Gymnophthalma*); Dowden, Buchanan and Carolin, 1948: 458, 461 (ex *Archips*); Reeks, Forbes and Cuming, 1948: 16, [maritime prov., Nfld.]; Daviault, 1946: 117, Que. (ex *Archips*); Bedard, 1938: 191 (as *A. pilipennis* ex *Cacoecia*).

Choristoneura pinus Freeman

Allen, D. C., Knight, Foltz and Mattson, 1969: 1471, 1475, Mich.; Dixon and Benjamin, 1963: 267, Wis.; Benjamin and Drooz, 1954: 589-590, Mich.

Choristoneura rosaceana (Harris)

All records ex *Archips rosaceana*.

Neunzig and Gyrisco, 1955: 450, N.Y.; Schuh and Mote, 1948: 26, 28, Oreg.

Croesia albicomana (Clemens)

Powell, 1962: 134, 136, Calif.

Phalonia oenotherana Riley

Dickerson and Weiss, 1920: 44, N.J. (as *A. pilipennis*).

Xenotemna pallorana (Robinson)

All records ex *Tortrix pallorana*.

Putman, 1935: 48, Ont.; Putman, 1935a: 108, Ont.

Unidentified Tortricidae

Aldrich, 1932: 2, Mass. (as *A. pilipennis* ex larva on *Baptisia tinctoria* (Linnaeus) Ventenat).

STAGE PARASITIZED: Larva, pupa.

IMMATURES: Pupa — Ross, 1952: (fig. 7); Greene, 1922: (descr., fig. 38 on pl. 8).

BIOLOGY: Miller, 1963: (229, 237, 239); Evans, 1962: (597);
Schaffner, 1959: (31, 48); Daviault, 1950a: (150).

Actia sp. near *interrupta* Curran

All records as *Actia* sp. near *pilipennis* (Fallén).

HOST: LEPIDOPTERA

Tortricidae

Archips cerasivoranus (Fitch)

Schaffner, 1959: 30, 48, [N.E. U.S.].

Archips fervidanus (Clemens)

Schaffner, 1959: 29, 48, [N.E. U.S.].

BIOLOGY: Schaffner, 1959: (29-30, 48).

Actia ontario Curran

HOST: LEPIDOPTERA

Geometridae

Ennomos subsignarius (Hübner)

Raizenne, 1952: 151, Ont.

Actia palloris (Coquillett)

All records as *Xanthoactia palloris*.

HOST: LEPIDOPTERA

Drepanidae

Drepana arcuata Walker

Raizenne, 1952: 105, Ont.

Drepana bilineata (Packard)

Raizenne, 1952: 106, Que.; Raizenne, 1952: 107, Que. (ex
D. bilineata form *levis* Hudson).

Actia sp. near *palloris* (Coquillett)

HOST: LEPIDOPTERA

Geometridae

Ennomos subsignarius (Hübner)

Schaffner and Griswold, 1934: 99-100, 107, Mass.

Eugonobapta nivosaria (Guenée)

Schaffner and Griswold, 1934: 99, 107, Mass.

BIOLOGY: Schaffner and Griswold, 1934: (107).

Actia pilipennis, see *Actia interrupta*

Actia sp. near *pilipennis*, see *Actia* sp. near *interrupta*

Actia tibialis, see *Strobliomyia tibialis*

Actia spp.

HOST: LEPIDOPTERA

Diptidae

Phryganidia californica Packard

Harville, 1955: 115-116, 129, 132-133, 143, 146, Calif.

Gelechiidae

Gnorimoschema spp.

Schaffner, 1959: 37, 49, Mass.

Noctuidae

Catocala relictata Walker

Schaffner and Griswold, 1934: 65, 107, Mass.

Unidentified *Catocalinae*

Schaffner and Griswold, 1934: 67, 107.

Notodontidae

Heterocampa guttivitta (Walker)

Collins, 1926: 693, 695, N.H.

Olethreutidae

Epinotia similana (Hübner)

Schaffner, 1959: 25, 48, Mass.

Episimus argutanus (Clemens)

Schaffner, 1959: 20, 48, Mass., N.H.

Tortricidae

Choristoneura pinus Freeman

Dixon and Benjamin, 1963: 267, Wis.

STAGE PARASITIZED: Larva.

BIOLOGY: Schaffner, 1959: (20, 25, 37, 48-49); Harville, 1955: (115-116, 129, 132-133, 143, 146); Schaffner and Griswold, 1934: (107).

Admontia Brauer and Bergenstamm (Goniinae: *Admontiini*)

Admontia "degeerioides," see *Admontia degeerioides*

Admontia degeerioides (Coquillett)

HOST: DIPTERA

Tipulidae

Tipula abdominalis (Say)

Fattig, 1949: 5-6, Ga.

HOST: LEPIDOPTERA

Noctuidae

Spodoptera frugiperda (J. E. Smith)Luginbill, 1928: 82, Tex. (ex *Laphygma*).

Olethreutidae

Grapholitha molesta (Busck)Muesebeck, 1936: 536, Ohio (as *A. "degeeriodes"*).

STAGE PARASITIZED: Larva (Diptera, Lepidoptera).

Admontia demylus, see *Spathimeigenia spinigera**Admontia hylotomae*, see *Spathimeigenia hylotomae**Admontia pergandei* Coquillett

HOST: DIPTERA

Tipulidae

Tipula abdominalis (Say)Boardman, 1939: 160 (as *Euadmontia*); Johannsen, 1935: 37, N.Y. (as *Euadmontia*).*Tipula cunctans* (Say)Most records ex *Tipula infuscata* Loew.Alexander, 1920: 731, Ill. (ex *T. cunctans*); Townsend, 1912b: 165, Tenn.; Hyslop, 1910: 128.

STAGE PARASITIZED: Larva.

IMMATURES: Third instar larva — Johannsen, 1935: (descr.).

BIOLOGY: Boardman, 1939: (160); Johannsen, 1935: (37); Townsend, 1912b: (165).

Admontia retiniae, see *Urophylloopsis retiniae**Admontia* spp.

HOST: DIPTERA

Tipulidae

Ctenophora spp.Alexander, 1920: 732 (as "*Trichoparia*")."Adomonita" *demylus*, see *Spathimeigenia spinigera**Adoryphorophaga* Townsend (Goniinae: Blondeliini)

Adoryphorophaga aberrans (Townsend)Most records as *Doryphorophaga aberrans*.

HOST: COLEOPTERA

Chrysomelidae

Blepharida rhois (Forster)

Townsend, 1916: 217, Va.

Leptinotarsa decemlineata (Say)

Anon., 1958c: 284, Ont. (as *Adoryphorophaga*); Briand, 1958: 57, Ont. (as *Adoryphorophaga*); Anon., 1957c: 273, Ont. (as *Adoryphorophaga*); Bruneteau, 1937: 131-135, figs. 9-14, N.Y. (ex *Doryphora*); Feytaud, 1937: 64-65; Feytaud, 1936: 26-27 (ex *Doryphora*); Smith, H. E., 1918: 124, Conn., Mass. (ex *Doryphora*); Townsend, 1916: 217, Va.

Chrysomelid spp. (including *Lema trilineata* (Olivier) and/or *Leptinotarsa decemlineata* (Say))

All records ex potato beetles.

*Procter, 1946: 431, Maine (Procter, 1938); Procter, 1938: 373, Maine; Johnson, 1927: 201, Maine; Johnson, 1925: 199, Conn., Maine, Mass., Vt.

STAGE PARASITIZED: Larva, adult.

IMMATURES: Egg — Bruneteau, 1937: (fig. 9).

First instar larva — Bruneteau, 1937: (descr., figs. 10-11).

Third instar larva — Bruneteau, 1937: (descr., figs. 11-13).

Pupa — Bruneteau, 1937: (fig. 14).

BIOLOGY: Anon., 1958c: (284); Briand, 1958: (57); Bruneteau, 1937: (131-135).

Adoryphorophaga spp. and/or **Doryphorophaga** spp.

HOST: COLEOPTERA

Chrysomelidae

Leptinotarsa decemlineata (Say)

Walsh and Riley, 1868: 46, Ill. (as *Tachina* fly [LeBaron places in *Lydella*] ex *Doryphora 10-lineata*).

STAGE PARASITIZED: Larva.

BIOLOGY: Walsh and Riley, 1868: (46).

“*Agrophyla*” sp., see *Argyrophylax* spp.*Alophora opaca*, see *Alophorella opaca**Alophora pulverea*, see *Alophorella pulverea**Alophorella* Townsend (Phasiinae: Phasiini)*Alophorella* “*aeneiventris*,” see *Alophorella aeneoventris*

Alophorella aeneoventris (Williston)
HOST: HEMIPTERA (HETEROPTERA)

Miridae

Lygus hesperus Knight probably
Clancy and Pierce, 1966: 855, Calif.

Lygus lineolaris (Palisot de Beauvois)
Clancy and Pierce, 1966: 855, 857, Ark., Mo., N.J.; Medler,
1961: 102, Wis. (as *A. "aeneiventris"*).

Pentatomidae

Cosmopepla bimaculata (Thomas)
Clancy and Pierce, 1966: 857, N.Y.
HOST: HEMIPTERA (HOMOPTERA)

Cercopidae

Philaenus spumarius (Linnaeus)
Clancy and Pierce, 1966: 857, Pa.
STAGE PARASITIZED: Adult.
BIOLOGY: Medler, 1961: (102).
Alophorella fumosa (Coquillett)
HOST: HEMIPTERA (HETEROPTERA)

Miridae

Lygocoris caryae (Knight)
Brooks, 1945a: 666, N.Y. (as *Oedematopteryx ex Lygus*).
Lygus lineolaris (Palisot de Beauvois)
Clancy and Pierce, 1966: 855, 857, N.J.
STAGE PARASITIZED: Adult.
Alophorella opaca (Coquillett)
HOST: HEMIPTERA (HETEROPTERA)

Miridae

Lygus pratensis (Linnaeus)
Painter, 1929: 105-106, Ont. (as *Alophora*).
STAGE PARASITIZED: Nymph, adult.
BIOLOGY: Painter, 1929: (105-106).
Alophorella pulverea (Coquillett)
HOST: HEMIPTERA (HETEROPTERA)

Miridae

Lygocoris caryae (Knight) probably

Clancy and Pierce, 1966: 855, N.Y. (ex *Lygus*).

Lygus lineolaris (Palisot de Beauvois)

Clancy and Pierce, 1966: 855, Ark.; Medler, 1961: 101-102,
Wis. (as *Oedematopteryx*).

Pentatomidae

Holcostethus limbolarius (Stål)

Clancy and Pierce, 1966: 857.

STAGE PARASITIZED: Adult.

IMMATURES: Pupa — Greene, 1922: (as *Alophora*; descr., fig.
45 on pl. 10).

BIOLOGY: Medler, 1961: (101-102).

Alophorella spp.

HOST: HEMIPTERA (HETEROPTERA)

Miridae

Lygus lineolaris (Palisot de Beauvois)

Clancy and Pierce, 1966: 855, Ont.; Medler, 1961: 102, Wis.;
Medler, 1961: 102, Wis. (as *Oedematopteryx*).

STAGE PARASITIZED: Adult.

Alophorellopsis purpurascens, see *Hyalomya purpurascens*

Amedoria luctuosa, see *Medina barbata*

Amedoria spinosa, see *Medina barbata*

"*Anachaetopsis tortricis*," see *Erynnia tortricis*

Anachaetopsis tortricis, see *Erynnia tortricis*

Anachaetopsis vagans, see *Erynnia tortricis*

Anachaetopsis sp., see *Erynnia* spp.

Anaporia Townsend (Dexiinae: Uramyini)

Anaporia limacodis (Townsend)

HOST: LEPIDOPTERA

Limacodidae

Limacodes spp.

*Brauer and Bergenstamm, 1895: 543, 597, N. Amer. (as
Aporia sp., Townsend, 1892c); Townsend, 1892c: 275,
N.Y. (as *Aporia*).

Lithacodes fasciola (Herrich-Schaeffer)

Schaffner, 1959: 7, 49, Conn.

Sisyrosea textula Herrich-Schaeffer

Schaffner, 1959: 5, 49, Conn.

Tortricidia flexuosa (Grote)Schaffner, 1959: 7, 49, Conn.; Johnson, 1925: 189, Conn., Mass., N.H. (as *Pseudeuantha coquilletti* Aldrich); Aldrich, 1921: 91, N.Y. (as *Pseudeuantha coquilletti*).

BIOLOGY: Schaffner, 1959: (5, 7, 49).

Anaporia pristis (Walker)

HOST: LEPIDOPTERA

Arctiidae

Halisidota argentata PackardCoquillett, 1897: 18, 25, Oreg. (as *Macquartia*).

Limacodidae

Euclea cippus CramerJohnson, 1925: 189, Conn., Maine, Mass., N.H., Vt. (as *Pseudeuantha* ex *E. "ceppus"*); *vide Aldrich, 1921: 90, Va. (as *Pseudeuantha*, Coquillett, 1897); Coquillett, 1897: 13, 25, Va. (as *Exorista blanda* (Osten Sacken)).*Phobetron pithecium* (J. E. Smith)

Schaffner, 1959: 6, 49, [N.E. U.S.].

Sibine stimulea (Clemens)All records ex *Empretia stimulea*.Johnson, 1925: 189, Conn., Maine, Mass., N.H., Vt. (as *Pseudeuantha*); Aldrich, 1921: 90, N.Y. (as *Pseudeuantha*).*Sisyrosea textula* Herrich-SchaefferCoquillett, 1897: 14, 25 (as *Exorista isae* Coquillett ex *Isa inornata* (Grote and Robinson)).

Megalopygidae

Megalopyge crispata (Packard)Schaffner, 1959: 7, 49, [N.E. U.S.] (ex *Lagoa*).

STAGE PARASITIZED: Larva.

BIOLOGY: Schaffner, 1959: (6-7, 49).

Anaporia sp.

HOST: LEPIDOPTERA

Limaconidae

Sisyrosea textula Herrich-Schaeffer

Schaffner, 1959: 5, 49, Conn.

BIOLOGY: Schaffner, 1959: (5, 49).

Andrina radialis, see *Lydella radialis*

Androeryops Beneway (Phasiinae: Phasiini)

Androeryops ecitonis (Townsend)

HOST: HYMENOPTERA

Formicidae

Eciton burchelli foreli Mayr probably

Townsend, 1897: 30, 32, Mexico (V.C.) (as *Hyalomyia* ex *E. foreli*, suggested host).

BIOLOGY: Townsend, 1897: (30, 32).

Anetia armigera, see *Eucelatoria armigera*

Anetia dimmocki, see *Eucelatoriopsis dimmocki*

Anetia eufitchiae, see *Blondelia eufitchiae*

Anetia hyphantriae, see *Blondelia hyphantriae*

Anetia sp. complex, see *Blondelia* spp.

Anisia Wulp (Goniinae: Blondeliini)

"*Anisia nigrocincta* Wulp" probably

HOST: LEPIDOPTERA

Pyralidae

Ostrinia penitalis (Grote)

Brauer and Bergenstamm, 1895: 543, 585, N. Amer. (ex *Botys*).

Anisia sp. near *variabilis*, see *Lixophaga* sp. near *variabilis*

Aphria Robineau-Desvoidy (Tachininae: Aphriini)

Aphria "cypterata," see *Aphria ocypterata*

Aphria ocypterata Townsend

HOST: LEPIDOPTERA

Noctuidae

Euxoa messoria (Harris)

*Crumb, 1929: 40, 94 (Coquillett, 1897); Essig, 1926: 580, Wash.; Johnson, 1925: 194, Maine, Mass., N.H. (ex *Carneades*); Coquillett, 1897: 10, 24, Wash. (ex *Carneades*).

Pyralidae

Nephopteryx subfuscella (Ragonot)

Schaffner, 1959: 17, 49, Maine (ex *Salebria semiobscura* Hulst).

Trascula reductella (Walker)

Raizenne, 1952: 174, Ont. (as *A. "cypterata"*).

BIOLOGY: Schaffner, 1959: (17, 49).

Aplomya Robineau-Desvoidy (Goniinae: Eryciini)

Aplomya caesar (Aldrich)

HOST: LEPIDOPTERA

Arctiidae

Diacrisia virginica (Fabricius)

Fattig, 1949: 6.

Gracillariidae

Gracillaria negundella Chambers

All records ex boxelder leaf roller.

Knowlton, Janes and Thomas, 1933: 155, Utah (as *Zenillia*); Knowlton and Janes, 1931: 147, Utah (as *Zenillia*).

Lymantriidae

Nygmia phaeorrhoea (Donovan)

Wishart, 1945: 157-167, figs. 2-8, 13 on pl. 7; Sellers, 1943: 89.

Noctuidae

Papaipema nebris (Guenée)

All records ex *Papaipema nitela* (Guenée).

Wishart, 1945: 157-167, figs. 2-8, 13 on pl. 7; Sellers, 1943: 89.

Olethreutidae

Laspeyresia nigricana (Stephens)

Anon., 1947: 180, 195, 200, Ont.

Pyralidae

Crambus mutabilis Clemens

Ainslie, G. G., 1927: 14-15, Ind. (as *Exorista nigripalpis* Townsend).

Crambus trisectus Walker

Ainslie, G. G., 1927: 14-15, Ind. (as *Exorista nigripalpis*); Aldrich and Webber, 1924: 28, Ind. (as *Zenillia*); Ainslie, G. G., 1923: 412, Ind. (as *Exorista nigripalpis*).

Crambus spp.

Noble, 1932: 3, U.S. (as *Zenillia*); Johnson, 1925: 196, Maine, Mass., N.H., Vt. (as *Zenillia*).

Dioryctria reniculella (Grote)

McLeod, J.M., and Daviault, 1963: 316, Que. (as *Eumea*).

Loxostege commixtalis (Walker)

Wishart, 1945: 157-167, figs. 2-8, 13 on pl. 7; Sellers, 1943: 89, Colo.; Hoerner, 1933: 12 (as *Zenillia*).

Loxostege sticticalis (Linnaeus)

Wishart, 1945: 157-167, figs. 2-8, 13 on pl. 7; Sellers, 1943: 89, Colo.; Essig, 1926: 581, Colo. (as *Zenillia*); Aldrich and Webber, 1924: 28, Colo. (as *Zenillia*).

Ostrinia nubilalis (Hübner)

Most records ex *Pyrausta nubilalis*.

Anon., 1963d: 1029, Wis. (ex *Ostrinia*); Hudon, 1960: 125-126, Que. (as *Eumea* ex *Ostrinia*); Pond, 1955: 106, N.B. (as *Eumea*); Blickenstaff, Arbuthnot and Harris, 1953: 368-370, Iowa; Boyes and Wilkes, 1953: 126, 138, Ont.; Hawkins and Devitt, 1953: 24, Maine (ex European corn borer); Arbuthnot and Wright, 1952: 233, Ill.; Arbuthnot, 1950: 45-46, N. Cent. states (ex European corn borer); Arbuthnot, Jones, Carter and Evans, 1949: 7-8, 12-13, 18-22, 24, Iowa, Maine, Mass., N.H., N.J., N.Y., Wis.; Baker, W. A., Bradley and Clark, 1949: 3, 174-175, fig. 38Db, Mich., Ohio (as *Aplomyia*); Arbuthnot, Jones, Carter and Evans, 1948: 3, 9-10, 13, Minn., Ohio, S. New England; Anon., 1947: 180, 192, 198, Ont.; Arbuthnot, 1947: 30, Ind. (as *Aplomyia* ex European corn borer); Deay, 1947: 68, Ind. (as "*Aplomyea*" ex European corn borer); Drake, 1947: 129, Iowa (ex European corn borer); Drake, 1946: 176, Iowa (ex European corn borer); Wishart, 1945: 157-167, figs. 2-8, 13 on pl. 7; Sellers, 1943: 89; Bradley, W.G., and Clark, 1941: 788, 792, Mich., Ohio (ex European corn

borer); Bradley, W.G., and Clark, 1940: 527, 531, Mich., N.Y., Ohio (ex European corn borer); Bradley, W.G., and Clark, 1939: 528, 534, Ind., Mich., Ohio (as *Zenillia* ex European corn borer); Baker, W.A., and Bradley, 1938: 345-346, 348, Mich., Ohio (as *Zenillia*); Bradley, W.G., and Clark, 1938: 650, 654, Mich., Ohio (as *Zenillia* ex European corn borer); Baird, 1937: 66, Canada (as *Zenillia*); Baird, 1929: 38-39, Ont. (as *Zenillia* ex [European] corn borer); Baird, 1926: 78, Ont. (as *Zenillia* ex European corn borer); Huber and Neiswander, 1924: 127, Ohio (as *Exorista nigripalpis*); Spencer, 1923: 19, Ont. (as *Exorista nigripalpis* ex European corn borer); Crawford and Spencer, 1922: 235, Ont. (as *Exorista nigripalpis* ex European corn borer); Caffrey, 1919: 21, Mass. (as *Exorista nigripalpis*); Vinal and Caffrey, 1919: 57-58, Mass. (as *Exorista nigripalpis*).

Ostrinia obliteralis (Walker)

All records ex *Pyrausta ainsliei* Heinrich.

Blickenstaff, Arbuthnot and Harris, 1953: 368, 370, Iowa;
Wishart, 1945: 157-167, figs. 2-8, 13 on pl. 7; Sellers,
1943: 89.

Ostrinia penitalis (Grote)

All records ex *Pyrausta penitalis*.

Wishart, 1945: 157-167, figs. 2-8, 13 on pl. 7; Sellers,
1943: 89.

Phlyctaenia coronata tertialis (Guenée)

All records ex *Phlyctaenia tertialis*.

Wishart, 1945: 157-167, figs. 2-8, 13 on pl. 7; Balduf,
1929: 238, 240, Ill., Ohio (as *Zenillia*).

Pyrausta spp.

Johnson, 1925: 196, Maine, Mass., N.H., Vt. (as *Zenillia*).

Tortricidae

Amorbia humerosana Clemens

Wishart, 1945: 157-167, figs. 2-8, 13 on pl. 7.

Archips argyrospilus (Walker)

Paradis and LeRoux, 1965: 44, 46-47, fig. 14D, Que. (as *Eumea*); Powell, 1962: 137, Calif.; Paradis, 1961: 63, Que. (as *Eumea*); Newcomer, 1958: 88, Wash.; Wishart, 1945: 157-167, figs. 2-8, 13 on pl. 7; Sellers, 1943: 89, Colo., Mont., Ont., Wash.; Harman, 1928: 9, N.Y. (as *Zenillia*); Essig, 1926: 581, Colo., N. Mex., Wash. (as

Zenillia); Aldrich and Webber, 1924: 28-29, Colo., Ont., Wash. (as *Zenillia ex Cacoecia*); Gillette, 1922: 15, Colo. (as *Exorista nigripalpis*); Aldrich, 1916: 21, Ont. (as *Exorista*); Gill, 1913: 102, Colo. or N. Mex. (as *Exorista nigripalpis*).

Archips purpuranus (Clemens)

Schaffner, 1959: 31, 49-50, [N.E. U.S.]; Wishart, 1945: 157-167, figs. 2-8, 13 on pl. 7; Sellers, 1943: 89.

Archips rosanus (Linnaeus)

Schaffner, 1959: 31, 49-50, [N.E. U.S.]; Putman, 1943: 222, Ont. (as *Zenillia*).

Archips semiferanus (Walker)

Schaffner, 1959: 30, 49-50, [N.E. U.S.]; Wishart, 1945: 157-167, figs. 2-8, 13 on pl. 7; Sellers, 1943: 89.

Archips spp.

Johnson, 1925: 196, Maine, Mass., N.H., Vt. (as *Zenillia ex Cacoecia*).

Argyrotaenia citrana (Fernald)

Basinger, 1935: 233, Calif. (as "*Zenilla*").

Argyrotaenia quadrifasciana (Fernald)

Wishart, 1945: 157-167, figs. 2-8, 13 on pl. 7 (ex. *Eulia*).

Choristoneura conflictana (Walker)

Prentice, 1955: 471, Man., Sask. (as *Eumea*).

Choristoneura fractivittana (Clemens)

Wishart, 1945: 157-167, figs. 2-8, 13 on pl. 7 (ex *Archips* "*practivittana*").

Choristoneura fumiferana (Clemens)

Blais, 1965: 131-134, Que. (as *Eumea*); MacDonald and Webb, 1963: 299, N.B.; Miller, 1963: 237, 239, N.B. (as *Eumea*); Blais, 1960: 388, 390-391, Que. (as *Eumea* ex spruce budworm); Coppel, 1960: 94, 97, B.C. (as *Eumea*); Carolin and Coulter, 1959: 553-554, Oreg.; McGugan and Blais, 1959: 765, Ont. (as *Eumea*); Boyes and Wilkes, 1953: 126, 140; Coppel, 1953: 191, B.C. (as *Zenillia* ex "*Chloristoneura*"); Dowden, Jaynes and Carolin, 1953: 309, Maine; Anon., 1952g: 289, 293, 295, B.C. (as *Zenillia*); Jaynes and Drooz, 1952: 1058, 1060, Maine, N.Y.; Raizenne, 1952: 219, Ont. (as *Epimasicera*); Ross, 1952: 112, fig. 11, Canada; Anon., 1950b: 210, maritime prov. (as *Epimasicera* ex spruce budworm); Anon., 1950c: 241, 246, 249, B.C. (as *Zenillia*); Daviault, 1950: 42-45, Que. (as *Aplomyia*); Dowden and Carolin, 1950: 778-781, N.Y. (ex *Archips*); Anon., 1949: 276, 283, 286, B.C. (as *Zenillia*); Wilkes,

Coppel and Mathers, 1949: 140, 146, 149, 151, B.C.; Dowden, Buchanan and Carolin, 1948: 458-461 (as *Aplomyia* ex *Archips*); Wilkes and Anderson, 1947: 41, 43, Ont., Que. (as *Zenillia* ex *Archips*); Reeks, 1946: 25, N.B., Nfld. (as *Epimasicera* ex *Archips*); Reeks, Smith and Forbes, 1946: 12, N.B. (as *Epimasicera* ex *Archips*); Wishart, 1945: 157-167, figs. 2-8, 13 on pl. 7 (ex *Archips*); Sellers, 1943: 89 (ex *Archips*); Essig, 1926: 581, B.C. (as *Zenillia* ex *Harmologa*); Tothill, 1913: 71, B.C., Que. (as *Exorista nigripalpis* ex *Tortrix*).

Choristoneura parallela (Robinson)

All records ex *Archips parallela*.

Wishart, 1945: 157-167, figs. 2-8, 13 on pl. 7; Sellers, 1943: 89.

Choristoneura pinus Freeman

Allen, D.C., Knight, Foltz and Mattson, 1969: 1471-1472, 1474-1475, Mich.; Dixon and Benjamin, 1963: 267-270, Wis.; Kulman and Hodson, 1961: 222, Minn.; Drooz and Benjamin, 1956: 412-413, Mich.

Choristoneura rosaceana (Harris)

All records ex *Archips rosaceana*.

Raizenne, 1952: 215, Ont. (as *Epimasicera*); Wishart, 1945: 157-167, figs. 2-8, 13 on pl. 7.

Pandemis limitata (Robinson)

Wishart, 1945: 157-167, figs. 2-8, 13 on pl. 7 (ex *P. "limitana"*).

Phaecasiophora niveiguttana Grote

Montgomery, 1933: 188, Ind. (as *Zenillia*).

Tortricid spp. (including *Amorbia humerosana* Clemens, *Archips argyrospilus* (Walker), *A. semiferanus* (Walker), *Argyrotaenia quadrifasciana* (Fernald), *Choristoneura fractivittana* (Clemens), *C. rosaceana* (Harris) and/or *Pandemis limitata* (Robinson))

Hall, 1929: 139, Ont. (as *Zenillia* ex *Amorbia humerosana*, *Archips argyrospila*, *A. fractivittana*, *A. rosaceana*, *A. semiferana*, *Eulia quadrifasciana* and/or *P. limitata*).

Tortricid spp. (including *Archips* spp. and/or *Choristoneura* spp.)

Caesar, 1916a: 172-173, unnumbered fig. on p. 172, Ont. (as *Exorista* ex *Tortrix* leaf rollers attacking apples).

STAGE PARASITIZED: Larva, pupa.

IMMATURES: Egg — Wishart, 1945: (descr., fig. 2 on pl. 7).

First instar larva — Wishart, 1945: (descr., figs. 3-4 on pl. 7).

Second instar larva — Wishart, 1945: (descr., fig. 5 on pl. 7).

Third instar larva — Wishart, 1945: (descr., fig. 6 on pl. 7).

Pupa — Paradis and LeRoux, 1965: (fig. 14D); Ross, 1952: (fig. 11); Baker, W. A., Bradley and Clark, 1949: (fig. 38Db); Wishart, 1945: (descr., figs. 7-8, 13 on pl. 7); Greene, 1922: (descr., fig. 83 on pl. 18); Caesar, 1916a: (unnumbered fig. on p. 172).

BIOLOGY: Paradis and LeRoux, 1965: (44, 46-47); Miller, 1963: (237, 239); Hudon, 1960: (125-126); Schaffner, 1959: (30-31, 49-50); Baker, W.A., Bradley and Clark, 1949: (3, 174-175); Wishart, 1945: (157-167); Sellers, 1943: (89); Noble, 1932: (3); Balduf, 1929: (238, 240); Ainslie, G.G., 1927: (14-15).

Aplomya caesar (Aldrich) probably

All records as *Carcelia "nigropalpus"* (Townsend), error for *nigripalpis* (Townsend)

HOST: LEPIDOPTERA

Tortricidae

Archips argyrospilus (Walker)

Herrick and Leiby, 1915: 260, N. Y.; Herrick, 1912: 286, N.Y.

STAGE PARASITIZED: Larva.

Aplomya cerurae Sellers

HOST: LEPIDOPTERA

Notodontidae

Cerura cinerea (Walker)

Vide Sellers, 1943: 95, Mass. (Schaffner and Griswold, 1934); Schaffner and Griswold, 1934: 82-83, 113, Mass. (as *Zenillia* n. sp. (3)).

BIOLOGY: Schaffner and Griswold, 1934: (82-83, 113).

Aplomya confusionis Sellers

HOST: LEPIDOPTERA

Pyralidae

Crambus inornatellus Walker

Bohart, 1947: 290, 292, Calif. (as *Aplomyia* ex *C. bonifatellus* Hulst).

Crambus mutabilis Clemens

*Vide Sellers, 1943: 87 (Ainslie, G. G., 1923); Ainslie, G. G., 1923: 411-412, Ind. (as *Exorista nigripalpis* Townsend).

Crambus sperryellus Klots

Bohart, 1947: 290, 292, Calif. (as *Aplomyia*).

Crambus trisectus Walker

Sellers, 1943: 87.

STAGE PARASITIZED: Larva.

IMMATURES: Pupa—Bohart, 1947: (descr.).

Aplomya crassiseta (Aldrich and Webber)

HOST: LEPIDOPTERA

Tortricidae

Choristoneura conflictana (Walker)

Prentice, 1955: 472, Alta.

Aplomya epicydes (Walker)

HOST: HYMENOPTERA

Cimbicidae

Cimbex americana Leach

Raizenne, 1957: 10, Ont.; Sellers, 1943: 93; Johnson, 1925: 196, Mass. (as *Zenillia coerulea* Aldrich and Webber); Aldrich and Webber, 1924: 23, Mass. (as *Zenillia coerulea*).

BIOLOGY: Sellers, 1943: (93).

Aplomya estigmenensis, see *Hubneria estigmenensis*

Aplomya fronto (Coquillett)

HOST: HYMENOPTERA

Pamphiliidae

Acantholyda spp.

Eidt, 1969: 54, Canada.

Neurotoma spp.

Eidt, 1969: 54, Canada.

STAGE PARASITIZED: Larva.

Aplomya helvina (Coquillett)

HOST: LEPIDOPTERA

Arctiidae

Ecpantheria deflorata (Fabricius)

Fattig, 1949: 6, Ga.

Geometridae

Biston cognataria (Guenée)

Sellers, 1943: 92 (ex *Lycia*); Schaffner and Griswold, 1934: 96, 112 (as *Zenillia* ex *Amphidasis*); Johnson, 1925: 196, Maine, Mass., N.H. (as *Zenillia* ex "Amphidosia"); Tohill, 1913: 71, B. C. (as *Exorista* ex *Lycia*).

Phaeoura quernaria (J. E. Smith)

All records ex *Nacophora quernaria*.

Sellers, 1943: 92; Schaffner and Griswold, 1934: 96, 112 (as *Zenillia*).

Unidentified Geometridae

Sellers, 1943: 92; Schaffner and Griswold, 1934: 102, 112 (as *Zenillia*).

STAGE PARASITIZED: Larva.

BIOLOGY: Sellers, 1943: (92); Schaffner and Griswold, 1934: (112).

Aplomya mitis (Meigen)

HOST: LEPIDOPTERA

Pyralidae

Ostrinia nubilalis (Hübner)

Baker, W. A., Bradley and Clark, 1949: 3, 7, 37, 53, 149-153, fig. 38Da (as *Aplomya* ex *Pyrausta*).

STAGE PARASITIZED: Larva.

IMMATURES: Pupa — Baker, W. A., Bradley and Clark, 1949: (fig. 38Da).

BIOLOGY: Baker, W. A., Bradley and Clark, 1949: (3, 7, 37, 53, 149-153).

Aplomya neurotomae Sellers

HOST: HYMENOPTERA

Pamphiliidae

Neurotoma fasciata (Norton)

Sellers, 1943: 86, Mass., N.H., N.J., Pa.

BIOLOGY: Sellers, 1943: (86).

Aplomya pheosiae Sellers

HOST: LEPIDOPTERA

Noctuidae

Alypia octomaculata (Fabricius)

Sellers, 1943: 96.

Notodontidae

Pheosia rimosa Packard

Vide Sellers, 1943: 95-96, Maine (ex *P. dimidiata* (Herrich-Schaeffer), Schaffner and Griswold, 1934); Schaffner and Griswold, 1934: 77, 114, Maine (as *Zenillia* n. sp. (5)).

BIOLOGY: Sellers, 1943: (95-96); Schaffner and Griswold, 1934: (77, 114).

Aplomya theclarum (Scudder)

HOST: LEPIDOPTERA

Lasiocampidae

Gloveria howardi Dyar

*Sellers, 1943: 78 (Coquillett, 1897; *Aldrich and Webber, 1924: 36 (as *Zenillia confinis* (Fallén), Coquillett, 1897); Coquillett, 1897: 13, 25, Ariz. (as *Exorista confinis* ex *Dendrobinus*).

Lycaenidae

Brephidium exilis (Boisduval)

Sellers, 1943: 78; Essig, 1926: 581, N. Mex. (as *Zenillia confinis*); *Aldrich and Webber, 1924: 36, N. Mex. (as *Zenillia confinis* ex "Bryshidium exile," Coquillett, 1897); Coquillett, 1897: 13, 26, N. Mex. (as *Exorista confinis* ex *Lycaena*).

Celastrina argiolus pseudargiolus (Boisduval and LeConte)

*Sellers, 1943: 78 (ex *Lycaenopsis pseudargiolus*, Williston, 1889); Clark, 1936a: 689 (as *Zenillia confinis* ex *Lycaenopsis*); Johnson, 1910: 780, N.J. (as *Exorista confinis* ex *Lycaena pseudargiolus*); *Brauer and Bergenstamm, 1895: 554, 590, N. Amer. (as *Exorista* ex *Cyaniris pseudargiolus*, Williston, 1889); Williston, 1889: 1914, 1920 (as *Exorista* ex *Cyaniris pseudargiolus*).

Lycaeides melissa (Edwards)

Most records ex *Plebeius melissa*.

Sellers, 1943: 78; Essig, 1926: 581, Colo. (as *Zenillia confinis*); Aldrich and Webber, 1924: 36, Colo. (as *Zenillia confinis* ex *Lycaena*).

Lycaena thoe Guérin-Ménéville

Sellers, 1943: 78; Aldrich and Webber, 1924: 36, Ont. (as *Zenillia confinis* ex *Heodes*).

Lycaena xanthoides (Boisduval)

Sellers, 1943: 78.

Lycaena xanthoides dione ScudderSellers, 1943: 78 (ex *L. dione*); Brauer and Bergenstamm, 1895: 574, 587, Iowa (as *Sisyropa "chrysophanae"* (Townsend) ex *Chrysophanus dione*); Townsend, 1891e: 197-198, Iowa (as *Exorista chrysophani* ex *Chrysophanus dione*).*Lycaena* spp.Sellers, 1943: 78; Coquillett, 1897: 13, 26, D.C. (as *Exorista confinis*).*Strymon calanus* HübnerMost records ex *Thecla calanus*.Sellers, 1943: 78 (ex *Strymon*); Aldrich and Webber, 1924: 36 (as *Zenillia confinis*); Johnson, 1910: 780, N.J. (as *Exorista confinis*); *Brauer and Bergenstamm, 1895: 554, 608, N. Amer. (as *Exorista*, Williston, 1889); Williston, 1889: 1914, 1920 (as *Exorista*).*Strymon falacer* (Godart)Raizenne, 1952: 6, Ont., Que.; Sellers, 1943: 78; *Brauer and Bergenstamm, 1895: 577, 608 (as *Tachina* ex *Thecla "inornata"* (Grote and Robinson), Scudder in Osten Sacken, 1887); Scudder in Osten Sacken, 1887: 166 (as *Tachina* ex *Thecla inornata*).*Strymon melinus* (Hübner)Fattig, 1949: 7; Sellers, 1943: 78; Little and Martin, 1941: 74, Tex. (as "*Zanillia*" *confinis*); Allen, 1929: 686, Miss. (as *Zenillia confinis* ex *Uranotes*); Reinhard, 1929: 33-34, Tex. (as *Zenillia confinis*); Aldrich and Webber, 1924: 36, Tex. (as *Zenillia confinis*); Reinhard, 1922: 72, Tex. (as *Exorista confinis* ex *Uranotes*).

Unidentified Lycaenidae

Sellers, 1943: 78, Colo. (ex larva on *Sedum stenopetalum* Pursh); Essig, 1926: 581, U.S. (as *Zenillia confinis*).

Unidentified Lepidoptera

Miscellaneous

Sellers, 1943: 78, Mass. (ex larva on apple); Aldrich, 1932: 2, Mass. (as *Zenillia confinis* ex larva on apple).

STAGE PARASITIZED: Larva, pupa.

IMMATURES: Pupa — Greene, 1922: (descr., fig. 21 on pl. 5).

BIOLOGY: Reinhard, 1929: (33-34).

Aplomya trichiosomae Sellers

HOST: HYMENOPTERA

Cimbicidae

Trichiosoma sp.

Sellers, 1943: 94, Maine.

STAGE PARASITIZED: Larva.

Aplomya trisetosa (Coquillett)

HOST: LEPIDOPTERA

Noctuidae

Autographa californica (Speyer)

Sellers, 1943: 79, Mont.

Nephelodes emmedonia (Cramer)Sellers, 1943: 79; Aldrich and Webber, 1924: 30, Ill. (as *Zenillia*).

Pyralidae

Loxostege commixtalis (Walker)Sellers, 1943: 79, Colo.; Hoerner, 1933: 12 (as *Zenillia*).

STAGE PARASITIZED: Larva.

"Aplomyea" caesar, see *Aplomya caesar**Aplomya caesar*, see *Aplomya caesar**Aplomya confusionis*, see *Aplomya confusionis**Aplomya mitis*, see *Aplomya mitis**Aplomyiopsis* Villeneuve (Goniinae: Blondeliini)*Aplomyiopsis epilachnae* (Aldrich)All records as *Paradexodes epilachnae*.

HOST: COLEOPTERA

Coccinellidae

Epilachna borealis Fabricius

Landis and Howard, 1940: 2 (lab.).

Epilachna defecta Mulsant

Landis and Howard, 1940: 2, Mexico.

Epilachna mexicana Guérin-Méneville

Landis and Howard, 1940: 2, Mexico.

Epilachna obscurella Mulsant

Landis and Howard, 1940: 2, 5, Mexico.

Epilachna varivestis Mulsant

Jimenez J., 1959: 193-194, 207, Mexico (Mex.); Clausen, 1956: 53-54, Mexico; White, 1941: 351, Mexico; Landis and Howard, 1940: 2, 5-6, 9-27, figs. 10, 14-18, Ala.,

Ky., N.C., Ohio, Tenn., Tex., Mexico; Howard, N.F., and Landis, 1936: 2, 6, Mexico; Anon., 1931b: 1114, Mexico (ex Mexican bean beetle); Howard, N. F., 1930: 8, Mexico (ex *E. corrupta* Mulsant); Howard, N. F., 1924: 7, Mexico (ex *E. corrupta*); Aldrich, 1923a: 95, Mexico (ex *E. corrupta*).

Epilachna virgata Mulsant

Landis and Howard, 1940: 2, Mexico.

STAGE PARASITIZED: Larva, pupa.

IMMATURES: Egg — Landis and Howard, 1940: (descr., fig. 14).

First instar larva — Landis and Howard, 1940: (descr., fig. 15).

Second instar larva — Landis and Howard, 1940: (descr., fig. 16).

Third instar larva — Landis and Howard, 1940: (descr., fig. 17).

Pupa — Landis and Howard, 1940: (descr., figs. 10, 18).

BIOLOGY: Landis and Howard, 1940: (2, 5-6, 9-27).

Aplomyiopsis epilachnae (Aldrich) probably

HOST: COLEOPTERA

Coccinellidae

Epilachna defecta Mulsant

Landis and Howard, 1940: 2, Guatemala (as apparently *Paradexodes*, incomplete rearing — pupa).

STAGE PARASITIZED: Larva.

Aplomyiopsis galerucellae Villeneuve

HOST: COLEOPTERA

Chrysomelidae

Pyrrhalta luteola (Müller)

Flanders, 1940: 948, Calif. (as *Synaplomyia* ex *Galerucella xanthomelaena* (Schrank)); Villeneuve, 1933: 125-126, Oreg. (ex *Galerucella*).

STAGE PARASITIZED: Larva.

BIOLOGY: Villeneuve, 1933: (125-126).

Aporia limacodis, see *Anaporia limacodis*

Aporia sp., see *Anaporia limacodis*

Aporotachina Meade (Goniinae: Blondeliini)

Aporotachina chaetoneura (Coquillett)

Most records as *Dexodes chaetoneura*.

HOST: HYMENOPTERA

Argidae

Arge clavicornis (Fabricius) complex

Raizenne, 1957: 8, Que. (as *Neothelaira* ex *A. clavicornis* abdomen yellow).

Tenthredinidae

Nematus ventralis Say

All records ex willow sawfly.

*Procter, 1946: 436, Maine (Procter, 1938); Procter, 1938: 377, Maine.

Aporotachina dexina (Townsend)

HOST: HYMENOPTERA

Argidae

Arge pectoralis (Leach)

Raizenne, 1957: 8, Ont. (as *Neothelaira aurifrons* (Coquillett)).

"Architas" *piliventris*, see *Archytas marmoratus*

Archytas Jaennicke (Tachininae: Dejeaniini)

Archytas "alterrima," see *Archytas aterrimus*

Archytas analis, see *Archytas* spp. (including *A. apicifer* and/or *A. californiae*)

Archytas "annalis," see *Archytas* spp. (including *A. apicifer* and/or *A. californiae*)

Archytas apicifer (Walker)

HOST: LEPIDOPTERA

Lasiocampidae

Malacosoma californicum (Packard)

Townsend, 1895: 70, Colo. (as *Jurinia* ex *Clisiocampa*).

Noctuidae

Peridroma saucia (Hübner)

Anon., 1969: 59, Okla.; Bottrell, 1969: 251, Okla.; Burrell, 1967: 114, La.; Montgomery, 1933: 186, Ind. (ex *Lycophotia*); Curran, 1928: 278, Miss. (as *A. vulgaris* Curran ex *P. margaritosa* (Haworth)).

Pseudaletia unipuncta (Haworth)

Burrell, 1967: 114, La.; Breeland, 1958: 323, 325, 332, fig.

25, Tenn.; Patton, 1958: 33, Fla.; Fattig, 1949: 7 (ex *Cirphis*); Brimley, 1938: 366, N.C. (as *A. vulgaris* ex armyworm); Maxson, 1920: 81, Colo. (as "*Archytus apicifere*" ex *Cirphis*).

Spodoptera frugiperda (J. E. Smith)

Wolcott, 1951a: 482, P.R. (as *A. basifulva* "(Walker)" ex *Laphygma*).

STAGE PARASITIZED: Larva, pupa.

IMMATURES: Pupa — Breeland, 1958: (fig. 25).

BIOLOGY: Burrell, 1967: (114); Patton, 1958: (33).

Archytas aterrimus (Robineau-Desvoidy)

HOST: LEPIDOPTERA

Arctiidae

Cycnia tenera Hübner

Schaffner and Griswold, 1934: 32, 125.

Hyphantria cunea (Drury)

Fattig, 1949: 7; Brimley, 1938: 366, N.C.; Brimley, 1922: 21, N.C. (ex fall webworm).

Lasiocampidae

Malacosoma americanum (Fabricius)

Patton, 1958: 33, Fla.

Megalopygidae

Megalopyge crispata (Packard)

All records ex *Lagoa crispata*.

Johnson, 1910: 784, N.J.; Coquillett, 1897: 10, 25, D.C., Iowa.

Noctuidae

Acronicta interrupta Guenée

All records ex *Acronicta occidentalis* Grote and Robinson.

Johnson, 1910: 784, N.J.; Coquillett, 1897: 10, 23, Va.

Acronicta ovata Grote

Coquillett, 1897: 10, 23, Mo.

Baileya dormitans (Guenée)

Raizenne, 1952: 66, Ont. (as *A. "alterrima"*).

Catocala neogama (J. E. Smith)

Schaffner and Griswold, 1934: 65, 125, N.J.

Peridroma saucia (Hübner)

Davis, J. J., 1912: 86, Ill. (as "*Archytes*" ex *P. margaritosa* (Haworth)).

Plathypena scabra (Fabricius)

Schaffner and Griswold, 1934: 71-72, 125, Mass.; Webber and Schaffner, 1926: 22, New England.

Unidentified Noctuidae

Schaffner and Griswold, 1934: 72, 125.

Notodontidae

Cerura sp.

Coquillett, 1897: 10, 24.

Psychidae

Thyridopteryx ephemeraeformis (Haworth)

Balduf, 1937: 173 (questionable host).

STAGE PARASITIZED: Larva, pupa.

BIOLOGY: Patton, 1958: (33); Schaffner and Griswold, 1934: (125).

Archytas basifulva, see *Archytas apicifer*

Archytas californiae (Walker)

HOST: LEPIDOPTERA

Arctiidae

Euchaetias oregonensis (Stretch)

Arnaud, 1959: 156, Calif.

Noctuidae

Spodoptera praefica (Grote)

All records ex *Prodenia praefica*.

Bosch and Hagen, 1966: 26, Calif.; Blanchard and Conger, 1932: 1065, 1067, Calif.

Trichoplusia ni (Hübner)

Bosch and Hagen, 1966: 25, Calif.

STAGE PARASITIZED: Pupa.

BIOLOGY: Arnaud, 1959: (156).

Archytas cirphis Curran

HOST: LEPIDOPTERA

Noctuidae

Agrotis ipsilon (Hufnagel)

Zimmerman, 1958: 253, Hawaii.

Cirphis spp.

Curran, 1928: 255, Mexico (Sin.).

Leucania cholica (Dyar)

All records ex *Cirphis cholica*.

Van Zwaluwenburg, 1951: 373, W. Mexico; Curran, 1927: 498, Mexico (Sin.).

Leucania latiuscula Herrich-Schaeffer

All records ex *Cirphis latiuscula*.

Van Zwaluwenburg, 1951: 373, Mexico; Curran, 1927: 498, Mexico (Sin.); Swezey, 1927a: 499, Mexico (Sin.); Swezey, 1927b: 214, Hawaii; Swezey, 1926: 226, Hawaii (as *Archytas* sp. (*A. "cirphisae"* in footnote)).

Peridroma saucia (Hübner)

Swezey, 1935a: 93, Hawaii (ex *Lycophotia margaritosa* (Haworth)).

Pseudaletia unipuncta (Haworth)

Most records ex *Cirphis unipuncta*.

Zimmerman, 1958: 305, Hawaii (ex *Pseudaletia*); Van Zwaluwenburg, 1951: 373, Mexico; Pemberton, 1948a: 72, Hawaii; Swezey, 1935a: 93, Hawaii; Williams, 1931: 292-294, Hawaii; Swezey, 1928a: 11, Hawaii.

Spodoptera exempta (Walker)

Most records ex *Laphygma exempta*.

Zimmerman, 1958: 336, Hawaii (ex *Spodoptera*); Bianchi, 1949a: 345, Hawaii; Pemberton, 1948a: 72, Hawaii; Pemberton, 1948b: 187-188, 200, Hawaii.

Spodoptera mauritia (Boisduval)

Sweetman, 1936: 355, Hawaii; Swezey, 1927a: 501-502, Hawaii; Swezey, 1927b: 216-217, Hawaii (lab.).

Armyworm(-s) (including *Pseudaletia unipuncta* (Haworth) and/or *Spodoptera mauritia* (Boisduval))

All records ex *Cirphis unipuncta* and/or *S. mauritia*.

Swezey, 1931c: 326, Hawaii; Swezey, 1928: 673, Hawaii.

Armyworm(-s)

Bianchi, 1940: 383, Hawaii; Osborn, H. T., 1938: 151-152, Hawaii; Bryan, 1934: 416, 451, Hawaii; Swezey, 1931b: 427, Hawaii; Swezey, 1927a: 503, Hawaii.

Cutworm(-s)

Bianchi, 1940: 383, Hawaii.

STAGE PARASITIZED: Larva, pupa.

IMMATURES: Pupa—Williams, 1931: (descr.).

BIOLOGY: Pemberton, 1948b: (187-188, 200); Williams, 1931: (292-294); Swezey, 1927a: (501-502); Swezey, 1927b: (214, 216-217).

Archytas cirphis Curran probably

HOST: LEPIDOPTERA

Noctuidae

Peridroma saucia (Hübner)

Zimmerman, 1958: 290, Hawaii (ex *P. porphyrea* (Denis and Schiffermüller)).

Archytas "cirphisae," see *Archytas cirphis*

Archytas hystrix, see *Archytas metallicus*

Archytas incertus, see *Archytas incertus* or *Archytas marmoratus*

Archytas incertus (Macquart)

HOST: LEPIDOPTERA

Noctuidae

Spodoptera frugiperda (J. E. Smith)

Hayslip, 1945: 198, Fla. (as *Pseudoarchytas* ex fall armyworm).

Archytas lateralis (Macquart)

HOST: LEPIDOPTERA

Lasiocampidae

Malacosoma americanum (Fabricius)

Patton, 1958: 33, Fla.; Dozier, 1920: 372, 378, Fla.

Malacosoma californicum (Packard)

Stehr and Cook, 1968: 263, 268, 300, Ariz., Wyo.

Malacosoma californicum fragile (Stretch)

Stehr and Cook, 1968: 279, 301, Ariz.

Malacosoma californicum lutescens (Neumoegen and Dyar)

Stehr and Cook, 1968: 275-276, 301, Alta., Mont.

Malacosoma constrictum constrictum (H. Edwards)

Stehr and Cook, 1968: 256, 299, Calif.

Malacosoma incurvum incurvum (H. Edwards)

Stehr and Cook, 1968: 280, 301, Ariz.

Malacosoma spp.

Ross, 1953: 20, 23, fig. 10, B.C. (as *Nemochaeta*).

STAGE PARASITIZED: Larva, pupa.

IMMATURES: Pupa — Ross, 1953: (descr., fig. 10);

Greene, 1922: (descr., fig. 65 on pl. 14).

Archytas marmoratus (Townsend)

Most records as *Archytas piliventris* Wulp.

HOST: LEPIDOPTERA

Noctuidae

Agrotis ipsilon (Hufnagel) probably

Curran, 1927: 498, Mexico (Sin.).

Heliophila spp.

Vickery, 1915: 391, Tex.

Heliothis zea (Boddie)

Fattig, 1949: 8 (ex *H. armigera* (Hübner)); Bibby, 1942: 943, Tex. (ex *H. armigera*); Curran, 1928: 222, Tex. (ex *H. armiger*); Gowdey, 1923a: 54, Jamaica (ex *Chloridea obsoleta* (Fabricius)); Currie, 1905: 42 (ex *H. obsoleta*); Quaintance and Brues, 1905: 123, Tex. (ex *H. obsoleta*); Titus and Pratt, 1904: 41-42 (ex *H. obscura* (Fabricius)).

Heliothis spp.

Bottrell, Young, Price and Adams, 1968: 1054, Okla.

Laphygma spp.

James, H. C., 1953: 27, Jamaica.

Leucania latiuscula Herrich-Schaeffer

All records ex *Cirphis latiuscula*.

Vickery, 1929: 49, Tex.; Vickery, 1926: 1109-1111, 1116, Tex.; Vickery, 1925: 140, Tex.

Mocis spp. (including *M. latipes* (Guenée) and/or *M. repanda* (Fabricius))

Scaramuzza, 1946: 549, Cuba (ex *M. repanda*).

Pseudaletia unipuncta (Haworth)

All records ex *Cirphis unipuncta*.

Fattig, 1949: 8, Ga.; Vickery, 1929: 49, Tex.; Vickery, 1925: 140, Tex.

Spodoptera frugiperda (J. E. Smith)

Most records ex *Laphygma frugiperda*.

Scaramuzza and Barry, 1960: 997, Cuba (as "Architas"); Gibson, W. W., and Carrillo S., 1959: 180, Mexico (V.C.) (as *A. marmoratus*); Butler, 1958: 562, Ariz. (as *A. marmoratus*); Patton, 1958: 33, Fla. (as *A. marmoratus*); James, H. C., 1952: 34, Jamaica; Scaramuzza, 1951a: 299, Cuba (as "Architas"); Luginbill, 1950: 6-7, U.S.; Scaramuzza, 1942: 148, Cuba; Bruner, S. C., 1934: 96, Cuba; Van Dine and Christenson, 1932: 2, Cuba; Scaramuzza, 1929: 122, Cuba (ex *Laphygma*); Scaramuzza, 1929a: 113, Cuba

(ex *Laphygma*); Vickery, 1929: 1, 15-21, 45-49, Tex.; Curran, 1927: 498, Mexico (Sin.); Gowdey, 1926: 83, Jamaica; Vickery, 1926: 1116, Tex.; Vickery, 1925: 140, Tex.; Wolcott, 1924: 223, P. R.; Gowdey, 1923a: 20, Jamaica; Jones, T.H., and Wolcott, 1922: 47, P.R.; Gowdey, 1921: 4, Jamaica; Walton and Luginbill, 1916: 11; Cardin, 1915: 140, Cuba (as "*Architas*" ex "*Laphygma*"); Vickery, 1915: 391, Tex.; Jones, T. H., 1913: 235, P.R.; Van Dine, 1913b: 202, P.R. (as *A. "pilventris"*).

Spodoptera latifascia (Walker)

Tissot, 1945: 64, Fla. (as *A. incerta* (Macquart) ex *Prodenia*).

Spodoptera latifascia (Walker) probably

Patton, 1958: 33, Fla. (as *A. marmoratus* ex *Prodenia*).

Armyworm(-s)

Hayslip, Genung, Kelsheimer and Wilson, 1953: 46-47, Fla. (as *A. "pilventris"*); Curran, 1928: 222, La. (ex Mexican army worm).

Cutworm(-s)

Gibson, W. W., and Carrillo S., 1959: 180, Mexico (Son.) (as *A. marmoratus* ex gusano cortador); Hayslip, Genung, Kelsheimer and Wilson, 1953: 46-47, Fla. (as *A. "pilventris"*).

Unidentified Lepidoptera

Corn stalk borer(-s)

Curran, 1928: 222, Calif.?

Miscellaneous

Wolcott, 1951a: 483, P.R. (as *A. incerta* ex caterpillars on beets and peas).

STAGE PARASITIZED: Larva, pupa.

BIOLOGY: Hayslip, Genung, Kelsheimer and Wilson, 1953: (46-47); Luginbill, 1950: (6-7); Vickery, 1929: (1, 15-21, 45-49); Vickery, 1926: (1109-1111, 1116); Walton and Luginbill, 1916: (11).

Archytas metallicus (Robineau-Desvoidy)

HOST: LEPIDOPTERA

Arctiidae

Ecpantheria deflorata (Fabricius)

Brauer and Bergenstamm, 1895: 558, 591, N. Amer. (as *Jurinia hystrix* Williston ex *E. scribonia* Stoll).

Noctuidae

Spodoptera frugiperda (J. E. Smith)
Fattig, 1949: 7, Ga. (ex *Laphygma*).

Notodontidae

Datana angusii Grote and Robinson
Schaffner and Griswold, 1934: 74-75, 125; Bruner, 1890:
51-52, Nebr. (as *Tachina datanae* Bruner).

Datana integerrima Grote and Robinson
Raizenne, 1952: 81-82, Ont.; Schaffner and Griswold, 1934:
76, 125.

Datana ministra (Drury)
Raizenne, 1952: 80, Ont.; Schaffner and Griswold, 1934: 74,
125.

Datana perspicua Grote and Robinson
Schaffner and Griswold, 1934: 75, 125; Brimley, 1922: 21,
N.C. (as *A. hystrix*).

STAGE PARASITIZED: Larva.

IMMATURES: Pupa — Greene, 1922: (descr., fig. 64 on pl. 14).

BIOLOGY: Schaffner and Griswold, 1934: (125); Bruner, 1890:
(51-52).

Archytas sp. near *metallicus* (Robineau-Desvoidy)

HOST: LEPIDOPTERA

Noctuidae

Unidentified Catocalinae

Schaffner and Griswold, 1934: 67, 125, N.J.

BIOLOGY: Schaffner and Griswold, 1934: (67, 125).

Archytas piliventris, see *Archytas marmoratus*

Archytas "piliventris," see *Archytas marmoratus*

Archytas smaragdinus (Macquart)

HOST: LEPIDOPTERA

Megalopygidae

Megalopyge crispata (Packard)

Brauer and Bergenstamm, 1895: 543, 595, N. Amer. (ex
Lagoa).

Archytas vulgaris, see *Archytas apicifer*

Archytas sp. (A. "cirphisae"), see *Archytas cirphis*

Archytas spp. (including *A. apicifer* (Walker) and/or *A. californiae* (Walker))

All records as *Archytas analis* (Fabricius).

HOST: LEPIDOPTERA

Lasiocampidae

Malacosoma californicum (Packard)

Essig, 1926: 585, Calif.; Chittenden, 1901a: 61 (ex *Clisiocampa*).

Malacosoma californicum fragile (Stretch)

Essig, 1926: 585, Colo. (ex *M. fragilis*).

Tent caterpillar(-s)

Aldrich, 1915: 83.

Noctuidae

Elaphria nucicolora (Guenée)

Vickery, 1925: 140, Tex. (ex *Monodes*).

Heliophila spp.

Vickery, 1915: 391, Tex.

Leucania juncicola (Guenée)

All records ex *Cirphis juncicola*.

Sweetman, 1936: 220, N. Amer. (lab.); Allen, 1926: 424, 428 (lab.).

Peridroma saucia (Hübner)

Most records ex *Lycophotia margaritosa* (Haworth).

Sweetman, 1936: 123-125, 220, fig. 38, N. Amer.; *Crumb, 1929: 40, 115 (ex *L. saucia*, Chittenden, 1901a); Allen, 1926: 417-435, figs. 1-5, pl. 1, Miss.; Essig, 1926: 585, Calif., Colo.; Wakeland, 1920: 18, Colo.; Chittenden, 1901a: 61, Ill. (ex *Peridroma saucia*).

Pseudaletia unipuncta (Haworth)

Most records ex *Cirphis unipuncta*.

Sweetman, 1936: 220, N. Amer. (ex armyworm); Poos, 1928: 146, Va.; Allen, 1926: 417-435, figs. 1-5, pl. 1, Miss.; Vickery, 1926: 1117-1118, Tex.; Vickery, 1925: 140, Tex.; Reinhard, 1922: 72, Tex.; Sherman, 1915: 301-302, N.C. (ex *Heliophila*); Chittenden, 1901a: 61, Kans. (ex *Leucania*).

Spodoptera frugiperda (J. E. Smith)

All records ex *Laphygma frugiperda*.

Sweetman, 1936: 220, N. Amer.; Luginbill, 1928: 81, Kans.; Allen, 1926: 427-428, Miss.

Spodoptera ornithogalli (Guenée)

All records ex *Prodenia ornithogalli*.

Sweetman, 1936: 220, N. Amer. (lab.); Allen, 1926: 424, 428, Miss. (ex *P. "ornithogalli"*).

Spodoptera ornithogalli (Guenée) probably

Wolcott, 1951a: 482, P.R. (ex *Prodenia*).

Armyworm(-s)

Brimley, 1922: 21, N.C.; Reinhard, 1919: 280, Tex.; Aldrich, 1915: 83.

Cutworm(-s)

Smith, R. C., 1943: 377, Kans.; Allen, 1924: 24, Miss. (as *A. "annalis"*); Wolcott, 1924: 223, P.R. (ex cutworm on tobacco).

Unidentified Lepidoptera

Miscellaneous

Smith, R. C., 1943: 377, Kans. (ex destructive caterpillars).

STAGE PARASITIZED: Larva, pupa.

IMMATURES: First instar larva — Sweetman, 1936: (fig. 38); Allen, 1926: (descr., figs. 1-2, 5, pl. 1).

Second instar larva — Allen, 1926: (descr.).

Third instar larva — Allen, 1926: (descr., fig. 3, pl. 1).

Pupa — Allen, 1926: (descr., fig. 4, pl. 1); Greene, 1922: (descr., fig. 63 on pl. 14).

BIOLOGY: Sweetman, 1936: (123-125, 220); Allen, 1926: (417-435).

Archytas sp. (*A. apicifer* (Walker) or *A. californiae* (Walker) probably)

HOST: LEPIDOPTERA

Noctuidae

Pseudaletia unipuncta (Haworth)

Gahan, 1917: 212, Tenn. (as probably "*Archytes*" *analis* (Fabricius) ex *Leucania*).

BIOLOGY: Gahan, 1917: (212).

Archytas spp.

HOST: LEPIDOPTERA

Noctuidae

Peridroma saucia (Hübner)

Walkden, 1943: 381 (ex *P. margaritosa* (Haworth)).

Spodoptera sp.

Rosa, 1927: 360, Hawaii.

Armyworm(-s)

Swezey, 1925: 372, Hawaii.

Notodontidae

Datana integerrima Grote and Robinson

*Haseman, 1940: 13 (Baerg, 1928); Baerg, 1928: 15, Ark.

STAGE PARASITIZED: Larva.

"Archytes" analis, see *Archytas* sp. (*A. apicifer* or *A. californiae* probably)

"Archytes" aterrimum, see *Archytas aterrimum*

"Archytus apicifere," see *Archytas apicifer*

"Argyrophyla" albincisa, see *Argyrophylax albincisa*

Argyrophylax Brauer and Bergenstamm (Goniinae: Eumasicerini)

Argyrophylax "albincisa," see *Argyrophylax albincisa*

Argyrophylax albincisa (Wiedemann)

HOST: LEPIDOPTERA

Pyralidae

Eulepte concordalis Hübner

Most records ex *Mesocondyla concordalis*.

Wolcott, 1951a: 480, P.R.; Martorell, 1945a: 515, P.R.
(as *A. "albincisa"* ex *Eulepte*); Wolcott, 1924: 222,
P.R. (as *Sturmia*).

Hedylepta indicata (Fabricius)

Wolcott, 1951a: 480, P.R. (ex "*Hydelepta*"); Bruner, S.C.,
1935: 12, 36-37, Cuba (as "*Argyrophyla*" ex
Lamprosema); Aldrich, 1925: 129, P.R. (ex *Nacoleia*);
Wolcott, 1924: 222, P.R. (as *Sturmia* ex *Nacoleia*);
Walton, 1916a: 189, 192, P.R. (ex "*Nacoleia*").

Hymenia perspectalis (Hübner)

All records ex *Zinckenia perspectalis*.

Wolcott, 1951a: 480, P.R.; Wolcott, 1924: 222, P.R. (as
Sturmia).

Hymenia recurvalis (Fabricius)

Aldrich, 1925: 129, Virgin Islands (ex *H. fascialis* (Cramer));
Wilson, C. E., 1923a: 7, Virgin Islands (ex *Zinckenia*
fascialis).

Maruca testulalis (Geyer)

Bruner, S. C., 1935: 12, Cuba (as "*Argyrophyla*"); Leonard
and Mills, 1931: 468, Cuba (as "*Argyrophyla*").

Pilemia periusalis (Walker)

Wolcott, 1951a: 480, P.R. (ex *Psara*).

Pilocrocis inguinalis (Guenée)

Martorell, 1945: 349, P.R.; Martorell, 1945a: 535, P.R. (as *A. "albicansa"*).

Psara pallicaudalis Snellen

All records ex *Pachyzancla periusalis* (Walker).

Aldrich, 1925: 129, P.R. (ex *Pachyzancla "pertusalis"*);

Wolcott, 1924: 222, P.R. (as *Sturmia*).

STAGE PARASITIZED: Larva, pupa.

Argyrophylax piperi, see *Lespesia schizurae*

Argyrophylax protoparcis, see *Drino rhoeo*

Argyrophylax protoparcis probably, see *Drino rhoeo* probably

Argyrophylax schizurae, see *Lespesia schizurae*

Argyrophylax spp.

HOST: LEPIDOPTERA

Pyralidae

Etiella zinckenella (Treitschke)

Anon., 1939a: 108, P.R. (as "*Agrophylax*").

Saturniidae

Hemileuca maia (Drury)

Brauer and Bergenstamm, 1895: 544, 594, 606, N. Amer.

Sphingidae

Agrius cingulatus (Fabricius)

Brauer and Bergenstamm, 1895: 544, 598 (ex *Macrosila*).

Ceratonia amyntor (Hübner)

Brauer and Bergenstamm, 1895: 544, 586, N. Amer.

Pholus achemon (Drury)

Townsend, 1895: 72, Colo. (ex *Philampelus*).

STAGE PARASITIZED: Larva.

Argyrophylax sp. probably

HOST: LEPIDOPTERA

Arctiidae

Estigmene acraea (Drury)

Brauer and Bergenstamm, 1895: 544, 597, N. Amer. (ex *Leucarctia acraea*).

Arrhinomyia tragica, see *Elodia tragica*

Atacta Schiner (Goniinae: Belvosini)

Atacta apicalis, see *Atacta brasiliensis*

Atacta brasiliensis Schiner

HOST: LEPIDOPTERA

Noctuidae

Mocis latipes (Guenée) probably

Aldrich, 1925b: 29-30, Panama (ex *Remigia repanda* (Fabricius)).

Sphingidae

Manduca spp.

Rowe, 1933:123, Ill. (as *A. apicalis* Coquillett ex *Protoparce*).

Atacta geminata, see *Drino rhoeo*

Ateloglossa Coquillett (Proseninae: Prosenini)

Ateloglossa cinerea Coquillett

HOST: COLEOPTERA

Elateridae

Melanotus spp.

Thomas, 1940: 30, Mass.; Packard, C. M., 1929: 598, Mass.
STAGE PARASITIZED: Larva.

BIOLOGY: Packard, C. M., 1929: (598).

Athanatus Reinhard (Dexiinae: Macquartiini)

Athanatus californicus (Coquillett)

HOST: LEPIDOPTERA

Noctuidae

Heliothis ononis (Denis and Schiffermüller)

McDonald, 1947: 239, W. Canada (as *Phytopsis*).

STAGE PARASITIZED: Larva.

Athrycia Robineau-Desvoidy (Tachininae: Voriini)

Athrycia cinerea (Coquillett)

Most records as *Blepharigena spinosula* (Bigot).

HOST: HYMENOPTERA

Unidentified Hymenoptera

Sawfly(-flies)

Webster, 1898: 19, Ohio (as *Paraplusia*).

HOST: LEPIDOPTERA

Noctuidae

Pseudaletia unipuncta (Haworth)

Guppy, J. C., 1967: 97, 99-100, 102, Ont. (as *A. cinerea*);
Breeland, 1958: 323, 334, Nebr. (as *B. "cineria"* and *B. "cinera"*);
Butler, 1958: 562, Ariz.; Schaffner and
Griswold, 1934: 47, 108, Mass. (ex *Cirphis*); Allen,
1929: 684, Miss. (ex *Cirphis*).

STAGE PARASITIZED: Larva (Hymenoptera, Lepidoptera),
pupa (Lepidoptera).

IMMATURES: Pupa — Greene, 1922: (as *Paraplagia "spinulosa"*;
descr., fig. 43 on pl. 9).

BIOLOGY: Guppy, J.C., 1967: (97, 99-100, 102); Schaffner
and Griswold, 1934: (47, 108).

Aubaeanetia anorbitalis, see *Oswaldia anorbitalis*

Aubaeanetia assimilis, see *Oswaldia assimilis*

"*Belvoisia*" *unifasciata*, see *Triachora unifasciata*

Belvosia Robineau-Desvoidy (Goniinae: Belvosiini)

Belvosia argentifrons Aldrich

HOST: LEPIDOPTERA

Unidentified Lepidoptera

Miscellaneous

Aldrich, 1928a: 33, Va. (ex pupa (Hopkins 14802 F)).

STAGE PARASITIZED: Pupa.

Belvosia bicincta Robineau-Desvoidy

HOST: LEPIDOPTERA

Noctuidae

Mocis latipes (Guenée) probably

All records ex *Mocis repanda* (Fabricius).

Anon., 1954: 99, Costa Rica; Anon., 1953: 91, Costa Rica
(as *B. "bicinta"*).

Sphingidae

Hawk moth(-s)

Aldrich, 1928a: 36, Jamaica.

STAGE PARASITIZED: Larva, pupa.

Belvosia "bicinta," see *Belvosia bicincta*

Belvosia bifasciata (Fabricius)

HOST: LEPIDOPTERA

Citheroniidae

Anisota rubicunda (Fabricius)

Schaffner and Griswold, 1934: 30-31, 109; Aldrich, 1928a: 31; Johnson, 1925: 193, Conn., Mass., N.H., R.I. (ex *Dryocampa*); Johnson, 1910: 779, N.J. (as *Latreillimyia* ex *Dryocampa*); Howard and Chittenden, 1909: 5, Mo.; Felt, 1908: 16, N.Y.; Coquillett, 1897: 10, 25, Mo. (ex *Dryocampa*); Brauer and Bergenstamm, 1895: 558, 582, N. Amer. (as *Latreillia*); Packard, 1890: 393 (as *Tachina* ex *Dryocampa*); Riley, 1873: 140-141, N.Y. (as *Tachina*).

Anisota sp. near *rubicunda* (Fabricius)

Patton, 1958: 33, Fla.

Anisota senatoria J. E. Smith

Brimley, 1922: 21, N.C.

Citheronia regalis (Fabricius)

Fattig, 1949: 8-9 (ex "*Citherona*"); Aldrich, 1928a: 31; Johnson, 1925: 193, Conn., Mass., N.H., R.I.; Brimley, 1922: 21, N.C.; Johnson, 1910: 779, N.J. (as *Latreillimyia*); Coquillett, 1897: 10, 24; Brauer and Bergenstamm, 1895: 558, 587, N. Amer. (as *Latreillia*); Brauer and Bergenstamm, 1895: 558, 586 (as *Latreillia* ex *Cerocampa*); Riley, 1873: 141, Pa.

Eacles imperialis (Drury)

Most records ex *Basilona imperialis*.

Fattig, 1949: 8-9; Brimley, 1922: 21, N.C.; Walton, 1913b: 24, 27, Pa.; Britton, 1912: 341, Conn. (as *Latreillimyia*); Johnson, 1910: 779, N.J. (as *Latreillimyia* ex *Eacles*).

Notodontidae

Datana integerrima Grote and Robinson

Smith, R. C., 1943: 377, Kans. (ex walnut caterpillar).

Saturniidae

Antheraea polyphemus (Cramer)

All records ex *Telea polyphemus*.

Fattig, 1949: 8-9, Ga.; Aldrich, 1928a: 31, Mass.

Hemileuca spp. (including *H. lucina* H. Edwards and/or *H. maia* (Drury))

Schaffner and Griswold, 1934: 29, 109.

Hemileuca sp.

Coquillett, 1897: 10, 25, Calif.

Hemileuca sp. probably

Aldrich, 1928a: 31, Calif.

Sphingidae

Agrius cingulatus (Fabricius)All records ex *Herse cingulata*.

Wolcott, 1951a: 477-478, P.R.; Wolcott, 1924: 222, P.R.

Ceratomia amyntor (Hübner)

Aldrich, 1928a: 32, N.C.

Ceratomia undulosa (Walker)

Brimley, 1922: 21, N.C.

Unidentified Lepidoptera

Miscellaneous

Smith, R. C., 1943: 377, Kans.

STAGE PARASITIZED: Larva, pupa.

IMMATURES: Pupa — Greene, 1922: (descr., fig. 15 on pl. 4).

BIOLOGY: Patton, 1958: (33); Wolcott, 1951a: (477-478);

Schaffner and Griswold, 1934: (109); Britton, 1912: (341).

Belvosia borealis Aldrich

HOST: LEPIDOPTERA

Sphingidae

Ceratomia catalpae (Boisduval)

Fattig, 1949: 9, Ga.

Ceratomia undulosa (Walker)Fattig, 1949: 9; Brimley, 1938: 361, [N.C.]; Brimley, 1928: 205, N.C. (as *B. orion* Brimley).

STAGE PARASITIZED: Pupa.

Belvosia ciliata Aldrich

HOST: LEPIDOPTERA

Hesperiidae

Unidentified Hesperiidae

Arnaud, 1963: 108, Mexico.

Saturniidae

Automeris nyctimene leucane Geyer

Aldrich, 1928a: 23, Mexico (ex *A. leucane*).

Copaxa lavendera (Westwood) probably

Aldrich, 1928a: 23, Mexico.

Belvosia ciliata var. *formosa*, see *Belvosia formosa*

Belvosia ferruginosa Townsend

HOST: LEPIDOPTERA

Unidentified Lepidoptera

Miscellaneous

Johnson, 1919: 436, Jamaica (ex chrysalis); Townsend,
1895: 71, Jamaica (ex chrysalis).

STAGE PARASITIZED: Pupa.

Belvosia formosa Aldrich

HOST: LEPIDOPTERA

Saturniidae

Rothschildia orizaba (Westwood)

Aldrich, 1928a: 23, Mexico (as *B. ciliata* var. *formosa* ex
Saturnia).

Belvosia nigrifrons Aldrich

HOST: LEPIDOPTERA

Saturniidae

Hyalophora calleta (Westwood)

Aldrich, 1928a: 39, El Salvador (ex *Callosamia* "colleta").

Rothschildia aroma Schaus probably

Quezada, 1967: 595, 597-598, El Salvador.

Rothschildia orizaba (Westwood)

Aldrich, 1928a: 39, El Salvador (ex "Attacas").

Rothschildia sp.

Aldrich, 1928a: 39, Honduras.

STAGE PARASITIZED: Larva, pupa.

Belvosia ochriventris, see *Triachora ochriventris*

Belvosia omissa, see *Triachora omissa*

Belvosia orion, see *Belvosia borealis*

Belvosia recticornis Macquart

HOST: LEPIDOPTERA

Saturniidae

Hylesia alinda Druce

Aldrich, 1928a: 15, Mexico.

Hylesia darlingi Dyar

Aldrich, 1928a: 15, Panama.

Hylesia sp.

Aldrich, 1928a: 15, Panama.

Belvosia townsendi Aldrich

HOST: LEPIDOPTERA

Citheroniidae

Citheronia regalis (Fabricius)

Aldrich, 1928a: 34.

Eacles imperialis (Drury)

Schaffner and Griswold, 1934: 31, 109, N.J.; Aldrich,
1928a: 34, N.Y., Pa.

Sphingidae

Ceratonia amyntor (Hübner)

Aldrich, 1928a: 34, N.C.

Unidentified Lepidoptera

Miscellaneous

Aldrich, 1928a: 34, Va. (ex large pupa (Hopkins 14802 F)).

STAGE PARASITIZED: Larva, pupa.

BIOLOGY: Schaffner and Griswold, 1934: (31, 109).

Belvosia unifasciata, see *Triachora unifasciata*

Belvosia sp.

HOST: LEPIDOPTERA

Citheroniidae

Anisota rubicunda (Fabricius)

Brauer and Bergenstamm, 1895: 558, 591, N. Amer. (as
Latreillia bifasciata or *Willistonina bicincta* ex
Dryocampa).

Beskia Brauer and Bergenstamm (Phasiinae: Cylindromyiini)

Beskia aelops (Walker)

HOST: HEMIPTERA (HETEROPTERA)

Pentatomidae

Pentatomid spp. (including *Euschistus ictericus* (Linnaeus),
Mormidea pictiventris Stål, *Nezara viridula* (Linnaeus)
and/or *Oebalus pugnax* (Fabricius))

Thames, 1954: 236, Fla. (ex *E. ictericus*, *M. pictiventris*,
"Nezera" *viridula* and/or *Solubea pugnax*).

HOST: HEMIPTERA (HOMOPTERA)

Aphididae

Macrosiphum avenae (Fabricius) probably

Townsend, 1928: 150-151, Okla. (ex grain aphid on wheat,
probably *M. granaria* (Kirby)).

HOST: LEPIDOPTERA

Noctuidae

Alabama argillacea (Hübner)

Creighton, 1936: 93, Fla.

STAGE PARASITIZED: Adult (Hemiptera—Heteroptera).

IMMATURES: Pupa — Greene, 1922: (descr., fig. 11 on pl. 3).

Beskia aelops (Walker) probably

HOST: ORTHOPTERA

Tettigoniidae

Neoconocephalus robustus robustus (Scudder)

Nutting, 1953: 76, Mass.

STAGE PARASITIZED: Adult.

Bessa Robineau-Desvoidy (Goniinae: Exoristini)

Bessa harveyi (Townsend)

HOST: HYMENOPTERA

Argidae

Arge clavicornis (Fabricius) complex

Turnock and Melvin, 1963: 648, Que.; Raizenne, 1957: 8,
Que. (ex *A. clavicornis* abdomen yellow).

Arge pectoralis (Leach)

Turnock and Melvin, 1963: 648, Man., Mass.

Arge rubiginosa (Palisot de Beauvois)

Turnock and Melvin, 1963: 648, R.I.

Arge sp.

Turnock and Melvin, 1963: 648, Ont.

Diprionidae

Diprion hercyniae (Hartig)

Turnock and Melvin, 1963: 647-648, Maine, N.B., Nfld., N.S., Ont., P.E.I., Que. (ex *Gilpinia*); Finlayson, T., 1960a: 923, 936-937, fig. 12, Canada.

Neodiprion abietis (Harris)

Finlayson, T., 1963: 477, 502.

Neodiprion abietis (Harris) complex

Turnock and Melvin, 1963: 647-648, Man., N.S., Ont., Que., Sask.

Neodiprion lecontei (Fitch)

Finlayson, T., 1963: 479, 502; Turnock and Melvin, 1963: 648, Ont.; Monteith, 1962: 881-883, Ont. (lab.); Raizenne, 1957: 20, Ont.; Benjamin, 1955: 20, Canada.

Neodiprion maurus Rohwer

Turnock and Melvin, 1963: 648, Ont.

Neodiprion nanulus nanulus Schedl

Turnock and Melvin, 1963: 648, Ont.

Neodiprion pratti banksianae Rohwer

Turnock and Melvin, 1963: 648, Man., Ont.

Neodiprion swaini Middleton

Turnock and Melvin, 1963: 648, Ont., Que.

Neodiprion virginiana Rohwer complex

Finlayson, T., 1963: 485, 502; Turnock and Melvin, 1963: 648.

Neodiprion spp.

Turnock and Melvin, 1963: 648, B.C., Ont., S.C.

Tenthredinidae

Amauronematus spp.

Turnock and Melvin, 1963: 648, Man., Sask.

Ametastegia pallipes (Spinola)

Tothill, 1913: 72, Ont. (as *Frontina tenthredinidarum* (Townsend) ex *Emphytus canadensis* Kirby).

Anoplonyx canadensis Harrington

Turnock and Melvin, 1963: 648, Ont.

Anoplonyx luteipes (Cresson)

Turnock and Melvin, 1963: 648, Man., Nfld.

Anoplonyx spp.

Turnock and Melvin, 1963: 648, B.C.

Cladius isomerus Norton

Turnock and Melvin, 1963: 648, Ont.

Cladius pectinicornis Fourcroy

Tothill, 1913: 72, Ont. (as *Frontina tenthredinidarum*).

Croesus latitarsus Norton

Turnock and Melvin, 1963: 648, Man., Ont., Sask.

Hemichroa crocea (Fourcroy)

Turnock and Melvin, 1963: 647-648, B.C., Man., N.B., Ont., Sask.

Macremphytus varianus (Norton)

Turnock and Melvin, 1963: 648, Man., Sask.

Macremphytus spp.

Turnock and Melvin, 1963: 648, Man.

Nematus chalceus (Marlatt) probably

Turnock and Melvin, 1963: 648, Sask.

Nematus hyalinus (Marlatt)

Turnock and Melvin, 1963: 648, Man.

Nematus limbatus Cresson

Turnock and Melvin, 1963: 648, Man., Ont.

Nematus ribesii (Scopoli)

Turnock and Melvin, 1963: 648, Maine, Mich., N.H., Ont.

Nematus ventralis Say

Turnock and Melvin, 1963: 648, B.C., Conn., Man., Wis.

Pikonema alaskensis (Rohwer)

Monteith, 1964: 1480-1481 (lab.); Turnock and Melvin, 1963: 647, 649, Alta., Maine, Man., Minn., Ont., Que., Sask.; Raizenne, 1957: 32, Ont.; Bradley, 1951: 130, Sask.

Pikonema dimmockii (Cresson)

Turnock and Melvin, 1963: 649, Man., Que.

Pikonema sp.

Turnock and Melvin, 1963: 649, Maine.

Platycampus spp.

Turnock and Melvin, 1963: 649, Idaho, Sask.

Priophorus morio (Lepeletier)

Turnock and Melvin, 1963: 649, Calif.

Pristiphora erichsonii (Hartig)

Ives, Turnock, Buckner, Heron and Muldrew, 1968: 14, 25-27, 32-35, Man.; Ives, 1964: 39, Man. (ex larch sawfly); Monteith, 1964: 1480-1481 (lab.); Turnock and Muldrew, 1964: 95-96, Man. (ex larch sawfly); Turnock and Melvin, 1963: 647, 649, Alta., B.C., Maine, Man., Mich., Minn., Mont., N.B., Nfld. N.S., N.W.T., Ont., Que., Sask., Wis.; Monteith, 1962: 881, Ont.; Turnock and Ives, 1962: 898, 900, Man.; Bracken and Martineau, 1961: 259, Que.; Heron, 1961: 431-433 (lab.); Nairn,

Turnock, Ives and Buckner, 1961: 37-39, 41-42, 45, Cent. Canada, Man.; Heron, 1960: 476-480, Canada; Monteith, 1960: 641-646, 648-651, Ont.; Turnock, 1960: 513-514, Man., Sask.; Buckner, 1959: 537-542, Man.; Ives and Prentice, 1959: 496-500, Man.; Ives and Turnock, 1959: 652, 656-658, Man.; Raizenne, 1957: 30, Ont.; Graham, S. A., 1956: 152-154, Minn.; Ives, Muldrew, Turnock and Nairn, 1956: 13, 15, 17-18, Man. (ex larch sawfly); Muldrew, 1956: 13, Man. (ex larch sawfly); Anon., 1955c: 196, St. Lawrence Gulf prov.; Graham, 1955: 66 (lab.); Gardiner, 1954: 18, Man., Sask. (ex larch sawfly); Lejeune and Hildahl, 1954: 337, 339-340, Man., Sask.; McLeod, 1954: 23, B.C.; Reeks, 1954: 474-479, N.B., N.S.; Drooz, 1953: 828, Minn.; Anon., 1952d: 209-210, Man., Sask.; Lejeune, 1951: 153-154, Man., Sask.; Lejeune, 1951a: 22-23, Man., Sask.; MacNay, 1951: 127, Man., Sask. (ex larch sawfly); Hildahl, 1950: 109, Man., Sask.; Gardiner, 1949: 97, B.C. (ex larch sawfly); Gardiner, 1948: 89, Man. (ex larch sawfly); Gardiner, 1947: 93, Man. (ex larch sawfly); McGuffin and Barker, 1947: 61, Man. (ex "European" larch sawfly); Reeks, Forbes and Cuming, 1947: 9, Nfld.; Anon., 1946: 195, maritime area; Gardiner, 1946: 74, Man. (ex larch sawfly); Procter, 1946: 435, Maine (as *B. tenthredinidarum* ex "*Lygaeonematis ericksonii*"); *Procter, 1946: 434, Maine (as *Achaetoneura tenthredinidarum* ex "*Lygaeonematis ericksonii*," Procter, 1938); Reeks, 1946: 26, Nfld.; Baird, 1944: 119, 129, 137, N.B., Ont.; Baird, 1943: 110, 121, 127, N.B., Ont.; Procter, 1938: 375, Maine (as *Achaetoneura tenthredinidarum* ex "*Lygaeonematis ericksonii*"); Baird, 1923: 159, 163-164, 166, N.B. (as *Frontina tenthredinidarum* ex *Lygaeonematus*); Britton, 1916a: 131, N.B. (as *Frontina tenthredinidarum* ex *Lygaeonematus*); Hewitt, 1912: 33, N.B. (as *Frontina tenthredinidarum* ex *Nematus*).

Pristiphora ericksonii (Hartig) probably

*Vide Hewitt, 1912: 33, Ont. (as *Frontina tenthredinidarum* ex *Nematus*, Brauer and Bergenstamm, 1895; Townsend, 1892a); *Brauer and Bergenstamm, 1895: 561, 600 (as *Masicera tenthredinidarum* ex *Nematus?* sp., Townsend, 1892a); Townsend, 1892a: 285-286, Canada (as *Masicera tenthredinidarum* ex *Nematus?* [sp.]).

Pristiphora geniculata (Hartig)

Forbes and Daviault, 1964: 1128, N.E. states; Turnock and Melvin, 1963: 647, 649, N.B., N.S., Ont., Que.; Raizenne, 1957: 30, Ont.

Pristiphora leechi Wong and Ross

Turnock and Melvin, 1963: 649, B.C.

Pristiphora lena Kincaid

Turnock and Melvin, 1963: 649, Man.

Strongylogaster sp.

Turnock and Melvin, 1963: 649, Man.

Trichiocampus irregularis (Dyar)

Turnock and Melvin, 1963: 649, Ont.

Trichiocampus viminalis (Fallén)

Turnock and Melvin, 1963: 649, 652, Maine, Man., Ont., Que.; Raizenne, 1957: 27, Ont., Que.

Trichiocampus sp.

Turnock and Melvin, 1963: 649, Sask.

Unidentified Tenthredinidae

Turnock and Melvin, 1963: 649, Alta., B.C., Man., Sask.

HOST: LEPIDOPTERA

Geometridae

Eupithecia luteata Packard

Turnock and Melvin, 1963: 649, Man.

Nepytia canosaria (Walker)

Turnock and Melvin, 1963: 649, N.B.

Semiothisa granitata Guenée complex

Turnock and Melvin, 1963: 649, B.C.

Unidentified Geometridae

Turnock and Melvin, 1963: 649, Que.

Nymphalidae

Nymphalis antiopa (Linnaeus)

Turnock and Melvin, 1963: 649, N.B.

Tortricidae

Archips sp.

Turnock and Melvin, 1963: 649, Sask.

Unidentified Tortricidae

Turnock and Melvin, 1963: 649, Man.

STAGE PARASITIZED: Larva (Hymenoptera), pupa (Hymenoptera).

IMMATURES: Egg — Monteith, 1960: (descr.).

Second instar larva — Finlayson, T., 1960a: (descr., fig. 12).

Pupa — Greene, 1922: (descr., fig. 82 on pl. 17).

BIOLOGY: Ives, Turnock, Buckner, Heron and Muldrew, 1968: (14, 25-27, 32-35); Monteith, 1964: (1480-1481); Monteith, 1962: (881-883); Heron, 1961: (431-433); Nairn, Turnock, Ives and Buckner, 1961: (37-39, 41-42, 45); Finlayson, T., 1960a: (923, 936-937); Heron, 1960: (476-480); Monteith, 1960: (641-646, 648-651); Turnock, 1960: (513-514); Ives, Muldrew, Turnock and Nairn, 1956: (13, 15, 17-18); Muldrew, 1956: (13); Baird, 1923: (159, 163-164, 166).

Bessa selecta (Meigen)

HOST: HYMENOPTERA

Diprionidae

Diprion hercyniae (Hartig)

Briand, 1949: 56-57, Que. (ex [European] spruce sawfly);
Hawboldt, 1947: 84-103, pls. 8-9, N.B., N.S. (ex
Gilpinia); Lambert, 1941: 36 (ex European spruce
sawfly).

Diprion polytomum (Hartig)

Reeks, 1938: 25-26, E. Canada; Balch, 1936: 28, Que.

Neodiprion lecontei (Fitch)

Anon., 1959a: 272, Minn. (ex red-headed pine sawfly).

Neodiprion swaini Middleton

Anon., 1959a: 272, Minn. (ex Swaine jack-pine sawfly).

Tenthredinidae

Hemichroa crocea (Fourcroy)

Hopping, 1937: 248-249, B.C.

Pikonema alaskensis (Rohwer)

Anon., 1959a: 272, Minn. (ex yellow-headed spruce sawfly);
Rendall, 1943: 39, Maine; Nash, 1939: 333, Maine.

Pikonema dimmockii (Cresson)

Reeks, 1938: 26 (lab.).

Priophorus morio (Lepeletier)

Smith, L.M., and Kido, 1949: 50, Calif. (ex *P. rubivorus*
Rohwer).

Pristiphora erichsonii (Hartig)

Drooz, 1960: 32-33, 45, Minn.; Anon., 1959a: 272, Minn.
(ex larch sawfly); Anon., 1959b: 711, Minn.; Hawboldt,
1947: 84; Hopping, Leech and Morgan, 1943: 59, 61,

B.C.; Baird, 1942: 115, 127, 132-133, N.B., Ont.; Baird, 1941a: 97, 118, 121-122, 124, N.B.; Balch, 1941: 37, N.B.; Balch and Hawboldt, 1941: 24, N.B.; Lambert, 1941: 36, Que.; Twinn, 1941: 12, N.B., N.S.; Atwood, 1940: 211, Ont. (ex larch sawfly); Baird, 1940: 97, 117, 122-123, N.B., Que.; Balch, 1940: 22, N.S.; Richmond, 1940: 52, Man.; Atwood, 1939: 38, Ont., Que.; Baird, 1939: 105, 121, 125, 127, N.B., Que.; Balch, 1939: 30, N.B., N.S.; Hopping, 1939: 277, B.C.; Twinn, 1939: 18, B.C., Man., N.B., N.S., Ont.; Atwood, 1938: 296, Ont. (ex larch sawfly, incomplete rearing — egg); Balch, 1938: 26, N.B., N.S., Que. (ex *Lygaeonematus*); Brown, A.W.A., 1938: 17, E. Canada; Reeks, 1938: 25, N.B.; Twinn, 1938: 132, N.B., N.S.; Baird, 1937: 67, Canada (ex *Lygaeonematus*); Dowden, 1937: 381, Great Lakes region, Mont. (ex *Nematus "ericksonii"*); Hopping, 1937: 248-249, B.C. (ex "European" larch sawfly); Hopping, R., 1937: 63, B.C. (ex *Lygaeonematus*); Twinn, 1937: 85, N.B. (ex *Lygaeonematus*); Balch, 1936a: 144, N.B. (ex *Lygaeonematus*); Balch, 1936b: 197, N.B., N.S. (ex *Lygaeonematus*).

Pristiphora geniculata (Hartig)

Forbes and Daviault, 1964: 1128, N.E. U.S. (as "*Ptychomia*").

HOST: LEPIDOPTERA

Lymantriidae

Hemerocampa leucostigma (J. E. Smith)

Beaulne, 1953: 195, Que.

STAGE PARASITIZED: Larva (Hymenoptera), pupa (Hymenoptera).

IMMATURES: Egg — Hawboldt, 1947: (descr., figs. 1-2 on pl. 8).

First instar larva — Hawboldt, 1947: (descr., figs. 3-6, 9-10 on pl. 8).

Second instar larva — Hawboldt, 1947: (descr., figs. 7-8, 11, 13, 19 on pl. 8).

Third instar larva — Hawboldt, 1947: (descr., figs. 12, 14 on pl. 8, 20-24 on pl. 9).

Pupa — Hawboldt, 1947: (descr., figs. 15-18 on pl. 8, 25-30 on pl. 9).

BIOLOGY: Drooz, 1960: (32-33, 45); Smith, L. M., and Kido, 1949: (50); Hopping, Leech and Morgan, 1943: (59, 61); Reeks, 1938: (25-26).

Bessa selecta (Meigen) probably
HOST: HYMENOPTERA

Tenthredinidae

Pristiphora erichsonii (Hartig)

Leech, 1936: 99, B.C. (ex larch sawfly).

STAGE PARASITIZED: Pupa.

BIOLOGY: Leech, 1936: (99).

Bessa tenthredinidarum, see *Bessa harveyi*

Bessa spp.

HOST: HYMENOPTERA

Tenthredinidae

Pristiphora erichsonii (Hartig)

Hawboldt, 1940: 155, N.B.

STAGE PARASITIZED: Pupa.

"*Bessaria*" (*Celatoria*) *crawii*, see *Celatoria diabroticae*

Besseria crawii, see *Celatoria diabroticae*

Bigonichaeta setipennis, see *Bigonicheta spinipennis*

Bigonichaeta spp., see *Bigonicheta spinipennis*

Bigonicheta Rondani (Tachininae: Germariini)

Bigonicheta setipennis, see *Bigonicheta spinipennis*

Bigonicheta spinipennis (Meigen)

Most records as *Bigonicheta setipennis* (Fallén).

HOST: DERMAPTERA

Forficulidae

Forficula auricularia Linnaeus

Sabrosky and Arnaud, 1965: 1012 (as *B. spinipennis* ex [European] earwigs); Anon., 1958: 126, Nfld. (as *B. spinipennis*); Johansen, 1957: 62-63, Wash.; Clausen, 1956: 5, 71, 85-86, Idaho, Oreg., Utah, Wash.; MacNay, 1956: 295-296, 299, 301, B.C. (as *Bigonichaeta* spp.); MacNay, 1955: 321, 326, 328, B.C. (as *Bigonichaeta spinipennis*); Anon., 1954f: 300, B.C. (lab.); Davis, J. E., 1954: 204, Oreg. (probable host); MacNay, 1954: 332, 336, 339, B.C.; McLeod, 1954: 20-21, B.C. (as *Bigonichaeta*); MacNay, 1953a: 307, 312, 316, B.C.; Anon., 1952b: 175, 181, 186, B.C.; Anon., 1952g: 284, 291, 295, B.C.; McLeod and Chant, 1952: 343-345, B.C.; McLeod, 1951: 28, 30-32, B.C. (as *Bigonichaeta*);

Spencer, 1947: 8-9, B.C. (as *Digonochaeta*); Baird, 1942: 115, 126, 133, Ont. (as *Bigonichaeta*); Baird, 1941a: 97, 114, 123, B.C., Ont.; Smith, R. W., 1941: 29-32, B.C.; Baird, 1940: 99-100, 114, 121, B.C. (as *Digonochaeta*); Eddy, 1940: 60-62, Wash. (ex European earwig); Gardiner, 1940: 23, Ont. (as *Digonochaeta* ex European earwig, lab.); Knowlton, 1940: 204, Utah (incomplete rearing — pupa); Whitcomb, 1940: 65, Wash. (as *Bigonichaeta* ex European earwig); [Gardiner], 1939: 16, B.C. (as *Digonichaeta* ex [European] earwig, lab.); Baird, 1938: 88, 127, 140, U.S. (as *Digonochaeta*); Downes, 1938: 73, B.C. (as *Digonichaeta* ex European earwig); Gardiner, 1938: 101-102, B.C. (as *Digonichaeta* ex [European] earwig); Getzendaner, 1937: 9-12, Wash. (as *Digonichaeta*); Getzendaner, 1936: 1105-1114, Wash. (ex European earwig, lab.); Sweetman, 1936: 119-120, 233, 238, 311, N. Amer. (as *Digonichaeta*); Twinn, 1936: 85, B.C. (as *Digonochaeta*); Downes, 1935: 181, B.C. (as *Digonochaeta* ex [European] earwig); Dimick and Mote, 1934: 863-865, Oreg. (as *Digonichaeta* ex European earwig, lab.); Mote, 1933: 259, Oreg. (as *Digonichaeta*); Dimick, 1931: 962-964, Oreg. (as *Digonichaeta* ex European earwig, lab.); Mote, 1931: 948-956, Oreg. (as *Digonichaeta*); Mote, 1931a: 230, Oreg. (as *Digonichaeta*); Mote, Stearns and Dimick, 1931: 957-961, Oreg. (as *Digonichaeta* ex European earwig); Atwell, 1927: 101-102, Oreg. (as *Digonochaeta* [sp.], lab.); Jardine, 1926: 105, Oreg. (as *Digonochaeta* ex European earwig, lab.).

STAGE PARASITIZED: Adult.

BIOLOGY: Anon., 1958: (126); Clausen, 1956: (5, 71, 85-86); Smith, R. W., 1941: (29-32); Sweetman, 1936: (119-120, 233, 238, 311); Mote, Stearns and Dimick, 1931: (957-961); Atwell, 1927: (101-102).

Biomyia eleodivora, see *Sitophaga eleodivora*

Biomyia georgiae, see *Sitophaga georgiae*

Biomyia georgiae probably, see *Sitophaga georgiae* probably

Biomyia lachnosternae, see *Sitophaga georgiae*

Blepharidea hirsuta, see *Phryxe vulgaris* or *Phryxe pecosensis* probably

Blepharigena "cinera," see *Athrycia cinerea*

Blepharigena "cineria," see *Athrycia cinerea*

Blepharigena spinosula, see *Athrycia cinerea*

Blepharipa Rondani (Goniinae: Sturmiini)

Blepharipa schineri (Mesnil)

HOST: LEPIDOPTERA

Lymantriidae

Porthetria dispar (Linnaeus)Thompson, 1910: 284 (as *Crossocosmia sericariae* (Rondani)
ex "gipsy" moth, lab.).

BIOLOGY: Thompson, 1910: (284)

Blepharipa scutellata (Robineau-Desvoidy)Most records as *Sturmia scutellata*.

HOST: LEPIDOPTERA

Lasiocampidae

Malacosoma americanum (Fabricius)

Bess, 1936: 605.

Malacosoma disstria HübnerBess, 1936: 605; Schaffner, 1934: 588; Schaffner and
Griswold, 1934: 89, 118.

Lymantriidae

Euproctis chrysorrhoea (Linnaeus)Burgess, 1924: 34, Mass. (as *Blepharipa*).*Porthetria dispar* (Linnaeus)Dowden, 1962: 32-33, 50-51, New England (as
Blepharipoda); Nichols, 1962: 20, Pa.; Dowden, 1961:
875, Mass.; Clausen, 1956: 6-7, 63, 86-87 (as
Blepharipoda); Anon., 1955a: 5, N.H. (ex gypsy moth);
Anon., 1955b: 13, 15, 17, N.H. (ex gypsy moth); Anon.,
1954g: 309, E. U.S. (ex *Lymantria*); Anon., 1951: 1, 4,
Mass. (ex gypsy moth); Anon., 1950a: 2, Mass. (ex gypsy
moth); Franklin, 1950: 57, 62, fig. 8B; Friend, 1945:
621, Conn.; Brown and Sheals, 1944: 400; Sweetman,
1936: 120-121, 221, 232-233, fig. 35, E. N. Amer.;
Britton, 1935: 637, Conn.; Collins, 1935: 38, Mass. (ex
gypsy moth); Dowden, 1935: 495, 509, 511, 521, Maine,
Mass., N.H.; Collins, 1934: 31, Mass., New England (ex
gypsy moth); Proper, 1934: 372, New England (ex gypsy
moth); Schaffner, 1934: 588, New England, N.J., Pa.;
Britton and Ashworth, 1933: 407, Conn. (ex "gipsy"
moth); Collins, 1933: 27, Mass. (ex gypsy moth); Collins
and Jones, 1932: 237 (ex "gipsy" moth); Webber, 1932:

205-206; Anon., 1931a: 775, N.Y. (ex "gipsy" moth, incomplete rearing — pupa); Anon., 1931c: 124, N.Y. (incomplete rearing — pupa); Burgess, 1930: 15-16, Mass.; Burgess and Crossman, 1929: 63-65, 67-76, 142, Conn., Mass., N.H., Vt.; Crowell, 1929: 220-227, figs. 1-4, N.H. (as *Blepharipa*); Burgess, 1924: 34, Mass. (as *Blepharipa*); Britton and Ashworth, 1923: 316, Conn. (as *Blepharipa* ex "gipsy" moth); Lochhead, 1919: 200, 406 (as *Blepharipa*); Howard, 1918: 45-46, Mass. (as *Blepharipa* ex gipsy moth); Pouillaude, 1913: 82, Mass. (as *Blepharipa* ex "Porthesia"); Howard and Fiske, 1911: 213-218, 308, figs. 36-40, pl. 18 (as *Blepharipa*); Thompson, 1910: 284, 289 (as *Blepharipa* ex "gipsy" moth).

Stilpnotia salicis (Linnaeus)

Burgess and Crossman, 1927: 16, New England.

Noctuidae

Catocala sp.

Schaffner, 1934: 588.

Lithophane sp.

Schaffner, 1934: 588 (ex *Graptolitha*).

Lithophane spp. (including *L. disposita* Morrison, *L. innominata* (Smith) and/or *L. petulca* Grote)

Schaffner and Griswold, 1934: 48, 118 (ex *Graptolitha* spp. (including *G. disposita*, *G. innominata* and/or *G. petulca*)).

Unidentified Catocalinae

Schaffner and Griswold, 1934: 67, 118-119.

Unidentified Noctuidae

Schaffner, 1934: 588; Schaffner and Griswold, 1934: 72, 118.

Notodontidae

Datana integerrima Grote and Robinson

Schaffner, 1934: 588; Schaffner and Griswold, 1934: 76, 118.

Symmerista albifrons (J. E. Smith)

Schaffner, 1934: 588; Schaffner and Griswold, 1934: 78, 118.

STAGE PARASITIZED: Larva, pupa.

IMMATURES: Egg — Franklin, 1950: (descr., fig. 8B); Sweetman, 1936: (fig. 35); Howard and Fiske, 1911: (figs. 36-37).

First instar larva — Howard and Fiske, 1911: (descr., fig. 38).

Second instar larva — Howard and Fiske, 1911: (descr., figs. 39-40).

Third instar larva — Crowell, 1929: (descr., figs. 1-4); Howard and Fiske, 1911: (pl. 18).

Pupa — Howard and Fiske, 1911: (pl. 18).

BIOLOGY: Dowden, 1962: (32-33, 50-51); Clausen, 1956: (6-7, 63, 86-87); Franklin, 1950: (57, 62); Sweetman, 1936: (120-121, 221, 232-233); Dowden, 1935: (495, 509, 511, 521); Schaffner, 1934: (588); Schaffner and Griswold, 1934: (118); Collins and Jones, 1932: (237); Webber, 1932: (205-206); Burgess and Crossman, 1929: (63-65, 67-76, 142); Howard and Fiske, 1911: (213-218, 308); Thompson, 1910: (284, 289).

Blepharipa sp.

HOST: LEPIDOPTERA

Lymantriidae

Porthetria dispar (Linnaeus)

Howard, 1913: 72 (ex gypsy moth).

Blepharipeza adusta, see *Leschenaultia adusta*

Blepharipeza exul, see *Leschenaultia exul*

Blepharipeza fulvipes, see *Leschenaultia fulvipes*

Blepharipeza leucophrys, see *Leschenaultia leucophrys*

Blepharipeza sp., see *Leschenaultia* spp.

Blepharipoda scutellata, see *Blepharipa scutellata*

"*Blepharopeza*" *adusta*, see *Leschenaultia adusta*

Blondelia Robineau-Desvoidy (Goniinae: Blondeliini)

Blondelia armigera, see *Eucelatoria armigera*

Blondelia eufitchiae (Townsend)

Most records as *Phrynolydella eufitchiae*.

HOST: LEPIDOPTERA

Arctiidae

Halisidota tessellaris (J. E. Smith)

Tothill, 1913: 72, Ont. (as *Masicera*).

Hyphantria cunea (Drury)

Essig, 1926: 583, Calif. (as *Masicera*); Johnson, 1910: 781, N.J. (as *Masicera*).

Hyphantria textor (Harris)

Raizenne, 1952: 29, Ont.; *Brauer and Bergenstamm, 1895: 560, 594, N. Amer. (as *Masicera* "eufitchii," Townsend, 1893); Townsend, 1893: 467, Ill. (as *Masicera*).

Geometridae

Alsophila pometaria (Harris)

Raizenne, 1952: 108, Ont.; Schaffner and Griswold, 1934: 90, 120 (as *Anetia*).

Caripeta angustiorata Walker

Raizenne, 1952: 154, Que.

Ennomos subsignarius (Hübner)

Raizenne, 1952: 151, Ont.

Eupithecia gibsonata Taylor

Raizenne, 1952: 114, Ont.

Itame flavicaria (Packard)

Essig, 1926: 583, Colo. (as *Masicera*).

Itame quadrilinearia (Packard)

Essig, 1926: 583, Colo. (as *Masicera*).

Itame ribearia (Fitch)

Most records ex *Eufitchia ribearia*.

Essig, 1926: 583 (as *Masicera* ex *Itame*); Johnson, 1910: 781, N.J. (as *Masicera*); *Brauer and Bergenstamm, 1895: 560, 592, N. Amer. (as *Masicera* "eufitchii," Townsend, 1892a); Townsend, 1892a: 286-287, Colo. (as *Masicera*).

Lambdina fiscellaria (Guenée)

Schaffner and Griswold, 1934: 99, 120 (as *Anetia* ex *Ellopi*).

Lambdina fiscellaria fiscellaria (Guenée)

Carroll, 1956: 594, Nfld. (as *Anetia*); Raizenne, 1952: 156, Ont.

Nepytia canosaria (Walker)

Raizenne, 1952: 152, Ont.

Semiothisa bisignata (Walker)

Raizenne, 1952: 122, Ont., Que.

Semiothisa granitata Guenée complex

Raizenne, 1952: 124, Ont.

Semiothisa minorata (Packard)

Raizenne, 1952: 121, Que.

Semiothisa orillata (Walker)

Raizenne, 1952: 129, Ont.

Semiothisa oweni (Swett)

Raizenne, 1952: 126, Que.

Semiothisa sexmaculata (Packard)

Raizenne, 1952: 127, Ont.

Thera juniperata (Linnaeus)

Raizenne, 1952: 117, Ont. (ex *T. procteri* Brower).

Noctuidae

Hypena humuli (Harris)

Hawley, 1918: 196, N.Y. (as *Masicera*).

Olethreutidae

Spilonota ocellana (Denis and Schiffermüller)

Stultz, 1955: 80, N.S.

Tortricidae

Acleris minuta (Robinson)

Webster, 1898: 19, Ohio (as *Masicera* ex *Teras minuta* var. *cinderella* Riley).

Sparganothis distincta (Walsingham)

Webster, 1898: 19, Ohio (as *Masicera* ex *Oenectra*).

Unidentified Tortricidae

Caesar, 1916a: 172-173, unnumbered fig. on p. 172, Ont. (as *Masicera* ex *Tortrix* spp.).

STAGE PARASITIZED: Larva, pupa.

IMMATURES: Pupa — Greene, 1922: (descr., fig. 69 on pl. 15);

Caesar, 1916a: (unnumbered fig. on p. 172).

BIOLOGY: Schaffner and Griswold, 1934: (120).

Blondelia hyphantriae (Tothill)

Most records as *Anetia hyphantriae*.

HOST: LEPIDOPTERA

Arctiidae

Diacrisia virginica (Fabricius)

Schaffner and Griswold, 1934: 36, 120.

Halisidota harrisii Walsh

Raizenne, 1952: 25, Ont. (as *Phrynolydella*); Schaffner and Griswold, 1934: 34-35, 120.

Halisidota tessellaris (J. E. Smith)

Raizenne, 1952: 24, Ont. (as *Phrynolydella*); Schaffner and Griswold, 1934: 34, 120.

Hyphantria cunea (Drury)

Schaffner and Griswold, 1934: 37, 120; Tothill, 1922: 29-31, 43-48, 107, figs. 9-15, B.C. (as *Lydella*).

Hyphantria textor (Harris)

Raizenne, 1952: 29, Ont. (as *Phrynolydella*).

Noctuidae

Bomolocha abalienalis (Walker)

Raizenne, 1952: 75, Ont. (as *Phrynolydella*).

Bomolocha deceptalis (Walker)

Raizenne, 1952: 76, Ont. (as *Phrynolydella*).

Ceramica picta (Harris)

Schaffner and Griswold, 1934: 47, 120.

Lithophane antennata (Walker)

Schaffner and Griswold, 1934: 49, 120 (ex *Graptolitha*).

Lithophane spp. (including *L. disposita* Morrison, *L. innominata* (Smith) and/or *L. petulca* Grote)

Schaffner and Griswold, 1934: 48, 120 (ex *Graptolitha* spp. (including *G. disposita*, *G. innominata* and/or *G. petulca*)).

Notodontidae

Heterocampa guttivitta (Walker)

Raizenne, 1952: 88, Ont. (as *Phrynolydella*).

STAGE PARASITIZED: Larva, pupa.

IMMATURES: First instar larva — Tothill, 1922: (descr., figs. 9-12).

Second instar larva — Tothill, 1922: (descr., figs. 13-15).

BIOLOGY: Schaffner and Griswold, 1934: (120); Tothill, 1922: (29-31, 43-48, 107).

Blondelia polita (Townsend)

All records as *Sturmia strigata* Wulp.

HOST: LEPIDOPTERA

Pyralidae

Loxostege mancalis (Lederer)

Fattig, 1949: 32; Brimley, 1938: 364; Brimley, 1922: 25,
N.C.

Blondelia sp. near *polita* (Townsend)

HOST: LEPIDOPTERA

Tortricidae

Archips infumatanus (Zeller)

Schaffner, 1959: 29, 54, Mass. (as *Sturmia* sp. near *strigata*
Wulp).

BIOLOGY: Schaffner, 1959: (29,54).

Blondelia spp.

HOST: LEPIDOPTERA

Geometridae

Anacamptodes clivinaria profanata (Barnes and McDunnough)

Furniss, M. M., and Barr, 1967: 9, Idaho (as *Blondelia* (sp.
n.?).)

Coryphista meadii (Packard)

Neiswander, 1941: 388, Ohio (as *Phrynolydella*).

Tortricidae

Archips spp.

*Procter, 1946: 431, Maine (Procter, 1938); Procter, 1938:
372, Maine.

STAGE PARASITIZED: Pupa.

Blondelia spp. probably

HOST: HYMENOPTERA

Diprionidae

Diprion hercyniae (Hartig)

Briand, 1949: 56-57, Que. (as *Lydella* sp. ex [European]
spruce sawfly).

HOST: LEPIDOPTERA

Geometridae

Cingilia catenaria (Cramer)

Schaffner and Griswold, 1934: 98, 120, Mass., N.H., N.Y.,
R.I. (as *Anetia* sp. (complex)).

STAGE PARASITIZED: Larva (Hymenoptera).

BIOLOGY: Schaffner and Griswold, 1934: (98, 120).

Bombyliomyia abrupta, see *Bombyliopsis abrupta*

Bombyliopsis Townsend (Tachininae: Macromyini)

Bombyliopsis abrupta (Wiedemann)

Most records as *Bombyliomyia abrupta*.

HOST: LEPIDOPTERA

Arctiidae

Diacrisia virginica (Fabricius)

Guppy, R., 1950: 4, B.C. (as *Bombyliopsis*); Schaffner and
Griswold, 1934: 36, 125.

Euchaetias egle (Drury)

Schaffner and Griswold, 1934: 39, 125.

Halisidota caryae (Harris)

Schaffner and Griswold, 1934: 33, 125; Townsend, 1892:
88, N.Y. (as *Hystricia*, probable host).

Halisidota maculata (Harris)

Schaffner and Griswold, 1934: 33-34, 125; Hine, 1904: 87,
B.C.

Halisidota tessellaris (J. E. Smith)

Schaffner and Griswold, 1934: 34, 125.

Hyphantria cunea (Drury)

Schaffner and Griswold, 1934: 37, 125.

Unidentified Arctiidae

Schaffner and Griswold, 1934: 40-41, 125.

STAGE PARASITIZED: Larva, pupa.

BIOLOGY: Schaffner and Griswold, 1934: (125).

Bonellimya tessellata, see *Linnaemya tessellata*

"Bonnetia" comta, see *Bonnetia comta*

Bonnetia Robineau-Desvoidy (Tachininae: Linnaemyini)

Bonnetia "comia," see *Bonnetia comta*

Bonnetia "compta," see *Bonnetia comta*

Bonnetia comta (Fallén)

HOST: COLEOPTERA

Scarabaeidae

Phyllophaga sp.

Brooks, 1944: 197.

HOST: LEPIDOPTERA

Noctuidae

Agrotis gladiaria Morrison

Fattig, 1949: 9 (ex *Feltia*).

Agrotis ipsilon (Hufnagel)

All records ex *Agrotis ypsilon*.

Crumb, 1929: 40, 59 (as "*Bonnettia compta*"); Essig, 1926: 580, Calif. (as "*Bonneta*"); Johnson, 1925: 194, Conn., Maine, Mass., N.H., R.I., Vt.; Coquillett, 1897: 18, 23, Calif. (as *Linnaemyia*).

Agrotis malefida Guenée

All records ex *Feltia malefida*.

Chapman, Fife and Cavitt, 1935: 305, Tex.; Crumb, 1929: 40 (as "*Bonnettia compta*").

Agrotis orthogonia Morrison

Most records ex *Porosagrotis orthogonia*.

Thornley and Sorenson, 1940: 48, Utah (ex *Agrotis*, probable host); Seamans and McMillan, 1935: 424 (ex *Agrotis*); Cook, W. C., 1930: 31, Mont.; Cook, W. C., 1923: 327, Mont. (as *B. "compta"*); Seamans, 1923: 53, Alta.; Strickland, 1923: 29-37, figs. 16-19, Alta.; Parker, J. R., Strand and Seamans, 1921: 313, Mont. (as *B. "compta"*); Strickland, 1921: 98-99, Alta. (as "*Bonneta*").

Agrotis venerabilis Walker

Crumb, 1929: 40 (as "*Bonnettia compta*" ex *Feltia*).

Copablepharon viridisparsa Dod

Strickland, 1923: 35, 37, figs. 16-19, Alta.

Euxoa auxiliaris (Grote)

Cook, W. C., 1930: 31, Mont. (ex *Chorizagrotis*).

Euxoa messoria (Harris)

Crumb, 1929: 40, 94 (as "*Bonnettia compta*"); Essig, 1926: 580, Wash. (as "*Bonneta*"); Johnson, 1925: 194, Conn., Maine, Mass., N.H., R.I., Vt. (ex *Carneades*); Coquillett, 1897: 18, 24, Wash. (as *Linnaemyia* ex *Carneades*).

Euxoa messoria (Harris) probably

*Vide Brauer and Bergenstamm, 1895: 562, 581, N. Amer. (as *Micropalpus fulgens* (Meigen) ex *Agrotis cochrani* (Riley), Townsend, 1893); Townsend, 1893: 467, Ill. (as *Micropalpus fulgens* ex cutworms).

Euxoa ochrogaster (Guenée)

King, K.M., and Atkinson, 1928: 170, 176, Sask.

Euxoa tristicula (Morrison)

Cook, W. C., 1930: 31, Mont.; Strickland, 1923: 36, figs. 16-20, Mont.

Euxoa spp. (including *E. messoria* (Harris) and/or *E. ochrogaster* (Guenée))

*Vide pers. comm. Handford, 1970: [B.C.] (Handford, 1962); Anon., 1962: 121, B.C. (as *B. "compta"*); Handford, 1962: 19, B.C. (as *B. "compta"*).

Feltia subterranea (Fabricius)

Patton, 1958: 34, Fla. (incomplete rearing — pupa); Tissot, 1945: 64, Fla. (as *B. "comia"*); Crumb, 1929: 40, 81, Va. (as "*Bonnettia compta*" ex *F. annexa* Treitschke); Jones, T. H., 1918a: 12, La. (as *Linnaemyia* ex *F. annexa*).

Peridroma saucia (Hübner)

*Crumb, 1929: 40, 115 (as "*Bonnettia compta*" ex *Lycophotia*, Wakeland, 1920); Allen, 1926a: 283-286, fig. 1, Miss. (as *Linnaemyia* ex "varigated" cutworms); Essig, 1926: 580, Colo. (as "*Bonneta*" ex *Lycophotia margaritosa* (Haworth)); Wakeland, 1920: 18, Colo. (as *Linnaemyia*).

Polia acutermis (Smith)

Strickland, 1923: 33, 37, Alta.

Spodoptera frugiperda (J. E. Smith)

Aldrich, 1915: 83 (as *Linnaemyia* ex *Laphygma*).

Armyworm(-s)

Allen, 1926a: 283-286, fig. 1, Miss. (as *Linnaemyia*).

Cutworm(-s)

Strickland, 1938: 216, Alta. (as *Linnaemyia "compta"*); Knowlton and Harmston, 1937: 148, Utah; Aldrich, 1915: 83 (as *Linnaemyia*).

Unidentified Noctuidae

Schaffner and Griswold, 1934: 72, 109, Maine (as *Linnaemyia*); Knowlton and Janes, 1931: 147, Utah (as "*Bonneta*"); Essig, 1926: 580, U.S. (as "*Bonneta*").

STAGE PARASITIZED: Larva, pupa.

IMMATURES: Egg — Crumb, 1929: (descr.); Strickland, 1923: (descr., fig. 16).

First instar larva — Allen, 1926a: (descr., fig. 1); Strickland, 1923: (descr., fig. 17).

Second instar larva — Strickland, 1923: (descr., fig. 18).

Third instar larva — Strickland, 1923: (descr., fig. 19).

Pupa — Crumb, 1929: (descr.); Strickland, 1923: (descr., figs. 19-20); Greene, 1922: (descr., fig. 70 on pl. 15).

BIOLOGY: Seamans and McMillan, 1935: (424); Schaffner and Griswold, 1934: (72, 109); Allen, 1926a: (283-286); Strickland, 1923: (29-37).

Bonnetia spp.

HOST: LEPIDOPTERA

Aegeriidae

Sylvora acerni (Clemens)

Brauer and Bergenstamm, 1895: 562, 605, N. Amer. (as
Micropalpus ex Sesia).

"*Bonnetia compta*," see *Bonnetia compta*

Bucentes ceres, see *Siphona ceres*

Bucentes geniculata, see *Siphona geniculata*

Buquetia Robineau-Desvoidy (Goniinae: Eryciini)

Buquetia musca Robineau-Desvoidy

HOST: LEPIDOPTERA

Papilionidae

Papilio machaon Linnaeus

Curran, 1927b: 149 (as *Eipogona setifacies* Rondani).

Buquetia obscura (Coquillett)

Most records as *Eupogona obscura*.

HOST: LEPIDOPTERA

Danaiidae

Danaus plexippus plexippus (Linnaeus)

Schaffner and Griswold, 1934: 8, 124, Mass. (ex *D. menippe*
(Hübner)); Webber and Schaffner, 1926: 23, New
England (as *Winthemia ex Anosia plexippus*).

Papilionidae

Papilio polyxenes asterius Stoll

Schaffner and Griswold, 1934: 6, 124, Mass. (ex *P. ajax*
Linnaeus); Curran, 1927b: 149, Iowa (as *Eipogona*
americana Curran ex *P. "asterias"*); Aldrich, 1928b: 44,
Mass. (ex *P. polyxenes* Fabricius).

STAGE PARASITIZED: Larva.

BIOLOGY: Schaffner and Griswold, 1934: (124).

"*Caetogaedia*" *monticola*, see *Chaetogaedia monticola*

Calodexia Wulp (Dexiinae: Calodexiini)

Calodexia sp. near *agilis* Curran or *interrupta* Curran

HOST: ORTHOPTERA

Gryllidae

Eneoptera spp.

Rettenmeyer, 1961: 1027, Panama.

Calodexia dives Curran

HOST: ORTHOPTERA

Blattidae

Cockroach(-es)

Rettenmeyer, 1961: 1027-1028, Panama.

STAGE PARASITIZED: Nymph.

Calodexia fumosa (Townsend)

HOST: ORTHOPTERA

Tettigoniidae

Unidentified Tettigoniidae

Rettenmeyer, 1961: 1028-1029, Panama (unsuccessful lab. rearing).

Calodexia interrupta Curran

HOST: ORTHOPTERA

Gryllidae

Ponca venosa Hebard

Rettenmeyer, 1961: 1026-1027, 1049, Panama.

BIOLOGY: Rettenmeyer, 1961: (1026-1027, 1049).

Calodexia venteris Curran

HOST: ORTHOPTERA

Blattidae

Eublaberus posticus (Erickson)

Rettenmeyer, 1961: 1028, 1059, Panama (lab.).

Calodexia spp.

HOST: ORTHOPTERA

Blattidae

Epilampra azteca Saussure

Rettenmeyer, 1961: 1027, Panama.

Cockroach(-es)

Rettenmeyer, 1961: 1027, Panama; Rettenmeyer, 1961a: 15, Panama.

Gryllidae

Cricket(-s)

Rettenmeyer, 1961a: 15, Panama.

BIOLOGY: Rettenmeyer, 1961a: (15).

Cane borer parasite, see *Lixophaga sphenophori*

Cane borer tachinid, see *Lixophaga sphenophori*

Carcelia Robineau-Desvoidy (Goniinae: Carceliini)

Carcelia amplexa (Coquillett)

HOST: LEPIDOPTERA

Arctiidae

Ecpantheria eridanus (Cramer)

*Sellers, 1943: 65 (Wolcott, 1924); Wolcott, 1924: 222, P.R.
(as *Exorista*).

Ecpantheria icasia (Cramer)

Wolcott, 1951a: 479, P.R.; Martorell, 1945a: 503, P.R.

Lymantriidae

Euproctis chrysorrhoea (Linnaeus)

Howard and Fiske, 1911: 93 (as *Exorista griseomicans* Wulp).

Hemerocampa leucostigma (J. E. Smith)

Raizenne, 1952: 95, Ont.; Sellers, 1943: 65; Dowden, 1935: 509 (as *Zenillia*); Schaffner and Griswold, 1934: 85, 111 (as *Zenillia*); Webber and Schaffner, 1926: 12-13, Mass., New England, N.Y., Pa. (as *Zenillia*); Britton, 1917: 110 (as *Exorista griseomicans*); Johnson, 1910: 780, N.J. (as *Exorista griseomicans* ex *Orgyia*); Townsend, 1909: 248, D.C., Mass. (as *Sisyropa hemerocampae* Townsend); Howard, 1898: 88, D.C. (as *Exorista griseomicans* ex *Orgyia*); Coquillett, 1897: 14, 26, D.C. (as *Exorista griseomicans* ex *Orgyia*); Howard, 1897a: 16, D.C. (as *Exorista griseomicans* ex *Orgyia*); Howard, 1897b: 7, 44, 49, D.C. (as *Exorista griseomicans* ex *Orgyia*).

Orgyia antiqua (Linnaeus)

All records ex *Notolophus antiqua*.

Sellers, 1943: 65; Schaffner and Griswold, 1934: 84, 111
(as *Zenillia*).

Porthetria dispar (Linnaeus)

Sellers, 1943: 65.

Tussock moth(-s)

Thompson, 1910: 291 (as *Exorista*).

Megalopygidae

Megalopyge costaricensis Schaus

Ballou, 1936: 142, Costa Rica (as *Zenillia griseomicans*);

Ballou, 1936a: 48, Costa Rica (as *Zenillia griseomicans*).

Megalopyge crispata (Packard)

Brimley, 1922: 22, N.C. (as *Exorista griseomicans* ex *Lagoa*).

Megalopyge opercularis (J. E. Smith)

Patton, 1958: 34, Fla.; Fattig, 1949: 10, Ga.; Brimley, 1938:

362, N.C. (as *Zenillia*); Allen, 1929: 686, Miss. (as *Zenillia*).

Noctuidae

Plathypena scabra (Fabricius)

*Sellers, 1943: 65 (Hill, 1925); Hill, 1925: 17, Md. (as

Exorista).

Spodoptera frugiperda (J.E. Smith)

Brimley, 1922: 22, N.C. (as *Exorista griseomicans* ex fall army worm).

Sphingidae

Manduca sexta jamaicensis (Butler)

Gilmore, 1938: 711, Ky. and/or Tenn. (as *Zenillia* ex

Protoparce sexta (Johanssen)).

STAGE PARASITIZED: Larva, pupa.

IMMATURES: Pupa — Greene, 1922: (descr., fig. 61 on pl. 13 and fig. 84 on pl. 18).

BIOLOGY: Sellers, 1943: (65); Dowden, 1935: (509); Schaffner and Griswold, 1934: (111); Thompson, 1910: (291).

Carcelia diacrisiae Sellers

HOST: LEPIDOPTERA

Arctiidae

Diacrisia virginica (Fabricius)

Vide Sellers, 1943: 63, N.J. (Schaffner and Griswold, 1934);

Schaffner and Griswold, 1934: 36, 113-114 (as *Zenillia* n. sp. (4)).

Estigmene acrea (Drury)

Patton, 1958: 34, Fla.; vide Sellers, 1943: 63 (ex *E. acrea*,

Schaffner and Griswold, 1934); Schaffner and Griswold, 1934: 38, 113-114 (as *Zenillia* n. sp. (4)).

STAGE PARASITIZED: Larva.

BIOLOGY: Sellers, 1943: (63).

Carcelia flavirostris, see *Carcelia lagoae*

Carcelia formosa (Aldrich and Webber)

Most records as *Zenillia formosa*.

HOST: LEPIDOPTERA

Noctuidae

Acronicta hamamelis Guenée

*Vide Sellers, 1943: 44 (as *Carcelia*, Coquillett, 1897);
Coquillett, 1897: 13, 23, Va. (as *Exorista eudryae*
Townsend).

Agrotis ipsilon (Hufnagel)

All records ex *Agrotis ypsilon*.

Vide Sellers, 1943: 44, Mo. (as *Carcelia*, Coquillett,
1897); *Crumb, 1929: 40, 59 (Aldrich and Webber,
1924); Johnson, 1925: 196, Conn., Mass.; Aldrich
and Webber, 1924: 24, Mo.; Coquillett, 1897: 13, 23
(as *Exorista eudryae*).

Lithophane spp. (including *L. disposita* Morrison and/or *L.*
innominata (Smith))

Sellers, 1943: 44 (as *Carcelia* ex *Graptolitha* (including *G.*
disposita and/or *G. innominata*)).

Saturniidae

Automeris io (Fabricius)

Patton, 1958: 34, Fla. (as *Carcelia*); Hammond, 1949a:
236, Ont. (as "*Cercelia*"); Sellers, 1943: 44, Mass. (as
Carcelia); Schaffner and Griswold, 1934: 28, 112,
Conn., Mass., N.J., N.Y., Pa., R.I.; Johnson, 1925:
196, Conn., Mass.; Aldrich and Webber, 1924: 24,
Mass., Md.

STAGE PARASITIZED: Larva.

BIOLOGY: Sellers, 1943: (44); Schaffner and Griswold, 1934:
(28, 112).

Carcelia illota (Curran)

HOST: LEPIDOPTERA

Noctuidae

Heliothis virescens (Fabricius)

Bryan, D. E., Jackson and Stoner, 1969: 1, 8, 12, Ariz. (lab.).

Heliothis zea (Boddie)

Bryan, D. E., Jackson and Stoner, 1969: 1, 8, 12, Ariz. (lab.).

STAGE PARASITIZED: Larva, pupa.

BIOLOGY: Bryan, D. E., Jackson and Stoner, 1969: (1, 8, 12).

Carcelia lagoae (Townsend)

HOST: LEPIDOPTERA

Arctiidae

Ecpantheria icasia (Cramer)

Wolcott, 1951a: 479, P.R. (as *C. flavirostris* (Wulp)).

Halisidota spp.

All records ex red "*Halesidota*."

Johnson, 1919: 437, Jamaica; Townsend, 1893c: 314, Jamaica (as *Exorista*).

Megalopygidae

Lagoa sp.

Sellers, 1943: 59.

Megalopyge crispata (Packard)

Brimley, 1922: 22, N.C. (as *Exorista flavirostris* ex *Lagoa*).

Megalopyge krugii (Dewitz)

Sellers, 1943: 61 (as *C. flavirostris*).

Megalopyge opercularis (J. E. Smith)

Patton, 1958: 34, Fla.; Patton, 1958: 34, Fla. (as *C. flavirostris*); Micks, 1956: 38-39, fig. 3, Tex.; Sellers, 1943: 59; Bishopp, 1923: 13, Tex. (as *Exorista flavirostris*); Reinhard, 1922: 72, Tex. (as *Exorista flavirostris*); Coquillett, 1897: 14, 25, Fla., Miss., Tex., Va. (as *Exorista flavirostris* ex *Lagoa*); *Brauer and Bergenstamm, 1895: 574, 595, N. Mex. (as *Sisyropa* ex *Lagoa*, Townsend, 1891d); Townsend, 1891d: 159, Mexico (as *Exorista* ex *Lagoa*).

Megalopyge opercularis (J. E. Smith) probably

Sellers, 1943: 59.

Pyralidae

Eulepte concordalis Hübner probably

*Vide Wolcott, 1951a: 479, P.R. (as *C. flavirostris* ex presumably *Mesocondyla*, Sellers, 1943); Martorell, 1945a: 515, P.R. (as *C. flavirostris*); Sellers, 1943: 61 (as *C. flavirostris* ex larva on calabash tree).

Omphalocera cariosa Lederer

Coquillett, 1897: 14, 26 (as *Exorista flavirostris*).

STAGE PARASITIZED: Larva, pupa.

IMMATURES: Pupa — Micks, 1956: (fig. 3).

BIOLOGY: Sellers, 1943: (60); Bishopp, 1923: (13).

Carcelia laxiformis Villeneuve

HOST: LEPIDOPTERA

Arctiidae

Apantesis proxima (Guérin-Méneville)

Coquillett, 1897: 13, 24, Calif. (as *Exorista cheloniae*
Rondani ex *Arctia docta* (Walker)).

Unidentified Arctiidae

Ingles, 1929: 60, Calif. (as *Exorista cheloniae*).

Lasiocampidae

Malacosoma americanum (Fabricius)

Sellers, 1943: 55-56; Howard and Fiske, 1911: 299 (as
Parexorista cheloniae).

Malacosoma disstria Hübner

Sellers, 1943: 55-56; Tothill, 1913: 70, N.B., Ont. (as
Exorista cheloniae); Howard and Fiske, 1911: 299 (as
Parexorista cheloniae).

Lymantriidae

Euproctis chrysorrhoea (Linnaeus)

Burgess, 1924: 34, Mass. (as *Parexorista cheloniae*); Howard
and Fiske, 1911: 297-300, 310, fig. 4 on pl. 20, probably
Mass. (as *Parexorista cheloniae*).

Nygmia phaeorrhoea (Donovan)

Dowden, 1962: 29, 52, New England; Clausen, 1956: 6, 64,
89; Sellers, 1943: 55-56; Dowden, 1935: 509; Proper,
1934: 373-374, New England (ex brown-tail moth);
Collins and Jones, 1932: 237 (ex brown-tail moth);
Burgess and Crossman, 1929: 134-138, 142, Maine,
Mass., N.H.

Porthetria dispar (Linnaeus)

Sellers, 1943: 55; Burgess, 1924: 34, Mass. (as *Parexorista*
cheloniae).

Notodontidae

Schizura concinna (J. E. Smith)

Hitchings, 1908: 11, Maine (as *Exorista* "chelonia" ex red-humped caterpillars).

STAGE PARASITIZED: Larva.

IMMATURES: Pupa — Howard and Fiske, 1911: (fig. 4 on pl. 20).

BIOLOGY: Dowden, 1962: (29, 52); Clausen, 1956: (6, 64, 89); Sellers, 1943: (55-56); Dowden, 1935: (509); Proper, 1934: (373-374); Burgess and Crossman, 1929: (134-138, 142); Howard and Fiske, 1911: (297-300, 310).

Carcelia leucanae, see *Winthemia leucanae*

Carcelia "leucaniae," see *Winthemia leucanae*

Carcelia malacosomae Sellers

HOST: HYMENOPTERA

Tenthredinidae

Unidentified Tenthredinidae

Sellers, 1943: 52.

HOST: LEPIDOPTERA

Arctiidae

Hyphantria cunea (Drury)

Sellers, 1943: 52.

Turuptiana permaculata (Packard)

Sellers, 1943: 52.

Lasiocampidae

Malacosoma americanum (Fabricius)

Sellers, 1943: 52-53.

Malacosoma disstria Hübner

Sippell, 1961: 435-436, fig. 8, Ont.; Cumming, 1954: 183, Alta.; Raizenne, 1952: 103, Ont.; Sellers, 1943: 52-53.

Malacosoma spp.

Ross, 1953: 19, 21, fig. 4, B.C.

Notodontidae

Datana integerrima Grote and Robinson

*Vide Sellers, 1943: 52 (Schaffner and Griswold, 1934);
Schaffner and Griswold, 1934: 76, 112 (as *Zenillia*
protuberans Aldrich and Webber).

Saturniidae

Hemileuca spp. (including *H. lucina* H. Edwards and/or *H. maia*
(Drury))

Sellers, 1943: 52.

IMMATURES: Pupa — Sippell, 1961: (descr., fig. 8); Ross,
1953: (descr., fig. 4).

BIOLOGY: Sellers, 1943: (54).

Carcelia malacosomae Sellers probably

Most records as *Zenillia cheloniae* (Rondani).

HOST: LEPIDOPTERA

Arctiidae

Apantesis ornata (Packard)

*Vide Sellers, 1943: 52 (as *Carcelia malacosomae* probably
ex *Callarctia*, Tothill, 1913); Essig, 1926: 581, B.C.;
Tothill, 1913: 70, B.C. (as *Exorista* ex *A. ornata* var.
achaia (Grote and Robinson)).

Apantesis proxima (Guérin-Méneville)

Sellers, 1943: 52 (as *Carcelia malacosomae* probably ex
Callarctia).

Arachnis picta Packard

*Vide Sellers, 1943: 52 (as *Carcelia malacosomae* probably,
Coquillett, 1897); Essig, 1926: 581, Calif.; Coquillett,
1897: 13, 24, Calif. (as *Exorista*).

Turuptiana permaculata (Packard)

Essig, 1926: 581, Colo.; Aldrich and Webber, 1924: 32, Colo.

Lasiocampidae

Malacosoma americanum (Fabricius)

Burgess and Crossman, 1929: 136; Essig, 1926: 581, Ariz.,
Utah; Aldrich and Webber, 1924: 10, 32, probably Mass.

Malacosoma disstria Hübner

Burgess and Crossman, 1929: 136.

American tent caterpillar(-s)

Johnson, 1925: 197, Conn., Mass., N.H.

Tortricidae

Archips argyrospilus (Walker)

*Vide Sellers, 1943: 52 (Gill, 1913); Aldrich and Webber, 1924: 32, Colo. (ex *Cacoecia*); Gill, 1913: 102, Colo. or N. Mex. (as *E[xorista]*).

STAGE PARASITIZED: Larva, pupa.

Carcelia "nigropalpus," see *Aplomya caesar* probably

"*Carcelia ochracea* Wulp"

HOST: LEPIDOPTERA

Pyralidae

Ostrinia nubilalis (Hübner)

Bradley, W.G., and Clark, 1941: 792, E. U.S. (ex European corn borer).

STAGE PARASITIZED: Larva.

Carcelia olenensis Sellers

HOST: LEPIDOPTERA

Lymantriidae

Dasychira atomeria Walker

All records ex *Olene* "*atomaria*."

Vide Sellers, 1943: 68-69, Mass. (Schaffner and Griswold, 1934); Schaffner and Griswold, 1934: 86, 113 (as *Zenillia* n. sp. (2)).

Dasychira plagiata (Walker)

Raizenne, 1952: 99, Que. (ex *Olene*).

BIOLOGY: Sellers, 1943: (68-69).

Carcelia perplexa Sellers

HOST: LEPIDOPTERA

Lymantriidae

Dasychira plagiata (Walker)

Vide Sellers, 1943: 67 (ex *Olene*, Schaffner and Griswold, 1934); Schaffner and Griswold, 1934: 87, 113 (as *Zenillia* n. sp. (2) ex *Olene pini* (Dyar)).

Hemerocampa leucostigma (J. E. Smith)

Sellers, 1943: 67.

BIOLOGY: Sellers, 1943: (67).

Carcelia protuberans (Aldrich and Webber)

Most records as *Zenillia protuberans*.

HOST: LEPIDOPTERA

Arctiidae

Hyphantria cunea (Drury)

Schaffner and Griswold, 1934: 37, 112.

Lasiocampidae

Malacosoma americanum (Fabricius)

Schaffner and Griswold, 1934: 88, 112.

Malacosoma disstria Hübner

Sellers, 1943: 54 (as *Carcelia*); Bess, 1936: 605; Schaffner and Griswold, 1934: 89, 112; Johnson, 1925: 196, Vt.; Aldrich and Webber, 1924: 15, Vt.

BIOLOGY: Schaffner and Griswold, 1934: (112).

Carcelia pyste, see *Nemorilla pyste*

Carcelia reclinata (Aldrich and Webber)

Most records as *Zenillia reclinata*.

HOST: LEPIDOPTERA

Arctiidae

Diacrisia virginica (Fabricius)

Sellers, 1943: 49 (as *Carcelia*); Walker, 1936: 19, Kans. (as *Z. "rectinata"*); Schaffner and Griswold, 1934: 36, 112-113.

Ecpantheria deflorata (Fabricius)

Patton, 1958: 34-35, Fla. (as *Carcelia*).

Estigmene acraea (Drury)

Patton, 1958: 34-35, Fla. (as *Carcelia*); Fattig, 1949: 10 (as *Carcelia*); Sellers, 1943: 49-50 (as *Carcelia ex E. acraea*); Schaffner and Griswold, 1934: 38, 112-113; Essig, 1926: 581, Ariz., N. Mex. (ex *E. acraea*); Aldrich and Webber, 1924: 33, New England (ex *E. acraea*).

Phragmatobia assimilans Walker

All records ex *Phragmatobia assimilans* var. *franconia* Slosson.

Sellers, 1943: 49-50, Ont. (as *Carcelia*); Aldrich and Webber, 1924: 33, Ont. (ex *P. assimilans* var. "*franconica*").

Unidentified Arctiidae

Sellers, 1943: 49-50 (as *Carcelia*); Schaffner and Griswold, 1934: 40-41, 112-113.

Unidentified Lepidoptera

Miscellaneous

Knowlton and Harmston, 1937: 149, Utah; Essig, 1926: 581, E. U.S.

STAGE PARASITIZED: Larva, pupa.

BIOLOGY: Patton, 1958: (34-35); Sellers, 1943: (50); Schaffner and Griswold, 1934: (112).

Carcelia reclinata (Aldrich and Webber) probably

HOST: LEPIDOPTERA

Arctiidae

Estigmene acrea (Drury)

*Vide Sellers, 1943: 101, N.C. (Brimley, 1922); Brimley, 1922: 22, N.C. (as *Exorista eudryae* Townsend).

Phragmatobia assimilans Walker

Most records ex *Phragmatobia assimilans* var. *franconia* Slosson.

*Vide Sellers, 1943: 52 (ex *P. assimilans*, Gibson, 1911; Gibson, 1911a); Gibson, 1911: 127, Ont. (as *Exorista cheloniae* Rondani); Gibson, 1911a: 117, Ont. (as *Exorista cheloniae*).

STAGE PARASITIZED: Larva, pupa.

Carcelia yalensis Sellers

HOST: LEPIDOPTERA

Arctiidae

Halisidota tessellaris (J. E. Smith)

Sellers, 1943: 70.

Lymantriidae

Hemerocampa leucostigma (J. E. Smith)

Vide Sellers, 1943: 70 (in part Schaffner and Griswold, 1934); Schaffner and Griswold, 1934: 85, 113 (as *Zenillia* n. sp. (2)).

Orgyia antiqua (Linnaeus)

All records ex *Notolophus antiqua*.

Raizenne, 1952: 93, Ont.; vide Sellers, 1943: 70 (in part Schaffner and Griswold, 1934); Schaffner and Griswold, 1934: 84, 113 (as *Zenillia* n. sp. (2)).

Unidentified Lepidoptera

Miscellaneous

Sellers, 1943: 70 (ex cocoon of caterpillar on *Abies concolor* (Gordon and Glendinning) Lindley).

Carcelia spp.

Most records as *Chaetolyga* sp.

HOST: LEPIDOPTERA

Arctiidae

Ecpantheria icasia (Cramer)

Harrison and Stephens, 1966: 673-674, Costa Rica (as *Carcelia*).

Unidentified Arctiidae

Brauer and Bergenstamm, 1895: 568, 583, N. Amer. (as *Parexorista*).

Limacodidae

Adoneta spinuloides Herrich-Schaeffer

Brauer and Bergenstamm, 1895: 547, 581, N. Amer.

Phobetron pithecium (J. E. Smith)

Patton, 1958: 35, Fla. (as *Carcelia* ex hag-moth).

Lymantriidae

Dasychira plagiata (Walker)

Walgenbach and Benjamin, 1963: 15, Wis. (as *Carcelia*).

Noctuidae

Alypia octomaculata (Fabricius)

Brauer and Bergenstamm, 1895: 547, 582.

Peridroma saucia (Hübner)

Brauer and Bergenstamm, 1895: 547, 581 (ex *Agrotis inermis* (Harris)).

Pseudaletia unipuncta (Haworth)

Brauer and Bergenstamm, 1895: 547, 596 (ex *Leucania*).

Notodontidae

Datana integerrima Grote and Robinson

Brauer and Bergenstamm, 1895: 547, 590 (ex *D. ministra* (Drury) schwarze Var. auf Wallnüssen).

Olethreutidae

Melissopus latiferreanus (Walsingham)

*Vide Murtfeldt, 1894a: 322, Mo. (as small tachinid, Murtfeldt, 1892); Murtfeldt, 1892: 135, Mo. (as *Parexorista* sp.? ex acorn *Carpocapsa* [sp.]).

Pyralidae

Pantographa limata Grote and Robinson

Murtfeldt, 1894: 45, Mo. (as *Parexorista*).

Sphingidae

Manduca sexta (Linnaeus)

Alwood, 1898: 74, Va. (ex *Protoparce carolina* (Linnaeus)).

STAGE PARASITIZED: Larva, pupa.

Carinosillus Reinhard (Proseninae: Theresiini)*Carinosillus novaeangliae* (West)

HOST: DIPTERA

Tabanidae

Hybomitra trispila sodalis (Williston) probably

James, H. G., 1963: 1228-1229, Ont. (as *Phorostoma*).

STAGE PARASITIZED: Larva.

BIOLOGY: James, H. G., 1963: (1228-1229).

Carinosillus tabanivorus (Hall)

HOST: DIPTERA

Tabanidae

Chrysops sp.

Hays, 1958: 552, Mich. (as *Phorostoma*).

Hybomitra epistates (Osten Sacken)

Teskey, 1969: 114-115.

Hybomitra metabola (McDunnough)

Teskey, 1969: 114-115.

Hybomitra nuda (McDunnough)

Teskey, 1969: 114-115, Ont.

Hybomitra trepida (McDunnough)

Teskey, 1969: 114-115, Ont.; Hays, 1958: 552-553, figs.

A-D, Mich. (as *Phorostoma*).

Hybomitra spp.

Hays, 1958: 552-553, figs. A-D, Mich. (as *Phorostoma*).

Tabanus fulvicallus Philip

Teskey, 1969: 114-115, Ont.

Tabanus marginalis Fabricius

Teskey, 1969: 114-115.

Tabanus reinwardtii Wiedemann

Teskey, 1969: 114-115.

Tabanus trimaculatus Palisot de Beauvois

Most records ex tabanid larva.

Greene, 1937: 217-218, fig. 27, Minn. (as *Myocera*); Hall, 1937: 207, Minn. (as *Myocera*); Philip, 1931: 68, Minn. (as *Phorostoma* n. sp. ex *T. trimaculatus*).

Horse fly (flies)

Sabrosky and Arnaud, 1965: 989 (as *Carinosillus* sp.).

STAGE PARASITIZED: Larva.

IMMATURES: Third instar larva — Hays, 1958: (descr., figs. A-D).

Pupa — Greene, 1937: (descr., fig. 27).

BIOLOGY: Teskey, 1969: (114-115).

Carinosillus sp., see *Carinosillus tabanivorus*

Cartocometes Aldrich (Goniinae: Hyperecteinini)

Cartocometes io Aldrich

HOST: LEPIDOPTERA

Lasiocampidae

Malacosoma americanum (Fabricius)

Bess, 1936: 605; Schaffner and Griswold, 1934: 88, 107.

Malacosoma disstria Hübner

Sippell, 1961: 436, Ont.; Raizenne, 1952: 103, Ont.; Bess, 1936: 605 (as "*Cartometes*"); Schaffner and Griswold, 1934: 89, 107.

BIOLOGY: Schaffner and Griswold, 1934: (107).

"*Cartometes*" *io*, see *Cartocometes io*

Catagoniopsis Townsend (Goniinae: Eryciini)

Catagoniopsis meracanthae (Greene)

All records as *Phorocera meracanthae*.

HOST: COLEOPTERA

Tenebrionidae

Meracantha contracta (Palisot de Beauvois)

Aldrich and Webber, 1924: 69, Md.; Greene, 1921a: 127, Md.

STAGE PARASITIZED: Larva.

IMMATURES: Pupa — Greene, 1922: (descr., fig. 3 on pl. 1).

Catagoniopsis sp. near *specularis* (Aldrich and Webber)

HOST: HYMENOPTERA

Diprionidae

Diprion polytomum (Hartig)

Reeks, 1938: 26, N.B. (as *Phorocera*).

STAGE PARASITIZED: Larva.

Catharosia Rondani (Phasiinae: Catharosiini)

Catharosia lustrans (Reinhard)

HOST: HEMIPTERA (HETEROPTERA)

Lygaeidae

Eremocoris opacus Van Duzee

Ashlock and O'Brien, 1964: 98-100, Calif.

Eremocoris sp.

Ashlock and O'Brien, 1964: 99-100, Calif. (lab.).

Thylochromus nitidulus Barber

Ashlock and O'Brien, 1964: 98-100, Calif.

STAGE PARASITIZED: Adult.

BIOLOGY: Ashlock and O'Brien, 1964: (98-100).

Celatoria Coquillett (Goniinae: Blondeliini)

Celatoria crawii, see *Celatoria diabroticae*

Celatoria "diabrotica," see *Celatoria diabroticae*

Celatoria diabroticae (Shimer)

HOST: COLEOPTERA

Chrysomelidae

Acalymma trivittata (Mannerheim)

All records ex *Diabrotica trivittata*.

Essig, 1926: 579; Craw, 1891: 27-28, fig. 31, Calif. (as *C. crawii* Coquillett).

Acalymma vittata (Fabricius)

All records ex *Diabrotica vittata*.

Fattig, 1949: 10, Ga. (as *C. "diabrotica"*); Bussart, 1937: 286, Ill.; Landis, 1935: 345, Ohio (probable parasite, incomplete rearing — pupa); Poos, 1928: 146, 148, Va.; Essig, 1926: 579; Balduf, 1925: 285, 305; Johnson, 1925: 187, Mass., N.H. (as *C. "diabrotica"*); Chittenden, 1923: 6; Britton, 1919: 35, Calif., Tex.;

Chittenden, 1919: 9; Lowry, Q. S., 1918: 267, Conn.; Girault, 1913: 55, Tex. (as *C. "diabrotica"*); Chittenden, 1898: 29, D.C. or Md.; Chittenden, 1898a: 79, D.C.; Coquillett, 1897: 11, 23, Md.; *Brauer and Bergenstamm, 1895: 562, 590 (as (?*Melanophora*), Riley, 1884); Riley, 1884: 77 (as *Melanophora?*); Shimer, 1871: 219 (as *Tachina* ex "*Diabrothica*").

Cerotoma trifurcata (Forster)

Fronk, 1950: 24 Va.; Eddy, C.O., and Nettles, 1930: 20, S.C.; Isely, 1930: 19, Ark.; Allen, 1929: 683, Miss.; McConnell, W.R., 1915: 264, [probably Miss.].

Diabrotica undecimpunctata Mannerheim

Essig, 1926: 579 (ex *D. duodecimpunctata* (Fabricius)); Essig, 1926: 579 (ex *D. soror* LeConte); Sweetman, 1926: 490 (as *C. "diabrotica"* ex *D. 12-punctata*); Johnson, 1925: 187, Mass., N.H. (as *C. "diabrotica"* ex *D. 12-punctata*); Currie, 1905: 46 (ex *D. duodecimpunctata*); Titus and Pratt, 1904: 78 (ex *D. "duodecempunctata"*); Chittenden, 1898: 30 (ex *D. 12-punctata*); Chittenden, 1898a: 79 (ex *D. 12-punctata*).

Diabrotica undecimpunctata howardi Barber

Most records ex *Diabrotica duodecimpunctata* (Fabricius).

Anon., 1961: 7 (ex *D. undecimpunctata howardi*); Summers and Stafford, 1953: 180, Miss. (ex *D. undecimpunctata howardi*); Fronk, 1950: 23-24, Va. (ex *D. undecimpunctata howardi*); Fattig, 1949: 10, Ga. (as *C. "diabrotica"*); Allen, 1929: 683, Miss. (ex *D. "duodecempunctata"*); Arant, 1929: 8-10, Ala.; Isely, 1929: 26, Ark.; Luginbill, 1918: 8-9; Coquillett, 1897: 10, 23, D.C. (ex *D. 12-punctata*).

Diabrotica undecimpunctata undecimpunctata Mannerheim

Most records ex *Diabrotica soror* LeConte.

Michelbacher, Middlekauff, Bacon and Swift, 1955: 13-14, fig. 13b (ex *D. undecimpunctata*); Michelbacher, MacLeod and Smith, 1943: 31, Calif. (ex *D. 11-punctata*); Rockwood and Chamberlin, 1943: 840, Calif., Oreg. (ex *D. 11-punctata*); Anon., 1931d: 459, Oreg.; Balduf, 1925: 305, Calif.; *Essig, 1915: 271, [Calif.] (as *C. "diabrotica,"* Coquillett, 1890); Sell, R. A., 1915: 518, Calif.; Essig, 1913: 229, 259, Calif.; Chittenden, 1910: 72, 75; Chittenden, 1898: 30, Calif. (as *C. crawii*); Coquillett, 1897: 10, 23, Calif.; *Brauer and Bergenstamm, 1895: 544, 590 (as "*Bessaria*")

crawii, Coquillett, 1890); Coquillett, 1895: 251, Calif. (as *C. crawii*); Craw, 1891: 27-28, fig. 31, Calif. (as *C. crawii* ex 12-spotted squash beetle); Coquillett, 1890: 234-236, Calif. (as *C. crawii*); *Koebele, 1890: 87, Calif. (as dipterous larva, Coquillett, 1890).

Diabrotica spp.

Carrillo S., Ortega C., and Gibson, 1966: 94, Mexico (V.C.); Gibson, W.W., and Carrillo S., 1959: 181, Mexico (V.C.); Sweetman, 1936: 220, N. Amer.

Trirhabda bacharidis (Weber)

Johnson, 1925a: 86, New England (as *C. "diabrotica"* ex *Galeruca "bachardis"*).

STAGE PARASITIZED: Larva, adult.

IMMATURES: Third instar larva — Arant, 1929: (descr.); Craw, 1891: (fig. 31).

Pupa — Michelbacher, Middlekauff, Bacon and Swift, 1955: (fig. 13b); Arant, 1929: (descr.); Greene, 1922: (descr., fig. 5 on pl. 1); Luginbill, 1918: (descr.); Craw, 1891: (descr., fig. 31); Coquillett, 1890: (descr.).

BIOLOGY: Michelbacher, Middlekauff, Bacon and Swift, 1955: (13-14); Michelbacher, MacLeod and Smith, 1943: (31); Sweetman, 1936: (220); Eddy, C.O., and Nettles, 1930: (20); Arant, 1929: (8-10); Luginbill, 1918: (8-9); Essig, 1913: (229, 259); Craw, 1891: (27-28); Coquillett, 1890: (234-236); Shimer, 1871: (219).

Celatoria setosa (Coquillett)

Most records as *Chaetophleps setosa*.

HOST: COLEOPTERA

Chrysomelidae

Acalymma vittata (Fabricius)

Most records ex *Diabrotica vittata*.

Gould, G. E., 1944: 11, Ind.; Gould, G. E., 1943: 61, Ind. (ex striped cucumber beetle); Bussart, 1937: 286-295, pl. 2, Ill.; Allen, 1929: 682, Miss.; Isely, 1927: 25-26, 33, Ark.; Balduf, 1925: 285-292, figs. 8-9, Ohio; Chittenden, 1923: 6; Chittenden, 1919: 9; Walton, 1914: 11-12, Md. (as *Neocelatoria ferox* Walton).

Diabrotica undecimpunctata Mannerheim

Bussart, 1937: 286-295, pl. 2, Ill. (ex *D. "duodecempunctata"* Fabricius).

Diabrotica undecimpunctata howardi Barber

Fattig, 1949: 11, Ga. (ex *D. duodecimpunctata*).

Cucumber beetle(-s)

Gould, G. E., 1936: 734, Ind.

STAGE PARASITIZED: Adult.

IMMATURES: Egg — Bussart, 1937: (descr., pl. 2); Balduf, 1925: (descr.).

First instar larva — Bussart, 1937: (descr., pl. 2); Balduf, 1925: (descr.).

Second instar larva — Bussart, 1937: (descr., pl. 2).

Third instar larva — Bussart, 1937: (descr., pl. 2); Balduf, 1925: (descr., fig. 8).

Pupa — Bussart, 1937: (descr., pl. 2); Balduf, 1925: (descr., fig. 9); Walton, 1914: (descr.).

BIOLOGY: Gould, G. E., 1944: (11); Bussart, 1937: (286-295); Isely, 1927: (25-26, 33); Balduf, 1925: (285-292).

Celatoria spinosa, see *Medina barbata*

Centeter cinerea, see *Hyperecteina aldrichi*

Centeter sp., see *Hyperecteina aldrichi*

Ceracia Rondani (Goniinae: Acemyini)

Ceracia dentata (Coquillett)

HOST: ORTHOPTERA

Acrididae

Camnula pellucida (Scudder)

Newton, 1954: 936.

Chimarocephala pacifica pacifica (Thomas)

Arnaud and Rentz, 1965: 204-206, fig. 1, Calif.

Chortophaga viridifasciata (De Geer)

Coquillett, 1897: 9, 23, Mo. (as *Acemyia*).

Chortophaga spp.

Townsend, 1912b: 164, N. Amer. (as *Acemyia*).

Melanoplus bivittatus (Say)

Newton, 1954: 936.

Melanoplus bruneri Scudder

Newton, 1954: 936.

Melanoplus confusus Scudder

Newton, 1954: 936.

Melanoplus differentialis (Thomas)

Morgan, 1901: 25, Miss. (as *Acemyia*).

Melanoplus femurrubrum (De Geer)

Smith, R. W., 1944: 29, Canada (ex *M. "f. rubrum"*).

Melanoplus packardii Scudder

Newton, 1954: 936, Mont.

Melanoplus sanguinipes (Fabricius)

Smith, 1965: 187-190, 194, 197, Ont.; Anon., 1963a: 8, Ont.; Beirne, 1962a: 19, Ont.; Smith, R. W., 1944: 29, Canada (ex *M. mexicanus* Saussure).

Melanoplus spp.

Boyes and Wilkes, 1953: 126, 142, Ont.; Townsend, 1912b: 164, N. Amer. (as *Acemyia*).

Grasshopper(-s)

Smith, R. W., and Finlayson, 1950: 98-101, figs. 26, 50, Canada; Strickland, 1946: 173, Alta.

STAGE PARASITIZED: Nymph, adult.

IMMATURES: Egg — Arnaud and Rentz, 1965: (fig. 1).

Second instar larva — Smith, R. W., and Finlayson, 1950: (descr., figs. 26, 50).

Pupa — Greene, 1922: (descr., fig. 50 on pl. 11).

BIOLOGY: Arnaud and Rentz, 1965: (204-206); Beirne, 1962a: (19); Anon., 1963a: (8); Smith, R. W., and Finlayson, 1950: (98-101).

Ceracia dentata (Coquillett) probably

HOST: ORTHOPTERA

Acrididae

Chortophaga viridifasciata (De Geer)

St. Amand and Cloyd, 1954: 83-87, figs. 1-14, Tenn.

Psoloessa delicatula (Scudder)

Newton, 1954: 936, N. Dak.

STAGE PARASITIZED: Nymph, adult.

IMMATURES: First instar larva — St. Amand and Cloyd, 1954: (descr., figs. 1-3).

Second instar larva — St. Amand and Cloyd, 1954: (descr., figs. 4-7).

Third instar larva — St. Amand and Cloyd, 1954: (descr., figs. 8-10, 14).

Pupa — St. Amand and Cloyd, 1954: (descr., figs. 11-13).

BIOLOGY: St. Amand and Cloyd, 1954: (83-87).

Ceracia spp.

HOST: ORTHOPTERA

Acrididae

Melanoplus foedus Scudder

Lavigne and Pfadt, 1966: 24, Wyo.

Melanoplus occidentalis (Thomas)

Lavigne and Pfadt, 1966: 24, Wyo.

Grasshopper(-s)

Smith, R. W., 1939: 61, Ont.

Ceratochaeta tortricis, see *Hemisturmia tortricis*

"*Cercelia*" *formosa*, see *Carcelia formosa*

Ceromasia Rondani (Goniinae: Eryciini)

Ceromasia auricaudata Townsend

HOST: LEPIDOPTERA

Arctiidae

Hyphantria cunea (Drury)

Coppel and Maw, 1954: 153, 155 (lab.); Coppel, 1947: 39, B.C.

Olethreutidae

Sciaphila duplex (Walsingham)

McGregor, 1967: 1215.

Pieridae

Pieris rapae (Linnaeus)

Coppel and Maw, 1954: 153-154 (lab.).

Pyralidae

Galleria mellonella (Linnaeus)

Coppel and Maw, 1954: 153 (lab.).

Ostrinia nubilalis (Hübner)

Coppel and Maw, 1954: 153 (ex *Pyrausta*, lab.).

Tortricidae

Archips cerasivoranus (Fitch)

Coppel and Maw, 1954: 153 (lab.).

Choristoneura fumiferana (Clemens)

Blais, 1960: 388, 391, Que. (ex spruce budworm); Coppel, 1960: 94, 97, B.C.; Carolin and Coulter, 1959: 552-554, Oreg.; Anon., 1954b: 147, B.C.; Coppel and Maw, 1954: 144-155, figs. 2-14, B.C.; Boyes and Wilkes, 1953: 126, 143, B.C.; Coppel, 1953: 191, B.C. (ex

"*Chloristoneura*"); MacNay, 1953a: 308, 313, 315-316, B.C.; Anon., 1952b: 175, 184-186, B.C.; Anon., 1952g: 285, 292, 294, B.C.; Ross, 1952: 112, fig. 15, Canada; Anon., 1950c: 233, 245, 248, B.C.; Anon., 1949: 268, 282, 285, 287-288, B.C., Ont.; Wilkes, 1949: 111-112, B.C. (ex *Archips*); Wilkes, Coppel and Mathers, 1949: 140, 146, 149, 151, 153, B.C.; Anon., 1948: 281, 293, 295, B.C.; Anon., 1948a: 350, 366, 368-371, B.C.; Annand, 1948: 206, Colo. (ex spruce budworm); Dowden, Buchanan and Carolin, 1948: 458, 462-463 (ex *Archips*); Anon., 1947: 181, 196-201, B.C.; Coppel, 1947: 39, B.C. (ex *Archips*); Baird, 1946: 135, 148, 150-151, B.C. (ex *Archips*); Wilkes, 1946: 83-86, B.C. (ex *Archips*); Baird, 1945: 122, 133, 137, B.C. (ex *Archips*).

STAGE PARASITIZED: Larva, pupa.

IMMATURES: Egg — Coppel and Maw, 1954: (descr., fig. 2).

First instar larva — Coppel and Maw, 1954: (descr., figs. 3-7).

Second instar larva — Coppel and Maw, 1954: (descr., figs. 8-9).

Third instar larva — Coppel and Maw, 1954: (descr., figs. 10-11).

Pupa — Coppel and Maw, 1954: (descr., figs. 12-14); Ross, 1952: (fig. 15).

BIOLOGY: Coppel and Maw, 1954: (144-155); Coppel, 1953: (191); Coppel, 1947: (39).

Ceromasia aurifrons Townsend

Most records as *Masicera rutila* (Meigen).

HOST: LEPIDOPTERA

Geometridae

Cingilia catenaria (Cramer)

Schaffner and Griswold, 1934: 98, 119, Conn., Mass., N.H., N.Y. (as *C. "rutila"*); Aldrich, 1932: 2, N.H.; Curran, 1927a: 17, N.S. (as *Erycia* ex *C. "catanaria"*).

Noctuidae

Hypena humuli (Harris)

Hawley, 1918: 196, N.Y.

Tortricidae

Choristoneura fumiferana (Clemens)

Ross, 1952: 112, fig. 14, Canada (as *C. aurifrons*); Dowden,

Buchanan and Carolin, 1948: 458 (ex *Archips*); Tothill, 1913: 73, B.C. (ex *Tortrix*).

Unidentified Lepidoptera

Miscellaneous

Townsend, 1908: 102, Mass. (as *C. aurifrons*).

STAGE PARASITIZED: Larva, pupa.

IMMATURES: Pupa — Ross, 1952: (fig. 14); Greene, 1922: (descr., fig. 89 on pl. 19).

BIOLOGY: Schaffner and Griswold, 1934: (98, 119).

Ceromasia lepida, see *Lydella thompsoni*

Ceromasia rutila, see *Ceromasia aurifrons*

Ceromasia "rutila," see *Ceromasia aurifrons*

Ceromasia senilis, see *Lydella radialis*

Ceromasia sphenophori, see *Lixophaga sphenophori*

Ceromasia sp. (1) or (2), see *Ceromasia* spp.

Ceromasia spp.

HOST: LEPIDOPTERA

Geometridae

Lycia ursaria (Walker)

Schaffner and Griswold, 1934: 95-96, 119, Conn. (as *Ceromasia* sp. (2)).

Phigalia titea (Cramer)

Schaffner and Griswold, 1934: 97, 119, Mass., N.H., N.J. (as *Ceromasia* sp. (1)).

Noctuidae

Archanara oblonga (Grote)

Walton, 1908: 298, Pa. (ex *Nonagria*).

Papaipema purpurifascia (Grote and Robinson)

Bird, H., 1914: 69 (as *Ceromasia* fly).

Tortricidae

Choristoneura fumiferana (Clemens)

Powell, 1962: 137, Calif.

STAGE PARASITIZED: Larva.

BIOLOGY: Schaffner and Griswold, 1934: (95-97, 119).

"*Ceromesia sphenophori*," see *Lixophaga sphenophori*

Chaetexorista Brauer and Bergenstamm (Goniinae: Exoristini)

Chaetexorista javana Brauer and Bergenstamm

HOST: LEPIDOPTERA

Limacodidae

Cnidocampa flavescens (Walker)

Sabrosky and Arnaud, 1965: 1053, Mass.; Dowden, 1962: 3, 8, 52-53, Mass.; Schaffner, 1959: 6, 50, Mass.; Clausen, 1956: 4, 65, 89-90, Mass.; Dowden, 1946: 225-241, Mass.; Brown, 1937: 626, Mass. (ex oriental moth); Collins, 1935: 38, Mass. (as "*Chootexorista*" ex oriental moth); Collins, 1934: 32, Mass. (ex oriental moth); Proper, 1934: 374, Mass., New England (ex oriental moth); Collins, 1933: 28, [probably Mass.] (ex oriental moth); Anon., 1930: 894, Mass.

STAGE PARASITIZED: Larva, pupa.

BIOLOGY: Dowden, 1962: (3, 8, 52-53); Schaffner, 1959: (6, 50); Clausen, 1956: (4, 65, 89-90); Dowden, 1946: (225-241).

Chaetogaedia Brauer and Bergenstamm (Goniinae: Goniini)**Chaetogaedia analis** (Wulp)

HOST: LEPIDOPTERA

Arctiidae

Unidentified Arctiidae

Schaffner and Griswold, 1934: 40-41, 125, N.H.

Geometridae

Ennomos subsignarius (Hübner)

Davis, R., 1961: 248, Ga.

Noctuidae

Lacinipolia renigera (Stephens)Crumb, 1929: 40, 128, Va. (ex *Polia*).

STAGE PARASITIZED: Larva, pupa.

IMMATURES: Pupa — Greene, 1922: (descr., fig. 62 on pl. 13).

BIOLOGY: Schaffner and Griswold, 1934: (40-41, 125).

Chaetogaedia sp. near *analis* (Wulp)

HOST: LEPIDOPTERA

Noctuidae

Heliothis spp.

Butler, 1958: 562, Ariz.

STAGE PARASITIZED: Larva.

Chaetogaedia crebra (Wulp)

HOST: LEPIDOPTERA

Noctuidae

Agrotis sp.

Coquillett, 1897: 11, 23, Calif.

Protorthodes rufula (Grote)Essig, 1926: 584, Calif. (ex *Eriopyga*); Coquillett, 1897: 11,
27, Calif. (ex *Taeniocampa*).

Cutworm(-s)

Essig, 1926: 584, Calif.

Pieridae

Pieris occidentalis ReakirtEssig, 1926: 584, N. Mex. (ex *Neophasia*).

STAGE PARASITIZED: Larva.

Chaetogaedia monticola (Bigot)

HOST: LEPIDOPTERA

Danaiidae

Danaus plexippus plexippus (Linnaeus)Severin, H. H. P., Severin and Hartung, 1915: 134 (ex *Anosia*
plexippus, lab.).

Geometridae

Eupithecia monticolans ButlerZimmerman, 1958: 158, Hawaii; Swezey, 1943: 279, Hawaii
(ex *Eucymatoge*).*Scotorythra paludicola* (Butler) probably

Swezey, 1926a: 361, Hawaii.

Scotorythra rara (Butler)Zimmerman, 1958: 135, Hawaii; Swezey, 1912: 160, Hawaii;
Swezey, 1907: 49, Hawaii.*Scotorythra trapezias* Meyrick

Zimmerman, 1958: 140, Hawaii; Swezey, 1943: 280, Hawaii.

Noctuidae

- Achaea janata* (Linnaeus)
 Zimmerman, 1958: 353, Hawaii; Bianchi, 1946: 469, Hawaii;
 Van Zwaluwenburg, 1946: 14, Hawaii.
- Actebia fennica* (Tauscher)
 Swezey, 1907: 49, Hawaii (ex black army worm).
- Agrotis dislocata* (Walker)
 Zimmerman, 1958: 239, Hawaii; Swezey, 1907: 49, Hawaii.
- Agrotis ipsilon* (Hufnagel)
 Zimmerman, 1958: 253, Hawaii.
- Anomis hawaiiensis* (Butler)
 Zimmerman, 1958: 381, Hawaii; Swezey, 1907: 49, Hawaii
 (ex *Cosmophila sabulifera* (Guenée)).
- Anomis noctivolans* (Butler)
 Zimmerman, 1958: 385, Hawaii.
- Autographa californica* (Speyer)
 Clancy, 1969: 1079, Calif.
- Autoplusia egena* (Guenée)
 Lange, 1945: 13, Calif. (ex *Autographa*).
- Carneades* sp.
 Coquillett, 1897: 11, 24, Hawaii.
- Chrysodeixis calcites* (Esper)
 All records ex *Plusia* "chalcites."
 Zimmerman, 1958: 371, Hawaii; Bianchi, 1940: 385,
 Hawaii; Illingworth, 1937: 354, Hawaii; Swezey, 1936:
 117, Hawaii; Severin, H. H. P., Severin and Hartung,
 1915: 134 (lab.); Swezey, 1907: 49, Hawaii.
- Elydna nonagrica* (Walker)
 Fullaway, 1914: 8, Hawaii (ex *Caradrina reculsa* Hampson).
- Heliothis virescens* (Fabricius)
 Beardsley, 1957: 190, Hawaii
- Peridroma cinctipennis cinctipennis* (Butler)
 Most records ex *Agrotis cinctipennis*.
 Zimmerman, 1958: 279, Hawaii (ex *P. cinctipennis*
cinctipennis); Swezey, 1908: 163-164, Hawaii;
 Swezey, 1908a: 7-9, Hawaii; Swezey, 1908b: 17-19,
 pl. 3, Hawaii; Swezey, 1907: 49, Hawaii.
- Peridroma coniotis coniotis* (Hampson)
 Zimmerman, 1958: 283, Hawaii; Look, 1948: 222, Hawaii
 (ex *Agrotis coniotis*).
- Peridroma saucia* (Hübner)
 Swezey, 1935a: 93, Hawaii (ex *Lycophotia margaritosa*
 (Haworth)); *Crumb, 1929: 40, 115 (ex *Lycophotia*,

Wakeland, 1920); Essig, 1926: 584-585, Calif., Colo. (ex *Lycophotia margaritosa*); Wakeland, 1920: 18, Colo.; Chittenden, 1901a: 62; Coquillett, 1897: 11, 26, Calif.

Polydesma umbricola Boisduval

Kamran, 1968: 1010-1011, Hawaii; Zimmerman, 1958: 392, Hawaii; Rosa, 1946: 478, Hawaii; Van Zwaluwenburg, 1946: 16, Hawaii.

Pseudaletia macrosaris (Meyrick)

Zimmerman, 1958: 303, Hawaii; Swezey, 1907: 49, Hawaii (ex *Leucania pyrrhias* Meyrick).

Pseudaletia unipuncta (Haworth)

Zimmerman, 1958: 305, Hawaii; Severin, H. H. P., Severin and Hartung, 1915: 134, 136 (ex *Heliophila*, lab.); Swezey, 1907: 49, Hawaii (ex *Leucania*).

Spodoptera exempta (Walker)

Most records ex *Laphygma exempta*.

Zimmerman, 1958: 336, Hawaii (ex *Spodoptera*); Bess, 1951: 212, Hawaii; Bianchi, 1949a: 345, Hawaii; Pemberton, 1948b: 183-184, 200, figs. 2-3, Hawaii.

Spodoptera exigua (Hübner)

Swezey, 1935a: 93, Hawaii (as *Chaetogaedia* sp.).

Spodoptera mauritia (Boisduval)

Beardsley, 1955: 386, Hawaii; Swezey, 1907: 49, Hawaii.

Armyworm(-s) (including *Pseudaletia unipuncta* (Haworth) and/or *Spodoptera mauritia* (Boisduval))

All records ex armyworms (*Cirphis unipuncta* and/or *S. mauritia*).

Swezey, 1931c: 326, Hawaii; Swezey, 1928: 673, Hawaii.

Armyworm(-s)

Bianchi, 1940: 383, Hawaii; Bryan, 1934: 415, Hawaii; Swezey, 1931b: 427, Hawaii; Swezey, 1927a: 503, Hawaii; Swezey, 1927b: 217, Hawaii; Swezey, 1917: 432-433, figs. 9-13, Hawaii; Swezey, 1909: 26, 28-30, pl. 3, Hawaii (as "*Caetogaedia*"); Swezey, 1908: 163-164, Hawaii; Swezey, 1908b: 16-17, Hawaii.

Cutworm(-s) (including *Agrotis* spp. and/or *Leucania* spp.)

Swezey, 1908b: 16-17, Hawaii.

Cutworm(-s)

Bianchi, 1940: 383, Hawaii; Bryan, 1934: 415, Hawaii; Illingworth, 1923: 277, Hawaii; Fullaway, 1914: 8, Hawaii; Swezey, 1909: 26, 28-30, pl. 3, Hawaii (as "*Caetogaedia*"); Swezey, 1908: 163-164, Hawaii.

Unidentified Noctuidae

Swezey, 1944a: 4, Hawaii.

Nymphalidae

Vanessa cardui (Linnaeus)

Zimmerman, 1958: 457, Hawaii; Essig, 1926: 584, Calif., Oreg.; Severin, H. H. P., Severin and Hartung, 1915: 134 (lab.); Swezey, 1907: 49, Hawaii; Coquillett, 1897: 11, 27, Calif. (ex *Pyrameis*).

Vanessa tammeamea Eschscholtz

Williams, 1931: 291-292, figs. 8-13 on pl. 19, Hawaii; Swezey, 1927: 356, Hawaii.

Pieridae

Pieris rapae (Linnaeus)

Severin, H. H. P., Severin and Hartung, 1915: 134 (ex *Pontia*, lab.)

Pyralidae

Hedylepta accepta (Butler)

Swezey, 1907: 49, Hawaii (ex *Omiodes*).

Hedylepta blackburni (Butler)

Williams, 1931: 148, 291-292, figs. 8-13 on pl. 19, Hawaii (ex *Omiodes*).

Hedylepta continuatalis (Wallengren)

Swezey, 1907: 49, Hawaii (ex *Omiodes*).

Hymenia recurvalis (Fabricius)

Swezey, 1907: 49, Hawaii.

Oeobia monticolans (Butler)

Swezey, 1907: 49, Hawaii (ex *Phlyctaenia*).

Sphingidae

Agrius cingulatus (Fabricius)

Severin, H. H. P., Severin and Hartung, 1915: 134 (ex *Herse*, lab.).

Unidentified Lepidoptera

Miscellaneous

Swezey, 1931a: 495, Hawaii (ex caterpillars feeding on *Acacia koa* (A. Gray)).

STAGE PARASITIZED: Larva, pupa.

IMMATURES: Egg — Pemberton, 1948b: (descr., figs. 2-3); Williams, 1931: (figs. 12-13 on pl. 19); Swezey, 1917: (figs. 12-13; Swezey, 1909: (descr., pl. 3); Swezey, 1908a: (descr.); Swezey, 1908b: (descr., pl. 3).

Larva — Williams, 1931: (fig. 9 on pl. 19); Swezey, 1917: (fig. 9).

Third instar larva — Swezey, 1909: (pl. 3); Swezey, 1908b: (pl. 3).

Pupa — Williams, 1931: (descr., figs. 10-11 on pl. 19); Swezey, 1917: (figs. 10-11); Swezey, 1909: (pl. 3); Swezey, 1908a: (descr.); Swezey, 1908b: (descr., pl. 3).

BIOLOGY: Kamran, 1968: (1010-1011); Pemberton, 1948b: (183-184, 200); Swezey, 1944a: (4); Williams, 1931: (148, 291-292); Severin, H. H. P., Severin and Hartung, 1915: (134, 136); Swezey, 1908a: (7-9); Swezey, 1908b: (17-19).

Chaetogaedia monticola (Bigot) probably

HOST: LEPIDOPTERA

Noctuidae

Peridroma saucia (Hübner)

Zimmerman, 1958: 290, Hawaii (ex *P. porphyrea* (Denis and Schiffermüller)).

Chaetogaedia sp. near *monticola* (Bigot)

HOST: LEPIDOPTERA

Lasiocampidae

Malacosoma californicum fragile (Stretch)

Stehr and Cook, 1968: 278-279, 301, Ariz., Calif.

Malacosoma constrictum austrinum Stehr

Stehr and Cook, 1968: 299, Calif.

Malacosoma incurvum discoloratum (Neumoegen)

Stehr and Cook, 1968: 280, 301, Nev.

STAGE PARASITIZED: Larva.

Chaetogaedia sp., see *Chaetogaedia monticola*

Gen. and sp. near *Chaetogaedia*

HOST: LEPIDOPTERA

Noctuidae

Litoprosopus sp.

Patton, 1958: 35, Fla. (incomplete rearing — pupa).

STAGE PARASITIZED: Larva.

Chaetolyga sp., see *Carcelia* spp.

Chaetolyga deilephilae, see *Winthemia deilephilae*

Chaetolyga leucanae, see *Winthemia leucanae*

Chaetolyga "leucaniae," see *Winthemia leucanae*

Chaetona Wulp (Dexiinae: Urodexiini)

Chaetona sp.

HOST: LEPIDOPTERA

Arctiidae

Turuptiana permaculata (Packard)

Townsend, 1895: 78, Colo. (ex *Ecpantheria*).

Chaetonodexodes Townsend (Dexiinae: Urodexiini)

Chaetonodexodes vanderwulpi (Townsend)

HOST: COLEOPTERA

Chrysomelidae

Monoxia consputa (LeConte)

Essig, 1926: 579, Utah (as *Tachinophyto*).

STAGE PARASITIZED: Larva.

Chaetophleps setosa, see *Celatoria setosa*

Chaetophleps n. sp., see *Miamimyia cincta*

Chaetophlepsis Townsend (Tachininae: Campylochetini)

Chaetophlepsis eudryae (Smith)

Most records as *Chaetophlepsis tarsalis* Townsend.

HOST: LEPIDOPTERA

Aegeriidae

Melittia cucurbitae (Harris)

Fattig, 1949: 11 (ex "*Melitta*").

Melittia cucurbitae (Harris) probably

Brimley, 1938: 359, N.C. (ex squash borer moth).

Noctuidae

Alypia octomaculata (Fabricius)

Schaffner and Griswold, 1934: 41, 108; Webber, 1931: 4;

Webber and Schaffner, 1926: 22, New England.

Autographa sp.

Vickery, 1925: 140, Tex.

Euthisanotia grata (Fabricius)

All records ex *Eudryas grata*.

Johnson, 1925: 189, Maine, Mass., N.H., R.I.; Smith, H. E.,
1916: 94, Mass. (as *Hypochoeta*).

Plathypena scabra (Fabricius)

Hill, 1925: 17, Tenn. (as *Hypochoeta*).

Pseudoplusia includens (Walker)

Allen, 1929: 683, Miss. (ex *Autographa oo* (Cramer)).

Unidentified Noctuidae

Aldrich, 1923b: 162, Miss.

Notodontidae

Pheosia rimosa Packard

Schaffner and Griswold, 1934: 77, 108.

STAGE PARASITIZED: Larva.

BIOLOGY: Schaffner and Griswold, 1934: (108).

Chaetophlepsis orbitalis Webber

HOST: LEPIDOPTERA

Geometridae

Cingilia catenaria (Cramer)

Wood, 1951: 243, N.B.; Schaffner and Griswold, 1934: 98, 107, Mass., N.H., R.I.; *vide Webber, 1931: 3-4, Mass., N.H., R.I. (Greene, 1922).

Protitame virginalis (Hulst)

Raizenne, 1952: 132, Ont.

Noctuidae

Cryphia pervertens Barnes and McDunnough

Raizenne, 1952: 63, Ont.

IMMATURES: Pupa — Greene, 1922: (descr., fig. 74 on pl. 16).

BIOLOGY: Schaffner and Griswold, 1934: (98, 107); Webber, 1931: (3-4).

Chaetophlepsis rindgei Reinhard

HOST: LEPIDOPTERA

Geometridae

Drepanulatrix monicaria (Guenée)

*Vide Reinhard, 1952: 21, Calif. (Rindge, 1949); Rindge, 1949: 285, 288, Calif. (as *Chaetophlepsis* sp. probably undescribed).

Drepanulatrix spp. (including *D. baueraria* Sperry and/or *D. monicaria* (Guenée))

[*Vide Reinhard, 1952: 21, Calif. (Rindge, 1949)]; Rindge, 1949: 285, 288, Calif. (as *Chaetophlepsis* sp. probably undescribed).

STAGE PARASITIZED: Larva.
Chaetophlepsis semiothisae Brooks
 HOST: LEPIDOPTERA

Geometridae

- Itame fulvaria* Villers probably
 Brooks, 1945: 82, Ont.
Lambdina fiscellaria fiscellaria (Guenée)
 Brooks, 1945: 82, Ont. (ex "*Lamlidina*" *fiscellaria*).
Nepytia canosaria (Walker)
 Brooks, 1945: 82, B.C. (ex "*Nephytia*").
Pero sp.
 Brooks, 1945: 82, Ont.
Semiothisa bisignata (Walker)
 Raizenne, 1952: 122, Ont.
Semiothisa granitata Guenée complex
 All records ex *Semiothisa granitata*.
 Raizenne, 1952: 124, Ont.; Brooks, 1945: 82-83, B.C.,
 N.B., N.S., Ont., Que.
Semiothisa minorata (Packard)
 Raizenne, 1952: 121, Ont.; Brooks, 1945: 82-83, Que.
Semiothisa oweni (Swett)
 Raizenne, 1952: 126, Ont.
Semiothisa sexmaculata (Packard)
 Brooks, 1945: 83, N.B.

Tortricidae

- Acleris variana* (Fernald)
 Miller, 1966: 603, N.S.
 STAGE PARASITIZED: Larva.
Chaetophlepsis tarsalis, see *Chaetophlepsis eudryae*
Chaetophlepsis sp. probably undescribed [of Rindge, 1949], see
Chaetophlepsis rindgei
Chaetophlepsis spp.
 HOST: LEPIDOPTERA

Aegeriidae

- Melittia satyriniformis* Hübner
 Brimley, 1922: 23, N.C. (as *Hypochoaeta longicornis*).

Noctuidae

Catocala antinympha (Hübner)

Schaffner and Griswold, 1934: 64, 108, Maine (as *Chaetophlepsis*).

Plathypena scabra (Fabricius)

Whiteside, Burbutis and Kelsey, 1967: 327, Del. (as *Chaetophlepsis*); Hill, 1925: 17, Miss. (as *Hypochoaeta longicornis*).

Tortricidae

Choristoneura conflictana (Walker)

Prentice, 1955: 472, Alta. (as *Chaetophlepsis*).

STAGE PARASITIZED: Larva.

BIOLOGY: Schaffner and Griswold, 1934: (64, 108).

Chaetotachina simulans, see *Guerinia rufostomata*

"*Chalomyia*" *longipes*, see *Cholomyia inaequipes*

Choeteprosopa Schiner (Goniinae: Belvosiini)

Choeteprosopa hedemanni Brauer and Bergenstamm

All records as *Paragaedia hedemanni*.

HOST: LEPIDOPTERA

Noctuidae

Spodoptera eridania (Cramer)

Aldrich, 1925a: 463, Panama (ex *Xylomyges*).

Spodoptera ornithogalli (Guenée)

Crumb, 1929: 40 (ex *Prodenia*).

STAGE PARASITIZED: Larva.

Cholomyia Bigot (Dexiinae: Urodexiini)

Cholomyia inaequipes Bigot

HOST: COLEOPTERA

Curculionidae

Conotrachelus affinis Boheman

Mutchler and Weiss, 1925: 10-11 (as *C. longipes* (Fabricius));
Quaintance and Jenne, 1912: 150, Ark.; Brooks, F. E.,
1911a: 190, W. Va. (as *Metadexia basalis* Giglio-Tos);
Brooks, F. E., 1910: 182, W. Va. (as *Metadexia basalis* ex
hickorynut curculio).

Conotrachelus aratus Germar

Mutchler and Weiss, 1925: 13 (as "*Chalomyia*" *longipes*).

Conotrachelus elegans (Say)

Pierce, 1908: 381, 390, Tex.

Conotrachelus juglandis LeConte

Johnson, 1912: 102-103, La. (as *C. longipes*); Pierce, Cushman and Hood, 1912: 78; Pierce, Cushman and Hood, 1912: 78 (as *Metadexia basalis*); Quaintance and Jenne, 1912: 150, La., Tex.; Quaintance and Jenne, 1912: 150, W. Va. (as *Metadexia basalis*); Pierce, 1908: 381, 390, La.; Pierce, 1908: 381, 390, W. Va. (as *Metadexia basalis*); Pierce, 1908a: 42 (as *Metadexia basalis*).

Conotrachelus naso LeConte

Gibson, L., 1964: 525, Ala., Ga., Ohio.

Conotrachelus nenuphar (Herbst)

Armstrong, 1958: 17, Ont.; Fattig, 1949: 12; Snapp, 1930: 77, 79, Ga. (as *C. longipes*); Pierce, Cushman and Hood, 1912: 78; Quaintance and Jenne, 1912: 150, fig. 30, Ark., Ga., Md., Mo., Pa., Va. (ex *Curculio*); Pierce, 1908: 381, 390, Mo.

Conotrachelus posticatus Boheman

Gibson, L., 1964: 525, Ala., Ga., Ohio.

Conotrachelus spp.

Johnson, 1912: 102, Mo. (as *C. longipes*); Pierce, 1910: 453; Pierce, 1910: 453 (as *Metadexia basalis*).

STAGE PARASITIZED: Larva.

IMMATURES: Pupa — Quaintance and Jenne, 1912: (fig. 30).

BIOLOGY: Pierce, 1910: (453).

Cholomyia longipes, see *Cholomyia inaequipes*

Cholomyia sp.

HOST: COLEOPTERA

Curculionidae

Conotrachelus naso LeConte

Gibson, L., 1964: 525, Ala.

“*Chootexorista*” javana, see *Chaetexorista javana*

Chrysoexorista Townsend (Goniinae: Eryciini)

Chrysoexorista dawsoni (Sellers)

All records as *Zenillia dawsoni*.

HOST: LEPIDOPTERA

Geometridae

Lambdina fiscellaria laeta (Hulst) probably

Sellers, 1943: 29, N. Mex. (ex *Ellopia* “*lactea*”).

Olethreutidae

Laspeyresia pomonella (Linnaeus)

Sellers, 1943: 29, Ga. (ex *Carpocapsa*).

Chrysoexorista marginata (Aldrich and Webber)

HOST: LEPIDOPTERA

Geometridae

Hydria spp. (including *H. prunivorata* (Ferguson) and/or *H. undulata* (Linnaeus)).

Raizenne, 1952: 113, Ont. (ex *Calocalpe undulata*).

Olethreutidae

Laspeyresia pomonella (Linnaeus)

Aldrich and Webber, 1924: 17-18, Ga. (as *Zenillia* ex *Carpocapsa*, probable host).

Chrysoexorista ochracea (Wulp)

HOST: LEPIDOPTERA

Olethreutidae

Laspeyresia pomonella (Linnaeus)

Van Leeuwen, 1929: 78, Ga. (as *Exorista* "achracea" ex *Carpocapsa*).

STAGE PARASITIZED: Larva.

Chrysotachina Brauer and Bergenstamm (Tachininae: Linnaemyini)

Chrysotachina alcedo (Loew)

HOST: LEPIDOPTERA

Hesperiidae

Urbanus proteus (Linnaeus)

Patton, 1958: 36, Fla.; Aldrich, 1926b: 57, Fla. (ex *Eudamus*).

Lacosomidae

Cicinnus melsheimeri (Harris)

Schaffner and Griswold, 1934: 102, 111, Conn.

STAGE PARASITIZED: Larva.

BIOLOGY: Schaffner and Griswold, 1934: (102, 111).

Cistogaster immaculata, see *Gymnoclytia immaculata*

"*Cistogoster*" *immaculata*, see *Gymnoclytia immaculata*

Clausicella Rondani (Goniinae: Siphonini)

Clausicella floridensis (Townsend)

Most records as *Siphophyto floridensis*.

HOST: LEPIDOPTERA

Olethreutidae

Endothenia hebesana Walker

Satterthwait, 1948: 729 (ex *Olethreutes*); Satterthwait and Swain, 1946: 578.

Epiblema tripartitana (Zeller)

Bottimer, 1926: 816, Tex.

Suleima helianthana (Riley)

Satterthwait, 1948: 726, 729; Satterthwait and Swain, 1946: 578.

Pyralidae

Alberada parabates (Dyar)

Mann, 1969: 14, 43, Calif. (as *Clausicella*).

Homoeosoma electellum (Hulst)

Satterthwait, 1948: 729 (ex sunflower moth); Satterthwait and Swain, 1946: 578, Mo.

Plodia interpunctella (Hübner)

Morrill, 1940: 126 (as *Epigrymyia* ex Indian meal moth); Hamlin, Reed and Phillips, 1931: 23, Calif. (as *Drepanoglossa*); Simmons, Reed and M'Gregor, 1931: 42, Calif. (as *Drepanoglossa*).

Plodia sp.

Essig, 1926: 579, N. Mex. (as "*Epigrymyia*").

STAGE PARASITIZED: Larva, pupa.

IMMATURES: Pupa — Satterthwait and Swain, 1946: (descr.).

BIOLOGY: Satterthwait and Swain, 1946: (578).

Clausicella neomexicana (Townsend)

Most records as *Siphophyto neomexicana*.

HOST: COLEOPTERA

Curculionidae

Curculio sp.

Reinhard, 1946: 84, Calif. (ex *Balaninus* larvae on acorn, questionable host).

HOST: HYMENOPTERA

Apidae

Bombus fervidus (Fabricius)

Reinhard, 1946: 83-84, N. Mex. (ex nest of *B. fervidus*, questionable host).

HOST: LEPIDOPTERA

Pyralidae

Homoeosoma electellum (Hulst)

Teetes and Randolph, 1969: 1463-1464, Tex. (as *Clausicella*).

Plodia interpunctella (Hübner)

Reinhard, 1946: 83, Calif., Kans.

Plodia spp.

Vide Reinhard, 1946: 83-84, N. Mex. (Coquillett, 1897);
Coquillett, 1897: 11, 26, N. Mex. (as *Epigrimyia floridensis* (Townsend)).

STAGE PARASITIZED: Larva (Coleoptera, Lepidoptera).

Clausicella opaca (Coquillett)

HOST: LEPIDOPTERA

Noctuidae

Peridroma saucia (Hübner)

Anon., 1969: 59, Okla.

Pyralidae

Homoeosoma electellum (Hulst)

Reinhard, 1946: 89, Tex. (as *Siphophyto* ex "*Homoesoma*").

Clausicella tarsalis, see *Elfia tarsalis*

Clausicella turmalis (Reinhard)

HOST: LEPIDOPTERA

Geometridae

Eupithecia coagulata Guenée probably

All records ex *Tephroclystis absynthiata* Clerck.

Vide Reinhard, 1946: 87, N.H. (as *Siphophyto*, Aldrich, 1932); Aldrich, 1932: 2, N.H. (as *Siphophyto floridensis* Townsend).

Clytiomyia flava, see *Euclytia flava*

Coloradomyia Arnaud (Tachininae: Germariini)

Coloradomyia eucosmaphaga Arnaud

HOST: LEPIDOPTERA

Olethreutidae

Eucosma ridingsana (Robinson)

Arnaud, 1963b: 2, 19-20, figs. 21-22, Colo.

IMMATURES: Pupa — Arnaud, 1963b: (descr., figs. 21-22).

"Compilura" *concinata*, see *Compsilura concinnata**Compsilura* Bouché (Goniinae: Blondeliini)*Compsilura concinnata* (Meigen)

HOST: COLEOPTERA

Curculionidae

Pissodes strobi (Peck)Mott, 1930: 34, N.E. U.S.; Webber and Schaffner, 1926: 27,
New England, N.H.; Howard, 1922: 137 (probable
parasite).

HOST: HYMENOPTERA

Cimbicidae

Cimbex americana LeachWebber and Schaffner, 1926: 2, 8, 26, New England;
Howard, 1922: 137; Culver, 1919: 5, New England.

Diprionidae

Diprion similis (Hartig)

Webber and Schaffner, 1926: 27, Conn., New England.

Pamphiliidae

Neurotoma fasciata (Norton)

Webber and Schaffner, 1926: 3, 10, 26, New England.

Tenthredinidae

Croesus latitarsus NortonWebber and Schaffner, 1926: 2, 8, 26, New England;
Howard, 1922: 136.*Nematus erythrogaster* NortonWebber and Schaffner, 1926: 3, 10, 27, New England (ex
Pteronidea coryla Cresson).

Nematus ribesii (Scopoli)

Webber and Schaffner, 1926: 3, 9, 27, New England (ex *Pteronidea*).

Pristiphora geniculata (Hartig)

Forbes and Daviault, 1964: 1128; Raizenne, 1957: 30, Ont.

Trichiocampus viminalis (Fallén)

Beigué, 1961: 1093, Que.

Unidentified Hymenoptera

Sawfly (-flies)

Howard, 1922: 137.

Tenthredinoidea

Schaffner, 1934: 586, 589.

HOST: LEPIDOPTERA

Arctiidae

Diacrisia virginica (Fabricius)

Schaffner and Griswold, 1934: 36, 115; Webber and Schaffner, 1926: 2, 5-6, 9, 14, 20, New England; Howard, 1922: 135; Culver, 1919: 5, 22, New England; Smith, H. E., 1917: 58, Mass.

Estigmene acraea (Drury)

Most records ex *Estigmene acraea*.

Schaffner and Griswold, 1934: 38, 115 (ex *E. acraea*); Webber and Schaffner, 1926: 3, 9, 14, 20, New England; Howard, 1922: 135; Culver, 1919: 5, New England.

Euchaetias egle (Drury)

Webber, 1937: 144-149 (lab.); Dowden, 1935: 509, 517 (lab.); Schaffner and Griswold, 1934: 39, 115; Schaffner, 1927: 729; Webber and Schaffner, 1926: 3, 6, 9, 20, New England; Howard, 1922: 135; Culver, 1919: 5, New England.

Halisidota argentata Packard

Silver, 1958: 78, B.C.

Halisidota caryae (Harris)

Raizenne, 1952: 21, Ont.; Webber, 1937: 144-149 (lab.); Schaffner and Griswold, 1934: 33, 115; Schaffner, 1927: 729; Webber and Schaffner, 1926: 3, 9, 20, New England; Howard, 1922: 136.

Halisidota harrisii Walsh

Raizenne, 1952: 25, Ont.; Schaffner and Griswold, 1934: 34-35, 115.

Halisidota maculata (Harris)

Schaffner and Griswold, 1934: 33-34, 115.

Halisidota tessellaris (J. E. Smith)

Schaffner and Griswold, 1934: 34, 115; Webber and Schaffner, 1926: 3, 9, 20, New England; Howard, 1922: 136.

Hyphantria cunea (Drury)

Webber, 1937: 144-149 (lab.); Dowden, 1935: 517 (lab.); Schaffner, 1934: 589; Schaffner and Griswold, 1934: 37, 115; Webber and Schaffner, 1926: 3, 6, 9, 13-14, New England; Tothill, 1922: 9, 29, 36-43, 107, maritime prov., N.B.; Culver, 1919: 5, New England; Tothill, 1916: 114, N.B. or N.S.; Tothill, 1913a: 57, N.B.; Howard, 1911: 96 (as *Compsilura* [sp.] ex fall webworm); Howard and Fiske, 1911: 203 (ex fall webworm).

Hyphantria textor (Harris)

Reeks and Smith, 1956: 575, maritime prov.; Raizenne, 1952: 29, Ont.; Reeks, Smith and Forbes, 1946: 14, maritime prov.; Daviault, 1942: 398, Que. (as *C. "conncinata"*); Howard, 1922: 136; Thompson, 1910: 292.

Hyphantria sp.

Tothill, 1919: 13, N.B.

Unidentified Arctiidae

Schaffner, 1934: 589-590; Schaffner and Griswold, 1934: 40-41, 115; Webber and Schaffner, 1926: 2, 8, New England; Culver, 1919: 5, New England.

Bombycidae

Bombyx mori Linnaeus

Webber, 1937: 144-149 (lab.); Dowden, 1935: 509 (lab.); Webber and Schaffner, 1926: 10 (lab.); Culver, 1919: 1-27, figs. 1, 5-8, fig. 2 on pl. 1, Mass. (lab.).

Citheroniidae

Anisota rubicunda (Fabricius)

Reeks and Smith, 1956: 575, maritime prov.; Reeks, Smith and Forbes, 1946: 14, maritime prov.; Schaffner and Griswold, 1934: 30-31, 115; Webber and Schaffner, 1926: 2, 8, 26, New England.

Anisota senatoria J. E. Smith

Schaffner and Griswold, 1934: 29-30, 115; Webber and Schaffner, 1926: 2, 8, 25, New England.

Anisota virginiensis (Drury)

Schaffner and Griswold, 1934: 30, 115.

Anisota spp.

Schaffner, 1934: 589.

Ctenuchidae

Ctenucha virginica Charpentier

Schaffner and Griswold, 1934: 32, 115.

Danaiidae

Danaus plexippus plexippus (Linnaeus)

Ehrlich, 1948: 92 (ex *D. plexippus*); Webber, 1937: 144-149 (ex *D. menippe* (Hübner), lab.); Schaffner and Griswold, 1934: 8, 115 (ex *D. menippe*); Webber and Schaffner, 1926: 2, 8, 23, New England (ex *Anosia plexippus*); Howard, 1922: 135 (ex *Anosia plexippus*).

Geometridae

Biston cognataria (Guenée)

Schaffner and Griswold, 1934: 96, 115-116 (ex *Amphidasis*); Webber and Schaffner, 1926: 3, 9, 23, New England (ex *Lycia*).

Cingilia catenaria (Cramer)

Franklin, 1950: 57; Webber, 1937: 144-149 (lab.); Schaffner and Griswold, 1934: 98, 115-116; Webber and Schaffner, 1926: 2, 8, 23, New England.

Ennomos subsignarius (Hübner)

Schaffner and Griswold, 1934: 99-100, 115-116, Mass.; Webber and Schaffner, 1926: 2, 9, 23, New England; Culver, 1919: 5, New England.

Eugonobapta nivosaria (Guenée)

Schaffner and Griswold, 1934: 99, 115-116, Mass.

Hydria spp. (including *H. prunivorata* (Ferguson) and/or *H. undulata* (Linnaeus)).

Most records ex *Hydria undulata*.

Webber, 1937: 144-149 (lab.); Schaffner and Griswold, 1934: 91-92, 115-116 (ex *Calocalpe undulata*); Webber and Schaffner, 1926: 3, 9, 23, New England.

Itame ribearia (Fitch)

Schaffner and Griswold, 1934: 94, 115-116.

Lambdina fiscellaria fiscellaria (Guenée)

All records ex *Lambdina fiscellaria*.

Reeks and Smith, 1956: 575, maritime prov.; Reeks, Smith, and Forbes, 1946: 14, maritime prov.

Melanolophia imitata (Walker)

Evans, 1962: 597, B.C.

Nematocampa filamentaria Guenée

Reeks and Smith, 1956: 575, maritime prov. (ex *N. limbata* Haworth).

Phigalia titea (Cramer)

Schaffner and Griswold, 1934: 97, 115-116; Webber and Schaffner, 1926: 3, 9, 23, New England.

Unidentified Geometridae

Culver, 1919: 5, 22, New England.

Hesperiidae

Epargyreus clarus (Cramer)

All records ex *Epargyreus tityrus* (Fabricius).

Webber, 1937: 144-149 (lab.); Schaffner and Griswold, 1934: 15, 115; Schaffner, 1927: 729; Webber and Schaffner, 1926: 2, 9, 25, New England; Howard, 1922: 135.

Erynnis brizo (Boisduval and LeConte)

Webber and Schaffner, 1926: 27, New England (ex *Thanaos*).

Erynnis icelus (Scudder and Burgess)

Schaffner and Griswold, 1934: 15, 115.

Erynnis spp. (including *E. horatius* (Scudder and Burgess) and/or *E. juvenalis* (Fabricius))

Schaffner and Griswold, 1934: 15-16, 115, Mass.

Erynnis spp.

Webber and Schaffner, 1926: 3, 7, New England (ex *Thanaos*).

Thymelicus lineola (Ochsenheimer)

Arthur, 1962: 1083, 1085, Ont.

Unidentified Hesperiidae

Schaffner, 1934: 589-590.

Lasiocampidae

Epicnaptera americana (Harris)

Schaffner and Griswold, 1934: 89, 115-116; Webber and Schaffner, 1926: 3, 9, 13, New England.

Malacosoma americanum (Fabricius)

Webber, 1937: 144-149 (lab.); Bess, 1936: 605-607; Schaffner, 1934: 589; Schaffner and Griswold, 1934: 88, 115-116; Webber and Schaffner, 1926: 3, 9, 13, New England; Howard, 1922: 136; Culver, 1919: 5, New England.

Malacosoma californicum pluviale (Dyar)

Johansen, 1957: 70, Wash. (ex *M. pluviale*).

Malacosoma disstria Hübner

Sippell, 1961: 436, Ont.; Johansen, 1957: 70, Wash.; Webber, 1937: 144-149 (lab.); Bess, 1936: 605-607; Schaffner and Griswold, 1934: 89, 115-116; Webber and Schaffner, 1926: 3, 9, 13, New England; Howard, 1922: 136; Culver, 1919: 5, New England; Howard, 1914: 83-84 (ex forest tent caterpillar).

Malacosoma spp.

Ross, 1953: 20, 22, fig. 8, B.C.

American tent caterpillar(-s)

Howard, 1914: 83-84.

Limacodidae

Cnidocampa flavescens (Walker)

Schaffner, 1959: 6, 50, [N.E. U.S.]; Schaffner and Griswold, 1934: 104-105, 115-116, Mass.; Webber and Schaffner, 1926: 2, 8, 26, New England.

Lymantriidae

Dasychira basiflava (Packard)

All records ex *Olene basiflava*.

Schaffner and Griswold, 1934: 86, 115-116; Webber and Schaffner, 1926: 3, 9-10, New England.

Dasychira spp.

All records ex *Olene* spp.

Schaffner and Griswold, 1934: 87, 115-116; Webber and Schaffner, 1926: 3, 9, New England.

Euproctis chrysorrhoea (Linnaeus)

Webber and Schaffner, 1926: 1, 11, 30, Mass., New England; Burgess, 1924: 34-35, Mass.; Lochhead, 1919: 202; Burgess, 1917: 9, Mass.; Smith, H. E., 1917: 57; Burgess, 1915a: 6-7, 9-11, fig. 2 on pl. 3; Burgess, 1914: 6; Pouillaude, 1913: 82-83, Mass.; Tothill, 1913a: 57-58; Howard and Fiske, 1911: 263, 309, fig. 1 on pl. 20; Britton, 1910: 20.

Hemerocampa leucostigma (J. E. Smith)

Reeks and Smith, 1956: 575, maritime prov.; Franklin, 1950: 57; Reeks, Forbes and Cuming, 1947: 10, N.S., P.E.I.; Webber, 1937: 144-149 (lab.); Dowden, 1935: 509; Schaffner and Griswold, 1934: 85, 115-116; Webber and Schaffner, 1926: 3, 9, 11-12, 20, Mass., New England, R.I.; Howard, 1922: 135; Culver, 1919: 1-27, figs. 1, 5-8, fig. 2 on pl. 1, New England; Tothill and McLaine, 1919: 37-39, N.B. (ex white marked tussock moth).

Nygmia phaeorrhoea (Donovan)

Most records ex brown-tail moth.

Dowden, 1962: 27, 29, 54, U.S. (ex *N. phaeorrhoea*); Reeks and Smith, 1956: 570, 575, maritime prov. (ex *N. phaeorrhoea*); Conklin, 1952: 78, N.H.; Procter, 1946: 431, Maine (as "*Compsilura*"); Sellers, 1943: 56 (ex *N. phaeorrhoea*); Baird, 1939a: 52, E. Canada (ex *N. phaeorrhoea*); Baird, 1938: 85, 124, 141, 143, U.S. (ex *N. phaeorrhoea*); Webber, 1937: 144-149 (ex *N. phaeorrhoea*, lab.); Sweetman, 1936: 121-123, 221, 269, N. Amer.; Dowden, 1935: 509, 520 (ex *N. phaeorrhoea*); deGryse, 1935: 9, [N.B., N.S.]; Collins, 1934: 31, [Maine, Mass, and/or N.H.]; Proper, 1934: 369-370, 375-376, New England; Schaffner, 1934: 588-590, 592, New England (ex *N. phaeorrhoea*); Collins, 1933: 28, Mass. (ex *N. phaeorrhoea*); Collins and Jones, 1932: 237; Burgess, 1930: 28, Mass. (ex *N. phaeorrhoea*); Burgess and Crossman, 1929: 66, 134, 136, 142, Mass., N.H. (ex *N. phaeorrhoea*); Curran, 1927a: 24; Gilliatt, 1926: 64-65, N.S.; Britton and Ashworth, 1923: 316, Conn.; Gilliatt, 1921: 78, N.S.; Culver, 1919: 1-27, figs. 1, 5-8, fig. 2 on pl. 1, New England; Tothill and McLaine, 1919: 36-39, Mass., N.B.; Philbrook, 1918: 60, Maine; Britton and Irving, 1916: 113, Conn.; Burgess, 1916: 154, New England; McLaine, 1916: 22, 24, fig. 4; O'Kane, 1916: 132, N.H.; Britton, 1915: 19; Burgess, 1915: 85; Howard, 1913: 72-73, Mass. (as *Compsilura* [sp.]); Howard, 1911: 96 (as *Compsilura* [sp.]); Thompson, 1910: 292.

Orgyia antiqua (Linnaeus)

All records ex *Notolophus antiqua*.

Schaffner and Griswold, 1934: 84, 115-116; Webber and Schaffner, 1926: 3, 9, 11-12, New England; Howard, 1922: 136; Culver, 1919: 5, New England.

Porthetria dispar (Linnaeus)

Dowden, 1962: 30, 32, 34, 54, New England; Nichols, 1962: 20, Pa.; Dowden, 1961: 875, Mass.; Clausen, 1956: 6-7, 63, 97-98; Cooke, 1953: 152, Conn. (as *C. "concinati"*); Conklin, 1952: 78, N.H. (ex gypsy moth); Franklin, 1950: 57, 60; Friend, 1945: 620-621, Conn.; Brown and Sheals, 1944: 400; Felker, 1940: 122, N.H. (as *C. "concinati"* ex "gypsy" moth); Webber, 1937: 144-149 (lab.); *Sweetman, 1936: 121-123, 221, 269, N. Amer. (ex gypsy moth, Proper, 1934); Britton, 1935: 637, Conn.; Collins, 1935: 38, Mass. (ex gypsy moth); Dowden, 1935: 495, 509, 511, 517, 521; Collins, 1934: 31, Mass., New England (ex gypsy moth); Proper, 1934: 369-371; 375-376, New England (ex gypsy moth); Schaffner, 1934: 588-592, New England, N.J., Pa.; Britton and Ashworth, 1933: 407, Conn. (ex "gypsy" moth); Collins, 1933: 27, Mass. (ex gypsy moth); Collins and Jones, 1932: 237 (ex "gypsy" moth); Webber, 1932: 205; Anon., 1931a: 775, N.Y. (ex gypsy moth, incomplete rearing—pupa); Anon., 1931c: 124 (incomplete rearing—pupa); Burgess, 1930: 15-16, Mass.; Burgess and Crossman, 1929: 52-67, 142, fig. 24, Conn., Maine, Mass., N.H., Vt.; Curran, 1927a: 24 (ex gypsy moth); Webber and Schaffner, 1926: 1, 11, 30, Mass., New England; Burgess, 1924: 34-35, Mass.; Britton and Ashworth, 1923: 316, Conn. (ex "gypsy" moth); Culver, 1919: 1-27, figs. 1, 5-8, fig. 2 on pl. 1, New England; Lochhead, 1919: 200, 406; Tothill and McLaine, 1919: 36, 39 (ex "gypsy" moth); Britton and Davis, 1918: 250, Conn.; Howard, 1918: 45 (ex gypsy moth); Philbrook, 1918: 60, Maine (ex gypsy moth); Burgess, 1917: 9, Mass.; Smith, H. E., 1917: 57; Britton and Irving, 1916: 113, Conn. (ex "gypsy" moth); Burgess, 1916: 154, New England (ex "gypsy" moth); McLaine, 1916: 24-25, fig. 4 (ex "gypsy" moth); O'Kane, 1916: 132, pl. 4, N.H. (ex "gypsy" moth); Burgess, 1915a: 6-7, 9-11, fig. 2 on pl. 3; Hewitt, 1915: 12-13, 15, Mass. (ex "gypsy" moth); Burgess, 1914: 6; Howard, 1914: 83-84, Mass. (ex gypsy moth); Howard, 1914a: 86-87 (ex "gypsy" moth); Howard, 1913: 73, Mass. (ex gypsy moth); Pouillaude, 1913: 82-83, Mass. (ex "*Porthesia*" *dispar*); Tothill, 1913a: 57; Tothill, 1913a: 57, Mass. (ex "gypsy" moth); Howard, 1911: 96 (as *Compsilura* [sp.] ex gypsy moth); Howard and Fiske, 1911: 218-225, 308, fig. 1 on pl. 20, Mass.

Stilpnotia salicis (Linnaeus)

Dowden, 1962: 43-45, 54, Pacific N.W.; Lejeune and Silver, 1961: 456, 459-460, 462, 466, B.C.; Reeks and Smith, 1956: 569-571, 575-576, maritime prov.; Clausen, 1956: 65, 97-98; McLeod, 1954: 25, B.C.; Anon., 1952a: 164, B.C.; Anon., 1952e: 211, B.C.; Beaulne, 1949: 206, Que.; Reeks, Forbes and Cuming, 1946: 16, N.B.; Reeks, Smith and Forbes, 1946: 14, maritime prov.; Gauthier, 1945: 27, Que.; Prebble, 1945: 112, B.C.; Prebble, 1942: 111, B.C.; Baird, 1939a: 52, E. Canada; Balch, 1939: 31, maritime prov., Que.; Twinn, 1939: 21, N.B., N.S.; Baird, 1938: 86, 137, 140-141, 143, U.S.; Balch, 1938: 26, maritime prov.; Baird, 1937: 67, Canada; Balch, 1937: 26, N.B., N.S.; Webber, 1937: 144-149 (lab.); *Sweetman, 1936: 121-123, 221, 269, N. Amer. (ex satin moth, Proper, 1934); Collins, 1935: 38, Mass. (ex satin moth); deGryse, 1935: 9, maritime prov. (ex satin moth); Dowden, 1935: 509, 517; Webber, 1935: 460, Wash. (ex satin moth, probable host); Proper, 1934: 369-371, 375-376, New England (ex satin moth); Schaffner, 1934: 588-589, Mass.; Collins and Jones, 1932: 239 (ex satin moth); Weir, 1932: 178, B.C., N.S. (ex satin moth); Gilliatt, 1931: 13, N.S.; Glendenning, 1931: 32, E. U.S. (ex satin moth); Burgess and Crossman, 1927: 16-18, New England; Howard, 1926: 279-280, New England (ex "European" satin moth); Webber and Schaffner, 1926: 3, 10-11, 30, Mass., New England; Howard, 1922: 135.

Tussock moth(-s)

Howard, 1911: 96 (as *Compsilura* [sp.]); Howard and Fiske, 1911: 221-224, Mass.

Unidentified Lymantriidae

Schaffner, 1934: 589 (ex Liparidae).

Megalopygidae

Megalopyge crispata (Packard)

Schaffner, 1959: 7, 50, [N.E. U.S.] (ex *Lagoa*).

Noctuidae

Acronicta americana (Harris)

Schaffner and Griswold, 1934: 57-58, 115 (ex *Acronycta*); Webber and Schaffner, 1926: 2, 5, 8, 20, New England (ex *Apatela*).

Acronicta brumosa Guenée

All records ex *Apatela brumosa*.

Webber and Schaffner, 1926: 2, 8, 21, New England;
Howard, 1922: 135.

Acronicta dactylina (Grote)

Schaffner and Griswold, 1934: 58, 115 (ex *Acronycta*).

Acronicta fragilis (Guenée)

Reeks and Smith, 1956: 575, maritime prov.

Acronicta furcifera Guenée

Schaffner and Griswold, 1934: 55-56, 115 (ex *Acronycta*);
Webber and Schaffner, 1926: 2, 8, 21, New England (ex
Apatela).

Acronicta grisea Walker

Reeks, Smith and Forbes, 1946: 14, maritime prov.

Acronicta hasta Guenée

All records ex *Apatela hasta*.

Howard, 1922: 135; Culver, 1919: 5, New England.

Acronicta impleta Walker

Schaffner and Griswold, 1934: 55, 115 (ex *Acronycta*).

Acronicta impressa Walker

Schaffner and Griswold, 1934: 58-59, 115 (ex *Acronycta*).

Acronicta lanceolaria (Grote)

Schaffner and Griswold, 1934: 60, 115 (ex *Acronycta*).

Acronicta oblinita (J. E. Smith)

Reeks and Smith, 1956: 575, maritime prov. (ex *A.*
"obliniata"); Reeks, Smith and Forbes, 1946: 14,
maritime prov. (ex *A.* "obliniata"); Schaffner and
Griswold, 1934: 60, 115 (ex *Acronycta*).

Acronicta superans Guenée

Schaffner and Griswold, 1934: 55, 115 (ex *Acronycta*).

Acronicta spp.

Most records ex *Apatela* spp.

*Procter, 1946: 431, Maine (as "*Compsiluria*," Procter,
1938); Procter, 1938: 372, [Maine] (as "*Comp-*
siluria"); Schaffner and Griswold, 1934: 60, 115 (ex
Acronycta); Webber and Schaffner, 1926: 2, 8, New
England.

Alypia octomaculata (Fabricius)

Schaffner and Griswold, 1934: 41, 115; Webber and
Schaffner, 1926: 2, 8, 22, New England; Howard, 1922:
135.

Amphipyra pyramidoides Guenée

Schaffner and Griswold, 1934: 52, 115; Webber and
Schaffner, 1926: 3, 10, 21, New England (ex *Pyrophila*).

Calpe canadensis Bethune

Schaffner and Griswold, 1934: 70-71, 115, Mass.; Webber and Schaffner, 1926: 2, 8, 21, New England.

Catocala spp.

Webber and Schaffner, 1926: 2, 8, 22, New England.

Ceramica picta (Harris)

Most records ex *Mamestra picta*.

Schaffner and Griswold, 1934: 47, 115 (ex *Ceramica*); Webber and Schaffner, 1926: 3, 5-6, 9, 21, New England; Howard, 1922: 136; Culver, 1919: 5, New England.

Euthisanotia grata (Fabricius)

Webber and Schaffner, 1926: 3, 9, 20, New England.

Lithophane antennata (Walker)

Schaffner and Griswold, 1934: 49, 115 (ex *Graptolitha*).

Lithophane spp. (including *L. disposita* Morrison, *L. innominata* (Smith) and/or *L. petulca* Grote)

Schaffner and Griswold, 1934: 48, 115 (ex *Graptolitha* spp. (including *G. disposita*, *G. innominata* and/or *G. petulca*)).

Lithophane sp.

Reeks and Smith, 1956: 575, maritime prov. (ex *Graptolitha*).

Phosphila turbulenta Hübner

Webber, 1937: 144-149 (ex *Hadena*, lab.); Schaffner and Griswold, 1934: 52-53, 115 (ex *Trachea*).

Plathypena scabra (Fabricius)

Whiteside, Burbutis and Kelsey, 1967: 327, Del. (as "*Compilura*"); Schaffner and Griswold, 1934: 71-72, 115, Mass.; Webber and Schaffner, 1926: 3, 9, 22, New England; Hill, 1925: 17, Mass.

Plusiodonta compressipalpis Guenée

Webber and Schaffner, 1926: 3, 5, 9, 22, New England; Culver, 1919: 5, Mass.

Polia adjuncta (Boisduval)

Most records ex *Mamestra adjuncta*.

Schaffner and Griswold, 1934: 44-45, 115 (ex *Polia*); Webber and Schaffner, 1926: 3, 9, 21, New England; Howard, 1922: 135; Culver, 1919: 5, New England.

Polia legitima (Grote)

Schaffner and Griswold, 1934: 45, 115; Webber and Schaffner, 1926: 3, 5, 9, 21, New England (ex *Mamestra*).

Pseudaletia unipuncta (Haworth)

All records ex *Cirphis unipuncta*.

Franklin, 1950: 57; Schaffner and Griswold, 1934: 47, 115; Webber and Schaffner, 1926: 2, 8, 22, New England; Howard, 1922: 135; Culver, 1919: 5, New England.

Rhodophora florida Guenée

Webber and Schaffner, 1926: 3, 9, 22, New England; Culver, 1919: 5, Mass.

Scoliopteryx libatrix (Linnaeus)

Schaffner and Griswold, 1934: 71, 115; Webber and Schaffner, 1926: 3, 9, 21, New England.

Simyra henrici (Grote)

Schaffner and Griswold, 1934: 61, 115; Webber and Schaffner, 1926: 2, 6, 8, 21, New England (ex *Arsilonche albovenosa* (Goeze)).

Trichoplusia ni (Hübner)

Most records ex *Autographa brassicae* Riley.

Sutherland, 1966: 38-39, N.Y. (ex *Trichoplusia ni*); Harcourt, 1963: 72, Ont. (ex *Trichoplusia ni*); Webber, 1937: 144-149 (lab.); Schaffner and Griswold, 1934: 69, 115; Webber and Schaffner, 1926: 2, 8, 20, New England; Culver, 1919: 5, New England.

Xylena sp.

Culver, 1919: 5, New England (ex *Xylina*).

Unidentified Catocalinae

Schaffner and Griswold, 1934: 67, 115.

Unidentified Noctuidae

Schaffner, 1934: 589-590; Schaffner and Griswold, 1934: 72, 115; Culver, 1919: 5, New England.

Notodontidae

Cerura occidentalis Linter

Schaffner and Griswold, 1934: 82, 115-116; Webber and Schaffner, 1926: 2, 8, 19, New England.

Datana angusii Grote and Robinson

Schaffner and Griswold, 1934: 74-75, 115; Webber and Schaffner, 1926: 2, 9, 18, New England.

Datana integerrima Grote and Robinson

Raizenne, 1952: 82, Ont.; Schaffner, 1934: 591, Pa.; Schaffner and Griswold, 1934: 76, 115-116; Schaffner, 1927: 729; Webber and Schaffner, 1926: 2, 9, 18, New England.

Datana major Grote and Robinson

Schaffner and Griswold, 1934: 75, 115-116; Webber and Schaffner, 1926: 2, 9, 18, New England.

Datana ministra (Drury)

Reeks and Smith, 1956: 575, maritime prov.; Raizenne, 1952: 80, Ont.; Reeks, Smith and Forbes, 1946: 14, maritime prov.; Beaulne, 1939: 120, Que.; Webber, 1937: 144-149 (lab.); Schaffner and Griswold, 1934: 74, 115; Schaffner, 1927: 729; Webber and Schaffner, 1926: 2, 9, 18, New England; Howard, 1922: 135.

Datana perspicua Grote and Robinson

Schaffner, 1934: 591, Pa.; Schaffner and Griswold, 1934: 75, 115-116; Webber and Schaffner, 1926: 2, 9, 18, New England.

Fentonia marthesia (Cramer)

Schaffner and Griswold, 1934: 83, 115-116.

Heterocampa guttivitta (Walker)

Schaffner and Griswold, 1934: 79-80, 115-116; Collins, 1926: 693, 695, Mass., N.H.; Webber and Schaffner, 1926: 3, 9, 19, New England.

Heterocampa umbrata Walker

Schaffner and Griswold, 1934: 79, 115-116; Webber and Schaffner, 1926: 3, 9, 19, New England.

Ichthyura inclusa Hübner

All records ex *Melalopha inclusa*.

Webber, 1937: 144-149 (lab.); Schaffner, 1934: 589; Schaffner and Griswold, 1934: 73, 115; Webber and Schaffner, 1926: 3, 9, 19, New England; Howard, 1922: 136; Culver, 1919: 5, New England.

Nadata gibbosa (J. E. Smith)

Schaffner and Griswold, 1934: 78, 115-116; Webber and Schaffner, 1926: 3, 9, 21-22, New England.

Pheosia rimosa Packard

Schaffner and Griswold, 1934: 77, 115-116; Webber and Schaffner, 1926: 3, 9, 19, New England.

Schizura badia (Packard)

Schaffner and Griswold, 1934: 81, 115-116, Mass.

Schizura concinna (J. E. Smith)

Raizenne, 1952: 90, Ont.; Webber, 1937: 144-149 (ex *S. "cincinna,"* lab.); Schaffner and Griswold, 1934: 81, 115-116; Schaffner, 1927: 730; Webber and Schaffner, 1926: 3, 6, 10, 19, New England; Howard, 1922: 135; Culver, 1919: 5, New England.

Schizura ipomaeae Doubleday

Raizenne, 1952: 89, Ont. (ex *S. "ipomoeae"*).

Schizura unicornis (J. E. Smith)

Schaffner and Griswold, 1934: 81-82, 115-116; Webber and Schaffner, 1926: 3, 5-6, 10, 19, New England.

Symmerista albifrons (J. E. Smith)

Webber, 1937: 144-149 (lab.); Schaffner and Griswold, 1934: 115-116; Webber and Schaffner, 1926: 26, Mass., New England.

Symmerista canicosta Franclemont

Raizenne, 1952: 85, Ont.

Unidentified Notodontidae

Schaffner, 1934: 589-590; Culver, 1919: 5, New England.

Nymphalidae

Agraulis vanillae incarnata (Riley)

Arnaud, 1969: 77, Calif.

Chlosyne harrisii (Scudder)

Schaffner and Griswold, 1934: 9, 115 (ex *Lemonias*).

Chlosyne nycteis (Doubleday)

Webber and Schaffner, 1926: 2, 8, 14, 17, New England (ex *Charidryas*).

Euphydryas phaeton (Drury)

Webber, 1937: 144-149 (lab.); Schaffner and Griswold, 1934: 9, 115.

Limenitis archippus (Cramer)

All records ex *Basilarchia archippus*.

Schaffner and Griswold, 1934: 13-14, 115; Webber and Schaffner, 1926: 2, 8, 18, New England; Howard, 1922: 135.

Limenitis astyanax (Fabricius)

Schaffner and Griswold, 1934: 13, 115 (ex *Basilarchia arthemis astyanax*); Webber and Schaffner, 1926: 2, 8, 18, New England (ex *Basilarchia*).

Nymphalis antiopa (Linnaeus)

Reeks and Smith, 1956: 575, maritime prov.; Raizenne, 1952: 4, Ont.; Reeks, Smith and Forbes, 1946: 14, maritime prov.; Webber, 1937: 144-149 (ex *Hamadryas*, lab.); Dowden, 1935: 509 (ex *Hamadryas*); Schaffner and Griswold, 1934: 11, 115 (ex *Hamadryas*); Schaffner, 1927: 729 (ex *Euvanessa*); Webber and Schaffner, 1926: 3, 9, 14-15, New England (ex *Euvanessa*); Howard, 1922: 135 (ex *Euvanessa*); Culver, 1919: 5, New England (ex

Vanessa); Smith, H. E., 1917: 57-58, New England (ex *Euvanessa*).

Nymphalis milberti (Godart)

Most records ex *Aglais milberti*.

Schaffner and Griswold, 1934: 11, 115 (ex *Hamadryas*);
Webber and Schaffner, 1926: 2, 8, 16, New England;
Howard, 1922: 135.

Nymphalis vau-album j-album (Boisduval and LeConte)

Schaffner and Griswold, 1934: 10-11, 115, Mass., N.H. (ex *Hamadryas j-album*); Webber and Schaffner, 1926: 3, 9, 14, New England (ex *Eugonia j-album*).

Polygonia comma (Harris)

Raizenne, 1952: 2, Ont.; Schaffner and Griswold, 1934: 10, 115; Webber and Schaffner, 1926: 3, 9, 16, New England.

Polygonia interrogationis (Fabricius)

Raizenne, 1952: 2, Ont.; Schaffner and Griswold, 1934: 9-10, 115; Webber and Schaffner, 1926: 3, 9, 14-15, New England; Howard, 1922: 135.

Vanessa atalanta (Linnaeus)

Schaffner and Griswold, 1934: 12, 115 (ex *Cynthia*); Webber and Schaffner, 1926: 3, 10, 16-17, New England, N.H.; Howard, 1922: 135; Culver, 1919: 5, New England.

Vanessa virginiensis (Drury)

Most records ex *Vanessa huntera* (Fabricius).

Schaffner and Griswold, 1934: 12, 115 (ex *Cynthia virginiensis*); Webber and Schaffner, 1926: 3, 10, 16, New England; Culver, 1919: 5, New England.

Unidentified Nymphalidae

Schaffner, 1934: 589.

Oecophoridae

Dasytoma salicellum (Hübner)

Raine, 1966: 334, B.C.

Papilionidae

Battus philenor (Linnaeus)

Schaffner and Griswold, 1934: 6, 115, Conn., Mass., N.J., R.I. (ex *Papilio*).

Papilio glaucus Linnaeus

Most records ex *Papilio turnus* Linnaeus.

Schaffner and Griswold, 1934: 7, 115 (ex *P. glaucus* form

turnus); Webber and Schaffner, 1926: 3, 9, 24, New England; Howard, 1922: 135.

Papilio polyxenes polyxenes Fabricius

Most records ex *Papilio polyxenes*.

Schaffner and Griswold, 1934: 6, 115 (ex *P. ajax* Linnaeus); Webber and Schaffner, 1926: 3, 5, 7, 9, 24, New England; Howard, 1922: 135; Culver, 1919: 5, 22, New England.

Papilio troilus Linnaeus

Webber, 1937: 144-149 (lab.); Schaffner and Griswold, 1934: 7, 115; Webber and Schaffner, 1926: 3, 5, 7, 9, 24, New England; Howard, 1922: 135.

Unidentified Papilionidae

Schaffner, 1934: 589-590.

Pieridae

Pieris rapae (Linnaeus)

Fitzpatrick, 1967: 15, B.C.; Wilkinson, 1966: 1012-1013, B.C.; Anon., 1963b: 19, Ont.; Harcourt, 1963: 65, Ont.; Beirne, 1962: 19, Ont.; Webber, 1937: 144-149 (ex *Ascia*, lab.); Schaffner and Griswold, 1934: 7-8, 115 (ex *Ascia*); Webber and Schaffner, 1926: 3, 5-6, 9-10, 20, 24, New England (ex *Pontia*); Howard, 1922: 136 (ex *Pontia*); Culver, 1919: 1-27, figs. 1, 5-8, fig. 2 on pl. 1, New England (ex *Pontia*); Howard and Fiske, 1911: 223-224 (ex *Pontia*).

Pterophoridae

Oidaematophorus homodactylus (Walker)

Schaffner, 1959: 19, 50, [N.E. U.S.].

Pyralidae

Evergestis straminealis (Hübner)

Schaffner, 1959: 8, 50, [N.E. U.S.]; Webber and Schaffner, 1926: 3, 9, 22, New England.

Ostrinia nubilalis (Hübner)

Webber and Schaffner, 1926: 27, Mass., New England (ex *Pyrausta*).

Saturniidae

Antheraea polyphemus (Cramer)

Most records ex *Telea polyphemus*.

Reeks and Smith, 1956: 575, maritime prov. (ex *Antheraea*); Schaffner and Griswold, 1934: 28, 115;
Webber and Schaffner, 1926: 3, 10, 24, New England.

Automeris io (Fabricius)

Raizenne, 1952: 16, Ont.; Schaffner and Griswold, 1934: 28, 115; Schaffner, 1927: 729; Webber and Schaffner, 1926: 2, 8, 24, New England, N.Y.

Callosamia promethea (Drury)

Webber, 1937: 144-149 (lab.); Schaffner and Griswold, 1934: 27, 115; Webber and Schaffner, 1926: 2, 5-6, 8, 24, New England; Culver, 1919: 1-27, figs. 1, 5-8, fig. 2 on pl. 1, Mass.; Smith, H. E., 1917: 58, Mass.

Hemileuca maia (Drury)

Webber and Schaffner, 1926: 3, 9, 24, New England.

Hemieuca oliviae Cockerell

Caffrey in Britton, 1916: 250, N. Mex. (ex range caterpillar).

Hemileuca spp. (including *H. lucina* H. Edwards and/or *H. maia* (Drury))

Schaffner and Griswold, 1934: 29, 115.

Platysamia cecropia (Linnaeus)

Webber, 1937: 144-149 (lab.); Schaffner and Griswold, 1934: 26-27, 115; Webber and Schaffner, 1926: 3, 10, 24, 29, New England (ex *Samia*).

Unidentified Saturniidae

Schaffner, 1934: 589-590.

Sphingidae

Ampeloeca myron (Cramer)

Schaffner and Griswold, 1934: 23-24, 115; Webber and Schaffner, 1926: 2, 5-6, 8, New England (ex *Ampelophaga*).

Amphion nessus (Cramer)

Schaffner and Griswold, 1934: 25, 115.

Atreides plebeius (Fabricius)

Aldrich, 1932: 3, Mass. (ex *Atreus* "plebeja").

Ceratomia catalpae (Boisduval)

Schaffner and Griswold, 1934: 17, 115, N.J.

Ceratomia undulosa (Walker)

Schaffner and Griswold, 1934: 17, 115.

Deidamia inscriptum (Harris)

Schaffner and Griswold, 1934: 24, 115; Webber and Schaffner, 1926: 2, 5, 9, 25, New England; Culver, 1919: 5, 22, New England.

Hemaris thysbe (Fabricius)

Schaffner and Griswold, 1934: 22, 115; Webber and Schaffner, 1926: 3, 9, 25, New England.

Hyles gallii (Rottenburg)

Most records ex *Deilephila gallii*.

Schaffner and Griswold, 1934: 25, 115 (ex *Sphinx gallii intermedia* Kirby); Webber and Schaffner, 1926: 2, 5, 9, 25, New England; Culver, 1919: 5, 22, Mass.

Hyles lineata (Fabricius)

Schaffner and Griswold, 1934: 25, 115 (ex *Sphinx*); Webber and Schaffner, 1926: 2, 9, 25, New England (ex *Deilephila*).

Manduca quinquemaculata (Haworth)

Most records ex *Phlegethontius quinquemaculata*.

Baird, 1943: 111, 125, 128, Que. (ex *Protoparce*); Baird, 1942: 116, 130, 133, U.S.; Baird, 1940: 99, 118, 123, N.B., N.S.; Baird, 1939: 106, 122, 126, N.B., N.S.; Gilmore, 1938: 711, Ky. and/or Tenn. (ex *Protoparce*); Schaffner and Griswold, 1934: 16, 115; Webber and Schaffner, 1926: 3, 9, 25, New England; Howard, 1922: 136; Culver, 1919: 5, New England.

Paonias excaecata (J. E. Smith)

Webber, 1937: 144-149 (lab.); Schaffner and Griswold, 1934: 21, 115; Schaffner, 1927: 730.

Paonias myops (J. E. Smith)

Webber, 1937: 144-149 (lab.); Schaffner and Griswold, 1934: 21, 115; Webber and Schaffner, 1926: 3, 5-6, 9, New England.

Pholus achemon (Drury)

Schaffner and Griswold, 1934: 23, 115; Webber and Schaffner, 1926: 3, 9, New England.

Pholus satellitia pandorus (Hübner)

Schaffner and Griswold, 1934: 23, 115.

Smerinthus cerisyi Kirby

Schaffner and Griswold, 1934: 21, 115, Maine.

Smerinthus jamaicensis (Drury)

Schaffner and Griswold, 1934: 20, 115 (ex *S. geminatus* Say).

Sphecodina abbottii Swainson

Schaffner and Griswold, 1934: 24, 115; Webber and Schaffner, 1926: 3, 10, New England.

Sphinx chersis (Hübner)

Schaffner and Griswold, 1934: 18, 115 (ex *Hyloicus*).

Sphinx drupiferarum J. E. Smith

Schaffner and Griswold, 1934: 20, 115 (ex *Hyloicus*).

Sphinx gordius Cramer

Webber, 1937: 144-149 (lab.); Schaffner and Griswold, 1934: 19, 115 (ex *Hyloicus*); Webber and Schaffner, 1926: 3, 5-6, 10, New England.

Unidentified Sphingidae

Schaffner, 1934: 589-590; Schaffner and Griswold, 1934: 26, 115; Webber and Schaffner, 1926: 3, 10, New England.

Tortricidae

Archips argyrospilus (Walker)

Paradis and LeRoux, 1965: 44, 46, 49-50, fig. 14F, Que.;
Paradis, 1961: 63, Que.

Archips cerasivoranus (Fitch)

Schaffner, 1959: 30, 50, [N.E. U.S.].

Choristoneura fumiferana (Clemens)

Dowden, Buchanan and Carolin, 1948: 459 (ex *Archips*, lab.).

Unidentified Lepidoptera

Cabbage worm(-s)

O'Kane, 1916: 132, N.H.

Miscellaneous

Crossman and Webber, 1924: 67, 69, Mass.; Philbrook, 1918: 60, Maine (ex injurious caterpillars).

STAGE PARASITIZED: Egg (Lepidoptera [Pouillaude, 1913: 82-83]), larva (Hymenoptera, Lepidoptera), pupa (Lepidoptera).

IMMATURES: First instar larva — Culver, 1919: (descr., figs. 1, 5).

Second instar larva — Culver, 1919: (descr., fig. 6).

Third instar larva — Culver, 1919: (descr., fig. 7).

Pupa — Paradis and LeRoux, 1965: (fig. 14F); Ross, 1953: (descr., fig. 8); Burgess and Crossman, 1929: (fig. 24); Culver, 1919: (descr., fig. 8, fig. 2 on pl. 1); McLaine, 1916: (fig. 4); O'Kane, 1916: (pl. 4); Burgess, 1915a: (fig. 2 on pl. 3); Howard and Fiske, 1911: (fig. 1 on pl. 20); Thompson, 1910: (descr.).

BIOLOGY: Paradis and LeRoux, 1965: (44, 46, 49-50); Beirne, 1962: (19); Dowden, 1962: (30, 32, 34, 43-45, 54); Evans, 1962: (597); Schaffner, 1959: (6-8, 19, 30, 50); Clausen, 1956: (6-7, 63, 97-98); Franklin, 1950: (57, 60); Sellers, 1943: (56); Sweetman, 1936: (121-123, 221, 269); Dowden, 1935: (509); Proper, 1934: (369-371, 375-376); Schaffner, 1934: (588-592); Schaffner and Griswold, 1934: (115); Burgess and Crossman, 1929: (52-67, 142); Burgess and Crossman, 1927: (16-18); Webber and Schaffner, 1926: (3, 5-6, 9-10, 20, 24); Burgess, 1924: (34-35); Culver, 1919: (1-27); Tothill, 1922: (9, 29, 36-43, 107); Smith, H. E., 1917: (57-58); McLaine, 1916: (22, 24); Burgess, 1915: (85); Tothill, 1913a: (57); Howard and Fiske, 1911: (218-225, 308); Thompson, 1910: (292).

Compsilura "concinati," see *Compsilura concinnata*

Compsilura "conncinata," see *Compsilura concinnata*

Compsilura oppugnator, see *Eucelatoria australis*

Compsilura [sp.], see *Compsilura concinnata* or *Compsilura* sp. probably

Compsilura sp. probably

HOST: LEPIDOPTERA

Lasiocampidae

Malacosoma disstria Hübner

Hawboldt, 1944: 23, N.B.; Reeks, 1943: 213, N.B.

STAGE PARASITIZED: Larva

"*Compsiluria*" *concinata*, see *Compsilura concinnata*

"*Consilura* *opugnator*," see *Eucelatoria australis*

Copecrypta Townsend (Tachininae: Cuphocerini)

Copecrypta ruficauda (Wulp)

HOST: LEPIDOPTERA

Noctuidae

Ogdoconta cinereola (Guenée)

Chittenden, 1923a: 156, D.C. (as *Cuphocera*).

Plathypena scabra (Fabricius)

Hill, 1925: 17, fig. 15, Miss., Mo., N.C., Tenn. (as *Trichophora*).

STAGE PARASITIZED: Larva, pupa.

IMMATURES: Pupa — Hill, 1925: (fig. 15); Greene, 1922: (descr., fig. 97 on pl. 20).

BIOLOGY: Hill, 1925: (17)

Copecrypta sp.

HOST: LEPIDOPTERA

Aegeriidae

Sylvora acerni (Clemens)Brauer and Bergenstamm, 1895: 552, 581, N. Amer. (as
Elachipalpus ex Sesia).**Coquillettina plankii**, see *Hemithrixion plankii***Coronimyia Townsend** (Goniinae: Siphonini)**Coronimyia geniculata** Townsend

HOST: LEPIDOPTERA

Pyralidae

Moodna ostrinella (Clemens)

Reinhard, 1946: 91, Ont.

Unidentified Phycitinae

Reinhard, 1946: 91, N.C. (ex phycid in dried peach).

Pyralidae or Tortricidae

Unidentified Pyralidae or Tortricidae

Reinhard, 1946: 91, Miss.; Allen, 1929: 684, Miss. (as
Epigrammyia).

STAGE PARASITIZED: Larva.

Coronimyia melitarae Reinhard

HOST: LEPIDOPTERA

Pyralidae

Melitara sp.

Reinhard, 1946: 92, Calif.

Crossocosmia sericariae, see *Blepharipa schineri***Cryptomeigenia** Brauer and Bergenstamm (Goniinae: Blondeliini)**Cryptomeigenia aurifacies** Walton

HOST: COLEOPTERA

Scarabaeidae

Chemarachis vandinei (Smyth)Most records ex *Phyllophaga vandinei*.Martorell, 1945a: 431-432, P.R. (as *Emphanopteryx*);Wolcott, 1924: 221, P.R. (ex *Lachnosterna*); Smyth,

1917: 82, 86-87, pl. 7, P.R.; Smyth, 1917a: 151, P.R.

Lachnosterna crenaticollis Blanchard probably
Walton, 1913: 200, P.R. (ex *L. "crenatocollis"* probably).

Phyllophaga portoricensis Smyth

Wolcott, 1933: 186, P.R.; Wolcott, 1924: 221, P.R. (ex
Lachnosterna); Smyth, 1917: 86, pl. 7, P.R.; Smyth,
1917a: 151, P.R.

Phyllophaga spp.

Most records ex *Lachnosterna* spp.

Wolcott, 1951a: 473-474, P.R.; Wolcott, 1935: 449, P.R.;
Wolcott, 1924a: 27, P.R.; Wolcott, 1922: 15-16, P.R.;
Van Zwaluwenburg, 1917: 517, P.R. (ex *Phyllophaga*);
Van Dine, 1913a: 37, P.R.; Van Dine, 1913b: 200,
P.R.

Scarabaeid spp. (including *Phyllophaga* spp. or *Phytalis* spp.)

Smyth, 1917: 55-56, pl. 7, P.R. (ex white grubs).

STAGE PARASITIZED: Adult.

IMMATURES: Pupa — Greene, 1922: (descr., fig. 55 on pl. 12);
Smyth, 1917: (pl. 7).

BIOLOGY: Wolcott, 1951a: (473-474); Martorell, 1945a:
(431-432); Wolcott, 1935: (449); Smyth, 1917: (82, 86-87).

Cryptomeigenia prisca (Walker)

HOST: COLEOPTERA

Scarabaeidae

May beetle(-s)

Johnson, 1925: 185, Maine, Mass., N.H. (as *Emphanop-
teryx*).

Cryptomeigenia simplex Curran

HOST: COLEOPTERA

Scarabaeidae

Phyllophaga spp.

Hammond, 1945: 159, Ont.

Cryptomeigenia theutis (Walker)

HOST: COLEOPTERA

Scarabaeidae

Phyllophaga crassissima (Blanchard)

Hayes, 1920: 317, Kans. (ex *Lachnosterna*).

Phyllophaga crenulata (Frölich)

Hayes, 1920: 317, Kans. (ex *Lachnosterna*).

Phyllophaga futilis (LeConte)Hayes, 1920: 317, Kans. (ex *Lachnosterna*).*Phyllophaga implicita* (Horn)Hayes, 1920: 317, Kans. (ex *Lachnosterna* "implicita").*Phyllophaga inversa* (Horn)All records ex *Lachnosterna inversa*.Walton, 1913: 198, D.C.; Johnson, 1910: 775, N.J.;
Chittenden, 1899: 79; Coquillett, 1897: 11, 23, D.C.*Phyllophaga rugosa* (Melsheimer)Hayes, 1920: 317, Kans. (ex *Lachnosterna*).*Phyllophaga* spp.Hammond, 1945a: 159, Ont.; Hammond, 1934: 66, Que. (ex
June beetle); Hammond, 1934b: 130, Que. (ex June
beetles); Criddle, 1918: 454, Man. (ex *Lachnosterna*);
Davis, J. J., 1918: 14 (ex *Lachnosterna*); Smyth, 1917:
59, U.S.; Davis, J. J., 1916: 271, N.E. U.S. (ex
Lachnosterna); Aldrich, 1915: 83 (ex *Lachnosterna*);
Shannon, 1914: 182, Md. (ex *Lachnosterna*).

STAGE PARASITIZED: Adult.

IMMATURES: Pupa — Greene, 1922: (descr., fig. 53 on pl. 12).

BIOLOGY: Hammond, 1934: (66); Hammond, 1934b: (130);
Criddle, 1918: (454).*Cryptomeigenia triangularis* Curran

HOST: COLEOPTERA

Scarabaeidae

Phyllophaga spp. (including *P. anxia* (LeConte) and/or *P. fusca*
(Frölich))

Hammond, 1945b: 159, Ont.

Cryptomeigenia spp.

HOST: COLEOPTERA

Scarabaeidae

Phyllophaga anxia (LeConte)Hammond, 1936a: 84 (ex white grub); Hammond, 1934a:
99, Que. (ex common white grub).*Phyllophaga* spp.Hammond, 1941: 186-187, Ont., Que. (ex June beetles);
Hammond 1936a: 84, Ont. (ex June beetles);
Hammond, 1934c: 130, Que. (ex June beetles).

May beetle(-s)

Hooker, 1913: 35, P.R.

HOST: HYMENOPTERA

Diprionidae

Diprionid spp. (including *Diprion* spp. and/or *Neodiprion* spp.)
Brauer and Bergenstamm, 1895: 552, 597 (as *Eumyothyria*
ex *Lophyrus*).

STAGE PARASITIZED: Larva (Coleoptera), adult (Coleoptera).

BIOLOGY: Hammond, 1936a: (84); Hammond, 1934c: (130).

Ctenophorocera sp., see *Pales* spp.

Cuphocera miscelli, see *Deopalpus miscelli*

Cuphocera parksi, see *Deopalpus parksi*

Cuphocera ruficauda, see *Copecrypta ruficauda*

Cylindromyia Meigen (Phasiinae: Cylindromyiini)

Cylindromyia armata Aldrich

HOST: HEMIPTERA (HETEROPTERA)

Pentatomidae

Chlorochroa sayi Stål

Strand, 1937: 26, Mont. (ex Say's plant bug).

Cylindromyia binotata (Bigot)

HOST: HEMIPTERA (HETEROPTERA)

Pentatomidae

Euschistus servus (Say)

Rolston and Kendrick, 1961: 156, Ark.

Euschistus variolarius (Palisot de Beauvois)

Rings and Brooks, 1958: 15-16, Ohio.

HOST: LEPIDOPTERA

Saturniidae

Actias luna (Linnaeus)

Fattig, 1949: 14, Ga.

STAGE PARASITIZED: Pupa (Lepidoptera), adult (Hemiptera).

BIOLOGY: Rolston and Kendrick, 1961: (156).

Cylindromyia "carolinae (Robineau-Desvoidy)"

All records as *Ocyptera carolinae*.

HOST: LEPIDOPTERA

Noctuidae

Pseudaletia unipuncta (Haworth)

Johnson, 1910: 779, N.J. (ex *Leucania*).

HOST: ORTHOPTERA

Acrididae

Grasshopper(-s)

Aldrich, 1915: 82.

IMMATURES: Pupa — Greene, 1922: (descr., fig. 20 on pl. 5).

Cylindromyia euchenor (Walker)Most records as *Ocyptera euchenor*.

HOST: HEMIPTERA (HETEROPTERA)

Pentatomidae

Chlorochroa ligata (Say)Russell, E. E., 1952: 15, Ariz. (as *Cylindromyia*).*Chlorochroa sayi* StålRussell, E. E., 1952: 15, Ariz. (as *Cylindromyia*).Pentatomid spp. (including *Euschistus ictericus* (Linnaeus),
Mormidea pictiventris Stål, *Nezara viridula* (Linnaeus)
and/or *Oebalus pugnax* (Fabricius))Thames, 1954: 236, Fla. (as *Cylindromyia* ex pentatomids,
E. ictericus, *M. pictiventris*, "Nezera" *viridula* and/or
Solubea pugnax).

HOST: LEPIDOPTERA

Noctuidae

Pseudaletia unipuncta (Haworth)All records ex *Leucania unipuncta*.Lintner, 1897: 210; *Brauer and Bergenstamm, 1895: 565,
596 (Townsend, 1893); Townsend, 1893: 466, Ill.

HOST: ORTHOPTERA

Acrididae

Unidentified Acrididae

*Brauer and Bergenstamm, 1895: 565, 580 (ex Acridiodea
G[enus], Townsend, 1893); Townsend, 1893: 466, Ill.

STAGE PARASITIZED: Larva (Lepidoptera), adult (Hemiptera).

Cylindromyia fumipennis (Bigot)

HOST: HEMIPTERA (HETEROPTERA)

Pentatomidae

Euschistus variolarius (Palisot de Beauvois)

Rings and Brooks, 1958: 15-16, Ohio.

STAGE PARASITIZED: Adult.

Cyrtophleba Rondani (Tachininae: Voriini)

Cyrtophleba coquilletti Aldrich

All records as *Cyrtophloeba coquilletti*.

HOST: LEPIDOPTERA

Noctuidae

Epiglaea apiata (Grote)

Schaffner and Griswold, 1934: 50, 108, Mass.

Eupsilia spp. (including *E. morrisoni* (Grote) and/or *E. sidus* (Guenée))

Schaffner and Griswold, 1934: 50-51, 108 (ex *Conistra* spp. (including *C. morrisoni* and/or *C. sidus*)).

Unidentified Noctuidae

Schaffner and Griswold, 1934: 72, 108, Mass.

BIOLOGY: Schaffner and Griswold, 1934: (108).

Cyrtophloeba coquilletti, see *Cyrtophleba coquilletti*

"*Cyzenia*" *albicans*, see *Cyzenis albicans*

Cyzenis Robineau-Desvoidy (Goniinae: Eryciini)

Cyzenis albicans (Fallén)

HOST: LEPIDOPTERA

Geometridae

Alsophila pometaria (Harris)

Embree and Sisojevic, 1965: 634, N.S. (unsuccessful lab. rearing).

Operophtera bruceata (Hulst)

Embree and Sisojevic, 1965: 631, 634, N.S. (lab.).

Operophtera brumata (Linnaeus)

Embree, 1966: 1159-1167, N.S.; Embree, 1965: 5, 21, 41, 51, 53-54, N.S.; Embree and Sisojevic, 1965: 631-639, N.S.; Cuming, 1964: 38, N.S. (ex winter moth); Embree, 1964: 40, N.S. (ex winter moth); Williamson, 1963: 139, 145, 147, N.S.; MacPhee, 1962: 106, N.S. (as "*Cyzenia*" ex winter moth); Maybee and Wylie, 1961: 851-855, Canada; Embree, 1960: 862-864, N.S.; Anon., 1958b: 139, N.S.; Graham, 1958: 595-596, N.S.; Anon., 1957a: 176, N.S.; Anon., 1956a: 198, N.S.; Anon., 1956b: 284, N.S.

Lasiocampidae

Malacosoma americanum (Fabricius)

Embree and Sisojevic, 1965: 634, N.S. (unsuccessful lab. rearing).

Olethreutidae

Pseudexentera improbana (Walker)

Embree and Sisojevic, 1965: 634, N.S. (ex *P. cressoniana* Clemens, unsuccessful lab. rearing).

STAGE PARASITIZED: Larva, pupa.

BIOLOGY: Embree, 1966: (1159-1167); Embree, 1965: (5, 21, 41, 51, 53-54); Embree and Sisojevic, 1965: (631-639); Maybee and Wylie, 1961: (851-855); Embree, 1960: (862-864); Anon., 1956b: (284).

"*Cyzenis festinans* (Aldrich and Webber)" probably

HOST: LEPIDOPTERA

Olethreutidae

Pseudexentera improbana oregonana (Walsingham)

Wong, H. R., and Melvin, 1967: 4, Man. (ex *Pseudexentera oregonana*).

STAGE PARASITIZED: Larva.

Cyzenis spp.

HOST: LEPIDOPTERA

Olethreutidae

Pseudexentera improbana oregonana (Walsingham)

Wong, H.R., and Melvin, 1967: 4, Man. (ex *Pseudexentera oregonana*).

STAGE PARASITIZED: Larva.

Dejeaniopalpus Townsend (Dexiinae: Leskiini)

Dejeaniopalpus texensis Townsend

HOST: LEPIDOPTERA

Pyralidae

Desmia funeralis (Hübner)

James, M. T., 1947: 108, D.C., Va.

Pyrausta spp. probably

All records ex *Botys* sp.

James, M. T., 1947: 108; Aldrich, 1924: 213, Va. (as *Genea*).

Tortricidae

Leaf roller(-s)

James, M. T., 1947: 108, Pa. (ex leaf roller on *Helianthus*).

STAGE PARASITIZED: Larva.
 Deopalpus Townsend (Tachininae: Cuphocerini)
 Deopalpus miscelli (Coquillett)
 HOST: LEPIDOPTERA

Noctuidae

Catabena lineolata Walker

All records ex *Adisophanes miscellus* (Grote).

*Reinhard, 1934a: 50, Calif. (as *Cuphocera*, Coquillett, 1897); Coquillett, 1897: 21, 23, Calif. (as *Trichophora*).

Deopalpus parksi (Reinhard)
 HOST: LEPIDOPTERA

Unidentified Lepidoptera

Miscellaneous

Reinhard, 1934a: 52, Tex. (as *Cuphocera*).

STAGE PARASITIZED: Larva.
 Dexia Meigen (Dexiinae: Dexiini)
 Dexia limata (Coquillett)
 HOST: DIPTERA

Tipulidae

Unidentified Tipulidae

Johnson, 1925: 188, Mass. (as *Neadmontia*).

STAGE PARASITIZED: Larva.
 Dexia ventralis, see *Dexilla ventralis*
 Dexia vertebrata, see *Zelia vertebrata*
 Dexilla Westwood (Proseninae: Dexillini)
 Dexilla ventralis (Aldrich)
 Most records as *Dexia ventralis*.
 HOST: COLEOPTERA

Scarabaeidae

Maladera castanea (Arrow)

Smith, L. B., 1929: 125, N.J. (ex Asiatic beetle).

Phyllophaga spp.

Hadley, 1938: 211, U.S. (lab.).

Popillia japonica Newman

Fleming, 1963: 25, N.J. (as *Dexilla*); Clausen, 1956: 6, 23, 102, N.J. (as *Dexilla*); King and Parker, 1941: 127, 131, N.J. (ex Japanese beetle); Gardner and Parker, 1940: 3,

8-10, 32, U.S.; King, 1939: 25, N.J. (ex Japanese beetle); Hadley, 1938: 211, N.J.; Sweetman, 1936: 124-126, 243, 259, fig. 105b, N. Amer.; Burrell, 1931: 323-336, fig. 7, N.J.; King, 1931: 453, 459-460, N.J.; King, 1930: 160, N.J.; Smith, L. B., 1930: 500, U.S. (ex Japanese beetle); Smith, L. B., 1929: 125, N.J. (ex Japanese beetle); Clausen, King and Teranishi, 1927: 4, 26-31, 54, figs. 14A, 20A-E, 21B, non-N. Amer.

STAGE PARASITIZED: Larva.

IMMATURES: First instar larva — Sweetman, 1936: (fig. 105b); Clausen, King and Teranishi, 1927: (descr., figs. 20A-E).

Third instar larva — Burrell, 1931: (fig. 7); Clausen, King and Teranishi, 1927: (descr., fig. 21B).

Pupa — Clausen, King and Teranishi, 1927: (descr., fig. 14A).

BIOLOGY: Clausen, 1956: (6, 23, 102); King and Parker, 1941: (127, 131); King, 1939: (25); Sweetman, 1936: (124-126, 243, 259); Burrell, 1931: (323-336); King, 1931: (453, 459-460); King, 1930: (160); Smith, L. B., 1930: (500).

Dexodes aurifrons, see *Oswaldia aurifrons*

Dexodes chaetoneura, see *Aporotachina chaetoneura*

Dexodes tenella, see *Lydellohoughia tenella*

Dexodes sp. (= ? *Masicera luctuosa*), see *Oswaldia* sp. probably

Dichaetoneura Johnson (Goniinae: Siphonini)

Dichaetoneura leucoptera Johnson

HOST: LEPIDOPTERA

Nymphalidae

Nymphalis antiopa (Linnaeus)

Knowlton and Hanson, 1949: 3, Utah (ex "*Euvanesia*").

Oecophoridae

Depressaria pastinacella (Duponchel)

All records ex *Depressaria heracliana* (Linnaeus).

Schaffner, 1959: 39, 50, [N.E. U.S.]; Thompson, 1953a: 391-392, fig. 1 (ex *D. "heracleana"*); Leonard, 1920: 491-492, N.Y.

Olethreutidae

Epiblema strenuanum (Walker)

Putman, 1935a: 107, Ont. (as *Phytomyptera*).

Grapholitha molesta (Busck)

Allen, 1962: 13, 112, Md.

Tortricidae

Archips cerasivoranus (Fitch)

Allen, 1962: 12; Schaffner, 1959: 30, 50, [N.E. U.S.]; Thompson, 1953: 19-30, pls. 1-3; Thompson, 1953a: 391-392, fig. 1; Raizenne, 1952: 210, Ont.; *Procter, 1946: 435, Maine (as *Phytomyptera*, Procter, 1938); Beaulne, 1939: 120, Que. (as "*Phytomyptera*"); Beaulne, 1939a: 95, Que. (as *Phytomyptera* ex "*Caecacia*"); Daviault, 1939: 184, [Que.] (as *Phytomyptera* ex *Cacoecia*); Procter, 1938: 376, Maine (as *Phytomyptera*); Putman, 1935a: 108, Ont. (as *Phytomyptera* ex *Cacoecia*); Greene, 1934: 29-30, Conn., Maine, N.H., probably N.Y.; Johnson, 1927: 198, Maine; Johnson, 1925: 187, Conn., Maine, Mass., N.H., Vt.; Leonard, 1920: 492, Maine, Mass., N.Y.; Baird, 1918: 768, probably Alta. or N.B.; Gibson, 1916: 216, Ont.; Johnson, 1916: 81, Vt.; Herrick, 1912: 291, N.Y.; Johnson, 1907: 9-10, Maine, Mass.; Patch, 1907: 265, Maine.

Archips fervidanus (Clemens)

Strohecker and Yackley, 1963: 1354, Pa.; Schaffner, 1959: 29, 50, [N.E. U.S.]; *Procter, 1946: 435, Maine (as *Phytomyptera* ex *A. "fervadana,"* Procter, 1938); Procter, 1938: 376, Maine (as *Phytomyptera* ex *A. "fervadana"*); Johnson, 1927: 198, Maine (ex *A. "fervadana"*); Johnson, 1925: 187, Conn., Maine, Mass., N.H., Vt. (ex *A. "fervadana"*); Leonard, 1920: 492, Maine; Patch, 1907: 265, Maine.

STAGE PARASITIZED: Larva, pupa.

IMMATURES: First instar larva — Thompson, 1953: (descr., figs. 1-2 on pl. 1, figs. 11-13 on pl. 3); Thompson, 1953a: (descr., fig. 1).

Second instar larva — Thompson, 1953: (descr., fig. 4 on pl. 2, fig. 14 on pl. 3).

Third instar larva — Thompson, 1953: (descr., figs. 5-7 on pl. 2, fig. 15 on pl. 3).

Pupa — Thompson, 1953: (descr., fig. 17 on pl. 3); Greene, 1922: (descr., fig. 77 on pl. 16).

BIOLOGY: Allen, 1962: (12-13, 112); Schaffner, 1959: (29-30, 39, 50); Thompson, 1953: (19-30); Baird, 1918: (768).

Didyma Wulp (Goniinae: Blondeliini)

Didyma exigua, see *Didyma exigua* or *Hyphantrophaga hyphantriae*

Didyma exigua Wulp

HOST: LEPIDOPTERA

Pyralidae

Acrobasis caryae Grote

Gill, 1930: 72 (ex pecan nut case-bearer).

Ostrinia obliteralis (Walker)

Rowe, 1933: 122, Ill. (ex "*Pipausta ainslei*" Heinrich).

Didyma inconspicua, see *Paralipse infernalis*

Digonichaeta setipennis, see *Bigonicheta spinipennis*

Digonochaeta setipennis, see *Bigonicheta spinipennis*

Digonochaeta sp., see *Bigonicheta spinipennis*

Dinera Robineau-Desvoidy (Proseninae: Prosenini)

Dinera grisescens (Fallén) probably

HOST: COLEOPTERA

Scarabaeidae

Phyllophaga spp.

Criddle, 1918: 454, Man. (as *Myiocera "cremiodes"* (Walker) ex *Lachnosterna*).

STAGE PARASITIZED: Larva.

Diplostichus Brauer and Bergenstamm (Goniinae: Exoristini)

Diplostichus hamata, see *Diplostichus lophyri*

Diplostichus lophyri (Townsend)

HOST: HYMENOPTERA

Diprionidae

Diprion frutetorum (Fabricius)

Raizenne, 1957: 24, Ont. (as *Neophorocera hamata* (Aldrich and Webber)).

Diprion hercyniae (Hartig)

Bird and Burk, 1961: 235, Ont. (as *D. hamatus*); Finlayson, T., 1960a: 923, 937, Que. (as *D. hamatus*); Raizenne, 1957: 24, Ont. (as *Neophorocera hamata*); Daviault, 1944: 40, Que. (as *Phorocera hamata* ex *G[ilpinia]*); Daviault, 1941: 15, Que. (as *Phorocera hamata* ex [European spruce sawfly]).

Diprion polytomum (Hartig)

Reeks, 1938: 26, N.B. (as *Phorocera hamata*).

Diprion similis (Hartig)

Finlayson, T., 1962: 273, 280, Canada (as *D. hamatus*); Johnson, 1925: 198, Conn., Mass. (as *Phorocera hamata*); Aldrich and Webber, 1924: 63, Conn. (as *Phorocera hamata*); Hartley, 1923: 386, Pa. (as *Exorista petiolata* Coquillett); Britton and Zappe, 1918: 283, Conn. (as *Exorista petiolata*); Britton, 1916: 282, Conn. (as *Exorista petiolata*); Britton, 1916a: 122, Conn. (as *Exorista petiolata*).

Neodiprion abbotii (Leach)

Finlayson, T., 1963: 476, 502 (as *D. hamatus*); Patton, 1958: 80, Fla. (as *Phorocera hamata* ex *N. fabricii* (Leach)); Raizenne, 1957: 21, Ont. (as *Neophorocera hamata*).

Neodiprion abietis (Harris)

Finlayson, T., 1963: 477, 502 (as *D. hamatus*); *Brauer and Bergenstamm, 1895: 569, 597 (as (*Phorocera*) ex *Lophyrus*, Townsend, 1892a); Townsend, 1892a: 289, Ont. (as *Phorocera* ex *Lophyrus*).

Neodiprion abietis (Harris) complex

Raizenne, 1957: 13, Ont. (as *Neophorocera hamata*).

Neodiprion lecontei (Fitch)

Finlayson, T., 1963: 479, 502 (as *D. hamatus*); Griffiths, 1961: 196-197, Ont. (as *D. hamatus*); Raizenne, 1957: 19-20, Ont., Que. (as *Neophorocera hamata*); Benjamin, 1955: 20, 23-24, 53, Ill., Mich., Que. (as *Phorocera hamata*); Boyes and Wilkes, 1953: 126, 150 (as *Neophorocera hamata*); Daviault, 1953: 181-184, Que. (as *Neophorocera hamata*); Baldwin, W.F., and Coppel, 1949: 222-223, fig. 1, Canada (as *Phorocera hamata*); Baldwin, W. F., and Coppel, 1949a: 237-245, figs. 1-14, Ont. (as *Phorocera hamata*); Baird, 1944a: 157, Ont. (as *Neophorocera hamata*); Daviault, 1942: 397, Que. (as *Phorocera hamata*).

Neodiprion nanulus nanulus Schedl

Finlayson, T., 1963: 480, 502 (as *D. hamatus*); Kapler and Benjamin, 1960: 262, Wis. (as *Phorocera hamata*); Raizenne, 1957: 13, Ont. (as *Neophorocera hamata*); Coppel, 1954: 168, Ont. (as *Neophorocera hamata* ex *N. nanulus*); Baldwin, W. F., and Coppel, 1949a: 237-245, figs. 1-14, Ont. (as *Phorocera hamata* ex *N. nanulus*).

Neodiprion nigroscutum Middleton

Becker, Jr., and Benjamin, 1967: 158-159, Wis. (as *D. hamata*); Becker, Jr., and Benjamin, 1964a: 20, Wis. (as *D. hamata*).

Neodiprion pinetum (Norton)

Finlayson, T., 1963: 482, 502 (as *D. hamatus*); Raizenne, 1957: 21, Ont. (as *Neophorocera hamata*).

Neodiprion pratti banksianae Rohwer

Finlayson, T., 1963: 483, 502 (as *D. hamatus*); Griffiths, 1960: 654-656, Ont. (as *D. hamatus*); Raizenne, 1957: 15, Ont. (as *Neophorocera hamata*); Baldwin, W. F., and Coppel, 1949: 222-223, fig. 1, Canada (as *Phorocera hamata* ex *N. banksianae*); Baldwin, W. F., and Coppel, 1949a: 237-245, figs. 1-14, Ont. (as *Phorocera hamata* ex *N. banksianae*).

Neodiprion pratti pratti (Dyar)

Bobb, 1963: 619-620, Va. (as *Phorocera hamata*).

Neodiprion rugifrons Middleton

Wilkinson, R. C., Becker, and Benjamin, 1966: 791, Wis. (as *D. hamata*); Raizenne, 1957: 22, Ont. (as *Neophorocera hamata*).

Neodiprion sp. near *scutellatis* Rohwer

Smith, 1962: 189, Idaho (as *Phorocera hamata* ex *N.* near "scutellaris").

Neodiprion sertifer (Geoffroy)

Lyons, 1964: 21 (as *D. hamatus*); Finlayson, T., 1960: 24, 42, figs. 21, 24, Ont. (as *D. hamata*); Finlayson and Finlayson, 1958: 223-224, Ont. (as *Neophorocera hamata*); Raizenne, 1957: 14, Ont. (as *Neophorocera hamata*).

Neodiprion swaini Middleton

Becker, Jr., and Benjamin, 1964: 356, Wis. (as *D. hamatus*); Finlayson, T., 1963: 484, 502 (as *D. hamatus*); Raizenne, 1957: 12, Que. (as *Neophorocera hamata*).

Neodiprion swaini Middleton probably

Martineau, 1959: 230, Que. (as *D. hamatus* ex *N. swaini*).

Neodiprion taedae linearis Ross

Warren, L. O., and Coyne, 1958: 20, Ark. (as *Phorocera* "namata").

Neodiprion virginiana Rohwer complex

Finlayson, T., 1963: 485, 502 (as *D. hamatus*).

Neodiprion spp.

Dahlsten, 1961: 193-194, Calif. (as *D. hamatus*); Patton, 1958: 80, Fla. (as *Phorocera hamata*); Anon., 1953c: 137, B.C. (as *Neophorocera hamata*); Greene, 1921: 43, Conn. (as *Exorista petiolata*).

Neodiprion sp. probably

Patton, 1958: 80, Fla. (as *Phorocera hamata*).

Diprionid spp. (including *Diprion* spp. and/or *Neodiprion* spp.)
Coquillett, 1897: 14, 22, Va. (as *Exorista petiolata* ex
Lophyrus).

Tenthredinidae

Pikonema spp.

Anon., 1953c: 137, B.C. (as *Neophorocera hamata*).

Pristiphora erichsonii (Hartig)

*Drooz, 1960: 34, Canada (as "*Neophorocera*" *hamata*,
Graham, 1955); Raizenne, 1957: 30, Ont. (as
Neophorocera hamata); Graham, 1955: 66 (as
Neophorocera hamata).

Pristiphora geniculata (Hartig)

Raizenne, 1957: 30, Ont. (as *Neophorocera hamata*).

HOST: LEPIDOPTERA

Lasiocampidae

Malacosoma americanum (Fabricius)

Raizenne, 1952: 101, Ont. (as *Neophorocera hamata*).

STAGE PARASITIZED: Larva (Hymenoptera), pupa
(Hymenoptera).

IMMATURES: Egg — Baldwin, W.F., and Coppel, 1949a: (descr.,
figs. 1-2).

First instar larva — Baldwin, W.F., and Coppel, 1949a: (descr.,
figs. 3-6).

Second instar larva — Baldwin, W.F., and Coppel, 1949a:
(descr., fig. 7).

Third instar larva — Finlayson, T., 1963: (descr.); Finlayson,
T., 1962: (descr.); Finlayson, T., 1960: (descr., fig. 21);
Finlayson, T., 1960a: (descr.); Baldwin, W.F., and Coppel,
1949a: (descr., figs. 8-10).

Pupa — Finlayson, T., 1960: (descr., fig. 24); Baldwin, W.F.,
and Coppel, 1949: (descr., fig. 1); Baldwin, W.F., and
Coppel, 1949a: (descr., figs. 11-14).

BIOLOGY: Bobb, 1963: (619-620); Dahlsten, 1961: (193-194);
Finlayson, T., 1960a: (923, 937); Daviault, 1953: (181-184).

Diplostichus sp. near *lophyri* (Townsend)

HOST: HYMENOPTERA

Diprionidae

Neodiprion abietis (Harris) complex

Struble, 1957a: 312, Calif. (as *Phorocera* sp., near *hamata*
Aldrich and Webber).

STAGE PARASITIZED: Larva.

Diplostichus sellersi Hall

HOST: HYMENOPTERA

Diprionidae

Neodiprion tsugae Middleton

Furniss and Dowden, 1941: 49-50, Oreg.; Hall, 1939: 243, Oreg.

Neodiprion spp.

Hall, 1939: 243, Calif.

STAGE PARASITIZED: Larva.

Dipterous larva [*Celatoria crawii*], see *Celatoria diabroticae*

Doryphorophaga Townsend (Goniinae: Blondeliini)

Doryphorophaga aberrans, see *Adoryphorophaga aberrans*

Doryphorophaga australis Reinhard

HOST: COLEOPTERA

Chrysomelidae

Leptinotarsa decemlineata (Say)

Kelleher, 1966: 1059, Man.; Reinhard, 1935b: 390, Tex.

Leptinotarsa defecta Stål

Reinhard, 1935a: 390, Tex.

Chrysomelid spp. (including *Lema trilineata* (Olivier) and/or *Leptinotarsa decemlineata* (Say))

Patton, 1958: 36, Fla. (ex potato beetle).

STAGE PARASITIZED: Larva.

Doryphorophaga dorsalis, see *Myiopharus dorsalis*

Doryphorophaga "dorypharae," see *Doryphorophaga doryphorae*

Doryphorophaga doryphorae (Riley)

HOST: COLEOPTERA

Chrysomelidae

Leptinotarsa decemlineata (Say)

Williamson, 1969: 16, Ont.; Williamson, 1968: 16, Ont.; Kelleher, 1966: 1059-1061, Man.; Anon., 1964a: 388, Wash.; Landis, 1963: 714, Wash. (as *D. "dorypharae"*); Kelleher, 1960: 122, Man.; MacNay, 1960: 221, Man. (ex Colorado potato beetle); MacNay, 1960a: 1083, Man.; Anon., 1958a: 134, Man.; Maybee, 1958: 312, Man. (ex *L. "decimlineata"*); Anon, 1957b: 259, Man.; Anon., 1957c: 273, Ont.; Anon., 1954d: 250, Man.;

Neilson, 1954: CC67, B.C. (as *Phorocera*); Anon., 1953b: 135, B.C. (ex Colorado potato beetle); Handford and Neilson, 1953: 6, B.C.; Anon., 1952c: 204, Man.; Bird, R. D., and Mitchener, 1952: 47, Man.; MacNay, 1952: 318, B.C. (ex Colorado potato beetle); Bird, R. D., and Mitchener, 1951: 50, Man.; Fattig, 1949: 15, Ga. (ex *L. "decimlineata"*); Bird, R. D., and Mitchener, 1947: 37, Man.; Hammond, 1947: 37, Ont.; Munro and Telford, 1943: 9, N. Dak.; Drake, 1942: 171, Iowa; Munro, 1939: 563, N. Dak.; Bird, R. D., and Mitchener, 1938: 49, Man. (as *D. "doryphorea"*); Bird, R. D., and Mitchener, 1937: 44, Man. (as *D. "doryphorea"*); Brunteau, 1937: 117-119, 128-134, fig. 6, N.Y. (ex *Doryphora*); Feytaud, 1937: 64-65; Strickland, 1937: 449-450, Alta.; Feytaud, 1936: 26-27, U.S. (ex *Doryphora*); Hammond, 1936: 38, Ont.; Landis, 1936: 284, Ohio; Allen, 1929: 687, Miss.; Lochhead, 1919: 309 (as *Lydella*); Smith, H. E., 1918: 123, U.S.; Caesar, 1916: 32-33, Ont. (as *Phorocera* ex Colorado potato beetle); Britton, 1915a: 186, 190, Conn. (ex Colorado potato beetle); Sanderson, 1913: 294, Mo. (as *Lydella*); Townsend, 1912b: 164, N. and E. U.S. to Mo.; Conradi, 1907: 33-34, Tex. (as *Lydella* ex *L. 10-lineata*); Coquillett, 1897: 19, 23, Md. (as *Phorocera* ex *Doryphora 10-lineata*); Brauer and Bergenstamm, 1895: 558, 591 (as *Lydella* ex *Doryphora*); Fuller, 1880: 190 (as *Lydella* ex Colorado potato beetle); Brodie, 1878: 9-10, Ont. (as *Lydella "doryphora"* ex *Doryphora 10-lineata*); Riley, 1877: 40 (as *Lydella* ex *Doryphora 10-lineata*); Riley, 1876a: 52-53, N. Cent. U.S., N.E. U.S. (as *Lydella* ex *Doryphora 10-lineata*); Riley, 1875a: 769 (as *Lydella "doryphora"* ex *Doryphora 10-lineata*); Riley, 1875b: 706, U.S. (as *Lydella* ex "*doryphora*"); Reed, 1872: 71, 74 (as *Lydella* ex *Doryphora 10-lineata*); LeBaron, 1871a: 64-65, Ill. (as *Lydella* ex *Doryphora 10-lineata*); Packard, 1869: 408 (as *Tachina* ex Colorado potato beetle); Riley, 1869: 111-112, Mo. (as *Lydella* ex "*Doryphorae*" *10-lineata*); Walsh and Riley, 1868: 46, Ill. (as *Lydella* [sp.] ex *Doryphora 10-lineata*).

Chrysomelid spp. (including *Lema trilineata* (Olivier) and/or *Leptinotarsa decemlineata* (Say))

All records ex potato beetle.

*Procter, 1946: 431, Maine (Procter, 1938); Procter, 1938: 373, [Maine]; Strickland, 1938: 217, Alta.; Briand,

1936: 32, 34, Ont.; Johnson, 1927: 201, Maine; Johnson, 1925: 199, Conn., Maine, Mass., N.H.; Johnson, 1910: 780, N.J. (as *Phorocera*); Reed, 1883: 52 (as *Tachina*).

Coccinellidae

Epilachna varivestis Mulsant

Howard, N.F., and Landis, 1936: 2, 6, Ohio.

HOST: LEPIDOPTERA

Nymphalidae

Nymphalis antiopa (Linnaeus)

Coquillett, 1897: 19, 27, Mo. (as *Phorocera ex Vanessa*).

STAGE PARASITIZED: Larva (Coleoptera), adult (Coleoptera).

IMMATURES: Egg — Riley, 1875a: (descr.).

First instar larva — Bruneteau, 1937: (descr., fig. 6).

Pupa — Bruneteau, 1937: (descr.).

BIOLOGY: Kelleher, 1966: (1059-1061); Munro and Telford, 1943: (9); Feytaud, 1937: (64-65); Caesar, 1916: (32-33); Conradi, 1907: (33-34); Riley, 1876a: (52-53); Riley, 1875a: (769); Riley, 1875b: (706); Reed, 1872: (71, 74); Riley, 1869: (111-112); Walsh and Riley, 1868: (46).

Doryphorophaga "doryphorea," see *Doryphorophaga doryphorae*

Doryphorophaga macella Reinhard

HOST: COLEOPTERA

Chrysomelidae

Leptinotarsa decemlineata (Say)

Keller, 1966: 1059, Man.

STAGE PARASITIZED: Larva.

Doryphorophaga spp.

HOST: COLEOPTERA

Chrysomelidae

Leptinotarsa decemlineata (Say)

Williamson, 1963: 150, Ont.; Williamson, 1962: 158, Ont.; Williamson, 1961: 268, Ont.; Kelleher, 1960a: 280, Canada; Maybee, 1959: 263, Man.; Drake, 1941: 151-152, Iowa.

Doryphorophaga spp. and/or *Adorophorophaga* spp., see *Adorophorophaga* spp. and/or *Doryphorophaga* spp.

Drepanoglossa Townsend (Goniinae: Siphonini)

Drepanoglossa amydriae Townsend

HOST: LEPIDOPTERA

Tineidae

Amydria sp.

Townsend, 1908: 76, Mexico (Mor.).

STAGE PARASITIZED: Pupa.

Drepanoglossa floridensis, see *Clausicella floridensis***Drino Robineau-Desvoidy** (Goniinae: Sturmiini)**Drino austrina** (Coquillett)

HOST: LEPIDOPTERA

Pyralidae

Unidentified Pyralidae

Johnson, 1908: 75, Bahamas (as *Sturmia*).**Drino** sp. near *bakeri* (Coquillett)

HOST: LEPIDOPTERA

Sphingidae

Amphion nesus (Cramer)Schaffner and Griswold, 1934: 25, 118, Mass. (as *Sturmia*).

BIOLOGY: Schaffner and Griswold, 1934: (25, 118).

Drino balloui (Curran)All records as *Sturmia balloui*.

HOST: LEPIDOPTERA

Sphingidae

Sphinx moth(-s)

Arnaud, 1963: 128, Costa Rica (ex "sphinx" moth).

Unidentified Sphingidae

Curran, 1935: 22, Costa Rica.

STAGE PARASITIZED: Pupa.

Drino bohémica Mesnil

HOST: HYMENOPTERA

Diprionidae

Diprion hercyniae (Hartig)

Warren, 1967: 11, Nfld.; Monteith, 1964: 1480-1481 (lab.);

Reeks, 1964: 31, Ont.; Rose, 1964: 79, Ont. (ex European spruce sawfly); Monteith, 1963: 163-166;

Dowden, 1962: 16, 56, N.B.; Balch, 1960: 303, Canada; Finlayson, T., 1960a: 923, 937-938, fig. 13, Que.; Monteith, 1960: 641, 643-644, 647-651 (lab.); Bird and Elgee, 1957: 375, 377, N.B.; Raizenne, 1957: 24, Ont.; Monteith, 1955: 510; Anon., 1954c: 148, Que.; Boyes and Wilkes, 1953: 126, 145 (lab.); Reeks, 1953: 406, 409, 414, 417-420, 423-427, N.B.; Anon., 1952b: 176, 182, 186-187 (lab.); Anon., 1952f: 256, Que.; Anon., 1950d: 234, 242, 251 (ex *Gilpinia*, lab.); Anon., 1949: 274-275, 280, 286, 288 (as *Sturmia* sp. ex *Gilpinia*, lab.); Briand, 1949: 56-57, 59, Que. (as *Sturmia* sp. ex [European] spruce sawfly); Briand, 1949a: 125-127, Que. (as *Sturmia* sp. ex European spruce sawfly); Briand, 1949b: 180-181, Que. (as *Sturmia* sp. ex *Gilpinia*); Hawboldt, 1949: 37, Ont. (as *Sturmia* sp. ex *Gilpinia*); House, 1949: 60-62 (as *Sturmia* sp. ex *Gilpinia*, lab.); Reeks, Forbes and Cuming, 1949: 15, N.B. (as *Zygobothria gilva* (Hartig) ex *Gilpinia*); Anon., 1948: 287-288, 291, 295-297, Ont. (as *Sturmia* sp. ex *Gilpinia*); Anon., 1948a: 357-358, 362, 369-371 (as *Sturmia* sp. ex *Gilpinia*, lab.); Reeks, Forbes and Cuming, 1948: 15, N.B. (as *Zygobothria gilva* ex *Gilpinia*); Anon., 1947: 187-188, 192, 197-198, 200-201 (as *Sturmia* sp. ex *Gilpinia*, lab.); Coppel and House, 1947: 74 (as *Sturmia* sp. ex *Gilpinia*, lab.); Reeks, Forbes and Cuming, 1947: 11, [N.B., N.S.] (as *Sturmia* sp. ex *Gilpinia*); Baird, 1946: 141-142, 145, 149-150, 152 (as *Sturmia* sp. ex *Gilpinia*, lab.); Reeks, 1946: 25, N.B. (as *Sturmia* sp. ex *Gilpinia*); Anon., 1945: 224 (as *Sturmia* sp. ex *Gilpinia*); Anon., 1945a: 224, Que. (as *Sturmia* sp. ex [European] spruce sawfly); Baird, 1945: 127, 131, 137-139, Ont. (as *Sturmia* sp. ex *Gilpinia*); Paquet, 1945: 94 (as *Sturmia* sp. ex *Gilpinia*); Baird, 1944: 126-127, 131, 135-137, Ont. (as *Sturmia* sp. ex *Gilpinia*); Gardiner, 1942: 57, maritime prov., Ont., Que. (as *Sturmia* sp. ex *Gilpinia*).

Diprion polytomum (Hartig)

Gardiner, 1943: 55, N.B., Nfld., N.S., Ont., P.E.I., Que. (as *Sturmia* sp. ex spruce sawfly).

Neodiprion abietis (Harris)

Finlayson, T., 1963: 477, 503; Anon., 1948b: 356, 360, 369 (as *Sturmia* sp., lab.).

Neodiprion lecontei (Fitch)

Monteith, 1966: 1205-1207 (lab.); Monteith, 1964:

1479-1481 (lab.); Monteith, 1963: 162-166; Monteith, 1963a: 418-425 (lab.); Monteith, 1960: 641, 643-644, 647-651 (lab.); Benjamin, 1955: 26 (as *Sturmia* sp., lab.); Monteith, 1955: 510; Anon., 1952b: 176, 182, 187 (lab.); Anon., 1948: 288, 292, 296 (as *Sturmia* sp., lab.); Anon., 1948a: 359, 364, 369 (as *Sturmia* sp., lab.); Anon., 1947: 189, 194, 199 (as *Sturmia* sp., lab.); Baird, 1946: 142, 146, 151 (as *Sturmia* sp., lab.).

Neodiprion nanulus nanulus Schedl

Baird, 1946: 142, 147, 151 (as *Sturmia* sp. ex *N. nanulus*, lab.).

Neodiprion pratti banksianae Rohwer

Monteith, 1964: 1479-1481 (lab.); Finlayson, T., 1963: 482, 503; Raizenne, 1957: 15, Ont.; Anon., 1952b: 176, 182, 187 (ex *N. americanus banksianae*, lab.); Anon., 1948a: 358, 363, 369 (as *Sturmia* sp. ex *N. banksianae*, lab.); Baird, 1946: 142, 146, 151 (as *Sturmia* sp. ex *N. banksianae*, lab.); Baird, 1944: 127, 132, 137 (as *Sturmia* sp. ex *N. banksianae*, lab.).

Neodiprion sertifer (Geoffroy)

Lyons, 1964: 21, 31, Ont.; Monteith, 1963: 163-164, 166; Monteith, 1955: 510; Anon., 1948: 287, 290, 296 (as *Sturmia* sp., lab.); Anon., 1948a: 357, 362, 369 (as *Sturmia* sp., lab.); Anon., 1947: 187, 192, 199 (as *Sturmia* sp., lab.); Baird, 1946: 140, 145, 150 (as *Sturmia* sp., lab.); Baird, 1945: 127, 131, 138 (as *Sturmia* sp. ex *Diprion*, lab.).

Neodiprion swaini Middleton

Monteith, 1963: 163-164, 166; Monteith, 1955: 510; Anon., 1952b: 176, 183, 187 (lab.); Anon., 1948a: 358, 363, 369 (as *Sturmia* sp., lab.).

Neodiprion virginiana Rohwer complex

Finlayson, T., 1963: 485, 503.

Tenthredinidae

Hemichroa crocea (Fourcroy)

Anon., 1948c: 356, 360, 369 (as *Sturmia* sp., lab.).

Pikonema alaskensis (Rohwer)

Monteith, 1964: 1479-1481 (lab.); Dowden, 1962: 56, N.B.; Finlayson, T., 1960a: 923, 938, fig. 13; Raizenne, 1957: 32, Ont.; Reeks, 1953: 426; Anon., 1952b: 176, 184, 187 (lab.); Briand, 1949: 58 (as *Sturmia* [sp.]); Briand, 1949a: 125, Que. (as *Sturmia* sp.); Anon., 1948: 289, 293, 296 (as *Sturmia* sp., lab.); Anon., 1948a: 359, 366,

371 (as *Sturmia* sp., lab.); Anon., 1947: 189, 196-197 (as *Sturmia* sp., lab.).

Pikonema dimmockii (Cresson)

Dowden, 1962: 56, N.B.; Finlayson, T., 1960a: 923, 938, fig. 13; Reeks, 1953: 426.

Pristiphora erichsonii (Hartig)

Anon., 1948: 288, 292, 296 (as *Sturmia* sp., lab.).

Pristiphora geniculata (Hartig)

Anon., 1948: 288, 292, 296 (as *Sturmia* sp., lab.).

Unidentified Hymenoptera

Sawfly (-flies)

Anon., 1948: 288, 292, 296, Ont. (as *Sturmia* sp.); Anon., 1947: 189, 195, 199-200 (as *Sturmia* sp., lab.).

STAGE PARASITIZED: Larva, pupa.

IMMATURES: Second instar larva — Finlayson, T., 1960a: (descr., fig. 13).

BIOLOGY: Monteith, 1966: (1205-1207); Monteith, 1964: (1479-1481); Monteith, 1963: (162-166); Monteith, 1963a: (418-425); Dowden, 1962: (16, 56); Finlayson, T., 1960a: (923, 937-938); Monteith, 1960: (641, 643-644, 647-651); Reeks, 1953: (406, 409, 414, 417-420, 423-427); Briand, 1949a: (125-127); House, 1949: (60-62).

Drino cubaecola (Jaennicke)

HOST: LEPIDOPTERA

Pyralidae

Hedylepta indicata (Fabricius)

Plank, 1945: 26-27, P.R. (as *Sturmia*).

Drino fraudulenta (Wulp)

HOST: LEPIDOPTERA

Hesperiidae

Erynnis icelus (Scudder and Burgess)

Schaffner and Griswold, 1934: 15, 118 (as *Sturmia*).

Drino inca (Townsend)

HOST: LEPIDOPTERA

Sphingidae

Agrius cingulatus (Fabricius)

All records ex *Herse cingulata*.

Patton, 1958: 86, Fla. (as *Zygoturmia*); Fattig, 1949: 38, Ga. (as *Zenillia*); Wylie, 1946: 90, Fla. (as *Sturmia*);
Brimley, 1938: 364, N.C. (as *Sturmia*); Aldrich, 1927b: 19, La., Tex. (as *Sturmia*).

Agrius cingulatus (Fabricius) probably

Aldrich, 1927b: 19, Tex. (as *Sturmia* ex *Herse*).

Unidentified Sphingidae

Tucker, R. W. E., 1952: 351, Barbados (as *Zygoturmia* ex larvae feeding on sweetpotato).

STAGE PARASITIZED: Larva.

BIOLOGY: Patton, 1958: (86).

Drino incompta (Wulp)

Most records as *Sturmia inquinata* (Wulp).

HOST: LEPIDOPTERA

Citheroniidae

Eacles imperialis (Drury)

Johnson, 1910: 781, N.J.

Saturniidae

Hemileuca maia (Drury)

Essig, 1926: 583, [W. U.S.]; Packard, 1914: 269; Coquillett, 1897: 20, 25.

Sphingidae

Agrius cingulatus (Fabricius)

Brimley, 1922: 25, N.C. (ex *Phlegethontius*); Coquillett, 1897: 20, 26, Fla. (ex *Protoparce*).

Ceratomia amyntor (Hübner)

Schaffner and Griswold, 1934: 17, 118, N.H. (as *S. incompta*); Coquillett, 1897: 20, 24.

Ceratomia catalpae (Boisduval)

Fattig, 1949: 32, Ga. (as *S. incompta*).

Ceratomia undulosa (Walker)

Coquillett, 1897: 20, 24, Mo.

Ceratomia undulosa (Walker) probably

Brimley, 1922: 25, N.C.

Ceratonia spp.

Brimley, 1938: 364, N.C.

Hyles lineata (Fabricius)

Most records ex *Deilephila lineata*.

Essig, 1926: 583, [W. U.S.] (ex *Celerio*); Johnson, 1910: 781, N.J.; Coquillett, 1897: 20, 25, La.

Manduca quinquemaculata (Haworth)

Most records ex *Protoparce quinquemaculata*.

Gilmore, 1938: 710-711, Ky. and/or Tenn. (as *S. incompta*); Essig, 1926: 583, [W. U.S.]; Lovett, 1915a: 171, fig. 3 on pl. 7, Oreg. (ex *Phlegethontius*); Johnson, 1910: 781, N.J. (ex *P. celeus* Hübner); Coquillett, 1897: 20, 26, Mo. (ex *P. celeus*).

Manduca sexta (Linnaeus)

Most records ex *Protoparce sexta*.

*Madden and Chamberlin, 1945: 48 (as *S. incompta*, Gilmore, 1938); Gilmore, 1938: 710-711, Ky. and/or Tenn. (as *S. incompta*); Essig, 1926: 583, [W. U.S.]; Johnson, 1910: 781, N.J. (ex *P. carolina* (Linnaeus)); Coquillett, 1897: 20, 26, Colo., Ill. (ex *P. carolina*).

Manduca spp.

Garman, 1897: 29 (ex *Phlegethontius*).

Pholus achemon (Drury)

Coquillett, 1897: 20, 26, Mo. (ex *Philampelus*).

Pholus vitis (Linnaeus)

All records ex *Pholus fasciatus* Sulzer.

Patton, 1958: 83, Fla. (as *S. incompta*); Patton, 1957: 69 (as *S. incompta*).

Sphinx chersis (Hübner)

Aldrich, 1932: 1-2, Mass., N.H. (as *S. incompta*); Essig, 1926: 583, [W. U.S.]; Tothill, 1913: 74, Ont.; Eliot and Soule, 1902: 58.

Unidentified Sphingidae

*Procter, 1946: 434, Maine (as *S. incompta*, Procter, 1938); Procter, 1938: 375, Maine (as *S. incompta*).

STAGE PARASITIZED: Larva, pupa.

IMMATURES: Pupa — Greene, 1922: (descr., fig. 4 on pl. 1); Lovett, 1915a: (fig. 3 on pl. 7).

BIOLOGY: Patton, 1958: (83); Patton, 1957: (69); Schaffner and Griswold, 1934: (17, 118); Lovett, 1915a: (171).

Drino inconspicua (Meigen)

Most records as *Sturmia inconspicua*.

HOST: HYMENOPTERA

Diprionidae

Diprion frutetorum (Fabricius)

Baird, 1943: 119, 122, 129 (ex *Gilpinia*, lab.).

Diprion hercyniae (Hartig)

Baird, 1943: 118-119, 123, 127-130 (ex *Gilpinia*, lab); Baird,
1941: 160-161, Que. (ex [European] spruce sawfly).

Diprion polytomum (Hartig)

Finlayson and Reeks, 1936: 164, Que. (lab.).

Diprion similis (Hartig)

Webber, 1932: 193, 202, 204, 206-207.

Neodiprion lecontei (Fitch)

Webber, 1932: 204, 207, Mass.

Neodiprion pratti pratti (Dyar)

Webber, 1932: 194-204, 207, figs. 3-6 (ex *N. dyari* Rohwer).

Neodiprion sertifer (Geoffroy)

Baird, 1943: 118, 122, 129 (lab.).

HOST: LEPIDOPTERA

Geometridae

Hydria undulata (Linnaeus)

Webber, 1932: 207.

Lasiocampidae

Malacosoma americanum (Fabricius)

Webber, 1932: 204, 207.

Malacosoma disstria Hübner

Webber, 1932: 204, 207.

Lymantriidae

Hemerocampa leucostigma (J. E. Smith)

Webber, 1932: 207.

Porthetria dispar (Linnaeus)

Clausen, 1956: 63; Webber, 1932: 193-208, figs. 3-6, Mass.;
Howard and Fiske, 1911: 308 (as *Zygobothria gilva*
Hartig).

Noctuidae

Alypia octomaculata (Fabricius)

Webber, 1932: 207.

STAGE PARASITIZED: Larva (Hymenoptera, Lepidoptera).

IMMATURES: Egg — Webber, 1932: (descr.).

First instar larva — Webber, 1932: (descr., fig. 3).

Second instar larva — Webber, 1932: (descr., fig. 4).

Third instar larva — Webber, 1932: (descr., fig. 5).

Pupa — Webber, 1932: (descr., fig. 6).

BIOLOGY: Webber, 1932: (193-208).

Drino inconspicua (Meigen) probably

HOST: HYMENOPTERA

Diprionidae

Diprion hercyniae (Hartig)

Reeks, Smith and Forbes, 1946: 11, N.B. (as *Sturmia* ex *Gilpinia*).

STAGE PARASITIZED: Larva.

Drino munda, see *Palexorista laxa*

Drino pilatei (Coquillett)

IMMATURES: Pupa — Greene, 1922: (as *Sturmia*; descr., fig. 33 on pl. 7).

Drino rhoeo (Walker)

Most records as *Sturmia distincta* Wiedemann.

HOST: LEPIDOPTERA

Psychidae

Thyridopteryx ephemeraeformis (Haworth)

Webster and Mally, 1897: 42, Ohio.

Sphingidae

Agrius cingulatus (Fabricius)

Most records ex *Protoparce cingulata*.

Tucker, R. W. E., 1952: 351, Barbados (ex *Herse*); Catoni, 1922: 25-26, P.R.; Bourne, 1921: 13, Barbados.

Agrius convolvuli (Linnaeus)

Reinhard, 1922: 73, Tex. (ex *Herse*).

Manduca quinquemaculata (Haworth)

Fattig, 1949: 32 (as *S. protoparcis* (Townsend) ex *Protoparce*); Brimley, 1938: 364, N.C. (ex *Phlegethontius*); Brimley, 1922: 25, N.C. (ex *Phlegethontius*); Coquillett, 1897: 20, 26, Ill. (ex *Protoparce celeus* Hübner).

Manduca sexta (Linnaeus)

Most records ex *Protoparce sexta*.

Comstock and Vazquez Garcia, 1964: 221, Mexico (as *Argyrophylax protoparcis* ex *Phlegethontius*); Comstock and Vazquez Garcia, 1964: 221, Mexico (ex *Phlegethontius*); Lawson, 1959: 741, 744-745, N.C. (as *Zygosturmia protoparcis*); Patton, 1958: 86, Fla. (as *Zygosturmia protoparcis*); Fattig, 1949: 32, Ga. (as *S. protoparcis*); Chamberlin, F.S., and Madden, 1942: 18, Fla. (as *S. protoparcis*); Brimley, 1942: 27, N.C. (as *S. protoparcis*); Chamberlin, F.S., and Madden, 1942: 18, Fla., Ga. (as *S. protoparcis*); Brimley, 1938: 364, N.C. (ex *Phlegethontius*); Gilmore, 1938: 710-711, Ky. and/or Tenn. (as *S. protoparcis*); Madden and Chamberlin, 1938: 705, Fla. (as *S. protoparcis*); Allen, 1929: 688, Miss.; Russo, 1927a: 186-187, Fla., Jamaica, Lesser Antilles; Brimley, 1922: 25, N.C. (ex *Phlegethontius*).

Manduca sexta jamaicensis (Butler)

Most records ex *Protoparce sexta jamaicensis*.

Curran, 1926: 113, Jamaica (as *Atacta geminata* Curran); Gowdey, 1926: 82, Jamaica (as *Atacta geminata*); Gowdey, 1926: 84, Jamaica (as *Zenillia* ex *P. sexta* "jamaicensis"); Gowdey, 1923: 2, Jamaica; Gowdey, 1923a: 59, Jamaica; Johnson, 1919: 437, Jamaica (ex *P. jamaicensis*); Coquillett, 1897: 20, 26, Jamaica (ex *P. jamaicensis*); *Brauer and Bergenstamm, 1895: 561, 604 (as *Masicera protoparcis* ex *P. jamaicensis*, Townsend, 1892d); Cockerell, 1892: 43, Jamaica (as *Masicera protoparcis* ex *P. jamaicensis*); Cockerell, 1892a: 56, Jamaica (as *Masicera protoparcis* ex *P. jamaicensis*); Townsend, 1892d: 70, Jamaica (as *Masicera protoparcis* ex *P. jamaicensis*).

Pholus achemon (Drury)

Essig, 1926: 583, Calif.; Nougaret, 1919: 579, Calif.

Sphinx moth(-s)

Aldrich, 1927b: 19, U.S.

Unidentified Sphingidae

Coquillett, 1897: 20, 27, Tex.

STAGE PARASITIZED: Larva, pupa.

IMMATURES: Egg — Madden and Chamberlin, 1945: (descr.).

Pupa — Greene, 1922: (descr., fig. 2 on pl. 1).

BIOLOGY: Madden and Chamberlin, 1945: (46-48); Allen, 1929: (688); Nougaret, 1919: (579).

Drino rhoeo (Walker) probably
 Most records as *Sturmia trifida* Wulp.
 HOST: LEPIDOPTERA

Sphingidae

Manduca quinquemaculata (Haworth)

Garman, 1897: 29 (ex *Phlegethontius celeus* (Hübner)).

Manduca sexta (Linnaeus)

Kuitert and Tissot, 1957: 94, Fla. (as *S. protoparcis* (Townsend) ex [tobacco] hornworms); Girault, 1908a: 235-236, Tex. (as *Argyrophylax protoparcis* (or a n. sp.?) ex *Phlegethontius*); Garman, 1897: 29 (ex *Phlegethontius carolina* (Linnaeus)).

STAGE PARASITIZED: Larva.

IMMATURES: Pupa — Girault, 1908a: (descr.).

Drino sociabilis (Greene)

IMMATURES: Pupa — Greene, 1922: (as *Sturmia*; descr., fig. 1 on pl. 1).

Drino sp.

All records as *Zygosturmia* sp.

HOST: LEPIDOPTERA

Sphingidae

Sphinx moth(-s)

Wolcott, 1951a: 480, P.R. (ex caterpillar on *Cordia*);

Wolcott, 1924: 222, P.R. (ex larva on *Cordia*).

STAGE PARASITIZED: Larva.

Echinomyia algens, see *Nowickia algens*

Echinomyia thomsoni, see *Peleteria thomsoni*

Eipogona americana, see *Buquetia obscura*

Eipogona obscura, see *Buquetia obscura*

Eipogona setifacies, see *Buquetia musca*

Elachipalpus sp., see *Copecrypta* sp.

Eleodiphaga Walton (Goniinae: Frontinini)

Eleodiphaga caffreyi Walton

HOST: COLEOPTERA

Tenebrionidae

Eleodes extricata (Say)

Essig, 1926: 579, Ariz., N. Mex.; Walton, 1918: 24, N. Mex.

Eleodes obsoleta (Say)

Essig, 1926: 579, Ariz., N. Mex.; Walton, 1918: 24, Ariz.

STAGE PARASITIZED: Adult.

Eleodiphaga pollinosa Walton

HOST: COLEOPTERA

Tenebrionidae

Eleodes hispilabris (Say)

Essig, 1926: 579, N. Mex.; Walton, 1918: 24, N. Mex.

STAGE PARASITIZED: Adult

"*Elephantocera greeni*," see *Elephantocera greeni*

Elephantocera Townsend (Goniinae: Blondeliini)

Elephantocera greeni Townsend

All records as *Elephantocera* "greeni."

HOST: LEPIDOPTERA

Olethreutidae

Grapholitha molesta (Busck)

All records ex *Laspeyresia molesta*.

Steenburgh and Boyce, 1938: 73, Ont. (as "*Elephantocera greeni*"); Steenburgh, 1935: 71, Ont.; Steenburgh, 1932: 67-68, Ont.

STAGE PARASITIZED: Larva.

Elephantocera "greeni," see *Elephantocera greeni*

Elfia Robineau-Desvoidy (Goniinae: Siphonini)

Elfia aenea (Coquillett)

HOST: LEPIDOPTERA

Tortricidae

Phalonia oenotherana Riley

Dickerson and Weiss, 1920: 44, N.J. (as "*Tachniphyto*").

STAGE PARASITIZED: Larva.

Elfia sp. near *aenea* (Coquillett)

HOST: LEPIDOPTERA

Pyralidae

Pyrausta futilalis (Lederer)

Schaffner, 1959: 11, 53, N.J. (as *Phyllomya*).

BIOLOGY: Schaffner, 1959: (11, 53).

Elfia amplicornis (James)

HOST: LEPIDOPTERA

Glyphipterygidae

Lotisma trigonana Walsingham

James, M. T., 1955: 84, Wash. (as *Plectops*).

Elfia curriei (Townsend)

HOST: LEPIDOPTERA

Pyralidae

Dioryctria reniculella (Grote)

Daviault, 1946: 117, Que. (as *Plectops* "currei").

Elfia melissopodis (Coquillett)

All records as *Plectops melissopodis*.

HOST: COLEOPTERA

Curculionidae

Curculio sp.

Brooks, F. E., and Cotton, 1929: 14, Mass. (ex *Curculio* larvae in acorns of *Quercus velutina* Lamarck, questionable parasite).

HOST: LEPIDOPTERA

Olethreutidae

Melissopus latiferreanus (Walsingham)

Coquillett, 1897: 19, 26, D.C., Mo.

STAGE PARASITIZED: Larva (Lepidoptera).

Elfia palpigera (Coquillett)

HOST: LEPIDOPTERA

Pyralidae

Acrobasis comtoniella Hulst

Schaffner, 1959: 15, 51, N.H. (as *Lispidea*).

BIOLOGY: Schaffner, 1959: (15, 51).

Elfia sp. near *palpigera* (Coquillett)

HOST: LEPIDOPTERA

Tortricidae

Argyrotaenia citrana (Fernald)

Basinger, 1935: 233, Calif. (as "Lispidia").

STAGE PARASITIZED: Larva.

Elfia setigera (Thomson)All records as *Goliathocera setigera*.

HOST: LEPIDOPTERA

Pyralidae

Dioryctria abietella (Denis and Schiffermüller)

Keen, 1958: 70.

Cone moth(-s)

Keen, 1958: 70, Oreg.

STAGE PARASITIZED: Pupa.

Elfia tarsalis (Coquillett)All records as *Clausicella tarsalis*.

HOST: LEPIDOPTERA

Pyralidae

Pyralis farinalis Linnaeus probably

Chittenden, 1897: 42-43, Md. (probable parasite).

IMMATURES: Pupa — Greene, 1922: (descr., fig. 31 on pl. 7).

Elfia sp. near *usitata* (Coquillett)

HOST: LEPIDOPTERA

Oecophoridae

Psilocorsis reflexella ClemensSchaffner, 1959: 40, 53, N.H. (as *Plectops*).

BIOLOGY: Schaffner, 1959: (40, 53).

Elfia sp. (10), see *Elfia* spp.**Elfia** spp.

HOST: LEPIDOPTERA

Gelechiidae

Coleotechnites canusella (Freeman)McLeod, 1969: 168, Que. (as "*Elphia*" ex *Eucordylea*).*Coleotechnites thujella* (Kearfott)Procter, 1946: 436, Maine (as *Lispidea* ex *Recurvaria*).

Oecophoridae

Psilocorsis faginella (Chambers)Embree, 1958: 170, N.S. (as *Lophosiocera*).*Psilocorsis quercicella* ClemensCochaux, 1969: 69, 79-81, Que. (as *Elfia* sp. (10)).

Psilocorsis reflexella Clemens

Cochaux, 1969: 69, 79-81, Que. (as *Elfia* sp. (10)).

Olethreutidae

Epinotia aceriella (Clemens)

Raizenne, 1952: 198, Que. (as *Plectops*).

Melissopus latiferreanus (Walsingham)

Dohanian, 1942: 837, 841, Oreg. (as *Plectops*, probable rearing).

Pterophoridae

Platyptilia carduidactyla (Riley)

Lange, 1950: 592, Calif. (as *Plectops*); Lange, 1950: 592, Calif. (as *Lispidea* spp. (two spp.)); Lange, 1941: 24, 67, Calif. (as *Plectops*); Lange, 1941: 24, 67, Calif. (as *Lispidea* sp. [1]); Lange, 1941: 24, 67, Calif. (as *Lispidea* sp. [2]).

STAGE PARASITIZED: Larva.

BIOLOGY: McLeod, 1969: (168).

Elodia Robineau-Desvoidy (Goniinae: Erythrocerini)

Elodia flavipalpis Aldrich

HOST: LEPIDOPTERA

Olethreutidae

Grapholitha molesta (Busck)

Allen, Holloway and Haeussler, 1940: 4, 33, 40-43, 49 (lab.).

STAGE PARASITIZED: Larva.

Elodia subfasciata Aldrich

HOST: LEPIDOPTERA

Olethreutidae

Grapholitha molesta (Busck)

Allen, Holloway and Haeussler, 1940: 4, 33, 40-43, 49 (lab.).

STAGE PARASITIZED: Larva.

Elodia tragica (Meigen)

HOST: LEPIDOPTERA

Olethreutidae

Grapholitha molesta (Busck)

Allen, Holloway and Haeussler, 1940: 3, 40-41, 60 (as *Arrhinomyia*, lab.).

STAGE PARASITIZED: Larva.

"Elphia" sp., see *Elfia* spp.

Emphanopteryx aurifacies, see *Cryptomeigenia aurifacies*

Emphanopteryx prisca, see *Cryptomeigenia prisca*

"Encelatoria" rubentis, see *Eucelatoria rubentis*

Ennyomma clistoides, see *Myiophasia clistoides*

Ennyomma globosa, see *Myiophasia globosa*

Ennyommopsis nigrifrons, see *Myiophasia nigrifrons*

Eohyria pellucida, see *Pelatachina pellucida*

Epalpus Rondani (Tachininae: Juriniini)

Epalpus signifer (Walker)

HOST: LEPIDOPTERA

Noctuidae

Lithophane spp. (including *L. disposita* Morrison, *L. innominata* (Smith) and/or *L. petulca* Grote)

Schaffner and Griswold, 1934: 48, 125, N.J. (ex *Graptolitha* spp. (including *G. disposita*, *G. innominata* and/or *G. petulca*)).

BIOLOGY: Schaffner and Griswold, 1934: (48, 125).

Epigrimyia floridensis, see *Clausicella floridensis*

Epigrimyia geniculata, see *Coronimyia geniculata*

"*Epigrymia*" floridensis, see *Clausicella floridensis*

Epimasicera caesar, see *Aplomya caesar*

Epiplagiops Blanchard (Goniinae: Exoristini)

Epiplagiops littoralis Blanchard

HOST: COLEOPTERA

Curculionidae

Listroderes costirostris obliquus (Klug)

Compere, 1946: 671-672 (ex *L. obliquus*, lab.).

STAGE PARASITIZED: Larva.

Eribella Mesnil (Goniinae: Blondeliini)

Eribella exilis (Coquillett)

All records as *Masicera exilis*.

HOST: COLEOPTERA

Chrysomelidae

Chelymorpha cassidea (Fabricius)

Chittenden, 1924: 49, D.C., Va.

Plagiometriona clavata (Fabricius)

Brimley, 1922: 24, N.C. (ex *Coptocycla*).

STAGE PARASITIZED: Larva.

Eribella sp. near *exilis* (Coquillett)

Most records as *Masicera* sp. near *exilis*.

HOST: HYMENOPTERA

Diprionidae

Neodiprion lecontei (Fitch)

Greene, 1921: 43, Va.

HOST: LEPIDOPTERA

Olethreutidae

Barbara colfaxiana (Kearfott)

Hedlin, 1960: 832, B.C. (as "*Pseudoeribia*" sp. near *tohilli* (Curran)).

IMMATURES: Pupa — Greene, 1922: (descr., fig. 58 on pl. 13).

Eribella spp.

HOST: LEPIDOPTERA

Pterophoridae

Unidentified Pterophoridae

Sabrosky and Arnaud, 1965: 1042.

STAGE PARASITIZED: Larva.

Ernestia ampelus, see *Mericia ampelus*

Ernestia sp. (1) (? *arcuata*), see *Mericia arcuata* probably

Ernestia sp. (10) (? *bicarina*), see *Mericia bicarina* probably

Ernestia johnsoni, see *Mericia johnsoni*

Ernestia radicum, see *Mericia* spp.

Ernestia spp. (2), (3), (4), (5), (6), (7), (8), (9) or (10), see *Mericia* spp.

Ernestia sp., see *Mericia* spp.

Ervia Robineau-Desvoidy (Dexiinae: Iceliini)

Ervia triquetra (Olivier)

HOST: COLEOPTERA

Chrysomelidae

Monocesta coryli (Say)

Fattig, 1949: 15, Ga. (ex "*Monocesta*").

STAGE PARASITIZED: Larva.

Erycia arator, see *Pilatea arator*

Erycia deckeri, see *Lydella deckeri*

Erycia leechi, see *Sitophaga leechi*

Erycia myoidaea, see *Lydella radialis*

Erycia rutila, see *Ceromasia aurifrons*

Erynnia Robineau-Desvoidy (Goniinae: Erythrocerini)

Erynnia nitida, see *Erynniopsis rondanii*

Erynnia tortricis (Coquillett)

Most records as *Anachaetopsis tortricis*.

HOST: LEPIDOPTERA

Coleophoridae

Coleophora fuscadinella Zeller

Reeks, 1951: 213, maritime prov. (as *Tortriciophaga* ex *C. salmani* Heinrich).

Coleophora malivorella Riley

Beacher, 1947: 532, 535-536, 539, Pa., W. Va.; Steiner and Worthley, 1941: 12, Pa.; Gould and Geissler, 1940: 815, W. Va.

Gelechiidae

Anarsia lineatella Zeller

Allen, 1962: 11, 114; Hardy, 1957: 195, probably Calif.; Basinger, 1935a: 246, Calif.

Pectinophora gossypiella (Saunders)

Allen, 1962: 114.

Noctuidae

Arzama obliqua (Walker)

Brehme, 1915: 473, N.J. (as *Hypostena* ex *Bellura*).

Olethreutidae

Ancylis comptana (Frölich)

Procter, 1946: 433, Maine (as *Phyllomyia columbia* (Curran)).

Ancylis comptana floridana (Zeller)

All records ex *Ancylis floridana*.

*Procter, 1946: 435, Maine (Procter, 1938); Procter, 1938: 375, Maine.

Ancylis comptana fragariae (Walsh and Riley)

Allen, 1962: 11, 115; Neiswander, 1944: 8, Ohio.

Evora hemidesma (Zeller)

Roberts, 1966: 169, Iowa.

Grapholitha molesta (Busck)

Most records ex *Laspeyresia molesta*.

Allen, 1962: 12, Ill., Ind., Md., Mo., N.Y., Ohio, Va. (ex *Grapholitha*); Steenburgh and Boyce, 1938: 73, Ont. (as "*Anachaetapsis tortiricus*"); Steenburgh, 1935: 71, Ont.

Laspeyresia pomonella (Linnaeus)

Most records ex *Carpocapsa pomonella*.

Allen, 1962: 12, 115; Lloyd, 1944: 461-462, 468, Calif. (as "*Achaetopsis*" ex codling moth); Yothers and Van Leeuwen, 1931: 26, Oreg.; Aldrich, 1923: 54, Oreg. (as *A. vagans* Aldrich).

Melissopus latiferreanus (Walsingham)

Dohanian, 1942: 837, 840, Oreg., Wash.

Proteoteras willingana (Kearfott)

Peterson, L. O. T., 1958: 646, Sask.

Rhyacionia buoliana (Schifferrmüller)

Torgersen and Coppel, 1969: 164-165, Wis. (as *Erynnia*).

Psychidae

Thyridopteryx ephemeraeformis (Haworth)

Kulman, 1965a: 864, W. Va.; Schaffner, 1959: 46, 49, [N.E. U.S.]; Schaffner and Griswold, 1934: 102-103, 107, N.J.

Pyralidae

Desmia funeralis (Hübner)

Allen, 1962: 117; Donohoe and Kaloostian, 1937: 967-968, Calif.

Etiella zinckenella (Treitschke)

Stone, 1965: 24, Calif. (as *Erynnia*).

Homoeosoma electellum (Hulst)

Satterthwait and Swain, 1946: 577, Mo.

Moodna ostrinella (Clemens)

Raizenne, 1952: 179, Ont. (as *Tortriciophaga*).

Tortricidae

Acleris variana (Fernald)

Reeks, Forbes and Cuming, 1947: 12, [maritime prov.] (as *Tortriciophaga*).

Archips cerasivoranus (Fitch)

Schaffner, 1959: 30, 49, [N.E. U.S.].

Archips fervidanus (Clemens)

Schaffner, 1959: 29, 49, [N.E. U.S.].

Archips rosanus (Linnaeus)

Schaffner, 1959: 31, 49, [N.E. U.S.].

Choristoneura rosaceana (Harris)Raizenne, 1952: 214, Ont. (as *Tortriciophaga* ex *Archips*).*Platynota flavedana* Clemens

Allen, 1962: 115; Summerland and Hamilton, 1954: 941, Ind.

Sparganothis sulfureana (Clemens)Allen, 1962: 115 (ex "*Sparganothis sulfurana*"); Beckwith, 1938: 255, N.J.

Unidentified Tortricidae

Coquillett, 1897: 17, 27, Calif. (as *Hypostena*).

Unidentified Tortricidae probably

Coquillett, 1895a: 56, Calif. (as *Pseudomyothyria* ex tortricid probably on *Solanum* "*douglassi*" Dunal).

Unidentified Lepidoptera

Miscellaneous

Aldrich, 1928b: 41, N. Amer. (ex Microlepidoptera).

STAGE PARASITIZED: Larva, pupa.

IMMATURES: Egg — Satterthwait and Swain, 1946: (descr.).

Pupa — Greene, 1922: (descr., fig. 32 on pl. 7).

BIOLOGY: Satterthwait and Swain, 1946: (577); Schaffner, 1959: (29-31, 46, 49); Schaffner and Griswold, 1934: (102-103, 107).

Erynnia spp.

HOST: LEPIDOPTERA

Tortricidae

Acleris variana (Fernald)Brown, A. W. A., 1938: 16, Ont. (as *Anachaetopsis* [sp.] ex *Peronea*).

STAGE PARASITIZED: Larva.

Erynniopsis Townsend (Goniinae: Blondeliini)*Erynniopsis rondanii* TownsendMost records as *Erynnia nitida* Robineau-Desvoidy.

HOST: COLEOPTERA

Chrysomelidae

Pyrrhata luteola (Müller)

Most records ex *Galerucella xanthomelaena* (Schrank).

Dowden, 1962: 16, 18, 56-57, Calif. (as *Erynniopsis rondani*); Anon., 1960c: 985, Calif. (probable parasite); Clausen, 1956: 5, 67, 102-103, Calif.; Keifer, 1945: 249, Calif. (ex elm leaf beetle); Flanders, 1942: 607, Calif. (ex elm leaf beetle); Flanders, 1940: 947-948, Calif.; Flanders, 1940a: 412, Calif. (ex elm leaf beetle).

STAGE PARASITIZED: Larva, pupa, adult.

BIOLOGY: Clausen, 1956: (5, 67, 102-103); Flanders, 1942: (607); Flanders, 1940: (947-948); Flanders, 1940a: (412).

Euacemyia tibialis, see *Acemya tibialis*

Euacemyia spp., see *Acemya* spp.

Euadmontia pergandei, see *Admontia pergandei*

Eubiomyia calosomae, see *Sitophaga calosomae*

Eubrachymera Townsend (Dexiinae: Macquartiini)

Eubrachymera debilis Townsend

HOST: HYMENOPTERA

Pamphiliidae

Neurotoma inconspicua (Norton)

Townsend, 1919a: 162, S. Dak.

"*Eucelaria*" *armigera*, see *Eucelatoria armigera*

Eucelatoria Townsend (Goniinae: Blondeliini)

Eucelatoria "amigera," see *Eucelatoria armigera*

Eucelatoria armigera (Coquillett)

HOST: HYMENOPTERA

Diprionidae

Neodiprion spp.

Greene, 1921: 43, Ga. (as *Frontina*).

HOST: LEPIDOPTERA

Arctiidae

Estigmene acrea (Drury)

Carrillo S., Ortega C., and Gibson, W. W., 1966: 94, Mexico (Son.) (ex *E. acrea*).

Geometridae

Unidentified Geometridae

Nakao, 1963a: 535, Hawaii.

Noctuidae

Achaea janata (Linnaeus)

Zimmerman, 1958: 353, Hawaii; Pemberton, 1948: 209, Hawaii; Bianchi, 1946: 469, Hawaii (incomplete rearing — pupa); Van Zwaluwenburg, 1946: 14, Hawaii.

Agrotis ipsilon (Hufnagel)

Most records ex *Agrotis ypsilon*.

Patton, 1958: 37, Fla. (incomplete rearing — pupa); Zimmerman, 1958: 253, Hawaii (ex *A. ipsilon*); Tissot, 1945: 64, Fla.

Amyna natalis (Walker)

Zimmerman, 1958: 348, Hawaii; Pemberton, 1948: 209, Hawaii; Van Zwaluwenburg, 1946: 15, Hawaii.

Autographa californica (Speyer)

Clancy, 1969: 1079, Calif.

Chrysodeixis calcites (Esper)

All records ex *Plusia "chalcites."*

Zimmerman, 1958: 371, Hawaii; Pemberton, 1948: 209, Hawaii; Swezey, 1944: 3, Hawaii; Pemberton, 1943: 16, Hawaii; Van Zwaluwenburg, 1943: 82, Hawaii; Zimmerman, 1943: 267, Hawaii.

Heliothis hawaiiensis (Quaintance and Brues)

Zimmerman, 1958: 211, Hawaii.

Heliothis phloxiphaga Grote and Robinson

Jackson, Bryan and Patana, 1969: 907-910 (lab.).

Heliothis virescens (Fabricius)

Bryan, D. E., Jackson and Stoner, 1969: 1, 5-6, 12 (lab.); Jackson, C. G., Bryan and Patana, 1969: 907-910 (lab.).

Heliothis zea (Boddie)

Bryan, D. E., Jackson and Stoner, 1969: 1, 5-6, 12 (lab.); Jackson, C. G., Bryan and Patana, 1969: 907-910, Ariz.; Bottrell, Young, Price and Adams, 1968: 1054, Okla.; Bosch and Hagen, 1966: 25, Calif.; Zimmerman, 1958: 214, Hawaii; Pemberton, 1948: 209, Hawaii (ex *H. armigera* (Hübner)); Van Zwaluwenburg, 1945: 215, Hawaii (ex *H. armigera*); Pemberton, 1943: 16 (ex *H. armigera*, lab.); Bibby, 1942: 943, Tex. (as *Anetia* ex *H. armigera*); Essig, 1926: 582, Calif. (as *Frontina* ex *H.*

obsoleta (Fabricius)); Essig, 1915: 397, Calif. (as *Frontina* ex *Chloridea obsoleta*); Essig, 1913: 165, Calif. (as *Frontina* ex *H. obsoleta*); Quaintance and Brues, 1905: 123, Calif. (as *Frontina* ex *H. obsoleta*); Coquillett, 1897: 15, 25, Calif. (as *Frontina* ex *H. armiger*); Coquillett, 1889: 331-332, Calif. (as *Tachina* ex *H. armigera*).

Heliothis spp. (including *H. hawaiiensis* (Quaintance and Brues) and/or *H. zea* (Boddie))

Zimmerman and Fletcher, 1956: 173, Hawaii.

Heliothis spp.

Bottrell, Young, Price and Adams, 1968: 1054, Okla.; Butler, 1958: 562, Ariz.; Van Zwaluwenburg, 1943: 82, 85 (lab.).

Leucania cholica (Dyar)

Van Zwaluwenburg, 1951: 373, W. Mexico (ex *Cirphis*).

Leucania inconspicua Herrich-Schaeffer

Most records ex *Cirphis inconspicua*.

Scaramuzza and Barry, 1960: 997, Cuba (as *Blondelia* ex *Pseudaletia*); Scaramuzza, 1951a: 297-298, Cuba (as *Frontina*); Van Dine and Christenson, 1932: 2, Cuba (as *Frontina* ex *C. "inconspicua"*); Cardin, 1915: 113, Cuba (as *Frontina* ex *C. "inconspicua"*).

Leucania latiuscula Herrich-Schaeffer

All records ex *Cirphis latiuscula*.

Van Zwaluwenburg, 1951: 373, Mexico; Van Zwaluwenburg, 1943: 81, Mexico (V.C.).

Mocis spp. (including *M. latipes* (Guenée) and/or *M. repanda* (Fabricius))

Most records ex *Mocis repanda*.

Scaramuzza and Barry, 1960: 997, Cuba (as *Blondelia*); Scaramuzza, 1951a: 300, Cuba (as *Frontina*); Scaramuzza, 1946: 549, Cuba (as *Frontina*); Van Dine and Christenson, 1932: 2, Cuba (as *Frontina*); Scaramuzza, 1929: 123, Cuba (as *Frontina*); Scaramuzza, 1929a: 113, Cuba (as *Frontina*); Cardin, 1915: 113, Cuba (as *Frontina* ex "*Remigera*").

Plusia spp.

Pemberton, 1943: 16, Hawaii; Van Zwaluwenburg, 1943: 83-84, Hawaii.

Polydesma umbricola Boisduval

Miyahira and Nakao, 1969: 439, Hawaii (probable parasite); Kamran, 1968: 1011, Hawaii; Zimmerman, 1958: 392, Hawaii; Pemberton, 1948: 209, Hawaii; Pemberton,

1948b: 195-196, 200, Hawaii; Van Zwaluwenburg, 1946: 16, Hawaii; Van Zwaluwenburg, 1946a: 476, Hawaii; Williams, 1946: 479, Hawaii.

Pseudaletia unipuncta (Haworth)

Most records ex *Cirphis unipuncta*.

Tanada, 1966: 307-308, Hawaii (ex *Pseudaletia*); Scaramuzza and Barry, 1960: 997, Cuba (as *Blondelia* ex *Pseudaletia*); Zimmerman, 1958: 305, Hawaii (ex *Pseudaletia*); Pemberton, 1951: 402, Hawaii; Scaramuzza, 1951a: 298, Cuba (as *Frontina*); Van Zwaluwenburg, 1951: 373, Mexico; Pemberton, 1945: 220, Hawaii; Pemberton, 1944: 18, Hawaii; Van Dine and Christenson, 1932: 2, Cuba (as *Frontina*).

Spodoptera exempta (Walker)

Most records ex *Laphygma exempta*.

Zimmerman, 1958: 336, Hawaii (ex *Spodoptera*); Van Zwaluwenburg, 1955: 375, Hawaii; Bess, 1951: 212, Hawaii; Pemberton, 1951: 402, Hawaii; Bianchi, 1949a: 345, Hawaii; Pemberton, 1948: 209, Hawaii; Pemberton, 1948b: 195-196, 200, Hawaii; Pemberton, 1944: 18, Hawaii; Van Zwaluwenburg, 1944: 4 (as "*Eucelaria*," lab.); Van Zwaluwenburg, 1944a: 12 (lab.); Pemberton, 1943: 16 (lab.); Van Zwaluwenburg, 1943: 82-87 (lab.).

Spodoptera exigua (Hübner)

Bosch and Hagen, 1966: 24, Calif.

Spodoptera frugiperda (J. E. Smith)

Most records ex *Laphygma frugiperda*.

Scaramuzza and Barry, 1960: 997, Cuba (as *Blondelia*); Scaramuzza, 1951a: 299-300, Cuba (as *Frontina*); *Van Zwaluwenburg, 1943: 82, Cuba (Bruner, S. C., 1934); Scaramuzza, 1942: 148, Cuba (as *Frontina*); Bruner, S. C., 1934: 96, Cuba (as *Frontina*); Van Dine and Christenson, 1932: 2, Cuba (as *Frontina*); Scaramuzza, 1929: 122, Cuba (as *Frontina* ex *L. [frugiperda]*); Scaramuzza, 1929a: 113, Cuba (as *Frontina* ex *L. [frugiperda]*); Cardin, 1915: 113, Cuba (as *Frontina*).

Spodoptera mauritia (Boisduval)

Tanada and Beardsley, 1958: 434, Hawaii.

Spodoptera praefica (Grote)

All records ex *Prodenia praefica*.

Bosch and Hagen, 1966: 26, Calif.; *Van Zwaluwenburg, 1943: 82, Calif. (Blanchard and Conger, 1932); Blanchard and Conger, 1932: 1065-1067, Calif.

Trichoplusia ni (Hübner)

Clancy, 1969: 1079, Calif.; Jackson, Bryan and Patana, 1969: 907-910 (lab.); Bosch and Hagen, 1966: 25, Calif.; Oatman, 1966: 1137-1138, Calif.; Butler, 1958: 562, Ariz.

Armyworm(-s)

Baver, 1950: 24, Hawaii; Bianchi, 1949: 328, Hawaii (ex armyworm in sugarcane); Pemberton, 1948a: 73, Hawaii.

Pieridae

Pieris rapae (Linnaeus)

Oatman, 1966: 1138, Calif.

Pyralidae

Hedylepta accepta (Butler)

All records ex *Omiodes accepta*.

Baver, 1950: 24, Hawaii; Pemberton, 1948: 209, Hawaii.

Hedylepta blackburni (Butler)

Van Zwaluwenburg, 1950: 7, Hawaii (ex *Omiodes*).

Herpetogramma licarsisalis (Walker)

Anon., 1969c: 260, Hawaii; Davis, 1969: 315, Hawaii; Davis and Chong, 1969: 320, Hawaii; Au, 1968: 17, Hawaii (as *E. "amigera"* ex *Pachyzancla*); Miyahira, 1968a: 562, Hawaii (probable parasite).

Hymenia recurvalis (Fabricius)

Pemberton, 1948: 209, Hawaii; Rosa, 1947: 6, Hawaii.

STAGE PARASITIZED: Larva, pupa.

IMMATURES: Larva — Van Zwaluwenburg, 1943: (descr.).

Pupa — Greene, 1922: (descr., fig. 59 on pl. 13).

BIOLOGY: Bryan, D. E., Jackson and Stoner, 1969: (1, 5-6, 12); Jackson, C. G., Bryan and Patana, 1969: (907-910); Kamran, 1968: (1011); Pemberton, 1948b: (195-196, 200); Van Zwaluwenburg, 1944a: (12); Van Zwaluwenburg, 1943: (82-87).

Eucelatoria armigera (Coquillett) probably

HOST: LEPIDOPTERA

Noctuidae

Peridroma saucia (Hübner)

Zimmerman, 1958: 290, Hawaii (ex *P. porphyrea* (Denis and Schiffermüller)).

Eucelatoria australis TownsendMost records as *Compsilura oppugnator* Walton.

HOST: LEPIDOPTERA

Hesperiidae

Calpodes ethlius (Stoll)Aldrich, 1927b: 19, West Indies (as *E. australis*).

Noctuidae

Leucania latiuscula Herrich-SchaefferMost records ex *Cirphis latiuscula*.Scaramuzza and Barry, 1960: 997, Cuba (ex *Pseudaletia*);

Scaramuzza, 1951a: 298, Cuba; Wolcott, 1951a: 473,

P.R. (as *E. australis*); Wolcott, 1924: 161, 220, P.R.;

Walton, 1914a: 95, P.R.

Pyralidae

Agathodes monstralis GuenéeCardin, 1915: 156, Cuba (as "*Consilura opugnator*").

STAGE PARASITIZED: Larva.

Eucelatoria comosa (Wulp)

HOST: LEPIDOPTERA

Noctuidae

Cutworm(-s)

Ingram, Jaynes and Lobdell, 1939: 97, Fla.

Eucelatoria rubentis (Coquillett)Most records as *Eucelatoria* "*rubentris*."

HOST: LEPIDOPTERA

Noctuidae

Pseudaletia unipuncta (Haworth)Breeland, 1958: 323, 334, Tenn. (as "*Encelatoria*" *rubentis*).*Spodoptera exigua* (Hübner)All records ex *Laphygma exigua*.

Van Zwaluwenburg, 1943: 82, Fla.; Wilson, J. W., 1933: 3,

13-14, Fla.; Wilson, J. W., 1932: 39, Fla.

Armyworm(-s)

Hayslip, Genung, Kelsheimer and Wilson, 1953: 47, Fla. (as

E. rubentis ex army worms and related forms).

STAGE PARASITIZED: Larva.

Eucelatoria rubentis (Coquillett) probably

HOST: LEPIDOPTERA

Noctuidae

Pseudaletia unipuncta (Haworth)

Patton, 1958: 37, Fla. (as *E. sp.* (?dark *rubentis*),
incomplete rearing — pupa).

Eucelatoria "rubentris," see *Eucelatoria rubentis*

Eucelatoria spp.

HOST: LEPIDOPTERA

Noctuidae

Heliothis zea (Boddie)

Sifuentes and Young, 1960: 344, Mexico (as "*Eucletoria*"
(n. sp.?).)

Pyralidae

Hedylepta indicata (Fabricius)

Plank, 1945: 26-27, P.R.

Hymenia recurvalis (Fabricius)

Plank, 1945: 27, P.R. (ex *H. fascialis* (Cramer)).

Eucelatoriopsis Townsend (Goniinae: Blondeliini)

Eucelatoriopsis dimmocki (Aldrich)

Most records as *Anetia dimmocki*.

HOST: COLEOPTERA

Chrysomelidae

Agrioconota bivittata (Say)

Stearns, 1933: 151, 153, Del. (ex *Metriona*).

Chelymorpha cassidea (Fabricius)

Aldrich, 1932: 4, 6-7, Fla., Mass., N.H., Va.

Coptocycla sp.

Aldrich, 1932: 7, Md.

Metriona bicolor (Fabricius)

Aldrich, 1932: 7, La.

Plagiometriona clavata (Fabricius)

Most records ex *Deloyala clavata*.

Fattig, 1949: 20 (as *Lydellohoughia ex Plagiometriona*);
 Brimley, 1938: 363, N.C.; Aldrich, 1932: 7, probably
 N.C. (ex "*Deloyola*").

STAGE PARASITIZED: Larva.

Eucelatoriopsis dimmocki (Aldrich) probably

HOST: COLEOPTERA

Chrysomelidae

Plagiometriona clavata (Fabricius)

All records ex *Coptocycla clavata*.

*Vide Aldrich, 1932: 7, Md. (as *Anetia*, Coquillett, 1897);
 Coquillett, 1897: 17, 23, Md. (as *Hypostena barbata*
 Coquillett).

STAGE PARASITIZED: Larva.

Euceromasia Townsend (Goniinae: Eryciini)

Euceromasia floridensis Reinhard

HOST: LEPIDOPTERA

Yponomeutidae

Urodus parvulus (H. Edwards)

Reinhard, 1957: 111, Fla. (ex *Trichostiba*).

Eucheirophaga James (Tachininae: Ernestiini)

Eucheirophaga lugubris James

HOST: LEPIDOPTERA

Pieridae

Eucheira socialis Westwood

Shaw, 1947: 436-437, Mexico; James, M. T., 1945: 330,
 Mexico (Mor.).

STAGE PARASITIZED: Larva.

"*Eucletoria*" (n. sp.), see *Eucelatoria* spp.

Euclytia Townsend (Phasiinae: Phasiini)

Euclytia flava (Townsend)

IMMATURES: Pupa — Greene, 1922: (as *Clytiomyia*; descr., fig.
 48 on pl. 10).

Eucnephalia Townsend (Goniinae: Goniini)

Eucnephalia sp. probably

HOST: LEPIDOPTERA

Noctuidae

Protorthodes rufula (Grote)Coquillett, 1894: 26, Calif. (ex *Taeniocampa*).

STAGE PARASITIZED: Larva.

Eucoronimyia Townsend (Goniinae: Siphonini)*Eucoronimyia hastata* (Coquillett)All records as *Isoglossa hastata*.

HOST: LEPIDOPTERA

Pyralidae

Titanio belialis DruceEssig, 1926: 579, Colo.; Dyar, 1902: 397, Colo. (ex *Pionea*).

STAGE PARASITIZED: Larva.

Euexorista Townsend (Goniinae: Eryciini)*Euexorista futilis* (Osten Sacken)Most records as *Zenillia futilis*.

HOST: LEPIDOPTERA

Arctiidae

Apantesis oithona Strecker

Learned, 1927: 219 (lab.).

Apantesis phalerata (Harris)

*Sellers, 1943: 12 (Learned, 1927); Learned, 1927: 219 (lab.).

Euchaetias egle (Drury)

Sellers, 1943: 12; Schaffner and Griswold, 1934: 39, 112.

Isia isabella (J. E. Smith)

*Sellers, 1943: 12 (Tothill, 1913); Essig, 1926: 581;

*Aldrich and Webber, 1924: 43, Ont. (Tothill, 1913);
Tothill, 1913: 71, Ont. (as *Exorista*).

Geometridae

Ennomos subsignarius (Hübner)Sellers, 1943: 12; Schaffner and Griswold, 1934: 99-100,
112, Mass.; Aldrich and Webber, 1924: 43.

Hesperiidae

Thymelicus lineola (Ochsenheimer)Arthur, 1962: 1085, Ont. (as *Euexorista*).

Lasiocampidae

Malacosoma americanum (Fabricius)

Kulman, 1965: 68, W. Va. (as *Euexorista*); Sellers, 1943: 12;
Bess, 1936: 605; Schaffner and Griswold, 1934: 88, 112;
West, 1928: 814, N.Y.

Malacosoma californicum californicum (Packard)

*Aldrich and Webber, 1924: 43, Calif. (ex *M. thoracica*
(Stretch), Coquillett, 1897); Coquillett, 1897: 14, 24,
Calif. (as *Exorista ex Clisiocampa thoracica*).

Malacosoma californicum pluviale (Dyar)

Stehr and Cook, 1968: 274-275, 300, Oreg., Wash. (as
Euexorista).

Malacosoma disstria Hübner

Sippell, 1961: 433, 435-436, fig. 6, Ont. (as *Euexorista*);
McGugan, Haliburton and MacDonald, 1952: 45, Ont.
(as *Euexorista*); Raizenne, 1952: 103, Ont. (as
Euexorista); Sellers, 1943: 12; Sellers, 1943: 12 (ex *M.*
disstria erosa form *thoracica*); Bess, 1936: 605; Schaffner
and Griswold, 1934: 89, 112; Essig, 1926: 581, Calif.;
Aldrich and Webber, 1924: 43, N.Y.

Malacosoma spp.

Ross, 1953: 20, 23, fig. 11, B.C. (as *Euexorista*).

Noctuidae

Apamea apamiformis (Guenée)

Most records ex *Hadena apamiformis*.

Sellers, 1943: 12 (ex *Septis*); Johnson, 1910: 780, N.J. (as
Exorista); Coquillett, 1897: 14, 25 (as *Exorista*);
Brauer and Bergenstamm, 1895: 567, 593, N. Amer.
(as *Parexorista*).

Autographa californica (Speyer)

*Sellers, 1943: 12 (Essig, 1926); Essig, 1926: 581, Calif.

Hyppa xylinoides (Guenée)

Tothill, 1913: 71, Ont. (as *Exorista*).

Nephelodes emmedonia (Cramer)

Sellers, 1943: 12; Schaffner and Griswold, 1934: 46, 112.

Nycteola spp. (including *N. cinereana* Neumoegen and Dyar, *N.*
columbiana H. Edwards and/or *N. frigidana* Walker)

Sellers, 1943: 12 (ex *Sarrothripus revayanus* (Scopoli));
Schaffner and Griswold, 1934: 63-64, 112 (ex *N.*
revayana).

Sideridis maryx (Guenée)

Brauer and Bergenstamm, 1895: 567, 599 (as *Parexorista* ex *Mamestra vindemialis* Grote).

Nymphalidae

Nymphalis milberti (Godart)

Sellers, 1943: 12; Schaffner and Griswold, 1934: 11, 112 (ex *Hamadryas*).

Vanessa atalanta (Linnaeus)

Sellers, 1943: 12; Schaffner and Griswold, 1934: 12, 112 (ex *Cynthia*); Aldrich, 1932: 3, Mass.; Essig, 1926: 581; Webber and Schaffner, 1926: 17, New England; Johnson, 1925: 197, Conn., Maine, Mass., N.H., R.I., Vt. (ex "*Vanassa*"); Johnson, 1925a: 87, New England; Aldrich and Webber, 1924: 43; Johnson, 1910: 780, N.J. (as *Exorista* ex *Pyrameis*); Hitchings, 1908: 11, Maine (as *Exorista*); *Brauer and Bergenstamm, 1895: 567, 609 (as *Parexorista*, Williston, 1889); Williston, 1889: 1914, 1918, Conn. (as *Exorista*); Osten Sacken, 1887: 162 (as *Tachina*).

Vanessa cardui (Linnaeus)

*Sellers, 1943: 12 (Schaffner and Griswold, 1934); Schaffner and Griswold, 1934: 13, 112 (ex *Cynthia*).

Vanessa spp.

Sellers, 1943: 12.

Pyralidae

Ostrinia nubilalis (Hübner)

All records ex *Pyrausta nubilalis*.

Sellers, 1943: 12; Aldrich and Webber, 1924: 43, Mass.

Saturniidae

Antheraea polyphemus (Cramer)

Hitchings, 1908: 11, Maine (as *Exorista* ex *polyphemus* moth).

Unidentified Lepidoptera

Miscellaneous

Essig, 1926: 581, Oreg., N. Amer. (ex many moths and butterflies).

STAGE PARASITIZED: Larva, pupa.

IMMATURES: Pupa — Sippell, 1961: (descr., fig. 6); Ross, 1953: (descr., fig. 11); Greene, 1922: (descr., fig. 93 on pl. 19).

BIOLOGY: Sellers, 1943: (12); Schaffner and Griswold, 1934: (112); Learned, 1927: (219).

Eufrontina Brooks (Goniinae: Eryciini)

Eufrontina ethniae Brooks

HOST: LEPIDOPTERA

Ethmiidae

Ethmia longimaculella (Clemens)

Brooks, 1945: 91, Ont. (ex "*Ethnia*").

Eufrontina spectabilis (Aldrich)

HOST: LEPIDOPTERA

Pyralidae

Canarsia ulmiarrosorella (Clemens)

Raizenne, 1952: 178, Ont.

Tetralopha asperatella (Clemens)

Brooks, 1945: 91, Que.

Tetralopha robustella Zeller

Schaffner, 1959: 13, 48, Mass., N.H. (as *Achaetoneura*);

Webber, 1930: 4 (as *Achaetoneura*).

IMMATURES: Pupa — Brooks, 1945: (descr.).

BIOLOGY: Schaffner, 1959: (13, 48).

Euhaliidaya Walton (Goniinae: Acemyini)

Euhaliidaya genalis (Coquillett)

HOST: ORTHOPTERA

Phasmatidae

Diapheromera femorata (Say)

Walton, 1914b: 132, Wis. (as "*Euhaliidaya*" *severinii* Walton).

Diapheromera spp.

Walton, 1918: 23 (as *E. severinii*).

STAGE PARASITIZED: Adult.

Euhaliidaya severinii, see *Euhaliidaya genalis*

"*Euhaliidaya*" *severinii*, see *Euhaliidaya genalis*

Eulasiona Townsend (Dexiinae: Macquartiini)

Eulasiona comstocki Townsend

HOST: LEPIDOPTERA

Oecophoridae

Psilocorsis sp.

Daviault, 1935: 245, Que.

Olethreutidae

Epinotia sollicitana (Walker)

Smith, C. C., 1946: 120, N.B.

Pseudexentera improbana oregonana (Walsingham)Wong, H. R., and Melvin, 1967: 4, Man. (ex *P. oregonana*).

STAGE PARASITIZED: Larva.

Eulasiona genalis (Townsend)All records as *Paramuscopteryx genalis*.

HOST: LEPIDOPTERA

Aegeriidae

Unidentified Aegeriidae

Tilden, 1954: 43, Calif.; Tilden, 1951: 164, 166, Calif. (ex host feeding on *Baccharis pilularis* DeCandolle).

STAGE PARASITIZED: Larva.

Eulasiona nigra Curran

HOST: LEPIDOPTERA

Tortricidae

Archips rosanus (Linnaeus)Curran, 1924: 194, B.C. (ex *Cacoecia*).

STAGE PARASITIZED: Larva.

Euloewia globosa, see *Myiophasia globosa**Eumasicera* Townsend (Goniinae: Eumasicerini)*Eumasicera sternalis* (Coquillett)

HOST: LEPIDOPTERA

Citheroniidae

Anisota rubicunda (Fabricius)

Schaffner and Griswold, 1934: 30-31, 119.

Anisota senatoria J. E. Smith

Schaffner and Griswold, 1934: 29-30, 119.

STAGE PARASITIZED: Pupa.

BIOLOGY: Schaffner and Griswold, 1934: (119).

Eumasicera sp. near *sternalis* (Coquillett)

HOST: LEPIDOPTERA

Noctuidae

Acronicta impleta Walker

Schaffner and Griswold, 1934: 55, 119 (ex *Acronycta*).

Acronicta sp.

Schaffner and Griswold, 1934: 60, 119 (ex *Acronycta*).

STAGE PARASITIZED: Pupa.

BIOLOGY: Schaffner and Griswold, 1934: (119).

Eumea caesar, see *Aplomya caesar*

Eumyothyria sp., see *Cryptomeigenia* spp.

Eupelecotheca Townsend (Goniinae: Neominthoini)

Eupelecotheca celer Townsend

HOST: LEPIDOPTERA

Limacodidae

Packardia ceanothi Dyar

Raizenne, 1952: 160, Ont.

Euphasiopteryx Townsend (Proseninae: Ormiini)

Euphasiopteryx brevicornis nuttingi Sabrosky

HOST: ORTHOPTERA

Tettigoniidae

Neoconocephalus ensiger (Harris)

Nutting, 1953: 79, Mass. (as *E. brevicornis* (Townsend)).

Neoconocephalus robustus (Scudder)

Sabrosky, 1953: 293, Mass.

Neoconocephalus robustus robustus (Scudder)

Nutting, 1953: 69-81, fig. 3, figs. 2-10 on pl. 4, Mass. (as *E. brevicornis*).

Unidentified Orthoptera

Miscellaneous

Sabrosky, 1953: 293.

STAGE PARASITIZED: Adult.

IMMATURES: First instar larva — Nutting, 1953: (descr., figs. 2-5 on pl. 4).

Second instar larva — Nutting, 1953: (descr., fig. 10 on pl. 4).

Third instar larva — Nutting, 1953: (descr., figs. 6, 9 on pl. 4).

Pupa — Nutting, 1953: (descr., fig. 3, figs. 7-8 on pl. 4).

BIOLOGY: Nutting, 1953: (69-81).

Euphasiopteryx ochracea (Bigot)

HOST: ORTHOPTERA

Gryllidae

Gryllus sp.Reinhard, 1922: 72-73, Tex. (as *Ormia* ex *G. assimilis* (Fabricius), incomplete rearing — pupa).

STAGE PARASITIZED: Adult.

IMMATURES: Pupa — Greene, 1922: (descr., fig. 12 on pl. 3).

Euphorocera Townsend (Goniinae: Exoristini)**Euphorocera claripennis** (Macquart)Most records as *Phorocera claripennis*.

HOST: COLEOPTERA

Chrysomelidae

Calligrapha multipunctata (Say)Coquillett, 1897: 11, 23, Mo. (as *Euphorocera* ex *Chrysomela*).

Coccinellidae

Epilachna borealis FabriciusJohnson, 1910: 780, N.J. (as *Euphorocera* ex "*Epilachnae*");Chittenden, 1899: 16 (as *Euphorocera*); Coquillett,1897: 12, 23, N.J. or N.Y. (as *Euphorocera*).*Epilachna varivestis* Mulsant

Howard, N. F., and Landis, 1936: 2, 4-5, fig. 1B, Ala., Ky.,

Md., N.C., N.J., Ohio, Tenn., Va.; Friend and Turner,

1931: 92, Ala. (ex *E. corrupta* Mulsant); Howard, N. F.,1930: 7, Ala. (ex *E. corrupta*); Howard, N. F., 1924: 7,Ala. (ex *E. corrupta*).

Unidentified Coleoptera

Miscellaneous

Balduf, 1937: 173; Aldrich and Webber, 1924: 59.

HOST: HYMENOPTERA

Diprionidae

Diprion similis (Hartig)

Webber, 1932: 206.

Neodiprion abbotii (Leach)

Hetrick, 1941: 376, Va. (ex *N. americanus* (Leach)).

Neodiprion lecontei (Fitch)

Benjamin, 1955: 20, U.S.; Webber and Schaffner, 1926: 29, New England (ex *Lophyrus*); Greene, 1921: 41, Wis.; Middleton, 1921: 757.

Neodiprion sertifer (Geoffroy)

Raizenne, 1957: 14, Ont. (as *Neophorocera*).

Neodiprion virginiana Rohwer

Greene, 1921: 43, W. Va.

Neodiprion sp.

Greene, 1921: 41, W. Va.

Diprionid spp. (including *Diprion* spp. and/or *Neodiprion* spp.)

All records ex *Lophyrus* sp.

Fattig, 1949: 26; Brimley, 1922: 24-25, N.C.; Coquillett, 1897: 12, 22, Tenn. (as *Euphorocera*).

Unidentified Hymenoptera

Sawfly(-flies)

Balduf, 1937: 173; Aldrich, 1932: 4, N.H. (ex larvae on *Pinus rigida* Miller); Aldrich and Webber, 1924: 59.

HOST: LEPIDOPTERA

Arctiidae

Apantesis oithona Strecker

Allen, 1929: 686, Miss.

Apantesis proxima (Guérin-Méneville)

Essig, 1926: 582, Calif.; Coquillett, 1897: 11, 24, Calif. (as *Euphorocera* ex *Arctia docta* (Walker)).

Eubaphe aurantiaca rubicundaria (Hübner)

Coquillett, 1897: 11, 24, Fla. (as *Euphorocera* ex *Crocota rubicundaria*).

Euchaetias egle (Drury)

Schaffner and Griswold, 1934: 39, 114; Webber and Schaffner, 1926: 20, New England.

Halisidota tessellaris (J. E. Smith)

Schaffner and Griswold, 1934: 34, 114; Coquillett, 1897: 12, 25, N.Y. (as *Euphorocera*).

Hyphantria cunea (Drury)

Schaffner and Griswold, 1934: 37, 114; Essig, 1926: 582, Calif.; Johnson, 1925: 198, Conn., Maine, Mass., N.H.,

R.I. (ex fall webworm); Howard, 1897b: 43, fig. 23, D.C.
(as *Euphorocera*).

Utetheisa ornatrix bella (Linnaeus)

Schaffner and Griswold, 1934: 39-40, 114, N.J. (ex *U. bella*).

Citheroniidae

Anisota rubicunda (Fabricius)

Schaffner and Griswold, 1934: 30-31, 114.

Anisota senatoria J. E. Smith

Schaffner and Griswold, 1934: 29-30, 114; Johnson, 1910:
780, N.J. (as *Euphorocera*); Coquillett, 1897: 11, 23 (as
Euphorocera).

Ctenuchidae

Lymire edwardsii (Grote)

Bratley, 1929: 44, Fla.

Gelechiidae

Anarsia lineatella Zeller

Essig, 1926: 582; Howard, 1897b: 43, fig. 23, D.C. (as
Euphorocera).

Geometridae

Cingilia catenaria (Cramer)

Schaffner and Griswold, 1934: 98, 114; Aldrich, 1932: 2,
N.H.

Itame sulphurea (Packard)

Franklin, 1950: 57.

Philtraea elegantaria (H. Edwards)

Tucker, E. S., 1917: 395, La. (as *Phorocera*).

Semiothisa neptaria (Guenée)

Coquillett, 1897: 12, 26, Calif. (as *Euphorocera* ex *Phasiana*
"neptata").

Hesperiidae

Epargyreus clarus (Cramer)

All records ex *Epargyreus tityrus* (Fabricius).

Schaffner and Griswold, 1934: 15, 114; Allen, 1929: 686,
Miss.

Lasiocampidae

Malacosoma americanum (Fabricius)

Bess, 1936: 605; Schaffner and Griswold, 1934: 88, 114.

Malacosoma disstria Hübner

Bess, 1936: 605; Schaffner and Griswold, 1934: 89, 114;
Essig, 1926: 582, Calif.; Johnson, 1910: 780, N.J. (as
Euphorocera ex Clisiocampa); Coquillett, 1897: 11, 24,
Mass. (as *Euphorocera ex Clisiocampa*).

Malacosoma sp.

Tothill, 1913: 70 (as *Euphorocera*).

Apple tree tent caterpillar(-s)

Brimley, 1922: 24-25, N.C.

Limacodidae

Sibine stimulea (Clemens)

All records ex *Empretia stimulea*.

Johnson, 1910: 780, N.J. (as *Euphorocera*); Coquillett,
1897: 11, 25, Ind. (as *Euphorocera*).

Lymantriidae

Euproctis chrysorrhoea (Linnaeus)

Howard and Fiske, 1911: 93 (as *Euphorocera*); Fernald,
1898: 28, Mass. (as *Euphorocera*).

Hemerocampa leucostigma (J. E. Smith)

Franklin, 1950: 57; Dowden, 1935: 509; Schaffner and
Griswold, 1934: 85, 114; Essig, 1926: 582; Webber and
Schaffner, 1926: 12-13, 20, Mass., New England, N.Y.,
Pa., R.I.; Britton, 1917: 110 (as *Euphorocera*); Johnson,
1910: 780, N.J. (as *Euphorocera ex Orgyia*); Howard,
1898: 88, D.C. (as *Euphorocera ex Orgyia*); Coquillett,
1897: 12, 26, D.C. (as *Euphorocera ex Orgyia*); Howard,
1897a: 16-17, D.C. (as *Euphorocera ex Orgyia*); Howard,
1897b: 7, 43, 49, fig. 23, D.C. (as *Euphorocera ex
Orgyia*).

Nygmia phaeorrhoea (Donovan)

Burgess and Crossman, 1929: 143.

Porthetria dispar (Linnaeus)

Burgess and Crossman, 1929: 143.

Stilpnotia salicis (Linnaeus)

Burgess and Crossman, 1927: 14, New England.

Tussock moth(-s)

Johnson, 1925: 198, Conn., Maine, Mass., N.H., R.I.

Megalopygidae

Lagoa sp.

Coquillett, 1897: 12, 25 (as *Euphorocera*).

Megalopyge crispata (Packard)

All records ex *Lagoa crispata*.

Schaffner, 1959: 7, 52, [N.E. U.S.]; Schaffner and
Griswold, 1934: 105-106, 114; Aldrich, 1932: 3, Mass.

Megalopyge opercularis (J. E. Smith)

Bishop, 1923: 13, Tex.; Reinhard, 1922: 73, Tex.

Megalopyge pyxidifera (J. E. Smith)

Patton, 1958: 80, Fla.

Noctuidae

Acronicta hamamelis Guenée

Coquillett, 1897: 11, 23, Va. (as *Euphorocera* ex *Acronycta*).

Actebia fennica (Tauscher)

Phipps, 1927: 212 (ex *Agrotis*); Britton, 1926: 324, Conn.
(ex *Noctua*).

Agrotis ipsilon (Hufnagel)

All records ex *Agrotis ypsilon*.

Franklin, 1950: 57; Crumb, 1929: 40; Essig, 1926: 582,
Calif.; Howard, 1897b: 43, fig. 23, D.C. (as
Euphorocera).

Alabama argillacea (Hübner)

Most records ex *Aletia argillacea*.

Essig, 1926: 582 (ex *Alabama*); Coquillett, 1897: 11, 23
(as *Euphorocera*); Howard, 1897b: 43, fig. 23, D.C. (as
Euphorocera).

Noctuidae

Amathes c-nigrum (Linnaeus)

Most records ex *Agrotis c-nigrum*.

Franklin, 1950: 57 (ex *Amathes*); *Crumb, 1929: 40, 140
(Franklin and Lacroix, 1924); Franklin and Lacroix,
1924: 408, Mass.

Eubolina sp. probably

Coquillett, 1897: 12, 25, Calif. (as *Euphorocera*).

Euxoa auxiliaris (Grote)

Walkden, 1943: 381 (ex *Chorizagrotis*).

Faronta diffusa (Walker)

Essig, 1926: 582, B.C. (ex *Neleucania albilinea* (Hübner));
Webster, R. L., 1911: 181, Iowa (as *Euphorocera* ex
Meliana albilinea); Howard, 1897b: 43, fig. 23, D.C. (as
Euphorocera ex *Leucania albilinea*).

Feltia ducens Walker — *Feltia subgothica* (Haworth) complex
Crumb, 1929: 40, 68.

Feltia herilis (Grote)

Coquillett, 1897: 12, 25, Mo., Tenn. (as *Euphorocera*);
Howard, 1897b: 43, fig. 23, D.C. (as *Euphorocera*).

Feltia subgothica (Haworth)

Brimley, 1922: 24-25, N.C.

Heliothis virescens (Fabricius)

Patton, 1958: 80, Fla.

Heliothis zea (Boddie)

All records ex *Heliothis obsoleta* (Fabricius).

Essig, 1926: 582; Quaintance and Brues, 1905: 123, Tex.
(as *Euphorocera*).

Hydroecia immanis Guenée

Howard, 1897b: 43, fig. 23, D.C. (as *Euphorocera*).

Lithophane spp. (including *L. disposita* Morrison, *L.*
innominata (Smith) and/or *L. petulca* Grote)

Schaffner and Griswold, 1934: 48, 114 (ex *Graptolitha* spp.
(including *G. disposita*, *G. innominata* and/or *G.*
petulca)).

Mocis spp. (including *M. latipes* (Guenée) and/or *M. repanda*
(Fabricius))

Most records ex *Remigia repanda*.

Wolcott, 1951a: 478, P.R. (as *Euphorocera* ex *Mocis*);
Wolcott, 1924: 222, P.R. (as *Euphorocera*); Jones, T.
H., and Wolcott, 1922: 49, P.R.

Peridroma saucia (Hübner)

Crumb, 1929: 40 (ex *Lycophotia*); Essig, 1926: 582, Oreg.
(ex *Lycophotia margaritosa* (Haworth)); Lovett, 1915:
145, Oreg. (as *Euphorocera* ex *P. margaritosa saucia*).

Plathypena scabra (Fabricius)

Fattig, 1949: 26; Smith, R. C., 1943: 378, Kans. (ex green
clover worm); Hill, 1925: 17, Md.; Brimley, 1922: 24-25,
N.C. (as green clover worm); Sherman, 1920: 298-299,
N.C.

Plathypena scabra (Fabricius) probably

Patton, 1958: 80, Fla.

Pseudaletia unipuncta (Haworth)

Most records ex *Cirphis unipuncta*.

Fattig, 1949: 26, Ga.; Schaffner and Griswold, 1934: 47, 114; Essig, 1926: 582, Canada; Baker, 1915: 86, Ont. (ex *Heliophila*); Gibson, 1915: 14, Ont.; Sherman, 1915: 299-302, N.C. (ex *Heliophila*); Tothill, 1913: 70, Ont. (as *Euphorocera* ex *Heliophila*).

Scotogramma trifolii (Rottenburg)

Most records ex *Mamestra trifolii*.

Essig, 1926: 582, Colo. (ex *Scotogramma*); Marsh, 1913: 17, Colo. or Kans.; Titus and Pratt, 1904: 77 (as *Euphorocera* ex *M. "tripolii"*); Coquillett, 1897: 12, 26, D.C., Mo. (as *Euphorocera*); Howard, 1897b: 43, fig. 23, D.C. (as *Euphorocera*).

Spodoptera eridania (Cramer)

Patton, 1958: 80, Fla. (ex *Prodenia*).

Spodoptera frugiperda (J. E. Smith)

All records ex *Laphygma frugiperda*.

Luginbill, 1928: 81, Ala., S.C.; Reinhard, 1922: 73, Tex.

Spodoptera praefica (Grote)

All records ex *Prodenia praefica*.

Bosch and Hagen, 1966: 26, Calif.; Blanchard and Conger, 1932: 1067, Calif.

Armyworm(-s)

Patton, 1958: 80, Fla.; Brimley, 1922: 24-25, N.C.

Cutworm(-s)

Knowlton and Harmston, 1937: 148, Utah; Johnson, 1925: 198, Conn., Maine, Mass., N.H., R.I.; Chittenden, 1903: 34, fig. 31 (as "*Euphrocera*").

Noctuidae and/or Pyralidae

Noctuid and/or pyralid spp. (including *Scotogramma trifolii* (Rottenburg) and/or *Loxostege sticticalis* (Linnaeus))

Anon., 1956d: 1044, Idaho (ex clover cutworm and/or beet webworm).

Notodontidae

Datana angusii Grote and Robinson

Schaffner and Griswold, 1934: 74-75, 114.

Datana contracta Walker

Webster, 1899a: 224, Ind. (as *Euphorocera*); Coquillett, 1897: 11, 24, Ind. (as *Euphorocera*).

Datana integerrima Grote and Robinson

Fattig, 1949: 26; Hixson, 1941: 20, Okla.; Haseman, 1940:

12-13, fig. 5, Mo.; Schaffner and Griswold, 1934: 76, 114; Baerg, 1928: 15, Ark.; Leiby, 1925: 23, N.C. (as *Euphorocera* ex "*Dantana*"); Brimley, 1922: 24-25, N.C.

Datana ministra (Drury)

Schaffner and Griswold, 1934: 74, 114; Allen, 1929: 686, Miss.; Essig, 1926: 582; Webster, 1898: 19, Ohio (as *Euphorocera* "*clavipennis*"); Howard, 1897b: 43, fig. 23, D.C. (as *Euphorocera*).

Datana perspicua Grote and Robinson

Fattig, 1949: 26, Ga.; Schaffner and Griswold, 1934: 75, 114.

Datana sp.

Allen, 1924: 24, Miss. (ex *Datana* on oak).

Ichthyura inclusa Hübner

All records ex *Melalopha inclusa*.

Allen, 1929: 686, Miss.; Brimley, 1922: 24-25, N.C.

Schizura concinna (J. E. Smith)

Schaffner and Griswold, 1934: 81, 114; Allen, 1929: 686, Miss.; Essig, 1926: 582; Webber and Schaffner, 1926: 19, New England; Cole, F. R., and Lovett, 1921: 302, Oreg.; Coquillett, 1897: 12, 26, Mo. (as *Euphorocera* ex *Oedemasia*).

Schizura unicornis (J. E. Smith)

Webster and Mally, 1897: 42-43, Ohio (as *Euphorocera* ex *Coelodasys*).

Nymphalidae

Agraulis vanillae Linnaeus

Essig, 1926: 582, Calif. (ex *Dione*); Coquillett, 1897: 11, 23, Calif. (as *Euphorocera*).

Asterocampa clyton (Boisduval and LeConte)

Coquillett, 1897: 11, 24, La. (as *Euphorocera* ex *Apatura*).

Chlosyne lacinia (Geyer)

Reinhard, 1922: 73, Tex. (ex *Synchloe*).

Nymphalis antiopa (Linnaeus)

Schaffner and Griswold, 1934: 11, 114 (ex *Hamadryas*); Essig, 1926: 582 (ex *Aglais*); Johnson, 1910: 780, N.J. (as *Euphorocera* ex *Vanessa*).

Polygonia comma (Harris)

Schaffner and Griswold, 1934: 10, 114.

Polygonia interrogationis (Fabricius)

Schaffner and Griswold, 1934: 9-10, 114.

Pieridae

Colias eurytheme Boisduval

Most records ex *Eurymus eurytheme*.

Anon., 1963: 5 (ex *Colias*); Anon., 1952: 4 (ex *C. philodice eurytheme*); Smith, R. C., 1943: 378, Kans. (ex alfalfa caterpillar); Essig, 1926: 582, Calif., Utah; Wildermuth, 1920: 8; Wildermuth, 1914: 24, fig. 14, Ariz., Calif., Kans., Tenn., Utah; Wildermuth, 1911: 7-8, fig. 6 (as *Euphorocera*).

Pieris rapae (Linnaeus)

Poos, 1928: 147, Va. (ex *Pontia*).

Psychidae

Thyridopteryx ephemeraeformis (Haworth)

Kulman, 1965a: 864, W. Va.; Schaffner, 1959: 46, 52, [N.E. U.S.]; Patton, 1958: 80, Fla.; Schaffner and Griswold, 1934: 102-103, 114.

Pyralidae

Crambus mutabilis Clemens

Ainslie, G. G., 1927: 14-15, Ind.; Ainslie, G. G., 1923: 411-412, Ind.

Crambus spp.

Noble, 1932: 3, U.S.

Pyralidae and/or Noctuidae,
see Noctuidae and/or Pyralidae

Saturniidae

Automeris io (Fabricius)

Schaffner and Griswold, 1934: 28, 114; Webber and Schaffner, 1926: 25, New England.

Callosamia promethea (Drury)

Schaffner and Griswold, 1934: 27, 114.

Hemileuca electra Wright

Essig, 1926: 582, Calif.; Coquillett, 1897: 12, 25, Calif. (as *Euphorocera*).

Hemileuca nevadensis Stretch

Essig, 1926: 582, N. Mex.; Coquillett, 1897: 12, 25, N. Mex. (as *Euphorocera* ex *H. artemis* Packard).

Hemileuca oliviae Cockerell

Wildermuth and Frankenfeld, 1933: 797, [N. Mex.]; Ainslie, 1910: 92, fig. 52, N. Mex. (as *Euphorocera*).

Hemileuca spp. (including *H. lucina* H. Edwards and/or *H. maia* (Drury))

Schaffner and Griswold, 1934: 29, 114.

Platysamia cecropia (Linnaeus)

Schaffner and Griswold, 1934: 26-27, 114.

Sphingidae

Ceratomia catalpae (Boisduval)

Schaffner and Griswold, 1934: 17, 114, N.J.; Howard and Chittenden, 1916: 6, fig. 4; Howard and Chittenden, 1907: 5, fig. 2 (as *Euphorocera*); Coquillett, 1897: 11, 24 (as *Euphorocera*); Howard, 1897b: 43, fig. 23, D.C. (as *Euphorocera*).

Ceratomia undulosa (Walker)

Brimley, 1922: 24-25, N.C.

Sphinx chersis (Hübner)

Essig, 1926: 582, Canada.

Zygaenidae

Harrisina americana (Guérin-Méneville)

Coquillett, 1897: 12, 25, Ga. (as *Euphorocera*).

Unidentified Lepidoptera

Miscellaneous

Smith, R. C., 1943: 378, Kans.; Knowlton and Janes, 1931: 147, Utah; Aldrich and Webber, 1924: 59; Aldrich, 1915: 83 (as *Euphorocera*); Howard and Chittenden, 1907: 5, fig. 2 (as *Euphorocera*).

HOST: ORTHOPTERA

Acrididae

Melanoplus differentialis (Thomas)

Essig, 1926: 582; Morgan, 1901: 25, fig. 11, Miss. delta (as *Euphorocera*).

Grasshopper(-s)

Knowlton and Harmston, 1937: 148, Utah.

STAGE PARASITIZED: Larva (Coleoptera, Hymenoptera,

Lepidoptera), pupa (Coleoptera, Hymenoptera, Lepidoptera), adult (Coleoptera).

IMMATURES: Egg — Haseman, 1940: (fig. 5); Howard, N.F., and Landis, 1936: (descr., fig. 1B); Wildermuth, 1920: (descr.).

Third instar larva — Howard, N. F., and Landis, 1936: (descr.).

Pupa — Howard, N. F., and Landis, 1936: (descr.); Greene, 1922: (descr., fig. 73 on pl. 16); Howard and Chittenden, 1916: (fig. 4); Wildermuth, 1914: (fig. 14); Wildermuth, 1911: (fig. 6); Ainslie, 1910: (fig. 52); Howard and Chittenden, 1907: (fig. 2); Chittenden, 1903: (fig. 31); Morgan, 1901: (fig. 11); Howard, 1897b: (fig. 23).

BIOLOGY: Anon., 1963: (5); Schaffner, 1959: (7, 46, 52); Anon., 1952: (4); Hixson, 1941: (20); Haseman, 1940: (12-13); Howard, N. F., and Landis, 1936: (2, 4-5); Dowden, 1935: (509); Schaffner and Griswold, 1934: (114); Ainslie, G. G., 1923: (411-412); Bishopp, 1923: (13); Wildermuth, 1920: (8); Webster, 1899a: (224); Coquillett, 1897: (11, 24); Howard, 1897b: (7, 43, 49).

Euphorocera claripennis (Macquart) probably

HOST: LEPIDOPTERA

Megalopygidae

Megalopyge opercularis (J. E. Smith)

Micks, 1956: 38-39, Tex. (as *Phorocera*).

STAGE PARASITIZED: Larva.

Euphorocera sp. near *claripennis* (Macquart)

HOST: HYMENOPTERA

Diprionidae

Diprion polytomum (Hartig)

Reeks, 1938: 26, N.B. (as *Phorocera*).

STAGE PARASITIZED: Pupa.

Euphorocera "clavipennis," see *Euphorocera claripennis*

Euphorocera edwardsii (Williston)

Most records as *Neophorocera edwardsii*.

HOST: HYMENOPTERA

Diprionidae

Diprion hercyniae (Hartig)

Finlayson, T., 1960a: 923, 938.

Diprion similis (Hartig)

Finlayson, T., 1962: 273, 280, Canada.

Neodiprion abbotii (Leach)

Finlayson, T., 1963: 476, 503.

Neodiprion lecontei (Fitch)

Finlayson, T., 1963: 479, 503.

Neodiprion sertifer (Geoffroy)

Finlayson, T., 1960: 25, 42, fig. 22.

HOST: LEPIDOPTERA

Lasiocampidae

Malacosoma americanum (Fabricius)

Kulman, 1965: 68, W. Va. (as *Phorocera*).

Malacosoma disstria Hübner

Sippell, 1961: 434, 436, fig. 4, Ont.; Raizenne, 1952: 102, Ont.

Lymantriidae

Hemerocampa leucostigma (J. E. Smith)

Raizenne, 1952: 95, Ont.

Noctuidae

Euxoa messoria (Harris) probably

*Vide Brauer and Bergenstamm, 1895: 569, 581 (as *Phorocera*) ex *Agrotis cochrani* (Riley), Townsend, 1893); Townsend, 1893: 467 (as *Phorocera* ex cutworm).

Euxoa spp. (including *E. messoria* (Harris) and/or *E. ochrogaster* (Guenée)

*Vide pers. comm. Handford, 1970: [B.C.] (Handford, 1962); Anon., 1962: 121, B.C.; Handford, 1962: 19, B.C.

Notodontidae

Datana integerrima Grote and Robinson

Raizenne, 1952: 81, Ont.

Datana ministra (Drury)

Raizenne, 1952: 80, Ont.

Symmerista canicosta Franclemont

Raizenne, 1952: 85, Ont.

Nymphalidae

Anaea andria Scudder

*Brauer and Bergenstamm, 1895: 569, 582 (as *Phorocera*),
Williston, 1889); Williston, 1889: 1914, 1921-1922 (as
Phorocera).

Asterocampa celtis (Boisduval and LeConte)

All records ex *Chlorippe celtis*.

*Brauer and Bergenstamm, 1895: 569, 587 (as *Phorocera*,
Williston, 1889); Williston, 1889: 1914, 1921-1922 (as
Phorocera).

Nymphalis antiopa (Linnaeus)

*Brauer and Bergenstamm, 1895: 569, 609 (as *Phorocera*)
ex *Vanessa*, Williston, 1889); Williston, 1889: 1914,
1921-1922 (as *Phorocera* ex *Euvanessa*).

STAGE PARASITIZED: Larva (Lepidoptera), pupa (Hymen-
optera).

IMMATURES: Third instar larva — Finlayson, T., 1963: (descr.);
Finlayson, T., 1962: (descr.); Finlayson, T., 1960: (descr.,
fig. 22); Finlayson, T., 1960a: (descr.).

Pupa — Sippell, 1961: (descr., fig. 4); Williston, 1889: (descr.).

BIOLOGY: Finlayson, T., 1960a: (923, 938).

Euphorocera floridensis Townsend

Most records as *Phorocera floridensis*.

HOST: COLEOPTERA

Cerambycidae

Saperda sp.

Aldrich and Webber, 1924: 61, Ark.

Coccinellidae

Epilachna varivestis Mulsant

Fattig, 1949: 16, Ga. (as *Euphorocera* ex "*Epilachne*
varivestris").

HOST: LEPIDOPTERA

Arctiidae

Apantesis oithona Strecker

Allen, 1929: 686, Miss.

Hyphantria cunea (Drury)

Swain, 1937: 245, Colo.

Citheroniidae

Anisota sp. near *rubicunda* (Fabricius)

Patton, 1958: 80, Fla.

Ctenuchidae

Lymire edwardsii (Grote)

Patton, 1958: 80, Fla.

Syntomeida epilais (Walker)

Patton, 1958: 80, Fla.

Geometridae

Ennomos subsignarius (Hübner)

Davis, R., 1961: 248, Ga. (as *Euphorocera*).

Melanchroia cephise (Cramer)

Patton, 1958: 80, Fla.

Hesperiidae

Epargyreus clarus (Cramer)

Allen, 1929: 686, Miss. (ex *E. tityrus* (Fabricius)).

Noctuidae

Alabama argillacea (Hübner)

Fattig, 1949: 16, Ga. (as *Euphorocera*).

Anticarsia gemmatilis Hübner

All records ex *Anticarsia* "gemmatilis."

Patton, 1958: 80, Fla.; *Aldrich and Webber, 1924: 61, Fla. (Townsend, 1916); Townsend, 1916: 217, Fla. (as *Euphorocera*); Watson, J. R., 1916: 527, fig. 5 on pl. 39 (as *Euphorocera*).

Heliothis zea (Boddie)

Fattig, 1949: 16 (as *Euphorocera* ex *H. armigera* (Hübner)).

Plathypena scabra (Fabricius)

Whiteside, Burbutis and Kelsey, 1967: 327, Del. (as *Euphorocera*); Fattig, 1949: 16, Ga. (as *Euphorocera*); Aldrich and Webber, 1924: 61, N.C.; Sherman, 1920: 298-299, N.C. (as *Euphorocera*).

Pseudaletia unipuncta (Haworth)

Fattig, 1949: 16, Ga. (as *Euphorocera* ex *Cirphis*).

Pseudoplusia includens (Walker)

Allen, 1929: 686, Miss. (ex *Autographa oo* (Cramer)).

Spodoptera frugiperda (J. E. Smith)All records ex *Laphygma frugiperda*.

Patton, 1958: 80, Fla.; Luginbill, 1928: 81, S.C.; Aldrich and Webber, 1924: 61, Md., S.C.

Notodontidae

Datana integerrima Grote and RobinsonFattig, 1949: 16, Ga. (as *Euphorocera*).*Datana major* Grote and Robinson

Patton, 1958: 80, Fla.

Datana ministra (Drury)

Allen, 1929: 686, Miss.

Ichthyura inclusa HübnerMost records ex *Melalopha inclusa*.Fattig, 1949: 16, Ga. (as *Euphorocera* ex *Ichthyura*);

Allen, 1929: 686, Miss.; Allen, 1926b: 194, Miss.

Ichthyura spp. (including *I. apicalis* Walker or *I. inclusa* Hübner)Allen, 1924: 24, Miss. (ex "*Melanopha*" *apicalis* or "*Melanopha*" *inclusa*).

Sphingidae

Ceratomia catalpae (Boisduval)Fattig, 1949: 16, Ga. (as *Euphorocera*).

Zygaenidae

Harrisina americana (Guérin-Méneville)

Patton, 1958: 80, Fla. (ex grape leaf skeletonizer).

STAGE PARASITIZED: Larva (Lepidoptera), pupa (Lepidoptera).

IMMATURES: Pupa — Watson, J. R., 1916: (fig. 5 on pl. 39).

BIOLOGY: Patton, 1958: (80).

Euphorocera sp. near *floridensis* TownsendAll records as *Phorocera* sp. near *floridensis*.

HOST: LEPIDOPTERA

Pieridae

Pieris protodice Boisduval and LeConte

Bibby, 1961: 328, Ariz.

Saturniidae

Saturnia galbina (Clemens)

Buck and Keister, 1956: 94-96 (ex *Agapema*).

STAGE PARASITIZED: Larva, pupa.

BIOLOGY: Buck and Keister, 1956: (94-96).

Euphorocera gelida, see *Spoggosia gelida*

Euphorocera omissa (Reinhard)

HOST: LEPIDOPTERA

Noctuidae

Heliothis spp.

Bottrell, Young, Price and Adams, 1968: 1054, Okla.

Peridroma saucia (Hübner)

Anon., 1969: 59, Okla.

Spodoptera ornithogalli (Guenée)

Bottrell, 1969: 251, Okla. (ex *Prodenia*).

Pieridae

Colias eurytheme Boisduval

Reinhard, 1934b: 195, Ark. (as *Phorocera* ex *Eurymus*).

Pyralidae

Loxostege sticticalis (Linnaeus)

Strickland, 1946: 172, Alta.

STAGE PARASITIZED: Pupa.

Euphorocera subnitens (Aldrich and Webber) probably

HOST: LEPIDOPTERA

Zygaenidae

Gen. and sp. near *Triplocris*

Patton, 1958: 80, Fla. (as *Phorocera* ex larva on cherry laurel).

STAGE PARASITIZED: Larva.

Euphorocera tachinomoides Townsend

HOST: LEPIDOPTERA

Hesperiidae

Lerodea eufala (Edwards)

Butler, 1958: 562, Ariz. (as *Phorocera*).

Lasiocampidae

Malacosoma americanum (Fabricius)

Stehr and Cook, 1968: 257-258, 299, Tex.; Essig, 1926: 582, Ariz., N. Mex., Mexico (as *Phorocera*); Aldrich and Webber, 1924: 56, Ariz. (as *Phorocera*).

Malacosoma californicum (Packard)

Stehr and Cook, 1968: 264, 300, Ariz., N. Mex.

Malacosoma californicum fragile (Stretch)

Stehr and Cook, 1968: 278-279, 301, Calif., Utah.

Malacosoma incurvum discoloratum (Neumoegen)

Stehr and Cook, 1968: 280, 301, Nev., Utah.

Malacosoma incurvum incurvum (H. Edwards)

Stehr and Cook, 1968: 279-280, 301, Ariz.

Malacosoma tigris (Dyar)

Stehr and Cook, 1968: 257, 299, Tex.

Apple tree tent caterpillar(-s)

Knowlton and Harmston, 1937: 149, Utah (as *Phorocera*).

Noctuidae

Heliothis zea (Boddie)

Bottrell, Young, Price and Adams, 1968: 1054, Okla.

Heliothis spp.

Bottrell, Young, Price and Adams, 1968: 1054, Okla.

Melipotis famelica (Guenée)

Wolcott, 1951a: 478, P.R.

Spodoptera ornithogalli (Guenée)

Bottrell, 1969: 251, Okla. (ex *Prodenia*).

Pieridae

Pieris rapae (Linnaeus)

Oatman, 1966: 1138, Calif.

Psychidae

Thyridopteryx meadii H. Edwards

Comstock, 1937: 23, Calif. (as *Phorocera*).

Unidentified Lepidoptera

Green looper(-s)

Butler, 1958: 562, Ariz. (as *Phorocera* ex green looper on acacia).

STAGE PARASITIZED: Larva.

IMMATURES: Pupa — Greene, 1922: (descr., fig. 94 on pl. 20).
Euphorocera sp. near *tachinomoides* Townsend
 HOST: LEPIDOPTERA

Zygaenidae

Harrisina brillians Barnes and McDunnough

Clausen, 1961: 618, Ariz. (as *Phorocera* sp. near
 “*tachinomiodes*”).

STAGE PARASITIZED: Larva.

Euphorocera sp.

HOST: HYMENOPTERA

Diprionidae

Neodiprion sertifer (Geoffroy)

Raizenne, 1957: 14, Ont. (as *Neophorocera*).

“*Euphorocera*” *claripennis*, see *Euphorocera claripennis*

Eupogona obscura, see *Buquetia obscura*

Eusisyropa Townsend (Goniinae: Carceliini)

Eusisyropa autographae (Sellers)

HOST: LEPIDOPTERA

Noctuidae

Trichoplusia ni (Hübner)

All records ex *Autographa brassicae* Riley.

*Vide Sellers, 1943: 23-24, Cuba (as *Zenillia*, Aldrich and
 Webber, 1924); Aldrich and Webber, 1924: 40, Cuba
 (as *Zenillia blanda blanda* (Osten Sacken)).

Eusisyropa blanda (Osten Sacken)

Most records as *Zenillia blanda*.

HOST: LEPIDOPTERA

Arctiidae

Hyphantria cunea (Drury)

Sellers, 1943: 20; Beaulne, 1939: 120, Que. (as “*Zenilla*”);
 Beaulne, 1939a: 96, Que. (as “*Zenilla*”); Schaffner and
 Griswold, 1934: 37, 111; Sellers, 1930: 572-573.

Hyphantria textor (Harris)

Raizenne, 1952: 29, Ont. (as *Eusisyropa*); Townsend, 1908:
 98, D.C. (as *Eusisyropa*).

Citheroniidae

Anisota rubicunda (Fabricius)

Beaulne, 1939: 120, Que. (as "*Zenilla*"); Beaulne, 1939a: 96, Que. (as "*Zenilla*").

Gelechiidae

Gelechiid webber(-s)

Sellers, 1943: 20 (ex gelechiid webber on willow).

Geometridae

Cingilia catenaria (Cramer)

Sellers, 1943: 20; Schaffner and Griswold, 1934: 98, 111; Aldrich, 1932: 2, N.H.; Sellers, 1930: 572-573; Aldrich and Webber, 1924: 39 (as *Z. blanda blanda*).

Ennomos subsignarius (Hübner)

Ciesla, 1964: 595, Appalachians (as *Eusisyropa*); Davis, R., 1961: 248, Ga. (as *Eusisyropa*); Sellers, 1943: 20; Schaffner and Griswold, 1934: 99-100, 111, Mass.; Knoll, 1932: 1202, Pa.; Sellers, 1930: 572.

Hydria undulata (Linnaeus)

Most records ex *Calocalpe undulata*.

Raizenne, 1952: 113, Ont. (as *Eusisyropa*); Sellers, 1943: 20; Schaffner and Griswold, 1934: 91-92, 111; Sellers, 1930: 572 (ex *Hydria*).

Isturgia truncataria (Walker)

Vide Sellers, 1943: 19-20, 22 (Coquillett, 1897); Aldrich and Webber, 1924: 39, Mass.; Coquillett, 1897: 13, 24, Mass. (as *Exorista boarmiae* Coquillett ex *Boarmia pampinaria* Guenée).

Hopping, 1930: 64, B.C. (ex hemlock looper).

Nepytia phantasmaria (Strecker)

Hopping, 1930: 64, B.C.

Hesperiidae

Calpodes ethlius (Stoll)

Allen, 1929: 686, Miss. (as *Z. blanda blanda*).

Epargyreus clarus (Cramer)

Most records ex *Epargyreus tityrus* (Fabricius).

Sellers, 1943: 20 (ex *Proteides*); Schaffner and Griswold, 1934: 15, 111; Sellers, 1930: 572-573.

Erynnis brizo (Boisduval and LeConte)

Most records ex *Thanaos brizo*.

Sellers, 1943: 19-20 (ex *Erynnis*); Sellers, 1930: 574-575; Essig, 1926: 581; Johnson, 1925: 197, Maine, Mass., N.H., R.I. (as *Z. blanda blanda*); Aldrich and Webber, 1924: 39 (as *Z. blanda blanda*); Johnson, 1910: 779, N.J. (as *Eusisyropa* ex *Nisoniades*); *Brauer and Bergenstamm, 1895: 553, 608, N. Amer. (as (*Exorista*), Williston, 1889); Williston, 1889: 1914, 1919 (as *Exorista blanda* var. *proserpina* Williston).

Limacodidae

Euclea cippus Cramer

Johnson, 1910: 779, N.J. (as *Eusisyropa*).

Euclea delphinii Boisduval

Essig, 1926: 581.

Lymantriidae

Euproctis chrysorrhoea (Linnaeus) probably

Townsend, 1908: 98, Mass. (as *Eusisyropa*).

Nygmia phaeorrhoea (Donovan)

Sellers, 1943: 20; Sellers, 1930: 572; Burgess and Crossman, 1929: 143.

Orgyia antiqua (Linnaeus)

All records ex *Notolophus antiqua*.

Sellers, 1943: 20; Schaffner and Griswold, 1934: 84, 111.

Stilpnotia salicis (Linnaeus)

*Sellers, 1943: 20 (Burgess and Crossman, 1927); Burgess and Crossman, 1927: 14, New England.

Noctuidae

Alabama argillacea (Hübner)

Essig, 1926: 581.

Catocala spp.

Sellers, 1943: 20; Sellers, 1930: 572.

Hypena humuli (Harris)

Essig, 1926: 581; Aldrich and Webber, 1924: 39 (as *Z. blanda blanda*).

Mocis spp. (including *M. latipes* (Guenée) and/or *M. repanda* (Fabricius))

Scaramuzza, 1946: 549, Cuba (ex *M. repanda*).

Phosphila turbulenta Hübner

Sellers, 1943: 20; Schaffner and Griswold, 1934: 52-53, 111
(ex *Trachea*); Sellers, 1930: 572-573 (ex *Hadena*).

Plathypena scabra (Fabricius)

Aldrich and Webber, 1924: 40, N.C. (as *Z. blanda blanda*);
Chittenden, 1901b: 50, D.C. (as *Exorista*).

Spodoptera frugiperda (J. E. Smith)

Most records ex *Laphygma frugiperda*.

Scaramuzza and Barry, 1960: 997, Cuba; Scaramuzza,
1951a: 299-300, Cuba; Scaramuzza, 1942: 148, Cuba;
Bruner, S. C., 1934: 96, Cuba; Van Dine and
Christenson, 1932: 2, Cuba; Scaramuzza, 1929: 122,
Cuba (ex *L. [frugiperda]*); Scaramuzza, 1929a: 113,
Cuba (ex *L. [frugiperda]*).

Unidentified Catocalinae

Schaffner and Griswold, 1934: 67, 111.

Nymphalidae

Nymphalis antiopa (Linnaeus)

Raizenne, 1952: 4, Ont. (as *Eusisyropa*).

Polygona interrogationis (Fabricius)

*Sellers, 1943: 20 (Schaffner and Griswold, 1934); Schaffner
and Griswold, 1934: 9-10, 111.

Vanessa atalanta (Linnaeus)

Sellers, 1943: 20; Schaffner and Griswold, 1934: 12, 111 (ex
Cynthia); Sellers, 1930: 572-573.

Vanessa cardui (Linnaeus)

Sellers, 1943: 19-20; Sellers, 1930: 575 (ex *Cynthia cardui*
"*J. rufipes*"); Essig, 1926: 581; Johnson, 1925: 197,
Maine, Mass., N.H., R.I. (as *Z. blanda blanda* ex
"*Vanassa*"); Aldrich and Webber, 1924: 39 (as *Z. blanda*
blanda); Johnson, 1910: 779, N.J. (as *Eusisyropa* ex
Pyrameis); *Brauer and Bergenstamm, 1895: 553, 609,
N. Amer. (as (*Exorista*), Williston, 1889); Williston,
1889: 1914, 1919 (as *Exorista*); Osten Sacken,
1887: 163 (as *Tachina* ex *Cynthia*).

Oecophoridae

Depressaria pastinacella (Duponchel)

All records ex *Depressaria heracliiana* (Linnaeus).

Sellers, 1943: 20; Sellers, 1930: 575, Maine.

Depressaria spp.

All records ex *Depressaria* sp. infesting cow parsnip.

*Procter, 1946: 432, Maine (Procter, 1938); Procter, 1938: 373, Maine; Johnson, 1927: 201, Maine.

Psilocorsis faginella (Chambers)

Embree, 1958: 170, N.S.

Psilocorsis fletcherella Gibson

Raizenne, 1952: 243, Ont. (as *Eusisyropa* ex "*Psilocorsis*").

Olethreutidae

Gretchena bolliana (Slingerland)

Sellers, 1943: 20; Aldrich and Webber, 1924: 40, Fla. (as *Z. blanda blanda* ex *Proteopteryx*).

Psychidae

Thyridopteryx ephemeraeformis (Haworth)

Kulman, 1965a: 864-865, W. Va.; Schaffner, 1959: 46, 54, [N.E. U.S.]; Sellers, 1943: 20; Schaffner and Griswold, 1934: 102-103, 111; Sellers, 1930: 572.

Pyralidae

Acrobasis comptoniella Hulst

Sellers, 1943: 20, N.H.; Essig, 1926: 581; Aldrich and Webber, 1924: 40, N.H. (as *Z. blanda blanda*).

Alatuncusia bergi (Möschler)

Aldrich and Webber, 1924: 40, Fla. (as *Z. blanda blanda* ex "*Dichogamma*").

Dichogama redtenbacheri Lederer

Aldrich and Webber, 1924: 40, Fla. (as *Z. blanda blanda* ex "*Dichogamma*").

Loxostege similalis (Guenée)

Essig, 1926: 581.

Nephopteryx subcaesiella (Clemens)

Schaffner, 1959: 17, 54, [N.E. U.S.] (ex *Salebria*).

Nephopteryx subfuscella (Ragonot)

Schaffner, 1959: 17, 54, [N.E. U.S.] (ex *Salebria semiobscura* Hulst).

Tetralopha robustella Zeller

Schaffner, 1959: 13, 54, [N.E. U.S.].

Sphingidae

Smerinthus jamaicensis (Drury)

All records ex *Smerinthus geminatus* Say.

*Sellers, 1943: 20 (Schaffner and Griswold, 1934);
Schaffner and Griswold, 1934: 20, 111.

Sphecodina abbottii Swainson

Sellers, 1943: 20; Schaffner and Griswold, 1934: 24, 111;
Sellers, 1930: 572 (ex *S. "abbottii"*).

Tortricidae

Archips argyrospilus (Walker)

*Sellers, 1943: 20 (Gill, 1913); Essig, 1926: 581, Colo.;
Aldrich and Webber, 1924: 40, Colo. (as *Z. blanda blanda* ex *Cacoecia*); Gill, 1913: 102, Colo. or N. Mex. (as *E[xorista]*).

Archips cerasivoranus (Fitch)

Schaffner, 1959: 30, 54, [N.E. U.S.]; Thompson, 1953: 19;
Thompson, 1953b: 393-400, figs. 2-11, Ont. (as *Eusisyropa*); Sellers, 1943: 20; Daviault, 1939: 184, [Que.] (ex *Cacoecia*); Sellers, 1930: 572; Aldrich and Webber, 1924: 39 (as *Z. blanda blanda* ex *Cacoecia*); Baird, 1918: 769, N.B. or probably Alta. (as *Exorista boarmiae*).

Archips fervidanus (Clemens)

Schaffner, 1959: 29, 54, [N.E. U.S.]; Sellers, 1943: 20;
Sellers, 1930: 572; Aldrich and Webber, 1924: 39 (as *Z. blanda blanda* ex *Cacoecia*).

Argyrotaenia mariana (Fernald)

Gilliatt, 1937: 145, N.S. (ex *Eulia*).

Unidentified Tortricidae

*Sellers, 1943: 20 (ex tortricid on poplar, Sellers, 1930);
Sellers, 1943: 20 (ex tortricid on sumac); Sellers, 1930:
572 (ex tortricid on poplar); Sellers, 1930: 572 (ex
tortricid on sumac).

Unidentified Lepidoptera

Miscellaneous

Sellers, 1943: 20 (ex larva on *Comptonia peregrina* (Linnaeus) Coulter (*Myrica asplenifolia* Linnaeus));
Aldrich, 1932: 2, Mass. (ex mixed spp. of Microlepidoptera on *Myrica asplenifolia* Linnaeus); Essig, 1926: 581.

Oakworm(-s)

Sellers, 1943: 20.

STAGE PARASITIZED: Larva, pupa.

IMMATURES: Egg — Thompson, 1953b: (descr., fig. 5).

First instar larva — Thompson, 1953b: (descr., fig. 3).

Second instar larva — Thompson, 1953b: (descr., fig. 4).

Third instar larva — Thompson, 1953b: (descr., figs. 2, 6-7, 9-10).

Pupa — Thompson, 1953b: (descr., figs. 8, 11); Sellers, 1930: (fig. 2).

BIOLOGY: Schaffner, 1959: (13, 17, 29-30, 46, 54); Sellers, 1943: (20); Schaffner and Griswold, 1934: (111).

Eusisyropa blanda (Osten Sacken) probably

All records as *Zenillia blanda* probably.

HOST: LEPIDOPTERA

Gelechiidae

Filatima monotaeniella (Bottimer)

*Sellers, 1943: 20 (Bottimer, 1926); Bottimer, 1926: 814, Tex. (ex *Gelechia*).

Noctuidae

Anomis erosa Hübner

*Sellers, 1943: 20 (Bottimer, 1926); Bottimer, 1926: 800, Tex.

STAGE PARASITIZED: Larva, pupa.

BIOLOGY: Bottimer, 1926: (814).

Eusisyropa boarmiae (Coquillett)

Most records as *Zenillia boarmiae*.

HOST: LEPIDOPTERA

Arctiidae

Hyphantria cunea (Drury)

Sellers, 1943: 22.

Hyphantria textor (Harris)

Sellers, 1943: 23.

Geometridae

Isturgia truncataria (Walker)

Chittenden, 1907: 26, Mass. (as *Exorista ex Cleora*).

Noctuidae

Alabama argillacea (Hübner)

Most records ex *Aletia argillacea*.

Sellers, 1943: 22 (ex *Alabama*); *vide Aldrich and Webber, 1924: 39, Miss. (as *Z. blanda blanda* (Osten Sacken), Townsend, 1908); Townsend, 1908: 98, Miss. (as *Eusisyropa*).

Hypena humuli (Harris)

Sellers, 1943: 22-23; *Hawley, 1918: 196 (as *Exorista hypenae* Coquillett, Howard, 1897); Howard, 1897: 47, Mass. (as *Exorista hypenae*).

Plathypena scabra (Fabricius)

Sellers, 1943: 22-23; Sherman, 1920: 298-299, N.C. (as *Exorista*).

Pyralidae

Acrobasis caryae Grote

Nickels, Pierce and Pinkney, 1950: 2, 19, Tex.

Acrobasis indigenella (Zeller)

Leiby, 1925: 9, N.C. (as *Exorista* ex *A. nebulella* Riley).

Acrobasis indigenella (Zeller) probably

*Sellers, 1943: 22 (ex *A. juglandis* (LeBaron), Brimley, 1922); Brimley, 1922: 22, N.C. (as *Exorista* ex *A. "nubilella"*).

Dichogama redtenbacheri Lederer

Sellers, 1943: 22-23.

Loxostege similalis (Guenée)

*Vide Sellers, 1943: 22-23 (Aldrich and Webber, 1924); *Aldrich and Webber, 1924: 40, Ark. (as *Z. blanda blanda*, Coquillett, 1897); Coquillett, 1897: 13, 26, Ark. (as *Exorista*).

Tortricidae

Archips fervidanus (Clemens)

Johnson, 1910: 780, N.J. (as *Eusisyropa* ex *Cacoecia*).

Yponomeutidae

Urodus parvulus (H. Edwards)

Patton, 1958: 85, Fla.

STAGE PARASITIZED: Larva, pupa.

IMMATURES: Pupa — Greene, 1922: (descr., fig. 25 on pl. 6).

BIOLOGY: Patton, 1958: (85).

Eusisyropa boarmiae (Coquillett) probably

Most records as *Zenillia boarmiae* probably.

HOST: LEPIDOPTERA

Noctuidae

Plathypena scabra (Fabricius)

All records ex green clover worm.

*Sellers, 1943: 22 (Brimley, 1922); Brimley, 1922: 22,
N.C. (as *Exorista boarmiae*).

Pyralidae

Acrobasis sp.

Sellers, 1943: 22.

STAGE PARASITIZED: Pupa.

Eusisyropa virilis (Aldrich and Webber)

Most records as *Zenillia virilis*.

HOST: LEPIDOPTERA

Arctiidae

Hyphantria cunea (Drury)

Sellers, 1943: 18; Swain, 1937: 245, Colo. (as *Z. blanda virilis*); Schaffner and Griswold, 1934: 37, 113; Sellers, 1930: 572-573.

Danaidae

Danaus plexippus plexippus (Linnaeus)

Sellers, 1943: 17 - 18 (ex *D. plexippus*).

Diptoridae

Phryganidia californica Packard

Harville, 1955: 115, 129, 132-133, Calif.

Geometridae

Anacamptodes ephyraria (Walker)

Sellers, 1943: 18; Schaffner and Griswold, 1934: 95, 113,
Mass.; Sellers, 1930: 573.

Cingilia catenaria (Cramer)

Sellers, 1943: 18; Schaffner and Griswold, 1934: 98, 113;
Sellers, 1930: 572-573.

Ennomos subsignarius (Hübner)

Sellers, 1943: 17-18; Aldrich and Webber, 1924: 40 (as *Z. blanda virilis*).

Melanolophia imitata (Walker)

Evans, 1962: 597, B.C. (as *Eusisyropa*).

Phigalia titea (Cramer)

Sellers, 1943: 18; Schaffner and Griswold, 1934: 97, 113;
Sellers, 1930: 573.

Hesperiidae

Epargyreus clarus (Cramer)

Most records ex *Epargyreus tityrus* (Fabricius).

Sellers, 1943: 18 (ex *Proteides clarus*); Schaffner and
Griswold, 1934: 15, 113; Sellers, 1930: 572-573.

Limaconidae

Euclea delphinii Boisduval

*Vide Sellers, 1943: 18 (Coquillett, 1897); *Aldrich and
Webber, 1924: 40, Va. (as *Z. blanda blanda* (Osten
Sacken), Coquillett, 1897); Coquillett, 1897: 13, 25, Va.
(as *Exorista blanda* ex *E. cippus* Cramer).

Lymantriidae

Porthetria dispar (Linnaeus)

Sellers, 1943: 17-18; Sellers, 1930: 573; Burgess and
Crossman, 1929: 143 (as *Zenillia blanda*); Howard and
Fiske, 1911: 90 (as *Exorista blanda*).

Noctuidae

Acronicta dactylina (Grote)

Most records ex *Acronycta dactylina*.

Sellers, 1943: 17 (ex *Acronicta*); Schaffner and Griswold,
1934: 58, 113; Sellers, 1930: 573.

Acronicta grisea Walker

Most records ex *Acronycta grisea*.

Sellers, 1943: 17 (ex *Acronicta*); Schaffner and Griswold,
1934: 56, 113, Maine; Sellers, 1930: 573.

Acronicta impressa Walker

Sellers, 1943: 17; Schaffner and Griswold, 1934: 58-59, 113
(ex *Acronycta*).

Acronicta leporina vulpina (Grote)

Sellers, 1943: 17-18; Schaffner and Griswold, 1934: 57, 113
(ex *Acronycta*).

Acronicta obliterata (J. E. Smith)

*Sellers, 1943: 18 (ex *A. "oblinita,"* Schaffner and
Griswold, 1934); Schaffner and Griswold, 1934: 60, 113
(ex *Acronycta*).

Acronicta spp. (including *A. leporina* (Linnaeus) and/or *A. lepusculina* Guenée)

Sellers, 1930: 573 (ex *Acronycta* spp. (including *A. leporina*
and/or *A. chionochroa* Hampson)).

Papaipema harrisii (Grote)

*Arnaud, 1963c: 116, Ill. (as *Eusisyropa*, Aldrich and
Webber, 1924); Sellers, 1943: 17-18; Aldrich and
Webber, 1924: 40, N.Y. (as *Z. blanda virilis*).

Phosphila turbulenta Hübner

Sellers, 1943: 18; Schaffner and Griswold, 1934: 52-53, 113
(ex *Trachea*); Sellers, 1930: 572-573 (ex *Hadena*).

Polia legitima (Grote)

Sellers, 1943: 18; Schaffner and Griswold, 1934: 45, 113;
Sellers, 1930: 573 (ex *Mamestra "legitima"*).

Simyra henrici (Grote)

Sellers, 1943: 17; Sellers, 1930: 573 (ex *Arsilonche*
albovenosa (Goeze)).

Notodontidae

Gluphisia septentrionalis Walker

Sellers, 1943: 18 (ex *G. "septentrionis"*); Schaffner and
Griswold, 1934: 83-84, 113; Sellers, 1930: 573.

Ichthyura albosigma (Fitch)

Most records ex *Melalopha albosigma*.

Sellers, 1943: 18 (ex *Ichthyura*); Schaffner and Griswold,
1934: 73-74, 113; Sellers, 1930: 573.

Ichthyura strigosa Grote

Most records ex *Melalopha strigosa*.

Sellers, 1943: 18 (ex *Ichthyura*); Schaffner and Griswold,
1934: 73, 113; Sellers, 1930: 573.

Nymphalidae

Polygonia interrogationis (Fabricius)

*Sellers, 1943: 18 (Schaffner and Griswold, 1934); Schaffner and Griswold, 1934: 9-10, 113.

Vanessa atalanta (Linnaeus)

Sellers, 1943: 18; Schaffner and Griswold, 1934: 12, 113 (ex *Cynthia*); Sellers, 1930: 572-573.

Pieridae

Eucheira socialis Westwood

Shaw, 1947: 436-437, Mexico.

Psychidae

Thyridopteryx ephemeraeformis (Haworth)

Kulman, 1965a: 864-865, W. Va. (as *Eusisyropa*).

Saturniidae

Callosamia promethea (Drury)

Sellers, 1943: 18; Schaffner and Griswold, 1934: 27, 113; Sellers, 1930: 573.

Platysamia cecropia (Linnaeus)

*Sellers, 1943: 18 (Schaffner and Griswold, 1934); Schaffner and Griswold, 1934: 26-27, 113.

Sphingidae

Ceratomia catalpae (Boisduval)

Sellers, 1943: 18; Schaffner and Griswold, 1934: 17, 113, N.J.; Sellers, 1930: 573.

STAGE PARASITIZED: Larva.

IMMATURES: Pupa — Sellers, 1930: (fig. 1).

BIOLOGY: Evans, 1962: (597); Sellers, 1943: (18); Schaffner and Griswold, 1934: (113).

Euthera Loew (Phasiinae: Eutherini)

Euthera tentatrix Loew

HOST: HEMIPTERA (HETEROPTERA)

Pentatomidae

Euschistus ictericus (Linnaeus)

Sabrosky, 1965: 61, Fla.

Euschistus servus (Say)

*Sabrosky, 1965: 61, Fla., Ohio (Patton, 1958); Patton, 1958: 37, Fla.

Euschistus tristigmus (Say)

Sabrosky, 1965: 61, Ohio; Sabrosky, 1965: 61, Ohio (ex *E. tristigmus* var. *luridus* Dallas).

Euschistus variolarius (Palisot de Beauvois)

Sabrosky, 1965: 61, Ohio; Rings and Brooks, 1958: 15-16, Ohio.

Oebalus pugnax (Fabricius)

Sabrosky, 1965: 61, Fla., La.

Pentatomid spp. (including *Euschistus ictericus* (Linnaeus), *Mormidea pictiventris* Stål, *Nezara viridula* (Linnaeus) and/or *Oebalus pugnax* (Fabricius))

Thames, 1954: 236, Fla. (ex pentatomids, *Solubea pugnax*, *E. ictericus*, *M. pictiventris* and/or "Nezera" *viridula*).

STAGE PARASITIZED: Adult.

Eutheresia Townsend (Proseninae: Theresiini)

Eutheresia canescens, see *Ptilodexia canescens*

Eutheresia monohammi Townsend

HOST: COLEOPTERA

Cerambycidae

Monochamus notatus (Drury)

Soper and Olson, 1963: 89-91, Maine; Townsend, 1912: 117, Ga. (ex "*Monohammus*" *confusor* Kirby).

Monochamus scutellatus (Say)

Soper and Olson, 1963: 89, Maine; Aldrich, 1932: 4, Mass.

Monochamus titillator (Fabricius)

Fattig, 1949: 16-17, Ga.; Savely, 1939: 380, N.C. (as *Theresia*).

Monochamus spp.

Soper and Olson, 1963: 89, Maine.

Lucanidae

Ceruchus piceus (Weber)

Hoffmann, 1937: 281, 283, Minn. (as *Theresia*).

STAGE PARASITIZED: Larva, adult probably.

IMMATURES: Egg — Townsend, 1912: (descr.).

First instar larva — Townsend, 1912: (descr.).

Third instar larva — Hoffmann, 1937: (descr.).

Pupa — Hoffmann, 1937: (descr.).

BIOLOGY: Soper and Olson, 1963: (89-91).

Eutheresia monohammi Townsend probably

HOST: COLEOPTERA

Cerambycidae

Unidentified Cerambycidae

Townsend, 1912: 117 (ex cerambycid in chestnut).

Lucanidae

Ceruchus piceus (Weber)

Aldrich, 1932: 3-4, Mass.

STAGE PARASITIZED: Larva.

Eutheresia nipigonensis (Curran)

HOST: COLEOPTERA

Cerambycidae

Stenocorus inquisitor inquisitor (Linnaeus)

Thomas, J. B., 1955: 340, 343, Ont. (ex *S. inquisitor*).

STAGE PARASITIZED: Larva, pupa.

Eutheresia satisfacta West probably

HOST: COLEOPTERA

Cerambycidae

Stenocorus inquisitor inquisitor (Linnaeus)

Soper and Olson, 1963: 89-90, Maine (ex *Rhagium inquisitor*).

STAGE PARASITIZED: Larva.

Eutheresia trivittata Curran

HOST: COLEOPTERA

Cerambycidae

Monochamus notatus (Drury)

Thomas, J. B., 1955: 340, Ont.

Monochamus scutellatus (Say)

Thomas, J. B., 1955: 340, 343, Ont.

Stenocorus inquisitor inquisitor (Linnaeus)

Thomas, J. B., 1955: 343, Ont. (ex *S. inquisitor*).

Eutritochaeta carpocapsae, see *Nilea carpocapsae*

Eutrixa Coquillett (Proseninae: Aulacephalini)

Eutrixia "exile," see *Eutrixia exilis*

Eutrixia exilis (Coquillett)

Most records as *Eutrixia* "exile."

HOST: COLEOPTERA

Scarabaeidae

Eutheola rugiceps (LeConte)

Fattig, 1949: 17, Ga. (as *E. exilis* ex "*Eutheola*").

Ligyris gibbosus (De Geer)

Fattig, 1949: 17, Ga. (as *E. exilis*).

Phyllophaga anxia (LeConte)

Melvin and Nairn, 1967: 43, Man. (as *E. exilis*); Melvin and Nairn, 1967a: 27, Man. (as *E. exilis*).

Phyllophaga drakii (Kirby)

Melvin and Nairn, 1967: 43, Man. (as *E. exilis*); Melvin and Nairn, 1967a: 27, Man. (as *E. exilis*).

Phyllophaga fervida (Fabricius)

All records ex *Lachnosterna arcuata* Smith.

Shannon, 1914: 182; Chittenden, 1899: 79, D.C. (as *E. masuria* (Walker)); Chittenden, 1898a: 76, D.C. (as *E. masuria*); Coquillett, 1897: 12, 23, D.C. (as *E. masuria*).

Phyllophaga nitida (LeConte)

Melvin and Nairn, 1967: 43, Man. (as *E. exilis*); Melvin and Nairn, 1967a: 27, Man. (as *E. exilis*).

Phyllophaga rubiginosa (LeConte)

Hayes, 1920: 317, Kans. (as "*Eutrixia*" ex *Lachnosterna*).

Phyllophaga spp.

Most records ex *Lachnosterna* spp.

Davis, J. J., 1918: 14; Smyth, 1917: 59 (ex *Phyllophaga*); Davis, J. J., 1916: 271, N.E. U.S.; Walton, 1913: 198, N. Amer.

June beetle(-s)

Hammond, 1942: 147, Ont. (as *Eutrixia* sp.); Hammond, 1941a: 187, Ont.

May beetle(-s)

*Procter, 1946: 431, Maine (Procter, 1938); Procter, 1938: 372, Maine; Johnson, 1927: 199, Maine; Johnson, 1925: 191, Conn., Maine, Mass., N.H.

STAGE PARASITIZED: Adult.

IMMATURES: Pupa — Greene, 1922: (descr., fig. 51 on pl. 11).

BIOLOGY: Melvin and Nairn, 1967: (43); Melvin and Nairn, 1967a: (27); Hammond, 1941a: (187).

Eutrixa masuria, see *Eutrixa exilis*

Eutrixa sp., see *Eutrixa exilis*

"*Eutrixia exile*," see *Eutrixa exilis*

Eutrixoides Walton (Proseninae: Dexillini)

Eutrixoides jonesii Walton

HOST: COLEOPTERA

Scarabaeidae

Chemarachis vandinei (Smyth)

Most records ex *Phyllophaga vandinei*.

Martorell, 1945a: 432, P.R.; Wolcott, 1924: 221, P.R. (ex *Lachnosterna*); Smyth, 1917: 82, 86-87, P.R.

Phyllophaga portoricensis Smyth

Wolcott, 1924: 221, P.R. (ex *Lachnosterna*); Smyth, 1917: 86, P.R.

Phyllophaga spp.

Most records ex *Lachnosterna* spp.

Wolcott, 1935: 449, P.R.; Wolcott, 1924a: 27, P.R.;
Wolcott, 1922: 15-16, P.R.; Smyth, 1917a: 151, P.R.
(ex *Phyllophaga*); Van Zwaluwenburg, 1917: 517, P.R.
(ex *Phyllophaga*); Van Dine, 1913a: 37, P.R.; Van
Dine, 1913b: 200, P.R.; Walton, 1913d: 49, 51, P.R.

Scarabaeid spp. (including *Phyllophaga* spp. or *Phytalis* spp.)

Smyth, 1917: 55-56, P.R. (ex white grubs).

STAGE PARASITIZED: Adult.

BIOLOGY: Wolcott, 1935: (449); Smyth, 1917: (82, 86-87).

Euzenillia variabilis, see *Lixophaga variabilis*

Euzenilliopsis "diatraea," see *Lixophaga diatraeae*

Euzenilliopsis diatraeae, see *Lixophaga diatraeae*

Euzenilliopsis diatraeae probably, see *Lixophaga diatraeae* probably

Exorista Meigen (Goniinae: Exoristini)

Exorista "achracea," see *Chrysoexorista ochracea*

Exorista affinis, see *Hubneria estigmenensis*

Exorista amplexa, see *Carcelia amplexa*

Exorista arvicola, see *Nemorilla pyste*

Exorista blanda, see *Anaporia pristis*, *Eusisyropa blanda* or
Eusisyropa virilis

Exorista boarmiae, see *Eusisyropa blanda* or *Eusisyropa boarmiae*

Exorista caesar, see *Aplomya caesar*

Exorista cecropia, see *Winthemia cecropia*

Exorista "cecropiae," see *Winthemia cecropia*

Exorista ceratomiae, see *Hyphantrophaga hyphantriae*

Exorista "chelonia," see *Carcelia laxifrons*

- Exorista cheloniae*, see *Carcelia laxifrons*, *Carcelia malacosomae*
 probably or *Carcelia reclinata* probably
Exorista chrysophani, see *Aplomya theclarum*
Exorista ciliata, see *Winthemia rufopicta*
Exorista confinis, see *Aplomya theclarum*
Exorista datanae, see *Winthemia datanae*
Exorista eudryae, see *Carcelia formosa*, *Carcelia reclinata* probably,
Sisyropa alypiae probably or *Sisyropa eudryae*
Exorista fasciata fasciata (Fallén)
 HOST: LEPIDOPTERA

Lymantriidae

Byrdia groenlandica Wocke

Henriksen and Lundbeck, 1917: 622, E. Greenland (as
Tachina fasciata ex *Dasychira*).

STAGE PARASITIZED: Larva.

- Exorista fernaldi*, see *Exorista mella*
Exorista flavicauda, see *Triachora unifasciata*
Exorista flavirostris, see *Carcelia lagoae*
Exorista futilis, see *Euexorista futilis*
Exorista griseomicans, see *Carcelia amplexa*
Exorista helvina, see *Aplomya helvina*
Exorista hirsuta, see *Phryxe vulgaris*
Exorista hypenae, see *Eusisyropa boarmiae*
Exorista infesta, see *Winthemia infesta*
Exorista isae, see *Anaporia pristis*
Exorista lagoae, see *Carcelia lagoae*
Exorista larvarum, see *Exorista mella* or *Exorista* spp.
Exorista larvarum probably, see *Exorista mella* probably
Exorista leucanae, see *Winthemia leucanae*
Exorista leucanae lacking red at tip, see *Winthemia leucanae* lacking
 red at tip
Exorista "leucania," see *Winthemia leucanae*
Exorista "leucania" probably, see *Winthemia leucanae* probably
Exorista "leucaniae," see *Winthemia leucanae*
Exorista "leucaniae" lacking red at tip, see *Winthemia leucanae*
 lacking red at tip
Exorista "leucaniae var. cecropiae," see *Winthemia cecropia*
 "Exorista" *lobeliae* Coquillett
 Most records as *Zenillia lobeliae*.
 HOST: LEPIDOPTERA

Lymantriidae

Hemerocampa leucostigma (J. E. Smith)

Coquillett, 1897: 14, 26, D.C. (as *Exorista* ex *Orgyia*).

Noctuidae

Acronicta sp. near *afflicta* Grote

Patton, 1958: 85, Fla.

Acronicta grisea Walker

Schaffner and Griswold, 1934: 56, 112, Maine (ex *Acronycta*).

Acronicta hamamelis Guenée

All records ex *Acronycta hamamelis*.

Aldrich and Webber, 1924: 20, Va.; Coquillett, 1897: 14, 23, Va. (as *Exorista*).

Acronicta lobeliae Guenée

All records ex *Acronycta lobeliae*.

Aldrich and Webber, 1924: 20; Bischoff, 1909: 76 (as *Exorista*); Coquillett, 1897: 14, 23 (as *Exorista*).

Acronicta spp.

All records ex *Acronycta* sp.

Aldrich and Webber, 1924: 20, Va.; Aldrich and Webber, 1924: 20, Md. (ex *Acronycta* sp. on oak); Coquillett, 1897: 14, 23, Md. (as *Exorista*).

Alabama argillacea (Hübner)

Aldrich and Webber, 1924: 20, Tex.; Reinhard, 1922: 72, Tex. (as *Exorista*).

STAGE PARASITIZED: Larva, pupa.

IMMATURES: Pupa — Greene, 1922: (descr., fig. 18 on pl. 4).

BIOLOGY: Patton, 1958: (85); Schaffner and Griswold, 1934: (56, 112); Bischoff, 1909: (76).

"*Exorista*" sp. near *lobeliae* Coquillett

HOST: LEPIDOPTERA

Noctuidae

Acronicta betulae Riley

Schaffner and Griswold, 1934: 58, 115, N.J. (as *Phorocera* near *Zenillia* ex *Acronycta*).

BIOLOGY: Schaffner and Griswold, 1934: (58, 115).

Exorista loxostegeae, see *Pseudoperichaeta erecta*

Exorista "melba," see *Exorista mella*

Exorista mella (Walker)

HOST: HYMENOPTERA

Tenthredinidae

Macremphytus varianus (Norton)

Greene, 1921: 41, Md. (as *Tachina*).

HOST: LEPIDOPTERA

Arctiidae

Apantesis nevadensis superba (Stretch)

Essig, 1926: 583, Colo. (as *Tachina larvarum* (Linnaeus) ex *A. blakei superba*).

Apantesis oithona Strecker

Allen, 1929: 689, Miss. (as *E. larvarum*); Reinhard, 1922: 73, Tex. (as *Tachina* ex *A. rectilinea* French).

Apantesis phyllira (Drury)

Patton, 1958: 38, Fla.; Coquillett, 1897: 21, 24, Fla. (as *Tachina* ex *Arctia*).

Cycnia tenera Hübner

Schaffner and Griswold, 1934: 32, 120.

Diacrisia virginica (Fabricius)

Schaffner and Griswold, 1934: 36, 120.

Estigmene acraea (Drury)

Bryan, D. E., Jackson and Stoner, 1969: 1, 7, Ariz. (lab.);
Butler, Bryan and Jackson, 1968: 161-162, Ariz. (lab.);
Bosch, and Hagen, 1966: 25, Calif. (as *E. larvarum*);
Bibby, 1961: 327, Ariz. (as *E. larvarum* ex salt marsh caterpillar); Patton, 1958: 38, Fla.; Stevenson, W. A., Kaufman and Sheets, 1957: 280, Ariz. (as *E. larvarum*); Taylor, E. A., 1954: 526-529, Ariz. (as *E. larvarum*); Schaffner and Griswold, 1934: 38, 120; Allen, 1929: 689, Miss. (as *E. larvarum* ex "*Estigeme*" *acraea*); Essig, 1926: 583, Calif. (as *Tachina larvarum* ex *E. acraea*); Reinhard, 1922: 73, Tex. (as *Tachina* ex *E. acraea*); Johnson, 1910: 781, N.J. (as *Tachina* ex *Leucarctia acraea*); Coquillett, 1897: 21, 26, Kans. (as *Tachina* ex *Leucarctia acraea*).

Euchaetias egle (Drury)

Schaffner and Griswold, 1934: 39, 120.

Halisidota harrisii Walsh

Schaffner and Griswold, 1934: 34-35, 120.

Halisidota tessellaris (J. E. Smith)

Schaffner and Griswold, 1934: 34, 120.

Hyphantria textor (Harris)

Raizenne, 1952: 29, Ont.

Isia isabella (J. E. Smith)

Schaffner and Griswold, 1934: 36-37, 120; Essig, 1926: 583, Calif. (as *Tachina larvarum*); Johnson, 1910: 781, N.J. (as *Tachina ex Pyrrharctia*); Coquillett, 1897: 21, 27, Ont. (as *Tachina ex Pyrrharctia*).

Bombycidae

Unidentified Bombycidae

Howard, 1897b: 42-43, D.C. (as *Tachina*).

Danaidae

Danaus plexippus plexippus (Linnaeus)

Essig, 1926: 583, Colo. (as *Tachina larvarum ex D. menippe* (Hübner)); Townsend, 1895: 73, Colo. (as *Tachina clisiocampae* Townsend ex "*Danais*" *archippus* (Fabricius)).

Lasiocampidae

Epicnaptera americana (Harris)

Raizenne, 1952: 104, Ont.; Schaffner and Griswold, 1934: 89, 120.

Malacosoma americanum (Fabricius)

Stehr and Cook, 1968: 258, 299, Tex., Wis.; Fattig, 1949: 17, Ga. (as *E. larvarum*); Bess, 1936: 605 (as "*Exoristia*"); Schaffner and Griswold, 1934: 88, 120; West, 1928: 817, N.Y. (as *E. larvarum*).

Malacosoma californicum (Packard)

Stehr and Cook, 1968: 263, 265, 267-268, 270, 300, Colo., Idaho, Oreg., Wash., Wyo.; Essig, 1926: 583, Calif., Colo. (as *Tachina larvarum*); Townsend, 1895: 73, Colo. (as *Tachina clisiocampae ex Clisiocampa*, questionable host).

Malacosoma californicum californicum (Packard)

Coquillett, 1897: 21, 24, Calif. (as *Tachina ex Clisiocampa thoracica* Stretch).

Malacosoma californicum fragile (Stretch)

Essig, 1926: 583, Colo. (as *Tachina larvarum ex M. fragilis*).

Malacosoma californicum lutescens (Neumoegen and Dyar)

Stehr and Cook, 1968: 275, 301, Alta.

Malacosoma californicum pluviale (Dyar)

Stehr and Cook, 1968: 274-275, 300, Alta., B.C., Mont.

Malacosoma constrictum (H. Edwards)

Essig, 1926: 583, Calif. (as *Tachina larvarum*).

Malacosoma disstria Hübner

Sippell, 1961: 434, 436, fig. 3, Ont.; Raizenne, 1952: 103, Ont.; Strickland, 1938: 218, Alta. (ex forest tent caterpillar); Bess, 1936: 605 (as "*Exoristia*"); Knowlton, 1936: 242, Utah (as *E. larvarum* ex forest tent caterpillar); Schaffner and Griswold, 1934: 89, 120; Essig, 1926: 583, Calif. (as *Tachina larvarum*); Lochhead, 1919: 204 (as *Tachina*); Essig, 1915: 331, 412, Calif. (as *Tachina*); Essig, 1913: 259, Calif. (as *Tachina*); Felt, 1905: 436, N.Y. (as *Tachina*); Felt, 1902: 861-862, N.Y. (as *Tachina ex Clisiocampa*); Felt, 1910: 997, N.Y. (as *Tachina ex Clisiocampa*); *Brauer and Bergenstamm, 1895: 575, 588 (as (*Tachina clisiocampae* ex *Clisiocampa sylvatica* (Harris), Townsend, 1891a); Townsend, 1891a: 84, Maine (as *Tachina clisiocampae* ex *Clisiocampa sylvatica*).

Malacosoma incurvum incurvum (H. Edwards)

Stehr and Cook, 1968: 279, 301, Ariz.

Malacosoma spp.

Ross, 1953: 19, 22, fig. 6, B.C.; Essig, 1915: 331, Calif. (as *Tachina*); Essig, 1913: 259, Calif. (as *Tachina*); Fletcher, 1905: 78, Alta. (as *Tachina ex Clisiocampa*); Coquillett, 1897: 21, 24, Calif., Colo., Wyo. (as *Tachina ex Clisiocampa*).

Tolyte laricis (Fitch)

Raizenne, 1952: 100, Ont.

Tolyte velleda (Stoll)

Raizenne, 1952: 99, Que.

Apple tree tent caterpillar(-s)

Brimley, 1922: 26, N.C. (as *Tachina*).

Tent caterpillar(-s)

Brimley, 1938: 365, N.C. (as *E. larvarum*); Hitchings, 1908: 11, Maine (as *Tachina*).

Lymantriidae

Dasychira basiflava (Packard)

Schaffner and Griswold, 1934: 86, 120 (ex *Olene*).

Dasychira vagans (Barnes and McDunnough)

Brittain and Payne, 1919: 68, N.S. (as *Tachina ex Olene*).

Euproctis chrysorrhoea (Linnaeus)

Howard and Fiske, 1911: 93, 227 (as *Tachina*).

Hemerocampa leucostigma (J. E. Smith)

Raizenne, 1952: 95, Ont.; Dowden, 1935: 509 (as *Tachina*); Schaffner and Griswold, 1934: 85, 120; Essig, 1926: 583, Calif. (as *Tachina larvarum* ex *H. oslari* (Barnes)); Lochhead, 1919: 203 (as *Tachina*); Payne, 1918: 67, [N.S.] (as *Tachina*); Britton, 1917: 110 (as *Tachina*); Britton, 1917: 110 (as *Tachina clisiocampae*); Johnson, 1910: 781, N.J. (as *Tachina* ex *Orgyia*); Schoene, 1909: 47, N.Y. (as *Tachina clisiocampae* ex "*Hemerocampe*"); Felt, 1905: 436, N.Y. (as *Tachina* ex *Notolophus*); Titus and Pratt, 1904: 28 (as *Tachina*); Felt, 1902: 864, N.Y. (as *Tachina* ex *Notolophus*); Howard, 1898: 88, D.C. (as *Tachina* ex *Orgyia*); Coquillett, 1897: 21, 26, D.C. (as *Tachina* ex *Orgyia*); Howard, 1897a: 16-17, D.C. (as *Tachina* ex *Orgyia*); Howard, 1897b: 7, 42-43, 49, D.C. (as *Tachina* ex *Orgyia*); *Brauer and Bergenstamm, 1895: 576, 601 (as (*Tachina*) *orgyiae* Townsend ex *Orgyia*, Townsend, 1892a); Hopkins, A.D., 1893a: 238, W. Va. (as *Tachina* "*orgyia*" ex *Orgyia* "*leugcostigma*"); Townsend, 1892a: 284-285, W. Va. (as *Tachina orgyiae* ex *Orgyia*); LeBaron, 1871: 16, Ill. (as *Tachina orgyiae* LeBaron ex *Orgyia*).

Hemerocampa pseudotsugata McDunnough

Davis and Knowlton, 1963: 1399, Utah.

Hemerocampa vetusta (Boisduval)

Essig, 1926: 583, Calif. (as *Tachina larvarum*); Essig, 1915: 331, 410, Calif. (as *Tachina*); Essig, 1913: 180, 259, Calif. (as *Tachina*).

Nygmia phaeorrhoea (Donovan)

Proper, 1934: 374, New England (as *Tachina* ex brown-tail moth); Burgess and Crossman, 1929: 143 (as *Tachina*).

Orgyia antiqua (Linnaeus)

All records ex *Notolophus antiqua*.

Raizenne, 1952: 93, Ont.; Schaffner and Griswold, 1934: 84, 120; Tothill, 1913: 74, N.S. (as *Tachina*).

Porthetria dispar (Linnaeus)

Burgess and Crossman, 1929: 143 (as *Tachina*); Burgess, 1924: 35, Mass. (as *Tachina*); Britton, 1915: 17, Conn. (as *Tachina*); Britton, 1911: 680, Conn. (as *Tachina* ex gypsy moth); Howard and Fiske, 1911: 90, 225, 227 (as *Tachina*); Howard and Fiske, 1911: 90 (as *E. fernaldi* (Williston)).

Stilpnotia salicis (Linnaeus)

Lejeune and Silver, 1961: 459-460, 462, 466, B.C.; Reeks and Smith, 1956: 569, maritime prov.; Beaulne, 1949: 206, Que.; vide Burgess and Crossman, 1927: 14-15, New England (as *Tachina*, Glendenning, 1924); Glendenning, 1924: 13, B.C. (as *Tachina*).

Tussock moth(-s)

Howard and Fiske, 1911: 223, Mass. (as *Tachina*); Thompson, 1910: 291 (as *Tachina*, incomplete rearing — pupa).

Noctuidae

Acronicta americana (Harris)

Schaffner and Griswold, 1934: 57-58, 120 (ex *Acronycta*).

Acronicta lepusculina Guenée

Coquillett, 1897: 21, 23 (as *Tachina* ex *Acronycta populi* Riley).

Acronicta rubricoma Guenée

Schaffner and Griswold, 1934: 59, 120, N.J. (ex *Acronycta*).

Acronicta sp.

Schaffner and Griswold, 1934: 60, 120 (ex *Acronycta*).

Pseudaletia unipuncta (Haworth)

Guppy, J. C., 1967: 97, 99, 102, Ont. (as *E. larvarum*); Pond, 1960: 655, N.B. (as *E. "melba"*).

Pyreferra hesperidago (Guenée)

Schaffner and Griswold, 1934: 51, 120 (ex *Conistra indirecta* Walker).

Simyra henrici (Grote)

Schaffner and Griswold, 1934: 61, 120; Cole, 1931: 9, 38, Idaho (as *E. larvarum* ex *Arsilonche albovenosa* (Goeze)).

Notodontidae

Datana integerrima Grote and Robinson

Schaffner and Griswold, 1934: 76, 120.

Datana ministra (Drury)

Schaffner and Griswold, 1934: 74, 120; Essig, 1926: 583, B.C. (as *Tachina larvarum*); Tothill, 1913: 74, B.C. (as *Tachina*).

Ichthyura inclusa Hübner

Schaffner and Griswold, 1934: 73, 120 (ex *Melalopha*).

Schizura concinna (J. E. Smith)

Schaffner and Griswold, 1934: 81, 120.

Nymphalidae

Nymphalis antiopa (Linnaeus)

Schaffner and Griswold, 1934: 11, 120 (ex *Hamadryas*).

Polygonia interrogationis (Fabricius)

Schaffner and Griswold, 1934: 9-10, 120.

Saturniidae

Callosamia promethea (Drury)

Schaffner and Griswold, 1934: 27, 120.

Hemileuca oliviae Cockerell

Wildermuth and Frankenfeld, 1933: 797, [N. Mex.] (as *Tachina*); Essig, 1926: 583, N. Mex. (as *Tachina larvarum*); Wildermuth and Caffrey, 1916: 8, N. Mex. (as *Tachina*); *Packard, 1914: 269 (as *Tachina*, Ainslie, 1910); Ainslie, 1910: 88-91, N. Mex. (as *Tachina*).

STAGE PARASITIZED: Larva (Lepidoptera), pupa (Lepidoptera).

IMMATURES: Third instar larva — LeBaron, 1871: (descr.).

Pupa - Sippell, 1961: (descr., fig. 3); Ross, 1953: (descr., fig. 6); Greene, 1922: (descr., fig. 72 on pl. 16); LeBaron, 1871: (descr.).

BIOLOGY: Bryan, D. E., Jackson and Stoner, 1969: (1, 7); Butler, Bryan and Jackson, 1968: (161-162); Lejeune and Silver, 1961: (459-460, 462, 466); Patton, 1958: (38); Taylor, E. A., 1954: (526-529); Dowden, 1935: (509); Schaffner and Griswold, 1934: (120); Burgess, 1924: (35); Glendenning, 1924: (13); Britton, 1915: (17); Britton, 1911: (680); Ainslie, 1910: (88-91); Thompson, 1910: (291).

Exorista mella (Walker) probably

HOST: LEPIDOPTERA

Arctiidae

Estigmene acrea (Drury)

Carrillo S., Ortega C., and Gibson, 1966: 94, Mexico (Son.)
(as *E. larvarum* (Linnaeus) probably ex *E. acrea*).

Exorista militaris, see *Winthemia militaris*

Exorista nigripalpis, see *Aplomya caesar* or *Aplomya confusionis*

Exorista ochracea, see *Chrysoexorista ochracea*

Exorista ostensackenii, see *Winthemia ostensackenii*

Exorista petiolata, see *Diplostichus lophyri*

- Exorista phycitae*, see *Nemorilla pyste*
Exorista "piste," see *Nemorilla pyste*
Exorista platysamiae, see *Winthemia cecropia*
Exorista pyste, see *Nemorilla pyste*
Exorista sp. near *pyste*, see *Nemorilla* sp. near *pyste*
Exorista scudderi, see *Nemorilla pyste*
Exorista simulans, see *Guerinia dydas*
Exorista theclarum, see *Aplomya theclarum*
Exorista trivittata probably, see "*Nemorilla trivittata*" probably
Exorista vulgaris, see *Phryxe vulgaris*
Exorista spp.

HOST: LEPIDOPTERA

Arctiidae

Diacrisia virginica (Fabricius)

Aldrich, 1932: 3, Mass. (as *E. larvarum* (Linnaeus)).

Estigmene acrea (Drury)

Anon., 1954h: 47, Ariz.

Euchaetias egle (Drury)

Webber and Schaffner, 1926: 20, New England (as *Tachina mella* Walker).

Danaidae

Danaus plexippus plexippus (Linnaeus)

Urquhart, 1960: 207, Ont. (ex *D. plexippus*).

Geometridae

Biston cognataria (Guenée)

Aldrich, 1932: 1, Mass. (as *E. larvarum* ex *Amphidasis*, incomplete rearing — pupa).

Lasiocampidae

Malacosoma disstria Hübner

Johnson, 1925: 204, Conn., Maine, Mass., N.H., R.I., Vt. (as *E. larvarum* ex forest tent caterpillar); Tothill, 1923: 180, Canada; Tothill, 1919: 12, N.B.

Lymantriidae

Dasychira basiflava (Packard)

Felt and Bromley, 1941: 384, Conn. (as *E. larvarum* ex *Olene*).

Euproctis chrysorrhoea (Linnaeus)

Howard and Fiske, 1911: 225, 309, pl. 20 (as *Tachina larvarum*).

Hemerocampa leucostigma (J. E. Smith)

Most records ex *Orgyia leucostigma*.

Webber and Schaffner, 1926: 12-13, 20, Mass., New England, N.Y., Pa., R.I. (as *Tachina mella* ex *Hemerocampa*); Howard, 1898: 88, D.C.; Howard, 1897a: 17, D.C.

Nygmia phaeorrhoea (Donovan)

Dowden, 1962: 27, 29, 58, New England (as *E. larvarum*); Clausen, 1956: 6, 105-106, N.H. (as *E. larvarum*).

Porthetria dispar (Linnaeus)

Dowden, 1962: 30, 32, 35, 58, New England (as *E. larvarum*); Dowden, 1961: 875, Mass. (as *E. larvarum*); Clausen, 1956: 6-7, 63, 105-106, Conn., Mass. (as *E. larvarum*); Brown and Sheals, 1944: 400 (as *E. larvarum*); Britton, 1935: 636, Conn. (as *E. larvarum*); Johnson, 1925: 204, Conn., Maine, Mass., N.H., R.I., Vt. (as *E. larvarum* ex "gipsy" moth); Howard and Fiske, 1911: 225-227, 308, pl. 20 (as *Tachina larvarum*).

Tussock moth(-s)

Johnson, 1925: 204, Conn., Maine, Mass., N.H., R.I., Vt. (as *E. larvarum*).

Noctuidae

Acronicta americana (Harris)

Webber and Schaffner, 1926: 20-21, New England (as *Tachina mella* ex *Apatela*).

Acronicta lepusculina Guenée

Brauer and Bergenstamm, 1895: 579-580 (as *Tricholyga* ex *Acronycta populi* Riley).

Papaipema cataphracta (Grote)

Washburn, 1910a: 94, Minn.

Papaipema nebris (Guenée)

All records ex *Papaipema nitela* (Guenée).

Franklin, 1910: 94, Minn.; Washburn, 1910: 168; Washburn, 1910a: 94, Minn.

Notodontidae

Cerura spp.Aldrich, 1932: 3, Mass. (as *E. larvarum*).

Olethreutidae and/or Pyralidae

Olethreutid and/or pyralid spp. (including *Laspeyresia* spp.
and/or *Dioryctria* spp. probably)

Keen, 1958: 70.

Pyralidae

Acrobasis indigenella (Zeller)Gill, 1917a: 21 (ex *A. nebulella* Riley).*Desmia funeralis* (Hübner)

Lochhead, 1919: 209.

Tortricidae

Archips fervidanus (Clemens)Schaffner, 1959: 29, 50, N.E. U.S. (as *E. larvarum*).

STAGE PARASITIZED: Larva, pupa.

IMMATURES: Pupa — Greene, 1922: (descr., fig. 88 on pl. 19);

Howard and Fiske, 1911: (pl. 20).

BIOLOGY: Schaffner, 1959: (29, 50); Clausen, 1956: (6-7, 63,
105-106); Britton, 1935: (636).*Exorista* spp. probablyMost records as *Tricholyga* sp. probably.

HOST: LEPIDOPTERA

Notodontidae

Datana contracta Walker

Brauer and Bergenstamm, 1895: 579, 590.

Datana ministra (Drury)

Brauer and Bergenstamm, 1895: 579, 590.

Unidentified Lepidoptera

Miscellaneous

Townsend, 1893c: 315, Jamaica (as *Exorista* ex moth allied
to "*Euchaetes*," but with an *Orgyia*-like larva).

STAGE PARASITIZED: Larva, pupa.

“Exoriste” *pyste*, see *Nemorilla pyste*

“Exoristia” *mella*, see *Exorista mella*

Exoristoides Coquillett (Tachininae: Germariini)

Exoristoides harrisi Reinhard

HOST: GEOPHILOMORPHA (CHILOPODA)

Geophilidae

Geophilus sp. probably

Reinhard, 1935: 161, Iowa.

Exoristoides johnsoni Coquillett

HOST: ORTHOPTERA

Gryllidae

Gryllus pennsylvanicus Burmeister

Aldrich, 1932a: 24, S. Dak. (ex *G. abbreviatus* Serville).

Gryllus spp.

All records ex *Gryllus assimilis* (Fabricius).

Aldrich, 1932a: 24, Calif.; Severin, 1926: 224, S. Dak.

STAGE PARASITIZED: Nymph, adult.

Exoristoides slossonae, see *Helioplagia slossonae*

Exoristoides sp.

HOST: ORTHOPTERA

Gryllidae

Anurogryllus muticus (De Geer)

Weaver and Sommers, 1969: 342, La.

Fabricia? *algens*, see *Nowickia algens*

Fabriciella hispida, see *Nowickia hispida*

Fabriciella latifrons, see *Norwickia latifrons*

Feather-legged fly, see *Trichopoda* spp.

“*Fortina*” sp., see *Lespesia* spp.

“*Frontia*” *frenchii*, see *Lespesia frenchii*

Frontina aletiae, see *Lespesia aletiae*

Frontina “*aletinae*,” see *Lespesia aletiae*

Frontina ancilla, see *Pseudomyoethyria ancilla*

Frontina “*archippiva*,” see *Lespesia archippivora*

Frontina archippivora, see *Lespesia archippivora*

Frontina armigera, see *Eucelatoria armigera*

Frontina frenchii, see *Lespesia frenchii*

Frontina irrequieta, see *Tachinomyia panaetius*

Frontina malacosomae, see *Lespesia frenchii*

- Frontina rileyi*, see *Lespesia rileyi*
Frontina tenthredinidarum, see *Bessa harveyi*
Frontina violenta, see *Lespesia testacea*
Frontina sp., see *Lespesia* spp.
Frontiniella Townsend (Goniinae: Eryciini)
Frontiniella parancilla Townsend
 HOST: LEPIDOPTERA

Gelechiidae

- Gnorimoschema* sp.
 Schaffner, 1959: 37, 51, Mass.

Pyralidae

- Tetralopha scortealis* (Lederer)
 Patton, 1958: 38, Fla.; Poos and Hetrick, 1945: 315, Va.
 STAGE PARASITIZED: Larva, pupa.
 BIOLOGY: Schaffner, 1959: (37, 51); Patton, 1958: (38).
Genea Rondani (Dexiinae: Leskiini)
Genea aurea James
 HOST: LEPIDOPTERA

Pyralidae

- Tetralopha subcanalis* (Walker)
 James, M. T., 1947: 113, Fla.
Genea texensis, see *Dejeaniopalpus texensis*
Gilvella Mesnil (Goniinae: Hyperecteinini)
Gilvella gilvipes (Coquillett)
 HOST: ORTHOPTERA

Gryllidae

- Cricket(-s)
 Aldrich, 1928: 301, Kans. (as *Oedematocera*).
Ginglymia Townsend (Goniinae: Siphonini)
Ginglymia acrirostris Townsend
 All records as *Ginglymyia acrirostris*.
 HOST: LEPIDOPTERA

Pyralidae

Parargyractis fulicalis (Clemens)

All records ex *Elophila fulicalis*.

Boardman, 1939: 160-161; Johannsen, 1935: 37-38, figs. 73-74, N.Y.; Lloyd, J. T., 1920: 263-265, figs. 3-4 on pl. 25, N.Y.

STAGE PARASITIZED: Larva.

IMMATURES: Third instar larva — Lloyd, J. T., 1920: (fig. 4 on pl. 25).

Pupa — Boardman, 1939: (descr.); Johannsen, 1935: (descr., figs. 73-74); Lloyd, J. T., 1920: (descr., fig. 3 on pl. 25).

BIOLOGY: Boardman, 1939: (160-161); Johannsen, 1935: (37-38); Lloyd, J. T., 1920: (263-265).

Ginglymyia acirostris, see *Ginglymyia acirostris*

Goliathocera setigera, see *Elfia setigera*

Gonia Meigen (Goniinae: Goniini)

Gonia aldrichi Tothill

HOST: LEPIDOPTERA

Noctuidae

Agrotis orthogonia Morrison

Strickland, 1938: 217, Alta. (ex pale western cutworm).

Crymodes devastator (Brace)

*Vide Crumb, 1929: 40, 103 (ex *Sidemia*, Coquillett, 1897); Coquillett, 1897: 17, 25, Iowa (as *G. capitata* (De Geer) ex *Hadena devastatrix*).

Euxoa ochrogaster (Guenée)

Strickland, 1938: 217, Alta. (ex redbacked cutworm); King, K.M., and Atkinson, 1928: 170-173, Sask.

STAGE PARASITIZED: Larva.

Gonia aldrichi Tothill probably

HOST: LEPIDOPTERA

Noctuidae

Euxoa spp. (including *E. messoria* (Harris) and/or *E. ochrogaster* (Guenée))

*Vide pers. comm. Handford, 1970: [B.C.] (Handford, 1962); Anon., 1962: 121, B.C.; Handford, 1962: 19, B.C.

STAGE PARASITIZED: Larva.

Gonia sp. near *aldrichi* Tothill

HOST: LEPIDOPTERA

Noctuidae

Agrotis orthogonia Morrison

Thornley and Sorenson, 1940: 48, Utah.

STAGE PARASITIZED: Larva.

BIOLOGY: Thornley and Sorenson, 1940: (48).

Gonia angusta, see *Gonia crassicornis*

Gonia breviporceps Tothill

HOST: LEPIDOPTERA

Noctuidae

Euxoa ochrogaster (Guenée)

Morrison, 1940: 348, Alta., Sask.

Gonia "capitaia," see *Gonia* spp.

Gonia capitata, see *Gonia aldrichi* or *Gonia* spp.

Gonia capitata types B, C, or D, see *Gonia* spp.

Gonia crassicornis (Fabricius)

HOST: COLEOPTERA

Scarabaeidae

Phyllophaga spp.

All records ex *Lachnosterna* spp.

Wolcott, 1951a: 482, P.R. (questionable host); Wolcott,
1924: 223, P.R. (as *G. angusta* Macquart).

HOST: LEPIDOPTERA

Noctuidae

Feltia subterranea (Fabricius)

Patton, 1958: 82, Fla. (as *Salmacia*); Tissot, 1945: 64, Fla.
(as "*Phosococephalus*").

Prodenia sunia (Guenée)

All records ex "*Xylomiges*" *sunia*.

Wolcott, 1951a: 482, P.R. (as *G. pallens* Wiedemann);
Gowdey, 1926: 83, Jamaica (as *G. pallens*); Wolcott,
1924: 223, P.R. (as *G. pallens*).

Spodoptera eridania (Cramer)

Patton, 1958: 82, Fla. (as *Salmacia* ex *Prodenia*); Wolcott,
1951a: 482, P.R. (ex "*Xylomiges*").

Spodoptera exigua (Hübner)

All records ex *Laphygma exigua*.

Wilson, J. W., 1933: 3, 13, Fla.; Wilson, J. W., 1932: 39,
Fla.

Spodoptera frugiperda (J. E. Smith)All records ex *Laphygma frugiperda*.

*Wolcott, 1951a: 481-482, P.R. (Jones, T. H., and Wolcott, 1922; Jones, T. H., 1913; Van Dine, 1913b); Wolcott, 1924: 223, P.R.; Jones, T.H., and Wolcott, 1922: 47, P.R. (as *G. "grassicornis"*); Jones, T. H., 1913: 235, P.R.; Van Dine, 1913b: 202, P.R.

STAGE PARASITIZED: Larva (Lepidoptera), pupa (Lepidoptera), adult (Coleoptera).

Gonia exul, see *Gonia sequax*

Gonia "frontalis," see *Gonia frontosa*

Gonia frontosa Say

HOST: COLEOPTERA

Meloidae

Tetraonyx quadrimaculata (Fabricius)

Fattig, 1949: 30, Ga. (as *Salmacia*).

HOST: LEPIDOPTERA

Noctuidae

Crymodes devastator (Brace)

Crumb, 1929: 40, 103, Iowa (ex *Sidemia*); Brauer and Bergenstamm, 1895: 555, 593, N. Amer. (as *G. "frontalis"* ex *Hadena devastatrix*).

Feltia ducens Walker - *Feltia subgothica* (Haworth) complex

Crumb, 1929: 40, 66-67 (ex *F. ducens-subgothica*).

STAGE PARASITIZED: Larva (Lepidoptera), pupa (Lepidoptera).

BIOLOGY: Crumb, 1929: (40, 66-67).

Gonia fuscicollis Tothill

HOST: LEPIDOPTERA

Noctuidae

Euxoa ochrogaster (Guenée)

King, K. M., and Atkinson, 1928: 170-173, Sask. (as *G. "fuscicollis"*).

Feltia ducens Walker

Crumb, 1929: 40.

STAGE PARASITIZED: Larva.

Gonia "fuscicollis," see *Gonia fuscicollis*

Gonia "grassicornis," see *Gonia crassicornis*

Gonia longiforceps Tothill

HOST: LEPIDOPTERA

Noctuidae

Agrotis orthogonia Morrison

Morrison, 1940: 347, Alta.; Thornley and Sorenson, 1940: 48, Utah; Strickland, 1938: 217, Alta. (ex pale western cutworm); Cook, W. C., 1930: 31, Mont. (ex *Porosagrotis*).

Euxoa tristicula (Morrison)

Cook, W. C., 1930: 31, Mont.

STAGE PARASITIZED: Larva.

Gonia longipulvilli Tothill

HOST: LEPIDOPTERA

Noctuidae

Agrotis ipsilon (Hufnagel)

Gressitt, 1959: 4, Hawaii (as *Salmacia*).

Agrotis malefida Guenée

Chapman, Fife and Cavitt, 1935: 305, Tex. (ex *Feltia*).

Agrotis orthogonia Morrison

*Arnaud, 1957: 262, Sask. (as *Salmacia*, Morrison, 1940); Morrison, 1940: 354, Sask.

Feltia subterranea (Fabricius)

Arnaud, 1957: 262, Ariz. (as *Salmacia*); Crumb, 1929: 40 (ex *F. annexa* Treitschke).

Peridroma saucia (Hübner)

Crumb, 1929: 40 (ex *Lycophotia*).

STAGE PARASITIZED: Larva, pupa.

Gonia pallens, see *Gonia crassicornis***Gonia porca** Williston

HOST: LEPIDOPTERA

Noctuidae

Peridroma saucia (Hübner)

All records ex *Lycophotia margaritosa* (Haworth).

Essig, 1926: 584; Essig, 1915: 401, Calif.

Pseudaletia unipuncta (Haworth)

Essig, 1926: 584 (ex *Cirphis*).

Cutworm(-s)

Essig, 1926: 584.

STAGE PARASITIZED: Larva.

Gonia sagax Townsend

HOST: LEPIDOPTERA

Geometridae

Alsophila pometaria (Harris)

Schaffner and Griswold, 1934: 90, 124, N.J.

STAGE PARASITIZED: Pupa.

BIOLOGY: Schaffner and Griswold, 1934: (90, 124).

"*Gonia sagax* Williston"

HOST: LEPIDOPTERA

Arctiidae

Unidentified Arctiidae

Allen, 1929: 689, Miss.

STAGE PARASITIZED: Pupa.

Gonia sequax Williston

HOST: LEPIDOPTERA

Noctuidae

Agrotis ipsilon (Hufnagel)

Rockwood, 1925: 720, Oreg. (as *G. exul* Williston ex
A. ypsilon).

Peridroma saucia (Hübner)

Anon., 1969: 59, Okla.; *Crumb, 1929: 40, 115 (ex
Lycophotia, Wakeland, 1920); Wakeland, 1920: 18,
Colo.

Spodoptera frugiperda (J. E. Smith)

Brauer and Bergenstamm, 1895: 555, 595, N. Amer. (as *G.*
exul ex *Laphygma*).

STAGE PARASITIZED: Larva, pupa.

IMMATURES: Pupa — Greene, 1922: (descr., fig. 67 on pl. 15).

BIOLOGY: Rockwood, 1925: (720).

Gonia texensis Reinhard

HOST: LEPIDOPTERA

Noctuidae

Mocis spp. (including *M. latipes* (Guenée) and/or *M. repanda*
(Fabricius))

Scaramuzza, 1946: 549, Cuba (ex *M. repanda*).

Spodoptera frugiperda (J. E. Smith)

Most records ex *Laphygma frugiperda*.

Scaramuzza and Barry, 1960: 997, Cuba; Scaramuzza, 1951a: 299, Cuba; Scaramuzza, 1942: 148, Cuba; Bruner, S. C., 1934: 96, Cuba; Van Dine and Christenson, 1932: 2, Cuba; Scaramuzza, 1929: 122, Cuba (ex *Laphygma* [sp.]); Scaramuzza, 1929a: 113, Cuba (ex *Laphygma* [sp.]).

STAGE PARASITIZED: Larva.

Gonia spp. (1) or (2), see *Gonia* spp.

Gonia spp.

Most records as *Gonia capitata* (De Geer).

HOST: LEPIDOPTERA

Lymantriidae

Nygmia phaeorrhoea (Donovan)

Burgess and Crossman, 1929: 143.

Noctuidae

Agrotis ipsilon (Hufnagel)

All records ex *Agrotis ypsilon*.

Crumb, 1929: 40 (as *Gonia* sp.); *Crumb, 1929: 59 (Sanderson, 1906); Sanderson, 1906: 8, Tex.

Agrotis orthogonia Morrison

Most records ex *Porosagrotis orthogonia*.

*Sweetman, 1936: 232, Canada (Strickland, 1923); Seamans and McMillan, 1935: 424 (as *Gonia* spp. ex *Agrotis*); Essig, 1926: 584, Canada; Seamans, 1923: 53, Alta.; Strickland, 1923: 22, figs. 8-12, 14-15 (as *G. capitata* type B, lab.); Strickland, 1923: 22, figs. 8-12, 14-15 (as *G. capitata* type D, unsuccessful lab. rearing); Strickland, 1923: 18-29, 39, figs. 8-12, 14-15, Alta. (as *G. capitata* et al.); Strickland, 1923: 22, figs. 8-12, 14-15, Alta. (as *G. capitata* type C); Strickland, 1921: 98-99, Alta.

Agrotis sp.

Tothill, 1913: 72, Ont. (ex agrotid sp.).

Anticarsia gemmatalis Hübner

All records ex *Anticarsia* "gemmatilis."

Wolcott, 1951a: 481, P.R. (as *Gonia* sp.); Wolcott, 1924: 223, P.R. (as *Gonia* sp.).

Crymodes devastator (Brace)

Johnson, 1910: 783, N.J. (ex *Hadena devastatrix*).

Euxoa ochrogaster (Guenée)

Schaffner and Griswold, 1934: 42, 124, Mass. (as *Gonia* sp. (1)); King, K. M., and Atkinson, 1928: 170-173, Sask. (as *Gonia* spp.); Essig, 1926: 584, Canada; Strickland, 1923: 39, figs. 8-12, 14-15, Alta. (as *G. capitata* et al.); Strickland, 1923: 22, figs. 8-12, 14-15 (as *G. capitata* type B, unsuccessful lab. rearing); Tothill, 1913: 72, "CowBay, C.B." (ex *Paragrotis*).

Heliothis zea (Boddie)

Most records ex *Heliothis obsoleta* (Fabricius).

Bosch and Hagen, 1966: 25, Calif. (ex *H. zea*); Essig, 1926: 584, U.S.; Quaintance and Brues, 1905: 123, Tex. (as *G. "capitaia"*).

Lithophane antennata (Walker)

Schaffner and Griswold, 1934: 49, 124, N.J. (as *Gonia* sp. (2) ex *Graptolitha*).

Peridroma saucia (Hübner)

*Crumb, 1929: 40, 115 (ex *Lycophotia*, Coquillett, 1897); Essig, 1926: 584, Calif., Colo. (ex *Lycophotia margaritosa* (Haworth)); Johnson, 1910: 783, N.J.; Chittenden, 1901a: 62; Coquillett, 1897: 17, 26, Calif.

Scotogramma trifolii (Rottenburg) probably

Lintner, 1889: 323, Conn. (as *Gonia* sp. ex *Mamestra*).

Spodoptera frugiperda (J. E. Smith)

All records ex *Laphygma frugiperda*.

Patton, 1958: 82, Fla. (as *Salmacia* sp.); Coquillett, 1897: 17, 25, Ark.

Cabbage cutworm(-s)

Lintner, 1887: 90, N.Y. (as *Gonia* sp.).

Cutworm(-s)

Aldrich, 1915: 82.

STAGE PARASITIZED: Larva, pupa.

IMMATURES: Egg — Strickland, 1923: (descr., figs. 8, 14).

First instar larva — Strickland, 1923: (descr., figs. 9, 14-15).

Second instar larva — Strickland, 1923: (descr., fig. 10).

Third instar larva — Strickland, 1923: (descr., fig. 11).

Pupa — Strickland, 1923: (descr., fig. 12); Greene, 1922: (descr., fig. 66 on pl. 14).

BIOLOGY: Sweetman, 1936: (232); Seamans and McMillan, 1935: (424); Schaffner and Griswold, 1934: (42, 49, 124); Strickland, 1923: (18-29, 39).

Goniomima unifasciata, see *Triachora unifasciata*

"*Goniomyia*" *unifasciata*, see *Triachora unifasciata*

Grisdalemyia Curran (Dexiinae: Macquartiini)

Grisdalemyia spp.

HOST: COLEOPTERA

Chrysomelidae

Chrysomela crotchii Brown

Smereka, 1965: 548, Ont.

STAGE PARASITIZED: Larva

Guerinia Robineau-Desvoidy (Goniinae: Exoristini)**Guerinia dydas** (Walker)

HOST: HYMENOPTERA

Diprionidae

Diprion hercyniae (Hartig)Daviault, 1944: 40, Que. (as *G. "spinosila"* (Townsend) ex *Gilpinia*).

Tenthredinidae

Allantus cinctus (Linnaeus)Chittenden, 1908: 12, Pa. (as *Tachina rustica* Fallén ex *Emphytus*).*Macremphytus intermedius* (Dyar)All records ex "*Macrephytus*" *intermedius*.*Procter, 1946: 434, Maine (as *Exorista simulans* Meigen, Procter, 1938); Procter, 1938: 375, Maine (as *Exorista simulans*).*Macremphytus tarsatus* (Say)Tothill, 1913: 74, Ont. (as *Tachina simulans* ex *Harpiphorus*).*Macremphytus testaceus* (Norton)

Raizenne, 1957: 38, Ont.

Macremphytus varianus (Norton)Raizenne, 1957: 38, Ont.; Greene, 1921: 41, Md. (as *Tachina rustica*).*Pristiphora geniculata* (Hartig)Forbes and Daviault, 1964: 1128 (as *Exorista simulans*).

Unidentified Hymenoptera

Sawfly(-flies)

Patton, 1958: 38, Fla. (as *G. simulans* ex small sawfly on *Fraxinus* sp.); Essig, 1926: 583 (as *Tachina rustica*);

*Brauer and Bergenstamm, 1895: 577, 608, N. Amer. (as

(*Tachina*) *tenthredinivora* Townsend ex *Tenthredo* sp.,
Townsend, 1892a); Townsend, 1892a: 285, Canada (as
Tachina tenthredinivora).

STAGE PARASITIZED: Larva, pupa.

IMMATURES: Pupa — Greene, 1922: (descr., fig. 78 on pl. 17).

Guerinia rufostomata (Bigot)

HOST: HYMENOPTERA

Unidentified Hymenoptera

Sawfly(-flies)

Taylor, R. L., 1929: 223, Mass. (as *Chaetotachina simulans*
(Meigen), probable host).

BIOLOGY: Taylor, R. L., 1929: (223).

Guerinia simulans, see *Guerinia dydas*

Guerinia "spinosila," see *Guerinia dydas*

Guerinia spinosula, see *Guerinia dydas*

Gymnocarcelia Townsend (Goniinae: Carceliini)

Gymnocarcelia ricinorum Townsend

Most records as *Sturmia albifrons* (Walker).

HOST: LEPIDOPTERA

Arctiidae

Apantesis phyllira (Drury)

Townsend, 1919: 582, Fla. (as *G. ricinorum* ex
"Apantesia").

Diacrisia virginica (Fabricius)

Schaffner and Griswold, 1934: 36, 118 (as *S. ricinorum*).

Ecpantheria deflorata (Fabricius)

All records ex *Ecpantheria scribonia* Stoll.

Johnson, 1910: 781, N.J.; Coquillett, 1897: 20, 25, Fla.

Estigmene acraea (Drury)

Most records ex *Estigmene acraea*.

Carrillo S., Ortega C., and Gibson, 1966: 94, Mexico (Son.)

(as *G. ricinorum*); Patton, 1958: 38-39, Fla. (as *G. ricinorum* ex *E. acraea*); Schaffner and Griswold, 1934: 38, 118 (as *S. ricinorum* ex *E. acraea*); Essig, 1926: 583, Calif.; Reinhard, 1922: 73, Tex.; Winn and Beaulieu, 1915: 142, Que. (ex *Leucarctia* "acraca"); Johnson, 1910: 781, N.J. (ex *Leucarctia*); Coquillett, 1897: 20, 26, Calif., Conn., S.C., Tex. (ex *Leucarctia*).

Utetheisa ornatrix bella (Linnaeus)

Schaffner and Griswold, 1934: 39-40, 118, N.J. (as *S. ricinorum* ex *U. bella*).

Unidentified Arctiidae

Schaffner and Griswold, 1934: 40-41, 118 (as *S. ricinorum*);
Aldrich, 1932: 3, Mass. (as *S. ricinorum*).

Noctuidae

Pseudaletia unipuncta (Haworth)

Essig, 1926: 583, Calif. (ex *Cirphis*); Tothill, 1913: 74, Ont.
(ex *Heliophila*).

Unidentified Noctuidae

Johnson, 1925a: 87, N.H. (as *G. ricinorum* ex *Noctua* sp.).

STAGE PARASITIZED: Larva, pupa.

IMMATURES: Pupa — Greene, 1922: (descr., fig. 99 on pl. 20).

BIOLOGY: Patton, 1958: (38-39); Schaffner and Griswold,
1934: (118).

Gymnocarcelia ricinorum Townsend probably

All records as *Hemimasicera* spp.

HOST: LEPIDOPTERA

Arctiidae

Ecpantheria deflorata (Fabricius)

Brauer and Bergenstamm, 1895: 556, 591, N. Amer. (ex *E.*
scribonia Stoll).

Estigmene acraea (Drury)

Brauer and Bergenstamm, 1895: 556, 597, N. Amer. (ex
Leucarctia acraea).

Gymnocarcelia sp.

HOST: LEPIDOPTERA

Arctiidae

Estigmene acraea (Drury)

Taylor, E. A., 1954: 526, Ariz.

STAGE PARASITIZED: Larva.

Gymnochaeta ruficornis, see *Gymnochaeta ruficornis*

Gymnochaetopsis Townsend (Tachininae: Linnaemyini)

Gymnochaetopsis fulvicauda (Walton)

Most records as *Linnaemyia fulvicauda*.

HOST: LEPIDOPTERA

Noctuidae

Heliothis zea (Boddie)

Gowdey, 1923a: 54, Jamaica (ex *Chloridea obsoleta*
(Fabricius)).

Mocis spp. (including *M. latipes* (Guenée) and/or *M. repanda* (Fabricius))

Wolcott, 1951a: 481, P.R. (ex *M.* (or *Remigia*) *repanda*);
Gowdey, 1926: 83, Jamaica (ex *M. punctularis* (Hübner)); Vickery, 1924: 405, P.R. (as *Linnaemya* ex *M. repanda*); Wolcott, 1924: 223, P.R. (ex *Remigia repanda*); Gowdey, 1923a: 21, Jamaica (ex "*Mocia*" *repanda*); Jones, T. H., and Wolcott, 1922: 49, P.R. (ex *Remigia repanda*); Walton, 1914a: 93, P.R. (ex *Remigia repanda*).

STAGE PARASITIZED: Larva.

IMMATURES: Pupa — Greene, 1922: (descr., fig. 29 on pl. 7).

Gymnocheta Robineau-Desvoidy (Tachininae: Linnaemyini)

Gymnocheta ruficornis Williston

All records as *Gymnochaeta ruficornis*.

HOST: LEPIDOPTERA

Noctuidae

Papaipema arctivorens Hampson

Decker, 1935: 569, 578, Iowa.

Papaipema cataphracta (Grote)

Decker, 1935: 569, 578, Iowa.

Papaipema nebris (Guenée)

Decker, 1935: 568-569, 578, Iowa; Decker, 1931: 344, Iowa.

STAGE PARASITIZED: Larva.

BIOLOGY: Decker, 1935: (568-569, 578); Decker, 1931: (344).

Gymnoclytia Brauer and Bergenstamm (Phasiinae: Gymnosomatini)

Gymnoclytia dubia (West)

HOST: HEMIPTERA (HETEROPTERA)

Pentatomidae

Cosmopepla bimaculata (Thomas)

West, 1925: 122, N.Y. (as *Gymnosoma*).

Gymnoclytia immaculata (Macquart)

All records as *Cistogaster immaculata*.

HOST: HEMIPTERA (HETEROPTERA)

Pentatomidae

Euschistus variolarius (Palisot de Beauvois)

Parish, 1934: 54, Iowa (as "*Cistogaster*").

HOST: LEPIDOPTERA

Noctuidae

Pseudaletia unipuncta (Haworth)

All records ex *Leucania unipuncta*.

Lintner, 1897: 210; *Brauer and Bergenstamm, 1895: 548, 596, Carolina (Townsend, 1893); Townsend, 1893: 466, Ill.

STAGE PARASITIZED: Larva (Lepidoptera), adult (Hemiptera-Heteroptera).

Gymnoclytia occidua (Walker)

HOST: HEMIPTERA (HETEROPTERA)

Pentatomidae

Euschistus variolarius (Palisot de Beauvois)

Rings and Brooks, 1958: 15-16, Ohio.

HOST: LEPIDOPTERA

Pieridae

Pieris protodice Boisduval and LeConte

Fattig, 1949: 18-19, Ga.

Pieris rapae (Linnaeus)

Fattig, 1949: 18-19, Ga.

STAGE PARASTIZED: Larva (Lepidoptera), adult (Hemiptera-Heteroptera).

Gymnophthalma americana, see *Actia americana*

Gymnophthalma interrupta, see *Actia interrupta*

Gymnosoma Meigen (Phasiinae: Gymnosomatini)

Gymnosoma dubia, see *Gymnoclytia dubia*

Gymnosoma filiola Loew

HOST: HEMIPTERA (HETEROPTERA)

Pentatomidae

Chlorochroa ligata (Say)

Brooks, 1946a: 218 (as *Rhodogyne*).

Gymnosoma fuliginosum Robineau-Desvoidy

HOST: HEMIPTERA (HETEROPTERA)

Pentatomidae

Chlorochroa ligata (Say)

Most records ex *Pentatoma ligata*.

Russell, E. E., 1952: 15, Ariz. (as *Rhodogyne*); Morrill, 1910: 64, figs. 9-10 on pl. 3, Tex., Mexico (Dgo.); Morrill, 1907: 11, Tex.

Chlorochroa sayi Stål

Russell, E. E., 1952: 15, Ariz. (as *Rhodogyne*); Clancy, 1946: 327-328, Ariz.; Strand, 1937: 26, Mont. (ex Say's plant bug); Knowlton and Janes, 1931: 147, Utah; Aldrich, 1915: 81, Ariz., N. Mex. (ex *Pentatoma*).

Chlorochroa uhleri Stål

Walster, 1942: 9, N. Dak.

Eurygaster spp.

Vojdani, 1961: 576, Calif.

Euschistus impictiventris Stål

Russell, E. E., 1952: 15, Ariz. (as *Rhodogyne*, incomplete rearing — [larva]).

Euschistus servus (Say)

Rolston and Kendrick, 1961: 156, Ark.

Euschistus variolarius (Palisot de Beauvois)

Parish, 1934: 54, Iowa.

Wheat bug(-s)

Strickland, 1938: 216.

STAGE PARASITIZED: Nymph, adult.

IMMATURES: Egg — Morrill, 1910: (fig. 9 on pl. 3).

Pupa — Greene, 1922: (descr., fig. 19 on pl. 5); Morrill, 1910: (fig. 10 on pl. 3).

BIOLOGY: Rolston and Kendrick, 1961: (156); Clancy, 1946: (327-328); Morrill, 1910: (64).

Gymnostylia Macquart (Goniinae: Blondeliini)

"*Gymnostylia*" sp.

HOST: HYMENOPTERA

Tenthredinidae

Unidentified Tenthredinidae

Brauer and Bergenstamm, 1895: 556, 608, N. Amer. (ex Tenthrediniden-Raupen auf Erlen).

STAGE PARASITIZED: Larva.

Hamaxia Walker (Phasiinae: Palpostomatini)

Hamaxia incongrua Walker

All records as *Ochromeigenia ormioides* Townsend.

HOST: COLEOPTERA

Scarabaeidae

Popillia japonica Newman

Sweetman, 1936: 121, 236, U.S.

Unidentified Scarabaeidae

Sweetman, 1936: 121, U.S. (ex Scarabaeidae (other than *Popillia japonica*)).

STAGE PARASITIZED: Adult.

BIOLOGY: Sweetman, 1936: (121, 236).

Helioplagia Townsend (Tachininae: Germariini)

Helioplagia slossonae (Coquillett)

HOST: COLEOPTERA

Coccinellidae

Epilachna varivestis Mulsant

Aldrich, 1932a: 25, Ala. (as *Exoristoides* ex *E. corrupta* Mulsant).

HOST: LEPIDOPTERA

Hesperiidae

Thymelicus lineola (Ochsenheimer)

Arthur, 1966: 625, [Canada]; Arthur, 1962: 1083, 1085, Ont.

STAGE PARASITIZED: Pupa (Lepidoptera).

Hemimasicerca sp., see *Gymnocarcelia ricinorum* probably

Hemisturmia Townsend (Goniinae: Winthemiini)

Hemisturmia tortricis (Coquillett)

Most records as *Phorocera tortricis*.

HOST: LEPIDOPTERA

Glyphipterygidae

Anthophila pariana (Clerck)

All records ex *Hemerophila pariana*.

*Cameron, 1931: 41, Conn. (Porter and Garman, 1923);
Porter and Garman, 1923: 261, Conn.

Nymphalidae

Nymphalis antiopa (Linnaeus)

Knowlton and Hanson, 1949: 6, Utah (ex "*Euvanesia*").

Olethreutidae

Ancylis comptana (Frölich)

Summerland, 1938: 172-173, Kans. (as *P. "torticis"*).

Pterophoridae

Pterophorus periscelidactylus Fitch

Putman, 1935a: 105, Ont.

Pyralidae

Acrobasis indigenella (Zeller)

Aldrich and Webber, 1924: 76, Mo. (ex *Mineola "indiginella"*).

Tortricidae

Acleris minuta (Robinson)

All records ex *Peronea minuta*.

*Aldrich and Webber, 1924: 76, N.J. (Weiss, 1917); Weiss, 1917: 220, N.J. (as *Neopales*).

Acleris variana (Fernald)

Miller, C. A., 1966: 603, N.S.

Archips argyrospilus (Walker)

Paradis and LeRoux, 1965: 44, 46, 48-49, fig. 14E, Que. (as *Ceratochaeta*); Aldrich and Webber, 1924: 76, Ark. (ex *Cacoecia*).

Archips cerasivoranus (Fitch)

Powell, 1962: 133, 136, Calif.; Schaffner, 1959: 30, 53, Mass., N.H., Vt.; Thompson, 1953: 19; Hoffman, 1936: 211, Minn. (ex *Cacoecia*); Putman, 1935a: 108, Ont. (ex *Cacoecia*); Johnson, 1925: 199, Vt. (ex *Cacoecia*); Aldrich and Webber, 1924: 76, Mass., Vt. (ex *Cacoecia*); Johnson, 1916: 81, Vt. (as *Neopales*); Herrick, 1912: 291, N.Y.

Archips rileyanus (Grote)

Wylie and Tadic, 1963: 165, Ark. (as *P. "torbricis"* ex *A. cerasivorana rileyana*).

Argyrotaenia velutinana (Walker)

Anon., 1963c: 20, Ont.

Choristoneura fumiferana (Clemens)

Coppel, 1960: 94, 97, B.C. (as *Ceratochaeta*); Ross, 1952: 112, fig. 13, Canada; Dowden and Carolin, 1950: 777,

N.Y. (ex *Archips*); Wilkes, Coppel, and Mathers, 1949: 146, 149, B.C.

Choristoneura pinus Freeman

Dixon and Benjamin, 1963: 267-270, Wis.

Choristoneura rosaceana (Harris)

Schuh and Mote, 1948: 26, 28, Oreg. (ex *Archips*).

Unidentified Tortricidae

Coquillett, 1897: 19, 27, Mich.

STAGE PARASITIZED: Larva, pupa.

IMMATURES: Pupa — Paradis and LeRoux, 1965: (fig. 14E); Ross, 1952: (fig. 13); Greene, 1922: (descr., fig. 75 on pl. 16).

BIOLOGY: Paradis and LeRoux, 1965: (44, 46, 48-49); Schaffner, 1959: (30, 53).

Hemithrixion Brauer and Bergenstamm (Goniinae: Acemyini)

Hemithrixion oestriforme, see *Hemithrixion oestriforme* or *Hemithrixion plankii*

Hemithrixion oestriforme Brauer and Bergenstamm

HOST: ORTHOPTERA

Acrididae

Boopedon nubilum (Say)

Newton, 1954: 936, Mont.

Camnula pellucida (Scudder)

Newton, 1954: 936, Mont.

Melanoplus bivittatus (Say)

Newton, 1954: 936, Mont.; Smith, R. W., 1944: 29, Canada.

Melanoplus bruneri Scudder

Newton, 1954: 936, Mont.

Melanoplus sanguinipes (Fabricius)

Paul and Putnam, 1960: 492, Sask. (ex *M. bilituratus* (Walker), probable host); Smith, R. W., 1944: 29, Canada (ex *M. mexicanus* Saussure).

Grasshopper(-s)

Anon., 1956: 196, W. Canada; Smith, R. W., and Finlayson, 1950: 98-99, figs. 24, 48, Canada; Strickland, 1946: 172, Alta.

STAGE PARASITIZED: Nymph, adult.

IMMATURES: Second instar larva — Smith, R. W., and Finlayson, 1950: (descr., figs. 24, 48).

BIOLOGY: Smith, R. W., and Finlayson, 1950: (98-99).

Hemithrixion plankii (Walton)

HOST: ORTHOPTERA

Acrididae

Grasshopper(-s)

*Greene, 1934: 31, N.J. (as *H. oestriforme* Brauer and Bergenstamm ex Quaintance No. 11808, Walton, 1915);
Walton, 1915: 105, 107, figs. 4-5 on pl. 13, N.J. (as *Coquilletina*).

IMMATURES: Pupa — Greene, 1922: (descr., fig. 14 on pl. 3);
Walton, 1915: (descr., figs. 4-5 on pl. 13).

Hemithrixion sp.

HOST: ORTHOPTERA

Acrididae

Grasshopper(-s)

Smith, R. W., 1939: 60, Sask.

Hemithrixion sp. probably

HOST: ORTHOPTERA

Acrididae

Dissosteira spurcata Saussure

Kelly and Middlekauff, 1961: 422, Calif.

STAGE PARASITIZED: Adult.

“Hemorilla” maculosa, see *Nemorilla pyste*

Hemyda Robineau-Desvoidy (Phasiinae: Cylindromyiini)

Hemyda aurata Robineau-Desvoidy

HOST: HEMIPTERA (HETEROPTERA)

Pentatomidae

Podisus maculiventris (Say)

Coppel and Jones, 1964: 191-194, figs. 1-8, Wis.

STAGE PARASITIZED: Adult.

IMMATURES: First instar larva — Coppel and Jones, 1964:
(descr., figs. 1-2).

Second instar larva — Coppel and Jones, 1964: (descr., fig. 3).

Third instar larva — Coppel and Jones, 1964: (descr., fig. 4).

Pupa — Coppel and Jones, 1964: (descr., figs. 5-8).

Homoeonychia rapae, see *Obolocera rapae*

Hubneria Robineau-Desvoidy (Goniinae: Carceliini)

Hubneria estigmenensis (Sellers)

HOST: LEPIDOPTERA

Arctiidae

Diacrisia virginica (Fabricius)

Sellers, 1943: 81 (as *Aplomya*).

Estigmene acraea (Drury)

Sellers, 1943: 81 (as *Aplomya* ex *E. acraea*); Schaffner and Griswold, 1934: 38, 111 (as *Zenillia affinis* (Fallén)).

Isia isabella (J. E. Smith)

Sellers, 1943: 81 (as *Aplomya*); Schaffner and Griswold, 1934: 36-37, 111 (as *Zenillia affinis*).

Phragmatobia fuliginosa (Linnaeus)

*Sellers, 1943: 81 (as *Aplomya*, Tothill, 1913); *Aldrich and Webber, 1924: 35 (as *Zenillia affinis*, Tothill, 1913); Tothill, 1913: 70, Ont. (as *Exorista affinis*).

Phragmatobia fuliginosa rubricosa (Harris)

Sellers, 1943: 81 (as *Aplomya*); Schaffner and Griswold, 1934: 35-36, 111 (as *Zenillia affinis*).

Unidentified Arctiidae

Essig, 1926: 581 (as *Zenillia affinis*).

STAGE PARASITIZED: Pupa.

IMMATURES: Pupa — Sellers, 1943: (descr.).

BIOLOGY: Sellers, 1943: (81); Schaffner and Griswold, 1934: (111).

Hyalomya Robineau-Desvoidy (Phasiinae: Phasiini)

Hyalomya aldrichii Townsend

HOST: HEMIPTERA (HETEROPTERA)

Lygaeidae

Emblethis vicarius Horvath

Scudder, 1966: 286, B.C.

Geocoris bullatus (Say)

Clancy and Pierce, 1966: 857-858, Calif.

Geocoris punctipes (Say)

Clancy and Pierce, 1966: 857-858, Calif.

Geocoris uliginosus speculator Montandon

Clancy and Pierce, 1966: 857-858, Calif.

Nysius ericae (Schilling)

Bird, R. D., and Mitchener, 1953: 44, Man. (as "*Hyalomyopsis*," probable host).

Nysius raphanus Howard

Clancy and Pierce, 1966: 858, Calif.

Unidentified Hemiptera (Heteroptera)

Miscellaneous

Bird, R. D., and Mitchener, 1953: 44 (as "*Hyalomyopsis*").

STAGE PARASITIZED: Adult.

Hyalomya purpurascens Townsend

HOST: HEMIPTERA (HETEROPTERA)

Cydniidae

Sehirus cinctus Palisot de Beauvois

Brooks, 1945a: 676, Ga. (as *Alophorellopsis*).

Hyalomyia ecitonis, see *Androeurops ecitonis*

Hyalomyiopsis aldrichii, see *Hyalomya aldrichii*

Hyalomyodes Townsend (Phasiinae: Strongygastrini)

Hyalomyodes "triangularis," see *Hyalomyodes triangulifer*

Hyalomyodes triangulifer (Loew)

HOST: COLEOPTERA

Alleculidae

Chromatia amoena (Say)

Thompson, 1954: 137-143, figs. 1-7, 10-12, 16, 20, 23 on
pls. 1-2, Mass.

Chrysomelidae

Altica bimarginata Say

Thompson, 1954: 139-142, figs. 1-7, 10-12, 16, 20, 23 on
pls. 1-2 (ex *Haltica* "*bimarginata*"); Johnson, 1925: 191,
Conn., Maine, Mass., N.H., R.I., Vt. (ex *Haltica*); Woods,
W. C., 1917: 273-274, Maine (as *H.* "*triangularis*").

Chrysomela interrupta Fabricius

Thompson, 1954: 139-143, figs. 1-7, 10-12, 16, 20, 23 on
pls. 1-2, Ont. (ex *Lina*).

Chrysomela knabi Brown

Brown, W. J., 1957: 11, Ont.

Oulema melanopus (Linnaeus)

Wellso and Hoxie, 1969: 923-924, Mich.

Coccinellidae

Coccinella trifasciata perplexa Mulsant

Smith, B. C., 1960: 652, Ont.

Coleomegilla maculata (De Geer)

All records ex *Ceratomegilla fuscilabris* Mulsant.

Thompson, 1954: 139-143, figs. 1-7, 10-12, 16, 20, 23 on pls. 1-2; Howard, N. F., and Landis, 1936: 6, N.Y.

Coleomegilla maculata lengi Timberlake

Smith, B. C., 1960: 652, Ont.

Epilachna varivestis Mulsant

Thompson, 1954: 139-143, figs. 1-7, 10-12, 16, 20, 23 on pls. 1-2, Mexico (Mor.); Howard, N. F., and Landis, 1936: 3, 6, Mexico.

Curculionidae

Hypera postica (Gyllenhal)

Wellso and Hoxie, 1969: 924, Mich.; Brunson and Coles, 1968: 6, E. U.S. (as "*Hyaolmyodes*").

Hypera punctata (Fabricius)

Wellso and Hoxie, 1969: 924, Mich.

Listroderes costirostris obliquus (Klug)

Beckham, 1955: 10, Ga.; Beckham, 1953: 20, Ga.; Beckham, 1952: 541, Ga.; Beckham, 1952a: 38-39, Ga. (ex vegetable weevil).

Sitona hispidula (Fabricius)

Thompson, 1954: 137-143, figs. 1-7, 10-12, 16, 20, 23 on pls. 1-2, Ont.

HOST: LEPIDOPTERA

Pterophoridae

Platyptilia carduidactyla (Riley)

Lange, 1950: 592, Calif.; Lange, 1941: 24, 67, Calif.

STAGE PARASITIZED: Larva (Coleoptera, Lepidoptera), adult (Coleoptera).

IMMATURES: First instar larva — Thompson, 1954: (descr., figs. 3, 6, 10, 20 on pls. 1-2).

Second instar larva — Thompson, 1954: (descr., figs. 1, 5, 11-12 on pls. 1-2).

Third instar larva — Thompson, 1954: (descr., figs. 2, 7, 16, 23 on pls. 1-2).

Pupa — Thompson, 1954: (descr., fig. 4 on pl. 1); Greene, 1922: (as *H. "triangulifera"*; descr., fig. 56 on pl. 12).

BIOLOGY: Wellso and Hoxie, 1969: (923-924); Beckham, 1953: (20).

Hyalomyodes "triangulifera," see *Hyalomyodes triangulifer*

"*Hyalomyopsis*" *aldrichii*, see *Hyalomya aldrichii*

Hyalurgus Brauer and Bergenstamm (Dexiinae: Hyalurgini)

Hyalurgus sp.

HOST: HYMENOPTERA

Tenthredinidae

Tomostethus multicoloratus (Rohwer)

Anon., 1954e: 253, Ont.

STAGE PARASITIZED: Larva.

"*Hyalomyodes*" *triangulifera*, see *Hyalomyodes triangulifer*

Hylotomomyia hylotomae, see *Spathimeigenia hylotomae*

"*Hylotomomyia hylotomae*," see *Spathimeigenia hylotomae*

"*Hypercteina*" *aldrichi*, see *Hyperecteina aldrichi*

Hyperecteina Schiner (Goniinae: Hyperecteinini)

Hyperecteina aldrichi Mesnil

Most records as *Centeter cinerea* Aldrich.

HOST: COLEOPTERA

Scarabaeidae

Maladera castanea (Arrow) probably

Smith, L. B., 1929: 124-125, Conn., N.J., Pa. (ex Asiatic beetle).

Popillia japonica Newman

Fleming, 1963: 25, N.J. (as "*Hypercteina*" *aldrichi*); Clausen, 1956: 23, 107-108, Conn., Mass., N.J., Pa. (as *H. aldrichi*); Cory and Langford, 1955: 11, 20, Md. (ex Japanese beetle); Schread, 1948: 8, Conn. (ex Japanese beetle); Schread, 1947: 13, 17-19, Conn.; Schread, 1946: 49, Conn. (ex Japanese beetle); Schread, 1945: 339, Conn. (ex Japanese beetle); Slate, 1945: 60, Conn. (ex Japanese beetle); Adams, 1944: 9, fig. 4, N.Y. (as *Centeter* fly ex Japanese beetle); Adams and Wheeler, 1944: 25, N.Y. (ex Japanese beetle); Rex, 1944: 25, N.J.; French, 1943: 62, W. Va. (ex Japanese beetle); King and Parker, 1941: 127, 130, 138, Conn., N.J., Pa. (ex Japanese beetle); Vosburgh, 1941: 227, 236, 240, unnumbered fig. on p. 227, N.J. (ex Japanese beetle); Rex, 1940: 25, N.J.; King, 1939: 26, Conn., N.H., N.J., Pa. (ex Japanese beetle); Britton and Johnson, 1938: 468, Conn.; Hadley, 1938: 211; King, 1937: 475, 482, Conn., N.H., N.J., Pa. (ex Japanese beetle); Breshears and Dawson, 1936: 9, U.S.; Sweetman, 1936: 119, 121-122,

236, 245, figs. 33, 36, 102, U.S.; Rex, 1934: 24-25; Bade, 1933: 16; King, 1931: 453, 455-457, Conn., N.J.; King, 1930: 158-160, fig. 20, Conn., N.J., Pa.; Rex, 1930: 22-23, figs. 5-6 on fig. 9, N.J.; Smith, L. B., 1930: 500, Conn., Pa. (ex Japanese beetle); Showalter, 1929: pl. 17; Smith, L. B., 1929: 124-125, Conn., N.J., Pa. (ex Japanese beetle); Anon., 1928: 296-297, Conn., Pa.; Anon., 1928a: 943, Conn., Pa. (ex Japanese beetle); Allen and Jaynes, 1928: 823-824, 828-829, 831-832, N.J., Pa.; King, 1928: 404-405, fig. 131, N.J.; Britton, 1927: 246, Conn.; Clausen, King and Teranishi, 1927: 4-17, 54, figs. 11A-C, 12, 14D on pl. 1, N.J.; King and Hallock, 1925: 352-353, N.J.

STAGE PARASITIZED: Larva, adult.

IMMATURES: Egg — Adams, 1944: (fig. 4); Vosburgh, 1941: (unnumbered fig. on p. 227); Sweetman, 1936: (figs. 33, 102); King, 1930: (fig. 20); Rex, 1930: (descr., figs. 5-6 on fig. 9); Showalter, 1929: (pl. 17); King, 1928: (fig. 131); Clausen, King and Teranishi, 1927: (fig. 12 on pl. 1).

Larva — King, 1930: (fig. 20).

First instar larva — Sweetman, 1936: (fig. 36); Clausen, King and Teranishi, 1927: (descr., fig. 11A on pl. 1).

Second instar larva — Clausen, King and Teranishi, 1927: (descr., fig. 11B on pl. 1).

Third instar larva — Sweetman, 1936: (fig. 36); Rex, 1930: (fig. 5 on fig. 9); Clausen, King and Teranishi, 1927: (descr., fig. 11C on pl. 1).

Pupa — Clausen, King and Teranishi, 1927: (descr., fig. 14D on pl. 1).

BIOLOGY: Clausen, 1956: (6, 23, 107-108); Cory and Langford, 1955: (11, 20); Schread, 1947: (13, 17-19); Adams, 1944: (9); Rex, 1944: (25); King and Parker, 1941: (127, 130, 138); King, 1939: (26); Britton and Johnson, 1938: (468); Hadley, 1938: (211); King, 1937: (475, 482); Sweetman, 1936: (119, 121-122, 236, 245); Rex, 1934: (24-25); King, 1931: (453, 455-457); Rex, 1930: (22-23); Anon., 1928: (296-297); Anon., 1928a: (943); King, 1928: (404-405); Britton, 1927: (246); Clausen, King and Teranishi, 1927: (4-17, 54).

Hyperecteina hylotomae, see *Spathimeigenia hylotomae*

Hyperecteina "retinae," see *Urophyllopsis retiniae*

Hyperecteina retiniae, see *Urophyllopsis retiniae*

Hyphantrophaga Townsend (Goniinae: Carceliini)

Hyphantrophaga desmiae (Sellers)

HOST: LEPIDOPTERA

Pyralidae

Desmia funeralis (Hübner)Sellers, 1943: 17, Calif. (as *Zenillia*).

STAGE PARASITIZED: Pupa.

Hyphantrophaga euchaetiae (Sellers)All records as *Zenillia euchaetiae*.

HOST: LEPIDOPTERA

Arctiidae

Cycnia inopinatus (H. Edwards)

Sellers, 1943: 14.

Cycnia tenera Hübner

Sellers, 1943: 14, N.J.

Euchaetias egle (Drury)

Sellers, 1943: 14, N.J., N.Y.

Pyralidae

Pyrausta futilalis (Lederer)

Schaffner, 1959: 11, 54, N.J.; Sellers, 1943: 14.

STAGE PARASITIZED: Pupa.

BIOLOGY: Schaffner, 1959: (11, 54); Sellers, 1943: (15).

Hyphantrophaga "hyphantria," see *Hyphantrophaga hyphantriae***Hyphantrophaga hyphantriae** (Townsend)

HOST: LEPIDOPTERA

Arctiidae

Cycnia inopinatus (H. Edwards)Schaffner and Griswold, 1934: 33, 111 (as *Zenillia ceratomiae* (Coquillett)).*Cycnia tenera* HübnerSchaffner and Griswold, 1934: 32, 111 (as *Zenillia ceratomiae*).*Euchaetias egle* (Drury)Schaffner and Griswold, 1934: 39, 111 (as *Zenillia ceratomiae*).*Hyphantria cunea* (Drury)Sellers, 1943: 16 (as *Zenillia*); *Essig, 1926: 580 (Coquillett, 1897); Coquillett, 1897: 17, 25, N. Mex.; *Brauer and

Bergenstamm, 1895: 566, 594 (as *H. "hyphantria,"*
Townsend, 1892b; Townsend, 1891b); Townsend,
1892b: 258-259 (ex fall webworm); Townsend, 1891b:
176-177, N. Mex. (as *Meigenia*).

Hyphantria textor (Harris)

*Brauer and Bergenstamm, 1895: 564, 594 (as *Nemoraea*,
Townsend, 1893); Townsend, 1893: 467, Ill. (as
Nemoraea).

Citheroniidae

Anisota senatoria J. E. Smith

Webber and Schaffner, 1926: 25, New England (as *Zenillia*
ceratomiae).

Geometridae

Eucaterva variaria Grote

*Sellers, 1943: 16 (as *Zenillia*, Townsend, 1892b); *Essig,
1926: 580-581, N. Mex. (Townsend, 1892b); Townsend,
1892b: 258-259, N. Mex.

Lygris diversilineata (Hübner)

Sellers, 1943: 16, Tex. (as *Zenillia*).

Lasiocampidae

Gloveria howardi Dyar

Sellers, 1943: 16 (as *Zenillia*).

Malacosoma americanum (Fabricius)

Stehr and Cook, 1968: 258, 299-300, Tex.

Malacosoma californicum (Packard)

Stehr and Cook, 1968: 264, 300, Ariz.

Malacosoma incurvum discoloratum (Neumoegen)

Stehr and Cook, 1968: 280-281, 301, Nev., Utah.

Malacosoma incurvum incurvum (H. Edwards)

Stehr and Cook, 1968: 280, 301, Ariz.

Megalopygidae

Megalopyge crispata (Packard)

All records ex *Lagoa crispata*.

Sellers, 1943: 16, Fla. (as *Zenillia*); Aldrich and Webber,
1924: 42, Fla. (as *Zenillia ceratomiae*).

Noctuidae

Heliopsis zea (Boddie)

Quaintance and Brues, 1905: 123, Tex. (as *Exorista ceratomiae* ex *H. obsoleta* (Fabricius)).

Spodoptera frugiperda (J. E. Smith)

All records ex *Laphygma frugiperda*.

Sellers, 1943: 16 (as *Zenillia*); Luginbill, 1928: 81, Tex.;
Luginbill, 1928: 82, Tenn. (as *Exorista ceratomiae*).

Nymphalidae

Nymphalis milberti (Godart)

Sellers, 1943: 16 (as *Zenillia*); Essig, 1926: 580-581, Colo.
(ex *Aglais*); Townsend, 1895: 75, Colo. (ex *Vanessa*).

Pyralidae

Acrobasis caryae Grote

*Vide Sellers, 1943: 16 (as *Zenillia*, Leiby, 1925); Brimley,
1938: 362, N.C.; Leiby, 1925: 18, N.C. (as *Didyma exigua* Wulp).

Desmia funeralis (Hübner)

Donohoe and Kaloostian, 1937: 967-968, Calif. (as *Zenillia*
"ceratomyiae").

Loxostege similalis (Guenée)

Butler, 1958: 562, Ariz. (as *Zenillia*); Fattig, 1949: 37 (as
Zenillia); Sellers, 1943: 16, Okla., Tex. (as *Zenillia*);
Aldrich and Webber, 1924: 42, Okla., Tex. (as *Zenillia*
ceratomiae); Reinhard, 1922: 72, Tex. (as *Exorista*
ceratomyiae).

Omphalocera cariosa Lederer

Sellers, 1943: 16, Kans. (as *Zenillia*); Aldrich and Webber,
1924: 42, Kans. (as *Zenillia ceratomyiae*); Coquillett,
1897: 13, 26, Kans. (as *Exorista ceratomyiae*).

Pempelia spp.

*Sellers, 1943: 16, Tex. (as *Zenillia*, Coquillett, 1897);
*Aldrich and Webber, 1924: 42, Tex. (as *Zenillia*
ceratomyiae, Coquillett, 1897); Coquillett, 1897: 13, 26,
Tex. (as *Exorista ceratomyiae*).

Unidentified Pyralidae

*Sellers, 1943: 16, Mo. (as *Zenillia*, Coquillett, 1897);
*Aldrich and Webber, 1924: 42, Mo. (as *Zenillia*

ceratomiae, Coquillett, 1897); Coquillett, 1897: 13, 26, Mo. (as *Exorista ceratomiae*).

Sphingidae

Ceratomia catalpae (Boisduval)

Fattig, 1949: 37, Ga. (as *Zenillia*).

Ceratomia undulosa (Walker)

*Sellers, 1943: 16 (as *Zenillia*, Coquillett, 1897); *Aldrich and Webber, 1924: 42, Mo. (as *Zenillia ceratomiae*, Coquillett, 1897); Coquillett, 1897: 13, 24, Mo. (as *Exorista ceratomiae*).

STAGE PARASITIZED: Larva, pupa.

IMMATURES: Egg — Townsend, 1892b: (descr.).

Pupa — Greene, 1922: (descr., fig. 26 on pl. 6).

BIOLOGY: Schaffner and Griswold, 1934: (111); Townsend, 1892b: (258-259).

Hyphantrophaga sp.

HOST: LEPIDOPTERA

Nymphalidae

Nymphalis milberti (Godart)

Townsend, 1895: 75, Colo. (ex *Vanessa*).

Hypochoaeta eudryae, see *Chaetophlepsis eudryae*

Hypochoaeta longicornis, see *Chaetophlepsis orbitalis* or *Chaetophlepsis* spp.

Hypostena Meigen (Dexiinae: Dexiini)

Hypostena barbata, see *Eucelatoriopsis dimmocki* probably or *Medina barbata*

“*Hypostena floridensis* Townsend”

HOST: HYMENOPTERA

Argidae

Sphacophilus cellularis (Say)

All records ex *Schizocerus ebenus* Norton.

Chittenden, 1901: 107 (ex “*Shizocera*”); Coquillett, 1897: 17, 22, Miss.

Sterictiphora sp.

Johnson, 1913: 70-71, Fla. (ex *Schizocerus*).

HOST: LEPIDOPTERA

Blastobasidae

Valentinia glandulella (Riley)

All records ex *Blastobasis nubilella* Zeller.

Chittenden, 1901: 107; Coquillett, 1897: 17, 24.

Pyralidae

Diatraea saccharalis (Fabricius)

Gowdey, 1926: 81, Jamaica.

Ostrinia obliteralis (Walker)

Chittenden, 1901: 107, D.C., Md. or Va. (ex *Loxostege*).

STAGE PARASITIZED: Larva (Hymenoptera), pupa (Lepidoptera).

Hypostena grisea, see *Lixophaga grisea*

Hypostena tortricis, see *Erynnia tortricis*

Hypostena variabilis, see *Lixophaga variabilis*

Hypostena "variabilis," see *Lixophaga variabilis*

"*Hypostena*" sp.

HOST: COLEOPTERA

Chrysomelidae

Erynephila maritima (LeConte)

Chittenden and Marsh, 1920a: 17 (ex *Monoxia puncticollis* Say).

STAGE PARASITIZED: Larva.

Hypostena sp. probably, see *Lixophaga diatraeae*

Hystricia abrupta, see *Bombyliopsis abrupta*

"*Ictericophyta*" tibialis, see *Ictericophyto tibialis*

Ictericophyto Townsend (Dexiinae: Macquartiini)

Ictericophyto spinosa (Coquillett)

HOST: LEPIDOPTERA

Geometridae

Alsophila pometaria (Harris)

Schaffner and Griswold, 1934: 90, 106.

Brephos infans Möschler

Schaffner and Griswold, 1934: 90, 106.

Erannis tiliaria (Harris)

Raizenne, 1952: 142, Ont.; Schaffner and Griswold, 1934: 97-98, 196.

Noctuidae

Pyreferra hesperidago (Guenée)

Schaffner and Griswold, 1934: 51, 106 (ex *Conistra indirecta* Walker).

Tortricidae

Pandemis limitata (Robinson)

Raizenne, 1952: 208, Ont.

BIOLOGY: Schaffner and Griswold, 1934: (106).

Ictericophyto tibialis (Curran)

HOST: LEPIDOPTERA

Geometridae

Melanolopia imitata (Walker)

Evans, 1962: 597, B.C. (as "*Ictericophyta*").

STAGE PARASITIZED: Larva.

BIOLOGY: Evans, 1962: (597).

"*Icterophyto*" *spinosa*, see *Ictericophyto spinosa*

Isoglossa hastata, see *Eucoronimyia hastata*

"*Jurinea*" *algens*, see *Nowickia algens*

Jurinia algens, see *Nowickia algens*

Jurinia apicifer, see *Archytas apicifer*

Jurinia hystrix, see *Archytas metallicus*

Jurinia metallicus, see *Archytas metallicus* or *Juriniopsis floridensis* probably

Juriniopsis Townsend (Tachininae: Dejeaniini)

Juriniopsis adusta, see *Juriniopsis adusta* or *Juriniopsis floridensis* probably

Juriniopsis adusta (Wulp)

HOST: LEPIDOPTERA

Arctiidae

Ecpantheria deflorata (Fabricius)

Sabrosky, 1969: 82, S.C. (ex *E. scribonia* Stoll).

Hesperiidae

Epargyreus clarus (Cramer)

Sabrosky, 1969: 82, Pa.

Juriniopsis aurifrons Brooks

HOST: LEPIDOPTERA

Arctiidae

Halisidota sp.

Sabrosky, 1969: 82, Nev.

Juriniopsis floridensis Townsend probably

HOST: LEPIDOPTERA

Arctiidae

Ecpantheria deflorata (Fabricius)

Most records ex *Ecpantheria scribonia* Stoll.

*Vide Sabrosky, 1969: 82, Fla. (Coquillett, 1897); *vide Sabrosky, 1969: 82, Fla. (Patton, 1958); Patton, 1958: 39, Fla. (as *J. adusta* (Wulp) ex *E. deflorata*, incomplete rearing — pupa); Coquillett, 1897: 18, 25, Fla. (as *Jurinia metallica* Robineau-Desvoidy).

STAGE PARASITIZED: Larva.

Larvaevoridae, see Tachinidae page 537.

Latreillia bifasciata, see *Belvosia bifasciata*

Latreillia bifasciata or *Willistonina bicincta*, see *Belvosia* sp.

Latreillimylia bifasciata, see *Belvosia bifasciata*

Leschenaultia Robineau-Desvoidy (Goniinae: Harrisini)

Leschenaultia adusta (Loew)

Most records as *Blepharipeza adusta*.

HOST: LEPIDOPTERA

Arctiidae

Arachnis picta Packard

Essig, 1926: 583, Calif.; Coquillett, 1897: 10, 24.

Ecpantheria deflorata (Fabricius)

Daecke, 1916: 286, Pa. (as "*Blepharopeza*").

Estigmene acraea (Drury)

Bryan, D. E., Jackson and Stoner, 1969: 1, 7-8, Ariz. (as *Leschenaultia*, lab.); Stevenson, W. A., Kaufman and Sheets, 1957: 280, Wash. (as *Rileymylia*); Taylor, E. A., 1954: 526-529, Ariz. (as *Rileymylia*); Brooks, 1947: 172, 180, fig. 6 on pl. 14 (as *Rileymylia* ex *E. acraea*); Essig, 1926: 583, Calif. (ex *E. acraea*); Johnson, 1925: 202, Conn., Maine, Mass., N.H., R.I., Vt. (ex *Leucarctia acraea*); *Brauer and Bergenstamm, 1895: 545, 607 (ex *Spilosoma*, Osten Sacken, 1875); Osten Sacken, 1875: 72 (as "*Blepharopeza*" ex *Spilosoma*).

Halisidota argentata Packard

Silver, 1958: 76, B.C. (as "*Rileymylia*"); *Brauer and

Bergenstamm, 1895: 545, 593 (ex "*Halesidota*,"
Townsend, 1893a); Townsend, 1893a: 403-404.

Halisidota caryae (Harris)

Schaffner and Griswold, 1934: 33, 121; Essig, 1926: 583,
Calif.; Tothill, 1913: 69, Ont.

Halisidota harrisii Walsh

Schaffner and Griswold, 1934: 34-35, 121.

Halisidota maculata (Harris)

Schaffner and Griswold, 1934: 33-34, 121; Essig, 1926: 583,
Calif.; Tothill, 1913: 69, Ont.

Halisidota tessellaris (J. E. Smith)

Schaffner and Griswold, 1934: 34, 121.

Halisidota spp.

Winn and Beaulieu, 1915: 142, Que. (ex "*Halesidota*").

Hemihyalea edwardsi (Packard)

Essig, 1926: 583, Calif.; Coquillett, 1897: 10, 25, Calif. (ex
Halisidota).

Unidentified Arctiidae

Johnson, 1925a: 87, New England (probable host).

Lasiocampidae

Malacosoma californicum californicum (Packard)

Coquillett, 1897: 10, 24, Calif. (ex *Clisiocampa thoracica*
Stretch).

Malacosoma constrictum (H. Edwards)

Essig, 1926: 583, Calif.; Coquillett, 1897: 10, 24, Calif. (ex
Clisiocampa).

Malacosoma disstria Hübner

Essig, 1926: 583, Calif.; Tothill, 1913: 69-70, N.B., Ont.

Saturniidae

Coloradia pandora Blake

Patterson, 1929: 17, Oreg.

STAGE PARASITIZED: Larva, pupa.

IMMATURES: Egg — Brooks, 1947: (descr., fig. 6 on pl. 14).

Pupa — Greene, 1922: (descr., fig. 68 on pl. 15); Townsend,
1893a: (descr.).

BIOLOGY: Bryan, D. E., Jackson and Stoner, 1969: (1, 7-8);

Taylor, E. A., 1954: (526-529); Schaffner and Griswold, 1934:
(121); Patterson, 1929: (17).

Leschenaultia americana (Brauer and Bergenstamm)

HOST: LEPIDOPTERA

Arctiidae

Halisidota argentata Packard

Silver, 1958: 76, B.C. (as "*Rileymia*" *triseta* Brooks).

Lasiocampidae

Malacosoma californicum californicum (Packard)

Brauer and Bergenstamm, 1895: 571, 588 (as *Rileya* ex *Clisiocampa thoracica* Stretch).

Malacosoma californicum pluviale (Dyar)

Johansen, 1957: 70, Wash. (as *Rileymia triseta* ex *M. pluviale*).

Malacosoma constrictum (H. Edwards)

Brauer and Bergenstamm, 1895: 571, 588 (as *Rileya* ex *Clisiocampa*).

Malacosoma disstria Hübner

Johansen, 1957: 70, Wash. (as *Rileymia triseta*).

Malacosoma spp. (including *M. californicum pluviale* (Dyar) and/or *M. disstria* Hübner)

Dailey, 1958: 269, Wash. (as *Rileymia* ex western tent caterpillar and/or forest tent caterpillar); Dailey, 1957: 628, Wash. (as *Rileymia* ex *M. disstria* and/or *M. pluviale*).

Malacosoma spp.

Ross, 1953: 20, 23, fig. 9, B.C. (as *Rileymia*); Brooks, 1947: 172, 182, fig. 9 on pl. 14, B.C. (as *Rileymia triseta*).

Noctuidae

Pyrophila sp.

Brauer and Bergenstamm, 1895: 571, 604 (as *Rileya*).

STAGE PARASITIZED: Larva, pupa.

IMMATURES: Egg — Brooks, 1947: (descr., fig. 9 on pl. 14).

Pupa — Ross, 1953: (descr., fig. 9).

Leschenaultia cilipes, see *Leschenaultia leucophrys*

Leschenaultia exul (Townsend)

HOST: LEPIDOPTERA

Lasiocampidae

Malacosoma americanum (Fabricius)

Brooks, 1947: 172, 181, fig. 3 on pl. 14 (as *Rileymia*); Bess, 1937: 411-412 (lab.); Bess, 1936: 594-612, figs. 1-18 on

pls. 1-4; Schaffner and Griswold, 1934: 88, 121 (as *Blepharipeza*).

Malacosoma disstria Hübner

Sippell, 1961: 434, 436, fig. 5, Ont.; McGugan, Haliburton and MacDonald, 1952: 45, Ont.; Brooks, 1947: 172, 181, fig. 3 on pl. 14 (as *Rileymyia*); Bess, 1936: 594-612, figs. 1-18 on pls. 1-4; Schaffner and Griswold, 1934: 89, 121 (as *Blepharipeza*).

Tent caterpillar(-s)

LeRoux, 1961: 114, Que. (as *Rileymyia*).

STAGE PARASITIZED: Larva.

IMMATURES: Egg — Brooks, 1947: (descr., fig. 3 on pl. 14); Bess, 1936: (descr., fig. 1 on pl. 1).

First instar larva — Bess, 1936: (descr., figs. 2-6 on pls. 1-2).

Second instar larva — Bess, 1936: (descr., figs. 7-9 on pl. 2).

Third instar larva — Bess, 1936: (descr., figs. 10-13 on pl. 3).

Pupa — Sippell, 1961: (descr., fig. 5); Bess, 1936: (descr., figs. 14-18 on pl. 4).

BIOLOGY: Bess, 1937: (411-412); Bess, 1936: (594-612); Schaffner and Griswold, 1934: (121).

Leschenaultia fulvipes (Bigot)

HOST: LEPIDOPTERA

Lasiocampidae

Malacosoma californicum (Packard)

Stehr and Cook, 1968: 267, 300, N. Mex.

Malacosoma californicum fragile (Stretch)

Stehr and Cook, 1968: 278, 301, Calif.

Malacosoma californicum ssp. (intermediate between *M. californicum lutescens* (Neumoegen and Dyar) and *M. californicum pluviale* (Dyar))

Stehr and Cook, 1968: 277-278, 301, Mont.

Malacosoma incurvum incurvum (H. Edwards)

Stehr and Cook, 1968: 280, 301, Ariz.

Saturniidae

Hemileuca lucina H. Edwards

Brooks, 1947: 172, 181 (as *Rileymyia*).

Hemileuca maia (Drury)

Brooks, 1947: 172, 181 (as *Rileymyia*).

Hemileuca spp. (including *H. lucina* H. Edwards and/or *H. maia* (Drury))

Schaffner and Griswold, 1934: 29, 122 (as *Blepharipeza*).

STAGE PARASITIZED: Larva.

IMMATURES: Egg — Brooks, 1947: (descr.).

BIOLOGY: Schaffner and Griswold, 1934: (122).

Leschenaultia grossa Brooks

HOST: LEPIDOPTERA

Arctiidae

Unidentified Arctiidae

Brooks, 1947: 172, 178, fig. 7 on pl. 14.

IMMATURES: Egg — Brooks, 1947: (descr., fig. 7 on pl. 14).

Leschenaultia halisidotae Brooks

HOST: LEPIDOPTERA

Arctiidae

Halisidota argentata Packard

Brooks, 1947: 172, 176, fig. 5 on pl. 14.

Halisidota caryae (Harris)

Raizenne, 1952: 22, Ont.; Brooks, 1947: 172, 176, fig. 5 on pl. 14, Ont., Que.

Halisidota harrisii Walsh

Brooks, 1947: 172, 176, fig. 5 on pl. 14.

Halisidota maculata (Harris)

Brooks, 1947: 172, 176, fig. 5 on pl. 14.

Halisidota tessellaris (J. E. Smith)

Raizenne, 1952: 24, Ont., Que. (as "*Leschenaultis*"); Brooks, 1947: 172, 176, fig. 5 on pl. 14.

IMMATURES: Egg — Brooks, 1947: (descr., fig. 5 on pl. 14).

Leschenaultia jurinoides, see *Leschenaultia leucophrys*

Leschenaultia leucophrys (Wiedemann)

HOST: LEPIDOPTERA

Arctiidae

Epantheria icasia (Cramer)

Harrison and Stephens, 1966: 673, Costa Rica.

Estigmene acrea (Drury)

Patton, 1958: 39, Fla.

Euchaetias egle (Drury)

Patton, 1958: 39, Fla.

Lasiocampidae

Malacosoma disstria Hübner

Baird, 1938: 81, 137, 140, B.C. (as *Blepharipeza*); Balch and Simpson, 1935: 36, N.B. (as *Blepharipeza*).

Lymantriidae

Euproctis chrysorrhoea (Linnaeus)

Howard and Fiske, 1911: 93 (as *Blepharipeza*).

Sphingidae

Paonias excaecata (J. E. Smith)

Brooks, 1947: 172, 174, fig. 8 on pl. 14; Schaffner and Griswold, 1934: 21, 122 (as *Blepharipeza*); Aldrich, 1932: 2, N.H. (as *Blepharipeza* ex *P. "excoecata"*).

Sphinx chersis (Hübner)

Brooks, 1947: 172, 174, fig. 8 on pl. 14; Schaffner and Griswold, 1934: 18, 122 (as *Blepharipeza* ex *Hyloicus*); Tothill, 1913: 70, Ont. (as *Blepharipeza*).

Unidentified Sphingidae

Procter, 1946: 433, Maine (as *L. jurinoides* Townsend); Ballou, 1936: 142, Costa Rica (as *L. cilipes* Robineau-Desvoidy ex esfingido from *Annona cherimola* Miller); Ballou, 1936a: 48, Costa Rica (as *L. cilipes* ex esfingido from *Annona cherimola*).

HOST: RODENTIA (MAMMALIA)

Unidentified Rodentia

Mouse (mice) probably

Wolcott, 1951a: 481, P.R.; Wolcott, 1924: 223, P.R. (as *Blepharipeza*).

STAGE PARASITIZED: Larva (Lepidoptera), pupa (Lepidoptera).

IMMATURES: Egg — Brooks, 1947: (descr., fig. 8 on pl. 14).

BIOLOGY: Schaffner and Griswold, 1934: (122).

Leschenaultia spp.

HOST: LEPIDOPTERA

Arctiidae

Halisidota argentata Packard

Guppy, R., 1950: 4, B.C. (as *Rileymyia*).

Halisidota caryae (Harris)

Brooks, 1946: 185, Ont. (as "*Leschnaultia*").

Woolly bear(-s)

Duncan and Pickwell, 1939: 344, fig. 177, [Calif.] (as tachinid fly); Duncan, 1933: 240, fig. 123, [Calif.] (as tachinid fly).

Lasiocampidae

Malacosoma disstria Hübner

Tothill, 1923: 180, Canada (as *Blepharipeza*); Tothill, 1919: 12, N.B. (as *Blepharipeza*).

Malacosoma sp.

Guppy, R., 1950: 4 (as *Rileymyia*).

Lymantriidae

Hemerocampa leucostigma (J. E. Smith)

Schaffner and Griswold, 1934: 85 (as *Blepharipeza*).

Unidentified Lepidoptera

Miscellaneous

Townsend, 1893c: 314, Jamaica (as *Blepharipeza* sp. ? ex moth allied to "*Euchaetes*," but with an *Orgyia*-like larva).

STAGE PARASITIZED: Larva, pupa.

IMMATURES: Third instar larva — Duncan and Pickwell, 1939: (fig. 177); Duncan, 1933: (fig. 123).

Pupa — Duncan and Pickwell, 1939: (fig. 177); Duncan, 1933: (fig. 123).

BIOLOGY: Guppy, R., 1950: (4).

"*Leschenaultis*" *halisidotae*, see *Leschenaultia halisidotae*

"*Leschnaultia*" sp., see *Leschenaultia* spp.

Leskia Robineau-Desvoidy (Dexiinae: Leskiini)

"*Leskia analis* Say"

HOST: LEPIDOPTERA

Pyralidae

Diaphania hyalinata (Linnaeus)

Wolcott, 1924: 221, P.R.

Leskia gilensis, see *Sipholeskia occidentalis*

Leskiella James (Dexiinae: Leskiini)

Leskiella brevirostris James

HOST: LEPIDOPTERA

Olethreutidae

- Laspeyresia caryana* (Fitch)
Phillips, A. M., 1957: 89, Fla. (ex hickory shuckworm).
Laspeyresia pomonella (Linnaeus)
James, M. T., 1947: 98, Ill. (ex *Carpocapsa*).
Unidentified Olethreutidae
Krombein, 1967: 69, Fla.
STAGE PARASITIZED: Larva.
Leskiella brevis James probably
HOST: LEPIDOPTERA

Olethreutidae

- Grapholitha molesta* (Busck)
*Vide James, M. T., 1947: 98, 103, Md. (Haeussler, 1930);
Haeussler, 1930: 367, Md. (as *Leskiomima tenera*
(Wiedemann) ex *Laspeyresia*).
STAGE PARASITIZED: Larva.
Leskiomima Brauer and Bergenstamm (Dexiinae: Leskiini)
Leskiomima tenera, see *Leskiomima brevis* probably or
Leskiomima tenera
Leskiomima tenera (Wiedemann)
HOST: LEPIDOPTERA

Olethreutidae

- Eucosma* sp.
Ciesla, Rauschenberger and Bell, 1968: 1130, S.C. (as
"Leskiomina").
Grapholitha molesta (Busck)
All records ex *Laspeyresia molesta*.
McConnell, 1932: 367-368, Md.; Stearns, 1928: 7, Md.;
Garman, P., 1918: 117, Md.
Laspeyresia pomonella (Linnaeus)
James, M. T., 1947: 103, Ill. (ex *Carpocapsa*).

Pyralidae

- Acrobasis indigenella* (Zeller)
Most records ex *Acrobasis nebulella* Riley.
Allen, 1929: 683, Miss. (ex *Mineola* "indigenella" *nebulella*,
probable host); Gill, 1917: 15, Fla.; Gill, 1917a: 21.
Acrobasis juglandis (LeBaron)
James, M. T., 1947: 103, Fla., Miss.

Desmia funeralis (Hübner)

James, M. T., 1947: 103, D.C.; Strauss, 1916: 9, fig. k on pl. 4, D.C.

Dioryctria abietella (Denis and Schiffermüller)

Ebel, 1965: 629, Fla. (as "*Leskiomina*").

Homoeosoma electellum (Hulst)

Satterthwait and Swain, 1946: 577-578, Mo.

Macrobotys pertextalis (Lederer)

Schaffner, 1959: 10, 51, [N.E. U.S.].

Tetralopha sp. near *asperatella* (Clemens)

Schaffner, 1959: 13, 51, [N.E. U.S.].

Unidentified Pyralidae

Coquillett, 1897: 18, 27, Va.

STAGE PARASITIZED: Larva.

IMMATURES: Pupa — Satterthwait and Swain, 1946: (descr.); Greene, 1922: (descr., fig. 42 on pl. 9); Strauss, 1916: (fig. k on pl. 4).

BIOLOGY: Schaffner, 1959: (10, 13, 51); Satterthwait and Swain, 1946: (577-578).

Leskiomima sp.

HOST: LEPIDOPTERA

Pyralidae

Dioryctria reniculella (Grote)

Daviault, 1946: 117, Que. (as "*Leskiomina*").

"*Leskiomina*" tenera, see *Leskiomima tenera*

"*Leskiomina*" sp., see *Leskiomima* sp.

Leskiopalpus depilis, see *Myobiopsis depilis*

Leskiopalpus flavipennis, see *Myobiopsis flavipennis*

Lespesia Robineau-Desvoidy (Goniinae: Eryciini)

Lespesia "achaetoneura (Williston)," see *Lespesia* spp.

Lespesia aletiae (Riley)

HOST: COLEOPTERA

Coccinellidae

Epilachna varivestis Mulsant

Beneway, 1963: 638 (ex *E. corrupta* Mulsant).

HOST: LEPIDOPTERA

Arctiidae

Diacrisia virginica (Fabricius)

Beneway, 1963: 637-638.

Estigmene acrea (Drury)

Beneway, 1963: 638; Patton, 1958: 31, Fla. (as *Achaetoneura*).

Estigmene sp.

Beneway, 1963: 638.

Halisidota harrisii Walsh

Beneway, 1963: 638; Schaffner and Griswold, 1934: 34-35, 116 (as *Achaetoneura*).

Halisidota maculata (Harris)

Essig, 1926: 582, Calif. (as *Frontina*); Coquillett, 1897: 15, 25, Calif. (as *Frontina*); Howard, 1897b: 41, D.C. (as *Frontina*).

Halisidota tessellaris (J. E. Smith)

Most records ex *Halisidota* "tessellata."

Schaffner and Griswold, 1934: 34, 116 (as *Achaetoneura* ex *H. tessellaris*); Webber, 1930: 4 (as *Achaetoneura* ex "Halisidota" *tessellaris*); Johnson, 1910: 781, N.J. (as *Frontina*); Coquillett, 1897: 15, N.Y. (as *Frontina*); Howard, 1897b: 41, D.C. (as *Frontina*).

Halisidota sp.

Beneway, 1963: 638.

Hyphantria cunea (Drury)

Beneway, 1963: 638; Fattig, 1949: 4, Ga. (as *Achaetoneura* "aletia"); Brimley, 1938: 363, N.C. (as *Achaetoneura*); Schaffner and Griswold, 1934: 37, 116 (as *Achaetoneura*); Webber, 1930: 4 (as *Achaetoneura*); Brimley, 1922: 23, N.C. (as *Frontina* ex fall-webworm).

Hyphantria sp.

Beneway, 1963: 638.

Seirarctia echo (J. E. Smith)

Beneway, 1963: 638.

Utetheisa ornatrix bella (Linnaeus)

Beneway, 1963: 638 (ex *U. bella*); Patton, 1958: 31, Fla. (as *Achaetoneura* ex *bella* moth).

Unidentified Arctiidae

Webber, 1930: 4 (as *Achaetoneura*).

Brassolidae

Opsiphanes tamarindi Felder

Harrison, 1963: 282-283, Costa Rica (as *Achaetoneura*);
Harrison, 1963a: 92, Costa Rica (as *Achaetoneura*).

Ctenuchidae

Ceramidia butleri Möschler

Harrison, 1963: 282-283, Costa Rica (as *Achaetoneura*).

Syntomeida epilais (Walker)

Beneway, 1963: 637; Patton, 1958: 32, Fla. (as *Achaetoneura*).

Geometridae

Ennomos subsignarius (Hübner)

Davis, R., 1961: 248, Ga. (as *Achaetoneura*).

Hesperiidae

Autochton cellus (Boisduval and LeConte)

Clark, 1936: 47 (as *Achaetoneura* ex *Rhabdoides*).

Epargyreus clarus (Cramer)

All records ex *Epargyreus tityrus* (Fabricius).

Beneway, 1963: 638; Schaffner and Griswold, 1934: 15, 116 (as *Achaetoneura*); Webber, 1930: 4 (as *Achaetoneura*); Allen, 1926b: 189-191, pl. 9, Miss. (as *Achaetoneura*).

Urbanus proteus (Linnaeus)

Beneway, 1963: 638; Patton, 1958: 32, Fla. (as *Achaetoneura*).

Lasiocampidae

Malacosoma americanum (Fabricius)

Beneway, 1963: 638; Patton, 1958: 31, Fla. (as *Achaetoneura*); Dozier, 1920: 372, 378, Fla. (as *Frontina*).

Limacodidae

Sibine apicalis Dyar

Harrison, 1963: 282-283, Costa Rica (as *Achaetoneura*);
Harrison, 1963a: 94, Costa Rica (as *Achaetoneura* ex
"Sabine").

Lymantriidae

Hemerocampa leucostigma (J. E. Smith)

Beneway, 1963: 638; Raizenne, 1952: 95, Ont. (as *Achaetoneura*); Dowden, 1935: 509 (as *Achaetoneura*); Schaffner and Griswold, 1934: 85, 116 (as *Achaetoneura*); Webber, 1930: 4 (as *Achaetoneura*); Webber and Schaffner, 1926: 12-13, Mass., New England, N.Y., Pa., R.I. (as *Frontina*); Britton, 1917: 110 (as *Frontina*); Johnson, 1910: 781, N.J. (as *Frontina* ex *Orgyia*); Titus and Pratt, 1904: 28 (as *Frontina*); Howard, 1898: 88, D.C. (as *Frontina* ex *Orgyia*); Coquillett, 1897: 15, 26, D.C. (as *Frontina* ex *Orgyia*); Howard, 1897a: 16, D.C. (as *Frontina* ex *Orgyia*); Howard, 1897b: 7, 41-42, 49, D.C. (as *Frontina* ex *Orgyia*).

Orgyia sp.

Beneway, 1963: 638 (ex *Orgyia*).

Porthetria dispar (Linnaeus)

Webber, 1930: 4 (as *Achaetoneura*).

Stilpnotia salicis (Linnaeus)

Webber, 1930: 4 (as *Achaetoneura*).

Tussock moth(-s)

Johnson, 1925: 199, Conn., Maine, Mass., R.I., Vt. (as *Frontina*); Thompson, 1910: 291 (as *Frontina*).

Megalopygidae

Megalopyge opercularis (J. E. Smith)

Beneway, 1963: 638; Patton, 1958: 31, Fla. (as *Achaetoneura*); Allen, 1926b: 188-191, pl. 9, Miss. (as *Achaetoneura*); Coquillett, 1897: 15, 25, Tex. (as *Frontina* ex *Lagoa*); Howard, 1897b: 41, D.C. (as *Frontina* ex *Lagoa*).

Megalopyge pyxidifera (J. E. Smith)

Patton, 1958: 31, Fla. (as *Achaetoneura*).

Megalopyge sp.

Beneway, 1963: 638.

Noctuidae

Alabama argillacea (Hübner)

Beneway, 1963: 638; Webber, 1930: 4 (as *Achaetoneura*); Allen, 1926b: 188-191, pl. 9, Miss. (as *Achaetoneura* ex

A. "argillicea"); Coquillett, 1897: 15, 23, Fla. (as *Frontina ex Aletia*); Howard, 1897b: 41, Ala., D.C., Ga., Miss., Tex. (as *Frontina ex Aletia*); *Brauer and Bergenstamm, 1895: 575, 581 (as *Tachina ex Aletia*, Riley, 1879); Riley, 1885: 101, 109, 377, Ala., Ga., Miss., Tex. (as *Tachina ex Aletia xylina* (Say)); Riley, 1885: 101, 109 (as *Tachina fraterna* Comstock ex *Aletia xylina*); Riley, 1880: 93-94 (as *Tachina ex cotton worm*); Comstock, J. H., 1879: 203-204, Ala. (as *Tachina ex Aletia*); Riley, 1879: 161-162 (as *Tachina ex Aletia*).

Heliothis zea (Boddie)

Most records ex *Heliothis obsoleta* (Fabricius).

Allen, 1926b: 188-191, pl. 9, Miss. (as *Achaetoneura*); Essig, 1926: 582 (as *Frontina*); Gowdey, 1923a: 54, Jamaica (as *Frontina "aletinae" ex Chloridea*); Quaintance and Brues, 1905: 123 (as *Frontina*); Riley, 1885: 377, Ala. (as *Tachina ex H. armigera* (Hübner)).

Heliothis sp.

Beneway, 1963: 638.

Mocis latipes (Guenée) probably

Patton, 1958: 32, Fla. (as *Achaetoneura ex M. repanda* (Fabricius)).

Plathypena scabra (Fabricius)

Fattig, 1949: 4 (as *Achaetoneura "aletia"*); Brimley, 1938: 363, N.C. (as *Achaetoneura*); Brimley, 1922: 23, N.C. (as *Frontina ex green clover worm*); Sherman, 1920: 298-299, N.C. (as *Frontina*).

Pseudaletia unipuncta (Haworth)

Burrell, 1967: 114, La. (as *Achaetoneura*); Beneway, 1963: 638; Breeland, 1958: 320, 323, 333, Tenn. (as *Achaetoneura*); Poos, 1928: 146, Va. (as *Achaetoneura ex Cirphis*); King, V., and Barber, 1921: 487, Mo. (as *Frontina ex Heliophila*, probable host).

Pseudoplusia includens (Walker)

Allen, 1926b: 188-191, pl. 9, Miss. (as *Achaetoneura ex Autographa oo* (Cramer)).

Spodoptera frugiperda (J. E. Smith)

All records ex *Laphygma frugiperda*.

Beneway, 1963: 638; Allen, 1926b: 189-191, pl. 9, Miss. (as *Achaetoneura*); Gowdey, 1926: 82, Jamaica (as *Frontina*); Gowdey, 1923a: 20, Jamaica (as *Frontina*); Gowdey, 1921: 4, Jamaica (as *Frontina*).

Spodoptera ornithogalli (Guenée)

Bottrell, 1969: 251, Okla. (ex *Prodenia*).

Trichoplusia ni (Hübner)

Beneway, 1963: 638.

Xanthopastis timais (Cramer)

Beneway, 1963: 638; Patton, 1958: 32, Fla. (as *Achaetoneura* ex "*Xanthopastes*").

Notodontidae

Cerura spp.

Coquillett, 1897: 15, 24, Calif. (as *Frontina*); Howard, 1897b: 41, D.C. (as *Frontina*).

Dasylophia anguina (J. E. Smith)

Johnson, 1910: 781, N.J. (as *Frontina*); Howard, 1897b: 41, D.C. (as *Frontina* ex "*Dasylopha*").

Datana ministra (Drury)

Beneway, 1963: 638; Patton, 1958: 32, Fla. (as *Achaetoneura*).

Heterocampa manteo (Doubleday)

Howard, 1897b: 41, D.C. (as *Frontina*).

Nymphalidae

Asterocampa sp.

Beneway, 1963: 638 (ex *Apatura*).

Nymphalis antiopa (Linnaeus)

Beneway, 1963: 638; Schaffner and Griswold, 1934: 11, 116 (as *Achaetoneura* ex *Hamadryas*); Webber, 1930: 4 (as *Achaetoneura* ex *Euvanessa*); Allen, 1926b: 188-191, pl. 9, Miss. (as *Achaetoneura* ex *Aglais*).

Polygonia interrogationis (Fabricius)

Beneway, 1963: 638.

Vanessa cardui (Linnaeus)

Beneway, 1963: 638.

Pieridae

Pieris protodice Boisduval and LeConte

Beneway, 1963: 638.

Pieris rapae (Linnaeus)

Beneway, 1963: 638.

Pyralidae

Evergestis rimosalis (Guenée)

Beneway, 1963: 638.

Olycella junctolineella (Hulst)Mann, 1969: 14, 39; Hamlin, 1925: 10-11, Tex. (as *Frontina* ex *Olyca*).

Saturniidae

Hemileuca maia (Drury)Webber, 1930: 4 (as *Achaetoneura*).

Sphingidae

Ceratonia catalpae (Boisduval)

Beneway, 1963: 638.

Pholus sp.Webber, 1930: 4 (as *Achaetoneura*).

Unidentified Lepidoptera

Miscellaneous

Wolcott, 1951a: 479, P.R. (as *Achaetoneura* ex chrysalis on *Inga laurina* (Swartz) Willdenow); Essig, 1926: 582, U.S. (as *Frontina* ex many moths); Wolcott, 1924: 222, P.R. (as *Frontina* ex larva on *Inga laurina*).

STAGE PARASITIZED: Larva (Lepidoptera), pupa (Lepidoptera).

IMMATURES: Egg — Allen, 1926b: (descr., pl. 9).

First instar larva — Allen, 1926b: (descr., pl. 9).

Second instar larva — Allen, 1926b: (descr., pl. 9).

Third instar larva — Allen, 1926b: (descr.).

Pupa — Allen, 1926b: (descr.); Greene, 1922: (descr., fig. 30 on pl. 7).

BIOLOGY: Patton, 1958: (31); Clark, 1936: (47); Dowden, 1935: (509); Schaffner and Griswold, 1934: (116); Allen, 1926b: (188-191); Hamlin, 1925: (10-11); Thompson, 1910: (291); Howard, 1897b: (7, 41-42, 49); Riley, 1885: (101, 109, 377); Comstock, J. H., 1879: (203-204).

Lespesia aletiae (Riley) probably

HOST: LEPIDOPTERA

Ctenuchidae

Lymire edwardsii (Grote)

Genung, 1959: 40, 41, Fla. (as *Achaetoneura*).

STAGE PARASITIZED: Pupa.

Lespesia anonyma, see *Lespesia* spp.

Lespesia "archaetoneura," see *Lespesia* spp.

Lespesia archippivora (Riley)

HOST: HYMENOPTERA

Tenthredinidae

Pristiphora erichsonii (Hartig)

Beneway, 1963: 640.

HOST: LEPIDOPTERA

Arctiidae

Estigmene acrea (Drury)

Bryan, D. E., Jackson and Patana, 1968: 819-823 (lab.);
Beneway, 1963: 640; Butler, 1958: 562, Ariz. (as
Achaetoneura); Taylor, E. A., 1954: 526, Ariz. (as
Achaetoneura).

Estigmene sp.

Beneway, 1963: 640.

Euchaetias egle (Drury)

Webber, 1930: 4 (as *Achaetoneura*); Townsend, 1893: 467,
Ill. (as *Prospberrya promiscua* (Townsend) ex
"Euchaetes").

Halisidota caryae (Harris)

Webber, 1930: 4 (as *Achaetoneura*).

Tyria jacobaeae (Linnaeus)

Hawkes, 1968: 500, Calif.

Citheroniidae

Citheronia regalis (Fabricius)

Riley, 1871: 151 (as *Tachina*).

Danaiidae

Danaus gilippus berenice (Cramer)

Beneway, 1963: 640 (ex *D. berenice*).

Danaus plexippus plexippus (Linnaeus)

Weaver, 1969: 71, W. Va. (ex *D. plexippus*); Anon., 1968: 85, Hawaii (ex *D. plexippus*); Hale, 1967: 180, Hawaii (ex *D. plexippus*); Brewer and Thomas, 1966: 235, Mass. (ex *D. plexippus*); Mitchell, 1966: 129, Hawaii (as *Achaetoneura* ex *D. plexippus*); Mitchell, 1965: 525, Hawaii (as *Achaetoneura* ex *D. plexippus*); Beneway, 1963: 640 (ex *D. plexippus*); Urquhart, 1960: 207-208, Ont. (as *Achaetoneura* ex *D. plexippus*); Zimmerman, 1958: 473, Hawaii (as *Frontina* ex *D. plexippus*); Teale, 1948: 356-359, 4 unnumbered figs. on p. 358, E. U.S. (as *Achaetoneura* ex monarch); Teale, 1946: 25, New England (as *Achaetoneura* ex monarch butterfly); Schaffner and Griswold, 1934: 8, 116 (as *Achaetoneura* ex *D. menippe* (Hübner)); Aldrich, 1932: 1, Mass. (as *Achaetoneura* ex *D. archippus* (Fabricius)); Williams, 1931: 292, Hawaii (as *Frontina* ex *D. archippus*); Webber, 1930: 4 (as *Achaetoneura* ex *Anosia plexippus*); Swezey, 1929: 211, Hawaii (as *Frontina* ex *D. archippus*); Essig, 1926: 582, Colo., Idaho, Wash. (as *Frontina* ex *D. menippe*); Webber and Schaffner, 1926: 17, 23, New England (as *Frontina* ex *Anosia plexippus*); Johnson, 1925: 200, Conn., Maine, Mass., N.H., R.I. (as *Frontina* ex monarch butterfly); Johannsen, 1910: 38, Maine (as *Frontina* ex *Anosia plexippus*); Swezey, 1907: 48, Hawaii (as *Frontina* ex *Anosia plexippus*); Coquillett, 1897: 15, 24, Pa., Wash. (as *Frontina* ex "*Danais*" *archippus*); *Brauer and Bergenstamm, 1895: 560, 582, N. Amer. (as (*Masicera*) ex *Anosia plexippus*, Williston, 1889); Townsend, 1893: 467, Ill. (as *Masicera* ex "*Danais*" *archippus*); Williston, 1889: 1914, 1923 (as *Masicera* ex *Anosia plexippus*); Riley, 1873: 149 (as *Tachina* ex "*Danais*" *archippus*); Riley, 1871: 116, 149-150, Mo., Pa. (as *Tachina* ex "*Danais*" *archippus*).

Geometridae

Anacamptodes fragilaria (Grossbeck)

Van Zwaluwenburg, 1946: 13, Hawaii (as *Frontina* ex *Anacamptodes* [sp.]).

Scotorythra paludicola (Butler)

Zimmerman, 1958: 125, Hawaii (as *Frontina*); Swezey, 1926a: 360, Hawaii (as *Frontina*).

Scotorythra rara (Butler)

Swezey, 1907: 48, Hawaii (as *Frontina*).

Lasiocampidae

Malacosoma americanum (Fabricius)

Stehr and Cook, 1968: 258, 299-300, Tex.; Beneway, 1963: 640; Bess, 1936: 605 (as *Achaetoneura*); Schaffner and Griswold, 1934: 88, 116 (as *Achaetoneura*); Webber, 1930: 4 (as *Achaetoneura*); West, 1928: 815, N.Y. (as *Frontina*); Townsend, 1916c: 75-76, N.C. (as *Ypophaemyia malacosomae* Townsend).

Malacosoma californicum (Packard)

Stehr and Cook, 1968: 259-261, 267, 299-300, Calif., Colo.

Malacosoma californicum ambisimile (Dyar)

Stehr and Cook, 1968: 271, 299-300, Calif.

Malacosoma californicum californicum (Packard)

Stehr and Cook, 1968: 271, 299-300, Calif.

Malacosoma californicum fragile (Stretch)

Stehr and Cook, 1968: 278-279, 299, 301, Ariz., Calif., Nev., Utah.

Malacosoma californicum pluviale (Dyar)

Stehr and Cook, 1968: 275, 299-300, Mont., Wash.; Beneway, 1963: 640 (ex *M. pluvialis*); Essig, 1926: 582, Calif. (as *Frontina* ex *M. pluvialis*); Coquillett, 1897: 15, 24, Wash. (as *Frontina* ex *Clisiocampa pluvialis*).

Malacosoma californicum recenseo Dyar

Stehr and Cook, 1968: 273, 299-300, Calif.

Malacosoma californicum ssp. (intermediate between *M. californicum ambisimile* (Dyar) and *M. californicum californicum* (Packard))

Stehr and Cook, 1968: 272-273, 299-300, Calif.

Malacosoma constrictum (H. Edwards)

Beneway, 1963: 640; Essig, 1926: 582, Colo., Idaho, Wash. (as *Frontina*); Coquillett, 1897: 15, 24, Calif. (as *Frontina* ex *Clisiocampa*).

Malacosoma disstria Hübner

Beneway, 1963: 640; Sippell, 1961: 436, Ont. (as *Achaetoneura*); *Brauer and Bergenstamm, 1895: 542, 588, Maine (as *Achaetoneura promiscua* ex *Clisiocampa sylvatica* (Harris), Townsend, 1891a); Townsend, 1891a: 85, Maine (as *Phorocera promiscua* ex *Clisiocampa sylvatica*).

Malacosoma incurvum discoloratum (Neumoegen)

Stehr and Cook, 1968: 280, 299, 301, Nev.

Malacosoma incurvum incurvum (H. Edwards)

Stehr and Cook, 1968: 279-280, 299, 301, Ariz.

Malacosoma spp.

Beneway, 1963: 640; Gibson, W. W., and Carrillo S., 1959: 180, Mexico (D.F.) (as *Achaetoneura* ex "*Malocosoma*"); *Brauer and Bergenstamm, 1895: 560, 588, N. Amer. (as (*Masicera*) ex *Clisiocampa*, Townsend, 1893); Townsend, 1893: 467, Ill. (as *Masicera* ex *Clisiocampa*).

Tent caterpillar(-s)

Brimley, 1922: 26, N.C. (as *Ypophemyia malacosomae*).

Lycaenidae

Lampides boeticus (Linnaeus)

Zimmerman, 1958: 490, Hawaii (as *Frontina*); Williams, 1931: 292, Hawaii (as *Frontina* ex *Polyommatus*); Swezey, 1922: 8-9, Hawaii (as *Frontina* ex *Lycaena*).

Noctuidae

Agrotis crinigera (Butler)

Zimmerman, 1958: 237, Hawaii (as *Frontina*); Swezey, 1907: 48, Hawaii (as *Frontina*).

Agrotis dislocata (Walker)

Zimmerman, 1958: 239, Hawaii (as *Frontina*); Swezey, 1907: 48, Hawaii (as *Frontina*).

Agrotis ipsilon (Hufnagel)

All records ex *Agrotis ypsilon*.

Beneway, 1963: 640; *Crumb, 1929: 40, 59 (as *Frontina*, Coquillett, 1897); Essig, 1926: 582, Calif. (as *Frontina*); Coquillett, 1897: 15, 23 (as *Frontina*).

Alypia octomaculata (Fabricius)

Schaffner and Griswold, 1934: 41, 116 (as *Achaetoneura*); Webber, 1930: 4 (as *Achaetoneura*).

Anomis hawaiiensis (Butler)

Zimmerman, 1958: 381, Hawaii (as *Frontina*).

Anomis noctivolans (Butler)

Zimmerman, 1958: 385, Hawaii (as *Frontina*).

Autographa californica (Speyer)

Seidel, 1954: 186, Wash. (as "*Achaetoneura*").

Elydna nonagrica (Walker)

Fullaway, 1914: 8, Hawaii (as *Frontina* ex *Caradrina reculsa* Hampson).

Euxoa vestigialis (Rottenburg)

Beneway, 1963: 640.

Feltia sp. near *subgothica* (Haworth)

Riley, 1871: 151 (as *Tachina* ex *Agrotis*).

Feltia subterranea (Fabricius)

Beneway, 1963: 640; Wolcott, 1951a: 479, P.R. (as *Achaetoneura*).

Heliothis hawaiiensis (Quaintance and Brues)

Zimmerman, 1958: 211, Hawaii (as *Frontina*).

Heliothis zea (Boddie)

Bottrell, Young, Price and Adams, 1968: 1054, Okla.; Bryan, D. E., Jackson and Patana, 1968: 819-823 (lab.); Bosch and Hagen, 1966: 25, Calif. (as *Achaetoneura*); Zimmerman, 1958: 214, Hawaii (as *Frontina*); Bibby, 1942: 943, Tex. (as "*Achaetonura*" ex *H. armigera* (Hübner)); Swezey, 1937: 359, Hawaii (as *Frontina* ex *Chloridea obsoleta* (Fabricius)); Swezey, 1923: 184-185, Hawaii (as *Frontina* ex *H. obsoleta*); Fullaway, 1914: 13, Hawaii (as *Frontina* ex *H. obsoleta*).

Heliothis spp. (including *H. hawaiiensis* (Quaintance and Brues) and/or *H. zea* (Boddie))

Zimmerman and Fletcher, 1956: 173, Hawaii (as *Frontina*).

Heliothis spp.

Bottrell, Young, Price and Adams, 1968: 1054, Okla.; Butler, 1958: 562, Ariz. (as *Achaetoneura*).

Peridroma coniotis (Hampson)

Look, 1948: 222, Hawaii (as *Frontina* ex *Agrotis*).

Peridroma coniotis coniotis (Hampson)

Zimmerman, 1958: 283, Hawaii (as *Frontina*).

Peridroma saucia (Hübner)

Swezey, 1935a: 93, Hawaii (as *Frontina* ex *Lycophotia margaritosa* (Haworth)).

Polydesma umbricola Boisduval

Kamran, 1968: 1011, Hawaii (as *Achaetoneura*).

Pseudaletia unipuncta (Haworth)

Breeland, 1958: 320, 323, 334, Nebr. (as *Achaetoneura*); Zimmerman, 1958: 305, Hawaii (as *Frontina*); Swezey, 1935a: 93, Hawaii (as *Frontina* ex *Cirphis*); Pemberton, 1923: 258, Hawaii (as *Frontina* ex *Cirphis*); Swezey, 1907: 48, Hawaii (as *Frontina* ex *Leucania*).

Scotogramma trifolii (Rottenburg)

Beneway, 1963: 640.

Spodoptera exempta (Walker)

All records ex *Laphygma exempta*.

Bianchi, 1958: 319, Hawaii (as *Achaetoneura*); Bianchi, 1949a: 345, Hawaii (as *Frontina*); Pemberton, 1948b: 185, 200, Hawaii (as *Frontina*).

Spodoptera exigua (Hübner)

Bryan, D. E., Jackson and Patana, 1969: 765-767, Ariz. (lab.); Bryan, D. E., Jackson and Stoner, 1969: 1, 4-5, 12, Ariz. (lab.); Bryan, D. E., Jackson and Patana, 1968: 819-823, Ariz. (lab.); Bosch and Hagen, 1966: 24, Calif. (as *Achaetoneura*); Beneway, 1963: 640 (ex *Laphygma*); Butler, 1958: 562, Ariz. (as *Achaetoneura* ex *Laphygma*); Zimmerman, 1958: 340, Hawaii (as *Frontina*); Swezey, 1935a: 93, Hawaii (as *Frontina*); Luginbill, 1928: 82, Idaho (as *Frontina* ex *Laphygma flavimaculata* Harvey); Essig, 1926: 582, Calif. (as *Frontina* ex *Laphygma*); Swezey, 1907: 48, Hawaii (as *Frontina*); Chittenden, 1902: 45, Calif. (as *Frontina* ex *Laphygma*); Coquillett, 1897: 15, 25, Calif. (as *Frontina* ex *Laphygma flavimaculata*).

Spodoptera frugiperda (J. E. Smith)

Most records ex *Laphygma frugiperda*.

Scaramuzza and Barry, 1960: 997, Cuba (as *Achaetoneura*); Butler, 1958: 562, Ariz. (as *Achaetoneura*); Painter, 1955: 39, 41, Guatemala (as *Achaetoneura*); Vickery, 1929: 1, 16, 20-21, 45, Tex. (as *Frontina*); Luginbill, 1928: 82, Tex. (as *Frontina*); Vickery, 1925: 140, Tex. (as *Frontina*); Wolcott, 1924: 222, P.R. (as *Frontina*); Jones, T.H., and Wolcott, 1922: 47, P.R. (as *Frontina*); Jones, T. H., 1913: 235, P.R. (as *Frontina*); Van Dine, 1913b: 202, P.R. (as *Frontina*); Riley, 1871: 116, 150-151 (as *Tachina* ex *Prodenia autumnalis* (Riley)).

Spodoptera mauritia (Boisduval)

*Vide pers. comm. Todd, 1970: [Hawaii] (Clagg, 1962); Einmo, 1963: 196, Hawaii (as *Achaetoneura*); Clagg, 1962: 107-108, Hawaii (as *Achaetoneura* ex *S. mauritia acronyctoides* (Guenée)); Beardsley, 1955: 386, Hawaii (as *Achaetoneura*); Swezey, 1935a: 93, Hawaii (as *Frontina*); Swezey, 1930: 280, Hawaii (as *Frontina*); Swezey, 1907: 48, Hawaii (as *Frontina*).

Spodoptera ornithogalli (Guenée)

All records ex *Prodenia ornithogalli*.

Bottrell, 1969: 251, Okla.; Butler, 1958: 562, Ariz. (as *Achaetoneura*).

Spodoptera praefica (Grote)

Beneway, 1963: 640 (ex *Prodenia*).

Spodoptera spp.

Beneway, 1963: 640 (ex *Prodenia*).

Trichoplusia ni (Hübner)

Brubaker, 1968: 307, Ariz.; Bryan, D. E., Jackson and Patana, 1968: 819-823 (lab.); Butler, 1958: 562, Ariz. (as *Achaetoneura*).

Armyworms (including *Pseudaletia unipuncta* (Haworth) and/or *Spodoptera mauritia* (Boisduval))

All records ex armyworms (*Cirphis unipuncta* and/or *Spodoptera mauritia*).

Swezey, 1931c: 326, Hawaii (as *Frontina*); Swezey, 1928: 673, Hawaii (as *Frontina*).

Armyworm(-s)

Bianchi, 1940: 383, Hawaii (as *Frontina*); Holdaway and Suehiro, 1940: 37, Hawaii (as *Frontina*); Bryan, 1934: 415, Hawaii (as *Frontina*); Swezey, 1931b: 427, Hawaii (as *Frontina*); Swezey, 1927a: 503, Hawaii (as *Frontina*); Swezey, 1927b: 217, Hawaii (as *Frontina*); Swezey, 1925: 372, Hawaii (as *Frontina*); Swezey, 1923a: 303, Hawaii (as *Frontina*); Swezey, 1909: 26, Hawaii (as *Frontina*); Swezey, 1908b: 16-17, Hawaii (as *Frontina*).

Cutworms (including *Agrotis* spp. and/or *Leucania* spp.)

Swezey, 1908b: 16-17, Hawaii (as *Frontina*).

Cutworm(-s)

Bianchi, 1940: 383, Hawaii (as *Frontina*); Illingworth, 1923: 277, Hawaii (as *Frontina*); Fullaway, 1914: 8, Hawaii (as *Frontina*); Swezey, 1909: 26, Hawaii (as *Frontina*); Lintner, 1893: 238, N.Y. (as "*Masicora*").

Noctuid spp. (including *Mamestra configurata* Walker and/or *Peridroma saucia* (Hübner))

*Vide pers. comm. Handford, 1970: [B.C.] (Handford, 1962); Anon., 1962: 121, B.C. (as *Achaetoneura*); Handford, 1962: 19, B.C. (as *Achaetoneura*).

Noctuidae and/or Pyralidae

Noctuid and/or pyralid spp. (including *Scotogramma trifolii* (Rottenburg) and/or *Loxostege sticticalis* (Linnaeus))

Anon., 1956d: 1044, Idaho (as *Achaetoneura* "*achippivora*" ex beet webworm and/or clover cutworm).

Notodontidae

Schizura concinna (J. E. Smith)

Beneway, 1963: 640.

Nymphalidae

Anaea glycerium (Doubleday)

Riley, 1873: 149 (as *Tachina* ex *Paphia* [sp.]).

Nymphalis antiopa (Linnaeus)

Beneway, 1963: 640; Schaffner and Griswold, 1934: 11, 116 (as *Achaetoneura* ex *Hamadryas*); Webber, 1930: 4 (as *Achaetoneura* ex *Euvanessa*); Luginbill, 1928: 82, Idaho (as *Frontina* ex *Vanessa*); Johnson, 1925: 200, Conn., Maine, Mass., N.H., R.I. (as *Frontina* ex mourning-cloak).

Nymphalis milberti (Godart)

Schaffner and Griswold, 1934: 11, 116 (as *Achaetoneura* ex *Hamadryas*); Webber, 1930: 4 (as *Achaetoneura* ex *Aglais*).

Vanessa atalanta (Linnaeus)

Beneway, 1963: 640; Schaffner and Griswold, 1934: 12, 116 (as *Achaetoneura* ex *Cynthia*); Webber, 1930: 4 (as *Achaetoneura*); Webber and Schaffner, 1926: 17, New England (as *Frontina*).

Vanessa cardui (Linnaeus)

Beneway, 1963: 640; Butler, 1958: 562, Ariz. (as *Achaetoneura*); Zimmerman, 1958: 457, Hawaii (as *Frontina*); Swezey, 1947: 5, Hawaii (as *Frontina*); Swezey, 1935a: 94, Hawaii (as *Frontina*); Schaffner and Griswold, 1934: 13, 116 (as *Achaetoneura* ex *Cynthia*); Luginbill, 1928: 82, Idaho (as *Frontina* ex *Pyrameis*); Essig, 1926: 582, Calif. (as *Frontina*); Swezey, 1907: 48-49, Hawaii (as *Frontina*); Coquillett, 1897: 15, 27, Calif. (as *Frontina* ex *Pyrameis*); *Brauer and Bergenstamm, 1895: 542, 609 (as *Achaetoneura websteri* (Townsend), Townsend, 1891); Coquillett, 1894: 30, [Calif.] (as *Masicera* ex *Pyrameis*); *vide Townsend, 1892b: 259 (as *Meigenia websteri* ex *Pyrameis*, Townsend, 1891); Townsend, 1891: 207, Ind. (as *Meigenia websteri* ex chrysalis); Riley, 1871: 151 (as *Tachina* ex *Cynthia*).

Vanessa carye Hübner

Beneway, 1963: 640 (ex *V. "caryae"*); Essig, 1926: 582, Calif. (as *Frontina*); Coquillett, 1897: 15, 27, Calif. (as *Frontina* ex *Pyrameis*); Coquillett, 1894: 30, Calif. (as *Masicera* ex *Pyrameis* "caryae").

Papilionidae

Papilio spp.

Beneway, 1963: 640; *Brauer and Bergenstamm, 1895: 560, 602, N. Amer. (as *Masicera*), Williston, 1889); Williston, 1889: 1914, 1923, Colo. (as *Masicera*).

Pieridae

Colias eurytheme Boisduval

Most records ex *Eurymus eurytheme*.

Butler, 1958: 562, Ariz. (as *Achaetoneura* ex *C. philodice eurytheme*); Essig, 1926: 582, Calif., Utah (as *Frontina*); *Essig, 1915: 460 (as *Frontina*, Wildermuth, 1914); Wildermuth, 1914: 24, Calif., Utah (as *Frontina*).

Pieris protodice Boisduval and LeConte

Beneway, 1963: 640.

Pieris rapae (Linnaeus)

Anon., 1968: 85, Hawaii; Miyahira, 1967a: 1048, Hawaii; Beneway, 1963: 640; Zimmerman, 1958: 450, Hawaii (as *Frontina*); Holdaway and Lucas, 1941: 41-42, Hawaii (as *Frontina*); Holdaway and Suehiro, 1940: 37, Hawaii (as *Frontina* ex *Pontia*); Pemberton and Williams, 1938: 222, Hawaii (as *Frontina*); Schaffner and Griswold, 1934: 7-8, 116 (as *Achaetoneura* ex *Ascia*); Williams, 1931: 292, Hawaii (as *Frontina* ex *Pontia*); Webber 1930: 4 (as *Achaetoneura* ex *Pontia*); Essig, 1926: 582, Calif. (as *Frontina*); Chittenden, 1916: 9 (as *Frontina* ex *Pontia*); Swezey, 1907: 48-49, Hawaii probably (as *Frontina*).

Pieris sp.

Beneway, 1963: 640.

Pyralidae

Hedylepta accepta (Butler)

All records ex *Omiodes accepta*.

Swezey, 1943: 280, Hawaii (as *Frontina*); Bianchi, 1940: 384, Hawaii (as *Frontina*); Van Dine, 1911: 43, Hawaii (as *Frontina*); Swezey, 1907: 48, Hawaii (as *Frontina*).

Hedylepta blackburni (Butler)

All records ex *Omiodes blackburni*.

Bianchi, 1940: 384, Hawaii (as *Frontina*); Williams, 1931: 148, 292, Hawaii (as *Frontina*); Swezey, 1907: 48, Hawaii (as *Frontina*).

Loxostege commixtalis (Walker)

Hoerner, 1933: 12 (as *Achaetoneura*).

Loxostege similalis (Guenée)

Butler, 1958: 562, Ariz. (as *Achaetoneura*).

Loxostege sticticalis (Linnaeus)

Beneway, 1963: 640; Knowlton, Janes and Thomas, 1933: 156, Utah (as *Achaetoneura* ex sugar-beet webworm);

Hawley, 1925: 12, Utah (as *Frontina* "archippiva").

Pyralidae and/or Noctuidae,
see Noctuidae and/or Pyralidae

Yponomeutidae

Mapsidius iridescens Walsingham

Swezey, 1943: 283, Hawaii (as *Frontina*).

Unidentified Lepidoptera

Cabbage worm(-s)

Swezey, 1935: 26, Hawaii (as *Frontina*).

Miscellaneous

Swezey, 1931a: 495, Hawaii (as *Frontina* ex caterpillars feeding on *Acacia koa* A. Gray).

STAGE PARASITIZED: Larva (Lepidoptera), pupa (Lepidoptera), adult (Lepidoptera [Weaver, 1969: 71]).

IMMATURES: Third instar larva — Teale, 1948: (unnumbered fig. on p. 358); Riley, 1871: (descr.).

Pupa — Teale, 1948: (unnumbered fig. on p. 358); Williams, 1931: (descr.); Greene, 1922: (descr., fig. 30 on pl. 7).

BIOLOGY: Bryan, D. E., Jackson and Patana, 1969: (765-767); Bryan, D. E., Jackson and Stoner, 1969: (1, 4-5, 12); Bryan, D. E., Jackson and Patana, 1968: (819-823); Kamran, 1968: (1011); Brewer and Thomas, 1966: (235); Einmo, 1963: (196); Pemberton, 1948b: (185, 200); Teale, 1948: (356-359); Holdaway and Lucas, 1941: (41-42); Swezey, 1937: (359); Williams, 1931: (148, 292); Hawley, 1925: (12); Pemberton, 1923: (258); Johannsen, 1910: (38); Riley, 1873: (149); Riley, 1871: (116, 149-150).

Lespesia archippivora (Riley) probably

HOST: LEPIDOPTERA

Noctuidae

Spodoptera frugiperda (J. E. Smith)

Patton, 1958: 32, Fla. (as *Achaetoneura ex Laphygma*).

Nymphalidae

Vanessa cardui (Linnaeus)

Munson, 1966: 979, Mo.

Sphingidae

Sphinx sp.

Brauer and Bergenstamm, 1895: 566, 606, N. Amer. (as
Parafrontina (? *apicalis*) Brauer and Bergenstamm).

STAGE PARASITIZED: Larva.

Lespesia callosamiae Beneway

HOST: LEPIDOPTERA

Saturniidae

Callosamia promethea (Drury)

Beneway, 1963: 642-643, Conn., N.J., N.Y., Ohio (ex
Hyalophora).

Lespesia ciliata (Macquart)

HOST: LEPIDOPTERA

Bombycidae

Bombyx sp.

Robineau-Desvoidy, 1863: 569.

Lasiocampidae

Malacosoma americanum (Fabricius)

Bess, 1936: 605 (as *Achaetoneura samiae* Webber); Schaffner
and Griswold, 1934: 88, 117 (as *Achaetoneura samiae*);
Webber, 1930: 4 (as *Achaetoneura samiae*).

Malacosoma californicum (Packard)

Stehr and Cook, 1968: 264, 299-300, Ariz., N. Mex.

Malacosoma californicum lutescens (Neumoegen and Dyar)

Stehr and Cook, 1968: 275, 299, 301, Alta.

Malacosoma incurvum incurvum (H. Edwards)

Stehr and Cook, 1968: 280, 299, 301, Ariz.

Pyralidae

Loxostege sticticalis (Linnaeus)

Strickland, 1946: 172, Alta. (as *Achaetoneura samiae*).

Saturniidae

Hyalophora gloveri (Strecker)

Beneway, 1963: 645.

Hyalophora nokomis Brodie

Beneway, 1963: 645 (ex *H. columbia nokomis*).

Hyalophora sp.

Beneway, 1963: 645.

Platysamia cecropia (Linnaeus)

Beneway, 1963: 645; Raizenne, 1952: 14, Ont. (as *Achaetoneura samiae*); Procter, 1946: 434, Maine (as *Achaetoneura samiae* ex *Samia*); Breland, 1940: 261, N.Y. (as *Achaetoneura samiae* ex *cecropia* moth); Schaffner and Griswold, 1934: 26-27, 117 (as *Achaetoneura samiae*); Webber, 1930: 4, 17, 31, fig. 14, Maine, Mass., N.H., N.Y., R.I. (as *Achaetoneura samiae* ex *Samia*).

Platysamia euryalus (Boisduval)

Beneway, 1963: 645; Webber, 1930: 17, Calif. (as *Achaetoneura samiae* ex *Samia rubra* Behr).

STAGE PARASITIZED: Larva, pupa.

IMMATURES: Pupa — Beneway, 1963: (descr.); Webber, 1930: (descr., fig. 14).

BIOLOGY: Schaffner and Griswold, 1934: (117); Webber, 1930: (4, 17, 31).

Lespesia cuculliae (Webber)

HOST: LEPIDOPTERA

Noctuidae

Cucullia spp.

Beneway, 1963: 647; Webber, 1930: 4, 19, Va. (as *Achaetoneura*).

Notodontidae

Dasylophia anguina (J. E. Smith)

Beneway, 1963: 647; Schaffner and Griswold, 1934: 78-79, 116, N.J., N.Y. (as *Achaetoneura*).

Datana angusii Grote and Robinson

Beneway, 1963: 647.

IMMATURES: Pupa — Webber, 1930: (descr.).

BIOLOGY: Schaffner and Griswold, 1934: (78-79, 116).

Lespesia cuculliae (Webber) probably

HOST: LEPIDOPTERA

Notodontidae

Dasylophia anguina (J. E. Smith)

Vide Webber, 1930: 4, 19, Mass., N.Y. (as *Achaetoneura*,
Coquillett, 1897); Coquillett, 1897: 15, 24, Mass. (as
Frontina aletiae (Riley) or *Frontina frenchii* (Williston)
ex "*Dasylopha*").

STAGE PARASITIZED: Larva.

Lespesia sp. near *cuculliae* (Webber)

HOST: LEPIDOPTERA

Arctiidae

Estigmene acrea (Drury)Patton, 1958: 32, Fla. (as *Achaetoneura*).*Lespesia datanarum* (Townsend)

HOST: LEPIDOPTERA

Citheroniidae

Anisota rubicunda (Fabricius)

Beneway, 1963: 649; Schaffner and Griswold, 1934: 30-31,
116 (as *Achaetoneura anisotae* Webber); Webber, 1930:
4 (as *Achaetoneura anisotae*).

Anisota senatoria J. E. Smith

Beneway, 1963: 649; Raizenne, 1952: 17, Ont. (as
Achaetoneura anisotae); Schaffner and Griswold, 1934:
29-30, 116 (as *Achaetoneura anisotae*); Webber, 1930: 4,
14-15, Conn., N.J., N.Y., R.I. (as *Achaetoneura*
anisotae).

Anisota virginensis (Drury)

Beneway, 1963: 649; Raizenne, 1952: 18, Ont. (as
Achaetoneura anisotae).

Anisota sp.

Beneway, 1963: 649.

Notodontidae

Datana integerrima Grote and Robinson

Beneway, 1963: 649.

Datana ministra (Drury)

Beneway, 1963: 649.

Datana spp.

*Beneway, 1963: 649, N.Y. (Townsend, 1892a); *Brauer and Bergenstamm, 1895: 560, 590, N. Amer. (as (*Masicera*), Townsend, 1892a); Townsend, 1892a: 287, N.Y. (as *Masicera*).

Saturniidae

Antheraea polyphemus (Cramer)

Leech and Farris, 1959: 22, B.C. (as *Achaetoneura*).

STAGE PARASITIZED: Pupa.

IMMATURES: Pupa — Webber, 1930: (descr.).

BIOLOGY: Schaffner and Griswold, 1934: (116).

Lespesia dimmocki (Webber)

Most records as *Achaetoneura dimmocki*.

HOST: LEPIDOPTERA

Noctuidae

Simyra henrici (Grote)

Beneway, 1963: 651 (as *Lespesia*).

Saturniidae

Automeris io (Fabricius)

Beneway, 1963: 651, Mass. (as *Lespesia*); Aldrich, 1932: 3, Mass.; Webber, 1930: 4, 20, Mass.

Unidentified Lepidoptera

Miscellaneous

Aldrich, 1932: 3, Mass. (ex larva on *Rhus copallina* Linnaeus).

STAGE PARASITIZED: Larva.

Lespesia dubia (Williston)

HOST: LEPIDOPTERA

Nymphalidae

Nymphalis antiopa (Linnaeus)

*Brauer and Bergenstamm, 1895: 560, 609, N. Amer. (as *Masicera*) (?*Phorocera*) ex *Vanessa*, Williston, 1889);
Williston, 1889: 1914, 1924 (as *Masicera* ex *Euvanessa*).

Lespesia euchaetiae (Webber)

Most records as *Achaetoneura euchaetiae*.

HOST: LEPIDOPTERA

Arctiidae

Euchaetias egle (Drury)

Beneway, 1963: 652-653, fig. 34A-B, Mass., N.Y. (as *Lespesia*); Schaffner and Griswold, 1934: 39, 116;
Webber, 1930: 4, 12-13, 31, fig. 13, Mass., N.J., N.Y., Pa.

Halisidota tessellaris (J. E. Smith)

Webber, 1930: 4.

STAGE PARASITIZED: Larva.

IMMATURES: Pupa — Beneway, 1963: (fig. 34A-B); Webber, 1930: (descr., fig. 13).

BIOLOGY: Webber, 1930: (4, 12-13, 31).

Lespesia flavifrons Beneway

HOST: LEPIDOPTERA

Sphingidae

Paonias myops (J. E. Smith)

Beneway, 1963: 656-657, Mass., Pa.

Smerinthus cerisyi Kirby

Beneway, 1963: 657.

Smerinthus jamaicensis (Drury)

Beneway, 1963: 656-657, Mass.

Smerinthus sp.

Beneway, 1963: 657.

Lespesia frenchii (Williston)

HOST: HYMENOPTERA

Apidae

Bombus sp.

Coquillett, 1897: 16, 22, Canada (as *Frontina*).

HOST: LEPIDOPTERA

Arctiidae

Diacrisia virginica (Fabricius)

Schaffner and Griswold, 1934: 36, 117 (as *Achaetoneura*);
Webber, 1930: 4 (as *Achaetoneura*).

Euchaetias egle (Drury)

Webber and Schaffner, 1926: 20, New England (as *Frontina*).

Halisidota caryae (Harris)

Schaffner and Griswold, 1934: 33, 117 (as *Achaetoneura*).

Halisidota tessellaris (J. E. Smith)

Schaffner and Griswold, 1934: 34, 117 (as *Achaetoneura*).

Hyphantria cunea (Drury)

Schaffner and Griswold, 1934: 37, 117 (as *Achaetoneura*);
Webber, 1930: 4 (as *Achaetoneura*).

Bombycidae

Unidentified Bombycidae

Howard, 1897b: 42, D.C. (as *Frontina*).

Citheroniidae

Anisota rubicunda (Fabricius)

Webber and Schaffner, 1926: 26, New England (as *Frontina*);
Howard and Chittenden, 1909: 5, D.C. (as *Frontina*);
Felt, 1908: 16, N.Y. (as *Frontina*); Coquillett, 1897: 16,
25, Md. (as *Frontina ex Dryocampa*).

Anisota senatoria J. E. Smith

Webber and Schaffner, 1926: 25, New England (as *Frontina*);
Johnson, 1910: 780, N.J. (as *Frontina*); Coquillett,
1897: 15, 23, Md. (as *Frontina*).

Anisota virginiensis (Drury)

Coquillett, 1897: 16, 24 (as *Frontina*).

Citheronia regalis (Fabricius)

Coquillett, 1897: 16, 24 (as *Frontina*).

Eacles imperialis (Drury)

Fattig, 1949: 4, Ga. (as *Achaetoneura ex Basilonia*).

Geometridae

Unidentified Geometridae

Schaffner and Griswold, 1934: 102, 117 (as *Achaetoneura*).

Lasiocampidae

Malacosoma americanum (Fabricius)

Stehr and Cook, 1968: 299-300, Minn.; Kulman, 1965: 68-69, W. Va.; Beneway, 1963: 658, fig. 33; Raizenne, 1952: 101, Ont. (as *Achaetoneura*); *Procter, 1946: 434, Maine (as *Achaetoneura* ex "*Malacosomoa*," Procter, 1938); Procter, 1938: 375, Maine (as *Achaetoneura*); Bess, 1936: 605 (as *Achaetoneura*); Britton, 1935a: 77, (as *Frontina*); Schaffner and Griswold, 1934: 88, 117 (as *Achaetoneura*); Webber, 1930: 4 (as *Achaetoneura*); Quaintance, 1915: 7 (as *Frontina*); Johnson, 1910: 780, N.J. (as *Frontina* ex *Clisiocampa*); Quaintance, 1908a: 6, E. U.S. (as *Frontina*); Surface, 1904: 43, N.E. U.S. (as "*Frontia*" ex *Clisiocampa*); Howard, 1902: 109, W. Va. (as *Frontina* ex *Clisiocampa*); Coquillett, 1897: 16, 24, Mass. (as *Frontina* ex *Clisiocampa*).

Malacosoma californicum (Packard)

Stehr and Cook, 1968: 260, 262-263, 266, 268, 270, 299-300, Calif., Oreg., Utah, Wash., Wyo.; Essig, 1926: 582, Calif. (as *Frontina*); Coquillett, 1897: 16, 24, Calif. (as *Frontina* ex *Clisiocampa*).

Malacosoma californicum californicum (Packard)

Coquillett, 1897: 16, 24, Calif. (as *Frontina* ex *Clisiocampa thoracica* Stretch).

Malacosoma californicum fragile (Stretch)

Stehr and Cook, 1968: 279, 299, 301, Utah; Beneway, 1963: 658, fig. 33 (ex *M. fragilis*).

Malacosoma californicum lutescens (Neumoegen and Dyar)

Stehr and Cook, 1968: 275, 299, 301, Alta.

Malacosoma californicum pluviale (Dyar)

Stehr and Cook, 1968: 274-275, 299-300, Alta., B.C., Mont., Oreg., Wash.; Beneway, 1963: 658, fig. 33 (ex *M. pluvialis*); Johansen, 1957: 70, Wash. (as *Achaetoneura* ex *M. pluviale*).

Malacosoma constrictum (H. Edwards)

Most records ex *Clisiocampa constricta*.

Essig, 1926: 582, B.C. (as *Frontina* ex *M. constricta*); Coquillett, 1897: 16, 24, Calif. (as *Frontina*); Coquillett, 1894: 30, Calif. (as *Masicera*).

Malacosoma disstria Hübner

Stehr and Cook, 1968: 255, 299, Alta.; Beneway, 1963: 658, fig. 33; Beneway, 1963: 658, fig. 33 (ex *M. disstria erosa* Stretch); Sippell, 1961: 435-436, fig. 10, Ont. (as

Achaetoneura); Johansen, 1957: 70, Wash. (as *Achaetoneura*); Raizenne, 1952: 103, Ont. (as *Achaetoneura*); Strickland, 1938: 218, Alta. (as *Achaetoneura* ex forest tent caterpillar); Bess, 1936: 605 (as *Achaetoneura*); Schaffner and Griswold, 1934: 89, 117 (as *Achaetoneura*); Webber, 1930: 4 (as *Achaetoneura*); Essig, 1926: 582, Calif. (as *Frontina*).

Malacosoma spp.

Beneway, 1963: 658, fig. 33; Ross, 1953: 19, 22, fig. 5, B.C. (as *Achaetoneura*).

American tent caterpillar(-s)

Curran, 1925a: 156, N.S. (as *Frontina malacosomae* Curran).

Tent caterpillar(-s)

Brimley, 1922: 23, N.C. (as *Frontina*).

Lymantriidae

Hemerocampa leucostigma (J. E. Smith)

Most records ex *Orgyia leucostigma*.

Schaffner and Griswold, 1934: 85, 117 (as *Achaetoneura* ex *Hemerocampa*); Webber and Schaffner, 1926: 12-13, 20, Mass., New England, N.Y., Pa., R.I. (as *Frontina* ex *Hemerocampa*); Britton, 1917: 110 (as *Frontina* ex *Hemerocampa*); Johnson, 1910: 780, N.J. (as *Frontina*); Howard, 1898: 88, D.C. (as *Frontina*); Coquillett, 1897: 16, 26, D.C. (as *Frontina*); Howard, 1897a: 16-17, D.C. (as *Frontina*); Howard, 1897b: 7, 42, 49, D.C. (as *Frontina*).

Hemerocampa vetusta (Boisduval)

Essig, 1926: 582, Calif. (as *Frontina*).

Nygmia phaeorrhoea (Donovan)

Webber, 1930: 4 (as *Achaetoneura*); Burgess and Crossman, 1929: 143 (as *Achaetoneura*).

Porthetria dispar (Linnaeus)

Webber, 1930: 4 (as *Achaetoneura*); Burgess and Crossman, 1929: 143 (as *Achaetoneura*).

Stilpnotia salicis (Linnaeus)

Beneway, 1963: 658, fig. 33; Lejeune and Silver, 1961: 459-460, 462, B.C. (as *Achaetoneura*); Reeks and Smith, 1956: 569, maritime prov. (as *Achaetoneura*); Burgess and Crossman, 1927: 14, New England (as *Frontina*).

Noctuidae

Faronta diffusa (Walker)

Essig, 1926: 583, Colo. (as *Frontina* ex *Neleucania albilinea* (Hübner)); Webster, R. L., 1911: 181 (as *Frontina* ex *Meliana albilinea*).

Heliothis zea (Boddie)

Most records ex *Heliothis armigera* (Hübner).

Bibby, 1942: 943, Tex. (as *Achaetoneura*); Quaintance and Brues, 1905: 123 (as *Frontina* ex *H. obsoleta* (Fabricius)); Coquillett, 1897: 16, 25 (as *Frontina*).

Hydroecia immanis Guenée

Hawley, 1918: 169, N.Y. (as *Frontina* ex *Gortyna*).

Hypsoropha hormos Hübner

Coquillett, 1897: 16, 25 (as *Frontina*).

Unidentified Noctuidae

Howard, 1897b: 42, D.C. (as *Frontina*).

Notodontidae

Datana angusii Grote and Robinson

Schaffner and Griswold, 1934: 74-75, 117 (as *Achaetoneura*); Webber, 1930: 4 (as *Achaetoneura*).

Datana integerrima Grote and Robinson

Hixson, 1941: 20, Okla. (as *Achaetoneura*); Schaffner and Griswold, 1934: 76, 117 (as *Achaetoneura*); Webber, 1930: 4 (as *Achaetoneura*).

Datana ministra (Drury)

Schaffner and Griswold, 1934: 74, 117 (as *Achaetoneura*); Webber, 1930: 4 (as *Achaetoneura*); Essig, 1926: 582-583, B.C. (as *Frontina*); Coquillett, 1897: 16, 24, Ind. (as *Frontina*).

Datana sp.

Coquillett, 1897: 16, 24 (as *Frontina*).

Ichthyura inclusa Hübner

Dowden, 1935: 509 (as *Achaetoneura* ex *Melalopha*); Webber and Schaffner, 1926: 19, New England (as *Frontina* ex *Melalopha*); Webster, 1899a: 224, Ind. (as *Frontina*); Coquillett, 1897: 16, 25, Ind., Mo. (as *Frontina*).

Nymphalidae

Nymphalis antiopa (Linnaeus)

Schaffner and Griswold, 1934: 11, 117 (as *Achaetoneura* ex *Hamadryas*).

Vanessa atalanta (Linnaeus)

Schaffner and Griswold, 1934: 12, 117 (as *Achaetoneura* ex *Cynthia*); Webber, 1930: 4 (as *Achaetoneura*).

Vanessa cardui (Linnaeus)

All records ex *Pyrameis cardui*.

Webster, 1899a: 224, Ind. (as *Frontina*); Howard, 1897b: 42, D.C. (as *Frontina*).

Papilionidae

Papilio bairdii oregonius Edwards

Beneway, 1963: 658, fig. 33.

Papilio eurymedon Lucas

Beneway, 1963: 658, fig. 33; Tothill, 1913: 71-72, B.C. (as *Frontina*).

Papilio glaucus Linnaeus

Beneway, 1963: 658, fig. 33; Beneway, 1963: 658, fig. 33 (ex *P. glaucus turnus* Linnaeus); Schaffner and Griswold, 1934: 7, 117 (as *Achaetoneura* ex *P. glaucus* form *turnus*); Webber, 1930: 4 (as *Achaetoneura* ex *P. turnus*); Johnson, 1925: 200, Conn., Maine, Mass., N.H., Vt. (as *Frontina*); Johnson, 1910: 780, N.J. (as *Frontina* ex *P. turnus*); Howard, 1897b: 42, D.C. (as *Frontina* ex *P. turnus*); *Brauer and Bergenstamm, 1895: 560, 595, N. Amer. (as (*Masicera*) ex *Jasoniades*, Williston, 1889); Coquillett, 1894: 30, Maine (as *Masicera* ex *P. turnus* var. *glaucus*); Williston, 1889: 1914, 1923, Maine (as *Masicera* ex *Jasoniades*).

Papilio glaucus canadensis Rothschild and Jordan

Beneway, 1963: 658, fig. 33.

Papilio multicaudata Kirby

All records ex *Papilio daunus* Boisduval.

Beneway, 1963: 658, fig. 33; Tothill, 1913: 71, B.C. (as *Frontina*).

Papilio polyxenes Fabricius

Webber, 1930: 4 (as *Achaetoneura*).

Papilio polyxenes asterius Stoll

Schaffner and Griswold, 1934: 6, 117 (as *Achaetoneura* ex *P. ajax* Linnaeus).

Papilio rutulus Lucas

Beneway, 1963: 658, fig. 33; Essig, 1926: 582, B.C. (as *Frontina*).

Papilio troilus Linnaeus

Schaffner and Griswold, 1934: 7, 117 (as *Achaetoneura*); Webber, 1930: 4 (as *Achaetoneura*).

Papilio sp.

Beneway, 1963: 658, fig. 33.

Pieridae

Pieris rapae (Linnaeus)

Schaffner and Griswold, 1934: 7-8, 117 (as *Achaetoneura* ex *Ascia*); Webber, 1930: 4 (as *Achaetoneura* ex *Pontia*).

Pyralidae

Unidentified Pyralidae

Coquillett, 1897: 17, 26, Va. (as *Frontina*).

Saturniidae

Actias luna (Linnaeus)

Webber, 1930: 4 (as *Achaetoneura* ex *Tropaea*).

Antheraea polyphemus (Cramer)

Most records ex *Telea polyphemus*.

Collins, M. M., and Weast, 1961: 93-94, unnumbered fig. on p. 96 (as *Frontina* ex *Antheraea*); Fattig, 1949: 4-5 (as *Achaetoneura*); Brimley, 1938, N.C. (as *Achaetoneura*); Schaffner and Griswold, 1934: 28, 117 (as *Achaetoneura*); Webber, 1930: 4 (as *Achaetoneura*); Allen, 1929: 687, Miss. (as *Achaetoneura*); Essig, 1926: 582, Calif. (as *Frontina*); Johnson, 1925: 200, Conn., Maine, Mass., N.H., Vt. (as *Frontina*); Brimley, 1922: 23, N.C. (as *Frontina* ex *Polyphemus* moth); Packard, 1914: 268 (as *Frontina*); Johnson, 1910: 780, N.J. (as *Frontina*); Hine, 1904: 87, B.C. (as *Frontina*); Coquillett, 1897: 17, 27, Pa. (as *Frontina*).

Attacus sp.

Coquillett, 1897: 16, 24, Calif. (as *Frontina*).

Automeris io (Fabricius)

Collins, M. M., and Weast, 1961: 93-94, unnumbered fig. on p. 96 (as *Frontina*); Coquillett, 1897: 16, 25, Ill. (as *Frontina* ex *Hyperchiria*).

Callosamia promethea (Drury)

Schaffner and Griswold, 1934: 27, 117 (as *Achaetoneura*); Webber, 1930: 4 (as *Achaetoneura*).

Hyalophora calleta (Westwood)

Collins, M. M., and Weast, 1961: 93-94, unnumbered fig. on p. 96 (as *Frontina*).

Hyalophora columbia Smith

Tothill, 1913: 72, Man. (as *Frontina* ex *Samia*).

Platysamia cecropia (Linnaeus)

Collins, M.M., and Weast, 1961: 93-94, unnumbered figs. on pp. 96-97 (as *Frontina* ex *Hyalophora*); Raizenne, 1952: 14, Ont. (as *Achaetoneura* ex *Hyalophora*); Schaffner and Griswold, 1934: 26-27, 117 (as *Achaetoneura*); Webber, 1930: 4 (as *Achaetoneura* ex *Samia*); Johnson, 1925: 200, Conn., Maine, N.H., Vt. (as *Frontina* ex *Samia*); Curran, 1925a: 156 (as *Frontina* ex *Cecropia*); *Packard, 1914: 268 (as *Frontina* ex *Samia*, Felt, 1912); Felt, 1912: 116, 191, fig. 3 on pl. 27, N.Y. (as *Frontina* ex *Samia*); Johnson, 1910: 780, N.J. (as *Frontina* ex *Attacus*); Thompson, 1910: 289, 293 (as *Frontina* ex *Cecropia*); Fiske and Thompson, 1909: 453-454 (as *Achaetoneura* ex *Cecropia*); Hitchings, 1908: 12, Maine (as *Frontina* ex *Hyalophora*); Coquillett, 1897: 16, 24, N.J. (as *Frontina* ex *Attacus*).

Platysamia euryalus (Boisduval)

Collins, M. M., and Weast, 1961: 93-94, unnumbered fig. on p. 96 (as *Frontina* ex *Hyalophora rubra* Behr).

Samia cynthia (Drury)

Webber, 1930: 4 (as *Achaetoneura* ex *Philosamia*).

Samia cynthia advena (Packard)

Collins, M. M., and Weast, 1961: 93-94, unnumbered fig. on p. 96 (as *Frontina*); Schaffner and Griswold, 1934: 26, 117 (as *Achaetoneura* ex *S. walkeri advena*).

Syssphinx bicolor (Harris)

Eliot and Soule, 1902: 58 (as *Frontina* ex *Sphingicampa*).

Sphingidae

Ceratonia amyntor (Hübner)

Beneway, 1963: 658, fig. 33.

Ceratonia catalpae (Boisduval)

Beneway, 1963: 658, fig. 33; Schaffner and Griswold, 1934: 17, 117, N.J. (as *Achaetoneura*); Webber, 1930: 4 (as *Achaetoneura*); Howard and Chittenden, 1916: 6 (as *Frontina*); Howard and Chittenden, 1907: 5 (as *Frontina*); Coquillett, 1897: 16, 24, D.C. (as *Frontina*).

Ceratonia undulosa (Walker)

Beneway, 1963: 658, fig. 33; Schaffner and Griswold, 1934: 17, 117 (as *Achaetoneura*); Webber, 1930: 4 (as *Achaetoneura*).

Manduca quinquemaculata (Haworth)

Schaffner and Griswold, 1934: 16, 117 (as *Achaetoneura* ex *Phlegethontius*).

Pachysphinx modesta (Harris)

*Vide Essig, 1926: 582-583, Oreg. (as *Frontina*, Cole, F. R., and Lovett, 1921); Cole, F. R., and Lovett, 1921: 302, Oreg. (as *Frontina* ex sphingid on *Populus trichocarpa* Torrey and Gray).

Paonias myops (J. E. Smith)

Schaffner and Griswold, 1934: 21, 117 (as *Achaetoneura*); Webber, 1930: 4 (as *Achaetoneura*).

Smerinthus cerisyi Kirby

Essig, 1926: 583, Oreg. (as *Frontina*); Coquillett, 1897: 17, 27, Calif. (as *Frontina*).

Smerinthus jamaicensis (Drury)

All records ex *Smerinthus geminatus* Say.

Schaffner and Griswold, 1934: 20, 117 (as *Achaetoneura*); Aldrich, 1932: 3, Mass. (as *Achaetoneura*); Webber, 1930: 4 (as *Achaetoneura*).

Sphinx chersis (Hübner)

Beneway, 1963: 658, fig. 33; Raizenne, 1952: 10, Ont. (as *Achaetoneura*); Schaffner and Griswold, 1934: 18, 117 (as *Achaetoneura* ex *Hyloicus*); Aldrich, 1932: 1, Mass. (as *Achaetoneura*, incomplete rearing — pupa); Webber, 1930: 4 (as *Achaetoneura*).

Sphinx kalmiae J. E. Smith

Beneway, 1963: 658, fig. 33; Schaffner and Griswold, 1934: 18-19, 117 (as *Achaetoneura* ex *Hyloicus*); Webber, 1930: 4 (as *Achaetoneura*).

Unidentified Sphingidae

Schaffner and Griswold, 1934: 26, 117 (as *Achaetoneura*); *Brauer and Bergenstamm, 1895: 561, 606, N. Amer. (as *Masicera sphingivora* Townsend ex *Sphinx* sp., Townsend, 1892a); Townsend, 1892a: 286, W. Va. (as *Masicera sphingivora*).

Unidentified Lepidoptera

Miscellaneous

Essig, 1926: 583, N. Amer. (as *Frontina* ex great variety of caterpillars); Howard and Chittenden, 1907: 5 (as *Frontina*).

HOST: ORTHOPTERA

Acrididae

Dissosteira carolina (Linnaeus)

Coquillett, 1897: 16, 23, Mo. (as *Frontina*).

STAGE PARASITIZED: Larva (Lepidoptera), pupa (Lepidoptera), adult (Orthoptera).

IMMATURES: Egg — Collins, M. M., and Weast, 1961: (unnumbered fig. on p. 97).

Third instar larva — Beneway, 1963: (fig. 33); Collins, M. M., and Weast, 1961: (descr.).

Pupa — Collins, M. M., and Weast, 1961: (descr., unnumbered fig. on p. 96); Sippell, 1961: (descr., fig. 10); Ross, 1953: (descr., fig. 5); Webber, 1930: (fig. 12); Greene, 1922: (descr., fig. 28 on pl. 6); Felt, 1912: (fig. 3 on pl. 27); Thompson, 1910 (descr.).

BIOLOGY: Collins, M. M., and Weast, 1961: (93-94); Hixson, 1941: (20); Proctor, 1938: (375); Dowden, 1935: (509); Aldrich, 1932: (1); Allen, 1929: (687); Felt, 1912: (116, 191); Thompson, 1910: (289, 293); Hitchings, 1908: (12).

Lespesia frenchii (Williston) complex

All records as *Achaetoneura frenchii* complex.

HOST: LEPIDOPTERA

Notodontidae

Datana integerrima Grote and Robinson

Patton, 1958: 32, Fla.

Datana ministra (Drury)

Patton, 1958: 32, Fla.

STAGE PARASITIZED: Pupa.

BIOLOGY: Schaffner and Griswold, 1934: (117).

Lespesia laniiferae (Webber)

HOST: LEPIDOPTERA

Pyralidae

Laniifera cyclades Druce

Beneway, 1963: 660; Webber, 1930: 4, 33, Mexico (as *Achaetoneura*).

STAGE PARASITIZED: Pupa.

Lespesia sp. near *laniiferae* (Webber)

HOST: LEPIDOPTERA

Arctiidae

Eupseudosoma involutum floridanum Grote

Patton, 1958: 32, Fla. (as *Achaetoneura*, incomplete rearing
— pupa).

STAGE PARASITIZED: Larva.

BIOLOGY: Patton, 1958: (32).

Lespesia melalophae (Allen)

Most records as *Achaetoneura melalophae*.

HOST: LEPIDOPTERA

Geometridae

Metanema inatomaria Guenée

Raizenne, 1952: 150, Ont.

Lasiocampidae

Malacosoma disstria Hübner

Beneway, 1963: 662 (as *Lespesia*); Sippell, 1961: 435-436,
fig. 9, Ont.

Noctuidae

Nycteola cinereana Neumoegen and Dyar

Beneway, 1963: 662 (as *Lespesia*).

Notodontidae

Ichthyura inclusa Hübner

Most records ex *Melalopha inclusa*.

Beneway, 1963: 662 (as *Lespesia* ex "*Ichthyura*"); Fattig,
1949: 5 (ex *Ichthyura*); Brimley, 1938: 363, probably
N.C.; Schaffner and Griswold, 1934: 73, 117, Conn.,
N.J., N.Y., R.I.; Webber, 1930: 4; Allen, 1926b:
192-196, Miss.

STAGE PARASITIZED: Larva, pupa.

IMMATURES: Pupa — Sippell, 1961: (descr., fig. 9).

BIOLOGY: Schaffner and Griswold, 1934: (73, 117); Allen,
1926b: (192-196).

Lespesia pholi (Webber)

HOST: LEPIDOPTERA

Sphingidae

Pholus satellitia pandorus (Hübner)

Beneway, 1963: 664.

Pholus spp.Beneway, 1963: 664; Aldrich, 1932: 3, Mass. (as *Achaetoneura*); Webber, 1930: 4, 28, Mass. (as *Achaetoneura*).

STAGE PARASITIZED: Larva.

Lespesia rileyi (Williston)

HOST: LEPIDOPTERA

Papilionidae

Papilio cresphontes CramerBeneway, 1963: 666; Patton, 1958: 32, Fla. (as *Achaetoneura*); *Brauer and Bergenstamm, 1895: 561, 594 (as *Masicera* ex *Heraclides*, Williston, 1889); Williston, 1889: 1914, 1924 (as *Masicera* ex *Heraclides*).*Papilio thoas* Linnaeus*Vide Webber, 1930: 4, 32 (as *Achaetoneura* ex "Papilo," Coquillett, 1897); Allen, 1929: 687, Miss. (as *Frontina*); Coquillett, 1897: 17, 26, Fla. (as *Frontina irrequieta* (Walker)).

STAGE PARASITIZED: Pupa.

Lespesia sabroskyi Beneway

HOST: LEPIDOPTERA

Saturniidae

Antheraea polyphemus (Cramer)Beneway, 1963: 642, 667; *Brauer and Bergenstamm, 1895: 560, 584, N. Amer. (as (*Masicera*) *datanarum* Townsend ex *Attacus*, Townsend, 1892a); Townsend, 1892a: 287, Ont. (as *Masicera datanarum* ex *Attacus*).*Automeris io* (Fabricius)

Beneway, 1963: 642, 667.

Platysamia euryalus (Boisduval)Beneway, 1963: 642, 667 (ex *Hyalophora*).

STAGE PARASITIZED: Pupa.

Lespesia schizurae (Townsend)

HOST: LEPIDOPTERA

Arctiidae

Euchaetias egle (Drury)

*Brauer and Bergenstamm, 1895: 544, 592 (as *Argyrophylax* ex "*Euchaetes*," Townsend, 1893); Townsend, 1893: 467, Ill. (as *Masicera* ex "*Euchaetes*").

Danaiidae

Danaus plexippus plexippus (Linnaeus)

Urquhart, 1960: 207, Ont. (as *Achaetoneura* ex *D. plexippus*).

Lasiocampidae

Malacosoma americanum (Fabricius)

Dozier, 1920: 372, 378, Fla. (as *Achaetoneura*).

Notodontidae

Heterocampa biundata Walker

Beneway, 1963: 670.

Heterocampa sp.

Beneway, 1963: 670.

Schizura ipomaeae Doubleday

Beneway, 1963: 670 (ex *S. "ipomoeae"*); Webber, 1930: 4, 30, Wash. (as *Achaetoneura piperi* (Townsend) ex *S. "ipomoeae"*); Essig, 1926: 583, Wash. (as *Sturmia*); Johnson, 1925: 200, Mass. (as *Argyrophylax*); Townsend, 1908: 98, Wash. (as *Argyrophylax piperi*); Coquillett, 1897: 21, 27, Wash. (as *Sturmia* ex *S. "ipomoeae"*).

Schizura unicornis (J. E. Smith)

Beneway, 1963: 670; Schaffner and Griswold, 1934: 81-82, 117, N.J. (as *Achaetoneura piperi*); *Webber, 1930: 4, 29, Kans. (as *Achaetoneura*, Townsend, 1891c); *Brauer and Bergenstamm, 1895: 544, 605, Kans. (as *Argyrophylax*, Townsend, 1891c); Townsend, 1891c: 187, Kans. (as *Masicera*).

Sphingidae

Pholus satellitia pandorus (Hübner)

Fattig, 1949: 5, Ga. (as *Achaetoneura* ex *P. pandorus*).

STAGE PARASITIZED: Larva, pupa.

BIOLOGY: Schaffner and Griswold, 1934: (81-82, 117).

Lespesia testacea (Webber)

HOST: LEPIDOPTERA

Noctuidae

Xerociris wilsoni (Grote)

Beneway, 1963: 672.

Sphingidae

Pholus achemon (Drury)

Beneway, 1963: 672; Fattig, 1949: 5 (as *Achaetoneura*);

Brimley, 1938: 363, N.C. (as *Achaetoneura*); Brimley,

1922: 23, N.C. (as *Frontina violenta* (Walker)).

Pholus vitis (Linnaeus)

Beneway, 1963: 672; *Webber, 1930: 4, 26, D.C. (as

Achaetoneura, Coquillett, 1897); Coquillett, 1897: 17,

26, D.C. (as *Frontina violenta ex Philampelus*).

STAGE PARASITIZED: Larva.

Lespesia spp.

HOST: LEPIDOPTERA

Arctiidae

Ecpantheria icasia (Cramer)

Harrison and Stephens, 1966: 673-674, Costa Rica (as
Achaetoneura).

Citheroniidae

Anisota rubicunda (Fabricius)

Riley, 1873: 139-140, Kans. (as *Tachina anonyma* Riley ex
Dryocampa).

Anisota senatoria J. E. Smith

Brauer and Bergenstamm, 1895: 542, 582, N. Amer. (as
Achaetoneura).

Citheronia regalis (Fabricius)

*Brauer and Bergenstamm, 1895: 541, 587 (as *Achaetoneura*
anonyma, Riley, 1872); Riley, 1872: 129 (as *Tachina*
anonyma).

Ctenuchidae

Syntomeida epilais jucundissima Dyar

Fernald, H. T., 1934: 23-24, Fla. (as *Achaetoneura*).

Danaiidae

Danaus plexippus plexippus (Linnaeus)

All records ex *Danaus archippus* (Fabricius).

Brauer and Bergenstamm, 1895: 541, 590 (as *Achaetoneura anonyma* ex "Danais"); Riley, 1872: 129 (as small *Tachina anonyma* ex "Danais").

Lasiocampidae

Malacosoma californicum (Packard)

Brauer and Bergenstamm, 1895: 542, 588 (as *Achaetoneura* ex *Clisiocampa*).

Malacosoma disstria Hübner

Anon., 1960d: 1026, S. and S.E. U.S. (as *Achaetoneura*).

Malacosoma sp.

Gibson, W. W., and Carrillo S., 1959: 180, Mexico (D. F.) (as *Achaetoneura* ex "*Malocosoma*").

Lycaenidae

Strymon melinus (Hübner)

Little and Martin, 1941: 74, Tex. (as *Frontina*); Reinhard, 1929: 33-34, Tex. (as *Frontina*).

Lymantriidae

Hemerocampa leucostigma (J. E. Smith)

Schaffner and Griswold, 1934: 85 (as *Achaetoneura*).

Megathymidae

Megathymus yuccae (Boisduval and LeConte)

Riley, 1876: 179 (as [*Tachina*] *anonyma*); Riley, 1876b: 339, S.C. (as [*Tachina*] *anonyma*).

Noctuidae

Acronicta lanceolaria (Grote)

Schaffner and Griswold, 1934: 60, 117, Mass. (as near *Achaetoneura* (n. sp.) (1) ex *Acronycta*).

Autographa californica (Speyer)

Clancy, 1969: 1079, Calif.

Celama sorghiella (Riley)Reinhard, 1938: 32, Tex. (as *Achaetoneura*).*Faronta diffusa* (Walker)Titus and Pratt, 1904: 62 (as *Tachina anonyma* ex *Heliophila albilinea* (Hübner)); Riley, 1877: 54, probably Mo. (as *Tachina anonyma* ex *Leucania albilinea*).*Heliothis virescens* (Fabricius)

Lewis, W. J., and Brazzel, 1968: 674-675, Miss.

Heliothis zea (Boddie)Most records ex *Heliothis armigera* (Hübner).Lewis, W. J., and Brazzel, 1968: 674-675, Miss. (ex *H. zea*); Brauer and Bergenstamm, 1895: 542, 593 (as *Achaetoneura anonyma*); Brauer and Bergenstamm, 1895: 542, 594, N. Amer. (as *Achaetoneura*); Riley, 1872: 129 (as small *Tachina anonyma*).*Hypsoropha hormos* HübnerBrauer and Bergenstamm, 1895: 542, 595, N. Amer. (as *Achaetoneura*).*Simyra henrici* (Grote)Schaffner and Griswold, 1934: 61, 117, Mass. (as near *Achaetoneura* (n. sp.) (2)).*Spodoptera frugiperda* (J. E. Smith)All records ex *Prodenia autumnalis* Riley.Brauer and Bergenstamm, 1895: 542, 604 (as *Achaetoneura anonyma*); Riley, 1872: 129 (as small *Tachina anonyma*).*Trichoplusia ni* (Hübner)Clancy, 1969: 1079-1080, Calif.; Oatman, 1966: 1138, Calif. (as *L. "achaetoneura* (Williston)").

Unidentified Noctuidae

Riley, 1872: 129 (as small *Tachina anonyma*).

Notodontidae

Datana ministra (Drury)Brauer and Bergenstamm, 1895: 541, 590 (as *Achaetoneura anonyma*); Brauer and Bergenstamm, 1895: 542, 590, N. Amer. (as *Achaetoneura*); Riley, 1872: 129 (as small *Tachina anonyma*).*Heterocampa umbrata* WalkerSchaffner and Griswold, 1934: 79, 117, Mass. (as *Achaetoneura*).

Ichthyura inclusa Hübner

Brauer and Bergenstamm, 1895: 542, 595, N. Amer. (as *Achaetoneura*).

Ichthyura spp. (including *I. apicalis* Walker or *I. inclusa* Hübner)

Allen, 1924: 24, Miss. (as *Frontina* undescribed sp. ex "*Melanopha*" *inclusa* or *apicalis*).

Nymphalidae

Vanessa cardui (Linnaeus)

Brauer and Bergenstamm, 1895: 542, 609 (as *Achaetoneura anonyma*); Riley, 1872: 129 (as small *Tachina anonyma* ex *Cynthia*).

Olethreutidae

Eucosma scudderiana (Clemens)

Brauer and Bergenstamm, 1895: 542, 592, N. Amer. (as *Achaetoneura* ex *Euryptychia saligneana* Clemens).

Grapholitha molesta (Busck)

Stearns and Amos, 1941: 247, Del. (as *Achaetoneura*).

Pieridae

Colias eurytheme Boisduval

Lochhead, 1919: 176 (as "*Formtina*" ex *Eurymus*).

Pieris rapae (Linnaeus)

Oatman, 1966: 1137-1138, Calif. (as *L. "Achaetoneura (Williston)"*).

Psychidae

Oiketiscus kirbyi Guilding

Stephens, 1962: 385, Costa Rica (as *Achaetoneura*).

Pyralidae

Diatraea sp.

Gibson, W. W., and Carrillo S., 1959: 180, Mexico (V.C.) (as *Achaetoneura*).

Laniifera cyclades Druce

Mann, 1969: 14, 77, probably Mexico (ex *Megastes*).

Melitara prodenialis Walker

Mann, 1969: 14, 32, Fla.

Pyrausta signatalis (Walker)Schaffner, 1959: 12, 48, Mass. (as *Achaetoneura*).

Unidentified Pyralidae

Brauer and Bergenstamm, 1895: 542, 604, N. Amer. (as *Achaetoneura* ex pyralid on alder tree).

Saturniidae

Antheraea polyphemus (Cramer)Most records ex *Attacus polyphemus*.Powell, 1958: 48, Calif. (as *Achaetoneura* ex "Antheracea"); *Brauer and Bergenstamm, 1895: 541, 584 (as *Achaetoneura anonyma*, Riley, 1872); Brauer and Bergenstamm, 1895: 542, 584, N. Amer. (as *Achaetoneura*); Riley, 1872: 129, Mo. (as *Tachina anonyma*).*Hemileuca maia* (Drury)Riley, 1873: 133 (as [*Tachina*] *anonyma* ex *Saturnia*).*Rothschildia aroma* Schaus probably

Quezada, 1967: 595, 597, El Salvador.

Sphingidae

Ceratonia catalpae (Boisduval)Schaffner and Griswold, 1934: 17, 117, N.J. (as *Achaetoneura*).*Manduca sexta* (Linnaeus)All records ex *Sphinx carolina* Linnaeus.Brauer and Bergenstamm, 1895: 542, 606 (as *Achaetoneura anonyma*); Riley, 1872: 129 (as *Tachina anonyma*, var.).

HOST: ORTHOPTERA

Acrididae

Dissosteira carolina (Linnaeus)Riley, 1875: 178, Mo. (as *Tachina anonyma* ex *Oedipoda*).*Melanoplus differentialis* (Thomas)Riley, 1875: 178, Mo. (as *Tachina anonyma* ex differential locust).*Melanoplus spretus* (Walsh)All records ex *Caloptenus spretus*.

Brauer and Bergenstamm, 1895: 541, 586 (as *Achaetoneura anonyma*); Riley, Packard and Thomas, 1883: 40, Mont., Wyo. (as *Tachina anonyma*); Riley, 1875: 178, Colo. (as *Tachina anonyma*).

Locust(-s)

Riley, Packard and Thomas, 1878: 323 (as *Tachina anonyma*).

STAGE PARASITIZED: Larva (Lepidoptera), pupa (Lepidoptera), adult (Orthoptera).

BIOLOGY: Clancy, 1969: (1079-1080); Schaffner, 1959: (12, 48); Powell, 1958: (48); Schaffner and Griswold, 1934: (60-61, 79, 117); Riley, 1873: (133).

Leucostoma Meigen (Phasiinae: Leucostomatini)

Leucostoma acirostre Reinhard

HOST: HEMIPTERA (HETEROPTERA)

Coreidae

Harmostes reflexulus (Say)

Reinhard, 1956: 163, D.C.

STAGE PARASITIZED: Adult

Leucostoma acuminata, see *Leucostoma aterrimum*

Leucostoma aterrimum (Villers)

HOST: HEMIPTERA (HETEROPTERA)

Rhopalidae

Liorhyssus hyalinus (Fabricius)

All records ex *Corizus hyalinus*.

Bryan, 1934: 416, Hawaii; Allen, 1929: 684, Miss. (as *L. acuminata* Townsend).

Leucostoma atra, see *Leucostoma simplex*

Leucostoma gravipes (Wulp)

HOST: HEMIPTERA (HETEROPTERA)

Lygaeidae

Oncopeltus fasciatus (Dallas)

Prine, 1967: 92, Calif.

Leucostoma simplex (Fallén)

Most records as *Leucostoma atra* Townsend.

HOST: HEMIPTERA (HETEROPTERA)

Nabidae

Nabis alternatus Parshley

Morrill, W. L., 1969: 240, S. Dak. (as *L. simplex*); Clancy and Pierce, 1966: 857-858, Calif. (as *L. simplex*); Taylor, 1951: 133, probably Utah (probable host); Taylor, 1949: 991, Utah.

Nabis americanoferus Carayon

Clancy and Pierce, 1966: 857-858, Calif. (as *L. simplex*); Reinhard, 1956: 160, Wis. (as *L. simplex* ex *N. ferus* (Linnaeus)); Swezey, 1922a: 30-31 (ex *Reduviolus ferus*).

Nabis roseipennis Reuter

Shannon, 1914: 182, Md. (ex *Reduviolus*).

Rhopalidae

Liorhyssus hyalinus (Fabricius)

All records ex *Corizus hyalinus*.

Williams, 1931: 100, 296-297, Hawaii; Swezey, 1922a: 30-31, Hawaii.

HOST: ORTHOPTERA

Acrididae

Melanoplus sanguinipes (Fabricius)

Smith, R. W., 1944: 29, Ont. (ex *M. mexicanus* Saussure).

Grasshopper(-s)

Smith, R. W., and Finlayson, 1950: 98-100, figs. 27, 51, Canada (as *Paradionaea*).

STAGE PARASITIZED: Nymph (Hemiptera-Heteroptera), adult (Hemiptera-Heteroptera).

IMMATURES: Second instar larva — Smith, R. W., and Finlayson, 1950: (descr., figs. 27, 51).

Pupa — Greene, 1922: (descr., fig. 16 on pl. 4).

BIOLOGY: Morrill, W. L., 1969: (240); Smith, R. W., and Finlayson, 1950: (98-100); Taylor, 1949: (991).

Leucostoma spp.

HOST: HEMIPTERA (HETEROPTERA)

Lygaeidae

Lygaeus kalmii Stål

Simanton and Andre, 1936: 106, Iowa.

STAGE PARASITIZED: Adult.

Linnaemya Robineau-Desvoidy (Tachininae: *Linnaemyini*)

Linnaemya anthracina, see *Thompsonomyia anthracina*

Linnaemya comta, see *Bonnetia comta*

Linnaemya fulvicauda, see *Gymnochaetopsis fulvicauda*

"*Linnaemya picta* Meigen"

HOST: LEPIDOPTERA

Noctuidae

Agrotis vetusta (Walker)

All records ex *Porosagrotis vetusta*.

*Crumb, 1929: 40, 89 (as *Linnaemyia*, Chittenden, 1912);
Chittenden, 1912: 51, N.C., Va.

STAGE PARASITIZED: Larva.

Linnaemya tessellata (Brooks)

All records as *Bonellimyia tessellata*.

HOST: LEPIDOPTERA

Noctuidae

Amathes smithii (Snellen)

Wood and Neilson, 1957: 503, N.B. (ex *Graphiphora*);
Brooks, 1944: 198-200, fig. 3, Wash. (ex *Agrotis*).

IMMATURES: First instar larva — Brooks, 1944: (descr., fig. 3).

Linnaemyia "compta," see *Bonnetia comta*

Linnaemyia comta, see *Bonnetia comta*

Linnaemyia fulvicauda, see *Gymnochaetopsis fulvicauda*

Linnaemyia picta, see "*Linnaemya picta*"

Lispidea palpigera, see *Elfia palpigera*

Lispidea sp., see *Elfia* spp.

"*Lispidia*" sp. near *palpigera*, see *Elfia* sp. near *palpigera*

"*Lixephaga distraeae*," see *Lixophaga diatraeae*

Lixophaga Townsend (Goniinae: *Blondeliini*)

Lixophaga "diatraea," see *Lixophaga diatraeae*

Lixophaga diatraeae (Townsend)

HOST: LEPIDOPTERA

Arctiidae

Utetheisa venusta Dalman

Scaramuzza, 1932: 90 (unsuccessful lab. rearing).

Hesperiidae

Calpodes ethlius (Stoll)

Scaramuzza, 1932: 90 (lab.)

Noctuidae

Spodoptera frugiperda (J. E. Smith)

Scaramuzza, 1932: 90 (ex *Laphygma*, lab.).

Pyralidae

Diatraea canella Hampson

Box, 1939: 228 (lab.).

Diatraea crambidoides (Grote)

Holloway and Loftin, 1919: 41, 61-62, La., Cuba (as *Euzenilliopsis* ex *D. saccharalis* var. *crambidoides*).

Diatraea saccharalis (Fabricius)

Allen, 1962: 9; Miskimen, 1962: 138, Virgin Islands; Anon., 1960: 211 (lab.); Anon., 1960a: 58-61, Antigua, Jamaica, St. Kitts (as *L. "diatraea"*); Anon., 1960b: 8, Fla. (as "*Lixophoga*"); Box, 1960: 901-913, Antigua, Haiti, St. Kitts, St. Lucia; Charpentier, McCormick and Mathes, 1960: 865-868, La.; Scaramuzza, 1960: 939-941, Cuba; Scaramuzza and Barry, 1960: 998, Cuba; Charpentier, 1958: 163-164, La.; Charpentier, 1956: 267-268, La.; Clausen, 1956: 4-5, 44-45, 110-111, Fla.; Dugas, Miller and Concienne, 1956: 93, La. (ex sugarcane borer); Simmonds, 1956: 554-555, St. Kitts; Wolcott, 1956: 311, P.R.; Dugas, Concienne, Sibley and Joyner, 1955: 88, La. (ex sugarcane borer); Scaramuzza, 1955: 319-320, Cuba (probable host); Simmonds, 1955: 199, Dominica; Mathes, Ingram, Questel, Thames and Dugas, 1954: 562, Fla.; Simmonds, 1953: 122-124, 126, Guadeloupe; Thames, Mathes, Bregger and Questel, 1952: 208, Fla. (ex sugarcane moth borer); Hayslip, Mathes and Bregger, 1951: 190, Fla.; Ingram, Bynum and Mathes, 1951: 395, Fla.; Ingram, Mathes, Haley and Charpentier, 1951: 7-8, Fla.; Ingram, Mathes, Wilson and Hayslip, 1951: 356-360, Fla., La.; Plank, 1951: 15, P.R.; Scaramuzza, 1951: 282-291, Cuba; Scaramuzza, 1951a: 298, Cuba; Wolcott, 1951: 9-10, P.R.; Wolcott, 1951a: 474-475, P.R.; Anon., 1950: 4 (as *L. "diatraea,"* lab.); Beauchamp, 1950: 1151-1161, Fla., Cuba (ex sugarcane

borer); Plank, 1950: 14, P.R.; Scaramuzza, 1950: 99-100, Cuba; Anon., 1948b: 50, Cuba; Scaramuzza, Fernández Artiles and Fernández Perez, 1948: 37-40 (lab.); Simmonds, 1948: 68, U.S., Antigua, St. Kitts; Scaramuzza, Fernández Artiles and Fernández Perez, 1947: 1-5 (lab.); Wilson, J. W., and Bregger, 1947: 189, Fla. (as *L. "diatraea"* ex sugarcane moth borer); Scaramuzza, 1946: 544, 546, 548, Cuba; Wilson, J. W., 1946: 179, Fla. (as *L. "diatraea"*); Plank, 1945a: 353-355, Cuba; Turner, 1945: 24, Antigua; Annand, 1944: 134, 146, Fla., Cuba (ex sugarcane borer); Hayslip, 1944a: 128, Fla.; Lee, 1942: 67, P.R. (as *L. "diatraea"*); Scaramuzza, 1942: 121-123, Cuba; Scaramuzza and Ingram, 1942: 642-645, Fla.; Wilson, J. W., 1942: 164, Fla. (as *L. "diatraea"*); Ingram, 1941: 800, Fla. (ex sugarcane borer); Ingram and Bynum, 1941: 9, Fla., La.; Wolcott, 1941: 116, P.R.; Tucker, R. W. E., 1940: 462, Cuba, Haiti, Jamaica, Santo Domingo (as *L. "diatraea"*); Box, 1939: 227-228, St. Lucia; Holloway, 1939: 260, Fla., La.; Ingram, Jaynes and Lobdell, 1939: 90, Fla.; Jaynes, 1939: 245-253, Fla., La.; Tucker, R. W. E., 1939: 241 (as *L. "saccharalis" Tucker*, lab.); Lee, 1938: 96-97, P.R.; Bartlett, 1937: 8, P.R. (as *L. "variabilis"* (Coquillett)); Lobdell, 1937: 146, Fla. (as *L. "diatraea,"* lab.); Box, 1935: 474, Cuba (as *Lixophaga* [sp.]); Myers, 1935: 382, Antigua, St. Kitts, St. Lucia; Scaramuzza, 1934: 14, Cuba; Leonard and Séin, 1932: 2, P.R. (as *Lixophaga* [sp.]); Plank, 1932: 1-8, Cuba; Scaramuzza, 1932: 87-91, Cuba; Scaramuzza, 1932a: 60, Cuba; Van Dine and Christenson, 1932: 2, Cuba; Scaramuzza, 1930: 999-1004, Cuba; Plank, 1929: 116-117, Cuba; Plank, 1929a: 107-108, Cuba; Plank, 1929b: 622-629, figs. 2-3, Cuba; Holloway, 1928: 241, La., Cuba (ex sugarcane moth borer); Holloway, Haley, Loftin and Heinrich, 1928: 43, 64-65, La., Cuba; *vide Holloway, Haley, Loftin and Heinrich, 1928: 43, P.R. (as *Tachinophyto* sp., Van Dine, 1913b); Russo, 1927: 91, Cuba or Dominican Republic (as *Euzenilliopsis "diatraea"* ex "*Diatrea*"); Russo, 1927a: 138, Cuba, P.R., Santo Domingo (as "*Lixephaga distraeae*"); Aldrich, 1925c: 134, La., Cuba, Mexico; Menedez Ramos, 1924: 92, P.R. (as *Euzenilliopsis "diatraea"*); Wolcott, 1924: 221, P.R. (as *Euzenilliopsis*); Wolcott, 1924a: 19, Cuba, Jamaica, P.R.

(as *Euzenillioptis*); Van Zwaluwenburg, 1923: 227, Cuba (as *Euzenillioptis* ex *D. "saccharalia"*); Holloway, 1919: 176-177, Cuba (as *Euzenillioptis*, lab.); *vide Holloway and Loftin, 1919: 41, P.R. (as *Tachinophyto* sp. ex *D. saccharalis* var. *crambidoides*, Van Dine, 1913b); Townsend, 1916c: 76, Cuba (as *Euzenillioptis*); Wolcott, 1916: 19, P.R. (as probably *Hypostena* sp.); Van Dine, 1913b: 199, P.R. (as "*Tachinophyta*" sp.).

Diatraea spp.

Bartlett, 1941: 8; Jaynes, 1933: 2, La., Cuba; Van Zwaluwenburg, 1926: 667-668, Cuba, P.R. (as *Euzenillioptis*).

Zeadiatraea lineolata (Walker)

All records ex *Diatraea lineolata*.

Box, 1935: 474, Cuba (as *Lixophaga* [sp.]); Scaramuzza, 1932: 90 (lab.); Aldrich, 1925c: 134, Mexico (probable host).

Unidentified Lepidoptera

Miscellaneous

Flores Caceres, 1958: 245, Cuba (ex barrenadores); James, H. C., 1952: 34, Jamaica (ex small moth borer caterpillars attacking sugar-cane); Townsend, 1916c: 76, La. (as *Euzenillioptis* ex pupa found in cane).

Moth stalkborer(-s)

Holloway, 1927: 67, La.; Plank, 1927: 68, Cuba.

STAGE PARASITIZED: Larva, pupa.

IMMATURES: First instar larva — Scaramuzza, 1930: (descr.).

Second instar larva — Scaramuzza, 1930: (descr.).

Third instar larva — Scaramuzza, 1930: (descr.).

Pupa — Scaramuzza, 1951: (descr.); Bartlett, 1941: (descr.); Scaramuzza, 1930: (descr.); Plank, 1929: (descr.); Plank, 1929a: (descr.); Plank, 1929b: (figs. 2-3).

BIOLOGY: Allen, 1962: (9); Charpentier, 1956: (267-268); Clausen, 1956: (4-5, 44-45, 110-111); Simmonds, 1953: (122-124, 126); Scaramuzza, 1951: (282-291); Scaramuzza, Fernández Artilés and Fernández Perez, 1947: (1-5); Scaramuzza, 1942: (121-123); Jaynes, 1939: (245-253); Scaramuzza, 1934: (14); Scaramuzza, 1930: (999-1004); Holloway, 1928: (241); Wolcott, 1924: (221).

Lixophaga sp. near *fasciata* Curran

HOST: LEPIDOPTERA

Olethreutidae

Petrova albicapitana (Busck)

Turnock, 1953: 242, Man., Minn., Sask.

Lixophaga grisea (Curran)

All records as *Hypostena grisea*.

HOST: LEPIDOPTERA

Pyralidae

Diatraea saccharalis (Fabricius)

Russo, 1927a: 138-139, Jamaica; Curran, 1926: 106-107,
Jamaica.

STAGE PARASITIZED: Larva.

Lixophaga jennei Aldrich

HOST: LEPIDOPTERA

Olethreutidae

Laspeyresia pomonella (Linnaeus)

Aldrich, 1926a: 19, Ark. (ex *Carpocapsa*).

Lixophaga mediocris Aldrich

HOST: LEPIDOPTERA

Gelechiidae

Frumenta nundinella (Zeller)

Fattig, 1949: 20, Ga.

Limaecodidae

Cnidocampa flavescens (Walker)

Brimley, 1938: 359, N.C. (ex Oriental moth).

Olethreutidae

Grapholitha molesta (Busck)

Allen, 1962: 11, Ill., Ind., Ky., Ohio, S.C., Tenn., Va.; Butler,
H. G., 1933: 984-985, Tenn.; Driggers, 1929: 169-170,
N.J. (ex Oriental peach moth).

Laspeyresia caryana (Fitch)

Patton, 1958: 77, Fla.; Phillips, A. M., 1957: 89, Fla. (ex
hickory shuckworm).

Rhyacionia buoliana (Schiffermüller)

Schaffner, 1959: 23, 51, [N.E. U.S.].

Rhyacionia frustrana (Comstock)

DuRant and Hyche, 1967: 406-409, fig. 1, Ala.; Schaffner, 1959: 24, 51, [N.E. U.S.]; Cushman, 1927: 618, Va.; Aldrich, 1925c: 136, Va.

Rhyacionia rigidana (Fernald)

Allen, 1962: 11, N.J.

Suleima helianthana (Riley)

Allen, 1962: 116.

STAGE PARASITIZED: Larva.

IMMATURES: Egg — DuRant and Hyche, 1967: (descr., fig. 1A).

First instar larva — DuRant and Hyche, 1967: (descr., fig. 1A-D).

Second instar larva — DuRant and Hyche, 1967: (descr., fig. 1E, G).

Third instar larva — DuRant and Hyche, 1967: (descr., fig. 1F, H-J).

Pupa — DuRant and Hyche, 1967: (descr., fig. 1K).

BIOLOGY: Allen, 1962: (11); Schaffner, 1959: (23-24, 51); Cushman, 1927: (618).

Lixophaga orbitalis Aldrich

HOST: LEPIDOPTERA

Olethreutidae

Laspeyresia pomonella (Linnaeus)

Most records ex *Carpocapsa pomonella*.

Lloyd, 1944: 461-464, 466-467, 470-471, Calif. (ex codling moth); *vide Aldrich, 1926a: 18, Calif. (Aldrich, 1925c; Townsend, 1916b; Coquillett, 1897); *vide Aldrich, 1925c: 134 (as undescribed *Lixophaga* sp. ex *Carpocapsa* [sp.], Townsend, 1916b; Coquillett, 1897); Townsend, 1916b: 32, Calif. (as possibly western ssp. of *Euzenillia variabilis* (Coquillett)); Coquillett, 1897: 17, 24 (as *Hypostena variabilis*).

STAGE PARASITIZED: Larva.

BIOLOGY: Lloyd, 1944: (461-464, 466-467, 470-471).

Lixophaga "parva," see *Lixophaga parva*

Lixophaga parva Townsend

HOST: COLEOPTERA

Curculionidae

Lixus scrobicollis Boheman

Pierce, Cushman and Hood, 1912: 75, Tex.; Pierce, 1908: 382, 391, Tex.; Townsend, 1908: 86, Tex.

HOST: LEPIDOPTERA

Olethreutidae

Grapholitha molesta (Busck)

Steenburgh and Boyce, 1938: 73, Ont. (as *L. "parua"* ex *Laspeyresia*).

Lixophaga "plumbeae," see *Lixophaga plumbea*

Lixophaga plumbea Aldrich

HOST: LEPIDOPTERA

Olethreutidae

Grapholitha molesta (Busck)

Most records ex *Laspeyresia molesta*.

Allen, 1962: 11, Conn., Ind., N.J., N.Y., Ohio (ex *Grapholitha*); Stearns and Amos, 1941: 247, Del. (ex *Grapholitha*); Steenburgh and Boyce, 1938: 73, Ont.; Steenburgh, 1935: 70-71, Ont.; Steenburgh, 1932: 67-68, Ont. (as *L. "plumbeae"*); Steenburgh, 1931: 59, Ont.; Haeussler, 1930: 367, Va.; Driggers, 1929: 169-170, N.J. (ex Oriental peach moth); Aldrich, 1925c: 135, Va.

Rhyacionia frustrana (Comstock)

Allen, 1962: 11, 116; Cushman, 1927: 618, Va.; Aldrich, 1925c: 135, Va.

Phaloniidae

Phalonia oenotherana Riley

Allen, 1962: 11, 115; Aldrich, 1925c: 135, N.J.

Pyralidae

Pyrausta futilalis (Lederer)

Schaffner, 1959: 11, 51, Mass., N.H., N.J.

Tortricidae

Xenotemna pallorana (Robinson)

Martin, 1958: 47, Ont. (as *Prolixophaga* ex *Tortrix*).

STAGE PARASITIZED: Larva.

BIOLOGY: Schaffner, 1959: (11, 51).

Lixophaga "saccharalis Tucker," see *Lixophaga diatraeae*

Lixophaga sphenophori (Villeneuve)

Most records as *Ceromasia sphenophori*.

HOST: COLEOPTERA

Curculionidae

Rhabdoscelus obscurus (Boisduval)

Most records ex *Rhabdocnemis obscura*.

- Chong, 1969: 278 (ex New Guinea sugarcane borer, lab.); Anon., 1966d: 502, Hawaii (as *Microceromasia* ex *Rhabdoscelus*); Pemberton, 1951: 401, Hawaii (as *Microceromasia* ex *Rhabdoscelus*); Pemberton, 1948a: 63-64, Hawaii (as *Microceromasia* ex cane borer); Pemberton, 1942: 18, Hawaii (as *Microceromasia*); Pemberton, 1941: 22, Hawaii (as *Microceromasia* "spherophori"); Pemberton, 1940: 17-18, Hawaii (as *Microceromasia*); Box, 1939a: 501, Hawaii; Pemberton, 1939: 19-20, Hawaii; Pemberton, 1938: 20, Hawaii (as New Guinea tachinid fly); Pemberton, 1936: 21, Hawaii; Sweetman, 1936: 355, Hawaii; Bryan, 1934: 416-417, 451, Hawaii (as *Microceromasia*); Pemberton, 1934: 22-23, Hawaii; Swezey, 1934: 340, Hawaii; Pemberton, 1933: 15, Hawaii (as New Guinea parasite); Swezey, 1933: 429, Hawaii; Holloway and Rands, 1932: 464, Hawaii; Pemberton, 1932: 47-48, Hawaii; Pemberton, 1932a: 19, Hawaii (as New Guinea parasite); Swezey, 1932: 680, Hawaii; Anon., 1931: 81, Hawaii; Swezey, 1931: 391, Hawaii; Swezey, 1931b: 425-426, Hawaii; Swezey, 1931c: 325, Hawaii; Williams, 1931: 213-216, 294-296, figs. 11-12 on pl. 33, Hawaii; Swezey, 1930: 278, Hawaii; Swezey, 1929a: 269, Hawaii; Swezey, 1928: 671-672, Hawaii; Swezey, 1928b: 118, Hawaii; Agee, 1927: 73, Hawaii (as Tachinid fly); Holloway, 1927: 67, Hawaii (as "*Ceromesia sphenopheri*" ex weevil borer); Timberlake, 1927: 540-544, figs. 8-9 on pl. 21, Hawaii; Muir and Swezey, 1926: 16, Hawaii (as New Guinea tachinid fly ex cane borer); Muir and Swezey, 1925: 13-14, Hawaii (as New Guinea tachinid ex cane borer); Pemberton, 1925: 174-181, 183-185, Hawaii; Swezey, 1925: 372, Hawaii; Pemberton, 1924: 28-29, Hawaii (ex cane borer); Swezey, 1924: 14, Hawaii; Swezey, 1924a: 16-17, Hawaii (as New Guinea Tachinid ex cane borer); Swezey, 1923a: 303, Hawaii; Agee, 1922: 15, Hawaii (as New Guinea Tachinid ex cane borer); Agee, 1919: 163, Hawaii (as Tachinid parasite ex cane borer); Bridwell, 1919: 166, Hawaii; Swezey, 1918: 380, Hawaii; Agee, 1917: 13-14,

Hawaii (as Tachinid fly ex cane borer); Agee, 1916: 12, Hawaii (as Tachinid fly ex cane borer); Agee, 1916a: 341, Hawaii (as Tachinid fly ex cane borer); Agee, 1916b: 4-5, Hawaii (as Tachinid fly ex cane borer); Howard, 1916: 174-176, Hawaii; Howard, 1916a: 293-294, Hawaii (ex "*Rhabadocnemis*"); Howard, 1916b: 367, 369, Hawaii (ex "*Rhabadocnemis*"); Muir and Swezey, 1916: 16-19, 34-43, frontis., pl. 3, Hawaii; Agee, 1915: 16-17, Hawaii (as cane borer tachinid ex cane borer); Agee, 1915a: 5, Hawaii (as Tachinid fly ex cane borer); Agee, 1915b: 337, Hawaii (as Tachinid fly ex cane borer); Swezey, 1915: 455-456, Hawaii; Agee, 1914: 21, Hawaii (as cane borer Tachinid ex cane borer); Agee, 1914a: 5-6, Hawaii (as cane borer Tachinid ex cane borer); Agee, 1914b: 305, 308 [=306], Hawaii (as cane borer Tachinid ex cane borer); Ilingworth, 1914: 390-398, Hawaii; Swezey, 1914: 455-457, Hawaii; Agee, 1913: 9, Hawaii (as cane borer Tachinid ex cane borer); Agee, 1913a: 656, Hawaii (as cane borer Tachinid ex cane borer); Eckart, 1913: 18-20, Hawaii (as cane borer Tachinid ex cane borer); Eckart, 1912: 20-21, Hawaii (as cane borer parasites ex cane borer); Eckart, 1912a: 345-347, Hawaii (as cane borer Tachinid ex cane borer); Eckart, 1912b: 9-11, Hawaii (as cane borer Tachinid ex cane borer); Eckart, 1911: 325-327, Hawaii (as cane borer parasite ex cane borer); Paxton, Waldron, Bottomley, Dowsett and Gartley, 1911: 322, Hawaii (as cane borer parasites ex cane borer); Swezey, 1911: 185, Hawaii (as cane borer parasite ex cane borer); Perkins, R. C. L., 1910: 292-295 (as parasitic fly ex cane borer).

HOST: LEPIDOPTERA

Pyralidae

Diatraea crambidoides (Grote)

Holloway and Loftin, 1919: 60-61, La. (ex *D. saccharalis* var. *crambidoides*, unsuccessful lab. rearing).

STAGE PARASITIZED: Larva (Coleoptera), pupa (Coleoptera).

IMMATURES: Third instar larva — Williams, 1931: (fig. 11 on pl. 33); Timberlake, 1927: (fig. 8 on pl. 21); Muir and Swezey, 1916: (frontis.).

Pupa — Williams, 1931: (fig. 12 on pl. 33); Timberlake, 1927: (fig. 9 on pl. 21); Muir and Swezey, 1916: (frontis., pl. 3).
 BIOLOGY: Williams, 1931: (213-216, 294-296); Howard, 1916: (174-176); Howard, 1916a: (293-294); Howard, 1916b: (367, 369); Muir and Swezey, 1916: (16-19, 34-43); Illingworth, 1914: (390-398); Swezey, 1914: (455-457).

Lixophaga stenomae Curran
 HOST: LEPIDOPTERA

Stenomidae

Stenoma spp.

All records ex *Stenoma* sp. on avocado.

Arnaud, 1963: 120, Costa Rica; Curran, 1935: 24, Costa Rica.

Lixophaga "variabilis," see *Lixophaga diatraeae*

Lixophaga variabilis (Coquillett)

HOST: COLEOPTERA

Bostrichidae

Amphicerus bicaudatus (Say)

Allen, 1962: 113; Aldrich, 1925c: 134, Md. (ex apple twig borer).

Cerambycidae

Oberea bimaculata (Olivier)

Allen, 1962: 113; Aldrich, 1925c: 134, D.C. (ex *O. "maculata"*).

Curculionidae

Lixus scrobicollis Boheman

Allen, 1962: 113 (ex *L. "scrobicollis"*).

Sphenophorus pontederiae Chittenden

Aldrich, 1925c: 134, Mass.

HOST: HYMENOPTERA

Tenthredinidae

Unidentified Tenthredinidae

Coquillett, 1897: 17, 22, Mo. (as *Hypostena*).

Unidentified Hymenoptera

Sawfly(-flies)

Essig, 1926: 579, Canada.

HOST: LEPIDOPTERA

Noctuidae

Anomis erosa Hübner

Allen, 1962: 118; Satterthwait, 1948: 729.

Papaipema cataphracta (Grote)

Franklin, 1910: 94, Minn. (as *Hypostena*); Washburn, 1910: 168, [Minn.] (as *Hypostena*); Washburn, 1910a: 94, Minn. (as *Hypostena*).

Papaipema nebris (Guenée)

Allen, 1962: 118; Allen, 1962: 118 (ex *P. nebris nitela* (Guenée)); Decker, 1935: 570, 578, Iowa; Decker, 1931: 344, Iowa (as "*Lixus variabili*" ex *Papaipema* [sp.]); Lowry, 1927: 22 (as *Hypostena*); Washburn, 1910a: 94, Minn. (as *Hypostena* ex *P. nitela*).

Stibadium spumosum Grote

Allen, 1962: 118; Satterthwait, 1948: 727, 729.

Trichoplusia ni (Hübner)

Allen, 1962: 118.

Olethreutidae

Ancylis comptana (Frölich)

McConnell, 1932: 368-369, Md.

Epiblema otiosanum (Clemens)

Decker, 1935: 570, 578, Iowa; Decker, 1933: 509, Iowa.

Epiblema otiosanum (Clemens) probably

Allen, 1962: 10, 116, N.J.

Epiblema scudderiana (Clemens)

Allen, 1962: 10, N.J.

Epiblema strenuanum (Walker)

Allen, 1962: 10, 116, N.J.; Decker, 1935: 570, 578, Iowa; Montgomery, 1933: 189, Ind.; McConnell, 1932: 368, Md.; Allen, 1929: 683, Miss. (ex "*Epiblana*"); Aldrich, 1925c: 134, Ga.

Eumaroza malachitana (Zeller)

Bottimer, 1926: 817, Tex. (ex *Olethreutes*).

Grapholitha molesta (Busck)

Most records ex *Laspeyresia molesta*.

Allen, 1962: 9-10, Conn., Ill., Ky., Mo., N.Y., Ohio, S.C. (ex *Grapholitha*); Fattig, 1949: 20 (ex *Grapholitha*); Stearns and Amos, 1941: 247, Del. (ex *Grapholitha*); McConnell, 1932: 367-369, Md.; Haeussler, 1930: 367, Ga., Md., N.J., Va.; Snapp and Swingle, 1929: 12, Ga.; Stearns, 1928: 4-5, 7, Ga., Md., N.J., Va.; Aldrich, 1925c: 133, Va.; Stearns, 1919: 348 (as *Euzenillia*); Garman, P., 1918: 117, Md. (as *Euzenillia*); Wood, W. B., 1918: 527, E. U.S. (as *Hypostena*); Wood, W. B., and Selkregg, 1918: 70, Va. (as *Hypostena*).

Laspeyresia caryana (Fitch)

Allen, 1962: 116.

Laspeyresia pomonella (Linnaeus)

Most records ex *Carpocapsa pomonella*.

Allen, 1962: 115; Lloyd, 1944: 468-469, Calif., cent. Appalachians (ex codling moth); Hensill, 1933: 603, Calif. (ex codling moth); Stearns, 1928: 4; Essig, 1926: 579, Calif., Canada (ex codling moth); Aldrich, 1925c: 134, Calif.; Johnson, 1925: 188, Conn., Maine, Mass., N.H. (ex "coddling" moth); Brooks, F. E., and Blakeslee, 1915: 47, W. Va. (as *Tachinophyto*); Quaintance, 1908: 443 (as *Hypostena* ex codling moth); Simpson, 1903: 59 (as *Hypostena*).

Rhyacionia rigidana (Fernald)

Allen, 1962: 10, N.J.

Suleima helianthana (Riley)

Satterthwait, 1948: 726, 729; Satterthwait and Swain, 1946: 577, Mo.

Pyralidae

Desmia funeralis (Hübner)

Allen, 1962: 117; Essig, 1926: 579 (ex grape leaf folder); Strauss, 1916: 9, D.C. (as *Tachinophyto*).

Homoeosoma electellum (Hulst)

Satterthwait and Swain, 1946: 577, Mo.

Loxostege similalis (Guenée)

Allen, 1929: 683, Miss.

Ostrinia nubilalis (Hübner)

All records ex *Pyrausta nubilalis*.

Allen, 1962: 117; Wishart, 1943: 27, Ont.

Ostrinia penitalis (Grote)

All records ex *Pyrausta penitalis*.

Allen, 1962: 117; Decker, 1935: 570, Iowa; Poos, 1927: 83, Ohio; Johnson, 1925: 188, Conn., Maine, Mass., N.H.; Chittenden, 1918: 456 (as *Hypostena*); Johnson, 1910: 776, N.J. (as *Hypostena*).

Ostrinia spp. (including *O. oblitalis* (Walker) or *O. penitalis* (Grote))

*Ainslie, G. G., and Cartwright, 1921: 843 (as *Hypostena* ex *Pyrausta penitalis* or *P. ainsliei* Heinrich, Coquillett, 1897); Coquillett, 1897: 17, 27, D.C. (as *Hypostena* "variabilis" ex *Pyrausta penitalis*).

Unidentified Pyralidae

Essig, 1926: 579, Canada.

Unidentified Lepidoptera

Miscellaneous

Aldrich, 1925c: 133, Tenn.

STAGE PARASITIZED: Larva (Hymenoptera, Lepidoptera), pupa (Lepidoptera)

IMMATURES: Third instar larva — Satterthwait and Swain, 1946: (descr.).

Pupa — Satterthwait and Swain, 1946: (descr.); Greene, 1922: (descr., fig. 35 on pl. 8).

BIOLOGY: Allen, 1962: (9-10, 116).

Lixophaga sp. near *variabilis* (Coquillett)

HOST: COLEOPTERA

Curculionidae

Hypera nigrirostris (Fabricius)

Webster, 1909: 11, Md. (as *Anisia* ex *Phytonomus*).

STAGE PARASITIZED: Larva, pupa.

Lixophaga spp., see *Lixophaga orbitalis* or *Lixophaga* spp.

Lixophaga spp.

HOST: COLEOPTERA

Unidentified Coleoptera

Rhynchophora

Townsend, 1913a: 303-304.

HOST: LEPIDOPTERA

Olethreutidae

Laspeyresia pomonella (Linnaeus)

All records ex codling moth.

*Lloyd, 1944: 470, Calif. (Flanders, 1926a); Flanders, 1926: 158, Calif.; Flanders, 1926a: 44, Calif.

Diatraea impersonatella (Walker)

Box, 1960: 910, Guadeloupe (probable parasite); Fretay, 1960: 142, 145, Guadeloupe (ex "*Diatrea*"); Fretay, 1960a: 148, 151, Guadeloupe.

Diatraea saccharalis (Fabricius)

Fretay, 1960: 142, 145, Guadeloupe (ex "*Diatrea*"); Fretay, 1960a: 148, 151, Guadeloupe; Box, 1939a: 497-513, St. Kitts.

STAGE PARASITIZED: Larva (Coleoptera, Lepidoptera), adult (Coleoptera).

BIOLOGY: Flanders, 1926: (158); Townsend, 1913a: (303-304).

Gen. and sp. near *Lixophaga*

HOST: HYMENOPTERA

Unidentified Hymenoptera

Sawfly(-flies)

Procter, 1946: 436, Maine.

"*Lixophoga*" diatraeae, see *Lixophaga diatraeae*

"*Lixus variabili*," see *Lixophaga variabilis*

Long-legged Tachina fly, see Tachinidae

Lophosiocera sp., see *Elfia* spp.

Lydella Robineau-Desvoidy (Goniinae: Eryciini)

Lydella deckeri (Curran)

All records as *Erycia deckeri*.

HOST: LEPIDOPTERA

Noctuidae

Papaipema nebris (Guenée)

Arnaud, 1963: 113, Iowa; Decker, 1935: 569, 578, Iowa; Curran, 1929: 13, Iowa.

STAGE PARASITIZED: Larva.

Lydella delecta, see *Pilatea delecta*

Lydella "doryphora," see *Doryphorophaga doryphorae*

Lydella doryphorae, see *Doryphorophaga doryphorae*

Lydella grisescens, see *Lydella thompsoni*

Lydella nigrita, see *Lydella radialis*

Lydella radialis (Townsend)

HOST: LEPIDOPTERA

Noctuidae

Achatodes zae (Harris)

Decker, 1935: 568, 578, Iowa (as *Masicera senilis* Meigen).

Archanara oblonga (Grote)

Most records ex *Archanara subcarnea* (Kellcott).

Decker, 1935: 568, 578, Iowa (as *Masicera senilis*); Cole, 1931: 11, 39, Mich., Ohio (as *Masicera senilis*); Cole, 1929: 225, Mich. (as *Masicera senilis* ex "Arachnara"); Townsend, 1916a: 19, Pa. (as *Andrina* ex *Nonagria oblonga*).

Arzama densa Walker

Vogel and Oliver, 1969: 752, La.

Arzama obliqua (Walker)

Guppy, R., 1950: 4, B.C. (as *L. nigrita* Townsend); Schaffner and Griswold, 1934: 62-63, 119, Mass. (as *Ceromasia senilis*); Cole, 1931: 11, 39, Idaho (as *Sturmia nigrita*); Claassen, 1921: 503, N.Y. (as *Sturmia nigrita*); Coquillett, 1897: 18, 24 (as *Masicera myoidea* (Robineau-Desvoidy) ex *A. "obliquata"*).

Hydroecia immanis Guenée

Hawley, 1918: 169, N.Y. (as *Masicera "myoidea"* ex *Gortyna*).

Luperina stipata (Morrison)

Baker, W. A., Bradley and Clark, 1949: 53-54 (as *L. nigrita*); Decker, 1935: 568, 578, Iowa (as *Masicera senilis*); Decker, 1930: 159, Iowa (as *Masicera senilis*).

Macronoctua onusta Grote

*Vide Baker, W. A., Bradley and Clark, 1949: 53-54 (as *L. nigrita* ex "Macronoctus," Breakey, 1931); Decker, 1935: 568, 578, Iowa (as *Masicera senilis*); Schaffner and Griswold, 1934: 53, 119, Mass. (as *Ceromasia senilis*); Breakey, 1931: 43, Wis. (as *Masicera senilis*); Decker, 1930: 159, Iowa (as *Masicera senilis*); Breakey, 1929: 460, 462, Wis. (as *Masicera senilis*).

Papaipema appassionata (Harvey)

Most records ex *Papaipema* "appassionata."

Curran, 1927a: 19, Ont. (as *Erycia myoidaea* ex *P. appassionata*); Winn and Beaulieu, 1915: 142, Que. (as *Masicera* "myoidea"); Fletcher, 1907: 102, Ont. (as *Masicera myoidaea*).

Papaipema araliae Bird and Jones

Bird, H., and Jones, 1921: 139, Del. (as *Masicera* "seniles").

Papaipema arctivorens Hampson

Decker, 1935: 568, 578, Iowa (as *Masicera senilis*).

Papaipema cataphracta (Grote)

Decker, 1935: 568, 578, Iowa (as *Masicera senilis*); Decker, 1930: 159, Iowa (as *Masicera senilis*); Washburn, 1910a: 94, Minn. (as *Masicera myoidaea*).

Papaipema frigida (Smith)

Decker, 1935: 568, 578, Iowa (as *Masicera senilis*).

Papaipema nebris (Guenée)

*Vide Baker, W. A., Bradley and Clark, 1949: 53-54 (as *L. nigrita*, Decker, 1931; Lowry, 1927); Decker, 1935: 567-569, 578, Iowa (as *Masicera senilis*); Schaffner and Griswold, 1934: 62, 119 (as *Ceromasia senilis*); Schaffner and Griswold, 1934: 62, 118, Mass., N.H. (as *Sturmia nigrita*); Decker, 1931: 343-344, Iowa (as *Masicera senilis*); Decker, 1930: 159, Iowa (as *Masicera senilis*); Lowry, 1927: 22, N.H. (as *Masicera senilis*); Lowry, 1927: 22 (as *Masicera myoidaea*); Bird, H., 1925: 305 (as *Masicera senilis*); Johnson, 1925: 201, Maine, Mass. (as *Paraphorocera myoidaea* ex *P. nitela* (Guenée)); Britton, 1920: 182, Conn. (as *Masicera* "myoidea" ex *P. nitela*); Vinal and Caffrey, 1919: 58, Mass. (as *Masicera* "myoidea" ex *P. nitela*); Franklin, 1912: 94, Minn. (as *Masicera myoidaea* ex *P. nitela*); Washburn, 1910: 168 (as *Masicera myoidaea* ex *P. nitela*); Washburn, 1910a: 94, Minn. (as *Masicera myoidaea* ex *P. nitela*); Coquillett, 1897: 18, 25, Mass. (as *Masicera myoidaea* ex *Hydroecia nitela*); Brauer and Bergenstamm, 1895: 561, 593, N. Amer. (as (*Masicera*) *nigrita* ex *Gortyna nitela*).

Papaipema pterisii Bird

Bird, H., 1927: 95 (as *Masicera senilis*).

Papaipema purpurifascia (Grote and Robinson)

Decker, 1935: 568, 578, Iowa (as *Masicera senilis*); Tothill, 1913: 73, Ont. (as *Masicera* "myoidea").

Papaipema stenoscielis Dyar

Bird, H., 1936: 120 (as *L. senilis*).

Papaipema spp.

Decker, 1930: 159, Iowa (as *Masicera senilis*).

Olethreutidae

Epiblema otiosanum (Clemens)

Decker, 1935: 568, 578, Iowa (as *Masicera senilis* ex "*Epibleme*"); Decker, 1933: 509, Iowa (as *Masicera senilis*).

Pyralidae

Ostrinia nubilalis (Hübner)

Most records ex *Pyrausta nubilalis*.

Johnson, 1925: 201, Maine, Mass. (as *Paraphorocera myoidea* ex European corn-borer); Bird, H., 1923a: 200 (as *Masicera senilis* ex European corn borer); Thompson, 1923: 34, Mass. (as *Andrina* ex *P. nubilalis*"); Caffrey, 1919: 21, Mass. (as *Masicera "myoidea"*); Vinal and Caffrey, 1919: 57-58, Mass. (as *Masicera "myoidea"*).

Unidentified Lepidoptera

Miscellaneous

Townsend, 1916a: 19 (as *Andrina* ex larvae boring in roots of plants).

STAGE PARASITIZED: Larva.

IMMATURES: Pupa — Greene, 1922: (descr., figs. 80-81 on pl. 17).

BIOLOGY: Guppy, R., 1950: (4); Decker, 1935: (567-569, 578); Schaffner and Griswold, 1934: (62, 118-119); Decker, 1931: (343-344); Breakey, 1929: (460, 462); Lowry, 1927: (22); Claassen, 1921: (503).

Lydella radialis (Townsend) probably

HOST: LEPIDOPTERA

Noctuidae

Achatodes zea (Harris)

Silver, J. C., 1933: 17, Mich. (as *Masicera* sp. probably "*myoidea*" (Robineau-Desvoidy)).

Papaipema nebris (Guenée)

Townsend, 1893: 467, Ill. (as *Masicera nigrita* Townsend ex *Gortyna nitela* Guenée).

Papaipema spp.

Bird, H., 1909: 66 (as *Masicera myoidaea*).

STAGE PARASITIZED: Larva.

BIOLOGY: Silver, J.C., 1933: (17).

Lydella senilis, see *Lydella radialis*

Lydella stabulans, see *Lydella thompsoni*

Lydella stabulans "gresecens," see *Lydella thompsoni*

Lydella stabulans *grisescens*, see *Lydella thompsoni*

Lydella thompsoni Herting

HOST: LEPIDOPTERA

Noctuidae

Papaipema nebris (Guenée)

Hsiao and Holdaway, 1966: 125-133, Minn. (as *L. grisescens* Robineau-Desvoidy); Hsiao and Holdaway, 1962: 144 (as *L. grisescens*); York, Schaffner and Brindley, 1955: 765, Iowa (as *L. stabulans grisescens*).

Pyralidae

Galleria mellonella (Linnaeus)

Hsiao and Holdaway, 1962: 144 (as *L. grisescens*, lab.).

Ostrinia nubilalis (Hübner)

Most records ex European corn borer.

Petty, 1967: 29, Ill. (as *Lydella* sp. ex *O. nubilalis*); Franklin, R. T., and Holdaway, 1966: 440-441, Minn. (as *L. grisescens* ex *O. nubilalis*); Hsiao and Holdaway, 1966: 125-133, Minn. (as *L. grisescens* ex *O. nubilalis*); Anon., 1963d: 1029, Wis. (as *L. grisescens* ex *O. nubilalis*); Anon., 1963e: 1337, Md. (as *L. grisescens*); Sparks, Raun and Carter, 1963: 20-21, Iowa (as *L. grisescens* ex *O. nubilalis*); Anon., 1962a: 16-17, U.S. (as *L. grisescens* ex *O. nubilalis*); Anon., 1962b: 266, Kans. (as *L. grisescens*); Hsiao and Holdaway, 1962: 144 (as *L. grisescens* ex *O. nubilalis*); Van Denburgh, Burbutis and York, 1962: 11-14, Del. (as *L. stabulans grisescens* ex *O. nubilalis*); Anon., 1961c: 280, Del. (as *L. grisescens*); Anon., 1961d: 639, Wis. (as *L. grisescens*); Jarvis and York, 1961: 213-214, Iowa (as *L. grisescens* ex *Pyrausta nubilalis*); Manson, Wressell, Begg and Miller, 1960: 53, Ont. (as *Lydella* sp. ex *O. nubilalis*); Wressell, Driscoll and Svec, 1959: 240, Ont. (as *Lydella* sp. ex *O. nubilalis*); Anon., 1958d: 194,

Wis. (as *L. grisescens* ex *Pyrausta nubilalis*); Anon., 1958f: 668, Wis. (as *L. grisescens* ex *Pyrausta nubilalis*); Anon., 1958g: 755, Wis. (as *L. grisescens* ex *Pyrausta nubilalis*); Jackson and Peters, 1958: 626, Mo. (as *Lydella* sp. ex *Pyrausta nubilalis*); Anon., 1957d: 240, Wis. (as *L. grisescens* ex *Pyrausta nubilalis*); Anon., 1957e: 460, Wis. (as *L. grisescens* ex *Pyrausta nubilalis*); Anon., 1956c: 274, Wis. (as *L. stabulans grisescens* ex *Pyrausta nubilalis*); Clausen, 1956: 7, 42-43, 111-112, E., mid-Atlantic, N. cent. U.S. (as *L. stabulans grisescens* ex *Pyrausta nubilalis*); Anon., 1955: 17 (as *L. stabulans* var. *grisescens* ex *Pyrausta nubilalis*); Arbuthnot, 1955: 91-93, Conn. (as *L. stabulans grisescens* ex *Pyrausta nubilalis*); Udine, 1955: 722, Pa. (as *L. stabulans*, probable parasite, incomplete rearing — pupa); York, 1955: 65, Ark., N. Cent. U.S., Okla., Tenn. (as *L. grisescens* ex *Pyrausta nubilalis*); York, Schaffner and Brindley, 1955: 765-766, Iowa (as *L. stabulans grisescens* ex *Pyrausta nubilalis*); Anon., 1954i: 438, Minn. (as *L. stabulans grisescens*); Harris, 1954: 152, Iowa (as *Lydella* flies); Haut, 1954: 61-62, Md. (as *L. stabulans grisescens*); Lannon, 1954: 70, R.I. (as *L. stabulans grisescens* ex *Pyrausta nubilalis*); Parks and Goleman, 1954: 500, Ohio (as *L. stabulans grisescens* ex *Pyrausta nubilalis*); Weiss, 1954: 196, N.J. (as *L. grisescens*); Welch, 1954: 54-55, Ky. (as *L. stabulans* “*gresecens*”); Arbuthnot, 1953: 30-31 (as *L. stabulans grisescens*); Bissell, 1953: 127, Md. (as *L. stabulans grisescens* ex *Pyrausta* “*nubilalis*”); Blickenstaff, Arbuthnot and Harris, 1953: 339-353, Iowa (as *L. stabulans grisescens* ex *Pyrausta nubilalis*); Boyes and Wilkes, 1953: 126, 146, Ont. (as *L. grisescens* ex *Pyrausta nubilalis*); Burkhardt, 1953: 67, Kans. (as *L. stabulans grisescens*); Hibbs, 1953: 58-59, Ohio (as *L. stabulans grisescens*); Parks, 1953: 670, Ohio (as *L. stabulans grisescens* ex *Pyrausta nubilalis*, incomplete rearing — pupa); Weiss, 1953: 192, N.J. (as *L. grisescens*); Welch, 1953: 49, Ky. (as *L. stabulans grisescens*); Anon., 1952a: 287, 290, 294, E. U.S. (as *L. grisescens* ex *Pyrausta nubilalis*); Arbuthnot, 1952: 19-21, Ill., Ind., Iowa, Kans., Ky., Minn., Mo., N. Dak., Nebr., Ohio, S. Dak., Wis. (as *L. stabulans grisescens*); Arbuthnot and Wright, 1952: 223, 225-230, Ill., N.J. (as *L. stabulans grisescens* ex

Pyrausta nubilalis); Chambers, 1952: 233, Wis. (as *Lydella stabulans grisescens* ex *Pyrausta nubilalis*); Chambers, 1952a: 267, Wis. (as *Lydella stabulans grisescens* ex *Pyrausta nubilalis*); Chambers, 1952b: 357, Wis. (as *L. stabulans grisescens* ex *Pyrausta nubilalis*); Cory, McConnell, Arbuthnot and Jones, 1952: 2-7, Md. (as *L. stabulans grisescens* ex *Pyrausta nubilalis*); Decker, 1952: 431, Ill. (as *L. stabulans grisescens* ex *Pyrausta nubilalis*); Harris, 1952: 143-144, Iowa (as *L. stabulans grisescens*); Haut, 1952: 50, Md. (as *L. stabulans grisescens*); Lannon, 1952: 16, R.I. (as *L. stabulans grisescens* ex *Pyrausta nubilalis*); Munro, Post and Nelson, 1952: 128, N. Dak. (as *Lydella* [sp.]); Weiss, 1952: 181, N.J. (as *L. grisescens*); Welch, 1952: 38, Ky. (as *L. stabulans grisescens*); Wilson, M. C., 1952: 161, Ind. (as *L. stabulans grisescens* ex *Pyrausta nubilalis*); Andre, 1951: 39-40, Iowa (as *L. stabulans grisescens*); Cooper, 1951: 18, Ky. (as *L. stabulans grisescens*); Harris, 1951: 138-141, Iowa (as *L. stabulans grisescens*); Haut, 1951: 48, Md. (as *L. stabulans grisescens*); Weiss, 1951: 163, N.J. (as *L. grisescens*); Wilson, M. C., 1951: 222-225, Ind. (as *L. stabulans grisescens*); Wilson, M. C., 1951a: 41, Ind. (as *L. grisescens*); Arbuthnot, 1950: 44-46, Ill., Ind., Mich., Minn., Mo., Wis. (as *L. stabulans grisescens*); Arbuthnot, 1950a: 423-424 (as *L. stabulans grisescens* ex *Pyrausta nubilalis*); Cooper, 1950: 14, Ky. (as *L. stabulans grisescens*); Cory, Foster, McConnell, Bissell, Langford and Ditman, 1950: 22-24, fig. 14, Md. (as *L. stabulans grisescens* ex *Pyrausta "vubilalis"*); Harris, 1950: 141, 146-159, Iowa (as *L. grisescens*); Harris and Lilly, 1950: 247, Iowa (as *L. grisescens*); Weiss, 1950: 170, N.J. (as *L. grisescens*); Wilson, M. C., 1950: 11, Ind. (as *L. grisescens* ex corn borer); Wishart, 1950: 48, Ont. (as *L. grisescens*); Arbuthnot, Jones, Carter and Evans, 1949: 2-14, 16-24, Conn., Del., Ill., Ind., Iowa, Ky., Mass., Md., Mich., Minn., Mo., N.C., N.J., N.Y., Ohio, Pa., R.I., Va., Wis. (as *L. stabulans grisescens* ex *Pyrausta nubilalis*); Baker, W. A., Bradley and Clark, 1949: 3, 6, 8-9, 12, 21, 23-24, 26-27, 38-40, 53-56, 166-170, 182, fig. 38B, Conn., Mass., Ohio (as *L. stabulans grisescens* ex *Pyrausta nubilalis*); Cooper, 1949: 26, Ky. (as *L. stabulans* var. *grisescens*); Harris,

1949: 129-131, Iowa (as *L. grisescens*); Jones, D. W., and Arbuthnot, 1949: 57, Ill., Iowa, Ky., Ohio (as *L. stabulans grisescens*); MacCreary, 1949: 22, Del. (as *L. grisescens*); MacCreary and Rice, 1949: 141-153, Del. (as *L. stabulans grisescens ex Pyrausta nubilalis*); Weiss, 1949: 157, N.J. (as *L. grisescens*); Wilson, M.C., 1949: 83, Ind. (as *L. stabulans grisescens*); Wilson, M.C., 1949a: 157-159, Ind. (as *L. grisescens*); Anon., 1948: 284, 290, 294, Ont., U.S. (as *L. grisescens ex Pyrausta nubilalis*); Anon., 1948a: 354, 361, 368, U.S. (as *L. grisescens ex Pyrausta nubilalis*); Arbuthnot, Jones, Carter and Evans, 1948: 1-13, Del., Ill., Ind., Iowa, Ky., Md., Mich., Minn., Mo., N.C., New England, N.J., N.Y., Ohio, Pa., Va., Wis. (as *L. grisescens ex Pyrausta nubilalis*); Cooper, 1948: 41, Ky. (as *L. stabulans var. grisescens*); Harris, 1948: 129-138, Iowa (as *L. grisescens*); MacCreary, 1948: 21-22, Del. (as *L. grisescens*); Weiss, 1948: 170, N.J. (as *L. grisescens*); Wilson, M. C., 1948: 10-11, Ind. (as *L. grisescens*); Anon., 1947: 185, 192, 198, U.S. (as *L. grisescens ex Pyrausta nubilalis*); Arbuthnot, 1947: 30-32, Ind., Iowa, Ky., Mich., Ohio (as *L. grisescens*); Arbuthnot, Jones, Carter and Cosenza, 1947: 1, E. U.S. (as *L. stabulans var. grisescens*); Clark, N., and Hoveland, 1947: 38, Wis. (as *L. grisescens ex corn borer*); Cooper, 1947: 28, Ky. (as *L. stabulans grisescens*); Deay, 1947: 68, Ind. (as *L. grisescens*); Drake, 1947: 121-129, Iowa (as *L. grisescens*); Jones, D. W., Carter and Cosenza, 1947: 2-6, Del., Ill., Ind., Ky., Mass., Mich., New England, N.J., N.Y., Ohio, Va., Wis. (as *L. grisescens*); MacCreary, 1947: 26, Del. (as *L. grisescens*); Ritcher, 1947: 19-20, 22, fig. 17, Ky. (as *L. stabulans grisescens ex Pyrausta nubilalis*); Weiss, 1947: 161, N.J. (as *L. grisescens*); Wright, 1947: 33, Ill. (as *L. grisescens*); Clark, C. A., and Jones, 1946: 2-10, Conn., Del., Ill., Ind., Iowa, Ky., Mass., Mich., N.C., N.J., N.Y., Ohio, Pa., Va., Wis. (as *L. grisescens*); Cooper, 1946: 30, Ky. (as *L. grisescens*); Cory, 1946: 147-148 (as *L. grisescens*); Deay, 1946: 11, Ind. (as *L. grisescens*); Drake, 1946: 171-176, Iowa (as *L. grisescens*); Weiss, 1946: 153, N.J. (as *L. grisescens*); Clark, C. A., 1945: 1-5, Conn., Del., Mass., Mich., N.J., Ohio, R.I., Va. (as *L. grisescens*); Clark, C. A., 1945a: 1-2, Conn., Mass., N.J. (as *L. grisescens*); Clark, C. A., and Jones, 1945:

1-2, Conn., Mass., N.J. (as *L. grisescens*); Drake, Harris and Blickenstaff, 1945: 100-107, Iowa (as *L. grisescens*); Drake, Harris and Tauber, 1945: 68, Iowa (as *L. grisescens*); Gardiner, 1945: 66, Ont. (as *L. stabulans* Robineau-Desvoidy ex *Pyrausta nubilalis*); Rice and Wolfenbarger, 1945: 21, Del. (as *L. grisescens*); Gardiner, 1944: 66, Ont. (as *L. stabulans* ex *Pyrausta nubilalis*); Rice, Wolfenbarger and Chada, 1944: 23-24, Del. (as *L. grisescens* ex *Pyrausta nubilalis*); Wishart, 1944: 12-13, Ont. (as *L. grisescens* ex *Pyrausta nubilalis*); Clark, C. A., 1943: 1-3, Conn., Mass., N.J., Ohio, Va. (as *L. grisescens* ex *Pyrausta nubilalis*); Clark, C. A., 1943a: 1-2, Mass., N.J. (as *L. grisescens*); Clark, C. A., 1943b: 2-8, Conn., Mass., Mich., N.J., Ohio, S.E. New England, Va. (as *L. grisescens*); Clark, C. A., and Carter, 1943: 120, Conn., Mass. (as *L. grisescens*); Gardiner, 1943: 55, Ont. (as *L. stabulans* ex *Pyrausta nubilalis*); Wishart, 1943: 26-29, Ont. (as *L. stabulans* ex *Pyrausta nubilalis*); Gardiner, 1942: 58, Ont., Que. (as *L. stabulans* var. *grisescens* ex *Pyrausta nubilalis*); Bradley, W. G., and Clark, 1941: 788-790, 793-794, Conn., Mass., Mich., N.J., Ohio, Va. (as *L. stabulans* var. *grisescens*); Clark, C. A., 1941: 621, Conn., Mass. (as *L. grisescens*); Bradley, W. G., and Clark, 1940: 526, 528-532, Conn., Mass., Mich., N.J., Ohio, Va. (as *L. stabulans* var. *grisescens*); Bradley, W. G., and Clark, 1939: 528, 530-532, 534, Conn., Mass., Mich., N.J., Ohio, Va. (as *L. stabulans* var. *grisescens*); Bradley, W. G., and Clark, 1939: 528, Ind. (as *Lydella* sp.); Jones, D. W., Walker and Anderson, 1939: 1645-1646, Va. (as *L. stabulans* var. *grisescens* ex *Pyrausta nubilalis*); Baker, W. A., and Bradley, 1938: 345-347, Conn., Mass., Md., Mich., N.J., Ohio (as *L. stabulans* var. *grisescens* ex *Pyrausta nubilalis*); Bradley, W. G., and Clark, 1938: 650-652, 655-656, Conn., Mass., Mich., N.J., Ohio, Va. (as *L. stabulans grisescens*); Baird, 1937: 66, Canada (as *L. stabulans grisescens* ex *Pyrausta nubilalis*); Baird, 1937: 66, Canada (as *Ceromasia lepida* (Meigen) ex *Pyrausta nubilalis*); Clark, C. A., 1934: 18-20, 31, 35, U.S. (as *L. grisescens* ex *Pyrausta nubilalis*); Wishart and Thomas, 1933: 40, E. Canada (as *Masicera senilis* Meigen ex *Pyrausta nubilalis*); Jones, D. W., 1929: 2-3, 24-26, Mass. (as *Masicera senilis* ex *Pyrausta nubilalis*).

STAGE PARASITIZED: Larva, pupa.

IMMATURES: Pupa — Cory, Foster, McConnell, Bissell, Langford and Ditman, 1950: (descr., fig. 14); Baker, W. A., Bradley and Clark, 1949: (fig. 38B); Ritcher, 1947: (fig. 17).

BIOLOGY: Hsiao and Holdaway, 1966: (125-133); Jarvis and York, 1961: (213-214); Clausen, 1956: (7, 42-43, 111-112); Anon., 1955: (17); Andre, 1951: (39-40); Cory, Foster, McConnell, Bissell, Langford and Ditman, 1950: (22-24); Baker, W. A., Bradley and Clark, 1949: (3, 6, 8-9, 12, 21, 23-24, 26-27, 38-40, 53-56, 166-170, 182); Jones, D. W., 1929: (2-3, 24-26).

Lydella sp., see *Blondelia* spp. probably or *Lydella thompsoni*

Lydella sp. probably

HOST: HYMENOPTERA

Argidae

Arge clavicornis (Fabricius) complex

Gobeil, 1937: 87, Ont. (ex *Hylotoma macleayi* Leach).

STAGE PARASITIZED: Pupa.

Lydellohoughia Townsend (Goniinae: Blondeliini)

Lydellohoughia dimmocki, see *Eucelatoriopsis dimmocki*

Lydellohoughia tenella (Reinhard)

HOST: LEPIDOPTERA

Unidentified Lepidoptera

Tuber moth(-s)

Reinhard, 1937: 66, Calif. (as *Dexodes* ex tuber moth on potato tops, probable host).

Lydellohoughia sp.

HOST: LEPIDOPTERA

Pyralidae

Cone moth(-s) probably

Keen, 1958: 70.

Lydinolydella Townsend (Goniinae: Blondeliini)

Gen. and sp. near *Lydinolydella*

HOST: LEPIDOPTERA

Tortricidae

Archips rosanus (Linnaeus)

Schaffner, 1959: 31, 50, Mass. (as *Compsilurini*, near *Lydinolydella* sp. (?)).

BIOLOGY: Schaffner, 1959: (31, 50).

Lypha Robineau-Desvoidy (Tachininae: Lyphini)

Lypha intermedia Brooks

HOST: LEPIDOPTERA

Gelechiidae

Anacamptis innocuella (Zeller)

Raizenne, 1952: 237, Ont.

Olethreutidae

Sciaphila duplex (Walsingham)

Brooks, 1945: 86, Ont.

Lypha setifacies (West)

HOST: LEPIDOPTERA

Tortricidae

Choristoneura conflictana (Walker)

Prentice, 1955: 471, Man., Sask.

Choristoneura fumiferana (Clemens)

Blais, 1965: 131-134, Que.; MacDonald and Webb, 1963: 299, N.B.; Miller, 1963: 237, 239, N.B.; Blais, 1960: 388, 390-392, Que. (ex spruce budworm); Coppel, 1960: 94, 97, B.C.; Carolin and Coulter, 1959: 552-553, Oreg.; McGugan and Blais, 1959: 763, 765, 768, 774-775, Ont.; Coppel, 1953: 191-192, B.C.; Dowden, Jaynes and Carolin, 1953: 309, Maine; Anon., 1952g: 287, 292, 296, B.C.; Jaynes and Drooz, 1952: 1058, 1060, Maine, N.Y.; Ross, 1952: 108, fig. 9, Canada; Dowden and Carolin, 1950: 778-779, N.Y. (ex *Archips*); Wilkes, Coppel and Mathers, 1949: 140, 146, 149, 151, 153, B.C.; Dowden, Buchanan and Carolin, 1948: 458 (ex *Archips*); Coppel, 1947: 40, B.C. (ex *Archips*); Wilkes and Anderson, 1947: 41, Ont. (ex *Archips*); Brooks, 1945: 84, Ont., Que. (ex *Archips*).

Choristoneura pinus Freeman

Allen, D. C., Knight, Foltz and Mattson, 1969: 1471-1472, 1474-1475, Mich.; Dixon and Benjamin, 1963: 267-270, Wis.; Benjamin and Drooz, 1954: 589-590, Mich.

Choristoneura rosaceana (Harris)

Brooks, 1945: 84, Que. (ex *Archips*).

Croesia albicomana (Clemens)

Cochaux, 1968: 104, 106, Que. (ex *C. semipurpurana* (Kearfott)).

STAGE PARASITIZED: Larva, pupa.

IMMATURES: Pupa — Ross, 1952: (fig. 9).

BIOLOGY: Cochaux, 1968: (104, 106); Miller, 1963: (237, 239); Coppel, 1947: (40).

Lypha sp.

HOST: LEPIDOPTERA

Tortricidae

Sparganothis pettitana (Robinson)

Raizenne, 1952: 203, Que.

Macquartia Robineau-Desvoidy (Dexiinae: Macquartiini)

Macquartia pristis, see *Anaporia pristis*

"*Macquartia*" spp.

HOST: COLEOPTERA

Unidentified Coleoptera

Miscellaneous

*Brauer and Bergenstamm, 1895: 560, 588, N. Amer. (Townsend, 1893); Townsend, 1893: 467, Ill. (as *Macquartia* sp. ?).

HOST: LEPIDOPTERA

Arctiidae

Halisidota argentata Packard

Rawlings, 1960: 56, Calif. (ex "*Euchausia*").

STAGE PARASITIZED: Larva (Coleoptera).

Gen. and sp. of Marquartiidae, see Tachinidae

Macromeigenia chrysoprocta, see *Mystacella chrysoprocta*

"*Macrophthalma*" *disjuncta*, see *Microphthalma disjuncta*

"*Madremeyia*" *saundersii*, see *Madremyia saundersii*

Madremyia saundersii, see *Madremyia saundersii*

Madremyia Townsend (Goniinae: Eryciini)

Madremyia saundersii (Williston)

HOST: COLEOPTERA

Curculionidae

Hypera postica (Gyllenhal)

Michelbacher, 1943: 20, Calif.

HOST: LEPIDOPTERA

Danaiidae

Danaus plexippus plexippus (Linnaeus)

Weaver, 1969: 71, W. Va. (ex *D. plexippus*).

Geometridae

Cingilia catenaria (Cramer)

Franklin, 1950: 57; Schaffner and Griswold, 1934: 98, 115.

Coryphista meadii (Packard)

Neiswander, 1941: 388, Ohio.

Deuteronomos magnarius (Guenée)

All records ex *Ennomos magnarius*.

*Aldrich and Webber, 1924: 88 (Tothill, 1913); Tothill, 1913: 73, N.B. (as *Phorocera*).

Ennomos subsignarius (Hübner)

Schaffner and Griswold, 1934: 99-100, 115, Mass.

Lambdina fiscellaria (Guenée)

Schaffner and Griswold, 1934: 99, 115 (ex *Ellopia*).

Lambdina fiscellaria fiscellaria (Guenée)

Carroll, 1956: 594, Nfld.; Reeks, Forbes and Cuming, 1947: 13, Nfld. (ex *L. fiscellaria*).

Lambdina fiscellaria somnaria (Hulst)

Chamberlin, 1931: 1038, Oreg. (ex *Ellopia somnaria*).

Lasiocampidae

Malacosoma disstria Hübner

Schaffner and Griswold, 1934: 89, 115.

Lymantriidae

Euproctis chrysorrhoea (Linnaeus)

Aldrich and Webber, 1924: 88 (ex *E. "chrysorrhoea"*).

Nygmia phaeorrhoea (Donovan)

Burgess and Crossman, 1929: 143 (as *Phorocera*).

Noctuidae

Agrotis ipsilon (Hufnagel)

All records ex *Agrotis ypsilon*.

Franklin, 1950: 57; *Crumb, 1929: 59 (Aldrich and Webber, 1924); Aldrich and Webber, 1924: 88, Calif.

Autographa californica (Speyer)

Clancy, 1969: 1079-1080, Calif.; Seidel, 1954: 186, Wash. (ex *A. "california"*); Essig, 1926: 582, Wash; *Aldrich and Webber, 1924: 88 (Hyslop, 1912); Hyslop, 1912: 117, probably Wash. (as *Phorocera* ex *A. gamma californica*).

Epiglaea apiata (Grote)

Franklin, 1950: 57.

Peridroma saucia (Hübner)

*Crumb, 1929: 40, 115 (ex *Lycophotia*, Aldrich and Webber, 1924); Essig, 1926: 582, Calif., Oreg. (ex *Lycophotia margaritosa* (Haworth)); Aldrich and Webber, 1924: 88 (ex *Lycophotia margaritosa*); Essig, 1915: 329, 401, Calif. (as *Phorocera* ex *Lycophotia margaritosa*); Lovett, 1915: 145, Oreg. (as *Phorocera* ex *P. margaritosa saucia*); Chittenden, 1901a: 61, Oreg. (as *Phorocera*).

Pseudaletia unipuncta (Haworth)

Pond, 1960: 665, N.B.

Trichoplusia ni (Hübner)

Oatman, 1966: 1138, Calif.

Xylena nupera (Lintner)

Franklin, 1950: 57.

Nymphalidae

Nymphalis antiopa (Linnaeus)

All records ex *Euvanessa antiopa*.

*Aldrich and Webber, 1924: 88 (Tothill, 1913); Tothill, 1913: 73-74, Ont. (as *Phorocera*).

Speyeria cybele (Fabricius)

All records ex *Argynnis cybele*.

*Aldrich and Webber, 1924: 88 (Williston, 1889); *Brauer and Bergenstamm, 1895: 569, 583 (as (*Phorocera*), Williston, 1889); Williston, 1889: 1914, 1922 (as *Phorocera*).

Vanessa cardui (Linnaeus)

Lange, 1941: 67, Calif.

Vanessa carye Hübner

Essig, 1926: 582, Calif.; Coolidge, 1925: 147, Calif. (as *Phorocera*).

Olethreutidae

Sciaphila duplex (Walsingham)

McGregor, 1967: 1215.

Pieridae

Pieris occidentalis Reakirt

Aldrich and Webber, 1924: 88, Idaho.

Pieris rapae (Linnaeus)

Oatman, 1966: 1136-1138, Calif.; Harcourt, 1963: 65, Ont.;
Coppel and Maw, 1954a: 321 (lab.).

Pyralidae

Ostrinia nubilalis (Hübner)

Coppel and Maw, 1954a: 321-322 (ex *Pyrausta*, lab.).

Tortricidae

Acleris variana (Fernald)

Miller, C. A., 1966: 603, N.B.

Choristoneura conflictana (Walker)

Prentice, 1955: 471, Man., Sask.

Choristoneura fumiferana (Clemens)

Blais, 1965: 131, Que. (as *Madremya*); Miller, 1963: 237, 239, N.B.; Blais, 1960: 388, 390, Que. (as *Madremya* ex spruce budworm); Coppel, 1960: 94, 97, B.C.; Carolin and Coulter, 1959: 552-554, Oreg.; Coppel and Maw, 1954a: 314-322, figs. 1-16, B.C.; Boyes and Wilkes, 1953: 126, 147; Coppel, 1953: 191-192, B.C.; Anon., 1952b: 178, 184-185, B.C.; Anon., 1952g: 287, 292, 295, B.C.; Ross, 1952: 108, fig. 6, Canada; Anon., 1950c: 236, 245, 248, B.C.; Dowden and Carolin, 1950: 778-779, N.Y. (ex *Archips*); Anon., 1949: 270, 282, 285, B.C.; Wilkes, Coppel and Mathers, 1949: 146, 149, 151, 153, B.C.; Anon., 1948: 284, 293, 295, 297, B.C.; Anon., 1948a: 354, 366, 368-371, B.C.; Dowden, Buchanan and Carolin, 1948: 458-463 (as "*Madremeyia*" ex *Archips*); Bedard, 1938: 191 (ex *Cacoecia*).

Choristoneura parallela (Robinson)

Franklin, 1950: 57 (ex *Archips*); Johnson, 1925: 199, Mass. (ex *Cacoecia*).

Choristoneura pinus Freeman

Allen, D. C., Knight, Foltz and Mattson, 1969: 1471-1472, 1474-1475, Mich.; Dixon and Benjamin, 1963: 267, Wis.; Kulman and Hodson, 1961: 221-222, Minn.

Unidentified Lepidoptera

Looper(-s)

Knowlton, 1936: 242, Utah.

Miscellaneous

Essig, 1926: 582.

STAGE PARASITIZED: Larva (Coleoptera, Lepidoptera), pupa (Lepidoptera), adult (Lepidoptera [Weaver, 1969: 71]).

IMMATURES: Egg — Coppel and Maw, 1954a: (descr., fig. 1)

First instar larva — Coppel and Maw, 1954a: (descr., figs. 2-5, 7).

Second instar larva — Coppel and Maw, 1954a: (descr., figs. 6, 8-9).

Third instar larva — Coppel and Maw, 1954a: (descr., figs. 10-13).

Pupa — Coppel and Maw, 1954a: (descr., figs. 14-16); Ross, 1952: (fig. 6); Greene, 1922: (descr., fig. 24 on pl. 6).

BIOLOGY: Clancy, 1969: (1079-1080); Miller, 1963: (237, 239); Coppel and Maw, 1954a: (314-322); Seidel, 1954: (186); Schaffner and Griswold, 1934: (115).

“Masicera” sp., see “*Masicera*” spp.

Masicera Macquart (Goniinae: Eryciini)

Masicera albifacies, see *Oswaldia albifacies*

Masicera “albifacies,” see *Oswaldia albifacies*

Masicera arator, see *Pilatea arator*

Masicera archippivora, see *Lespesia archippivora*

Masicera datanarum, see *Lespesia datanarum*

Masicera dubia, see *Lespesia dubia*

Masicera eufitchiae, see *Blondelia eufitchiae*

Masicera “eufitchii,” see *Blondelia eufitchiae*

Masicera exilis, see *Eribella exilis*

Masicera sp. near *exilis*, see *Eribella* sp. near *exilis*

Masicera frenchii, see *Lespesia frenchii*

Masicera myoidea, see *Lydella radialis*

Masicera “myoidea,” see *Lydella radialis*

Masicera “myoidea” probably, see *Lydella radialis* probably

Masicera nigrita, see *Lydella radialis*

Masicera protoparcis, see *Drino rhoeo*

Masicera rileyi, see *Lespesia rileyi*

Masicera rutila, see *Ceromasia aurifrons*

Masicera “seniles,” see *Lydella radialis*

Masicera senilis, see *Lydella radialis* or *Lydella thompsoni*

Masicera schizurae, see *Lespesia schizurae*

Masicera sphingivora, see *Lespesia frenchii*

“*Masicera tantilla* Wulp”

HOST: LEPIDOPTERA

Pyralidae

Acigona dentilineatella (Barnes and McDunnough)

Van Zwaluwenburg, 1951: 375, Mexico (V.C.) (ex *Xubida*).

Masicera tenthredinidarum, see *Bessa harveyi*

Masicera unispinosa, see *Sturmia harrisinae*

Masicera sp., see “*Masicera*” spp. or *Sitophaga* sp.

“*Masicera*” spp.

HOST: LEPIDOPTERA

Citheroniidae

Anisota rubicunda (Fabricius)

Schaffner and Griswold, 1934: 30-31.

Noctuidae

Simyra henrici (Grote)

Webber and Schaffner, 1926: 21, New England (ex
Arsilonche albovenosa (Goeze)).

Pieridae

Colias eurytheme (Boisduval)

All records ex *Eurymus eurytheme*.

Wildermuth, 1914: 24, Calif.; Wildermuth, 1911: 7.

Sphingidae

Manduca sexta (Linnaeus)

Ashmead, 1887: 17, “the North” (as “*Mascicera*” ex *Sphinx carolina* Linnaeus).

STAGE PARASITIZED: Pupa.

“*Masicora*” *archippivora*, see *Lespesia archippivora*

Medina Robineau-Desvoidy (Goniinae: Blondeliini)

Medina barbata (Coquillett)

HOST: COLEOPTERA

Chrysomelidae

Altica chalybea Illiger

Loan, 1963: 538, Ont. (as *Amedoria spinosa* (Coquillett)).

Altica corni Woods

Loan, 1963: 538, Ont. (as *Amedoria spinosa*); Woods, W. C., 1918: 173, Maine (as *Celatoria spinosa*).

Altica ulmi Woods

Loan, 1963: 538, Ont. (as *Amedoria spinosa*); Woods, W. C., 1918: 173, 193, Maine (as *Celatoria spinosa*).

Altica spp.

Woods, W. C., 1917: 274, Maine (as *Celatoria spinosa* ex *Altica* sp. 1); Woods, W. C., 1917: 274, Maine (as *Celatoria spinosa* ex *Altica* sp. 2).

Disonycha xanthomelas (Dalman)

Most records ex *Disonycha* "*xanthomelaena*."

Johnson, 1925: 188, Conn., Maine, Mass., N.H. (as *Methypostena*); Knab, 1909: 34, D.C. (as *Hypostena* ex "*Disomycha*"); Chittenden, 1899: 85 (as *Hypostena* ex *D.* "*xanthomeioena*"); Chittenden, 1898a: 78-79 (as *Hypostena*); Coquillett, 1897: 17, 23, D.C. (as *Hypostena*).

"*Luperodes*" spp.

Eddy, C. O., 1941: 420, La. (as *Amedoria luctuosa* (Meigen)); Eddy, C. O., 1941a: 509, La. (as *Amedoria luctuosa*).

Plagiometriona clavata (Fabricius)

Johnson, 1925: 188, Conn., Maine, Mass., N.H. (as *Methypostena* ex *Deloyala*).

STAGE PARASITIZED: Adult.

BIOLOGY: Loan, 1963: (538); Woods, W. C., 1918: (173).

Medina spp.

HOST: COLEOPTERA

Chrysomelidae

Unidentified Chrysomelidae

Townsend, 1913a: 304 (as *Methypostena*).

STAGE PARASITIZED: Larva, adult.

BIOLOGY: Townsend, 1913a: (304).

Megaparia opaca, see *Megapariopsis opaca**Megapariopsis* Townsend (Proseninae: Dexillini)*Megapariopsis opaca* (Coquillett)

HOST: COLEOPTERA

Scarabaeidae

Euetheola rugiceps (LeConte)

Phillips and Fox, 1924: 30, Va.

STAGE PARASITIZED: Larva.

IMMATURES: Pupa — Greene, 1922: (descr., fig. 9 on pl. 2).

Meigenia Robineau-Desvoidy (Goniinae: Blondeliini)*Meigenia hyphantriae*, see *Hyphantrophaga hyphantriae**Meigenia mutabilis* (Fallén)

HOST: COLEOPTERA

Chrysomelidae

Crioceris asparagi (Linnaeus)

Clausen, 1956: 53, N.J.

Meigenia websteri, see *Lespesia archippivora**Melanophora Meigen* (Rhinophorinae: Rhinophorini)*Melanophora diabroticae*, see *Celatoria diabroticae**Melanophora roralis* (Linnaeus)

HOST: ISOPODA (CRUSTACEA)

Oniscidae

Oniscus asellus Linnaeus

Jones, F. M., 1948: 32-33, Mass.

Porcellio scaber Latreille

Jones, F. M., 1948: 32-33, Mass.

Porcellio scaber Latreille probably

Brues, 1903: 291, Mass.

Porcellio spp.

Johnson, 1925: 210, Conn., Mass., N.H.

STAGE PARASITIZED: Adult.

IMMATURES: Pupa — Brues, 1903: (descr.).

"Memorilla" *maculosa*, see *Nemorilla pyste*"Mericea" *ampelus*, see *Mericia ampelus**Mericia Robineau-Desvoidy* (Tachininae: Ernestiini)*Mericia ampelus* (Walker)

HOST: LEPIDOPTERA

Arctiidae

Diacrisia virginica (Fabricius)Aldrich, 1932: 2, Mass. (as *Ernestia*, incomplete rearing — pupa).

Hyphantria cunea (Drury)

Williamson, 1969: 17, Ont.; Williamson, 1968: 17, Ont.; Williamson, 1963: 150, Ont.; Williamson, 1962: 158, Ont.; Kelleher, 1960a: 280, Canada; Maybee, 1959: 264, Ont. (as "*Mericea*"); Maybee, 1958: 312, B.C., N.B., N.S., Ont.; Boyes and Wilkes, 1953: 126; Fattig, 1949: 21; *Strickland, 1938: 216 (ex fall webworm, Tothill, 1921); Schaffner and Griswold, 1934: 37, 109 (as *Ernestia*); Allen, 1929: 685, Miss. (as *Ernestia*); Johnson, 1925: 195, Conn., Maine, Mass., N.H., Vt. (as *Ernestia* ex fall webworm); Tothill, 1922: 10-14, 17-20, 22-24, 31-32, 34, 44-45, 49-59, 103-105, 107, figs. 18-28, B.C., N.B. (as *Ernestia*); Tothill, 1921: 230, 274, Canada (as *Ernestia*); Johnson, 1910: 779, N.J. (as *Panzeria radicum* (Fabricius)); Coquillett, 1897: 19, 25, D.C., Mo. (as *Panzeria radicum*).

Hyphantria textor (Harris)

Raizenne, 1952: 29, Ont.; Daviault, 1942: 398, Que.; Sweetman, 1936: 231-232, N. Amer. (as *Ernestia*); *Brauer and Bergenstamm, 1895: 564, 594 (as (*Nemoraea*) *nigricornis* Townsend, Townsend, 1893); Townsend, 1893: 467, Ill. (as *Nemoraea nigricornis*).

Hyphantria sp.

Coquillett, 1897: 19, 25, Md. (as *Panzeria radicum*).

Lasiocampidae

Malacosoma americanum (Fabricius)

Bess, 1936: 605 (as *Ernestia*); Schaffner and Griswold, 1934: 88, 109 (as *Ernestia*); Hine, 1904: 87, B.C. (as *Panzeria* ex *Clisiocampa* (?)).

Malacosoma disstria Hübner

Berry, P. A., 1939: 718 (as *Ernestia*, lab.); Bess, 1936: 605 (as *Ernestia*); Schaffner and Griswold, 1934: 89, 109 (as *Ernestia*).

Notodontidae

Datana integerrima Grote and Robinson

Raizenne, 1952: 81, Ont.; Schaffner and Griswold, 1934: 76, 109 (as *Ernestia*).

Datana ministra (Drury)

Tothill, 1922: 10-12, 17-20, 22-24, 31-32, 44-45, 49-59, 103-105, 107, figs. 18-28, B.C., Mass., N.B., N.S., Ont., Que. (as *Ernestia*).

Unidentified Lepidoptera

Miscellaneous

Aldrich, 1915: 82 (as *Panzeria radicum*).

STAGE PARASITIZED: Larva, pupa.

IMMATURES: First instar larva — Tothill, 1922: (descr., figs. 18-19, 26-27).

Second instar larva — Tothill, 1922: (descr., figs. 20-21, 24-25, 28).

Third instar larva — Tothill, 1922: (descr., figs. 22-23).

Pupa — Greene, 1922: (descr., fig. 86 on pl. 18 and fig. 92 on pl. 19); Tothill, 1922: (descr.).

BIOLOGY: Schaffner and Griswold, 1934: (109); Tothill, 1922: (10-14, 17-20, 22-24, 31-32, 34, 44-45, 49-59, 103-105, 107); Aldrich, 1915: (82).

Mericia arcuata (Tothill)

HOST: LEPIDOPTERA

Arctiidae

Hyphantria cunea (Drury)

Strickland, 1938: 216 (ex fall webworm).

Mericia arcuata (Tothill) probably

HOST: LEPIDOPTERA

Noctuidae

Eupsilia devia (Grote)

Schaffner and Griswold, 1934: 51, 109, Maine (as *Ernestia* sp. (1) (?*arcuata*) ex *Conistra*).

BIOLOGY: Schaffner and Griswold, 1934: (51, 109).

Mericia bicarina (Tothill) probably

HOST: LEPIDOPTERA

Noctuidae

Barathra curialis (Smith)

Schaffner and Griswold, 1934: 44, 111, Maine (as *Ernestia* sp. (10) (? *bicarina*)).

STAGE PARASITIZED: Pupa.

BIOLOGY: Schaffner and Griswold, 1934: (44, 111).

Mericia johnsoni (Tothill)

HOST: LEPIDOPTERA

Arctiidae

Hyphantria cunea (Drury)

Tothill, 1921: 230 (as *Ernestia*).

Mericia spp.

HOST: LEPIDOPTERA

Arctiidae

Diacrisia virginica (Fabricius)

Schaffner and Griswold, 1934: 36, 110 (as *Ernestia* sp. (7)).

Estigmene acrea (Drury)

Schaffner and Griswold, 1934: 38, 110 (as *Ernestia* sp. (7)).

Euchaetias egle (Drury)

Schaffner and Griswold, 1934: 39, 110, N.Y. (as *Ernestia* sp. (5)).

Hyphantria cunea (Drury)

Howard and Fiske, 1911: 203 (as *Varichaeta aldrichi* (Townsend) ex fall webworm).

Geometridae

Ennomos subsignarius Hübner

Schaffner and Griswold, 1934: 99-100, 109, Mass. (as *Ernestia* sp. (2)).

Noctuidae

Agrotis orthogonia Morrison

Strickland, 1921: 98, Alta. (as *Ernestia radicum* (Fabricius) ex *Porosagrotis*).

Amphipyra pyramidoides Guenée

Schaffner and Griswold, 1934: 52, 110 (as *Ernestia* sp. (10)).

Cosmia canescens (Behr)

Schaffner and Griswold, 1934: 62, 110, N.J. (as *Ernestia* sp. (3)).

Euxoa auxiliaris (Grote)

Crumb, 1929: 40 (as *Ernestia* ex *Chorizagrotis*); Snow, 1925: 605, 608, Utah (as *Ernestia*).

Lithophane antennata (Walker)

Schaffner and Griswold, 1934: 49, 110, N.J. (as *Ernestia* sp. (4) ex *Graptolitha*).

Lithophane spp. (including *L. disposita* Morrison, *L. innominata* (Smith) and/or *L. petulca* Grote)

Schaffner and Griswold, 1934: 48, 110 (as *Ernestia* sp. (9) ex *Graptolitha* (including *G. disposita*, *G. innominata* and/or *G. petulca*)).

Unidentified Noctuidae

Schaffner and Griswold, 1934: 72, 110, Maine (as *Ernestia* sp. (8)).

Notodontidae

Datana ministra (Drury)

Schaffner and Griswold, 1934: 74, 110, Maine, N.Y. (as *Ernestia* sp. (6)).

Olethreutidae

Spilonota ocellana (Denis and Schiffermüller)

Stultz, 1955: 80, N.S.

STAGE PARASITIZED: Larva, pupa.

BIOLOGY: Schaffner and Griswold, 1934: (39, 49, 62, 72, 74, 99-100, 109-110); Howard and Fiske, 1911: (203).

Mericia spp. probably

All records as *Varichaeta* [sp.].

HOST: LEPIDOPTERA

Arctiidae

Hyphantria cunea (Drury)

Tothill, 1920: 32, N.B. (ex fall webworm).

Hyphantria sp.

Tothill, 1919: 13, N.B.

STAGE PARASITIZED: Larva.

Metachaeta helymus, see *Periscepsia helymus*

Metadexia Coquillett (Dexiinae: Zeliini)

Metadexia basalis, see *Cholomyia inaequipis*

Metadexia tricolor Coquillett

HOST: COLEOPTERA

Curculionidae

Rhabdoscelus obscurus (Boisduval)

Wray, 1950: 31, N.C. (ex "*Rhodobaemus*" [sp.] in dahlia stem).

STAGE PARASITIZED: Larva.

Metagonistylum Townsend (Goniinae: Eryciini)

Metagonistylum minense Townsend

HOST: LEPIDOPTERA

Pyralidae

Diatraea canella Hampson

Bartlett, 1941: 2, St. Lucia; Box, 1939: 230, 233, 235-236, 238-239, St. Lucia.

Diatraea impersonatella (Walker)

Box, 1960: 910, Guadeloupe; Fretay, 1960: 145, Guadeloupe (as *Metagonistylum* [sp.]); Fretay, 1960a: 151, Guadeloupe (as *Metagonistylum* [sp.]).

Diatraea saccharalis (Fabricius)

Anon., 1960: 211, Barbados; Box, 1960: 904, 909, St. Lucia; Charpentier, McCormick and Mathes, 1960: 865-867, La.; Fretay, 1960: 145, Guadeloupe (ex "*Diatrea*"); Fretay, 1960a: 151, Guadeloupe (as *Metagonistylum* [sp.]); Charpentier, 1956: 267-268, La.; Dugas, Miller and Conciencie, 1956: 93, La. (ex sugarcane borer); Dugas, Conciencie, Sibley and Joyner, 1955: 88, La. (ex sugarcane borer); Box, 1954: 551, Guadeloupe, Martinique, St. Lucia; Mathes, Ingram, Questel, Thames and Dugas, 1954: 562, Fla., La.; Simmonds, 1953: 122-127, Guadeloupe, Martinique; Ingram, Mathes, Wilson and Hayslip, 1951: 356, La.; Wolcott, 1951a: 476, P.R. (ex sugarcane moth borer); Annand, 1944: 134, Fla. (ex sugarcane borer); Bartlett, 1943: 16-17, P.R.; Bartlett, 1942: 19, P.R.; Lee, 1942: 65-67, P.R.; Scaramuzza, 1942: 123, Fla.; Scaramuzza and Ingram, 1942: 644, Fla.; Wilson, J. W., 1942: 164, Fla.; Bartlett, 1941: 1-18, figs. 2-6a, P.R.; Ingram and Bynum, 1941: 9, La.; Wilson, J. W., 1941: 54-55, La.; Holloway and Mathes, 1940: 738-742, La.; Lee, 1940: 102-105 (lab.); Wolcott, 1940: 31-32 (lab.); Bartlett, 1939: 243-245, P.R.; Box, 1939: 228-239, St. Lucia; Holloway, 1939: 260, La., P.R.; Lee, 1939: 95, P.R.; Bartlett, 1938: 86, P.R.; Bartlett, 1938a: 8 (lab.); Bartlett, 1938b: 1-4 (lab.); Bartlett, 1937: 3-4 (lab.); Scaramuzza, 1934: 14, Antiqua, Cuba, St. Kitts.

Diatraea spp.

Simmonds, 1948: 68, U.S., Guadeloupe, St. Lucia.

STAGE PARASITIZED: Larva, pupa.

IMMATURES: Egg — Bartlett, 1941; (descr., fig. 2a).

First instar larva — Bartlett, 1941: (descr., figs. 2b, 3a).

Second instar larva — Bartlett, 1941: (descr., fig. 3b).

Third instar larva — Bartlett, 1941: (descr., figs. 3c, 4).

Pupa — Bartlett, 1941: (descr., figs. 5, 6a).

BIOLOGY: Simmonds, 1953: (122-127); Bartlett, 1941: (1-18);

Bartlett, 1939: (243-245); Bartlett, 1938a: (8).

Metagonistylum sp., see *Metagonistylum minense*

Metaplagia Coquillett (Tachininae: Voriini)

Metaplagia occidentalis Coquillett

HOST: LEPIDOPTERA

Noctuidae

Heliothis virescens (Fabricius)

Fattig, 1949: 21-22.

Heliothis zea (Boddie)

Fattig, 1949: 21-22 (ex *H. armigera* (Hübner)); Brimley, 1938: 360, N.C. (ex *H. armiger*); Brimley, 1922: 24, N.C. (ex *H. obscura* (Fabricius)).

Sphingidae

Agrius convolvuli (Linnaeus)

Reinhard, 1922: 72, Tex. (ex *Herse*).

Sphinx luscitiosa (Clemens)

Schaffner and Griswold, 1934: 19, 108, Mass. (ex *Hyloicus*)

BIOLOGY: Schaffner and Griswold, 1934: (19, 108).

Metavoria Townsend (Tachininae: Voriini)

Metavoria orientalis Townsend

HOST: LEPIDOPTERA

Sphingidae

Manduca sexta (Linnaeus)

All records ex *Protoparce sexta*.

Allen, 1929: 684, Miss.; Allen, 1924: 24, Miss.

Methypostena barbata, see *Medina barbata*

Methypostena sp., see *Medina* spp.

Metopiops Townsend (Goniinae: Eumasicerini)

Metopiops pyralidis (Coquillett)

All records as *Pseudochaeta pyralidis*.

HOST: LEPIDOPTERA

Nymphalidae

Vanessa cardui (Linnaeus)

Rowe, 1933: 125, Ill. (ex thistle butterfly).

Pyralidae

Unidentified Pyralidae

Coquillett, 1897: 20, 27.

STAGE PARASITIZED: Larva.

Metopiops sp. near *pyralidis* (Coquillett)

HOST: LEPIDOPTERA

Pyralidae

Tetralopha asperatella (Clemens)

Schaffner, 1959: 13, 51, Mass.

BIOLOGY: Schaffner, 1959: (13, 51).

Miamimyia Townsend (Goniinae: Hyperecteinini)

Miamimyia cincta Townsend

HOST: ORTHOPTERA

Blattidae

Periplaneta americana (Linnaeus)

Patton, 1958: 35, Fla. (as *Chaetophleps* n. sp.).

BIOLOGY: Patton, 1958: (35).

Microceromasia sphenophori, see *Lixophaga sphenophori*

Microceromasia "spherophori," see *Lixophaga sphenophori*

Micropalpus fulgens, see *Bonnetia comta*

Micropalpus sp., see *Bonnetia* sp.

"*Microphthalma*" *disjuncta*, see *Microphthalma disjuncta*

Microphthalma Macquart (Proseninae: Dexillini)

Microphthalma disjuncta (Wiedemann)

HOST: COLEOPTERA

Scarabaeidae

Euetheola rugiceps (LeConte)

Douglas, W. A., and Ingram, 1942: 9, S. U.S.; Ingram, Bynum
and Douglas, 1939: 480, [S. U.S.] (as "*Microphthalma*");

Ingram and Bynum, 1932: 847, La.

Phyllophaga fervida (Fabricius)

All records ex *Lachnosterna arcuata* Smith.

Walton, 1913: 198; Johnson, 1910: 783, N.J.; Chittenden, 1899: 79, D.C.; Coquillett, 1897: 18, 23, D.C.

Phyllophaga spp.

Fattig, 1949: 22, Ga.; Hammond, 1948: 175, Ont.; Fattig, 1944: 4, Ga.; Fattig, 1944: 4, Ga. (as *M. "divisa"* Wiedemann); Johnson, 1925: 208, Conn., Maine, Mass., N.H., Vt.; Hayes, 1920: 317, Kans. (ex *Lachnosterna*); Hudson, 1920: 83, Ont. (ex *Lachnosterna*); Criddle, 1918: 454, Man. (ex *Lachnosterna*); Davis, J. J., 1918: 13-14 (ex *Lachnosterna*); Smyth, 1917: 59, Cent. W. U.S.; Davis, J. J., 1916: 271, Cent. W. U.S. (ex *Lachnosterna*); Aldrich, 1915: 82 (ex *Lachnosterna*); Forbes, S. A., 1908: 160-161, Ill. (as "*Macrophthalma*" ex *Lachnosterna*).

June beetle(-s)

Felt, 1908: 43, [Ill.] (as "*Macrophthalma*").

Scarabaeid spp. (including *Cyclocephala immaculata* (Olivier) or *Phyllophaga* spp.)

Forbes, S. A., 1891: 44-45, 52, Ill. (as *M. nigra* Macquart ex white grubs (*Lachnosterna* spp. or *C. immaculata*)).

Scarabaeid spp. (including *Cyclocephala* spp. and/or *Phyllophaga* spp.)

Forbes, S. A., 1907: 472-473, Ill. (as "*Macrophthalma*" ex white grubs (*Lachnosterna* spp. or *Cyclocephala* spp.)).

Scarabaeid spp. (allied genera of *Phyllophaga* spp.)

Criddle, 1918: 454, Man. (ex allied genera of *Lachnosterna*).

White grub(-s)

Hammond, 1949: 161, Ont.

STAGE PARASITIZED: Larva.

IMMATURES: Pupa — Greene, 1922: (descr., fig. 17 on pl. 4).

Microphthalma "divisa," see *Microphthalma disjuncta*

Microphthalma michiganensis (Townsend)

HOST: COLEOPTERA

Scarabaeidae

Phyllophaga anxia (LeConte)

Jarvis, 1966: 406-407, 409, Nebr.; Hammond, 1931: 71-72, Que.; Hammond, 1930: 34, Que.; Hammond, 1930a: 60, Que. (ex "*Phyllophaga*"); Aldrich, 1926: 5, Que.; Petch and Hammond, 1926: 85, 89-90, Que.; Petch and Hammond, 1925: 25-27, Que. (as *M. phyllophagae* Curran).

Phyllophaga fervida (Fabricius)

Aldrich, 1926: 5, D.C. (ex *P. arcuata* (Smith)).

Phyllophaga spp.

Hammond, 1942a: 148, Ont., Que.; Aldrich, 1926: 5, Md.

White grub(-s)

Hammond, 1937: 163, Ont.; Anon., 1932: 1251, Que.; Weir, 1932: 178, 185, [Que.]; Hammond, 1929: 39, Que.; Motherwell, 1928: 117 (as "*Microthalma*"); Anon., 1927: 440, Canada; Gibson, 1927: 57, Que.; Petch, 1925: 6, Que. (as *M. phyllophagae*).

STAGE PARASITIZED: Larva.

IMMATURES: Egg — Petch and Hammond, 1925: (descr.).

First instar larva — Petch and Hammond, 1925: (descr.).

Third instar larva — Petch and Hammond, 1926: (descr.).

BIOLOGY: Petch and Hammond, 1926: (85, 89-90).

Microphthalma nigra, see *Microphthalma disjuncta*

Microphthalma obsoleta (Wulp)

All records as *Microphthalma pruinosa* Coquillett.

HOST: COLEOPTERA

Scarabaeidae

Phyllophaga spp.

Johnson, 1925: 208, Mass., Vt.; Davis, J. J., 1916: 271, New England (ex *Lachnosterna*).

STAGE PARASITIZED: Larva.

Microphthalma phyllophagae, see *Microphthalma michiganensis*

Microphthalma pruinosa, see *Microphthalma obsoleta*

Microphthalma spp.

HOST: COLEOPTERA

Scarabaeidae

White grub(-s)

Curran, 1925: 16, Que.

STAGE PARASITIZED: Larva.

"*Microthalma*" *michiganensis*, see *Microphthalma michiganensis*

"*Miophasia*" *metallica*, see *Myiophasia metallica*

Mochlosoma Brauer and Bergenstamm (Proseninae: Prosenini)

Mochlosoma lacertosum (Wulp)

HOST: COLEOPTERA

Scarabaeidae

Scarabaeid spp. (including *Phyllophaga* spp. or other Scarabaeidae)

Townsend, 1912a: 39-40, Mexico (Chih.) (ex *Lachnosterna* or other "scarbaeid" beetles, suggested host).

STAGE PARASITIZED: Larva.

BIOLOGY: Townsend, 1912a: (39-40).

Mochlosoma mexicanum (Macquart)

HOST: LEPIDOPTERA

Unidentified Lepidoptera

Miscellaneous

Gibson, W. W., and Carrillo S., 1959: 183, Mexico (Mex.) (ex gusanos de maíz).

Muscadae, see Tachinidae

Myiocera "cremiodes," see *Dinera grisescens* probably

Myiopharus Brauer and Bergenstamm (Goniinae: Blondeliini)

Myiopharus canadensis Reinhard

HOST: COLEOPTERA

Chrysomelidae

Calligrapha multipunctata bigsbyana (Kirby)

Reinhard, 1945: 32, Que. (ex *C. bigsbyana*).

STAGE PARASITIZED: Adult.

Myiopharus dorsalis (Coquillett)

HOST: COLEOPTERA

Chrysomelidae

Calligrapha multipunctata bigsbyana (Kirby)

Daviault, 1941a: 106-107 (as *Doryphorophaga*).

BIOLOGY: Daviault, 1941a: (106-107).

Myiophasia Brauer and Bergenstamm (Proseninae: Myiophasiini)

Myiophasia "aenea Wiedemann"
HOST: COLEOPTERA

Curculionidae

Ampelogypter sesostris (LeConte)

Pierce, Cushman and Hood, 1912: 48, 80 (ex *A. "sesostris"*);
Quaintance and Jenne, 1912: 149, Ohio; Brooks, F.E.,
1911: 68, Ohio; Brooks, F. E., 1909: 329; Pierce, 1908:
381, 388; Pierce, 1908a: 42; Webster, 1899: 54, pl. 3,
Ohio.

Anthonomus grandis Boheman

Hunter and Pierce, 1912: 142, Tex.; Pierce, Cushman and
Hood, 1912: 48; Quaintance and Jenne, 1912: 149,
Tex.; Pierce, 1910: 453 (ex boll weevil); Pierce, 1908:
381, 389; Pierce, 1908a: 28, 30, 42, La., Tex.

Chalcodermus aeneus Boheman

Thomas, W. A., 1931: 519, N.C.; Walton, 1913b: 25, S.C. (as
M. "aenia"); Quaintance and Jenne, 1912: 149, S.C.;
Ainslie, G. G., 1910: 140-141, S.C.; Pierce, 1908a: 42,
47.

Chalcodermus spp.

Quaintance and Jenne, 1912: 149, La.; Coquillett, 1897: 18,
23, La.

Conotrachelus affinis Boheman

Mutchler and Weiss, 1925: 11; Pierce, Cushman and Hood,
1912: 48, 77; Pierce, 1908: 381, 390; Pierce, 1908a: 42,
46.

Conotrachelus elegans (Say)

Pierce, Cushman and Hood, 1912: 48, 78; Quaintance and
Jenne, 1912: 149, D.C., Tex.; Pierce, 1908: 381, 390.

Conotrachelus juglandis LeConte

Pierce, Cushman and Hood, 1912: 48, 78; Johnson, 1910:
775, N.J.; Pierce, 1908: 381, 390; Pierce, 1908a: 42;
Coquillett, 1897: 18, 23, Mo., Wash.

Conotrachelus nenuphar (Herbst)

Quaintance and Jenne, 1912: 149-150, Ga., Md., Mich., Va.
(ex *Curculio*).

Conotrachelus spp. (including *C. affinis* Boheman or *C.*
posticatus Boheman)

Pierce, 1907: 362, La.

Conotrachelus spp.

Pierce, 1910: 453.

Curculio nasicus (Say)All records ex *Balaninus nasicus*.

Pierce, Cushman and Hood, 1912: 48, 80; Johnson, 1910: 775, N.J.; Pierce, 1908: 381, 390; Pierce, 1908a: 42; Coquillett, 1897: 18, 23.

Curculio uniformis (LeConte)Quaintance and Jenne, 1912: 149, Mo. (ex *Balaninus*).*Sphenophorus parvulus* GyllenhalQuaintance and Jenne, 1912: 149; Johnson, 1910: 775, N.J.; Pierce, 1908: 381, 392; Pierce, 1908a: 42; *vide Brauer and Bergenstamm, 1895: 563, 606, N. Amer. (Townsend, 1893); Townsend, 1893: 467, Ill. (as *Phasioclista metallica* Townsend).*Trichobaris compacta* Casey

Pierce, Cushman and Hood, 1912: 48, 79.

Curculionid spp. (including *Conotrachelus naso* LeConte or *Curculio pardalis* (Chittenden))Brooks, F. E., 1911a: 190, W. Va. (as "*Myophasia*" ex *Balaninus* "*pardalus*" or *Curculio naso*); Brooks, F. E., 1910: 182, W. Va. (as "*Myophasia*" ex *Balaninus* "*pardulus*" or *Curculio naso*).

Unidentified Coleoptera

Miscellaneous

Aldrich, 1915: 82.

HOST: LEPIDOPTERA

Noctuidae

Pseudaletia unipuncta (Haworth)Lintner, 1897: 210 (as "*Myophasia*" ex *Leucania*).

STAGE PARASITIZED: Larva (Coleoptera, Lepidoptera), pupa (Coleoptera).

IMMATURES: Pupa — Webster, 1899: (pl. 3).

BIOLOGY: Pierce, Cushman and Hood, 1912: (48, 77-80);

Ainslie, G. G., 1910: (140-141).

Myiophasia "*aenea* Wiedemann" probably

HOST: COLEOPTERA

Curculionidae

Conotrachelus nenuphar (Herbst)

Armstrong, 1958: 17, Ont.

Myiophasia "aenia," see *Myiophasia* "aenea"

"*Myiophasia atra* (Robineau-Desvoidy)"

HOST: COLEOPTERA

Curculionidae

Curculio uniformis (LeConte)

Keen, 1958: 70, Oreg.

HOST: LEPIDOPTERA

Arctiidae

Halisidota caryae (Harris)

Fattig, 1949: 22.

Olethreutidae

Laspeyresia caryana (Fitch)

Fattig, 1949: 22, Ga.

Pyralidae

Acrobasis caryae (Grote)

Fattig, 1949: 22.

Myiophasia clistoides (Townsend)

All records as *Ennyomma clistoides*.

HOST: COLEOPTERA

Curculionidae

Chalcodermus aeneus Boheman

Pierce, Cushman and Hood, 1912: 77; Pierce, 1908: 382,
390; Howard, 1895: 280, La.

Myiophasia globosa (Townsend)

HOST: COLEOPTERA

Curculionidae

Anthonomus grandis Boheman

Fattig, 1949: 22-23; Bissell, 1945: 420-440, pls. 1-2, 4, Tex.;
Stuckey, 1941: 116 (ex cotton boll weevil); Allen, 1929:

682, Miss.; Townsend, 1915: 113, Ark., La., Okla. (as *Euloewia*); Hunter and Pierce, 1912: 142, fig. d on pl. 18, La. (as *Ennyomma*); Pierce, Cushman and Hood, 1912: 48, La. (as *Ennyomma*); Pierce, 1910: 453 (as *Ennyomma* ex boll weevil); Pierce, 1908: 382, 389, La. (as *Ennyomma*); Townsend, 1908: 58, La. (as *Ennyomma*).

Chalcodermus aeneus Boheman

Dupree and Beckham, 1955: 12, Ga.; Fattig, 1949: 22-23, Ga.; Hetrick, 1947: 11, 20, 22, Va. (as "Myophasia"); Bissell, 1945: 417-440, pls. 1-2, 4, Ga.; Stuckey, 1943: 49-50, Ga. (ex cowpea curculio); Bissell, 1942: 8-9, Ga. (as *Pseudoclista* ex cowpea curculio); Stuckey, 1942: 78, Ga. (ex cowpea curculio); Stuckey, 1941: 115-116, unnumbered fig. on p. 115, Ga. (ex cowpea curculio); Bissell, 1940: 845, Ga.; Stuckey, 1940: 60, Ga.; Arant, 1938: 28, Ala. (as "Myophasia"); Stuckey, 1938: 66, Ga.; Arant, 1933: 28, Ala.; Townsend, 1915: 113, S.C. (as *Euloewia*); Townsend, 1913b: 115, S.C. (as *Ennyomma*); Walton, 1913a: 204, S.C.; Hunter and Pierce, 1912: 142, fig. d on pl. 18 (as *Ennyomma* ex cowpea curculio); Pierce, Cushman and Hood, 1912: 48, 77, S. U.S. (as *Ennyomma*).

Chalcodermus collaris Horn

Bissell, 1940: 845, Ga.

Chalcodermus inaequicollis Horn

Bissell, 1940: 845, Ga.

Chalcodermus spp.

Townsend, 1915: 113, La. (as *Euloewia*).

Conotrachelus affinis Boheman

Mutchler and Weiss, 1925: 11.

Conotrachelus anaglypticus (Say)

Mutchler and Weiss, 1925: 15.

Conotrachelus aratus Germar

Mutchler and Weiss, 1925: 13.

Conotrachelus nenuphar (Herbst)

Fattig, 1949: 22-23; Stuckey, 1941: 116 (ex plum curculio); Snapp, 1930: 77, 79-80, Ga.

Trichobaris trinotata (Say)

Allen, 1929: 682, Miss.

Unidentified Curculionidae

Townsend, 1915: 113, Ark., La., Okla. (as *Euloewia* ex weevils other than *Anthonomus grandis*).

STAGE PARASITIZED: Larva.

IMMATURES: First instar larva — Bissell, 1945: (descr., pls. 1-2, 4); Stuckey, 1941: (unnumbered fig. on p. 115).

Second instar larva — Bissell, 1945: (descr., pls. 1-2, 4).

Third instar larva — Bissell, 1945: (descr., pls. 1-2, 4).

Pupa — Bissell, 1945: (descr., pls. 1-2, 4); Hunter and Pierce, 1912: (fig. d on pl. 18).

BIOLOGY: Bissell, 1945: (417-440); Stuckey, 1943: (49-50); Bissell, 1942: (8-9); Stuckey, 1941: (115-116).

Myiophasia metallica (Townsend)

HOST: COLEOPTERA

Curculionidae

Conotrachelus spp.

Johnson, 1925: 184, Conn., Mass., N.H., R.I.

Curculio spp.

Johnson, 1925: 184, Conn., Mass., N.H., R.I. (ex "*Balanius*").

Sphenophorus incurrens Gyllenhal

Van Zwaluwenburg, 1951: 375, Mexico (Pue., V.C.) (as "*Miophasia*" ex *Calendra*).

Caneborer(-s)

Pemberton, 1948a: 64, Mexico.

HOST: LEPIDOPTERA

Noctuidae

Pseudaletia unipuncta (Haworth)

Most records ex *Leucania unipuncta*.

*Quaintance and Jenne, 1912: 149 (ex *Heliophila*, Townsend, 1893, questionable host); *Brauer and Bergenstamm, 1895: 563, 597, N. Amer. (Townsend, 1893, questionable host); Townsend, 1893: 467, Ill. (as *Phasioclista*, questionable host).

Myiophasia nigrifrons (Townsend)

HOST: COLEOPTERA

Curculionidae

Conotrachelus affinis Boheman

Townsend, 1915: 113, Mo. (as *Ennyommopsis* ex hickory nut curculio).

Curculio auriger (Casey)

Brooks, F. E., and Cotton, 1929: 15, W. Va.

Curculio proboscideus (Fabricius)

Brooks, F. E., and Cotton, 1929: 15, fig. B on pl. 5, W. Va.

Curculio strictus (Casey) probably

Brooks, F. E., and Cotton, 1929: 15.

Curculio spp.

Brooks, F. E., and Cotton, 1929: 15, Md. (ex *Curculio* larvae in acorns of *Quercus nana* (Marshall) Sargent).

STAGE PARASITIZED: Larva.

IMMATURES: Pupa — Brooks, F. E., and Cotton, 1929: (fig. B on pl. 5).

Myiophasia robusta Coquillett

HOST: COLEOPTERA

Curculionidae

Sphenophorus robustus Horn

Essig, 1926: 578, Calif.; Pierce, 1908: 381, 392; Pierce, 1908a: 42; Coquillett, 1897: 18, 23, Calif.

STAGE PARASITIZED: Larva.

Myobiopsis Townsend (Dexiinae: Leskiini)*Myobiopsis depilis* (Coquillett)

HOST: LEPIDOPTERA

Pyralidae

Macrobotys aeglealis (Walker)

Judd, 1954: 403, Ont. (as *Leskiopalpus*).

STAGE PARASITIZED: Larva.

Myobiopsis flavipennis (Wiedemann)

All records as *Leskiopalpus flavipennis*.

HOST: LEPIDOPTERA

Pyralidae

"Blepharomastix" *ebulialis* Guenée

Wolcott, 1951a: 477, P.R. (ex *Lamprosema* (or *Blepharomastix*)).

Unidentified Lepidoptera

Leaf-folder caterpillar(-s)

Martorell, 1945: 213, P.R.

STAGE PARASITIZED: Larva.

Miocera cremides, see *Dinera grisescens*

Myocera tabanivora, see *Carinosillus tabanivorus*

"*Myophasia*" *aenea*, see *Myiophasia* "*aenea*"

"*Myophasia*" *globosa*, see *Myiophasia globosa*

Mystacella Wulp (Goniinae: Sturmiini)

Mystacella chrysoprocta (Wiedemann)

HOST: COLEOPTERA

Scarabaeidae

Xyloryctes sp.

Aldrich, 1932: 4, Mass. (as *Macromeigenia*).

Mystacella spp.

HOST: LEPIDOPTERA

Arctiidae

Isia isabella (J. E. Smith)

Townsend, 1895: 75, Colo. (as *Mystacella* n. sp. ex
Pyrrharctia).

Noctuidae

Acronicta americana (Harris) probably

Townsend, 1895: 75, Colo. (ex *Acronycta*).

Myxexorista sp., see "*Zenillia*" spp.

"*Myxexorsita*" sp., see "*Zenillia*" spp.

Neadmontia limata, see *Dexia limata*

Nemochaeta lateralis, see *Archytas lateralis*

Nemoraea Macquart (Tachininae: Nemoraeni)

Nemoraea hyphantriae, see *Hyphantrophaga hyphantriae*

Nemoraea leucanae, see *Winthemia leucanae*

Nemoraea sp. near *leucanae*, see *Winthemia* sp. near *leucanae*

Nemoraea "*leucaniae*," see *Winthemia leucanae* or *Winthemia* spp.

(including *W. leucanae* probably or *W. militaris*)

Nemoraea nigricornis, see *Mericia ampelus*

"*Nemoraea*" sp.

HOST: LEPIDOPTERA

Arctiidae

Turuptiana permaculata (Packard)

Townsend, 1895: 75, Colo. (as *Nemoraea* n. sp. ex
Ecpantheria).

- "Nemorea leucaniae," see *Winthemia leucanae*
Nemorilla Rondani (Goniinae: Winthemiini)
Nemorilla floralis, see *Nemorilla pyste*
Nemorilla sp. near *floralis*, see *Nemorilla* sp. near *pyste*
Nemorilla "floralis," see *Nemorilla pyste*
Nemorilla maculosa, see *Nemorilla pyste*
Nemorilla maculosa probably, see *Nemorilla pyste* probably
Nemorilla "maculosa," see *Nemorilla pyste*
Nemorilla phycitae, see *Nemorilla pyste*
Nemorilla pyste (Walker)
 HOST: COLEOPTERA

Coccinellidae

Epilachna varivestis Mulsant

Douglass, 1933: 23, N. Mex. (as *N. maculosa* (Meigen) ex *E. corrupta* Mulsant).

HOST: LEPIDOPTERA

Gelechiidae

Filatima persicaeella (Murtfeldt)

Ingerson, 1918: 12, Mich. (as *Exorista* ex *Gelechia confusella* Chambers).

Glyphipterygidae

Anthophila pariana (Clerck)

Most records ex *Hemerophila pariana*.

Schaffner, 1959: 41, 52, [N.E. U.S.] (as *N. floralis* (Fallén) ex *Anthophila*); *Cameron, 1931: 41, Conn. (as *Exorista*, Porter and Garman, 1923); *Cameron, 1931: 41, Conn. (as *N. maculosa*, Porter and Garman, 1923); Cameron, 1931: 42, N.S. (as *N. floralis*); Porter and Garman, 1923: 260-261, Conn., N.Y. (as *Exorista*); Porter and Garman, 1923: 261, Conn. (as *N. maculosa*); Britton, 1922: 188, Conn. (as *Exorista*).

Hesperiidae

Urbanus proteus (Linnaeus)

Wilson, C. E., 1923a: 5, Virgin Islands (as *Exorista* ex *Eudamus*).

Lycaenidae

Strymon ontario autolyucus (Edwards)

Williston, 1889: 1914, 1921, Tex. (as *Exorista scudderi*
Williston ex *Thecla autolyucus*).

Lymantriidae

Porthetria dispar (Linnaeus)

Howard and Fiske, 1911: 90 (as *Exorista*).

Noctuidae

Achatodes zea (Harris)

Silver, J. C., 1933: 18, La. (as *N. floralis*).

Heliothis zea (Boddie)

Wilson, C. E., 1923a: 15, Virgin Islands (as *Exorista* ex *H. obsoleta* (Fabricius)).

Prodenia sunia (Guenée)

Wilson, C. E., 1923: 7, Virgin Islands (as *Exorista* ex
"Xylomiges").

Spodoptera frugiperda (J. E. Smith)

All records ex *Laphygma frugiperda*.

Wilson, C. E., 1923: 5, Virgin Islands (as *Exorista*); Wilson,
C. E., 1923a: 19, Virgin Islands (as *Exorista*).

Trichoplusia ni (Hübner)

Wilson, C. E., 1923a: 22, Virgin Islands (as *Exorista* ex
Plusia).

Oecophoridae

Depressaria pastinacella (Duponchel)

Schaffner, 1959: 39, 52, [N.E. U.S.] (as *N. floralis* ex *D. heracliana* (Linnaeus)).

Olethreutidae

Ancylis comptana (Frölich)

Summerland, 1938: 172-173, Kans. (as *N. floralis*); Fink,
1932: 555-556, fig. 9c, N.J. (as *Exorista*); Lewis, 1925:
611, Ohio (as "Exoriste").

Ancylis comptana fragariae (Walsh and Riley)

Petterson, 1946: 323, Ohio (as *N. "floralis"*); Neiswander,
R. B., 1944: 8, Ohio (as *N. floralis*).

Episimus argutanus (Clemens)

Schaffner, 1959: 20, 52, [N.E. U.S.] (as *N. floralis*).

Evora hemidesma (Zeller)

Roberts, 1966: 169, Iowa (as *N. floralis*).

Exartema sericoranum Walsingham

Schaffner, 1959: 22, 52, R.I. (as *N. floralis*).

Grapholitha molesta (Busck)

Most records ex *Laspeyresia molesta*.

Allen, 1962: 11, 112, Mass., N.J., N.Y. (as *N. floralis* ex *Grapholitha*); Brunson and Allen, 1948: 447 (as *N. floralis* ex *Grapholitha*); Steenburgh and Boyce, 1938: 73, Ont. (as *N. maculosa*); Steenburgh, 1935: 71, Ont. (as *N. maculosa*); McConnell, 1932: 367-368, Md. (as *N. phycitae* (LeBaron)); Steenburgh, 1931: 59, Ont. (as *N. maculosa*); Haeussler, 1930: 367, Md. (as *N. phycitae*); Stearns, 1928: 7, Md. (as *N. phycitae*); Garman, P., 1918: 117, Md. (as *N. phycitae*).

Rhopobota naevana (Hübner)

Franklin, 1950: 57, 60, fig. 4B (as *N. maculosa* ex *R. vacciniana* (Packard)).

Rhyacionia buoliana (Schifferrmüller)

Schaffner, 1959: 23, 52, [N.E. U.S.] (as *N. floralis*).

Spilonota ocellana (Denis and Schifferrmüller)

Schaffner, 1959: 25, 52, [N.E. U.S.] (as *N. floralis*); Stultz, 1955: 80, N.S.

Pyralidae

Acrobasis betulella Hulst

Daviault, 1935: 318-319, Que. (as *N. maculosa*).

Acrobasis caryae Grote

Nickels, Pierce and Pinkney, 1950: 2, 19-20, Tex. (as *N. floralis*); Fattig, 1949: 23 (as *N. floralis*); Gill, 1930: 71-72 (as *N. "mocolosa"* ex pecan nut case-bearer); Gill, 1930: 71-72 (as *Exorista* ex pecan nut case-bearer); Leiby, 1925: 18-19, N.C. (as *N. maculosa*); Brimley, 1922: 22, N.C. (as *Exorista*); Kirk, 1913: 258, Conn. (as *Exorista*).

Acrobasis caryivorella Ragonot

Schaffner, 1959: 14, 52, [N.E. U.S.] (as *N. floralis*); Nickels, 1951: 434, Tex. (as *N. maculosa*); Reinhard, 1922: 72, Tex. (as *Exorista*).

Acrobasis comptoniella Hulst

Schaffner, 1959: 15, 52, [N.E. U.S.] (as *N. floralis*).

Acrobasis coryliella Dyar

Schaffner, 1959: 14, 52, Mass. (as *N. floralis*).

Acrobasis indigenella (Zeller)

Leiby, 1925: 9, N.C. (as *Exorista* ex *A. nebulella* Riley);
 Brimley, 1922: 22, N.C. (as *Exorista* ex *A. nebulella*);
 Coquillett, 1897: 14, 26, Ind., Mo. (as *Exorista* ex
Mineola "indiginella"); *Brauer and Bergenstamm, 1895:
 554, 580, N. Amer. (as *Exorista phycitae*, Townsend,
 1893; Riley, 1872); *Brauer and Bergenstamm, 1895:
 554, 577, 602, Ill. (as (*Tachina*) *phycitae* ex *Phycita*
nebulo Walsh, Riley, 1872); Townsend, 1893: 467, Ill.
 (as *Exorista phycitae*); LeBaron, 1872: 123, Ill. (as
Tachina phycitae ex *Phycita nebulo*); Riley, 1872: 40-41
 (as *Tachina phycitae* ex *Phycita nebulo*).

Acrobasis juglandis (LeBaron)

Gill, 1930: 71 (as *Exorista* ex leaf case-bearer).

Asciodes anormalis (Guenée) probably

Wilson, C. E., 1923a: 27, Virgin Islands (as *Exorista* ex
Sylepta helcitalis Walker).

Desmia funeralis (Hübner)

Donohoe and Kaloostian, 1937: 967-968, Calif. (as *N.*
maculosa); Knowlton and Janes, 1931: 147, Utah (as *N.*
floralis ex grape leaf-folder); Allen, 1929: 686, Miss. (as
N. floralis); Strauss, 1916: 9, 12, D.C. (as "Exoriste").

Diaphania hyalinata (Linnaeus)

Wolcott, 1951a: 478, P.R. (as *N. maculosa* ex *Margaronia*);
 Russo, 1927a: 63, 76, Dominican Republic (as *N. floralis*
 ex *Margaronia*); Wolcott, 1924: 222, P.R. (as *N.*
maculosa).

Diatraea saccharalis (Fabricius)

Wilson, C. E., 1923a: 18, Virgin Islands (as *Exorista*).

Hedylepta indicata (Fabricius)

Wolcott, 1951a: 478, P.R. (as *N. floralis* ex "Hydelepta");
 Plank, 1945: 26-27, P.R. (as *N. floralis*); Wilson, C. E.,
 1923a: 3, Virgin Islands (as *Exorista* ex *Nacoleia*).

Hellula phidilealis (Walker)

Wilson, C. E., 1923a: 23, Virgin Islands (as *Exorista*).

Hellula rogatalis (Hulst)

Most records ex *Hellula undalis* (Fabricius).

Chittenden and Marsh, 1912: 31 (as *Exorista*);
 *Chittenden, 1900: 60, S.C. (as *Exorista*, Coquillett,
 1897); Forbes, S. A., and Hart, 1900: 460, Ill. (as
Exorista "piste"); Chittenden, 1899: 56, Ga. (as
Exorista "piste"); Coquillett, 1897: 14, 26, S.C. (as
Exorista ex pyralid).

Herpetogramma bipunctalis (Fabricius)

Russo, 1927a: 183, Dominican Republic (as *N. floralis* ex *Psara* "*bispunctalis*").

Hulstia undulatella (Clemens)

*Essig, 1926: 581, Calif. (as *N. maculosa*, Titus, 1905); Titus, 1905: 38-39, Calif. (as *Exorista* ex "*Hulstea*").

Hymenia recurvalis (Fabricius)

Poos, 1928: 146, Va. (as *N. maculosa* ex *H. fascialis* (Cramer)); Wilson, C. E., 1923a: 7-8, Virgin Islands (as *Exorista* ex *Zinckenia fascialis*).

Loxostege similalis (Guenée)

Butler, 1958: 562, Ariz. (as *N. floralis*); Smith, R. C., and Franklin, 1954: 31, Kans. (as *N. maculosa*); Fattig, 1949: 23 (as *N. floralis*); Reinhard, 1922: 72, Tex. (as *Exorista*).

Maruca testulalis (Geyer)

Bruner, S. C., 1935: 12, Cuba (as *N. floralis*).

Nephopteryx subcaesiella (Clemens)

Schaffner, 1959: 17, 52, [N.E. U.S.] (as *N. floralis* ex *Salebria*).

Ostrinia nubilalis (Hübner)

All records ex *Pyrausta nubilalis*.

Baker, W. A., Bradley and Clark, 1949: 8, 38-39, 53, fig. 38Ab (as *N. floralis*); Baker, W. A., Bradley and Clark, 1949: 176, fig. 38Aa, New England (as *N. maculosa*); Clark, C. A., 1934: 18, 20, 31, 36 (as *N. floralis*, lab.); Johnson, 1925: 196, Conn., Maine, Mass., N.H., R.I. (as *N. maculosa*); Caffrey, 1919: 21, Mass. (as *Exorista*); Vinal and Caffrey, 1919: 57-58, Mass. (as *Exorista*).

Ostrinia spp. (including *O. obliteralis* (Walker) and/or *O. penitalis* Grote)

Poos, 1927: 82-83, Ohio (as *Exorista* ex *Pyrausta penitalis* or *P. ainsliei* Heinrich).

Pachyzancla bipunctalis (Fabricius)

Wilson, C. E., 1923a: 9, Virgin Islands (as *Exorista*); Jones, T. H., 1915: 8, P.R. (as *Exorista*).

Phlyctaenia coronata tertialis (Guenée)

All records ex *Phlyctaenia tertialis*.

Schaffner, 1959: 10, 52, [N.E. U.S.] (as *N. floralis*); Balduf, 1929: 239-240, Ill., Ohio (as *N. floralis*).

Pilocrocis tripunctata (Fabricius)

Jones, T. H., 1917: 9, La. (as *Exorista*).

Pococera scabridella (Ragonot)

Wolcott, 1951a: 478, P.R. (as *N. maculosa* ex *Tetralopha*).

Psara pallicaudalis Snellen

Fattig, 1949: 23 (as *N. floralis* ex *Pachyzancla periusalis* (Walker)).

Pyrausta futilalis (Lederer)

Schaffner, 1959: 11, 52, [N.E. U.S.] (as *N. floralis*).

Pyrausta signatalis (Walker)

Schaffner, 1959: 12, 52, Mass. (as *N. floralis*).

Pyrausta tyralis (Guenée)

Allen, 1929: 686, Miss. (as *N. floralis*); Allen, 1924: 24, Miss. (as *N. maculosa*).

Tetralopha asperatella (Clemens)

Schaffner, 1959: 13, 52, [N.E. U.S.] (as *N. floralis*).

Unidentified Pyralidae

Coquillett, 1897: 14, 26, D.C. (as *Exorista*).

Tortricidae

Acleris minuta (Robinson)

Franklin, 1950: 57, 60, fig. 4B (as *N. maculosa* ex *Peronea*);
Johnson, 1925: 196, Conn., Maine, Mass., N.H., R.I. (as
N. maculosa ex "*Peronia*").

Acleris variana (Fernald)

Miller, C. A., 1966: 603, N.S.; Reeks, Forbes and Cuming,
1947: 12, [maritime prov.]; Schedl, 1932: 1-2, N.S. (as
"*Memorilla*" *maculosa* ex *Peronea*).

Adoxophyes furcatana (Walker)

Schaffner, 1959: 27, 52, Mass. (as *N. floralis*).

Amorbia emigratella Busck

Vickery, 1925: 141, Tex. (as *N. maculosa*).

Aphelia alleniana (Fernald)

Martin, 1958: 46, Ont. (ex *Tortrix*).

Archippus packardianus (Fernald)

Daviault, 1950a: 150, Que. (ex *Tortrix*); Daviault, 1946:
117, Que. (ex *Tortrix* "*parckardiana*").

Archips argyrospilus (Walker)

Prokopy, 1968: 350, Conn.; Schaffner, 1959: 30, 52, [N.E.
U.S.] (as *N. floralis*); *Essig, 1926: 581, Colo., N. Mex.
(as *N. maculosa*, Gill, 1913); Gill, 1913: 102, Colo. or N.
Mex. (as *Exorista*).

Archips cerasivoranus (Fitch)

Schaffner, 1959: 30, 52, [N.E. U.S.] (as *N. floralis*);
Thompson, 1953: 19-20 (as *N. maculosa*); Hoffman,

1936: 210-211, Minn. (as *N. maculosa* ex *Cacoecia*);
Herrick, 1912: 291, N.Y. (as *Carcelia*).

Archips fervidanus (Clemens)

Schaffner, 1959: 29, 52, [N.E. U.S.] (as *N. floralis*).

Archips griseus (Robinson)

Prokopy, 1968: 350-351, Conn.

Archips purpuranus (Clemens)

Schaffner, 1959: 31, 52, [N.E. U.S.] (as *N. floralis*).

Archips rosanus (Linnaeus)

Schaffner, 1959: 31, 52, Mass. (as *N. floralis*); Johnson,
1925: 196, Conn., Maine, Mass., N.H., R.I. (as *N. maculosa*); Whitehead, 1925: 79, N.S. (as *Exorista* ex
Cacoecia); Walden, 1914: 226, Conn. (as *Exorista*).

Argyrotaenia velutinana (Walker)

Anon., 1963c: 20, Ont.

Choristoneura fumiferana (Clemens)

Blais, 1965: 131, Que.; Blais, 1960: 388, 390, Que. (ex
spruce budworm); Coppel, 1960: 94, 97, B.C.; Carolin
and Coulter, 1959: 553, Oreg. (as *N. floralis*); McGugan
and Blais, 1959: 765, Ont.; Schaffner, 1959: 32, 52,
[N.E. U.S.] (as *N. floralis*); Boyes and Wilkes, 1953:
126, 149 (lab.); Raizenne, 1952: 219, Ont.; Ross, 1952:
112, fig. 8, Canada; Dowden and Carolin, 1950: 778,
N.Y. (as *N. floralis* ex *Archips*); Wilkes, Coppel and
Mathers, 1949: 140, 146, 149, 151, B.C.; Dowden,
Buchanan and Carolin, 1948: 458 (as *N. floralis* ex
Archips); Daviault, 1946: 117, Que. (as *N. maculosa* ex
Archips); Richmond, 1941: 57, Sask. (as *N. maculosa* ex
Archips); Richmond, 1940: 50, Man., Ont., Sask. (as *N. maculosa*
ex *Cacoecia*); Bedard, 1938: 191 (as *N. floralis*
ex *Cacoecia*); Richmond, 1938: 45, Man., Ont. (as
"Hemorilla" *maculosa* ex *Cacoecia*); Tothill, 1913: 71,
Que. (as *Exorista* ex *Tortrix*).

Choristoneura houstonana (Grote)

Anon., 1966c: 323, Kans.

Choristoneura parallela (Robinson)

Franklin, 1943: 26, Mass. (as *N. floralis* ex *Cacoecia*).

Choristoneura pinus Freeman

Dixon and Benjamin, 1963: 267, Wis. (as *N. floralis*).

Choristoneura rosaceana (Harris)

All records ex *Archips rosaceana*.

Schaffner, 1959: 31, 52, [N.E. U.S.] (as *N. floralis*); Davis,
J. J., 1912: 97, Ill. (as *Exorista*).

Pandemis pyrusana (Kearfott)

Newcomer and Carlson, 1953: 1080, Wash. (as *N. floralis*).

Sparganothis sulfureana (Clemens)

Beckwith, 1938: 255, N.J. (as *N. floralis*).

Unidentified Tortricidae

Krombein, 1967: 69, Fla. (as *N. floralis*); Titus, 1905: 39, Calif. (as *Exorista*); Coquillett, 1897: 15, 27, Tex. (as *Exorista*).

Yponomeutidae

Yponomeuta malinella Zeller

Parrott and Schoene, 1912: 30, figs. 8-9, N.Y. (as *Exorista arvicola* (Meigen)).

Yponomeuta multipunctella Clemens

Most records ex *Hyponomeuta multipunctella*.

Judd, 1955: 267, Ont. (as *N. floralis*); Judd, 1954: 403, Ont. (as *N. floralis*); Putman, 1943: 223, Ont. (as *N. maculosa* ex *Yponomeuta*); Coquillett, 1897: 14, 25, D.C. (as *Exorista*); Brauer and Bergenstamm, 1895: 564, 595, N. Amer. (as *Nemorilla* sp.).

Yponomeuta padella (Linnaeus)

Parrott and Schoene, 1912: 30, N.Y. (as *Exorista arvicola*).

Unidentified Lepidoptera

Miscellaneous

Aldrich and Webber, 1924: 6, Canada, U.S. (as *N. maculosa* ex moths and butterflies).

STAGE PARASITIZED: Larva (Coleoptera, Lepidoptera), pupa (Lepidoptera)

IMMATURES: Egg — Franklin, 1950: (fig. 4B); Fink, 1932: (descr., fig. 9c); Lewis, 1925: (descr.); Parrott and Schoene, 1912: (descr., fig. 8).

Pupa — Ross, 1952: (fig. 8); Baker, W. A., Bradley and Clark, 1949: (fig. 38Aa-Ab); Fink, 1932: (descr.); Lewis, 1925: (descr.); Greene, 1922: (descr., fig. 34 on pl. 8); Parrott and Schoene, 1912: (fig. 9).

BIOLOGY: Allen, 1962: (11, 112); Schaffner, 1959: (10-15, 17, 20, 22-23, 25, 27, 29-32, 39, 41, 52); Judd, 1955: (267); Smith, R. C., and Franklin, 1954: (31); Franklin, 1950: (57, 60); Nickels, Pierce and Pinkney, 1950: (2, 19-20); Summerland, 1938: (172-173); Fink, 1932: (555-556); Gill, 1930: (71-72); Balduf, 1929: (239-240); Lewis, 1925: (611); Parrott and Schoene, 1912: (30).

Nemorilla pyste (Walker) probably
HOST: LEPIDOPTERA

Pyralidae

Acrobasis rubrifasciella Packard

Balduf, 1968: 474-475, Minn. (as *N. maculosa* (Meigen) probably).

STAGE PARASITIZED: Pupa.

Nemorilla sp. near pyste (Walker)
HOST: LEPIDOPTERA

Pyralidae

Acrobasis indigenella (Zeller)

Gill, 1917: 15, Fla. (as *Exorista* ex *A. nebulella* Riley).

Asciodes gordialis (Guenée)

Patton, 1958: 78, Fla. (as gen. sp. ? near *N. floralis* (Fallén), incomplete rearing — pupa).

“*Nemorilla trivittata* (Wulp)” probably
All records as *Exorista trivittata* probably.
HOST: LEPIDOPTERA

Noctuidae

Acronicta oblinita (J. E. Smith)

All records ex *Apatela oblinita*.

*Brauer and Bergenstamm, 1895: 554, 583, N. Amer.
(Townsend, 1893); Townsend, 1893: 467, Ill.

Nemorilla sp., see *Nemorilla pyste*

Nemorilla spp.

HOST: LEPIDOPTERA

Tortricidae

Choristoneura pinus Freeman

Richmond, 1941a: 209, Man. (ex jack pine budworm).

Neocelatoria ferox, see *Celatoria setosa*

Neopales macra, see “*Phorocera*” spp.

Neopales “maera,” see “*Phorocera*” spp.

Neopales tortricis, see *Hemisturmia tortricis*

Neopales sp., see *Pales* spp.

“*Neophorocera*” hamata, see *Diplostichus lophyri*

Neophorocera claripennis, see *Euphorocera claripennis*

- Neophorocera edwardsii*, see *Euphorocera edwardsii*
Neophorocera hamata, see *Diplostichus lophyri*
Neophorocera sp., see *Euphorocera* sp.
Neothelaira aurifrons, see *Aporotachina dexina*
Neothelaira chaetoneura, see *Aporotachina chaetoneura*
 New Guinea parasite, see *Lixophaga sphenophori*
 New Guinea tachinid, see *Lixophaga sphenophori*
Nilea Robineau-Desvoidy (Goniinae: Eryciini)
Nilea carpocapsae (Townsend)
 HOST: LEPIDOPTERA

Olethreutidae

Laspeyresia pomonella (Linnaeus)

Townsend, 1919: 580, Ark. (as *Eutritochaeta* ex codling moth).

STAGE PARASITIZED: Pupa.

Nilea noctuiformis (Smith)

Most records as *Phorocera noctuiformis*.

HOST: LEPIDOPTERA

Noctuidae

Unidentified Noctuidae

*Aldrich and Webber, 1924: 75, Mass. (Smith, H. E., 1915);
 Smith, H. E., 1915: 102, Mass. (as *Neopales*).

Olethreutidae

Laspeyresia pomonella (Linnaeus)

Aldrich and Webber, 1924: 76, Ark. (ex *Carpocapsa*).

STAGE PARASITIZED: Larva, pupa.

Nilea sp. near *noctuiformis* (Smith)

HOST: LEPIDOPTERA

Noctuidae

Acronicta interrupta Guenée

Raizenne, 1952: 41, Ont. (as *Phorocera*).

Nilea sternalis (Coquillett)

All records as *Phorocera sternalis*.

HOST: LEPIDOPTERA

Noctuidae

Acronicta impressa Walker

Schaffner and Griswold, 1934: 58-59, 115, Maine (ex *Acronycta*).

Tortricidae

Archips cerasivoranus (Fitch)

Powell, 1962: 136, Calif.

BIOLOGY: Schaffner and Griswold, 1934: (58-59, 115).

Nowickia Wachtl (Tachininae: Tachinini)*Nowickia algens* (Wiedemann)

HOST: LEPIDOPTERA

Noctuidae

Apamea lignicolor (Guenée)

All records ex *Hadena lignicolor*.

*Brauer and Bergenstamm, 1895: 558, 593, N. Amer. (as "*Jurinea*," Townsend, 1893b); Townsend, 1895: 70, Colo. (as *Jurinia*); Townsend, 1893b: 576-577, Colo. (as *Jurinia*).

Euxoa messoria (Harris)

Brauer and Bergenstamm, 1895: 554, 581, N. Amer. (as (?*Fabricia*) ex *Agrotis cochrani* (Riley)).

Cutworm(-s)

Aldrich, 1915: 83 (as *Echinomyia*).

Nowickia hispida (Tothill)

HOST: LEPIDOPTERA

Noctuidae

Polia purpurissata (Grote)

Wood, Neilson, Maxwell and McKiel, 1954: 173, N.B. (as *Fabriciella*).

STAGE PARASITIZED: Larva.

Nowickia latifrons (Tothill)

HOST: LEPIDOPTERA

Noctuidae

Crymodes devastator (Brace)

King, K. M., 1925: [14], Sask. (as *Fabriciella* ex *Sidemia*).

STAGE PARASITIZED: Larva.

Obolocera Townsend (Goniinae: Eryciini)

Obolocera rapae (Smith)

HOST: LEPIDOPTERA

Pieridae

Pieris rapae (Linnaeus)

All records ex *Pontia rapae*.

Johnson, 1925: 204, Mass.; Smith, H. E., 1917a: 140,
Mass. (as *Homoeonychia*).

Ochromeigenia ormoides, see *Hamaxia incongrua*

Ocyptera carolinae, see "*Cylindromyia carolinae*"

Ocyptera euchenor, see *Cylindromyia euchenor*

Ocytata Gistel (Goniinae: Erythrocerini)

Ocytata pallipes (Fallén)

All records as *Rhacodineura antiqua* (Meigen).

HOST: DERMAPTERA

Forficulidae

Forficula auricularia Linnaeus

Sweetman, 1936: 233, N. Amer.; Mote, 1931: 948-950,
952-955 (lab.).

BIOLOGY: Sweetman, 1936: (233).

Oedematocera Townsend (Goniinae: Hyperecteinini)

Oedematocera dampfi Aldrich

HOST: ORTHOPTERA

Acrididae

Schistocerca cancellata (Serville)

All records ex *Schistocerca paranensis* (Burmeister).

Greene, 1956: 22, fig. 13, Guatemala, Mexico; Aldrich,
1928: 301; Townsend, 1928a: 152 (as *Schistocercophaga*);
Aldrich, 1927: 589-590, Guatemala, Mexico;
Aldrich, 1927a: 17, Mexico.

STAGE PARASITIZED: Adult.

IMMATURES: Third instar larva — Greene, 1956: (descr., fig.
13).

BIOLOGY: Aldrich, 1927: (589-590).

Oedematocera flaveola (Coquillett)

HOST: ORTHOPTERA

Gryllacrididae

Ceuthophilus latibuli Scudder

Hubbell, 1936: 507, Fla.

Oedematocera flaveola (Coquillett) probably

HOST: ORTHOPTERA

Gryllacrididae

Ceuthophilus spp.

Hubbell, 1936: 507, Mich., N.C., Ohio.

BIOLOGY: Hubbell, 1936: (507).

Oedematocera gilvipes, see *Gilvella gilvipes**Oedematopteryx fumosa*, see *Alophorella fumosa**Oedematopteryx pulverea*, see *Alophorella pulverea**Oedematopteryx* sp., see *Alophorella* spp.*Oestrophasia ochracea*, see *Euphasiopteryx ochracea**Omotoma* Lioy (Goniinae: Winthemiini)*Omotoma fumiferanae* (Tothill)

HOST: LEPIDOPTERA

Geometridae

Alsophila pometaria (Harris)Schaffner and Griswold, 1934: 90, 122 (as *Winthemia*).*Phigalia titea* (Cramer)Schaffner and Griswold, 1934: 97, 122 (as *Winthemia*).

Noctuidae

Amphipyra pyramidoides GuenéeSchaffner and Griswold, 1934: 52, 122 (as *Winthemia*).*Lithophane antennata* (Walker)Schaffner and Griswold, 1934: 49, 122 (as *Winthemia* ex *Graptolitha*).*Lithophane* spp. (including *L. disposita* Morrison, *L. innominata* (Smith) and/or *L. petulca* Grote)Schaffner and Griswold, 1934: 48, 122 (as *Winthemia* ex *Graptolitha* (including *G. disposita*, *G. innominata* and/or *G. petulca*)).*Pyreferra hesperidago* (Guenée)Schaffner and Griswold, 1934: 51, 122 (as *Winthemia* ex *Conistra indirecta* Walker).*Spodoptera ornithogalli* (Guenée)Fattig, 1949: 35, Ga. (as *Winthemia amoena* (Meigen) ex *Prodenia*).

Tortricidae

Choristoneura conflictana (Walker)

Prentice, 1955: 471, Man., Sask.

Choristoneura fumiferana (Clemens)

Blais, 1965: 131, Que. (as *Winthemia amoena*); MacDonald and Webb, 1963: 299, N.B.; Miller, 1963: 229, 237, 239, N.B. (as *Winthemia amoena*); Blais, 1960: 388, 390-392, Que. (ex spruce budworm); Coppel, 1960: 94, 97, B.C.; Carolin and Coulter, 1959: 552-554, Oreg.; McGugan and Blais, 1959: 765, 768, 774-775, Ont.; Boyes and Wilkes, 1953: 126, 151; Coppel, 1953: 191-192, B.C.; MacNay, 1953a: 310, 314, 316, B.C.; Anon., 1952b: 178, 184, 186, B.C.; Anon., 1952g: 288, 292, 295-296, B.C.; Raizenne, 1952: 219, Ont.; Ross, 1952: 112, fig. 12, Canada; Anon., 1950c: 238, 246, 248, B.C.; Dowden and Carolin, 1950: 778-779, N.Y. (ex *Archips*); Anon., 1949: 272, 283, 285-286, B.C., Ont.; Wilkes, Coppel and Mathers, 1949: 140, 146, 149, 151, 153, B.C.; Anon., 1948: 285, 293-295, B.C.; Dowden, Buchanan and Carolin, 1948: 458-463 (ex *Archips*); Coppel, 1947: 40, B.C. (ex *Archips*); Wilkes and Anderson, 1947: 41, 43, Ont., Que. (ex *Archips*); Reinhard, 1931: 14, B.C. (as *Winthemia ex Tortrix*); Essig, 1926: 584, B.C. (as *Winthemia ex Harmologa*); Johnson, 1925: 203, N.H. (as *Winthemia ex spruce budworm*); Lochhead, 1919: 231 (as *Winthemia ex Harmologa*); Johannsen, 1913: 25-26, Canada (as *Winthemia ex Tortrix*); Tothill, 1912: 3, Canada (as *Winthemia ex Tortrix*).

STAGE PARASITIZED: Larva, pupa.

IMMATURES: Pupa — Ross, 1952: (fig. 12).

BIOLOGY: Miller, 1963: (229, 237, 239); Coppel, 1953: (191-192); Coppel, 1947: (40); Schaffner and Griswold, 1934: (122).

Opsotheresia Townsend (Proseninae: Theresiini)*Opsotheresia bigelowi* (Curran)

HOST: DIPTERA

Tabanidae

Unidentified Tabanidae

West, 1928: 821, N.Y. (as *Paraprosena apicalis* (Robineau-Desvoidy)).

STAGE PARASITIZED: Larva.

Ormia Robineau-Desvoidy (Proseninae: Ormiini)

Ormia ochracea, see *Euphasiopteryx ochracea*

Ormia punctata Robineau-Desvoidy

HOST: DIPTERA

Tabanidae

Tabanus sp.

Jones and Anthony, 1964: 57, Fla.

STAGE PARASITIZED: Larva.

Ormia spp. probably

HOST: LEPIDOPTERA

Pyralidae

Crambus spp.

*Brauer and Bergenstamm, 1895: 564, 589, N. Amer.
(Townsend, 1893); Townsend, 1895: 78, Ill.; Townsend,
1893: 468, Ill.

Oswaldia Robineau-Desvoidy (Goniinae: Blondeliini)

Oswaldia albifacies (Townsend)

HOST: LEPIDOPTERA

Noctuidae

Parallelia bistriaris Hübner

Raizenne, 1952: 70, Ont. (as *Parameigenia*).

Pyralidae

Loxostege mancalis (Lederer)

Brimley, 1938: 363, N.C. (as *Masicera* "albifascies");

Brimley, 1922: 23, N.C. (as *Masicera*).

Oswaldia anorbitalis (Brooks)

HOST: LEPIDOPTERA

Noctuidae

Parallelia bistriaris Hübner

Raizenne, 1952: 70, Ont. (as *Aubaeanetia*).

Oswaldia assimilis (Townsend)

HOST: LEPIDOPTERA

Noctuidae

Parallelia bistriaris Hübner

Schaffner and Griswold, 1934: 68, 120, Mass. (as
Aubaeanetia).

BIOLOGY: Schaffner and Griswold, 1934: (68, 120).

Oswaldia aurifrons (Townsend)

All records as *Dexodes aurifrons*.

HOST: HYMENOPTERA

Tenthredinidae

Nematus ventralis Say

*Procter, 1946: 436, Maine (ex willow sawfly, Procter,
1938); Procter, 1938: 377, Maine (ex willow sawfly).

HOST: LEPIDOPTERA

Citheroniidae

Anisota senatoria J. E. Smith

Schaffner and Griswold, 1934: 29-30, 120, N.Y.

BIOLOGY: Schaffner and Griswold, 1934: (29-30, 120).

Oswaldia sp. probably

HOST: LEPIDOPTERA

Noctuidae

Arzama obliqua (Walker)

Brauer and Bergenstamm, 1895: 551, 583 (as *Dexodes* sp.
(? = *Masicera luctuosa* Wulp) ex A. "obliquata").

Oxynops Townsend (Goniinae: Blondeliini)*Oxynops anthracinus* (Bigot)

HOST: LEPIDOPTERA

Pterophoridae

Oidaematophorus homodactylus (Walker)

Schaffner, 1959: 19, 52, [N.E. U.S.].

Oidaematophorus monodactylus (Linnaeus)

Schaffner, 1959: 19, 52, [N.E. U.S.].

BIOLOGY: Schaffner, 1959: (19, 52).

Gen. and sp. near *Oxynops*

HOST: LEPIDOPTERA

Olethreutidae

Petrova albicapitana (Busck)

Lyon, 1957: 75.

Pyralidae

Dioryctria abietella (Denis and Schiffermüller)

Lyon, 1957: 77-78, Ont.

Dioryctria cambiicola (Dyar)

Lyon, 1957: 78, Ont.

Dioryctria disclusa Heinrich

Lyon, 1957: 75, 77-78, Ont.

STAGE PARASITIZED: Larva.

Pacidianus Reinhard (Goniinae: Eumasicerini)

Pacidianus persimilis Reinhard

HOST: LEPIDOPTERA

Citheroniidae

Anisota rubicunda (Fabricius)

Reinhard, 1943: 90, Ont.

Pacidianus sp.

HOST: LEPIDOPTERA

Citheroniidae

Anisota virginensis (Drury)

Patton, 1958: 79, Fla.

STAGE PARASITIZED: Larva.

Pales Robineau-Desvoidy (Goniinae: Sturmiini)

Pales spp.

HOST: COLEOPTERA

Tenebrionidae

Meracantha contracta (Palisot de Beauvois)

Hyslop, 1915: 48, Md. (as *Neopales*).

HOST: LEPIDOPTERA

Notodontidae

Cerura sp.

Brauer and Bergenstamm, 1895: 549, 586, N. Amer. (as
Ctenophorocera ex *Cerura* sp. on willows).

STAGE PARASITIZED: Larva (Coleoptera).

BIOLOGY: Hyslop, 1915: (48).

Palexorista Townsend (Goniinae: Sturmiini)

Palexorista laxa (Curran)

Most records as *Drino munda* (Wiedemann).

HOST: LEPIDOPTERA

Noctuidae

Heliothis virescens (Fabricius)

Bryan, D. E., Jackson and Stoner, 1969: 1, 8, 12 (as *P. laxa*, lab.); Chauthani and Hamm, 1967: 373 (lab.).

Heliothis zea (Boddie)

Bryan, D. E., Jackson and Stoner, 1969: 1, 8, 12 (as *P. laxa*, lab.); Chauthani and Hamm, 1967: 373-376, fig. 2 (lab.).

Spodoptera frugiperda (J. E. Smith)

Chauthani and Hamm, 1967: 373-376, fig. 2 (lab.).

STAGE PARASITIZED: Larva.

IMMATURES: Larva — Chauthani and Hamm, 1967: (fig. 2).

BIOLOGY: Bryan, D. E., Jackson and Stoner, 1969: (1, 8, 12);

Chauthani and Hamm, 1967: (373-376).

Palpexorista Townsend (Goniinae: Exoristini)

Palpexorista coccyx (Aldrich and Webber)

Most records as *Phorocera longiuscula* Walker.

HOST: LEIPDOPTERA

Limacodidae

Phobetron pithecium (J. E. Smith)

Schaffner, 1959: 6, 52, [N.E. U.S.]; Schaffner and Griswold, 1934: 104, 114, Conn.

Sibine stimulea (Clemens)

Schaffner, 1959: 4, 52, [N.E. U.S.]; Aldrich and Webber, 1924: 65, Ind. (as *Phorocera coccyx*).

Notodontidae

Datana ministra (Drury)

Fattig, 1949: 27, Ga.

STAGE PARASITIZED: Larva.

BIOLOGY: Schaffner, 1959: (4, 6, 52); Schaffner and Griswold, 1934: (104, 114).

Palpozenillia Townsend (Goniinae: Eryciini)

Palpozenillia palpalis (Aldrich)

HOST: LEPIDOPTERA

Castniidae

Castnia licoides Boisduval

Box, 1952: 485-490, fig. 2 (non N. Amer.).

Pyralidae

Diatraea considerata Heinrich

Vide Box, 1952: 487-488, Mexico (Nay.) (Van Zwaluwenburg, 1926); Van Zwaluwenburg, 1926: 667, Mexico (Nay.) (as *Phorostoma* sp. ex *Diatraea* in sugar cane).

STAGE PARASITIZED: Larva.

IMMATURES: Egg — Box, 1952: (descr., fig. 2).

First instar larva — Box, 1952: (fig. 2).

Third instar larva — Box, 1952: (fig. 2).

Pupa — Box, 1952: (fig. 2).

BIOLOGY: Box, 1952: (485-490).

Panzeria ampelus, see *Mericia ampelus*

Panzeria penitalis, see *Pyraustomyia penitalis*

Panzeria radicum, see *Mericia ampelus*

Parachaeta Coquillett (Goniinae: Harrisini)

Parachaeta "bicolor," see *Parachaeta* spp.

Parachaeta bicolor, see *Parachaeta* spp.

Parachaeta fusca Townsend

HOST: LEPIDOPTERA

Arctiidae

Isia isabella (J. E. Smith)

Schaffner and Griswold, 1934: 36-37, 122, Conn., N.H., N.J.

STAGE PARASITIZED: Larva.

BIOLOGY: Schaffner and Griswold, 1934: (36-37, 122).

Parachaeta spp.

All records as *Parachaeta bicolor* (Macquart).

HOST: LEPIDOPTERA

Arctiidae

Isia isabella (J. E. Smith)

Brooks, 1947: 172-173, fig. 4 on pl. 14.

HOST: RODENTIA (MAMMALIA)

Unidentified Rodentia

Mouse (Mice)

Wolcott, 1951a: 481, P.R. (as *P. "bicolor"* ex dead mouse).

Mouse (mice) probably

Wolcott, 1924: 223, P.R. (ex possibly from dead mouse).

IMMATURES: Egg — Brooks, 1947: (descr., fig. 4 on pl. 14).

Paradejeania Brauer and Bergenstamm (Tachininae: Dejeaniini)

Paradejeania rutilioides, see *Paradejeania rutilioides nigrescens*

Paradejeania rutilioides nigrescens Arnaud

All records as *Paradejeania rutilioides* (Jaennicke).

HOST: LEPIDOPTERA

Unidentified Lepidoptera

Miscellaneous

Essig, 1915: 330, Calif. (ex caterpillars of various species);

Essig, 1913: 261, Calif. (ex caterpillars).

STAGE PARASITIZED: Larva.

IMMATURES: First instar larva — Arnaud, 1951: (descr., fig. 4).

Paradexodes epilachnae, see *Aplomyiopsis epilachnae*

Paradidyma Brauer and Bergenstamm (Dexiinae: Eriothrixini)

Paradidyma singularis (Townsend)

HOST: LEPIDOPTERA

Unidentified Lepidoptera

Burrowing webworm(-s)

Aldrich, 1915: 82, Ill.

Paradionaea atra, see *Leucostoma simplex*

Parafischeria Townsend (Tachininae: Aphriini)

Parafischeria eucerata (Bigot)

HOST: LEPIDOPTERA

Aegeriidae

Synanthedon bibionipennis (Boisduval)

Wilcox, Gray and Mote, 1932: 17, Oreg. (as *P. venatoris* (Coquillett)).

STAGE PARASITIZED: Pupa.

Parafischeria venatoris, see *Parafischeria eucerata*

Parafrontina apicalis probably, see *Lespesia archippivora* probably

Paragaedia hedemanni, see *Choeteprosopa hedemanni*

Paralispe Brauer and Bergenstamm (Goniinae: Blondeliini)

Paralispe infernalis (Townsend)Most records as *Stomatolydella infernalis*.

HOST: COLEOPTERA

Chrysomelidae

Crioceris asparagi (Linnaeus)Puttler, 1966: 475, Md., S.C. (as *Paralispe*); Watts, 1939: 566, S.C.; Watts, 1938: 192, S.C.*Lema nigrovittata* (Guérin-Ménéville)Terrazas Loyola, 1961: 35-36, Mexico (Mex.) (as *S. "infernalis"*).*Lema sexnotata* ChevrolatPuttler, 1966: 475, Va. (as *Paralispe*).*Lema trilineata californica* Schaeffer

Krauss, 1947: 4, Calif.

Lema trilineata trilineata (Olivier)Aldrich, 1925: 113, D.C. (as *Paralispe* ex *L. trilineata*).*Lema* sp.Puttler, 1966: 475-476, Del. (as *Paralispe*).

HOST: LEPIDOPTERA

Tortricidae

Archips argyrospilus (Walker)All records ex *A. argyrospila*.Newcomer, 1958: 88, Wash.; Johansen, 1957: 74, Wash. (as *Didyma inconspicua* Wulp).

STAGE PARASITIZED: Larva (Coleoptera).

BIOLOGY: Puttler, 1966: (475-476); Terrazas Loyola, 1961: (35-36).

Parameigenia albifacies, see *Oswaldia albifacies*Paramuscopteryx genalis, see *Eulasiona genalis*Paraphasia nigra, see *Phasia nigra*

Paraphoranthia Townsend (Phasiinae: Phasiini)

Paraphoranthia pollinosa Brooks

HOST: HEMIPTERA (HETEROPTERA)

Pentatomidae

Chlorochroa ligata (Say)Brooks, 1945a: 660, Tex. (ex "*Chlorachroa*").Paraphorocera myoidaea, see *Lydella radialis*Paraplagia spinosula, see *Athrycia cinerea*

Paraplagia "spinulosa," see *Athrycia cinerea*

Paraprosena apicalis, see *Opsotheresia bigelowi*

Parasetigena Brauer and Bergenstamm (Goniinae: Exoristini)

Parasetigena agilis (Robineau-Desvoidy)

HOST: LEPIDOPTERA

Lasiocampidae

Malacosoma americanum (Fabricius)

Burgess and Crossman, 1929: 108 (as *Phorocera*, lab.).

Lymantriidae

Porthetria dispar (Linnaeus)

Dowden, 1962: 31-32, 35, 62-63, New England (as *P. silvestris* (Robineau-Desvoidy)); Dowden, 1961: 875, Mass. (as *P. silvestris*); Clausen, 1956: 6-7, 63, 118-119 (as *P. silvestris*); Friend, 1945: 621, Conn. (as *Phorocera*); Brown and Sheals, 1944: 400 (as *P. silvestris*); O'Kane and Conklin, 1942: 53, N.H. (as *P. silvestris* ex gypsy moth); Brown, 1940: 585, Mass. (as *P. silvestris* ex gypsy moth); Dowden, 1935: 509 (as *Phorocera*, lab.); Webber, 1932: 205-206 (as *Phorocera*); Burgess and Crossman, 1929: 104-111, 142, fig. 45, Mass. (as *Phorocera*); Howard and Fiske, 1911: 229-231, 308 (as *P. segregata* (Rondani)); Thompson, 1910: 284 (as *P. segregata* ex "gypsy" moth, lab.).

Stilpnotia salicis (Linnaeus)

Burgess and Crossman, 1929: 108 (as *Phorocera*, lab.).

STAGE PARASITIZED: Larva, pupa.

IMMATURES: Egg — Burgess and Crossman, 1929: (descr., fig. 45).

BIOLOGY: Clausen, 1956: (6-7, 63, 118-119); Dowden, 1935: (509); Webber, 1932: (205-206); Burgess and Crossman, 1929: (104-111, 142); Howard and Fiske, 1911: (229-231, 308); Thompson, 1910: (284).

Parasetigena segregata, see *Parasetigena agilis*

Parasetigena silvestris, see *Parasetigena agilis*

Paratheresia Townsend (Proseninae: Theresiini)

Paratheresia claripalpis (Wulp)

HOST: LEPIDOPTERA

Pyralidae

Chilo plejadellus Zincken

Jaynes, 1939: 254 (as *Theresia*, lab.).

Diatraea considerata Heinrich

Box, 1953: 29, 87, N.W. Mexico; Box, 1952: 487-488, Mexico (Nay.).

Diatraea grandiosella Dyar

Box, 1953: 30, 87, Mexico (Sin.).

Diatraea magnifactella Dyar

Box, 1953: 30, 87, Cent. and W. Mexico.

Diatraea saccharalis (Fabricius)

Anon., 1960: 211, Barbados; Anon., 1960a: 58-59, Barbados; Gibson, W. W., and Carrillo S., 1959: 183, Mexico (Camp.); Clausen, 1956: 4-5, 44-45, 119-120, Fla.; Simmonds, 1955: 199-200, Dominica, Guadeloupe; Anon., 1954: 99, Costa Rica (as "*Paratheresis*" ex "*Ditraea*"); Mathes, Ingram, Questel, Thames and Dugas, 1954: 562, Fla., La.; Box, 1953: 30, 87, Guatemala, Panama; Simmonds, 1953: 126 (non N. Amer., lab.); Van Zwaluwenburg, 1953: 280-282, Mexico (as *Theresia*); Ingram, Mathes, Wilson and Hayslip, 1951: 356, Fla., La. (as *Theresia*); Van Zwaluwenburg, 1951: 374, Mexico (as *Theresia*); Wolcott, 1951a: 475, P.R.; Bartlett, 1943: 16-17 (as *P. diatraeae* (Brèthes), lab.); Bartlett, 1942: 19 (as *P. diatraeae*, lab.); Lee, 1942: 65, 67-68 (as *Theresia*, lab.); Ingram and Bynum, 1941: 9, Fla., La. (as *Theresia*); Lee, 1940: 102-104, 106 (as *Theresia*, lab.); Holloway, 1939: 261, Fla., La. (as *Theresia*); Ingram, Jaynes and Lobdell, 1939: 90-91, Fla. (as *Theresia*); Jaynes, 1939: 254-256, Fla., La. (as *Theresia*); Scaramuzza, 1939: 589-595 (as *Theresia*, lab.); Jaynes, 1938: 93-95, Fla., La. (as *Theresia*); Lobdell, 1937: 146, Fla. (as *Theresia* "*clarapalpis*"); Scaramuzza, 1934: 13-17 (lab.); Jaynes, 1933: 3-10, figs. 2-4 (non N. Amer.); Aldrich, 1929: 33, Honduras, Mexico (as *P. signifera* Townsend); Holloway, Haley, Loftin and Heinrich, 1928: 44, Mexico (V.C.) (as "*Parathesia*" *signifera*); Russo, 1927a: 138, Cent. Amer., Mexico (as *P. signifera*).

Diatraea tabernella Dyar

Box, 1953: 31, 87, Costa Rica, Panama.

Diatraea spp.

Gibson, W. W., and Carrillo S., 1959: 183, Mexico (V.C.);

Box, 1953: 31, 87, Mexico (V.C.); Simmonds, 1948: 68, U.S.; Bartlett, 1941: 8, fig. 6B (as *Theresia*).

Diatraea spp. probably

Painter, 1955: 41-42, Guatemala.

Ostrinia nubilalis (Hübner)

Baker, W. A., Bradley and Clark, 1949: 165 (ex *Pyrausta*, unsuccessful lab. rearing).

Zeadiatraea lineolata (Walker)

Richardson, 1959: 151, Mexico (V.C.); Scaramuzza, 1939: 591 (as *Theresia* ex *Diatraea*, unsuccessful lab. rearing).

Zeadiatraea lineolata (Walker) probably

Russo, 1927a: 138, Cent. Amer.; Mexico (as *P. signifera* ex *Diatraea*).

STAGE PARASITIZED: Larva, pupa.

IMMATURES: First instar larva — Jaynes, 1933: (descr., fig. 2A-B).

Second instar larva — Jaynes, 1933: (descr., fig. 3).

Third instar larva — Jaynes, 1933: (descr., fig. 2C-E).

Pupa — Bartlett, 1941: (fig. 6B); Jaynes, 1933: (descr., fig. 4).

BIOLOGY: Clausen, 1956: (4-5, 44-45, 119-120); Simmonds, 1953: (126); Scaramuzza, 1939: (591); Scaramuzza, 1934: (13-17); Jaynes, 1933 (3-10).

Paratheresia diatraeae, see *Paratheresia claripalpis*

Paratheresia signifera, see *Paratheresia claripalpis*

Paratheresia spp.

HOST: LEPIDOPTERA

Pyralidae

Diatraea saccharalis (Fabricius)

Gibson, W. W., and Carrillo S., 1959: 183, Mexico (Camp., V.C.).

Diatraea spp. (including *D. impersonatella* (Walker) and/or *D. saccharalis* (Fabricius))

Fretay, 1960: 145, Guadeloupe (ex "*Diatrea*" (*saccharalis* or *impersonatella*)); Fretay, 1960a: 151, Guadeloupe.

Diatraea spp.

Gibson, W. W., and Carrillo S., 1959: 183, Mexico (V.C.).

Zeadiatraea lineolata (Walker)

Wellhausen, 1957: 268-269, Mexico (V.C.).

STAGE PARASITIZED: Larva, pupa.

"Paratheresis" *claripalpis*, see *Paratheresia claripalpis*

"Parathesia" *signifera*, see *Paratheresia claripalpis*

Parexorista cheloniae, see *Carcelia laxifrons*

Parexorista futilis, see *Euexorista futilis*

Parexorista sp., see *Carcelia* spp.

Patelloa Townsend (Goniinae: Eryciini)

Patelloa fuscimacula (Aldrich and Webber)

All records as *Phorocera fuscimacula*.

HOST: LEPIDOPTERA

Lymantriidae

Hemerocampa vetusta (Boisduval)

Atkins, 1958: 764, Calif.; Essig, 1926: 582, Calif.; Aldrich and Webber, 1924: 74, Calif.

Saturniidae

Antheraea polyphemus (Cramer)

Most records ex *Telea polyphemus*.

Collins, M. M., and Weast, 1961: 94 (ex *Antheraea*); Essig, 1926: 582, Calif.; Aldrich and Webber, 1924: 74, Calif.

STAGE PARASITIZED: Larva.

Patelloa leucaniae (Coquillett)

Most records as *Phorocera leucaniae*.

HOST: LEPIDOPTERA

Lymantriidae

Euproctis chrysorrhoea (Linnaeus)

*Aldrich and Webber, 1924: 45, 68, N.B. (ex *E. "chrysorrhoea,"* Tothill, 1913a); Tothill, 1913a: 58-59, N.B.

Hemerocampa leucostigma (J. E. Smith)

Schaffner and Griswold, 1934: 85, 114.

Noctuidae

Alabama argillacea (Hübner)

Allen, 1929: 686, Miss. (ex *A. "argillicae"*).

Pseudaletia unipuncta (Haworth)

Most records ex *Leucania unipuncta*.

*Aldrich and Webber, 1924: 45, 67 (ex *Cirphis*, Coquillett, 1897); Johnson, 1910: 780, N.J.; Coquillett, 1897: 19, 26, Tenn.

Armyworm(-s)

Johnson, 1925: 198, Conn., Mass.

Notodontidae

Datana ministra (Drury)

Patton, 1958: 79, Fla. (as *Patelloa*, incomplete rearing — pupa); Schaffner and Griswold, 1934: 74, 114.

Heterocampa guttivitta (Walker)

Raizenne, 1952: 88, Ont. (as *Patelloa*).

Psychidae

Thyridopteryx ephemeraeformis (Haworth)

Fattig, 1949: 26; Brimley, 1938: 363, N.C. (ex evergreen bagworm).

Pyralidae

Loxostege similalis (Guenée)

Aldrich and Webber, 1924: 45, 68; Coquillett, 1897: 19, 26, Ark.

Unidentified Lepidoptera

Miscellaneous

Rowe, 1933: 124, Ill. (ex cocoon).

STAGE PARASITIZED: Larva, pupa.

BIOLOGY: Schaffner and Griswold, 1934: (114); Aldrich and Webber, 1924: (45, 67-68).

Patelloa leucaniae (Coquillett) probably

HOST: LEPIDOPTERA

Lymantriidae

Euproctis chrysorrhoea (Linnaeus)

Howard and Fiske, 1911: 93 (as ? *Phorocera*, incomplete rearing — pupa).

Patelloa sp. near *leucaniae* (Coquillett)

HOST: COLEOPTERA

Chrysomelidae

Lema trilineata trilineata (Olivier)

Fattig, 1949: 26-27 (as *Phorocera* ex *L. trilineata*).

Patelloa pachypyga (Aldrich and Webber)Most records as *Phorocera pachypyga*.

HOST: LEPIDOPTERA

Lasiocampidae

Malacosoma disstria HübnerSippell, 1961: 433, 435-436, fig. 7, Ont. (as *Patelloa*); Bess, 1936: 605; Schaffner and Griswold, 1934: 89, 115, N.J.*Malacosoma* spp.Ross, 1953: 19-20, 23, fig. 12, B.C. (as *Patelloa*).

Lymantriidae

Euproctis chrysorrhoea (Linnaeus)Vide Aldrich and Webber, 1924: 71, New England (ex *E. "chrysorrhoea,"* Tothill, 1913); Tothill, 1913: 73, N.B. (as *Phorocera leucaniae* Coquillett).*Nygmia phaeorrhoea* (Donovan)

Burgess and Crossman, 1929: 143.

STAGE PARASITIZED: Larva.

IMMATURES: Pupa — Sippell, 1961: (descr., fig. 7); Ross, 1953: (descr., fig. 12).

BIOLOGY: Schaffner and Griswold, 1934: (89, 115).

Patelloa pachypyga (Aldrich and Webber) probably

HOST: LEPIDOPTERA

Nymphalidae

Nymphalis antiopa (Linnaeus)Thompson, 1953b: 399 (as *Phorocera*).

STAGE PARASITIZED: Larva.

Patelloa spp.

HOST: LEPIDOPTERA

Geometridae

Drepanulatrix spp. (including *D. baueraria* Sperry and/or *D. monicaria* (Guenée))

Rindge, 1949: 285, 288, Calif.

Noctuidae

Crocigrapha normani (Grote)

Raizenne, 1952: 50, Que.

STAGE PARASITIZED: Pupa.

Pelatachina Meade (Dexiinae: Macquartiini)

Pelatachina pellucida Coquillett

HOST: LEPIDOPTERA

Nymphalidae

Nymphalis antiopa (Linnaeus)

Raizenne, 1952: 4, Ont. (as *Eohyria*); Dowden, 1935: 509 (ex *Hamadryas*); Schaffner and Griswold, 1934: 11, 108 (ex *Hamadryas*); Webber and Schaffner, 1926: 14-15, 29-30, New England (ex *Euvanessa*); Johnson, 1925: 189, Mass. (ex *Euvanessa*); Smith, H. E., 1917: 57, Mass. (ex *Euvanessa*).

Nymphalis milberti (Godart)

Schaffner and Griswold, 1934: 11, 108 (ex *Hamadryas*).

Vanessa atalanta (Linnaeus)

Schaffner and Griswold, 1934: 12, 108 (ex *Cynthia*).

STAGE PARASITIZED: Larva.

IMMATURES: Pupa — Smith, H. E., 1917: (descr.).

BIOLOGY: Dowden, 1935: (509); Schaffner and Griswold, 1934: (108); Webber and Schaffner, 1926: (14-15, 29-30); Smith, H. E., 1917: (57).

Pelecotheca macra, see "*Phorocera*" spp.

Peleteria Robineau-Desvoidy (Tachininae: Tachinini)

Peleteria aenea (Staeger)

HOST: LEPIDOPTERA

Lymantriidae

Byrdia groenlandica Wocke

Henriksen and Lundbeck, 1917: 621, Greenland (ex *Dasychira*, probable host).

Peleteria obsoleta Curran

HOST: LEPIDOPTERA

Noctuidae

Unidentified Noctuidae

Guppy, R., 1950: 4, B.C. (ex unknown phalaenid).

STAGE PARASITIZED: Larva.

Peleteria robusta, see *Peleteria texensis*

Peleteria "tessellata (Fabricius)"

HOST: LEPIDOPTERA

Noctuidae

Euxoa auxiliaris (Grote)

All records ex *Chorizagrotis auxiliaris*.

Crumb, 1929: 40, 99; Essig, 1926: 585, Mont.

Peleteria "tessellata (Fabricius)" probably

HOST: LEPIDOPTERA

Noctuidae

Euxoa auxiliaris (Grote)

Cooley, 1908: 146, Mont. (ex *Chorizagrotis agrestis* (Grote)).

STAGE PARASITIZED: Larva.

Peleteria texensis Curran

Most records as *Peleteria robusta* (Wiedemann).

HOST: LEPIDOPTERA

Lymantriidae

Hemerocampa vetusta (Boisduval)

Essig, 1926: 585, Calif.; Essig, 1915: 330, Calif.; Essig, 1913: 260, Calif.

Noctuidae

Agrotis orthogonia Morrison

All records ex *Porosagrotis orthogonia*.

Essig, 1926: 585, Alta.; Cook, W. C., 1923: 327, Mont. (as "*Pleteria*"); Parker, J. R., Strand and Seamans, 1921: 313, Mont.; Strickland, 1921: 98, Alta.

Euxoa auxiliaris (Grote)

*Vide Crumb, 1929: 40, 99 (ex *Chorizagrotis*, Aldrich, 1915); Aldrich, 1915: 84, Kans. (ex *C. agrestis* (Grote)).

Euxoa ridingsiana (Grote)

Aldrich, 1915: 84, Nev. (ex *Paragrotis*).

Heliophila spp.

Vickery, 1915: 391, Tex.

Leucania latiuscula Herrich-Schaeffer

All records ex *Cirphis latiuscula*.

Vickery, 1926: 1109-1111, 1116, fig. 3D, Tex.; Vickery, 1925: 140, Tex.

Leucania microgonia (Hampson)

Box, 1954: 551, Barbados (as "*Peletieria*" ex *Cirphis*); Tucker, R. W. E., 1952: 351, Barbados.

Peridroma saucia (Hübner)

Anon., 1969: 59, Okla. (as *P. texensis*).

Pseudaletia unipuncta (Haworth)

Most records ex *Cirphis unipuncta*.

Vickery, 1926: 1116, Tex.; Vickery, 1925: 140, Tex.;
Reinhard, 1922: 73, Tex.; Aldrich, 1915: 83-84, Tex.
(ex *Leucania*).

Armyworm(-s)

Reinhard, 1919: 283, Tex.

STAGE PARASITIZED: Larva, pupa.

IMMATURES: Pupa — Vickery, 1926: (fig. 3D); Greene, 1922:
(descr., fig. 71 on pl. 15).

BIOLOGY: Vickery, 1926: (1109-1111, 1116).

Peleteria thomsoni (Williston)

HOST: LEPIDOPTERA

Lepidoptera probably

Miscellaneous

Townsend, 1895: 72, Colo. (as *Echinomyia* ex unknown
larva on clover).

STAGE PARASITIZED: Larva.

Peleteria spp.

HOST: LEPIDOPTERA

Noctuidae

Agrotis orthogonia Morrison

Cook, W. C., 1930: 31, Mont. (ex *Porosagrotis*).

Pseudaletia unipuncta (Haworth)

Burrell, 1967: 114, La.

"*Peletieria*" *robusta*, see *Peleteria texensis*

Periscepsia Gistel (Dexiinae: Eriothrixini)

Periscepsia cinerosa (Coquillett)

All records as *Phorichaeta cinerosa*.

HOST: LEPIDOPTERA

Noctuidae

Euxoa auxiliaris (Grote)

*Crumb, 1929: 40, 99 (ex *Chorizagrotis*, Snow, 1925);
Snow, 1925: 606, 609, Utah.

STAGE PARASITIZED: Larva.

IMMATURES: Pupa — Snow, 1925: (descr.); Greene, 1922:
(descr., fig. 37 on pl. 8).

BIOLOGY: Snow, 1925: (606, 609).

Periscepsia helymus (Walker)

HOST: LEPIDOPTERA

Noctuidae

Actebia fennica (Tauscher)

Most records ex *Agrotis fennica*.

Phipps, 1927: 212 (as *Phorichaeta sequax* (Williston));

Gibson, 1915: 14, Mich. (as *Wagneria sequax* ex *Noctua*); Cook, 1884: 5-6, Mich. (as *Scopolia sequax*);

Cook, 1884a: 424-425, Mich. (as *Scopolia sequax*).

Agrotis sp.

Coquillett, 1897: 19, 23 (as *Phorichaeta sequax*).

Amathes smithii (Snellen)

Wood and Neilson, 1957: 503, N.B. (as *Wagneria* ex *Graphiphora*).

Autographa californica (Speyer)

Clancy, 1969: 1079, Calif.

Cirphis sp.

Reinhard, 1955: 57, Wash. (as *Wagneria*).

Elaphria nucicolora (Guenée)

Vickery, 1925: 141, Tex. (as *Metachaeta* ex *Monodes*).

Epiglaea apiata (Grote)

Franklin, 1950: 57 (as *Phorichaeta sequax*).

Eupsilia devia (Grote)

Schaffner and Griswold, 1934: 51, 124 (as *Wagneria sequax* ex *Conistra*).

Euxoa auxiliaris (Grote)

Strickland, 1938: 218, Alta. (as *Wagneria sequax* ex army cutworm); *Crumb, 1929: 40, 99 (as *Phorichaeta sequax* ex *Chorizagrotis*, Strickland, 1916); Strickland, 1916: 22, Alta. (as *Phorichaeta sequax*).

Euxoa ochrogaster (Guenée)

King, K. M., and Atkinson, 1928: 170, 176, Sask. (as *Phorichaeta sequax*).

Faronta diffusa (Walker)

Webster, R. L., 1911: 181, Iowa (as *Metachaeta* ex *Meliana albilinea* (Hübner)).

Leucania commoides Guenée

Tothill, 1913: 73, Ont. (as *Phorichaeta sequax* ex *Heliophila*).

Lithophane innominata (Smith)

Reinhard, 1955: 58, Maine (as *Wagneria*).

Lithophane spp. (including *L. disposita* Morrison, *L. innominata* (Smith) and/or *L. petulca* Grote)

All records ex *Graptolitha* spp. (including *G. disposita*, *G. innominata* and/or *G. petulca*).

Schaffner and Griswold, 1934: 48, 124 (as *Wagneria*);

Schaffner and Griswold, 1934: 48, 124 (as *Wagneria sequax*).

Peridroma saucia (Hübner)

Reinhard, 1955: 57, Calif. (as *Wagneria*).

Polia adjuncta (Boisduval)

Schaffner and Griswold, 1934: 44-45, 124 (as *Wagneria sequax*).

Pseudaletia unipuncta (Haworth)

Guppy, J. C., 1967: 97, 99, 102, Ont.; Baker, 1915: 86, Ont. (as *Wagneria sequax* ex *Heliophila*); Gahan, 1915: 24,

Md. (as *Metachaeta* ex *Leucania*); Gibson, 1915: 14, Ont. (as *Wagneria sequax* ex *Cirphis*).

Trichoplusia ni (Hübner)

Clancy, 1969: 1079, Calif.

Armyworm (-s)

Aldrich, 1915: 83, Md. (as *Metachaeta*).

Cutworm (-s)

Lintner, 1893: 238, N.Y. (as *Scopolia sequax*).

Unidentified Noctuidae

Schaffner and Griswold, 1934: 72, 124 (as *Wagneria*);

Schaffner and Griswold, 1934: 72, 124 (as *Wagneria sequax*); Coquillett, 1897: 19, 26, Calif. (as *Phorichaeta sequax*).

STAGE PARASITIZED: Larva.

IMMATURES: Pupa — Greene, 1922: (descr., fig. 44 on pl. 10 and fig. 76 on pl. 16).

BIOLOGY: Guppy, J. C., 1967: (97, 99, 102); Schaffner and Griswold, 1934: (124).

Periscepsia laevigata (Wulp)

Most records as *Wagneria laevigata*.

HOST: LEPIDOPTERA

Noctuidae

Elaphria nucicolora (Guenée)

Reinhard, 1955: 54.

Euxoa auxiliaris (Grote)

Reinhard, 1955: 54, Guatemala to Canada.

Lascoria ambigualis (Walker)

Reinhard, 1955: 54.

Lithophane sp.

Reinhard, 1955: 54 (ex *Graptolitha*).

Peridroma saucia (Hübner)

Anon., 1969: 59, Okla. (as *Periscepsia*).

Pseudaletia unipuncta (Haworth)

Guppy, J. C., 1967: 97, 99, 102, Ont. (as *Periscepsia*);

Breeland, 1958: 323, 333, Tenn.; Reinhard, 1955: 54,

Guatemala to Canada (ex *Cirphis*).

STAGE PARASITIZED: Larva.

Periscepsia rohweri (Townsend)

All records as *Wagneria rohweri*.

HOST: LEPIDOPTERA

Noctuidae

Agrotis orthogonia Morrison

Reinhard, 1955: 59, Colo. or Idaho or Wash.; Cook, W. C.,

1930: 31, Mont. (ex *Porosagrotis*).

Euxoa pallipennis (Smith)

Cook, W. C., 1930: 31, Mont.

Euxoa tristicula (Morrison)

Cook, W. C., 1930: 31, Mont.

Spaelotis clandestina (Harris)

Cook, W. C., 1930: 31, Mont. (ex *Agrotis unicolor* (Walker)).

Periscepsia spp.

Most records as *Phorichaeta* sp.

HOST: LEPIDOPTERA

Noctuidae

Agrotis sp. probably

Brauer and Bergenstamm, 1895: 569, 600.

Glaea sp.

Brauer and Bergenstamm, 1895: 569, 593.

Unidentified Lepidoptera

Leaf roller(-s)

Townsend, 1895: 78, Colo. (as *Scopolia* ex leaf-roller on choke-cherry).

Peteina stylata, see *Petinarctia stylata*

"Petina" stylata, see *Petinarctia stylata*

Petinarctia Villeneuve (Tachininae: Germariini)
 Petinarctia stylata (Brauer and Bergenstamm)
 HOST: LEPIDOPTERA

Lymantriidae

Byrdia groenlandica Wocke

Henriksen and Lundbeck, 1917: 622, E. Greenland (as
 "Petina" ex *Dasychira*).

STAGE PARASITIZED: Larva.

Phasia Latreille (Phasiinae: Phasiini)

Phasia nigra (Brooks)

HOST: HEMIPTERA (HETEROPTERA)

Pentatomidae

Eurygaster spp.

Vojdani, 1961: 576, Calif. (as *Paraphasia*).

Phasia occidentis, see "*Phorantha occidentis*"

Phasia nearly allied to *Trichopoda pennipes*, see *Trichopoda* spp.

Phasioclista metallica, see *Myiophasia* "aenea" or *Myiophasia metallica*

Phasiops Coquillett (Proseninae: Trixini)

Phasiops flavus Coquillett

HOST: DIPTERA

Tabanidae

Tabanus petiolatus Hine

Patton, 1958: 79, Fla.

Tabanus petiolatus Hine probably

Jones and Anthony, 1964: 56-57, Fla.

Tabanus trimaculatus Palisot de Beauvois

Jones, T. H., and Bradley, 1923: 312.

Tabanus trimaculatus Palisot de Beauvois probably

Patton, 1958: 79, La.

Tabanus sp.

Jones, 1953: 1109, Fla.

HOST: LEPIDOPTERA

Pyralidae

Crambus spp.

Sabrosky and Arnaud, 1965: 980 (as *Phasiops* [sp.]).

STAGE PARASITIZED: Larva (Diptera, Lepidoptera).

Phasiops flavus Coquillett probably

HOST: DIPTERA

Tabanidae

Tabanus fumipennis Wiedemann

Patton, 1958: 79, Fla.

Phasiops sp., see *Phasiops flavus*

Phasmophaga Townsend (Goniinae: Hyperecteinini)

Phasmophaga antennalis Townsend

HOST: ORTHOPTERA

Phasmatidae

Diapheromera femorata (Say)

Severin, H. H. P., Severin and Hartung, 1915: 135 (lab.);

Townsend, 1909: 244, Wis. (ex "*Diapheromerata*").

IMMATURES: Pupa — Greene, 1922: (descr., fig. 13 on pl. 3).

BIOLOGY: Severin, H. H. P., Severin and Hartung, 1915: (135);

Townsend, 1909: (244).

Phasmophaga antennalis Townsend probably

HOST: ORTHOPTERA

Phasmatidae

Anisomorpha buprestoides (Stoll)

Neff and Eisner, 1960: 102, Fla. (as "*Phasmophora*");

Patton, 1958: 79, Fla.

STAGE PARASITIZED: Adult.

Phasmophaga meridionalis Townsend

HOST: ORTHOPTERA

Phasmatidae

Anisomorpha buprestoides (Stoll)

Patton, 1958: 79, Fla.; Russell, 1912: 117, Fla.; Townsend,

1909: 244, Fla.

STAGE PARASITIZED: Adult.

"*Phasmophora*" *antennalis* probably, see *Phasmophaga antennalis* probably

"*Pherostoma*" sp., see "*Phorostoma*" spp.

Philocalia Reinhard (Goniinae: Siphonini)

Philocalia tenuirostris Reinhard

HOST: LEPIDOPTERA

Tineidae

Amydria effrentella Clemens

Arnaud, 1959a: 186, Ill. (ex A. "effrenatella").

Phorantha Rondani (Phasiinae: Phasiini)

"*Phorantha occidentis* (Walker)"

HOST: HEMIPTERA (HETEROPTERA)

Lygaeidae

Nysius angustatus Uhler

Aldrich, 1915: 81.

Nysius ericae (Schilling)

Milliken and Wadley, 1923: 28-31, Kans. (as *Phasia*).

Miridae

Leptopterna dolabrata (Linnaeus)

All records ex *Miris dolabratus*.

West, 1928: 806, N.Y. (as *Phoranthella*); Leonard, 1916: 236.

STAGE PARASITIZED: Nymph, adult.

IMMATURES: Third instar larva — Milliken and Wadley, 1923: (descr.).

Pupa — Milliken and Wadley, 1923: (descr.); Greene, 1922: (descr., fig. 46 on pl. 10).

BIOLOGY: Milliken and Wadley, 1923: (28-31).

Phoranthella occidentis, see "*Phorantha occidentis*"

Phorichaeta cinerosa, see *Periscepsia cinerosa*

Phorichaeta sequax, see *Periscepsia helymus*

Phorichaeta sp., see *Periscepsia* spp.

Phorocera Robineau-Desvoidy (Goniinae: Exoristini)

Phorocera agilis, see *Parasetigena agilis*

Phorocera claripennis, see *Euphorocera claripennis*

Phorocera coccyx, see *Palpexorista coccyx*

"*Phorocera*" *comstocki* Williston

HOST: HYMENOPTERA

Diprionidae

Diprionid spp. (including *Diprion* spp. and/or *Neodiprion* spp.)

All records ex *Lophyrus* spp.

*Aldrich and Webber, 1924: 79, Ill. (Townsend, 1893);

Brauer and Bergenstamm, 1895: 569, 598; Townsend, 1893: 467, Ill.

HOST: LEPIDOPTERA

Cossidae

Cossula magnifica (Strecker)

Fattig, 1949: 26; Brimley, 1938: 363, N.C.; Leiby, 1925: 49, N.C.; *Aldrich and Webber, 1924: 79, N.C. (Brimley, 1922); Brimley, 1922: 25, N.C.

Megathymidae

Megathymus yuccae (Boisduval and LeConte)

Remington, 1959: 179-181, 183, Ga. (as *Pales*); Aldrich and Webber, 1924: 78, S.C.; Coquillett, 1897: 19, 26, S.C.; *Brauer and Bergenstamm, 1895: 569, 599 (Williston, 1889); Williston, 1889: 1914, 1922.

Pyralidae

Melitara dentata (Grote)

Mann, 1969: 14, 33; Mann, 1969: 14, 35 (ex *M. doddalis* Dyar).

Olycella junctolineella (Hulst)

Hunter, Pratt and Mitchell, 1912: 27, 46, Tex. (ex *Melitara*).

Ostrinia spp. (including *O. obliteralis* (Walker) and/or *O. penitalis* Grote)

*Aldrich and Webber, 1924: 79, Ill. (ex *Pyrausta penitalis*, Townsend, 1893); *Ainslie, G. G., and Cartwright, 1921: 843 (ex *Pyrausta penitalis* or *P. ainsliei* Heinrich, Townsend, 1893); *Brauer and Bergenstamm, 1895: 569, 585, N. Amer. (ex *Botys penitalis*, Townsend, 1893); Townsend, 1893: 467, Ill. (ex "*Botis*" *penitalis*).

STAGE PARASITIZED: Larva (Lepidoptera), pupa (Lepidoptera).

BIOLOGY: Remington, 1959: (179-181, 183).

Phorocera doryphorae, see *Doryphorophaga doryphorae*

Phorocera edwardsii, see *Euphorocera edwardsii*

Phorocera erecta, see *Pseudoperichaeta erecta*

Phorocera sp. near *erecta*, see *Pseudoperichaeta* sp. near *erecta*

Phorocera flavicauda, see *Prophryno parviteres*

Phorocera floridensis, see *Euphorocera floridensis*

Phorocera sp. near *floridensis*, see *Euphorocera* sp. near *floridensis*

Phorocera fuscimacula, see *Patelloa fuscimacula*

"*Phorocera*" *halisidotae* Aldrich and Webber

HOST: LEPIDOPTERA

Arctiidae

Halisidota maculata (Harris)

Essig, 1926: 582, Idaho; Aldrich and Webber, 1924: 85, Idaho (ex "*Halisidota*").

Phorocera hamata, see *Diplostichus lophyri*

Phorocera sp. near *hamata*, see *Diplostichus* sp. near *lophyri*

"*Phorocera*" *incrassata* Smith

HOST: LEPIDOPTERA

Tortricidae

Choristoneura fumiferana (Clemens)

Coppel, 1960: 94, 97, B.C.; Carolin and Coulter, 1959: 552-554, Oreg.; Ross, 1952: 108, fig. 3, Canada; Anon., 1950d: 238, 246, 248, B.C.; Wilkes, 1949: 111, B.C. (ex *Archips*); Wilkes, Coppel and Mathers, 1949: 140, 146, 149, 151, 153, B.C.; Anon., 1948b: 355, 366, 371, B.C.; Dowden, Buchanan and Carolin, 1948: 458 (ex *Archips*); Coppel, 1947: 39, B.C. (ex *Archips*); Wilkes, 1946: 83-86, B.C. (ex *Archips*); Baird, 1945: 126, 133, 137, B.C. (ex *Archips*).

STAGE PARASITIZED: Larva, pupa.

IMMATURES: Pupa — Ross, 1952: (fig. 3).

BIOLOGY: Coppel, 1947: (39).

Phorocera leucaniae, see *Patelloa leucaniae*

Phorocera sp. near *leucaniae*, see *Patelloa* sp. near *leucaniae*

Phorocera longiuscula, see *Palpexorista coccyx*

Phorocera lophyri, see *Diplostichus lophyri*

Phorocera macra, see "*Phorocera*" spp.

Phorocera sp. near *macra*, see "*Phorocera*" spp.

Phorocera marginalis, see *Prophryno marginalis*

Phorocera meracanthae, see *Catagoniopsis meracanthae*

Phorocera "namata," see *Diplostichus lophyri*

Phorocera noctuiformis, see *Nilea noctuiformis*

Phorocera sp. near *noctuiformis*, see *Nilea* sp. near *noctuiformis*

Phorocera omissa, see *Euphorocera omissa*

Phorocera pachypyga, see *Patelloa pachypyga*

"*Phorocera parva* Bigot"

HOST: LEPIDOPTERA

Pyralidae

Loxostege similalis (Guenée)

Coquillett, 1897: 19, 26.

Udea rubigalis (Guenée)

All records ex *Phlyctaenia ferrugalis* Hübner.

Gibson, 1919: 629, Ill., Pa. (ex *P. "ferrugalis"*); Davis, J. J., 1912: 100, Ill.

Tortricidae

Argyrotaenia citrana (Fernald)

All records ex *Tortrix citrana*.

Essig, 1915: 442, Calif.; Essig, 1913: 176, Calif.; Coquillett, 1897: 19, 27, Calif.

STAGE PARASITIZED: Larva.

Phorocera parviteres, see *Prophryno parviteres*

Phorocera promiscua, see *Lespesia archippivora*

Phorocera pumilio, see *Pseudoperichaeta insidiosa*

Phorocera saundersii, see *Madremyia saundersii*

"*Phorocera*" *signata* Aldrich and Webber

HOST: LEPIDOPTERA

Cossidae

Cossula magnifica (Strecker)

Aldrich and Webber, 1924: 87, S.C.

Noctuidae

Papaipema nebris (Guenée)

Aldrich and Webber, 1924: 82, Tenn. (as *P. xanthura* Aldrich and Webber ex *P. nitela* (Guenée)).

Phorocera sp. near *specularis*, see *Catagoniopsis* sp. near *specularis*

Phorocera sternalis, see *Nilea sternalis*

Phorocera subnitens probably, see *Euphorocera subnitens* probably

Phorocera sp. near "tachinomiodes," see *Euphorocera* sp. near *tachinomoides*

Phorocera tachinomoides, see *Euphorocera tachinomoides*

Phorocera tenuiseta, see "*Phorocera*" *victoria*

"*Phorocera*" *texana* Aldrich and Webber

HOST: LEPIDOPTERA

Megathymidae

Agathymus baueri (Stallings and Turner)

Roever, 1964: 116, 119, Ariz.

Agathymus chisosensis (Freeman)

Roever, 1964: 116.

Agathymus evansi (Freeman)

Roever, 1964: 116, Ariz.

Agathymus freemani Stallings and Turner

Roever, 1964: 116, 119, Ariz.

Agathymus neumogeni (Edwards)

Roever, 1964: 116, 119, Ariz. (ex *A. aryxna* (Dyar)).

Pyralidae

Melitara dentata (Grote)

Mann, 1969: 14, 33; Mann, 1969: 14, 35 (ex *M. doddalis* Dyar).

Melitara prodenialis Walker

Mann, 1969: 14, 32, Tex.

Melitara spp.

Mann, 1969: 14, 37, Tex.; Aldrich and Webber, 1924: 79, Tex. (ex *Melitara* in *Opuntia* [sp.]).

Olycella junctolineella (Hulst)

Mann, 1969: 14, 39.

Olycella nephelepasa (Dyar)

Mann, 1969: 14, 40, Mexico (S.L.P.).

STAGE PARASITIZED: Larva, pupa.

Phorocera "torbricis," see *Hemisturmia tortricis*

Phorocera "torticis," see *Hemisturmia tortricis*

Phorocera tortricis, see *Hemisturmia tortricis*

"*Phorocera victoria* Aldrich and Webber"

HOST: LEPIDOPTERA

Arctiidae

Unidentified Arctiidae probably

Aldrich and Webber, 1924: 83, B.C. (as *P. tenuiseta* Aldrich and Webber).

STAGE PARASITIZED: Pupa.

Phorocera xanthura, see "*Phorocera*" *signata*

Phorocera sp. near *Zenillia lobeliae*, see "*Exorista*" sp. near *lobeliae*

"*Phorocera*" spp.

HOST: HYMENOPTERA

Diprionidae

Neodiprion excitans Rohwer

Hetrick, 1959: 161, Fla.

Neodiprion lecontei (Fitch)

*Vide Arnaud, 1963a: 4, 53, Va. (as *P. macra* Wulp, Benjamin, 1955; Middleton, 1921); Benjamin, 1955: 20, U.S. (as *Pelecotheca macra*); Bentley, 1940: 191, Ala.; Greene, 1921: 41-42, D.C. (as *P.* sp. (near *macra*)); Middleton, 1921: 757 (as *Neopales* "maera").

Neodiprion swaini Middleton

Atwood, 1944: 59, Que.

Neodiprion spp.Greene, 1921: 41, Va.; Greene, 1921: 42, Conn. (as *P. sp.*, near *macra*).

Tenthredinidae

Pristiphora erichsonii (Hartig)

Hopping, Leech and Morgan, 1943: 59, B.C.

Unidentified Hymenoptera

Pitch pine sawfly(-flies)

*Procter, 1946: 432, Maine (Procter, 1938); Procter, 1938: 373, Maine.

HOST: LEPIDOPTERA

Arctiidae

Estigmene acrea (Drury)

Bibby, 1961: 327, Ariz. (ex salt-marsh caterpillar).

Lymantriidae

Hemerocampa leucostigma (J. E. Smith)All records ex *Orgyia leucostigma*.

Howard, 1898: 88, D.C.; Howard, 1897a: 17, D.C.

Noctuidae

Trichoplusia ni (Hübner)

Sutherland, 1966: 38-39, N.Y.

Armyworm(-s)

Annand, 1945: 146, Fla.

Nymphalidae

Chlosyne harrisii (Scudder)All records ex *Cinclidia harrisii*.

*Brauer and Bergenstamm, 1895: 569, 587 (Williston, 1889); Williston, 1889: 1913-1914.

Nymphalis antiopa (Linnaeus)Schaffner and Griswold, 1934: 10 (ex *Hamadryas*).

Pieridae

Colias eurytheme Boisduval

Lochhead, 1919: 176 (ex *Eurymus*).

Pyralidae

Diatraea sp.

Gibson, W. W., and Carrillo S., 1959: 183 (ex "*Diarnea*," lab.).

Laniifera cyclades Druce

Mann, 1969: 14, 77, Mexico (ex *Megastes*).

Loxostege similalis (Guenée)

Brauer and Bergenstamm, 1895: 569, 598.

Zygaenidae

Harrisina brillians Barnes and McDunnough

Clausen, 1955: 78, Ariz., Mexico.

Harrisina spp.

Clausen, 1961: 618.

Unidentified Lepidoptera

Hog moth(-s)

Fattig, 1949: 27.

STAGE PARASITIZED: Larva (Hymenoptera, Lepidoptera),
pupa (Hymenoptera, Lepidoptera).

IMMATURES: Pupa — Greene, 1922: (descr., fig. 91 on pl. 19).

BIOLOGY: Bentley, 1940: (191).

Phorostoma Robineau-Desvoidy (Proseninae: Prosenini)

Phorostoma novaeangliae, see *Carinosillus novaeangliae*

Phorostoma tabanivorus, see *Carinosillus tabanivorus*

Phorostoma n. sp., see *Carinosillus tabanivorus*

Phorostoma sp., see *Palpozenillia palpalis*

"*Phorostoma*" spp.

HOST: DIPTERA

Tabanidae

Tabanus trimaculatus Palisot de Beauvois

Philip, 1931: 68, Minn.

HOST: LEPIDOPTERA

Pyralidae

Diatraea saccharalis (Fabricius)

Holloway, Haley, Loftin and Heinrich, 1928: 44, Mexico (Nay.); Russo, 1927a: 138, Mexico (as "*Pherostoma*").

Diatraea spp.

Van Zwaluwenburg, 1951: 374, Mexico (Nay.).

Zeadiatraea lineolata (Walker) probably

Russo, 1927a: 138, Mexico (as "*Pherostoma*" ex *Diatraea*).

STAGE PARASITIZED: Larva (Diptera, Lepidoptera).

Phosococephalops crassicornis, see *Gonia crassicornis*

"*Phosococephalus*" *crassicornis*, see *Gonia crassicornis*

"*Phryx*" *vulgaris*, see *Phryxe vulgaris*

Phrynofrontina Townsend (Goniinae: Blondeliini)

Phrynofrontina discalis (Coquillett)

HOST: LEPIDOPTERA

Lymantriidae

Euproctis chrysorrhoea (Linnaeus)

Howard and Fiske, 1911: 93 (as *Sturmia*).

Phrynofrontina spp.

HOST: LEPIDOPTERA

Olethreutidae

Petrova metallica (Busck)

Stark, 1959: 162, Rocky Mt. region of Canada.

Pyralidae

Dioryctria abietella (Denis and Schiffermüller) probably

Ebel, 1965: 629, Fla.

Dioryctria amatella (Hulst)

Ebel, 1965: 628, Fla.

Dioryctria clarioralis (Walker)

Ebel, 1965: 630, Fla.

STAGE PARASITIZED: Larva.

Phrynolydella eufitchiae, see *Blondelia eufitchiae*

Phrynolydella hyphantriae, see *Blondelia hyphantriae*

Phrynolydella sp., see *Blondelia* spp.

"*Phryxa*" *pecosensis*, see *Phryxe pecosensis*

Phryxe Robineau-Desvoidy (Goniinae: Eryciini)

Phryxe pecosensis (Townsend)

HOST: HYMENOPTERA

Tenthredinidae

Pikonema alaskensis (Rohwer)

Raizenne, 1957: 32, Ont.

HOST: LEPIDOPTERA

Danaidae

Danaus plexippus plexippus (Linnaeus)

Sellers, 1943: 36 (ex *D. plexippus*).

Geometridae

Alsophila pometaria (Harris)

Sellers, 1943: 37.

Cingilia catenaria (Cramer)

Sellers, 1943: 36.

Ennomos subsignarius (Hübner)

Sellers, 1943: 36.

Erannis tiliaria (Harris)

Raizenne, 1952: 142, Ont., Que.; Sellers, 1943: 36.

Lambdina fiscellaria fiscellaria (Guenée)

Most records ex *Lambdina fiscellaria*.

Carroll, 1956: 594, Nfld. (ex *L. fiscellaria fiscellaria*);

Reeks, Forbes and Cuming, 1947: 13, Nfld.; Reeks,

Smith and Forbes, 1946: 13, Nfld. (as "*Phryxa*").

Operophtera brumata (Linnaeus)

Smith, C. C., 1950: 106, N.S. (questionable host).

Hesperiidae

Thymelicus lineola (Ochsenheimer)

Arthur, 1962: 1085, Ont.

Noctuidae

Heptagrotis phyllophora (Grote)

Wood and Neilson, 1957: 505, N.B.

Lithophane antennata (Walker)

Sellers, 1943: 37 (ex *Graptolitha*).

Pseudaletia unipuncta (Haworth)

Pond, 1960: 665, N.B.; Sellers, 1943: 36-37 (ex *Cirphis*).

Noctuid spp. (including *Mamestra configurata* Walker and/or

Peridroma saucia (Hübner))

Anon., 1962: 121, B.C.; Handford, 1962: 19, B.C.
Unidentified Noctuidae
Sellers, 1943: 36.

Pieridae

Pieris rapae (Linnaeus)
Maw and Coppel, 1953: 392-403, figs. 1-16 (lab.).

Pyralidae

Dioryctria reniculella (Grote)
McLeod, J. M., and Daviault, 1963: 316, Que.
Nephoteryx subcaesiella (Clemens)
Schaffner, 1959: 17, 53, Maine (ex *Salebria*).

Saturniidae

Hemileuca spp. (including *H. lucina* H. Edwards or *H. maia*
(Drury))
Sellers, 1943: 37.

Tortricidae

Acleris variana (Fernald)
Miller, C. A., 1966: 603, N.B.; Reeks, Forbes and Cuming,
1947: 12, [maritime prov.].
Aphelia alleniana (Fernald)
Martin, 1958: 46, Ont. (ex *Tortrix*).
Archips argyrospilus (Walker)
LeRoux, 1961: 115, Que.; vide Sellers, 1943: 36-37 (Regan,
1923); Regan, 1923: 53, Mont. (as *Zenillia vulgaris*
(Fallén) ex *Cacoecia*).
Archips cerasivoranus (Fitch)
Sellers, 1943: 36.
Archips infumatanus (Zeller)
Sellers, 1943: 37.
Choristoneura conflictana (Walker)
Schaffner, 1959: 32, 53, Maine (ex *Archips*); Prentice, 1955:
471, Man., Sask.
Choristoneura fumiferana (Clemens)
Blais, 1965: 131-134, Que.; MacDonald and Webb, 1963:
299, N.B.; Miller, 1963: 237-239, N.B.; Blais, 1960:
387-388, 390-392, Que. (ex spruce budworm); Coppel,

1960: 94, 97, B.C.; Carolin and Coulter, 1959: 553, Oreg.; McGugan and Blais, 1959: 765, Ont.; Boyes and Wilkes, 1953: 126, 152; Dowden, Jaynes and Carolin, 1953: 309, Maine; Maw and Coppel, 1953: 392, Nfld., Que.; Anon., 1952b: 178, 184, 186, B.C.; Anon., 1952g: 288, 292, 295, B.C.; Jaynes and Drooz, 1952: 1058, 1060, Maine, N.Y.; Raizenne, 1952: 219, Ont.; Ross, 1952: 108, fig. 5, Canada; Anon., 1950c: 239, 246, 248, B.C.; Daviault, 1950: 42-45, Que.; Dowden and Carolin, 1950: 778-781, N.Y. (ex *Archips*); Anon., 1949: 272, 283, 285, B.C.; Wilkes, Coppel and Mathers, 1949: 140, 146, 149, 151, 153, B.C.; Anon., 1948: 285, 293, 295, B.C.; Dowden, Buchanan and Carolin, 1948: 458-461 (ex *Archips*); Reeks, Forbes and Cuming, 1947: 12, Nfld. (ex *Archips*); Wilkes and Anderson, 1947: 41, 43, Ont., Que. (ex *Archips*); Reeks, 1946: 25, N.B., Nfld. (ex *Archips*); Reeks, Smith and Forbes, 1946: 12, Nfld. (as "*Phryxa*" ex *Archips*).

Choristoneura pinus Freeman

Allen, D.C., Knight, Foltz and Mattson, 1969: 1471-1472, 1474-1475, Mich.; Dixon and Benjamin, 1963: 267-270, Wis.; Kulman and Hodson, 1961: 222, Minn.; Drooz and Benjamin, 1956: 412-413, Mich.; Benjamin and Drooz, 1954: 589-590, Mich.

Choristoneura rosaceana (Harris)

Raizenne, 1952: 214, Ont. (ex *Archips*).

Unidentified Lepidoptera

Miscellaneous

Sellers, 1943: 36.

STAGE PARASITIZED: Larva (Lepidoptera), pupa (Lepidoptera).

IMMATURES: Egg — Maw and Coppel, 1953: (descr., fig. 1).

First instar larva — Maw and Coppel, 1953: (descr., figs. 2-5).

Second instar larva — Maw and Coppel, 1953: (descr., figs. 6-8).

Third instar larva — Maw and Coppel, 1953: (descr., figs. 9-11).

Pupa — Maw and Coppel, 1953: (descr., figs. 12-16); Ross, 1952: (fig. 5).

BIOLOGY: Schaffner, 1959: (17, 32, 53); Maw and Coppel, 1953: (392-403); Sellers, 1943: (37).

Phryxe pecosensis (Townsend) probably

HOST: LEPIDOPTERA

Geometridae

Synchlora spp. (including *S. aerata* (Fabricius) and/or *S. liquoraria albolineata* Packard)

All records ex *Synchlora rubrifrontaria* Packard.

*Vide Sellers, 1943: 37 (ex "*Synchlora*," Schaffner and Griswold, 1934); Schaffner and Griswold, 1934: 91, 113 (as *Zenillia vulgaris* (Fallén)).

Noctuidae

Nephelodes emmedonia (Cramer)

Sellers, 1943: 37.

Pyralidae

Ostrinia spp. (including *O. obliteralis* (Walker) and/or *O. penitalis* Grote)

*Vide Sellers, 1943: 37 (ex *Pyrausta penitalis*, Townsend, 1893); *vide Ainslie, G. G., and Cartwright, 1921: 843 (as *Exorista vulgaris* ex *Pyrausta penitalis* or *P. ainsliei* Heinrich, Townsend, 1893); *Chittenden, 1918: 456, Ill. (as *Exorista vulgaris* ex *Pyrausta penitalis*, Townsend, 1893); *Brauer and Bergenstamm, 1895: 545, 585 (as *Blepharidea hirsuta* (Osten Sacken) ex *Botys penitalis*, Townsend, 1893); Townsend, 1893: 467, Ill. (as *Exorista hirsuta* ex "*Botis*" *penitalis*).

Tortricidae

Choristoneura fumiferana (Clemens)

*Vide Sellers, 1943: 37 (ex *Archips*, Essig, 1926; Johnson, 1925; Tothill, 1913); Essig, 1926: 581 (as *Zenillia vulgaris* ex *Harmologa*); Johnson, 1925: 196, Maine, Mass., N.H. (as *Zenillia vulgaris* ex *Harmologa*); Tothill, 1913: 71, Que. (as *Exorista vulgaris* ex *Tortrix*).

STAGE PARASITIZED: Larva.

Phryxe vulgaris (Fallén)

HOST: LEPIDOPTERA

Danaidae

Danaus plexippus plexippus (Linnaeus)

Schaffner and Griswold, 1934: 8, 113 (as *Zenillia* ex *D.*

menippe (Hübner)); Webber and Schaffner, 1926: 23, New England (as *Zenillia ex Anosia plexippus*).

Geometridae

Alsophila pometaria (Harris)

Schaffner and Griswold, 1934: 90, 113 (as *Zenillia*).

Cingilia catenaria (Cramer)

Schaffner and Griswold, 1934: 98, 113 (as *Zenillia*).

Ennomos subsignarius (Hübner)

Schaffner and Griswold, 1934: 99-100, 113, Mass. (as *Zenillia*).

Erannis tiliaria (Harris)

Schaffner and Griswold, 1934: 97-98, 113 (as *Zenillia*).

Lymantriidae

Porthetria dispar (Linnaeus)

Sellers, 1943: 34.

Noctuidae

Lithophane antennata (Walker)

Schaffner and Griswold, 1934: 49, 113 (as *Zenillia ex Graptolitha*).

Pseudaletia unipuncta (Haworth)

All records ex *Cirphis unipuncta*.

Schaffner and Griswold, 1934: 47, 113 (as *Zenillia*); Essig, 1926: 581 (as *Zenillia*); Gibson, 1915: 14, N.B., N.S.

Trichoplusia ni (Hübner)

Sutherland, 1966: 38-39, N.Y.

Unidentified Noctuidae

Schaffner and Griswold, 1934: 72, 113 (as *Zenillia*).

Pieridae

Pieris rapae (Linnaeus)

Fitzpatrick, 1967: 15, B.C.; Wilkinson, 1966: 1012-1013, B.C.; Anon, 1963b: 19, Ont.; Harcourt, 1963: 65, Ont.; Beirne, 1962: 19, Ont.; Anon., 1961a: 299, B.C.; Pimentel, 1961: 889-891, N.Y. (as "*Phryex*"); Anon., 1957: 172, E. Canada; Hudon, 1957: 546, Que.; Sellers, 1943: 34-35; Schaffner and Griswold, 1934: 7-8, 113 (as *Zenillia ex Ascia*); Essig, 1926: 581 (as *Zenillia*);

Johnson, 1925: 196, Maine, Mass., N.H. (as *Zenillia* ex *Pontia*); Aldrich and Webber, 1924: 27-28, probably N.Y. (as *Zenillia*); Chittenden, 1916: 9 (as *Exorista* ex *Pontia*); Essig, 1915: 458, Calif. (as *Exorista* ex *Pontia*); Brauer and Bergenstamm, 1895: 545, 602 (as *Blepharidea hirsuta* (Osten Sacken)); Williston, 1889: 1914, 1920 (as *Exorista hirsuta*); Osten Sacken, 1887: 164 (as *Tachina hirsuta*).

Pyralidae

Evergestis straminealis (Hübner)

Schaffner, 1959: 8, 53, Mass.; Sellers, 1943: 34.

Loxostege commixtalis (Walker)

Hoerner, 1933: 12 (as *Zenillia*).

Saturniidae

Hemileuca spp. (including *H. lucina* H. Edwards and/or *H. maia* (Drury))

Schaffner and Griswold, 1934: 29, 113 (as *Zenillia*).

Tortricidae

Archips argyrospilus (Walker)

Schaffner, 1959: 30, 53, Mass.

Choristoneura fumiferana (Clemens)

Most records ex *Tortrix fumiferana*.

Schaffner, 1959: 32, 53, Mass. (ex *Choristoneura*); Winn and Beaulieu, 1915: 142, Que. (as *Exorista* ex *T. "fum-ferana"*); Johannsen, 1913: 25, Maine (as *Exorista*).

STAGE PARASITIZED: Larva, pupa.

BIOLOGY: Harcourt, 1963: (65); Schaffner, 1959: (8, 30, 32, 53); Sellers, 1943: (35); Schaffner and Griswold, 1934: (113).

Phryxe sp.

HOST: LEPIDOPTERA

Pyralidae

Loxostege commixtalis (Walker)

Maxson, 1920: 85, 144, [Colo.].

STAGE PARASITIZED: Larva.

Phyllomya sp. near *aenea*, see *Elfia* sp. near *aenea*
 Phyllomyia columbia, see *Erynnia tortricis*
 Phyllophilopsis Townsend (Dexiinae: Urodexiini)
 Phyllophilopsis evanida Reinhard
 HOST: COLEOPTERA

Endomychidae

Endomychus biguttatus Say
 Reinhard, 1958: 234, Md.
 STAGE PARASITIZED: Larva.
 Phytomyptera Rondani (Goniinae: Siphonini)
 Phytomyptera leucoptera, see *Dichaetoneura leucoptera*
 Phytomyptera walleyi Brooks
 HOST: LEPIDOPTERA

Olethreutidae

Zeiraphera ratzeburgiana (Ratzeburg)
 Brooks, 1945: 92, Ont.
 "Phytomyptera" leucoptera, see *Dichaetoneura leucoptera*
 Phytopsis californica, see *Athanatus californicus*
 Pilatea Townsend (Goniinae: Eryciini)
 Pilatea arator (Aldrich)
 Most records as *Masicera arator*.
 HOST: DIPTERA

Tipulidae

Unidentified Tipulidae
 Curran, 1927a: 17, Que. (as *Erycia*); Aldrich, 1925b: 33,
 Que. (ex large *Tipula* larva).
 HOST: LEPIDOPTERA

Lasiocampidae

Malacosoma americanum (Fabricius)
 West, 1928: 816, N.Y.
 STAGE PARASITIZED: Larva (Diptera).
 BIOLOGY: Aldrich, 1925b: (33).
 Pilatea delecta (Curran)
 HOST: LEPIDOPTERA

Noctuidae

Autographa californica (Speyer)

Spencer, 1936: 28, B.C. (as *Lydella*).

STAGE PARASITIZED: Larva.

Pilatea occidentalis (Coquillett)

IMMATURES: Pupa — Greene, 1922: (as *Sturmia*; descr., fig. 98 on pl. 20).

Plagia americana, see *Voria ruralis*

Plagia ruralis probably, see *Voria ruralis* probably

Plagiomima Brauer and Bergenstamm (Tachininae: Voriini)

Plagiomima cognata Aldrich

HOST: LEPIDOPTERA

Noctuidae

Heliothis spp.

Butler, 1958: 562, Ariz.

STAGE PARASITIZED: Larva.

Plagiomima spinosula (Bigot) probably

HOST: LEPIDOPTERA

Noctuidae

Heliothis spp.

Bottrell, Young, Price and Adams, 1968: 1054, Okla.

"*Plagioprospherysa*" *parvipalpis*, see *Stomatomyia parvipalpis*

Plagiprospherysa parvipalpis, see *Stomatomyia floridensis*
or *Stomatomyia parvipalpis*

Plagiprospherysa sp., see *Stomatomyia* sp.

Plectops amplicornis, see *Elfia amplicornis*

Plectops "currei", see *Elfia curriei*

Plectops curriei, see *Elfia curriei*

Plectops melissopodis, see *Elfia melissopodis*

Plectops sp. near *usitata*, see *Elfia* sp. near *usitata*

Plectops sp., see *Elfia* spp.

"*Pleteria*" *robusta*, see *Peleteria texensis*

Polistomyia subdivisa, see *Trichopoda subdivisa*

Polychaetoneura elyii, see *Actia americana*

Prolixophaga plumbea, see *Lixophaga plumbea*

Prophryno Townsend (Goniinae: Eryciini)

Prophryno marginalis (Aldrich and Webber)

All records as *Phorocera marginalis*.

HOST: LEPIDOPTERA

Geometridae

Semiothisa punctolineata (Packard)

All records ex *Macaria punctolineata*.

Vickery, 1925: 141, Tex.; Aldrich and Webber, 1924: 84, Tex.

STAGE PARASITIZED: Larva.

Prophryno parviteres (Aldrich and Webber)

HOST: LEPIDOPTERA

Geometridae

Melanchroia cephise (Cramer)

Wolcott, 1951a: 479, P.R. (as *Phorocera*).

Noctuidae

Plathypena scabra (Fabricius)

Hill, 1925: 17, Miss., Tenn. (as *Phorocera flavicauda* Wulp).

Pieridae

Ascia monuste (Linnaeus)

Wolcott, 1951a: 479, P.R. (as *Phorocera*).

STAGE PARASITIZED: Larva.

IMMATURES: Pupa — Greene, 1922: (descr., fig. 85 on pl. 18).

Prosenia Lepeletier and Serville (Proseninae: Prosenini)

Prosenia siberita (Fabricius)

HOST: COLEOPTERA

Scarabaeidae

Popillia japonica Newman

Fleming, 1963: 25, N.J.; Clausen, 1956: 6, 23, 123-124, N.J. (as *P. "sibirita"*); King and Parker, 1941: 127, 131, N.J. (as *P. "sibirita"* ex Japanese beetle); Gardner and Parker, 1940: 3, 10-11, 32, U.S. (as *P. "sibirita"*); King, 1939: 26, N.J. (ex Japanese beetle); Hadley, 1938: 211-213, N.J.; Sweetman, 1936: 122-123, 233, 243, fig. 105a, N. Amer.; King, 1931: 453, 457-458, N.J., Pa.; King, 1930: 160, N.J.; Smith, L. B., 1930: 500 (ex Japanese beetle); King, 1928: 405-406, fig. 133; Clausen, King and Teranishi, 1927: 4, 20-25, 54, figs. 14B, 20F-20I, 21A (non N. Amer.).

STAGE PARASITIZED: Larva.

IMMATURES: First instar larva — Clausen, King and Teranishi, 1927: (descr., fig. 20F-20H).

Third instar larva — King, 1928: (fig. 133); Clausen, King and Teranishi, 1927: (descr., fig. 20I, 21A).

Larva — Sweetman, 1936: (fig. 105a).

Pupa — Clausen, King and Teranishi, 1927: (descr., fig. 14B).

BIOLOGY: Clausen, 1956: (6, 23, 123-124); King, 1939: (26); Hadley, 1938: (211-212); Sweetman, 1936: (122-123, 233, 243); King, 1931: (453, 457-458); King, 1928: (405-406); Clausen, King and Teranishi, 1927: (4, 20-25, 54).

Prosena "sibirita," see *Prosena siberita*

Prosena spp.

HOST: COLEOPTERA

Scarabaeidae

Adoretus spp.

Agee, 1917: 17 (lab.); Agee, 1916a: 344-345 (lab.); Agee, 1916b: 8 (lab).

Anomala spp.

Agee, 1918: 35 (lab.); Agee, 1918a: 13 (lab.); Agee, 1917: 17 (lab.); Agee, 1916a: 344-345 (lab.); Agee, 1916b: 8 (lab.).

STAGE PARASITIZED: Larva.

BIOLOGY: Agee, 1917: (17); Agee, 1916a: (344-345); Agee, 1916b: (8).

Prosopaea Brauer and Bergenstamm (Goniinae: Eryciini)

"*Prosopaea*" spp.

HOST: LEPIDOPTERA

Noctuidae

Aletia sp.

Brauer and Bergenstamm, 1895: 570, 581.

Notodontidae

Cerura sp.

Brauer and Bergenstamm, 1895: 570, 586, N. Amer.

Sphingidae

Smerinthus ophthalmicus Boisduval

Brauer and Bergenstamm, 1895: 571, 606, N. Amer.

Prospberrya Wulp (Goniinae: Blondeliini)

Prospberrya promiscua, see *Lespesia archippivora*

Prospberrya similis, see *Tachinomyia similis*

"*Prospberrya*" sp.

HOST: HYMENOPTERA

Sphecidae

Bembix nubilipennis (Cresson)

Lafler, 1896: 62-63, Nebr. (as "*Prospberrya*" ex stores in nest of "*Bembex*").

"*Prospberrya*" sp., see "*Prospberrya*" sp.

Psalidopteryx Townsend (Goniinae: Graphogastrini)

Psalidopteryx macdunnoughi Brooks

HOST: LEPIDOPTERA

Scythridae

Scythis epilobiella McDunnough

McDunnough, 1942: 224, N.B., P.E.I.

Scythis sp.

Brooks, 1942: 146, P.E.I.

STAGE PARASITIZED: Larva.

Psalidopteryx nuda Brooks

HOST: LEPIDOPTERA

Olethreutidae

Griselda radicana (Walsingham) probably

Brooks, 1942: 147, B.C.

Psalidopteryx pollinosa Brooks

HOST: LEPIDOPTERA

Gelechiidae

Anarsia lineatella Zeller

Brooks, 1942: 148, B.C.

Psalidopteryx pseudonuda Brooks

HOST: LEPIDOPTERA

Tortricidae

Acleris variana (Fernald)

Brooks, 1942: 147, B.C. (ex *Peronea*).

***Psalidopteryx psilocorsiphaga* Brooks**

HOST: LEPIDOPTERA

Oecophoridae

Psilocorsis faginella (Chambers)

Embree, 1958: 170, N.S.

Psilocorsis spp.

Brooks, 1942: 150, Que.

Olethreutidae

Epinotia solicitana (Walker)

Smith, C. C., 1946: 120, N.B.

Eucosma rescissoriana Heinrich

Goyer and Schenk, 1969: 1063, 1065-1066, Idaho; Goyer, 1966: 17-18, Idaho.

STAGE PARASITIZED: Larva, pupa.

BIOLOGY: Goyer, 1966: (17-18).

***Psalidopteryx slossonae* Townsend**Most records as *Psalidopteryx "slossoni."*

HOST: LEPIDOPTERA

Gelechiidae

"Telphusa" sp. probably

Daviault, 1935: 252, Que. (as *P. slossonae*).

Unidentified Lepidoptera

Leaf tier(-s)

*Procter, 1946: 435, Maine (ex leaf tier, white birch, Procter, 1938); Procter, 1938: 376, Maine (ex leaf tier, white birch).

STAGE PARASITIZED: Larva.

Psalidopteryx "slossoni,"* see *Psalidopteryx slossonae***Psalidopteryx* sp.**

HOST: LEPIDOPTERA

Olethreutidae

Eucosma rescissoriana Heinrich

Ollieu and Schenk, 1966: 273, Idaho (probable host).

Pseudatractocera neomexicana*, see *Sitophaga neomexicana

- Pseudeuantha coquilletti*, see *Anaporia limacodis*
Pseudeuantha pristis, see *Anaporia pristis*
Pseudoarchytas incertus, see *Archytas incertus*
Pseudoarchytopsis sp., see *Archytas* spp.
Pseudochaeta Coquillett (Goniinae: Eumasicerini)
Pseudochaeta argentifrons Coquillett
 HOST: LEPIDOPTERA

Arctiidae

- Cisthene nexa* (Boisduval)
 Flanders, 1926: 158, Calif. (ex *Illice*).
Tigrioides bicolor (Grote)
 Brooks, 1945: 94, Que. (as *P. canadensis* Brooks ex *Lexis*).

Bombycidae

- Unidentified Bombycidae
 Coquillett, 1897: 20, 24, Calif.

Pyralidae

- Loxostege mancalis* (Lederer)
 Fattig, 1949: 28-29; Brimley, 1938: 364, N.C.; Brimley,
 1922: 25, N.C.
 STAGE PARASITIZED: Larva, pupa.
Pseudochaeta canadensis, see *Pseudochaeta argentifrons*
Pseudochaeta pyralidis, see *Metopiops pyralidis*
Pseudochaeta siminina Reinhard
 HOST: LEPIDOPTERA

Pyralidae

- Macrobotys pertextalis* (Lederer)
 Schaffner, 1959: 10, 53, Mass.
Palpita magniferalis (Walker)
 Raizenne, 1952: 162, Ont.
 BIOLOGY: Schaffner, 1959: (10, 53).
Pseudoclista globosa, see *Myiophasia globosa*
Pseudoeribea sp. near *tothilli*, see *Eribella* sp. near *exilis*
 "Pseudoeribia" sp. near *tothilli*, see *Eribella* sp. near *exilis*
 "Pseudomyotheria" *ancilla*, see *Pseudomyotheria ancilla*
Pseudomyotheria Townsend (Goniinae: Blondeliini)
Pseudomyotheria ancilla (Walker)
 HOST: COLEOPTERA

Chrysomelidae

Diabrotica undecimpunctata Mannerheim

Brauer and Bergenstamm, 1895: 571, 590 (as *P. indecisa*
Townsend ex *D. 12-punctata* Fabricius).

Scarabaeidae

Serica iricolor (Say)

Fattig, 1949: 29, Ga. (as "*Pseudomyotheria*").

HOST: LEPIDOPTERA

Gelechiidae

Filatima persicaeella (Murtfeldt)

Ingerson, 1918: 12, Mich. (as *Frontina* ex *Gelechia confusella*
Chambers).

STAGE PARASITIZED: Larva (Lepidoptera), pupa (Lepi-
doptera), adult (Coleoptera).

IMMATURES: Pupa — Greene, 1922: (descr., fig. 47 on pl. 10).

BIOLOGY: Fattig, 1949: (29).

Pseudomyotheria indecisa, see *Pseudomyotheria ancilla*

Pseudomyotheria tortricis, see *Erynnia tortricis*

Pseudomyotheria spp.

HOST: HYMENOPTERA

Tenthredinidae

Unidentified Tenthredinidae

Townsend, 1913a: 304.

HOST: LEPIDOPTERA

Unidentified Lepidoptera

Miscellaneous

Townsend, 1913a: 304.

STAGE PARASITIZED: Larva (Hymenoptera, Lepidoptera).

Pseudomyotheria spp. probably

HOST: LEPIDOPTERA

Unidentified Lepidoptera

Miscellaneous

Brauer and Bergenstamm, 1895: 571, 596 (ex Raupe auf
Viburnum nudum Linnaeus).

STAGE PARASITIZED: Larva.

Pseudoperichaeta Brauer and Bergenstamm (Goniinae: Eryciini)

Pseudoperichaeta erecta (Coquillett)

Most records as *Phorocera erecta*.

HOST: COLEOPTERA

Meloidae

Epicauta sp.

Aldrich and Webber, 1924: 77, Fla. (ex *Epicauta* (?) on beet, questionable host).

HOST: LEPIDOPTERA

Noctuidae

Nycteola spp. (including *N. cinereana* Neumoegen and Dyar, *N. columbiana* H. Edwards and/or *N. frigidana* Walker)

Schaffner and Griswold, 1934: 63-64, 114, Maine (ex *N. revayana* (Scopoli)).

Notodontidae

Schizura concinna (J. E. Smith)

Webber and Schaffner, 1926: 19, New England.

Olethreutidae

Episimus argutus (Clemens)

Schaffner, 1959: 20, 52, [N.E. U.S.]; Putman, 1943: 220, Ont.

Evora hemidesma (Zeller)

Roberts, 1966: 169, Iowa.

Grapholitha molesta (Busck)

All records ex *Laspeyresia molesta*.

Steenburgh and Boyce, 1938: 73, Ont.; Steenburgh, 1935: 71, Ont.

Melissopus latiferreanus (Walsingham)

Dohanian, 1942: 837, 840, Oreg. (probable host).

Spilonota ocellana (Denis and Schiffermüller)

Stultz, 1955: 80, N.S.

Pyralidae

Desmia funeralis (Hübner)

Essig, 1926: 582; Johnson, 1925: 198, Conn.; Aldrich and Webber, 1924: 78, Conn.

Loxostege similalis (Guenée)

Essig, 1926: 582; Aldrich and Webber, 1924: 77, Ark., Tex.; Reinhard, 1922: 72, Tex. (as *Exorista loxostegeae* Reinhard); Chittenden, 1911: 21, Tex.

Macrobotys pertextalis (Lederer)

Schaffner, 1959: 10, 52, [N.E. U.S.]; Putman, 1943: 219, Ont. (ex *Pyrausta*).

Ostrinia nubilalis (Hübner)

Most records ex *Pyrausta nubilalis*.

Baker, W. A., Bradley and Clark, 1949: 8, 39, 53, 158-159, 176, fig. 38Ca, U.S. (as *Pseudoperichaeta*); Clark, C. A., 1934: 18, 20, 31 (lab.); Baird, 1929: 39, Ont. (ex corn borer); Johnson, 1925: 198, Mass.; Aldrich and Webber, 1924: 78, Mass.; Caffrey, 1919: 21, Mass.; Vinal and Caffrey, 1919: 57-58, Mass.

Pachyzancla bipunctalis (Fabricius)

Chittenden, 1911: 21, D.C., Fla., Ga. or Tex.

Pantographa limata Grote and Robinson

Schaffner, 1959: 8, 52, [N.E. U.S.]; Putman, 1943: 219, Ont.

Phlyctaenia coronata tertialis (Guenée)

Schaffner, 1959: 10, 52, [N.E. U.S.] (ex *P. tertialis*).

Pyrausta futilalis (Lederer)

Schaffner, 1959: 11, 52, [N.E. U.S.].

Pyrausta signatalis (Walker)

Schaffner, 1959: 12, 52, [N.E. U.S.].

Tortricidae

Amorbia essigana Busck

Weber, 1953: 129, Calif.

Archips argyrospilus (Walker)

Prokopy, 1968: 350, Conn.; Schaffner, 1959: 30, 52, [N.E. U.S.].

Archips cerasivoranus (Fitch)

Schaffner, 1959: 30, 52, [N.E. U.S.].

Archips purpuranus (Clemens)

Schaffner, 1959: 31, 52, [N.E. U.S.].

Archips rosanus (Linnaeus)

Schaffner, 1959: 31, 52, Mass.; Putman, 1943: 222, Ont.

Argyrotaenia citrana (Fernald)

Essig, 1926: 582 (ex *Tortrix*).

Argyrotaenia mariana (Fernald)

Gilliatt, 1937: 145, N.S. (ex *Eulia*).

Choristoneura fumiferana (Clemens)

Blais, 1960: 388, 390, Que. (as *Pseudoperichaeta* ex spruce budworm); Coppel, 1960: 94, 97, B.C. (as *Pseudoperichaeta*); Carolin and Coulter, 1959: 553, Oreg.; McGugan and Blais, 1959: 765, Ont. (as *Pseudoperichaeta*); Anon., 1954b: 147, B.C. (as *Pseudoperichaeta*); MacNay, 1953a: 310, 314, 316, B.C.; Anon., 1952g: 288, 292, 295, B.C.; Ross, 1952: 108, fig. 4, Canada; Wilkes, Coppel and Mathers, 1949: 146, 149, 151-153, B.C.; Dowden, Buchanan and Carolin, 1948: 458 (ex *Archips*).

Choristoneura pinus Freeman

Dixon and Benjamin, 1963: 267, Wis.

Choristoneura rosaceana (Harris)

Most records ex *Archips rosaceana*.

Schaffner, 1959: 31, 52, [N.E. U.S.]; Raizenne, 1952: 214, Ont. (as *Pseudoperichaeta*); Schuh and Mote, 1948: 26, 28, 30, Oreg. (as *Pseudoperichaeta*); Zeller and Schuh, 1944: 48, Oreg.; Knowlton and Hansen, 1938: 130, Utah (ex *Cacoecia*); Knowlton and Allen, 1937: 784, Utah (ex *Cacoecia*).

Tortricid spp. (including *Amorbia humerosana* Clemens, *Archips argyrospilus* (Walker), *A. semiferanus* (Walker), *Argyrotaenia quadrifasciana* Fernald, *Choristoneura fractivittana* (Clemens), *C. rosaceana* (Harris) and/or *Pandemis limitata* (Robinson))

Hall, J. A., 1929: 139, Ont. (ex *Archips semiferana*, *A. argyrospila*, *A. rosaceana*, *A. fractivittana*, *P. limitata*, *Amorbia humerosana* and/or *Eulia quadrifasciana*).

Unidentified Lepidoptera

Leaf roller(-s)

Aldrich and Webber, 1924: 77, Wash.

STAGE PARASITIZED: Larva (Lepidoptera), pupa (Lepidoptera).

IMMATURES: Pupa — Ross, 1952: (fig. 4); Baker, W. A., Bradley and Clark, 1949: (fig. 38 Ca); Greene, 1922: (descr., fig. 52 on pl. 11).

BIOLOGY: Schaffner, 1959: (8, 10-12, 20, 30-31, 52);
Schaffner and Griswold, 1934: (63-64, 114).

Pseudoperichaeta erecta (Coquillett) probably
HOST: LEPIDOPTERA

Tortricidae

Amorbia essigana Busck

Ebeling, 1950: 664, Calif. (as tentatively *Phorocera*).

STAGE PARASITIZED: Larva.

Pseudoperichaeta sp. near *erecta* (Coquillett)
HOST: LEPIDOPTERA

Olethreutidae

Melissopus latiferreanus (Walsingham)

Dohanian, 1940: 856, Oreg. (as *Phorocera*).

STAGE PARASITIZED: Larva, pupa.

Pseudoperichaeta insidiosa (Robineau-Desvoidy)
Most records as *Zenillia roseanae* (Brauer and Bergenstamm).
HOST: LEPIDOPTERA

Olethreutidae

Grapholitha molesta (Busck)

Allen, Holloway and Haeussler, 1940: 4, 33, 40-43, 49 (as
Phorocera pumilio Aldrich, lab.); Allen, Holloway and
Haeussler, 1940: 3, 40-41, 60 (lab.).

Pyralidae

Ostrinia nubilalis (Hübner)

All records ex *Pyrausta nubilalis*.

Baker, W. A., Bradley and Clark, 1949: 3, 7, 12, 39, 53,
151, 154-158, fig. 38Cb, Mass. (as *Pseudoperichaeta*);
Caffrey and Worthley, 1927: 21, Mich., New England,
N. Y., Ohio; Thompson and Thompson,
1921: 127-139, figs. 1-17, 20 (lab.).

STAGE PARASITIZED: Larva.

IMMATURES: Egg — Thompson and Thompson, 1921: (descr.).

First instar larva — Thompson and Thompson, 1921: (descr.,
figs. 1-5).

Second instar larva — Thompson and Thompson,
1921: (descr., figs. 6-9, 20).

Third instar larva — Thompson and Thompson, 1921: (descr., figs. 10-15).

Pupa — Baker, W. A., Bradley and Clark, 1949: (fig. 38Cb); Thompson and Thompson, 1921: (descr., figs. 16-17).

BIOLOGY: Baker, W. A., Bradley and Clark, 1949: (3, 7, 12, 39, 53, 151, 154-158); Thompson and Thompson, 1921: (127-139).

Pseudoperichaeta roseanae, see *Pseudoperichaeta insidiosa*

Pseudosiphona Townsend (Goniinae: Siphonini)

Pseudosiphona brevirostris (Coquillett)

HOST: LEPIDOPTERA

Pterophoridae

Oidaematophorus homodactylus (Walker)

Schaffner, 1959: 19, 53, Mass.

BIOLOGY: Schaffner, 1959: (19, 53).

Pseudotachinomyia Smith (Goniinae: Exoristini)

Pseudotachinomyia slossonae (Townsend)

Most records as *Pseudotachinomyia webberi* Smith.

HOST: LEPIDOPTERA

Geometridae

Alsophila pometaria (Harris)

Schaffner and Griswold, 1934: 90, 121.

Erannis tiliaria (Harris)

Raizenne, 1952: 142, Ont. (as *P. slossonae*); Schaffner and Griswold, 1934: 97-98, 121.

BIOLOGY: Schaffner and Griswold, 1934: (121).

Pseudotachinomyia webberi, see *Pseudotachinomyia slossonae*

“*Pseudotraetocera*” *calosomae*, see *Sitophaga calosomae*

“*Pseudotraetocera*” *neomexicana*, see *Sitophaga neomexicana*

Ptilodexia Brauer and Bergenstamm (Proseninae: Prosenini)

“*Ptilodexia abdominalis* (Robineau-Desvoidy)”

HOST: COLEOPTERA

Scarabaeidae

Phyllophaga spp.

Criddle, 1918: 454, Man. (ex *Lachnosterna*).

STAGE PARASITIZED: Larva.

Ptilodexia canescens (Walker)

All records as *Eutheresia canescens*.

HOST: COLEOPTERA

Cerambycidae

Saperda calcarata Say

Peterson, 1948: 60-61, Ont.

Stenocorus inquisitor lineatus (Olivier) probablyChamplain and Knull, 1923: 214, Pa. (ex *Rhagium* sp.
probably *lineatum*).

STAGE PARASITIZED: Larva.

Ptilodexia cerata (Walker)

HOST: COLEOPTERA

Scarabaeidae

Phyllophaga spp.

Fattig, 1944: 3, Ga.

STAGE PARASITIZED: Larva.

Ptilodexia harpasa (Walker)

HOST: COLEOPTERA

Scarabaeidae

Chemarachis hogardi (Blanchard)Box, 1925: 351, Haiti (ex *Lachnosterna*).*Phyllophaga* spp.*Procter, 1946: 429, Maine (as *Rhamphinina*, Procter,
1938); Procter, 1938: 371, Maine (as *Rhynchioidexia*);
Johnson, 1927: 203, Maine.

May beetle(-s)

Johnson, 1925: 208, Conn., Maine, Mass., N.H., R.I., Vt.

STAGE PARASITIZED: Larva.

“*Ptilodexia tibialis* (Robineau-Desvoidy)”

HOST: COLEOPTERA

Scarabaeidae

Phyllophaga spp.All records ex *Lachnosterna* spp.Criddle, 1918: 454, Man.; Davis, J. J., 1918: 13-14; Davis,
J. J., 1916: 271, Tex.

STAGE PARASITIZED: Larva.

IMMATURES: Pupa — Greene, 1922: (descr., fig. 6 on pl. 2).

Ptilodexia sp.

HOST: COLEOPTERA

Scarabaeidae

Popillia japonica Newman

Fleming, 1963: 24, N.J.

"*Ptychomyia*" *selecta*, see *Bessa selecta*

Ptychomyia selecta, see *Bessa selecta*

"*Pyraustomyea*" *penitalis*, see *Pyraustomyia penitalis*

Pyraustomyia Townsend (Tachininae: Aphriini)

Pyraustomyia penitalis (Coquillett)

HOST: LEPIDOPTERA

Pyralidae

Ostrinia nubilalis (Hübner)

Most records ex European corn borer.

Anon., 1963e: 1337, Md.; Anon., 1961d: 639, Wis.; Blickenstaff, Arbuthnot and Harris, 1953: 366-369, Iowa (ex *Pyrausta nubilalis*); Arbuthnot and Wright, 1952: 233, Ill. (ex *Pyrausta* "*nubilalis*"); Cory, McConnell, Arbuthnot and Jones, 1952: 1, 11-12, Md. (ex *Pyrausta nubilalis*); Welch, 1952: 38, Ky.; Harris, 1951: 139-141, Iowa; Wilson, M. C., 1951: 224, Ind.; Arbuthnot, 1950: 45-46, S. part of N. Cent. U.S.; Arbuthnot, Jones, Carter and Evans, 1949: 2-5, 12, 16-24, Del., Ill., Ind., Iowa, Ky., Md., N.C., Nebr., N.J., Ohio, Pa., Va. (ex *Pyrausta nubilalis*); Baker, W. A., Bradley and Clark, 1949: 175, Mich., Ohio (ex *Pyrausta nubilalis*); Harris, 1949: 131, Iowa; MacCreary and Rice, 1949: 148, Del. (ex *Pyrausta nubilalis*); Arbuthnot, Jones, Carter and Evans, 1948: 2-3, 8-12, Ill., Iowa, Kans., Mo., N.J., Ohio, Pa., Va. (ex *Pyrausta nubilalis*); Arbuthnot, 1947: 30-31, Ill., Ind., Iowa; Deay, 1947: 68, Ind. (as "*Pyraustomyea*"); Drake, 1947: 127, 129, Iowa; Clark, C. A., and Jones, 1946: 6, 10, Ill., Wis.; Drake, 1946: 176, Iowa; Bradley, W. G., and Clark, 1940: 527, 531, Mich., Ohio (as *Panzeria*); Bradley, W. G., and Clark, 1939: 528, 534, Ind., Mich., Ohio (as *Panzeria*); Bradley, W. G., and Clark, 1938: 650, 654, Mich., Ohio (as *Panzeria*).

Ostrinia obliteralis (Walker)

All records ex *Pyrausta ainsliei* Heinrich.

Blickenstaff, Arbuthnot and Harris, 1953: 368, Iowa; Schopp, 1931: 34-36, Kans.; Allen, 1929: 685, Miss.;

Ainslie, G. G., and Cartwright, 1922: 11 (as *Panzeria*);
Ainslie, G. G., and Cartwright, 1921: 843, Tenn.

Ostrinia penitalis (Grote)

All records ex *Pyrausta penitalis*.

Poos, 1927: 82-83, Ohio (as *Panzeria*); *Ainslie, G. G., and
Cartwright, 1922: 11 (as *Panzeria*, questionable host,
Coquillett, 1897); Chittenden, 1918: 456 (as
Panzeria); Johnson, 1910: 779, N.J. (as *Panzeria*);
Coquillett, 1897: 19, 27, Mo. (as *Panzeria*).

Pyrausta spp.

Blickenstaff, Arbuthnot and Harris, 1953: 368, Iowa.

STAGE PARASITIZED: Larva.

IMMATURES: Pupa — Greene, 1922: (descr., fig. 41 on pl. 9).

BIOLOGY: Blickenstaff, Arbuthnot and Harris, 1953:
(366-369); Ainslie, G. G., and Cartwright, 1921: (843).

Red-tailed Tachina, see *Winthemia* spp. (including *W. leucanae*
probably or *W. militaris*)

Red-tailed Tachina fly, see *Winthemia quadripustulata*

Rhacodineura antiqua, see *Ocytata pallipes*

Rhamphinina harpasa, see *Ptilodexia harpasa*

Rhodogyne filiola, see *Gymnosoma filiola*

Rhodogyne fuliginosa, see *Gymnosoma fuliginosum*

Rhynchiodexia harpasa, see *Ptilodexia harpasa*

Rileya americana, see *Leschenaultia americana*

"*Rileymia*" *adusta*, see *Leschenaultia adusta*

"*Rileymia*" *triseta*, see *Leschenaultia americana*

Rileymyia adusta, see *Leschenaultia adusta*

Rileymyia americana, see *Leschenaultia americana*

Rileymyia exul, see *Leschenaultia exul*

Rileymyia fulvipes, see *Leschenaultia fulvipes*

Rileymyia triseta, see *Leschenaultia americana*

Rileymyia sp., see *Leschenaultia* spp.

Roeseliopsis Townsend (Goniinae: Hyperecteinini)

Roeseliopsis americana (Coquillett)

HOST: ORTHOPTERA

Phasmatidae

Anisomorpha buprestoides (Stoll)

Neff and Eisner, 1960: 102-103, figs. 1-2, 4, Fla.

STAGE PARASITIZED: Adult.

IMMATURES: Third instar larva — Neff and Eisner, 1960: (figs.
2, 4).

- Pupa — Neff and Eisner, 1960: (descr., figs. 1, 4).
Salmacia crassicornis, see *Gonia crassicornis*
Salmacia frontosa, see *Gonia frontosa*
Salmacia longipulvilli, see *Gonia longipulvilli*
Salmacia sp., see *Gonia* spp.
Schistocercophaga dampfi, see *Oedematocera dampfi*
Schizactia Townsend (Goniinae: Siphonini)
Schizactia vitinervis (Thompson)
 HOST: LEPIDOPTERA

Gelechiidae

- Aristotelia roseosuffusella* Clemens
 Thompson, 1911: 271 (as *Schizotachina*).
Coleotechnites coniferella (Kearfott)
 Raizenne, 1952: 232, Ont. (ex *Recurvaria*).
Coleotechnites piceaella (Kearfott)
 McLeod, J. M., 1964: 33, Que. (ex *Pulicalvaria*).
Coleotechnites sp.
 Renault and McDougall, 1961: 198-199, N.B. (ex *Evagora*).
Exoteleia pinifoliella (Chambers)
 Thompson, 1911: 271 (as *Schizotachina* ex *Paralechia*).
Exoteleia pinifoliella (Chambers) probably
 Champlain and Knull, 1923: 214, Pa. (as *Schizotachina* ex
Paralechia).

Olethreutidae

- Epinotia nanana* (Treitschke)
 Daviault and Ducharme, 1966: 697, Que.

Tineidae

- Unidentified Tineidae
 All records ex Tineidae on *Pinus australis* A. M. Michaux.
 Greene, 1934: 33 (as *Schizotachina*); Thompson, 1911:
 271 (as *Schizotachina*).
 STAGE PARASITIZED: Larva.
Schizocero-phaga Townsend (Goniinae: Blondeliini)
Schizocero-phaga leibyi Townsend
 HOST: HYMENOPTERA

Argidae

Sphacophilus cellularis (Say)

Most records ex *Schizocerus privatus* (Norton).

Thompson, 1953: 19 (ex *Sterictiphora cellularis*); Gould, G. E., and Walker, 1931: 379, Va. (ex *Sterictiphora collaris* Say); Brimley, 1928: 364, N.C.; Brimley, 1922: 25, N.C.; Townsend, 1916c: 77, N.C.

HOST: LEPIDOPTERA

Tortricidae

Archips cerasivoranus (Fitch)

*Thompson, 1953: 19 (questionable host, Hoffman, 1936); Hoffman, 1936: 211, Minn. (ex *Cacoecia*).

STAGE PARASITIZED: Larva (Hymenoptera), pupa (Hymenoptera).

Schizotachina Walker (Goniinae: Siphonini)

Schizotachina convecta (Walker)

HOST: LEPIDOPTERA

Pyralidae

Acrobasis comptoniella Hulst

Schaffner, 1959: 15, 54, Maine.

Tortricidae

Croesia albicomana (Clemens) probably

Thompson, 1911: 272 (ex *Tortrix*).

BIOLOGY: Schaffner, 1959: (15, 54).

Schizotachina vitinervis, see *Schizactia vitinervis*

Schizotachina sp.

HOST: LEPIDOPTERA

Gelechiidae

Chionodes sp.

Patton, 1958: 82, Fla. (ex *Chionodes* sp. on seagrape).

Scopolia sequax, see *Periscepsia helymus*

Scopolia sp., see *Periscepsia* spp.

Sipholeskia Townsend (Dexiinae: Leskiini)

Sipholeskia occidentalis (Coquillett)

HOST: LEPIDOPTERA

Aegeriidae

Synanthedon bibionipennis (Boisduval)

Cole, F. R., and Lovett, 1921: 300, Oreg. (as *Leskia gilensis* (Townsend) ex *Sesia rutilans* H. Edwards).

Siphona Meigen (Goniinae: Siphonini)

Siphona ceres (Curran)

All records as *Bucentes ceres*.

HOST: LEPIDOTPERA

Noctuidae

Unidentified Noctuidae

All records ex larva feeding on alfalfa.

Arnaud, 1963: 109, Guatemala; Curran, 1932: 14, Guatemala.

STAGE PARASITIZED: Larva.

Siphona conata (Reinhard)

HOST: LEPIDOPTERA

Diptoridae

Phryganidia californica Packard

Reinhard, 1959: 163, Calif. (as *Siphonopsis* ex *Phryganidia* [sp.]).

STAGE PARASITIZED: Pupa.

Siphona geniculata (De Geer)

HOST: DIPTERA

Tipulidae

Tipula caloptera Loew

Fattig, 1949: 10, Ga. (as *Bucentes*).

Tipula paludosa Meigen

Fitzpatrick, 1969: 355 (lab.); Williamson, 1969: 5, 8 (lab.).

STAGE PARASITIZED: Larva.

IMMATURES: Pupa — Greene, 1922: (descr., fig. 39 on pl. 9).

Siphona plusiae Coquillett

HOST: LEPIDOPTERA

Diptoridae

Phryganidia californica Packard

Bloeser, 1914: 301-309, fig. 1, pls. 39-40, Calif.

Noctuidae

Autographa californica (Speyer)

Clancy, 1969: 1079, Calif.; Essig, 1926: 579; Cole, F. R., 1915: 559, Calif.; Coquillett, 1897: 20, 26, Calif. (ex *Plusia*); Coquillett, 1895: 125, Calif. (ex *Plusia*).

Trichoplusia ni (Hübner)

Clancy, 1969: 1079, Calif.; Cole, F. R., 1915: 559, Calif. (ex *Autographa brassicae* Riley).

Unidentified Noctuidae

Coquillett, 1895: 125, Calif.

STAGE PARASITIZED: Larva.

IMMATURES: Third instar larva — Bloeser, 1914: (descr., fig. 1, figs. 1-6 on pl. 39 and figs. 1-2 on pl. 40).

Pupa — Greene, 1922: (descr., fig. 40 on pl. 9); Bloeser, 1914: (descr., fig. 1).

BIOLOGY: Bloeser, 1914: (301-309).

Siphona spp.

HOST: DIPTERA

Tipulidae

Crane fly (flies)

Anon., 1954a: 130, Que.

HOST: LEPIDOPTERA

Noctuidae

Autographa gamma (Linnaeus)

Brauer and Bergenstamm, 1895: 574, 603, N. Amer. (ex *Plusia*).

Trichoplusia ni (Hübner)

Oatman, 1966: 1138, Calif.

STAGE PARASITIZED: Larva (Diptera, Lepidoptera).

Siphonopsis conata, see *Siphona conata*

Siphophyto floridensis, see *Clausicella floridensis*

Siphophyto neomexicana, see *Clausicella neomexicana*

Siphophyto opaca, see *Clausicella opaca*

Siphophyto turmalis, see *Clausicella turmalis*

Siphosturmia Coquillett (Goniinae: Sturmiini)

Siphosturmia confusa Reinhard

HOST: LEPIDOPTERA

Nymphalidae

Euphydryas editha taylori (Edwards)

Tothill, 1913: 74, B.C. (as *Sturmia normula* Wulp ex
Lemonias taylori).

STAGE PARASITIZED: Larva.

Siphosturmia phyciodis (Coquillett)

Most records as *Sturmia phyciodis*.

HOST: LEPIDOPTERA

Nymphalidae

Phyciodes tharos (Drury)

Tothill, 1913: 74, Ont.

Phyciodes spp.

Reinhard, 1934: 19, Mass., Md. (as *Siphosturmiopsis*);
Coquillett, 1897: 21, 26, Md.

STAGE PARASITIZED: Larva.

Siphosturmiopsis phyciodis, see *Siphosturmia phyciodis*

Sisyropa Brauer and *Bergenstamm* (Goniinae: Carceliini)

Sisyropa alypiae Sellers

HOST: LEPIDOPTERA

Noctuidae

Alypia octomaculata (Fabricius)

Vide Sellers, 1943: 103, Mass. (Schaffner and Griswold,
1934); Schaffner and Griswold, 1934: 41, 113 (as
Zenillia n. sp. (1)).

Psychomorpha epimenis (Drury)

Sellers, 1943: 103 (ex *P. "epimensis"*).

BIOLOGY: Sellers, 1943: (103).

Sisyropa alypiae Sellers probably

HOST: LEPIDOPTERA

Noctuidae

Alypia octomaculata (Fabricius)

*Vide Sellers, 1943: 103 (Aldrich and Webber, 1924;
Coquillett, 1897); Webber and Schaffner, 1926: 22,
New England (as *Zenillia eudryae* (Townsend)); Aldrich
and Webber, 1924: 22, Mass. (as *Zenillia eudryae*);
Coquillett, 1897: 14, 23, Mo. (as *Exorista eudryae*).

STAGE PARASITIZED: Larva.

Sisyropa "chrysophanae," see *Aplomya theclarum*

Sisyropa chrysophani, see *Aplomya theclarum*

Sisyropa eudryae (Townsend)

HOST: LEPIDOPTERA

Arctiidae

Estigmene acraea (Drury)

Essig, 1926: 581 (as *Zenillia* ex *E. acraea*).

Unidentified Arctiidae

Aldrich, 1932: 2, N.H. (as *Exorista*).

Noctuidae

Acronycta hamamelis Guenée

Johnson, 1910: 780, N.J. (as *Exorista* ex *Acronycta*).

Acronycta sp.

Aldrich and Webber, 1924: 22, Mo. (as *Zenillia* ex *Acronycta*).

Agrotis ipsilon (Hufnagel)

All records ex *Agrotis ipsilon*.

*Crumb, 1929: 40, 59 (as *Zenillia*, Coquillett, 1897);
Johnson, 1910: 780, N.J. (as *Exorista*); Coquillett,
1897: 13 (as *Exorista*).

Alypia spp.

Johnson, 1925: 196, Conn., Maine, Mass., N.H., R.I. (as *Zenillia*).

Euthisanotia grata (Fabricius)

Fattig, 1949: 31; Sellers, 1943: 101, N.C., Vt.; Brimley,
1938: 362, N.C. (as *Zenillia*); Schaffner and Griswold,
1934: 63, 112, Mass., N.H., N.J., N.Y., Vt. (as *Zenillia*);
Aldrich, 1932: 3, Mass. (as *Zenillia*); Johnson,
1925: 196, Conn., Maine, Mass., N.H., R.I. (as *Zenillia*
ex *Eudryas*); Aldrich and Webber, 1924: 22, N.C., Pa.
(as *Zenillia*); Brimley, 1922: 22, N.C. (as *Exorista*);
Tothill, 1913: 70, Ont. (as *Exorista*).

Euthisanotia unio Hübner

Fattig, 1949: 31, Ga.; Sellers, 1943: 101, Ohio; Brimley,
1938: 362, N.C. (as *Zenillia*); Johnson, 1925: 196,
Conn., Maine, Mass., N.H., R.I. (as *Zenillia* ex *Eudryas*);
Aldrich and Webber, 1924: 22, Ohio (as *Zenillia*);
Brimley, 1922: 22, N.C. (as *Exorista*); Johnson,
1910: 780, N.J. (as *Exorista* ex *Eudryas*); Coquillett,
1897: 14, 25, Ohio (as *Exorista* ex *Eudryas*).

Euthisanotia spp.

Sellers, 1943: 101; Aldrich, 1932: 3, Mass. (as *Zenillia*);
 Aldrich and Webber, 1924: 22, N.Y. (as *Zenillia*);
 *Brauer and Bergenstamm, 1895: 553, 592, N. Amer.
 (as (*Exorista*) ex *Eudryas*, Townsend, 1892a); Townsend,
 1892a: 287-288, N.Y. (as *Exorista* ex *Eudryas*).

Nymphalidae

Vanessa atalanta (Linnaeus)

Johnson, 1910: 780, N.J. (as *Exorista* ex *Pyrameis*).

Saturniidae

Automeris io (Fabricius)

Collins, M. M., and Weast, 1961: 94 (as *Exorista*); Sellers,
 1943: 101; Aldrich and Webber, 1924: 23, Conn. (as
Zenillia); Johnson, 1910: 780, N.J. (as *Exorista* ex
Hyperchiria); Coquillett, 1897: 14, 25 (as *Exorista* ex
Hyperchiria).

Automeris spp.

Johnson, 1925: 196, Conn., Maine, Mass., N.H., R.I. (as
Zenillia).

Unidentified Lepidoptera

Moth(-s)

Essig, 1926: 581 (as *Zenillia*).

STAGE PARASITIZED: Larva.

IMMATURES: Pupa — Greene, 1922: (descr., fig. 27 on pl. 6).

BIOLOGY: Sellers, 1943: (101); Schaffner and Griswold,
 1934: (63, 112).

Sisyropa eudryae (Townsend) probably

HOST: LEPIDOPTERA

Noctuidae

Acronicta impleta Walker

All records ex *Acronycta luteicoma* Grote and Robinson.

*Vide Sellers, 1943: 101 (Coquillett, 1897); Coquillett,
 1897: 13, 23, Mo. (as *Exorista*).

Acronicta spp.

All records ex *Acronycta* sp.

*Vide Sellers, 1943: 101 (Coquillett, 1897); Coquillett,
 1897: 13, 23, Md., Mo. (as *Exorista*).

Notodontidae

Fentonia marthesia (Cramer)

All records ex *Heterocampa marthesia*.

*Vide Sellers, 1943: 101 (Coquillett, 1897); Coquillett, 1897: 14, 25, Mo. (as *Exorista*).

Nymphalidae

Vanessa atalanta (Linnaeus)

*Vide Sellers, 1943: 101 (Coquillett, 1897); Coquillett, 1897: 14, 27, Mass. (as *Exorista ex Pyrameis*).

STAGE PARASITIZED: Larva.

Sisyropa hemerocampae, see *Carcelia amplexa*

Sisyropa lagoae, see *Carcelia lagoae*

Sisyropa spp.

HOST: LEPIDOPTERA

Arctiidae

Arachnis picta Packard

Brauer and Bergenstamm, 1895: 574, 583.

Lymantriidae

Hemerocampa leucostigma (J. E. Smith)

Britton, 1917: 110; Schoene, 1909: 47, N.Y. (ex "*Hemerocampe*").

Megalopygidae

Megalopyge opercularis (J. E. Smith)

Brauer and Bergenstamm, 1895: 574, 595 (ex *Lagoa*).

Megalopyge pyxidifera (J. E. Smith)

Brauer and Bergenstamm, 1895: 574, 595 (ex *Lagoa*).

Noctuidae

Acronicta hastulifera (J. E. Smith)

Brauer and Bergenstamm, 1895: 574, 580 (ex *Acronycta*).

Agrotis ipsilon (Hufnagel)

Brauer and Bergenstamm, 1895: 574, 581 (ex *A. telifera* Harris).

Notodontidae

Heterocampa bilineata (Packard)

Brauer and Bergenstamm, 1895: 574, 600 (ex *Notodonta ulmi* Harris).

STAGE PARASITIZED: Pupa.

Sitophaga Gistel (Goniinae: Blondeliini)

Sitophaga calosomae (Townsend)

Most records as *Eubiomyia calosomae*.

HOST: COLEOPTERA

Carabidae

Calosoma calidum Fabricius

Collins and Hood, 1920: 488, 490, 495-497, [New England]; Townsend, 1916c: 74, Mass.; Burgess, 1897: 431, Mass. (as "*Pseudotraetocera*").

Calosoma frigidum Kirby

Collins and Hood, 1920: 490, 495, [Mass. or S. N.H.].

Calosoma peregrinator Guérin-Méneville

Burgess, 1897: 431, Calif. (as "*Pseudotraetocera*").

Calosoma sycophanta Linnaeus

Collins and Hood, 1920: 483-497, pls. 58-59, Maine, Mass., N.H., R.I.

Calosoma willcoxi LeConte

Collins and Hood, 1920: 495 (ex *C. "wilcoxi,"* unsuccessful lab. rearing).

Carabus nemoralis Müller

Collins and Hood, 1920: 490, 495, Mass.

STAGE PARASITIZED: Adult.

IMMATURES: Egg — Collins and Hood, 1920: (descr., pl. 58A).

First instar larva — Collins and Hood, 1920: (descr.).

Second instar larva — Collins and Hood, 1920: (descr., pl. 58D).

Third instar larva — Collins and Hood, 1920: (descr., pl. 58B-58C).

Pupa — Collins and Hood, 1920: (descr., pl. 59A-59C).

BIOLOGY: Collins and Hood, 1920: (483-497); Burgess, 1897: (431).

Sitophaga eleodivora (Walton)

HOST: COLEOPTERA

Tenebrionidae

Eleodes tricostata (Say)Walton, 1918: 25, Nebr. (as *Biomyia*).

STAGE PARASITIZED: Adult.

Sitophaga georgiae (Brauer and Bergenstamm)Most records as *Biomyia georgiae*.

HOST: COLEOPTERA

Carabidae

Calosoma calidum Fabricius

Burgess and Collins, 1917: 11-12 (incomplete rearing — pupa); Chittenden, 1898a: 76; Coquillett, 1897: 10, 23, Mass.

Calosoma frigidum Kirby

Burgess and Collins, 1917: 12 (incomplete rearing — pupa).

Calosoma peregrinator Guérin-Méneville*Vide Chittenden, 1898a: 76 (Coquillett, 1890); Coquillett, 1897: 10, 23, Calif.; Coquillett, 1890: 234 (as *Masicera* sp.).*Calosoma sycophanta* Linnaeus

Tothill, 1913a: 58, Mass.

Calosoma spp.Johnson, 1925: 193, Mass. (as *Viviania*).

Scarabaeidae

Phyllophaga crenulata (Frölich)*Vide Smyth, 1917: 59 (as *B. lachnosternae* (Townsend), Davis, J. J., 1916; Forbes, S. A., 1907); *vide Davis, J. J., 1916: 271, 279, N.E. U.S. (as *B. lachnosternae* ex *Lachnosterna* spp., Forbes, S. A., 1907); Forbes, S. A., 1907: 475, Ill. (as "*Viviania*" sp. ex *Lachnosterna*).*Phyllophaga* spp.Most records ex *Lachnosterna* spp.Fattig, 1949: 34-35, Ga. (as *Viviania* ex *Phyllophaga*); Reinhard, 1930: 104 (as *Viviania* ex *Phyllophaga*); Aldrich, 1925: 107, Miss. (as *Viviania*); Davis, J. J., 1918: 14 (as *B. lachnosternae*); Townsend, 1908: 106, Ill. (as *Viviania lachnosternae*, probable host).

STAGE PARASITIZED: Adult.

IMMATURES: Pupa — Greene, 1922: (descr., fig. 54 on pl. 12).

BIOLOGY: Tothill, 1913a: (58); Forbes, S. A., 1907: (475).

Sitophaga georgiae (Brauer and Bergenstamm) probably
 Most records as *Biomyia georgiae* probably.
 HOST: COLEOPTERA

Carabidae

Calosoma calidum Fabricius

Felt, 1913: 74 (incomplete rearing — pupa).

Calosoma sycophanta Linnaeus

Burgess and Collins, 1917: 12, pl. 2b, Mass.; Burgess and
 Collins, 1915: 19, [New England] (as *Viviania*).

STAGE PARASITIZED: Adult.

IMMATURES: Pupa — Burgess and Collins, 1917: (pl. 2b).

BIOLOGY: Felt, 1913: (74).

Sitophaga leechi (Curran)

All records as *Erycia leechi*.

HOST: COLEOPTERA

Carabidae

Carabus taedatus taedatus Fabricius

Arnaud, 1963: 113, B.C. (ex *C. taedatus* var. near
oregonensis LeConte); Curran, 1932a: 12-13, B.C. (ex *C.*
taedatus variety).

STAGE PARASITIZED: Adult.

Sitophaga neomexicana (Townsend)

HOST: COLEOPTERA

Scarabaeidae

Serica serotina LeConte

Essig, 1926: 580, Calif. (as "*Pseudotractocera*").

STAGE PARASITIZED: Adult.

Sitophaga nocturnalis (Reinhard)

HOST: COLEOPTERA

Scarabaeidae

Phyllophaga spp.

Reinhard, 1930: 104, Tex. (as *Viviania*, suggested host).

BIOLOGY: Reinhard, 1930: (104).

Sitophaga sp.

HOST: COLEOPTERA

Carabidae

Calosoma peregrinator Guérin-Méneville

Coquillett, 1890: 234, Calif. (as *Masicera*).

STAGE PARASITIZED: Adult.

BIOLOGY: Coquillett, 1890: (234).

Small tachinid [of Murtfeldt, 1894a], see *Carcelia* spp.

Spallanzania Robineau-Desvoidy (Goniinae: Goniini)

Spallanzania bucephala, see *Spallanzania hebes*

Spallanzania hebes (Fallén)

All records as *Spallanzania bucephala* (Meigen).

HOST: LEPIDOPTERA

Noctuidae

Feltia subterranea (Fabricius)

Patton, 1958: 83, Fla.; Tissot, 1945: 64, Fla.

Spallanzania hesperidarum, see *Acroglossa hesperidarum*

Spathidexia Townsend (Dexiinae: Thelairini)

Spathidexia dunningii (Coquillett)

HOST: LEPIDOPTERA

Hesperiidae

Ochlodes yuma (Edwards)

Arnaud, 1960: 1-2, 22, 25, fig. 8, Calif.

STAGE PARASITIZED: Larva.

IMMATURES: Pupa — Arnaud, 1960: (descr., fig. 8).

BIOLOGY: Arnaud, 1960: (1-2, 22, 25).

Spathimeigenia Townsend (Goniinae: Blondeliini)

Spathimeigenia aurifrons Curran

HOST: HYMENOPTERA

Diprionidae

Diprion polytomum (Hartig)

Reeks, 1938: 26, Que.

Monoctenus fulvus (Norton)

Raizenne, 1957: 11, Ont.

Neodiprion abietis (Harris) complex

Raizenne, 1957: 13, Ont.

Neodiprion lecontei (Fitch)

Raizenne, 1957: 20, Ont.; Benjamin, 1955: 21, 23, Canada,
Que., U.S.; Daviault, 1953: 181-184, Que.; Daviault,

1942: 397, Que.; Reeks, 1938: 26, E. Canada; Aldrich, 1931: 7, Pa.

Neodiprion maurus Rohwer

Raizenne, 1957: 16, Ont.

Neodiprion pinirigidae (Norton)

*Procter, 1946: 433, Maine (Procter, 1938); Procter, 1938: 375, Maine.

Neodiprion pratti banksianae Rohwer

Raizenne, 1957: 15, Ont.; Reeks, 1938: 26, E. Canada (ex jack pine sawflies).

Neodiprion rugifrons Middleton

Wilkinson, R. C., Becker and Benjamin, 1966: 791, Wis.; Raizenne, 1957: 22, Ont.

Neodiprion swaini Middleton

Reinhard, 1958a: 209, Que.

Neodiprion spp.

*Vide Aldrich, 1931: 7, Calif. (ex *Neodiprion* sp. (1), Greene, 1921); *vide Aldrich, 1931: 7, Calif. (ex *N. edwardsii* (Norton), Greene, 1921); Curran, 1930: 247, Ont.; Greene, 1921: 43, Calif. (as *Admontia hylotomae* Coquillett ex probably a *Diprion* sp.); Greene, 1921: 42, Calif. (as *S. spinigera* Townsend ex *N. edwardsii*).

Tenthredinidae

Pikonema alaskensis (Rohwer)

Raizenne, 1957: 32, Ont.

Pristiphora erichsonii (Hartig)

*Drooz, 1960: 34, Canada (Raizenne, 1957); Raizenne, 1957: 30, Ont.

Unidentified Hymenoptera

Sawfly(-flies) probably

Reinhard, 1958a: 209, Calif. (ex sawfly (?) on cypress).

STAGE PARASITIZED: Larva, pupa.

BIOLOGY: Daviault, 1953: (181-184).

Spathimeigenia spp. (*aurifrons* Curran-*erecta* Aldrich complex)

HOST: HYMENOPTERA

Diprionidae

Neodiprion nigroscutum Middleton

Becker, Jr., and Benjamin, 1967: 158-159, Wis.; Becker, Jr., and Benjamin, 1964a: 20, Wis.

Neodiprion swainei Middleton

Becker, Jr., and Benjamin, 1964: 356, Wis.

STAGE PARASITIZED: Larva, pupa.

Spathimeigenia erecta Aldrich

HOST: HYMENOPTERA

Diprionidae

Neodiprion lecontei (Fitch)Raizenne, 1957: 20, Ont., Que.; Benjamin, 1955: 21,
Canada; Daviault, 1953: 181, Que.*Neodiprion pratti banksianae* Rohwer

Raizenne, 1957: 15, Ont.

Neodiprion rugifrons Middleton

Raizenne, 1957: 22, Ont.

Neodiprion sertifer (Geoffroy)

Finlayson and Finlayson, 1958: 223-224, Ont.

Neodiprion sp.

Aldrich, 1931: 8, Mich.

STAGE PARASITIZED: Pupa.

Spathimeigenia erecta Aldrich probably

HOST: HYMENOPTERA

Diprionidae

Neodiprion lecontei (Fitch)

Reinhard, 1958a: 210, Canada.

Neodiprion pinetum (Norton)

Reinhard, 1958a: 210, Mich.

Spathimeigenia hylotomae (Coquillett)

HOST: HYMENOPTERA

Argidae

Arge humeralis (Palisot de Beauvois)All records ex *Hylotoma humeralis*.Aldrich, 1931: 8, Mass.; Johnson, 1925: 188, Mass. (as
Hylotomomyia); *Smith, H. E., 1912: 123 (as
Hyperecteina, Coquillett, 1898); Coquillett, 1898:
233, Mass. (as *Admontia* ex *H. "hemeralis"*).*Arge rubiginosa* (Palisot de Beauvois)

Patton, 1958: 83, Fla.

Arge spp.

Aldrich, 1931: 8, Conn., Va. (as "*Spathmeigenia*"); Greene, 1921: 42, Conn. (as *Admontia*); Greene, 1921: 42, Va. (as *Admontia*).

Diprionidae

Neodiprion lecontei (Fitch)

Benjamin, 1955: 20, U.S. (as "*Hylotomoyia hyletomae*"); Greene, 1921: 43, Va. (as *Admontia* ex N. "*lecontei*").

Unidentified Hymenoptera

Sawfly(-flies)

Essig, 1926: 579, Calif. and "east" (as *Admontia*).

IMMATURES: Pupa — Greene, 1922: (descr., fig. 57 on pl. 12).

Spathimeigenia hylotomae (Coquillett) probably

HOST: LEPIDOPTERA

Notodontidae

Heterocampa guttivitta (Walker)

Collins, 1926: 693, 695, N.H. (as *Admontia*).

STAGE PARASITIZED: Larva.

BIOLOGY: Collins, 1926: (693, 695).

Spathimeigenia mexicana Aldrich

HOST: HYMENOPTERA

Diprionidae

Zadiprion vallicola (Rohwer)

Aldrich, 1931: 6, Mexico (Micho.) (ex *Neodiprion*).

Spathimeigenia "spiniger," see *Spathimeigenia spinigera*

Spathimeigenia spinigera Townsend

Argidae

Arge pectoralis (Leach)

Tripp, 1960: 348.

Diprionidae

Diprion hercyniae (Hartig)

Finlayson, T., 1960a: 923, 938, Ont., Que.; Raizenne, 1957: 24, Ont.

Monoctenus fulvus (Norton)

Tripp, 1960: 348.

Neodiprion abbotii (Leach)

Patton, 1958: 83, Fla. (ex *N. fabricii* (Leach)); Hetrick, 1941: 376, Va. (as "*Spathimeigentia*" ex *N. americanum* (Leach)); Aldrich, 1931: 4, Va. (ex *N. "abbottii"*); Johnson, 1925: 186, Conn. (as *S. "spiniger"* ex *Lophyrus "abbotti"*); Johnson, 1910: 776, N.J. (as *Admontia demylus* (Walker) ex *Lophyrus*); Coquillett, 1897: 9, 22, Md. (as *Admontia demylus* ex *Lophyrus*).

Neodiprion abietis (Harris)

Tripp, 1960: 348; Reeks, Forbes and Cuming, 1947: 13, [maritime prov., Nfld.].

Neodiprion excitans Rohwer

Hetrick, 1959: 161, Fla.

Neodiprion lecontei (Fitch)

Griffiths, 1961: 196-197, Ont.; Patton, 1958: 83, Fla.; Reinhard, 1958a: 207, Tex.; Raizenne, 1957: 20, Ont.; Benjamin, 1955: 21, 24, 53, Canada, Ill., U.S.; Bentley, 1940: 191, Ala.; *Aldrich, 1931: 4, Va. (Coquillett, 1897); Johnson, 1925: 186, Conn. (as *S. "spiniger"* ex *Lophyrus*); Greene, 1921: 42, Pa.; Middleton, 1921: 757 (as "*Spathimeitenis*"); Middleton, 1921: 757 (as "*Adomonita demylus*"); Coquillett, 1897: 9, 22, Va. (as *Admontia demylus* ex *Lophyrus*).

Neodiprion maurus Rohwer

Tripp, 1960: 348.

Neodiprion nanulus nanulus Schedl

Tripp, 1960: 348.

Neodiprion pinetum (Norton)

Raizenne, 1957: 21, Ont.

Neodiprion pratti banksianae Rohwer

Griffiths, 1960: 654-656, Ont.; Raizenne, 1957: 15, Ont.

Neodiprion sertifer (Geoffroy)

Finlayson, T., 1960: 24, 45, figs. 23, 25, Ont.

Neodiprion swaini Middleton

Tripp, 1965: 96, Que.; Tripp, 1962: 809-817, Que.; Tripp, 1962a: 1250, 1265, Que.; Tripp, 1960: 347-359, figs. 1-12, Que.; Reinhard, 1958a: 207, Pa.; Raizenne, 1957: 12, Ont.

Neodiprion taedae linearis Ross

Warren, L. O., and Coyne, 1958: 20, Ark.

Neodiprion virginiana Rohwer

Greene, 1921: 42, Va. (ex *N. affinis* Rohwer).

Neodiprion spp.

- Aldrich, 1931: 5, Va.; Greene, 1921: 42, Pa. (ex *Neodiprion* sp. (1)); Greene, 1921: 42, Va. (ex *Neodiprion* sp. (2)).
 Diprionid spp. (including *Diprion* spp. and/or *Neodiprion* spp.)
 Townsend, 1912b: 165 (as *Admontia demylus* ex "Cophyrus" larvae).

Tenthredinidae

Pikonema alaskensis (Rohwer)

Tripp, 1960: 348.

Pristiphora erichsonii (Hartig)

Tripp, 1960: 348.

Unidentified Hymenoptera

Pine sawfly (-flies)

Reinhard, 1958a: 207.

Sawfly (-flies)

Aldrich, 1931: 5; Aldrich, 1931: 5, Pa.

HOST: LEPIDOPTERA

Unidentified Lepidoptera

Miscellaneous

*Procter, 1946: 433, Maine (Procter, 1938); Procter, 1938: 375, Maine.

STAGE PARASITIZED: Larva (Hymenoptera, Lepidoptera), pupa (Hymenoptera).

IMMATURES: First instar larva — Tripp, 1960: (descr., figs. 1-2, 8).

Second instar larva — Tripp, 1960: (descr., figs. 3-4, 6, 10).

Third instar larva — Finlayson, T., 1960: (descr., fig. 23); Tripp, 1960: (descr., figs. 5, 7, 9).

Pupa — Finlayson, T., 1960: (descr., fig. 25); Tripp, 1960: (descr., figs. 11-12); Greene, 1922: (descr., fig. 90 on pl. 19).

BIOLOGY: Tripp, 1965: (96); Tripp, 1962: (809-817); Finlayson, T., 1960a: (923, 938); Tripp, 1960: (347-359); Bentley, 1940: (191).

Spathimeigenia spinigera Townsend complex

HOST: HYMENOPTERA

Diprionidae

Neodiprion abbotii (Leach)

Finlayson, T., 1963: 476, 503.

Neodiprion abietis (Harris)

Finlayson, T., 1963: 477, 503.

Neodiprion lecontei (Fitch)

Finlayson, T., 1963: 479, 503.

Neodiprion nanulus nanulus Schedl

Finlayson, T., 1963: 481, 503.

Neodiprion pinetum (Norton)

Finlayson, T., 1963: 482, 503.

Neodiprion pratti banksianae Rohwer

Finlayson, T., 1963: 483, 503.

Neodiprion swaini Middleton

Finlayson, T., 1963: 484, 503.

Neodiprion virginiana Rohwer complex

Finlayson, T., 1963: 485, 503.

Spathimeigenia spp.

HOST: HYMENOPTERA

Diprionidae

Diprion hercyniae (Hartig)

Pepler, 1938: 17, Que. (ex European spruce sawfly).

Diprion polytomum (Hartig)

Brown, A. W. A., 1938: 16, Que.

Monoctenus fulvus (Norton)

Raizenne, 1957: 11, Ont., Que.

Neodiprion lecontei (Fitch)Finnegan, 1964: 29, Que.; Raizenne, 1957: 20, Ont., Que.;
Benjamin, 1955: 21, U.S.; Boyes and Wilkes, 1953: 126,
154; Baird, 1944a: 157, Ont.*Neodiprion maurus* Rohwer

Raizenne, 1957: 16, Ont.

Neodiprion pinetum (Norton)

Raizenne, 1957: 21, Ont.

Neodiprion rugifrons Middleton

Raizenne, 1957: 22, Ont.

Neodiprion sertifer (Geoffroy)

Raizenne, 1957: 14, Ont.

Neodiprion swaini Middleton

Atwood, 1944: 59, Que.

Neodiprion swaini Middleton probably

Martineau, 1959: 230, Que.

Neodiprion taedae linearis Ross

Warren, L. O., and Coyne, 1958: 20, Ark.

STAGE PARASITIZED: Larva, pupa.

"*Spathimeigientia*" *spinigera*, see *Spathimeigenia spinigera*

"*Spathimeitenis*" *spinigera*, see *Spathimeigenia spinigera*

"*Spathimeigenia*" *hylotomae*, see *Spathimeigenia hylotomae*

Spoggosia Rondani (Goniinae: Exoristini)

Spoggosia gelida (Coquillett)

HOST: LEPIDOPTERA

Lymantriidae

Dasychira spp. probably

Malloch, 1919: 57c, fig. 15 on pl. 8, Alaska (as *Euphorocera*
ex "*Dasychirus*" sp. (?)).

STAGE PARASITIZED: Pupa.

IMMATURES: Pupa — Malloch, 1919: (descr., fig. 15 on pl. 8).

BIOLOGY: Malloch, 1919: (57c).

Stomatodexia Brauer and Bergenstamm (Dexiinae: Leskiini)

Stomatodexia cothurnata (Wiedemann)

HOST: LEPIDOPTERA

Pyralidae

Diaphania hyalinata (Linnaeus)

Wolcott, 1951a: 477, P.R. (ex *Margaronia*).

Stomatolydella "infernalis," see *Paralispe infernalis*

Stomatolydella infernalis, see *Paralispe infernalis*

Statomyia Brauer and Bergenstamm (Goniinae: Exoristini)

Statomyia floridensis (Townsend)

HOST: LEPIDOPTERA

Pyralidae

Crambus trisectus Walker

Banerjee, 1967: 1173, Ill.

Elasmopalpus lignosellus (Zeller)

Leuck, 1966: 799-800, Ga.; Leuck and Dupree, 1965:
779-780, Ga. (part as *Plagiprospherysa parvipalpis* Wulp).

Pyrausta tyralis (Guenée)

Krombein, 1967: 150, Fla.

STAGE PARASITIZED: Larva, pupa.

BIOLOGY: Banerjee, 1967: (1173).

Stomatomyia parvipalpis (Wulp)

Most records as "*Plagioprospherysa*" *parvipalpis*.

HOST: LEPIDOPTERA

Hesperiidae

Hylephila phyleus (Drury)

Butler, 1961: 202, Ariz.

Pyralidae

Hulstia undulatella (Clemens)

Banerjee, 1967: 1173, western [U.S.] (as *Stomatomyia*).

Loxostege similalis (Guenée)

Reinhard, 1922: 73, Tex. (as *Plagioprospherysa*).

Loxostege sticticalis (Linnaeus)

Banerjee, 1967: 1173, western [U.S.] (as *Stomatomyia*).

HOST: ORTHOPTERA

Gryllacrididae

Stenopelmatus pictus Scudder

Duncan, 1923: 76-77, Calif.

Unidentified Orthoptera

Pictured sand cricket(-s)

Knowlton and Janes, 1931: 147, Utah.

STAGE PARASITIZED: Larva (Lepidoptera), pupa (Lepidoptera), adult (Orthoptera).

IMMATURES: Egg — Butler, 1961: (descr.).

BIOLOGY: Duncan, 1923: (76-77).

Stomatomyia sp.

HOST: LEPIDOPTERA

Pyralidae

Hellula rogatalis (Hulst)

Chittenden and Marsh, 1912: 31, Tex. (as *Plagioprospherysa* ex *H. undalis* (Fabricius)).

Strobliomyia Townsend (Goniinae: Siphonini)

Strobliomyia tibialis (Robineau-Desvoidy)

HOST: LEPIDOPTERA

Olethreutidae

Grapholitha molesta (Busck)

Allen, Holloway and Haeussler, 1940: 3, 40-41, 60 (as *Actia*, lab.).

STAGE PARASITIZED: Larva.

Sturmia Robineau-Desvoidy (Goniinae: Sturmiini)

Sturmia albifrons, see *Gymnocarcelia ricinorum*

Sturmia albincisa, see *Argyrophylax albincisa*

Sturmia austrina, see *Drino austrina*

Sturmia sp. near *bakeri*, see *Drino* sp. near *bakeri*

Sturmia balloui, see *Drino balloui*

Sturmia cubaecola, see *Drino cubaecola*

Sturmia discalis, see *Phrynofrontina discalis*

Sturmia distincta, see *Drino rhoeo*

Sturmia fraudulenta, see *Drino fraudulenta*

Sturmia harrisinae Coquillett

HOST: LEPIDOPTERA

Zygaenidae

Harrisina americana (Guérin-Méneville)

Patton, 1958: 83, Fla.; Coquillett, 1897: 20, 25.

Harrisina brillians Barnes and McDunnough

Clausen, 1961: 617-629, Ariz., Calif., Utah, Mexico (Chih.);

Clausen, 1956: 22, Ariz., Calif. (as *Sturmia* spp.); Jones,

M. L., and Forbes, 1956: 156, Calif. (as *Sturmia* sp.);

Jones, M. L., 1955: 175-176, Calif.; Smith, O. J., 1955:

7, Calif. (ex western grape leaf skeletonizer); Armitage,

1954: 209, Calif. (as *Sturmia* sp.); Smith, O. J., 1953: 9,

Ariz., Calif.; Smith, O. J., and Langston, 1953: 478, 483,

Ariz.; Jones, M. L., and Forbes, 1952: 161-162, Calif. (as

Masicera unispinosa Reinhard); Steinhaus and Hughes,

1952: 744, Ariz.

STAGE PARASITIZED: Larva.

BIOLOGY: Clausen, 1961: (617-629); Jones, M. L., 1955: (175-176); Smith, O. J., 1955: (7); Smith, O. J., and Langston, 1953: (478, 483).

Sturmia harrisinae Coquillett probably

HOST: LEPIDOPTERA

Zygaenidae

Acoloithus sp.

Clausen, 1961: 628, Fla.

Harrisina americana (Guérin-Méneville)

Clausen, 1961: 618, 628, Mo. east to N.Y., coastal states to Fla.

Harrisina brillians Barnes and McDunnough

Anon., 1957f: 828, Calif. (ex western grape leaf skeletonizer); Knowlton, Dorst, Lieberman, Thornley and Orr, 1953: 865, Utah.

Harrisina spp.

Clausen, 1961: 618, Mo. east to N.Y., coastal states to Fla.

STAGE PARASITIZED: Larva, pupa.

Sturmia sp. near *harrisinae* Coquillett

HOST: LEPIDOPTERA

Zygaenidae

Harrisina americana (Guérin-Méneville)

Patton, 1958: 83, Fla.

STAGE PARASITIZED: Larva.

Sturmia inca, see *Drino inca*

Sturmia incompta, see *Drino incompta*

Sturmia inconspicua, see *Drino inconspicua*

Sturmia inquinata, see *Drino incompta*

Sturmia nidicola, see *Townsendiellomyia nidicola*

Sturmia nigrita, see *Lydella radialis*

Sturmia normula, see *Siphosturmia confusa*

Sturmia occidentalis, see *Pilatea occidentalis*

Sturmia phyciodis, see *Siphosturmia phyciodis*

Sturmia pilatei, see *Drino pilatei*

Sturmia protoparcis, see *Drino rhoeo*

Sturmia ricinorum, see *Gymnocarcelia ricinorum*

Sturmia schizurae, see *Lespesia schizurae*

Sturmia scutellata, see *Blepharipa scutellata*

Sturmia sociabilis, see *Drino sociabilis*

Sturmia strigata, see *Blondelia polita*

Sturmia sp. near *strigata*, see *Blondelia* sp. near *polita*

"*Sturmia tipulensis* Tothill"

HOST: DIPTERA

Tipulidae

Unidentified Tipulini

Alexander, 1920: 731-732, Ont. (ex large tipuline crane fly).

STAGE PARASITIZED: Larva.

Sturmia trifida, see *Drino rhoeo* probably

Sturmia sp., see *Drino bohémica*, *Sturmia harrisinae* or "*Sturmia*" spp.

"*Sturmia*" spp.

HOST: HYMENOPTERA

Argidae

Arge sp.

Greene, 1921: 43, Conn.

HOST: LEPIDOPTERA

Citheroniidae

Anisota rubicunda (Fabricius)

Webber and Schaffner, 1926: 26, New England.

Anisota senatoria J. E. Smith

Webber and Schaffner, 1926: 25, New England.

Sphingidae

Manduca quinquemaculata (Haworth)

Gilmore, 1938: 710-711, Ky. and/or Tenn. (ex *Protoparce*).

Tortricidae

Choristoneura fumiferana (Clemens)

Gardiner, 1946: 80, N.B., Que. (ex *Archips*).

STAGE PARASITIZED: Larva (Lepidoptera).

"*Sturmia*" sp. probably
HOST: LEPIDOPTERA

Saturniidae

Hemileuca nevadensis Stretch

Packard, 1914: 502 (as dipterous parasite probably *Sturmia*
ex *H. maia nevadensis*).

STAGE PARASITIZED: Pupa.

Gen. and sp. of Sturmiini
HOST: LEPIDOPTERA

Noctuidae

Anticarsia gemmatalis Hübner

Plank, 1945: 26, P.R. (ex *A. "gemmatilis"*).

STAGE PARASITIZED: Larva.

Synaplomyia galerucellae, see *Aplomyiopsis galerucellae*

Tachina Meigen (Tachininae: Tachinini)

Tachina aletiae, see *Lespesia aletiae*

Tachina anonyma, see *Lespesia* spp.

Tachina archippivora, see *Lespesia archippivora*

Tachina armigera, see *Eucelatoria armigera*

Tachina bifasciata, see *Belvosia bifasciata*

Tachina blanda, see *Eusisyropa blanda*

Tachina clisiocampae, see *Exorista mella*

Tachina datanae, see *Archytas metallicus*

Tachina deilephilae, see *Winthemia deilephilae*

"*Tachina desmiae* Riley"

HOST: LEPIDOPTERA

Pyralidae

Desmia funeralis (Hübner)

All records ex *Desmia maculalis* Westwood.

Brauer and Bergenstamm, 1895: 575, 590; Riley, 1870b:
209.

STAGE PARASITIZED: Larva.

Tachina diabroticae, see *Celatoria diabroticae*

Tachina doryphorae, see *Doryphorophaga doryphorae*

Tachina fasciata, see *Exorista fasciata fasciata*

Tachina flavicauda, see *Triachora unifasciata*

Tachina fly, see Tachinidae page 537.

Tachina fraterna, see *Lespesia aletiae*

Tachina futilis, see *Euexorista futilis*

- Tachina hirsuta*, see *Phryxe vulgaris*
Tachina larvarum, see *Exorista mella* or *Exorista larvarum*
Tachina mella, see *Exorista mella*
Tachina militaris, see *Winthemia militaris*
Tachina "orgyia," see *Exorista mella*
Tachina orgyiae, see *Exorista mella*
Tachina phycitae, see *Nemorilla pyste*
Tachina robusta, see *Tachinomyia panaetius* or *Tachinomyia* spp.
Tachina rustica, see *Guerinia dydas*
Tachina simulans, see *Guerinia dydas*
Tachina tenthredinivora, see *Guerinia dydas*
Tachina theclarum, see *Aplomya theclarum*
 "Tachina" spp.

HOST: COLEOPTERA

Chrysomelidae

Coptocycla sp.

Townsend, 1895: 74, Colo.

HOST: HEMIPTERA (HOMOPTERA)

Cicadidae

Magiccada septendecim (Linnaeus)

Marlatt, 1898: 96-97, U.S. (ex *Cicada*, questionable parasite).

HOST: HYMENOPTERA

Apidae

Bombus spp.

Ashmead, 1894: 23.

Unidentified Hymenoptera

Bee(-s)

Packard, 1869: 407.

HOST: LEPIDOPTERA

Aegeriidae

Vespamima pini (Kellicott)

All records ex *Harmonia pini*.

*Packard, 1890: 729, N.Y. (Kellicott, 1885); Kellicott, 1885: 171-173, N.Y.

Arctiidae

Hyphantria textor (Harris)
Lochhead, 1919: 182.

Geometridae

Cankerworm(-s) (including *Alsophila pometaria* (Harris) and/or
Paleacrita vernata (Peck))
Harris, T. W., 1841: 340, Mass.

Lasiocampidae

Malacosoma americanum (Fabricius)
Surface, 1904: 43-44, N.E. U.S. (ex *Clisiocampa*).

Noctuidae

Alabama argillacea (Hübner)
Comstock, 1879: 204, Ala. (ex *Aletia*).
Papaipema nebris (Guenée)
All records ex "*Gortina*" *nitela* Guenée.
Gillette, 1890: 280, Iowa (as *Tachina* fly); *vide Riley and
Howard, 1890: 281, Iowa (Gillette, 1890).
Pseudaletia unipuncta (Haworth)
Kirkpatrick, 1861: 355, Ohio (as *Tachina* sp. (larger) ex
Leucania extranea Guenée).
Trichoplusia ni (Hübner)
Townsend, 1895: 73, Colo. (ex *Plusia brassicae* (Riley)).
Cutworm(-s)
Gillette, 1890: 280, Iowa (as *Tachina* fly); *vide Riley and
Howard, 1890: 281, Iowa (Gillette, 1890).

Psychidae

Thyridopteryx ephemeraeformis (Haworth)
Riley, 1887: 28 (ex *T. "ephemeroeformis"*); Lintner, 1883:
87, N.Y.

Pyralidae

Acrobasis indigenella (Zeller)
All records ex *Phycita nebulo* Walsh.

LeBaron, 1872: 121, Ill.; LeBaron, 1872: 121, Ill. (as small
Tachina sp.).

Crambus vulgivagellus Clemens
Lintner, 1883: 145-146, N.Y.

Saturniidae

Antheraea polyphemus (Cramer)
Fiske and Thompson, 1909: 453-454 (ex polyphemus).
Callosamia promethea (Drury)
Fiske and Thompson, 1909: 453-454 (ex promethea).
Platysamia cecropia (Linnaeus)
Fiske and Thompson, 1909: 453-454 (ex cecropia).

Sphingidae

Hyles lineata (Fabricius)
Lugger, 1899: 37, Minn. (ex *Deilephila*).

Unidentified Lepidoptera

Miscellaneous

Packard, 1869: 407 (ex caterpillars).

Moth(-s)

Morris, 1874: 177, Canada.

HOST: ORTHOPTERA

Phasmatidae

Bacillus rossia (Rossi)

All records ex *Bacillus rossii* Fabricius.

*Brauer and Bergenstamm, 1895: 575, 584 (Osten Sacken,
1877); Osten Sacken, 1877: 23.

Diapheromera femorata (Say)

*Brauer and Bergenstamm, 1895: 575, 591 (Osten Sacken,
1877); Osten Sacken, 1877: 23.

STAGE PARASITIZED: Larva (Coleoptera, Lepidoptera), pupa
(Lepidoptera), adult (Hemiptera).

IMMATURES: Egg — Walsh, 1867a: (descr.).

Third instar larva — Kirkpatrick, 1861: (descr.).

Pupa — Fiske and Thompson, 1909: (descr.); Kirkpatrick,
1861: (descr.).

BIOLOGY: Marlatt, 1898: (96-97); Riley, 1887: (28); Lintner,
1883: (145-146); Riley, 1871: (157).

Tachinidae, see page 537.

"Tachiniphyto" aenea, see *Elfia aenea*

Tachinomyia Townsend (Goniinae: Exoristini)

Tachinomyia acosta Webber

HOST: LEPIDOPTERA

Noctuidae

Lithophane antennata (Walker)

Webber, 1941: 300 (ex *Graptolitha*).

Lithophane disposita Morrison

Webber, 1941: 300 (ex *Graptolitha*).

Lithophane innominata (Smith)

Webber, 1941: 300 (ex *Graptolitha*).

Lithophane sp.

Webber, 1941: 300 (ex *Graptolitha*).

Tachinomyia nigricans Webber

HOST: HYMENOPTERA

Diprionidae

Monoctenus fulvus (Norton)

Raizenne, 1957: 11, Ont.

HOST: LEPIDOPTERA

Geometridae

Alsophila pometaria (Harris)

Raizenne, 1952: 108, Ont.; Webber, 1941: 302, Conn.

Ennomos subsignarius (Hübner)

Raizenne, 1952: 151, Ont.

Erannis tiliaria (Harris)

Webber, 1941: 302.

Itame pustularia (Guenée)

Webber, 1941: 302 (ex *Physostegania*).

Phigalia titea (Cramer)

Raizenne, 1952: 140, Ont.; Webber, 1941: 302.

Lasiocampidae

Malacosoma disstria Hübner

Sippell, 1961: 436, Ont.

Lymantriidae

Porthetria dispar (Linnaeus)

Webber, 1941: 302.

Noctuidae

Acronicta leporina (Linnaeus)

Raizenne, 1952: 38, Ont.

Baileya dormitans (Guenée)

Raizenne, 1952: 66, Ont. (as "*Tachinonyia*").

Catocala micronympha Guenée

Webber, 1941: 302 (ex *C. micronympha* form *fratercula*
Grote and Robinson).

Lithophane antennata (Walker)

Webber, 1941: 302 (ex *Graptolitha*).

Tortricidae

Choristoneura fumiferana (Clemens)

Ross, 1952: 108, fig. 10, Canada.

IMMATURES: Pupa — Ross, 1952: (fig. 10).

Tachinomyia panaetius (Walker)

HOST: LEPIDOPTERA

Lasiocampidae

Malacosoma disstria Hübner

Essig, 1926: 583, Calif. (as *Tachina robusta* (Townsend)).

Malacosoma spp.

Coquillett, 1897: 21, 24, Calif. (as *Tachina robusta* ex
Clisiocampa).

Lymantriidae

Hemerocampa vetusta (Boisduval)

Essig, 1926: 583, Calif. (as *Tachina robusta*).

Noctuidae

Agrotis ipsilon (Hufnagel)

All records ex *Agrotis ypsilon*.

Franklin, 1950: 57 (as *T. robusta*); *Crumb, 1929: 40, 59
(as *T. robusta*, Coquillett, 1897); Essig, 1926: 583,

Calif. (as *Tachina robusta*); Johnson, 1925: 204, Conn., Maine, Mass., N.H., R.I., Vt. (as *T. robusta*); Johnson, 1910: 781, N.J. (as *T. robusta*); Coquillett, 1897: 21, 23 (as *Tachina robusta*).

Epiglaea apiata (Grote)

Franklin, 1950: 57 (as *T. robusta*).

Cutworm(-s)

Fattig, 1949: 33; Smith, R. C., 1943: 377, Kans. (as *T. robusta*).

Unidentified Lepidoptera

Moth(-s)

Aldrich, 1915: 82 (as *Tachina robusta*).

STAGE PARASITIZED: Larva.

IMMATURES: Pupa — Greene, 1922: (descr., fig. 79 on pl. 17).

Tachinomyia sp. near *panaetius* (Walker)

HOST: LEPIDOPTERA

Noctuidae

Acronicta betulae Riley

Webber, 1941: 293 (ex *Acronycta*, lab.).

Catocala sp.

Webber, 1941: 293 (lab.).

Lithophane antennata (Walker)

Webber, 1941: 293 (ex *Graptolitha*, lab.).

Orthosia hibisci (Guenée)

Webber, 1941: 293 (ex *O. hibisci* form *insciens* Walker, lab.).

Tachinomyia robusta, see *Tachinomyia panaetius*

Tachinomyia similis (Williston)

HOST: LEPIDOPTERA

Arctiidae

Halisidota argentata Packard

Silver, 1958: 78, B.C.; Webber, 1941: 295.

Lasiocampidae

Malacosoma californicum californicum (Packard)

Webber, 1941: 295 (ex *Clisiocampa* "thoracia" Stretch).

Malacosoma californicum pluviale (Dyar)

Johansen, 1957: 70, Wash. (ex *M. pluviale*).

Malacosoma disstria Hübner

Johansen, 1957: 70, Wash.

Malacosoma spp. (including *M. californicum pluviale* (Dyar) and/or *M. disstria* Hübner)

Dailey, 1958: 269, Wash. (ex western tent caterpillar and/or forest tent caterpillar); Dailey, 1957: 628, Wash. (ex *M. disstria* and/or *M. pluviale*).

Malacosoma spp.

Ross, 1953: 19, 22, fig. 7, B.C.; *Brauer and Bergenstamm, 1895: 542, 588, Calif. (as *Achaetoneura* ex *Clisiocampa*, Williston, 1893); Williston, 1893: 256, Calif. (as *Prospheya* ex *Clisiocampa*).

Lymantriidae

Hemerocampa vetusta (Boisduval)

Webber, 1941: 295.

Stilpnotia salicis (Linnaeus)

Lejeune and Silver, 1961: 459-460, 462, 466, B.C.; Johansen, 1957: 70, Wash.; Webber, 1941: 295, Wash.

STAGE PARASITIZED: Larva, pupa.

IMMATURES: Pupa — Ross, 1953: (descr., fig. 7).

BIOLOGY: Lejeune and Silver, 1961: (459-460, 462, 466).

Tachinomyia variata Curran

HOST: LEPIDOPTERA

Lasiocampidae

Malacosoma californicum pluviale (Dyar)

Guppy, R., 1950: 4, B.C. (ex *M. pluviale*).

Lymantriidae

Porthetria dispar (Linnaeus)

Webber, 1941: 298, Mass.

Noctuidae

Nephelodes emmedonia (Cramer)

Webber, 1941: 298, N.B.

STAGE PARASITIZED: Pupa.

BIOLOGY: Guppy, R., 1950: (4).

Tachinomyia spp. (1), (2), (3) or (4), see *Tachinomyia* spp.

Tachinomyia spp.

HOST: LEPIDOPTERA

Geometridae

Alsophila pometaria (Harris)

Schaffner and Griswold, 1934: 90, 121, Maine, Mass., N.J.,
R.I. (as *Tachinomyia* sp. (3)).

Phigalia titea (Cramer)

Talerico, 1968: 561, Va.; Schaffner and Griswold, 1934: 97,
121, Mass., N.Y. (as *Tachinomyia* sp. (4)).

Lasiocampidae

Malacosoma disstria Hübner

Schaffner and Griswold, 1934: 89, 121 (as *Tachinomyia* sp.
(1)).

Lymantriidae

Stilpnotia salicis (Linnaeus)

*Burgess and Crossman, 1927: 14-15 (as *Tachina robusta*
(Townsend), Glendenning, 1924); Glendenning, 1924:
13, B.C. (as *Tachina robusta*).

Noctuidae

Acronicta betulae Riley

Schaffner and Griswold, 1934: 58, 121, N.J. (as *Tachinomyia*
sp. (1) ex *Acronycta*).

Catocala micronympha Guenée

Schaffner and Griswold, 1934: 67, 121, N.J. (as *Tachinomyia*
sp. (2) ex *C. micronympha* form *fratercula* Grote and
Robinson).

Lithophane antennata (Walker)

Schaffner and Griswold, 1934: 49, 121 (as *Tachinomyia* sp.
(1) ex *Graptolitha*).

Lithophane spp. (including *L. disposita* Morrison, *L.*
innominata (Smith) and/or *L. petulca* Grote)

Schaffner and Griswold, 1934: 48, 121 (as *Tachinomyia* sp.
(1) ex *Graptolitha* (including *G. disposita*, *G. innominata*
and/or *G. petulca*)).

Orthosia hibisci (Guenée)

Schaffner and Griswold, 1934: 46, 121 (as *Tachinomyia* sp.
(1) ex *O. hibisci* form normal *insciens* Walker).

Raphia frater Grote

Raizenne, 1952: 35, Ont.

Unidentified Catocalinae

Schaffner and Griswold, 1934: 67, 121, N.J. (as *Tachinomyia* sp. (2)).

Unidentified Noctuidae

Schaffner and Griswold, 1934: 72, 121 (as *Tachinomyia* sp. (1)).

STAGE PARASITIZED: Larva, pupa.

BIOLOGY: Schaffner and Griswold, 1934: (90, 97, 121);
Glendenning, 1924: (13).

"*Tachinomyia*" *nigricans*, see *Tachinomyia nigricans*

"*Tachinophyto*" sp., see *Lixophaga diatraeae*

Tachinophyto Townsend (Goniinae: Blondeliini)

Tachinophyto aenea, see *Elfia aenea*

Tachinophyto floridensis Townsend

HOST: LEPIDOPTERA

Pterophoridae

Oidaematophorus homodactylus (Walker)

Schaffner, 1959: 19, 54, Mass.

IMMATURES: Pupa — Greene, 1922: (descr., fig. 36 on pl. 8).

BIOLOGY: Schaffner, 1959: (19, 54).

Tachinophyto tortricis, see *Erynnia tortricis*

Tachinophyto vanderwulpi, see *Chaetonodexodes vanderwulpi*

Tachinophyto variabilis, see *Lixophaga variabilis*

Tachinophyto sp., see *Lixophaga diatraeae* or "*Tachinophyto*" spp.

"*Tachinophyto*" spp.

HOST: LEPIDOPTERA

Gelechiidae

Isophrictis similiella denotata Bottimer

Bottimer, 1926: 810, Tex.

Olethreutidae

Epiblema tripartitana (Zeller)

Bottimer, 1926: 816, Tex.

Laspeyresia pomonella (Linnaeus)

All records ex *Carpocapsa pomonella*.

Lochhead, 1919: 224; Jenne, 1909: 30, Ark.

Pyralidae

Desmia funeralis (Hübner)

Lochhead, 1919: 209.

Diatraea saccharalis (Fabricius)

Wolcott, 1915: 32, Cuba; Van Dine, 1913: 254, P.R.; Van Dine, 1913c: 29, P.R.

Tineidae

Unidentified Tineidae

Townsend, 1913a: 304.

STAGE PARASITIZED: Larva.

Thelaira Robineau-Desvoidy (Dexiinae: Thelairini)*Thelaira americana* BrooksAll records as *Thelaira leucozona* (Panzer).

HOST: LEPIDOPTERA

Arctiidae

Diacrisia virginica (Fabricius)

Schaffner and Griswold, 1934: 36, 125-126.

Estigmene acrea (Drury)

Schaffner and Griswold, 1934: 38, 125-126.

BIOLOGY: Schaffner and Griswold, 1934: (125).

Thelaira leucozona, see *Thelaira americana**Thelaira* spp.

HOST: LEPIDOPTERA

Arctiidae

Isia isabella (J. E. Smith)All records ex *Pyrrharctia isabella*.Townsend, 1895: 78, Colo.; Townsend, 1895: 78, Colo. (as *Thelaira* n. sp.).*Turuptiana permaculata* (Packard)Townsend, 1895: 78, Colo. (ex *Ecpantheria*).

Unidentified Arctiidae

Brauer and Bergenstamm, 1895: 578, 583, N. Amer.

Pyralidae

Diaphania nitidalis (Stoll) probablyBrauer and Bergenstamm, 1895: 578, 602, N. Amer. (ex *Phakellura*).

Thelymyia Brauer and Bergenstamm (Goniinae: Eryciini)

Thelymyia mathesoni (Reinhard)

HOST: LEPIDOPTERA

Lymantriidae

Tussock moth(-s) probably

Patton, 1958: 86, Fla. (as *Zenillia* ex caterpillar on poison ivy and *Viburnum semitomentosus* Small).

STAGE PARASITIZED: Larva.

Theresia Robineau-Desvoidy (Proseninae: Theresiini)

Theresia "claripalpis," see *Paratheresia claripalpis*

Theresia claripalpis, see *Paratheresia claripalpis*

Theresia diatraeae, see *Paratheresia claripalpis*

Theresia monohammi, see *Eutheresia monohammi*

Theresia rutilans (Fabricius)

HOST: COLEOPTERA

Cerambycidae

Enaphalodes atomarius (Drury)

Fattig, 1949: 33-34 (ex *Romaleum*).

Gen. and sp. of Theresiini

HOST: ORTHOPTERA

Gryllidae

Anurogryllus muticus (De Geer)

Weaver and Sommers, 1969: 342, La.

Thompsonomyia Brooks (Tachininae: Linnaemyini)

Thompsonomyia anthracina (Thompson)

HOST: LEPIDOPTERA

Arctiidae

Parasemia parthenos Harris

Thompson, 1911: 268, Ont. (as *Linnaemya* ex *Hyphoraia*).

STAGE PARASITIZED: Larva.

Thryptocera americana, see *Actia americana*

Thryptocera flavipes, see *Actia flavipes*

Thysanomyia Brauer and Bergenstamm (Goniinae: Sturmiini)

Gen. and sp. near *Thysanomyia*

HOST: LEPIDOPTERA

Sphingidae

Hemaris thysbe (Fabricius)

Schaffner and Griswold, 1934: 22, 118, Mass.

STAGE PARASITIZED: Pupa.

BIOLOGY: Schaffner and Griswold, 1934: (22, 118).

Tortriciophaga tortricis, see *Erynnia tortricis*

Townsendiellomyia Baranov (Goniinae: Sturmiini)

Townsendiellomyia nidicola (Townsend)

HOST: LEPIDOPTERA

Lymantriidae

Euproctis chrysorrhoea (Linnaeus)

Lochhead, 1919: 202, 406 (as *Zygobothria*); Howard and

Fiske, 1911: 262, 289-295, 309 (as *Zygobothria*);

Britton, 1910: 20, U.S. (as *Zygobothria*).

Nygmia phaeorrhoea (Donovan)

Most records ex brown-tail moth.

Dowden, 1962: 27, 29, 66, New England (ex *N.*

phaeorrhoea); Clausen, 1956: 6, 64, 133-134 (ex *N.*

phaeorrhoea); Dowden, 1935: 509 (as *Sturmia* ex *N.*

phaeorrhoea); Collins, 1934: 31, [Maine, Mass., N.H.]

(as *Sturmia*); Proper, 1934: 371-372, New England (as

Sturmia); Collins and Jones, 1932: 237 (as *Sturmia*);

Burgess and Crossman, 1929: 131-135, 142, Maine,

Mass., N.H. (as *Sturmia* ex *N. phaeorrhoea*); Britton

and Ashworth, 1923: 316, Conn. (as *Sturmia*);

Howard, 1918: 46 (as *Zygobothria*); Thompson, 1910:

290 (as *Zygobothria*).

Porthetria dispar (Linnaeus)

Britton and Ashworth, 1923: 316, Conn. (as *Sturmia* ex

"gipsy" moth); Britton and Davis, 1918: 250, Conn. (as

Zygobothria); Howard and Fiske, 1911: 232, probably

Mass. (as *Zygobothria*).

STAGE PARASITIZED: Larva.

IMMATURES: Pupa — Howard and Fiske, 1911: (descr.);

Thompson, 1910: (descr.).

BIOLOGY: Clausen, 1956: (6, 64, 133-134); Dowden, 1935:

(509); Proper, 1934: (371-372); Burgess and Crossman,

1929: (131-135, 142); Howard and Fiske, 1911: (262,

289-295, 309); Thompson, 1910: (290).

Trepophrys Townsend (Goniinae: Eryciini)

Trepophrys cinerea Townsend

HOST: LEPIDOPTERA

Tineidae

Amydria sp.

Townsend, 1908: 96, Mexico (Mor.).

STAGE PARASITIZED: Pupa.

Triachora Townsend (Goniinae: Belvosiini)*Triachora luteola* (Coquillett)

HOST: LEPIDOPTERA

Noctuidae

Leucania sp.Patton, 1958: 38, Fla. (as *Goniomima* ex *Leucania* on sugar
cane).*Triachora ochriventris* (Wulp)

HOST: LEPIDOPTERA

Noctuidae

Armyworm(-s)

Aldrich, 1928a: 14, Mexico (as *Belvosia*).*Triachora omissa* (Aldrich)

HOST: LEPIDOPTERA

Noctuidae

Feltia subgothica (Haworth)Fattig, 1949: 9, Ga. (as *Belvosia*).

STAGE PARASITIZED: Pupa.

Triachora unifasciata (Robineau-Desvoidy)

HOST: LEPIDOPTERA

Noctuidae

Alabama argillacea (Hübner)Brauer and Bergenstamm, 1895: 576 (as *Tachina flavicauda*
(Riley) ex *Aletia*).*Pseudaletia unipuncta* (Haworth)Guppy, J. C., 1967: 97, 99, 102, Ont.; Breeland, 1958: 320,
323, 333-334, Tenn. (as *Belvosia*); Franklin, 1950: 57
(ex *Cirphis*); Fattig, 1949: 9 (as *Belvosia* ex *Cirphis*);
Brimley, 1938: 361, N.C. (as *Belvosia* ex *Cirphis*);
Schaffner and Griswold, 1934: 47, 109, Mass. (as
Belvosia ex *Cirphis*); Johnson, 1925: 193, Conn., Mass

(ex *Heliophila* "unipunctata"); Allen, 1924: 24, Miss. (as *Belvosia* ex "Criphis"); Knight, 1916: 760, fig. 1 on pl. 40, N.Y. (as *Goniomima* ex *Leucania*); Britton, 1915a: 167, Conn. (as *Goniomima* ex *Heliophila*); Sherman, 1915: 299-302, N.C. (as "Goniomyia" ex *Heliophila*); Johnson, 1910: 778, N.J. (ex *Leucania*); Felt, 1906: 65, N.Y. (as "Belvoisia" ex *Heliophila* "unipunctata"); Coquillett, 1897: 10, 26, N.Y. (as *Belvosia* ex *Leucania*); Lintner, 1897: 210, N.Y. (as "Belvoisia" ex *Leucania*); Riley, 1883: 126 (as *Exorista flavicauda* ex *Leucania*); Thomas, C., 1881: 37-38, Ill. (as *Exorista flavicauda* ex *Leucania*); Riley, 1876: 53 (as *Exorista flavicauda* ex *Leucania*); Riley, 1870: 51-52, [Mo.] (as *Exorista flavicauda* ex *Leucania*, probable host).

Armyworm(-s)

Brimley, 1922: 21, N.C. (as *Belvosia*); Riley, Packard and Thomas, 1878: 323 (as *Exorista flavicauda*).

STAGE PARASITIZED: Larva, pupa.

IMMATURES: Pupa — Knight, 1916: (fig. 1 on pl. 40).

BIOLOGY: Schaffner and Griswold, 1934: (47, 109); Knight, 1916: (760); Lintner, 1897: (210); Riley, 1883: (126).

Trichiopoda pennipes, see *Trichopoda pennipes*

Trichiopoda sp., see *Trichopoda* spp.

Tricholyga sp. probably, see *Exorista* spp. probably

Trichopareia sp., see *Admontia* spp.

"Trichoparia" sp., see *Admontia* spp.

Trichophora miscelli, see *Deopalpus miscelli*

Trichophora ruficauda, see *Copecrypta ruficauda*

Trichopoda Berthold (Phasiinae: Trichopodini)

Trichopoda lanipes (Fabricius)

HOST: HEMIPTERA (HETEROPTERA)

Coreidae

Acanthocephala femorata (Fabricius)

Patton, 1958: 84, Fla.

Archimerus alternatus (Say)

Fattig, 1949: 34, Ga.

Pentatomidae

Nezara viridula (Linnaeus)

Drake, 1920: 75, Fla.

Podisus maculiventris (Say)

Fattig, 1949: 34, Ga. (as *T. radiata* Loew).

STAGE PARASITIZED: Adult.

IMMATURES: Pupa — Greene, 1922: (descr., fig. 23 on pl. 5).

Trichopoda pennipes (Fabricius)

HOST: HEMIPTERA (HETEROPTERA)

Coreidae

Acanthocephala declivis (Say)

Drake, 1920: 68, Fla. (incomplete rearing — egg).

Acanthocephala femorata (Fabricius)

Watson, J. R., 1922: 59R, Fla. (ex "*Acanthocephalus*");

Drake, 1920: 68, Fla. (incomplete rearing — egg).

Acanthocephala spp.

Bratley, 1936: 66, Fla.

Anasa armigera (Say)

Beard, 1940a: 625, Conn.

Anasa tristis (De Geer)

Yonke and Medler, 1969: 170-171, Wis.; Daugherty, 1959: 888, N.C. (ex squash bug); Dietrick and Bosch, 1957: 627-629 (lab.); Johansen, 1957: 64, Wash.; Clausen, 1956: 13, 57, Wash.; Milliron, 1956: 860, Del. (ex squash bug); Clausen, 1956: 13, 57, Wash.; Clausen, 1955: 78, Conn. (as *T. pennipes* squash bug strain); Fattig, 1949: 34, Ga.; MacCreary and Parker, 1948: 20, Del. (ex squash bug); Annand, 1946: 137 (ex squash bug, lab.); Leaver, 1946: 34, R.I.; Annand, 1945: 146, Conn., N.J. (ex squash bug); Eichmann, 1944: 36, Wash.; Eichmann, 1943: 39, N.J., Wash.; Beard, 1940: 269-272, fig. 1, Conn.; Beard, 1940a: 607, 624-672, figs. 10-13, pl. 1E-1F, Conn.; Beard, 1936: 335-336, Conn.; Allen, 1929: 682, Miss.; Essig, 1926: 578 (ex squash bug); Worthley, 1924: 7-16, Mass.; Worthley, 1924a: 57-77, figs. 16-21 on pl. 3, Mass. (ex squash bug); Worthley, 1923: 77-79, Mass.; Girault, 1913: 56, Tex.; Johnson, 1910: 775, N.J.; Weed and Conradi, 1902: 20-21, N.H.; Chittenden, 1899: 26; Coquillett, 1897: 21-22, Pa.; Brauer and Bergenstamm, 1895: 579, 582, Cent. Amer.; Cook, 1889: 95-96, Mich.

Anasa spp.

Aldrich, 1915: 83.

Archimerus alternatus (Say)

Yonke and Medler, 1969: 176-177, Wis.

Archimerus calcarator (Fabricius)

Drake, 1920: 68, Fla.

Chelinidea canyona Hamlin

Mann, 1969: 14, 127, Tex.; Hamlin, 1924: 207, Tex.

Chelinidea tabulata (Burmeister)

Mann, 1969: 14, 127, Tex.

Chelinidea vittiger Uhler

Mann, 1969: 14, 127, Tex.

Chelinidae vittiger aequoris McAtee

Hamlin, 1924: 199, Fla., Tex.; Hamlin, 1924: 201, Fla. (ex
C. vittiger aequoris var. *artuatra* McAtee).

Euthochtha galeator (Fabricius)

Yonke and Medler, 1969: 183-184, Wis.

Leptoglossus oppositus Say

Fattig, 1949: 34; Morrill, 1910: 92, Tex. (incomplete rearing
— egg); Chittenden, 1902: 25.

Leptoglossus phyllopus (Linnaeus)

Capelouto, 1949: 33, Fla. (incomplete rearing — egg); Fattig,
1949: 34; Bratley, 1936: 65-66, Fla. (incomplete rearing
— egg); Allen, 1929: 682, Miss.; Worthley, 1924: 10,
Tex.; Reinhard, 1922: 73, Tex. (as *Trichiopoda*);
Watson, J. R., 1922: 59R, Fla.; Drake, 1920: 68, Fla.

Leptoglossus spp.

Essig, 1926: 578 (ex leaf-footed plant bugs); Aldrich, 1915:
83.

Narnia pallidicornis Stål

Mann, 1969: 14, 135.

Largidae

Largus cinctus californicus (Van Duzee)

Gerhardt, 1942: 43 (ex *Euryophthalmus*).

Largus sp.

Sabrosky, 1955: 711, Calif.

Pentatomidae

Acrosternum hilare (Say)

Underhill, 1934: 17, Va.; Drake, 1920: 45, 68, Fla.

Acrosternum pennsylvanicum (De Geer)

Drake, 1920: 68, Fla.

Alcaeorrhynchus grandis (Dallas)

Watson, J. R., 1922: 59R, Fla.

Euschistus servus (Say)

Capelouto, 1949: 33, Fla. (ex brown stink bug, incomplete
rearing — egg); Fattig, 1949: 34, Ga.

Euschistus variolarius (Palisot de Beauvois)

Parish, 1934: 54, Iowa.

Murgantia histrionica (Hahn)

Davis, 1964a: 370 (lab.); Worthley, 1924: 10, Tex. (incomplete rearing — egg).

Nezara viridula (Linnaeus)

Funasaki, 1969: 555, Hawaii (as *T. pennipes* var. *pilipes* (Fabricius), incomplete rearing — egg); Miyahira, 1969: 241, Hawaii (incomplete rearing — egg); Mitchell, 1968: 18, Hawaii (incomplete rearing — egg); Shahjahan, 1968: 1088-1090, Hawaii (as *T. pennipes pilipes*); Shahjahan, 1968a: 1102, Hawaii (as *T. pennipes pilipes*); Clausen, 1956: 57; Hayslip, Genung, Kelsheimer and Wilson, 1953: 47, fig. 30, Fla. (ex southern green stink bug); Simmonds, 1953a: 782, Fla., Ga.; Wolcott, 1951a: 471, P.R. (ex southern green stink bug); Capelouto, 1949: 31-33, unnumbered fig. on p. 32, Fla.; Fattig, 1949: 34, Ga.; Hayslip, 1945: 199, Fla.; Hayslip, 1944: 124, Fla.; Bratley, 1939: 95, Fla.; Bratley, 1937: 72, Fla.; Bratley, 1936: 65-66, Fla.; Watson, J. R., 1936: 406, Fla.; Douglas, 1935: 686, La.; Watson, J. R., 1935: 244, Fla.; Bratley, 1934: 52, Fla.; Watson, 1934: 43, Fla.; Bratley, 1933: 77, Fla.; Watson, J. R., and Berger, 1932: 101-103, 132, fig. 65, Fla.; Anon., 1930a: 400, Fla. (ex southern green stink bug); Watson, J. R., 1928: 46R, Fla.; Wilson, C. E., 1923: 14, Virgin Islands; Wilson, C.E., 1923a: 29, Virgin Islands; Reinhard, 1922: 73, Tex. (as *Trichiopoda*); Watson, J. R., 1922: 59R, Fla.; Watson, J. R., 1921: 32R, Fla. (as "*Trichopodes*" ex "*Nezara*," incomplete rearing — egg); Dozier, 1920: 372, Fla.; Drake, 1920: 67-74, figs. 21, 23-24, Fla.; Jones, T. H., 1918: 22, La. (incomplete rearing — egg); Watson, J. R., 1918: 261-262, Fla. (ex pumpkin bug).

Nezara viridula smaragdula (Fabricius)

Most records ex *Nezara viridula*.

Anon., 1969a: 80, Hawaii (incomplete rearing — egg); Davis and Chong, 1969: 319, Hawaii (as *T. pennipes*, Florida strain); Anon., 1968: 85, Hawaii (incomplete rearing — egg); Miyahira, 1968: 503, Hawaii (incomplete rearing — egg); Miyahira, 1968b: 624, Hawaii; Miyahira and Ah Sam, 1968: 966, Hawaii (incomplete rearing — egg); Anon., 1966: 75, Hawaii (ex *N. viridula smaragdula*); Anon., 1966b: 103, Hawaii (ex *N. viridula smaragdula*, incomplete rearing — egg);

Mitchell and Yoshioka, 1966: 286, Hawaii (incomplete rearing — egg); Miyahira, 1966: 1069, Hawaii (incomplete rearing — egg); Yoshioka and Miyahira, 1966: 231, Hawaii (ex *N. viridula smaragdula*, incomplete rearing — egg); Anon., 1965: 79, Hawaii; Davis, 1965: 8-9, Hawaii (ex southern green stink bug); Davis and Krauss, 1965: 89, Hawaii (ex *N. viridula smaragdula*); Miyahira, 1965: 28, Hawaii (ex *N. viridula smaragdula*); Anon., 1964: 221-222, Hawaii (ex *N. viridula smaragdula*); Davis, 1964b: 414, Hawaii; Kim and Miyahira, 1964: 1155, Hawaii; Davis, 1963: 203, Trinidad (ex *N. viridula smaragdula*); Davis, 1963a: 1324, Hawaii; Kim, 1963: 186-187, Hawaii (incomplete rearing — egg); Kim, 1963a: 535, Hawaii.

Plautia stali Scott

Mau, 1969: 267, Hawaii (incomplete rearing — egg); Anon., 1968a: 316, Hawaii (incomplete rearing — egg).

Thyanta accerra McAtee

Beardsley, 1966: 146, Hawaii (incomplete rearing — egg).

Thyanta perditor (Fabricius)

Wilson, C. E., 1923a: 34, Virgin Islands.

Scutelleridae

Coleotichus blackburniae White

Beardsley, 1965a: 8, Hawaii; Davis, 1965: 8-9, Hawaii; Beardsley, 1964a: 414, Hawaii; Davis, 1964a: 370 (lab.); Davis, 1964b: 414, Hawaii; Perkins, 1963: 1416, Hawaii (incomplete rearing — egg).

STAGE PARASITIZED: Nymph, adult.

IMMATURES: Egg — Hayslip, Genung, Kelsheimer and Wilson, 1953: (descr., fig. 30); Capelouto, 1949: (unnumbered fig. on p. 32); Beard, 1940: (fig. 1); Beard, 1940a: (descr., fig. 12, pl. 1E); Watson, J. R., and Berger, 1932: (fig. 65); Worthley, 1924a: (descr., fig. 16 on pl. 3); Drake, 1920: (descr., fig. 21); Jones, T. H., 1918: (descr.); Weed and Conradi, 1902: (descr.).

First instar larva — Beard, 1940a: (descr., figs. 10-13).

Second instar larva — Beard, 1940a: (descr., figs. 10, 13).

Third instar larva — Beard, 1940a: (descr., figs. 11, 13, pl. 1F); Worthley, 1924a: (descr., figs. 17-18 on pl. 3); Drake, 1920: (descr., fig. 23).

Pupa — Beard, 1940a: (descr.); Worthley, 1924a: (descr., figs. 19-21 on pl. 3); Greene, 1922: (descr., fig. 22 on pl. 5);

Drake, 1920: (descr., fig. 24); Jones, T. H., 1918: (descr.);
Weed and Conradi, 1902: (descr.).

BIOLOGY: Yonke and Medler, 1969: (176-177, 183-184);
Miyahira and Ah Sam, 1968: (966); Shahjahan, 1968:
(1088-1090); Davis, 1963a: (1324); Hayslip, Genung,
Kelsheimer and Wilson, 1953: (47); Capelouto, 1949:
(31-33); Bratley, 1943: (6); Beard, 1940: (269-272); Beard,
1940a: (607, 624-672); Beard, 1936: (335-336); Watson, J.
R., and Berger, 1932: (101-103, 132); Worthley, 1924:
(7-16); Worthley, 1923: (77-79); Drake, 1920: (67-74);
Jones, T. H., 1918: (22); Weed and Conradi, 1902: (20-21).

Trichopoda pennipes (Fabricius) probably
HOST: HEMIPTERA (HETEROPTERA)

Coreidae

Anasa armigera (Say)
Chittenden, 1899: 34.

HOST: ORTHOPTERA

Mantidae

Tenodera australasiae (Leach)

Beardsley, 1965: 3, Hawaii (incomplete rearing — egg);

Beardsley, 1964: 150, Hawaii (incomplete rearing — egg).

STAGE PARASITIZED: Adult (Hemiptera — Heteroptera,
Orthoptera).

Trichopoda pennipes (Fabricius) complex
HOST: HEMIPTERA (HETEROPTERA)

Coreidae

Archimerus alternatus (Say)

Patton, 1958: 84, Fla.

Leptoglossus phyllopus (Linnaeus)

Patton, 1958: 84, Fla.

STAGE PARASITIZED: Adult.

Trichopoda pilipes (Fabricius)
All records as *Trichopoda pennipes pilipes*.
HOST: HEMIPTERA (HETEROPTERA)

Pentatomidae

Nezara viridula smaragdula (Fabricius)
Most records ex *Nezara viridula*.

Anon., 1969a: 80, Hawaii (incomplete rearing — egg); Davis and Chong, 1969: 319, Hawaii; Kawamura, 1969: 726, Hawaii (incomplete rearing — egg); Kawamura and Funasaki, 1969: 632, Hawaii (incomplete rearing — egg); Wong, 1969: 217, Hawaii (incomplete rearing — egg); Yamamoto, 1969: 301, Hawaii (incomplete rearing — egg); Miyahira, 1968: 503, Hawaii (incomplete rearing — egg); Miyahira and Ah Sam, 1968: 966, Hawaii (incomplete rearing — egg); Yoshioka, 1968: 652, Hawaii (incomplete rearing — egg); Funasaki, 1967: 450, Hawaii (incomplete rearing — egg); "Jackson et al.," 1967: 822, Hawaii (incomplete rearing — egg); Miyahira, 1967: 473, Hawaii (incomplete rearing — egg); Nakao and Miyahira, 1967: 345, Hawaii (incomplete rearing — egg); Yoshioka and Miyahira, 1966a: 444, Hawaii (ex *N. viridula smaragdula*); Davis and Krauss, 1965: 89, Hawaii (ex *N. viridula smaragdula*); Davis, 1964: 347, Fla., Hawaii (ex *N. viridula smaragdula*); Davis, 1964a: 369-375, Fla., Hawaii (ex *N. viridula smaragdula*); Davis and Krauss, 1964: 393, Hawaii (ex *N. viridula smaragdula*); Kim, 1964: 328, Hawaii (ex *N. viridula smaragdula*); Kim, 1964a: 333, Hawaii; Mitchell, 1964: 345, Hawaii (ex southern green stink bug); Davis and Krauss, 1963: 247, Hawaii, Trinidad (ex *N. viridula smaragdula*).

Scutelleridae

Coleotichus blackburniae White

Perkins, 1964: 351, Hawaii (incomplete rearing — egg).

STAGE PARASITIZED: Adult.

BIOLOGY: Kawamura and Funasaki, 1969: (632); Davis, 1964a: (369-375); Kim, 1964: (328); Mitchell, 1964: (345); Perkins, 1964: (351).

Trichopoda plumipes (Fabricius)

HOST: HEMIPTERA (HETEROPTERA)

Coreidae

Acanthocephala terminalis (Dallas)

Fattig, 1949: 34, Ga.

Leptoglossus phyllopus (Linnaeus)

Fattig, 1949: 34, Ga.

HOST: ORTHOPTERA

Acrididae

Dissosteira pictipennis (Bruner)

All records ex *Dissosteira venusta* Stål.

Essig, 1926: 578, Calif.; Coquillett, 1897: 21, 23, Calif.

Unidentified Acrididae

Coquillett, 1897: 21, 23.

STAGE PARASITIZED: Adult (Orthoptera).

Trichopoda plumipes (Fabricius) probably

HOST: ORTHOPTERA

Acrididae

Dissosteira pictipennis (Bruner)

Brauer and Bergenstamm, 1895: 579, 591, N. Amer. (as *T.* (?
trifasciata Loew) ex *D. cornuta* Stål).

Trichopoda radiata, see *Trichopoda lanipes*

Trichopoda subdivisa (Townsend)

HOST: ORTHOPTERA

Acrididae

Dissosteira pictipennis (Bruner)

Townsend, 1908: 133, Calif. (as *Polistomyia* ex *D. venusta*
Stål).

Trichopoda trifasciata probably, see *Trichopoda plumipes* probably

Trichopoda sp., see *Trichopoda pennipes* or *Trichopoda* spp.

Trichopoda spp. (including *T. pennipes* and/or *T. pilipes*)

HOST: HEMIPTERA (HETEROPTERA)

Pentatomidae

Nezara viridula smaragdula (Fabricius)

Miyahira and Ah Sam, 1969: 72, Hawaii (as *T. pennipes*
and/or *T. pennipes* var. *pilipes* ex *N. viridula*, incomplete
rearing — egg).

STAGE PARASITIZED: Adult.

Trichopoda spp.

HOST: HEMIPTERA (HETEROPTERA)

Coreidae

Acanthocephala terminalis (Dallas)

Shannon, 1914: 182, D.C. (ex *Metapodius*); Shannon, 1914:
182, Va. (ex *Metapodius instabilis* Uhler).

Anasa spp.

Townsend, 1913: 148 (as *Trichopodopsis*); Townsend, 1908: 135.

Leptoglossus spp.

Townsend, 1913: 148 (as *Trichopodopsis*); Townsend, 1908: 135.

Pentatomidae

Chlorochroa spp.

Cook, 1889: 96 (as *Phasia* nearly allied to *T. pennipes* (Fabricius) ex "*Pentatomia*").

Nezara viridula (Linnaeus)

All records ex southern green stink bug.

Bratley, 1943: 6, Fla. (as feather-legged fly); Bratley, 1938: 13-14, Fla. (as feather-legged fly).

Nezara viridula smaragdula (Fabricius)

Anon., 1966e: 995, Hawaii.

Plant bug(-s)

Bratley, 1938: 13-14, Fla. (as feather-legged fly).

HOST: ORTHOPTERA

Acrididae

Dissosteira spp.

Townsend, 1913: 148 (as *Trichiopoda*).

STAGE PARASITIZED: Adult (Hemiptera-Heteroptera).

IMMATURES: Egg — Bratley, 1938: (descr.); Townsend, 1913: (descr.).

BIOLOGY: Bratley, 1943: (6); Bratley, 1938: (13-14); Shannon, 1914: (182).

"*Trichopodes*" *pennipes*, see *Trichopoda pennipes*

Trichopodopsis sp., see *Trichopoda* spp.

Trisisyropa platysamiae, see *Winthemia cecropia*

Tsugaea Hall (Goniinae: Eryciini)

Tsugaea nox Hall

HOST: HYMENOPTERA

Diprionidae

Neodiprion sp. near *scutellatis* Rohwer

Smith, 1962: 189, Idaho (ex *N.* near "*scutellaris*").

Neodiprion tsugae Middleton

Anon., 1953c: 137, B.C.; Furniss and Dowden, 1941: 49-50, Oreg.; Hall, 1939: 242, Oreg. (ex "*Neodipiron*").

STAGE PARASITIZED: Larva, pupa.

BIOLOGY: Furniss and Dowden, 1941: (49-50).

Uramya Robineau-Desvoidy (Dexiinae: Uramyini)

Uramya acuminata, see *Uramya halisidotae*

Uramya halisidotae (Townsend)

HOST: LEPIDOPTERA

Arctiidae

Halisidota argentata Packard

Silver, 1958: 76, B.C. (as *Uromacquartia*); Guppy, R., 1950: 4, B.C. (as *Uromacquartia*); Prebble, 1942a: 165, B.C. (as *Uramya*); Aldrich, 1921: 86, Calif., Oreg. (as *Uramya*); Cole, F. R., and Lovett, 1921: 300, Oreg. (as *Uramya acuminata* (Bigot)).

Lasiocampidae

Malacosoma sp.

Aldrich, 1921: 86, Calif. (as *Uramya*).

STAGE PARASITIZED: Larva.

IMMATURES: Pupa — Greene, 1922: (descr., fig. 96 on pl. 20).

BIOLOGY: Guppy, R., 1950: (4).

Uramya acuminata, see *Uramya halisidotae*

Uramya halisidotae, see *Uramya halisidotae*

Uromacquartia halisidotae, see *Uramya halisidotae*

Urophyllopsis Townsend (Goniinae: Blondeliini)

Urophyllopsis retiniae (Coquillett)

Most records as *Admontia retiniae*.

HOST: LEPIDOPTERA

Olethreutidae

Rhyacionia pasadenana (Kearfott)

Rawlings, 1960: 56, Calif. (ex *Evetria*); Lange, 1937: 29, Calif.

Rhyacionia spp.

All records ex *Retinia* sp.

Smith, H. E., 1912: 127 (as *Hyperecteina* "retinae");
Coquillett, 1897: 9, 27, Calif.

Rhyacionia sp. probably

Essig, 1926: 579, Calif. (ex *Evetria*).

STAGE PARASITIZED: Larva, pupa.

IMMATURES: Third instar larva — Lange, 1937: (descr.).

Pupa — Lange, 1937: (descr.).

BIOLOGY: Lange, 1937: (29).

Varichaeta aldrichi, see *Mericia* spp.

Varichaeta sp., see *Mericia* spp. probably

Vibrissotheresia Reinhard (Proseninae: Theresiini)

Vibrissotheresia pechumani Reinhard

HOST: DIPTERA

Tabanidae

Tabanus reinwardtii Wiedemann probably

Tashiro and Schwardt, 1953: 681, N.Y.

STAGE PARASITIZED: Larva.

"*Viviana*" sp., see *Sitophaga georgiae*

Viviania georgiae, see *Sitophaga georgiae*

Viviania lachnosternae, see *Sitophaga georgiae*

Viviania nocturnalis, see *Sitophaga nocturnalis*

Viviania sp., see *Sitophaga* spp.

Voria Robineau-Desvoidy (Tachininae: Voriini)

Voria aurifrons (Townsend)

HOST: LEPIDOPTERA

Noctuidae

Autographa californica (Speyer) probably

Anon., 1961a: 299, B.C.; MacNay, 1961: 274, B.C. (ex alfalfa looper).

Heliothis zea (Boddie)

Bottrell, Young, Price and Adams, 1968: 1054, Okla.

STAGE PARASITIZED: Larva.

Voria ruralis (Fallén)

HOST: LEPIDOPTERA

Noctuidae

Anticarsia gemmatalis Hübner

All records ex *Anticarsia "gemmatilis."*

*Wolcott, 1951a: 481, P.R. (Plank, 1945); Plank, 1945: 26, P.R.

Autographa californica (Speyer)

Clancy, 1969: 1079, Calif.; Jackson, C. G., Butler and Bryan, 1969: 70, Ariz.; Brubaker, 1968: 308, Ariz.; Puttarudriah, 1953: 723, Calif.; Spencer, 1936: 28, B.C.; Essig, 1926: 580, Wash. (as *Plagia americana* Wulp); Childs, 1915: 185, Oreg. (as *Plagia americana* ex *Plusia*);

Hyslop, 1912: 116, probably Wash. (as *Plagia americana* ex *A. gamma californica*).

Autographa sp.

Vickery, 1925: 140, Tex. (as *Plagia americana*).

Autoplusia egena (Guenée)

Clancy, 1969: 1079, Calif.

Peridroma saucia (Hübner)

Jackson, C. G., Butler and Bryan, 1969: 69, Ariz.;
Montgomery, 1933: 186, Ind. (ex *Lycophotia*).

Plusia aereoides Grote

Tothill, 1913: 74, Ont. (as *Plagia americana*).

Spodoptera frugiperda (J. E. Smith)

Jackson, C. G., Butler and Bryan, 1969: 69, Ariz.; Butler,
1958: 562, Ariz. (ex *Laphygma*).

Trichoplusia ni (Hübner)

Bryan, D. E., Jackson and Stoner, 1969: 1, 6-7 (lab.); Clancy,
1969: 1079-1082, Calif.; Jackson, C. G., Butler and
Bryan, 1969: 69-70 (lab.); Brubaker, 1968: 306-309,
Ariz.; Bosch and Hagen, 1966: 25, Calif.; Oatman, 1966:
1136-1138, Calif.; Anon., 1962b: 269, Ariz. (ex cabbage
looper); Anon., 1961b: 18, Ariz.; Anon., 1961e: 1049,
Ariz. (ex cabbage looper); Bibby, 1961: 328, Ariz.;
Gibson, W. W., and Carrillo S., 1959: 184, Mexico (Son.);
Butler, 1958: 562, Ariz.; McKinney, 1944: 14, Ariz. (ex
Autographa brassicae Riley); Schaffner and Griswold,
1934: 69, 108 (ex *Autographa brassicae*).

Phalaenid moth(-s)

Tilden, 1951: 164-165, Calif.

Unidentified Noctuidae

Schaffner and Griswold, 1934: 72, 108.

Noctuidae and/or Pyralidae

Noctuid and/or pyralid spp. (including *Scotogramma trifolii*
(Rottenburg) and/or *Loxostege sticticalis* (Linnaeus))
Anon., 1956d: 1044, Idaho (ex clover cutworm and/or beet
webworm).

Pyralidae and/or Noctuidae,
see Noctuidae and/or Pyralidae

Unidentified Lepidoptera

Green looper(-s)

Butler, 1958: 562, Ariz. (ex green looper on mesquite).

STAGE PARASITIZED: Larva, pupa.

IMMATURES: Egg — Brubaker, 1968: (descr.).

Pupa — Greene, 1922: (descr., fig. 49 on pl. 11).

BIOLOGY: Bryan, D. E., Jackson and Stoner, 1969: (1, 6-7);
Clancy, 1969: (1079-1082); Jackson, C. G., Butler and
Bryan, 1969: (69-70); Brubaker, 1968: (306-309); Schaffner
and Griswold, 1934: (108); Hyslop, 1912: (116).*Voria ruralis* (Fallén) probably

HOST: LEPIDOPTERA

Noctuidae

Autographa gamma (Linnaeus)Brauer and Bergenstamm, 1895: 570, 603 (as *Plagia* ex
Plusia).*Trichoplusia ni* (Hübner)

Anon., 1960e: 1115, Ariz. (ex cabbage looper).

STAGE PARASITIZED: Larva.

BIOLOGY: Anon., 1960e: (1115).

Wagneria Robineau-Desvoidy (Tachininae: Voriini)*Wagneria helymus*, see *Periscepsia helymus**Wagneria laevigata*, see *Periscepsia laevigata**Wagneria major* Curran

HOST: LEPIDOPTERA

Noctuidae

Choephora fungorum Grote and Robinson

Reinhard, 1955: 55, Miss.

Wagneria rohweri, see *Periscepsia rohweri**Wagneria sequax*, see *Periscepsia helymus**Wagneria vernata* West

HOST: LEPIDOPTERA

Noctuidae

Protorthodes oviducta (Guenée)

Reinhard, 1955: 54, N.J.

Xylena sp.

Reinhard, 1955: 54, N.J.

Wagneria vernata West probably
HOST: LEPIDOPTERA

Noctuidae

Orthosia hibisci (Guenée)

Schaffner and Griswold, 1934: 46, 124, Mass. (ex *O. hibisci*
form normal *insciens* Walker).

BIOLOGY: Schaffner and Griswold, 1934: (46, 124).

Wagneria sp.

HOST: LEPIDOPTERA

Noctuidae

Choephora fungorum Grote and Robinson

Allen, 1929: 689, Miss. (ex *Epsilia*).

"*Winthema*" *quadripustulata*, see *Winthemia quadripustulata*

Winthemia Robineau-Desvoidy (Goniinae: Winthemiini)

Winthemia abdominalis (Townsend)

HOST: LEPIDOPTERA

Noctuidae

Spodoptera frugiperda (J. E. Smith)

Fattig, 1949: 35, Ga. (ex *Laphygma*).

Winthemia amoena, see *Omotoma fumiferanae*

Winthemia borealis Reinhard

HOST: LEPIDOPTERA

Noctuidae

Scoliopteryx libatrix (Linnaeus)

Raizenne, 1952: 74, Ont.

Winthemia cecropia (Riley)

HOST: LEPIDOPTERA

Saturniidae

Actias luna (Linnaeus)

Raizenne, 1952: 15, Ont.

Antheraea polyphemus (Cramer)

Hagen, 1875: 205, New England (as *Exorista "leucaniae* var.
cecropiae" ex *polyphemus*).

Platysamia cecropia (Linnaeus)

Collins, M. M., and Weast, 1961: 94 (as *Exorista* "cecropiae" ex *Hyalophora*); Raizenne, 1952: 14, Ont. (as *Trisisyropha platysamiae* (Townsend) ex *Hyalophora*); Sabrosky, 1948: 64, Mich., Pa. (ex *Samia*); Reinhard, 1931: 35, Ill. (as *W. cecropiae* Reinhard); Reinhard 1931: 54, N.Y. (as *W. platysamiae*); Packard, 1914: 269 (as *Exorista* "cecropiae" ex *Samia*); *Brauer and Bergenstamm, 1895: 554, 605, N. Amer. (as *Exorista platysamiae* ex *Saturnia*, Townsend, 1892a); Townsend, 1892a: 288, N.Y. (as *Exorista platysamiae*); Hagen, 1875: 205, New England (as *Exorista* "leucaniae var. cecropiae" ex cecropia); Riley, 1872: 108-109, Ill. (as *Exorista* "leucaniae var. cecropiae" ex *Attacus*); Riley, 1870a: 101, Ill. (as *Exorista* ex *Attacus*).

STAGE PARASITIZED: Larva.

IMMATURES: Third instar larva — Riley, 1872: (descr.); Riley, 1870a: (descr.).

BIOLOGY: Riley, 1872: (108-109); Riley, 1870a: (101).

Winthemia cecropiae, see *Winthemia cecropia*

Winthemia cilitibia, see *Winthemia venusta*

Winthemia citheroniae Sabrosky

HOST: LEPIDOPTERA

Citheroniidae

Citheronia regalis Fabricius

Patton, 1958: 84, Fla.; Sabrosky, 1948: 65, Fla., Md.

Eacles imperialis (Drury)

Patton, 1958: 84, Fla.; Patton, 1957: 70 (ex "*Eacles imperialis*").

STAGE PARASITIZED: Pupa.

BIOLOGY: Patton, 1958: (84); Patton, 1957: (70).

Winthemia datanae (Townsend)

HOST: LEPIDOPTERA

Arctiidae

Diacrisia virginica (Fabricius)

Schaffner and Griswold, 1934: 36, 123 (as *W. datanae* (var. C)).

Estigmene acrea (Drury)

Schaffner and Griswold, 1934: 38, 123 (as *W. datanae* (var. C)).

Euchaetias egle (Drury)

Schaffner and Griswold, 1934: 39, 123 (as *W. datanae* (var. C)).

Unidentified Arctiidae

Schaffner and Griswold, 1934: 40-41, 123 (as *W. datanae* (var. C)).

Citheroniidae

Anisota rubicunda (Fabricius)

Schaffner and Griswold, 1934: 30-31, 123 (as *W. datanae* (var. B)).

Anisota senatoria J. E. Smith

Hitchcock, 1961: 963, Conn.; Schaffner and Griswold, 1934: 29-30, 123 (as *W. datanae* (var. B)).

Lasiocampidae

Malacosoma disstria Hübner

Sippell, 1961: 436, Ont.

Lymantriidae

Hemerocampa leucostigma (J. E. Smith)

Schaffner and Griswold, 1934: 85, 123 (as *W. datanae* (var. C)).

Noctuidae

Pseudaletia unipuncta (Haworth)

Boyes and Wilkes, 1953: 126 (ex *P. "unipunctata"*).

Unidentified Catocalinae

Schaffner and Griswold, 1934: 67, 123 (as *W. datanae* (var. A)).

Notodontidae

Datana angusii Grote and Robinson

Schaffner and Griswold, 1934: 74-75, 122.

Datana integerrima Grote and Robinson

Schaffner and Griswold, 1934: 76, 122.

Datana ministra (Drury)

Patton, 1958: 84, Fla.; Fattig, 1949: 35, Ga.; *Procter, 1946: 431, Maine (Procter, 1938); Procter, 1938: 372,

Maine (as *W. "detana"* ex "*Detana*"); Schaffner and Griswold, 1934: 74, 122.

Datana perspicua Grote and Robinson
Schaffner and Griswold, 1934: 75, 122.

Datana spp.

Reinhard, 1931: 30; *Brauer and Bergenstamm, 1895: 553, 590, N. Amer. (as *Exorista*), Townsend, 1892a); Townsend, 1892a: 288-289, N.Y. (as *Exorista*).

Dicentria lignicolor (Walker)

Schaffner and Griswold, 1934: 80, 123 (as *W. datanae* (var. A) ex *Ianassa*).

Hyperaeschra stragula (Grote)

Schaffner and Griswold, 1934: 76-77, 123, Conn., Mass. (as *W. datanae* (var. D)).

Pheosia rimosa Packard

Raizenne, 1952: 83, Ont.; Schaffner and Griswold, 1934: 77, 122.

Schizura badia (Packard)

Schaffner and Griswold, 1934: 81, 123, Mass. (as *W. datanae* (var. A)).

Schizura concinna (J. E. Smith)

Schaffner and Griswold, 1934: 81, 123 (as *W. datanae* (var. A)).

Schizura ipomaeae Doubleday

Schaffner and Griswold, 1934: 80, 123 (as *W. datanae* (var. A) ex *S. "ipomoeae"*).

Schizura leptinoides (Grote)

Schaffner and Griswold, 1934: 82, 123 (as *W. datanae* (var. A)).

Schizura unicornis (J. E. Smith)

Schaffner and Griswold, 1934: 81-82, 123 (as *W. datanae* (var. A)).

Saturniidae

Platysamia cecropia (Linnaeus)

Most records ex *Samia cecropia*.

Marsh, F. L., 1941: 336, Ill.; Marsh, F. L., 1938: 138-140, Ill. (ex *cecropia*); Marsh, F. L., 1937: 107-108, Ill.

Sphingidae

Ceratonia catalpae (Boisduval)

Schaffner and Griswold, 1934: 17, 122, N.J.

Cressonia juglandis (J. E. Smith)

Schaffner and Griswold, 1934: 22, 123 (as *W. datanae* (var. B)).

Paonias excaecata (J. E. Smith)

Schaffner and Griswold, 1934: 21, 123 (as *W. datanae* (var. B)).

Sphinx drupiferarum J. E. Smith

Schaffner and Griswold, 1934: 20, 123 (as *W. datanae* (var. B) ex *Hyloicus*).

Sphinx gordius Cramer

Schaffner and Griswold, 1934: 19, 123 (as *W. datanae* (var. B) ex *Hyloicus*).

STAGE PARASITIZED: Larva.

IMMATURES: Egg — Marsh, F.L., 1938: (descr.).

BIOLOGY: Patton, 1958: (84); Marsh, F. L., 1938: (138-140); Schaffner and Griswold, 1934: (76-77, 122-123).

Winthemia datanae (vars. A, B, C, D), see *Winthemia datanae*

Winthemia datanae (Townsend) probably

HOST: LEPIDOPTERA

Arctiidae

Isia isabella (J. E. Smith)

Schaffner and Griswold, 1934: 36-37, 123.

Noctuidae

Acronicta impleta Walker

Schaffner and Griswold, 1934: 55, 122 (ex *Acronycta*).

Saturniidae

Samia cynthia advena (Packard)

Schaffner and Griswold, 1934: 26, 123 (ex *S. walkeri advena*).

Winthemia sp. near *datanae* (Townsend)

HOST: LEPIDOPTERA

Citheroniidae

Anisota senatoria J. E. Smith

Hitchcock, 1961: 963, Conn.

STAGE PARASITIZED: Larva.

Winthemia deilephilae (Osten Sacken)Most records as *Tachina deilephilae*.

HOST: LEPIDOPTERA

Sphingidae

Hyles lineata (Fabricius)Reinhard, 1931: 34 (as *Winthemia ex Celerio*); *Brauer and Bergenstamm, 1895: 575, 607 (ex *Sphinx*, Osten Sacken, 1887); Osten Sacken, 1887: 165 (ex *Deilephila*).**Winthemia deilephilae** (Osten Sacken) probably

HOST: LEPIDOPTERA

Sphingidae

Sphinx sp.Brauer and Bergenstamm, 1895: 548, 606, N. Amer. (as *Chaetolyga*).**Winthemia "detana,"** see *Winthemia datanae***"Winthemia diversoides Baranov"**

HOST: LEPIDOPTERA

Geometridae

Anacampodes fragilaria (Grossbeck)*Vide Chong, 1965: 15, Hawaii (ex "*Anacampodes*," probable parasite, Nakao, 1964); Nakao, 1964: 333 (as *Eucelatoria armigera* (Coquillett), lab.).

Noctuidae

Achaea janata (Linnaeus)*Vide Chong, 1965: 15, Hawaii (probable parasite, Nakao, 1964); Nakao, 1964: 333 (as *Eucelatoria armigera*, lab.).

STAGE PARASITIZED: Larva.

Winthemia fumiferanae, see *Omotoma fumiferanae***Winthemia infesta** (Williston)Most records as *Exorista infesta*.

HOST: LEPIDOPTERA

Noctuidae

Spodoptera frugiperda (J. E. Smith)All records ex *Laphygma frugiperda*.Reinhard, 1931: 54, Ill. (as *Winthemia*); Forbes, 1885: 65, Ill.

Armyworm(-s)

Chittenden, 1901a: 35.

Winthemia intermedia Reinhard

HOST: LEPIDOPTERA

Noctuidae

Pseudoplusia includens (Walker)Fattig, 1949: 35, Ga. (ex *Autographa rogationis* Guenée).*Winthemia leucanae* (Kirkpatrick)All but four records as "*leucaniae*."

HOST: LEPIDOPTERA

Noctuidae

Euthisanotia grata (Fabricius)All records ex *Eudryas grata*.Brauer and Bergenstamm, 1895: 554, 592, Ont., N. Amer. (as *Exorista*); Saunders, 1875: 43-44, Canada (as *Exorista* "*leucania*"); Saunders, 1871: 34-35, Ont. (as *Exorista* "*leucania*").*Peridroma saucia* (Hübner)Riley, 1870: 50 (as *Exorista* ex *Agrotis inermis* (Harris)).*Pseudaletia unipuncta* (Haworth)Most records ex *Leucania unipuncta*.Reinhard, 1931: 53, Ohio; Fyles, 1897: 102, Que. (as *Nemoraea*); Lugger, 1897: 16, Minn. (as *Exorista*); Panton, 1897: 51, fig. 50, Ont. (as *Nemoraea*); Lintner, 1896: 56, N.Y. (as *Nemoraea*); Howard, 1894: 4-5, fig. 3, Va. (as *Nemoraea*); Webster, 1890: 46, Ind. (as *Nemoraea*); Bruner, 1889: 32, Nebr. (as *Exorista*); Lintner, 1883: 146, fig. 37 (as *Nemoraea*); Reed, 1883: 52, fig. 48 (as *Nemoraea*); Riley, 1883: 126 (as *Nemoraea leucanae*); Riley, Packard and Thomas, 1883: 105, 135, Ill. (as *Nemoraea*); Coquillett, 1882b: 52-54, 63, Ill. (as *Exorista*); Thomas, C., 1881: 36-38, Ill. (as *Nemoraea*); Riley, 1876: 53 (as *Exorista*); Riley, 1871: 116 (as *Exorista* ex true army-worm); Riley, 1870c: 266 (as *Exorista* ex army-worm); Kirkpatrick, 1861: 358, Ohio (as *Exorista leucanae* ex *L. extranea* Guenée).*Spodoptera frugiperda* (J. E. Smith)Most records ex *Laphygma frugiperda*.

Hinds and Dew, 1915: 88-90, Ala. (as "*Nemorea*"); Dew, 1913: 265, [Ala.] (as "*Nemorea*"); Riley, 1871: 116, Mo. (as *Exorista* ex *Prodenia autumnalis* Riley).

Armyworm(-s)

Riley, 1871: 129 (as *Exorista*).

Cutworm(-s)

Lintner, 1893: 238, N.Y. (as *Nemoraea*).

Pyralidae

Pyralis farinalis Linnaeus

Chittenden, 1897: 42-43, Md. (as *Carcelia*, probable host).

Saturniidae

Platysamia cecropia (Linnaeus)

Riley, 1870: 50 (as *Exorista* ex common large Cecropia worm).

Sphingidae

Manduca quinquemaculata (Haworth)

Brauer and Bergenstamm, 1895: 547, 604 (as *Chaetolyga* ex *Protoparce celeus* Hübner).

STAGE PARASITIZED: Larva, pupa.

IMMATURES: Egg — Panton, 1897: (fig. 50); Howard, 1894: (descr., fig. 3).

Larva — Lintner, 1883: (fig. 37).

Third instar larva — Panton, 1897: (fig. 50); Howard, 1894: (fig. 3); Reed, 1883: (fig. 48); Saunders, 1875: (descr.).

Pupa — Panton, 1897: (fig. 50); Howard, 1894: (fig. 3); Lintner, 1883: (fig. 37); Reed, 1883: (fig. 48); Thomas, C., 1881: (descr.); Saunders, 1875: (descr.); Saunders, 1871: (descr.).

BIOLOGY: Hinds and Dew, 1915: (88-90); Fyles, 1897: (102); Panton, 1897: (51); Howard, 1894: (4-5); Riley, 1883: (126); Riley, Packard and Thomas, 1883: (105, 135); Thomas, C., 1881: (36-38); Saunders, 1875: (43-44); Saunders, 1871: (34-35).

Winthemia leucanae (Kirkpatrick) lacking red at tip

All records as *Exorista* "*leucaniae*" lacking red at tip.

HOST: LEPIDOPTERA

Lasiocampidae

Malacosoma disstria Hübner

All records ex *Clisiocampa sylvatica* (Harris).

Riley, 1871: 129; Riley, 1870c: 266, Mo.

Noctuidae

Peridroma saucia (Hübner)

All records ex *Agrotis inermis* (Harris).

Riley, 1871: 129; Riley, 1870c: 266.

Agrotidian(-s)

Riley, 1871: 129; Riley, 1870c: 266.

Notodontidae

Datana ministra (Drury)

Riley, 1871: 129; Riley, 1870c: 266.

Saturniidae

Platysamia cecropia (Linnaeus)

All records ex *Attacus cecropia*.

Riley, 1871: 129; Riley, 1870c: 266 (ex *A. "cecropiae"*).

STAGE PARASITIZED: Larva.

Winthemia leucanae (Kirkpatrick) probably

HOST: LEPIDOPTERA

Noctuidae

Euthisanotia grata (Fabricius)

Saunders, 1876: 38-39 (as *Exorista "leucania"* ex *Eudryas*).

Pseudaletia unipuncta (Haworth)

Riley, Packard and Thomas, 1883: 96, Md. (as *Nemoraea "leucaniae"* ex *Leucania*).

STAGE PARASITIZED: Larva.

IMMATURES: Pupa — Saunders, 1876: (descr.).

BIOLOGY: Saunders, 1876: (38-39).

Winthemia militaris (Walsh)

HOST: LEPIDOPTERA

Hesperiidae

Thymelicus lineola (Ochsenheimer)

Arthur, 1962: 1085, Ont.

Noctuidae

Pseudaletia unipuncta (Haworth)

Most records ex *Leucania unipuncta*.

Reinhard, 1931: 54, Ill. (ex armyworm); Wood, J. G., 1874: 770, U.S. (as *Exorista* ex "*Leucaria unipunctata*"); Packard, 1869: 407-408 (as *Tachina*); Walsh and Riley, 1869a: 216-217, Mo. (as *Exorista* ex *L. "unipunctata"*); Walsh, 1867: 114, N.Y. (as *Exorista*); Walsh, 1861: 361-362, 367, fig. 5, Ill., Md., Ohio (as *Senometopia*).

Armyworm(-s)

Johannsen, Patch, Morse and Lewis, 1911: 376 (as *Tachina*); Packard, 1890: 16 (as *Tachina*).

Saturniidae

Platysamia cecropia (Linnaeus)

Brauer and Bergenstamm, 1895: 554, 605 (as *Exorista* ex *Saturnia*); Gentry, 1877: 50, Pa. (as *Exorista*); Saunders, 1875a: 25-26, Ont. (as *Exorista* ex *Attacus*); Wood, J. E., 1874: 770-771, U.S. (as *Exorista* ex *Attacus*).

HOST: ORTHOPTERA

Acrididae

Grasshopper(-s)

Johannsen, Patch, Morse and Lewis, 1911: 376 (as *Tachina*, questionable host).

STAGE PARASITIZED: Larva (Lepidoptera), pupa (Lepidoptera).

IMMATURES: Egg — Saunders, 1875a: (descr.); Walsh and Riley, 1869a: (descr.).

Third instar larva — Saunders, 1875a: (descr.).

Pupa — Walsh, 1861: (descr., fig. 5).

BIOLOGY: Johannsen, Patch, Morse and Lewis, 1911: (376); Saunders, 1875a: (25-26); Packard, 1869: (407-408); Walsh, 1861: (361-362, 367).

Winthemia militaris (Walsh) probably

HOST: LEPIDOPTERA

Noctuidae

Pseudaletia unipuncta (Haworth)

Pond, 1960: 665, N.B.

STAGE PARASITIZED: Larva.
Winthemia sp. near *militaris* (Walsh)
 HOST: LEPIDOPTERA

Lasiocampidae

Malacosoma disstria Hübner
 Walsh, 1867: 113-114, N.Y. (as *Exorista ex Clisiocampa sylvatica* (Harris)).

Noctuidae

Pseudaletia unipuncta (Haworth)
 Maxson, 1920: 81, Colo. (ex *Cirphis*).
 STAGE PARASITIZED: Larva.
Winthemia obscura, see *Buquetia obscura*
Winthemia occidentis Reinhard
 HOST: LEPIDOPTERA

Geometridae

Lambdina fiscellaria (Guenée)
 Anon., 1952b: 180, 182, 186, B.C.; Anon., 1950c: 241, 243, 249-250, B.C.; Anon., 1949: 276, 280, 286, B.C.
Lambdina fiscellaria fiscellaria (Guenée)
 Carroll, 1956: 594, B.C.
Lambdina fiscellaria lugubrosa (Hulst)
 All records ex western hemlock looper.
 Arnaud, 1963: 131, B.C.; Reinhard, 1931: 23, B.C.
Lambdina fiscellaria somniaria (Hulst)
 Boyes and Wilkes, 1953: 126, 157, B.C. (ex *L. fiscellaria* "somniaria").
 STAGE PARASITIZED: Pupa.
Winthemia ostensackenii (Kirkpatrick)
 HOST: LEPIDOPTERA

Noctuidae

Pseudaletia unipuncta (Haworth)
 Reinhard, 1931: 53, Ohio (ex *Leucania*); Kirkpatrick, 1861: 358, Ohio (as *Exorista ex Leucania extranea* Guenée).
 STAGE PARASITIZED: Larva.
Winthemia platysamiae, see *Winthemia cecropia*
Winthemia "pustulata," see *Winthemia quadripustulata*

Winthemia "quadripunctata," see *Winthemia quadripustulata*

Winthemia quadripustulata (Fabricius)

HOST: COLEOPTERA

Curculionidae

Curculio proboscideus (Fabricius)

Brooks, F. E., and Cotton, 1929: 15, W. Va.

Unidentified Curculionidae

Parker, 1916: 236, Md.

HOST: HYMENOPTERA

Diprionidae

Diprion similis (Hartig)

Webber, 1932: 206, Mass. (ex *D. "simile"*).

HOST: LEPIDOPTERA

Arctiidae

Halisidota tessellaris (J. E. Smith)

Allen, 1925: 7, figs. 2-7; Johnson, 1910: 782, N.J. (ex *H. "tessellata"*); Coquillett, 1897: 22, 25, N.Y. (as *W. 4-pustulata* ex *H. "tessellata"*).

White woolly bear(-s)

Rowe, 1933: 125, Ill.

Bombycidae

Unidentified Bombycidae

Coquillett, 1897: 21, 24 (as *W. 4-pustulata*).

Citheroniidae

Anisota senatoria J. E. Smith

Webber and Schaffner, 1926: 25, New England.

Geometridae

Cingilia catenaria (Cramer)

Franklin, 1950: 57.

Ematurga amitaria (Guenée)

Franklin, 1950: 57.

Lymantriidae

Hemerocampa leucostigma (J. E. Smith)

Franklin, 1950: 57; Essig, 1926: 584; Webber and Schaffner, 1926: 12-13, Mass., New England, N.Y., Pa.; Allen, 1925: 7-8, figs. 2-7; Britton, 1917: 110; Johnson, 1910: 782, N.J. (ex *Orgyia*); Coquillett, 1897: 22, 26, D.C. (as *W. 4-pustulata* ex *Orgyia*); Howard, 1897b: 7, 44, D.C. (ex *Orgyia*).

Stilpnotia salicis (Linnaeus)

Burgess and Crossman, 1927: 14, New England.

Tussock moth(-s)

Johnson, 1925: 203, Conn., Maine, Mass., N.H., R.I., Vt.

Noctuidae

Alypia octomaculata (Fabricius)

Essig, 1926: 584; Webber and Schaffner, 1926: 22, New England; Allen, 1925: 6, figs. 2-7; Lochhead, 1919: 184 (as *W. 4-pustulata*); Lowry, Q. S., 1917: 48, Conn.; Lowry, Q. S., 1917a: 122, Conn.; Coquillett, 1897: 21, 23, N.Y. (as *W. 4-pustulata*).

Amathes c-nigrum (Linnaeus)

Most records ex *Agrotis c-nigrum*.

Franklin, 1950: 57 (ex *Amathes*); *Crumb, 1929: 40, 140 (Franklin and Lacroix, 1924); Franklin and Lacroix, 1924: 407-408, Mass.

Caenurgina erechtea (Cramer)

Smith, R. C., 1924: 318, Kans. (ex "*Caenurgia*").

Cucullia convexipennis Grote and Robinson

Tothill, 1913: 75, Ont.

Euxoa ochrogaster (Guenée)

Strickland, 1938: 217, Alta. (ex redbacked cutworm).

Faronta diffusa (Walker)

Most records ex *Meliana albilinea* (Hübner).

Allen, 1925: 6, figs. 2-7; Webster, R. L., 1911: 181, Iowa; Ainslie, 1910: 92, N. Mex. (ex *Heliophila*).

Feltia ducens Walker-*subgothica* (Haworth) complex

Crumb, 1929: 40, 68.

Feltia herilis (Grote)

Allen, 1925: 6, figs. 2-7; Coquillett, 1897: 21, 25, Mo. (as *W. 4-pustulata*).

Heliothis zea (Boddie)

Most records ex *Heliothis obsoleta* (Fabricius).

Bosch and Hagen, 1966: 25, Calif. (ex *H. zea*); Fattig, 1949: 35-36 (ex *H. armigera* (Hübner)); Winburn and Painter, 1932: 5, 19, Kans.; Phillips and Barber, 1931: 11-12, E. U.S.; Essig, 1926: 584; Allen, 1925: 5-6, 10, 29, figs. 2-7, Miss.; Luginbill and Beyer, 1921: 13 (ex corn earworm); Quaintance and Brues, 1905: 123-124, Tex. (as *W. 4-pustulata*).

Leucania latiuscula Herrich-Schaeffer probably
Wolcott, 1951a: 481, P.R. (ex *Cirphis*).

Lithophane antennata (Walker)

All records ex *Xylina antennata*.

Felt, 1912: 49-50, N.Y.; Felt, 1898: 211, N.Y. (as *W. 4-pustulata*).

Papaipema nebris (Guenée)

Decker, 1931: 344, Iowa (as "*Winthema*").

Peridroma saucia (Hübner)

Crumb, 1929: 40, 114-115 (ex *Lycophotia*); Essig, 1926: 584, Oreg. (ex *Lycophotia margaritosa* (Haworth)); Allen, 1925: 6, 9-10, 24, 29, figs. 2-9, Miss. (ex *Lycophotia margaritosa*); Cook, 1912: 273, fig. 118, Calif. (as *W. "4-postulata"*); Chittenden, 1901a: 62 (as *W. 4-pustulata*); Coquillett, 1897: 22, 26 (as *W. 4-pustulata*).

Plathypena scabra (Fabricius)

Webber and Schaffner, 1926: 22, New England.

Platysenta sutor (Guenée)

All records ex *Perigea sutor*.

Allen, 1925: 6, figs. 2-7; Brimley, 1922: 26, N.C. (ex "*Perigaea*").

Platysenta videns (Guenée)

Seifert, 1901: 13, N.Y.

Pseudaletia unipuncta (Haworth)

Anon., 1969b: 637, Wis.; Anon., 1966a: 92, Mich.; Anon., 1965a: 402, Wis.; Guyer et al., 1965: 670, Mich.; Anon., 1964b: 779, Wis. (incomplete rearing — egg, larva); Anon., 1959: 241-242, Wis.; Anon., 1958e: 648, Wis.; Franklin, 1950: 57-58 (ex *Cirphis*); Walton and Packard, 1940: 7, U.S. (ex *Cirphis*); Luginbill, 1928: 78 (ex true armyworm); Poos, 1928: 146, Va. (ex *Cirphis*); Essig, 1926: 584, Oreg. (ex *Cirphis*); Webber and Schaffner, 1926: 22, New England (ex *Cirphis*); King, V., and Barber, 1921: 487, Mo. (as *W. 4-pustulata* ex *Heliophila*, probable host); Severin, 1921: 6-7, S. Dak. (ex "*Cirphus*"); Lochhead, 1919: 190, fig. 129 (as *W.*

4-pustulata ex *Cirphis*); Felt, 1916: 37, N.Y. (ex *Heliophila*); Knight, 1916: 760-761, fig. 1 on pl. 40, N.Y. (ex *Leucania*); Walton, 1916: 9, fig. 2b (ex *Cirphis*); Baker, 1915: 86, Ont. (as *W. 4-pustulata* ex *Heliophila*); Britton, 1915a: 167, Conn. (ex *Heliophila*); Essig, 1915: 396, Calif. (as *W. 4-pustulata* ex *Cirphis*); Gibson, 1915: 13, N.S., Ont. (ex *Cirphis*); Sherman, 1915: 299-302, N.C. (as *W. 4-pustulata* ex *Heliophila*); Baldwin, 1914: 47, Ind. (ex *Leucania*); Sanderson, 1913: 117-118, fig. 87c-87d (as *W. 4-pustulata* ex *Leucania*); Felt, 1912: 49-50, N.Y. (ex *Heliophila*); Johnson, 1910: 782, N.J. (ex *Leucania*); Metcalf, 1908: 354-355, S.C. (ex *Heliophila*); Webster, 1908: 247, fig. 25 (ex *Heliophila*); Felt, 1906: 65, N.Y. (ex *Heliophila* "unipunctata"); Washburn, 1906: 61, fig. 51c-51d, Minn. (ex *Leucania* "unipunctata"); Kellogg, 1905: 347, fig. 493c-493d (as *W. "4-pustulata"* ex *Leucania*); Coquillett, 1897: 22, 26, Mass., N.Y., Pa., Va. (as *W. 4-pustulata* ex *Leucania*); Howard, 1897b: 44, D.C. (ex *Leucania*); Lintner, 1897: 210, 310, 353-354, N.Y. (as *W. 4-pustulata* ex *Leucania*).

Schinia brevis (Grote)

Bird, H., 1923: 199 (ex *Lygranthoecia*).

Spodoptera dolichos (Fabricius)

Crumb, 1929: 40, 148 (ex *Prodenia*).

Spodoptera eridania (Cramer)

All records ex *Prodenia eridania*.

Patton, 1958: 85, Fla.; *Crumb, 1929: 40, 145 (Chittenden and Russell, 1909); Allen, 1925: 6, 10, 24, 26, 28-29, figs. 2-7, Miss.; Chittenden and Russell, 1909: 63, Fla.

Spodoptera frugiperda (J. E. Smith)

Most records ex *Laphygma frugiperda*.

Franklin, 1950: 57; Luginbill, 1950: 6-7, U.S.; Smith, R.C., 1943: 376, fig. 409, Kans. (ex fall armyworm, incomplete rearing — egg); Luginbill, 1928: 78-80, S.C.; Allen, 1925: 6, 10, 24, 29, figs. 2-7, Miss.; Smith, R. C., 1921: 304-305, Kans.; Walton and Luginbill, 1916: 10-11; Britton, 1913: 286, Conn.; Walton, 1913b: 25, D.C.; Walton, 1913c: 129-130, D.C.; Chittenden, 1901a: 35, fig. 5, Ill., Mass., Md.; Chittenden, 1900: 84 (as *W. "quadripunctata"*); Coquillett, 1897: 22, 25 (as *W. 4-pustulata*); Howard, 1897b: 44, D.C.

Spodoptera ornithogalli (Guenée)

All records ex *Prodenia ornithogalli*.

Crumb, 1929: 40, 155-156; Allen, 1925: 6, figs. 2-7.

Trichoplusia ni (Hübner)

Bosch and Hagen, 1966: 25, Calif.; Allen, 1925: 7-8, figs. 2-7

(ex *Autographa brassicae* Riley).

Xylena nupera (Lintner)

Franklin, 1950: 57, fig. 3B.

Armyworm(-s)

Fattig, 1949: 36; Smith, R. C., 1943: 376, Kans.; Johnson, 1925: 203, Conn., Maine, Mass., N.H., R.I., Vt.; Essig, 1913: 160-161, fig. 142, [Calif.] (as *W. 4-pustulata*); Garman, 1908: 435-436, Ky. (as *W. 4-pustulata*); Washburn, 1905: 156, fig. 150c-150d, Minn. (as *W. "pustulata"*); Howard, 1897b: 51, D.C. (as *W. 4-pustulata*); Riley, 1875: 179 (as red-tailed Tachina fly, incomplete rearing — egg).

Cutworm(-s)

Fattig, 1949: 36; Smith, R. C., 1943: 376, Kans.; Allen, 1925: 8, 25, 27, figs. 2-9; Lochhead, 1919: 185 (as *W. 4-pustulata*); Gibson, 1915a: 11, fig. 7, E. Canada; Essig, 1913: 160-161, fig. 142, [Calif.] (as *W. 4-pustulata*); Howard, 1897b: 44, D.C.

Cutworm(-s) and related Noctuidae

Allen, 1924: 24, Miss.

Unidentified Noctuidae

Wolcott, 1924: 223, P.R. (as *W. "quadripustulata"* ex noctuid on sugarcane).

Notodontidae

Datana ministra (Drury)

Essig, 1926: 584; Allen, 1925: 7, figs. 2-7; Smith, J. B., 1910: 146 (as *W. IV-pustulata*); Coquillett, 1897: 21, 24 (as *W. 4-pustulata*).

Heterocampa guttivitta (Walker)

Collins, 1926: 693, 695, Mass.

Schizura concinna (J. E. Smith)

Webber and Schaffner, 1926: 19, New England.

Olethreutidae

Spilonota ocellana (Denis and Schiffermüller)

Porter, 1924: 16, Conn.

Pyralidae

Loxostege similalis (Guenée)

Smith, R. C., and Franklin, 1954: 31, Kans.

Saturniidae

Antheraea polyphemus (Cramer)

Most records ex *Telea polyphemus*.

Collins, M. M., and Weast, 1961: 94 (ex *Antheraea*); Essig, 1926: 584; Allen, 1925: 6, figs. 2-7; Johnson, 1910: 782, N.J.; Coquillett, 1897: 22, 27, D.C. (as *W. 4-pustulata*).

Callosamia promethea (Drury)

Howard, 1897b: 44, D.C. (ex *Attacus*).

Platysamia cecropia (Linnaeus)

Collins, M. M., and Weast, 1961: 94 (ex *Hyalophora*); Allen, 1925: 6, figs. 2-7 (ex *Samia*); Watson, 1921: 233-234, N.Y. (ex *Samia*); Johnson, 1910: 782, N.J. (ex *Attacus*); Coquillett, 1897: 21, 24, Mo. (as *W. 4-pustulata* ex *Attacus*).

Sphingidae

Hemaris diffinis (Boisduval)

Allen, 1925: 7, figs. 2-7 (ex *Haemorrhagia*).

Hyles lineata (Fabricius)

All records ex *Celerio lineata*.

Essig, 1926: 584; Allen, 1925: 7, figs. 2-7.

Manduca quinquemaculata (Haworth)

Most records ex *Protoparce quinquemaculata*.

Gilmore, 1938: 710-711, Ky. and/or Tenn.; Hefley, 1928: 213-221; Hefley, 1926: 77-80, Okla. (as *W. "4-pustulata"*); Allen, 1925: 7, figs. 2-7; Johnson, 1910: 782, N.J. (ex *P. celeus* Hübner); Coquillett, 1897: 22, 26, Mo. (as *W. 4-pustulata* ex *P. celeus*).

Manduca sexta (Linnaeus)

Most records ex *Protoparce sexta*.

Gilmore, 1938: 710-711, Ky. and/or Tenn.; Essig, 1926: 584; Garman, 1897: 29 (as *W. 4-pustulata* ex *Phlegthontius carolina* (Linnaeus)).

Pachysphinx modesta (Harris)

Fletcher, 1904: 99, Ont. (ex *Marumba*).

Pholus achemon (Drury)

Tothill, 1913: 75, Ont.

Sphinx gordius Cramer probably

Aldrich, 1932: 2, N.H.

Unidentified Sphingidae

Howard, 1897b: 44, D.C.

Unidentified Lepidoptera

Miscellaneous

Aldrich, 1915: 82.

STAGE PARASITIZED: Larva (Coleoptera, Hymenoptera, Lepidoptera), pupa (Lepidoptera).

IMMATURES: Egg — Franklin, 1950: (fig. 3B); Smith, R. C., 1943: (fig. 409); Allen, 1925: (descr., figs. 2, 8); Lochhead, 1919: (fig. 129); Felt, 1916: (descr.); Walton, 1916: (fig. 2b); Gibson, 1915: (fig. 7); Essig, 1913: (fig. 142); Sanderson, 1913: (fig. 87c-87d); Cook, 1912: (fig. 118); Garman, 1908: (descr.); Webster, 1908: (fig. 25); Felt, 1906: (descr.); Washburn, 1906: (fig. 51c-51d); Kellogg, 1905: (fig. 493c-493d); Quaintance and Brues, 1905: (descr.); Washburn, 1905: (fig. 150c-150d); Chittenden, 1901a: (fig. 5); Riley, 1875: (descr.).

Larva — Chittenden, 1901a: (fig. 5).

First instar larva — Hefley, 1926: (descr.); Allen, 1925: (descr., fig. 3).

Second instar larva — Allen, 1925: (descr., figs. 4, 9).

Third instar larva — Hefley, 1926: (descr.); Allen, 1925: (descr., figs. 5-6); Webster, 1908: (fig. 25).

Pupa — Allen, 1925: (descr., fig. 7); Greene, 1922: (descr., fig. 95 on pl. 20); Knight, 1916: (fig. 1 on pl. 40); Webster, 1908: (fig. 25); Chittenden, 1901a: (fig. 5).

BIOLOGY: Smith, R. C., and Franklin, 1954: (31); Franklin, 1950: (57-58); Luginbill, 1950: (6-7); Walton and Packard, 1940: (7); Decker, 1931: (344); Phillips and Barber, 1931: (11-12); Hefley, 1928: (213-221); Luginbill, 1928: (78-80); Hefley, 1926: (77-80); Allen, 1925: (1-32); Luginbill and Beyer, 1921: (13); Watson, 1921: (233-234); Knight, 1916: (760-761); Walton, 1916: (9); Walton and Luginbill, 1916: (10-11); Baker, 1915: (86); Gibson, 1915: (13); Walton, 1913c: (129-130); Felt, 1906: (65); Quaintance and Brues, 1905: (123-124); Chittenden, 1901a: (35); Chittenden, 1900: (84); Howard, 1897b: (7, 44); Lintner, 1897: (210, 310, 353-354); Riley, 1875: (179).

Winthemia quadripustulata (Fabricius) probably

HOST: LEPIDOPTERA

Noctuidae

Ceramica picta (Harris)

Schaffner and Griswold, 1934: 47, 123.

Epiglaea apiata (Grote)

Schaffner and Griswold, 1934: 50, 122, Mass.

Pseudaletia unipuncta (Haworth)

Dillier and Seibels, 1965: 457, Ala. (incomplete rearing — egg); Thompson, R. W., 1944: 30, Ont. (ex *Cirphis*).

Nymphalidae

Nymphalis milberti (Godart)

Schaffner and Griswold, 1934: 11, 122 (ex *Hamadryas*).

STAGE PARASITIZED: Larva.

BIOLOGY: Schaffner and Griswold, 1934: (122).

Winthemia "quadripustulata," see *Winthemia quadripustulata*

Winthemia "rufopicta," see *Winthemia rufopicta*

Winthemia rufopicta (Bigot)

HOST: LEPIDOPTERA

Geometridae

Alsophila pometaria (Harris)

Schaffner and Griswold, 1934: 90, 123.

Cankerworm(-s)

Decker, 1935: 570, Iowa.

Lasiocampidae

Tent caterpillar(-s)

LeRoux, 1961: 114, Que.

Noctuidae

Alypia octomaculata (Fabricius)

Schaffner and Griswold, 1934: 41, 123.

Anticarsia gemmatilis Hübner

Hinds and Osterberger, 1931: 1172, La. (ex *A. "gemmatilis"*).

Ceramica picta (Harris)

Schaffner and Griswold, 1934: 47, 123.

Cucullia convexipennis Grote and Robinson

Schaffner and Griswold, 1934: 48, 123.

Heliothis zea (Boddie)

Bottrell, Young, Price and Adams, 1968: 1054, Okla.

Heliothis spp.

Bottrell, Young, Price and Adams, 1968: 1054, Okla.

Lithophane antennata (Walker)

Schaffner and Griswold, 1934: 49, 123 (ex *Graptolitha*).

Lithophane spp. (including *L. disposita* Morrison, *L. innominata* (Smith) and/or *L. petulca* Grote)

Schaffner and Griswold, 1934: 48, 123 (ex *Graptolitha* (including *G. disposita*, *G. innominata* and/or *G. petulca*)).

Luperina stipata (Morrison)

Decker, 1935: 569-570, 578, Iowa.

Oligia fractilinea (Grote)

Decker, 1935: 569-570, 578, Iowa.

Papaipema cataphracta (Grote)

Decker, 1935: 569-570, 578, Iowa.

Papaipema nebris (Guenée)

Decker, 1935: 569-570, 578, Iowa; Reinhard, 1931: 32, Iowa.

Peridroma saucia (Hübner)

Anon., 1969: 59, Okla.; Montgomery, 1933: 186, Ind. (ex *Lycophotia*).

Platysenta sutor (Guenée)

Brimley, 1938: 364, N.C. (ex "*Perigaea*").

Pseudaletia unipuncta (Haworth)

Burrell, 1967: 114, La.; Guppy, J. C., 1967: 97, 99, 102-103, Ont.; Pond, 1960: 665, N.B.; Breeland, 1958: 321, 323-325, 331-332, figs. 23-24, Tenn.; Fattig, 1949: 36, Ga. (ex *Cirphis*); Beaulne, 1939: 120, Que. (as *W. "rufopecta"* ex *Leucania*); Beaulne, 1939a: 95, Que. (as *W. "rufopecta"* ex *Leucania*); Schaffner and Griswold, 1934: 47, 123 (ex *Cirphis*).

Spaelotis clandestina (Harris)

LeRoux, 1961: 115, Que.

Spodoptera eridania (Cramer)

Patton, 1958: 85, Fla. (ex *Prodenia*).

Spodoptera frugiperda (J. E. Smith)

Most records ex *Laphygma frugiperda*.

Patton, 1958: 85, Fla.; Hofmaster and Greenwood, 1949: 505, Va.; Brimley, 1938: 364, N.C. (ex fall army worm).

Spodoptera ornithogalli (Guenée)

Bottrell, 1969: 251, Okla. (ex *Prodenia*).

Trichoplusia ni (Hübner)

Schaffner and Griswold, 1934: 69, 123 (ex *Autographa brassicae* Riley).

Xylena nupera (Lintner)

Schaffner and Griswold, 1934: 50, 123, Mass.

Armyworm(-s)

Brimley, 1938: 364, N.C.; Reinhard, 1931: 33.

Noctuid spp. (including *Mamestra configurata* Walker and/or *Peridroma saucia* (Hübner))

Anon., 1962: 121, B.C.; Handford, 1962: 19, B.C.

Noctuid spp. (including *Heliothis zea* (Boddie), *Prodenia dolichos* (Fabricius), *Spodoptera eridania* (Cramer), *S. frugiperda* (J. E. Smith) and/or *S. ornithogalli* (Guenée))

Patton, 1958: 85, Fla. (ex *P. eridania*, *P. ornithogalli*, *P. dolichos*, *H. armigera* (Hübner) and/or *Laphygma frugiperda*).

Sphingidae

Hemaris diffinis (Boisduval)

*Brauer and Bergenstamm, 1895: 553, 594, N. Amer. (as *Exorista ciliata* Townsend, Townsend, 1893); Townsend, 1893: 467, Ill. (as *Exorista ciliata*).

Unidentified Sphingidae

Schaffner and Griswold, 1934: 26, 123.

STAGE PARASITIZED: Larva.

IMMATURES: Third instar larva — Breeland, 1958: (fig. 24).

Pupa — Breeland, 1958: (fig. 23).

BIOLOGY: Guppy, J. C., 1967: (97, 99, 102-103); Patton, 1958: (85); Schaffner and Griswold, 1934: (123); Hinds and Osterberger, 1931: (1172).

Winthemia rufopicta (Bigot) probably

HOST: LEPIDOPTERA

Noctuidae

Amathes c-nigrum (Linnaeus)

Schaffner and Griswold, 1934: 42-43, 123 (ex *Agrotis*).

Amphipyra pyramidoides Guenée

Schaffner and Griswold, 1934: 52, 123.

Epiglaea apiata (Grote)

Schaffner and Griswold, 1934: 50, 123, Mass.

Notodontidae

Heterocampa guttivitta (Walker)

Schaffner and Griswold, 1934: 79-80, 123.

Winthemia sinuata Reinhard

HOST: LEPIDOPTERA

Geometridae

Cingilia catenaria (Cramer) probablyReinhard, 1931: 26, Maine (ex *C. latenaria* Drury).*Erannis tiliaria* (Harris)

Schaffner and Griswold, 1934: 97-98, 122.

Hesperiidae

Thymelicus lineola (Ochsenheimer)

Arthur, 1962: 1085-1086, Ont.

Lasiocampidae

Tolype laricis (Fitch)

Raizenne, 1952: 100, Ont.

Noctuidae

Luperina stipata (Morrison)*Vide Reinhard, 1931: 26-27, Iowa (Decker, 1930); Decker, 1930: 159, Iowa (as *W. quadripustulata* (Fabricius)).*Plathypena scabra* (Fabricius)Whiteside, Burbutis and Kelsey, 1967: 326-328, fig. 3, Del.;
Schaffner and Griswold, 1934: 71-72, 122, Mass.*Spodoptera frugiperda* (J. E. Smith)Fattig, 1949: 36, Ga. (ex "*Laphygma*").

Notodontidae

Schizura unicornis (J. E. Smith)

Raizenne, 1952: 91, Ont.

STAGE PARASITIZED: Larva, pupa.

BIOLOGY: Schaffner and Griswold, 1934: (122); Decker, 1930:
(159).*Winthemia sinuata* Reinhard probably

HOST: LEPIDOPTERA

Geometridae

Cingilia catenaria (Cramer)

Schaffner and Griswold, 1934: 98, 122.

Unidentified Geometridae

Schaffner and Griswold, 1934: 102, 122.

Noctuidae

Calpe canadensis Bethune

Schaffner and Griswold, 1934: 70-71, 122, Mass.

Nymphalidae

Nymphalis antiopa (Linnaeus)

Schaffner and Griswold, 1934: 11, 122 (ex *Hamadryas*).

Winthemia tricolor (Wulp)

HOST: LEPIDOPTERA

Saturniidae

Samia cynthia (Drury)

Reinhard, 1931: 44, Panama (ex *Arsenura erythrinae* Fabricius).

Winthemia venusta (Meigen) probably

HOST: LEPIDOPTERA

Geometridae

Lambdina fiscellaria lugubrosa (Hulst)

Anon., 1946a: 336, B.C. (as *W. cilitibia* (Rondani) probably).

Winthemia spp. (including *W. leucanae* probably or *W. militaris*)

HOST: LEPIDOPTERA

Noctuidae

Pseudaletia unipuncta (Haworth)

Dearness, 1898: 15, fig. 4, Ont. (as red-tailed Tachina or possibly *Nemoraea "leucaniae"* ex *Leucania*).

STAGE PARASITIZED: Larva.

IMMATURES: Egg — Dearness, 1898: (fig. 4).

Third instar larva — Dearness, 1898: (fig. 4).

Pupa — Dearness, 1898: (fig. 4).

Winthemia spp. (including *W. quadripustulata* or allied sp.: *W. datanae*, *W. sinuata* or *W. vesiculata*)

HOST: COLEOPTERA

Chrysomelidae

Leptinotarsa decemlineata (Say)

Feytaud, 1937: 65 (incomplete rearing — egg).

STAGE PARASITIZED: Larva.

BIOLOGY: Feytaud, 1937: (65).

Winthemia "4-postulata," 4-pustulata, "4-pustuluta," or IV-pustulata, see *Winthemia quadripustulata*

Winthemia spp.

HOST: LEPIDOPTERA

Arctiidae

Hyphantria cunea (Drury)

Schaffner and Griswold, 1934: 37, 123 (incomplete rearing — pupa).

Geometridae

Lambdina fiscellaria (Guenée)

Schaffner and Griswold, 1934: 99, 123 (ex *Ellopia*, incomplete rearing — pupa).

Megalopygidae

Megalopyge opercularis (J. E. Smith)

Snyder, 1936: 423, La.

Noctuidae

Catocala antinympa (Hübner)

Schaffner and Griswold, 1934: 64, 123 (incomplete rearing — pupa).

Catocala minuta Edwards

Schaffner and Griswold, 1934: 66-67, 123, N.J. (incomplete rearing — pupa).

Heliothis zea (Boddie)

Lewis, W. J., and Brazzel, 1968: 674-675, Miss.

Pseudaletia unipuncta (Haworth)

Goble, 1965: 12, Ont.; Baker, 1938: 97-98, Ont. (ex *Leucania*, incomplete rearing — egg).

- Spodoptera ornithogalli* (Guenée)
 Bottrell, 1969: 251, Okla. (ex *Prodenia*).
 Cutworm(-s)
 Shaw, 1947: 437, Mexico.

Notodontidae

- Ichthyura inclusa* Hübner
 Schaffner and Griswold, 1934: 73, 123 (ex *Melalopha*,
 incomplete rearing — pupa).
Schizura spp.
 All records ex "*Shizura*" sp.
 *Procter, 1946: 431, Maine (Procter, 1938); Procter, 1938:
 372, Maine.
Symmerista albifrons (J. E. Smith)
 Schaffner and Griswold, 1934: 78, 123 (incomplete rearing —
 pupa).

Pieridae

- Eucheira socialis* Westwood
 Shaw, 1947: 437, Mexico.

Saturniidae

- Callosamia promethea* (Drury)
 Schaffner and Griswold, 1934: 27, 123 (incomplete rearing —
 pupa).
Hemileuca spp. (including *H. lucina* H. Edwards and/or *H. maia*
 (Drury))
 Schaffner and Griswold, 1934: 29, 123 (incomplete rearing —
 pupa).
Platysamia cecropia (Linnaeus)
 Schaffner and Griswold, 1934: 26-27, 123 (incomplete
 rearing — pupa).

Sphingidae

- Manduca sexta* (Linnaeus)
 Lawson, 1959: 741, 744-745, N.C. (ex *Protoparce*).

Tortricidae

- Choristoneura fumiferana* (Clemens)
 Tothill, 1923: 175-176, B.C., N.B. (ex spruce budworm).

STAGE PARASITIZED: Larva, pupa.

BIOLOGY: Baker, 1938: (97-98); Schaffner and Griswold, 1934: (123).

Winthemia spp. probably

HOST: LEPIDOPTERA

Noctuidae

Heliothis spp.

Tower, 1913: 13, probably Tenn.

STAGE PARASITIZED: Pupa.

Xanthoactia palloris, see *Actia palloris*

Xanthoepalpus Townsend (Tachininae: Juriniini)

Xanthoepalpus bicolor (Williston)

HOST: LEPIDOPTERA

Geometridae

Unidentified Geometridae

Anon., 1953c: 137, B.C.

STAGE PARASITIZED: Pupa.

Xanthoernestia sp., see *Xanthophyto* sp. probably

Xanthophyto Townsend (Tachininae: Ernestiini)

Xanthophyto labis (Coquillett)

HOST: LEPIDOPTERA

Aegeriidae

Vespa mima pini (Kellicott)

Raizenne, 1952: 246, Ont. (ex *Parharmonia*).

Lasiocampidae

Malacosoma incurvum incurvum (H. Edwards)

Stehr and Cook, 1968: 280, 301, Ariz.

Olethreutidae

Sciaphila duplex (Walsingham)

McGregor, 1967: 1215.

STAGE PARASITIZED: Larva, pupa.

Xanthophyto spp.

HOST: LEPIDOPTERA

Geometridae

Sabulodes truxaliata (Guenée)

Tilden, 1951: 165, Calif. (ex *Prochoerodes*).

Oecophoridae

Depressaria pastinacella (Duponchel)

Schaffner, 1959: 39, 54, N.J., Vt. (as *Xanthoernestia* ex *D. heracliana* (Linnaeus)).

Pyralidae

Dioryctria amatella (Hulst)

Ebel, 1965: 628, Fla.

Tortricidae

Choristoneura fumiferana (Clemens)

Dowden, Buchanan and Carolin, 1948: 458 (ex *Archips*).

STAGE PARASITIZED: Larva.

BIOLOGY: Schaffner, 1959: (39, 54).

Xanthophyto sp. probably

HOST: LEPIDOPTERA

Geometridae

Ennomos subsignarius (Hübner)

Davis, R., 1961: 248, Ga. (as *Xanthoernestia*).

STAGE PARASITIZED: Larva, pupa.

Ypophaemyia malacosomae, see *Lespesia archippivora*

"*Zanillia*" *confinis*, see *Aplomya theclarum*

Zelia Robineau-Desvoidy (Dexiinae: Zeliini)

"*Zelia* sp. near *genuina* Wulp"

IMMATURES: Pupa — Greene, 1922: (descr., fig. 87 on pl. 18).

Zelia vertebrata (Say)

HOST: COLEOPTERA

Cerambycidae

Unidentified Cerambycidae

Brauer and Bergenstamm, 1895: 551, 586, N. Amer. (as
(*Dexia*) ex larva on chestnut).

Passalidae

Popilius disjunctus (Illiger)

Most records ex *Passalus cornutus* Fabricius.

Patton, 1958: 85, Fla.; Fattig, 1949: 37; Gray, 1946:
740-741, N.C. (ex *P. disjunctus*); Mangrum, 1942:
73-75, pl. 1.

Scarabaeidae

Osmoderma eremicola Knoch

Stevenson, 1903: 58, Que.

Tenebrionidae

Meracantha contracta (Palisot de Beauvois)

Hyslop, 1916: 24-25, Md.

STAGE PARASITIZED: Larva, pupa.

IMMATURES: First instar larva — Mangrum, 1942: (descr.).

Third instar larva — Mangrum, 1942: (descr., pl. 1).

Pupa — Mangrum, 1942: (descr., pl. 1); Greene, 1922: (descr.,
fig. 10 on pl. 2).

BIOLOGY: Gray, 1946: (740-741).

"Zenilla" *blanda*, see *Eusisyropa blanda*

"Zenilla" *caesar*, see *Aplomya caesar*

Zenillia Robineau-Desvoidy (Goniinae: Eryciini)

Zenillia affinis, see *Hubneria estigmenensis*

Zenillia amplexa, see *Carcelia amplexa*

Zenillia autographae, see *Eusisyropa autographae*

Zenillia blanda, see *Eusisyropa blanda*

Zenillia blanda blanda, see *Eusisyropa autographae*, *Eusisyropa
blanda*, *Eusisyropa boarmiae* or *Eusisyropa virilis*

Zenillia blanda virilis, see *Eusisyropa virilis*

Zenillia blandita (Coquillett)

HOST: LEPIDOPTERA

Noctuidae

Nycteola spp. (including *N. cinereana* Neumoegen and Dyar, *N.
columbiana* H. Edwards and/or *N. frigidana* Walker)

All records ex *Sarothripus revayanus* (Scopoli).

*Sellers, 1943: 18 (Aldrich and Webber, 1924); Aldrich
and Webber, 1924: 41, D.C.

Zenillia boarmiae, see *Eusisyropa boarmiae*

- Zenillia caesar*, see *Aplomya caesar*
Zenillia ceratomiae, see *Hyphantrophaga hyphantriae*
Zenillia "ceratomyiae," see *Hyphantrophaga hyphantriae*
Zenillia cheloniae, see *Carcelia malacosomae* probably
Zenillia coerulea, see *Aplomya epicydes*
Zenillia confinis, see *Aplomya theclarum*
Zenillia dawsoni, see *Chrysoexorista dawsoni*
Zenillia desmiae, see *Hyphantrophaga desmiae*
Zenillia distincta, see *Drino rhoeo*
Zenillia euchaetiae, see *Hyphantrophaga euchaetiae*
Zenillia eudryae, see *Sisyropa alypiae* probably or *Sisyropa eudryae*
Zenillia formosa, see *Carcelia formosa*
Zenillia futilis, see *Euexorista futilis*
Zenillia griseomicans, see *Carcelia amplexa*
Zenillia helvina, see *Aplomya helvina*
Zenillia hyphantriae, see *Hyphantrophaga hyphantriae*
Zenillia inca, see *Drino inca*
Zenillia libatrix (Panzer)
 HOST: LEPIDOPTERA

Arctiidae

Euchaetias egle (Drury)

Sellers, 1943: 10 (lab.); Dowden, 1934: 98, 109, figs. 1-4 (lab.).

Bombycidae

Bombyx mori Linnaeus

Sellers, 1943: 10 (lab.); Dowden, 1934: 98, 112, figs. 1-4 (lab.).

Lymantriidae

Euproctis chrysorrhoea (Linnaeus)

Howard and Fiske, 1911: 302-303.

Notodontidae

Ichthyura inclusa Hübner

All records ex *Melalopha inclusa*.

Sellers, 1943: 10 (lab.); Dowden, 1934: 98, 113, figs. 1-4 (lab.).

STAGE PARASITIZED: Pupa.

IMMATURES: Egg — Dowden, 1934: (descr., figs. 1, 4).

First instar larva — Dowden, 1934: (descr., fig. 1).

Second instar larva — Dowden, 1934: (descr., fig. 2).

Third instar larva — Dowden, 1934: (descr., fig. 3).

Pupa — Dowden, 1934: (descr., fig. 3).

BIOLOGY: Sellers, 1943: (10); Dowden, 1934: (98, 112-113);

Howard and Fiske, 1911: (302-303).

Zenillia lobeliae, see "*Exorista*" *lobeliae*

Zenillia sp. near *lobeliae*, see "*Exorista*" sp. near *lobeliae*

Zenillia marginata, see *Chrysoexorista marginata*

Zenillia mathesoni, see *Thelymyia mathesoni*

Zenillia protuberans, see *Carcelia malacosomae* or *Carcelia protuberans*

Zenillia reclinata, see *Carcelia reclinata*

Zenillia "rectinata," see *Carcelia reclinata*

Zenillia roseanae, see *Pseudoperichaeta insidiosa*

Zenillia trisetosa, see *Aplomya trisetosa*

Zenillia virilis, see *Eusisyropa virilis*

Zenillia vulgaris, see *Phryxe vulgaris*

Zenillia n. sp. (1), see *Sisyropa alypiae* or "*Zenillia*" spp.

Zenillia n. sp. (2), see *Carcelia olenensis*, *Carcelia perplexa* or *Carcelia yalensis*

Zenillia n. sp. (3), see *Aplomya cerurae*

Zenillia n. sp. (4), see *Carcelia diacrisiae* or "*Zenillia*" spp.

Zenillia n. sp. (5), see *Aplomya pheosiae*

"*Zenillia*" spp.

HOST: HYMENOPTERA

Diprionidae

Neodiprion lecontei (Fitch)

Benjamin, 1955: 21, Canada; Daviault, 1953: 181, Que.

HOST: LEPIDOPTERA

Arctiidae

Halisidota tessellaris (J. E. Smith)

Schaffner and Griswold, 1934: 34, 113-114 (as *Zenillia* n. sp. (4)).

Haploa lecontei Guérin-Méneville

Schaffner and Griswold, 1934: 40 (ex *H. lecontei* including form *militaris* Harris).

Megalopygidae

Megalopyge opercularis (J. E. Smith)

Brauer and Bergenstamm, 1895: 564, 595, N. Amer. (as
"Myxexorsita" ex *Lagoa*).

Noctuidae

Acronicta betulae Riley

Brauer and Bergenstamm, 1895: 564, 580, N. Amer. (as
Myxexorista ex *Acronycta*).

Acronicta furcifera Guenée

Schaffner and Griswold, 1934: 55-56 (ex *Acronycta*).

Acronicta longa Guenée

Schaffner and Griswold, 1934: 59, N.J. (ex *Acronycta*).

Acronicta sp.

Brauer and Bergenstamm, 1895: 564, 580, N. Amer. (as
Myxexorista ex *Acronycta* sp. auf Eichen).

Psychomorpha epimenis (Drury)

Schaffner and Griswold, 1934: 63, 113, Mass. (as *Zenillia* n.
sp. (1)).

Pyralidae

Acrobasis caryae Grote

Nickels, Pierce and Pinkney, 1950: 2, 19, Tex.

Ostrinia nubilalis (Hübner)

Bradley, W. G., and Clark, 1938: 651, Mass. (ex European
corn borer).

Sphingidae

Unidentified Sphingidae

Schaffner and Griswold, 1934: 26.

STAGE PARASITIZED: Larva (Lepidoptera), pupa
(Lepidoptera).

BIOLOGY: Schaffner and Griswold, 1934: (113).

Gen. and sp. near *Zenillia*

HOST: LEPIDOPTERA

Noctuidae

Unidentified Noctuidae

Schaffner and Griswold, 1934: 72, 114, N.J.

BIOLOGY: Schaffner and Griswold, 1934: (72, 114).

Zygobothria gilva, see *Drino bohémica* or *Drino inconspicua*

Zygobothria nidicola, see *Townsendiellomyia nidicola*

Zygosturmia inca, see *Drino inca*

Zygosturmia protoparcis, see *Drino rhoeo*

Zygosturmia sp., see *Drino* sp.

Tachinidae (including Larvaevoridae and tachina fly)

HOST: COLEOPTERA

Buprestidae

Buprestis aurulenta Linnaeus

Chamberlin, 1932: 834, fig. 52, Oreg. (incomplete rearing — egg).

Carabidae

Carabus sp.

Curran, 1934: 138, B.C.

Pterostichus lucublandus (Say)

Rivard, 1964: 919, Ont.

Pterostichus melanarius Illiger

Rivard, 1964: 919, Ont.

Cerambycidae

Enaphalodes atomarius (Drury)

Savely, 1939: 384, N.C. (ex *Romaleum*).

Monochamus scutellatus (Say)

Hopkins, A. D., 1893a: 239, W. Va. (ex "*Monohamus*").

Saperda obliqua Say

Felt and Joutel, 1904: 20, N.E. N. Amer.

Chrysomelidae

Acalymma vittata (Fabricius)

Headlee, 1908: 207-209, N.H. (ex *Diabrotica*).

Altica probata Fall

Moznette, 1917: 16, Oreg. (ex *Haltica*).

Chrysomela scripta Fabricius

Knab, 1909: 35 (ex *Lina*, incomplete rearing — egg).

Diabrotica undecimpunctata Mannerheim

Garman, 1891: 80 (ex *D. 12-punctata* Mannerheim).

Diabrotica undecimpunctata howardi Barber

Riley, 1891a: 108, S.E. U.S. (ex *D. 12-punctata*).

Diabrotica undecimpunctata undecimpunctata Mannerheim

All records ex *Diabrotica soror* LeConte.

Kellogg, 1905: 346, Calif. (as *Tachina*-fly [sp. 2]);

Coquillett, 1889a: 74, Calif.

Leptinotarsa decemlineata (Say)

Popenoe, 1909: 3, Va.

Physonota helianthi (Randall)

Caulfield, 1887: 75, Que.

Cleridae

Thanasimus dubius (Fabricius)

Hopkins, A. D., 1893: 150, W. Va. (as *Tachina* sp.);

*Hopkins, A. D., 1893a: 239, W. Va. (as undetermined sp. b, Hopkins, A. D., 1893).

Coccinellidae

Epilachna varivestis Mulsant

Howard, N.F., and Landis, 1936: 2, N. Mex.; Howard, N. F., and Landis, 1936: 2, 6, Ala., Ohio (incomplete rearing — egg); Becker, 1922: 107, Mexico (ex Mexican bean beetle); Chittenden and Marsh, 1920: 14, Colo. (ex *E. corrupta* Mulsant, incomplete rearing — egg).

Curculionidae

Conotrachelus elegans (Say)

Brauer and Bergenstamm, 1895: 560, 588, N. Amer. (as Macquartiidae N.G. or N.G. Macquartiidarum).

Rhodobaenus tredecimpunctatus (Illiger)

Lugger, 1888: 50 (ex *R. 13-punctatus*, incomplete rearing — egg).

Sphenophorus incurrens Gyllenhal

Muir, 1923: 12, Mexico.

Scarabaeidae

Phyllophaga fusca (Frölich)

All records ex *Lachnosterna fusca*.

Howard, 1899: 198-199, N.Y. (incomplete rearing — egg);

Riley, 1891: 134.

Phyllophaga inversa (Horn)

Howard, 1899: 199, D.C. (ex *Lachnosterna*, probable host).

Phyllophaga spp.

All records ex *Lachnosterna* spp.

Wolcott, 1922: 15, Ill.; Howard, 1904: 98; Howard, 1899: 199.

May beetle(-s)

Forbes, S. A., 1907: 475, Ill.

Tenebrionidae

Eleodes opacus (Say)

Bruner, 1892: 12, Nebr. (incomplete rearing — egg).

Eleodes tricostata (Say)

Bruner, 1892: 12, Nebr. (incomplete rearing — egg).

HOST: DERMAPTERA

Carcinophoridae

Anisolabis maritima (Géné) probably

Spencer, 1926: 66, B.C. (incomplete rearing — larva).

HOST: DIPTERA

Cecidomyiidae

Rhabdophaga salicisbrassicoides (Packard)

Walsh, 1864: 633, Ill. (as robust blackish tachinid ex gall of *Cecidomyia* “*salicis brassicoides*”).

Tabanidae

Hybomitra lasiophthalma (Macquart)

Philip, 1931: 68, Minn. (ex *Tabanus*, incomplete rearing — larva).

Unidentified Tabanidae

Segal, 1936: 55.

Tipulidae

Tipula simplex Doane

Carnes and Newcomer, 1912: 279, Calif.

HOST: HEMIPTERA (HETEROPTERA)

Coreidae

Anasa tristis (De Geer)

Packard and Putman, 1875: 519 (ex *Coreus*).

Pentatomidae

Nezara viridula (Linnaeus)

Drake, 1918: 91.

Plant bug(-s)

Bratley, 1938: 14, Fla.

Rhopalidae

Liorhyssus hyalinus (Fabricius)

All records ex *Corizus hyalinus*.

Swezey, 1921: 467, Hawaii; Bridwell, 1920: 339-340, Hawaii.

HOST: HEMIPTERA (HOMOPTERA)

Cicadidae

Magiccada septendecim (Linnaeus)

Hyslop, 1936: 676 (probable host); Marlatt, 1907: 129 (ex *Tibicen*, probable host); Lintner, 1886: 178 (ex *Cicada*).

HOST: HYMENOPTERA

Argidae

Sphacophilus cellularis (Say)

Riley and Howard, 1888: 44, Miss. (ex *Schizocerus ebenus* Norton).

Cimbicidae

Cimbex americana Leach

Washburn, F. L., 1905: 158, Minn.; Cockerell, 1890: 76, Colo.; Riley and Howard, 1889: 164, Minn.

Diprionidae

Diprion similis (Hartig)

All records ex *Diprion* "simile."

Hartley, 1923: 387, Pa.; Britton, 1916: 282, Conn. (incomplete rearing — egg).

Tenthredinidae

Nematus ribesii (Scopoli)

Riley, 1877: 17 (ex *N. ventricosus* Klug).

Tethida cordigera (Palisot de Beauvois)

Osborne, 1884: 151, Iowa (ex *Selandria barda* Say).

HOST: LEPIDOPTERA

Arctiidae

Cycnia tenera Hübner

Schaffner and Griswold, 1934: 32 (incomplete rearing — pupa).

Diacrisia virginica (Fabricius)

Most records ex *Spilosoma virginica*.

Schaffner and Griswold, 1934: 36 (ex *Diacrisia*, incomplete rearing — pupa); Maxson, 1920: 87, Colo. (ex *Diacrisia*); Dimmock and Ashmead, 1898: 149, Mass. (incomplete rearing — pupa); Riley, 1871: 69, Mo. (as Muscadae probably); Riley, 1870d: 273 (as Muscadae probably).

Estigmene acrea (Drury)

Hills and Taylor, 1952: 126, Ariz.

Euchaetias egle (Drury)

Schaffner and Griswold, 1934: 39 (incomplete rearing — pupa); Thompson, 1910: 290.

Halisidota caryae (Harris)

Schaffner and Griswold, 1934: 33 (incomplete rearing — pupa).

Halisidota maculata (Harris)

Schaffner and Griswold, 1934: 33-34 (incomplete rearing — pupa).

Halisidota tessellaris (J. E. Smith)

Schaffner and Griswold, 1934: 34 (incomplete rearing — pupa).

Hyphantria cunea (Drury)

Oliver, 1964: 316-317, La.; Swain, 1937a: 43, Colo. (ex fall webworm); Schaffner and Griswold, 1934: 37 (incomplete rearing — pupa); Packard, 1890: 256; Riley, 1887: 52-53.

Isia isabella (J. E. Smith)

Schaffner and Griswold, 1934: 36-37 (incomplete rearing — pupa).

Phragmatobia fuliginosa rubricosa (Harris)

Schaffner and Griswold, 1934: 35-36 (incomplete rearing — pupa).

Utetheisa ornatrix bella (Linnaeus)

Bratley, 1933: 77, Fla. (ex "*Utethesia*" *bella*).

Brassolidae

Brassolis isthmia Bates

Dunn, 1917: 484, Panama.

Citheroniidae

Anisota senatoria J. E. Smith

Schaffner and Griswold, 1934: 29-30 (incomplete rearing — pupa).

Anisota virginiensis (Drury)

Schaffner and Griswold, 1934: 30 (incomplete rearing — pupa).

Eacles imperialis (Drury)

Lintner, 1889: 324, N.Y. (incomplete rearing — pupa).

Ctenuchidae

Ctenucha virginica Charpentier

Schaffner and Griswold, 1934: 32 (incomplete rearing — pupa).

Danaiidae

Danaus plexippus plexippus (Linnaeus)

Arnaud, 1960a: 51, Calif. (incomplete rearing — larva); Schaffner and Griswold, 1934: 8 (ex *D. menippe* (Hübner), incomplete rearing — pupa); Weeks, 1890: 82 (ex *D. archippus* (Fabricius)); Anon., 1876: 15 (ex "*Danais*" *archippus*).

Dioptidae

Phryganidia californica Packard

Keen, 1958: 90, Calif.; Van Duzee, 1926: 180, Calif. (ex "*Phryganidea*").

Drepanidae

Drepana bilineata (Packard)

Aldrich, 1932: 1, Mass. (incomplete rearing — pupa).

Geometridae

- Alsophila pometaria* (Harris)
Schaffner and Griswold, 1934: 90 (incomplete rearing — pupa).
- Cingilia catenaria* (Cramer)
Schaffner and Griswold, 1934: 98 (incomplete rearing — pupa).
- Coniodes plumogeraria* Hulst
Coquillett, 1897a: 66, Calif. (ex *Boarmia* “*plumigeraria*”).
- Deilinia erythemaria* (Guenée)
Schaffner and Griswold, 1934: 93-94, Maine (ex *Cabera* “*erythemaria*,” incomplete rearing — pupa).
- Ennomos subsignarius* (Hübner)
Schaffner and Griswold, 1934: 99-100, Mass. (incomplete rearing — pupa).
- Erannis tiliaria* (Harris)
Schaffner and Griswold, 1934: 97-98 (incomplete rearing — pupa); Coquillett, 1882a: 25, 28-29, Ill. (ex *Hibernia*).
- Eucaterva variaria* (Grote)
Brauer and Bergenstamm, 1895: 575, 592.
- Eulype hastata* (Linnaeus)
Schaffner and Griswold, 1934: 92 (incomplete rearing — pupa).
- Itame pustularia* (Guenée)
Schaffner and Griswold, 1934: 93 (ex *Physostegania*, incomplete rearing — pupa).
- Itame ribearia* (Fitch)
Riley, 1877: 6, Mo. (ex *Eufitchia*).
- Lambdina fiscellaria* (Guenée)
Schaffner and Griswold, 1934: 99 (ex *Ellopia*, incomplete rearing — pupa).
- Nematocampa filamentaria* Guenée
Schaffner and Griswold, 1934: 101 (ex *N. limbata* Haworth, incomplete rearing — pupa).
- Nepytiya phantasmaria* (Strecker)
Wickman and Hunt, 1969: 1049, Calif.
- Paleacrita vernata* (Peck)
Riley, Packard and Thomas, 1883: 176, Mass.; Riley, 1870: 103, Mass. (ex *Anisopteryx*).
- Phaeoura quernaria* (J. E. Smith)
Schaffner and Griswold, 1934: 96 (ex *Nacophora*, incomplete rearing — pupa).

Phigalia titea (Cramer)

Schaffner and Griswold, 1934: 97 (incomplete rearing — pupa).

Sabulodes transversata (Drury)

Schaffner and Griswold, 1934: 101 (incomplete rearing — pupa).

Hepialidae

Hepialus pharus Druce

Williams, 1935: 297, Guatemala.

Hesperiidae

Cymaenes tripunctus (Herrich-Schaeffer)

Scaramuzza and Barry, 1960: 997, Cuba (ex *Lerodea tripuncta tripuncta*); Scaramuzza, 1951a: 300, Cuba (ex *Lerodea tripuncta*).

Epargyreus clarus (Cramer)

Schaffner and Griswold, 1934: 15 (ex *E. tityrus* (Fabricius), incomplete rearing — pupa).

Erynnis icelus (Scudder and Burgess)

Schaffner and Griswold, 1934: 15 (incomplete rearing — pupa).

Thymelicus lineola (Ochsenheimer)

Pengelly, 1961: 195, Ont.

Hesperiid spp. (including *Cymaenes tripunctus* (Herrich-Schaeffer), *Nyctelius nyctelius* (Latreille) and/or *Panoquina sylvicola* (Herrich-Schaeffer))

Scaramuzza, 1946: 549, Cuba (ex *Lerodea tripunctus*, *Prenes nyctelius coscinia* (Herrich-Schaeffer) and/or *Prenes nero sylvicola*).

Lasiocampidae

Epicnaptera americana (Harris)

Schaffner and Griswold, 1934: 89 (incomplete rearing — pupa).

Malacosoma americanum (Fabricius)

Most records ex *Clisiocampa americana*.

Britton, 1935a: 77, Conn. (ex *Malacosoma*); Felt, 1901: 1007, N.Y.; Bethune, 1871: 17, Ont.

Malacosoma californicum californicum (Packard)

Koebele, 1891: 43, Calif. (ex *Clisiocampa thoracica* Stretch).

Malacosoma constrictum (H. Edwards)

All records ex *Clisiocampa constricta*.

Coquillett, 1894: 30, Calif.; Koebele, 1891: 43, Calif.

Malacosoma disstria Hübner

Hodson, 1941: 39, 41, Minn. (incomplete rearing — ?); Balch and Simpson, 1935: 36, N.B.; Schaffner and Griswold, 1934: 89 (incomplete rearing — pupa); Felt, 1901: 997, N.Y. (ex *Clisiocampa*); Bethune, 1871: 17, Ont. (ex *Clisiocampa sylvatica* (Harris)); Walsh and Riley, 1869: 208, N.Y. (ex *Clisiocampa sylvatica*).

American tent caterpillar(-s)

Baird, 1917: 86-87, B.C.

Limaecodidae

Euclea delphinii Boisduval

Schaffner, 1959: 5, [N.E. U.S.] (as Larvaevoridae, incomplete rearing — larva).

Limaecodes sp. probably

Lintner, 1889: 192.

Phobetron pithecium (J. E. Smith)

Schaffner, 1959: 6, [N.E. U.S.] (as Larvaevoridae, incomplete rearing — larva); Schaffner and Griswold, 1934: 104 (incomplete rearing — pupa); Lintner, 1889: 192, N.Y.

Prolimaecodes badia (Hübner)

Schaffner, 1959: 6, [N.E. U.S.] (as Larvaevoridae, incomplete rearing — pupa).

Sibine stimulea (Clemens)

Schaffner, 1959: 4, [N.E. U.S.] (as Larvaevoridae, incomplete rearing — pupa); Schaffner and Griswold, 1934: 103 (incomplete rearing — pupa).

Lycaenidae

Brephidium exilis (Boisduval)

Coolidge, 1924: 116, Calif.

Plebejus icarioides (Boisduval)

Newcomer, 1912: 33, Calif. (ex *Lycaena fulla* Edwards).

Lymantriidae

Hemerocampa leucostigma (J. E. Smith)

Schaffner and Griswold, 1934: 85 (incomplete rearing — pupa); Dustan, 1923: 117-118, N.B.; Dustan, 1923: 115,

N.B. (incomplete rearing — larva); Woods, 1907: 297-298, Maine (ex *Notolophus*); Panton, 1897: 53, Ont. (ex *Orgyia*); Riley, 1887: 32 (ex *Orgyia*); Lintner, 1886: 79, N.Y. (ex *Orgyia*).

Hemerocampa pseudotsugata McDunnough

Keen, 1958: 92, N.W. U.S.

Orgyia antiqua (Linnaeus)

All records ex *Notolophus antiqua*.

Schaffner and Griswold, 1934: 84 (incomplete rearing — pupa); Woods, 1907: 297-298, Maine.

Orgyia sp.

Lintner, 1886: 79.

Porthetria dispar (Linnaeus)

Collins, 1932: 26, Mass. (ex gypsy moth); Hitchings, 1908: 12, Maine (ex "gypsy" caterpillars, incomplete rearing — egg); Henshaw, 1892: 81, Mass. (ex *Ocneria*).

Stilpnotia salicis (Linnaeus)

Keen, 1958: 93, U.S.

Tussock moth(-s)

Snodgrass, 1917: 174, 189, Ind.

Megalopygidae

Megalopyge crispata (Packard)

All records ex *Lagoa crispata*.

Schaffner, 1959: 7, [N.E. U.S.] (as Larvaevoridae, incomplete rearing — pupa); Dillon, 1944: 176, Pa.; Schaffner and Griswold, 1934: 105-106 (incomplete rearing — pupa).

Noctuidae

Acronicta furcifera Guenée

Schaffner and Griswold, 1934: 55-56 (ex *Acronycta*, incomplete rearing — pupa).

Acronicta impressa Walker

Schaffner and Griswold, 1934: 58-59 (ex *Acronycta*).

Acronicta lepusculina Guenée

Riley, 1870: 120 (ex *Acronycta populi* Riley, incomplete rearing — larva).

Acronicta superans Guenée

Schaffner and Griswold, 1934: 55 (ex *Acronycta*, incomplete rearing — pupa).

- Actebia fennica* (Tauscher)
Phipps, 1927: 212, fig. 29D, Maine (ex *Agrotis*, incomplete rearing — egg); Lugger, 1897: 21, Minn. (ex *Noctua*).
- Agrotis dislocata* (Walker)
Nakao, 1963: 535, Hawaii.
- Alabama argillacea* (Hübner)
Jones, R. W., 1885: 19-20, Miss. (ex *Aletia xyliana* (Say)).
- Aletia* spp.
Hubbard, 1885: 16, Fla.
- Alypia octomaculata* (Fabricius)
Schaffner and Griswold, 1934: 41 (incomplete rearing — pupa).
- Amphipyra pyramidoides* Guenée
Schaffner and Griswold, 1934: 52 (incomplete rearing — pupa).
- Calloplistria floridensis* (Guenée)
Chittenden, 1913: 10, Fla. (ex *Eriopus*).
- Catocala parta* Guenée
Schaffner and Griswold, 1934: 65-66 (incomplete rearing — pupa).
- Ceramica picta* (Harris)
Schaffner and Griswold, 1934: 47 (incomplete rearing — pupa).
- Chrysodeixis calcites* (Esper)
All records ex *Plusia* "chalcites."
Williams, 1931: 142, Hawaii; Swezey, 1917: 428, Hawaii.
- Cirphis* spp.
Osborn, H. T., 1938: 150-151, Mexico (V.C.).
- Cosmia canescens* (Behr)
Schaffner and Griswold, 1934: 62 (incomplete rearing — pupa).
- Crymodes devastator* (Brace)
Schaffner and Griswold, 1934: 53-54 (ex *Sidemia*, incomplete rearing — pupa).
- Cucullia convexipennis* Grote and Robinson
Schaffner and Griswold, 1934: 49 (incomplete rearing — larva).
- Epiglaea apiata* (Grote)
Beckwith, 1930: 416, N.J.
- Euparthenos nubilis* (Hübner)
Schaffner and Griswold, 1934: 68 (incomplete rearing — pupa).
- Eupsilia* spp. (including *E. morrisoni* (Grote) and/or *E. sidus* (Guenée))

Schaffner and Griswold, 1934: 50-51 (ex *Conistra* spp. (including *C. sidus* and/or *C. morrisoni*), incomplete rearing — pupa).

Euthisanotia grata (Fabricius)

Lugger, 1899: 71, Minn. (ex *Eudryas*).

Euxoa auxiliaris (Grote)

Manis, Gittins, Portman, Walz, Barr, Douglass, Evans, Smith and Davis, 1956: 78, Idaho (ex *Chorizagrotis*).

Euxoa scandens (Riley)

Hudson and Wood, 1929: 69-70, Ont.

Faronta diffusa (Walker)

Smith, J. B., 1893: 190, N.J. (ex *Leucania albilinea* (Hübner)).

Heliothis zea (Boddie)

Most records ex *Heliothis armigera* (Hübner).

Anon., 1965b: 1091, Okla. (ex *H. zea*); Mally, 1893: 26;

Mally, 1891: 28; Hubbard, 1885: 16, Fla.

Heliothis spp.

Jones, R. W., 1885: 19, Miss.

Lithophane antennata (Walker)

Schaffner and Griswold, 1934: 49 (ex *Graptolitha*, incomplete rearing — pupa).

Lithophane spp. (including *L. disposita* Morrison, *L. innominata* (Smith) and/or *L. petulca* Grote)

Schaffner and Griswold, 1934: 48 (ex *Graptolitha* spp. (including *G. disposita*, *G. innominata* and/or *G. petulca*), incomplete rearing — pupa).

Mamestra configurata Walker

King, K. M., 1928: 288, Sask. (ex *Barathra*).

Nephelodes emmedonia (Cramer)

Lintner, 1883: 109, N.Y. (ex *N. violans* Guenée, incomplete rearing — pupa); Lintner, 1883: 103, Mo. (ex *N. violans*, incomplete rearing — egg).

Orthodes cynica Guenée

Schaffner and Griswold, 1934: 45, Mass. (ex *Eriopyga*, incomplete rearing — larva).

Papaipema nebris (Guenée)

Felt, 1904: 190, 216, N.Y. (ex *P. nitela* (Guenée)).

Parallelia bistriaris Hübner

Schaffner and Griswold, 1934: 68 (incomplete rearing — pupa).

Peridroma saucia (Hübner)

Davis, J. J., 1912: 86 (ex *P. margaritosa* (Haworth)); Chittenden, 1901a: 62, B.C.; Doane and Brodie, 1901: 10, Wash. (incomplete rearing — larva).

Pseudaletia unipuncta (Haworth)

Breeland, 1958: 323, 325, Tenn. (as "*Larvaevorida*"); Handford, Painter and Moore, 1937: 159, Man. (ex armyworms); Poos, 1928: 148, Va. (ex *Cirphis*, incomplete rearing — egg); Webster, 1890: 45, Ind. (as *Tachinae* ex *Leucania*); Kirkpatrick, 1861: 355, Ohio (ex *Leucania extranea* Guenée).

Pyreferra hesperidago (Guenée)

Schaffner and Griswold, 1934: 51 (ex *Conistra indirecta* Walker, incomplete rearing — pupa).

Spodoptera dolichos (Fabricius)

Chittenden, 1901: 64 (ex *Prodenia commelinae* (J. E. Smith), incomplete rearing — egg).

Spodoptera eridania (Cramer)

Anon., 1928b: 40, Fla. (ex *Xylomyges*).

Spodoptera frugiperda (J. E. Smith)

Most records ex fall armyworm.

Richardson, 1959: 152, Mexico (V.C.); Arant, 1955: 994, Ala.; Forbes, S. A., and Hart, 1900: 498, Ill. (ex *Laphygma frugiperda*).

Spodoptera ornithogalli (Guenée)

Chittenden, 1901: 68, 70, Ga. (ex *Prodenia*, incomplete rearing — egg).

Trichoplusia ni (Hübner)

McKinney, 1944: 15, Ariz. (ex *Autographa brassicae* Riley).

Armyworm(-s)

Kellogg, 1905: 346, Ala. (as *Tachina*-fly [sp. 1]); Walsh, 1867a: 119.

Cutworm(-s)

Gillette, 1891: 259, Iowa.

Unidentified Noctuidae

Smith, M. E., 1958: 123, Mass. (incomplete rearing — larva); Walsh, 1867a: 119, Iowa (ex "Miller" moth of the group of owlet-moths, incomplete rearing — egg).

Notodontidae

Cerura cinerea (Walker)

Baker, C. F., 1910: 382, Calif. (ex *Harpyia*).

Dasylophia anguina (J. E. Smith)

Schaffner and Griswold, 1934: 78-79 (incomplete rearing — pupa).

Datana integerrima Grote and Robinson

Schaffner and Griswold, 1934: 76 (incomplete rearing — pupa); Harned, 1927: 27, Miss.; vide Dyar, 1891: 414,

[N.Y.] (incomplete rearing — egg, Wier, 1890); Wier, 1890: 27 (as *Tachina* fly ex *D. ministra* (Drury) hairy, black walnut form, incomplete rearing — egg).

Datana ministra (Drury)

Schaffner and Griswold, 1934: 74 (incomplete rearing — pupa); Girault, 1912: 399-400, Va.; Lugger, 1899: 107, Minn.; Dyar, 1891: 414, [N.Y.] (incomplete rearing — egg); Webster, 1890: 45 (as *Tachinae*); Webster, 1890a: 257, Ind.; Wier, 1890: 26-27 (as *Tachina* fly, incomplete rearing — egg).

Datana perspicua Grote and Robinson

Schaffner and Griswold, 1934: 75 (incomplete rearing — pupa); Dyar, 1891: 414, [N.Y.] (incomplete rearing — egg).

Datana spp.

Hooker, W. A., 1908: 156, Mass. (ex walnut *Datana*); Weeks, 1890: 82.

Heterocampa guttivitta (Walker)

Schaffner and Griswold, 1934: 79-80 (incomplete rearing — pupa).

Hyperaeschra stragula (Grote)

Schaffner and Griswold, 1934: 76-77 (incomplete rearing — pupa).

Ichthyura inclusa Hübner

Schaffner and Griswold, 1934: 73 (ex *Melalopha*, incomplete rearing — pupa).

Nadata gibbosa (J. E. Smith)

Schaffner and Griswold, 1934: 78 (incomplete rearing — pupa).

Schizura concinna (J. E. Smith)

Schaffner and Griswold, 1934: 81 (incomplete rearing — pupa).

Schizura ipomaeae Doubleday

Schaffner and Griswold, 1934: 80 (ex *S. "ipomoeae,"* incomplete rearing — pupa).

Schizura unicornis (J. E. Smith)

Schaffner and Griswold, 1934: 81-82 (incomplete rearing — pupa); Lugger, 1899: 114, Minn.

Symmerista albifrons (J. E. Smith)

Schaffner and Griswold, 1934: 78 (incomplete rearing — pupa).

Nymphalidae

Chlosyne harrisii (Scudder)

Schaffner and Griswold, 1934: 9 (ex *Lemonias*, incomplete rearing — larva).

Limenitis archippus (Cramer)

All records ex *Limenitis disippus* Godart.

Saunders, 1873: 56, Ont.; Riley, 1871: 157 (incomplete rearing — egg).

Nymphalis antiopa (Linnaeus)

Patch, 1906: 225, Maine (ex *Euvanessa*); Weed, 1899: 139, N.H. (ex *Vanessa*).

Nymphalis milberti (Godart)

Guppy, R., 1951: 69, B.C.; Schaffner and Griswold, 1934: 11 (ex *Hamadryas*, incomplete rearing — pupa).

Polygonia comma (Harris)

Schaffner and Griswold, 1934: 10 (incomplete rearing — pupa); Howard, 1897: 51; *Brauer and Bergenstamm, 1895: 575, 603 ([incomplete rearing — pupa], Williston, 1889); Williston, 1889: 1913-1914 (incomplete rearing — pupa).

Oecophoridae

Depressaria pastinacella (Duponchel)

Schaffner, 1959: 39, [N.E. U.S.] (as Larvaevoridae ex *D. heracliana* (Linnaeus)).

Psilocorsis reflexella Clemens

Schaffner, 1959: 40, [N.E. U.S.] (as Larvaevoridae, incomplete rearing — pupa).

Olethreutidae

Grapholitha molesta (Busck)

Smith, R. W., 1929: 75, Ont. (ex *Laspeyresia*).

Rhyacionia frustrana (Comstock)

Swenk, 1927: 4, 46, Nebr.

Rhyacionia frustrana bushnelli Busck

Graham, S. A., and Baumhofer, 1927: 331, Nebr.

Papilionidae

Papilio glaucus Linnaeus

Schaffner and Griswold, 1934: 7 (ex *P. glaucus* form *turnus* Linnaeus, incomplete rearing — pupa).

Papilio polyxenes asterius Stoll

Schaffner and Griswold, 1934: 6 (ex *P. ajax* Linnaeus, incomplete rearing — pupa); Soule, 1899: 436, Vt. (ex *P. polyxenes* Fabricius, incomplete rearing — ?).

Papilio zelicaon Lucas

*Essig, 1915: 455, Calif. (ex *P. "zelicaon,"* Coolidge, 1910); Coolidge, 1910: 334, Calif. (ex *P. "zelicayn"*).

Pieridae

Pieris protodice Boisduval and LeConte

Hopkins, 1953: 794, Ariz.

Pieris rapae (Linnaeus)

Blunck, 1957: 836, Calif.; Schaffner and Griswold, 1934: 7-8 (ex *Ascia*, incomplete rearing — pupa); Packard, 1873: 58, N. Amer.; *vide Riley, 1870: 109-110, Que. (Provancher, 1869); Provancher, 1869: 18, Que. (as genre sarcophage).

Zerene "catenaria"

Dimmock and Ashmead, 1898: 149, N.H.

Plutellidae

Harpiteryx xylostella (Linnaeus)

Schaffner, 1959: 42, [N.E. U.S.] (as Larvaevoridae).

Psychidae

Thyridopteryx ephemeraeformis (Haworth)

Schaffner, 1959: 46, [N.E. U.S.] (as Larvaevoridae, incomplete rearing — larva); Balduf, 1937: 173, Ill. (incomplete rearing — pupa).

Pterophoridae

Oidaematophorus monodactylus (Linnaeus)

Schaffner, 1959: 19, [N.E. U.S.] (as Larvaevoridae, incomplete rearing — pupa).

Trichoptilus lobidactylus (Fitch)

Schaffner, 1959: 18, [N.E. U.S.] (as Larvaevoridae, incomplete rearing — pupa).

Pyralidae

Acrobasis betulella Hulst

Schaffner, 1959: 14, [N.E. U.S.] (as Larvaevoridae, incomplete rearing — pupa).

Acrobasis caryae Grote

All records ex case bearer.

Burkett, 1932: 212, Tex.; Burkett, 1925: 182-183, Tex.

Crambus spp.

Felt, 1906: 70, N.Y. (as small "Tachinid").

Diaphania hyalinata (Linnaeus)

Ashmead, 1887: 27 (ex *Eudiotis*).

Diatraea saccharalis (Fabricius)

Koebele, 1908: 509, Mexico (V.C.) (ex moth borer of the sugar cane, incomplete rearing — pupa).

Diatraea spp.

Wolcott, 1915: 37, Jamaica.

Dioryctria abietella (Denis and Schiffermüller)

Keen, 1958: 152.

Hedylepta accepta (Butler)

Swezey, 1906: 17, Hawaii (ex *Omiodes*).

Loxostege similalis (Guenée)

Forbes, S. A., and Hart, 1900: 457, Ill.

Loxostege sticticalis (Linnaeus)

Maxson, 1920: 77, fig. 12 on pl. 5, [Colo.] (ex *L. "sticticalis"*).

Macrobotys pertextalis (Lederer)

Schaffner, 1959: 10, [N.E. U.S.] (as Larvaevoridae, incomplete rearing — pupa).

Nephopteryx subcaesiella (Clemens) probably

Balduf, 1961: 82, Ill.

Phlyctaenia coronata tertialis (Guenée)

Schaffner, 1959: 10, [N.E. U.S.] (as Larvaevoridae ex *P. tertialis*, incomplete rearing — pupa).

Pyrausta futilalis (Lederer)

Schaffner, 1959: 11, [N.E. U.S.] (as Larvaevoridae, incomplete rearing — pupa).

Tetralopha subcanalis (Walker)

Leiby, 1925: 30, N.C.

Saturniidae

Actias luna (Linnaeus)

Schaffner and Griswold, 1934: 27-28 (incomplete rearing —

pupa); Riley, 1872: 125, Ky. (ex *Attacus*, incomplete rearing — egg).

Antheraea polyphemus (Cramer)

All records ex *Telea polyphemus*.

Schaffner and Griswold, 1934: 28 (incomplete rearing — pupa); Coolidge, 1908: 13, Calif.; Weed, 1897: 203-205.

Automeris io (Fabricius)

Schaffner and Griswold, 1934: 28 (incomplete rearing — pupa).

Callosamia promethea (Drury)

Schaffner and Griswold, 1934: 27 (incomplete rearing — pupa); Riley, 1872: 123 (ex *Attacus*).

Hemileuca spp. (including *H. lucina* H. Edwards and/or *H. maia* (Drury))

Schaffner and Griswold, 1934: 29 (incomplete rearing — pupa).

Platysamia cecropia (Linnaeus)

West, 1923: 23-25, N.Y. (ex *Samia*, incomplete rearing — larva); Lugger, 1899: 122, fig. 124 on pl. 14, Minn. (ex *Attacus*); Bruner, 1890: 18, Nebr.; Reed, 1873: 41, Ont.; Bethune, 1871: 24, Ont. (ex *Samia*).

Sphingidae

Agrius convolvuli (Linnaeus)

Perkins, R. C. L., 1907: 114, Hawaii (ex native species of the same group as *Sphinx*).

Ceratonia catalpae (Boisduval)

Schaffner and Griswold, 1934: 17, N.J. (incomplete rearing — pupa); Girault, 1908: 198, Tex.

Ceratonia undulosa (Walker)

Schaffner and Griswold, 1934: 17 (incomplete rearing — pupa).

Cressonia juglandis (J. E. Smith)

Schaffner and Griswold, 1934: 22 (incomplete rearing — larva).

Hemaris thysbe (Fabricius)

Schaffner and Griswold, 1934: 22 (incomplete rearing — pupa).

Hyles lineata (Fabricius)

Schaffner and Griswold, 1934: 25 (ex *Sphinx*, incomplete rearing — pupa).

Pachysphinx modesta occidentalis (H. Edwards)

Comstock and Dammers, 1943: 41, [Calif.].

Paonias myops (J. E. Smith)

Schaffner and Griswold, 1934: 21 (incomplete rearing — pupa).

Pholus satellitia pandorus (Hübner)

Hefley, 1926: 79, Okla. (ex *P. pandorus*).

Smerinthus jamaicensis (Drury)

Schaffner and Griswold, 1934: 20 (ex *S. geminatus* Say, incomplete rearing — pupa).

Sphinx chersis (Hübner)

Schaffner and Griswold, 1934: 18 (ex *Hyloicus*, incomplete rearing — pupa).

Sphinx drupiferarum J. E. Smith

Schaffner and Griswold, 1934: 20 (ex *Hyloicus*, incomplete rearing — pupa).

Sphinx gordius Cramer

Schaffner and Griswold, 1934: 19 (ex *Hyloicus*, incomplete rearing — pupa).

Sphinx kalmiae J. E. Smith

Schaffner and Griswold, 1934: 18-19 (ex *Hyloicus*, incomplete rearing — pupa).

Stenomidae

Antaeotricha leucillana (Zeller)

Frost, 1931: 1213, Pa. (ex *Stenoma algidella* Walker).

Thyatiridae

Pseudothyatira cymatophoroides (Guenée)

Schaffner and Griswold, 1934: 84 (incomplete rearing — pupa).

Tortricidae

Archips argyrospilus (Walker)

Prokopy, 1968: 350, Conn.; Schaffner, 1959: 30, [N.E. U.S.] (as Larvaevoridae).

Archips spp. (including *A. argyrospilus* (Walker) and/or *A. griseus* (Robinson))

Prokopy, 1968: 350, Conn.

Archips spp. probably

Dudley, 1919: 47, Maine (ex cherry ugly moths).

Choristoneura conflictana (Walker)

Wickman, 1963: 596, Calif.

Choristoneura fumiferana (Clemens)

Bean, 1958: 400-403, Minn. (as larvaevorids); Barker, 1947: 2-3, Man. (ex spruce budworm).

Choristoneura occidentalis Freeman

Williams, C. B., Walton and Tiernan, 1969: 310-311, Mont.

Choristoneura pinus Freeman

Barker, 1947: 3, Man. (ex jack pine budworm).

Choristoneura rosaceana (Harris)Ross, W. A., 1939: 52, Ont. (ex *Archips*); Sanderson and Jackson, 1909: 398-399 (ex *Archips*, incomplete rearing — egg); Coquillett, 1882: 14, Ill. (ex "*Laxotaenia*," incomplete rearing — egg); Snow, F. H., 1880: 202, Kans. (ex *Loxotaenia*).*Sparganothis pettitana* (Robinson)

Schaffner, 1959: 28, [N.E. U.S.] (as Larvaevoridae, incomplete rearing — pupa).

Yponomeutidae

Urodus parvulus (H. Edwards)Patton, 1958: 86, Fla. (ex *Trichostiba*).

Unidentified Lepidoptera

Miscellaneous

Walsh, 1866: 46, Maine (ex caterpillar on apple tree).

HOST: ORTHOPTERA

Acrididae

Ageneotettix deorum (Scudder)Smith, R. W., 1944: 29, Canada (as undetermined tachinid sp. may be *Acemya tibialis* Coquillett); Smith, R. W., 1944: 29, Canada (as undetermined tachinid sp. may be *Hemithrixion oestriforme* Brauer and Bergenstamm).*Camnula pellucida* (Scudder)Smith, R. W., 1944: 29, Canada (as undetermined tachinid sp. may be *Acemya tibialis*); Smith, R. W., 1944: 29, Canada (as undetermined tachinid sp. may be *Hemithrixion oestriforme*); Bruner, 1890a: 139, Idaho (incomplete rearing — larva).

Chimarocephala pacifica pacifica (Thomas)

Rentz, 1961: 60, Calif.

Dissosteira longipennis (Thomas)

Wakeland, 1958: 94, Colo.; *Wakeland, 1958: 92 (as long-legged *Tachina* fly, probable host, Bruner, 1897); Bruner, 1897: 38-39, Nebr. (as long-legged *Tachina* fly, probable host).

Melanoplus bivittatus (Say)

Washburn, 1912: 37, Minn.

Melanoplus dawsoni (Scudder)

Smith, R. W., 1944: 29, Canada (as undetermined tachinid sp. may be *Acemya tibialis*); Smith, R. W., 1944: 29, Canada (as undetermined tachinid sp. may be *Hemithrixion oestriforme*); Washburn, 1912: 37, Minn.

Melanoplus femurrubrum (De Geer)

Smith, R. W., 1944: 29, Canada (as undetermined tachinid sp. may be *Hemithrixion oestriforme* ex *M. "f. rubrum"*).

Melanoplus foedus Scudder

Smith, R. W., 1944: 29, Canada (as undetermined tachinid sp. may be *Acemya tibialis*); Smith, R. W., 1944: 29, Canada (as undetermined tachinid sp. may be *Hemithrixion oestriforme*).

Melanoplus packardii Scudder

Smith, R. W., 1944: 29, Canada (as undetermined tachinid sp. may be *Acemya tibialis*); Smith, R. W., 1944: 29, Canada (as undetermined tachinid sp. may be *Hemithrixion oestriforme*).

Melanoplus sanguinipes (Fabricius)

Smith, R. W., 1965: 190, 197, Ont.; Washburn, 1912: 37, Minn. (ex *Dissosteira carolina* (Linnaeus)); Washburn, 1912: 37, Minn. (ex *M. atlantis* (Riley)); Marlatt, 1889: 68, 70, N.H. (ex "*Melanopus*" *atlantis*, incomplete rearing — larva).

Melanoplus spretus (Walsh)

All records ex *Caloptenus spretus*.

Riley, Packard and Thomas, 1883: 319-323, Colo., Man., Minn.; Riley, 1876: 107-108, Kans., Mo. (incomplete rearing — egg or larva).

Melanoplus spp. (including *M. bivittatus* (Say) and/or *M. differentialis* (Thomas))

Osborn, 1891: 70, 74, Kans. (ex *Caloptenus differentialis* or *bivittatus*); *Osborn, 1891a: 51, 55-56, Kans. (ex *Caloptenus* spp. (*C. differentialis* and/or *C. bivittatus*), Osborn, 1891).

Melanoplus spp.

Criddle, 1911: 239, Man.

Orphulella speciosa (Scudder)

Washburn, 1912: 37, Minn.

Spharagemon collare (Scudder)

Washburn, 1912: 37, Minn.

Grasshopper(-s)

Corkins, 1924: 47; Washburn, 1902: 28, Minn.

Locust(-s)

Fletcher, 1897: 60-61, Ont. (ex red-legged locust, two-striped locust and/or lesser migratory locust); Riley, Packard and Thomas, 1883: 18-19, N.W. [U.S.].

Unidentified Acrididae

Milliken, R., 1893: 19, Idaho (incomplete rearing — larva).

Mantidae

Stagmomantis carolina (Linnaeus)

Riley, 1885: 100 (ex *Mantis*).

Phasmatidae

Bacillus rossia (Rossi)

Osten Sacken, 1877: 23 (ex *B. rossii* Fabricius, incomplete rearing — larva).

Diapheromera femorata (Say)

Osten Sacken, 1877: 23.

SYNOPSIS OF HOST-PARASITE RELATIONSHIPS

This section is based entirely on information contained in the next section on hosts. The class Insecta is listed first, followed by non-Insecta. The host orders, families, and genera are arranged alphabetically, followed by nonscientific names. The parasite names are arranged alphabetically, followed by the unassigned Tachinidae.

Class Insecta**Order COLEOPTERA****Family ALLECULIDAE**

Chromatia: *Hyalomyodes*.

Family BOSTRICHIDAE

Amphicerus: *Lixophaga*.

Family BUPRESTIDAE

Buprestis: Tachinidae.

Family CARABIDAE

Calosoma: *Sitophaga*.

Carabus: *Sitophaga*, Tachinidae.

Pterostichus: Tachinidae.

Family CERAMBYCIDAE

Enaphalodes: *Theresia*, Tachinidae.

Monochamus: *Eutheresia*, Tachinidae.

Oberea: *Lixophaga*.

Saperda: *Euphorocera*, *Ptilodexia*, Tachinidae.

Stenocorus: *Eutheresia*, *Ptilodexia*.

Unidentified: *Eutheresia*, *Zelia*.

Family CHRYSOMELIDAE

Acalymma: *Celatoria*, Tachinidae.

Agrioconota: *Eucelatoriopsis*.

- Altica: *Hyalomyodes*, *Medina*, Tachinidae.
 Blepharida: *Adoryphorophaga*.
 Calligrapha: *Euphorocera*, *Myiopharus*.
 Cerotoma: *Celatoria*.
 Chelymorpha: *Eribella*, *Eucelatoriopsis*.
 Chrysomela: *Grisdalemyia*, *Hyalomyodes*, Tachinidae.
 Coptocycla: *Eucelatoriopsis*, "Tachina."
 Crioceris: *Meigenia*, *Paralispe*.
 Diabrotica: *Celatoria*, *Pseudomyothyria*, Tachinidae.
 Disonycha: *Medina*.
 Erynephila: "Hypostena."
 Lema: *Paralispe*, *Patelloa*.
 Leptinotarsa: *Adoryphorophaga*, *Doryphorophaga*, *Winthemia*,
 Tachinidae.
 "Luperodes": *Medina*.
 Metriona: *Eucelatoriopsis*.
 Monocesta: *Ervia*.
 Monoxia: *Chaetonodexodes*.
 Oulema: *Hyalomyodes*.
 Physonota: Tachinidae.
 Plagiometriona: *Eribella*, *Eucelatoriopsis*, *Medina*.
 Pyrrhalta: *Aplomyiopsis*, *Erynniopsis*.
 Trirhabda: *Celatoria*.
 Chrysomelid (including *Lema* and/or *Leptinotarsa*): *Adoryphoro-*
phaga, *Doryphorophaga*.
 Cucumber beetle(-s): *Celatoria*.
 Unidentified: *Medina*.

Family CLERIDAE

- Thanasimus: Tachinidae.

Family COCCINELLIDAE

- Coccinella: *Hyalomyodes*.
 Coleomegilla: *Hyalomyodes*.
 Epilachna: *Aplomyiopsis*, *Doryphorophaga*, *Euphorocera*, *Helio-*
plagia, *Hyalomyodes*, *Lespesia*, *Nemorilla*, Tachinidae.

Family CURCULIONIDAE

- Ampelogypter: *Myiophasia*.
 Anthonomus: *Myiophasia*.
 Chalcodermus: *Myiophasia*.

- Conotrachelus:** *Cholomyia*, *Myiophasia*, Tachinidae.
Curculio: *Clausicella*, *Elfia*, *Myiophasia*, "Myiophasia," *Winthemia*.
Hypera: *Hyalomyodes*, *Lixophaga*, *Madremyia*.
Listroderes: *Epiplagiops*, *Hyalomyodes*.
Lixus: *Lixophaga*.
Pissodes: *Compsilura*.
Rhabdoscelus: *Lixophaga*, *Metadexia*.
Rhodoaenus: Tachinidae.
Sitona: *Hyalomyodes*.
Sphenophorus: *Lixophaga*, *Myiophasia*, Tachinidae.
Trichobaris: *Myiophasia*.
Caneborer(-s): *Myiophasia*.
Curculionid (including Conotrachelus or Curculio): *Myiophasia*.
Unidentified: *Myiophasia*, *Winthemia*.

Family ELATERIDAE

- Melanotus:** *Ateloglossa*.

Family ENDOMYCHIDAE

- Endomychus:** *Phyllophilopsis*.

Family LUCANIDAE

- Ceruchus:** *Eutheresia*.

Family MELOIDAE

- Epicauta:** *Pseudoperichaeta*.
Tetraonyx: *Gonia*.

Family PASSALIDAE

- Popilius:** *Zelia*.

Family SCARABAEIDAE

- Adoretus:** *Prosenia*.
Anomala: *Prosenia*.
Chemarachis: *Cryptomeigenia*, *Eutrixoides*, *Ptilodexia*.
Euetheola: *Eutrixia*, *Megapariopsis*, *Microphthalma*.
Lachnosterna: *Cryptomeigenia*.
Ligyris: *Eutrixia*.

Maladera: *Dexilla*, *Hyperecteina*.

Osmoderma: *Zelia*.

Phyllophaga: *Bonnetia*, *Cryptomeigenia*, *Dexilla*, *Dinera*, *Eutrixia*, *Eutrixoides*, *Gonia*, *Microphthalma*, *Ptilodexia*, "Ptilodexia," *Sitophaga*, Tachinidae.

Popillia: *Dexilla*, *Hamaxia*, *Hyperecteina*, *Prosenia*, *Ptilodexia*.

Serica: *Pseudomyothyria*, *Sitophaga*.

Xyloryctes: *Mystacella*.

June beetle(-s): *Eutrixia*, *Microphthalma*.

May beetle(-s): *Cryptomeigenia*, *Eutrixia*, *Ptilodexia*, Tachinidae.

Scarabaeid (allied genera of Phyllophaga): *Microphthalma*.

Scarabaeid (including Cyclocephala and/or Phyllophaga): *Microphthalma*.

Scarabaeid (including Phyllophaga or other Scarabaeidae): *Mochlosoma*.

Scarabaeid (including Phyllophaga or Phytalis): *Cryptomeigenia*, *Eutrixoides*.

Unidentified: *Hamaxia*.

White grub(-s): *Microphthalma*.

Family TENEBRIONIDAE

Eleodes: *Eleodiphaga*, *Sitophaga*, Tachinidae.

Meracantha: *Catagoniopsis*, *Pales*, *Zelia*.

UNIDENTIFIED COLEOPTERA

Rhynchophora: *Lixophaga*.

Unidentified: *Euphorocera*, "Macquartia," *Myiophasia*.

Order DERMAPTERA

Family CARCINOPHORIDAE

Anisolabis: Tachinidae.

Family FORFICULIDAE

Forficula: *Bigonicheta*, *Ocytata*.

Order DIPTERA

Family CECIDOMYIIDAE

Rhabdophaga: Tachinidae.

Family TABANIDAE

Chrysops: *Carinosillus*.

Hybomitra: *Carinosillus*, Tachinidae.

Tabanus: *Carinosillus*, *Ormia*, *Phasiops*, "*Phorostoma*," *Vibrissotheresia*.

Horse fly (flies): *Carinosillus*.

Unidentified: *Opsotheresia*, Tachinidae.

Family TIPULIDAE

Ctenophora: *Admontia*.

Tipula: *Admontia*, *Siphona*, Tachinidae.

Crane fly (flies): *Siphona*.

Tipulini: "*Sturmia*."

Unidentified: *Dexia*, *Pilatea*.

Order HEMIPTERA (HETEROPTERA)

Family COREIDAE

Acanthocephala: *Trichopoda*.

Anasa: *Trichopoda*, Tachinidae.

Archimerus: *Trichopoda*.

Chelinidea: *Trichopoda*.

Euthochtha: *Trichopoda*.

Harmostes: *Leucostoma*.

Leptoglossus: *Trichopoda*.

Narnia: *Trichopoda*.

Family CYDNIDAE

Sehirus: *Hyalomya*.

Family LARGIDAE

Largus: *Trichopoda*.

Family LYGAEIDAE

Emblethis: *Hyalomya*.

Eremocoris: *Catharosia*.

Geocoris: *Hyalomya*.

Lygaeus: *Leucostoma*.

Nysius: *Hyalomya*, "Phorantha."

Oncopeltus: *Leucostoma*.

Thylochromus: *Catharosia*.

Family MIRIDAE

Leptopterna: "Phorantha."

Lygocoris: *Alophorella*.

Lygus: *Alophorella*.

Family NABIDAE

Nabis: *Leucostoma*.

Family PENTATOMIDAE

Acrosternum: *Trichopoda*.

Alcaeorrhynchus: *Trichopoda*.

Chlorochroa: *Cylindromyia*, *Gymnosoma*, *Paraphorantha*, *Trichopoda*.

Cosmopepla: *Alophorella*, *Gymnoclytia*.

Eurygaster: *Gymnosoma*, *Phasia*.

Euschistus: *Cylindromyia*, *Euthera*, *Gymnoclytia*, *Gymnosoma*, *Trichopoda*.

Holcostethus: *Alophorella*.

Murgantia: *Trichopoda*.

Nezara: *Trichopoda*, Tachinidae.

Oebalus: *Euthera*.

Pitedia: *Gymnosoma*, *Trichopoda*.

Plautia: *Trichopoda*.

Podisus: *Hemyda*, *Trichopoda*.

Thyanta: *Trichopoda*.

Pentatomid (including *Euschistus*, *Mormidea*, *Nezara* and/or *Oebalus*): *Beskia*, *Cylindromyia*, *Euthera*.

Plant bug(-s): *Trichopoda*, Tachinidae.

Wheat bug(-s): *Gymnosoma*.

Family PYRRHOCORIDAE

Dysdercus: *Acaulona*.

Family RHOPALIDAE

Liorhyssus: *Leucostoma*, Tachinidae.

Family SCUTELLERIDAE

Coleotichus: *Trichopoda*.

UNIDENTIFIED HEMIPTERA (HETEROPTERA)

Unidentified: *Hyalomya*.

Order HEMIPTERA (HOMOPTERA)

Family APHIDIDAE

Macrosiphum: *Beskia*.

Family CERCOPIDAE

Philaenus: *Alophorella*.

Family CICADIDAE

Magiccicada: "*Tachina*," Tachinidae.

Order HYMENOPTERA

Family APIDAE

Bombus: *Clausicella*, *Lespesia*, "*Tachina*."

Family ARGIDAE

Arge: *Aporotachina*, *Bessa*, *Lydella* probably, *Spathimeigenia*, "*Sturmia*."

Sphacophilus: "*Hypostena*," *Schizocerophaga*, Tachinidae.

Sterictiphora: "*Hypostena*."

Family CIMBICIDAE

Cimbex: *Aplomya*, *Compsilura*, Tachinidae.

Trichiosoma: *Aplomya*.

Family DIPRIONIDAE

Diprion: *Bessa*, *Blondelia* probably, *Catagoniopsis*, *Compsilura*, *Diplostichus*, *Drino*, *Euphorocera*, *Guerinia*, *Spathimeigenia*, *Winthemia*, Tachinidae.

Monoctenus: *Spathimeigenia*, *Tachinomyia*.

Neodiprion: *Bessa*, *Diplostichus*, *Drino*, *Eribella*, *Eucelatoria*, *Euphorocera*, "Phorocera," *Spathimeigenia*, *Tsugaea*, "Zenilia."

Zadiprion: *Spathimeigenia*.

Diprionid (including Diprion and/or Neodiprion): *Cryptomeigenia*, *Diplostichus*, *Euphorocera*, "Phorocera," *Spathimeigenia*.

Family FORMICIDAE

Eciton: *Androeuropyops*.

Family PAMPHILIIDAE

Acantholyda: *Aplomya*.

Neurotoma: *Aplomya*, *Compsilura*, *Eubrachymera*.

Family SPHECIDAE

Bembix: "Prospheysa."

Family TENTHREDINIDAE

Allantus: *Guerinia*.

Amauronematus: *Bessa*.

Ametastegia: *Bessa*.

Anoplonyx: *Bessa*.

Cladius: *Bessa*.

Croesus: *Bessa*, *Compsilura*.

Hemichroa: *Bessa*, *Drino*.

Macremphytus: *Bessa*, *Exorista*, *Guerinia*.

Nematus: *Aporotachina*, *Bessa*, *Compsilura*, *Oswaldia*, Tachinidae.

Pikonema: *Bessa*, *Diplostichus*, *Drino*, *Phryxe*, *Spathimeigenia*.

Platycampus: *Bessa*.

Priophorus: *Bessa*.

Pristiphora: *Bessa*, *Compsilura*, *Diplostichus*, *Drino*, *Guerinia*, *Lespesia*, "Phorocera," *Spathimeigenia*.

Strongylogaster: *Bessa*.

Tethida: Tachinidae.

Tomostethus: *Hyalurgus*.

Trichiocampus: *Bessa*, *Compsilura*.

Unidentified: *Bessa*, *Carcelia*, "Gymnostylia," *Lixophaga*, *Pseudomyothyria*.

UNIDENTIFIED HYMENOPTERA

- Bee(-s): "*Tachina*."
 Pine sawfly(-flies): *Spathimeigenia*.
 Pitch pine sawfly(-flies): "*Phorocera*."
 Sawfly(-flies): *Athrycia*, *Compsilura*, *Drino*, *Euphorocera*, *Guerinia*, *Lixophaga*, Gen. near *Lixophaga*, *Spathimeigenia*.
 Tenthredinoidea: *Compsilura*.

Order LEPIDOPTERA

Family AEGERIIDAE

- Melittia: *Chaetophlepsis*.
 Sylvora: *Bonnetia*, *Copecrypta*.
 Synanthedon: *Parafischeria*, *Sipholeskia*.
 Vespamima: "*Tachina*," *Xanthophyto*.
 Unidentified: *Eulasiona*.

Family ARCTIIDAE

- Apantesis: *Carcelia*, *Euexorista*, *Euphorocera*, *Exorista*, *Gymnocarcelia*.
 Arachnis: *Carcelia*, *Leschenaultia*, *Sisyropa*.
 Cisthene: *Pseudochaeta*.
 Cycnia: *Archytas*, *Exorista*, *Hyphantrophaga*, Tachinidae.
 Diacrisia: *Aplomya*, *Blondelia*, *Bombyliopsis*, *Carcelia*, *Compsilura*, *Exorista*, *Gymnocarcelia*, *Hubneria*, *Lespesia*, *Mericia*, *Thelaira*, *Winthemia*, Tachinidae.
 Ecpantheria: *Aplomya*, *Archytas*, *Carcelia*, *Gymnocarcelia*, *Juriniopsis*, *Leschenaultia*, *Lespesia*.
 Estigmene: *Argyrophylax* probably, *Carcelia*, *Compsilura*, *Eucelatoria*, *Exorista*, *Gymnocarcelia*, *Hubneria*, *Leschenaultia*, *Lespesia*, *Mericia*, "*Phorocera*," *Sisyropa*, *Thelaira*, *Winthemia*, Tachinidae.
 Eubaphe: *Euphorocera*.
 Euchaetias: *Archytas*, *Bombyliopsis*, *Compsilura*, *Euexorista*, *Euphorocera*, *Exorista*, *Hyphantrophaga*, *Leschenaultia*, *Lespesia*, *Mericia*, *Winthemia*, *Zenillia*, Tachinidae.
 Eupseudosoma: *Lespesia*.
 Halisidota: *Anaporia*, *Blondelia*, *Bombyliopsis*, *Carcelia*, *Compsilura*, *Euphorocera*, *Exorista*, *Juriniopsis*, *Leschenaultia*, *Lespesia*, "*Macquartia*," "*Myiophasia*," "*Phorocera*," *Tachinomyia*, *Uramya*, *Winthemia*, "*Zenillia*," Tachinidae.

Haploa: "*Zenillia*."

Hemihyalea: *Leschenaultia*.

Hyphantria: *Archytas*, *Blondelia*, *Bombyliopsis*, *Carcelia*, *Ceromasia*, *Compsilura*, *Euphorocera*, *Eusisyropa*, *Exorista*, *Hyphantrophaga*, *Lespesia*, *Mericia*, "*Tachina*," *Winthemia*, Tachinidae.

Isia: *Euexorista*, *Exorista*, *Hubneria*, *Mystacella*, *Parachaeta*, *Thelaira*, *Winthemia*, Tachinidae.

Parasemia: *Thompsonomyia*.

Phragmatobia: *Carcelia*, *Hubneria*, Tachinidae.

Seirarctia: *Lespesia*.

Tigrioides: *Pseudochaeta*.

Turuptiana: *Carcelia*, *Chaetona*, "*Nemoraea*," *Thelaira*.

Tyria: *Lespesia*.

Utetheisa: *Euphorocera*, *Gymnocarcelia*, *Lespesia*, *Lixophaga*, Tachinidae.

Unidentified: *Bombyliopsis*, *Carcelia*, *Chaetogaedia*, *Compsilura*, "*Gonia*," *Gymnocarcelia*, *Hubneria*, *Leschenaultia*, *Lespesia*, "*Phorocera*," *Sisyropa*, *Thelaira*, *Winthemia*.

White woolly bear(-s): *Winthemia*.

Woolly bear(-s): *Leschenaultia*.

Family BLASTOBASIDAE

Valentinia: "*Hypostena*."

Family BOMBYCIDAE

Bombyx: *Compsilura*, *Lespesia*, *Zenillia*.

Unidentified: *Exorista*, *Lespesia*, *Pseudochaeta*, *Winthemia*.

Family BRASSOLIDAE

Brassolis: Tachinidae.

Opsiphanes: *Lespesia*.

Family CASTNIIDAE

Castnia: *Palpozenillia*.

Family CITHERONIIDAE

Anisota: *Belvosia*, *Compsilura*, *Eumasicera*, *Euphorocera*, *Eusisyropa*, *Hyphantrophaga*, *Lespesia*, "*Masicera*," *Oswaldia*, *Pacidianus*, "*Sturmia*," *Winthemia*, Tachinidae.

Citheronia: *Belvosia*, *Lespesia*, *Winthemia*.

Eacles: *Belvosia*, *Drino*, *Lespesia*, *Winthemia*, Tachinidae.

Family COLEOPHORIDAE

Coleophora: *Erynnia*.

Family COSSIDAE

Cossula: "*Phorocera*."

Family CTENUCHIDAE

Ceramidia: *Lespesia*.

Ctenucha: *Compsilura*, Tachinidae.

Lymire: *Euphorocera*, *Lespesia*.

Syntomeida: *Euphorocera*, *Lespesia*.

Family DANAIDAE

Danaus: *Buquetia*, *Chaetogaedia*, *Compsilura*, *Eusisyropa*, *Exorista*,
Lespesia, *Madremyia*, *Phryxe*, Tachinidae.

Family DIOPTIDAE

Phryganidia: *Actia*, *Eusisyropa*, *Siphona*, Tachinidae.

Family DREPANIDAE

Drepana: *Actia*, Tachinidae.

Family ETHMIIDAE

Ethmia: *Eufrontina*.

Family GELECHIIDAE

Anacamptis: *Lypha*.

Anarsia: *Erynnia*, *Euphorocera*, *Psalidopteryx*.

Aristotelia: *Schizactia*.

Chionodes: *Schizotachina*.

Coleotechnites: *Elfia*, *Schizactia*.

Exoteleia: *Schizactia*.

Filatima: *Eusisyropa*, *Nemorilla*, *Pseudomyothyria*.

Frumenta: *Lixophaga*.
 Gnorimoschema: *Actia*, *Frontiniella*.
 Isophrictis: "*Tachinophyto*."
 Pectinophora: *Erynnia*.
 "Telphusa" probably: *Psalidopteryx*.
 Gelechiid webber(-s): *Eusisyropa*.

Family GEOMETRIDAE

Alsophila: *Blondelia*, *Cyzenis*, *Gonia*, *Ictericophyto*, *Omotoma*,
Phryxe, *Pseudotachinomyia*, *Tachinomyia*, *Winthemia*,
 Tachinidae.
 Anacamptodes: *Blondelia*, *Eusisyropa*, *Lespesia*, "*Winthemia*."
 Biston: *Aplomya*, *Compsilura*, *Exorista*.
 Brephos: *Actia*, *Ictericophyto*.
 Caripeta: *Blondelia*.
 Cingilia: *Blondelia* probably, *Ceromasia*, *Chaetophlepsis*, *Compsi-*
lura, *Euphorocera*, *Eusisyropa*, *Madremyia*, *Phryxe*, *Winthemia*,
 Tachinidae.
 Coniodes: Tachinidae.
 Coryphista: *Blondelia*, *Madremyia*.
 Cosymbia: *Actia*.
 Deilinia: Tachinidae.
 Deuteronomos: *Madremyia*.
 Drepanulatrix: *Chaetophlepsis*, *Patelloa*.
 Ematurga: *Winthemia*.
 Ennomos: *Actia*, *Blondelia*, *Chaetogaedia*, *Compsilura*, *Euexorista*,
Euphorocera, *Eusisyropa*, *Lespesia*, *Madremyia*, *Mericia*,
Phryxe, *Tachinomyia*, *Xanthophyto* probably, Tachinidae.
 Erannis: *Ictericophyto*, *Phryxe*, *Pseudotachinomyia*, *Tachinomyia*,
Winthemia, Tachinidae.
 Eucaterva: *Hyphantrophaga*, Tachinidae.
 Eugonobapta: *Actia*, *Compsilura*.
 Eulype: Tachinidae.
 Eupithecia: *Bessa*, *Blondelia*, *Chaetogaedia*, *Clausicella*.
 Hydria: *Chrysoexorista*, *Compsilura*, *Drino*, *Eusisyropa*.
 Isturgia: *Eusisyropa*.
 Itame: *Blondelia*, *Chaetophlepsis*, *Compsilura*, *Euphorocera*,
Tachinomyia, Tachinidae.
 Lambdina: *Blondelia*, *Chaetophlepsis*, *Chrysoexorista*, *Compsilura*,
Eusisyropa, *Madremyia*, *Phryxe*, *Winthemia*, Tachinidae.
 Lycia: *Ceromasia*.
 Lygris: *Hyphantrophaga*.
 Melanchroia: *Euphorocera*, *Prophryno*.

- Melanolophia: *Actia*, *Compsilura*, *Eusisyropa*, *Ictericophyto*.
Metanema: *Lespesia*.
Nematocampa: *Compsilura*, Tachinidae.
Nepytiya: *Bessa*, *Blondelia*, *Chaetophlepsis*, *Eusisyropa*, Tachinidae.
Operophtera: *Cyzenis*, *Phryxe*.
Paleacrita: Tachinidae.
Pero: *Chaetophlepsis*.
Phaeoura: *Aplomya*, Tachinidae.
Phigalia: *Ceromasia*, *Compsilura*, *Eusisyropa*, *Omotoma*,
Tachinomyia, Tachinidae.
Philtraea: *Euphorocera*.
Protitame: *Chaetophlepsis*.
Sabulodes: *Xanthophyto*, Tachinidae.
Scotorythra: *Chaetogaedia*, *Lespesia*.
Semiothisa: *Bessa*, *Blondelia*, *Chaetophlepsis*, *Euphorocera*,
Prophryno.
Synchlora: *Phryxe*.
Thera: *Blondelia*.
Cankerworm(-s): *Winthemia*.
Cankerworm(-s) (including *Alsophila* and/or *Paleacrita*): "*Tachina*."
Unidentified: *Aplomya*, *Bessa*, *Compsilura*, *Eucelatoria*, *Lespesia*,
Winthemia, *Xanthoepalpus*.

Family GLYPHIPTERYGIDAE

- Anthophila: *Hemisturmia*, *Nemorilla*.
Lotisma: *Elfia*.

Family GRACILLARIIDAE

- Gracillaria: *Aplomya*.

Family HEPIALIDAE

- Hepialus: Tachinidae.

Family HESPERIIDAE

- Autochton: *Lespesia*.
Calpodes: *Eucelatoria*, *Eusisyropa*, *Lixophaga*.
Cymaenes: Tachinidae.
Epargyreus: *Acroglossa*, *Compsilura*, *Euphorocera*, *Eusisyropa*,
Juriniopsis, *Lespesia*, Tachinidae.
Erynnis: *Compsilura*, *Drino*, *Eusisyropa*, Tachinidae.

Hylephila: *Stomatomyia*.

Lerodea: *Euphorocera*.

Ochlodes: *Spathidexia*.

Thymelicus: *Compsilura*, *Euexorista*, *Helioplagia*, *Phryxe*,
Winthemia, Tachinidae.

Urbanus: *Chrysotachina*, *Lespesia*, *Nemorilla*.

Hesperiid (including *Cymaenes*, *Nyctelius* and/or *Pan-*
oquina): Tachinidae.

Unidentified: *Belvosia*, *Compsilura*.

Family LACOSOMIDAE

Cicinnus: *Chrysotachina*.

Family LASIOCAMPIDAE

Epicnaptera: *Compsilura*, *Exorista*, Tachinidae.

Gloveria: *Aplomya*, *Hyphantrophaga*.

Malacosoma: *Archytas*, *Blepharipa*, *Carcelia*, *Cartocometes*,
Chaetogaedia, *Compsilura*, *Cyzenis*, *Diplostichus*, *Drino*,
Euexorista, *Euphorocera*, *Exorista*, *Hyphantrophaga*,
Leschenaultia, *Lespesia*, *Madremyia*, *Mericia*, *Parasetigena*,
Patelloa, *Pilatea*, "*Tachina*," *Tachinomyia*, *Uramya*, *Winthemia*,
Xanthophyto, Tachinidae.

Tolype: *Exorista*, *Winthemia*.

American tent caterpillar(-s): *Carcelia*, *Compsilura*, *Lespesia*,
Tachinidae.

Apple tree tent caterpillar(-s): *Euphorocera*, *Exorista*.

Tent caterpillar(-s): *Archytas*, *Exorista*, *Leschenaultia*, *Lespesia*,
Winthemia.

Family LIMACODIDAE

Adoneta: *Carcelia*.

Cnidocampa: *Chaetexorista*, *Compsilura*, *Lixophaga*.

Euclea: *Anaporia*, *Eusisyropa*, Tachinidae.

Limacodes: *Anaporia*, Tachinidae.

Lithacodes: *Anaporia*.

Packardia: *Eupelecotheca*.

Phobetron: *Anaporia*, *Carcelia*, *Palpexorista*, Tachinidae.

Prolimacodes: Tachinidae.

Sibine: *Anaporia*, *Euphorocera*, *Lespesia*, *Palpexorista*, Tachinidae.

Sisyrosea: *Anaporia*.

Tortricidia: *Anaporia*.

Family LYCAENIDAE

- Brephidium:** *Aplomya*, Tachinidae.
Celastrina: *Aplomya*.
Lampides: *Lespesia*.
Lycaeides: *Aplomya*.
Lycaena: *Aplomya*.
Plebejus: Tachinidae.
Strymon: *Aplomya*, *Lespesia*, *Nemorilla*.
Unidentified: *Aplomya*.

Family LYMANTRIIDAE

- Byrdia:** *Exorista*, *Peleteria*, *Petinarctia*.
Dasychira: *Carcelia*, *Compsilura*, *Exorista*, *Spoggosia*.
Euproctis: *Blepharipa*, *Carcelia*, *Compsilura*, *Euphorocera*,
Eusisyropa, *Exorista*, *Leschenaultia*, *Madremyia*, *Patelloa*,
Phrynofrontina, *Townsendiellomyia*, *Zenillia*.
Hemerocampa: *Bessa*, *Carcelia*, *Compsilura*, *Drino*, *Euphorocera*,
Exorista, "Exorista," *Leschenaultia*, *Lespesia*, *Patelloa*, *Pele-*
teria, "Phorocera," *Sisyropa*, *Tachinomyia*, *Winthemia*,
Tachinidae.
Nygmia: *Aplomya*, *Carcelia*, *Compsilura*, *Euphorocera*, *Eusisyropa*,
Exorista, *Gonia*, *Lespesia*, *Madremyia*, *Patelloa*, *Townsend-*
iellomyia.
Orgyia: *Carcelia*, *Compsilura*, *Eusisyropa*, *Exorista*, *Lespesia*,
Tachinidae.
Porthetria: *Blepharipa*, *Carcelia*, *Compsilura*, *Drino*, *Euphorocera*,
Eusisyropa, *Exorista*, *Lespesia*, *Nemorilla*, *Parasetigena*, *Phryxe*,
Tachinomyia, *Townsendiellomyia*, Tachinidae.
Stilpnobia: *Blepharipa*, *Compsilura*, *Euphorocera*, *Eusisyropa*,
Exorista, *Lespesia*, *Parasetigena*, *Tachinomyia*, *Winthemia*,
Tachinidae.
Tussock moth(-s): *Carcelia*, *Compsilura*, *Euphorocera*, *Exorista*,
Lespesia, *Thelymyia*, *Winthemia*, Tachinidae.
Unidentified: *Compsilura*.

Family MEGALOPYGIDAE

- Lagoa:** *Carcelia*, *Euphorocera*.
Megalopyge: *Anaporina*, *Archytas*, *Carcelia*, *Compsilura*, *Euphoro-*
cera, *Hyphantrophaga*, *Lespesia*, *Sisyropa*, *Winthemia*,
"Zenillia," Tachinidae.

Family MEGATHYMIDAE

Agathymus: "*Phorocera*."

Megathymus: *Lespesia*, "*Phorocera*."

Family NOCTUIDAE

Achaea: *Chaetogaedia*, *Eucelatoria*, "*Winthemia*."

Achatodes: *Lydella*, *Nemorilla*.

Acronicta: *Archytas*, *Carcelia*, *Compsilura*, *Eumasicera*, *Euphorocera*, *Eusisyropa*, *Exorista*, "*Exorista*," *Lespesia*, *Mystacella*, "*Nemorilla*," *Nilea*, *Sisyropa*, *Tachinomyia*, *Winthemia*, "*Zenillia*," Tachinidae.

Actebia: *Chaetogaedia*, *Euphorocera*, *Periscepsia*, Tachinidae.

Agrotis: *Archytas*, *Bonnetia*, *Carcelia*, *Chaetogaedia*, *Eucelatoria*, *Euphorocera*, *Gonia*, *Lespesia*, "*Linnaemya*," *Madremyia*, *Mericia*, *Peleteria*, *Periscepsia*, *Sisyropa*, *Tachinomyia*, Tachinidae.

Alabama: *Beskia*, *Euphorocera*, *Eusisyropa*, "*Exorista*," *Lespesia*, *Patelloa*, "*Tachina*," *Triachora*, Tachinidae.

Aletia: "*Prosopaea*," Tachinidae.

Alypia: *Aplomya*, *Carcelia*, *Chaetophlepsis*, *Compsilura*, *Drino*, *Lespesia*, *Sisyropa*, *Winthemia*, Tachinidae.

Amathes: *Euphorocera*, *Linnaemya*, *Periscepsia*, *Winthemia*.

Amphipyra: *Compsilura*, *Mericia*, *Omotoma*, *Winthemia*, Tachinidae.

Amyna: *Eucelatoria*.

Anomis: *Chaetogaedia*, *Eusisyropa*, *Lespesia*, *Lixophaga*.

Anticarsia: *Euphorocera*, *Gonia*, *Voria*, *Winthemia*, Sturmiini.

Apamea: *Euexorista*, *Nowickia*.

Archanara: *Ceromasia*, *Lydella*.

Arzama: *Erynnia*, *Lydella*, *Oswaldia* probably.

Autographa: *Aplomya*, *Chaetogaedia*, *Chaetophlepsis*, *Eucelatoria*, *Euexorista*, *Lespesia*, *Madremyia*, *Periscepsia*, *Pilatea*, *Siphona*, *Voria*.

Autoplusia: *Chaetogaedia*, *Voria*.

Baileya: *Tachinomyia*.

Barathra: *Mericia*.

Bomolocha: *Blondelia*.

Caenurgina: *Winthemia*.

Callopistria: Tachinidae.

Calpe: *Compsilura*, *Winthemia*.

Carneades: *Chaetogaedia*.

Catabena: *Deopalpus*.

- Catocala: *Actia*, *Archytas*, *Blepharipa*, *Chaetophlepsis*, *Compsilura*, *Eusisyropa*, *Tachinomyia*, *Winthemia*, Tachinidae.
- Celama: *Lespesia*.
- Ceramica: *Blondelia*, *Compsilura*, *Winthemia*, Tachinidae.
- Choephora: *Wagneria*.
- Chrysodeixis: *Chaetogaedia*, *Eucelatoria*, Tachinidae.
- Cirphis: *Archytas*, *Periscepsia*, Tachinidae.
- Copablepharon: *Bonnetia*.
- Cosmia: *Mericia*, Tachinidae.
- Crocigrapha: *Patelloa*.
- Crymodes: *Gonia*, *Nowickia*, Tachinidae.
- Cryphia: *Chaetophlepsis*.
- Cucullia: *Lespesia*, *Winthemia*, Tachinidae.
- Elaphria: *Archytas*, *Periscepsia*.
- Elydna: *Chaetogaedia*, *Lespesia*.
- Epiglaea: *Cyrtophleba*, *Madremyia*, *Periscepsia*, *Tachinomyia*, *Winthemia*, Tachinidae.
- Eubolina probably: *Euphorocera*.
- Euparthenos: Tachinidae.
- Eupsilia: *Cyrtophleba*, *Mericia*, *Periscepsia*, Tachinidae.
- Euthisanotia: *Chaetophlepsis*, *Compsilura*, *Sisyropa*, *Winthemia*, Tachinidae.
- Euxoa: *Aphria*, *Bonnetia*, *Euphorocera*, *Gonia*, *Lespesia*, *Mericia*, *Nowickia*, *Peleteria*, *Periscepsia*, *Winthemia*, Tachinidae.
- Faronta: *Euphorocera*, *Lespesia*, *Periscepsia*, *Winthemia*, Tachinidae.
- Feltia: *Bonnetia*, *Euphorocera*, *Gonia*, *Lespesia*, *Spallanzania*, *Triachora*, *Winthemia*.
- Glaea: *Periscepsia*.
- Heliophila: *Archytas*, *Peleteria*.
- Heliothis: *Archytas*, *Athanatus*, *Carcelia*, *Chaetogaedia*, *Eucelatoria*, *Euphorocera*, *Gonia*, *Gymnochaetopsis*, *Hyphantrophaga*, *Lespesia*, *Metaplagia*, *Nemorilla*, *Palexorista*, *Plagiomima*, *Voria*, *Winthemia*, Tachinidae.
- Heptagrotis: *Phryxe*.
- Hydroecia: *Euphorocera*, *Lespesia*, *Lydella*.
- Hypena: *Blondelia*, *Ceromasia*, *Eusisyropa*.
- Hyppa: *Euexorista*.
- Hypsoropha: *Lespesia*.
- Lacinipolia: *Chaetogaedia*.
- Laphygma: *Archytas*.
- Lascoria: *Periscepsia*.
- Leucania: *Archytas*, *Eucelatoria*, *Peleteria*, *Periscepsia*, *Triachora*, *Winthemia*.

- Lithophane:** *Blepharipa*, *Blondelia*, *Carcelia*, *Compsilura*, *Epalpus*, *Euphorocera*, *Gonia*, *Mericia*, *Omotoma*, *Periscepsia*, *Phryxe*, *Tachinomyia*, *Winthemia*, Tachinidae.
- Litoprosopus:** Gen. near *Chaetogaedia*.
- Luperina:** *Lydella*, *Winthemia*.
- Macronoctua:** *Lydella*.
- Mamestra:** Tachinidae.
- Melipotis:** *Euphorocera*.
- Mocis:** *Archytas*, *Atacta*, *Belvosia*, *Eucelatoria*, *Euphorocera*, *Eusisyropa*, *Gonia*, *Gymnochaetopsis*, *Lespesia*.
- Nephelodes:** *Aplomya*, *Euexorista*, *Phryxe*, *Tachinomyia*, Tachinidae.
- Nycteola:** *Euexorista*, *Lespesia*, *Pseudoperichaeta*, *Zenillia*.
- Ogdoconta:** *Copecrypta*.
- Oligia:** *Winthemia*.
- Orthodes:** Tachinidae.
- Orthosia:** *Tachinomyia*, *Wagneria*.
- Papaipema:** *Aplomya*, *Ceromasia*, *Eusisyropa*, *Exorista*, *Gymnocheta*, *Lixophaga*, *Lydella*, "Phorocera," "Tachina," *Winthemia*, Tachinidae.
- Parallelia:** *Oswaldia*, Tachinidae.
- Peridroma:** *Archytas*, *Bonnetia*, *Carcelia*, *Chaetogaedia*, *Clausicella*, *Eucelatoria*, *Euphorocera*, *Gonia*, *Lespesia*, *Madremyia*, *Peleteria*, *Periscepsia*, *Voria*, *Winthemia*, Tachinidae.
- Phosphila:** *Compsilura*, *Eusisyropa*.
- Plathypena:** *Archytas*, *Carcelia*, *Chaetophlepsis*, *Compsilura*, *Copecrypta*, *Euphorocera*, *Eusisyropa*, *Lespesia*, *Prophryno*, *Winthemia*.
- Playtysenta:** *Winthemia*.
- Plusia:** *Eucelatoria*, *Voria*.
- Plusiodonta:** *Compsilura*.
- Polia:** *Bonnetia*, *Compsilura*, *Eusisyropa*, *Nowickia*, *Periscepsia*.
- Polydesma:** *Chaetogaedia*, *Eucelatoria*, *Lespesia*.
- Prodenia:** *Gonia*, *Nemorilla*.
- Protorthodes:** *Chaetogaedia*, *Eucnephalia* probably, *Wagneria*.
- Pseudaletia:** *Archytas*, *Athrycia*, *Carcelia*, *Chaetogaedia*, *Compsilura*, *Cylindromyia*, *Eucelatoria*, *Euphorocera*, *Exorista*, *Gonia*, *Gymnocarcelia*, *Gymnoclytia*, *Lespesia*, *Madremyia*, *Myiophasia*, *Patelloa*, *Peleteria*, *Periscepsia*, *Phryxe*, "Tachina," *Triachora*, *Winthemia*, Tachinidae.
- Pseudoplusia:** *Chaetophlepsis*, *Euphorocera*, *Lespesia*, *Winthemia*.
- Psychomorpha:** *Sisyropa*, "Zenillia."
- Pyreferra:** *Exorista*, *Ictericophyto*, *Omotoma*, Tachinidae.
- Pyrophila:** *Leschenaultia*.

- Raphia: *Tachinomyia*.
Rhodophora: *Compsilura*.
Schinia: *Winthemia*.
Scoliopteryx: *Compsilura*, *Winthemia*.
Scotogramma: *Euphorocera*, *Gonia*, *Lespesia*.
Sideridis: *Euexorista*.
Simyra: *Compsilura*, *Eusisyropa*, *Exorista*, *Lespesia*, "Masicera."
Spaelotis: *Periscepsia*, *Winthemia*.
Spodoptera: *Admontia*, *Archytas*, *Bonnetia*, *Carcelia*, *Chaetogaedia*, *Choeteprosopa*, *Eucelatoria*, *Euphorocera*, *Eusisyropa*, *Gonia*, *Hyphantrophaga*, *Lespesia*, *Lixophaga*, *Nemorilla*, *Omotoma*, *Palexorista*, *Voria*, *Winthemia*, Tachinidae.
Stibadium: *Lixophaga*.
Trichoplusia: *Archytas*, *Compsilura*, *Eucelatoria*, *Eusisyropa*, *Lespesia*, *Lixophaga*, *Madremyia*, *Nemorilla*, *Periscepsia*, "Phorocera," *Phryxe*, *Siphona*, "Tachina," *Voria*, *Winthemia*, Tachinidae.
Xanthopastis: *Lespesia*.
Xerociris: *Lespesia*.
Xylena: *Compsilura*, *Madremyia*, *Wagneria*, *Winthemia*.
Agrotidian(-s): *Winthemia*.
Armyworm(-s): *Archytas*, *Bonnetia*, *Chaetogaedia*, *Eucelatoria*, *Euphorocera*, *Lespesia*, *Patelloa*, *Peleteria*, *Periscepsia*, "Phorocera," *Triachora*, *Winthemia*, Tachinidae.
Armyworm(-s) (including *Pseudaletia* and/or *Spodoptera*): *Archytas*, *Chaetogaedia*, *Lespesia*.
Cabbage cutworm(-s): *Gonia*.
Catocalinae: *Actia*, *Archytas*, *Blepharipa*, *Compsilura*, *Eusisyropa*, *Tachinomyia*, *Winthemia*.
Cutworm(-s): *Archytas*, *Bonnetia*, *Chaetogaedia*, *Eucelatoria*, *Euphorocera*, *Gonia*, *Lespesia*, *Nowickia*, *Periscepsia*, "Tachina," *Tachinomyia*, *Winthemia*, Tachinidae.
Cutworm(-s) and related Noctuidae: *Winthemia*.
Cutworm(-s) (including *Agrotis* and/or *Leucania*): *Chaetogaedia*, *Lespesia*.
Noctuid (including *Heliopsis*, *Prodenia* and/or *Spodoptera*): *Winthemia*.
Noctuid (including *Mamestra* and/or *Peridroma*): *Lespesia*, *Phryxe*, *Winthemia*.
Phalaenid moth(-s): *Voria*.
Unidentified: *Archytas*, *Blepharipa*, *Bonnetia*, *Chaetogaedia*, *Chaetophlepsis*, *Compsilura*, *Cyrtophleba*, *Gymnocarcelia*, *Lespesia*, *Mericia*, *Nilea*, *Peleteria*, *Periscepsia*, *Phryxe*, *Siphona*, *Tachinomyia*, *Voria*, *Winthemia*, Gen. near *Zenillia*, Tachinidae.

Family NOCTUIDAE and/or PYRALIDAE

Noctuid and/or pyralid (including *Scotogramma* and/or *Loxostege*): *Euphorocera*, *Lespesia*, *Voria*.

Family NOTODONTIDAE

Cerura: *Aplomya*, *Archytas*, *Compsilura*, *Exorista*, *Lespesia*, *Pales*, "*Prosopaea*," Tachinidae.

Dasylophia: *Lespesia*, Tachinidae.

Datana: *Archytas*, *Belvosia*, *Blepharipa*, *Carcelia*, *Compsilura*, *Euphorocera*, *Exorista*, *Lespesia*, *Mericia*, *Palpexorista*, *Patelloa*, *Winthemia*, Tachinidae.

Dicentria: *Winthemia*.

Fentonia: *Compsilura*, *Sisyropa*.

Gluphisia: *Eusisyropa*.

Heterocampa: *Actia*, *Blondelia*, *Compsilura*, *Lespesia*, *Patelloa*, *Sisyropa*, *Spathimeigenia*, *Winthemia*, Tachinidae.

Hyperaeschra: *Winthemia*, Tachinidae.

Ichthyura: *Compsilura*, *Euphorocera*, *Eusisyropa*, *Exorista*, *Lespesia*, *Winthemia*, *Zenillia*, Tachinidae.

Nadata: *Compsilura*, Tachinidae.

Pheosia: *Aplomya*, *Chaetophlepsis*, *Compsilura*, *Winthemia*.

Schizura: *Actia*, *Carcelia*, *Compsilura*, *Euphorocera*, *Exorista*, *Lespesia*, *Pseudoperichaeta*, *Winthemia*, Tachinidae.

Symmerista: *Blepharipa*, *Compsilura*, *Euphorocera*, *Winthemia*, Tachinidae.

Unidentified: *Compsilura*.

Family NYMPHALIDAE

Agraulis: *Compsilura*, *Euphorocera*.

Anaea: *Euphorocera*, *Lespesia*.

Asterocampa: *Euphorocera*, *Lespesia*.

Chlosyne: *Compsilura*, *Euphorocera*, "*Phorocera*," Tachinidae.

Euphydryas: *Compsilura*, *Siphosturmia*.

Limenitis: *Compsilura*, Tachinidae.

Nymphalis: *Bessa*, *Compsilura*, *Dichaetoneura*, *Doryphorophaga*, *Euexorista*, *Euphorocera*, *Eusisyropa*, *Exorista*, *Hemisturmia*, *Hyphantrophaga*, *Lespesia*, *Madremyia*, *Patelloa*, *Pelatachina*, "*Phorocera*," *Winthemia*, Tachinidae.

Phyciodes: *Siphosturmia*.

Polygonia: *Compsilura*, *Euphorocera*, *Eusisyropa*, *Exorista*, *Lespesia*, Tachinidae.

Speyeria: *Madremyia*.

Vanessa: *Chaetogaedia*, *Compsilura*, *Euexorista*, *Eusisyropa*,
Lespesia, *Madremyia*, *Metopiops*, *Pelatachina*, *Sisyropa*.
 Unidentified: *Compsilura*.

Family OECOPHORIDAE

Dasystoma: *Compsilura*.
Depressaria: *Dichaetoneura*, *Eusisyropa*, *Nemorilla*, *Xanthophyto*,
 Tachinidae.
Psilocorsis: *Elfia*, *Eulasiona*, *Eusisyropa*, *Psalidopteryx*, Tachinidae.

Family OLETHREUTIDAE

Ancylis: *Erynnia*, *Hemisturmia*, *Lixophaga*, *Nemorilla*.
Barbara: *Eribella*.
Endothenia: *Clausicella*.
Epiblema: *Clausicella*, *Dichaetoneura*, *Lixophaga*, *Lydella*,
 "Tachinophyto."
Epinotia: *Actia*, *Elfia*, *Eulasiona*, *Psalidopteryx*, *Schizactia*.
Episimus: *Actia*, *Nemorilla*, *Pseudoperichaeta*.
Eucosma: *Coloradomyia*, *Leskiomima*, *Lespesia*, *Psalidopteryx*.
Eumarozia: *Lixophaga*.
Evora: *Erynnia*, *Nemorilla*, *Pseudoperichaeta*.
Exartema: *Nemorilla*.
Grapholitha: *Actia*, *Admontia*, *Dichaetoneura*, *Elephantocera*,
Elodia, *Erynnia*, *Leskiella*, *Leskiomima*, *Lespesia*, *Lixophaga*,
Nemorilla, *Pseudoperichaeta*, *Strobliomyia*, Tachinidae.
Gretchena: *Eusisyropa*.
Griselda: *Psalidopteryx*.
Laspeyresia: *Aplomya*, *Chrysoexorista*, *Erynnia*, *Leskiella*,
Leskiomima, *Lixophaga*, "Myiophasia," *Nilea*, "Tachino-
 phyto."
Melissopus: *Carcelia*, *Elfia*, *Erynnia*, *Pseudoperichaeta*.
Petrova: *Lixophaga*, Gen. near *Oxynops*, *Phrynofrontina*.
Proteoteras: *Erynnia*.
Pseudexentera: *Cyzenis*, "Cyzenis," *Eulasiona*.
Rhopobota: *Nemorilla*.
Rhyacionia: *Erynnia*, *Lixophaga*, *Nemorilla*, *Urophylopsis*,
 Tachinidae.
Sciaphila: *Ceromasia*, *Lypha*, *Madremyia*, *Xanthophyto*.
Silonota: *Actia*, *Blondelia*, *Mericia*, *Nemorilla*, *Pseudoperichaeta*,
Winthemia.
Suleima: *Clausicella*, *Lixophaga*.
Zeiraphera: *Phytomyptera*.
 Unidentified: *Leskiella*.

Family OLETHREUTIDAE and/or PYRALIDAE

Olethreutid and/or pyralid (including *Laspeyresia* and/or *Dioryctria* probably): *Exorista*.

Family PAPILIONIDAE

Battus: *Compsilura*.

Papilio: *Buquetia*, *Compsilura*, *Lespesia*, Tachinidae.

Unidentified: *Compsilura*.

Family PIERIDAE

Ascia: *Prophryno*.

Colias: *Euphorocera*, *Lespesia*, "*Masicera*," "*Phorocera*."

Eucheira: *Eucheirophaga*, *Eusisyropa*, *Winthemia*.

Pieris: *Ceromasia*, *Chaetogaedia*, *Compsilura*, *Eucelatoria*, *Euphorocera*, *Gymnoclytia*, *Lespesia*, *Madremyia*, *Obolocera*, *Phryxe*, Tachinidae.

Zerene: Tachinidae.

Family PLUTELLIDAE

Harpipteryx: Tachinidae.

Family PSYCHIDAE

Oiketicus: *Lespesia*.

Thyridopteryx: *Archytas*, *Drino*, *Erynnia*, *Euphorocera*, *Eusisyropa*, *Patelloa*, "*Tachina*," Tachinidae.

Family PTEROPHORIDAE

Oidaematophorus: *Compsilura*, *Oxynops*, *Pseudosiphona*, *Tachinophyto*, Tachinidae.

Platyptilia: *Elfia*, *Hyalomyodes*.

Pterophorus: *Hemisturmia*.

Trichoptilus: Tachinidae.

Unidentified: *Eribella*.

Family PYRALIDAE

Acigona: "*Masicera*."

Acrobasis: *Didyma*, *Elfia*, *Eusisyropa*, *Exorista*, *Hemisturmia*,

- Hyphantrophaga*, *Leskiomima*, "Myiophasia," *Nemorilla*,
Schizotachina, "Tachina," "Zenillia," Tachinidae.
- Agathodes:** *Eucelatoria*.
- Alatuncusia:** *Eusisyropa*.
- Alberada:** *Clausicella*.
- Asciodes:** *Nemorilla*.
- "Blepharomastix":** *Myobiopsis*.
- Canarsia:** *Eufrontina*.
- Chilo:** *Paratheresia*.
- Crambus:** *Aplomya*, *Euphorocera*, *Ormia* probably, *Phasiops*,
Stomatomyia, "Tachina," Tachinidae.
- Desmia:** *Dejeaniopalpus*, *Erynnia*, *Exorista*, *Hyphantrophaga*,
Leskiomima, *Lixophaga*, *Nemorilla*, *Pseudoperichaeta*,
"Tachina," "Tachinophyto."
- Diaphania:** "Leskia," *Nemorilla*, *Stomatodexia*, *Thelaira*, Tachini-
dae.
- Diatraea:** "Hypostena," *Lespesia*, *Lixophaga*, *Metagonistylum*,
Nemorilla, *Palpozenillia*, *Paratheresia*, "Phorocera," "Phoros-
toma," "Tachinophyto," Tachinidae.
- Dichogama:** *Eusisyropa*.
- Dioryctria:** *Aplomya*, *Elfia*, *Leskiomima*, Gen. near *Oxynops*,
Phrynofrontina, *Phryxe*, *Xanthophyto*, Tachinidae.
- Elasmopalpus:** *Stomatomyia*.
- Etiella:** *Argyrophylax*, *Erynnia*.
- Eulepte:** *Argyrophylax*, *Carcelia*.
- Evergestis:** *Compsilura*, *Lespesia*, *Phryxe*.
- Galleria:** *Ceromasia*, *Lydella*.
- Hedylepta:** *Argyrophylax*, *Chaetogaedia*, *Drino*, *Eucelatoria*,
Lespesia, *Nemorilla*, Tachinidae.
- Hellula:** *Nemorilla*, *Stomatomyia*.
- Herpetogramma:** *Eucelatoria*, *Nemorilla*.
- Homoesoma:** *Clausicella*, *Erynnia*, *Leskiomima*, *Lixophaga*.
- Hulstia:** *Nemorilla*, *Stomatomyia*.
- Hymenia:** *Argyrophylax*, *Chaetogaedia*, *Eucelatoria*, *Nemorilla*.
- Laniifera:** *Lespesia*, "Phorocera."
- Loxostege:** *Aplomya*, *Blondelia*, *Euphorocera*, *Eusisyropa*,
Hyphantrophaga, *Lespesia*, *Lixophaga*, *Nemorilla*, *Oswaldia*,
Patelloa, "Phorocera," *Phryxe*, *Pseudochaeta*, *Pseudoperi-
chaeta*, *Stomatomyia*, *Winthemia*, Tachinidae.
- Macrobotys:** *Leskiomima*, *Myobiopsis*, *Pseudochaeta*, *Pseudo-
perichaeta*, Tachinidae.
- Maruca:** *Argyrophylax*, *Nemorilla*.
- Melitara:** *Coronimyia*, *Lespesia*, "Phorocera."
- Moodna:** *Coronimyia*, *Erynnia*.

- Nephopteryx: *Aphria*, *Eusisyropa*, *Nemorilla*, *Phryxe*, Tachinidae.
 Oebia: *Chaetogaedia*.
 Olycella: *Lespesia*, "Phorocera."
 Omphalocera: *Carcelia*, *Hyphantrophaga*.
 Ostrinia; "Anisia," *Aplomya*, "Carcelia," *Ceromasia*, *Compsilura*,
Didyma, *Euexorista*, "Hypostena," *Lixophaga*, *Lydella*,
Madremyia, *Nemorilla*, *Paratheresia*, "Phorocera," *Phryxe*,
Pseudoperichaeta, *Pyraustomyia*, "Zenillia."
 Pachyzancla: *Nemorilla*, *Pseudoperichaeta*.
 Palpita: *Pseudochaeta*.
 Pantographa: *Carcelia*, *Pseudoperichaeta*.
 Parargyractis: *Ginglymia*.
 Pempelia: *Hyphantrophaga*.
 Phlyctaenia: *Aplomya*, *Nemorilla*, *Pseudoperichaeta*, Tachinidae.
 Pilemia: *Argyrophylax*.
 Pilocrocis: *Argyrophylax*, *Nemorilla*.
 Plodia: *Clausicella*.
 Pococera: *Nemorilla*.
 Psara: *Argyrophylax*, *Nemorilla*.
 Pyralis: *Elfia*, *Winthemia*.
 Pyrausta: *Aplomya*, *Dejeaniopalpus*, *Elfia*, *Hyphantrophaga*,
Lespesia, *Lixophaga*, *Nemorilla*, *Pseudoperichaeta*, *Pyrausto-*
myia, *Stomatomyia*, Tachinidae.
 Tetralopha: *Eufrentina*, *Eusisyropa*, *Frontiniella*, *Genea*,
Leskiomima, *Metopiops*, *Nemorilla*, Tachinidae.
 Titanio: *Eucoronimyia*.
 Tlascala: *Aphria*.
 Udea: "Phorocera."
 Zeadiatraea: *Lixophaga*, *Paratheresia*, "Phorostoma."
 Cone moth(-s): *Elfia*, *Lydellohoughia*.
 Phycitinae: *Coronimyia*.
 Unidentified: *Drino*, *Hyphantrophaga*, *Leskiomima*, *Lespesia*,
Lixophaga, *Metopiops*, *Nemorilla*.

Family PYRALIDAE or TORTRICIDAE

Unidentified: *Coronimyia*.

Family SATURNIIDAE

- Actias: *Cylindromyia*, *Lespesia*, *Winthemia*, Tachinidae.
 Antheraea: *Belvosia*, *Compsilura*, *Euexorista*, *Lespesia*, *Patelloa*,
 "Tachina," *Winthemia*, Tachinidae.

Attacus: *Lespesia*.

Automeris: *Belvosia*, *Carcelia*, *Compsilura*, *Euphorocera*, *Lespesia*, *Sisyropa*, Tachinidae.

Callosamia: *Compsilura*, *Euphorocera*, *Eusisyropa*, *Exorista*, *Lespesia*, "*Tachina*," *Winthemia*, Tachinidae.

Coloradia: *Leschenaultia*.

Copaxa: *Belvosia*.

Hemileuca: *Argyrophylax*, *Belvosia*, *Carcelia*, *Compsilura*, *Drino*, *Euphorocera*, *Exorista*, *Leschenaultia*, *Lespesia*, *Phryxe*, "*Sturmia*" probably, *Winthemia*, Tachinidae.

Hyalophora: *Belvosia*, *Lespesia*.

Hylesia: *Belvosia*.

Platysamia: *Compsilura*, *Euphorocera*, *Eusisyropa*, *Lespesia*, "*Tachina*," *Winthemia*, Tachinidae.

Rothschildia: *Belvosia*, *Lespesia*.

Samia: *Lespesia*, *Winthemia*.

Saturnia: *Euphorocera*.

Syssphinx: *Lespesia*.

Unidentified: *Compsilura*.

Family SCYTHRIDAE

Scythris: *Psalipteryx*.

Family SPHINGIDAE

Agrius: *Argyrophylax*, *Belvosia*, *Chaetogaedia*, *Drino*, *Metaplagia*, Tachinidae.

Ampeloeca: *Compsilura*.

Amphion: *Compsilura*, *Drino*.

Atreides: *Compsilura*.

Ceratomia: *Argyrophylax*, *Belvosia*, *Compsilura*, *Drino*, *Euphorocera*, *Eusisyropa*, *Hyphantrophaga*, *Lespesia*, *Winthemia*, Tachinidae.

Cressonia: *Winthemia*, Tachinidae.

Deidamia: *Compsilura*.

Hemaris: *Compsilura*, Gen. near *Thysanomyia*, *Winthemia*, Tachinidae.

Hyles: *Compsilura*, *Drino*, "*Tachina*," *Winthemia*, Tachinidae.

Manduca: *Atacta*, *Carcelia*, *Compsilura*, *Drino*, *Lespesia*, "*Masicera*," *Metavoria*, "*Sturmia*," *Winthemia*.

Pachysphinx: *Lespesia*, *Winthemia*, Tachinidae.

Paonias: *Compsilura*, *Leschenaultia*, *Lespesia*, *Winthemia*, Tachinidae.

Pholus: *Argyrophylax*, *Compsilura*, *Drino*, *Lespesia*, *Winthemia*,
Tachinidae.

Smerinthus: *Compsilura*, *Eusisyropa*, *Lespesia*, "Prosopaea,"
Tachinidae.

Sphecodina: *Compsilura*, *Eusisyropa*.

Sphinx: *Compsilura*, *Drino*, *Euphorocera*, *Leschenaultia*, *Lespesia*,
Metaplagia, *Winthemia*, Tachinidae.

Hawk moth(-s): *Belvosia*.

Sphinx moth(-s): *Drino*.

Unidentified: *Compsilura*, *Drino*, *Leschenaultia*, *Lespesia*,
Winthemia, "Zenillia."

Family STENOMIDAE

Antaeotricha: Tachinidae.

Stenoma: *Lixophaga*.

Family THYATIRIDAE

Pseudothyatira: Tachinidae.

Family TINEIDAE

Amydria: *Drepanoglossa*, *Philocalia*, *Trepophrys*.

Unidentified: *Schizactia*, "Tachinophyto."

Family TORTRICIDAE

Acleris: *Actia*, *Blondelia*, *Chaetophlepsis*, *Erynnia*, *Hemisturmia*,
Madremyia, *Nemorilla*, *Phryxe*, *Psalidopteryx*.

Adoxophyes: *Nemorilla*.

Amorbia: *Actia*, *Aplomya*, *Nemorilla*, *Pseudoperichaeta*.

Aphelia: *Nemorilla*, *Phryxe*.

Archippus: *Actia*, *Nemorilla*.

Archips: *Actia*, *Aplomya*, *Bessa*, *Blondelia*, *Carcelia*, *Ceromasia*,
Compsilura, *Dichaetoneura*, *Erynnia*, *Eulasiona*, *Eusisyropa*,
Exorista, *Hemisturmia*, Gen. near *Lydinolydella*, *Nemorilla*,
Nilea, *Paralispe*, *Phryxe*, *Pseudoperichaeta*, *Schizocerophaga*,
Tachinidae.

Argyrotaenia: *Aplomya*, *Elfia*, *Eusisyropa*, *Hemisturmia*, *Nemorilla*,
"Phorocera," *Pseudoperichaeta*.

Choristoneura: *Actia*, *Aplomya*, *Ceromasia*, *Chaetophlepsis*,
Compsilura, *Erynnia*, *Hemisturmia*, *Lypa*, *Madremyia*,
Nemorilla, *Omotoma*, "Phorocera," *Phryxe*, *Pseudoperichaeta*,

"*Sturmia*," *Tachinomyia*, *Winthemia*, *Xanthophyto*, Tachinidae.

Croesia: *Actia*, *Lypha*, *Schizotachina*.

Pandemis: *Aplomya*, *Ictericophyto*, *Nemorilla*.

Phaecasiophora: *Aplomya*.

Phalonia: *Actia*, *Elfia*, *Lixophaga*.

Platynota: *Erynnia*.

Sparganothis: *Blondelia*, *Erynnia*, *Lypha*, *Nemorilla*, Tachinidae.

Xenotemna: *Actia*, *Lixophaga*.

Leaf roller(-s): *Dejeaniopalpus*, *Periscepsia*, *Pseudoperichaeta*.

Tortricid (including Amorbia, Archips, Argyrotaenia, Choristoneura and/or Pandemis): *Aplomya*, *Pseudoperichaeta*.

Tortricid (including Archips and/or Choristoneura): *Aplomya*.

Unidentified: *Actia*, *Bessa*, *Blondelia*, *Erynnia*, *Eusisyropa*, *Hemisturmia*, *Nemorilla*.

Family YPONOMEUTIDAE

Mapsidius: *Lespesia*.

Urodus: *Euceromasia*, *Eusisyropa*, Tachinidae.

Yponomeuta: *Nemorilla*.

Family ZYGAENIDAE

Acoloithus: *Sturmia*.

Harrisina: *Euphorocera*, "*Phorocera*," *Sturmia*.

Unidentified: *Euphorocera*.

UNIDENTIFIED LEPIDOPTERA

Burrowing webworm(-s): *Paradidyma*.

Butterfly(-flies): *Acroglossa*.

Cabbage worm(-s): *Compsilura*, *Lespesia*.

Corn stalk borer(-s): *Archytas*.

Green looper(-s): *Euphorocera*, *Voria*.

Hog moth(-s): "*Phorocera*."

Leaf-folder caterpillar(-s): *Myobiopsis*.

Leaf tier(-s): *Psalidopteryx*.

Looper(-s): *Madremyia*.

Moth(-s): *Sisyropa*, "*Tachina*," *Tachinomyia*.

Moth stalkborer(-s): *Lixophaga*.

Oakworm(-s): *Eusisyropa*.

Tuber moth(-s): *Lydellohoughia*.

Unidentified: *Aplomya*, *Archytas*, *Belvosia*, *Carcelia*, *Ceromasia*, *Chaetogaedia*, *Compsilura*, *Deopalpus*, *Erynnia*, *Euexorista*,

Euphorocera, *Eusisyropa*, *Exorista* probably, *Leschenaultia*, *Lespesia*, *Lixophaga*, *Lydella*, *Madremyia*, *Mericia*, *Mochlosoma*, *Nemorilla*, *Paradejeania*, *Patelloa*, *Peleteria*, *Phryxe*, *Pseudomyothyria*, *Spathimeigenia*, "Tachina," *Winthemia*, Tachinidae.

Order ORTHOPTERA

Family ACRIDIDAE

Ageneotettix: Tachinidae.

Boopedon: *Hemithrixion*.

Camnula: *Ceracia*, *Hemithrixion*, Tachinidae.

Chimarocephala: *Ceracia*, Tachinidae.

Chortophaga: *Ceracia*.

Dissosteira: *Hemithrixion* probably, *Lespesia*, *Trichopoda*, Tachinidae.

Melanoplus: *Acemya*, *Ceracia*, *Euphorocera*, *Hemithrixion*, *Lespesia*, *Leucostoma*, Tachinidae.

Orphulella: Tachinidae.

Psoloessa: *Ceracia*.

Schistocerca: *Oedematocera*.

Spharagemon: Tachinidae.

Grasshopper(-s): *Acemya*, *Ceracia*, *Cylindromyia*, *Euphorocera*, *Hemithrixion*, *Leucostoma*, *Winthemia*, Tachinidae.

Locust(-s): *Lespesia*, Tachinidae.

Unidentified: *Cylindromyia*, *Trichopoda*, Tachinidae.

Family BLATTIDAE

Epilampra: *Calodexia*.

Eublaberus: *Calodexia*.

Periplaneta: *Miamimyia*.

Cockroach(-es): *Calodexia*.

Family GRYLLACRIDIDAE

Ceuthophilus: *Oedematocera*.

Stenopelmatus: *Stomatomyia*.

Family GRYLLIDAE

Anurogryllus: *Exoristoides*, *Theresiini*.

Eneoptera: *Calodexia*.

Gryllus: *Euphasiopteryx*, *Exoristoides*.

Ponca: *Calodexia*.

Cricket(-s): *Calodexia*, *Gilvella*.

Family MANTIDAE

Stagmomantis: Tachinidae.

Tenodera: *Trichopoda*.

Family PHASMATIDAE

Anisomorpha: *Phasmophaga*, *Roeseliopsis*.

Bacillus: "Tachina," Tachinidae.

Diapheromera: *Euhalidaya*, *Phasmophaga*, "Tachina," Tachinidae

Family TETTIGONIIDAE

Neoconocephalus: *Beskia*, *Euphasiopteryx*.

Unidentified: *Calodexia*.

UNIDENTIFIED ORTHOPTERA

Pictured sand cricket(-s): *Stomatomyia*.

Unidentified: *Euphasiopteryx*.

Class Chilopoda

Order GEOPHILOMORPHA

Family GEOPHILIDAE

Geophilus probably: *Exoristoides*.

Class Crustacea

Order ISOPODA

Family ONISCIDAE

Oniscus: *Melanophora*.

Porcellio: *Melanophora*.

Class Mammalia

Order RODENTIA

UNIDENTIFIED RODENTIA

Mouse (mice): *Leschenaultia*, *Parachaeta*.

HOSTS

The hosts are arranged alphabetically by generic and nonscientific names. Following these names, the order and family assignments are in parentheses. Specific names of hosts are listed alphabetically within each genus. Parasite names follow the host names. Synonyms, obsolete names, and incorrect spellings of host names are cross indexed to currently accepted names, with misspellings enclosed by quotation marks. For convenience, the common names of insects approved by the Entomological Society of America (Blickenstaff, 1970) for all host scientific names used in this catalog are cross indexed.

Acalymma (Col.: Chrysomelidae)

trivittata: *Celatoria diabroticae*.

vittata: *Celatoria diabroticae*, *C. setosa*, Tachinidae.

Acanthocephala (Hem. (Heter.): Coreidae)

declivis: *Trichopoda pennipes*.

femorata: *Trichopoda lanipes*, *T. pennipes*.

terminalis: *Trichopoda plumipes*, *T. spp.*

spp.: *Trichopoda pennipes*.

"Acanthocephalus" *femorata*, see *Acanthocephala femorata*

Acantholyda (Hym.: Pamphiliidae)

spp.: *Aplomya fronto*.

Achaea (Lep.: Noctuidae)

janata: *Chaetogaedia monticola*, *Eucelatoria armigera*,
"Winthemia diversoides."

Achatodes (Lep.: Noctuidae)

zeae: *Lydella radialis*, *L. radialis* probably, *Nemorilla pyste*.

Acigona (Lep.: Pyralidae)

dentilineatella: "Masicera tantilla."

Acleris (Lep.: Tortricidae)

minuta: *Blondelia eufitchiae*, *Hemisturmia tortricis*, *Nemorilla pyste*.

variana: *Actia diffidens*, *A. interrupta*, *Chaetophlepsis semiothisae*, *Erynnia tortricis*, *E. spp.*, *Hemisturmia tortricis*, *Madremyia saundersii*, *Nemorilla pyste*, *Phryxe pecosensis*, *Psaliapteryx pseudonuda*.

variegana: *Actia interrupta*.

Acoloithus (Lep.: Zygaenidae)

sp.: *Sturmia harrisinae* probably.

Acorn Carpocapsa [sp.], see *Melissopus latiferreanus*

Acrididae (Orth.: Acrididae)

Unidentified: *Cylindromyia euchenor*, *Trichopoda plumipes*,
Tachinidae.

Acridiodes G[enus], see Acrididae, unidentified

Acrobasis (Lep.: Pyralidae)

betulella: *Nemorilla pyste*, Tachinidae.

caryae: *Didyma exigua*, *Eusisyropa boarmiae*, *Hyphantrophaga hyphantriae*, "Myiophasia atra," *Nemorilla pyste*, "Zenillia" spp., Tachinidae.

caryvorella: *Nemorilla pyste*.

comptoniella: *Elfia palpigera*, *Eusisyropa blanda*, *Nemorilla pyste*, *Schizotachina convecta*.

coryliella: *Nemorilla pyste*.

indigenella: *Eusisyropa boarmiae*, *Exorista* spp., *Hemisturmia tortricis*, *Leskiomima tenera*, *Nemorilla pyste*, *N.* sp. near *pyste*, "Tachina" spp.

indigenella probably: *Eusisyropa boarmiae*.

juglandis, see *A. indigenella* probably or *A. juglandis*.

juglandis: *Leskiomima tenera*, *Nemorilla pyste*.

nebullella, see *A. indigenella*.

"nubilella," see *A. indigenella* probably.

rubrifasciella: *Nemorilla pyste* probably.

sp.: *Eusisyropa boarmiae* probably.

Acronicta (Lep.: Noctuidae)

sp. near *afflicta*: "Exorista" *lobeliae*.

americana: *Compsilura concinnata*, *Exorista mella*, *E.* spp.

americana probably: *Mystacella* spp.

betulae: "Exorista" sp. near *lobeliae*, *Tachinomyia* sp. near *panaetius*, *T.* spp., "Zenillia" spp.

brumosa: *Compsilura concinnata*.

dactylina: *Compsilura concinnata*, *Eusisyropa virilis*.

fragilis: *Compsilura concinnata*.

furcifera: *Compsilura concinnata*, "Zenillia" spp., Tachinidae.

grisea: *Compsilura concinnata*, *Eusisyropa virilis*, "Exorista" *lobeliae*.

hamamelis: *Carcelia formosa*, *Euphorocera claripennis*, "Exorista" *lobeliae*, *Sisyropa eudryae*.

hasta: *Compsilura concinnata*.

hastulifera: *Sisyropa* spp.

impleta: *Compsilura concinnata*, *Eumasicera* sp. near *sternalis*, *Sisyropa eudryae* probably, *Winthemia datanae* probably.

impressa: *Compsilura concinnata*, *Eusisyropa virilis*, *Nilea sternalis*, Tachinidae.

- interrupta*: *Archytas aterrimus*, *Nilea* sp. near *noctuiiformis*.
lanceolaria: *Compsilura concinnata*, *Lespesia* spp.
leporina: *Tachinomyia nigricans*.
leporina vulpina: *Eusisyropa virilis*.
lepusculina: *Exorista mella*, *E.* spp., Tachinidae.
lobeliae: "*Exorista*" *lobeliae*.
longa: "*Zenillia*" spp.
"oblinata," see *A. oblinita*.
"obliniata," see *A. oblinita*.
oblinita: *Compsilura concinnata*, *Eusisyropa virilis*, "*Nemorilla trivittata*" probably.
occidentalis, see *A. interrupta*.
ovata: *Archytas aterrimus*.
rubricoma: *Exorista mella*.
superans: *Compsilura concinnata*, Tachinidae.
spp. (including *A. leporina* and/or *A. lepusculina*): *Eusisyropa virilis*.
spp.: *Compsilura concinnata*, *Eumasicera* sp. near *sternalis*, "*Exorista*" *lobeliae*, *E. mella*, *Sisyropa eudryae*, *S. eudryae* probably, "*Zenillia*" spp.

Acronycta, see *Acronicta* species of same name except as noted

Acronycta chionochroa and/or *A. leporina*, see *Acronicta* spp.
(including *A. leporina* and/or *A. lepusculina*)

Acronycta luteicoma, see *Acronicta impleta*

Acronycta populi, see *Acronicta lepusculina*

Acrosternum (Hem. (Heter.): Pentatomidae)

hilare: *Trichopoda pennipes*.

pennsylvanicum: *Trichopoda pennipes*.

Actebia (Lep.: Noctuidae)

fennica: *Chaetogaedia monticola*, *Euphorocera claripennis*,
Periscepsia helymus, Tachinidae.

Actias (Lep.: Saturniidae)

luna: *Cylindromyia binotata*, *Lespesia frenchii*, *Winthemia cecropia*, Tachinidae.

Adisophanes miscellus, see *Catabena lineolata*

Adoneta (Lep.: Limacodidae)

spinuloides: *Carcelia* spp.

Adoretus (Col.: Scarabaeidae)

spp.: *Prosenia* spp.

Adoxophyes (Lep.: Tortricidae)

furcatana: *Nemorilla pyste*.

Aegeriidae (Lep.: Aegeriidae)

Unidentified: *Eulasiona genalis*.

Agapema galbina, see *Saturnia galbina*

Agathodes (Lep.: Pyralidae)

monstralis: *Eucelatoria australis*.

Agathymus (Lep.: Megathymidae)

aryxna, see *A. neumogeni*.

baueri: "Phorocera" *texana*.

chisosensis: "Phorocera" *texana*.

evansi: "Phorocera" *texana*.

freemani: "Phorocera" *texana*.

neumogeni: "Phorocera" *texana*.

Ageneotettix (Orth.: Acrididae)

deorum: Tachinidae.

Aglais antiopa, see *Nymphalis antiopa***Aglais milberti**, see *Nymphalis milberti***Agraulis** (Lep.: Nymphalidae)

vanillae: *Euphorocera claripennis*.

vanillae incarnata: *Compsilura concinnata*, *Euphorocera claripennis*.

Agrioconota (Col.: Chrysomelidae)

bivittata: *Eucelatoriopsis dimmocki*.

Agrius (Lep.: Sphingidae)

cingulatus: *Agryrophylax* spp., *Belvosia bifasciata*, *Chaetogaedia monticola*, *Drino inca*, *D. incompta*, *D. rhoeo*.

cingulatus probably: *Drino inca*.

convolvuli: *Drino rhoeo*, *Metaplagia occidentalis*, Tachinidae.

Agrotid sp., see *Agrotis* spp.**Agrotidian(-s)** (Lep.: Noctuidae)

Unidentified: *Winthemia leucanae* lacking red at tip.

Agrotis (Lep.: Noctuidae)

cinctipennis, see *Peridroma cinctipennis cinctipennis*.

c-nigrum, see *Amathes c-nigrum*.

cochrani, see *Euxoa messoria* or *E. messoria* probably.

coniotis, see *Peridroma coniotis* or *P. coniotis coniotis*.

criniger: *Lespesia archippivora*.

dislocata: *Chaetogaedia monticola*, *Lespesia archippivora*, Tachinidae.

fennica, see *Actebia fennica*.

gladiaria: *Bonnetia comta*.

inermis, see *Peridroma saucia*.

ipilon: *Archytas cirphis*, *Bonnetia comta*, *Carcelia formosa*, *Chaetogaedia monticola*, *Eucelatoria armigera*, *Euphorocera claripennis*, *Gonia longipulvilli*, *G. sequax*, *G. spp.*, *Lespesia archippivora*, *Madremyia saundersii*, *Sisyropa eudryae*, *S. spp.*, *Tachinomyia panaetius*.

ipsilon probably: *Archytas marmoratus*.

malefida: *Bonnetia comta*, *Gonia longipulvilli*.

orthogonia: *Bonnetia comta*, *Gonia aldrichi*, *G.* sp. near *aldrichi*, *G. longiforceps*, *G. longipulvilli*, *G.* spp., *Mericia* spp., *Peleteria texensis*, *P.* spp., *Periscepsia rohweri*.

smithii, see *Amathes smithii*.

sp. near *subgothica*, see *Feltia* sp. near *subgothica*.

telifera, see *A. ipsilon*.

unicolor, see *Spaelotis clandestina*.

venerabilis: *Bonnetia comta*.

vetusta: "Linnaemya picta."

ypsilon, see *A. ipsilon*.

spp.: *Chaetogaedia crebra*, *Gonia* spp., *Periscepsia helymus*.

sp. probably: *Periscepsia* spp.

Alabama (Lep.: Noctuidae)

argillacea: *Beskia aelops*, *Euphorocera claripennis*, *E. floridensis*, *Eusisyropa blanda*, *E. boarmiae*, "Exorista" *lobeliae*, *Lespesia aletiae*, *Patelloa leucaniae*, "Tachina" spp., *Triachora unifasciata*, Tachinidae.

"argillicae," see *A. argillacea*.

"argillicea," see *A. argillacea*.

Alatuncusia (Lep.: Pyralidae)

bergi: *Eusisyropa blanda*.

Alberada (Lep.: Pyralidae)

parabates: *Clausicella floridensis*.

Alcaeorrhynchus (Hem. (Heter.): Pentatomidae)

grandis: *Trichopoda pennipes*.

Aletia (Lep.: Noctuidae)

argillacea, see *Alabama argillacea*.

xylina, see *Alabama argillacea*.

spp.: "Prosopaea" sp., Tachinidae.

Alfalfa caterpillar, see *Colias eurytheme*

Alfalfa looper, see *Autographa californica*

Alfalfa webworm, see *Loxostege commixtalis*

Alfalfa weevil, see *Hypera postica*

Allantus (Hym.: Tenthredinidae)

cinctus: *Guerinia dydas*.

Alsophila (Lep.: Geometridae)

pometaria: *Blondelia eufitchiae*, *Cyzenis albicans*, *Gonia sagax*, *Ictericophyto spinosa*, *Omotoma fumiferanae*, *Phryxe pecosensis*, *P. vulgaris*, *Pseudotachinomyia slossonae*, *Tachinomyia nigricans*, *T.* spp., *Winthemia rufopicta*, Tachinidae.

Altica (Col.: Chrysomelidae)

bimarginata: *Hyalomyodes triangulifer*.

chalybea: *Medina barbata*.

corni: *Medina barbata*.

probata: Tachinidae.

ulmi: *Medina barbata*.

spp.: *Medina barbata*.

Alypia (Lep.: Noctuidae)

octomaculata: *Aplomya pheosiae*, *Carcelia* spp., *Chaetophlepsis eudryae*, *Compsilura concinnata*, *Drino inconspicua*, *Lespesia archippivora*, *Sisyropa alypiae*, *S. alypiae* probably, *Winthemia quadripustulata*, *W. rufopicta*, Tachinidae.

spp.: *Sisyropa eudryae*.

Amathes (Lep.: Noctuidae)

c-nigrum: *Euphorocera claripennis*, *Winthemia quadripustulata*, *W. rufopicta* probably.

smithii: *Linnaemya tessellata*, *Periscepsia helymus*.

Amauronematus (Hym.: Tenthredinidae)

spp.: *Bessa harveyi*.

American cockroach, see *Periplaneta americana*

American dagger moth, see *Acronicta americana*

American tent caterpillar(-s) (Lep.: Lasiocampidae)

American tent caterpillar(-s): *Carcelia malacosomae* probably, *Compsilura concinnata*, *Lespesia frenchii*, Tachinidae.

Ametastegia (Hym.: Tenthredinidae)

pallipes: *Bessa harveyi*.

Amorbia (Lep.: Tortricidae)

cuneana: *Actia interrupta*.

emigratella: *Nemorilla pyste*.

essigana: *Pseudoperichaeta erecta*, *P. erecta* probably.

humerosana: *Aplomya caesar*.

humerosana, *Archips argyrospila*, *A. fractivittana*, *A. rosaceana*, *A. semiferana*, *Eulia quadrifasciana* and/or *Pandemis limitata*, see Tortricid spp. (including *Amorbia humerosana*, *Archips argyrospilus*, *A. semiferanus*, *Argyrotaenia quadrifasciana*, *Choristoneura fractivittana*, *C. rosaceana* and/or *P. limitata*).

Ampeloeca (Lep.: Sphingidae)

myron: *Compsilura concinnata*.

Ampelogypter (Col: Curculionidae)

sesostris: *Myiophasia "aenea."*

"*sesostris*," see *A. sesostris*.

Ampelophaga myron, see *Ampeloeca myron*

Amphicerus (Col.: Bostrichidae)

bicaudatus: *Lixophaga variabilis*.

Amphidasis cognataria, see *Biston cognataria*

"Amphidosia" *cognataria*, see *Biston cognataria*

Amphion (Lep.: Sphingidae)

nessus: *Compsilura concinnata*, *Drino* sp. near *bakeri*.

Amphipyra (Lep.: Noctuidae)

pyramidoides: *Compsilura concinnata*, *Mericia* spp., *Omotoma fumiferanae*, *Winthemia rufopicta* probably, Tachinidae.

Amydria (Lep.: Tineidae)

"*effrenatella*," see *A. effrenatella*.

effrenatella: *Philocalia tenuirostris*.

spp.: *Drepanoglossa amydriae*, *Trepophrys cinerea*.

Amyna (Lep.: Noctuidae)

natalis: *Eucelatoria armigera*.

Anacampsis (Lep.: Gelechiidae)

innocuella: *Lypha intermedia*.

Anacamptodes (Lep.: Geometridae)

clivinaria profanata: *Blondelia* spp.

ephyraria: *Eusisyropa virilis*.

fragilaria: *Lespesia archippivora*, "*Winthemia diversoides*."

[sp.], see *A. fragilaria*.

"Anacaptodes" *fragilaria*, see *Anacamptodes fragilaria*

Anaea (Lep.: Nymphalidae)

andria: *Euphorocera edwardsii*.

glycerium: *Lespesia archippivora*.

Anarsia (Lep.: Gelechiidae)

lineatella: *Erynnia tortricis*, *Euphorocera claripennis*, *Psaliapteryx pollinosa*.

Anasa (Hem. (Heter.): Coreidae)

armigera: *Trichopoda pennipes*, *T. pennipes* probably.

tristis: *Trichopoda pennipes*, Tachinidae.

spp.: *Trichopoda pennipes*, *T.* spp.

Ancylis (Lep.: Olethreutidae)

comptana: *Erynnia tortricis*, *Hemisturmia tortricis*, *Lixophaga variabilis*, *Nemorilla pyste*.

comptana floridana: *Erynnia tortricis*.

comptana fragariae: *Erynnia tortricis*, *Nemorilla pyste*.

floridana, see *A. comptana floridana*.

Anisolabis (Dermaptera: Carcinophoridae)

maritima probably: Tachinidae.

Anisomorpha (Orth.: Phasmatidae)

buprestoides: *Phasmophaga antennalis* probably, *P. meridionalis*, *Roeseliopsis americana*.

Anisopteryx vernata, see *Paleacrita vernata*

Anisota (Lep.: Citheroniidae)

rubicunda: *Belvosia bifasciata*, *B. sp.*, *Compsilura concinnata*, *Eumasicera sternalis*, *Euphorocera claripennis*, *Eusisyropa blanda*, *Lespesia datanarum*, *L. frenchii*, *L. spp.*, "Masicera" spp., *Pacidianus persimilis*, "Sturmia" spp., *Winthemia datanae*.

sp. near rubicunda: *Belvosia bifasciata*, *Euphorocera floridensis*.

senatoria: *Belvosia bifasciata*, *Compsilura concinnata*, *Eumasicera sternalis*, *Euphorocera claripennis*, *Hyphantrophaga hyphantriae*, *Lespesia datanarum*, *L. frenchii*, *L. spp.*, *Oswaldia aurifrons*, "Sturmia" spp., *Winthemia datanae*, *W. sp. near datanae*, *W. quadripustulata*, Tachinidae.

virginiensis: *Compsilura concinnata*, *Lespesia datanarum*, *L. frenchii*, *Pacidianus sp.*, Tachinidae.

spp.: *Compsilura concinnata*, *Lespesia datanarum*.

Anomala (Col.: Scarabaeidae)

spp.: *Prosenia spp.*

Anomis (Lep.: Noctuidae)

erosa: *Eusisyropa blanda* probably, *Lixophaga variabilis*.

hawaiiensis: *Chaetogaedia monticola*, *Lespesia archippivora*.

noctivolans: *Chaetogaedia monticola*, *Lespesia archippivora*.

Anoplonyx (Hym.: Tenthredinidae)

canadensis: *Bessa harveyi*.

luteipes: *Bessa harveyi*.

spp.: *Bessa harveyi*.

Anosia plexippus, see *Danaus plexippus plexippus*

Antaeotricha (Lep.: Stenomidae)

leucillana: Tachinidae.

"**Antheracea**" *polyphemus*, see *Antheraea polyphemus*

Antheraea (Lep.: Saturniidae)

polyphemus: *Belvosia bifasciata*, *Compsilura concinnata*, *Euexorista futilis*, *Lespesia datanarum*, *L. frenchii*, *L. sabroskyi*, *L. spp.*, *Patelloa fuscimacula*, "Tachina" spp., *Winthemia cecropia*, *W. quadripustulata*, Tachinidae.

Anthonomus (Col.: Curculionidae)

grandis: *Myiophasia "aenea," M. globosa*.

Anthophila (Lep.: Glyphipterygidae)

pariana: *Hemisturmia tortricis*, *Nemorilla pyste*.

Anticarsia (Lep.: Noctuidae)

gemmatalis: *Euphorocera floridensis*, *Gonia spp.*, Gen. and sp.

of Sturmiini, *Voria ruralis*, *Winthemia rufopicta*.

“*gemmatilis*,” see *A. gemmatalis*.

Anurogryllus (Orth.: Gryllidae)

muticus: *Exoristoides* sp., Gen. and sp. of Theresiini.

Apamea (Lep.: Noctuidae)

apamiformis: *Euexorista futilis*.

lignicolor: *Nowickia algens*.

“*Apantesia*” *phyllira*, see *Apantesis phyllira*

Apantesis (Lep.: Arctiidae)

blakei superba, see *A. nevadensis superba*.

nevadensis superba: *Exorista mella*.

oithona: *Euexorista futilis*, *Euphorocera claripennis*, *E. floridensis*, *Exorista mella*.

ornata: *Carcelia malacosomae* probably.

ornata var. *achaia*, see *A. ornata*.

phalerata: *Euexorista futilis*.

phyllira: *Exorista mella*, *Gymnocarcelia ricinorum*.

proxima: *Carcelia laxifrons*, *C. malacosoma* probably, *Euphorocera claripennis*.

rectilinea, see *A. oithona*.

Apatela, see *Acrionicta* species of same name

Apatura clyton, see *Asterocampa clyton*

Apatura sp., see *Asterocampa* sp.

Aphelia (Lep.: Tortricidae)

alleniana: *Nemorilla pyste*, *Phryxe pecosensis*.

Apple-and-thorn skeletonizer, see *Anthophila pariana*

Apple tree tent caterpillar(-s) (Lep.: Lasiocampidae)

Apple tree tent caterpillar(-s): *Euphorocera claripennis*, *E. tachinomoides*, *Exorista mella*.

Apple twig borer, see *Amphicerus bicaudatus*

“*Arachnara*” *oblonga*, see *Archanara oblonga*

Arachnis (Lep.: Arctiidae)

picta: *Carcelia malacosomae* probably, *Leschenaultia adusta*, *Sisyropa* spp.

Archanara (Lep.: Noctuidae)

oblonga: *Ceromasia* spp., *Lydella radialis*.

subcarnea, see *A. oblonga*.

Archimerus (Hem. (Heter.): Coreidae)

alternatus: *Trichopoda lanipes*, *T. pennipes*, *T. pennipes* complex.

calcarator: *Trichopoda pennipes*.

Archippus (Lep.: Tortricidae)

packardianus: *Actia interrupta*, *Nemorilla pyste*.

Archips (Lep.: Tortricidae)

argyrospilus: *Actia interrupta*, *Aplomya caesar*, *A. caesar* probably, *Carcelia malacosomae* probably, *Compsilura concinnata*, *Eusisyropa blanda*, *Hemisturmia tortricis*, *Nemorilla pyste*, *Paralipse infernalis*, *Phryxe pecosensis*, *P. vulgaris*, *Pseudoperichaeta erecta*, Tachinidae.

cerasivorana rileyana, see *A. rileyanus*.

cerasivoranus: *Actia* sp. near *interrupta*, *Ceromasia auricaudata*, *Compsilura concinnata*, *Dichaetoneura leucoptera*, *Erynnia tortricis*, *Eusisyropa blanda*, *Hemisturmia tortricis*, *Nemorilla pyste*, *Nilea sternalis*, *Phryxe pecosensis*, *Pseudoperichaeta erecta*, *Schizocerophaga leiby*.

conflictana, see *Choristoneura conflictana*.

"*fervadana*," see *A. fervidanus*.

"*fervadona*," see *A. fervidanus*.

fervidanus: *Actia* sp. near *interrupta*, *Dichaetoneura leucoptera*, *Erynnia tortricis*, *Eusisyropa blanda*, *E. boarmiae*, *Exorista* spp., *Nemorilla pyste*.

fumiferana, see *Choristoneura fumiferana*.

griseus: *Nemorilla pyste*.

infumatanus: *Blondelia* sp. near *polita*, *Phryxe pecosensis*.

parallela, see *Choristoneura parallela*.

"*practivittana*," see *Choristoneura fractivittana*.

purpuranus: *Aplomya caesar*, *Nemorilla pyste*, *Pseudoperichaeta erecta*.

rileyanus: *Hemisturmia tortricis*.

rosaceana, see *Choristoneura rosaceana*.

rosanus: *Actia interrupta*, *Aplomya caesar*, *Erynnia tortricis*, *Eulasiona nigra*, Gen. and sp. near *Lydinolydella*, *Nemorilla pyste*, *Pseudoperichaeta erecta*.

semiferanus: *Aplomya caesar*.

spp. (including *A. argyrospilus* and/or *A. griseus*): *Actia interrupta*, Tachinidae.

spp.: *Aplomya caesar*, *Bessa harveyi*, *Blondelia* spp.

spp. probably: Tachinidae.

Arctia docta, see *Apantesis proxima*

Arctia phyllira, see *Apantesis phyllira*

Arctiidae (Lep.: Arctiidae)

Unidentified: *Bombyliopsis abrupta*, *Carcelia laxifrons*, *C. reclinata*, *C.* spp., *Chaetogaedia analis*, *Compsilura concinnata*, "*Gonia sagax*," *Gymnocarcelia ricinorum*, *Hubneria estigmenensis*, *Leschenaultia adusta*,

L. grossa, *Lespesia aletiae*, *Sisyropa eudryae*, *Thelaira* spp., *Winthemia datanae*.

Unidentified probably: "*Phorocera victoria*."

Arge (Hym.: Argidae)

clavicornis abdomen yellow, see *A. clavicornis* complex.

clavicornis complex: *Aporotachina chaetoneura*, *Bessa harveyi*, *Lydella* sp. probably.

humeralis: *Spathimeigenia hylotomae*.

pectoralis: *Aporotachina dexina*, *Bessa harveyi*, *Spathimeigenia spinigera*.

rubiginosa: *Bessa harveyi*, *Spathimeigenia hylotomae*.

spp.: *Bessa harveyi*, *Spathimeigenia hylotomae*, "*Sturmia*" sp.

Argus tortoise beetle, see *Chelymorpha cassidea*

Argynnis cybele, see *Speyeria cybele*

Argyrotaenia (Lep.: Tortricidae)

citrana: *Aplomya caesar*, *Elfia* sp. near *palpiger*, "*Phorocera parva*," *Pseudoperichaeta erecta*.

mariana: *Eusisyropa blanda*, *Pseudoperichaeta erecta*.

quadrifasciana: *Aplomya caesar*.

velutinana: *Hemisturmia tortricis*, *Nemorilla pyste*.

Aristotelia (Lep.: Gelechiidae)

roseosuffusella: *Schizactia vitinervis*.

Army cutworm, see *Euxoa auxiliaris*

Armyworm, see *Pseudaletia unipuncta*

Armyworm(-s) (Lep.: Noctuidae)

Armyworm(-s) (including *Pseudaletia unipuncta* and/or *Spodoptera mauritia*): *Archytas cirphis*, *Chaetogaedia monticola*, *Lespesia archippivora*.

Armyworm(-s): *Archytas cirphis*, *A. marmoratus*, *A.* spp. (including *A. apicifer* and/or *A. californiae*), *A.* spp., *Bonnetia comta*, *Chaetogaedia monticola*, *Eucelatoria armigera*, *E. rubentis*, *Euphorocera claripennis*, *Lespesia archippivora*, *Patelloa leucaniae*, *Peleteria texensis*, *Periscepsia helymus*, "*Phorocera*" spp., *Triachora ochriventris*, *T. unifasciata*, *Winthemia infesta*, *W. leucanae*, *W. militaris*, *W. quadripustulata*, *W. rufopicta*, Tachinidae.

Arsenura erythrinae, see *Samia cynthia*

Arsilonche albovenosa, see *Simyra henrici*

Artichoke plume moth, see *Platyptilia carduidactyla*

Arzama (Lep.: Noctuidae)

densa: *Lydella radialis*.

obliqua: *Erynnia tortricis*, *Lydella radialis*, *Oswaldia* sp. probably.

"*obliquata*," see *A. obliqua*.

Ascia (Lep.: Pieridae)

monuste: *Prophryno parviteres*.

rapae, see *Pieris rapae*.

Asciodes (Lep.: Pyralidae)

anormalis probably: *Nemorilla pyste*.

gordialis: *Nemorilla* sp. near *pyste*.

Asiatic beetle, see *Maladera castanea* or *M. castanea* probably

Asiatic garden beetle, see *Maladera castanea*

Asparagus beetle, see *Crioceris asparagi*

Aspen leaf beetle, see *Chrysomela crotchii*

Asterocampa (Lep.: Nymphalidae)

celtis: *Euphorocera edwardsii*.

clyton: *Euphorocera claripennis*.

sp.: *Lespesia aletiae*.

Atreides (Lep.: Sphingidae)

plebeius: *Compsilura concinnata*.

Atreus "*plebeja*," see *Atreides plebeius*

"**Attacas**" *orizaba*, see *Rothschildia orizaba*

Attacus (Lep.: Saturniidae)

cecropia, see *Platysamia cecropia*.

"*cecropiae*," see *Platysamia cecropia*.

luna, see *Actias luna*.

polyphemus, see *Antheraea polyphemus*.

promethea, see *Platysamia promethea*.

sp.: *Lespesia frenchii*.

Autochton (Lep.: Hesperiiidae)

cellus: *Lespesia aletiae*.

Autographa (Lep.: Noctuidae)

brassicae, see *Trichoplusia ni*.

"*california*," see *A. californica*.

californica: *Aplomya trisetosa*, *Chaetogaedia monticola*, *Eucelatoria armigera*, *Euexorista futilis*, *Lespesia archippivora*, *L. spp.*, *Madremyia saundersii*, *Periscepsia helymus*, *Pilatea delecta*, *Siphona plusiae*, *Voria ruralis*.

californica probably: *Voria aurifrons*.

egena, see *Autoplusia egena*.

gamma: *Siphona spp.*, *Voria ruralis* probably.

gamma californica, see *A. californica*.

oo, see *Pseudoplusia includens*.

rogationis, see *Pseudoplusia includens*.

spp.: *Chaetophlepsis eudryae*, *Voria ruralis*.

Automeris (Lep.: Saturniidae)

io: *Carcelia formosa*, *Compsilura concinnata*, *Euphorocera*

claripennis, *Lespesia dimmocki*, *L. frenchii*, *L. sabroskyi*, *Sisyropa eudryae*, Tachinidae.

leucane, see *A. nyctimene leucane*.

nyctimene leucane: *Belvosia ciliata*.

spp.: *Sisyropa eudryae*.

Autoplusia (Lep.: Noctuidae)

egena: *Chaetogaedia monticola*, *Voria ruralis*.

Bacillus (Orth.: Phasmatidae)

rossia: "*Tachina*" spp., Tachinidae.

rossii, see *B. rossia*.

Bagworm, see *Thyridopteryx ephemeraeformis*

Baileya (Lep.: Noctuidae)

dormitans: *Tachinomyia nigricans*.

Balaninus nasicus, see *Curculio nasicus*

Balaninus "pardalus" or *Curculio naso*, see Curculionid spp.
(including *Conotrachelus naso* or *Curculio pardalis*)

Balaninus "pardulus" or *Curculio naso*, see Curculionid spp.
(including *Conotrachelus naso* or *Curculio pardalis*)

Balaninus uniformis, see *Curculio uniformis*

Balaninus [sp.], see *Curculio* spp.

"**Balanus**" spp., see *Curculio* spp.

Balsam fir sawfly, see *Neodiprion abietis*

Banded woollybear, see *Isia isabella*

Barathra (Lep.: Noctuidae)

configurata, see *Mamestra configurata*.

curialis: *Mericia bicarina* probably.

Barbara (Lep.: Olethreutidae)

colfaxiana: *Eribella* sp. near *exilis*.

Barrenadores, see Lepidoptera, miscellaneous

Basilarchia archippus, see *Limenitis archippus*

Basilarchia arthemis astyanax, see *Limenitis astyanax*

Basilarchia astyanax, see *Limenitis astyanax*

Basilona imperialis, see *Eacles imperialis*

Basswood leafroller, see *Pantographa limata*

Battus (Lep.: Papilionidae)

philenor: *Compsilura concinnata*.

Bean leaf beetle, see *Cerotoma trifurcata*

Bean leafroller, see *Urbanus proteus*

Bean leafskeletonizer, see *Autoplusia egena*

Bee(-s) (Hym.: Unidentified Hymenoptera)

Bee(-s): "*Tachina*" spp.

Beet armyworm, see *Spodoptera exigua*

Beet webworm, see *Loxostege sticticalis*

Beet webworm and/or clover cutworm, see Noctuid and/or pyralid

- spp. (including *Scotogramma trifolii* and/or *Loxostege sticticalis*)
- Bella moth, see *Utetheisa ornatrix bella*
- Bellura obliqua, see *Arzama obliqua*
- "Bembex" *nubilipennis*, see *Bembix nubilipennis*
- Bembix (Hym.: Sphecidae)
- nubilipennis*: "*Prospherysa*" sp.
- Bertha armyworm, see *Mamestra configurata*
- Bidens borer, see *Epiblema otiosanum*
- Birch casebearer, see *Coleophora fuscedinella*
- Birch sawfly, see *Arge pectoralis*
- Birch tubemaker, see *Acrobasis betulella*
- Biston (Lep.: Geometridae)
- cognataria*: *Aplomya helvina*, *Compsilura concinnata*, *Exorista* spp.
- Black army cutworm, see *Actebia fennica*
- Black army worm, see *Actebia fennica*
- Black cutworm, see *Agrotis ipsilon*
- Blackheaded ash sawfly, see *Tethida cordigera*
- Blackheaded fireworm, see *Rhopobota naevana*
- Black swallowtail, see *Papilio polyxenes asterius*
- Blastobasis *nubilella*, see *Valentinia glandulella*
- Blepharida (Col.: Chrysomelidae)
- rhois*: *Adoryphorophaga aberrans*.
- "Blepharomastix" (Lep.: Pyralidae)
- ebulialis*: *Myobiopsis flavipennis*.
- Blue cactus borer, see *Melitara dentata*
- Bluegrass billbug, see *Sphenophorus parvulus*
- Boarmia *pampinaria*, see *Isturgia truncataria*
- Boarmia "*plumigeraria*," see *Coniodes plumogeraria*
- Boll weevil, see *Anthonomus grandis*
- Bollworm, see *Heliothis zea*
- Bombus (Hym.: Apidae)
- fervidus*: *Clausicella neomexicana*.
- spp.: *Lespesia frenchii*, "*Tachina*" spp.
- Bombycidae (Lep.: Bombycidae)
- Unidentified: *Exorista mella*, *Lespesia frenchii*, *Pseudochaeta argentifrons*, *Winthemia quadripustulata*.
- Bombyx (Lep.: Bombycidae)
- mori*: *Compsilura concinnata*, *Zenillia libatrix*.
- sp.: *Lespesia ciliata*.
- Bomolocha (Lep.: Noctuidae)
- abalienalis*: *Blondelia hyphantriae*.
- deceptalis*: *Blondelia hyphantriae*.

- Boopedon** (Orth.: Acrididae)
nubilum: *Hemithrixion oestriforme*.
- “**Botis**” *penitalis*, see *Ostrinia* spp. (including *O. oblitalis* and/or *O. penitalis*)
- Botys** *penitalis*, see *Ostrinia penitalis* or *Ostrinia* spp. (including *O. oblitalis* and/or *O. penitalis*)
- Botys** sp., see *Pyrausta* spp. probably
- Boxelder leaf roller**, see *Gracillaria negundella*
- Boxelder twig borer**, see *Proteoteras willingana*
- Brassolis** (Lep.: Brassolidae)
isthmia: Tachinidae.
- Brephidium** (Lep.: Lycaenidae)
exilis: *Aplomya theclarum*, Tachinidae.
- Brephos** (Lep.: Geometridae)
infans: *Actia flavipes*, *Ictericophyto spinosa*.
- Bristly cutworm**, see *Lacinipolia renigera*
- Bristly roseslug**, see *Cladius isomerus*
- Brownheaded ash sawfly**, see *Tomostethus multicinctus*
- Brown stink bug**, see *Euschistus servus*
- Browntail moth**, see *Nygmia phaeorrhoea*
- Bruce spanworm**, see *Operophtera bruceata*
- “**Bryshidium exile**,” see *Brephidium exilis*
- Buck moth**, see *Hemileuca maia*
- Buprestis** (Col.: Buprestidae)
aurulenta: Tachinidae.
- Burdock borer**, see *Papaipema cataphracta*
- Burrowing webworm(-s)** (Lep.: Unidentified Lepidoptera)
Burrowing webworm(-s): *Paradidyma singularis*.
- Butterfly(-flies)** (Lep.: Unidentified Lepidoptera)
Butterfly(-flies): *Acroglossa hesperidarum*.
- Byrdia** (Lep.: Lymantriidae)
groenlandica: *Exorista fasciata fasciata*, *Peleteria aenea*, *Petin-arctia stylata*.
- Cabbage cutworm(-s)** (Lep.: Noctuidae)
Cabbage cutworm(-s): *Gonia* spp.
- Cabbage looper**, see *Trichoplusia ni*
- Cabbage webworm**, see *Hellula rogatalis*
- Cabbage worm(-s)** (Lep.: Unidentified Lepidoptera)
Cabbage worm(-s): *Compsilura concinnata*, *Lespesia archippivora*.
- Cabera** “*erythremaria*,” see *Deilinia erythemaria*
- Cacoecia argyrospilus**, see *Archips argyrospilus*
- Cacoecia cerasivoranus**, see *Archips cerasivoranus*
- Cacoecia fervidanus**, see *Archips fervidanus*

- Cacoecia fumiferana*, see *Choristoneura fumiferana*
Cacoecia parallela, see *Choristoneura parallela*
Cacoecia rosaceana, see *Choristoneura rosaceana*
Cacoecia rosanus, see *Archips rosanus*
Cacoecia spp., see *Archips* spp.
 "Caecacia" *cerasivoranus*, see *Archips cerasivoranus*
 "Caenurgia" *erechtea*, see *Caenurgina erechtea*
Caenurgina (Lep.: Noctuidae)
 erechtea: *Winthemia quadripustulata*.
Calendra incurrens, see *Sphenophorus incurrens*
 California oakworm, see *Phryganidia californica*
Callarctia ornata, see *Apantesis ornata*
Callarctia proxima, see *Apantesis proxima*
Calligrapha (Col.: Chrysomelidae)
 bigbyana, see *C. multipunctata bigbyana*.
 multipunctata: *Euphorocera claripennis*.
 multipunctata bigbyana: *Myiopharus canadensis*, *M. dorsalis*.
Callopietria (Lep.: Noctuidae)
 floridensis: Tachinidae.
Callosamia (Lep.: Saturniidae)
 "colleta," see *Hyalophora calleta*.
 promethea: *Compsilura concinnata*, *Euphorocera claripennis*,
 Eusisyropa virilis, *Exorista mella*, *Lespesia callosamiae*,
 L. frenchii, "Tachina" spp., *Winthemia quadripustulata*,
 W. spp., Tachinidae.
Calocalpe undulata, see *Hydria undulata* or *Hydria* spp. (including
 H. prunivorata and/or *H. undulata*)
Caloptenus bivittatus or *C. differentialis*, see *Melanoplus* spp.
 (including *M. bivittatus* and/or *M. differentialis*)
Caloptenus spretus, see *Melanoplus spretus*
Calosoma (Col.: Carabidae)
 calidum: *Sitophaga calosomae*, *S. georgiae*, *S. georgiae*
 probably.
 frigidum: *Sitophaga calosomae*, *S. georgiae*.
 peregrinator: *Sitophaga calosomae*, *S. georgiae*, *S. sp.*
 sycophanta: *Sitophaga calosomae*, *S. georgiae*, *S. georgiae*
 probably.
 "wilcoxi," see *C. willcoxi*.
 willcoxi: *Sitophaga calosomae*.
 spp.: *Sitophaga georgiae*.
Calpe (Lep.: Noctuidae)
 canadensis: *Compsilura concinnata*, *Winthemia sinuata* prob-
 ably.

Calpodes (Lep.: Hesperiiidae)

ethlius: *Eucelatoria australis*, *Eusisyropa blanda*, *Lixophaga diatraeae*.

Camnula (Orth.: Acrididae)

pellucida: *Ceracia dentata*, *Hemithrixion oestriforme*, Tachinidae.

Canarsia (Lep.: Pyralidae)

ulmiarrosorella: *Eufrontina spectabilis*.

Cane borer, see *Rhabdoscelus obscurus*

Caneborer(-s) (Col.: Curculionidae)

Caneborer(-s): *Myiophasia metallica*.

Cankerworm(-s) (Lep.: Geometridae)

Cankerworm(-s) (including *Alsophila pometaria* and/or *Paleacrita vernata*): "Tachina" spp.

Cankerworm(-s): *Winthemia rufopicta*.

Carabus (Col.: Carabidae)

nemoralis: *Sitophaga calosomae*.

taedatus variety, see *C. taedatus taedatus*.

taedatus var. near *oregonensis*, see *C. taedatus taedatus*.

taedatus taedatus: *Sitophaga leechi*.

sp.: Tachinidae.

Caradrina reculsa, see *Elydna nonagricola*

Caripeta (Lep.: Geometridae)

angustiorata: *Blondelia eufitchiae*.

Carneades (Lep.: Noctuidae)

messoria, see *Euxoa messoria*.

sp.: *Chaetogaedia monticola*.

Carolina grasshopper, see *Dissosteira carolina*

Carolina mantid, see *Stagmomantis carolina*

Carpocapsa pomonella, see *Laspeyresia pomonella*

Carpocapsa [sp.], see *Laspeyresia pomonella* or *Melissopus latiferreanus*

Case bearer, see *Acrobasis caryae*

Castnia (Lep.: Castniidae)

licoides: *Palpozenillia palpalis*.

Catabena (Lep.: Noctuidae)

lineolata: *Deopalpus miscelli*.

Catalpa sphinx, see *Ceratomia catalpae*

Catocala (Lep.: Noctuidae)

antinympha: *Chaetophlepsis* spp., *Winthemia* spp.

micronympha: *Tachinomyia nigricans*, *T.* spp.

micronympha form *fratercula*, see *C. micronympha*.

minuta: *Winthemia* spp.

neogama: *Archytas aterrimus*.

parta: Tachinidae.

relicta: *Actia* spp.

spp.: *Blepharipa scutellata*, *Compsilura concinnata*, *Eusisyropa blanda*, *Tachinomyia* sp. near *panaetius*.

Catocalinae (Lep.: Noctuidae)

Unidentified: *Actia* spp., *Archytas* sp. near *metallicus*, *Blepharipa scutellata*, *Compsilura concinnata*, *Eusisyropa blanda*, *Tachinomyia* spp., *Winthemia datanae*.

Ceanothus silk moth, see *Platysamia euryalus*

Cecidomyia "salicis brassicoides," see *Rhabdophaga salicisbrassicoides*

Cecropia moth, see *Platysamia cecropia*

Celama (Lep.: Noctuidae)

sorghiella: *Lespesia* spp.

Celastrina (Lep.: Lycaenidae)

argiolus pseudargiolus: *Aplomya theclarum*.

Celerio lineata, see *Hyles lineata*

Celery leaftier, see *Udea rubigalis*

Cerambycidae (Col.: Cerambycidae)

Unidentified: *Eutheresia monohammi* probably, *Zelia vertebrata*.

Ceramica (Lep.: Noctuidae)

picta: *Blondelia hyphantriae*, *Compsilura concinnata*, *Winthemia quadripustulata* probably, *W. rufopicta*, Tachinidae.

Ceramidia (Lep.: Ctenuchidae)

butleri: *Lespesia aletiae*.

Ceratomegilla fuscilabris, see *Coleomegilla maculata*

Ceratomia (Lep.: Sphingidae)

amyntor: *Argyrophylax* spp., *Belvosia bifasciata*, *B. townsendi*, *Drino incompta*, *Lespesia frenchii*.

catalpae: *Belvosia borealis*, *Compsilura concinnata*, *Drino incompta*, *Euphorocera claripennis*, *E. floridensis*, *Eusisyropa virilis*, *Hyphantrophaga hyphantriae*, *Lespesia aletiae*, *L. frenchii*, *L. spp.*, *Winthemia datanae*, Tachinidae.

undulosa: *Belvosia bifasciata*, *B. borealis*, *Compsilura concinnata*, *Drino incompta*, *Euphorocera claripennis*, *Hyphantrophaga hyphantriae*, *Lespesia frenchii*, Tachinidae.

undulosa probably: *Drino incompta*.

spp.: *Drino incompta*.

Cereal leaf beetle, see *Oulema melanopus*

- Cerocampa regalis*, see *Citheronia regalis*
- Cerotoma** (Col.: Chrysomelidae)
trifurcata: *Celatoria diabroticae*.
- Ceruchus** (Col.: Lucanidae)
piceus: *Eutheresia monohammi*, *E. monohammi* probably.
- Cerura** (Lep.: Notodontidae)
cinerea: *Aplomya cerurae*, Tachinidae.
occidentalis: *Compsilura concinnata*.
spp.: *Archytas aterrimus*, *Exorista* spp., *Lespesia aletiae*, *Pales* spp., "*Prosopaea*" sp.
- Ceuthophilus** (Orth.: Gryllacrididae)
latibuli: *Oedematocera flaveola*.
spp.: *Oedematocera flaveola* probably.
- Chainspotted geometer, see *Cingilia catenaria*
- Chalcodermus** (Col.: Curculionidae)
aeneus: *Myiophasia "aenea," M. clistoides*, *M. globosa*.
collaris: *Myiophasia globosa*.
inaequicollis: *Myiophasia globosa*.
spp.: *Myiophasia "aenea," M. globosa*.
- Charidryas nycteis**, see *Chlosyne nycteis*
- Chelinidea** (Hem. (Heter.): Coreidae)
canyona: *Trichopoda pennipes*.
tabulata: *Trichopoda pennipes*.
vittiger: *Trichopoda pennipes*.
vittiger aequoris: *Trichopoda pennipes*.
vittiger aequoris var. *artuatra*, see *C. vittiger aequoris*.
- Chelymorpha** (Col.: Chrysomelidae)
cassidea: *Eribella exilis*, *Eucelatoriopsis dimmocki*.
- Chemarachis** (Col.: Scarabaeidae)
hogardi: *Ptilodexia harpasa*.
vandinei: *Cryptomeigenia aurifacies*, *Eutrixoides jonesii*.
- Cherry ugly moths, see *Archips* spp. probably
- Chilo** (Lep.: Pyralidae)
plejadellus: *Paratheresia claripalpis*.
- Chimarocephala** (Orth.: Acrididae)
pacifica pacifica: *Ceracia dentata*, Tachinidae.
- Chionodes** (Lep.: Gelechiidae)
sp.: *Schizotachina* sp.
- "*Chlorachroa*" *ligata*, see *Chlorochroa ligata*
- Chloridea obsoleta*, see *Heliothis zea*
- Chlorippe celtis*, see *Asterocampa celtis*
- "*Chloristoneura*" *fumiferana*, see *Choristoneura fumiferana*
- Chlorochroa** (Hem. (Heter.): Pentatomidae)
ligata: *Cylindromyia euchenor*, *Gymnosoma filiola*, *G. fuliginosum*, *Paraphoranthia pollinosa*.

sayi: *Cylindromyia armata*, *C. euchenor*, *Gymnosoma fuliginosum*.

uhleri: *Gymnosoma fuliginosum*.

spp.: *Trichopoda* spp.

Chlosyne (Lep.: Nymphalidae)

harrisii: *Compsilura concinnata*, "Phorocera" spp., Tachinidae.

lacinia: *Euphorocera claripennis*.

nycteis: *Compsilura concinnata*.

Choephora (Lep.: Noctuidae)

fungorum: *Wagneria major*, *W. sp.*

Choristoneura (Lep.: Tortricidae)

conflictana: *Actia diffidens*, *A. interrupta*, *Aplomya caesar*, *A. crassiseta*, *Chaetophlepsis* spp., *Lypha setifacies*, *Madremyia saundersii*, *Omotoma fumiferanae*, *Phryxe pecosensis*, Tachinidae.

fractivittana: *Aplomya caesar*.

fumiferana: *Actia interrupta*, *Aplomya caesar*, *Ceromasia auricaudata*, *C. aurifrons*, *C. spp.*, *Compsilura concinnata*, *Hemisturmia tortricis*, *Lypha setifacies*, *Madremyia saundersii*, *Nemorilla pyste*, *Omotoma fumiferanae*, "Phorocera" *incrassata*, *Phryxe pecosensis*, *P. pecosensis* probably, *P. vulgaris*, *Pseudoperichaeta erecta*, "Sturmia" spp., *Tachinomyia nigricans*, *Winthemia* spp., *Xanthophyto* spp., Tachinidae

houstonana: *Nemorilla pyste*.

occidentalis: Tachinidae.

parallela: *Aplomya caesar*, *Madremyia saundersii*, *Nemorilla pyste*.

pinus: *Actia interrupta*, *A. spp.*, *Aplomya caesar*, *Hemisturmia tortricis*, *Lypha setifacies*, *Madremyia saundersii*, *Nemorilla pyste*, *N. spp.*, *Phryxe pecosensis*, *Pseudoperichaeta erecta*, Tachinidae.

rosaceana: *Actia interrupta*, *Aplomya caesar*, *Erynnia tortricis*, *Hemisturmia tortricis*, *Lypha setifacies*, *Nemorilla pyste*, *Phryxe pecosensis*, *Pseudoperichaeta erecta*, Tachinidae.

Chorizagrotis agrestis, see *Euxoa auxiliaris*

Chorizagrotis auxiliaris, see *Euxoa auxiliaris*

Chortophaga (Orth.: Acrididae)

viridifasciata: *Ceracia dentata*, *C. dentata* probably.

spp.: *Ceracia dentata*.

Chromatia (Col.: Alleculidae)

amoena: *Hyalomyodes triangulifer*.

Chrysodeixis (Lep.: Noctuidae)

calcites: *Chaetogaedia monticola*, *Eucelatoria armigera*,
Tachinidae.

Chrysomela (Col.: Chrysomelidae)

crotchi: *Grisdalemyia* spp.

interrupta: *Hyalomyodes triangulifer*.

knabi: *Hyalomyodes triangulifer*.

multipunctata, see *Calligrapha multipunctata*.

scripta: Tachinidae.

Chrysomelid (Col.: Chrysomelidae)

spp. (including *Lema trilineata* and/or *Leptinotarsa decemlineata*): *Adoryphorophaga aberrans*, *Doryphorophaga australis*, *D. doryphorae*.

Chrysomelidae (Col.: Chrysomelidae)

Unidentified: *Medina* spp.

Chrysophanus *dione*, see *Lycaena xanthoides dione***Chrysops** (Dipt.: Tabanidae)

sp.: *Carinosillus tabanivorus*.

Cicada septendecim, see *Magicicada septendecim***Cicinnus** (Lep.: Lacosomidae)

melsheimeri: *Chrysotachina alcedo*.

Cimbex (Hym.: Cimbicidae)

americana: *Aplomya epicydes*, *Compsilura concinnata*, Tachinidae.

Cinclidia harrisii, see *Chlosyne harrisii***Cingilia** (Lep.: Geometridae)

"*catanaria*," see *C. catanaria*.

catanaria: *Blondelia* spp. probably, *Ceromasia aurifrons*,
Chaetophlepsis orbitalis, *Compsilura concinnata*,
Euphorocera claripennis, *Eusisyropa blanda*, *E. virilis*,
Madremyia saundersii, *Phryxe pecosensis*, *P. vulgaris*,
Winthemia quadripustulata, *W. sinuata* probably,
Tachinidae.

catanaria probably: *Winthemia sinuata*.

latanaria, see *C. catanaria* probably.

Cirphis (Lep.: Noctuidae)

cholica, see *Leucania cholica*.

inconspicua, see *Leucania inconspicua*.

"*inconspicua*," see *Leucania inconspicua*.

juncicola, see *Leucania juncicola*.

latiuscula, see *Leucania latiuscula*.

latiuscula probably, see *Leucania latiuscula* probably.

microgonia, see *Leucania microgonia*.

unipuncta, see *Pseudaletia unipuncta*.

unipuncta and/or *Spodoptera mauritia*, see Armyworm(-s)
(including *Pseudaletia unipuncta* and/or *Spodoptera mauritia*).

spp.: *Archytas cirphis*, *Periscepsia helymus*, Tachinidae.

“Cirphus” *unipuncta*, see *Pseudaletia unipuncta*

Cisthene (Lep.: Arctiidae)

nexa: *Pseudochaeta argentifrons*.

“*Citherona*” *regalis*, see *Citheronia regalis*

Citheronia (Lep.: Citheroniidae)

regalis: *Belvosia bifasciata*, *B. townsendi*, *Lespesia archippi-
vora*, *L. frenchii*, *L. spp.*, *Winthemia citheroniae*.

Cladius (Hym.: Tenthredinidae)

isomerus: *Bessa harveyi*.

pectinicornis: *Bessa harveyi*.

Claybacked cutworm, see *Agrotis gladiara*

Clearwinged grasshopper, see *Camnula pellucida*

Cleora pampinaria, see *Isturgia truncataria*

Clisiocampa americanum, see *Malacosoma americanum*

Clisiocampa californicum, see *Malacosoma californicum*

Clisiocampa constricta, see *Malacosoma constrictum*

Clisiocampa disstria, see *Malacosoma disstria*

Clisiocampa pluvialis, see *Malacosoma californicum pluviale*

Clisiocampa sylvatica, see *Malacosoma disstria*

Clisiocampa “thoracia,” see *Malacosoma californicum californicum*

Clisiocampa thoracica, see *Malacosoma californicum californicum*

Clisiocampa spp., see *Malacosoma spp.*

Clover cutworm, see *Scotogramma trifolii*

Clover leaf weevil, see *Hypera punctata*

Clover root curculio, see *Sitona hispidula*

Cnidocampa (Lep.: Limacodidae)

flavescens: *Chaetexorista javana*, *Compsilura concinnata*,
Lixophaga mediocris.

Coccinella (Col.: Coccinellidae)

trifasciata perplexa: *Hyalomyodes triangulifer*.

Cocklebur weevil, see *Rhodoaenus tredecimpunctatus*

Cockroach(-es) (Orth.: Blattidae)

Cockroach(-es): *Calodexia dives*, *C. spp.*

Coconut leafroller, see *Hedylepta blackburni*

“Coddling” moth, see *Laspeyresia pomonella*

Codling moth, see *Laspeyresia pomonella*

Coelodasys unicornis, see *Schizura unicornis*

Coleomegilla (Col.: Coccinellidae)

maculata: *Hyalomyodes triangulifer*.

maculata lengi: *Hyalomyodes triangulifer*.

Coleophora (Lep.: Coleophoridae)*fuscedinella*: *Erynnia tortricis*.*malivorella*: *Erynnia tortricis*.*salmani*, see *C. fuscedinella*.**Coleoptera** (Col.: Unidentified Coleoptera)Miscellaneous: *Euphorocera claripennis*, "Macquartia" spp.,
*Myiophasia "aenea."***Coleotechnites** (Lep.: Gelechiidae)*canusella*: *Elfia* spp.*coniferella*: *Schizactia vitinervis*.*piceaella*: *Schizactia vitinervis*.*thujella*: *Elfia* spp.sp.: *Schizactia vitinervis*.**Coleotichus** (Hem. (Heter.): Scutelleridae)*blackburniae*: *Trichopoda pennipes*, *T. pilipes*.**Colias** (Lep.: Pieridae)*eurytheme*: *Euphorocera claripennis*, *E. omissa*, *Lespesia*
archippivora, *L.* spp., "Masicera" spp., "Phorocera" spp.*philodice eurytheme*, see *C. eurytheme*.**Coloradia** (Lep.: Saturniidae)*pandora*: *Leschenaultia adusta*.Colorado potato beetle, see *Leptinotarsa decemlineata*Columbine borer, see *Papaipema purpurifascia*Conchuela, see *Chlorochroa ligata***Cone moth(-s)** (Lep.: Pyralidae)Cone moth(-s): *Elfia setigera*.Cone moth(-s) probably: *Lydellohoughia* sp.**Coniodes** (Lep.: Geometridae)*plumogeraria*: Tachinidae.**Conistra devia**, see *Eupsilia devia***Conistra indirecta**, see *Pyreferra hesperidago***Conistra morrisoni** and/or *C. sidus*, see *Eupsilia* spp. (including *E.*
morrisoni and/or *E. sidus*)**Conotrachelus** (Col.: Curculionidae)*affinis*: *Cholomyia inaequipes*, *Myiophasia "aenea," M. globosa*,
M. nigrifrons.*anaglypticus*: *Myiophasia globosa*.*aratus*: *Cholomyia inaequipes*, *Myiophasia globosa*.*elegans*: *Cholomyia inaequipes*, *Myiophasia "aenea,"* Tachini-
dae.*juglandis*: *Cholomyia inaequipes*, *Myiophasia "aenea."**naso*: *Cholomyia inaequipes*, *C.* sp.*nenuphar*: *Cholomyia inaequipes*, *Myiophasia "aenea," M.*
"aenea" probably, *M. globosa*.

- posticatus*: *Cholomyia inaequipes*.
 spp. (including *C. affinis* or *C. posticatus*): *Myiophasia*
 “*aenea*.”
 spp.: *Cholomyia inaequipes*, *Myiophasia* “*aenea*,” *M. metallica*.
- Copablepharon (Lep.: Noctuidae)
 viridisparsa: *Bonnetia comta*.
- Copaxa (Lep.: Saturniidae)
 lavendera probably: *Belvosia ciliata*.
- “Cophyrus” larvae, see Diprionid spp. (including *Diprion* spp.
 and/or *Neodiprion* spp.)
- Coptocyclus (Col.: Chrysomelidae)
 clavata, see *Plagiometriona clavata*.
 spp.: *Eucelatoriopsis dimmocki*, “*Tachina*” spp.
- Coreus tristis*, see *Anasa tristis*
- Corizus hyalinus*, see *Liorhyssus hyalinus*
- Corn borer, see *Ostrinia nubilalis*
- Corn earworm, see *Heliothis zea*
- Corn stalk borer(-s) (Lep.: Unidentified Lepidoptera)
 Corn stalk borer(-s): *Archytas marmoratus*.
- Coryphista (Lep.: Geometridae)
 meadii: *Blondelia* spp., *Madremyia saundersii*.
- Cosmia (Lep.: Noctuidae)
 canescens: *Mericia* spp., Tachinidae.
- Cosmopepla (Hem. (Heter.): Pentatomidae)
 bimaculata: *Alophorella aeneoventris*, *Gymnoclytia dubia*.
- Cosmophila sabulifera*, see *Anomis hawaiiensis*
- Cossula (Lep.: Cossidae)
 magnifica: “*Phorocera*” *comstocki*, “*P.*” *signata*.
- Cosymbia (Lep.: Geometridae)
 lumenaria, see *C. pendulinaria*.
 pendulinaria: *Actia flavipes*.
- Cotton boll weevil, see *Anthonomus grandis*
- Cotton leafworm, see *Alabama argillacea*
- Cotton square borer, see *Strymon melinus*
- Cottonwood dagger moth, see *Acronicta lepusculina*
- Cottonwood leaf beetle, see *Chrysomela scripta*
- Cotton worm, see *Alabama argillacea*
- Cowpea curculio, see *Chalcodermus aeneus*
- Crambus (Lep.: Pyralidae)
 bonifatellus, see *C. inornatellus*.
 inornatellus: *Aplomya confusionis*.
 mutabilis: *Aplomya caesar*, *A. confusionis*, *Euphorocera*
 claripennis.
 sperryellus: *Aplomya confusionis*.

- trisectus*: *Aplomya caesar*, *A. confusionis*, *Stomatomyia floridensis*.
- vulgivagellus*: "Tachina" spp.
- spp.: *Aplomya caesar*, *Euphorocera claripennis*, *Ormia* spp. probably, *Phasiops flavus*, Tachinidae.
- Cranberry spanworm, see *Anavitrinella pampinaria*
- Crane fly (flies) (Dipt.: Tipulidae)
- Crane fly (flies): *Siphona* spp.
- Cressonia (Lep.: Sphingidae)
- juglandis*: *Winthemia datanae*, Tachinidae.
- Cricket(-s) (Orth.: Gryllidae)
- Cricket(-s): *Calodexia* spp., *Gilvella gilvipes*.
- Crinkled flannel moth, see *Megalopyge crispata*
- Crioceris (Col.: Chrysomelidae)
- asparagi*: *Meigenia mutabilis*, *Paralispe infernalis*.
- "Criphis" *unipuncta*, see *Pseudaletia unipuncta*
- Crocigrapha (Lep.: Noctuidae)
- normani*: *Patelloa* spp.
- Crocota *rubicundaria*, see *Eubaphe aurantiaca rubicundaria*
- Croesia (Lep.: Tortricidae)
- albicomana*: *Actia interrupta*, *Lypha setifacies*.
- albicomana* probably: *Schizotachina convecta*.
- semipurpurana*, see *C. albicomana*.
- Croesus (Hym.: Tenthredinidae)
- latitarsus*: *Bessa harveyi*, *Compsilura concinnata*.
- Crossstriped cabbageworm, see *Evergestis rimosalis*
- Crymodes (Lep.: Noctuidae)
- devastator*: *Gonia aldrichi*, *G. frontosa*, *G.* spp., *Nowickia latifrons*, Tachinidae.
- Cryphia (Lep.: Noctuidae)
- pervertens*: *Chaetophlepsis orbitalis*.
- Ctenophora (Dipt.: Tipulidae)
- spp.: *Admontia* spp.
- Ctenucha (Lep.: Ctenuchidae)
- virginica*: *Compsilura concinnata*, Tachinidae.
- Cucullia (Lep.: Noctuidae)
- convexipennis*: *Winthemia quadripustulata*, *W. rufopicta*, Tachinidae.
- spp.: *Lespesia cuculliae*.
- Cucumber beetle(-s) (Col.: Chrysomelidae)
- Cucumber beetle(-s): *Celatoria setosa*.
- Curculio (Col.: Curculionidae)
- auriger*: *Myiophasia nigrifrons*.
- nasicus*: *Myiophasia "aenea."*

nenuphar, see *Conotrachelus nenuphar*.

proboscideus: *Myiophasia nigrifrons*, *Winthemia quadripustulata*.

strictus probably: *Myiophasia nigrifrons*.

uniformis: *Myiophasia* "aenea," "M. atra."

spp.: *Clausicella neomexicana*, *Elfia melissopodis*, *Myiophasia metallica*, *M. nigrifrons*.

Curculionid (Col.: Curculionidae)

spp. (including *Conotrachelus naso* or *Curculio pardalis*):
Myiophasia "aenea."

Curculionidae (Col: Curculionidae)

Unidentified: *Myiophasia globosa*, *Winthemia quadripustulata*.

Curled rose sawfly, see *Allantus cinctus*

Currant spanworm, see *Itame ribearia*

Cutworm, see *Euxoa messoria* probably

Cutworm(-s) (Lep.: Noctuidae)

Cutworm(-s) (including *Agrotis* spp. and/or *Leucania* spp.): *Chaetogaedia monticola*, *Lespesia archippivora*.

Cutworm(-s) and related Noctuidae: *Winthemia quadripustulata*.

Cutworm(-s): *Archytas cirphis*, *A. marmoratus*, *A.* spp. (including *A. apicifer* and/or *A. californiae*), *Bonnetia comta*, *Chaetogaedia crebra*, *C. monticola*, *Eucelatoria comosa*, *Euphorocera claripennis*, *Gonia porca*, *G.* spp., *Lespesia archippivora*, *Nowickia algens*, *Periscepsia helymus*, "Tachina" spp., *Tachinomyia panaetius*, *Winthemia leucanae*, *W. quadripustulata*, *W.* spp., Tachinidae.

Cyaniris pseudargiolus, see *Celastrina argiolus pseudargiolus*

Cyclocephala immaculata or *Lachnosterna* spp., see Scarabaeid spp. (including *C. immaculata* or *Phyllophaga* spp.)

Cyclocephala spp. or *Lachnosterna* spp., see Scarabaeid spp. (including *Cyclocephala* spp. and/or *Phyllophaga* spp.)

Cycnia (Lep.: Arctiidae)

inopinatus: *Hyphantrophaga euchaetiae*, *H. hyphantriae*.

tenera: *Archytas aterrimus*, *Exorista mella*, *Hyphantrophaga euchaetiae*, *H. hyphantriae*, Tachinidae.

Cymaenes (Lep.: Hesperiiidae)

tripunctus: Tachinidae.

Cynthia atalanta, see *Vanessa atalanta*

Cynthia cardui, see *Vanessa cardui*

Cynthia cardui "J. rufipes," see *Vanessa cardui*

Cynthia moth, see *Samia cynthia*

Cynthia virginiana, see *Vanessa virginiana*

"Danais" *archippus*, see *Danaus plexippus plexippus*

Danaus (Lep.: Danaidae)

archippus, see *D. plexippus plexippus*.

berenice, see *D. gilippus berenice*.

gilippus berenice: *Lespesia archippivora*.

menippe, see *D. plexippus plexippus*.

plexippus, see *D. plexippus plexippus*.

plexippus plexippus: *Buquetia obscura*, *Chaetogaedia monticola*, *Compsilura concinnata*, *Eusisyropa virilis*, *Exorista mella*, *E. spp.*, *Lespesia archippivora*, *L. schizurae*, *L. spp.*, *Madremyia saundersii*, *Phryxe pecosensis*, *P. vulgaris*, Tachinidae.

"Dantana" *integerrima*, see *Datana integerrima*

Darksided cutworm, see *Euxoa messoria*

Dasychira (Lep.: Lymantriidae)

atomeria: *Carcelia olenensis*.

basiflava: *Compsilura concinnata*, *Exorista mella*, *E. spp.*

groenlandica, see *Byrdia groenlandica*.

plagiata: *Carcelia olenensis*, *C. perplexa*, *C. spp.*

vagans: *Exorista mella*.

spp.: *Compsilura concinnata*.

spp. probably: *Spoggosia gelida*.

"Dasychirus" sp. (?), see *Dasychira spp. probably*

"Dasylopha" *anguina*, see *Dasylophia anguina*

Dasylophia (Lep.: Notodontidae)

anguina: *Lespesia aletiae*, *L. cuculliae*, *L. cuculliae probably*, Tachinidae.

Dasystoma (Lep.: Oecophoridae)

salicellum: *Compsilura concinnata*.

Datana (Lep.: Notodontidae)

angusii: *Archytas metallicus*, *Compsilura concinnata*, *Euphorocera claripennis*, *Lespesia cuculliae*, *L. frenchii*, *Winthemia datanae*.

contracta: *Euphorocera claripennis*, *Exorista spp. probably*.

integerrima: *Archytas metallicus*, *A. spp.*, *Belvosia bifasciata*, *Blepharipa scutellata*, *Carcelia malacosomae*, *C. spp.*, *Compsilura concinnata*, *Euphorocera claripennis*, *E. edwardsii*, *E. floridensis*, *Exorista mella*, *Lespesia datanarum*, *L. frenchii*, *L. frenchii complex*, *Mercia ampelus*, *Winthemia datanae*, Tachinidae.

major: *Compsilura concinnata*, *Euphorocera floridensis*.

ministra: *Archytas metallicus*, *Compsilura concinnata*, *Euphorocera claripennis*, *E. edwardsii*, *E. floridensis*, *Exorista mella*, *E. spp. probably*, *Lespesia aletiae*, *L.*

datanarum, *L. frenchii*, *L. frenchii* complex, *L. spp.*,
Mericia ampelus, *M. spp.*, *Palpexorista coccyx*, *Patelloa*
leucaniae, *Winthemia datanae*, *W. leucanae* lacking red
 at tip, *W. quadripustulata*, Tachinidae.

perspicua: *Archytas metallicus*, *Compsilura concinnata*,
Euphorocera claripennis, *Winthemia datanae*, Tachi-
 nidae.

spp.: *Euphorocera claripennis*, *Lespesia datanarum*, *L. frenchii*,
Winthemia datanae, Tachinidae.

Deidamia (Lep.: Sphingidae)

inscriptum: *Compsilura concinnata*.

Deilephila gallii, see *Hyles gallii*

Deilephila lineata, see *Hyles lineata*

Deilinia (Lep.: Geometridae)

erythemaria: Tachinidae.

Deloyala clavata, see *Plagiometriona clavata*

"**Deloyola**" *clavata*, see *Plagiometriona clavata*

Dendrobinus howardi, see *Gloveria howardi*

Depressaria (Lep.: Oecophoridae)

"*heracleana*," see *D. pastinacella*.

heracliana, see *D. pastinacella*.

pastinacella: *Dichaetoneura leucoptera*, *Eusisyropa blanda*,
Nemorilla pyste, *Xanthophyto spp.*, Tachinidae.

spp.: *Eusisyropa blanda*.

Desmia (Lep.: Pyralidae)

funeralis: *Dejeaniopalpus texensis*, *Erynnia tortricis*, *Exorista*
spp., *Hyphantrophaga desmiae*, *H. hyphantriae*,
Leskiomima tenera, *Lixophaga variabilis*, *Nemorilla*
pyste, *Pseudoperichaeta erecta*, "*Tachina desmiae*,"
 "*Tachinophyto*" *spp.*

maculalis, see *D. funeralis*.

"**Detana**" *ministra*, see *Datana ministra*

Deuteronomos (Lep.: Geometridae)

magnarius: *Madremyia saundersii*.

"**Diabrothica**" *vittata*, see *Acalymma vittata*

Diabrotica (Col.: Chrysomelidae)

"*duodecimpunctata*," see *D. undecimpunctata* or *D. undecim-*
punctata howardi.

duodecimpunctata, see *D. undecimpunctata* or *D. undecim-*
punctata howardi.

soror, see *D. undecimpunctata* or *D. undecimpunctata*
undecimpunctata.

trivittata, see *Acalymma trivittata*.

undecimpunctata, see *D. undecimpunctata* or *D. undecimpunctata undecimpunctata*.

undecimpunctata: *Celatoria diabroticae*, *C. setosa*, *Pseudomyothyria ancilla*, Tachinidae.

undecimpunctata howardi: *Celatoria diabroticae*, *C. setosa*, Tachinidae.

undecimpunctata undecimpunctata: *Celatoria diabroticae*, Tachinidae.

vittata, see *Acalymma vittata*.

spp.: *Celatoria diabroticae*.

11-punctata, see *D. undecimpunctata undecimpunctata*.

12-punctata, see *D. undecimpunctata* or *D. undecimpunctata howardi*.

Diacrisia (Lep.: Arctiidae)

virginica: *Aplomya caesar*, *Blondelia hyphantriae*, *Bombyliopsis abrupta*, *Carcelia diacrisiae*, *C. reclinata*, *Compsilura concinnata*, *Exorista mella*, *E. spp.*, *Gymnocarcelia ricinorum*, *Hubneria estigmenensis*, *Lespesia aletiae*, *L. frenchii*, *Mericia ampelus*, *M. spp.*, *Thelaira americana*, *Winthemia datanae*, Tachinidae.

Diaphania (Lep.: Pyralidae)

hyalinata: "*Leskia analis*," *Nemorilla pyste*, *Stomatodexia cothurnata*, Tachinidae.

nitidalis probably: *Thelaira* spp.

Diapheromera (Orth.: Phasmatidae)

femorata: *Euhalidaya genalis*, *Phasmophaga antennalis*, "*Tachina*" spp., Tachinidae.

spp.: *Euhalidaya genalis*.

"*Diapheromerata*" *femorata*, see *Diapheromera femorata*

"*Diarnea*" sp., see *Diatraea* spp.

Diatraea (Lep.: Pyralidae)

canella: *Lixophaga diatraeae*, *Metagonistylum minense*.

considerata: *Palpozenillia palpalis*, *Paratheresia claripalpis*.

crambidoides: *Lixophaga diatraeae*, *L. sphenophori*.

grandiosella: *Paratheresia claripalpis*.

impersonatella: *Lixophaga* spp., *Metagonistylum minense*.

lineolata, see *Zeadiatraea lineolata*.

lineolata probably, see *Zeadiatraea lineolata* probably.

magnifactella: *Paratheresia claripalpis*.

"*saccharalia*," see *D. saccharalis*.

saccharalis: "*Hypostena floridensis*," *Lixophaga diatraeae*, *L. grisea*, *L. spp.*, *Metagonistylum minense*, *Nemorilla pyste*, *Paratheresia claripalpis*, *P. spp.*, "*Phorostoma*" spp., "*Tachinophyto*" spp., Tachinidae.

saccharalis var. *crambidoides*, see *D. crambidoides* or *D. saccharalis*.

tabernella: *Paratheresia claripalpis*.

[sp.] in sugar cane, see *D. considerata*.

spp. (including *D. impersonatella* and/or *D. saccharalis*):
Paratheresia spp.

spp.: *Lespesia* sp., *Lixophaga diatraeae*, *Metagonistylum minense*, *Paratheresia claripalpis*, *P. spp.*, "Phorocera" sp., "Phorostoma" spp., Tachinidae.

spp. probably: *Paratheresia claripalpis*.

"Diatrea" *impersonatella*, see *Diatraea impersonatella*

"Diatrea" *impersonatella* or "D." *saccharalis*, see *Diatraea* spp.
(including *D. impersonatella* and/or *D. saccharalis*)

"Diatrea" *saccharalis*, see *Diatraea saccharalis*

Dicentria (Lep.: Notodontidae)

lignicolor: *Winthemia datanae*.

Dichogama (Lep.: Pyralidae)

redtenbacheri: *Eusisyropa blanda*, *E. boarmiae*.

"Dichogamma" *bergi*, see *Alatuncusia bergi*

"Dichogamma" *redtenbacheri*, see *Dichogama redtenbacheri*

Differential grasshopper, see *Melanoplus differentialis*

Differential locust, see *Melanoplus differentialis*

Dingy cutworm, see *Feltia subgothica*

Dione vanillae, see *Agraulis vanillae*

Dioryctria (Lep.: Pyralidae)

abietella: *Elfia setigera*, *Leskiomima tenera*, Gen. and sp. near *Oxynops*, Tachinidae.

abietella probably: *Phrynofrontina* spp.

amatella: *Phrynofrontina* spp., *Xanthophyto* spp.

cambiicola: Gen. and sp. near *Oxynops*.

clarioralis: *Phrynofrontina* spp.

disclusa: Gen. and sp. near *Oxynops*.

reniculella: *Aplomya caesar*, *Elfia curriei*, *Leskiomima* sp., *Phryxe pecosensis*.

spp. and/or *Laspeyresia* spp. probably, see Olethreutid and/or pyralid spp. (including *Laspeyresia* spp. and/or *Dioryctria* spp. probably).

Diprion (Hym.: Diprionidae)

frutetorum: *Diplostichus lophyri*, *Drino inconspicua*.

hercyniae: *Bessa harveyi*, *B. selecta*, *Blondelia* spp. probably, *Diplostichus lophyri*, *Drino bohémica*, *D. inconspicua*, *D. inconspicua* probably, *Euphorocera edwardsii*, *Guerinia dydas*, *Spathimeigenia spinigera*, *S. spp.*

polytomum: *Bessa selecta*, *Catagoniopsis* sp. near *specularis*,
Diplostichus lophyri, *Drino bohémica*, *D. inconspicua*,
Euphorocera sp. near *claripennis*, *Spathimeigenia*
aurifrons, *S.* spp.

sertifer, see *Neodiprion sertifer*.

"*simile*," see *D. similis*.

similis: *Compsilura concinnata*, *Diplostichus lophyri*, *Drino*
inconspicua, *Euphorocera claripennis*, *E. edwardsii*,
Winthemia quadripustulata, Tachinidae.

sp. probably, see *Neodiprion* spp.

Diprionid (Hym.: Diprionidae)

spp. (including *Diprion* spp. and/or *Neodiprion* spp.): *Crypto-*
meigenia spp., *Diplostichus lophyri*, *Euphorocera*
claripennis, "*Phorocera*" *comstocki*, *Spathimeigenia*
spinigera.

"*Disomycha xanthomelaena*," see *Disonycha xanthomelas*

Disonycha (Col.: Chrysomelidae)

"*xanthomelaena*," see *D. xanthomelas*.

xanthomelas: *Medina barbata*.

"*xanthomeloena*," see *D. xanthomelas*.

Dissosteira (Orth.: Acrididae)

carolina, see *D. carolina* or *Melanoplus sanguinipes*.

carolina: *Lespesia frenchii*, *L.* spp.

cornuta, see *D. pictipennis*.

longipennis: Tachinidae.

pictipennis: *Trichopoda plumipes*, *T. plumipes* probably, *T.*
subdivisa.

spurcata: *Hemithrixion* sp. probably.

venusta, see *D. pictipennis*.

spp.: *Trichopoda* spp.

"*Dittraea*" *saccharalis*, see *Dittraea saccharalis*

Doryphora decemlineata, see *Leptinotarsa decemlineata*

Doryphora 10-lineata, see *Leptinotarsa decemlineata*

"*Doryphorae*" *10-lineata*, see *Leptinotarsa decemlineata*

Doryphore, see *Leptinotarsa decemlineata*

Douglas fir cone moth, see *Barbara colfaxiana*

Douglas fir tussock moth, see *Hemerocampa pseudotsugata*

Drepana (Lep.: Drepanidae)

arcuata: *Actia palloris*.

bilineata: *Actia palloris*, Tachinidae.

bilineata form *levis*, see *D. bilineata*.

Drepanulatrix (Lep.: Geometridae)

monicaria: *Chaetophlepsis rindgei*.

spp. (including *D. baueraria* and/or *D. monicaria*): *Chaeto-*
phlepsis rindgei, *Patelloa* spp.

- Dryocampa rubicunda*, see *Anisota rubicunda*
 Dusky birch sawfly, see *Croesus latitarsus*
 Dusky stink bug, see *Euschistus tristigmus*
Dysdercus (Hem. (Heter.): Pyrrhocoridae)
 andreae: *Acaulona erythropyla*, *A. peruviana*.
 sanguinarius: *Acaulona erythropyla*, *A. peruviana*.
 spp.: *Acaulona erythropyla*.
 "Eacle *ismperialis*," see *Eacles imperialis*
Eacles (Lep.: Citheroniidae)
 imperialis: *Belvosia bifasciata*, *B. townsendi*, *Drino incompta*,
 Lespesia frenchii, *Winthemia citheroniae*, Tachinidae.
 Earwig, see *Forficula auricularia*
 Eastern blackheaded budworm, see *Acleris variana*
 Eastern tent caterpillar, see *Malacosoma americanum*
Eciton (Hym.: Formicidae)
 burchelli foreli probably: *Androeuryps ecitonis*.
 foreli probably, see *E. burchelli foreli* probably.
Ecpantheria (Lep.: Arctiidae)
 deflorata: *Aplomya helvina*, *Archytas metallicus*, *Carcelia*
 reclinata, *Gymnocarcelia ricinorum*, *G. ricinorum*
 probably, *Juriniopsis adusta*, *J. floridensis* probably,
 Leschenaultia adusta.
 eridanus: *Carcelia amplexa*.
 icasia: *Carcelia amplexa*, *C. lagoae*, *C. spp.*, *Leschenaultia*
 leucophrys, *Lespesia spp.*
 permaculata, see *Turuptiana permaculata*.
 scribonia, see *E. deflorata*.
 Eightspotted forester, see *Alypia octomaculata*
Elaphria (Lep.: Noctuidae)
 nucicolora: *Archytas spp.* (including *A. apicifer* and/or *A.*
 californiae), *Periscepsia helymus*, *P. laevigata*.
Elasmopalpus (Lep.: Pyralidae)
 lignosellus: *Stomatomyia floridensis*.
 Elder shoot borer, see *Achatodes zeae*
Eleodes (Col.: Tenebrionidae)
 extricata: *Eleodiphaga caffreyi*.
 hispilabris: *Eleodiphaga pollinosa*.
 obsoleta: *Eleodiphaga caffreyi*.
 opacus: Tachinidae.
 tricostata: *Sitophaga eleodivora*, Tachinidae.
Ellopia fiscellaria, see *Lambdina fiscellaria*
Ellopia "lactea," see *Lambdina fiscellaria laeta* probably
Ellopia somniaria, see *Lambdina fiscellaria somniaria*
 Elm leaf beetle, see *Pyrrhalta luteola*

- Elm sawfly, see *Cimbex americana*
- Elm spanworm, see *Ennomos subsignarius*
- Elm sphinx, see *Ceratonia amyntor*
- Elophila fulicalis*, see *Parargyractis fulicalis*
- Elydna (Lep.: Noctuidae)
nonagricola: *Chaetogaedia monticola*, *Lespesia archippivora*.
- Ematurga (Lep.: Geometridae)
amitaria: *Winthemia quadripustulata*.
- Emblethis (Hem. (Heter.): Lygaeidae)
vicarius: *Hyalomya aldrichii*.
- Emphytus canadensis, see *Ametastegia pallipes*
- Emphytus cinctus, see *Allantus cinctus*
- Empretia stimulea, see *Sibine stimulea*
- Enaphalodes (Col.: Cerambycidae)
atomarius: *Theresia rutilans*, Tachinidae.
- Endomychus (Col.: Endomychidae)
biguttatus: *Phyllophilopsis evanida*.
- Endothenia (Lep.: Olethreutidae)
hebesana: *Clausicella floridensis*.
- Eneoptera (Orth.: Gryllidae)
spp.: *Calodexia* sp. near *agilis* or *interrupta*.
- English grain aphid, see *Macrosiphum avenae*
- Ennomos (Lep.: Geometridae)
magnarius, see *Deuteronomos magnarius*.
subsignarius: *Actia ontario*, *A.* sp. near *palloris*, *Blondelia eufitchiae*, *Chaetogaedia analis*, *Compsilura concinnata*, *Euexorista futilis*, *Euphorocera floridensis*, *Eusisyropa blanda*, *E. virilis*, *Lespesia aletiae*, *Madremyia saundersii*, *Mericia* spp., *Phryxe pecosensis*, *P. vulgaris*, *Tachinomyia nigricans*, *Xanthophyto* sp. probably, Tachinidae.
- Epargyreus (Lep.: Hesperidae)
clarus: *Acroglossa hesperidarum*, *Compsilura concinnata*, *Euphorocera claripennis*, *E. floridensis*, *Eusisyropa blanda*, *E. virilis*, *Juriniopsis adusta*, *Lespesia aletiae*, Tachinidae.
tityrus, see *E. clarus*.
- "Epiblana" strenuanum, see *Epiblema strenuanum*
- Epiblema (Lep.: Olethreutidae)
otiosanum: *Lixophaga variabilis*, *Lydella radialis*.
otiosanum probably: *Lixophaga variabilis*.
scudderiana: *Lixophaga variabilis*.
strenuanum: *Dichaetoneura leucoptera*, *Lixophaga variabilis*.
tripartitana: *Clausicella floridensis*, "Tachinophyto" spp.

“*Epibleme*” *otiosanum*, see *Epiblema otiosanum*

Epicauta (Col.: Meloidae)

sp.: *Pseudoperichaeta erecta*.

Epicnaptera (Lep.: Lasiocampidae)

americana: *Compsilura concinnata*, *Exorista mella*, Tachinidae.

Epiglaea (Lep.: Noctuidae)

apiata: *Cyrtophleba coquilletti*, *Madremyia saundersii*,
Periscepsia helymus, *Tachinomyia panaetius*,
Winthemia quadripustulata probably, *W. rufopicta*
probably, Tachinidae.

Epilachna (Col.: Coccinellidae)

borealis: *Aplomyiopsis epilachnae*, *Euphorocera claripennis*.
corrupta, see *E. varivestis*.

defecta: *Aplomyiopsis epilachnae*, *A. epilachnae* probably.

mexicana: *Aplomyiopsis epilachnae*.

obscurella: *Aplomyiopsis epilachnae*.

varivestis: *Aplomyiopsis epilachnae*, *Doryphorophaga*
doryphorae, *Euphorocera claripennis*, *E. floridensis*,
Helioplaga slossonae, *Hyalomyodes triangulifer*,
Lespesia aletiae, *Nemorilla pyste*, Tachinidae.

virgata: *Aplomyiopsis epilachnae*.

“*Epilachnae*” *borealis*, see *Epilachna borealis*

“*Epilachne varivestris*,” see *Epilachna varivestis*

Epilampra (Orth.: Blattidae)

azteca: *Calodexia* spp.

Epinotia (Lep.: Olethreutidae)

aceriella: *Elfia* spp.

crenana probably: *Actia interrupta*.

emarginana: *Actia interrupta*.

nanana: *Schizactia vitinervis*.

similana: *Actia* spp.

solicitana: *Eulasiona comstocki*, *Psilidopteryx psilocorsiphaga*.

Episimus (Lep.: Olethreutidae)

argutanus: *Actia* spp., *Nemorilla pyste*, *Pseudoperichaeta*
erecta.

Epsilia fungorum, see *Choephora fungorum*

Erannis (Lep.: Geometridae)

tiliaria: *Ictericophyto spinosa*, *Phryxe pecosensis*, *P. vulgaris*,
Pseudotachinomyia slossonae, *Tachinomyia nigricans*,
Winthemia sinuata, Tachinidae.

Eremocoris (Hem. (Heter.): Lygaeidae)

opacus: *Catharosia lustrans*.

sp.: *Catharosia lustrans*.

Eriopus floridensis, see *Callopietria floridensis*

Eriopyga cynica, see *Orthodes cynica*

Eriopyga rufula, see *Protorthodes rufula*

Ermine moth, see *Yponomeuta padella*

Erynephila (Col.: Chrysomelidae)

maritima: "*Hypostena*" sp.

Erynnis (Lep.: Hesperiiidae)

brizo: *Compsilura concinnata*, *Eusisyropa blanda*.

icelus: *Compsilura concinnata*, *Drino fraudulenta*, Tachinidae.

spp. (including *E. horatius* and/or *E. juvenalis*): *Compsilura concinnata*.

spp.: *Compsilura concinnata*.

"**Estigeme**" *acraea*, see *Estigmene acraea*

Estigmene (Lep.: Arctiidae)

acraea, see *E. acraea*.

acraea: *Argyrophylax* sp. probably, *Carcelia diacrisiae*, *C. reclinata*, *C. reclinata* probably, *Compsilura concinnata*, *Eucelatoria armigera*, *Exorista mella*, *E. mella* probably, *E. spp.*, *Gymnocarcelia ricinorum*, *G. ricinorum* probably, *G. sp.*, *Hubneria estigmenensis*, *Leschenaultia adusta*, *L. leucophrys*, *Lespesia aletiae*, *L. archippivora*, *L. sp.* near *cuculliae*, *Mericia spp.*, "*Phorocera*" spp., *Sisyropa eudryae*, *Thelaira americana*, *Winthemia datanae*, Tachinidae.

spp.: *Lespesia aletiae*, *L. archippivora*.

Ethmia (Lep.: Ethmiidae)

longimaculella: *Eufantina ethniae*.

"**Ethnia**" *longimaculella*, see *Ethmia longimaculella*

Etiella (Lep.: Pyralidae)

zinckenella: *Argyrophylax* spp., *Erynnia tortricis*.

Eubaphe (Lep.: Arctiidae)

aurantiaca rubicundaria: *Euphorocera claripennis*.

Eublaberus (Orth.: Blattidae)

posticus: *Calodexia venteris*.

Eubolina (Lep.: Noctuidae)

sp. probably: *Euphorocera claripennis*.

Eucaterva (Lep.: Geometridae)

variaria: *Hyphantrophaga hyphantriae*, Tachinidae.

"**Euchaetes**" *egle*, see *Euchaetias egle*

Euchaetias (Lep.: Arctiidae)

egle: *Bombyliopsis abrupta*, *Compsilura concinnata*, *Euexorista futilis*, *Euphorocera claripennis*, *Exorista mella*, *E. spp.*, *Hyphantrophaga euchaetiae*, *H. hyphantriae*, *Leschenaultia leucophrys*, *Lespesia archippivora*, *L. euchaetiae*, *L. frenchii*, *L. schizurae*,

Mericia spp., *Winthemia datanae*, *Zenillia libatrix*,
Tachinidae.

oregonensis: *Archytas californiae*.

“*Euchausia*” *argentata*, see *Halisidota argentata*

Eucheira (Lep.: Pieridae)

socialis: *Eucheirophaga lugubris*, *Eusisyropa virilis*, *Winthemia*
spp.

Euclea (Lep.: Limacodidae)

“*ceppus*,” see *E. cippus*.

cippus, see *E. cippus* or *E. delphinii*.

cippus: *Anaporia pristis*, *Eusisyropa blanda*.

delphinii: *Eusisyropa blanda*, *E. virilis*, Tachinidae.

Eucordylea canusella, see *Coleotechnites canusella*

Eucosma (Lep.: Olethreutidae)

rescissoriana: *Psilidopteryx psilocorsiphaga*, *P.* sp.

ridingsana: *Coloradomyia eucosmaphaga*.

scudderiana: *Lespesia* spp.

sp.: *Leskiomima tenera*.

Eucymatoge monticolans, see *Eupithecia monticolans*

Eudamus proteus, see *Urbanus proteus*

Eudamus tityrus, see *Epargyreus clarus*

Eudioptis hyalinata, see *Diaphania hyalinata*

Eudryas grata, see *Euthisanotia grata*

Eudryas unio, see *Euthisanotia unio*

Eudryas spp., see *Euthisanotia* spp.

Euetheola (Col.: Scarabaeidae)

rugiceps: *Eutrixa exilis*, *Megapariopsis opaca*, *Microphthalma*
disjuncta.

Eufitchia ribearia, see *Itame ribearia*

Eugonia j-album, see *Nymphalis vau-album j-album*

Eugonobapta (Lep.: Geometridae)

nivosaria: *Actia* sp. near *palloris*, *Compsilura concinnata*.

Eulepte (Lep.: Pyralidae)

concordalis: *Argyrophylax albincisa*.

concordalis probably: *Carcelia lagoae*.

Eulia mariana, see *Argyrotaenia mariana*

Eulia quadrifasciana, see *Argyrotaenia quadrifasciana*

Eulype (Lep.: Geometridae)

hastata: Tachinidae.

Eumarozia (Lep.: Olethreutidae)

malachitana: *Lixophaga variabilis*.

Euparthenos (Lep.: Noctuidae)

nubilis: Tachinidae.

Euphydryas (Lep.: Nymphalidae)*editha taylori*: *Siphosturmia confusa*.*phaeton*: *Compsilura concinnata*.**Eupithecia** (Lep.: Geometridae)*coagulata* probably: *Clausicella turmalis*.*gibsonata*: *Blondelia eufitchiae*.*luteata*: *Bessa harveyi*.*monticolans*: *Chaetogaedia monticola*.**Euproctis** (Lep.: Lymantriidae)"chrysorrhoea," see *E. chrysorrhoea*.*chrysorrhoea*: *Blepharipa scutellata*, *Carcelia amplexa*, *C. laxifrons*, *Compsilura concinnata*, *Euphorocera claripennis*, *Exorista mella*, *E. spp.*, *Leschenaultia leucophrys*, *Madremyia saundersii*, *Patelloa leucaniae*, *P. leucaniae* probably, *P. pachypyga*, *Phrynofrontina discalis*, *Townsendiellomyia nidicola*, *Zenillia libatrix*.*chrysorrhoea* probably: *Eusisyropa blanda*.**Eupseudosoma** (Lep.: Arctiidae)*involutum floridanum*: *Lespesia* sp. near *laniiferae*.**Eupsilia** (Lep.: Noctuidae)*devia*: *Mericia arcuata* probably, *Periscepsia helymus*.spp. (including *E. morrisoni* and/or *E. sidus*): *Cyrtophleba coquilletti*, Tachinidae.European corn borer, see *Ostrinia nubilalis*European crane fly, see *Tipula paludosa*European earwig, see *Forficula auricularia*European honeysuckle leafroller, see *Harpiteryx xylostella*"European" larch sawfly, see *Pristiphora erichsonii*European pine sawfly, see *Neodiprion sertifer*European pine shoot moth, see *Rhyacionia buoliana*"European" satin moth, see *Stilpnotia salicis*European spruce sawfly, see *Diprion hercyniae***Eurygaster** (Hem. (Heter.): Pentatomidae)spp.: *Gymnosoma fuliginosum*, *Phasia nigra*.**Eurymus eurytheme**, see *Colias eurytheme***Euryophthalmus cinctus californicus**, see *Largus cinctus californicus***Euryptychia saligneana**, see *Eucosma scudderiana***Euschistus** (Hem. (Heter.): Pentatomidae)*ictericus*: *Euthera tentatrix*.*ictericus*, *Mormidea pictiventris*, "Nezera" *viridula* and/or *Solubea pugnax*, see Pentatomid spp. (including *E. ictericus*, *M. pictiventris*, *Nezara viridula* and/or *Oebalus pugnax*).*impictiventris*: *Gymnosoma fuliginosum*.

servus: *Cylindromyia binotata*, *Euthera tentatrix*, *Gymnosoma fuliginosum*, *Trichopoda pennipes*.

tristigmus: *Euthera tentatrix*.

tristigmus var. *luridus*, see *E. tristigmus*.

variolarius: *Cylindromyia binotata*, *C. fumipennis*, *Euthera tentatrix*, *Gymnoclytia immaculata*, *G. occidua*, *Gymnosoma fuliginosum*, *Trichopoda pennipes*.

“*Eutheola*” *rugiceps*, see *Eutheola rugiceps*

Euthisanotia (Lep.: Noctuidae)

grata: *Chaetophlepsis eudryae*, *Compsilura concinnata*, *Sisyropa eudryae*, *Winthemia leucanae*, *W. leucanae* probably, Tachinidae.

unio: *Sisyropa eudryae*.

spp.: *Sisyropa eudryae*.

Euthochtha (Hem. (Heter.): Coreidae)

galeator: *Trichopoda pennipes*.

“*Euvanesia*” *antiopa*, see *Nymphalis antiopa*

Euvanessa antiopa, see *Nymphalis antiopa*

Euxoa (Lep.: Noctuidae)

auxiliaris: *Bonnetia comta*, *Euphorocera claripennis*, *Mericia* spp., *Peleteria* “*tessellata*,” *P.* “*tessellata*” probably, *P. texensis*, *Periscepsia cinerosa*, *P. helymus*, *P. laevigata*, Tachinidae.

messoria: *Aphria ocypterata*, *Bonnetia comta*, *Nowickia algens*.

messoria probably: *Bonnetia comta*, *Euphorocera edwardsii*.

ochrogaster: *Bonnetia comta*, *Gonia aldrichi*, *G. breviforceps*, *G. fuscicollis*, *G.* spp., *Periscepsia helymus*, *Winthemia quadripustulata*.

pallipennis: *Periscepsia rohweri*.

ridingsiana: *Peleteria texensis*.

scandens: Tachinidae.

tristicula: *Bonnetia comta*, *Gonia longiforceps*, *Periscepsia rohweri*.

vestigialis: *Lespesia archippivora*.

spp. (including *E. messoria* and/or *E. ochrogaster*): *Bonnetia comta*, *Euphorocera edwardsii*, *Gonia aldrichi* probably.

Evagora sp., see *Coleotechnites* sp.

Evergestis (Lep.: Pyralidae)

rimosalis: *Lespesia aletiae*.

straminalis: *Compsilura concinnata*, *Phryxe vulgaris*.

Evergreen bagworm, see *Thyridopteryx ephemeraeformis*

Evetria pasadenana, see *Rhyacionia pasadenana*

Evetria sp. probably, see *Rhyacionia* sp. probably

Evora (Lep.: Olethreutidae)

hemidesma: *Erynnia tortricis*, *Nemorilla pyste*, *Pseudo-perichaeta erecta*.

Exartema (Lep.: Olethreutidae)

sericoranum: *Nemorilla pyste*.

Exoteleia (Lep.: Gelechiidae)

pinifoliella: *Schizactia vitinervis*.

pinifoliella probably: *Schizactia vitinervis*.

Eyespotted bud moth, see *Spilonota ocellana*

Fall armyworm, see *Spodoptera frugiperda*

Fall cankerworm, see *Alsophila pometaria*

Fall webworm, see *Hyphantria cunea*

False chinch bug, see *Nysius ericae*

False hemlock looper, see *Nepytia canosaria*

Faronta (Lep.: Noctuidae)

diffusa: *Euphorocera claripennis*, *Lespesia frenchii*, L. spp.,
Periscepsia helymus, *Winthemia quadripustulata*,
Tachinidae.

Feltia (Lep.: Noctuidae)

annexa, see *F. subterranea*.

ducens: *Gonia fuscicollis*.

ducens-subgothica, see *F. ducens-F. subgothica* complex.

ducens-F. subgothica complex: *Euphorocera claripennis*,
Gonia frontosa, *Winthemia quadripustulata*.

gladiaria, see *Agrotis gladiaria*.

herilis: *Euphorocera claripennis*, *Winthemia quadripustulata*.

malefida, see *Agrotis malefida*.

subgothica: *Euphorocera claripennis*, *Triachora omissa*.

sp. near *subgothica*: *Lespesia archippivora*.

subterranea: *Bonnetia comta*, *Gonia crassicornis*, *G. longipulvilli*, *Lespesia archippivora*, *Spallanzania hebes*.

venerabilis, see *Agrotis venerabilis*.

Fentonia (Lep.: Notodontidae)

marthesia: *Compsilura concinnata*, *Sisyropa eudryae* probably.

Fiery hunter, see *Calosoma calidum*

Filament borer, see *Nematocampa filamentaria*

Filatima (Lep.: Gelechiidae)

monotaeniella: *Eusisyropa blanda* probably.

persicaeella: *Nemorilla pyste*, *Pseudomyothyria ancilla*.

Filbertworm, see *Melissopus latiferreanus*

Flax bollworm, see *Heliothis ononis*

Florida fern caterpillar, see *Callopistria floridensis*

Forage looper, see *Caenurgina erechtea*

- Forest tent caterpillar, see *Malacosoma disstria*
- Forest tent caterpillar and/or western tent caterpillar, see *Malacosoma* spp. (including *M. californicum pluviale* and/or *M. disstria*)
- Forficula (Dermaptera: Forficulidae)
auricularia: *Bigonicheta spinipennis*, *Ocytata pallipes*.
- Fruittree leafroller, see *Archips argyrospilus*
- Frumenta (Lep.: Gelechiidae)
nundinella: *Lixophaga mediocris*.
- Galeruca "bachardis," see *Trirhabda bacharidis*
- Galerucella *luteola*, see *Pyrrhalta luteola*
- Galerucella *xanthomelaena*, see *Pyrrhalta luteola*
- Galleria (Lep.: Pyralidae)
mellonella: *Ceromasia auricaudata*, *Lydella thompsoni*.
- Gelechia *confusella*, see *Filatima persicaeella*
- Gelechia *monotaeniella*, see *Filatima monotaeniella*
- Gelechiid webber(-s) (Lep.: Gelechiidae)
 Gelechiid webber(-s): *Eusisyropa blanda*.
- Geocoris (Hem. (Heter.): Lygaeidae)
bullatus: *Hyalomya aldrichii*.
punctipes: *Hyalomya aldrichii*.
uliginosus speculator: *Hyalomya aldrichii*.
- Geometridae (Lep.: Geometridae)
 Unidentified: *Aplomya helvina*, *Bessa harveyi*, *Compsilura concinnata*, *Eucelatoria armigera*, *Lespesia frenchii*, *Winthemia sinuata* probably, *Xanthoepalpus bicolor*.
- Geophilus (Geophilomorpha (Chilopoda): Geophilidae)
 sp. probably: *Exoristoides harrisi*.
- Gilpinia *frutetorum*, see *Diprion frutetorum*
- Gilpinia *hercyniae*, see *Diprion hercyniae*
- "Gipsy" moth, see *Porthetria dispar*
- Glaea (Lep.: Noctuidae)
 sp.: *Periscepsia* sp.
- Glassy cutworm, see *Crymodes devastator*
- Gloveria (Lep.: Lasiocampidae)
howardi: *Aplomya theclarum*, *Hyphantrophaga hyphantriae*.
- Gluphisia (Lep.: Notodontidae)
septentrionalis: *Eusisyropa virilis*.
 "septentrionis," see *G. septentrionalis*.
- Gnorimoschema (Lep.: Gelechiidae)
 spp.: *Actia* spp., *Frontiniella parancilla*.
- Golden buprestid, see *Buprestis aurulenta*
- Golden tortoise beetle, see *Metriona bicolor*
- "Gortina" *nitela*, see *Papaipema nebris*

- Gortyna immanis*, see *Hydroecia immanis*
Gortyna nitela, see *Papaipema nebris*
Gracillaria (Lep.: Gracillariidae)
negundella: *Aplomya caesar*.
Granulate cutworm, see *Feltia subterranea*
Grape cane gallmaker, see *Ampelogypter sesostris*
Grape flea beetle, see *Altica chalybea*
Grape leaffolder, see *Desmia funeralis*
Grapeleaf skeletonizer, see *Harrisina americana*
Grape plume moth, see *Pterophorus periscelidactylus*
Grapevine looper, see *Lygris diversilineata*
Graphiphora smithii, see *Amathes smithii*
Grapholitha (Lep.: Olethreutidae)
molesta: *Actia interrupta*, *Admontia degeeriodes*, *Dichae-
toneura leucoptera*, *Elephantocera greeni*, *Elodia
flavipalpis*, *E. subfasciata*, *E. tragica*, *Erynnia tortricis*,
Leskiella brevirostris probably, *Leskiomima tenera*,
Lespesia spp., *Lixophaga mediocris*, *L. parva*, *L.
plumbea*, *L. variabilis*, *Nemorilla pyste*, *Pseudoperi-
chaeta erecta*, *P. insidiosa*, *Strobliomyia tibialis*, Tach-
inidae.
Graptolitha antennata, see *Lithophane antennata*
Graptolitha disposita, see *Lithophane disposita*
Graptolitha disposita and/or *G. innominata*, see *Lithophane* sp.
(including *L. disposita* and/or *L. innominata*)
Graptolitha disposita, *G. innominata* and/or *G. petulca*, see *Litho-
phane* spp. (including *L. disposita*, *L. innominata* and/or *L.
petulca*)
Graptolitha innominata, see *Lithophane innominata*
Graptolitha sp., see *Lithophane* spp.
Grasshopper(-s) (Orth.: Acrididae)
Grasshopper(-s): *Acemya tibialis*, *A.* spp., *Ceracia dentata*, *C.
spp.*, *Cylindromyia "carolinae," Euphorocera clari-
pennis*, *Hemithrixion oestriforme*, *H. plankii*, *H. sp.*,
Leucostoma simplex, *Winthemia militaris*, Tachinidae.
Graybanded leafroller, see *Argyrotaenia mariana*
Great ash sphinx, see *Sphinx chersis*
Greater wax moth, see *Galleria mellonella*
Green cloverworm, see *Plathypena scabra*
Green fruitworm, see *Lithophane antennata*
Greenheaded spruce sawfly, see *Pikonema dimmockii*
Greenhouse leaf-tier, see *Udea rubigalis*
Green looper(-s) (Lep.: Unidentified Lepidoptera)
Green looper(-s): *Euphorocera tachinomoides*, *Voria ruralis*.

- Green stink bug, see *Acrosternum hilare*
- Greenstriped grasshopper, see *Chortophaga viridifasciata*
- Greenstriped mapleworm, see *Anisota rubicunda*
- Gretchena (Lep.: Olethreutidae)
bolliana: *Eusisyropa blanda*.
- Griselda (Lep.: Olethreutidae)
radicana probably: *Psaliapteryx nuda*.
- Gryllus (Orth.: Gryllidae)
abbreviatus, see *G. pennsylvanicus*.
assimilis, see *G. spp.*
pennsylvanicus: *Exoristoides johnsoni*.
spp.: *Euphasiopteryx ochracea*, *Exoristoides johnsoni*.
- Gusano cortador, see Cutworm(-s)
- Gusanos de maiz, see Lepidoptera, miscellaneous
- Gypsy moth, see *Porthetria dispar*
- Hadena apamiformis*, see *Apamea apamiformis*
- Hadena devastatrix*, see *Crymodes devastator*
- Hadena lignicolor*, see *Apamea lignicolor*
- Hadena turbulenta*, see *Phosphila turbulenta*
- Haemorrhagia diffinis*, see *Hemaris diffinis*
- Hag moth, see *Phobetron pithecium*
- "*Halesidota*" *argentata*, see *Halisidota argentata*
- "*Halesidota*" *spp.*, see *Halisidota spp.*
- "*Halisidota*" *maculata*, see *Halisidota maculata*
- "*Halisidota*" *tessellaris*, see *Halisidota tessellaris*
- Halisidota* (Lep.: Arctiidae)
argentata: *Anaporia pristis*, *Compsilura concinnata*, *Leschenaultia adusta*, *L. americana*, *L. halisidotae*, *L. spp.*,
"*Macquartia*" *spp.*, *Tachinomyia similis*, *Uramya halisidotae*.
caryae: *Bombyliopsis abrupta*, *Compsilura concinnata*,
Leschenaultia adusta, *L. halisidotae*, *L. spp.*, *Lespesia archippivora*, *L. frenchii*, "*Myiophasia atra*,"
Tachinidae.
edwardsi, see *Hemihyalea edwardsi*.
harrisii: *Blondelia hyphantriae*, *Compsilura concinnata*, *Exorista mella*, *Leschenaultia adusta*, *L. halisidotae*, *Lespesia aletiae*.
maculata: *Bombyliopsis abrupta*, *Compsilura concinnata*,
Leschenaultia adusta, *L. halisidotae*, *Lespesia aletiae*,
"*Phorocera*" *halisidotae*, Tachinidae.
tessellaris: *Blondelia eufitchiae*, *B. hyphantriae*, *Bombyliopsis abrupta*, *Carcelia yalensis*, *Compsilura concinnata*,
Euphorocera claripennis, *Exorista mella*, *Leschenaultia*

adusta, *L. halisidotae*, *Lespesia aletiae*, *L. euchaetiae*,
L. frenchii, *Winthemia quadripustulata*, "Zenillia"
spp., Tachinidae.

"tessellata," see *H. tessellaris*.

spp.: *Carcelia lagoae*, *Juriniopsis aurifrons*, *Leschenaultia*
adusta, *Lespesia aletiae*.

Haltica bimarginata, see *Altica bimarginata*

Haltica "bimarginata," see *Altica bimarginata*

Haltica probata, see *Altica probata*

Hamadryas antiopa, see *Nymphalis antiopa*

Hamadryas j-album, see *Nymphalis vau-album j-album*

Hamadryas milberti, see *Nymphalis milberti*

Haploa (Lep.: Arctiidae)

lecontei: "Zenillia" spp.

lecontei including form *militaris*, see *H. lecontei*.

Harlequin bug, see *Murgantia histrionica*

Harmologa fumiferana, see *Choristoneura fumiferana*

Harmonia pini, see *Vespamima pini*

Harmostes (Hem. (Heter.): Coreidae)

reflexulus: *Leucostoma acirostre*.

Harpiphorus tarsatus, see *Macremphytus tarsatus*

Harpipteryx (Lep.: Plutellidae)

xylostella: Tachinidae.

Harpyia cinerea, see *Cerura cinerea*

Harrisina (Lep.: Zygaenidae)

americana: *Euphorocera claripennis*, *E. floridensis*, *Sturmia*
harrisinae, *S. harrisinae* probably, *S. sp.* near *harrisinae*.

brillians: *Euphorocera sp.* near *tachinomoides*, "Phorocera"
spp., *Sturmia harrisinae*, *S. harrisinae* probably.

spp.: "Phorocera" spp., *Sturmia harrisinae* probably.

Hawaiian beet webworm, see *Hymenia recurvalis*

Hawk moth(-s) (Lep.: Sphingidae)

Hawk moth(-s): *Belvosia bicincta*.

Hedylepta (Lep.: Pyralidae)

accepta: *Chaetogaedia monticola*, *Eucelatoria armigera*,
Lespesia archippivora, Tachinidae.

blackburni: *Chaetogaedia monticola*, *Eucelatoria armigera*,
Lespesia archippivora.

continuatalis: *Chaetogaedia monticola*.

indicata: *Argyrophylax albincisa*, *Drino cubaecola*, *Eucelatoria*
spp., *Nemorilla pyste*.

Heliophila (Lep.: Noctuidae)

albilinea, see *Faronta diffusa*.

commoides, see *Leucania commoides*.

unipuncta, see *Pseudaletia unipuncta*.

"*unipunctata*," see *Pseudaletia unipuncta*.

spp.: *Archytas marmoratus*, *A. spp.* (including *A. apicifer* and/or *A. californiae*), *Peleteria texensis*.

Heliothis (Lep.: Noctuidae)

armigera, see *H. zea*.

armigera, *Laphygma frugiperda*, *Prodenia dolichos*, *P. eridania* and/or *P. ornithogalli*, see Noctuid spp. (including *H. zea*, *P. dolichos*, *Spodoptera eridania*, *S. frugiperda* and/or *S. ornithogalli*).

hawaiiensis: *Eucelatoria armigera*, *Lespesia archippivora*.

obscura, see *H. zea*.

obsoleta, see *H. zea*.

ononis: *Athanatus californicus*.

phloxiphaga: *Eucelatoria armigera*.

virescens: *Carcelia illota*, *Chaetogaedia monticola*, *Eucelatoria armigera*, *Euphorocera claripennis*, *Lespesia spp.*, *Metaplagia occidentalis*, *Palexorista laxa*.

zea: *Archytas marmoratus*, *Carcelia illota*, *Eucelatoria armigera*, *E. spp.*, *Euphorocera claripennis*, *E. floridensis*, *E. tachinomoides*, *Gonia spp.*, *Gymnochaetopsis fulvicauda*, *Hyphantrophaga hyphantriae*, *Lespesia aletiae*, *L. archippivora*, *L. frenchii*, *L. spp.*, *Metaplagia occidentalis*, *Nemorilla pyste*, *Voria aurifrons*, *Winthemia quadripustulata*, *W. rufopicta*, *W. spp.*, Tachinidae.

spp. (including *H. hawaiiensis* and/or *H. zea*): *Eucelatoria armigera*, *Lespesia archippivora*.

spp.: *Archytas marmoratus*, *Chaetogaedia sp. near analis*, *Eucelatoria armigera*, *Euphorocera omissa*, *E. tachinomoides*, *Lespesia aletiae*, *L. archippivora*, *Plagiomima cognata*, *P. spinosula* probably, *Winthemia rufopicta*, *W. spp.* probably, Tachinidae.

Hellula (Lep.: Pyralidae)

phidilealis: *Nemorilla pyste*.

rogatalis: *Nemorilla pyste*, *Stomatomyia sp.*

undalis, see *H. rogatalis*.

Hemaris (Lep.: Sphingidae)

diffinis: *Winthemia quadripustulata*, *W. rufopicta*.

thysbe: *Compsilura concinnata*, Gen. and sp. near *Thysanomyia*, Tachinidae.

Hemerocampa (Lep.: Lymantriidae)

leucostigma: *Bessa selecta*, *Carcelia amplexa*, *C. perplexa*, *C. yalensis*, *Compsilura concinnata*, *Drino inconspicua*,

Euphorocera claripennis, *E. edwardsii*, "Exorista" *lobeliae*, *E. mella*, *E. spp.*, *Leschenaultia spp.*, *Lespesia aletiae*, *L. frenchii*, *L. spp.*, *Patelloa leucaniae*, "Phorocera" spp., *Sisyropa spp.*, *Winthemia datanae*, *W. quadripustulata*, Tachinidae.

oslari, see *H. leucostigma*.

pseudotsugata: *Exorista mella*, Tachinidae.

vetusta: *Exorista mella*, *Lespesia frenchii*, *Patelloa fuscimacula*, *Peleteria texensis*, *Tachinomyia panaetius*, *T. similis*.

"Hemerocampe" *leucostigma*, see *Hemerocampa leucostigma*

Hemerophila pariana, see *Anthophila pariana*

Hemichroa (Hym.: Tenthredinidae)

crocea: *Bessa harveyi*, *B. selecta*, *Drino bohemica*.

Hemihyalea (Lep.: Arctiidae)

edwardsi: *Leschenaultia adusta*.

Hemileuca (Lep.: Saturniidae)

artemis, see *H. nevadensis*.

electra: *Euphorocera claripennis*.

lucina: *Leschenaultia fulvipes*.

maia: *Argyrophylax spp.*, *Compsilura concinnata*, *Drino incompta*, *Leschenaultia fulvipes*, *Lespesia aletiae*, *L. spp.*

maia nevadensis, see *H. nevadensis*.

nevadensis: *Euphorocera claripennis*, "Sturmia" sp. probably.

oliviae: *Compsilura concinnata*, *Euphorocera claripennis*, *Exorista mella*.

spp. (including *H. lucina* and/or *H. maia*): *Belvosia bifasciata*, *Carcelia malacosomae*, *Compsilura concinnata*, *Euphorocera claripennis*, *Leschenaultia fulvipes*, *Phryxe pecosensis*, *P. vulgaris*, *Winthemia spp.*, Tachinidae.

sp.: *Belvosia bifasciata*.

sp. probably: *Belvosia bifasciata*.

Hemiptera (Heteroptera) (Hem. (Heter.): Unidentified Hemiptera (Heteroptera))

Miscellaneous: *Hyalomya aldrichii*.

Hemlock looper, see *Lambdina fiscellaria*

Hemlock sawfly, see *Neodiprion tsugae*

Heodes thoe, see *Lycaena thoe*

Hepialus (Lep.: Hepialidae)

pharus: Tachinidae.

Heptagrotis (Lep.: Noctuidae)

phyllophora: *Phryxe pecosensis*.

Heraclides cresphontes, see *Papilio cresphontes*

Herpetogramma (Lep.: Pyralidae)*bipunctalis*: *Nemorilla pyste*.*licarsisalis*: *Eucelatoria armigera*.**Herse cingulata**, see *Agrius cingulatus***Herse cingulata** probably, see *Agrius cingulatus* probably**Herse convolvuli**, see *Agrius convolvuli***Hesperia tityrus**, see *Epargyreus clarus***Hesperiid** (Lep.: HesperIIDae)spp. (including *Cymaenes tripunctus*, *Nyctelius nyctelius*, and/or *Panoquina sylvicola*): Tachinidae.**Hesperiidae** (Lep.: HesperIIDae)Unidentified: *Belvosia ciliata*, *Compsilura concinnata*.**Heterocampa** (Lep.: Notodontidae)*bilineata*: *Sisyropa* spp.*biundata*: *Lespesia schizurae*.*guttivitta*: *Actia* spp., *Blondelia hyphantriae*, *Compsilura concinnata*, *Patelloa leucaniae*, *Spathimeigenia hylotomae* probably, *Winthemia quadripustulata*, *W. rufopicta* probably, Tachinidae.*manteo*: *Lespesia aletiae*.*marthesia*, see *Fentonia marthesia*.*umbrata*: *Compsilura concinnata*, *Lespesia* spp.sp.: *Lespesia schizurae*.**Hibernia tiliaria**, see *Erannis tiliaria***Hickory horned devil**, see *Citheronia regalis***Hickorynut curculio**, see *Conotrachelus affinis***Hickory plant bug**, see *Lygocoris caryae***Hickory shuckworm**, see *Laspeyresia caryana***Hickory tussock moth**, see *Halisidota caryae***High Plains grasshopper**, see *Dissosteira longipennis***Hog moth(-s)** (Lep.: Unidentified Lepidoptera)Hog moth(-s): "*Phorocera*" spp.**Holcostethus** (Hem. (Heter.): Pentatomidae)*limbolarius*: *Alophorella pulverea*.**Homoeosoma** (Lep.: Pyralidae)*electellum*: *Clausicella floridensis*, *C. neomexicana*, *C. opaca*, *Erynnia tortricis*, *Leskiomima tenera*, *Lixophaga variabilis*."Homoesoma" *electellum*, see *Homoeosoma electellum***Hop looper**, see *Hypena humuli***Horned passalus**, see *Popilius disjunctus***Horned squash bug**, see *Anasa armigera***Hornworm**, see *Manduca sexta***Horse fly (flies)** (Dipt.: Tabanidae)Horse fly (flies): *Carinosillus tabanivorus*.

“*Hulstea*” *undulatella*, see *Hulstia undulatella*

Hulstia (Lep.: Pyralidae)

undulatella: *Nemorilla pyste*, *Stomatomyia parvipalpis*.

Hyalophora (Lep.: Saturniidae)

calleta: *Belvosia nigrifrons*, *Lespesia frenchii*.

cecropia, see *Platysamia cecropia*.

columbia: *Lespesia frenchii*.

columbia nokomis, see *H. nokomis*.

euryalus, see *Platysamia euryalus*.

gloveri: *Lespesia ciliata*.

nokomis: *Lespesia ciliata*.

promethea, see *Callosamia promethea*.

rubra, see *Platysamia euryalus*.

sp.: *Lespesia ciliata*.

Hybomitra (Dipt.: Tabanidae)

epistates: *Carinosillus tabanivorus*.

lasiophthalma: Tachinidae.

metabola: *Carinosillus tabanivorus*.

nuda: *Carinosillus tabanivorus*.

trepida: *Carinosillus tabanivorus*.

trispila sodalis probably: *Carinosillus novaeangliae*.

spp.: *Carinosillus tabanivorus*.

“*Hydelepta*” *indicata*, see *Hedylepta indicata*

Hydria (Lep.: Geometridae)

undulata, see *H. undulata* or *H. spp.* (including *H. prunivorata* and/or *H. undulata*).

undulata: *Drino inconspicua*, *Eusisyropa blanda*.

spp. (including *H. prunivorata* and/or *H. undulata*): *Chryso-exorista marginata*, *Compsilura concinnata*.

Hydroecia (Lep.: Noctuidae)

immanis: *Euphorocera claripennis*, *Lespesia frenchii*, *Lydella radialis*.

nitela, see *Papaipema nebris*.

Hylephila (Lep.: Hesperidae)

phyleus: *Stomatomyia parvipalpis*.

Hyles (Lep.: Sphingidae)

gallii: *Compsilura concinnata*.

lineata: *Compsilura concinnata*, *Drino incompta*, “*Tachina*” spp., *Winthemia deilephila*, *W. quadripustulata*, Tachinidae.

Hylesia (Lep.: Saturniidae)

alinda: *Belvosia recticornis*.

darlingi: *Belvosia recticornis*.

sp.: *Belvosia recticornis*.

- Hyloicus chersis*, see *Sphinx chersis*
Hyloicus drupiferarum, see *Sphinx drupiferarum*
Hyloicus gordius, see *Sphinx gordius*
Hyloicus kalmiae, see *Sphinx kalmiae*
Hyloicus luscitiosa, see *Sphinx luscitiosa*
Hylotoma "hemeralis," see *Arge humeralis*
Hylotoma humeralis, see *Arge humeralis*
Hylotoma macleayi, see *Arge clavicornis* complex
Hymenia (Lep.: Pyralidae)
 fascialis, see *H. recurvalis*.
 perspectalis: *Argyrophylox albincisa*.
 recurvalis: *Argyrophylox albincisa*, *Chaetogaedia monticola*,
 Eucelatoria armigera, *E. spp.*, *Nemorilla pyste*.
Hypena (Lep.: Noctuidae)
 humuli: *Blondelia eufitchiae*, *Ceromasia aurifrons*, *Eusisyropa*
 blanda, *E. boarmiae*.
Hypera (Col.: Curculionidae)
 nigrirostris: *Lixophaga* sp. near *variabilis*.
 postica: *Hyalomyodes triangulifer*, *Madremyia saundersii*.
 punctata: *Hyalomyodes triangulifer*.
Hyperaeschra (Lep.: Notodontidae)
 stragula: *Winthemia datanae*, Tachinidae.
Hyperchiria io, see *Automeris io*
Hyphantria (Lep.: Arctiidae)
 cunea: *Archytas aterrimus*, *Blondelia eufitchiae*, *B. hyphan-*
 triae, *Bombyliopsis abrupta*, *Carcelia malacosomae*, *C.*
 protuberans, *Ceromasia auricaudata*, *Compsilura*
 concinata, *Euphorocera claripennis*, *E. floridensis*,
 Eusisyropa blanda, *E. boarmiae*, *E. virilis*, *Hyphantro-*
 phaga hyphantriae, *Lespesia aletiae*, *L. frenchii*,
 Mericia ampelus, *M. arcuata*, *M. johnsoni*, *M. spp.*, *M.*
 spp. probably, *Winthemia spp.*, Tachinidae.
 textor: *Blondelia eufitchiae*, *B. hyphantriae*, *Compsilura*
 concinata, *Eusisyropa blanda*, *E. boarmiae*, *Exorista*
 mella, *Hyphantrophaga hyphantriae*, *Mericia ampelus*,
 "*Tachina*" spp.
 spp.: *Compsilura concinnata*, *Lespesia aletiae*, *Mericia ampelus*,
 M. spp. probably.
Hyphoraia parthenos, see *Parasemia parthenos*
Hyponomeuta multipunctella, see *Yponomeuta multipunctella*
Hyppa (Lep.: Noctuidae)
 xylinoides: *Euexorista inutilis*.
Hypsoropha (Lep.: Noctuidae)
 hormos: *Lespesia frenchii*, *L. spp.*

Ianassa lignicolor, see *Dicentria lignicolor*

Ichthyura (Lep.: Notodontidae)

albosigma: *Eusisyropa virilis*.

inclusa: *Compsilura concinnata*, *Euphorocera claripennis*, *E. floridensis*, *Exorista mella*, *Lespesia frenchii*, *L. melalophae*, *L. spp.*, *Winthemia spp.*, *Zenillia libatrix*, Tachinidae.

strigosa: *Eusisyropa virilis*.

spp. (including *I. apicalis* or *I. inclusa*): *Euphorocera floridensis*, *Lespesia spp.*

"*Ichthyura*" *inclusa*, see *Ichthyura inclusa*

Illice nexa, see *Cisthene nexa*

Imperial moth, see *Eacles imperialis*

Imported cabbageworm, see *Pieris rapae*

Imported currantworm, see *Nematus ribesii*

Indian meal moth, see *Plodia interpunctella*

Introduced pine sawfly, see *Diprion similis*

Io moth, see *Automeris io*

Iris borer, see *Macronoctua onusta*

Isa inornata, see *Sisyrosea textula*

Isia (Lep.: Arctiidae)

isabella: *Euexorista futilis*, *Exorista mella*, *Hubneria estigmenensis*, *Mystacella spp.*, *Parachaeta fusca*, *P. spp.*, *Thelaira spp.*, *Winthemia datanae* probably, Tachinidae.

Isophrictis (Lep.: Gelechiidae)

similiella denotata: "*Tachinophyto*" spp.

Isturgia (Lep.: Geometridae)

truncataria: *Eusisyropa blanda*, *E. boarmiae*.

Itame (Lep.: Geometridae)

flavicaria: *Blondelia eufitchiae*.

fulvaria probably: *Chaetophlepsis semiothisae*.

pustularia: *Tachinomyia nigricans*, Tachinidae.

quadrilinearia: *Blondelia eufitchiae*.

ribearia: *Blondelia eufitchiae*, *Compsilura concinnata*, Tachinidae.

sulphurea: *Euphorocera claripennis*.

Jack pine budworm, see *Choristoneura pinus*

Jack pine sawfly, see *Neodiprion pratti banksianae*

Japanese beetle, see *Popillia japonica*

Jasoniades glaucus, see *Papilio glaucus*

June beetles, see June beetle(-s) or *Phyllophaga spp.*

June beetle(-s) (Col.: Scarabaeidae)

June beetle(-s): *Eutrixa exilis*, *Microphthalma disjuncta*.

Lachnosterna (Col.: Scarabaeidae)

allied genera of, see Scarabaeid spp. (allied genera of *Phyllophaga* spp.).

arcuata, see *Phyllophaga fervida*.

crassissima, see *Phyllophaga crassissima*.

crenaticollis probably: *Cryptomeigenia aurifacies*.

"*crenatocollis*" probably, see *L. crenaticollis* probably.

crenulata, see *Phyllophaga crenulata*.

fusca, see *Phyllophaga fusca*.

futilis, see *Phyllophaga futilis*.

hogardi, see *Chemarachis hogardi*.

"*implicata*," see *Phyllophaga implicita*.

inversa, see *Phyllophaga inversa*.

portoricensis, see *Phyllophaga portoricensis*.

rubiginosa, see *Phyllophaga rubiginosa*.

rugosa, see *Phyllophaga rugosa*.

vandinei, see *Chemarachis vandinei*.

or other "scarbaeid" beetles, see Scarabaeid spp. (including *Phyllophaga* spp. or other Scarabaeidae).

spp., see *Phyllophaga crenulata* or *Phyllophaga* spp.

Lacinipolia (Lep.: Noctuidae)

renigera: *Chaetogaedia analis*.

Lagoa (Lep.: Megalopygidae)

crispata, see *Megalopyge crispata*.

opercularis, see *Megalopyge opercularis*.

pyxidifera, see *Megalopyge pyxidifera*.

spp.: *Carcelia lagoae*, *Euphorocera claripennis*.

Lambdina (Lep.: Geometridae)

fiscellaria, see *L. fiscellaria* or *L. fiscellaria fiscellaria*.

fiscellaria: *Blondelia eufitchiae*, *Eusisyropa blanda*, *Madremyia saundersii*, *Winthemia occidentis*, *W. spp.*, Tachinidae.

fiscellaria fiscellaria: *Blondelia eufitchiae*, *Chaetophlepsis semi-othisae*, *Compsilura concinnata*, *Madremyia saundersii*, *Phryxe pecosensis*, *Winthemia occidentis*.

fiscellaria laeta probably: *Chrysoexorista dawsoni*.

fiscellaria lugubrosa: *Winthemia occidentis*, *W. venusta* probably.

fiscellaria "somniara", see *L. fiscellaria somniaria*.

fiscellaria somniaria: *Madremyia saundersii*, *Winthemia occidentis*.

"**Lamlidina**" *fiscellaria*, see *Lambdina fiscellaria fiscellaria*

Lampides: (Lep.: Lycaenidae)

boeticus: *Lespesia archippivora*.

Lamprosema ebullialis, see "*Blepharomastix*" *ebullialis*

Lamprosema indicata, see *Hedylepta indicata*

Laniifera (Lep.: Pyralidae)

cyclades: *Lespesia laniiferae*, L. spp., "Phorocera" spp.

"Laphigma" *frugiperda*, see *Spodoptera frugiperda*

Laphygma (Lep.: Noctuidae)

exempta, see *Spodoptera exempta*.

exigua, see *Spodoptera exigua*.

flavimaculata, see *Spodoptera exigua*.

frugiperda, see *Spodoptera frugiperda*.

[sp.], see *Spodoptera frugiperda*.

spp.: *Archytas marmoratus*.

"Laphygma" *frugiperda*, see *Spodoptera frugiperda*

Lappet moth, see *Epicnaptera americana*

Larch sawfly, see *Pristiphora erichsonii*

Large aspen tortrix, see *Choristoneura conflictana*

Large milkweed bug, see *Oncopeltus fasciatus*

Larger canna leafroller, see *Calpodes ethlius*

Larger elm leaf beetle, see *Monocesta coryli*

Largus (Hem. (Heter.): Largidae)

cinctus californicus: *Trichopoda pennipes*.

sp.: *Trichopoda pennipes*.

Lascoria (Lep.: Noctuidae)

ambigualis: *Periscepsia laevigata*.

Laspeyresia (Lep.: Olethreutidae)

caryana: *Leskiella brevirostris*, *Lixophaga mediocris*, *L. variabilis*, "Myiophasia atra."

molesta, see *Grapholitha molesta*.

nigricana: *Aplomya caesar*.

pomonella: *Chrysoexorista dawsoni*, *C. marginata*, *C. ochracea*, *Erynnia tortricis*, *Leskiella brevirostris*, *Leskiomima tenera*, *Lixophaga jennei*, *L. orbitalis*, *L. variabilis*, *L.* spp., *Nilea carpocapsae*, *N. noctuiformis*, "Tachinophyto" spp.

spp. and/or *Dioryctria* spp. probably, see Olethreutid and/or pyralid spp. (including *Laspeyresia* spp. and/or *Dioryctria* spp. probably).

Lawn armyworm, see *Spodoptera mauritia*

"Laxotaenia" *rosaceana*, see *Choristoneura rosaceana*

Leaf casebearer, see *Acrobasis juglandis*

Leaf crumpler, see *Acrobasis indigenella*

Leaf-folder caterpillar(-s) (Lep.: Unidentified Lepidoptera)

Leaf-folder caterpillar(-s): *Myobiopsis flavipennis*.

Leaf-footed bug, see *Leptoglossus phyllopus*

Leaf-footed plant bugs, see *Leptoglossus* spp.

Leaf roller(-s) (Lep.: Tortricidae)

Leaf roller(-s): *Dejeaniopalpus texensis*, *Periscepsia* spp.,
Pseudoperichaeta erecta.

Leaf tier(-s) (Lep.: Unidentified Lepidoptera)

Leaf tier(-s): *Psalidopteryx slossonae*.

Lema (Col.: Chrysomelidae)

nigrovittata: *Paralipse infernalis*.

sexnotata: *Paralipse infernalis*.

trilineata, see *L. trilineata trilineata*.

trilineata californica: *Paralipse infernalis*.

trilineata trilineata: *Paralipse infernalis*, *Patelloa* sp. near
leucaniae.

sp.: *Paralipse infernalis*.

Lemonias *harrisii*, see *Chlosyne harrisii*Lemonias *taylori*, see *Euphydryas editha taylori*

Lepidoptera (Lep.: Unidentified Lepidoptera)

Miscellaneous: *Aplomya theclarum*, *Archytas marmoratus*, *A.*
spp. (including *A. apicifer* and/or *A. californiae*),
Belvosia argentifrons, *B. bifasciata*, *B. ferruginosa*, *B.*
townsendi, *Carcelia reclinata*, *C. yalensis*, *Ceromasia*
aurifrons, *Chaetogaedia monticola*, *Compsilura concin-*
nata, *Deopalpus parksi*, *Erynnia tortricis*, *Euexorista*
futilis, *Euphorocera claripennis*, *Eusisyropa blanda*,
Exorista spp. probably, *Leschenaultia* spp., *Lespesia*
aletiae, *L. archippivora*, *L. dimmocki*, *L. frenchii*,
Lixophaga diatraeae, *L. variabilis*, *Lydella radialis*,
Madremyia saundersii, *Mericia ampelus*, *Mochlosoma*
mexicanum, *Nemorilla pyste*, *Paradejeania rutilioides*
nigrescens, *Patelloa leucaniae*, *Phryxe pecosensis*,
Pseudomyothyria spp., *P.* spp. probably, *Spathi-*
meigenia spinigera, "Tachina" spp., *Winthemia quad-*
ripustulata, Tachinidae.

Miscellaneous, Lepidoptera probably: *Peleteria thomsoni*.

Leptinotarsa (Col.: Chrysomelidae)

decemlineata: *Adoryphorophaga aberrans*, *A.* spp. and/or *Dory-*
phorophaga spp., *Doryphorophaga australis*, *D. dory-*
phorae, *D. macella*, *D.* spp., *Winthemia* spp. (including
W. quadripustulata or allied sp.: *W. datanae*, *W. sinuata*
or *W. vesiculata*), Tachinidae.

"decimlineata," see *L. decemlineata*.

defecta: *Doryphorophaga australis*.

10-lineata, see *L. decemlineata*.

Leptoglossus (Hem. (Heter.): Coreidae)

oppositus: *Trichopoda pennipes*.

phyllopus: *Trichopoda pennipes*, *T. pennipes* complex, *T. plumipes*.

spp.: *Trichopoda pennipes*, *T. spp.*

Leptopterna (Hem. (Heter.): Miridae)

dolabrata: "*Phorantha occidentis*."

Lerodea (Lep.: Hesperidae)

eufala: *Euphorocera tachinomoides*.

tripuncta, see *Cymaenes tripunctus*.

tripuncta tripuncta, see *Cymaenes tripunctus*.

tripuncta, *Prenes nero sylvicola* and/or *P. nyctelius coscinia*, see Hesperiid spp. (including *Cymaenes tripunctus*, *Nyctelius nyctelius* and/or *Panoquina sylvicola*).

Lespedeza webworm, see *Tetralopha scortealis*

Lesser clover leaf weevil, see *Hypera nigrirostris*

Lesser cornstalk borer, see *Elasmopalpus lignosellus*

Lesser migratory locust, red-legged locust and/or two-striped locust, see Locust(-s)

Leucania (Lep.: Noctuidae)

albilinea, see *Faronta diffusa*.

cholica: *Archytas cirphis*, *Eucelatoria armigera*.

commoides: *Periscepsia helymus*.

extranea, see *Pseudaletia unipuncta*.

inconspicua: *Eucelatoria armigera*.

juncicola: *Archytas* spp. (including *A. apicifer* and/or *A. californiae*).

latiuscula: *Archytas cirphis*, *A. marmoratus*, *Eucelatoria armigera*, *E. australis*, *Peleteria texensis*.

latiuscula probably: *Winthemia quadripustulata*.

microgonia: *Peleteria texensis*.

pyrrhias, see *Pseudaletia macrosaris*.

unipuncta, see *Pseudaletia unipuncta*.

"*unipunctata*," see *Pseudaletia unipuncta*.

sp.: *Triachora luteola*.

Leucarctia "*acraca*," see *Estigmene acraea*

Leucarctia *acraea*, see *Estigmene acraea*

"**Leucaria** *unipunctata*," see *Pseudaletia unipuncta*

Lexis *bicolor*, see *Tigrioides bicolor*

Ligyris (Col.: Scarabaeidae)

gibbosus: *Eutrixia exilis*.

Limabean pod borer, see *Etiella zinckenella*

Limacodes (Lep.: Limacodidae)

spp.: *Anaporia limacodis*.

sp. probably: Tachinidae.

Limenitis (Lep.: Nymphalidae)

archippus: *Compsilura concinnata*, Tachinidae.

astyanax: *Compsilura concinnata*.

disippus, see *L. archippus*.

Lina interrupta, see *Chrysomela interrupta*

Lina scripta, see *Chrysomela scripta*

Linden looper, see *Erannis tiliaria*

Lined stalk borer, see *Oligia fractilinea*

Liorhyssus (Hem. (Heter.): Rhopalidae)

hyalinus: *Leucostoma aterrimum*, *L. simplex*, Tachinidae.

Liparidae, see Lymantriidae, unidentified

Listroderes (Col.: Curculionidae)

costirostris obliquus: *Epiplagiops littoralis*, *Hyalomyodes triangulifer*.

obliquus, see *L. costirostris obliquus*.

Lithacodes (Lep.: Limacodidae)

fasciola: *Anaporia limacodis*.

Lithophane (Lep.: Noctuidae)

antennata: *Blondelia hyphantriae*, *Compsilura concinnata*, *Gonia* spp., *Mericia* spp., *Omotoma fumiferanae*, *Phryxe pecosensis*, *P. vulgaris*, *Tachinomyia acosta*, *T. nigricans*, *T. sp.* near *panaetius*, *T. spp.*, *Winthemia quadripustulata*, *W. rufopicta*, Tachinidae.

disposita: *Tachinomyia acosta*.

innominata: *Periscepsia helymus*, *Tachinomyia acosta*.

spp. (including *L. disposita* and/or *L. innominata*): *Carcelia formosa*.

spp. (including *L. disposita*, *L. innominata* and/or *L. petulca*): *Blepharipa scutellata*, *Blondelia hyphantriae*, *Compsilura concinnata*, *Epalpus signifer*, *Euphorocera claripennis*, *Mericia* spp., *Omotoma fumiferanae*, *Periscepsia helymus*, *Tachinomyia* spp., *Winthemia rufopicta*, Tachinidae.

spp.: *Blepharipa scutellata*, *Compsilura concinnata*, *Periscepsia laevigata*, *Tachinomyia acosta*.

Litoprosopus (Lep.: Noctuidae)

sp.: Gen. and sp. near *Chaetogaedia*.

Lixus (Col.: Curculionidae)

"*scrobicollis*," see *L. scrobicollis*.

scrobicollis: *Lixophaga parva*, *L. variabilis*.

Locust leafroller, see *Nephopteryx subcaesiella*

Locust(-s) (Orth.: Acrididae)

Locust(-s): *Lespesia* spp., Tachinidae.

Looper(-s) (Lep.: Unidentified Lepidoptera)

- Looper(-s): *Madremyia saundersii*.
- Lophyrus abbotii*, see *Neodiprion abbotii*
- Lophyrus "abbotii,"* see *Neodiprion abbotii*
- Lophyrus abietis*, see *Neodiprion abietis*
- Lophyrus lecontei*, see *Neodiprion lecontei*
- Lophyrus* spp., see Diprionid spp. (including *Diprion* spp. and/or *Neodiprion* spp.)
- Lotisma* (Lep.: Glyphipterygidae)
trigonana: *Elfia amplicornis*.
- Loxostege* (Lep.: Pyralidae)
commixtalis: *Aplomya caesar*, *A. trisetosa*, *Lespesia archippivora*, *Phryxe vulgaris*, *P.* sp.
mancalis: *Blondelia polita*, *Oswaldia albifacies*, *Pseudochaeta argentifrons*.
obliteralis, see *Ostrinia obliteralis*.
similalis: *Eusisyropa blanda*, *E. boarmiae*, *Hyphantrophaga hyphantriae*, *Lespesia archippivora*, *Lixophaga variabilis*, *Nemorilla pyste*, *Patelloa leucaniae*, "Phorocera parva," "P." spp., *Pseudoperichaeta erecta*, *Stomatomyia parvipalpis*, *Winthemia quadripustulata*, Tachinidae.
sticticalis: *Aplomya caesar*, *Euphorocera omissa*, *Lespesia archippivora*, *L. ciliata*, *Stomatomyia parvipalpis*, Tachinidae.
 "sticticolis," see *L. sticticalis*.
- Loxotaenia rosaceana*, see *Choristoneura rosaceana*
- Luna moth, see *Actias luna*
- Luperina* (Lep.: Noctuidae)
stipata: *Lydella radialis*, *Winthemia rufopicta*, *W. sinuata*.
- "Luperodes" (Col.: Chrysomelidae)
 spp.: *Medina barbata*.
- Lycaeides* (Lep.: Lycaenidae)
melissa: *Aplomya theclarum*.
- Lycaena* (Lep.: Lycaenidae)
boeticus, see *Lampides boeticus*.
dione, see *L. xanthoides dione*.
exilis, see *Brephidium exilis*.
fulla, see *Plebejus icarioides*.
melissa, see *Lycaeides melissa*.
pseudargiolus, see *Celastrina argiolus pseudargiolus*.
thoe: *Aplomya theclarum*.
xanthoides: *Aplomya theclarum*.
xanthoides dione: *Aplomya theclarum*.
 spp.: *Aplomya theclarum*.

- Lycaenidae (Lep.: Lycaenidae)
 Unidentified: *Aplomya theclarum*.
- Lycaenopsis *argiolus pseudargiolus*, see *Celastrina argiolus pseudargiolus*
- Lycaenopsis *pseudargiolus*, see *Celastrina argiolus pseudargiolus*
- Lycia (Lep.: Geometridae)
cognataria, see *Biston cognataria*.
ursaria: *Ceromasia* spp.
- Lycophotia *margaritosa*, see *Peridroma saucia*
- Lycophotia *saucia*, see *Peridroma saucia*
- "Lygaeonematis *erichsonii*," see *Pristiphora erichsonii*
- Lygaeonematus *erichsonii*, see *Pristiphora erichsonii*
- Lygaeus (Hem. (Heter.): Lygaeidae)
kalmii: *Leucostoma* spp.
- Lygocoris (Hem. (Heter.): Miridae)
caryae: *Alophorella fumosa*.
caryae probably: *Alophorella pulverea*.
- Lygranthoecia *brevis*, see *Schinia brevis*
- Lygris (Lep.: Geometridae)
diversilineata: *Hyphantrophaga hyphantriae*.
- Lygus (Hem. (Heter.): Miridae)
caryae, see *Lygocoris caryae*.
caryae probably, see *Lygocoris caryae* probably.
hesperus probably: *Alophorella aeneoventris*.
lineolaris: *Alophorella aeneoventris*, *A. fumosa*, *A. pulverea*, *A.*
 spp.
pratensis: *Alophorella opaca*.
- Lymantria *dispar*, see *Porthetria dispar*
- Lymantriidae (Lep.: Lymantriidae)
 Unidentified: *Compsilura concinnata*.
- Lymire (Lep.: Ctenuchidae)
edwardsii: *Euphorocera claripennis*, *E. floridensis*, *Lespesia aletiae* probably.
- Macaria *punctolineata*, see *Semiothisa punctolineata*
- Macremphytus (Hym.: Tenthredinidae)
intermedius: *Guerinia dydas*.
tarsatus: *Guerinia dydas*.
testaceus: *Guerinia dydas*.
varianus: *Bessa harveyi*, *Exorista mella*, *Guerinia dydas*.
 spp.: *Bessa harveyi*.
- "Macrephytus" *intermedius*, see *Macremphytus intermedius*

Macrobotys (Lep.: Pyralidae)*aeglealis*: *Myobiopsis depilis*.*pertextalis*: *Leskiomima tenera*, *Pseudochaeta siminina*, *P. erecta*, Tachinidae.**Macronoctua** (Lep.: Noctuidae)*onusta*: *Lydella radialis*."Macronoctus" *onusta*, see *Macronoctua onusta**Macrosila cingulata*, see *Agrius cingulatus***Macrosiphum** (Hem. (Hom.): Aphididae)*avenae* probably: *Beskia aelops*.*granaria* probably, see *M. avenae* probably.**Magisicada** (Hem. (Hom.): Cicadidae)*septendecim*: "Tachina" spp., Tachinidae.**Malacosoma** (Lep.: Lasiocampidae)*americanum*: *Archytas aterrimus*, *A. lateralis*, *Blepharipa scutellata*, *Carcelia laxifrons*, *C. malacosomae*, *C. malacosomae* probably, *C. protuberans*, *Cartocometes io*, *Compsilura concinnata*, *Cyzenis albicans*, *Diplostichus lophyri*, *Drino inconspicua*, *Euexorista futilis*, *Euphorocera claripennis*, *E. edwardsii*, *E. tachinomoides*, *Exorista mella*, *Hyphantrophaga hyphantriae*, *Leschenaultia exul*, *Lespesia aletiae*, *L. archippivora*, *L. ciliata*, *L. frenchii*, *L. schizurae*, *Mericia ampelus*, *Parasetigena agilis*, *Pilatea arator*, "Tachina" spp., Tachinidae.*californicum*: *Archytas apicifer*, *A. lateralis*, *A. spp.* (including *A. apicifer* and/or *A. californiae*), *Euphorocera tachinomoides*, *Exorista mella*, *Hyphantrophaga hyphantriae*, *Leschenaultia fulvipes*, *Lespesia archippivora*, *L. ciliata*, *L. frenchii*, *L. spp.**californicum ambisimile*: *Lespesia archippivora*.*californicum californicum*: *Euexorista futilis*, *Exorista mella*, *Leschenaultia adusta*, *L. americana*, *Lespesia archippivora*, *L. frenchii*, *Tachinomyia similis*, Tachinidae.*californicum fragile*: *Archytas lateralis*, *A. spp.* (including *A. apicifer* and/or *A. californiae*), *Chaetogaedia sp. near monticola*, *Euphorocera tachinomoides*, *Exorista mella*, *Leschenaultia fulvipes*, *Lespesia archippivora*, *L. frenchii*.*californicum lutescens*: *Archytas lateralis*, *Exorista mella*, *Lespesia ciliata*, *L. frenchii*.*californicum pluviale*: *Compsilura concinnata*, *Euexorista futilis*, *Exorista mella*, *Leschenaultia americana*, *Lespesia archippivora*, *L. frenchii*, *Tachinomyia similis*, *T. variata*.

californicum recenseo: *Lespesia archippivora*.

californicum ssp. (intermediate between *M. californicum ambisimile* and *M. californicum californicum*): *Lespesia archippivora*.

californicum ssp. (intermediate between *M. californicum lutescens* and *M. californicum pluviale*): *Leschenaultia fulvipes*.

constrictum: *Exorista mella*, *Leschenaultia adusta*, *L. americana*, *Lespesia archippivora*, *L. frenchii*, Tachinidae.

constrictum austrinum: *Chaetogaedia* sp. near *monticola*.

constrictum constrictum: *Archytas lateralis*.

disstria: *Blepharipa scutellata*, *Carcelia laxifrons*, *C. malacosomae*, *C. malacosomae* probably, *C. protuberans*, *Cartocometes io*, *Compsilura concinnata*, *C.* sp. probably, *Drino inconspicua*, *Euexorista futilis*, *Euphorocera claripennis*, *E. edwardsii*, *Exorista mella*, *E.* spp., *Leschenaultia adusta*, *L. americana*, *L. exul*, *L. leucophrys*, *L.* spp., *Lespesia archippivora*, *L. frenchii*, *L. melalophae*, *L.* spp., *Madremyia saundersii*, *Mericia ampelus*, *Patelloa pachypyga*, *Tachinomyia nigricans*, *T. panaetius*, *T. similis*, *T.* spp., *Winthemia datanae*, *W. leucanae* lacking red at tip, *W.* sp. near *militaris*, Tachinidae.

disstria erosa, see *M. disstria*.

disstria erosa form *thoracica*, see *M. disstria*.

disstria and/or *M. pluviale*, see *M.* spp. (including *M. californicum pluviale* and/or *M. disstria*).

fragilis, see *M. californicum fragile*.

incurvum discoloratum: *Chaetogaedia* sp. near *monticola*, *Euphorocera tachinomoides*, *Hyphantrophaga hyphantriae*, *Lespesia archippivora*.

incurvum incurvum: *Archytas lateralis*, *Euphorocera tachinomoides*, *Exorista mella*, *Hyphantrophaga hyphantriae*, *Leschenaultia fulvipes*, *Lespesia archippivora*, *L. ciliata*, *Xanthophyto labis*.

pluviale, see *M. californicum pluviale*.

thoracica, see *M. californicum californicum*.

tigris: *Euphorocera tachinomoides*.

spp. (including *M. californicum pluviale* and/or *M. disstria*): *Leschenaultia americana*, *Tachinomyia similis*.

spp.: *Archytas lateralis*, *Carcelia malacosomae*, *Compsilura concinnata*, *Euexorista futilis*, *Euphorocera claripennis*, *Exorista mella*, *Leschenaultia americana*, *L.* spp., *Lespesia archippivora*, *L. frenchii*, *L.* sp., *Patel-*

loa pachypyga, *Tachinomyia panaetius*, *T. similis*,
Uramya halisidotae.

“*Malacosomoa*” *americanum*, see *Malacosoma americanum*

Maladera (Col.: Scarabaeidae)

castanea: *Dexilla ventralis*.

castanea probably: *Hyperecteina aldrichi*.

“*Malocosoma*” sp., see *Malacosoma* spp.

Mamestra (Lep.: Noctuidae)

adjuncta, see *Polia adjuncta*.

configurata: Tachinidae.

legitima, see *Polia legitima*.

“*ligitima*,” see *Polia legitima*.

picta, see *Ceramica picta*.

trifolii, see *Scotogramma trifolii*.

trifolii probably, see *Scotogramma trifolii* probably.

“*tripolii*,” see *Scotogramma trifolii*.

vindemialis, see *Sideridis maryx*.

Manduca (Lep.: Sphingidae)

quinquemaculata: *Compsilura concinnata*, *Drino incompta*, *D. rhoeo*, *D. rhoeo* probably, *Lespesia frenchii*, “*Sturmia*” spp., *Winthemia leucanae*, *W. quadripustulata*.

sexta: *Carcelia* spp., *Drino incompta*, *D. rhoeo*, *D. rhoeo* probably, *Lespesia* spp., “*Masicera*” spp., *Metavoria orientalis*, *Winthemia quadripustulata*, *W.* spp.

sexta jamaicensis: *Carcelia amplexa*, *Drino rhoeo*.

spp.: *Atacta brasiliensis*, *Drino incompta*.

Mantis carolina, see *Stagmomantis carolina*

Maple callus borer, see *Sylvora acerni*

Maple trumpet skeletonizer, see *Epinotia aceriella*

Mapsidius (Lep.: Yponomeutidae)

iridescens: *Lespesia archippivora*.

Margaronia hyalinata, see *Diaphania hyalinata*

Maruca (Lep.: Pyralidae)

testulalis: *Argyrophyllax albincisa*, *Nemorilla pyste*.

Marumba modesta, see *Pachysphinx modesta*

May beetle(-s) (Col.: Scarabaeidae)

May beetle(-s): *Cryptomeigenia prisca*, *C.* spp., *Eutrixa exilis*,
Ptilodexia harpasa, Tachinidae.

Meadow plant bug, see *Leptopterna dolabrata*

Meal moth, see *Pyralis farinalis*

Megalopyge (Lep.: Megalopygidae)

costaricensis: *Carcelia amplexa*.

crispata: *Anaporis pristis*, *Archytas aterrimus*, *A. smaragdinus*,
Carcelia amplexa, *C. lagoae*, *Compsilura concinnata*,

Euphorocera claripennis, *Hyphantrophaga hyphantriæ*, Tachinidae.

krugii: *Carcelia lagoae*.

opercularis: *Carcelia amplexa*, *C. lagoae*, *Euphorocera claripennis*, *E. claripennis* probably, *Lespesia aletiae*, *Sisyropa* spp., *Winthemia* spp., "Zenillia" spp.

opercularis probably: *Carcelia lagoae*.

pyxidifera: *Euphorocera claripennis*, *Lespesia aletiae*, *Sisyropa* spp.

sp.: *Lespesia aletiae*.

Megastes cyclades, see *Laniifera cyclades*

Megathymus (Lep.: Megathymidae)

yuccae: *Lespesia* spp., "Phorocera" *comstocki*.

Melalopha albosigma, see *Ichthyura albosigma*

Melalopha inclusa, see *Ichthyura inclusa*

Melalopha strigosa, see *Ichthyura strigosa*

Melanchroia (Lep.: Geometridae)

cephise: *Euphorocera floridensis*, *Prophryno parviteres*.

Melanolopia (Lep.: Geometridae)

imitata: *Actia interrupta*, *Compsilura concinnata*, *Eusisyropa virilis*, *Ictericophyto tibialis*.

"Melanopha" *apicalis* or "M." *inclusa*, see *Ichthyura* spp. (including *I. apicalis* or *I. inclusa*)

Melanoplus (Orth.: Acrididae)

atlanis, see *M. sanguinipes*.

bilituratus, see *M. sanguinipes*.

bivittatus: *Acemya tibialis*, *Ceracia dentata*, *Hemithrixion oestriforme*, Tachinidae.

bruneri: *Acemya tibialis*, *Ceracia dentata*, *Hemithrixion oestriforme*.

confusus: *Ceracia dentata*.

dawsoni: Tachinidae.

differentialis: *Ceracia dentata*, *Euphorocera claripennis*, *Lespesia* spp.

femurrubrum: *Acemya tibialis*, *Ceracia dentata*, Tachinidae.

foedus: *Ceracia* spp., Tachinidae.

"f. rubrum," see *M. femurrubrum*.

mexicanus, see *M. sanguinipes*.

occidentalis: *Ceracia* spp.

packardii: *Ceracia dentata*, Tachinidae.

sanguinipes: *Acemya tibialis*, *Ceracia dentata*, *Hemithrixion oestriforme*, *Leucostoma simplex*, Tachinidae.

spretus: *Lespesia* spp., Tachinidae.

spp. (including *M. bivittatus* and/or *M. differentialis*): Tachinidae.

- spp.: *Ceracia dentata*, Tachinidae.
- “*Melanopus*” *atlanis*, see *Melanoplus sanguinipes*
- Melanotus (Col.: Elateridae)
- spp.: *Ateloglossa cinerea*.
- Meliana *albilinea*, see *Faronta diffusa*
- Melipotis (Lep.: Noctuidae)
- famelica*: *Euphorocera tachinomoides*.
- Melissopus (Lep.: Olethreutidae)
- latiferranus*: *Carcelia* spp., *Elfia melissopodis*, *E.* spp., *Erynnia tortricis*, *Pseudoperichaeta erecta*, *P.* sp. near *erecta*.
- Melitara (Lep.: Pyralidae)
- dentata*: “*Phorocera*” *comstocki*, “*P.*” *texana*.
- doddalis*, see *M. dentata*.
- junctolineella*, see *Olycella junctolineella*.
- prodenialis*: *Lespesia* spp., “*Phorocera*” *texana*.
- spp.: *Coronimyia melitarae*, “*Phorocera*” *texana*.
- “*Melitta*” *cucurbitae*, see *Melittia cucurbitae*
- Melittia (Lep.: Aegeriidae)
- cucurbitae*: *Chaetophlepsis eudryae*.
- cucurbitae* probably: *Chaetophlepsis eudryae*.
- satyriniformis*: *Chaetophlepsis* spp.
- Melonworm, see *Diaphania hyalinata*
- Meracantha (Col.: Tenebrionidae)
- contracta*: *Catagoniopsis meracanthae*, *Pales* spp., *Zelia vertebrata*.
- Mesocondyla *concordalis*, see *Eulepte concordalis*
- Mesocondyla *concordalis* probably, see *Eulepte concordalis* probably
- Metanema (Lep.: Geometridae)
- inatomaria*: *Lespesia melalophae*.
- Metapodius *instabilis*, see *Acanthocephala terminalis*
- Metapodius *terminalis*, see *Acanthocephala terminalis*
- Metriona (Col.: Chrysomelidae)
- bicolor*: *Eucelatoriopsis dimmocki*.
- bivittata*, see *Agrioconota bivittata*.
- Mexican army worm, see Armyworm(-s)
- Mexican bean beetle, see *Epilachna varivestis*
- Mexican leafroller, see *Amorbia emigratella*
- Microlepidoptera, see Lepidoptera, miscellaneous
- Migratory grasshopper, see *Melanoplus sanguinipes*
- “Miller” moth of the group of owlet-moths, see Noctuidae, unidentified
- Mineola “*indiginella*,” see *Acrobasis indigenella*
- Mineola “*indiginella*” *nebullella*, see *Acrobasis indigenella*

Miris *dolabratus*, see *Leptopterna dolabrata*

"Mocia" *repanda*, see *Mocis* spp. (including *M. latipes* and/or *M. repanda*)

Mocis (Lep.: Noctuidae)

latipes probably: *Atracta brasiliensis*, *Belvosia bicincta*, *Lespesia aletiae*.

punctularis, see *M.* spp. (including *M. latipes* and/or *M. repanda*).

repanda, see *M. latipes* probably or *M.* spp. (including *M. latipes* and/or *M. repanda*).

spp. (including *M. latipes* and/or *M. repanda*): *Archytas marmoratus*, *Eucelatoria armigera*, *Euphorocera claripennis*, *Eusisyropa blanda*, *Gonia texensis*, *Gymnochaetopsis fulvicauda*.

Monarch butterfly, see *Danaus plexippus*

Monocesta (Col.: Chrysomelidae)

coryli: *Ervia triquetra*.

"*Monocestra*" *coryli*, see *Monocesta coryli*

Monochamus (Col.: Cerambycidae)

notatus: *Eutheresia monohammi*, *E. trivittata*.

scutellatus: *Eutheresia monohammi*, *E. trivittata*, Tachinidae.

titillator: *Eutheresia monohammi*.

spp.: *Eutheresia monohammi*.

Monoctenus (Hym.: Diprionidae)

fulvus: *Spathimeigenia aurifrons*, *S. spinigera*, *S.* spp., *Tachinomyia nigricans*.

Monodes nucicolora, see *Elaphria nucicolora*

"*Monohammus*" *confusor*, see *Monochamus notatus*

"*Monohamus*" *scutellatus*, see *Monochamus scutellatus*

Monoxia (Col.: Chrysomelidae)

consputa: *Chaetonodexodes vanderwulpi*.

puncticollis, see *Erynephila maritima*.

Moodna (Lep.: Pyralidae)

ostrinella: *Coronimymia geniculata*, *Erynnia tortricis*.

Moth borer of the sugar cane, see *Diatraea saccharalis*

Moth stalkborer(-s) (Lep.: Unidentified Lepidoptera)

Moth stalkborer(-s): *Lixophaga diatraeae*.

Moth(-s) (Lep.: Unidentified Lepidoptera)

Moth(-s): *Sisyropa eudryae*, "Tachina" spp., *Tachinomyia panaetius*.

Mountain ash sawfly, see *Pristiphora geniculata*

Mourningcloak butterfly, see *Nymphalis antiopa*

Mouse (mice) (Rodentia (Mammalia): Unidentified Rodentia)

Mouse (mice): *Parachaeta* spp.

Mouse (mice) probably: *Leschenaultia leucophrvs*. *Parachaeta* spp.

Murgantia (Hem. (Heter.): Pentatomidae)*histrionica*: *Trichopoda pennipes*.**Nabis** (Hem. (Heter.): Nabidae)*alternatus*: *Leucostoma simplex*.*americoferus*: *Leucostoma simplex*.*ferus*, see *N. americoferus*.*roseipennis*: *Leucostoma simplex*."Nacaleia" *indicata*, see *Hedylepta indicata**Nacoleia indicata*, see *Hedylepta indicata**Nacophora quernaria*, see *Phaeoura quernaria***Nadata** (Lep.: Notodontidae)*gibbosa*: *Compsilura concinnata*, Tachinidae.Nantucket pine tip moth, see *Rhyacionia frustrana***Narnia** (Hem. (Heter.): Coreidae)*pallidicornis*: *Trichopoda pennipes*.*Neleucania albilinea*, see *Faronta diffusa***Nematocampa** (Lep.: Geometridae)*filamentaria*: *Compsilura concinnata*, Tachinidae.*limbata*, see *N. filamentaria*.**Nematus** (Hym.: Tenthredinidae)*chalceus* probably: *Bessa harveyi*.*erichsonii*, see *Pristiphora erichsonii*.*erichsonii* probably, see *Pristiphora erichsonii* probably."ericksonii," see *Pristiphora erichsonii*.*erythrogaster*: *Compsilura concinnata*.*hyalinus*: *Bessa harveyi*.*limbatus*: *Bessa harveyi*.*ribesii*: *Bessa harveyi*, *Compsilura concinnata*, Tachinidae.*ventralis*: *Aporotachina chaetoneura*, *Bessa harveyi*, *Oswaldia aurifrons*.*ventricosus*, see *N. ribesii*.sp., see *Pristiphora erichsonii* probably.**Neoconocephalus** (Orth.: Tettigoniidae)*ensiger*: *Euphasiopteryx brevicornis nuttingi*.*robustus*: *Euphasiopteryx brevicornis nuttingi*.*robustus robustus*: *Beskia aelops* probably, *Euphasiopteryx brevicornis nuttingi*."Neodipiron" *tsugae*, see *Neodiprion tsugae***Neodiprion** (Hym.: Diprionidae)*abbottii*: *Diplostichus lophyri*, *Euphorocera claripennis*, *E. edwardsii*, *Spathimeigenia spinigera*, *S. spinigera* complex."abbottii," see *Neodiprion abbotii*.*abietis*: *Bessa harveyi*, *Diplostichus lophyri*, *Drino bohémica*, *Spathimeigenia spinigera*, *S. spinigera* complex.

- abietis* complex: *Bessa harveyi*, *Diplostichus lophyri*, *D. sp.* near *lophyri*, *Spathimeigenia aurifrons*.
- affinis*, see *N. virginiana*.
- americanus*, see *N. abbotii*.
- americanus banksianae*, see *N. pratti banksianae*.
- banksianae*, see *N. pratti banksianae*.
- dyari*, see *N. pratti pratti*.
- edwardsii*, see *N. spp.*
- excitans*: “*Phorocera*” spp., *Spathimeigenia spinigera*.
- fabricii*, see *N. abbotii*.
- lecontei*: *Bessa harveyi*, *B. selecta*, *Diplostichus lophyri*, *Drino bohémica*, *D. inconspicua*, *Eribella sp.* near *exilis*, *Euphorocera claripennis*, *E. edwardsii*, “*Phorocera*” spp., *Spathimeigenia aurifrons*, *S. erecta*, *S. erecta* probably, *S. hylotomae*, *S. spinigera*, *S. spinigera* complex, *S. spp.*, “*Zenillia*” spp.
- “*leconti*,” see *N. lecontei*.
- maurus*: *Bessa harveyi*, *Spathimeigenia aurifrons*, *S. spinigera*, *S. spp.*
- nanulus*, see *N. nanulus nanulus*.
- nanulus nanulus*: *Bessa harveyi*, *Diplostichus lophyri*, *Drino bohémica*, *Spathimeigenia spinigera*, *S. spinigera* complex.
- nigroscutum*: *Diplostichus lophyri*, *Spathimeigenia spp.* (*aurifrons-erecta* complex).
- pinetum*: *Diplostichus lophyri*, *Spathimeigenia erecta* probably, *S. spinigera*, *S. spinigera* complex, *S. spp.*
- pinirigidae*: *Spathimeigenia aurifrons*.
- pratti banksianae*: *Bessa harveyi*, *Diplostichus lophyri*, *Drino bohémica*, *Spathimeigenia aurifrons*, *S. erecta*, *S. spinigera*, *S. spinigera* complex.
- pratti pratti*: *Diplostichus lophyri*, *Drino inconspicua*.
- rugifrons*: *Diplostichus lophyri*, *Spathimeigenia aurifrons*, *S. erecta*, *S. spp.*
- sp.* near “*scutellaris*,” see *N. sp.* near *scutellatis*.
- sp.* near *scutellatis*: *Diplostichus lophyri*, *Tsugaea nox*.
- sertifer*: *Diplostichus lophyri*, *Drino bohémica*, *D. inconspicua*, *Euphorocera claripennis*, *E. edwardsii*, *E. sp.*, *Spathimeigenia erecta*, *S. spinigera*, *S. spp.*
- swaini*, see *N. swaini* or *N. swaini* probably.
- swaini*: *Bessa harveyi*, *B. selecta*, *Diplostichus lophyri*, *Drino bohémica*, “*Phorocera*” spp., *Spathimeigenia aurifrons*, *S. spp.* (*aurifrons-erecta* complex), *S. spinigera*, *S. spinigera* complex, *S. spp.*

swainei probably: *Diplostichus lophyri*, *Spathimeigenia* spp.
taedae linearis: *Diplostichus lophyri*, *Spathimeigenia spinigera*,
S. spp.

tsugae: *Diplostichus sellersi*, *Tsugaea nox*.

vallicola, see *Zadiprion vallicola*.

virginiana: *Euphorocera claripennis*, *Spathimeigenia spinigera*.

virginiana complex: *Bessa harveyi*, *Diplostichus lophyri*,
Drino bohémica, *Spathimeigenia spinigera* complex.

spp.: *Bessa harveyi*, *Diplostichus lophyri*, *D. sellersi*,
Eucelatoria armigera, *Euphorocera claripennis*,
“*Phorocera*” spp., *Spathimeigenia aurifrons*, *S. erecta*,
S. spinigera.

sp. probably: *Diplostichus lophyri*.

Neophasia occidentalis, see *Pieris occidentalis*

Nephelodes (Lep.: Noctuidae)

emmedonia: *Aplomya trisetosa*, *Euexorista futilis*, *Phryxe*
pecosensis probably, *Tachinomyia variata*, Tachinidae.

violans, see *N. emmedonia*.

Nephopteryx (Lep.: Pyralidae)

subcaesiella: *Eusisyropa blanda*, *Nemorilla pyste*, *Phryxe*
pecosensis.

subcaesiella probably: Tachinidae.

subfuscilla: *Aphria ocypterata*, *Eusisyropa blanda*.

“*Nephytia*” *canosaria*, see *Nepytia canosaria*

Nepytia (Lep.: Geometridae)

canosaria: *Bessa harveyi*, *Blondelia eufitchiae*, *Chaetophlepsis*
semiothisae.

phantasmaria: *Eusisyropa blanda*, Tachinidae.

Neurotoma (Hym.: Pamphiliidae)

fasciata: *Aplomya neurotomae*, *Compsilura concinnata*.

inconspicua: *Eubrachymera debilis*.

spp.: *Aplomya fronto*.

New Guinea sugarcane borer, see *Rhabdoscelus obscurus*

New Guinea sugarcane weevil, see *Rhabdoscelus obscurus*

Nezara (Hem. (Heter.): Pentatomidae)

viridula, see *N. viridula* or *N. viridula smaragdula*.

viridula: *Trichopoda lanipes*, *T. pennipes*, *T.* spp., Tachinidae.

viridula smaragdula: *Trichopoda pennipes*, *T. pilipes*, *T.* spp.
(including *T. pennipes* and/or *T. pilipes*), *T.* spp.

“*Nezarra*” *viridula*, see *Nezara viridula*

Nisoniades brizo, see *Erynnis brizo*

Noctua fennica, see *Actebia fennica*

Noctua sp., see Noctuidae, unidentified

Noctuid (Lep.: Noctuidae)

spp. (including *Mamestra configurata* and/or *Peridroma saucia*):
Lespesia archippivora, *Phryxe pecosensis*, *Winthemia rufopicta*.

spp. (including *Heliothis zea*, *Prodenia dolichos*, *Spodoptera eridania*, *S. frugiperda* and/or *S. ornithogalli*):
Winthemia rufopicta.

Noctuid and/or pyralid (Lep.: Noctuidae and/or Pyralidae)

spp. (including *Scotogramma trifolii* and/or *Loxostege sticticalis*): *Euphorocera claripennis*, *Lespesia archippivora*, *Voria ruralis*.

Noctuidae (Lep.: Noctuidae)

Unidentified: *Archytas aterrimus*, *Blepharipa scutellata*, *Bonnetia comta*, *Chaetogaedia monticola*, *Chaetophlepsis eudryae*, *Compsilura concinnata*, *Cyrtophleba coquilletti*, *Gymnocarcelia ricinorum*, *Lespesia frenchii*, *L. spp.*, *Mericia spp.*, *Nilea noctuiformis*, *Peleteria obsoleta*, *Periscepsia helymus*, *Phryxe pecosensis*, *P. vulgaris*, *Siphona ceres*, *S. plusiae*, *Tachinomyia spp.*, *Voria ruralis*, *Winthemia quadripustulata*, Gen. and sp. near *Zenillia*, Tachinidae.

Nonagria oblonga, see *Archanara oblonga*

Northeastern sawyer, see *Monochamus notatus*

Northern pitch twig moth, see *Petrova albicapitana*

Notodonta ulmi, see *Heterocampa bilineata*

Notodontidae (Lep.: Notodontidae)

Unidentified: *Compsilura concinnata*.

Notolophus antiqua, see *Orgyia antiqua*

Notolophus leucostigma, see *Hemerocampa leucostigma*

Nutgrass armyworm, see *Spodoptera exempta*

Nycteola (Lep.: Noctuidae)

cinereana: *Lespesia melalophae*.

revayana, see *N. spp.* (including *N. cinereana*, *N. columbiana* and/or *N. frigidana*).

spp. (including *N. cinereana*, *N. columbiana* and/or *N. frigidana*): *Euexorista futilis*, *Pseudoperichaeta erecta*, *Zenillia blandita*.

Nygmia (Lep.: Lymantriidae)

phaeorrhoea: *Aplomya caesar*, *Carcelia laxifrons*, *Compsilura concinnata*, *Euphorocera claripennis*, *Eusisyropa blanda*, *Exorista mella*, *E. spp.*, *Gonia spp.*, *Lespesia frenchii*, *Madremyia saundersii*, *Patelloa pachypyga*, *Townsendiellomyia nidicola*.

Nymphalidae (Lep.: Nymphalidae)

Unidentified: *Compsilura concinnata*.

Nymphalis (Lep.: Nymphalidae)

antiopa: *Bessa harveyi*, *Compsilura concinnata*, *Dichaetoneura leucoptera*, *Doryphorophaga doryphorae*, *Euphorocera claripennis*, *E. edwardsii*, *Eusisyropa blanda*, *Exorista mella*, *Hemisturmia tortricis*, *Lespesia aletiae*, *L. archippivora*, *L. dubia*, *L. frenchii*, *Madremyia saundersii*, *Patelloa pachypyga* probably, *Pelatachina pellucida*, "Phorocera" spp., *Winthemia sinuata* probably, Tachinidae.

milberti: *Compsilura concinnata*, *Euexorista futilis*, *Hyphantrophaga hyphantriae*, *H. sp.*, *Lespesia archippivora*, *Pelatachina pellucida*, *Winthemia quadripustulata* probably, Tachinidae.

vau-album j-album: *Compsilura concinnata*.

Nysius (Hem. (Heter.): Lygaeidae)

angustatus: "*Phorantha occidentis*."

ericae: *Hyalomya aldrichii*, "*Phorantha occidentis*."

raphanus: *Hyalomya aldrichii*.

Oak leaftier, see *Croesia albicomana*

Oak webworm, see *Archips fervidanus*

Oakworm(-s) (Lep.: Unidentified Lepidoptera)

Oakworm(-s): *Eusisyropa blanda*.

Oberea (Col.: Cerambycidae)

bimaculata: *Lixophaga variabilis*.

"*maculata*," see *O. bimaculata*.

Obliquebanded leafroller, see *Choristoneura rosaceana*

Ochlodes (Lep.: Hesperidae)

yuma: *Spathidexia dunningii*.

Ocneria dispar, see *Porthetria dispar*

Oebalus (Hem. (Heter.): Pentatomidae)

pugnax: *Euthera tentatrix*.

Oedemasia concinna, see *Schizura concinna*

Oedipoda carolina, see *Dissosteira carolina*

Oenectra distincta, see *Sparganothis distincta*

Oeobia (Lep.: Pyralidae)

monticolans: *Chaetogaedia monticola*.

Ogdoconta (Lep.: Noctuidae)

cinereola: *Copecrypta ruficauda*.

Oidaematophorus (Lep.: Pterophoridae)

homodactylus: *Compsilura concinnata*, *Oxynops anthracinus*, *Pseudosiphona brevirrostris*, *Tachinophyto floridensis*.

monodactylus: *Oxynops anthracinus*, Tachinidae.

Oiketicus (Lep.: Psychidae)

kirbyi: *Lespesia* spp.

- Olene* "atomaria," see *Dasychira atomaria*
Olene basiflava, see *Dasychira basiflava*
Olene pini, see *Dasychira plagiata*
Olene plagiata, see *Dasychira plagiata*
Olene vagans, see *Dasychira vagans*
Olene spp., see *Dasychira* spp.
Olethreutes hebesana, see *Endothenia hebesana*
Olethreutes malachitana, see *Eumarozia malachitana*
 Olethreutid and/or pyralid (Lep.: Olethreutidae and/or Pyralidae)
 spp. (including *Laspeyresia* spp. and/or *Dioryctria* spp.
 probably): *Exorista* spp.
 Olethreutidae (Lep.: Olethreutidae)
 Unidentified: *Leskiella brevirostris*.
Oligia (Lep.: Noctuidae)
 fractilinea: *Winthemia rufopicta*.
Olyca junctolineella, see *Olycella junctolineella*
Olycella (Lep.: Pyralidae)
 junctolineella: *Lespesia aletiae*, "Phorocera" *comstocki*, "P."
 texana.
 nephelepassa: "Phorocera" *texana*.
Omiodes accepta, see *Hedylepta accepta*
Omiodes blackburni, see *Hedylepta blackburni*
Omiodes continuatalis, see *Hedylepta continuatalis*
Omphalocera (Lep.: Pyralidae)
 cariosa: *Carcelia lagoae*, *Hyphantrophaga hyphantriae*.
Oncopeltus (Hem. (Heter.): Lygaeidae)
 fasciatus: *Leucostoma gravipes*.
 Onespot stink bug, see *Euschistus variolarius*
Oniscus (Isopoda (Crustacea): Oniscidae)
 asellus: *Melanophora roralis*.
Operophtera (Lep.: Geometridae)
 bruceata: *Cyzenis albicans*.
 brumata: *Cyzenis albicans*, *Phryxe pecosensis*.
Opsiphanes (Lep.: Brassolidae)
 tamarindi: *Lespesia aletiae*.
 Orangedog, see *Papilio cresphontes*
 Orangestriped oakworm, see *Anisota senatoria*
 Orange tortrix, see *Argyrotaenia citrana*
Orgyia (Lep.: Lymantriidae)
 antiqua: *Carcelia amplexa*, *C. yalensis*, *Compsilura concinnata*,
 Eusisyropa blanda, *Exorista mella*, Tachinidae.
 leucostigma, see *Hemerocampa leucostigma*.
 "leugcostigma," see *Hemerocampa leucostigma*.
 spp.: *Lespesia aletiae*, Tachinidae.

- Oriental fruit moth, see *Grapholitha molesta*
Oriental moth, see *Cnidocampa flavescens*
Oriental peach moth, see *Grapholitha molesta*
Orphulella (Orth: Acrididae)
 speciosa: Tachinidae.
Orthodes (Lep.: Noctuidae)
 cynica: Tachinidae.
Orthoptera (Orth: Unidentified Orthoptera)
 Miscellaneous: *Euphasiopteryx brevicornis nuttingi*.
Orthosia (Lep.: Noctuidae)
 hibisci: *Tachinomyia* sp. near *panaetius*, *T.* spp., *Wagneria vernata* probably.
 hibisci form *insciens*, see *O. hibisci*.
 “*Orygia*” spp., see *Orgyia* spp.
Osmoderma (Col.: Scarabaeidae)
 eremicola: *Zelia vertebrata*.
Ostrinia (Lep.: Pyralidae)
 nubilalis: *Aplomya caesar*, *A. mitis*, “*Carcelia ochracea*,”
 Ceromasia auricaudata, *Compsilura concinnata*,
 Euexorista futilis, *Lixophaga variabilis*, *Lydella radialis*,
 L. thompsoni, *Madremyia saundersii*, *Nemorilla pyste*,
 Paratheresia claripalpis, *Pseudoperichaeta erecta*, *P.*
 insidiosa, *Pyraustomyia penitalis*, “*Zenillia*” spp.
 obliteralis: *Aplomya caesar*, *Didyma exigua*, “*Hypostena floridensis*,” *Pyraustomyia penitalis*.
 penitalis: “*Anisia nigrocincta*” probably, *Aplomya caesar*,
 Lixophaga variabilis, *Pyraustomyia penitalis*.
 spp. (including *O. obliteralis* and/or *O. penitalis*): *Lixophaga variabilis*, *Nemorilla pyste*, “*Phorocera*” *comstocki*,
 Phryxe pecosensis probably.
Oulema (Col.: Chrysomelidae)
 melanopus: *Hyalomyodes triangulifer*.
Pachysphinx (Lep.: Sphingidae)
 modesta: *Lespesia frenchii*, *Winthemia quadripustulata*.
 modesta occidentalis: Tachinidae.
Pachyzancla (Lep.: Pyralidae)
 bipunctalis: *Nemorilla pyste*, *Pseudoperichaeta erecta*.
 licarsisalis, see *Herpetogramma licarsisalis*.
 periusalis, see *Psara pallicaudalis*.
 “*pertusalis*,” see *Psara pallicaudalis*.
Pacific tent caterpillar, see *Malacosoma constrictum*
Packard grasshopper, see *Melanoplus packardii*
Packardia (Lep.: Limacodidae)
 ceanothi: *Eupelecotheca celer*.

Painted beauty, see *Vanessa virginiensis*

Painted lady, see *Vanessa cardui*

Paleacrita (Lep.: Geometridae)

vernata: Tachinidae.

Palesided cutworm, see *Agrotis malefida*

Pale tussock moth, see *Halisidota tessellaris*

Pale western cutworm, see *Agrotis orthogonia*

Palpita (Lep.: Pyralidae)

magniferalis: *Pseudochaeta siminina*.

Pandemis (Lep.: Tortricidae)

"*limitana*," see *P. limitata*.

limitata: *Aplomya caesar*, *Ictericophyto spinosa*.

pyrusana: *Nemorilla pyste*.

Pandora moth, see *Coloradia pandora*

Pantographa (Lep.: Pyralidae)

limata: *Carcelia* spp., *Pseudoperichaeta erecta*.

Paonias (Lep.: Sphingidae)

excaecata: *Compsilura concinnata*, *Leschenaultia leucophrys*,
Winthemia datanae.

"*excoecata*," see *P. excaecata*.

myops: *Compsilura concinnata*, *Lespesia flavifrons*, *L. frenchii*,
Tachinidae.

Papaipema (Lep.: Noctuidae)

appassionata: *Lydella radialis*.

"*appassionata*," see *P. appassionata*.

araliae: *Lydella radialis*.

arctivorens: *Gymnocheta ruficornis*, *Lydella radialis*.

cataphracta: *Exorista* spp., *Gymnocheta ruficornis*, *Lixophaga*
variabilis, *Lydella radialis*, *Winthemia rufopicta*.

frigida: *Lydella radialis*.

harrisii: *Eusisyropa virilis*.

nebris: *Aplomya caesar*, *Exorista* spp., *Gymnocheta ruficornis*,
Lixophaga variabilis, *Lydella deckeri*, *L. radialis*, *L.*
radialis probably, *L. thompsoni*, "*Phorocera*" *signata*,
"*Tachina*" spp., *Winthemia quadripustulata*, *W.*
rufopicta, Tachinidae.

nebris nitela, see *P. nebris*.

nitela, see *P. nebris*.

pterisii: *Lydella radialis*.

purpurifascia: *Ceromasia* spp., *Lydella radialis*.

stenoscelis: *Lydella radialis*.

[sp.], see *P. nebris*.

spp.: *Lydella radialis*, *L. radialis* probably.

Paphia [sp.], see *Anaea glycerium*

Papilio (Lep.: Papilionidae)

ajax, see *P. polyxenes asterius* or *P. polyxenes polyxenes*.

"*asterias*," see *P. polyxenes asterius*.

bairdii oregonius: *Lespesia frenchii*.

crephontes: *Lespesia rileyi*.

daunus, see *P. multicaudata*.

eurymedon: *Lespesia frenchii*.

glaucus: *Compsilura concinnata*, *Lespesia frenchii*, Tachinidae.

glaucus canadensis: *Lespesia frenchii*.

glaucus turnus, see *P. glaucus*.

machaon: *Buquetia musca*.

multicaudata: *Lespesia frenchii*.

philenor, see *Battus philenor*.

polyxenes, see *P. polyxenes*, *P. polyxenes asterius* or *P. polyxenes polyxenes*.

polyxenes: *Lespesia frenchii*.

polyxenes asterius: *Buquetia obscura*, *Lespesia frenchii*, Tachinidae.

polyxenes polyxenes: *Compsilura concinnata*.

rutulus: *Lespesia frenchii*.

thoas: *Lespesia rileyi*.

troilus: *Compsilura concinnata*, *Lespesia frenchii*.

turnus, see *P. glaucus*.

turnus var. glaucus, see *P. glaucus*.

zelicaon: Tachinidae.

"*zelicayn*," see *P. zelicaon*.

"*zolicaon*," see *P. zelicaon*.

spp.: *Lespesia archippivora*, *L. frenchii*.

Papilionidae (Lep.: Papilionidae)

Unidentified: *Compsilura concinnata*.

"**Papilo**" *thoas*, see *Papilio thoas*

Paragrotis ochrogaster, see *Euxoa ochrogaster*

Paragrotis ridingsiana, see *Euxoa ridingsiana*

Paralechia pinifoliella, see *Exoteleia pinifoliella*

Paralechia pinifoliella probably, see *Exoteleia pinifoliella* probably

Parallelia (Lep.: Noctuidae)

bistriaris: *Oswaldia albifacies*, *O. anorbitalis*, *O. assimilis*, Tachinidae.

Parargyractis (Lep.: Pyralidae)

fulicalis: *Ginglymia acirostris*.

Parasemia (Lep.: Arctiidae)

parthenos: *Thompsonomyia anthracina*.

Parharmonia pini, see *Vespamima pini*

Parsleyworm, see *Papilio polyxenes asterius*

- Parsnip webworm, see *Depressaria pastinacella*
- Passalus cornutus, see *Popilius disjunctus*
- Peach twig borer, see *Anarsia lineatella*
- Pea moth, see *Laspeyresia nigricana*
- Pecan bud moth, see *Gretchena bolliana*
- Pecan carpenterworm, see *Cossula magnifica*
- Pecan leaf casebearer, see *Acrobasis juglandis*
- Pecan nut casebearer, see *Acrobasis caryae*
- Pectinophora (Lep.: Gelechiidae)
gossypiella: *Erynnia tortricis*.
- Pempelia (Lep.: Pyralidae)
 spp.: *Hyphantrophaga hyphantriae*.
- Pentatoma ligata, see *Chlorochroa ligata*
- Pentatoma sayi, see *Chlorochroa sayi*
- "Pentato mia," see *Chlorochroa* spp.
- Pentatomid (Hem. (Heter.): Pentatomidae)
 spp. (including *Euschistus ictericus*, *Mormidea pictiventris*,
Nezara viridula and/or *Oebalus pugnax*): *Beskia aelops*,
Cylindromyia euchenor, *Euthera tentatrix*.
- Pepper-and-salt moth, see *Biston cognataria*
- Peridroma (Lep.: Noctuidae)
cinctipennis cinctipennis: *Chaetogaedia monticola*.
coniotis: *Lespesia archippivora*.
coniotis coniotis: *Chaetogaedia monticola*, *Lespesia archippi-*
vora.
margaritosa, see *P. saucia*.
margaritosa saucia, see *P. saucia*.
porphyrea, see *P. saucia*.
saucia: *Archytas apicifer*, *A. aterrimus*, *A. cirphis*, *A. cirphis*
 probably, *A. spp.* (including *A. apicifer* and/or *A.*
californiae), *A. spp.*, *Bonnetia comta*, *Carcelia spp.*,
Chaetogaedia monticola, *C. monticola* probably,
Clausicella opaca, *Eucelatoria armigera* probably,
Euphorocera claripennis, *E. omissa*, *Gonia longipulvilli*,
G. porca, *G. sequax*, *G. spp.*, *Lespesia archippivora*,
Madremyia saundersii, *Peleteria texensis*, *Periscepsia*
helymus, *P. laevigata*, *Voria ruralis*, *Winthemia*
leucanae, *W. leucanae* lacking red at tip, *W.*
quadripustulata, *W. rufopicta*, Tachinidae.
- "Perigaea" sutor, see *Platysenta sutor*
- Perigea sutor, see *Platysenta sutor*
- Periodical cicada, see *Magicicada septendecim*
- Periplaneta (Orth.: Blattidae)
americana: *Miamimyia cincta*.

Pero (Lep.: Geometridae)

sp.: *Chaetophlepsis semiothisae*.

Peronea minuta, see *Acleris minuta*

Peronea variana, see *Acleris variana*

"*Peronia*" *minuta*, see *Acleris minuta*

Petrova (Lep: Olethreutidae)

albicapitana: *Lixophaga* sp. near *fasciata*.

metallica: *Phrynofrontina* spp.

Phaecasiophora (Lep.: Tortricidae)

niveiguttana: *Aplomya caesar*.

Phaeoura (Lep.: Geometridae)

quernaria: *Aplomya helvina*, Tachinidae.

Phakellura nitidalis probably, see *Diaphania nitidalis* probably

Phalaenid, see Noctuidae, unidentified

Phalaenid moth(-s) (Lep.: Noctuidae)

Phalaenid moth(-s): *Voria ruralis*.

Phalonia (Lep.: Tortricidae)

oenotherana: *Actia interrupta*, *Elfia aenea*, *Lixophaga plumbea*.

Phantom hemlock looper, see *Nepytia phantasmaria*

Phasiana "neptata," see *Semiothisa neptaria*

Pheosia (Lep.: Notodontidae)

dimidiata, see *P. rimosa*.

rimosa: *Aplomya pheosiae*, *Chaetophlepsis eudryae*, *Compsilura concinnata*, *Winthemia datanae*.

Phigalia (Lep.: Geometridae)

titea: *Ceromasia* spp., *Compsilura concinnata*, *Eusisyropa virilis*, *Omotoma fumiferanae*, *Tachinomyia nigricans*, *Tachinomyia* spp., Tachinidae.

Philaenus (Hem. (Hom.): Cercopidae)

spumarius: *Alophorella aeneoventris*.

Philampelus achemon, see *Pholus achemon*

Philampelus vitis, see *Pholus vitis*

Philosamia cynthia, see *Samia cynthia*

Philtraea (Lep.: Geometridae)

elegantaria: *Euphorocera claripennis*.

Phlegethontius carolina, see *Manduca sexta*

Phlegethontius celeus, see *Manduca quinquemaculata*

Phlegethontius cingulatus, see *Agrius cingulatus*

Phlegethontius quinquemaculata, see *Manduca quinquemaculata*

Phlegethontius sexta, see *Manduca sexta*

Phlegethontius spp., see *Manduca* spp.

Phlyctaenia (Lep.: Pyralidae)

coronata tertialis: *Aplomya caesar*, *Nemorilla pyste*, *Pseudoperichaeta erecta*, Tachinidae.

"*ferrugalis*," see *Udea rubigalis*.

- ferrygalis*, see *Udea rubigalis*.
monticolans, see *Oebia monticolans*.
tertialis, see *P. coronata tertialis*.
- Phobetron** (Lep.: Limacodidae)
pithecium: *Anaporia pristis*, *Carcelia* spp., *Palpexorista coccyx*,
Tachinidae.
- Pholus** (Lep.: Sphingidae)
achemon: *Argyrophylax* spp., *Compsilura concinnata*, *Drino incompta*, *D. rhoeo*, *Lespesia testacea*, *Winthemia quadripustulata*.
fasciatus, see *P. vitis*.
pandorus, see *P. satellitia pandorus*.
satellititia pandorus: *Compsilura concinnata*, *Lespesia pholi*, *L. schizurae*, Tachinidae.
vitis: *Drino incompta*, *Lespesia testacea*.
spp.: *Lespesia aletiae*, *L. pholi*.
- Phosphila** (Lep.: Noctuidae)
turbulenta: *Compsilura concinnata*, *Eusisyropa blanda*, *E. virilis*.
- Phragmatobia** (Lep.: Arctiidae)
assimilans: *Carcelia reclinata*, *C. reclinata* probably.
assimilans var. *franconia*, see *P. assimilans*.
assimilans var. "franconica," see *P. assimilans*.
fuliginosa: *Hubneria estigmenensis*.
fuliginosa rubricosa: *Hubneria estigmenensis*, Tachinidae.
- "Phryganidea" *californica*, see *Phryganidia californica*
- Phryganidia** (Lep.: Diopsideae)
californica: *Actia flavipes*, *A.* spp., *Eusisyropa virilis*, *Siphona conata*, *S. plusiae*, Tachinidae.
[sp.], see *P. californica*.
- Phycid in dried peach, see Phycitinae, unidentified
- Phyciodes** (Lep.: Nymphalidae)
tharos: *Siphosturmia phyciodis*.
spp.: *Siphosturmia phyciodis*.
- Phycita nebulo**, see *Acrobasis indigenella*
- Phycitinae** (Lep.: Pyralidae)
Unidentified: *Coronimyia geniculata*.
- Phyllophaga** (Col.: Scarabaeidae)
anxia: *Cryptomeigenia* spp., *Eutrixa exilis*, *Microphthalma michiganensis*.
arcuata, see *P. fervida*.
crassissima: *Cryptomeigenia theutis*.
crenulata: *Cryptomeigenia theutis*, *Sitophaga georgiae*.
drakii: *Eutrixa exilis*.

fervida: *Eutrixa exilis*, *Microphthalma disjuncta*, *M. michiganensis*.

fusca: Tachinidae.

futilis: *Cryptomeigenia theutis*.

implicita: *Cryptomeigenia theutis*.

inversa: *Cryptomeigenia theutis*, Tachinidae.

nitida: *Eutrixa exilis*.

portoricensis: *Cryptomeigenia aurifacies*, *Eutrixoides jonesii*.

rubiginosa: *Eutrixa exilis*.

rugosa: *Cryptomeigenia theutis*.

vandinei, see *Chemarachis vandinei*.

spp. (including *P. anxia* and/or *P. fusca*): *Cryptomeigenia triangularis*.

spp.: *Bonnetia comta*, *Cryptomeigenia aurifacies*, *C. simplex*, *C. theutis*, *C. spp.*, *Dexilla ventralis*, *Dinera grisescens* probably, *Eutrixa exilis*, *Eutrixoides jonesii*, *Gonia crassicornis*, *Microphthalma disjuncta*, *M. michiganensis*, *M. obsoleta*, "*Ptilodexia abdominalis*," *P. cerata*, *P. harpasa*, "*P. tibialis*," *Sitophaga georgiae*, *S. nocturnalis*, Tachinidae.

"*Phyllophoga*" *anxia*, see *Phyllophaga anxia*

Physonota (Col.: Chrysomelidae)

helianthi: Tachinidae.

Physostegania pustularia, see *Itame pustularia*

Phytonomus nigrirostris, see *Hypera nigrirostris*

Pickleworm, see *Diaphania nitidalis*

Pictured sand cricket(-s) (Orth: Unidentified Orthoptera)

Pictured sand cricket(-s): *Stomatomyia parvipalpis*.

Pieris (Lep.: Pieridae)

occidentalis: *Chaetogaedia crebra*, *Madremyia saundersii*.

protodice: *Euphorocera* sp. near *floridensis*, *Gymnoclytia occidua*, *Lespesia aletiae*, *L. archippivora*, Tachinidae.

rapae: *Ceromasia auricaudata*, *Chaetogaedia monticola*, *Compsilura concinnata*, *Eucelatoria armigera*, *Euphorocera claripennis*, *E. tachinomoides*, *Gymnoclytia occidua*, *Lespesia aletiae*, *L. archippivora*, *L. frenchii*, *L. spp.*, *Madremyia saundersii*, *Obolocera rapae*, *Phryxe pecosensis*, *P. vulgaris*, Tachinidae.

sp.: *Lespesia archippivora*.

Pikonema (Hym.: Tenthredinidae)

alaskensis: *Bessa harveyi*, *B. selecta*, *Drino bohémica*, *Phryxe pecosensis*, *Spathimeigenia aurifrons*, *S. spinigera*.

dimmockii: *Bessa harveyi*, *B. selecta*, *Drino bohémica*.

spp.: *Bessa harveyi*, *Diplostichus lophyri*.

- Pilemia** (Lep.: Pyralidae)
periusalis: *Argyrophylax albincisa*.
- Pilocrocis** (Lep.: Pyralidae)
inguinalis: *Argyrophylax albincisa*.
tripunctata: *Nemorilla pyste*.
- Pine needleminer, see *Exoteleia pinifoliella*
- Pine sawfly(-flies) (Hym.: Unidentified Hymenoptera)
 Pine sawfly(-flies): *Spathimeigenia spinigera*.
- Pine tussock moth, see *Dasychira plagiata*
- Pine webworm, see *Tetralopha robustella*
- Pink bollworm, see *Pectinophora gossypiella*
- Pinkstriped oakworm, see *Anisota virginensis*
- Pionea belialis**, see *Titanio belialis*
- “**Pipausta ainslei**,” see *Ostrinia obliteralis*
- Pipevine swallowtail, see *Battus philenor*
- Pissodes** (Col.: Curculionidae)
strobi: *Compsilura concinnata*.
- Pistol casebearer, see *Coleophora malivorella*
- Pitch mass borer, see *Vespamima pini*
- Pitch pine sawfly(-flies) (Hym.: Unidentified Hymenoptera)
 Pitch pine sawfly(-flies): “*Phorocera*” spp.
- Pitch pine tip moth, see *Rhyacionia rigidana*
- Plagiometriona** (Col.: Chrysomelidae)
clavata: *Eribella exilis*, *Eucelatoriopsis dimmocki*, *E. dimmocki*
 probably, *Medina barbata*.
- Plains false wireworm, see *Eleodes opacus*
- Plant bug(-s) (Hem. (Heter.): Pentatomidae)
 Plant bug(-s): *Trichopoda* spp., Tachinidae.
- Plathypena** (Lep.: Noctuidae)
scabra: *Archytas aterrimus*, *Carcelia amplexa*, *Chaetophlepsis eudryae*, *C. spp.*, *Compsilura concinnata*, *Copecrypta ruficauda*, *Euphorocera claripennis*, *E. floridensis*, *Eusisyropa blanda*, *Lespesia aletiae*, *Prophryno parviteres*, *Winthemia quadripustulata*, *W. sinuata*.
scabra probably: *Euphorocera claripennis*, *Eusisyropa boarmiae*, *E. boarmiae* probably.
- Platycampus** (Hym.: Tenthredinidae)
 spp.: *Bessa harveyi*.
- Platynota** (Lep.: Tortricidae)
flavedana: *Erynnia tortricis*.
- Platyptilia** (Lep.: Pterophoridae)
carduidactyla: *Elfia* spp., *Hyalomyodes triangulifer*.

Platysamia¹ (Lep.: Saturniidae)

cecropia: *Compsilura concinnata*, *Euphorocera claripennis*, *Eusisyropa virilis*, *Lespesia ciliata*, *L. frenchii*, "Tachina" spp., *Winthemia cecropia*, *W. datanae*, *W. leucanae*, *W. leucanae* lacking red at tip, *W. militaris*, *W. quadripustulata*, *W. spp.*, Tachinidae.

euryalus: *Lespesia ciliata*, *L. frenchii*, *L. sabroskyi*.

Platysenta (Lep.: Noctuidae)

sutor: *Winthemia quadripustulata*, *W. rufopicta*.

videns: *Winthemia quadripustulata*.

Plautia (Hem. (Heter.): Pentatomidae)

stali: *Trichopoda pennipes*.

Plebeius melissa, see *Lycaeides melissa***Plebejus** (Lep.: Lycaenidae)

icarioides: Tachinidae.

Plodia (Lep.: Pyralidae)

interpunctella: *Clausicella floridensis*, *C. neomexicana*.

spp.: *Clausicella floridensis*, *C. neomexicana*.

Plum curculio, see *Conotrachelus nenuphar***Plum webspinning sawfly**, see *Neurotoma inconspicua***Plusia** (Lep.: Noctuidae)

aereoides: *Voria ruralis*.

brassicae, see *Trichoplusia ni*.

californica, see *Autographa californica*.

"*chalcites*," see *Chrysodeixis calcites*.

gamma, see *Autographa gamma*.

ni, see *Trichoplusia ni*.

spp.: *Eucelatoria armigera*.

Plusiodonta (Lep.: Noctuidae)

compressipalpis: *Compsilura concinnata*.

Pococera (Lep.: Pyralidae)

scabridella: *Nemorilla pyste*.

Podisus (Hem. (Heter.): Pentatomidae)

maculiventris: *Hemyda aurata*, *Trichopoda lanipes*.

Polia (Lep.: Noctuidae)

acutermis: *Bonnetia comta*.

adjuncta: *Compsilura concinnata*, *Periscepsia helymus*.

¹ As this Catalog was being prepared, *Platysamia* was synonymized under *Hyalophora* by Ferguson, 1972, Fascicle 20.2B Saturniidae, in Dominick et al., "Moths of North America."

legitima: *Compsilura concinnata*, *Eusisyropa virilis*.

purpurissata: *Nowickia hispida*.

renigera, see *Lacinipolia renigera*.

Polydesma (Lep.: Noctuidae)

umbricola: *Chaetogaedia monticola*, *Eucelatoria armigera*,
Lespesia archippivora.

Polygonia (Lep.: Nymphalidae)

comma: *Compsilura concinnata*, *Euphorocera claripennis*,
Tachinidae.

interrogationis: *Compsilura concinnata*, *Euphorocera clari-*
pennis, *Eusisyropa blanda*, *E. virilis*, *Exorista mella*,
Lespesia aletiae.

Polyommatus boeticus, see *Lampides boeticus*

Polyphemus moth, see *Antheraea polyphemus*

Ponca (Orth: Gryllidae)

venosa: *Calodexia interrupta*.

Pontia rapae, see *Pieris rapae*

Popilius (Col.: Passalidae)

disjunctus: *Zelia vertebrata*.

Popillia (Col.: Scarabaeidae)

japonica: *Dexilla ventralis*, *Hamaxia incongrua*, *Hyperecteina*
aldrichi, *Prosenia siberita*, *Ptilodexia* sp.

Poplar borer, see *Saperda calcarata*

Poplar tentmaker, see *Ichthyura inclusa*

Porcellio (Isopoda (Crustacea): Oniscidae)

scaber: *Melanophora roralis*.

scaber probably: *Melanophora roralis*.

spp.: *Melanophora roralis*.

Porosagrotis orthogonia, see *Agrotis orthogonia*

Porosagrotis vetusta, see *Agrotis vetusta*

“**Porthesia**” *dispar*, see *Porthetria dispar*

Porthetria (Lep.: Lymantriidae)

dispar: *Blepharipa schineri*, *B. scutellata*, *B. sp.*, *Carcelia*
amplexa, *C. laxifrons*, *Compsilura concinnata*, *Drino*
inconspicua, *Euphorocera claripennis*, *Eusisyropa*
virilis, *Exorista mella*, *E. spp.*, *Lespesia aletiae*, *L.*
frenchii, *Nemorilla pyste*, *Parasetigena agilis*, *Phryxe*
vulgaris, *Tachinomyia nigricans*, *T. variata*, *Townsend-*
iellomyia nidicola, Tachinidae.

Potato beetles, see Chrysomelid spp. (including *Lema trilineata*
and/or *Leptinotarsa decemlineata*)

Potato stalk borer, see *Trichobaris trinotata*

Priophorus (Hym.: Tenthredinidae)*morio*: *Bessa harveyi*, *B. selecta*.*rubivorus*, see *P. morio*.**Pristiphora** (Hym.: Tenthredinidae)*erichsonii*: *Bessa harveyi*, *B. selecta*, *B. selecta* probably, *B. spp.*, *Diplostichus lophyri*, *Drino bohémica*, *Lespesia archippivora*, "Phorocera" spp., *Spathimeigenia aurifrons*, *S. spinigera*.*erichsonii* probably: *Bessa harveyi*.*geniculata*: *Bessa harveyi*, *B. selecta*, *Compsilura concinnata*, *Diplostichus lophyri*, *Drino bohémica*, *Guerinia dydas*.*leechi*: *Bessa harveyi*.*lena*: *Bessa harveyi*.**Prochoerodes truxaliata**, see *Sabulodes truxaliata***Prodenia** (Lep.: Noctuidae)*autumnalis*, see *Spodoptera frugiperda*.*commelinae*, see *Spodoptera dolichos*.*dolichos*, see *Spodoptera dolichos*.*eridania*, see *Spodoptera eridania*.*latifascia*, see *Spodoptera latifascia*.*latifascia* probably, see *Spodoptera latifascia* probably.*ornithogalli*, see *Spodoptera ornithogalli*.*ornithogalli* probably, see *Spodoptera ornithogalli* probably."ornithogalli," see *Spodoptera ornithogalli*.*praefica*, see *Spodoptera praefica*.*sunia*: *Gonia crassicornis*, *Nemorilla pyste*.sp., see *Spodoptera* spp.**Prolimacodes** (Lep.: Limacodidae)*badia*: Tachinidae.**Promethea**, see *Calosamia promethea***Proteides clarus**, see *Epargyreus clarus***Proteides tityrus**, see *Epargyreus clarus***Proteopteryx bolliana**, see *Gretchena bolliana***Proteoteras** (Lep.: Olethreutidae)*willingana*: *Erynnia tortricis*.**Protitame** (Lep.: Geometridae)*virginalis*: *Chaetophlepsis orbitalis*.**Protoparce carolina**, see *Manduca sexta***Protoparce celeus**, see *Manduca quinquemaculata***Protoparce cingulata**, see *Agrius cingulatus***Protoparce jamaicensis**, see *Manduca sexta jamaicensis***Protoparce quinquemaculata**, see *Manduca quinquemaculata***Protoparce sexta**, see *Manduca sexta* or *Manduca sexta jamaicensis***Protoparce sexta jamaicensis**, see *Manduca sexta jamaicensis*

Protoparce sexta "jamaicensis," see *Manduca sexta jamaicensis*
Protoparce spp., see *Manduca* spp.

Protorthodes (Lep.: Noctuidae)

oviducta: *Wagneria vernata*.

rufula: *Chaetogaedia crebra*, *Eucnephalia* sp. probably.

Psara (Lep.: Pyralidae)

"*bispunctalis*," see *Herpetogramma bipunctalis*.

pallicaudalis: *Argyrophylax albincisa*, *Nemorilla pyste*.

periusalis, see *Pilemia periusalis*.

Pseudaletia (Lep.: Noctuidae)

inconspicua, see *Leucania inconspicua*.

latiuscula, see *Leucania latiuscula*.

macrosaris: *Chaetogaedia monticola*.

unipuncta: *Archytas apicifer*, *A. cirphis*, *A. marmoratus*, *A.*
 spp. (including *A. apicifer* and/or *A. californiae*), *A.* sp.
 (*A. apicifer* or *A. californiae*) probably, *Athrycia*
cinerea, *Carcelia* spp., *Chaetogaedia monticola*,
Compsilura concinnata, *Cylindromyia "carolinae," C.*
euchenor, *Eucelatoria armigera*, *E. rubentis*, *E.*
rubentis probably, *Euphorocera claripennis*, *E.*
floridensis, *Exorista mella*, *Gonia porca*, *Gymnocar-*
celia ricinorum, *Gymnoclytia immaculata*, *Lespesia*
aletiae, *L. archippivora*, *Madremyia saundersii*,
Myiophasia "aenea," M. metallica, *Patelloa leucaniae*,
Peleteria texensis, *P.* spp., *Periscepsia helymus*, *P.*
laevigata, *Phryxe pecosensis*, *P. vulgaris*, "Tachina"
 spp., *Triachora unifasciata*, *Winthemia datanae*, *W.*
leucanae, *W. leucanae* probably, *W. militaris*, *W.*
militaris probably, *W.* sp. near *militaris*, *W. osten-*
sackenii, *W. quadripustulata*, *W. quadripustulata*
 probably, *W. rufopicta*, *W.* spp. (including *W. leucanae*
 probably or *W. militaris*), *W.* spp., Tachinidae.

"*unipunctata*," see *P. unipuncta*.

Pseudexentera (Lep.: Olethreutidae)

cressoniana, see *P. improbana*.

improbana: *Cyzenis albicans*.

improbana oregonana: "*Cyzenis festinans*" probably, *C.* spp.,

Eulasiona comstocki.

oregonana, see *improbana oregonana*.

Pseudoplusia (Lep.: Noctuidae)

includens: *Chaetophlepsis eudryae*, *Euphorocera floridensis*,
Lespesia aletiae, *Winthemia intermedia*.

Pseudothyatira (Lep.: Thyatiridae)

cymatophoroides: Tachinidae.

"*Psilocorsis*" *fletcherella*, see *Psilocorsis fletcherella*

Psilocorsis (Lep.: Oecophoridae)

faginella: *Elfia* spp., *Eusisyropa blanda*, *Psalidopteryx psilocorsiphaga*.

fletcherella: *Eusisyropa blanda*.

quercicella: *Elfia* spp.

reflexella: *Elfia* sp. near *usitata*, *E.* spp., Tachinidae.

spp.: *Eulasiona comstocki*, *Psalidopteryx psilocorsiphaga*.

Psoloessa (Orth.: Acrididae)

delicatula: *Ceracia dentata* probably.

Psychomorpha (Lep.: Noctuidae)

epimenis: *Sisyropa alypiae*, "*Zenillia*" spp.

"*epimensis*," see *P. epimenis*.

Pteronidea coryla, see *Nematus erythrogaster*

Pteronidea ribesii, see *Nematus ribesii*

Pterophoridae (Lep.: Pterophoridae)

Unidentified: *Eribella* spp.

Pterophorus (Lep.: Pterophoridae)

periscelidactylus: *Hemisturmia tortricis*.

Pterostichus (Col.: Carabidae)

lucublandus: Tachinidae.

melanarius: Tachinidae.

Pulicalvaria piceaella, see *Coleotechnites piceaella*

Pumpkin bug, see *Nezara viridula*

Puss caterpillar, see *Megalopyge opercularis*

Pyralid, see *Hellula rogatalis*

Pyralidae (Lep.: Pyralidae)

Unidentified: *Drino austrina*, *Hyphantrophaga hyphantriae*, *Leskiomima tenera*, *Lespesia frenchii*, *L.* spp., *Lixophaga variabilis*, *Metopiops pyralidis*, *Nemorilla pyste*.

Pyralidae or Tortricidae (Lep.: Pyralidae or Tortricidae)

Unidentified: *Coronimya geniculata*.

Pyralis (Lep.: Pyralidae)

farinalis: *Winthemia leucanae*.

farinalis probably: *Elfia tarsalis*.

Pyrameis atalanta, see *Vanessa atalanta*

Pyrameis cardui, see *Vanessa cardui*

Pyrameis "caryae," see *Vanessa carye*

Pyrameis carye, see *Vanessa carye*

Pyrausta (Lep.: Pyralidae)

ainsliei, see *Ostrinia obliteralis*.

ainsliei or *P. penitalis*, see *Ostrinia* spp. (including *O. obliteralis* and/or *O. penitalis*).

futilalis: *Elfia* sp. near *aenea*, *Hyphantrophaga euchaetiae*, *Lixophaga plumbea*, *Nemorilla pyste*, *Pseudoperichaeta erecta*, Tachinidae.

"*nubilalis*," see *Ostrinia nubilalis*.

nubilalis, see *Ostrinia nubilalis*.

"*nubililalis*," see *Ostrinia nubilalis*.

penitalis, see *Ostrinia penitalis* or *Ostrinia* spp. (including *O. obliteralis* and/or *O. penitalis*).

pertextalis, see *Macrobotys pertextalis*.

signatalis: *Lespesia* spp., *Nemorilla pyste*, *Pseudoperichaeta erecta*.

tyralis: *Nemorilla pyste*, *Stomatomyia floridensis*.

"*vubilalis*," see *Ostrinia nubilalis*.

spp.: *Aplomya caesar*, *Pyraustomyia penitalis*.

spp. probably: *Dejeaniopalpus texensis*.

Pyreferra (Lep.: Noctuidae)

hesperidago: *Exorista mella*, *Ictericophyto spinosa*, *Omotoma fumiferanae*, Tachinidae.

Pyrophila (Lep.: Noctuidae)

pyramidoides, see *Amphipyra pyramidoides*.

sp.: *Leschenaultia americana*.

Pyrrhalla (Col.: Chrysomelidae)

luteola: *Aplomyiopsis galerucellae*, *Erynniopsis rondanii*.

Pyrrharctia isabella, see *Isia isabella*

Ragweed borer, see *Epiblema strenuanum*

Range caterpillar, see *Hemileuca oliviae*

Range crane fly, see *Tipula simplex*

Raphia (Lep.: Noctuidae)

frater: *Tachinomyia* spp.

Raspberry cane borer, see *Oberea bimaculata*

Recurvaria coniferella, see *Coleotechnites coniferella*

Recurvaria thujella, see *Coleotechnites thujella*

Red admiral, see *Vanessa atalanta*

Redbacked cutworm, see *Euxoa ochrogaster*

Redbanded leafroller, see *Argyrotaenia velutinana*

Redheaded pine sawfly, see *Neodiprion lecontei*

Redhumped caterpillar, see *Schizura concinna*

Redlegged grasshopper, see *Melanoplus femurrubrum*

Red pine sawfly, see *Neodiprion nanulus nanulus*

Reduviolus ferus, see *Nabis americanoferus*

Reduviolus roseipennis, see *Nabis roseipennis*

Regal moth, see *Citheronia regalis*

- “*Remigera*” *repanda*, see *Mocis* spp. (including *M. latipes* and/or *M. repanda*)
- Remigia repanda*, see *Mocis latipes* probably or *Mocis* spp. (including *M. latipes* and/or *M. repanda*)
- Retinia* sp., see *Rhyacionia* spp.
- “*Rhabdocnemis*” *obscurus*, see *Rhabdoscelus obscurus*
- Rhabdocnemis obscura*, see *Rhabdoscelus obscurus*
- Rhabdoides cellus*, see *Autochton cellus*
- Rhabdophaga* (Dipt.: Cecidomyiidae)
salicisbrassicoides: Tachinidae.
- Rhabdoscelus* (Col.: Curculionidae)
obscurus: *Lixophaga sphenophori*, *Metadexia tricolor*.
- Rhagium inquisitor*, see *Stenocorus inquisitor inquisitor*
- Rhagium lineatum* probably, see *Stenocorus inquisitor lineatus* probably
- “*Rhodoabaemus*” sp. in dahlia stem, see *Rhabdoscelus obscurus*
- Rhodoabaenus* (Col.: Curculionidae)
tredecimpunctatus: Tachinidae.
13-punctatus, see *R. tredecimpunctatus*.
- Rhodophora* (Lep.: Noctuidae)
florida: *Compsilura concinnata*.
- Rhopobota* (Lep.: Olethreutidae)
naevana: *Nemorilla pyste*.
vacciniana, see *R. naevana*.
- Rhyacionia* (Lep.: Olethreutidae)
buoliana: *Erynnia tortricis*, *Lixophaga mediocris*, *Nemorilla pyste*.
frustrana: *Lixophaga mediocris*, *L. plumbea*, Tachinidae.
frustrana bushnelli: Tachinidae.
pasadenana: *Urophylloopsis retiniae*.
rigidana: *Lixophaga mediocris*, *L. variabilis*.
spp.: *Urophylloopsis retiniae*.
sp. probably: *Urophylloopsis retiniae*.
- Rhynchophora* (Col.: Unidentified Coleoptera)
Miscellaneous: *Lixophaga* spp.
- Rice stalk borer, see *Chilo plejadellus*
- Rice stink bug, see *Oebalus pugnax*
- Rocky Mountain grasshopper, see *Melanoplus spretus*
- Romaleum atomarius*, see *Enaphalodes atomarius*
- Rothschildia* (Lep.: Saturniidae)
aroma probably: *Belvosia nigrifrons*, *Lespesia* spp.
orizaba: *Belvosia formosa*, *B. nigrifrons*.
sp.: *Belvosia nigrifrons*.
- Rusty tussock moth, see *Orgyia antiqua*

“Sabine” *apicalis*, see *Sibine apicalis*

Sabulodes (Lep.: Geometridae)

transversata: Tachinidae.

truxaliata: *Xanthophyto* spp.

Saddleback caterpillar, see *Sibine stimulea*

Saddled prominent, see *Heterocampa guttivitta*

Salebria *semiobscura*, see *Nephopteryx subfuscella*

Salebria *subcaesiella*, see *Nephopteryx subcaesiella*

Saltmarsh caterpillar, see *Estigmene acrea*

Samia (Lep.: Saturniidae)

cecropia, see *Platysamia cecropia*.

columbia, see *Hyalophora columbia*.

cynthia: *Lespesia frenchii*, *Winthemia tricolor*.

cynthia advena: *Lespesia frenchii*, *Winthemia datanae* probably.

rubra, see *Platysamia euryalus*.

walkeri advena, see *S. cynthia advena*.

Saperda (Col.: Cerambycidae)

calcarata: *Ptilodexia canescens*.

obliqua: Tachinidae.

sp.: *Euphorocera floridensis*.

Sarothrips *revayanus*, see *Nycteola* spp. (including *N. cinereana*,
N. columbiana and/or *N. frigidana*)

Satin moth, see *Stilpnotia salicis*

Saturnia (Lep.: Saturniidae)

cecropia, see *Platysamia cecropia*.

galbina: *Euphorocera* sp. near *floridensis*.

maia, see *Hemileuca maia*.

orizaba, see *Rothschildia orizaba*.

Saturniidae (Lep.: Saturniidae)

Unidentified: *Compsilura concinnata*.

Sawfly(-flies) (Hym.: Unidentified Hymenoptera)

Sawfly(-flies): *Athrycia cinerea*, *Compsilura concinnata*, *Drino bohemica*, *Euphorocera claripennis*, *Guerinia dydas*, *G. rufostomata*, *Lixophaga variabilis*, Gen. and sp. near *Lixophaga*, *Spathimeigenia hylotomae*, *S. spinigera*.

Sawfly(-flies) probably: *Spathimeigenia aurifrons*.

Say's plant bug, see *Chlorochroa sayi*

Say stink bug, see *Chlorochroa sayi*

Scarabaeid (Col.: Scarabaeidae)

spp. (including *Cyclocephala immaculata* or *Phyllophaga* spp.):
Microphthalma disjuncta.

spp. (including *Cyclocephala* spp. and/or *Phyllophaga* spp.):
Microphthalma disjuncta.

spp. (including *Phyllophaga* spp. or *Phytalis* spp.): *Cryptomei-
genia aurifacies*, *Eutrixoides jonesii*.

- spp. (including *Phyllophaga* spp. or other Scarabaeidae):
Mochlosoma lacertosum.
- spp. (allied genera of *Phyllophaga* spp.): *Microphthalma disjuncta*.
- Scarabaeidae (Col.: Scarabaeidae)
Unidentified: *Hamaxia incongrua*.
- Schinia (Lep.: Noctuidae)
brevis: *Winthemia quadripustulata*.
- Schistocerca (Orth.: Acrididae)
cancellata: *Oedematocera dampfi*.
paranensis, see *S. cancellata*.
- Schizocerus *ebenus*, see *Sphacophilus cellularis*
- Schizocerus *privatus*, see *Sphacophilus cellularis*
- Schizocerus sp., see *Sterictiphora* sp.
- Schizura (Lep.: Notodontidae)
badia: *Compsilura concinnata*, *Winthemia datanae*.
"cincinna," see *S. concinna*.
"concina," see *S. concinna*.
concinna: *Actia americana*, *A. interrupta*, *Carcelia laxifrons*,
Compsilura concinnata, *Euphorocera claripennis*,
Exorista mella, *Lespesia archippivora*, *Pseudoperi-
chaeta erecta*, *Winthemia datanae*, *W. quadripustulata*,
Tachinidae.
ipomaeae: *Compsilura concinnata*, *Lespesia schizurae*, *Winthe-
mia datanae*, Tachinidae.
"ipomoeae," see *S. ipomaeae*.
leptinoides: *Winthemia datanae*.
unicornis: *Compsilura concinnata*, *Euphorocera claripennis*,
Lespesia schizurae, *Winthemia datanae*, *W. sinuata*,
Tachinidae.
spp.: *Winthemia* spp.
- Sciaphila (Lep.: Olethreutidae)
duplex: *Ceromasia auricaudata*, *Lypha intermedia*, *Madremyia
saundersii*, *Xanthophyto labis*.
- Scoliopteryx (Lep.: Noctuidae)
libatrix: *Compsilura concinnata*, *Winthemia borealis*.
- Scotogramma (Lep.: Noctuidae)
trifolii: *Euphorocera claripennis*, *Lespesia archippivora*.
trifolii probably: *Gonia* spp.
- Scotorythra (Lep.: Geometridae)
paludicola probably: *Chaetogaedia monticola*, *Lespesia
archippivora*.
rara: *Chaetogaedia monticola*, *Lespesia archippivora*.
trapezias: *Chaetogaedia monticola*.

Scythris (Lep.: Scythridae)*epilobiella*: *Psalidopteryx macdunnoughi*.sp.: *Psalidopteryx macdunnoughi*.**Sehirus** (Hem. (Heter.): Cydnidae)*cinctus*: *Hyalomya purpurascens*.**Seirarctia** (Lep.: Arctiidae)*echo*: *Lespesia aletiae*.**Selandria barda**, see *Tethida cordigera***Semiothisa** (Lep.: Geometridae)*bisignata*: *Blondelia eufitchiae*, *Chaetophlepsis semiothisae*.*granitata*, see *S. granitata* complex.*granitata* complex: *Bessa harveyi*, *Blondelia eufitchiae*,
Chaetophlepsis semiothisae.*minorata*: *Blondelia eufitchiae*, *Chaetophlepsis semiothisae*.*neptaria*: *Euphorocera claripennis*.*orillata*: *Blondelia eufitchiae*.*oweni*: *Blondelia eufitchiae*, *Chaetophlepsis semiothisae*.*punctolineata*: *Prophryno marginalis*.*sexmaculata*: *Blondelia eufitchiae*, *Chaetophlepsis semiothisae*.**Septis apamiformis**, see *Apamea apamiformis***Serica** (Col.: Scarabaeidae)*iricolor*: *Pseudomyothyria ancilla*.*serotina*: *Sitophaga neomexicana*.**Sesia acerni**, see *Sylvora acerni***Sesia rutilans**, see *Synanthedon bibionipennis***"Shizocera" ebenus**, see *Sphacophilus cellularis***"Shizura" sp.**, see *Schizura* spp.**Sibine** (Lep.: Limacodidae)*apicalis*: *Lespesia aletiae*.*stimulea*: *Anaporia pristis*, *Euphorocera claripennis*, *Palpexorista coccyx*, Tachinidae.**Sidemia devastator**, see *Crymodes devastator***Sideridis** (Lep.: Noctuidae)*maryx*: *Euexorista futilis*.**Silkworm**, see *Bombyx mori***Silverspotted skipper**, see *Epargyreus clarus***Silverspotted tiger moth**, see *Halisidota argentata***Simyra** (Lep.: Noctuidae)*henrici*: *Compsilura concinnata*, *Eusisyropa virilis*, *Exorista mella*, *Lespesia dimmocki*, *L. spp.*, "Masicera" spp.**Sisyrosea** (Lep.: Limacodidae)*textula*: *Anaporia limacodis*, *A. pristis*, *A. sp.***Sitona** (Col.: Curculionidae)*hispidula*: *Hyalomyodes triangulifer*.

Small milkweed bug, see *Lygaeus kalmii*

Smearred dagger moth, see *Acronicta oblinita*

Smerinthus (Lep.: Sphingidae)

cerisyi: *Compsilura concinnata*, *Lespesia flavifrons*, *L. frenchii*.
geminatus, see *S. jamaicensis*.

jamaicensis: *Compsilura concinnata*, *Eusisyropa blanda*, *Lespesia flavifrons*, *L. frenchii*, Tachinidae.

ophthalmicus: "Prosopaea" spp.

sp.: *Lespesia flavifrons*.

Sonoran tent caterpillar, see *Malacosoma tigris*

Sorghum webworm, see *Celama sorghiella*

Southern armyworm, see *Spodoptera eridania*

Southern beet webworm, see *Herpetogramma bipunctalis*

Southern cabbageworm, see *Pieris protodice*

Southern corn rootworm, see *Diabrotica undecimpunctata howardi*

Southern cornstalk borer, see *Diatraea crambidoides*

Southern green stink bug, see *Nezara viridula* or *Nezara viridula smaragdula*

Southern pine sawyer, see *Monochamus titillator*

Southwestern corn borer, see *Diatraea grandiosella*

Southwestern tent caterpillar, see *Malacosoma incurvum*

Soybean looper, see *Pseudoplusia includens*

Spaelotis (Lep.: Noctuidae)

clandestina: *Periscepsia rohweri*, *Winthemia rufopicta*.

Sparganothis (Lep.: Tortricidae)

distincta: *Blondelia eufitchiae*.

pettitana: *Lypha* sp., Tachinidae.

sulfureana: *Erynnia tortricis*, *Nemorilla pyste*.

"*Sparganothis sulfurana*," see *Sparganothis sulfureana*

Speyeria (Lep.: Nymphalidae)

cybele: *Madremyia saundersii*.

Sphacophilus (Hym.: Argidae)

cellularis: "*Hypostena floridensis*," *Schizocerophaga leiby*,
Tachinidae.

Spharagemon (Orth.: Acrididae)

collare: Tachinidae.

Sphecodina (Lep.: Sphingidae)

"*abbottii*," see *S. abbottii*.

abbottii: *Compsilura concinnata*, *Eusisyropa blanda*.

Sphenophorus (Col.: Curculionidae)

incurrens: *Myiophasia metallica*, Tachinidae.

parvulus: *Myiophasia "aenea."*

pontederiae: *Lixophaga variabilis*.

robustus: *Myiophasia robusta*.

Sphingicampa bicolor, see *Syssphinx bicolor*

Sphingid on *Populus trichocarpa*, see *Pachysphinx modesta*

Sphingidae (Lep.: Sphingidae)

Unidentified: *Compsilura concinnata*, *Drino balloui*, *D. inca*, *D. incompta*, *D. rhoeo*, *Leschenaultia leucophrys*, *Lespesia frenchii*, *Winthemia quadripustulata*, *W. rufopicta*, "Zenillia" spp.

Sphinx (Lep.: Sphingidae)

carolina, see *Manduca sexta*.

chersis: *Compsilura concinnata*, *Drino incompta*, *Euphorocera claripennis*, *Leschenaultia leucophrys*, *Lespesia frenchii*, Tachinidae.

convolvuli, native species of the same group as, see *Agrius convolvuli*.

drupiferarum: *Compsilura concinnata*, *Winthemia datanae*, Tachinidae.

gallii intermedia, see *Hyles gallii*.

gordius: *Compsilura concinnata*, *Winthemia datanae*, Tachinidae.

gordius probably: *Winthemia quadripustulata*.

kalmiae: *Lespesia frenchii*, Tachinidae.

lineata, see *Hyles lineata*.

luscitiosa: *Metaplagia occidentalis*.

spp., see Sphingidae, unidentified or *S.* spp.

spp.: *Lespesia archippivora* probably, *Winthemia deilephilae* probably.

Sphinx moth(-s) (Lep.: Sphingidae)

Sphinx moth(-s): *Drino balloui*, *D. rhoeo*, *D.* sp.

"Sphynx" moth, see Sphinx moth(-s)

Spicebush swallowtail, see *Papilio troilus*

Spilonota (Lep.: Olethreutidae)

ocellana: *Actia diffidens*, *Blondelia eufitchiae*, *Mericia* spp., *Nemorilla pyste*, *Pseudoperichaeta erecta*, *Winthemia quadripustulata*.

Spilosoma acrea, see *Estigmene acrea*

Spilosoma virginica, see *Diacrisia virginica*

Spinach flea beetle, see *Disonycha xanthomelas*

Spined soldier bug, see *Podisus maculiventris*

Spodoptera (Lep.: Noctuidae)

dolichos: *Winthemia quadripustulata*, Tachinidae.

eridania: *Choeteprosopa hedemanni*, *Euphorocera claripennis*, *Gonia crassicornis*, *Winthemia quadripustulata*, *W. rufopicta*, Tachinidae.

exempta: *Archytas cirphis*, *Chaetogaedia monticola*, *Eucelatoria armigera*, *Lespesia archippivora*.

exigua: *Chaetogaedia monticola*, *Eucelatoria armigera*, *E. rubentis*, *Gonia crassicornis*, *Lespesia archippivora*.

frugiperda: *Admontia degeerioides*, *Archytas apicifer*, *A. incertus*, *A. marmoratus*, *A. metallicus*, *A. spp.* (including *A. apicifer* and/or *A. californiae*), *Bonnetia comta*, *Carcelia amplexa*, *Eucelatoria armigera*, *Euphorocera claripennis*, *E. floridensis*, *Eusisyropa blanda*, *Gonia crassicornis*, *G. sequax*, *G. texensis*, *G. spp.*, *Hyphanthropaga hyphantriae*, *Lespesia aletiae*, *L. archippivora*, *L. archippivora* probably, *L. spp.*, *Lixophaga diatraeae*, *Nemorilla pyste*, *Palexorista laxa*, *Voria ruralis*, *Winthemia abdominalis*, *W. infesta*, *W. leucanae*, *W. quadripustulata*, *W. rufopicta*, *W. sinuata*, Tachinidae.

latifascia: *Archytas marmoratus*.

latifascia probably: *Archytas marmoratus*.

mauritiae: *Archytas cirphis*, *Chaetogaedia monticola*, *Eucelatoria armigera*, *Lespesia archippivora*.

mauritiae acronyctoides, see *S. mauritiae*.

ornithogalli: *Archytas spp.* (including *A. apicifer* and/or *A. californiae*), *Choeteprosopa hedemanni*, *Euphorocera omissa*, *E. tachinomoides*, *Lespesia aletiae*, *L. archippivora*, *Omotoma fumiferanae*, *Winthemia quadripustulata*, *W. rufopicta*, *W. spp.*, Tachinidae.

ornithogalli probably: *Archytas spp.* (including *A. apicifer* and/or *A. californiae*).

praefica: *Archytas californiae*, *Eucelatoria armigera*, *Euphorocera claripennis*, *Lespesia archippivora*.

spp.: *Archytas spp.*, *Lespesia archippivora*.

Spotted beet webworm, see *Hymenia perspectalis*

Spotted cucumber beetle, see *Diabrotica undecimpunctata howardi*

Spotted cutworm, see *Amathes c-nigrum*

Spotted tussock moth, see *Halisidota maculata*

Spring cankerworm, see *Paleacrita vernata*

Spruce budworm, see *Choristoneura fumiferana*

Spruce coneworm, see *Dioryctria reniculella*

Spruce sawfly, see *Diprion hercyniae* or *Diprion polytomum*

Squash beetle, see *Epilachna borealis*

Squash borer moth, see *Melittia cucurbitae* probably

Squash bug, see *Anasa tristis*

Squash vine borer, see *Melittia cucurbitae*

Stagmomantis (Orth.: Mantidae)

carolina: Tachinidae.

Stalk borer, see *Papaipema nebris*

Stenocorus (Col.: Cerambycidae)

inquisitor, see *S. inquisitor inquisitor*.

inquisitor inquisitor: *Eutheresia nipigonensis*, *E. satisfacta* probably, *E. trivittata*.

inquisitor lineatus probably: *Ptilodexia canescens*.

Stenoma (Lep.: Stenomidae)

algidella, see *Antaeotricha leucillana*.

spp.: *Lixophaga stenomae*.

Stenopelmatus (Orth.: Gryllacrididae)

pictus: *Stomatomyia parvipalpis*.

Sterictiphora (Hym.: Argidae)

cellularis, see *Sphacophilus cellularis*.

collaris, see *Sphacophilus cellularis*.

sp.: "*Hypostena floridensis*."

Stibadium (Lep.: Noctuidae)

spumosum: *Lixophaga variabilis*.

Stilpnotia (Lep.: Lymantriidae)

salicis: *Blepharipa scutellata*, *Compsilura concinnata*, *Euphorocera claripennis*, *Eusisyropa blanda*, *Exorista mella*, *Lespesia aletiae*, *L. frenchii*, *Parasetigena agilis*, *Tachinomyia similis*, *T. spp.*, *Winthemia quadripustulata*, Tachinidae.

Strawberry crown moth, see *Synanthedon bibionipennis*

Strawberry leafroller, see *Ancylis comptana fragariae*

Striped alder sawfly, see *Hemichroa crocea*

Striped cucumber beetle, see *Acalymma vittata*

Striped garden caterpillar, see *Polia legitima*

Strongylogaster (Hym.: Tenthredinidae)

sp.: *Bessa harveyi*.

Strymon (Lep.: Lycaenidae)

calanus: *Aplomya theclarum*.

falacer: *Aplomya theclarum*.

melinus: *Aplomya theclarum*, *Lespesia spp.*

ontario autolytus: *Nemorilla pyste*.

Sugarbeet crown borer, see *Hulstia undulatella*

Sugarbeet webworm, see *Loxostege sticticalis*

Sugarcane beetle, see *Euetheola rugiceps*

Sugarcane borer, see *Diatraea saccharalis*

Sugarcane leafroller, see *Hedylepta accepta*

Sugarcane moth borer, see *Diatraea saccharalis*

Suleima (Lep.: Olethreutidae)

helianthana: *Clausicella floridensis*, *Lixophaga mediocris*, *L. variabilis*.

- Sunflower moth, see *Homoeosoma electellum*
 Swaine jack pine sawfly, see *Neodiprion swainei*
 Sweetpotato hornworm, see *Agrius cingulatus*
 Sweetpotato leafroller, see *Pilocrocis tripunctata*
 Sycamore tussock moth, see *Halisidota harrisii*
Sylepta helcitalis, see *Asciodes anormalis* probably
Sylvora (Lep.: Aegeriidae)
 acerni: *Bonnetia* sp., *Copecrypta* sp.
Symmerista (Lep.: Notodontidae)
 albifrons: *Blepharipa scutellata*, *Compsilura concinnata*,
 Winthemia spp., Tachinidae.
 canicosta: *Compsilura concinnata*, *Euphorocera edwardsii*.
Synanthedon (Lep.: Aegeriidae)
 bibionipennis: *Parafischeria eucerata*, *Sipholeskia occidentalis*.
Synchloe lacinia, see *Chlosyne lacinia*
Synchlora (Lep.: Geometridae)
 rubrifrontaria, see *S.* spp. (including *S. aerata* and/or *S.*
 liquoraria albolineata).
 spp. (including *S. aerata* and/or *S. liquoraria albolineata*):
 Phryxe pecosensis probably.
 "Synclora" *rubrifrontaria*, see *Synchlora* spp. (including *S. aerata*
 and/or *S. liquoraria albolineata*)
Syntomeida (Lep.: Ctenuchidae)
 epilais: *Euphorocera floridensis*, *Lespesia aletiae*.
 epilais jucundissima: *Lespesia* spp.
Syssphinx (Lep.: Saturniidae)
 bicolor: *Lespesia frenchii*.
 Tabanid larva, see *Tabanus trimaculatus*
 Tabanidae (Dipt.: Tabanidae)
 Unidentified: *Opsotheresia bigelowi*, Tachinidae.
Tabanus (Dipt.: Tabanidae)
 fulvicallus: *Carinosillus tabanivorus*.
 fumipennis: *Phasiops flavus* probably.
 lasiophthalma, see *Hybomitra lasiophthalma*.
 marginalis: *Carinosillus tabanivorus*.
 petiolatus: *Phasiops flavus*.
 petiolatus probably: *Phasiops flavus*.
 reinwardtii: *Carinosillus tabanivorus*.
 reinwardtii probably: *Vibrissotheresia pechumani*.
 trimaculatus: *Carinosillus tabanivorus*, *Phasiops flavus*,
 "Phorostoma" spp.
 trimaculatus probably: *Phasiops flavus*.
 spp.: *Ormia punctata*, *Phasiops flavus*.
Taeniocampa rufula, see *Protorthodes rufula*

- Tarnished plant bug, see *Lygus lineolaris*
- Telea polyphemus*, see *Antheraea polyphemus*
- “*Telphusa*” (Lep.: Gelechiidae)
 sp. probably: *Psalipteryx slossonae*.
- Tenodera (Orth: Mantidae)
australasiae: *Trichopoda pennipes* probably.
- Tent caterpillar(-s) (Lep.: Lasiocampidae)
 Tent caterpillar(-s): *Archytas* spp. (including *A. apicifer* and/or *A. californiae*), *Exorista mella*, *Leschenaultia exul*, *Lespesia archippivora*, *L. frenchii*, *Winthemia rufopicta*.
- Tenthredinidae (Hym.: Tenthredinidae)
 Unidentified: *Bessa harveyi*, *Carcelia malacosomae*, “*Gymnostylia*” sp., *Lixophaga variabilis*, *Pseudomyothyria* spp.
- Tenthredinoidea (Hym.: Unidentified Hymenoptera)
 Tenthredinoidea: *Compsilura concinnata*.
- Tenthredo sp., see Sawfly(-flies)
- Tephroclystis *absynthiata*, see *Eupithecia coagulata* probably
- Teras minuta* var. *cinderella*, see *Acleris minuta*
- Tethida (Hym.: Tenthredinidae)
cordigera: Tachinidae.
- Tetralopha (Lep.: Pyralidae)
asperatella: *Eufrontina spectabilis*, *Metopiops* sp. near *pyralidis*, *Nemorilla pyste*.
 sp. near *asperatella*: *Leskiomima tenera*.
robustella: *Eufrontina spectabilis*, *Eusisyropa blanda*.
scabridella, see *Pococera scabridella*.
scortealis: *Frontiniella parancilla*.
subcanalis: *Genea aurea*, Tachinidae.
- Tetraonyx (Col.: Meloidae)
quadrifasciata: *Gonia frontosa*.
- Tettigoniidae (Orth.: Tettigoniidae)
 Unidentified: *Calodexia fumosa*.
- Thanaos brizo*, see *Erynnis brizo*
- Thanaos* spp., see *Erynnis* spp.
- Thanasimus* (Col.: Cleridae)
dubius: Tachinidae.
- Thecla autolycus*, see *Strymon ontario autolycus*
- Thecla calanus*, see *Strymon calanus*
- Thecla inornata*, see *Strymon falacer*
- Thecla* “*inornata*,” see *Strymon falacer*
- Thera* (Lep.: Geometridae)
juniperata: *Blondelia eufitchiae*.

- procteri*, see *T. juniperata*.
- Thistle butterfly, see *Vanessa cardui*
- Threelined leafroller, see *Pandemis limitata*
- Thyanta** (Hem. (Heter.): Pentatomidae)
accerra: *Trichopoda pennipes*.
perditor: *Trichopoda pennipes*.
- Thylochromus** (Hem. (Heter.): Lygaeidae)
nitidulus: *Catharosia lustrans*.
- Thymelicus** (Lep.: Hesperidae)
lineola: *Compsilura concinnata*, *Euexorista futilis*, *Helioplagia slossonae*, *Phryxe pecosensis*, *Winthemia militaris*, *W. sinuata*, Tachinidae.
- Thyridopteryx** (Lep.: Psychidae)
"ephemerae formis," see *T. ephemeraeformis*.
ephemeraeformis: *Archytas aterrimus*, *Drino rhoeo*, *Erynnia tortricis*, *Euphorocera claripennis*, *Eusisyropa blanda*, *E. virilis*, *Patelloa leucaniae*, "*Tachina*" spp., Tachinidae.
"ephemeroeformis," see *T. ephemeraeformis*.
meadii: *Euphorocera tachinomoides*.
- Tibicen septendecim**, see *Magisicada septendecim*
- Tiger swallowtail, see *Papilio glaucus*
- Tigrioides** (Lep.: Arctiidae)
bicolor: *Pseudochaeta argentifrons*.
- Tineidae** (Lep.: Tineidae)
 Unidentified: *Schizactia vitinervis*, "*Tachinophyto*" spp.
- Tipula** (Dipt.: Tipulidae)
abdominalis: *Admontia degeerioides*, *A. pergandei*.
caloptera: *Siphona geniculata*.
cunctans: *Admontia pergandei*.
infuscata, see *T. cunctans*.
paludosa: *Siphona geniculata*.
simplex: Tachinidae.
 [sp.], see Tipulidae, unidentified.
- Tipulidae** (Dipt.: Tipulidae)
 Unidentified: *Dexia limata*, *Pilatea arator*.
- Tipulini** (Dipt.: Tipulidae)
 Unidentified: "*Sturmia tipulensis*."
- Titanio** (Lep.: Pyralidae)
belialis: *Eucoronimyia hastata*.
- Tlascala** (Lep.: Pyralidae)
reductella: *Aphria ocypterata*.
- Tobacco budworm, see *Heliothis virescens*
 [Tobacco] hornworm, see *Manduca sexta*

Tolyte (Lep.: Lasiocampidae)

laricis: *Exorista mella*, *Winthemia sinuata*.

velleda: *Exorista mella*.

Tomato fruitworm, see *Heliothis zea*

Tomato hornworm, see *Manduca quinquemaculata*

Tomostethus (Hym.: Tenthredinidae)

multicinctus: *Hyalurgus* sp.

Tortricid (Lep.: Tortricidae)

spp. (including *Amorbia humerosana*, *Archips argyrospilus*, *A. semiferanus*, *Argyrotaenia quadrifasciana*, *Choristoneura fractivittana*, *C. rosaceana* and/or *Pandemis limitata*): *Aplomya caesar*, *Pseudoperichaeta erecta*.

spp. (including *Archips* spp. and/or *Choristoneura* spp.): *Aplomya caesar*.

Tortricidae (Lep.: Tortricidae)

Unidentified: *Actia interrupta*, *Bessa harveyi*, *Blondelia eufitchiae*, *Erynnia tortricis*, *Eusisyropa blanda*, *Hemisturmia tortricis*, *Nemorilla pyste*.

Unidentified probably: *Erynnia tortricis*.

Tortricidia (Lep.: Limacodidae)

flexuosa: *Anaporia limacodis*.

Tortrix albicomana probably, see *Croesia albicomana* probably

Tortrix alleniana, see *Aphelia alleniana*

Tortrix citrana, see *Argyrotaenia citrana*

Tortrix "fum-ferana," see *Choristoneura fumiferana*

Tortrix fumiferana, see *Choristoneura fumiferana*

Tortrix packardianus, see *Archippus packardianus*

Tortrix pallorana, see *Xenotemna pallorana*

Tortrix "parckardiana," see *Archippus packardianus*

Tortrix spp., see *Archips* spp. and/or *Choristoneura* spp., or Tortricidae, unidentified

Trachea turbulenta, see *Phosphila turbulenta*

Trichiocampus (Hym.: Tenthredinidae)

irregularis: *Bessa harveyi*.

viminalis: *Bessa harveyi*, *Compsilura concinnata*.

sp.: *Bessa harveyi*.

Trichiosoma (Hym.: Cimbicidae)

sp.: *Aplomya trichiosomae*.

Trichobaris (Col.: Curculionidae)

compacta: *Myiophasia "aenea."*

trinotata: *Myiophasia globosa*.

Trichoplusia (Lep.: Noctuidae)

ni: *Archytas californiae*, *Compsilura concinnata*, *Eucelatoria armigera*, *Eusisyropa autographae*, *Lespesia aletiae*, *L.*

archippivora, *L.* spp., *Lixophaga variabilis*, *Madremyia saundersii*, *Nemorilla pyste*, *Periscepsia helymus*, "Phorocera" spp., *Phryxe vulgaris*, *Siphona plusiae*, *S.* spp., "Tachina" spp., *Voria ruralis*, *V. ruralis* probably, *Winthemia quadripustulata*, *W. rufopicta*, Tachinidae.

Trichoptilus (Lep.: Pterophoridae)

lobidactylus: Tachinidae.

Trichostiba parvula, see *Urodus parvulus*

Trirhabda (Col.: Chrysomelidae)

bacharidis: *Celatoria diabroticae*.

Tropaea luna, see *Actias luna*

True armyworm, see *Pseudaletia unipuncta*

Tuber moth(-s) (Lep.: Unidentified Lepidoptera)

Tuber moth(-s): *Lydellohoughia tennella*.

Turuptiana (Lep.: Arctiidae)

permaculata: *Carcelia malacosomae*, *C. malacosomae* probably, *Chaetona* sp., "Nemoraea" sp., *Thelaira* spp.

Tussock moth(-s) (Lep.: Lymantriidae)

Tussock moth(-s): *Carcelia amplexa*, *Compsilura concinnata*, *Euphorocera claripennis*, *Exorista mella*, *E.* spp., *Lespesia aletiae*, *Winthemia quadripustulata*, Tachinidae.

Tussock moth(-s) probably: *Thelymyia mathesoni*.

12-spotted squash beetle, see *Diabrotica undecimpunctata undecimpunctata*

Twostriped grasshopper, see *Melanoplus bivittatus*

Twostriped walkingstick, see *Anisomorpha buprestoides*

Tyria (Lep.: Arctiidae)

jacobaeae: *Lespesia archippivora*.

Udea (Lep.: Pyralidae)

rubigalis: "Phorocera parva."

Uglynest caterpillar, see *Archips cerasivoranus*

Unicorn caterpillar, see *Schizura unicornis*

Uranotes melinus, see *Strymon melinus*

Urbanus (Lep.: Hesperidae)

proteus: *Chrysotachina alcedo*, *Lespesia aletiae*, *Nemorilla pyste*.

Urodus (Lep.: Yponomeutidae)

parvulus: *Euceromasia floridensis*, *Eusisyropa boarmiae*, Tachinidae.

Utetheisa (Lep.: Arctiidae)

bella, see *U. ornatrix bella*.

ornatrix bella: *Euphorocera claripennis*, *Gymnocarcelia ricinorum*, *Lespesia aletiae*, Tachinidae.

- venusta*: *Lixophaga diatraeae*.
- “*Utethesia*” *bella*, see *Utetheisa ornatrix bella*
- Vagabond crambus, see *Crambus vulgivagellus*
- Valentinia (Lep.: Blastobasidae)
- glandulella*: “*Hypostena floridensis*.”
- “*Vanassa*” *atalanta*, see *Vanessa atalanta*
- “*Vanassa*” *cardui*, see *Vanessa cardui*
- Vanessa (Lep.: Nymphalidae)
- antiopa*, see *Nymphalis antiopa*.
- atalanta*: *Compsilura concinnata*, *Euexorista futilis*, *Eusisyropa blanda*, *E. virilis*, *Lespesia archippivora*, *L. frenchii*, *Pelatachina pellucida*, *Sisyropa eudryae*, *S. eudryae* probably.
- cardui*: *Chaetogaedia monticola*, *Euexorista futilis*, *Eusisyropa blanda*, *Lespesia aletiae*, *L. archippivora*, *L. archippivora* probably *L. frenchii*, *L. spp.*, *Madremyia saundersii*, *Metopiops pyralidis*.
- “*caryae*,” see *V. carye*.
- carye*: *Lespesia archippivora*, *Madremyia saundersii*.
- huntera*, see *V. virginensis*.
- milberti*, see *Nymphalis milberti*.
- tammeamea*: *Chaetogaedia monticola*.
- virginensis*: *Compsilura concinnata*.
- spp.*: *Euexorista futilis*.
- Variable oakleaf caterpillar, see *Heterocampa manteo*
- Variiegated cutworm, see *Peridroma saucia*
- “Varigated” cutworm, see *Peridroma saucia*
- Vegetable weevil, see *Listroderes costirostris obliquus*
- Velvetbean caterpillar, see *Anticarsia gemmatalis*
- Verbena bud moth, see *Endothenia hebesana*
- Vespa mima (Lep.: Aegeriidae)
- pini*: “*Tachina*” *spp.*, *Xanthophyto labis*.
- Viceroy, see *Limenitis archippus*
- Violet sawfly, see *Ametastegia pallipes*
- Walkingstick, see *Diapheromera femorata*
- Walnut caterpillar, see *Datana integerrima*
- Walnut *Datana*, see *Datana spp.*
- Walnut sphinx, see *Cressonia juglandis*
- Weevil borer, see *Rhabdoscelus obscurus*
- Weevils other than *Anthonomus grandis*, see Curculionidae, unidentified
- Western brown stink bug, see *Euschistus impictiventris*
- Western grapeleaf skeletonizer, see *Harrisina brillians*
- Western hemlock looper, see *Lambdina fiscellaria lugubrosa*

- Western oak looper, see *Lambdina fiscellaria somniaria*
- Western spotted cucumber beetle, see *Diabrotica undecimpunctata undecimpunctata*
- Western spruce budworm, see *Choristoneura occidentalis*
- Western striped cucumber beetle, see *Acalymma trivittata*
- Western tent caterpillar, see *Malacosoma californicum*
- Western tussock moth, see *Hemerocampa vetusta*
- Western yellowstriped armyworm, see *Spodoptera praefica*
- Wheat bug(-s) (Hem. (Heter.): Pentatomidae)
Wheat bug(-s): *Gymnosoma fuliginosum*.
- Wheat head armyworm, see *Faronta diffusa*
- White cutworm, see *Euxoa scandens*
- White grub, see *Phyllophaga anxia* or White grub(-s)
- White grub(-s) (Col.: Scarabaeidae)
White grub(-s): *Microphthalma disjuncta*, *M. michiganensis*, *M. spp.*
- Whitemarked tussock moth, see *Hemerocampa leucostigma*
- White pine sawfly, see *Neodiprion pinetum*
- White pine weevil, see *Pissodes strobi*
- Whitespotted sawyer, see *Monochamus scutellatus*
- White woolly bear(-s) (Lep.: Arctiidae)
White woolly bear(-s): *Winthemia quadripustulata*.
- Willow sawfly, see *Nematus ventralis*
- Winter moth, see *Operophtera brumata*
- W-marked cutworm, see *Spaelotis clandestina*
- Woolly bear(-s) (Lep.: Arctiidae)
Woolly bear(-s): *Leschenaultia spp.*
- "Xanthopastes" *timais*, see *Xanthopastis timais*
- Xanthopastis (Lep.: Noctuidae)
timais: *Lespesia aletiae*.
- Xenotemna (Lep.: Tortricidae)
pallorana: *Actia interrupta*, *Lixophaga plumbea*.
- Xerociris (Lep.: Noctuidae)
wilsoni: *Lespesia testacea*.
- Xubida *dentilineatella*, see *Acigona dentilineatella*
- Xylena (Lep.: Noctuidae)
nupera: *Madremyia saundersii*, *Winthemia quadripustulata*, *W. rufopicta*.
spp.: *Compsilura concinnata*, *Wagneria vernata*.
- Xylina *antennata*, see *Lithophane antennata*
- Xylina *sp.*, see *Xylena spp.*
- "Xylomiges" *eridania*, see *Spodoptera eridania*
- "Xylomiges" *sunia*, see *Prodenia sunia*
- Xylomyges *eridania*, see *Spodoptera eridania*

Xyloryctes (Col.: Scarabaeidae)sp.: *Mystacella chrysoprocta*.Yellowheaded fireworm, see *Acleris minuta*Yellowheaded spruce sawfly, see *Pikonema alaskensis*Yellownecked caterpillar, see *Datana ministra*Yellowstriped armyworm, see *Spodoptera ornithogalli*Yellow woollybear, see *Diacrisia virginica***Yponomeuta** (Lep.: Yponomeutidae)*malinella*: *Nemorilla pyste*.*multipunctella*: *Nemorilla pyste*.*padella*: *Nemorilla pyste*.**Zadiprion** (Hym.: Diprionidae)*vallicola*: *Spathimeigenia mexicana*.**Zeadiatraea** (Lep.: Pyralidae)*lineolata*: *Lixophaga diatraeae*, *Paratheresia claripalpis*, *P.* spp.*lineolata* probably: *Paratheresia claripalpis*, "Phorostoma" spp.Zebra caterpillar, see *Ceramica picta***Zeiraphera** (Lep.: Olethreutidae)*ratzeburgiana*: *Phytomyptera walleyi*.**Zerene** (Lep.: Pieridae)

"catenaria": Tachinidae.

Zinckenia fascialis, see *Hymenia recurvalis***Zinckenia perspectalis**, see *Hymenia perspectalis***Zygaenidae** (Lep.: Zygaenidae)Unidentified: *Euphorocera subnitens* probably.

BIBLIOGRAPHY

This bibliography includes over 2,200 references. Under each author(s) name, references are arranged chronologically. Within each year they are grouped by journal. A year enclosed by parentheses following a journal abbreviation indicates a date that appeared on the publication but is different from the actual year of publication.

When two or more authors have the same surname, an asterisk precedes the name of the author whose article was first indexed. No initials are given in the bibliographic citations for this author in the section on parasites. The second and subsequent authors with the same surname do not have an asterisk preceding their names, but their initials are given in the bibliographic citations.

Anonymous.

- 1876 Meetings of the Entomological Club of the American Association for the Advancement of Science. Ontario Ent. Soc. Ann. Rpt. 1875: 13-17.
- 1927 Current notes. Jour. Econ. Ent. 20: 436-442.
- 1928 Japanese beetle (*Popillia japonica* Newm.). Insect Pest Survey Bul. 8 (8): 296-297.
- 1928a Current notes. Jour. Econ. Ent. 21: 942-947.
- 1928b An outbreak of the semi-tropical army worm. Fla. Ent. 12 (3): 40.
- 1930 Current notes. Jour. Econ. Ent. 23: 891-898.
- 1930a Outstanding entomological features in the United States for October, 1930. Insect Pest Survey Bul. 10 (9): 399-401.
- 1931 [Frederick A. G. Muir]. Hawaii. Planters' Rec. 35 (2): 81-82.
- 1931a Current notes. Jour. Econ. Ent. 24: 770-776.
- 1931b Current notes. Jour. Econ. Ent. 24: 1113-1115.
- 1931c Gypsy moth (*Porthetria dispar* L.). Insect Pest Survey Bul. 11 (3): 124.
- 1931d Western spotted cucumber beetle (*Diabrotica soror* Lec.). Insect Pest Survey Bul. 11 (7): 459.
- 1932 Current notes. Jour. Econ. Ent. 25: 1251-1253.
- 1945 [*Sturmia* sp. parasite of *Gilpinia hercyniae* (Htg.)]. Canad. Insect Pest Rev. 23 (3): 223-224.

- 1945a [*Sturmia* sp. parasite of spruce sawfly]. Canad. Insect Pest Rev. 23 (3): 224.
- 1946 [*Bessa harveyi* parasite of *Pristophora erichsonii* Htg.]. Canad. Insect Pest Rev. 24 (2): 195.
- 1946a [*Winthemia cilitibia* Rond. tentatively identified parasite of *Lambdina fiscellaria* var. *lugubrosa* Hulst]. Canad. Insect Pest Rev. 24 (6): 336.
- 1947 Summary of parasite and predator liberations in Canada in 1946. Canad. Insect Pest Rev. 25 (3) Sup.: 180-201.
- 1948 Summary of parasite and predator liberations in Canada in 1948. Canad. Insect Pest Rev. 26 (8): 281-297.
- [1948a] Summary of parasite and predator liberations in Canada in 1947. Canad. Insect Pest Rev. 25 (8) Sup.: 348-371.
- 1948b Notes on biological control in Cuba. Fla. Ent. 31 (2): 50.
- 1949 Summary of parasites and predator liberations in Canada in 1949. Canad. Insect Pest Rev. 27 (8): 266-289.
- 1950 Sugar cane investigational work in the British West Indies. Brit. W. Indies Sugar Assoc. Res. Work Sugar-Cane Agr. 1950: 1-13.
- 1950a Pupal collection program. U.S. Dept. Agr. Gypsy and Brown-Tail Moths Control Ann. Rpt. 1950: 2.
- 1950b [*Epimasicera caesar* Ald. parasite of the spruce budworm]. Canad. Insect Pest Rev. 28 (6): 210.
- 1950c Summary of parasites and predator liberations in Canada in 1950. Canad. Insect Pest Rev. 28 (8): 232-251.
- 1951 Foreword. Pupal collection program. U.S. Dept. Agr. Gypsy and Brown-Tail Moths Control Ann. Rpt. 1951: 1, 4-8.
- 1952 Control of the alfalfa caterpillar. U.S. Dept. Agr. Div. Cereal and Forage Insect Invest. Leaflet 325: 1-5, 4 figs.
- 1952a Satin moth (*Stilpnotia salicis* L.). Canad. Insect Pest Rev. 30 (3): 164.
- 1952b A summary of parasite and predator liberations in Canada in 1951. Canad. Insect Pest Rev. 30 (4): 174-187.

- 1952c Colorado potato beetle (*Leptinotarsa decemlineata* Say). *Canad. Insect Pest Rev.* 30 (5): 204-205.
- 1952d Larch sawfly (*Pristiphora erichsonii* Htg.). *Canad. Insect Pest Rev.* 30 (5): 209-210.
- 1952e Satin moth (*Stilpnotia salicis* L.). *Canad. Insect Pest Rev.* 30 (5): 211.
- 1952f European spruce sawfly (*Diprion hercyniae* Htg.). *Canad. Insect Pest Rev.* 30 (7): 256.
- 1952g A summary of parasite and predator liberations in Canada in 1952. *Canad. Insect Pest Rev.* 30 (9): 284-296.
- 1953 Departamento de defensa agropecuaria. Costa Rica Min. de Agr. Indus. Mem. 1952: 81-96.
- 1953a [*Euacemyia* spp. parasite of grasshoppers]. *Canad. Insect Pest Rev.* 31 (2): 128.
- 1953b Colorado potato beetle parasite (*Doryphorophaga doryphorae* Riley). *Canad. Insect Pest Rev.* 31 (2): 135.
- 1953c Dipterous parasite recoveries. *Canad. Insect Pest Rev.* 31 (2): 137.
- 1954 Departamento de defensa agro-pecuaria. Costa Rica Min. de Agr. Indus. Mem. 1953: 79-112.
- 1954a Crane flies (Tipulidae). *Canad. Insect Pest Rev.* 32 (2): 130.
- 1954b Parasites of the spruce budworm (*Choristoneura fumiferana* Clem.). *Canad. Insect Pest Rev.* 32 (2): 147.
- 1954c Parasites of the European spruce sawfly (*Diprion hercyniae* Htg.). *Canad. Insect Pest Rev.* 32 (2): 148.
- 1954d Colorado potato beetle (*Leptinotarsa decemlineata* Say). *Canad. Insect Pest Rev.* 32 (5): 250.
- 1954e Brown-headed ash sawfly (*Tomostethus multicinctus* Roh.). *Canad. Insect Pest Rev.* 32 (5): 253.
- 1954f European earwig (*Forficula auricularia* L.). *Canad. Insect Pest Rev.* 32 (7): 300.
- 1954g Gypsy moth (*Lymantria dispar* L.). *Canad. Insect Pest Rev.* 32 (7): 309.
- 1954h Salt-marsh caterpillar (*Estigmene acrea*). (U.S. Dept. Agr.) *Coop. Econ. Insect Rpt.* 4 (3): 47.
- 1954i European corn borer parasites in Minnesota. (U.S. Dept. Agr.) *Coop. Econ. Insect Rpt.* 4 (21): 438.

- 1955 The European corn borer and its control. U.S. Dept. Agr. Farmers' Bul. 2084: 1-20, figs. 1-23.
- 1955a Summary. U.S. Dept. Agr. Gypsy Moth Control Ann. Rpt. 1955: 1-11.
- 1955b Program activities. U.S. Dept. Agr. Gypsy Moth Control Ann. Rpt. 1955: 12-62.
- 1955c Larch sawfly (*Pristiphora erichsonii* Htg.). Canad. Insect Pest Rev. 33 (3): 195-196.
- 1956 Parasites of grasshoppers. Canad. Insect Pest Rev. 34 (4): 196.
- 1956a Winter moth (*Operophtera brumata* L.). Canad. Insect Pest Rev. 34 (4): 198.
- 1956b Winter moth (*Operophtera brumata* L.). Canad. Insect Pest Rev. 34 (7): 284.
- 1956c Summary of insect conditions — 1955: Wisconsin. (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 6 (13): 274.
- 1956d Parasites — Ida. (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 6 (45): 1044.
- 1957 Imported cabbageworm (*Pieris rapae* L.). Canad. Insect Pest Rev. 35 (3): 172.
- 1957a Winter moth (*Operophtera brumata* L.). Canad. Insect Pest Rev. 35 (3): 176.
- 1957b Colorado potato beetle (*Leptinotarsa decemlineata* Say). Canad. Insect Pest Rev. 35 (6): 259.
- 1957c Colorado potato beetle (*Leptinotarsa decemlineata* Say). Canad. Insect Pest Rev. 35 (7): 273.
- 1957d Summary of insect conditions — 1956: Wisconsin. (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 7 (13): 240-244.
- 1957e European corn borer (*Pyrausta nubilalis*) — Wisconsin. (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 7 (24): 460.
- 1957f Parasites — California. (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 7 (42): 828.
- 1958 European earwig (*Forficula auricularia* L.). Canad. Insect Pest Rev. 36 (2): 126.
- 1958a Colorado potato beetle (*Leptinotarsa decemlineata* Say). Canad. Insect Pest Rev. 36 (2): 134.
- 1958b Winter moth (*Operophtera brumata* L.). Canad. Insect Pest Rev. 36 (2): 139.
- 1958c A parasite of the Colorado potato beetle (*Adoryphorophaga aberrans* Tnsd.). Canad. Insect Pest Rev. 36 (7): 284.

- 1958d Summary of insect conditions — 1957: Wisconsin. (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 8 (11): 193-198.
- 1958e Armyworm (*Pseudaletia unipuncta*) — Wisconsin. (U.S., Dept. Agr.) Coop. Econ. Insect Rpt. 8 (30): 648.
- 1958f European corn borer (*Pyrausta nubilalis*) — Wisconsin. (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 8 (31): 668.
- 1958g European corn borer (*Pyrausta nubilalis*) — Wisconsin. (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 8 (35): 755.
- 1959 Summary of insect conditions — 1958[:] Wisconsin. (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 9 (14): 241-245.
- 1959a Summary of insect conditions — 1958[:] Minnesota. (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 9 (15): 268-274.
- 1959b Larch sawfly (*Pristiphora erichsonii*) — Minnesota. (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 9 (31): 711.
- 1960 Barbados. Review of work in economic entomology. Commonwealth Ent. Conf. Rpt. 7: 211-213.
- 1960a IV. Sugar industry research, including a survey of pests and diseases and notes on irrigation. Brit. W. Indies Sugar Assoc. Handb. 1961: 43-65.
- 1960b The sugarcane borer: How to control it. U.S. Dept. Agr. Leaflet 479: 1-8, 7 figs.
- 1960c Summary of insect conditions in the United States [-] 1959. (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 10 (42): 975-988.
- 1960d Forest insect conditions in the United States, 1959. (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 10 (43): 1003-1032.
- 1960e A tachinid (*Voria ruralis*) — Arizona. (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 10 (49): 1115.
- 1961 The southern corn rootworm, how to control it. U.S. Dept. Agr. Leaflet 391: 1-8, 8 figs.
- 1961a Insects of the season 1961 in the Lower Fraser Valley. Canad. Insect Pest Rev. 39 (10): 298-301.
- 1961b A tachina fly (*Voria ruralis*) — Arizona. (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 11 (2): 18.
- 1961c Summary of insect conditions — 1960. (Cont.). (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 11 (14): 274-284.

- 1961d Parasites — Wisconsin. (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 11 (28): 639.
- 1961e A parasitic tachinid (*Voria ruralis*) — Arizona. (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 11 (46): 1049.
- 1962 Cutworms (Noctuidae). Canad. Insect Pest Rev. 40 (6): 120-121.
- 1962a The European corn borer, how to control it. U.S. Dept. Agr. Farmers' Bul. 2190: 1-20, figs. 1-20.
- 1962b Summary of insect conditions — 1961. (Cont.). (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 12 (12): 244-274.
- 1963 Control of the alfalfa caterpillar. U.S. Dept. Agr. Leaflet 325 (rev.): 1-8, 7 figs.
- 1963a Grasshopper parasitism. Canad. Insect Pest Rev. 41 (1): 8.
- 1963b Imported cabbageworm (*Pieris rapae* (L.)). Canad. Insect Pest Rev. 41 (1): 19.
- 1963c Red-banded leaf roller (*Argyrotaenia velutinana* (Walker)). Canad. Insect Pest Rev. 41 (1): 20.
- 1963d European corn borer parasites — Wisconsin. (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 13 (35): 1029.
- 1963e Parasites of European corn borer in Maryland. (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 13 (46): 1337.
- 1964 Summary of insect conditions in Hawaii — 1963. (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 14 (12): 220-223.
- 1964a Summary of insect conditions in the United States — 1963. (Cont.). (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 14 (17): 384-394.
- 1964b Parasites — Wisconsin. (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 14 (28): 779.
- 1965 Summary of insect conditions in Hawaii — 1964. (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 15 (6): 77-79.
- 1965a Summary of insect conditions in the United States — 1964. (Cont.). (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 15 (17): 397-406.

- 1965b Corn earworm (*Heliothis zea*) — Oklahoma. (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 15 (38): 1091.
- 1966 Summary of insect conditions in Hawaii — 1965. (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 16 (5): 73-76.
- 1966a Summary of insect conditions in the United States — 1965. (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 16 (6): 87-96.
- 1966b A tachina fly (*Trichopoda pennipes*). (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 16 (7): 103.
- 1966c Summary of insect conditions in the United States — 1965. (Cont.). (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 16 (15): 319-329.
- 1966d New Guinea surgarcane [sic] weevil (*Rhabdoscelus obscurus*). (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 16 (22): 502.
- 1966e Southern green stink bug (*Nezara viridula* var. *smaragdula*). (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 16 (42): 995.
- 1968 Summary of insect conditions in Hawaii — 1967. (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 18 (7): 82-85.
- 1968a A tachina fly (*Trichopoda pennipes*). (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 18 (16): 316.
- 1969 Tachina flies — Oklahoma. (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 19 (5): 59.
- 1969a Summary of insect conditions in Hawaii — 1968. (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 19 (6): 77-80.
- 1969b Armyworm (*Pseudaletia unipuncta*) — Wisconsin. (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 19 (33): 637.
- 1969c Parasites of *Herpetogramma licarsisalis* (Walker). Hawaii. Ent. Soc. Proc. (1968) 20 (2): 260.

Adams, J. A.

- 1944 Japanese beetle control by parasitic insects. An Asiatic wasp and a fly from Japan which are natural enemies of the Jap beetle show promise in fight on pest. N.Y. Agr. Expt. Sta. (Geneva) Farm Res. 10 (4): 1, 9, figs. 1-4.

— and Wheeler, E. H.

- [1944] Biological control of the Japanese beetle by parasitic insects. N.Y. State Agr. Expt. Sta. Ann. Rpt. 63: 25.

Adams, R. H. (See Bottrell et al.)

Agee, H. P.

- 1913 Report of director of the Experiment Station. Hawaii. Sugar Planters' Assoc. Expt. Sta. Committee Rpt. 1913: 9-23.
- 1913a Report of the director of the Experiment Station. Hawaii. Planters' Rec. 9 (6): 656-668.
- 1914 [Report of director of the Experiment Station]. Hawaii. Sugar Planters' Assoc. Proc. 33rd Ann. Mtg. 1913 33: 21-35.
- 1914a Report of the director of the Experiment Station. Hawaii. Sugar Planters' Assoc. Expt. Sta. Committee Rpt. 1914: 5-17.
- 1914b Report of the director of the Experiment Station. Hawaii. Planters' Rec. 11 (6): 305-317.
- 1915 Director's report. Hawaii. Sugar Planters' Assoc. Proc. 34th Ann. Mtg. 1914 34: 16-28.
- 1915a Director's report. Hawaii. Sugar Planters' Assoc. Expt. Sta. Committee Rpt. 1915: 5-21.
- 1915b Director's annual report. Hawaii. Planters' Rec. 13 (6): 337-352.
- 1916 Director's report. Hawaii. Sugar Planters' Assoc. Proc. 35th Ann. Mtg. 1915 35: 11-28.
- 1916a Director's report. Hawaii. Planters' Rec. 15 (6): 340-362.
- 1916b Director's report. Hawaii. Sugar Planters' Assoc. Expt. Sta. Committee Rpt. 1916: 4-26.
- 1917 Director's report. Hawaii. Sugar Planters' Assoc. Proc. 36th Ann. Mtg. 1916 36: 13-35.
- 1918 Director's report. Hawaii. Sugar Planters' Assoc. Proc. 37th Ann. Mtg. 1917 37: 33-52.
- 1918a Director's report. Hawaii. Planters' Rec. 18 (1): 11-31.
- 1919 Director's report. Hawaii. Sugar Planters' Assoc. Proc. 38th Ann. Mtg. 1918 38: 153-188.
- 1922 Director's report. Hawaii. Sugar Planters' Assoc. Rpt. Committee Charge Expt. Sta. 1922: 6-64.
- 1927 Work with parasites of sugar cane insects in Hawaii. Internatl. Soc. Sugar Cane Technol. Cong. Proc. 2: 72-74.

Ah Sam, J. (See Miyahira and Ah Sam)

*Ainslie, C. N.

- 1910 The New Mexico range caterpillar. U.S. Dept. Agr. Bur. Ent. Bul., n. ser., 85 (5): 59-96, figs. 32-53, pls. 3-4.

Ainslie, G. G.

- 1910 The cowpea curculio. U.S. Dept. Agr. Bur. Ent. Bul., n. ser., 85 (8): 129-142, figs. 67-69.
- 1923 Striped sod webworm, *Crambus mutabilis* Clemens. Jour. Agr. Res. 24 (5): 399-414, figs. 1-2, pls. 1-2.
- 1927 The larger sod webworm. U.S. Dept. Agr. Tech. Bul. 31: 1-18, figs. 1-2.

— and Cartwright, W. B.

- 1921 Biology of the smartweed borer, *Pyrausta ainsliei* Heinrich. Jour. Agr. Res. 20 (11): 837-844.
- 1922 Biology of the lotus borer (*Pyrausta penitalis* Grote). U.S. Dept. Agr. Bul. 1076: 1-14, pls. 1-4.

Aldrich, J. M.

- 1915 Results of twenty-five years' collecting in the Tachinidae, with notes on some common species. Ent. Soc. Amer. Ann. 8 (1): 79-84.
- 1916 Two new Canadian Diptera. Canad. Ent. 48 (1): 20-22.
- 1921 The muscoid genera *Pseudeuanta* and *Uramyia*. Insecutor Inscitiae Menstruus 9 (4-6): 83-92.
- 1923 A new tachinid parasite of the codling moth (Dip.). Ent. News 34 (2): 53-54.
- 1923a A new parasitic fly bred from the bean beetle. Ent. Soc. Wash. Proc. 25 (4): 95-96.
- 1923b The present status of Coquillett's *Hypochoaeta longicornis* Schiner (Diptera). Ent. Soc. Wash. Proc. 25 (7-8): 161-162.
- 1924 The muscoid genus *Genea* in North America (Dipt.). Ent. News 35 (6): 210-214.
- 1925 Notes on some types of American muscoid Diptera in the collection of the Vienna Natural History Museum. [Cont.]. Ent. Soc. Amer. Ann. 18 (1): 107-130, figs. 1-2 [Cont.].
- 1925a Notes on some types of American muscoid Diptera in the collection of the Vienna Natural History Museum. [Concl.]. Ent. Soc. Amer. Ann. 18 (4): 456-469, fig. 1.
- 1925b New Diptera or two-winged flies in the United States National Museum. U.S. Natl. Mus. Proc. 66 (18): 1-36, fig. 1.
- 1925c Two new species of the tachinid genus *Lixophaga*, with notes and key (Diptera). Ent. Soc. Wash. Proc. 27 (6): 132-136.

- 1926 American two-winged flies of the genus *Microphthalma* Macquart, with notes on related forms. U.S. Natl. Mus. Proc. 69 (13): 1-8.
- 1926a Descriptions of new and little known Diptera or two-winged flies. U.S. Natl. Mus. Proc. 69 (22): 1-26.
- 1926b Notes on the metallic green tachinids allied to *Gymnochaeta*, with keys and one new Chinese genus (Diptera). Insecutor Inscitiae Menstruus 14 (4-6): 51-58.
- 1927 The dipterous parasites of the migratory locust of tropical America, *Schistocerca paranensis* Burmeister. Jour. Econ. Ent. 20: 588-593.
- 1927a A new species of *Oedematocera* reared from the tropical migratory locust (Diptera). Ent. Soc. Wash. Proc. 29 (1): 17-18.
- 1927b Notes on muscoid synonymy. Brooklyn Ent. Soc. Bul. 22 (1): 18-25.
- 1928 A new species of *Oedematocera* with notes on *Schistocercophaga* Townsend (Dipt.: Tachinidae). Ent. News 39 (10): 301-304.
- 1928a A revision of the American parasitic flies belonging to the genus *Belvosia*. U.S. Natl. Mus. Proc. 73 (8): 1-45.
- 1928b Synonymic notes on Diptera. Ent. Soc. Wash. Proc. 30 (3): 41-45.
- 1929 Notes on synonymy of Diptera, no. 3. Ent. Soc. Wash. Proc. 31 (2): 32-36.
- 1931 North American two-winged flies of the genus *Spathimeigenia*, with descriptions of five new species. U.S. Natl. Mus. Proc. 80 (11): 1-10.
- 1932 Records of dipterous insects of the family Tachinidae reared by the late George Dimmock, with description of one new species and notes on the genus *Anetia* Robineau-Desvoidy. U.S. Natl. Mus. Proc. 80 (20): 1-8.
- 1932a New Diptera, or two-winged flies, from America, Asia, and Java, with additional notes. U.S. Natl. Mus. Proc. 81 (9): 1-28, pl. 1.

— and Webber, R. T.

- 1924 The North American species of parasitic two-winged flies belonging to the genus *Phorocera* and allied genera. U.S. Natl. Mus. Proc. 63 (17): 1-90, fig. 1.

Alexander, C. P.

- 1920 The crane-flies of New York. Part II. Biology and phylogeny. N.Y. (Cornell) Agr. Expt. Sta. Mem. 38: 691-1133, figs. 1-539, pls. 11-97.

Allen, D. C., Knight, F. B., Foltz, J. L., and Mattson, W. J.

- 1969 Influence of parasites on two populations of *Choristoneura pinus* (Lepidoptera: Tortricidae) in Michigan. Ent. Soc. Amer. Ann. 62 (6): 1469-1475, fig. 1.

*Allen, H. W. (See also Brunson and Allen)

- 1924 A biological and ecological study of Tachinidae parasitic on insect pests of southern crops. Miss. Agr. Expt. Sta. Ann. Rept. 37: 24-25.
- [1925] Biology of the red-tailed tachina-fly, *Winthemia quadripustulata* Fabr. Miss. Agr. Expt. Sta. Tech. Bul. 12: 1-32, figs. 1-9.
- 1926 Life history of the variegated cutworm tachina fly, *Archytas analis*. Jour. Agr. Res. 32 (5): 417-435, figs. 1-6, pl. 1.
- 1926a Observations upon the early maggot stage of *Linnaemyia comta* Fall. (Diptera: Tachinidae). Ent. News 37 (9): 283-286, fig. 1.
- 1926b Notes on some North American species of *Achaetoneura* with a description of one new species (Diptera: Tachinidae). Amer. Ent. Soc. Trans. 52 (3): 187-198, pl. 9.
- 1929 An annotated list of the Tachinidae of Mississippi. Ent. Soc. Amer. Ann. 22 (4): 676-690, fig. 1.
- 1962 Parasites of the oriental fruit moth in the Eastern United States. U.S. Dept. Agr. Tech. Bul. 1265: 1-139, figs. 1-28.

—— Holloway, J. K., and Haeussler, G. J.

- 1940 Importation, rearing, and colonization of parasites of the oriental fruit moth. U.S. Dept. Agr., Dept. Cir. 561: 1-62, figs. 1-15.

—— and Jaynes, H. A.

- 1928 Progress with the imported parasites of the Japanese beetle during 1927. Jour. Econ. Ent. 21: 823-832, fig. 55.

Allen, M. W. (See Knowlton and Allen)

Alwood, W. B.

- 1898 On the life history of *Protoparce carolina*. U.S. Dept. Agr. Div. Ent. Bul., n. ser., 17: 72-74.

Amos, J. M. (See Stearns and Amos)

Anderson, L. D. (See Jones, D. W., Walker, and Anderson)

Anderson, M. (See Wilkes and Anderson)

Andre, F. (See also Simanton and Andre)

- 1951 New findings for farm folk. Part I. Iowa Agr. Expt. Sta. Rpt. 1949-1951: 1-74, figs. 1-8, 69 figs.

Annand, P. N.

- 1944 Report of the chief of the Bureau of Entomology and Plant Quarantine, Agricultural Research Administration, 1943. U.S. Dept. Agr., Agr. Res. Admin., Adminr. Rpt. 1943: 123-180.
- 1945 Report of the chief of the Bureau of Entomology and Plant Quarantine, Agricultural Research Administration, 1944. U.S. Dept. Agr., Agr. Res. Admin., Adminr. Rpt. 1944: 123-178.
- 1946 Report of the chief of the Bureau of Entomology and Plant Quarantine, Agricultural Research Administration, 1945. U.S. Dept. Agr., Agr. Res. Admin., Adminr. Rpt. 1945: 117-177.
- 1948 Report of the chief of the Bureau of Entomology and Plant Quarantine, Agricultural Research Administration, 1947. U.S. Dept. Agr., Agr. Res. Admin., Adminr. Rpt. 1947: 177-251.

Anthony, D. W. (See Jones, C. M., and Anthony)

Arant

- 1955 Tachina flies — Alabama. (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 5 (43): 994.

Arant, F. S.

- 1929 Biology and control of the southern corn rootworm. Ala. Agr. Expt. Sta. Bul. 230: 1-46, figs. 1-14.
- 1933 Life history and control of the cowpea curculio, *Chalcodermus aeneus*, Boh. Ala. Agr. Expt. Sta. Ann. Rpt. 44: 28-29.
- 1938 Life history and control of the cowpea curculio. Ala. Agr. Expt. Sta. Bul. 246: 1-34, figs. 1-14.

Arbuthnot, K. D. (See also Blickenstaff, Arbuthnot, and Harris; Cory et al.; Jones, D. W., and Arbuthnot)

- 1947 Present status of European corn borer parasites in the north central states, and reports from Indiana, Illinois, and Iowa. [Part] (a). Ent. Soc. Amer., No. Cent. Branch, Proc. 2: 30-33.
- 1950 The status of European corn borer parasitism in the north central states at the close of 1949. Ent. Soc. Amer., No. Cent. Branch, Proc. 5: 44-46.

- 1950a Status of European corn borer parasites in the United States. Jour. Econ. Ent. 43: 422-426, figs. 1-3.
- 1952 The release and establishment of European corn borer parasites in the corn belt states and the effect of the spray program on their abundance. Ent. Soc. Amer., No. Cent. Branch, Proc. 7: 18-22.
- 1953 Present status and potential value of parasites of the European corn borer. Ent. Soc. Amer., No. Cent. Branch, Proc. 8: 30-32.
- 1955 European corn borer parasite complex near East Hartford, Connecticut. Jour. Econ. Ent. 48: 91-93, figs. 1-2.
- Jones, D. W., Carter, S. W., and Cosenza, A. V.
- 1947 The colonization of parasites of the European corn borer in the United States in 1946. U.S. Dept. Agr., Agr. Res. Admin., Bur. Ent. and Plant Quar., Insect Pest Survey, Spec. Sup. 1947, No. 6: 1-4.
- Jones, D. W., Carter, S. W., and Evans, R. W.
- 1948 The field status of parasites of the European corn borer at the close of 1947. U.S. Dept. Agr., Agr. Res. Admin., Bur. Ent. and Plant Quar., Insect Pest Survey, Spec. Sup. 1948, No. 5: 1-14, fig. 1.
- 1949 The field status of parasites of the European corn borer at the close of 1948. U.S. Dept. Agr., Agr. Res. Admin., Bur. Ent. and Plant Quar., Insect Pest Survey, Spec. Sup. 1949, No. 7: 1-24, figs. 1-2.
- and Wright, J. M.
- 1952 Parasites of the European corn borer in Illinois. Ill. State Acad. Sci. Trans. (1951) 44: 222-234, figs. 1-2.
- Armitage, H. M.
- 1954 Bureau of entomology. Calif. State Dept. Agr. Bul. (1953) 42 (4): 199-233, figs. 1-7.
- Armstrong, T.
- 1958 Life-history and ecology of the plum curculio, *Conotrachelus nenuphar* (Hbst.) (Coleoptera: Curculionidae), in the Niagara Peninsula, Ontario. Canad. Ent. 90 (1): 8-17, fig. 1.
- Arnaud, P. H. (Jr.) (See also Sabrosky and Arnaud)
- 1951 A study of the genus *Paradejeania* Brauer and

- Bergentamm (Diptera: Tachinidae or Larvaevoridae). *Canad. Ent.* 83 (12): 317-329, figs. 1-9.
- 1957 The occurrence of *Salmacia longipulvilli* in the Hawaiian Islands (Diptera: Larvaevoridae). *Ent. News* 68 (10): 259-263, figs. 1-5.
- 1959 A new host record for *Archytas californiae* (Walker) in California (Diptera: Tachinidae). *Pan-Pacific Ent.* 35 (3): 156.
- 1959a A first host record and new distribution records for *Philocalia tenuirostris* Reinhard (Diptera, Tachinidae). *Ent. Soc. Wash. Proc.* 61 (4): 186.
- 1960 A review of the genus *Spathidexia* Townsend (Diptera: Tachinidae). *Wasmann Jour. Biol.* 18 (1): 1-36, figs. 1-33.
- 1960a [Monarch butterfly chrysalis parasitized]. *Pan-Pacific Ent.* 36 (1): 51.
- 1963 Types of the Tachinidae (Diptera) in the American Museum of Natural History. *Amer. Mus. Nat. Hist. Bul.* 125 (2): 101-137.
- 1963a Systematic studies in the tribe Neominthoini (Diptera, Tachinidae). *Amer. Mus. Nat. Hist., Amer. Mus. Novitates* 2135: 1-55, figs. 1-44.
- 1963b Two new genera of the Germariini from the western Nearctic, with notes on related genera (Diptera, Tachinidae). *Amer. Mus. Nat. Hist., Amer. Mus. Novitates* 2157: 1-20, figs. 1-22.
- 1963c The type locality of *Eusisyropa virilis* (Aldrich and Webber) (Diptera: Tachinidae). *Ent. Soc. Wash. Proc.* 65 (2): 116.
- 1969 *Compsilura concinnata* (Meigen) (Diptera: Tachinidae) reared from *Agraulis vanillae incarnata* (Riley) (Lepidoptera: Nymphalidae) at San Leandro, California. *Pan-Pacific Ent.* 45 (1): 77.
- and Rentz, D. C.
- 1965 *Ceracia dentata* a parasite of *Chimarocephala pacifica pacifica* in California (Diptera: Tachinidae and Orthoptera: Acrididae). *Pan-Pacific Ent.* 41 (3): 204-206, fig. 1.
- Arthur, A. P.
- 1962 A skipper, *Thymelicus lineola* (Ochs.) (Lepidoptera: HesperIIDae) and its parasites in Ontario. *Canad. Ent.* 94 (10): 1082-1089, fig. 1.

- 1966 The present status of the introduced skipper, *Thymelicus lineola* (Ochs.) (Lepidoptera: HesperIIDae), in North America and possible methods of control. *Canad. Ent.* 98 (6): 622-626, fig. 1.
- Artiles, R. Fernández (See Scaramuzza, Fernández Artiles, and Fernández Perez)
- Ashlock, P. D., and O'Brien, C. W.
- 1964 *Catharosia lustrans*, a tachinid parasite of some drymine Lygaeidae (Diptera and Hemiptera-Heteroptera). *Pan-Pacific Ent.* 40 (2): 98-100.
- Ashmead, W. H. (See also Dimmock and Ashmead)
- 1887 Report on insects injurious to garden crops in Florida. U.S. Dept. Agr., Div. Ent. Bul. 14: 9-29.
- 1894 The habits of the aculeate Hymenoptera. — I. *Psyche* 7 (214): 19-26.
- Ashworth, J. T. (See Britton and Ashworth)
- Atkins, E. L. (Jr.)
- 1959 The western tussock moth, *Hemerocampa vetusta* (Bdv.), on citrus in southern California. *Jour. Econ. Ent.* (1958) 51: 762-765.
- Atkinson, N. J. (See King, K. M., and Atkinson)
- Atwell, H. C.
- 1927 The European earwig (*Forficula auricularia* Linn). *Oreg. Bd. Hort. Bien. Rpt.* 19: 86-103, figs. 1-5.
- Atwood, C. E.
- 1938 [Eggs of *Bessa selecta* found on larch sawfly larvae]. *Canad. Insect Pest Rev.* 16 (2): 295-296.
- 1939 Insects of the season 1938 in the Laniel District, P.Q., and other territory served by the Laniel field station. *Canad. Insect Pest Rev.* 17 (1): 37-39.
- 1940 [*Bessa selecta* parasite of *Pristiphora erichsoni* Hartig]. *Canad. Insect Pest Rev.* 18 (3): 210-211.
- 1944 Forest insect conditions 1943 in eastern Ontario and western Quebec. *Canad. Insect Pest Rev.* 22 (1): 52-61.
- Au, S. A.
- 1968 *Pachyzancla licarsisalis* (Walker). *Hawaii. Ent. Soc. Proc.* 1967 20 (1): 17.
- Bacon, O. G. (See Michelbacher et al.)
- Bade, E.
- 1933 The Japanese beetle, a pest. *Gard. Chron. Amer.* 37 (1): 12, 16, figs. 1-2.

- Baerg, W. J.
1928 Three shade tree insects. Ark. Agr. Expt. Sta. Bul. 224: 1-25, figs. 1-13.
- Baird, A. B.
1917 An historical account of the forest tent caterpillar and of the fall webworm in North America. Ontario Ent. Soc. Ann. Rpt. 47: 73-87.
1918 Some notes on the natural control of the cherry-tree ugly nest tortricid, *Archips cerasivorana* Fitch. Agr. Gaz. Canada 5 (8): 766-771, figs. 1-6.
[1923] Some notes on the natural control of the larch sawfly and larch case bearer in New Brunswick in 1922. Acadian Ent. Soc. Proc. (1922) 8: 158-171.
1926 Recent developments in the introduction of parasites of the European corn borer in Ontario. Ontario Ent. Soc. Ann. Rpt. 56: 78.
1929 The present status of corn borer parasites in Canada. Ontario Ent. Soc. Ann. Rept. (1928) 59: 38-40.
1937 Summary statement of parasites liberated or distributed in Canada prior to June, 1936. Canad. Insect Pest Rev. 15 (1): 66-68.
1938 Summary of insect parasites and predators liberated in Canada up to December 31, 1937. Canad. Insect Pest Rev. 16 (1) Sup.: 77-154.
1939 Summary of parasite liberations in Canada during the year 1938. Canad. Insect Pest Rev. 17 (1): 102-128.
1939a Biological control of insect pests in Canada with special reference to the control of the European spruce sawfly *Gilpinia polytoma* Htg. Ontario Ent. Soc. Ann. Rpt. 70: 51-56.
1940 Summary of insect parasites and predators liberated in Canada and Newfoundland 1939. Canad. Insect Pest Rev. 18 (1): 94-126.
1941 The spruce sawfly parasite situation in 1940. Pulp and Paper Mag. Canada 42 (2): 159-161.
1941a Summary of parasite and predator liberations in Canada and Newfoundland in 1940. Canad. Insect Pest Rev. 19 (1): 94-125.
1942 Summary of parasite and predator liberations in Canada in 1941. Canad. Insect Pest Rev. 20 (1): 112-135.

- 1943 Summary of parasite and predator liberations in Canada in 1942. *Canad. Insect Pest Rev.* 21 (1): 107-130.
- 1944 Summary of parasite and predator liberations in Canada in 1943. *Canad. Insect Pest Rev.* 22 (1): 117-138.
- 1944a [A summary of parasites recovered from a collection of *Neodiprion lecontei* in Ontario]. *Canad. Insect Pest Rev.* 22 (2): 157.
- 1945 Summary of parasite and predator liberations in Canada in 1944. *Canad. Insect Pest Rev.* 23 (1): 119-139.
- 1946 Summary of parasite and predator liberations in Canada in 1945. *Canad. Insect Pest Rev.* 24 (1): 134-152.
- *Baker, A. W.**
- 1915 The army worm in Ontario in 1914. *Ontario Ent. Soc. Ann. Rpt.* 45: 75-90, figs. 13-18.
- 1938 Notes on the armyworm, *Leucania unipuncta* Haw. outbreak in Ontario in 1938. *Ontario Ent. Soc. Ann. Rpt.* 69: 96-99.
- Baker, C. F.**
- 1910 *Harpyia cinerea* Wlk. near Spadra, Cal. *Pomona Col. Jour. Ent.* 2 (4): 382.
- Baker, W. A., and Bradley, W. G.**
- 1938 Field status of European corn borer parasites at the close of 1936. *Jour. Econ. Ent.* 31: 345-348.
- **Bradley, W. G., and Clark, C. A.**
- 1949 Biological control of the European corn borer in the United States. *U.S. Dept. Agr. Tech. Bul.* 983: 1-185, figs. 1-40.
- Balch, R. E.**
- 1936 The European spruce sawfly outbreak in 1935. *Canad. Ent.* 68 (2): 23-31.
- 1936a [*Bessa selecta* parasite of larch sawfly]. *Canad. Insect Pest Rev.* 14 (3): 144.
- 1936b [*Bessa selecta* parasite of *Lygaeonematus erichsoni* Hartig]. *Canad. Insect Pest Rev.* 14 (4): 197.
- 1937 Forest insects in the maritime provinces and eastern Quebec in 1936. *Canad. Insect Pest Rev.* 15 (1): 24-26.
- 1938 Forest insects in the maritime provinces and eastern Quebec in 1937. *Canad. Insect Pest Rev.* 16 (1): 25-27.

- 1939 Forest insects of the season 1938 in the maritime provinces and eastern Quebec. *Canad. Insect Pest Rev.* 17 (1): 29-32.
- 1940 Forest insects of the season 1939 in the maritime provinces and eastern Quebec. *Canad. Insect Pest Rev.* 18 (1): 20-23.
- 1941 Entomological investigation. Report on forest insect conditions in Nova Scotia in 1940. Nova Scotia Dept. Lands and Forests Rpt. 1940: 36-39.
- 1960 The approach to biological control in forest entomology. *Canad. Ent.* 92 (4): 297-310.
- and Hawboldt, L. S.
- 1941 Forest insects of the season 1940 in the maritime provinces and eastern Quebec. *Canad. Insect Pest Rev.* 19 (1): 22-26.
- and Simpson, L. J.
- 1935 Forest insects in the maritime provinces and Gaspé Peninsula, 1934. *Canad. Insect Pest Rev.* 13 (1): 35-37.
- Balduf, W. V.**
- 1925 The striped cucumber beetle *Diabrotica vittata* Fabr., order Coleoptera, family Chrysomelidae. *Ohio Agr. Expt. Sta. Bul.* 388: 241-310, figs. 1-14.
- 1929 Bionomic notes on some parasites of *Achatodes zea* Harris (Noctuidae, Lep.) and *Phlyctaenia tertialis* (Guen.) (Pyralidae, Lep.). *Ohio Jour. Sci.* 29 (5): 218-242.
- 1937 Bionomic notes on the common bagworm, *Thyridopteryx ephemeraeformis* Haw., (Lepid., Psychidae) and its insect enemies (Hym., Lepid.). *Ent. Soc. Wash. Proc.* 39 (7): 169-184, figs. 1-3.
- 1961 Insects from tunnels of *Xylocopa virginica* (Linn.) (Col., Lep., Dipt., Hym.). *Brooklyn Ent. Soc. Bul.* 56 (3): 81-85.
- 1968 Bionomic notes on the hexapodous parasites of *Acrobasis rubrifasciella*. *Ent. Soc. Amer. Ann.* 61 (2): 463-476, figs. 1-11.
- *Baldwin, C. H.**
- 1914 Insects of the year 1913-1914. *Ind. State Ent. Ann. Rpt.* 7: 13-58, 40 figs.
- Baldwin, W. F., and Coppel, H. C.**
- 1949 Observations on the emergence of *Phorocera hamata* A. & W. (Tachinidae) from sawfly cocoons. *Canad. Ent.* (1947) 79 (11-12): 221-223, fig. 1.

- 1949a The biology of *Phorocera hamata* A. & W., a tachinid parasite of sawflies. *Canad. Ent.* 81 (10): 237-245, figs. 1-16.
- Ballou, C. H.
 1936 *Entomologia*. [Costa Rica] *Cent. Nac. de Agr. Bol.* 18: 95-155, 3 figs.
 1936a *Insectos observados durante el año 1934*. [Costa Rica] *Cent. Nac. de Agr. Bol.* 20: 1-60, 3 figs.
- Banerjee, A. C.
 1967 Sod webworm parasites in Illinois. *Jour. Econ. Ent.* 60: 1173-1174.
- Barber, G. W. (See King, V., and Barber; Phillips, W. J., and Barber)
- Barker, R. B. (See also McGuffin and Barker)
 1947 Budworm pupal parasite survey. *Canada Dept. Agr., Div. Forest Biol., Bi-monthly Prog. Rpt.* 3 (1): 2-3.
- Barr, W. F. (See Furniss, M. M., and Barr; Manis et al.)
- Bartlett, K. A.
 1937 The introduction and colonization in Puerto Rico of beneficial insects parasitic on the sugarcane moth borer. *Puerto Rico Agr. Expt. Sta. (Mayaguez) Agr. Notes* 78: 1-8.
 1938 The introduction and colonization of the Amazon fly, *Metagonistylum minense* Tns., in Puerto Rico. *Internatl. Soc. Sugar Cane Technol. Rpt. Committee, Abs. Papers* 6: 86.
 1938a Biological control of the sugarcane moth borer in Puerto Rico. *Puerto Rico Sugar Manual* 1938: 7-9.
 1938b The second introduction of the Amazon fly from British Guiana into Puerto Rico. *Puerto Rico Agr. Expt. Sta. (Mayaguez) Agr. Notes* 86: 1-4.
 1939 The introduction and colonization of the Amazon fly, *Metagonistylum minense* Tns., in Puerto Rico. *Internatl. Soc. Sugar Cane Technol. Cong. Proc.* 6: 243-245.
 1941 The biology of *Metagonistylum minense* Tns., a parasite of the sugarcane borer. *Puerto Rico Expt. Sta. (Mayaguez) Bul.* 40: 1-20, figs. 1-10.
 1942 Biological control activities. *Puerto Rico Expt. Sta. (Mayaguez) Rpt.* 1941: 19-20.
 1943 Biological control activities. *Puerto Rico Expt. Sta. (Mayaguez) Rpt.* 1942: 15-17.

Basinger, A. J.

1935 Parasites reared from *Argyrotaenia (Tortrix) citrana* Fernald. Calif. Dept. Agr. Monthly Bul. 24 (4-6): 233-234.

1935a Notes on the parasites of the peach twig borer in southern California (*Anarsia lineatella* Zell[er]). Calif. Dept. Agr. Monthly Bul. 24 (4-6): 245-246.

Baumhofer, L. G. (See Graham, S. A., and Baumhofer)

Baver, L. D.

1950 Insects. Hawaii. Sugar Planters' Assoc. Expt. Sta. Rpt. 1950: 22-25, figs. 13-18.

Beacher, J. H.

1947 Studies of pistol case-bearer parasites. Ent. Soc. Amer. Ann. 40 (3): 530-544.

Bean, J. L.

1958 The use of larvaevorid maggot drop in measuring trends in spruce budworm populations. Ent. Soc. Amer. Ann. 51 (4): 400-403, figs. 1-2.

Beard, R. L.

1936 Further observations on the squash bug in Connecticut. Conn. Agr. Expt. Sta. Bul. 383: 333-339, figs. 107-109.

1940 Parasitic castration of *Anasa tristis* DeG. by *Trichopoda pennipes* Fab., and its effect on reproduction. Jour. Econ. Ent. 33: 269-272, figs. 1-2.

1940a The biology of *Anasa tristis* DeGeer with particular reference to the tachinid parasite, *Trichopoda pennipes* Fabr. Conn. Agr. Expt. Sta. Bul. 440: 595-679, figs. 1-18, pls. 1-3.

Beardsley, J. W. (Jr.) (See also Tanada and Beardsley)

1955 *Spodoptera mauritia* (Boisduval). Hawaii. Ent. Soc. Proc. 1954 15 (3): 386.

1957 *Heliothis virescens* (F.). Hawaii. Ent. Soc. Proc. 1956 16 (2): 190.

1964 A parasitic tachina fly (*Trichopoda pennipes* (Fabr.)). (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 14 (10): 150.

1964a A parasitic tachina fly (*Trichopoda pennipes* (F.)). (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 14 (18): 414.

1965 Tachinid eggs on Australian mantis. Hawaii. Ent. Soc. Proc. 1964 19 (1): 3.

- 1965a *Trichopoda pennipes* (F.) parasitizing koa bugs. Hawaii. Ent. Soc. Proc. 1964 19 (1): 8.
- 1966 *Thyanta accerra* (McAtee) a newly established immigrant. Hawaii. Ent. Soc. Proc. 1965 19 (2): 146.
- Beauchamp, C. E.
1950 El control del borer por medio de la mosca cubana *Lixophaga diatraeae*. Inst. Cubano de Suelos y Abanos Inform. Téc. Bol. 8 (3): 1151-1161.
- Beaulieu, G. (See Winn and Beaulieu)
- Beaulne, J. I.
1939 Parasites and predators reared at Quebec. Canad. Ent. 71 (5): 120.
- 1939a Liste des parasites et prédateurs récoltés au laboratoire d'Entomologie à Québec. Nat. Canad. 66 (3): 95-96.
- [1949] Notes sur quelques prédateurs et parasites d'importance économique. Quebec Soc. Protect. Plants Rpt. 1945-1947 30: 206-209.
- [1953] Amis et ennemis des cultures. Quebec Soc. Protect. Plants Rpt. 1950-1951 32-33: 194-197.
- *Becker, G. C.
1922 Report of meeting of cotton states entomologists. Jour. Econ. Ent. 15: 105-112.
- Becker, G. C., Jr. (See also Wilkinson, R. C., Becker, and Benjamin)
— and Benjamin, D. M.
1964 Notes on insect parasites and predators of the Swaine jack-pine sawfly in Wisconsin. Jour. Econ. Ent. 57: 355-357.
- 1964a A preliminary report on *Neodiprion nigroscutum* Midd., a closely related member of the *N. abbotii* complex. Ent. Soc. Amer., No. Cent. Branch, Proc. 19: 20-21.
- 1967 The biology of *Neodiprion nigroscutum* (Hymenoptera: Diprionidae) in Wisconsin. Canad. Ent. 99 (2): 146-159, figs. 1-8.
- Beckham, C. M. (See also Dupree and Beckham)
1952 A parasite of the vegetable weevil. Jour. Econ. Ent. 45: 541.
- 1952a Parasite of vegetable weevil discovered. Ga. Agr. Expt. Sta. Ann. Rpt. 1951 63: 38-39.
- 1953 Biology and control of the vegetable weevil in

- Georgia. Ga. Expt. Sta. Tech. Bul. 2: [1]-36, figs. 1-3.
- 1955 Biology and control of the vegetable weevil. Ohio State Univ., Abs. Doctoral Diss. 66: 9-11.
- Beckwith, C. S.**
- 1930 The blossom worm, a cranberry pest. N.Y. Ent. Soc. Jour. (1929) 37 (4): 409-416, 1 fig.
- 1938 *Sparganothis sulfureana* Clem., a cranberry pest in New Jersey. Jour. Econ. Ent. 31: 253-256.
- Bedard, W. D.**
- 1938 An annotated list of the insect fauna of Douglas fir (*Pseudotsuga mucronata* Rafinesque) in the northern Rocky Mountain region. Canad. Ent. 70 (9): 188-197.
- Begg, J. A.** (See Manson et al.)
- Béigüe, R.**
- 1961 Etude sur la mouche-à-scie du peuplier, *Trichio-campus viminalis* (Fall.) (Hyménoptère: Tenthredinidae). Canad. Ent. 93 (12): 1085-1097, figs. 1-11.
- Beirne, B. P.**
- 1962 Parasites of the cabbage butterfly. Canada Dept. Agr., Res. Branch, Ent. Res. Inst. Biol. Control, Belleville, Ontario, Res. Rpt. 1960-1961: 19.
- 1962a Parasitism of grasshoppers. Canada Dept. Agr., Res. Branch, Ent. Res. Inst. Biol. Control, Belleville, Ontario, Res. Rpt. 1960-1961: 19.
- Bell, J. C., Jr.** (See Ciesla, Rauschenberger, and Bell)
- Beneway, D. F.**
- 1963 A revision of the flies of the genus *Lespesia* (= *Achaetoneura*) in North America (Diptera: Tachinidae). Kans. Univ. Sci. Bul. 44 (13): 627-686, figs. 1-34.
- Benjamin, D. M.** (See also Becker, G. C., Jr., and Benjamin; Dixon and Benjamin; Drooz and Benjamin; Kapler and Benjamin; Walgenbach and Benjamin; Wilkinson, R. C., Becker, and Benjamin)
- 1955 The biology and ecology of the red-headed pine sawfly. U.S. Dept. Agr. Tech. Bul. 1118: 1-57, figs. 1-24.
- and Drooz, A. T.
- 1954 Parasites affecting the jack-pine budworm in Michigan. Jour. Econ. Ent. 47: 588-591.

Bentley, G. M.

- 1940 Red-headed pine sawfly (*Neodiprion lecontei* Fitch).
Insect Pest Survey Bul. 20 (4): 191.

Bergenstamm, J. E. v. (See Brauer and Bergenstamm)**Berger, E. W.** (See Watson and Berger)**Berry, P. A.**

- 1939 Biology and habits of *Ephialtes turionellae* (L.), a pupal parasite of the European pine shoot moth.
Jour. Econ. Ent. 32: 717-721.

Bess, H. A.

- 1936 The biology of *Leschenaultia exul* Townsend, a tachinid parasite of *Malacosoma americana* Fabricius and *Malacosoma disstria* Hubner. Ent. Soc. Amer. Ann. 29 (4): 593-613, pls. 1-5.

- 1937 Family Tachinidae. Methods used in rearing *Leschenaultia exul*. Pp. 411-412. In Needham, J. G., Galtsoff, P. S., Lutz, F. E., and Welch, P. S., Culture methods for invertebrate animals. xxxii + 590 pp., 84 figs. Ithaca, N.Y.

- 1951 *Laphygma exempta* (Walker). Hawaii. Ent. Soc. Proc. 1950 14 (2): 212.

Bethune, C. J. S.

- 1871 Insects affecting the apple. Ontario, Noxious Insects, Ann. Rpt. 1: 3-28, figs. 1-33.

Beyer, A. H. (See Luginbill and Beyer)**Bianchi, F. A.**

- 1940 Notes on the role of the self-introduced insects in the economic entomology of Hawaii. Hawaii. Ent. Soc. Proc. 1939 10 (3): 377-388.

- 1946 Parasites of *Achaea janata* (Linn.). Hawaii. Ent. Soc. Proc. 1945 12 (3): 469.

- 1949 *Chelonus texanus* Cresson. Hawaii. Ent. Soc. Proc. 1948 13 (3): 328.

- 1949a Recent changes in the parasite complex of armyworms. Hawaii. Ent. Soc. Proc. 1948 13 (3): 345.

- 1958 *Laphygma exempta* (Walker). Hawaii. Ent. Soc. Proc. 1957 16 (3): 319.

Bibby, F. F.

- 1942 Some parasites of *Heliothis armigera* (Hbn.) in Texas. Jour. Econ. Ent. 35: 943-944.

- 1961 Notes on miscellaneous insects of Arizona. Jour. Econ. Ent. 54: 324-333.

*Bird, F. T., and Burk, J. M.

- 1961 Artificially disseminated virus as a factor controlling the European spruce sawfly, *Diprion hercyniae* (Htg.) in the absence of introduced parasites. *Canad. Ent.* 93 (3): 228-238.

— and Elgee, D. E.

- 1957 A virus disease and introduced parasites as factors controlling the European spruce sawfly, *Diprion hercyniae* (Htg.), in central New Brunswick. *Canad. Ent.* 89 (8): 371-378, figs. 1-3.

Bird, H.

- 1909 New histories and species in *Papaipema* (*Hydroecia*). *Canad. Ent.* 41 (2): 57-68.
- 1914 New species and histories in *Papaipema* Sm. (Lepid.). *Canad. Ent.* 46 (2): 69-73, pl. 3.
- 1923 Life-histories in the genus *Schinia* and allies (Lepid.: Noctuidae) No. 1. *Ent. News* 34 (7): 193-199.
- 1923a Some adaptation of indigenous parasites to introduced pests. *N.Y. Ent. Soc. Jour.* (1922) 30 (4): 200.
- 1925 New life histories in *Papaipema* No. 23 (Lepidoptera). *Canad. Ent.* 57 (12): 303-306.
- 1927 The fern-feeding species of *Papaipema*; No. 25 (Lepidoptera). *Canad. Ent.* 59 (4): 92-96.
- 1936 *Papaipema stenoscelis* Dyar, a rare noctuid of the eastern United States (Lepidoptera). *Canad. Ent.* 68 (6): 119-121, 1 pl.

— and Jones, F. M.

- 1921 A new *Papaipema* from Delaware (Lepidoptera, Noctuidae). *Canad. Ent.* 53 (6): 137-139, pl. 6.

Bird, R. D., and Mitchener, A. V.

- 1937 Insects of the season 1936 in Manitoba. *Canad. Insect Pest Rev.* 15 (1): 42-46.
- 1938 Insects of the season 1937 in Manitoba. *Canad. Insect Pest Rev.* 16 (1): 47-52.
- 1947 Colorado potato beetle (*Leptinotarsa decemlineata* Say) Manitoba. *Canad. Insect Pest Rev.* 25 (1): 37.
- 1951 Insects of the season 1950 in Manitoba. *Canad. Insect Pest Rev.* 29 (1): 46-57.
- 1952 Insects of the season 1951 in Manitoba. *Canad. Insect Pest Rev.* 30 (1): 43-55.
- 1953 Insects of the season 1952 in Manitoba. *Canad. Insect Pest Rev.* 31 (1): 40-52.

- Bischoff, H.
1909 Neue Beiträge zur Lebensweise der Trigonaloiden. Berlin. Ent. Ztschr. 54: 76-80, 1 fig.
- Bishopp, F. C.
1923 The puss caterpillar and the effects of its sting on man. U.S. Dept. Agr., Dept. Cir. 288: 1-14, figs. 1-12.
- Bissell, T. L. (See also Cory et al.)
1940 Curculionidae, Bruchidae, Lepidoptera, and their parasites, infesting the seed pods of cowpea and various wild plants. Jour. Econ. Ent. 33: 844-847, fig. 1.
1942 Life history of *Pseudoclista globosa* Tns., parasite of the cowpea curculio. Ga. Ent. 1 (1): 8-9.
1945 *Myiophasia globosa* (Tns.), tachinid parasite of the cowpea curculio. Ent. Soc. Amer. Ann. 38 (3): 417-440, pls. 1-5.
1953 Summary of insect conditions — 1952: Maryland. (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 3 (8): 126-129.
- Blais, J. R. (See also McGugan and Blais)
1960 Spruce budworm parasite investigations in the lower St. Lawrence and Gaspé regions of Quebec. Canad. Ent. 92 (5): 384-396, fig. 1.
1965 Parasite studies in two residual spruce budworm (*Choristoneura fumiferana* (Clem.)) outbreaks in Quebec. Canad. Ent. 97 (2): 129-136.
- Blakeslee, E. B. (See Brooks, F. E., and Blakeslee)
- Blanchard, R. A., and Conger, C. B.
1932 Notes on *Prodenia praefica* Grote. Jour. Econ. Ent. 25: 1059-1070.
- Blickenstaff, C. C. (See also Drake, Harris, and Blickenstaff)
1970 Common names of insects approved by the Entomological Society of America. 36 pp. [College Park, Md.].
- Arbuthnot, K. D., and Harris, H. M.
1953 Parasites of the European corn borer in Iowa. Iowa State Col. Jour. Sci. 27 (3): 335-379, figs. 1-9.
- Bloeser, W.
1914 Notes on the life history and anatomy of *Siphona plusiae* Coq. Ent. Soc. Amer. Ann. 7 (4): 301-309, figs. 1-2, pls. 39-40.

- Blunck, H.
1957 *Pieris rapae* (L.), its parasites and predators in Canada and the United States. Jour. Econ. Ent. 50: 835-836.
- Boardman, E. T.
1939 Field guide to lower aquarium animals. Cranbrook Inst. Sci. Bul. 16: 1-186, figs. 1-51.
- Bobb, M. L.
1963 Insect parasites of the Virginia pine sawfly, *Neodiprion pratti pratti*. Jour. Econ. Ent. 56: 618-621, fig. 1.
- Bohart, R. M.
1947 Sod webworms and other lawn pests in California. Hilgardia 17 (8): 267-308, figs. 1-20.
- Bosch, R. van den (See also Dietrick and Bosch)
—— and Hagen, K. S.
1966 Predaceous and parasitic arthropods in California cotton fields. Calif. Univ. Agr. Expt. Sta. Bul. 820: 1-32, figs. 1-20.
- Bottimer, L. J.
1926 Notes on some Lepidoptera from eastern Texas. Jour. Agr. Res. 33 (9): 797-819, figs. 1-3.
- Bottomley, A. W. T. (See Paxton et al.)
- Bottrell, D. G.
1969 Notes on parasites attacking the yellow-striped armyworm in Oklahoma. Ent. Soc. Amer. Ann. 62 (1): 250-252.
- Young, J. H., Price, R. G., and Adams, R. H.
1968 Parasites reared from *Heliothis* spp. in Oklahoma in 1965 and 1966. Ent. Soc. Amer. Ann. 61 (5): 1053-1055.
- Bourne, B. A.
1921 Insect attacks reported or observed. Barbados Dept. Agr. Rpt. 1919-1920: 12-14.
- Box, H. E.
1925 Porto Rican cane-grubs and their natural enemies, with suggestions for the control of lamellicorn larvae by means of wasp-parasites (Scoliidae). Porto Rico Dept. Agr. Jour. 9 (4): 291-356, figs. 1-21.
- 1935 The species of *Diatraea* attacking sugar cane in the New World. Internatl. Soc. Sugar Cane Technol. Cong. Proc. 5: 470-476.

- 1939 Biological control of *Diatraea saccharalis* (Fabricius) in St. Lucia, B.W.I. Internatl. Soc. Sugar Cane Technol. Cong. Proc. 6: 223-240, fig. 1.
- 1939a Some aspects of the campaign against the moth borer (*Diatraea saccharalis* Fabr.) in Antigua and St. Kitts, 1931-1938. Internatl. Soc. Sugar Cane Technol. Cong. Proc. 6: 495-513.
- 1952 *Palpozenillia palpalis* (Aldr.), a tachinid parasite of sugar cane moth borers (*Castnia* and *Diatraea*). Hawaii. Ent. Soc. Proc. 1951 14 (3): 485-490, figs. 1-2.
- 1953 List of sugar-cane insects. 101 pp. London.
- 1954 A preliminary list of the insects affecting sugarcane in the Lesser Antilles and Trinidad. Internatl. Soc. Sugar Cane Technol. Cong. Proc. 8: 549-553.
- 1960 Status of the moth borer, *Diatraea saccharalis* (F.), and its parasites in St. Kitts, Antigua and St. Lucia; with observations on Guadeloupe and an account of the situation in Haiti. Internatl. Soc. Sugar Cane Technol. Cong. Proc. 10: 901-914.
- Boyce, H. R. (See Steenburgh and Boyce)
- Boyes, J. W., and Wilkes, A.
- 1953 Somatic chromosomes of higher Diptera. I. Differentiation of tachinid parasites. Canad. Jour. Zool. 31 (2): 125-165, figs. 1-14, pls. 1-2.
- Bracken, D. F., and Martineau, R.
- [1961] Larch sawfly (*Pristiphora erichsonii* Htg.). Canad. Insect Pest Rev. (1960) 38 (7): 258-260.
- *Bradley, G. A.
- 1951 Notes on the parasitism of the yellow-headed spruce sawfly, *Pikonema alaskensis* (Roh.) (Hymenoptera: Tenthredinidae). Canad. Ent. 83 (5): 130-131.
- Bradley, W. G. (See also Baker, W. A., and Bradley; Baker, W. A., Bradley, and Clark; Jones, T. H., and Bradley)
- and Clark, C. A.
- 1938 The field status of parasites of the European corn borer in the fall of 1937. Insect Pest Survey Bul. 18 (9) Sup.: 649-656.
- 1939 The field status of parasites of the European corn borer in the fall of 1938. Insect Pest Survey Bul. 19 (8) Sup.: 527-534.

- 1940 The field status of parasites of the European corn borer at the close of the 1939 season. *Insect Pest Survey Bul.* 20 (9) Sup.: 525-532.
- 1941 The field status of parasites of the European corn borer at the close of the 1940 season. *Insect Pest Survey Bul.* 21 (9) Sup.: 787-794.
- Bratley, H. E.**
- 1929 Notes on *Lymire edwardsi* Grote, the rubber tree caterpillar. *Fla. Ent.* 13 (3): 44.
- 1933 Larger plant bugs on citrus and truck crops. *Fla. Agr. Expt. Sta. Ann. Rpt.* 1933: 77.
- 1934 The larger plant bugs. *Fla. Agr. Expt. Sta. Ann. Rpt.* 1934: 52.
- 1936 The larger plant bugs. *Fla. Agr. Expt. Sta. Ann. Rpt.* 1936: 65-66.
- 1937 The larger plant bugs. *Fla. Agr. Expt. Sta. Ann. Rpt.* 1937: 72.
- 1938 The feather-legged fly — characterized as a friend of the farmer after extensive study of its capabilities as a destroyer of plant bugs. *Citrus Indus.* 19 (9): 13-14.
- 1939 The larger plant bugs. *Fla. Agr. Expt. Sta. Ann. Rpt.* 1939: 95.
- 1943 Feather-legged fly aids grower. Winged "commando" attacks leading plant enemy. *Fla. Grower* 51 (8): 6.
- Brauer, F., and Bergenstamm, J. E. v.**
- 1895 Die Zweiflügler des Kaiserlichen Museums zu Wien. VII. Vorarbeiten zu einer Monographie der Muscaria Schizometopa (exclusive Anthomyidae). Pars IV. K. Akad. der Wiss. Wien, Math.-Nat. Cl. Denkschr. 61: 537-624. (Also published as an 88 page separate in 1894. Since the 1894 separate is rare the 1895 publication has been cited in this work.)
- Brazzel, J. R.** (See Lewis, W. J., and Brazzel)
- Breakey, E. P.**
- 1929 Notes on the natural enemies of the iris borer, *Macronoctua onusta* Grote (Lepidoptera). *Ent. Soc. Amer. Ann.* 22 (3): 459-464.
- 1931 Additional notes on the natural enemies of the iris borer, *Macronoctua onusta* Grote (Lepidoptera). *Ent. Soc. Amer. Ann.* 24 (1): 40-44.

Breeland, S. G.

- 1958 Biological studies on the armyworm, *Pseudaletia unipuncta* (Haworth), in Tennessee (Lepidoptera: Noctuidae). Jour. Tenn. Acad. Sci. 33 (4): 263-347, figs. 1-26.

Bregger, T. (See Hayslip, Mathes, and Bregger; Thames et al.; Wilson, J. W., and Bregger)

Brehme, H. H.

- 1915 Parasite of *Bellura obliqua* G. and R. (Lep., Dip.). Ent. News 26 (10): 473.

Breland, O. P.

- 1940 Some parasites and hyperparasites of the cecropia moth. N.Y. Ent. Soc. Jour. 48 (3): 259-264.

Breshears, J. C., and Dawson, J. C.

- 1936 The Japanese beetle including report of progress in safeguarding Missouri agriculture from this destructive oriental insect pest. Mo. State Dept. Agr. Bul. 34 (8): 1-23, figs. 1-12.

Brewer, J., and Thomas, G. M.

- 1966 Causes of death encountered during rearing of *Danaus plexippus* (Danaidae). Lepidopterists' Soc. Jour. 20 (4): 235-238.

Briand, L. J.

- 1936 Shipping of potato beetle parasites and predators to France, with notes on the species involved. Ontario Ent. Soc. Ann. Rpt. 1935 66: 31-34.
- [1949] Spruce sawfly parasite investigations Parke Reserve, St. Alexandre, Kamouraska, Que. Quebec Soc. Protect. Plants Rpt. 1945-1947 30: 54-59.
- [1949a] Field studies on *Sturmia* sp., an important parasite of the European spruce sawfly. Quebec Soc. Protect. Plants Rpt. 1945-1947 30: 125-127.
- [1949b] Notes on the spruce sawfly (*Gilpinia hercyniae* Htg.) and parasitism in the Parke Reserve, Quebec, area. Quebec Soc. Protect. Plants Rpt. 1945-1947 30: 180-181.
- 1958 Note on overwintering of *Adoryphorophaga aberrans* (Townsend), a tachinid parasite of the Colorado potato beetle. Ontario Ent. Soc. Ann. Rpt. 88: 57.

Bridwell, J. C.

- 1919 Descriptions of new species of hymenopterous parasites of muscoid Diptera with notes on their

- habits. Hawaii. Ent. Soc. Proc. 1918 4 (1): 166-179.
- 1920 A dipterous parasite of *Corizus hyalinus*. Hawaii. Ent. Soc. Proc. 1919 4 (2): 339-340.
- Brimley, C. S.**
- 1922 List of the Tachinidae (Diptera) of North Carolina. Ent. News 33 (1): 20-26.
- 1928 Some new wasps (Hymenoptera) and two new Diptera from North Carolina. Elisha Mitchell Sci. Soc. Jour. 43 (3-4): 199-206.
- 1938 The insects of North Carolina. 560 pp. Raleigh.
- 1942 Supplement to insects of North Carolina. 39 pp. Raleigh.
- Brindley, T. A.** (See York, Schaffner, and Brindley)
- Brittain, W. H., and Payne, H. G.**
- [1919] Some notes on *Olene vagans* B. and McD. in Nova Scotia. Ent. Soc. Nova Scotia, Proc. (1918) 4: 62-68, pl. 4.
- Britton, W. E.**
- 1910 The brown-tail moth, *Euproctis chrysorrhoea* Linn. Conn. Agr. Expt. Sta. Bul. 182: 1-26, figs. 1-16.
- 1911 Tenth report of the state entomologist of Connecticut for the year 1910. Conn. Agr. Expt. Sta. Bien. Rpt. 1909-1910 9: i-vi, 657-712, figs. 10-23, pls. 25-32.
- 1912 Eleventh report of the state entomologist of Connecticut. Conn. Agr. Expt. Sta. Rpt. 1911 35: 259-346, figs. 1-10, pls. 1-16.
- 1913 Outbreak of the fall army worm *Laphygma frugiperda* S. & A. Conn. Agr. Expt. Sta. Ann. Rpt. 1912 36: 284-287.
- 1915 The gypsy moth. Conn. Agr. Expt. Sta. Bul. 186: 1-24, figs. 1-16.
- 1915a Fourteenth report of the state entomologist of Connecticut. Conn. Agr. Expt. Sta. Ann. Rpt. 1914 38: 113-198, pls. 9-24.
- 1916 Further notes on *Diprion simile* Hartig. Jour. Econ. Ent. 9: 281-282.
- 1916a Fifteenth report of the state entomologist of Connecticut. Conn. Agr. Expt. Sta. Ann. Rpt. 1915 39: 81-192, figs. 1-6, pls. 1-16.
- 1917 Sixteenth report of the state entomologist of Connecticut. Conn. Agr. Expt. Sta. Ann. Rpt. 1916 40: 65-146, figs. 1-3, pls. 1-16.

- 1919 Insects attacking squash, cucumber, and allied plants in Connecticut. Conn. Agr. Expt. Sta. Bul. 216: 31-51.
- 1920 Nineteenth report of the state entomologist of Connecticut for the year 1919. Conn. Agr. Expt. Sta. Bul. 218: 107-208, pls. 9-32.
- 1922 Twenty-first report of the state entomologist of Connecticut for the year 1921. Conn. Agr. Expt. Sta. Bul. 234: 111-204, figs. 1-6, pls. 1-16.
- 1926 Twenty-fifth report of the state entomologist of Connecticut 1925. Conn. Agr. Expt. Sta. Bul. 275: 212-330, figs. 38-50, pls. 1-20.
- 1927 Twenty-sixth report of the state entomologist of Connecticut 1926. Conn. Agr. Expt. Sta. Bul. 285: 161-283, figs. 1-11, pls. 1-16.
- 1935 The gypsy moth. Conn. Agr. Expt. Sta. Bul. 375: 621-647, figs. 77-94.
- 1935a The eastern tent caterpillar *Malacosoma americana* Fabr. Conn. Agr. Expt. Sta. Bul. 378: 63-82, figs. 1-22.
- and Ashworth, J. T.
- 1923 Report of work in suppressing the gypsy and brown-tail moths. Conn. Agr. Expt. Sta. Bul. 247: 290-326, figs. 4-5.
- 1933 Report of gypsy moth control in Connecticut in 1932. Conn. Agr. Expt. Sta. Bul. 349: 400-407, figs. 106-107.
- and Davis, I. W.
- 1916 Brown-tail moth work, season of 1914-1915. Conn. Agr. Expt. Sta. Ann. Rpt. 1915 39: 111-114.
- 1918 Suppression work against the gypsy and brown-tail moths. Conn. Agr. Expt. Sta. Bul. 203: 246-258.
- and Johnson, J. P.
- 1938 The Japanese beetle in Connecticut. Conn. Agr. Expt. Sta. Bul. 411: [450]-486, figs. 69-85.
- and Zappe, M. P.
- 1918 The imported pine sawfly, *Diprion (Lophyrus) simile* Hartig. Conn. Agr. Expt. Sta. Bul. 203: 273-290, pls. 6-9.
- Brodie, D. A. (See Doane and Brodie)
- Brodie, W.
- 1878 Experiments on the Colorado potato beetle. Ontario Ent. Soc. Ann. Rpt. 1877: 9-10.
- Bromley, S. W. (See Felt and Bromley)

***Brooks, A. R.**

- 1942 *Clistomorpha, Psalidopteryx* and allies (Diptera, Tachinidae). *Canad. Ent.* 74 (8): 140-150, figs. 1-4.
- 1944 A review of the North American species of *Linnaemya* sens. lat. (Diptera, Tachinidae). *Canad. Ent.* 76 (10): 193-206, pl. 12.
- 1945 New Canadian Diptera (Tachinidae). *Canad. Ent.* 77 (5): 78-96.
- 1945a A revision of the North American species of the *Phasia* complex (Diptera, Tachinidae). *Sci. Agr.* 25 (11): 647-679, pls. 1-4.
- 1946 The genus *Girschneria* Townsend (Diptera, Tachinidae). *Canad. Ent.* (1945) 77 (10): 184-185, fig. 1.
- 1946a A revision of the North American species of the *Rhodogyne* complex (Diptera, Larvaevoridae). *Canad. Ent.* (1945) 77 (12): 218-230, pls. 10-11.
- 1947 A revision of the North American species of *Leschenaultia* sens. lat. (Diptera, Larvaevoridae). *Canad. Ent.* (1946) 78 (9-10): 169-182, pls. 14-16.

Brooks, F. E.

- 1909 The grapecane gall-maker and the grapecane girdler. *W. Va. Agr. Expt. Sta. Bul.* 119: 319-339, frontis., 5 pls.
- 1910 Snout beetles that injure nuts. *W. Va. Agr. Expt. Sta. Bul.* 128: 143-185, frontis., 10 figs., 6 pls.
- 1911 The grapecane gall-maker and the grapecane girdler. *W. Va. Expt. Sta. Rpt.* 1909-1910: 62-79, 7 figs.
- 1911a Snout beetles that injure nuts. *W. Va. Expt. Sta. Rpt.* 1909-1910: 160-193, 15 figs.

—— and Blakeslee, E. B.

- 1915 Studies of the codling moth in the central Appalachian region. *U.S. Dept. Agr. Bul.* 189: 1-49, figs. 1-23, pl. 1.

—— and Cotton, R. T.

- 1929 The chestnut curculios. *U.S. Dept. Agr. Tech. Bul.* 130: 1-23, fig. 1, pls. 1-6.

Brooks, R. F. (See Rings and Brooks)**Brown, A. W. A.**

- 1938 The forest insect survey. *Ontario Ent. Soc. Ann. Rpt.* 1937 68: 13-18.

***Brown, R. C.**

- 1937 Oriental moth. Insect Pest Survey Bul. 17
(10): 626.
- 1940 [Recovery of *Parasetigena silvestris* R. D. in
Massachusetts]. Insect Pest Survey Bul. 20
(10): 585.

— and Sheals, R. A.

- 1944 The present outlook on the gypsy moth problem.
Jour. Forestry 42 (6): 393-407, figs. 1-9.

Brown, W. J.

- 1957 The New World species of *Chrysomela* L. (Coleop-
tera: Chrysomelidae). Canad. Ent. (1956) 88 (3)
Sup.: 1-54, figs. 1-40.

Brubaker, R. W.

- 1968 Seasonal occurrence of *Voria ruralis*, a parasite of
the cabbage looper, in Arizona, and its behavior
and development in laboratory culture. Jour.
Econ. Ent. 61: 306-309.

Brues, C. T. (See also Quaintance and Brues)

- 1903 A dexiid parasite of the sow-bug. Ent. News 14
(9): 291.

***Bruner, L.**

- 1889 On certain injurious insects of the year 1888. Nebr.
Agr. Expt. Sta. Bul. 1 (5): 135-172, figs. 1-19.
- 1890 Insects injurious to young trees on tree claims. Nebr.
Agr. Expt. Sta. Bul. 3 (14): 83-149, figs. 1-98.
- 1890a Report on a local outbreak of grasshoppers in Idaho.
Insect Life (U.S. Dept. Agr., Div. Ent., Period.
Bul.) 3 (4): 135-141.
- 1892 Report upon insect depredations in Nebraska for
1891. U.S. Dept. Agr., Div. Ent. Bul. 26: 9-12.
- 1897 Grasshopper report for 1896. U.S. Dept. Agr. Div.
Ent. Bul., n. ser., 7: 36-39.

Bruner, S. C.

- 1934 La palomilla del maiz. Nota de divulgacion agricola.
[Cuba] Rev. de Agr., Com. y Trab. 14
(52): 94-97, 2 figs.
- 1935 La maruca y otros insectos de las habas de lima.
Cuba Estac. Expt. Agron. Bol. 56: 1-52, 24 figs.

Bruneteau, J.

- 1937 Recherches sur les ennemis naturels du doryphore
en Amérique. Ann. des Epiphyt. et de Phyto-
génét., n. ser., 3 (1): 113-135, figs. 1-14.

Brunson, M. H., and Allen, H. W.

1948 Oriental fruit moth cocoon parasites. Jour. Econ. Ent. 41: 446-450.

— and Coles, L. W.

1968 The introduction, release, and recovery of parasites of the alfalfa weevil in eastern United States. U.S. Dept. Agr., Prod. Res. Rpt. 101: 1-12.

Bryan, D. E. (See also Butler, G. D., Jr., Bryan, and Jackson; Jackson, C. G., Bryan, and Patana; Jackson, C. G., Butler, and Bryan)

— Jackson, C. G., and Patana, R.

1968 Laboratory studies of *Lespesia archippivora* in four lepidopterous hosts. Jour. Econ. Ent. 61: 819-823.

1969 Effect of temperature on the progeny production and longevity of *Lespesia archippivora* in the laboratory. Jour. Econ. Ent. 62: 765-767.

— Jackson, C. G., and Stoner, A.

1969 Rearing cotton insect parasites in the laboratory. U.S. Dept. Agr., Prod. Res. Rpt. 109: 1-13, figs. 1-10.

*Bryan, E. H., Jr.

1934 A review of the Hawaiian Diptera, with descriptions of new species. Hawaii. Ent. Soc. Proc. 1933 8 (3): 399-468.

Buchanan, W. D. (See Dowden, Buchanan, and Carolin)

Buck, J., and Keister, M.

1956 Host-parasite relations in *Agapema* pupae (Lepidoptera, Saturniidae). Ent. Soc. Amer. Ann. 49 (1): 94-97.

Buckner, C. H. (See also Ives et al.; Nairn et al.)

1959 Mortality of cocoons of the larch sawfly, *Pristiphora erichsonii* (Htg.), in relation to distance from small-mammal tunnels. Canad. Ent. 91 (9): 535-542, figs. 1-6.

Burbutis, P. P. (See Van Denburgh, Burbutis, and York; White-side, Burbutis, and Kelsey)

Burgess, A. F.

1897 Notes on certain Coleoptera known to attack the gypsy moth. Mass. State Bd. Agr. Ann. Rpt. Sec. 1896 44: 412-433, pls. 1-5.

1914 The gypsy moth and the brown-tail moth, with suggestions for their control. U.S. Dept. Agr. Farmers' Bul. 564: 1-24, figs. 1-10.

- 1915 Parasite work. Report of Mr. A. F. Burgess, in charge of moth work, parasite laboratory, Melrose Highlands, Mass. Mass. State Forester Ann. Rpt. 1914 11: 84-86.
- 1915a Report on the gipsy moth work in New England. U.S. Dept. Agr. Bul. 204: 1-32, figs. 1-3, pls. 1-8.
- 1916 The work carried on in the United States against the gipsy and brown-tail moths. Ontario Ent. Soc. Ann. Rpt. 1915 46: 153-155.
- 1917 The gipsy moth and the brown-tail moth and their control. U.S. Dept. Agr. Farmers' Bul. 845: 1-28, figs. 1-13.
- 1924 The value of natural enemies of injurious insects. Ontario Ent. Soc. Ann. Rpt. 54: 30-36.
- 1930 The gipsy moth and the brown-tail moth. U.S. Dept. Agr. Farmers' Bul. 1623: i-ii, 1-33, figs. 1-19.
- and Collins, C. W.
- 1915 The calosoma beetle (*Calosoma sycophanta*) in New England. U.S. Dept. Agr. Bul. 251: 1-40, figs. 1-3, pls. 1-7.
- 1917 The genus *Calosoma*: including studies of seasonal histories, habits, and economic importance of American species north of Mexico and of several introduced species. U.S. Dept. Agr., Dept. Bul. 417: 1-124, figs. 1-5, pls. 1-19.
- and Crossman, S. S.
- 1927 The satin moth, a recently introduced pest. U.S. Dept. Agr., Dept. Bul. 1469: 1-22, figs. 1-5, pl. 1.
- 1929 Imported insect enemies of the gipsy moth and the brown-tail moth. U.S. Dept. Agr. Tech. Bul. 86: 1-147, figs. 1-55, pls. 1-6.
- Burk, J. M. (See Bird, F. T., and Burk)
- Burke, H. E.
- 1919 Notes on the California oak worm, *Phryganidia californica*, (Lepid.). Ent. Soc. Wash. Proc. 21 (6): 124-125.
- and Herbert, F. B.
- 1920 California oak worm. U.S. Dept. Agr. Farmers' Bul. 1076: [1]-14, figs. 1-9.
- Burkett, J. H.
- 1932 The pecan in Texas pecan soils, districts, streams, orchards, groves, care, propagation, cultivation, covercrops, irrigation, production, insects,

- diseases, etc. Tex. Dept. Agr. Bul. 111: 1-239, figs. 1-79.
- Burkhardt, C. C.
1953 Insects injurious to corn. Kans. Agr. Expt. Sta. Dir. Rpt. 16: 67.
- Burrell, R. W.
1931 *Dexia ventralis* Aldrich, an imported parasite of the Japanese beetle. Jour. Agr. Res. 43 (4): 323-336, figs. 1-7.
1967 Parasites of the armyworm in Louisiana. Jour. Econ. Ent. 60: 111-114.
- Bussart, J. E.
1937 The bionomics of *Chaetophleps setosa* Coquillett (Diptera: Tachinidae). Ent. Soc. Amer. Ann. 30 (2): 285-295, pls. 1-2.
- *Butler, G. D., Jr. (See also Jackson, C. G., Butler, and Bryan)
1958 Tachinid flies reared from lepidopterous larvae in Arizona, 1957. Jour. Econ. Ent. 51: 561-562.
1961 Two parasites of *Hylephila phyleus* Drury in Arizona (Lepidoptera: Hesperidae). Pan-Pacific Ent. 37 (4): 202.
- Bryan, D. E., and Jackson, C. G.
1968 Development of the salt-marsh caterpillar parasite, *Exorista mella* at controlled constant and variable temperatures in the laboratory. Jour. Econ. Ent. 61: 161-162.
- Butler, H. G.
1933 Larval parasites of the oriental fruit moth in Roane County, Tennessee. Jour. Econ. Ent. 26: 982-987.
- Bynum, E. K. (See Ingram and Bynum; Ingram, Bynum, and Douglas; Ingram, Bynum, and Mathes; Ingram et al.)
- Caceres, S. Flores (See Flores Caceres, S.)
- Caesar, L.
1916 Insects of the season in Ontario. Ontario Ent. Soc. Ann. Rpt. 1915 46: 29-33.
1916a Leaf-rollers attacking apples. Ontario Ent. Soc. Ann. Rpt. 1915 46: 163-178, 10 figs.
- Caffrey, D. J. (See also Vinal and Caffrey; Wildermuth and Caffrey)
1916 Current notes. Jour. Econ. Ent. 9: 250.
1919 The European corn borer: a menace to the country's corn crop. U.S. Dept. Agr. Farmers' Bul. 1046: 1-28, figs. 1-16.

— and Worthley, L. H.

- 1927 The European corn borer: its present status and methods of control. U.S. Dept. Agr. Farmers' Bul. 1548: 1-48, figs. 1-42.

Cameron, J. M.

- 1931 The apple and thorn-leaf skeletonizer (*Hemerophila pariana* Clerck). Quebec Soc. Protect. Plants Ann. Rpt. 22: 37-43.

Capelouto, R.

- 1949 Some studies on the parasitic relationship between the feather-legged tachina fly, *Trichopoda pennipes* Fab. and the southern green stink bug, *Nezara viridula* Linne. Newell Ent. Soc. Jour. 4 (1): 31-33, 1 fig.

Cardin, P.

- 1915 Informe del departamento de patologia vegetal y entomologia. Cuba Estac. Expt. Agron. Informe 3: 98-173, 5 pls.

Carlson, F. W. (See Newcomer and Carlson)

Carnes, E. K., and Newcomer, E. J.

- 1912 An insect enemy of pasture-land and growing alfalfa. Calif. State Comn. Hort. Monthly Bul. 1 (7): 275-280, figs. 120-122.

Carolin, V. M. (See also Dowden, Buchanan, and Carolin; Dowden and Carolin; Dowden, Jaynes, and Carolin)

— and Coulter, W. K.

- 1959 The occurrence of insect parasites of *Choristoneura fumiferana* (Clem.), in Oregon. Jour. Econ. Ent. 52: 550-555.

Carrillo S., J. L. (See also Gibson, W. W., and Carrillo S.)

— Ortega C., A., and Gibson, W. W.

- 1966 Lista de insectos en la colección entomológica del Instituto Nacional de Investigaciones Agrícolas. Primer suplemento a la "Lista de insectos de la colección entomológica de la Oficina de Estudios Especiales, S. A. G.". Mex., S A G, Inst. Nac. de Invest. Agr., Fol. Misc. 14: i-viii, 1-133.

Carroll, W. J.

- 1956 History of the hemlock looper, *Lambdina fiscellaria fiscellaria* (Guen.), (Lepidoptera: Geometridae) in Newfoundland, and notes on its biology. Canad. Ent. 88 (10): 587-599, figs. 1-3.

Carter, S. W. (See Arbuthnot et al.; Arbuthnot et al.; Clark, C. A., and Carter; Jones, D. W., Carter, and Cosenza; Sparks, Raun, and Carter)

Cartwright, W. B. (See Ainslie, G. G., and Cartwright)

Catoni, L. A.

- 1922 Plagas de insectos que atacan a las plantaciones de batata. Rev. Agr. de Puerto Rico 9 (3): 25-28.

Caulfield, F. B.

- 1887 Some further notes on *Physonota*. Canad. Ent. 19 (4): 73-76.

Cavitt, H. S. (See Chapman, Fife, and Cavitt)

Chada, H. L. (See Rice, Wolfenbarger, and Chada)

Chamberlin, F. S. (See also Madden and Chamberlin)

— and Madden, A. H.

- 1942 Insect pests of cigar-type tobaccos in the southern districts. U.S. Dept. Agr., Dept. Cir. 639: 1-54, figs. 1-57.

Chamberlin, T. R. (See Rockwood and Chamberlin)

*Chamberlin, W. J.

- 1931 Remarks on the genus *Ellopiia* (order Lepidoptera, family Geometridae) with special reference to the oak looper, *E. somniaria* Hulst and the hemlock looper, *E. fervidaria* Hubner. Jour. Econ. Ent. 24: 1036-1041, pls. 42-43.
- 1932 Notes on the synonymy [sic], hosts and distribution of some Buprestidae (Coleoptera). Jour. Econ. Ent. 25: 833-836, fig. 52.

Chambers, E. L.

- 1952 European corn borer (*Pyrausta nubilalis*) — Wisconsin. (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 2 (17): 233.
- 1952a European corn borer (*Pyrausta nubilalis*) — Wisconsin. (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 2 (19): 267.
- 1952b European corn borer (*Pyrausta nubilalis*) — Wisconsin. (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 2 (27): 357.

Champlain, A. B., and Knull, J. N.

- 1923 Notes on Pennsylvania Diptera. Ent. News 34 (7): 211-215.

Chant, D. A. (See McLeod, J. H., and Chant)

Chapman, A. J., Fife, L. C., and Cavitt, H. S.

- 1935 Cutworms (Noctuidae). Insect Pest Survey Bul. 15 (6): 305.

Charpentier, L. J. (See also Ingram et al.)

- 1956 1954 studies of parasites for sugarcane borer control in Louisiana. Jour. Econ. Ent. 49: 267-268.

- 1958 Recent attempts to establish sugarcane borer parasites in Louisiana. Jour. Econ. Ent. 51: 163-164.
- McCormick, W. J., and Mathes, R.
- 1960 Biological control of the sugarcane borer in Louisiana. Internatl. Soc. Sugar Cane Technol. Cong. Proc. 10: 865-869.
- Chauthani, A. R., and Hamm, J. J.
- 1967 Biology of the exotic parasite *Drino munda* (Diptera: Tachinidae). Ent. Soc. Amer. Ann. 60 (2): 373-376, figs. 1-2.
- Childs, L.
- 1915 The alfalfa looper as a truck crop pest. *Plusia californica*. Oreg. Agr. Col. Expt. Sta. Bien. Crop Pest and Hort. Rpt. 1913-1914 2: 184-188, figs. 41-42.
- Chittenden, F. H. (See also Howard, L. O., and Chittenden)
- 1897 Some little-known insects affecting stored vegetable products: a collection of articles detailing certain original observations made upon insects of this class. U.S. Dept. Agr. Div. Ent. Bul., n. ser., 8: 1-45, figs. 1-10.
- 1898 Notes on cucumber beetles. U.S. Dept. Agr. Div. Ent. Bul., n. ser., 10: 26-31, figs. 7-8.
- 1898a On the parasites of adult Coleoptera. Ent. Soc. Wash. Proc. 4 (2): 75-79, figs. 2-3.
- 1899 Some insects injurious to garden and orchard crops. U.S. Dept. Agr. Div. Ent. Bul., n. ser., 19: 1-99, 20 figs.
- 1900 Some insects injurious to garden crops. U.S. Dept. Agr. Div. Ent. Bul., n. ser., 23: 1-92, 23 figs.
- 1901 Some insects injurious to the violet, rose, and other ornamental plants. U.S. Dept. Agr. Div. Ent. Bul., n. ser., 27: 1-114, 29 figs., 4 pls.
- 1901a The fall army worm and variegated cutworm. U.S. Dept. Agr. Div. Ent. Bul., n. ser., 29: 1-64, 11 figs.
- 1901b The green clover worm. U.S. Dept. Agr. Div. Ent. Bul., n. ser., 30: 45-50, fig. 26.
- 1902 Some insects injurious to vegetable crops. U.S. Dept. Agr. Div. Ent. Bul., n. ser., 33: 1-117, figs. 1-30.
- 1903 A brief account of the principal insect enemies of the sugar beet. U.S. Dept. Agr. Div. Ent. Bul., n. ser., 43: 1-71, figs. 1-65.

- 1907 Some insects injurious to truck crops. The cranberry spanworm. The striped garden caterpillar. U.S. Dept. Agr. Bur. Ent. Bul., n. ser., 66 (3): 21-32, figs. 6-7.
- 1908 The rose slugs. U.S. Dept. Agr., Bur. Ent. Cir. 105: 1-12, figs. 1-5.
- 1910 Notes on the cucumber beetles. U.S. Dept. Agr. Bur. Ent. Bul., n. ser., 82 (6): 67-75, figs. 19-23.
- 1911 Papers on insects affecting vegetables. The southern beet webworm. U.S. Dept. Agr. Bur. Ent. Bul., n. ser., 109 (2): 17-22, fig. 1.
- 1912 Papers on insects affecting vegetables. A little-known cutworm. U.S. Dept. Agr. Bur. Ent. Bul., n. ser., 109 (4): 47-51, fig. 8.
- 1913 The Florida fern caterpillar. U.S. Dept. Agr. Bur. Ent. Bul., n. ser., 125: 1-11, fig. 1.
- 1916 The common cabbage worm. U.S. Dept. Agr. Farmers' Bul. 766: 1-14, figs. 1-10.
- 1918 The lotus borer. Jour. Econ. Ent. 11: 453-457, pl. 16.
- 1919 The striped cucumber beetle and its control. U.S. Dept. Agr. Farmers' Bul. 1038: 1-20, figs. 1-15.
- 1923 The striped cucumber beetle and how to control it. U.S. Dept. Agr. Farmers' Bul. 1322: 1-16, figs. 1-16.
- 1923a Notes on *Ogdoconta cinereola* Guen. Brooklyn Ent. Soc. Bul. 18 (5): 155-156, 1 fig.
- 1924 The argus tortoise beetle. Jour. Agr. Res. 27 (1): 43-52, figs. 1-4, pl. 1.
- and Marsh, H. O.
- 1912 Papers on insects affecting vegetables. The imported cabbage webworm. U.S. Dept. Agr. Bur. Ent. Bul., n. ser., 109 (3): 23-45, figs. 4-7.
- 1920 The bean ladybird. U.S. Dept. Agr. Bul. 843: 1-21, figs. 1-5, pls. 1-6.
- 1920a The beet leaf-beetle. U.S. Dept. Agr. Bul. 892: 1-24, figs. 1-4, pls. 1-9.
- and Russell, H. M.
- 1909 Some insects injurious to truck crops. The semitropical army worm. U.S. Dept. Agr. Bur. Ent. Bul., n. ser., 66 (5): 53-70, figs. 8-11.
- Chong, [M.] (See also Davis, C. J., and Chong)
- 1965 *Winthemia* sp. (*diversoides* Bar.?). Hawaii. Ent. Soc. Proc. 1964 19 (1): 15.

- 1969 *Lixophaga sphenophori* (Villeneuve). Hawaii. Ent. Soc. Proc. 1968 20 (2): 278.
- Christenson, L. D. (See Van Dine and Christenson)
- Ciesla, W. M.
- 1964 Life history and habits of the elm spanworm, *Ennomos subsignarius*, in the southern Appalachian Mountains (Lepidoptera: Geometridae). Ent. Soc. Amer. Ann. 57 (5): 591-596, figs. 1-6.
- Rauschenberger, J. L., and Bell, J. C., Jr.
- 1968 *Eucosma* infestations in shortleaf pine cones. Jour. Econ. Ent. 61: 1129-1130, figs. 1-2.
- Claassen, P. W.
- 1921 *Typha* insects: their ecological relationships. N.Y. (Cornell) Agr. Expt. Sta. Mem. 47: 457-531, pls. 39-69.
- Clagg, [C. F. or H. B.]
- 1962 [A tachinid fly (*Achaetoneura archippivora*) — Hawaii]. (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 12 (8): 107-108.
- Clancy, D. W.
- 1946 Natural enemies of some Arizona cotton insects. Jour. Econ. Ent. 39: 326-328.
- 1969 Parasitization of cabbage and alfalfa loopers in southern California. Jour. Econ. Ent. 62: 1078-1083.
- and Pierce, H. D.
- 1966 Natural enemies of some lygus bugs. Jour. Econ. Ent. 59: 853-858.
- *Clark, A. H.
- 1936 The gold-banded skipper (*Rhabdoides cellus*). Smithsn. Inst., Smithsn. Misc. Collect. 95 (7): 1-50, figs. A-H, pls. 1-8.
- 1936a Who's who among the butterflies. Natl. Geog. Mag. 79 (5): 679-692, 10 pls.
- Clark, C. A. (See also Baker, W. A., Bradley, and Clark; Bradley, W. G., and Clark)
- 1934 The European corn borer and its controlling factors in the Orient. U.S. Dept. Agr. Tech. Bul. 455: 1-37, figs. 1-8.
- 1941 Liberations of European corn borer parasites in the United States in 1941. Insect Pest Survey Bul. 21 (8) Sup.: 621-624.

- 1943 The field status of parasites of the European corn borer at the close of 1941. U.S. Dept. Agr., Bur. Ent. and Plant Quar. E-588: 1-4.
- 1943a The collection, emergence, and release of parasites of the European corn borer, season of 1943. U.S. Dept. Agr., Bur. Ent. and Plant Quar., Insect Pest Survey, Spec. Sup. October 1, 1943: 1-3, errata.
- 1943b The field status of parasites of the European corn borer at the close of the 1942 season. U.S. Dept. Agr., Bur. Ent. and Plant Quar., Insect Pest Survey, Spec. Sup. December 27, 1943: 1-9.
- 1945 The field status of parasites of the European corn borer at the close of the 1943 season. U.S. Dept. Agr., Bur. Ent. and Plant Quar., Insect Pest Survey, Spec. Sup. 1945 (1): 1-5.
- 1945a The collection, emergence, and release of parasites of the European corn borer, season of 1944. U.S. Dept. Agr., Bur. Ent. and Plant Quar., Insect Pest Survey, Spec. Sup. 1945 (3): 1-4.
- and Carter, S. W.
- 1943 The collection, rearing, and release of parasites of the European corn borer in 1942. Jour. Econ. Ent. 36: 120.
- and Jones, D. W.
- 1945 Parasitization of the summer generation of the European corn borer in five eastern localities in 1944. U.S. Dept. Agr., Bur. Ent. and Plant Quar., Insect Pest Survey, Spec. Sup. 1945 (4): 1-2.
- 1946 The field status of parasites of the European corn borer at the close of the 1944 season. U.S. Dept. Agr., Bur. Ent. and Plant Quar., Insect Pest Survey, Spec. Sup. 1946 (2): 1-10.
- Clark, N., and Hoveland, N.
- 1947 What's new in farm science. Wis. Agr. Expt. Sta. Bul. 472: 1-63, 14 figs.
- Clausen, C. P.
- 1955 Releases of recently imported insect parasites and predators in California, 1952-53. Pan-Pacific Ent. 31 (2): 77-79.
- 1956 Biological control of insect pests in the continental United States. U.S. Dept. Agr. Tech. Bul. 1139: i-vi, 1-151.
- 1961 Biological control of western grape leaf skeletonizer (*Harrisina brillians* B. and McD.) in California. Hilgardia 31 (16): 613-638, 7 figs.

- King, J. L., and Teranishi, C.
1927 The parasites of *Popillia japonica* in Japan and Chosen (Korea), and their introduction into the United States. U.S. Dept. Agr., Dept. Bul. 1429: 1-55, figs. 1-35, pl. 1.
- Cloyd, W. J. (See St. Amand and Cloyd)
Cochaux, P.
1968 La tordeuse printanière du chêne, *Croesia semipurana* (Kft.) (Lepidoptera: Tortricidae) dans la région de Québec. Québec Ent. Soc. Ann. 13 (2): 98-107, figs. 1-3.
1969 Observations sur un groupe de défoliateurs du chêne rouge dans la région de Québec dont *Psilocorsis quercicella* Clem. (Lepidoptera: Oecophoridae). Québec Ent. Soc. Ann. 14 (2-3): 69-82, figs. 1-8.
- Cockerell, T. D. A.
1890 The tachinid parasite of *Cimbex americana*. Canad. Ent. 22 (4): 76.
1892 [Specimens recently added to the Museum]. Inst. Jamaica, Jour. 1 (2): 43.
1892a Additions to the Museum. Inst. Jamaica, Jour. 1 (2): 54-56.
- *Cole, A. C.
1929 *Arachnara subcarnea* Kell. (Lepidop.: Noctuidae) a host of *Masicera senilis* Rond. (Dipt.: Tachinidae). Ent. News 40 (7): 225.
1931 *Typha* insects and their parasites. Ent. News 42 (1): 6-11; (2): 35-43.
- Cole, F. R.
1915 Current notes. Jour. Econ. Ent. 8: 559.
- and Lovett, A. L.
1921 An annotated list of the Diptera (flies) of Oregon. Calif. Acad. Sci. Proc. ser. 4, 11 (15): 197-344, figs. 1-54.
- Coles, L. W. (See Brunson and Coles)
- *Collins, C. W. (See also Burgess and Collins)
1926 Observations on a recurring outbreak of *Heterocampa guttivitta* Walker and natural enemies controlling it. Jour. Agr. Res. 32 (7): 689-699.
1932 Natural enemies of the gypsy moth and the brown-tail moth in Massachusetts. Mass. Comnr. Conserv. and State Forester Ann. Rpt. 1931: 25-26.

- 1933 Natural enemies of the gypsy moth, brown-tail moth and a few other pests in Massachusetts. Mass. Comnr. Conserv. and State Forester Ann. Rpt. 1932: 27-28.
- 1934 Natural enemies of the gypsy moth, brown-tail moth, and some other forest and shade tree pests in Massachusetts. Mass. Comnr. Conserv., State Forester and Dir. Parks Ann. Rpt. 1933: 31-32.
- 1935 Natural enemies of the gypsy moth, brown-tail moth, and some other forest and shade tree pests in Massachusetts. Mass. Comnr. Conserv., State Forester and Dir. Parks Ann. Rpt. 1934: 38-39.
- and Hood, C. E.
1920 Life history of *Eubiomyia calosomae*, a tachinid parasite of *Calosoma* beetles. Jour. Agr. Res. 18 (9): 483-498, pls. 58-59.
- and Jones, T. H.
1932 Gipsy and brown-tail moth infestations are checked by imported parasites. (U.S. Dept. Agr.) Yearb. Agr. 1932: 236-239, fig. 68.
- Collins, M. M., and Weast, R. D.
1961 Wild silk moths of the United States. iii + 138 pp., 96 figs. Cedar Rapids, Iowa.
- Compere, H.
1946 Vegetable weevil parasites from South America and their colonization in California. Jour. Econ. Ent. 39: 671-672.
- *Comstock, J. A.
1937 Miscellaneous notes on western Lepidoptera. South. Calif. Acad. Sci. Bul. 36 (1): 19-23, pls. 6-9.
- and Dammers, C. M.
1943 Western poplar sphinx. South. Calif. Acad. Sci. Bul. 42 (1): 39-41, pl. 1.
- and Vazquez Garcia, L.
1964 Estudios de los ciclos biologicos en lepidopteros Mexicanos. II. Heterocera. [Mex.] Univ. Nac., Inst. de Biol. An. (1963) 34 (1-2): 217-273, figs. 1-22.
- Comstock, J. H.
1879 Report upon cotton insects. v + 511 pp., 77 figs., 3 pls. Washington.
- Concienne, E. J. (See Dugas et al.; Dugas, Miller, and Concienne)
- Conger, C. B. (See Blanchard and Conger)

- Conklin, J. G. (See also O'Kane and Conklin)
1952 Gypsy and browntail moths. N.H. Dept. Agr., Div. Insect and Plant Dis. Suppr. and Control Cir. 1951-1952 30: 76-79.
- Conradi, A. F. (See also Weed and Conradi)
1907 Insects of the garden. Tex. Agr. Expt. Sta. Bul. 89: 1-52, figs. 1-44.
- *Cook, A. J.
1884 Practical entomology. Notes from the entomological laboratory of the Michigan Agricultural College. 32 pp., 14 figs. Michigan Agricultural College.
1884a Practical entomology. Notes from the entomological laboratory of the Michigan Agricultural College. Mich. State Bd. Agr. Ann. Rpt. Sec. 1882-1883 22: 422-450, 14 figs.
1889 Report of the entomological department. Mich. Agr. Expt. Sta. Ann. Rpt. 2: 88-103, figs. 1-4.
1912 Cutworms. Calif. State Comnr. Hort. Monthly Bul. 1 (7): 271-274, figs. 116-119.
- Cook, E. F. (See Stehr and Cook)
- Cook, W. C.
1923 Note on a bombyliid parasite of the pale western cutworm, (*Porosagrotis orthogonia* Morr.). Jour. Econ. Ent. 16: 327.
1930 Field studies of the pale western cutworm (*Porosagrotis orthogonia* Morr.). Mont. Agr. Expt. Sta. Bul. 225: 1-79, figs. 1-13, pl. A.
- Cooke, O. B.
1953 Gypsy moth control in Connecticut — 1952. (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 3 (10): 151-152.
- Cooley, R. A.
1908 Fifth annual report of the state entomologist of Montana. Mont. Agr. Expt. Sta. Bul. 71: 131-152, 3 figs.
- Coolidge, K. R.
1908 *Telea polyphemus*. Ent. News 19 (1): 13.
1910 A California orange dog. Pomona Col. Jour. Ent. 2 (4): 333-334.

- 1924 The life-history of *Brephidium exilis* Bdv. (Lepid.: Lycaenidae). Ent. News 35 (4): 115-121.
- 1925 California butterfly notes — III. Brooklyn Ent. Soc. Bul. 20 (3): 146-147.
- Cooper, T. P.
- 1946 European corn borer parasites. Ky. Agr. Expt. Sta. Ann. Rpt. 1945 58: 30.
- 1947 European corn borer parasites. Ky. Agr. Expt. Sta. Ann. Rpt. 1946 59: 28.
- 1948 European corn borer parasites. Ky. Agr. Expt. Sta. Ann. Rpt. 1947 60: 41.
- 1949 European corn borer parasites. Ky. Agr. Expt. Sta. Ann. Rpt. 1948 61: 25-26.
- 1950 European corn borer parasites. Ky. Agr. Expt. Sta. Ann. Rpt. 1949 62: 13-14.
- 1951 European corn borer in Kentucky. Ky. Agr. Expt. Sta. Ann. Rpt. 1950 63: 18.
- Coppel, H. C. (See also Baldwin, W. F., and Coppel; Maw and Coppel; Torgersen and Coppel; Wilkes, Coppel, and Mathers)
- 1947 The collection of spruce budworm parasites in British Columbia with notes on their over-wintering habits. Ontario Ent. Soc. Ann. Rpt. 1946 77: 38-40.
- 1953 Effects of super-, multi-, and hyper-parasitism on the abundance of *Phytodietus fumiferanae* Rohw. (Hymenoptera: Ichneumonidae), a primary parasite of the spruce budworm, *Choristoneura fumiferana* (Clem.) (Lepidoptera: Tortricidae). Quebec Soc. Protect. Plants Rpt. [1950-1951] 32-33: 189-193, fig. 1.
- 1954 Notes on the parasites of *Neodiprion nanulus* Schedl (Hymenoptera: Tenthredinidae). Canad. Ent. 86 (4): 167-168.
- 1960 Key to adults of dipterous parasites of spruce budworm, *Choristoneura fumiferana* (Clem.) (Lepidoptera, Tortricidae). Ent. Soc. Amer. Ann. 53 (1): 94-97, pls. 1-2.
- and House, H. [L.]
- 1947 A description of cages used for the large-scale breeding of *Sturmia* sp., a tachinid parasite of sawflies. Canad. Ent. 79 (4): 74-80, figs. 1-5.

— and Jones, P. A.

- 1964 A note on *Hemyda aurata* R. D. (Diptera: Tachinidae), a parasite of *Podisus maculiventris* (Say) (Hemiptera: Pentatomidae). Ent. News 75 (7): 191-194, figs. 1-8.

— and Maw, M. G.

- 1954 Studies on dipterous parasites of the spruce budworm, *Choristoneura fumiferana* (Clem.) (Lepidoptera: Tortricidae). III. *Ceromasia auricaudata* Tns. (Diptera: Tachinidae). Canad. Jour. Zool. 32 (3): 144-156, figs. 1-16.
- 1954a Studies on dipterous parasites of the spruce budworm, *Choristoneura fumiferana* (Clem.) (Lepidoptera: Tortricidae). IV. *Madremyia saundersii* (Will.) (Diptera: Tachinidae). Canad. Jour. Zool. 32 (4): 314-323, figs. 1-16.

Coquillett, D. W.

- 1882 The greater leaf-roller. Noxious and Beneficial Insects of State of Ill., State Ent. Rpt. 11:10-15.
- 1882a The yellow canker worm. Noxious and Beneficial Insects of State of Ill., State Ent. Rpt. 11: 25-31.
- 1882b The army worm. Noxious and Beneficial Insects of State of Ill., State Ent. Rpt. 11: 49-64.
- 1889 The corn worm or boll worm in California. Insect Life (U.S. Dept. Agr., Div. Ent., Period. Bul.) 1 (11): 331-332.
- 1889a Enemies of *Diabrotica*. Insect Life (U.S. Dept. Agr., Div. Ent., Period. Bul.) 2 (3): 74.
- 1890 The dipterous parasite of *Diabrotica soror*. Insect Life (U.S. Dept. Agr., Div. Ent., Period. Bul.) 2 (7-8): 233-236, fig. 46.
- 1894 Report on some of the injurious insects of California. U.S. Dept. Agr., Div. Ent. Bul. 32: 22-32.
- 1895 New Tachinidae with a slender proboscis. Canad. Ent. 27 (5): 125-128.
- 1895a Notes and descriptions of Tachinidae. N.Y. Ent. Soc. Jour. 3 (2): 49-58.
- 1895b On the validity of the tachinid genus *Celatoria*. Psyche 7 (230): 251-252.
- 1897 Revision of the Tachinidae of America north of Mexico. A family of parasitic two-winged insects. U.S. Dept. Agr., Div. Ent. Tech. Ser. 7: 1-156.

- 1897a The walnut spanworm. (*Boarmia plumigeraria* Hulst). U.S. Dept. Agr. Div. Ent. Bul., n. ser., 7: 64-66, figs. 40-41.
- 1898 Additions to my synopsis of the Tachinidae. Canad. Ent. 30 (9): 233-237.
- Corkins, C. L.
1924 Grasshoppers. Colo. State Ent. Cir. 43: 44-48.
- Cory, E. N.
1946 Status of the European corn borer in Maryland. Md. Agr. Soc. Rpt. 1945 30: 146-148.
- Foster, J. R., McConnell, H. S., Bissell, T. L., Langford, G. S., and Ditman, L. P.
1950 European corn borer in Maryland. Md. Univ. Ext. Serv. Bul. 135: 1-32, figs. 1-18.
- and Langford, G. S.
1955 The Japanese beetle retardation program in Maryland. Md. Univ. Ext. Bul. 156: 1-20, 10 figs.
- McConnell, H. S., Arbuthnot, K. D., and Jones, D. W.
1952 Parasites of the European corn borer in Maryland. Md. Agr. Expt. Sta. Bul. A72: 1-12, figs. 1-2.
- Cosenza, A. V. (See Arbuthnot et al.; Jones, D. W., Carter, and Cosenza)
- Cotton, R. T. (See Brooks, F. E., and Cotton)
- Coulter, W. K. (See Carolin and Coulter)
- Coyne, J. F. (See Warren, L. O., and Coyne)
- Craw, A.
1891 Destructive insects, their natural enemies, remedies and recommendations. 51 pp., 62 figs. Sacramento, Calif.
- Crawford, H. G., and Spencer, G. J.
1922 The European corn borer control measures. Jour. Econ. Ent. 15: 231-236.
- Creighton, J. T.
1936 Report on cotton leaf-worm investigations in Florida, including temperature and hibernation studies. Jour. Econ. Ent. 29: 88-94.
- Criddle, N.
1911 Injurious insects of 1910 at Treesbank, Manitoba. Jour. Econ. Ent. 4: 236-241.
1918 The habits and control of white grubs in Manitoba. Agr. Gaz. Canada 5 (5): 449-454, 4 figs.
- Crossman, S. S. (See also Burgess and Crossman)
—and Webber, R. T.
1924 Recent European investigations of parasites of the gipsy moth, *Porthetria dispar* L., and the

- brown-tail moth, *Euproctis chrysorrhoea* L. Jour. Econ. Ent. 17: 67-76.
- Crowell, M. F.
1929 A preliminary study of the tracheal system of the mature larva of *Blepharipa scutellata* R-Desvoidy. Psyche 36 (3): 220-227, figs. 1-4.
- Crumb, S. E.
1929 Tobacco cutworms. U.S. Dept. Agr. Tech. Bul. 88: 1-179, figs. 1-19, pls. 1-9.
- Culver, J. J.
1919 A study of *Compsilura concinnata*, an imported tachinid parasite of the gipsy moth and the brown-tail moth. U.S. Dept. Agr., Dept. Bul. 766: 1-27, figs. 1-10, pl. 1.
- Cuming, F. G. (See also Reeks, Forbes, and Cuming)
1964 The biology, morphology, and biological control of the winter moth. Canada Dept. Forestry, Forest and Ent. Path. Branch, Ann. Rpt. 1964: 38.
- Cumming, M. E. P.
1954 [*Carcelia malacosomae* Sell. parasite of *Malacosoma disstria* Hbn.]. Canad. Insect Pest Rev. 32 (3): 182-183.
- Curran, C. H.
1924 New Canadian Diptera, with synopsis of the genus *Cynorhina*. Canad. Ent. 56 (8): 193-196.
1925 Three new Nearctic Tachinidae (Dipt.). Ent. News 36 (1): 13-18.
1925a New Tachinidae in the Canadian National Collection (Diptera). Canad. Ent. 57 (6): 150-156.
1926 New Diptera from Jamaica. Jamaica Dept. Agr. Ent. Bul. 4 (1-2): 102-114.
1927 A new tachinid parasitic on armyworms in Mexico. Hawaii. Ent. Soc. Proc. 1926 6 (3): 497-498.
1927a Studies in Canadian Diptera—III. The species of the tachinid genera related to *Lydella*, as represented in the Canadian National Collection. Canad. Ent. 59 (1): 11-24.
1927b Some new American Tachinidae (Diptera). Brooklyn Ent. Soc. Bul. 22 (3): 144-154.
1928 Revision of the American species of *Archytas* (Tachinidae, Diptera). Canad. Ent. 60 (8): 201-208; (9): 218-226, pl. 17; (10): 249-256, pl. 18; (11): 275-282, pl. 19.

- 1929 New Diptera in the American Museum of Natural History. Amer. Mus. Nat. Hist., Amer. Mus. Novitates 339: 1-13, figs. 1-2.
- 1930 A new tachinid parasitic on a sawfly. Canad. Ent. 62 (11): 246-247.
- 1932 New American Diptera. Amer. Mus. Nat. Hist., Amer. Mus. Novitates 534: 1-15.
- 1932a New North American Diptera, with notes on others. Amer. Mus. Nat. Hist., Amer. Mus. Novitates 526: 1-13, figs. 1-9.
- 1934 [*Carabus* from British Columbia parasitized]. N.Y. Ent. Soc. Jour. 42 (1): 138.
- 1935 New American Diptera. Amer. Mus. Nat. Hist., Amer. Mus. Novitates 812: 1-24, figs. 1-4.
- Currie, R. P.
1905 Catalogue of the exhibit of economic entomology at the Lewis and Clark Centennial Exposition, Portland, Oregon, 1905. U.S. Dept. Agr. Bur. Ent. Bul., n. ser., 53: 1-127.
- Cushman, R. A. (See also Pierce, Cushman, and Hood)
1927 The parasites of the pine tip moth, *Rhyacionia frustrana* (Comstock). Jour. Agr. Res. 34 (7): 615-622, fig. 1.
- Daecke, [V. A. E.]
1916 Lepidoptera and Diptera. Ent. News 27 (6): 286.
- Dahlsten, D. L.
1961 Life history of a pine sawfly, *Neodiprion* sp., at Willits, California (Hymenoptera: Diprionidae). Canad. Ent. 93 (3): 182-195, figs. 1-4.
- Dailey, E. F.
1957 Tent caterpillars — Washington. (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 7 (31): 628.
1958 Summary of insect conditions — 1957: Washington. (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 8 (14): 264-269.
- Dammers, C. M. (See Comstock, J. A., and Dammers)
- Daugherty, [D. M.]
1959 A tachinid (*Trichopoda pennipes*) — North Carolina. (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 9 (39): 888.
- Daviault, L. (See also Forbes, R. S., and Daviault; McLeod, J. M., and Daviault)
1935 Contribution à l'étude des insectes du bouleau. Nat. Canad. 62 (6-7): 201-214, figs. 1-8;

- (8-9): 237-256, figs. 9-20; (10): 282-297, figs. 21-33; (11): 315-330, figs. 34-39.
- 1939 Notes sur l'action des parasites de la chenille a tente du cerisier: *Cacoecia cerasivorana* Fitch. Nat. Canad. 66 (6-7): 179-187.
- 1941 La mouche à scie européenne de l'épinette dans les forêts du nord du St-Laurent. Forêt Québécoise 3 (7): 12-16.
- 1941a La chrysomèle du saule *Calligrapha multipunctata* var. *bigsbyana* Kby. Nat. Canad. 68 (3): 57-81, figs. 1-3; (4): 89-111, figs. 4-7.
- 1942 L'organisation et les travaux du laboratoire d'entomologie forestière de Berthierville. Forêt Québécoise 4 (8): 387-406, figs. 1-7.
- 1944 Nouvelles observations sur la mouche à scie européenne de l'épinette dans les forêts du nord du St-Laurent. Quebec Soc. Protect. Plants Ann. Rpt. 28: 39-43.
- 1946 Forest entomology. Quebec Min. Lands and Forests Rpt. 1945: 113-122.
- 1950 Les parasites de la tordeuse des bourgeons de l'épinette (*Choristoneura fumiferana* Clem.) dans la province de Québec. Quebec Soc. Protect. Plants Ann. Rpt. 31: 41-45.
- 1950a La tordeuse à tête verte de l'épinette: *Tortrix packardiana* Fern. Assoc. Canad.-Franc. pour l'Avanc. des Sci. Ann. 16: 147-151.
- 1953 La tenthrède de Leconte (*Neodiprion lecontei* (Fitch)) dans la province de Québec. Quebec Soc. Protect. Plants Rpt. [1950-1951] 32-33: 165-184, figs. 1-4.

— and Ducharme, R.

- 1966 Life history and habits of the green spruce leaf miner, *Epinotia nanana* (Treitschke) (Lepidoptera: Tortricidae). Canad. Ent. 98 (7): 693-699, figs. 1-3.

*Davis, C. J.

- 1963 *Trichopoda pennipes* (Fabricius). Hawaii. Ent. Soc. Proc. 1962 18 (2): 203.
- 1963a A parasitic encyrtid (*Exoristobia philippinensis* Ashmead). (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 13 (45): 1324.
- 1964 *Trichopoda pennipes pilipes* Fabricius. Hawaii. Ent. Soc. Proc. 1963 18 (3): 347.

- 1964a The introduction, propagation, liberation, and establishment of parasites to control *Nezara viridula* variety *smaragdula* (Fabricius) in Hawaii (Heteroptera: Pentatomidae). Hawaii. Ent. Soc. Proc. 1963 18 (3): 369-375.
- 1964b A parasitic tachina fly (*Trichopoda pennipes* (F.)). (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 14 (18): 414.
- 1965 *Trichopoda pennipes* (F.). Hawaii. Ent. Soc. Proc. 1964 19 (1): 8-9.
- 1969 Notes on the grass webworm, *Herpetogramma licarsisalis* (Walker) (Lepidoptera: Pyraustidae), a new pest of turfgrass in Hawaii and its enemies. Hawaii. Ent. Soc. Proc. 1968 20 (2): 311-316, figs. 1-2.
- and Chong, M.
- 1969 Recent introductions for biological control in Hawaii XIV. Hawaii. Ent. Soc. Proc. 1968 20 (2): 317-322.
- and Knowlton, G. F.
- 1963 A tachina fly (*Exorista mella*) — Utah. (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 13 (50): 1399.
- and Krauss, N. L. H.
- 1963 Recent introductions for biological control in Hawaii — VIII. Hawaii. Ent. Soc. Proc. 1962 18 (2): 245-249.
- 1964 Recent introductions for biological control in Hawaii — IX. Hawaii. Ent. Soc. Proc. 1963 18 (3): 391-397.
- 1965 Recent introductions for biological control in Hawaii — X. Hawaii. Ent. Soc. Proc. 1964 19 (1): 87-90.
- Davis, E. G. (See Manis et al.)
- Davis, I. W. (See Britton and Davis)
- Davis, J. E.
- 1954 Summary of insect conditions — 1953: Oregon. (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 4(10): 203-204.
- Davis, J. J.
- 1912 Report on insects injurious to flowering and ornamental greenhouse plants in Illinois. Noxious and Beneficial Insects of State of Ill., State Ent. Rpt. 27: 83-143, figs. 1-52.

- 1916 A progress report on white grub investigations. Jour. Econ. Ent. 9: 261-281, pls. 14-16.
- 1918 Common white grubs. U.S. Dept. Agr. Farmers' Bul. 940: 1-28, figs. 1-21.
- Davis, R.
[1961] Parasites of the elm spanworm, *Ennomos subsignarius* (Hbn.), in Georgia (Lepidoptera: Geometridae). Ent. Soc. Wash. Proc. (1960) 62 (4): 247-248.
- Dawson, J. C. (See Breshears and Dawson)
- Dearness, J.
1898 Annual address of the president. Ontario Ent. Soc. Ann. Rpt. 28: 14-23, figs. 1-17.
- Deay, H. O.
1946 Exotic parasites of European corn borer have become established in two counties. Ind. Agr. Expt. Sta. Dir. Rpt. 59: 11.
- 1947 European corn borer — parasite releases and recoveries. Ind. Agr. Expt. Sta. Ann. Rpt. 60: 68.
- Decker, G. C.
1930 The biology of the four-lined borer *Luperina stipata* (Morr.). Iowa Agr. Expt. Sta. Res. Bul. 125: 125-164, figs. 1-22.
- 1931 The biology of the stalk borer *Papaipema nebris* (Gn.). Iowa Agr. Expt. Sta. Res. Bul. 143: 289-351, figs. 1-22.
- 1933 Biology of the bidens borer, *Epiblema otiosana* (Clemens) (Lepidoptera, Olethreutidae). N.Y. Ent. Soc. Jour. (1932) 40 (4): 503-509, fig. 1.
- 1935 The parasites of some lepidopterous stalk borers in Iowa. Iowa State Col. Jour. Sci. 9 (4): 567-580.
- 1952 Summary of insect conditions — 1952: Illinois. (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 2 (33): 431-434.
- deGryse, J. J.
1935 On the use of parasites in the control of forest insects. Forestry Chron. 11 (3): 8-11.
- Devitt, J. J. (See Hawkins and Devitt)
- Dew, J. A. (See also Hinds and Dew)
1913 Fall army worm *Laphygma frugiperda* (S. & A.). Jour. Econ. Ent. 6: 361-366.
- Dickerson, E. L., and Weiss, H. B.
1920 The insects of the evening primroses in New Jersey. N.Y. Ent. Soc. Jour. 28 (1): 32-74, pls. 1-3.

- Dietrick, E. J., and Bosch, R. van den
 1957 Insectary propagation of the squash bug and its
 parasite *Trichopoda pennipes* Fabr. Jour. Econ.
 Ent. 50: 627-629.
- Dillier, [J. H.], and Seibels, [W. T.]
 1965 A tachina fly (*Winthemia quadripustulata*)
 Alabama. (U.S. Dept. Agr.) Coop. Econ. Insect
 Rpt. 15 (19): 457.
- Dillon, L. S.
 1944 Post-mortem "stings" of the white flannel moth.
 Brooklyn Ent. Soc. Bul. (1943) 38 (5): 176.
- Dimick, R. E. (See also Mote, Stearns, and Dimick)
 1931 Hand-parasitizing the European earwig with larvae
 of *Digonichaeta setipennis* Fall. Jour. Econ. Ent.
 24: 962-964.
- and Mote, D. C.
 1934 Progress report regarding the introduction in Oregon
 of *Digonichaeta setipennis* Fall., a tachinid
 parasite of the European earwig. Jour. Econ. Ent.
 27: 863-865.
- Dimmock, G., and Ashmead, W. H.
 1898 Notes on parasitic Hymenoptera, with descriptions
 of some new species. Ent. Soc. Wash. Proc. 4
 (2): 148-171.
- Ditman, L. P. (See Cory et al.)
- Dixon, J. C., and Benjamin, D. M.
 1963 Natural control factors associated with the jack-pine
 budworm, *Choristoneura pinus*. Jour. Econ.
 Ent. 56: 266-270.
- Doane, R. W., and Brodie, D. A.
 1901 The variegated cutworm (*Peridroma saucia* Hubner).
 Wash. Agr. Expt. Sta. Bul. 47: 1-16, figs. 1-5.
- Dohanian, S. M.
 1940 *Melissopus latiferreanus* as a pest of filberts in the
 Northwest. Jour. Econ. Ent. 33: 852-856.
 1942 Parasites of the filbert worm. Jour. Econ. Ent.
 35: 836-841.
- Donohoe, H. C., and Kaloostian, G. H.
 1937 Notes on the grape leaf folder and its parasites in the
 San Joaquin Valley. Jour. Econ. Ent.
 30: 967-968.
- Dorst, H. E. (See Knowlton et al.)
- Douglas, W. A. (See also Ingram, Bynum, and Douglas)
 1935 Insects observed attacking *Crotalaria* in Louisiana in

- 1932 and 1933. Jour. Econ. Ent. 28: 686-687, fig. 102.
- and Ingram, J. W.
1942 The sugarcane beetle from rice-field insects. Rice Jour. 45 (7): 8-10.
- Douglass, J. R. (See also Manis et al.)
1933 Habits, life history, and control of the Mexican bean beetle in New Mexico. U.S. Dept. Agr. Tech. Bul. 376: 1-46, figs. 1-30.
- Dowden, P. B. (See also Furniss, R. L., and Dowden)
1934 *Zenillia libatrix* Panzer, a tachinid parasite of the gypsy moth and the brown-tail moth. Jour. Agr. Res. 48 (2): 97-114, figs. 1-4.
1935 *Brachymeria intermedia* (Nees), a primary parasite, and *B. compsilurae* (Cwfd.), a secondary parasite, of the gypsy moth. Jour. Agr. Res. 50 (6): 495-523, figs. 1-5.
1937 Parasites of larch sawfly in western Montana and Great Lakes region. Jour. Econ. Ent. 30: 381.
1946 Parasitization of the oriental moth (*Cnidocampa flavescens* (Walk.)) by *Chaetexorista javana* B. and B. Ent. Soc. Amer. Ann. 39 (2): 225-241, figs. 1-4.
1961 The persistence of gypsy moth parasites in heavily sprayed areas on Cape Cod, Massachusetts. Jour. Econ. Ent. 54: 873-875.
1962 Parasites and predators of forest insects liberated in the United States through 1960. U.S. Dept. Agr., Agr. Handb. 226: i-iv, 1-70.
- Buchanan, W. D., and Carolin, V. M.
1948 Natural-control factors affecting the spruce budworm. Jour. Econ. Ent. 41: 457-464.
- and Carolin, V. M.
1950 Natural control factors affecting the spruce budworm in the Adirondacks during 1946-1948. Jour. Econ. Ent. 43: 774-783.
- Jaynes, H. A., and Carolin, V. M.
1953 The role of birds in a spruce budworm outbreak in Maine. Jour. Econ. Ent. 46: 307-312.
- Downes, W.
1935 [*Digonichaeta setipennis* parasite of earwigs]. Canad. Insect Pest Rev. 13 (4): 181.
1938 Insect conditions on Vancouver Island during 1937. Canad. Insect Pest Rev. 16 (1): 72-73.

- Dowsett, J. M. (See Paxton et al.)
- Dozier, H. L.
 1920 An ecological study of hammock and piney woods insects in Florida. Ent. Soc. Amer. Ann. 13 (4): 325-380, figs. 1-22.
- Drake, C. J.
 1918 Pumpkin bugs. Fla. Buggist 2 (2): 91.
 1920 The southern green stink-bug in Florida. Fla. State Plant Bd. Quart. Bul. 4 (3): 41-94, figs. 6-38.
 1941 Survey of potato insects. Iowa Agr. Expt. Sta. Rpt. Agr. Res. 1940-1941 (pt. 1): 151-152.
 1942 Survey of potato insects. Iowa Agr. Expt. Sta. Rpt. Agr. Res. 1941-1942 (pt. 1): 170-171.
 1946 Part 4. Division of Entomology and Apiculture. Division of Entomology. Iowa Agr. Yearb. 1945 46: 161-176, figs. 1-5, 1 fig.
 1947 Part 4. Report of the state entomologist. Division of Entomology. Iowa Agr. Yearb. 1946 47: 107-130, 11 figs.
- Harris, H. M., and Blickenstaff, C. C.
 1945 The European corn borer situation in Iowa. Iowa Dept. Agr. Yearb. 1944 45: 91-107, 14 figs.
- Harris, H. M., and Tauber, O. E.
 1945 Field and laboratory studies on the ecology and behavior of the European corn borer in Iowa. Iowa Corn Res. Inst. Ann. Rpt. 10: 64-68, figs. 15-17.
- Driggers, B. F.
 1929 Some parasites of the oriental peach moth in New Jersey. N.Y. Ent. Soc. Jour. 37 (2): 169-170.
- Driscoll (See Wressell, Driscoll, and Svec)
- Drooz, A. T. (See also Benjamin and Drooz; Jaynes and Drooz)
 1953 Larch sawfly investigations in Minnesota, 1952. Jour. Econ. Ent. 46: 826-828.
 1960 The larch sawfly, its biology and control. U.S. Dept. Agr. Tech. Bul. 1212: i-iii, 1-52, figs. 1-13.
- and Benjamin, D. M.
 1956 Parasites from two jack-pine budworm outbreaks on the upper peninsula of Michigan. Jour. Econ. Ent. 49: 412-413.
- Ducharme, R. (See Daviault and Ducharme)
- Dudley, F. H.
 1919 Report of the state horticulturist. Maine Comnr. Agr. Rpt. 18: 30-69, 11 figs.

Dugas, A. L. (See also Mathes et al.)

———Concienne, E. J., Sibley, L. M., and Joyner, H. L.

1955 Sugar cane insects. La. Agr. Expt. Sta. Ann. Rpt. 1953-1954: 87-89.

———Miller, G. C., and Concienne, E. J.

1956 Larval parasites. La. Agr. Expt. Sta. Ann. Rpt. 1954-1955: 93.

Duncan, C. D.

1923 Notes on the biology of two species of *Stenopelmatus* (Orth.: Tettigoniidae). Ent. News 34 (3): 73-77, figs. A-C.

1933 The control of harmful insects. Pp. 236-247, figs. 120-124. In Pickwell, G., Duncan, C. D., Hazeltine, K. S., and Smith, E., eds., Insects. xv + 304 pp., 139 figs. Los Angeles, San Jose, and San Francisco.

———and Pickwell, G.

1939 The world of insects. ix + 409 pp., 194 figs. New York and London.

Dunn, L. H.

1917 The cocoanut-tree caterpillar (*Brassolis isthmia*) of Panama. Jour. Econ. Ent. 10: 473-488, pls. 22-23.

Dupree, M. (See also Leuck and Dupree)

———and Beckham, C. M.

1955 The cowpea curculio - a pest of southern field peas. Ga. Agr. Expt. Sta. Bul., n. ser., 6: 1-32, frontis., figs. 1-2.

DuRant, J. A., and Hyche, L. L.

1967 Descriptions of the immature stages and internal reproductive structures of *Lixophaga mediocris* (Diptera: Tachinidae). Ent. Soc. Amer. Ann. 60 (2): 406-409, figs. 1-2.

Dustan, A. G.

1923 The natural control of the white marked tussock moth under city and forest conditions. Acadian Ent. Soc. Proc. (1922) 8: 109-127, pls. 1-2.

Dyar, H. G.

1891 On the oviposition of *Tachina*. Insect Life (U.S. Dept. Agr., Div. Ent., Period. Bul.) 3 (9-10): 414.

1902 Descriptions of the larvae of some moths from Colorado. U.S. Natl. Mus. Proc. 25 (1290): 369-412.

- Ebel, B. H.
1965 The *Dioryctria* coneworms of north Florida pines (Lepidoptera: Phycitidae). Ent. Soc. Amer. Ann. 58 (5): 623-630, figs. 1-3.
- Ebeling, W.
1950 Subtropical entomology. xi + 747 pp., 570 figs. San Francisco.
- Eckart, C. F.
1911 Report of director of the Experiment Station. Hawaii. Planters' Rec. 5 (6): 325-349.
1912 [Report of director of the Experiment Station]. Hawaii. Sugar Planters' Assoc. Proc. 31st Ann. Mtg. 1911 31: 19-44.
1912a Report of director of the Experiment Station. Hawaii. Planters' Rec. 7 (6): 345-365.
1912b Report of director of the Experiment Station. Hawaii. Sugar Planters' Assoc. Expt. Sta. Committee Rpt. 1912: 9-32.
1913 Report of director of the Experiment Station. Hawaii. Sugar Planters' Assoc. Proc. 32d Ann. Mtg. 1912 32: 18-40.
- *Eddy, B.
1940 Office of Entomology and Plant Industry. R.I. Dept. Agr. and Conserv. Ann. Rpt. 5: 56-83, 14 figs.
1941 Office of Entomology and Plant Industry. R.I. Dept. Agr. and Conserv. Ann. Rpt. 6: 49-67, 1 fig.
1943 Division of Entomology and Plant Industry. R.I. Dept. Agr. and Conserv. Ann. Rpt. 9: 33-50, 2 figs.
- Eddy, C. O.
1941 Silk beetles (*Luperodes* spp.). Insect Pest Survey Bul. 21 (6): 420.
1941a A silk beetle (*Luperodes* sp.). Insect Pest Survey Bul. 21 (7): 509.
- and Nettles, W. C.
1930 The bean leaf beetle. S.C. Agr. Expt. Sta. Bul. 265: 1-25, figs. 1-8.
- Ehrlich, P. R.
1948 On the parasitization of *Danaus plexippus*. Lepidopterists' News 2 (8): 92.
- Eichmann, R. D.
1943 Control of the squash bug (*Anasa tristis* De Geer) in eastern Washington. Wash. Agr. Expt. Sta. Bul. 435: 38-39.

- 1944 Control of the squash bug (*Anasa tristis*) in eastern Washington. Wash. Agr. Expt. Sta. Bul. 455: 36.
- Eidt, D. C.
1969 The life histories, distribution, and immature forms of the North American sawflies of the genus *Cephalcia* (Hymenoptera: Pamphiliidae). Ent. Soc. Canada, Mem. 59: 1-56, figs. 1-107.
- Einmo, [J. R.]
1963 *Achaetoneura archippivora* (Williston). Hawaii. Ent. Soc. Proc. 1962 18 (2): 196.
- Eisner, T. (See Neff and Eisner)
- Elgee, D. E. (See Bird, F. T., and Elgee)
- Eliot, I. M., and Soule, C. G.
1902 Caterpillars and their moths. xiii + 302 pp., 132 figs. New York.
- Embree, D. G.
1958 The external morphology of the immature stages of the beech leaf tier, *Psilocorsis faginella* (Chamb.) (Lepidoptera: Oecophoridae), with notes on its biology in Nova Scotia. Canad. Ent. 90 (3): 166-174, figs. 1-32.
1960 Observations on the spread of *Cyzenis albicans* (Fall.) (Tachinidae: Diptera), an introduced parasite of the winter moth, *Operophtera brumata* (L.), (Geometridae: Lepidoptera), in Nova Scotia. Canad. Ent. 92 (11): 862-864.
1964 The population dynamics of the winter moth. Canada Dept. Forestry, Forest Ent. and Path. Branch, Ann. Rpt. 1964: 40.
1965 The population dynamics of the winter moth in Nova Scotia, 1954-1962. Ent. Soc. Canada, Mem. 46: 1-57, figs. 1-27.
1966 The role of introduced parasites in the control of the winter moth in Nova Scotia. Canad. Ent. 98 (11): 1159-1168, figs. 1-7.
- and Sisojevic, P.
1965 The bionomics and population density of *Cyzenis albicans* (Fall.) (Tachinidae: Diptera) in Nova Scotia. Canad. Ent. 97 (6): 631-639, figs. 1-7.
- Essig, E. O.
1913 Injurious and beneficial insects of California. Calif. State Comnr. Hort. Monthly Bul. 2 (1-2): i-xxxii, 1-367, figs. 1-321.

- 1915 Injurious and beneficial insects of California (second edition). Calif. State Comnr. Hort. Monthly Bul. 4 Sup.: i-lxxxii, 1-541, figs. 1-503.
- 1926 Insects of western North America. xi + 1035 pp., 766 figs. New York.
- Evans, D.
1962 Descriptions and life history of *Melanolophia imitata* (Walker) (Lepidoptera: Geometridae). Canad. Ent. 94 (6): 594-605, figs. 1-15.
- Evans, K. E. (See Manis et al.)
- Evans, R. W. (See Arbuthnot et al.)
- Farris, S. H. (See Leech and Farris)
- Fattig, P. W.
1944 The *Phyllophaga* or May beetles of Georgia. Emory Univ. (Ga.) Mus. Bul. 2: 1-32.
- 1949 The Larvaevoridae (Tachinidae) or parasitic flies of Georgia. Emory Univ. (Ga.) Mus. Bul. 8: 1-40.
- Felker, A. L.
1940 Report for the biennial period July 1, 1938 to June 30, 1940. (Div. of Insect Suppression). N.H. Dept. Agr. Rpt. (1938-1940) 1940: 120-148.
- Felt, E. P.
1898 14th report of the state entomologist on injurious and other insects of the state of New York 1898. N.Y. State Mus. Bul. 5 (23): 153-295, figs. 1-20, pls. 1-9.
- 1901 16th report of the state entomologist on injurious and other insects of the state of New York 1900. N.Y. State Mus. Bul. 7 (36): 949-1063, figs. 1-3, pls. 1-16.
- 1902 17th report of the state entomologist on injurious and other insects of the state of New York 1901. N.Y. State Mus. Bul. 53: 699-925, figs. 1-29, pls. 1-6.
- 1904 19th report of the state entomologist on injurious and other insects of the state of New York 1903. N.Y. State Mus. Bul. 76: 91-235, pls. 1-4.
- 1905 20th report of the state entomologist on injurious and other insects of the state of New York 1904. N.Y. State Mus. Bul. 97: 359-597, figs. 1-24, pls. 1-19.
- 1906 21st report of the state entomologist on injurious and other insects of the state of New York 1905.

- N.Y. State Mus. Bul. 104: 49-186, figs. 1-48, pls. 1-10.
- 1908 23rd report of the state entomologist on injurious and other insects of the state of New York 1907. N.Y. State Mus. Bul. 124: 1-541, figs. 1-49, pls. 1-44.
- 1912 27th report of the state entomologist on injurious and other insects of the state of New York 1911. N.Y. State Mus. Bul. 155: 1-198, figs. 1-5, pls. 1-27.
- 1913 [Experiences during 1912]. N.Y. Ent. Soc. Jour. 21 (1): 74.
- 1916 Thirtieth report of the state entomologist on injurious and other insects of the State of New York 1914. N.Y. State Mus. Bul. 180: 1-336, figs. 1-101, pls. 1-19.
- and Bromley, S. W.
- 1941 New and unusual shade tree pests. Jour. Econ. Ent. 34: 383-386.
- and Joutel, L. H.
- 1904 Monograph of the genus *Saperda*. N.Y. State Mus. Bul. 74: 1-86, figs. 1-7, pls. 1-14.
- *Fernald, C. H.
- 1898 The brown-tail moth. U.S. Dept. Agr. Div. Ent. Bul., n. ser., 17: 24-32.
- Fernald, H. T.
- 1934 [Polka dot wasp moth (*Syntomeida epilais* Walk. var. *jucundissima* Dyar) on oleanders in Florida]. Insect Pest Survey Bul. 14 (1): 23-24.
- Fernández Artiles, R. (See Scaramuzza, Fernández Artiles, and Fernández Perez)
- Fernández Perez, J. (See Scaramuzza, Fernández Artiles, and Fernández Perez)
- Feytaud, J.
- [1936] Exposé de M. Feytaud (France). Pp. 19-31. In Anon., Compte rendu et rapports de la conférence internationale pour l'étude de la lutte en commun contre le doryphore — Bruxelles, Palais des Académies, 22 et 23 Janvier, 1936. 62 pp. Brussels.
- 1937 Recherches sur le doryphore. III. — Causes de réduction naturelles (milieu — maladies — ennemis). Ann. des Epiphyt. et de Phytogénét., n. ser., 3 (1): 35-97, figs. 1-4, pls. 1-2.

Fife, L. C. (See Chapman, Fife, and Cavitt)

Fink, D. E.

- 1932 Biology and habits of the strawberry leaf roller, *Ancylis comptana* (Froel.), in New Jersey. Jour. Agr. Res. 44 (7): 541-558, figs. 1-9.

*Finlayson, L. R., and Finlayson, T. [U.]

- 1958 Parasitism of the European pine sawfly, *Neodiprion sertifer* (Geoff.) (Hymenoptera: Diprionidae), in southwestern Ontario. Canad. Ent. 90 (4): 223-225.

— and Reeks, W. A.

- 1936 Notes on the introduction of *Diprion* parasites to Canada. Canad. Ent. 68 (7): 160-166.

Finlayson, T. [U.] (See also Finlayson, L. R., and Finlayson; Smith, R. W., and Finlayson)

- 1960 Taxonomy of cocoons and puparia, and their contents, of Canadian parasites of *Neodiprion sertifer* (Geoff.) (Hymenoptera: Diprionidae). Canad. Ent. 92 (1): 20-47, figs. 1-43.

- 1960a Taxonomy of cocoons and puparia, and their contents, of Canadian parasites of *Diprion hercyniae* (Htg.) (Hymenoptera: Diprionidae). Canad. Ent. 92 (12): 922-941, figs. 1-24.

- 1962 Taxonomy of cocoons and puparia, and their contents, of Canadian parasites of *Diprion similis* (Htg.) (Hymenoptera: Diprionidae). Canad. Ent. 94 (3): 271-282, figs. 1-10.

- 1963 Taxonomy of cocoons and puparia, and their contents, of Canadian parasites of some native Diprionidae (Hymenoptera). Canad. Ent. 95 (5): 475-507, figs. 1-22.

Finnegan, R. J.

- 1964 Red-headed pine sawfly (*Neodiprion lecontei* (Fitch)). Canad. Insect Pest Rev. 42 (2): 29.

Fiske, W. F. (See also Howard, L. O., and Fiske)

— and Thompson, W. R.

- 1909 Notes on the parasites of the Saturniidae. Jour. Econ. Ent. 2: 450-460.

Fitzpatrick, R. E.

- 1967 [Research report]. Canada Dept. Agr., Res. Branch, Res. Sta., Vancouver, Brit. Columbia, Res. Rpt. 1967: 1-22.

- [1969] [Research report]. Canada Dept. Agr., Res. Branch, Res. Sta., Vancouver, Brit. Columbia, Res. Rpt. (1968): 349-357.

Flanders, S. E.

- 1926 Notes on parasites at Saticoy, California, during the year 1925. *Pan-Pacific Ent.* 2 (3): 157-158.
- 1926a Insect enemies of the codling moth in the vicinity of Ventura, California. *Pan-Pacific Ent.* 3(1): 44-45.
- 1940 Observations on the biology of the elm leaf beetle parasite, *Erynnia nitida* R.-Desv. *Jour. Econ. Ent.* 33: 947-948.
- 1940a Elm leaf beetle parasite on the job. *South. Pacific Rural Press and Calif. Farmer* 140 (13): 412.
- 1942 An additional observation on the biology of *Erynnia nitida*. *Jour. Econ. Ent.* 35: 607.

Fleming, W. E.

- 1963 The Japanese beetle in the United States. U.S. Dept. Agr., *Agr. Handb.* 236: 1-30, figs. 1-21.

Fletcher, D. S. (See Zimmerman, E. C., and Fletcher)

Fletcher, J.

- 1897 Insect injuries to Ontario crops in 1896. *Ontario Ent. Soc. Ann. Rpt.* 27: 58-69, figs. 58-70, 1 pl.
- 1904 Entomological record, 1903. *Ontario Ent. Soc. Ann. Rpt.* 1903 34: 85-99.
- 1905 Entomological record, 1904. *Ontario Ent. Soc. Ann. Rpt.* 1904 35: 56-78.
- 1907 Entomological record, 1906. *Ontario Ent. Soc. Ann. Rpt.* 1906 37: 102.

Flores Caceres, S.

- 1958 Principales plagas de la caña de azucar en Mexico. (Distribucion geográfica). (Mex.) *Cong. Nac. de Ent. y Fitopat. Mem.* 1: 243-259, 2 figs.

Foltz, J. L. (See Allen, D. C. et al.)

Forbes, A. G. (See Jones, M. L., and Forbes)

Forbes, R. S. (See also Reeks, Forbes, and Cuming; Reeks, Smith, and Forbes)

— and Daviault, L.

- 1964 The biology of the mountain-ash sawfly, *Pristiphora geniculata* (Htg.) (Hymenoptera: Tenthredinidae), in eastern Canada. *Canad. Ent.* 96 (8): 1117-1133, figs. 1-7.

Forbes, S. A.

- 1885 The grass worm or fall army worm. *Noxious and Beneficial Insects of State of Ill., State Ent. Rpt.* 14: 55-67, 120-121, pl. 5.

- 1891 On the common white grubs (*Lachnosterna* and *Cyclocephala*). Noxious and Beneficial Insects of State of Ill., State Ent. Rpt. 1889-1890 17: 30-53, pl. 4.
- 1907 On the life history, habits, and economic relations of the white-grubs and May-beetles (*Lachnosterna*). Ill. Agr. Expt. Sta. Bul. 116: 445-480.
- 1908 On the life history, habits, and economic relations of the white-grubs and May-beetles (*Lachnosterna*). Noxious and Beneficial Insects of State of Ill., State Ent. Rpt. 24: 135-168, errata, 1 fig., pls. 9-11.
- and Hart, C. A.
- 1900 The economic entomology of the sugar beet. Ill. Univ. Agr. Expt. Sta. Bul. 60: 397-532, figs. 1-97, pls. 1-9.
- Foster, J. R. (See Cory et al.)
- Fox, H. (See Phillips, W. J., and Fox)
- Frankenfeld, J. C. (See Wildermuth and Frankenfeld)
- *Franklin, H. J.
- 1910 Further work upon the stalk borers. *Papaipema* sp. Minn. Expt. Sta. State Ent. Rpt. 13: 85-94, figs. 38-46.
- 1943 Injurious and beneficial insects affecting the cranberry. Mass. Agr. Expt. Sta. Bul. 398: 26-28.
- 1950 Cranberry insects in Massachusetts. Part V. Insects and other animals beneficial in cranberry growing. Mass. Agr. Expt. Sta. Bul. 445: 55-76, figs. 1-18.
- and Lacroix, D. S.
- 1924 The spotted cutworm, *Agrotis c-nigrum* (L), a cranberry pest. Jour. Econ. Ent. 17: 406-408.
- Franklin, R. T., and Holdaway, F. G.
- 1966 A relationship of the plant to parasitism of European corn borer by the tachinid parasite *Lydella grisescens*. Jour. Econ. Ent. 59: 440-441.
- Franklin, W. W. (See Smith, R. C., and Franklin)
- French, G. T.
- 1943 Division of Plant Industry, part II, plant pest control — insects and plant diseases. Va. Dept. Agr. and Immigr., Ann. Rpt. Comnr. 1942-43: 56-66.
- Fretay, E.
- 1960 Contribution a la lutte contre le borer de la canne (*Diatrea* [sic] *saccharalis*). Brit. W. Indies Sugar Technol. Proc. 1960: 141-146.

- 1960a [Translation] Contribution to the fight against sugar cane borer. Brit. W. Indies Sugar Technol. Proc. 1960: 147-151.
- Friend, R. B.
1945 The gypsy moth in Connecticut. Conn. Acad. Arts and Sci., Trans. 36: 607-629.
- and Turner, N.
1931 The Mexican bean beetle in Connecticut. Conn. Agr. Expt. Sta. Bul. 332: 72-108, figs. 1-14.
- Fronk, W. D.
1950 Cultural and biological control of the southern corn rootworm in peanuts. Jour. Econ. Ent. 43: 22-24.
- Frost, S. W.
1931 The well-marked fruit-worm, *Stenomoma algidella* Walk. Jour. Econ. Ent. 24: 1208-1213, figs. 81-82, pl. 50.
- Fullaway, D. T.
1914 Tobacco insects in Hawaii. Hawaii Agr. Expt. Sta. Bul. 34: 1-20, figs. 1-9.
- F[uller], A. S.
1880 The Colorado potato-beetle [concl.]. Amer. Ent. 3 (8): 190-191, figs. 99-102.
- Funasaki, [G.] (See also Kawamura and Funasaki)
1967 Southern green stink bug (*Nezara viridula*). (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 17 (21): 450.
1969 Southern green stink bug (*Nezara viridula*). (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 19 (29): 555.
- Furniss, M. M., and Barr, W. F.
1967 Bionomics of *Anacamptodes clivinaria profanata* (Lepidoptera: Geometridae) on mountain mahogany in Idaho. Idaho Agr. Expt. Res. Bul. 73: 1-24, figs. 1-15.
- *Furniss, R. L., and Dowden, P. B.
1941 Western hemlock sawfly, *Neodiprion tsugae* Middleton, and its parasites in Oregon. Jour. Econ. Ent. 34: 46-52.
- Fyles, T. W.
1897 Lepidopterous pests of the meadow and the lawn. Ontario Ent. Soc. Ann. Rpt. 1896 27: 97-104, figs. 85-89.

Gahan, A. B.

- 1915 Notes on two parasitic Diptera. Ent. Soc. Wash. Proc. 17 (1): 24.
- 1917 Descriptions of some new parasitic Hymenoptera. U.S. Natl. Mus. Proc. 53 (2197): 195-217.

Garcia, L. Vazquez (See Comstock, J. A., and Vazquez Garcia)

Gardiner, J. G.

- 1938 [Report]. Canada [Dept. Agr.] Min. Agr. Rpt. 1938: 1-117.
- 1939 [Report]. Canada [Dept. Agr.] Min. Agr. Rpt. 1939: 1-138.
- 1940 [Report]. Canada [Dept. Agr.] Min. Agr. Rpt. 1940: 1-154.
- 1942 [Report]. Canada [Dept. Agr.] Min. Agr. Rpt. 1942: 1-162.
- 1943 [Report]. Canada [Dept. Agr.] Min. Agr. Rpt. 1943: 1-155.
- 1944 [Report]. Canada [Dept. Agr.] Min. Agr. Rpt. 1944: 1-186.
- 1945 [Report]. Canada [Dept. Agr.] Min. Agr. Rpt. 1945: 1-212.
- 1946 [Report]. Canada [Dept. Agr.] Min. Agr. Rpt. 1946: 1-235.
- 1947 [Report]. Canada [Dept. Agr.] Min. Agr. Rpt. 1947: 1-257.
- 1948 [Report]. Canada [Dept. Agr.] Min. Agr. Rpt. 1948: 1-245.
- 1949 [Report]. Canada [Dept. Agr.] Min. Agr. Rpt. 1949: 1-260.
- 1950 [Report]. Canada [Dept. Agr.] Min. Agr. Rpt. 1950: 1-287.
- 1954 [Report]. Canada [Dept. Agr.] Min. Agr. Rpt. 1954: 1-151.

Gardner, T. R., and Parker, L. B.

- 1940 Investigations of the parasites of *Popillia japonica* and related Scarabaeidae in the Far East from 1929 to 1933, inclusive. U.S. Dept. Agr. Tech. Bul. 738: 1-36, fig. 1.

***Garman, H.**

- 1891 A supplementary note on *Diabrotica 12-punctata*. Psyche 6 (181): 78-80.
- 1897 Notes on tobacco worms, from observations made in 1896. Ky. Agr. Expt. Sta. Bul. 66: 6-32, figs. 1-4.
- 1908 The army-worm. Ky. Agr. Expt. Sta. Bul. 137: 429-449, figs. 1-16.

- Garman, P. (See also Porter and Garman)
1918 A comparison of several species of Lepidoptera infesting peach and apple in Maryland, with additional notes on the oriental peach moth. Md. Agr. Expt. Sta. Bul. 223: 103-126, figs. 30-35, pls. 1-4.
- Gartley, A. (See Paxton et al.)
- Gauthier, G.
1945 Insects of the season 1944 in Quebec. Canad. Insect Pest Rev. 23 (1): 25-28.
- Geissler, G. H. (See Gould, E., and Geissler)
- Gentry, T. G.
1877 Metamorphic changes of *Platysamia cecropia*. Canad. Ent. 9 (3): 41-50.
- Genung, W. G. (See also Hayslip et al.)
1959 Notes on the syntomid moth *Lymire edwardsi* (Grote) and its control as pest of *Ficus* in south Florida. Fla. Ent. 42 (1): 39-42.
- Gerhardt, [P. D.]
1942 [*Trichopoda pennipes* parasitic on *Euryophthalmus cinctus californicus*]. Pan-Pacific Ent. 18 (1): 42-43.
- Getzendaner, C. W.
1936 Parasitizing European earwig with *Bigonicheta setipennis* Fall. Recent developments in methods and equipment. Jour. Econ. Ent. 29: 1105-1114, figs. 1-7.
[1937] The introduction and propagation of *Digonichaeta setipennis* Fall., a parasite of the European earwig. Ent. Soc. Brit. Columbia, Proc. 1936 33: 8-12.
- *Gibson, A.
1911 The preparatory stages of *Phragmatobia assimilans* Walker, variety *franconia* Slosson. Canad. Ent. 43 (4): 125-128.
1911a The entomological record for 1910. Ontario Ent. Soc. Ann. Rpt. 41: 101-120.
1915 The army-worm *Cirphus (Leucania) unipuncta* Haw. Canada Dept. Agr. Ent. Bul. 9: 1-34, figs. 1-19.
1915a Cutworms and their control. Canada Dept. Agr. Ent. Bul. 10: 1-31, figs. 1-20.
1916 The entomological record, 1915. Ontario Ent. Soc. Ann. Rpt. 46: 194-230.

- 1919 The greenhouse leaf-tyer (*Phlyctaenia ferrugalis* Hbn.). Agr. Gaz. Canada 6 (7): 626-629, 2 figs.
- 1927 International entomology - retrospective and prospective. Jour. Econ. Ent. 20: 47-62.
- Gibson, L. P.
- 1964 Biology and life history of acorn-infesting weevils of the genus *Conotrachelus* (Coleoptera: Curculionidae). Ent. Soc. Amer. Ann. 57 (5): 521-526.
- Gibson, W. W. (See also Carrillo S., Ortega C., and Gibson)
 — and Carrillo S., J. L.
- 1959 Lista de insectos en la coleccion entomologica de la Oficina de Estudios Especiales, S. A. G. Mex., Sec. de Agr. y Ganaderia, Fol. Misc. 9: i-xviii, 1-254.
- Gill, J. B.
- 1913 Papers on deciduous fruit insects and insecticides. The fruit-tree leaf-roller. U.S. Dept. Agr., Bur. Ent. Bul. 116 (5): 91-110, pls. 12-16.
- 1917 The pecan leaf case-bearer. U.S. Dept. Agr. Bul. 571: 1-28, pls. 1-3.
- 1917a Important pecan insects and their control. U.S. Dept. Agr. Farmers' Bul. 843: 1-48, figs. 1-58.
- 1930 Notes on parasites of pecan nut case-bearer. Natl. Pecan Assoc. Bul. 4 (1): 70-75.
- Gillette, C. P.
- 1890 Insects from Iowa. Insect Life (U.S. Dept. Agr., Div. Ent., Period. Bul.) 2 (9): 280-281.
- 1891 Notes on certain experiments and observations at the Iowa experiment station. Insect Life (U.S. Dept. Agr., Div. Ent., Period. Bul.) 3 (6): 258-260.
- 1922 Notes on insect pests. Colo. State Ent. Cir. 36: 14-26, fig. 1, pl. 2.
- Gilliatt, F. C.
- [1921] The brown-tail moth situation in Nova Scotia. Nova Scotia Ent. Soc. Proc. (1920) 6: 74-80.
- 1926 Controlling the brown tail moth in Nova Scotia. Ontario Ent. Soc. Ann. Rpt. 56: 63-67.
- 1931 Insects of the season 1930 in Nova Scotia. Ontario Ent. Soc. Ann. Rpt. 61: 10-13.
- 1937 Natural control of the grey banded leaf roller, *Eulia mariana* Fern, in Nova Scotia orchards. Canad. Ent. 69 (7): 145-146.

Gilmore, J. U.

- 1938 Observations on the hornworms attacking tobacco in Tennessee and Kentucky. *Jour. Econ. Ent.* 31: 706-712.

Girault, A. A.

- 1908 Texas, Virginia and Maryland notes on the catalpa sphinx, *Ceratonia catalpae* Boisduval. *Ent. News* 19 (5): 197-199.
- 1908a Effect of the parasitism of tachina flies on the larvae of *Phlegethontius sexta* Johanssen. *Ent. News* 19 (5): 235-236.
- 1912 Fragments on North American insects — I. *Ent. News* 23 (9): 399-411.
- 1913 Fragments on North American insects — III. *Ent. News* 24 (2): 53-63.

Gittins, A. R. (See Manis et al.)

Glendenning, R.

- 1924 The satin moth in British Columbia. *Canad. Dept. Agr. Pam., n. ser.*, 50: 1-14, 8 figs.
- 1931 The progress of parasite introduction in British Columbia. *Ent. Soc. Brit. Columbia, Proc.* 28: 29-32.

Gobeil, A.-R.

- 1937 Observations sur la mouche à scie, *Hylotoma macleayi* Leach. *Nat. Canad.* 64 (3): 81-88, figs. 1-6.

Goble, H. W.

- 1965 The armyworm *Pseudaletia unipuncta* (Haworth) in Ontario in 1964. *Ontario Ent. Soc. Ann. Rpt.* 95: 11-14, figs. 1-3.

Goleman, [D. L.] (See Parks and Goleman)

*Gould, E., and Geissler, G. H.

- 1940 Parasites of the pistol case-bearer. *Jour. Econ. Ent.* 33: 814-815.

Gould, G. E.

- 1936 Studies on cucumber beetle control in 1935. *Jour. Econ. Ent.* 29: 731-735.
- 1943 Investigate the biology and means for control of the striped cucumber beetle. *Ind. Agr. Expt. Sta. Ann. Rpt.* 56: 61.
- 1944 The biology and control of the striped cucumber beetle. *Ind. Agr. Expt. Sta. Bul.* 490: 1-28, frontis., fig. 1.

- and Walker, H. G.
 1931 Sweetpotato sawfly (*Sterictiphora collaris* Say).
 Insect Pest Survey Bul. 11 (6): 379.
- Gowdey, C. C.
 1921 The fall army worm, *Laphygma frugiperda* (S. & A.). Jamaica Dept. Agr. Ent. Cir. 4: 1-4.
 1923 The principal insect pests of tobacco in Jamaica. Jamaica Dept. Agr. Ent. Cir. 9: 1-6.
 1923a The principal agricultural pests of Jamaica. Jamaica Dept. Agr. Ent. Bul. 2: i-vi, 1-80.
 1926 Catalogus insectorum jamaicensis. Jamaica Dept. Agr. Ent. Bul. 4 (1-2): 1-114.
- Goyer, R. A.
 1966 (Abstract) The parasite complex associated with *Eucosma rescissoriana* Heinrich (Lepidoptera: Olethreutidae) in northern Idaho. Idaho Univ. Ent. Club Newsletter 5: 17-18.
- and Schenk, J. A.
 1969 Parasitism of the cone moth *Eucosma rescissoriana* (Lepidoptera: Olethreutidae) in northern Idaho. Canad. Ent. 101 (10): 1063-1069.
- *Graham, A. R.
 1955 Biology and establishment in Canada of *Mesoleius tenthredinis* Morley (Hymenoptera: Ichneumonidae), a parasite of the larch sawfly, *Pristiphora erichsonii* (Hartig) (Hymenoptera: Tenthredinidae). Quebec Soc. Protect. Plants Rpt. 35: 61-75, figs. 1-22.
 1958 Recoveries of introduced species of parasites of the winter moth, *Operophtera brumata* (L.) (Lepidoptera: Geometridae), in Nova Scotia. Canad. Ent. 90 (10): 595-596.
 1965 A preliminary list of the natural enemies of Canadian agricultural pests. Canada Dept. Agr. Res. Inst. Inform. Bul. 4: 1-179.
- Graham, S. A.
 1956 The larch sawfly in the Lake states. Forest Sci. 2 (2): 132-160.
- and Baumhofer, L. G.
 1927 The pine tip moth in the Nebraska National Forest. Jour. Agr. Res. 35 (4): 323-333, figs. 1-3.
- Gray, I. E.
 1946 Observations on the life history of the horned passalus. Amer. Midland Nat. 35 (3): 728-746, figs. 1-3.

- Gray, K. W. (See Wilcox, Gray, and Mote)
- Greene, C. T.
- 1921 Dipterous parasites of sawflies. Ent. Soc. Wash. Proc. 23 (2): 41-43.
- 1921a Two new species of Diptera. Ent. Soc. Wash. Proc. 23 (6): 125-127, fig. 1.
- 1922 An illustrated synopsis of the puparia of 100 muscoid flies (Diptera). U.S. Natl. Mus. Proc. 60 (10): 1-39, pls. 1-20.
- 1934 Tachinid flies with an evanescent fourth vein, including a new genus and five new species. Ent. Soc. Wash. Proc. 36 (2): 27-40, figs. 1-37.
- 1937 The pupa of *Myocera tabanivora* Hall (Diptera). U.S. Natl. Mus. Proc. 84 (3012): 217-218, fig. 27.
- 1956 Dipterous larvae parasitic on animals and man and some dipterous larvae causing myiasis in man. Amer. Ent. Soc. Trans. 82 (1): 17-34, figs. 1-21.
- Greenwood, D. E. (See Hofmaster and Greenwood)
- Gressitt, J. L.
- 1959 *Salmacia longipulvilli* (Tothill). Hawaii. Ent. Soc. Proc. 1958 17 (1): 4.
- Griffiths, K. J.
- 1960 Parasites of *Neodiprion pratti banksianae* Rohwer in northern Ontario. Canad. Ent. 92 (9): 653-658.
- 1961 Red-headed pine sawfly (*Neodiprion lecontei* Fitch). Canad. Insect Pest Rev. 39 (5): 196-197.
- Griswold, C. L. (See Schaffner and Griswold)
- Gryse, J. J. de (See deGryse, J. J.)
- *Guppy, J. C.
- 1967 Insect parasites of the armyworm, *Pseudaletia unipuncta* (Lepidoptera: Noctuidae), with notes on species observed in Ontario. Canad. Ent. 99 (1): 94-106.
- Guppy, R.
- [1950] Some records of parasitic Diptera from Wellington, B.C. Ent. Soc. Brit. Columbia, Proc. (1949) 46: 4.
- 1951 Mortality of *Nymphalis milberti* larvae. Lepidopterists' News 5 (6-7): 69.
- Guyer, [G. G.] et al.
- 1965 Armyworm (*Pseudaletia unipuncta*) — Michigan. (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 15 (26): 670.
- Gyrisco, G. G. (See Neunzig and Gyrisco)

- Hadley, C. H.
1938 Progress of Japanese beetle investigations. N.Y. Ent. Soc. Jour. 46 (2): 203-216.
- Haeussler, G. J. (See also Allen, H. W., Holloway, and Haeussler)
1930 Parasites of the oriental peach moth, *Laspeyresia molesta* Busck, in North America. Jour. Agr. Res. 41 (5): 365-377, figs. 1-2.
- Hagen, H. A.
1875 On *Attacus (Samia) columbia* and its parasites. Buffalo Soc. Nat. Sci. Bul. 2: 201-208.
- Hagen, K. S. (See Bosch and Hagen)
- Hale
1967 A tachina fly (*Lespesia archippivora*). (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 17 (11): 180.
- Haley, W. E. (See Holloway et al.; Ingram et al.)
- Haliburton, W. H. (See McGugan, Haliburton, and MacDonald)
- *Hall, D. G.
1937 New muscoid flies (Diptera) in the United States National Museum. U.S. Natl. Mus. Proc. 84 (3011): 201-216, figs. 19-26.
1939 Two new species of Tachinidae parasitic upon hemlock sawfly larvae in North America (Diptera: Tachinidae). Ent. Soc. Wash. Proc. 41 (8): 239-243.
- Hall, J. A.
1929 Leaf rollers attacking the apple in Norfolk County, Ontario. Ontario Ent. Soc. Ann. Rpt. 60: 137-139.
- Hallock, H. C. (See King, J. L., and Hallock)
- Hamilton, D. W. (See Summerland and Hamilton)
- Hamlin, J. C.
1924 A review of the genus *Chelinidea* (Hemiptera-Heteroptera) with biological data. Ent. Soc. Amer. Ann. 17 (2): 193-208.
1925 Important *Opuntia* insects of the United States: biological notes on the moth-borers (Pyralidae). Pan-Pacific Ent. 2 (1): 1-11.
- Reed, W. D., and Phillips, M. E.
1931 Biology of the Indian-meal moth on dried fruits in California. U.S. Dept. Agr. Tech. Bul. 242: 1-26, fig. 1.
- Hamm, J. J. (See Chauthani and Hamm)

Hammond, G. H. (See also Petch and Hammond)

- 1929 [Parasites of white grubs reared in Quebec]. *Canad. Insect Pest Rev.* 7 (4): 39.
- 1930 [Parasites of *Phyllophaga anxia* LeConte in Quebec]. *Canad. Insect Pest Rev.* 8 (3): 34.
- 1930a [Control of *Phyllophaga anxia* LeConte in Quebec by parasites]. *Canad. Insect Pest Rev.* 8 (5): 60.
- 1931 The white grub situation in eastern Canada. *Ontario Ent. Soc. Ann. Rpt.* 1930 61: 69-73, fig. 1.
- 1934 [*Cryptomeigenia theutis* parasite of June beetle]. *Canad. Insect Pest Rev.* 12 (2): 66.
- 1934a [*Cryptomeigenia* parasite of common white grub]. *Canad. Insect Pest Rev.* 12 (3): 99.
- 1934b [*Cryptomeigenia theutis* parasite of June beetle]. *Canad. Insect Pest Rev.* 12 (4): 130.
- 1934c [*Cryptomeigenia* sp. parasite of June beetle]. *Canad. Insect Pest Rev.* 12 (4): 130.
- 1936 Insects of the season in the Apple Hill district and other localities in Ontario, in 1935. *Canad. Insect Pest Rev.* 14 (1): 38-40.
- 1936a [White grub parasite, *Cryptomeigenia* sp.]. *Canad. Insect Pest Rev.* 14 (2): 84.
- 1937 [Parasitism by the white grub parasite, *Microphthalma michiganensis* Tns.]. *Canad. Insect Pest Rev.* 15 (4): 163.
- 1941 [*Cryptomeigenia* sp. parasite of June beetles]. *Canad. Insect Pest Rev.* 19 (3): 186-187.
- 1941a [*Eutrixia exile* Coquillett parasite of adult June beetles]. *Canad. Insect Pest Rev.* 19 (3): 187.
- 1942 [*Eutrixia* spp. parasites of June beetles]. *Canad. Insect Pest Rev.* 20 (2): 147.
- 1942a [*Microphthalma michiganensis* Townsend parasite of *Phyllophaga* sp.]. *Canad. Insect Pest Rev.* 20 (2): 148.
- 1945 [*Cryptomeigenia simplex* Curran parasite of *Phyllophaga* spp.]. *Canad. Insect Pest Rev.* 23 (2): 159.
- 1945a [*Cryptomeigenia theutis* Walker parasite of *Phyllophaga* spp.]. *Canad. Insect Pest Rev.* 23 (2): 159.
- 1945b [*Cryptomeigenia triangularis* Curran parasite of *Phyllophaga* spp.]. *Canad. Insect Pest Rev.* 23 (2): 159.

- 1947 Colorado potato beetle (*Leptinotarsa decemlineata* Say) — Ontario. *Canad. Insect Pest Rev.* 25 (1): 37.
- 1948 White grub parasites (*Microphthalma* spp.). *Canad. Insect Pest Rev.* 26 (4): 175.
- 1949 A white grub parasite (*Microphthalma disjuncta* Wied.). *Canad. Insect Pest Rev.* 27 (3): 161.
- 1949a Io moth (*Automeris io* F.). *Canad. Insect Pest Rev.* 27 (6): 236.
- Handford, R. H.
- 1962 Research report 1928-1961 entomology laboratory, Kamloops, British Columbia. Canada Dept. Agr., Res. Branch, Kamloops, Brit. Columbia, Ent. Lab. Res. Rpt. 1928-1961: 1-36, fig. 1.
- and Neilson, C. L.
- 1953 Insects of the season 1952 in the interior of British Columbia. *Canad. Insect Pest Rev.* 31 (1): 4-10.
- Painter, Richard H., and Moore, H. W.
- 1937 [Tachinid on armyworms]. *Canad. Insect Pest Rev.* 15 (4): 159.
- Hansen, L. L. (See Knowlton and Hansen)
- Hanson, W. J. (See Knowlton and Hanson)
- Harcourt, D. G.
- 1963 Biology of cabbage caterpillars in eastern Ontario. *Ontario Ent. Soc. Ann. Rpt.* 93: 61-73, figs. 1-15.
- Hardy, [D. E.]
- 1957 *Anachaetopsis tortricis* (Coquillett). *Hawaii. Ent. Soc. Proc.* 1956 16 (2): 195.
- Harman, S. W.
- 1928 The fruit tree leaf roller in western New York. N.Y. (Geneva) Agr. Expt. Sta. Bul. 561: 1-31, figs. 1-6.
- Harmston, F. C. (See Knowlton and Harmston)
- Harned, R. W.
- 1927 Pecan insect project. *Miss. Agr. Expt. Sta. Ann. Rpt.* 40: 26-28.
- *Harris, H. M. (See also Blickenstaff, Arbuthnot, and Harris; Drake, Harris, and Blickenstaff; Drake, Harris, and Tauber)
- 1948 Part 4. Report of the state entomologist. Division of Entomology. *Iowa Dept. Agr. Yearb.* 1947 48: 119-138, 5 figs.
- 1949 Part 4. Division of Entomolgy [sic] and Apiculture. *Iowa Dept. Agr. Yearb.* 1948 49: 117-131, 10 figs.

- 1950 Part 4. Report of the state entomologist. Division of Entomology. Iowa Dept. Agr. Yearb. 1949 50: 127-159, 15 figs.
- 1951 Part 4. Report of the state entomologist. Division of Entomology. Iowa Dept. Agr. Yearb. 1950 51: 123-154, 9 figs.
- 1952 Part IV. Report of the state entomologist. Division of Entomology. Iowa Dept. Agr. Yearb. 1951 52: 123-146, figs. 1-4, 1 fig.
- 1954 Part IV. Report of the state entomologist. Division of Entomology. Iowa Dept. Agr. Book Bien. Rpt. 1: 139-154, figs. 1-3, 4 figs.

— and Lilly, J. H.

- [1950] The pattern of attack and survival of the European corn borer on various plant growth stages and its relation to performance in testing inbred lines and hybrid combinations of corn for borer resistance, tolerance and susceptibility. Iowa Agr. Expt. Sta. Rpt. Agr. Res. 1949: 245-247.

Harris, T. W.

- 1841 A report on the insects of Massachusetts, injurious to vegetation. viii + 459 pp. Cambridge, Mass.

Harrison, J. O.

- 1963 The natural enemies of some banana insect pests in Costa Rica. Jour. Econ. Ent. 56: 282-285.
- 1963a On the biology of three banana pests in Costa Rica (Lepidoptera: Limacodidae, Nymphalidae). Ent. Soc. Amer. Ann. 56 (1): 87-94, pls. 1-3.

— and Stephens, C. S.

- 1966 Notes on the biology of *Ecpantheria icasia* (Lepidoptera: Arctiidae), a pest of bananas. Ent. Soc. Amer. Ann. 59 (4): 671-674, figs. 1-5.

Hart, C. A. (See Forbes, S. A., and Hart)

Hartley, E. A.

- 1923 Parasitism of the European pine sawfly, *Diprion (Lophyrus) simile* Hartig, Hymenoptera, Tenthredinidae, in Pennsylvania. Jour. Econ. Ent. 16: 386-388.

Hartung, W. (See Severin, H. H. P., Severin, and Hartung)

Harville, J. P.

- 1955 Ecology and population dynamics of the California oak moth *Phryganidia californica* Packard (Lepidoptera: Dioptriidae). Microentomology 20 (4): 83-166, figs. 53-58.

- Haseman, L.
1940 The walnut caterpillar. Mo. Agr. Expt. Sta. Bul. 418: 1-14, figs. 1-11.
- Haut, I. C.
1951 Research leads the way to agricultural progress, 64th annual report. Md. Agr. Expt. Sta. Bul. A-67: 1-88, fig. 1, 76 figs.
1952 Research the key to agricultural progress, sixty-fifth annual report, 1951-1952. Md. Agr. Expt. Sta. Bul. A-74: 1-88, 65 figs.
1954 The progress of research in serving agriculture, sixty-sixth annual report, 1952-1953. Md. Agr. Expt. Sta. Bul. A-78: 1-96, 59 figs.
- Hawboldt, L. S. (See also Balch and Hawboldt)
1940 [*Pristiphora erichsonii* Hartig multiply parasitized by *Bessa* and *Mesoleius*]. Canad. Insect Pest Rev. 18 (2): 155.
1944 Forest insects of the season 1943 in the maritime provinces, Gaspé, and Newfoundland. Canad. Insect Pest Rev. 22 (1): 19-24.
1947 *Bessa selecta* (Meigen) (Diptera: Tachinidae) as a parasite of *Gilpinia hercyniae* (Hartig) (Hymenoptera: Diprionidae). Canad. Ent. 79 (5): 84-104, pls. 8-9.
1949 Forest insects of the season. Nova Scotia Dept. Lands and Forests Rpt. 1948: 35-39, figs. A-B.
- Hawkes, R. B.
1968 The cinnabar moth, *Tyria jacobaeae*, for control of tansy ragwort. Jour. Econ. Ent. 61: 499-501.
- Hawkins, J. H., and Devitt, J. J.
1953 The European corn borer in Maine — its biology and life history. Maine Agr. Expt. Sta. Bul. 522: 1-26, figs. 1-17.
- Hawley, I. M.
1918 Insects injurious to the hop in New York with special reference to the hop grub and the hop redbug. N.Y. (Cornell) Agr. Expt. Sta. Mem. 15: 141-224, figs. 9-70.
1925 The more important insects injurious to the sugar-beet in Utah. Utah Agr. Expt. Sta. Cir. 54: 1-45, figs. 1-33.
- Hayes, W. P.
1920 The life histories of some Kansas *Lachnosterna*. Jour. Econ. Ent. 13: 303-318, figs. 11-12.

- Hays, K. L.
 1958 Descriptions of the larva and adult female of *Phorostoma tabanivora* (Hall) (Diptera: Larvaevoridae) and notes on the biology of the species. Ent. Soc. Amer. Ann. 51 (6): 552-553, figs. A-E.
- Hayslip, N. C. (See also Ingram et al.)
 1944 Insect pests and their control. Fla. Agr. Expt. Sta. Ann. Rpt. 1944: 124-125.
 1944a Prevalence and control of sugarcane moth borer. Fla. Agr. Expt. Sta. Ann. Rpt. 1944: 128.
 1945 Insect pests and their control. Fla. Agr. Expt. Sta. Ann. Rpt. 1945: 195-199.
- Genung, W. G., Kelsheimer, E. G., and Wilson, J. W.
 1953 Insects attacking cabbage and other crucifers in Florida. Fla. Agr. Expt. Sta. Bul. 534: 1-57, figs. 1-38.
- Mathes, R., and Bregger, T.
 1951 Studies upon the prevalence and control of the sugarcane moth borer in south Florida. Fla. Agr. Expt. Sta. Ann. Rpt. 1951: 190-191.
- Headlee, T. J.
 1908 Life history of the striped cucumber beetle with a brief account of some experiments for its control. Jour. Econ. Ent. 1: 203-209.
- Hedlin, A. F.
 1960 On the life history of the Douglas-fir cone moth, *Barbara colfaxiana* (Kft.) (Lepidoptera: Olethreutidae), and one of its parasites, *Glypta evetriae* Cush. (Hymenoptera: Ichneumonidae). Canad. Ent. 92 (11): 826-834, figs. 1-11.
- Hefley, H. M., Jr.
 1926 IX. Parasite of the larva of the tomato worm moth: *Protoparce quinque maculatus*. Okla. Acad. Sci. Proc. 5: 77-80.
 1928 Differential effects of constant humidities on *Protoparce quinque maculatus* Haworth, and its parasite, *Winthemia quadripustulata* Fabricus [sic]. Jour. Econ. Ent. 21: 213-221, figs. 17-18.
- Heinrich, C. (See Holloway et al.; Holloway, Loftin, and Heinrich)
- Henriksen, K. L., and Lundbeck, W.
 1917 Grønlands Landarthropoder (Insecta et Arachnida Groenlandicae). Meddel. om Grønland (Copenhagen) 22: 481-823, figs. 1-4.

- Henshaw, S.
1892 Report upon the gypsy moth in Massachusetts. U.S. Dept. Agr., Div. Ent. Bul. 26: 75-82, 1 fig.
- Hensill, G. S.
1933 Larval parasites of the codling moth in the Santa Clara Valley, California. Jour. Econ. Ent. 26: 603.
- Herbert, F. B. (See Burke and Herbert)
- Heron, R. J. (See also Ives et al.)
1960 The relative effects of cocoon submergence on the mortality of the larch sawfly, *Pristiphora erichsonii* (Hymenoptera: Tenthredinidae) and its parasite *Bessa harveyi* (Diptera: Tachinidae). Ent. Soc. Amer. Ann. 53 (4): 476-481, fig. 1.
1961 A note on temperature and postdiapause development of the larch sawfly and its parasite *Bessa harveyi* (Tnsd.). Canad. Ent. 93 (6): 431-433, figs. 1-2.
- Herrick, G. W.
1912 The fruit-tree leaf-roller. N.Y. (Cornell) Agr. Expt. Sta. Bul. 311: 277-292, frontis., figs. 59-76.
- and Leiby, R. W.
1915 The fruit-tree leaf roller (*Archips argyrospila* Walk.), order Lepidoptera, family Tortricidae. N.Y. (Cornell) Agr. Expt. Sta. Bul. 367: 246-279, figs. 44-61.
- Hetrick, L. A. (See also Poos and Hetrick)
1941 Life history studies of *Neodiprion americanum* (Leach). Jour. Econ. Ent. 34: 373-377, figs. 1-4.
1947 The cowpea curculio: its life history and control. Va. Agr. Expt. Sta. Bul. 409: 1-23, figs. 1-2.
1959 Ecology of the pine sawfly, *Neodiprion excitans* (Rohwer) (Hymenoptera, Diprionidae). Fla. Ent. 42 (4): 159-162.
- Hewitt, C. G.
1912 The large larch sawfly [*Nematus erichsonii*] with an account of its parasites, other natural enemies and means of control. Canada Dept. Agr., Div. Ent., Ent. Bul. 5: 1-42, figs. 1-26, pl. 1.
1915 Report of the dominion entomologist C. Gordon Hewitt, D. Sc., F. R. S. C. for the year ending March 31, 1915. Canada Dept. Agr., Dominion Ent. Rpt. 1915: 1-40, figs. 1-4.

- Hibbs, E. T.
1953 Parasites aid in European corn borer control. Ohio Agr. Expt. Sta. Bul. 725: 58-59.
- Hildahl, V. (See also Lejeune and Hildahl)
1950 Larch sawfly (*Pristiphora erichsonii* Htg.). Canad. Insect Pest Rev. 28 (2): 109.
- Hill, C. C.
1925 Biological studies of the green clover worm. U.S. Dept. Agr., Dept. Bul. 1336: 1-20, figs. 1-15.
- Hills, O. A., and Taylor, E. A.
[1952] Salt-marsh caterpillar (*Estigmene acrea* (Drury)). (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 1 (7): 126.
- Hinds, W. E., and Dew, J. A.
1915 The grass worm or fall army worm (*Laphygma frugiperda*, S. A.). Ala. Agr. Expt. Sta. Bul. 186: 59-92, pls. I-IV.
- and Osterberger, B. A.
1931 The soybean caterpillar in Louisiana. Jour. Econ. Ent. 24: 1168-1173.
- Hine, J. S.
1904 The Diptera of British Columbia. (First part). Canad. Ent. 36 (4): 85-92.
- Hitchcock, S. W.
1961 Pupal mortality of the orange-striped oakworm. Jour. Econ. Ent. 54: 962-964.
- Hitchings, E. F.
1908 Third annual report of the state entomologist on the gipsy and brown-tail moths and other insect pests of the state of Maine. Maine Dept. Agr., Bur. Ent., State Ent. Ann. Rpt. 1907 3: 1-105, figs. 1-17, 20 pls.
- Hixson, E.
1941 The walnut datana. Okla. Agr. Expt. Sta. Bul. B-246: 1-29, frontis., figs. 1-8.
- Hodson, A. C. (See also Kulman and Hodson)
1941 An ecological study of the forest tent caterpillar, *Malacosoma disstria* Hbn., in northern Minnesota. Minn. Agr. Expt. Sta. Tech. Bul. 148: 1-55, figs. 1-6.
- Hoerner, J. L.
1933 The alfalfa webworm (*Loxostege commixtalis* Walker). Colo. State Ent. Cir. 58: 1-12, 5 figs.

Hoffman, C. H.

1936 A population study of *Cacoecia cerasivorana* Fitch with special reference to its insect parasites (Tortricidae-Lepidoptera). Brooklyn Ent. Soc. Bul. 31 (5): 209-211.

1937 Biological notes on *Pseudolucanus placidus* Say, *Platycerus quercus* Weber and *Ceruchus piceus* Weber (Lucanidae — Coleoptera). Ent. News 48 (10): 281-284.

Hofmaster, R. N., and Greenwood, D. E.

1949 Fall armyworm control on forage and truck crops. Jour. Econ. Ent. 42: 502-506.

Holdaway, F. G. (See also Franklin, R. T., and Holdaway; Hsiao and Holdaway)

—— and Lucas, [E.]

1941 Biological control of cabbage worm. Hawaii Agr. Expt. Sta. Rpt. 1940: 41-42.

—— and Suehiro, [Amy]

1940 Biological control of cabbage worm, *Pontia rapae*. Hawaii Agr. Expt. Sta. Rpt. 1939: 37.

Holloway, J. K. (See Allen, H. W., Holloway, and Haeussler)

Holloway, T. E.

1919 Parasite introduction as a means of saving sugar. Jour. Econ. Ent. 12: 175-178.

1927 Parasites of the moth stalkborer. Internatl. Soc. Sugar Cane Technol. Cong. Proc. 2: 67-68.

1928 Parasites of the sugar cane moth borer, and their introduction into Louisiana. Planter and Sugar Mfr. 80 (13): 241-242.

1939 Introductions and recoveries of parasites of sugarcane insects in continental United States. Internatl. Soc. Sugar Cane Technol. Cong. Proc. 6: 258-263.

—— Haley, W. E., Loftin, U. C., and Heinrich, C.

1928 The sugar-cane moth borer in the United States. U.S. Dept. Agr. Tech. Bul. 41: 1-76, frontis., figs. 1-25, pl. 1.

—— Loftin, U. C., and Heinrich, C.

1919 The sugar-cane moth borer. U.S. Dept. Agr., Dept. Bul. 746: 1-74, frontis., figs. 1-12, pls. 1-9.

—— and Mathes, R.

1940 The Amazon fly, *Metagonistylum minense*, a parasite of the sugarcane borer. Jour. Econ. Ent. 33: 738-742.

- and Rands, R. D.
 1932 Sugarcane insect pests and diseases with special reference to United States plant quarantines. Jour. Econ. Ent. 25: 461-470.
- Hood, C. E. (See Collins, C. W., and Hood; Pierce, Cushman, and Hood)
- *Hooker, C. W.
 1913 Report of the entomologist. Porto Rico Agr. Expt. Sta. Ann. Rpt. 1912: 34-38.
- Hooker, [W. A.]
 1908 [Tachinid flies parasitic on walnut *Datana*]. Jour. Econ. Ent. 1: 156.
- Hopkins, A. D.
 1893 Catalogue of West Virginia Scolytidae and their enemies. With list of trees and shrubs attacked. W. Va. Agr. Expt. Sta. Bul. 31: 121-168.
 1893a Catalogue of West Virginia forest and shade tree insects. Collected in 1890-1893, including injurious, beneficial and other insects taken on or in some part of the tree examined. W. Va. Agr. Expt. Sta. Bul. 32: 171-251.
- *Hopkins, [L.]
 1953 A tachina fly — Arizona. (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 3 (47[46]): 794.
- *Hopping, G. R.
 1930 [*Zenillia blanda* Osten Sacken reared from *Nepytia phantasmaria* Strecker]. Canad. Insect Pest Rev. 8 (5): 64.
 1937 Sawfly biologies no. 2, *Hemichroa crocea* Geoffroy. Canad. Ent. 69 (11): 243-249, pl. 13.
 1939 [*Pristiphora erichsonii* parasitized by *Bessa selecta*]. Canad. Insect Pest Rev. 17 (4): 277.
- Leech, H. B., and Morgan, C. V. G.
 1943 The larch sawfly, *Pristiphora erichsonii* (Hartig) in British Columbia, with special reference to the cocoon parasites *Mesoleius tenthredinis* Morley and *Tritneptis klugii* (Ratzeburg). Sci. Agr. 24 (2): 53-63, fig. 1.
- Hopping, R.
 1937 Forest insect conditions in British Columbia, in 1936. Canad. Insect Pest Rev. 15 (1): 63-65.
- House, H. L. (See also Coppel and House)
 1949 The laboratory propagation of *Sturmia* sp., a parasite of sawflies. Quebec Soc. Protect. Plants Rpt. 1945-1947 30: 60-62.

Hoveland, N. (See Clark, N., and Hoveland)

*Howard, L. O. (See also Riley and Howard)

- 1894 The army worm (*Leucania unipuncta* Haw.). U.S. Dept. Agr., Dept. Cir., ser. 2, 4: 1-5, figs. 1-3.
- 1895 Some rearings of parasites. Insect Life (U.S. Dept. Agr., Div. Ent., Period. Bul.) (1894) 7 (3): 280.
- 1897 Some insects affecting the hop plant. U.S. Dept. Agr. Div. Ent. Bul., n. ser., 7: 40-51, figs. 35-38.
- 1897a Additional observations on the parasites of *Orygia leucostigma*. U.S. Dept. Agr. Div. Ent. Bul., n. ser., 9: 15-18.
- 1897b A study in insect parasitism. U.S. Dept. Agr., Div. Ent. Tech. Ser. 5: 1-57, figs. 1-24.
- 1898 Additional observations on the parasites of *Orygia leucostigma*. Ontario Ent. Soc. Ann. Rpt. 28: 87-89.
- 1899 A dipterous parasite of *Lachnosterna*. Ent. Soc. Wash. Proc. 4 (3): 198-199.
- 1902 Parasites of the tent caterpillar (*Clisiocampa americana* Harr.). U.S. Dept. Agr. Div. Ent. Bul., n. ser., 38: 109.
- 1904 A tachina parasite of May beetles. U.S. Dept. Agr. Div. Ent. Bul., n. ser., 44: 98.
- 1911 Parasite work. Report of Dr. L. O. Howard, chief of the Bureau of Entomology, Washington, D. C. Mass. State Forester Ann. Rpt. 1910 7: 94-98.
- 1913 Parasite work. Report of Dr. L. O. Howard, chief of the Bureau of Entomology, Washington, D.C. Mass. State Forester Ann. Rpt. 1912 9: 70-73.
- 1914 Parasite work. Report of Dr. L. O. Howard, chief of the Bureau of Entomology, Washington, D.C. Mass. State Forester Ann. Rpt. 1913 10: 83-85.
- 1914a Report on parasites. Ent. Soc. Amer. Ann. 7 (1): 86-88.
- 1916 On the Hawaiian work in introducing beneficial insects. Jour. Econ. Ent. 9: 172-179.
- 1916a On the Hawaiian work in introducing beneficial insects. Hawaii. Planters' Rec. 14 (5): 290-297.
- 1916b Cane borer parasites. Tachinid fly successfully used to fight weevil in Hawaii. Facts about Sugar 3 (23): 367, 369.

- 1918 Report of Dr. L. O. Howard, United States entomologist. Mass. State Forester Ann. Rpt. 1917 14: 45-46.
- 1922 A side line in the importation of insect parasites of injurious insects from one country to another. Natl. Acad. Sci. Proc. 8 (6): 133-139.
- 1926 The parasite element of natural control of injurious insects and its control by man. Jour. Econ. Ent. 19: 271-282.
- and Chittenden, F. H.
- 1907 The catalpa sphinx, (*Ceratomia catalpae* Bdv.). U.S. Dept. Agr., Bur. Ent. Cir. 96: 1-7, figs. 1-2.
- 1909 The green-striped maple worm, (*Anisota rubicunda* Fab.). U.S. Dept. Agr., Bur. Ent. Cir. 110: 1-7, figs. 1-3.
- 1916 The catalpa sphinx. U.S. Dept. Agr. Farmers' Bul. 705: 1-9, figs. 1-5.
- and Fiske, W. F.
- 1911 The importation into the United States of the parasites of the gypsy moth and the brown-tail moth: A report of progress, with some consideration of previous and concurrent efforts of this kind. U.S. Dept. Agr. Bur. Ent. Bul., n. ser., 91: 1-312, figs. 1-74, pls. 1-28.
- Howard, N. F. (See also Landis, B. J., and Howard)
- 1924 The Mexican bean beetle in the East. U.S. Dept. Agr. Farmers' Bul. 1407: 1-14, figs. 1-10.
- 1930 The Mexican bean beetle in the East and its control. U.S. Dept. Agr. Farmers' Bul. 1624: 1-14, figs. 1-10.
- and Landis, B. J.
- 1936 Parasites and predators of the Mexican bean beetle in the United States. U.S. Dept. Agr., Dept. Cir. 418: 1-12, figs. 1-3.
- Hoxie, R. P. (See Wellso and Hoxie)
- Hsiao, T.-H., and Holdaway, F. G.
- 1962 A method of rearing a tachinid parasite, *Lydella grisescens* R-D, in an unnatural host. Ent. Soc. Amer., No. Cent. Branch, Proc. 17: 144-145.
- 1966 Seasonal history and host synchronization of *Lydella grisescens* (Diptera: Tachinidae) in Minnesota. Ent. Soc. Amer. Ann. 59 (1): 125-133, figs. 1-4.
- Hubbard, H. G.
- 1885 Appendix I. Report of H. G. Hubbard. U.S. Dept. Agr., Ent. Comn. Rpt. 4: 5-16.

- Hubbell, T. H.
 1936 A monographic revision of the genus *Ceuthophilus* (Orthoptera, Gryllacrididae, Rhabdophorinae). Fla. Univ. Pub. Biol. Sci. Ser. 2 (1): 1-551, frontis., pls. 1-38.
- Huber, L. L., and Neiswander, C. R.
 1924 European corn borer investigations in Ohio. Jour. Econ. Ent. 17: 125-128.
- Hudon, M.
 1957 *Phryxe vulgaris* (Fall.) (Diptera: Tachinidae), a new parasite of *Pieris rapae* (L.) (Lepidoptera: Pieridae) in Canada. Canad. Ent. 89 (12): 546.
 1960 Some aspects of seasonal development of the European corn borer, *Ostrinia nubilalis* (Hubn.) (Lepidoptera: Pyralidae), in southwestern Quebec. Quebec Soc. Protect. Plants Ann. Rpt. 41: 121-128.
- Hudson, H. F.
 1920 Some notes on the life history of our common June beetles. Ontario Ent. Soc. Ann. Rpt. 50: 81-83.
- and Wood, A. A.
 1929 The life-history of the white cutworm, *Euxoa scandens* Riley. Ontario Ent. Soc. Ann. Rpt. 60: 67-70.
- Hughes, K. M. (See Steinhaus and Hughes)
- Hunt, R. H. (See Wickman and Hunt)
- Hunter, W. D., and Pierce, W. D.
 1912 Mexican cotton-boll weevil. [U.S.] 62d Cong., 2d Session, Senate Doc. 305: 1-188, figs. 1-34, pls. 1-22. [U.S. Dept. Agr. Bur. Ent. Bul., n. ser. 114: 1-188, figs. 1-34, pls. 1-22].
- Pratt, F. C., and Mitchell, J. D.
 1912 The principal cactus insects of the United States. U.S. Dept. Agr. Bur. Ent. Bul., n. ser., 113: 1-71, figs. 1-8, pls. 1-7.
- Hyche, L. L. (See DuRant and Hyche)
- Hyslop, J. A.
 1910 The smoky crane-fly. U.S. Dept. Agr. Bur. Ent. Bul., n. ser., 85 (7): 119-132, figs. 60-66.
 1912 The alfalfa looper. U.S. Dept. Agr. Bur. Ent. Bul., n. ser., 95 (7): 109-118, figs. 45-51, pl. 11.
 1915 Observations on the life history of *Meracantha contracta* (Beauv.). Psyche 22 (2): 44-48, figs. 1-2, pl. 4.

- 1916 The host of *Zelia vertebrata* (Diptera, Dexiidae). *Psyche* 23 (1): 24-25.
- 1936 The periodical cicada. *Jour. Econ. Ent.* 29: 671-676.
- Illingworth, J. F.
- 1914 Further notes on the breeding of the tachinid fly, parasitic on the cane beetle borer. *Jour. Econ. Ent.* 7: 390-398, pl. 10.
- 1923 A list with notes of the insects found on the Parker Ranch, June, 1922. *Hawaii. Ent. Soc. Proc.* 1922 5 (2): 277-279.
- 1937 *Plusia chalcites* Esp. *Hawaii. Ent. Soc. Proc.* 1936 9 (3): 354.
- Ingerson, H. G.
- 1918 The striped peach worm. U.S. Dept. Agr. Bul. 599: 1-14, pls. 1-4.
- Ingles, L. G.
- 1929 The seasonal and associational distribution of the fauna of the upper Santa Ana River wash. *Jour. Ent. and Zool.* 21 (2): 57-96.
- Ingram, J. W. (See also Douglas and Ingram; Mathes et al.; Scaramuzza and Ingram)
- 1941 Sugarcane borer. *Insect Pest Survey Bul.* 21 (10): 799-800.
- and Bynum, E. K.
- 1932 Observations on the sugarcane beetle in Louisiana. *Jour. Econ. Ent.* 25: 844-849.
- 1941 The sugarcane borer. U.S. Dept. Agr. Farmers' Bul. 1884: 1-17, figs. 1-5.
- Bynum, E. K., and Douglas, W. A.
- 1939 A summary of investigations on the sugarcane beetle in Louisiana. *Internatl. Soc. Sugar Cane Technol. Cong. Proc.* 6: 478-482.
- Bynum, E. K., and Mathes, R.
- 1951 Insect pests of sugar cane in continental United States. *Internatl. Soc. Sugar Cane Technol. Cong. Proc.* 7: 395-401.
- Bynum, E. K., Mathes, R., Haley, W. E., and Charpentier, L. J.
- 1951 Pests of sugarcane and their control. U.S. Dept. Agr., Dept. Cir. 878: 1-38, figs. 1-19.
- Jaynes, H. A., and Lobdell, R. N.
- 1939 Sugarcane pests in Florida. *Internatl. Soc. Sugar Cane Technol. Cong. Proc.* 6: 89-98.

- Mathes, R., Wilson, J. W., and Hayslip, N. C.
 1951 Biological control of the sugar-cane borer in continental United States. *Internatl. Soc. Sugar Cane Technol. Cong. Proc.* 7: 354-360.
- Isely, D.
 1927 The striped cucumber beetle. *Ark. Agr. Expt. Sta. Bul.* 216: 1-36, pls. 1-2.
 1929 The southern corn rootworm. *Ark. Agr. Expt. Sta. Bul.* 232: 1-31, figs. 1-11, pl. 1.
 1930 The biology of the bean leaf-beetle. *Ark. Agr. Expt. Sta. Bul.* 248: 1-20, figs. 1-10.
- Ives, W. G. H. (See also Nairn et al.; Turnock and Ives)
 1964 Problems encountered in the development of life tables for insects. *Manitoba Ent. Soc. Proc.* 20: 34-44.
- Muldrew, J. A., Turnock, W. J., and Nairn, L. D.
 1956 Summary of a panel discussion of larch sawfly problems. *Manitoba Ent. Soc. Proc.* 12: 11-20.
- and Prentice, R. M.
 1959 Estimation of parasitism of larch sawfly cocoons by *Bessa harveyi* Tnsd. in survey collections. *Canad. Ent.* 91 (8): 496-500.
- and Turnock, W. J.
 1959 Estimation of cocoon populations of the larch sawfly, *Pristiphora erichsonii* (Hartig). *Canad. Ent.* 91 (10): 650-661, figs. 1-5.
- Turnock, W. J., Buckner, C. H., Heron, R. J., and Muldrew, J. A.
 1968 Larch sawfly population dynamics: techniques. *Manitoba Ent.* 2: 5-36, figs. 1-22.
- Jackson et al.
 1967 Southern green stink bug (*Nezara viridula*). (U.S. Dept. Agr.) *Coop. Econ. Insect Rpt.* 17 (35): 822.
- *Jackson and Peters, [L. L.]
 1958 European corn borer (*Pyrausta nubilalis*) — Missouri. (U.S. Dept. Agr.) *Coop. Econ. Insect Rpt.* 8 (29): 626.
- Jackson, A. D. (See Sanderson and Jackson)
- Jackson, C. G. (See also Bryan, D. E., Jackson, and Patana; Bryan, D. E., Jackson, and Stoner; Butler, G. D., Jr., Bryan, and Jackson)

- Bryan, D. E., and Patana, R.
1969 Laboratory studies of *Eucelatoria armigera*, a tachinid parasite of *Heliothis* spp. Jour. Econ. Ent. 62: 907-910.
- Butler, G. D., Jr., and Bryan, D. E.
1969 Time required for development of *Voria ruralis* and its host, the cabbage looper, at different temperatures. Jour. Econ. Ent. 62: 69-70.
- *James, H. C.
1952 Biological control. Investigations and application of results (EE.bc.1). Jamaica Dept. Agr. Bul., n. ser., 50: 34.
1953 Biological control. Investigations and application of results (EE.bc.1). Jamaica Dept. Agr. Bul., n. ser., 53: 26-27.
- James, H. G.
1963 Larval habitats, development, and parasites of some Tabanidae (Diptera) in southern Ontario. Canad. Ent. 95 (11): 1223-1232, figs. 1-3.
- James, M. T.
1945 A new larvaevorid parasite of the social butterfly *Eucheira socialis* Westwood (Diptera). Wash. Acad. Sci. Jour. 35 (10): 328-330, fig. 1.
1947 A review of the larvaevorid flies of the tribe Leskiini with the setulose first vein (R_1). U.S. Natl. Mus. Proc. 97 (3212): 91-115, fig. 3.
1955 A new tachina fly of economic importance. Pan-Pacific Ent. 31 (2): 83-85, fig. 1.
- Janes, M. J. (See Knowlton and Janes; Knowlton, Janes, and Thomas)
- Jardine, J. T.
1926 Director's biennial report. Oreg. Agr. Expt. Sta. Dir. Bien. Rpt. 1924-1926: 1-133.
1928 Director's biennial report. Oreg. Agr. Expt. Sta. Dir. Bien. Rpt. 1926-1928: 1-133.
- Jarvis, J. L.
1966 Studies of *Phyllophaga anxia* (Coleoptera: Scarabaeidae) in the sandhills area of Nebraska. Kans. Ent. Soc. Jour. 39 (3): 401-409.
- and York, G. T.
1961 Population fluctuations of *Lydella grisescens*, a parasite of the European corn borer. Jour. Econ. Ent. 54: 213-214.

- Jaynes, H. A. (See also Allen, H. W., and Jaynes; Dowden, Jaynes, and Carolin; Ingram, Jaynes, and Lobdell)
- 1933 The parasites of the sugarcane borer in Argentina and Peru, and their introduction into the United States. U.S. Dept. Agr. Tech. Bul. 363: 1-26, figs. 1-10.
- 1938 Introduction and recovery in Florida and Louisiana of parasites of the sugarcane borer. Jour. Econ. Ent. 31: 93-95.
- 1939 Further attempts to establish *Lixophaga diatraeae* (Townsend) and other sugarcane borer parasites in Louisiana and Florida, with recoveries in 1936 and 1937. Internatl. Soc. Sugar Cane Technol. Cong. Proc. 6: 245-258.
- and Drooz, A. T.
- 1952 The importance of parasites in the spruce budworm infestations in New York and Maine. Jour. Econ. Ent. 45: 1057-1061.
- Jenne, E. L. (See also Quaintance and Jenne)
- 1909 The codling moth in the Ozarks. U.S. Dept. Agr. Bur. Ent. Bul., n. ser., 80 (1): 1-31, 8 figs.
- Jimenez J., E.
- 1959 El empleo de enemigos naturales para el control de insectos que constituyen plagas agrícolas en la República Mexicana. Chapingo 12 (73): 191-208.
- Johannsen, O. A.
- 1910 Insect notes for 1909. Maine Agr. Expt. Sta. Bul. 177: 21-44, figs. 10-57.
- 1913 Spruce bud worm and spruce leaf miners. Maine Agr. Expt. Sta. Bul. 210: 13-36, figs. 1-9.
- 1935 Aquatic Diptera. Part II. Orthorrhapha-Brachycera and Cyclorrhapha. N.Y. (Cornell) Agr. Expt. Sta. Mem. 177: 1-62, pls. 1-12.
- Patch, E. M., Morse, W. J., and Lewis, C. E.
- 1911 Apple tree enemies of Maine. Maine Comnr. Agr. Rpt. 9: 317-379, figs. 1-28, 30-55.
- Johansen, C. A.
- 1957 History of biological control of insects in Washington. Northwest Sci. 31 (2): 57-79.
- Johnson, C. W.
- 1907 A new genus and species of the family Tachinidae, parasitic on *Archips cerasivorana*. Psyche 14 (1): 9-10.

- 1908 The Diptera of the Bahamas, with notes and description of one new species. *Psyche* 15 (4): 69-80.
- 1910 Order Diptera. N.J. State Mus. Ann. Rpt. 1909: 703-814, figs. 293-340.
- 1912 Dipterological notes. *Psyche* 19 (3): 102-104.
- 1913 Insects of Florida. I. Diptera. Amer. Mus. Nat. Hist. Bul. 32 (3): 37-90.
- 1916 Parasites of *Archips cerasivorana* Fitch. *Psyche* 23 (3): 81.
- 1919 A revised list of the Diptera of Jamaica. Amer. Mus. Nat. Hist. Bul. 41 (8): 421-449.
- 1925 Fauna of New England. 15. List of the Diptera or two-winged flies. Boston Soc. Nat. Hist. Occas. Papers 7: 1-326.
- 1925a Diptera of the Harris collection. Boston Soc. Nat. Hist. Proc. 38 (2): 57-99.
- 1927 The insect fauna with reference to the flora and other biological features. Pt. I, 247 pp., 1 fig. In Procter, W., ed., Biological survey of the Mount Desert region. Philadelphia.
- Johnson, J. P. (See Britton and Johnson)
- *Jones, C. M.
- 1953 Biology of Tabanidae in Florida. *Jour. Econ. Ent.* 46: 1108-1109.
- and Anthony, D. W.
- 1964 The Tabanidae (Diptera) of Florida. U.S. Dept. Agr. Tech. Bul. 1295: i-ii, 1-85, figs. 1-18.
- Jones, D. W. (See also Arbuthnot et al.; Arbuthnot et al.; Clark, S. W., and Jones; Cory et al.)
- 1929 Imported parasites of the European corn borer in America. U.S. Dept. Agr. Tech. Bul. 98: 1-27, figs. 1-24.
- and Arbuthnot, K. D.
- 1949 Report of European corn borer parasite recoveries. *Ent. Soc. Amer.*, No. Cent. Branch, Proc. 4: 56-57.
- Carter, S. W., and Cosenza, A. V.
- 1947 The field status of parasites of the European corn borer at the close of the 1945 season. U.S. Dept. Agr., Bur. Ent. and Plant Quar., Insect Pest Survey, Spec. Sup. 1947 (3): 1-6.
- Walker, H. G., and Anderson, L. D.
- 1939 The European corn borer on the eastern shore of

- Virginia. Va. Truck Expt. Sta. Bul. 102: 1621-1648, figs. 289-292.
- Jones, F. M. (See also Bird, H., and Jones)
 1948 Notes on *Melanophora roralis* (Linn.) (Diptera).
 Psyche 55 (1): 31-34.
- Jones, M. L.
 1955 Eradication of incipient infestations of new insect
 pests. Calif. Dept. Agr. Bul. (1954) 43
 (4): 173-179, 8 figs.
- and Forbes, A. G.
 1952 Western grape skeletonizer (*Harrisina brillians*).
 Pp. 159-162, 1 fig. In Armitage, H. M. Bureau of
 Entomology. Calif. Dept. Agr. Bul. (1951) 40
 (4): 147-185, 9 figs.
- 1956 Eradication of incipient infestations of new insect
 pests. Calif. Dept. Agr. Bul. 45 (2): 149-156.
- Jones, P. A. (See Coppel and Jones)
- Jones, R. W.
 1885 Appendix II. Report of Prof. R. W. Jones. U.S.
 Dept. Agr., Ent. Comn. Rpt. 4: 17-23.
- Jones, T. H. (See also Collins, C. W., and Jones)
 1913 Some notes on *Laphygma frugiperda* S. and A. in
 Porto Rico. Jour. Econ. Ent. 6: 230-236.
- 1915 Insects affecting vegetable crops in Porto Rico. U.S.
 Dept. Agr. Bul. 192: 1-11, pls. 1-4.
- 1917 The sweet-potato leaf-folder. U.S. Dept. Agr. Bul.
 609: 1-12, figs. 1-4.
- 1918 The southern green plant-bug. U.S. Dept. Agr.,
 Dept. Bul. 689: 1-27, figs. 1-14.
- 1918a Miscellaneous truck-crop insects in Louisiana. U.S.
 Dept. Agr. Bul. 703: 1-19, figs. 1-5, pls. 1-5.
- and Bradley, W. G.
 1923 Observations on Tabanidae (horseflies) in Loui-
 siana. Jour. Econ. Ent. 16: 307-312.
- and Wolcott, G. N.
 1922 The caterpillars which eat the leaves of sugar cane in
 Porto Rico. Porto Rico Dept. Agr. Jour. 6
 (1): 38-50, figs. 10-19.
- Joutel, L. H. (See Felt and Joutel)
- Joyner, H. L. (See Dugas et al.)
- Judd, W. W.
 1954 Four species of leaf-tying moths and their dipterous
 and hymenopterous parasites reared from may-
 apple, *Podophyllum peltatum* L. Amer. Micros.
 Soc. Trans. 73 (4): 401-404.

- 1955 *Hyponomeuta multipunctella* Clemens (Lepidoptera: Hyponomeutidae) and its dipterous and hymenopterous parasites. Amer. Micros. Soc. Trans. 74 (3): 265-267.
- Kaloostian, G. H. (See Donohoe and Kaloostian)
- Kamran, M. A.
- 1968 Seasonal fluctuations in the abundance of the monkey pod moth, *Polydesma umbricola*, in Hawaii. Jour. Econ. Ent. 61: 1007-1012, figs. 1-5.
- Kapler, J. E., and Benjamin, D. M.
- 1969 The biology and ecology of the red-pine sawfly in Wisconsin. Forest Sci. 6 (3): 253-268, 14 figs.
- Kaufman, W. (See Stevenson, W. A., Kaufman, and Sheets)
- Kawamura, [K. F.]
- 1960 Southern green stink bug (*Nezara viridula*). (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 19 (37): 726.
- and Funasaki, [G.]
- 1969 [Southern green stink bug (*Nezara viridula*)]. (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 19 (32): 632.
- Keen, F. P.
- 1952 Insect enemies of western forests. U.S. Dept. Agr. Misc. Pub. 273: 1-280, figs. 1-111.
- 1958 Cone and seed insects of western forest trees. U.S. Dept. Agr. Tech. Bul. 1169: i-vi, 1-168, figs. 1-57.
- Keifer, H. H.
- 1945 Systematic entomology. Calif. Dept. Agr. Monthly Bul. (1944) 33 (4): 248-252.
- Keister, M. (See Buck and Keister)
- Kelleher, J. S.
- 1960 Colorado potato beetle (*Leptinotarsa decemlineata* Say). Canad. Insect Pest Rev. 38 (2): 122.
- 1960a A summary of parasite and predator liberations in Canada and of insect shipments from Canada in 1960. Canad. Insect Pest Rev. 38 (8): 266-281.
- 1966 The parasite *Doryphorophaga doryphorae* (Diptera: Tachinidae) in relation to populations of the Colorado potato beetle in Manitoba. Ent. Soc. Amer. Ann. 59 (6): 1059-1061, figs. 1-2.

Kellicott, [D. S.]

- 1885 On the larval period of *Harmonia pini*, and a parasite of same. Ent. Amer. 1 (9): 171-173.

Kellogg, V. L.

- 1905 American insects. vii + 674 pp., 812 figs., 1 pl. New York.

Kelly, G. D., and Middlekauff, W. W.

- 1961 Biological studies of *Dissosteira spurcata* Saussure with distributional notes on related California species (Orthoptera, Acrididae). Hilgardia 30 (14): 395-424, figs. 1-13.

Kelsey, L. P. (See Whiteside, Burbutis, and Kelsey)

Kelsheimer, E. G. (See Hayslip et al.)

Kendrick, R. L. (See Rolston and Kendrick)

Kido, G. S. (See Smith, L. M., and Kido)

Kim, J.

- 1963 A parasitic tachina fly (*Trichopoda pennipes*). (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 13 (10): 186-187.

- 1963a A parasitic tachina fly (*Trichopoda pennipes*). (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 13 (20): 535.

- 1964 *Trichopoda pennipes pilipes* Fabricius. Hawaii. Ent. Soc. Proc. 1963 18 (3): 328.

- 1964a *Trichopoda pennipes* var. *pilipes* Fabricius. Hawaii. Ent. Soc. Proc. 1963 18 (3): 333.

—— and Miyahira, N.

- 1964 A parasitic tachina fly (*Trichopoda pennipes* (F.)). (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 14 (42): 1155.

*King, J. L. (See also Clausen, King, and Teranishi)

- 1928 Japanese beetle now faces attack by five imported parasites. (U.S. Dept. Agr.) Yearb. Agr. 1927: 403-407, figs. 130-136.

- 1930 Imported parasites of *Popillia japonica* and *Aserica castanea*. N.J. State Dept. Agr. Ann. Rpt. 1929-1930 15: 158-162, fig. 20.

- 1931 The present status of the established parasites of *Popillia japonica* Newman. Jour. Econ. Ent. 24: 453-462, figs. 34-37.

- 1937 Liberations of Japanese beetle parasites in the eastern states in 1937. Insect Pest Survey Bul. 17 (9) Sup.: 473-482, 2 figs.

- 1939 Colonization of Japanese beetle parasites in the eastern states in 1938. *Insect Pest Survey Bul.* 19 (1) Sup.: 25-28.
- and Hallock, H. C.
- 1925 A report on certain parasites of *Popillia japonica* Newm. *Jour. Econ. Ent.* 18: 351-356, figs. 20-21.
- and Parker, L. B.
- 1941 Summary of Japanese beetle parasite liberations, including the year 1940. *Insect Pest Survey Bul.* 21 (3) Sup.: 127-138.
- King, K. M.
- 1925 [Reduction of the population of *Sidemia devastator* Brace in Saskatchewan by parasites in 1924]. *Canad. Insect Pest Rev.* 3 (2): [14].
- 1928 *Barathra configurata* Wlk., an armyworm with important potentialities on the northern prairies. *Jour. Econ. Ent.* 21: 279-293, figs. 19-20, pl. 6.
- and Atkinson, N. J.
- 1928 The biological control factors of the immature stages of *Euxoa ochrogaster* Gn. (Lepidoptera, Phalaenidae) in Saskatchewan. *Ent. Soc. Amer. Ann.* 21 (2): 167-188, fig. 1.
- King, V., and Barber, G. W.
- 1921 Controlling the army-worm in southeast Missouri. *Jour. Econ. Ent.* 14: 486-488, pl. 6.
- Kirk, H. B.
- 1913 The walnut bud-moth. *Acrobasis caryae* Grote. *Conn. Agr. Expt. Sta. Ann. Rpt.* 1912 36: 253-258.
- Kirkpatrick, J.
- 1861 The army worm. *Ohio State Bd. Agr. Ann. Rpt.* 1860 15: 350-358.
- Knab, F.
- 1909 Notes on tachinid parasites of Chrysomelidae. *Psyche* 16 (2): 34-35.
- Knight, F. B. (See Allen, D. C. et al.)
- Knight, H. H.
- 1916 The army-worm in New York in 1914. *Leucania unipuncta* Haworth. Order, Lepidoptera. Family, Noctuidae. N.Y. (Cornell) *Agr. Expt. Sta. Bul.* 376: 749-765, pls. 37-44.
- Knowlton, G. F. (See also Davis, C. J., and Knowlton)
- 1936 Utah Tachinidae — notes. *Utah Acad. Sci., Arts and Letters, Proc.* 13: 241-244.

- 1940 The European earwig in Utah. Jour. Econ. Ent. 33: 204.
- and Allen, M. W.
- 1937 Oblique-banded leaf roller, a dewberry pest in Utah. Jour. Econ. Ent. 30: 780-785, fig. 1.
- Dorst, H. E., Lieberman, F. V., Thornley, H. F., and Orr, L. W.
- 1953 Summary of insect conditions — 1953: Utah. (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 3 (52): 863-869.
- and Hansen, L. L.
- 1938 Notes on berry insects of Utah. Utah Acad. Sci., Arts and Letters, Proc. 15: 127-130, fig. 1.
- and Hanson, W. J.
- 1949 Some Utah Tachinidae. Utah Agr. Expt. Sta. Mimeograph Ser. 361: 1-7.
- and Harmston, F. C.
- 1937 Utah Diptera. Utah Acad. Sci., Arts and Letters, Proc. 14: 141-149.
- and Janes, M. J.
- 1931 Notes on some beneficial Utah Diptera. Utah Acad. Sci., Arts and Letters, Proc. 8: 147-148.
- Janes, M. J., and Thomas, W. L.
- 1933 Some Utah Diptera. Utah Acad. Sci. Proc. 10: 155-157.
- Knoll, J. N. (See also Champlain and Knoll)
- 1932 Observations on three important forest insects. Jour. Econ. Ent. 25: 1196-1203, pls. 38-39.
- Koebele, A.
- 1890 Report on California insects. U.S. Dept. Agr., Div. Ent. Bul. 22: 85-94.
- 1891 Report of experiments with resin compounds on *Phylloxera*, and general notes on California insects. U.S. Dept. Agr., Div. Ent. Bul. 23: 37-44.
- 1908 Insect investigations in Mexico. Hawaii. Planters' Monthly 27 (12): 507-513.
- Krauss, [N. L. H.] (See also Davis, C. J., and Krauss)
- 1947 *Lema trilineata californica* Schaeffer. Hawaii. Ent. Soc. Proc. 1946 13 (1): 4.
- Krombein, K. V.
- 1967 Trap-nesting wasps and bees: life histories, nests, and associates. vi + 570 pp., frontis., 29 pls. Washington.

Kuitert, L. C., and Tissot, A. N.

- 1957 Influence of cultural practices on the incidence and control of insect infestations in flue-cured tobacco. Fla. Agr. Expt. Sta. 1957: 93-94.

Kulman, H. M.

- 1965 Natural control of the eastern tent caterpillar and notes on its status as a forest pest. Jour. Econ. Ent. 58: 66-70.
- 1965a Natural control of the bagworm and notes on its status as a forest pest. Jour. Econ. Ent. 58: 863-866.

— and Hodson, A. C.

- 1961 Parasites of the jack-pine budworm, *Choristoneura pinus*, with special reference to parasitism at particular stand locations. Jour. Econ. Ent. 54: 221-224.

Lacroix, D. S. (See Franklin, H. J., and Lacroix)

Lafler, H. A.

- 1896 A new parasite. Ent. News 7 (2): 62-63.

Lambert, R.

- 1941 Les insectes forestiers du Québec 1940. Forêt Québécoise 3 (3): 29-45, figs. 4-9.

Landis, B. J. (See also Howard, N. F., and Landis)

- 1935 Striped cucumber beetle (*Diabrotica vittata* Fab.). Insect Pest Survey Bul. 15 (7): 345.
- 1936 Colorado potato beetle (*Leptinotarsa decemlineata* Say). Insect Pest Survey Bul. 16 (6): 284.
- 1963 Colorado potato beetle (*Leptinotarsa decemlineata*) — Washington. (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 13 (26): 714.

— and Howard, N. F.

- 1940 *Paradexodes epilachnae*, a tachinid parasite of the Mexican bean beetle. U.S. Dept. Agr. Tech. Bul. 721: 1-31, figs. 1-23.

Lange, W. H. (Jr.)

- 1937 California pine tip moths of the genus *Rhyacionia*. South. Calif. Acad. Sci. Bul. 36 (1): 25-34, pls. 11-15.
- 1941 The artichoke plume moth and other pests injurious to the globe artichoke. Calif. Univ. Agr. Expt. Sta. Bul. 653: 1-71, figs. 1-29.
- 1945 *Autographa egena* (Guen.) a periodic pest of beans in California. Pan-Pacific Ent. 21 (1): 13.

- 1950 Biology and systematics of plume moths of the genus *Platyptilia* in California. *Hilgardia* 19 (19): 561-668, figs. 1-8, pls. 1-16.
- Langford, G. S. (See Cory et al.; Cory and Langford)
- Langston, R. L. (See Smith, O. J., and Langston)
- Lannon, A. J.
- 1952 Division of Entomology and Plant Industry. R.I. Dept. Agr. and Conserv. Ann. Rpt. 17: 7-37.
- 1954 Division of Entomology. R.I. Dept. Agr. and Conserv. Ann. Rpt. 18: 51-71.
- Lavigne, R. J., and Pfadt, R. E.
- 1966 Parasites and predators of Wyoming rangeland grasshoppers. *Wyo. Univ. Agr. Expt. Sta. Sci. Monog.* 3: 1-31, figs. 1-2.
- Lawson, F. R.
- 1959 The natural enemies of the hornworms on tobacco (Lepidoptera: Sphingidae). *Ent. Soc. Amer. Ann.* 52 (6): 741-755, figs. 1-3.
- Learned, E. T.
- 1927 Note on leaf-oviposition of *Zenillia futilis* (Dipt., Tachinidae). *Brooklyn Ent. Soc. Bul.* 22 (4): 219.
- Leaver, F. S.
- 1946 Division of Entomology and Plant Industry. R.I. Dept. Agr. and Conserv. Ann. Rpt. 11: 30-43.
- LeBaron, W.
- [1871] Insects injurious to the apple tree. Noxious [and Beneficial] Insects of State of Ill., [State Ent.] Rpt. [2]: 13-46, 4 figs.
- [1871a] Insects injurious to the potato. Noxious [and Beneficial] Insects of State of Ill., [State Ent.] Rpt. [2]: 63-78, 1 fig.
- 1872 Insects injurious to the apple. Noxious [and Beneficial] Insects of State of Ill., [State Ent.] Rpt. [3]: 99-133, 3 figs.
- Lee, A.
- 1938 Report of the Puerto Rico Experiment Station, 1937. Puerto Rico Expt. Sta. (Mayaguez) Rpt. 1937: i-ii, 1-115, figs. 1-28.
- 1939 Report of the Puerto Rico Experiment Station, 1938. Puerto Rico Expt. Sta. (Mayaguez) Rpt. 1938: i-ii, 1-137, figs. 1-36.
- 1940 Report of the Puerto Rico Experiment Station, 1939. Puerto Rico Expt. Sta. (Mayaguez) Rpt. 1939: i-ii, 1-126, figs. 1-43.

- 1942 Report of the Puerto Rico Experiment Station, 1940. Puerto Rico Expt. Sta. (Mayaguez) Rpt. 1940: i-ii, 1-103, figs. 1-38.
- Leech, H. B. (See also Hopping, G. R., Leech, and Morgan)
1936 [*Bessa selecta* possible parasite of larch sawfly].
Canad. Insect Pest Rev. 14(2): 99.
- and Farris, S. H.
1959 *Achaetoneura datanarum* reared from *Antheraea polyphemus* in British Columbia (Diptera: Tachinidae). Ent. Soc. Brit. Columbia, Proc. 56: 22.
- Leiby, R. W. (See also Herrick and Leiby)
1925 Insect enemies of the pecan in North Carolina. N.C. Dept. Agr. Bul. 1925 (Feb.): 1-67, figs. 1-63.
- Lejeune, R. R.
1951 Some ecological factors governing populations of the larch sawfly, *Pristiphora erichsonii* (Htg.).
Canad. Ent. 83 (6): 152-156.
1951a Part IV. The larch sawfly problem. Forestry Chron. 27 (1): 22-25.
- and Hildahl, V.
1954 A survey of parasites of the larch sawfly (*Pristiphora erichsonii* (Hartig)) in Manitoba and Saskatchewan. Canad. Ent. 86 (8): 337-345.
- and Silver, G. T.
1961 Parasites and hyperparasites of the satin moth, *Stilpnotia salicis* Linnaeus, (Lymantriidae) in British Columbia. Canad. Ent. 93 (6): 456-467.
- Leonard, M. D.
1916 A tachinid parasite reared from an adult capsid (Dip., Hom.). Ent. News 27 (5): 236.
1920 A dipterous parasite of the parsnip webworm (*Depressaria heracliana* Linn.). Jour. Econ. Ent. 13: 491-492.
- and Mills, A. S.
1931 A preliminary report on the lima bean pod-borer and other legume pod-borers in Porto Rico. Jour. Econ. Ent. 24: 466-473.
- and Seín, F., Jr.
1932 Observations on some factors which may affect the abundance of *Diatraea saccharalis* in Porto Rico. Internatl. Soc. Sugar Cane Technol. Cong. Proc. 4, Bul. 92: 1-2.
- LeRoux, E. J. (See also Paradis and LeRoux)
1961 Effects of "modified" and "commercial" spray

- programs on the fauna of apple orchards in Quebec. Soc. Ent. du Quebec, Ann. (1960) 6: 87-121, figs. 1-2.
- Leuck, D. B.
 1966 Biology of the lesser cornstalk borer in south Georgia. Jour. Econ. Ent. 59: 797-801.
- and Dupree, M.
 1965 Parasites of the lesser cornstalk borer. Jour. Econ. Ent. 58: 779-780.
- Lewis, C. E. (See Johannsen et al.)
- *Lewis, H. C.
 1925 Parasitism of the strawberry leaf-roller *Ancylis comptana* Frohl. Jour. Econ. Ent. 18: 609-612.
- Lewis, W. J., and Brazzel, J. R.
 1968 A three-year study of parasites of the bollworm and the tobacco budworm in Mississippi. Jour. Econ. Ent. 61: 673-676.
- Lieberman, F. V. (See Knowlton et al.)
- Lilly, J. H. (See Harris and Lilly)
- Lintner, J. A.
 1883 Reports on the injurious and other insects of the State of New York. Rpt. 1: First annual report [for 1882], xxii + 381 pp., 84 figs. Albany, N.Y.
- 1886 Reports on the injurious and other insects of the State of New York. Rpt. 2: [for 1885], xiv + 265 pp., 68 figs. Albany, N.Y.
- 1887 Report of the state entomologist to the regents of the University of the State of New York, for the year 1885. N.Y. State Mus. Ann. Rpt. [1885] 39: 77-125.
- 1889 Reports on the injurious and other insects of the State of New York, Rpt. 5: [for 1888], pp. 145-347, figs. 1-50. Albany, N. Y.
- 1893 Eighth report on the injurious and other insects of the State of New York for the year 1891. N.Y. State Mus. Rpt. 45: 103-320, figs. 1-53.
- 1896 Notes on some of the insects of the year in the State of New York. U.S. Dept. Agr. Div. Ent. Bul., n. ser., 6: 54-63.
- 1897 Twelfth report on the injurious and other insects of the State of New York for the year 1896. N.Y. State Mus. Rpt. 50: 163-399, figs. 1-10, pls. 1-15.

- Little, V. A., and Martin, D. F.
 1941 Cotton insects of the United States. 130 pp., 12
 figs. Minneapolis, Minn.
- *Lloyd, D. C.
 1944 A study of the codling moth and its parasites in
 California. Sci. Agr. 24 (10): 456-473, fig. 1.
- Lloyd, J. T.
 1920 An aquatic dipterous parasite, *Ginglymyia acirostris*
 Towns., and additional notes on its lepidopterous
 host, *Elophila fulicalis*. N.Y. Ent. Soc. Jour.
 (1919) 27 (4): 263-265, pl. 25.
- Loan, C. C.
 1963 Parasitism of the dogwood flea-beetle, *Altica corni*,
 in Ontario. Jour. Econ. Ent. 56: 537-538.
- Lobdell, R. N. (See also Ingram, Jaynes, and Lobdell)
 1937 Studies upon the prevalence and control of the
 sugarcane moth borer, *Diatraea saccharalis* Fab.,
 in south Florida. Fla. Agr. Expt. Sta. Ann. Rpt.
 1937: 146.
- Lochhead, W.
 1919 Class book of economic entomology with special
 reference to the economic insects of the northern
 United States and Canada. xiv + 436 pp., 257
 figs. Philadelphia.
- Loftin, U. C. (See Holloway et al.; Holloway, Loftin, and
 Heinrich)
- Look, [W. C.]
 1948 *Agrotis coniotis* Hampson. Hawaii. Ent. Soc. Proc.
 1947 13 (2): 222.
- Lovett, A. L. (See also Cole, F. R., and Lovett)
 1915 The variegated cutworm. *Peridroma margaritosa*
saucia Hubn. Oreg. Agr. Col. Expt. Sta. Bien.
 Crop Pest and Hort. Rpt. (1913-1914)
 2: 141-149, figs. 28-30, pls. 2-3.
 1915a The tomato worms. *Phlegethontius sexta* Johan.
Phlegethontius quinquemaculata Haw. Oreg. Agr.
 Col. Expt. Sta. Bien. Crop Pest and Hort. Rpt.
 (1913-1914) 2: 170-172, fig. 38, pl. 7.
- *Lowry, P. R.
 1927 The stalk borer. Life history in New Hampshire,
 1925-1926. N.H. Agr. Expt. Sta. Tech. Bul.
 34: 1-23.

Lowry, Q. S.

- 1917 An outbreak of the eight-spotted forester, *Alypia octomaculata* Fabr., in New Haven, Conn. Jour. Econ. Ent. 10: 47-48.
- 1917a Outbreak of the eight-spotted forester. *Alypia octomaculata* Fabr. Order Lepidoptera, family Zygaenidae. In Britton, W. E. Sixteenth report of the state entomologist of Connecticut. Conn. Agr. Expt. Sta. Ann. Rpt. 1916 40: 118-122, pls. 10-11.
- 1918 The striped cucumber beetle. *Diabrotica vittata* Fabr. Order Coleoptera; family Chrysomelidae. Conn. Agr. Expt. Sta. Bul. 203: 262-273, fig. 2.

Loyola, J. Terrazas (See Terrazas Loyola, J.)

Lucas, E. (See Holdaway and Lucas)

Lugger, O.

- 1888 On a case of faulty instinct in one of our tachina flies. Ent. Soc. Wash. Proc. (1885-1887) 1 (2): 50.
- 1897 Insects injurious in 1896. Minn. State Expt. Sta. Ann. Rpt. 1896 2: 3-43, figs. 1-23, pls. 1-7.
- 1899 Butterflies and moths injurious to our fruit-producing plants. Minn. State Expt. Sta., Ent. Ann. Rpt. 1898 4: i-xv, 1-280, figs. 1-237, pls. 1-24.

Luginbill, P. (See also Walton and Luginbill)

- 1918 The southern corn rootworm and farm practices to control it. U.S. Dept. Agr. Farmers' Bul. 950: 1-10, figs. 1-7.
- 1928 The fall army worm. U.S. Dept. Agr. Tech. Bul. 34: 1-91, figs. 1-49.
- 1950 Habits and control of the fall armyworm. U.S. Dept. Agr. Farmers' Bul. 1990: 1-11, figs. 1-12.

—— and Beyer, A. H.

- 1921 The corn earworm as an enemy of vetch. U.S. Dept. Agr. Farmers' Bul. 1206: 1-18, figs. 1-13.

Lundbeck, W. (See Henriksen and Lundbeck)

Lyons, L. A.

- 1957 Insects affecting seed production in red pine II. *Dioryctria disclusa* Heinrich, *D. abietella* (D. and S.), and *D. cambiicola* (Dyar) (Lepidoptera: Phycitidae). Canad. Ent. 89 (2): 70-79, figs. 1-3.
- 1964 The European pine sawfly, *Neodiprion sertifer* (Geoff.) (Hymenoptera: Diprionidae). A review

with emphasis on studies in Ontario. Ontario Ent. Soc. Proc. (1963) 94: 5-37, figs. 1-4.

MacCreary, D.

1947 Control of European corn borer. Del. Agr. Expt. Sta. Bul. 263: 26.

1948 European corn borer. Del. Agr. Expt. Sta. Bul. 271: 21-22.

1949 Control of the European corn borer by the use of natural enemies. Del. Agr. Expt. Sta. Bul. 276: 22.

— and Parker, W. L.

1948 Cucurbit insects. Del. Agr. Expt. Sta. Bul. 271: 20.

— and Rice, P. L.

1949 Parasites of the European corn borer in Delaware. Ent. Soc. Amer. Ann. 42 (2): 141-153, figs. 1-5.

MacDonald, D. R., and Webb, F. E.

1963 Insecticides and the spruce budworm. Canad. Ent. Soc. Mem. 31: 288-310, figs. 39.1-39.6.

MacDonald, J. E. (See McGugan, Haliburton, and MacDonald)

MacLeod, G. F. (See Michelbacher, MacLeod, and Smith)

MacNay, C. G.

1951 Forest and shade-tree insects. Canad. Insect Pest Rev. 29 (2): 123-129.

1952 Summary of important insect infestations, occurrences, and damage in Canada in 1952. Canad. Insect Pest Rev. 30 (10): 297-341.

1953 Summary of parasite and predator liberation in Canada in 1953. Canad. Insect Pest Rev. 31 (8): 307-317.

1954 Summary of parasite and predator liberations in Canada in 1954. Canad. Insect Pest Rev. 32 (8): 332-341.

1955 Summary of parasite and predator liberations in Canada in 1955. Canad. Insect Pest Rev. 33 (8): 321-330.

1956 Summary of parasite and predator liberations in Canada in 1956. Canad. Insect Pest Rev. 34 (8): 295-302.

1960 Some highlights of insect and mite occurrences in Canada, 1960. Canad. Insect Pest Rev. 38 (6): 219-222.

1960a Some highlights of insect and mite occurrences in Canada 1960. (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 10 (46): 1082-1084.

- 1961 Summary of important insect infestations, occurrences, and damage in agricultural areas of Canada in 1961. *Canad. Insect Pest Rev.* 39 (9): 269-296.
- MacPhee, A. W.
1962 [Winter moth in Nova Scotia in 1962]. *Canad. Insect Pest Rev.* 40 (5): 106.
- Madden, A. H. (See also Chamberlin, F. S., and Madden)
——— and Chamberlin, F. S.
1938 Biological studies of the tomato worm on tobacco in Florida, 1936 and 1937. *Jour. Econ. Ent.* 31: 703-706.
1945 Biology of the tobacco hornworm in the southern cigar-tobacco district. U.S. Dept. Agr. Tech. Bul. 896: 1-51, figs. 1-24.
- Malloch, J. R.
1919 The Diptera collected by the Canadian expedition, 1913-1918 (excluding the Tipulidae and Culiidae). Pp. 34c-90c, pls. 7-10. In Anderson, R. M., ed., Report of the Canadian Arctic Expedition 1913-18. Vol. 3: Insects, Pt. C: Diptera, 90 pp., 2 figs., 10 pls. Ottawa.
- Mally, C. W. (See Webster, F. M., and Mally)
Mally, F. W.
1891 The boll worm of cotton. A report of progress in a supplementary investigation of this insect. U.S. Dept. Agr., Div. Ent. Bul. 24: 1-50, figs. 1-2.
1893 Report on the boll worm of cotton (*Heliothis armiger* Hübn.). U.S. Dept. Agr., Div. Ent. Bul. 29: 1-73, pls. 1-2.
- Mangrum, J. F.
1942 The parasitic fly, *Zelia vertebrata* Say (Diptera, Dexiidae). *Ent. Soc. Amer. Ann.* 35 (1): 73-75, pl. 1.
- Manis, H. C., Gittins, A. R., Portman, R. W., Walz, A. J., Barr, W. F., Douglass, J. R., Evans, K. E., Smith, H. W., and Davis, E. G.
1956 Summary of insect conditions — 1955: Idaho. (U.S. Dept. Agr.) *Coop. Econ. Insect Rpt.* 6 (5): 78-82.
- Mann, J.
1969 Cactus-feeding insects and mites. U.S. Natl. Mus. Bul. 256: i-x, 1-158, pls. 1-8.
- Manson, G. F., Wressell, H. B., Begg, J. A., and Miller, L. A.
1960 Insects of the season 1959 in southwestern Ontario. *Canad. Insect Pest Rev.* 38 (1): 52-59.

- Marlatt, C. L.
 1889 A report on the lesser migratory locust. *Insect Life* (U.S. Dept. Agr., Div. Ent., Period. Bul.) 2 (3): 66-70.
 1898 The periodical cicada. U.S. Dept. Agr. Div. Ent. Bul., n. ser., 14: 1-148, 57 figs.
 1907 The periodical cicada. U.S. Dept. Agr. Bur. Ent. Bul., n. ser., 71: 1-181, 68 figs., 6 pls.
- Marsh, F. L.
 1937 Ecological observations upon the enemies of *Cecropia*, with particular reference to its hymenopterous parasites. *Ecology* 18 (1): 106-112, figs. 1-2.
 [1938] Biology of the tachinid *Winthemia datanae* Tns. *Psyche* (1937) 44 (4): 138-140.
 1941 A few life-history details of *Samia cecropia* within the southwestern limits of Chicago. *Ecology* 22 (3): 331-337, figs. 1-3.
- *Marsh, H. O. (See also Chittenden and Marsh)
 1913 Papers on insects affecting vegetable and truck crops. The striped beet caterpillar. U.S. Dept. Agr. Bur. Ent. Bul., n. ser., 127 (2): 13-18, figs. 4-5, pl. 5.
- Martin, D. F. (See Little and Martin)
- Martin, J. L.
 1958 Observations on the biology of certain tortricids in young coniferous plantations in southern Ontario. *Canad. Ent.* 90 (1): 44-53, figs. 1-2.
- Martineau, R. (See also Bracken and Martineau)
 1959 Jack-pine sawflies (*Neodiprion swainei* Midd. and *virginianus* complex). *Canad. Insect Pest Rev.* 37 (6): 229-230.
- Martorell, L. F.
 1945 A survey of the forest insects of Puerto Rico. Part I. (An annotated list of the insects affecting forest, shade and ornamental trees in Puerto Rico). *Puerto Rico Univ. Jour. Agr.* 29 (3): 69-354.
 1945a A survey of the forest insects of Puerto Rico. Part II. (A discussion of the most important insects affecting forest, shade and ornamental trees in Puerto Rico). *Puerto Rico Univ. Jour. Agr.* 29 (4): 355-608, figs. 1-18, pls. 1-21.
- Mathers, W. G. (See Wilkes, Coppel, and Mathers)

- Mathes, R. (See also Charpentier, McCormick, and Mathes; Hayslip, Mathes, and Bregger; Holloway and Mathes; Ingram, Bynum, and Mathes; Ingram et al.; Ingram et al.; Thames et al.)
- Ingram, J. W., Questel, D. D., Thames, W. H. (Jr.), and Dugas, A. L.
 1954 Current status of sugarcane-insect investigations in the United States. Internatl. Soc. Sugar Cane Technol. Cong. Proc. 8: 560-567.
- Mattson, W. J. (See Allen, D. C. et al.)
- Mau, R.
 1969 *Plautia stali* Scott. Hawaii. Ent. Soc. Proc. 1968 20 (2): 267.
- Maw, M. G. (See also Coppel and Maw)
 — and Coppel, H. C.
 1953 Studies on dipterous parasites of the spruce budworm, *Choristoneura fumiferana* (Clem.) (Lepidoptera: Tortricidae). II. *Phryxe pecosensis* (Tns.) (Diptera: Tachinidae). Canad. Jour. Zool. 31 (4): 392-403, figs. 1-18.
- Maxson, A. C.
 1920 Principal insect enemies of the sugar beet in the territories served by the Great Western Sugar Company. vii + 157 pp., 30 figs., 9 pls. Denver, Colo.
- Maxwell, C. W. (See Wood, G. W. et al.)
- Maybee, G. E.
 1958 Summary of parasite and predator liberations in Canada and of insect shipments from Canada in 1958. Canad. Insect Pest Rev. 36 (8): 300-313.
 1959 Summary of parasite and predator liberations in Canada and of insect shipments from Canada in 1959. Canad. Insect Pest Rev. 37 (8): 252-264.
- *McConnell, H. S. (See also Cory et al.; Cory et al.)
 1932 Oriental fruit moth parasites in Maryland. Jour. Econ. Ent. 25: 367-370.
- McConnell, W. R.
 1915 A unique type of insect injury. Jour. Econ. Ent 8: 261-267.
- McCormick, W. J. (See Charpentier, McCormick, and Mathes)
- McDonald, H.
 1947 Biology and control of *Heliothis ononis* Schiff., an important new pest of flax in western Canada.

- Ohio State Univ., Abs. Doctors' Diss.
(1946-1947) 53: 231-240, figs. 1-5.
- McDougall, G. A. (See Renault and McDougall)
- McDunnough, J.
1943 Two undescribed Canadian Microlepidoptera.
Canad. Ent. (1942) 74(12): 224.
- McGregor, M. D.
1967 Biology and natural enemies of an aspen leaf tier,
Sciaphila duplex, in the intermountain region.
Jour. Econ. Ent. 60: 1213-1216, figs. 1-3.
- McGuffin, W. C., and Barker, R. B.
1947 Prairie provinces forested area. Canada Dept. Agr.,
Forest Insect and Dis. Survey, Ann. Rpt.
1946: 54-65.
- McGugan, B. M., and Blais, J. R.
1959 Spruce budworm parasite studies in northwestern
Ontario. Canad. Ent. 91 (12): 758-783, figs.
1-3.
- Haliburton, W. H., and MacDonald, J. E.
1952 [Forest insect survey]. Province of Ontario. Canada
Dept. Agr., Forest Insect and Dis. Survey, Ann.
Rpt. 1951: 39-65.
- McKiel, J. A. (See Wood, G. W. et al.)
- McKinney, K. B.
1944 The cabbage looper as a pest of lettuce in the
Southwest. U.S. Dept. Agr. Tech. Bul. 846:
1-30, figs. 1-10.
- McLaine, L. S. (See also Tothill and McLaine)
1916 Rearing the parasites of the brown-tail moth in New
England for colonization in Canada. Agr. Gaz.
Canada 3(1): 22-25, figs. 1-5.
- *McLeod, J. H.
1951 Biological control investigations in British Columbia.
Ent. Soc. Brit. Columbia, Proc. (1950)
47: 27-36.
1954 Statuses of some introduced parasites and their
hosts in British Columbia. Ent. Soc. Brit. Colum-
bia, Proc. (1953) 50: 19-27.
- and Chant, D. A.
1952 Notes on the parasitism and food habits of the
European earwig, *Forficula auricularia* L.
(Dermaptera: Forficulidae). Canad. Ent. 84
(11): 343-345.

McLeod, J. M.

1964 Spruce needle miner parasites. *Canad. Insect Pest Rev.* 42 (2): 32-33.

1969 On the habits of a jack pine needle-miner, *Eucordylea canusella* (Lepidoptera: Gelechiidae), with special reference to its association with a fungus, *Aureobasidium pullulans* (Maniliales (Deuteromycetes) Dematiaceae). *Canad. Ent.* 101 (2): 166-179, figs. 1-12.

— and Daviault, L.

1963 Notes on the life history and habits of the spruce cone worm, *Dioryctria reniculella* (Grt.) (Lepidoptera: Pyralidae). *Canad. Ent.* 95 (3): 309-316, figs. 1-9.

McMillan, E. (See Seamans and McMillan)

Medler, J. T. (See also Yonke and Medler)

1961 A new record of parasitism of *Lygus lineolaris* (P. De. B.) (Hemiptera) by Tachinidae (Diptera). *Ent. Soc. Wash. Proc.* 63 (2): 101-102.

Melvin, J. C. E. (See also Turnock and Melvin; Wong, H. R., and Melvin)

— and Nairn, L. D.

1967 Parasites and predators of *Phyllophaga* spp. in southeastern Manitoba. *Canada Dept. Forestry and Rural Devlpmt., Bi-monthly Res. Notes* 23 (6): 43-44.

1967a Parasites et prédateurs de *Phyllophaga* dans le sud-est du Manitoba. *Canada Min. Forêts et Dévlpmt. Rural, Rev. Bimestrielle Recherches* 23 (4): 27.

[Menéndez Ramos, R.]

1924 Report of the Division of Entomology, fiscal year 1923-1924. *Porto Rico Insular Expt. Sta. (Rio Piedras), Ann. Rpt. 1923-1924:* 88-103.

Metcalf, Z. P.

1908 The army worm at Durham, North Carolina. *Jour. Econ. Ent.* 1: 354-355.

M'Gregor, E. A. (See Simmons, Reed, and M'Gregor)

Michelbacher, A. E.

1943 The present status of the alfalfa weevil in California. *Calif. Agr. Expt. Sta. Bul.* 677: 1-24, figs. 1-12.

— MacLeod, G. F., and Smith, R. F.

1943 Control of diabrotica, or western spotted cucumber

- beetle, in deciduous fruit orchards. Calif. Agr. Expt. Sta. Bul. 681: 1-34, figs. 1-20.
- Middlekauff, W. W., Bacon, O. G., and Swift, J. E.
1955 Controlling melon insects and spider mites. Calif. Agr. Expt. Sta. Bul. 749: 1-46, figs. 1-21, 4 figs.
- Micks, D. W.
1956 Laboratory rearing of the puss caterpillar, with notes on the incidence of parasitism. Jour. Econ. Ent. 49: 37-39, figs. 1-3.
- Middlekauff, W. W. (See Kelly and Middlekauff; Michelbacher et al.)
- Middleton, W.
1921 Leconte's sawfly, an enemy of young pines. Jour. Agr. Res. 20 (10): 741-760, figs. 1-3, pls. 88-92.
- Miller, C. A.
1963 Parasites and the spruce budworm. Canad. Ent. Soc. Mem. 31: 228-244, figs. 34.1-34.7.
1966 The black-headed budworm in eastern Canada. Canad. Ent. 98 (6): 592-613, figs. 1-12.
- Miller, G. C. (See Dugas, Miller, and Concienne)
- Miller, L. A. (See Manson et al.)
- *Milliken, F. B., and Wadley, F. M.
1923 *Phasia (Phorantha) occidentis* Walker, an internal parasite of the false chinch bug. Brooklyn Ent. Soc. Bul. 18 (1): 28-31.
- Milliken, R.
1893 Report on outbreaks of the western cricket and of certain locusts in Idaho. Insect Life (U.S. Dept. Agr., Div. Ent., Period. Bul.) 6 (1): 17-24.
- Milliron, [H. E.]
1956 A parasite (*Trichopoda pennipes*) — Delaware. (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 6 (34): 860.
- Mills, A. S. (See Leonard and Mills)
- Miskimen, G. W.
1962 Studies of the biological control of *Diatraea saccharalis* F. (Lepidoptera: Crambidae) on St. Croix, U.S. Virgin Islands. Puerto Rico Univ. Jour. Agr. 46 (2): 135-139.
- Mitchell, J. D. (See Hunter, Pratt, and Mitchell)
- Mitchell, [W. C.]
1964 *Trichopoda pennipes* var. *pilipes* Fabricius. Hawaii. Ent. Soc. Proc. 1963 18 (3): 345.

- 1965 Monarch butterfly (*Danaus plexippus* (L.)). (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 15 (21): 525.
- 1966 *Danaus plexippus* (L.). Hawaii. Ent. Soc. Proc. 1965 19 (2): 129.
- 1968 *Eumerus figurans* (Walker), subterranean termite and *Nezara viridula*. Hawaii. Ent. Soc. Proc. 1967 20 (1): 18.
- and Yoshioka, [E.]
- 1966 A tachina fly (*Trichopoda pennipes*). (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 16 (14): 286.
- Mitchener, A. V. (See Bird, R. D., and Mitchener)
- Miyahira, N. (See also Kim and Miyahira; Nakao and Miyahira; Yoshioka and Miyahira)
- 1965 *Trichopoda pennipes* Fabricius. Hawaii. Ent. Soc. Proc. 1964 19 (1): 28.
- 1966 Southern green stink bug (*Nezara viridula*). (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 16 (46): 1069.
- 1967 [A tachina fly (*Trichopoda pennipes* var. *pilipes*)]. (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 17 (22): 473.
- 1967a A tachina fly (*Lespesia archippivora*). (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 17 (49): 1048.
- 1968 [Southern green stink bug (*Nezara viridula*)]. (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 18 (23): 503
- 1968a A tachina fly (*Eucelatoria armigera*). (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 18 (25): 562.
- 1968b A tachina fly (*Trichopoda pennipes*). (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 18 (27): 624.
- 1969 Southern green stink bug (*Nezara viridula*). (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 19 (14): 241.
- and Ah Sam, [J.]
- 1968 Tachina flies (*Trichopoda pennipes* and *T. pennipes* var. *pilipes*). (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 18 (41): 966.
- 1969 [Southern green stink bug (*Nezara viridula*)]. (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 19 (6): 72.
- and Nakao, [H.]
- 1969 Monkeypod moth (*Polydesma umbricola*). (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 19 (24): 439.

Monteith, L. G.

- 1955 Host preferences of *Drino bohémica* Mesn. (Diptera: Tachinidae), with particular reference to olfactory responses. *Canad. Ent.* 87 (12): 509-530, figs. 1-3.
- 1960 Influence of plants other than the food plants of their host on host-finding by tachinid parasites. *Canad. Ent.* 92 (9): 641-652.
- 1962 Loss of attractiveness to *Bessa harveyi* (Tns.) (Diptera: Tachinidae) of host larvae during cold storage. *Canad. Ent.* 94 (8): 881-883.
- 1963 Daily selection by *Drino bohémica* Mesn. (Diptera: Tachinidae) of four species of hosts. *Canad. Ent.* 95 (2): 162-166, fig. 1.
- 1963a Habituation and associative learning in *Drino bohémica* Mesn. (Diptera: Tachinidae). *Canad. Ent.* 95 (4): 418-426.
- 1964 Influence of the health of the food plant of the host on host-finding by tachinid parasites. *Canad. Ent.* 96 (11): 1477-1482.
- 1966 Influence of new growth on the food plant of the host on host-finding by *Drino bohémica* Mesnil (Diptera: Tachinidae). *Canad. Ent.* 98 (11): 1205-1207.

Montgomery, B. E.

- 1933 Preliminary studies of insect parasites in Indiana. *Canad. Ent.* 65 (8): 185-190.

Moore, H. W. (See Handford, Painter, and Moore)

Morgan, C. V. G. (See Hopping, G. R., Leech, and Morgan)

Morgan, H. A.

- 1901 The differential grasshopper in the Mississippi Delta—other common species. U.S. Dept. Agr. Div. Ent. Bul., n. ser., 30: 7-33, 19 figs.

*Morrill, A. W.

- 1907 The Mexican conchuela in western Texas in 1905. U.S. Dept. Agr. Bur. Ent. Bul., n. ser., 64 (1): 1-14, 2 figs., 1 pl.
- 1910 Plant-bugs injurious to cotton bolls. U.S. Dept. Agr. Bur. Ent. Bul., n. ser., 86: 1-110, figs. 1-25, pls. 1-5.
- 1940 Living parasitic and predaceous insects for classroom use. *Jour. Econ. Ent.* 33: 123-128.

- Morrill, W. L.
1969 Parasitization of *Nabis alternatus* by *Leucostoma simplex* in eastern South Dakota. Ent. Soc. Amer. Ann. 62 (1): 240.
- Morris, B. R.
1874 Stray notes on Canadian Diptera. Canad. Ent. 6 (9): 176-178.
- Morrison, F. O.
1940 A revision of the American species of *Gonia* Meigen (Diptera: Tachinidae). Canad. Jour. Res. ser. D, 18 (9): 336-362, figs. 1-22a.
- Morse, W. J. (See Johannsen et al.)
- Mote, D. C. (See also Dimick and Mote; Schuh and Mote; Wilcox, Gray, and Mote)
1931 The introduction of the tachinid parasites of the European earwig in Oregon. Jour. Econ. Ent. 24: 948-956.
1931a European earwig (*Forficula auricularia* L.). Insect Pest Survey Bul. 11 (4): 230.
1933 [Emergence date of the second generation of *Digonichaeta setipennis* (Fallen) in Oregon]. Insect Pest Survey Bul. 13 (7): 259.
- Stearns, H. C., and Dimick, R. E.
1931 The biology of *Digonichaeta setipennis* Fall., a tachinid parasite of the European earwig, as observed primarily under western Oregon conditions. Jour. Econ. Ent. 24: 957-961.
- Motherwell, W. R.
1928 Report of the minister of agriculture for the Dominion of Canada for the year ended March 31, 1928. 141 pp. Ottawa.
- Mott, P. B.
1930 An annotated bibliography of the white pine weevil, *Pissodes strobi* (Peck), for white pine blister rust workers and others. N.J. Dept. Agr. Cir. 177: 1-37.
- Moznette, G. F.
1917 The rose flea-beetle (*Haltica probata* Fall). Jour. Ent. and Zool. 9 (1): 13-18, 1 pl.
- Muesebeck, C. F. W.
1936 New records of insects. Insect Pest Survey Bul. 16 (10): 535-536.
- Muir, F.
1923 Entomology. Hawaii. Sugar Planters' Assoc., Committee Charge Expt. Sta. Rpt. 1923: 10-18.

- and Swezey, O. H.
 1916 The cane borer beetle in Hawaii and its control by natural enemies. Hawaii. Sugar Planters' Assoc. Expt. Sta., Div. Ent. Bul. 13: 1-45, frontis., figs. 1-4, pls. 1-4.
 1925 Entomology. Hawaii. Sugar Planters' Assoc., Committee Charge Expt. Sta. Rpt. 1925: 12-23.
 1926 Entomology. Hawaii. Sugar Planters' Assoc., Committee Charge Expt. Sta. Rpt. 1926: 16-29.
- Muldrew, J. A. (See also Ives et al.; Ives et al.; Turnock and Muldrew)
 1956 Biological and natural control. Manitoba Ent. Soc. Proc. 12: 12-13.
- Munro, J. A.
 1939 Colorado potato beetle (*Leptinotarsa decemlineata* Say). Insect Pest Survey Bul. 19 (9): 563.
 1951 Drafting insect "soldiers" to war on plant parasites. How parasites of the sweet clover weevil and European corn borer are being established in North Dakota. N. Dak. Agr. Expt. Sta. Bi-monthly Bul. 13 (6): 231-236, figs. 1-3.
- Post, R. L., and Nelson, D. B.
 1952 2. Introduction of insect parasites for European corn borer control. N. Dak. Agr. Expt. Sta. Bi-monthly Bul. 14 (4): 128.
- and Telford, H. S.
 1943 Spraying and dusting potatoes. Part I. To control potato insects. N. Dak. Agr. Expt. Sta. Bul. 319: 1-13, figs. 1-8.
- Munson, [R. E.]
 1966 A tachina fly (*Lespesia archippivora*) — Missouri. (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 16 (41): 979.
- Murtfeldt, M. E.
 1892 Notes from Missouri. Insect Life (U.S. Dept. Agr., Div. Ent., Period. Bul.) 5 (2): 135-136.
 1894 Notes on the insects of Missouri for 1893. U.S. Dept. Agr., Div. Ent. Bul. 32: 37-45.
 1894a Acorn insects, primary and secondary. Insect Life (U.S. Dept. Agr., Div. Ent., Period. Bul.) 6 (4): 318-324.
- Mutchler, A. J., and Weiss, H. B.
 1925 Beetles of the genus *Conotrachelus* known to occur

- in New Jersey. N.J. Dept. Agr. Cir. 87: 1-22, pls. 1-6.
- Myers, J. G.
 1935 Biological control of sugar-cane pests in the British West Indies and British Guiana. Internatl. Soc. Sugar Cane Technol. Cong. Proc. 5: 381-385.
- Nairn, L. D. (See also Ives et al.; Melvin and Nairn)
 ——— Turnock, W. J., Ives, W. G. H., and Buckner, C. H.
 1961 Investigations of the population dynamics of the larch sawfly in Manitoba. Manitoba Ent. Soc. Proc. 17: 31-36.
- Nakao, H. (See also Miyahira and Nakao)
 1963 A parasitic braconid (*Apanteles militaris*). (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 13 (20): 535.
 1963a A parasitic tachina fly (*Eucelatoria armigera*). (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 13 (20): 535.
 1964 *Eucelatoria armigera* (Coquillett) [sic]. Hawaii. Ent. Soc. Proc. 1963 18 (3): 333.
- and Miyahira, [N.]
 1967 [Southern green stink bug (*Nezara viridula*)]. (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 17 (17): 345.
- Nash, R. W.
 1939 The yellow-headed spruce sawfly in Maine. Jour. Econ. Ent. 32: 330-334, fig. 1.
- Neff, S. E., and Eisner, T.
 1960 Note on two tachinid parasites of the walking stick, *Anisomorpha buprestoides* (Stoll). Brooklyn Ent. Soc. Bul. 55 (4): 101-103, figs. 1-4.
- Neilson, C. L. (See also Handford and Neilson)
 1954 Report of provincial entomologist. Brit. Columbia Dept. Agr. Ann. Rpt. 48: CC67-CC73.
- Neilson, W. T. A. (See Wood, G. W., and Neilson; Wood, G. W. et al.)
- *Neiswander, C. R. (See also Huber and Neiswander)
 1941 *Coryphista meadii* (Packard), a new pest of Japanese barberry. Jour. Econ. Ent. 34: 386-389, figs. 1-4.
- Neiswander, R. B.
 1944 Insect pests of strawberries in Ohio. Ohio Agr. Expt. Sta. Bul. 651: 1-37, figs. 1-15.
- Nelson, D. B. (See Munro, Post, and Nelson)
- Nettles, W. C. (See Eddy, C. O., and Nettles)

- Neunzig, H. H., and Gyrisco, G. G.
 1955 Some insects injurious to birdsfoot trefoil in New York. Jour. Econ. Ent. 48: 447-450.
- Newcomer, E. J. (See also Carnes and Newcomer)
 1912 Some observations on the relations of ants and lycaenid caterpillars, and a description of the relational organs of the latter. N.Y. Ent. Soc. Jour. 20 (4): 31-36, pls. 2-3.
 1958 Some parasites and predators of fruit pests in the Pacific Northwest. Pan-Pacific Ent. 34 (2): 87-91.
- and Carlson, F. W.
 1953 The leaf roller moth *Pandemis pyrusana*. Jour. Econ. Ent. (1952) 45: 1079-1081.
- Newton, R. C.
 1954 New records of dipterous parasites of grasshoppers. Jour. Econ. Ent. 47: 935-936.
- Nichols, J. O.
 1962 The gypsy moth in Pennsylvania — its history and eradication. Pa. Dept. Agr. Misc. Bul. 4404: i-ii, 1-82, figs. 1-39.
- Nickels, C. B.
 1951 Notes on the life history and habits of the pecan nursery casebearer in Texas. Jour. Econ. Ent. 44: 433-434.
- Pierce, W. C., and Pinkney, C. C.
 1950 Parasites of the pecan nut casebearer in Texas. U.S. Dept. Agri. Tech. Bul. 1011: 1-21, figs. 1-7.
- Noble, W. B.
 1932 Sod webworms and their control in lawns and golf greens. U.S. Dept. Agr., Dept. Cir. 248: 1-4.
- Nougaret, R. L.
 1919 The achemon sphinx moth. (*Pholus achemon* Drury.) A pest of the grape in California. Calif. Dept. Agr. Monthly Bul. 8 (10): 560-584, figs. 146-162.
- Nutting, W. L.
 1953 The biology of *Euphasiopteryx brevicornis* (Townsend) (Diptera, Tachinidae), parasitic in the cone-headed grasshoppers (Orthoptera, Copiphorinae). Psyche 60 (2): 69-81, figs. 1-3, pl. 4.
- Oatman, E. R.
 1966 An ecological study of cabbage looper and imported

- cabbageworm populations on cruciferous crops in southern California. Jour. Econ. Ent. 59: 1134-1139, figs. 1-2.
- O'Brien, C. W. (See Ashlock and O'Brien)
- O'Kane, W. C.
 1916 Plan and progress of work, 1915-1916. [Division of Insect Suppression]. N.H. Dept. Agr. Rpt. (1914-1916) 1916: 123-144, pls. 1-8.
- and Conklin, J. C. [sic]
 1942 Sawflies, gipsy moths, and apple maggots. N.H. Agr. Expt. Sta. Bul. 345: 53-54.
- Oliver, A. D. (Jr.) (See also Vogel and Oliver)
 1964 Studies on the biological control of the fall webworm, *Hyphantria cunea*, in Louisiana. Jour. Econ. Ent. 57: 314-318, figs. 1-2.
- Ollieu, M. M., and Schenk, J. A.
 1966 The biology of *Eucosma rescissoriana* Heinrich in western white pine in Idaho (Lepidoptera: Olethreutidae). Canad. Ent. 98 (3): 268-274, figs. 1-6.
- Olson, R. E. (See Soper and Olson)
- Orr, L. W. (See Knowlton et al.)
- Ortega C., A. (See Carrillo S., Ortega C., and Gibson)
- *Osborn, H.
 1884 The ash sawfly (*Selandria barda* Say). Canad. Ent. 16 (8): 148-152, fig. 6.
 1891 Report of a trip to Kansas to investigate reported damages from grasshoppers. Ontario Ent. Soc. Ann. Rpt. 22: 69-74.
 1891a Report of a trip to Kansas to investigate reported damages from grasshoppers. Insect Life (U.S. Dept. Agr., Div. Ent., Period. Bul.) 4 (1-2): 49-56.
- Osborn, H. T.
 1938 The introduction into Hawaii from Mexico of insect parasites to control armyworms (1923-1924). Hawaii. Planters' Rec. 42 (2): 147-152, figs. 1-5, 1 fig.
- *Osborne [sic], H. (See *Osborn, H.)
- Osten Sacken, [C. R.]
 1875 Parasitic Diptera. Canad. Ent. 7 (4): 72.
 1877 Tachina parasite on Phasmidae. Psyche 2 (35-36): 23.
 1887 Some North American Tachinae. Canad. Ent. 19 (9): 161-166.

Osterberger, B. A. (See Hinds and Osterberger)

*Packard, A. S., Jr. (See also Riley, Packard, and Thomas).

- 1869 Guide to the study of insects, and a treatise on those injurious and beneficial to crops: for the use of colleges, farm-schools, and agriculturists. ix + 702 pp., 652 figs., 11 pls. Salem, Mass.
- 1873 Half-hour recreations in natural history, division first. Half-hours with insects. Twelve parts. Part 2. Insects of the garden, their habits, &c. Pp. 33-64, figs. 22-53. Boston.
- 1890 Fifth report of the United States Entomological Commission, being a revised and enlarged edition of bulletin number 7, on insects injurious to forest and shade trees. V. 5, viii + 957 pp., 306 figs., 40 pls. Washington.
- 1914 Monograph of the bombycine moths of North America, including their transformations and origin of the larval markings and armature. Part III. Families Ceratocampidae (exclusive of Ceratocampinae), Saturniidae, Hemileucidae, and Brahmaeidae. Natl. Acad. Sci. Mem. 12 (1): i-ix, 1-516, figs. 1-34, pls. 1-113.

[*Packard, A. S., Jr., and Putnam, F. W., eds.]

- 1875 A tachina parasite of the squash bug. Amer. Nat. 9 (9): 519.

Packard, C. M. (See also Walton and Packard)

- 1929 *Ateloglossa cinerea* Coq., a parasite of *Melanotus* sp. Jour. Econ. Ent. 22: 598.

Painter, Reginald H. (See also Winburn and Painter)

- 1955 Insects on corn and teosinte in Guatemala. Jour. Econ. Ent. 48: 36-42.

Painter, Richard H. (See also Handford, Painter, and Moore)

- 1929 The tarnished plant bug *Lygus pratensis* L.: a progress report. Ontario Ent. Soc. Ann. Rpt. 60: 102-107.

Panton, J. H.

- 1897 Two insect pests of 1896. Ontario Ent. Soc. Ann. Rpt. 27: 44-54, figs. 45-55.

Paquet, G.

- 1945 Liberation of parasites. Quebec Min. Lands and Forests Rpt. 1943-1944: 93-94.

Paradis, R.-O.

- 1961 Essai d'analyse des facteurs de mortalité chez *Archips argyrospilus* (Wlk.) (Lepidoptères:

- Tortricidae). Soc. Ent. du Québec, Ann. (1960) 6: 59-69.
- and LeRoux, E. J.
 1965 Recherches sur la biologie et la dynamique des populations naturelles d'*Archips argyrospilus* (Wlk.) (Lépidoptères: Tortricidae) dans le sud-ouest du Québec. Soc. Ent. du Canada Mém. 43: 1-77, figs. 1-22.
- Parish, H. E.
 1934 Biology of *Euschistus variolarius* P. De B. (Family Pentatomidae; Order Hemiptera). Ent. Soc. Amer. Ann. 27(1): 50-54.
- *Parker, H. L.
 1916 Rearing of *Winthemia quadripustulata* from rhyngophorous larva (Dip., Col.). Ent. News 27 (5): 236.
- Parker, J. R., Strand, A. L., and Seamans, H. L.
 1921 Pale western cutworm (*Porosagrotis orthogonia* Morr.). Jour. Agr. Res. 22 (6): 289-322, pls. C, 28-30.
- Parker, L. B. (See Gardner and Parker; King, J. L., and Parker)
 Parker, W. L. (See MacCreary and Parker)
- Parks, [T. H.]
 1953 European corn borer (*Pyrausta nubilalis*) — Ohio. (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 3 (37): 670.
- and Goleman, [D. L.]
 1954 European corn borer (*Pyrausta nubilalis*) — Ohio. (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 4 (24): 500.
- Parrott, P. J., and Schoene, W. J.
 1912 The apple and cherry ermine moths. N.Y. (Geneva) Agr. Expt. Sta. Tech. Bul. 24: 1-40, figs. 1-10, pls. 1-9.
- Patana, R. (See Bryan, D. E., Jackson, and Patana; Jackson, C. G., Bryan, and Patana)
- Patch, E. M. (See also Johannsen et al.)
 1906 Insect notes for 1906. Maine Agr. Expt. Sta. Bul. 134: 209-228, figs. 16-23.
 1907 Insect notes for 1907. Maine Agr. Expt. Sta. Bul. 148: 258-285, figs. 34-44.
- Patterson, J. E.
 1929 The pandora moth, a periodic pest of western pine forests. U.S. Dept. Agr. Tech. Bul. 137: 1-19, figs. 1-8.

Patton, C. N.

- 1957 Notes on the biology of *Sturmia incompta* (van der Wulp) and *Winthemia citheroniae* Sabrosky (Diptera: Larvaevoridae). Fla. Ent. 40 (2): 69-70.
- 1958 A catalogue of the Larvaevoridae of Florida. Fla. Ent. 41 (1): 29-39; (2): 77-89.

Paul, L. C., and Putnam, L. G.

- 1960 Morphometrics, parasites, and predators of migrant *Melanoplus bilituratus* (Wlk.) (Orthoptera: Acrididae) in Saskatchewan in 1940. Canad. Ent. 92 (7): 488-493, fig. 1.

Paxton, E. E., Waldron, J. W., Bottomley, A. W. T., Dowsett, J. M., and Gartley, A.

- 1911 Report of the Experiment Station Committee. Hawaii. Planters' Rec. 5 (6): 322-325.

Payne, H. G. (See also Brittain and Payne)

- [1918] The white marked tussock moth, (*Hemerocampa leucostigma* Smith & Abbot). Ent. Soc. Nova Scotia, Proc. (1917) 3: 62-68, pl. 13.

P[emberton], C. E.

- 1923 Notes on armyworms and cutworms. Hawaii. Planters' Rec. 27 (4): 257-258, 2 figs.
- 1924 Rat control and other projects in Hamakua. Hawaii. Sugar Planters' Assoc. Committee Charge Expt. Sta. Rpt. 1924: 25-32.
- 1925 A study of the cane borer, *R. obscura*, and its parasite, *C. sphenophori*, at Paauhau Sugar Plantation Company. Hawaii. Planters' Rec. 29 (2): 174-185.
- 1932 The present status of sugar cane insects in Hawaii. Assoc. Hawaii. Sugar Technol. Rpt. 11: 47-49.
- 1932a Entomology. Hawaii. Sugar Planters' Assoc. Committee Charge Expt. Sta. Ann. Rpt. 1932: 18-22.
- 1933 Entomology. Hawaii. Sugar Planters' Assoc. Committee Charge Expt. Sta. Ann. Rpt. 1933: 14-19.
- 1934 Entomology. Hawaii. Sugar Planters' Assoc. Committee Charge Expt. Sta. Ann. Rpt. 1934: 19-26.
- 1936 Entomology. Hawaii. Sugar Planters' Assoc. Committee Charge Expt. Sta. Ann. Rpt. 1936: 20-27.
- 1938 Entomology. Hawaii. Sugar Planters' Assoc. Committee Charge Expt. Sta. Rpt. 1938: 19-29.
- 1939 Entomology. Hawaii. Sugar Planters' Assoc. Committee Charge Expt. Sta. Rpt. 1939: 19-27.
- 1940 Entomology. Hawaii. Sugar Planters' Assoc. Committee Charge Expt. Sta. Rpt. 1940: 17-21.

- 1941 Entomology. Hawaii. Sugar Planters' Assoc. Committee Charge Expt. Sta. Rpt. 1941: 21-27.
- 1942 Entomology. Hawaii. Sugar Planters' Assoc. Committee Charge Expt. Sta. Rpt. 1942: 18-22.
- 1943 Entomology. Hawaii. Sugar Planters' Assoc. Committee Charge Expt. Sta. Rpt. 1943: 14-18.
- 1944 Entomology. Hawaii. Sugar Planters' Assoc. Committee Charge Expt. Sta. Rpt. 1944: 17-21.
- 1945 *Eucelatoria armigera* (Coq.). Hawaii. Ent. Soc. Proc. 1944 12 (2): 220.
- 1948 *Eucelatoria armigera* (Coq.). Hawaii. Ent. Soc. Proc. 1947 13 (2): 209.
- 1948a History of the entomology department Experiment Station, H.S.P.A. 1904-1945. Hawaii. Planters' Rec. 52(1): 53-90, figs. 1-6.
- 1948b The control of the grass armyworm, *Laphygma exempta* (Walker), in Hawaii by parasites. Hawaii. Planters' Rec. 52 (3-4): 181-200, figs. 1-21.
- 1951 The present status of the insect pests of sugar cane in Hawaii and the prospect of new invasions. Internatl. Soc. Sugar Cane Technol. Cong. Proc. 7: 401-404.
- and Williams, F. X.
- 1938 Some insect and other animal pests in Hawaii not under satisfactory biological control. Hawaii. Planters' Rec. 42 (3): 211-230.
- Pengelly, D. H.
- 1961 *Thymelicus lineola* (Ochs.) (Lepidoptera: Hesperidae) a pest of hay and pasture grasses in southern Ontario. Ontario Ent. Soc. Ann. Rpt. 91: 189-197, figs. 1-8.
- [Pepler, W. A.]
- 1938 Report of the sub-committee on entomology. Forestry Chron. 14 (1): 15-18.
- Perez, J. Fernández (See Scaramuzza, Fernández Artiles, and Fernández Perez)
- *Perkins, B. D. (Jr.)
- 1963 A parasitic tachina fly (*Trichopoda pennipes* F.). (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 13 (51): 1416.
- 1964 *Coleotichus blackburniae* White. Hawaii. Ent. Soc. Proc. 1963 18 (3): 351.

Perkins, [R. C. L.]

1907 [Parasitism of young caterpillars of *Sphinx convulvi*]. Hawaii. Ent. Soc. Proc. 1 (4): 114.

1910 Work on borer. Pp. 292-295. In Eckart, C. F., Report of director of the Experiment Station. Hawaii. Planters' Rec. 3 (6): 292-317.

Petch, C. E.

1925 [White grub parasites at Hemmingford, P. Q.]. Canad. Insect Pest Rev. 3 (1): 6.

—— and Hammond, G. H.

1925 Notes on insect parasites of *Phyllophaga anxia* in the Province of Quebec. Ontario Ent. Soc. Ann. Rpt. 55: 24-28.

1926 Parasites of white grubs in southern Quebec. A progress report. Ontario Ent. Soc. Ann. Rpt. 56: 85-91.

Peters, L. L. (See Jackson and Peters)

*Peterson, A.

1946 Laboratory tests showing the effect of DDT on several important parasitic insects. Ohio Jour. Sci. 46 (6): 323-326.

Peterson, L. O. T.

1948 Some aspects of poplar borer, *Saperda calcarata* Say, (Cerambycidae) infestations under parkbelt conditions. Ontario Ent. Soc. Ann. Rpt. 78: 56-61.

1958 The boxelder twig borer, *Proteoteras willingana* (Kearfott), (Lepidoptera: Olethreutidae). Canad. Ent. 90 (11): 639-646, figs. 10-12.

Petty, [H. B., Jr.]

1967 European corn borer (*Ostrinia nubilalis*) — Illinois. (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 17 (3): 29.

Pfadt, R. E. (See Lavigne and Pfadt)

Philbrook, E. E.

1918 Report of special field agent, gypsy moth work. Maine Comnr. Agr. Ann. Rpt. 1917 16: 57-61, 3 pls.

Philip, C. B.

1931 The Tabanidae (horseflies) of Minnesota with special reference to their biologies and taxonomy. Minn. Univ. Agr. Expt. Sta. Tech. Bul. 80: 1-132, figs. 1-3, pls. 1-4.

Phillips, A. M.

1957 Control of hickory shuckworm on pecans. Fla. Agr. Expt. Sta. Ann. Rpt. 1957: 89.

Phillips, M. E. (See Hamlin, Reed, and Phillips)

*Phillips, W. J., and Barber, G. W.

- 1931 The corn earworm as an enemy of field corn in the eastern states. U.S. Dept. Agr. Farmers' Bul. 1651: i-ii, 1-18, figs. 1-17.

——— and Fox, H.

- 1924 The rough-headed corn stalk-beetle. U.S. Dept. Agr. Bul. 1267: 1-34, figs. 1-16, pls. 1-4.

Phipps, C. R.

- 1927 The black army cutworm. A blueberry pest. Maine Agr. Expt. Sta. Bul. 340: 201-216, figs. 29-30.

Pickwell, G. (See Duncan and Pickwell)

Pierce, H. D. (See Clancy and Pierce)

Pierce, W. C. (See Nickels, Pierce, and Pinkney)

Pierce, W. D. (See also Hunter and Pierce)

- 1907 Contributions to the knowledge of Rhynchophora. Ent. News 18 (8): 356-363.

- 1908 A list of parasites known to attack American Rhynchophora. Jour. Econ. Ent. 1: 380-396.

- 1908a Studies of parasites of the cotton boll weevil. U.S. Dept. Agr. Bur. Ent. Bul., n. ser., 73: 1-63, figs. 1-6, pls. 1-3.

- 1910 On some phases of parasitism displayed by insect enemies of weevils. Jour. Econ. Ent. 3: 451-458.

——— Cushman, R. A., and Hood, C. E.

- 1912 The insect enemies of the cotton boll weevil. U.S. Dept. Agr. Bur. Ent. Bul., n. ser., 100: 1-99, figs. 1-26, pls. 1-3.

Pimentel, D.

- 1961 The natural control of caterpillar populations on cole crops. Jour. Econ. Ent. 54: 889-892, figs. 1-6.

Pinkney, C. C. (See Nickels, Pierce, and Pinkney)

Plank, H. K.

- 1927 Cuban parasites of the moth stalkborer. Internatl. Soc. Sugar Cane Technol. Cong. Proc. 2: 68-69.

- 1929 Algunas observaciones acerca del control del "borer" de la caña de azúcar. Asoc. de Téc. Azucareros Cuba, Mem. de la Conf. An. 3: 114-119.

- 1929a Some observations on the control of the sugar cane moth stalkborer in Cuba. Asoc. de Téc. Azucareros Cuba, Proc. 3: 106-110.

- 1929b Natural enemies of the sugar cane moth stalkborer in Cuba. Ent. Soc. Amer. Ann. 22 (4): 621-640, figs. 1-7.
- 1932 The seasonal development of *Diatraea saccharalis* Fabr. and its principal parasites in Cuba. Internatl. Soc. Sugar Cane Technol. Cong. Proc. 4, Bul. 113: 1-8.
- 1945 Insect pests of food crops. Puerto Rico Fed. Expt. Sta. (Mayaguez) Rpt. 1944: 26-27.
- 1945a Algunas observaciones acerca del control del "borer" de la cana de azucar. Asoc. de Téc. Azucareros Cuba, Bol. Ofic. 4 (9): 351-356.
- 1950 Establishment of introduced beneficial insects. Puerto Rico Fed. Expt. Sta. (Mayaguez) Rpt. 1950: 13-16.
- 1951 Establishment of introduced beneficial insects. Puerto Rico Fed. Expt. Sta. (Mayaguez) Rpt. 1951: 15-19.
- Pond, D. D.
- 1955 Insects of the season in New Brunswick, 1954. Canad. Insect Pest Rev. 33 (1): 105-108.
- 1960 Life history studies of the armyworm, *Pseudaletia unipuncta* (Lepidoptera: Noctuidae), in New Brunswick. Ent. Soc. Amer. Ann. 53 (5): 661-665, fig. 1.
- Poos, F. W.
- 1927 Biology of the European corn borer (*Pyrausta nubilalis* Hubn.) and two closely related species in northern Ohio. Ohio Jour. Sci. 27 (2): 47-94, pls. 1-6.
- 1928 An annotated list of some parasitic insects. Ent. Soc. Wash. Proc. 30 (8): 145-150.
- and Hetrick, L. A.
- 1945 *Tetralopha scortealis* (Led.) a new insect pest of lespedeza. Jour. Econ. Ent. 38: 312-315, figs. 1-2.
- Popenoe, C. H.
- 1909 The Colorado potato beetle in Virginia in 1908. U.S. Dept. Agr. Bur. Ent. Bul, n. ser., 82 (1): 1-8, pls. 1-2.
- Porter, B. A.
- 1924 The bud moth. U.S. Dept. Agr., Dept. Bul. 1273: 1-20, pls. 1-2.

—— and Garman, P.

- 1923 The apple and thorn skeletonizer. Conn. (New Haven) Agr. Expt. Sta. Bul. 246: 245-264, figs. 1-3, pls. 1-4.

Portman, R. W. (See Manis et al.)

Post, R. L. (See Munro, Post, and Nelson)

Pouillaude, I.

- 1913 Les parasites de "*Porthesia dispar*" et de "*Euproctis chrysorrhoea*" aux Etats-Unis en 1912. Soc. Sci. et Méd. de l'Ouest (Rennes), Bul. 22 (3): 81-86.

Powell, J. A.

- 1958 [*Achaetoneura* reared from *Antheraea (Telea) polyphemus* Cramer]. Pan-Pacific Ent. 34 (1): 48.
- 1962 Host-parasite relationships of California Tortricinae (Lepidoptera: Tortricidae). Pan-Pacific Ent. 38 (2): 131-140.

Pratt, F. C. (See Hunter, Pratt, and Mitchell; Titus and Pratt)

Prebble, M. L.

- 1935 *Actia diffidens* Curran, a parasite of *Peronea variana* (Fernald) in Cape Breton, Nova Scotia. Canad. Jour. Res. 12 (2): 216-227, figs. 1-31.
- 1942 Forest insects of the season 1941 in British Columbia. Canad. Insect Pest Rev. 20 (1): 108-111.
- 1942a [*Uramyia halisidotae* (Townsend) parasite of *Halisidota argentata* Packard]. Canad. Insect Pest Rev. 20 (2): 165.
- 1945 Forest insects of the season 1944 in Vancouver Island and western British Columbia. Canad. Insect Pest Rev. 23 (1): 110-112.

Prentice, R. M. (See also Ives and Prentice)

- 1955 The life history and some aspects of the ecology of the large aspen tortrix, *Choristoneura conflictana* (Wlkr.) (N. Comb.) (Lepidoptera: Tortricidae). Canad. Ent. 87 (11): 461-473, figs. 1-18.

Price, R. G. (See Bottrell et al.)

Prine, J. E.

- 1967 *Leucostoma gravipes* van der Wulp from *Oncopeltus fasciatus* Dallas. Pan-Pacific Ent. 43 (1): 92.

Procter, W.

- 1938 The insect fauna with references to methods of capture, food plants, the flora and other biological features. Pt. VI, 496 pp., 12 figs. In his, ed., Biological survey of the Mount Desert region. Philadelphia.

- 1946 The insect fauna with references to methods of capture, food plants, the flora and other biological features. Pt. VII, 566 pp., 10 figs. *In his*, ed., Biological survey of the Mount Desert region. Philadelphia.
- Prokopy, R. J.
1968 Parasites of the leaf rollers *Archips argyrospilus* and *A. griseus* in Connecticut. *Jour. Econ. Ent.* 61: 348-352.
- Proper, A. B.
1934 Hyperparasitism in the case of some introduced lepidopterous tree defoliators. *Jour. Agr. Res.* 48 (4): 359-376.
- [Provancher, L.]
1869 La Piéride de la rave, *Pieris rapae* Schrank. *Nat. Canad.* 2 (1): 13-18, figs. 3-4.
- Putman, W. L.
1935 Notes on the native hosts of some oriental fruit moth parasites. *Canad. Ent.* 67 (3): 46-49.
1935a Notes on the hosts and parasites of some lepidopterous larvae. *Canad. Ent.* 67 (5): 105-109.
1943 Host plants and parasites of some lepidopterous larvae. *Canad. Ent.* (1942) 74 (12): 219-224.
- Putnam, F. W. (See Packard, A. S., and Putnam, eds.)
Putnam, L. G. (See Paul and Putnam)
- Puttarudriah, M.
1953 The natural control of the alfalfa looper in central California. *Jour. Econ. Ent.* 46: 723.
- Puttler, B.
1966 Notes on two parasites attacking a *Lema* sp. (Coleoptera: Chrysomelidae). *Jour. Econ. Ent.* 59: 475-476.
- Quaintance, A. L.
1908 The codling moth or apple worm. (U.S. Dept. Agr.) *Yearb. Agr.* 1907: 435-450, pls. 52-55.
1908a The apple-tree tent caterpillar (*Malacosoma americana* Fab.). U.S. Dept. Agr., Dept. Cir. 98: 1-8, figs. 1-4.
1915 The apple-tree tent caterpillar. U.S. Dept. Agr. *Farmers' Bul.* 662: 1-10, figs. 1-7.
- and Brues, C. T.
1905 The cotton bollworm. U.S. Dept. Agr. *Bur. Ent. Bul.*, n. ser., 50: 1-155, figs. 1-27, pls. 1-25.

- and Jenne, E. L.
 1912 The plum curculio. U.S. Dept. Agr. Bur. Ent. Bul.,
 n. ser., 103: 1-250, figs. 1-36, pls. 1-20.
- Questel, D. D. (See Mathes et al.; Thames et al.)
- Quezada, J. R.
 1967 Notes on the biology of *Rothschildia ?aroma*
 (Lepidoptera: Saturniidae), with special refer-
 ence to its control by pupal parasites in El
 Salvador. Ent. Soc. Amer. Ann. 60 (3): 595-599,
 figs. 1-3.
- Raine, J.
 1966 Life history of *Dasystoma salicellum* Hbn. (Lepidop-
 tera: Oecophoridae), a new pest of blueberries in
 British Columbia. Canad. Ent. 98
 (31): 331-334, figs. 1-11.
- Raizenne, H.
 1952 Forest Lepidoptera of southern Ontario and their
 parasites received and reared at the Ottawa Forest
 Insect Survey Laboratory from 1937 to 1948. v
 + 277 pp. Canada Dept. Agr., Sci. Serv., Div.
 Forest Biol.
 1957 Forest sawflies of southern Ontario and their
 parasites. (Received and reared at the Ottawa
 Laboratory from 1936 to 1952 inclu-
 sive). Canada Dept. Agr. Pub. 1009: 1-45.
- Ramos, R. Menéndez (See Menéndez Ramos, R.)
- Randolph, N. M. (See Teetes and Randolph)
- Rands, R. D. (See Holloway and Rands)
- Raun, E. S. (See Sparks, Raun, and Carter)
- Rauschenberger, J. L. (See Ciesla, Rauschenberger, and Bell)
- Rawlings, G. B.
 1960 Fungi and insects associated with *Pinus radiata*. Pp.
 250-320. In Scott, C. W., *Pinus radiata*. FAO
 Forestry and Forest Prod. Studies 14: i-xii,
 1-328, frontis.
- Reed, E. B.
 1872 Insects injurious to the potato. Attacking the foliage
 only. Prov. Ontario Ent. Soc. Rpt. 1871: 65-81,
 figs. 65-83.
 1873 Insects infesting maple trees. Ontario Ent. Soc. Ann.
 Rpt. 1872: 35-43, figs. 26-33.
 1883 Diptera. Two-winged flies. Ontario Ent. Soc. Rpt.
 1882: 45-53, figs. 17-54.
- Reed, W. D. (See Hamlin, Reed, and Phillips; Simmons, Reed, and
 M'Gregor)

- Reeks, W. A. (See also Finlayson, L. R., and Reeks)
- 1938 Native insect parasites and predators attacking *Diprion polytomum* (Hartig) in Canada. Ontario Ent. Soc. Ann. Rpt. 69: 25-28.
- 1943 [*Compsilura* sp. parasite of *Malacosoma disstria* Hbn.]. Canad. Insect Pest Rev. 21 (3): 213.
- 1946 Forest insects of the season 1945 in the maritime provinces and Newfoundland. Canad. Insect Pest Rev. 24 (1): 25-29.
- 1951 Birch. Birch casebearer (*Coleophora salmani* Heinr.). Canad. Insect Pest Rev. 29(5): 212-213.
- 1953 The establishment of introduced parasites of the European spruce sawfly (*Diprion hercyniae* (Htg.)) (Hymenoptera: Diprionidae) in the maritime provinces. Canad. Jour. Agr. Sci. 33 (5): 405-429, figs. 1-3.
- 1954 An outbreak of the larch sawfly (*Pristiphora erichsonii* (Htg.)) in the maritime provinces (Hymenoptera: Tenthredinidae) and the role of parasites in its control. Canad. Ent. 86 (10): 471-480, figs. 1-2.
- 1964 European spruce sawfly (*Diprion hercyniae* (Hartig)). Canad. Insect Pest Rev. 42 (2): 29-32.
- Forbes, R. S., and Cuming, F. G.
- 1947 Maritime provinces and Newfoundland. Canada Dept. Agr., Forest Insect and Dis. Survey, Ann. Rpt. 1946: 7-19.
- 1948 Maritime provinces and Newfoundland. Canada Dept. Agr., Forest Insect and Dis. Survey, Ann. Rpt. 1947: 9-23.
- 1949 Maritime provinces and Newfoundland. Canada Dept. Agr., Forest Insect and Dis. Survey, Ann. Rpt. 1948: 9-26.
- and Smith, C. C.
- 1956 The satin moth *Stilpnotia salicis* (L.), in the maritime provinces and observations on its control by parasites and spraying. Canad. Ent. 88 (10): 565-579.
- Smith, C. C., and Forbes, R. S.
- 1946 Maritime provinces and Newfoundland. Canada Dept. Agr., Forest Insect and Dis. Survey, Ann. Rpt. 1945: 7-19.
- Regan, W. S.
- 1923 The fruit-tree leaf-roller in the Bitter Root Valley. Mont. Agr. Expt. Sta. Bul. 154: 1-56, 7 figs.

Reinhard, H. J.

- 1919 Preliminary notes on Texas Tachinidae (Diptera).
Ent. News 30 (10): 279-285.
- 1922 Host records of some Texas Tachinidae (Diptera).
Ent. News 33 (3): 72-73.
- 1929 The cotton-square borer. Tex. Agr. Expt. Sta. Bul.
401: 1-36, figs. 1-4.
- 1930 On the genus *Viviania* with the description of two
new species from Texas. (Tachinidae, Diptera).
Brooklyn Ent. Soc. Bul. 25 (2): 102-107.
- 1931 Revision of the American parasitic flies belonging to
the genus *Winthemia*. U.S. Natl. Mus. Proc. 79
(20): 1-54, pl. 1.
- 1934 Two new species of the tachinid genus *Siphostur-*
miopsis with key and notes (Diptera). Ent. News
45 (1): 15-19.
- 1934a Revision of the American two-winged flies belonging
to the genus *Cuphocera*. U.S. Natl. Mus. Proc.
83 (2974): 45-70.
- 1934b New North American Tachinidae. Brooklyn Ent.
Soc. Bul. 29 (5): 186-195.
- 1935 New genera and species of American muscoid flies
(Tachinidae: Diptera). Ent. Soc. Amer. Ann. 28
(1): 160-173.
- 1935a North American two-winged flies of the genus
Doryphorophaga, (Tachinidae, Diptera). N.Y.
Ent. Soc. Jour. 43 (4): 387-394.
- 1937 New North American muscoid Diptera. Brooklyn
Ent. Soc. Bul. 32 (2): 62-74.
- 1938 The sorghum webworm (*Celama sorghiella* Riley).
Tex. Agr. Expt. Sta. Bul. 559: 1-35, figs. 1-3.
- 1943 New Tachinidae from northeastern United States
(Diptera). Brooklyn Ent. Soc. Bul. 38 (3): 78-90.
- 1945 New genera and species of North American Tach-
inidae (Diptera). Canad. Ent. 77 (2): 28-36.
- 1946 A review of the tachinid genera *Siphophyto* and
Coronimyia (Diptera). Ent. Soc. Wash. Proc. 48
(4): 79-92.
- 1952 Muscoid flies of the genus *Chaetophlepsis* (Diptera).
Kans. Ent. Soc. Jour. 25 (1): 13-21.
- 1955 North American tachinid flies of the genus *Wagneria*
(Diptera). Kans. Ent. Soc. Jour. 28 (2): 49-59.
- 1956 A synopsis of the tachinid genus *Leucostoma*
(Diptera). Kans. Ent. Soc. Jour. 29 (4): 155-168.

- 1957 New American muscoid Diptera (Sarcophagidae, Tachinidae). Ent. News 68 (4): 99-111.
- 1958 New American Tachinidae (Diptera). Ent. News 69 (9): 233-242.
- 1958a Notes on *Spathimeigenia* with descriptions of four new species (Diptera, Tachinidae). Ent. Soc. Wash. Proc. 60 (5): 207-212.
- 1959 New Nearctic Tachinidae (Diptera). Pan-Pacific Ent. 35 (3): 157-163.
- Remington, C. L.
- 1959 On the autecology of *Megathymus yuccae* in Florida, with notes on foodplant specificity (Hesperioidea). Lepidopterists' News (1958) 12 (5-6): 175-185.
- Renault, T. R., and McDougall, G. A.
- 1961 Spruce budworm (*Choristoneura fumiferana* Clemens). Canad. Insect Pest Rev. 39 (5): 198-199.
- Rendall, R. E.
- 1943 Maine Forest Comnr. Bien. Rpt. 1941-1942 24: 1-78.
- Rentz, D. C. (See also Arnaud and Rentz)
- 1961 [*Chimarocephala pacifica pacifica* (Thomas) parasitism]. Pan-Pacific Ent. 37 (1): 60.
- Rettenmeyer, C. W.
- 1961 Observations on the biology and taxonomy of flies found over swarm raids of army ants. (Diptera: Tachinidae, Conopidae). Kans. Univ. Sci. Bul. 42 (8): 993-1066.
- 1961a Unusual host-seeking behavior by flies found over swarm raids of Neotropical army ants. Ent. Soc. Amer., No. Cent. Branch, Proc. 16: 15-16.
- Rex, E. G.
- 1930 Facts pertaining to the Japanese beetle. N.J. Dept. Agr. Cir. 180: 1-31, figs. 1-10.
- 1934 Information on the Japanese beetle. N.J. Dept. Agr. Cir. 242: 1-34, figs. 1-15.
- 1940 Information on the Japanese beetle. N.J. Dept. Agr. Cir. 316: 1-34, figs. 1-15.
- 1944 Information on the Japanese beetle. N.J. Dept. Agr. Cir. 349: 1-34, figs. 1-15.
- Rice, P. L. (See also MacCreary and Rice)

- and Wolfenbarger, D. O.
 1945 Control of the European corn borer through the utilization of natural enemies. Del. Agr. Expt. Sta. Bul. 259: 21.
- Wolfenbarger, D. O., and Chada, H. L.
 1944 European corn borer studies. Del. Agr. Expt. Sta. Bul. 251: 23-24.
- [Richardson, R. W.]
 1959 Biological control studies. Rockefeller Found., Mex. Agr. Program, Dir. Ann. Rpt. 1958-1959: 151-153.
- Richmond, H. A.
 1938 Forest insects of the season 1937 in Manitoba and northern Ontario. Canad. Insect Pest Rev. 16 (1): 45-46.
 1940 Review of forest insects of the season 1939 from the Winnipeg laboratory. Canad. Insect Pest Rev. 18 (1): 49-54.
 1941 Forest insects of the season 1940 in central Canada. Canad. Insect Pest Rev. 19 (1): 57-60.
 1941a [*Nemorilla* parasite of jack pine budworm in Manitoba]. Canad. Insect Pest Rev. 19 (3): 209.
- Riley, C. V. (See also Walsh and Riley)
 1869 First annual report of the noxious, beneficial, and other insects, of the State of Missouri, made to the State Board of Agriculture, pursuant to an appropriation for this purpose from the legislature of the state. 181 + 7 + 1 pp., 98 figs. Jefferson City, Mo.
 1870 Second annual report on the noxious, beneficial, and other insects, of the State of Missouri, made to the State Board of Agriculture, pursuant to an appropriation for this purpose from the legislature of the state. 135 + 6 + 1 pp., 99 figs. Jefferson City, Mo.
 1870a The cecropia moth (*Attacus cecropia*, Linn.). Amer. Ent. 2 (4): 97-102, figs. 59-67.
 1870b Insects injurious to the grape-vine. —No. 8. Amer. Ent. and Bot. 2 (7): 208-209, fig. 127.
 1870c The tent-caterpillar of the forest, (*Clisiocampa sylvatica*, Harr.). Amer. Ent. and Bot. 2 (9): 261-266, figs. 165-168.
 1870d Insects injurious to the grape-vine. —No. 10, the common yellow bear, (*Spilosoma virginica*, Fabr.). Amer. Ent. and Bot. 2 (9): 272-273, fig. 170.

- 1871 Third annual report on the noxious, beneficial, and other insects, of the State of Missouri, made to the State Board of Agriculture, pursuant to an appropriation for this purpose from the legislature of the state. 175 + 7 + 1 pp., 73 figs. Jefferson City, Mo.
- 1872 Fourth annual report on the noxious, beneficial, and other insects, of the State of Missouri, made to the State Board of Agriculture, pursuant to an appropriation for this purpose from the legislature of the state. 145 + 6 pp., 66 figs. Jefferson City, Mo.
- 1873 Fifth annual report on the noxious, beneficial, and other insects, of the State of Missouri, made to the State Board of Agriculture, pursuant to an appropriation for this purpose from the legislature of the state. 160 + VIII pp., 75 figs. Jefferson City, Mo.
- 1875 Seventh annual report on the noxious, beneficial, and other insects of the State of Missouri, made to the State Board of Agriculture, pursuant to an appropriation for this purpose from the legislature of the state. 196 + III + 1 pp., 39 figs. Jefferson City, Mo.
- 1875a Potato bug. Pp. 768-771, figs. 1-7. In Ripley, G., and Dana, C. A., eds., *The American cyclopaedia: a popular dictionary of general knowledge*. Vol. 13 Palestine — printing. vi + 854 + vii pp., numerous figs. New York.
- 1875b The Colorado potato beetle. *Pop. Sci. Monthly* 7: 704-709, figs. 1-6.
- 1876 Eighth annual report on noxious, beneficial, and other insects of the State of Missouri, made to the State Board of Agriculture, pursuant to an appropriation for this purpose from the legislature of the state. 185 + IV + 1 pp., 55 figs. Jefferson City, Mo.
- 1876a Potato pests. Being an illustrated account of the Colorado potato-beetle and other insect foes of the potato in North America. With suggestions for their repression and methods for their destruction. 108 pp., 49 figs. New York.
- 1876b Notes on the yucca borer, *Megathymus yuccae* (Walk.). *Acad. Sci. St. Louis, Trans.* (1868-1877) 3: 323-344, figs. 25-31.

- 1877 Ninth annual report on the noxious, beneficial, and other insects of the State of Missouri, made to the State Board of Agriculture, pursuant to an appropriation for this purpose from the legislature of the state. 129 + III + 1 pp., 33 figs. Jefferson City, Mo.
- 1879 Parasites of the cotton worm. *Canad. Ent.* 11 (9): 161-162.
- 1880 The cotton worm in the United States. *Amer. Ent.* 3 (4): 93-95.
- 1884 Larval habits of the dipterous family Dexidae. *Amer. Nat.* 18 (1): 76-77.
- 1885 Fourth report of the United States Entomological Commission, being a revised edition of Bulletin No. 3, and the final report on the cotton worm, together with a chapter on the boll worm. XXXVIII + 399 + 147 pp., 45 figs., 64 pls. Washington.
- 1887 Our shade trees and their insect defoliators. Being a consideration of the four most injurious species which affect the trees of the capital; with means of destroying them. U.S. Dept. Agr., Div. Ent. Bul. 10: 1-69, figs. 1-27, pl. 1.
- [1891] On the time of transformation in the genus *Lachnosterna*. *Ent. Soc. Wash. Proc.* (1890) 2 (1): 132-134.
- 1891a On the habits and life history of *Diabrotica 12-punctata* Oliv. *Insect Life* (U.S. Dept. Agr., Div. Ent., Period. Bul.) 4 (3-4): 104-108, fig. 5.
- [Riley, C. V., and Howard, L. O.]
- 1888 The sweet-potato saw-fly. (*Schizocerus ebenus* Norton). [Order Hymenoptera; family Tenthredinidae]. *Insect Life* (U.S. Dept. Agr., Div. Ent., Period. Bul.) 1 (2): 43-45, figs. 7-8.
- 1889 Willow and poplar insects. *Insect Life* (U.S. Dept. Agr., Div. Ent., Period. Bul.) 2 (6): 164.
- 1890 [Reply to Mr. C. P. Gillette's letter of August 28, 1889]. *Insect Life* (U.S. Dept. Agr., Div. Ent., Period. Bul.) 2 (9): 281.
- Packard, A. S., Jr., and Thomas, C.
- 1878 First annual report of the United States Entomological Commission for the year 1877 relating to the Rocky Mountain locust and the best methods of preventing its injuries and of guarding against its

invasions, in pursuance of an appropriation made by Congress for this purpose. XVI + 477 + 295 pp., 111 figs., 5 pls. Washington.

- 1883 Third report of the United States Entomological Commission, relating to the Rocky Mountain locust, the western cricket, the armyworm, canker worms, and the Hessian fly; together with descriptions of the larvae of injurious forest insects, studies of the embryological development of the locust and other insects, and on the systematic position of the Orthoptera in relation to the other orders of insects. XIV + 347 + 12 + 92 pp., 64 pls. Washington.

Rindge, F. H.

- 1949 A revision of the geometrid moths formerly assigned to *Drepanulatrix* (Lepidoptera). Amer. Mus. Nat. Hist. Bul. 94 (5): 231-298, figs. 1-14.

Rings, R. W., and Brooks, R. F.

- 1958 Bionomics of the one-spot stink bug, *Euschistus variolarius* (Palisot de Beauvois), in Ohio. Ohio Agr. Expt. Sta. Res. Cir. 50: 1-16, figs. 1-3.

Ritcher, P. O.

- 1947 European corn borer in Kentucky. Ky. Agr. Expt. Sta. Bul. 502: 1-23, figs. 1-18.

Rivard, I.

- 1964 Notes on parasitism of ground beetles (Coleoptera: Carabidae) in Ontario. Canad. Jour. Zool. 42 (5): 919-920.

Roberts, D. W.

- 1966 Life history and parasites of *Evora hemidesma* (Zeller) (Lepidoptera: Olethreutidae). Boyce Thompson Inst. Contrib. 23 (5): 165-170, figs. 2-4.

Robineau-Desvoidy, J. B.

- 1863 Histoire naturelle des Diptères des environs de Paris. Vol. 1. xvi + 1143 pp., 1 pl. Auxerre.

Rockwood, L. P.

- 1925 An outbreak of *Agrotis ypsilon* Rott. on overflow land in western Oregon. Jour. Econ. Ent. 18: 717-721.

— and Chamberlin, T. R.

- 1943 The western spotted cucumber beetle as a pest of forage crops in the Pacific Northwest. Jour. Econ. Ent. 36: 837-842.

- Roever, K.
1964 Bionomics of *Agathymus* (Megathymidae). Jour. Res. Lepidoptera 3 (2): 103-120, figs. 1-5.
- Rolston, L. H., and Kendrick, R. L.
1961 Biology of the brown stink bug, *Euschistus servus* Say. Kans. Ent. Soc. Jour. 34 (3): 151-157, figs. 1-2.
- Rosa, [J. S.]
[1927] [*Archytas cirphis* reared from *Spodoptera* caterpillars.] Hawaii. Ent. Soc. Proc. 1926 6 (3): 360.
1946 *Chaetogaedia monticola* (Bigot). Hawaii. Ent. Soc. Proc. 1945 12 (3): 478.
1947 *Eucelatoria armigera* (Coquillett). Hawaii. Ent. Soc. Proc. 1946 13 (1): 6.
- Rose, A. H.
1964 Distribution of introduced parasites. Canada Dept. Forestry, Forest Ent. and Path. Branch, Ann. Rpt. 1964: 79.
- *Ross, D. A.
1952 Key to the puparia of the dipterous parasites of *Choristoneura fumiferana* Clem. Canad. Ent. 84 (4): 108-112, figs. 1-15.
1953 Key to puparia of the dipterous parasites of *Malacosoma* spp. in British Columbia. Ent. Soc. Brit. Columbia, Proc. (1952) 49: 19-23, figs. 1-12.
- Ross, W. A.
1939 Fruit insects of the season 1938 in Ontario. Canad. Insect Pest Rev. 17 (1): 44-52.
- Rowe, J. A.
1933 Records of Tachinidae from Illinois with description of one new species (Diptera). Ent. News 44 (5): 122-126, 1 fig.
- Russell, E. E.
1952 Stink bugs on seed alfalfa in southern Arizona. U.S. Dept. Agr., Dept. Cir. 903: 1-19, 3 figs.
- *Russell, H. M. (See also Chittenden and Russell)
1912 A note on the southern walking-stick (*Anisomorpha buprestoides* Stoll) and a tachinid parasite. Ent. Soc. Wash. Proc. 14 (3): 117.
- [Russo, G.]
1927 Informe sobre la actividad del laboratorio de entomologia. [Dominican Repub.] Estac. Agron. de Moca Informe 2: 75-107, 1 pl.

- 1927a Informe de entomologia agricola dominicana. [Dominican Repub.] Estac. Agron. de Moca Informe 3: 31-244, 32 pls.

Sabrosky, C. W.

- 1948 *Winthemia citheroniae*, new species, with notes on the correct name of *W. cecropia* (Diptera, Larvaevoridae). Ent. Soc. Wash. Proc. 50 (3): 63-67.
- 1950 Notes on Trichopodini (Diptera, Larvaevoridae), with description of a new parasite of cotton stainers in Puerto Rico. Wash. Acad. Sci. Jour. 40 (11): 361-371, figs. 1-11.
- 1953 Taxonomy and host relations of the tribe Ormiini in the Western Hemisphere, II. (Diptera, Larvaevoridae). Ent. Soc. Wash. Proc. 55 (6): 289-305.
- 1955 The interrelations of biological control and taxonomy. Jour. Econ. Ent. 48: 710-714.
- 1965 Hosts of the tachinid tribe Eutherini (Diptera). Ent. Soc. Wash. Proc. 67 (1): 61.
- 1969 A review of the genus *Juriniopsis* Townsend (Diptera: Tachinidae). Fla. Ent. 52 (2): 79-90, figs. 1-10.

— and Arnaud, P. H., Jr.

- 1965 Family Tachinidae (Larvaevoridae). In Stone, A., Sabrosky, C. W., Wirth, W. W., Foote, R. H., and Coulson, J. R., A Catalog of the Diptera of America North of Mexico, pp. 961-1108. U.S. Dept. Agr. Agr. Handb. 276: I-IV, 1-1696.

Sam, J. Ah (See Miyahira and Ah Sam)

Sanderson, E. D.

- 1906 Report on miscellaneous cotton insects in Texas. U.S. Dept. Agr. Div. Ent. Bul. 57: 1-63, figs. 1-33, pl. 1.
- 1913 Insect pests of farm, garden and orchard. XII + 648 pp., 513 figs. New York.

— and Jackson, A. D.

- 1909 The oblique-banded leafroller *Archips rosaceana* Harris. Jour. Econ. Ent. 2: 391-403, pls. 15-18.

Satterthwait, A. F.

- 1948 Important sunflower insects and their insect enemies. Jour. Econ. Ent. 41: 725-731.

— and Swain, R. B.

- 1946 The sunflower moth and some of its natural enemies. Jour. Econ. Ent. 39: 575-580.

Saunders, W.

- 1871 Insects injurious to the grape. Ontario, Noxious Insects, Ann. Rpt. 1: 29-52, figs. 34-52.
- 1873 On some innocuous insects. Ontario Ent. Soc. Ann. Rpt. 1872: 51-58, figs. 39-49.
- 1875 On some of our common insects. The beautiful wood nymph — *Eudryas grata*. Canad. Ent. 7 (3): 41-44.
- 1875a On some of our common insects. Ontario Ent. Soc. Ann. Rpt. 1874: 22-28, figs. 17-29.
- 1876 On some of our common insects. Ontario Ent. Soc. Ann. Rpt. 1875: 36-42, figs. 17-29.

Savely, H. E., Jr.

- 1939 Ecological relations of certain animals in dead pine and oak logs. Ecol. Monog. 9 (3): 321-385, figs. 1-8.

Scaramuzza, L. C.

- 1929 Los gusanos de las hierbas que atacan la caña de azúcar en Cuba. Asoc. de Téc. Azucareros Cuba, Mem. de la Conf. An. 3: 119-124.
- 1929a Grass worms attacking sugar cane in Cuba. Asoc. de Téc. Azucareros Cuba, Proc. 3: 110-115.
- 1930 Preliminary report on a study of the biology of *Lixophaga diatraeae* Tns. Jour. Econ. Ent. 23: 999-1004.
- 1932 Prospects for the control of the sugar cane moth stalk-borer (*Diatraea saccharalis* Fab.) in Cuba by means of natural enemies. Asoc. de Téc. Azucareros Cuba, Proc. 6: 87-93.
- 1932a Beneficios que puede reportar a la caña el cultivo del arroz. Rev. de Agr. Com. y Trab. Cuba 14 (4): 58-60, 2 figs.
- 1934 La primera introducción en Cuba de un parásito contra el "borer" o perforador de la caña de azúcar (*Diatraea saccharalis* Fabr.). Rev. de Agr. [Cuba] 15 (6): 13-17.
- 1939 The introductions of *Theresia claripalpis* V. d. W., into Cuba, and its artificial multiplication. Internatl. Soc. Sugar Cane Technol. Cong. Proc. 6: 589-595, 2 figs.
- 1942 Los insectos y otros animales que atacan a la caña de azúcar en Cuba. Asoc. de Téc. Azucareros de Cuba, Bol. Ofic. 1 (1): 22-24; (4): 50-51; (5): 64-67; (7): 95-98; (9): 121-123; (10): 146-151.

- 1946 Capitulo XII. Los insectos y otros animales que atacan a la caña de azucar en Cuba. Pp. 529-563, figs. 1-19, 1 pl. *In* Agete y Pinero, F., La cana azucar en Cuba. Vol. 2. Pp. 297-602, numerous figs. Havana.
- 1950 Comentarios acerca del control del borer o perforador de la caña, por medio de insecticidas. *Asoc. Téc. Azucareros Cuba, Bol. Ofic.* 9 (3): 97-100.
- 1951 El control biologico y sus resultados en la lucha contra el barrenador o perforador de la caña, *Diatraea saccharalis* (Fabr.) en Cuba, por medio de la mosca *Lixophaga diatraeae* (Town.). *Mex., Sec. de Agr. y Ganaderia, Fol. Misc.* 4: 282-292.
- 1951a Relacion de los insectos y otros animales que atacan a la caña de azucar en Cuba. *Mex., Sec. de Agr. y Ganaderia, Fol. Misc.* 4: 292-302.
- 1955 El borer y la restriccion de la zafra. *Asoc. de Téc. Azucareros de Cuba, Bol. Ofic.* 14 (6): 319-321.
- 1960 Damage by the sugarcane borer in Louisiana and Cuba. The importance of biological control. *Internatl. Soc. Sugar Cane Technol. Cong. Proc.* 10: 938-942.
- and Barry, F. V.
- 1960 A list of insects and animals affecting sugarcane in Cuba. *Internatl. Soc. Sugar Cane Technol. Cong. Proc.* 10: 994-1000.
- Fernández Artilles, R., and Fernández Perez, J.
- 1947 Results obtained with the biological control of the sugar cane borer in Cuba. *Asoc. de Téc. Azucareros Cuba, Proc.* 21: 1-6.
- 1948 Controlling the sugar cane borer in Cuba by biological control. *Asoc. de Téc. Azucareros Cuba, Proc.* 22: 37-40.
- and Ingram, J. W.
- 1942 Results attained in the biological control of *Diatraea saccharalis* (F.) in Florida. *Jour. Econ. Ent.* 35: 642-645.
- Schaffner, J. C. (See York, Schaffner, and Brindley)
- Schaffner, J. V., Jr. (See also Webber and Schaffner)
- 1927 Dispersion of *Compsilura concinnata* Meig. beyond the limits of the gipsy moth and the brown-tail moth infestation. *Jour. Econ. Ent.* 20: 725-732, fig. 25.

- 1934 Introduced parasites of the brown-tail and gipsy moths reared from native hosts. Ent. Soc. Amer. Ann. 27 (4): 585-592.
- 1959 Microlepidoptera and their parasites reared from field collections in the northeastern United States. U.S. Dept. Agr. Misc. Pub. 767: i-iv, 1-97.
- and Griswold, C. L.
- 1934 Macrolepidoptera and their parasites reared from field collections in the northeastern part of the United States. U.S. Dept. Agr. Misc. Pub. 188: 1-160.
- Schedl, K. E.
- 1932 Parasites reared from forest insects in 1929. Canad. Ent. 64 (1): 1-2.
- Schenk, J. A. (See Goyer and Schenk; Ollieu and Schenk)
- Schoene, W. J. (See also Parrott and Schoene)
- 1909 The tussock moth in orchards. N.Y. (Geneva) Agr. Expt. Sta. Bul. 312: 39-49, pls. 1-3.
- Schopp, R.
- 1931 The smartweed borer (*Pyrausta ainsliei* Heinrich, Lepidoptera) in Kansas. Kans. Ent. Soc. Jour. 4 (2): 25-38.
- Schread, J. C.
- 1945 Status of the natural enemies of the Japanese beetle in Connecticut. Conn. Agr. Expt. Sta. Bul. 488: 331-339.
- 1946 Japanese beetle investigations. Conn. Agr. Expt. Sta. Bul. 512: 49-50.
- 1947 Japanese beetle — life history and control. Conn. Agr. Expt. Sta. Bul. 505: 1-20, figs. 1-5.
- 1948 Control of the Japanese beetle. Conn. Agr. Expt. Sta. Cir. 166: 1-8, figs. 1-2, 2 figs.
- Schuh, J. (See also Zeller and Schuh)
- and Mote, D. C.
- 1948 The oblique-banded leaf roller on red raspberries [*Archips rosaceana* (Harris)]. Oreg. Agr. Expt. Sta. Tech. Bul. 13: 1-43, figs. 1-12.
- Schwardt, H. H. (See Tashiro and Schwardt)
- Scudder, G. G. E.
- [1966] *Hyalomya aldrichii* Townsend (Dipt., Tachinidae), a parasite of *Emblethis vicarius* Horvath (Hem., Lygaeidae). Ent. Monthly Mag. (1965) 101(1217-1219): 286.

- Seamans, H. L. (See also Parker, J. R., Strand, and Seamans)
 1923 Popular and practical entomology. Forecasting outbreaks of the pale western cutworm in Alberta. *Canad. Ent.* 55 (3): 51-53.
- and McMillan, E.
 1935 The effect of food plants on the development of the pale western cutworm (*Agrotis orthogonia* Morr.). *Jour. Econ. Ent.* 28: 421-425.
- Segal, B.
 1936 Synopsis of the Tabanidae of New York, their biology and taxonomy. I. The genus *Chrysops* Meigen. *N.Y. Ent. Soc. Jour.* 44 (1): 51-78.
- Seibels, [W. T.]. (See Dillier and Seibels)
- Seidel, D. R.
 1954 Two new dipteran parasites of *Autographa californica*. *Pan-Pacific Ent.* 30 (3): 186.
- Seifert, O.
 1901 Life-history of *Platysenta videns* Guen. *N.Y. Ent. Soc. Jour.* 9 (1): 12-19.
- Seín, F., Jr. (See Leonard and Seín)
- Sell, R. A.
 1915 Some notes on the western twelve-spotted and the western striped cucumber beetles. *Jour. Econ. Ent.* 8: 515-520.
- Sellers, W. F.
 1930 The identity of *Zenillia blanda* O. S. and *Zenillia virilis* A. & W. with notes on *Zenillia blandita* Coq. (Diptera, Tachinidae). *Ent. Soc. Amer. Ann.* 23 (3): 569-576, figs. 1-2.
 1943 The Nearctic species of parasitic flies belonging to *Zenillia* and allied genera. *U.S. Natl. Mus. Proc.* 93(3157): 1-108.
- *Severin, H. C. (See also Severin, H. H. P., Severin, and Hartung)
 1921 The army worm. *S. Dak. Off. State Ent. Cir.* 24: 1-10, figs. 1-4.
 1926 The common black field cricket, *Gryllus assimilis* (Fab.) and its control. *Jour. Econ. Ent.* 19: 218-227.
- Severin, H. H. P., Severin, H. C., and Hartung, W.
 1915 The stimuli which cause the eggs of the leaf-ovipositing Tachinidae to hatch. *Psyche* 22 (4): 132-137.

Shahjahan, M.

- 1968 Superparasitization of the southern green stink bug by the tachinid parasite *Trichopoda pennipes pilipes* and its effect on the host and parasite survival. Jour. Econ. Ent. 61: 1088-1091.
- 1968a Effect of diet on the longevity and fecundity of the adults of the tachinid parasite *Trichopoda pennipes pilipes*. Jour. Econ. Ent. 61: 1102-1103.

Shannon, R. C.

- 1914 Habits of some Tachinidae. Ent. Soc. Wash. Proc. 16 (4): 182.

Shaw, J. G.

- 1947 Parasites of a bag-making pierid, *Eucheira socialis* in Morelos, Mexico. Jour. Econ. Ent. 40: 436-437.

Sheals, R. A. (See Brown, R. C., and Sheals)

Sheets, L. W. (See Stevenson, W. A., Kaufman, and Sheets)

Sherman, F., Jr.

- 1915 Rearing of moths and tachina-flies from larvae and pupae of army-worm in North Carolina in 1914. Jour. Econ. Ent. 8: 299-302.
- 1920 The green clover worm (*Plathypena scabra* Fabr.) as a pest of soy beans. Jour. Econ. Ent. 13: 295-303.

Shimer, H.

- 1871 Additional notes on the striped squash beetle. Amer. Nat. 5 (4): 217-220, figs. 60-61.

Showalter, W. J.

- 1929 Exploring the wonders of the insect world. Natl. Geog. Mag. 56 (1): 1-90, 59 figs., pls. 1-24.

Sibley, L. M. (See Dugas et al.)

Sifuentes, J. A., and Young, W. R.

- 1960 Summary of insect conditions in the Yaqui Valley, Sonora, Mexico, 1959. (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 10 (18): 343-344.

*Silver, G. T. (See also Lejeune and Silver)

- 1958 Studies on the silver-spotted tiger moth, *Halisidota argentata* Pack. (Lepidoptera: Arctiidae), in British Columbia. Canad. Ent. 90 (2): 65-80, figs. 1-16.

Silver, J. C.

- 1933 Biology and morphology of the spindle worm, or elder borer. U.S. Dept. Agr. Tech. Bul. 345: 1-19, figs. 1-9.

- Simanton, W. A., and Andre, F.
 1936 A biological study of *Lygaeus kalmii* Stål (Hemiptera-Lygaeidae). Brooklyn Ent. Soc. Bul. 31 (3): 99-106, pl. 5.
- Simmonds, F. J.
 1948 The biological control of insect pests of sugarcane in the West Indies. Brit. W. Indies Sugar Technol. Proc. 1948: 66-73.
 1953 Insect pests of sugar-cane in the French Antilles. Trop. Agr. [Trinidad] 30 (4-6): 122-127.
 1953a [Southern green stink bug (*Nezara viridula*)]. (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 3 (45): 782.
 1955 Establishment of parasites of *Diatraea saccharalis* F. in Dominica (British West Indies) and Guadeloupe (French West Indies). Trop. Agr. [Trinidad] 32 (3): 198-200.
 1956 The present status of biological control. Canad. Ent. 88 (9): 553-563.
- Simmons, P., Reed, W. D., and M'Gregor, E. A.
 1931 Fig insects in California. U.S. Dept. Agr., Dept. Cir. 157: 1-72, figs. 1-38.
- Simpson, C. B.
 1903 The codling moth. U.S. Dept. Agr. Bul. 41: 1-105, figs. 1-19, pls. 1-16.
- Simpson, L. J. (See Balch and Simpson)
- Sippell, W. L.
 1961 Key to the puparia of common dipterous parasites of the forest tent caterpillar, *Malacosoma disstria* Hbn., in Ontario. Canad. Ent. 93 (6): 433-436, figs. 1-10.
- Sisojevic, P. (See Embree and Sisojevic)
- Skelkregg, E. R. (See Wood, W. B., and Skelkregg)
- Slate, W. L.
 1945 Progress of the station's work entomology. Conn. Agr. Expt. Sta. Bul. 484: 59-68.
- Smereka, E. P.
 1965 The life history and habits of *Chrysomela crotchii* Brown (Coleoptera: Chrysomelidae) in northwestern Ontario. Canad. Ent. 97 (5): 541-549, figs. 1-8.
- Smith, B. C.
 1960 Note on parasitism of two coccinellids, *Coccinella trifasciata perplexa* Muls. and *Coleomegilla maculata lengi* Timb. (Coleoptera: Coccinellidae) in Ontario. Canad. Ent. 92 (9): 652.

- Smith, C. C. (See also Reeks and Smith; Reeks, Smith, and Forbes)
- 1946 Notes on the birch shoot borer, *Epinotia solicitana* Walker (Olethreutidae, Lepidoptera). *Acadian Nat.* 2 (6): 114-121, pl. 5.
- 1950 [*Phryxe pecosensis* Tns. probable parasite of *Operophtera brumata* Linn.]. *Canad. Insect Pest Rev.* 28 (2): 105-106.
- Smith, C. W. (See Smith, R. W.)
- *Smith, D. R.
- 1962 Parasites reared from a species of *Neodiprion* found on Douglas-fir in Idaho (Hymenoptera: Diprionidae). *Pan-Pacific Ent.* 38 (3): 189.
- Smith, H. E.
- 1912 A contribution to North American dipterology. *Ent. Soc. Wash. Proc.* 14 (3): 118-127.
- 1915 New species of Tachinidae (Diptera) from New England. *Psyche* 22 (3): 98-102.
- 1916 New Tachinidae from North America. *Ent. Soc. Wash. Proc.* 18 (2): 94-98.
- 1917 Notes on New England Tachinidae, with the description of one new genus and two new species. *Psyche* 24 (2): 54-58.
- 1917a Five new species of North American Tachinidae. *Psyche* 24 (5): 137-141.
- 1918 Notes on North American Tachinidae, including the description of one new genus. *Ent. Soc. Wash. Proc.* (1917) 19 (1-4): 122-126.
- Smith, H. W. (See Manis et al.)
- Smith, J. B.
- 1893 Insects of the year in New Jersey. *Insect Life* (U.S. Dept. Agr., Div. Ent., Period. Bul.) 6 (2): 187-192.
- 1910 [*Datana ministra* Dru. with parasitic eggs]. *Ent. News* 21 (3): 146.
- Smith, L. B.
- 1929 Japanese and Asiatic beetle report. *N.J. State Dept. Agr. Ann. Rpt. 1928-1929* 14: 120-125.
- 1930 The Japanese beetle in 1929. *Jour. Econ. Ent.* 23: 495-501.
- Smith, L. M., and Kido, G. S.
- 1949 The raspberry leaf sawfly. *Hilgardia* 19 (2): 43-54, figs. 1-7.

- Smith, M. E.
1958 Predatism in a parasitized caterpillar. Brooklyn Ent. Soc. Bul. 53 (5): 123.
- Smith, O. J.
1953 Grape leaf skeletonizer. Two parasites of the western skeletonizer colonized in successful search for natural enemies of pest. Calif. Agr. 7 (5): 9, 3 figs.
1955 Western grape leaf skeletonizer. 1954 biological control program indicates parasitism plus virus disease registering important reduction of vineyard pest. Calif. Agr. 9 (8): 7, 3 figs.
- and Langston, R. L.
1953 Continuous laboratory propagation of western grape leaf skeletonizer and parasites by prevention of diapause. Jour. Econ. Ent. 46: 477-484.
- Smith, R. C.
1921 Observations on the fall army worm (*Laphygma frugiperda* Smith and Abbott) and some control experiments. Jour. Econ. Ent. 14: 300-305.
1924 *Caenurgia* [sic] *erechtea* (Cram.) (Noctuidae) as an alfalfa pest in Kansas. Jour. Econ. Ent. 17: 312-319, fig. 11.
1943 Descriptions of Kansas insects. A list and brief description of the most important or interesting species of insects in Kansas by orders and families. Kans. State Bd. Agr. Rpt. 62 (255): 117-414, figs. 1-464, pls. 1-6.
- and Franklin, W. W.
1954 The garden webworm — *Loxostege similalis* Guen. as an alfalfa pest in Kansas. Kans. Ent. Soc. Jour. 27 (1): 27-38, figs. 1-7.
- Smith, R. F. (See Michelbacher, MacLeod, and Smith)
- Smith, R. W. (Published also as C. W. Smith)
1929 Parasitism of the oriental peach moth in Ontario with special reference to biological control experiments with *Trichogramma minutum* Riley. Ontario Ent. Soc. Ann. Rpt. 59: 72-80, figs. 3-6.
1939 An exchange of grasshopper parasites between Argentina and Canada with notes on parasitism of native grasshoppers. Ontario Ent. Soc. Ann. Rpt. 70: 57-62.
1941 Successful hibernation of the earwig parasite *Bigonicheta setipennis* Fall. in Ontario. Ontario Ent. Soc. Ann. Rpt. 1940 71: 29-32.

- 1944 Observations on parasites of some Canadian grasshoppers. *Canad. Ent.* 76 (2): 28-33.
- 1965 A field population of *Melanoplus sanguinipes* (Fab.) (Orthoptera: Acrididae) and its parasites. *Canad. Jour. Zool.* 43 (1): 179-201, 7 figs.
- and Finlayson, T. U.
- 1950 Larvae of dipterous parasites of nymphal and adult grasshoppers. *Canad. Jour. Res., Sect. D, Zool. Sci.* 28 (2): 81-117, figs. A-J, 1-51.
- Smyth, E. G.
- 1917 The white-grubs injuring sugar cane in Porto Rico. *Porto Rico Dept. Agr. Jour.* 1 (2): 47-92, pls. 2-9.
- 1917a The white-grubs injuring the sugar cane in Porto Rico. 1. Life-histories of May-beetles --continued. *Porto Rico Dept. Agr. Jour.* 1 (3): 141-169.
- Snapp, O. I.
- 1930 Life history and habits of the plum curculio in the Georgia peach belt. *U.S. Dept. Agr. Tech. Bul.* 188: 1-90, figs. 1-7, pls. 1-10.
- and Swingle, H. S.
- 1929 Life history of the oriental peach moth in Georgia. *U.S. Dept. Agr. Tech. Bul.* 152: 1-16, figs. 1-3.
- Snodgrass, R. E.
- 1917 Some of the important insect pests of Indiana. *Ind. State Ent. Ann. Rpt. 1915-1916* 9: 105-225, figs. 1-60.
- Snow, F. H.
- 1880 Letter. *In* The oblique-banded moth of the rose tribe. (*Loxotaenia rosaceana*, Harris.) [by] Sec'y. [G. C. Brackett]. Pp. 201-202. *Kans. State Hort. Soc. Rpt.* (1879) 9: 201-202.
- *Snow, S. J.
- 1925 Observations on the cutworm, *Euxoa auxiliaris* Grote, and its principal parasites. *Jour. Econ. Ent.* 18: 602-609.
- Snyder, T. E.
- 1936 Puss caterpillar (*Megalopyge opercularis* S. and A.). *Insect Pest Survey Bul.* 16 (9): 423.
- Sommers, R. A. (See Weaver, J. E., and Sommers)
- Soper, R. S., Jr., and Olson, R. E.
- 1963 Survey of biota associated with *Monochamus* (Coleoptera: Cerambycidae) in Maine. *Canad. Ent.* 95 (1): 83-95, figs. 1-4.

- Sorenson, C. J. (See Thornley and Sorenson)
- Soule, C. G. (See also Eliot and Soule)
- 1899 Color-variation in larvae of *Papilio polyxenes*, and other notes. *Psyche* 8 (284): 435-436.
- Sparks, A. N., Raun, E. S., and Carter, S. W.
- 1963 *Lydella grisescens* R. D. populations in corn borers of Boone County, Iowa. *Ent. Soc. Amer., No. Cent. Branch, Proc.* 18: 20-21, fig. 1.
- Spencer, G. J. (See also Crawford and Spencer)
- 1923 Further notes on the life history of the European corn borer in Ontario. *Ontario Ent. Soc. Ann. Rpt.* 53: 18-25.
- 1926 An ear-wig new to British Columbia. *Pacific Coast Ent. Soc. Proc.* 2 (4): 65-66.
- 1936 An outbreak of *Autographa californica* Speyer around Kamloops, summer 1934. *Ent. Soc. Brit. Columbia, Proc.* (1935) 32: 26-28.
- 1947 The 1945 status of *Digonochaeta setipennis*, tachinid parasite of the European earwig *Forficula auricularia* Linn. in West Point Grey, Vancouver, B.C. *Ent. Soc. Brit. Columbia, Proc.* (1946) 43: 8-9.
- Stafford, E. W. (See Summers and Stafford)
- St. Amand, W., and Cloyd, W. J.
- 1954 Parasitism of the grasshopper, *Chortophaga viridifasciata* (DeGeer) (Orthoptera: Locustidae), by dipterous larvae. *Jour. Parasitol.* 40 (1): 83-87, 2 pls.
- Stark, R. W.
- 1959 Parasites of *Petrova metallica* Busck. *Canad. Insect Pest Rev.* 37 (3): 162-163.
- Stearns, H. C. (See Mote, Stearns, and Dimick)
- Stearns, L. A.
- 1919 Some recently recorded parasites of the oriental peach moth. *Jour. Econ. Ent.* 12: 347-348.
- 1928 The larval parasites of the oriental peach moth (*Laspeyresia molesta* Busck) with special reference to the biology of *Macrocentrus ancylihora* Rohwer. *N.J. Agr. Expt. Sta. Bul.* 460: 1-24, figs. 1-3, 1 fig.
- 1933 Observations on the biology and control of *Metriona bivittata* Say. *Jour. Econ. Ent.* 26: 151-154, pls. 4-5.

- and Amos, J. M.
 1941 Ten-year record of oriental fruit moth parasitism in Delaware. Jour. Econ. Ent. 34: 245-248, fig. 1.
- Steenburgh, W. E. [van]
 1931 The biological control factors affecting the abundance of the oriental peach moth (*Laspeyresia molesta* Busck) in Ontario during 1930. Ontario Ent. Soc. Ann. Rpt. 1930 61: 57-65.
 1932 The parasites of the oriental fruit moth (*Laspeyresia molesta* Busck) in Ontario 1931. Ontario Ent. Soc. Ann. Rpt. 1931 62: 66-69.
 1935 Parasites of the oriental fruit moth (*Laspeyresia molesta* Busck) in Ontario. A summary 1932-33-34. Ontario Ent. Soc. Ann. Rpt. 1934 65: 68-72.
- and Boyce, H. R.
 1938 Biological control of the oriental fruit moth *Laspeyresia molesta* Busck in Ontario: A review of ten years' work. Ontario Ent. Soc. Ann. Rpt. 69: 65-74.
- Stehr, F. W., and Cook, E. F.
 1968 A revision of the genus *Malacosoma* Hübner in North America (Lepidoptera: Lasiocampidae): systematics, biology, immatures, and parasites. U.S. Natl. Mus. Bul. 276: i-vi, 1-321, figs. 1-399.
- Steiner, H. M., and Worthley, H. N.
 1941 The pistol case-bearer and its control in Pennsylvania orchards. Pa. Agr. Expt. Sta. Bul. 406: 1-26, figs. 1-9.
- Steinhaus, E. A., and Hughes, K. M.
 1952 A granulosis of the western grape leaf skeletonizer. Jour. Econ. Ent. 45: 744-745, figs. 1-3.
- Stephens, C. S. (See also Harrison and Stephens)
 1962 *Oiketicus kirbyi* (Lepidoptera: Psychidae) a pest of bananas in Costa Rica. Jour. Econ. Ent. 55: 381-386, figs. 1-2.
- *Stevenson, C.
 1903 Notes on the season of 1902. (Western Quebec). Ontario Ent. Soc. Ann. Rpt. 33: 57-58, fig. 35.
- Stevenson, W. A., Kaufman, W., and Sheets, L. W.
 1957 The salt-marsh caterpillar and its control in Arizona. Jour. Econ. Ent. 50: 279-280.
- Stone, M. W.
 1965 Biology and control of the lima-bean pod borer in southern California. U.S. Dept. Agr. Tech. Bul. 1321: 1-46, 15 figs.

- Stoner, A. (See Bryan, D. E., Jackson, and Stoner)
- Strand, A. L. (See also Parker, J. R., Strand, and Seamans)
- 1937 Montana insect pests for 1935 and 1936. The twenty-sixth report of the state entomologist of Montana. Mont. Agr. Expt. Sta. Bul. 333: 1-39, figs. 1-7.
- Strauss, J. F.
- 1916 The grape leaf-folder. U.S. Dept. Agr., Dept. Bul. 419: 1-16, figs. 1-6, pls. 1-4.
- Strickland, E. H.
- 1916 The army cutworm *Euxoa (Chorizagrotis) auxiliaris* Grote. Canada Dept. Agr. Ent. Bul. 13: 1-31, figs. 1-15.
- 1921 Popular and practical entomology. Parasites of the pale western cutworm in Alberta. Canad. Ent. 53 (5): 97-100.
- 1923 Biological notes on parasites of prairie cutworms. Canada Dept. Agr. Bul. 26: 1-40, 20 figs.
- 1937 The northern limits for potato beetle infestation. Sci. Agr. 17 (7): 447-450.
- 1938 An annotated list of the Diptera (flies) of Alberta. Canad. Jour. Res., Sect. D, Zool. Sci. 16: 175-219, 1 fig.
- 1946 An annotated list of the Diptera (flies) of Alberta. Additions and corrections. Canad. Jour. Res., Sect. D, Zool. Sci. 24: 157-173.
- Strohecker, [H. A.], and Yackley, [W. H.]
- 1963 Parasites in Pennsylvania. (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 13 (47): 1354.
- Struble, G. R.
- 1957 Biology and control of the white-fir sawfly. Forest Sci. 3 (4): 306-313, 5 figs.
- Stuckey, H. P.
- 1938 Entomology. Ga. Expt. Sta. Ann. Rpt. 50: 65-71.
- 1940 Entomology. Ga. Expt. Sta. Ann. Rpt. 52: 58-64.
- 1941 Entomology. Ga. Expt. Sta. Ann. Rpt. 53: 112-119, 2 figs.
- 1942 Entomology. Ga. Expt. Sta. Ann. Rpt. 54: 77-83, figs. 12-14.
- 1943 Entomology. Ga. Expt. Sta. Ann. Rpt. 55: 49-52, fig. 8.
- Stultz, H. T.
- 1955 The influence of spray programs on the fauna of apple orchards in Nova Scotia. VIII Natural

- enemies of the eye-spotted bud moth, *Spilonota ocellana* (D. & S.) (Lepidoptera: Olethreutidae).
 Canad. Ent. 87 (2): 79-85.
- Suehiro, [A.] (See Holdaway and Suehiro)
- Summerland, S. A.
 1938 The biology and synonymy [sic] of the parasites of the strawberry leaf roller, *Ancylis comptana* Froel. (Lepidoptera, Tortricidae), found in Kansas. Kans. Acad. Sci. Trans. (1937) 40: 161-178.
- and Hamilton, D. W.
 1954 A leaf-roller, *Platynota flavedana* Clem., attacking peaches. Jour. Econ. Ent. 47: 941.
- Summers, T. E., and Stafford, E. W.
 1953 *Diabrotica undecimpunctata howardi* parasitized by *Celatoria diabroticae*. Jour. Econ. Ent. 46: 180.
- Surface, H. A.
 1904 Special article: The apple-tree tent-caterpillar (*Clisiocampa americana*). Pa. State Dept. Agr., Div. Zool., Monthly Bul. 1 (11-12): 35-49, figs. 1-9.
- Sutherland, D. W. S.
 1966 Biological investigations of *Trichoplusia ni* (Hübner) and other Lepidoptera damaging cruciferous crops on Long Island, New York. N.Y. (Cornell) Agr. Expt. Sta. Mem. 399: 1-99, figs. 1-25.
- Svec (See Wressell, Driscoll, and Svec)
- Swain, R. B. (See also Satterthwait and Swain)
 1937 The parasites of the fall webworm. *Hyphantria cunea* Drury. (Lep.: Arctiidae). Ent. News 48 (9): 244-248.
 1937a Notes in the natural control of the fall webworm in Colorado. Colo.-Wyo. Acad. Sci. Jour. 2 (2-3): 43-44.
- Sweetman, H. L.
 1926 Results of life history studies of *Diabrotica 12-punctata* Fabr. (Chrysomelidae, Coleoptera). Jour. Econ. Ent. 19: 484-490, fig. 25.
 1936 The biological control of insects. With a chapter on weed control. xii + 461 pp., 142 + 2 figs. Ithaca, N. Y.
- Swenk, M. H.
 1927 The pine tipmoth in the Nebraska National Forest. Nebr. Univ. Agr. Expt. Sta. Res. Bul. 40: 1-50, figs. 1-9.

Swezey, O. H. (See also Muir and Swezey)

- 1906 Observations on insects during a recent trip on Hawaii. Hawaii. Ent. Soc. Proc. 1905 1 (1): 16-19.
- 1907 The sugar cane leaf-roller (*Omiodes accepta*) with an account of allied species and natural enemies. Hawaii. Sugar Planters' Assoc. Expt. Sta. Rpt., Ent. Bul. 5: 1-60, pls. 1-6.
- 1908 Notes on *Chaetogaedia monticola* Bigot. Hawaii. Ent. Soc. Proc. 1907 1 (5): 163-164.
- 1908a Observations of the life-history of *Chaetogaedia monticola* Bigot. Hawaii. Ent. Soc. Proc. 2 (1): 7-9.
- 1908b Army worms and cut worms attacking sugar cane. Hawaii. Sugar Planters' Assoc. Expt. Sta., Div. Ent. Cir. 5: 1-21, pls. 1-3.
- 1909 Army worms and cut worms of sugar cane in the Hawaiian Islands. Hawaii. Sugar Planters' Assoc. Expt. Sta., Div. Ent. Bul. 7: 1-32, pls. 1-3.
- 1911 The cane borer parasite colonies. Hawaii. Planters' Rec. 4 (4): 185.
- 1912 Insects associated with "mamake" (*Pipturus albidus*), a native Hawaiian tree. Hawaii. Ent. Soc. Proc. 1910 2 (4): 153-163.
- 1914 The introduction of a tachinid parasite of the sugar cane weevil borer in Hawaii. Jour. Econ. Ent. 7: 455-457.
- 1915 Some results of the introduction of beneficial insects in the Hawaiian Islands. Jour. Econ. Ent. 8: 450-457.
- 1917 Insect pests of truck crops. Hawaii. Planters' Rec. 16 (6): 426-435, 14 figs.
- 1918 New records of insects on Kauai. Hawaii. Ent. Soc. Proc. 1917 3 (5): 379-380.
- 1921 Tachinid parasite of *Corizus*. Hawaii. Ent. Soc. Proc. 1920 4 (3): 467.
- 1922 *Frontina archippivora*. Hawaii. Ent. Soc. Proc. 1921 5 (1): 8-9.
- 1922a *Leucostoma atra*. Hawaii. Ent. Soc. Proc. 1921 5 (1): 30-31.
- 1923 *Heliothis obsoleta*. Hawaii. Ent. Soc. Proc. 1922 5 (2): 184-185.
- 1923a Records of introduction of beneficial insects into the Hawaiian Islands. Hawaii. Ent. Soc. Proc. 1922 5 (2): 299-304.

- 1924 Sugar cane entomology. Internatl. Soc. Sugar Cane Technol. Cong. Proc. 1: 13-16.
- 1924a Entomological projects in general. Hawaii. Sugar Planters' Assoc. Committee Charge Expt. Sta. Rpt. 1924: 16-25.
- 1925 Records of introduction of beneficial insects into the Hawaiian Islands. Hawaii. Planters' Rec. 29 (4): 369-376.
- 1926 *Archytas* sp. Hawaii. Ent. Soc. Proc. 1925 6 (2): 226.
- 1926a Reports on defoliation of Maui Koa forests by caterpillars. Hawaii. Planters' Rec. 30 (3): 354-362, 4 figs.
- 1927 *Chaetogaedia monticola* (Bigot). Hawaii. Ent. Soc. Proc. 1926 6 (3): 356.
- 1927a Notes on the Mexican tachinid, *Archytas cirphis* Curran, introduced into Hawaii as an armyworm parasite. (Diptera). Hawaii. Ent. Soc. Proc. 1926 6 (3): 499-503.
- 1927b Notes on the Mexican tachinid, *Archytas cirphis* Curran, introduced into Hawaii as an armyworm parasite. Hawaii. Planters' Rec. 31 (3): 214-217, 1 fig.
- 1928 Present status of certain insect pests under biological control in Hawaii. Jour. Econ. Ent. 21: 669-676.
- 1928a *Archytas cirphis* Curran. Hawaii. Ent. Soc. Proc. 1927 7 (1): 11.
- 1928b Hawaiian sugarcane borer (*Rhabdocnemis obscura* Boisd.). Insect Pest Survey Bul. 8 (4): 118.
- 1929 *Brachymeria obscurata* (Walker). Hawaii. Ent. Soc. Proc. 1928 7 (2): 211.
- 1929a Status of sugar cane pests in Hawaii in the year 1927. Hawaii. Ent. Soc. Proc. 1928 7 (2): 269.
- 1930 Entomology. Hawaii. Sugar Planters' Assoc. Proc. 49th Ann. Mtg. 1929 49: 278-287.
- 1931 *Rhabdocnemis obscura* (Boisd.). Hawaii. Ent. Soc. Proc. 1929-1930 7 (3): 391.
- 1931a Some observations on the insect faunas of native forest trees in the Olinda forest on Maui. Hawaii. Ent. Soc. Proc. 1929-1930 7 (3): 493-504.
- 1931b Entomology. Hawaii. Sugar Planters' Assoc. Proc. 50th Ann. Mtg. 1930 50: 425-432.
- 1931c Insect pests of sugar cane in Hawaii. Mid-Pacific Mag. 17 (4): 324-327, figs. 1-5.

- 1932 Summary of insect conditions in Hawaii for 1931.
Insect Pest Survey Bul. 11 (10): 680.
- [1933] Summary of insect conditions in Hawaii for 1932.
Insect Pest Survey Bul. (1932) 12 (10): 429-431.
- [1934] Summary of insect conditions in Hawaii for
1933. Insect Pest Survey Bul. (1933) 13
(10): 340-341.
- 1935 *Apanteles glomeratus* (Linn.). Hawaii. Ent. Soc.
Proc. 1934 9 (1): 25-26.
- 1935a The winter revival of insect life in the arid region at
Koko Head, Oahu. Hawaii. Ent. Soc. Proc. 1934 9
(1): 93-96.
- 1936 *Chaetogaedia monticola* (Bigot). Hawaii. Ent. Soc.
Proc. 1935 9 (2): 117.
- 1937 *Chloridea obsoleta* (Fab.). Hawaii. Ent. Soc. Proc.
1936 9 (3): 359.
- 1943 Host plant records of some Kauai Lepidoptera.
Hawaii. Ent. Soc. Proc. 1942 11 (3): 279-284.
- 1944 *Eucelatoria armigera* (Coq.). Hawaii. Ent. Soc. Proc.
1943 12 (1): 3.
- 1944a [*Chaetogaedia monticola* (Bigot)]. Hawaii. Ent. Soc.
Proc. 1943 12 (1): 4.
- 1947 *Vanessa cardui* (L.). Hawaii. Ent. Soc. Proc. 1946 13
(1): 5.
- Swift, J. E. (See Michelbacher et al.)
- Swingle, H. S. (See Snapp and Swingle)
- Tadic, M. (See Wylie and Tadic)
- Talerico, R. L.
- 1968 Life history of the looper *Phigalia titea* in Virginia.
Ent. Soc. Amer. Ann. 61 (3): 557-561, figs. 1-7.
- Tanada, Y.
- 1966 Field observation of the biotic factors regulating the
population of the armyworm, *Pseudaletia
unipuncta* (Haworth). Hawaii. Ent. Soc. Proc.
1965 19 (2): 302-308, fig. 1.
- and Beardsley, J. W., Jr.
- 1958 A biological study of the lawn armyworm, *Spodo-
ptera mauritia* (Boisduval), in Hawaii
(Lepidoptera: Phalaenidae). Hawaii. Ent. Soc.
Proc. 1957 16 (3): 411-436, figs. 1-7.
- Tashiro, H., and Schwardt, H. H.
- 1953 Some natural enemies of horse flies in New York.
Jour. Econ. Ent. 46: 680-681, figs. 1-2.
- Tauber, O. E. (See Drake, Harris, and Tauber).

- Taylor, E. A. (See also Hills and Taylor)
 1954 Parasitization of the salt-marsh caterpillar in Arizona. *Jour. Econ. Ent.* 47: 525-530.
- *Taylor, E. J.
 1949 A life history study of *Nabis alternatus*. *Jour. Econ. Ent.* 42: 991.
 1951 Biology of the damsel bug — *Nabis alternatus*. *Utah Acad. Sci., Arts and Letters, Proc.* 26: 132-133.
- Taylor, R. L.
 1929 The arthropod fauna of coniferous leaders weeviled by *Pissodes strobi* (Peck). *Psyche* 35 (4): 217-225.
- Teale, [E. W.]
 1946 [Summer collecting experiences]. *Brooklyn Ent. Soc. Bul.* 41 (1): 24-25.
 1948 Scourge of the monarch. The deadly larva of a fly brings tragedy to its powerless host. *Nat. Hist.* 57 (8): 356-359, 11 figs.
- Teetes, G. L., and Randolph, N. M.
 1969 Seasonal abundance and parasitism of the sunflower moth, *Homoeosoma electellum*, in Texas. *Ent. Soc. Amer. Ann.* 62 (6): 1461-1464, fig. 1.
- Telford, H. S. (See Munro and Telford)
- Teranishi, C. (See Clausen, King, and Teranishi)
- Terrazas Loyola, J.
 1961 Introduccion al estudio de la biologia y control de *Lema nigrovittata* (Guerim) [sic] en Chapingo, Mex. *Fitofilo* 14(31): 33-39.
- Teskey, H. J.
 1969 Larvae and pupae of some eastern North American Tabanidae (Diptera). *Ent. Soc. Canada, Mem.* 63: 1-147, figs. 1-148.
- Thames, W. H., Jr. (See also Mathes et al.)
 1954 Insect investigations. *Fla. Agr. Expt. Sta. Ann. Rpt.* 1954: 236.
- Mathes, R., Bregger, T., and Questal, D. D.
 1952 Studies upon the prevalence and control of the sugarcane moth borer in south Florida. *Fla. Agr. Expt. Sta. Ann. Rpt.* 1952: 207-208.
- Thomas, C. (See also Riley, Packard, and Thomas)
 1881 The army worm. *Noxious and Beneficial Insects of State of Ill., State Ent. Rpt.* 10: 5-43, figs. 1-5.
- *Thomas, C. A.
 1940 The biology and control of wireworms. A review of

- the literature. Pa. Agr. Expt. Sta. Bul. 392: 1-90.
- Thomas, G. M. (See Brewer and Thomas)
- Thomas, I. E. (See Wishart and Thomas)
- Thomas, J. B.
1955 Notes on insects and other arthropods in red and white pine logging slash. *Canad. Ent.* 87 (8): 338-343.
- Thomas, W. A.
1931 [Cowpea curculio (*Chalcodermus aeneus* Boh.) parasitized by *Myiophasia aenea* Wied.]. *Insect Pest Surv. Bul.* 11 (8): 519.
- Thomas, W. L. (See Knowlton, Janes, and Thomas)
- Thompson, M. C. (See Thompson, W. R., and Thompson)
- Thompson, R. W.
1944 Insects of the season 1943 in Ontario. *Canad. Insect Pest Rev.* 22 (1): 29-44.
- *Thompson, W. R. (See also Fiske and Thompson)
- 1910 Notes on the pupation and hibernation of tachinid parasites. *Jour. Econ. Ent.* 3: 283-295.
- 1911 Tachinidae, new and old. *Canad. Ent.* 43 (8): 265-272, pl. 4.
- 1943—
1958 A catalogue of the parasites and predators of insect pests. 2 sects. Belleville, Ottawa, Ont.
- 1953 The tachinid parasites of *Archips cerasivorana* Fitch. (1) *Dichaetoneura leucoptera* Johns. (Diptera). *Canad. Ent.* 85 (1): 19-30, pls. 1-3.
- 1953a Note on *Dichaetoneura leucoptera* Johns. (Diptera, Tachinidae). *Canad. Ent.* 85 (10): 391-392, fig. 1.
- 1953b The tachinid parasites of *Archips cerasivorana* Fitch. (2) *Eusisyropa blanda* O. S. (Diptera). *Canad. Ent.* 85 (11): 393-404, figs. 1-19.
- 1954 *Hyalomyodes triangulifera* Loew. (Diptera, Tachinidae). *Canad. Ent.* 86 (3): 137-144, pls. 1-2.
- and Thompson, M. C.
1921 Studies of *Zenillia roseanae* B. & B. a parasite of the European corn borer. (*Pyrausta nubilalis* Hb.). *Ent. Soc. Wash. Proc.* 23 (6): 127-139, pls. 8-10.
- 1923 *Masicera senilis*, a parasite of the European corn borer (*Pyrausta nubilalis*). *Ent. Soc. Wash. Proc.* 25 (2): 33-44, figs. 1-21, pls. 1-3.
- Thomley, H. F. (See also Knowlton et al.)

——— and Sorenson, C. J.

- 1940 Some insect pests of farm crops in the juniper-pinon belt of Utah. Utah Acad. Sci., Arts and Letters, Proc. 17: 47-48.

Tiernan, C. F. (See Williams, C. B., Jr., Walton, and Tiernan)

Tilden, J. W.

- 1951 The insect associates of *Baccharis pilularis* De Candolle. Microentomology 16 (1): 149-188.
- 1954 Microlepidoptera associated with *Baccharis pilularis*. III. Aegeriidae, Coleophoridae. Wasmann Jour. Biol. 12 (1): 43-52.

Timberlake, P. H.

- 1927 Biological control of insect pests in the Hawaiian Islands. Hawaii. Ent. Soc. Proc. 1926 6 (3): 529-556, 5 figs., pls. 19-23.

Tissot, A. N. (See also Kuitert and Tissot)

- 1945 Biology and control of cutworms and armyworms in Florida. Fla. Agr. Expt. Sta. Ann. Rpt. 1945: 64.

Titus, E. S. G.

- 1905 The sugar-beet crown-borer. U.S. Dept. Agr. Bur. Ent. Bul., n. ser., 54: 34-40, figs. 9-14.

——— and Pratt, F. C.

- 1904 Catalogue of the exhibit of economic entomology at the Louisiana Purchase Exposition, St. Louis, Mo., 1904. U.S. Dept. Agr. Div. Ent. Bul., n. ser., 47: 1-155.

Torgersen, T. R., and Coppel, H. C.

- 1969 Parasites of *Rhyacionia buoliana* and their occurrence at different crown levels of red pine in Wisconsin. Ent. Soc. Amer. Ann. 62 (1): 163-169.

Tothill, J. D.

- 1912 Systematic notes on North American Tachinidae. Canad. Ent. 44 (1): 1-5.
- 1913 Tachinidae and some Canadian hosts. Canad. Ent. 45 (3): 69-75.
- 1913a Progress of the introduction of the insect enemies of the brown-tail moth, *Euproctis chrysorrhoea* Linn. into New Brunswick and some biological notes on the host. Ontario Ent. Soc. Ann. Rpt. 1912 43: 57-61.
- 1916 The introduction and establishment in Canada of the natural enemies of the brown-tail and gipsy moths. Agr. Gaz. Canada 3 (2): 111-116.

- 1919 The meaning of natural control. Ent. Soc. Nova Scotia, Proc. (1918) 4: 10-14.
- 1920 Insect outbreaks and their causes. Ontario Ent. Soc. Ann. Rpt. 50: 31-33.
- 1921 A revision of the Nearctic species of the tachinid genus *Ernestia* R. D. (Diptera). Canad. Ent. 53 (10): 226-230; (12): 270-274.
- 1922 The natural control of the fall webworm [*Hyphantria cunea* Drury] in Canada together with an account of its several parasites. Canada Dept. Agr. Bul., n. ser., 3: 1-107, figs. 1-99, pls. 1-6.
- [1923] Notes on the outbreaks of spruce budworm, forest tent caterpillar and larch sawfly in New Brunswick. Acadian Ent. Soc. Proc. (1922) 8: 172-182.

— and McLaine, L. S.

- 1919 The recovery in Canada of the brown-tail moth parasite *Compsilura concinnata* (Diptera, Tachinidae). Ontario Ent. Soc. Ann. Rpt. 1918 49: 35-39.

Tower, W. V.

- 1913 Report of the Board of Commissioners of Agriculture of Porto Rico for the period from January 1st, 1912, to June 30, 1913. Porto Rico Bd. Comnr. Agr. Rpt. 2: 7-24.

Townsend, C. H. T.

- 1891 A tachinid bred from a chrysalis. Canad. Ent. 23 (10): 206-207.
- 1891a Two new tachinids. Psyche 6 (181): 83-85.
- 1891b A parasite of the fall web-worm. Psyche 6 (187): 176-177.
- 1891c A tachinid parasite of the oak unicorn prominent. Psyche 6 (188): 187-188.
- 1891d An *Exorista* parasitic on *Lagoa opercularis*. Ent. News 2 (8): 159-160.
- 1891e A tachinid parasite of *Chrysophanus dione*. *Exorista chrysophani* n. sp. Ent. News 2 (10): 197-199.
- 1892 Notes on North American Tachinidae sens. str. with descriptions of new genera and species. Paper III. Amer. Ent. Soc. Trans. 19: 88-132, pls. 1-3.
- 1892a Notes on North American Tachinidae, with descriptions of new species. Paper VII. Amer. Ent. Soc. Trans. 19: 284-289.

- 1892b Tachinid parasite of *Eucaterva variaria* Grote, and other notes. *Psyche* 6 (193): 258-259.
- 1892c An *Aporia* bred from *Limacodes* sp. *Psyche* 6 (194): 275-276.
- 1892d A tachinid bred from larva of *Protoparce jamaicensis* (Butl.) in Jamaica. *Inst. Jamaica, Jour.* 1 (2): 70-71.
- 1893 Hosts of North American Tachinidae, etc., I. *Psyche* 6 (206): 466-468.
- 1893a The puparium of *Blepharipeza*. *Amer. Nat.* 27 (316): 402-404.
- 1893b The puparium of *Jurinia*. *Amer. Nat.* 27 (318): 576-577.
- 1893c Notes on some Jamaican Muscidae, with brief anonymous descriptions of doubtful forms. *Inst. Jamaica, Jour.* 1 (7): 314-317.
- 1895 Contributions to the dipterology of North America. II.—Tabanidae, Conopidae, Tachinidae, etc. *Amer. Ent. Soc. Trans.* 22: 55-80.
- 1897 Contributions from the New Mexico Biological Station. — No. II. (continued). On a collection of Diptera from the lowlands of the Rio Nautla, in the State of Vera Cruz. II. *Ann. and Mag. Nat. Hist., ser. 6*, 20: 19-33.
- 1908 The taxonomy of the muscoidean flies, including descriptions of new genera and species. *Smithsn. Inst., Smithsn. Misc. Collect.* 51 (2[=pub. 1803]): 1-138, 1 fig.
- 1909 Descriptions of some new Tachinidae. *Ent. Soc. Amer. Ann.* 2 (4): 243-250.
- 1912 Foundation of some new genera and species of muscoid flies mainly on reproductive and early-stage characters. *N.Y. Ent. Soc. Jour.* 20 (2): 107-119.
- 1912a [A dexiine parasite of *Lachnosterna* from Mexico]. Letter, pp. 39-40. In Van Dine, D. L., Progress report on introductions of beneficial parasites into Porto Rico. *Porto Rico Bd. Comnr. Agr. Rpt.* 1: 31-47.
- 1912b Six new genera of Nearctic Muscoidea. *Ent. Soc. Wash. Proc.* 14 (3): 163-166.
- 1913 On *Trichiopoda* Latreille, *Polistomyia* Townsend and *Trichopodopsis* new genus. *N.Y. Ent. Soc. Jour.* 21 (2): 147-148.

- 1913a Notes on Exoristidae (Tachinidae auctt.) and allies. N.Y. Ent. Soc. Jour. 21 (4): 301-305.
- 1913b Criticism and muscoid taxonomy. Insecutor Inscitiae Menstruus 1 (9): 115-117.
- 1915 Revision of *Myiophasia*. Ent. Soc. Wash. Proc. 17 (3): 107-114.
- 1916 Description of two new tachinids (Dip.). Ent. News 27 (5): 217.
- 1916a *Andrina radialis* Townsend, new name. Canad. Ent. 48 (1): 19.
- 1916b Elucidations of New England Muscoidea. Insecutor Inscitiae Menstruus 4 (1-3): 17-33.
- 1916c Some new North American muscoid forms. Insecutor Inscitiae Menstruus 4 (7-9): 73-78.
- 1919 New genera and species of muscoid flies. U.S. Natl. Mus. Proc. 56(2301): 541-592.
- 1919a New muscoid genera, species and synonymy (Diptera). Insecutor Inscitiae Menstruus (1918) 6 (10-12): 157-182.
- 1928 On the structure, host habit, affinity and geographic range of *Beskia* (Dipt.: Tachinidae). Ent. News 39 (5): 150-151.
- 1928a *Schistocercophaga*, new genus of locust parasites (Larvaevoridae) (Dipt.: Tachinidae). Ent. News 39 (5): 152.
- 1942 Manual of myiology in twelve parts. Part XII. General consideration of the Oestromuscaria. 365 pp., frontis. + 84 pls. Sao Paulo.

Tripp, H. A.

- 1960 *Spathimeigenia spinigera* Townsend (Diptera: Tachinidae), a parasite of *Neodiprion swaini* Middleton (Hymenoptera: Tenthredinidae). Canad. Ent. 92 (5): 347-359, figs. 1-15.
- 1962 The relationship of *Spathimeigenia spinigera* Townsend (Diptera: Tachinidae) to its host, *Neodiprion swaini* Midd. (Hymenoptera: Diprionidae). Canad. Ent. 94 (8): 809-818, fig. 1.
- 1962a The biology of *Perilampus hyalinus* Say (Hymenoptera: Perilampidae), a primary parasite of *Neodiprion swaini* Midd. (Hymenoptera: Diprionidae) in Quebec, with descriptions of the

- egg and larval stages. *Canad. Ent.* 94 (12): 1250-1270, figs. 1-12.
- 1965 The development of *Neodiprion swaini* Middleton (Hymenoptera: Diprionidae) in the Province of Quebec. *Canad. Ent.* 97 (1): 92-107, figs. 1-8.
- *Tucker, E. S.
- 1917 Outbreaks of the elegant looper (*Philtraea elegantaria* Hy. Edw.) on privet in Louisiana (Lep.). *Ent. News* 28 (9): 394-396.
- Tucker, R. W. E.
- 1939 Some aspects of the control of the sugarcane moth borer, *Diatraea saccharalis* F. *Internatl. Soc. Sugar Cane Technol. Cong. Proc.* 6: 240-243.
- 1940 '*Diatraea saccharalis*' un insecto de la caña de azúcar — cómo se le combate en la Isla Barbada. Parte II. *Hacienda* 35: 462-463, 3 figs.
- 1952 The insects of Barbados. *Puerto Rico Univ. Jour. Agr.* 36 (4): 330-363.
- Turner, N. (See Friend and Turner)
- Turner, P. E.
- 1945 Researches on sugar-cane agriculture in the British West Indies and British Guiana, 1945. *Brit. W. Indies Sugar Assoc. Rpt. Res. Work* 1945: 5-25.
- Turnock, W. J. (See also Ives et al.; Ives and Turnock; Ives et al.; Nairn et al.)
- 1953 Some aspects of the life history and ecology of the pitch nodule maker, *Petrova albicapitana* (Busck) (Lepidoptera — Olethreutidae). *Canad. Ent.* 85 (7): 233-243, figs. 1-6.
- 1960 Ecological life-history of the larch sawfly, *Pristiphora erichsonii* (Htg.) (Hymenoptera: Tenthredinidae), in Manitoba and Saskatchewan. *Canad. Ent.* 92 (7): 500-516, figs. 1-3.
- and Ives, W. G. H.
- 1962 Evaluation of mortality during the cocoon stage of the larch sawfly, *Pristiphora erichsonii* (Htg.). *Canad. Ent.* 94 (9): 897-902.
- and Melvin, J. C. E.
- 1963 The status of *Bessa harveyi* (Tnsd.) (Diptera: Tachinidae). *Canad. Ent.* 95 (6): 646-654.
- and Muldrew, J. A.
- 1964 Biological control of the larch sawfly. *Canada Dept. Forestry, Forest Ent. and Path. Branch, Ann. Rpt.* 1964: 95-96.

Twinn, C. R.

- 1936 A summary of insect conditions in Canada in 1935. Ontario Ent. Soc. Ann. Rpt. 1935 66: 80-95.
- 1937 A summary of the insect pest situation in Canada in 1936. Ontario Ent. Soc. Ann. Rpt. 1936 67: 73-87.
- 1938 A summary of the insect pest situation in Canada in 1938. Ontario Ent. Soc. Ann. Rpt. 69: 121-134.
- 1939 A summary of the insect pest situation in Canada in 1938. Canad. Insect Pest Rev. 17(1): 1-21.
- 1941 A summary statement concerning some of the more important insect pests in Canada in 1940. Canad. Insect Pest Rev. 19 (1): 1-15.

Udine, [E. J.]

- 1955 A European corn borer parasite (probably *Lydella stabulans*) — Pennsylvania. (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 5 (30): 722.

Underhill, G. W.

- 1934 The green stinkbug. Va. Polytechnic Inst. Agr. Expt. Sta. Bul. 294: 1-26, frontis., figs. 1-9.

Urquhart, F. A.

- 1960 The monarch butterfly. xxiv + 361 pp., 79 figs., 12 pls. Toronto, Ont.

Van Denburgh, R. S., Burbutis, P. P., and York, G. T.

- 1962 The re-introduction and recovery of *Lydella stabulans grisescens*, a parasite of the European corn borer in Delaware. Jour. Econ. Ent. 55: 11-14.

Van Dine, D. L.

- 1911 The sugar-cane insects of Hawaii. U.S. Dept. Agr. Bur. Ent. Bul., n. ser., 93: 1-54, figs. 1-5, pls. 1-4.
- 1913 The insects affecting sugar cane in Porto Rico. Jour. Econ. Ent. 6: 251-257.
- 1913a The introduction of parasites of May-beetles into Porto Rico. Porto Rico Bd. Comnr. Agr. Rpt. 2: 36-47.
- 1913b Insects injurious to sugar cane in Porto Rico, and their natural enemies. Brit. Guiana Bd. Agr. Jour. 6 (4): 199-203.
- 1913c Report of the entomologist. Porto Rico Sugar Prod. Assoc. Expt. Sta. (Rio Piedras) Bul. 5: 25-46.

— and Christenson, L. D.

- 1932 A revised list of the insects affecting sugar-cane in Cuba. Internatl. Soc. Sugar Cane Technol. Cong. Proc. 4, Bul. 116: 1-3.

Van Duzee, E. P.

- 1926 [Oak moth (*Phryganidea* [sic] *californica*) at Mill Valley, California]. Pan-Pacific Ent. 2 (4): 180.

Van Leeuwen, E. R. (See also Yothers and Van Leeuwen)

- 1929 Life history of the codling moth in northern Georgia. U.S. Dept. Agr. Tech. Bul. 90: 1-94, figs. 1-21.

Van Zwaluwenburg, R. H.

- 1917 Insects affecting coffee in Porto Rico. Jour. Econ. Ent. 10: 513-517.
- 1923 Tachinids and sarcophagids established in Mexico. Jour. Econ. Ent. 16: 227.
- 1926 Insect enemies of sugarcane in western Mexico. Jour. Econ. Ent. 19: 664-669.
- 1943 A recent immigrant tachinid fly parasite of noctuid caterpillars in Hawaii. Hawaii. Planters' Rec. 47 (2): 81-87, figs. 1-4.
- 1944 *Eucelatoria armigera* (Coq.). Hawaii. Ent. Soc. Proc. 1943 12 (1): 4.
- 1944a *Eucelatoria armigera* (Coq.). Hawaii. Ent. Soc. Proc. 1943 12 (1): 12.
- 1945 *Eucelatoria armigera* (Coquillett). Hawaii. Ent. Soc. Proc. 1944 12 (2): 215.
- 1946 Recent immigrant insects. Hawaii. Planters' Rec. 50 (1): 11-17, figs. 1-4.
- 1946a *Polydesma umbricola* Boisduval. Hawaii. Ent. Soc. Proc. 1945 12 (3): 475-476.
- 1950 *Eucelatoria armigera* (Coquillett). Hawaii. Ent. Soc. Proc. 1949 14 (1): 7.
- 1951 The insects affecting sugar cane in Mexico. Internatl. Soc. Sugar Cane Technol. Cong. Proc. 7: 373-377.
- 1953 Insectos y otras plagas de la caña de azúcar. Banco de Méx., Indus. Azucarera de Méx. 2 (1): 269-315, 10 figs.
- 1955 *Laphygma exempta* (Walker). Hawaii. Ent. Soc. Proc. 1954 15 (3): 375.

Vazquez Garcia, L. (See Comstock, J. A., and Vazquez Garcia)

Vickery, R. A.

- 1915 Notes on three species of *Heliophila* which injure cereal and forage crops at Brownsville, Texas. Jour. Econ. Ent. 8: 389-392.
- 1924 The striped grass looper, *Mocis repanda* Fab., in Texas. Jour. Econ. Ent. 17: 401-406, pl. 5.

- 1925 List of parasitic insects reared from host insects collected in the vicinity of Brownsville, Texas. Ent. Soc. Wash. Proc. 27 (7): 137-141.
- 1926 Observations on *Cirphis latiuscula* H. Sch. in the Gulf coast region of Texas. Jour. Agr. Res. 32 (12): 1099-1119, figs. 1-3.
- 1929 Studies on the fall army worm in the Gulf coast district of Texas. U.S. Dept. Agr. Tech. Bul. 138: 1-63.
- Villeneuve, J.
- 1933 Description de *Aplomyiopsis galerucellae* n. gen., n. sp. (Tachinidae), parasite de *Galerucella luteola* (F. Müll.) en Amerique du Nord. Portici, R. Ist. Super. Agr., Lab. Zool. Gen. e Agr. Bol. 27: 125-126.
- Vinal, S. C., and Caffrey, D. J.
- 1919 The European corn borer and its control. Mass. Agr. Expt. Sta. Bul. 189: i-iv, 1-71, pls. 1-2.
- Vogel, E., and Oliver, A. D., Jr.
- 1969 Life history and some factors affecting the population of *Arzama densa* in Louisiana. Ent. Soc. Amer. Ann. 62 (4): 749-752, figs. 1-3.
- Vojdani, S.
- 1961 Bio-ecology of some *Eurygaster* species in central California (Pentatomidae-Scutellerinae). Ent. Soc. Amer. Ann. 54 (4): 567-578, figs. 1-6.
- Vosburgh, F. G.
- 1941 Our insect fifth column. Natl. Geo. Mag. 80 (2): 225-248, 14 figs., pls. 1-8.
- Wadley, F. M. (See Milliken, F. B., and Wadley)
- Wakeland, C. C.
- 1920 Variegated cutworm, *Peridroma saucia*, Hubner. Colo. State Ent. Cir. 28: 17-19.
- 1958 The high plains grasshopper, a compilation of facts about its occurrence and control. U.S. Dept. Agr. Tech. Bul. 1167: i-iv, 1-168, figs. 1-29.
- Walden, B. H.
- 1914 A lepidopterous leaf-folder on privet. *Archips rosana* Linn. Conn. Agr. Expt. Sta. Ann. Rpt. 1913 37: 223-226, pls. 3-4.
- Waldron, J. W. (See Paxton et al.)
- Walgenbach, D. D., and Benjamin, D. M.
- 1963 Natural controlling factors affecting the pine tussock moth. Ent. Soc. Amer., No. Cent. Branch, Proc. 18: 15-16.

Walkden, H. H.

- 1943 Cutworm and armyworm populations in pasture grasses, waste lands, and forage crops. Jour. Econ. Ent. 36: 376-381.

Walker, F. H., Jr.

- 1936 Observations on sunflower insects in Kansas. Kans. Ent. Soc. Jour. 9 (1): 16-25.

Walker, H. G. (See Gould, G. E., and Walker; Jones, D. W., Walker, and Anderson)

Walsh, B. D.

- 1861 Insects injurious to vegetation in Illinois. Ill. State Agr. Soc. Trans. (1859-1860) 4: 335-372, figs. 1-3, 2 pls.
- 1864 On the insects, coleopterous, hymenopterous and dipterous, inhabiting the galls of certain species of willow. Part 1st. — Diptera. Ent. Soc. Phila. Proc. 3: 543-644.
- 1866 Answers to correspondents. Geo. E. Brackett, Maine. Pract. Ent. 1 (6): 46.
- 1867 The three so-called army worms. Pract. Ent. 2 (11-12): 111-114.
- 1867a Answers to correspondents. John Edgerton, Iowa. Pract. Ent. 2 (11-12): 119.

— and Riley, C. V.

- 1868 Potato bugs [concl.]. Amer. Ent. 1 (3): 41-49, figs. 33-48.
- 1869 Tent-caterpillar of the forest. Amer. Ent. 1 (10): 208, fig. 146.
- 1869a The true army-worm (*Leucania unipunctata* [sic], Haworth). Amer. Ent. 1 (11): 214-217, figs. 152-155.

Walster, H. L.

- 1942 The battle against insects. N. Dak. Agr. Expt. Sta. Bimonthly Bul. 4 (4): 6-10.

Walton, G. S. (See Williams, C. B., Jr., Walton, and Tiernan)

Walton, W. R.

- 1908 Notes on the life history of *Nonagria oblonga* Gr. Ent. News 19 (7): 295-299, pl. 12.
- [1913] A new species of Tachinidae from Porto Rico. Ent. Soc. Wash. Proc. (1912) 14 (4): 198-200, pl. 10.
- [1913a] Meeting of October 3, 1912. [Comments by Mr. Walton on discussion of Mr. Caudell's paper]. Ent. Soc. Wash. Proc. 14 (4): 204.

- 1913b The variation of structural characters used in the classification of some muscoidean flies. Ent. Soc. Wash. Proc. 15 (1): 21-28.
- 1913c Efficiency of a tachinid parasite on the last instar of *Laphygma*. Ent. Soc. Wash. Proc. 15 (3): 128-131.
- 1913d New North American Tachinidae (Dipt.). Ent. News 24 (2): 49-52, pl. 3.
- 1914 A new tachinid parasite of *Diabrotica vittata*. Ent. Soc. Wash. Proc. 16 (1): 11-14, pl. 1.
- 1914a Four new species of Tachinidae from North America. Ent. Soc. Wash. Proc. 16 (2): 90-95, figs. 1-7.
- 1914b A new tachinid parasite of *Diapheromera femorata* Say. Ent. Soc. Wash. Proc. 16 (3): 129-132, figs. 1-6.
- 1915 A new and interesting genus of North American Tachinidae. Ent. Soc. Wash. Proc. 17 (3): 104-107, pl. 13.
- 1916 The true army worm and its control. U.S. Dept. Agr. Farmers' Bul. 731: 1-12, figs. 1-8.
- 1916a The tachinid genus *Argyrophylax* B. & B. Ent. Soc. Wash. Proc. 18 (3): 189-192, pl. 14.
- 1918 Three new tachinid parasites of *Eleodes*. Ent. Soc. Wash. Proc. (1917) 19 (1-4): 22-26, pl. 5.
- and Luginbill, P.
- 1916 The fall army worm, or "grass worm," and its control. U.S. Dept. Agr. Farmers' Bul. 752: 1-16, figs. 1-13.
- and Packard, C. M.
- 1940 The armyworm and its control. U.S. Dept. Agr. Farmers' Bul. 1850: 1-11, figs. 1-8.
- Walz, A. J. (See Manis et al.)
- *Warren, G. L.
- 1967 Newfoundland region. Canada Dept. Agr., Forest Insect and Dis. Survey, Ann. Rpt. 1966: 9-17.
- Warren, L. O., and Coyne, J. F.
- 1958 The pine sawfly, *Neodiprion taedae linearis* Ross, in Arkansas. Ark. Agr. Expt. Sta. Bul. 602: 1-23, figs. 1-6.
- Washburn, F. L.
- 1902 Insects notably injurious in 1902. Minn. State Expt. Sta., Ent. Ann. Rpt. 7: 1-74, figs. 1-62, 1 pl.

- 1905 The Diptera of Minnesota. Two-winged flies affecting the farm, garden, stock and household. Minn. Agr. Expt. Sta. Ann. Rpt. 10: 19-168, figs. 1-163, pls. 1-2.
- 1906 The cabbage maggot and other injurious insects of 1906. Minn. State Expt. Sta., Ent. Ann. Rpt. 11: i-vi, 1-87, figs. 1-64, 1 pl.
- 1910 1. Further observations on *Empoasca mali*;
2. Notes on *Papaipema nitela* and *P. cataphracta*. Jour. Econ. Ent. 3: 162-169.
- 1910a Further work upon the stalk borers. *Papaipema* sp. Minn. State Agr. Expt. Sta., Ent. Ann. Rpt. 13: 85-94, figs. 38-46.
- 1912 Grasshoppers, and other injurious insects of 1911 and 1912. Minn. State Expt. Sta., Ent. Ann. Rpt. 14: i-xiii, 1-114, figs. 1-73, pls. 1-3.
- *Watson, F. E.
1921 Miscellaneous notes and records of local Lepidoptera, with the description of a new form. N.Y. Ent. Soc. Jour. (1920) 28 (3-4): 227-235.
- Watson, J. R.
1916 Life-history of the velvet-bean caterpillar (*Anticarsia gemmatilis* [sic] Hübner). Jour. Econ. Ent. 9: 521-528, figs. 36-37, pls. 39-40.
- 1918 Insects of a citrus grove. Fla. Agr. Expt. Sta. Bul. 148: 165-267, figs. 74-142.
- 1921 Report of the entomologist. Fla. Agr. Expt. Sta. Rpt. 1921: 29R-33R.
- 1922 Report of entomologist. Fla. Agr. Expt. Sta. Rpt. 1922: 56R-59R.
- 1928 Entomology. Fla. Agr. Expt. Sta. Rpt. 1928: 42R-49R.
- 1934 Notes on *Nezara viridula* (L.). Fla. Ent. 18 (3): 43.
- 1935 Southern green stink bug (*Nezara viridula* L.). Insect Pest Survey Bul. 15 (5): 244.
- 1936 Southern green stinkbug (*Nezara viridula* L.). Insect Pest Survey Bul. 16 (9): 406.
- and Berger, E. W.
1932 Citrus insects and their control. Fla. Univ. Agr. Ext. Bul. 67: 1-140, figs. 1-90.
- Watts, J. G.
1938 Insect control studies. S. C. Agr. Expt. Sta. Clemson Agr. Col. Ann. Rpt. 51: 189-194.
- 1939 Asparagus beetle (*Crioceris asparagi* L.). Insect Pest Survey Bul. 19 (9): 566.

- Weast, R. D. (See Collins, M.M., and Weast)
- Weaver, [J. E.]
 1969 Tachina flies — West Virginia. (U.S. Dept. Agr.)
 Coop. Econ. Insect Rpt. 19 (6): 71.
- Weaver, J. E., and Sommers, R. A.
 1969 Life history and habits of the short-tailed cricket,
Anurogryllus muticus, in central Louisiana. Ent.
 Soc. Amer. Ann. 62 (2): 337-342, figs. 1-5.
- Webb, F. E. (See MacDonald and Webb)
- Webber, R. T. (See also Aldrich and Webber; Crossman and
 Webber)
 1930 A revision of the North American tachinid flies of
 the genus *Achaetoneura*. U.S. Natl. Mus. Proc. 78
 (10): 1-37, figs. 1-14.
 1931 A new parasitic fly of the genus *Chaetophlepsis*.
 U.S. Natl. Mus. Proc. 78 (20): 1-4.
 1932 *Sturmia inconspicua* Meigen, a tachinid parasite of
 the gipsy moth. Jour. Agr. Res. 45 (4): 193-208,
 figs. 1-6.
 1935 Satin moth. Insect Pest Survey Bul. 15 (10): 460.
 1937 Laboratory propagation of *Compsilura concinnata*
 Meigen. Jour. Econ. Ent. 30: 144-149.
 1941 Synopsis of the tachinid flies of the genus *Tach-*
inomyia, with descriptions of new species. U.S.
 Natl. Mus. Proc. 90(3108): 287-304, fig. 17.
- and Schaffner, J. V., Jr.
 1926 Host relations of *Compsilura concinnata* Meigen, an
 important tachinid parasite of the gipsy moth and
 the brown-tail moth. U.S. Dept. Agr., Dept. Bul.
 1363: 1-31.
- Weber, P. W.
 1953 Recent liberations of beneficial insects in Hawaii —
 II. Hawaii. Ent. Soc. Proc. 1952 15 (1):
 127-130.
- *Webster, F. M.
 1890 Report of observations upon insects affecting grains.
 U.S. Dept. Agr., Div. Ent. Bul. 22: 42-72, figs.
 1-2.
 1890a A case of excessive parasitism. Insect Life (U.S.
 Dept. Agr., Div. Ent., Period. Bul.) 2 (7-8):
 256-257.
 1898 Notes and observations on several species of Diptera.
 Canad. Ent. 30 (1): 18-19.

- 1899 Some notes on the grape-cane gall-maker, *Ampelolypter sesostris* (Coleoptera). Ent. News 10 (3): 53-55, pl. 3.
- 1899a Species of Diptera reared in Indiana during the years 1884 to 1890. Ind. Acad. Sci. Proc. 1898: 224.
- 1908 The value of insect parasitism to the American farmer. (U.S. Dept. Agr.) Yearb. 1907: 237-256, figs. 7-30.
- 1909 The lesser clover-leaf weevil. U.S. Dept. Agr. Bur. Ent. Bul., n. ser., 85 (1): 1-12, figs. 1-8.
- and Mally, C. W.
- 1897 Insects of the year in Ohio. U.S. Dept. Agr. Div. Ent. Bul., n. ser., 9: 40-46.
- Webster, R. L.
- 1911 Notes on the wheat-head army-worm (*Meliana albilinea* Hübner) as a timothy pest. Jour. Econ. Ent. 4: 179-184.
- Weed, C. M.
- 1897 Life histories of American insects. xii + 272 pp., 94 figs., 21 pls. New York.
- 1899 The spiny elm caterpillar. N.H. Col. Agr. Expt. Sta. Bul. 67: 124-141, figs. 40-51, 1 fig.
- and Conradi, A. F.
- 1902 The squash bug. N.H. Agr. Expt. Sta. Bul. 89: 13-28, figs. 1-2.
- Weeks, A. C.
- 1890 III. Utility of dragon flies as destroyers of mosquitoes. Pp. 69-95. In Lamborn, R. H., ed., Dragon flies vs. mosquitoes. Can the mosquito pest be mitigated? Studies in the life history of irritating insects, their natural enemies, and artificial checks, by working entomologists. 202 pp., 9 pls. New York.
- Weir, R.
- 1932 Report of the minister of agriculture for the Dominion of Canada for the year ended March 31, 1932. 225 pp. Ottawa.
- Weiss, H. B. (See also Dickerson and Weiss; Mutchler and Weiss)
- 1917 Additions to insects of New Jersey, No. 5. Ent. News 28 (5): 214-221.
- 1946 Status of parasites of the European corn borer in New Jersey. N.J. Dept. Agr. Ann. Rpt. 1945-1946 31: 152-153.

- 1947 Status of parasites of the European corn borer in New Jersey. N.J. Dept. Agr. Ann. Rpt. 1946-1947 32: 161.
- 1948 Parasites of the European corn borer. N.J. Dept. Agr. Ann. Rpt. 1947-1948 33: 170-171.
- 1949 Parasites of the European corn borer. N.J. Dept. Agr. Ann. Rpt. 1948-1949 34: 157.
- 1950 Parasites of the European corn borer. N.J. Dept. Agr. Ann. Rpt. 1949-1950 35: 169-170.
- 1951 Parasites of the European corn borer. N.J. Dept. Agr. Ann. Rpt. 1950-1951 36: 163.
- 1952 Parasites of the European corn borer. N.J. Dept. Agr. Ann. Rpt. 1951-1952 37: 181-182.
- 1953 Parasites of the European corn borer. N.J. Dept. Agr. Ann. Rpt. 1952-1953 38: 192.
- 1954 Parasites of the European corn borer. N.J. State Dept. Agr. Ann. Rpt. 39: 196.
- Welch, F. J.
- 1952 Effect of parasites on the European corn borer. Ky. Agr. Expt. Sta. Ann. Rpt. 1951 64: 38.
- 1953 Effect of parasites on the European corn borer. Ky. Agr. Expt. Sta. Ann. Rpt. 1952 65: 49.
- 1954 Effect of parasites on the European corn borer. Ky. Agr. Expt. Sta. Ann. Rpt. 1953 66: 54-55.
- [Wellhausen, E. J.]
- 1957 Cotaxtla Experiment Station, Veracruz. Pests of corn. Rockefeller Found., Mex. Agr. Program, Dir. Ann. Rpt. 1956-1957: 264-269.
- Wellso, S. G., and Hoxie, R. P.
- 1969 *Hyalomyodes triangulifer* parasitizing the cereal leaf beetle, *Oulema melanopus*. Ent. Soc. Amer. Ann. 62 (4): 923-924.
- West, L. S.
- 1923 Immunity to parasitism in *Samia cecropia* Linn. (Lep.: Saturniidae; Dip.: Tachinidae). Ent. News 34 (1): 23-25.
- 1925 New Phasiidae and Tachinidae from New York state. N.Y. Ent. Soc. Jour. 33: 121-135.
- 1928 Family Phasiidae, family Tachinidae. Family Megaprosopidae. Family Dexiidae. Pp. 805-823. In Leonard, M. D., ed., A list of the insects of New York with a list of the spiders and certain other allied groups. N.Y. (Cornell) Agr. Expt. Sta. Mem. 101: 1-1121, fig. 1.

Wheeler, E. H. (See Adams and Wheeler)

Whitcomb, W. D.

1940 Liberation of parasites of the European earwig. Mass. Agr. Expt. Sta. Bul. 369: 64.

1941 Parasites of the European earwig. Mass. Agr. Expt. Sta. Bul. 378: 71.

White, W. H.

1941 The Mexican bean beetle. Smithsn. Inst. Ann. Rpt. 1940: 343-356, fig. 1, pls. 1-6.

Whitehead, W. E.

1925 Notes on the currant leaf-roller (*Cacoecia rosana* Linn.) in Nova Scotia. Acadian Ent. Soc. Proc. (1924) 10: 76-79, pl. 2.

Whiteside, R. C., Burbutis, P. P., and Kelsey, L. P.

1967 Insect parasites of the green cloverworm in Delaware. Jour. Econ. Ent. 60: 326-328, figs. 1-3.

Whitlock, C.

1939 Abbreviations used in the Department of Agriculture for titles of publications. U.S. Dept. Agr. Misc. Pub. 337: 1-278.

Wickman, B. E.

1963 The large aspen tortrix, *Choristoneura conflictana*, in California (Lepidoptera: Tortricidae). Jour. Econ. Ent. 56: 593-596, figs. 1-3.

——— and Hunt, R. H.

1969 Biology of the phantom hemlock looper on Douglas-fir in California. Jour. Econ. Ent. 62: 1046-1050, figs. 1-5.

Wier, D. B.

1890 Parasites on *Datana ministra*. Insect Life (U.S. Dept. Agr., Div. Ent., Period. Bul.) 3 (1): 26-27.

Wilcox, J., Gray, K. W., and Mote, D. C.

1932 The strawberry crown moth. Oreg. Agr. Expt. Sta. Bul. 296: 1-30, figs. 1-11.

Wildermuth, V. L.

1911 The alfalfa caterpillar (*Eurymus eurytheme* Bdv.). U.S. Dept. Agr., Bur. Ent. Cir. 133: 1-14, figs. 1-8.

1914 The alfalfa caterpillar. U.S. Dept. Agr. Bul. 124: 1-40, figs. 1-20, pls. 1-2.

1920 The alfalfa caterpillar. U.S. Dept. Agr. Farmers' Bul. 1094: 1-16, figs. 1-14.

- and Caffrey, D. J.
1916 The New Mexico range caterpillar and its control.
U.S. Dept. Agr., Dept. Bul. 443: 1-12, figs. 1-12.
- and Frankenfeld, J. C.
1933 The New Mexico range caterpillar and its natural
control. Jour. Econ. Ent. 26: 794-798.
- Wilkes, A. (See also Boyes and Wilkes)
1946 The introduction of insect parasites of the spruce
budworm, *Archips fumiferana* Clem., into eastern
Canada. Canad. Ent. 78 (4): 82-86.
1949 Notes on the transfer of budworm parasites from
British Columbia to eastern Canada. Quebec Soc.
Protect. Plants Rpt. 1945-1947 30: 111-112.
- and Anderson, M.
1947 Notes on recovery of the introduced spruce bud-
worm parasite, *Phytodietus fumiferanae* Rohw. in
eastern Canada. Ontario Ent. Soc. Ann. Rpt.
77: 40-44.
- Coppel, H. C., and Mathers, W. G.
1949 Notes on the insect parasites of the spruce budworm
Choristoneura fumiferana (Clem.) in British
Columbia. Canad. Ent. (1948) 80 (1-12):
138-155, fig. 1.
- *Wilkinson, A. T. S.
1966 *Apanteles rubecula* Marsh and other parasites of
Pieris rapae in British Columbia. Jour. Econ. Ent.
59: 1012-1013.
- Wilkinson, R. C., Becker, G. C., [Jr.], and Benjamin, D. M.
1966 The biology of *Neodiprion rugifrons* (Hymen-
optera: Diprionidae), a sawfly infesting jack pine
in Wisconsin. Ent. Soc. Amer. Ann. 59
(4): 786-792, figs. 1-6.
- Williams, C. B., Jr., Walton, G. S., and Tiernan, C. F.
1969 Zectran and naled affect incidence of parasitism of
the budworm *Choristoneura occidentalis* in
Montana. Jour. Econ. Ent. 62: 310-312.
- *Williams, F. X. (See also Pemberton and Williams)
1931 Handbook of the insects and other invertebrates of
Hawaiian sugar cane fields. 400 pp., 190 figs.,
41 pls. Honolulu, Hawaii.
1935 *Hepialus pharus* Druce a moth borer attacking sugar
cane in Guatemala. Hawaii. Planters' Rec. 39
(4): 292-297, figs. 1-6, 2-3.

- 1946 *Polydesma umbricola* Boisduval. Hawaii. Ent. Soc. Proc. 1945 12 (3): 479.
- Williamson, G. D.
- 1961 Summary of parasite and predator liberations in Canada and of insect shipments from Canada in 1961. *Canad. Insect Pest Rev.* 39 (8): 257-268.
- 1962 Summary of parasite and predator liberations in Canada and of insect shipments from Canada in 1962. *Canad. Insect Pest Rev.* 40 (7): 147-158.
- 1963 Summary of parasite and predator liberations in Canada and of insect shipments from Canada in 1963. *Canad. Insect Pest Rev.* 41 (7): 137-151.
- 1968 Insect liberations in Canada. Parasites and predators 1967. 18 pp. Canada Dept. Agr., Res. Inst., Res. Branch, Belleville, Ont.
- 1969 Insect liberations in Canada — parasites and predators 1968. *Canada Dept. Agr. Liberation Bul.* 32: 1-19.
- Williston, S. W.
- 1889 The dipterous parasites of North American butterflies. Pp. 1912-1924, pl. 89. *In* Scudder, S. H., *Butterflies of the eastern United States and Canada with special reference to New England.* Vol. 3. Pp. i-vii, 1775-1958, frontis., pls. 1-89. Cambridge, Mass.
- 1893 List of Diptera of the Death Valley Expedition. Pp. 253-259. *In* Riley, C. V., *The Death Valley Expedition. A biological survey of parts of California, Nevada, Arizona, and Utah. Part II. 4. Report on a small collection of insects made during the Death Valley Expedition.* No. Amer. Fauna 7: 235-268.
- *Wilson, C. E.
- 1923 Insect pests of cotton in St. Croix and means of combating them. *Virgin Isl. Agr. Expt. Sta. Bul.* 3: 1-20, figs. 1-21.
- 1923a Truck-crop insect pests in the Virgin Islands and methods of combating them. *Virgin Isl. Agr. Expt. Sta. Bul.* 4: 1-35, figs. 1-24.
- Wilson, J. W. (See also Hayslip et al.; Ingram et al.)
- 1932 Notes on the biology of *Laphygma exigua* Huebner. *Fla. Ent.* 16 (3): 33-39.

- 1933 The biology of parasites and predators of *Laphygma exigua* Huebner reared during the season of 1932. Fla. Ent. 17 (1): 1-15, figs. 1-10.
- 1941 Biological control of *Diatraea saccharalis* in the Florida Everglades during 1940 and 1941. Fla. Ent. 24 (3): 52-57.
- 1942 Studies upon the prevalence and control of the sugarcane moth borer *Diatraea saccharalis* (F.). Fla. Agr. Expt. Sta. Ann. Rpt. 1942: 164.
- 1946 Studies upon the prevalence and control of the sugarcane moth borer. Fla. Agr. Expt. Sta. Ann. Rpt. 1946: 179.
- and Bregger, T.
- 1947 Studies upon the prevalence and control of the sugarcane moth borer. Fla. Agr. Expt. Sta. Ann. Rpt. 1947: 189.
- Wilson, M. C.
- 1948 Biological control of the European corn borer. Exotic parasites of European corn borer show rapid increase in Tipton County. Ind. Agr. Expt. Sta. Ann. Rpt. 61: 10-11, fig. 3.
- 1949 Biological control of the European corn borer. Another parasite recovered for the first time in Steuben, Kosciusko, and Jasper Counties. Ind. Agr. Expt. Sta. Ann. Rpt. 62: 83-84.
- 1949a The introduction of exotic species of European corn borer parasites in Indiana. Ind. Acad. Sci. Proc. 58: 157-159.
- 1950 Parasites of corn borer extending coverage. Ind. Agr. Expt. Sta. Ann. Rpt. 63: 11.
- 1951 The status of exotic European corn borer parasites in Indiana. Ind. Acad. Sci. Proc. 60: 222-225, fig. 1.
- 1951a Continue release of corn borer parasites. Ind. Agr. Expt. Sta. Ann. Rpt. 64: 41.
- 1952 European corn borer (*Pyrausta nubilalis*) — Indiana. (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 2 (12): 161.
- Winburn, T. F., and Painter, Reginald H.
- 1932 Insect enemies of the corn earworm (*Heliothis obsoleta* Fabr.). Kans. Ent. Soc. Jour. 5 (1): 1-28.
- Winn, A. F., and Beaulieu, G.
- 1915 A preliminary list of the insects of the Province of

Quebec. Part II — Diptera (two-winged flies).
Quebec Soc. Protect. Plants Rpt. 7 Sup.:
108-159.

Wishart, G.

- 1943 Important developments in the corn borer parasite situation. Ontario Ent. Soc. Ann. Rpt. 73: 26-30, fig. 1.
- 1944 An increase in the multiple generation of the European corn borer in Ontario and its relation to parasite establishment. Ontario Ent. Soc. Ann. Rpt. 74: 11-13.
- 1945 *Aplomya caesar* (Aldrich), a tachinid parasite of the European corn borer. Canad. Ent. 77 (9): 157-167, figs. 1-14.
- 1950 Biological control of the European corn borer. Quebec Soc. Protect. Plants Ann. Rpt. 31: 48-49.

— and Thomas, I. E.

- 1933 Recent developments in the corn borer parasite situation in eastern Canada. Ontario Ent. Soc. Ann. Rpt. 1932 63: 39-41.

Wolcott, G. N. (See also Jones, T. H., and Wolcott)


- 1915 Report of the traveling entomologist. Porto Rico Bd. Comnr. Agr. Rpt. 3: 25-40.
- 1916 Report of the entomologist. Porto Rico Bd. Comnr. Agr. Rpt. 4: 17-22.
- 1922 Insect parasite introduction in Porto Rico. Porto Rico Dept. Agr. Jour. 6 (1): 5-20, figs. 1-7.
- 1924 "Insectae Portoricensis." A preliminary annotated check-list of the insects of Porto Rico, with descriptions of some new species. Porto Rico Dept. Agr. Jour. (1923) 7 (1): 1-313, pls. 1-2.
- 1924a Entomología económica Puertorriqueña. Puerto Rico Estac. Expt. Insular (Rio Piedras) Bul. 32: 1-176, figs. 1-98, pls. 1-12.
- 1933 [Parasitism of *Phyllophaga portoricensis* Smyth by *Cryptomeigenia aurifacies* Walton]. Insect Pest Survey Bul. 13 (5): 186.
- 1935 The white grub problem in Puerto Rico. Internatl. Soc. Sugar Cane Technol. Cong. Proc. 5: 445-456, 5 figs.
- 1940 To rear large numbers of the caterpillars of *Diatraea saccharalis* F., as hosts for the Amazon fly. Puerto Rico Univ., Agr. Expt. Sta. Ann. Rpt. 1939-1940: 31-32.

- 1941 A supplement to "Insectae Borinquenses". Puerto Rico Univ. Jour. Agr. 25 (2): 33-158.
- 1951 The present status of economic entomology in Puerto Rico. Puerto Rico Agr. Expt. Sta. (Rio Piedras) Bul. 99: 1-21, figs. 1-7.
- 1951a The insects of Puerto Rico. Puerto Rico Univ. Jour. Agr. (1948) 32 (3): 417-748, 79 figs.
- 1956 Insect conditions in Puerto Rico. (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 6 (15): 311.
- Wolfenbarger, D. O. (See Rice and Wolfenbarger; Rice, Wolfenbarger, and Chada)
- *Wong
1969 Southern green stink bug (*Nezara viridula*). (U.S. Dept. Agr.) Coop. Econ. Insect Rpt. 19 (13): 217.
- Wong, H. R., and Melvin, J. C. E.
1967 The leaf roller *Pseudexentera oregonana* Wlsh. Canada Dept. Forestry and Rural Devlpmt., Bi-monthly Res. Notes 23 (1): 3-4.
- Wood, A. A. (See Hudson and Wood)
- *Wood, G. W.
1951 An annotated list of lepidopterous larvae from commercial blueberry fields, Charlotte County, N.B. Canad. Ent. 83 (9): 241-244.
- and Neilson, W. T. A.
1957 Notes on life-histories of four species of climbing cutworms collected from low-bush blueberry fields in New Brunswick (Lepidoptera: Phalaenidae). Canad. Ent. 89 (11): 502-506.
- Neilson, W. T. A., Maxwell, C. W., and McKiel, J. A.
1954 Life-history studies of *Spaelotis clandestina* (Harr.) and *Polia purpurissata* (Grt.) (Lepidoptera: Phalaenidae) in low-bush blueberry areas in New Brunswick. Canad. Ent. 86 (4): 169-173.
- Wood, J. G.
1874 Insects abroad. [A companion volume to "Insects at home."] Being a popular account of foreign insects, their structure, habits, and transformations. xii + 780 pp., 520 figs., 20 pls. London.
- Wood, W. B.
1918 The oriental peach moth: a Japanese fruit insect recently introduced into the United States. Calif. State Comn. Hort. Monthly Bul. 7 (9): 523-528, figs. 76-83.

- and Selkregg, E. R.
 1918 Further notes on *Laspeyresia molesta*. Jour. Agr. Res. 13 (1): 59-72, pls. 5-10.
- Woodard, J. S.
 1925 Pecan insects. Tex. Dept. Agr. Bul. 81: 165-183, figs. 1-13 (including two fig. 2's).
- *Woods, C. D.
 1907 Newspaper bulletins and circulars published in 1906. Maine Agr. Expt. Sta. Ann. Rpt. 1906 22: 283-304, figs. 4 [sic], 25-46.
- Woods, W. C.
 1917 The biology of the alder flea-beetle, *Altica bimarginata* Say. Maine Agr. Expt. Sta. Bul. 265: 249-284, figs. 18-31
 1918 The biology of Maine species of *Altica*. Maine Agr. Expt. Sta. Bul. 273: 149-204, figs. 10-13.
- Worthley, H. N. (See also Steiner and Worthley)
 1923 The squash bug in Massachusetts. Jour. Econ. Ent. 16: 73-79, figs. 2-3.
 1924 The biology of *Trichopoda pennipes* Fab. (Diptera, Tachinidae), a parasite of the common squash bug. Part I. Introduction. Psyche 31 (1): 7-16.
 1924a The biology of *Trichopoda pennipes* Fab. (Diptera, Tachinidae), a parasite of the common squash bug. Part II. Morphology. Psyche 31 (2): 57-77, pls. 1-4.
- Worthley, L. H. (See Caffrey and Worthley)
- Wray, D. L.
 1950 Insects of North Carolina. Second supplement. 59 pp. N.C. Dept. Agr., Raleigh.
- Wressell, H. B. (See also Manson et al.)
- Driscoll, and Svec
 1959 European corn borer (*Ostrinia nubilalis* Hbn.). Canad. Insect Pest Rev. 37 (7): 240.
- Wright, J. M. (See also Arbuthnot and Wright)
 1947 Present status of European corn borer parasites in the north central states, and reports from Indiana, Illinois, and Iowa (b). Ent. Soc. Amer., No. Cent. Branch, Proc. 2: 33.
- Wylie, W. D.
 1946 Insects affecting sweet potatoes in the Everglades. Fla. State Hort. Soc. Proc. 59: 87-91.

- and Tadic, M.
1963 Two tortricids and their parasites in Arkansas. *Kans. Ent. Soc. Jour.* 36 (3): 165-166, 1 fig.
- Yackley, [W. H.] (See Strohecker and Yackley)
- Yamamoto, [H. Y.]
1969 Southern green stink bug (*Nezara viridula*). (U.S. Dept. Agr.) *Coop. Econ. Insect Rpt.* 19 (17): 301.
- Yonke, T. R., and Medler, J. T.
1969 Biology of the Coreidae in Wisconsin. *Wis. Acad. Sci., Arts and Letters, Trans.* 57: 163-188, figs. 1-7.
- York, G. T. (See also Jarvis and York; Van Denburgh, Burbutis, and York)
1955 Present status of parasites of the European corn borer. *Ent. Soc. Amer., No. Cent. Branch, Proc.* 10: 65.
- Schaffner, J. C., and Brindley, T. A.
1955 Parasites of the European corn borer found infesting the stalk borer. *Jour. Econ. Ent.* 48: 765-766.
- Yoshioka, [E.] (See also Mitchell and Yoshioka)
1968 Southern green stink bug (*Nezara viridula*). (U.S. Dept. Agr.) *Coop. Econ. Insect Rpt.* 18 (28): 652.
- Yoshioka, [E.], and Miyahira, [N.]
1966 A tachina fly (*Trichopoda pennipes*). (U.S. Dept. Agr.) *Coop. Econ. Insect Rpt.* 16 (12): 231.
1966a Southern green stink bug (*Nezara viridula* var. *smaragdula*). (U.S. Dept. Agr.) *Coop. Econ. Insect Rpt.* 16 (20): 444.
- Yothers, M. A., and Van Leeuwen, E. R.
1931 Life history of the codling moth in the Rogue River Valley of Oregon. *U.S. Dept. Agr. Tech. Bul.* 255: 1-34, figs. 1-18.
- Young, J. H. (See Bottrell et al.)
- Young, W. R. (See Sifuentes and Young)
- Zappe, M. P. (See Britton and Zappe)
- Zeller, S. M., and Schuh, J.
1944 Diseases and insect pests of cane fruits in Oregon. *Oreg. Agr. Expt. Sta. Bul.* 418: 1-58, figs. 1-27, cover fig.
- Zimmerman, E. C.
1943 *Eucelatoria armigera* (Coq.). *Hawaii. Ent. Soc. Proc.* 1942 11 (3): 267-268.

- 1958 Insects of Hawaii. A manual of the insects of the Hawaiian Islands, including an enumeration of the species and notes on their origin, distribution, hosts, parasites, etc. Vol. 7. Macrolepidoptera. ix + 542 pp., 423 figs. Honolulu, Hawaii.
- and Fletcher, D. S.
- 1956 *Heliothis* in Hawaii (Lepidoptera: Phalaenidae). Hawaii Ent. Soc. Proc. 1955 16 (1): 170-176.



**U. S. DEPARTMENT OF AGRICULTURE
SCIENCE AND EDUCATION
ADMINISTRATION
HYATTSVILLE, MARYLAND 20782**

**OFFICIAL BUSINESS
PENALTY FOR PRIVATE USE, \$300**

**POSTAGE AND FEES PAID
U. S. DEPARTMENT OF
AGRICULTURE
AGR 101**

