Historic, Archive Document

Do not assume content reflects current scientific knowledge, policies, or practices.

d States Department of Agriculture

6

Soil Conservation Service

Bozeman, Montana



MONTANA WATER SUPPLY OUTLOOK

April 1, 1986



	Foreword								
How Forecasts Are Made	Most of the annual s lates high in the mo estimate the runoff t water equivalent at so data are viewed in o comprehensive pict includes selected st data and narratives	treamflow in the Western United States originates as snowfall. This snowfall accumu- buntains during winter and early spring. As the snowpack accumulates, hydrologists hat will occur when it melts. Predictions are based on careful measurements of snow elected index points. Precipitation, temperature, soil moisture and antecedent streamflow conjunction with snowpack data to prepare runoff forecasts. This report presents a ure of water supply outlook conditions for areas dependent upon surface runoff. It reamflow forecasts, summarized snowpack and precipitation data, reservoir storage describing current conditions.							
	Streamflow forecasi Service hydrologists For this reason, fore Average, and Above N maximum. Actual st out of ten years.	ts are cooperatively generated by Soil Conservation Service and National Weather a. Forecasts become more accurate as more data affecting runoff becomes known. accasts are issued that refiect three future precipitation conditions — Below Normal, Jormal. These forecasts are termed reasonable minimum. most probable, and reasonable reamflow can be expected to fall between the lower and upper forecast values eight							
	Snowpack data are Manual readings of monthly or semi-mo temperature, and ot to central data colle	obtained by using a combination of manual and automated measurement methods. snow depth and water equivalent are taken at locations called snow courses on a inthly schedule during the winter. In addition, snow water equivalent, precipitation, her parameters are monitored on a daily basis and transmitted via radio telemetry action facilities. Both monthly and daily data are used to project snowmelt runoff.							
For More Information	Copies of Monthly Water Supply Outlook Reports and other reports may be obtained from the states listed below. Because of the limited space, snow survey measurements are not published in monthly reports. An annual snow survey data summary is published by the Soil Conservation Service for each of the western states. Historical snow survey data may be obtained at those same offices.								
	STATE	ADDRESS							
	Alaska	201 East 9th Ave., Suite 300, Anchorage, AK 99501-3687							
	Arizona	201 East Indianola, Suite 200, Phoenix, AZ 85012							
	Colorado (New Mexico)	2490 West 26th Ave., Denver, CO 80211							
	Idaho	304 North 8th Street, Room 345, Boise, ID 83702							
	Montana	10 East Babcock, Room 443, Federal Building, Bozeman, MT 59715							
	Nevada	50 South Virginia Street, Third Floor, Reno, NV 89505							
	Oregon	1220 Southwest 3rd Ave., 16th Floor, Portland, OR 97204							
	Utah	4402 Federal Building, 125 South State Street, Salt Lake City, UT 84147							
	Washington	360 U.S. Court House, Spokane, WA 99201							
	Wyoming	Federal Building, 100 East "B" Street, Casper, WY 82602							
	In addition to state Soil Conservation So obtained from the S Room 547, Portland	reports, a Water Supply Outlook for the Western United States is published by the ervice and National Weather Service monthly, January through May. Reports may be tool Conservation Service, West National Technical Center, 511 Northwest Broadway, , OR 97209.							

Published by other agencies:

2 (n) 21

Water Supply Outlook Reports prepared by other agencies include: California — Snow Survey Branch, California Department of Water Resources, P.O. Box 388, Sacramento, CA 98502; British Columbia — The Ministry of Environment, Water Investigations Branch, Parliament Buildings, Victoria, British Columbia, V8V 1X5; Yukon Territory — Department of Indian and Northern Affairs, Northern Operations Branch, 200 Range Road, Whitehorse, Yukon Territory, Y1A 3V1; Alberta, Saskatchewan, and N.W.T. — The Water Survey of Canada, Inland Waters Branch, 110-12 Avenue S.W., Calgary, Alberta, T3C 1A6.

Montana Water Supply Outlook

and

Federal – State – Private Cooperative Snow Surveys

Issued by

Wilson Scaling Chief Soil Conservation Service Washington, D.C.

Released by

Glen H. Loomis State Conservationist Soil Conservation Service Bozeman, Montana

Prepared by

Phillip E. Farnes Snow Survey Supervisor Soil Conservation Service 10 E. Babcock Bozeman, Montana 59715

Programs and assistance of the United States Department of Agriculture are available without regard to race, creed, color, sex, age, or national origin.



2

SDA SCS FORT WOR

GENERAL OUTLOOK

SUMMARY:

Below average March precipitation and melt generated by warm temperatures have reduced the snowpack levels reported on March 1. Snowpacks in extreme southwest Montana and near the Montana-Wyoming border are near normal, but all other areas have below to well below average snow cover. Generally, the northern areas have the poorest snowpack. Many areas have less snowpack now than was measured a year ago. Temperatures and melt conditions seem to be about a month earlier than normal. Streamflows are forecast to be near to a little below average in the Jefferson, Madison and Yellowstone River drainages but below to well below average in other areas. Some streams with lower elevation headwaters have already reached their peak snowmelt runoff. Widespread irrigation water shortages are expected by late June to early July over most of the state for irrigators not having stored water.

SNOWPACK:

Snowpack levels are about 10 percent less than reported on March 1. Warm temperatures created melt at low and mid-elevations. Also, mountain precipitation was below average in all areas. Snowpack is well below average in northern areas increasing to below average through most of central Montana. The only areas reporting near average snowpack are along the Continental Divide from southwest of Helena to Yellowstone National Park, throughout most of the Yellowstone River headwaters and in the headwaters of the Clarks Fork of the Yellowstone, Bighorn, Little Bighorn, Tongue and Powder Rivers in Wyoming.

PRECIPITATION:

Mountain precipitation during March was generally about 60 to 70 percent of average over most of the state. The lower Clark Fork area, west of Missoula, was a little better but still only around 90 percent of average. The St. Mary and Milk River headwaters also reported about 80 percent of average moisture in March. Many valley locations recorded well below average precipitation for the month. If the weather patterns do not improve, Montana can expect another dry spring and summer. For some areas, this could be the fifth consecutive year of below average precipitation.

RESERVOIRS:

Most irrigation reservoirs across the state have average or above average storage due to good carryover from August and September rains and early season runoff that started in late February. Storage in most large and multipurpose reservoirs is near or above average.

STREAMFLOW:

Except for average or above average runoff from streams with headwaters in Wyoming, below average streamflows are forecast for all streams and rivers in Montana. Well below average runoff is expected from streams in the Gallatin Valley and most streams in northwest and north central Montana. Except for areas in extreme southwest Montana and near the Montana-Wyoming border, most areas can expect shortages of irrigation water supplies by late June to early July. If above normal temperatures continue, runoff will occur earlier than usual and will create additional water shortage problems during the main irrigation season.

Kootenai Basin



WATER SUPPLY OUTLOOK:

Snowpack conditions deteriorated in March as a result of below average mountain precipitation and melt created by warm temperatures. Snowpack is better in British Columbia than in Montana. Streamflow on the Kootenai River is forecast to be below average while tributary streams in Montana are predicted to have well below average runoff for the spring and summer months. Some smaller streams with low elevation headwaters may have already reached their peak snowmelt runoff.

FORECAST POINT	FORECAST	20 YR. AVE.	MOST PROBABLE	MOST PROBABLE	REAS. MAX.	REAS. MIN.	PEAK FLOW	PEAK	LOH FLOH	LOH
	PERIOD	(1000AF)	(1000AF)	(% AVE.)	(% AVE.)	(% AVE.)	(CFS)	DATE	(CFS)	DATE
KOOTENAI RIVER blw Libby Dam #	APR-JUL	6020.0	5310.0	88	110	66				
	APR-SEP	7041.0	6210.0	88	110	66				
FISHER RIVER near Libby	APR-JUL	248.0	162.0	65	90	41				
	AFR-SEP	264.0	174.0	65	90	42				
YAAK RIVER near Troy	APR-JUL	500.0	345.0	69	93	45				
	APR-SEP	523.0	373.0	71	95	47				
KOOTENAI RIVER at Leonia *	APR-JUL	7498.0	6220.0	82	103	63				
	APR-SEP	8602.0	7130.0	82	103	63				
	APR-JUN	6051.0	4930.0	81	101	61				

STREAMFLOW FORECASTS

1 RESERVOIR STORAGE (1000AF) WATERSHED SNOWPACK ANALYSIS 1 -----!-----! USEABLE I ** USEABLE STORAGE ** I THIS YEAR AS % OF NO. CAPACITYI THIS LAST I WATERSHED I YEAR YEAR AVE.I COURSES ------RESERVOIR -----_____ ------5748.0 2238.0 1801.0 1694.0 | EAST KODTENAI in B.C. 29 100 83 LAKE KOOCANUSA KOOTENAI in MONTANA 31 67 65 1 KODTENAI ab BONNERS FERRY 60 78 71 1 Т

*Corrected for upstream diversions or changes in reservoir storage.

Average is for 1961-80 period.

Flathead Basin

Mountain snowpack* (inches)



Precipitation* (percent of normal)

WATER SUPPLY OUTLOOK:

Below average mountain precipitation and warmer temperatures during March have lowered snowpack percentages. Some higher elevations have fair snowpack but most areas including lower elevations have well below average amounts of snow cover. Spring and summer streamflows are forecast to be below average. Some low elevation streams have already had their peak snowmelt runoff.

FORECAST POINT	FORECAST	20 YR. AVE.	MOST PROBABLE	HOST FROBAELE	REAS. MAX.	REAS. HIN.	PEAK FLOW	PEAK	LOW FLOW	LOW
	PERI00	(1000AF)	(1000AF)	(% AVE.)	(% AVE.)	(% AVE.)	(CFS)	OATE	(CFS)	DATE
NE ELATHEAN moon Columbia Falle	APR 111	1732.0	1300.0	75	89	61				
AF FEMINENO NEAL COTOMDIA FAIIS	APR-SEP	1913.0	1440.0	75	89	61				
	APR-JUN	1471.0	1120.0	76	90	62				
F FLATHFAD open West Glacier	APREJUL	1713.0	1410.0	82	96	68				
	APR-SEP	1867.0	1510.0	80	95	67				
	APR-JUN	1453.0	1220.0	83	98	70			,	
F FLATHEAD near Columbia Falls ¥	APR-JUL	2142.0	1750.0	81	100	64				
	APR-SEP	2278.0	1870.0	82	101	63				
	APR-JUN	1886.0	1550.0	82	100	64				
LATHEAD at Columbia Falls *	ARP-JUL	5721.0	4600.0	80	94	66				
	APR-SEP	6208.0	4950.0	79	94	66				
	APR-JUN	4921.0	4020.0	81	96	68				
WAN RIVER near Big Fork	APR-JUL	604.0	530.0	87	102	74				
	APR-SEP	689.0	600.0	87	101	73				
LATHEAD RIVER near Polson ¥	APR-JUL	6712.0	5400.0	80	94	66				
	APR-SEP	7278.0	5830.0	80	94	66				
	APR-JUN	5759.0	4685.0	81	95	67				

	RESERVOIR STORAGE		(1000AF)		WATERSHED SNOWPACK ANALYSIS					
ŔESERVOIŔ	USEABLE I CAPACITYI I	** US THIS YEAR	EABLE STO LAST YEAR	AGE ** AVE.	HATERSHED	NO, COURSES AVE,O	THIS Y	EAR AS % OF R. AVERAGE		
CAMAS (4)	45.2	31.3	18.0	23.1	NORTH FORK FLATHEAO	16	69	65		
MISSION VALLEY (8)	100.0	50.3	37.6	41.1	MIDDLE FORK FLATHEAD	12	78	73		
HUNGRY HORSE	3451.0	2515.0	1796.0	2054.0	SOUTH FORK FLATHEAO	13	73	70		
FLATHEAO LAKE	1791.0	805.3	649.3	762.0	STILLWATER-WHITEFISH	9	73	65		
					SHAN	11	79	76		
					LITTLE BITTERROOT	9	61	66		
				l	FLATHEAO	50	73	70		

STREAMFLOW FORECASTS

Clark Fork Basin above Missoula



WATER SUPPLY OUTLOOK:

Snowpack conditions deteriorated in March. Below average mountain precipitation and above average melt dropped the snowpack 5 to 10 percent since March 1. The mountains around Butte and Anaconda have a little better snowpack than other areas. The Blackfoot has less snow than a year ago while the Clark Fork has about the same. Spring and summer runoff is expected to be 15 to 20 percent below average on most streams. Shortages of irrigation water supplies can be expected by late June or early July.

	FORECAST	20 YR.	MOST	HOST	REAS.	REAS.	PEAK	PEAF	LOW	LOW
FURELASI FUINI	PERIOD	(1000AF)	(1000AF)	(% AVE.)	(% AVE.)	(% AVE.)	(CFS)	DATE	CFS)	DATE
OULTON RESERVOIR Inflow (MG)*	APR-JUL	263.0	215.0	81	106	58				
	APR-JUN	237.0	195.0	82	106	58				
ARM SPRINGS CR at Meyers Dam ×	APR-JUL	37.8	32.0	84	108	61				
	APR-SEP	46.8	39.7	84	109	60				
LINT CREEK near Southern Cross I	APR-JUL	15.4	12.9	83	117	52				
	APR-SEP	18.3	15.3	83	120	49				
LINT CREEK below Boulder Creek #	APR-JUL	59.9	48.5	80	115	47				
	APR-SEP	75.8	. 61.5	81	115	47				
DWER WILLOW CR RES Inflow *	APR-JUL	14.9	10.8	72	107	40				
	APR-SEP	15.7	11.5	73	108	38				
• FK. ROCK CRK near Philipsburg	APR-JUL	70.5	61.7	87	112	64				
	APR-SEP	78.2	68+3	87	111	64				
EVADA CREEK near Finn	APR-JUL	21.3	14.8	69	103	38				
	APR-SEP	23.0	16.0	69	104	35				
LACKFOOT RIVER near Bonner	APR-JUL	904.0	658.0	72	87	59				
	APR-SEP	999.0	745.0	74	89	61				
	APR-JUN	782.0	565.0	72	86	58				
LARK FORK RIVER above Milltown *	APR-JUL	708.0	555.0	78	108	48				
•	APR-SEP	816.0	644.0	78	109	49				
	APR-JUN	597.0	470.0	78	109	49				
LARK FORK RIVER above Missoula	APR-JUL	1612.0	1210.0	75	99	51				
	APR-SEP	1815.0	1400.0	77	101	53				
	APR-JUN	1379.0	1040.0	75	99	51				

STREAMFLOW	FORECASTS
------------	-----------

	RESERVOIR STORAGE	(1000AF) I			I I WATERSHED SNOWPACK ANALYSIS I					
RESERVOIR	USEABLE CAPACITY 	XX USE THIS YEAR	ABLE STORA LAST YEAR	AGE XX I AVE. I	WATERSHED	ND. COURSES AVE.0	THIS YEA	R AS % OF AVERAGE		
GEORGETOWN LAKE	31.0	25.4	25.8	23.7	CLARK FORK ab BLACKFOOT	46	99	85		
LOWER WILLOW CREEK	4.9	4.9	1.3	1.8	BLACKFOOT	22	78	66		
NEVADA CREEK	12.6	11.0		7.4 I 1	CLARK FORK above MISSOULA	62	93	79		

≭Corrected for upstream diversions or changes in reservoir storage. Average is for 1961-80 period.

Clark Fork Basin below Missoula



Precipitation* (percent of normal)

WATER SUPPLY OUTLOOK:

Snowpack percentages have dropped about 10 percent since March 1. This is a result of below average mountain precipitation and melt caused by warm temperatures during March. There is less water stored in the snowpack than there was last year at Spring and summer streamflows are this time. forecast to be below average in all drainages. Shortages of irrigation water can be expected by late June to early July.

	FORFRACE	20 VD	HOCT	NOCT	5.5 A.C	D.C.A.C	E.E.A.V	E.E.A.V		
FORECACT DOTNE	FORECAST	20 YR.	PDODADLE	MUST EEOEADLE	REAS.	REAS.	PEAK	FEAF	LOW	LOn
FURELAST PUINT	PERIOO	(1000AF)	(1000AF)	(% AVE.)	(% AVE.)	(% AVE.)	(CFS)	OATE	(CFS)	OATE
	ADD 110	1/12 0	1210.0	75		F.				
TAKK FORK KINER SDONE HISSOUIS	APK-JUL	1012+0	1210.0	75	101	51				
	APE-JUN	1379.0	1040.0	75	99	51				
	HIN OOK	10/ / 10	104010	/0	<i>,,</i>	51				
.F. BITTERROOT RIVER or Conner #	APR-JUL	164.0	135.0	82	106	59				
	APR-SEP	178.0	145.0	81	106	57				
	45.5. 11.11	F 2 2 4		05						
ITTEFROOT FIVER near Uarby	APR-JUL	532.0	455.0	85	110	61				
	APR-SEP	580.0	490.0	84	108	61				
	APR-JUN	464.0	400.0	86	110	62				
KALKAHD CREEK pear Hamilton	APR-JU	48.7	43.2	88	103	74				
	APR-SEP	56.0	49.5	88	102	75				
URNT FORK CR nr Stevensville 🗷	AF:R-JUL	32.2	27.7	86	109	62				
	APR-SEP	37.4	32.0	85	110	61				
TITEREONT RIVER of Miscoulo B		1384.0	1165.0	84	108	60				
THERROOM RIVER at HISSOUR -	APE-SEP	1504.0	1260.0	83	108	60				
	APR-JUN	1191.0	1010.0	84	109	61				
				•						
LARK FORK RIVER below Missoula	APR-JUL	2996.0	2375.0	79	95	63				
	APR-SEP	3319.0	2650.0	79	96	64				
	APR-JUN	2570.0	2050.0	79	96	64				
	455 88	2020 0	21.00	70	101					
LARK FURK RIVER at St. Regis	APR-JUL	3928.0	3100.0	/8	101	57				
	APR-SEP	4411.0	3480.0	/8	101	57				
	APK-JUN	3428+0	2/10.0	/4	101	3/				
APK FORK RIVER near Plains #	APR-JUL	11071.0	8450.0	76	91	61				
	APR-SEP	12153.0	9280.0	76	91	61				
	APR-JUN	9459.0	7050.0	74	90	60				
	400 111	222.0	457 0	. 7		45				
numrouw KIVEK near Inompson Falls	APR-JUL	233.0	15/10	6/	89	40				
	AFK-SEF	261.0	180.0	68	91	4/				
ROSPECT CREEK at Thompson Falls	APR-JUI	132.0	100.0	75	100	52				
	APR-SEP	142.0	110.0	77	101	54				
	CAR OL	17210		, ,	1.11					
LARK FORK at Whitehorse Rapids #	APR-JUL	12351.0	9370.0	75	92	60				
	APR-SEP	13575.0	10300.0	75	92	60				
	ADD HIM	10570 0	0025 0	75	02	40				

	RESERVOIR STORAGE	(1000AF)			WATERSHED SN	SNOWPACK ANALYSIS			
RESERVOIR	USEABLE CAPACITY	** US THIS YEAR	EABLE STOR LAST YEAR	AGE II I	WATERSHED	ND. COURSES AVE.0	THIS YEAR	AS % OF	
PAINTED ROCKS LAKE		NO REPO) f: T		CLARK FORK above MISSOULA	62	93	79	
NOXON RAPIOS	335.0	299.8	156.2	197.6	BITTERRODT	22	93	80	
COHO	34.9	23.0	10.1	14.6	LWR CLARK FK blw MISSOULA	20	86	85	
					EITTERROOT & LWR C.F.	41	89	84	
					CLARK FORK TOTAL	97	¢٥	81	
					FLATHEAO	50	73	70	
					PENO D'REILLE	141	83	77	

STREAMFLOW FORECASTS

#Corrected for upstream diversions or changes in reservoir storage. Average is for 1961-80 period.

Jefferson Basin

Precipitation* (percent of normal)

Mountain snowpack* (inches)



WATER SUPPLY OUTLOOK:

Snowpack in the Beaverhead and upper Big Hole is near to a little below average and a little below average in the lower Big Hole, Ruby and Boulder headwaters. Melt and below average mountain precipitation have decreased the snowpack percentages about 10 percent since March 1. Streamflow for the spring and summer is forecast to be a little below average for most drainages. Irrigation water supplies should be near to a little below average for most streams.

JEFFERSON RIVER BASIN

FORECAST POINT	FORECAST	20 YR. AVE.	MOST PROBABLE	MOST PROBAGLE	REAS. MAX.	REAS. MIN.	PEAK FLOW	PEAK	LO¥ FLO¥	LOW
	PERIOD	(1000AF)	(1000AF)	(% AVE.)	(% AVE.)	(% AVE.)	(CFS)	DATE	(CFS)	DATE
RED ROCK RIVER near Monida *	APR-JUL	96.0	87.0	90	121	60				
	APR-SEP	103.0	93.0	90	120	60				
BEAVERHEAD RIVER near Grant #	APR-JUL	137.0	128.0	93	123	64				
	APR-SEP	158.0	142.0	89	120	60				
BEAVERHEAD RIVER at Barratts 🛎	APR-JUL	180.0	165.0	91	122	62				
	APR-SEP	209.0	190.0	90	121	61				
RUBY RIVER near Alder	APR-JUL	85.0	77.0	90	115	66				
	APR-SEP	101.0	91.0	90	115	65				
GIG HOLE RIVER near Melrose	APR-JUL	698.0	655.0	93	119	69				
	APR-SEP	760.0	705.0	92	118	68				
HILLOW CREEK near Harrison	APR-JUL	17.9	17.2	96	128	67				
	APR-SEP	20.0	19.3	96	125	65				

STREAMFLOW FORECASTS

	RESERVOIR STORAGE	 (1000AF) 			WATERS	WATERSHED SNOWPACK ANALYSIS			
RESERVOIR	USEABLE I CAPACITYI I	** USI THIS YEAR	EAGLE STOR LAST YEAR	AGE ## I	HATERSHED	NO. COURSES AVE.D	THIS YE	AR AS % OF	
LIMA	84.0	29.2	31.9	38.0 1	BEAVERHEAD	32	110	96	
CLARK CANYON	257.0	158.3	151.8	147.6	RUBY	13	103	85	
RUBY RIVER	38.8	34.0	33.3	30.3	BIGHOLE	29	105	91	
				1	BOULDER	15	98	88	
					JEFFERSON	71	105	92	

≭Corrected for upstream diversions or changes in reservoir storage. Average is for 1961-80 period.

Madison Basin



Mountain snowpack* (inches)

WATER SUPPLY OUTLOOK:

The snowpack is about 10 percent less than recorded During March, the mountain precipitation on March 1. was below average and melt was caused by warm temperatures. Spring and summer streamflows are forecast to be above average upstream for Hebgen Downstream, runoff from tributary streams is Lake, predicted to be below average. Some late season irrigation shortages can be expected along these smaller streams.

For more information contact your local Soil Conservation Service office.

135;

STREAMFLOW FORECASTS

FORECAST POINT	FORECAST PERIOD	20 YR. AVE. (1000AF)	NOST PROBABLE (1000AF)	MOST PROBABLE (% AVE.)	REAS. MAX. (% AVE.)	REAS. MIN. (% AVE.)	PEAK FLOW (CFS)	PEAK DATE	LOW FLOW (CFS)	LOW DATE
MADISON RIVER near Grayling ¥	APR-JUL APR-SEP	388.0 496.0	420.0 530.0	108 106	123 122	93 92				
MADISON RIVER near McAllister ¥	APR-JUL APR-SEP	672.0 848.0	650.0 810.0	96 95	113 112	81 79				

	RESERVOIR STORAGE		(1000AF)		WATERSHED	SNOWPACK ANA	LYSIS	
RESERVOIR	USEABLE I CAPACITYI I	XX USE THIS YEAR	ABLE STORI LAST YEAR	AGE XX I I AVE+ I	WATERSHED	NO. COURSES AVE.D	THIS YEA	R AS % OF AVERAGE
ENNIS LAKE	41.0	31.4	32.3	35.0	MADISON above HEBGEN	18	119	104
HEBGEN LAKE	378.0	278.5	297.0	233.6	LOWER MADISON	20	101	84
				1	MADISON	38	110	95

≖Corrected for upstream diversions or changes in reservoir storage. Average is for 1961-80 period.

16

-

Gallatin Basin



Mountain snowpack* (inches)

WATER SUPPLY OUTLOOK:

Mountain snowpack continues to be well below average particularly in the Bridger Range and Bozeman-Hyalite Creek areas south of Bozeman. Mountain precipitation was below average for March and some melt occurred at the lower and mid-elevations. Spring and summer streamflows are forecast to be well below average in all drainages. Shortages of irrigation supplies can be expected by late June on smaller low elevation streams and by July on the Gallatin River.

GALLATIN RIVER BASIN

	DIREM	INFLOW FORE	CH315						
FORECAST	20 YR. AVE. (1000AF)	MOST PROBABLE (1000AF)	MOST PROBABLE (% AVE.)	REAS. MAX. (X AVF.)	REAS. MIN. (Z AVE.)	PEAK FLOW (CES)	PEAK	LOW FLOW (CES)	
APR-JUL	464.0	370.0	79	96	64				
APR-SEP	545.0	430.0	78	95	63				
APR-JUL	25.0	19.9	79	96	64				
APR-SEP	29.0	22.4	77	93	62				
APR-JUL	39.0	30.6	78	97	59				
APR-SEP	45.0	35.2	78	98	58				
APR-JUL	523.0	380.0	72	98	48				
APR-SEP	611.0	445.0	72	98	48				
	FORECAST PERIOD APR-JUL APR-SEP APR-JUL APR-SEP APR-JUL APR-SEP	FORECAST 20 YR. AVE. PERIOD (1000AF) APR-JUL 464.0 APR-SEP 545.0 APR-JUL 25.0 APR-JUL 25.0 APR-JUL 39.0 APR-JUL 39.0 APR-SEP 45.0 APR-JUL 523.0 APR-SEP 611.0	FORECAST 20 YR. AVE. PROBABLE MOST PROBABLE PERIOD (1000AF) (1000AF) APR-JUL 464.0 370.0 APR-JUL 25.0 19.9 APR-JUL 25.0 19.9 APR-JUL 37.0 30.6 APR-JUL 39.0 30.6 APR-JUL 35.2 380.0 APR-SEP 611.0 445.0	FORECAST 20 YR. MOST MOST PROBABLE PROBABLE PROBABLE PROBABLE YROBABLE YROBABABLE YROBABABA YROBABA <td>FORECAST 20 YR. AVE. MOST PROBABLE MOST PROBABLE REAS. MAX. PERIOD (1000AF) (1000AF) (2 AVE.) (2 AVE.) APR-JUL 464.0 370.0 79 96 APR-SEP 545.0 430.0 78 95 APR-JUL 25.0 19.9 79 96 APR-SEP 29.0 22.4 77 93 APR-SEP 45.0 35.2 78 98 APR-JUL 523.0 380.0 72 98</td> <td>FORECAST 20 YR. MOST PROEABLE PROEABLE PROEABLE PROEABLE PROEABLE PAN MIN. PERIOD (1000AF) (1000AF) (1000AF) (2 AVE.) (2<td>FORECAST 20 YR. AVE. MOST PROBABLE MOST PROBABLE REAS. PROBABLE REAS. MAX. REAS. HIN. REAS. FLOH (Z AVE.) REAS. HIN. PEAK FLOH (Z AVE.) APR-JUL 464.0 370.0 79 76 64 APR-JUL 464.0 370.0 78 95 63 APR-JUL 25.0 19.9 79 76 64 APR-SEP 29.0 22.4 77 93 62 APR-SEP 45.0 35.2 78 98 58 APR-JUL 523.0 380.0 72 98 48 APR-SEP 611.0 445.0 72 98 48</td><td>FORECAST 20 YR. AVE. MOST PROBABLE MOST PROBABLE REAS. PROBABLE REAS. MAX. REAS. HIN. PEAK FLOH (Z AVE.) PEAK FLOH (Z AVE.)<td>STREMPLOW FORECASTS FORECAST 20 YR. MOST PROBABLE PROBABLE PROBABLE PROBABLE PROBABLE HAX. MIN. FLOH FLOH O(1000AF) PERIOD (1000AF) (1000AF) (2 AVE.) (2 AVE.)<</td></td></td>	FORECAST 20 YR. AVE. MOST PROBABLE MOST PROBABLE REAS. MAX. PERIOD (1000AF) (1000AF) (2 AVE.) (2 AVE.) APR-JUL 464.0 370.0 79 96 APR-SEP 545.0 430.0 78 95 APR-JUL 25.0 19.9 79 96 APR-SEP 29.0 22.4 77 93 APR-SEP 45.0 35.2 78 98 APR-JUL 523.0 380.0 72 98	FORECAST 20 YR. MOST PROEABLE PROEABLE PROEABLE PROEABLE PROEABLE PAN MIN. PERIOD (1000AF) (1000AF) (1000AF) (2 AVE.) (2 <td>FORECAST 20 YR. AVE. MOST PROBABLE MOST PROBABLE REAS. PROBABLE REAS. MAX. REAS. HIN. REAS. FLOH (Z AVE.) REAS. HIN. PEAK FLOH (Z AVE.) APR-JUL 464.0 370.0 79 76 64 APR-JUL 464.0 370.0 78 95 63 APR-JUL 25.0 19.9 79 76 64 APR-SEP 29.0 22.4 77 93 62 APR-SEP 45.0 35.2 78 98 58 APR-JUL 523.0 380.0 72 98 48 APR-SEP 611.0 445.0 72 98 48</td> <td>FORECAST 20 YR. AVE. MOST PROBABLE MOST PROBABLE REAS. PROBABLE REAS. MAX. REAS. HIN. PEAK FLOH (Z AVE.) PEAK FLOH (Z AVE.)<td>STREMPLOW FORECASTS FORECAST 20 YR. MOST PROBABLE PROBABLE PROBABLE PROBABLE PROBABLE HAX. MIN. FLOH FLOH O(1000AF) PERIOD (1000AF) (1000AF) (2 AVE.) (2 AVE.)<</td></td>	FORECAST 20 YR. AVE. MOST PROBABLE MOST PROBABLE REAS. PROBABLE REAS. MAX. REAS. HIN. REAS. FLOH (Z AVE.) REAS. HIN. PEAK FLOH (Z AVE.) APR-JUL 464.0 370.0 79 76 64 APR-JUL 464.0 370.0 78 95 63 APR-JUL 25.0 19.9 79 76 64 APR-SEP 29.0 22.4 77 93 62 APR-SEP 45.0 35.2 78 98 58 APR-JUL 523.0 380.0 72 98 48 APR-SEP 611.0 445.0 72 98 48	FORECAST 20 YR. AVE. MOST PROBABLE MOST PROBABLE REAS. PROBABLE REAS. MAX. REAS. HIN. PEAK FLOH (Z AVE.) PEAK FLOH (Z AVE.) <td>STREMPLOW FORECASTS FORECAST 20 YR. MOST PROBABLE PROBABLE PROBABLE PROBABLE PROBABLE HAX. MIN. FLOH FLOH O(1000AF) PERIOD (1000AF) (1000AF) (2 AVE.) (2 AVE.)<</td>	STREMPLOW FORECASTS FORECAST 20 YR. MOST PROBABLE PROBABLE PROBABLE PROBABLE PROBABLE HAX. MIN. FLOH FLOH O(1000AF) PERIOD (1000AF) (1000AF) (2 AVE.) (2 AVE.)<

STREAMELOW FORECASTS

	RESERVOIR STORAGE		(1000AF)		WATERSHE) SNOWPACK AN	ALYSIS	
RESERVOIR	USEABLE CAPACITY 	XX US THIS YEAR	EA&LE STOR Last Year	AGE XX	WATERSHED	NO. COURSES AVE.D	THIS YE	AR AS % OF
MIDDLE CREEK	8.0	5.9	3.7	3.9	UPPER GALLATIN EAST GALLATIN GALLATIN	14 13 24	102 90 97	80 67 74

≭Corrected for upstream diversions or changes in reservoir storage. Average is for 1961-80 period.

Missouri Basin



Precipitation* (percent of normal)

WATER SUPPLY OUTLOOK:

Warm temperatures causing snowmelt and below average mountain precipitation during March combined to drop snowpack percentages about 10 percent since March 1. Snowpacks vary from near average to well below average. Spring and summer streamflows are forecast to be below average from all drainages. Shortages of irrigation water supplies can be expected by late June on lower elevation streams and by early July on most other drainages for those not having stored water.

FOREPACT POINT	FORECAST	ZO YR.	MOST	MOST	FEAS.	REAS.	PEAK	PEAK	LOW	LOw
FUNELASI FUINI	FERIOO	(1000AF)	(1000AF)	(% AVE.)	(% AVE.)	(% AVE.)	(CFS)	DATE	FLON (CFS)	OATE
MISSOURI RIVER at Toston ×	APR-JUL	2196.0	1900.0	86	123	60				
	APR-SEP	2545.0	2235.0	87	125	61				
SHEEP CREEK or White Sulphur Spgs.	APR-JUL	19.0	17.3	91	132	53				
	APR-SEP	22.0	20.0	90	127	55				
BELT CREEK near Monarch	APR~JUL	123.0	101.0	82	116	48				
	AFR-SEP	134.0	110.0	82	116	48				
MISSOURI RIVER at Fort Benton #	APR-JUL	3468.0	2825.0	81	125	51				
	APR-SEP	3980.0	3345.0	84	128	54				
MISSOURI RIVER at Virgelle #	APR-JUL	4030.0	3180.0	78	126	48				
HISSUURI RIVER at VIIgelle =	APR-SEP	4570.0	3735.0	81	129	51				
MISSOURI RIVER near Landusky #	APR-JUL	4383.0	3512.0	- 80	129	48				
	APR-SEP	4980.0	4135.0	83	132	51				
N.F. MUSSELSHELL near Delpine	APR-JUL	5.4	5.3	98	130	56				
	APR-SEP	6.4	6.2	96	141	63				
S.F. MUSSELSHELL above Martinsdale	APR-JUL	59.0	50.0	84	122	47				
	APR-SEP	63.0	52.0	82	121	44				
MISSOURI FIVER below Fort Peck *	APR-JUL	4428.0	3454.0	78	125	47				
	AFR-SEP	4961.0	4030.0	81	132	48				
LAKE SAKAKAWEA Inflow *	APR-JUN	12239+0	11385.0	93	135	63				
	APR-SEP	12775.0	11860.0	92	136	63				

STREAMFLOW FORECASTS

	RESERVOIR STORAGE		(1000AF)	
RESERVOIR	USEA&LE I ** USEA&LE STOFAGE ** I CAPACITYI THIS LAST I WATERSHED			
		TEAR	YEAR	AVE.
CANYON FERRY LAKE	2043.0	1487.0	1394.0	1498.0
HELENA VALLEY	10.4	3.3	3.2	4.9
LAKE HELENA	10.4	10.9	10.7	9,8
HAUSER & HELENA	61.9	63.0	62.4	60.0
HOLTER LAKE	81.9	80.5	78.1	64.9
SMITH RIVER	10.6	7.5	9.6	7.6
NEWLAN CREEK	12.4	10.0	9.0	9.1
BAIR	7.0	3.2	1.2	5.2
MARTINSOALE	23.1	9.8	4.8	9.6
DEADMAN'S BASIN	72.2	37.4	48.0	49.7
FORT PECK LAKE*	18.9	14.2	15.7	15.0
*Million Acre Feet				

*Corrected for upstream diversions or changes in reservoir storage. Average is for 1961-80 period.

Sun, Teton and Marias Basins



WATER SUPPLY OUTLOOK:

Snowpack conditions deteriorated during March. Mountain precipitation for March was below average and melt was occurring at low and mid-elevations. The snowpack is presently well below average and less than it was a year ago. Spring and summer streamflows are forecast to be well below average on all drainages. Shortages of irrigation water supplies can be expected to develop by late June or early July for those users not having stored water.

FORFLAST POINT	FORECAST	20 YR. AVE.	MOST PROBABLE	MOST PROBABLE	REAS.	REAS. MIN.	PEAK	PEAK	LOW	LOW
	PERIOD	(1000AF)	(1000AF)	(% AVE.)	(% AVE.)	(% AVE.)	(CFS)	DATE	(CFS)	DATE
SUN RIVER at Gibson Dam *	APR-JUL	522.0	407.0	77	100	56				
	APR-SEP	570.0	455.0	79	102	58				
NO MEDICINE CREEK near Browning *	APR-JUL	235.0	173.0	73	108	40				
	APR-SEP	248.0	190.0	76	108	45				
ADGER CREEK near Browning	APR-JUL	113.0	87.0	76	111	43				
	APR-SEP	130.0	102.0	78	111	46				
WIFT RESERVOIR Inflow or Dupuyer	APR-JUL	74.7	59.0	78	112	46				
	APR-SEP	86.7	69.0	79	112	47				
UT BANK CREEK at Cut Bank	APR-JUL	108.0	75.5	69	104	36				
	APR-SEP	114.0	82.0	71	104	40				
MARIAS RIVER near Shelby	APR-JUL	518.0	365.0	70	103	38				
	APR-SEP	542.0	385.0	71	103	39				

STREAMFLOW FORECASTS

	RESERVOIR STORAGE		(1000AF)		WATERSHED	SNOWPACK ANA	ALYSIS	
RESERVOIR	USEABLE I CAPACITYI I	** USE THIS YEAR	ABLE STOR LAST YEAR	AGE ##	WATERSHED	ND. COURSES AVE.D	THIS Y	EAR AS % DF
GIBSON	99.1	72.6	55,2	46.2	SUN-TETON	12	71	67
PISHKUN	32.0	18.0	18.5	18,2	MARIAS	7	84	76
WILLOW CREEK	32.2	26.2	13.4	22.1	SUN-TETON-MARIAS	18	78	72
LOWER TWO MEDICINE LAKE		NO REPO	RT					
FOUR HORNS LAKE		NO REPO	RT	1				
SWIFT	30.0	9,9	10.7	16.8				
LAKE FRANCES	112.0	94.5	24.8	71.2				
LAKE ELWELL (TIBER)	1347.0	784.8	680.9	562.3 				

*Corrected for upstream diversions or changes in reservoir storage. Average is for 1961-80 period.

St. Mary and Milk Basins



WATER SUPPLY OUTLOOK:

Below average March precipitation and warm temperatures reduced snowpack levels. Snow in mountains away from the Continental Divide has melted except for shaded high elevation areas. Spring and summer runoff is forecast to be well below average. However, reservoir storage is above average as a result of earlier runoff. Shortages of irrigation water supplies can be expected by mid to late June for those users not having stored water.

STREAMFLOW FORECASTS										
FORECAST POINT	FORECAST PERIOD	20 YR. AVE. (1000AF)	MOST PROEABLE (1000AF)	MOST FROEAELE (% AVE.)	REAS. MAX. (% AVE.)	REAS. MIN. (% AVE.)	PEAK FLOW (CFS)	PEAK DATE	LOW FLOW (CFS)	LOM DATE
SWIFTCURRENT CREEK at Sherburne ¥	APR-JUL APR-SEP	112.0 128.0	78.1 95.5	69 74	90 95	50 55				
ST. MARY RIVER near Babb ≖	APR-JUL APR-SEP	416.0 487.0	288.0 345.0	69 70	83 85	55 57				
MILK RIVER at Eastern Crossing #	APR-SEP	248.0	218.0	87	124	75				
MILK RIVER at Eastern Crossing	APR-SEP	81.7	33.7	41	77	28				

	RESERVOIR STORAGE	(1000AF) 		1	WATERSHED SNOWPACK ANALYSIS					
RESERVOIR	USEABLE CAPACITY 	** USE THIS YEAR	EABLE STOR LAST YEAR	AGE ## i i AVE. i	WATERSHED	NO. COURSES AVE.D	THIS YEA	R AS % OF AVERAGE		
LAKE SHERBURNE	64.3	50.5	31.9	24.0	MILK HEADWATERS	5	57	56		
FRESNO	127.0	99.7	16.3	86.7 i	BEAR PAN	6	4	6		
BEAVER CREEK	3.5	3.3	1+1	2.1	MILK RIVER	11	45	48		
NELSON	66.8	49.4	15.9	38.7	ST. MARY	12	54	52		
				1	ST. MARY and MILK	18	48	48		
				1	BOW RIVER in ALBERTA	18	136	117		
				i i	OLDMAN RIVER in ALBERTA	11	55	56		

■Corrected for upstream diversions or changes in reservoir storage. Average is for 1961-80 period.

Yellowstone Basin

Precipitation* (percent of normal)

Mountain snowpack* (inches)



WATER SUPPLY OUTLOOK:

Snowpack deteriorated during March because of below average mountain precipitation and some snowmelt. Southern drainages have near average snowpack, decreasing to well below average in the northern drainages. Except for below average runoff from streams flowing out of the Crazy and Bridger Mountains, streamflows are expected to be near to a little below average. Irrigation water is expected to be short from streams out of the Crazy and Bridger Mountains but adequate elsewhere.

YELLOWSTONE RIVER BASIN

FORECAST POINT	FORECAST	20 YR. AVE.	MOST PROBABLE	HOST PROBABLE	REAS. MAX.	REAS. MIN.	FEAK FLOW	PEAK	LOH FLOH	LOW
	PERIOO	(1000AF)	(1000AF)	(% AVE.)	(% AVE.)	(% AVE.)	(CFS)	DATE	(CFS)	DATE
YELLOWSTONE at Lake Outlet	APR-SEP	826.0	900.0	108	122	96				
YELLOWSTONE at Corwin Springs	APR-JUL	1686.0	1590.0	94	108	80				
	APR-SEP	2027.0	1900.0	93	108	80				
YELLOWSTONE near Livingston	APR-JUL	1969.0	1817.0	92	106	78				
	APR-SEP	2379.0	2190.0	92	106	78				
BOULDER RIVER at Big Timber	APR-JUL	366.0	348.0	95	117	73				
	APR-SEP	398.0	370.0	92	115	71				
STILLWATER RIVER nr Absarokee 🕷	APR-JUL	528.0	547.0	103	134	74				
	APR-SEP	632.0	650.0	102	133	73				
CLARKS FORK RIVER near Belfry	APR-JUL	563.0	615.0	109	134	84				
	APR-SEP	628.0	700.0	111	136	86				
CODNEY RESERVOIR Inflow	APR-JUL	49.5	41.4	83	113	55				
	APR-SEP	60.5	50.8	83	114	55				
YELLOWSTONE RIVER at Billings	APR-JUL	3833.0	3710.0	96	119	79				
	APR-SEP	4516.0	4340.0	96	118	78				
BIGHORN RIVER near St. Xavier 🕷	APR-JUL	1794.0	2370.0	132	179	100				
	APR-SEP	1976.0	2620.0	132	180	101				
LITTLE BIGHORN RIVER near Hardin	APR-JUL	162.0	185.0	114	172	69				
	APR-SEP	182.0	207.0	113	171	69				
TONGUE RIVER near Oecker	APR-JUL	244.0	260.0	106	159	48				
	APR-SEP	269.0	290.0	107	161	49				
YELLOWSTONE RIVER at Miles City 🕷	APR-JUL	5906.0	6200.0	104	137	82				
	APR-SEP	6787.0	7180.0	105	138	83				
POWDER RIVER at Moorehead	APR-JUL	243.0	255.0	104	160	42				
	APR-SEP	263.0	276.0	104	160	42				
YELLOWSTONE RIVER near Sidney #	APR-JUL	6544.0	6870.0	104	141	79				
	APR-SEP	7518.0	7930.0	105	141	79				

STREAMFLOW FORECASTS

	RESERVOIR STORAGE (1000AF) I HATERSHED						NOWPACK ANALYSIS			
RESERVOIR	USEABLE I ** USEABLE STORAGE ** I RESERVOIR CAPACITYI THIS LAST I WATERSHED I YEAR YEAR AVE. I		WATERSHED	ND. Courses Ave.d	THIS YEA	R AS % OF				
MYSTIC LAKE	21.0	0.3	1.0	4.2	YELLOWSTONE ab LIVINGSTON	26	126	101		
COONEY	27.4	22.0	21.7	15.8	SHIELDS	10	86	65		
BIGHORN LAKE	1356.0	709.6	866.7	607.2	BOULDER-STILLWATER	12	104	88		
TONGUE RIVER	68.0	30.2	16.2	41.6	CLARK'S FORK-ROCK CREEK	22	128	102		
				į	YELLOWSTONE above BIGHORN	56	113	91		
				ļ	LITTLE BIGHORN	5	135	103		
				ļ	WIND RIVER (Wyoming)	28	197	147		
					BIGHORN RIVER (Hyoming)	34	150	116		
				1	BIGHORN BASIN (Total)	58	160	123		
					TONGUE RIVER (Hvoming)	15	133	108		
				1	FOWDER RIVER (Hvoming)	15	148	110		
				1	YELLOWSTONE RIVER	125	133	104		

*Corrected for upstream diversions or changes in reservoir storage. Average is for 1961-80 period.





The Following Organizations Cooperate With The Soil Conservation Service In Snow Survey Work

Canadian	Department of the Environment Atmospheric Environment Service Water Management Service British Columbia Ministry of Environment Inventory and Engineering Branch, Hydrology Section Alberta Environment Technical Services Division
Federal	 U.S. Department of Agriculture Forest Service U.S. Department of the Army Corps of Engineers U.S. Department of Commerce NOAA, National Weather Service NAtional Environmental Satellite Service U.S. Department of the Interior Bureau of Indian Affairs Fish and Wildlife Service Geological Survey National Park Service Bureau of Reclamation U.S. Department of Energy Bonneville Power Administration
State	Montana Conservation Districts Montana Department of Fish, Wildlife, and Parks Montana Department of Natural Resources and Conservation Montana Department of State Lands Montana State University - Agricultural Experiment Station University of Montana - School of Forestry
Private	Big Sky of Montana Butte Water Company Flathead Valley Community College Montana Power Company Pondera County Canal & Reservoir Company
	Other organizations and individuals furnish information for the snow survey reports. Their cooperation is gratefully acknowledged.

UNITED STATES DEPARTMENT OF AGRICULTURE

SOIL CONSERVATION SERVICE

SNOW SURVEY UNIT

Federal Bldg., Rm. 443 10 East Babcock Street Bozeman, MT 59715

OFFICIAL BUSINESS PENALTY FOR PRIVATE USE, \$300 THIRD-CLASS BULK RATE POSTAGE AND FEES PAID USDA - SCS

PERMIT NO G-267

THIRD CLASS MAIL

Montana Water Supply Outlook

and

Federal-State-Private Cooperative Snow Surveys



SOIL CONSERVATION SERVICE