

Research



Cite this article: Rojas-Sereno ZE, Streicker DG, Suarez-Yana T, Lineros M, Yung V, Godreuil S, Benavides JA. 2023 Detection of antimicrobial-resistant Enterobacterales in insectivorous bats from Chile. *R. Soc. Open Sci.* **10**: 231177. <https://doi.org/10.1098/rsos.231177>

Received: 9 August 2023

Accepted: 3 October 2023

Subject Category:

Ecology, conservation and global change biology

Subject Areas:

health and disease and epidemiology/
microbiology

Keywords:

Tadarida brasiliensis, antimicrobial resistance, *Rahnella aquatilis*, urban, extended-spectrum beta-lactamase, Latin America

Authors for correspondence:

Zulma Esperanza Rojas-Sereno

e-mail: z.rojassereno@uandresbello.edu

Julio A. Benavides

e-mail: julio.benavides@ird.fr

Detection of antimicrobial-resistant Enterobacterales in insectivorous bats from Chile

Zulma Esperanza Rojas-Sereno¹, Daniel G. Streicker^{2,3}, Tania Suarez-Yana¹, Michelle Lineros⁴, Verónica Yung⁴, Sylvain Godreuil^{5,6,7} and Julio A. Benavides^{1,7}

¹Centro de Investigación para la Sustentabilidad y Doctorado en Medicina de la Conservación /Facultad Ciencias de la Vida, Universidad Andrés Bello, República 440, Santiago 8320000, Chile

²School of Biodiversity, One Health and Veterinary Medicine, University of Glasgow, Glasgow G12 8QQ, UK

³MRC-University of Glasgow Centre for Virus Research, Glasgow G61 1QH, UK

⁴Sección Rabia, Departamento Laboratorio Biomédico, Instituto de Salud Pública de Chile, Santiago 8320000, Chile

⁵Laboratoire de Bactériologie, Centre Hospitalier Universitaire de Montpellier, Montpellier 34295, France

⁶Laboratoire Mixte International, DRISA, IRD, Montpellier 34394, France

⁷MIVEGEC, IRD, CNRS, Université de Montpellier, Montpellier 34394, France

ZER-S, 0000-0003-1030-3876; DGS, 0000-0001-7475-2705; JAB, 0000-0002-3920-9165

Enterobacterales of clinical importance for humans and domestic animals are now commonly detected among wildlife worldwide. However, few studies have investigated their prevalence among bats, particularly in bat species living near humans. In this study, we assessed the occurrence of Extended-spectrum beta-lactamase-producing (ESBL) and carbapenemase-resistant (CR) Enterobacterales in rectal swabs of bats submitted to the Chilean national rabies surveillance program from 2021 to 2022. From the 307 swabs screened, 47 (15%) harboured cefotaxime-resistant Enterobacterales. Bats carrying these bacteria originated from 9 out of the 14 Chilean regions. Most positive samples were obtained from *Tadarida brasiliensis* ($n = 42$), but also *Lasiurus varius*, *L. cinereus* and *Histiotus macrotus*. No Enterobacterales were resistant to imipenem. All ESBL-Enterobacterales were confirmed as *Rahnella aquatilis* by MALDI-TOF. No other ESBL or CR Enterobacterales were detected. To our knowledge, this is the first screening of antibiotic-resistant bacteria in wild bats of Chile, showing the bat faecal carriage of *R. aquatilis* naturally resistant to cephalosporins, but also including acquired resistance to important antibiotics for public health such as amoxicillin with clavulanic acid. Our results suggest unknown selective pressures on *R. aquatilis*, but low or no carriage of ESBL or CR *Escherichia coli* and *Klebsiella*

spp. Future studies should assess the zoonotic and environmental implications of *R. aquatilis*, which are likely present in the guano left by bats roosting in human infrastructures.

1. Introduction

Antimicrobial resistance (AMR) circulating at the human–domestic–wildlife interface is a global threat to public health, causing at least 1.27 million human deaths per year [1,2]. Extended-spectrum beta-lactamase-producing (ESBL) and carbapenemase-resistant (CR) Enterobacterales including *Escherichia coli* and *Klebsiella pneumoniae* represent the highest burden of deaths attributed to AMR in humans. Antimicrobial-resistant bacteria are carried by wildlife worldwide, and are often considered as an indicator of wildlife exposure to anthropogenic pathogen sources [2–9]. Wildlife can act as sentinels of ‘human pathogen pollution’ in natural environments, but could also become potential reservoirs of AMR to humans or domestic animals [4,7,10,11]. Bats (order Chiroptera) are considered one of the main reservoirs of human infectious diseases, because they harbour a large number of zoonotic viral pathogens, and their global distribution widely overlaps with humans [12]. However, the role of bats in the dissemination of AMR remains still poorly understood [7].

Recent studies have investigated the occurrence of AMR among bats worldwide, mostly reporting the faecal carriage of ESBL-*E. coli* [7,13–19]. For example, ESBL-*E. coli* have been reported in *Eidolon helvum*, *Megaloglossus woermanni* and *Nycteris hypsipida* in Africa and *Tadarida teniotis* in Europe [18,20,21]. In Latin America, ESBL-*E. coli* have been reported in frugivorous bats including *Artibeus planirostris* and *Sturnina lilium* of Brazil, the omnivorous *Phyllostomus hastatus* in Trinidad, insectivorous bats (*Pteronotus parnelli*, *Eptesicus tadeii* and *Molossus* spp.) in Brazil and Trinidad, and the hematophagous *Desmodus rotundus* in Peru and Brazil [7,13–15,22]. ESBL and CR-*E. coli* have also been reported in the frugivorous bats *Artibeus lituratus* and *Carollia perspicillata* in Brazil [22]. However, little is known about ESBL-*E. coli* in other non-tropical regions of Latin America including Chile.

Antimicrobial-resistant bacteria found in bats are commonly hypothesized to be attributable to contamination from humans or livestock [13,19]. However, few studies have compared the prevalence of these bacteria among natural, rural and urban areas [7,18,20–22]. As such, the role of human activities and the ability of bats to spread AMR within their populations remains poorly understood. In Chile, ESBL-*E. coli* have been isolated from wild rodents, gulls, and condors [23–26], but no study has been conducted on bats. Chile hosts several insectivorous bats inhabiting urban and rural areas that are subject to the passive rabies surveillance program of the National Institute of Health [27]. Among insectivorous bats submitted to the surveillance program, *T. brasiliensis* is one of the bat species most closely associated with humans [28]. In this study, we aimed to estimate and compare the prevalence of ESBL and CR-Enterobacterales among Chilean bats from rural and urban areas.

2. Material and methods

We collected 307 rectal samples from dead bats that were submitted to the National Rabies Surveillance Program of the Institut of Public Health (Instituto de Salud Pública, ISP) in Chile from February 2021 to August 2022. Bats studied originated from 14 geographical regions and belonged to the genera *Tadarida*, *Histiotus*, *Lasiurus*, and *Myotis*. Morphological identification of bat individuals was performed by ISP staff, and was based on key distinctive features of each species. All bats studied here were negative for rabies according to the direct fluorescent antibody test. Samples were stored in Stuart transport media at 4°C until being screened in MacConkey selective media supplemented with antibiotics to isolate ESBL and CR Enterobacterales. Samples were also grown on a MacConkey media without antibiotics (negative control), to confirm the presence of Enterobacterales on the faecal sample. Screening was made by direct incubation at 37°C for 24 h on MacConkey media supplemented with cefotaxime sodium salt (CTX) at 2 mg/ml for ESBL-Enterobacterales and imipenem (IM) at 4 mg/ml for CR-Enterobacterales screening, following previously published methods [7,13]. Samples presenting bacterial growth were purified in the same medium to confirm growth and stored at –80°C. Bacterial species confirmation was performed on 32 randomly chosen ESBL isolates among a total of isolates with similar morphology and antibiotic-resistant phenotypic profile, using the matrix-assisted laser desorption ionization-time of flight (MALDI-TOF) mass spectrometry using the MALDI Biotyper database DB 8326 MSP (Bruker Daltonics, Bremen, Germany) in the Laboratory of Microbiology of the Hospital Arnaud

de Villeneuve in Montpellier (France). Susceptibility tests were performed using the disc diffusion method on Müller-Hinton agar following the European Committee on Antimicrobial Susceptibility Testing (EUCAST) guidelines (Version 7.1, 2017). Twenty-seven antibiotics were analysed including penicillin (ampicillin, amoxicillin, ticarcillin, piperacillin and temocillin), penicillin with β -lactamase inhibitors (amoxicillin-clavulanic acid), antipseudomonal penicillin with β -lactamase inhibitors (ticarcillin-clavulanic acid and piperacillin-tazobactam), non-extended spectrum cephalosporins (cephalexin), extended-spectrum cephalosporins (cefotaxime, ceftazidime, cefepime and cefpodoxime), cephamycin (cefotixin), carbapenems (imipenem, ertapenem, meropenem), monobactams (aztreonam), quinolones (ofloxacin, ciprofloxacin and levofloxacin), aminoglycosides (gentamicin, tobramycin and amikacin), folate pathway inhibitors (trimethoprim-sulfamethoxazole), phenicol (chloramphenicol) and phosphonic acids (fosfomycin). We divided antibiotics into the 'categories' defined in Magiorakos *et al.* [29].

To evaluate the difference in occurrence and prevalence between bats inhabiting rural and urban areas, we conducted two complementary statistical tests. First, each municipality was assigned as rural or urban following the Chilean 'Comisión Interministerial de Ciudad, Vivienda y Territorio-COMICIVYT' (<https://www.masvidarural.gob.cl/ruralidad-en-chile/>). A municipality was assigned as urban if more than 75% of its population were living in human densities higher than 150 inhabitants by km². A Chi-square test was then used to test the overall difference in prevalence between samples originated from urban or rural municipalities using the `prop.test` function in R, or a Fisher's exact-test if the number of samples was small (`fisher.test` function). We also used a continuous variable of population density to evaluate the effect of urban areas by conducting a general linear mixed-effect model with binomial residual distribution with the `glmer` function [30]. The model included the dependent variable 'presence/absence of AMR', and year, the municipality's population density (log-transformed to increase normality and avoid outliers) or total population (square-root transformation) as independent variables (each tested in a separated model given their correlation), and the municipality within region as random effects. Chile was divided by regions and each municipality was assigned to a region using the shapefile from Biblioteca Del Congreso Nacional de Chile (https://www.bcn.cl/siit/mapas_vectoriales/index_html). The human population size and density per municipality were obtained from the Comisión Interministerial de Ciudad, Vivienda y Territorio-COMICIVYT. Since bats were collected on private property, no GPS coordinates were available in order to comply with the confidential agreement established with the ISP.

3. Results

From 307 faecal samples collected out of 14 Chilean regions ($n = 238$ from urban municipalities, $n = 69$ from rural municipalities, figure 1a), 47 bats (15%) carried CTX-resistant Enterobacterales, including 42 individuals of *Tadarida brasiliensis* (15% of AMR prevalence, 42 out of 272 *T. brasiliensis* individuals), defined as at least one isolate of CTX-resistant Enterobacterales. Two individuals of *Lasiurus varius* and one individual of *Histiotus macrotus* and *L. cinereus* also carried these bacteria (table 1). No lactose-positive Enterobacterales were resistant to imipenem. AMR-positive samples were obtained from 9 out of the 14 (64.3%) Chilean regions sampled. There was no significant difference in the prevalence between regions (Fisher's exact test: p -value = 0.07, figure 1b). AMR prevalence in urban municipalities was 11% (35 out of 307) and 17% (12 out of 69) in rural municipalities, with no significant difference between them (Chi-squared test: $\chi^2 = 0.13$, p -value = 0.72, figure 1c). Neither municipality's human population size (GLMM, AIC = 249, Estimate = -1.38×10^{-3} , p -value = 0.16) or population density (GLMM, AIC = 251, Estimate = 0.01×10^{-5} , p -value = 0.87) was significantly correlated to the presence/absence of AMR in bats. All 32 ESBL-Enterobacterales isolates (out of 47 isolates having similar morphology and antibiotic-resistant profile) were identified as *Rahnella aquatilis*. No ESBL-*E. coli* or *Klebsiella* sp. were detected. *Rahnella aquatilis* isolates were resistant to seven antimicrobial categories; 65% were resistant to five antimicrobial categories, followed by six (19%) and four (16%) categories. Resistance to antimicrobial categories included β -lactams and cephalosporins (100% to penicillin and non-extended β -lactams cephalosporins; 60% to extended-spectrum-beta-lactamases), penicillin with β -lactamase inhibitors (68% to amoxicillin with clavulanic acid), phosphonic acid (60%) and chloramphenicol (13%) (figure 2). One isolate was resistant to piperacillin with tazobactam.

4. Discussion

The circulation of AMR in bats remains poorly understood, but recent reports have detected the presence of AMR in several bat species living in close proximity to humans and domestic animals [7,13,18]. In this study, we found cephalosporin-resistant *R. aquatilis* in three of the four genera of bats collected, mostly




Figure 1. Spatial distribution of antimicrobial-resistant Enterobacteriales among insectivorous bats from Chile: (a). Map of Chile including sampled municipalities (b). Bar plot represents the proportion of AMR Enterobacteriales in insectivorous bats per region. The total number of individual bats sampled per region (n) is shown at the top of each confidence interval bar (estimated using the fisher.test function in R). (c) Zoom map of central Chile showing municipalities with bat samples that harboured antimicrobial resistant bacteria (purple) or not (yellow). (d). Bar plot comparing the proportion of AMR between rural and urban municipalities. A municipality was assigned as urban if more than 75% of its population was living in densities higher than 150 inhabitants per km².

Table 1. Prevalence of cefotaxime (CTX)-resistant Enterobacteriales among insectivorous bats of Chile.

bat genera		bat species			CTX-resistant	
name	total	species	total	%	number	prevalence
<i>Tadarida</i>	272	<i>T. brasiliensis</i>	272	88.3	42	13.6
<i>Histiotus</i>	26	<i>H. macrotus</i>	24	7.8	1	0.3
		<i>H. montanus</i>	1	0.3	0	0.0
		<i>Histiotus</i> sp.	1	0.3	0	0.0
<i>Lasiurus</i>	6	<i>L. cineris</i>	1	0.3	1	0.3
		<i>L. varius</i>	4	1.3	2	0.6
		<i>L. villosissimus</i>	1	0.3	0	0.0
<i>Myotis</i>	3	<i>M. atacamensis</i>	1	0.3	0	0.0
		<i>M. chiloensis</i>	2	0.6	0	0.0
unknown	1	-	1	0.3	1	0.3
total			308		47	
%					15.3	

belonging to *T. brasiliensis*, one of the bat species with the closest association with humans in Chile and Latin America [28]. However, the lack of isolation of ESBL-*E. coli* or *Klebsiella* suggests a low or non-existent circulation of these bacteria among insectivorous bats of Chile living in proximity to humans.

Additionally, field sample collection of live bats is required to confirm whether the observed pattern detected among dead bats of the surveillance program reflects the circulation of AMR among healthy wild bats. Finally, given our inability to obtain GPS coordinates for the location of the surveyed bats, a more refined definition of an urban/rural area where bats were found (e.g. households) was not possible. Thus, our definition of a bat as collected on an ‘urban’ or ‘rural’ area could be poorly resolved for municipalities with high spatial internal heterogeneity, but this will also depend on the extent of bat foraging within a municipality, which could expose them to most of this heterogeneity.

5. Conclusion

To our knowledge, this is the first screening of antibiotic-resistant bacteria in bats from Chile. We identified the faecal carriage of ESBL-*R. aquatilis* in bats, but the absence of faecal carriage of ESBL or CR-*E. coli*. Although *R. aquatilis* has been reported in bats and is naturally resistant to certain antimicrobials, resistance to β -lactamase inhibitors and chloramphenicol suggests potential mechanisms selecting for AMR on these urban and rural bats. Future studies should assess the zoonotic and public health implications of bacteria such as *R. aquatilis*, which could be largely present in the guano left by these bats in the environment.

Ethics. This study was approved by the Ethics Committee from the Facultad Ciencias de la Vida, Universidad Andrés Bello, Santiago, Chile (036-2020).

Data accessibility. Rural or urban landscape, total population number and population density per municipality: Chilean ‘Comisión Interministerial de Ciudad, Vivienda y Territorio-COMICIVYT’ (<https://www.masvidarural.gob.cl/ruralidad-en-chile/>). Region shapefile: Biblioteca Del Congreso Nacional de Chile (https://www.bcn.cl/siit/mapas_vectoriales/index.html). Bat information was obtained from the National Rabies surveillance program of the Instituto de Salud Pública (ISP) in Chile. Data can be publicly requested directly from ISP.

Declaration of AI use. We have not used AI-assisted technologies in creating this article.

Authors’ contributions. Z.E.R.-S.: conceptualization, data curation, formal analysis, funding acquisition, investigation, methodology, project administration, resources, software, validation, visualization, writing—original draft, writing—review and editing; D.G.S.: conceptualization, supervision, writing—original draft, writing—review and editing; T.S.-Y.: methodology, writing—review and editing; M.L.: data curation, resources, writing—review and editing; V.Y.: data curation, resources, writing—review and editing; S.G.: methodology, writing—review and editing; J.A.B.: conceptualization, formal analysis, funding acquisition, methodology, resources, software, supervision, validation, writing—original draft, writing—review and editing.

All authors gave final approval for publication and agreed to be held accountable for the work performed therein.

Conflict of interest declaration. We have no competing interests.

Funding. Laboratory work and writing work were funded by the National Agency for Research and Development (ANID) FONDECYT Iniciación 11181017, awarded to J.B. and Beca Doctorado Nacional 21201076, awarded to Z.E.R.-S., D.G.S. was supported by a Wellcome Trust Senior Research Fellowship (grant no. 217221/Z/19/Z).

Acknowledgements. We thank members of ISP for contributing to the collection of samples. We thank Marilia Salgado-Caxito and members of the Monkey Lab for useful help and comments on microbiological methods.

References

- Stanford K, Zaheer R, Klima C, McAllister T, Peters D, Niu YD, Ralston B. 2020 Antimicrobial resistance in members of the bacterial bovine respiratory disease complex isolated from lung tissue of cattle mortalities managed with or without the use of antimicrobials. *Microorganisms* **8**, 288. (doi:10.3390/MICROORGANISMS8020288)
- Murray CJ *et al.* 2022 Global burden of bacterial antimicrobial resistance in 2019: a systematic analysis. *Lancet* **399**, 629–655. (doi:10.1016/S0140-6736(21)02724-0)
- Allen HK, Donato J, Wang HH, Cloud-Hansen KA, Davies J, Handelsman J. 2010 Call of the wild: antibiotic resistance genes in natural environments. *Nat. Rev. Microbiol.* **8**, 251–259. (doi:10.1038/nrmicro2312)
- Vittecoq M *et al.* 2016 Antimicrobial resistance in wildlife. *J. Appl. Ecol.* **53**, 519–529. (doi:10.1111/1365-2664.12596)
- Ramey AM, Ahlstrom CA. 2020 Antibiotic resistance bacteria in wildlife Perspectives on trend, acquisition and dissemination, data gaps, and future directions. *J. Wildl. Dis.* **56**, 1. (doi:10.7589/2019-04-099)
- Darwich L, Vidal A, Seminati C, Albamonte A, Casado A, López F, Molina-López RA, Migura-García L. 2019 High prevalence and diversity of extended-spectrum β -lactamase and emergence of OXA-48 producing Enterobacterales in wildlife in Catalonia. *PLoS ONE* **14**, e0210686. (doi:10.1371/JOURNAL.PONE.0210686)
- Benavides JA, Godreuil S, Opazo-Capurro A, Mahamat OO, Falcon N, Oravcova K, Streicker DG, Shiva C. 2022 Long-term maintenance of multidrug-resistant *Escherichia coli* carried by vampire bats and shared with livestock in Peru. *Sci. Total Environ.* **810**, 152045. (doi:10.1016/j.scitotenv.2021.152045)
- Ballash GA, Dennis PM, Mollenkopf DF, Albers AL, Robison TL, Adams RJ, Li C, Tyson GH, Wittum TE. 2022 Colonization of White-Tailed Deer (*Odocoileus virginianus*) from Urban and Suburban Environments with Cephalosporinase- and Carbapenemase-Producing Enterobacterales. *Appl. Environ. Microbiol.* **88**, e00465-22. (doi:10.1128/aem.00465-22)
- Guenther S, Grobbel M, Lübke-Becker A, Goedecke A, Friedrich ND, Wieler LH, Ewers C. 2010 Antimicrobial resistance profiles of *Escherichia coli* from common European wild bird species. *Vet. Microbiol.* **144**, 219–225. (doi:10.1016/j.vetmic.2009.12.016)

10. Zeballos-Gross D, Rojas-Sereno Z, Salgado-Caxito M, Poeta P, Torres C, Benavides JA. 2021 The Role of Gulls as Reservoirs of Antibiotic Resistance in Aquatic Environments: A Scoping Review. *Front. Microbiol.* **12**, 1938. (doi:10.3389/FMICB.2021.703886/BIBTEX)
11. Ahlstrom CA *et al.* 2021 Evidence for continental-scale dispersal of antimicrobial resistant bacteria by landfill-foraging gulls. *Sci. Total Environ.* **764**, 144551. (doi:10.1016/j.scitotenv.2020.144551)
12. Olival KJ, Hosseini PR, Zambrana-Torrel C, Ross N, Bogich TL, Daszak P. 2017 Host and viral traits predict zoonotic spillover from mammals. *Nature* **546**, 646–650. (doi:10.1038/nature22975)
13. Benavides JA, Shiva C, Virhuez M, Tello C, Appelgren A, Vendrell J, Solassol J, Godreuil S, Streicker DG. 2018 Extended-spectrum beta-lactamase-producing *Escherichia coli* in common vampire bats *Desmodus rotundus* and livestock in Peru. *Zoonoses Public Health* **65**, 454–458. (doi:10.1111/zph.12456)
14. Cláudio VC, Gonzalez I, Barbosa G, Rocha V, Moratelli R, Rassy F. 2018 Bacteria richness and antibiotic-resistance in bats from a protected area in the Atlantic Forest of Southeastern Brazil. *PLoS ONE* **13**, e0203411. (doi:10.1371/journal.pone.0203411)
15. Adesiyun AA, Stewart-Johnson A, Thompson NN. 2009 Isolation of enteric pathogens from bats in trinidad. *J. Wildl. Dis.* **45**, 952–961. (doi:10.7589/0090-3558-45.4.952)
16. Devnath P, Karah N, Graham JP, Rose ES, Asaduzzaman M. 2023 Evidence of antimicrobial resistance in bats and its planetary health impact for surveillance of zoonotic spillover events: a scoping review. *Int. J. Environ. Res. Public Heal.* **20**, 243. (doi:10.3390/IJERPH20010243)
17. Garcês A, Correia S, Amorim F, Pereira JE, Igrejas G, Poeta P. 2019 First report on extended-spectrum beta-lactamase (ESBL) producing *Escherichia coli* from European free-tailed bats (*Tadarida teniotis*) in Portugal: a one-health approach of a hidden contamination problem. *J. Hazard. Mater.* **370**, 219–224. (doi:10.1016/j.jhazmat.2017.12.053)
18. Obodoechi LO, Carvalho I, Chenouf NS, Martínez-Álvarez S, Sadi M, Nwanta JA, Chah KF, Torres C. 2021 Antimicrobial resistance in *Escherichia coli* isolates from frugivorous (*Eidolon helvum*) and insectivorous (*Myotis hispida*) bats in Southeast Nigeria, with detection of CTX-M-15 producing isolates. *Comp. Immunol. Microbiol. Infect. Dis.* **75**, 101613. (doi:10.1016/J.CIMID.2021.101613)
19. McDougall FK, Boardman WSJ, Power ML. 2021 Characterization of beta-lactam-resistant *Escherichia coli* from Australian fruit bats indicates anthropogenic origins. *Microb. Genomics* **7**, 571. (doi:10.1099/MGEN.0.000571)
20. Nowakiewicz A, Zięba P, Gnat S, Trościańczyk A, Osińska M, Łagowski D, Kosior-Korzecka U, Puzio I. 2020 Bats as a reservoir of resistant *Escherichia coli*: A methodical view. Can we fully estimate the scale of resistance in the reservoirs of free-living animals? *Res. Vet. Sci.* **128**, 49–58. (doi:10.1016/J.RVSC.2019.10.017)
21. Nguema PPM *et al.* 2020 Characterization of ESBL-Producing Enterobacteria from Fruit Bats in an Unprotected Area of Makokou, Gabon. *Microorganisms* **8**, 138. (doi:10.3390/MICROORGANISMS8010138)
22. Sens-Junior H, Trindade WA, Oliveira AF, Zaniolo MM, Serenini GF, Araujo-Ceranto JB, Gonçalves DD, Germano RM. 2018 Bacterial resistance in bats from the Phyllostomidae family and its relationship with unique health. *Pesqui. Veterinária Bras.* **38**, 1207–1216. (doi:10.1590/1678-5150-PVB-5185)
23. Benavides JA, Salgado-Caxito M, Opazo-Capurro A, Muñoz PG, Piñeiro A, Medina MO, Rivas L, Munita J, Millán J. 2021 ESBL-Producing *Escherichia coli* Carrying CTX-M Genes Circulating among Livestock, Dogs, and Wild Mammals in Small-Scale Farms of Central Chile. *Antibiotics* **10**, 510. (doi:10.3390/ANTIBIOTICS10050510)
24. Fuentes-Castillo D *et al.* 2019 Wild owls colonized by international clones of extended-spectrum β -lactamase (CTX-M)-producing *Escherichia coli* and *Salmonella* Infantis in the Southern Cone of America. *Sci. Total Environ.* **674**, 554–562. (doi:10.1016/j.scitotenv.2019.04.149)
25. Hernandez J *et al.* 2013 Characterization and Comparison of Extended-Spectrum β -Lactamase (ESBL) Resistance Genotypes and Population Structure of *Escherichia coli* Isolated from Franklin's Gulls (*Leucophaeus pipixcan*) and Humans in Chile. *PLoS ONE* **8**, e76150. (doi:10.1371/journal.pone.0076150)
26. Báez J *et al.* 2015 Molecular Characterization and Genetic Diversity of ESBL-Producing *Escherichia coli* Colonizing the Migratory Franklin's Gulls (*Leucophaeus pipixcan*) in Antofagasta, North of Chile. *Microb. Drug Resist.* **21**, 111–116. (doi:10.1089/mdr.2014.0158)
27. Escobar LE, Restif O, Yung V, Favi M, Pons DJ, Medina-Vogel G. 2015 Spatial and temporal trends of bat-borne rabies in Chile. *Epidemiol. Infect.* **143**, 1486–1494. (doi:10.1017/S095026881400226X)
28. Rodríguez-San Pedro A, Allendes JL, Ossa G. 2016 Lista actualizada de los murciélagos de Chile con comentarios sobre taxonomía, ecología, y distribución. *Biodivers. Nat. Hist* **2**, 16–39.
29. Magiorakos AP *et al.* 2012 Multidrug-resistant, extensively drug-resistant and pandrug-resistant bacteria: An international expert proposal for interim standard definitions for acquired resistance. *Clin. Microbiol. Infect.* **18**, 268–281. (doi:10.1111/j.1469-0691.2011.03570.x)
30. Bates D, Mächler M, Bolker BM, Walker SC. 2015 Fitting Linear Mixed-Effects Models Using lme4. *J. Stat. Softw.* **67**, 1–48. (doi:10.18637/JSS.V067.i01)
31. Benavides JA, Rojas Paniagua E, Hampson K, Valderrama W, Streicker DG. 2017 Quantifying the burden of vampire bat rabies in Peruvian livestock. *PLoS Negl. Trop. Dis.* **11**, e0006105. (doi:10.1371/journal.pntd.0006105)
32. Walkers-Rooijackers JCM, Rebmann K, Bosch T, Hazeleger WC. 2019 Fecal Bacterial Communities in Insectivorous Bats from the Netherlands and Their Role as a Possible Vector for Foodborne Diseases. *Acta Chiropterologica* **20**, 475–483. (doi:10.3161/15081109ACC2018.20.2.017)
33. Gerbáčová K, Maliničová L, Kisková J, Maslišová V, Uhrin M, Pistaš P. 2020 The Faecal Microbiome of Building-Dwelling Insectivorous Bats (*Myotis myotis* and *Rhinolophus hipposideros*) also Contains Antibiotic-Resistant Bacterial Representatives. *Curr. Microbiol.* **77**, 2333–2344. (doi:10.1007/S00284-020-02095-Z/TABLES/3)
34. Stock I, Grüger T, Wiedemann B. 2000 Natural Antibiotic Susceptibility of *Rahnella aquatilis* and *R. aquatilis*-Related Strains. *J. Chemother.* **12**, 30–39. (doi:10.1179/JOC.2000.12.1.30)
35. Koczura R, Mokracka J, Makowska N. 2016 Environmental Isolate of *Rahnella aquatilis* Harbors Class 1 Integron. *Curr. Microbiol.* **72**, 64. (doi:10.1007/S00284-015-0917-4)
36. Tash K. 2005 *Rahnella aquatilis* Bacteremia from a Suspected Urinary Source. *J. Clin. Microbiol.* **43**, 2526. (doi:10.1128/JCM.43.5.2526-2528.2005)
37. Ruimy R, Meziane-Cherif D, Momcilovic S, Arlet G, Andreumont A, Courvalin P. 2010 RAHN-2, a chromosomal extended-spectrum class A β -lactamase from *Rahnella aquatilis*. *J. Antimicrob. Chemother.* **65**, 1619–1623. (doi:10.1093/JAC/DKQ178)